

REPELLENCY OF AROMATIC MEDICINAL PLANT EXTRACTS AND A STEAM DISTILLATE TO *Aedes aegypti*

YOUNG-CHEOL YANG,¹ EUN-HAE LEE,² HOI-SEON LEE,³ DONG-KYU LEE⁴ AND YOUNG-JOON AHN^{2,5}

ABSTRACT. The repellent activity of methanol extracts from 23 aromatic medicinal plant species and a steam distillate against female blood-starved *Aedes aegypti* was examined in the laboratory by skin test and compared with that of *N,N*-diethyl-*m*-toluamide (deet). Responses varied according to plant species. At a dose of 0.1 mg/cm², the repellency of extracts of *Cinnamomum cassia* bark (91%), *Nardostachys chinensis* rhizome (81%), *Paeonia suffruticosa* root bark (80%), and *Cinnamomum camphora* steam distillate (94%) was comparable to deet (82%). The duration of the effectiveness for extracts from *C. cassia* bark and *N. chinensis* rhizome was comparable to deet and lasted for ~1 h. Relatively short duration of repellency was observed in *P. suffruticosa* root bark extract and *C. camphora* steam distillate. The plants described merit further study as potential mosquito repellent agents.

KEY WORDS Natural repellent, mosquito, *Aedes aegypti*, aromatic plant, deet, skin test

INTRODUCTION

Mosquito repellents could be one of the most effective tools for protecting humans from mosquito attack and from mosquito-borne diseases, such as dengue hemorrhagic fever, malaria, encephalitis, and filariasis (Curtis et al. 1990). Repeated use of synthetic insecticides for mosquito control has disrupted natural biological control systems and led to resurgences in mosquito populations (Croft and Brown 1975). It has also resulted in the development of resistance (Brown 1986), had undesirable effects on nontarget organisms, and fostered environmental and human health concerns (Hayes and Laws 1991). The most commonly used mosquito repellent is *N,N*-diethyl-*m*-toluamide (deet), which is still most effective. However, this compound has an unpleasant odor, can damage plastics and synthetic rubber, and has high skin penetration characteristics (Qiu et al. 1998). These problems indicate a need for new and improved repellents and strategies for protection from mosquito attack.

Plants could be an alternative source for mosquito repellents because they constitute a potential source of bioactive chemicals (Wink 1993) and typically are free from harmful effects (Isman 1995). Because of this, much interest has focused on plant extracts, or plant essential oils, as potential mosquito repellent agents. The effectiveness and duration of repellency depend on the type of repellent (active ingredient and formulation), mode of application, local conditions, attractiveness of individual people to insects, loss of repellent with removal by

perspiration and abrasion, sensitivity of the insects to repellents, and biting density (Rozendaal 1997).

This paper describes a laboratory study that was made to assess the potential of plant extracts for use as commercial mosquito repellents. Repellent activity of methanol extracts from 23 aromatic medicinal plant species and a steam distillate were assessed against female blood-starved *Aedes aegypti* (L.) and compared with that of deet.

MATERIALS AND METHODS

Insects: The *Ae. aegypti* used in this study were from cultures maintained in the laboratory for 7 years without exposure to insecticide. Adult mosquitoes were reared on a 10% sucrose solution and blood-fed on live mice. Larvae were reared in plastic trays (24 × 35 × 5 cm) containing 2,000 ml of water supplied with 0.5 g of sterilized diet (40-mesh chick chow powder and yeast, 4:1 by weight). They were held at 28 ± 2°C and 75 ± 5% relative humidity (RH) and a 16:8 h photoperiod.

Plants and sample preparation: A total of 24 aromatic medicinal plant species were selected (Namba 1993). The parts of each plant that have been used in Chinese medicine (Namba 1993) were purchased from Boeun medicinal herb shop, Kyungdong market, Seoul, and used in extractions (Table 1). With the exception of *Chaenomeles sinensis* (Thouin) Koehn. and *Cinnamomum camphora* Presl, the plants were dried in an oven at 40°C for 2 days and finely powdered with a blender. Each sample (50 g) was extracted twice with 300 ml of methanol at room temperature for 2 days and filtered. Slices (200 g) of the fresh *Chaenomeles* fruits were ground in a blender, extracted twice each with 900 ml of methanol at room temperature for 1 day, and filtered. The combined filtrate was concentrated to dryness by rotary evaporation at 40°C. The yield of each methanolic extraction is given in Table 1. *Cinnamomum camphora* was purchased as a steam distillate.

Bioassay: The method of Frances et al. (1996)

¹ Department of Advanced Organic Materials Engineering, Chonbuk National University, Chonju 561-756, Republic of Korea.

² School of Agricultural Biotechnology, Seoul National University, Seoul 151-742, Republic of Korea.

³ Faculty of Biotechnology, Chonbuk National University, Chonju 561-756, Republic of Korea.

⁴ Department of Biological Sciences, Kosin University, Busan 606-701, Republic of Korea.

⁵ Author to whom correspondence should be addressed.

Table 1. Aromatic medicinal plants tested.

Family	Species	Tissue used ¹	Yield (%) ²
Apiaceae	<i>Angelica dahurica</i>	Ro	17.7
Araceae	<i>Acorus calamus</i> var. <i>angustatus</i>	Rh	10.1
	<i>Acorus gramineus</i>	Rh	9.5
Compositae	<i>Artemisia princeps</i> var. <i>orientalis</i>	Wp	6.6
	<i>Inula helenium</i>	Ro	16.3
Dioscoreaceae	<i>Dioscorea batatas</i>	Rh	2.4
Fabaceae	<i>Gleditsia horrida</i>	Fr	17.3
	<i>Glycyrrhiza glabra</i>	Ro	21.9
Labiatae	<i>Agastache rugosa</i>	Wp	9.5
	<i>Schizonepeta tenuifolia</i>	Wp	8.1
Lauraceae	<i>Cinnamomum camphora</i> ³	—	—
	<i>Cinnamomum cassia</i>	Ba	5.1
Magnoliaceae	<i>Magnolia obovata</i>	Ba	5.8
Myrtaceae	<i>Eugenia caryophyllata</i>	Fb	37.8
Paeoniaceae	<i>Paeonia suffruticosa</i>	Rb	18.6
Piperaceae	<i>Piper nigrum</i>	Fr	10.1
Polygonaceae	<i>Rheum coreanum</i>	Rh	41.6
Prumulaceae	<i>Lysimachia davurica</i>	Wp	9.0
Rosaceae	<i>Chaenomeles sinensis</i>	Fr	60.8
Rutaceae	<i>Evodia rutaecarpa</i>	Fr	9.5
Solanaceae	<i>Solanum melongena</i>	Fr	47.7
Stemonaceae	<i>Stemona japonica</i>	Ro	15.2
Thymelaeaceae	<i>Aquilaria agallocha</i>	Hw	6.6
Valerianaceae	<i>Nardostachys chinensis</i>	Rh	12.9

¹ Ba, bark; Fb, flower bud; Fr, fruit; Hw, heart wood; Rb, root bark; Rh, rhizome; Ro, root; and Wp, whole plant.

² (Weight of crude methanol extract/weight of dried test material) × 100; except *Chaenomeles sinensis*, (weight of crude methanol extract/weight of fresh fruits) × 100.

³ Steam distillate.

with a slight modification was used to determine the repellent activity of test samples against female blood-starved *Ae. aegypti*. Every bioassay was conducted from 1200 to 1700 h. In a preliminary test, 0.1 mg of each plant extract was solubilized in 20 µl of ethanol by sonication for 10 sec and provided an appropriate amount for repellent bioassays. Ethanol (20 µl) was applied directly to exposed skin through a 5-cm-diameter hole on the back of a rubber glove and dried for 3 min. Because biting density plays an important role in studies of repellency (Rozendaal 1997), skin was exposed for 5 min in a screen wire cage (30 × 30 × 30 cm) containing 300 blood-starved females (6–8 days old). Immediately after the control exposure, the hand was removed from the cage and a dose of 0.1 mg/cm² of each test plant material and deet (Sigma, St. Louis, MO) in 20 µl of ethanol were applied evenly over the skin surface. After air drying for 3 min, the treated hand was exposed to mosquitoes in the same test cage for 5 min at 30-min intervals. The number of test mosquitoes biting on the skin was recorded. Experiments were conducted at 28 ± 2°C and 75 ± 5% RH. Each assay was replicated 3 times.

Repellency was calculated according to the formula from Schreck et al. (1977): % repellency = [(Ta - Tb)/Ta] × 100, where Ta is the number of mosquitoes in the control and Tb is the number of mosquitoes in the treated group.

Statistical analyses: Percent repellency was de-

termined and transformed to arcsine square root values for ANOVA. Treatment means were compared and separated by the Scheffe test at *P* = 0.05 (SAS Institute 1990). Means ± SE of untransformed data are reported.

RESULTS AND DISCUSSION

The repellent activity of methanol extracts from 23 aromatic medicinal plant species and a steam distillate against starved *Ae. aegypti* females varied according to plant species (Table 2). At a dose of 0.1 mg/cm², potent repellency against mosquito adults was obtained with the extracts of *Cinnamomum cassia* Blume bark (91%), *Nardostachys chinensis* Batalin rhizome (81%), *Paeonia suffruticosa* Andrews root bark (80%), and *C. camphora* steam distillate (94%). Repellency in each case was comparable to that of deet (82%). *Eugenia caryophyllata* Thunb. extract provided 75% repellency. The other 19 plant extracts exhibited <70% repellency.

We know that plant-derived insect repellent agents are selective, have no or little harmful effect on nontarget organisms or the environment, and can be applied to human skin and clothing in the same way as conventional repellents (Curtis et al. 1990, Isman 1995, Rozendaal 1997). Furthermore, many plant extracts and essential oils manifest repellent activity against different mosquito species (Curtis et al. 1990, Sukumar et al. 1991, Rozendaal 1997). Sukumar et al. (1991) noted that the most promis-

Table 2. Repellent activities of aromatic medicinal plants against female *Aedes aegypti* skin test.

Plant species ¹	% repellency ²
<i>Acorus calamus</i> var. <i>angustatus</i>	56 ± 9.6 CDEF
<i>Acorus gramineus</i>	44 ± 1.8 DEFG
<i>Agatache rugosa</i>	54 ± 1.6 CDEF
<i>Angelica dahurica</i>	50 ± 1.5 CDEFG
<i>Aquilaria agallocha</i>	62 ± 2.2 BCDEF
<i>Artemisia princeps</i> var. <i>orientalis</i>	60 ± 3.6 BCDEF
<i>Chaenomeles sinensis</i>	49 ± 2.5 CDEFG
<i>Cinnamomum camphora</i>	94 ± 1.8 A
<i>Cinnamomum cassia</i>	91 ± 3.9 AB
<i>Dioscorea batatas</i>	39 ± 1.6 EFG
<i>Eugenia caryophyllata</i>	75 ± 3.3 ABCDE
<i>Evodia rutaecarpa</i>	32 ± 3.2 FG
<i>Gleditsia horrida</i>	57 ± 3.6 CDEF
<i>Glycyrrhiza glabra</i>	28 ± 4.5 FG
<i>Inula helenium</i>	50 ± 3.7 CDEFG
<i>Lysimachia davurica</i>	50 ± 2.7 CDEFG
<i>Magnolia obovata</i>	59 ± 2.3 CDEF
<i>Nardostachys chinensis</i>	81 ± 5.8 ABCD
<i>Paeonia suffruticosa</i>	80 ± 4.3 ABCD
<i>Piper nigrum</i>	52 ± 3.9 CDEFG
<i>Rheum coreanum</i>	51 ± 1.9 CDEFG
<i>Schizonepeta tenuifolia</i>	53 ± 4.0 CDEF
<i>Solanum melongena</i>	15 ± 3.2 G
<i>Stemona japonica</i>	61 ± 1.3 BCDEF
Deet	82 ± 3.7 ABC

¹ Exposed to 0.1 mg/cm² for 15 min.

² Means within a column followed by the same letter are not significantly different ($P = 0.05$, Scheffe test; SAS Institute 1990). Repellency was transformed to arcsine square root values before ANOVA. Means ± SE of untransformed data are reported.

ing botanical mosquito control agents are in the families Asteraceae, Cladophoraceae, Labiatae, Meliaceae, Oocystaceae, and Rutaceae. In this study, repellency against female *Ae. aegypti* comparable to deet was observed for extracts from plants in the Lauraceae, Myrtaceae, Paeoniaceae, and Valerianaceae families. Repellent activity of the extracts of *C. cassia* bark, *E. caryophyllata* flower bud, *N. chinensis* rhizome, and *P. suffruticosa* root bark, as well as the steam distillate of *C. camphora*, against female *Ae. aegypti* was comparable to that of deet.

The repellent activity of extracts from *C. cassia* bark, *N. chinensis* rhizome, *P. suffruticosa* root bark, and *C. camphora* steam distillate against female *Ae. aegypti* at 0.1 mg/cm² was comparable to that of deet (Fig. 1). Their efficacy lasted for 1 h. Relatively short duration of repellency (30 min) was observed in *P. suffruticosa* extract and *C. camphora* steam distillate.

Many plant extracts and essential oils with high volatility, such as alkanes, terpenoids, alcohols, and aldehydes are repellent to mosquitoes for periods ranging from 15 min to 10 h (Rozendaal 1997). Furthermore, various formulations for controlled release have been developed to increase the protection period provided by repellents (Gupta and Rutledge 1989). Sharma and Ansari (1994), for ex-




Fig. 1. Duration of protection of methanol extracts from 3 plant species, a steam distillate of *C. camphora*, and deet against female *Ae. aegypti* when dosed at a rate of 0.1 mg/cm². Bar represents standard error.

ample, reported that a 1% neem oil-kerosene mixture could provide economical personal protection from mosquito bites. *Lantana camara* L. flower extract in coconut oil provided 94.5% protection from *Aedes albopictus* (Skuse) and *Ae. aegypti* without adverse effects on the human volunteers for a 3-month period after the application (Dua et al. 1996).

Results of this study indicate that some plant extracts could be useful for protecting human and domestic animals from mosquito attack, provided a slow-release effect for the repellents can be developed. For practical use of these plants as novel mosquito repellents to proceed, however, further research on their safety in human health, as well as formulations that improve repellent potency and stability, is necessary.

ACKNOWLEDGMENTS

This work was supported by grants from the Korea Science and Engineering Foundation (R01-2000-000088-0) and the Ministry of Education (Brain Korea 21 Project) to YJA.

REFERENCES CITED

- Brown AWA. 1986. Insecticide resistance in mosquitoes: pragmatic review. *J Am Mosq Control Assoc* 2:123-140.
- Croft BA, Brown AWA. 1975. Responses of arthropod natural enemies to insecticides. *Ann Rev Entomol* 20: 285-335.
- Curtis CF, Lives JD, Baolih LU, Renz A. 1990. Natural and synthetic repellents. In: Curtis CF, ed. *Appropriate technology in vector control* Boca Raton, FL: CRC Press. p 75-92.
- Dua VK, Gupta NC, Pandey AC, Sharma VP. 1996. Repellency of *Lantana camara* (Verbenaceae) flowers against *Aedes* mosquitoes. *J Am Mosq Control Assoc* 12:406-408.

- Frances SP, Klein TA, Hildebrandt DW, Burge R, Noigamol C, Eikarat N, Sripongsai B, Wirtz RA. 1996. Laboratory and field evaluation of deet, CIC-4 and AI3-37220 against *Anopheles dirus* (Diptera: Culicidae) in Thailand. *J Med Entomol* 33:511-515.
- Gupta RK, Rutledge LC. 1989. Laboratory evaluation of controlled release repellent formulations on human volunteers under three climatic regimens. *J Am Mosq Control Assoc* 5:52-55.
- Hayes JB Jr, Laws ER Jr. 1991. *Handbook of pesticide toxicology* Volume 1. San Diego, CA: Academic Press.
- Isman MB. 1995. Leads and prospects for the development of new botanical insecticides. *Rev Pestic Toxicol* 3:1-20.
- Namba T. 1993. *The encyclopedia of wakan-yaku (traditional Sino-Japanese medicines) with color pictures* Osaka, Japan: Hoikusha.
- Qiu H, Jun HW, John WM. 1998. Pharmacokinetic, formulation, and safety of insect repellent *N,N*-diethyl-3-methylbenzamide (DEET): a review. *J Am Mosq Control Assoc* 14:12-27.
- Rozendaal JA. 1997. *Vector control* Geneva, Switzerland: World Health Organization. p 7-177.
- SAS Institute. 1990. *SAS/STAT user's guide* Version 6. Cary, NC: SAS Institute, Inc.
- Schreck CE, Posey K, Smith D. 1977. Repellent activity of compounds submitted by Walter Reed Army Institute of Research. 1. Protection time and minimum effective dosage against *Aedes aegypti* mosquitoes. *Tech Bull US Dept Agric* 1549:215.
- Sharma VP, Ansari MA. 1994. Personal protection from mosquitoes (Diptera: Culicidae) by burning neem oil in kerosene. *J Med Entomol* 31:505-507.
- Sukumar K, Perich MJ, Boobar LR. 1991. Botanical derivatives in mosquito control: a review. *J Am Mosq Control Assoc* 7:210-237.
- Wink M. 1993. Production and application of phytochemicals from an agricultural perspective. In: van Beek TA, Breteler H, eds. *Phytochemistry and agriculture* Oxford, United Kingdom: Clarendon Press. p 171-213.