PHOTO ANALYSIS OF SEA FLOOR MICRORELIEF

On the basis of over 700 NEL sea floor photographs, numerical microroughness ratings have been predicted for normal bottom environments, and zones of equal microroughness mapped for large portions of Pacific and adjacent ocean areas.

C. J. Shipek • Research Report
U. S. NAVY ELECTRONICS LABORATORY, SAN DIEGO, CALIFORNIA

26 April 1966

T12
7855
105
no. 1374

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

PROBLEM

Determine and study those characteristics of the sea floor that affect the propagation of acoustic energy in the sea. Specifically, investigate the character, magnitude, and distribution of sea floor microrelief and its superposition on intermediate and major underwater topographic features.

RESULTS

1. More than 700 NEL sea floor photographs, obtained over the past ten years, in a wide variety of environments, have been analyzed. The bottom microrelief has been related to the major agencies responsible for its formation and, on this basis, five predominating types of microrelief have been defined.

2. General ranges in the vertical height and horizontal extent of the microrelief have been determined for the upper exposed surfaces of 39 intermediate underwater features that are, in turn, related to 12 major features or provinces.

3. Microroughness ratings, on a numerical scale of 1 through 5, are predicted for normal bottom environments. These ratings should prove useful in acoustic problems.

4. Isoroughness zones have been mapped for large portions of the Pacific and adjacent ocean areas.

RECOMMENDATIONS

1. Continue combined photographic, sediment, water-sampling, water-current, and temperature studies at ocean stations to improve the reliability of microroughness predictions for the sea floor.

2. Construct microprofiles of the sea floor covering vertical, horizontal, and slope measurements of micro-relief for correlation analyses. The principles of photogrammetry should be applied if stereo cameras are utilized to obtain the needed data.

3. Encourage the study of micro and intermediate relief as an aid to the future operation of submersibles and tracked vehicles on the sea floor.

4. Encourage further use of portable, subbottom, acoustic profiling methods to help establish the relationships of microrelief at the sediment-water interface to deeper substrata features.

ADMINISTRATIVE INFORMATION

Field work for this project has extended over the past 10 years on an intermittent basis. This report was prepared under BUSHIPS Problem SR 104 03 01, Task 0539 (NEL Problem L4-1 (now L40161)). It was approved for publication on 26 April 1966.

Many scientists and crew members on numerous oceanographic cruises helped the author lower cameras, collect samples, and process film; they are too numerous to be named here. Special thanks are due to R. E. Boyce for the preparation of photographic grid scales and help in the analysis of bottom photographs; to D. Muerdter for mathematical assistance; and G. L. Prible, D. Bachman, and Mrs. Renee Forsyth for the preparation of maps and illustrations.

CONTENTS

INTRODUCTION...	page 7
HISTORICAL BACKGROUND...	8
DEFINITION OF MICRORELIEF...	10
SUMMARY OF MICRORELIEF PHOTOGRAPHIC DATA...	13
THE CAMERAS AND TECHNIQUES USED...	20
INTERPRETATION OF BOTTOM PHOTOGRAPHS...	25
Value and Limitations of Bottom Photography...	25
Methods Used in the Interpretation of the Photographs...	26
DISCUSSION OF THE ORIGIN AND OCCURRENCE OF MICRORELIEF...	28
General...	28
Sediment and its Formation...	32
Rock Outcrops...	33
Microrelief Resulting from Physical Processes...	36
Microrelief Resulting from Chemical Precipitation...	40
Microrelief Caused by the Bodies of Benthonic Organisms...	44
Microrelief Caused by the Activity of Benthonic Organisms...	47
Summary of the Origin and Occurrence of Microrelief...	50
A SEA FLOOR MICROROUGHNESS SCALE AND WORLDWIDE ZONES OF ISOROUGHNESS...	53
Microrelief Numerical Ratings...	53
Summary of Microrelief Characteristics...	55
Microrelief Interrelationships and Distribution...	56
Zones of Isoroughness...	63
CONCLUSIONS...	65
REFERENCES...	67-70

TABLES

- 1 Underwater topographic relationships and photo analysis of sea floor microrelief... *page 16-17*
- 2 Distribution and general relationships of underwater topographic relief... *19*
- 3 Numerical scale of sea floor microroughness and descriptive notes... *53-55*

ILLUSTRATIONS

- 1 Sea floor topographic relief... *page 12*
- 2 Map showing NEL sea floor camera stations and underwater topographic features... *15*
- 3 Photo and laboratory analysis card... *21*
- 4 Camera, Type I, low oblique photographic coverage of sea floor... *22*
- 5 Camera, Type II, high oblique photographic coverage of the sea floor... *23*
- 6 Camera, Type III, vertical stereophotographic coverage of the sea floor... *24*
- 7 Microrelief on sea floor... *29*
- 8 Microrelief on sea floor - east equatorial Pacific... *30*
- 9 Manganese-encrusted bedrock, boulders, or nodules on a topographic high, Rift Mountains, Indian Ocean... *35*
- 10 Exposed bedrock enclosed by coarse, rippled calcareous sand on abyssal plain south of Java Trench, Indian Ocean... *35*
- 11 Manganese-encrusted boulders in coarse sand and gravel, located on ridge above elongated basin, Gulf of California... *35*
- 12 Exposed boulders imbedded in thin silty-sediment cover on upper flanks of Coronado Bank... *35*
- 13 Cross rippling in sorted calcareous sand on a topographic high, Rift Mountains, Indian Ocean... *37*

ILLUSTRATIONS (Continued)

- 14 Cross rippling in sorted calcareous sand on a topographic high, Rift Mountains, Indian Ocean... 37
- 15 Sediment slumping in calcareous clay on slope of inter-mountain valley, Rift Mountains, Indian Ocean... 37
- 16 Manganese-encrusted symmetrical ripple marks in calcareous sand on southwest slope of Eniwetok Atoll... 39
- 17 Free, spherical manganese nodules, 3.48 cm diameter, resting on *Globigerina* ooze. South central Pacific Ocean, northwest of Tahiti... 42
- 18 Partially buried manganese nodules in clay sediment, Pacific Ocean south of New Zealand... 42
- 19 Manganese-impregnated pumice slabs resting on brown clay sediment; some clay capping is visible. East North Pacific... 42
- 20 Scattered manganese nodules resting on red clay sediment, East Indian Ocean. Rough surface of clay indicates churning by benthonic organisms... 42
- 21 A high concentration of evenly spaced spherical manganese nodules, 3.48 cm diameter, resting on a *Globigerina* ooze. Animal churning has caused the partial covering of some nodules. South central Pacific Ocean, northwest of Tahiti... 43
- 22 Coral growing on encrusted volcanic boulders, inshore slope on east side of Mauritius Island, Indian Ocean... 45
- 23 Holothurian (sea cucumber) and small ophiuroids (brittle stars) on surface of brown silty clay in San Diego Trough... 45
- 24 Numerous worms protruding from churned silty clay on east slope of Guaymas Basin, Gulf of California... 45
- 25 Ophiuroids (brittle stars), holothurians (spiny sea cucumbers), on churned surface of brown silty clay in San Diego Trough... 45
- 26 Sea pens growing on well churned clay in north Guaymas Basin, Gulf of California... 46
- 27 Large sea urchin on rippled gray sand, marine slope of Saint Paul Island, Indian Ocean... 46
- 28 Large ophiuroid (brittle star) and a number of small, spiny holothurians (sea cucumbers) on silty clay in protected Flores Sea basin north of Soembawa Island, Indonesia... 46

ILLUSTRATIONS (Continued)

- 29 Sutured animal mounds, in churned red clay, intermountain valley of Rift Mountains, Indian Ocean...48
- 30 Worm and excrement on mottled clay near Cocos Keeling Island, Indian Ocean... 49
- 31 Starfish, brittle star, and sea pen in churned silty clay, west slope of Guaymas Basin, Gulf of California... 49
- 32 World maps showing NEL sea floor camera stations and underwater topographic features:
 - A. Zones of isoroughness...57
 - B. Roughness due to ripples and bedrock...58
 - C. Roughness due to manganese nodules ...59
 - D. Roughness due to benthonic organisms ...60
 - E. Roughness due to biological churning of sediment ...61

INTRODUCTION

The purpose of this report is to describe photographic samplings of very small bottom-relief features found at different depths in the ocean and in different environments; and to show how photographic and related information from the surfaces of intermediate topographic features can be interpreted to predict the general nature of the microrelief and project microrelief characteristics from area to area. Approximately 700 bottom photographs of the surfaces of selected underwater features in the Pacific, Indian, and Arctic Oceans, and in the Gulf of California, have made this study and analysis possible.

HISTORICAL BACKGROUND

Early mariners probed the sea bottom in shallow nearshore waters and brought up samples of sand, clay, and rock. Their major interest was in making safe passages across the oceans and in and out of harbors and channels. The shortness of their sounding lines prevented them from reaching the deeper sea floor surfaces. Consequently little was learned about the detailed microcharacter of the deep sea floor. On the basis of meager knowledge and scattered samples, many thought of the sea floor as an extremely flat, monotonous plain composed of a soft, clayey upper surface that blended gradually upward into seawater. In recent times, sonic depth finders, deep-sea cameras, modern sampling equipment, acoustic probes, and manned diving submersibles have revealed the structure and character of what is referred to as microrelief at the sea floor.

Major underwater topographic relief features have now been located and mapped. Distinct provinces encompassing similar features have been recognized and delineated.^{1,2} Intermediate bottom features are now being studied to make our knowledge of the sea floor more comprehensive and to provide a basis for the examination of microroughness and internal structures.³ It has been shown, for example, that abyssal hills cover 80 to 85 percent of the Pacific floor alone while, in other areas, bottom environments vary from smooth plains to steep eroded escarpments, mountain ranges, and fracture zones of irregular roughness.⁴

For many years the very small vertical relief on the sea floor was not thought to be significant to mankind. Recent advances in methods of propagating and receiving underwater sound signals, however, have shown that the roughness formed by natural chemical, physical, and biological forces at the sea floor has sufficient height and extent to affect the reflection and scattering of certain sound frequencies therefrom. Thus, Urick⁵ made back-scattering studies of the bottom in 1954 with a tiltable transducer at several locations in a harbor on the East Coast.

¹Superscript numbers denote references in the list at the end of this report.

His results suggested that backscattering of sound from a natural bottom is due more to its roughness than to the sedimentary particles that make it up, and that at normal incidence the bottom is sufficiently rough so that little sound is returned by specular reflection. The increase in scattering for sand-rock bottom seemed to be due to the greater roughness. Again, in 1960, Urick⁶ utilized two transducers to study side scattering over a uniform bottom. The results indicated that isotropic bottom scattering may be safely assumed in applications utilizing separated transmitting and receiving transducers at kilocycle frequencies. As examples of sea scatterers of sound, Urick listed marine life, the air bubbles existing near the sea surface in bad weather, and the roughnesses of the sea floor. The latter are known to be the dominant sound scatterers in shallow water when the bottom is hard and when downward refraction exists.

Mackenzie⁷ made bottom reverberation measurements with 530 and 1030 c/s omnidirectional sound sources in 2100-fathom water near San Diego. An analysis of nonspecular reflections to obtain a scattering constant for the bottom revealed that, for clays, muds, or fine-grained sands, there appears to be no significant frequency dependence over a range of seven octaves. Mackenzie⁸ stated that, for frequencies from 200 to 3000 c/s in long-range shallow-water propagation, bottom reverberation is probably the only important type of reverberation. He pointed out that accurate close-range reverberation studies over different bottoms and especially at lower frequencies are needed for grazing angles from 1° to 10° , and he noted that reverberation strength is more significant than reverberation level.

In discussing the theory of sound reflection from flat and uniform fluid bottoms, Mackenzie⁹ stated that more measurements should be made of the bottom reflection properties, especially for grazing angles of incidence from zero to 30° where the results are more critically dependent on the bottom properties.

Lieberman¹⁰ measured the reflection coefficient of the ocean bottom for supersonic sound (24 kc/s) at grazing

incidence, and found that it depended not only on the median size of the particles making up the bottom, but also on the sorting of the particles and on the bottom topography.

Working at 17 stations near Woods Hole, Massachusetts, he obtained the following *in situ* reflection coefficients for five gradations of materials at the sediment-water interface:

mud	0 to 0.20
mud-sand	0.20 to 0.40
sand-mud	0.40 to 0.60
sand	0.40 to 0.85
stony	0.50 to 0.85

It can be seen that sand and stony bottoms have the best reflecting ability. Lieberman stated that an exact treatment of the reflection from an irregular surface may be given only if the geometry of the surface is completely specified.

DEFINITION OF MICRORELIEF

Earth forces and water movements have created great trenches, rises, mountain ranges, seamounts, scarps, and other significant relief features in deep-sea environments. The sediments that are continually forming at the sea floor are also not without relief, even though this relief is of very small magnitude compared to that of the larger features.

The term "underwater microrelief" usually refers to those very small features not ordinarily detected by modern echo-sounders. Laughton¹¹ used the term "microtopography" and defined it as referring to those features of the sea floor that can be visually observed in the scale range of 50 meters to 1 mm. (He regarded 50 meters as approximately the limit of visibility in the clearest of oceanic water.) Carsola¹² defined microrelief on the arctic sea floor as consisting of the smallest features that can be resolved at depths between 0 and 200 fathoms (366 meters) with the NMC and NMC-2

echo-sounders. He added that the pits and mounds of the microrelief areas are in places 60 ft (18+ meters) from bottom of depression to top of rise, but usually only 6 to 30 ft (1.8 to 9.1 meters).

The author of this report has defined microrelief¹³ as the very small surficial topographic features that are superimposed on both major and intermediate relief and are distinguishable by examination of bottom photographs (fig. 1). Vertical heights are measured in centimeters and meters, and lengths in meters and occasionally tens of meters. Major underwater relief, at the other extreme, encompasses the larger earth features such as deep trenches, high plateaus, rises, and mountain ranges. Intermediate relief consists of the smaller underwater topographic features such as low hills, valleys, banks, small basins, gullies, and island slopes that may be parts of major features, but are still too large to be photographed in their entirety with present-day underwater cameras. Microrelief, as defined by the author, covers 100 percent of the sea floor, whereas intermediate and major relief features are localized and discontinuous.

Figure 1. Sea floor topographic relief.

SUMMARY OF MICRORELIEF PHOTOGRAPHIC DATA

Various parts of the surfaces of 39 intermediate topographic features of the Pacific Ocean, Indian Ocean, Arctic Ocean, and attached water bodies were sampled photographically. These intermediate features were parts of 12 major topographic features or provinces. (The photographic techniques and cameras used are described in the next section.) Figure 2 is a world map showing the locations of the NEL deep-sea camera stations occupied for this purpose, and their relation to the major underwater topographic features.

These camera stations were not all preselected to obtain the greatest number of similar or dissimilar sea floor characteristics. Many photographs were taken from oceanographic vessels moving along tracks designated for other scientific programs. Available vessel time along these routes, however, made it possible to obtain photo samplings of a wide variety of sea floor features.

Table 1 is an overall summary of the photographic data. It relates the 12 major features, the 39 intermediate features, and the associated microrelief as found by random sampling in the areas surveyed. It also gives the heights, lengths, and widths of the visible targets at the sediment-water interface, and the dimensions of free chemical and biological specimens. Although the latter are not permanent parts of the relief, they are included here because of their significance in acoustics. Many crest lengths of physically formed ripples were not determinable from the small-area bottom photographs.

Table 2, like table 1, relates the major and intermediate topographic features and the associated microrelief, but classifies the microrelief into five types based on the five principal agencies responsible for its formation.

TABLE 1 UNDERWATER TOPOGRAPHIC RELATIONSHIPS AND PHOTO ANALYSIS OF SEA FLOOR MICRORELIEF

MAJOR TOPOGRAPHIC FEATURE	INTERMEDIATE TOPOGRAPHIC FEATURE	BOTTOM SEGMENT	PHOTO DEPTHS (meters)	MICRORELIEF (FROM PHOTO INTERPRETATION)			USDEGRA OF PHOTO (sp. cm)	NUMBER OF BOTTOM PHOTOS
				TYPE	SIZE (cm) Width x Length	HEIGHT (cm)		
CONTINENTAL TERRACE	BARNA	Sandy silt	495	Out-rops	27 x 28	32	35	10
		Pebbles, shales	1154	BrITTLE stars, sea urchins, sponges	8 x 8	3	20	10
	BASIN	Silty clay	600	Churning	11 x 25	7	—	10
		Excavated	1800	Worms, brittle stars, starfish, sea pens, sea cucumbers, corals	4 x 7	1	1	52
	CANYON OUTER	Silty fine sand	620	Chemical reduction	—	1	20	10
		Debris	1100	Churning	8 x 15	4	—	51
	ISLAND SHELF	Silty fine sand	24	BrITTLE stars, sea pens, crustaceans, worms, gastropods, fish	4 x 9	1	7	27
		Coarse material	100	Churning	12 x 23	9	—	27
	CONTINENTAL SLOPE	Silty clay	158	Rock fragments, animal shells, worms, sea urchins, polychaetes	33 x 23	4	1	5
			618	Churning	3 x 8	16	19	26
	SILL	Silty clay	1174	BrITTLE stars, gastropods, sea pens, starfish	8 x 12	3	4	26
				Churning	4 x 10	2	—	2
	SADDLE	Silty clay	542	BrITTLE stars	4 x 4	3	13	34
			902	Churning	4 x 8	2	—	13
RIDGE	TROUGH	Silty clay	140	Outcrops or rock fragments	10 x 16	6	32	13
		Debris	1080	BrITTLE stars, sea cucumbers, worms, sea urchins, corals, crustaceans	8 x 10	3	21	27
	VALLEY	Silty clay	490	Churning	5 x 8	3	—	26
		Debris	522	Sea urchins, sea cucumbers, worms, sea pens, fish	5 x 7	1	1	16
	RIFT VALLEY	Silty clay and sand	7025	Churning-glow trails	5 x 12	1	—	37
		Eastward	7080	BrITTLE stars, sea urchins	3 x 13	6	—	11
	BASIN	Clay	4080	Churning-gps	7 x 20	4	—	21
		Debris		Manganese nodules	2 x 2	3	5	2
	GENTLE RELIEF 275 Meter Slope	Calcareous sand	3180	Sea cucumbers, worms	5 x 1	2	1	1
		Debris	3520	Churning	1 x 3	5	—	1
	ISLAND SHELF	Silt	495	Sea urchin, worm, sponge	4 x 4	10	1	1
		Debris		Churning-glow trails	5 x 21	4	—	2
	SEAMOUNT SURFACE	Calc. ooze	1320	Out-rops or boulders	12 x 15	8	63	9
				Sea pens, worms, sponges, brittle stars	2 x 2	5	24	15
SULF OF CALIFORNIA	BASIN	Silty clay and pebbles	810	Churning-gps	20 x 20	3	—	12
		Debris	1480	Ripple marks	4 x 25	4	—	7
	CANYON INNER	Coarse sand and silt	732	Manganese nodules	1 x 1	1	10	2
		Debris	1866	Ripple marks, sponges	16 x 40	5	80	2
	ISLAND SHELF	Sand and silt	72	Outcrops, boulder	20 x 20	35	1	15
		Debris	1700	Corals, starfish, sponges, worm, fish	4 x 6	7	12	44
	SMALL LOCAL RIDGE	Coarse gravel	72	Churning	4 x 16	5	—	19
		Manganese grain		Out-rops	22 x 34	48	25	8
		Debris		BrITTLE stars, fish, worms, sponges	13 x 13	5	10	18
				Churning	—	14	—	13
				Ripple marks	5 x 20	2	60	8
				Out-rops, boulders	4 x 1	1	9	9
				BrITTLE stars, sea pens, animal shells	4 x 9	16	5	20
				Sea cucumbers, corals, worms	5 x 9	2	—	1

TABLE 1 (CONTINUED)

MAJOR TOPOGRAPHIC FEATURE	IMMEDIATE TOPOGRAPHIC FEATURE	BOTTOM SEDIMENT	PHOTO DEPTHS (meters)	MICRORELIEF (FROM PHOTO INTERPRETATION)				
				TYPE	SIZE, m Width x Length	HEIGHT (cm)	COVERAGE OF PHOTO (percent)	NUMBER OF BOTTOM PHOTOS
FRACTURE ZONE EAST PACIFIC	BASIN	Fine clay Coarse particles	5180	Manganese nodules Sea cucumbers, at extremities Churning	3 x 4 2 x 5 1 x 3	3 2 —	50 1 —	7 1 —
		IRREGULAR HILLS 180 Meter Relief	Fine clay Coarse particles	4900	Manganese nodules Sea urchins, sea cucumbers (at extremities) Churning	2 x 4 12 x 12 2 x 2	2 15 2	5 1 —
	GENTLE TOPOGRAPHY	Clay Debris	5400	Manganese nodules Sponges, worms Churning	2 x 2 2 x 2 2 x 10	2 2 —	3 1 —	3 2 —
	IRREGULAR TOPOGRAPHY	Clay	4400	Manganese nodules Churning	3 x 3 1 x 1	2.5 2	100 —	2 —
	ABYSSAL PLAIN	GENTLE TOPOGRAPHY	Calcareous ooze Clay	3200 to 5100	Outcrop, boulders, manganese oxide nodules Coral, sea cucumbers, sponges, worms Churning Ripple marks	41 x 57 2 x 2 5 x 12 120 x 13	31 2 4 2	20 — — 70
RISE EAST PACIFIC	ISLAND SLOPE	Calcareous ooze Shells	274 to 437	Oolitic carbonate rock and coral Tube worm Churning	10 x 100 6 x 5 10 x 12	60 5 2	26 — —	3 — —
		IRREGULAR TOPOGRAPHY	Calcareous ooze Clay	3477	Manganese nodules Boulders Sea pens, sea cucumbers, worms Churning	16 x 21 3 x 5 5 x 13	11 1 —	11 1 —
RISE CENTRAL INDIAN	ISLAND SLOPE	Volcanic sand Cobbles and pebbles Coral debris	448	Outer speculum, boulders Sea urchins, sea cucumbers, coral, fish Churning Ripple marks	8 x 8 5 x 7 2 x 2 2 x 10	5 4 2 —	24 10 — —	15 17 — —
		LOW HILLS	Clay	1754 and 4940	Manganese nodules Bottle stars, coral Churning	7 x 8 0 x 9 2 x 5	6 2 —	10 5 —
DEEP SEA BASIN	GENTLE TOPOGRAPHY	Clay	2044 and 5130	Manganese nodules Sea urchins Churning-trails	1 x 3 15 x 15 2 x 2	— 5 —	— — —	— 40 45
		GENTLE TOPOGRAPHY	Fine mineral sand	6580	Sponges, sea anemones, coral Churning	10 x 10 5 x 13	10 —	— —
TRENCH	LOCAL RIDGE	Fine mineral sand Gravel Coarse material	1500	Crusting chemical Outcrop, boulders Churning-trails, mudcracks Ripple marks	80 x 120 40 x 70 8 x 200 2 x 100	1 1 2 2	70 10 — 70	— 11 — 1
		FLAT TOPOGRAPHY	Silt clay	5000	Worms Churning	2 x 2 1 x 4	2 2	10 —
	TOPOGRAPHIC HIGH	Gibberine ooze	2040	Manganese nodules and crusting Outcrop Churning Ripple marks	8 x 8 2 x 1 150 x 90	6 2 12	100 — 100	2 1 —
RIFT MOUNTAINS	INTERMOUNTAIN VALLEY	Clay	1760	Worms, sea anemones, gastropods Sediment elongation Churning-trails, pits	2 x 2 8 x 20 6 x 18	1 12 4	5 40 —	17 1 18
		ABYSSAL HILLS	Silt ooze	1440 to 4722	Manganese nodules Sponges, worms, sea urchins Churning	4 x 2 1 x 2 5 x 7	1 2 —	60 10 —
SEAMOUNT PROVINCE	SEAMOUNT SURFACE	Calcareous sand Debris	1490 to 2035	Manganese nodules and fragments Crusting chemical Ripple marks	1 x 5 — 100 x 18	2 — 10	18 10 80	10 10 33
		IRREGULAR TOPOGRAPHY	Clay	6520	Pumice slabs manganese rating	15 x 17	5	42
BEAUFORT SEA	SLOPE	Clay Debris Pebbles, scattered	549 to 2423	Brilliant stars Churning-trails, trails	13 x 15 2 x 15	2 2	— —	— 16
		SEA VALLEY	Clay Debris Pebbles, scattered	604 to 1831	Fan worms, brittle stars, crustaceans Churning	17 x 18 2 x 7	1 —	5 —
	SMOOTH TOPOGRAPHY		Silt clay Debris	1528 and 2423	Sea pens Churning-trails	1 x 4 2 x 30	8 1	2 —

TABLE 2. DISTRIBUTION AND GENERAL RELATIONSHIPS OF
UNDERWATER TOPOGRAPHIC RELIEF

MAJOR RELIEF	INTERMEDIATE RELIEF	MICRORELIEF				
		OUT- CROPS	CHEMICAL DISTURB.	BENTHONIC ANIMALS	BIOLOG. DISTURB.	PHYSICAL DISTURB.
CONTINENTAL TERRACE	BANK					
	BASIN					
	CANYON					
	ISLAND SHELF					
	CONTINENTAL SLOPE					
	SILL					
	SADDLE					
	TROUGH					
	VALLEY					
	RIFT VALLEY					
RIDGE	BASIN					
	GENTLE RELIEF					
	ISLAND SHELF					
	SEAMOUNT SURFACE					
GULF OF CALIF.	BASIN					
	CANYON					
	ISLAND SHELF					
	SMALL LOCAL RIDGE					
FRACTURE ZONE	BASIN					
	IRREGULAR HILLS					
	GENTLE TOPOG.					
	IRREG. TOPOG.					
ABYS. PLAIN	GENTLE TOPOG.					
EAST PAC. RISE	ISLAND SLOPE					
	LOW HILLS					
CNT. IND. RISE DEEP SEA BASIN	ISLAND SLOPE					
	LOW HILLS					
	GENTLE TOPOG.					
TRENCH	GENTLE TOPOG.					
	LOCAL RIDGE					
	FLAT TOPOG.					
RIFT MTS.	TOPOG. HIGH					
	INTERMOUNTAIN VALLEY					
SEAMOUNT PROVINCE	ABYSSAL HILLS					
	SEAMOUNT SURFACE					
	IRREG. TOPOG.					
BEAUFORT SEA	SLOPE					
	SEA VALLEY					
	SMOOTH TOPOGRAPHY					

THE CAMERAS AND TECHNIQUES USED

The type of form used for recording the results of the examination of each photograph is shown in figure 3A. Figure 3B shows the back of this form on which was listed the properties of any water, sediment, or animal sample taken in the area.

Three types of deep-sea camera systems were employed for the collection of the bottom photographs. Type I (fig. 4) has a tilted semiwide-angle lens of 30 mm focal length, and uses 35 mm film and a repeating electronic flash light mounted in an upright frame. A bottom contact switch activates the light and causes the film to advance each time a lowered weight touches the sea floor. All photos taken with this apparatus were low obliques (30° from the vertical) at a preset target distance of 6 feet underwater. Figure 4 also shows the grid scale used for making measurements on the photos taken with this camera system.

Type II camera (fig. 5) has a sled-type frame supporting a 35 mm camera and a flash unit which can rotate about a common axis to give vertical or oblique photos. Normally the camera angle was set close to the horizontal (15° from the horizontal) to permit greater bottom coverage even though the photo scale was more distorted. This camera was usually lowered to a preselected target distance and then allowed to drift over the sea floor using sonar to control the height above the bottom. Photographs were taken automatically at preset time intervals. Excessive heave and roll of the lowering vessel caused unwanted deviations from acceptable depths of focus. Figure 5 shows the optics and grid overlay for photo measurements.

The Type III camera is a 35 mm Edgerton, Germeshausen, and Grier stereo underwater system. Photo pairs were taken with the camera axes oriented vertically and set at a prefixed separation. Figure 6 shows the optics and field coverage for this system.

The grids for the three camera systems are constructed of clear plastic. They were applied as overlays in examining the bottom photos, and established scale for

A. FRONT OF CARD

PHOTO ANALYSIS

Station _____ Cruise _____ Vessel _____ Photographer _____ Date _____
 Photo No. _____ Total _____ Depth-M Uncorr. _____ to _____ Lat. _____
 Vault No. _____ Camera Type _____ Depth-M Corr. _____ to _____ Long. _____
 Area _____ Major Topog. _____ Intermediate Topog. _____
 Sediment Type _____ Color _____ Sample _____ Slope < _____

PHYSICAL DISTURBANCES: Ripple Marks _____ Pattern _____ Shape _____ Ht _____ L _____ Coverage _____
 Scour _____ Slumping _____ Slipping _____ Banking _____ Capping _____ Sorting _____
 Current Velocity _____ Direction _____ Transparency _____ Water Sample _____
 Bending of fauna and other _____ Orientation of fauna _____

ORGANIC PRODUCTIVITY: Benthic _____ Pelagic _____ Ht _____ L and W _____ Coverage _____
 Range H _____ L _____ W _____

ORGANIC DISTURBANCES: Debris _____ Tracks _____ Trails _____ Tubes _____ Pits _____ Mounds _____
 Excrement _____ Churning _____ Scratches _____ Old _____ New _____ Other _____
 Average H _____ L _____ W _____ Range H _____ L _____ W _____ Coverage _____

CHEMICAL DISTURBANCES: Nodules _____ Crusts _____ Slabs _____ Fragments _____ Other _____
 Composition _____ Hardness _____ Color _____ Coverage _____

ROCKS: Bedrock _____ Boulders _____ Cobbles _____ Pebbles _____ Fragments _____
 Average H _____ L _____ W _____ Range H _____ L _____ W _____
 Strike _____ Dip _____ Color _____ Hardness _____ Sample _____ Coverage _____

Remarks:

Single _____ Stereo _____ % overlap _____ Camera Elev. _____ Scale: 1 cm = _____ Contour Int. _____

B. BACK OF CARD

LABORATORY ANALYSIS

I. Sample Analysis

A. Sediment

1. Type _____
2. Med. Diam Gr. (mm) _____
3. % sand _____
 % silt _____
 % clay _____
4. Density, sat. gr/cc _____
5. Porosity _____ %

6. Bulk Density Mineral, gr/cc _____
7. Shear Strength, gr/cm² _____
8. Sound Speed, m/sec. _____
9. Attenuation, dB/m _____
10. Other Properties or Constituents: _____

11. Rate of Accumulation _____

a. Method _____

B. Bottom Water Analysis

1. Temp., deg. C. _____
2. Salinity, ppt. _____ Sigma-T _____
 a. Bottle No. _____ Sp. Vol. An. _____
3. Pressure, kg/cm² _____ Density _____
4. Sound Speed _____
 a. Ratio: $\frac{\text{sound speed sed.}}{\text{sound speed water}}$ _____

C. Summary of Microrelief and causes: _____

Figure 3. Photo and laboratory analysis card.

Figure 4. Camera, Type I, low oblique photographic coverage of sea floor.

Figure 5. Camera, Type II, high oblique photographic coverage of the sea floor.

Figure 6. Camera, Type III, vertical stereophotographic coverage of the sea floor.

all parts of each photo. With the proper grid, measurements of ripple marks, benthonic organisms, nodules, rocks, and other relief features were made on each photo when such were distinguishable. Composition of sediment samples and the water clarity were correlated with photo interpretation and measurements whenever possible. With all three camera systems, target distance seldom exceeded 12 feet (3.7 meters).

INTERPRETATION OF BOTTOM PHOTOGRAPHS

Value and Limitations of Bottom Photography

While the human eye has the ability to see and interpret, it is unreliable and it cannot retain accurate images for future use. In contrast to the eye, the camera faithfully records everything in its field of view. The image is substantially permanent, and variations can occur only when the photographic prints are being measured and interpreted. As a consequence, the development and application of the underwater camera have shed new light on the nature of the sediment-water interface.

Fortunately, many techniques used for measurement and interpretation in engineering photography can be, and are being, used in the study of underwater microrelief. Photography as applied to underwater surficial relief does not differ basically from subaerial photography, but the submarine environment is much more difficult to work in. Because of refraction, the angular coverage of the usual lens is restricted in water, and corrections must be made to get normal, undistorted views. Seawater also scatters and absorbs light much more than does dry, clear air. Except in the clearest of deep oceanic waters, target distances for underwater photography are limited to less than 12 feet (3.7 meters) for maximum resolution and detail. Photo coverage is thereby confined to relatively small areas. Electronic flash at 5000° Kelvin temperature gives the greatest light penetration under water.

In the study of the sea bottom, considerable application is being made of stereophotography, for which vertical camera axes are best. Microrelief viewed from directly above is in true plan view, but side detail is obscured. Conversely, when viewed from the side, the relief is clear but the plan view is highly distorted. For this reason oblique photography is used mainly for rapid exploratory work when side views of low relief features are desired and measurements are of secondary importance. When measurements of vertical height are the primary objective, modern methods of photogrammetry are being widely used.¹⁴

Methods Used in the Interpretation of the Photographs

The vastness of the oceans prohibits complete photographic and sediment sampling of the bottom surface with present-day oceanographic equipment. We can, however, build up composite pictures of local and regional bottom environments from collections of isolated camera samplings and associated information, such as sediment grain size, bottom currents, composition, nearness to land, topographic barriers, and availability of food supplies to benthonic organisms.

A complication in the study and interpretation of microrelief is that its structure is far from permanent. As Crozier has pointed out,¹⁵ very considerable sediment turnover goes on over relatively short periods of time. In the shallow waters around Bermuda, in an area of 1.7 square miles, he found that 500 to 1000 tons of sand passed through holothurians (sea cucumbers) annually. This represents a layer of sediment about 1 centimeter thick every 100 years, and sediments have been forming for millions of years.

To overcome the difficulties in deriving patterns of origin and change on the sea floor, investigators use many and varied techniques of photo interpretation. Some of the more important are discussed in this section.

Photographic contrast has been found to be an effective means of detecting and describing visible targets, and of relating them to one another. Sharpness of definition is required in aerial photographs, but is even more important

underwater where suspended matter tends to obliterate the sharp edges of all targets more than a few meters from the camera. Sharpness of definition depends on such factors as lens quality, water transparency, exposure and target distance, type of film, quality of look-through port, and film processing. Since contrast is often poor in bottom photographs, it is necessary to apply indirect methods to decipher unknown and obscure markings. Shadow effects are used extensively for this purpose.

A knowledge of the agencies that create the micro-relief can help in photo interpretation. For example, scratches, tracks, grooves, trails, impressions, pits, mounds, and holes can normally be related to some form of benthonic organism if enough photos of an area are examined. Obtaining actual specimens makes such identification easier and more reliable. Organisms in shallow environments sometimes can be viewed directly while creating their peculiar marks or disturbances. Patterns of organism distribution can be worked out from photos where scale, target distance, and angular field coverage are known or can be calculated.

Rock outcrops can usually be detected and delineated easily on photographs by means of contrast, resulting sharp boundaries, and relief with respect to the surrounding sediment.

Where ripple marks show up, it is safe to assume the presence of sandy materials, and of current or wave motion of sufficient strength to form the ripples. A knowledge of grain size and density, along with the velocity of water flow present, can be used to supplement the photographic data, furnishing descriptions of wave characteristics, estimations of length, and determinations of symmetry and steepness of ripple marks.

The firmness of the sediment-water interface can be determined indirectly from photographs. The generation of turbid clouds of fine-grained material by artificial agitation of the sea floor usually denotes a soft clay bottom. (On the other hand, such turbid clouds occasionally form over a hard bottom, the fine suspended material coming from elsewhere.) The presence of silt in clay hardens the bottom, and the effect is intensified when sand is present. Although these bottom materials are subject to movement

by water, the sediment-water interface is probably firm and sharply defined most of the time. The presence of coarse sand, gravel, pebbles, and boulders, of course, denotes a very hard bottom, capable of sustaining heavy underwater loads. Such bottom materials are usually easily determinable by photographic sampling. The freshness of the upper surfaces of manganese nodules, capping of sediment, banking of sediment, etc., give clues to the rate of sediment accumulation. Changes in bottom materials, outcrops, and chemical deposits can be studied photographically by tracing sediment boundary lines.

Repetition of patterns of sea floor features assists in extending the results of photo interpretation to other areas. Photographs can be compared on a geographic basis, by targets, by animal populations, or by micro-relief types.

DISCUSSION OF THE ORIGIN AND OCCURRENCE OF MICRORELIEF

General

The sea floor can be described as a huge jigsaw puzzle. Much remains to be done before all the pieces of information concerning the individual phenomena that contribute to the formation of the microrelief can be fitted into a single, overall pattern. Nevertheless, some general statements can be made with respect to the origin, distribution, and relationships of the microstructures.

Thus, examination of surfaces and smaller related parts of larger underwater features has shown that the same kind of environment may extend for great distances. For example, Mero¹⁵ states that sea floor sediments tend to be rather uniform over large regions and that a few chemical analyses of nodules can be averaged to give fairly reliable results. Averages of heights for visible targets distinguished in the photographic study described in this report are shown in figure 7. A specific area of the

Figure 7. Microrelief on sea floor.

Pacific Ocean, the East Equatorial Province, was examined separately, giving the results shown in figure 8. Comparison of the two figures reveals that the very small relief is rather uniform over large areas.

Figure 8. Microrelief on sea floor — east equatorial Pacific.

The NEL photographs have revealed similar environmental relationships in microrelief separated by hundreds of miles. Comparable microstructures also exist at different depths, presumably due to a depth-range toleration by many benthonic organisms.

Random sampling of irregularly distributed bottom targets makes it difficult to obtain accurate distribution counts. The photos revealed a range of distribution of small visible targets for all environments from less than 1 per square meter of sea floor to 5 or more. Tracks, markings, scratches, plow trails, impressions, pits, and mounds resulting from benthonic sediment churning did not vary greatly in magnitude and horizontal distribution considering the vast size of the oceans.

In shoal depths, organisms are normally plentiful because of rich food supplies. With greater depth there appears to be a linear decrease of the number of visible organisms, with a corresponding decrease of food. As an example of the numbers of organisms, Beagles and Shumway¹⁷ reported high concentrations of clams ($400/\text{m}^2$) in depths of 5 meters or less in Alaska waters. Worm mounds up to 5 cm high in the same area indicated concentrations up to $10 \text{ animals}/\text{m}^2$. In other nearshore shallow areas in Alaska, Beagles and Shumway reported concentrations of worm mounds at $20/\text{m}^2$. On the other hand, many photographs taken in deep abyssal areas indicate benthonic organism populations of less than $1/\text{m}^2$. Number of benthonic fauna and amount of churning of sea floor sediments are closely related, as is to be expected.

There does not appear to be a close relationship between bedrock outcrops and manganese nodule formation, but further investigation will probably show considerable manganese encrustation on the harder surfaces such as bedrock and coral boulders that exist on topographic highs. Table 2 indicates a distinct relationship between physical disturbances such as wave or current-formed ripple marks and outcrops of rock. Photographs have repeatedly shown that coarser sediments such as *Globigerina* ooze and coral sands are rippled in the vicinity of bedrock.

Sediment and its Formation

The materials that constitute the sea floor are derived from various sources. Organic and inorganic matter falls from the water column or moves in laterally to find places of accumulation. Most inorganic matter comes from the erosion of subaerial rocks; from atmospheric fallout including dust, ash, and meteoritic spherules; and from volcanic activity. Ice rafting drops variable amounts of inorganic and organic material in the high latitudes, but in tropical and midlatitudes the quantity is not significant. Most organic matter comes from the tests and debris of organisms living in the upper layers of the oceans. Deep-water pelagic fish supply a varying amount of organic material to bottom sediments.

Turbidity flows help to move sediments rapidly into deep water toward their final resting places.^{1a}

Normally, in deeper water, the organic content of sediments is highly variable. Sediments, if rich in organic matter, support large colonies of surface and subsurface feeders leading, in turn, to churning and the development of microrelief. Where little organic material is present, churning is insignificant, microrelief is minimal, and the bottom approaches flatness. Nearshore sediments are made up largely of subaerial runoff, are usually extremely rich in organic matter, and form maximum microrelief.

Sediments are usually characterized according to their composition. If 30 percent or more is calcium carbonate, the sediment is designated a calcareous ooze. If less than 30 percent carbonate is present, the sediment is called a clay. When the sediment consists mostly of silica, it is designated a siliceous ooze.

In spite of wide differences in environment and origin, bottom sediments may still be acoustically similar as far as their microrough upper surfaces are concerned. Differences in color have no influence on sound reflection. Organic content of sediments may have some effect, but this has not been determined.

Rock Outcrops

Soundings, dredge samples, rock-damaged core barrels, and photographs have proved the existence of hard rock and other hard surfaces on the sea floor even in deep water. Acoustic probing has verified the occurrence of isolated pinnacles, lava flows, flat-lying buried strata, and partially covered boulders and loose rock at, or not far below, the sediment-water interface. The presence of exposed rock is related to major earth movements, erosion, volcanic activity, changing rates of sediment accumulation, bottom current, and wave action, and it also depends on the major underwater topographic features and superimposed intermediate relief. Usually, rock appears on ledges, ridges, topographic highs, and upper surfaces of seamounts; in trenches, canyons, fault scarps, and rift mountains; and on nearshore shelves, banks, and terraces. But it is probable that outcroppings of nearly all magnitudes occur throughout the entire range of underwater environments.

Rock relief above existing sediment surfaces varies in height from a few centimeters to thousands of meters. Although still within the definition of microrelief, many outcroppings examined photographically were more than 50 cm high. Partially buried boulders can appear as outcrops in bottom photographs, but probably are not continuous for any distance. Distribution of rock outcrop is highly irregular. Laughton¹¹ found that surface features of seamounts varied considerably over short distances. On a topographic high in the Indian Ocean, the surface at one location changes from cross-rippled sands to a manganese-encrusted bed of nodules or bedrock in a distance of not over 70 meters (fig. 9). Many small isolated local outcrops of a few meters in extent exist in a variety of surrounding areas of different origins. They are in sharp contrast to great underwater mountain ranges many kilometers in length. Depth of water is not a limiting factor.

Unless turbidity flows are active in an area, currents, wave and eddy action, and gravity pull tend to remove the finer-grained materials from isolated highs by a constant washing action.¹⁹ It is not unusual for these agencies to keep areas clean of fine sediment cover for intermittent

lengths of time. In some instances remnant outcrops on abyssal plains are being kept free from fine sediments (fig. 10).

Gradually, the hollows, basins, lows, depressions, valleys, and other such areas along continental margins and in ocean deeps will fill up with sediment. The end result will be a smooth level plain, with the outcrops covered up.^{20,21} Marine-cut rock terraces have formed in the past along the Southern California coastline and then have been buried with subaerial deposits.²¹ Fortunately, both exposed and buried bedrock can be investigated with the subbottom acoustic profiler.

Seldom is it possible to detect freshly broken boulders or rocks on the sea floor. Exposure underwater eventually leads to crusting, and the process appears to go on endlessly.¹

Manganese-encrusted bedrock, boulders, or nodules above the sediment surface indicate ineffective sediment accumulation for the area (fig. 11), and suggest that volcanism has been locally active.²² In shallow areas of the lighted zone, organisms and plants attach themselves to hard surfaces forming colonies in a very short time (fig. 12). Busby²³ has discovered underwater outcrops of limestone in the Bahamas exhibiting microrelief features (cavities) that were formed subaerially and then submerged without appreciable change. Except in rare instances, rock outcrops tend to retain their original edges and faces. Unlike the rapid changes caused by subaerial erosion agents, changes to underwater rock occur very slowly unless the rock is exposed to earthquake and volcanic activity.

Figure 9. Manganese-encrusted bedrock, boulders, or nodules on a topographic high, Rift Mountains, Indian Ocean. Depth: 2940 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 10. Exposed bedrock enclosed by coarse, rippled calcareous sand on abyssal plain south of Java Trench, Indian Ocean. Depth: 3290 m. Photo area: 1.2 m x 3.7 m. Type II camera. 2.6 m target distance.

Figure 11. Manganese-encrusted boulders in coarse sand and gravel, located on ridge above elongated basin, Gulf of California. Depth: 720 m. Photo area: 1.2 m x 1.8 m. Type I camera. 2.4 m target distance.

Figure 12. Exposed boulders imbedded in thin silty-sediment cover on upper flanks of Coronado Bank. Depth: 495 m. Photo area: 1.2 m x 1.8 m. Type I camera. 2.4 m target distance.

Microrelief Resulting from Physical Processes

Major processes, other than biological, responsible for the formation and disruption of microstructures on and within the sea floor are gravitational downslope movements, waterwaves and currents, and turbidity flows. The magnitude of these forces is attested to by an ever-growing library of bottom photos showing wave- and current-derived ripple marks in sandy sediments (fig. 13), grain sorting (fig. 14), gravel beds or lag deposits (fig. 11), scour around bottom objects (fig. 10), slumping on unstable slopes (fig. 15), and current-distorted benthonic animals. Even weather factors including surface winds have an effect. The author²⁴ has described the formation of cross ripples by the action of currents on submerged seamounts when underwater obstructions are present (fig. 14). Menard²⁵ regarded internal waves as a possible source of oscillations in deep water. Inman²⁶ pointed out that symmetrical ripple marks require oscillatory currents for formation. Busby²⁷ considered that, at a depth of 1929 meters in the Tongue of the Ocean, Bahamas, ripples were probably caused by tidal oscillations rather than by internal waves. Laughton¹¹ stated that, in the light of our present knowledge of deep currents, it may be possible to explain the formation of deep ripple marks in terms of steady currents. Further, variations in the symmetry of the ripples may be ascribed to local current and tidal fluctuations, rather than to short-period oscillatory motion of the water. Deep-seated tidal-wave motion plays an important part in the movement and formation of sediments.¹⁹

An increasing amount of information as to its effects on bottom sedimentation indicates the importance of turbidity flow. The irregular and heterogeneous sediment surfaces often attributed to turbidity flow are difficult to detect in bottom photos, and physical and biological processes tend to modify near-surface microstructures in a short period of time anyway.

Biological activity, especially burrowing on unstable slopes, could initiate and accelerate turbidity flows and resultant slumping and disruption of microstructures. Marks, pits, mounds, grooves, scratches, and impressions are short-lived and ever-changing. The bottom surfaces

Figure 13. Cross rippling in sorted calcareous sand on a topographic high, Rift Mountains, Indian Ocean. Depth: 2940 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 14. Cross rippling in sorted calcareous sand on a topographic high, Rift Mountains, Indian Ocean. Depth: 2940 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance. This photograph was taken a short distance from the one shown as figure 10.

Figure 15. Sediment slumping in calcareous clay on slope of intermountain valley, Rift Mountains, Indian Ocean. Note sutured mounds and radial feeding pattern. Depth: 1760 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

involved tend to be soft and cohesive. Many abyssal deep-sea clay areas exhibit such softness, while other compacted and stable clay bottoms act as strong sound-reflecting surfaces. In some cases, subsurface reflecting surfaces are so hard that soft sediments at the water interface are masked out and become unrecognizable on sonic recorders. It would be difficult to measure the disruption that goes on during sediment accumulation before final consolidation takes place. The sediment surfaces exhibit some form and degree of microrelief throughout the period during which these processes occur.

The height of the microrelief caused by physical forces can be measured. Its extent can only be estimated from area to area, being beyond normal photographic limits. Laughton¹¹ has reported a wide range extending from ripples with wavelengths of a few centimeters to huge sand waves with wavelengths of a kilometer or more. Within photographic limits many ripple marks of symmetrical and asymmetrical shapes have been observed and recorded (fig. 16). The author²⁴ has reported wavelengths from 15 cm to 61 cm in calcareous sands on a submerged seamount surface at Eniwetok; ripple heights ranged from <2.5 cm to 15 cm. Vertical heights, averaged for the ripple marks noted in table 1 of this report, are between 2 and 6 cm (fig. 16).

Average values, of course, are of little use when analyzing the ripples on the sea floor because of the close relationship between type of microrelief and the intermediate or major feature on which it is superimposed. This does not mean, however, that every underwater topographic feature has its own kind of microrelief, since similar microreliefs exist in many different environments. Sand size, density, and chemical makeup, together with current or wave velocities, determine the ripple height and length. Variations in, or deflections to, the flow of water change or modify the shape and steepness of the ripple marks. As the velocity is increased, ripple mark formation continues until finally a high enough velocity is reached to destroy all the existing ripple marks. At such high velocities, grains move in sheet flow. Sand ripples normally form over a velocity range of 12 cm/sec in fine sand to 100 cm/sec in coarse granules.

Figure 16. Manganese-encrusted symmetrical ripple marks in calcareous sand on southwest slope of Eniwetok Atoll. Depth: 1650 m. Photo area: 1.2 m x 1.8 m. Type I camera. 2.4 m target distance.

Deep-sea ripple marks show as great a variety of forms as those in shallow water. Variations in wavelength, crest length, amplitude, and symmetry occur over very short distances.¹¹

Lack of ripple marks does not necessarily mean a lack of water motion; it could mean merely an absence of grains capable of being rippled by the moving water. Fine-grained silts and clays in abyssal deeps will not ripple regardless of the strength of the current. But scour marks can result, providing evidence of currents in such areas. Dzulynski and Sanders²⁸ hold that such marks are the result of scour of a cohesive mud bottom (clay) by turbulent eddies of sediment-laden water, and also that current marks can be made in any depositional environment where sand is transported over a cohesive mud bottom. These authors also consider that "tool" marks result from tractional movement of larger particles. Where the latter are lacking

and no sand grains are present, visible animal markings and impressions indicate a lack of current activity.

Currents are normally very small in deeper clay areas, but occasional banking of fine-grained sediments against nodules and rock outcrops indicates the presence of low-order water motion. Sometimes the bending of bottom-attached organisms and deformed bottom crawlers indicate weak sea floor currents.

Current meters attached to camera frames are now being used to measure water velocities at the sea floor in various environments.

Microrelief Resulting from Chemical Precipitation

Of all the microstructures that have been discovered on the sea floor, the manganese nodule, slab, or encrustation is perhaps the most interesting. It may have influence on sound reflectivity when concentrated, and it will probably have economic value to man in the future.

The manganese nodule, like phosphorite and ferric nodules, is a product of chemical precipitation that takes place at the sea floor (figs. 17 and 18). The manganese nodule usually contains small percentages of iron, cobalt, copper, and nickel oxides. Goldberg and Arrhenius²⁹ believe that organisms concentrate dissolved chemicals in the sea water column, after which chemical changes and attachment to various nuclei on the bottom occur. These authors have also suggested that there is a direct correlation between the iron/manganese ratio in nodules and the rate of sedimentation. Graham³⁰ believes that organic reactions in seawater form manganese nodules. Although the mode of formation of the nodules is not yet clear to the oceanographer, the deposits are substantial and form recognizable relief.

The heights and lengths of the nodules range from less than 1 mm to 50 cm or greater, but the nodules for a particular area usually fall within a narrow size range. The concentration (number of nodules per square meter of bottom) is difficult to determine. Sonar and TV scanning

promise to help solve this problem in the future. Sampling and extensive photography have disclosed the presence of different grades of manganese nodules over great areas, especially in the Pacific Ocean.¹ The distribution of the nodules is somewhat patchy which is to be expected in view of depth variations and the obstructing effect of major and intermediate underwater topographic features. However, regional patterns of surface nodules and encrustations do occur.^{1,31}

Menard has concluded from photographs and dredge hauls that almost all rock outcrops in the open ocean are covered with oxides.¹ Slabs of volcanic ash or consolidated sediment encrusted with ferromanganese oxides (fig. 19) are very common in the eastern Pacific.³²

Manganese nodules occur in a wide variety of underwater environments (table 1). In protected basins where water circulation is poor, nodules do not usually form. In fact it is not uncommon to find oxygen-starved sediments which form a reducing environment. Evidence of chemical reduction in sediments has been found in San Pedro Basin on the southern California Continental Terrace. The surface was mottled and smooth, and animal activity was lacking. Microrelief is very low in such areas because of the lack of organisms and of sediment-churning. Nodules are normally found on ridges, highs, hills, large mounds, in open deep-sea basins, and on abyssal plains (fig. 20) where oxygen is in plentiful supply, unless a very high sill cuts off water circulation. In some abyssal areas, nodules occur on highs and in lows between hills as a relatively uniform cover, but it is likely that most nodules are on hills.¹ Sediment accumulation would normally be greater in the lows than on the highs.

Cores from Pacific areas indicate that fewer nodules exist below the interface than at the interface.¹ To explain the presence of nodules at the surface, the sediment must accumulate at a slower rate than manganese oxide, although some workers now feel that organisms can keep the nodules at the surface in their search for food.¹ The author doubts the effectiveness of this mechanism, except in areas where nodules are small and closely spaced (fig. 21), since many nodules would be too large for any benthonic organisms to move except downward by undercutting.

Figure 17. Free, spherical manganese nodules, 3.48 cm diameter, resting on *Globigerina* ooze. South central Pacific Ocean, northwest of Tahiti. Depth: 3695 m. Photo area: 0.5 m x 0.3 m. Type II camera. < 1 m target distance.

Figure 18. Partially buried manganese nodules in clay sediment, Pacific Ocean south of New Zealand. Depth: 5160 m. Photo area: 1.2 m x 1.2 m. Type II camera. 2.6 m target distance.

Figure 19. Manganese-impregnated pumice slabs resting on brown clay sediment; some clay capping is visible. East North Pacific. Depth: 4302 m. Photo area: 1.2 m x 1.8 m. Type I camera. 2.4 m target distance.

Figure 20. Scattered manganese nodules resting on red clay sediment, East Indian Ocean. Rough surface of clay indicates churning by benthonic organisms. Depth: 4940 m. Photo area: 1.2 m x 2.4 m. Type II camera. 3 m target distance.

Figure 21. A high concentration of evenly spaced spherical manganese nodules, 3.48 cm diameter, resting on a Globigerina ooze. Animal churning has caused the partial covering of some nodules. South central Pacific Ocean, northwest of Tahiti. The freshness of the nodule surfaces indicates a low rate of sedimentation in this area. Depth: 3695 m. Photo area: 1.2 m x 1.2 m. Type II camera. 2.6 m target distance.

A new approach to the problem of manganese nodule concentration at or near the sediment surface has been made by Olausson and Uusitlalo³³ who experimented with artificial sands. They contend that seismic vibrations may cause a lifting effect to keep large particles at the surface for longer times. This, they say, would give rise to high concentrations of nodules at the sediment-water interface and in regions with a comparatively high rate of sedimentation. However, this process would probably not be effective in cohesive clays. In some areas manganese nodules appear to be almost buried as a result of a high rate of sedimentation (fig. 18).

Type of manganese nodule varies from closely packed spheroids (fig. 21) to widely scattered irregular chunks (fig. 20) and encrusted, irregularly shaped, volcanic rock fragments. In places, manganese oxides have cemented sand surfaces, and closely packed nodules have coalesced to form hard rock surfaces (fig. 16).

Microrelief Caused by the Bodies of Benthonic Organisms

That organisms exist on the sea floor has, of course, been known for many years.³³⁻³⁵ Strange looking creatures have been brought to the surface by dredge, trawl, sounding leads, nets, anchors, scoops, and other means in increasing numbers (figs. 22 through 27). Although many of these animals are alike in structure, they differ from species that live in shallow environments.

Because of the usual haphazard manner of collection and lack of detailed distribution information, little is known about these bottom feeders and burrowers. Sampling has been insufficient to establish the concentration of most of them. Exact identification from photographs, also, is difficult unless actual specimens are available for confirmation. However, increased use of bottom photography and coordinated sampling has somewhat alleviated these difficulties, and the concentration of a few identifiable animals has been obtained.³⁴

Few underwater photos of the sea floor exceed 5 square meters in coverage. Most present-day samplers collect from areas of about 0.4 to 1 square meter of the sea floor. And single grabs seldom bring up more than one organism or part of an organism, although bottom trawls and dredges screen more sediment and yield more specimens.³⁵ Thus, few of the deep-sea animals are ever visible for counting and examination. And counting is complicated by the fact that some organisms combine bottom and subbottom feeding habits.

In most deep-sea environments the food supply is sufficient to support only a few animals, from less than one per square meter in fine-grained red clays to approximately ten in richer nearshore silts and clays. In abyssal plains and interseamount areas where intermediate relief is slight, the numbers are even smaller. Those animals living at the interface and visible in photographs vary in size from less than a millimeter to 15 centimeters in length (fig. 28).

To constitute effective microrelief on the sea floor, many organisms would have to be concentrated there. In large enough numbers, as in shallow water, crustaceans,

Figure 22. Coral growing on encrusted volcanic boulders, inshore slope on east side of Mauritius Island, Indian Ocean. Depth: 479 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 23. Holothurian (sea cucumber) and small ophiuroids (brittle stars) on surface of brown silty clay in San Diego Trough. Depth: 905 m. Photo area: 1.2 x 1.8 m. Type I camera. 2.4 m target distance.

Figure 24. Numerous worms protruding from churned silty clay on east slope of Guaymas Basin, Gulf of California. Depth: 610 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 25. Ophiuroids (brittle stars), holothurians (spiny sea cucumbers), on churned surface of brown silty clay in San Diego Trough. Depth: 905 m. Photo area: 1.2 x 1.8 m. Type I camera. 2.4 m target distance.

Figure 26. Sea pens growing on well churned clay in north Guaymas Basin, Gulf of California. Depth: 894 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 27. Large sea urchin on rippled gray sand, marine slope of Saint Paul Island, Indian Ocean. Depth: 448 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance.

Figure 28. Large ophiuroid (brittle star) and a number of small, spiny holothurians (sea cucumbers) on silty clay in protected Flores Sea basin north of Soembawa Island, Indonesia. Depth: 2094 m. Photo area: 0.7 m x 0.9 m. Type III camera. 1.5 m target distance.

starfish, brittle stars, sea cucumbers, sponges, crinoids, worms, and urchins might have sufficient density and height to constitute microstructures capable of reflecting sound or changing coefficients of sound reflection.

It may be concluded that, except in shallow areas, the bodies of living organisms themselves hardly affect the reflection of sound.

Microrelief Caused by the Activity of Benthonic Organisms

During feeding, benthonic organisms churn up the softer sediments of the sea floor.^{11,13} Sokolova, working in the Kurile-Kamchatka Trench of the Pacific Ocean, found that most (about 55 percent) of deep-sea bottom invertebrates are deposit feeders, a smaller number (about 25 percent) are suspension feeders, and still fewer (about 20 percent) are carnivorous.³⁵ Further, the composition and quantitative abundance of deep-sea communities are largely determined by the amount of organic material on the sea floor and in suspension in the water layers just above the bottom. Sokolova adds that a zonation of communities arises, each zone extending over a region characterized by a distinct type of sedimentation. The dominant organisms in each area are those best adapted to it by their feeding habits.

Organisms living in the deep-sea environment must spend their entire lives seeking out and digesting food. Even where food is plentiful, for example, in nearshore areas fed by rich subaerial runoff, the organisms must spend most of their time seeking food.

Laughton¹¹ found more sediment disturbance and more animals in shallow water (fig. 26), and less disturbance and fewer animals in deep water (fig. 28). This relationship results from diminishing food supplies with increasing depth. Numerous bottom photographs on deep abyssal plains bear this out (fig. 29); here the main source of food is the surface water. If the latter is not rich in nutrients, as it seldom seems to be over abyssal areas of the midlatitudes, very little organic debris falls to the sea floor. Also, because of the great depth, a large percentage of the falling

Figure 29. Sutured animal mounds, in churned red clay, intermountain valley of Rift Mountains, Indian Ocean. Depth: 1760 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance. The mounds are 5 cm high and probably created or altered by crustaceans.

debris is apt to be consumed by pelagic fishes before reaching the sea floor.

Benthonic organisms of the deep barren areas are therefore forced to crawl, burrow, plow, hold fast, and dig in the sediment for the minute quantities of food available (fig. 30). More churning, of course, takes place in the organically rich, shallow coastal sediments (fig. 31), because of the greater numbers and varieties of organisms present. In the process of seeking food, the organisms build undersurface galleries, tubes, and tunnels.

Regardless of the total amount of churning, the vertical height of the resulting microrelief seems to reach an equilibrium at 6 centimeters above the sediment-water interface. Benthonic organisms such as brittle stars, holothurians, starfish, urchins, and even crustaceans seem unable to build up larger microstructures and, indeed, have no need to do so since their only objective is to obtain food.

Figure 30. Worm and excrement on mottled clay near Cocos Keeling Island, Indian Ocean. Depth: 5100 m. Photo area: 0.6 m x 4 m. Type II camera. 2.4 m target distance. Mounds are 3 cm high.

Figure 31. Starfish, brittle star, and sea pen on churned silty clay, west slope of Guaymas Basin, Gulf of California. Depth: 911 m. Photo area: 0.6 m x 3 m. Type II camera. 2.4 m target distance. Pits are approximately 15 cm deep and 30 cm wide across the openings.

Burrowing organisms, whether they feed at the surface or just below it, digest enormous quantities of sediment. They produce ribbons of excrement (fig. 30) and tunnels that wind around in crisscrossing patterns, evidence of effective extraction of organic matter. The activities of these organisms must have an overall effect on the strength, shear properties, and reflectivity of sediments. Tunnels, voids, changes in porosity, and disruption of grain alignment individually may not be of great importance, but collectively they contribute significantly to the general pattern of the microrelief.

Summary of the Origin and Occurrence of Microrelief

The summarized information in this section is arranged to correspond with the five microrelief types of table 2.

I. ROCK OUTCROPS AND BURIED BOULDERS

These usually occur on shoaler and irregular under-water topographic features such as banks, island shelves, sides of troughs, seamounts, mountains, cliffs, ridges, canyons, marine terraces, and shelf breaks. Medium-to-coarse sands are usually present near the outcrops and show ripple marks, scour, banking, and other signs of water activity. Fine-grained materials other than fine-grained sands, such as silts and clays, have usually been washed into deeper water leaving the rock outcrops and gravel deposits exposed. A variety of benthonic organisms, together with a few fish, live in or near the shallow rocky environments, whereas finer sediments, manganese coatings and crusts, and infrequent benthonic organisms occur in the deeper rocky environments. Rock outcrops usually consist of volcanics (including basalt flows and pumice slabs), coral boulders, and conglomeratic aggregates, and they are normally coated with manganese and iron oxides.

II. CHEMICAL DISTURBANCES

Manganese nodules are the most prominent free chemical precipitate on the sea floor. They occur over a wide range of depths, but are most noticeable on topographical highs, on red clays below 5000 meters, and on shoaler calcareous oozes. They are present in high concentrations on extended abyssal areas and in deep-sea basins of all major oceans, constitute strong microrelief, and represent a mineral resource of great potential economic value. Phosphorite slabs are found in high concentrations on coastal banks where they are localized. Both manganese

and phosphorite deposits may have a strong acoustic reflecting effect since they occur at or near the sediment-water interface. How these concentrations arose is not not clearly understood at the present time.

III. BENTHONIC ORGANISMS

Organisms living on or near the sea floor surface are nearly always associated with soft, organically rich clays, and nearshore silts and clayey silts. If burrowing organisms such as worms and sea cucumbers are present, the bottom materials are apt to be churned or disrupted by feeding, exploratory, or escape activities. Surface feeders such as sea urchins, crustaceans, and brittle stars account for a large portion of the tracks, trails, scratches, and marks found on bottom sediments. Brittle stars and sea cucumbers engage in both burrowing and surface feeding activities. Other organisms such as corals, sponges, sea pens, and some worms hold fast to the bottom and filter-feed directly from the water. Crustaceans and brittle stars normally move about over many bottom environments seeking detrital particles and modifying microstructures such as mounds and pits formed by burrowing organisms.

Fewer species of animals, but often of larger size, are present in deeper environments where food is relatively scarce. Animals are normally abundant in shoaler near-shore areas where surface plankton blooms and subaerial runoff from adjacent land areas provide large quantities of organic matter. Benthonic organisms are not present in high enough concentrations in deeper waters to greatly affect directed sonar signals.

IV. CHURNING OF SEA FLOOR SEDIMENTS

The churning is related directly to the abundance of burrowing organisms and the quantity of food matter present; and it is most noticeable in clays and clayey silts, and in shallow water. Water depth is related to the degree of churning, amount of resultant microrelief, and the predominant type of microstructure present. Churning takes

place in all sediments inhabited by benthonic organisms. Sometimes it is localized, but normally it exists over great areas and often with very little variation from environment to environment.

V. PHYSICAL DISTURBANCES OF BOTTOM SEDIMENTS

Waves and currents, together with earthquake jarring, faulting, gravitational slumping, turbidity flows, and tsunamis, all affect the sediment-water interface. Study of the types, numbers, and distribution patterns of the marks left on the interface helps determine the origin, strength, and mode of operation of the physical forces involved. Normally, the latter are active in canyons, on seamounts, on topographic highs, on marine terraces, in channels, on beaches, and in coarse sediments around rocky outcrops.

Symmetrical ripple profiles denote oscillatory wave motion, whereas asymmetrical profiles are indicative of currents coming strongly from one general direction. Visible ripple marks are evidence of the water velocities mentioned in a previous section, above and below which sediment grains move only in sheet flow without rippling. Clays do not ripple at any velocity because of the coherency of the fine grains, but gravels can be rippled by strong currents under certain conditions. Many ripple marks have been preserved in place by the cementing of sand grains after rippling. Water motion, sand grain size, and chemical makeup, rather than depth, are the limiting factors in the formation of ripple marks. The water velocities needed to initiate rippling are directly proportional to the density of the sand grains. Coarse sediments are most often detected where rocky outcrops or boulders are exposed, that is, usually on shoaler topographical features.

A SEA FLOOR MICROROUGHNESS SCALE AND WORLDWIDE ZONES OF ISOROUGHNESS

Microrelief Numerical Ratings

Individual descriptions of sea floor microrelief, and predictions of its environmental distribution, have little practical value unless the results can be expressed quantitatively. In a numerical form, however, roughness of the sea floor can be correlated with other factors influencing the transmission of underwater sound, such as sediment properties, sound velocity, temperature, salinity, and larger topographic features.

An arbitrary scale of sea floor roughness has been developed from the measurements and photo-interpretations of table 1, and is presented here as table 3. In addition to numerical ratings, table 3 includes an explanation or definition of the meaning of each rating.

This numerical scale of sea floor microroughness is a beginning attempt, with a minimum of data, to provide a means of predicting microrelief for acoustic purposes. It is hoped that roughness ratings for all sea floor areas can eventually be obtained for use in both military and commercial applications. The success of this endeavor will depend on the accumulation of more and better data, and particularly on accurate ship positioning, camera control, and depth measurement.

A summary of microrelief characteristics follows which, it is hoped, will make the ratings more useful.

TABLE 3. NUMERICAL SCALE OF SEA FLOOR MICROROUGHNESS AND DESCRIPTIVE NOTES

	Numerical Rating of the Relief
Canyon - inner	5
Bank	4
Irregular topography	4
Small local ridge	4,3
Island slope	4,2
Local ridge	3,4

TABLE 3. (Continued)

	Numerical Rating of the Relief
Side slopes of trough	3
Low hills	3
Topographic high	3
Island shelf	3,2
Seamount surface	3,2
Irregular topography	3,2
Island slope	2,4
Seamount surface	2,3
Island shelf	2,3
Irregular topography	2,3
Basin	2
Canyon - outer	2
Continental slope	2
Sill	2
Volley	2
Rift valley	2
Gentle relief (275 meter hills)	2
Irregular hills (180 meter relief)	2
Intermountain valley	2
Abyssal hills	2
Gentle topography	2,1
Gentle topography	1,2
Saddle	1
Slope	1
Sea valley	1
Smooth topography	1
Flat topography	1

EXPLANATION OF NUMERICAL RATINGS

Because the sea floor is never perfectly flat, the rating zero is not used. The rating 1 denotes minimum but recognizable roughness; the rating 5 denotes maximum roughness. Some descriptive notes on the ratings follow:

1. Almost smooth surfaces formed on clays, oozes, and silty clays in abyssal areas between, and on, major and intermediate topographic features. Visible evidence of churning is lacking, with a minimum of epifauna and infauna present. Rock fragments and manganese nodules occur in scattered patches, varied according to chemical composition of water.

Height of churning 3 cm.

Normal range of microrelief 0 to 10 cm.

2. Low-order bottom relief formed by fauna churning on clays, oozes, and silty clays in areas of gentle relief. Such low-order relief also occurs on marine slopes, valleys, basins, and other gentle topographic features. More epifauna visibly present but not in great numbers in deeper areas. Occasional occurrences of small manganese nodules and rock fragments in tightly packed or scattered patterns of distribution, again dependent on seawater conditions. Occasional occurrences of loosely scattered and larger manganese nodules with visible churning between targets. Oozes are normally in shoaler areas and are coarser grained.

Height of churning 3 to 6 cm.

Normal range of microrelief 3 to 20 cm.

3. Maximum churning of clay and silty sediments. Ripple marks occur where fine sands and sandy silts are present. Greater occurrence of larger manganese nodules, pumice slabs, and rock fragments. Chemical crusting of sediments sometimes present. Major and intermediate features predominantly island slopes, hills, ridges, highs, and irregular topography.

Height of churning 6 to 15 cm.

Normal range of microrelief 6 to 30 cm.

4. Rock fragments, outcrops, boulders, and coarse sediments predominate in this shoaler environment. Fauna churning maximum where silty sediments exist. Ripple marks often present in sandy sediments. Greater abundance of epifauna on rocky surfaces.

Height of churning, where present, 15 cm.

Normal range of microrelief 6 cm to 1 meter.

5. Jagged rocks, phosphorite nodules, large boulders, and coarse sediments on upper surfaces and slopes of underwater features such as ridges, mountains, banks, cliffs, walls of canyons, and other topographic highs. Fauna churning of coarse sediments variable and dependent on presence of organic matter. Predominance of attached and unattached epifauna on rock exposures.

Height of churning highly variable up to 15 cm.

Normal range of microrelief 3 cm to 2 or 3 meters.

Summary of Microrelief Characteristics

1. In general, microroughness is directly related to the intermediate and major topographic relief on which it is superimposed.

2. With some exceptions, microrelief diminishes in magnitude with increasing depth. Outcrops are fewer in deeper water where less infauna and epifauna are present

to disrupt normal processes of erosion and sedimentation. Surfaces of targets are cleaned or smoothed by the fine-grained materials falling through the water column. Iso-microroughness values continue largely unchanged for hundreds of square miles.

3. Whatever its origin, the magnitude of the roughness eventually reaches an equilibrium value.

4. There are many causes of microrelief, but they are of less importance acoustically than an ability to delineate the shape or pattern of the relief for a particular environment at a specific time.

5. Microroughness in deeper water is largely caused by infauna churning in clays and silts where organic matter is scarce.

6. Microroughness in shoaler areas, in contrast, is influenced mainly by the presence of rock outcrops and the effects of epifauna attached to hard surfaces. Ripple marks are present because of the greater occurrence of sandy sediments. Churning is highly variable in nearshore sediments because of the great influence of inflowing subaerial sediments, light, and plant life. The acoustic effects of nearshore sediment roughness are predictable. Sediment changes occur faster than in deeper water.

Microrelief Interrelationships and Distribution

Figures 32A through 32E have been prepared as an aid to the study of the origin and distribution of microrelief, and the interrelationships of different types of microrelief.

Figure 32A shows the distribution of sea floor roughness according to the numerical scale of table 3, and general zones of isoroughness. Figures 32B through 32E similarly show the distribution of ripple marks and bedrock, manganese nodules, benthonic organisms, and biological churning, respectively. A close relationship naturally exists between figures 32D and 32E. Exposure

Figure 32. World maps showing NEL sea floor camera stations and underwater topographic features (continued through page 61)

Figure 32 (Continued)

Figure 32 (Continued)

Figure 32 (Continued)

Figure 32 (Continued)

of bedrock (fig. 32B) normally means the presence of coarse sediments and bottom currents, with ripple marks often resulting. The great areal extent of manganese nodules (fig. 32C) in both the Indian and Pacific Oceans requires that this type of microrelief be studied for its possible influence on underwater sound transmission. (A chart showing the distribution of manganese nodules at the sediment-water interface is now in preparation. Source information includes photographs, samples, dredge hauls, and foreign and domestic scientific reports.)

When studied together these figures should permit microrelief interrelationships to be established. The value of such a series of figures will grow with the accumulation of microrelief data. It is believed that the correlation of type of microrelief with depth on specially constructed bathymetric charts will permit a more accurate numerical zoning of roughness and point up the effect of roughness on the bottom reflection of sound.

Zones of Isoroughness

The general zones of isoroughness delineated in figure 32A should be accepted with caution until much more photographic sampling of the oceans has been performed. At present there are many gaps in the data.

The Gulf of Alaska and the area southward to just north of the Mendocino Escarpment is largely an abyssal plain with many isolated seamounts. No camera stations have been occupied in this great area but, on the basis of studies in other abyssal plain areas, a roughness rating of 2 has been assigned to it. The East Pacific Fracture Zone and Baja Seamount Province constitute a fairly large area of generally irregular topography. Roughness rating here will vary from 2 to 3, and reach 4 in some coastal areas where there is a high percentage of bedrock outcroppings. The sediments in the southwestern Pacific Basin are largely of pelagic origin. Red clays predominate but grade into coarser calcareous oozes on the flanks. Microroughness rating varies between 2 and 3 because of the presence of manganese nodules in the deeper areas. Most of the semi-closed deep-sea basins in the Pacific Ocean are largely

supplied by pelagic materials from the water column, but they do not all have free manganese nodules present on their deeper floors. The Philippine Basin floor is marked by a series of low abyssal hills with red clays predominating. The South Australian Basin has a smooth red clay bottom with numerous scattered nodules. Microroughness rating will vary from 1 to 2 in most of these basins. The large abyssal area lying southwest of the Emperor Seamount Chain and north of the Marcus Necker Ridge is largely made up of abyssal aprons and has a roughness rating of 3 due to the presence of manganese nodules.

North of the Bering Straits and on into the Beaufort Basin, the bottom relief is of very low order. Smooth surfaces and minimum churning keep the roughness rating to 1. In the Gulf of California, the rating is high in the canyon and ridge areas but drops to 2 in the deeper basin areas. The Indian Ocean is roughly divided into a western shallow part of calcareous ooze and a deeper eastern part predominantly red clay. A median ridge trending northwest-southeast appears to be the dividing line. However, roughness rating varies from 2 to 3 over most of the Indian Ocean because of the presence of manganese nodules and irregular bottom topography. The roughness rating in the Indian Antarctic Ridge area drops to 2. Nodules are not present, and churning is not pronounced; numerous tracks and trails indicate a paucity of organic matter. Very few identifiable animals are present at the interface.

CONCLUSIONS

Microrelief is related to, and directly influenced by, volcanism, earthquakes, mass water movements, depth of water, chemistry of seawater, sediment accumulation, turbidity flows, atmospheric fallout, subaerial runoff, biological activities, geographical location, and intermediate and major underwater relief features. Microrelief is not confined to visible microstructures at the interface but also occurs as subsurface internal structures of varied origin. The importance of microrelief is only now being recognized because of its effect on underwater sound transmission.

Microrelief will have a bearing on the future construction of underwater launching pads, roads, industrial plants, towers, transducer platforms, dwellings, and farming areas, as well as on methods of waste disposal and on the underwater movements of submersibles and tracked vehicles.

Patterns of microrelief over thousands of square miles of sea floor are being recognized. From many photographic samplings and target measurements, a numerical scale of sea floor microroughness has been established that should prove useful for acoustic purposes.

Zones of equal sea floor roughness and general similarities have been delineated in parts of the Pacific, Indian, and Arctic Oceans. Because of the existence of multitudes of localized smaller features, it is impossible to show microrelief in its entirety on the small-scale charts normally employed. The average values that are shown should, however, permit predictions to be made to assist in the solving of acoustic problems.*

* Stereophotographic studies of sea floor roughness now in progress will provide greater resolution of microscopes.

REFERENCES

1. Menard, H. W., Marine Geology of the Pacific, McGraw-Hill, 1964
2. Hamilton, E. L., "Marine Geology of the Pacific Basin," p.135-154 in Pacific Science Association Standing Committee on Geology and Geophysics, Geology and Solid Earth Geophysics of the Pacific Basin, University of Hawaii Press, 1963
3. Hamilton, E. L., "Sea Floor Relief," American Geophysical Union. Transactions, v.44, p.493-494, June 1963
4. Shippek, C. J., "Photographic Study of Some Deep-Sea Floor Environments in the Eastern Pacific," Geological Society of America. Bulletin, v.71, p.1067-1074, July 1960
5. Urick, R. J., "The Backscattering of Sound From a Harbor Bottom," Acoustical Society of America. Journal, v.26, p.231-235, March 1954
6. Urick, R. J., "Side Scattering of Sound in Shallow Water," Acoustical Society of America. Journal, v.32, p.351-355, March 1960
7. Mackenzie, K. V., "Bottom Reverberation for 530- and 1030- cps Sound in Deep Water," Acoustical Society of America. Journal, v.33, p.1498-1504, November 1961
8. Mackenzie, K. V., "Long-Range Shallow-Water Reverberation," Acoustical Society of America. Journal, v.34, p.62-66, January 1962
9. Mackenzie, K. V., "Reflection of Sound From Coastal Bottoms," Acoustical Society of America. Journal, v.32, p.221-231, February 1960
10. Lieberman, L. N., "Reflection of Sound From Coastal Sea Bottoms," Acoustical Society of America. Journal, v.20, p.305-309, May 1948

References (Continued)

11. Laughton, A. S., "Microtopography," p. 437-472 in The Sea; Ideas and Observations on Progress in the Study of the Seas, v. 3, by M. N. Hill, Interscience, 1963
12. Carsola, A. J., "Microrelief on Arctic Sea Floor," American Association of Petroleum Geologists. Bulletin, v. 38, p. 1587-1601, July 1954
13. Shipek, C. J., "Microrelief on the Sea Floor," The Science Teacher, v. 28, p. 6-13, October 1961
14. Shipek, C. J., "Surveying the Deep-Sea Floor With Cameras," Photogrammetric Engineering, v. 27, p. 84-91, March 1961
15. Crozier, W. J., "The Amount of Bottom Material Ingested by Holothurians (Stichopus)," Journal of Experimental Psychology, v. 26, p. 379-389, 1918
16. Mero, J. L., "Minerals on the Ocean Floor," Scientific American, v. 203, p. 64-72, December 1960
17. Navy Electronics Laboratory Report 916, Arctic SCUBA Diving, 1958. I: Observations on the Alaskan Arctic Shelf and Under Ice in the Polar Sea; II: Notes on the Arctic Diving Operation, by G. Shumway and J. A. Beagles, 17 August 1959
18. Heezen, B. C. and Ewing, M., "Turbidity Currents and Submarine Slumps, and the 1929 Grand Banks Earthquake," American Journal of Science, v. 250, p. 849-873, December 1952
19. Heezen, B. C., "Dynamic Processes of Abyssal Sedimentation: Erosion, Transportation, and Redeposition on the Deep-Sea Floor," Geophysical Journal, v. 2, p. 142-163 June 1959

References (Continued)

20. Emery, K. O., The Sea Off Southern California, Wiley, 1960
21. Moore, D. G., "Acoustic-Reflection Studies of the Continental Shelf and Slope Off Southern California," Geological Society of America. Bulletin, v.71, p.1121-1136, August 1960
22. Arrhenius, G., "Pelagic Sediments in the Sea," p.655-727 in The Sea; Ideas and Observations on Progress in the Study of the Seas, v.3, by M. N. Hill, Interscience, 1963
23. Hydrographic Office Informal Manuscript Report 0-48-62, Subaerial Features of the Floor of the Tongue of the Oceans, Bahamas, by R. F. Busby, 1962
24. Shipek, C. J., "Photographic Survey of Sea Floor on Southwest Slope of Eniwetok Atoll," Geological Society of America. Bulletin, v.73, p.805-812, July 1962
25. Menard, H. W., "Deep Ripple Marks in the Sea," Journal of Sedimentary Petrology, v.22, p.3-9, March 1952
26. Army Department. Beach Erosion Board Technical Memorandum 100, Wave-Generated Ripples in Nearshore Sands, by D. L. Inman, October 1957
27. Naval Oceanographic Office Technical Report TR-108, Submarine Geology of the Tongue of the Ocean, Bahamas, by R. F. Busby, November 1962
28. Dzulynski, S. and Sanders, J. E., "Current Marks on Firm Mud Bottoms," Connecticut Academy of Arts and Sciences. Transactions, v.42, p.57-96, 1962
29. Goldberg, E. D. and Arrhenius, G. O. S., "Chemistry of Pacific Pelagic Sediments," Geochimica et Cosmochimica Acta, v.13, p.153-212, 1958

References (Continued)

30. Graham, J. W., "Metabolically Induced Precipitation of Trace Elements From Sea Water," Science, v.129, p.1428-1429, 22 May 1959
31. Mero, J. L., "Ocean-Floor Manganese Nodules," Economic Geology and the Bulletin of the Society of Economic Geologists, v.57, p.747-767, August 1962
32. Menard, H. W., "Consolidated Slabs on the Floor of the Eastern Pacific," Deep-Sea Research, v.7, p.35-41, 1960
33. Olausson, E. and Uusitalo, S., "On the Influence of Seismic Vibrations on Sediments," Société Géologique de Finlande. Comptes Rendus, v.35, p.101-114, 1963
34. Zenkevich, L. A. and Birstein, J. A., "Studies of the Deep Water Fauna and Related Problems," Deep-Sea Research, v.4, p.54-64, 1956
35. Spärck, R., "The Density of Animals on the Ocean Floor," p.196-201 in The Galathea Deep Sea Expedition, 1950-1952, Macmillan, 1956
36. Sokolova, M. N., "On the Distribution of Deep-Water Bottom Animals in Relation to Their Feeding Habits and the Character of Sedimentation," Deep-Sea Research, v.6, p.1-4, 1959

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY <i>(Corporate author)</i> Navy Electronics Laboratory San Diego, California 92152		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
		2b. GROUP
3. REPORT TITLE PHOTO ANALYSIS OF SEA FLOOR MICRORELIEF		
4. DESCRIPTIVE NOTES <i>(Type of report and inclusive dates)</i> Research Report		
5. AUTHOR(S) <i>(Last name, first name, initial)</i> Shipek, C. J.		
6. REPORT DATE 26 April 1966	7a. TOTAL NO. OF PAGES 70	7b. NO. OF REFS 36
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) 1374	
b. PROJECT NO. SR 104 03 01 c. Task 0539 NEL L40161 d.	9b. OTHER REPORT NO(S) <i>(Any other numbers that may be assigned this report)</i>	
10. AVAILABILITY/LIMITATION NOTICES DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Bureau of Ships, Department of the Navy Washington, D. C. 20360	
13. ABSTRACT Over 700 NEL sea floor photographs, obtained intermittently over a 10-year period, in a wide variety of environments, have been analyzed. Five major types of bottom microrelief, on the basis of origin, have been defined; a numerical scale of microroughness has been developed and ratings predicted for normal bottom environments; and zones of isoroughness have been delineated for large portions of the Pacific and adjacent ocean areas.		

DD FORM 1 JAN 64 1473 0101-807-6800

UNCLASSIFIED

Security Classification

UNCLASSIFIED
Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Ocean Bottom Photographic Analysis						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- "Qualified requesters may obtain copies of this report from DDC."
- "Foreign announcement and dissemination of this report by DDC is not authorized."
- "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

UNCLASSIFIED
Security Classification

<p>Navy Electronics Lab., San Diego, Calif. Report 1374</p> <p>PHOTO ANALYSIS OF SEA FLOOR MICRORELIEF, by C. J. Shipek 70p., 26 Apr 66.</p> <p>UNCLASSIFIED</p> <p>Over 700 NEL sea floor photographs, obtained intermittently over a 10-year period, in a wide variety of environments, have been analyzed. Five major types of bottom microrelief, on the basis of origin, have been defined; a numerical scale of micro-roughness has been developed and ratings predicted for normal bottom environments; and zones of isoroughness have been delineated for large portions of the Pacific and adjacent ocean areas.</p> <p>SR 104 03 01, Task 0539 (NEL L40161)</p> <p>This card is UNCLASSIFIED</p>	<p>1. Ocean Bottom - Photographic Analysis</p> <p>I. Shipek, C. J.</p> <p>SR 104 03 01, Task 0539 (NEL L40161)</p> <p>This card is UNCLASSIFIED</p>
<p>Navy Electronics Lab., San Diego, Calif. Report 1374</p> <p>PHOTO ANALYSIS OF SEA FLOOR MICRORELIEF, by C. J. Shipek 70p., 26 Apr 66.</p> <p>UNCLASSIFIED</p> <p>Over 700 NEL sea floor photographs, obtained intermittently over a 10-year period, in a wide variety of environments, have been analyzed. Five major types of bottom microrelief, on the basis of origin, have been defined; a numerical scale of micro-roughness has been developed and ratings predicted for normal bottom environments; and zones of isoroughness have been delineated for large portions of the Pacific and adjacent ocean areas.</p> <p>SR 104 03 01, Task 0539 (NEL L40161)</p> <p>This card is UNCLASSIFIED</p>	<p>1. Ocean Bottom - Photographic Analysis</p> <p>I. Shipek, C. J.</p> <p>SR 104 03 01, Task 0539 (NEL L40161)</p> <p>This card is UNCLASSIFIED</p>

INITIAL DISTRIBUTION LIST

COMMANDER, NAVAL SHIP SYSTEMS COMMAND
 CODE 1610
 CODE 1620
 CODE 1631
 CODE 2021 (2)
 CODE 204113
 COMMANDER, NAVAL SHIP ENGINEERING CENTER
 CODE 6420
 CODE 6440
 CODE 6452E
 CODE 6454
 CHIEF OF NAVAL MATERIAL
 WAC 033
 NAVAL ELECTRONIC SYSTEMS COMMAND HEADQUARTERS
 LIBRARY
 CHIEF, BUREAU OF NAVAL WEAPONS
 DL1-3
 DL1-304
 FASS
 R-56
 RU-222
 RUDC
 RUDC-2
 RUDC-3
 CHIEF, BUREAU OF YARDS AND DOCKS
 CODE 42310
 CHIEF OF NAVAL PERSONNEL
 PERS 118
 CHIEF OF NAVAL OPERATIONS
 OP-312 F
 OP-077
 OP-701
 OP-71
 OP-03EG
 OP-09B5
 OP-311
 OP-322C
 OP-702C
 OP 716
 OP-922YAC1
 CHIEF OF NAVAL RESEARCH
 CODE 416
 CODE 418
 CODE 427
 CODE 466
 CODE 468
 CODE 493
 COMMANDER IN CHIEF US PACIFIC FLEET
 COMMANDER IN CHIEF US ATLANTIC FLEET
 COMMANDER OPERATIONAL TEST AND
 EVALUATION FORCE
 DEPUTY COMMANDER OPERATIONAL TEST +
 EVALUATION FORCE, PACIFIC
 COMMANDER SUBMARINE FORCE
 US PACIFIC FLEET
 US ATLANTIC FLEET
 DEPUTY COMMANDER SUBMARINE FORCE,
 US ATLANTIC FLEET
 COMMANDER ANTISUBMARINE WARFARE FOR
 US PACIFIC FLEET
 COMMANDER FIRST FLEET
 COMMANDER SECOND FLEET
 COMMANDER TRAINING COMMAND
 US ATLANTIC FLEET
 OCEANOGRAPHIC SYSTEM PACIFIC
 COMMANDER SUBMARINE DEVELOPMENT
 GROUP TWO
 COMMANDER KEY WEST TEST +
 EVALUATION DETACHMENT
 DESTROYER DEVELOPMENT GROUP PACIFIC
 FLEET AIR WINGS, ATLANTIC FLEET
 SCIENTIFIC ADVISORY TEAM
 US NAVAL AIR DEVELOPMENT CENTER
 NADC LIBRARY
 US NAVAL MISSILE CENTER
 TECH. LIBRARY
 PACIFIC MISSILE RANGE /CODE 3250/
 US NAVAL ORDNANCE LABORATORY
 LIBRARY
 SYSTEMS ANALYSIS GROUP OF THE ASW
 R-D PLANNING COUNCIL,
 CODE RA
 US NAVAL ORDNANCE TEST STATION
 PASADENA ANNEX LIBRARY
 CHINA LAKE
 US NAVAL WEAPONS LABORATORY
 KXL
 LIBRARY
 PEARL HARBOR NAVAL SHIPYARD
 PORTSMOUTH NAVAL SHIPYARD
 PUGET SOUND NAVAL SHIPYARD
 SAN FRANCISCO NAVAL SHIPYARD
 USN RADIOLOGICAL DEFENSE LABORATORY
 DAVID TAYLOR MODEL BASIN
 /LIBRARY/
 US NAVY MINE DEFENSE LABORATORY
 US NAVAL TRAINING DEVICE CENTER
 CODE 365H, ASW DIVISION
 USN UNDERWATER SOUND LABORATORY
 LIBRARY
 CODE 905
 ATLANTIC FLEET ASW TACTICAL SCHOOL
 USN MARINE ENGINEERING LABORATORY
 US NAVAL CIVIL ENGINEERING LAB.
 L54
 US NAVAL RESEARCH LABORATORY
 CODE 2027
 CODE 5440
 US NAVAL ORDNANCE LABORATORY
 CORONA
 USN UNDERWATER SOUND REFERENCE LAB.
 US FLEET ASW SCHOOL
 US FLEET SONAR SCHOOL
 USN UNDERWATER WEAPONS RECH & ENG STATION
 OFFICE OF NAVAL RESEARCH
 PASADENA
 US NAVAL SHIP MISSILE SYSTEMS
 ENGINEERING STATION
 CHIEF OF NAVAL AIR TRAINING
 USN WEATHER RESEARCH FACILITY
 US NAVAL OCEANOGRAPHIC OFFICE
 SUPERVISOR OF SHIPBUILDING US NAVY
 GROTON
 US NAVAL POSTGRADUATE SCHOOL
 LIBRARY(CODE 0384)
 DEPT. OF ENVIRONMENTAL SCIENCES
 OFFICE OF NAVAL RESEARCH BR OFFICE
 LONDON
 BOSTON
 CHICAGO
 SAN FRANCISCO
 FLEET NUMERICAL WEATHER FACILITY
 US NAVAL APPLIED SCIENCE LABORATORY
 CODE 9200, ELECTRONICS DIVISION
 CODE 9832
 US NAVAL ACADEMY
 ASSISTANT SECRETARY OF THE NAVY R-D
 US NAVAL SECURITY GROUP HDQTRS(G43)
 ONR SCIENTIFIC LIAISON OFFICER
 WOODS HOLE OCEANOGRAPHIC INSTITUTION
 INSTITUTE OF NAVAL STUDIES
 LIBRARY
 AIR DEVELOPMENT SQUADRON ONE /VX-1/
 SUBMARINE FLOTILLA ONE
 DEFENSE DOCUMENTATION CENTER (20)
 DOD RESEARCH AND ENGINEERING
 WEAPONS SYSTEMS EVALUATION GROUP
 DEFENSE ATOMIC SUPPORT AGENCY
 NATIONAL OCEANOGRAPHIC DATA CENTER
 US COAST GUARD OCEANOGRAPHIC UNIT
 COMMITTEE ON UNDERSEA WARFARE
 US COAST GUARD HDGTRS(OSR-2)
 ARCTIC RESEARCH LABORATORY
 WOODS HOLE OCEANOGRAPHIC INSTITUTION
 US COAST AND GEODETIC SURVEY
 WASHINGTON SCIENCE CENTER - 23
 FEDERAL COMMUNICATIONS COMMISSION
 US WEATHER BUREAU
 DIRECTOR, METEOROLOGICAL RESEARCH
 LIBRARY
 NATIONAL SEVERE STORMS LABORATORY
 NORMAN, OKLAHOMA 73069
 NATIONAL BUREAU OF STANDARDS
 BOULDER LABORATORIES
 US GEOLOGICAL SURVEY LIBRARY
 DENVER SECTION
 US BUREAU OF COMMERCIAL FISHERIES
 LA JOLLA
 WASHINGTON, D. C. 20240
 WOODS HOLE, MASSACHUSETTS 02543
 HONOLULU, HAWAII 96812
 STANFORD, CALIFORNIA 94305
 TUNA RESOURCES LAB LA JOLLA
 ABERDEEN PROVING GROUND, MD. 21005
 REDSTONE SCIENTIFIC INFORMATION
 CENTER
 US ARMY ELECTRONICS R-O LABORATORY
 ANSEL-RD-MAT
 COASTAL ENGINEERING RESEARCH CENTER
 CORPS OF ENGINEERS, US ARMY
 HEADQUARTERS, US AIR FORCE
 AFRSTA
 AIR UNIVERSITY LIBRARY
 AIR FORCE EASTERN TEST RANGE
 /AFMTC TECH LIBRARY - MU-135/
 AIR PROVING GROUND CENTER, PG8PS-12
 HD AIR WEATHER SERVICE
 WRIGHT-PATTERSON AF BASE
 SYSTEMS ENGINEERING GROUP (RTD)
 UNIVERSITY OF MICHIGAN
 OFFICE OF RESEARCH ADMINISTRATION
 UNIVERSITY OF MIAMI
 THE MARINE LAB. LIBRARY
 MICHIGAN STATE UNIVERSITY
 COLUMBIA UNIVERSITY
 HUDSON LABORATORIES
 LAMONT GEOLOGICAL OBSERVATORY
 DARTMOUTH COLLEGE
 RADIOPHYSICS LABORATORY
 CALIFORNIA INSTITUTE OF TECHNOLOGY
 JET PROPULSION LABORATORY
 HARVARD COLLEGE OBSERVATORY
 OREGON STATE UNIVERSITY
 DEPARTMENT OF OCEANOGRAPHY
 UNIVERSITY OF WASHINGTON
 DEPARTMENT OF OCEANOGRAPHY
 FISHERIES-OCEANOGRAPHY LIBRARY
 NEW YORK UNIVERSITY
 DEPT OF METEOROLOGY + OCEANOGRAPHY
 UNIVERSITY OF MICHIGAN
 DIRECTOR, COLLEY ELECTRONICS LAB
 DR. JOHN C. AYERS
 UNIVERSITY OF WASHINGTON
 DIRECTOR, APPLIED PHYSICS LABORATORY
 OHIO STATE UNIVERSITY
 PROFESSOR L. E. BOLLINGER
 UNIVERSITY OF ALASKA
 GEOPHYSICAL INSTITUTE
 UNIVERSITY OF RHODE ISLAND
 NARRAGANSETT MARINE LABORATORY
 YALE UNIVERSITY
 BINGHAM OCEANOGRAPHIC LABORATORY
 FLORIDA STATE UNIVERSITY
 OCEANOGRAPHIC INSTITUTE
 UNIVERSITY OF HAWAII
 HAWAII INSTITUTE OF GEOPHYSICS
 ELECTRICAL ENGINEERING DEPT
 HARVARD UNIVERSITY
 GORDON MCKAY LIBRARY
 A-M COLLEGE OF TEXAS
 DEPARTMENT OF OCEANOGRAPHY
 THE UNIVERSITY OF TEXAS
 DEFENSE RESEARCH LABORATORY
 ELECTRICAL ENGINEERING RESEARCH LAB
 HARVARD UNIVERSITY
 UNIVERSITY OF CALIFORNIA-SAN DIEGO
 SCRIPPS INSTITUTION OF OCEANOGRAPHY
 MARINE PHYSICAL LAB
 PENNSYLVANIA STATE UNIVERSITY
 ORDNANCE RESEARCH LABORATORY
 NAVAL WARFARE RESEARCH CENTER
 STANFORD RESEARCH INSTITUTE
 MASSACHUSETTS INST OF TECHNOLOGY
 ENGINEERING LIBRARY
 MIT-LINCOLN LABORATORY
 RADIO PHYSICS DIVISION
 THE JOHNS HOPKINS UNIVERSITY
 APPLIED PHYSICS LABORATORY
 INSTITUTE FOR DEFENSE ANALYSES
 FLORIDA ATLANTIC UNIVERSITY