J-4

PL 242

Engraved by G. S. S. S.

Engraved by G. S. S. S.

Trachea of

1. 2. *Chloephaga Magellanica* 3. 4. *Tadorna Pelloni*

5. 6. *Carina Moschata* 7. *Melanitta fusca*.

G. Schaeffer del. et lith.

Printed by C. Hallman del.

Trachea of

Fig 1 *Aia sponsa*

2 *Querquedula formosa*

3 *Cyanopterus circeus*

4 *Micropterus latirostris*

5 *Bernicula antarctica*

All males

Chased
and Quarrel
Birds

A

MONOGRAPH

ON

THE ANATIDÆ,

OR

DUCK TRIBE.

BY

sonus
T. C. EYTON, ESQ. F.L.S. F.Z.S.

LONDON:

LONGMAN, ORME, BROWN, GREEN, & LONGMAN, PATERNOSTER-ROW;
AND EDDOWES, SHREWSBURY.

MDCCCXXXVIII.

Since the greater part of this work was printed we have received Mr. Swainson's volume of Dr. Lardner's Cyclopædia, entitled "Animals in Menageries." There is little new in it as regards the class we at present are endeavouring to illustrate, the greater part of the letter-press being copied from Latham. The following changes in nomenclature, however, occur, as regards which we shall leave our readers to judge for themselves, merely stating that we cannot agree with Mr. Swainson in them: viz. the specific name of the only known species of *Cereopsis* is changed from *Novæ Hollandiæ* to *Australis*; the semipalmated goose (*Anas semipalmata*, Lath. *Choristopus semipalmata nobis*) is placed in *Dendrocygna*, Sw. being the same genus in which *Anas*

PREFACE.

arcuata, *Horsf.* has been before placed by Mr. Swainson, and on which species he instituted the genus; the musk duck (*Carina moschata*, *Flem.*) is placed in the same genus with the common sheldrake (*Tadorna*); the blue-winged teal (*Anas discors*, *Linn.* *Cyanopterus discors nobis*), the Baikal teal (*Anas formosa*, *Linn.* *Querquedula formosa nobis*), and the Java teal (*Anas Javanensis*, *Lath.* *Querquedula Javanensis nobis*), are all placed in *Genus Boschus*, *Sw.* together with the common wild duck. The specific name also of *Javanensis* is altered to *Javanica*.

We have also, since the printing of our generic characters, seen an article in the "Penny Cyclopædia" (*Fuligulineæ*) relating to a portion of the *Anatidæ*, in which we are informed that the generic appellation of *Chauliodus* has been pre-occupied: as we find on reference this to have been the case, we venture to propose that of *Ktinorhynchus* instead, alluding to the comb-like appearance of the lamellæ on each side of the bill.

In the Catalogue of European and American Birds, just published by the Prince of Musignano, the generic name *Chauliodes* is altered to *Chaulelasmus*, *Gray*, which had priority to our name for the same genus, having been published first; and *Fuligula Americana*, *Bonap.* is altered to *Aythya Erythrocephala*, *Bonap.* which genus we do not think a good one; its true name, therefore, will be *Fuligula Erythrocephala*.

Brehm's work "Handbuech der Naturgeschite aller vogel Deutschlands" would have been scarcely worthy of notice, had it not been that some new genera had been instituted in it, and also the synonyms of Boie, which are valuable, as they forestall many of those given by authors in this country. Looking at the work, however, altogether, it is worse than those of the older authors as regards the creation of false species, five or six species in many instances being made of one.

CONTENTS.

PREFACE.	PAGE.
Introductory Chapter	1
Characters of Family Anatidæ	5
Sub-family Plectropterinae	7
Genus Choristopus	8
Plectropterus	10
Sub-family Anserinae	11
Genus Cereopsis	12
Chloephaga	13
Bernicla	14
Cheniscus	15
Anser	16
Cygnus	18
Sarkidiornis	20
Chenalopex	22
Sub-family Anatinae	24
Genus Tadorna	25
Casarka	27
Dendrocygna	28
Leptotarsis	29
Dafila	30
Pæcillonitta	32
Mareca	33
Aia	35
Querquedula	37
Cyanopterus	38
Rhynchaspis	40
Malacorhynchus	42
Chauliodes chauleasmus	43
Anas	45
Carina	47
Sub-family Fuligulinæ	49
Genus Micropterus	50
Melanitta	52
Somateria	54
Polysticta	55
Kamptorhynchus	57
Callichen	59
Fuligula	61
Nyroca	63
Harelda	65
Clangula	67
Sub-family Erismaturinae	69
Genus Thalassornis	70
Biziura	71
Erismatura	73

CONTENTS.

Characters of Sub-family Merginæ	75
Genus Mergus	76
Synopsis speciorum	78
Characters of Choristopus semipalmatus	78
Plectropterus Gambensis	79
Cereopsis Novæ Hollandiæ	80
Chloephaga Sandvicensis	81
Magellanicæ	82
Bernicla antarctica	84
ruficollis	85
leucopsis	86
jubata	87
Cheniscus Madagascariensis	88
Coromandelianus	89
Anser indicus	90
erythropus	90
segetum	91
hyperboreas	92
melanopterus	93
ferus	93
Guineensis	94
Hutchinsii	95
Canadensis	95
Cygnus olor	96
immutabilis	96
atratus	97
nigricollis	98
Bewickii	98
Americanus	99
buccinator	100
anatoides	101
ferus	101
Sarkidiornis regius	102
Africanus	103
Chenalopex Ægyptiaca	103
cana	104
Tadorna bellonii	105
radjah	106
Casarka rutila	106
tadanoides	107
castanea	108
Dendrocygna arcuata	108
autumnalis	109
viduata	110
arborea	110
Leptotarsis Eytoni	111
Dafila urophasianus	112
pyrogaster	113
caudacuta	113
marmorata	115
Pæcillonitta erythrorhyncha	115
Bahamensis	116

CONTENTS.

Characters of <i>Mareca Americana</i>		116
— <i>Chiloensis</i>		117
— <i>fistularis</i>		118
— <i>castanea</i>		119
<i>Aia galericulata</i>		119
— <i>sponsa</i>		120
<i>Querquedula Javanensis</i>		121
— <i>falcaria</i>		126
— <i>erythrorhyncha</i>		127
— <i>Carolinensis</i>		124
— <i>formosa</i>		124
— <i>Manillensis</i>		125
— <i>glocitans</i>		122
— <i>crecca</i>		123
— <i>creccoides</i>		128
— <i>Capensis</i>		128
— <i>Hottentota</i>		129
<i>Cyanopterus circa</i>		130
— <i>fretensis</i>		131
— <i>discors</i>		131
— <i>Rafflesii</i>		132
<i>Rhynchaspis rhynchotis</i>		133
— <i>maculatus</i>		134
— <i>clypeata</i>		134
— <i>Capensis</i>		135
<i>Malacorhynchus membrinacea</i>		136
<i>Ktinorhynchus (Chaulelasmus) strepera</i>		137
<i>Anas pæcilorhyncha</i>		138
— <i>specularis</i>		138
— <i>superciliosa</i>		139
— <i>obscura</i>		140
— <i>boschas</i>		140
— <i>flavirostris</i>		141
— <i>sparsa</i>		142
<i>Carina moschata</i>		142
<i>Micropterus patachonicus</i>		143
— <i>brachypterus</i>		144
<i>Melanitta Americana</i>		144
— <i>nigra</i>		145
— <i>perspicillata</i>		146
— <i>fusca</i>		147
<i>Somateria spectabilis</i>		148
— <i>St. Cuthberti</i>		149
<i>Polysticta Stellerii</i>		150
<i>Kamptorhynchus Labradorus</i>		151
<i>Callichen caryophylaceum</i>		152
— <i>rufinum</i>		152
<i>Fuligula valisneria</i>		153
— <i>ferina</i>		154
— <i>Americana</i>		155
— <i>marilla</i>		156

CONTENTS.

Characters of	Fuligula affinis					157
—	rufitorques				..	158
—	cristata					159
Nyroca	Australis				..	160
—	leucophthalmus					160
—	brunnea				..	161
Harelda	glacialis					162
Clangula	histrionica				..	163
—	albeola					164
—	Barrovii				..	165
—	vulgaris					166
—	Americana				..	167
—	leuconotus					168
Biziura	lobata				..	168
Erismatura	maccoa					169
—	ferruginea				..	170
—	leucocephala					170
—	rubida				..	171
—	Australis					172
—	Dominica				..	172
Mergus	merganser					173
—	serrator				..	175
—	Brazilianus					176
—	cucullatus				..	177
—	albellus					178

Appendix, No. 1.

Appendix, No. 2.

Supplement.

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

..

LIST OF PLATES.

DIRECTIONS TO THE BINDER.

1. Trachea of <i>Cleophaga Magellanica</i> , <i>Tadorna bellonii</i> . <i>Carina moschata</i> . <i>Melanitta fusca</i>	} to be placed before title page.
2. Trachea of <i>Aia sponsa</i> , <i>Querquedula formosa</i> , <i>Cyanopterus circia</i> , <i>Micropterus brachyptera</i> , <i>Bernicla antarctica</i>	
3. Posterior margins of the sterna of <i>Alca torda</i> , <i>Phalacrocorax cristatus</i> , <i>Podiceps cristatus</i> , <i>Clangula vulgaris</i> , <i>Fuligula cristata</i> , <i>Mergus albellus</i> , <i>Rhynchaspis clypeata</i> , <i>Carina mos-</i> <i>chata</i> , <i>Tadorna bellonii</i> , <i>Anser ferus</i> , <i>Cygnus olor</i>	to face p. 5.
4. <i>Œsophagus</i> , proventriculus, stomach, and cæca of <i>Anser segetum</i>	— p. 11.
5. Skeleton of <i>Anser cygnoides</i>	— p. 16.
6. Skeleton and trachea of <i>Cygnus ferus</i>	— p. 18.
7. ————— <i>Chenalopex Ægyptiaca</i>	— p. 22.
8. <i>Œsophagus</i> , proventriculus, stomach, and cæca of <i>Querquedula crecca</i> ; cæca of <i>Tadorna bellonii</i>	— p. 24.
9. Skeleton of <i>Tadorna bellonii</i>	— p. 25.
10. ————— and trachea of <i>Querquedula crecca</i>	— p. 37.
11. ————— <i>Anas boschas</i>	— p. 45.
12. <i>Œsophagus</i> , proventriculus, stomach, and cæca of <i>Melanitta nigra</i> ; cæca of <i>Fuligula ferina</i> , and <i>Clangula vulgaris</i>	— p. 49.
13. Skeleton and trachea of <i>Nyroca leucophthalmus</i>	— p. 63.
14. ————— <i>Harelda glacialis</i>	— p. 65.
15. Skeleton of <i>Clangula histrionica</i> ; trachea of <i>Clangula vulgaris</i>	— p. 67.
16. <i>Œsophagus</i> , proventriculus, stomach, and cæca of <i>Mergus serrator</i>	— p. 75.
17. Skeleton and trachea of <i>Mergus serrator</i>	— p. 76.
18. Trachea and muscles of voice in <i>Cygnus Bewickii</i>	— p. 98.
19. <i>Casarka castanea</i>	— p. 108.
20. <i>Dafila urophasianus</i> , male	— p. 112.
21. <i>Mareca Chiloensis</i> , male	— p. 117.
22. <i>Mareca castanea</i> , male	— p. 119.
23. <i>Nyroca brunnea</i> , male and female	— p. 161.
24. <i>Oxyura (Erisimatura) Maccoa</i> ; <i>Oxyura Dominica</i> , fem.	— p. 169.

CORRIGENDA.

p. 16.	1.	6.	<i>for</i>	instructis	<i>read</i>	instructa.
—	7.			inferiorum		inferiorem.
—	—			contractam		contracta.
22.	4.			calcaria		calcari.
—	—			una		uno.
24.	5.			processibusque		processibus.
28.	2.			armatæ		armati.
29.	2.			complanatâ		complanatum.
—	3.			instructo		instructum.
30.	6.			onga		longa.
33.	5.			instructis		instructa.
37.	19.			1½		2½.
43.	1.			Chauliodes		Chaulelasinus.
43.	3.		<i>dele</i>	vix elongatæ.		
—	7.		<i>for</i>	laterater		lateralliter.
45.	18.			boschas		pæcilorhyncha.
49.	9.			extremus		extremas.
—	10.			rami		ramorum.
54.	13.			mollissima		spectabilis.
61.	6.			instructis		instructa.
—	20.			marilla		cristata.
65.	3.		<i>after</i>	parte	<i>add</i>	anteriore.
80.	1.		<i>for</i>	Nova	<i>read</i>	Novæ.
82.	3.		<i>after collo dele que, and add que after pedibus.</i>			

INTRODUCTORY CHAPTER.

BEFORE entering upon a description of the group we propose to illustrate, it may not be amiss to make a few observations on some of the divisions at present in use among Ornithologists, more particularly Genera and Species, for we regard all other divisions, if indeed Genera are not so, as certainly to a great extent arbitrary, and constituted merely for the convenience of the student in Zoology, that he may at once turn to the required species or group he may wish to find, instead of, as would be the case if he consulted the works of Buffon, having to turn over volumes before he would be able to find the wished-for place. When we use the term arbitrary, as applied to divisions of the animal kingdom, let them be called either classes, families, sub-families, or by any other name, we mean, that in forming them there are generally rejected as many points of connection between contiguous groups, and often stronger ones, than those made use of for their division. Divisions and arrangements of some sort are, however, absolutely necessary for the Zoologist, under each of which should be stated the points of connection it bears to others. Much has been said and written on analogy and affinity, and the connection by one or the other of them between the groups and species of the animal kingdom. We have not, however, been able to distinguish between them in any other manner than that the former is generally applied when the groups or species between which a connexion is supposed to exist are far removed from each other, and the latter when nearly related, we shall use the terms indiscriminately, as convenient.

A definition of species again presents to the Zoologist in particular another difficulty.

The generally received opinion is that of John Hunter, viz. that hybrids between true species will not be productive; and this, we are inclined to believe, is partially correct, but not entirely so, as some birds, in a state of domestication, have bred together, and their offspring been productive, although differing most materially in external form.

It may be advanced, however, and with truth, that those animals upon which this experiment has been tried have invariably been brought from countries far apart, and that consequently in a wild state the experiment has never been tried; no fact, that we are aware of, can be brought forward towards an answer to this objection. Should it prove true, that animals inhabiting different countries, and with slightly different forms and colouring, are of the same species, and we confess there is good reason for supposing that this may be the case, it can only be accounted for in this mode, namely, that at the universal distribution of animals after the Deluge, those of the same species, and derived from the same parents, going to different localities, have in a succession of ages been influenced by various local circumstances, as climate, the plentitude, the want, or nature of food, which causes have changed their form, colouring, and, in many instances, their habits.

We have nothing whatever to urge against this theory, and we find much in favour of it, considering the different varieties of the human form and habits, although we cannot for a moment suppose that they are derived from other than the same common parents, and that the distinctions between the different races are caused otherwise than solely by climate, yet do we make

species of animals inhabiting the same countries, where man has undergone changes equal to any specific character we are able to give between his inferiors. As far, however, as our data at present go, the theory of Hunter, excepting those animals in a state of domestication, appears to be the true one, and we shall here take the liberty of referring our readers to some short papers by us in the Zoological proceedings for 1837, and to Loudon's Magazine of Natural History, No. VII. New Series, and also to a Note appended to a paper by Mr. Blyth, in volume IX. page 511, of the same Magazine.

That man is subjected to a change in complexion as well as in the formation of the skull, when acted upon by different climates, appears from the following quotation:—"In Cairo, and throughout the northern provinces, those (Mooslim Egyptians) who have not been much exposed to the sun have a yellowish but very clear complexion and soft skin, the rest are of a considerably darker and coarser complexion. The people of Middle Egypt are of a more tawny colour, and those of the more southern provinces are of a deep bronze or brown complexion, darkest towards Nubia, where the climate is hottest." Vide "An Account of the Manners and Customs of the Modern Egyptians, by W. Lane, Vol. I. Library of Entertaining Knowledge."

From a paper read at the meeting of the British Association, held at Liverpool, by Dr. Warren (of Boston, U. S.), it appears that the crania of Indians found buried in mounds in North America, and on that account called the Mound Indians, differ in form from those of the Indians at present inhabiting the same country, and agree with those of the Peruvians now inhabiting South America. Are the present North American Indians the posterity of these Mound Indians? What evidence is there that they are not?

The geographical distribution of animals is a subject of much interest to the Zoologist as well as to the Geologist, and very closely connected with the distinction of species, more nearly so, indeed, than is generally supposed.

Mr. Swainson divides the earth into five Zoological provinces, corresponding with the continents of Europe, Asia, Africa, America, and Australia; these are well known divisions of the world, and as such are convenient, but we must consider that those countries occupied by the different races of mankind (except where emigration of late date has taken place) form much more natural Zoological provinces than those mentioned by Mr. Swainson.

The division of the human species which appears to be most generally approved and adopted is that of Blumenbach: with the characters of these races we have at present nothing to do, but shall refer those of our readers who wish to investigate the matter further to the "Decades of Crania," by the author above quoted, to the "Regne animal," and Dr. Prichard's "Researches into the Physical History of Man."

We shall, however, make use of that division of the Mongolian variety slightly hinted at by Cuvier, separating from it the Samoides or Lapland race, and it is a singular fact that the Zoology of the country, as well as the human inhabitants of it, form the connecting links between the Zoology and human inhabitants of the three great continents, Europe, Asia, and America.

The following divisions or ranges are those which we think the most natural:—

The Caucassian range occupies the whole of Europe, excepting only Lapland and Finland, that part of Asia to the west of the Caspian sea, and the rivers Ob and Ganges.

The Mongolian range occupies that part of Asia not inhabited by the Caucassian races, except Malacca.

The Esquimaux range contains those most northern parts of the three continents of Europe, Asia, and America, not occupied by the other races.

The Ethiopian or Negro range is confined to Africa, together with the island of Madagascar.

The American range occupies the whole of that vast continent, except the parts north of a line drawn from the northern side of Hudson's Bay to Cook's Inlet, which forms a part of the Esquimaux range.

The Malay range contains Malacca, Australia, and the intermediate islands of the Pacific and Indian Oceans, extending as far northwards as the Phillipine Group, New Zealand, and New Guinea.

The numerous islands not mentioned are considered as belonging to the ranges to which they are contiguous,

The advantage of this arrangement of the globe is simply that the zoology of the country occupied by each range will, in a great measure, as far as we have been able to ascertain, be found to be peculiar, the central portions of each district being occupied by a race of animals as well as by a variety of man distinct from those of the other ranges, the borders of each district, or those points at which they approach one another, being as it were inhabited by a mixed zoology, many of the species and Genera of either of the contiguous ranges being found in it.

There are, however, certain birds of great power of flight found distributed to a very large extent over the surface of the globe, the Kestrel and Peregrine Falcon may be enumerated as examples.

The object we have in view in this detail is an attempt, however imperfect, to shew that the same districts occupied by the various races of man are also occupied by various races of animals, each differing from the races of man and animals in the other ranges, and that it is exceedingly probable that as the races of man are merely varieties, one of the other those of animals may also be so.

And why is it, we may ask, in any way improbable that, at the dispersion of man and animals after the flood, the same causes may not have influenced both? We can observe even in the present state of science and observation some instances in which a parallel between animals and the human race, inhabiting the same countries, can be traced, and it may be expected that as science advances these parallels will increase: any one acquainted with this subject, upon seeing a box of African bird skins, will at once, without examining each species, say from whence they came, by their general appearance, and the darkness of their hues; the negro race from the same countries is also dark. Mr. Swainson, in volume 66 of Lardner's Cyclopædia, observes, that European Ornithology is characterised by the great number of Genera it contains in proportion to the number of species. There are also, perhaps, more sub-varieties of the Caucasian race than of any other, caused, probably, by the great variety of the climates.

The birds and animals of the Esquimaux range are nearly all migratory; the human inhabitants of the country are also seldom stationary for long together in the same spot, for the same reason, the want of food.

Thus, in some instances at least, we see that man and animals are influenced by the same cause, and why, therefore, is it by any means improbable that many of the animals, inhabiting different countries, and allowed to be allied, are of the same species, but have, by local causes, through a succession of ages, become altered?

Although we have said much, perhaps too much, on this subject, and, we fear, tired our

readers, let it be understood that the whole is merely brought forward as a conjecture, to which, whatever data we have been able to collect being given, we leave our readers to decide with regard to the probability or the contrary of it. Should it, however, prove true that many now supposed species, inhabiting different countries, are the same, and merely varieties, it will not lessen in any degree, or make less interesting or important, the labours of the zoologist; for, in an intellectual point of view, can any thing be more interesting than to unravel what appears to be a mystery, and, in an useful one, to point out such animals as will breed together, and thereby open the way to improvement in our varieties of domesticated animals.

In the internal arrangements of the group we intend to illustrate we shall deviate from the course followed by other authors, and propose one which, as far as we have been able to carry our investigations, appears to be in harmony with both the external and internal organization of each genus. We must here mention it as a subject of regret, that, although our utmost endeavours have been used, we have been totally, on account of their rarity in this country, unable to obtain either subjects for dissection, or even a sight of the skeletons, of many Genera.

Many parts throughout the group appear to be in favour of Mons. Temminck's opinion, that there are no such divisions as Genera in nature, the transition from one extreme of form to another being so gradual that it is difficult to say where to draw the line of division.

Upon a minute examination, however, as far at least as we are at present acquainted with the species, there is always found some break as it were between the forms constituting contiguous Genera or Subgenera, and some tangible distinction between them, although in many particulars they appear closely to approach. It is probable, however, that many new forms will yet be discovered; therefore, in the present state of science, it is impossible to say whether this opinion of Mons. Temminck's will eventually prove true or not. Should it be correct, a serious difficulty will arise with regard to nomenclature, in designating the objects to be named by a specific appellation alone. If we do not admit Genera we cannot admit families, orders, or classes; the transition will, therefore, we contend, be complete through all, before we can abolish the one or the other.

We shall now add a few words on the characters we have employed. Measurements are always subject to a certain degree of variation; we have, therefore, when the opportunity has offered, given a mean of two or three. The number of tail feathers is far too variable a character to be admitted into any ornithological work. We have now before us three specimens of *Erismatura rubida*, in which, although in this genus the tail is an important member, the numbering of each differs. We have, for the first time, as far as we are aware of, made use of anatomical characters for divisions, taken from the trachea and skeleton principally; this has not been done without due consideration, and the examination of a great number of specimens.

We have no doubt that some will find fault with us on account of the shortness of the descriptions at the head of each specific character, the use of these, at least what we have supposed it to be, is, that in looking over a large number of descriptions we may readily catch some mark whereby the object we wish to name may be distinguished, perhaps, in common with one or two others from its congeners, and thus that the time taken to read over a large number of long descriptions will be saved.

The characters of the Genera, and the cuts of bills and feet, are invariably taken from the species we have named as the generic type, unless it is mentioned to the contrary.

Pl. Engr. 22

1	Posterior margin of the Sternum of <i>Alca Litta</i>	6	<i>Mergus albellus</i>
2	<i>Phalaropus cristatus</i>	7	<i>Rhynchopus duplicata</i>
3	<i>Podiceps cristatus</i>	8	<i>Corvus Moschata</i>
4	<i>Clanula vulgaris</i>	9	<i>Tadorna bellena</i>
5	<i>Fulicula cristata</i>	10	<i>Anser ferus</i>
		11	<i>Cyanus ster</i>

ORDER NATATOIRES.

FAM. ANATIDÆ.

EXT. CHAR. *Rostrum* magnum epidermide molli potius quam cornuo veritabili tectum; lateribus lamellis vel dentibus parvis armatis. *Lingua* magna carnosae lateribus setis instructis. *Alæ* mediocres. *Pedes* fere vel omnino in equilibrio locati, digitis tribus anterioribus membrano junctis.

ANAT. CHAR. *Sternum* et pelvis magna et lata. *Gigeria* magna carnosae. *Cæca* longa. *Larynx* inferior in maribus fere bulbosa. *Trachea* aliquando convoluta, sæpius bulbosa. *Habitatio* aquatica,

EXT. CHAR. *Bill* large, covered with a soft epidermis rather than true horn, the sides armed with lamellæ or small teeth. *Tongue* large fleshy, with the margins toothed. *Wings* moderate. *Feet* placed in or near the equilibrium. *Toes*, with the three anterior ones, joined by a membrane. *Neck* long.

ANAT. CHAR. *Sternum* and pelvis large and broad. *Gizzard* large and fleshy. *Cæca* long. *Trachea* sometimes convoluted, more often bulbous. The inferior *Larynx* in the male commonly bulbous. Inhabits the water.

The intestinal anatomy of the family *Anatidæ* is characterised by the very great length of the alimentary canal compared to that of the bird: in some instances it is as many as five times its length, or even more. The cæcal appendages are also much elongated, being often equal to more than one-third the length of the body. The stomach is always large and muscular, forming a true gizzard. The œsophagus varies little in its diameter, but is generally slightly enlarged a little above the gizzard, again contracted, and then of nearly uniform diameter to the glottis.

The skeleton may be known by the length of the neck, and the great number of vertebræ of which it is composed, the large size of the sternum, the great developement of its keel and the form of its posterior extremity, which is always perforated by two foramina or indented by two fissures; the great length as well as breadth of the pelvis; the laterally flattened and anteriorly arched form of the rami of the os furcatum. The truncate form and great depth of the occipital portion of the cranium in a vertical direction, and by the fibula not being entirely ankylosed to the tibia.

Affinities. Considerable resemblance to the Divers is shewn in the *Merginæ* and *Erismaturinæ* in outward form, in the structure of the tail and the dense oily nature of the plumage. The

latter make an evident approach to the cormorants in particular ; nothing, however, is known of their anatomy. The *Merginæ* in their anatomy shew an affinity to the guillemots and the genera allied to them, chiefly consisting in the great developement of the keel of the sternum, and in its posterior edge being armed with an ensiform process. The pelvis also is much narrower and more elongated than is the case among the rest of the *Anatidæ*. The elongated head to the tibia possessed by all true divers is also slightly apparent in the genus *Mergus*.

We are aware by some authors that the Flamingo has been admitted among the *anatidæ*, chiefly on account of its slightly laminated bill and palmated feet, and some portions of its intestinal anatomy resembling this family : we cannot, however, follow this arrangement for the following reasons.

Characters for the larger divisions of birds are taken from the external organs necessary to procure their food, especially those of progression. The divisions of each of these have been made, on the other hand, chiefly from the different modifications of the bill.* Let us, for instance, take the order *grallatores* : we have there divisions or families formed for feeding on fish, mollusca, earth worms, and small insects : there is one general character of the skeleton through all. Not so with the bill and organs of digestion, which are modified, the former to procure and the latter to assimilate the different kinds of food necessary for the subsistence of the bird.

The same rule we shall find hold throughout all the different orders of birds, namely, that the skeleton and structure of the legs and feet will invariably furnish a sure guide to the Order ; while, on the other hand, the bill and organs of digestion supply characters for further subdivision.

What we have said above is by no means meant to imply that no modifications of the skeleton exist in the different divisions of each order, for such is not by any means the case ; some slight one always being to be found where there is a change of habit,—in other words, in the modes in which the various kinds of food are obtained.

The application of what we have said to the case in question is self-evident, that the flamingoes, having the organs of progression and skeleton peculiar to *grallatorial* birds, consequently must be placed in the order *Grallatores*, and that the bill and organs of digestion are fitted, the one for procuring, and the other for assimilating its natural food,—minute animalculæ obtained at the edges of waters by wading and not by swimming. The webbed feet also possessed by this bird peculiarly fit it for walking on the soft mud and sand, commonly found on shores ; their structure, however, is not that of any modification among the *anatidæ*, but essentially that of a crane, which bird has them also slightly webbed.

* Where such is not the case the orders and families are evidently unnatural.

SYNOPSIS
FAMILIARUM ET GENERUM.

SUB FAM. I. PLECTROPTERINÆ.

EXT. CHAR. *Rostrum* magnum, ad apicem angustatum, et ungue magnâ instructâ. *Plumæ* inter oculos rostrumque carent. *Tibiæ* parte quarta denudatæ. *Pedes* validi, digitis unguibus magnis instructis.

ANAT. CHAR. *Trachea* aliquando convoluta maribus sed non in sterno penetrans. *Caca* mediocria.

EXT. CHAR. *Bill* large, narrowed at the apex, and armed with a large claw. The space between the eyes and bill devoid of feathers. *Thighs* denuded for a quarter of their length. *Feet* strong, armed with sharp claws.

ANAT. CHAR. *Trachea* sometimes convoluted in the males, but not penetrating the sternum. *Caca* moderate.

Affinities. This curious group appears to lead directly to the grallatorial birds, but with the flamingo in particular it coincides to a great degree in the intestinal anatomy, while its skeleton, at least the fragments of it we have inspected, is that of a true goose, that of the flamingo approaching most intimately the *Ardeidæ*, which in our opinion (although in some particulars its intestinal anatomy appears to accord with the *Anatidæ*) entirely precludes it from being admitted into the order.

GENUS I. CHORISTOPUS.

EXT. CHAR. *Pedes* digitis longis, semi-membrinatis, unguibus magnis incurvatis instructis. *Tarsi* magni. *Tibiae* multum denudatae. *Rostrum* longum acuminatum cerâ ad oculos tendente. *Caput* osseâ tuberculâ supra nares et ungue, magnâ.

ANAT. CHAR. *Trachea* longa convoluta.

CHORISTOPUS nobis.

ANSER. *Linn. Auct.*

TYPE. *Choristopus semipalmatus.*

Affinity. This genus certainly approaches most nearly of any among the *Anatidæ* to the order *Grallatores*. To the genus *Plectropterus* it is nearly allied in form.

GENUS II. PLECTROPTERUS.

EXT. CHAR. *Pedes* membrinati digitis posterioribus longis, magnis, sed non lobatis, unguibus magnis et incurvatis instructis. *Rostrum* latum ungue magnâ instructâ. *Nares* ovales, in medio rostri positæ. *Alæ* una calcariâ armatæ.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes long, large and not lobed, armed with large and curved claws. *Bill* broad, furnished with a large nail. *Nostrils* oval, placed in the middle of the bill. *Wings* armed with one spur.

ANAT. CHAR. Unknown.

ANAS. *Linn.*

PLECTROPTERUS. *Steph.*

TYPE. *Plectropterus gambersis.*

Affinity. We are ignorant that any anatomical character of the genus has been ascertained; in default of which we have placed it next to the genus it appeared to approach in external character.

L. F. Sc.

1. Esophage Proventriculus & Stomach of *Muscovy duck* from
2. Caeca of Fowl

SUB FAM. II. ANSERINÆ.

EXT. CHAR. *Rostrum* altius-quam latum. *Pedes* mediocres validi, digitis posterioribus non lobatis.

ANAT. CHAR. *Sternum* mediocre, postice fissuris magnis et apertis; carinâ non antice productâ, latâ postice et non ad ultimam marginem sterni continuatâ. *Pelvis* mediocris. *Foramina ischiadica* magna et ovata. *Os furcatorium* validum arcuatum. *Processus coracoideus* validus et brevis. *Alæ* longæ. *Costæ* latæ, processu longo instructæ. *Trachea* fere complanata, sed aliquando bulbosa, et in sterno penetrans. *Larynx inferior* aliquando bulbosa.

EXT. CHAR. *Bill* higher than broad. *Feet* moderate, strong, the posterior toes not lobed.

ANAT. CHAR. *Sternum* moderate, furnished with large and open fissures behind: keel not produced anteriorly, broad, not continued to the posterior margin. *Pelvis* of moderate size, with the ischiadic foramina large and ovate. *Os furcatorium* strong, arcuated. *The coracoid process* strong and short. *Wings* long. *Ribs* broad, furnished with a long process. *Trachea* commonly of equal size. *The inferior larynx* sometimes bulbous.

HABIT. The birds contained in this family feed equally on the land and water; some species, however, scarcely ever approach it; the whole live almost entirely on vegetables. They have a very wide geographical range, and are mostly migratory.

TYPE. *Anser*.

Affinities. The nearest affinity it appears to present to the following sub-family appears to be through the genus *Chenalopez*. On the other hand the genus *Cereopsis* appears to approach the *Plectropterinæ* in its anatomy and habit.

GENUS I. CEREOPSIS.

EXT. CHAR. *Pedes* similibus genere sequenti. *Rostrum* breve, obtusum; culmine arcuato. *Nares* cerâ tectæ, magnæ, et rotundatæ.

ANAT CHAR. *Trachea* magna, complanata. *Larynx inferior* anseribus similis.

EXT. CHAR. *Feet* as in the preceding. *Bill* with a cere covering the nostrils; the culmen arched. *Nostrils* large, rounded.

ANAT. CHAR. The *Trachea* large, uniform. *Inferior Larynx* like that of the geese.

CEREOPSIS. *Lath.*

TYPE. *Cereopsis Novæ Hollandiæ*.

The only anatomical notes that I know of in existence, with regard to this curious bird, are those of Mr. Yarrell, in the Zoological Proceedings (1830, p. 25), which in giving our specific account of it, we cannot do better than extract.

Affinities. Another rather anomalous genus, more closely connected, however, with the geese than the preceding.

GENUS II. CHLOEPHAGA.

EXT. CHAR. *Pedes robusti. Digitis posterioribus mediocribus, unguibus incurvatis instructis. Membrana inter digitos antevius concavatis. Tarsi robusti longi. Rostrum robustum, culmine vix arcuatum, ungue magnâ instructâ. Nares lineares in medio rostri positæ. Lamellæ parvæ antice oblitteratæ.*

ANAT. CHAR. *Trachea bulbo uno latere osseo.*

EXT. CHAR. *Feet robust. Posterior toes moderate, armed with a curved claw, with the Membranes between the toes scolloped out in front. Tarsi large and strong. Bill robust, with the culmen slightly arched, armed with a large nail. Nostrils linear, placed in the middle of the bill. Lamellæ small, obliterated anteriorly.*

ANAT. CHAR. *Trachea with one osseous bulb on the side.*

ANAS. *Linn.*

BERNICLA. *Steph.*

CHLOEPHAGA *nobis.*

TYPE. *Chloephaga Magellanica.*

Affinities. The above genus we have formed on *Bernicla Magellanica*, which bird in outward form appears to make a link between the genera *Bernicla* and *Cercopsis*, and internally, as far as regards the trachea, presents an affinity to the *Anatinæ*.

GENUS III. BERNICLA.

EXT. CHAR. *Pedes parvi*, digitis posterioribus non lobatis, unguibus magnis instructis. *Tarsi elongati*. *Rostrum parvum*, capite brevius. *Nares lineares*, in medio rostri positæ.

ANAT. CHAR. *Trachea* tubulâ equali. *Larynx inferior* ossea aliquando bulbosa.

EXT. CHAR. *Feet* small, with the posterior toes not lobated; the nails large. *Tarsi* elongate. *Bill* small, shorter than the head. *Nostrils* linear, placed in the middle of the bill.

ANAT. CHAR. *Trachea* with the tube of equal size. The *inferior Larynx* osseous but not bulbous.

BERNICLA. *Steph.*

ANAS. *Linn.*

TYPE. *Bernicla jubata*.

The proportions of the skeleton are as follows:—Depth of the keel to the length of the sternum, as 1 to $3\frac{1}{2}$. Breadth of the pelvis to its length, as 1 to 2. Length of sternum to that of tibia, as 1 to $1\frac{2}{3}$. Height of cranium to its length, as 1 to $2\frac{1}{3}$. The keel is not continued to the posterior extremity of the sternum.

Affinity. Nearly allied, both in anatomy and habit, to the preceding and following.

GENUS IV. CHENISCUS.

EXT. CHAR. *Rostrum* parvum, ad basin elevatum, valde altius-quam latum, et ungue magnâ instructâ. *Nares* rotundatæ, culmen versus positæ. *Pedes* magni. *Digiti* longi et acutis unguibus armati. *Tarsi* breves. *Cauda* rotundata.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Bill* small, elevated at the base, much higher than broad, and armed with a large nail. *Nostrils* rounded, placed towards the culmen. *Feet* large. *Toes* long, and armed with sharp claws. *Tarsi* short. The *tail* rounded.

ANAT. CHAR. Unknown.

ANSER. Linn. *Lâth*.

BERNICLA. Steph.

QUERQUEULA. Gray.

CHENISCUS. Brookes' MSS.

TYPE. *Cheniscus Coromandeliana*.

Affinity. In the structure of the feet this genus bears considerable resemblance to the species of the genus *Anser*, adapted for swimming, though of extremely diminutive size. The bill, on the contrary, is that of *Bernicla*.

GENUS V. ANSER.

EXT. CHAR. *Pedes* digitis posterioribus non lobatis, unguibus magnis arcuatis. *Tarsi* robusti et magni. *Tibiæ* parte tertiâ denudatæ. *Rostrum* magnum, validum, ungue magna instructum, lateribus plerumque hiantibus. *Nares* ovales in medio rostri positæ. *Lingua* obtusa non ad apicem appendiculata, sed subtus cornea, in latera unâ ordine setarum brevium, et supra has setas multis rigidis spinis instructis ad apicem tendentibus.

ANAT. CHAR. *Trachea* tubula complanata, sed versus larynginem inferiorum contractam. *Larynx* inferior ossea, sed non tuberculosa.

EXT. CHAR. *Feet* with the posterior toes not lobated, armed with large claws. *Tarsi* robust and large. *Tibiæ* for a third part denuded. *Bill* large, strong, furnished with a large nail, the sides gaping. *Nostrils* oval, placed in the middle of the bill. *Tongue* obtuse, without an appendage, but below corneous, the sides armed with one rank of bristles, above and on the sides is a row of many stiff bristles extending to the tip.

ANAT. CHAR. *Trachea* with the tube of equal size, but contracted towards the inferior larynx. The *inferior Larynx* long and not tuberculous.

ANSER. *Auct.*

ANAS. *Linn.*

TYPE. *Anser fesus.*

The proportions of the skeleton are as follows:—Depth of the keel to the length of the sternum, as 1 to 4. Breadth of the pelvis to its length, as 1 to $1\frac{1}{2}$. Length of sternum to that of tibiæ, as 1 to $1\frac{2}{3}$. Height of cranium to its length, as 1 to 3.

ANGER EYFOIDES.
Hay the Ant. St.

Affinities. Some of the species of this genus approach near the swans, as *Anser Canadensis*, while others present an affinity to the Bernicles, as *Anser indica*.

GENUS VI. CYGNUS.

EXT. CHAR. *Pedes* magni, digitis posterioribus parvis, gibbosis, mediis longissimis. *Rostrium* magnum, lateribus parallelis. *Nares* subovales, in medio rostri positæ. *Cervix* longa.

ANAT. CHAR. *Trachea* tubulâ equali, aliquando in sterno penetrante. *Larynx inferior* ossea. *Sternum* carinâ anticè incurvatâ.

EXT. CHAR. *Feet* large, with the posterior toes small, gibbous, the middle ones longest. *Bill* large, the sides parallel. *Nostrils* suboval, placed in the middle of the bill. *Neck* long.

ANAT. CHAR. *Trachea* with the tube of equal size, sometimes penetrating the sternum. *Inferior Larynx* osseous. *Sternum* with the keel anteriorly curved.

CYGNUS. *Ray. Briss.*

ANAS. *Linn. Lath.*

TYPE. *Cygnus ferus.*

FRIGATA ACADEMICA,
the Red-tail.

collected by J. A. Smith.

The proportions of the skeleton are as follow:—Depth of the keel to the length of the sternum, as 1 to 4. Breadth of pelvis to its length, as 1 to $2\frac{1}{4}$. Length of sternum to that of tibia, as 1 to 2. Depth of cranium to its length, as 1 to 3.

Affinities. Although we have placed the swans with the geese, we are not by any means certain that it is their proper place.

GENUS VII. SARKIDIORNIS.

EXT. CHAR. *Pedes* digitis posterioribus medioeribus, in tarsi altè locatis. *Tarsi* medioeres. *Rostrum* in maribus carunculâ magnâ compressâ culmine. *Nares* rotundatæ magnæ. *Alæ* una calcaria armatæ, et remigibus apices versus incurvatis. *Cauda* ad apicem truncata.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes moderate, placed high on the tarsi. *Tarsi* moderate. *Bill* in the males with a large compressed caruncle on the top. *Wings* each armed with a spur, and with the quills incurved towards the points. *Tail* squared at the end.

ANAT. CHAR. Unknown.

CARINA. Steph.

ANSER. Linn.

SARKIDIORNIS nobis.

PLECTROPTERUS. Steph.

TYPE. *Sarkidiornis melanotus*.

Various have been the genera in which the bird before us has been placed (*Anser melanotus*, Linn.), with none of which does it agree. We have therefore proposed a new one for it.

Affinities. We have placed it, as far as we have been able (no anatomical character being known), in what appears to us to be its proper place, namely between the genera *Chenulopez* and *Anser*. It agrees with the former, in having a spur on the wing.

GENUS VIII. CHENALOPEX.

EXT. CHAR. *Pedes* digitis posterioribus gibbosis, et unguibus robustis. *Tarsi* longi. *Rostrum* validum ad basum tuberculatum. *Lamellæ* non ad marginem rostri continuatæ. *Nares* subrotundatæ in medio rostri positæ. *Alæ* calcaria una armatæ. *Cauda* ad apicem truncata.

ANAT. CHAR. *Trachea* tubulâ equali. *Larynx inferior* bulbo laterali.

EXT. CHAR. *Feet* with the posterior toes gibbous, and the claws strong. *Tarsi* long. *Bill* strong, at the base tuberculated. *Lamellæ* not continued to the margin. *Nostrils* rather rounded, placed in the middle of the bill. *Wings* armed with one spur. *Tail* squared at the end.

ANAT. CHAR. *Trachea* with the tube of equal size. *Inferior larynx* with a lateral bulb.

ANSER. *Auct.*

CHENALOPEX. *Steph. Antiq.*

TYPE. *Chenalopez Egyptiaca.*

C. J. S. del. et lith.

Printed by J. S. S. S. S.

THE NALOPSE SDY TIAIA & TIAHEA.
Three fifths the Nat size

Affinity. This genus, in possessing a bulb on the trachea, presents an affinity to the ducks; while its form and habit evidently point out its connection with the geese.

SUB FAM. III. ANATINÆ.

EXT. CHAR. *Rostrum* latitudine altitudini æquante. *Pedes* medioeres, *digiti* posteriores nunquam lobati, sed aliquando gibbosi.

ANAT. CHAR. *Sternum* longum, angustatum, posticè latissimum, carinâ anticè productâ et latâ posticè, ad ultimam marginem sterni continuatâ. *Pelvis* lata, brevis. *Os furcatum* medioere lateribus compressis. *Processus coracoides* longus, sed non anticè productus. *Costæ* processibusque posterioribus compressis. *Trachea* tubulâ in omni parte ferè equali. *Larynx inferior* bulbosa ferèque omninò essea in maribus.

EXT. CHAR. *Bill* in breadth and height equal. *Feet* moderate, posterior toes never lobated, but sometimes gibbous.

ANAT. CHAR. *Sternum* long, narrowed, behind broadest, with keel anteriorly produced, broad posteriorly, continued to the margin of the sternum. *Pelvis* broad, short. *Os furcatum* moderate, compressed on the sides. *Coracoid process* long, and not anteriorly produced. *Ribs* and the posterior processes compressed. *Trachea* generally of equal size throughout. *Inferior Larynx* bulbous, and altogether osseous in the males.

TYPE. *Anas*.

HABIT. Aquatic, frequenting fresh water in preference to the sea, feeding generally along the edges of lakes and rivers, on small molluscous animals, in preference to diving for them; soft aquatic plants and weeds also constitute a considerable portion of their food: they often also feed on land. The species, having great power of flight, have a very wide geographical range.

Affinities. To the *Fuligulinx* this sub-family is connected by the musk duck (*Cariua Moschata*), which, besides having a largely developed and slightly lobated hind toe, has the bulb situated at the inferior larynx compressed, and not altogether osseous, as is the case in the rest of the family. The genus *Micropterus* has, on the other hand, a trachea very nearly resembling that of the true land duck. We prefer taking the genus *Anas*, as at present restricted, for the type of this sub-family, instead of *Rhynchaspis*, as proposed by Mr. Swainson, regarding that form and the nearly allied one of *Malacorhynchus*, rather as ultra than typical, and not so well suited to what may be termed the true habit of the sub-family as that we have chosen.

1 *Esophagus preventriculus* and *Stomach* of *Querquedula crecca* Fem.
 2 *Ceca* of *D.* 3 *Ceca* of *Tadorna Belloni* Male.

TAD OF SUELLON
Half the Volume

GENUS I. TADORNA.

EXT. CHAR. *Pedes* digitis posterioribus non lobatis, unguibus robustis armatis. *Tarsi* robusti et longi. *Rostrum* mediocre validum recurvatum. *Lamellæ* vix elongatæ. *Nares* ovales, magnæ, in dimidio basali rostri positæ. *Cauda* rotundata.

ANAT. CHAR. *Trachea* tubula æquali, sed ad larynginem inferiorem paululum expandente et tunc contracta. *Larynx inferior* duabus osseis bulbis utrinque, et anticæ extendentibus.

EXT. CHAR. *Feet* with the posterior toes not lobated, armed with thick claws. *Tarsi* thick and long. *Bill* moderate strong. *Lamellæ* scarcely elongated. *Nostrils* oval, large, placed in the basal half of the bill. *Tail* rounded.

ANAT. CHAR. *Trachea* with the tube of equal size, but a little expanded and again contracted towards the inferior larynx. The *inferior Larynx* with two osseous bulbs extending on each side and before.

ANAS. Linn. Gmel. Lath.

TADORNA. Leach. Flem. Steph.

TYPE. *Tadorna Bellonii*.

The depth of the keel to the length of the sternum is as 1 to 4. The breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. The length of the sternum to that of the tibia as 1 to 2. The breadth of the cranium to its length as 1 to $2\frac{3}{4}$.

Affinities. The common Sheldrake approaches near in external character to the following genus. Nothing being known of the anatomy of that genus, we cannot trace the affinity farther.—To the *Anserinæ* it also approaches in its osteology and general habit, particularly to the genus *Chenalopex*.

GENUS II. CASARKA.

EXT. CHAR. *Rostrum* mediocre lateribus parallelibus non recurvatum. *Nares* subovales, culminem versus positæ. *Lamellæ* elongatæ. *Pedes* robusti, digiti unguibus magnis armati.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Bill* moderate with the sides parallel, not recurved. *Nostrils* suboval, placed near the culmen. *Lamellæ* elongated. *Feet* robust, the toes armed with large claws.

ANAT. CHAR. Unknown.

ANAS. Linn. Lath.

TADORNA. Selby.

CASARKA. Bonap.

TYPE. *Casarka rutila*.

Affinity. This genus may be distinguished from *Tadorna*, to which it is closely allied, by the straighter and much shorter bill.

GENUS III. DENDROCYGNA.

EXT. CHAR. *Pedes* digitis posterioribus longis, non lobatis unguibus longis armatæ. *Tarsi* robusti et longi. *Rostrum* mediocre validum altius-quam latum. *Nares* ovales magnæ prope culmen in medio rostri positæ. *Lamellæ* abbreviatæ. *Cauda* flexilis rotundata. *Alæ* remigibus primariis magnâ incisurâ in medio parte.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes long, not lobated, and with long claws. *Tarsi* robust and long. *Bill* moderate strong, higher than broad. *Nostrils* placed near the culmen in the middle of the bill. *Lamellæ* abbreviated. *Tail* flexible, rounded. *Primaries* with a deep notch in the middle.

ANAT. CHAR. Unknown.

ANAS. Linn.

DENDROCYGNA. Swainson.

TYPE. *Dendrocygna arcuata*.

This genus will be easily recognized by the great length of the tarsi, and at the same time rounded form of the tail. *Anas arcuata*, Hors. or *Anas viduata*, Linn. may be considered as typical species.

Affinities. The genus appears to form a beautiful connecting link between the true fresh-water ducks and the foregoing genus.

GENUS IV. LEPTOTARSIS.

EXT. CHAR. *Rostrum* mediocre ad basin, altius-quam latum, culmine complanatâ et ungue magno instructo. *Nares* parvæ, culmen versus locatæ. *Tarsi* longi et graciles. *Digit*i unguibus paululum incurvatis acutisque armati.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Bill* moderate at the base, higher than broad, the culmen much flattened and with a large nail. *Nostrils* small, placed near the culmen. *Tarsi* long and slender. *Toes* armed with curved and sharp claws.

ANAT. CHAR. Unknown.

LEPTOTARSIS. *Gould*.

The only species of this genus we have seen is in the collection of the Zoological Society, and was received from New South Wales.

Affinity. The structure of the bill appears to ally this genus to that of *Dendrocygna*; while, on the other hand, that of the tail, which, though not elongated, is acuminate, to the genus *Dafila*.

GENUS V. DAFILA.

EXT. CHAR. *Pedes* digitis posterioribus, non lobatis, mediis longissimis. *Tibiæ* paululum denudatæ. *Rostrum* longius capite lateribus parallelibus. *Nares* magnæ ovales, prope culmen ad basin positæ. *Lamellæ* ad marginem rostri continuatæ. *Lingua* subobtusa ad apicem appendiculata lateribus una ordine setarum brevium, supra has setas in medio octo spinis rigidis instructa. *Cauda* longa.

ANAT. CHAR. *Trachea* tubula æquali sed larynginem inferiorem versus paululum contracta. *Larynx inferior* bulbo mediocri uno latere tendente.

EXT. CHAR. *Feet* with the posterior toe not lobed, the middle one longest. *Thighs* a little denuded. *Bill* longer than the head, the sides parallel. *Nostrils* large oval, placed near the culmen at the base. *Lamellæ* continued to the edge of the bill. *Tongue* rather obtuse, at the point with an appendage, the sides furnished with one row of bristles, and above the bristles in the middle with eight stiff spines.

ANAT. CHAR. *Trachea* with the tube equal, but towards the inferior larynx a little contracted. The *inferior Larynx* with a middle-sized bulb on one side.

ANAS. Linn.

DAFILA. Leach. Steph.

TYPE. *Dafila acuta*.

The proportions of the skeleton are: the depth of the keel to the length of the sternum as 1 to 4. Breadth of pelvis to its length as 1 to $1\frac{2}{3}$. Length of sternum to that of the tibia as 1 to $1\frac{7}{8}$. Height of cranium to its length as 1 to 3.

Affinity. The tail in this genus is much elongated, presenting in that particular an affinity to the genus *Mareca*, which has it slightly so.

GENUS VI. PÆCILONITTA.

EXT. CHAR. *Pedes* digitis posterioribus brevibus et parvis, non lobatis. *Rostrum* mediocre, altitudine latitudine æquante. *Nares* prope basi positæ, rotundatæ. *Cauda* medioeris, rotundata.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes short and small, not lobed. *Bill* moderate, the height equal to the breadth. *Nostrils* placed near the base, rounded. *Tail* moderate, rounded.

ANAT CHAR. Unknown.

ANAS. Linn.

MARECA. Steph.

PÆCILONITTA *nobis*

TYPE. *Pæcilonitta Bahamensis*.

We have formed this genus for the reception of *Anas Bahamensis* and its near ally *Anas Erythrorhyncha*.

Affinity. In default of anatomical character, and from general structure, this genus appears to be nearly allied to *Dafila*.

GENUS VII. MARECA.

EXT. CHAR. *Pedes* digitis posterioribus non lobatis mediis longissimis. *Tibiæ* paululum denudatæ. *Rostrum* brevius capite, prope basin latissimum. *Nares* parvæ, ovales. *Lamellæ* ad marginem rostri continuatæ. *Lingua* subobtusa, ad apicem appendiculata latera una ordine setarum brevium et in medio duplice ordine instructis.

ANAT. CHAR. *Trachea* in maribus tubulâ infra glottidem expandente, tunc contractâ, rursus expandente usque ad larynginem. *Larynx inferior* bulbo magno osseo latere et paululum antè tendente.

EXT. CHAR. *Feet* with the posterior toes not lobed, the middle one longest. The *Thighs* a little denuded. *Bill* shorter than the head, near the base broadest. *Nostrils* small, oval. *Lamellæ* continued to the margin of the bill. *Tongue* rather obtuse, with an appendage at the tip.

ANAT. CHAR. *Trachea* with the tube expanding below the glottis, then contracted, again expanded as far as the inferior larynx. *Inferior Larynx* with a large osseous bulb extended on the side and a little before.

ANAS. Linn.

MARECA. Steph.

TYPE. *Mareca fistularis*.

The proportions of the skeleton are:—Height of the keel to the length of the sternum as 1 to $3\frac{1}{3}$. Breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. Length of femur to that of the tibia as 1 to $1\frac{2}{3}$. Height of cranium to its length as 1 to $2\frac{1}{3}$.

Affinity. To the next genus this presents an approach in the formation of the tail, which in some of the species is pointed and slightly elongated. To the following the form of the bill presents an affinity.

GENUS VIII. AIA.

EXT. CHAR. *Pedes* digitis posterioribus non lobatis, mediis longissimis. *Tibiæ* plumatæ. *Rostrum* brevius capite, baso ad apicem angustatum. *Lamellæ* abbreviatæ. *Nares* subovales, parvæ. *Cauda* mediocris.

ANAT. CHAR. *Trachea* tubula equali. *Larynx inferior* bulbosa, ossca, similis precedenti.

EXT. CHAR. *Feet* with the posterior toes not lobated, the middle ones longest. *Thighs* feathered. *Bill* shorter than the head, narrowed from the base to the point. *Lamellæ* abbreviated. *Nostrils* suboval, small. *Tail* moderate.

ANAT. CHAR. *Trachea* with the tube equal. The *inferior larynx* bulbous, osseous like the preceding genus.

ANAS. *Linn.*

AIA. *Boie.*

QUERQUEDULA. *Briss. Steph.*

DENDRONESSA. *Swain.*

TYPE. *Aia sponsa.*

Its habit of sitting and making its nest in trees is curious.

Affinity. Certainly more nearly allied to *Querquedula* and *Mareca* than any other genus of this sub-family.

des. de l'auteur

color. by H. G. S. S. S.

QUERQUEDULA URECA & TRACHEA
Nat. size

GENUS IX. QUERQUEDULA.

EXT. CHAR. *Pedes* digitis posterioribus non lobatis. *Rostrum* lateribus parallelibus. *Lamellæ* paululum elongatæ. *Nares* rotundatæ mediocres. *Cauda* brevis. *Lingua* subobtusa, ad apicem appendiculata, unâ ordine setarum brevium, ad apicem aliquando pene oblitteratæ, supra hæ setæ, in medio, lateraliter, septem spinis instructâ.

ANAT. CHAR. *Trachea* tubulâ infra glottidem expandente, tunc contractâ, rursus expandente, et contractâ. *Larynx inferior* bulbo parvo osseo latere et anticé extendente.

EXT. CHAR. *Feet* with the posterior toes moderate, and not lobated. *Bill* with the sides equal. *Lamellæ* slightly elongated. *Nostrils* rounded, moderate. *Tail* short. *Tongue* rather obtuse, with an appendage at the tip, the sides armed with one rank of short bristles towards the apex very short; above these bristles, in the middle, on each side, armed with seven spines.

ANAT. CHAR. *Trachea* with the tube below the glottis expanding, then contracted, again expanding, and again contracted above the inferior larynx. The *inferior Larynx* with a small osseous bulb extending on one side and before.

ANAS. *Linn.*

QUERQUEDULA. *Ray, Briss.*

TYPE. *Querquedula crecca.*

The proportions of the skeleton are:—Breadth of the keel to the length of the sternum as 1 to $4\frac{1}{5}$. Breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. Length of femur to that of the tibia as 1 to $1\frac{2}{3}$. Height of cranium to its length as 1 to 3.

Affinity. More closely allied to the genera *Mareca* and *Aia* than any other in the sub-family.

GENUS X. CYANOPTERUS.

EXT. CHAR. *Pedes* digitis posterioribus mediocribus, et non lobatis. *Rostrum* latius ad apicem quam ad basin. *Lamellæ* paululum elongatæ. *Nares* rotundatæ, magnæ. *Cauda* vix elongata. *Alæ* ptilis cæruleis.

ANAT. CHAR. *Trachea* tubulâ infra glotidem expandente, tunc contractâ, rursusque expandente, et contractâ supra larynginem inferiorem. *Larynx inferior* bulbo magno osseo lateribus et antice tendente.

EXT. CHAR. *Feet* with the posterior toes moderate, and not lobated. *Bill* broader at the tip than at the base. *Lamellæ* a little elongated. *Nostrils* rounded, large. *Tail* a little elongated.

ANAT. CHAR. *Trachea* with the tube below the glottis expanding, then contracted, and again expanding, and contracted above the inferior larynx. The *inferior Larynx* with a large osseous bulb extending below and on both sides.

ANAS. Linn.

QUERQUEDULA. Briss.

CYANOPTERUS nobis.

TYPE. *Cyanopterus Rafflesii*.

The above genus consists of that group commonly called the blue-winged teal; nor do we know one instance of a bird agreeing with the character which has not bright azure blue wings, except the *Garganey*, which has, however, dull slate-coloured ones. *Anas Rafflesii*, King, may be considered as a good specimen of the form, which may be always distinguished by the peculiar abrupt termination to the upper mandible, which is bent as it were over the lower one, both in front

and on the sides at the apex. The *Garganey* comes nearer to the common teal than the rest of the genus in external form, but in the form of the trachea differs from it. We have never observed a skeleton in any collection except that of the *Garganey*, which we do not consider typical.

Affinities. In the form of the bill and trachea this genus approaches that of *Chauliodus*, but differs from it in the former not having the lamellæ so much elongated, and in the trachea being more enlarged above the inferior larynx. From the following genus it is very distinct in both particulars, but agrees with it in general colouring.

GENUS XI. RHYNCHASPIS.

EXT. CHAR. *Pedes* digitis posterioribus parvis, et non lobatis. *Tibiæ* parte quartâ denudatæ. *Rostrum* longum, ad apicem multum latissimum. *Lamellæ* elongatæ. *Nares* ovaes, prope culmine et baso locatæ. *Lingua* obtusa, ad apicem appendiculata, lateraliter una ordine setarum brevium, et supra has setas posticè novem spinis instructa; et duplici ordine centrali spinarum brevium ornata.

ANAT. CHAR. *Trachea* tubulâ inferiore parte maximâ. *Larynx inferior* bulbo parvo osseo latere.

EXT. CHAR. *Feet* with the hind toe small, and not lobated. *Thighs* with a fourth part naked. *Bill* long, much the broadest at the tip. *Nostrils* oval, placed near the culmen and base. *Tongue* obtuse, with an appendage at the tip, furnished with one row of short bristles on the sides, and above these bristles behind with nine spines; in the middle, above, with a double rank of stout spines.

ANAT. CHAR. *Trachea* with tube largest inferiorly. *Inferior Larynx* with a small bulb extending on one side and behind.

ANAS. *Lin.* Swainson.

RHYNCHASPIS. *Leach MSS.* Steph.

SPATHULEA. *Flem.*

TYPE. *Rhynchaspis clypeata.*

The proportions of the skeleton are:—Depth of the keel to the length of the sternum as 1 to 4. Breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. Length of femur to that of the tibia as 1 to $2\frac{2}{3}$. Height of cranium to its length as 1 to 5.

Affinities. This and the two following, as well as the preceding genera, are very nearly connected, both in form and habit.

GENUS XII. MALACORHYNCHUS.

EXT. CHAR. *Pedes* digitis posterioribus parvis, et non lobatis. *Tibiæ* paululum denudatæ. *Rostrum* longum, ad apicem multum latissimum, et lateribus membranis flexilibus instructis. *Lamellæ* longæ. *Nares* ovales, parvæ, prope basin locatæ.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toe small, and not lobated. *Thighs* little denuded. *Bill* long, at the apex much the broadest, and furnished on the sides with flexible membranes. *Lamellæ* long. *Nostrils* oval, small, placed near the base.

ANAT CHAR. Unknown.

ANAS. *Linn.*

RHYNCHASPIS. *Leach.*

MALACORHYNCHUS. *Swainson.*

TYPE. *Malacorhynchus membrinacea*.

Affinities. No skeleton is, that I am aware of, at present in England; but we should suppose that the bony part of the bill does not extend so far laterally at the point as might be supposed, or, in other words, is not so broad at the apex. Under this supposition, the bird on which Mr. Swainson formed the genus (*Malacorhynchus membrinacea*) forms a beautiful link between the true shovellers and the following genus.

GENUS XIII. CHAULIODES.

EXT. CHAR. *Pedes* digitis posterioribus mediocribus, et non lobatis. *Rostrum* non latius ad apicem quam ad basin. *Lamellæ* paululum elongatæ. *Nares* ovales, magnæ, vix elongatæ. *Lingua* obtusa, ad apicem appendiculata, lateribus unâ ordine setarum brevium, supra has setas laterater centrali parte quinque spinis, et in medio supra duplice ordine setarum brevium instructa.

ANAT. CHAR. *Trachea* tubulâ inferiore parte maximâ, sed supra larynginem inferiorem contractâ. *Larynx inferior* bulbo mediocri osseo latere et multum anticè tendente.

EXT. CHAR. *Feet* with the posterior toes of moderate size, and not lobated. *Bill* not broader at the tip than at the base. *Lamellæ* elongated. *Tongue* obtuse, with an appendage at the tip, the sides furnished with a rank of short bristles, above these in the middle of the sides with five spines, and in the middle of the upper surface with a double rank of short spines.

ANAT. CHAR. *Trachea* with the tube enlarged near the lower extremity, but above the inferior larynx contracted. *Inferior Larynx* with a middle-sized osseous bulb extending on one side and also much anteriorly.

ANAS. Linn. Gmel. Lath.

CHAULIODES. Swainson.

TYPE. *Chauliodus Strepera*.

The proportions of the skeleton are:—Depth of the keel to the length of the sternum as 1 to 5. Breadth of the pelvis to its length as 1 to 1 4-5ths. Length of femur to that of tibia as 1 to $1\frac{2}{3}$. Depth of cranium to its length as 1 to $2\frac{2}{3}$.

Affinity. The elongated lamellæ ally this genus to the two preceding. It appears, indeed, to be precisely intermediate between them and the following.

Fig. 1. Skull of a *Libinia*.

Printed by H. H. H. H. H.

ANAS BOSCHAS & TRACHEA.
I thank the Nat. Soc.

GENUS XIV. ANAS.

EXT. CHAR. *Pedes* digitis posterioribus simplicibus, mediis longissimis. *Tibiæ* parte tertiâ inferiore denudatâ. *Rostrum* lateribus parallelibus. *Nares* ovaies, prope culmen ad basin locatæ. *Lingua* obtusa, ad apicem appendiculata, lateraliter una ordine setarum longarum et flexilum instructa, supra has setas ad latera sex spinis instructa, et in medio suprà duplici ordine setarum brevium.

ANAT. CHAR. *Trachea* tubula equali. *Larynx inferior* unâ bulbâ osseâ latere et antice extendente in maribus.

EXT. CHAR. *Feet* with the posterior toes simple, the middle one longest. *Thighs* with a third part naked. *Bill* with the sides parallel. *Nostrils* oval, placed near the culmen at the base. *Tongue* blunt, at the point with an appendage, the sides furnished with one rank of long and flexible bristles, above these bristles in the middle of the sides furnished with six spines, and in the middle above with a double rank of short bristles.

ANAT. CHAR. *Trachea* with the tube of equal size. *Inferior Larynx* with one osseous bulb on the side and extending anteriorly.

ANAS. *Linn.*

BOSCHAS. *Swainson.*

TYPE. *Anas Boschas.*

We have here retained the old Linnean name for this genus, instead of that of *Boschas* proposed by Mr. Swainson. The proportions of the skeleton are:—The depth of the keel to the length of the sternum as 1 to 5. Breadth of the pelvis to its length as 1 to $1\frac{3}{4}$. Length of femur to that of tibia as 1 to $1\frac{2}{3}$. Depth of cranium to its length as 1 to 3.

Affinity. In size and the form of the tongue this genus approaches nearest to the last of any among the *Anatinæ*.

GENUS XV. CARINA.

EXT. CHAR. *Pedes* digitis posterioribus gibbosis, mediis longissimis. *Tibiæ* plumatæ. *Rostrum* equale, et carunculâ subglobosâ supra nares positâ. *Nares* ovales, magnæ, prope culmine basi locatæ, *Caput* ad oculos denudatum. *Lingua* obtusa, ad apicem appendiculata lateraliter unâ ordine setarum et supra has quatuoribus spinis instructis et in medio supra duplici ordine setarum brevium armata.

ANAT. CHAR. *Trachea* tubulâ equali. *Larynx inferior* unâ bulbâ compressâ latere partem osseâ partem membrinaccâ et postice tendente instructo.

EXT. CHAR. *Feet* with the posterior toes gibbous, the middle longest. *Thighs* feathered. *Bill* with the sides equal, and with a subglobose tubercule placed above. *Nostrils* oval, large, placed at the base of the bill near the culmen. *Head* naked to the eyes. *Tongue* obtuse, with an appendage at the point, the sides armed with one row of bristles, above these with four spines, and in the middle on the upper side armed with a double row of short bristles.

ANAT. CHAR. *Trachea* with the tube equal. The *inferior Larynx* with one compressed bulb, partly osseous, partly membrinaceous, extending on the side and behind.

ANAS. Linn.

CARINA. Flem.

TYPE. *Carina moschata*.

The proportions of the skeleton are:—Depth of the keel to the length of the sternum as 1 to $5\frac{1}{2}$. Breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. Length of femur to that of tibia as 1 to $\frac{2}{3}$. Depth of the cranium to its length as 1 to $2\frac{1}{3}$. The keel of the sternum produced in front. Ischiadic foramina large.

Affinities. To some of the genera in the next sub-family this genus presents an affinity in the form of the hind toe, which, though not strictly lobated, is thickened on its lower side. The trachea presents considerable affinity to both the sub-family we have placed it in, and also to the following. The bulb at the inferior larynx much more osseous on its external aspect than among the *Fuligulinæ*, and less so than among the *Anatinæ*: it is also laterally much compressed, as in the following sub-family.

T. E. del.

$\frac{2}{3}$ Nat. Size.

1 Esophagus Proventriculus and Stomach of *Melanitta nigra* Fem.
 2 Caeca of *D.* 3 Caeca of *Fuligula ferrina* Male 4 Caeca of *Clangula vulgaris* Fem.

SUB FAM. IV. FULIGULINÆ.

EXT. CHAR. *Rostrum* ad basin altius-quam latum. *Pedes* magni, digitis posterioribus lobatis, externis vix brevioribus intermediis. *Rectrices Caudæ* plerumque breves et flexiles. *Alæ* mediocres.

ANAT. CHAR. *Sternum* carinâ parte anteriore vix productâ; posticè non ad marginem sterni continuatâ margine duabus fissuris instructâ his posticè angustatis, intus magnis et ovalibus; nonnunquam autem fissuræ vicè foraminibus multum latioribus posticè quam anticè. *Costæ* longe in abdomine tendentes processibus marginalibus arcuatis vel ad apicem bifidis. *Processus coracoides* anticè vix ultra humeros elongatus. *Os furcatum* arcuatum præcipuè ad extremus partes rami. *Pelvis* latitudine mediocris, elongata. *Cranium* ad occiput altius-quam latum. *Larynx inferior* partim ossea, partim membrinacea.

EXT. CHAR. *Bill* higher than broad at the base. *Feet* large, posterior toe lobated, the outer slightly shorter than the middle one. *Tail feathers* generally short and flexible. *Wings* moderate.

ANAT. CHAR. *Sternum* with the keel slightly produced in front, and not continued to the posterior margin; posterior margin furnished with two fissures narrowed posteriorly, each interiorly large, and oval; or with two foramina on each side, much broader posteriorly than anteriorly. *Ribs* extending far backwards, with the processes on the posterior margin arcuated or bifid at the extremity. *Coracoid process* slightly elongated anteriorly. *Os furcatum* with the rami arched in front, more so above. *Pelvis* of moderate width, long. *Cranium* at the occiput higher than broad.

TYPE. *Fuligula*.

Habit. Aquatic and marine, feeding on small molusca, and occasionally small fish; some of the species possess considerable power of flight.

Affinity. The sterna of some genera of this family, *Clangula* and *Harelda*, are furnished with an ensiform process as in the *Mergansers*. In the former the trachea also much resembles that family. To the *Anatinæ*, as we have before stated, we believe it to be connected through the genus *Micropterus* and *Carina*.

GENUS I. MICROPTERUS.

EXT. CHAR. *Pedes* digitis posterioribus lobatis. *Rostrum* validum, rugosum. *Lamellæ* magnæ. *Nares* in medio rostri positæ. *Tibiæ* plumatæ. *Alæ* breves.

ANAT. CHAR. *Trachea* tubula equali. *Larynx inferior* bulbo laterali osseo.

EXT. CHAR. *Feet* with the posterior toes lobated. *Bill* strong, wrinkled. *Lamellæ* large. *Nostrils* placed in the middle of the bill. *Thighs* feathered. *Wings* short.

ANAT. CHAR. *Trachea* with the tube of equal size. The *inferior Larynx* with an osseous lateral bulb.

ANAS. *Linn.*

MICROPTERUS. *King.*

TYPE. *Micropterus Patachonichus*,

Affinity. The structure of the hind toe and foot unites this genus to the *Fuligulinæ*, while, on the other hand, that of the trachea bears a great resemblance to the *Anatinæ*.

GENUS II. MELANITTA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, et externis longissimis. *Rostrum* magnum, basi elevatum. *Lamellæ* magnæ, ad marginem continuatæ. *Nares* ovales, magnæ, in medio rostri positæ.

ANAT. CHAR. *Trachea* uno vel duabus bulbis tubulâ. *Larynx inferior* non bulbo instructo.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer ones longest. *Bill* large, elevated at the base. *Lamellæ* large, continued to the margin. *Nostrils* oval, large, placed in the middle of the bill.

ANAT. CHAR. *Trachea* with one or two bulbs on the trachea. The *inferior larynx* without a bulb.

MELANITTA. *Boie.*

ANAS. *Linn.*

OIDEMIA. *Flem.*

TYPE. *Melanitta fusca.*

The proportions of the skeleton are:—The depth of the keel to the length of the sternum, as 1 to $4\frac{3}{4}$. Breadth of the pelvis to its length, 1 to $3\frac{1}{5}$. Length of the femur to that of the tibia, as 1 to $1\frac{3}{4}$. Depth of cranium to its length, as 1 to $2\frac{1}{4}$. Keel of the sternum not produced to the posterior extremity.

Affinities. The form of the bill in some of the species of the genus presents an affinity to the genus *Somateria*. In its general form, however, and the structure of the feet, it bears a close resemblance to the foregoing genus,

GENUS III. SOMATERIA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis mediis equalibus. *Rostrum* medioere ad basin elevatum, et ad apicem angustatum. *Lamellæ* non ad apicem continuatæ.

ANAT. CHAR. *Trachea* tubulâ equali. *Larynx inferior* bulbosum osseum, bronchiâ sinistrâ maximâ.

EXT. CHAR. *Feet* with the hind toes lobated, the external ones equal to the middle. *Bill* moderate, with the base elevated, and narrowed towards the tip. The *Lamellæ* not continued to the apex.

ANAT. CHAR. *Trachea* with the tubes of equal size throughout. The *inferior Larynx* bulbous, the left bronchia largest.

ANAS. Linn.

SOMATERIA. Leach.

TYPE. *Somateria mollissima*.

Affinity. In the form of the inferior larynx this genus approaches near the *Anatinae*.

GENUS IV. POLYSTICTA.

EXT. CHAR. *Pedes* magni, *digitis* posterioribus lobatis. *Rostrum* ad basin elevatum, ad apicem angustatum. *Caput* parvum, et cristatum maribus. *Cauda* brevis et acutâ. *Remiges* *tertiae* longæ.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* large, with the posterior toes lobated. *Bill* elevated at the base, narrowed towards the point. *Head* small, and crested in the male. *Tail* short, sharp. *Tertiaries* long.

ANAT. CHAR. *Unknown*.

ANAS. *Linn*.

FULIGULA. *Gould*.

MACROPUS. *Nuttall*.

POLYSTICTA *nobis*.

TYPE. *Polysticta stelleri*.

After we had proposed *Polysticta* as a new generic name for the Western Duck, we found that Nuttall, in his American Ornithology, had also made it into a new genus. Which name has the priority we are uncertain; but that proposed by Mr. Nuttall (*Macropus*) having been used as a generic appellation to a genus of kangaroos, cannot stand. Our own, therefore, is that which we have placed at the head of this genus.

Affinity. The long tertiary feathers and the form of the bill in this genus evidently point out a near affinity to *Somateria*.

GENUS V. KAMPTORHYNCHUS.

EXT. CHAR. *Pedes* magni, digitis externis mediis æquantibus, posterioribus lobatis. *Rostrum* lateribus ad apicem latum, et flexile. *Nares* magnæ, ovales, prope basum locatæ. *Unguis* magna et lata.

ANAT. CHAR. *Trachea* maribus duabus bulbis tubulâ.

EXT. CHAR. *Feet* large, the outer toe equal to the middle one, posterior one lobated. *Bill* broad towards the point at the sides, flexible. *Nostrils* large, oval, placed near the base. *Nail* large and broad.

ANAT. CHAR. *Trachea* in the males with two bulbs on the tube.

ANAS. *Linn.*

FULIGULA. *Ray, Leach.*

KAMPTORHYNCHUS *nobis.*

TYPE. *Kamptorhynchus Labradoræ.*

This genus we have formed on *F. Labradoræ Bonap.* the variety it exhibits being amply sufficient to distinguish it from any other genus. Nothing that I am aware of is known concerning anatomical characters, besides those mentioned in Wilson, and quoted at length under the head *Kamp. Labradoræ.*

Affinities. We have never seen the skeleton of this bird, nor are we aware of one in existence in this kingdom. As far, however, as we are acquainted with it, from Wilson's account, its true place appears to be somewhere near that we have assigned to it.

GENUS VI. CALLICHEN.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, intermediis vix longissimis. *Tibiæ* plumatæ. *Rostrum* capite equalè, prope basum latissimum. *Lamellæ* validæ, et ad marginem continuatæ. *Nares* subovales, magnæ. *Cauda* mediocris. *Caput* cristatum.

ANAT. CHAR. *Trachea* duobus bulbis tubula. *Larynx inferior* bulbo laterali, partim membrinaceo, partim osseo, latere complanato.

EXT. CHAR. *Feet* with the posterior toes lobated, the middle ones scarcely the longest. *Thighs* feathered. *Bill* equal to the head, broadest at the base. *Lamellæ* strong, and continued to the margin. *Nostrils* suboval, large, *Tail* moderate. *Head* crested.

ANAT. CHAR. *Trachea* with two bulbs on the tube. *Inferior Larynx* with a lateral bulb, partly membrinaceous, partly osseous, on the side flattened.

ANAS. *Linn.*

FULIGULA. *Ray, Leach.*

GLAUCHUM, QUERQUEDULA. *Briss.*

CALLICHEN. *Brehm.*

MERGOIDES *nobis.*

TYPE. *Callichen rufina.*

We first instituted this genus on *F. rufina*, after having observed the trachea of that species. To this, however, will be added the beautiful species *Fuligula caryophylacea*. With any anatomical character further than that above given we are unacquainted, not having ever seen the skeleton.

Affinities. In having bulbs on the tube of the trachea, and also in general form, this bird presents considerable affinity to the *Mergansers*.

GENUS VII. FULIGULA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis inter mediis equantibus. *Tibiæ* plumatæ. *Rostrum* prope apicem latissimum, obtusum. *Lamellæ* abbreviatæ et non ad marginem continuatæ. *Nares* subovales, parvissimæ. *Cauda* brevissima. *Lingua* obtusa ad apicem appendiculata, lateribus unâ ordine setarum flexilum instructis; in medio ad latera supra has setas quatuoribus spinis instructis.

ANAT. CHAR. *Trachea* tubulâ inferiore parte contractâ. *Larynx inferior* bulbo laterali, partim membrinaceo, partim osseo, latere complanato.

EXT. CHAR. *Feet* with the hind toes lobated, the outer ones scarcely equal to the middle. *Thighs* feathered, *Bill* broadest at the point, blunt. *Lamellæ* abbreviated, and not continued to the margin. *Nostrils* suboval, very small. *Tail* very short. *Tongue* obtuse, with an appendage at the tip, and the sides furnished with one row of flexible bristles; in the middle with the sides above the bristles furnished with four spines.

ANAT. CHAR. *Trachea* with inferior portion of the tube contracted. The *inferior Larynx* with one lateral bulb, partly membrinaceous, partly osseous, on the sides compressed.

ANAS. Linn.

FULIGULA. Ray, Leach.

NYROCA. Fleming.

GLAUCIUM, QUERQUEDULA. Briss.

TYPE. *Fuligula Marilla*.

We here again assume the name to this genus given to it by Ray, as being the oldest. The proportions of the skeleton are:—Depth of the keel to the whole length of the sternum as 1 to 4. Breadth of the pelvis to its length as 1 to $1\frac{3}{4}$. Length of femur to that of tibia as 1 to $1\frac{7}{8}$. Depth of cranium to its length as 1 to $2\frac{3}{4}$. Sternum with the keel produced anteriorly. Ribs projecting far backwards.

Affinities. To the following genus this bears a near affinity in the external form and that of the trachea and skeleton, as well as to the foregoing.

NYROCA LEUCOPTHALMUS & TRACHEA.

in the Act. Soc.

GENUS VIII. NYROCA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis mediis vix æquantibus. *Tibiæ* plumatæ. *Rostrum* brevius capite, prope apicem latissimum, obtusum. *Lamellæ* abbreviatæ, et non ad marginem continuatæ. *Nares* subovales, parvæ. *Cauda* brevis. *Lingua* obtusa, ad apicem appendiculata, lateribus unâ et in medio duplici ordine setarum flexilium instructis, ad latera supra has setas quinque spinis instructis.

ANAT. CHAR. *Trachea* tubulâ in medio multum expandente, contractâ supra larynginem inferiorem. *Larynx inferior* partim ossea, partim membrinacea.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer ones scarcely equal to the middle. *Tibia* feathered. *Bill* shorter than the head, near the point broadest, obtuse. *Lamellæ* abbreviated, and not continued to the margin. *Nostrils* suboval, small. *Tail* short. *Tongue* obtuse, with an appendage at the point, the sides with one and the middle with a double rank of flexible bristles, above these bristles on the sides armed with five strong spines.

ANAT. CHAR. *Trachea* with the tube much expanded in the middle, again contracted above the inferior larynx. The *inferior Larynx* partly osseous, partly membrinaceous.

ANAS. *Linn.*

NYROCA. *Flem.*

TYPE. *Nyroca leucophthalmus*.

Dr. Fleming included the Scaup, Tufted Duck, and Pochards in his genus *Nyroca*. It is, however, now restricted to the Pochards which want the crested head.

The proportions of the skeleton are:—The depth of the keel to the whole length of the sternum, as 1 to 5. Breadth of the pelvis to its length, as 1 to 2. Length of femur to that of tibia, as 1 to $1\frac{2}{3}$. Depth of the cranium to its length, as 1 to $2\frac{2}{3}$. The keel of the sternum much produced anteriorly.

Affinities. This genus, as before stated, presents an affinity to the last in general form, and also in that of the inferior larynx.

Osseous skeleton

Trachea by G. H. M. M. M. M. M.

HARELDA GLACIALIS & TRACHEA.

2 This is the Nat Size

GENUS IX. HARELDA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis mediis vix equantibus. *Tibiæ* plumatæ. *Rostrum* mediocre, baso ad apicem angustatum. *Lamellæ* parte omnino oblitteratæ. *Nares* lineares, versus media rostri positæ. *Cauda* maribus elongata.

ANAT. CHAR. *Trachea* tubulâ inferiore parte tuberculosâ a posteriore membranâ tectâ. *Larynx inferior* bulbosa, partim ossea, partim membrinacea.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer one scarcely equal to the middle. *Thighs* feathered. *Bill* moderate, narrowed from the base to the apex. *Lamellæ* altogether obliterated at the points. *Nostrils* linear, situated about the middle of the Bill. *Tail* in the male elongated.

ANAT. CHAR. *Trachea* on the lower part posteriorly covered with a membrane. The *inferior Larynx* bulbous, partly osseous, partly membrinaceous.

HARELDA. Ray. Leach.

ANAS. Linn.

QUERQUEDULA. Briss.

CLANGULA. Flem.

TYPE. *Harelda Glacialis*.

This genus having been instituted by Ray, requires no alteration. The depth of the keel to the whole length of the sternum is as 1 to 4. The length of the femur to that of the tibia as 1 to $1\frac{2}{3}$. The height of the head to its length as 1 to $2\frac{5}{8}$. There is a small ensiform process attached to the posterior margin of the sternum.

Affinity. The sternum in this genus, as well as in the next, being furnished with an ensiform process, evidently points out an affinity between them. The trachea, however, differs most materially, and the elongated tail is unlike any other genus of the family: still, that this is its true place no doubt can remain, the ribs projecting far backward over the abdomen, and the hind toe being largely lobated.

Bohn del et litho

Scale 1/2 inch

1. SKELETON OF CLANGULA HISTRIONICA.
2. TRACHEA OF CLANGULA VULGARIS.

Booth's Nat. Hist.

GENUS X. CLANGULA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis mediis vix æquantibus. *Tibiæ* plumosæ. *Rostrum* parvum, capite brevius, a basi ad apicem angustatum. *Nares* ovales, medium versus rostri positæ. *Lamellæ* ad apicem pene oblitteratæ. *Cauda* rectricibus rigidis, sed ad apicem flexilibus. *Lingua* obtusa, ad apicem appendiculata, lateribus ordine uno setarum flexilium instructis.

ANAT. CHAR. *Trachea* tubulâ bulbo uno instructâ. *Larynx inferior* bulbo magno partim membrinaceo, partim osseo. *Costæ* a posticè longe tendentes. *Sternum* posticè appendicæ ensiformi parvo.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer ones scarcely equal to the middle. *Tibiæ* feathered. *Bill* small, shorter than the head, narrowed from the base to the point. *Nostrils* oval, placed towards the middle of the bill. *Lamellæ* almost obliterated at the top. *Tail* with the quills stiff, but flexible at the points. *Tongue* obtuse, with an appendage at the point, the sides garnished with one row of flexible bristles.

ANAT. CHAR. *Trachea* with one bulb on the tube. The *inferior Larynx* with a large bulb, partly membrinaceous, partly osseous. *Ribs* extending far backwards. *Sternum* behind with a small ensiform appendage.

ANAS. *Pars. Linn.*

CLANGULA. *Leach, Flem.*

QUERQUEDULA. *Briss.*

TYPE. *Clangula vulgaris.*

This genus was instituted by Dr. Leach. Brisson included the birds contained in it in the genus *Querquedula*, which is now applied solely to a genus of fresh-water ducks (*Anatidæ*). The depth of the keel to the length of the sternum in this genus is as 1 to $4\frac{1}{2}$. Breadth of the pelvis to its length as 1 to $1\frac{1}{2}$. Length of femur to that of tibia as 1 to $1\frac{1}{2}$. Depth of cranium to its length as 1 to $2\frac{1}{8}$.

Affinity. The disposition to rigidity, and the form of the tail, as well as that of the bill and feet, evidently point out an affinity to the next family; while the skeleton approaches to that of the genus *Mergus*, both in possessing an ensiform appendage to the sternum, and in the form of the inferior larynx.

SUB FAM. V. ERISMATURINÆ.

EXT. CHAR. *Rostrum* ad basin altius-quam latum. *Pedes* magni, digitis posterioribus lobatis, externis et intermediis inter se æqualibus. *Rectrices Caudæ* longæ et validæ. *Alæ* breves.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Bill* higher than broad. *Feet* large, posterior toe lobated, the outer toe equal to the middle. *Tail feathers* long and stiff.

ANAT. CHAR. Unknown.

TYPE. *Erismatura*.

Habit. Like the last, truly aquatic.

Affinity. The shortness of the wing deprives these birds of any great power of flight, consequently the species have not a wide geographical range. We are not aware that the use of the stiff tail feathers found on birds of this family and the cormorants has ever been ascertained, and which character appears to separate them so distinctly from the other families of ducks: they are certainly of great use to them when on shore and stationary, their position being necessarily upright, the legs being placed far backwards. We should also suppose them to be connected with their motion in the water while diving, perhaps giving them the power of raising or lowering themselves suddenly. If it should be eventually ascertained that fish constitute any considerable portion of their food, this power must be of great use to them, as well as to the cormorants, in following their prey through the water.

GENUS I. THALASSORNIS.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis inter-mediis vix æquantibus. *Tibiæ* plumosæ. *Rostrum* validum, latum; lateribus parallelis; et ungue magnâ armatum. *Nares* parvæ, ovales, in medio rostri culminem versus positæ. *Cauda* rotundata, rectricibus brevibus et subrigidis instructa. *Alæ* breves, remigibus ad apices incurvatis.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer ones scarcely equal to the middle. *Thighs* feathered. *Bill* strong, broad; the sides parallel, and armed with a large nail. *Nostrils* small, oval, placed in the middle of the bill towards the culmen. *Tail* rounded, furnished with short and slightly rigid feathers. *Wings* short, with the quills incurved at the apex.

ANAT. CHAR. Unknown.

THALASSORNIS *nobis*.

TYPE. *Thalassornis leuconotus*.

Affinity. The above genus appears to form a connecting link between the genus *Clangula* and *Biziura*, the structure of the tail being nearly that of the former genus, while the bill is that of the latter.

GENUS II. BIZIURA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis inter mediis æquantibus. *Tibiæ* plumosæ. *Rostrum* validum, latum, ad apicem angustatum. *Mandibulæ* ambæ unguiculatæ, inferior in maribus paleâ magnâ pendente instructa. *Nares* rotundatæ, in medio rostri, culminem versus positæ. *Cauda* rotundata, repticibus rigidis et acutis instructa. *Alæ* remigibus, ad apices incurvatis.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes lobated, the external ones equal to the middle. *Thighs* feathered. *Bill* strong, broad, narrowed towards the apex. Both *mandibles* furnished with nails, the inferior one in the males with a large hanging wattle. *Nostrils* rounded, placed in the middle of the bill towards the culmen. *Tail* rounded, furnished with stiff and acute quills. *Wings* with the quills at the points incurved.

ANAT. CHAR. Unknown.

BIZIURA. *Steph. Leach.*

TYPE. *Biziura lobata.*

Mons. Temminck included this genus, together with all the other aquatic ducks, in one large genus *Hydrobates*. The name of *Biziura* was first given to it, according to Stephens, in Shaw's Zoology, by Leach, in his MSS. and was afterwards published by Mr. Stephens in the above work. It is much to be regretted that no skeleton or parts to illustrate the anatomy of this extraordinary bird have hitherto been brought to England.

Affinity. It bears considerable resemblance to the other genera of the family, but differs from all in having the lobe under the chin.

GENUS III. ERISMATURA.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, externis mediis vix æquantibus. *Tibie* plumatæ. *Rostrum* validum, aut mediocre, prope apicem latissimum. *Mandibulæ* unguiculatæ. *Nares* ovales, parvæ, in medio rostri versus culminem positæ. *Cauda* rotundata, rectricibus rigidis. *Alæ* remigibus paululum incurvatis.

ANAT. CHAR. Ignoti.

EXT. CHAR. *Feet* with the posterior toes lobated, the outer one scarcely equal to the inner. *Thighs* feathered. *Bill* moderate, broadest near the tip. *Mandibles* furnished with nails. *Nostrils* oval, small, placed in the middle of the bill towards the culmen. *Tail* rounded, with the quills stiff. *Wings* with the quills a little incurved.

ANAT. CHAR. Unknown.

ANAS. *Lin.*

OXYURA. *Bonap.*

ERISMATURA. *Bonap.*

UNDINA. *Gould.*

GYMNURA. *Nuttall.*

TYPE. *Erismatura rubida.*

The genus *Oxyura* was first instituted by the Prince of Musignano, for the reception of *Anas rubida*, Wils. Mr. Gorler having overlooked it, formed his genus *Undina*, or *Anas Leucocephala*, Linn. a bird of the same conformation. Mr. Nuttall having discovered that the name of *Oxyura* had been before used for a genus of creepers, substituted that of *Gymnura* for it. The Prince of Musignano had, however, in the appendix to the same work in which he named the genus *Oxyura*, also called it *Erismatura*, which name will stand, that of *Oxyura* having been before employed for another genus.

Affinity. Some of the species in this genus appear to present a slight approach to the next sub-family, the tail being diminished in length and rigidity, and also in the wings not being so much curved inwards as in the last.

Fig. 87.

1. Esophagus proventriculus and Stomach of *Mercurius Serrator* fem.
2. Caeca et D^o

SUB FAM. VI. MERGINÆ.

EXT. CHAR. *Rostrum* mediocre, ad apicem angustatum acuminatum et dentibus ad latera instructum. *Digit*i posteriores lobati.

ANAT. CHAR. *Sternum* carinâ parte anteriore valde angustatâ et non ad marginem continuatâ, margine posteriore appendice ensiformi instructo, et duobus foraminibus ad latera. *Processus coracoides* antice elongatus. *Os furcatum* arcuatum, rami lateraliter compressi. *Trachea* uno vel duobus bulbis tubulâ. *Larynx inferior* in maribus bulbo osseo instructa. *Œsophagus* dilateri potest. *Gigeria* interne magna cavitione.

EXT. CHAR. *Bill* moderate, narrowed to the point, acuminate, and at the sides armed with teeth. Posterior *Toes* lobated.

ANAT. CHAR. *Sternum* with the keel anteriorly produced, posteriorly narrowed, and not continued to the margin, furnished with an ensiform process and two lateral foramina. *Coracoid process* elongated anteriorly. *Os furcatum* arched, the branches laterally compressed. *Trachea* with one or two enlargements on the tube. The *inferior Larynx* in the males furnished with an osseous bulb. *Œsophagus* dilatable. The *Gizzard* internally with a large cavity.

TYPE. *Mergus*.

Affinities. Having, under the head *Lamellirostres*, stated some of the affinities which exist between the family generally and others, we have merely, under the head sub-families, stated those which hold between each contained in the above family. In both external and internal character the strongest relation exists between the above and that of *Fuligulinæ*, the genera *Clangula* and *Harelda* in particular, both of which possess the ensiform process on the posterior margin of the sternum, and the latter a bulb on the tube of the trachea, formed by enlarged rings. The genus *Callichen* possesses two bulbs, but the skeleton does not approach so nearly as that of the two genera before mentioned. To the *Erismaturinæ* we suspect the affinity is not so great, certainly not so far as relates to external character, although we have placed them contiguous, there being evidently a double point of affinity to the Divers.

Habit. The habit of this sub-family is truly aquatic, its length of body, and legs being placed far backwards, give the birds contained in it great power of diving. Its food consists principally of small fish.

GENUS I. MERGUS.

EXT. CHAR. *Pedes* digitis posterioribus lobatis, mediis vix longissimis. *Tibiæ* plumatæ. *Rostrum* a baso ad apicem angustatum, lateribus serratis, apice unguiculato. *Nares* subovales, medioeres, versus medium rostri positæ. *Lingua* acuminata, lateribus ordine uno, parte centrali duplici ordine setarum reflexarum instructa.

ANAT. CHAR. *Trachea* uno vel duobus bulbis tubulâ in maribus instructa. *Larynx inferior* bulbo magno, parte membrinaceo, parte osseo. *Costæ* postice longè tendentes.

EXT. CHAR. *Feet* with the posterior toes lobated, the middle ones scarcely the longest. *Thighs* feathered. *Bill* narrowed from the base to the point, the apex with a nail. *Nostrils* oval, moderate, placed towards the middle of the bill. *Tongue* pointed, the sides armed with one and the middle with two rows of bristles.

ANAT. CHAR. *Trachea* with one or two bulbs on the tube in the males. The *inferior Larynx* with a large bulb, partly osseous, partly membrinaceous. *Ribs* extending far backwards.

MERGUS. Ray, Linn. Gmel. Briss.

MERGANSER. Briss.

TYPE. *Mergus serrator*.

PLATE 10. FIG. 1.

Drawn by H. J. G. S. 1861

STUDY OF SERPENT & TRAUMA
Two things the Art de

Mr. Stephens, in vol. XII. p. 156, of Shaw's Zoology, divides this genus, separating from it the Smew (*Mergus albellus*), and placing the other Mergansers under the genus *Merganser*, *Briss.* The depth of the keel to the length of the sternum in this genus is as 1 to 5. Breadth of the pelvis to its length as 1 to $2\frac{1}{4}$. Length of the cranium to its depth as 1 to $3\frac{1}{2}$. Length of the femur to the length of the tibia as 1 to $1\frac{3}{4}$.

We have spoken before of the affinities of this genus under the head of the sub-family *Merginæ*.

SYNOPSIS SPECIERUM.

CHORISTOPUS SEMIPALMATUS.

Cho.—Capite colloque nigro, dorso superiore albo.

SEMIPALMATED GOOSE.

CHO. with the head and neck black, the upper part of the back white.

	FT.	IN.		IN.
<i>Length</i>	3	4	<i>Tarsi</i>	4
<i>Bill</i>		4	<i>Middle toe</i>	4½

CHO.: with the upper part of the back, rump, and below, with the exception of the tibiæ, white; the remainder and the tibiæ glossy black; legs and bill orange; the nail of the latter horn colour.

Anas semipalmata *Linn. Trans.* iv. 103.

Lath. Ind. Orn. App. 69.

Semipalmated Goose..... *Lath. Gen. Syn. Sup.* 2, 347.

Anas melanoleuca *Lath. Ind. Orn. App.* 69.

Trachea *Lin. Trans. col.* xv. p. 383, *pl.* xiii, xiv.

Inhabits New Holland, and utters a shrill whistling note.

Mr. Yarrell, in vol. 15, p. 383, gives the following anatomical peculiarities of this bird. The trachea "was situated on the outside of the left pectoral muscle, under the skin, sufficiently raised under the wing that respiration would not be impeded when the bird rested with its breast on the ground, the parallel tubes being firmly attached both to the muscle and the skin by cellular tissue. The clavicle on the right side of the bird is of the usual character, but that on the left is both shorter and wider, having an aperture about the middle, the sides diverging, with a projecting point on the inner side, to which the tube of the trachea is firmly attached about two inches above the bone of divarication. The trachea lying on the left side of the bird, the lower portion of the tube, in its passage to the lungs, crosses the left branch of the fureula at a right angle, but becoming attached to this projection of the clavicle, receives from the point described its central direction into the body. The whole length of the windpipe in this bird is four feet eight inches." In a young bird dissected by Mr. Yarrell, the trachea was not nearly so much convoluted.

PLECTROPTERUS GAMBENSIS.

Plec.—Ater; subtus albus.

GAMBA GOOSE.

PLEC. black; below white.

	IN.		IN.		IN.
Length.....	39½	Naked part of the bill....	5	Tarsi.....	5

PLEC.: smaller wing coverts, belly, neck, and thighs white; the remainder deep purplish black; spur on the wing of the male one inch long; male with a large knob on the culmen, at the base of the bill.

Female smaller than the male, and with the knob at the culmen of the bill smaller.

<i>Plectropterus Gambensis</i>	<i>Steph. Shaw Zool.</i> 12, 7.
<i>Anas Gambenses</i>	<i>Linn. Syst. Nat.</i> 1, 195.
	<i>Gmel. Syst. Nat.</i> 1, 503.
	<i>Lath. Ind. Orn.</i> 1, 839.
	<i>Shaw, Mus. Ser. pl.</i> p. 231.

An African species, inhabiting the Gamba.

The following are some notes we took on the dissection of a female specimen. The length of the intestinal, from gizzard to anus, 6 feet 4 inches; from cæca to anus 4 inches; the length of cæca 5¾ inches. Trachea acted upon by two pairs of muscles; first pair the usual sterno tracheal ones; second pair diverging from the trachea higher than the first, and inserted in the coracoids. Tube of large calibre, contracted slightly just above the inferior larynx, which resembles in form that of the true geese. The bronchiæ of equal diameter. Œsophagus placed on the right side of the neck, and much dilated in the middle. The sternum long and narrow.

The above account appears to bear a considerable resemblance to that of the *Cereopsis* of Mr. Yarrell.

N. B. Our cut of the bill and foot illustrating the genus was taken from a female specimen.

CEREOPSIS NOVA HOLLANDIÆ,

Cer.—*Cinerius tectricibus alarum atribus maculis ornatis.*

NEW HOLLAND CEREOPSIS.

CER. blue grey, spotted on the wing coverts with black.

	IN.		IN.
<i>Length</i>	34	<i>Middle toe</i>	$3\frac{1}{4}$
<i>Tarsi</i>	4	<i>Outer</i>	$3-2\frac{3}{4}$
		<i>Hind toe</i>	$1\frac{1}{8}$

CER.: bluish grey, with the tail, under and upper tail coverts, and a few spots on the scapulars, black; bill black; cere covering the nostrils greenish yellow; webs scolloped, as in the Sandwich Island goose; nostrils large and round; legs orange; feet black.

Cereopsis Nova Hollandiæ..... *Lath. Ind. Orn. Sup.* 67.

Anser Griseus..... *Vieill. 2 edit. Dic. Hist. Nat.* 23—338.

Cereopsis cendrée *Temm. Pl. col.* 206.

Now found in most of our large museums; has also been brought over alive. Appears to dread the water, and feed chiefly on grass. Inhabits New Holland.

In the Zoological Proceedings for January 25, 1831, we find the following: Mr. Yarrell stated, that having examined the body of the bird, he had remarked that its trunk was much shorter than that of the true geese, and more triangular in its shape; the pectoral muscles were large and dark coloured; the trachea was of large and nearly uniform calibre, without convolution, and attached in its descent to the right side of the neck, as in the heron and bittern; in the form of its bone of divarication and bronchiæ it most resembled the same part in the geese. The muscles of voice were two pairs, one pair attached to the shafts of the os furcatorium, the other to the inner lateral surface of the sternum. The lobes of the liver were of large size, morbidly dark in colour; their substance broke down under the finger on the slightest pressure. The stomach, a true gizzard, was of small size as compared with the bulk of the bird. The first duplicature of intestine was six inches in length, at the returning portion of which the biliary and pancreatic ducts entered; from thence to the origin of the cæca four feet six inches; the cæca nine inches each; the colon and rectum together five inches: the whole length of the intestines was seven feet five inches. The stomach and intestinal viscera were loaded with fat; the other parts exhibited nothing remarkable. Internally this bird, which was a male, resembled the true geese, but externally, in the character of the bones, particularly in the rounded form of the edge, and great depth of the keel of the sternum, and the lateral situation of the trachea in reference to the cervical vertebræ, it was decidedly similar to the *Ardeidæ*.

CHLOEPHAGA SANDVICENSIS.

Chlo.—Brunnea, vertice, genis, colloque parte posteriore, atris.

SANDWICH ISLAND GOOSE.

CHLO. brown, with the crown, cheeks, and line down the back of the neck, black.

	IN.		IN.
<i>Length</i>	23	<i>Middle toe</i>	$3\frac{1}{4}$
<i>Tarsi</i>	$3\frac{1}{2}$	<i>Outer</i>	$2\frac{7}{8}$
<i>Bill</i>	$1\frac{1}{2}$	<i>Inner</i>	$2\frac{3}{8}$
<i>Hind toe</i>	$\frac{5}{8}$		

CHLO.: crown, cheeks, quills, and a line down the back of the neck black; remainder of the neck bright brownish grey; quills, tail, and upper tail coverts white; remainder ashy grey, darker on the back, with each feather tipped with brown or grey; ring round the neck brown; wings black, the edges of the feathers lighter; legs, feet, and bill black; webs deeply scalloped.

Bernicla Sandvicensis *Vig. Zool. Pro. May 27, 1834.*

Inhabits the Sandwich Islands.

An interesting account of the habits of this bird, while confined in Lord Derby's menagerie, by that nobleman, will be found in the Zoological Proceedings as above quoted, when he states they laid four eggs of a white colour, and large in proportion to the size of the bird, two of which were hatched, and one reared; since which time also another hatch has been reared by the Zoological Society.

CHLOEPHAGA MAGELLANICA.

Chlo. Mas.—Albus, dorso subtusque atro transversum strigato ; rostro, pedibus, tarsisque nigris.

Chlo. Fem.—Ferruginea; subtus albus atro strigato; colloque canescente; tarsi rubris; pedibus nigris.

UPLAND MAGELLANIC GOOSE.

CHLO. Mas. white, with the back and under surface transversely barred with black; the bill, feet, and legs black.

CHLO. Fem. ferruginous; below white barred with black; neck hoary; legs red; the feet black.

	IN.		IN.
Male: Length.....	26	Inner toe.....	$1\frac{5}{8}$
Bill.....	$1\frac{1}{4}$	Middle.....	$2\frac{1}{2}$
Tarsi.....	3	Outer toe.....	$2\frac{1}{8}$

Head, neck, under surface, and back white, the two latter barred with black; wing coverts and secondaries white; tertiaries and bastard wings blue black, tipped with white; bill and legs black, the latter approaching to orange in dried specimens.

	IN.		IN.
Fem. Length.....	25	Middle toe.....	$2\frac{1}{2}$
Bill.....	$6\frac{3}{4}$	Outer.....	2
Tarsi.....	$1\frac{1}{8}$	Inner.....	$1\frac{5}{8}$

Lower part of the neck, upper part of the back, and breast light ferruginous, barred with black; head and neck grey or hoary brown; quills, tail, upper tail coverts, and lower part of the back black, glossed with green inflections; tertiaries long, grey; secondaries white, behind which is a bar of black; wing spot (formed by the secondary coverts) metallic green; wing coverts, belly, and vent white; flanks white barred with black; bill black; legs orange, with the inside black.

Young: the young birds of both sexes have the head and upper part of the neck brownish ash; below greyish white lineated with dark brown; upper part of the back grey barred with black; rump and scapulars deep ash; thighs white barred with light brown; vent brownish black; middle of the belly and wing coverts white; bill as in the adults; legs greenish black.

<i>Bernicla leucoptera</i>	<i>Less. Traite d' Orn.</i> 627.
<i>Anas</i>	<i>Gmel. Syst.</i> 1. 505.
	<i>Lath. Ind. Orn.</i> 2. 835.
<i>Bernicla</i>	<i>Steph. Sh. Zool.</i> 12. 58.
<i>Anas Magellanica</i>	<i>Gmel. Syst. Nat.</i> 1. 505.
	<i>Lath. Ind. Orn.</i> 2. 836.
<i>Bernicla</i>	<i>Steph. Sh. Zool.</i> 12. 61.
<i>Anas Picta</i>	<i>Gmel. Syst.</i> 1. 504.
	<i>Lath. Ind. Orn.</i> 2. 836.

holocephala?

<i>Bernicla</i>	<i>Steph. Sh. Zool.</i> 12. 60.
<i>White-winged Antarctic Goose</i>	<i>Brown. Illust. Zool.</i> pl. 40.
<i>Bastard Goose</i>	<i>Lath. Syn.</i> 6. 440.
<i>Magellanica Goose</i>	<i>Lath. Syn.</i> 6. 443.
<i>Painted Goose</i>	<i>Lath. Syn.</i> 6. 443.
<i>Oie des iles Malouines</i>	<i>Buff. Ois.</i> ix. 69.
	<i>Less. Traite Orn.</i> 627.
<i>L' Oie des terres Magellaniques</i>	<i>Buff. Ois.</i> ix. p. 58, pl. En. 1006.

Inhabits the southern extremity of America and the Falkland Isles; found generally in flights, according to Lesson; though we are informed by Charles Darwin, Esq. that it is generally found in pairs.

The changes which the plumage of the female of this species undergoes are very extraordinary. With those of the male we are at present ignorant, as to whether it assumes the white state at once or not. The young female, which is the *Anas Picta* of authors, first in its approach to the adult state, has the black bars on the back and neck, which become ferruginous; those on the flanks become darker, and the intermediate spaces white; the head also becomes ashy grey, and the legs approach to deep orange. Before the time of incubation black bars again appear on the neck, breast, and the upper part of the back, and the ferruginous colouring becomes lighter; the vent and under tail coverts are in very old birds white; in younger ones ferruginous, the legs become of a brighter orange, and the head and neck become brown.

Such are what we believe to be the changes of this bird: for material assistance in its investigation we are indebted to Charles Darwin, Esq. and also for the trachea from which our figure is taken. The inferior larynx is furnished with a hollow rounded bulb, very similar to that of the common widgeon, but differing in its form, being flattened superiorly, and of a much stronger structure than is usual among the wading ducks. It may at once be distinguished from all the other tracheæ with which we are acquainted, by an indented line on the lower surface of the bulb.

(*Vide our plate.*)

BERNICLA ANTARCTICA.

Ber. Mas.—Albus, remigibus primariis apicibus nigris.

Ber. Fem.—Nigra, dorso subtusque albo transversim strigato; occipite brunneo: rostro aurantiaco.

ANTARCTIC GOOSE.

BER. Mas. white, with the ends of the primary quills black.

BER. Fem. black, with the back and beneath transversely striped with white; the occiput brown; the bill orange.

		IN.			IN.
Male:	<i>Length</i>	25		<i>Tarsi</i>	2 $\frac{1}{2}$
	<i>Bill</i>	1 7-12ths			

Entirely white except the apical half of the primaries, which are ashy black. The back of the head in young birds is tinged with cinereous; legs orange; bill (in the preserved specimens we have seen) black, with an orange patch over the nostrils: but Azara says it is orange.

		IN.			IN.
Fem.:	<i>Length</i>	26		<i>Inner toe</i>	2
	<i>Bill</i>	1 $\frac{1}{2}$		<i>Middle</i>	3 $\frac{1}{8}$
	<i>Tarsi</i>	2 $\frac{1}{2}$		<i>Outer</i>	3 $\frac{3}{4}$

Neck, breast, and belly black barred with white, the bars minute towards the head; back part of the head umber brown; scapulars and primaries very dark brown, sometimes black; smaller wing coverts and secondaries white; tertiaries blue green; tail, lower part of the back, under tail coverts, and thighs, white; bill and legs orange; toes and webs black in dried specimens.

<i>Anas Antarctica</i>	<i>Gmel. Syst.</i> 1. 505.
	<i>Lath. Ind. Orn.</i> 2. 835.
<i>Bernicla</i>	<i>Steph. Sh. Zool.</i> 12. 59.
<i>Anser candidus</i>	<i>Feill. Nouv. Dict. Hist. Nat.</i> 23. 331.
	<i>En Method.</i> 3. 351.
	<i>Steph. Sh. Zool.</i> 12. 38.
<i>Ganso Blanco</i>	<i>Azara Parax del Paraguay,</i> 3. 406.
<i>Die anarctque, fem.</i>	<i>Voy. de la Coquille,</i> pl. 50.
<i>Antarctic Goose</i>	<i>Lath. Gen. Syn.</i> 6. 442.

The male of this species appears to have been first discovered by Azara. A good plate of the female will be found in the "Voyage de la Coquille." It inhabits South America and the adjoining islands, frequenting the sea coast and the lowlands adjoining, from whence it is called by Mr. Darwin, in his amusing Journal, the Lowland Goose. For the trachea from which our figure is taken we are indebted to the above-named friend. The bulb of the inferior larynx is not flattened above as in the last bird, but altogether bears considerable resemblance to it and the wading ducks, from which latter, in common with all other geese, it may be distinguished by the rings of the trachea being more massive, and also by the form of the tube, which is always contracted towards the inferior larynx, while in the wading ducks it always expands in that part.

The flesh is not good food. Skins of both sexes are in the museum of the Zoological Society, brought home by Capt. King.

BERNICLA RUFICOLLIS.

Ber.—Vertice colloque posteriore atris; genis rufis.

RED-NECKED GOOSE.

BER. with the crown and neck posteriorly black; the cheeks red.

Length 21 inches.

BER.: crown, forehead, and back of the neck and throat, black; on the cheek a patch of brown margined by a line of white, which afterwards passes down the neck and breast, beyond which is a bar of white; belly, wings, and tail, black; abdomen, vent, upper and under tail coverts white; wing coverts margined with white; bill and legs fuscous.

<i>Anser ruficollis</i>	<i>Pall. Spic.</i> 6. 21.
<i>Anas ruficollis</i>	<i>Gmel. Syst.</i> 1. 511.
<i>Anas torquata</i>	<i>Gmel. Syst.</i> 1. 514.
<i>Bernicla ruficollis</i>	<i>Steph. Sh. Zool.</i> 12. 53.
<i>Red-breasted goose</i>	<i>Penn. Arct. Zool.</i> 4. 571.
	<i>Lath. Syst.</i> 6, 455.

A few specimens have at different times been killed in the British isles. It is a native of the arctic region, and is said to retire to Siberia and the Frozen Ocean to breed. Temminck informs us that it is in abundance near the rivers Ob and Lena during the breeding season. Nothing, however, appears to be known of its nidification or habits.

BERNICLA BRENTA.

Ber.—Cinerea, capite colloque nigro; hoc lateribus albo maculatis.

BRENT BERNICLE.

BER. cinereous, with the head and neck black; the latter on the sides with a white patch.

	IN.		IN.
<i>Length</i>	29	<i>Tarsi</i>	2 2-5ths
<i>Bill</i>	1 2-5ths	<i>Middle toe</i>	2

BER: head, neck, and breast black, on each side of the neck a patch of white; back and wing

coverts brown, with the edges of the feathers paler; below dirty white, with a yellow tinge; the margin of the feathers pure white; back and upper tail coverts black; neck and bill black.

<i>Bernicla Brenta</i>	<i>Steph. Sh. Zool.</i> 12. 46.
<i>Anas Bernicla</i>	<i>Lin. Syst.</i> 1. 198.
	<i>Gmel. Syst.</i> 1. 513.
	<i>Lath. Ind. Orn.</i> 2. 844.
<i>Brent Goose</i>	<i>Penn. Arct. Zool.</i> 2 N. 478.
<i>Le Cravant</i>	<i>Buff. Ois.</i> 9. pl. 87.

Inhabits the same countries with *B. Leucopsis*; its nest is made of grass and sea-weed in morasses; it lays ten or twelve eggs, perfectly white.

Like its congeners, it is easily domesticated. The trachea and inferior larynx is that of the typical geese, but slightly expanded below the glottis.

BERNICLA LEUCOPSIS.

Ber.—Cinerea, capite colloque atro; facie albâ.

COMMON BERNICLE.

BER. cinereous, with the head and neck black; the face white.

	IN.		IN.
<i>Length</i>	25	<i>Tarsi</i>	2½
<i>Bill</i>	1¼	<i>Middle toe</i>	2½

BER.: cheeks, forehead, and throat white; crown, neck, and breast black; above blue grey, with the margins of the feathers lighter, within which there is a line of black; tail and quills black; tail coverts and the under surface white; the flanks with some dark bars of grey.

<i>Anser Leucopsis</i>	<i>Bechs. Nat. Deut.</i> 4. 921.
<i>Anas Leucopsis</i>	<i>Temm. Mun.</i> 2. 828.
<i>Bernicle</i>	<i>Penn. Arct. Zool.</i> 2 N. 479.
	<i>Lath. Gen. Syn.</i> 6. 466.
<i>Le Bernache</i>	<i>Buff. Ois.</i> 9. 9. 3. f. 5.

An arctic species, and, like most of its genus, is only a winter visitant in these islands; is easily domesticated. Little is known of its nidification, it having only once or twice bred in confinement. The trachea is like that of *B. Brenta*, but of rather smaller diameter.

BERNICLA JUBATA.

Ber. Mas.—Capite cristato, colloque brunneis; corpore cinereo, atro alboque maculato; abdomine atro.

Ber. Fem.—Capite colloque brunneis albo sparsis; corpore cinereo-brunneo et albo maculato; abdomine albo.

HAWKSURY BERNICLE.

BER. Male, with the head crested, together with the neck brown; body cinereous, spotted with black and white; abdomen black.

BER. Fem. with the head and neck brown blotched with white; body cinereous, spotted with brown and white; abdomen white.

	IN.		IN.
<i>Length</i>	23	<i>Tarsi</i>	2
<i>Bill</i>	1 1-6th	<i>Middle toe</i>	2 1-6th

Male: with the head and upper part of the neck umber brown; the feathers on the occiput much elongated and darker; upper part of the back, wing coverts, and lower part of the neck, cinerous, in the latter each feather having on its lateral edges a spot of black, and being tipped with white: lower part of the back, tail, abdomen, and belly in the centre, black; quills brown; tertiary long, the largest cinereous, the smaller ones with the outer webs black with metallic green reflections; wing spot metallic green, bordered anteriorly with a band of white, then with one of black posteriorly with one of white; flanks white, beautifully barred and irrorated with narrow lines of black; bill and legs black.

Fem.: rather smaller than the male; head and neck as in the male, but spotted anteriorly with small white specks; a whitish line passes from the bill below the eye, and another above it; breast and flanks white, each feather having a transverse bar of brown, narrowest in the centre, and expanding at the edges; abdomen and vent white; tail coverts cinereous; remainder as in the male, with the exception of the bill, which is horn colour,

Bernicla Jubata *Steph. Sh. Zool.* 12. 63.

Anas Jubata *Lath. Ind. Orn. App.* 69.

Hawksbury Bernicle *Lath. Syn. Sup.* 2. 358.

Inhabits New South Wales, in the neighbourhood of the Hawksbury river; is said occasionally to perch in trees.

CHENISCUS MADAGASCARIENSIS.

Ber.—Dorso colloque maculâ viridi; subtus albus.

MADAGASCAR BERNICLE.

BER. with the back and neck with a patch of green; below white.

	IN.		IN.
<i>Length</i>	$14\frac{3}{4}$	<i>Middle toe</i>	7-12ths
<i>Bill</i>	$\frac{7}{8}$		

Male: occiput, a stripe down the back of the neck; scapulars and rump green, with purplish inflections; quills and primary coverts sooty black, the latter edged with white; upper part of the breast, tail coverts, and flanks, reddish yellow; lower part of the breast, belly, forehead, chin, cheeks, and front of the neck, white; an ovate patch on each side of the neck, light green, with a black edge irrorated with white; under tail coverts black; bill gamboge yellow; nail, legs, and feet, black.

Fem.: as in the male, but without the green marking on the neck, instead of which it is grey; the forehead is also grey; in other respects similar.

<i>Anas Madagascariensis</i>	<i>Gmel. Sys. Nat.</i> 1. 522.
	<i>Lath. Ind. Orn.</i> 2. 875.
<i>Madagascar Teal</i>	<i>Lath. Gen. Syn.</i> 6. 556.
<i>Sarcelle de Madagascar</i>	<i>Buff. Pl. Enl.</i> 770.

Inhabits Madagascar. Specimens are in the British Museum, and in that of the Zoological Society. Mr. Leadbeater also possesses two most beautiful specimens. We have also been so fortunate as to obtain a specimen of the female.

CHENISCUS COROMANDELIANUS.

Ber.—Vertice, alis, dorso, caudaque, æneis-fuscis ; colloque striis nigricantibus strigato.

COTTON BERNICLE.

BER. with the crown, wings, back, and tail, brassy brown ; neck streaked with black.

<i>Length</i>	IN. 13 $\frac{1}{4}$	<i>Bill</i>	IN. 1	<i>Tarsi</i>	IN. 1
---------------------	-------------------------	-------------------	----------	--------------------	----------

BER.: crown black or brassy brown ; the remainder of the head and neck white ; a bar of metallic green across the breast ; wings and body brown or black, glossed with green ; beneath white, with the lateral under-tail coverts ferruginous ; legs black ; bill lead-coloured.

The female differs from the male in the markings being less distinct, and being slightly smaller.

Both sexes have occasionally the basal half of the quills white.

<i>Anas Coromandeliana</i>	<i>Gmel. Syst. Nat.</i> 1. 522.
	<i>Lath. Ind. Orn.</i> 2. 875.
<i>Bernicla Coromandeliana</i>	<i>Steph. Sh. Zool.</i> 12. 56.
<i>Coromandel Teal</i>	<i>Lath. Gen. Syn.</i> 6. 556.
<i>Sarcelle de Coromandel</i>	<i>Buff. Pl. Enl.</i> 949-950.
<i>Bernicla Girra</i>	<i>Gray Ind. Zool.</i>

Is found in Coromandel, Australia, and India. Specimens are in the museum of the Zoological Society, and in the British Museum.

This bird, which is included in Col. Sykes's catalogue of birds found in the Deccan, contained in the "Proceedings of the Zoological Society," is said by that gentleman to be "monogamous: vegetable matter and gravel were found in the stomach. These birds, when wounded, dive, and on returning to the surface shew only the bill above water, keeping the body below at pleasure." It is called the "Cotton Teal" by the Europeans of the Deccan, from the quantity of white in the plumage.

ANSER INDICUS.

Ans.—Cinereus; capite albo; fasciâ lunatâ occipitali brunneâ.

BAR-HEADED GOOSE.

Cinereous goose; with the head white; a lunate fascia on the occiput brown.

	IN.		IN.
<i>Length</i>	27	<i>Tarsi</i>	3
<i>Bill</i>	$1\frac{3}{4}$	<i>Middle toe</i>	$2\frac{3}{4}$

Light grey goose; with the head and neck white; a brown lunule on the occiput, and an irregular fascia a little below it, from which extends down the back of the neck a broad streak, both of the same colour as the lunule; anterior part of the neck brown grey, having a space of white down each side between it and the brown at the back; centre tail feathers and quills brown; lateral tail feathers, upper and under tail coverts, and under surface, yellowish white; a brown patch undulated with grey on each side above the thighs; bill and legs orange.

<i>Anser Indica</i>	<i>Steph. Sh. Zool.</i> 12, 36.
<i>Anas Indica</i>	<i>Lath. Ind. Orn.</i> 2, 839.
<i>Bar-headed Goose</i>	<i>Lath. Gen. Syn. Sup.</i> 277.
<i>Anser Indica</i>	<i>Gould. Cent. Birds.</i>

Found in most parts of India during the winter months.

ANSER ERYTHROPUS.

Ans.—Cinereus; maculâ albâ basum rostri cingente.

LAUGHING GOOSE.

Cinereous goose; with a white patch surrounding the base of the bill.

	IN.		IN.
<i>Length</i>	27	<i>Tarsi</i>	$2\frac{1}{2}$
<i>Bill</i>	2	<i>Middle toe</i>	$2\frac{3}{4}$

Head, neck, chest, breast, and belly, grey brown, the latter marked with irregular patches and

bars of black; back brown, with each feather margined with lighter; quills dark hoary brown; vent, abdomen, and upper tail coverts white; tail grey brown tipped with white; the two outer tail feathers entirely white; a white patch on each side the base of the bill.

<i>Anser Erythropus</i>	<i>Flem. Brit. An.</i> 127.
<i>Anas</i>	<i>Linn. Syst.</i> 1. 197.
<i>Anser albifrons</i>	<i>Steph. Sh. Zool.</i> 12. 25.
<i>Laughing Goose</i>	<i>Edw. Gl. pl.</i> 158.
<i>White-fronted Goose</i>	<i>Penn. Brit. Zool.</i> 2. 575.
	<i>Penn. Arct. Zool.</i> 2. 476.
	<i>Lath. Syn.</i> 6. 463.
<i>L'Oie rieuse</i>	<i>Buff. Ois.</i> 9. 81.

A regular winter resident in the British islands, and is found in the northern portion of the continents of Europe, Asia, and America. Specimens are in our collection, received from the Himalayan range and from within the arctic circle. Dr. Richardson, in the "Fauna Boreali Americana," informs us that its breeding places are in the woody districts skirting the Mackenzie, to the north of 67 parallel.

ANSER SEGETUM.

Ans.—Cinereus; rostro nigro, fasciâ flavâ ad apicem.

BEAN GOOSE.

Cinereous goose; the bill black, with a yellow fascia towards the tip.

	IN.		IN.
<i>Length</i>	30	<i>Tarsi</i>	3½

Head and neck, back, wings, tail, breast, abdomen, and flank, grey brown, each feather edged with lighter, and sometimes with white; vent, under tail coverts, and rump white; bill black with a band of yellow, in some specimens strongly tinged with pink, surrounding both mandibles near the extremity.

<i>Anser segetum</i>	Steph. Sh. Zool. 12. 27.
<i>Anas segetum</i>	Gmel. Syst. 1. 512.
	Lath. Ind. Orn. 2. 843.

- Bean Goose* *Brit. Zool.* 2. 575.
Arct. Zool. 2. V. 472.
L'Oie sauvage *Buff. Ois.* 9. 30.

Visits the British isles regularly during winter. Easily domesticated, becoming tame very shortly after their capture; but I am not aware that they have ever bred in this state, or does any thing appear to be known of their nidification.

Trachea and inferior larynx typical, the former slightly expanding above the latter part.

ANSER HYPERBOREAS,

Ans.—Albus; remigibus atris.

SNOW GOOSE.

White goose; with the quills black.

	IN.		IN.		IN.
<i>Length</i>	31 $\frac{1}{2}$	<i>Bill</i>	2 $\frac{1}{4}$	<i>Tarsi</i>	2 $\frac{3}{4}$

White, with the quills black; the legs orange; bill black. In the young state this bird has the wing coverts blue, and the remainder of the plumage more or less tinged with cinereous. The bill of this species gapes more at the sides than in any other of the genus.

- Anas hyperborea* *Gmel. Syst. Nat.* 1. 504.
Lath. Ind. Orn. 2. 837.
Pall. Spic. vi. p. 26.
Anser niveus *Briss. Orn.* 6. 288.
Oie hyperborcé ou le neige *Temm. Man. d' Orn.* 525.
Cur. Reg. An. 1. 531.
Anas carulescens *Linn. Syst. Nat.* 1. 198.
Gmel. Syst. Nat. 1. 513.
Lath. Ind. Orn. 2. 836.
Oie des Esquimaux *Buff. Hist. Nat. Ois.* 9. 80.

Met with in the northern parts of America and Europe.

ANSER MELANOPTERUS.

Anser.—Albus; rectricibus majoribus, tertiis, et secundariis, atris.

BLACK AND WHITE GOOSE.

White goose; with the greater quills, tertiaries, and secondaries, black.

	IN.		IN.
<i>Length</i>	28 $\frac{1}{2}$	<i>Tarsi</i>	3 $\frac{1}{2}$
<i>Bill</i>	1 $\frac{5}{8}$	<i>Middle toe</i>	3 $\frac{1}{2}$

White, with the greater quills and adjoining secondaries brown black, glossed with brassy; a few of the tertiary coverts with a longitudinal stripe of brown on the shaft; head and neck inclining to grey; bill orange; nail black; legs orange.

Anser melanopterus, nobis.

Brought home by Capt. King, obtained from the lake of Titicaca, Chili. In the museum of the Zoological Society, We observed also specimens in the British Museum, and in that of Lord Derby, at Knowsley.

ANSER FERUS.

Ans.—Cinereus; rostro aurantiaco.

WILD GOOSE.

Cinereous goose; with the bill orange.

	IN.		IN.
<i>Length</i>	34	<i>Bill</i>	2 $\frac{3}{8}$

Head, neck, back, and wing coverts, grey brown, the former rather the lightest; all the feathers edged with light grey; breast and belly dirty white, crossed by a stripe of grey; rump, vent, and under tail coverts white; bill deep orange, with the nail white; legs, tarsi, and feet red.

<i>Anser ferus</i>	<i>Steph. Sh. Zool.</i> 12. 28.
<i>Anas ferus</i>	<i>Gmel. Syst.</i> 1. 510.
	<i>Lath. Ind. Orn.</i> 2. 841.
<i>Grey lag Goose</i>	<i>Penn. Brit. Zool.</i> 2. 570.
	<i>Penn. Arct. Zool.</i> 2. 473.
	<i>Lath. Syn.</i> 6. 459.
<i>Oie cendree ou premiere</i>	<i>Temm. Man. Orn.</i> 2. 818.

We have followed Mr. Stephens and Mr. Gould in preserving to this species the name of *ferus*, and to one of the foregoing *A. segetum*, as being better known than those of Dr. Fleming used by Mr. Selby. The above bird arrives here in the winter in great numbers, and, according to Temminck, the distribution of it is very wide, being over Europe, Nubia, and part of Western Africa, inhabiting lakes and marshes.

Breeds in the north, and formerly reared its young in this country, laying about ten eggs of a white colour.

ANSER GUINEENSIS.

Ans.—Griseo-cinereus: subtus albus, tuberculo ad basin rostri.

SWAN GOOSE.

Griseous-cinereous goose; below white, with a tubercle at the base of the bill.

Length 39 inches.

Above cinereous grey, with the edges of the feathers light grey; from the occiput down the back of the neck runs a dark brown streak to the back; fore part of the neck, breast, and abdomen, dirty white; bill (which has a large knob at its base) greenish white colour; legs orange.

The female is less than the male, and with the colours not so distinct; the knob at the base of the bill is also smaller.

<i>Anser Guineensis</i>	<i>Raii Syn.</i> 1. 38.
	<i>Briss. Orn.</i> 6. 280.
<i>Anser cygnoides</i>	<i>Linn. Syst.</i> 1. 194.
	<i>Lath. Ind. Orn.</i> 2. 838.
<i>Chinese Goose</i>	<i>Penn. Arct. Zool.</i> 2. 571.
<i>Swan Goose</i>	<i>Bewick Brit. Birds,</i> 2. 281.
<i>L' Oie de Guinée</i>	<i>Buff. Ois.</i> 9. 72, <i>pl.</i> 3, <i>pl. Ent.</i> 347.

Said to inhabit (in a wild state) some parts of Siberia and Kamschatka, and to have been originally brought to this country from Guinea, but are now to be found in a semi-domesticated state in most parts of Europe.

This species crosses freely with the common tame goose, and their offspring are again productive.

Trachea of large diameter, and very much in this part, as in its whole anatomy, resembling the true swans. The trachea descends between the rami of the os furcatum, and is joined to the point from whence they diverge by a strong ligament, as in the black swan.

ANSER HUTCHINSII.

Ans.—Minusquam sequenti, sed similis.

HUTCHINS GOOSE.

Goose similar to the following, but smaller.

	IN.		IN.
<i>Length</i>	25	<i>Tarsi</i>	$2\frac{1}{2}$
<i>Bill</i>	$1\frac{2}{3}$	<i>Middle toe</i>	$3\frac{1}{3}$

Goose with the head, neck, rump, and tail, black; back and wings brown, with the edges of the feathers lighter; a small mark anterior to the eye, and a kidney-shaped patch on the throat, white; below white, with the flank transversely barred with grey; bill and feet black.

Anser Hutchinsii..... *Faun. Bo. Am.* 2 p. 270.

Anas bernicla, *Var. B.*..... *Rich app. Parry 2 Voy.* 1. 368.

A common species in Hudson's Bay and the arctic parts of America, where it has, until separated by Dr. Richardson, been confounded with the Canada goose, which it much resembles in every respect except in size.

ANSER CANADENSIS.

Anser.—Cinereus; capite colloque superiore atris; genis guttureque maculâ albâ.

CANADA GOOSE.

Cinereous goose; with the head and upper part of the neck black; the cheeks and throat with a white patch.

	IN.		IN.
<i>Length</i>	42	<i>Tarsi</i>	$3\frac{1}{2}$
<i>Bill</i>	2 1-6th	<i>Middle toe</i>	$4\frac{1}{4}$

Goose with the head, upper half of the neck, quills, tail, and rump, velvet black; back and wings brown, with the edges of each feather lighter; the lower part of the neck anteriorly and under surface white, with a grey tinge at the root of the feathers; a large kidney-shaped patch on the throat and cheeks, with the small ends placed behind the eye, white; upper tail coverts also white; bill and feet black.

Anas Canadensis

Linn. Syst. Nat. 1. 198.

Anser Canadensis

Bonap, Syn. Bir. Am. 377.

Aud. Orn. Biog. 3. 18.

This species is easily kept in a semi-domesticated state, and breeds freely. In a wild state it breeds in the neighbourhood of Hudson's Bay, from whence they migrate southward over the American Continent during winter. Lays about five eggs.

CYGNUS OLOR.

Cyg.—Albus ; rostro aurantiaco, tuberculâ culmine ad basin pedibusque atris.

MUTE SWAN.

Cyg. white ; bill orange, with a knob on the culmen and feet black.

Length 5 feet 6 inches.

<i>Anas olor</i>	<i>Gmel. Syst.</i> 1. 501.
<i>Anas cygnus mansuetus</i>	<i>Linn. Syst. Nat.</i> 1. 194.
	<i>Raii Syn.</i> 133 <i>A.</i>
<i>Mute Swan</i>	<i>Penn. Arct. Zool.</i> 2. 470.
	<i>Lath. Gen. Syn.</i> 6. 436.
<i>Le Cygne</i>	<i>Buff. Ois.</i> 9. 3 <i>Pl.</i> 1.

Supposed to have been introduced into this country from Asia, and is said at present to exist in a wild state in Russia and Siberia ; now found commonly on large rivers and pieces of water, where it is half domesticated and breeds freely, laying six or seven eggs of a dirty white colour, or with a very slight tinge of greenish ; sits about six weeks. The trachea in this species does not enter the sternum, but has the tube of equal thickness for its whole length, being merely slightly contracted above the inferior larynx.

CYGNUS IMMUTABILIS.

Cyg.—Albus ; rostro aurantiaco, tuberculâque culmine ad basin atro ; pedibus flavicantibus.

POLISH SWAN.

White swan ; with the bill orange, and a tubercle on the culmen at the base black ; the feet yellowish.

Similar in size and in every other particular to the preceding, except in the colouring of the feet and legs, which are light drab colour.

The young of this species are white from the eggs, and do not exhibit any of the grey or brown colouring found invariably among the others.

Cygnus immutabilis *Yarrell.*

For what we have said concerning this species we are altogether indebted to the Earl of Derby

and William Yarrell, Esq. We never have had an opportunity of minutely examining a specimen for ourselves. Several specimens have been killed this winter (1837-8) both in Scotland and England, from flocks varying in number from seven to thirty.

Mr. Yarrell also informs us, that hitherto he was only acquainted with the bird as an article of commerce among the London dealers in live birds, who distinguished it by the name of the Polish Swan.

CYGNUS ATRATUS.

Cyg.—Niger, cinereo undulato.

BLACK SWAN.

	FT.	IN.		IN.
<i>Length</i>	4	2	<i>Outer toe</i>	4 $\frac{1}{2}$
<i>Tarsi</i>	4 $\frac{1}{4}$		<i>Inner toe</i>	3 $\frac{3}{4}$
<i>Centre toe</i>	5		<i>Hind toe</i>	$\frac{7}{8}$

Black, with the edges of the feathers cinereous; quills, secondaries, and greater coverts white tipped with black; bill carmine red, marked transversely near the tip with a white band; nail white; legs and feet dark yellowish ash, with the webs darkest.

<i>Anas atrata</i>	<i>Lath. Ind. Orn.</i> 2. 834.
<i>Anas platonia</i>	<i>Shaw, Nat. Misc. pl.</i> 108.
<i>Cygnus atratus</i>	<i>Steph. Sh. Zool.</i> 12. 18.
<i>Black Swan</i>	<i>Lath. Gen. Syn. Sup.</i> 2. 343.
<i>Le Cygne noir</i>	<i>Cuv. Reg. An.</i> 1. 529.
<i>Trachea</i>	<i>Linn. Trans. vol.</i> 15, <i>tab.</i> 12.

Entirely an Australian species. The tube of the trachea does not enter the sternum, but is attached to the *os furcatum* by a ligament; in other respects the anatomy resembles that of the true swans.

CYGNUS NIGRICOLLIS.

Cyg.—Albus; capite colloque atris.

BLACK-NECKED SWAN.

White swan; with the head and neck black.

	FT.	IN.		IN.
<i>Length</i>	4	2	<i>Bill</i>	3 $\frac{3}{4}$

White swan; with the head and neck black; bill black, with two knobs at the base; legs orange.

<i>Anas nigricollis</i>	<i>Gmel. Syst. Nat.</i> 1. 502.
	<i>Lath. Ind. Orn.</i> 2. 834.
<i>Anas melanoccephalus</i>	<i>Gmel. Syst. Nat.</i> 1. 502.
<i>Cygnus nigricollis</i>	<i>Steph. Sh. Zool.</i> 12. 17.
<i>Anas melancorypha</i>	<i>Moll. French edit.</i> 213.

Inhabits the Falkland Isles and the Straits of Magellan.

CYGNUS BEWICKII.

Cyg.—Albus; rostro atro, baso usque ad nares aurantiaco.

BEWICK'S SWAN.

White swan; with the bill black, the base orange to the nostrils.

	FT.	IN.		IN.
<i>Length</i>	3	10 $\frac{1}{2}$	<i>Tarsi</i>	4 $\frac{3}{4}$
<i>Bill</i>		3 $\frac{1}{2}$	<i>Middle toe</i>	5

White; with the bill black, the basal half yellow, the yellow marking truncated at the extremity; legs and toes black.

<i>Cygnus Bewickii</i> , and trachea	<i>Yarr. Linn. Trans.</i> vol. 10.
	<i>Eyton, Hist. R. Brit. Birds</i> , p. 86.

Discovered by Mr. Yarrell and Mr. R. Wingate to be an inhabitant of the British isles.

The trachea of this species during youth penetrates the keel of the sternum in a similar manner to that of *C. ferus*, but may always be distinguished by the greater length of the bronchiæ. As the bird approaches maturity the tube at the internal extremity of the portion penetrating the sternum takes a horizontal twist, increasing in size and the depth to which it penetrates with age.

The tube of the trachea is acted upon by two pairs of muscles, the first pair sterno-tracheal, the other having their attachment on the membrane stretched between the rami of the *os furcatum* and the *sternum*. (*Vide fig.*)

Engraved by J. B. S.

ANATOMY OF THE VOICE OF CYGNUS BERNICULA.

The Swans Voice.

J. B. S. del.

CYGNUS AMERICANUS.

Cyg.—*Cygnus Bewickii* similis, sed majore.

AMERICAN SWAN.

Swan like *Cygnus Bewickii*, but larger.

	FT.	IN.		IN.
<i>Length</i>	4	1	<i>Tarsi</i>	4
<i>Ext. Alæ</i>	7	0	<i>Outer toe</i>	5½
<i>Bill</i>		4	<i>Middle toe</i>	6

The only specimen of this bird we have seen was shewn to us by Mr. Audubon; it appeared to have the bill more expanded towards the tip than is the case in *Cygnus Bewickii*, and altogether to be of larger size. The Prince of Musignano assures us that he is quite satisfied as to its distinctness from *C. Bewickii*.

<i>Cygnus Americanus</i>	<i>Sharpless in Silliman's Journal.</i>
<i>Trachea</i>	<i>Ditto ditto.</i>
<i>Cygnus Bewickii</i>	<i>Faun. Bo. Am. 2. 465.</i>

Inhabits North America.

The following is an extract from Dr. Sharpless's paper above quoted. We must not, however, forget to mention that the tracheal anatomy of this species is so very similar to that of *C. Bewickii*, that Dr. Sharpless has copied the illustrations of Mr. Yarrell's paper on that species, published in the Linnæan Transactions, to illustrate his on *C. Americanus*.

SPECIFIC CHAR. Beak black and semi-cylindrical; sides of the base with a small orange or yellow spot; body white; tail feathers twenty; feet black.

	<i>Hooper.</i>	<i>Bewick.</i>	<i>American.</i>
Weight	24 lbs. ..	13¾ lbs. ..	21 lbs.
Point of bill to the end of tail	5ft. ..	3ft. 9in. ..	4ft. 6in.
———— edge of forehead	4¾in. ..	3½in. ..	4½in.
———— eye	5¼in. ..	4¾in. ..	5in.
———— occiput	7¼in. ..	6¼in. ..	7½in.
Width of the beak at the widest part near the point			{ 1½in. 1¾in.
———— with wings extended	7ft. 10in. ..	6ft. 1in. ..	7ft. 2in.
Carpus to the end of primaries	2ft. 1½in. ..	1ft. 8½in. ..	1ft. 11in.
Length of middle toe	6½in. ..	5¼in. ..	6in.
———— intestine	12ft. ..	10ft. 2in. ..	10ft. 7in.
———— breast bone	8½in. ..	6¾in. ..	7½in.
Depth of insertion of trachea	3in. ..	5¾in. ..	6½in.
Length of bronchial tubes	3½in. ..	1½in. ..	1½in.
Tail feathers in number	20 ..	18 ..	20

It is not improbable, as Mr. Yarrell suggests, that this species and *C. Buccinator* will be found to be confined to America; *C. Bewickii* and *C. ferus* representing them in Europe.

Dr. Richardson, in the "Fauna Boreali Americana," says that "this swan breeds on the sea coast within the arctic circle, and is seen in the fur countries on its passage only. It makes its appearance among the latest of the migratory birds in the spring, while the Trumpeter swans are, with the exception of the eagles, the earliest. Capt. Lyon describes its nest as built of moss peat, nearly six feet long, four feet and three quarters wide, and two feet high exteriorly; the cavity a foot and half in diameter, the eggs brownish white slightly coloured with darker tints."

CYGNUS BUCCINATOR.

Cyg.—Albus; rostro atro.

TRUMPETER SWAN.

White swan; with the bill black.

	IN.		IN.
<i>Length</i>	42	<i>Tarsi</i>	$3\frac{1}{2}$
<i>Bill</i>	$2\frac{1}{2}$	<i>Middle toe</i>	$4\frac{3}{8}$

Cygnus Buccinator *Faun. Bo. Am.* 2. 464.

Trachea and sternum *Linn. Trans. vol. xvii. p. 1, T. 1.*

Dr. Richardson informs us, in the "Fauna Boreali Americana," that "this is the most common of the genus in the interior of the fur countries. It breeds as far south as lat. 61, but principally within the arctic circle, and in its migrations generally precedes the geese a few days. A fold of its trachea enters a protuberance on the dorsal or interior aspect of the sternum at its upper part, which is wanting in *C. ferus* and *C. Bewickii*." It is to the Trumpeter that the bulk of the swan skins imported from Hudson's Bay belong.

CYGNUS ANATOIDES.

Cyg.—Albus; remigibus primariis. ad apicem nigris; rostro pedibusque rubris, illo lato, subdepresso, tuberculo nullo. (*King.*)

DUCK-LIKE SWAN.

White swan; with the primaries black at the apices; bill and feet red, the former broad, subdepressed, without a tubercle.

	IN.		IN.
<i>Length</i>	42	<i>Tarsi</i>	3½
<i>Bill</i>	2½	<i>Middle toe</i>	4½

Cygnus anatoides. *King* *Zool. Proc.* Dec. 14, 1830.

Habitat in sinibus interioribus apud extremitatis meridionalem America. (*King.*)

I am informed by C. Darwin, Esq. that he observed this swan in company with *C. nigricollis* in the Straits of Magellan. The bill in this species is broader in proportion to its length than is usual amongst the swans.

CYGNUS FERUS.

Cyg.—Albus: rostro atro, baso ultra nares aurantiaco.

HOOPER.

White swan; with the bill black, the base orange to beyond the nostrils.

	IN.		IN.
<i>Length</i>	60	<i>Tarsi</i>	4
<i>Bill</i>	4½	<i>Middle toe</i>	6½

White; with an occasional tinge of buff colour; bill black, cere yellow, and is carried to a point before the nostrils on the edge of the upper mandible; legs and feet black.

<i>Cygnus ferus</i>	<i>Raii Syn.</i> 130 A.
<i>Anas Cygnus ferus</i>	<i>Linn. Syst.</i> 1, 194.
	<i>Gmel. Syst.</i> 1, 501.
	<i>Lath. Ind. Orn.</i> 2, 833.
<i>Whistling Swan</i>	<i>Penn. Arct. Zool.</i> 2, 833.
<i>Trachea</i>	<i>Linn. Trans.</i> vol. 4, tab. 12, fig. 1, 2.

A winter visitant in the British isles, particularly during hard weather; its true habitation, however, is within the arctic circle. In Iceland it also occasionally breeds.

Lives well and is prolific on large pieces of water in a semi-domesticated state, sitting six weeks, and laying from four to six eggs of a greenish colour.

The trachea in this species perforates the keel of the sternum deeply in a vertical direction.

SARKIDIORNIS REGIUS.

Sar.—Supra niger-æneus, subtus albus; mas. carunculâ magnâ culmine rostri.

BLACK-BACKED GOOSE.

Goose with the upper part brassy-black; below white; the male with a large caruncle at the base of the bill.

	IN.		IN.
Male: <i>Length</i>	30	<i>Tarsi</i>	$2\frac{3}{4}$
<i>Bill</i>	$2\frac{1}{2}$	<i>Middle toe</i>	$3\frac{1}{4}$

Bill black, with a large caruncle on the culmen; head and neck white, spotted with metallic green; under parts white, and wings purplish black, glossed in some lights with brassy, a band of black extending from the back to the posterior part of the thighs; wings with a knob; lower part of the back white; tail coverts black.

Fem.: *Length* 27 inches *Bill*..... 2 inches

Bill without the caruncle, colouring similar, but with the white more or less tinged with brown, and without the band of black behind the thighs; bill and legs in both black.

<i>Anas regia</i>	<i>Gmel. Syst. Nat.</i> 1. 508.
	<i>Lath. Ind. Orn.</i> 2. 847.
	<i>Moll. Hist. du Chili.</i> p. 212.
<i>Carina</i>	<i>Steph. in the Zool.</i> vol. 12. 82.
<i>Anas melanotus</i>	<i>Gmel. Syst. Nat.</i> 1. 503.
	<i>Lath. Ind. Orn.</i> 2. 839.
<i>L' Oie bronzee d' Coromandel</i>	<i>Buff. Hist. Nat. des Ois.</i> pl. 937.
<i>Black-backed Goose</i>	<i>Lath. Gen. Syn.</i> 6. 449.
<i>Le cravant royal</i>	<i>Veil. En. Method. Orn.</i> vol. 1. 126.
<i>L' Oie bronzee</i>	<i>Buff. Ois.</i> 115.

Found in Cayenne and most parts of India; is also included in the list of birds of the Deccan, by Col. Sykes, who also says that seeds of water grasses, and the remarkable quadrangular hard seeds met with in the stomach of *Pterocles exustus*, were also found in the stomach of this bird. Col. Sykes found the digastric muscle of the remarkable thickness of $1\frac{1}{2}$ inch.

SARKIDIORNIS AFRICANUS.

Sar.—Minusquam precedenti, sed in colore similis.

AFRICAN BLACK-BACKED GOOSE.

Less than the preceding, but in colour similar.

		IN.			IN.
Male:	Length.....	23		Tarsi.....	2
	Bill.....	2		Middle toe.....	2½
Fem.:	Length.....	21		Tarsi.....	1¾
	Bill.....	1¾		Middle toe.....	2½

Both sexes resemble perfectly in colouring the preceding species, except in admeasurement, in which they differ considerably. We have never seen a specimen with the caruncle, our specimen having evidently had it cut off, probably from the difficulty of preserving it.

Sarkidiornis Africanus nobis.

The four specimens in our collection were received from the Gamba river. Specimens have also been brought home by Dr. Smith from Southern Africa, who will in his forthcoming work publish a figure of this species.

CHENALOPEX ÆGYPTIACA.

Chen.—Dorso castaneo; subtus leviter flavo-ferrugineo, sparsus lineatusque brunneo.

ÆGYPTIAN CHENALOPEX.

With the back castaneous; below slightly tinged with ferruginous yellow, sprinkled and lined with brown.

		IN.			IN.
	Length.....	27½		Tarsi.....	3
	Bill.....	2½		Middle toe.....	3

Male: head and neck hoary, with the occiput and a stripe down the back of the neck approaching to ferruginous; a patch round the eyes and ring round the lower part of the neck deep castaneous; upper part of the back, scapulars, tertiaries, and a triangular patch on the breast, also castaneous, some of the feathers in the former being irrorated with black; the inner webs also of some of the tertiaries hoary grey. The whole of the remainder of the breast and under surface is light yellow ferruginous irrorated with brown, except the abdomen, which is pure white; vent ferruginous; lower part of the back, tail, quills, and spurious wing, black; secondaries bright metallic green, forming the wing spot; wing coverts white, the greater ones with a bar of black near the tips; bill and legs orange.

Female: with the chesnut patches round the eyes and on the breast smaller,

<i>Anas Ægyptiaca</i>	<i>Linn. Syst.</i> 1. 197.
	<i>Gmel. Syst.</i> 1. 512.
	<i>Briss. Orn.</i> 6. 281. <i>pl.</i> 27.
	<i>Lath. Ind. Orn.</i> 2. 810.
<i>Chenalopex Ægyptiaca</i>	<i>Steph. Sh. Zool.</i> 12. 43.
<i>L' Oie d' Ægypte</i>	<i>Buff. Pl. En.</i> 379, 982, 983.
<i>Egyptian Goose</i>	<i>Lath. Syn.</i> 6. 155.

Inhabits most parts of Africa, especially Egypt and the Cape of Good Hope. We have figured the trachea and skeleton of this bird in illustration of the genus.

CHENALOPEX CANA.

Che.—Mas. castaneus capite albo; occipite colloque canis.

Che.—Fem. Mari similis sed capite omnino cano.

GREY-HEADED SPURWING.

	IN.		IN.
<i>Length</i>	25	<i>Tarsi</i>	2 $\frac{1}{4}$
<i>Bill</i>	1 $\frac{3}{4}$	<i>Middle toe</i>	2 $\frac{1}{4}$

Male: castaneous chenalopex, with the whole of the head excepting the crown white; neck and crown grey; wing coverts white; secondaries with the outer webs green; quills, tail, and vent hoary brown, the latter with a black band; bill and legs black; the bend of the wing armed with a spur.

Female: similar to the male, but with the whole of the head hoary.

<i>Anas Cana</i>	<i>Gmel. Syst.</i> 1. 510.
	<i>Lath. Ind. Orn.</i> 2. 840.
<i>Bernicla Cana</i>	<i>Steph. Sh. Zool.</i> 12. 61.
<i>Grey-headed Goose</i>	<i>Lath. Syn.</i> 6. 458.
<i>L' Oie sauvage a tête grise de Coromandel</i>	<i>Sonn Voy.</i> 2. 220.

The above descriptions were taken from two specimens brought home by Dr. Smith, and exhibited in the South African museum, having been obtained in the neighbourhood of the Cape. It has, however, been introduced alive into this country, and kept in confinement both by Lord Derby at Knowsley and in the gardens of the Zoological Society. Specimens are in the museum of that society.

TADORNA BELLONII.

Tad.—Albus; capite colloque atris; rostro cruribusque rubris.

COMMON SHELDRAKE,

White sheldrake; with the head and neck black; the bill and legs red,

	IN.		IN.
<i>Length</i>	23 $\frac{1}{2}$	<i>Tarsi</i>	2
<i>Bill</i>	2	<i>Middle toe</i>	2 $\frac{1}{4}$

Head and upper part of the neck black, glossed with green; fore part of the wings, back, tail, abdomen, and tail coverts, white; tail feathers tipped with black; a line down the middle of the abdomen black; upper part of the back and band across the breast deep ferruginous; quills black; wing spot green; bill and legs reddish orange, the latter in some specimens inclining to flesh colour.

<i>Anas Tadorna</i>	<i>Linn. Syst. Nat.</i> 1. 195-4.
	<i>Raii Syn.</i> 140 <i>A.</i> 1.
<i>Tadorna vulpanser</i>	<i>Flem. Brit. An.</i> 1. 122.
<i>Le Tadorne</i>	<i>Buff. Ois.</i> 9. 205.
<i>Canard Tadorne</i>	<i>Temm. Man.</i>
<i>Tadorna vulpanser</i>	<i>Selb. Illust. Brit. Orn.</i> 11. 289. .
	<i>Gould, Birds of Europe.</i>
<i>Tadorna Bellonii</i>	<i>Steph. Sh. Zool.</i> 12. 72.
<i>Trachea</i>	<i>Linn. Trans. vol. iv. tab.</i> 15, <i>fig.</i> 9, 10.

Found over nearly the whole of Europe and some parts of Asia and America; lays twelve eggs of a whitish colour, generally in rabbit holes; frequents the sea. Trachea of male slightly enlarged about an inch and half above the inferior larynx, and again contracted immediately above it, as among the *Anserinæ*. The inferior larynx is furnished with two large compressed bulbs, the one on the right side being nearly double the size of that on the left; but neither are of such strength as in those species of the last family which possess that adjunct.

TADORNA RADJAH.

Tad.—Capite, collo, ventre, ptilisque albis; collo inferiore lato castaneo cingulo ornato, sparso atro.

RADJAH SHELDRAKE.

Head, neck, belly, and anterior part of the wing white; the lower part of the neck surrounded by a band of deep castaneous irrorated with black.

<i>Length</i>	^{IN.} 22½	<i>Bill</i>	^{IN.} 1½	<i>Tarsi</i>	^{IN.} 2
-------------------------	-----------------------	-----------------------	----------------------	------------------------	---------------------

Head, neck, belly, and anterior part of the wing white; lower part of the neck surrounded by a collar of deep castaneous, thickly and beautifully irrorated with black; wing spot bright green, anterior to which there is a white stripe, then a very narrow one of black, posteriorly a narrow black stripe followed by one of white; scapulars dark sienna brown irrorated with black; tertiaries with the outer webs sienna brown; inner webs, quills, tail, back, and under tail coverts, black; vent black irrorated with white; bill and legs orange.

Anas Radjah *Duperrey, Voy. de la Coquille, pl. 49.*

Specimens are in most museums, received from Australia, but from what part we cannot say. Although we have examined many specimens, no sexual difference has been observed.

CASARKA RUTILA.

Cas.—Ferruginea; caudâ remigesque primariis atris; speculo viridi.

FERRUGINOUS SHELDRAKE.

Ferruginous sheldrake; with the tail and quills black; the wing spot green.

<i>Length</i>	^{IN.} 24½	<i>Bill</i>	^{IN.} 1½	<i>Tarsi</i>	^{IN.} 2
-------------------------	-----------------------	-----------------------	----------------------	------------------------	---------------------

Ruddy ferruginous, with the head and anterior part of the wing lighter, sometimes nearly white; tail and lower part of the back black glossed with green; quills black; secondaries bright glossy green, with the inner webs hoary; bill and legs black; sometimes a black ring round the neck is visible.

<i>Casarka rutila</i>	<i>Bonap. Cat. Am. and Europ. Birds.</i>
<i>Anas Casarka</i>	<i>Linn. Syst. 3 App. 224.</i>
<i>Anas rutila</i>	<i>Pall. Nov. Com. Petrop. 14. 579.</i>
<i>Tadorna rutila</i>	<i>Steph. Sh. Zool. 12. 71.</i>
	<i>Selb. Illust. Brit. Orn. 2. 293.</i>
<i>Canard Casarka</i>	<i>Temm. Man. Orn. 2. 832.</i>
<i>Ruddy Sheldrake</i>	<i>Gould, Birds of Europe.</i>
<i>Ferruginous Duck</i>	<i>Bew. Brit. Birds, 2. 313.</i>
<i>Ruddy Goose</i>	<i>Lath. Syn. 6. 456.</i>

Met with occasionally in most parts of the European continent, India, and Africa; lays eight white eggs; frequents chiefly fresh water. Col. Sykes found these birds for the most part of the year in pairs in the Deccan; but on the Nerbudda river, in Guzerat, he has seen them congregated in hundreds in April. Col. Sykes found the stomachs of many of them filled with grass seeds and vegetable matter only.

CASARKA TADONOIDES.

Cas.—Capite, caudâ, vertreque atris, hoc castaneo sparso; cingulo albo collo.

NEW HOLLAND SHELDRAKE.

Sheldrake, with the head, tail, back, and belly black, the latter irrorated with castaneous; a white ring round the neck.

	IN.		IN.		IN.
<i>Length</i>	27	<i>Bill</i>	1 $\frac{7}{8}$	<i>Tarsi</i>	2

Head, back, belly, and tail black; the back and belly irrorated with ferruginous; the lower part of the back and tail slightly tinged with green; upper part of the back, breast, and outer webs of the tertiaries ferruginous; a white collar round the neck; fore part of the wing white; wing spot light green; quills black; legs and bill black. In the female a white band surrounds the base of the bill, from which another passes through the eye.

Anas Tadonoides, *Jard. & Selb. Illus. Orn. pl. 411.*

Inhabits New Holland.

CASARKA CASTANEA.

Cas.—Castanea; capite colloque albo.

CASTANEOUS SHELDRAKE.

Castaneous sheldrake; with the head and neck white.

	IN.		IN.
<i>Length</i>	29	<i>Tarsi</i>	$2\frac{1}{2}$
<i>Bill</i>	$2\frac{1}{2}$	<i>Middle toe</i>	$2\frac{1}{2}$

Sheldrake, with the body below castaneous; the head, neck, and greater wing coverts white; the back, rump, and tail brown, the former speckled with white; tertiaries with the external webs castaneous; the internal ones and some of the secondary coverts blueish grey; bill black; legs crimson.

Casarka castanea nobis.

Purchased from Mr. Leadbeater, who received it from New Holland.

DENDROCYGNA ARCUATA.

Dend.—Fusca; marginibus pennarum vinaceis, subtus castanea; vertice brunneo.

ARCUATED WHISTLING DUCK.

Fusca duck; with the margins of the feathers vinaceous, below castaneous; the head brown.

	IN.		IN.
<i>Length</i>	19	<i>Tarsi</i>	$1\frac{3}{4}$
<i>Bill</i>	$\frac{5}{8}$	<i>Middle toe</i>	$2\frac{3}{4}$

Duck, with the head and neck whitish, slightly tinged with yellow vinaceous; crown brown; back hoary fuscous, the edges of each feather vinaceous, forming arcuated lines; wing and upper tail coverts deep chesnut; below ferruginous; under tail coverts white; quills and tail black; bill and feet also black.

Female smaller, and the colours less distinct.

<i>Anas arcuata</i>	<i>Hors. Zool. Res. Java.</i>
<i>Anas Javanica</i>	<i>Hors. Arrang. of the Birds of Java: Linn.</i>
	<i>Trans. vol. xiii. p. 199.</i>
<i>Mareca Ansuree</i>	<i>Sykes, Proc. Zool. Soc. 1832, p. 168.</i>

Tadorna erythrorhynchos. (india)

Red-billed Teal

Found by Dr. Horsfield in Java, and by Col. Sykes in the Deccan. It appears to be generally dispersed over India. Col. Sykes, in his Synopsis, says it is "gregarious, and abundant in the Deccan. Sexes alike in plumage," "These birds are characterised by a very peculiar whistle when disturbed, by a proportionate length of the intestine (one-third shorter than any other species of the *Anatidæ*), and by the inferior larynx being dilated into two oblong chambers, placed rather in front of, than lateral to, the trachea."

DENDROCYGNA AUTUMNALIS.

Dend.—Fusca; ventre atro; rostro pedibusque rubris.

AUTUMNAL WHISTLING DUCK.

Fuscous; with the belly black; bill and legs red.

	IN.		IN.
<i>Length</i>	24	<i>Tarsi</i>	2
<i>Bill</i>	2	<i>Middle toe</i>	2½

Back, crown, head, collar on the neck, and tertiary coverts, vinaceous; streak down the back of the neck black fuscous; throat, cheeks, and upper part of the breast grey, inclining to vinaceous on the former; secondaries, top of the primaries, tail, belly, and lower part of the back black; under tail coverts barred with black and dirty white; base of the primaries and some of the primary coverts white.

<i>Anas Autumnalis</i>	<i>Linn. Syst.</i> 1. 205.
	<i>Gmel. Syst.</i> 1. 537.
	<i>Lath. Ind. Orn.</i> 2. 852.
<i>Red-billed Whistling Duck</i>	<i>Edw. Glea. pl.</i> 194.
	<i>Lath. Gen. Syn.</i> 6. 498.
<i>Siffleur a bec rouge et narines jaune</i>	<i>Buff. Ois.</i> 9. 183.

Inhabits the West Indian Isles; is common also in South America. We have received it from Peru, where it is not uncommonly kept tame as in this country.

DENDROCYGNA VIDUATA.

Dend. Mas.—Capite anteriore guttureque albis; subtus atro.

Dend. Fem.—Mari similis, sed obscurior.

WIDOW WHISTLING DUCK.

Male duck, with the fore part of the head and throat white; below black.

Fem. duck, similar, but with the marking not so bright.

	IN.		IN.		IN.
<i>Length</i>	20	<i>Bill</i>	$1\frac{7}{8}$	<i>Tarsi</i>	$1\frac{3}{4}$

Fore part of the head and throat whitish; crown white, with the tip of the feathers stiff like hair, and olive brown; occiput and hind neck black; upper part of the neck and breast dark chesnut; back fuscous olive, the edge of each feather lighter; smaller wing coverts chesnut; quills and secondaries black; remainder of the wing fuscous, with a tinge of green; tail, middle of the abdomen, and thighs black; flanks yellowish white barred with black; legs and bill black.

Female similar, but with the colours not so bright.

Anas viduata *Linn. Syst.* 1. 205.

Gmel. Syst. 1. 536.

Lath. Ind. Orn. 2. 858.

Spanish Duck *Penn. Genera of Birds*, pl. 13.

Canard a face blanc de Maragnon *Buff. Ois.* 9. 255.

Inhabits both South America and Senegal. Is represented by travellers as having, like the rest of its genus, a peculiar whistling note.

DENDROCYGNA ARBOREA.

Dend.—Brunnea albo maculato; vertice nigro.

BLACK-BILLED WHISTLING DUCK.

Brown duck, mottled with white; the crown black.

	IN.		IN.
<i>Length</i>	$19\frac{1}{2}$	<i>Middle toe</i>	3
<i>Bill</i>	$2\frac{1}{8}$	<i>Outer toe</i>	2 3-5ths
<i>Tarsi</i>	3	<i>Hind toe</i>	1
<i>Inner toe</i>	2 1-5th		

Crown black; fore part of the head, back, breast, and wings, dark brown; lighter on the

edges of the feathers; breast and neck dirty white, at the lower part slightly spotted with black and brown; a streak down the back of the neck dark brown; under surface white, each feather barred with black and brown; thighs the same; tail and lower part of the back black; bill lead-coloured; legs and feet black.

<i>Anas Arborea</i>	Linn. Syst. Nat. 1. 207.
	Gmel. Syst. Nat. 1. 540.
	Lath. Ind. Orn. 2. 852.
<i>Anas fistularis Jamaicensis</i>	Raii Syn. 192.
<i>Black-billed Whistling Duck</i>	Edw. Glea. pl. 193.
<i>Canard Siffleur de St. Dominique</i>	Buff. Ois. 804.

Inhabits the West Indies and the adjacent portions of America.

LEPTOTARSIS EYTONI.

Lep.—Brunneus; lateribus axillaribusque ferrugineis.

BROWN SLENDERLEGS.

Brown duck, with the flanks and axillaries ferruginous.

	IN.		IN.
<i>Length</i>	21	<i>Tarsi</i>	2 1-5th
<i>Bill</i>	1½	<i>Middle toe</i>	2½

Duck, with the wings, back, and scapulars vandyke brown, inclining to slaty blue on the upper part of the back; flanks posteriorly bright ferruginous buff, ornamented with very long axillary feathers, each of which are edged with black, and entirely covered when the wing is closed; tail coverts also buff; the flanks anteriorly and under wing coverts bright ferruginous, the former beautifully barred with black; crown and streak down the back of the neck sienna brown, the feathers on the former rather stiff; below luteous, inclining to dirty lemon colour; the chin and throat lighter; two-thirds of the bill towards the base black; the remainder, with the legs, approaching to orange yellow.

Leptotarsis Eytleri,..... Gould. MSS.

This beautiful and unique bird, which Mr. Gould has done us the honour to call by our name, is in the possession of the Zoological Society, and was received from Australia.

DAFILA UROPHASIANUS.

Daf. Mas.—Caudâ pallidè rufis; dorso, capite, colloque brunneis, marginibus pennarum saturatoribus.
Daf. Fem.—Mare similis, sed subtus fuscus.

PINTAIL.

Pintail, with the tail pale rufous; the back, head, and neck brown; with the edges of the feathers lighter.

Fem. like the male, but below fuscous.

	IN.		IN.		IN.
<i>Length</i>	$24\frac{1}{2}$	<i>Bill</i>	2	<i>Tarsi</i>	$1\frac{3}{4}$

Male: with the head, neck, and body grey brown; the edges of the feathers light fuscous; cheeks, throat, and upper part of the fore neck, white; wing coverts fuscous; speculum bright green, anterior to which there is a stripe of pale red-fuscous; posteriorly a narrow one of black, then a broad one of pale reddish fuscous; tail and upper tail coverts pale reddish; bill and legs fuscous, the former with an orange patch on each side,

Female: similar to the male, but much smaller; and below brownish fuscous.

Anas urophasianus, King *Zool. Jour.* 4. 351.

Although some difference exists in length between our specimens of this bird and that described by Capt. King as above, and also a trifling one in colouring, we have been induced to place them as one, both having been received from the west coast of South America, placing ours the smallest as the female. Mr. Vigors, however, makes no mention of the orange marking on each side of the bill, which we suspect to be only apparent in old birds during the season of love. The buff colour on the wing also changes to white during certain times of the year in all the Pintails.

Frontalis, p. 1. (Mallard)

Dafila Urophasianus

DAFILA PYROGASTER.

Daf.—Subtus alba, singulis pennis marginatis apices versus lunulis brunneis.

BROWN-BELLIED PINTAIL.

Pintail; below white, with each feather edged near its tip with a lunule of bright brown.

	IN.		IN.
<i>Length</i>	24	<i>Inner toe</i>	$1\frac{3}{4}$
<i>Bill</i>	1 4-5ths	<i>Middle toe</i>	$2\frac{1}{8}$
<i>Tarsi</i>	2	<i>Outer toe</i>	2

Under parts, breast, and upper part of the back dirty white, each feather having near its tip a lunule of brown, brighter on the breast and shaded off at the edges; tail black; wing coverts brown; wing spot brassy purplish, edged posteriorly with brassy green, and then with a broad band of white, formed by the secondaries; neck dirty white, slightly spotted with brown and black; head slightly crested; crown and crest brown; bill lead coloured; legs black.

Anas pyrogaster *Meyen Nova acta.*

In the museum of the Zoological Society, and in Lord Derby's collection. Inhabits Chili.

DAFILA CAUDACUTA.

Daf. Mas.—Capite colloque anteriore brunneis; posterioriter atro, albo marginato.

Daf. Fem.—Brunnea, marginibus pennarum suturationibus; speculo flavo-rufo.

PINTAIL.

Male pintail, with the head and upper part of the neck brown; the posterior part black bordered with white.

Fem. brown, with the edges of the feathers lighter; wing spot buff.

	IN.		IN.		IN.
<i>Length</i>	24	<i>Bill</i>	2	<i>Tarsi</i>	$1\frac{3}{4}$

Male: crown, throat, cheeks, and upper part of the neck, brown. with purple inflections; back part of the neck black, bordered on each side with white; fore neck and below white; speculum green, bordered above with buff, below with white; scapulars black, with light buff

edges; back and flanks white, undulated with lines of black; middle tail feathers elongated grey black; bill lead coloured; feet and tarsi reddish; eyes brown.

Fem.: smaller than the male; brown, with the edge of the feathers light; lower parts lighter; scapulars buff, margined above with lighter, below with white; middle tail feathers slightly elongated. During the summer months the male assumes nearly the plumage of the female, being ferruginous below.

<i>Anas cauducuta</i>	<i>Raii Syn.</i> 147. <i>A</i> 5.
<i>Dafila cauducuta</i>	<i>Steph. Sh. Zool.</i> 12. 127.
<i>Anas acuta</i>	<i>Linn. Syst.</i> 1. 202.
	<i>Gmel. Syst.</i> 1. 528.
	<i>Wils. Am. Orn.</i> 8. 75.
<i>Querquedula acuta</i>	<i>Selb. Illust. Brit. Orn.</i> vol. ii. 311.
<i>Trachea</i>	<i>Linn. Trans.</i> vol. 4, tab. 13, fig. 6.

A very widely distributed species, being found over Asia, Europe, America, and the north of Africa as far as the Gamba, from whence we have received them in the summer state of plumage. Breeds in Holland and Germany in morasses, building its nest of flags and grass, and laying eight eggs of a greenish blue colour.

Trachea with the tube of equal diameter to one-third of its length inferiorly, where it is slightly expanded, and again contracted above the inferior larynx, which is (except in size) similar to that of *A. Boschas*.

DAFILA MARMORATA.

Daf.—Toto corpore marmorato obscuro et subfusco.

MARBLED DUCK.

Duck, with the whole body marbled with dark and light brown.

	IN.		IN.
<i>Length</i>	18	<i>Inner toe</i>	$1\frac{3}{4}$
<i>Bill</i>	1 4-5ths	<i>Middle toe</i>	$1\frac{7}{8}$
<i>Tarsi</i>	$1\frac{5}{8}$	<i>Outer toe</i>	$1\frac{3}{4}$

The whole body beautifully marbled with light sepia brown and dusky white; the edges of the feathers white, the base bright brown except those on the breast, which have the base lighter; head slightly crested; the forehead, crown, and occiput light brown, the former barred with black; neck white, with minute longitudinal black dashes; tail ashy brown, the tip of each feather lighter; bill and legs deep lead colour.

Fuligula marmorata..... Gould, *Birds of Europe*.
Anas marmorata Temm.

Sexes not known, or the same in plumage.

Inhabits southern Europe and the adjoining portions of Asia; it is, however, rare. We suspect the male has not yet been discovered.

PÆCILONITTA ERYTHRORHYNCHA.

Pecc.—Cinereo brunnea, jugulo genisque albis; rostro coccineo.

RED-BILLED MARBLED DUCK.

Cinereous brown duck, with the throat and cheeks white; the bill red.

	IN.		IN.
<i>Length</i>	17½	<i>Tarsi</i>	1½
<i>Bill</i>	2	<i>Middle toe</i>	2

Slaty brown duck, with the bill red ; at first sight very similar to *P. Bahamensis*—the white markings on the throat and cheeks are, however, more extensive, and the bird altogether rather larger ; the tertiaries and scapulars, instead of being elongate as in *Bahamensis*, are broad and short ; the tail has all the feathers of equal length ; the wing spot is white, bounded posteriorly with buff.

Anas erythrorhyncha Gmel. Syst. 1. 517.
Lath. Ind. Orn. 2. 855.
Crimson-billed Duck..... Lath. Gen. Syn. 6. 507.

Brought home from the Cape of Good Hope by Dr. Smith, to whom we are indebted for permission to take the above description.

PÆCILONITTA BAHAMENSIS.

Pæc.—Guttur albo; speculo violaceo utrinque subflavo marginato; rostro duabus coccineis maculis lateribus.

BAHAMA MARBLED DUCK.

Throat white; wing spot metallic violet, bounded on each side by a streak of buff; the bill with two red markings on the sides.

	IN.		IN.
<i>Length</i>	$18\frac{1}{2}$	<i>Tarsi</i>	$1\frac{1}{2}$
<i>Bill</i>	$1\frac{3}{4}$	<i>Middle toe</i>	$1\frac{3}{4}$

Bill lead coloured, lighter on the sides; tail and back brown, edged with lighter; crown immaculate sepia brown; breast grey, spotted with brown; wing spot metallic violet, bounded anteriorly with a narrow stripe of buff, posteriorly with a broad one; legs and bill fuscous, the latter with a triangular patch of red on each side behind the nostrils.

Sexes similar.

In the young bird the bill is reddish brown.

<i>Anas Bahamensis</i>	<i>Linn. Syst.</i> 1. 199.
	<i>Gmel. Syst.</i> 1. 516.
	<i>Lath. Ind. Orn.</i> 2. 835.
<i>Plathera Duck</i>	<i>Catesb. Carol.</i> 1. pl. 63.
	<i>Penn. Arct. Zool.</i> 2. pl. 95.
	<i>Lath. Gen. Syn.</i> 6. 507.
<i>Mareca Bahamensis</i>	<i>Steph. Sh. Zool.</i> 12. 137.
<i>Le Marec</i>	<i>Buff.</i>

Inhabits Brazil and the Bahamas.

MARECA AMERICANA.

Mar. Mas.—Speculo viridi, utrinque atro marginato, et fasciâ latâ viridi post oculari.

Mar. Fem.—Brunnea, marginibus pennarum suturationibus.

AMERICAN WIDGEON.

Mas. speculum metallic green, broadly bounded on each side by black; a broad fasciâ of green behind the eye.

Fem. brown, the edges of each feather lighter.

	IN.		IN.		IN.
<i>Length</i>	23	<i>Bill</i>	$1\frac{5}{8}$	<i>Tarsi</i>	$1\frac{5}{8}$

Mareca chloensis

Male: has a cream-coloured patch extending from the culmen of the bill to the crown; behind the eye a green patch extending to the occiput; remainder of the head and neck cream-colour, blotched and dotted with black; breast vinaceous, the edges of the feathers lightest; back grey, narrowly barred with black and cream-colour; wing coverts and abdomen white; under tail coverts black; quills and tail grey; tertiaries grey, with the outer webs black edged with cream-colour; bill lead colour tipped with black; legs reddish; two centre tail feathers elongated.

Fem.: with the upper surface brown; each feather edged with lighter; without the green on the head, and rather smaller than the male.

<i>Anas Americana</i>	Gmel. Syst. 1. 526.
	Lath. Ind. Orn. 2. 861.
<i>American Widgeon</i>	Penn. Arct. Zool. 2. 502.
	Wils. Am. Orn. p. 86, pl. 69, f. 4.
<i>Mareca Americana</i>	Steph. Sh. Zool. 12. 135.

Inhabits America, the West Indies, and Cayenne, frequenting inland lakes and rivers, as well as the sea coast; breeds in the northern part of America in swamps, laying six or eight eggs. This bird is curious, as presenting a connecting link between the widgeon and pintail. The bill is that of the former, and the two elongated tail feathers shew an affinity to the latter.

MARECA CHILOENSIS.

Mar.—Fronte genis pteromatibus subtusque albis, strigâ post oculari purpurecente-æneâ.

CHILOE WIDGEON.

Widgeon, with the forehead, cheeks, wing coverts, and under surface white; a post ocular streak æneous-purple.

	IN.		IN.
<i>Length</i>	19	<i>Middle toe</i>	$2\frac{1}{8}$
<i>Bill</i>	$1\frac{3}{8}$	<i>Inner toe</i>	$1\frac{1}{2}$
<i>Tarsi</i>	$1\frac{1}{2}$	<i>Outer toe</i>	$1\frac{7}{8}$

Widgeon; with the forehead, cheeks, abdomen, breast, upper and under wing coverts, and upper tail coverts, white; a streak from the eye to the occiput brassy purple; neck, remainder of the head, lower part of the back, and numerous transverse bars across the breast, dark brown; tertiaries and scapulars also dark brown, with the edges of the feathers white; quills light brown; speculum velvet black; flanks white clouded with ferruginous; tail brown; legs, feet, and bill brown.

Anas Chilensis, King Zool. Proc. 1831. 15.

Inhabits the Island of Chiloe. We have also received specimens from Cayenne.

MARECA FISTULARIS.

Mar. Mas.—Vertice flavo; capite cristâque castaneis.

Mar. Fem.—Brunnea, marginibus penarum suturationibus.

WIDGEON.

Male, crown cream-coloured; head crested chesnut.

Fem. brown, with the edges of the feathers lighter.

	<small>IN.</small>		<small>IN.</small>		<small>IN.</small>
<i>Length</i>	20	<i>Bill</i>	$1\frac{1}{2}$	<i>Tarsi</i>	$1\frac{5}{8}$

Widgeon, with the crown cream-coloured; cheeks, throat, and neck chesnut, occasionally spotted with black; breast vinaceous; back white, each feather minutely barred and waved with dark grey; flanks the same; belly white; under tail coverts black; speculum metallic green, bounded by black; wing coverts white; remainder of the quills and coverts grey; bill lead coloured; legs and feet black.

Fem.: with the back fuscous grey, the edges of the feathers lighter; head and neck fuscous spotted with black; below white; remainder as in the male.

Mareca fistularis *Steph. Sh. Zool.* 12. 131.

Anas fistularis *Briss. Orn.* 6. 391, *p.* 35, *f.* 2.

Anas Penelope *Linn. Syst.* 1. 202.

Gmel. Syst. 1. 527.

Lath. Ind. Orn. 2. 860.

Widgeon *Penn. Brit. Orn.* 2. 268.

Lath. Gen. Syn. 6. 518.

Canard siffleur *Buff. Ois.* 9. 169, *pl.* 10, 11.

Trachea *Linn. Trans. vol. iv. tab.* 13, *fig.* 9.

Common in the British islands during winter, frequenting both the sea and fresh water. It is a very widely distributed species, being found throughout Africa, America, and Asia. The trachea in the male of this species, like that of several of the genera nearly allied to it, is largest immediately below the glottis, and again slightly enlarged at about one-third of its length interiorly; the inferior larynx, except in size (being smaller) greatly resembles that of *Anas Boschas*.

Anas platyrhynchos

MARECA CASTANEA.

Mar. Mas.—Castaneus atro maculato; speculoque viridi.

Mar. Fem.—Brunnea cinerea, marginibus pennarum suturationibus.

CASTANEOUS WIDGEON.

Mas. castaneous spotted with black; the wing spot green.

Fem. cinereous brown, with the edges of the feathers lighter.

	IN.		IN.		IN.
<i>Length</i>	17 $\frac{1}{2}$	<i>Bill</i>	1 3-5ths	<i>Tarsi</i>	1 $\frac{1}{2}$

Male: under parts, neck, and upper part of the back castaneous, spotted with black; lower part of the back black, each feather edged with castaneous; wing coverts brown, slightly tinged with greenish; quills brown; secondary coverts tipped with white; tail and under coverts black glossed with green and purplish; wing spot bright green; head black glossed with green; bill, legs, and feet black.

Fem.: rather smaller than the male; light brownish grey, spotted with darker; wing spot green, edged anteriorly with white, and posteriorly with black; bill and legs as in the male.

Mareca castanea nobis.

Inhabits New Holland. The first specimens we observed were in the collection of the Natural History and Antiquarian Society of Shrewsbury, presented (together with many other birds from the same country) by Major Wakeman.

AIA GALERICULATA.

Ai. Mas.—Vertice occipiteque viridibus; fasciâ latâ supra-oculari albâ; tectricibus tertiis ad apices expandentibus.

Ai. Fem.—Faciis duabus alis.

MANDARIN DUCK.

Male, with the crown and occipital crest green; a broad white band over the eye; tertiaries expanded at the tip, and prominent.

Fem. with two white bars on the wing

	IN.		IN.
<i>Length</i>	18 $\frac{1}{2}$	<i>Inner toe</i>	1 $\frac{3}{8}$
<i>Bill</i>	1 $\frac{1}{8}$	<i>Middle toe</i>	1 $\frac{7}{8}$
<i>Tarsi</i>	1 $\frac{1}{2}$	<i>Outer toe</i>	1 $\frac{3}{4}$

Male: head with a long occipital crest and crown purplish green; above the eye a broad band of white; throat, cheeks, and remainder of the neck bright reddish ferruginous, the shaft of each feather lighter; breast bright purple; back, tail, and wings hair brown; scapulars with the outer webs black; the inner ones white; one tertiary prolonged and expanded on each side, prominent over the back; on the lower part of the breast a broad band of black not meeting in front, on which there are one or two narrow ones of white; flanks ferruginous, beautifully undulated with narrow bands of black; belly, vent, and under tail coverts white; bill and legs orange.

Fem.: similar to the summer duck *A. sponsa*, but may be distinguished from it (according to Latham) by having two bars of white on the wing, and by the white spots on the breast being round instead of triangular,

<i>Anas galericulata</i>	<i>Linn. Syst.</i> 1. 206.
	<i>Gmel. Syst.</i> 1. 539.
	<i>Lath. Ind. Orn.</i> 2. 871.
<i>Querquedula sinensis</i>	<i>Briss. Orn.</i> 6. 450.
<i>Chinese Teal</i>	<i>Edw. Gla. pl.</i> 102.
	<i>Lath. Gen. Syn.</i> 6. 548.
<i>Sarcelle de la Chine</i>	<i>Buff. Ois.</i> 9. 276.

Inhabits China and Japan, where it is kept (as in this country), for the sake of its beauty, in a state of domestication. Nothing is known of its habits in a wild state.

AIA SPONSA.

Ai. Mas.—Capite colloque viridibus; gutture albo.

Ai. Fem.—Brunnea, rostro cinguloque occulari etiam ad basin rostri, et fasciâ alis albis.

SUMMER DUCK.

Mas. head and upper part of the neck metallic green; throat white; crest long.

Fem. dark brown, with the belly, space round the eye, and bill white; one white band across the wing.

	IN.		IN.
<i>Length</i>	20	<i>Tarsi</i>	1 $\frac{3}{8}$
<i>Bill</i>	1 $\frac{3}{8}$	<i>Middle toe</i>	2

Crown and neck and occipital crest glossy green; line extending from the bill over the eye, another from the lower part of the eye to the occiput, together with the throat, white; neck and upper part of the breast purplish, and sprinkled with spots of white; scapulars, wing coverts, and tail black, beautifully glossed in some lights with green and purple inflections; on the lower part of the breast is a white bar, posteriorly to which is a black one; flanks yellowish ferruginous, barred and undulated with dusky; under tail coverts and quills sooty black; below white; bill and legs orange.

Fem.: dusky slate brown, with a space round the eye, a band round the bill, the throat, and belly white; breast dusky, marked with triangular spots of white; quills and tail as in the male; legs and bill dusky.

<i>Aia sponsa</i>	<i>Boie Isis.</i> 1826.
<i>Dendronessa sponsa</i>	<i>Faun. Bo. Am.</i> 2. 446.
<i>Anas sponsa</i>	<i>Linn. Syst.</i> 1. 207.
	<i>Gmel. Syst.</i> 1. 539.
	<i>Lath. Ind. Orn.</i> 2. 871.
<i>Anas Aestiva</i>	<i>Briss. Orn.</i> 6. 351. <i>pl.</i> 32, <i>fig.</i> 2.
<i>Summer Duck</i>	<i>Catesb. Carol.</i> 1. <i>pl.</i> 97.
	<i>Edw. Gla.</i> p. 101.
	<i>Lath. Gen. Syn.</i> 6. 546.
<i>Beau canard, Huppe</i>	<i>Buff. Ois.</i> 9. 254.
<i>Trachea</i>	<i>Mont. Orn. die, Sup.</i> <i>fig.</i> 5, <i>et nobis.</i>

Inhabits America, from Hudson's Bay to Mexico and the Antilles, throughout the whole of which space it breeds, seldom frequenting the sea. It generally makes its nest in decayed trees hanging over the water, laying twelve or thirteen eggs.

This species is often kept in confinement in this country, and breeds freely, as well as its congener. The inferior larynx is that of the true wading duck, and much resembles in miniature that of the widgeon; it is, however, more expanded anteriorly, the two ends of the bulb being of equal size.

QUERQUEDULA JAVANENSIS.

Quer.—Capite viridi, maculâ ad basum rostri; gutture et fasciâ occipitali albis.

JAVA TEAL.

Head green, with a spot at the base of the bill; throat and patch extending to the occiput white.

	IN.		IN.
<i>Length</i>	18	<i>Tarsi</i>	$1\frac{3}{4}$
<i>Exp. al</i>	27	<i>Middle toe</i>	2
<i>Bill</i>	$1\frac{1}{2}$		

Head bright glossy green; spot behind the culmen of the bill, throat, and streak towards the back of the neck from the latter white; collar white; breast with each feather beautifully marked with lunules of black; vent coverts white undulated with black; wing spot purplish green, above which is a conspicuous spot of velvet black; tertiaries very long, with the shaft and a narrow edging exteriorly white, and the edge of the posterior or inner web grey; quill and tail feathers grey; upper and under tail coverts black; vent buff, anteriorly to which there is a band of black; feet and bill black.

Anas Javanensis..... Veill. *En. Method. Orn.* 1. 160.

Sarcelle de l' Ile de Java..... Buff. *Ois. tom. xvii.* p. 395.

Inhabits Java. I have never seen but one specimen of this beautiful species; it is in Lord Derby's collection at Knowsley, and has been exhibited at one of the Zoological Society's evening meetings.

QUERQUEDULA GLOCITANS.

Quer. Mas.—Capite viridi, maculâ rotundatâ castaneâ ante oculari et post oculari ornato.

Quer. Fem.—Brunnea, marginibus pennarum suturatoribus.

BIMACULATED TEAL.

Male Teal, with the head green, ornamented with a rounded chesnut patch before and behind the eyes.

Fem. brown teal, with the margin of the feathers lighter.

	IN.		IN.		IN.
<i>Length</i>	15	<i>Bill</i>	1 $\frac{7}{8}$	<i>Tarsi</i>	1 9-20ths

Male: head and neck green, with a dark chesnut patch before and behind the eyes; the throat approaching to black; back and flanks white, undulated with black lines; breast and abdomen ferruginous, with a few black spots; wing coverts fuscous; speculum velvet black, bordered anteriorly with violaceous, posteriorly with white; quills brown, margined with lighter; bill, feet, and legs lead-coloured; the webs black.

Fem.: slightly smaller than the male; head and neck light ferruginous, with the crown and base of each feather darker; abdomen white; back and flanks fuscous ferruginous, with the base of each feather brown.

Anas glocitans..... Pall. *Acta Stock.* 1. 779, 40, tab. 33, fig. 1.

Gmel. Syst. 1. 526.

Lath. Ind. Orn. 862.

Querquedula glocitans..... Vig. *Lin. Trans.* vol. xiv. 559.

Gould, Birds of Europe.

Bimaculated Teal..... Penn. *Brit. Zool.* 2. 602.

Cunard Glousseur..... Veill. *En. Method. Orn.* 1. 126.

Said to inhabit Siberia and the neighbourhood of Lake Baikal. One or two specimens only have been met with in England.

QUERQUEDULA CRECCA.

Quer. Mas.—Capite colloque castaneis, fasciâ occipitali viridi.

Quer. Fem.—Brunnea, marginibus pennarum saturatioribus; subtus alba.

COMMON TEAL.

Teal, with the head and neck chesnut, with a green occipital fascia.

Fem. brown teal, with the margin of the feathers lighter; below white.

	IN.		IN.
<i>Length</i>	$14\frac{1}{2}$	<i>Tarsi</i>	1 1-6th
<i>Bill</i>	1 5-12ths	<i>Middle toe</i>	$1\frac{5}{8}$

Male: head, cheeks, and neck chesnut; from the eye to the occiput passes a band of metallic green, bordered by a stripe of yellowish; below white; the breast dotted with black; the scapulars, back, and flanks white undulated with black; wing coverts brown; quills hoary fuscous; speculum metallic green, bounded above and below by velvet black; lateral tail coverts with a band of black.

Fem.: brown, with the edge of the feathers lighter; below yellowish white; speculum green, without the black edging; crown brown.

<i>Querquedula crecca</i>	<i>Steph. Sh. Zool.</i> 12. 146.
<i>Anas erecca</i>	<i>Linn. Syst.</i> 1. 264.
	<i>Gmel. Syst.</i> 1. 532.
	<i>Lath. Ind. Orn.</i> 2. 872.
<i>Common Teal</i>	<i>Penn. Arct. Zool.</i> 2. 577.
<i>Green-winged Teal</i>	<i>Wils. Am. Orn.</i> 8. 106, pl. 70. fig. 40.
<i>Petite sarcelle</i>	<i>Buff. Ois.</i> 9. 265.
<i>Canard sarcelle d' Hiver</i>	<i>Temm. Man. d' Orn.</i> 2. 846.
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 12, fig. 1.</i>

Breeds in most marshy lakes in the north of Europe, America, Asia, and the British Isles; migrates southwards during winter, when it frequents almost entirely fresh water. Mr. Selby, in his "Illustrations of British Ornithology," mentions that he is inclined to think that our indigenous broods seldom quit the immediate neighbourhood of the place in which they were bred. The nest is formed of grass, and it lays from eight to ten eggs of a whitish colour.

Trachea with the tube largest immediately below the glottis, then contracted, then expanded, and gradually lessening to the inferior larynx, which is similar in form to that of *Anas Boschas*.

QUERQUEDULA CAROLINENSIS.

Quer.—Precedenti similis, sed distingueri potest fasciâ albâ ante alis.

AMERICAN TEAL.

Quer. similar to *Q. crecca*, but distinguished from it by a white bar on each side of the breast.

	^{18.}		^{18.}		^{18.}
<i>Length</i>	15	<i>Bill</i>	1 $\frac{5}{8}$	<i>Tarsi</i>	1

Precisely similar to *Q. crecca*, with the exception of the under tail coverts, which are black, and in having a white band on each side opposite the point of the wing.

<i>Anas Carolinensis</i>	<i>Gmel. Syst. Nat.</i> 1, 533.
	<i>Lath. Ind. Orn.</i> 2, 874.
	<i>Jard. Illus. Orn. pl.</i>
<i>American Teal</i>	<i>Penn. Arct. Zool.</i> 2, 584.

Inhabits America, where it takes the place of the preceding species.

QUERQUEDULA FORMOSA.

Quer.—Vertice, gutture, fasciâque ab hoc ad oculos atris, ntrinque albo marginato.

BAIKAL TEAL.

Teal, with the crown, throat, and a streak proceeding from the latter to the eye black, bounded on each side with white.

	^{18.}		^{18.}
<i>Length</i>	15 $\frac{1}{2}$	<i>Tarsi</i>	1 $\frac{3}{8}$
<i>Bill</i>	1 5-12ths	<i>Middle toe</i>	1 4-5ths

Male: crown and throat black, with each feather slightly tipped with ferruginous brown; from the latter proceeds a black waved streak to the eye, bounded on each side by a narrow one of white, and enclosing a patch of fawn-colour; from the bill through the eye proceeds a broad band of glossy green to the occiput, where it is broadest, and trifurcate below, the centre branch extending down to the back of the neck, the lateral ones curve to the fore neck, but do not meet, the whole bounded by a narrow edging of white; fore part of the neck bright reddish fawn colour, spotted with black; back brown; lower part of the neck behind black, thickly irrorated with white; flanks the same; belly white; quills and coverts brown; middle coverts tipped with brown; pri-

mary coverts glossy purplish black, tipped with white; tertiaries long, outer webs black edged with ferruginous; inner webs light fawn colour; tail brown; tail coverts the same, edged with lighter; under tail coverts black, having on the sides anteriorly a broad band of white; the under coverts bounded laterally with a bar of ferruginous; bill brown; legs light yellow; wing spot bounded anteriorly by brown, posteriorly by white.

Fem.: unknown.

Querquedula formosa Steph. Sh. Zool. vol. 12. 151.

Anas formosa Gmel. Syst. Nat. 1. 523.

Lath. Ind. Orn. 2. 876.

Baikal Teal Lath. Gen. Syn. 6. 557.

Inhabits Russia and Lake Baikal. (*Lath.*) Specimens are in the museum of the Zoological Society, brought over by R. Reeves, Esq. in 1832, from China; and also in our own. We have also very lately succeeded in obtaining the trachea, of which a figure is given: though that of a true teal, it is remarkable for the exceeding small size of the larynginal bulb, and also for a slight approach it makes in form to that of the next genus, *Cyanopterus*.

QUERQUEDULA MANILLENSIS.

Quer.—Genis albis, fasciâ brunneâ occipite ad collum inferiorem tendente.

MANILLA TEAL.

Teal, with the cheeks dirty white, with a streak from the occiput to the lower part of the neck, when it expands into a collar, brown.

	IN.		IN.
<i>Length</i>	15	<i>Tarsi</i>	$1\frac{3}{8}$
<i>Bill</i>	2	<i>Middle toe</i>	$1\frac{5}{8}$

Allied to *Q. formosa*; tertiaries not acuminate, and ferruginous; breast as in the last; upper parts of the back olive brown; tail and rump black glossed with green; belly white irrorated with black; middle vent coverts black; lateral coverts white; secondary quills with the outer webs brassy green; coverts white tipped with black; small coverts black; quills brown; head, neck, and crown dark brown; cheeks dirty white; from the occiput proceeds a streak of brown to the lower part of the neck, when it expands into a collar; bill lead colour; feet and legs light orange.

Anas Manillensis..... Gmel. Syst. Nat. 1. 523.

Lath. Ind. Orn. 2. 876.

Sarcelle de l' Isle de Lucon..... Sonn. Voy. Ind. 91, pl. 54.

Manilla Teal..... Lath. Gen. Syn. 6. 557.

Inhabits the Manillas. Specimens are in our collection, and in that of the Earl of Derby.

QUERQUEDULA FALCARIA.

Quer. Mas.—Capite brunneo; cingulo collari guttateque purpureis.

Quer. Fem.—Brunnea, marginibus pennarum suturationibus.

FALCATED TEAL.

Teal, with the head deep brown; ring round the neck and throat purple.

Fem. brown teal, with the margins of the feathers lighter.

	IN.		IN.
<i>Length</i>	17	<i>Tarsi</i>	1½
<i>Bill</i>	1½	<i>Middle toe</i>	2

Male: head and crown deep brown; ring round the neck and throat brassy purple; remainder of the neck and a spot behind the upper part of the base of the bill white; belly, breast, flanks, and upper part of the back white, each feather lunulated with ashy black; tail ashy; upper tail coverts and centre of the under ones black, laterally buff, anteriorly to which there is a black streak, then a patch of white; quills ashy brown; tertiaries long, inner webs brown grey, outer glossy green edged with light grey, the edging broader at the base; wing spot green; primary coverts light ashy brown, lightest on the edge.

Fem.: about half an inch shorter than the male; brown, each feather lunulated with darker; crown dark brown; neck light brown, streaked with dark brown; tertiaries long; wings and tail as in the male; coverts brown; belly light; legs and bill in both sexes lead coloured.

Anas falcaria *Gmel. Syst. Nat.* 8. 521.

Lath. Ind. Orn. 2. 860.

Falcated Duck *Penn. Arct. Zool.* 2. 574.

Lath. Gen. Syn. 6. 516.

Anas falcaria *Steph. Sh. Zool. vol. xii. p.* 100.

The specimens from which the above descriptions were taken are in the British Museum, and were brought home by R. Reeves. Esq. from China.

QUERQUEDULA ERYTHORRHYNCHA.

Quer.—Brunnea, ptilis atris, tectrices majores speculoque æneis.

CRIMSON-BILLED DUCK.

Brown, with the upper coverts velvet black, the greater ones and speculum brassy green.

	IN.		IN.		IN.
<i>Length</i>	18	<i>Bill</i>	$1\frac{1}{2}$	<i>Tarsi</i>	$1\frac{1}{4}$

Above brown, with the edges of the feathers brighter; throat inclining to grey; back, lesser wing coverts, and tail velvet black; below the upper tail coverts and flanks grey, the latter with two or three deep brown spots; the breast sometimes slightly waved with brown; the remainder of the wing beautiful æneous green, with the exception of the two outer quill feathers, which are black, and the terminal half of the secondaries, which are white; bill and legs bright red in the male, fuscous in the female; bill brown, with a red patch on each side at the base.

Anas Erythrorhyncha *Spir. N. Sp. avium*.

Inhabits Southern America.

We had an opportunity of dissecting this bird through the kindness of the Earl of Derby, who had a pair alive for some time.

The trachea and inferior larynx is that of a typical teal. It may, however, be distinguished from that of *Q. crecca* (which it in other respects resembles) by the bulb being more oblique, and swelling more in an upward direction.

The tube of the trachea in both sexes is acted upon by two pairs of muscles of voice, the first pair the usual sterno-tracheal ones; in the female the second pair diverge from the tube, where it penetrates the cavity of the chest, and are inserted into the membrane stretched between the *rami* of the *os furcatum* and the *coracoids*. In the male they diverge similarly, and are inserted in the *rami* of the *os furcatum*, having also a slight attachment along the internal surface of the membrane stretched between the *rami* of the *os furcatum*, which is not the case in the female.

QUERQUEDULA CRECCOIDES.

Quer.—Pallide brunnea-grisca, sparsa notataque dorso imo ptilisque fasciis hic apice rufo, speculo nigro, fulvo marginato.

MOTTLED TEAL.

	IN.		IN.		IN.
<i>Length</i>	$16\frac{1}{2}$	<i>Bill</i>	$1\frac{3}{4}$	<i>Tarsi</i>	$1\frac{1}{4}$

The head and neck of this bird are finely marked with fuscous dots, those of the neck being mostly in transverse lines, thus forming several undulating fascia; the feathers of the upper part of the back are of a brownish grey, and are marked in the centre by oval fuscous spots; the scapular feathers are similarly marked by long central black strigæ, which in some lights are of a beautiful green; the lower part of the back is of a uniform fuscous colour; the upper wing coverts are fuscous, the apices of the lowest range of feathers being also terminated with rufous, which colour forms a narrow fascia across the wing immediately over the speculum; the external secondary quill feathers are black on their outside margin, and are terminated with light buff; the internal webs are fuscous on their outside webs, with a shade of deep green, and slightly margined with buff; the breast and middle of the abdomen are whitish, the feathers of the former being strongly marked with round black spots in the centre, those of the latter being more strongly marked, so as to exhibit irregular fasciæ; bill yellow, with a black apex.

Anas Creccoides, King *Zool. Jour.* 4. 99.

Inhabits the Straights of Magellan: a specimen is in the museum of the Zoological Society.

QUERQUEDULA CAPENSIS.

Quer.—Rostro coccineo; marginibus capituloque basali atris.

CAPE TEAL.

Teal, with the bill crimson; the margins and a band round the base black.

	IN.		IN.
<i>Length</i>	$18\frac{1}{2}$	<i>Tarsi</i>	$2\frac{1}{2}$
<i>Bill</i>	$1\frac{1}{2}$	<i>Middle toe</i> . . .	$2\frac{3}{4}$

Light brown teal, with all the dorsal aspect with the feathers edged with lighter; those on the breast and under parts very light brown, with a lunule of darker on each feather; the throat

white; the head and neck spotted, and minutely streaked with dark and light brown; wing spot bright green, margined on every side but the upper with a band of white; wing coverts, quills, and tail blue slate-coloured, the latter with each feather edged with much lighter; legs reddish brown.

Female similar, but rather smaller.

Querquedula capensis Smith, *Cat. S. Afric. Mus.*

Inhabits Southern Africa, and brought home by Dr. Smith, by whom it will be figured in his forthcoming work on South African Zoology.

QUERQUEDULA HOTTENTOTA.

Quer. Mas.—Brunneus, marginibus penarum suturationibus; tectricibus caudæ juguloque pallidé brunneis, illo atro fasciato.

Quer. Fem.—Mari similis, sed tectricibus caudæ toto corpori similibus.

HOTTENTOT TEAL.

Male, brown teal, with the margins of the feathers lighter; the tail coverts and throat light brown; the latter fasciated with black.

Fem. like the male, but with the quill feathers like the rest of the body.

	IN.		IN.
<i>Length</i>	14½	<i>Tarsi</i>	1
<i>Bill</i>	1½	<i>Middle toe</i>	1½

Male; umber brown, with the edges of the feathers lighter; the crown, occiput, and quills dark brown; chin, throat, cheeks, rump, and under tail coverts light brown, the latter narrowly barred with darker; breast and lower part of the neck and under surface darker than on the rump; the abdomen barred with black; speculum and secondaries bright brassy green, the former edged posteriorly with black, then with white; coverts brown, slightly glossed with brassy; tail dark brown.

Fem.: similar to the male, but with the markings generally not so distinct, and the whole plumage below lighter; legs in both sexes brown; bill lead coloured, with the nail horn colour.

Querquedula Hottentota Smith, *Cat. S. Afric. Mus.*

Brought home by Dr. Smith from the western coast of South Africa, near the Orange river: a specimen is also in our collection, but we are not aware from what locality.

CYANOPTERUS CIRCIA.

Cyan. Mas.—Undulatus brunneo et albo; ptilis azureis-cinereis.

Cyan. Fem.—Brunnea, marginibus pennarum suturationibus.

GARGANEY TEAL.

Male teal, undulated with brown and white; lesser coverts blueish grey.

Fem. brown teal, with the margin of the feathers lighter; coverts as in the male.

	IN.		IN.
<i>Length</i>	$16\frac{1}{2}$	<i>Tarsi</i>	$1\frac{1}{10}$ th
<i>Bill</i>	$1\frac{1}{2}$	<i>Middle toe</i>	$1\frac{1}{2}$

Male: crown, back of the neck, cheeks, and sides of the neck brown, the latter inclining to chesnut in some specimens; a white stripe from the bill under each eye; lower part of the neck and breast whitish, marked with brown lunules; back brown, with the edges of the feathers lighter; scapulars brown, with a white stripe down the shaft of each; wing coverts blue grey; the rump and tail brown; below white, undulated with dark brown on the flanks; speculum green; bill, tarsi, and feet lead coloured; eyes brown.

Fem.: brown, each feather edged with lighter; throat white; below yellowish white; wings as in the male.

<i>Querquedula circia</i>	<i>Steph. Sh. Zool.</i> 12, 143.
<i>Anas circia</i>	<i>Linn. Syst.</i> 1, 201.
	<i>Gmel. Syst.</i> 1, 553.
	<i>Lath. Ind. Orn.</i> 2, 873.
<i>Anas Querquedula</i>	<i>Linn. Syst.</i> 1, 263-2.
	<i>Gmel. Syst.</i> 1, 531.
	<i>Raii Syn.</i> 148, 8.
<i>Garganey</i>	<i>Penn. Aret. Zool.</i> 2, 576.
	<i>Lath. Ind. Orn.</i> 6, 550.
<i>Le Sarcelle d'ete</i>	<i>Buff. Ois.</i> 9, 208.
<i>Le Sarcelle commun.</i>	<i>Buff. Ois.</i> 9, 260.
<i>Trachea</i>	<i>Linn. Trans. vol.</i> 4, <i>tab.</i> 13, <i>fig.</i> 2—3.

Found over the whole of Asia, the northern part of Africa, and Europe; visits the British isles in May, frequenting fresh water; breeds in Holland; its nest is constructed of grass placed by the water side; lays from ten to twelve white eggs.

Trachea with the tube smaller about an inch below the glottis, then slightly swelled, and again slightly contracted about one inch above the inferior larynx, from which point it gradually expands to the end. The inferior larynx with an osseous bulb extending anteriorly and on both sides, and not swelling out so abruptly as in the preceding genus or among the true teal.

CYANOPTERUS FRETENSIS.

Cyan. Mas.—Gulá colloque anteriore pallide vadiis; dorso, abdomineque, imis caudaque. albis, nigro fasciatis; ptilis cæruleis.

Cyan. Fem.—Gula colloque ut in mare; dorso imo ptilisque atro-cæruleis; abdomine ferrugineo, sparso et fasciato atro.

FASCIATED TEAL.

	IN.		IN.
<i>Length</i>	14	<i>Inner toe</i>	1 1-6th
<i>Bill</i>	1 $\frac{1}{2}$	<i>Middle toe</i>	1 $\frac{2}{3}$
<i>Tarsi</i>	1	<i>Outer toe</i>	1 5-12ths

Male Teal, with the back, breast, fore part of the neck, and cheeks light yellowish brown, the two former having in the centre of each feather a dark brown patch; flanks, abdomen, tail, and rump white, the former broadly and the latter narrowly barred transversely with black; vertex as far down as the eyes, and a streak down the back of the neck, deep brown; feet, legs, and bill blueish brown, the latter with a yellow patch on each side of the bill, extending beyond the nostrils; wing coverts slaty blue, the greater ones tipped with white; secondaries æneous green, also tipped with white; quills and tertiaries (which are long) dusky, the latter with the centre of each feather darker.

Female and young similar to the male, but with the abdomen and flanks ferruginous, irregularly barred and irrorated with dark brown; rump black; the bill without the yellow markings on the sides.

Anas fretensis, King *Zool. Proc. Dec. 14, 1830.*

Inhabits the Straits of Magellan and other localities on the western coast of South America.

CYANOPTERUS DISCORS.

Cyan.—Capite cinereo brunneo, lunulá albá ante oculos.

BLUE-WINGED TEAL.

Teal, with the head ashy brown; a large white lunule before the eye.

	IN.		IN.		IN.
<i>Length</i>	16	<i>Bill</i>	1 $\frac{1}{2}$	<i>Tarsi</i>	1 $\frac{3}{4}$

Head ashy brown, slightly glossed at the tip of each feather with brassy; crown darkest; between the eye and bill, and partially surrounding the former, a large lunate spot of white, the

horns projecting backwards; breast and neck ferruginous, beautifully spotted with dark brown; quills, tail, and back dark brown, the latter slightly lunated with ferruginous; under tail coverts deep black or brown, on the sides a patch of white; wing coverts bright blue, lower ones marked with a streak of white, then with one of green; anterior tertiaries with the outer webs blue; on the inner side of the shaft a streak of ferruginous; the remainder black.

Female: length 15 inches; wing coverts as in the male; tertiaries and remainder brown lunated with ferruginous; crown darker.

<i>Querquedula discors</i>	<i>Steph. in Sh. Zool. vol. 12.</i>	149.
	<i>Linn. Syst. Nat.</i>	1. 205.
	<i>Lath. Ind. Orn.</i>	2. 854.
<i>Sarcelle soucrourou</i>	<i>Buff. Pl. Enl.</i>	966, <i>mas.</i>
— <i>soucrouette</i>		403, <i>fem.</i>
<i>White-faced Duck</i>	<i>Catesb. Carol.</i>	1. 100, <i>mas.</i>
<i>Blue-winged Teal</i>		1. 99, <i>fem.</i>

Inhabits the American continent, from the 58th parallel to Guiana and the West Indies.

CYANOPTERUS RAFFLESII.

Cyan.—Castaneo-ferrugineus, speculo viridi.

RAFFLES'S TEAL.

Deep reddish chesnut teal, with the speculum green.

	IN.		IN.
<i>Length</i>	18	<i>Tarsi</i>	$1\frac{1}{4}$
<i>Bill</i>	2	<i>Middle toe</i>	2

Deep reddish chesnut teal, with the crown and vent inclining to dark brown; each of the feathers on the back with black lunules; tertiaries black and blue, with the shaft buff; wing coverts light blue; speculum metallic green; quills and tail black.

Anas Rafflesii, *King*

Zool. Journ. 4. 97.

First brought home by Capt. King from Chili; since which time we have received specimens from the same locality. Said by navigators to be exceedingly common on the west of South America.

RHYNCHASPIS RHYNCHOTIS.

Rhyn. Mas.—Ferrugineus; capite colloque superiore cæruleis; tectricibus caudæ nigris; speculo viridi.

Rhyn. Fem.—Brunnea, marginibus pennarum suturationibus; rostro pallido.

NEW HOLLAND SHOVELLER.

Male shoveller, with the head and upper part of the neck blue; under and upper tail coverts black; wing spot green.

Fem. shoveller, brown, each feather edged with lighter; bill horn colour.

	IN.		IN.
<i>Length</i>	20	<i>Tarsi</i>	$1\frac{1}{4}$
<i>Bill</i>	1 2-5ths	<i>Middle toe</i>	$1\frac{1}{8}$

Male: head and neck, lesser coverts, and tertiaries light blue, some of the latter with a stripe on the inner side of the shaft white; bar behind the base of the bill on each side white; centre of the back, quills, and tail white, with a lunule near the tip and two spots on the shaft of each feather brown; wing spot bright brassy green, anterior to which is a white stripe; flanks, belly, and breast deep ferruginous, the former barred and dotted, and the latter undulated with brown; above the thighs on each side a white spot; under and upper tail coverts black; under wing coverts white; top of the head and back of the neck darkest; bill black; legs orange.

Fem.: brown, with the tip of each feather lighter; below lightest; lesser coverts slightly inclining to blue; bill and legs horn colour in the dried specimen.

Young: male nearly intermediate between the two; bill black; legs orange.

Anas Rhynchotis *Lath. Ind. Orn. App. 1.*

Rhynchaspis Rhynchotis *Steph. in Sh. Zool. 12. 123.*

New Holland Shoveller *Lath. Gen. Syn. Sup. 359.*

Inhabits New Holland, and appears to be identical with specimens received from South America.

The bird described by Latham under the name of *Anas Rhynchotis* is the young. It is possible also that the white markings mentioned as present in our specimens of the old male, behind the base of the bill and down the fore neck, are not constant, but merely appear during the moult.

RHYNCHASPIS MACULATUS.

Rhyn.—Precedenti similis. sed corpore flavescens-ferrugineo, nigris maculis rotundatis ornatis; capite simili, nigro sparso.

SPOTTED SHOVELLER.

Shoveller like the preceding, but with the body yellowish-ferruginous, adorned with rounded black spots; head the same, sprinkled with black.

	IN.		IN.
<i>Length</i>	20	<i>Tarsi</i>	$1\frac{1}{2}$
<i>Bill</i>	2 3-10ths	<i>Middle toe</i>	2 1-10th

We have considerable doubts as to the validity of this species. The marks of distinction between it and *R. Rhynchotis* are the following: the nail on the bill is broader, the flattened space between the nostrils and behind them is also broader; the whole body is of a lighter colour, and spotted in the centre of each feather with round dots; the white spot behind the thighs is similar, but the admeasurements slightly differ. We suspect that this bird will eventually turn out to be the female of *Rhynchotis*; and that the birds we have described as female and young of that bird will turn out to be both young males.

Rhynchaspis maculatus Gould in *Jard. and Selb. Illus. Orn.* p. 147.

Mr. Gould received his bird from South America; ours was received from New Holland.

RHYNCHASPIS CLYPEATA.

Rhyn. Mas.—Capite colloque viridibus; ptilis cæruleis.

Rhyn. Fem.—Brunnea, marginibus pennarum suturatoribus.

COMMON SHOVELLER.

Male shoveller, with the head and neck green; the wing coverts blue.

Fem. shoveller, brown, with the margins of the feathers lighter.

	IN.		IN.
<i>Length</i>	20	<i>Tarsi</i>	$1\frac{3}{8}$
<i>Bill</i>	$2\frac{2}{3}$	<i>Middle toe</i>	2

Male: head and neck glossy metallic green; throat black; breast, smaller scapulars, and greater wing coverts white; larger scapulars and lesser wing coverts blue, the former with a light

stripe down the centre of each feather; speculum green; quills and tail brown; the outer tail feathers elongated, pointed; below chestnut; bill fuscous red.

Fem.: brown, having the edges of all the feathers lighter; wing as in the male.

<i>Rhynchaspis clypeata</i>	<i>Steph. Sh. Zool.</i> 12. 115.
<i>Anas clypeata</i>	<i>Linn. Syst.</i> 1. 200. 19.
	<i>Gmel. Syst.</i> 1. 518.
	<i>Lath. Ind. Orn.</i> 2. 856.
<i>Spathulea clypeata</i>	<i>Flem. Brit. An.</i> 123.
<i>Anas platyrhynchos</i>	<i>Rail Syn.</i> 144. 13.
<i>Shoreller</i>	<i>Penn. Arct. Zool.</i> 2 N. 489.
	<i>Penn. Brit. Zool.</i> 2. 596.
<i>Red-breasted Shoreller</i>	<i>Penn. Brit. Zool.</i> 2. 597.
<i>Souchet ou le rouge</i>	<i>Buff. Ois.</i> 9. 191.
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 13, fig. 4, 5.</i>

Was found in the Deccan by Col. Sykes: one specimen has been brought home by Dr. Smith from South Africa. Generally inhabits Asia, Europe, America, and North Africa; breeds in Holland, and has occasionally bred in Norfolk, laying twelve eggs; little, however, is known of its nidification.

Trachea with the tube of nearly equal size; the inferior larynx bulbous, bulb projecting both before and on the left side, anteriorly slightly acuminate; the bronchiæ far apart.

RHYNCHASPIS CAPENSIS.

Rhyn.—Brunneus; subtus singulis pennis lunulâ centrali, apiceque leviter ferrugineis.

CAPE SHOVELLER.

Brown shoveller; below with a lunule, and the tip of each feather light ferruginous.

	IN.		IN.
<i>Length</i>	22½	<i>Tarsi</i>	1½
<i>Bill</i>	2¾	<i>Middle toe</i>	2½

Head and neck light brown, minutely spotted with dark brown; crown and slightly elongated feathers on the occiput darker; below dark sepia brown, each feather having its tip and a transverse lunule in the centre light brownish buff; the back brown, slightly glossed with greenish; greater quills and tail deep brown, the latter edged with lighter; secondaries with the outer webs metallic green; tertiaries with the outer webs blue; lesser wing coverts azure, the greater ones white, forming a band of that colour across the wing; bill black; legs reddish orange.

Rhynchaspis Capensis

Smith, Cat. S. African Mus.

We are indebted to Dr. Smith for allowing us to take the above description from the specimens exhibited in the South African museum, of which he intends shortly to publish drawings.

MALACORHYNCHUS MEMBRINACEUS.

Mal. Mas.—Cinereus, maculâ caryophyllaceâ post oculari.

Mal. Fem.—Mare similis, sed macula post ocularis caret.

MEMBRINACEOUS DUCK.

Male, ashy, with the side of the bill towards the tip membrinaceous ; a pink spot behind the eye.

Fem. similar, but without the pink spot behind the eye.

	IN.		IN.		IN.
<i>Length</i>	16½	<i>Bill</i>	2¾	<i>Tarsi</i>	1½

Male: grey, transversely striped with brownish black ; stripes broadest on the flanks ; neck brown ; a stripe round the eye and down the back of the neck dark brown ; a little behind the eye a bright pink spot ; back brown, transversely irrorated with lighter ; rump black ; tail coverts white ; tail brown ; under tail coverts fawn colour ; quills dark brown ; coverts light brown, except the tip of the primary coverts, which are white, forming a white wing spot ; bill and legs brown, the former largely lobated.

Fem. : as above.

<i>Anas malacorhynchus</i>	<i>Gmel. Syst.</i> 1. 526.
	<i>Lath. Ind. Orn.</i> 2. 862.
<i>Anas membrinacca</i>	<i>Lath. Ind. Orn. App.</i> 11.
<i>Anas fasciata</i>	<i>Shar. N. Mis.</i> 17 pl. 697.
<i>Soft-billed Duck</i>	<i>Lath. Gen. Syn.</i> 6. 522.
<i>New Holland Duck</i>	<i>Lath. Gen. Syn. Sup.</i> 11. 359.
<i>Rhynchaspis malacorhynchus</i>	<i>Steph. Sh. Zool.</i> 12. 123.
————— <i>membrinacca</i>	<i>Steph. Sh. Zool.</i> 12. 121.

Inhabits New South Wales.

KTINORHYNCHUS STREPERA.

Kti. Mas.—Cinereus et albus; speculo atro; tectricibus castaneis.

Kti. Fem.—Brunnea, marginibus pennarum suturationibus.

GADWALL.

Male, grey and white; speculum velvet black; primary coverts chesnut.

Fem. brown, with the margins of the feathers lighter.

	IN.		IN.		IN.
<i>Length</i>	21	<i>Bill</i>	$1\frac{3}{4}$	<i>Tarsi</i>	$1\frac{1}{2}$

Male: head and neck brownish, marked with dots of grey brown; upper part of the breast and lower part of the neck grey brown, each feather marked with a lunule of white; back the same, thickly barred with white; tertiaries grey, the edges light; speculum black, anterior to which there is a reddish chesnut patch formed by the greater primary coverts, posteriorly one of white; lower part and under tail coverts black; quills and tail grey; bill black; legs fuscous.

Fem.: with the speculum as in the male; remainder of the plumage as in the female of the common wild duck.

Chauliodus strepera Gould, *European Birds*.

Anas strepera Linn. *Syst.* 1. 200.

Gmel. Syst. 1. 520.

Lath. Ind. Orn. 2. 859.

Ktinorhynchus strepera Nobis. (*See preface.*)

Gadwall Penn. *Arct. Zool.* 2. 575.

Chipeau Buff. *Ois.* 9. 187, *pl.* 12.

Trachea Linn. *Trans.* vol. 4, *tab.* 13, *fig.* 8.

A very widely distributed species. We have received specimens from India, Africa, and America. It is also enumerated in the British lists, and is not uncommon in Holland, where it breeds in marshes among rushes, laying eight or nine ashy green eggs.

Tube of the trachea enlarged about one-third of its whole length from the lower extremity, and again contracted above the inferior larynx. Inferior larynx with a projection on the left side and before; anteriorly very slightly acuminate.

ANAS PÆCILORHYNCHA.

Anas.—Brunnea, marginibus pennarum suturationibus; rostro culmine ad basin et apice aurantiacis.

SPOTTED-BILLED DUCK.

Brown duck, with the edges of the feathers lighter; bill with the culmen at the base and tip orange.

<i>Length</i>	^{IN.} 25	<i>Bill</i>	^{IN.} $2\frac{1}{2}$	<i>Tarsi</i>	^{IN.} 2
---------------------	----------------------	-------------------	----------------------------------	--------------------	---------------------

Slaty brown, with the feathers on the breast, belly, and lower part of the neck broadly edged with yellowish white, those on the back narrowly so; crown as far as the eyes umber brown; neck white, with the centre of each feather dashed with brown; wing spot metallic green, bordered with velvet black, anterior to which there is a white stripe; exterior webs with some of the tertiaries white; bill black, with the tip for about a quarter of an inch and the upper part of the base orange; legs orange.

<i>Anas pæcilorhyncha</i>	<i>Gmel. Syst. Nat.</i> 1. 535.
	<i>Lath. Ind. Orn.</i> 2. 850.
	<i>Gray, Ind. Zool.</i>
<i>Mareca pæcilorhyncha</i>	<i>Steph. Sh. Zool.</i> 12. 134.
<i>Spotted-billed Duck</i>	<i>Lath. Gen. Syn.</i> 6. 487.

Common in the East Indies and Ceylon.

ANAS SPECULARIS.

An.—Brunnea, maculâ magnâ et ovali utrinque ante oculos; rostro colloque superiore singulis albis.

BRONZE-WINGED DUCK.

Brown duck, with a large oval spot before the eyes; bill and the upper part of the neck surrounded by a collar of white.

<i>Length</i>	^{IN.} $25\frac{1}{2}$	<i>Inner toe</i>	^{IN.} $\frac{7}{8}$
<i>Bill</i>	$1\frac{7}{8}$	<i>Middle toe</i>	$2\frac{1}{2}$
<i>Tarsi</i>	$1\frac{7}{8}$	<i>Outer toe</i>	$2\frac{1}{8}$

Head and neck umber brown, with a large oval spot of white between the eye and bill, placed

ANAS OBSCURA.

An.—Brunnea, speculo violaceo atro marginato.

DUSKY DUCK.

Brown duck, with the speculum violet margined with black.

	IN.		IN.
<i>Length</i>	24	<i>Tarsi</i>	$1\frac{3}{4}$
<i>Bill</i>	$2\frac{1}{8}$	<i>Middle toe</i>	$2\frac{1}{2}$

Male: crown, lower part of the neck, body, tail, and wings, entirely of a deep dusky brown, each feather edged with lighter; throat, cheeks, and upper part of the neck light brown, each feather having on its shaft a dark brown streak; speculum deep glossy violet margined with black; bill greenish, and formed similar to that of the mallard; legs and feet dusky yellow.

Female similar.

<i>Anas obscura</i>	<i>Gmel. Syst. Nat.</i> 1. 541.
	<i>Lath. Ind. Orn.</i> 2. 871.
<i>Dusky Duck</i>	<i>Lath. Gen. Syn.</i> 6. 545.
	<i>Wils. Am. Orn.</i> 8. p. 141. pl. 1—11.
	<i>Penn. Arct. Zool.</i> N. 496.

Inhabits America; occurs chiefly on the coasts of Florida, New England, and New Jersey, where it breeds in sequestered places on the marshes, laying eight or ten eggs similar to those of *A. Boschas*.

ANAS BOSCHAS.

An. Mas.—Capite colloque superiore viridibus; singulo collari albo.

An. Fem.—Brunnea, marginibus pennarum saturatioribus; speculo viridi.

WILD DUCK.

Male duck, with the head and upper part of the neck green; collar white.

Fem. brown duck, with the margins lighter; speculum green.

	IN.		IN.		IN.
<i>Length</i>	24	<i>Bill</i>	$2\frac{1}{4}$	<i>Tarsi</i>	2

Male: head and upper part of the neck metallic green, below which is a white collar; breast vinaceous; back brown; scapulars and flanks white, undulated with black; speculum metallic

green, margined anteriorly and posteriorly with white; quills, rump, and tail coverts brown, in some lights greenish, middle tail feathers curling upwards; tail grey-white; bill yellow; legs orange.

Fem.: brown, with the edge of the feathers lighter; speculum similar to that of the male.

Var. A. *Anas adunca* (Linn.) — Hook-billed Duck.

This variety has the bill turned downwards.

Var. B. Penguin Duck.

The peculiar upright position of this variety when at rest and standing distinguishes it.

<i>Anas Boschas</i>	<i>Linn. Syst.</i> 1. 205.
	<i>Gmel. Syst.</i> 1. 538.
	<i>Lath. Ind. Orn.</i> 2. 850.
<i>Wild Duck</i>	<i>Penn. Arct. Zool.</i> 2. 494.
<i>Le canard sauvage</i>	<i>Buff. Ois.</i> 9. 115.
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 12, fig. 10.</i>

Inhabits universally the temperate regions of the northern hemisphere, and does not appear to be subject to any considerable migration; breeds commonly in all marshy districts, laying from ten to fifteen eggs of a greenish colour.

The trachea is of nearly equal diameter throughout; the inferior larynx is furnished on the left side with an osseous bulb; anteriorly slightly acuminate.

ANAS FLAVIROSTRIS.

An.—Cinereo-brunneo, rostro flavo, naribus maculâ atrâ cingentibus.

YELLOW-BILLED DUCK.

Cinereous-brown duck, with a yellow bill, and the nostrils surrounded by a black patch.

	IN.		IN.
<i>Length</i>	$22\frac{1}{2}$	<i>Tarsi</i>	2
<i>Bill</i>	$2\frac{1}{2}$	<i>Middle toe</i>	$2\frac{1}{2}$

Slaty-brown duck, above with the edges of the feathers narrowly margined with lighter, below broadly; wing spot brassy green, surrounded anteriorly and posteriorly with a narrow streak of black, superiorly with one of the same colour, formed by the outer webs of the tertiaries, beyond the black streak posteriorly and anteriorly a narrow one of white; tail feathers pointed, slightly edged with light brown; bill yellow, with a black patch covering the nostrils; nail black; legs and feet brownish lead colour.

Anas flavirostris

Smith, Cat. S. Afric. Mus.

This new species is an inhabitant of the neighbourhood of the Cape of Good Hope. Specimens are in the South African Museum, and in the collection of the Zoological Society.

ANAS SPARSA.

An.—Brunnea, scapularibus remigibusque interioribus tribus vel duobus transversis et fasciis ferrugineis ornatis.

SPECKLED DUCK.

Brown duck, with the scapulars and tertiaries banded with three or four transverse fasciæ.

	IN.		IN.
<i>Length</i>	$23\frac{1}{2}$	<i>Tarsi</i>	2
<i>Bill</i>	$1\frac{3}{4}$	<i>Middle toe</i>	$2\frac{1}{2}$

Deep sepia brown duck, with the fore part of the neck and throat lighter, but thickly and minutely spotted with deep brown; scapulars and upper tertiaries each with one, two, or three bands of buff, on the former interrupted by the shaft and a small space on each side, which are of the same colour as the greater portion of the plumage, and on the latter merely extending over the outer webs; wing spot bright metallic green, bounded posteriorly and anteriorly first with a band of black, then with one of white; tail and tail coverts marked as the scapulars, but more obscurely; bill lead coloured; legs and feet orange, with the webs dusky.

Anas sparsa *Smith, Cat. S. Afric. Mus.*

Inhabits Southern Africa, and was brought to England by Dr. A. Smith, to whom we are indebted for permission to take the above description.

CARINA MOSCHATA.

Car.—Niger, ptilis albis.

MUSK DUCK.

Black, with the coverts white.

	IN.		IN.		IN.
<i>Length</i>	$32\frac{1}{2}$	<i>Bill</i>	$2\frac{5}{8}$	<i>Tarsi</i>	$2\frac{1}{2}$

Glossy black, with the wing coverts white; the bill in the male carunculated, less so in the female; legs black; caruncle red. In a domesticated state a greater or less quantity of white feathers make their appearance.

<i>Anas moschata</i>	Linn. Syst. 1. 199.
	Gmel. Syst. Nat. 1. 515.
<i>Anas Sylvestris Brasiliensis</i>	Raii Syn. 148, 150.
<i>Carina Sylvestris</i>	Steph. Sh. Zool. 12. 78.
<i>Trachea</i>	Linn. Trans. vol. 4, tab. 14, fig. 1—2, and
tab. 16, fig. 56; vol. 15, tab. 15, fig. a.

Supposed to have been originally natives of South America, but have now been domesticated in many parts of the world. In the tame varieties many of the feathers are white.

The tube of the trachea is of nearly equal size throughout; the inferior larynx is furnished on the left side with a compressed osseous bulb.

MICROPTERUS PATACHONICUS.

Mic.—Cinereus; subtus et macula post oculo albis

PATAGONIAN MICROPTERUS.

Cinereous micropterus; below and with a patch behind the eye white.

	IN.		IN'
<i>Length</i>	24	<i>Inner toe</i>	2 $\frac{1}{2}$
<i>Bill</i>	2	<i>Outer toe</i>	3 $\frac{1}{2}$
<i>Tarsi</i>	2 $\frac{3}{4}$	<i>Hind toe</i>	1

Dark grey, with the edges of the feathers on the flanks, head, back, wing coverts, and breast light chocolate; lower part of the breast, belly, abdomen, under tail coverts, and secondaries white; secondary coverts longer than the primary quills; tail long, with the feathers rather stiff; bill dusky green; legs and toes orange; thighs and a small triangular mark behind the eye white.

Micropterus Patachonicus, King..... *Proc. Zool. Soc.* 1830, p. 15.

In the museum of the Zoological Society, brought by Capt. James from Pricket's Harbour.

MICROPTERUS BRACHYPTERUS.

Mic.—Cinereus, subtus albus; maculâ ferrugineâ gutture.

SHORT-WINGED MICROPTERUS.

Grey micropterus, with the belly white; a ferruginous patch on the throat.

	IN.		IN.		IN.
<i>Length</i>	27	<i>Bill</i>	2	<i>Tarsi</i>	2½

Grey, with the head and neck lighter; a ferruginous tinge on the throat; belly and wing spot white; legs and bill orange; nail black.

Anas brachyptera *Lath. Syn.* 6. 439.

Voyage autour du monde Frey, p. 39.

Oiseaux grises, ou ois. de plein *Pernet Voy.* II. ch. xix. p. 21.

The specimen from which the above description was taken is in the museum of the Zoological Society, and was obtained from the Straits of Magellan; they also inhabit the Falkland and Staaten Islands. We have considerable doubts as to the propriety of making this and the preceding distinct species. Both are destitute of the powers of flight, merely using their wings to flapper over the surface of the sea. They are also bad divers, obtaining their living almost entirely by breaking shell-fish along the water-mark: hence the head is so strong, that Mr. Darwin informs us he had great difficulty in breaking one with his zoological hammer.

The tube of the trachea much resembles that of a true pochard; the inferior larynx approaches, however, nearer to that of those geese which have bulbs on that part than to any other form with which we are acquainted; consequently also to the wading ducks, from which it may be distinguished by being much stronger.

MELANITTA AMERICANA.

Mel. Mas.—Sequenti similis, sed tuberculâ ad basim rostri aurantiacâ.

AMERICAN SCOTER.

Male, duck similar to the following, but with the tubercle at the base of the bill entirely orange.

	IN.		IN.
<i>Length</i>	19	<i>Middle toe</i>	3
<i>Bill</i>	1½	<i>Outer toe</i>	2½
<i>Tarsi</i>	2½	<i>Inner toe</i>	2½

Male distinguished from the following, which it in every other respect resembles, by the protuberance at the base of the bill and a small space anteriorly and laterally being entirely orange, and by the sides of the nail at the apex of the upper mandible being suddenly narrowed.

Fem. : a similar distinction holds between the females of this and the following as regards the nail.

Oidemia Americana *Faun. Bo. Am.* 2. 450.

Nutt. Orn. 2. 422.

Brought home by Dr. Richardson from the neighbourhood of Hudson's Bay ; specimens are in our collection.

MELANITTA NIGRA.

Mel. Mas.—Ater ; tuberculá bifida atra ad basin rostri.

Mel. Fem.—Brunnea ; subtus cinerea, maculá albicante gutture.

BLACK SCOTER.

Male, scoter, with a bifid black tubercle at the base of the bill.

Fem. brown ; beneath cinereous, with a white patch on the throat.

	IN.		IN.
<i>Length</i>	21	<i>Tarsi</i>	1 9-12ths
<i>Bill</i>	1 $\frac{1}{4}$	<i>Middle toe</i>	2 $\frac{5}{8}$

Male entirely black, glossy on the head and neck ; no speculum ; a bifid protuberance at the base of the bill, orange in the centre, black on the sides ; a small patch anterior to the tubercle, also orange ; remainder black ; legs dark lead coloured.

Fem. : sooty black or brown ; a large light coloured grey patch on the throat, extending backwards to the hind neck, and below about half way down the fore neck ; bill (devoid or nearly so of a tubercle) entirely black ; legs as in the male.

Young similar to the female.

Melanitta nigra *Brehm. Hand. der Nat. aller, Vo. Deut.* 903.

Anas nigra *Linn. Syst.* 1. 196.

Gmel. Syst. 1. 588.

Lath. Ind. Orn. 2. 818.

Anas nigra minor *Raü Syn.* 141 *A.*

Oidemia nigra *Flem. Brit. An.* 119.

Scoter or Black Diver *Penn. Arct. Zool.* 2. 484.

Le Macreuse *Buff. Ois.* 9. 234. *pl.* 16.

A native of the northern regions, but migrating southwards during winter down the European

continent; breeds, like its congeners, on the coast in northern latitudes, making its nest near the water mark of grass, sea weed, and other rubbish, and near the time of hatching lines its nest with down from the body; lays from six to ten white eggs.

The trachea is very dissimilar from that of the two following species, having no bulb below the glottis, and the tube only enlarged to about twice its usual diameter in place of the inferior bulb; the inferior larynx is similar to that of its congeners. The trachea of the female is similar to that of the male, but not quite so much enlarged; both sexes have the bronchiæ large, and the tube acted upon by two pair of muscles, situated as in the velvet scoter.

MELANITTA PERSPICILLATA.

Mel. Mas.—Rostro aurantiaco, maculâ quadrangulâ utrinque baso atro; speculo nullo.

Mel. Fem.—Brunnea, maculâ cinereâ utrinque ad basin rostri.

SURF DUCK.

Male scoter, with the bill orange, a quadrangular patch on each side of the bill at the base; no speculum.

Fem.: brown duck, with a cinereous patch on each side at the base of the bill.

	IN.		IN.
<i>Length</i>	19	<i>Inner toe</i>	2
<i>Bill</i>	1½	<i>Middle toe</i>	2¾
<i>Tarsi</i>	1¾	<i>Outer toe</i>	2½

Male: bill orange, with a four-cornered diamond-shaped patch of black on each side at the base of the upper mandible; bill (raised at the base) short and thick; plumage in general dead black, slightly glossed on the sides of the neck, with a patch of white on the occiput, and another elongated one down the back of the neck; legs and toes orange, the webs dusky.

Fem.: dusky brown, lighter on the neck and belly; the raised portion of the bill not so prominent as in the male; an obscure patch behind the base of the bill on each side, and another elongated one between it and the occiput.

<i>Anas perspicillata</i>	<i>Linn. Syst.</i> 1. 261.
	<i>Gmel. Syst.</i> 1. 524.
<i>Oidemia perspicillata</i>	<i>Steph. Sh. Zool.</i> 12. 219.
	<i>Eyton, Hist. R. Brit. Birds,</i> 81.
<i>Black or Surf Duck</i>	<i>Wils. Am. Orn.</i> viii. p. 49. pl. 67.
<i>Black Duck</i>	<i>Edw. Gla.</i> p. 155.
<i>Macreuse a large bec Marchand</i>	<i>Buff. Ois.</i> 9. 244.
<i>Canard Marchand</i>	<i>Temm. Man.</i> 2. 853.

Common in America along the coast, from the river St. Lawrence to Florida, and has also been occasionally met with in the Orkney and Shetland Isles; it almost exclusively inhabits the sea. The female of this species may be distinguished from that of *O. nigra*, which it somewhat resembles, by the superior length of the bill, and by the grey marking on the cheeks behind the eye, that colour in the above-mentioned bird being confined entirely to the throat. The following is Wilson's description of the trachea and anatomy of the bird: "On dissection the gullet was found to be gradually enlarged to the gizzard, which was altogether filled with broken shell-fish. There is a similar hard expansion at the commencement of the wind pipe, and another much larger about three quarters of an inch above, where it separates into the two lobes of the lungs; this last was larger than a Spanish hazel nut, flat on one side and convex on the other; the protuberance on each side of the bill communicated with the nostrils, and was hollow." This description agrees with that of the rest of the genus in general character.

MELANITTA FUSCA.

Mel.—Ater, speculo et maculâ lunatâ suboculari albis.

VELVET DUCK.

Black duck, with the speculum and a lunate mark under the eye white.

	IN.		IN.
<i>Length</i>	20	<i>Tarsi</i>	$1\frac{3}{4}$
<i>Bill</i>	$1\frac{7}{8}$	<i>Middle toe</i>	$1\frac{3}{8}$

Male: entirely deep velvet black, with a lunate spot and a mark under the eye white; bill and legs orange, the former with a tubercle at the base; a space before it and the margins black; webs black.

Fem.: has the plumage above sooty black, below dirty grey; the tubercle at the base of the bill smaller than in the male.

Young similar to the female.

<i>Melanitta fusca</i>	<i>Boie in Brehm. Hand. der Nat. aller Vo.</i> <i>Deut.</i> 905.
<i>Anas fusca</i>	<i>Linn. Syst.</i> 1. 196. <i>Gmel. Syst.</i> 1. 507. <i>Lath. Ind. Orn.</i> 2. 848.
<i>Oidemia fusca</i>	<i>Flem. Brit. An.</i> 119. <i>Steph. Sh. Zool.</i> 12. 216.
<i>Velvet Duck</i>	<i>Penn. Arct. Zool.</i> 2. 272, <i>pl.</i> 196.
<i>Grand ou double macreuse</i>	<i>Buff. Ois.</i> 9. 242.
<i>Trachea</i>	<i>Linn. Trans. vol.</i> 4, <i>tab.</i> 15, <i>fig.</i> 3, 4, 5, 6, 7, <i>and vol.</i> 16, <i>tab.</i> 21, <i>fig.</i> 2, 3.

Inhabits the arctic regions, migrating southwards in the winter, like several of the other known species of the genus, on both the American and European continents; breeds in the neighbourhood of Hudson's Bay, on the borders of fresh water lakes, laying eight or ten white eggs. On the commencement of incubation the males, like those of the eider, form themselves in large flocks and desert the females.

The trachea of this species is very curious: it has two bulbs on the tube, the lower one composed of tracheal rings firmly ossified together; the upper one, situated immediately below the glottis, instead of being composed like the lower one of tracheal rings, is separate from the trachea, which passes through it, communicating with it by an orifice on each side. The inferior larynx, unlike the diving ducks in general, is not bulbous; the trachea is acted upon by two pairs of muscles, the first the usual sterno-tracheal ones, the other (furculo-tracheal) have their attachment a little before the middle bulb. For a figure of this see Vol. 15, Part 21, of the "Linnæan Transactions," illustrative of a passage by Mr. Yarrell on the organs of voice in birds, and our own plate.

SOMATERIA SPECTABILIS.

Som. Mas.—Capite supra et occipite cinereis; genis viridibus; rostrum tuberculâ magnâ basali.

Som. Fem.—Sequenti similis, sed tuberculâ minore basali.

KING DUCK.

Male eider, with the head above and occiput cinereous; the cheeks green; the bill with a large tubercle at the base.

Fem. eider, like the following species, but with a smaller tubercle at the base of the bill.

	IN.		IN.
<i>Length</i>	24 $\frac{1}{2}$	<i>Inner toe</i>	1 $\frac{7}{8}$
<i>Bill</i>	1 1-6th	<i>Middle toe</i>	2 $\frac{5}{8}$
<i>Tarsi</i>	1 5-6ths	<i>Outer toe</i>	2 $\frac{5}{8}$

Male: with the crown and occiput light cinereous grey; cheeks light green; round the base of the bill a band, and on the chin a V shaped mark black; remainder of the neck, breast, and upper part of the back white, slightly tinged with purplish; lower part of the back, tail quills, tertiaries, scapulars, and under parts black; tail and greater wing coverts white; bill and legs red, the former with a large knob at the base, the top of which is covered with feathers.

Fem.: in colour resembling the female of the common eider; but can be distinguished by the knob at the base of the bill being larger.

<i>Somateria spectabilis</i>	<i>Steph. Sh. Zool.</i> 12. 229.
<i>Anas spectabilis</i>	<i>Linn. Syst.</i> 1. 195.
	<i>Gmel. Syst.</i> 1. 907.
	<i>Lath. Ind. Orn.</i> 2. 845.
<i>Grey-headed Duck</i>	<i>Edw. Gla. pl.</i> 154.
<i>Anas freti Hudsonis</i>	<i>Briss. Orn.</i> 365.
<i>King Duck</i>	<i>Penn. Arct. Zool.</i> 2 N. 481.
<i>Le Canard a tete grise</i>	<i>Buff. Ois.</i> 9. 258.
<i>Trachea</i>	<i>Linn. Trans. vol.</i> 15, <i>tab.</i> 15, c—d.

Inhabits the same countries as the preceding; builds its nest on rocks and islands on the sea coast, formed of sea weed, laying five or six eggs scarcely to be distinguished from the preceding. By the Greenlanders this bird is considered a delicacy, particularly the knob at the base of the bill.

The trachea is not dissimilar in form from that of the common eider, but the enlargement at the inferior larynx is smaller. We much regret we have been unable to obtain one for dissection.

SOMATERIA ST. CUTHBERTI.

Som. Mas.—Capite supra occipiteque viridibus.

Som. Fem.—Brunnea, transversim lineata atro.

COMMON EIDER DUCK,

Male eider, with the head above and occiput green.

Fem. brown eider, transversely lineated with black.

	IN.		IN.
<i>Length</i>	24	<i>Inner toe</i>	2 $\frac{1}{8}$
<i>Bill</i>	2 1-6th	<i>Outer toe</i>	2 $\frac{1}{8}$
<i>Tarsi</i>	2	<i>Middle toe</i>	2 $\frac{1}{8}$

Male: crown and occiput green; back, wing coverts, and remainder of the head and neck white, with a yellowish tinge; quills and tail grey below, with the secondaries and rump black; bill and legs black.

Fem.: light brown on the breast; a lunule of black on each feather near the tip dark brown, with the edges of the feathers lighter; belly obscure; lesser coverts, the back, tail, secondaries, and greater wing coverts grey brown; primaries dark brown, with the edges light grey brown; bill and legs black.

<i>Somateria mollissima</i>	<i>Steph. Sh. Zool.</i> 12. 224.
<i>Anas mollissima</i>	<i>Linn. Syst.</i> 1. 178.
	<i>Gmel. Syst.</i> 1. 511.
<i>Anas St. Cuthberti</i>	<i>Raii Syn.</i> 14. 1.
<i>Eider—St. Cuthbert's Eider</i>	<i>Penn. Arct. Zool.</i> 2 N. 480.

- Oie a duret ou Eider*..... Buff. 9. 103. t. 6.
L'Eider..... Feill. En. Method. Orn. 30. 119.
Trachea..... Linn. Trans. vol. 12, tab. 30, fig. 1.

Like its congener, an arctic species, seldom seen south of the 55th degree N. L. but in the parts within that on all three continents and the adjacent islands, where it breeds, building its nest of sea weeds, and laying four eggs of a greenish colour. The trachea of the male has the tube of nearly equal diameter throughout; the inferior larynx swells outwards in front, and has on the left side a small bulb; the left bronchia is the largest, and is dilated in the centre.

POLYSTICTA STELLERI.

- Poly. Mas.*—Albus, fronte occipiteque viridibus; collum cingulo atro.
Poly. Fem.—Ferruginea, atro et obscuro maculato, duabus maculis albis tectricibus.

WESTERN DUCK.

Male, white duck, with the forehead and occiput green, neck with a black collar.

Fem.: ferruginous duck, marked with black and dusky; with two white spots on the wing coverts.

Length 17 inches.

Male: chin, throat, fore part of the neck, and collar black; a black stripe extends from the collar down the neck to the back, which is also black; head sides and the upper part of the neck to the collar white; forehead and nape greenish; a black irregular patch behind the eyes, with a narrow band proceeding from it, and surrounding the orbits; breast and sides light ferruginous; vent and tail black, with the edges of the feathers lighter; a humeral spot black; quills dark brown; tertiaries (very long) black edged with light brown.

Fem.: ferruginous, marbled with dark and black, with two white spots upon the wing coverts, the feathers of which are straight and blackish. (*Nuttall.*)

<i>Anas dispar</i>	Gmel. Syst. 1. 535.
<i>Anas Stelleri</i>	Gmel. Syst. 1. 518.
	Pull. Spic. Zool. v. p. 35. tab. 5.
<i>Polysticta Stelleri</i>	Eyton. Hist. R. Brit. Birds, 79.
<i>Macropus Stelleri</i>	Nuttall, Orn. 2. 451.
<i>Fuligula Stelleri</i>	Bonap. Syn. 344.

First discovered by Steller in Kamschatka, and do not appear to stray far from their native place. A pair were shot in Oster Gothland, in Sweden, and are figured by Sparman. Another was killed in the British isles. Nothing is known of the anatomy of this beautiful species.

KAMPTORHYNCHUS LABRADORUS.

Kamp. Mas.—Dorso, primariis, collo, fasciâ occipitali, subtusque, atris.

Kamp. Fem.—Capite, mento, colloque cinereo; dorso alisque obscuris.

PIED DUCK.

Male duck, with the back, primaries, neck, occipital fascia, and below, black.

Fem. duck, with the head, chin, and neck cinereous; the back and wing dark,

	IN.		IN.
<i>Length</i>	21 $\frac{1}{2}$	<i>Outer toe</i>	2 $\frac{3}{8}$
<i>Bill</i>	1 $\frac{5}{8}$	<i>Middle toe</i>	2 $\frac{1}{2}$
<i>Tarsi</i>	1 $\frac{1}{2}$	<i>Inner toe</i>	2

Male: head, neck, breast, scapulars, wing coverts, and secondaries white; crown, a collar round the neck, belly, back, quills, and a streak on the occiput, black.

Fem.: about one inch shorter than the male; head, chin, and neck ashy grey; back and wings brownish slate colour; below ashy; legs in both sexes deep brown, with the legs black; bill black, with the base and edges of the mandibles orange.

<i>Fuligula Labrador</i>	<i>Bonap. Syn.</i> 337.
<i>Anas Labrador</i>	<i>Wils. Am. Orn. pl.</i> 69, fig. 6.
<i>Rhynchaspis Labrador</i>	<i>Steph. Sh. Zool.</i> 12. 121.
<i>Pied Duck</i>	<i>Penn. Arct. Zool.</i> 2. 282.
	<i>Lath. Gen. Syn.</i> iii. p. 497.

Found most plentifully on the western side of the American continent, but occasionally also on the eastern. Nothing is known of its nidification.

Wilson gives the following description of the trachea: "The windpipe of the male measures ten inches in length, and has four enlargements, viz. one immediately below the mouth at the interval of an inch; it then bends largely down to the breast bone, to which it adheres by two strong muscles, and has at that place a third expansion; it then becomes flattened, and before it separates into the lungs has a fourth enlargement much greater than the former, which is bony and round, puffing out from the left side. The bill of this species differs much in form from any in the family, being broad and thin at the tip, and having the lamellæ more elongated.

CALLICHEN CARYOPHYLLACEUM.

Cal.—Fuscum, capite colloque caryophyllaceis.

PINK-HEADED POCHARD.

Fuscous pochard, with the head and neck pink.

	IN.		IN.
<i>Length</i>	21	<i>Bill</i>	$2\frac{1}{2}$

Head and neck bright pink, the latter with a black streak down the back; lower part of the neck, flanks, back, and abdomen umber brown; speculum ferruginous; bill orange; legs lead coloured.

Anas caryophyllacea..... *Lath. Ind. Orn.* 2. 866.

Pink-headed Duck *Lath. Syn. Sup.* 1. 276.

Fuligula caryophyllacea *Steph. Sh. Zool.* 12. 207.

Inhabits several parts of India. Few specimens have been brought to this country: we only know of two at present existing; one is in our collection, the other in the British Museum; both were purchased at the sale of the late Col. Cobbe's collection.

CALLICHEN RUFINUM.

Cal. Mas.—Capite cristato, colloque anteriore castaneis; subtus obscurus, lateribus albis.

Cal. Fem.—Brunnea-cinerea, occipite obscuro.

RED-CRESTED POCHARD.

Male duck, with the head crested, and, together with the anterior part of the neck, chesnut; below obscure, with the flanks white.

Fem. cinereous brown duck, with the occiput dark.

	IN.		IN.
<i>Length</i>	21	<i>Outer toe</i>	$2\frac{1}{8}$
<i>Bill</i>	$2\frac{1}{8}$	<i>Middle toe</i>	$2\frac{1}{2}$
<i>Tarsi</i>	$1\frac{3}{4}$	<i>Inner toe</i>	$1\frac{3}{4}$

Male: head and upper part of the neck chesnut, the former crested; lower part of the neck, breast, and belly dirty brownish black; sometimes with a few of the feathers slightly tipped with

white; back light brown; shoulders and a large spot on the flanks white; tail feathers and quills cinereous; rump black; bill red, with the nail whitish horn colour; legs and toes red; webs black.

Fem.: slightly smaller than the male; light cinereous brown, with the crown darker; the throat and sides of the neck dark cinereous; bill and legs reddish brown.

<i>Callichen rufinus</i>	<i>Brehm. Hand. der. Nat. aller, Vo. Deut.</i> 924.
<i>Anas rufina</i>	<i>Gmel. Syst.</i> 1. 541.
	<i>Lath. Ind. Orn.</i> 2. 870.
<i>Anas capite rufo major</i>	<i>Raii Syn.</i> 140.
<i>Canard siffleur, Huppe</i>	<i>Buff. Ois.</i> 9. 282.
	<i>Temm. Mun.</i> 2. 864.
<i>Red-crested Pochard</i>	<i>Lath. Gen. Syn.</i> 6. 554.
<i>Mergoides rufina</i>	<i>Eyton, Hist. R. Brit. Birds,</i> 77.
<i>Trachea</i>	<i>Linn. Trans. Vol.</i> 15, <i>tab.</i> 15, <i>fig.</i> 2.

Inhabits the eastern portions of the north of Europe, and migrates to the Caspian Sea, to Hungary, Austria, and Turkey, and the great lakes of Switzerland; never found on the sea. Such is the account Temminck gives of the bird before us; and we are not aware that any thing besides is known concerning it, either as regards its breeding places or nest. Col. Sykes says it is rare in the Deccan.

The trachea of this species has the inferior larynx similar to that of the true pochards; the tube, however, differs much, having two flattened bulbs as in the *Merganser*; the inferior one is placed nearer to the inferior larynx than in the genus *Mergus*.

FULIGULA VALISNERIA.

Fu. Mas.—Dorso albo atro undulato; collo castaneo; capistroque atro.
Fu. Fem.—Capite, collo, pectoreque obscuris; subtus alba.

CANVAS-BACKED POCHARD.

Male pochard, with the back white, barred with black; neck chesnut; collar black.

Fem. pochard, with the head, neck, and breast obscure; below white.

	IN.		IN.
<i>Length</i>	24	<i>Tarsi</i>	1 $\frac{5}{8}$
<i>Bill</i>	2 $\frac{3}{4}$	<i>Middle toe</i>	3 1-5th

Male: a band round the base of the bill; top of the head, lower part of the neck and throat, rump, and tail coverts, black; head and neck reddish chesnut; back, scapulars, wing coverts, tip of the secondaries, and flanks white, barred with narrow bands of black and brown; primaries brown, darkest at their tips; belly white; bill and legs blackish brown.

Fem. somewhat smaller than the male; brown, with the sides of the head, neck, and breast ferruginous; scapulars and under plumage edged with the same; back and coverts brown, undulated with dirty white.

Anas valisneria..... *Wils.* 8. 103. *pl.* 70, *f.* 5.

Fuligula valisneria *Bonap. Syn.* N. 338.

Fawn. Bo. Am. 2 p. 451.

Steph. Sh. Zool. p. 196.

According to Dr. Richardson, this bird breeds in the fur countries from the 50th parallel to their most arctic limits. They arrive in the United States about the middle of October, frequenting the bay of Chesapeake, the sounds and bays of North Carolina, and the coast of Mexico. The canvas-backed duck may be distinguished from the following at once by its greater size and the superior length of the bill; it is said to be most delicious food. Of this, however, we have considerable doubts: none of the other species contained in this genus or even among the *Fuliginæ* are so.

FULIGULA FERINA.

Fu. Mas.—Capite, colloque, castaneis; nullo speculo.

Fu. Fem.—Brunnea; subtus alba.

RED-HEADED POCHARD.

Male pochard, with the head and neck chesnut; the speculum wanting.

Fem. brown: below white.

	IN.		IN.
<i>Length</i>	19	<i>Breadth of nail</i>	3-16ths
<i>Bill</i>	1 $\frac{7}{8}$	<i>Middle toe</i>	2 $\frac{3}{4}$

Male: head and neck chesnut; breast, upper part of the back, and rump, black; remainder of the back, wing coverts, thighs, flanks, and scapulars, white undulated with black; quills and tail grey; bill and legs lead coloured, the former with the tip black.

Fem.: rather smaller than the male; head and neck reddish brown; below white; back similar to the male, but more obscure.

Young similar to the female.

<i>Fuligula ferina</i>	<i>Steph. Sh. Zool.</i> 1. 193.
<i>Anas ferina</i>	<i>Linn. Syst.</i> 1. 203.
	<i>Gmel. Syst.</i> 1. 530.
	<i>Lath. Ind. Orn.</i> 2. 862.
<i>Anas rufa</i>	<i>Gmel. Syst.</i> 515.
	<i>Lath. Ind. Orn.</i> 2. 863.
<i>Aythya ferina</i>	<i>Boie in Brehm. Hand. der Nat. aller, Vo.</i>
	<i>Deut.</i> 920.
<i>Nyroca ferina</i>	<i>Flem. Brit. An.</i> 121.
<i>Pochard or Red-headed Pochard</i>	<i>Penn. Arct. Zool.</i> 2. 491.
<i>Millouin</i>	<i>Buff. Ois.</i> 216.
<i>Le canard a cou roux</i>	<i>Veill. En. Meth. Orn.</i> 1. 132.
<i>Le millouin</i>	<i>Veill. En. Meth. Orn.</i> 1. 136.
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 14, fig. 3—4.</i>

Common on the European and British coasts during winter. We have also received it from India. The sea appears to be its true habitation, but it is occasionally found frequenting fresh water, particularly in the breeding season, when it makes its nest on the borders of rivers, laying twelve eggs of a greenish white colour, and generally within the arctic circle.

The trachea of this and the two following species are so very similar that it can be only by a comparative description that we shall be able to make their distinctions understood. All have the tube of the trachea largest a little below the glottis; it is slightly contracted about one-third of its length from that part, again expanded, and again contracted immediately above the inferior larynx. The bulb in all is flattened, and mostly membranaceous, with merely just sufficient bone to stretch the membranes or tympanum. Two of the species, *F. cristata* and *F. ferina*, have a second swelling or small bulb entirely osseous on the right side, composed of enlarged rings of the lower part of the tube, through which the bronchia of that side passes; this bulb and the whole of the tracheal apparatus is smaller, as well as the tube shorter, in *F. cristata* than in *F. ferina*.

The trachea of *F. marilla* may be distinguished from the preceding by the inferior larynx being destitute of the bulb on the right side; in other respects it is similar.

FULIGULA AMERICANA.

Fu.—Similis precedenti, sed rostrum ungue latiore.

AMERICAN POCHARD.

Pochard like the preceding, but the bill with the nail broader.

	IN.		IN.
<i>Length</i>	$19\frac{1}{2}$	<i>Breadth of nail</i>	$\frac{1}{4}$
<i>Bill</i>	$1\frac{3}{4}$	<i>Tarsi</i>	$1\frac{1}{2}$

It is not without considerable diffidence that we have placed this bird as a distinct species from *F. ferina*. We have, however, examined a very large number of specimens, and find the following distinctions to hold good both in young and old birds. The bill in the present species is shorter than in *F. ferina*, the nail broader, and the whole bird rather larger. In form the nail is much more rounded at the sides, the bill more rounded immediately above the nostrils, and not so deeply channelled in the centre; also broader measuring across the culmen.

<i>Fuligula Americana</i>	Bonap.
<i>Fuligula ferina</i>	Faun. Bo. Am. vol. 2. p. 452.
<i>Red-headed Pochard</i>	American authors.

Inhabits North America.

FULIGULA MARILLA.

Fu. Mas..—Capite colloque nigris; dorso scapularibusque albis nigro lineato.
Fu. Fem..—Brunnea, cingulo albo basini rostri cingente.

SCAUP DUCK.

Male duck, with the head and neck black; the back scapulars white lineated with black.
Fem. brown duck, with a white band surrounding the base of the bill.

	IN.		IN.
<i>Length</i>	20	<i>Breadth of nail</i> . .	3-10ths
<i>Bill</i>	1 $\frac{3}{4}$	<i>Inner toe</i>	1 9-10ths
<i>Tarsi</i>	1 $\frac{1}{2}$	<i>Middle toe</i>	2 $\frac{1}{2}$

Male: head and neck black, with purplish inflections; back and scapulars greyish white, finely and transversely lineated with black; small wing coverts, primaries, and greater wing coverts deep brown, the former sprinkled with light brown; secondaries (forming the speculum), belly, and flanks pure white; lower part of the back and tail deep brown; bill, head, and legs lead coloured; feet the same, with the webs darker; irides yellow.

Fem.: nearly equal in size to the male; brown, with a white band surrounding the base of the bill, and expanding on the cheeks; the back and scapulars sprinkled with specks of light brown.

<i>Fuligula marilla</i>	<i>Steph. Sh. Zool.</i> 12, 198.
<i>Anas marilla</i>	<i>Linn. Syst.</i> 1, 196.
	<i>Gmel. Syst.</i> 1, 509.
	<i>Lath. Ind. Orn.</i> 2, 853.
<i>Fuligula Gesneri</i>	<i>Raii Syn.</i> 142, .A. 6.
<i>Nyroca marilla</i>	<i>Flem. Brit. An.</i> 122.
<i>Scaup Duck</i>	<i>Penn. Brit. Zool.</i> 2, 275.
	<i>Penn. Arct. Zool.</i> 2, 493.

<i>White-faced Duck</i>	<i>Son. Brit. Mus. t. 62.</i>
<i>Le Milouian</i>	<i>Buff. Ois. 9. 221.</i>
	<i>Veill. En. Method. Orn. 1. 152.</i>
<i>Le canard a bec cercle</i>	<i>Veill. En. Method. Orn. 1. 144.</i>
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 14, fig. 5—6.</i>

Met with in the same localities as *Fuligula ferina*, breeding near the arctic circle, and laying from five to eight greenish eggs.

FULIGULA AFFINIS.

Fu.—Precedenti similis, sed rostro brevior et ungue angustior armato.

AMERICAN SCAUP.

Duck like the preceding, but with the bill shorter and armed with a narrower nail.

	IN.		IN.
<i>Length</i>	19	<i>Breadth of nail</i> ..	1-5th
<i>Bill</i>	1 3-5ths	<i>Tarsi</i>	1 1-5th

The above bird may be distinguished from the preceding in both sexes by the following comparative marks: total length less; bill shorter and not so broad; nail much narrower, and not so much rounded at its sides; tarsi shorter.

Scaup Duck of American authors.
Fuligula affinis nobis.

Inhabits North America.

This is another bird of which we have entertained considerable doubts as to the propriety of making into a species: the above distinctions, however, appear through all the specimens we have examined to be constant. It is a curious fact that most of the water ducks* (*Fuligulinae*) of North America and Europe should be all distinct, while those land ducks (*Anatinae*) and the mergansers (*Merginae*) should in a great measure be identical. We can only attribute it to their being endowed with a greater power of wing, and to the former (*Fuligulinae*) following the sea coast in their migrations, and probably performing it by shorter flights, consequently being less liable, if I may be allowed the expression, to lose their road.

It is, however, yet a question whether, if the theory we have advanced in the introductory chapter of this work prove true, they will not eventually take the rank of varieties.

* We have never examined the eiders of North America.

FULIGULA RUFITORQUES.

Fu. Mas.—Ater: subtus albus, lateribus cinereo undulatis; capistro castaneo.

Fu. Fem.—Brunnea, fronte subtusque albis.

RING-NECKED POCHARD.

Male, black duck; below white, the sides undulated with cinereous; collar chesnut.

Fem. brown duck, with the forehead and below white.

	IN.		IN.
<i>Length</i>	$20\frac{1}{4}$	<i>Inner toe</i>	$1\frac{3}{4}$
<i>Bill</i>	$1\frac{7}{8}$	<i>Outer toe</i>	2 2-5ths
<i>Tarsi</i>	$1\frac{7}{8}$	<i>Middle toe</i>	2 3-10ths

Male: head purplish black; back, neck, and breast purplish brown, glossed with green inflections; belly white; flanks mottled with black; tail grey brown; primary coverts light grey; bill and legs deep lead colour, the former crossed with a band of greenish behind the nail; the base also surrounded with a narrow band of the same colour; head slightly crested.

Fem.: upper plumage dark brown, edged on the top of the head, scapulars, and breast with chesnut; flanks chesnut; bill, throat, and belly greyish white, speckled with brown; vent dark brown.

Young similar to the female.

Fuligula rufitorques Bonap. Syn. p. 393, N. 341.

This species was found by Dr. Richardson in the fur countries during winter; they frequent the rivers and estuaries of North America.

The above bird appears to hold an intermediate station between *F. cristata* and *F. marilla*—the crest is smaller than in the former, and larger than in the latter. The female may be distinguished from the female of the latter by the greater extent of the triangular space at the upper part of the base of the bill, and from that of *Cristata* by its larger dimensions.

FULIGULA CRISTATA.

Fu. Mas.—Supra ater, capite cristato; subtus albus.

Fu. Fem.—Mare similis, sed minore cristâ.

TUFTED POCHARD.

Male pochard; above black, with the head crested; below white.

Female similar to the male, but with the crest smaller.

	IN.		IN.
<i>Length</i>	13	<i>Tarsi</i>	$1\frac{3}{8}$
<i>Bill</i>	$2\frac{3}{8}$	<i>Middle toe</i>	$2\frac{1}{4}$

Male: head and crest (which is long) black glossed with violet; back and upper part of the plumage glossy brown-black; the scapulars undulated with narrow bars of whitish; below, together with the speculum, white; bill lead-coloured, with the nail black; feet and legs black.

Fem.: similar to the male, but not so glossy, and with the crest smaller.

<i>Fuligula cristata</i>	<i>Steph. Sh. Zool.</i> 12. 190.
<i>Anas cristata</i>	<i>Raii Syn.</i> 142 <i>A.</i>
<i>Anas fuligula</i>	<i>Linn. Syst.</i> 1. 207.
	<i>Gmel. Syst.</i> 1. 542.
	<i>Lath. Ind. Orn.</i> 2. 869.
<i>Aythya cristata</i>	<i>Brehm. Hand. der. Nat. aller, Vo. Deut.</i> 916.
<i>Anas scandiaca</i>	<i>Gmel. Syst.</i> 1. 520.
	<i>Lath. Ind. Orn.</i> 2. 859.
<i>Tufted Duck</i>	<i>Penn. Arct. Zool.</i> 2. 573.
<i>Lapmark Duck</i>	<i>Penn. Arct. Zool.</i> 2. 576.
<i>Morillon</i>	<i>Buff. Ois.</i> 9. 227.
<i>Le canard brun</i>	<i>Buff. Ois.</i> 9. 252.
<i>Le canard acrete</i>	<i>Teill. En. Method. Orn.</i> 1. 143.

A winter visitant in the British isles; is found also in Asia, migrating northwards to breed. In America this bird is not found, its place being supplied by the preceding species. Nothing is known of its nidification.

The trachea of the male in this species is furnished with two pair of muscles of voice, the first the usual sterno-tracheal ones, the second pair are furculo-tracheal, diverging from the tube at the point where it penetrates the cavity of the chest, and inserted between the rami of the os furcatum at the point from whence they branch to unite with the coracoids.

— NYROCA AUSTRALIS.

Ny.—Proximo similis, sed majore.

AUSTRALIAN POCHARD.

Duck like the next, but larger.

	IN.		IN.
<i>Length</i>	21	<i>Outer toe</i>	$2\frac{1}{2}$
<i>Bill</i>	$1\frac{7}{8}$	<i>Middle toe</i>	$2\frac{5}{8}$
<i>Tarsi</i>	$1\frac{1}{2}$	<i>Inner toe</i>	2

So extremely like *Nyroca Leucophthalmus*, that as far as colouring goes one description will serve. The greater size, much larger and robust bill, and the bill being deep lead coloured, with a fascia of lighter across just behind the nail, as well as being destitute of the white spot on the chin, serve to distinguish it.

Nyroca Australis Gould, MSS.

Our specimen was obtained from Mr. Gould, who received it from Australia.

NYROCA LEUCOPHTHALMUS.

Ny. Mas.—Ferrugineus; iridibus, speculo, mentoque maculâ, albis.

Ny. Fem.—Mare similis, sed obscurior.

WHITE-EYED DUCK.

Male, ferruginous duck; with the eyes, speculum, and patch on the chin, white.

Fem. like the male, but darker.

	IN.		IN.
<i>Length</i>	$17\frac{1}{2}$	<i>Tarsi</i>	$1\frac{1}{4}$
<i>Exp. alæ</i>	26	<i>Middle toe</i>	$2\frac{1}{4}$
<i>Bill</i>	$1\frac{1}{2}$	<i>Inner toe</i>	$1\frac{5}{8}$

Male: bright reddish chesnut, with a collar round the neck brown; speculum, a spot on the chin, and under parts pure white; the back and wings in some specimens inclining to brownish; bill and legs lead coloured; irides white.

Fem.: similar, but with the colours more obscure.

<i>Nyroca Leucophthalmus</i>	<i>Flem. Brit. An.</i> 121.
<i>Anas Africana</i>	<i>Gmel. Syst.</i> 1. 522.
	<i>Lath. Ind. Orn.</i> 2. 875.
<i>Anas Nyroca</i>	<i>Gmel. Syst.</i> 1. 542.
	<i>Lath. Ind. Orn.</i> 2. 869.
<i>Anas ferruginea</i>	<i>Gmel. Syst.</i> 1. 528.
	<i>Lath. Ind. Orn.</i> 2. 866.
<i>Aythya nyroca</i>	<i>Boie in Brehm. Hand. der. Nat. aller, Vo.</i>
	<i>Deut.</i> 918.
<i>African Teal</i>	<i>Lath. Syn.</i> 6. 555.
<i>Ferruginous Duck</i>	<i>Penn. Brit. Zool.</i> 2. 601.

Rare as the occurrence of this bird is in the British isles, it is not uncommon in some parts of France, Holland, and Germany, and is found also in India and North America. Mons. Temminck informs us in his Manual that it constructs its nest by the sides of rivers and morasses, laying eight or ten white eggs slightly tinged with greenish.

The form of the tube and inferior larynx in this species will be best understood by consulting the plate. It is acted upon by two pairs of muscles, the first the usual sterno-tracheal ones, the others furculo-tracheal. It differs from that of the foregoing genus in having the tube much more enlarged.

NYROCA BRUNNEA.

Ny. Mas.—Obscurus, capite cristato; genis colloque inferiore castaneis; dorso brunneo.

Ny. Fem.—Mare similis, sed mento basoque rostri fasciâ cinctâ, genisque lineâ albis.

BROWN POCHARD.

Male, obscure duck, with the head slightly crested; the cheeks and lower parts of the neck chesnut; the back brown.

Fem. duck similar to the male, but with the chin and the base of the bill surrounded with a fascia; and the cheeks with a line white.

	IN.		IN.
<i>Length</i>	20	<i>Outer toe</i>	$2\frac{1}{4}$
<i>Bill</i>	$1\frac{5}{8}$	<i>Middle toe</i>	$2\frac{3}{8}$
<i>Tarsi</i>	$1\frac{3}{8}$	<i>Inner toe</i>	$1\frac{3}{4}$

Male: sooty black, with the head slightly crested and glossy; flanks and back inclining to olive brown, and speckled with light brown and white; wing spot white; cheeks deep chesnut, which colour extends for some distance down the neck; chin black; bill blueish lead colour; legs the same.

Fem.: as in the male, but with a circle round the base of the bill; a line from the eye to the lower part of the throat, and the tips of all the feathers below white.

Nyroca brunnea nobis.

Specimens of this species are in our own collection, and were received from Southern Africa. Specimens are also in the South African Museum, brought home by Dr. Andrew Smith.

HARELDA GLACIALIS.

Ha. Mas.—Capite colloque albis; maculâ auriculari brunneâ ornatus.

Ha. Fem.—Brunnea; gutture maculâ albâ ornata.

LONG-TAILED DUCK.

Male duck, with the head and neck white; with a brown auricular patch.

Fem. brown duck; the throat with a white patch.

	IN.		IN.
<i>Length</i>	22	<i>Inner toe</i>	$1\frac{5}{8}$
<i>Bill</i>	$1\frac{1}{8}$	<i>Middle toe</i>	$2\frac{1}{4}$
<i>Tarsi</i>	$1\frac{1}{4}$	<i>Outer toe</i>	$2\frac{1}{4}$

Male; head and neck white, with an auricular spot brown; scapulars and tertiaries, abdomen, and outer tail feathers white; breast, back, wing and middle tail feathers brown; during summer the plumage becomes more obscure, and similar to that of the female; middle tail feathers and tertiaries much elongated; bill black, with a band of red; legs and toes yellow, with the membranes dusky.

Fem.: without the long tail and tertiary feathers; dark brown, darkest on the back, the edge of the feathers lightest; throat with a large patch, and the under parts, light grey.

The young similar to the female.

<i>Harelda glacialis</i>	<i>Steph. Sh. Zool.</i> 12. 175.
<i>Anas glacialis</i>	<i>Linn. Syst.</i> 1. 203.
	<i>Gmel. Syst.</i> 1. 529.
	<i>Lath. Ind. Orn.</i> 2. 864.
<i>Anas hyemalis</i>	<i>Linn. Syst.</i> 202. 29.
	<i>Gmel. Syst.</i> 529. 29.
<i>Anas caudata harelda</i>	<i>Raii Syn.</i> 145. 14.
<i>Querquedula ferroensis</i>	<i>Briss. Orn.</i> 6. 466.
<i>Anas longicauda islandica</i>	<i>Briss. Orn.</i> 7. 299.
<i>Long-tailed Duck</i>	<i>Penn. Brit. Zool.</i> 2. 599.

<i>Cunard a longuequeue</i>	<i>Buff. Ois.</i> 9. 202.
<i>Cunard de midon</i>	<i>Buff. Pl. Enl.</i> 1008.
	<i>Veill. En. Method. Orn.</i> 1. 129.
<i>Sarcelle de ferroe</i>	<i>Buff. Ois.</i> 9. 278.
<i>L'Angletaske</i>	<i>Veill. En. Method. Orn.</i> 1. 134.
<i>Trachea</i>	<i>Mont. Orn. Dict. Sup. fig.</i> 12.
	<i>Linn. Trans. vol. 4, tab. 30, fig.</i> 3—4.

An inhabitant of the arctic regions, not migrating farther south than the northern part of Scotland. Breeds on the sea coast, making its nest of grass and sea weed; lays generally about eight eggs of a blueish colour. We have given an account of the trachea under the genus *Harelda*.

CLANGULA HISTRIONICA.

Clang. Mas.—Capite atro; maculâ utrinque ante oculari auricularique albis.

Clang. Fem.—Brunnea; capite maculis albis obscuris ornatis mare similibus.

HARLEQUIN DUCK.

Male duck, with the head black; a spot on each side before the eyes and on the ears white.

Fem. brown duck, with marks on the head similar to the male, but obscure.

	IN.		IN.
<i>Length</i>	16 $\frac{1}{2}$	<i>Outer toe</i>	2 $\frac{1}{4}$
<i>Bill</i>	1	<i>Inner toe</i>	1 $\frac{3}{4}$
<i>Tarsi</i>	1 $\frac{1}{4}$	<i>Middle toe</i>	2 $\frac{3}{8}$

Male: crown black; between the base of the bill and the eye is a triangular patch of white, with the small end towards the bill, from the upper corner of which a streak varying in intensity proceeds over the eye to the occiput; back of the neck black, with an elongated white streak on each side; throat and upper part of the neck shining violet black; an auricular patch white; collar and a streak extending nearly from the point of one wing round the front of the breast to that of the other, wing coverts, and tertiaries white; breast, belly, and vent brown; tail and wings dark umber brown; wing spot purple; bill and legs lead coloured, tipped with red; irides hazel.

Fem.: brown, with the margins of the feathers paler; a spot between the base of the bill and the eye, and an auricular one, whitish; belly and vent white blotched with brown; rather smaller than the male.

<i>Clangula Histrionica</i>	Steph. Sh. Zool. 1. 180.
	Eyton, Hist. R. Brit. Birds, 81.
<i>Anas Histrionica</i>	Linn. Syst. 1. 204.
	Gmel. Syst. 1. 534.

<i>Anas minuta</i>	<i>Linn. Syst.</i> 1. 204.
	<i>Gmel. Syst.</i> 1. 534.
<i>Anas torquata</i>	<i>Gmel. Syst.</i> 1. 514.
<i>Harlequin Duck</i>	<i>Penn. Arct. Zool.</i> 2. 490.
<i>Dusky and spotted Duck</i>	<i>Edw. Gla.</i> pl. 99.
<i>Le canard a collier de terre neuve</i>	<i>Buff. Ois.</i> 9. 250.
<i>Canard a collier ou histrion</i>	<i>Temm. Man.</i> 2. 878.
<i>La Sarcelle brun et blanche</i>	<i>Buff. Ois.</i> 9. 287.
<i>Le canard de terre neuve</i>	<i>Veill. En. Method. Orn.</i>
<i>Le canard histrion</i>	<i>Veill. En. Method. Orn.</i> 1. 146.

Inhabits the northern regions of both the American and European continents. A few specimens have been occasionally killed in Orkney and the northern islands; migrates northwards to breed; the female lays ten white eggs.

CLANGULA ALBEOLA.

Clang. Mas.—Capite colloque atris; maculâ albâ suboculari occipite extendente.

Clang. Fem.—Atra-fusca, albâ maculâ genis.

SPIRIT DUCK.

Male duck, with the head and neck black; a white patch extending below the eye to the occiput.

Fem. sooty black duck, with a white patch on each side of the head.

	IN.		IN.
<i>Length</i>	16	<i>Tarsi</i>	$1\frac{3}{8}$
<i>Bill</i>	$1\frac{1}{4}$	<i>Middle toe</i>	

Male: head and upper part of the neck rich purple green, with a large white patch extending from below the eye to the occiput; back and scapulars black; tail and quills hoary black; greater wing coverts white; secondaries black, above which is another white patch formed by the tertiary coverts; legs and toes orange; webs black; bill lead coloured.

Fem.: smaller than the male: head and plumage dark blackish brown; the fore part of the neck side of the breast, flanks, and vent feathers blackish grey; breast and belly white tinged with brownish orange; the white band on the ears and occiput much narrower than in the male, and obscure; lesser coverts and scapulars blackish brown; bill and feet brownish.

<i>Anas albeola</i>	<i>Linn. Syst.</i> 1. 199.
	<i>Gmel. Syst.</i> 1. 517.
	<i>Lath. Ind. Orn.</i> 2. 866.
<i>Anas Bucephala</i>	<i>Linn. Syst.</i> 1. 200.
	<i>Gmel. Syst.</i> 1. 521.
<i>Anas hyberna</i>	<i>Briss. Orn.</i> 6. 349.
<i>Querquedula Ludoviciana</i>	<i>Briss. Orn.</i> 6. 349.
<i>Anas rustica</i>	<i>Linn. Syst.</i> 1. 201.
	<i>Gmel. Syst.</i> 1. 524.
<i>Sarcelle de la Caroline</i>	<i>Buff. Ois.</i> 9. 286.
<i>Sarcelle blanche et noir ou la religieuse</i>	<i>Buff. Ois.</i> 9. 284.
<i>Petit canard a grosse tete</i>	<i>Buff. Ois.</i> 9. 249.
<i>Buffle-headed Duck</i>	<i>Catesb. Carol.</i> 1. pl. 95.
	<i>Penn. Arct. Zool.</i> 2. 489.
	<i>Lath. Gen. Syn.</i> 6. 532.
<i>Little brown Duck</i>	<i>Catesb. Carol.</i> 1. pl. 98.
	<i>Lath. Gen. Syn.</i> 6. 534.
<i>Le canard Lhoora</i>	<i>Veill. En. Method. Orn.</i> 1. 134.

An abundant species on the fresh water lakes of arctic America, where nidification is perfected, their nests are made in hollow trees contiguous to water. During autumn and winter they are found over a considerable portion of North America.

CLANGULA BARROVII.

Clang. Mas.—Capite colloque superiore atro; genis lunatâ fasciâ ornatis.

Clang. Fem.—Vulgari similis, sed singulo albo collo.

BARROW'S DUCK.

Male clangula, with the head and upper part of the neck black; the cheeks ornamented with a lunate fascia.

Fem. similar to the female of *Clangula vulgaris*, but with a white collar.

	IN.		IN.
<i>Length</i>	22½.	<i>Middle toe</i>	2½
<i>Bill</i>	1 7-12ths	<i>Outer toe</i>	1 5-12ths
<i>Tarsi</i>	1 7-12ths		

Male: head and upper part of the neck black, with purple and metallic green inflections; a crescent-shaped patch from the gape to the crown; below, lesser wing coverts, tips of the scapulars, and greater coverts, together with the outer secondaries, white; lateral tail coverts brown; bill lead coloured; legs and feet orange, the webs black.

Fem.: very similar to the female of *Clan. Vulgaris*, but rather larger; with a white ring round the middle of the neck, and the back of a darker colour.

Clangula Barroetii *Faun. Bo. Am.* 2. 415.
Gould, Birds of Europe.

The true habitat of this species is North America, in the neighbourhood of the Rocky Mountains, where it was discovered by Dr. Richardson. A single individual has since been obtained from Iceland by T. C. Atkinson, Esq. and a female is in our own collection. Brisson, in his "Ornithologie," pl. 37, fig. 2, vol. 6, has figured this bird in mistake for *Clangula vulgaris*, quoting for it the synonyms of Willoughby (whose plate is that of *C. vulgaris*) and of Linnæus. The specimen from which his description and plate is taken was in the collection of M. de Reaumur; he does not, however, say from whence obtained.

CLANGULA VULGARIS.

Clang. Mas.—Alba maculâ rotundatâ suboculari et speculo albo.

Clang. Fem.—Capite colloque superiore brunneis.

COMMON GOLDEN EYE.

Male duck, with a white spot under the eye, and wing spot white.

Fem. duck, with the head and upper part of the neck brown.

	IN.		IN.
<i>Length</i>	19	<i>Tarsi</i>	1½
<i>Bill</i>	1¼	<i>Middle toe</i>	2 7-10ths

Male: head and upper part of the neck green with purple inflections; remainder of the neck, belly, breast, greater wing coverts, and a spot below the eye, white; back, rump, lesser wing and tail coverts black; quills and tail black; legs, toes, and bill lead coloured; irides golden yellow.

Fem.: with the head and upper part of the neck brown; back, wings, and tail dusky slate colour; wing spot white; bill lead coloured, with a yellowish band round the nail.

<i>Anas clangula</i>	<i>Linn. Syst.</i> 1. 201.
	<i>Gmel. Syst.</i> 1. 523.
	<i>Raii Syn.</i> 142 A.
	<i>Lath. Ind. Orn.</i> 2. 867.
<i>Anas glaucion</i>	<i>Linn. Syst.</i> 1. 401.
	<i>Gmel. Syst.</i> 1. 525.
	<i>Lath. Ind. Orn.</i> 2. 868.
<i>Le garrot</i>	<i>Buff. Ois.</i> 9. 222.
<i>Clangula vulgaris</i>	<i>Flem. Brit. An.</i> 120.

- Clangula chrysophthalmus* Steph. Sh. Zool. 12. 182.
Canard garrot Temm. Man. 2. 870.
Golden eye Penn. Arct. Zool. 2. 276.
Trachea..... Linn. Trans. vol. 4, tab. 15, fig. 1—5.

Breeds in the arctic regions, but regularly migrates southwards during winter; is found commonly in Europe, frequenting both fresh and sea water, but more commonly the latter; during the breeding season, however, they are stated by travellers to frequent exclusively fresh water, making their nest on the shores of lakes and rivers in the fur countries, and laying from seven to ten white eggs.

The trachea has a large contractile bulb on the tube; the inferior larynx is not very dissimilar from that of the *Mergansers*; the tube is acted upon by two pairs of muscles of voice, the first of which are the usual sterno-tracheal ones, the second pair (which are furculo-tracheal) diverge from the tube at a little distance below the contractile bulb, and have their insertion on the rami of the os furcatum near the middle of each.

CLANGULA AMERICANA.

Clang. Mas.—*Clangula vulgari similis, sed majore.*

AMERICAN GOLDEN EYE.

Male clangula, similar to the common golden eye, but larger.

	IN.		IN.
<i>Length</i>	21	<i>Tarsi</i>	$\frac{1}{2}$
<i>Bill</i>	$1\frac{1}{2}$	<i>Middle toe</i>	$2\frac{3}{4}$

Very like the common golden eye of Europe, but larger, with the spot at the base of the bill more ovate, and the bill longer, broader, and stouter.

- Clangula Americana* Bonap. MSS.
Common Golden Eye Nutt. Orn. 2. 441.

It is even now doubtful whether or not many of the birds of North America are distinct from our own. There is, however, between many of them found enough of distinction for the practised eye of an ornithologist to say with certainty, this bird is North American—this European.

THALASSORNIS LEUCONOTUS.

Thal.—Corpore colloque ferrugineis; illo atro transversim fasciato, dorsoque imo albo.

FASCIATED DUCK.

Duck, with the body and neck ferruginous; the former transversely fasciated with black, and the lower part of the back white.

	IN.		IN.
<i>Length</i>	18	<i>Outer toe</i>	$2\frac{5}{8}$
<i>Bill</i>	$1\frac{5}{8}$	<i>Middle toe</i>	$2\frac{3}{4}$
<i>Tarsi</i>	$1\frac{3}{8}$	<i>Inner toe</i>	2

Head, upper part of the back, and flanks black; the tip and a bar across each feather dull brown; wing coverts black barred with ferruginous; neck light yellowish brown; belly rather darker; back and rump dirty white; tail coverts and tail black, the former tipped with light brown; feet and bill black; tail short; very old birds have a whitish spot on each side behind the base of the bill, and another smaller one on the chin.

Clangula leuconotus *Smith. Cat. S. Afric. Mus.*

In the museum of the Zoological Society, said to have been received from the Cape of Good Hope.

The bill in this species approaches in form to that of *Biziura*. Since we described the above, under another name, we have seen the specimens brought home by Dr. Smith from the Cape, and now exhibiting. As our specific appellation was not published, we have therefore adopted his.

BIZIURA LOBATA.

Biz. Mas.—Ater, albo transversim et minuté fasciatus; alis bruneis; mandibulâ inferiore carunculâ compressâ.

Biz. Fem.—Mare similis, sed carunculâ caret.

LOBATED DUCK.

Male, black duck, minutely and transversely lineated with white; wings brown; inferior mandibles with a compressed caruncle.

Fem. duck similar to the male, but without the caruncle.

	IN.		IN.
<i>Length</i>	33	<i>Inner toe</i>	3
<i>Bill</i>	$1\frac{3}{4}$	<i>Outer toe</i>	$3\frac{1}{2}$
<i>Tarsi</i>	2	<i>Middle toe</i>	$3\frac{5}{8}$

+ 1 *Oxyura Unifasciata*
 2 - - - *Domenica fem*

Painted by E. L. L.

Printed by "H. L. L."

Male: crown and upper part of the back of the neck black, occasionally speckled with whitish; back, lower part of the neck, flanks, rump, and under tail coverts deep glossy black, each feather transversely streaked with one or two narrow lines of white or light brown; wings and tail sooty black; remainder grey or silvery white; bill and legs lead coloured, the former with a large compressed caruncle on the lower mandible.

Fem.: smaller than the male, but in colouring similar; without the caruncle.

<i>Biziura Nova Hollandiæ</i>	<i>Steph. Sh. Zool.</i> 12. 222.
<i>Hydrobates lobatus</i>	<i>Temm. Pl. En.</i> N. 68.
<i>Anas lobata</i>	<i>Shar. Nat. Mis.</i> 8. pl. 255.
<i>Lobated Duck</i>	<i>Lath. Gen. Syn. Sup.</i> 2. p. 349.
<i>Le canard caroneule</i>	<i>Veill. En. Method. Orn.</i> 356.

All that is known of the habits of this curious bird is contained in a short notice by Lieut. Breton, R. N. inserted in the Zoological Proceedings for 1834, p. 19, from which we extract the following: "He (Lieut. Breton) stated that these birds are so extremely rare that he saw only three of them during his various excursions, which extended over twelve hundred miles of country. He has never heard of any instance in which more than two were seen together. They are only met with on the rivers and in pools left in the otherwise dry beds of streams. It is extremely difficult to shoot on account of the readiness with which they dive. The instant the trigger is drawn the bird is under water."

ERISMATURA MACCOA.

Eris.—*Brunnea*, gutture lineâque infra oculos albis.

MACCOA DUCK.

Brown duck, with the throat and a streak below the eye dirty white.

	IN.		IN.
<i>Length</i>	14	<i>Tarsi</i>	$1\frac{3}{8}$
<i>Bill</i>	$1\frac{1}{2}$	<i>Outer toe</i>	$2\frac{1}{4}$

Back black barred with ferruginous; flanks brown barred with the same; tail brown; head dark brown, undulated with ferruginous; chin and streak from the bill below the eye to the occiput whitish grey; below silvery grey undulated with brown; bill lead coloured; legs with a tinge of greenish; speculum none.

Oxyura Maccoa

Smith, Cat. S. Afric. Mus.

A specimen is in our collection, brought home from the Indian Isles. Except from the difference in locality, we should have supposed this bird to have been the young of *Oxyura rubida*. Since writing the above we have seen a specimen brought home by Dr. Smith, and have adopted his name.

ERISMATURA FERRUGINEA.

Eris.—Ferruginea, capite atro.

FERRUGINOUS DUCK.

Male. ferruginous duck, with the head black.

	IN.		IN.
<i>Length</i>	$17\frac{1}{2}$	<i>Tarsi</i>	$1\frac{1}{4}$
<i>Bill</i>	$1\frac{7}{8}$	<i>Centre toe</i>	$2\frac{3}{4}$

Male: deep ferruginous, with the head and upper part of the neck black; quills and tail brown; vent and belly silvery ash, tinged with ferruginous.

Closely allied to *Erismatura Australis*. The only specimen we have seen is in the British Museum, obtained from Chili.

ERISMATURA LEUCOCEPHALA.

Eris. Mas.—Vertice atro; subtus castaneus; reliquo albo.

Eris. Fem.—Mare similis, sed coloribus obscuris.

WHITE-HEADED DUCK.

Male duck, with the crown black; remainder of the head white; below chesnut.

Fem. similar, but with the markings obscure.

	IN.		IN.
<i>Length</i>	17	<i>Outer toe</i>	$2\frac{3}{4}$
<i>Bill</i>	$1\frac{3}{4}$	<i>Centre toe</i>	$2\frac{7}{8}$
<i>Tarsi</i>	$1\frac{3}{8}$	<i>Inner toe</i>	$2\frac{1}{8}$

Male: crown black; remainder of the head pure white; body silvery brown, tinged with ferruginous on the breast, flanks, and back, on the latter irrorated with black; tail and wings brown; no wing spot; legs and bill lead coloured; the latter more taper at the base than is usual in the genus.

Fem.: similar to the male, but with the colours not so distinct.

<i>Anas leucocephala</i>	<i>Gmel. Syst.</i> 1. 516.
	<i>Lath. Ind. Orn.</i> 2. 858.
<i>Undina leucocephala</i>	<i>Gould, Birds of Europe.</i>
<i>Anas mersa</i>	<i>Gmel. Syst.</i> 1. 520.

<i>White-headed Duck</i>	<i>Lath. Gen. Syn.</i> 6. 478.
<i>Ural Duck</i>	<i>Lath. Gen. Syn.</i> 6. 478.
<i>La Maereuse de Russie</i>	<i>Teill. En. Method. Orn.</i> 1. 140.
<i>Le canard Ural</i>	<i>Teill. En. Method. Orn.</i> 1. 127.

Inhabits the eastern portions of Europe ; occasionally found during migration in Austria and Hungary.

ERISMATURA RUBIDA.

Eris. Mas.—Brunnea, capite supra, fronte, et occipite atris ; genis mentoque albis.

Eris. Fem.—Mare similis, sed coloribus obscuris.

RUDDY DUCK.

Male, brown duck, with the upper part of the head, the forehead, and occiput black ; cheeks and chin white.

Fem. like the male, but with the colours less distinct.

	IN.		IN.
<i>Length</i>	15	<i>Outer toe</i>	2½
<i>Bill</i>	1½	<i>Middle toe</i>	2½
<i>Tarsi</i>	1¼	<i>Inner toe</i>	2

Male : crown, head, and back of the neck black ; back and tail brown dotted with whitish ; wings brown ; throat and cheeks white ; flanks, breast, and lower surface also white tinged more or less with ferruginous ; bill lead coloured ; legs and feet brown ; irides hazel.

Fem. : similar to the male, but with the back part of the head and neck inclining to brown ; the throat and cheeks light slate colour.

Young similar to the female.

<i>Erismatura rubida</i>	<i>Bonap. Cat. Am. and Europ. Birds.</i>
<i>Anas rubida</i>	<i>Wils. Am. Orn.</i> 8 p. 128, pl. 71, fig. 56.
<i>Fuligula (oryzura) rubida</i>	<i>Bonap. Syn.</i> p. 390, N. 336.
<i>Fuligula rubida</i>	<i>Faun. Bo. Am.</i> 2. 455.
<i>Gymnura rubida</i>	<i>Nutt. Man. Orn.</i> 2. 426.

Inhabits North America, retiring northwards to breed ; frequents fresh water in preference to the sea, according to Nuttall. When swimming they have a habit of carrying the tail so erect that it appears of the same height with the head and neck.

ERISMATURA AUSTRALIS.

Eris. Mas.—Capite colloque atris; corpore castaneo.

Eris. Fem.—Ferruginea, fasciata et punctata atro et brunneo.

AUSTRALIAN DUCK.

Male duck, with the head and neck black; the body chesnut.

Fem. ferruginous duck, fasciated and punctated with black and brown.

	IN.		IN.		IN.
<i>Length</i>	15	<i>Bill</i>	$1\frac{7}{8}$	<i>Tarsi</i>	$1\frac{1}{8}$

Male: with the head, throat, and neck black; below with the back dark chesnut; tail, wings, and rump black, the latter irrorated with ferruginous; under tail coverts silvery grey; bill, feet, and legs lead coloured.

Fem.: brown ferruginous, fasciated transversely with black and dark brown; throat light brown; under surface silvery grey, slightly tinged with brown; tail and wings brown.

Oxyura Australis, Gould *Zool. Proc.* 1836, p. 85.

Inhabits Australia. First sent to England by Lieut. Breton from Swan River. The specimens are in the collection of the Zoological Society.

ERISMATURA DOMINICA.

Eris. Mas.—Ferrugineus, capite anteriore atro; speculo albo.

Eris. Fem.—Brunnea, lineis albis infra et per oculos, occipite tendentibus: speculo mare simili.

DOMINICAN DUCK.

Male, ferruginous duck, with the head anteriorly black; the wing spot white.

Fem. brown duck, with a white streak through the eye and another below it, extending to the occiput; wing spot as in the male.

	IN.		IN.
<i>Length</i>	$13\frac{1}{2}$	<i>Outer toe</i>	$1\frac{3}{4}$
<i>Bill</i>	$1\frac{1}{4}$	<i>Middle toe</i>	2
<i>Tarsi</i>	1	<i>Inner toe</i>	$1\frac{1}{2}$

Male: fore part of the head sooty black; the hind part of the neck, back, scapulars, flanks, and rump ferruginous; wing coverts brown; quills ashy black, with the wing spot white; tail

black; throat, streak through the eye, and another below it, light ferruginous; bill and legs lead coloured.

Fem: with the back deep brown; flanks light brown, barred with darker; head deep brown, slightly undulated with ferruginous; under parts silvery grey, undulated with brown; a dusky white streak passes through the eye, and another below it; wing spot white; bill lead coloured; legs brown.

<i>Anas Dominica</i>	<i>Linn. Syst.</i> 1. 201.
	<i>Gmel. Syst.</i> 1. 521.
	<i>Lath. Ind. Orn.</i> 2. 874.
<i>Fuligula Dominica</i>	<i>Steph. Sh. Zool.</i> 12. 203.
<i>Querquedula Dominicensis</i>	<i>Briss. Orn.</i> 6. 472.
<i>Anas spinosa</i>	<i>Gmel. Syst.</i> 1. 522.
	<i>Lath. Ind. Orn.</i> 2. 874.
<i>Anas Dominica</i>	<i>P. Max. Bat. Nat. Braz.</i> 4. 938.
<i>Sarcelle de la Guadeloupe</i>	<i>Buff. Pl. Enl.</i> 938.
<i>Sarcelle a queue epineuse de Cayenne</i>	<i>Buff. Pl. Enl.</i> 968.
<i>Spinous-tailed Duck</i>	<i>Lath. Gen. Syn.</i> 6. 554.

A female specimen is in the British Museum. Both male and female are in our own collection, obtained from Jamaica. It also, according to Prince Maximilian, inhabits Brazil.

MERGUS MERGANSER.

Mer. Mas.—Capite colloque superiore viridibus; scapularibus primoribusque atris; corpore subtus albo.

Mer. Fem.—Capite rufo; corpore subtus cinerascanti-albo.

COMMON MERGANSER.

Male mer. with the head and upper part of the neck green; the scapulars and primaries black; below white.

Female, with the head rufous; below ashy white.

	IN.		IN.
<i>Length</i>	29	<i>Tarsi</i>	$1\frac{3}{4}$
<i>Bill</i>	2	<i>Middle toe</i>	$2\frac{3}{4}$

Male: head and neck deep glossy green, the former crested; upper part of the back and scapulars black; lower part of the back, tail, and secondary coverts, hoary grey; quills and secondaries brown; tertiaries white, beautifully edged with black; bill red, with the margins and culmen black; legs, feet, and webs, orange; irides hazel.

Fem.: head and upper part of the neck rufous brown, the former crested on the occiput; throat white; lower part of the neck, breast, flanks, and thighs, whitish ash; belly and abdomen yellowish white; above dark ash; feet, bill, and legs as in the male, but not so bright.

Young similar to the female.

<i>Mergus Merganser</i>	<i>Linn. Syst.</i> 1. 208.
	<i>Raii Syn.</i> 1. 34. <i>A.</i> 1.
	<i>Gmel. Syst.</i> 1. 544.
	<i>Briss. Orn.</i> 6. 231. <i>t.</i> 32.
	<i>Lath. Ind. Orn.</i> 2. 828.
	<i>Wils. Am. Orn.</i> pl. 68.
<i>Merganser Raii</i>	<i>Steph. Shaw Zool.</i> 12. 161.
<i>Mergus castor</i>	<i>Linn. Syst.</i> 1. 209.
.....	<i>Gmel. Syst.</i> 1. 545.
.....	<i>Lath. Ind. Orn.</i> 2. 829.
<i>Mergus ruficapillus</i>	<i>Gmel. Syst.</i> 1. 545.
<i>Mergus serratus longirostris</i>	<i>Raii Syn.</i> 1. 34. <i>A.</i> 2.
<i>Gooseander</i>	<i>Penn. Arct. Zool.</i> 2 N. 465.
.....	<i>Lath. Syn.</i> 6. 418.
<i>Dundiver</i>	<i>Penn. Arct. Zool.</i> 2. 465.
<i>Le Harle</i>	<i>Buff. Ois.</i> 8. 267.
——— femelle	<i>Buff. Ois.</i> 8. 236.
<i>Le Bievre</i>	<i>Veill. En. Method. Orn.</i> 1. 103.
<i>Le Harle</i>	<i>Veill. En. Method. Orn.</i> 1. 101.
<i>Trachea</i>	<i>Linn. Trans.</i> vol. 15, <i>tab.</i> 15, <i>fig.</i> h.

Inhabits the neighbourhood of the arctic circle, breeding in Iceland, Siberia, Kamschatka, and the fur countries, making its nest of grass near the water, and laying from ten to fourteen eggs of a yellowish colour. During winter, in hard weather, these birds migrate southwards, and are found over considerable portions of the continents of Europe, Asia, and America. We have also received them from Peru.

The trachea in the male has two enlargements on the tube, each of which are compressed, and composed of alternating semi-lunar plates.

The inferior larynx is very large and strong, projects most on the left side, but slightly before, behind, and on the right side. The bronchiæ are far apart, the left the largest; it does not differ in form from that of *M. serrator*, which we have figured.

MERGUS SERRATOR.

Mer. Mas.—Capite cristato, hóc colloque superiore viridibus; collo inferiore capistro albo.

Mer. Fem.—Capite cristato, hóc colloque superiore rufis, collo inferiore cinereo; ventre albo; speculoque fasciâ atrâ diviso.

RED-BREASTED MERGANSER.

Male mer. with the head crested, and the upper part of the neck green; the inferior part with a white collar.

Fem. mer. with the head crested, together with the upper part of the neck rufous; the inferior part cinereous; below white; speculum divided by a black bar.

	IN.		IN.
<i>Length</i>	20	<i>Inner toe</i>	2
<i>Bill</i>	$2\frac{5}{8}$	<i>Middle toe</i>	$2\frac{5}{8}$
<i>Tarsi</i>	$2\frac{1}{8}$	<i>Outer toe</i>	$2\frac{1}{2}$

Male: head and upper part of the neck glossy greenish black, the former crested; middle of the neck with a white collar; lower part of the neck and breast ferruginous, spotted with black; upper part of the back, scapulars, and sides of the breast black; wing coverts white, with a black bar across the larger ones; tail grey; flanks and lower part of the back white, undulated with narrow black lines; below white; irides reddish; bill orange, with the culmen black.

Fem.: head and crest brown; remainder of the neck greyish brown; secondaries, abdomen, and lower part of the breast white.

Young similar to the female.

<i>Mergus serrator</i>	<i>Linn. Syst.</i> 1. 208.
	<i>Gmel. Syst.</i> 1. 545.
	<i>Lath. Ind. Orn.</i> 2. 829.
<i>Mergus serrator leucomelas</i>	<i>Gmel. Syst.</i> 1. 546.
<i>Mergus serratus</i>	<i>Gmel. Syst.</i> 1. 546 <i>A.</i>
——— <i>niger</i>	<i>Gmel. Syst.</i> 1. 546 <i>B.</i>
——— <i>cristatus</i>	<i>Briss. Orn.</i> 6. 237. 2. <i>t.</i> 23.
<i>Red-breasted Merganser</i>	<i>Penn. Aret. Zool.</i> 2. 466.
	<i>Lath. Gen. Syn.</i> 6. 423.
<i>Le Harle, Huppe</i>	<i>Buff. Ois.</i> 8. 273.
<i>Harle à manteau noir</i>	<i>Buff. Ois.</i> 8. 277.
<i>Harle, Huppe</i>	<i>Temm. Man. Orn.</i> 2. p. 884.
<i>Trachea</i>	<i>Linn. Trans.</i> vol. 4, tab. 16, fig. 1—2.

Inhabits nearly the same countries as the preceding; but does not extend so far southwards; breeds in the neighbourhood of the arctic circle and in the highlands of Scotland, on the borders of fresh water lakes, making its nest of dried herbage lined with down stripped from its own body; lays from eight to twelve eggs of a brownish yellow colour.

The trachea in the male of this species differs from that of the last, in having only one enlargement on the tube (in other respects similar), and in the left bronchia being much larger than the right. The comparatively smaller size of this bird at once distinguishes it from the preceding.

MERGUS BRAZILIANUS.

Mer.—Niger, capite colloque viridi-æneis.

BRAZILIAN MERGANSER.

Black merganser, with the head and neck brassy green.

	IN.		IN.		IN.
<i>Length of bill</i>	2	<i>Tarsi</i>	$1\frac{3}{4}$	<i>Middle toe</i>	$2\frac{1}{4}$

Merganser, with the head and neck glossy green; the occiput with a long pendent crest the same; the whole of the back, tail, and wings sooty black; wing spot white, divided by a black streak; under surface and the lower part of the neck brown, the latter, with the flanks, minutely spotted and barred with white; the abdomen broadly so; bill black; legs and feet orange, with the webs dusky.

Mergus Brazilianus *Veill. Gal. des Oiseaux*, 2. 209.

Le Harle à huit brins *Dict. d'Hist. Nat.* 2d edit. 14. 222.

In the collection of the Zoological Society; evidently a young bird.

MERGUS CUCULLATUS.

Mer. Mas.—Capite atro cristato; cristâ albâ, et maculâ triangulari notatâ.

Mer. Fem.—Brunnea, capite cristato; gutture albo.

HOODED MERGANSER.

Male mer. with the head crested black; crest with a large triangular patch of white.

Fem. brown merganser, with the head crested; the throat white.

	IN.		IN.
<i>Length</i>	19	<i>Inner toe</i>	1½
<i>Bill</i>	1½	<i>Middle toe</i>	2
<i>Tarsi</i>	1½		

Male: head and neck glossy black, with purple and green inflections, the former largely crested with long silky feathers; crest marked with a fan-shaped patch of white, the small end placed towards the eye, a little behind it, and extending backwards nearly to the edge of the hood; breast and under surface white, the former with two bands of velvet black, tapering to a point in front of the wings; quills and tertiaries deep brown, the latter with white shafts and dark edges; tail dark brown; back black; flanks dark chesnut, with narrow transverse undulated bars of black or brown; irides yellow; bill red.

Fem.: with the crest smaller than in the male, and the feathers not so numerous; neck, back, head, crest, and wings umber brown, varying in intensity; throat white; lower part of the neck light umber brown; belly white; feet and legs in both sexes red.

Young similar to the female in colouring, but the males may be distinguished by the crest being larger.

<i>Mergus cucullatus</i>	<i>Linn. Syst.</i> 1. 207.
	<i>Gmel. Syst.</i> 1. 544.
	<i>Lath. Ind. Orn.</i> 2. 830.
	<i>Eyton, Hist. R. Brit. Birds,</i> 75.
	<i>Aud. Am. Orn.</i> 3. 246.
<i>Mergus fuscus</i>	<i>Lath. Ind. Orn.</i> 2. 802.
<i>Hooded merganser</i>	<i>Penn. Arct. Zool.</i> 2. 467.
	<i>Lath. Gen. Syn.</i> 6. 426. pl. ci.
<i>Round-crested Duck</i>	<i>Catesb. Carol.</i> 1. pl. 91.
<i>Brown merganser</i>	<i>Penn. Arct. Zool.</i> 2d Sup. 74.
<i>Le Harle couronne</i>	<i>Buff. Ois.</i> 8. 280.
	<i>Veill. En. Method. Orn.</i> 1. 103.
<i>Le Harle a queue fourchue</i>	<i>Veill. En. Method. Orn.</i> 1. 105.

Rare in Europe; one or two specimens have, however, been killed in the British isles. North America is its true habitat, over the whole of which it appears during winter to be found; as spring approaches, the greater portion of them retire to the north, though some remain and breed in Pennsylvania, Kentucky, and the intermediate district. It builds in holes of trees and rocks where it can find them; but in districts devoid of such localities builds on the ground on the borders of lakes, laying from six to ten white eggs.

MERGUS ALBELLUS.

Mer. Mas.—Supra albus, genis et occipite atris: capite cristato.

Mer. Fem.—Subtus alba, genis et occipite rufis: capite vix cristato.

WHITE SMEW.

Male mer. above white, with the cheeks and occiput black; head crested.

Fem. mer. below white, with the cheeks and occiput red; the head slightly crested.

	IN.		IN.
<i>Length</i>	17	<i>Outer toe</i>	2
<i>Bill</i>	$1\frac{3}{4}$	<i>Middle toe</i>	$2\frac{1}{8}$
<i>Tarsi</i>	$1\frac{1}{2}$	<i>Inner toe</i>	$1\frac{3}{4}$

Male: above and below white, with a large patch on each side of the head, and another one down the back of the neck, black; crest white; on the upper part of the breast on each side a bar of black, not meeting in front; coverts, primaries, and secondaries black; scapulars white, edged on their outer webs with black; tail, rump, and upper tail coverts grey; bill and legs lead coloured.

Fem.: throat, neck, belly, and abdomen white; the breast, back, tail, and upper tail coverts ash coloured; wings grey, with the outer webs of the scapulars white.

Young similar to the female.

<i>Mergus albellus</i>	<i>Linn. Syst.</i> 1. 209.
	<i>Gmel. Syst.</i> 1. 517.
	<i>Lath. Ind. Orn.</i> 2. 831.
	<i>Steph. Sh. Zool.</i> 12. 157.
	<i>Wils. Am. Orn.</i> 8. 136. pl. 71, fig. 7.
<i>Mergus minutus</i>	<i>Linn. Syst.</i> 1. 209.
	<i>Lath. Ind. Orn.</i> 2. 832.
<i>Mergus Asiaticus</i>	<i>Gmel. Syst.</i> 1. 548.
——— <i>stellatus</i>	<i>Briss. Orn.</i> 6. 252. pl. 6.
<i>Smer</i>	<i>Penn. Arct. Zool.</i> 2. 559.
	<i>Lath. Gen. Syn.</i> 6. 428.
<i>Blue merganser</i>	<i>Penn. Arct. Zool.</i> 2. Sup. 74.
<i>Le petit harle ou la piette</i>	<i>Buff. Ois.</i> 8. 275.
<i>Le piette</i>	<i>Veill. En. Method. Orn.</i> 1. 104.
<i>Le harle étoile</i>	<i>Buff. Ois.</i> 8. 278.
	<i>Veill. En. Method. Orn.</i> 1. 105.
<i>Trachea</i>	<i>Linn. Trans. vol. 4, tab. 10, fig. 3—1.</i>

Found in Europe, America, and Asia, but does not migrate far southwards; breeds in Siberia, Kamschatka, Greenland, and Iceland, generally on the banks of fresh water rivers or lakes, laying from ten to twelve yellowish eggs. The female and young of this species may be distinguished from those of the hooded merganser by the shortness and stoutness of the bill, and by the serratures not being pointed backwards, and more resembling the lamellæ of the true ducks. The tube of the trachea has one enlargement towards the middle, and continues of large diameter to the inferior larynx, which differs in form from that of the other mergansers, the greatest diameter being transverse instead of vertical; in other respects it is similar.

APPENDIX.

MUCH additional evidence in favour of the theory we have advanced having been brought forward in the second volume of Dr. Prichard's Researches into the Physical History of Man, we have formed the following Table, in order to shew the geographical distribution of the *Anatida*, and consequently those species which will, perhaps, some time or other, be accounted varieties.

M. signifies migrant; S. M. and W. M. summer and winter migrants; S. stationary; O. V. occasional visitant.

NAME.	Caucasian Range.	Mongolian Range.	Esquimaux Range.	Ethiopian Range.	American Range.	Malay Range.
<i>Choristopus semipalmatus</i>	..	..	..	..	..	S.
<i>Plectropterus Gambensis</i>	..	..	..	S.	..	
<i>Cercopsis Novæ Hollandiæ</i>	..	..	..	..	..	S.
<i>Cleophaga Magellanica</i>	..	..	..	..	S.	
<i>Sandvicensis</i>	..	..	..	..	S.	
<i>Bernicla Antarctica</i>	..	..	..	..	S.?	S.?
<i>jubata</i>	..	..	..	..	..	S.
<i>brenta</i>	W. M.	..	S. M.	..	..	
<i>leucopsis</i>	W. M.	..	S. M.	..	..	
<i>ruficollis</i>	O. V.	S.	..	..	..	
<i>Cheniscus Madagascariensis</i>	..	..	..	S.	..	
<i>Coromandeliana</i>	..	S.?	..	..	..	S.?
<i>Anser ferus</i>	W. M.	W. M.	S. M.	..	W. M.	
<i>erythropus</i>	W. M.	S. M.	S. M.	..	..	
<i>melanopterus</i>	..	..	..	..	S.	
<i>hyperboreas</i>	W. M.	W. M.	S. M.	..	..	
<i>Indica</i>	..	S.	..	..	..	
<i>Canadensis</i>	..	..	S. M.	..	W. M.	
<i>Hutchinsii</i>	..	..	S. M.	..	W. M.	
<i>cygnoides</i>	..	S.?	..	..	..	
<i>Cygnus olor</i>	..	S.	..	..	..	S.
<i>atratus</i>	..	..	..	..	..	
<i>nigricollis</i>	..	..	..	..	S.	
<i>Bewickii</i>	W. M.	..	S. M.	..	W. M.	
<i>Americanus</i>	..	..	S. M.	..	W. M.	
<i>buccinator</i>	..	..	S. M.	..	W. M.	
<i>immutabilis</i>	W. M.	..	S. M.	..	..	
<i>anatoides</i>	..	..	..	..	S.	
<i>ferus</i>	W. M.	..	S. M.	..	W. M.	
<i>Sarkidiornis regia</i>	..	S.	..	..	..	
<i>Africanus</i>	..	..	..	S.	..	
<i>Chenalopex Ægyptiaca</i>	O. V.	..	..	S.	..	
<i>cana</i>	..	..	..	S.	..	
<i>Tadorna bellonii</i>	S.	..	..	..	..	
<i>radjah</i>	..	..	..	..	..	S.
<i>Casarka rutila</i>	O. V.	S.	..	..	..	
<i>tadonoides</i>	..	..	..	..	..	S.
<i>castanea</i>	..	..	..	..	..	S.
<i>Dendrocygna viduata</i>	..	..	..	S.?	..	
<i>autumnalis</i>	..	S.	..	..	..	O.V.
<i>arcuata</i>	..	S.	..	..	..	S.
<i>arborea</i>	..	S.	..	..	..	

APPENDIX.

NAME.	Caucasian Range.	Mongolian Range.	Esquimaux Range.	Ethiopian Range.	American Range.	Malay Range.
<i>Leptotarsis Eytoni</i>	W. M.	W. M.?	S. M.	S.?	S.?	S.
<i>Dafila acuta</i>	..	..	..	..	S.	
— <i>pyrogaster</i>	..	..	..	..	S.	
— <i>urophasianus</i>	O.V.	S.	..	..	S.	
— <i>marmorata</i>	..	..	..	..	S.	
<i>Pæcilonitta Bahamensis</i>	..	..	..	S.		
— <i>erythrorhyncha</i>	W. M.	..	S. M.	S.?	W. M.	
<i>Mareca fistularis</i>	..	..	..	..	S.	
— <i>Americana</i>	..	..	..	..	..	S.
— <i>castanea</i>	..	..	..	..	S.	
— <i>Chiloensis</i>	..	S.	..	..	S. M.	S.?
<i>Aia sponsa</i>	..	S.	..	..	..	
— <i>galericulata</i>	..	S.	..	..	S.	
<i>Querquedula creceoides</i>	..	..	..	S.	..	
— <i>Hottentota</i>	..	..	..	..	S.	
— <i>erythrorhyncha</i>	..	S.	..	..	..	
— <i>falcaria</i>	O.V.	S.	..	..	..	S.?
— <i>glocitans</i>	..	..	..	..	..	
— <i>Javanensis</i>	..	S.	..	..	..	
— <i>Manillensis</i>	..	S.	..	..	..	
— <i>formosa</i>	..	S.	..	..	..	
— <i>crecca</i>	S.	S.	..	..	S. M.	
— <i>Carolinensis</i>	..	..	..	S.	..	
— <i>Capensis</i>	S. M.	S.	..	S.?	..	
<i>Cyanopterus circa</i>	..	..	..	..	S.	
— <i>fretensis</i>	..	..	..	..	S.	
— <i>discors</i>	..	..	..	..	S.	
— <i>Rafflesii</i>	W. M.	W. M.	..	O.V.	S.	
<i>Rhynchaspis clypeata</i>	..	..	..	..	S.	S.
— <i>rhynchotis</i>	..	..	..	..	S.	S.
— <i>maculata</i>	..	..	..	S.	..	
— <i>Capensis</i>	..	..	..	..	..	S.
<i>Malacorhynchus membrinacea</i>	W. M.	S.?	..	S.?	S.	O.V.
<i>Chaulelasmus strepera</i>	S.	S.	S. M.	O.V.	S.	
<i>Anas boschas</i>	..	..	..	..	S.	S.
— <i>obscura</i>	..	..	..	..	..	
— <i>superciliosa</i>	..	..	..	..	S.	
— <i>specularis</i>	..	S.	..	..	..	O.V.
— <i>pæcilorhyncha</i>	..	..	..	S.	..	
— <i>sparsa</i>	..	..	..	S.	..	
— <i>flavirostris</i>	..	..	..	..	..	
<i>Carina moschata</i>	..	..	..	..	S.	
<i>Micropterus patachonicus</i>	..	..	..	..	S.	
— <i>brachyptera</i>	W. M.	..	S. M.	..	S.	
<i>Melanitta fusca</i>	O.V.	..	S. M.	..	S.	
— <i>Americana</i>	W. M.	..	S. M.	..	..	
— <i>nigra</i>	O.V.	..	S. M.	..	S.	
— <i>perspicillata</i>	S.	..	S.?	..	S.?	
<i>Somateria mollissima</i>	S.	..	S.?	..	S.?	
— <i>spectabilis</i>	O.V.	S.	..	..	S.	
<i>Polysticta Stelleri</i>	..	..	..	..	..	
<i>Kamptorhynchus Labradoris</i>	S.	S.	..	..	..	
<i>Callichen rufina</i>	..	S.	..	..	..	S.
— <i>caryophylacea</i>	W. M.	S.	S. M.	..	..	
<i>Fuligula ferina</i>	..	..	S. M.?	..	S.	
— <i>falisneria</i>	..	..	S. M.?	..	S.	
— <i>erythrocephala</i>	W. M.	..	S. M.	..	..	
— <i>marilla</i>	W. M.	..	S. M.	..	..	
— <i>cristata</i>	..	..	S. M.?	..	S.	
— <i>rufitorques</i>	..	..	S. M.?	..	S.	
— <i>affinis</i>	..	..	S. M.?	..	S.	
<i>Nyroca brunnea</i>	S.	S.	..	S.	..	
— <i>leucophthalmus</i>	..	..	..	S.	..	

APPENDIX.

NAME.	Caucasian Range.	Mongolian Range.	Esquimaux Range.	Ethiopian Range.	American Range.	Malay Range.
Nyroca Australis						S.
Harelda glacialis	W. M.		S. M.		W. M.	
Clangula vulgaris	W. M.		S. M.			
—— Americana			S. M.		W. M.	
—— Barrovii	O.V.		S. M.		W. M.	
—— albeola			S. M.		W. M.	
—— histrionica	O.V.		S. M.		W. M.	
Thalassornis leuconotus				S.		
Biziura lobata						S.
Erismatura Maccoa				S.		
—— Dominica					S.	
—— Australis						S.
—— rubida					S.	
—— leucocephala	O.V.	S.				
—— ferruginea					S.	
Mergus albellus	W. M.	S.	S. M.			
—— cucullatus	O.V.		S. M.		W. M.	
—— serrator	S.	S.	S. M.		W. M.	
—— Brazilianus					S.	
—— merganser	W. M.	W. M.	S. M.		W. M.	

APPENDIX, No. 2.

In the following appendix we have given the admeasurements of the intestinal canal and number of vertebræ in all the species throughout the family *Anatide*, where we have been able to obtain them: the former, as a specific character, is subject to very great variation, an increase in age invariably producing a more lengthened measurement. The numbering of vertebræ in birds generally is very difficult: the mode in which we have proceeded, though differing from other authors, has at all events the advantage of simplification. We consider all those vertebræ anterior to the ribs, and not having any attachment to them, cervical; the dorsal are those anterior to the pelvic bones, having ribs attached to them; the sacral are those anchylosed together, and immediately succeeding the dorsal,—to some of these ribs are attached, either true or false; the numbering of those, where this is the case, we have shewn by the smaller and last number in the column headed No. of sacral vertebræ; so that, in order to ascertain the true number, the smaller one must be added to the greater. The caudal vertebræ are those posterior to and not anchylosed to the sacral.—In the numbering of the false ribs we have shewn their position by two figures, the first referring to those anterior to the true ones, the latter to those placed posteriorly to them.

N. B. The admeasurements are given in inches.

NAME.	Length of intestine from gizzard to anus.	Ditto ditto from anus to circa.	Ditto of cæca.	No. of cervical vertebræ.	Ditto of dorsal.	Ditto of sacral.	Ditto of caudal.	Total number of vertebræ.	True ribs.	False ribs.
<i>Plectropterus Gambensis</i> , F.	76	4	5 $\frac{3}{4}$							
<i>Bernicla Brenta</i>				18	6	13 3	7	47	7	2 1
<i>Anser segetum</i> , F.	85	7	13	18	6	11 3	6	47	7	1 1
— <i>cygnoides</i>				18	7	15 3	6	49	7	1 0
<i>Cygnus Bewickii</i> , M.	143	10	14 $\frac{1}{2}$							
— <i>atratus</i> , F.	114	7	11	23	6	16 4	8	57	8	2 1
— <i>olor</i>										
— <i>ferus</i> , F.	127	9 $\frac{1}{2}$	11	23	6	15 4	7	55	8	2 1
<i>Tadorna bellonii</i> , M.	123	4 $\frac{1}{2}$	8 $\frac{1}{2}$	16	6	13 3	7	45	6	2 1
<i>Dafila caudacuta</i> , M.	74	4 $\frac{1}{2}$	8 $\frac{1}{2}$	16	6	14 3	6		6	2 1
<i>Mareca fistularis</i> , M.	59	3 $\frac{1}{2}$	9	14	6	12 3	7	42	6	2 1
<i>Aia sponsa</i>	40	3	3 $\frac{1}{4}$							
<i>Querquedula erythrorhyncha</i> , M.	31 $\frac{1}{2}$	2 $\frac{1}{2}$	2 $\frac{1}{2}$							
— <i>crecca</i> , M.	30	2 $\frac{1}{2}$	2 $\frac{1}{2}$							
— <i>crecca</i> , F.	48	2 $\frac{1}{2}$	5	14	7	13 3	6	43	6	2 1
— <i>crecca</i> , F.	50	3 $\frac{1}{2}$	5 $\frac{1}{2}$							
<i>Cyanopterus ciria</i>										
<i>Rhynchaspis clypeata</i> , M.	89	4	5 $\frac{1}{2}$	14	6	14 3	6	43	6	2 1
— <i>clypeata</i> , F.	83	3 $\frac{3}{4}$	3 $\frac{1}{2}$							
<i>Chaulelasmus strepera</i>				14	7	13 3	7	44	7	2 1
<i>Anas boschas</i> , M.	84	4	9	14	6	14 3	7	44	7	2 1
— <i>boschas</i> , F.	79	3 $\frac{1}{2}$	6 $\frac{1}{2}$							
<i>Carina moschata</i> , M.	78	7	6 $\frac{1}{2}$	13	6	12 3	8	42	6	2 1
<i>Melanitta nigra</i> , F.	62	5	1	12	8	16 3	7	46	7	2 1
<i>Fuligula Gesneri</i> , M.	49	3 $\frac{1}{2}$	6	15	7	13 3	8	46	7	2 1
— <i>marilla</i> , M.	69	5	6	13	7	16 3	7	46	7	2 1
— <i>cristata</i> , M.	77	4	4 $\frac{1}{2}$	15	7	14 3	7	46	7	2 1
— <i>cristata</i> , F.	80	4 $\frac{1}{2}$	4 $\frac{1}{4}$							
<i>Nyroca leucophthalmus</i> , M.	43	2	4	16	7	15 3	6	47	7	2 1
— <i>leucophthalmus</i> , F.	37 $\frac{1}{2}$	2 $\frac{1}{2}$	3 $\frac{1}{2}$							
<i>Harelda glacialis</i>				14	7	14 3	7	45		
<i>Clangula vulgaris</i> , M.	54	4	2 $\frac{1}{2}$	14	7	15 3	6	45	7	2 1
— <i>vulgaris</i> , F.	49	4	3							
— <i>histrionica</i>				14	7	17 3	7	46		
<i>Mergus serrator</i> , F.	55	4 $\frac{1}{2}$	1	14	7	15 3	7	46		
— <i>merganser</i> , M.	79	4 $\frac{1}{2}$	1 $\frac{1}{2}$	14	7	13 3	7	44	7	2 1
— <i>merganser</i> , F.	66 $\frac{1}{2}$	6	1 $\frac{1}{2}$							
— <i>albellus</i> , F.	53	3 $\frac{1}{2}$	1 $\frac{1}{2}$	14	7	14 3	7	45	7	3 1

S U P P L E M E N T .

ADDITIONAL CHARACTERS OF AIA.

Ext. Char.—Lingua subobtusa, ad apicem appendiculata, ad latera unâ ordine setarum, et supra has setas quatuoribus spinis instructa.

Anat. Char.—Œsophagus magnus, in mediâ parte multum expandens ; proventriculo magno.

Ext. Char.—Tongue subobtuse, with an appendage at the tip, on the sides armed with one row of bristles, above which are four short spines.

Anat. Char.—Œsophagus large, much expanded in the middle ; the proventriculus large.

June 1838.

NEW WORKS AND NEW EDITIONS

LATELY PUBLISHED BY

LONGMAN, ORME, BROWN, GREEN, AND LONGMANS,

PATERNOSTER ROW.

LIFE and ADMINISTRATION of

EDWARD, FIRST EARL of CLARENDON.

With Original Correspondence and Authentic Papers, never before published.

By THOMAS HENRY LISTER, Esq.

3 vols. 8vo. with Portrait and Fac-similes, price £2. 8s. cloth lettered.

"A valuable contribution to the history of our native country."—*LITERARY GAZETTE*.

A HISTORY of PRICES,

With reference to the Causes of their principal Variations, from 1792 to the present time.

Preceded by a brief Sketch of the HISTORY of the CORN TRADE
in the last two Centuries.

By THOMAS TOOKE, Esq. F.R.S.

2 vols. 8vo. price £1. 10s. cloth lettered.

"In a book lately published, written by a gentleman of the highest possible authority—Mr. Tooke, there appeared so perfect and accurate a description, not only of the existing state of the revenue, and of the expenditure of the country, but also of the causes which had led to that state, that with the permission," &c. &c. &c.—Extract from MR. SPRING RICE'S SPEECH on
[the 18th May.

THE DOCTRINE of the DELUGE.

Vindicating the Scriptural Account from the doubts which have recently been cast upon it by geological speculations.

By the Rev. LEVESON VERNON HARCOURT.

2 vols. 8vo.—*Nearly ready.*

LETTERS, on the Subject of the CATHOLICS,

To my brother Abraham, who lives in the Country.

By PETER PLYMLEY.

Post 8vo. 21st Edition, 7s. in cloth.

NATURAL HISTORY.

INTRODUCTION to ENTOMOLOGY;

Or, ELEMENTS of the NATURAL HISTORY of INSECTS.

By WM. KIRBY, M.A. F.R.S. & L.S., and WM. SPENCE, Esq. F.L.S.

5th edition, 4 thick vols. 8vo. Plates and Portraits, price £4. in boards.

INTRODUCTION to the

MODERN CLASSIFICATION of INSECTS:

Comprising an Account of the Habits and Transformations of the different Families;
A Synopsis of all the British, and a Notice of the more remarkable Foreign Genera.

By J. O. WESTWOOD, Sec. Ent. Soc. London, F.L.S. &c.

Parts 1 and 2, price 2s. 6d. each.

To be completed in about Ten Monthly Parts, containing upwards of a Thousand
Illustrations on Wood.

* * * This work is intended to form a Sequel to the Work of Messrs. Kirby and Spence,
who have kindly sanctioned the work being so called.

MANUAL of BRITISH VERTEBRATE ANIMALS:

Containing Descriptions and Measurements of all the British Animals belonging to
the Classes *Mammalia*, *Aves*, *Reptilia*, *Amphibia*, and *Pisces*.

By the Rev. LEONARD JENYNS, M.A. F.L.S. &c.

8vo. 13s. bds.

A HISTORY of the RARER BRITISH BIRDS.

By T. C. EYTON, Esq. F.L.S.

Being intended as a Supplement to Bewick's Birds.

This work contains descriptions of all the species, about 33 in number, discovered
since the time of Bewick, with woodcut Figures of nearly all, and with Vignettes; also,
a Systematic Catalogue of all the British Birds hitherto discovered.

8vo. 10s. 6d.; royal 8vo. 21s.

By the same Author,

A MONOGRAPH of the ANATIDÆ, or DUCK TRIBE;

Including the GEESE and SWANS.

Illustrated by 24 lithographic Plates, chiefly by Gould, Scharf, and Lear; and upwards
of 70 Woodcuts, illustrating the Forms of the Bills and Feet in the different Genera.

Just ready.

LETTERS to a YOUNG NATURALIST, ON THE STUDY of NATURE and NATURAL THEOLOGY.

By J. L. DRUMMOND, M.D. Author of "First Steps to Botany."

12mo. Cuts, 7s. 6d. bds.

BOOK of NATURE:

A Popular Illustration of the General Laws and Phenomena of Creation.

By JOHN MASON GOOD, M.D. F.R.S.

3d Edition, 3 vols. fcp. 8vo. 24s.

TAXIDERMY;

Or, the Art of Collecting and Preparing Objects of Natural History.

12mo. Plates, 4th Edition, 7s. 6d.

NATURAL HISTORY—CONTINUED.**ESSAYS on NATURAL HISTORY.**

By CHARLES WATERTON, Esq. of Walton Hall;
Author of "Wanderings in South America."

1 vol. fcp. 8vo. 8s. cloth lettered.

"Most heartily do we wish that every student of natural history would, as far as possible, ground his pursuits and exertions on the example of Mr. Waterton: science would be the gainer not less than humanity. This little book abounds with hints and instructions of the greatest value to every lover of natural history."—EXAMINER.

GEOLOGY, CONCHOLOGY, & MINERALOGY.

REPORT on the
**GEOLOGY of CORNWALL, DEVON, and
WEST SOMERSET,**

By Order of the Lords Commissioners of Her Majesty's Treasury.
By HENRY T. DE LA BECHE, F.R.S. &c. Director of the Ordnance Geological Survey.

1 vol. 8vo. with numerous Sections and Plans.—*Shortly.*

AN INTRODUCTION to GEOLOGY;

Intended to convey a Practical Knowledge of the Science; and comprising the most important recent Discoveries; with Explanations of the Facts and Phenomena which serve to confirm or invalidate various Geological Theories.

By ROBERT BAKEWELL.

5th Edition, considerably enlarged from the 4th Edition, and with new Section and Cuts, price One Guinea in cloth, lettered.

NEW SYSTEM of GEOLOGY;

In which the great Revolutions of the Earth and Animated Nature are reconciled to Modern Science and to Sacred History.

By A. Uae, M.D. F.R.S.

8vo. with 7 Plates and 51 Woodcuts, 21s.

CONVERSATIONS on MINERALOGY.

With 12 Plates, engraved by Mr. and Miss Loway.

3d Edition, enlarged, 2 vols. 12mo. price 14s. in cloth.

MR. JOHN EDWARD GRAY, of the British Museum, is preparing a New Edition, thoroughly revised, and with considerable Additions, of TURTON'S

**MANUAL of the LAND and FRESH-WATER SHELLS
Of the BRITISH ISLANDS.**

Also, by Mr. J. E. GRAY,

OUTLINES of CONCHOLOGY;

Or, a short Account of the Orders, Families, and Genera of Molluscous Animals and their Shells. Illustrated with 12 Plates.

INTRODUCTION to LAMARCK'S CONCHOLOGY.

By E. A. CROUCH, F.L.S.

4to. 22 Plates, 31s. 6d. plain; £3. 3s. coloured.

NATURAL HISTORY (BOTANY) — CONTINUED.**BOTANY, AGRICULTURE, &c.***By* **JOHN LINDLEY, Ph. D. F.R.S. L.S. &c.***Professor of Botany in the London University College and in the Royal Institution.***INTRODUCTION to BOTANY.**2d Edition, with Corrections and considerable Additions,
1 large vol. 8vo. with numerous Plates and Woodcuts, 18s. cloth.*"We have no hesitation in pronouncing the 'Introduction to Botany,' by Dr. Lindley, to be the most valuable and perfect in any language we are acquainted with."*—MED. GAZ.*"The most valuable work of the kind in our language."*—BRIT. AND FOR. MED. REV.**A NATURAL SYSTEM of BOTANY;**

Or, a Systematic View of the Organization, Natural Affinities, and Geographical Distribution of the whole Vegetable Kingdom; together with the Uses of the most important Species in Medicine, the Arts, &c.

2d Edition, with numerous Additions and Corrections, and a complete List of Genera, with their Synonyms. 1 vol. 8vo. 18s. cloth.

SYNOPSIS of the BRITISH FLORA,

Arranged according to the Natural Orders.

2d Edit. with numerous Additions, Corrections, and Improvements, 12mo. 10s. 6d. bds.

**A KEY to STRUCTURAL, PHYSIOLOGICAL, and
SYSTEMATIC BOTANY.**

For the use of Classes. 8vo. 4s. 6d.

FLORA MEDICA:

Or, a Botanical Account of all the most remarkable Plants applied to Medical Practice in Great Britain and other Countries.

1 vol. post 8vo.—*Nearly ready.***FIRST PRINCIPLES of HORTICULTURE.**

2s. sewed.

GUIDE to the ORCHARD and KITCHEN GARDEN.

By G. LINDLEY, C.M.D.S. Edited by J. LINDLEY, Ph. D. F.R.S. &c.

1 large volume, 8vo. 16s. bds.

By **SIR JAMES EDWARD SMITH, M.D. F.R.S.***Late President of the Linnean Society, &c.***THE ENGLISH FLORA.**

New Edition, in 4 vols. 8vo., containing the FLOWERING PLANTS and the FERNS, £2. 8s.

Vol. V. Part 1. 12s.—CRYPTOGAMIA: comprising the Mosses, Hepaticæ, Lichens, Characeæ, and Algæ.

By Sir W. J. HOOKER, LL.D. F.R.A. and L.S., &c. &c.

Vol. V. Part 2—The FUNGI—completing the work, by Sir W. J. HOOKER, and the Rev. M. J. BERKELEY, F.L.S. &c. 8vo. 12s.

Compendium of the English Flora.

2d Edition, with Additions and Corrections. By Sir W. J. HOOKER, LL.D. &c. 12mo. 7s. 6d.

The same in Latin. 5th Edition, 12mo. 7s. 6d.

**INTRODUCTION to the STUDY of
PHYSIOLOGICAL and SYSTEMATICAL BOTANY.**

New Edition, with Illustrations of the Natural Orders (combining the object of Sir J. Smith's "Grammar" with that of his "Introduction.")

By Sir W. J. HOOKER, LL.D. &c. 8vo. 36 Plates, 16s. cloth.

NATURAL HISTORY (BOTANY)—CONTINUED.

By *SIR WILLIAM JACKSON HOOKER, K.B. LL.D.*

Regius Professor of Botany in the University of Glasgow, &c.

THE BRITISH FLORA ;

Comprising the FLOWERING PLANTS and the FERNS.

Svo. 4th Edition, enlarged and corrected, with 4 Plates, 12s. plain ; 16s. coloured.

. In this edition all the newly-discovered Species are introduced. The Linnæan arrangement is followed in the body of the work ; but in the Appendix are given the Characters of all the Natural Orders, with a List of the Genera, referring to the pages where they are described.

Vol. II. Part 1 of the above (CRYPTOGAMIA), Svo. 12s.

Vol. II. Part 2 (FUNGI), completing the work, by Sir W. J. HOOKER, and the Rev. M. J. BERKELEY. Svo. 12s.

MUSCOLOGIA BRITANNICA.

Containing the Mosses of Great Britain and Ireland, systematically arranged and described ; with Plates.

By Sir W. J. HOOKER, LL.D. F.L.S. &c., and T. TAYLOR, M.D. F.L.S. &c.

Svo. 2d Edit. enlarged, 31s. 6d. plain ; 43s. 3s. coloured.

ICONES PLANTARUM ;

Or, FIGURES, with brief Descriptive Characters and Remarks, of NEW or RARE PLANTS, selected from the Author's Herbarium.

2 vols. Svo. with 200 Plates, price 42. 16s. cloth lettered.

By *J. C. LOUDON, F.L.S. &c. &c.*

ENCYCLOPÆDIA of PLANTS ;

Comprising the Description, Specific Character, Culture, History, Application in the Arts, and every other desirable particular respecting all the Plants Indigenous to, Cultivated in, or Introduced into Britain.

With nearly Ten Thousand Engravings on Wood. 2d Edition, corrected, 1 large vol. Svo. 43. 13s. 6d. bds.

"The most useful and popular botanical work that has ever appeared in the English language."—JAMESON'S PHIL. JOURNAL.

HORTUS BRITANNICUS :

A Catalogue of all the Plants Indigenous to, Cultivated in, or Introduced into Britain.

Part I.—The Linnæan Arrangement, in which nearly Thirty Thousand Species are enumerated, &c. : preceded by an Introduction to the Linnæan System.

Part II.—The Jussieuian Arrangement of nearly Four Thousand Genera ; with an Introduction to the Natural System, and a Description of each Order.

Svo. 23s. 6d. in cloth.

AN ENCYCLOPÆDIA of GARDENING ;

COMPRISING

The Theory and Practice of Horticulture, Floriculture, Arboriculture, and Landscape Gardening : including all the latest Improvements ; a General History of Gardening in all Countries ; and a Statistical View of its Present State : with Suggestions for its Future Progress in the British Isles.

New Edition, greatly enlarged and improved, with nearly 1000 Engravings on Wood, 1 vol. Svo. 42. 10s.

NATURAL HISTORY (BOTANY)—CONTINUED.

Also, by Mr. LOUDON,

ENCYCLOPÆDIA of AGRICULTURE;

COMPRISING

The Theory and Practice of the Valuation, Transfer, Laying out, Improvement, and Management of Landed Property; and the Cultivation and Economy of the Animal and Vegetable Productions of Agriculture, including the latest Improvements; a General History of Agriculture in all Countries; and a Statistical View of its Present State: with Suggestions for its Future Progress in the British Isles.

With nearly Thirteen Hundred Engravings on Wood. 1 large vol. 8vo.
3d Edit. with a SUPPLEMENT, containing all the recent Improvements, £1. 10s. bds.

** The Supplement may be had separately, price 5s.

ARBORETUM ET FRUTICETUM BRITANNICUM;

Or, the Hardy Trees of Great Britain, Native and Foreign, Pictorially and Botanically delineated, and Scientifically and Popularly described.

Nos. 1 to 63, price 2s. 6d. each.

Complete in 8 thick 8vo. vols. (4 of Plates and 4 of Letterpress.)

ELEMENTS of PRACTICAL AGRICULTURE;

Comprehending the Cultivation of Plants; the Husbandry of Domestic Animals; and the Economy of the Farm.

By DAVID LOW, Esq. F.R.S.E. Professor of Agriculture in the University of Edinburgh.

1 vol. 8vo. 2d Edition, with Alterations and Additions, and above 200 Woodcuts, 18s. cloth lettered.

"No work on agriculture has appeared in our time which will bear a comparison with this excellent, and we would say classical work, of Professor Low. It will become the manual of practical agriculture for the British empire; and the judicious views and sound rules of our author will unquestionably prove beneficial to the agriculturists of other countries."

JAMESON'S EDIN. PHILOSOPHICAL JOURN.

ELEMENTS of AGRICULTURAL CHEMISTRY.

By Sir HUMPHRY DAVY, Bart. LL.D. F.R.S. &c.

5th Edition, with Notes by his Brother, Dr. JOHN DAVY, 8vo. 15s.

The CULTIVATION of the

GRAPE VINE on OPEN WALLS.

By CLEMENT HOARE.

8vo. New Edition, with Additions, 7s. 6d. cloth.

"Mr. Hoare has thrown more light on the subject of vine culture than any British gardener who has written on the subject."—LOUDON'S GARDENER'S MAG.

"One of the best productions upon any horticultural subject which has been published for some years."—ATHENÆUM.

FIRST STEPS to BOTANY.

By J. L. DRUMMOND, M.D.

12mo. 100 Woodcuts. 3d Edition, 9s. bds.

CONVERSATIONS on BOTANY.

With 22 Engravings. 12mo. 8th Edition, enlarged, 7s. 6d. plain; 12s. coloured.

CONVERSATIONS on VEGETABLE PHYSIOLOGY.

Comprehending the Elements of Botany, with their Application to Agriculture.

By JANE MARCET. 2 vols. 12mo. with Plates, 2d Edition, 12s.

By the same Author,

CONVERSATIONS on NATURAL PHILOSOPHY.

8th Edition, 10s. 6d. bds. With 22 Engravings by Lowry.

CONVERSATIONS on CHEMISTRY,

In which the Elements of that Science are familiarly explained and illustrated by Experiments; with a Conversation on the Steam Engine.

2 vols. 12mo. 13th Edition, with Plates by Lowry, 14s. bds.

POETRY.

LALLA ROOKH: an Oriental Romance.

By THOMAS MOORE, Esq.

New Edition, with 4 Engravings, from Paintings by R. WESTALL, R.A.
Fcp. 8vo. 10s. 6d. bds.; demy 8vo. without Plates, 14s.

Also,

IRISH MELODIES.

New Edition, in the press.

JAMES MONTGOMERY'S POETICAL WORKS.

A complete collected Edition, comprising "The Wanderer of Switzerland," "The West Indies," "World before the Flood," "Greenland," "Pelican Island," "Songs of Zion," "Poet's Portfolio," and all his smaller Poems.

3 vols. small 8vo. 18s.

SOUTHEY'S POETICAL WORKS;

COLLECTED BY HIMSELF.

To be completed in Ten Monthly Volumes, fcp. 8vo. with Frontispiece and Vignette, price 5s. each, cloth lettered.

* * 9 Volumes are now published.

FAMILY SHAKSPEARE;

In which nothing is added to the Original Text; but those Words and Expressions are omitted which cannot with propriety be read aloud in a Family.

By T. BOWDLER, Esq. F.R.S.

New Edition, in one large volume, 8vo. with 36 Illustrations after Smirke, Howard, &c. 30s. cloth; with gilt edges, 31s. 6d. Or in large type, without Illustrations, in 8 vols. 8vo. £4. 14s. 6d. bds.

REMAINS of HENRY KIRKE WHITE,

Selected, with prefatory Remarks, by ROBERT SOUTHEY, Esq.

2 vols. 8vo. 24s.; and 1 vol. 24mo. with engraved Title and Vignette, 5s.

N.B. The property of the Family having been invaded, it is necessary to state that these are the ONLY EDITIONS which contain the Life by Mr. Southey, and the whole of the contents of the Third Volume.

SELECT WORKS of the BRITISH POETS,

From JONSON to BEATTIE. With Biographical and Critical Prefaces,

By DR. AIKIN.

1 vol. 8vo. 18s. in cloth; or neatly done up, gilt edges, 20s.

SELECT WORKS of the BRITISH POETS,

From CHAUCER to WITHERS. With Biographical Sketches,

By ROBERT SOUTHEY, LL.D.

1 vol. 8vo. uniform with "Aikin's Poets," 30s. in cloth; or with gilt edges, £1. 11s. 6d.

THE MORAL of FLOWERS.

With 24 beautifully-coloured Plates, royal 8vo. 3d Edition, 30s. half-bound.

"Full of exquisite poetry."—BLACKWOOD'S MAGAZINE.

By the same Author,

THE SPIRIT of the WOODS.

1 vol. royal 8vo. with 26 beautifully-coloured Engravings, price 36s. half-bound.

"The two elegant volumes with which Mrs. William Hey has favoured the public are well fitted to lend grace and ornament to that which is solid and useful."—MED GAZ.

JOANNA BAILLIE'S POEMS, &c.

New DRAMAS, 3 vols. 8vo. 36s.—A SERIES of PLAYS, in which it is attempted to delineate the stronger Passions of the Mind. 3 vols. 8vo. £1. 11s. 6d.—MISCELLANEOUS PLAYS, 8vo. 9s.—METRICAL LEGENDS, 8vo. 14s.

RELIGION, THEOLOGY, ETC.

COLLEGE and SCHOOL GREEK TESTAMENT, WITH ENGLISH NOTES.

1 vol. 12mo. price 12s. cloth lettered.

By the Rev. S. T. BLOOMFIELD, D.D. F.S.A. of Sidney College, Cambridge;
Vicar of Bishbrooke, Rutland.

"Dr. Bloomfield's New Testament for the use of schools is an invaluable work. It is also entitled to the distinction of being regarded as an original production, the notes and critical apparatus being in general constructed with great labour for the present edition. It is impossible to say how far the public are indebted to Dr. Bloomfield for these labours of his industrious pen: they will carry down his name with honour to posterity."—EVANGELICAL MAGAZINE.

By the same Editor,

THE GREEK TESTAMENT,

With English Notes, Critical, Philological, and Exegetical, partly selected and arranged from the best Commentators, Ancient and Modern, but chiefly Original.

2d Edition, considerably improved, in 2 thick vols. 8vo. price £2.

GREEK and ENGLISH LEXICON of the NEW TESTAMENT.

By E. ROBINSON, D.D. Prof. Biblical Lit. Theol. Seminary, New York.

Edited, with careful revision, corrections, &c. &c.

By the Rev. S. T. BLOOMFIELD, D.D. F.S.A.

1 vol. 8vo. price 28s. in cloth, lettered.

"We hesitate not to affirm, that Dr. Robinson's Greek and English Lexicon to the New Testament, as edited by Dr. Bloomfield, is the best that has hitherto been published in this country."—BAPTIST MAG.

THE SUNDAY LIBRARY;

A Selection of Sermons from Eminent Divines of the Church of England, chiefly within the last Half Century.

With Notes, &c. by the Rev. T. F. DIBDIN, D.D.

Complete in 6 vols. small 8vo. with 6 Portraits of Distinguished Prelates, 30s. cloth.

"A little library for a churchman, and a treasure for the pious among the laity."
LITERARY GAZ.

THE SACRED HISTORY of the WORLD;

Philosophically considered, in a Series of Letters to a Son.

By SHARON TURNER, Esq. F.S.A. and R.A.S.L. 5th Edition, 2 vols. 8vo. 28s.

The Third and concluding Volume is just published, price 14s.

PRINCIPLES of CHRISTIAN PHILOSOPHY;

Containing the Doctrines, Duties, Admonitions, & Consolations of the Christian Religion.

By JOHN BURNS, M.D.

Regius Professor of Surgery in the University of Glasgow, &c.

12mo. 4th Edition, 7s. bds.

LETTERS from an ABSENT GODFATHER;

Or, a Compendium of RELIGIOUS INSTRUCTION for Young Persons.

By the Rev. J. E. RIDDLE, M.A.

Curate of Harrow; Author of "First Sundays at Church," &c. &c.

"An admirable manual of Christian duties."—ST. JAMES'S CHRONICLE.

DISCOURSES on the SOCINIAN CONTROVERSY.

By RALPH WARDLAW, D.D. Glasgow. 8vo. 5th Edition, 15s. cloth lettered.

By the same Author,

SERMONS. 8vo. 12s.

ORIENTAL CUSTOMS:

Applied to the Illustration of the Sacred Scriptures.

By SAMUEL BURDER, A.M. &c. New Edition, 12mo. 8s. 6d. bds.

HISTORY AND BIOGRAPHY.

A VIEW of the REIGN of JAMES II.

From his Accession to the Enterprize of the Prince of Orange.

By the late Right Hon. SIR JAMES MACKINTOSH.

In 4to. with a Portrait, 31s. 6d.

"We have no hesitation in pronouncing this fragment decidedly the best history now extant of the reign of James the Second. It contains much new and curious information, of which excellent use has been made. The accuracy of the narrative is deserving of high admiration."

EDINBURGH REVIEW.

Also, the above, completed to the SETTLEMENT of the CROWN, by the EDITOR.

To which is prefixed,

A Notice of the Life, Writings, and Speeches of Sir James Mackintosh.

4to. £3. 3s.

LIFE and TIMES of WILLIAM III.

King of England, and Stadtholder of Holland.

By the Rt. Hon. ARTHUR TREVOR (now LORD DUNOANNON), M. P. F.A.S. &c.

2 vols. 8vo. with Portrait, &c. price 24s.

LIFE of FREDERIC the SECOND, KING of PRUSSIA.

By LORD DOVER.

2 vols. 8vo. with Portrait, 2d Edition, 28s.

"A most delightful and comprehensive work. . . Judicious in selection, intelligent in arrangement, and graceful in style."—LITERARY GAZETTE.

MEMOIRS of SIR WILLIAM TEMPLE.

By the Right Hon. THOMAS PEREGRINE COURTENAY.

2 vols. 8vo. with Portrait, 28s.

LIFE of SIR HUMPHRY DAVY, Bart.

By his Brother, Dr. JOHN DAVY.

2 vols. 8vo. with Portrait, 28s.

HISTORY of ENGLAND,

From the earliest Period to the Death of Elizabeth.

By SHARON TURNER, Esq. F.A.S. R.A.S.L. 12 vols. 8vo. £8. 3s. bds.

The same work may also be had in the following separate portions:—

THE ANGLO-SAXONS. 3 vols. 8vo. 6th Edit. £2. 5s. bds.	REIGN OF HENRY VIII. 2 vols. 8vo. 3d Edit. 26s. boards.
THE MIDDLE AGES. 5 vols. 8vo. 3d Edition, £3. boards.	REIGNS OF EDWARD VI., MARY, and ELIZABETH. 2 vols. 8vo. 3d Edition, 32s. boards.

GIBBON'S HISTORY of the ROMAN EMPIRE:

For the Use of Families and Young Persons.

Reprinted from the Original Text, with the careful Omission of all Passages of an irreligious or immoral tendency.

By THOMAS BOWDLER, Esq. F.R.S. &c.

5 vols. 8vo. £3. 3s. bds.

HISTORY of THUCYDIDES;

Newly translated into English, and illustrated with very copious Annotations, &c.

By the Rev. S. T. BLOOMFIELD, D.D. &c., Editor of the Greek Testament.

3 vols. 8vo. with Maps, £2. 5s. bds.

"A version as literal and as perspicuous as erudition and industry combined can render it."
ECLECTIC REVIEW.

HISTORY & BIOGRAPHY—CONTINUED; & VOYAGES & TRAVELS.

MEMOIRS of the
COURT and CHARACTER of CHARLES I.

By LUCY AIKIN.

2 vols. 8vo. with Portrait, 2d Edition, 28s. bds.

"Miss Aikin's present work, and her previous Memoirs of the Courts of Elizabeth and of James I., are very acceptable additions to our literature."—EDINBURGH REVIEW.

By the same Authoress,

Memoirs of the Court of Queen Elizabeth. 6th Edit. 2 vols. 25s.

Memoirs of the Court of King James I. 3d Edit. 2 vols. 24s.

COMPANION TO MAUNDER'S TREASURY OF KNOWLEDGE.

THE BIOGRAPHICAL TREASURY;

Containing Memoirs, Sketches, or brief Notices of the Lives of about 12,000 Eminent Persons, from the earliest periods of History to the Present Day: surrounded by nearly 3500 Maxims and Precepts, original and selected.

By SAMUEL MAUNDER.

8s. 6d. cloth; 10s. 6d. bound, gilt leaves.

"A work worthy of great and lasting favour."—LITERARY GAZETTE.*"An extraordinary book, whether we look at the labour necessary to its production, the quantity of matter it contains, or the price at which it is sold."*—SPECTATOR.

VOYAGES & TRAVELS, GEOGRAPHY,

&c.

JOURNAL of a RESIDENCE in NORWAY,

Made with a view to inquire into the Rural and Political Economy of that Country, and the Condition of its Inhabitants.

By SAMUEL LAING, Esq.

2d Edition, 1 vol. 8vo. 14s. cloth.

"A book which every body should read."—DUBLIN REVIEW.

SEVEN WEEKS TOUR on the CONTINENT.

By JOHN ROBY, Esq. M.R.S.L.

2 vols. with numerous Illustrations, 25s.

"Should be read by all who contemplate visiting the countries included in his tour."

NAVAL AND MILITARY GAZETTE.

"The best book of the kind that has issued from the press these fifty years."—STANDARD.*"A pleasant postchaise companion."*—ATLAS.*"Full of anecdotes and sketches, graphic, amusing, and enthusiastic."*—MONTH. CHRON.

RAMBLES in the PYRENEES,

And a VISIT to SAN SEBASTIAN.

By FREDERIC W. VAUX, Esq.

1 volume, post 8vo. with a Frontispiece, drawn by Rochard, from a sketch by the Author, a Map, and 6 Engravings on Wood by S. Williams, 10s. 6d. cl. lett'd.

IMAGERY of FOREIGN TRAVEL;

Or, Descriptive Extracts from Scenes and Impressions in Egypt, India, Spain, Portugal, Italy, Germany, &c. &c.

By Major SHERER. 1 vol. post 8vo.

VOYAGES, TRAVELS, ETC.—CONTINUED.

SKETCH of ANCIENT and MODERN GEOGRAPHY.

By SAMUEL BUTLER, D. D.

Lord Bishop of Lichfield and Coventry, late Head Master of Shrewsbury Royal Free Grammar School.

Svo. New Edition, corrected, 9s. bds.

An Abridgment of the same. 12mo. price 2s. cloth.

By the same Author,

Atlas of Modern Geography, consisting of 23 Coloured Maps, from a New Set of Plates, with an Index of all the Names. svo. 12s. hf-bd.

Atlas of Ancient Geography, consist-

ing of 22 Coloured Maps, with a complete accentuated Index. svo. 12s. hf-bd.

General Atlas of Ancient and Modern Geography, 45 Coloured Maps, and two Indexes. 4to. 24s. half-bound.

** The Latitude and Longitude are given in the Indexes to these Atlases.

ENCYCLOPÆDIA of GEOGRAPHY:

Comprising a complete Description of the Earth; exhibiting its Relation to the Heavenly Bodies, its Physical Structure, the Natural History of each Country, and the Industry, Commerce, Political Institutions, and Civil and Social State of all Nations.

By HUGH MURRAY, F.R.S.E.

Assisted in

ASTRONOMY, &c. by Professor Wallace; | BOTANY, &c. by Sir W. J. Hooker;
GEOLOGY, &c. by Professor Jameson; | ZOOLOGY, &c. by W. Swainson, Esq.

With 82 Maps, drawn by Sidney Hall; and upwards of 1000 other Engravings on Wood, from Drawings by Swainson, T. Landseer, Sowerby, Strutt, &c., representing the most remarkable objects of Nature and Art in every Region of the Globe. Complete in one thick volume of upwards of 1500 pages, svo. £3. hf-bd. vellum.

"It is without a rival."—ASIATIC JOURNAL.*"The most perfect book on its subject."*—ATLAS.

NEW GENERAL ATLAS of FIFTY-THREE MAPS,

ON COLOMBIER PAPER.

With the Divisions and Boundaries carefully COLOURED.

Constructed entirely from New Drawings, and engraved by SIDNEY HALL.

Corrected to the present time.

Folded in half, and bound in canvas.....	£8 18 6
Ditto, half-bound in Russia	9 9 0
In the full extended size of the Maps, half-bound in Russia.....	10 0 0

Alphabetical Index of all the Names contained in the above Atlas,

With References to the Number of the Maps, and to the Latitude and Longitude in which the Places are to be found. Royal svo. 21s. cloth.

EDINBURGH GAZETTEER;

Containing a Description of the various Countries, States, Cities, Towns, Mountains, Seas, Rivers, Harbours, &c. of the World.

2d Edit. svo. with an Abstract of the last Population Return, and Maps, 18s.

As an Accompaniment to the above Work,

New General Atlas, constructed by A. ARROWSMITH.

Comprehended in Fifty-four Maps, including Two New Maps, with Recent Discoveries.

Royal 4to. half-bound, £1. 16s. plain; coloured, £2. 12s. 6d.

COMMERCE, ETC.

A DICTIONARY, PRACTICAL, THEORETICAL, AND HISTORICAL, OF COMMERCE and COMMERCIAL NAVIGATION.

Illustrated with Maps and Plans.

A New Edition, corrected and improved, in 1 closely and beautifully-printed volume, containing upwards of 1350 pages, 50s. bds.

By J. R. McCulloch, Esq.

In correcting this edition, the author has been assisted by valuable information communicated by many private and some official gentlemen in different parts of the world. In this respect he is under especial obligations to the government of Prussia. With a liberality of which there are certainly few (if any) examples, it has not merely taken pains to supply him with ample and authentic details as to the Commerce, Finances, &c. of that flourishing kingdom, but has authorized him to make any use he pleased of the information so communicated, without stipulation or condition of any kind, or so much as insinuating, directly or indirectly, what might be agreeable to it.

LIBER MERCATORIS ; or, The Merchant's Manual:

Being a practical Treatise on Bills of Exchange, particularly as relating to the Customs of Merchants; with the French Code relating to Bills of Exchange.

By F. HOBLER, Jun. Attorney-at-Law;

Author of "Familiar Exercises between an Attorney and his Articled Clerk."

Fcp. 8vo. 6s. cloth lettered.

STEEL'S SHIP-MASTER'S ASSISTANT, and OWNER'S MANUAL;

Containing General and Legal Information necessary for Owners and Masters of Ships, Ship-Brokers, Pilots, and other persons connected with the Merchant Service. New Edition, newly arranged, and corrected (containing the New Customs Laws, &c.) by J. STRICKMAN, Custom-House Agent. With Tables of Weights, Measures, Monies, &c. by Dr. KELLY. 1 large and closely-printed vol. 21s. bds.; 22s. 6d. bd.

CONVERSATIONS on POLITICAL ECONOMY.

By JANE MARCET. 12mo. 6th Edition, 9s. bds.

By the same Authoress,

John Hopkins' Notions on Political Economy. 12mo. 4s. 6d.

THE HISTORY and PRINCIPLES of BANKING.

By J. W. GILBART. 4th Edit. enlarged, 8vo. 9s.

This book may be considered as a grammar of banking. The general reader may here acquire a competent knowledge of most of the facts and principles connected with the subject.

NOVELS & WORKS OF FICTION.

THE ROBBER: A Tale.

By the Author of "The Gipsy," &c. &c. &c.

3 vols. post 8vo. £1. 11s. 6d.

"It is as full of the deepest interest, the sweetest philosophy, admirable portraits of characters, and gorgeous and powerful descriptions, as any work we have read for many years."

COURT JOURNAL.

"The best of Mr. James's romances."—SPECTATOR.

NOVELS AND WORKS OF FICTION—CONTINUED.

UNIFORM WITH "MARRYAT'S PIRATE."

BULWER'S LEILA ; & CALDERON the COURTIER.

15 Plates and Portrait of the Author, rl. 8vo. £1. 11s. 6d. ; Proof Plates, £2. 12s. 6d.

"These tales of Spain, from the pen of the Author of 'Eugene Aram' and 'Rienzi,' are what might be expected from such a quarter—highly descriptive, yet distinct, vigorous, and impassioned."—LITERARY GAZETTE.

UNIFORM WITH "BULWER'S LEILA."

CAPT. MARRYAT'S PIRATE & THREE CUTTERS.

20 Plates, by Stanfield, and Portrait of the Author, royal 8vo. £1. 11s. 6d. ; Proofs, £2. 12s. 6d.

"Captain Marryat never produced anything better than 'The Pirate.'"

METROPOLITAN MAG.

CONFESSIONS of an ELDERLY LADY.

By Lady BLESSINGTON. Eight Plates, £1. 1s.

"No actual confessions, whosoever the autobiographer might be, ever interested us more ; nor were any ever made that proffer a finer and truer lesson to humanity."—COURT JOURN.**CONFESSIONS of an ELDERLY GENTLEMAN.**

By Lady BLESSINGTON. Six Plates, 14s.

"There are few fictions, though of thrice the exterior pretensions of these 'Confessions,' that possess so much weight."—MONTHLY REVIEW.**THE DOCTOR, &c.**

Vol. IV. post 8vo. 10s. 6d.

** Vols. 1 to 3 may be had, price £1. 11s. 6d.

"Rich, beyond almost any other of the time, in the best knowledge, and in the most beautiful literature."—QUARTERLY REVIEW, No. 104.**MISCELLANEOUS.****COLONIZATION and CHRISTIANITY :**

A Popular History of the Treatment of the Natives in all their Colonies by the Europeans.

Contents:—The Discovery of America.—The Spaniards in the West Indies.—The Spaniards in Mexico.—The Spaniards in Peru.—The Spaniards in Paraguay.—The Spaniards in Brazil.—The Portuguese in India.—The Dutch in India.—The English in India ; their Mode of Territorial Acquisition there ; their Treatment of the Natives ; their more recent Amelioration of their Condition.—The English in America.—The Treatment of the American Indians by the United States.—The French in their Colonies.—The English in New Zealand and Australia.—The English at the Cape, &c. &c.

By WILLIAM HOWITT, Author of "The Book of the Seasons," &c.

1 vol. post 8vo.

By the same Author,

THE RURAL LIFE of ENGLAND.

2 vols. post 8vo. beautifully illustrated with Woodcuts, by S. Williams, 24s. cloth lettered.

*Contents:—Life of the Aristocracy ; Life of the Agricultural Population ; Picturesque and Moral Features of the Country ; Causes of the strong Attachment of the English to Country Life ; Forests of England ; Habits, Amusements, and Condition of the People.**"One of the most healthful, rigorous, fresh, and English-spirited of Mr. Howitt's productions. It is a work for all classes, rich and poor. It is written with good sense and good feeling. Its sentiments are as unexceptionable as its descriptions ; and we shall be greatly mistaken if it do not pass through many editions, and obtain a wider circulation season by season."*—COURT JOURNAL.*"Beautiful in its subject, beautiful in its literary execution, and beautiful in its getting up."*

ST. JAMES'S CHRONICLE.

MISCELLANEOUS—CONTINUED.

WOMAN in her SOCIAL & DOMESTIC CHARACTER.

By Mrs. JOHN SANDFORD.

5th Edition, 12mo. 6s. in cloth.

By the same Authoress,

ON FEMALE IMPROVEMENT.

2 vols. 12mo. 12s. cloth.

"Mrs. Sandford's previous work entitles her name to respect, and her present performance strengthens her claim to it. It is an offering which her sex should receive with gratitude."
COURT JOURNAL.

LACON; or, Many Things in Few Words.

By the Rev. C. C. COLTON. New Edition, 8vo. 12s. cloth.

SELECTIONS from the EDINBURGH REVIEW;

With a preliminary Dissertation and Notes.

Edited by MAURICE CROSS, Esq. 4 large vols. £3. 3s. bds.

GENERAL INDEX to the EDINBURGH REVIEW,

From Vol. I. to XX. 15s. bds.

WORKS OF GENERAL UTILITY.

THE TREASURY of KNOWLEDGE,
And LIBRARY of REFERENCE.

By SAMUEL MAUNDER.

In Two Parts: the 1st Part consisting of a new and enlarged English Dictionary, a Grammar, Tables of Verbal Distinctions, with Examples, &c.; the 2d Part including, among many other things of value, a new Universal Gazetteer, a Classical Dictionary, a Compendium of Chronology and History, a Dictionary of Law Terms, and various useful Tables.

The whole uniquely surrounded by Moral Maxims and Proverbs. 9th Edition. 1 thick vol. royal 18mo. neatly printed in pearl, 8s. 6d. cl.; or 10s. 6d. embossed and gilt.

INSTRUCTIONS to YOUNG SPORTSMEN.

By Lieut.-Col. P. HAWKER.

A New Edition, thoroughly revised, with new Cuts, Views of the Heads of the Wild and Tame Swans, &c. is in preparation, and will be ready before the Sporting Season.

"Col. Hawker is one of the best shots in England, and his 'Instructions to Sportsmen' the very best book we have on the subject"—BLACKWOOD'S MAGAZINE.

THE GUN;

Or, a Treatise on the Nature, Principle, and Manufacture of the various descriptions of Small Fire Arms.

By WILLIAM GREENER. 1 vol. 8vo. Plates, 15s.

"I have no hesitation in saying that I consider it by far the best work ever written on the subject, and I should recommend the perusal of it to every gun-maker in the United Kingdom."
LETTER FROM COLONEL HAWKER.

HINTS to MOTHERS, for the Management of Health.

By THOS. BULL, M.D. Physician-Accoucheur to the Finsbury Midwifery Institution, &c.
Fcp. 8vo. 5s. cloth lettered.

"There is no mother that will not be heartily thankful that this book ever fell into her hands; and no husband who should not present it to his wife. We cannot urge its value too strongly on all whom it concerns."—ECLECTIC REVIEW.

WORKS OF GENERAL UTILITY—CONTINUED.

TOMLINS'S POPULAR LAW DICTIONARY.

1 thick volume, post 8vo. 18s. cloth lettered.

"A volume of the greatest utility, and affords a vast mass of information essentially necessary to almost every man in his ordinary occupations of life."—WEEKLY DISPATCH.

PLAIN DIRECTIONS for making WILLS

In conformity with the NEW ACT, which came into operation on the 1st January.

By J. C. HUDSON, of the Legacy Duty Office, London.

6th Edition, fcp. 8vo. price Half-a-Crown, neatly done up in cloth, gilt edges.

By the same Author,

THE EXECUTOR'S GUIDE.

Fcp. 8vo. 5s. in cloth, gilt edges.

HINTS on ETIQUETTE;

AND THE USAGES OF SOCIETY, &c.

By Αγαγός.

16th Edition, fcap. 8vo. 2s. 6d. gilt edges.

SHORT WHIST; its Rise, Progress, and Laws :

Together with Maxims for Beginners, and Observations to make any one a Whist Player.

By MAJOR A*****.

5th Edition, fcap. 8vo. with Frontispiece, 3s. in fancy cloth, gilt edges.

INSTRUCTIONS to EXECUTORS

And ADMINISTRATORS,

Shewing the Duties and Responsibilities incident to the due Performance of their Trusts; with Forms properly filled up for every Bequest.

By J. H. BRADY, late of the Legacy Duty Office, Somerset House.

8vo. 6th Edit. improved to the latest date, 8s.

*** This work contains information of a more Technical Nature than Mr. Hudson's work, and is therefore adapted to the use of Attorneys as well as the Public.

DOMESTIC DUTIES ;

Or, Instructions to Young Married Ladies on the Management of their Households, &c. &c.

By Mrs. WILLIAM PARKES. 12mo. 4th Edition, 10s. 6d. cloth lettered.

"The volume before us is a perfect VADE MECUM for the young married lady, who may resort to it on all questions of household economy and etiquette."—NEW MONTHLY MAG.

ON THE

VALUATION of PROPERTY for the POOR'S RATE.

By J. S. BAYLDON. 8vo. New Edition, enlarged, 7s. 6d.

By the same Author,

Art of Valuing Rents and Tillages, &c. New Edition, 7s. bds.

THE COMPLETE LATIN-ENGLISH DICTIONARY,

Compiled from the best sources, chiefly German, and adapted to the use of Colleges and Schools.

By the Rev. E. RIDDLE.

8vo. price £1. 1s. cloth boards.

"An admirable Dictionary, and promises more help to the Latin student than any thing hitherto offered him in this country."—EVANGELICAL MAGAZINE.*"We do not think we could use terms too high to express our entire approbation and admiration of it. It is a vast improvement in its class, and does infinite credit to the talents and learning of Mr. Riddle."*—LIT. GAZETTE.

§§§ The above is abridged for the use of Schools.

*** The ENGLISH-LATIN portion is nearly ready.

ARCHITECTURE, ROAD-MAKING,

&c.

DICTIONARY of the ARCHITECTURE and ARCHÆOLOGY of the MIDDLE AGES ;

Including the Words used by Old and Modern Authors.

By JOHN BRITTON, F.S.A. &c.

This work will be completed by the publication of the Fourth Part, on the 1st of July.

T. F. HUNT'S ARCHITECTURAL WORKS.

Hints on Picturesque Domestic Architecture; in a Series of Designs for Gate Lodges, Gamekeepers' Cottages, &c. 4to. New Edition, 21s.

Designs for Lodges, Gardeners' Houses, and other Buildings, in the Modern or Italian Style. 4to. 21s.

Designs for Parsonage-Houses, Almshouses, &c. 4to. 21s.

Exemplars of Tudor Architecture, adapted to Modern Habitations; with illustrative Details, &c. 4to. £2. 2s.

The above may be had with Proof Impressions of the Plates on India Paper.

ELEMENTS of MUSICAL COMPOSITION ;

Comprehending the Rules of Thorough Bass, and the Theory of Tuning.

By WILLIAM CROTCH, Mus. Doc. Professor of Music in the University of Oxford.

2d Edition, small 4to. with Plates, 12s. in cloth.

By the same Author,

LECTURES on MUSIC. 7s. 6d.

ENCYCLOPÆDIA of COTTAGE, FARM, & VILLA ARCHITECTURE and FURNITURE.

By J. C. LOUDON, F.L.S. &c.

New Edition, 1 large vol. 8vo. with above 11,000 pages of Letterpress, and illustrated with upwards of 2000 Engravings, £3. boards.

A TREATISE on ROADS :

Wherein the principles on which Roads should be made are explained and illustrated, by the Plans, Specifications, and Contracts, made use of by Thos. Telfourd, Esq. on the Holyhead Road.

By the Rt. Hon. Sir HENRY PARNELL, Bart. Hon. Mem. Inst. Civ. Engin. Lond.

2d Edition, greatly enlarged.

PRACTICAL TREATISE on RAILROADS

And INTERIOR COMMUNICATION in GENERAL.

Containing the Performances of the improved Locomotive Engines: with Tables of the Comparative Cost of Conveyance on Canals, Railways, and Turnpike Roads.

By NICHOLAS WOOD, Colliery Viewer, Mem. Inst. Civ. Engin. &c.

3d Edition, very greatly enlarged, and illustrated by several new Engravings.

TREATISE on the STEAM ENGINE ;

Historical, Practical, and Descriptive.

By JOHN FAREY, Engineer.

4to. illustrated by numerous Woodcuts, and 25 Copperplates, engraved by Wilson Lowry, from Drawings by Messrs. Farey.

£5. 5s. bds.—VOL. II. is in the press.

REMARKS on the PRESENT SYSTEM of ROAD-MAKING.

By JOHN LOUDON M'ADAM.

8vo. 7s. 6d. boards.

WILSON AND SON, PRINTERS, 57, SKINNER STREET, LONDON.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00089 9930