ADVANCES IN HERPETOLOGY AND EVOLUTIONARY BIOLOGY

Essays in Honor of
Ernest E. Williams

EDITED BY
ANDERS G.J. RHODIN
KENNETH MIYATA

ADVANCES IN HERPETOLOGY
AND
EVOLUTIONARY BIOLOGY

Ernest E. Williams

ADVANCES IN HERPETOLOGY AND EVOLUTIONARY BIOLOGY

Essays in Honor of
Ernest E. Williams

EDITED BY
ANDERS G.J. RHODIN
KENNETH MIYATA

1983
Museum of Comparative Zoology
Cambridge, Massachusetts

Museum of Comparative Zoology
Harvard University
Cambridge, Massachusetts 02138 U.S.A.

Copyright © 1983 by the Presidents and Fellows of Harvard College

ISBN 0-910999-00-7

Library of Congress Card Catalog Number 73722

All rights reserved
Printed in the United States

PREFACE

In July of 1980 Ernest E. Williams officially retired as Curator of Herpetology at the Museum of Comparative Zoology at Harvard University, a position he held for a quarter of a century. Ernest is particularly fond of parties, especially if champagne and various delicacies are served, and so we marked his retirement with a celebration in the Romer Hall of Fossil Vertebrates on a warm summer afternoon. A host of his close associates and former graduate students converged on Cambridge to have a good time (some may have regretted their efforts the next day) and to visit a man whom we feel is unique. On that day, we had the pleasure of springing a major surprise on Ernest: this volume in his honor.

This Festschrift was initially conceived as an informal homage to a man whom so many respect. We contacted as many of his former students and colleagues as possible and solicited papers for possible publication in the volume. The response was overwhelming, and it quickly became obvious that the Festschrift would be a tome of major proportions. The Museum of Comparative Zoology and its director, Dr. A. W. Crompton, then generously provided the support necessary for an undertaking of this magnitude, and at the party we were able to present Ernest with a prospectus of the forthcoming volume. It was one of the few times we have seen him show surprise.

Though the majority of his own work has been in herpetology, focusing primarily on museum and field investigations, Ernest's interests encompass much of modern evolutionary biology. This volume is a testament to the extraordi-

nary breadth of his interests in the evolutionary biology of both vertebrates and invertebrates. More than that, it is a testament to the enormous influence he has had on students and colleagues in fostering this wide range of interests. Ernest's ability to encourage and stimulate research is remarkable, and we can think of no one else who has influenced such a diversity of perspectives, approaches, and personalities. His openness to new ideas, his willingness to tolerate unusual approaches and behavior, and his ability to stimulate and guide without making it obvious are all unusual, if not unique, traits in a man of his stature. We doubt that a single one of his associates has ever regretted working with him, and more than one graduate student has characterized him as being the "best of all possible major professors."

Although he is now retired as Curator of Herpetology, Williams remains a Professor of Biology and his research continues unabated. This past year he received his second Guggenheim Fellowship, and he has been busy collaborating with colleagues across two continents. Undoubtedly, his influence will continue to grow in the years to come.

Reflecting Ernest's diverse interests, this volume in his honor encompasses a wide spectrum of research investigations. There are 50 papers by 69 authors, split into seven broad topics: Systematics, Comparative Morphology, Zoogeography, Ecology, Behavior, Evolution, and Conservation. The papers deal with an enormous array of animal taxa, from protozoans and arthropods to birds, bats and primates. However, the bulk of the volume deals with reptiles

and amphibians, as do most of Ernest's own research investigations. There are 49 genera and 144 species of amphibians discussed, including one new genus and three new species. There are 338 genera and 643 species of reptiles represented, including one new genus, four new species and one new subspecies. Within the lizard genus *Anolis*, central among Ernest's interests, there are 97 taxa mentioned in the text.

We offer this Festschrift volume in tribute to Ernest E. Williams and join with the authors represented here in expressing our thanks for his friendship, his support of the investigative process in biology, and his stimulation of zoological inquiry among those colleagues fortunate enough to be within his sphere of influence.

ACKNOWLEDGMENTS. We wish to thank all those individuals who helped with the enormous task of the review process: Robin M. Andrews, Stephen C. Ayala, John L. Behler, John W. Bickham, Robert Bleiweiss, Peter Brazaitis, Donald Brinkman, Donald G. Broadley, Lisa Brooks, Walter C. Brown, Charles C. Carpenter, Robert L. Carroll, Susan M. Case, William E. Cooper, James R. Dixon, C. Kenneth Dodd, Jr., William E. Duellman, Eugene S. Gaffney, Alfred L. Gardner, Alice G. C. Grandison, Neil Greenberg, Allen E. Greer, Max K. Hecht, Paul E. Hertz, Robert D. Holt, Raymond B. Huey, Robert F. Inger, Farish A. Jenkins, Jr., Thomas A. Jenssen, Kenneth V. Kardong, A. Ross Kiestner, F. Wayne King, John Kirsch, George Lauder, Raymond F. Laurent, John D. Lynch, Gregory C. Mayer, Timothy C. Moermond, Scott M. Moody, Charles W. Myers, Thomas S. Parsons, Raymond A. Paynter, Jr., Peter C. H. Pritchard, A. Stanley Rand, Jonathan Roughgarden, Rodolfo Ruibal, Anthony

P. Russell, Joseph J. Schall, Thomas W. Schoener, Albert Schwartz, Kathleen K. Smith, William Stubblefield, Richard Thomas, Garth Underwood, Romulus Whitaker, Don E. Wilson, Roger Conant Wood, and George R. Zug. We also wish to thank several colleagues of Williams who extended their encouragement in this commemorative effort, but were not among the contributors or reviewers: Archie Carr, David Crews, Arthur C. Echternacht, Richard Etheridge, Kurt Fristrup, Jose Maria Gallardo, William Garstka, Terry E. Graham, W. Ronald Heyer, Paul F. A. Maderson, Ernst Mayr, George Meszoly, Juan A. Rivero, George Gaylord Simpson, Hobart M. Smith, Barbara Wu, Rainer Zangerl, and Richard G. Zweifel.

In addition, we wish to express our gratitude to those people who helped in the overall production of this volume: A. W. Crompton, who provided welcome institutional and moral support; Raymond A. Paynter, Jr., who shepherded the volume through its early editorial organizational stages; Elizabeth Campbell Elliott, who did a superb job as technical editor for the volume; Jose Rosado, James Knight, and Catherine McGeary of the Herpetology Section, who assisted with certain technical aspects; Pere Alberch, Williams's successor as Curator of Herpetology, for his continuing support; Laszlo Meszoly, who designed the book cover; and Susan DeSanctis Rhodin, who provided most of the initial impetus to mobilize this massive effort, then prepared the feast which marked the retirement party and presentation of the prospectus, and finally provided the patient support and encouragement needed to bring the volume to its fruition.

Anders G. J. Rhodin
Kenneth Miyata

INTRODUCTION

Ernest Edward Williams was born in 1914 in Easton, Pennsylvania, the only child of middle-aged parents. He attended Lafayette College, where one of his teachers was B. W. Kunkel, a capable vertebrate anatomist and an important early influence. After obtaining his B.S. at the age of nineteen, financial difficulties forced Williams to partially interrupt his studies for several years. Fortunately, having fought in World War II, he was able to fully resume in 1946 thanks to a "G.I. Bill" studentship. He enrolled at Columbia University for a Ph.D. under W. K. Gregory, the father image of vertebrate morphology in those days. However, Gregory retired soon after, and Williams proceeded on his own.

Williams's thesis subject, originating from his association with Kunkel during the Lafayette days, was variation and selection in the cervical central articulations of living turtles. The work was done mostly at the American Museum of Natural History, where Williams met a number of his future colleagues and collaborators, including Samuel B. McDowell, Max K. Hecht, Karl Koopman and Rodolfo Ruibal.

The thesis was presented in 1949 and published in 1950. It is the work of a fine anatomist with unusually broad views and interests. Unfortunately, a myopic committee mutilated it, excising a final chapter on the relationships between selective pressures and variability that, in reality, had against itself being ahead of the times. The thesis contains, as an appendix, a new classification of turtles to the subfamily level; this type of "appendix" was an American Museum

trademark that was to become part of Williams's style.

Williams's first employment was as an instructor in A. S. Romer's course in Comparative Anatomy at Harvard (Romer was also a former Gregory student). This is how his association with the Museum of Comparative Zoology (and, incidentally, with me) began.

Kunkel had gotten Williams interested in turtles, and this was a field well suited to his temperament—inquisitive, shy, and independent. It was not crowded, materials being abundant and easily made available. It afforded access to a broad range of subjects, from embryology and comparative anatomy to paleontology, and leading into systematics above the species level and zoogeography. Between 1950 and 1961 Williams published 23 papers on chelonological matters; one was coauthored with McDowell and another with T. S. Parsons, both graduate students at the time. In 1954 he received a Guggenheim fellowship to study fossil turtles in European institutions, especially the British Museum. The results then obtained stand to this day, as witnessed, for example, by Gaffney's contribution to this volume, one of its eight papers on turtles.

The anatomy of the chelonians, especially the osteology of the head and neck, led Williams into similar investigations of other reptilian groups, a field in which he published some highly original papers (e.g., on Gadow's theory of arcualia and on the occipito-vertebral joint) and to which he has sporadically returned. The articles in this volume by Allen Greer, a former student, and by T. Frazzetta, a col-

laborator since 1970, are a reflection of this phase.

Some of the turtle work on fossils from Cuba and elsewhere started Williams's involvement with the West Indies. A 1950 trip to Jamaica strengthened his interest in the area and marked the beginning of his work on lizards, in collaboration initially with Garth Underwood, then teaching in Jamaica, and Rodolfo Ruibal, a Cuban who was living in the U.S. but keeping close ties with the island. Williams's interest in the Antilles in the beginning centered on paleontology: besides the turtle work there are early papers on fossil and sub-fossil mammals with Koopman and with Reynolds and Koopman. Another of the West Indian papers, on a fossil tortoise from Mona, contains the currently accepted classification of the Testudinidae which originated the famous footnote by Loveridge in the coauthored revision of the African cryptodires. In this footnote Loveridge contrasted himself, a "working herpetologist," with his coauthor, who represented a "current trend in herpetology." This was a perceptive statement, as the next phase of Williams's career consisted essentially of contributing to dilate the borders of lacertilian systematics to coincide with those of biology in general. In the process, the standards of "working herpetology" were considerably raised and strengthened.

The instructorship at Harvard was for a period of three years, after which Williams was promoted in 1952 to Assistant Professor, having added to his former duties those of coordinating Romer's Evolution course in General Education. Romer really did not care much for the course, and Williams's appointment was more than anything a device to keep him at Harvard until Loveridge's retirement from the herpetology curatorship. It was an obvious step. Williams had been working on the collection, both on recent turtles and on anatomical matters, and was beginning to become interested in lizards. Also, his curatorial instincts were awakening.

Loveridge had arrived at the MCZ in the early twenties, essentially to renovate the herpetological collections. This he rapidly did, and he also promoted a program of exchanges that could be called intensive for its time, emphasizing as complete a representation as possible of all faunas, but definitely not long series. An uncommon lover of tidiness, Loveridge had meticulously planned his life in order to have completed the corpus of a Herpetology of East Africa by the time he was due to retire. (That he was about one year late he stoutly credited to the coauthored cryptodiran monograph.) This meant full concentration on African matters, with only a few brief digressions into other faunas. As a result, Williams inherited in 1957 a collection that was systematically very complete, but not especially rich in adequate series or skeletal materials. The department library, successively kept and added to by Agassiz, Garman, and Barbour, and further enriched by the addition of Loveridge's library (purchased by Williams and donated to Harvard) constituted an extremely valuable resource, but it was severely restricted to systematics at the alpha level.

After taking over the curatorship, Williams immediately began a program of strengthening the general collections, of building and steadily adding to a representative skeletal collection. Concomitantly, the library grew both in size and scope, embracing much of ecology and evolutionary biology. In conjunction with this, he began taking on graduate students.

Having become a curator contributed somewhat to a cooling of Williams's enthusiasm for turtle taxonomy. For instance, he and I had planned a revision of the South American chelids; the available materials were borrowed, innumerable measurements taken, and graphs prepared. The project was dropped, however, due to the absence of significant series and the logistic problems in handling any quantity of these very awkward animals. More recently, this research

has been resurrected by Rhodin and Mittermeier (this volume).

Among the reputedly strongest parts of the MCZ herpetological collections was that of West Indian anoles, mostly assembled by Thomas Barbour. To it Williams turned his attention. His initial aim was properly broad: to study the genus "in depth and at several levels, not only in terms of the descriptive taxonomy but also in terms of ecology and behavior."

The first four years of anoline work consisted of intensive collecting in the Greater and Lesser Antilles, clearing up the alpha systematics as far as possible, and getting acquainted with the types of problems that mere work on museum collections cannot reveal. Research was also started on the continent of South America; as I see it, this rapidly became important not so much in itself, but chiefly in focusing attention on the problem of generic borders in the Iguanidae, a subject which to this day looms large in the work of Williams and his associates (Estes, Hall, Moody, in this volume). By this time (1964) the groupings within the anoles also began to be envisaged, stemming from Etheridge's initially unpretentious concept of alpha and beta fracture plane types in tail vertebrae.

Another initial preoccupation was with digital pads, so characteristic of anoles. This line of work has progressed with moderate impetus, but fine results (Peterson, this volume).

The next phase of work (say to 1970) brought about a sharpening of the focus. The main problems perceived were those related to competition: coadaptation, exclusion, fragmentation. A very important result of this phase was Williams's classic paper on the ecology of colonization, published in 1969 in the *Quarterly Review of Biology*. By this time Schoener had begun his detailed studies on resource sharing (Moermond, this volume). Cytogenetics and biochemistry were adopted as everyday tools, in the spirit of the times with a bit of excessive optimism ("the ultimate

tool"), that still lasts in some members of the group (Gorman *et al.*, this volume). Thermoregulation, a field that seemed at first extremely promising, as I see it, failed to fulfill its promise.

By 1972 Williams had performed the first test of a model of an intra-island radiation. Central to the analysis was the concept of "ecomorphs"—massive opportunism in diverse lineages leading to convergence and parallelism to an extreme degree. The concept, to which Williams had been led by his joint work with Stan Rand, was not intrinsically new, but had previously been applied only to groups differing at least at the family level and inhabiting broadly separated geographic areas. Its application to members of a single genus within a single zoogeographic subregion required an enormous fund of information, and I believe West Indian anoles constitute the only viable case so far.

The following (and current) steps of the concerted attack on West Indian anoles address still more basic biological problems: species borders and species interactions, radiation, and species packing. "Most species packing in West Indian *Anolis* has been achieved by speciation rather than by accumulation of already differentiated colonizing species, and incompletely speciated populations still exist on West Indian islands. A background for studies of speciation in West Indian anoles is provided by the considerable information at hand on the taxonomy, ecology, behavior, biogeography, and morphological and karyotypic evolution of West Indian anoles."

In 1970 Williams was appointed Professor of Biology, thus automatically receiving a Harvard M.A. and becoming, in Harvard parlance, a "child of the house," no longer a foreigner. In 1972 he was made Agassiz Professor, a position of real significance, as witnessed by a list of fellow incumbents, including, among others, A. S. Romer, P. J. Darlington, Jr., Bryan Patterson, and Frank Carpenter. Since his assumption of the curatorship, the Harvard system had also allowed him

to have graduate students. In keeping with Harvard administrative tradition, it is a moot point who was Williams's first student, Carl Gans or Stan Rand. However that may be, there has been a steady flow ever since, and the type of relationship between professor and students has always been to me a matter of admiration. The main characteristic of the group has been diversity in concentration, i.e., to bring all sorts of approaches to bear on a theoretical problem embodied in a systematic unit. Within this very broad mandate there has been full freedom to pursue individual lines of thought. In fact, I can think of no other environment where a man as singular as Robert Trivers could have matured his signal contributions to evolutionary theory.

I have stated above that in the fifties zoology, as practiced in museums, and especially in the U.S., underwent a marked change in emphasis and even in scope, embracing theoretical and experimental aspects that had previously been outside the range of the profession. A notable forerunner was Charles M. Bogert, of the American Museum of Natural History, who, besides impeccable field and taxonomic research, did basic work in such new areas as thermoregulation in reptiles and acoustic com-

munication in amphibians. The trend developed rapidly; notwithstanding all the favorable aspects, in one important respect the consequences have been less fortunate. Emphasis on the experimental approach, demanding a different type of training, entailing stiff academic competition and an element of fashion, has tended to lower the prestige and the quality of curatorship. It has become in many cases increasingly difficult to base museum policy on the good of the collections—even their intrinsic worth has been questioned. Williams has been a steadfast and articulate advocate of the collection as the heart and kernel of the museum department, and its most enduring legacy. For Williams, the basic task of a curator is still to enlarge, structure, and make available a collection. His career has been better known through a consistent broadening and deepening of scientific preoccupations; that this has gone hand in hand with a definite philosophy and practice of selfless curatorship rounds off a complete and entire scientific personality. The esteem and respect evinced in this volume bear ample witness to this personality.

P. E. Vanzolini

LIST OF CONTRIBUTORS

ANGEL C. ALCALA

Professor of Biology and Vice President for Academic Affairs, Silliman University, Dumaguete City, Philippines. M.A., Ph.D. (herpetology and ecology), Stanford University. Association with E. E. Williams indirect through W. C. Brown. Research interests include herpetology and ichthyology and conservation of Philippine land vertebrates and marine mammals.

ROBIN M. ANDREWS

Associate Professor of Biology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061; Research Associate, Smithsonian Tropical Research Institute, Balboa, Panama. Ph.D. (zoology), University of Kansas. Post-doctoral fellowship under E. E. Williams and A. S. Rand. Research interests include life history evolution of reptiles, physiological ecology of reptiles, especially with regards to eggs, energetics and thermoregulation.

WALTER AUFFENBERG

Curator in Herpetology, Florida State Museum, University of Florida, Gainesville, Florida 32611. M.S. (herpetology), Ph.D. (paleontology of reptiles), University of Florida. Former special student and teaching assistant under E. E. Williams at Harvard. Research interests include behavioral ecology of varanid lizards and paleontology and behavior of tortoises.

ANGUS D'A. BELLAIRS

Emeritus Professor of Vertebrate Morphology, St. Mary's Hospital Medical School, University of London, Paddington, London, W.2., United Kingdom. D.Sc., London; M.R.C.S. (Member of the Royal College of Surgeons), London. Professional associate of E. E. Williams for many years. Research interests include the development, structure and evolution of the vertebrate skeleton and biology of the Reptilia.

JUDITH A. BLAKE

Curatorial Assistant, Division of Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138; Biologist, Ruecker Wildlife Refuge, 137 Seapowet Ave., Tiverton, Rhode Island 02878. Ph.D. (evolutionary biology), Harvard University under E. E. Williams. Research interests include evolutionary biology, genetics of speciation and conservation biology.

DONALD G. BROADLEY

Curator of Herpetology, National Museum,

Bulawayo, Zimbabwe. M.Sc., Ph.D. (herpetology), University of Natal. Professional associate of E. E. Williams for many years. Research interests include taxonomy, ecology and zoogeography of the reptiles of Africa.

WALTER C. BROWN

Emeritus Professor of Biology, Menlo College, Menlo Park, California 94025; Fellow and Research Associate in Herpetology, California Academy of Science, San Francisco, California 94118. Ph.D. (biology), Stanford University. Professional associate of E. E. Williams for many years. Research interests include systematics, evolution and ecology of amphibians and reptiles, with special interest in the Philippines and the Pacific.

DONALD G. BUTH

Assistant Professor of Biology, Department of Biology, University of California, Los Angeles, California 90024. Ph.D. (ecology, ethology and evolution), University of Illinois under T. H. Frazzetta (former post-doctoral fellow under E. E. Williams at Harvard). Association with E. E. Williams through both T. H. Frazzetta and G. C. Gorman. Research interests include biochemical systematics of reptiles and fishes.

ROGER CONANT

Adjunct Professor, Department of Biology, University of New Mexico, Albuquerque, New Mexico 87131; Research Associate, Academy of Natural Sciences, Philadelphia, and American Museum of Natural History, New York; Past-President and former Secretary, American Society of Ichthyologists and Herpetologists; Director and Curator of Reptiles Emeritus, Philadelphia Zoological Garden. Honorary Sc.D., University of Colorado. Professional associate of E. E. Williams for many years. Research interests include herpetology of the Chihuahuan desert and biology of *Agkistrodon*.

ISABEL D. CONSTABLE

Undergraduate Student, Brown University, Providence, Rhode Island 02912. Association with E. E. Williams indirect through R. A. Mittermeier; also, father associated with A. Loveridge as undergraduate student in herpetology at the MCZ. Research interests include primates, tropical forests and wildlife conservation.

JAMES R. DIXON

Professor and Chief Curator, Texas Cooperative Wildlife Collections, Department of Wild-

life and Fisheries Sciences, Texas A & M University, College Station, Texas 77843; Past-President, Society for the Study of Amphibians and Reptiles; Board of Governors, American Society of Ichthyologists and Herpetologists; Chairman, Board of Scientists, Chihuahuan Desert Research Institute. Ph.D. (zoology), Texas A & M University. Professional associate of E. E. Williams for many years. Research interests include herpetology of the Americas.

LUDMELA DJATSCHENKO

Dentist, 282 West Acres Dr., Guelph, Ontario, N1H 7P1, Canada. D.D.S., University of Toronto. Association with E. E. Williams indirect through T. S. Parsons.

WILLIAM E. DUELLMAN

Curator, Division of Herpetology, Museum of Natural History, and Professor, Department of Systematics and Ecology, University of Kansas, Lawrence, Kansas 66045; Past-President, Herpetologists' League. Ph.D. (zoology), University of Michigan. Professional associate of E. E. Williams for many years. Research interests include evolutionary biology of amphibians with emphasis on systematics, biogeography, and reproductive biology of Neotropical anurans.

PAUL ELIAS

Field Editor for Europe, Advanced Book Program, Addison-Wesley Publishing Co., Reading, Massachusetts 01867. M.A. (zoology), University of California, Berkeley; A.M. (biology), Harvard University under P. Alberch (E. E. Williams's successor as Curator of Herpetology). Research interests include the constraints imposed upon evolution by development and their ramifications in systematics.

NELLY CARRILLO DE ESPINOZA

Chief, Section of Herpetology, Museo de Historia Natural "Javier Prado," Universidad Nacional Mayor de San Marcos, Avenida Arenales 1256, Lima, Peru. Doctor (herpetology), Universidad Nacional Mayor de San Marcos. Professional associate of E. E. Williams for many years. Research interests include Neotropical snakes and lizards.

RICHARD ESTES

Professor of Zoology, San Diego State University, San Diego, California 92182, and Research Associate, Museum of Paleontology, University of California, Berkeley, California 94720; Former Research Associate, Museum of Comparative Zoology, Harvard University. Ph.D. (paleontology), University of California, Berkeley. Association with E. E. Williams through research at the MCZ. Research interests include morphology and systematics of lower vertebrates, paleoherpetology, faunal evolution and biogeography.

KARL F. FORSGAARD

Attorney, Houghton Cluck Coughlin & Riley, 1111 Third Ave., Suite 1500, Seattle, Washing-

ton 98101. Association with E. E. Williams as undergraduate student at Harvard and former curatorial assistant in herpetology at the MCZ. Research interests include high-altitude and polar zoology, colubrid and iguanid ecology and taxonomy, and wildlife protection law.

THOMAS H. FRAZZETTA

Professor of Ecology, Ethology and Evolution, University of Illinois, Urbana, Illinois 61801; Associate in Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138. Ph.D. (vertebrate zoology), University of Washington. Post-doctoral fellowship under E. E. Williams at Harvard. Research interests include evolutionary biology, biomechanics of vertebrate jaws, especially reptiles and galeoid sharks, population biology.

EUGENE S. GAFFNEY

Curator, Department of Vertebrate Paleontology, American Museum of Natural History, 79th St. and Central Park West, New York, New York 10024. Ph.D. (geology), Columbia University. Professional associate of E. E. Williams for many years. Research interests include turtle systematics.

CARL GANS

Professor of Zoology, The University of Michigan, Ann Arbor, Michigan 48109; Research Associate, Carnegie Museum, Pittsburgh, and American Museum of Natural History, New York; former Guggenheim Fellow; Past-President, American Society of Ichthyologists and Herpetologists. M.S. (engineering), Columbia University; Ph.D. (biology), Harvard University under E. E. Williams (his first graduate student). Research interests include functional morphology, reptilian adaptations.

GEORGE C. GORMAN

Professor of Biology, University of California, Los Angeles, California 90024; former Research Associate, Museum of Comparative Zoology, Harvard University. Ph.D. (herpetology), Harvard University under E. E. Williams. Research interests include evolution of *Anolis*, population genetics, biochemical evolution, reproductive biology of reptiles.

ALLEN E. GREER

Curator of Reptiles and Amphibians, The Australian Museum, 6-8 College St., Sydney, New South Wales 2000, Australia. Ph.D. (biology), Harvard University under E. E. Williams. Post-doctoral fellowship under E. E. Williams at Harvard. Research interests include biology and evolution of reptiles, especially squamates.

STELLA GUERRERO

Graduate Student, Department of Entomology and Parasitology, University of California, Berkeley, California 94720; former Research Assistant, Smithsonian Tropical Research Institute, Balboa, Panama. Association with E. E. Williams indirect through A. S. Rand and R. M.

Andrews. Research interests include ecology and effects of parasites on wild populations, especially lizards.

WILLIAM P. HALL

Post-Doctoral Fellow, Queen's College, University of Melbourne, Parkville, Victoria 3052, Australia; former Visiting Assistant Professor of EPO Biology, University of Colorado, Boulder; former Visiting Lecturer, University of Maryland, College Park. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include functional relationships between fixation of chromosomal mutations and speciation in iguanid lizard radiations, and roles of hybrid zones in maintaining genetic isolation in lizards and frogs.

MAX K. HECHT

Professor of Biology, Queens College, City University of New York, Flushing, New York 11367. Ph.D. (vertebrate zoology), Cornell University. Association with E. E. Williams through field work in the West Indies. Research interests include vertebrate systematics.

PAUL E. HERTZ

Assistant Professor of Biological Sciences, Barnard College, Columbia University, 606 West 120th St., New York, New York 10027. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include reptile ecology, the evolution of thermal biology, and mechanisms of adaptation.

ROBERT A. HICKS

P.O. Box 420, Ipswich, Massachusetts 01938; former Employee, Massachusetts Audubon Society, National Marine Fisheries Service, and Marine Resources Conservation Foundation. B.A., Harvard University, field research assistant with E. E. Williams and R. L. Trivers in the West Indies.

ROBERT D. HOLT

Assistant Professor, Department of Systematics and Ecology, and Assistant Curator, Museum of Natural History, University of Kansas, Lawrence, Kansas 66045. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include theoretical ecology, island biology, predator-prey interactions and tropical ecology of birds and lizards.

RAYMOND B. HUEY

Associate Professor of Zoology, Department of Zoology NJ-15, University of Washington, Seattle, Washington 98195. M.A. (zoology), University of Texas; Ph.D. (biology), Harvard University under E. E. Williams. Research interests include physiological ecology, defensive behavior and interspecific competition.

ROBERT F. INGER

Curator, Amphibians and Reptiles, Field Museum of Natural History, Chicago, Illinois 60605. Ph.D. (zoology), University of Chicago. Professional associate of E. E. Williams for many years. Research interests include evolu-

tionary ecology and systematics of Southeast Asian amphibians and reptiles and structure and organization of communities of reptiles and amphibians in tropical forests.

THOMAS A. JENSSEN

Associate Professor of Zoology, Department of Biology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061. Ph.D. (zoology), University of Oklahoma. Post-doctoral fellowship under E. E. Williams at Harvard. Research interests include behavioral ecology of amphibians and reptiles, especially display behavior of *Anolis*.

A. ROSS KESTER

Assistant Professor of Biology, Department of Biology, Tulane University, New Orleans, Louisiana 70118. Ph.D. (biology), Harvard University under E. E. Williams. Junior Fellow, Society of Fellows, Harvard. Research interests include evolutionary and population biology.

KARL F. KOOPMAN

Curator, Department of Mammalogy, American Museum of Natural History, 79th St. at Central Park West, New York, New York 10024. Ph.D. (zoology), Columbia University. Fellow graduate student and teaching assistant with E. E. Williams at Columbia, professional associate for many years. Research interests include systematics and zoogeography of Chiroptera.

RAYMOND F. LAURENT

Professor and Principal Investigator, Consejo Nacional de Investigaciones Científicas y Técnicas, Fundación Lillo, Miguel Lillo 205, 4000 Tucumán, Argentina; former Associate Professor, Museum National d'Histoire Naturelle, Paris; former Research Zoologist with E. E. Williams at Harvard. Ph.D. (zoology), University of Bruxelles. Research interests include African and South American herpetology, especially systematics, evolution and zoogeography.

JAMES D. LAZELL, JR.

President, The Conservation Agency, 8 Swinburne St., Conanicut Island, Rhode Island 02835; Associate in Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138. M.A. (biology), Harvard University under E. E. Williams; M.S. (zoology), University of Illinois; Ph.D. (biology), University of Rhode Island. Research interests include field and evolutionary biology of reptiles, amphibians, and mammals.

JOHN D. LYNCH

Professor of Life Sciences, School of Life Sciences, University of Nebraska, Lincoln, Nebraska 68588. Ph.D. (zoology), University of Kansas. Professional associate of E. E. Williams for many years. Research interests include systematics and biogeography of neotropical frogs, especially *Eleutherodactylus*.

GREGORY C. MAYER

Graduate Student, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138. Ph.D. candidate (biology), Harvard University under E. E. Williams. Research interests include the genetics, ecology and geography of speciation.

SAMUEL B. McDOWELL

Professor of Zoology, Rutgers University, Newark, New Jersey 07102. Ph.D. (zoology), Columbia University under G. G. Simpson. Fellow graduate student with E. E. Williams, illustrated his Ph.D. thesis, co-authored an early paper on turtle morphology, continued professional association concerning relationships of West Indian insectivores. Research interests include New Guinea reptiles, evolution of the vertebrate skull, origin of snakes, notacanthiform fishes, lipotyphlan mammals.

FEDERICO MEDEM

Professor, Universidad Nacional de Colombia, Bogotá, and Director, Estación de Biología Tropical "Roberto Franco," Apartado aereo 22-61, Villavicencio (Meta), Colombia; Member, IUCN/SSC Steering Committee and Crocodile Specialist Group; former Guggenheim Fellow. Dr. rer. nat. (zoology, paleontology, geology), Humboldt University, Berlin. Professional associate of E. E. Williams for many years. Research interests include ecology and zoogeography of turtles and crocodiles.

RUSSELL A. MITTERMEIER

Chairman, IUCN/SSC Primate Specialist Group and Director, World Wildlife Fund—U.S. Primate Program, World Wildlife Fund—U.S., 1601 Connecticut Ave., NW, Washington, D.C. 20009; Associate in Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts; Vice-Chairman, IUCN/SSC Freshwater Chelonian Specialist Group. Ph.D. (biological anthropology), Harvard University under I. DeVore. Research interests include primatology, herpetology and conservation.

KENNETH MIYATA

Post-Doctoral Fellow, Division of Amphibians and Reptiles, Smithsonian Institution, National Museum of Natural History, Washington, D.C. 20560; Research Associate in Vertebrate Zoology, Smithsonian Institution; Associate in Herpetology, Museum of Comparative Zoology, Harvard University. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include biogeography and ecology of neotropical herpetofauna.

TIMOTHY C. MOERMOND

Associate Professor, Department of Zoology, University of Wisconsin, Madison, Wisconsin 53706. Ph.D. (evolutionary biology), Harvard University under E. E. Williams. Research interests include feeding behavior of animals, optimal foraging strategies and community

structure, especially in *Anolis* lizards and birds, co-evolution of frugivorous birds and bird-dispersed plants.

SCOTT M. MOODY

Assistant Professor of Biomedical Science, Department of Zoology and College of Osteopathic Medicine, Ohio University, Athens, Ohio 45701. M.S., Ph.D. (evolutionary and ecological biology), University of Michigan. Post-doctoral fellowship, Deutscher Akademisches Austauschdienst. Association with E. E. Williams as undergraduate student at Harvard and former curatorial assistant in herpetology at the MCZ. Research interests include comparative and functional morphology and phylogenetics, zoogeography and paleontology of lizards, especially Agamidae.

JÓSE ALBERTO OTTENWALDER

Curator, Department of Mammalogy, Museo Nacional de Historia Natural, Plaza de la Cultura, and Department of Zoology, Investigation and Conservation, Parque Zoológico Nacional, Santo Domingo, Dominican Republic; Member, IUCN/SSC Crocodile, Insectivore and Elephant Shrew Specialist Groups; Museum Specialist, Carnegie Museum of Natural History; Field Associate, Florida State Museum. Licenciado (mammalogy), University of Santo Domingo. Professional associate of E. E. Williams. Research interests include zoogeography, conservation and insular biology of West Indies vertebrates.

STEPHEN W. PACALA

Assistant Professor, Ecology Section, Biological Sciences Group, University of Connecticut, Storrs, Connecticut 06268. Ph.D. (ecology), Stanford University under J. Roughgarden. Association with E. E. Williams indirect through J. Roughgarden. Research interests include development and testing of population dynamics and niche theories.

FRED PARKER

717 Ross River Rd., Kirwan, Townsville, Queensland 4814, Australia; former District Officer, Papua New Guinea; former Assistant Secretary for Wildlife, Department of Natural Resources, Papua New Guinea; Member, IUCN/SSC Freshwater Chelonian Specialist Group and Snake Specialist Group. Association with E. E. Williams through many years of collecting New Guinean and Australian herpetofauna for the Museum of Comparative Zoology. Research interests include herpetology of Papua New Guinea and the Solomon Islands.

THOMAS S. PARSONS

Professor of Zoology, University of Toronto, Toronto, Ontario M5S 1M1, Canada; former Instructor in Biology, Harvard University. M.A., Ph.D. (comparative anatomy), Harvard University under A. S. Romer. E. E. Williams was on Ph.D. thesis committee, and an early collaborator on turtle morphology. Research

interests include comparative anatomy and paleontology of vertebrates, especially reptiles.

JANE A. PETERSON

Assistant Professor of Biology, University of California, Los Angeles, California 90024. Ph.D. (anatomy), University of Chicago under D. B. Wake. Post-doctoral fellowship at Harvard University under E. E. Williams and A. W. Crompton. Research interests include adaptation in locomotor behavior and morphology, particularly of lower tetrapods, biomechanics, evolution of complex functional units, scaling in the musculoskeletal system.

MARK J. PLOTKIN

Associate in Ethnobotanical Conservation, Harvard Botanical Museum, Oxford St., Cambridge, Massachusetts 02138, and Ethnobotany Project Director, World Wildlife Fund-U.S., 1601 Connecticut Ave., N.W., Washington, D.C. 20009. M.F.S. (forestry), Yale University. Association with E. E. Williams as former curatorial assistant in herpetology at the MCZ. Research interests include neotropical conservation, ethnobotany and herpetology.

A. STANLEY RAND

Senior Scientist, Smithsonian Tropical Research Institute, Balboa, Panama. Ph.D. (zoology), Harvard University under E. E. Williams. Research interests include behavior and ecology of tropical amphibians and reptiles.

ANDERS G. J. RHODIN

Instructor and Chief Resident, Section of Orthopaedic Surgery, Yale University School of Medicine, New Haven, Connecticut 06510 (present address: Orthopaedic Associates, P.C., Nichols Rd., Fitchburg, Massachusetts 01420), and Associate in Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138; Member, IUCN/SSC Freshwater Chelonian Specialist Group. M.D., University of Michigan. Association with E. E. Williams also as former curatorial assistant in herpetology at the MCZ. Research interests include systematics, phylogeny, biology and zoogeography of turtles, especially neotropical and Australo-New Guinean chelids, comparative chondroosseous development of vertebrates, particularly marine turtles.

MARC G. M. VAN ROOSMALEN

Ecologist and Primatologist, Research Institute for Nature Management, Broekhuizerlaan 2, P.O. Box 46, 3956ZR Leersum, Netherlands; Member, IUCN/SSC Primate Specialist Group. Ph.D. (ecology), Agricultural University of Wageningen, Netherlands. Association with E. E. Williams indirect through R. A. Mittermeier. Research interests include ecology and conservation of neotropical primates and tropical ecosystems.

FRANKLIN D. ROSS

Curatorial Assistant, Department of Reptiles

and Amphibians, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138. Research interests include ecology and systematics of reptiles and amphibians, in particular the paleontology and zoogeography of crocodilians.

JAMES PERRAN ROSS

Associate in Herpetology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138; Member, IUCN/SSC Marine Turtle Specialist Group. Ph.D. (zoology), University of Florida. Research interests include physiological ecology and distribution of sea turtles, and application of ecology to conservation of endangered species.

JONATHAN ROUGHGARDEN

Professor of Biological Sciences, Stanford University, Stanford, California 94305. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include theoretical and community ecology.

JOHN D. RUMMEL

Graduate Student, Department of Biological Sciences, Stanford University, Stanford, California 94305. Ph.D. candidate (ecology), Stanford University under J. Roughgarden. Association with E. E. Williams indirect through J. Roughgarden. Research interests include experimental ecology, niche theory and the effect of population processes on ecosystem characteristics.

AMY SCHOENER

Research Assistant Professor, Department of Oceanography, University of Washington, Seattle, Washington 98195. Ph.D. (biology), Harvard University. Participated with E. E. Williams and T. W. Schoener on lizard studies in the West Indies. Research interests include island biogeography and marine ecology.

THOMAS W. SCHOENER

Professor of Zoology and Environmental Studies, Zoologist in the Agricultural Experiment Station, University of California, Davis, California 95616; former Professor, Department of Zoology, University of Washington, Seattle, Washington; former Associate Professor, Department of Biology, Harvard University. Ph.D. (biology), Harvard University under E. E. Williams and E. O. Wilson. Junior Fellow, Harvard. Also associated with E. E. Williams as undergraduate student at Harvard. Research interests include ecology in general.

ALBERT SCHWARTZ

Professor, Department of Biology, Miami-Dade Community College, North Campus, Miami, Florida 33167; Research Associate, Department of Amphibians and Reptiles, Carnegie Museum of Natural History, Pittsburgh. M.S. (parasitology), University of Florida; Ph.D. (mammalogy), University of Michigan. Professional associate of E. E. Williams for many years. Research interests include taxonomy and

zoogeography of amphibians, reptiles, birds and mammals of the West Indies.

JON SEGER

Post-doctoral Fellow, Museum of Zoology, University of Michigan, Ann Arbor, Michigan 48109. Ph.D. (biology), Harvard University under E. E. Williams. Research interests include kin selection theory and evolution of genetic systems and sex ratios, especially in solitary Hymenoptera.

MICHAEL E. SOULÉ

Director, Institute for Transcultural Studies, 901 S. Normandie Ave., Los Angeles, California 90006; former Professor of Biology, University of California, San Diego. M.A., Ph.D. (population biology), Stanford University. Professional association with E. E. Williams mainly through G. C. Gorman. Research interests include variation in natural populations and conservation biology.

SAMUEL F. TARSITANO

Adjunct Assistant Professor, Department of Biology, Queens College, City University of New York, Flushing, New York 11367. Ph.D. (comparative and functional anatomy), City University of New York. Association with E. E. Williams indirect through M. K. Hecht. Research interests include functional morphology and vertebrate paleontology.

RICHARD THOMAS

Associate Professor, Department of Biology, University of Puerto Rico, Río Piedras, Puerto Rico 00931. Ph.D. (zoology), Louisiana State University. Professional associate of E. E. Williams for many years. Research interests include systematics of West Indian amphibians and reptiles.

ROBERT L. TRIVERS

Professor of Biology, University of California, Santa Cruz, California 95064; former Assistant and Associate Professor, Department of Biology, Harvard University. Ph.D. (biology), Harvard University under E. E. Williams. Regents post-doctoral fellowship, Smithsonian Institution. Research interests include social evolution in general.

LINDA TRUEB

Associate Curator, Division of Herpetology,

Museum of Natural History, and Associate Professor, Department of Systematics and Ecology, University of Kansas, Lawrence, Kansas 66045. Ph.D. (zoology, herpetology), University of Kansas. Professional associate of E. E. Williams. Research interests include evolutionary biology of amphibians with emphasis on osteology and systematics of anurans.

PAULO EMILIO VANZOLINI

Director, Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil. Ph.D. (zoology), Harvard University under A. S. Romer. Professional associate and collaborator with E. E. Williams for more than thirty years. Research interests include herpetology and evolution.

DAVID B. WAKE

Professor of Zoology and Director, Museum of Vertebrate Zoology, University of California, Berkeley, California 94720; President, Society for the Study of Evolution. Ph.D. (biology), University of Southern California. Professional associate of E. E. Williams for many years. Research interests include origin of evolutionary novelty and adaptive radiation, evolutionary biology of plethodontid salamanders, size and shape in ontogeny and phylogeny, and functional, developmental, and comparative morphology.

ROGER CONANT WOOD

Professor of Zoology, Stockton State College, Pomona, New Jersey 08240; Member IUCN/SSC Freshwater Chelonian Specialist Group. Ph.D. (vertebrate paleontology), Harvard University under B. Patterson and E. E. Williams. Post-doctoral fellowships at Harvard and Florida Audubon Society. Research interests include evolution, morphology, ecology and behavior of fossil and living turtles.

SUH YUNG YANG

Professor and Chairman, Department of Biology, Inha University, Inchon, Korea. Ph.D. (zoology), University of Texas. Association with E. E. Williams indirect through G. C. Gorman. Research interests include evolutionary biology in vertebrates.

TABLE OF CONTENTS

SYSTEMATICS

- ELIAS, PAUL, AND DAVID B. WAKE. *Nyctanolis pernix*, a new genus and species of plethodontid salamander from northwestern Guatemala and Chiapas, Mexico. 1
- INGER, ROBERT F. Larvae of southeast Asian species of *Leptobrachium* and *Leptobrachella* (Anura: Pelobatidae). 13
- DUELLMAN, WILLIAM E., AND LINDA TRUEB. Frogs of the *Hyla columbiana* group: taxonomy and phylogenetic relationships. 33
- LYNCH, JOHN D. A new leptodactylid frog from the Cordillera Oriental of Colombia. 52
- RHODIN, ANDERS G. J., AND RUSSELL A. MITTERMEIER. Description of *Phrynops williamsi*, a new species of chelid turtle of the South American *P. geoffroanus* complex. 58
- WOOD, ROGER CONANT. *Kenyemys williamsi*, a fossil pelomedusid turtle from the Pliocene of Kenya. 74
- THOMAS, RICHARD, AND ALBERT SCHWARTZ. Variation in Hispaniolian *Sphaerodactylus* (Sauria: Gekkonidae). 86
- LAZELL, JAMES D., JR. Biogeography of the herpetofauna of the British Virgin Islands, with description of a new anole (Sauria: Iguanidae). 99
- VANZOLINI, P. E. Guiano-Brasilian *Polychrus*: distribution and speciation (Sauria: Iguanidae). 118
- DIXON, JAMES R. Systematics of the Latin American snake, *Liophis epinephelus* (Serpentes: Colubridae). 132

COMPARATIVE MORPHOLOGY

- BELLAIRS, A. D'A. Partial cyclopia and monorhinia in turtles. 150
- BROADLEY, DONALD G. Neural pattern—a neglected taxonomic character in the genus *Pelusios* Wagler (Pleurodira: Pelomedusidae). 159
- MCDOWELL, SAMUEL B. The genus *Emydura* (Testudines: Chelidae) in New Guinea with notes on the penial morphology of Pleurodira. . 169
- GAFFNEY, EUGENE S. The basicranial articulation of the Triassic turtle, *Proganochelys*. 190
- MOODY, SCOTT M. The rectus abdominis muscle complex of the Lacertilia: terminology, homology and assumed presence in primitive iguanian lizards. 195

GREER, ALLEN E. On the adaptive significance of the reptilian spectacle: the evidence from scincid, teiid, and lacertid lizards.	213
FRAZZETTA, T. H. Adaptation and function of cranial kinesis in reptiles: a time-motion analysis of feeding in alligator lizards.	222
PETERSON, JANE A. The evolution of the subdigital pad in <i>Anolis</i> . I. Comparisons among the anoline genera.	245
FORSGAARD, KARL. The axial skeleton of <i>Chamaelinorops</i>	284
PARSONS, THOMAS S., AND LUDMELA DJATSCHENKO. Variation in the left lung and bronchus of <i>Thamnophis sirtalis parietalis</i>	296
ROSS, FRANKLIN D., AND GREGORY C. MAYER. On the dorsal armor of the Crocodilia	305
HECHT, MAX K., AND SAMUEL F. TARSITANO. The tarsus and metatarsus of <i>Protosuchus</i> and its phyletic implications.	332

ZOOGEOGRAPHY

LAURENT, R. F. About the herpetofauna of central African montane forest.	350
KIESTER, A. ROSS. Zoogeography of the skinks (Sauria: Scincidae) of Arno Atoll, Marshall Islands.	359
ESTES, RICHARD. The fossil record and early distribution of lizards. ..	365
CONANT, ROGER. Commentary on a frog and lizard newly recorded from central Durango, Mexico.	399
ESPINOZA, NELLY CARRILLO DE. List of Peruvian <i>Anolis</i> with distributional data (Sauria: Iguanidae).	406
KOOPMAN, KARL F. Two general problems involved in systematics and zoogeography of bats.	412

ECOLOGY

BROWN, WALTER C., AND ANGEL C. ALCALA. Modes of reproduction of Philippine anurans.	416
MEDEM, FEDERICO. Reproductive data on <i>Platemys platycephala</i> (Testudines: Chelidae) in Colombia.	429
PARKER, FRED. The prehensile-tailed skink (<i>Corucia zebrata</i>) on Bougainville Island, Papua New Guinea.	435
ANDREWS, ROBIN M., A. STANLEY RAND, AND STELLA GUERRERO. Seasonal and spatial variation in the annual cycle of a tropical lizard.	441
RAND, A. S., STELLA GUERRERO, AND ROBIN M. ANDREWS. The ecological effects of malaria on populations of the lizard <i>Anolis limifrons</i> on Barro Colorado Island, Panama.	455
HERTZ, PAUL E. Eurythermy and niche breadth in West Indian <i>Anolis</i> lizards: a reappraisal.	472
HUEY, RAYMOND B. Natural variation in body temperature and physiological performance in a lizard (<i>Anolis cristatellus</i>).	484

SCHOENER, THOMAS W., AND AMY SCHOENER. On the voluntary departure of lizards from very small islands.	491
ROUGHGARDEN, JONATHAN, JOHN RUMMEL, AND STEPHEN PACALA. Experimental evidence of strong present-day competition between the <i>Anolis</i> populations of the Anguilla Bank—a preliminary report. ...	499
MOERMOND, TIMOTHY C. Competition between <i>Anolis</i> and birds: a reassessment.	507
MITTERMEIER, RUSSELL A., AND MARC G. M. VAN ROOSMALEN. A synecological study of Surinam monkeys.	521

BEHAVIOR

AUFFENBERG, WALTER. Courtship behavior in <i>Varanus bengalensis</i> (Sauria: Varanidae).	535
JENSSEN, THOMAS A. Display behavior of two Haitian lizards, <i>Anolis cybotes</i> and <i>Anolis distichus</i>	552
HICKS, ROBERT A., AND ROBERT L. TRIVERS. The social behavior of <i>Anolis valencienni</i>	570

EVOLUTION

SEGER, JON. Conditional relatedness, recombination, and the chromosome number of insects.	596
GANS, CARL. Is <i>Sphenodon punctatus</i> a maladapted relict?	613
BLAKE, J. A. Chromosomal C-banding in <i>Anolis grahami</i>	621
GORMAN, GEORGE C., DONALD G. BUTH, MICHAEL SOULÉ, AND S. Y. YANG. The relationships of the Puerto Rican <i>Anolis</i> : electrophoretic and karyotypic studies.	626
HALL, WILLIAM P. Modes of speciation and evolution in the sceloporine iguanid lizards. I. Epistemology of the comparative approach and introduction to the problem.	643
HOLT, ROBERT D. Immigration and the dynamics of peripheral populations.	680

CONSERVATION

PLOTKIN, MARK J., FEDERICO MEDEM, RUSSELL A. MITTERMEIER, AND ISABEL D. CONSTABLE. Distribution and conservation of the black caiman (<i>Melanosuchus niger</i>).	695
ROSS, JAMES PERRAN, AND JOSE ALBERTO OTTENWALDER. The leatherback sea turtle, <i>Dermochelys coriacea</i> , nesting in the Dominican Republic.	706

Nyctanolis pernix, A New Genus and Species of Plethodontid Salamander from Northwestern Guatemala and Chiapas, Mexico

PAUL ELIAS¹

DAVID B. WAKE²

ABSTRACT. A new bolitoglossine salamander, *Nyctanolis pernix*, from the Cordillera de los Cuchumatanes of Guatemala and neighboring Mexico has been discovered. It differs from all other neotropical plethodontids in its spotted color pattern, long legs, and divided premaxilla. The osteology of *Nyctanolis* is the most plesiomorphous of any member of the supergenus *Bolitoglossa*. Except for the apomorphies characterizing the supergenus, *Nyctanolis* retains a morphology ancestral to the entire tribe Bolitoglossini and one more generally primitive than found in the other supergenera of the tribe, *Batrachoseps* and *Hydromantes*. The morphological features of the new genus place it as a sister group to the rest of the supergenus *Bolitoglossa*. The apparently ancient lineage represented by the new form occurs today in the oldest land-positive area in all of Central America.

INTRODUCTION

The remarkable radiation of the Bolitoglossini that has occurred in Mexico includes 7 genera and 63 species as at present understood. . . . Beyond question other genera and dozens of other species remain to be discovered. . . .

—Smith and Smith, 1976

Our understanding of the evolutionary history of tropical salamanders is still in

its infancy, and undescribed species are found with regularity. No new genera have been described for thirty years (Tanner, 1950), but despite this seeming stability several of the presumed lineages are poorly defined (Wake and Lynch, 1976). In the summer of 1974 the senior author visited a remote area on the eastern slopes of the Sierra de los Cuchumatanes, in extreme northwestern Guatemala. There, in a limited area, he discovered five species of salamanders living in sympatry in a cloud forest at intermediate elevations. Three of these species proved to be undescribed. In this paper we present a description of the most distinctive of the new species. It is so unusual in its combination of ancestral and derived morphology that it requires its own genus. It is the only tropical plethodontid that has two premaxillary bones, and it differs from all other tropical salamanders as well in its body form and behavior. This long-legged, large, colorful salamander, with its long, whip-like tail, and high level of scansorial activity reminds us more of an anoline lizard than a typical salamander.

We name the new genus for its anoline aspect and nocturnal habits (Gr., *nyktos*, night) and the species (L., quick, agile) for its gymnastic behavior.

^{1,2}Museum of Vertebrate Zoology, University of California, Berkeley, California 94720, U.S.A.

TAXONOMY AND MORPHOLOGY

Nyctanolis new genus

Type species. Nyctanolis pernix sp. nov.

Diagnosis. A plethodontid salamander belonging to the subfamily Plethodontinae, tribe Bolitoglossini, supergenus *Bolitoglossa*; distinguished from all other members of the supergenus by the presence of paired premaxillary bones and extremely long tail, limbs, and digits. The genus is easily distinguished from the diminutive species of *Parvimolge* and *Thorius* by its large size, and from the elongate species of *Lineatriton* and *Oedipina* by its long limbs and large, broad head. Most species of both *Chirop-terotriton* alpha and beta are much smaller, but it is further distinguished from the former by having a fifth distal tarsal that is smaller than the fourth, and from the latter in having a distinct lingual cartilage and a well-developed columella. It differs from *Bolitoglossa* in having a distinct sublingual fold, nine tarsal elements, a well-developed columella, and a lingual cartilage. It differs further from *Bolitoglossa* alpha in having a well-defined tibial spur, and from *Bolitoglossa* beta in having only a slightly constricted tail base and unspecialized transverse processes on the first caudal vertebra. *Nyctanolis* most closely resembles species of *Pseudoeurycea*, but is distinguished from that genus by being more extreme in limb, hand, and foot specializations and by having a more fully developed columellar stylus and lingual cartilage, in addition to its unique premaxillae.

Nyctanolis pernix sp. nov.

Figures 1-4

Holotype. Museum of Vertebrate Zoology (MVZ) 134641, an adult female from Finca Chiblac, 10 km (air) NE Barillas, Huehuetenango, Guatemala,

(91°16'W, 15°53'N), 1,370 m (4,500 ft) elevation, collected 29 August 1975, by P. Elias and J. Jackson.

Paratypes. MVZ 131583-85, 134639-40, 134642-44, 149370, 149372-73, 173062 (cleared and stained), MCZ 100154, same data as holotype but collected at different times by P. Elias, J. Jackson, H. B. Shaffer, and A. Diaz. United States National Museum 206925, cave near stream that empties the main lake, Lagunas de Montebello, Chiapas, Mexico, (91°32'W, 16°5'N), collected by Scott Belfit, 8 July 1972.

Diagnosis. A large, slender species (standard length, SL, in four adult males 43.2-68.1, $\bar{x}$ 54.9; nine adult females 57.8-73.6, $\bar{x}$ 67.9) with a very long tail (SL/tail length in three adult males is 0.73-0.84, $\bar{x}$ 0.79; three adult females 0.80-0.94, $\bar{x}$ 0.87) and long limbs and digits (when appressed fore and hind limbs overlap in four adult males by 3-4 costal grooves, $\bar{x}$ 3.6; in eight adult females by 1.5-4.0 costal grooves, $\bar{x}$ 2.5). The head is broad (SL/head width in four adult males 5.8-6.3, $\bar{x}$ 6.0, in nine adult females 6.0-6.6, $\bar{x}$ 6.2) and relatively flat, and contains large numbers of maxillary-premaxillary (88-119, $\bar{x}$ 103 in four adult males; 82-131, $\bar{x}$ 110 in nine adult females) and vomerine teeth (29-47, $\bar{x}$ 41.5 in four adult males; 38-53, $\bar{x}$ 45.7 in nine adult females). The species has highly distinctive coloration, and typically animals are shiny dark black with bright red spots on the eyelids that become red-orange, then orange, then yellow, and finally cream-colored posteriorly on the limbs, body, and tail.

Description. This large and long-legged but slender species has a short snout. Both males and females grow to a large size. The nostrils and labial protuberances are small in both sexes but are largest in adult males, whose snouts are generally expanded forward to considerably overhang the mandible. The mental hedonic gland is large and pronounced in adult males, attaining a width half that of the entire head. The head is wide and flattened, always markedly broader in dorsal aspect than the broadest point on the trunk. The ratio of maximum head width to head depth at the angle of the

jaw varies in adults from 2.1 to 2.3. A deep unpigmented groove extends below the eye, following its curvature, to near the posterior border of the eye opening, but does not communicate with the lip. A marked angle in the line of the mouth beneath the eye cocks the posteriormost section of the lip gently ventrad to the angle of the jaw. The large and prominent eyes bulge upward from the flattened head, especially in males, but do not extend lateral to the jaw margins. An indistinct postorbital groove extends posteriorly and gently ventrad from behind the eye as a shallow depression turning sharply directly ventrad as a deep crease which, passing behind the jaw articulation, disappears below the level of the mandible. No trace of a nuchal groove is evident, but the gular fold is strongly developed and arches slightly forward at the midline. Vomerine teeth increase in number with increasing size (Table 1) but there are many even in small individuals. These teeth are arranged in a long gently arched row which extends laterally to far beyond the small internal nares, nearly to the jaw line. The many maxillary teeth extend in a long row to the interior angle of the jaw and slightly beyond the posterior margin of the eye. The maxillary teeth are smoothly continuous with the premaxillary teeth so that there is no break except for a slight discontinuity in the largest male. The premaxillary teeth are numerous (up to 15) but are only slightly sexually differentiated and do not pierce the upper lip in males. The tongue pad is nearly round and lies at the end of a pedicel. There is no anterior attachment, and the tongue has the boletoid form characteristic of the supergenus. A large, fleshy sublingual fold is present. The trunk and tail are slender and cylindrical in this species and no strong basal constriction is evident. The tail may be autotomized at any point along its length. The tail is relatively long. No postiliac gland is visible. The limbs are extremely long, and when fore and hind limbs are stretched along

the flank they overlap extensively. Hands and feet are large and unwebbed, with distally expanded, quadrangular digital tips. The fingers are, in order of decreasing length, 3,4,2,1; the toes, 3,4,2,5,1.

Measurements of the Holotype (in mm). Head width 11.6; snout to gular fold (head length) 15.2; head depth at posterior angle of jaw 5.4; eyelid length 5.0; eyelid width 3.7; anterior rim of orbit to snout 4.4; horizontal orbit diameter 3.5; interorbital distance 3.5; distance between vomerine teeth and parasphenoid tooth patch 0.5; snout to forelimb 21.1; distance separating internal nares 3.3; distance separating external nares 3.4; snout projection beyond mandible 0.3; snout to posterior angle of vent (standard length, SL) 70.5; snout to anterior angle of vent 64.4; axilla to groin 36.9; tail length 79.7; tail width at base 4.8; tail depth at base 5.6; forelimb length (to tip of longest toe) 21.6; hindlimb length 23.7; width of hand (across from tip of innermost to tip of outermost toe when spread) 7.6; width of foot 9.7.

Coloration (in alcohol). All ventral surfaces are slate black except palmar and plantar surfaces and tips of the nasal cirri which are less saturated with pigment and thus appear lighter grey. One large adult male (MVZ 134642) is uniformly dark black dorsally as well as ventrally, with the exception of a small, indistinct spot middorsally. All other specimens have essentially the same, almost gaudy coloration. Bright, discrete spots extend from eyelids to tail tip in these 13 animals. One of these has been cleared and stained; variation in 12 other specimens is as follows. A series of large, round, pale yellow spots with irregular margins, each about equal in diameter to the eye, is present on the black dorsum. These spots are almost symmetrically arranged. One spot covers each eyelid in all animals. One spot covers the dorsal surface of each elbow in all but one animal, which has a spot on one side only. All animals have a large spot covering each knee. A pair of spots lies over the shoulders in 11

Figure 1. Dorsal view of the holotype of *Nyctanolis pernix*, MVZ 134641.

TABLE 1. VARIATION IN *NYCTANOLIS PERNIX*.

	Sex	Standard Length	Head Length	Head Width	Foot Width	Hind Limb Length	Fore Limb Length	Tail Length	Maxillary and Premaxillary Tooth Number	Vomerine Tooth Number	Limb Interval
MVZ 134642	M	68.1	16.8	11.8	9.3	23.1	20.6	86.4	119	45	-4.0
MVZ 134644	M	58.0	13.4	9.5	7.1	18.8	18.0	69.2	105	45	-3.0
MVZ 131585	M	50.1	11.3	8.6	6.4	16.2	16.0	68.3	88	29	-4.0
MVZ 149370	M	43.2	9.8	6.9	5.1	14.1	14.2	-	100	47	-3.5
MVZ 134640	F	73.6	15.9	12.3	9.9	25.2	23.5	-	117	43	-1.5
MVZ 134639	F	71.7	16.1	11.5	7.9	22.7	20.7	89.9	110	49	-2.0
MVZ 134641*	F	70.5	15.2	11.6	9.7	23.7	21.6	79.7	112	42	-3.0
MVZ 134643	F	69.9	15.3	11.4	9.8	21.2	19.9	-	131	53	-2.5
MVZ 149373	F	68.2	15.1	10.4	8.8	21.6	21.6	-	82	51	-4.0
MVZ 131583	F	67.6	15.4	10.9	8.3	20.8	19.6	72.1	115	47	-2.0
USNM 206925	F	67.3	15.5	10.9	9.1	22.0	21.4	-	107	40	-
MVZ 173062	F	64.1	13.5	10.2	8.4	20.3	19.9	-	104	38	-2.5
MVZ 149372	F	57.8	12.3	9.1	6.9	19.5	18.6	-	109	48	-2.5
MVZ 131584	J	27.5	7.3	5.7	3.2	9.1	7.5	-	50	13	-0.5

*Holotype.

animals, and the 12th has a single shoulder spot. Many individuals have one spot over the pelvis (five have one, one has two, and six are unspotted). Most individuals (nine) have some sort of spot, variable in size and position, in the caudosacral region. Full-sized (i.e., the size of the eyelids) dorsal spots on the trunk between shoulder and pelvis occur in all individuals, and range in number from two to eight. Included among the trunk spots of four specimens is a pair placed symmetrically behind the shoulder. Three animals show variable numbers of small, obscure spots on the trunk. Most individuals lack complete tails, but large spots occur only on the proximal one third of the tail and the first third is complete in 11 specimens. Of these 11, all have some distinct tail spots, ranging from one to eight in number. In addition four of the seven with over one-half the tail present show clusters of up to nine small indistinct spots on the second half of the tail. Miscellaneous small spots occur in many animals: seven have spots on the temple; two have parietal spots; two have paired wrist spots; one has a foreleg spot; one has a flank spot; one has a mandibular spot.

Coloration (in life). These animals were patterned as they are in preservation, but the coloration of the dorsal spotting differed. The spots on the eyelids and neck are a deep crimson, those on the elbows and knees are orange, and the trunk and tail spots are light yellow, becoming cream at the tail tip.

Habitat. The series from the type locality was collected in a large clearing surrounded by virgin cloud forest. The locality is extremely humid as indicated by rainfall records taken within 5 km of the spot. These records show rainfall of 5 to 6 m annually, but they were taken below the cloudline and thus may underestimate precipitation at the type locality.

The animals at the type locality were collected under both moss and bark on felled cloud forest hardwoods. They were active on rainy nights. Rarely were

more than two individuals taken per man-day collecting.

Also occurring at the site were four other salamanders: *Bolitoglossa hartwegi*, *Bolitoglossa cuchumatana*, a new species of *Bolitoglossa* (Elias, in preparation), and another undescribed species belonging to an as yet undetermined genus of plethodontid salamanders. The three *Bolitoglossa* were taken in the same microhabitats as those in which *Nyctanolis* was found, but the disturbed nature of the habitat leaves ecological relationships vague.

Various hyloid and leptodactylid frogs were also taken at the site as was one snake (*Leptodeira*) and lizards of the genera *Anolis*, *Sceloporus*, *Scincella*, *Barisia*, and *Lepidophyma*.

Range. The species is known from the type locality and one other locality about 20 km NW of Finca Chiblac just across the border in Chiapas, Mexico.

Osteology. Information concerning osteology has been derived from one cleared and stained specimen (MVZ 173062), an adult female, and from radiographs of all of the specimens. Dorsal and ventral views of the skull are illustrated (Fig. 2).

The relatively broad and low skull is compact and well ossified, with well-organized articulations and well-defined bones. Premaxillary bones are distinctly separate in all individuals in which they can be seen; the smallest individual is insufficiently ossified for the bones to be resolved in radiographs. The premaxillary bones are large, relative to the small, fused bones characteristic of other members of the supergenus *Bolitoglossa*. They are well articulated with each other, and are tightly articulated laterally to the maxillaries, which dorsally overlap the dental portions of the premaxillaries. The premaxillaries of both males and females are part of a continuous maxillary-premaxillary arcade, rather than being discontinuously projected anteriorly as in many other tropical salamanders. The palatal process of each premaxillary bone

Figure 2. Dorsal (left) and ventral (right) views of the skull of *Nyctanolis pernix*, drawn from cleared and stained specimen with aid of microprojector. Cartilage stippled.

Abbreviations: Fr, frontal; Ma, maxillary; Op, operculum; Os, orbitosphenoid; Ot, otic-occipital; Pa, parietal; Pf, prefrontal; Pm, premaxillary; Ps, parasphenoid; Pt, pterygoid portion of palatopterygoid cartilage; Pv, posterior vomerine tooth patch (outline only, teeth not drawn); Qa, quadrate; Sq, squamosal; Vo, vomer.

is narrow and well separated from the vomer behind it. Frontal processes of the premaxillaries are stout and well developed; they arise from a narrow base as columnar structures, then immediately diverge around the large, internasal fontanelle. As the processes rise dorsally and posteriorly they flatten and form the borders of the fontanelle. As they flatten they become somewhat expanded. Laterally the processes closely abut the large nasals, and posteriorly they broadly overlap the anterior extensions of the facial portions of the frontals. The frontal processes do not contact each other behind the fontanelle, but in some individuals they come very close. The frontal processes terminate rather far forward, well in advance of the orbit, and of the posterior tip of the nasals. Nasals are large, but they are nonprotuberant in all females and in all but the largest male; in that

individual the premaxillary bones are larger than in any other specimen and the nasal bones extend anteriorly slightly beyond the premaxillaries. The large nasals are triangular in shape, and they have strong, overlapping articulations with the facial processes of the frontals. The nasals slightly overlap the prefrontals, and are overlapped by the anterior part of the facial processes of the maxillaries. The nasals are not domed, but are rather flattened or slightly concave. No septomaxillaries can be seen in any specimen (a slight shadow on one side of the skull in radiographs of one of the smaller specimens might conceivably be a septomaxillary). Prefrontal bones are discrete but relatively small. The bones are only about one-quarter or less the area of the nasals. The prefrontals broadly overlap the facial parts of the frontals, and are overlapped rather narrowly by the pos-

terior and dorsalmost parts of the facial processes of the maxillaries. The foramen for the nasolacrimal duct lies at the extreme anterolateral tip of the prefrontals and is indented into the dorsal margin of the facial processes of the maxillaries. There is a slight indentation of the posterolateral border of the nasals as well. The nasolacrimal duct proceeds from the orbit across the lateral parts of the prefrontals, which are depressed and almost grooved. The maxillary bones are very large and well developed. The bones extend posteriorly to terminate in sharp points at the level of the posterior margin of the eye. The maxillaries bear small but well-developed, bicuspid teeth from the premaxillary nearly to the posterior tip; these teeth are about the same size as the premaxillary teeth in females, but are somewhat smaller in males. The facial processes of the maxillaries are relatively very large and well developed; they are larger in area than the nasals. The palatal processes are moderately well developed and extend as shelflike processes to the margins of the vomer bodies, against which they tightly abut.

The vomers are large bones which are well separated from each other by the intervomerine fontanelle anteriorly. This fontanelle contains a large glandular mass, present also in other plethodontid salamanders but frequently not so well developed. The vomers abut dorsally and posteriorly to this mass, and the posterior parts of the two vomers are tightly articulated. The preorbital processes of the vomers are long and stout; they extend beyond the lateral margins of the vomer bodies, and bear teeth in a long, curving series nearly to their tips, well beyond the lateral border of the internal nares.

Frontals and parietals are well developed and firmly articulated with each other in an extensive interlocking contact. The paired elements are tightly articulated to each other along the midline as well. Anterior parts of the frontals are moderately large and contribute importantly to the facial part of the skull. The

overlapping posterior lobes of the frontals are relatively very large. The frontals appear to be relatively narrow, but we believe that this is simply an illusion that results from the unusual (for a tropical salamander) full development of the outer bones of the skull. Parietals are well developed, but they do not have the distinct lateral spurs that are characteristically present (but sometimes very poorly developed) in tropical salamanders (Wake, 1966). The region of the lateral spur is somewhat expanded and ventrally directed, but is not spurlike. The otic capsules are large and well developed, but they do not appear to be relatively as large as in other tropical genera, in which the other portions of the skull are frequently reduced in size. Two or three spinelike projections arise from the anterodorsal parts of the capsules, and are directed laterally; these form a kind of rudimentary crest. The large parasphenoid is narrow anteriorly, but the orbitosphenoids are well separated from each other. The tip of the parasphenoid is blunt. Posterior vomerine teeth are borne in long, relatively narrow patches that are well separated from each other and from the anterior vomerine teeth. The teeth in these patches are somewhat smaller than the maxillary teeth. In the cleared and stained specimen there are 81 (left) and 92 (right) bicuspid, ankylosed teeth. The operculum has a small but well-developed stilus that is as large as this structure ever becomes in any of the other tropical genera (it is absent in the vast majority of tropical species). Well-developed, stout quadrates are connected to the otic capsules by the very large, broad squamosals, and by the cartilaginous suspensorium.

The lower jaw has a long, slender dentary bearing a very long series of small teeth, about the size of those on the maxillary. The prearticular is relatively small and low.

The hyobranchial apparatus (Fig. 3) is very generalized for the supergenus *Boli-*

toglossa (cf. Tanner, 1952; Lombard and Wake, 1977). It is typical of the group; there is no urohyal, and the radii are essentially continuous with the basibranchial. There is a well-developed lingual cartilage, but it is represented by what is essentially a direct anterior continuation of the basibranchial, although there is a change from hyaline to fibrocartilage in the joint region between the cartilage and the basibranchial. We have also had available one sectioned head of the species, and detailed examination of it reveals that in all respects the hyobranchial apparatus and associated musculature fits the morphological features of Mode VI of Lombard and Wake (1977). Specifically, the entire apparatus is greatly elongated, and the proportions of the elements are unequal. The epibranchials are very long, more than three times the length of the basibranchial (minus lingual cartilage), and about six times the length of the second ceratobranchials. The diameters of the basibranchial, second ceratobranchial, and base of the epibranchial are about equal, and all are greater than that of the relatively slender first ceratobranchial. The first ceratobranchial is not so slender as is characteristic of that element in many members of the supergenus *Bolitoglossa*. The radii are of moderate length for a member of the supergenus, but are relatively stout. The ceratobranchials are relatively long, but generalized in form; there is a distinct, but narrow, flattened blade distally, and there is no anterior filament. The point of attachment of the suprapeduncularis muscle is drawn medially into a broad-based process.

Vertebral structure is generalized. Individual vertebrae have centra with hollow, tapered, bony husks, and are joined to one another by the spindlelike intervertebral cartilages and the zygapophyses. There is a single atlas, 14 trunk vertebrae, one sacral vertebra, two caudosacral vertebrae, and a variable number of caudal vertebrae. The trunk vertebrae all have transverse processes that

Figure 3. Dorsal view of the cartilaginous hyobranchial apparatus of *Nyctanolis pernix*.

Abbreviations: Bb, basibranchial; CbI, first ceratobranchial; CbII, second ceratobranchial; Ch, ceratohyal; Eb, epibranchial.

arise and remain separated until their tips. These processes are relatively long, and extend well beyond the limits of the zygapophyses. The upper transverse process is a little anterior in placement relative to the lower one. Relatively long, slender ribs are borne on all but the last trunk vertebra; rib heads are relatively widely separated. Spinal nerve routes are similar to those reported by Edwards (1976) for this group: a ventral root issues from the neural arch of the atlas; dorsal and ventral roots issue from separate foramina in the anterior, pre-transverse process part of the neural arch of the first trunk vertebra; there is a very large foramen in the anterior, pre-transverse process part of the neural arch of the second trunk vertebra, and a set of dorsal and

ventral roots emerge through a common, small foramen in the posterior, post-transverse process part of the neural arch of the same vertebra; dorsal and ventral roots emerge through a common, small foramen in the posterior, post-transverse process part of the third and all succeeding trunk vertebrae.

The first caudosacral vertebra has relatively long, nearly straight transverse processes. The second is a smaller vertebra, with processes that are much shorter than those of the first; these processes may be straight, but are more frequently oriented anteriorly. The origin of the processes on the first caudosacral vertebra is about midcentral, but those of the second vertebra arise in advance of the midpoint of the centrum. The second caudosacral vertebra bears rudiments of the haemal arch, not joined to each other, on the posterior part of the ventral surface of the centrum. The foramen for the spinal nerves is located behind the processes, about midcentrally.

The tail base region is not nearly as specialized for tail autotomy as in most other members of the supergenus (Wake and Dresner, 1967), but fundamentally it qualifies as a constricted-based tail even if the constriction is only slight. The first caudal vertebra is slightly, but not obviously, shorter than the second. It differs from the second caudal in having a complete haemal arch that lacks an anterior keel. It differs from the second caudosacral immediately in front of it in having transverse processes that arise at the very anterior end of the vertebra, essentially as extensions of the bases of the prezygapophyses. The transverse processes are short, and extend only a short distance anterior and lateral to the edge of the prezygapophyses. The musculature extending between the first caudal and the second caudosacral vertebrae is visibly shortened, relative to adjacent segments, but the tail is slender and the shortened segment is not obviously constricted. The smallest juvenile lost its tail at the base during capture, between the second

caudosacral and the first caudal vertebrae. The skin of the shortened segment has collapsed over the wound in the preserved animal, in the typical wound-healing specialization that characterizes salamanders with constricted and slender-based tails (Wake and Dresner, 1967).

Complete tails are rare. The only large specimen with a complete tail has 39 caudal vertebrae, a very large number for the supergenus, with the exception of the exceptionally elongate species of *Lineatriton* and *Oedipina*. Other large specimens show signs of earlier breaks, always away from the basal region. We cannot rule out tail base breaks in juveniles, with subsequent regeneration. One specimen has a tail broken at vertebra 39. Another has a total of 45 vertebrae, regenerated from vertebra 24. There are 49 vertebrae in the tail of another specimen, regenerated from vertebra 24. The holotype has 34 caudal vertebrae, but its tip is missing; further, it shows signs of an initial break at vertebra 18, and a subsequent break at vertebra 27. The caudal vertebrae are narrow and elongate; transverse processes are very short and are little more than small projections of the prezygapophyses. The spinal foramina are located midcentrally.

The hands and feet are highly distinctive (Fig. 4). The digits are very long, with terminal expansion. All phalangeal elements are well developed. The phalangeal formula is 1, 2, 3, 2 for the hand and 1, 2, 3, 3, 2 for the foot; this is the primitive formula for the supergenus *Bolitoglossa*. The long metapodial elements are cylindrical, as are all but the terminal phalanges. Terminal phalanges are specialized, and resemble those of *Aneides lugubris* (Wake, 1963); they are greatly expanded distally, and the expanded portion is slightly recurved and nearly bifurcated in the larger specimens. The expanded portion of each terminal phalanx is distinctly flattened. On the ventral surface of each phalanx, near the base, a well-defined projection is

present which serves as the site of insertion of a digital tendon. Even the phalanx of the first digit displays some distal expansion.

The carpal and tarsal arrangements are those characteristic of primitive plethodontids. There are eight carpals and nine tarsals. Distal tarsal 5 is smaller than distal tarsal 4 and does not articulate with the centrale.

Limb bones are very elongate, slender, and cylindrical. There is a well-defined, distinct tibial spur near the proximal end of the tibia, and the bone has a sharply defined crest distally.

Relationships. The tribe Bolitoglossini is characterized by seven unique apomorphies relative to all other plethodontids. All members of the tribe have 1) lost the urohyal; 2) radii fused to the basi-branchial; 3) long epibranchials relative to the ceratobranchials; 4) a second ceratobranchial modified for force transmis-

sion; 5) a cylindrical muscle complex around the tongue; 6) lost the circumglossus muscle (all discussed in Lombard and Wake, 1977); and 7) a juvenile otic capsule configuration (Lombard, 1977).

The supergenus *Bolitoglossa* is characterized by having a specialized autotomy area in the tail base that follows two caudosacral vertebrae (Wake and Dresner, 1967), and by some modifications of the throat and tongue musculature (Lombard and Wake, 1977).

The morphology of *Nyctanolis* includes all of the unique apomorphies characteristic of both the tribe Bolitoglossini and the supergenus *Bolitoglossa* and we therefore consider it to be a member of both taxa.

In most of its characters *Nyctanolis* is generalized within the supergenus *Bolitoglossa* and is similar to generalized members of related genera. Because of its generalized structure the phylogenetic position of the new genus relative to the other members of the supergenus is suggested by only two characters: absence of septomaxillary bones, and presence of paired premaxillaries. The absence of septomaxillary bones from most specimens of *Nyctanolis* (one of fourteen may have one septomaxillary) is a derived state that would place the new genus somewhere among the other genera. However, the erratic occurrence of these bones in species in which they are typically found (Wake, 1966), combined with the possible occurrence of septomaxillaries in one *Nyctanolis* specimen, weakens the value of this character. In having paired premaxillaries *Nyctanolis* is more primitive than any other member of the supergenus *Bolitoglossa*. Among neotropical salamanders only *Nyctanolis* has the premaxillary divided. This division is occasionally seen in an individual or two of some other species (Wake, 1966), but nowhere else is it characteristic of a taxon. This plesiomorphy argues strongly for the cladistic isolation of *Nyctanolis*; the new genus appears to be a sister

Figure 4. Dorsal view of the left foot of *Nyctanolis pernix*, cartilage stippled.

group to the rest of the supergenus (Fig. 5).

In many respects *Nyctanolis* preserves more plesiomorphic character states than any other member of the tribe Bolitoglossini; only those characters which diagnose the supergenus *Bolitoglossa* distinguish *Nyctanolis* from the ancestral state of all bolitoglossines. Members of the genus *Hydromantes* have the most primitive tail base region in the tribe, and have paired premaxillaries and well-developed septomaxillaries; however, they have a host of apomorphic characters, including a highly specialized projectile tongue and associated features, and they have lost prefrontals (Wake, 1966; Lombard and Wake, 1977). The species of *Batrachoseps* mainly have apomorphic characters, including only four toes, widely separated frontals and

parietals, and a strongly developed lateral parietal spur. All *Batrachoseps* are plesiomorphic relative to the other bolitoglossine supergenera in retaining genioglossus muscles as tongue retractors (although in a highly specialized form, see Lombard and Wake, 1977). While most *Batrachoseps* lack prefrontals and have but a single premaxillary, *B. wrighti* gains prefrontals and divided premaxillaries when it gets very large, and *B. campi* has these characters almost from the time of hatching (Marlow, Brode, and Wake, 1979). *Batrachoseps* has two or three caudosacral vertebrae and a wound-healing specialization, and in these respects is more apomorphic than *Hydromantes* but less than supergenus *Bolitoglossa*, all members of which have two caudosacral vertebrae and provision for autotomy of the tail at the base. In fact, only in this respect, and in the absence of well-developed septomaxillaries, is *Nyctanolis* less plesiomorphic than *B. campi*.

A feature uniting *Batrachoseps* and the supergenus *Bolitoglossa* is the reduction in number of pairs of diploid chromosomes from 14 to 13; unfortunately, we have no chromosomal information for *Nyctanolis*.

Nyctanolis probably represents the earliest surviving offshoot from the ancient ancestral stock of the supergenus *Bolitoglossa*. From a biogeographic perspective this is especially interesting, for this ancient group occupies not the northernmost part of the tropical distribution of bolitoglossines, where the most generalized forms have been found previously, but the eastern slopes of the mountainous core of Nuclear Central America. This area is one of three foci of evolutionary activity in tropical salamanders, and geographic isolates associated with it are thought to have been important in the adaptive radiation of tropical plethodontids (Wake and Lynch, 1976). The area in which *Nyctanolis* survives, now in a highly specialized ecological and

Figure 5. Cladogram showing the relationships of the members of the tribe Bolitoglossini. Principal synapomorphies characterizing each branch are as follows: a. Tribe Bolitoglossini, seven apomorphies discussed in text. b. Radii lost. c. Reduction in number of caudosacral vertebrae to two (three are still found in certain *Batrachoseps* but in these same species animals with two are common). d. Toe number reduced to four. e. Supergenous *Bolitoglossa*, tail base with complex of breakage specializations. f. See generic and species diagnoses in text. g. Characterized within the supergenus *Bolitoglossa* by fused premaxillae.

behavioral form, is the oldest land-positive area in all of Central America (Wake and Lynch, 1976; Rosen, 1978).

This is approximately 125 km ESE of the type locality (by air).

ACKNOWLEDGMENTS

Jeremy L. Jackson and H. Bradley Shaffer provided assistance and companionship during the field work of the senior author in Guatemala. We thank James F. Lynch, James Hanken, and Arden H. Brame, Jr. for review of the manuscript, and Gene M. Christman for preparing the illustrations. We appreciate the assistance of officials representing INAFOR for permission to conduct field work in Guatemala. The research has been supported by the National Science Foundation (current grant DEB-78 03008), the Museum of Vertebrate Zoology, and the National Institutes of Health (training grant 5-T32-GM07117).

For many years the work of Ernest E. Williams on the evolution of anoline lizards has been an inspiration for our work on the evolution of tropical salamanders. It is a privilege to be able to describe this anoline-like salamander in a volume dedicated to Professor Williams.

ADDENDUM

Since preparation of this paper, *Nyctanolis pernix* has been discovered at a third locality. Jonathan Campbell has collected one juvenile specimen (KU 189586) 2.4 mi. SE Purulha, Depto. Baja Verapaz, Guatemala, 1,615 m elevation.

LITERATURE CITED

- EDWARDS, J. L. 1976. Spinal nerves and their bearing on salamander phylogeny. *J. Morphol.*, **148**: 305-327.
- LOMBARD, R. E. 1977. Comparative morphology of the inner ear of salamanders (Caudata: Amphibia). *Contrib. Vert. Evol.*, **2**: viii + 140 pp.
- LOMBARD, R. E., AND D. B. WAKE. 1977. Tongue evolution in the lungless salamanders, family Plethodontidae. *J. Morphol.*, **153**: 39-79.
- MARLOW, R. W., J. M. BRODE, AND D. B. WAKE. 1979. A new salamander, genus *Batrachoseps*, from the Inyo mountains of California, with a discussion of relationships in the genus. *Contrib. Sci. Nat. Hist. Los Angeles Co.*, **308**: 1-17.
- ROSEN, D. E. 1978. Vicariant patterns and historical explanation in biogeography. *Syst. Zool.*, **27**: 159-188.
- SMITH, H. M., AND R. B. SMITH. 1976. Synopsis of the herpetofauna of Mexico, Vol. IV. Source analysis and index for Mexican amphibians. North Bennington, Vermont, John Johnson.
- TANNER, W. W. 1950. A new genus of plethodontid salamander from Mexico. *Great Basin Nat.*, **10**: 37-44.
- . 1952. A comparative study of the throat musculature of the Plethodontidae of Mexico and Central America. *Univ. Kansas Sci. Bull.*, **34**, pt. 2(10): 583-677.
- WAKE, D. B. 1963. Comparative osteology of the plethodontid salamander genus *Aneides*. *J. Morphol.*, **113**: 77-118.
- . 1966. Comparative osteology and evolution of the lungless salamanders, family Plethodontidae. *Mem. So. Calif. Acad. Sci.*, **4**: 1-111.
- WAKE, D. B., AND I. G. DRESNER. 1967. Functional morphology and evolution of tail autotomy in salamanders. *J. Morphol.*, **122**: 265-306.
- WAKE, D. B., AND J. F. LYNCH. 1976. The distribution, ecology, and evolutionary history of plethodontid salamanders in tropical America. *Nat. Hist. Mus. Los Angeles Co., Sci. Bull.*, **25**: 1-65.

Larvae of Southeast Asian Species of *Leptobrachium* and *Leptobrachella* (Anura: Pelobatidae)

ROBERT F. INGER¹

ABSTRACT. Seven larval forms of *Leptobrachium* Tschudi and *Leptobrachella* Smith (Anura: Pelobatidae) are identified from Southeast Asia. In terms of their buccopharyngeal and external anatomy, they are divisible into three groups that correspond to taxa proposed by Dubois (1980) on the basis of adult morphology: *Leptobrachium* (*Leptobrachium*), *Leptobrachium* (*Leptolalax*), and *Leptobrachella*. Larvae of *Leptobrachium* (*Leptobrachium*) have spheroidal bodies, numerous rows of denticles on both lips, no more than four pairs of papillae medially on the ventral velum, and well-developed glandular zones in front of the dorsal velum. All of these larvae live exposed in pools of small streams. Larvae of *Leptobrachium* (*Leptolalax*) have slender bodies, fewer rows of denticles, more than eight pairs of papillae medially on the ventral velum, no evident spicules, and reduced or absent glandular zones. Larvae of *Leptobrachella mjobergi*, the only larval form of this genus known, are like the preceding group in velar papillae and spicules and in reduction of the glandular zone. However, they are more slender, completely lack denticles, have only a single gill cavity on each side and lack ruffled filter rows. Larvae of the last two groups live in stronger currents than the first, and at least three of them are known to use interstices of stream bottoms. Limited analysis of food revealed no association with morphology or habitat.

INTRODUCTION

Larvae of the pelobatid genus *Leptobrachium* constitute a conspicuous portion of the tadpole assemblages in small streams of Southeast Asia and the Greater Sunda Islands. Most occur in relatively

quiet reaches of these streams (see for example Smith, 1917; Inger, 1966) and have the slightly depressed, spheroidal bodies we have come to associate with tadpoles inhabiting such environments. Others having more elongate bodies live in currents (Smith, 1917) and in the interstices of gravel on the bottom of riffles (Inger, 1966). The latter group of larval forms belong to species that Dubois (1980) has assigned to the subgenus *Leptolalax*, whereas the first group belong to species of the subgenus *Leptobrachium* (*sensu* Dubois, 1980). The two groups of larvae also differ in the number of rows of denticles and, as will be shown below, in features of the buccopharyngeal cavity. This range of variation in body form and denticular development is extended by Bornean larvae assigned here to *Leptobrachella*, which are even more elongate than larval *Leptolalax* and completely lack denticles. With material assembled for this study, eight larval forms are now known from the area south of China (see Table 1).

Although descriptions of external features of some of these larvae have appeared elsewhere (e.g., Smith, 1917; Taylor, 1962; Inger, 1966), the literature does not contain information on any given set of characters for all these larval forms. This paper attempts to correct this deficiency. Descriptions of the buccopharyngeal cavity in two larval forms of *Leptobrachium* (Wassersug, 1980) suggested a new source of taxonomic in-

¹Field Museum of Natural History, Chicago, Illinois 60605, U.S.A.

TABLE 1. EXTERNAL FEATURES OF SOUTHEAST ASIAN LARVAE OF SPECIES OF *LEPTOBRACHIUM* AND *LEPTOBRACHELLA*.

	<i>Leptobrachium</i>					<i>Leptobrachella</i>		
	<i>(Leptobrachium)</i>					<i>(Leptolalax)</i>		
	montanum	nigrops	hendricksoni	pullum	sp (Thai.)	gracilis	pelodytoides	mjobergi
Body shape								
wide and deep, width $\geq$ 0.60 HBL	+	+	+	+	+			
slender, width $<$ 0.55 HBL						+	+	+
Oral disk								
only upper lip notched	+	+	+	+	+			
both lips notched						+	+	+
number of divided rows of denticles								
upper lip $\geq$ 5	+	+	+	+	+			
upper lip $<$ 5						+	+	
lower lip $\geq$ 4	+	+	+	+	+			
lower lip $<$ 4						+	+	
denticles absent								+
denticles clawlike	+	+	+	+	+	+	+	
marginal papillae of upper lip								
bases confluent	+		+	+				
bases separated		+			+	+	+	+
inframarginal papillae, lower lip								
present, with denticles	+	+	+	+				
present, no denticles					+	+	+	+
Narial rim								
one small middorsal projection	+	+	+	+		+		+
2-3 small middorsal projections			+					
6-9 small middorsal projections					+			
4 subequal lobes around margin							+	
Spiracular tube								
no free portion	+	+	+	+				
free at tip					+	+	+	
free portion 2X opening diam.								+
Glandular patches								
present	+	+			+			
absent			+	+		+	+	+

*Exceeded in 1 specimen.

formation, which has been pursued in this study. In addition, a start has been made on analyzing food of some of these larvae with the ultimate goal, not yet realized, of relating diets to morphology and habitat distribution.

It is a pleasure to dedicate this paper to Dr. Ernest E. Williams who, despite his refusal to work on tadpoles, has done so much to stimulate study of morphology of reptiles and amphibians.

MATERIALS AND METHODS

Descriptions are based upon specimens preserved in ethanol and formalin. Most are from the collections of Field Museum of Natural History (FMNH), though a few are from the Bernice P. Bishop Museum (BBM) and the United States National Museum of Natural History (USNM).

Complete descriptions are given for

one form of each of the three main types of larvae: *Leptobrachium montanum*, *Leptobrachium gracilis*, and *Leptobranchella mjobergi*. Descriptions of other forms are abbreviated, emphasizing

points of difference from the respective basic type. Information on all is summarized in Tables 1-3. Body proportions, except where noted, are given in terms of head-body length

TABLE 2. VENTRAL SURFACES OF BUCCOPHARYNGEAL CAVITY OF SOUTHEAST ASIAN LARVAE OF SPECIES OF *LEPTOBRACHIUM* AND *LEPTOBRACHELLA*.

	<i>Leptobrachium</i>					<i>Leptobranchella</i>		
	<i>(Leptobrachium)</i>					<i>(Leptolalax)</i>		
	montanum	nigrops	hendricksoni	pullum	sp (Thai.)	gracilis	pelodytoides	mjobergi
Infralabial papillae								
anterior—large, transverse	+	+	+	+	+	+	+	+
posterior—2 or 3 branches						+		+
—4 or more branches	+	+	+	+	+		+	
Lingual papillae (paired)								
present	+	+	+	+	+	+	+	
absent								+
Buccal floor arena								
long lateral papillae	+	+	+	+	+	+	+	+
papillae across posterior								
present			+			+		+
absent	+	+		+	+		+	
central area								
30 or more pustules	+	+		+	+	+	+	+
papillae present			+			+		+
Buccal pocket area								
pre-pocket pustules or papillae								
present		+	+			+	+	+
absent	+			+	+			
post-pocket pustules or papillae								
present	+	+	+	+				+
absent					*	+	+	
Ventral velum								
pustules, dorsal surface								
2 groups						+		+
1 group, > 20	+	+	+	+			+	
1 group, < 20					+			
spicules, present	+	+	+	+	+			
absent						+	+	+
median papillae, 2-4 pairs	+	+	+	+	+			
8 or more pairs						+	+	+
secretory pits, dorsal surface								
on median papillae	+	+	+	+	*			
on margin	+	+	+	+				
absent						+	+	+
gill cavities, 3	+	+	+	+	+	+	+	
1								+
gill filter rows, present	+	+	+	+	+	+	+	
absent								+

*Condition obscure in dissection.

TABLE 3. DORSAL SURFACES OF BUCCOPHARYNGEAL CAVITY OF SOUTHEAST ASIAN LARVAE OF SPECIES OF *LEPTOBRACHIUM* AND *LEPTOBRACHELLA*.

	<i>Leptobrachium</i>					<i>Leptobrachella</i>		
	<i>(Leptobrachium)</i>					<i>(Leptolalax)</i>		
	montanum	nigrops	hendricksoni	pullum	sp (Thai.)	gracilis	pelodytoides	mjobergi
Prenarial papillae								
present	+	+	+	+	+	+		+
absent							+	
Narial walls, papillae								
each with one papilla	+	+	+	+	+		+	
one or both without papilla						+		+
Postnarial papillae								
more than 2 pairs	+	+	+	+	+	+	+	
pustules only								+
Lateral ridge papillae								
more than 2 branches	+	+	+	+	+		+	+
only 2 branches						+		
Median ridge								
semilunar			+				+	
triangular	+	+			+			
rectangular			+					
divided to base						+		+
Buccal roof arena, central portion								
with papillae	+	+			+	+		+
with pustules	+	+	+	+	+		+	
Glandular zone								
extends across roof	+	+	+	+	+			
restricted, medial or lateral							+	+
not evident						+		
Dorsal velum, papillae								
fewer than 15	+	+	+	+	+		+	
more than 15						+		+

(HBL). Terminology and sequence of description of internal buccopharyngeal anatomy follow Wassersug (1980).

specific status as *L. montanum* Fitzinger and *L. pullum* Smith, respectively.

Larval material examined: Sabah—BBM 7362, 7365; FMNH 63531-41, 77220-22, 130857-59, 131244, 140216. Sarawak—FMNH 77517, 83017-18, 96022, 121858, 139375-80, 148286, 212388-91.

TAXONOMY AND MORPHOLOGY

Leptobrachium montanum Fitzinger

Dr. Julian Dring (*in litt.*) has argued convincingly that the populations in Borneo and Thailand assigned to *hasselti* differ sufficiently from topotypic (i.e., Javan) *hasselti* Tschudi to warrant full

EXTERNAL FEATURES

Body oval, widest at level of spiracle; maximum width about 0.60 of length; somewhat flattened above, rounded below.

Eyes dorsolateral, not visible from below, oriented laterally, line connecting

pupils about 0.20 of HBL behind snout; length of eye 0.09; interorbital 0.27. Nostrils open, narial rim low with one simple weak projection middorsally; line connecting nostrils 0.10 of HBL behind snout; internarial 0.20; eye-nostril 0.11.

Oral disk ventral, subterminal, width 0.23 of HBL; lips continuous around margin except for narrow notch in center of upper lip; thick, homogeneous papillae in single continuous row around entire margin, bases of papillae contiguous; five to seven elongate, inframarginal ridges in diagonal row lateral to lower rows of denticles; largest of inframarginal ridges with small denticles. Dental formula in larvae state 26 or older usually $I:6+6/5+5:I$ or $I:5+5/5+5:I$, rarely more than six divided rows on upper lip or fewer than five on lower; undivided rows short, that of upper lip only half length of outer lower row; denticles laterally compressed, curved and pointed; dental ridges high. Beaks very heavy, fully pigmented, coarsely serrated; median cusp usually larger than adjacent ones.

Spiracle sinistral, slightly above mid-lateral line, tube fused to body wall dorsally and ventrally its entire length; opening 0.45 of HBL behind snout. Anus dextral, tube attached to ventral fin.

Tail 1.50 to 1.75 of HBL; weakly convex above and below, tapering to broadly rounded tip; maximum depth just behind mid-length, 0.52 of HBL, 0.27 of tail length. Caudal muscle heavy, 0.33 of HBL at end of body, 0.20 at maximum tail depth. Dorsal fin origin at end of body; fleshy in anterior fourth; margin rising only slightly; maximum depth subequal to muscle at point of maximum tail depth. Ventral fin not as deep as dorsal except near tip.

Head and body without asperities or spines. Lateral line system visible on HB and tail. Three whitish, subcircular glandular areas ventrally on each side of HB; each patch about size of pupil; two close together ventrolaterally behind level of eyes; third about twice its diameter anterolateral to base of anal tube.

Young larvae (i.e., stage 25) light with a few large dark spots on HB dorsally, usually with one large spot across root of tail and a few spots on caudal muscle; no spots on ventral surface of HB. Older larvae becoming darker, in many HB brownish dorsally and ventrally with obscure darker spots dorsally; caudal muscle and fins dusky with obscure darker spots.

HBL in stages 26–28 is 11.1 to 17.3 mm, stages 35–37 is 18.2 to 27.5 mm; maximum width of HB 0.59 to 0.64, mean 0.609; oral disk width 0.20 to 0.26, mean 0.231. Denticle formula (stages > 25) upper lip $I:6+6$ (66 tadpoles), $I:6+5$ (14), $I:5+5$ (52), $I:5+4$ (1); lower lip $6+6:I$ (1), $5+5:I$ (118), $5+4:I$ (6), $4+4:I$ (8).

BUCCAL CAVITY

Based on FMNH 63537, stage 36, HBL 17.5 mm and FMNH 139380, stage 37, HBL 20.5 mm.

Ventral surfaces. Two large infralabial papillae or palps on each side; one just inside lateral corner of beak compressed, with four long, tapering, pustulose branches; second medial and posterior to first, with thick base and five to seven unequal fingerlike projections, lateral two longest. A pair of simple lingual papillae, half length of posterior infralabials and medial to them. Tongue anlage a low swelling. Buccal floor arena roughly hexagonal; 10 to 13 large pointed papillae on each lateral border, all except end ones branched, posterior two-thirds of central area with many (35–50) low pustules. Buccal pocket transversely oval, perforated; no pre-pocket papillae or pustules. A row of six to seven small, slender papillae beginning posteromesad from buccal pocket and curving behind last three buccal floor arena papillae.

Ventral velum wide, with obvious spicules; dorsal surface between velar margin and buccal floor arena with numerous rounded pustules. Median area of velum set off by notches and divided into dorsal and ventral portions; dorsal portion with two pairs of large pa-

pillae, median ones longer; all with small knobs; ventral portion with one pair of papillae subequal to lateral ones of dorsal portion; all surfaces of median area with glandular pits. Lateral sections of velar margin with shallow curves marked by short projections.

Three gill filter cavities; 7 to 11 filter rows on filter plates; filter rows complexly folded and dense, but filter canals not roofed over.

Glottis on raised disk, just visible between median papillae of velum.

Dorsal surfaces. Prenarial arena narrow, long, with one pair of simple, slender papillae. Nares transversely oval, with thin walls; anterior wall with one inwardly curved, pointed, pustulose papilla laterally; higher posterior wall with similar papilla medially; relative heights of narial papillae vary. Postnarial arena with varying number of papillae; in one specimen three long, slender, unbranched papillae on each side; in second specimen two oblique, parallel rows of papillae, those of posterior row longer and pustulose. Lateral ridge papilla long, with one short branch. Median ridge triangular, margin pustulose. Buccal roof arena with 8 to 10 long pointed papillae laterally in staggered row, those in center of row with short branches; central area with many short papillae or pustules, decreasing in height posteriorly. A row of three to four short papillae lateral to buccal roof arena. A transverse band of secretory pits across roof between buccal roof arena and dorsal velum; band widest medially.

Dorsal velum with ca. 12 papillate projections on each side; medial ones longest. One large lateral pressure cushion visible in one larva; area damaged in dissection in other.

Leptobranchium nigrops Berry and Hendrickson

Figures 1–5

Larval material examined: Sarawak—FMNH 77512–13, 77578, 148284–85; 151515–21, 158209–14.

Figure 1. *Leptobranchium nigrops* (FMNH 148285), anterior portion of roof of buccal cavity and upper portion of oral disk ($\times 17$). Anterior at top.

Abbreviations: LRP—lateral ridge papilla (slightly damaged); MR, median ridge; NPA, anterior narial papilla; NPP, posterior narial papilla; UB, upper beak; ULP, upper marginal labial papilla.

EXTERNAL FEATURES

Very similar to *L. montanum* larvae in general form and proportions, differing mainly in coloration in life and in shape of dorsal fin. Beak more coarsely serrated, and marginal papillae of upper lip more distinctly separated than in *montanum* (Fig. 1). Denticles (Fig. 2) as in *montanum*.

Dental formula I:6+6/5+5:I except in young larvae (stage < 26). Dorsal fin origin slightly behind end of body, fin rising abruptly, margin more convex than in *montanum*; otherwise tail form similar to that of latter species. HB (in life) with middorsal stripe of gold chromatophores; in preservative HB straw-colored, with a dark middorsal stripe of melanophores; a dark interorbital band; a dark transverse band or a pair of large, narrowly, sepa-

Figure 2. *Leptobrachium nigrops* (FMNH 77513), denticles of lower lip ($\times 963$).

Figure 3. *Leptobrachium nigrops* (FMNH 77513), anterior portion of floor of buccal cavity ($\times 42$). Anterior at top.

rated, dark spots at rear of HB; ventrally HB without markings.

HBL of two largest stage 25 tadpoles 17.5 and 18 mm, two in stages 31–32 24.5 and 26.6 mm, one stage 38 22 mm.

BUCCAL CAVITY

Based on FMNH 77513, stage 25, HBL 14.8 mm and FMNH 148285, stage 32, HBL 24.5 mm. Very similar to *L. montanum*.

Ventral surfaces (Figs. 3, 4). Buccal floor arena with 50 to 100 low pustules; anterior to lateral papillae a low, thick, transverse ridge bearing several short papillae. Several short papillae and pustules in front of buccal pockets.

Dorsal surfaces (Figs. 1, 5). Prenarial area with a large 3-branched papilla just behind lateral corner of beak. Postnarial arena with six to eight pustulose papillae on each side. Lateral ridge papilla with one short branch near base and two longer terminal ones. An elongate oval area of pustules and papillae lateral to buccal roof arena. Band of secretory pits widest laterally.

Abbreviations: ILPA, anterior infralabial papilla; ILPP, posterior infralabial papilla; LP, presumed lingual papilla.

Leptobrachium hendricksoni Taylor

Differences between larvae of *hendricksoni* and *montanum* support the differences between adults noted by Taylor (1962).

Larval material examined: Malaya—BBM 6965, 7137; FMNH 134544. Sarawak—119898–99, 129014, 129016–7, 129019–20, 148282–83, 158215–18.

EXTERNAL FEATURES

Very similar to larvae of *montanum* and *nigrops* in general body form and oral disk, but differing from them in coloration and in absence of glandular patches ventrally on HB.

Narial rim low, in Bornean larvae one small middorsal projection flanked by

Figure 4. *Leptobrachium nigrops* (FMNH 148285), rear portion of floor of buccopharyngeal cavity ($\times 18$). Anterior at top.

Abbreviations: BFA, buccal floor arena; FC, filter cavity; GL, glottis; VVMP, median papillae of ventral velum; VVP, pustules of ventral velum.

Figure 5. *Leptobrachium nigrops* (FMNH 148285), left rear portion of roof of buccal cavity ($\times 18$). Anterior at top.

Abbreviations: BRA, buccal roof arena; DV, dorsal velum; GZ, glandular zone.

several subsidiary ones giving impression of a low, pustulose triangular projection; Malayan larvae with simple, low middorsal projection. Oral disk resembling that of *montanum*; dental formula usually $I:7+7/6+6:I$ or $I:6+6/6+6:I$; denticles shaped like those of *montanum*. Tail form as in *montanum*. No glandular patches evident ventrally on HB. In preservative reddish brown above and on tail, with numerous small dark, round dots on all surfaces.

HLB of largest stage 25 tadpole 23.0 mm, stages 26–28 (8 individuals) 24.2 to 30.0 mm, stages 35–37 (6) 25.8 to 34.4 mm. Width of HB 0.54 to 0.66 (mean 0.600), eye length (stages > 25) 0.08 to 0.12 (mean 0.092) width of oral disk 0.19 to 0.23. Denticle formula (stages > 25) upper lip $I:7+7$ (4 tadpoles) $I:6+7$ (1), $I:6+6$ (13); lower lip $6+6:I$ (13), $6+5:I$ (1), $5+5:I$ (4).

BUCCAL CAVITY

The larva (FMNH 148282) described by Wassersug (1980) under the name *L. hasselti* is *L. hendricksoni*. A stage 32 larva (FMNH 148283), dissected in the present study, agrees in almost all particulars with Wassersug's description. This specimen has a trilobate papilla anterior to the infralabial papilla and lateral to the three pustulations noted by Wassersug. The buccal cavity is generally similar to that of *L. montanum*.

Ventral surfaces. Buccal floor arena with 12 to 20 short papillae across its posterior fourth; a few low pustules medial to lateral papillae. About six pre-pocket papillae.

Dorsal surfaces. Prenarial area with two to three low pustules on each side. Postnarial arena with three short papillae on each side. Lateral ridge papilla bifur-

cate near tip. Median ridge semilunar; margin with small projections. Buccal roof arena medially with many pustules, densest posteriorly. Distribution of pits in transverse glandular zone as in *nigrops*. Dorsal velum with seven to nine papillae on each margin.

Leptobrachium pullum Smith

Four samples of larvae from southern Thailand (Naknon Si Thammarat Province, Ron Phi Bun—FMNH 173481, 173583, Phangnga Province, Lam Pi—FMNH 206820; probably from Trang Province—FMNH 172253) have the general facies of larval *L. montanum* and appear to belong to a single species, which I tentatively assign to *L. pullum* Smith (see above, p. 16). Although these samples are from an area close to the range of *L. hendricksoni*, which Taylor (1962) reported from Yala Province, Thailand, and not too distant from that of *L. nigrops* in central Malaya, these putative *pullum* larvae differ from those of *hendricksoni* and *nigrops*. The coarse serration of their beaks, the thickness of their labial papillae, and their coloration are quite different from the conditions in larval *hendricksoni*. The confluence of the bases of their upper labial papillae, a light triangular area at the root of the tail, and the apparent absence of ventral glandular patches distinguish them from larval *nigrops*. In the last two characters they also differ from larval Bornean *montanum*.

As noted earlier, J. Dring (*in litt.*) has suggested that all Thai frogs previously assigned to *L. hasselti* Tschudi are probably conspecific with *L. hasselti pullum* Smith, which Dring believes warrants full species recognition.

EXTERNAL FEATURES

Body form in general as in *L. montanum*, i.e., oval, flattened above, spheroidal below; narial rim as in *montanum*; labial papillae thick, bases of those on

upper lip confluent; dental formula most often 1:6+6/5+5:I or 1:5+5/5+5:I; denticles shaped as in *montanum*; beaks coarsely serrated, upper with enlarged median cusp. Tail shaped as in *montanum*. Glandular patches not evident ventrally on HB.

Color (in alcohol) pale to dusky on HB with many small, irregular dark spots dorsally; a conspicuous triangular light area over root of tail bounded anteriorly by wide transverse dark band and posteriorly by an oblique band, the two bands joining laterally; ventrally HB unmarked or with a few obscure spots in anterior fourth; tail muscle with large dark spots mostly in anterior half; fins with smaller, less conspicuous spots.

HBL of largest stage 25 tadpole is 16 mm, stages 26–28 is 15.8 to 19.2 mm, stages 35–37 is 21.5 to 24.7 mm; maximum body width 0.51 to 0.64 (mean 0.563); eye 0.09 to 0.11; width of oral disk 0.22 to 0.28 (mean 0.253); denticle formula (stages > 25) of upper lip 1:6+6 (9), 1:6+5 (1), 1:5+5 (5), of lower lip 5+5:I (13), 4+4:I (2).

BUCCAL CAVITY

Based on FMNH 172253, stage 36, HBL 22.0 mm and FMNH 173481, stage 36, HBL 24.3 mm. Generally similar to *L. montanum*.

Ventral surfaces. Posterior infralabial papillae with three major branches, each further subdivided into pustulose twigs. Central area of buccal floor arena with 40 to 50 low pustules.

Dorsal surfaces. Lateral ridge papilla slender, unbranched. Median ridge low, rectangular; margin sinuate; one low projection at lateral corner. Central area of buccal roof arena with many low pustules. A cluster of 8 to 10 papillae lateral to posterior papillae of buccal roof arena. Glandular zone widest laterally.

Leptobrachium sp.

One sample of two tadpoles from northern Thailand (Chiang Mai Province,

Doi Suthep—FMNH 212387) and another slightly shriveled series of two from Annam, Indo-China (FMNH 26505) appear to represent another species of the *L. montanum* group. The well-preserved sample from Doi Suthep differs from those previously described in this paper in having conical, denticle-less inframarginal papillae laterally on the lower lip and in having seven to nine small projections on the dorsal rim of the nares. The Annam larvae have the same type of inframarginal papillae on the lower lip, but their unsatisfactory state of preservation makes count of narial projections uncertain, although they clearly have more than one small projection. Both series also differ from larval *pullum* and *montanum* in having the bases of the upper labial papillae slightly separated, and from larval *hendricksoni* in coloration, upper labial papillae, and in serration of the upper beak.

Because there still remains some doubt about definition of species in the Thai members of this group, these larval series are not assigned a specific name.

EXTERNAL FEATURES

Body form and oral disk generally as in *L. montanum*, though slightly more slender; HBL at stage 30 is 26.0 mm, at stage 38 is 28.3 mm. Narial rim low with six to nine low pustules dorsally. Papillae of upper lip with bases not confluent; inframarginal papillae of lower lip conical, without denticles; width of oral disk 0.27; dental formula I:5+5/4+4:I; denticles as in *montanum*; beaks coarsely serrated, upper with enlarged median cusp. Tail shaped as in *montanum*. Glandular patches ventrally on HB as in *montanum*. HB and caudal muscle with very fine colorless asperities.

Color in formalin dark grayish brown on HB and tail; HB without evident pattern on spotting; indistinct dark spots or bars on tail; fins with dense dusting of melanophores.

BUCCAL CAVITY

Based on FMNH 212387, stage 30, HBL 26.0 mm. Generally similar to *L. montanum*.

Ventral surfaces. Buccal floor arena with about 40 low pustules. Dorsal surface of ventral velum with 12 low papillae or pustules; secretory pits not evident on dorsal surfaces of velum.

Dorsal surfaces. Postnarial arena with four slender papillae on each side. Lateral ridge papilla with two anterior and one posterior conical projections. Median ridge triangular, margin with seven projections. Buccal roof arena medially with dozen papillae anteriorly and equal number of low pustules posteriorly.

Leptobrachium spp. (Malaya)

Two samples of larvae from West Malaysia, one from Johore and one from Pulo Tioman, resemble larval *L. montanum* in form, but differ from each other and other larvae described in this paper. The only *Leptobrachium* known from the Malay Peninsula to which larvae have not been assigned is *L. heteropus*. No species of *Leptobrachium* has been reported from Pulo Tioman (Hendrickson, 1966). None of these tadpoles in these two samples is developed to the point at which diagnostic features of adults can be discerned.

BBM 7162—Mawai Estate, ca. 2.5 km north of Kota Tinggi, Johore, West Malaysia. Seven individuals, stages 25–39.

EXTERNAL FEATURES

Shape of oral disk as in *L. montanum*, body more slender. HBL 15.6 to 18.5 mm, stages 36–39; maximum body width 0.35 to 0.49. Nares large, narial rim very low, with single very weak mid-dorsal projection present in two. Papillae of upper lip with bases separated; a few rounded inframarginal papillae on lower lip, several with small denticles; width of

oral disk 0.17 to 0.20; dental formula 3+3/3+3:I (stages 31–39); beaks black in marginal half; coarsely serrated, no enlarged median cusp. Tail with straight margins, tapering abruptly in final third. Spiracular tube free at tip. Vent median, opening long and exposed. Coloration obscured in preservative.

BUCCAL CAVITY

Based on one tadpole at stage 38 (HBL 18.5 mm).

Ventral surfaces. Anterior infralabial papillae well separated from beak, each with narrow base and three short branches; posterior infralabial papillae with thick cylindrical bases, each with three branches in terminal half. Buccal floor arena oval, bounded laterally by a partially doubled row of 12 long, tapering papillae; no pustules or papillae in central area. No pustules or papillae between buccal floor arena and margin of ventral velum. Ventral velum with well-developed spicules. Median fifth of velum divided into dorsal and ventral portions, each with two pairs of thick papillae. Three subequal gill cavities; filter rows complexly branched; filter canals canopied.

Dorsal surfaces. Prenarial arena as wide as long; one pair of short papillae pustulose. Nares slitlike, with low thick walls, the posterior one drawn out laterally into short, thick, valvelike projection; no narial papillae. Postnarial area with deep, narrow pit behind each naris, pits narrowly separated; two short papillae behind each pit. Lateral ridge papilla long, slender, with three subequal branches in distal half. Median ridge semilunar, low, margin pustulose. Buccal roof arena with 12 to 14 long papillae in partly doubled row laterally; a single pustule visible in central area. Three to five secretory pits in extreme posterolateral corner of buccal roof. Surface from center to rear of buccal roof "furry" but without distinct pustules or papillae. Dor-

sal velum deep; median half of left side with five short, thick projections; right side damaged.

BBM 6972—Ulu Lalang, Pulo Tioman. One individual, stage 26.

EXTERNAL FEATURES

Body form as in *L. montanum*, HBL 22.6 mm, maximum width 0.73. Nares as in *L. montanum*. Width of oral disk 0.32; both lips with deep median notch; marginal papillae of upper lip with separate bases; lower lip very broad anteroposteriorly, surface covered with pillowlike, circular swellings lacking denticles; swellings in three irregular rows, continuing around corners of disk to upper lip where they become limited to single row reaching median notch of lip. Dental formula 1:4+4/3+3:I, undivided upper row one-fourth length of corresponding lower row; beaks completely black, coarsely serrated, no median cusp. No glandular patches. The tail is damaged, obscuring the shape, but the muscle is heavy. HB and tail are dusky in formalin without evident markings.

Leptobranchium gracilis Günther Figures 6–8

Larval material examined: Sabah—FMNH 130860, 130864, 131245–46, 212381–82. Sarawak—FMNH 77509, 77511, 77516, 139381, 146323, 146341.

EXTERNAL FEATURES

Head and body forming elongate oval, flat above, slightly rounded below; maximum width about 0.50 of length.

Eyes dorsolateral, not visible from below, oriented laterally, line connecting pupils 0.20 of HBL behind snout; length of eye varying with development of larva, about 0.06 in stage 36; interorbital about 0.15. Nostrils open; narial rim low with one low, simple projection mid-dorsally;

nostrils 0.08 behind snout, internarial about 0.15; eye-nostril varying with development, about 0.09 in stage 36.

Oral disk ventral, subterminal, cuplike; width about 0.20; both lips with deep, median indentation; small homogenous papillae in single row around entire margin of lips; small round inframarginal papillae flank lower dental rows; inframarginal papillae without denticles; broad band of lower lip smooth between dental rows and marginal papillae. Dental formula variable, in stages > 25 upper lip 1:3+3 to 1:6+6, lower lip usually 2+2:1; undivided row of lower lip longer than first divided row; undivided row of upper lip around one-third length of lower one; denticles laterally compressed, curved and pointed (Fig. 6). Beaks heavy, fully pigmented, finely serrated; upper beak without enlarged median cusp.

Spiracle sinistral, mid-lateral; with raised ventrolateral edge, free only at end; opening 0.50 behind snout. Anal tubedextral, attached to ventral fin.

Tail about 2.0 of HBL, margin very slightly convex, tapering gradually in last third to rounded tip; maximum depth before mid-length, 0.24 of HBL, about 0.10 of tail length. Caudal muscle heavy, 0.26 of HBL at end of body, not tapering until point of maximum tail depth. Dorsal fin origin just beyond level of anal opening; very low in proximal fourth; equal to depth of muscle only in distal third. Ventral fin deeper than dorsal only in proximal fourth.

Head and body without asperities or glandular patches. Lateral line system visible on HB and tail.

Coloration (in preservative) with light or heavy dusting of melanophores dorsally on HB and proximal two-thirds of caudal muscle; lower half of body, fins, and distal third of caudal muscle without pigment.

HBL of two largest stage 25 larvae is 19.3 and 20.0 mm, two in stages 35–37 are 21.6 mm; maximum width of HB 0.42 to 0.51; width of oral disk 0.17 to 0.22. Denticle formula (stages > 25) of upper lip

Figure 6. *Leptobrachium gracilis* (FMNH 146323), denticles of upper lip (x2300).

I:6+6 (1 larva), I:4+4 (5), I:3+3 (1); lower lip 2+2:I (6), 3+3:I (1).

BUCCAL CAVITY

Based on FMNH 146323, stage 35, HBL 21.6 mm.

Ventral surfaces (Fig. 7). Anterior infralabial papilla a thick transverse flap just behind lateral corner of lower beak; separated from that of other side by about half its length; lateral corner low, bearing short, smooth, conical spur; rest of papilla with thick pustulose projections; median portion highest. Posterior infralabial papilla with thick base and three terminally pustulose branches. A laterally curved simple papilla between the infralabial papillae. A pair of conical, smooth lingual papillae between posterior infralabial papillae, their bases continuous with those of the latter. Tongue anlage low. Buccal floor arena (Fig. 7) a wide rectangle, bounded laterally by eight to nine papillae; several of largest papillae with confluent bases and three to four sharp, posteriorly directed spurs; midline of floor arena with three thick, short pustulose papillae in center; about 40 low pustules in transverse band across rear third of floor arena, continuous with pustulose area of velum. Buccal pocket an inverted U-shaped slit; perforation not evident, an oblique row of three smooth pre-pocket papillae. Area behind pocket without papillae or pustules.

Ventral velum (Fig. 7) wide, without evident spicules; a deep median notch; dorsal surface densely covered medially with pustules except for bare strip in mid-line; laterally dorsal surface of velum smooth; margin without secretory pits. Median area of velum divided into dorsal and ventral portions; dorsal portion with two pairs of papillae, outer pair longer; ventral portion a forest of filamentous papillae, at least 20 on each side; no secretory pits on velar papillae. Lateral area of velum sinuate; without projections or marginal pits.

Three filter cavities, lateral one half

Figure 7. *Leptobrachium gracilis* (FMNH 146323), floor of buccopharyngeal cavity behind tongue anlage ($\times 18$).

Abbreviations: Anterior at top. BP, buccal pocket; other abbreviations as in Fig. 4.

length of others; 8 to 11 filter rows on filter plates; filter plates bordering middle cavity with a few dorsally directed, branching projections (Fig. 7, lower left); filter canals open.

Glottis hidden by fringing papillae of velum.

Dorsal surfaces (Fig. 8). Prenarial arena narrow, long; one pair of thick, low papillae just behind center. Nares obliquely oval, separated by length of one opening; anterior wall thin, low, with pustulose margin; posterior wall slightly higher, with one long, smooth papilla medially. Postnarial area with row of three flattened, triangular papillae on each side; central one of each set about twice height of others. Lateral ridge papilla a simple flat, elongate triangle, bifurcate at tip. Median ridge directly be-

Figure 8. *Leptobrachium gracilis* (FMNH 146323), roof of buccal cavity behind internal nares ($\times 18$). Anterior at top.

Abbreviations: see Figs. 1, 5.

hind gap in postnarial papillae deeply divided into three smooth, triangular lobes. Buccal roof arena (Fig. 8) oval, with about 30 smooth, unbranched, pointed papillae. A transverse band of pustules or low papillae behind roof arena, extending beyond lateral margins of roof arena; posterior border of band curved and pointed forward at midline; pustules along posterior border higher; anterolateral corner of band with three to four long papillae; about 150 pustules on each side of midline. No secretory pits.

Dorsal velum with 18 to 20 filamentous papillae on margin of each side; papillae subequal to those of ventral velum.

Leptobrachium pelodytoides Boulenger

Dubois (1981) suggests that *L. minimum* Taylor (type locality Doi Suthep, Chiang Mai Province, Thailand) is con-

specific with *L. oshanense* Liu (type locality Mt. Emei, Sichuan). Dring (*in litt.*) has suggested that *minimum* is conspecific with *L. pelodytoides* Boulenger (type locality Karen Hills, Burma). The material available for this study is not adequate to resolve this issue.

The Thai larvae listed below differ from our Chinese tadpoles of *L. oshanense* Liu in two respects; in the former the beaks are more coarsely serrated, and the rows of denticles on the lower lip are flanked by several rounded inframarginal papillae, which are absent in the Chinese series (see Liu, 1950: Fig. 42). The Thai larvae have dark spots on the tail as figured by Smith (1917) in his description of larval *pelodytoides*. Smith also illustrated inframarginal papillae, though on both lips. Tentatively, given these slight differences between Thai and Chinese tadpoles, I assign the former samples to *pelodytoides*.

Larval material examined: Thailand—FMNH 174525, 212386, USNM 130402. In addition several samples of larval *L. oshanense* Liu from China—FMNH 24419, 49589.

EXTERNAL FEATURES

Body form similar to that of *L. gracilis*; maximum width 0.50 to 0.55 of HBL; eye length (stage 35) 0.09; interorbital about 0.20. Narial rim prominent, perimeter divided into four subequal lobes. Oral disk as in *gracilis*; marginal and inframarginal papillae as in *gracilis*; dental formula I:2+2/2+2:I (4) or I:3+3/2+3:I (1), proportions of rows as in *gracilis*; beaks coarsely serrated, median cusp of upper beak not enlarged. Spiracular tube as in *gracilis*. Tail shape as in *gracilis*. No spinules or glandular patches evident.

Color (in preservative) brown with irregular dark spots on caudal muscle and dorsal fin; ventral fin colorless proximally, becoming brownish as dorsal fin in distal half; ventrally HB without melanophores.

HBL of largest stage 25 is 13.0 mm, stage 35 is 20.5 mm.

BUCCAL CAVITY

Based on FMNH 212386, stage 35, HBL 20.5 mm. Generally similar to *L. gracilis*.

Ventral surfaces. Posterior infralabial papillae with three major branches, middle one complexly subdivided. Central portion of buccal floor arena with ca. 100 low pustules, densest posteriorly. Band of pustules on dorsal surface of ventral velum without median interruption.

Dorsal surfaces. Prenarial area without evident pustules or papillae. Anterior narial wall with one long papilla laterally. Postnarial arena with three triangular, flattened papillae on each side. Lateral ridge papilla with four or five divisions of varying lengths. Median ridge semi-lunar with 10 short marginal projections. Central portion of buccal roof arena with ca. 70 pustules. Glandular zone limited to extreme posterolateral corner of roof. Dorsal velum with 8–10 long flattened papillae on each side.

Leptobranchella mjobergi Smith
Figures 9–10

A number of Bornean samples of pelobatid larvae, erroneously assigned to *Leptobranchium gracilis* (Inger 1966), resemble that species in having deep cuplike oral disks, but differ from that form and all other larvae previously described in this paper in lacking denticles (Fig. 9). These larvae further differ from *gracilis* in having more slender bodies (width of HB in stages > 25 0.32 to 0.41, in *gracilis* 0.43 to 0.50) and in having the spiracular tube free of the body wall for a distance equal to about twice the diameter of the opening. In *gracilis* the spiracular tube is free only at the very tip. Association of these larvae with *Leptobranchella* is based on a stage 41 larva that has the pointed digits peculiar to this genus (Inger, 1966: Fig. 8A).

The samples listed below may comprise more than one species. The inframarginal papillae in those from eastern Sabah (FMNH 77503–04) are restricted

Figure 9. *Leptobranchella mjobergi* (FMNH 77503), anterior portion of floor of buccal cavity ($\times 44$). Anterior at top.

Abbreviations: IMP, inframarginal papilla of lower lip; LB, lower beak; other abbreviations as in Figs. 1, 3, and 4. Left ILPP cut and tip of right obscured by detritus. Left edge of BFA damaged.

to the vicinity of the beaks leaving a wider portion of the lips smooth than is the case in the other samples. Also one of these Sabah larvae (FMNH 77503, stage 36) differs from one from Mount Kinabalu (FMNH 130861) in the amount of pustulation on the ventral velum within the buccal cavity (see below). Although only two larvae were dissected, this variation exceeds that observed in the small samples of other species.

Larval material examined: Sabah—BBM 7363, FMNH 77503–04, 130861–63. Sarawak—FMNH 77515, 157998, 212384–85.

EXTERNAL FEATURES

Head and body extremely elongate, flat above, rounded below; maximum width about 0.35 of HBL.

Eyes dorsolateral, not visible from below, oriented laterally; eyes 0.18 behind snout; length of eye varying with development, about 0.07 in stages 35–37; interorbital about 0.10. Nostrils open; rim not raised except for small middorsal projection; nostrils about 0.07 behind snout; internarial about 0.10.

Oral disk ventral, subterminal, cuplike; width about 0.15; lower lip with deep notch, upper with shallow notch; very short, fine papillae in single row around margin of lips; round inframarginal knobs close to beaks (Fig. 9); area between margin of lips and inframarginal papillae variable. No denticles, no long dental ridges (Fig. 9). Beaks fully pigmented, finely serrated, median cusp of upper beak slightly enlarged.

Spiracle sinistral, mid-lateral; spiracular tube free for distance 2 to 2.5 times diameter of opening; opening usually less than 0.50 behind snout. Anal tube dextral, attached to fin.

Tail about 2.0 of HBL, margins roughly straight most of length, tapering abruptly near end to rounded point; maximum depth at beginning of distal third, about 0.40 of HBL. Caudal muscle very heavy, almost as wide as HB; not tapering appreciably until distal third; three to five times depth of fins at mid-length. Dorsal fin origin behind HB; thick and low in proximal half of tail; equal to depth of muscle only in distal third. Ventral fin deeper than dorsal in proximal half.

Color in preservative pale yellowish brown without markings, in life pinkish gray.

HBL of two largest in stage 25 are 12.5 and 13.8 mm, of three in stages 26–27 are 11.5 to 13.0 mm, of three in stages 35–37 11.5 to 13.2 mm; maximum width of HB 0.31 to 0.41; width of oral disk 0.13 to 0.17.

BUCCAL CAVITY

Based on FMNH 77503, stage 36, HBL 11.5 mm, and FMNH 130861, stage 30, HBL 13.8 mm. The narrow heads made dissection of these larvae difficult and both were slightly damaged.

Ventral surfaces (Fig. 9). Anterior infralabial papilla immediately inside lower beak, transversely oriented, thick, lowest laterally and thickly pustulose in median two-thirds; narrowly separated from flap of other side. Posterior infralabial papilla thick, flattened with two smooth, subequal branches in distal half. Two or three smooth conical papillae between and almost confluent with the infralabial papillae. No lingual papillae. Tongue anlage low. Buccal floor arena trapezoidal, wider posteriorly, bordered laterally by 10 to 12 long, slender, simple papillae, bases contiguous; central area of arena with three to four short, thick papillae in mid-line anteriorly, followed by an expanding triangular zone of longer papillae and pustules; between this zone and lateral papillae 20 to 30 low pustules on each side. Buccal pocket a broad inverted U, perforate; three to five short papillae anterior to pocket parallel to buccal floor arena. Area behind pocket with three to four papillae.

Ventral velum (Fig. 10) wide, without evident spicules; a deep median notch; dorsal surface with two large, rounded groups of pustules narrowly separated in mid-line, anterior pustules larger, ca. 75 in each group; laterally dorsal surface of velum smooth in FMNH 77503, with low pustules in FMNH 130861; margin divided into two lobes on each side; margin without secretory pits. Median lobe of each side divided horizontally; dorsal portion with two long papillae; ventral portion with more than 10 filamentous papillae. Lateral lobe with two to three short marginal papillae.

Only one filter cavity on each side; no ruffled filter rows.

Glottis in raised disk, under median notch of velum.

Dorsal surfaces. Prenarial area long and narrow, bounded at rear by row of three to four short papillae. Nares obliquely oval; narial walls thin, low; both walls with one or two pustules, no papillae. Postnarial area smooth or with several indistinct pustules. Lateral ridge

Figure 10. *Leptobranchella mjobergi* (FMNH 77503), median portion of right half of ventral velum ($\times 82$). Anterior towards top.

Abbreviations: see Fig. 4.

papilla with two long, subequal branches. Median ridge with three triangular lobes, middle one slightly wider in one specimen; margins without pustules; one short papilla narrowly separated from median ridge on each side. Buccal roof arena occupying full width of buccal cavity; with about 40 pointed papillae decreasing in size posteromedially; one near center bifurcate; rear of arena with small, smooth triangular area apointed forward in FMNH 130861. A large transverse band of postules behind buccal roof arena; ca. 100 to 125 pustules, largest ones posteriorly; rear boundary of pustulose area arched forward medially. Indistinct glandular zone behind center of pustulose band.

Dorsal velum with 15 to 20 long papillae on margin of each side.

HABITAT DISTRIBUTION AND FOOD

All of the larvae considered here were collected in streams, although that term encompasses a variety of microhabitats. Precise information on microhabitats is available for only a portion of the specimens examined and is summarized in Table 4.

The results are not surprising. The heavy-bodied forms are largely confined to areas of slow or no current and the slender-bodied ones (except for one sample of *Leptobranchium pelodytoides*) to areas of stronger current. At the time of collection of the *L. pelodytoides* sample, the stream had dried into a series of small pools (notes of D. L. Damman). The tabulation does not show, however, an important difference between *montanum*, *hendricksoni*, and *nigrops*, on the one hand, and *gracilis* and *Leptobranchella* on the other. Individuals of the former group, though feeding on the bottom, are usually exposed on the substrate or hidden among dead leaves covering the bottom, whereas those of the second group are rarely exposed on the substrate and are usually within the interstices of the bottom gravel or boulders. Liu (1950) observed *Leptobranchium oshanense*, which is generally similar to *gracilis* in body form and oral apparatus, both in crevices between rocks and on the surface of the substrate.

Gut contents of *Leptobranchium montanum* (3 individuals), *pullum* (2), *hendricksoni* (2), and *gracilis* (2) were studied by means of light microscope (in Hoyer's medium) and SEM. Only the most anterior portion of the gut was used. In all cases the most abundant type of food was plant fragments—either parts of leaves or vascular tubes. The SEM revealed diatoms and fragments of hyphae or other fungal parts in all except *gracilis*. Larvae of *montanum*, reared in a field laboratory, fed actively on dead leaves and fragments of prawn, but did not attack one another. Size range of food fragments, selected as representative and

TABLE 4. HABITAT DISTRIBUTION OF LARVAE OF CERTAIN SOUTHEAST ASIAN SPECIES OF PELOBATIDS.

Species	Quiet pools*		Side pools†		Shingle beds		Riffles	
	Ind.‡	Coll.§	Ind.	Coll.	Ind.	Coll.	Ind.	Coll.
<i>Leptobrachium montanum</i>	123	13	11	4	2	2	22	3
<i>L. hendricksoni</i>	15	2	12	1				
<i>L. nigrops</i>	66	3	6	2				
<i>L. sp</i>	2	1						
<i>L. gracilis</i>							18	6
<i>L. pelodytoides</i>	2	1						
<i>Leptobrachella mjobergi</i>	1	1			2	2	14	5

*Areas of slow to moderate current.

†Areas cut off from the current, usually by gravel bars and often the site of accumulations of dead leaves.

‡Number of individuals.

§Number of collections.

TABLE 5. SIZE DISTRIBUTION OF REPRESENTATIVE FOOD FRAGMENTS FROM STOMACHS OF LARVAL *LEPTOBRACHIUM*.

Species	Greatest dimension (mm) of food item					
	< 0.01	0.01–0.05	0.06–0.10	0.11–0.20	0.21–0.30	0.30+
<i>L. montanum</i>	1	6	0	2	1	4
<i>L. hendricksoni</i>	1	5	0	2	1	3
<i>L. pullum</i>	1	3	5	2	1	1
<i>L. gracilis</i>	0	0	4	3	4	2

measured on SEM photographs, shows little differentiation among species (Table 5). As this is not an exhaustive or random sampling, rigorous statistical analysis is not appropriate.

DISCUSSION

Except for the genus *Megophrys*, larvae of Asiatic megophryine genera have depressed, spheroidal bodies even though all live in streams (Smith, 1917; Pope, 1931; Liu, 1950; Inger, 1966). The slender bodies of larval *Leptobrachium* (*Leptolalax*) and *Leptobrachella* may consequently be considered as a derived, specialized condition. These two groups also share other characteristics that distinguish them from all other non-*Megophrys* megophryines; 1) expanded, cuplike lips; 2) reduced or no spicules in the ventral velum; 3) dense clusters of

filamentous papillae medially on the ventral velum; and 4) reduction of the glandular zone in front of the dorsal velum (Tables 2, 3). As well-developed glandular zones and spicules characterize not only other megophryines (including *Megophrys minor*) but also *Scaphiopus bombifrons* (Wassersug, 1980), character states (2) and (4) appear to be derived. The cluster of filamentous velar papillae (3) is unlike anything observed by Wassersug (1980) in other larvae, except *Leptobrachium* (*Leptolalax*) *oshanense*. The expanded lips, though their function is still not clear (Liu, 1950), also represent a derived condition within the subfamily. At this time only one of these character states, the slender body form which in several species is associated with the habit of using bottom interstitial spaces, has an apparent functional significance. The sharing of these derived character states by all known larvae

of species of *Leptobrachium* (*Leptolalax*) supports Dubois's (1980) recognition of this taxon.

Larvae of *Leptobrachella* have three additional, functionally puzzling specializations: a single filter cavity on each side, loss of the usual frilled gill filter rows, and the absence of denticles. The first seems to be a further development of a condition in *Leptobrachium gracilis*, which has one filter cavity much reduced in size. These two larval forms utilize a microhabitat different from those of other tadpoles that have filter cavities reduced in size (Wassersug, 1980: 106) or number (Wassersug *et al.*, 1981). The reduction may be related to life in well oxygenated waters. The other two specializations suggest modified feeding behavior, though no clear evidence for this speculation is at hand. Loss of denticles has occurred in larvae of diverse habits and families (Mertens, 1960) and even within the megophryines (*Megophrys*).

The tadpoles from Johore (BBM 7162) are intermediate between forms assigned in Table 1 to the subgenera *Leptobrachium* and *Leptolalax*. Character states shared with the latter are: 1) slender body form; 2) number of divided rows of denticles less than four on both lips; and 3) restriction of the glandular zone at the rear of the buccal roof. The first two states, while presumably derived within the megophryines, represent trends seen frequently in other groups of tadpoles and, in this case, may be instances of parallelism. The third state, since it also is a reduction, may be another parallelism. Significantly, the Johore tadpoles lack the one derived character that would clearly ally them with *Leptolalax*, namely, the numerous filamentous papillae at the center of the ventral velum. Instead, they share the primitive condition of the velum found in the subgenus *Leptobrachium*. The cylindrical, rather than flaplike, infralabial papillae and the absence of pustules down the center of the buccal floor differentiate them from both subgenera. The combination of pri-

mitive and derived, reduced states makes it difficult to place these larvae in relation to the others reviewed here.

Differences among the larvae described in this paper in buccal anatomy are not easily explained functionally as all of those studied seem to feed on the same kinds of material and on the same size range. However, a more thorough analysis of diets might suggest how the array of large palps and papillae are involved in processing food.

ACKNOWLEDGMENTS

I am grateful to Richard Wassersug for advice and criticism, to Doyle L. Damman for several series of larvae from Thailand, and to Carol Small-Kaplan for help with the SEM micrographs.

LITERATURE CITED

- DUBOIS, A. 1980. Notes sur la systématique et la répartition des amphibiens anoures de Chine et des régions voisantes. IV. Classification generique et subgenerique des Pelobatidae Megophryinae. Bull. Mens. Soc. Linn. Lyon, **49**: 469-482.
- . 1981. Notes sur la systématique et la répartition des amphibiens anoures de Chine et des régions avoisinantes. V. *Megophrys oshanensis* Liu, 1950 et *Leptobrachium minimum* Taylor, 1962. Bull. Mens. Soc. Linn. Lyon, **51**: 183-192.
- HENRICKSON, J. R. 1966. Observations on the fauna of Pulau Tioman and Pulau Tulai. The Amphibians. Bull. Nat. Mus. Singapore, **34**: 72-84.
- INGER, R. F. 1966. The systematics and zoogeography of the Amphibia of Borneo. Fieldiana: Zool., **52**: 1-402.
- LIU, C. C. 1950. Amphibians of Western China. Fieldiana: Zool. Mem., **2**: 1-400.
- MERTENS, R. 1960. Die Larven der Amphibien und ihre evolutive Bedeutung. Zool. Anz., **164**: 337-358.
- POPE, C. H. 1931. Notes on amphibians from Fukien, Hainan, and other parts of China. Bull. Amer. Mus. Nat. Hist., **61**: 397-611.
- SMITH, M. A. 1917. On tadpoles from Siam. Jour. Nat. Hist. Soc. Siam, **2**: 261-275.
- TAYLOR, E. H. 1962. The amphibian fauna of Thailand. Univ. Kansas Sci. Bull. **43**: 265-599.
- WASSERSUG, R. 1980. Internal oral features of larvae of eight anuran families: functional, system-

atic, evolutionary and ecological considerations. Univ. Kansas Mus. Nat. Hist. Misc. Publ. No. 68, 146 pp.

WASSERBURG, R., K. J. FROGNER, AND R. F. INGER.

1981. Adaptations for life in tree holes by rhacophorid tadpoles from Thailand. *J. Herpetol.*, **15**: 41-52.

Frogs of the *Hyla columbiana* Group: Taxonomy and Phylogenetic Relationships

WILLIAM E. DUELLMAN¹

LINDA TRUEB²

ABSTRACT. Three species comprise the *Hyla columbiana* group—*H. carnifex* on the Pacific slopes of Ecuador and the northern part of the Cordillera Central in Colombia, *H. columbiana* in the upper Río Cauca Valley in Colombia, and a new species, *H. praestans*, in the upper Río Magdalena Valley in Colombia. *Hyla variabilis* Boulenger, 1896, is placed in the synonymy of *H. columbiana* Boettger, 1892. The three species are readily distinguished by color pattern, morphometric features, and mating calls. The *Hyla columbiana* group belongs to a complex of Neotropical hylids characterized by 30 chromosomes and reduced larval mouthparts.

INTRODUCTION

Among the approximately 385 genera of Neotropical amphibians and reptiles, three—*Anolis*, *Eleutherodactylus*, and *Hyla*—contain more than 800 recognized species. These three megagenera represent a significant component of the entire Neotropical herpetofauna. The largest of these is the leptodactylid frog genus *Eleutherodactylus*, which contains approximately 360 species—115 in the West Indies, 70 in Middle America, and 175 in South America. The most widely distributed is the hylid frog genus *Hyla*; although most of the 230 species are Neotropical, including six in the West Indies, there are also 20 species in the Holarctic Realm. The 230 species of the

iguanid lizard genus *Anolis* are divided about equally between the continental Neotropics and the West Indies; only one species is native to North America.

While we have attempted to sort out relationships among the hylid frogs during the preceding two decades, John D. Lynch has applied himself to *Eleutherodactylus*. Our current understanding of the taxonomy, relationships, and biogeography of the complex genus *Anolis* is largely the result of the assiduous efforts of Ernest E. Williams. In recognition of his devotion to the study of one of these megagenera and the frustrations associated with such endeavors, we dedicate this modest effort and a new species of *Hyla* to Ernest E. Williams.

Throughout the tropical lowlands of Middle and South America there are many species of small, yellowish-tan or brown *Hyla* that have pond-dwelling tadpoles with xiphicercal tails and reduced mouthparts. All of these species that have been studied karyologically have 30 chromosomes. There are few species of *Hyla* in the Andes, but some of these (*H. carnifex*, *H. columbiana*, and *H. variabilis*) resemble, superficially at least, the lowland taxa.

In 1975 we discovered a previously unknown species of small *Hyla* presumably related to *H. columbiana*. This find prompted our investigation of the taxonomy of the montane species and the definition of a *Hyla columbiana* group. In this paper we present the results of

^{1,2}Museum of Natural History and Department of Systematics and Ecology, University of Kansas, Lawrence, Kansas 66045, U.S.A.

these studies and propose a phylogenetic arrangement of the group and its relatives.

MATERIALS AND METHODS

Systematic aspects of this study are based on the examination of 591 preserved specimens, 14 skeletons, 12 lots of tadpoles, and 1 clutch of eggs belonging to species in the *Hyla columbiana* group. Recordings were made on a Uher-4000 tape recorder and analyzed on a Vibralyzer (Kay Electric Company). All measurements of morphological features and calls were taken as described by Duellman (1970), and webbing formulae were ascertained following Savage and Heyer (1967). Tadpoles were staged according to Gosner's (1960) tables. Statistical analyses of morphometrics were based on measurements of uniformly preserved specimens from the vicinities of the type localities. Pairwise comparisons of measurements and proportions were accomplished by one-way analyses of variance (Sokal and Rohlf, 1969). Results of $F_{\max}$ tests indicated normal distribution of the data. In a multiple comparison of the means, the Student-Newman-Keuls test was used to determine levels of significance.

All specimens are referred to by the following abbreviations: AMNH = American Museum of Natural History; ANSP = Academy of Natural Sciences, Philadelphia; BMNH = British Museum (Natural History); FMNH = Field Museum of Natural History; INDERENA = Instituto Nacional de Recursos Naturales, Bogota; KU = Museum of Natural History, University of Kansas; LACM = Museum of Natural History, Los Angeles County; MCZ = Museum of Comparative Zoology, Harvard University; NHMW = Naturhistorisches Museum, Wien; NHRM = Naturhistoriska Riksmuseet, Stockholm; SMF = Senckenberg Museum, Frankfurt-am-Main; UMMZ = Museum of Zoology, University of Michi-

gan; USNM = National Museum of Natural History; ZMB = Zoologisches Museum, Berlin; ZSM = Zoologisches Sammlung, München.

TAXONOMY

The *Hyla columbiana* Group

Definition. 1) Moderate sexual dimorphism in size; snout-vent lengths of males about 85% those of females; 2) snout short, blunt; 3) tympanum visible; tympanic annulus distinct or not; 4) hands and feet moderately webbed; 5) axillary membrane moderately extensive; 6) thoracic glands absent; 7) calcars and tarsal tubercles absent; ulnar tubercles present or absent; 8) males having single, median, subgular vocal sacs and no nuptial excrescences; 9) dorsum tan or brown, with or without dark dorsal markings and pale dorsolateral stripes; 10) pale labial stripe or spots present or not; 11) axilla and groin marked with yellow, orange, red or bluish gray in life; 12) anal, ulnar, and tarsal stripes absent; 13) iris bronze to reddish copper; 14) quadratojugal absent; 15) prevomers not articulating with maxillary arch, bearing small, posteromedially inclined dentigerous processes; 16) crista parotica poorly to moderately ossified; 17) zygomatic ramus of squamosal extending about one-third distance to maxillary arch; 18) pterygoid poorly developed, anterior ramus articulating weakly with unexpanded pars palatina of maxillary at level of midorbit; medial ramus widely separated from otic region of neurocranium; 19) sacral diapophyses moderately expanded (45–60°), anterior edge forming acute angle with longitudinal axis of body; 20) ilia lacking obvious crests; preacetabular angle obtuse; 21) tadpoles having ovoid bodies and xiphicercal tails with moderately deep fins barely extending onto body; 22) larval mouths terminal with two rows of small papillae ventrally, robust serrate beaks, and one upper and two lower rows

of denticles; 23) mating call consisting of short, moderately pitched notes, followed or not by shorter secondary notes; 24) diploid chromosome number 30.

Content. Three species, which may be identified by the following key:

1. Broad pale dorsolateral stripe present; groin bluish gray mottled with black in life *H. praestans*
 – Pale dorsolateral stripe absent; groin yellow, orange, or red in life 2
2. Small yellow spots on upper lip; venter bright yellow in life *H. carnifex*
 – No yellow spots on upper lip; venter creamy yellow in life *H. columbiana*

Distribution. Moderate elevations (975–2,580 m) on the Pacific slopes of the Cordillera Occidental in Ecuador and in the upper Río Cauca and Magdalena valleys and the northern part of the Cordillera Central in Colombia (Fig. 1).

Comment. The chromosome number is known only for *Hyla carnifex* (KU 173114), and tadpoles are known only for *H. carnifex* and *H. columbiana*. The populations from the northern part of the Cordillera Central, Colombia, are tentatively assigned to *H. carnifex*.

Following is an account of each of the species in the *Hyla columbiana* group. The results of the analyses of the morphometric data are presented in Table 1. For illustrations of the adult and larva of *H. carnifex*, see Duellman (1969).

Hyla carnifex Duellman

Hyla carnifex Duellman, 1969: 242 (Holotype.—KU 117993 from Tandapi, 1,460 m, Provincia de Pichincha, Ecuador); 1974: 7; 1977: 43.

Hyla bogerti Cochran and Goin, 1970: 261 (Holotype.—USNM 118731 from Medellín, Departamento de Antioquia, Colombia). Synonymy *fide* Duellman (1974: 7).

Hyla charlesbogerti Goin, 1970: 788 (Substitute name for *Hyla bogerti* Cochran and Goin, 1970). Synonymy *fide* Duellman (1974: 7).

Diagnosis. 1) Size small, snout-vent length in males 24.6 to 27.7 mm, in females 29.2 to 32.5 mm; 2) head short, 28.5 to 31.3% of snout-vent length; 3)

Figure 1. Western Colombia and Ecuador showing distributions of *Hyla carnifex* (●), *H. columbiana* (■), and *H. praestans* (▲). The dashed line is the 1,000-m contour; shaded areas are above 3,000 m.

snout round in profile; 4) interorbital distance narrow, 29.5 to 35.4% of head width; 5) skin on dorsum finely shagreened, nearly smooth; 6) ulnar tubercles distinct; 7) distal subarticular tubercle on fourth finger strongly bifid; 8) subanal tubercles

TABLE 1. STATISTICAL COMPARISON OF MORPHOMETRIC CHARACTERS OF TOPOTYPIC MALES OF THREE SPECIES IN THE *HYLA COLUMBIANA* GROUP.

First line = mean $\pm$ 1 standard deviation; second line = observed range. Means of all characters of *H. columbiana* significantly different ($P < 0.05$) from those of the other species; significant differences ($P < 0.05$) between *H. carnifex* and *H. praestans* are indicated by an asterisk*.

Character	<i>H. carnifex</i>		<i>H. columbiana</i>		<i>H. praestans</i>	
N	20		20		8	
Snout-vent length (SVL)	26.1 $\pm$ 0.90*		27.5 $\pm$ 1.14		28.8 $\pm$ 1.76*	
	24.6 – 27.7		25.8 – 29.3		26.0 – 31.5	
Tibia/SVL	0.455 $\pm$ 0.015		0.482 $\pm$ 0.017		0.455 $\pm$ 0.016	
	0.433 – 0.481		0.454 – 0.515		0.424 $\pm$ 0.471	
Foot/SVL	0.424 $\pm$ 0.018		0.456 $\pm$ 0.014		0.424 $\pm$ 0.026	
	0.395 – 0.455		0.430 – 0.479		0.372 – 0.452	
Head length/SVL	0.199 $\pm$ 0.008		0.339 $\pm$ 0.013		0.346 $\pm$ 0.009	
	0.285 – 0.313		0.314 – 0.356		0.334 – 0.360	
Head width (HW)/SVL	0.327 $\pm$ 0.010		0.312 $\pm$ 0.010		0.331 $\pm$ 0.006	
	0.310 – 0.349		0.290 – 0.332		0.324 – 0.340	
Interorbital/HW	0.332 $\pm$ 0.015		0.416 $\pm$ 0.032		0.319 $\pm$ 0.021	
	0.195 – 0.354		0.358 – 0.471		0.300 – 0.347	
Total prevomerine teeth	9.1 $\pm$ 1.17*		7.3 $\pm$ 1.13		8.6 $\pm$ 1.51*	
	7 – 11		6 – 10		6 – 10	

large; 9) dorsum tan with brown markings and usually with creamy yellow suborbital spot; 10) venter and vocal sac bright yellow, with or without brown spots on belly; 11) groin deep yellow orange; 12) mating call consisting of primary note followed by two or three secondary notes.

The presence of small, yellow spots on the upper lip and the bright yellow venter immediately distinguishes *H. carnifex* from other members of the group. Members of the *Hyla parviceps* group have larger suborbital spots and have white or gray venters and either orange spots on the ventral surfaces of the shanks or yellow spots on the anterior or dorsal surfaces of the thighs. The dorsal color pattern of *H. carnifex* resembles that of *H. minuta*, a species with a creamy yellow venter and white anal and heel stripes.

Description. Body robust; head small, noticeably narrower than body; snout short, its length about equal to diameter of eye, round in dorsal view and in profile; nostril about four-fifths distance from eye to tip of snout; canthus rostralis rounded, barely evident; loreal region slightly concave; lips moderately thick, not flared; internarial area noticeably

depressed; interorbital area slightly convex, much wider than eyelid; supratympanic fold short, moderately weak, obscuring upper part of tympanum; tympanic annulus indistinct; tympanum separated from eye by distance slightly greater than diameter of tympanum.

Axillary membrane short, extending about one-fourth length of upper arm; forearm moderately slender, bearing row of distinct ulnar tubercles; fingers short, fourth distinctly longer than second; discs round, half again as wide as digits; fingers webbed basally; webbing absent between first two fingers; webbing formula for other fingers II2–3–III3–2IV; subarticular tubercles moderately large, low, subconical; distal tubercle on fourth finger strongly bifid; supernumerary tubercles small, subconical, numerous on basal segments of digits; palmar tubercle flat, bifid; prepollical tubercle flat, elliptical; nuptial excrescences absent. Hind limbs moderately short, slender; calcars, tarsal folds and tubercles absent; inner metatarsal tubercle low, flat, elliptical; outer metatarsal tubercle evident in some specimens; toes moderately long, third and fifth subequal in length; toes about three-fourths webbed; webbing formula

II—(1½–2) III—(1½–2) IIII—(1–2) IV(2–2—IV; subarticular tubercles small, round; supernumerary tubercles minute, present in single row on proximal segment of each digit.

Skin on dorsum weakly shagreened, nearly smooth; skin on throat, belly, and proximal posteroventral surfaces of thighs granular; skin on other surfaces smooth. Anal opening directed posteriorly at upper level of thighs; anal flap short; subanal tubercles large. Tongue elliptical, shallowly notched posteriorly, and barely free behind. Dentigerous processes short, posteromedially inclined, widely separated medially, between small ovoid choanae, bearing 3–4 to 5–6 teeth for totals of 7–11 ($\bar{x}$ = 9.2, 20 ♂♂) and 4–5 to 6–6 teeth for totals of 9–12 ($\bar{x}$ = 10.4, 15 ♀♀). Vocal slits short, extending from near midlateral base of tongue towards angles of jaws. Vocal sac large, single, median, subgular.

Coloration in preservative. Dorsum of head, body, and limbs dark brown or grayish brown with darker brown flecks; middorsal dark brown blotch extending from occiput or eyelids at least to sacral region in all females (30) and most males (83 of 100); venter cream with or without dark brown spots; variation in 100 males and 30 females—plain (68%, 27%), spots on posterior part of belly (24%, 50%), spots on belly and chest (8%, 13%), spots on belly, chest, and throat (0%, 10%) (Fig. 2). Flanks dark brown, usually enclosing cream lateral streak; groin mottled dark brown and cream; posterior thighs brown with longitudinal cream stripe; axilla, webbing, and spots on upper lip cream; variation in number of small spots on upper lip in 130 adults—none 6.9%, one 70.0%, two 17.7%, three 3.1%, four 2.3%.

Coloration in life. Dorsum of head and body pale creamy tan or grayish tan with median brown or grayish-brown blotch and numerous dark brown flecks; dorsal surfaces of limbs, exclusive of thighs, pale brown with or without faint darker brown transverse bars; upper lip brown, usually with small yellow spots, at least

one below eye; axilla, posterior flanks, and anterior and posterior surfaces of thighs deep yellow or orange bordered by dark brown or black; ventral surfaces of shanks, inner surfaces of feet, webbing, and ventral surfaces of upper arms deep yellow or orange (deepest colors only in larger individuals; webbing orange-red in some large females); throat, belly, and median part of flanks bright yellow with or without faint brown, lavender, or gray spots; iris gray, flecked with reddish bronze.

Tadpoles. The tadpoles were described and illustrated by Duellman (1969). In structure and coloration they are much like those of *H. columbiana* (described and figured herein), except that the upper beak has distinct lateral processes and white flecks are absent on the body and tail.

Mating call. The mating call of *Hyla carnifex* consists of a monophasic primary note followed by two or three shorter secondary notes—"Wraah-ack-ack." Analysis of recordings of two individuals shows that call groups are repeated at a rate of about 14 calls per minute. Primary notes have durations of 0.20 to 0.36 sec. The fundamental frequency is at about 500 Hz, and the dominant frequency is the fifth or sixth harmonic at about 2,500 or 3,000 Hz (Fig. 3).

Distribution. This species is definitely known from the Pacific slopes of the Cordillera Occidental in Ecuador, where it occurs at elevations of 1,140 to 2,150 m. Northern populations assigned to this species are known from elevations of 1,480 to 2,580 m in the northern part of the Cordillera Central and the eastern slopes of the Cordillera Occidental in Departamento de Antioquia, Colombia (Fig. 1).

Remarks. The status of the populations in northern Colombia named as *Hyla bogerti* by Cochran and Goin (1970) and recognized as *H. carnifex* by Duellman (1974) is questionable. Although many specimens are available from Departa-

Figure 2. Ventral color patterns. *Hyla carnifex*: A. KU 164301, ♂. B. KU 164238, ♀. *Hyla praestans*: C. KU 169578, ♂. D. KU 169575, ♂. *Hyla columbiana*: E. KU 169470, ♂. F. KU 169490, ♂. G. KU 181165, ♂. H. KU 169472, ♀.

mento de Antioquia, Colombia, most are faded and poorly preserved, thereby making detailed morphometric comparisons useless. We have seen only one living specimen from Antioquia, and it had the distinctive bright yellow venter and flank pattern of *H. carnifex* from Ecuador. Although a gap of nearly 700 km separates the Ecuadorian localities from

those in Colombia, we anticipate that the species will be discovered along the intervening Pacific slopes of the Cordillera Occidental in Colombia. Until series of fresh, well-preserved specimens and recordings of mating calls are available from localities in northern Colombia, we can say only that living frogs from the widely separated regions have the same

Figure 3. Audiospectrograms of mating calls of: A. *Hyla carnifex* (KU Tape 1237, 19°C). B. *H. columbiana* (KU Tape 1337, 17°C). C. *H. praestans* (KU Tape 1348, 17.5°C). Narrow band (45 Hz) analysis.

distinctive coloration; therefore, we assign them to the same species.

Only a few specimens of *H. carnifex* have been found by day—under the bark of logs, in bromeliads, and in axils of

leaves of elephant-ear plants (*Xanthosoma*). Calling males have been found between January and September. Males call from low (<50 cm), emergent herbs and grasses in marshy, temporary pools

in clearings in cloud forest. Tadpoles have been found in these pools from April to July, and metamorphosing young have been collected from May until August. Five young with tail stubs of <2 mm from Tandapi, Ecuador, collected on 18 July 1970, have snout-vent lengths of 12.7 to 14.7 ($\bar{x}$ = 13.8) mm. Three young collected on 3 April 1975, 5 km east-southeast of Chiriboga, Ecuador, have snout-vent lengths of 15.0 to 15.5 ($\bar{x}$ = 15.3) mm, and four obtained on 11 May 1975 at the same locality have snout-vent lengths of 16.1 to 18.2 ($\bar{x}$ = 16.9) mm. In recently metamorphosed young, the dorsum is tan with a bronze tint; the anterior and posterior surfaces of the thighs and ventral surfaces of the shanks are dark gray, and the throat and belly are grayish white. In half-grown individuals (20–22 mm) the belly is white with a faint tint of yellow on the chin and chest; the flanks and anterior and posterior surfaces of the thighs are dark brown. A middorsal dark blotch is evident in these individuals, but the bright yellow or orange colors on the limbs and in the axilla and groin, and the spots on the upper lip do not develop until the frogs attain sexual maturity (snout-vent lengths of >23 mm in males and >27.5 mm in females).

Hyla columbiana Boettger

Hyla columbiana Boettger, 1892: 41 (Lectotype.—SMF 2365 [designated by Mertens, 1967: 4] from Popayán, Departamento de Cauca, Colombia); Nieden, 1923: 261; Cochran and Goin, 1970: 264; Duellman, 1977: 48.

Hyla variabilis Boulenger, 1896: 20 (Syntypes.—BMNH 1947.2.13.15–21; FMNH 3565; MCZ 2606; UMMZ 46464, 51269, 58908 [8 specimens]; USNM 71115 from Cali, Departamento de Valle, Colombia); Nieden, 1923: 262; Cochran and Goin, 1970: 258; Duellman, 1977: 108. NEW SYNONYMY.

Diagnosis. 1) Size small, snout-vent length in males 25.8 to 29.3 mm, in females 30.6 to 35.4 mm; 2) head moderately long, 31.4 to 35.6% of snout-vent length; 3) snout round in profile; 4) interorbital distance wide, 35.8 to 47.1% of

head width; 5) skin on dorsum smooth with scattered, small tubercles; 6) ulnar tubercles low, indistinct; 7) distal subarticular tubercle on fourth finger weakly bifid; 8) subanal tubercles small; 9) dorsum tan with brown markings; 10) venter creamy yellow with variable amount of brown or black markings (flecks to reticulations); vocal sac bright yellow in life; 11) groin yellow-orange to tomato-red in life; 12) mating call consisting of groups of 2 to 6 notes of equal duration.

The combination of small tubercles on the dorsum and absence of distinct labial spots and dorsolateral stripes distinguishes *H. columbiana* from other members of the group. The deep yellow, orange, or red axilla and groin and the extensive flecking or reticulations on the venter distinguish *H. columbiana* from all other species of small Neotropical *Hyla*.

Description. Body moderately robust; head as wide as body; snout moderately long, half again length of eye, round in dorsal view and in profile; nostril about two-thirds distance from eye to tip of snout; canthus rostralis acutely rounded; loreal region barely concave; lips thick, not flared; internarial area slightly depressed; interorbital area flat, nearly twice width of eyelid; supratympanic fold moderately weak, curving posteriorly, obscuring upper edge of tympanum; tympanic annulus distinct; tympanum separated from eye by distance only slightly greater than diameter of tympanum.

Axillary membrane moderately long, extending about two-fifths length of upper arm; forearm moderately robust; ulnar tubercles low, indistinct; fingers short, fourth slightly longer than second; discs round, slightly wider than digit; fingers webbed basally; webbing absent between first two fingers; webbing formula for other fingers II2[−]—3—III2½—2IV; subarticular tubercles small, subconical; distal tubercle on fourth finger weakly bifid; supernumerary tubercles small, subconical, numerous on basal segments; palmar tubercle low, flat,

weakly bifid; prepollical tubercle flat, elliptical; nuptial excrescences absent. Hind limbs moderately long, slender; calcars, tarsal folds and tubercles absent; inner metatarsal tubercle flat, narrowly elliptical; outer metatarsal tubercle absent; toes moderately long, third and fifth subequal in length; toes about three-fourths webbed; webbing formula $I(1-1\frac{1}{2})-2III1\frac{1}{2}-(2-2+III(1+-1\frac{1}{2})-2IV2-(1-1+)V$; subarticular tubercles small, subconical; supernumerary tubercles minute, subconical, numerous on proximal segments of each digit.

Skin on dorsal surfaces of head, body, and limbs smooth with scattered, small tubercles; skin on throat, belly, and proximal posteroventral surfaces of thighs coarsely granular. Anal opening directed posteriorly at upper level of thighs; anal flap short; subanal tubercles small. Tongue ovoid, barely notched behind, only slightly free posteriorly. Dentigerous processes of prevomers small, posteromedially inclined, moderately separated medially, between posterior borders of small, ovoid choanae, bearing 3-3 to 5-5 teeth for totals of 6-10 ($\bar{x} = 7.3$, 20 ♂♂) and 4-5 to 6-7 teeth for totals of 9-13 ($\bar{x} = 11.0$, 3 ♀♀). Vocal slits long, extending from midlateral base of tongue to angles of jaws. Vocal sac large, single, median, subgular.

Coloration in preservative. Dorsal surfaces of head, body, and limbs tan to brown with three types of patterns of darker brown markings in 40 recently preserved specimens: 1) middorsal blotch extending from eyelids to middle of dorsum and a smaller blotch usually evident in sacral region (50%); 2) narrow, diffuse stripe middorsally (25%); and 3) irregular flecks and/or reticulations (25%). Dorsal surfaces of limbs plain, flecked, or barred (in some specimens having dorsal blotches). Pale canthal, labial, and dorsolateral stripes absent; faint dark canthal and postorbital stripes evident in some specimens. Venter creamy tan with a variable amount of flecking and reticulation in 40 specimens (Fig. 2): 1) no flecks (35%); 2) flecks and reticulations

only in groin (30%); 3) reticulations in groin; flecks on belly, especially posteriorly (20%); and 4) reticulations in groin and on belly (15%). All females and some males in last category; females also having reticulations on throat. Anterior and posterior surfaces of thighs brown or tan with or without irregular brown markings; webbing and discs brown.

Coloration in life. At night dorsum yellowish tan; axilla, groin, proximal anterior surfaces of thighs, and inner surfaces of shanks pink to orange or yellow. By day, dorsum yellow, bronze, tan, or brown, with or without distinct brown blotches; venter yellow, with or without dark brown flecks and/or reticulations on belly; axilla, groin, proximal anterior surfaces of thighs, and inner surfaces of shanks deep yellow, orange, or red. Iris reddish copper. In some individuals, a faint dark canthal stripe and a dark brown postorbital stripe extending onto the anterior flanks are present; also, in some individuals a diffuse creamy yellow suborbital bar is present (Fig. 4).

Tadpoles. Among a series of tadpoles (KU 170202) in developmental stages 25 through 41, a typical tadpole in stage 38 is described. Body length 13.2 mm; total length 35.5 mm; body as wide as deep, sides nearly parallel for most of the length of body posterior to eyes, deepest posteriorly; snout in dorsal view broadly rounded, in profile more acutely rounded; eyes large, directed laterally; nostril about midway between eye and tip of snout, directed anteriorly; spiracle sinistral, directed posterodorsally just below midline at point about two-thirds length of body; anal opening dextral. Caudal musculature moderately robust anteriorly, tapering to attenuated point posteriorly; fins deepest at about mid-length of tail, tapering abruptly to xiphocercal tail; depth of either fin at mid-length of tail about equal to depth of caudal musculature; dorsal fin barely extending onto body (Fig. 5).

Mouth small (about one-fourth greatest width of body), terminal, directed anteroventrally; upper lip bare; two rows of

Figure 4. *Hyla columbiana*, KU 169463, ♂, 28.1 mm snout-vent length.

Figure 5. Tadpole of *Hyla columbiana*, KU 170202, 35.5 mm total length.

small, pointed papillae laterally and ventrally; lateral labial folds absent; beaks moderately massive, bearing fine serrations; upper beak broadly arch-shaped with no narrowed, lateral processes; lower beak broadly V-shaped; one upper and two lower complete rows of denticles; upper row just inside upper lip; second lower row having smallest denticles.

In preservative, dorsum of body and snout dull brown; sides and venter transparent with bluish sheen and numerous white flecks; caudal musculature orangetan; caudal fins translucent; caudal musculature and fins bearing minute white flecks; in larger specimens (stage 30 and beyond), brown blotches on caudal fins, becoming larger and darker in larger tadpoles. In life, dorsum olive-

green with broad, tan dorsolateral stripe; tail cream with minute brown flecks in small individuals and brown spots on fins in larger individuals; venter silvery white; iris pale bronze.

Mating call. The mating call of *Hyla columbiana* consists of short series of similar notes. Analysis of recordings of five individuals reveals that each call consists of groups of 2 to 6 ($\bar{x}$ = 3.6) notes repeated at a rate of 12 to 16 ($\bar{x}$ = 14.4) notes per minute. Each note has a duration of 0.1 to 0.3 ($\bar{x}$ = 0.16) sec. The fundamental frequency is at 240 to 320 ($\bar{x}$ = 265) Hz. The eighth or ninth harmonic is emphasized, giving a dominant frequency of 2,000 to 2,560 ($\bar{x}$ = 2,480) Hz (Fig. 3).

Remarks. Examination of the type specimens of *Hyla columbiana* and *H. variabilis* and study of living and preserved specimens from the regions of the type localities of the two named taxa, as well as analyses of mating calls, reveal the presence of only one species in the upper Río Cauca Valley. Cochran and Goin (1970: 260, 266) noted colors of a few living specimens from Cali and Popayán; they emphasized that the axilla was red in specimens from Popayán and yellow in specimens from Cali. In a series of 38 adults collected 8 km east of Popayán, the color of the axilla varied from deep yellow to orange and tomato-red, in living frogs from Lago de Calima (north of Cali) the axilla was pink.

Notable ontogenetic, sexual, and geographic variation exists in the amount of dark pigmentation on the venter. Juveniles lack dark pigment ventrally, whereas large adult females of all populations have reticulate venters, a condition also seen in many males from Cali and a few males from Popayán (Fig. 2).

We have observed this species only in disturbed areas that formerly supported cloud forest. Three individuals were found in *Agave* plants by day. All others were in shallow, grassy marshes or pools at night. Males call from clumps of grass in water or from grasses within 5 cm of

the surface. On 20 September 1974 we found a single clutch of eggs attached to grass on the surface of the water in a temporary pool. Tadpoles were found in the same pool.

Five recently metamorphosed young (KU 170201) have snout-vent lengths of 13.7 to 18.6 ($\bar{x}$ = 16.1) mm. In life, the dorsum was bronze tan; the anterior and posterior surfaces of the thighs were dark brown. The upper arms and suborbital bar were creamy yellow, and the throat and belly were white. The iris was reddish bronze. In preservative, the juveniles are pale tan dorsally and creamy white ventrally; there is no indication of flecks or reticulations on the belly or in the groin.

Distribution. *Hyla columbiana* is widespread in the upper Río Cauca Valley, Colombia, where it occurs on the valley floor at elevations of 975 to 1,000 m and on the lower eastern slopes of the Cordillera Occidental and western slopes of the Cordillera Central to elevations of 2,250 m. It also occurs in the upper drainage of the Río Patía, where it is known from elevations of 1,800 to 2,350 m (Fig. 1).

Hyla praestans new species

Holotype. KU 169575, an adult male, from the Parque Arqueológico San Agustín, 3 km southwest of the village of San Agustín, Departamento de Huila, Colombia (1°53'N, 76°16'W), 1,750 m, one of a series obtained on 27 May 1975 by William E. Duellman, John E. Simmons, and Linda Trueb.

Paratypes. KU 169574, 169576–80, MCZ 100216, INDERENA (2 specimens) collected with the holotype.

Referred Specimens. ICN 7556–58 from 7.4 km (by road) NW of San José de Isnos, Departamento de Huila, Colombia, 1,970 m (John D. Lynch, personal communication).

Diagnosis. 1) Size small, snout-vent length in males 26.0 to 31.5 mm, females 31.2 mm; 2) head moderately long, 33.4 to 36.0% of snout-vent length; 3) snout in profile truncate, rounded above; 4) interorbital distance narrow, 30.0 to 34.7% of head width; 5) skin on dorsum finely

shagreened, nearly smooth; 6) ulnar tubercles absent; 7) distal subarticular tubercle on fourth finger strongly bifid; 8) subanal tubercles small; 9) dorsum tan with creamy yellow labial stripe and orange-tan dorsolateral stripe; 10) venter and vocal sac creamy white with brown flecks on belly; 11) groin pale bluish gray mottled with black; 12) mating call consisting of primary note followed by one or two secondary notes.

Hyla praestans differs from the other members of the *Hyla columbiana* group by having pale labial and dorsolateral stripes; the latter is characteristic of females of many species in the *Hyla parviceps* group, all of which have yellow spots on the thighs or orange spots on the ventral surfaces of the shanks. Another distinguishing feature of *H. praestans* is the bluish-gray groin with black mottling, a pattern occurring in some populations of *H. labialis*, a much larger frog that is uniformly dark green dorsally.

Description. Body robust; head slightly narrower than body; snout moderately short, only slightly longer than diameter of eye, round in dorsal view, truncate and rounded above in profile; nostril about two-thirds distance from eye to tip of snout; canthus rostralis rounded; loreal region slightly concave; lips moderately thick, not flared; internarial area slightly depressed; interorbital area flat, much wider than eyelid; supratympanic fold short, weak, obscuring upper part of tympanum; tympanic annulus distinct; tympanum separated from eye by distance one third greater than diameter of tympanum.

Axillary membrane short, extending about one-fourth length of upper arm; forearm moderately slender, lacking ulnar tubercles; fingers moderately long, fourth slightly longer than second; discs round, slightly wider than digit; fingers webbed basally; webbing absent between first two fingers; webbing formula for other fingers $II2^{+}-(3^{-}-3^{+})III3-2\frac{1}{2}IV$; subarticular tubercles moderately

large, subconical; distal tubercle on fourth finger strongly bifid; supernumerary tubercles low, flat; palmar tubercle low, flat, ovoid; prepollical tubercle flat, elliptical; nuptial excrescences absent. Hind limbs moderately long, slender; calcars, tarsal folds and tubercles absent; inner metatarsal tubercle small, flat, elliptical; outer metatarsal tubercle absent; toes moderately long, fifth barely longer than third; toes about two-thirds webbed; webbing formula $I2^{+}-2III1-(2^{-}-2)III(1-1\frac{1}{2})-2^{-}-2)IV(2^{-}-2\frac{1}{2})-(1-1\frac{1}{2})V$; subarticular tubercles small, subconical; supernumerary tubercles absent.

Skin on dorsum weakly shagreened, nearly smooth; skin on throat, belly, and proximal posteroventral surfaces of thighs weakly granular; skin on other surfaces smooth. Anal opening directed posteriorly at upper level of thighs; anal flap short; subanal tubercles small. Tongue elliptical, shallowly notched posteriorly, barely free behind. Dentigerous processes of prevomers small, posteromedially inclined, narrowly separated medially, between small, ovoid choanae, bearing 3-3 to 5-5 teeth for totals of 6-10 ($\bar{x} = 8.6$) teeth in males. Vocal slits long, extending from midlateral base of tongue to angles of jaws. Vocal sac large, single, median, subgular.

Coloration in preservative. Dorsal surfaces of head, body, and limbs brown with darker brown flecks on body and limbs and dark brown longitudinal streaks of variable size and distinctness on body. Narrow, cream to orange-tan canthal line from tip of snout to eye; broad, cream to orange-tan dorsolateral strip from eye to groin; flanks dark brown. Venter cream with variable amount of brown flecking or spotting on throat, belly, and thighs (Fig. 2); anterior and posterior surfaces of thighs creamy tan with dark brown mottling; webbing and discs brown.

Coloration in life. At night, dorsum tan with creamy yellow dorsolateral stripes. By day, dorsum and flanks dark brown with orange-tan dorsolateral stripes,

creamy tan canthal stripes, and creamy yellow labial stripe; groin mottled bluish gray and black; posterior surfaces of thighs dark brown; throat and belly creamy white with brown flecks; ventral surfaces of limbs tan; iris coppery brown (Fig. 6).

Measurements of holotype (in mm). Snout-vent length 28.3, tibia length 13.3, foot length 12.6, head length 9.8, head width 9.3, eye-nostril 2.5, interorbital distance 3.2, eyelid width 2.1, horizontal diameter of eye 2.8, horizontal diameter of tympanum 1.3.

Tadpoles. Unknown.

Mating call. The call of *Hyla praestans* consists of a primary note followed by one or two, shorter, secondary notes. Analysis of a single recording shows that the call groups are repeated at a rate of 4.5 calls per minute. The duration of the

primary note is 0.08 sec. The fundamental frequency is at 450 Hz, and the fifth harmonic at 2,250 Hz is dominant (Fig. 3).

Distribution. This species is known from elevations of 1,750 to 1,970 m on the lower eastern slopes of the Cordillera Central next to the upper Río Magdalena Valley, Colombia (Fig. 1).

Remarks. The region of the type locality is humid lower montane forest with bamboo, *Heliconia*, and some bromeliads. The exact type locality is a small, temporary pond immediately behind (± 100 m) the Museo Arqueológico San Agustín. Males were calling from a tangle of brambles over (up to 1 m above) the water at night following a light rain (13 mm).

Etymology. The specific name, *praestans*, is Latin meaning preeminent or

Figure 6. *Hyla praestans*, KU 169574, ♂, 27.3 mm snout-vent length.

distinguished and is proposed in recognition of Ernest E. Williams's recognized preeminence in the field of herpetology.

PHYLOGENETIC RELATIONSHIPS

The phylogenetic relationships of groups of taxa may be inferred by determining shared-derived characters in defining monophyletic taxa; determination of sequential sets of shared-derived character states provides a stepwise association of character states defining branching sequences in a phylogenetic arrangement and classification reflecting these phylogenetic relationships. This approach has become known as phylogenetic systematics (see Wiley, 1980, for terms and methodology).

Although we can provide a reasonable hypothesis of the phylogenetic relationships of the *Hyla columbiana* group with other groups of *Hyla*, the distribution of character states among the species within the group and the absence of meaningful interpretation of evolutionary trends of many characters, such as skin texture and coloration, defies a reconstruction of the phylogeny of the three species. Using 10 of the characters listed in the diagnoses (all except snout-vent length and proportion of head width) plus six cranial characters (nature of frontoparietal fontanelle, degree of ossification of sphenethmoid, shape of parasphenoid alae, serration of coronoid, and amount of separation of prevomers and of palatines), we found that there was little concordance in shared character states. *Hyla carnifex* shares six states with *H. columbiana* and seven with *H. praestans*, which shares only three with *H. columbiana*. Thus, even a phenetic analysis reveals only that *H. columbiana* has more unique character states than either of the other two species.

Various apparently related groups of Neotropical hylid frogs have been defined as a matter of convenience by Bokermann (1964), Duellman (1970, 1974), and Duellman and Crump (1974).

Informally recognized assemblages of species include the *leucophyllata* (6 species), *marmorata* (4), *microcephala* (± 12), *minuta* (2), and *parviceps* (7) groups. For the purposes of the following discussion, we add *H. labialis* and the *columbiana* group.

In attempting to ascertain the phylogenetic relationships among these groups, we utilized eight character states; 2 from adults, 5 from larvae and the karyotype. These states were selected because they are ones in which we place confidence in our knowledge of phylogenetic trends within the Hyliidae. Other characters are highly variable within, as well as between, groups, or are restricted to members of only one group; therefore, they are useless in determining inter-group relationships. The characters, their states, and direction of evolutionary change (0 = primitive state; 1 = derived state; 2 = secondarily or independently derived state) are as follows:

A. *Chromosome number*. Hyliid frogs have diploid numbers of 22, 24, 26, 28, or 30 chromosomes. Bogart (1973) showed that the primitive number of chromosomes in leptodactylids, the presumed ancestors of hyliids, is 26—a number occurring in phyllomedusine, some pelodryadine, and some amphignathodontine hyliids. Most hyline frogs have 24 chromosomes, but all species so far examined of the groups of *Hyla* listed above have 30 chromosomes, a number unknown in other groups of hyliids. Bogart (1973) and Morescalchi (1973) suggested that the evolution of 30 chromosomes occurred by centric fission, and Bogart (1973) noted variable numbers of pairs of telocentric chromosomes in those species having a diploid number of 30. Because this number is unique to the groups of *Hyla* under consideration, we assume that this chromosome number is a uniquely derived character state in these frogs. The coded character states and polarity are:

0 = <30 chromosomes; 1 = 30 chromosomes

0 $\rightarrow$ 1

B. *Quadratojugal arch*. Primitively in anurans, the quadratojugal forms a complete arch between the quadrate and the maxillary. This bone is reduced or absent in a number of primitive and advanced families (Lynch, 1973), including some leptodactylids (Lynch, 1971) and some hylids (Duellman, 1970). The presence of a complete quadratojugal arch is primitive (Lynch, 1973; Trueb, 1973). The coded character states and polarity are:

0 = arch complete; 1 = quadratojugal reduced or absent

0 → 1

C. *Nuptial excrescences*. Nuptial excrescences on the prepollex of breeding males usually are associated with species that amplex in water, and the absence of excrescences is characteristic of those that amplex on land or on vegetation (Parker, 1940; Lynch, 1971). Presence of excrescences is considered to be primitive. The coded character states and polarity are:

0 = excrescences present; 1 = excrescences absent

0 → 1

D. *Larval body shape*. The general configuration of the body of tadpoles is a reflection of the major habitats in which they develop. Tadpoles in torrential streams tend to have depressed bodies, whereas those in ponds generally have ovoid bodies. Modifications of pond tadpoles are associated with the level in which they feed in the water column; both surface-feeders and bottom-feeders are more depressed in comparison with midwater-feeders. In dorsal view, tadpoles of the *Hyla leucophyllata* group have a lateral constriction at midbody, resulting in a violin shape. The coded character states and polarity are:

0 = ovoid; 1 = violin; 2 = depressed

0 → 1; 0 → 2

E. *Larval tail shape*. The tadpoles of most frogs in primitive and advanced families have a tail that terminates in a rounded or acuminate tip; the caudal musculature terminates anterior to the margin of the fin. In some of the hylids under consideration, the caudal muscula-

ture extends posterior to the margin of the fins, which extend as a narrow fringe along the terminal part of the musculature. This type of tail with a distal filament is termed xiphicercal and is a derived feature. The coded character states and polarity are:

0 = acute; 1 = xiphicercal

0 → 1

F. *Larval mouth position*. The generalized tadpole mouth is directed anteroventrally. Stream-adapted tadpoles have ventral mouths, and surface-feeding tadpoles have mouths directed dorsally. Some midwater tadpoles have mouths directed anteriorly at the tip of the snout. The coded character states and polarity are:

0 = anteroventrally; 1 = anteriorly

0 → 1

G. *Larval labial papillae*. In pond-dwelling tadpoles of the Leptodactylidae and Hylidae, the mouth usually is bordered ventrally by two rows of small papillae. Within the groups under consideration, we find one or two rows of small papillae, one row of large papillae, or no papillae. The large papillae seem to represent a fusion of smaller papillae, and the absence of papillae seems to result from the fusion of large papillae into a dermal fold, as seen laterally in members of the *Hyla parviceps* group. The coded character states and polarity are:

0 = 2 small rows; 1 = 1 small row;

2 = 1 large row; 3 = none

0 → 1; 0 → 2 → 3

H. *Larval labial denticles*. The standard denticle formula in tadpoles is two upper rows and three lower rows (2/3). This number is increased in many stream-adapted tadpoles and to a lesser extent in some pond-dwelling tadpoles. Within the Hylidae reduction of the number of rows of denticles is uncommon in free swimming and feeding tadpoles and is most extreme in the groups considered here. The coded character states and polarity are:

0 = 2/3; 1 = 1/2; 2 = 0/1; 3 = 0/0

0 → 1 → 2 → 3

In order to ascertain the evolutionary

trends within the groups being considered, it is necessary to have an outgroup for comparison. Ideally such an outgroup should be the most closely related taxon, but such a determination is not presently possible because of our inadequate knowledge of many Neotropical hylids. Thus we have chosen to make the comparisons with *Smilisca*, a genus of six generalized species of Neotropical hylids (Duellman and Trueb, 1966). The character states for the taxa are given in Table 2, and the resulting phylogenetic arrangement is shown in Figure 7.

Thus, if our assumption that the presence of 30 chromosomes defines a monophyletic complex of hylid frogs is correct, then it is evident that the major evolutionary trends in this complex involve larval structure, especially mouthparts. Wassersug (1980) emphasized that larval buccal morphology in the *Hyla leucophyllata* group was specialized for macrophagy and that this specialization

was best developed in the *Hyla microcephala* group. The ecological and biogeographical significance of these modifications are not well understood. In those few species that have been studied (Crump, 1974; Duellman, 1978), the reproductive strategy is to deposit multiple clutches of small eggs in, or on, vegetation above, temporary ponds.

Throughout most of the lowlands in tropical America, three or four species of this complex occur in sympatry, but 11 species occur sympatrically in Amazonian Ecuador. Only two of the groups occur in the Andes. The three species of the *Hyla columbiana* group have allopatric distributions in the western and central cordilleras in Colombia and Ecuador, whereas *Hyla labialis* inhabits higher elevations (2,400–3,000 m) in the Cordillera Oriental in Colombia and the Mérida Andes in Venezuela. The diversity of species in this complex in the tropical lowlands may be the result of

Figure 7. Reconstruction of the phylogeny of groups of Neotropical *Hyla* with 30 chromosomes. Shifts in character states are indicated by the horizontal bars and identified by the letters and numbers given in Table 2 and text.

TABLE 2. CHARACTER STATES OF GROUPS OF *HYLA* HAVING 30 CHROMOSOMES.

Group	Characters							
	A	B	C	D	E	F	G	H
<i>H. labialis</i>	1	1	0	0	0	0	0	1
<i>H. columbiana</i>	1	1	1	0	1	0	0	1
<i>H. minuta</i>	1	1	1	0	1	1	0	2
<i>H. marmorata</i>	1	1	1	0	1	1	1	3
<i>H. parviceps</i>	1	1	1	0	1	1	2	3
<i>H. leucophyllata</i>	1	1	1	1	1	1	2	3
<i>H. microcephala</i>	1	1	1	2	1	1	3	3
Outgroup (<i>Smilisca</i>)	0	0	0	0	0	0	0	0

late Cenozoic and Quaternary climatic-ecological fluctuations in the tropical lowlands that resulted in the alternating fragmentation and reunification of forest and nonforest habitats (Haffer, 1979; Duellman, 1982a). On the other hand, the allopatric distributions of the few species in the Andes may reflect isolation of previously more widespread species with the uplift of the Andes or as a consequence of the vicissitudes of Pleistocene climatic change in the Andes (Simpson, 1975, 1979; Duellman, 1979, 1982b).

ACKNOWLEDGMENTS

We are indebted to the following persons for the loan of specimens or for the provision of working space in their respective institutions: Josef Eiselt, Alice G. C. Grandison, the late Walter Hellmich, Jorge Hernández C., W. Ronald Heyer, Konrad Klemmer, Edmond Malnate, Hymen Marx, Charles W. Myers, Günther Peters, the late James A. Peters, Greta Vestergren, the late Charles F. Walker, Ernest E. Williams, and John W. Wright. Our field work in 1974–75, which was supported by grants from the National Geographic Society (No. 1304) and the National Science Foundation (DEB 74-01998), profited from the able assistance of Dana T. Duellman and John E. Simmons. Collecting permits for Colombia were generously issued by Jorge Hernández C. of Inderena. We

are grateful to Guillermo Gutierrez, Director of the Parque Arqueológico San Agustín, for his gracious hospitality.

APPENDIX: SPECIMENS EXAMINED

Only specimens of the *Hyla columbiana* group are listed.

Hyla carnifex (427). COLOMBIA: Antioquia: Envigado, AMNH 39796, 39798–99; Finca San José de Bella Vista, LACM 47036–52, 47139; Jericó, 1,970 m, KU 98139–41, 98142 (skeleton), MCZ 15076–100, 24888–93, UMMZ 90601, USNM 118243; La Ceja, 2,180 m, FMNH 63883; Medellín, 1,480 m, AMNH 1357, ANSP 25005, FMNH 17098–101, MCZ 7664–65, UMMZ 56508 (38), 60251 (2), 92167, USNM 75989, 118729–31; Río Porce, ANSP 25773; San José de los Andes, LACM 72999; San Pedro de Osa, 2,450 m, AMNH 38713–14, 39153–77, ANSP 21029, FMNH 30567–69, 63884, 63897, MCZ 24911–15, UMMZ 71214, 71215 (2), 78301 (17), USNM 152021–29; Santa Rosa, 2,580 m, AMNH 39453, 39458, Yarumal, 2,265 m, USNM 152030–31. ECUADOR: Azuay: Cuenca (? locality), AMNH 23703, SMF 2368, ZMB 28023. Imabura: Apuela, 1550 m, KU 117992, 132428–41, 132544 (tadpoles). Pichincha: 5 km ESE Chiriboga, 2,010 m, KU 164302–05, 166189–92, 166311–14 (skeletons), 173113–15; 4 km W Chiriboga, 2,120 m, KU 142638–44; 14 km W Chiriboga, 1,960 m, KU 164306–07, 166193–95 (tadpoles); 4 km NE Dos Ríos, 1,140 m, KU 164283–301, 166188 (tadpoles); Hwy 28, 22 km from Hwy 30, 1,770 m, MCZ 91202; Miligal (? = Milligalli), USNM-GOV 8041–47; 3.5 km NE Mindo, 1,540 m, KU 164308–11; Tandapi, 1,460 m, AMNH 81399–403, BMNH 1969.640–44, FMNH 170746–50, KU 109557 (skeleton), 111838–66, 111867–70 (skeletons), 111871–90, 112360 (tadpoles), 117993–8001, 132442, 136199–254, 138775–76 (tadpoles), 178720–22, 180352 (tadpoles), MCZ 75062–66, UMMZ 129016 (8), USNM 166543–50; 6.3 km E Tandapi, 1,770 m, MCZ 94728; 9 km SE Tandayapa, 2,150 m, KU 164312.

Hyla columbiana (180). COLOMBIA: Cauca: El Tambo, 1,800 m, ANSP 25148, FMNH 54776, KU 145075–79, NHRM 2015 (11); road to Munchique, 2,350 m, KU 148493 (tadpoles); Popayán, 1,740 m, ANSP 25677, FMNH 43933–80, 43982–91, 44050, 44109, 82005–07, SMF 2365, USNM 152081–82, 152146–51; 8 km E Popayán, 2,110 m, Inderena (2), KU 169464–98, 170110–12 (skeletons), 170201–02 (tadpoles), 170203 (eggs); Quintana, FMNH 54719; 4 km S Silvia, 2,200 m, KU 169461–62; 6 km S Silvia, 2,250 m, KU 169463. Quindío: Boquía, 25 km N Armenia, 1,710 m, KU 133449, 139524. Valle: Cali, 1,000 m, AMNH 10674–77, BMNH 1947.2.13.15–21, FMNH 3565, MCZ 2606, NHMW 6198, 19438, SMF 2642–44, UMMZ 46464, 51269, 58908 (8), USNM 71115, ZSM 1182/0 (3); Candelaria, 975 m, USNM 151983–84; Cerro La Herrera, USNM 137770–72;

Finca Wesfalia, ZSM 315/1937; Lago de Calima, 1,350 m, KU 181165-66.

Hyla praestans (11). COLOMBIA: Huila: Parque Arqueológico San Agustín, 3 km SW San Agustín, 1,750 m, INDERENA (2), KU 169574-80, 170113 (skeleton), MCZ 100216.

LITERATURE CITED

- BOETTGER, O. 1892. Katalog der Batrachier-Sammlung im Museum Senckenbergischen Naturforschenden Gesellschaft in Frankfurt-am-Main. Senckenberg. Natur. Ges. Ber., 1892: 1-73.
- BOGART, J. P. 1973. Evolution of anuran karyotypes, pp. 337-349. In J. L. Vial (ed.), *Evolutionary biology of the anurans: Contemporary research on major problems*. Columbia, Univ. Missouri Press.
- BOKERMANN, W. C. A. 1964. Notes on tree frogs of the *Hyla marmorata* group with description of a new species (Amphibia, Hylidae). Senckenberg. Biol., 45: 243-254.
- BOULENGER, G. A. 1896. Descriptions of new reptiles and batrachians from Colombia. Ann. Mag. Nat. Hist., (6) 17: 16-21.
- COCHRAN, D. M., AND C. J. GOIN. 1970. Frogs of Colombia. U.S. Natl. Mus. Bull., 288: 1-655.
- CRUMP, M. L. 1974. Reproductive strategies in a tropical anuran community. Univ. Kansas Mus. Nat. Hist. Misc. Publ., 61: 1-68.
- DUELLMAN, W. E. 1969. A new species of frog in the *Hyla parviceps* group. Herpetologica, 25: 241-247.
- . 1970. The hylid frogs of Middle America. Univ. Kansas Mus. Nat. Hist. Monogr., 1: 1-753.
- . 1974. A reassessment of the taxonomic status of some Neotropical hylid frogs. Univ. Kansas Mus. Nat. Hist. Occ. Pap., 27: 1-27.
- . 1977. Liste der rezenten Amphibien und Reptilien, Hylidae, Centrolenidae, Pseudidae. Das Tierreich, 95: 1-225.
- . 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. Univ. Kansas Mus. Nat. Hist. Misc. Publ., 65: 1-352.
- . 1979. The herpetofauna of the Andes: Patterns of distribution, origin, differentiation, and present communities, pp. 371-459. In W. E. Duellman (ed.), *The South American herpetofauna: Its origin, evolution, and dispersal*. Univ. Kansas Mus. Nat. Hist. Monogr., 7: 1-485.
- . 1982a. Quaternary climatic-ecological fluctuations in the lowland tropics: Frogs and forests, pp. 389-402. In G. T. Prance (ed.), *Biological diversification in the tropics*. New York, Columbia Univ. Press.
- . 1982b. Compresión climática cuaternaria en los Andes: Efectos sobre la especiación. Actas VIII Congr. Latinoamer. Zool. (in press).
- DUELLMAN, W. E., AND M. L. CRUMP. 1974. Speciation in frogs of the *Hyla parviceps* group in the upper Amazon Basin. Univ. Kansas Mus. Nat. Hist. Occ. Pap., 23: 1-40.
- DUELLMAN, W. E., AND L. TRUEB. 1966. Neotropical hylid frogs, genus *Smilisca*. Univ. Kansas Mus. Nat. Hist. Publ., 17: 281-375.
- GOIN, C. J. 1970. A new name for *Hyla bogerti* Cochran and Goin. Copeia, (4)1970: 788.
- GOSNER, K. L. 1960. A simplified table for staging anuran embryos and larvae with notes on identification. Herpetologica, 16: 183-190.
- HAFFER, J. 1979. Quaternary biogeography of tropical lowland South America, pp. 107-140. In W. E. Duellman (ed.), *The South American herpetofauna: Its origin, evolution, and dispersal*. Univ. Kansas Mus. Nat. Hist. Monogr., 7: 1-485.
- LYNCH, J. D. 1971. Evolutionary relationships, osteology, and zoogeography of leptodactyloid frogs. Univ. Kansas Mus. Nat. Hist. Misc. Publ., 53: 1-238.
- . 1973. The transition from archaic to advanced frogs, pp. 133-182. In J. L. Vial (ed.), *Evolutionary biology of the anurans: Contemporary research on major problems*. Columbia, Univ. Missouri Press.
- MERTENS, R. 1967. Die herpetologische Sektion des Natur-Museums und Forschungs-Instituts Senckenberg in Frankfurt a. M. nebst einem Verzeichnis ihrer Typen. I. Senckenber. Biol., 48: 1-106.
- MORESCALCHI, A. 1973. Amphibia, pp. 233-248. In A. B. Chiarelli and E. Capanna (eds.), *Cytotaxonomy and vertebrate evolution*. New York, Academic Press.
- NIEDEN, F. 1923. Anura I. Subordo Aglossa und Phaneroglossa. Section 1, Arcifera. Das Tierreich, 46: 1-584.
- PARKER, H. W. 1940. The Australasian frogs of the family Leptodactylidae. Novit. Zool., 42: 1-106.
- SAVAGE, J. M., AND W. R. HEYER. 1967. Variation and distribution in the tree-frog genus *Phyllomedusa* in Costa Rica, Central America. Beitr. Neotrop. Fauna, 5: 111-131.
- SIMPSON, B. B. 1975. Pleistocene changes in the flora of the high tropical Andes. Paleobiology, 1: 273-294.
- . 1979. Quaternary biogeography of the high montane regions of South America, pp. 157-188. In W. E. Duellman (ed.), *The South American herpetofauna: Its origin, evolution, and dispersal*. Univ. Kansas Mus. Nat. Hist. Monogr., 7: 1-485.
- SOKAL, R. R., AND F. J. ROHLF. 1969. *Biometry*. San Francisco, W. H. Freeman Co.
- TRUEB, L. 1973. Bones, frogs, and evolution, pp. 65-132. In J. L. Vial (ed.), *Evolutionary biology of the anurans: Contemporary research on major problems*. Columbia, Univ. Missouri Press.
- WASSERSUG, R. 1980. Internal oral features of larvae from eight anuran families: Functional, sys-

tematic, evolutionary and ecological considerations. Univ. Kansas Mus. Nat. Hist. Misc. Publ., **68**: 1-146.

WILEY, E. O. 1980. Phylogenetic systematics and vicariance biogeography. Syst. Bot., **5**: 194-220.

A New Leptodactylid Frog from the Cordillera Oriental of Colombia

JOHN D. LYNCH¹

ABSTRACT. *Eleutherodactylus anolirex* is named from the northern Cordillera Oriental of Colombia (2,830–3,400 m). This species was previously confused with *E. supernatis* and *E. vertebralis*. It is most closely related to *E. devillei*, *E. supernatis*, and *E. vertebralis*, species of the western Andean cordilleras of Colombia and Ecuador.

INTRODUCTION

Although Cochran and Goin (1970) reported the leptodactylid frog from *Eleutherodactylus vertebralis* (Boulenger) to be widely distributed in Andean Colombia, Lynch (1980) showed *E. vertebralis* to be a species of relatively restricted distribution found on the high Pacific slopes of the Andes in central Ecuador and described *E. supernatis* from the Cordillera Central of Colombia and the northernmost Cordillera Real of Ecuador. Cochran and Goin (1970) confused several species with *E. vertebralis*. Except for noting the misidentifications of specimens from the Sierra Nevada de Santa Marta, Lynch (1980) deferred taxonomic comment on other purported populations of "*E. vertebralis*." Adequate material is now available to describe the frog from the Cordillera Oriental of Colombia reported by Duellman and Simmons (1977) as *E. vertebralis*.

TAXONOMY AND MORPHOLOGY

Eleutherodactylus anolirex sp. nov.

Figure 1

Holotype. KU 168626, an adult female, obtained 18.5 km (by road) S Chitagá, Departamento de Norte de Santander, Colombia, 2,850 m, on 23 August 1974 by William E. Duellman.

Paratypes. Topotypes, KU 168627–30; 32 km (by road) S Chitagá, Depto. Norte de Santander, 3,400 m, KU 168631–39; Presidente, 23 km S Chitagá, Depto. Norte de Santander, KU 150727–30; 35 km (by road) ENE Bucaramanga, road to Cucuta, Depto. Santander, 2,830 m, KU 132722–23; Páramo del Almorzadero, Depto. Santander, KU 150731–32, and 17 uncatalogued specimens in the Museo La Salle (Bogotá).

Diagnosis. 1) Skin of dorsum bearing flattened warts (least obvious anteriorly), that of venter coarsely areolate; dorso-lateral folds most distinct on posterior part of body; 2) tympanum prominent, its length one-third to one-half eye length; 3) snout subacuminate in dorsal view, round in profile; eye-nostril (E-N) < eye length; canthus rostralis sharp; 4) upper eyelid width narrower than interorbital distance (IOD); low cranial crests present; 5) vomerine odontophores low, oval in outline; 6) males with vocal sac and slits, non-spinous nuptial pads on thumbs; 7) first finger shorter than second; moderate pads on fingers II–IV; 8) fingers bearing lateral fringes; 9) ulnar tubercles present; 10) small tubercle on heel; elongate tubercle on inner edge of tarsus; 11) two metatarsal tubercles, inner oval, three to four times size of sub-conical outer; numerous supernumerary plantar tubercles; 12) toes bearing lateral

¹School of Life Sciences, University of Nebraska, Lincoln, Nebraska 68588, U.S.A.

fringes; no webs; toe pads slightly smaller than those of fingers; 13) brown above with faint cream dorsolateral stripes; lip bearing cream stripe; venter heavily flecked with brown; groin and concealed surfaces of limbs brown with small cream spots; 14) adults moderate-sized, males 24.0 to 31.8 mm, females 35.5 to 40.0 mm snout-vent length (SVL).

Eleutherodactylus anolirex is a member of the *vertebralis* assembly of Lynch and Duellman (1980), most similar to *E. bricensi* (Boulenger), but differs from all other species of the assembly in that males have vocal slits and a vocal sac.

Description. Head narrower than or equal to width of body; head width (HW) 36.4 to 39.4% ($\bar{x}$ = 37.8 n = 14) SVL; snout short, E-N 71.8 to 93.8% ($\bar{x}$ = 83.8, n = 14) eye length; snout subacuminate in dorsal view, rounded in lateral profile; nostrils weakly protuberant, directed dorsolaterally; canthus rostralis sharp, feebly sinuous; loreal region nearly flat, sloping abruptly to non-flared lips; upper eyelid width 69.7 to 87.4% ($\bar{x}$ = 80.6, n = 14) IOD; no tubercles on eyelids; edges of frontoparietals weakly upturned, producing frontoparietal furrow; supratympanic fold thick, glandular, obscuring upper and posterior edges of tympanic annulus: tympanum distinct, superficial, round, length 37.8 to 47.7% ($\bar{x}$ = 42.5, n = 8) length in males, 32.4 to 45.2% ($\bar{x}$ = 36.9, n = 6) in females; tympanum separated from eye by distance equal to 1 to 1.5 tympanic diameters; two prominent round postrictal tubercles; choanae small, round, not concealed by palatal shelf of maxillary arch when roof of mouth is viewed from directly below; vomerine odontophores low, medial and posterior to choanae, separated medially by distance twice the width of a tooth clump, each bearing a clump of three to five teeth; tongue longer than wide, posterior margin notched; posterior one-third of tongue not adherent to floor of mouth; males with vocal slits, subgular vocal sac.

Skin of dorsum appears smooth an-

teriorly but is pitted (here interpreted as bearing low, flattened warts), whereas on lower back and upper flanks warts are more pungent; flanks areolate; low dorsolateral folds (aligned warts) most evident posteriorly; similar fold above coccyx; venter coarsely areolate; discoidal folds present; not distinct; forearm bearing ulnar tubercles, antebrachial largest; palmar tubercle bifid, much larger than oval thenar tubercle; numerous supernumerary palmar tubercles; subarticular tubercles round, moderately pungent; fingers flattened, bearing lateral fringes; fingers bearing broad discs on relatively small, rounded pads; pads of fingers III and IV larger than tympanum; first finger shorter than second (when equally adpressed, tip of I reaches base of disc of II); male with diffuse non-spinous nuptial pads on dorsal surfaces of thumbs.

Heel bearing distinct but small tubercles; an elongate tubercle on inner edge of tarsus (or a fold along distal two-fifths of tarsus); no tubercles on outer edge of tarsus; outer metatarsal tubercle subconical, one-third to one quarter size of oval (length twice width) inner metatarsal tubercle; numerous supernumerary plantar tubercles arranged in rows; subarticular tubercles round, low; toes lacking webbing (basal between IV and V) but bearing distinct lateral fringes; pads bearing broad discs, slightly smaller than those of fingers: hind legs moderately long; shank 48.4 to 56.1% ($\bar{x}$ = 52.0, n = 15) SVL; heels overlapping when legs are flexed and held at right angles to sagittal plane; heel of adpressed hind leg reaching eye.

Dorsum dark brown with pale cream dorsolateral stripes; flanks brown with numerous cream flecks; labial stripe cream; canthal-supratympanic stripe brown to black; groin, anterior and posterior surfaces of thighs, ventral surfaces of shanks dark brown with small cream flecks or spots; venter cream with dense brown flecking to brown with numerous minute cream flecks (general appearance brown).

Figure 1. Hand and head of *Eleutherodactylus anolirex* sp. nov. (KU 168626, holotype). Scale for hand equals 2 mm, for head 5 mm.

Unfortunately few color notes on living specimens exist. Living specimens are described as dark brown with a distinct white stripe on the lip, venter gray-brown; iris bronze with median reddish-brown streak. The failure to mention the pale dorsolateral stripes apparent in preserved examples may mean that the stripes are not evident in living frogs.

Measurements of the holotype in mm. SVL 35.8, shank 18.4, HW 14.0, head length 11.6, upper eyelid width 3.8, IOD 4.4, tympanum length 1.4, eye length 4.2, E-N 3.7.

Etymology. The trivial name is Latin meaning King of Anoles and is used in loose reference to Ernest E. Williams. The name might also refer to the ease with which frogs of the genus *Eleutherodactylus* sit astride their lofty Andean

thrones gazing down toward the Andean slopes and lowland environments with their populations of Ernest's favorite animals, the anoles.

Relationships. *Eleutherodactylus anolirex* is a member of the *unistrigatus* group as defined by Lynch (1976). This species group is very large (Lynch listed about 100 species) and is centered distributionally in northwestern South America, especially in the Andes of Colombia and Ecuador (Duellman, 1979; Lynch, 1979). The high species densities in Ecuador are partially real and partially a product of intense systematic research there between 1966 and 1978. Lynch and Duellman (1980) advocated the use of assemblies (a subunit of a species group) in discussing relationships and distributions within the *unistrigatus* group.

Figure 2. Distributions of members of the *Eleutherodactylus devillei* assembly. *E. anolirex* (▲), *E. briceni* (■), *E. supernatis* (●). The distributions of *E. devillei* (eastern Andean slopes, Ecuador) and *E. vertebralis* (western Andean slopes, Ecuador) are outlined (see Lynch, 1980, for dot map).

Eleutherodactylus anolirex is a member of Lynch and Duellman's *devillei* assembly. The five named species of the assembly are all moderate-sized, cloud forest-dwelling *Eleutherodactylus* with moderately developed digital pads, broad interorbital spaces, and low cranial crests. The distribution of the five species (Fig. 2) is disjunct and encompasses both modes cited by Lynch (1981) for the páramo and subpáramo assemblies found in northern Ecuador. Similar disjunct distributions seen in more vagile groups of organisms often prompt evocations of dispersal as a means to explain the biogeographic pattern. However, equally likely explanations include 1) extinctions of geographically annectant populations and 2) inadequate collecting. Whatever explanation is advocated must be consistent with the phylogenetic statement for the assembly.

My hypothesis of the relationships of the five taxa is expressed in Figure 3. The characteristics and polarities employed are as follows: A—vocal slits (present = 0, absent = 1), B—eyelid tubercle (absent = 0, present = 1), C—separation of vomerine odontophores (wide = 0, narrow = 1), D—heel tubercle (absent = 0, present = 1), E—inner tarsal tubercle or fold (absent = 0, present = 1), F—toe tips black (no = 0, yes = 1), G—dorso-lateral folds (present = 0, absent = 1), H—canthus shape (sharp = 0, rounded = 1), I—occipital folds (absent = 0, present = 1), J—outer tarsal tubercles (absent = 0, present = 1). In each case, the

Figure 3. Cladogram of relationships in the *Eleutherodactylus devillei* assembly. Shifts from primitive to derived character-states are indicated by lettered horizontal lines (letters correspond to characters, see Table 1 and text).

derived state is scored as 1 and the primitive state as 0. The character-states for each species are summarized in Table 1.

The resultant cladogram (Fig. 3) appears to be geographically consistent (nearest relatives are geographically proximate for at least the three western and southernmost taxa). Unfortunately, further analysis is impeded by a dearth of knowledge about the eleutherodactyline frogs of the Cordillera Oriental of Colombia and the Venezuelan Andes. The frogs of the Ecuadorian Andes are better known, but phylogenetic statements about them are not available. The alleged impoverishment of the Cordillera Oriental is more apparent than real

TABLE 1. CHARACTER-STATES OF SPECIES OF THE *ELEUTHERODACTYLUS DEVILLEI* ASSEMBLY.

Character	<i>briceni</i>	<i>anolirex</i>	<i>devillei</i>	<i>supernatis</i>	<i>vertebralis</i>
A	1	0	1	1	1
B	0	0	0	1	1
C	1	0	1	0	1
D	0	1	0	1	1
E	0	1	1	1	1
F	0	0	0	1	1
G	0	0	0	1	0
H	0	0	0	1	0
I	0	0	0	0	1
J	0	0	0	0	1

(Pedro M. Ruíz, personal communication) and for the present prevents further analysis.

ACKNOWLEDGMENTS

Specimens were loaned and facilities for study made available by Hernando Daniel (Museo La Salle, Bogotá), William E. Duellman (Museum of Natural History, University of Kansas—KU), Pedro M. Ruíz (Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá), and Ernest E. Williams (Museum of Comparative Zoology, Harvard University). Special thanks are extended to Ken Miyata and to Susan and Anders Rhodin for the opportunity to participate in honoring a giant in our field.

LITERATURE CITED

- COCHRAN, D. M., AND C. J. GOIN. 1970. Frogs of Colombia. U.S. Natl. Mus. Bull., (288): 1–655.
- DUELLMAN, W. E. 1979. The herpetofauna of the Andes, pp. 371–459. In W. E. Duellman (ed.), The South American Herpetofauna: its origin, evolution, and dispersal. Univ. Kansas Mus. Nat. Hist. Monogr., (7): 1–485.
- DUELLMAN, W. E., AND J. E. SIMMONS. 1977. A new species of *Eleutherodactylus* (Anura, Leptodactylidae) from the Cordillera Oriental of Colombia. Proc. Biol. Soc. Washington, 90: 60–65.
- LYNCH, J. D. 1976. The species groups of the South American frogs of the genus *Eleutherodactylus* (Leptodactylidae). Univ. Kansas, Mus. Nat. Hist. Occas. Pap. (61): 1–24.
- . 1979. The amphibians of the lowland tropical forests, pp. 189–215. In Duellman, W. E. (ed.), The South American herpetofauna: its origin, evolution, and dispersal. Univ. Kansas, Mus. Nat. Hist. Monogr. (7): 1–485.
- . 1980. The identity of *Eleutherodactylus vertebralis* (Boulenger) with the description of a new species from Colombia and Ecuador (Amphibia: Leptodactylidae). J. Herpetol., 13: 411–418.
- . 1981. The eleutherodactyline frogs from the high Andes of northern Ecuador and adjacent Colombia. Univ. Kansas, Mus. Nat. Hist. Misc. Publ. (72): 1–46.
- LYNCH, J. D., AND W. E. DUELLMAN. 1980. The *Eleutherodactylus* of the Amazonian slopes of the Ecuadorian Andes (Anura: Leptodactylidae). Univ. Kansas, Mus. Nat. Hist. Misc. Publ. (69): 1–86.

Description of *Phrynops williamsi*, A New Species of Chelid Turtle of the South American *P. geoffroanus* Complex

ANDERS G. J. RHODIN¹

RUSSELL A. MITTERMEIER²

ABSTRACT. *Phrynops williamsi*, sp. nov., is a distinctive member of the *P. geoffroanus* complex (Pleurodira: Chelidae). The species is characterized by heavy black facial bands with a separate, thick, horseshoe-shaped band on the ventral surface of the neck, fine well-delineated radiating carapace reticulations, and no plastral markings. The skull differs markedly from other species of the *P. geoffroanus* complex and from most *Phrynops* in general. A wide parietal roof, lack of exoccipital contact above the foramen magnum, and widely divergent trochlear processes appear to be primitive features. Robust anterior maxillary triturating surfaces with lingual ridges and a shovel-shaped mandible adapted for bottom feeding appear to be unique derived characters. The species has a limited distribution in southern Brazil (Rio Grande do Sul and Santa Catarina) and adjacent Uruguay.

INTRODUCTION

The South American side-necked turtles of the family Chelidae are among the most poorly known of all turtles. No comprehensive review has ever appeared, and much of the scattered litera-

ture has suffered from inaccuracies based on inadequate sample analysis. Elusive and often rare in the wild, museum collections usually include only haphazardly or incidentally collected specimens. Many descriptions of new taxa have been based on single specimens, often without comparative material at hand. No clear definition exists of the valid taxa and their distribution, variation, morphology and ecology. Only Gaffney (1977) has looked at the family in a relatively comprehensive manner, but his work dealt with supraspecific relationships based on cranial osteology and examined less than half the known species.

In an effort to clarify the taxonomy of the South American Chelidae, Ernest E. Williams and Paulo E. Vanzolini began an extensive review of the family during the late 1950's, but abandoned the project due to inadequate material. They turned their preliminary work over to us several years ago, and this data base has served as the foundation for our work. It is therefore particularly fitting that the first description of one of the new species of chelid turtles to arise from this material appear in a volume dedicated to Dr. Williams. In addition, it is with great pleasure that we name this new species in honor of Ernest E. Williams, both in appreciation of his wide-reaching contributions to herpetology and the study of turtles, as well as in gratitude for his un-

¹Department of Surgery (Orthopaedics), Yale University School of Medicine, New Haven, Connecticut 06510 and Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, U.S.A. (Present address: Orthopaedic Associates, P.C., Nichols Rd., Fitchburg, Massachusetts 01420, U.S.A.)

²World Wildlife Fund—U.S. Primate Program, 1601 Connecticut Avenue, NW, Washington, D.C. 20009, U.S.A.

failing support, his friendship, and his guidance in our studies of systematic biology over the years. This has been a most rewarding relationship, undiminished by our respective professional divergences into the fields of orthopaedic surgery and primate conservation, and we take this opportunity to offer him our thanks.

Both Wermuth and Mertens (1977) and Pritchard (1979) recognize 18 taxa of South American chelids, though they differ in their interpretation of subspecific rank in three cases. Of the four currently recognized genera (*Chelus*, *Hydro-medusa*, *Phrynopis*, *Platemys*), *Phrynopis* is the most complex taxonomically, with eleven currently recognized taxa: *P. dahli*, *geoffroanus*, *gibbus*, *hilarii*, *hogeii*, *nasutus*, *rufipes*, *tuberculatus*, *tuberosus*, *vanderhaegei*, and *wermuthi*. Three of these taxa (*P. geoffroanus*, *hilarii*, *tuberosus*) form a natural superspecies complex based on external and osteological similarities and will hereafter be referred to as the *P. geoffroanus* complex. Briefly, the forms within the complex share the general features of large, broad shells with moderately wide heads, parietal roof moderately broad, neurals usually numbering six or seven and contacting the nuchal bone broadly, carapace color either dark brown or reticulated with radial black markings, plastron color usually reddish, yellow or white and either immaculate or with scattered dark vermiculations or spots, head with characteristic lateral black stripe from snout through the eye and tympanic membrane and extending along the lateral surface of the neck, ventral surface of the neck with either broad black bands, scattered thinner vermiculations or spots.

During the course of our revision of the South American Chelidae, we have personally examined 353 specimens of members of the *P. geoffroanus* complex. Of this series, 54 represent the readily distinguishable *P. hilarii*, with its immaculate dark brown or gray carapace, black-spotted white plastron, and re-

duced head markings. Of the 299 other specimens of *P. geoffroanus* and *P. tuberosus*, several distinct geographic populations are identifiable. A complete analysis of the systematic status of these populations is beyond the scope of the present paper, but will be presented in a future publication. The most distinctive of the populations is represented by six specimens from Rio Grande do Sul and Santa Catarina in Brazil. Though allopatric in respect to other *P. geoffroanus* and *tuberosus*, the combination of external and osteological features present in this population warrant its recognition as a new species within the *P. geoffroanus* complex.

TAXONOMY AND MORPHOLOGY

Phrynopis williamsii sp. nov.

Figures 1–6

Holotype. MCZ 64135, Rio Grande do Sul, Brazil, collected by H. von Ihering.

Paratypes. BMNH 84.2.5.1, 84.2.5.3, Rio Grande do Sul, Brazil, collected by H. von Ihering; MZUSP 2675, Pôrto Alegre, Rio Grande do Sul, Brazil; MNRJ 3146, Tubarão, Santa Catarina, Brazil; ZMB 6858, "Estancia Velha," Rio Grande do Sul, Brazil, collected by R. Hensel (probably collected at Pikada do Café, Rio Cadéa).

Type Locality. Since the holotype specimen has no specific locality, we hereby designate the type locality as Rio Cadéa, Rio Grande do Sul, Brazil, where the species was first collected by R. Hensel in 1865.

Synonymy

- Platemys geoffreyana* (partim) Hensel, 1868:350
- Platemys geoffroyana* (partim) Hensel, 1868:354; Boulenger, 1885:191, 1886:424; Strauch, 1890:104; Lema, 1958:11
- Hydraspis geoffroyana* (partim) Boulenger, 1889:223; Siebenrock, 1904:23; Goeldi, 1906:751; Siebenrock, 1909:576; Luederwaldt, 1926:433
- Phrynopis geoffroyana* (partim) Vaz-Ferreira, 1955:xxv; Froes, 1957:19
- Phrynopis geoffroana geoffroana* (partim) Mertens and Wermuth, 1955:404; Vaz-Ferreira and Sierra de Soriano, 1960:14
- Phrynopis geoffroanus geoffroanus* (partim)

- Wermuth and Mertens, 1961:333, 1977:130;
 Pritchard, 1967:234, 1979:787
Phrynops geoffroanus (partim) Freiberg, 1970:190,
 1971:92, 1972:248, 1975:92; Lema and Fabian-
 Beurmann, 1977:65
Phrynops geoffroana (partim) Achaval, 1976:26
Phrynops geoffroanus spp. (partim) Mittermeier,
 Medem, and Rhodin, 1980:15

Distribution. Low-lying areas (below 500 m elevation) of eastern coastal Santa Catarina and Rio Grande do Sul in Brazil as well as the northern half of Uruguay as far west as the Rio Uruguay, possibly including eastern portions of Entre Rios and southeastern Corrientes in Argentina as well as southwestern inland Rio Grande do Sul (Fig. 9). Apparently absent from southern coastal Uruguay and northwestern inland Rio Grande do Sul in the upper Rio Uruguay and Rio Pelotas drainages.

Diagnosis. A moderately-sized member of the *P. geoffroanus* complex characterized by three heavy black facial stripes including a separate thick horse-shoe-shaped stripe on the ventral neck, well-delineated thin radiating carapacial reticulations and no plastral markings. Skull characterized by a wide parietal roof, lack of exoccipital contact above the foramen magnum, widely divergent trochlear processes, and robust anterior maxillary triturating surfaces with lingual ridges and shovel-shaped mandible.

EXTERNAL MORPHOLOGY

Carapace. Carapace (Fig. 1) broadly oval, almost subcircular in juveniles, carapace length averaging 1.15 times width, becoming relatively narrower with increasing size, carapace length averaging 1.33 times width in subadults and smaller adults, 1.46 in larger adults (see Table 1 for all measurements and ratios of external features). Shell moderately deep, length averaging 3.39 times height in adults and subadults. No marginal flaring, recurving or broad expansion. Mid-lateral marginals slightly narrower than anterior and posterior ones. Entire marginal rim mildly serrate in

juveniles, partially retained posteriorly in subadults and adults but less distinct. Small supracaudal notch. Nuchal approximately twice as long as broad, projecting slightly anterior to carapace margin. Vertebrae generally wider than long, including V4 which is notably wide. Intercoastal lateral seam contacts at M2, 5, 7, 9, and 11. Vertebrae without furrow, trough, keel or knobs in subadults and adults. Very vague, low mid-line bulge posteriorly on V5, extending toward supracaudal notch, causing very mild keeling at posterior end of shell, more prominent in large adults than subadults. Juveniles with low inconspicuous vertebral bulges on V1-5 and costals.

Carapace color in preserved specimens medium to light brown with extensive thin black reticulations radiating in a well-delineated radial pattern on all vertebrae, costals and marginals. Center of radiating pattern located at site of original juvenile scute: posteriorly in mid-line on vertebrae, posteromedially on costals, and posterolaterally on marginals. Pattern as distinct in large adults as in juveniles. Black lines generally same thickness or thinner than interspersed brown base color. Color demarcation sharp and clear, with pattern extending fully to edge of marginal rim. Carapace color in live specimens brown with black reticulations and thin yellowish orange carapacial edge (Vaz-Ferreira and Sierra de Soriano, 1960).

Plastron. Plastron (Fig. 2) broad, length averaging 1.82 times width, carapace width averaging 1.60 times plastron width, anteriorly truncate, posteriorly slightly narrower. Anal notch moderately deep. Very small axillary and inguinal scutes present. Intergular short and broad. Plastral seam length formula $Fem > Ig \geq An > Abd \geq Pect \geq Hum$.

Plastron color in preserved specimens yellowish brown, occasionally oxidized to darker brown. Immaculate on all ventral surfaces, though occasional large specimens with indistinct dorsal type color pattern on posterior ventral margin-

Figure 1. Dorsal view of carapace of ZMB 6858, a paratype of *P. williamsii*. Note the clearly delineated fine radiating reticulations.

als. Plastron color yellow in live specimens (Vaz-Ferreira and Sierra de Soriano, 1960).

Head and Neck. Head width moderately narrow, becoming relatively narrower as compared to carapace length with allometric growth (Fig. 8). Neck length fairly short.

Head and neck with distinctive color pattern (Fig. 3), primarily dark dorsally and light ventrally, characterized by three subparallel broad black bands. The uppermost band serves as the ventral border of the dark dorsal head and neck pattern, extending from the nostrils through the eye, through the upper one

Figure 2. Ventral view of BMNH 84.2.5.3, a paratype of *P. williamsi*. Indistinct reticulations can be seen on marginals posterior to bridge.

Figure 3. Ventral and lateral views of head and neck of MCZ 64135, the holotype of *P. williamsii*. Ventral photograph slightly retouched to obscure small laceration present on specimen itself.

third to one half of the tympanum, and then along the mid-lateral surface of the neck, gradually fading caudally. The lowermost band forms a posteriorly directed horseshoe-shaped figure on the ventral chin, extending anteriorly to the base of the barbels, usually sharply discontinuous posteriorly, with an interruption at the level of the posterior border of the tympanum, before continuing for a short distance as short subparallel bands or elongate spots. The intermediate band extends caudally from the angle of the jaws, serving as a continuation of thin bands of dark pigment on the external tomial surfaces of the inferior portion of the maxillary and superior portion of the mandibular horny sheaths. The band then continues along the inferior border of the tympanic membrane and ventrolaterally along the neck, ending abruptly at approximately the same level as the

last band or spot in the lowermost band. These three broad bands are usually totally separate from one another. Of seven specimens examined (including specimens figured in the literature), none had a connection between the upper and intermediate bands. Three out of seven specimens had very thin connections between the lower and intermediate bands posterior to the tympanum, two of them only unilateral and one with thin bilateral connections. One out of seven specimens had a discontinuous ventral horseshoe, narrowly lacking midline contact under the chin. The dorsal head pattern is relatively indistinct, composed of a dark background with narrow indistinct lighter lines subparallel to the uppermost dark band.

Live color of dorsal surface of head black with whitish lines, ventral surface of chin and neck reddish yellow or yellow with black bands (Vaz-Ferreira and Sierra de Soriano, 1960).

Two barbels present on the ventral surface of the chin, arising from the anterior portion of the horseshoe band. Barbels often unpigmented, though one sometimes the same color as the band.

Skin of top of snout, interorbital region and middle third of top of head adherent to bone, incompletely divided into apparent scales. Remaining dorsal surface of head (above temporal muscles) with more regular polygonal scales. Skin of dorsal aspect of neck plicate with only occasional more or less well defined papillae. Skin of ventral aspect of neck shallowly plicate, reticulate.

Limbs. Dorsal aspect of limbs with indistinct dark vermicular pattern. Ventral aspect light colored, juvenile with few scattered spots and vermiculations. Enlarged scales forming prominent swim flap on free ulnar skin edge of anterior limbs. Three pre-tibial scales enlarged, with the distal one prominent and cornified, resembling a blunt claw. No pre-tibial flap or ischial tuberosities. Claws numbering five on forelimbs, four on hindlimbs. Color of limbs in life blackish

TABLE 1. *PHRYNOPS WILLIAMSII* EXTERNAL MEASUREMENTS AND BASIC RATIOS.*

Specimen Number	Sex	CL	CW	CD	PL	PW	HW	CL/CW	CL/CD	CL/HW	CW/PW	PL/PW
MNRJ 3146	juv	77	67	28	65	40	16.0	1.15	2.75	4.81	1.68	1.63
MZUSP 2675	p ♀	140	108	40	119	70	24.0	1.30	3.50	5.83	1.54	1.70
MZVC-R 193†	♂	141	117					1.21				
MCZ 64135	p ♀	180	138	59	162	89	29.0	1.30	3.05	6.21	1.55	1.82
ZMB 6858	p ♂	201	154	54	171	92	31.5	1.31	3.72	6.38	1.67	1.86
BMNH 84.2.5.1	♀	243	179	70	206	112	38.0	1.36	3.47	6.39	1.60	1.84
BMNH 84.2.5.3	♀	252	183	79	212	112	37.5	1.38	3.19	6.72	1.64	1.89
MZVC-R 192†	♀	305	214					1.43				
Freiberg s/n†	♀	330	220					1.50				

*CL = carapace length, CW = carapace width, CD = carapace depth, PL = plastron length, PW = plastron width, HW = head width (tympenic). All measurements in mm.
 †Recorded from the literature (Vaz-Ferreira and Sierra de Soriano, 1960; Freiberg, 1970).

TABLE 2. *PHRYNOPS WILLIAMSII* SKULL MEASUREMENTS AND BASIC RATIOS.*

Specimen Number	CL	SL	SWT	SWM	SD	IOW	OW	PW	PtW	CL/SL	SL/SWT	SL/SWM	IOW/OW	IOW/PW	PW/PtW	SL/SD	PW/SWT
ZMB 6858	201	35.8	30.0	20.2	11.8	6.0	7.5	11.8	12.8	5.61	1.19	1.77	0.80	0.51	0.92	3.03	0.39
BMNH 84.2.5.3	252	44.5	35.5	24.5	14.5	7.6	8.6	12.8	16.0	5.66	1.25	1.82	0.88	0.59	0.80	3.07	0.36

*CL = carapace length, SL = skull length (condylobasilar), SWT = skull width (tympenic), SWM = skull width (maxillary), SD = skull depth (midline, posteriorly), IOW = interorbital width, OW = orbital width, PW = parietal width (minimal), PtW = pterygoid width (minimal). All measurements in mm.

brown dorsally, reddish yellow or yellow ventrally, with red webbing on feet and red swim flap on forelimb (Vaz-Ferreira and Sierra de Soriano, 1960).

Sexual Dimorphism and Size. Sexual dimorphism not apparent from series examined. Large specimens with deep shells, short tails and flat plastrons represent females. No obvious males with shallow shells, long tails and concave plastrons present in series, but Vaz-Ferreira and Sierra de Soriano (1960) figure a 141 mm male with a long, thick tail. The largest specimen examined measured 252 mm carapace length, but Vaz-Ferreira and Sierra de Soriano (1960) recorded a specimen measuring 305 mm and Freiberg (1970) measured a 330 mm individual.

OSTEOLOGY

Skull. Rather than describe the skull of *P. williamsi* in detail, we prefer to discuss it in relation to other *Phrynops* and chelid skulls. We have examined two skulls of *P. williamsi* and have compared them to 47 skulls of other *Phrynops* (representing all known species), of which 22 represent other members of the *P. Geoffroanus* complex. Most of this comparative material represents our unpublished data. However, Gaffney (1977) has discussed and figured skulls of *P. hylarii* (as "*P. Geoffroanus*") and *P. gibbus*, and we provide figures of *P. williamsi* (Figs. 4, 5) and *P. Geoffroanus* (Figs. 5, 7) in this work. Other species mentioned (e.g., *P. rufipes*, *hogeii*, *vanderhaegei*) will be fully discussed and figured in future publications. Table 2 lists all skull measurements and ratios for the two skulls of *P. williamsi* examined.

A number of features distinguish the skull of *P. williamsi* from the other members of the *Geoffroanus* complex. In fact, it is among the most distinctive of all *Phrynops* skulls in many respects.

The lateral extent of the parietal roof is greater than in most other *Phrynops*, the sides being nearly parallel. There is considerably less posterior temporal emargi-

nation, with relative loss of overlap between the anterior and posterior temporal emarginations (slight overlap in the larger specimen). The temporal arch is shorter and thicker than in other *Phrynops*. There is very little lateral maxillary protrusion, but two other *Phrynops* (*hogeii* and *rufipes*) show less protrusion, and other young *Geoffroanus* complex species can have the same extent. This characteristic is ontogenetically variable and larger *williamsi* have a slightly more developed protrusion.

The two processi trochlearis pterygoidei show wide divergence from the longitudinal axis of the skull, making a ca. 70 to 90° angle with each other when viewed from above. No other *Phrynops* shows this condition, though *hogeii* reaches 50 to 60° and *rufipes* 60 to 70°. All other *Phrynops* have an angle of 45° or less. The maxillary triturating surface has a moderately well-developed short lingual ridge anterior to the choanae, a condition not seen in any other *Phrynops*. Compared to other *Geoffroanus* complex species, the anterior maxillary region of *williamsi* is much more pronounced and robust with a thicker, heavier maxilla, an angular rather than rounded snout, a wider maxillary triturating surface especially anteriorly and the presence of short lingual ridges. The only other *Phrynops* with prominent ventral snout development are *hogeii* and *rufipes*, with *gibbus* and *vanderhaegei* showing an intermediate condition.

The mandible of *williamsi* is very distinctive, with the triturating surface being markedly widened and extremely flattened, coming very close to the horizontal. This is totally unlike other *Geoffroanus* complex species or any other *Phrynops*, including *hogeii* and *rufipes*. It most closely resembles the condition seen in the South American pelomedusine turtle *Podocnemis sextuberculata*. This shovel-like appearance of the mandible may represent a specialized adaptation to bottom feeding.

The exoccipitals of *williamsi* do not

Figure 4. Osteology of skull and mandible of *P. williamsi* (ZMB 6858). A. dorsal view; B. ventral view; C. lateral view; D. posterior view; E. lateral view of mandible; F. medial and slightly dorsal view of right half of mandible. Refer to Gaffney (1979) and Rhodin and Mittermeier (1976) for skull nomenclature.

meet above the foramen magnum, being moderately separated by the supraoccipital. This feature is also shared by *hogeii* and *rufipes*. The same feature, though less pronounced with only narrow separation, is variably present in *vanderhaegei*, *gibbus*, *hilarii* and *geoffroanus*.

The normal condition in these species is either a narrow or moderate exoccipital contact. No other South American chelid has separated exoccipitals, the feature being shared only by the Australopapuan genera *Emydura*, *Elseya*, and *Pseudemydura* (Gaffney, 1977).

Figure 5. Dorsal views of mandibles of *P. williamsi* (left, BMNH 84.2.5.3) and *P. geoffroanus* (right, MZUSP 50; Rio Grande, Sao Paulo, Brazil; carapace length 250 mm, skull condylobasilar length 44 mm). Note markedly widened and flatter triturating surface in *P. williamsi*. Photographic enlargement slightly greater for *P. geoffroanus*.

Figure 6. Neural bones of *P. williamsi* (BMNH 84.2.5.1).

The foramina nervi trigemini of *williamsi* face laterally, and are not visible from the dorsal aspect of the skull. In other *geoffroanus* complex species the foramina face dorsally and are easily visible from the dorsal aspect. The flattened horizontal portion of the pterygoid flare is relatively reduced in *williamsi* so that only a very small portion of the pterygoid is visible from the dorsal aspect of the skull. In other *geoffroanus* complex species the pterygoid flare is quite wide so that a broad expanse of pterygoid is visible from the dorsal aspect.

The postorbital wall of *williamsi* has a distinct sulcus jugalis and the medial portions of the jugal and postorbital face more posteriorly than all other *Phrynops*, where this wall faces more dorsolaterally and has a less well-developed sulcus jugalis.

The features of widely divergent trochleas, deep sulcus jugalis with overlying parietal roof, posteriorly facing postorbital wall, shovel-like appearance of

Figure 7. Osteology of skull and mandible of *P. geoffroanus* (MZUSP 2680; Ilha Solteira, Rio Parana, São Paulo, Brazil; carapace length 172 mm, skull condylobasilar length 33.7 mm). Views same as in Fig. 4.

the mandible and robust widened anterior maxillary triturating surface with lingual ridges appear to all represent feeding adaptations. The power of jaw closure is heightened through increased muscle mass in the anterior portions of the jaw adductors combined with divergent trochleas which improve leverage.

This increased muscle mass is most prominent in the relatively enlarged area of the postorbital wall and sulcus jugalis. The modified triturating surfaces appear adapted for bottom feeding and crushing of large hard food such as snails or small bivalves. Other *Phrynops* do not possess this spectrum of characteristics

Figure 8. Head width in *P. williamsi*. Plot of carapace length in mm (abscissa) versus carapace length divided by head width (ordinate).

and many of them are known to be predatory feeders. Unfortunately, nothing is known of the feeding habits of *williamsi*.

The wide parietal roof, lack of exoccipital contact above the foramen magnum and widely divergent trochlear processes of *P. williamsi* appear to be primitive features within the genus *Phrynops*. These features show a similarity to the pelomedusine turtles of the genus *Podocnemis* as well as the Australopapuan chelids of the genera *Elseya*, *Emydura*, and *Pseudemydura* (see Gaffney, 1977 and McDowell, 1983). The robust anterior maxillary triturating surfaces with lingual ridges and shovel-shaped mandible may be primitive features, but more likely represent unique derived characters within the genus *Phrynops*, showing secondary convergence with *Podocnemis sextuberculata* due to similar feeding strategy.

Cervicals. One specimen examined has the typical central articulation pattern of all Chelidae as described by Williams (1950): (2)3(4(5)6)7(8).

Neurals. Six large contiguous neurals present (Fig. 6). N1 ca. twice as long as N2-5. N6 half as small as N2-5. N1 contac-

ting nuchal widely, broadly rectangular. N2-5 roughly hexagonal, tapering posteriorly. N6 small and pentagonal, only partly separating C6. Pattern identical in the two specimens examined.

REPRODUCTION

No field data are available on eggs, nests or hatchlings of *P. williamsi*. One specimen, BMNH 84.2.5.3, a 252 mm female collected somewhere in Rio Grande do Sul, contained nine white, oval, brittle shelled oviductal eggs (four in the left oviduct, five in the right). The average egg size in this series was 33.3×27.0 mm with length ranging from 32.9–34.2 mm and width ranging from 26.7–27.6 mm. It is unclear when the specimen was collected, but egg deposition probably occurs in either November or December. *Phrynops hilarii* in northeastern Argentina nests in November (Gallardo, 1980) and in Rio Grande do Sul nests from late October to early January, with the peak activity occurring in November (Reischl *et al.*, 1979). *Pseudemys dorbignyi* (Emydidae) from the same area has the same nesting season but peaks primarily in December (Reischl *et al.*, 1979).

Clutch size in *P. hilarii* is apparently somewhat larger than in *P. williamsi*. In Rio Grande do Sul clutch size of *P. hilarii* averages 11, with the incubation period taking 105 to 140 days (Reischl *et al.*, 1979). Serrano (1977) notes that *P. hilarii* in Rio Grande do Sul lays clutches averaging 13.4 eggs (range 1–20) with spherical eggs measuring 33 ± 4 mm in diameter. Saporiti (1960) indicates that *P. hilarii* in Buenos Aires lays 10 to 14 subspherical eggs averaging 32.9×30.8 mm in size. Cohen (personal communication) had a captive specimen lay 17 eggs averaging 36.1×33.8 mm in size.

Few comparative data are available for *P. geoffroanus* and *tuberosus*. Wied (1825) noted that populations of *P. geoffroanus* inhabiting the rivers of eastern coastal Brazil (e.g., Rios Pardo, Jequitinhonha, and Mucuri) laid from 12 to 18 spherical eggs from December through

February. Medem (1969) recorded three nests of *P. tuberosus* from Colombia as having from 10 to 18 subspherical eggs with average size measuring 33.9×32.5 mm. We have examined a specimen of *P. geoffroanus* from the Rio Tapajós, Pará, Brazil (MZUSP 2682) containing 13 subspherical shelled eggs, average size measuring 30.5×29.0 mm.

HABITAT

Vaz-Ferreira and Sierra de Soriano (1960) obtained two specimens at Picada del Negro Muerto, Uruguay, where the Rio Cuareim flows relatively rapidly over a rocky streambed. Hensel (1868) noted that the species occurred in the "rushing and stony brooks of the forest" and was not to be found in the slower lowland rivers with muddy bottoms.

DISCUSSION

The species here recognized as *Phrynops williamsi* was first collected and described as *Platemys geoffreyana* (Schweigger, 1812) by Hensel (1868). His description of an animal he collected in 1865 at Pikada do Café, Rio Cadéa, Rio Grande do Sul fits perfectly our description of *P. williamsi*. Another specimen of the same species may have been observed at Estancia Velha, but Hensel's description of that animal is not as clear and it may have been based on a specimen of *P. hylarii*. Hensel states that he was able to obtain only a single specimen of "*P. williamsi*" on his trip. This specimen was collected at Pikada do Café, Rio Cadéa and had a carapace length of 202 mm (Hensel's measurement). The *P. williamsi* paratype allegedly collected by Hensel at Estancia Velha (ZMB 6858) has a carapace length of 201 mm, and if indeed collected by Hensel, probably represents the original Rio Cadéa specimen, though now mislabeled as coming from Estancia Velha.

The species was next collected by von Ihering, whose specimens served as the

basis for Boulenger's (1889) description of *Hydraspis geoffroyana* (Schweigger). One of these specimens is our *P. williamsi* holotype, two others are paratypes.

Clear descriptions of animals referable to *P. williamsi* did not re-appear in the literature until Vaz-Ferreira and Sierra de Soriano (1960) figured and described specimens of *P. geoffroana geoffroana* from Uruguay. Some of these and others as well were described and figured by Freiberg (1970) and served as his basis for the specific separation of *P. hylarii* from *P. geoffroanus* based on their Uruguayan sympatry.

The characteristics of *P. williamsi* discussed above readily distinguish it from any other *Phrynops*, as well as from any other member of the *P. geoffroanus* complex. The black bands on the head and neck coupled with the reticulate carapace pattern immediately differentiate it from *P. gibbus*, *rufipes*, *hogeii*, *nasutus*, *dahli*, *wermuthi*, *vanderhaegei*, and *tuberculatus*. *Phrynops hylarii* is distinguished by its black spots on a white plastron, immaculate carapace and markedly reduced head and neck bands (see photos in Freiberg, 1970, 1975). *Phrynops geoffroanus* and *P. tuberosus* represent two recognized populations of a very wide-spread species complex with marked geographic variation, consisting of several previously unrecognized and undescribed forms. In general, *P. williamsi* is distinguished from all of these populations by the combination of features listed. The most marked distinction is in skull morphology, but this is a feature not always available for identification purposes. The carapacial reticulations of *P. williamsi* are thinner, finer and more clearly well developed than in *P. geoffroanus* or *tuberosus* where they tend to be either broad and thick, irregular, unclear, or reduced. Most populations of *P. geoffroanus* and *tuberosus* have plastrons with varying degrees of a dark vermiculate pattern. However, some also have immaculate plastrons, so this

Figure 9. Map showing distribution of *P. williamsi* (▲), *P. geoffroanus* (●) and *P. hilarii* (⊛). Geographic ranges of the latter two species only partially shown. Stippled areas correspond approximately to elevations greater than 500 m.

feature in itself is not diagnostic for *P. williamsi*. The black head and neck bands of *P. geoffroanus* and *tuberosus* are almost always relatively thin and usually confluent in several places. Some populations have very thin bands, others somewhat thicker ones, but none are as prominent as in *P. williamsi*. Though occasionally separated from each other, the facial bands of *P. geoffroanus* and *tuberosus* usually connect broadly either in front of the tympanum (intermediate and lower bands connecting), behind the tympanum (upper and intermediate bands connecting) or at several places

along their course. Many populations have only two bands plus irregular vermiculations. A separate ventral horse-shoe band is usually not present, though some specimens of one population of *P. geoffroanus* from eastern coastal Brazil occasionally exhibit this character as well.

Based on external morphology alone it would be difficult to determine whether *P. williamsi* is distinct from other *P. geoffroanus* and *P. tuberosus* at a specific or subspecific level. Though sympatric with *P. hilarii*, it is allopatric with respect to *P. geoffroanus*, the closest known

populations occurring approximately 300 km to the north in the Rio Pelotas drainage (Fig. 9). However, the intervening region has not been well collected, and a zone of sympatry may well exist. The distinct skull morphology of *P. williamsi* argues strongly for specific status. The unique combination of presumably primitive and derived features is not shared even by other *Phrynops*, let alone other members of the *P. geoffroanus* complex.

ACKNOWLEDGMENTS

We thank the following people for assistance in various aspects of this work: F. Achaval, E. N. Arnold, A. L. de Carvalho, H. J. Cohen, G. Peters, S. D. Rhodin, J. Rosado, P. E. Vanzolini and E. E. Williams. Illustrations were prepared by J. Braslin and A. G. J. Rhodin. Rhodin's studies were partly funded by the American Philosophical Society, and Mittermeier's by the World Wildlife Fund—U.S.

APPENDIX: LOCALITY DATA

Italicized numbers represent specimens of *Phrynops williamsi* examined or confirmable literature records. BMNH = British Museum of Natural History; MCZ = Museum of Comparative Zoology, Harvard University; MHNM = Museo Nacional de Historia Natural, Montevideo; MNRJ = Museu Nacional, Rio de Janeiro; MZUSP = Museu de Zoologia da Universidade de São Paulo; MZVC-R = Museo de Departamento de Zoologia Vertebrados, Facultad de Ciencias, Universidad de la Republica, Montevideo; ZMB = Zoologisches Museum, Berlin.

BRAZIL: *Rio Grande do Sul*: MCZ 64135, BMNH 84.2.5.1, 84.2.5.3, Boulenger 1885, 1886, 1889; Estancia Velha (29°39'S, 51°11'W): ZMB 6858, Hensel 1868; Pikada do Café, Rio Cadéa (29°37'S, 51°15'W): Hensel 1868; Porto Alegre

(30°03'S, 51°10'W): MZUSP 2675; *Santa Catarina*: Tubarão (28°29'S, 49°00'W): MNRJ 3146. URUGUAY: *Artigas*: Freiberg 1970; *Picada del Negro Muerto*, Rio Cuareim (30°45'S, 56°05'W): MZVC-R 192-3, Vaz-Ferreira and Sierra de Soriano 1960; *Salto de Agua del Penitente*: MHNM 1582, Lema and Fabian-Beurmann 1977; *Paysandu*: Arroyo Chapiçuy Grande (31°42'S, 57°55'W): Freiberg 1970; *Cerro Largo*: Arroyo Yaguarón (31°55'S, 53°55'W): MHNM s/n; *Rivera*: Arroyo Cunapirú (31°30'S, 55°35'W): MZVC-R s/n; *Rio Negro*: Rincón del Bonete, Represa del Rio Negro (32°52'S, 56°27'W): Freiberg 1970; *Salto*: Rio Arapey (30°55'S, 57°50'W): MHNM s/n.

LITERATURE CITED

- ACHAVAL, F. 1976. Lista de las especies de vertebrados del Uruguay. Reptiles. Montevideo: Museo Nacional de Historia Natural, pp. 26-29.
- BOULENGER, G. A. 1885. A list of reptiles and batrachians from the Province Rio Grande do Sul, Brazil, sent to the Natural-History Museum by Dr. H. von Ihering. *Ann. Mag. Nat. Hist.*, (5)15: 191-196.
- . 1886. A synopsis of the reptiles and batrachians of the Province Rio Grande do Sul, Brazil. *Ann. Mag. Nat. Hist.*, (5)18: 423-445.
- . 1889. Catalogue of the Chelonians, Rhynchocephalians, and Crocodiles in the British Museum (Natural History). London: Trustees of the Museum, pp. 311.
- FREIBERG, M. A. 1970. Validez específica de "*Phrynops hilarii*" (D. & B.) (Testudines, Chelidae). *Rev. Mus. Argent. Cienc. Nat. Zool.*, 10: 189-197.
- . 1971. El mundo de las tortugas. Buenos Aires: Albatros, 139 pp.
- . 1972. Validez específica de *Phrynops hilarii* (D. & B.) (Testudines, Chelidae). *Acta Zool. Lilloano*, 29: 247-260.
- . 1975. El mundo de las tortugas. 2a. Edición. Buenos Aires: Albatros, 134 pp.
- FROES, O. M. 1957. Atualização da nomenclatura dos quelônios Brasileiros. *Iheringia Zool.*, 2: 1-24.
- GAFFNEY, E. S. 1977. The side-necked turtle family Chelidae: a theory of relationships using shared derived characters. *Amer. Mus. Novitates* No. 2620, pp. 1-28.
- . 1979. Comparative cranial morphology of recent and fossil turtles. *Bull. Amer. Mus. Nat. Hist.*, 164: 65-376.
- GALLARDO, J. M. 1980. Estudio ecológico sobre los anfibios y reptiles del noreste de la provincia de Buenos Aires (Argentina). I. Reunión Iberoamer. Zool. Vert., La Rabida, 1977(1980): 331-349.

- GOELDI, E. A. 1906. Chelonios do Brazil (Jabotys—Kagados—Tartarugas). Bol. Mus. Goeldi, 1906: 699–756.
- HENSEL, R. 1868. Beitrage zur Kenntniss der Wirbelthiere Sudbrasilien. Arch. Naturg., 1868(1): 323–375.
- LEMA, T. DE 1958. Notas sobre os repteis do Estado do Rio Grande do Sul—Brasil. I. Introducao ao estudo dos repteis do Estado do Rio Grande do Sul. Iheringia Zool., 10: 1–18.
- LEMA T. DE AND M. E. FABIAN-BEURMANN. 1977. Levantamento preliminar dos repteis da regio da fronteira Brasil-Uruguai. Iheringia Zool., 50: 61–92.
- LUEDERWALDT, H. 1926. Os chelonios brasileiros. Rev. Mus. Paulista, 14: 403–470.
- MCDOWELL, S. B. 1983. The genus *Emydura* (Testudines: Chelidae) in New Guinea with notes on the penial morphology of Pleurodira. This Volume, pp. 169–189.
- MEDEM, F. 1969. Estudios adicionales sobre los Crocodylia y Testudinata del alto Caqueta y Rio Caguan. Caldasia, 10: 329–353.
- MERTENS, R., AND H. WERMUTH. 1955. Die rezenten Schildkroten, Krokodile und Bruckenechsen. Zool. Jahrb., 83: 323–440.
- MITTERMEIER, R. A., F. MEDEM, AND A. G. J. RHODIN. 1980. Vernacular names of South American turtles. Soc. Stud. Amph. Rept. Herpetol. Circ., 10: 1–44.
- PRITCHARD, P. C. H. 1967. Living turtles of the world. Jersey City, T.F.H., 288 pp.
- . 1979. Encyclopedia of turtles. Hong Kong, T.F.H., 895 pp.
- REISCHL, E., C. O. D. C. DIEFENBACH, AND C. V. TONDO. 1979. Ontogenetic variation of hemoglobins from South American Chelonia (Reptilia). Comp. Biochem. Physiol., 62B: 539–544.
- RHODIN, A. G. J., AND R. A. MITTERMEIER. 1976. *Chelodina parkeri*, a new species of chelid turtle from New Guinea, with a discussion of *Chelodina siebenrocki* Werner, 1901. Bull. Mus. Comp. Zool., 147: 465–488.
- SAPORITI, E. J. 1960. Iconografia de las tortugas que vivieron y/o viven en el Jardin Zoologico de la Ciudad de Buenos Aires. Diana; Revista Caza-Tiro-Canofilia-Pesca, 236, 238, 240, 246: 40–46, 62–65, 68–70.
- SCHWEIGGER, A. F. 1812. Prodromus monographiae Cheloniorum. Königsb. Arch. Naturwiss. Math., 1: 271–368, 406–458.
- SERRANO, L. L. 1977. Nota sobre uma membrana extra-embionaria precoce, no embriao de *Phrynops hilari* Dumeril e Bibron, 1835 (nota previa). Pesqui. Zool. (Porto Alegre), 29: 7–13.
- SIEBENROCK, F. 1904. Schildkroten von Brasilien. Akad. Wiss. Wien. Math. Naturwiss. Kl. Denkschr., 76: 1–28.
- . 1909. Synopsis der rezenten Schildkroten. Zool. Jahrb. Suppl., 10: 427–618.
- STRAUCH, A. 1890. Bemerkungen über die Schildkrotensammlung im Zoologischen Museum. Mem. Ac. St. Petersburg, (7)38: 1–127.
- VAZ-FERREIRA, R. 1955. La fauna en el manuscrito de Toller. In Toller, W. 1955. Viaje de William Toller a la Banda Oriental y Rio de la Plata en 1715. Montevideo, Fac. Human. Cienc., 82 pp.
- VAZ-FERREIRA, R., AND B. SIERRA DE SORIANO. 1960. Notas sobre reptiles del Uruguay. Rev. Fac. Human. Cienc. Montevideo, 18: 133–206.
- WERMUTH, H., AND R. MERTENS. 1961. Schildkroten, Krokodile, Bruckenechsen. Jena, Gustav Fischer, 422 pp.
- AND —. 1977. Liste der rezenten Amphibien und Reptilien: Testudines, Crocodylia, Rhynchocephalia. Tierreich, 100: 1–174.
- WIED, M. ZU. 1825. Beitrage zur Naturgeschichte von Brasilien. I. Band. Weimar, Landes, 614 pp.
- WILLIAMS, E. E. 1950. Variation and selection in the cervical central articulations of living turtles. Bull. Amer. Mus. Nat. Hist., 94: 510–561.

Kenyemys williamsi, a Fossil Pelomedusid Turtle from the Pliocene of Kenya

ROGER CONANT WOOD¹

ABSTRACT. *Kenyemys williamsi*, a new genus and species of pelomedusid turtle from the Pliocene of northwestern Kenya, is described. *Kenyemys* was a freshwater turtle whose relationships to other African pelomedusids cannot at present be determined.

INTRODUCTION

During the mid 1960's, a series of Harvard paleontological expeditions to the Turkana District of northwestern Kenya, led by the late Professor Bryan Patterson, recovered a large quantity of fossil chelonian material. The majority of these specimens represent two new types of pelomedusid turtles, one common and the other rare (Wood, 1971). This festschrift provides an appropriate occasion for formally naming the latter taxon in honor of Ernest E. Williams.

SYSTEMATICS

Order Testudines
Suborder Pleurodira
Family Pelomedusidae
Kenyemys gen. nov.
Figures 1-5

Type species. *K. williamsi* sp. nov.

Distribution. Pliocene, northwestern Kenya

Diagnosis. Differing from all other members of the family by the following combination of characters: 1) a series of elongate tuberosities forming an interrupted keel extending along the midline rearward from the dorsal surface of the second neural bone; 2) six neural bones forming a continuous series, the anterior end of the first abutting directly against the rear margin of the nuchal bone and the sixth one being heptagonal; 3) outer corners of nuchal bone extending beyond lateral margins of first vertebral scute; 4) pentagonal shape of first vertebral scute; 5) only eighth and posterior part of seventh pairs of pleural bones meeting at midline of carapace; 6) anterior plastral lobe truncated; 7) triangular intergular scute not overlapping anterior end of entoplastron and only partially separating the gular scutes along the midline axis of the plastron.

Etymology. The generic name refers to Kenya, the nation within whose borders this turtle was found.

Discussion. Two pelomedusid subfamilies, the Podocneminae and Pelomedusinae, can be recognized. However, their critical distinguishing characters are based primarily on skull morphology and cervical vertebra structure (Wood, 1971). Because these parts of the skeleton are not known for *Kenyemys williamsi*, its subfamilial allocation is not at present possible.

In the past, many African pelomedusids have been referred to the South American genus *Podocnemis*. Such iden-

¹Faculty of Science and Mathematics, Stockton State College, Pomona, New Jersey 08240, U.S.A.

tifications have generally been based on just a few shell characters (particularly the shape and arrangement of the neural bones as well as the presence of subcircular, laterally-placed mesoplastra) which, by themselves, are generally of limited taxonomic importance at the generic level.

Invariably, when better material becomes available or more thorough studies are carried out, these purported African representatives of *Podocnemis* prove referable to some other genus. Thus, "*Podocnemis*" *congolensis* is now *Taphrosphys* (Wood, 1975), "*Podocnemis*" *antiqua* is *Shweboemys* (Wood, 1970), and "*Podocnemis*" *madagascariensis*, whose origins surely lie within Africa on the basis of fossil material from the Miocene of central Africa that I currently have under study, is now generally accepted as representing a distinct genus, *Erymnochelys* (Pritchard, 1979). Moreover, it seems very possible that shell material from the Oligocene Fayum Depression of Egypt which has been described as *Podocnemis fajumensis* (including its synonym *P. blanckenhorni*; see Wood, 1971) probably represents the same taxon as the single skull which has been designated the type of *Dacquemys* (Williams, 1954).

The name *Podocnemis*, therefore, has been used in Africa much as it has been applied to other fossil pelomedusids elsewhere (see, for example, Wood and Gamaro's 1971 description of "*Podocnemis*" *venezuelensis*), as a catch-all taxon to which a variety of pelomedusids have been referred more for convenience than anything else. (Recently-collected additional material of "*Podocnemis*" *venezuelensis*, for that matter, clearly indicates that it is a new genus.) As the fossil record of pelomedusids continues to improve, it becomes increasingly apparent that there is no compelling reason to believe that the South American genus *Podocnemis* (*sensu strictu*) has ever reached other continents.

The suite of features which character-

izes the pelomedusid described here clearly precludes its reference to *Podocnemis*. Moreover, its shell is markedly different from that of other African pelomedusids including the African precursors of *Erymnochelys madagascariensis* (see Table 1).

On this basis, recognition of a new genus seems appropriate.

Kenyemys williamsi sp. nov.

Type. National Museum of Kenya (NMK) LT 127, a nearly complete but somewhat crushed shell.

Hypodigm. The type and NMK LT 128, the anterior half of a plastron.

Horizon and locality. Pliocene (Lothagam—1 [Patterson, Behrensmeyer, and Sill, 1970], now regarded as being somewhere in age between 5 and 6 million years B. P. [Maglio, 1974: Fig. 2]), Lothagam Hill, southwestern Turkana District, Kenya (approximately 2°53'N, 36°04'E).

Diagnosis. As for the genus.

Etymology. I take great pleasure in naming this species in honor of Ernest E. Williams, in recognition of his significant contributions to knowledge of both fossil and living turtles, not to mention herpetology in general.

Description. Essentially all of the shell of *Kenyemys williamsi* has been preserved. Despite some distortion resulting from cracking and crushing, it is apparent that the carapace was roughly oval in outline and probably moderately arched. Restorations of the shell are shown in Figures 3–5.

At the front of the carapace, the nuchal bone is hexagonal and slightly broader than long. There are six neurals arranged along the midline in an uninterrupted series. The fusiform first neural abuts squarely against the rear margin of the nuchal; posteriorly it is rounded to fit into a semicircular notch at the front of the second neural. Neurals two through five are of approximately the same size

TABLE 1. SHELL CHARACTERS OF KENYEMYS WILLIAMSII COMPARED WITH THOSE OF OTHER SOMEWHAT SIMILAR AFRICAN PELOMEDUSID TURTLES.

CHARACTER	<i>Kenyemys williamsii</i>	<i>Shweboemys antiqua</i>	<i>Stereogenys*</i>	<i>Taphrosiphys congolensis†</i>	" <i>Podocnemis</i> "†	<i>Erymnochelys§</i>	<i>Pelomedusa</i>
1) Keel present?	yes	no	no	no	no	no	no
2) Shape of last neural bone	heptagonal	pentagonal	pentagonal	pentagonal	pentagonal	pentagonal	somewhat variable, although usually pentagonal
3) total number of neural bones	6	6	7	7	6	6	7-8
4) does first neural bone abut directly against posterior end of nuchal bone?	yes	yes	no	yes	yes	yes	usually but not always
5) does nuchal bone extend beyond lateral margins of first vertebral scute?	yes	no	yes	no	no	no	no
6) shape of first vertebral scute	pentagonal	hexagonal	pentagonal	pentagonal	heptagonal	pentagonal	pentagonal
7) number of pairs of pleurals meeting at mid-line of carapace between last neural and suprapygals	1½	2½	½	1½	2½	2½	variable: ½-2
8) shape of carapace	oval	cordiform	oval	?	oval	oval	oval
9) shape of anterior plastral lobe	truncated	rounded	truncated	rounded	variable: rounded to truncated	rounded	rounded
10) shape of intergular scute	triangular	variable: triangular to pentagonal	pentagonal (<i>S. libyca</i>); heptagonal (<i>S. cromeri</i>)	heptagonal	truncated triangular (except pentagonal in <i>P. bramlyi</i>)	triangular	pentagonal

11) does intergular overlap onto anterior end of entoplastron?	no	yes	yes	variable: may or may not overlap	usually not; if so, only slightly	yes
12) does intergular separate either the gular or humeral scutes in midline?	anterior part of gulars only	gulars always completely separated, and anterior part of humerals sometimes	both gulars and humerals completely	variable: may separate only anterior part of gulars, all of gulars, or even some or all of the humerals as well	anterior part of gulars only	gulars always and part or all of the humerals as well
13) additional features	—	a) first vertebral scute reaches anterior margin of carapace, preventing first pair of marginals from meeting in midline b) posterior plastral lobe 1½ times longer than anterior lobe c) pleurals flat rather than curved, sloping away from blunt continuous midline ridge (not, however, a keel)	a) axial and inguinal buttresses weakly developed a) posterior plastral lobe roughly 2 times longer than anterior lobe b) low ridge on visceral surface of posterior plastral lobe running parallel to outer rim.	—	a) seventh pair of pleurals parallel-sided or even slightly constricted laterally rather than expanded	a) fenestration of the central portion of the plastron often persists in mature adults

*Two species have been described, *S. cromerti* and *S. libyca*, material of the latter being considerably better preserved than that of the former.

†Part of the data for this column is derived from previously unreported specimens whose description I am currently preparing.

‡I tentatively refer four species to this name of convenience: "p." *aegyptiaca*, "p." *bramlii*, "p." *fejmunensis*, and "p." *nodocnemoides*.

§This genus is at present monotypic and confined to Madagascar. However, fossil material from Africa's Central Rift Valley which I currently am studying is clearly referable to the genus. Hence, inclusion of this genus here seems appropriate.

and roughly hexagonal, having antero-lateral sides that are much shorter than the postero-lateral ones. The sixth neural is considerably longer than wide and instead of being pentagonal, as is typically the case for the last neural in podocnemine pelomedusids, it is heptagonal. In contrast to the common fossil pelomedusid from Lothagam, in which three pairs of pleurals are involved, separation between this terminal neural and the triangular suprapygal is effected only by the midline union of part of the seventh and all of the eighth pair of pleurals. Extending along the antero-posterior ridge of the carapace is a series of keels which become progressively more prominent towards the rear of the shell. The first of these protrudes only slightly from the surface of the second neural and the front half of the third. The second, arising from the posterior surface of the third neural and continuing back onto the front half of the fifth, has a higher crest than the preceding one, while the third and highest of the keels extends from the rear portion of the fifth neural back onto the eighth pair of pleurals. A median bulge on the suprapygal suggests that a fourth keel may also have been present. Carapacial keels are unusual features in pelomedusid turtles. Immature examples of *Podocnemis unifilis* and *P. sextuberculata* can have quite prominent tubercular keels (P. C. H. Pritchard, personal communication). Similar but less pronounced keeling is also characteristic of adults of various species of *Pelusios* (Loveridge, 1941; Pritchard, 1979).

Owing to damage at the posterior end of the carapace, it is not possible to determine the shape of the pygal, but it was probably rectangular as in all other pelomedusids. Nine peripheral bones remain attached to the carapace on either side and, in addition, two complete but disarticulated peripherals as well as several fragments have also been preserved, so that it appears likely that the actual complement of peripherals was the normal pelomedusid number of

eleven pairs. Flanking the neurals are eight pairs of pleurals which decrease in size from front to rear.

Except for the shape of the first vertebral, the scute pattern on the carapace of *Kenyemys williamsi* is not exceptional. The first vertebral was pentagonal, with its apex directed forward and its lateral sides more or less parallel. The lateral ends of the first pair of marginal scutes extended beyond the greatest width of the first vertebral. The nuchal bone at its greatest breadth extends beyond the lateral boundaries of the first vertebral.

The anterior plastral lobe of *Kenyemys williamsi* is truncated, with pronounced notches occurring at the junction between the gular and humeral scutes along the epiplastral margin to mark the position where the sides begin to curve laterally. The small, triangular intergular scute did not extend back onto the entoplastron in either of the two known specimens of *Kenyemys*. The plastral scute sulci and suture patterns conform to the typical pelomedusid configuration.

Most of the pelvis is preserved in the type but is somewhat crushed and distorted. From what can be seen, it appears to be similar to those of other podocnemine pelomedusids. The upper half of the acetabulum is formed by an excavation in the outer surface of the ilium, while the ischium and pubis each contribute about equally to the formation of the lower half of this socket. A low ridge on the visceral surface of the plastron, formed by projections running inward from the base of the ischia, serves to connect these elements at the midline.

Estimated measurements for the shell are: length of carapace, 32 cm; length of plastron, 28 cm. The plastral formula is: abdominal>pectoral>femoral>anal>gular>intergular>humeral.

DISCUSSION

Kenyemys was clearly a fresh water turtle. The sediments of the Lothagam-1

unit as well as the fauna recovered from it both indicate a mixture of channel, floodplain, and backwater depositional environments (Patterson, Behrensmeier, and Sill, 1970). Most of the Lothagam pelomedusids were recovered from sandstones, the type of *Kenyemys* being the only one to have been found in a rather limey clay. This might indicate that *Kenyemys* occupied a somewhat different kind of habitat than did the other, much more abundantly represented Lothagam pelomedusid. Whatever the case, it seems reasonable to suppose that these two species were broadly sympatric, as both pelomedusid taxa have been discovered at essentially the same stratigraphic level.

The relationship of *Kenyemys* to other African pelomedusids is at present obscure. Perhaps the most distinctive feature of its shell is the midline keel of the carapace, a character not present on any other adequately known podocnemine from the continent or adjacent island of Madagascar.

Rather tenuous evidence exists to suggest that there may have been another African fossil pelomedusid characterized by a keeled carapace. An isolated nuchal bone has been referred to *Latisternon microsulcae*, the type of which is a single left epiplastron from Olduvai Gorge, Tanzania (Auffenberg, 1981). This nuchal bone was found at a different site within Olduvai Gorge than the type specimen, so the association of the fragments is questionable. Whether or not the association is correct, the fact remains that a median keel is reported to rise from the dorsal surface of the nuchal. The available evidence is insufficient to determine either the extent of this keel or whether it was continuous or interrupted along its length. What little is known, does, however, clearly reveal differences in detail, since the keel of *Kenyemys* first appears farther back along the midline of the carapace. Moreover, the proportions of the nuchal bones in *Kenyemys* and *Latisternon* differ markedly, that of the

former being broader than long whereas that of the latter is considerably longer than broad. No reason exists, therefore, to believe that there is any close relationship between these two taxa based on their common possession of a carapacial keel.

With respect to another notable feature of the carapace of *Kenyemys*, only *Stereogenys libyca*, among other African podocnemines, also has a first vertebral scute whose lateral borders lie within the confines of the nuchal bone. But in other features (e.g., number, shape, and arrangement of neural bones, shape of anterior plastral lobe and patterns of scutes thereon) *S. libyca* differs markedly from *Kenyemys* and there is no reason to postulate a close relationship between these species either.

The truncated anterior plastral lobe of *Kenyemys* is superficially similar to those of several other African fossil pelomedusids (e.g., "*Podocnemis*" *podocnemoides* from the late Eocene of Egypt, "*Podocnemis*" *fajumensis* from the Oligocene of Egypt, "*Podocnemis*" *aegyptiaca* from the Miocene of Egypt, both species of *Stereogenys* and *Latisternon microsulcae* from the Pleistocene of Tanzania) but differs in detail. For example, the intergular scute of "*Podocnemis*" *podocnemoides*, while triangular, is relatively large and extends well back onto the surface of the entoplastron. Moreover, the gulars are small, triangular scutes confined to the epiplastra (Wood, 1971: Fig. 5). "*Podocnemis*" *fajumensis* and "*Podocnemis*" *aegyptiaca* have virtually identical anterior plastral lobes. In both, the triangular intergular scute apparently varied in size, in some cases being confined to the median portions of the epiplastra while in others extending slightly or even considerably back onto the entoplastron. Further, in all cases, the lateral extent of the anterior truncation is not proportionately so broad as in *Kenyemys* (Wood, 1971: Figs. 7, 8, 10). The lateral extent of the truncated part of the anterior lobe of *Latisternon* is also considerably

less than in *Kenyemys*. Moreover, *Lati-sternon*'s intergular was pentagonal and reached back onto the entoplastron, while its gulars were triangular and confined to the surface of the epiplastra. Both species of *Stereogenys* also had large intergulars, pentagonal in *S. libyca* and heptagonal in *S. cromeri*. Finally, for all of these turtles, other characteristics of the shell preclude the likelihood of any particularly close relationship.

In sum, no probable precursors of *Kenyemys williamsi* can as yet be identified, nor is it at present possible to determine whether this species gave rise to any descendants.

ACKNOWLEDGMENTS

I am indebted to the late Bryan Patterson, who not only made it possible for me to study the material described in this paper but who also provided me the opportunity to be a member of his paleontological expeditions to Kenya during the summers of 1965, 1966, and 1967. Ernest Williams deserves much of the credit (perhaps blame might be a more appropriate term) for encouraging my interest in turtles while I was a graduate student at Harvard; I thank him not only for his freely given advice but also for access to the resources of his herpetology department at the MCZ. The type specimen of *Kenyemys williamsi* was collected in the field during the summer of 1967 and then subsequently prepared in the laboratory by Arnie Lewis, whose work in both places is unsurpassed by anyone else I have ever met. The illustrative skills of Laszlo Meszoly are responsible for the shell reconstructions depicted in Figures 3-5. Peter Pritchard was kind enough to review a manuscript version of this paper.

Professor Patterson's 1967 expedition to Kenya was supported by National

Science Foundation (NSF) Grant GA-425. Several grants from the National Geographic Society enabled me to study relevant collections of fossil and recent turtles in Africa, Europe, and the United States. An NSF Science Faculty Professional Development Award for the academic year 1980-81 enabled me to prepare this paper for publication. During this period I was a visiting research scientist with the Florida Audubon Society, for whose hospitality I am most grateful.

LITERATURE CITED

- AUFFENBERG, W. 1981. The fossil turtles of Olduvai Gorge, Tanzania, Africa. *Copeia*, (3)1981: 509-522.
- LOVERIDGE, A. 1941. Revision of the African terrapin of the family Pelomedusidae. *Bull. Mus. Comp. Zool.*, (6)88: 467-524.
- MAGLIO, V. J. 1974. Late Tertiary fossil vertebrate successions in the northern Gregory rift, east Africa. *Ann. Geol. Surv. Egypt*, 4: 269-286.
- PATTERSON, B., A. K. BEHRENSMEYER, AND W. D. SILL. 1970. Geology and Fauna of a new Pliocene locality in northwestern Kenya. *Nature*, 226, nr. 5249: 918-921.
- PRITCHARD, P. C. H. 1979. *Encyclopedia of Turtles*. Neptune, New Jersey, T.F.H. Publications, 895 pp.
- WILLIAMS, E. E. 1954. New or redescribed pelomedusid skulls from the Tertiary of Africa and Asia (Testudines, Pelomedusidae), 1. *Dacquemys paleomorpha*, new genus, new species from the lower Oligocene of the Fayum, Egypt. *Breviora Mus. Comp. Zool.* No. 35, pp. 1-8.
- WOOD, R. C. 1970. A review of the fossil Pelomedusidae (Testudines, Pleurodira) of Asia. *Breviora Mus. Comp. Zool.* No. 357, pp. 1-24.
- . 1971. The fossil Pelomedusidae (Testudines, Pleurodira) of Africa. Ph.D. thesis, Harvard University.
- . 1975. Redescription of "*Bantuchelys*" *congolensis*, a fossil pelomedusid turtle from the Paleocene of Africa. *Rev. Zool. Africaine*, 89: 127-144.
- WOOD, R. C., AND M. L. DIAZ DE GAMERO. 1971. *Podocnemis venezuelensis*, a new fossil pelomedusid (Testudines, Pleurodira) from the Pliocene of Venezuela and a review of the history of *Podocnemis* in South America. *Breviora Mus. Comp. Zool.* No. 376, pp. 1-23.

Figure 1. Carapace of *Kenyemys williamsi* (type specimen, NMK LT 127).

Figure 2. Plastron of *Kenyemys williamsi* (type specimen, NMK LT 127).

Figure 3. Reconstruction of the carapace of *Kenyemys williamsi*.

Figure 4. Reconstruction of the plastron of *Kenyemys williamsi*.

Figure 5. Reconstruction of the shell of *Kenyemys williamsi* in lateral view.

Variation in Hispaniolian *Sphaerodactylus* (Sauria: Gekkonidae)

RICHARD THOMAS¹

ALBERT SCHWARTZ²

ABSTRACT. A review of the gekkonid species *Sphaerodactylus altavelensis* assesses variation and clarifies the nomenclature. Four subspecies (one described herein) are recognized. A second species, apparently related to *S. altavelensis*, is described as *S. williamsi*.

INTRODUCTION

The West Indian island of Hispaniola is the center of radiation of members of the *notatus* group of the gekkonid lizard genus *Sphaerodactylus*. The *difficilis* complex (of the *notatus* group) is the most widespread and diverse; this complex is named for *S. difficilis* Barbour, the first species in the complex to be described. The last revision of the complex was that of Shreve (1968); since that time we have named four new species (*S. cryphius*, *S. ocoae*, *S. nycteropus*, *S. zygaena*; Thomas and Schwartz, 1977; Schwartz and Thomas, 1977), and removed a subspecies of *S. difficilis* as a separate species (*S. randi*; Schwartz, 1977). This then is our fourth paper dealing with the Hispaniolan *difficilis* complex. There remain eight nominal species in the complex (Schwartz and Thomas, 1975); among these species we have made nomenclatural changes that require justification. The present paper justifies

these changes and assesses the variation in one of the remaining species, *S. altavelensis* Noble and Hassler and describes another. Our arrangement is due primarily to having collected and examined in detail many more specimens than were available to Shreve (1968) and to having visited many areas in Hispaniola whose *Sphaerodactylus* fauna had been either poorly known or altogether unknown. In the following accounts, measurements and counts in parentheses represent mean values.

TAXONOMIC ACCOUNTS

Sphaerodactylus altavelensis Noble and Hassler

Definition. A species of *Sphaerodactylus* with large acute, strongly keeled, flattened, imbricate dorsal scales, axilla to groin 20 to 36; no area of middorsal granules or granular scales; dorsal body scales with four to six hair-bearing scale organs around apex and distal edge, each with one hair. Dorsal scales of tail keeled, acute, imbricate, flat-lying; ventral scales of tail smooth, rounded, enlarged midventrally. Snout blunt, not depressed or decurved; snout scales moderate in size, rounded, keeled, juxtaposed, cobblelike to weakly raised on posterior edge, or weakly imbricate; 2 postnasals; 0 to 3 internasals; upper labials to mid-eye 3; dorsal head scales

¹Biology Department, University of Puerto Rico, Río Piedras, Puerto Rico 00931.

²Miami-Dade Community College, North Campus, Miami, Florida 33167, U.S.A.

small, rounded to subhexagonal, keeled, juxtaposed and cobblelike to weakly raised on posterior edge; temporal scales broad, rounded, keeled, raised, weakly imbricate; first infralabial subrectangular to subtriangular; gular scales between-infralabial rami rounded to hexagonal, moderately-sized, smooth to keeled, juxtaposed to weakly imbricate; central gulars small, acute, imbricate, smooth to keeled; some gular scales keeled in nearly all specimens; chest scales smooth; ventral scales smooth, rounded, imbricate, axilla to groin 22 to 32; scales around midbody 38 to 62; fourth toe lamellae 6 to 13; escutcheon with broad central area and extensions onto thighs to underside of knee area (3–8 × 11–27).

Color pattern in life weakly dichromatic; males with yellow throat ground color and reduced pattern, occasionally unicolor. Pattern of paired light areas on upper surface of head, followed by second pair on occiput and neck; dark scapular patch and scapular ocelli surrounded by continuous light border concave anteriorly and convex posteriorly; dorsal body pattern of dark spotting or lines or combination of both; venter pale with dark lines; throat with dark lines or mottling; sacral pattern of dorsolateral light and dark lines; and tail with dark middorsal zone and series of irregular ocelli. We have examined 847 *S. alta-velensis*.

Basic pattern. In order to facilitate descriptions of variation in the subspecies of *S. alta-velensis*, we will describe in some detail here the basic pattern for the species. The fullest pattern development is found in some females of two of the subspecies. For convenience of discussion the basic pattern is broken down into the following elements:

Anterior cephalic figure. A posteriorly directed, bilobed, light figure outlined in dark pigment extending to about the level of the ear opening; the breadth covers most of the dorsal surface of the head; the lines forming the lateral edges originate as superior postocular stripes.

Posterior cephalic figure. A bipartite, dark-edged figure in the occipitonuchal area having posteriorly directed lobes, the anterior margins being formed by the posterior edges of the anterior cephalic figure.

Scapular figure. A complex figure consisting of 1) a black, roughly rectangular scapular patch enclosing; 2) a pair of pale ocelli; and 3) a narrow light border of greater extent than the patch (i.e., including an area of ground color peripheral to the patch), in roughly the shape of an inverse concave-convex trapezoid, and formed of a dark-edged light line.

Body pattern. About ten longitudinal dark lines occur on the dorsum (3) and sides (2) and venter (about 5); the dorsal series extends from the scapular figure to the sacral region; the lateral lines extend from the post-axillary region to the groin, and the ventral lines, which are faintest, extend irregularly from about midbody to the groin and vent. The lineate pattern is rarely as fully expressed as described, but many females show indications of it. Often, however, the dorsal body pattern is one of small dark spots about one to three scales in area, irregularly distributed. The spotted pattern seems to be derived from fragmentation of the linear one, in which even those individuals with spotted patterns have lines in the sacral region and on the venter.

Sacral figure. A pair of dorsolateral stripes on each side form the dark edges of a light dorsolateral sacral zone, which extends beyond the sacral region and fades on the posterior body and base of the tail.

Tail pattern. Six very dark longitudinal lines around the circumference form a lineate pattern: a dorsolateral pair of lines which is invaded by several pairs of pale ocelli, a lateral pair of lines, and a ventrolateral pair; the spaces between the dorsolateral and lateral, and the lateral and ventrolateral lines are very light, forming a contrastingly striped lateral tail pattern. The lateral and ventrolateral stripes with their pale inter-

spaces continue anteriorly onto the posterior surface of the thigh to form a prominent zonation there.

Male pattern. Males lose the basic pattern as they mature and assume either a nearly uniform brown or a uniformly spotted pattern. Faint remnants of the basic pattern may persist, and in a few individuals (and in one subspecies) apparently mature males retain rather fully developed basic patterns. The scapular patch is present in males of three subspecies (see below). Some males show an intensification of the head markings in which the edges of the markings become darkened and intensified with eventual loss of the lighter areas resulting in a vermiculate pattern.

Distribution. Isla Alto Velo; on the

mainland of Hispaniola, from the Llanos de Azua in the east, west continuously to the Baie de Port-au-Prince in xeric to semi-xeric situations, along the north coast of the basal Tiburon Peninsula to the vicinity of Petit-Goâve and south through the Vallée de Trouin (presumptive) to the vicinity of Cayes Jacmel on the south coast. A population occurs in the vicinity of Jérémie on the distal Tiburon Peninsula, and another on tiny Ile à Cabrit in the Golfe de la Gonâve (Schwartz, 1979). On the north island, *Sphaerodactylus altavelensis* is known from the coast north of Port-au-Prince to Pierre Payen and inland in the vicinity of Lascahobas and in the northwest from Gonaïves to Port-de-Paix and Ennery in Haiti (Fig. 1).

Figure 1. Map of western and central Hispaniola, showing the known distributions of *S. altavelensis* (●) and *S. williamsi* (▲). Subspecies of *S. altavelensis* are noted and their ranges shaded.

Sphaerodactylus altavelensis altavelensis
Noble and Hassler
Figures 2A, 3A

Sphaerodactylus altavelensis Noble and Hassler, 1933. Amer. Mus. Novitates, 652: 7.

Type locality. Isla Alto Velo, República Dominicana.

Holotype. AMNH 51488.

Definition. An insular subspecies of *Sphaerodactylus altavelensis* characterized by large size, high midbody scale counts (44–62), and very bold contrasting basic pattern (females).

Distribution. Isla Alto Velo, República Dominicana.

Variation. We have examined 22 specimens. Largest male (AMNH 51490) 28 mm snout-vent length (SVL), largest female (MCZ 45947) 29 mm SVL; dorsal scales between axilla and groin 26 to 36 (30.1); ventral scales axilla to groin 22 to 32 (27.9); scales around midbody 44 to 62 (51.7); supralabials to mid-eye 3/3; internasals 1 (N=21), 3 (N=1); fourth toe lamellae 8 to 11 (mode 10); throat scales keeled; escutcheon 3 to 6 × 22 to 27.

Males have a uniform grayish ground color and a nearly uniform scattering of dark brown spots over the dorsal surface of the head and body; the scapular patch and ocelli are absent or reduced to minute ocelli connected by a small scapular patch that does not include the ocelli. Most females show a bold development of the basic pattern in which the anterior and posterior cephalic figures are heavily outlined in dark pigment; the scapular patch is large and encloses the pair of ocelli, and the light scapular figure border is markedly concave-convex producing a comic-mask mouth appearance (Fig. 3A). The dorsal body pattern of females is heavily mottled or vermiculate with a tendency towards formation of transverse dark markings reaching an extreme in AMNH 51478. Some females show a simplification and fragmentation of pattern that ap-

proaches the spotted male pattern; the anterior and posterior cephalic figures are fragmented; the scapular patch and ocelli are reduced; and the scapular figure border is absent. Lateral and sacral lines are prominent in consonance with the boldness of the rest of the pattern; ventral lines are present as in the other populations of the species.

Remarks. Shreve (1968) regarded *S. altavelensis* as a species confined to Isla Alto Velo and suggested that it was related to *S. difficilis randi* (= *S. randi*; Schwartz, 1977). The bold scapular figure and dorsal spotting of some *randi* populations do bear a resemblance to *altavelensis*. However, the trilineate head pattern of *randi* plus the presence of a completely expressed basic pattern such as is found in Shreve's *S. brevirostratus* (and the lack of these pattern elements in *randi*) leave no question of the relationship of the Alto Velo sphaerodactyls to *brevirostratus*. In fact, the scapular pattern of *S. randi* only superficially resembles that of *S. altavelensis*. Shreve examined only the two MCZ paratypes of *altavelensis*, in which the pattern is not so well expressed as in some of the other specimens.

Sphaerodactylus a. altavelensis is separated from other populations of the species by not only a water gap but also, as far as is known, by a significant geographic hiatus that includes all of the Peninsula de Barahona. The nearest records are the region just to the south of the city of Barahona on the east, and the region of Cayes Jacmel to the west. Although the species may occur farther east along the south coast of Haiti than Cayes Jacmel, it apparently does not extend to the region of Pedernales on the Dominican-Haitian border—an area that has been relatively well-collected and from which other species of *Sphaerodactylus* are known. Another similarly relict member of the Alto Velo fauna is *Leiocephalus vinculum altavelensis* Noble and Hassler (Schwartz, 1967).

Figure 2. Dorsal views of subspecies of *S. altavelensis* and *S. williamsi* as follow: A. *S. a. altavelensis* (AMNH 51485). B. *S. a. enriqueilloensis* (ASFS V23288). C. *S. a. breviostratus* (ASFS X2376). D. *S. a. luciai* (MCZ 156208—holotype). E. *S. williamsi* (MCZ 156209—holotype).

Figure 3. Diagrammatic dorsal views of cephalic and scapular patterns of *S. altavelensis* (see text for discussions of variation), as follow: A. *S. a. altavelensis*. B. *S. a. enriquilloensis*, C. *S. a. brevisrostratus*. D. *S. a. lucioi*.

Specimens examined. REPÚBLICA DOMINICANA: *Isla Alto Velo*: AMNH 51472–86, AMNH 51489–91, MCZ 45946–47, AMNH 51488, ASFS V26910.

Sphaerodactylus altavelensis enriquilloensis Shreve

Figures 2B, 3B

Sphaerodactylus brevisrostratus enriquilloensis Shreve, 1968, *Breviora*, 280:14.

Type locality. 4 km E La Descubierta, near Lago Enriquillo, Independencia Province, República Dominicana.

Holotype. MCZ 57846.

Definition. A subspecies of *Sphaerodactylus altavelensis* characterized by lack or very weak development of the scapular patch and ocelli.

Distribution. The Valle de Neiba east of the Dominico-Haitian border, south around the eastern edge of the Sierra de Baoruco to slightly beyond the city of Barahona, eastward to the Llanos de Azua (17 km E Azua), south to Punta Martín García and north at least to the Azua-San Juan province border (southeast of Guanito). A specimen from near Vallejuelo within the Sierra de Neiba is tentatively referred to this subspecies and may indicate a more widespread

occurrence of the species in the Valle de San Juan and associated areas.

Variation. We have examined 408 specimens and have taken full or partial counts on 314 of these. Largest males 26 mm, largest female 28 mm; dorsal scales axilla to groin 23 to 31 (26.1); ventral scales axilla to groin 23 to 32 (28.0); scales around midbody 38 to 50 (46.0); supralabials to mid-eye usually 3/3, rarely 3/4 or 4/4; internasals 1 (N=205), 2 (N=28), 3 (N=1); fourth toe lamellae 7 to 11 (mode 9 and 10); throat scales keeled; escutcheon 4 to 8 × 11 to 27.

The anterior and posterior cephalic figures are absent in virtually all males, although some show a residual trace of them. Scapular patches and ocelli are present (not necessarily in the same specimen) in scattered males (the frequency is less than it is for the same pattern elements in females). Ocelli in males are very reduced, often to the size of a single scale; and the patch, if present, may be no more than a small area of black pigment between the two ocelli.

The anterior cephalic figure is well developed in females; the posterior figure is not so strongly developed and is frequently represented by only a pair of

light crescents in the occipital region demarcating the posterior end of the posterior figure (Fig. 3B). Eight of 86 Valle de Neiba females and juveniles have posterior extensions of the anterior cephalic figure from roughly the middle of each lobe, which may fuse with the posterior figure or fade out in the parietal region. Although absent or very reduced, scapular ocelli and patches are present in a small proportion of all specimens throughout the range of the subspecies. For example, 17 of 86 Valle de Neiba females and juveniles have discernible ocelli; among these, some lack scapular patches of any noticeable extent, and others have them weakly developed, two having what might be termed moderate development of the patch. Eight of the ocellate females from the Valle de Neiba (28 females) occur in the Las Clavellinas series. None of the Duvergé vicinity specimens has ocelli, and only one of the females from Mella (28 females from both localities combined) has ocelli. Thus the frequency of scapular ocelli seems lower in the females from the southern side of the Valle de Neiba, although the frequency of scapular ocelli, even though very reduced in expression, is as high or higher (15 of 50) in the Barahona-and-vicinity sample as in the Valle de Neiba one. In the long series (159 specimens) from the northeastern part of Barahona Province, indications of ocelli are found on but two females (although six males have faint small ocelli). The scapular figure border is complete or nearly complete in 22 of 86 Valle de Neiba females and juveniles; others have some indication of this feature, most frequently the concave anterior border. Of the five (of 50) females from the Barahona region having indications of the scapular figure border, it is complete in only one. Approximately the same situation holds for the more eastern population of *S. a. enriquilloensis*: indications of the scapular figure border are seen, but in no specimens are they complete. Dorsal body coloration in the subspecies varies from

nearly unicolor to heavily spotted or vermiculate with dark brown; dorso-lateral and lateral stripes extend the length, or nearly the length, of the body in some specimens. Ventral lines are present to a variable extent in nearly all specimens.

The 13 Punta Martín García specimens are noteworthy for their virtual lack of any pattern development in males, females, and juveniles. Two show ghost remnants of the posterior cephalic figure and scapular figure border, and others show brief ghost traces of the posterior cephalic figure; spotting and flecking is very reduced, although ventral lines are present.

The single specimen from 7 km NW Vallejuelo (between the north and south ranges of the Sierra de Neiba) clearly shows the anterior cephalic figure, traces of the posterior figure, and the anterior and posterior edges of the scapular figure border; there is a faint smudge that may represent the scapular patch and a small light marking that gives the appearance of a diverticulum of the anterior edge of the scapular figure border, but which may be a scapular ocellus. The dorsum is mottled with irregular dark markings tending to form transverse light bars in the middorsal zone between indications of dorsolateral lines (a pattern sometimes seen in other *altavelensis*); ventral lines are present but faint. We tentatively assign this specimen to the subspecies *enriquilloensis*, but the final allocation of the population represented by it must await the acquisition of more specimens.

Specimens examined. REPÚBLICA DOMINICANA: *Provincia de Independencia*: Boca de Cachón, ASFS V4381-83, ASFS V39707; 2 km E La Descubierta, ASFS V45007-08; 5 km N Jimaní, ASFS V35480-502; 10.4 km NE Jimaní, ASFS X9488-501; 5 km SE La Florida, ASFS V23303-08; 9 km W Duvergé, ASFS V20508; 7.0 km W Duvergé, ASFS V4401-06; 6.7 km NW Duvergé, ASFS V4343-47; 6 km NW Duvergé, ASFS V17170-71; La Source, about 5 km W Duvergé, ASFS V20803-06; just E Duvergé, ASFS V23276-302; outskirts of Mella, ASFS V30671-89; 8 km N Colonia Mixta, ASFS V35508; 3 km SE Angostura, ASFS V41761-64; 11 km SE

Angostura, ASFS V41299-307; *Provincia de Barahona*: 1.4 km WNW Las Clavellinas, ASFS V30377-85; Jaragua, ASFS X9787; 5.4 km ENE Neiba, ASFS V228-43; 5.9 km ENE Neiba, ASFS V30334-70; 4 km SW Galván, ASFS V40753-57; *Provincia de Barahona*: 11 km NE Cabral, ASFS V35509-10; 7.5 km E Cabral, ASFS X9623-25, ASFS V219-20, ASFS V350, ASFS V1450; 5 km S Cabral, MCZ 51820-21; 5 km N Barahona, ASFS V20515-37; Barahona, ASFS X9451, ASFS X9525; Barahona, southern outskirts, ASFS V30988-1011, ASFS V31013; 2 km SE Barahona, ASFS X9520; 4.8 km S Barahona, ASFS V14058, ASFS V14060, LDO 7-5344-50, LDO 7-5352-54, LDO 7-5361-63, LDO 7-5365-66, LDO 7-5522; 0.6 km NW Palo Alto, 31 m, ASFS V30543-72; 4.2 km NW Palo Alto, 31 m, ASFS V30530-40, ASFS V31515; Fondo Negro, ASFS X9708; 1 km NE Fondo Negro, ASFS V35667-78; El Iguito, 2.6 km NE Fondo Negro, ASFS V30495-522, ASFS V31429; 3.2 km NE Fondo Negro, ASFS X9678-81; 1.1 km NW Puerto Alejandro, ASFS V30637-47; west side, Punta Martín García, ASFS V117-29; *Provincia de Azua*: Barreras, ASFS V31176-78; 7.4 km NE Barreras, ASFS V31035-36; 19.8 km SE Guanito, 275 m, ASFS V31335-37; 15.2 km E Azua, ASFS X8095-101, ASFS V19352-54; 17 km E Azua, ASFS V21101; *Provincia de San Juan*: 7 km NW Vallejuelo, 793 m, ASFS V303.

Sphaerodactylus altavelensis brevirostratus Shreve

Figures 2C, 3C

Sphaerodactylus brevirostratus brevirostratus Shreve, 1968, *Breviora* 280:10.

Type locality. 5 km S Dufort, south of Léogâne, Département de l'Ouest, Haiti.

Holotype. MCZ 63234.

Definition. A subspecies of *S. altavelensis* characterized by well-developed anterior and posterior cephalic figures in females; scapular figure present as a black patch and ocelli in all females and the majority of males; scapular figure border present in all individuals but incomplete in most.

Distribution. Haiti, from the region of Lascahobas and St. Marc on the north island, south to the Cul de Sac Plain onto the north slopes of the Massif de la Selle (Morne l'Hôpital) in the vicinity of Pétionville (and apparently as high as Furcy) and west along the base of the Tiburon Peninsula at least to between Petit-Goâve and Grand-Goâve, and south to the south coast of the peninsula to the

vicinity of Cayes Jacmel. A population at Jérémie is assigned to this subspecies; it averages lower in scale counts but agrees in coloration. Although collecting on the Tiburon Peninsula has failed to reveal any *altavelensis* between Jacmel and the Petit-Goâve records and those from Jérémie, we would not be surprised at the eventual discovery of populations in this area.

Variation. We have examined 394 specimens and taken counts on 103 of these. Largest males 25 mm SVL, largest female 28 mm dorsal scales between axilla and groin 20 to 29 (24.7); ventral scales axilla to groin 22 to 32 (26.2); mid-body scales 38 to 48 (42.9); supralabials to mid-eye 3/3 (85), 3/4 (2), 4/4 (1); internasals 0 (3), 1 (83), 2 (7); fourth toe lamellae 5 to 11 (mode 9 but 10 nearly equimodal); throat scales keeled; es-cutcheon 4 to 8 × 11 to 24.

Males are gray to gray-brown or brown and may or may not be spotted or flecked with dark brown; heads may be unicolor in combination with a spotted dorsal body pattern or may have contrasting vermiculate patterns (few specimens); the scapular patch is reduced compared with that of females and may be nearly pin-head in size or absent. The anterior cephalic figure is weakly developed in most males; the posterior cephalic figure is very weakly indicated or absent with the exception of the males having well-developed basic patterns (Fig. 2C).

Most females have well-developed anterior cephalic figures and posterior figures (Fig. 3C) but show considerable variation in expression. All females have at least some indication of a scapular patch and ocelli; the extent of the patch varies from reduced (about four scales in width by three to four scales in length at greatest length) to fully developed. Fully developed patches may be 12 scales wide and six scales in length. In the reduced form the ocelli occur at the lateral edges of the patch and may not be completely included in it.

In life, the ground color of *S. a. brev-*

irostratus is tan to brown with dark brown markings. Anterior and posterior cephalic figures are paler (central portion) than the rest of the ground color. Scapular ocelli are white or whitish; venters are gray with brown lines. Males have yellow or yellowish heads and throats.

Specimens examined. HAITI. *Dépt. de l'Artibonite:* Pierre Payen, 14.4 km S St. Marc, ASFS V39457, ASFS V43777; *Dépt. de l'Ouest:* 3.5 km SW Lascabobas, 275 m, ASFS V26539, ASFS V26592; 10.1 km SE Montrouis, ASFS V39461-63; Trou Forban, USNM 117157-58, USNM 118894; 3.5 km SE Trou Forban, ASFS X3994-99; Ile-à-Cabrit, ASFS V40441-43; Sources Puantes, USNM 117159-64; Tabarre, 5.1 km SE François Duvalier airport, ASFS V36220; 20.2 km SE Mirebalais, 366 m, ASFS V35702; 6.2 km NW Ganthier, ASFS X2088-100, LDO 7-5799-800, LDO 7-5825-39, LDO 7-5842-43, LDO 7-5853-54, LDO 7-5801; 5.6 km E Croix des Bouquets, ASFS X2101; 12.6 km E Croix des Bouquets, ASFS V36906; 13.1 km E Croix des Bouquets, ASFS V42563-64, ASFS V8139-42, ASFS V8298-303; 13.4 km E Croix des Bouquets, ASFS V40464, ASFS 44852-55; Tête Source, 1.4 km NNE Thomazeau, ASFS V8177-78; Source Fond Parisien, ASFS V36974-77; Pétienville, ASFS V13688; Morne Calvaire, 1.6 km SW Pétienville, ASFS X1302; Furey, AMNH 124158; Diquni, ASFS X2375-79; Mariani, 11.2 km E Gressier, ASFS V8462; 6.4 km SW Léogâne, ASFS V8316; 1.6 km SE Fauché, ASFS V37270-73; $\pm$ 6.4 km SE Fauché, ASFS V37267; 9.9 km E Fauché, ASFS X2051; beach just W Grand-Goâve, ASFS V36569-71, ASFS V42557-61; 8.8 km W Grand-Goâve, ASFS V45218-346; 4.0 km E Petit-Goâve, ASFS V43311-16, ASFS V43605-711; 7.5 km E Petit-Goâve, ASFS V43317, ASFS V43586-89, ASFS V44073-83; 1.6 km N Jacmel, ASFS V9798; La Fond, near Jacmel, MCZ 64822; less than 1.6 km W Cayes Jacmel, ASFS V9704-09; *Dépt. du Sud:* beach area within 1 km E Jérémie, ASFS V25217-29; ca. 5 km SE Jérémie, ASFS V9392; Pistache, SW Jérémie, ASFS V9126-28.

Sphaerodactylus altavelensis lucioi new subspecies

Figures 2D, 3D

Holotype. MCZ 156208, adult male, from Terre Sonnain, 1.6 km N Les Poteux, 132 m, Département de l'Artibonite, Haiti, one of a series collected 9 July 1978 by native collectors. Original number ASFS V46397.

Paratypes. ASFS V46398-404, same data as holotype; ASFS V40263-65, same locality as holotype, 9 August 1977, native collectors; ASFS V49900-01,

Balladé, 8.8 km S Port-de-Paix, *Dépt. du Nord-Ouest*, Haiti, 18-19 July 1979, native collectors; ASFS V46984-86, Balladé, 8.8 km S Port-de-Paix, *Dépt. du Nord-Ouest*, Haiti, 22 July 1978, native collectors; ASFS V47797-98, 1.9 km W Ennery, 336 m, *Dépt. de l'Artibonite*, Haiti, 5 August 1978, native collectors; ASFS V50141-43, 1.9 km W Ennery, 336 m, *Dept. de l'Artibonite*, Haiti, 22 July 1979, native collectors; ASFS V46664-66, Gonaïves, *Dépt. de l'Artibonite*, Haiti, 12 July 1978, native collector.

Definition. A subspecies of *Sphaerodactylus altavelensis* characterized by a prominent anterior cephalic figure in both sexes, a very obscure to absent posterior cephalic figure in both sexes (Fig. 3D), and scapular figure present as a black patch and usually two white ocelli in almost all specimens of both sexes, the patch without edging.

Distribution. Northeastern Haiti, from the vicinity of Port-de-Paix in the north to Gonaïves in the south, and inland as far as the vicinity of Ennery.

Description of holotype. Adult male; 23 mm SVL, tail length 21 mm, tip regenerated; dorsal scales axilla to groin 29, ventral scales axilla to groin 27, scales around midbody 41; fourth toe lamellae 9; escutcheon 5×22 . Snout moderately blunt in dorsal aspect, very slightly depressed; snout scales moderately-sized, somewhat flattened, smooth, not imbricate; 1 internasal; 2 postnasals; supralabials to mid-eye 3/3; dorsal head scales small, elongate, keeled, almost imbricate to juxtaposed; temporal scales small, oval, keeled, subimbricate. First infralabial broader anteriorly than posteriorly, roughly rectangular in shape; gular series between infralabial rami large, juxtaposed, smooth; central gulars small, smooth, subimbricate; chest and ventral scales smooth, with about 1 lateral row of throat scales on each side keeled.

Dorsal ground color in life dark brown, dorsum finely peppered with slightly darker (grayish brown) scales; anterior cephalic figure buff, outlined with very dark brown to black, bilobed and prominent; posterior cephalic figure reduced to a series of dark flecks or spots; scapular

patch present and prominent, not bordered, black with a pair of included white ocelli, and constricted medially; tail vaguely lineate dorsally and without obvious ocellate pattern; venter creamy white, with chin and throat rather densely dotted with dark brown; a series of about four or five fragmented and irregular dark longitudinal lines on venter.

Variation. Of the 23 specimens, the largest male is 24 mm SVL, largest females 24 mm SVL; dorsal scales axilla to groin 22 to 29 (27.1), ventral scales axilla to groin 24 to 29 (26.7); midbody scales 37 to 47 (42.5); supralabials to mid-eye 3/3; internasal 1; fourth toe lamellae 8 to 13; throat scales smooth in all but three specimens, although a few lateral throat scales in lateral sequence may be keeled, whereas all other throat scales on the same specimen are smooth.

The dorsal ground color in all specimens was brown, dark brown, or almost black. Even darker dorsal body pattern elements are usually more or less isolated dark scales which give a dotted pattern. These dots may at times be aligned into a very vague series of dorsal lines. The anterior cephalic figure is bilobed and always prominent, whereas the posterior cephalic figure is absent and represented merely by a series of dark fragments. The black scapular patch is present but variable in extent in all specimens but one, although it was recorded as present in life on that specimen. Two white ocelli are present and may be included in the dark patch or occasionally lie just peripheral to it. Unregenerated tails continue the more or less lineate dorsal pattern, although the dorsal surface of the tail may have merely a random series of dark scales without any lineate effect. Sacral stripes are present in some specimens, absent in others. Throats are dotted with dark brown and venters either rather lineate with dark or not.

Females are colored and patterned dorsally like males; the anterior cephalic figure is prominent and the scapular patch is variable in expression, with

ocelli present in all specimens. Juveniles are patterned like females. Venters are like those of males except that they seem slightly more strongly lineate and there is a clearer indication of ventrolateral longitudinal stripes than in males (Fig. 3D).

Comparisons. *Sphaerodactylus a. lucioi* differs from the other subspecies of *S. altavelensis* in smaller size (although it is closest to *brevirostratus*), in having the black scapular patch and ocelli present in both sexes (the patch without a clear outline edging) and in the expression of the anterior cephalic figure and absence of the posterior figure in both sexes. The other subspecies usually have the throat scales keeled, whereas *S. a. lucioi* usually has them smooth, with at best a few lateral keeled scales. The mean of dorsal scales (27.1) is greater than those of *enriquilloensis* and *brevirostratus* (26.1 and 24.7), less than that of *altavelensis* (30.1). The mean of midbody scales is less (42.5) than that of any other subspecies (42.9 to 51.7), and that of ventral scales (26.7) is intermediate between that of *brevirostratus* (26.2) and *altavelensis* and *enriquilloensis* (27.9 and 28.0).

Remarks. We have no data on the precise niche which *S. a. lucioi* occupies. The general habitats are quite variable. The type locality is xeric, with cacti and *Agave* as typical plants, on a gravelly and rocky substrate. The specimens from Gonaïves presumably came from the vicinity of the city itself; Gonaïves lies in a xeric region. Extending inland from Gonaïves is the valley of the Rivière d'Ennery. The Ennery locality for the specimens of *S. a. lucioi* is mesic and at an elevation of 336 m; this locality is especially interesting in that there is a combination of a mesic interior and xeric exterior (= coastal) faunas there. The locality at Balladé is extremely mesic, in fact a coffee-cacao grove. Syntopic congeners of *S. a. lucioi* vary with locality: at the type-locality they are found with *S. cinereus* Wagler *sensu* Graham and Schwartz, 1978 (common) and *S. asterulus* Schwartz and Graham (rare). At

Ennery they were also taken with *S. difficilis* Barbour and *S. cinereus* in about equal frequency. Balladé yielded only *S. difficilis* in addition to *S. a. lucioi*. At any locality, the numbers of specimens of other species far outnumbered those of *S. a. lucioi*. Whether this is due to its scarcity or to its relatively smaller size is unknown; collecting was presumably non-selective.

We are pleased to name this subspecies in honor of John C. Lucio, who accompanied the junior author in Haiti for two successive summers (1978–79) and who contributed greatly to the success of both trips.

Sphaerodactylus williamsi new species
Figure 2E

Holotype. MCZ 156209, adult female, from 12.2 km W Ça Soleil, Département de l'Artibonite, Haiti, taken by native collector on 14 July 1978. Original number ASFS V46794.

Definition. An apparently small species of *Sphaerodactylus* (only known specimen 22 mm SVL), with small scales (33 keeled dorsal scales and 33 smooth ventral scales between axilla and groin, 52 scales around midbody), keeled throat scales; no area of middorsal granules or granular scales; dorsal body scales with three or four hair-bearing scale organs, each with one hair, on the posterior periphery of the scale, one organ at the apex; no head or scapular pattern or patch or ocelli, but dorsum with a series of four well-defined pale buffy lines alternating with broader dark brown lines, plus two pale dorsolateral lines; venter pale, throat with a few fine dark brown stipples and very vague indication of ventral longitudinal lines, most prominent posteriorly; head small and relatively narrow.

Distribution. Known only from the type locality (Fig. 1).

Description of holotype. Gravid female; 22 mm SVL and 17 mm tail length (mostly regenerated); dorsal scales

keeled and imbricate, small, axilla to groin 33, ventral scales axilla to groin 33, scales around midbody 52, fourth toe lamellae 10. Snout attenuate, head small and relatively narrow; snout scales moderately large, juxtaposed, smooth anteriorly and keeled posteriorly; 1 internasal; 2 postnasals; supralabials to mid-eye 3/3; dorsal head scales small, keeled, subimbricate; temporal scales small, juxtaposed, weakly keeled. First infralabial broader anteriorly than posteriorly, rectangular; gular series between rami large, smooth, juxtaposed; central gulars very small, smooth, imbricate, grading quickly to much larger keeled throat scales; chest and ventral scales smooth.

Dorsal ground color in life dark brown with four longitudinal pale buffy lines (Fig. 2E); head and neck scales without cephalic or nuchal pattern but with isolated scattered buffy scales; a pair of slightly more prominent pale buffy lines on the sides; venter creamy white, with a few scattered dark brown dots on the throat and vague indications on the posterior portion of the belly of dark longitudinal lines; no scapular patch or ocelli.

Comparisons. *Sphaerodactylus williamsi* seems most closely related to *S. altavelensis*; the basic dorsal color (dark brown) is similar in *S. williamsi* and *S. a. lucioi*. But direct comparison of the single *williamsi* with many comparably sized female *altavelensis* of any subspecies shows the much narrower and more acuminate snout of the former. Although some *S. altavelensis* are vaguely lineate, none shows the distinct lineation of *S. williamsi*. The total absence of cephalic patterns (except for the scattered pale buffy scales) and of a scapular patch and ocelli are also distinctive, especially in contrast to local *S. a. lucioi* which, in both sexes, have prominent anterior cephalic patterns and scapular patches and ocelli in both sexes. The high counts of the holotype of *S. williamsi* (dorsals 33, ventrals 33, midbody 52) are all greater than the highest

counts in local *S. a. lucioi* (29, 29, 47); the keeled throat scales of *S. williamsi* likewise are distinctive. The scale counts fall within (dorsals, midbody) or just above (ventrals) those of *S. altavelensis*; but it is the far-removed subspecies *S. a. altavelensis*, not the closer *brevirostratus*, *enriquilloensis*, or *lucioi*, whose upper parameters exceed or almost equal the counts of *S. williamsi*. The smaller scales of *S. williamsi* are obvious upon inspection and comparison with similarly sized *S. a. lucioi*, with which *S. williamsi* might be confused.

Remarks. The type locality of *S. williamsi* is a coastal oasis; the general region is called Lapierre. The surrounding area is xeric and barren (cacti, *Agave*), in strong contrast to the luxuriant growth in the oasis which is fed by a small source. Associated congeners here are *S. asterulus* and a second, as yet unnamed, species. Both these congeners are presumably more abundant in the adjacent xeric areas than within the oasis; this is certainly true for *S. asterulus*. Haitians living within the oasis are relatively independent and are not easily encouraged to collect; this may account for the lone specimen of *S. williamsi* (and perhaps even for the lack of *S. altavelensis*, which occurs only 19 km to the east at the type-locality of *lucioi* and 10 km to the south east at Gonaïves, across the Baie de Gonaïves). On the other hand, it is possible that *S. williamsi* is here at the very periphery of its range, which may encompass the herpetologically largely unknown Presqu'île du Nord-Ouest. From that peninsula proper, *S. shrevei* Lazell and *S. elegans* MacLeay are known (both from Môle St. Nicholas, far to the northwest and on the northern side of the Massif du Nord-Ouest). We call attention to this phenomenon once more: the line of demarcation between the peninsular fauna and the "mainland" fauna appears to be sharp. In the matter of *Sphaerodactylus*, *S. cinereus* and *S. altavelensis* are moderately common on the "mainland," but appear not to occur

on the peninsula. *Sphaerodactylus asterulus* is peninsular but a southern cognate of the related *S. shrevei* in the north. *Sphaerodactylus difficilis* occurs only along the northern slopes of the Massif du Nord-Ouest, as far west as Bombardopolis and also on the adjacent "mainland." The ecological relationships among this suite of congeners are indeed puzzling, and their geographical relationships are just beginning to be clarified.

We take very great pleasure in naming this new species in honor of Ernest E. Williams, who, over the years, has been a constant source of friendship, advice, cooperation, and loans of pertinent materials for our research in the Antilles.

ACKNOWLEDGMENTS

Most specimens examined for the present paper are in the Albert Schwartz Field Series (ASFS); many of them were collected under the sponsorship of National Science Foundation Grants G-7977 and B-02603 to the junior author. We thank the following field workers, both friends and students, who have helped make such a mass of material available to us: Patricia A. Adams, Robert K. Bobilin, Jeff R. Buffett, David A. Daniels, James R. Dennis, Danny C. Fowler, Eugene D. Graham, Jr., Ronald F. Klinikowski, Mark D. Lavrich, David C. Leber, James K. Lewis, John C. Lucio, Gary C. Mosely, James W. Norton, Lewis D. Ober, S. Craig Rhodes, James A. Rodgers, Jr., Bruce R. Sheplan, William W. Sommer, Michael H. Strahm, Oscar Vargas, and C. Rhea Warren. Specimens, in addition to those in the ASFS, have been examined in the following collections, to whose curators we are grateful for loans: American Museum of Natural History (AMNH—Richard G. Zweifel and George W. Foley); Museum of Comparative Zoology (MCZ—Ernest E. Williams); National Museum of Natural History (USNM—George R. Zug). Lewis

D. Ober (LDO) also allowed us to examine material in his personal collection. Finally, we are grateful to Fred G. Thompson of the Florida State Museum for allowing the junior author to retain one specimen of *S. a. altavelensis*. The illustrations are the work of David C. Leber and Alvis Gineika, to whom we are grateful, and the junior author.

LITERATURE CITED

- GRAHAM, E. D., JR., AND A. SCHWARTZ. 1978. Status of the name *Sphaerodactylus cinereus* Wagler and variation in "*Sphaerodactylus stejnegeri*" Cochran. *Florida Sci.*, (4)41: 243-521.
- NOBLE, G. K., AND W. G. HASSLER. 1933. Two new species of frogs, five new species and a new race of lizards from the Dominican Republic. *Amer. Mus. Novitates*, 652: 1-17.
- SCHWARTZ, A. 1967. The *Leiocephalus* (Lacertilia, Iguanidae) of Hispaniola. II. The *Leiocephalus personatus* complex. *Tulane Stud. Zool.*, 14(1): 1-53.
- . 1977. The geckoes (Sauria, Gekkonidae) of the genus *Sphaerodactylus* of the Dominican Peninsula de Barahona, Hispaniola. *Proc. Biol. Soc. Washington*, (2)90:243-254.
- . 1979. The herpetofauna of Ile à Cabrit, Haiti, with the description of two new subspecies. *Herpetologica* (3)35: 248-255.
- SCHWARTZ, A., AND R. THOMAS. 1975. A check-list of West Indian amphibians and reptiles. *Carnegie Mus. Nat. Hist. Spec. Publ.*, 1: 216 pp.
- , AND ———. 1977. Two new species of *Sphaerodactylus* (Reptilia, Lacertilia, Gekkonidae) from Hispaniola. *J. Herpetol.* (1)11: 61-68.
- SHREVE, B. 1968. The *notatus* group of *Sphaerodactylus* (Sauria, Gekkonidae) in Hispaniola. *Breviora Mus. Comp. Zool.*, 280: 1-28.
- THOMAS, R., AND A. SCHWARTZ. 1977. Three new species of *Sphaerodactylus* (Sauria: Gekkonidae) from Hispaniola. *Ann. Carnegie Mus. Nat. Hist.*, (4)46: 33-43.

Biogeography of the Herpetofauna of the British Virgin Islands, with Description of a New Anole (Sauria: Iguanidae)

JAMES D. LAZELL, JR.¹

ABSTRACT. Specimens and field data were collected for 44 of 46 named islands in this group. Published lists provide additional data for 24 of the islands visited and both that were not. The islands consistently have more species than areas, elevations, and distances from propagule sources might lead one to predict. There seems to be a basic, minimal number of three species per island, regardless of physical parameters. These are generally an *Anolis*, a member of the family Gekkonidae, and some other reptile. Even tiny rocks and overwashed bars usually support at least an *Anolis*. One such, Carrot Rock, 1.2 ha and 26 m high, supports a remarkable new species of the alpha section of *Anolis*. It is a member of the *crisatellus* group *sensu* Williams attaining very large size, combining two scale characters: 1) dorsals small; 2) digital lamellae count high; and three color characters: 1) chin tri-colored; 2) latero-dorsal trunk and neck boldly patterned; 3) a dark bar or blotch on each side of the sacrum. Islands, species, areas, heights, and distances are tabulated and discussed.

INTRODUCTION

The British Virgin Islands constitute the fragmented northeastern extremity of the Greater Antilles. For the most part they are old, folded, and buckled continental strata of grano-diorites, basalt, conglomerates, and slates and shales (Martin-Kaye, 1959). The nether isle of Anegada and some strata scattered on some of the other cays are oceanic limestones deposited during Pleistocene in-

terglacials. These same interglacial seas, standing dozens of meters higher than the sea today, loosened, rolled, and tumbled the rocks in many areas, leaving piles of huge boulders—some as big as a house—signalled today by names like Fallen and Broken Jerusalem.

As sea level fell following the Sangamon Interglacial, some hundred thousand years ago, more and more land was exposed. At the Wurm glacial maximum, the entire Puerto Rico Bank was dry land all the way to Anegada. In the last ten thousand years the rising sea has refragmented the land into fifty or more islands, rocks, and cays. Since the Wurm, sea level has probably never stood higher than it does today; it continues to rise (Morris *et al.*, 1977).

The British Virgin Islands are an artificial entity, separated from their American neighbors by a line passing between Pelican (British) and Flanagan (U.S.) Islands on the south, curving west through The Narrows between the Thatches (British) and St. John (U.S.), and running westward into the North Atlantic south of the Tobagos (British). My study of these islands involved 40 days of field work in March and April, 1980. Not counting rocks with no more than herb-stage vegetation, I tallied 46 islands suitable for reptiles and/or amphibians (see Fig. 1). I visited all of these except Jost Van Dyke and Mosquito, both large islands, often and easily visited, and both with good species lists. It is un-

¹The Conservation Agency, 8 Swinburne Street, Conanicut Island, Rhode Island 02835, U.S.A.

Figure 1. The British Virgin Islands. Four nameless cays are numbered. (Modified from Marler, 1973.)

likely I could have expanded either list, except by collecting the gecko, *Hemidactylus mabouia*, which I presume is present, if unrecorded, on both.

My count of 46 is too low. An unnamed but well-vegetated island of more than a hectare lies off Key Point, south of the western peninsula of Peter Island. Most people assume it to be part of that island, but I am assured (Marler, personal communication) that it is disjunct. Two rocks south of Dead Man's Chest appear to have shrub-stage vegetation as viewed from the heights of Peter Island. A rock separated from the southwest tip of Norman Island requires investigation. A separated, southern chunk of the Carrot Rock scarp should be climbed. There may be more.

Within historic times (though no one

living seems sure just when) "Pelican Cay" (not Pelican Island) was amalgamated with Little Jost Van Dyke by a (now) well-vegetated tombolo of cobbles and sediment; it lost its status as an island. In temporal similarity (and uncertainty), however, Green Cay was fragmented in two; its erstwhile southern terminus is today fully disjunct and called Sandy Spit. That is one of the smaller land areas on earth with two species of native reptiles. Man's activities have connected Wickhams Cay and Nannie Cay to the Tortola mainland with solid fill; these are not given separate consideration. Frenchmans Cay was separated from Tortola historically by mangrove swamp. Fill was built up to make a causeway from Tortola almost to the cay, and a channel ca. 2 m wide was

dredged (and bridged); Frenchmans Cay has been considered here as a separate island.

An island measuring ca. 50×100 m, showing the symbol for scrub vegetation, is a prominent feature on Directorate of Overseas Surveys (1959) map of Anegada, labelled "Little Anegada." It is imaginary. The real Little Anegada is a tiny (ca. 3×5 m), wave-washed block of limestone, ca. 30 cm above water at mean low tide. It is vegetated only with the maritime plants *Sesuvium portulacastrum* and *Rhizophora mangle* (one). "In before time," states Mr. Lance Vanterpool (personal communication), "it was no bigger, but it was more prominent because here were not all these mangrove groves." The more recent map, Directorate of Overseas Surveys (1977) quite correctly shows the region as mangrove continuous with Anegada itself. The place where the imaginary "Little Anegada" was located in 1959, is, quite correctly, open water.

Much herpetological work had been done in these islands before I arrived. Lists were available, prepared by Maclean *et al.* (1977) from Schwartz and Thomas (1975) and Philiposian and Yntema (1976, 1977, 1978). Heatwole (1976) had examined *Anolis cristatellus wileyae* from eight additional islands. Nevertheless, there were no data for 18 islands. I obtained 13 new records for 10 islands with extant lists, and 44 new records altogether. Nevertheless, I must have failed to find many populations, and there is undoubted necessity for far more work.

I missed populations for three major reasons: 1) I had to average over a cay per day in my field inventory; this is obviously far too little time for even the tiniest land areas. 2) I was on the ground at the height of the dry season in an unusually arid year; many species were cryptic, especially by midday, and some may have been in brumation. 3) I am me. I have poor vision and a gimp leg, and some say I move more slowly than I did a quarter-

of-a-century ago—and climb less high. And, I do not like to hunt frogs too small to eat (all in these islands are that). However, from a theoretical point of view, the fact that I missed lots of populations will, I hope to demonstrate, prove trivial.

THE FAUNA

Maclean *et al.* (1977) list 23 species of reptiles and amphibians occurring in the British Virgin Islands. Of these, at least one, *Anolis cuvieri*, is not known from specimens; Schwartz and Thomas (1975) note that its existence on Tortola has remained "unverified for over a century." Perhaps it should be deleted, but I saw fine habitat for this species in the big ghuts draining Mount Sage, on Tortola, and on the rugged eastern slopes of adjacent Beef Island. *Anolis cuvieri* could occur here.

Another species, the tortoise *Geochelone carbonaria*, is a suspected introduction. It is today unknown from Peter Island, whence listed, and rarely seen on Tortola. It is reported to occur on Virgin Gorda. Tortoises of one sort or another were widespread in the pre-Columbian Antilles (Auffenberg, 1967, 1974); no one has done the sort of study of geographic variation in *G. carbonaria* needed to demonstrate its true status, spurious or otherwise. I leave it in.

I have previously (Lazell, 1973) discussed the notion that *Iguana iguana* was introduced to these islands, the northern limit of its range. The *I. iguana* seen and collected during the present investigation support my earlier view: they are native.

On three rock cays, where I believe reptiles probably occur, I failed to obtain specimens: Carval, North Cockroach, and the Indians. All support sea grape (*Coccoloba uvifera*) clumps and other vegetation. I believe *Anolis* (probably *A. cristatellus wileyae*) live on them, but weather conditions prevented me finding the animals on North Cockroach. I was

unable to climb to the best vegetation on Carval and the Indians.

Mirecki (1977) led a group of four ornithologists and a botanist to the islands in 1976. I visited all the islands they failed to, and 11 more they did not consider. However, Dr. J. D. H. Smith (personal communication) even went onto rocks with a few sedges where he reports seeing no lizards.

The results of all this are presented in Table 1, where the islands are ranked in order according to area. Despite the numerous new island records, the fauna presented no surprises whatsoever until 4 April, 1980, when I climbed the sheer face on the northeast side, tunnelled through the *Coccoloba*, and entered the strange world of Carrot Rock. Apart from the previous three weeks, I had been out of *Anolis* work in the Antilles for over a decade, but I there saw my next new one:

*Anolis ernestwilliamsi*¹ sp. nov.

Figures 2, 3

Holotype. MCZ 158395, an adult male, J. Lazell coll. 4 April, 1980.

Type locality. Carrot Rock, south of Peter Island, British Virgin Islands, 18°19'45" N, 64°34'18" W, Caribbean Sea.

Diagnosis. An alpha section *Anolis* of the *cristatellus* group *sensu* Williams (1976:17), attaining very large size (to at least 82 mm SVL), combining the following two scale characters: 1) scales small, 34 to 45 (av. 40) middorsals contained in the standard distance at midbody; 2) digital pads large, males with 24 to 27 lamellae, females with 20 to 22 lamellae, under the second and third phalanges of the fourth toe; and combining the following three color characters: 1) chin tricolored, with a bold, reticulate or barred pattern of near-black to grey on lighter blue-grey,

with egg-shell white to creamy blotches. In the male the dark chin elements extend onto the anterior throat fan; in the female they extend posterior to at least the level of the eye. 2) Latero-dorsal trunk and neck boldly patterned with near-white to yellow spots and marblings, coalescing to form stripes, and isolating spots of darker pigment ventrolaterally. 3) A dark bar or blotch on each side of the sacrum, set off posteriorly by a pale, ashy or tan border. In the females the middorsal stripe is irregularly edged, often fragmented, and in poor contrast to the bold latero-dorsal pattern.

Description of the type. MCZ 158395 is an adult male 81 mm SVL, with a standard distance 13.8 mm. There are 42 middorsals, 33 ventrals, and 53 dorsals (eight rows lateral to the middorsal line) contained in the standard distance at midbody. There are 26 lamellae under phalanges II and III of the fourth toe. There is an enlarged, double row of bluntly tectiform middorsals. The ventrals are smooth posteriorly and become tectiform to keeled anteriorly (chest region). There are seven loreal rows (left side); seven scales border the interparietal; there are seven plates across the snout between second canthals; there are 22 mental rows between third infra-labials.

In life the type was somber, dark grey-brown with olive tints on the limbs, tail, and lateral body. Near-white to cream-yellow spots coalesced to form streaks and marbling which became especially bold ventrolaterally. Slaty to sooty blotches and bars were scattered over the dorsal surfaces and were boldly set off on the ashy venter. There were yellow tones ventrally on the limbs and abdomen. The throat fan was deep crimson with a dark green center and scales that could change from white to grey. The anterior 30° of the extended throat fan was boldly reticulated with slate- and blue-grey with cream white. This tricolored pattern covers the entire chin region back to the jowls.

¹This name commemorates the man who, in 1958, agreed to pay my way back to Dominica if I could prove that island was occupied by one, not four, species of *Anolis*.

Figure 2. *Anolis ernestwilliamsi* sp. nov., from Carrot Rock, British Virgin Islands. The type, MCZ 158395, above; an adult female paratype, MCZ 158402, below.

TABLE 1. HERPETOFAUNA OF THE BRITISH VIRGIN ISLANDS AS KNOWN IN APRIL 1980.

Island or Cay	Area: ha	Elevation: m	Number of species	1. <i>Anolis cristatellus</i>	2. <i>Sphaerodactylus macrolepis</i>	3. <i>Ameiva exsul</i>	4. <i>Anolis stratulus</i>	5. <i>Mabuya sloanei</i>	6. <i>Hemidactylus mabouia</i>	7. <i>Anolis pulchellus</i>	8. <i>Liophis portoricensis</i>	9. <i>Typhlops richardi</i>	10. <i>Iguana iguana</i>	11. <i>Amphisbaena fenestrata</i>	12. <i>Leptodactylus albilabris</i>	13. <i>Liophis exiguus</i>	14. <i>Eleutherodactylus schwartzi</i>	15. <i>Epicrates monensis</i>	16. <i>Geochelone carbonaria</i>	17. <i>Eleutherodactylus antillensis</i>	18. <i>Anolis Ernestiwilliamsi</i>	19. <i>Thecadactylus rapicaudus</i>	20. <i>Eleutherodactylus cochranae</i>	21. <i>Iguana pinguis</i>	22. <i>Bufo lenur</i>	23. <i>Sphaerodactylus parthenopion</i>	24. <i>Anolis cuvieri</i>
1 Tortola	5444	521	19	●	●	●	●	○	●	●	●	○	○	○	○	○	●	●	○	●	○	○	●	○	○	○	
2 Anegada	3872	8.5	11	●	●	●	○	○	○	○	○	○	○	○	○	○	○		○	○				○	○		
3 Virgin Gorda	2130	414	16	○	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
4 Jost Van Dyke	840	398	8	○	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
5 Peter	429	177	11	●	○	○	○	●	●	○	○	○	●	○	○	○	○		○	○							
6 Beef	372	244	6	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
7 Great Camanoe	337	187	8	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
8 Guana	297	266	8	●	○	○	○	○	○	○	○	○	○	○	○	○	○	S	○	○							
9 Norman	257	131	6	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
10 Cooper	138	155	4	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
11 Great Thatch	123	187	3	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
12 Ginger	105	164	5	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
13 Scrub	97	141	3	○	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
14 Great Tobago	87	147	2	H	X	X	X	○	○	○	○	○	○	○	○	○	○		○	○							
15 Salt	78	124	7	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							
16 Prickly Pear	70	52	4	●	○	○	○	○	○	○	○	○	○	○	○	○	○		○	○							

The sacro-iliac region was pigmented on each side in a rough triangle of sooty color, set off posteriorly by ashy-white.

Variation. Ten paratypes, six males and four females, were collected by me and George Marler on Carrot Rock, 18 April, 1980. The males are MCZ 158396-8 and 158400-1; the females 158402-4. A male, MCZ 158399, and a female, MCZ 158405, have been deposited in the British Museum (BMNH 1980.1-2).

The largest adult male, MCZ 158396, was 82 mm SVL, fresh. The smallest, judged adult in the field on the basis of territorial defense, courtship, and display, was MCZ 158399: 71 mm SVL. Two juvenile males, MCZ 158400-1, measure 63 and 59 mm SVL, respectively. Females measure 52 to 60 mm SVL; MCZ 158402 is the largest. Two 52 mm females are apparently immature. One, MCZ 158404, shows some beginnings of ovarian activity on both sides, with larger yolks present on the right, but seems

never to have laid an egg. The other, MCZ 158405, is definitely immature. A 57 mm female, MCZ 158403, has a well-developed ovarian egg on the left and an apparently recently spent oviduct on the right; she was mature.

The standard distance varied from 17.0 to 18.3 (av. 17.4) percent of SVL in males, and from 17.2 to 19.0 (av. 18.0) in females. The difference is not significant with the small sample sizes available. Except for lamellae and ventrals, discussed below, meristics were combined for both sexes.

Dorsals counted in the standard distance at midbody eight rows lateral to the midline varied from 45 to 53 (av. 49). Mid-dorsals in the same distance varied from 34 to 45 (av. 40). There were 6 to 8 (av. 7) loreal rows, 5 to 7 (av. 6) postrostrals, 20 to 28 (av. 25) scales between third infralabials, 8 to 11 (av. 10) suboculars, 5 to 8 (av. 7) scales across the snout between second canthals, and 12 to 18 (av. 15) circumparietals. I do not know how Heatwole (1976) quantified forearm scales, but I can see no obvious difference between the new species and *A. cristatellus* in this regard, so have omitted the character.

Sexual dimorphism in lamellae count is striking. Males have 24 to 27 (av. 26), females 20 to 22 (av. 21). There is also strong dimorphism in ventral scale size, quantified as midventrals in the standard distance at midbody. Males have 28 to 34 (av. 31), females 23 to 27 (av. 25). All specimens are plotted for these counts in Figure 4.

There is variation in the placement of the spots, streaks, mottlings, and marblings on the dorsal and ventro-lateral surfaces. However, these are always bold and prominent. Color change is from darker to lighter, and often enhances the markings in the disturbed extreme. There is real variation from a browner extreme (MCZ 158396) to a greyer extreme (MCZ 158399). The light pattern elements may approach white (MCZ 158400) or be quite bright yellow (MCZ

Figure 3. Head shape in two *Anolis*. A. *A. ernestwilliamsi* sp. nov., type-specimen, MCZ 158395, from Carrot Rock. B. *A. cristatellus wileyae*, type-specimen, MCZ 34792, from Culebra.

Figure 4. Sexual dimorphism in *Anolis ernestwilliamsi* sp. nov. from Carrot Rock. ● = type-specimen. See text.

158397). In both males and females the throat fan is deep crimson with a green center and white to grey scales. The female fan is quite small. The large male fan is variably invaded by the tricolor chin pattern elements anteriorly. In MCZ 158397, only about 20° of the extended fan is grey, blue, and cream. In MCZ 158396 nearly 40° of the fan is so patterned.

The sooty sacral blotch is prominent on all specimens, varying slightly in size and considerably in shape. The irregular, ashy middorsal stripe fades with age. The two smallest males retain it. Probably females retain portions of it throughout life, but in the large female (MCZ 158402) it was fading to obsolescence.

Comparisons. Heatwole (1976) provides an exhaustive account of geographic variation in *Anolis* of the *crisatellus* group on the Puerto Rico Bank and its satellites. He did not separate the sexes even with respect to lamellae count, and I do not find sexual dimorphism as pronounced in *A. crisatellus*—even *A. c. wileyae*—as in *ernestwilliamsi*. I counted 30 female *wileyae* and got 16 to 19 lamellae (av. 17); no overlap with female *ernestwilliamsi*. In males overlap occurs only in the Ramos,

Isleta Marina, and Villa del Mar populations where there may be 25, 24, or 25 lamellae, respectively. Two of these Puerto Rican coastal cays, Ramos and Isleta Marina, are populated by *A. c. crisatellus* × *wileyae* intergrades which cannot look much like *A. ernestwilliamsi*. At both Isleta Marina and Villa del Mar the scales are much larger than in *A. ernestwilliamsi*.

Anolis crisatellus has a shorter snout, on the average, than does *A. ernestwilliamsi*; in males it is 15.3 to 17.8% SVL (av. 17.0) and in females 16.4 to 18.2 (av. 17.2). Once again, the difference between sexes is insignificant, but the difference between species is significant at the 95% level of confidence. This difference does bias counts higher for *A. ernestwilliamsi* whenever the standard distance is used, but the scale size difference is real nonetheless, as can be seen from comparing like-sized individuals (e.g., MCZ Z-08348: *A. c. wileyae*; MCZ 158399: *ernestwilliamsi*).

Heatwole (1976:10) comments on the cline in scale size seen in *A. c. wileyae*: as one proceeds eastward from Puerto Rico through the Virgins, the anoles have lower counts—therefore larger scales. *A. ernestwilliamsi*, residing about 76% of the way along that cline, stands in bold contrast to it, with the smallest scales seen in any member of the complex.

I am not enamored of size as a taxonomic character in reptiles. Nevertheless I am compelled to admit the importance of size in the relationships of *A. ernestwilliamsi*. I measured 135 females of *A. c. wileyae*, and the largest I can find are 50 mm SVL (MCZ Z-08400, George Dog; MCZ 60635, Palominos, E of Puerto Rico; and MCZ 127725, Seal Dog). Some females of *A. c. wileyae* are producing eggs at 32 mm SVL: MCZ 35734, Vieques. Females of *A. ernestwilliamsi* from Carrot Rock may still be immature at 52 mm SVL (see above). Thus there is no evidence of overlap of sizes of mature females.

There is some overlap in sizes of males

that seem mature. Thus, large males of *A. c. wileyae*, MCZ Z-08348, from Broken Jerusalem, and MCZ 128484, from Culebra, are both 72 mm SVL, just larger than MCZ 158399, the smallest mature *ernestwilliamsi*. One male *wileyae*, MCZ Z-08429, from South Cockroach is a giant for his kind, measuring 75 mm SVL; he has 31 middorsals and 23 lamellae. MCZ Z-08429 has the shortest snout measured, 15.3% SVL, and was nearly patternless brown in life. None of the large *wileyae* approaches the color characters of *ernestwilliamsi*; none has more than 33 middorsals in the standard distance or 23 lamellae.

The three color characteristics which taken together are diagnostic of *ernestwilliamsi* can all be seen singly or weakly developed in occasional specimens of *A. c. wileyae*. MCZ 138543 (Estate Tutu, St. Thomas) combines a bold chin pattern with sacral marking like that of *ernestwilliamsi*; however, it lacks the dorsal and latero-ventral pattern. Conversely, MCZ Z-08409 (Eustatia) has fine trunk pattern but lacks the tricolored chin and sacral blotch. An adult male from Beef Island, MCZ Z-08436, was less boldly marked but a close approach to the color characters of *ernestwilliamsi* in life. The chin pattern was bicolor—dark grey and pale blue-grey—however, and there are only 33 middorsals and 22 lamellae. The closest approach to *ernestwilliamsi* I can find is MCZ 35952, from Culebra. This specimen is faded and may not have had a tricolor chin. Nevertheless, the pattern elements that remain closely resemble those of *ernestwilliamsi* in all three respects. The specimen just overlaps *ernestwilliamsi* in middorsal count with 34, but has only 23 lamellae. It is a seemingly adult male, but very small: 48 mm SVL.

There is a strong distinction in head shape of *A. ernestwilliamsi* relative to all *A. cristatellus*, but difficult to quantify. Basically the head of the Carrot Rock species is longer, lower in profile, the rostral bulge is longer, lower, and more gently tapered, and the angle the rostral

makes with the mouth is oblique, not rectilinear. These differences are shown in Figure 3.

In summary, *Anolis ernestwilliamsi* is closely allied to *Anolis cristatellus* and shares its presacral vertebral count of 23, as do *monensis*, *desechensis*, and *scriptus*. *A. ernestwilliamsi* differs from all of these, and from Puerto Rican *A. c. cristatellus* in throat fan color, just as does *A. c. wileyae*. From this last form, its closest neighbor, *A. ernestwilliamsi* is distinct in combination of color characters, small scales, high lamellae counts, large size, and head shape. For those of us who like quick, quantitative ways to separate species, *A. ernestwilliamsi* may be separated from *A. cristatellus* by simply dividing the lamellae count into one hundred minus the middorsal count. In Figure 5 I have plotted this against snout-vent length to graphically portray the distinction.

Habitat and behavior. Carrot Rock is steep-to and cliffed for virtually all of its perimeter. The vegetated top of the ca. 1.2 ha islet is a canted plateau rising from ca. 6 m above sea level on the windward side to 27.6 m on the lee. The surface is a deceptively uneven boulder field.

The deception is perpetrated by the vegetation, which covers the surface rather evenly. The dominants are the polygonaceous sea grape, *Coccoloba uvifera*, and leguminous vines. Main stems of these plants may reach 30 cm and 5 cm, respectively. The plants grow prostrate and sprawling over high points on the boulder substrate and canopy pockets or low areas rather like caclin bush (*Clusia mangle*) does in the high Lesser Antilles (Lazell, 1972). There are small areas with soil in which grow grasses and sedges, especially near the crest of the cay.

The anoles live primarily on the rocks (to a lesser extent on the vegetation, especially sea grape trunks) under the canopy. This is a shady, dimly lit zone, cooler and somewhat protected from the full force of the tradewinds.

Anolis ernestwilliamsi is quite com-

Figure 5. A graphic depiction of the morphometric distinction between *Anolis ernestwilliamsi* sp. nov. (solid symbols), and *A. cristatellus wileyae* (males, $\circ$; females $+$). M = middorsals counted in the standard distance. L = subdigital lamellae.

mon in its habitat. Sitting quietly in a pocket below the vegetation, I could count two to four in ca. 100 m². I estimate about two-thirds of the island provides good habitat, so a population of two to three thousand individuals is not unreasonable.

My observations indicate a balanced sex ratio. Both sexes bob and fan in the manner of the widespread *A. c. wileyae*. Apparently sexual maturity is attained at sizes larger than 63 mm SVL in males and 52 mm SVL in females. The largest female, MCZ 08460, 60 mm SVL, has a large egg in the left oviduct and a smaller one in the right. This implies the alternating oviduct, single egg laying strategy common in Antillean anoles (Lazell, 1972). The 57 mm female (MCZ 158403) was yolking up a large egg on the right

and seemed to have just laid an egg from the left at time of capture.

Small islands in the British Virgins (like those elsewhere in the Antilles) are occasionally burned over and frequently used for raising such livestock as goats. No evidence of fire or goats was visible on Carrot Rock, and its difficulty of access may eliminate the temptation to so abuse it. While I have no evidence that either burning or goat browsing extirpates lizards, I would certainly regard either or both with grave trepidation on Carrot Rock. In view of Carrot Rock's remarkable nesting seabirds (Lazell *et al.*, 1982)—none of which utilize the vegetated upland—and *Anolis ernestwilliamsi*, this islet should be set aside as a sanctuary or preserve. Limited collecting of the anole, especially by the technique of noosing individuals around the edge of the vegetated top, can do no serious harm, but such collecting should be regulated (see Lazell, 1980a).

Discussion. Heatwole's (1976) taxonomic conclusions are difficult for me to understand. It is unclear to me why the karyotypic difference between *A. monensis* and *A. cristatellus* augurs for the species status of *A. deseichensis* (Heatwole, 1976:13). In any case, karyotypic differences are not always indicative of species status in *Anolis* (or other vertebrates): Hall (1974). All the scale counts provided for *deseichensis* are included within the range of variation of Puerto Rican *cristatellus*, leaving no device for separating *deseichensis* from the yellow-fanned form in southwestern Puerto Rico. Scale characters are not specified (on p. 7 Heatwole alludes to a difference in postrostrals but his Figure 4 belies it), and probably only modal.

I agree with Heatwole (1976:12-13) that *A. c. wileyae* is an indivisible entity. The statistically significant differences Heatwole found fall far short of taxonomic significance. I find it impossible to frame diagnoses that would separate 75% of Vieques specimens, for example, from those in the Anegada population—at the opposite extreme of the range. (I have

examined 21 Anegada specimens: U.S. National Museum of Natural History 140309-10; MCZ 12145-56 and 35704-10.)

Against this background it is easy to defend *Anolis ernestwilliamsi* as a full species. Even granting that the minor karyotypic difference between *crisatellus* and *monensis* bequeaths species status to them, and assuming that *ernestwilliamsi* has the *crisatellus*-like complement of 27 (as one suspects *desechenensis* does), *ernestwilliamsi* is far more different from any of these previously named forms than they are from each other. It does not fit into the pattern of geographic variation shown by the intergrading forms of *Anolis crisatellus*, and geographically proximate populations of *A. c. wileyae* show no hint of character approach.

I fought my way through dense scrub on Peter Island to reach Stoney Bay, the southern edge of which is the closest land to Carrot Rock, and there collected a series of typical *A. c. wileyae* (MCZ 158927-31). This most proximate habitat is a xeric woodland dominated by gumbo-limbo (*Bursera simarouba*), figs (*Ficus* sp.), and mangroves (*Rhizophora mangle* and *Avicennia rhizophorarum*). A bay head barrier of cobbles, coral, and slabs that look like limestone fronts the sea. Anoles were scarce, but there were geckos (*Sphaerodactylus macrolepis*), iguanas (*I. iguana*), skinks (*Mabuya sloanei*), and kestrels (*Falco sparverius*). Standing on the cobbles I could identify to species the tropic birds (*Phaethon aethureus*) wheeling over their nest sites on Carrot Rock, a scant 450 m away. I wondered if the tropic birds ever alighted over here, or the kestrels over there. So far as I know, I am the only living thing which has ever been both places.

Anolis ernestwilliamsi is an autochthonous endemic on its peculiar little cay. It must have evolved very rapidly (see Lazell, 1972: 102-103) in a few thousand years of isolation.

Electrophoretic, karyotypic, and ethological studies of *Anolis ernestwilliamsi* are certainly called for. On present evidence, its status as a full species seems to me more secure than that of *monensis* or *desechenensis*. Granting that tastes differ on these points, dialogue and exposition will prove useful to those of us particularly interested in insular speciation and geographic variation.

OTHER FORMS

I suggest some nomenclatural changes not commonly in use today. Previously (Lazell, 1973) I commented on the trivial nature of the putative distinctions between "*Cyclura*," "*Brachylophus*," and *Iguana*. Examination of specimens reveals that even these differences are modal; I refer to all species in the complex as *Iguana*.

Maglio (1970) regarded the Caribbean colubrids as "oversplit." I concur. The entire complex of "*Alsophis*," "*Dromicus*," "*Arrhyton*," and *Liophis* are closely comparable to *Anolis* or *Eleutherodactylus* (or *Iguana sensu lato*) in diversity. Differences in dentition, cranial bone proportions, coloration, and hemipenes are, I suggest, reflections of character divergence among very closely allied species. Certainly the two Virgin Island species are congeneric; they seem a near perfect analog of the sympatric lizard pair *Anolis crisatellus* and *A. pulchellus*. The oldest name for this assemblage of snakes seems to be *Liophis*; it is used herein.

The species and their distributions are tabulated (Table 1). I have previously (Lazell, 1980a) listed four origins for members of this herpetofauna: 1) Species widely distributed on the greater Puerto Rico Bank stranded today in the Virgins by post-Wurm sea level rise. 2) Species coming from the west (Greater Antilles and other Puerto Rico Bank islands) across water. 3) Species coming from the east and south (Lesser Antilles) across water. 4) Autochthonous endemics

whose original ancestors may have come by any of the first three methods.

I cannot envision a way to separate examples of the first and second categories. I suspect most of the herpetofauna belongs to the first category, but it is truly unfortunate that I cannot perceive examples of the second, for they would be most interesting.

Four species (ca. 17%) came by the third method from the Lesser Antilles: *Hemidactylus mabouia*, *Thecadactylus rapicaudus*, *Iguana iguana*, and *Geochelone carbonaria*.

Five species (ca. 25%) are endemic to the islands east of Puerto Rico at full species level and are not known to have ever occurred on Puerto Rico: *Amphisbaena fenestrata*, *Sphaerodactylus parthenopion*, *Iguana pinguis* (?), *Anolis ernestwilliamsi*, and *Eleutherodactylus schwartzi*. There are bones of an *Iguana* near *pinguis* from Puerto Rico. Of these forms the last four are today known only from the British Virgin Islands (Lazell, 1980a). All of these species seem most likely to have been derived from Greater Antillean (in fact Puerto Rican) stocks. If one assumes that speciation began after the retreat of the Wurm, the processes have been rapid indeed. This is not, of course, necessarily the case, but seems most likely for at least *Anolis ernestwilliamsi*.

THEORIES OF BIOGEOGRAPHY

Two disparate and sharply contrasting theories of island biogeography have emerged in recent years. The first, published by MacArthur and Wilson (1967), is essentially abiological. It attempts to explain faunas (or floras) entirely in terms of measurable physical and/or temporal parameters such as island area, elevation, distance from a source of propagules, or length of time of island separation. The implicit philosophical basis of this theory is that biological phenomena and situations, like those in chemistry and

physics, should be reducible to symbols and numbers, which, in turn, can be arranged in mathematical formulae that "predict" (or postdict) similar biological phenomena or situations.

A necessary adjunct to the MacArthur and Wilson view is that time of island separation cannot matter very much because a large fauna stranded by rising sea level must necessarily dwindle by extinctions to the "right" number of species for an island with similar physical parameters which was never part of a larger land area. This notion has been hotly debated (see Simberloff, 1976, and works cited therein).

It is interesting that the MacArthur and Wilson theory germinated from Darlington's (1957) little rule of thumb which stated that Antillean herpetofaunas increase by *roughly* a factor of two for every increase in island size of a factor of ten. MacArthur and Wilson explicitly believed that, by measuring other parameters and coalescing them into ever more complex formulae, the roughness in Darlington's rule could be eliminated: the biota of an island should be calculated with little more difficulty than Boyle calculated pressure, given volume and temperature.

Lack's (1976) view of islands is considerably more complicated. While he did not doubt the importance in a general way of obvious physical features like island size, he was deeply impressed by biotic factors as well. He was impressed by the fact that, cumbersome as the formulae became, they still failed to "predict" (or, correctly, postdict to the time the data were collected) the right numbers in many cases. Most importantly, however, Lack was impressed by the resilience of insular faunas: apart from the artificial and edificarian catastrophes wrought by man, and often in spite of them, the predicted extinction rates were simply way out of line with apparent reality. Far from becoming extinct or replaced by new colonizers, insular populations often tended to expand their

niches and resist propagule input for so long that evolution carried them to the far realms of the bizarre. The dodo, Lack might have said, was just a pigeon which lived alone too long—but very, very long indeed. Many recent data on population genetics within the well-studied species *Felis catus* support this view of impenetrable matrices: Blumenberg (1977); Todd and Blumenberg (1978).

Lack's view may have grown from his field work in the Galapagos. Insular fragmentation may not result in diminished faunas at all. Quite the contrary, it may well lead to spectacular radiations.

The number of forms of finches, or tortoises, or lizards increases (very roughly) as a function of the number of islands. Since rising sea level increases the number of islands, the fact that it decreases their size may utterly fail to reverse the trend of species proliferation. *Anolis ernestwilliamsi*, possibly *Iguana pinguis*, and the subspecies of *Typhlops richardi* and *Liophis portoricensis* are examples of this trend in the British Virgins.

ANALYSIS

When a person has a poor ear for music he will flat and sharp right along without knowing it. He keeps near the tune, but it is *not* the tune.

—Mark Twain, 1895

Of the 43 islands with presently demonstrated herpetofaunas, only two—Great Tobago (two species; 88.6 ha) and West Dog (one species; 12.5 ha) fit the area-species curve of MacArthur and Wilson (1967:Fig. 2). All the rest have “too many” species. Tortola, Virgin Gorda, and Peter Island, for example, all have about three times the predicted number. Extrapolating the MacArthur and Wilson curve indicates that islands of less than ten hectares should have no species at all. More than 30% of the British Virgin Islands diverge from MacArthur and Wilson's predictions,

therefore, by an infinite factor. The relationship of area to number of species is shown in Figure 6.

Using the basic formula of MacArthur and Wilson (1967:8), but with new numbers, we find:

$$S = 1.63 A^{0.233}$$

Spearman $r = 0.797$, $p = 0.01$, which is reasonable: larger islands do tend to have more species. Extrapolating this curve into the realm of no species present indicates that at least one species should be present on any island of ca. 50 m² or greater. While this prediction will be most interesting to test, it is certainly less unreasonable than the previous 10 ha prediction.

The value $Z = 0.233$ is far lower than MacArthur and Wilson cited for any other fauna except land and fresh-water birds in the West Indies (1967:9). As Z approaches zero, the importance of area diminishes (at $Z = 0$ the formula becomes $S = C$).

Using the new, “improved” formula, 14 islands (nearly a third: 32 percent) have “too many” species and 11 islands (more than a quarter: 26 percent) have “too few.” Therefore, 58% of the islands fail to fit even the new predictions. It is important to understand that a biogeographic formula with a positive correlation of 80% is still *wrong most of the time*.

The cases involving too few species are, I must admit, probably the result of collecting failure. Great Thatch, Scrub, Great Tobago, West Dog, and Pelican are all “missing” two species. Cooper, Little Jost Van Dyke, West Seal Dog, East Seal Dog, Carrot Rock, and Broken Jerusalem are all “missing” one species each. Of these only Carrot Rock was examined carefully by me. Nevertheless, I believe additional search at a wetter season will reveal at least one more species there, and I believe all these islands are good for more species than are known at present. Finding the “missing” species will

Figure 6. Area-species relationships. Dots represent the 43 British Virgin Islands with presently known herpetofaunas. The solid line is the curve shown by MacArthur and Wilson (1967: Fig. 2). The dashed line is the extrapolation of this curve to the point where no species would be predicted present by MacArthur and Wilson (1967).

have a profound effect on the formula and the next generation of predictions. The value of C will be elevated; therefore the predicted size for a minimal area island harboring one species will be even less than 50 m². The value of Z will be reduced. Therefore the importance of area in the calculation will diminish.

Islands with too many species are far more embarrassing for advocates of a MacArthur and Wilson approach to biogeography. Of the islands with only one species "too many" the error is only about one-fifth for Mosquito and Necker (each with five species). On Fallen Jerusalem the error approaches 30% and on Round Rock it is more than a third (both have four species). For Saba Rock and Sandy Spit the failure of prediction is about 50%.

Of islands with two species "too many," Great Camanoe and Guana are not egregiously in error at about 20 to 25%. Sandy Cay (with four species) is about half off predicted, but Marina Cay (with five species on one hectare) is "wrong" by a factor of more than two-thirds.

The larger islands are little closer to

predicted values. All have counts 36 to 39% too high. This means that Salt Island has three species "too many," Peter has four, Virgin Gorda six, and Tortola has seven species more than predicted by the formula. Errors of this magnitude and frequency render the formulation useless to me as a practicing field biologist.

There is a weaker but "good" correlation of elevation (E) to species number: Spearman $r = 0.586$, $p = 0.01$. Using both kinds of data and searching for the best formulation in terms of least root mean square error (Smith, 1977: 257), one gleans:

$$S = C_1A + C_2E + C_3$$

$$\text{Where: } \begin{array}{ll} C_1 = 0.00198 & C_2 = 0.016 \\ C_3 = 2.1 & \text{RMS} = 1.65. \end{array}$$

This formula is strongly appealing mathematically because it represents a canted, but flat, plane rising in three dimensions. The fact that it rises from $A = 0$, $E = 0$, $S = 2$ —an island with neither area nor elevation, but with two (sodden) species—is a mathematically

inconsequential problem easily relegated to some lizard hunter for solution.

An interesting feature of this formula is that it values area less by an order of magnitude than it does elevation. This reverses the order of correlation produced when each factor was independently compared to species number *via* Spearman rank coefficients.

For islands > 100 ha the percent error for this formula is > 30 for only two islands: Peter (ca. 47% or five species "too many") and Great Thatch (78% or two species "too few"). Virgin Gorda has ca. 19 percent, or three species "too many." Of the twelve largest islands, the remainder are either off by one or two species (seven) and low percentages (10 to 30%) or are right on: two, Beef and Ginger Islands. In my opinion, both Beef and Ginger are undercollected and will prove to support several additional species in each case.

Of the 18 medium-sized (> 10, < 100 ha) islands, 13 have numbers of species differing from the prediction. The worst is West Dog, ca. 200% off with two species "too few." Great Tobago is more than 100% off also, and "missing" two species. Necker Island is more than 40% off with two species "too many," and Scrub errs about the same extent with two "too few." Salt Island is about 40% "wrong," with three species "too many." Buck, Little Camanoe, George Dog, Dead Mans Chest, and Eustatia are right on. The rest (eight) are off by one species; this amounts to an error of more than a third for Little Jost Van Dyke and Frenchmans Cay (both "too few"). For the remainder the percentage error is lower.

Of the 13 small islands (< 10 ha), ten have numbers of species differing from the prediction. Five of these have errors between 100 and 200%; of course the number of species "wrong" is usually just one, but both Carrot Rock and West Seal Dog are "missing" two species. Sandy Cay has two species "too many" and Marina Cay is the champion with

three species "too many." Bellamy Cay, Saba Rock, and Sandy Spit are right on.

Thus, using the new formula we get a much tighter fit in terms of percentage error in the larger islands (especially), but for 33 of the 43 islands—*more than three-quarters*—the formula is still *wrong*. As before, errors of "too few" will predictably be corrected by better hunting.

In attempting to find a fit that avoided the problem of $A = 0$, $E = 0$, $S = 2$ we fed the computer data for Carval, Indians, and North Cockroach, which have fine areas and elevations, but no known species. We also fed it the open ocean at $A = 0$, $E = 0$, $S = 0$. The computer could deal with this in the linear equation, but it did not help. It increased RMS but did not bring $C_3 = S$ below two significant figures. In exponential arrangements the log of zero cannot compute, of course. Giving tiny, false values permitted computation, but no exponential arrangement, with or without the $S = 0$ data points, was as good as the linear one.

A slight further refinement can be made to this formula utilizing the fact that there is some relationship between area and elevation (i.e., big islands tend to be higher than small islands):

$$S = C_1A + C_2E + C_3AE + C_4$$

$$\text{Where } C_1 = 0.0025 \quad C_2 = 0.019$$

$$C_3 = -1.97 \times 10^{-6} \quad C_4 = 1.8$$

$$\text{RMS} = 1.60$$

The effect of the admittedly tiny, negative value of C_3 is to dip the plain for large areas and high elevations. The biggest and smallest islands are now right on. Discounting islands with "too few" species as examples of collector failure, however, there are still 13 islands—ca. 30%—with "too many." Peter Island is the worst by a factor of nearly 50% and five species. Virgin Gorda, Salt, Necker, and Marina Cay all have an excess of three species. Great Camanoe and Sandy Cay are off by two. Norman, Prickly Pear,

Mosquito, Little Camanoë, Fallen Jerusalem, and Round Rock are all overstocked by one species.

The computer still asks us to believe that the open ocean has two species (at $A = 0$, $E = 0$, $C_4 = 1.8$, which is two). Perhaps it wants us to acknowledge the existence of marine turtles. In all the formula provides the wrong answer for 31 of 43 islands, or 72%.

Increasing the number of points to 47 by feeding in the three $S = 0$ islets and the open ocean brought C_4 to one—which is only half as insane, one might say—but increased RMS to 1.63. There was no improvement: the formula was still wrong for three-quarters of the islands.

CONCLUSIONS

Statistics is a game played by computers which are just as intelligent as bowling balls. My acquaintance is not large, but I have yet to meet a computer which knew that 16 (the number of known species on Virgin Gorda) is not at all the same thing as 16.00000. I believe that if mathematically inclined biogeographers would be more sanguine about “good” correlations and actually calculate the percentage error between the observed species numbers and their formulas’ “predictions” (and pay attention to which percentage errors amount to species numbers) they would take a far less enthusiastic view of computer time and costs. As I have pointed out before (Lazell, 1976, 1979), to determine the number of species on an island one must at the very least go to the island and count.

One striking fact was available to MacArthur and Wilson (1967) before they published: I had already shown that Sombbrero—anchor point in Darlington’s (1957) table from which MacArthur and Wilson selected their data (1967: Fig. 2)—had three species present (Lazell, 1964). Most West Indian islands, in fact, have at least three species comprising a

herpetofauna standardly involving a member of the genus *Anolis*, a member of the family Gekkonidae, and some other form.

The Rule of Three in the Virgin Islands holds for 22 islands over three orders of magnitude in area: Table 2. All of these islands support *Anolis c. wileyae*. All but three (the American islands of Leduck Cay, Flanagan, and Salt Cay)—86%—also support the gecko *Sphaerodactylus m. macrolepis*. The teiid lizard *Ameiva exsul* is present on 15, or 68%. Other species making up the complement of three include a second *Anolis* (*A. stratus*): five for ca. 23%; a skink (*Mabuya sloanei*): two for nine percent; a snake (*Liophis portoricensis*): two for nine percent; and a second gecko (*Hemidactylus mabouia*): one for less than five percent.

The Rule of the Irreducible Anole states simply that any West Indian islet, cay, rock, or spit which supports more than herb-stage vegetation—i.e., bushes or shrubs—will also support a member of the genus *Anolis*. This is not to be confused with Marler’s (1980) Mexican bush rule, albeit not wholly dissimilar. In my work in the Lesser Antilles (Lazell, 1972) I found but one islet—Green Cay off Saba—that had bushes (two *Coccoloba uvifera*) but no *Anolis* I could find. I believe every case of an islet in the British Virgins where I failed to find *Anolis c. wileyae* (Carval, Indians, North Cockroach) is simply that: where I failed.

I predict that subsequent work in these islands will add at least seven cays to those fitting the Rule of the Irreducible Anole (the three I failed to find *Anolis* on plus four unnamed ones). I predict that many islands now thought to fit the Rule of Three will turn out to have additional species, and many where less than three are known today will move up to fill the ranks. In short, I believe the fauna will certainly prove to be richer than is now known.

No extinction or extirpation need take place in the British Virgin Islands if reasonable conservation measures are en-

TABLE 2. THREE SPECIES ISLANDS IN THE VIRGINS.

Island	<i>Anolis c. wileyae</i>	<i>Sph. m. macrolepis</i>	<i>Ameiva exsul</i>	Other species	Area (ha)
1 Rotto Cay, A	X	X	X		0.8
2 Steven Cay, A	X	X		<i>Anolis stratulus</i>	2.4
3 Henley Cay, A	X	X		<i>Hemidactylus mabouia</i>	5.3
4 Cas Cay, A	X	X	X		7.1
5 Leduck Cay, A	X		X	<i>Anolis stratulus</i>	7.3
6 Cockroach I., A	X	X		<i>Liophis portoricensis</i>	7.7
7 Congo Cay, A	X	X	X		10.3
8 Flanagan, A	X		X	<i>Anolis stratulus</i>	10.5
9 Eustatia	X	X	X		10.5
10 Dead Mans Chest	X	X	X		14.1
11 George Dog	X	X	X		15.4
12 Little Camanoe	X	X	X		16.2
13 Buck	X	X	X		17.0
14 Little Tobago	X	X		<i>Mabuya sloanei</i>	22.3
15 Little Thatch	X	X		<i>Anolis stratulus</i>	23.5
16 Frenchmans	X	X	X		24.1
17 Salt Cay, A	X		X	<i>Mabuya sloanei</i>	25.6
18 Great Dog	X	X	X		30.3
19 Little Jost Van Dyke	X	X	X		57.2
20 Scrub	X	X	X		88.6
21 Great Thatch	X	X		<i>Anolis stratulus</i>	112
22 Hans Lollik, A	X	X		<i>Liophis portoricensis</i>	128

A = American islands with data from Maclean *et al.* (1977).

acted. I have already listed a number of endangered, threatened, and/or declining species and outlined plans to conserve them (Lazell, 1980a). In every case the decline, threat, or endangerment is strictly artificial: the direct result of human activities. In no case is a possible extinction or extirpation the foreseeable result of any natural phenomenon. Even tiny, overwashed Sandy Spit appears likely to accrete sediment on growing coral about as rapidly as sea level rises. I expect it to retain its two species and—with better search—prove to fit the Rule of Three, at least.

ACKNOWLEDGMENTS

I am most grateful to Gregory Mayer and Christopher Lutman for assistance in analyzing the data. William Geizentanner generously prepared Figures 4, 5,

and 6. Numi Spitzer made the map, Figure 1.

Numerous people assisted me in the field; at risk of omitting someone, I especially thank George and Luana Marler, Robert Chipley, John D. H. Smith, Gerald Christian, Leonard Dawson, Mary Randall, Herman and Marilyn Groezinger, Clement and Gracita Faulkner, Lance Vanterpool, Wallace Vanterpool, Paul Backshall, Grant Bongiorno, Hank Valdes, George and Jane Mitchell, Rowan Roy, Norman Thomas, Ishma Christopher, and the staff of the Peter Island Yacht Club. Robert Jenkins and Robert Creque handled the administrative end; Jose Rosado curated the collection.

My work was funded by the Department of Natural Resources, Government of the British Virgin Islands, and the International Program of The Nature Conservancy.

LITERATURE CITED

- AUFFENBERG, W. 1967. Notes on West Indian tortoises. *Herpetologica*, **23**: 34-44.
- . 1974. Checklist of fossil tortoises (Testudinidae). *Bull. Florida State Mus.*, **18**: 121-251.
- BLUMENBERG, B. 1977. Genetic differences and selection in domestic cat populations of the United Kingdom and former British colonies. *Theoret. Appl. Genetics*, **49**: 243-247.
- DARLINGTON, P. J. 1957. Zoogeography: the geographical distribution of animals. New York, John Wiley & Sons.
- DIRECTORATE OF OVERSEAS SURVEYS. 1959. Anegada. D.O.S. 346 (E 837). London, Edward Stanford Ltd.
- . 1977. Anegada. E 803 (D.O.S. 346P). London, Edward Stanford Ltd.
- HALL, W. P. 1974. Report in E. E. Williams (ed.). *The Second Anolis Newsletter*, Mus. Comp. Zool.: 1-2.
- HEATWOLE, H. 1976. Herpetogeography of Puerto Rico. VII. Geographic variation in the *Anolis cristatellus* complex in Puerto Rico and the Virgin Islands. *Univ. Kansas Mus. Nat. Hist. Occ. Pap.*, **46**: 1-18.
- LACK, D. 1976. Island biology illustrated by the land birds of Jamaica. *Studies in Ecology* 3, Berkeley, Univ. California Press.
- LAZELL, J. D. 1964. The reptiles of Sombrero, West Indies. *Copeia*, **1964**: 716-718.
- . 1972. The anoles (Sauria: Iguanidae) of the Lesser Antilles. *Bull. Mus. Comp. Zool.*, **143**: 1-115.
- . 1973. The lizard genus *Iguana* in the Lesser Antilles. *Bull. Mus. Comp. Zool.*, **145**: 1-28.
- . 1976. This broken archipelago. New York, Quadrangle, Harper and Row.
- . 1979. Deployment, dispersal, and adaptive strategies of land vertebrates on Atlantic and Gulf barrier islands. *Proc. First Conf. Sci. Res. Nat. Parks*, **1**: 415-419.
- . 1980a. Report. British Virgin Islands, 1980. Privately printed.
- . 1980b. Lesser Antillean *Anolis* (Sauria: Iguanidae) in the British Museum. *J. Herpetol.*, **14**: 194-195.
- LAZELL, J. D., R. CHIPLEY, AND G. MARLER. 1982. Contribution to the ornithology of the British Virgin Islands: in press.
- MACARTHUR, R., AND E. O. WILSON. 1967. *The theory of island biogeography*. New Jersey, Princeton Univ. Press.
- MACLEAN, W. P., R. KELLNER, AND H. DENNIS. 1977. Island lists of West Indian amphibians and reptiles. *Smithsonian Herp. Info. Service*, **40**: 1-47.
- MAGLIO, V. J. 1970. West Indian xenodontine colubrid snakes: their probable origin, phylogeny, and zoogeography. *Bull. Mus. Comp. Zool.*, **141**: 1-54.
- MARLER, G. 1973. *The undersea world of the British Virgin Islands*. Tortola, Caribbean Printing Co. Ltd.
- . 1980. In Mexico, no matter how small it is, if it's got a bush, that bush will be full of crabs. *Field Cat.*, **19**: 390-391.
- MARTIN-KAYE, P. 1959. Reports on the geology of the Leeward and British Virgin Islands. St. Lucia, Voice Publ. Co.
- MIRECKI, D. N. 1977. Report of the Cambridge ornithological expedition to the British Virgin Islands. Cambridge, Bluebell.
- MORRIS, B., J. BARNES, F. BROWN, AND J. MARKHAM. 1977. The Bermuda marine environment. *Bermuda Biol. Sta. Spec. Publ.*, **5**: 1-120.
- PHILIBOSIAN, R., AND J. A. YNTEMA. 1976. Records and status of some reptiles and amphibians in the Virgin Islands. I. 1968-1975. *Herpetologica*, **32**: 81-85.
- . AND ———. 1977. Annotated checklist of the birds, mammals, reptiles, and amphibians of the Virgin Islands and Puerto Rico. Fredriksted, St. Croix, Information Services.
- . AND ———. 1978. Records and status of some reptiles and amphibians in the Virgin Islands. II. 1975-1976. *Herpetologica*, **34**: 47-51.
- SCHWARTZ, A., AND R. THOMAS. 1975. A check-list of West Indian amphibians and reptiles. *Carnegie Mus. Nat. Hist. Spec. Publ.*, **1**: 1-216.
- SIMBERLOFF, D. 1976. Species turnover and equilibrium island biogeography. *Science*, **194**: 572-579.
- SMITH, J. M. 1977. *Scientific analysis on the pocket calculator*. Second edition. John Wiley & Sons, New York.
- TODD, B. B., AND B. BLUMENBERG. 1978. Mutant allele frequencies and genetic distance relationships in domestic cat populations of Lower Egypt and the eastern Mediterranean. *Theoret. Appl. Genetics* **52**: 257-262.
- TWAIN, M. 1895. Fenimore Cooper's literary offenses. In C. Neider (ed.), *The complete humorous sketches and tales of Mark Twain*. Garden City, N.Y., Doubleday.
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, **44**: 345-389.
- . 1976. West Indian anoles: a taxonomic and evolutionary summary I. Introduction and a species list. *Breviora Mus. Comp. Zool. No.* **440**, pp. 1-121.

Guiano-Brasilian *Polychrus*: Distribution and Speciation (Sauria: Iguanidae)

P. E. VANZOLINI¹

ABSTRACT. A historical résumé and formal synonymies are presented for *Polychrus marmoratus* and *P. acutirostris*. The two species are compared from the viewpoint of scale characters, color pattern, body proportions, and some aspects of ecology and behavior. Their distribution is analyzed and possible mechanisms of speciation discussed.

INTRODUCTION

Polychrus is a well-defined lizard genus, but the relationships between its species are not simple. The two forms that occur in the Guiano-Brasilian region, *P. marmoratus* (L., 1758) (Fig. 1) and *P. acutirostris* Spix, 1825, present very interesting problems of geographical-ecological distribution and consequently of speciation mechanisms. The data at hand permit a statement of such problems, and some measure of speculation.

HISTORICAL RÉSUMÉ

The history of *Polychrus*, as that of any common genus, is long, voluminous, full of irrelevance and redundancy, as well as of overlooked valuable information. At the species level the main taxonomic references are:

Lacerta marmorata was diagnosed by Linné (1758: 208) in the tenth edition of the *Systema Naturae*. He had seen specimens, and described them well, in the *Amoenitates Academicæ* (*Amphibia gyllenborgiana* and *Museum Principis*),

Figure 1. *Polychrus marmoratus* from Puruzinho, Rio Madeira, Amazonas, Brazil. Photo by W. R. Heyer.

¹Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil.

and in the *Museum Regis*; in the *Systema* he also cited a Seba plate. In view of the existence of actual specimens, this Seba citation as well as others included in the *Amoenitates*, but not in the *Systema*, may well be dismissed.

In his review of the Linnean materials in Uppsala, Lönnberg (1896: 14) stated that the specimen described in *Museum Principis* was still extant and was without doubt *Polychrus marmoratus* as then accepted, i.e., in the concept of Boulenger's catalogue. The specimen from the *Amphibia gyllenborgiana*, however, had disappeared. The Stockholm specimen (*Museum Regis*) had already been lost by 1802 (Andersson, 1900: 10). Holm (1957), reviewing the Uppsala materials, found four *Amphibia gyllenborgiana* specimens of *Polychrus* overlooked by Lönnberg (1896), and considered them as undoubted *marmoratus*. This name is then solidly founded on specimens.

Linnaeus, following Seba, initially gave "Gallaecia" (Galicia, in northern Spain) as the type locality of *L. marmorata*, broadening it to "Hispania" in the *Systema*. Gmelin (1789: 1065), in what is called the thirteenth edition of the *Systema*, added "America" to the distribution.

The first figure I found of *Lacerta marmorata* is Lacépède's (1788) Plate 26, a rough but perfectly recognizable woodcut. His description is adequate, and was adopted by subsequent writers, who also copied the figure.

Daudin (1802: 433) gave a good description and recorded the species, which he called *Agama marmorata*, from Surinam. I believe it is convenient to adopt this, and indeed Paramaribo, where the species is common, as the type locality, as proposed by Hoogmoed (1973); this is of course very plausible for a Linnean species.

Cuvier (1817: 41) erected the genus *Polychrus*, monobasic (monotypic at creation), for *Lacerta marmorata*, cited from Lacépède and thus recognizable beyond doubt.

Raddi (1820: 32; 1822: 58) listed *Agama marmorata* Daudin from Rio de Janeiro; this is the first mention of a specimen from the Atlantic forest, and the first reference to its being a forest animal.

Wied collected *P. marmoratus*, but did not refer to it in his "Reise" (1820). However, he made his notes available to Schinz who, in 1822 (p. 65), described, without locality, "*Polychrus virescens* Pr. Max." In the *Beiträge*, Wied (1825: 120) gave the exact locality of *virescens*, "Villa Viçosa am Flusse Peruhype," the present Nova Viçosa, Bahia; he had meanwhile come to the conclusion that *virescens* was a synonym of *marmoratus*. Wied's specimen is not cited by Burt and Burt (1931) in their report on the collection of the American Museum of Natural History, where Wied's surviving materials are kept. In the *Abbildungen* (1822–1831) Wied included a plate, published in 1829, with a good figure of the specimen, labelled as *P. marmoratus*.

In 1821 the Isis von Oken (pp. 337–342; see Vanzolini, 1977: 26) published an advertisement, signed by Wagler, of a forthcoming book. I have not been able to find any further reference to this work, but in the pages following the advertisement there is an anonymous review of the sample fascicle, certainly by Oken himself, mentioning a new generic name, *Psilocercus*, proposed by Wagler, apparently without further explanation, for *Lacerta marmorata*. This name seems to have never appeared again in the herpetological literature. It is, in any case, a clear junior synonym of *Polychrus* Cuvier, 1817.

Spix (1825: 14, Pl. 14) included *Polychrus marmoratus* from Rio de Janeiro among his new Brazilian species, as he did to a few other forms already known: this is not to be construed as the description of new taxa. On the next page and plate Spix described the new species *Polychrus acutirostris*, from "sylvia Bahiae." I take the word Bahia to mean the state (then province), not the hom-

onymous city, as the latter is always mentioned by Spix as "urbs." Spix's description, and especially the plate, are sufficient to identify the form. In fact, H. Boie (1826: 119) challenged the new species, saying that it was merely *P. marmoratus* stripped of its epidermis, but Spix (1826: 603) replied quite adequately, stating that his was a good specimen and mentioning the shape of the snout and the absence of a gular crest as differential characters.

Fitzinger (1826), in the *Neue Classification*, cited *Polychrus marmoratus*, proposed, as he was wont to do, a *nomen nudum* (*Polychrus geometricus*, "patria ignota") and placed *acutirostris* in his new genus *Ecphymotes*, which otherwise contained *E. plica* (= *Plica plica*), *E. undulatus* (*Anisolepis undulatus*), and *E. pictus* (*Enyalius pictus*). In his review of Spix's lizards, published after or about the time of Spix's death, Fitzinger (1827) maintained the same scheme.

Spix's materials were later reviewed again by W. Peters (1877: 410), who confirmed the identification of *marmoratus* and the validity of *acutirostris*.

Delaporte (1826), the future Count of Castelnau, described *Polychrus fasciatus*, without definite locality, but thought "with good reason" to come from either the Philippines or the Moluccas. Duméril and Bibron (1837: 69) noted that the type was a stuffed *P. marmoratus* on which a vertebral stripe had been painted. The specimen has not been cited either by Duméril and Duméril (1851) or by Guibé (1954) as extant in the Paris Museum collection.

In his *Icones*, Wagler (1833a: Pl. 12) figured and described two species. The first was *Polychrus virescens* Wied in Schinz, to which he attributed Wied's and Spix's Atlantic Forest specimens. The other, *P. strigiventris*, he with some hesitation described as new. Both plates are very good and clearly of *marmoratus*. *P. acutirostris* was mentioned as a good species.

Wagler (1833b), in a paper on Seba's lizard plates, published a synopsis of *Polychrus*, including: (i) *acutirostris* Spix; (ii) *marmoratus* Linné; (iii) *geometricus* Fitzinger (in doubt; not diagnosed and so still a *nomen nudum*); and (iv) *neovidanus*, new. The latter was based on a figure of Seba (Vol. 2, Plate 76, Fig. 4) and on Spix's *marmoratus*. The name should be attached to Spix's specimen, which has a definite locality (Rio de Janeiro), and was later examined by W. Peters; thus *neovidanus* definitely applies to the Atlantic forest form.

Wiegmann (1834: 16) in the *Herpetologia Mexicana* described a new species from "Brazil," *Polychrus anomalus*, very recognizably *acutirostris*. Gray (1845: 184) proposed the new genus *Sphaerops* for *P. anomalus*.

Hallowell (1845) described from "Colombia" (actually Venezuela: "within 200 miles from Caracas") *Leiolepis auduboni*, recognized by Roze (1958), who saw the presumptive type, as *Polychrus marmoratus*. Berthold (1846) described *P. gutturosus* from Popayán, Colombia.

The next significant contribution, and a very important one, was made by Reinhardt and Lütken (1861). They had both species, *marmoratus* without provenance data, and *acutirostris* (under the name *anomalus*) from several unspecified localities in Minas Gerais, but also definitely from Lagoa Santa, the first well defined locality for the species. They presented an excellent differential diagnosis, which has helped to orient the present work, and noted that the genus *Sphaerops* could not subsist on the basis of Gray's diagnosis, although it might eventually come to be justified on other grounds.

Boulenger (1885: 98-100), in the British Museum Catalogue, using good materials of *marmoratus* (including one specimen from Bahia) and only one *acutirostris* from "Brazil," presented a reasonable key, and adequate descrip-

tions, but was not as incisive as Reinhardt and Lütken (l.c.).

Following Boulenger's Catalogue came the usual rash of papers identifying specimens in collections from several parts of South America, some fortunately with field data. Cope (1899: 20) in a bare list of "a collection of Reptilia from Argentina" cited one specimen of "*Polychrus angustirostris* Wagl.," an obvious *lapsus calami* that should not be taken into account.

The first interesting contribution of this period was Peracca's (1897) observation that two young *marmoratus* from Amazonian Ecuador had smooth ventrals. Boulenger (1908: 113), using this character as diagnostic, described *Polychrus liogaster*, from two localities: Provincia del Sara in Bolivia and "Chancamayo" in Peru (actually Chanchamayo, beyond doubt since Schunke was the collector; Vanzolini, 1978).

In 1910 Werner described *P. femoralis* from Guayaquil, Ecuador, and Boulenger, in 1914, *P. spurrelli*, from the Chocó, in Colombia.

In 1924 Noble described the new genus and species *Polychroides peruvianus*. The genus was later synonymized with *Polychrus* by Gorman, Huey, and Williams (1969).

Mertens (1930: 269) reviewed the differences between *P. marmoratus* and *acutirostris*. He was the first to mention what the accumulation of locality records already permitted to infer: that *P. acutirostris* is an open formation form, and *marmoratus* a forest one. He also discussed the meaning of differences in color pattern, metachrosis, and relative eye size in terms of the ecological preferences.

Burt and Burt (1931), in their report on the American Museum collection, presented a key to the genus and made some confusing statements. They initially gave the range of *marmoratus* as "northeastern South America" and that of *acutirostris* as "southern Brazil, Uruguay, and

Argentina," all of which goes against most of the previous records in the literature for both species. Next they referred to the "intermediate region of Sao Paulo," which does not make sense either in view of the literature or of their own distributional data. They also recognized the trans-Andean *P. spurrelli* as a probable race of *marmoratus*, and determined as such a Manaus specimen, which I have seen and is plain *marmoratus*.

In their Check List, Burt and Burt (1933) assembled all species so far described, except *gutturosus*, as subspecies of *Polychrus marmoratus*, with the following geographical scheme: *acutirostris*, southern Brasil, Uruguay, Paraguay and eastern Bolivia; *femoralis*, southern Ecuador; *liogaster*, Bolivia and Peru; *marmoratus*, northeastern South America; *spurrelli*, Colombia and northwestern Brasil. This scheme is just as bad as the preceding one.

Parker (1935: 515), in his herpetology of Guyana, criticized the Burts's scheme, made some comments on the several forms, and proposed the presently accepted arrangement; his key, adapted from the Burts's and not very good, has subsequently been the basis for identification, and was adopted, in a simplified version, by Peters and Donoso-Barros (1970).

Amaral (1937), in his Check-list, proposed a new and totally gratuitous distributional pattern: *Polychrus m. marmoratus*, "equatorial and northeastern districts (from Amazonas to Pernambuco)"; *P.m. acutirostris*, "western, southern and southeastern districts (from Bahia to Rio Grande do Sul and neighboring countries)."

I (Vanzolini, 1948: 386) commented that all materials then at my disposal could be unambiguously assigned either to *acutirostris* or *marmoratus* (curiously enough, I failed to say why) and, unaware of Mertens (1930), remarked on the ecological distribution (open formation

versus forest). Schmidt and Inger (1951: 451), in their turn not aware of my paper, also came to the conclusion that *acutirostris* was a savanna (= open formation) animal.

Peters (1959: 119) commented on the difficulty of separating *marmoratus* and *liogaster* in Amazonian Ecuador. In 1967 he gave a key to Ecuadorian forms, based on Parker (1935), stressing body squamation and shape of the canthus rostralis.

SYNONYMY

Polychrus marmoratus (L., 1758)

Lacerta marmorata Linnaeus, 1758: 208.

Polychrus (marmoratus): Cuvier, 1817: 41.

Psilocercus marmoratus: Wagler in Oken, 1821: 341.

Polychrus virescens Schinz, 1822: 65. Brasil (Wied leg.).

Polychrus fasciatus Delaporte, 1826: 110. Philippine Islands or Moluccas.

Polychrus strigiventris Wagler, 1830: Pl. 12 and text. No locality given.

Polychrus neovidanus Wagler, 1833: 897. Rio de Janeiro.

Leiolepis auduboni Hallowell, 1845: 246. "Colombia": within 200 miles of Caracas.

Polychrus acutirostris Spix, 1825

Polychrus acutirostris Spix, 1825: 15, Pl. Sylvis Bahiae.

Polychrus anomalus Wiegmann, 1834: 16. Brasil.

Sphaerops anomalus: Gray, 1845: 154.

EXTERNAL MORPHOLOGY

Pholidosis (Hoogmoed, 1973: Figs. 29a-d; Vanzolini, Ramos-Costa, and Vitt, 1980: Figs. 71-74). The species of *Polychrus* are very similar in external appearance. They are characteristically arboricolous, with a laterally compressed body; the tail is long, prehensile and has all the scales strongly keeled, the keels forming longitudinal ridges.

The head is prismatic in shape, with a definite, if rounded, canthus rostralis. The upper head scales are irregular, without even proper supraocular semi-

circles or well differentiated supraorbitals. The gulars are arranged in characteristic longitudinal rows of narrow and elongate scales. The skin between the rows permits a considerable amount of stretching, the whole functioning as a throat fan.

The eyelids are also extremely characteristic. They are formed by concentric rows of granules, the upper and lower lids being fused together in front and behind, the seams remaining evident.

The dorsal scales, especially on the anterior half of the body are irregular, arranged in irregular oblique rows; on the flanks much wrinkled skin appears between the rows, allowing for stretching in display. The ventrals are lanceolate, keeled or smooth.

Femoral pores are always present, but vary much in degree of development, as well as in number (11-24 on a side). At times they are inconspicuous, or much deformed.

Although *marmoratus* and *acutirostris* are very similar in physiognomy, they differ in important respects in their externals, as first noted by Reinhardt and Lütken (1861).

The snout of *acutirostris*, as the name indicates, is narrow and pointed, compared to that of *marmoratus*. This is quite obvious at inspection, but I have not been able to devise a morphometric comparison that would clearly demonstrate the fact.

The scales of the top of the head of *marmoratus* are larger and nearly smooth, those of *acutirostris* smaller and of a very irregular texture.

The nostril, which is large, opens on the middle of a single scale, and its position differs in the two species. In *marmoratus* the nasal scale rests on the middle of the first labial, at most on the suture between the first and the second. In *acutirostris* it is on the suture between the second and third labials, or squarely on the third. Correspondingly, there is one scale between nasal and rostral in *marmoratus*, two in *acutirostris*.

As noted by Mertens (1930), the opening between the partly fused eyelids is relatively larger in *marmoratus*. This is not an easy character to measure, but the opening in this species approaches 50% of the diameter of the orbit, being 30% or less in *acutirostris*.

In the gular region there are two very noticeable differences. Starting a little behind the post-symphysials, there is in *marmoratus* a series of 10 to 15 thin raised scales (the "crista palaris" of the older authors); these scales are pliable but form a very obvious crest, no sign of which is seen in *acutirostris*. In the latter the rows of narrow elongate gulars abut on the labials; in *marmoratus* there are in between two or three rows of flat, well imbricate scales.

The middorsals of *marmoratus* are lanceolate, regular, and have a low, even keel, ending in a small swelling that may approximate a mucro. The scale rows are less regularly arranged toward the sacrum, but the shape of the individual scales varies little. In *acutirostris* the scales are irregular, very frequently truncated, with blunt, even swollen keels, or keelless; towards the back they may be practically smooth.

Much importance has been attached in the literature to the scales on the anterior part of the flanks. In typical cases *marmoratus* has more regular scales, with much less exposed skin between the rows. There is much variation in this character, though, and its use in keys has led to confusion.

The ventrals of *marmoratus* are regular, and have a low keel on the distal end, ending in a moderate mucro. In *acutirostris* they are narrower, with a complete high keel, but a less evident mucro. The general aspect is of a smooth surface in *marmoratus*, a ridged one in *acutirostris*.

A very interesting difference, apparently first noticed by Reinhardt and Lütken (1861), is in the shape of the femoral pores. In *marmoratus* each pore typically opens on the middle of a flat

scale; in *acutirostris* it notches the edge of the scale. The basic pattern is always obvious, but there is some variation in both species. In *marmoratus* this is not much: a few pores may sit on the edge of the scale. In *acutirostris* there is much more variation. In some extreme cases the pore-bearing scale is conical, raised, volcanolike; the pore, with prominent secretion plugs, is completely encircled, but a suture marks the place of the notch. In other cases the notch is open, narrow and deep, the scales assuming an irregular V shape; plugs of secretion may be present or not. In other cases the emarginations are so feeble that it is understandable that some authors have considered the pores as absent.

Color pattern. It appears that *P. marmoratus* has no sex dimorphism in color. There is considerable variation, even within single localities, but a general pattern is clearly discernible. I shall describe a live, healthy adult female from Oriximiná, Pará (not all authors have followed the same scheme in describing pattern, but all descriptions agree broadly).

The dorsal parts of the resting lizard were predominantly brown, the flanks barred with black, green and yellow, and the under parts greenish. The top of the head was havana brown with darker flecks, turning greenish towards the snout. The sides of the head were green. There were two black lines starting from the middle of the eye and running, respectively, towards the temporal edge and the corner of the mouth (these are very obvious in preserved animals).

The scapular region was dark green with dark brown spotting. On each flank there were six sets of oblique bars. The anterior element of each set was a thin black stripe meeting its fellow on the middle of the back. Each black stripe melted posteriorly into a green zone; the green zones grew broader (from six to twelve scales wide) towards the sacral region. The skin between the green scales was black. Behind each green

band there was a yellow one, two to four scales wide, with orange skin between the scales. On the middle of the back there were short dark bars between those that ran down the flanks. The members were greenish brown or vice-versa. The ventral parts were lettuce green, the color extending onto the base of the tail, the remainder of which was brown. The skin between the gulars was gray.

The tongue was orange, the mouth gray and the throat black. In a specimen described by Hoogmoed (1973) the whole mouth was black and the tongue pale orange.

Changes in the emotional or reproductive state of the animal cause the pattern to brighten or fade, from bright green to somber brown. The underparts may become whitish or light tan. Authors that have described metachrosis in this animal are Peracca (1895), Cott (1926), Mertens (1930), and Hoogmoed (1975).

Polychrus acutirostris (color Plates XXI and XXII in Vanzolini, Ramos-Costa, and Vitt, 1980) is a generally gray animal, with more or less distinct oblique bars in different tones of gray and brown, with wavy and frayed edges. The skin is the color of the surrounding scales. The two black lines found in *P. marmoratus*, radiating from the eye, also appear in *acutirostris*, but are usually very thin, at times almost imperceptible. I have not noted the color of the inside of the mouth.

A most important difference between the two forms is the fact that *acutirostris* presents sex dimorphism: the male in nuptial livery has two deep black spots surrounded with yellow on the anterior flank (Plate XXII in Vanzolini, Ramos-Costa, and Vitt, 1980).

Size and body proportions. Both species show sex dimorphism in snout to vent length, the females being larger, as noticed by Beebe (1944) and commented upon by Underwood (1962) and by Vitt and Lacher (1981). There seem to be no interspecific differences in body length. My largest *marmoratus* are, male

126 mm, female 144. Vitt and Lacher (1981) record respectively 124 and 145 mm for *acutirostris*.

The two species differ in relative tail length (as noted by Wagler, 1830; Wiegmann, 1834; Mertens, 1930) and in hind limb length (Boulenger, 1885; Mertens, 1930). The differences are quite real: for the same snout to vent length *marmoratus* has higher values of both measurements. The matter, however, is complicated by the presence of sex dimorphism in tail length in *marmoratus* and of geographic variation within both species; this should be the subject of specific research.

ECOLOGY AND BEHAVIOR

Habitat. It has been generally recognized that *P. marmoratus* is associated with rain forests, and the distributional data (see Fig. 2) permit no other interpretation. However, it is not absolutely certain what kind of habitat it prefers. It has been found inside the forest, on low trees and vines (Duellman, 1978; personal observation), and at least once in a crown situation (Rand apud Rand and Humphrey, 1968), but very frequently in the ecotone (Rand and Humphrey, 1968; Crump, 1971; Hoogmoed, 1973; Dixon and Soini, 1975; Duellman, 1978), in deciduous forest (Test, Sexton, and Heatwole, 1966), in secondary growth (Crump, 1971) and associated with water courses (Hoogmoed, 1973; Duellman, 1978).

The irregularity with which this lizard is taken makes me think it is primarily a crown animal, which at times comes down to the ecotone, which is ecologically similar to the crown environment, and where the animal is of course much more conspicuous.

Polychrus acutirostris, on the contrary has been collected in all types of open formations: cerrados, caatingas, fallow land, etc. It is usually found on low tree branches (Vitt and Lacher, 1981), but I

have captured one specimen on the ground.

It would thus seem that the essential ecological difference between the two species lies in the association with forest (*marmoratus*) as opposed to open formations (*acutirostris*), not in a preference for shade instead of exposed situations, as supposed by Mertens (1930, 1970).

Locomotion. The environments in which the two species have been collected have one thing in common: the substrate is predominantly twigs. It is thus not surprising that the two species are very similar in their locomotor behavior, which is rather striking. *P. marmoratus* has been studied in detail by Böker (1929, 1935), and many authors have figured and described the curious stances this lizard takes in the vegetation, especially on thin twigs.

One primary adaptation is the grasping capability of the feet (Böker, 11.cc.). Movement along or between branches may be remarkably slow (thence the Brazilian common names "sloth beast" and "blind lizard") or occasionally by means of very agile leaps. The animal may go through long periods of immobility, hanging in grotesque positions, with one or more limbs, or the tail, completely loose. On the ground the lizard may also be exasperatingly slow or unexpectedly agile (Test, Sexton, and Heatwole, 1966; Nicéforo, 1930; personal observation).

There is no comparable study of *acutirostris* in the literature, but Cott (1926), who wrote on *marmoratus*, noted that a specimen of *acutirostris* from the cerrado of Minas Gerais behaved in exactly the same way. I have had the opportunity of observing both *marmoratus* and *acutirostris* and, at the level of incidental observation, saw no major differences in locomotor behavior.

Food. *Polychrus marmoratus* has been said in the literature to be "omnivorous"; this is too broad a term, but in reality it has been known to eat small arthropods (insects of several orders, spiders) and vegetable food (leaves, fruits, seed—most

probably also flowers) (Beebe, 1944; Underwood, 1962; Rand and Humphrey, 1968; Hoogmoed, 1973; Duellman, 1978). *P. acutirostris* (see Vitt and Lacher, 1981 for a detailed study) has fundamentally the same habits.

Agonistic display. Full agonistic display in *Polychrus* in general (Breder, 1946; Underwood, 1962; Gorman, Huey, and Williams, 1969) comprises lateral flattening of the body, both of the dewlap and of the thorax (the latter aided by air sacs connected to the lungs), metachrosis, wide opening of the mouth, lunging and eventually biting. The same elements, and additionally bobbing are described for *P. acutirostris* by Vitt and Lacher (1981).

DISTRIBUTION

Materials. I found in the literature 111 definite locality citations of *P. marmoratus* and *acutirostris* (my own papers not included) which seemed reliable enough for mapping. I did not include localities in the Andes of Ecuador and Colombia because I feel that the situation there is not well understood (Peters, 1959; Gorman, Huey, and Williams, 1969; materials under study by myself). These areas, in any case, are not relevant to the present argument.

Discounting from the 111 citations the cases in which the same materials were cited twice and the localities represented by more than one collection, the literature yielded 95 individual localities.

In our own collection we have 364 *P. acutirostris* and 125 *marmoratus*, respectively from 114 and 45 localities. Some of our localities have already been cited in the literature. We therefore end up with 129 localities for *P. acutirostris* and 102 for *marmoratus*.

In drawing the map (Fig. 2), some areas were too densely covered for adequate representation: I had to omit many localities in the Distrito Federal of Venezuela, in the Peninsula de Paria, in northern Guyana, around Paramaribo, in the

Figure 2. Approximate distribution of *Polychrus marmoratus* (●) and *P. acutirostris* (○).

Iquitos area and especially in the state of São Paulo.

An important qualification refers to the southern limit of *acutirostris*. Koslowsky (1898) cites the species for Argentina (Gran Chaco, Misiones, Corrientes); he obviously saw specimens, but the references are vague. The only actual Argentinian specimens mentioned in the literature seem to be those from the Paraguayan border collected by the Deutsche Gran-Chaco-Expedition (Hellmich, 1960) and one from Misiones (no further data) cited by Burt and Burt (1931). The occurrence of the species in Uruguay is likewise in doubt, no specimens being mentioned by Vaz-Ferreira and Sierra de Soriano (1960) and Lema and Fabián-

Beurmann (1977). Finally, I know of no specimens from the three southernmost states of Brasil, Paraná, Santa Catarina, and Rio Grande do Sul.

Description. Inspection of the map shows one first important feature: the two species are parapatric. In fact, there are no intergrading specimens and no cases of sympatry. The complementarity between open and closed formations is practically perfect. The apparent mixture of localities in northeastern Brasil is due to the clearing of the Atlantic forest for agriculture. *P. acutirostris* very quickly occupies such spaces; this can also be very clearly seen in the state of São Paulo.

A second feature is that the distribution

of *marmoratus* is disjunct. It comprises two extensive continuous areas, one in the hylaea, up to the Caribbean coast, the other in the Atlantic forest, from Recife, in Pernambuco, to São Sebastião, in São Paulo. Between the two areas there is a broad gap in northeastern and central Brasil.

On the contrary, *P. acutirostris* occurs continuously from the northernmost cerrados and caatingas (Maranhão to Mato Grosso) to the Chaco, but is not known from the great savanna isolates of the North.

DISCUSSION

The problem posed by *Polychrus acutirostris* is by which mechanisms a species adapted and restricted to open formations arises from a forest-dwelling ancestor with which it maintains a parapatric relationship.

Vanzolini and Williams (1981) have proposed a variation of the orthodox model of geographic speciation, the "vanishing refuge," to account for such cases. The model starts with a widespread forest form at the humid peak of a climatic cycle. Deterioration of the climate brings about the dissection of the continuous forested areas by inroads of open formations. The forest isolates, initially large, progressively shrink in area. Some of them last the whole of the dry leg of the climatic cycle, constituting refuges in which forest fauna survives and undergoes geographic differentiation. Other forest isolates dwindle and finally vanish; normally the forest fauna would disappear with the refuge, but under certain circumstances, some such populations might survive, adapted to open formations and isolated from their relatives still faithful to the forest and inhabiting refuges. The populations surviving in open formations would be geographically isolated and under strong selective pressures, i.e., in ideal conditions for fast differentiation.

Let us see to what extent a simple model of this type would contribute to explain the appearance of *Polychrus acutirostris*. I begin by taking as putative parent the geographically most plausible relative, *P. marmoratus*.

The main prerequisite for a scenario such as the one outlined above is that the population outliving the vanishing refuge be pre-adapted to life in open situations. In the paper in which the model was first proposed (Vanzolini and Williams, 1981) and in others (e.g., Williams and Vanzolini, 1980) examples are given of incidental populations of forest-dwelling, umbrophilous species, living in ecotone or edge situations. In the case of *P. marmoratus* ecotonal populations seem to be at least extremely frequent, as stressed by authors working in the most diverse areas: the Napo in Ecuador (Duellman, 1978), the Iquitos region in Peru (Dixon and Soini, 1975), eastern Amazonia (Rand and Humphrey, 1968; Crump, 1971) and Surinam (Hoogmoed, 1973).

If the model is indeed viable, it is hard to see how *Polychrus* could avoid it.

The nature of the differences between *marmoratus* and *acutirostris* is not really essential to the argument, but affords a measure of support to it. The majority of these differences—body proportions, roughness of ventrals, color pattern—are obvious adaptations to life in cerrados and caatingas.

In these two types of plant formations the twig and crown niche is not otherwise occupied. The only lizard of similar ecology sympatric with *P. acutirostris* is the iguanid *Anisolepis grillii*, but the area of sympatry is the "cultural steppe" of the state of São Paulo, clearly under human influence. Another possibly syntopic species, *Anolis meridionalis*, a much smaller animal, is infrequent in the cerrados and absent from the caatingas.

Karyotypes. There is a fair amount of cytological work on *Polychrus*. The situation is highly interesting: "*Polychrus* remains a puzzle within the iguanids.

Now that three species have been studied it is quite clear that they are close to each other *inter se*, but distinctly different from anoles and anoline genera . . ." (Gorman, Huey, and Williams, 1969: 7). The three species then studied were *P. marmoratus*, *femoralis* and *peruvianus*. Later the karyotype of *acutirostris* was described by Peccinini (1970), under the name *marmoratus* (specimens from São Paulo) and by Beçak *et al.* (1972) under the name *Polychrus* sp. (In fact, it was the uncertainty in identifying these specimens with the help of current literature that led me to start this review; the problem of identification turned out to be exceedingly trivial, but the distribution captivating).

Paull, Williams, and Hall (1976: Table 4, p. 24) call *Polychrus* "the one egregious example of chromosomal diversity in a small genus" and thus summarize the situation: the diploid number varies from 20 (*acutirostris*) to 26 (*femoralis*) to 28 (*peruvianus*) to 30 (*marmoratus*).

Gorman, Atkins, and Holzinger (1967) demonstrated that the mechanism of sex determination in *P. marmoratus* is of the type $X_1X_2Y/X_1X_1X_2X_2$. Peccinini, Frota-Pessoa, and Ferrari (1971) showed that in *P. acutirostris* (also under the name *Polychrus* sp.) there was a particular case of the same mechanism ("pseudo XO/XX"), in which the Y chromosome is indistinguishable from X_2 . This is again strong indication that chromosomal rearrangements are frequent and obviously important in *Polychrus*.

It is possible, then, that we have chromosomal lability and extrinsic barriers to gene flow acting concurrently, a most favorable conjunction for fast rates of evolution. This is the mechanism I propose for consideration.

SOME OPEN QUESTIONS

This is a preliminary and speculative work that leaves many problems to be addressed.

An important problem is the nature of *P. marmoratus*, which brings forth the vexing question of disjunct populations that cannot be sorted out morphologically. In reality, we are talking about evolution at the species level, but our concept of species is extremely pragmatic and its application never more than provisional.

Another problem is that of time scale. If there has been differentiation within *Polychrus* dependent on climatic cycles, then it is certainly a Quaternary event, and this should be taken into account in discussions of evolutionary rates concerning chromosomal rearrangements (Williams and Hall, in Paull, Williams, and Hall, 1976).

That *acutirostris* does not reach the open formations north of the Amazon is probably due to its having arisen south of it. There are many open formation forms (e.g., *Crotalus*, *Tropidurus*) that occur on both sides of the valley, and even in small enclaves of open formations within the hylaea. Others (e.g., the extremely successful bird *Cariama*) do not. Why this should be so remains a puzzle.

Finally, it becomes mandatory to explore in depth the differentiation of *Polychrus* in the western end of its range, where the influence of Andean refuges (Haffer, 1969; Vanzolini and Williams, 1970) is apt to be strongly felt, and where the relationships between the several forms are anything but clear. There is in fact no reason to believe *a priori* that at any given time an animal group is subjected to only one speciation mechanism.

ACKNOWLEDGMENTS

This is one of the very few pieces of research in the last thirty-odd years which I did not do in close rapport with Ernest E. Williams. That the stamp of our collaboration is clearly on it enhances my pleasure in this dedication. I hope Ernest, while as usual disbelieving the results, will in some measure enjoy in the

paper the elements of uncertainty, ecotones, karyotypes and some contrariness.

LITERATURE CITED

- AMARAL, A. 1937. Check-list of the Lacertilia of Brazil. C.R. XII Congr. Int. Zool. (Lisbonne 1935), 3: 1733-1743.
- ANDERSSON, L. G. 1900. Catalogue of the Linnean type specimens of Linnaeus's Reptilia in the Royal Museum in Stockholm. Bih. t. K. Svenska Vet.-Akad. Handl. 24 (4) art. 1: 29 pp.
- BEČAK, M. L., W. BEČAK, AND L. DENARO. 1972. Chromosome polymorphism, geographical variation and karyotypes in Sauria. Caryologia, 25(3): 313-326.
- BEEBE, W. 1944. Field notes on the lizards of Katabo, British Guiana, and Caripito, Venezuela. II. Iguanidae. Zoologica, New York, 29: 195-216.
- BERTHOLD, A. A. 1846. Ueber verschiedene neue oder seltene Reptilien aus Neu-Granada und Crustaceen aus China. Abh. K. Ges. Wiss. Goettingen, 3: 1-16.
- BÖKER, H. 1929. Einige Ergebnisse einer biologisch-anatomisch Forschungsreise nach Nord-Brasilien. Anat. Anz., 67 (Ergänzungsheft): 9-20.
- . 1935. Einführung in die vergleichende biologische Anatomie der Wirbeltiere. 2 volumes. Jena, Gustav Fischer.
- BOIE, H. 1826. Bemerkungen über die von Hr. von Spix abgebildeten brasilianischen Saurier. Isis von Oken, 18: 117-120.
- BOULENGER, G. A. 1885. Catalogue of the lizards in the British Museum (Natural History). Second edition. Vol. 2. London, Trustees of the British Museum, xiii + 497 pp., pls.
- . 1908. Descriptions of new South-American reptiles. Ann. Mag. Nat. Hist., 1(8): 111-115.
- . 1914. On a second collection of batrachians and reptiles made by Dr. H.G.F. Spurrell, F.Z.S., in the Choco, Colombia. Proc. Zool. Soc. London, 1914: 813-817.
- BURT, C. E., AND M. D. BURT. 1931. South American lizards in the collection of the American Museum of Natural History. Bull. Amer. Mus. Nat. Hist., 61: 227-395.
- BURT, C. E., AND M. D. BURT. 1933. A preliminary check list of the lizards of South America. Trans. Acad. Sci. St. Louis, 28(1-2): 1-104.
- BREder, C. M., JR. 1946. Amphibians and reptiles of the Rio Chucunague drainage, Darien, Panama, with notes on their life histories. Bull. Amer. Mus. Nat. Hist., 86(8): 375-436.
- COPE, E. D. 1899. Catalogue of a collection of Reptilia from Argentina. Sci. Bull. Philadelphia Comm. Mus., 1: 19-21.
- COTT, H. B. 1926. Observations on the life-habits of some batrachians and reptiles from the lower Amazon; and a note on some mammals from Marajó Island. Proc. Zool. Soc. London, 1926: 1159-1178.
- CRUMP, M. 1971. Quantitative analysis of the ecological distribution of a tropical herpetofauna. Univ. Kansas, Mus. Nat. Hist. Occ. Pap. 3: 62 pp.
- CUVIER, G. 1817. Le Règne Animal, distribué d'après son organisation. Tome II. Paris, Deterville, xviii + 532 pp.
- DAUDIN, F. M. 1802. Histoire Naturelle générale et particulière des reptiles. Vol. 3. Paris, F. Dufart, 452 pp. (For date see Vanzolini, 1977).
- DELAPORTE, F. 1826. Description d'une nouvelle espèce de reptile du genre Marbré (*Polychrus*). Ann. Sci. Nat., Paris, 9: 110-111.
- DIXON, J. R., AND P. SOINI. 1975. The reptiles of the Upper Amazon Basin, Iquitos Region, Peru. I. Lizards and amphisbaenians. Milwaukee Publ. Mus. Contrib. Biol. Geol., 4: 58 pp.
- DUELLMAN, W. E. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. Univ. Kansas Mus. Nat. Hist., Misc. Publ. 65: 352 pp.
- DUMÉRIL, A. M. C. AND G. BIBRON. 1837. Erpétologie générale. Vol. 4. Paris, Librairie Encyclopédique de Roret, ii + 571 pp.
- DUMÉRIL, C., AND A. DUMÉRIL. 1851. Catalogue méthodique de la collection des Reptiles (Muséum d'Histoire Naturelle de Paris). Paris, Gide et Baudry, iv + 224 pp.
- FITZINGER, L. 1826. Neue Classification der Reptilien... Wien, J. G. Heubner, 66 pp.
- . 1827. Recension des Spixischen Eidechsenwerkes. Isis von Oken, 20: 741-750.
- GMELIN, J. F. 1789. Caroli a Linné Systema Naturae... editio decima tertia, aucta, reformata, cura... Tomus I Pars III: 1035-1514 Lipsiae: G. E. Beer. (For date see Hopkinson, 1908.)
- GORMAN, G. C., L. ATKINS, AND T. HOLZINGER. 1967. New karyotypic data on 15 genera of lizards in the family Iguanidae with a discussion of taxonomic and cytological implications. Cytogenetics, 6: 286-299.
- GORMAN, G. C., R. B. HUEY, AND E. E. WILLIAMS. 1969. Cytotaxonomic studies on some unusual iguanid lizards assigned to the genera *Chamaeleolis*, *Polychrus*, *Polychroides*, and *Phenacosaurus*, with behavioral notes. Breviora Mus. Comp. Zool. No. 316, pp. 1-17.
- GRAY, J. E. 1845. Catalogue of the specimens of lizards in the collection of the British Museum. London, Trustees of the British Museum, xxviii + 289 pp.
- GUIBÉ, J. 1954. Catalogue des types de lézards du Muséum National d'Histoire Naturelle. Bayeux, Colas, 119 pp.
- HAFFER, J. 1969. Speciation in Amazonian forest birds. Science, 165 (3889): 131-137.
- HALLOWELL, E. 1845. Description of reptiles from South America, supposed to be new. Proc. Acad. Nat. Sci. Philadelphia, 2: 241-247.
- HELLMICH, W. 1960. Die Sauria der Gran-Chaco

- und seiner Randgebiete. Abh. Bayer. Akad. Wiss. (N.F.), **101**: 1-131.
- HOLM, A. 1957. Specimina Linnaeana i Uppsala bevarade zoologiska sammlingar fran Linnés tid. Uppsala Univ. Arsskr., **1957** (6): 1-68.
- HOOGMOED, M. S. 1973. Notes on the herpetofauna of Surinam IV. The lizards and amphisbaenians of Surinam. The Hague, W. Junk, ix + 419 pp.
- . 1975. De hagedissen en wormhagedissen van Suriname II: de leguanen, de skinken. Lacerta, **33** (6): 87-104.
- HOPKINSON, J. 1908. Dates of publication of the separate parts of Gmelin's edition (13th) of the 'Systema Naturae' of Linnaeus. Proc. Zool. Soc. London, **1907**: 1035-1037.
- KOSLOWSKY, J. 1898. Enumeración sistemática y distribución geográfica de los réptiles argentinos. Rev. Mus. La Plata **8**: 161-200, pls.
- LACÉPÈDE, B. G. E. L., COMTE DE, 1788. Histoire naturelle des quadrupèdes ovipares et des serpens. Tome premier. Paris, Imprimerie du Roi, xviii + 651 pp.
- LEMA, T., AND M. E. FABIÁN-BEURMANN, 1977. Levantamento preliminar dos répteis da região da fronteira Brasil-Uruguai. Iheringia (Zool.), Porto Alegre, **50**: 61-92.
- LINNÉ, C. 1758. Systema Naturae, per regna tria naturae . . . Editio decima, reformata. Holmiae, Laurentii Salvii (Cura Societatis Zoologicae Germanicae iterum edita, 1894. Lipsiae, Wilhelm Engelmann), 824 pp.
- LÖNNBERG, L. G. 1896. Linnean type-specimens of birds, reptiles, batrachians and fishes in the Zoological Museum of the R. University in Upsala. Bih. t. K. Svenska Vet.-Akad. Handl., **22** (4) art., 1: 45 pp.
- MERTENS, R. 1930. Zur Biologie der Iguanidengattung *Polychrus* G. Cuvier. Zool. Gart., **3** (9-10): 269-279.
- . 1970. Reptilienauge und Umwelt. Natur und Museum, **100** (10): 435-446.
- NICÉFORO MARIA. 1930. Los reptiles de Villavicencio en el museo de La Salle. Rev. Soc. Colomb. Ci. Nat., **19** (195): 40-54.
- NOBLE, G. K. 1924. New lizards from northwestern Peru. Occ. Pap. Boston Soc. Nat. Hist., **5**: 107-113.
- PARKER, H. W. 1935. The frogs, lizards and snakes of British Guiana. Proc. Zool. Soc. London, **1935**: 505-530.
- PAULL, D., E. E. WILLIAMS, AND W. P. HALL. 1976. Lizard karyotypes from the Galapagos Islands: chromosomes in phylogeny and evolution. Breviora Mus. Comp. Zool. No. 441, pp. 1-31.
- PECCININI, D. 1970. Variação dos cromossomos do lagarto *Polychrus marmoratus* (Sauria, Iguanidae) de diferentes localidades (nota preliminar). Rev. Brasil. Biol., **30** (1): 1-4.
- PECCININI, D., O. FROTA-PESSOA, AND I. FERRARI. 1971. Sex determination of the "pseudo XO/XX" type in the Brazilian lizard *Polychrus* sp. (Sauria, Iguanidae). Caryologia, **24** (1): 129-139.
- PERACCA, M. G. 1895. Viaggio del dott. Alfredo Borelli nella Repubblica Argentina e nel Paraguay. X. Rettili ed anfibi. Boll. Mus. Torino, **10** (195): 32 pp.
- PERRACCA, M. G. 1897. Viaggio del Dr. Enrico Festa nell'Ecuador e regioni vicine. IV. Rettili. Boll. Mus. Torino, **12** (300): 20 pp.
- PETERS, J. A. 1959. Notas miscelâneas sobre saurios del Ecuador. Ci. Nat., Quito, **2** (3): 118-124.
- PETERS, J. A., AND R. DONOSO-BARROS. 1970. Catalogue of the Neotropical Squamata. Part II. Lizards and amphisbaenians. Bull. U.S. Nat. Mus., **297**: viii + 293 pp.
- PETERS, W. 1877. Über die von Spix in Brasilien gesammelten Eidechsen des Königlichen Naturalien-Kabinetts zu München. Monatsber. Akad. Wiss. Berlin, **1877**: 407-414.
- RADDI, G. 1820. Di alcune specie nuovi di rettili e piante brasiliane. Atti. Soc. Ital. Sci., Modena, **18**: 1-39, pls.
- . 1822. Continuazione della descrizione dei rettili brasiliani indicati nella memoria inserita nello secondo fascicolo delle memorie di fisica del precedente volume XVIII. Atti Soc. Ital. Sci., Modena, **19**: 56-73.
- RAND, A. S., AND S. S. HUMPHREY. 1968. Interspecific competition in the tropical rain forest: ecological distribution among lizards at Belém, Pará. Proc. U.S. Nat. Mus., **125** (3658): 17 pp.
- REINHARDT, J., AND C. LÜTKEN. 1861. Bidrag til Kundskab om Brasiliens Padder og Krybdyr. Første Afdeling, Padderne og Öglerne. Vid. Medd. Nat. For. Copenhagen, **1861**: 143-242, pls.
- ROZE, J. A. 1958. On Hallowell's type specimens of reptiles from Venezuela in the collection of the Academy of Natural Sciences of Philadelphia. Notulae Naturae, **309**: 4 pp.
- SCHINZ, H. R. 1822. Das Thierreich eingetheilt nach dem Bau der Thiere als Grundlage ihrer Naturgeschichte und der vergleichende Anatomie von dem Herrn Ritter von Cuvier. Vol. 2: 1-189. Stuttgart und Tübingen: J. G. Cotta.
- SCHMIDT, K. P., AND R. INGER. 1951. Amphibians and reptiles of the Hopkins-Branner Expedition to Brazil. Fieldiana Zool., **31** (42): 439-465.
- SPIX, J. B. 1825. Animalia nova sive Species novae lacertarum quas in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspicio Maximiliani Josephi I Bavariae Regis suscepto collegit et descripsit Dr. J. B. de Spix. Lipsiae, T. O. Weigel, 26 pp., pls.
- . Berichtigung der Bemerkungen des Herrn Dr. Boie in Leyden ueber die von Herrn Dr. Spix abgebildet Saurier (Isis 1826 Heft 1. S. 117). Isis von Oken, **18**: 601-604.
- TEST, F. H., O. J. SEXTON, AND H. HEATWOLE. 1966.

- Reptiles of Rancho Grande and vicinity, Estado Aragua, Venezuela. Misc. Publ. Mus. Zool. Univ. Michigan 128: 63 p.
- UNDERWOOD, G. 1962. Reptiles of the Eastern Caribbean. Caribbean Affairs (N.S.), 1: 192 pp.
- VANZOLINI, P. E. 1948. Notas sobre os ofidios e lagartos da Cachoeira de Emas, no municipio de Pirassununga, estado de S. Paulo. Rev. Brasil. Biol., 8 (3): 377-400.
- . 1977. An annotated bibliography of the land and fresh-water reptiles of South America. Vol. 1 (1758-1900). São Paulo, Museu de Zoologia da Universidade de São Paulo, iv + 186 pp.
- . 1978. An annotated bibliography of the land and fresh-water reptiles of South America. Vol. 2 (1901-1975). São Paulo, Museu de Zoologia da Universidade de São Paulo, 316 pp.
- VANZOLINI, P. E., A. M. M. RAMOS-COSTA, AND L. J. VITT. 1980. Repteis das caatingas. Rio de Janeiro, Academia Brasileira de Ciências, 161 pp., pls.
- VANZOLINI, P. E., AND E. E. WILLIAMS. 1970. South American anoles: the geographic differentiation and evolution of the *Anolis chrysolepis* species group (Sauria, Iguanidae). Arq. Zool. S. Paulo, 19 (1-4): 298 pp., pl.
- AND ———. 1981. The vanishing refuge: a mechanism for ecogeographic speciation. Papéis Avulsos Zool., S. Paulo, 34 (23): 251-256.
- VAZ-FERREIRA, R., AND B. SIERRA DE SORIANO. 1960. Notas sobre reptiles del Uruguay. Rev. Fac. Human. Ci. Montevideo, 18: 133-206.
- VITT, L. J., AND T. E. LACHER JR. 1981. Behavior, habitat, diet, and reproduction of the iguanid lizard *Polychrus acutirostris* in the caatinga of northeastern Brazil. Herpetologica, 37 (1): 53-63.
- WAGLER, J. 1821. Ankündigung: Die Amphibien. Isis von Oken, 8: 337-342.
- . 1830. Natürliches System der Amphibien . . . München, Stuttgart und Tübingen, J. G. Cotta, vi + 354 pp.
- . 1833a. Descriptiones et icones amphibiorum. Pars tertia. Monachii, Stuttgartiae et Tubingae, J. G. Cotta, (For date see Vanzolini, 1977).
- . 1833b. Deutung die in Sebas's Thesauri Rerum Naturalium T. 1. et 2. enthaltenen Abbildungen von Lurchen, mit kritische Bemerkungen. Isis von Oken, 1833 (9): 896-905.
- WERNER, F. 1910. Ueber neue oder seltene Reptilien des Naturhistorischen Museums in Hamburg II. Eidechsen. Mitteil. Naturhist. Mus. Hamburg, 27: 1-46.
- WIED-NEUWIED, MAXIMILIAN, PRINZ ZU. 1820. Reise nach Brasilien in den Jahren 1815 bis 1817. Vol. 1. Frankfurt a.M., H. L. Brönnner.
- . 1825. Beiträge zur Naturgeschichte von Brasilien. Vol. 1. Weimar, G.R.H.S. priv. Landes-Industrie-Comptoirs.
- . 1822-1831. Abbildungen zur Naturgeschichte Brasiliens: Weimar. 2 volumes (For dates see Vanzolini, 1977).
- WIEGMANN, A. F. A. 1834. Herpetologia Mexicana . . . Berlin, C. G. Lüderits, vi + 54 pp.
- WILLIAMS, E. E., AND P. E. VANZOLINI. 1980. Notes and biogeographic comments on anoles from Brasil. Papéis Avulsos Zool. S. Paulo, 34 (6): 99-108.

Systematics of the Latin American Snake, *Liophis epinephelus* (Serpentes: Colubridae)

JAMES R. DIXON¹

ABSTRACT. *Liophis epinephelus* is discussed in light of the 14 names associated with the species. Four currently recognized species, *L. bimaculatus*, *L. pseudocobella*, *L. fraseri*, and *L. albiventris* are reduced to subspecies of *L. epinephelus*. The known variation and distribution of *L. epinephelus* is presented; its possible origin is discussed considering past historical events.

INTRODUCTION

The widespread and highly variable Latin American snake, *Liophis epinephelus*, is the most enigmatic of all *Liophis* species. It has the greatest altitudinal distribution of the genus, sea level to 3,400 m. The species is primarily Trans-Andean, occurring in the lowlands of Panama, Colombia, and Ecuador; but also in foothills and mountains of Costa Rica, Panama, Colombia, Venezuela, Ecuador, and Peru (Figs. 1-4). Each mountain range and/or major vegetation community may contain a population of *L. epinephelus* with somewhat different characteristics than adjacent ones. This is evidenced by the various color pattern morphs from the northern, northwestern, and western Andes, the Trans-Andean lowlands of Colombia and Ecuador, lowlands of Panama, and mountains of Costa Rica (Figs. 5, 6). These morphs resulted in published descriptions of *Liophis epinephelus* (Cope, 1862), *L. reginae albiventris* (Jan, 1863), *L. reginae quadri-*

lineata (Jan, 1863), *L. alticolus* (Cope, 1868), *Zamensis ater* (Günther, 1872), *L. fraseri* (Boulenger, 1894), *L. bimaculatus* (Cope, 1899), *L. opisthotaenia* (Boulenger, 1908), *L. pseudocobella* (Peracca, 1914), *L. cobella alticolus* (Amaral, 1931), *L. taeniurus juvenalis* (Dunn, 1937), and *L. bimaculatus lamonae* (Dunn, 1944). Some quantitative characters (preoculars, ventrals) associated with changes in elevation resulted in the descriptions of *L. bipraeocularis* (Boulenger, 1903) and *L. epinephelus ecuadorensis* (Laurent, 1949).

Of the 14 published names, five are currently recognized as full species: *albiventris*, *bimaculatus*, *fraseri*, *pseudocobella*, and *epinephelus*. In this paper, I reduce the first four to subspecies of *epinephelus*.

A definition of *L. epinephelus* includes most quantitative and color pattern characteristics of all species of *Liophis*. Scale rows are typically 17-17-15, smooth, one apical pit; usually one preocular (occasionally two); two postoculars (occasionally one or three); one loreal; one anterior temporal (occasionally none); two posterior temporals (occasionally one or three); fourth and fifth supralabial entering orbit, infrequently third and fourth or 10 other combinations; supralabials normally eight, occasionally nine other combinations; infralabials typically ten, occasionally 11 other combinations (Tables 1, 2). The number of ventrals vary from 128 to 191; subcaudals 44 to 80; maxillary teeth 17 to 29; total length/tail ratios .17 to .27; hemipenial length 7 to 13

¹Department of Wildlife & Fisheries Sciences, Texas A&M University, College Station, Texas 77843, U.S.A.

Figure 1. Distribution of *Liophis epinephelus juvenalis* (black circle), *L. e. epinephelus* (open circles), and their intergrades (open circles with X) in Costa Rica and Panama. Insert illustrates the zone of contact between the two subspecies along the Costa Rica/Panama border. The dashed line represents the approximate position of the 1,500 m contour line.

subcaudals; total maximum length 775+ mm, ♀♀, 700+ mm, ♂♂ (both with incomplete tails).

Color patterns vary by population. Some are banded red and black dorsally and ventrally; some are striped pos-

teriorly with two or four black stripes on an olive, green, or brown ground color; some are totally leaf green dorsally and pale yellow ventrally; others may have combinations of black and yellow or red checkered venters with two black dorsal

Figure 2. Distribution of *L. epinephelus* and its subspecies in the vicinity of the Andes of Colombia and western Venezuela. Asterisk = *L. e. opisthotaenia*; Black square = *L. e. lamonae*; Black star = *L. e. epinephelus*; Black circle = *L. e. bimaculatus*; Black eight point star = *L. e. pseudocobella*; Black circle with a white star = sympatry of two and four-lined patterns and possible intergrades between the subspecies *bimaculatus* and *lamonae*; Open square with a black dot = intergrades between the subspecies *pseudocobella* and *lamonae*; Open square with open star = intergrades between subspecies *pseudocobella* and *epinephelus*; Open square with black star = intergrades between the subspecies *pseudocobella* and *bimaculatus*; stippled zone represents elevations above 1,400 m.

Figure 3. Ecuadorian distribution of *L. epinephelus fraseri* (black square), *L. e. lamonae* (open circle with black dot), *L. e. albiventris* (black circle), *L. e. bimaculatus* (open circle with black triangle), *L. e. epinephelus* × *e. albiventris* (open circle with black star), and *L. e. bimaculatus* × *e. fraseri* or *e. albiventris* (half white/black circle). Dotted line represents the approximate 1,500 m contour.

Figure 4. Distribution of samples of *L. epinephelus* by latitude in Latin America. Numbers in circles correspond to map code numbers and subspecies given in Table 2 and Figure 5.

TABLE 1. VARIATION IN SUPRALABIALS, INFRALABIALS, AND SUPRALABIALS ENTERING ORBIT OF *L. EPINEPHELUS*.

SUPRALABIALS	5-6	6-6	6-7	6-8	7-7	7-8	8-8	8-9	9-9	10-10		
<i>e. juvenalis</i>							49	1	1	1		
<i>e. juvenalis</i> × <i>epinephelus</i>							33					
<i>e. epinephelus</i>							99	4				
<i>e. pseudocobella</i>						2	29					
<i>e. opisthotaenia</i>			1		8	9	14					
<i>e. bimaculatus</i>	1		1		8	5	22					
<i>e. albiventris</i>		1			3		86	1				
<i>e. lamonae</i>	1		1			1	39					
<i>e. fraseri</i>			1	1	2	6	33					
INFRALABIALS	7-7	7-8	7-9	8-8	8-9	8-10	9-9	9-10	9-11	10-10	10-11	11-11
<i>e. juvenalis</i>						1		5		42	2	3
<i>e. juvenalis</i> × <i>epinephelus</i>								2		28	2	1
<i>e. epinephelus</i>					2	1	5	14		76	2	2
<i>e. pseudocobella</i>								11	4	1	13	1
<i>e. opisthotaenia</i>				5	5			19	2		1	
<i>e. bimaculatus</i>	1	1	1	6	3			17	5		4	
<i>e. albiventris</i>		1		2	3			10	7		66	2
<i>e. lamonae</i>			1	3	5			6	7		17	1
<i>e. fraseri</i>				2	1	1	19	11		12		
SUPRAL/ORBIT	3rd	3+4	$\frac{3+4}{4+5}$	$\frac{3+4+5}{4+5}$	3+4+5	4+5	$\frac{4+5}{4+5+6}$	4+5+6	$\frac{4+5}{5+6}$	$\frac{4th}{4+5}$	5+6	
<i>e. juvenalis</i>						48			2	1		1
<i>e. juvenalis</i> × <i>epinephelus</i>						33						
<i>e. epinephelus</i>						87				3		
<i>e. pseudocobella</i>						29						
<i>e. opisthotaenia</i>		6	9		2	14						
<i>e. bimaculatus</i>	1	5	4	1		29					1	
<i>e. albiventris</i>		1				92	1					
<i>e. lamonae</i>			3			46						
<i>e. fraseri</i>		4	6			36						

nape spots, obscure dorsal diagonal dark markings, and black tail stripes. Some high altitudinal populations tend to have 75% of the venter black. Intergrade zones between populations typically have combinations of patterns and these frequently aid in allocating the intergrades.

nephelus. To avoid confusion and properly allocate all published names from the past 100 years, the following synonymies, available names, and distributions are listed for the subspecies of *L. epinephelus*.

1) *Liophis epinephelus epinephelus* Cope 1862

Lower elevations of mountains of western Panama east to Colombian lowlands and most Colombian inter-Andean valleys to elevations of 1,500 m; also south to extreme northern Ecuador along Pacific coast.

ALLOCATION OF NAMES

Rather than discussing each subspecies individually, I have discussed them collectively under an intraspecies variation section. There are eight recognizable taxa associated with the species *epi-*

TABLE 2. RANGE OF VARIATION OF VENTRALS AND SUBCAUDALS OF *L. EPINEPHELUS* BY LATITUDE.

LATITUDE		N	R	M	SE	N	R	M	SE	MAP CODE
10-11° N	(J)*	18	139-148	143.9	0.57	18	52-62	54.5	0.68	1a
	(E)	4	138-152	144.8	3.00	4	60-66	63.5	1.50	1b
9-10° N	(J)	34	139-152	145.7	0.50	30	51-61	56.5	0.42	2a
	(E)	35	133-142	136.9	0.60	33	54-67	60.0	0.54	2b
	(O)	29	142-152	147.0	0.53	29	53-78	64.7	0.92	2c
8-9° N	(J)	2	139-140	139.5	—	2	59-63	61.0	—	3a
	(J/E)	33	132-148	140.0	0.50	30	53-62	58.3	0.50	3b
	(E)	13	136-147	139.7	0.94	14	55-64	60.6	0.64	3c
	(E)	3	134-144	138.0	—	3	56-66	60.0	—	3d
6-7° N	(E)	7	132-141	137.9	1.10	7	61-69	64.3	1.00	4a
	(L)	5	141-150	144.8	1.60	5	54-61	57.4	1.10	4b
	(P)	38	143-151	146.4	0.35	35	48-54	51.2	0.31	4c
5-6° N	(E)	24	128-149	137.1	0.82	23	49-72	62.6	1.10	5a
	(L)	3	148-153	150.3	—	3	53-55	54.0	—	5b
	(P)	1	149	—	—	1	45	—	—	5c
	(B)	1	171	—	—	1	68	—	—	5d
4-5° N	(E)	2	142	—	—	2	61-67	64.0	—	6a
	(L)	15	146-156	150.0	0.96	15	57-67	63.4	0.70	6b
	(P)	7	135-146	141.8	1.50	5	47-53	48.8	1.10	6c
	(B)	30	165-184	174.2	0.90	24	56-80	63.3	0.97	6d
3-4° N	(E)	5	134-143	138.6	1.60	5	64-71	68.6	1.00	7a
	(L)	1	156	—	—	1	64	—	—	7b
2-3° N	(L)	2	147-149	148.0	—	2	59	—	—	8a
	(P)	9	149-158	151.8	1.00	9	51-57	54.0	0.78	8b
	(B)	1	169	—	—	1	62	—	—	8c
1-2° N	(E)	2	141-151	147.5	—	2	66-70	68.0	—	9
0-1° N	(A)	20	145-164	152.3	1.10	19	55-70	61.4	1.00	10a
	(L)	1	146	—	—	1	68	—	—	10b
	(B)	3	185-188	186.0	—	1	69	—	—	10c
0-1° S	(A)	39	143-165	151.7	0.80	36	53-68	59.9	0.65	11a
	(L)	2	154-156	155.0	—	2	60-61	60.5	—	11b
2-3° S	(A)	6	143-154	149.2	1.60	6	55-66	60.1	1.70	12a
	(L)	2	143-146	144.5	—	2	51-57	54.0	—	12b
3-4° S	(A)	7	141-148	143.7	1.10	7	54-68	61.1	1.70	13
4-5° S	(F)	28	151-162	155.5	0.50	28	61-74	67.7	0.70	14
7-8° S	(F)	11	153-164	159.2	1.00	10	61-76	67.2	1.40	15
12° S	(F)	2	145-154	149.5	—	2	53-56	54.5	—	16

*Letter in parenthesis indicates first letter of subspecies name to which the sample is referred.

A = *albiventris*; B = *bimaculatus*; E = *epinephelus*; F = *fraseri*; J = *juvenalis*; L = *lamonae*; O = *opisthotaenia*; P = *pseudocobella*; J/E = *juvenalis* × *epinephelus*. (See Fig. 4 for map localities.)

2) *Liophis ephinephelus albiventris* Jan 1863

Liophis reginae quadrilineata Jan 1863
Opheomorphus alticolus Cope 1868
Zamensis ater Günther 1872

Western Ecuador from sea level to 2,600 m in the Andean valleys.

3) *Liophis epinephelus fraseri* Boulenger 1894

Middle elevations of the eastern and

western Andean slopes of southern Ecuador, south to central Peru.

4) *Liophis epinephelus bimaculatus* Cope 1899

Liophis bipraeocularis Boulenger 1903
Leimadophis epinephelus ecuadorensis Laurent 1949

High (2,600-3,300 m) Andean valleys of western Venezuela, central Colombia, south to northern Peru.

5) *Liophis epinephelus opisthotaenia*
Boulenger 1908

Merida region of Venezuela and
Páramo de Tama region of Venezuela
and Colombia.

6) *Liophis epinephelus pseudocobella*
Peracca 1914

Liophis cobella alticolus Amaral 1931

Middle elevations of central and
western portions of Colombian
Andes south to Ecuador border.

7) *Liophis epinephelus juvenalis* Dunn
1937

Middle slopes of most Costa Rica
mountains southeast to western
Panama.

8) *Liophis epinephelus lamonae* Dunn
1944

Eastern slopes of Andes of Colombia
from 1,500 to 2,600 m, southward to
east central Ecuador.

Liophis reginae albiventris and *L. reginae quadrilineata* represent color pattern morphs of west Ecuadorian *L. epinephelus*; the morphs show a complex pattern of sympatry and allopatry in Ecuador. About 98% of western Ecuadorian lowland *epinephelus* have immaculate cream or yellow venters and a black lateral posterior body stripe extending to the tip of the tail. Occasionally juveniles and adults possess paravertebral black stripes beginning at midbody or more posteriorly, in addition to the lateral black stripe characteristic of the *quadrilineata* morph. With increasing altitude, the frequency of four black stripes does not increase, whereas the number of ventrals do. *Liophis alticolis* is identical to *L. albiventris* with the exception of the abnormal single postocular. *Zamensis ater*, purportedly from Biscar, Algeria, represents "blackened" specimens of *Liophis albiventris*, probably from western Ecuador.

Liophis bimaculatus from the Sabana region of Bogota, Colombia, normally has four posterior black stripes, occasionally

two, infrequently none. The venter may be immaculate, dusky, black and yellow checkered, or almost entirely black. White ventered *bimaculatus* are very similar to *albiventris* in color pattern but are distinguished by their high numbers of ventrals and persistence of four black stripes.

Liophis fraseri and *L. bimaculatus lamonae* represent allopatric populations of *L. epinephelus* from alternate slopes of the middle elevations of the eastern, western, and/or southern flanks of the Andes.

The holotype of *L. bipraeocularis* is a two preocular variation of *L. e. bimaculatus*, and the holotype of *L. epinephelus ecuadorensis* is a patternless example of *L. e. bimaculatus*.

Liophis opisthotaenia and *L. taeniurus juvenalis* are examples of two (semi-isolated) mountain populations of *L. epinephelus* to the west and east of the nominate subspecies' major distribution.

Liophis pseudocobella and *L. cobella alticolus* represent the same population of *L. e. pseudocobella* from near Medellín, Colombia.

ZOOGEOGRAPHICAL COMMENTS

The evidence at hand suggests there were two separate invasions of *propinephelus* stock in the western and central Andes of Colombia. The *propinephelus* stock probably crossed the Andes in the Quaternary, at or near the Huancabamba Deflection zones of northern Peru. This invasion through the Andes left some populations behind that became isolated and/or adapted for higher elevations. Those that passed through the portal and distributed themselves along the Pacific lowlands moved northward into favorable environments, eventually reaching northern Costa Rica. The harsh deserts to the south prevented distribution in that direction. Those populations that remained at or near the upper slopes of the Andes moved both

north and south, but at a slower rate than the Pacific lowland population. The extremely cool environments of the upper slopes probably delayed adaptation to and reproduction in such environments. By the time the upper slope populations reached the highlands of northern Colombia, the lowland Pacific populations had established themselves along the middle slopes, eventually evolving into the *pseudocobella* and *juvenalis* races of the western slopes of Colombia and Costa Rica. The lowland populations adapted to the warmer wet and dry environments, eventually evolving into the *epinephelus* (warm-wet) and *albiventris* (warm-dry) subspecies.

The populations of the higher slopes evolved into a two and four-lined pattern population; those that continued to expand their range into the very high slopes evolved into the four-lined subspecies with high numbers of ventrals (*bimaculatus*), and those that remained along the higher middle slopes evolved into the subspecies *fraseri*, *lamonae* and *opisthotenia*. The latter two races were probably isolated by the Tachira Depression in northern Venezuela and evolved allopatrically.

INTRASPECIES VARIATION

The eight populations are recognized as subspecies by a combination of color, pattern, squamation, and to some extent, maxillary teeth (Fig. 7). Each population exhibits a unique feature which may represent a major environmental adaptation. The major shift in squamation, such as the number of ventrals and subcaudals, or meristic features such as tail length, are associated with cooler environments (higher elevations) and Pleistocene orogeny that resulted in geographic isolation. For example, *L. e. juvenalis* and *L. e. pseudocobella* possess a very similar color pattern of red and black bands encircling the body and the absence of a posterior lateral black line

that extends to the tip of the tail. There are 32 to 49 ($\bar{x}$ 41.4) black body bands in *juvenalis* and 28 to 42 ($\bar{x}$ 37.4) in *pseudocobella*. However similar they are in color pattern, number of ventrals, maxillary teeth and tail/total length ratios, the two taxa are still significantly distinct in the number of subcaudals (Table 3, Fig. 8). *Liophis e. pseudocobella* occurs between 1,500 and 2,000 m along the middle slopes of the central and western Andes of Colombia; *L. e. juvenalis* occurs between 650 and 2,100 m along both slopes of the mountains of Costa Rica and western Panama.

There is little doubt that *L. e. epinephelus* gave rise to both populations. *L. e. epinephelus* occurs in the lowlands between *e. juvenalis* and *e. pseudocobella* and intergrades with *juvenalis* along the lower slopes of the mountains of eastern Costa Rica and western Panama and with *pseudocobella* along the western and eastern slopes of the Andes of Colombia (Figs. 1, 2, 5d). The color pattern of individuals from the southern regions of Cartago and Punta Arenas Provinces of Costa Rica are distinctly banded but have less numerous black ventral marks and the subcaudal area is red, with only a few black flecks or spots laterally. A black lateral tail stripe is evident above the anus and continues to the tip of the tail. These individuals appear to have about 75% of the *juvenalis* features. Individuals from Boquete region of Chiriquí, Panama, have completely red or orange red subcaudals without black edging, the dorsal banding is less distinct posteriorly and the black ventral marks vary from almost absent to moderately dense. A black lateral caudal stripe is present in all specimens and occasionally begins in front of the vent. These individuals are almost perfect intermediates between *juvenalis* and *epinephelus*. Similar intermediates between *epinephelus* and *pseudocobella* occur along the upper slopes of the Andean foothills near Pueblo Rico and San Antonio, Pacific Colombia (also see Dunn, 1944).

TABLE 3. VARIATION OF *L. EPINEPHELUS* SUBSPECIES FOR VENTRALS, SUBCAUDALS, AND TAIL/TOTAL LENGTH RATIOS (LATTER CHARACTER BASED ON ADULTS ONLY).

	VENTRALS				SUBCAUDALS				TAIL/TOTAL LENGTH RATIOS (%)			
	N	Range	M	SE	N	Range	M	SE	N	Range	M	SE
<i>e. juvenalis</i>	51	139–152	145.0	0.4	45	51–62	56.0	0.4	25	18.9–25.2	21.5	0.2
<i>e. pseudocobella</i>	50	135–158	146.6	0.6	46	45–57	51.2	0.4	14	17.1–20.9	19.2	0.3
<i>e. epinephelus</i>	103	128–152	138.4	0.4	100	49–72	61.4	0.5	74	19.0–26.0	23.1	0.2
<i>e. albiventris</i>	93	141–165	150.7	0.5	83	52–70	60.3	0.5	63	19.1–24.2	22.0	0.2
<i>e. fraseri</i>	43	143–164	153.8	1.3	43	51–75	66.6	0.8	17	18.5–26.6	23.7	0.6
<i>e. lamonae</i>	48	141–157	149.7	0.7	41	51–67	59.2	0.7	22	20.2–24.1	22.2	0.2
<i>e. opisthotaenia</i>	32	142–152	147.0	0.5	29	53–78	64.7	0.9	17	20.2–25.6	23.6	0.3
<i>e. bimaculatus</i>	41	162–191	174.9	1.0	34	44–80	63.2	1.0	17	18.9–24.0	21.3	0.3

Liophis e. epinephelus from the lowlands of Panama and western Colombia have moderately defined black and red bands from the nape to midbody with the posterior part of the dorsum obscurely marked. The venter and subcaudals are orange or reddish. The black lateral stripe begins just anterior to the vent and continues to the tip of the tail. This pattern is typical of all samples of *e. epinephelus* in lowland Panama, Colombia, and extreme northwestern Ecuador. In some areas of its range (Cali, Pamplona, Colombia), *e. epinephelus* reaches an elevation of 1,800 m, where it does not contact another taxon of *epinephelus*.

Liophis e. epinephelus probably intergrades with *L. e. lamonae* along the upper slopes of the Río Cauca and Río Magdalena valleys. I have examined a single specimen from the state of Boyaca, Colombia, that shares characters of both taxa.

Color pattern morphs in the Andes of Colombia and Ecuador are complicated because there are three additional taxa that are parapatric with each other and infrequently intergrade with *e. epinephelus* and *e. pseudocobella*. These taxa, *e. lamonae*, *e. fraseri*, and *e. bimaculatus* (Figs. 5, 6), are similar in appearance. They tend to lose all indications of black dorsal banding in adults except for an occasional pair of black spots on the nape and have two to four

well-defined black stripes (two lateral and/or two dorsolateral). These stripes may extend from midbody or the posterior one-third of the body to the tip of the tail, with the two dorsolateral stripes merging as one on the median line of the tail.

Both the two and four-lined patterns may be found in the same population, but usually one has a higher frequency. The population of the Sabana region of Bogotá, Colombia and areas between 2,000–3,400 m are usually four-lined, whereas populations to the east, south, and west at elevations of 1,300–2,900 m are usually two-lined. The black lateral line is usually broad, occasionally occurring on part of the second, usually all or part of the third, and one-half to nearly all of the fourth scale rows. The lateral black line is usually bordered above by an ill-defined yellowish line about one to two scale rows wide, from midbody to the tip of the tail. The middorsal coloration is usually dark olive or dark brown, with or without a black line edging the union between the sixth and seventh (occasionally the fifth and sixth) scale rows. The ventral color is highly variable from almost immaculate yellow or salmon with a few scattered black spots to almost completely black with the lateral edges salmon or yellow. The usual condition is a yellow/black checkered pattern of about equal density. The checkered ven-

Figure 5. Subspecies of *Liophis epinephelus*. A. Cotypes of *L. bimaculatus* (AMNH 17532, 17604) "Colombia." B. *L. pseudocobella* (BMMH 1910.7.11.25) Pueblo Rico, Colombia. C. Syntype of *L. opisthotaenia* (BMNH 1946.1.4.55) Merida, Venezuela. D. Intergrade between *L. e. epinephelus* and *L. e. pseudocobella* (BMNH 1910.7.11.26) slopes above Rio San Juan, Colombia. E. Holotype, *L. taeniurus juvenalis* (ANSP 3687) San Jose, Costa Rica. F. Holotype, *L. epinephelus* (ANSP 3688) Truando, Colombia.

Figure 6. Subspecies of *L. epinephelus*. A and C represent *L. e. bimaculatus*; B and D, *L. e. albiventris*; E and F, *L. e. fraseri*. A. Holotype of *L. bipraeocularis* (BMNH 1946.1.4.93) Facatativa, Colombia. B. Syntype of *Zamensis ater* (BMNH 1946.1.21.52) Biscra, Algeria (in error). C. Holotype of *L. e. ecuadorensis* (MRHN 3267) "Ecuador." D. *Liophis e. albiventris* (KU 164215) 9.5 km NW Nono, Ecuador. E. Holotype of *Liophis bimaculatus lamonae* (MLS no number, photo courtesy of Charles Myers) Sonson, Colombia. F. Holotype of *Liophis fraseri* (BMNH 1946.1.6.63, photo courtesy of British Museum of Natural History) "West Ecuador."

ter is normal for *L. e. fraseri* and *L. e. lamonae*. The two-lined pattern is prevalent between elevations of 1,600–2,200 m in the eastern, central, and western Colombian Andes and the four-lined pattern between 2,200–3,400 m in the eastern Andes of Colombia and the central Andes of Ecuador and Peru.

Liophis e. bimaculatus is primarily a high altitude taxon (mean elevation of 2,600 m, Figs 2, 5a) with 162 to 191 ($\bar{x}$ 174.9) ventrals, whereas *L. e. fraseri* usually occurs at lower elevations (mean elevation of 1,900 m) with 141 to 164 ($\bar{x}$

152.8) ventrals. *Liophis e. bimaculatus*, *e. lamonae*, and *e. fraseri* tend to have 21 to 30% of their populations with seven supralabials and 76 to 80% with nine or less infralabials, respectively. In addition, approximately 20% of *L. e. bimaculatus* populations have two preoculars on each side. The latter taxon appears to have the highest number of ventrals in Ecuador with a slight decrease both north and south of the Ibarra region. On the other hand, the number of ventrals of *L. e. lamonae* averages about 150 in Colombia and 149 in northern Ecuador, while *e. fraseri* averages 155 ventrals in southern Ecuador, and decreases to a mean of 151 in central Peru (Fig. 8).

The nominate taxon intergrades with *L. e. albiventris* in the lowlands of southwestern Colombia and northwestern Ecuador. Two specimens from the lowlands of northwestern Ecuador have the dorsal and ventral pattern of *L. e. epinephelus* but with the number of ventrals normally found in *L. e. albiventris*. Typically patterned individuals of *L. e. albiventris* (Fig. 6d) occur within 70 km of these two specimens, suggesting that the zone of intergradation may be narrow. The majority of the specimens of *L. e. albiventris* (91%) occur above 500 m elevation, and most are found between 1,000 and 2,000 m along the western face of the Andes in Ecuador (Fig. 3). *Liophis e. albiventris* has an immaculate cream to yellow venter and subcaudal area. Only six of 93 specimens had any indication of black marks on the ventrals.

The dorsum of adult *L. e. albiventris* is leaf green with none to few obscure black marks on the nape and/or anterolaterally on the trunk. Juveniles (150 to 190 mm total length) are similar but have a black nape band or two black nape spots followed by a narrow white or cream collar. The light collar and dark nape spots disappear as the individual reaches 230 to 260 mm in total length. Young *L. e. fraseri* and *e. lamonae* have similar nape markings that fade almost as rapidly, but

Figure 7. Variation in the numbers of maxillary teeth in *L. epinephelus*. The irregular black line represents the number of individuals in a particular subspecies with a particular number of maxillary teeth. The horizontal line represents the total range of variation, vertical line, the mean; open rectangle one standard deviation on either side of the mean; black rectangle two standard errors on either side of the mean. Sample sizes are given in Table 2 and Figure 5.

Figure 8. Variation in the numbers of ventrals and subcaudals in *L. epinephelus*. Numbers and letters in left hand column refer to subspecies, latitude, and geographic location (see Table 2, Fig. 3). Horizontal line = range; vertical bar = mean; black rectangle = two SE on either side of mean; open rectangle = one SD on either side of mean.

occasional adults retain some indication of the juvenile pattern. All *L. e. albiventris* have a black lateral posterior stripe that continues to the tip of the tail. Infrequently, young and adults have black paravertebral posterior stripes that converge into one dorsomedian tail stripe. Juveniles and hatchlings occasionally have a diffuse light stripe immediately above the black lateral posterior stripe, resembling the typical pattern seen in *L. e. fraseri* and *e. lamonae* except for the immaculate venter.

Liophis e. albiventris and possibly *L. e. lamonae* seem to merge with populations of *L. e. bimaculatus* at elevations between 2,600–2,800 m in northern Ecuador. The number of ventrals increase from an average of 150 between 1,000–2,000 m to 156 between 2,000–2,600 m and 174 ventrals above 2,800 m. *Liophis*

e. albiventris also intergrades with *L. e. fraseri* in the vicinity of Celica, southern Ecuador, where the latter taxon switches from having an eastern slope distribution to a western one (Fig. 3). An individual from near Celica, 1,900 m, has a color pattern similar to most *L. e. fraseri* and the squamation of *L. e. albiventris*. Specimens from Alamor, 1,100 m, some 10 km northwest of Celica, are typical *L. e. albiventris*. The Alamor locality seems to be the southern limit of the distribution of *L. e. albiventris*, whereas *L. e. fraseri* continues southward along the middle slopes of the western face of the Andes to above Trujillo, Peru, switching again to the eastern slopes and continuing southward to Ayacucho, Peru. The two specimens of *L. e. fraseri* from central Peru are allopatric from other samples by 325 km. These two differ in

having almost black venters; white subcaudals, few ventrals and subcaudals (Sample 16 of Fig. 8). The dorsal body pattern resembles *L. williamsi* from Venezuela, and one of the two specimens has a similar head pattern. However, *L. williamsi* has no scale row reductions while *L. fraseri* has a 17-17-15 pattern. With additional material, the central Peru population may prove to represent a new taxon.

The only subspecies of *L. epinephelus* that appears to be allopatric is the Venezuelan *L. e. opisthotaenia*. This population occurs from the upper drainage of the Río Chama, 2,000 m, just east of Mérida, west-southwest to the vicinity of Las Delicias, 2,000 m, Páramo de Tama, on the Colombian border. Although there is no evidence of intergradation of this race with adjacent subspecies, it may occur because the subspecies crosses the Tachira Depression that separates two major Andean formations.

Most *Liophis e. opisthotaenia* (Fig. 5) have immaculate venters (26 of 32). Six show scattered to many black marks on the lateral edges of the ventrals. The dorsum is olive green, usually with two obscure black nape spots and occasionally dorsolateral dark streaking anteriorly. The juvenile pattern is similar to that of other *L. epinephelus*. The black postocular bar is usually present, similar to most individuals of *L. e. bimaculatus*. The basic body pattern is very similar to that of *L. e. bimaculatus*, *L. e. fraseri*, and *L. e. lamonae*. However, occasional individuals resemble *L. e. albiventris*, except for the slightly longer tail and usually 21 or fewer maxillary teeth (Fig. 7).

KEY TO THE SUBSPECIES OF *LIOPHIS EPINEPHELUS*

The following key is provided to aid in the identification of the taxa of this difficult species. The characters used are those that best exemplify the differences between the subspecies, but the geo-

graphic area of the samples is also important.

1. Lateral black tail stripe absent, dorsal and ventral surfaces of body and tail red and black banded 2
- Lateral black tail stripe present, dorsum may be banded, spotted, flecked, or almost unicolor, belly checkered with black or not 3
2. Caudals usually 54 or less (Colombia) *e. pseudocobella*
- Caudals usually 54 or more (Costa Rica) *e. juvenalis*
3. Ventrals usually 165 or less 4
- Ventrals usually 165 or more (high Andes, Venezuela to Peru) *e. bimaculatus*
4. Anterior half of dorsum with or without black flecks, streaks, or dark spots, never banded in adults, ventrals 143 or more 5
- Anterior fourth of dorsum banded with black or brownish-black bands, venter immaculate white, yellow or reddish, ventrals usually 143 or less (Panama to Ecuador) ... *e. epinephelus*
5. Venter usually immaculate white, yellow, or pinkish 6
- Venter yellow and black checkered 7
6. Dorsum leaf green, posterior dorsolateral black stripe usually absent (coastal and foothill zones of western Ecuador) *e. albiventris*
- Dorsum olive, olive brown, or grayish brown, posterior dorsolateral black stripe usually present (Páramo de Tama, east to Trujillo, Venezuela) *e. opisthotaenia*
7. Ventrals range from 141 to 156 ($\bar{x}$ 150), subcaudals 51 to 67 ($\bar{x}$ 59) (Andean slopes of central Colombia to central Ecuador) *e. lamonae*
- Ventrals range from 151 to 164 ($\bar{x}$ 157), subcaudals 61 to 76 ($\bar{x}$ 67) (Andean slopes of southern Ecuador to central Peru) *e. fraseri*

ACKNOWLEDGMENTS

My interest in South American herpetology was stimulated by the astute Ernest E. Williams, who on several occasions presented a view different from my own, and to my dismay, more frequently correct. His expertise, thoughtfulness, and willingness to help has been a rock upon which herpetologists have obtained a better understanding of evolutionary principles. Thanks, Ernest, for the more than 25 years of correspondence and friendship.

I heartedly thank those curators who willfully, but sometimes grudgingly, allowed me access to their specimens. I

offer my condolences to those curators who did not, for their *Liophis* specimens may forever remain misidentified. The collections examined, their acronyms and respective curators are: American Museum of Natural History (AMNH), R. G. Zweifel, C. M. Myers; Academy of Natural Sciences, Philadelphia (ANSP), E. Malnate; British Museum (Natural History) (BMNH), A. G. C. Grandison, A. F. Stimson; California Academy of Sciences (CAS), A. E. Leviton, R. C. Drewes; University of Southern California Costa Rica Series (CRE), J. M. Savage; Collection of Vertebrates, Universidad Los Andes, Mérida (UAM), J. Pefaur; Field Museum of Natural History (FMNH), R. Inger, H. Marx; private collection of Jay M. Savage (JMS); University of Kansas Museum of Natural History (KU), W. E. Duellman; Los Angeles County Natural History Museum (LACM), J. W. Wright; Louisiana State University Museum of Zoology (LSUMZ), D. A. Rossman; Museo de Historia Natural La Salle, Caracas (MHNLS), A. Paolillo; Harvard University Museum of Comparative Zoology (MCZ), E. E. Williams; Museum d'Histoire Naturelle, Geneva (MHNG), V. Mahnert; Museum d'Histoire Naturelle, Paris (MNHP), R. Roux-Esteve; Institut Royal des Sciences Naturelles de Belgique (MRHN), G. F. de Witte, J. P. Gosse; University of California Museum of Vertebrate Zoology (MVZ), D. B. Wake; Universidade de São Paulo Museum de Zoologia (MZUSP), P. E. Vanzolini; University of Texas Natural History Museum (TNHC), R. F. Martin; University of Illinois Museum of Natural History (UIMNH), D. F. Hoffmeister; University of Michigan Museum of Zoology (UMMZ), A. G. Kluge, R. A. Nussbaum; National Museum of Natural History (USNM), W. R. Heyer, G. Zug; and Federico Medem Collection (WWL), W. W. Lamar.

I thank Michael McCoid, Jack Sites, Jr., and Robert Dean for reading all or parts of the manuscript. I especially thank the administration of Texas A&M University for allowing me the time and

funds to examine some of the type material in European Museums. I especially acknowledge the perseverance of my wife, Mary, who, for the first time, found out what it was like to take squamative data from snakes.

APPENDIX: SPECIMENS EXAMINED

Liophis epinephelus albiventris. ECUADOR: doubtful or no specific localities, AMNH 17498, 17634, 49104, 35807, UIMNH 63524, UMMZ 88970, USNM (GOV) 7316-17, 7291, BMNH 1946.1.21.51-53, MHNG 1367.47; Alamor, AMNH 18320, 22230, 22095, 22231, 22235, 56168; Aluriguín, USNM (GOV) 7429-30; Balzapamba, UMMZ 88959; Balzar, USNM (GOV) 7330, 7423; Baños, UMMZ 88962-63; Camino de Guala, USNM un-numbered; Chaucha Valley, UMMZ 63121; 8 km W Chiriboga, USNM (JAP) 4109-10; 4 km W Chiriboga, KU 142807; 7.7 km E Chiriboga, KU 142805; 3.7 km E. Dos Rios, KU 142807; 14 km NE Dos Rios, KU 164214; Guayaquil, USNM 62792; Hacienda Carcelén, 15 km NE Quito, USNM (GOV) 7341-42; Intac, BMNH 78.1.25.52; Lita, USNM (GOV) 7425; Llambo, USNM (no number); Los Rios, Hda. Clementina, MHNG 1337.36; 10 km SE Machala, AMNH 113021; Mapoto, UMMZ 88960-61; Milligalli, MCZ 8394-95, USNM (GOV) 7424; Milpe, USNM (JAP) 9024; Mindo, BMNH 1916.5.23.3-4, USNM (GOV) 7426-27; 4 km N Mindo, USNM (JAP) 8278; 1 km E Mindo, USNM (JAP) 8273; 1.5 km E Nanegal Chico, USNM (JAP) 6285; 9.5 km NE Nono, KU 164215; 13 km NW Nono, KU 158534; Olalla, BMNH 82.7.24.2; near Oyacachi, USNM (GOV) 7396; below Pacto USNM (GOV) 7422; Paramba, BMNH 98.4.28.68; 56 km N Quevedo, KU 152605; Quito, BMNH 72.2.26.7; near mouth of Río Aguas Claras, USNM (GOV) 7333; 24 km S Río Baba, UIMNH 92256; Río Blanco, 20 km from Mindo, USNM (JAP) 7331 Río Mindo, USNM (JAP) 7428; Río Mulaule, USNM (GOV) 7299-300; Río Saboya, BMNH 1940.2.20.30; Río Silanchi, USNM (GOV) 7819; San José de Chimbo, AMNH 17498, 17634; Santo Domingo AMNH 27144, BMNH 1918.1.15.3; 5-10 km SSW Santo Domingo de los Colorados, AMNH 110588, UIMNH 92256, USNM (GOV) 6006; between km markers 38/29, NNW Santo Domingo de los Colorados, USNM (JAP) 1927, Tandapi, KU 112272-73, 121319-21; 9 km E Tandapi, AMNH 113020; Tandayapa, USNM (GOV) 7329; Tanti, MCZ 8400; Tumbaco, USNM (GOV) 7332.

Liophis epinephelus bimaculatus. COLOMBIA: no specific locality, AMNH 27615, 17532, 17604; Bogotá, AMNH 7648, 14029-31, 14035, 24227, 24232, 24251, 67151, 67157-58, 91813, CAS-SU 8274, BMNH 1919.3.6.21, MCZ 17133-34, MVZ 71534, USNM 95180; Bolívar, Vereda de San Juan

MZUSP 6131; Facatativá, BMNH 1946.1.4.93; La Calera, MVZ 71535-36; La Uvita, MCZ 66382; Moscopán, UMMZ 121039; 23 km E Pacho, AMNH 104679; Pueblo Viejo, MCZ 6551; Usaquén, LACM 28856; Villavicencio, (in error?) BMNH 1915.3.11.3; Vitelma, AMNH 67152-56. ECUADOR: no specific locality, MRHN 3267; Ibarra, BMNH 96.4.28.74-75; Otavalo, KU 135182. PERU: Cajamarca, UMMZ 59846; Huagal, UMMZ 59147. VENEZUELA: 17 km N. Guaraque, UAM 1160.

Liophis epinephelus epinephelus. COLOMBIA: no specific locality, AMNH 177556-58; Andagoya, BMNH 1915.10.21.13-14, 1916.4.25.7, MCZ 32733-35, UMMZ 48409, 121050, 121057; 100 km N Bogotá, MHNG 1367.55; Buenaventura, USNM 151724, 154037; 13 km from Buenaventura, FMNH 165441, 165451, 165453, 165564; Cali, USNM 151749; Caño Docordo, CAS 119599; Chigorodó, AMNH 97276; Cisneros, FMNH 43742-44; Condoto, BMNH 1913.11.12.48, 1914.5.21.43-45; Imbibi, FMNH 166024; Pueblo Rico, FMNH 54940, 54950-56; Río San Juan, USNM 72364, 72368, 159487-88; Río Sinu, UMMZ 126722; San Sebastián de Rábago, CAS 113880; Santa Rosa de Osa, AMNH 35784; Sierra Negra, USNM 117505; Truandó, ANSP 3688; near Turbo, AMNH 94343-44, AMNH 158946; Vista Nieve, UMMZ 63776 (3). ECUADOR: Borbón, BMNH 1902.5.27.19; near Esmeraldas, USNM (OVS) 981. PANAMA: no specific locality, FMNH 6113, 31211-13, 31721, 31725, 31728, 83546, 154513-14, 154527, USNM 13565; UMMZ 90338; Darien, AMNH 36205; Agua Clara, MCZ 45372-73; Albbrook Air Force Base, TNNH 23713-15; Barro Colorado Island, AMNH 80037, 77575, FMNH 178599, KU 75707-08, 80595, LACM 9278-79, MCZ 20593, UMMZ 63745, 76022, 124160-61, 124186, USNM 203834; near Camarones, KU 110712; Camp Chagres, KU 75709; Caña, USNM 50118-20; Caña Mines, UMMZ 83157 (6), 117703; Cerro la Campana, KU 75710; Chagres River, FMNH 154489, MCZ 34378, 37107-11, UMMZ 76721 (3), 83145; El Valle Río Anton, AMNH 76018, 108706, 108968, FMNH 68074, KU 110720-21; Empire, KU 110717, USNM 54228, 60028; Esperanza Ridge, MCZ 42721; Fort Clayton, KU 110714-16, UIMNH 41737; Fort Frank S Todd, USNM 140690-91; Fort Gulick, KU 110719; Fort Kobbe, CAS 85222; Fort Randolph, MCZ 20592; Fort Sherman, MCZ 22239, 22242, USNM 192283; Gamboa, AMNH 71672, 73323, KU 110713, MCZ 23972; Gatuncillo and Chagres fork, MCZ 37872; Las Cumbres, KU 110710-11, 110724; near Limón, MCZ 23984; Madden Dam, MCZ 38231, UMMZ 76022, 76724 (2); Minas de Caña, MCZ 42752; Pihulla-lucuti fork of Río Tuira, MCZ 37119-20; upper Río Chiriquí, AMNH 114320-22; Río Grande-Río Cocle, MCZ 37879; Río Sardinillo, KU 110718; Río Tuira at Río Mono, KU 110725; San Pablo, MCZ 3930-31, USNM 120826-27; Three

Falls Creek, AMNH 65300; Venado Beach, USNM 193374, MCZ 38215.

Liophis epinephelus fraseri. ECUADOR: "West Ecuador," BMNH 1946.1.6.63; 10 km N Celica, MCZ 93590; Loja, BMNH 1935.11.3.67-68, KU 125415; 2 km E Loja, KU 121322-23; 12.2 km S Loja, KU 141269-87, 141289; 6.5 km N Loja, KU 142809-11 7.3 km N Loja, KU 142808; Río Piuntza, KU 147199; San Ramon, USNM (GOV) 7328. PERU: 15 km E Canchaque, LSUMZ 32554-59; Chanchamayo, FMNH 40639; Huanacabaniha, MCZ 17398; 45 km NE Tarma, MCZ 45902; Trujillo, USNM 192303 (in error?); just below Yuracyacu, on Tambo-Valle del Apurimac path, LSUMZ 27138.

Liophis epinephelus juvenalis.—COSTA RICA: no specific locality, CRE 2996; vic. Alajuelita, CRE 8483-85; Barba, AMNH 17281; Capellades, CRE 7057; Cartago, CRE 6401, UMMZ 131415; Cervantes de Cartago, CRE 6393; Cinchona de Sarapiquí, CRE 6427, KU 31899, 31901, 35519-21, 63826; Concordia, KU 63827; between Curridabat and San Antonio de Desamparados, CRE 2725; El Angel, UMMZ 136999; El Silencio, CRE 3054-55; El Tablazo de la Carpintera, CRE 3031; Finca El Helechales, CRE 8278; Guadalupe, KU 31900; Guayapiles, UMMZ 83181; Isla Bonita, FMNH 101243-44, 188808, KU 25750-52; La Estrella, MCZ 28072; La Gloria, CRE 3012; La Palma, CRE 243, Limón, AMNH 5304; Monteverde, CRE 3178; Navarro, MCZ 15297-99, UMMZ 74302; 1.5 km NW Pastora, CRE 34; Puerto Veijo, CRE 3030 (in error?); Puriscal Centro, Santiago, CRE 6335; Rama sur Río Las Vueltas, KU 103885; Río Claro, JMS 1763; Río Coton, KU 100637; Río Torres in Guadalupe, CRE 908, 7170, KU 31901; Sabanilla, JMS 1160; San Antonio de Belem, MCZ 32041; San Ignacio de Acosta, CRE 3015 (2); San Isidro de Coronado, MCZ 55116, FMNH 37231; San Isidro del General, UMMZ 117729; San Jose, AMNH 17298, 17377, ANSP 3687, CRE 6336, 7003; San Pedro, in San Jose, CRE 2956, 6421, 6428-29, UMMZ 125508, 131414, 131416-20; San Pedro de Montes de Oca, CRE 6331; San Ramón de Tres Ríos, LSUMZ 15655; 4.5 km S San Vito de Java, LACM 114075; 6 km S San Vito de Java, LACM 114074; Tapanti, CRE 6314; vic. Tapanti, 0.8 km from Puente de Río Grande, CRE 3246; Tres Ríos, CRE 2946; Turrialba, MCZ 55025; Varablanca, KU 125479; near Villa Quesada, CRE 707; Volcán Poas, east slope, UMMZ 117828 (3). PANAMA: Santa Clara, AMNH 80085.

Liophis epinephelus lamonaie. COLOMBIA: no specific locality, MHNG 840.59; Aguadita, AMNH 35241; Bogotá, AMNH 24236 (in error?); Chusacá, KU 124939; Las Palmas, FMNH 63759; near Manzanares, WWL 747-48, 780-82, 802-06, 989-90, 1001, 1006; Medellín, AMNH 35492, 35721; Pensilvania, AMNH 35250; Río San Joaquín, FMNH

54957; San Agustín, FMNH 69666; San Francisco, AMNH 106660; San Pedro, AMNH 35745; Sinaí, MCZ 150201, MZUSP 6128; Villavicencio, BMNH 1915.3.11.2. ECUADOR: no specific locality, AMNH 2793, 35980, 35897–99, BMNH 1946.1.6.63; Baeza, USNM (JAP) 5758; Cayambe, UIMNH 82483 (in error ?); La Alegria, USNM (JAP) 4501; La Bonita, USNM (JAP) 4502–05, 4917; Mt. Tungurahua, FMNH 36623, KU 121329; ca. Sevilla de Oro, USNM (JAP) 6691.

Liophis epinephelus opisthotaenia. VENEZUELA: Betania, Páramo de Tama, MCZ 112415; Chama, BMNH 1905.6.30.59–60; Mérida, BMNH 1905.5.31.56–58, 1908.5.29.146–147, 1912.11.1.73–76, 1946.1.4.54–55, UAM 602, 771, 1046–48, 1050–51, 1162, 1697, 1710, 2318; Misisi, 14 km E Trujillo, USNM 162827; Río Albarregas, BMNH 1903.6.30.37–38, USNM 162822; Río Albarregas y Milla, AMNH 13418–19; San Cristóbal, MHNLS 1096.

Liophis epinephelus pseudocobella. COLOMBIA: no specific locality, AMNH 35263–65, 35267–76; 10 km NW Andes, LACM 72760; Barbacoas, FMNH 54938; Bogotá, MCZ 19210–11 (3) (in error); Jericó, AMNH 35246, FMNH 27015–16, UMMZ 84088, 78280 (2), USNM 86820; Medellín, AMNH 35539–44, 35553–56, 35560–65, 35623, 35627–28, 35639, 35672, 4492; Pasto, AMNH 91814; Popayán, FMNH 54931–35, 54985–95, MHNG 1532.13; 15 km N Popayán, MVZ 68705; Pueblo Rico, BMNH 1910.7.1.25; Río Cofre, KU 140411; 4 km NW San Antonio, MVZ 68697–703; ca. 5 km NW San José, AMNH 106655; Sonsón, FMNH 63780, MCZ 21991.

Liophis epinephelus epinephelus × *e. juvenalis*. PANAMA: no specific locality, AMNH 75637; Boquete, CAS 71340; north of Boquete, CAS 78922–36; Chiriquí, BMNH 93.11.22.5–10; El Hato, FMNH 68072–73, 154479–80; El Volcán, AMNH 63596, 75633, KU 75711, 110722–23.

Liophis epinephelus epinephelus × *e. lamonaе*. COLOMBIA: Junín, LACM 36780.

Liophis epinephelus bimaculatus × *e. pseudocobella*. COLOMBIA: 13 km SE Pedrancha, AMNH 106659.

Liophis epinephelus epinephelus × *e. pseudocobella*. COLOMBIA: slopes above Río San Juan, BMNH 1910.7.11.26.

LITERATURE CITED

- AMARAL, A. DO. 1931. Studies of neotropical ophidia XXIII. Additional notes on Colombian Snakes. *Bul. Antivenin Inst. Am.*, 4: 85–89.
- BOULENGER, G. A. 1894. Catalogue of the snakes in the British Museum (Natural History). Vol. 2. London, Trustees of Museum, 382 pp.
- . 1903. Descriptions of new snakes in the collection of the British Museum. *Ann. Mag. Nat. Hist.*, 12(7): 350–354.
- . 1908. Descriptions of new South American reptiles. *Ann. Mag. Nat. Hist.*, 1(8): 111–115.
- COPE, E. D. 1862. Synopsis of the species of *Holcosus* and *Ameiva*, with diagnosis of new West Indian and South American Colubridae. *Proc. Acad. Nat. Sci. Phila.*, 13: 60–82.
- . 1868. An examination of the Reptilia and Batrachia obtained by the Orton Expedition to Ecuador and the upper Amazon, with notes on other species. *Proc. Acad. Nat. Sci.*, 20: 96–140.
- . 1899. Contributions to the herpetology of New Granada and Argentina. *Philadelphia Mus. Sci. Bull.*, (1): 1–22.
- DUNN, E. R. 1937. New or unnamed snakes from Costa Rica. *Copeia*, 1937: 213–215.
- . 1944. A revision of the Colombian snakes of the genera *Leimadophis*, *Liophis*, *Lygophis*, *Rhadinaea*, and *Pliocercus*, with a note on Colombian *Coniophanes*. *Caldasia*, 11(10): 479–495.
- GÜNTHER, A. 1872. Seventh account of new species of snakes in the collection of the British Museum. *Ann. Mag. Nat. Hist.*, 9(4): 13–37.
- JAN, G. 1863. Enumerazione sistematica degli ofidi appartenenti al gruppo Coronellidae. *Arch. Zool. Anat. Fisiol.*, 2(2): 213–330.
- LAURENT, R. 1949. Notes sur quelques reptile appartenant a la collection de l'Institut Royal des Sciences Naturelles de Belgique. *Bull. Inst. Roy. Sci. Nat. Belgium*, 25(9): 1–20.
- PERACCA, M. G. 1914. Reptiles et batrachiens de Colombia. In O. Fuhrmann and E. Mayor 1914, *Voyage d'Exploration Scientifique en Colombie*. *Mem. Soc. Neuchatel*, 5: 1–1090.

Partial Cyclopia and Monorhinia in Turtles

A. D'A. BELLAIRS¹

ABSTRACT. Three albinoid embryos of the loggerhead turtle (*Caretta caretta*) with partial cyclopia and one with a single nostril are described. The two specimens sectioned show many other abnormalities of the head including virtual absence of the midline components of the anterior chondrocranium.

INTRODUCTION

The congenital malformation of cyclopia is well known in man and other mammals (Schwalbe and Josephy, ca. 1913; Willis, 1958) and has also been reported in birds (Romanoff, 1972), in reptiles (Ewert, 1979), and in amphibians and fish (Adelmann, 1936). It is characterized by the fusion or partial fusion of the eyes in the midline and by abnormalities of the brain and craniofacial structures. In amniotes the nose is often represented by a proboscis-like structure above the single or partially fused eyes. The malformation has been produced experimentally by various chemical and physical procedures (Johnston *et al.*, 1977), but may also occur spontaneously in the absence of any obvious environmental cause.

Among reptiles, cyclopia appears only to have been reported in chelonians (Ewert, 1979), though Bellairs (1965) described a Siamese twin embryo of *Lacerta agilis* showing a kind of "pseudocyclopia" caused by fusion of the two heads creating a third, single eye in the midline. Although Ewert figured

cyclopic near-hatching embryos of *Chrysemys picta* and *Pseudemys scripta elegans*, the anatomy of the condition remains undescribed. The present contribution seems appropriate in this collection of essays in honor of Dr. Ernest E. Williams, who has made important studies on the normal anatomy of chelonians.

MATERIALS AND METHODS

The four specimens described are very late embryos of the Atlantic loggerhead turtle, *Caretta caretta caretta*, with yolk sacs attached. Three specimens show partial cyclopia, while the fourth shows a malformation which I have termed monorhinia (single nose), though I know of no precedent for this term. These specimens, kindly sent me by Dr. P. C. H. Pritchard of the Florida Audubon Society, were obtained in the course of a study of the effects of environmental conditions on the hatchability of turtle eggs (McGehee, 1979). Both unsuitable temperatures and mechanical disturbance during incubation are known to have deleterious effects upon development (Ewert, 1979), and were among the variables investigated. It is by no means certain, however, that the malformations were caused by the experiments, since the majority of eggs were unaffected by them, and abnormal individuals also occur in natural nests. Specimens B and C were preserved intact while the heads of A and D were serially sectioned and stained with Masson and Casson's triple

¹St. Mary's Hospital Medical School, London, W.2., U.K.

stains. Some normal material was also available for comparison.

EXTERNAL APPEARANCE

Figures 1, 2

All four specimens were albinoid. None of them possessed an egg-caruncle, a defect which in itself might have prevented hatching. Apart from showing duplication of some of the epidermal scutes (notably the vertebrae in specimens B and C) and the reduction of pigmentation, these turtles show no obvious abnormalities outside the head region.

Specimen A. Carapace-length 36 mm. Egg turned through 360° four hours after laying in McGehee's experiments.

There is a single slitlike orbital opening within which the eyes are barely visible externally. Above this opening is a well-developed proboscis 2.5 mm long, bearing a single narial opening at its tip. The upper jaw is truncated, and the apparently normal lower jaw projects forward well beyond it. A small portion of the hindbrain is herniated from the back of the head.

Specimen B. Carapace-length 36 mm. Egg incubated at 32°C. instead of the optimum 30°. The eyes are not markedly reduced in size but are fused or contiguous medially. The proboscis is turned back over the head. The upper jaw is truncated.

Specimen C. Carapace-length 40 mm. Egg turned through 360° 48 hours after laying. As in B, the eyes are fused or contiguous medially. The lower jaw projects anteriorly and dorsally, covering the truncated upper jaw. The proboscis is directed forward, as in A.

Specimen D. Carapace-length 40 mm. Egg turned through 360°/40 hours after laying. The eyes are not markedly reduced and hardly closer together than normal. The upper jaw is shortened, though not to the same extent as in the previous specimens. There is no probos-

cis, but there is a single median nostril between the eyes.

MICROSCOPIC FINDINGS

Specimen A. Study of sections shows that the proboscis is traversed by a single nasal tube which widens out posteriorly, forming a single blind sac which has no connection with the mouth or pharynx. A similar absence of the choanal opening is mentioned by Schwalbe and Josephy (ca. 1913) in cases of mammalian cyclopia. The nasal tube is lined anteriorly by pseudostratified squamous epithelium. Posteriorly the nasal sac contains areas of olfactory epithelium with associated Bowman's glands. The nasal tube and sac are partly invested by a tube of cartilage which represents the nasal capsule (Fig. 3A-C). A small process projects dorsally from the floor of the capsule, as if it were an attempt to form a nasal septum (Fig. 3C); however, it shows little resemblance to the normal structure.

The bones are difficult to interpret. The bones within the base of the proboscis and around the nasal tube and sac might be the prefrontals as labelled here (Fig. 3); these are normally the most anterior bones of the cranial roof since there are no nasals in Recent cryptodiran Testudines. The identity of the more lateral bones which extend ventral to the partly fused eyes (Fig. 3C) is problematical; their relations are very different from those of normal frontals. An alternative possibility is that the bones around the nasal tube and sac are abnormally large frontals which have grown anteriorly and ventrally, while the problematical bones are prefrontals which have been displaced ventrolaterally, still retaining their normal contact with the maxillae. The maxillae themselves are partly fused in the midline and the premaxillae appear to be absent, as in most cases of mammalian cyclopia (Schwalbe and Josephy, ca. 1913).

Olfactory nerves arise from the pos-

Figure 1. Abnormal late embryos of *Caretta*. A, B, and C (partially cyclopic) and D (with monorhinia) from left to right.

terior region of the nasal sac (Fig. 4A), pass back as a large bundle beneath the forebrain and appear to end blindly among the eye muscles. These nerves arise from neuroblasts in the olfactory epithelium, and it would seem that in this specimen they have grown back but have failed to enter the olfactory bulb in the normal fashion. The brain itself is grossly malformed and difficult to interpret. There appears, however, to be only one olfactory bulb (possibly fused bulbs), and the cerebral hemispheres are more or less fused. The forebrain is supported by wings of cartilage which probably represent the planum supraseptale. There is no indication of the interorbital septum, and no parts of the trabeculae can be identified.

The eyes are considerably smaller than

normal and are fused medially (Figs. 3C 4A). Each eye consists of a much folded layer of retinal tissue surrounded by a layer of pigment, probably of retinal origin. It is interesting that no lenses are present. The lens is normally induced by the optic vesicle or cup, and it would seem that in this specimen there was a failure of induction, either because of the small size of the vesicles, or their failure to invaginate. This last possibility is suggested by the folded appearance of the retinae.

The two eyes are surrounded by a single capsule of rather thick scleral cartilage. Between the eyes and the proboscis is a flattened gland which discharges anteriorly through a long duct beneath a "palpebral" fold of tissue dorsal to the eyes. Probably this represents either an

Figure 2. Abnormal late embryos of *Caretta*. A (left) and B are at top, C (left) and D at bottom. A, B, and C are partially cyclopic, proboscis is turned back in B. D has monorhinia.

Harderian or a lacrimal gland. Massive but disorganized eye muscles are attached to the scleral cartilage.

A remarkable feature is the presence of a third eyelike structure which is connected with the two fused eyes by a narrow stalk which issues from a cleft in the median ventral aspect of the scleral cartilage (Figs. 3C, 4A). Anteriorly the stalk is pigmented; posteriorly it expands into

a dilated vesicle composed of apparently retinal tissue. The vesicle extends caudally for some distance and ends blindly without reaching the brain. It seems possible that this vesicle was derived from a single optic stalk which has become dilated and differentiated into retinal tissue, but which has lost its original connection with the brain.

The specimen shows many other cra-

Figure 3. Transverse sections through front of head of partially cyclopic specimen A of *Caretta*. A. Through proboscis. B. Through base of proboscis. C. Through nasal sac and fused eyes. A strand of retinal pigment associated with the "third eye" is issuing through the cleft in the scleral cartilage.

Abbreviations: b, bone of problematic identity; e, eye; g, eye gland; lj, lower jaw; mx, maxilla; nc, nasal capsule (cartilaginous); ns, nasal sac; nt, nasal tube; p, pigment; pf, ? prefrontal; re, retina; sc, scleral cartilage.

nial malformations. These are more extensive than the external appearance of the head would suggest. They include extreme narrowing of the pharynx, absence of inner ear structures (presupposing absence of otocysts), and absence of obvious pituitary elements. This last abnormality has been reported in other embryos with cranial malformations (Bellairs, 1965; Johnston *et al.*, 1977).

Specimen D. This specimen has a single tubular nasal vestibule (Fig. 4B) which leads back into the main part of the more or less single nasal sac. The most anterior part of this sac is paired. Throughout most of its extent, however, it is only partly divided by a broad but incomplete septum which lacks a cartilaginous component (Fig. 4C). As in specimen A, the sac ends blindly posteriorly and there is no internal choana. Both vestibule and main nasal sac are more or less surrounded by a single cartilaginous nasal capsule.

The front of the upper jaw has a sharp, beaklike appearance in transverse section (Fig. 4C.) There are no premaxillae, but maxillae, prefrontals and frontals are recognizable. In the absence of choanae, the bones of the palate are abnormal and the vomers cannot be identified. The palatal shelves of the maxillae form a median symphysis throughout much of their length (Fig. 4C,D) and articulate posteriorly with the palatines (Fig. 4E). A median diverticulum from the mouth, anterior to and quite separate from the supposed pituitary (see below), extends upwards between the palatines (Fig. 4E). It is possible that this diverticulum is a vestige of the choana.

The olfactory nerves pass caudally to enter the olfactory bulbs in more or less normal fashion. The nerves are, however, displaced to one side by a large cyst, presumably pathological, which lies above the posterior part of the nasal sac (Fig. 4C). The cerebral hemispheres are partly fused with each other and obviously abnormal.

The eyes are quite separate and rela-

tively normal in general appearance (though not perhaps in detailed structure), and possess lenses. The two optic nerves, however, appear to be directly continuous with each other across the midline (Fig. 4E), and do not form a chiasma. They have no connection with the brain. This would suggest that the proximal part of the optic stalk was originally single, and that this portion disappeared at some later stage of development.

No neural components of the pituitary can be identified. There is, however, a median diverticulum from the pharynx which extends dorsally to end within the substance of the ossifying basisphenoid. It seems possible that this represents the pharyngeal component of the pituitary, or Rathke's pouch.

A remarkable feature of this specimen is the deficiency of the midline structures of the anterior part of the chondrocranium, i.e., the structures which are partly or completely of trabecular origin. As previously mentioned, there is no cartilaginous median nasal septum of normal type. The lateral walls of the posterior part of the nasal capsule, however, become approximated almost in the midline for a short distance behind the blind nasal sac, and give the appearance of a paired septum. More posteriorly, there is no interorbital septum, although the eyes are not especially close together. The two Harderian glands, which normally lie one on either side of the interorbital septum (Fig. 4F), are contiguous with each other (Fig. 4D).

The anterior orbital cartilages are represented by a well-developed planum suprasedale which lies in the normal position beneath the forebrain. The paired parts of the trabeculae, which normally flank the pituitary fenestra, are not apparent, and there is no fenestra. Instead there is a pointed median rod of cartilage which projects anteriorly from the front of the parachordal or basal plate. It is partly ossified and merges posteriorly with the basisphenoid ossification

Figure 4. A. Transverse section through head of partially cyclopic specimen A of *Caretta* showing fused eyes and "third eye." Level is posterior to Fig. 3C. B–E. Transverse sections through head of specimen D of *Caretta* with monorhinia. B. Through front of snout. C. Through nasal sac showing incomplete septum. D. Through middle of eyes showing absence of interorbital septum. E. Through posterior parts of eyes showing optic nerve. F. Transverse section through interorbital septum of normal *Caretta* embryo.

Abbreviations as in Fig. 3, plus new terms: br, brain; cs, conjunctival space; cy, cyst; d, diverticulum from mouth (not pituitary); e 3, "third eye"; fr, frontal; Hg, Harderian gland; is, interorbital septum; j, jugal; le, lens; m, eye muscles; nse, nasal septum; ob, olfactory bulb and peduncle; p, parietal; pf, prefrontal; po, postorbital; ps, planum supraseptale; pt, palatine; so, scleral ossicle; I, olfactory nerves; II, optic nerve.

developing in the basal plate on either side of the notochord. This rod might represent either a taenia intertrabecularis or an intertrabecula; these structures are characteristic of the developing skulls of some Testudines (Bellairs and Kamal, 1981). Another possibility is that it represents the posterior parts of the trabeculae, which have undergone abnormal fusion in this region since abnormal trabecular fusion has been noted in amphibian embryos with experimentally produced cyclopia (de Beer, 1937). The more posterior parts of the skull of this specimen, the otic capsule and inner ear structures, show no gross malformations.

DISCUSSION AND CONCLUSIONS

Of the two specimens examined histologically, A is by far the more abnormal, showing partial fusion of the eyes with presence of a third eyelike structure, a well-developed proboscis as in typical mammalian cyclopia, absence of the pituitary and of inner ear structures, as well as other gross abnormalities which affect almost every part of the head. It is impossible from external examination to state whether similar abnormalities are present in the unsectioned specimens B and C.

Specimen D, though not actually cyclopic, seems to show a less extreme expression of the cyclopic trend. Although the eyes are quite separate, the two nasal sacs and their capsules appear to have fused into a largely single structure. As in A, the premaxillary component of the upper jaw is missing and there is virtual absence of the midline elements of the anterior chondrocranium.

The ontogenetic development of cyclopia and related conditions seems to stem from some deficiency of the front of the neural plate (or the neural tube) and of associated mesoderm manifesting itself at a very early stage in development. Cyclopia has been produced experimentally by surgical procedures de-

signed to produce such defects (Johnston *et al.*, 1977). Coalescence or approximation of the eyes and of the nasal placodes might be expected to occur if there was a reduction in the amount of tissue present in the anterior midline of the early embryo. Moreover, such approximation of paired structures might well prevent the tissues of the frontonasal process (which forms the anterior and median part of the face) from growing ventrally and giving rise to the premaxillary region of the upper jaw. Such material, prevented from reaching its normal destination, may produce instead an abnormal, proboscis-like structure.

Deficiencies of the cranial neural plate would also involve deficiencies of the cranial part of the neural crest. It is well known that neural crest tissue normally gives rise to branchial arch elements, including the trabeculae, and to other parts of the facial skeleton (Le Lièvre, 1978). It is thus possible to explain the absence of nasal and interorbital septa, which are at least partly of trabecular origin (Bellairs and Kamal, 1981), as well as the more posterior parts of the trabeculae which normally remain paired.

The albinoid condition of these embryos suggests some generalized, rather than a purely cranial, deficiency of the neural crest. It is noteworthy that Caldwell (1959) found albinoid embryos with malformations of the jaws and eyes in natural nests of *Caretta*, though cyclopia is not explicitly mentioned in his account.

ACKNOWLEDGMENTS

I am most grateful to P. C. H. Pritchard for supplying these specimens and relevant information about them, to A. McGehee for allowing me access to data in her unpublished thesis, and to M. W. Ferguson, A. D. Hoyes, and G. M. Morriss-Kay for helpful comments. My thanks are also due to B. G. H. Martin and to the Audio-visual Department of St.

Mary's Hospital Medical School for the photographs.

LITERATURE CITED

- ADELMANN, H. B. 1936. The problem of cyclopia. *Q. Rev. Biol.*, **11**: 161-182 and 284-304.
- BELLAIRS, A. D'A. 1965. Cleft palate, microphthalmia and other malformations in embryos of lizards and snakes. *Proc. Zool. Soc. Lond.*, **144**: 239-251.
- BELLAIRS, A. D'A., AND A. M. KAMAL. 1981. The chondrocranium and the development of the skull in Recent reptiles. In C. Gans and T. S. Parsons (eds.), *Biology of the Reptilia*. London and New York, Academic Press, **11**: 1-263.
- CALDWELL, D. K. 1959. The Atlantic loggerhead sea turtle, *Caretta caretta caretta* (L.), in America. III. The loggerhead turtles of Cape Romain, South Carolina. *Bull. Fla. St. Mus. Biol. Sci.*, **4**(10): 319-348.
- DE BEER, G. R. 1937. *The development of the vertebrate skull*. Oxford, Clarendon Press.
- EWERT, M. A. 1979. The embryo and its egg: development and natural history pp. 333-413. In M. Harless and H. Morlock (eds.), *Turtles. Perspectives and research*. New York, John Wiley.
- JOHNSTON, M. C., G. M. MORRISS, D. C. KUSHNER, AND G. J. BINGLE. 1977. Abnormal organogenesis of facial structures, pp. 421-451. In J. G. Wilson and F. C. Fraser, (eds.), *Handbook of teratology*. New York, Plenum Press.
- LE LIÈVRE, C. S. 1978. Participation of neural crest-derived cells in the genesis of the skull in birds. *J. Embryol. Exp. Morphol.*, **47**: 17-37.
- MCGEHEE, M. A. 1979. Factors affecting the hatching success of loggerhead sea turtles. Master's thesis, University of Central Florida, 1-252.
- MCGRATH, P. 1981. Absence of the presphenoid in relation to the neural and facial abnormalities in human cyclopia. *J. Anat.* **133**: 473-474.
- ROMANOFF, A. L. 1972. *Pathogenesis of the avian embryo*. New York, Wiley-Interscience.
- SCHWALBE, E., AND H. JOSEPHY. ca. 1913. Die Missbildungen des Kopfes. II. Die Cyclopie pp. 205-246. In *Die Morphologies der Missbildungen des Menschen und der Tiere*. (1906-1958) G. Fischer, Jena, 3 Teil, Kapitel 5.
- WILLIS, R. A. 1958. *The borderland of embryology and pathology*. London, Butterworth.

ADDENDUM

An association between craniofacial defects and albinism in turtles has also been noted by Ewert (1979) and by J. D. Miller (personal communication) and may reflect neural crest deficiency. McGrath (1981) reports absence of the presphenoid in cases of human cyclopia. This bone would probably correspond with part of the unfossified interorbital septum of reptiles.

Neural Pattern—A Neglected Taxonomic Character in the Genus *Pelusios* Wagler (Pleurodira: Pelomedusidae)

DONALD G. BROADLEY¹

ABSTRACT. Variation in the pattern of neural bones in the genus *Pelusios* is documented and shown to be a useful taxonomic character at the species level with respect to both Recent and fossil material. Neural pattern is also documented for *Pelomedusa subrufa*.

INTRODUCTION

The taxonomy of the African hinged terrapins of the genus *Pelusios* Wagler has long been a source of controversy, and numerous taxonomic characters have been employed by various workers. In the last comprehensive revision of the Recent species of the genus, Loveridge (1941) recognized only four species, i.e., *P. adansonii* (Schweigger), *P. gabonensis* (Duméril), *P. subniger* (Lacépède), and *P. sinuatus* (A. Smith). Müller and Hellmich (1954) subsequently reinstated *P. niger* (Duméril and Bibron) as a valid species. Laurent (1956, 1964, 1965) resurrected two more species, *castaneus* (Schweigger) and *bechuanicus* Fitz-Simons, from the synonymy of *P. subniger* and also described three new species, *nanus*, *carinatus*, and *williamsi* (with a subspecies *lutescens*) from Central Africa. He also revived the taxa *derbianus* (Gray) and *rhodesianus* Hewitt as subspecies of *P. castaneus* and described a new subspecies *P. c. chapini*. Raw

(1978) restored *P. rhodesianus* to specific rank when he found it to be sympatric with *P. castaneus* in Zululand. Bour (1978) proposed that the name *P. castaneus castaneus* should be restricted to the West African populations previously included under *P. c. derbianus* and he erected the name *P. castaneus kapika* for the Malagasy populations. Broadley (1981) confirmed the status of *P. rhodesianus* as a full species, sympatric with *P. castaneus castanoides* Hewitt in Zululand, where the two species are readily distinguished on neural pattern.

Laurent (1965) described *P. williamsi* as a full species because its yellowish western race *lutescens* is sympatric with the blackish *P. castaneus chapini* in the Lake Albert (Lac Mobutu Sese Seko) basin. Broadley (1981) noted sympatry between blackish *P. rhodesianus* and yellowish (but usually appearing black through soil staining) *P. w. williamsi* at Entebbe, Uganda. The neural patterns of the few specimens examined from this region suggest that *P. williamsi* is actually conspecific with *P. castaneus* and the subspecies *chapini* is conspecific with *P. rhodesianus*.

A review of the six southern African species of *Pelusios* (Broadley, 1981) generally confirms the taxonomy established by previous authors (Laurent, 1956, 1964, 1965; Raw, 1978). It also shows that the pattern of neural bones in the carapace provides a useful taxonomic character neglected by previous workers

¹National Museum, P.O. Box 240, Bulawayo, Zimbabwe.

on this genus. In the present paper, the survey of neural patterns is extended to cover the remaining species of *Pelusios*, plus *Pelomedusa subrufa* (Lacépède).

This paper is dedicated to Dr. Ernest E. Williams, who made a major contribution to our understanding of the evolution of the genus *Pelusios* in his 1954 paper.

MATERIALS AND METHODS

The initial review of neural patterns in *Pelusios* was based on osteological specimens in the National Museum collection. The neural arrangement was also checked on many complete specimens (both wet and dry) by stripping the vertebral shields from the carapace, as neural patterns cannot be identified on X-rays. With the cooperation of several Curators, I was able to extend the investigation to additional specimens on loan from other institutions.

The nomenclature of bones of the shell follows Zangerl (1969). Neurals are numbered from the front N1 to N8.

Most abbreviations for institutional prefixes are indicated under Acknowledgments. UM and NMZB (formerly NMSR) indicate two sets of catalogue numbers in the Department of Herpetology at the National Museum of Zimbabwe. † indicates extinct forms.

NEURAL BONES IN THE PLEURODIRA

Zangerl (1948) noted that the Chelidae differed from the Pelomedusidae in showing a strong tendency towards reduction of neural bones. The Neotropical *Phrynops nasutus* has only four neurals (N1, N6, N7, and N8 absent; Boulenger, 1889: Fig. 58) and *Platemys platycephala* lacks neurals. *Chelodina oblonga*, isolated in the southwest corner of Australia, is the only Australasian chelid with five to eight neurals (Burbidge *et al.*, 1974). Although the other taxa normally lack

neurals, one to four vestigial and usually isolated neurals are occasionally present in three other species of *Chelodina* and two species of *Elseya* (Rhodin and Mittermeier, 1977).

Recent Pelomedusidae do not show such striking reductions in neurals, *Podocnemis* has seven (rarely 6), *Erymnochelys* six, *Pelomedusa* seven (rarely 6 or 8), and *Pelusios* five to eight (rarely 4). However, the Pliocene *Podocnemis venezuelensis* lacked neurals (Wood and Gamero, 1971) as did the Eocene *Eusarkia rotundiformis* Bergounioux of Tunisia (Moody, 1972).

Although no species of *Pelusios* has less than four neurals, and there is intra-specific variation in the number and size of those present, neurals may be lost at either or both ends of the series, so that sympatric taxa may be distinguishable on neural pattern, e.g., *P. rhodesianus*, *P. castaneus castanoides*, and *P. sinuatus* in Zululand. Neural patterns are apparently nonadaptive and are therefore particularly useful taxonomic characters at the species level.

NEURAL PATTERNS IN PELOMEDUSA AND PELUSIOS

Boulenger (1889) stated that in the Pleurodira the neural series rarely contains more than seven bones. A single 'pygal' (=suprapygal) is present, and in none of the Recent forms does it make contact with the neurals. He illustrated the neural pattern of a *Sternothaerus derbianus* (= *Pelusios c. castaneus*) with a continuous series of eight neurals, separated from the single suprapygal (Boulenger, 1889: Fig. 47).

The first author to report a reduction in the number of neural bones in *Pelusios* was Hewitt (1933), who illustrated a paratype of *P. sinuatus leptus* with only five neurals, N1 failing to contact the nuchal and N5, N7, and N8 absent. He also recorded a Zululand specimen of *P. sinuatus zuluensis* Hewitt with a continuous series of seven neurals.

In his diagnosis of the family Pelomedusidae, Zangerl (1948) noted the presence of six to eight neural bones.

Broin (1969) illustrated the variation in neural pattern in nine Recent and fossil specimens of *P. sinuatus*, including the type of *P. rudolphi*† Arambourg. These have five to seven neurals: all lack N8, most lack N7, one lacks N6, and five have N5 reduced or absent, so that N6 is isolated.

In a review of the genus *Pelusios* south of Latitude 8°S Broadley (1981) checked the neural arrangement in 136 specimens from this region. In order to expand the study to cover the remaining species in the genus I have examined an additional 33 specimens from north of Latitude 8°S and incorporated data from the literature (Hewitt, 1933; Broin, 1969). For comparison, neural patterns for 25 specimens of *Pelomedusa subrufa* were also recorded. The neural patterns found in *Pelomedusa* and *Pelusios* are described below and summarized in Table 1.

Pelomedusa subrufa (Lacépède)

Twenty-five specimens were examined. N8 is absent in 20 and reduced in three others; two specimens also lack N7. N1 is narrowed anteriorly in some of these specimens, but it is reduced and widely separated from the nuchal in FMNH 17160 (Zangerl, 1948).

Genus *Pelusios*

P. rusingae† Williams

The type and only known specimen of this species is the earliest known shell belonging to the genus *Pelusios* (Williams, 1954, Pl. 1). The neural pattern is complete, consisting of an unbroken series of eight neurals, N1 in good contact with the nuchal and N8 well separated from the single suprapygal (Fig. 1A).

P. gabonensis (Duméril)

Five specimens have been examined. BM 1912.6.27.38 has a similar arrangement to *P. rusingae*. BM 1912.6.27.40 has N1 reduced, but N8 elongate and in contact with the suprapygal. Three others all lack N1, but the first of these specimens has a small N9 (Fig. 1B).

P. adansonii (Schweigger)

All seven specimens examined lack N1 (Fig. 1C). BM 1900.9.22.7 also lacks N8, but BM 1900.9.22.6 has a small N9.

P. nanus Laurent

Neural patterns were determined on four specimens. AMNH 50757 has eight neurals, but N8 is reduced in size and meets N7 at a point (Fig. 1D); three others lack N8. N1 always tapers anteriorly, meeting the nuchal at a point or just failing to make contact.

P. niger (Duméril and Bibron)

Three specimens were examined. Two have N1 reduced or absent and N8 elongate and in contact with, or narrowly separated from, the suprapygal (which is divided into three in one specimen, Fig. 1E). BM 1927.9.27.237 lacks N1 and N8.

P. subniger (Lacépède)

Of 25 specimens examined for neural pattern, 23 have a continuous series of eight neurals, N1 in good contact with the nuchal, N8 well separated from the suprapygal. Three of these specimens (UM 850; 3181–2) have two superposed suprapygals (Fig. 1F). Two specimens (AM 5432; UM 9720) lack N8.

P. bechuanicus bechuanicus FitzSimons

All 13 specimens examined have a continuous series of eight neurals separated

from the suprapygal (Fig. 1G). In UM 9718 and UM 29157, N1 is reduced and does not contact the nuchal; in addition, the latter specimen has N8 transversely divided.

P. bechuanicus upembae Broadley

Both specimens examined have eight neurals, but MRAC 11460 has N8 transversely divided.

P. rhodesianus Hewitt

Twenty-eight specimens were examined. Nine have a continuous series of eight neurals with N1 in good contact with the nuchal and N8 contacting the suprapygal (Fig. 1H), 15 others have N8 separated from the suprapygal. Two have N7 and N8 reduced in size and isolated, another has only N8 isolated. In MRAC 16962 N8 is absent.

P. castaneus castaneus (Schweigger)

Five specimens were examined, including the specimen illustrated by Boulenger (1889: Fig. 47). All have a continuous series of eight neurals with N1 in good contact with the nuchal and N8 well separated from the suprapygal (Fig. 2A).

P. castaneus williamsi Laurent

Two specimens from the type locality were examined. One lacks N1, and in both specimens N8 is elongate and only narrowly separated from the suprapygal (Fig. 2B).

P. castaneus lutescens Laurent

The paratype examined has seven neurals, N1 absent and N8 reduced in size.

P. castaneus castanoides Hewitt

Eighteen specimens were examined. The number of neurals varies from five to

eight, but N1, N7, and N8 are always reduced in size or absent. The Malagasy population (*P. castaneus kapika* Bour) seems to be indistinguishable from mainland populations. Geographical variation is as follows:

Malawi: N1, N7 and N8 absent in six specimens.

Moçambique: One specimen has N1 reduced, N7 absent and N8 minute (Fig. 2d).

Zululand: Five neurals in three specimens, six in two (Fig. 2C), and seven in two.

Madagascar: One specimen has N1, N7, and N8 absent.

Seychelles: Three specimens have seven or eight neurals, with N1 and N7 small to minute, N8 minute or absent.

P. carinatus Laurent

The paratype examined has eight neurals, but N7 and N8 are both very small and N8 is isolated (Fig. 2E).

P. sinuatus (A. Smith)

Neural patterns were examined in 55 specimens and complete patterns for an additional four Recent and four fossil specimens are illustrated by Broin (1969). There are four to seven neurals, N8 always absent. N1 is reduced (not contacting the nuchal) in six specimens, absent in only one (UM 32996). N5 is absent in five, reduced in 13 and divided in one (Fig. 2G); N6 is absent in five and reduced in four; N7 is absent in 52 specimens and reduced in four others.

Specimens with seven neurals (Fig. 2F) are more common in the south. A reduction to five (rarely 4) neurals seems to be restricted to populations in Lake Tanganyika and the Luangwa Valley in eastern Zambia (Fig. 2H), which attain the maximum sizes recorded for any *Pelusios* (maximum carapace length 465 mm in Lake Tanganyika; Witte, 1952).

Usually it is N5 that is missing or reduced, less commonly N6, rarely both (MCZ 48019). It should be noted that three specimens from the early Pleistocene of Omo, north of Lake Turkana, dated at 3.75 to 1.8 million years B.P., all have N5 absent (type of *P. rudolphi* Arambourg) or reduced and N7 and N8 absent (Broin, 1969). A specimen from Bed I, Pleistocene of Olduvai (BM R5761) also has six neurals, with N5 reduced and N1 not contacting the nuchal. The Recent Somali specimen of *P. sinuatus* illustrated by Broin (1969) has N5 reduced, but another from Uebi Shabeli River (MF 20647) shows no reduction of N5.

PHYLOGENY OF *PELUSIOS*

The earliest known fossil *Pelusios* shell is the type specimen of *P. rusingae*[†], described from the early Miocene of Rusinga Island, Lake Victoria, Kenya, by Ernest E. Williams (1954). The other Miocene species, *P. blanckenhorni*[†] Dacqué, is known only from a skull and hence could equally prove to be a *Pelomedusa* (Williams, 1954; Wood, 1974). *P. dewitzianus*[†] von Reinach is known only from fragments from the middle Pliocene of Egypt. *P. rudolphi* Arambourg from the early Pleistocene of Omo, Ethiopia, appears to be indistinguishable from Recent *P. sinuatus* (Broin, 1969).

Williams (1954) suggested that small lateral mesoplastra are primitive for the Pelomedusidae, as this condition is found in all known Cretaceous pelomedusids. This in turn implies that the large mesoplastra with median contact in *Pelusios* is a secondarily derived condition associated with the development of the plastral hinge. *Pelomedusa subrufa*, the only other African pelomedusid species, retains small lateral mesoplastra.

In the first couplet of his key to the genus *Pelusios*, Loveridge (1941) separated the species *adansonii* and *gabonensis*, which have the anterior lobe of

the plastron more than twice the length of the suture between the abdominal shields, from *subniger* and *sinuatus*, in which the anterior plastral lobe is less than twice the length of the abdominal suture. Williams (1954) endorsed this division into two species groups when he described *P. rusingae*[†].

The four species of what may be called the *P. gabonensis* group form an evolutionary sequence based on shape of mesoplastra. In the Miocene species *P. rusingae*[†] the mesoplastra are strongly tapered medially both anteriorly and posteriorly, hardly meeting. In *P. gabonensis* the mesoplastra are moderately tapered medially both anteriorly and posteriorly, forming oblique sutures with both hyo- and hypoplastra. In *P. adansonii* the mesoplastra are tapered only posteriorly, anteriorly forming a straight transverse hinge with the hypoplastra. In *P. nanus* the mesoplastra are only slightly tapered posteriorly.

In the *P. subniger* group the mesoplastra are not tapered, forming straight transverse sutures with both hyo- and hypoplastra. The taxonomy of this group has always presented problems because of the shortage of clear-cut morphological characters.

P. niger has a distinctive skull with a narrow snout and well-developed beak (Boulenger, 1889: Fig. 46; Müller and Hellmich, 1954: Figs. 3, 4; Gaffney, 1979: Figs. 132, 133). Although Wood (1974) placed this species in the *P. gabonensis* group, it has a short anterior plastral lobe, the mesoplastra do not taper medially, and it is a relatively large species (four adults in the British Museum are 250 to 277 mm in carapace length). *P. subniger* and *P. bechuanicus* may be distinguished on the relatively large head size (Broadley, 1981). The other species in this group are largely distinguished on the proportions of the epidermal shields and coloration (Laurent, 1956, 1964, 1965; Broadley, 1981), but neural patterns provide a useful new character.

Figure 1. Neural patterns in *Pelusios*. A. *P. rusingaet* (Type—Rusinga Island, Lake Victoria). B. *P. gabonensis* (CM 39662—Nyabessem, Cameroun). C. *P. adansonii* (UM 33631—Bahr el Ghazel, Sudan). D. *P. nanus* (AMNH 50757—Chitau, Angola). E. *P. niger* (BM 1974.3012—Nko, Nigeria). F. *P. subniger* (UM 850—Lake Kariba, Zimbabwe). G. *P. bechuanicus* (UM 32985—Kasane, Botswana). H. *P. rhodesianus* (Type—Mpika District, Zambia). Thick lines indicate sulci between epidermal shields, thin lines indicate sutures between bones. The horizontal lines equal 1 cm to scale.

Figure 2. Neural patterns in *Pelusios*. A. *P. castaneus castaneus* (UM 33495—Ghana). B. *P. castaneus williamsi* (UM 33165—Kaimosi, Kenya). C. *P. castaneus castanoides* (TM 52160—Lake St Lucia, Zululand). D. *P. castaneus castanoides* (UM 27828—Tando, Mocambique). E. *P. carinatus* (UM 33158—Eala, Zaire); F. *P. sinuatus* (UM 12076—Lundi River, Zimbabwe). G. *P. sinuatus* (UM 5278—Bumi Confluence, Lake Kariba, Zimbabwe). H. *P. sinuatus* (AM 5432—Mpika District, Zambia). Conventions as in Fig. 1.

TABLE 1. AFRICAN PELOMEDUSIDAE: DEVIATIONS FROM THE ANCESTRAL CONDITION OF 8 NEURALS, THE FIRST IN CONTACT WITH THE NUCHAL, THE LAST SEPARATED FROM THE SUPRAPYGAL.

TAXON	N	No. of neurals*	N1		N5		N6		N7		N8		
			Reduced	Absent	Reduced	Absent	Reduced	Absent	Reduced	Absent	Enlarged	Reduced	Absent
PELOMEDUSA													
<i>subrufa</i>	25	(6)7(8)	—	—	—	—	—	—	—	2	—	3	20
PELUSIOS													
<i>rusingae</i> [†]	1	8	—	—	—	—	—	—	—	—	—	—	—
<i>gabonensis</i>	5	7-8	1	3	—	—	—	—	—	—	1	—	—
<i>adansonii</i>	7	(6)7(8)	—	7	—	—	—	—	—	—	—	—	1
<i>nanus</i>	4	7(8)	—	—	—	—	—	—	—	—	—	1	3
<i>niger</i>	3	6-8	1	2	—	—	—	—	—	—	2	—	1
<i>subniger</i>	25	(7)8	—	—	—	—	—	—	—	—	—	—	2
<i>bechuanicus bechuanicus</i>	13	8	2	—	—	—	—	—	—	—	—	—	—
<i>bechuanicus upembae</i>	2	8	—	—	—	—	—	—	—	—	—	—	—
<i>rhodesianus</i>	28	(7)8	—	—	—	—	—	—	3	—	9	3	1
<i>castaneus castaneus</i>	5	8	—	—	—	—	—	—	—	—	—	—	—
<i>castaneus williamsi</i>	2	7-8	—	1	—	—	—	—	—	—	2	—	—
<i>castaneus lutescens</i>	1	7	—	1	—	—	—	—	—	—	—	1	—
<i>castaneus castanoides</i>	18	5-8	6	12	—	—	—	—	5	13	—	5	13
<i>carinatus</i>	1	8	—	—	—	—	—	—	1	—	—	1	—
<i>sinuatus</i>	63	(4)5-7	5	1	10	5	4	4	4	53	—	—	63

*Rare conditions are placed in parentheses. Numbers of specimens with various neurals enlarged (i.e., N8 contacting or nearly contacting suprapyg), reduced in size or absent, are indicated for each taxon.

Zangerl (1948) considered that the primitive pelomedusid carapace had eight neural bones, the first adjoining the nuchal. The limited number of neural patterns examined for the *P. gabonensis* group indicate a primitive condition for *P. rusingae*[†] with no reduction in neurals; *P. gabonensis* and *P. adansonii* usually lack N1, whereas in *P. nanus* N1 is narrowed anteriorly and N8 is reduced or absent, placing this species on a different evolutionary line.

In the *P. subniger* group, the two broad-headed species *P. subniger* and *P. bechuanicus* retain the full complement of 8 neurals, but *P. niger* has N1 reduced or absent.

The *P. rhodesianus*/*P. castaneus* assemblage exhibits a variety of patterns, but in the latter species there seems to be a stepped east-west cline in reduction of neurals, with N1 being lost first, then N7

and N8 simultaneously. The neural pattern of *P. castaneus castanoides*, with N1, N7, and N8 all reduced or absent, is very distinctive.

The *P. carinatus* neural pattern (N7 and N8 reduced) appears to represent a stage in the evolution of the *P. sinuatus* condition (N8 absent, other neurals lost in the sequence N7, N5, rarely N6). The two species are allopatric, with *P. carinatus* in the Zaire (Congo) basin, and *P. sinuatus* widespread in eastern Africa, but formerly extending westward to Chad (Broin, 1969).

It is hoped that this brief review of neural patterns in *Pelusios* will encourage future workers to explore the range of variation in the poorly known northern forms and also the fossil material from the fluvial and lacustrine deposits east of Lake Turkana (Rudolf) (Behrensmeyer, 1975).

ACKNOWLEDGMENTS

I am grateful to the following colleagues for assistance: P. H. Skelton and J. C. Greig of the Albany Museum, Grahamstown (AM); W. D. Haacke and Ms. L. Wessels of the Transvaal Museum (TM); D. F. E. Thys van den Audenaerde of the Musée Royal de l'Afrique Centrale, Tervuren (MRAC); R. G. Zweifel of the American Museum of Natural History (AMNH); C. J. McCoy and A. V. Bianculli of the Carnegie Museum (CM); E. E. Williams and J. P. Rosado of the Museum of Comparative Zoology (MCZ); R. H. Parker of FitzSimons' Snake Park, Durban (DSP); Lynn Raw of Pietermaritzburg (LR); C. J. McCarthy of the British Museum Natural History (BM); J. P. Gosse of the Institut Royal des Sciences Naturelles de Belgique (IRScNB); and B. Lanza of the Museo Zoologico de 'La Specola', Florence (MF).

I wish to thank M. L. Angell for photographic assistance, A. Spector for the donation of X-ray photographs, and M. Clift and B. L. Bennefield for typing various versions of the manuscript.

I am grateful to J. B. Iverson for commenting on the first draft of this paper and to the reviewers for their helpful suggestions.

APPENDIX: SPECIMENS EXAMINED FOR NEURAL PATTERN

Pelomedusa subrufa. GHANA: NMZB 6004. ZAMBIA: NMZB 2629. MALAWI: UM 3184, 33022. BOTSWANA: UM 7430, 11603-4, 14708, 29152-6, 32968. ZIMBABWE: UM 3174, 5277, 20584, 32950, 32983, 33021, 33029, 33130. SOUTH AFRICA: NMZB 6020; UM 33423-4.

GENUS *PELUSIOS*

Pelusios gabonensis. CAMEROUN: BM 1912.6. 27.38 and 40; CM 39662-3. GABON: UM 33431.

Pelusios adansonii. NIGERIA: BM 1970.1797. SUDAN: BM 1900.9.22.6 and 7; UM 33631; USNM 75099, 75105, 75107.

Pelusios nanus. ANGOLA: AMNH 50757, 50760; UM 33262. ZAMBIA: NMZB 3958.

Pelusios niger. GHANA: BM 1927.9.27.237. NIGERIA: BM 1974.3012-3.

Pelusios subniger. ZAMBIA: AM 5144, 6574, 6876; KM—; NMZB 3150; TM 38181; UM 6574. MALAWI: MCZ 51100; UM 25462. BOTSWANA: TM 46229. ZIMBABWE: UM 849, 850, 3181, 3182, 9720, 14517, 20370, 32966, 32986, 32987, 33170. MOÇAMBIQUE: UM 9713, 9714, 28345, 28842.

Pelusios bechuanicus bechuanicus. BOTSWANA: UM 12938, 16191, 22338, 29150, 29151, 29157, 32984, 33317. ZAMBIA: UM 791, 9718, 10673, 10677, 15594.

Pelusios bechuanicus upembae. ZAIRE (Shaba Province): MRAC 11460 (paratype); TM 38178 (holotype).

Pelusios rhodesianus. UGANDA: AM—(2); BM 1906.5.30.1; UM 33389. ANGOLA: AMNH 50752, 50753. ZAIRE: MRAC 16962, 16964, 16965; UM 33392. ZAMBIA: AM—(holotype), 5432 (2); NMZB 2625, 2626; UM 47, 49. ZIMBABWE: UM 9715, 14516, 21654, 33033, 33038. MOÇAMBIQUE: UM 9717. ZULULAND: TM 52341, 52342. NATAL (Durban): DSP 86, 87; UM 33621.

Pelusios castaneus castaneus. 'WEST AFRICA': BM 63.10.8.5. SENEGAL: CM 24785. GHANA: UM 33495. NIGERIA: AM 7282; BM 1974.3014.

Pelusios castaneus williamsi. KENYA: TM 16433; UM 33165 (paratype).

Pelusios castaneus lutescens. ZAIRE: CM 62248 (paratype).

Pelusios castaneus castanoides. MALAWI: AM T15, T16, T18, T19; UM 32990, 33390. MOÇAMBIQUE: UM 27828. ZULULAND: LR 1038, TM 13433 (holotype), 45644, 48318, 52160, 52544; UM 33634. MADAGASCAR: UM 33166. SEYCHELLES: AM—; BM 1911.4.7.2; TM 49338.

Pelusios carinatus. ZAIRE: UM 33158 (paratype).

Pelusios sinuatus. SOMALIA: MF 20647. TANZANIA: MCZ 30013, 48018, 48019. ZAIRE: BM 1953.1.11.48; IRScNB 3, 33, 99. ZAMBIA: AM—(holotype of *P. s. leptus* Hewitt), 5432, 5794, 6696, IRScNB 7143; NMZB 3151; UM 762, 17528, 23583, 32996, 32998, 33028. MALAWI: UM 4851, 25461, 33031, 33040. ZIMBABWE: NMZB 1210, 3700, 3751; UM 3178, 5278, 5856, 5867, 9799, 12076, 32975, 32976, 33029, 33030, 33032, 33039, 33386-8. MOÇAMBIQUE: UM 3157, 10343, 27585, 30436. BOTSWANA: UM 11255. TRANSVAAL: AM 7361; TM 15076. ZULULAND: TM 34682, 34684.

LITERATURE CITED

- BEHRENSMEYER, A. K. 1975. The Taphonomy and Paleocology of Plio-Pleistocene Vertebrate assemblages east of Lake Rudolf, Kenya. *Bull. Mus. Comp. Zool.*, 146: 473-578.
- BOULENGER, G. A. 1889. Catalogue of the Chelonians, Rynchocephalians and Crocodiles in the British Museum (Natural History). London, British Museum (Nat. Hist.), 311 pp.

- BOUR, R. 1978. Les tortues actuelles de Madagascar (République malagache): liste systématique et description de deux sous-espèces nouvelles (Reptilia—Testudines). *Bull. Soc. Étud. scient. Anjou N.S.*, **10**: 141–154.
- BROADLEY, D. G. 1981. A review of the genus *Pelusios* Wagler in southern Africa (Pleurodira: Pelomedusidae). *Occas. Pap. Natl. Mus. Zimbabwe, B.*, **6**: 635–683.
- BROIN, F. DE. 1969. Sur la présence d'une Tortue, *Pelusios sinuatus* (A. Smith) au Villafranchien inférieur du Tchad. *Bull. Soc. géol. Fr.*, **11**: 909–916.
- BURBIDGE, A. A., J. A. W. KIRSH, AND A. R. MAIN. 1974. Relationships within the Chelidae (Testudines: Pleurodira) of Australia and New Guinea. *Copeia*, **1974**: 392–409.
- GAFFNEY, E. S. 1979. Comparative cranial morphology of Recent and fossil turtles. *Bull. Amer. Mus. Nat. Hist.*, **164**: 65–376.
- HEWITT, J. 1933. Descriptions of some new reptiles and a frog from Rhodesia. *Occ. Pap. Rhod. Mus.*, **1**: 45–50.
- LAURENT, R. F. 1956. Contribution à l'Herpetologie de la Région des Grands Lacs de l'Afrique centrale. *Annls Mus. r. Congo belge, Sér. 8vo*, **48**: 1–390.
- . 1964. Reptiles et amphibiens de l'Angola (troisième contribution). *Publcoes cult. Co. Diam. Angola*, **67**: 1–165.
- . 1965. A contribution to the knowledge of the genus *Pelusios* (Wagler). *Annls Mus. r. Afr. cent., Sér. 8vo*, **135**: 1–33.
- LOVERIDGE, A. 1941. Revision of the African terrapin of the family Pelomedusidae. *Bull. Mus. Comp. Zool.*, **88**: 462–524.
- MOODY, R. T. J. 1972. The turtle fauna of the Eocene phosphates of Metlaoui, Tunisia. *Proc. Geol. Ass.*, **83**: 327–335.
- MÜLLER, L., AND W. HELLMICH. 1954. Zur Kenntnis einiger *Pelusios*-Arten (Testudines). *Veröff. zool. St. Samml. Münch.*, **3**: 51–79.
- RAW, L. R. G. 1978. Taxonomic notes on the hinged terrapins, genus *Pelusios*, of Natal. *Durban Mus. Novit.*, **11**: 287–294.
- RHODIN, A. G. J., AND R. A. MITTERMEIER. 1977. Neural bones in chelid turtles from Australia and New Guinea. *Copeia*, **1977**: 370–372.
- WILLIAMS, E. E. 1954. A new Miocene species of *Pelusios* and the evolution of that genus. *Breviora Mus. Comp. zool. No. 25*, pp. 1–7.
- WITTE, G.-F. DE. 1952. Amphibians and Reptiles. *Explor. hydrobiol. Lac. Tanganyika (1946–1947)*, **3(3)**: 1–22.
- WOOD, R. C. 1974. The systematics, evolution and zoogeography of African turtles. *Natn. geogr. Soc. Res. Rep. 1967 projects*: 301–306.
- WOOD, R. C., AND M. L. D. DE GAMERO. 1971. *Podocnemis venezuelensis*, a new fossil pelomedusid (Testudines, Pleurodira) from the Pliocene of Venezuela and a review of the history of *Podocnemis* in South America. *Breviora Mus. Comp. Zool. No. 376*, pp. 1–23.
- ZANGERL, R. 1948. The vertebrate fauna of the Selma formation of Alabama. Part II. The Pleurodiran turtles. *Fieldiana, Geol. Mem.*, **3**: 17–56.
- . 1969. The turtle shell, pp. 311–339. *In* C. Gans (ed.), *Biology of the Reptilia. 1, Morphology*. A. London & New York, Academic Press.

The Genus *Emydura* (Testudines: Chelidae) in New Guinea with Notes on the Penial Morphology of *Pleurodira*

SAMUEL B. McDOWELL¹

ABSTRACT. Two species of *Emydura* occur in New Guinea: *E. dentata* (synonyms: *Elseya dentata*, *Emydura novaeguineae*, *Elseya novaeguineae*) and *E. australis* (synonyms: *Emydura krefftii*, *E. subglobosa*, *E. albertisii*). The latter is closely related to *E. dentata* and probably derived from it. *E. dentata* probably also lies close to the ancestry of *E. latisternum* (northeast Australian), which in turn, appears closely related to *E. macquaria* (southeast Australian) and *Pseudemydura umbrina* (western Australian). *Emydura dentata* is the most primitive and pelomedusid-like of Australian Chelidae. The other Australo-New Guinean chelid genus, *Chelodina*, is not morphologically close to *Emydura*, but instead seems closer to South American forms. The penis of *Pleurodira* is intermediate between the protrusile erectile organ of Cryptodira and cloacal eversion as seen in caecilians. The pleurodiran penis, formed of an almost cartilagelike corpus fibrosum tightly bound to a highly expansible corpus cavernosum, warps when the corpus cavernosum expands while the corpus fibrosum remains of constant size. This warping doubles the penis over and acts as a lever to evert the cloaca as a sheath around the base of the protruded penis. This is interpreted as the primitive form of the penis for Testudines. Penial morphology offers useful taxonomic characters in *Pleurodira*, and the two New Guinea species of *Emydura* differ in penial structure.

INTRODUCTION

It seems fitting that any commemorative volume to Ernest E. Williams should include some discussion of turtles, and since it was during his curatorship of herpetology at the Museum of Comparative

Zoology that the MCZ acquired the largest collection of New Guinea reptiles in the world, it seems equally fitting to discuss New Guinea turtles. Accordingly, I here report on the side-neck terrapin of the genus *Emydura* in New Guinea, based mainly on those in the collection of the American Museum of Natural History (AMNH). I have purposely omitted use of the MCZ collection for framing the hypothesis presented below, so that the MCZ collection may be used for the testing of this hypothesis.

Prior to Goode (1967), the taxonomic arrangement and diagnostic characters used for *Emydura* (and the supposedly distinct genus *Elseya*) followed the lines laid down by Boulenger (1889) and modified to accommodate additional named forms by de Rooij (1915). This arrangement recognized: *Elseya*, with one species, *E. dentata*, characterized by a longitudinal ridge on the maxillary triturating surface; and *Emydura*, for the remaining forms (other than *Pseudemydura umbrina* of Western Australia, not described until 1901), characterized by lack of a ridge on the triturating surface of the maxillary bone.

Goode (1967) radically redefined *Emydura* and *Elseya*, the latter diagnosed by a horny shield on the crown of the head which developed at an early age, and by presence of erect tubercles on the dorsal surface of the neck; *Emydura* was restricted to those forms with soft skin on the top of the head (except in

¹Department of Zoology and Physiology, Rutgers University, Newark, New Jersey 07102, U.S.A.

some old adults) and with low and inconspicuous tubercles on the neck. By this redefinition, the species *novaeguineae* and *latisternum* were transferred to *Elseya* and *Emydura* was restricted to *E. krefftii*, *E. australis*, and *E. macquaria* (the type). Goode also much simplified the taxonomy by reducing many names to synonymy (my own findings support all of Goode's synonymizing, and I would go still further).

Burbidge, Kirsch, and Main (1974) examined osteological features and used serological tests with elaborate statistical analysis of their serological results. These serological comparisons had the defect that only a small number of individuals (sometimes only one) represented each (supposed) species in the comparison and thus, it is impossible to say what might be the range in genetic diversity within a single deme, much less within a broadly distributed species. However, the results of Burbidge *et al.* are extremely important in giving an independent estimate of relationships between the taxa. All previous authors used a series of morphological characters traditionally used in taxonomic studies of Testudines, but for which the utility in determining biological relationships was largely untested. Burbidge *et al.* found *Elseya* (*sensu* Goode) and *Emydura* closely related to one another and quite different from *Chelodina*. *Pseudemydura umbrina* was well isolated from both the *Elseya-Emydura* group and from *Chelodina* but, to judge from their graphs (their Figs. 8, 9), closer to *Elseya* (*sensu* Goode) than to anything else. Their initial computation of similarity on the basis of serology showed close relationship between all *Emydura* (*sensu* Goode), but the second computation (removing distortions produced by gross differences between groups in the first computation) showed *Elseya dentata* and *E. novaeguineae* close to each other and *E. latisternum* well separated. To judge from their graphs, an *Emydura australis* from Wyndham, Western Australia, and an

Emydura "subglobosa" from Lake Murray, Western Province, Papua New Guinea, came out extremely close, while an *E. krefftii* from Woodstock, Queensland was close but well separated and *E. macquaria* from Patho, Victoria was still farther separated. These results supported the general features of Goode's arrangement, but not all details of his taxonomy: Goode considered the New Guinea *Emydura* ("*E. subglobosa*") as a synonym of *E. krefftii*, but suspected that *E. australis* might not even belong in the genus *Emydura*.

It is my own finding that if certain morphological traits (particularly, form of the horny rhamphotheca, details of base of skull, form of ossified hyoid, coloration of soft parts) are given high weight, and certain other characters (particularly, details of external shell structure) are given low weight, then a morphological grouping emerges that is fully consistent with Burbidge, Kirsch, and Main's serological groupings.

It also would appear from my material that "*Elseya*" *dentata* and "*Elseya novaeguineae*" are so closely related that no character will consistently separate them and, since they are allopatric, there is no evidence that any speciation has occurred and these two named taxa should be considered a single species, *Emydura dentata*. Similarly, there seem to be no constant characters for distinguishing *Emydura australis*, *E. "subglobosa"* and *E. "krefftii"*, and all three names appear to refer to a single species, *Emydura australis*.

THE NEW GUINEA SPECIES OF EMYDURA

Study of the American Museum of Natural History (AMNH) *Emydura* from New Guinea convinces me that two species occur there, each also occurring in Australia:

1) *Emydura dentata* (Gray) (*Elseya dentata* and *Elseya novaeguineae* of

Goode [1967], which see for full references). Waigoe I. and nearly all of New Guinea (at least as far east as Popondetta, Northern Province, Papua New Guinea), but not known from the Huon Peninsula or Milne Bay Province (I cannot find a definite record for the Port Moresby region); and Victoria River, northern Australia to the Cairns and Gayndah regions of Queensland.

2) *Emydura australis* (Gray) (*Emydura kreffti* and *Emydura australis* of Goode [1967], which see for full references). Southern coast of New Guinea from Mimika and Setekwa Rivers (Irian Jaya) to the Aird Hills (Gulf District, Papua New Guinea) and the elevated region around Port Moresby (Central District); tropical and subtropical Queensland, Northern Australia, and northern Western Australia.

These two species, as here expanded, are largely sympatric and clearly distinct; but not all the characters supposedly distinguishing them will work on my material, and some features suggest these two forms are most closely related to each other. The special differences and resemblances need discussion.

Head pattern (see Fig. 1). Both New Guinea species differ from other Australo-New Guinean short-necked chelids in having a pale mark on the upper temporal region (1). (The numbers in parentheses refer to correspondingly numbered characters depicted in the figures.) In *E. australis* this mark is always conspicuous and in sharp contrast to its general surroundings. In *E. dentata* from New Guinea the mark is not very distinct and appears to be only one of several paler mottlings of the head (2), and (*fide* Goode, 1967) in Northern Australian *E. dentata* this mark may merge with a whitening of the entire temporal region. In *E. dentata* this pale temporal mark is continued through the eyelid to join its counterpart as a transverse pale bar or arch (3) between the fronts of the orbits. In *E. australis*, this anterior continuation of the pale temporal mark is either absent

(Western and Northern Australia) or extends forward on the snout to meet its counterpart near the nostrils (Queensland and New Guinea).

Head scutellation (see Fig. 1). As pointed out by Goode (1967), *E. dentata* resembles the Australian *E. latisternum* (and also *P. umbrina*) in having well-defined convex scutes on the temporal region (4), whereas *E. australis* resembles the Australian *E. macquaria* in having the temporal scutes flat and imperfectly defined from one another. If there is any difference between *E. dentata* and *E. australis* in the thickness of the cornification on the crown, the difference is too subtle for me to discern.

Neck tubercles. I am unable to find any constant difference between *E. dentata* and *E. australis* in the development of tubercles on the neck; *E. dentata* from Papua New Guinea north of the central watershed have the neck tubercles less well developed than in the *E. australis* from Central District, Papua New Guinea, shown in Figure 1. (Further, all northern New Guinea *E. dentata* have rounded, rather than pointed, neck tubercles and would key out as "*Elseya dentata*" rather than "*Elseya novae-guineae*" by Goode's [1967] key.)

Penis (see Fig. 2). The terminology of the penis used here is explained below (pp. 180–185). In *E. australis* (Fig. 2A,C) the penis is relatively longer and narrower than in *E. dentata* (Fig. 2B). This is most conspicuous in the proportions of the free tip, which is longer than wide in *E. australis*, but as wide as long in *E. dentata* (5). In *E. australis* there is a row of horny papillae (6) on the anterior plica externa, sometimes on the posterior plica externa, as well, but in *E. dentata* these papillae are absent. (The single Australian male of *E. dentata* examined, BM [18]75.5.1.8, Gayndah, is similar in penial structure to New Guinea specimens; the penis of *E. latisternum* is similar to that of *E. dentata*; I have not seen the penis of *E. macquaria* or *Pseudemydura umbrina*.)

Figure 1. Heads of New Guinea *Emydura*. A. *E. dentata*, AMNH 59927. B. *E. australis*, AMNH 59910. Numbers refer to text discussion.

Figure 2. Penes of *Emydura*. A. *E. australis*, unprotruded, AMNH 69578. B. *E. dentata*, unprotruded, AMNH 103629. C. *E. australis*, protruded, AMNH 57898. Numbers refer to text discussion.

Abbreviations: ape, anterior plica externa; ecl, everted cloacal lining; ft, free tip; pi, plica interna; pm, plica medialis; ppe, posterior plica externa; sc, sinus communis.

Chelid foramen (see Fig. 3). *Emydura dentata* and *E. australis* agree with each other and differ from other Chelidae (except *Chelodina novaeguineae*) in the closure of the foramen (7) just antero-lateral to the carotid foramen. In other Chelidae this foramen is patent and transmits a small vein from the eustachian tube (which has its pharyngeal orifice immediately ventral to the foramen) to the vena capitis lateralis within the cavum epiptericum. Siebenrock (1897) erroneously identified this foramen as

containing the palatine branch of the internal carotid, but the palatine artery of Chelidae branches from the carotid within the carotid canal of the pterygoid, as in other Testudines.

Triturating surfaces and associated skull characters (see Fig. 4). In all *E. dentata* examined, excepting AMNH 66733 (a hatchling from Lae, still with egg-caruncle), the horny sheath of the maxilla has a longitudinal row of tubercles along the center of the triturating surface (8), sometimes accompanied by

Figure 3. Ear regions (not to same scale) of *Emydura* and *Pseudemydura*. A. *E. dentata*, AMNH 62612. B. *E. australis*, AMNH 104011. C. *E. latisternum*, AMNH 27287. D. *E. macquaria*, AMNH 66737. E. *P. umbrina*, WAM R29338. Number (7), chelid foramen (see text discussion).

Figure 4. Skulls of *Emydura dentata* (A, B, C, all AMNH 62612) and *E. australis* (D, from AMNH 57586; E, F, G, from AMNH 104011), showing horny maxillary sheath (A, D, E), palatal view, horny sheath removed (B, F): and medial view of mandible as if cut on symphysis (C, G). Numbers refer to text discussion.

an additional, more medial row of tubercles. In *E. australis* the horny triturating surface may be worn smooth (as in Fig. 4E), but usually has randomly scattered small tubercles (which may, however, suggest two rows in juveniles, as in Fig. 4D). In the largest *E. dentata* skull examined (AMNH 62611, Bernhard Camp, Idenburg River, Irian Jaya; condylobasal length 44 mm) there is a distinct crest on the maxilla underlying the central tubercle row (9). In three Papua New Guinea skulls with condylobasal lengths of 39 to 43 mm (AMNH 92675, Lae; 104007, Abam, Oriomo River; 99613, Ambunti, West Sepik Province); there is a feeble, but discernible, maxillary crest, although smaller skulls (condylobasal length 18 to 28 mm) show no crest. In *E. australis* the crest is absent in even the largest skulls examined (AMNH 104011, Oriomo River, Papua New Guinea; condylobasal length 43 mm; 108957, Darwin area, Northern Australia, 46 mm).

The Australian *E. dentata* examined, BM [18]75.5.1.8 (adult male, spirit specimen), BM[18]76.5.4.7 (juvenile female, spirit), and BM[18]77.5.19.77 (large adult skeleton) were all from Gayndah, southeastern Queensland. The juvenile (externally measured headlength 31.4 mm; carapace length 124.6 mm) has the ridge on the triturating surface represented only by a row of tubercles on the rhamphotheca, without modification of the underlying maxillary bone. The adult skeleton (condylobasal length 44.2 mm), which was the basis for Boulenger's (1889) diagnosis and figure of *Elseya*, shows a distinct ridge on the maxilla, as do New Guinea skulls of the same size.

Although the difference is slight in small juveniles, *E. australis* has broader triturating surfaces than those of *E. dentata*, this being very marked in the largest skulls (AMNH 104011 and 108957, where I can detect no difference between the New Guinea specimen ["*subglobosa*"] and the Darwin area specimen ["*australis*"]). As a consequence, the median posterior emargination of the upper horny jaw sheath is longer than its

width in *E. australis*, but no longer than wide in *E. dentata* (10). Further, the choanae are more posterior in *E. australis* than in *E. dentata*, with the anterior choanal rim opposite the center of the eye in *E. australis* and opposite the anterior margin of the eye in *E. dentata*. Additionally, in *E. dentata*, the crests of the prefrontals separating the olfactory and orbital chambers are visible from below through the choanae (11), but in *E. australis* the triturating surfaces conceal these crests. The vomer is stouter in *E. australis* than it is in *E. dentata*, forming a strong supporting strut for the crushing triturating surface. In *E. australis* the vomer conceals the median contact of the palatines from beneath (12), whereas the palatine contact is exposed ventrally in *E. dentata* (the value of this character is put in question by variability in *E. latisternum*, where AMNH 69576, Pascoe River, Queensland, has the interpalatine suture exposed, but AMNH 26287 [Ravenshoe, Queensland] and 69572 [Iron Range, Queensland] have the contact concealed).

The broadening of the mandibular triturating surface in *E. australis* results in an horizontal platform behind the (rounded) symphyseal hook (13), whereas the triturating surface is confined to the sloping posterior face of the (pointed) symphyseal hook in *E. dentata*.

In the Australian *E. latisternum* and *P. umbrina*, the triturating surfaces are narrow and smooth, with the choanae anterior in position. *E. macquaria* has scattered inconspicuous tubercles on triturating surfaces that are narrower than in *E. australis* but broader than in the other forms.

Hyoid. *Emydura dentata* agrees with *E. australis* (and *E. latisternum*) in having ossification of the body of the hyoid confined to the region from the attachment of the first ceratobranchial to the attachment of the second ceratobranchial. In *E. macquaria* and *P. umbrina* ossification extends farther anteriorly, reaching the rim of the hyoid fenestra.

Shell. I am unable to find any com-

pletely reliable external character of the shell distinguishing adult *Emydura dentata* from *E. australis*, and both species seem quite variable in shell structure. In both, the intergular may meet the pectorals and completely separate the humerals, as in *Pseudemydura* (seen in *E. australis*: AMNH 59052, Mafulu, and 57586, Daru I.; in *E. dentata*: AMNH 66734, 66736, Lae). Both usually have a nuchal (absent in 3 of 45 New Guinea *E. dentata* but absent in all 3 Australian specimens seen), an axillary (absent in 4 of 44 *E. dentata* and 5 of 15 *E. australis*) and an inguinal (absent in 7 of 15 *E. australis*). Contrary to Goode's (1967) characterization of "*kreffti*," all my *E. australis* have the carapace distinctly flared over the hind legs and widest across the eighth marginals except for AMNH 104339 (Manning Creek near Mt. Bernard Homestead, Western Australia), which has the carapace unflared and widest across the seventh marginals. Most *E. australis* show the transparency of the carapacial scutes noted by Goode (1967) for "*kreffti*," but this is even more marked in most New Guinea *E. dentata*. I have been unable to find any constant differences between the *Emydura* available to me in the proportions of the intergular.

Juvenile *E. australis* (AMNH 57586, Daru I., carapace length 90 mm; 57587, same, 109 mm; 92967, Fly R., 108 mm; 104339, Western Australia, 105 mm) agree with adults in having a smooth rim of the carapace. But in *E. dentata*, the edge of the carapace is conspicuously serrated in 17 specimens with a carapace of 58 to 92 mm (except AMNH 99621, W. Sepik Prov., 79 mm), serrated or not in 10 specimens (including a strongly serrated Queensland specimen, 2 strongly serrated New Guinea specimens) with carapace 104 to 126 mm, smooth or only faintly serrated in 17 specimens with carapace 156 to 252 mm (including a nearly smooth Queensland specimen; most nearly "serrate" is AMNH 104837, Western Prov., 184 mm). The three smallest specimens (AMNH 66733, 66735,

99622) have a nearly smooth carapace rim (carapace length 31 to 52 mm).

E. latisternum, supposedly the closest relative of *E. dentata*, is less like New Guinea *E. dentata* than are New Guinea *E. australis* in several shell features: the nuchal is absent (a vestige, excluded from margin, in AMNH 69576, Pascoe River, Queensland); the inguinal is absent (vestige on right in AMNH 69573, Pascoe River); axillary absent except in 3 of 13 specimens (all three, AMNH 69569–71, from Iron Range, Queensland); the carapace is conspicuously serrated at all ages; and the bridge is much narrower than New Guinea *E. australis* and *E. dentata*. With B = width of left plastral bridge and P = maximum length of plastron (measurements to nearest mm), regression equations are as follows:

New Guinea *E. dentata* (N = 43) $\hat{B} = -1.7 + 0.35P$ ($s_{\hat{B}} = 1.6$)

New Guinea *E. australis* (N = 11) $\hat{B} = -5.7 + 0.36P$ ($s_{\hat{B}} = 2.1$)

Queensland *E. latisternum* (N = 13) $\hat{B} = -5.9 + 0.30P$ ($s_{\hat{B}} = 1.8$)

RELATIONSHIP OF *EMYDURA DENTATA* AND *E. AUSTRALIS*

Based on my investigation, *Emydura australis* (including "*kreffti*" and "*subglobosa*") seems the closest relative of *E. dentata* (the type species of "*Elseya*"). Because of this, the generic recognition of *Elseya* seems unwarranted and does not appear to express the apparent phylogenetic relationships of these turtles. I therefore synonymize *Elseya* with *Emydura*.

Burbidge *et al.* (1974: 401, 402) give stereo pair graphs of the relationships of Australo-New Guinean chelids, first on the basis of a correlation matrix of all their serological data, then with data on anti-"*Elseya*" *latisternum* sera omitted. In both graphs the cluster of *E. australis* (including "*E. kreffti*" and "*E. subglobosa*") seems nearest to that for *E. dentata* ("*Elseya dentata*" and "*E.*

novaeguineae"), a result in accord with my morphological findings, provided great weight is given to the pale temporal spot and to occlusion of the chelid (eustachian tube vein) foramen.

The serological findings of Burbidge *et al.* do not appear to support Goode's (1967) division into *Elseya* and *Emydura* at the generic level. However, when anti-"*Elseya*" *latisternum* serum data were omitted, *E. latisternum* fell close to "*Elseya dentata*" and "*E. novaeguineae*," a result in accord with my own observations of penial structure (unfortunately, I do not have data on the penis of *E. macquaria* or *P. umbrina*). That their preliminary analysis showed *E. latisternum* to be quite distant from all other *Emydura* (*sensu lato*) may not be the result of experimental artifact, for it agrees with my morphological findings, that *E. latisternum* differs sharply from *E. dentata* in having an open chelid foramen, in lacking a pale temporal mark, lacking axillary and inguinal scutes and in having smooth horny triturating surfaces at all ages. The resemblances between *E. dentata* and *E. latisternum* stressed by Goode (1967) in his concept of "*Elseya*" may simply be primitive characters retained by these species but lost in the species constituting Goode's *Emydura*. This is supported by the fact that *P. umbrina*, omitting its unique features, would key out as "*Elseya*." Although paleontological evidence is lacking, the large number of unique specializations and the isolated distribution (around Perth) of *P. umbrina* suggest that it separated phylogenetically from other *Emydura*-like terrapins long ago and might be expected to retain characteristics of early, ancestral *Emydura*.

In both graphs given by Burbidge *et al.*, *E. macquaria* seems isolated, but is nearer to *E. australis* than to *E. dentata* (in keeping with the morphology of the triturating surfaces and the flat skin of the temporal region). In the first graph, including data on anti-*E. latisternum* fac-

tors, *E. macquaria* agrees with *E. latisternum* in position on the left-right axis, but widely separated on other axes. My own observations suggest that *E. latisternum* is closely related to *E. macquaria*. Of 13 *E. latisternum* examined, the four smallest (AMNH 69567, 69573-4, 69576; carapace length 97 to 144 mm) show a strongly defined ventrolateral white stripe on the neck, cited by Goode (1967) as a characteristic of *E. macquaria*, while Goode notes that older specimens of the latter have a thickly cornified crown (as in *E. dentata*, *E. latisternum*, and *P. umbrina*). In having a patent chelid foramen and a dark and unicolor temporal region, *E. latisternum* and *E. macquaria* resemble each other (and *P. umbrina*) and differ from *E. dentata* and *E. australis*.

Pseudemydura umbrina agrees with *E. macquaria* in the extensive ossification of the body of the hyoid and in its southern distribution, although *P. umbrina* is southwestern and *E. macquaria* southeastern in distribution. *Pseudemydura* is strikingly characterized by several unique features. These include: extreme flattening of the skull that causes the parietal to rest upon the quadrate (but without an interdigitating suture); contact of the supratemporal ("squamosal" of most authors) with the transversely expanded supraoccipital; and constant enlargement of the intergular to separate the humerals. In spite of these peculiarities, it seems closely related to *Emydura* and I am not certain that full generic (as opposed to subgeneric) rank is necessary for *Pseudemydura*. The graphs of Burbidge *et al.* show *P. umbrina* far separated from all *Emydura*, but the nearest approach is to *E. dentata*, a species showing separation of the humerals by the intergular as a variation (see above) and with narrow triturating surfaces (as in *P. umbrina*). I believe this is best explained by considering *E. dentata* the most primitive species of *Emydura*.

THE BEARING OF *EMYDURA DENTATA* ON CHELID PHYLOGENY

In the classification of Gaffney (1975), the Infraorder Pleurodira contains two families, Pelomedusidae and Chelidae. The former seems the more primitive in retaining mesoplastra in the shell and a squamosal (=quadratojugal of most authors) in the skull. In Pelomedusidae (*Podocnemis unifilis*, *Pelomedusa sub-rufa*, *Pelusios derbianus* examined) the retractor penis muscles arise from the sacral region, as in Cryptodira and probably the primitive condition. However, in Chelidae the retractor penis is reduced and arises from the transverse ventral musculature anterior to the cloaca (*Emydura dentata*, *E. australis*, *E. latisternum*, *Phrynops gibbus*, *P. geoffroanus*, *Platemys platycephala*, *Chelus fimbriatus*, *Chelodina novaeguineae*, *C. expansa* [= *C. "rugosa"*] examined). On the other hand, in having nasal bones (except in *Chelus*) and in usually having a suturally distinct splenial, the Chelidae seem more primitive than the Pelomedusidae. (The splenial is variable within New Guinea *Emydura dentata* and was absent in three skulls examined [AMNH 92965 and 92971, Fly R., and 104007, Oriomo R.] but present in five others.)

The genera *Emydura* and *Pseudemydura* are apparently more primitive than other Chelidae, more like the Pelomedusidae in several features: 1) the dentaries are fused at the symphysis; 2) the supraoccipital forms the dorsal rim of the foramen magnum and is the most backwardly projecting bone of the skull; 3) the anteriormost part of the temporal fossa is roofed over by a transverse shell of parietal from the ventral surface of which short and vertical jaw adductor fibres arise (seen also in *Platemys*; in other Chelidae the jaw adductor is formed entirely of long and stretchable fibres, allowing a wide gape of the mouth at the expense of lessened crushing force of the jaws); 4) there is a scar on the

pharyngeal surface of the pterygoid bone for the ventral pterygoideus muscle (this muscle seems to be absent in other Chelidae except *Platemys*); 5) the flange of the quadrate suspending the eustachian tube is extended posteromedial to the tube to meet the cartilaginous border of the fenestra ovalis (not extended beyond the eustachian tube in other Chelidae; *Pelomedusa* and *Pelusios* are like *Emydura*, but in *Podocnemis* and its close relatives, this flange is still better developed as a firm brace against the basisphenoid and basioccipital); and 6) the cervical vertebrae have normal zygapophysial articulations, oriented horizontally (cylindroid or vertically oriented in other Chelidae).

Among *Emydura*, *E. dentata* shows special similarity to *Podocnemis* and its close relatives in having a longitudinal ridge on the maxillary triturating surface, present as a tuberculate keel on the horny sheath in all but hatchlings, and with bony ridging of the maxilla in large adults. *E. dentata* seems unique among Chelidae in its ridged triturating surface and, as the most *Podocnemis*-like of Chelidae, seems the most primitive species of the family structurally, except in lacking neural bones (true of all *Emydura*). As in the other *Emydura* examined (*E. australis*, *E. latisternum*), the cloacal bursa of *E. dentata* is formed as in *Podocnemis*: a bilobate sac with a single median dorsal aperture into the cloaca and with a lining of coarse papillae. The simplest explanation for special resemblances between *Podocnemis* and *E. dentata* is that these are primitive features for Pleurodira that have been lost or modified in other forms.

For geographical reasons the Australo-New Guinean genus *Chelodina* might be expected to show special relationship to *Emydura* and *Pseudemydura*, but no morphological evidence corroborates this. Rather, as shown convincingly by Gaffney (1977), *Emydura* and *Chelodina* seem at opposite extremes, with the

South American Chelidae serving as intermediates. *Chelodina* (*C. longicollis*, *C. novaeguineae*, *C. expansa* [*C. "rugosa"*], *C. steindachneri* examined) has all of the specializations that set South American chelids apart from *Emydura* and *Pseudemydura* (persistent suture between dentaries, supraoccipital excluded from foramen magnum and exceeded by exoccipitals in backward extent, vertical cervical zygapophysial articulations, etc.). In addition, *Chelodina* adds some peculiarities of its own: 1) fusion of the frontals (perhaps unique in Testudines); 2) only four claws on hand (seen also in the South American *Hydromedusa*); 3) greatly elongated cervical vertebrae (as in the South American *Chelus* and *Hydromedusa*); 4) total reduction of roofing of the temporal fossa (approached by some South American forms, e.g., *Hydromedusa*, and the exact opposite of *Emydura* and *Pseudemydura*); and 5) cloacal bursae paired, completely separated, and with smooth lining (most nearly approached by *Chelus*, where the lining is smooth and the bursae fused only at the point of entry into the cloaca). The only special resemblance between *Emydura* and *Chelodina*, absence of neural bones, is also seen in some South American forms (e.g., *Platemys*) and is probably convergent, since Burbidge *et al.* (1974) report that *Chelodina oblonga* has five to eight neural bones. Two possible interpretations of these morphological resemblances and differences are:

1) *Chelodina* arose in Australia from *Emydura* through a series of intermediates that have become extinct. However, one of these intermediates invaded South America to give rise to the South American Chelidae. Any special resemblances between some South American forms and *Chelodina* are fortuitous.

2) Some primitive *Emydura*-like form invaded South America from Australia, and gave rise to the South American chelid radiation, including a long-necked line (*Hydromedusa* and *Chelus*). A primi-

tive member of the long-necked line reinvaded Australia and gave rise to *Chelodina*. Special resemblances between *Chelodina* and certain South American Chelidae are the result of close common ancestry.

The latter alternative seems now to have greater evidence, but the first alternative cannot yet be excluded. Real fossils, rather than hypothetical common ancestors, will have to be brought into the discussion before the decision can be considered a realistic examination of nature rather than a statement of the rules of irreversibility and minimum coincidence that we impose (as a necessary control) on evolutionary hypotheses.

THE MORPHOLOGY OF THE PENIS OF PLEURODIRA

Although Zug (1966) has described the penis of cryptodires and shown the usefulness of penial morphology in the classification of that group, there appears to be no account of the morphology of the penis of Pleurodira, aside from Schmidtgen's (1907) account of the unprotruded organ of *Chelodina longicollis*. (Goode [1967] has figured what I would consider the partial protrusion of the penis of *Emydura australis*.) As noted in the previous section of this paper, the penis appears to offer very useful taxonomic characters within the pleurodires. Further, the morphology of the pleurodiran penis is quite different (but *not* fundamentally) from that of cryptodires and suggests how the protrusile penis of cryptodires might have arisen from an ornamented cloacal evert, such as that of living gymnophione Lissamphibia (and, presumably, the earliest reptiles).

As in Cryptodira, the most ventral portion of the penis of Pleurodira is formed by the corpus fibrosum. The corpus fibrosum of pleurodires differs from that of cryptodires in not being cavernous or erectile in texture; rather, it is dense and cartilagelike in consistency (but not hy-

aline cartilage). For most or all of its length, the corpus fibrosum of pleurodires is attached to the floor of the cloaca. This attachment is complete in *Podocnemis*, *Chelodina*, and *Phrynos geoffroanus*. The extreme posterior end of the corpus fibrosum projects as a short free tip (as in *Emydura dentata* and *E. latisternum*, as described in the previous section) in *Pelusios*, *Chelus*, and *Phrynos gibbus*. There is a longer free posterior tip (as in *Emydura australis*, as described in the previous section) in *Pelomedusa*. In *Platemys* the free tip of the corpus fibrosum is much longer than in the other forms examined, but still does not include the portion of the organ bearing the seminal groove.

The very spongy and obviously erectile tissue of the corpus cavernosum is only sparsely distributed over the corpus fibrosum of pleurodires, as compared with cryptodires, a point noted by Schmidtgen (1907: 391) for *Chelodina longicollis*. However, the corpora cavernosa are continued forward into the body cavity as a pair of very large proximal bulbs of the cavernosum that lie anterior (proximal) to the corpus fibrosum. The paired and narrow peritoneal canals (essentially similar to those of cryptodires) pass dorsomedial to the bulbs of the cavernosum to lie close to, and parallel with, the seminal groove.

The seminal groove is simple in all the pleurodires examined and runs posteriorly from the urodaeal chamber of the cloaca along the dorsal surface of the penis to about the middle of the corpus fibrosum. The seminal groove lies in a longitudinal trough of cavernosum tissue, but this is not a conspicuous mass of cavernous tissue (there seems to be just enough to form lips defining the seminal groove). I feel reasonably confident that this trough of erectile tissue defining the seminal groove is homologous to the plica interna of Cryptodira, in the terminology of Zug (1966). I am also confident of the homology of the corpus fibrosum, the corpus cavernosum (as an

entity—not its parts), the seminal groove, and the peritoneal canals of pleurodires with the correspondingly named parts of the cryptodiran penis. Beyond these, I have very little confidence in the homologies of pleurodiran to cryptodiran penial structures, because of the very different appearance of the organ in the two groups (particularly when protruded), and my use of Zug's terminology must be considered very tentative.

The distal (posterior) end of the seminal groove opens (in the unprotruded organ) into a U-shaped basin that lies distal as well as lateral to the seminal groove and plica interna. Zug (1966) finds three separate and paired pits or sinuses in cryptodires, but I find only this U-shaped depression in pleurodires, although this depression may show pocketlike extensions that would seem to correspond to the clearly defined sinuses of cryptodires. I term this single and U-shaped depression of pleurodires the *sinus communis*. The main portion of cavernous tissue of the pleurodiran penis lies on the outer rim of the sinus communis; thus the sinus communis isolates the plica interna (and seminal groove) from the remainder of the corpus cavernosum. The anterior (i.e., proximal) ends of the sinus communis lie near the posterior (i.e., distal) ends of the paired peritoneal canals and it is tempting (but without any supporting developmental evidence) to regard the sinus communis as a depression resulting from the collapse of an underlying posterior continuation (and median fusion, distal to the tip of the seminal groove) of the peritoneal canals.

Schmidtgen (1907) found the peritoneal canal to end blindly in all Testudines he studied (both male and female), but the point of approximation of the tip of the peritoneal canal to the surface of the penis to be marked by a pale spot in *Trionyx*. I also have failed to find any patent opening of peritoneal canal, but found a yellow spot in *Podocnemis*. Thus, present evidence (including all of my own observations) favors Schmidt-

gen's belief that the peritoneal canal ends blindly and that reports of an open pore are based on artifacts, but this question must still be considered open. It is possible that a patent pore may be of seasonal occurrence, or perhaps only develops during copulation. A pore might be confined to females (with a clitoris very similar to the penis of males, but much smaller), since my own and Schmidtgen's dissections and histological sections have been carried out on males (this appears to be the case with Zug's dissections, as well). If the peritoneal canal is ever open to the outside, then the sinus communis might serve as the receiving chamber for whatever fluid might issue from this pore. It should be remembered that the precise stimulus to ejaculation by the copulating male is unknown for any turtle.

The cavernous tissue forming the outer rim of the sinus communis is differentiated at its middle—that is, at the level of the distal end of the seminal groove and plica interna—into a distinct mound, usually (*Platemys* is the exception) bearing a horn-like prominence. Tentatively, I identify this mound as the homologue in Pleurodira of the plica medialis (Zug's terminology) of testudinoids and term the hornlike prominence the *cornu of the plica medialis*. In the Pelomedusidae examined, the corpus fibrosum has a distinct lateral lobe supporting the plica medialis and its cornu and the cornu is outwardly directed in the unprotruded organ. The cornu of *Podocnemis* appears to be bifid, with a longer and more acute anterior lobe exactly opposite the tip of

the seminal groove, and with a blunter posterior lobe on the lateral border of the corpus fibrosum distinctly posterior (distal) to the level of the tip of the seminal groove (but this posterior lobe perhaps belongs to the *posterior plica externa*, to be discussed below). In *Pelomedusa* there is a single cornu of the plica medialis, supported by a hornlike process of the corpus fibrosum and extending anterolaterally in the unprotruded organ, but an oblique fleshy fold extends postero-medially from the apex of the cornu to end blindly in the sinus communis, well posterior (distal) to the tip of the seminal groove. This oblique ridge may be the homologue of the posterior lobe of the cornu of the plica medialis of *Podocnemis*; the sinus communis forms a deep, posteriorly directed pocket between the posterior end of this oblique ridge and the posterior plica externa (this pocket perhaps represents the "posterior sinus of the glans" of testudinoids, in Zug's terminology). In *Pelusios* the plica medialis is a longitudinally ovoid swelling lateral to the distal end of the seminal groove and plica interna, with a blunt free posterior end that probably represents the cornu.

In Chelidae, the corpus fibrosum does not have any supporting lobe for the plica medialis or its cornu. In *Platemys*, the plica medialis is simply a low, longitudinal ridge without a projecting cornu. In the others examined, the cornu is a quite distinct free process, but it lies (in the unprotruded organ) in the sinus communis, with its free end directed medially (rather than laterally as seen in

Figure 5. Penis of: A. *Podocnemis unifilis* (AMNH 87966), unprotruded. B. *Pelomedusa subrufa* (AMNH 50744), unprotruded. C. *Pelusios derbianus* (AMNH 50746), unprotruded. D. *Chelodina expansa* ("rugosa") (AMNH 86548), unprotruded. E. *Phrynops gibbus* (AMNH 61521), protruded. F. *Platemys platycephala* (AMNH 61529), protruded.

Abbreviations as in Fig. 2, as well as: alc, anterior lobe of cornu of plica medialis; c, tip of cornu of plica medialis; plc, posterior lobe of cornu of plica medialis; psg, posterior sinus of glans; ssg, sinus of seminal groove; ys, yellow spot marking termination of coelomic canal. The plica interna is present in F, but concealed by the plicae mediales and everted cloacal lining.

Pelomedusa and *Podocnemis* or dorso-laterally in *Pelusios*). However, in the naturally protruded organs of *Phrynos gibus* and *Emydura australis*, the cornu is rotated to extend laterally, and this is probably true of the other Chelidae, as well (manipulation of preserved chelid penes, by bending them at the level of the tip of the seminal groove, as in the naturally protruded specimens available, produces the lateral rotation of the cornu). Probably the cornu of the plica medialis functions as a holdfast to prevent dislodgement of the penis before the completion of copulation.

In most pleurodires, the apparent homologue of the cryptodiran plica externa is divided by the plica medialis into two separate portions: 1) the plica externa anterior, proximal to the plica medialis and with its posterior (distal) end joined to the anterior end of the plica medialis; and 2) the plica externa posterior, lying on the lateral edges of the most posterior (distal) part of the corpus fibrosum, with its anterior (proximal) end joined to the plica medialis. All Chelidae examined and *Pelomedusa* show this linear order of corpus cavernosum lobes on the lateral rim of the sinus communis, arranged (from anterior to posterior); plica externa anterior, plica medialis, plica externa posterior. In *Pelusios*, however, the plica externa anterior and plica externa posterior join lateral to the plica medialis. I am uncertain about the correct interpretation of *Podocnemis*, where the "posterior lobe of the cornu of the plica medialis" is joined basally to the plica externa anterior, lateral to the "anterior lobe of the cornu of the plica medialis." If this "posterior lobe of the cornu" is really a lateral wing of the proximal part of the plica externa posterior, then *Podocnemis* would be like *Pelusios* and cryptodires in having the two portions of the plica externa continuous lateral to the plica medialis.

In all pleurodires examined except *Platemys*, the proximal end of the plica externa posterior has a blunt lobe that

lies in the more distal (posterior) part of the sinus communis of the unprotruded organ. In at least *Phrynos gibus* and *Emydura* (probably in the others, as well), this lobe swings to a ventrally directed vertical position in the protruded organ. The lobe probably functions, along with the cornu of the plica medialis, as a holdfast during copulation (in *Phrynos gibus* and *P. geoffroanus* there are coarse and hard denticulations on this lobe of plica externa posterior, further suggesting that it acts as a holdfast).

In all the pleurodires examined except the juvenile *Chelus* (the adult agreed with the majority), a paired anterior extension of the sinus communis extends proximally between the plica interna and the plica externa anterior. This would seem to represent the "sinus of the seminal groove" of cryptodires, using Zug's terminology. In *Podocnemis*, there is a shallower and much less clearly defined depression (continuous with the sinus communis) between the plica medialis and plica externa anterior. This perhaps represents the "anterior sinus of the glans" of Zug's terminology for cryptodires.

Zug's terminology was intended for the description of only a small terminal part of the cryptodiran penis, the "glans." Much of the length of the cryptodiran penis, quite apart from the proximal bulbs of the corpora cavernosa, is formed by a shaft region in which the only structure formed by the corpus cavernosum is a paired seminal ridge (Zug's term), flanking the seminal groove. In Pleurodira, the penis has no shaft and the entire structure, apart from the bulbs of the corpora cavernosa, seems to represent the "glans" of the cryptodiran organ (in this respect it resembles the clitoris of cryptodires).

I have seen the naturally protruded penis of only *Emydura australis*, *Phrynos gibus*, and *Platemys platycephala* among pleurodires, but some morphological points lead me to believe

that the organ must function in a similar manner in the other Pleurodira. In the naturally protruded pleurodiran penis seen, the organ is doubled over, so that the tip of the seminal groove and plica interna lie at the tip of the organ, flanked by the plicae mediales, with the plica externa posterior now inverted to lie on the ventral side. The tip of the corpus fibrosum, directed posteriorly in the unprotruded state, is now pointed forward because of the doubling-over of the organ that has taken place at the level of the plicae mediales (slightly behind the level of the tip of the seminal groove of the unprotruded organ, with some enlargement of the plicae internae that bear the tip of the seminal groove accounting for the extreme terminal position of the tip of the seminal groove in the fully protruded organ). That the seminal groove extends only about halfway along the penis in all pleurodires examined can easily be accounted for if doubling-over of the penis is characteristic of pleurodires generally.

The protruded penis is only a part of the copulatory organ of pleurodires. At least to judge from *Emydura*, *Phrynops*, and *Platemys*, the entire copulatory organ of pleurodires is an eversion of the cloacal lining, to which the penis is entirely, or almost entirely, attached. The dorsal cloacal lining is drawn out with the penis and forms a fold over the seminal groove that is a nearly complete roof for the groove. Thus, although the penis of Cryptodira is extruded (Zug's term) to form the copulatory organ as in crocodilians, the copulatory organ of Pleurodira is a cloacal evert as in Gymnophiona, with the penis forming only a specialized part of this evert.

The mechanism of penial protrusion in Pleurodira. So far as I can determine from the study of dead material, the penis of pleurodires acts as a lever to evert the cloaca; protrusion of the penis must necessarily produce eversion of the cloaca because the penis is attached to the cloacal wall for almost its entire length. The mechanism for protrusion of

the penis, in turn, would seem to be warping. The penis is formed of two layers that are very tightly bound together: a ventral layer, the corpus fibrosum, that is flexible but probably unexpandable because it is densely fibrous without blood spaces; and a dorsal layer of erectile tissue, the corpus cavernosum, that has large blood spaces and can increase in size if the rate of blood flow into it exceeds the rate of outward blood flow. When the corpus cavernosum expands and the corpus fibrosum does not, the penis must bend like a heated thermocouple, with the more elongate corpus cavernosum on the outside of the curvature and the shorter corpus fibrosum on the inside of the curve.

Because the blood spaces within the corpus cavernosum are connected with one another, they would make up collectively a fluid chamber subject to Pascal's law, with pressure exerted equally in all directions. Therefore, the pressure at the proximal bulbs of the corpus cavernosum would be equal to that at the flexure-region of the penis, and pressure on these bulbs from the surrounding coelomic fluid could increase the difference in length between corpus cavernosum and corpus fibrosum (thus increasing flexure of the entire organ), as well as pressing the entire penis—and cloacal lining bound to the penis—outward. I do not know of any special mechanism for increasing coelomic pressure, although contraction of the transverse musculature in the region of the cloaca could do so.

The cryptodiran penis is freed at its posterior end from the cloacal wall, so that the cloaca is not everted when the penis protrudes, and penial protrusion appears to depend mainly or entirely on the enlargement of the organ through distension of cavernous spaces by blood. The difference from pleurodires is not quite absolute, since in a number of cryptodires preserved with the penis protruded, I have found a small eversion of the cloacal lining at the base of the organ, although not enough tissue is everted to

roof the seminal groove and there is a vast difference in degree of cloacal eversion from that observed in pleurodires; examples of cryptodires with a small eversion of the cloacal lining at the base of the protruded penis are: *Trionyx ater* (AMNH 88879); *Platysternon megacephalum* (AMNH 30123); *Clemmys guttata* (AMNH 7453). Because the corpus fibrosum is not as expansible as the corpus cavernosum in cryptodires, a number of protruded cryptodiran penes examined show a warping that causes the distal tip to be bent downward (but not the doubling over of the organ observed in pleurodires); examples are: *Clemmys guttata* (AMNH 7453), *Ocadia sinensis* (AMNH 30190), *Kinosternon hirtipes* (AMNH 71363). In *Trionyx*, the four terminal branches of the seminal groove (characteristic of the family) are contained in four corresponding lobes of the corpus fibrosum, with corpus cavernosum tissue firmly attached to the ventral side of each lobe; accordingly, the warping produced by the differential expansion of the two tissues causes the sulcus-bearing lobes to arch upward at their tips (this gives some support to the mechanism here suggested).

I have not seen a protruded chelonioid penis and the only chelonioid penis examined (*Chelonia mydas*, AMNH 107114) suggests that the mechanism may be somewhat different from that in other Cryptodira. The free tip of the organ, formed from the corpus cavernosum and bearing the seminal groove, lies above (but free of) a distal (posterior) projection of corpus fibrosum that is entirely attached to the cloacal lining. I am unable to guess whether: a) penial protrusion is entirely the result of enlargement of the corpus cavernosum, with the corpus fibrosum acting as an immobile base; or b) full protrusion involves an external (backward) displacement of the corpus fibrosum, which, because of its total attachment to the cloacal lining, would drag the cloacal lining with it and produce an extensive cloacal eversion, perhaps approaching that of pleurodires.

The phylogenetic position of the pleurodiran penis. A number of morphologists, but most unequivocally Tonutti (1932), have presented arguments that the most primitive copulatory organ among tetrapods is the eversible cloaca of male caecilians (=Gymnophiona). The cleidoic egg, usually considered the distinctive specialization of the earliest reptiles, implies the existence of internal fertilization, and thus makes it likely some sort of copulatory organ was present. The only living Rhynchocephalian, *Sphenodon*, has no copulatory organ and is alleged to evert the cloaca (but I have been unable to find a circumstantial account of copulation). Most birds lack a copulatory organ, but some (Struthionidae, Rheidae*, Casuariidae* including *Dromaeus*, Apterygidae, Tinamidae, Cracidae*, Anatidae*, Otididae, Burhinidae, Threskiornithidae, and Phoenicopteridae) have a solid protrusible penis on the floor of the cloaca (this penis, in turn, contains an eversible sac, the *Blindslauch*, in the families marked with an asterisk [*] above). Probably the absence of a penis in most birds is a secondary reduction. Crocodilia, cryptodiran turtles and mammals also have a solid and protrusible penis formed from the cloacal floor. Squamata, on the other hand, have a paired eversible sac (the hemipenis) formed from the posterolateral cloacal wall. A considerable part of the attractiveness of considering the caecilian cloacal evert as the primitive tetrapod copulatory organ stems from the fact that by specializations that are easily imagined, the simple cloacal evert can give rise to the different kinds of copulatory organs known in amniotes, but it is difficult to imagine how the squamatan hemipenis could either arise from or give rise to the median protrusible penis of crocodilians, cryptodires, or mammals.

The copulatory organ of at least *Emydura*, *Phrynops*, and *Platemys* (and probably all Pleurodira) is a cloacal evert, essentially like that of caecilians but with the simple holdfast welts and tubercles of

the caecilian evert replaced by a unified ventral structure, the penis, containing a differentiated corpus fibrosum, paired corpus cavernosum, and paired coelomic diverticula. It is relatively easy to derive the cryptodiran penis from that of pleurodires, and thus from a cloacal evert; and so, the pleurodiran penis gives some support to the general theorem of comparative anatomists that the primitive copulatory organ of tetrapods is a cloacal evert. If this phylogenetic interpretation is accepted, then the resemblance in general penial structure between crocodilians and cryptodires must be considered convergent, since this interpretation would mean the protrusible (as opposed to eversible) organ of cryptodires arose within the Testudines, after the many specializations (e.g., shell, middle ear, lack of lachrymal duct) that rule out Testudines as ancestors of any archosaurian group. By this interpretation, conversion of a cloacal evert to a protrusible organ took place at least twice, and quite possibly more often than that.

This interpretation is not bothersome to me, but it should be remembered that there is no proof the phylogeny did not go the other way, from a protrusible cryptodire-like penis to a secondarily developed cloacal evert in Pleurodira. It is very unlikely that any documentation from the fossil record will ever indicate the direction of evolution of the penis in Testudines. This is the fundamental difficulty in the use of soft tissue anatomy in higher classification. Though the characters of the reproductive system, circulation, and pathways in the central nervous system may have been much more important to the organisms than the details of skeletal structure, and may have been the most important factors in determining the success or failure of evolutionary lineages, they are rarely testable against the fossil record. This much does seem to fit with what can be tested (at least for general plausibility): 1) pleurodires and cryptodires diverged very early in the history of Testudines and this divergence is associated among living forms with

marked differences in the copulatory organ (that is, structure of the copulatory organ proves its usefulness as a taxonomic character in turtles by indicating a difference between two groups that are known from other evidence to be quite different); 2) just as Zug's (1966) survey of the penis of cryptodires divides that group concordantly with Gaffney's (1975) division of the cryptodires on skull anatomy, there appear to be penial differences between the two (osteologically) defined families of living pleurodires; and 3) species can be sorted out—at least in some cases, such as the New Guinea species of *Emydura*—by differences in penial morphology. It seems a reasonable expectation, then, that penial anatomy will be as useful to the taxonomy of Testudines, at all levels, as has been the anatomy of the hemipenis to squamatan taxonomy.

ACKNOWLEDGMENTS

I am most grateful to E. S. Gaffney (Vertebrate Paleontology) and R. G. Zweifel (Herpetology) for access to material in their care at the American Museum of Natural History (AMNH), including certain specimens on loan to them, from the Western Australian Museum (WAM) and the Institut Royale des Sciences Naturelles, Bruxelles (IRSN). I am further indebted to A. Grandison of the British Museum (Natural History) (BM) for use of material in her care.

APPENDIX: SPECIMENS EXAMINED

EMYDURA AND *PSEUDEMYDURA* EXAMINED

Emydura dentata. IRIAN JAYA: Kurik, near Merauke: IRSN unnumb., 3; Bernard Camp: AMNH 62611-13: "Dutch New Guinea": AMNH 62043. PAPUA NEW GUINEA: (Western Province): Fly River: AMNH 92964-66, 92968-74; above d'Albertis Junction: AMNH 59927; Abam, Oriomo R.: AMNH 104007-08; Zim, Oriomo R.: AMNH 104009; Wipim: AMNH 104837-38; (West Sepik Province): Ambunti: AMNH 99612-14,

99618-20; Wagu: AMNH 99615-16, 99621-22; Hunstein R.: AMNH 99617; Lumi: AMNH 100002-04; (Madang Province): near Alexishafen: Peabody Mus. Field no. 0103; (Morobe Province): Oomsis Creek: AMNH 92676; Lae: AMNH 66733-36, 92675, 103629; Wau: AMNH 103630. QUEENSLAND: Gayndah: BM [18]75.5.1.8, [18]76.5.4.7, [18]77.5.19.77.

Emydura australis. PAPUA NEW GUINEA: (Western Province): Daru: AMNH 57586-88; Fly River: AMNH 92967; Zim, Oriomo R.: AMNH 104839; Oriomo, Oriomo R.: AMNH 104011; (Central Province): Mafulu: AMNH 59050-52, 59910; Baroko: AMNH 57898. *Australia*: (Western Australia): Manning Creek: AMNH 104339; (Northern Australia): Mainoru Station: AMNH 87878; Darwin area: AMNH 108957; (Queensland): Wenlock: AMNH 69578.

Emydura latisternum. AUSTRALIA (Queensland): Larradeenya Creek: AMNH 69567; Brown's Creek, Pascoe R.: AMNH 69573-77; Iron Range: AMNH 69569-72; Speewah: AMNH 69566; near Ravenshoe: AMNH 27287-88.

Emydura macquaria. AUSTRALIA: (Queensland): Brisbane: AMNH 103701; (Victoria): Patho, 20 mi. W of Echuca: AMNH 108961; Nancy R.: AMNH 103702; 40 mi. SE of Mildura: AMNH 110488; (South Australia): Cooper's Creek, near Innamincka: AMNH 110486-87. No data: AMNH 77637, 77648.

Pseudemydura umbrina. AUSTRALIA (Western Australia): Ellenbrook Reserve: WAM R29338; Warbrook (Twin Swamp) Reserve: WAM R29341; Bullsbrook Reserve: WAM R21859.

Osteological Material. (*=hyoid examined as well as skull; **=complete skeleton; CBL=condylobasal length in mm.)

Emydura dentata. AMNH 99618 (CBL 18.3); AMNH 92971 (CBL 20.0); *AMNH 92965 (CBL 25.6); IRSN [McD tag 46] (CBL 28.0); **AMNH 92675 (CBL 39.0); AMNH 104007 (CBL 40.8); *AMNH 99613 (CBL 43.0); AMNH 62612 (CBL 44.0); **BM(NH) [18]77.5.19.77 (CBL 44.2); also hyoid only on AMNH 100003.

Emydura australis. AMNH 57586 (CBL 19.8); AMNH 59090 (CBL 31.7); AMNH 104011 (CBL 42.7); **AMNH 108957 (CBL 45.6); also hyoid only on AMNH 92967.

Emydura latisternum. AMNH 69576 (CBL 35.4); *AMNH 27287 (CBL 56.6); also hyoid only on AMNH 69567 and 69566.

Emydura macquaria. **AMNH 103701 (CBL 38.5 [rearticulated]); **AMNH 103702 (CBL 45.6); **AMNH 77636 (CBL 46.3); also hyoid only on AMNH 108961 and 110487.

Pseudemydura umbrina. WAM R29338; *WAM R29341. ([Disarticulated], but smaller than any *E. macquaria* or *E. latisternum* examined and smaller than most *E. dentata* and *E. australis* examined); WAM R21859.

PLEURODIRAN PENES EXAMINED

Pelomedusidae: *Podocnemis unifilis* (AMNH 87966; Brazil: Mato Grosso: confluence of R. Araguaia and R. Tapirapa). *Pelomedusa subrufa* (AMNH 50744; Tanzania: Chai Camp). *Pelusios derbianus* (AMNH 50746; Angola: Huambo).

Chelidae: *Chelodina novaeguineae* (AMNH 57589; Papua New Guinea: Western Province: Mabaduane. [No no., Bruxelles IRSN]; Indonesia: Irian Jaya: Kurik.). *C. expansa* ["rugosa"] (AMNH 86548, 86549; Queensland: Normanton, lower Norman R.). *Chelus fimbriatus* (AMNH 107378 [adult] and 76190 [juvenile]; no data). *Phrynops geoffroanus* (AMNH 90674; Peru: Cuzco, R. Apurimac). *P. gibbus* (AMNH 64721; Guyana: Kartabo [organ unprotruded]. AMNH 60521; Guyana: Parabam [naturally protruded organ]). *Platemys platycephala* (AMNH 61529; Guyana: Shudikarwan R. [naturally protruded organ]). Also *Emydura australis* (AMNH 57898, 59090-1, 57588, 104339, 69578). *E. dentata* (AMNH 103629, 104008, 104837, BM [18]75.5.1.8). *E. latisternum* (AMNH 69566, 69576).

LITERATURE CITED

- BOULENGER, G. A. 1889. Catalogue of the chelonians, rhynchocephalians, and crocodiles in the British Museum (Natural History). New Edition. Trustees, British Museum, x + 311 pp., 6 pls.
- BURBIDGE, A. A., J. A. W. KIRSCH, AND W. R. MAIN. 1974. Relationships within the Chelidae (Testudines: Pleurodira) of Australia and New Guinea. *Copeia*, 1974: 392-409.
- GAFFNEY, E. S. 1975. A phylogeny and classification of the higher categories of turtles. *Bull. Am. Mus. nat. Hist.*, 155(3): 387-436.
- . 1977. The side-necked turtle Family Chelidae: a theory of relationships using shared derived characters. *Amer. Mus. Novitates*, No. 2620, pp. 1-28.
- GOODE, J. 1967. Freshwater tortoises of Australia and New Guinea (in the family Chelidae). Lansdowne Press, x + 154 pp.
- ROOIJ, N. DE. 1915. The reptiles of the Indo-Australian Archipelago. I. Lacertilia, Chelonina, Emydosauria. E. J. Brill, xiv + 384 pp.
- SCHMIDTGEN, O. 1907. Die Cloake und ihre Organe bei den Schildkröten. *Zool. Jahrb. (Abt. f. Anat.)*, 24: 354-414, Taf. 32-33.
- SIEBENROCK, F. 1897. Das Kopskelet der Schildkröten. *Sitzungsber. Akad. Wiss. Wien (math.-naturw. cl.)*, 106: 1-84, taf. i-vi.
- TONUTTI, E. 1932. Vergleichend-morphologische Studie über die Phylogenie des Enddarmes und des Kopulationsorganes der männlichen

Amnioten, ausgehend von den Gymnophion-
nen. Morphol. Jahrb. (Jahrb. Morphol. und
Mikr. Anat. 1st Abt.), 70: 101-130, Taf. 1.

ZUG, G. R. 1966. The penial morphology and the
relationships of cryptodiran turtles. Occ. Pap.
Mus. Zool. Univ. Mich. No. 647, pp. 1-24.

The Basicranial Articulation of the Triassic Turtle, *Proganochelys*

EUGENE S. GAFFNEY¹

ABSTRACT. The Triassic turtle *Proganochelys quenstedti* (Keuper, Trossingen, West Germany) has a movable basicranial articulation formed by the basicranial tubercle of the basisphenoid medially and the basicranial recess of the pterygoid and/or epipterygoid laterally. This is interpreted as the generalized amniote condition, while the akinesis of all remaining turtles (the Casichelydia) is interpreted as a shared derived character testing their monophyly.

INTRODUCTION

Ernest E. Williams performed a singular service for turtle fanciers when he revealed that the skull morphology of the Triassic turtle *Proganochelys* "... approaches the primitive reptilian condition and is separated by a very sizeable morphological gap from such modernized turtles as the Jurassic ones we have described" (Parsons and Williams, 1961: 91). Previous to Williams's revelation the only information about Triassic turtle skulls came from Jaekel's (1916) description of a partially crushed skull. Jaekel understated the degree of damage to his specimen and produced a palatal restoration showing an apparently akinetic skull with a broad, median tooth plate. Williams examined a specimen in Stuttgart that was well preserved and clearly showed significant differences from Jaekel's descriptions. Williams recognized that these differences gave a new

basis for interpreting the Triassic turtles and argued that, except for the shell, *Proganochelys* was a relatively generalized reptile.

The Stuttgart specimens are now the focus of a renewed effort to study the Triassic turtles, and it is singularly appropriate to pay homage to Ernest E. Williams's many contributions to chelonian systematics by dedicating this paper to him. Although the major descriptive work on *Proganochelys* is still in preparation, I thought it would be useful to describe one particular morphologic feature of the skull that has figured in discussions concerning the oldest turtles.

AVAILABLE MATERIAL AND PREVIOUS WORK

To the best of my knowledge, there are three Triassic turtle skulls known at present. All three are, in my opinion, the same taxon, *Proganochelys quenstedti*, and all are middle Keuper in age. In order of discovery, they are as follows:

1) Museum für Naturkunde, Berlin. Collected in 1912, Halberstadt, East Germany. Consists of skull, lower jaws, cervical vertebrae, shoulder girdle, carapace, and plastron. Type of *Stegochelys (Triasochelys) dux* (Jaekel), described and figured in Jaekel (1916), skull also figured in Gaffney (1979b: Figs. 117–124).

2) Staatliches Museum für Naturkunde in Stuttgart, 15759. Collected in 1927, Neuhaus near Aichheim, West Germany. Consists of skull, lower jaws, forefoot, miscellaneous fragments of limbs and cervicals, and a partial carapace. Skull figured in Berckhemer (1931: 7, 1951: 36) and Kuhn (1956: 69).

¹American Museum of Natural History, New York, New York 10024, U.S.A.

3) Staatliches Museum für Naturkunde in Stuttgart, 16980. Collected in 1932, Trossingen, West Germany. Consists of a nearly complete skeleton. Skull along with shell and part of the skeleton figured in Berckhemer (1938: Pl. 4), skull also figured in Parsons and Williams (1961: Pls. 5, 6), and in Gaffney (1979b: Figs. 114, 115).

Jaekel (1916) published information on the basicranium of *Proganochelys* ("*Triassochelys*"), but in retrospect it has proven to be misleading. Although Jaekel's specimen from Halberstadt is in general well preserved (the cervicals, for example, are still the best known), the skull has been subjected to lateral crushing. The anterior, denticle bearing portions of the pterygoids have been pushed together at the midline, producing what Jaekel interpreted as a median toothbearing element covering the area where the basiptyergoid articulation might be found, resulting in an apparently akinetic skull. Although this reconstruction was reproduced from time to time (cf. Romer, 1956), it did not stimulate very much discussion or comment.

The discovery of the Aixheim specimen in 1927 would not have significantly aided basicranial studies, even had it been described, because it is also laterally crushed and lacks the posterior portion of the braincase. Nonetheless, it is the only specimen preserving the major part of the epiptyergoid, and it reveals a number of other important features. The discovery of the best preserved Triassic turtle skull was the result of a combined University of Tübingen-Staatliches Museum für Naturkunde, Stuttgart, effort to collect *Plateosaurus* skeletons from a quarry in Trossingen. Near the end of the 1932 season they discovered three turtle skeletons. Only one of the specimens, SMNS 16980, has a skull, but it is relatively well preserved. These three specimens were prepared at Stuttgart under the direction of Dr. F. Berckhemer, and were later in the care of Professor K. Staesche. It was during Staesche's stewardship of the material in the 1950's that Ernest E. Williams examined the skull

and presented his conclusions (Parsons and Williams, 1961) that the cranium possessed many generalized amniote features. In 1975 I presented a cladistic view of his information and figures and argued that *Proganochelys* was the sister group to all other turtles. The nature of the basicranial articulation was dubious from available photographs, and I suggested that the articulation might actually be fused rather than movable (Gaffney, 1975: 424). The actual material, however, had still not been studied in detail, so it was with great elation that in late 1978 I was given permission to work on the Stuttgart material.

DESCRIPTION

In the generalized amniote condition, such as seen in *Captorhinus*, the palatoquadrate is articulated to the braincase by means of a movable basiptyergoid articulation. In turtles of the suborder Casichelydia this kinesis is lost, and extensive sutural contact between the pterygoid and basisphenoid obliterates the articular area. As I have discussed in detail elsewhere (Gaffney, 1975, 1979b), the basicranium of turtles is an important source of characters differentiating the two main infraorders of casichelydian turtles, Cryptodira and Pleurodira. Embryonic stages of turtles have rudiments of a palatobasal articulation in the form of the cartilago articularis and the processus basiptyergoideus (Kunkel, 1912), demonstrating that the generalized amniote condition can also be found in turtles. Pleurodires have a medial process of the quadrate that forms the main brace to the braincase, whereas in cryptodires the pterygoid extends posteriorly to form the brace between braincase and quadrate. In adults of both groups the basiptyergoid articulation is fused and the area of the cranio-quadrate space is nearly obliterated by the broad sutural contact of the braincase (prootic, opisthotic) and palatoquadrate (quadrate,

pterygoid). The akinetic condition of casichelydian turtles, therefore, not only includes fusion of the basiptyergoid articulation, but also the development of these subsidiary contacts anterior, dorsal, and posterior, to the position of the articulation.

In known captorhinids (*Captorhinus*, *Eocaptorhinus*, see Heaton, 1979; Warren, 1961) the basicranial articulation is formed between the epiptyergoid and basisphenoid. The epiptyergoid has a well-developed ventral expansion that sends a curved process into a pocket of the pterygoid so that the actual articulation is borne by the epiptyergoid.

In *Proganochelys* the epiptyergoid is poorly known; the most complete one is in SMNS 15759, but crushing has distorted its position, and the ventral limits are broken. However, in SMNS 16980, on the left side, there is what appears to be a fragment of curved bone closely applied to the pterygoid, and I interpret this as the epiptyergoid contribution to the articulation (see Fig. 1). The sutures on the ventral surface are not clear but seem to indicate that the epiptyergoid forms most

of the articular area. It is possible, however, that the posterior part of the articulation is actually borne on the pterygoid.

The basisphenoid of *Proganochelys* has well-developed, paired basiptyergoid tubercles that extend ventrolaterally (see Fig. 2), rather than anterolaterally as in *Captorhinus* (Price, 1935). The articular surface is nearly flat and faces anterolaterally to fit into the basicranial recess. The tubercles are posterior to the dorsum sellae and sella turcica rather than anterior as in *Captorhinus*. In another paper (Gaffney, 1979a) I argued that a structure on the basisphenoid of *Mesochelys* was not a remnant basiptyergoid process as identified by Evans and Kemp (1975) because it is posterior to the dorsum sellae rather than anterior as in *Captorhinus*. This argument no longer has validity because the tubercles are also posterior in *Proganochelys* thus reopening the question of the identity of these structures in *Mesochelys* and *Glyptops* (I hope to deal with this problem elsewhere).

The pterygoid of *Proganochelys* has a basicranial recess that probably includes

Figure 1. *Proganochelys quenstedtii*, SMNS 16980, stereo photograph of ventral view of skull.

Figure 2. *Proganochelys quenstedti*, SMNS 16980, ventral view of basicranium.

at least part of the epipterygoid, but because the suture is dubious, particularly in the ventral view, I am unable to describe the portions formed by each bone. The recess has the form of a trough that is oriented anterodorsomedially and closely fits the opposing tubercle on the basisphenoid. The general form of the recess is similar to *Captorhinus*, but the recess faces more posteriorly in *Proganochelys*. The question of actual movement in the skull of *Proganochelys* must await the complete description of the specimens, but it is clear that some sort of movement was possible in the basicranial articulation.

The akinesis of all turtles except *Proganochelys* results from a broad sutural fusion of the basisphenoid and pterygoid that I have interpreted (Gaffney, 1975) as a shared derived character testing the monophyly of the Casichelydia. The recognition of the generalized amniote condition of the basicranial articulation in *Proganochelys* supports the argument that it (as the suborder Proganochelydia) is the sister group of the Casichelydia.

ACKNOWLEDGMENTS

The opportunity to study the Triassic turtles in Stuttgart is due to the cooperation and support of Rupert Wild, Curator, and Bernhard Ziegler, Director, Staatliches Museum für Naturkunde in Stuttgart (SMNS). Wild spent many hours aiding me and my colleagues, and I am very grateful for his help and hospitality. I would also like to thank Karl-Heinz Fischer, Museum für Naturkunde, Berlin, for giving me access to the Jaekel Triassic material in East Berlin. Otto Simonis, Department of Vertebrate Paleontology, American Museum of Natural History (AMNH), performed the superb preparation necessary to expose the basicranial morphology of the Stuttgart *Proganochelys* skull. Chester Tarka, also of the Department of Vertebrate Paleontology, AMNH, took the photographs. This paper is part of a project funded by National Science Foundation Grant DEB 8002885.

LITERATURE CITED

- BERKHEMER, F. 1931. Die Saurierfunde aus den Keuperablagerungen der Baar. Tuttlinger Heimatblätter, Heft 14, 12 pp.
- . 1938. 25 Jahre Verein zur Förderung der Württ. Naturaliensammlung in Stuttgart. Stuttgart, Ernst Klett, pp. 1-18.
- . 1951. Die Sprachen der Steine. Verlag "Die Schönen Bücher," Reihe C, "Natur" Band 2, Stuttgart, pp. 9, 36.
- EVANS, J., AND T. S. KEMP. 1975. The cranial morphology of a new lower Cretaceous turtle from southern England. *Palaeontology*, 18: 25-40.
- GAFFNEY, E. S. 1975. A phylogeny and classification of the higher categories of turtles. *Bull. Amer. Mus. Nat. Hist.*, 155(5): 387-436.
- . 1979a. The Jurassic turtles of North America. *Bull. Amer. Mus. Nat. Hist.*, 162(3): 91-136.
- . 1979b. Comparative cranial morphology of recent and fossil turtles. *Bull. Amer. Mus. Nat. Hist.*, 164(2): 65-375.
- HEATON, M. J. 1979. Cranial anatomy of primitive captorhinid reptiles from the late Pennsylvanian and early Permian Oklahoma and Texas. *Oklahoma Geol. Surv., Bull.* 127, pp. 1-84.
- JAEKEL, O. 1916. Die Wirbeltierfunde aus dem Keuper von Halberstadt. Serie II. Testudinata, Teil 1, *Stegochelys dux*, n.g., n. sp. *Paläont. Zeitschr.*, 2: 88-214.

- KUHN, O. 1956. Deutschlands vorzeitliche Tierwelt (2nd ed.). München, Bayerischer Landwirtschaftsverlag, 126 pp.
- KUNKEL, B. W. 1912. The development of the skull of *Emys lutaria*. J. Morphol., 23: 693-780.
- PARSONS, T. S., AND E. WILLIAMS. 1961. Two Jurassic turtle skulls: a morphological study. Bull. Mus. Comp. Zool., 125: 43-107.
- PRICE, L. I. 1935. Notes on the braincase of *Captorhinus*. Proc. Boston Soc. Nat. Hist., 40(7): 377-386.
- ROMER, A. S. 1956. The osteology of the reptiles. Univ. Chicago Press, 772 pp.
- WARREN, J. W. 1961. The basicranial articulation of the early Permian cotylosaur, *Captorhinus*. J. Paleont., 35(3): 561-563.

The Rectus Abdominis Muscle Complex of the Lacertilia: Terminology, Homology, and Assumed Presence in Primitive Iguanian Lizards

SCOTT M. MOODY¹

ABSTRACT. Camp (1923) divided the lacertilians into two groups, Autarchoglossa and Ascalabota, on the basis of the presence or absence, respectively, of the Muscle rectus abdominis superficialis. His recognition of the "superficialis" is both anatomically and terminologically incorrect. The M. rectus abdominis has only three parts in the Autarchoglossa: medialis, lateralis, and pyramidalis. In the Ascalabota, it has only two parts: medialis and pyramidalis. Nearly all genera of the iguanian families Agamidae, Chamaeleonidae, Uromastycidae, and Iguanidae (including Oplurinae) were surveyed for the third part of the M. rectus abdominis. I confirmed the earlier observations by anatomists of a small muscle lying superficial to the M. pectoralis and inserting into the skin at the axilla in the uromastycids *Leiolepis* and *Uromastyx* and the primitive agamid *Physignathus*, and I additionally observed it in the agamid *Hydrosaurus*. Observed developmental variations, comparative morphology and innervation demonstrate that this muscle is part of the M. pectoralis and is named the M. pectoralis pars cutaneous. Camp believed this muscle to be instead a vestigial M. rectus abdominis lateralis and considered this evidence that the Ascalabota were evolutionarily advanced lizards which had lost the autarchoglossan attributes of terrestrial slinking made possible by this muscle. My reanalysis suggests instead that the Ascalabota are primitive and that the ancestral lacertilians possessed only two parts of the M. rectus abdominis, with the third part evolving later with the ancestor of the Autarchoglossa.

INTRODUCTION

The goals of this paper are to: 1) review the terminology and anatomy of the ventral and lateral trunk musculature, with emphasis on the M. rectus abdominis complex, among lacertilians on the basis of both original dissections and translations of the often cited but seldom read publications of the comparative anatomists Gadow, Maurer, Hoffmann, and Fürbringer working in the 19th century of Germany; 2) survey the lizard infraorder Iguania (Agamidae, Chamaeleonidae, Uromastycidae, and Iguanidae) defined by Camp (1923) and widely believed to be monophyletic (Underwood, 1971; Northcutt, 1978) for a double-layered M. rectus abdominis; 3) determine the homology of the small cutaneous muscle slip lying lateral to the M. pectoralis which is present in the primitive agamids *Physignathus* and *Hydrosaurus* (infrequently observed) and uromastycids *Leiolepis* and *Uromastyx* and was assumed by Camp (1923) to be a vestigial superficial layer of the M. rectus abdominis; and 4) assess the value of the variable M. rectus abdominis as a phylogenetic character in hypothesizing cladistic relationships among lizard families because this character was used by Camp (1923) for the first dichotomy in his classification of lizard families into higher taxa.

¹Department of Zoology and College of Osteopathic Medicine, Ohio University, Athens, Ohio 45701, U.S.A.

TERMINOLOGY

Herpetologists and reptilian anatomists currently follow Camp's (1923: 377-378) arrangement of the *M. rectus abdominis*:

The Rectus abdominis muscle in autarchoglossid lizards can usually be subdivided into four parts . . . 1) the Rectus profundus or main median portion of the Rectus, segmented . . . 2) the Rectus medianus, segmented, often intimately connected with the skin over the whole belly . . . 3) the Rectus lateralis, unsegmented, and forming the lateral continuation of the Rectus medianus, 4) the Rectus internus, unsegmented, arising on the pubis and passing forwards sometimes as far as the sternum and dorsal to the Rectus profundus . . . the medianus and lateralis may together be called the Rectus superficialis.

Camp attributed the observation of these four divisions to Maurer (1896). However, careful reading of Maurer's monograph reveals that: 1) Maurer recognized only three parts of the *M. rectus abdominis*; 2) the *M. rec. abd. medianus* and *M. rec. abd. profundus*¹ distinguished by Camp are in fact the same muscle; and 3) the *M. rec. abd. medianus* and *M. rec. abd. lateralis* are not embryonically or anatomically primarily continuous. Gadow (1882) earlier also recognized only three divisions of the *M. rectus abdominis*.

Maurer (1896: 228-229, my translation) summarized the variation of the *M. rectus abdominis* among lacertilians:

The *M. rectus abdominis* of lizards is complicated . . . in *Lacerta* a *M. rec. abd. medialis* and *lateralis* can be distinguished. The latter is missing in *Sphenodon*. Both extend completely together from the pelvis to the posterior edge of the sternum. Here the medial part terminates, as does the entire *M. rectus abdominis* in *Sphenodon*. However the *M. rec. abd. lateralis* of lizards is continued further anterior as a ventral slip

superficial to the *M. pectoralis* and is overlapped along its lateral edge by the *M. obliquus abdominis externus superficialis*. The boundary between the *M. rec. abd. medialis* and *lateralis* is easily recognized if the *M. rec. abd. lateralis* is first severed and reflected from its lateral edge. The *M. obliquus externus profundus* can then be seen to meet along a line with the *M. rec. abd. medialis*, just as in *Sphenodon*. This line demarcates the boundary between both *M. rectus abdominis* portions, as well as the lateral-most boundary of the entire *M. rectus abdominis* of *Sphenodon* . . . The superficial layer of *M. rectus abdominis* connects with the integument without aid of inscriptional ribs or gastralia . . . along the margins of the large transverse series of ventral scutes, by inserting into the skin by fibers coming from both the *M. rec. abd. medialis* as well as the *M. rec. abd. lateralis* muscles. Because of this manner of attachment, the *M. rectus abdominis* is necessarily segmented, similar to *Sphenodon* which is segmented by inscriptional ribs, but in the case of the lizard by a series of fibrous bundles inserting into the skin. However these bundles are comprised only of the superficial layer of fibers. Therefore in *Lacerta* superficial and deep portions of the *M. rec. abd. medialis* may be distinguished, but they are continuously adherent to each other. The *M. rec. abd. lateralis* of *Lacerta* is connected with the skin to an even greater extent than the *M. rec. abd. medialis*, especially at its anterior end which ends bluntly at the base of the neck as a skin muscle. A triangular muscle (*M. rec. abd. pyramidalis* after Maurer, *M. rec. abd. internus* after Gadow) develops deep to the *M. rec. abd. medialis* and immediately anterior to the pelvis with fibers radiating anteriorly and medially and inserting into the linea alba.

I have examined within the Autarchoglossa, a scincid *Tiliqua scincoides* (UMMZ 65915), the lacertids *Lacerta agilis* (UMMZ 119991), *L. lepida* (UMMZ 72473), *L. viridis* (UMMZ 65756), and *Podarcis muralis* (UMMZ 65947), a teiid *Ameiva ameiva* (personal collection) and a cordylid *Cordylus giganteus* (personal collection), and have confirmed the observations of Maurer and Gadow. Ventral and anterolateral views of the musculature of *Ameiva ameiva* are illustrated in Figures 1 and 2.

Throughout his paper Maurer unfortunately changes his terminology and switches back and forth from formal latinized names to descriptive anatomical

¹For sake of brevity, the name *M. rectus abdominis* will be abbreviated to *M. rec. abd.* when preceding the descriptors *superficialis*, *lateralis*, *profundus*, etc.

Figure 1. Ventral view of the trunk musculature of *Ameiva ameiva* (Teiidae). Specimen is superficially dissected on the right with a window cut in the M. rectus abdominis, and deeply dissected on the left with portions of the M. obliquus abdominis externus and M. obl. abd. internus removed at different levels to reveal underlying layers.

Abbreviations: abd lat, M. rectus abdominis lateralis; abd med, M. rectus abdominis medialis; abd pyr, M. rectus abdominis pyramidalis; ext obl, M. obliquus abdominis externus; int obl, M. obliquus abdominis internus; myccom, Myocommata; pect, M. pectoralis; trn abd, M. abdominis transversus.

terms. These changes were probably the cause of Camp's misinterpretation of the anatomical arrangement and homology of

the different parts of the M. rectus abdominis. For example, Maurer describes the transverse series of fibrous bundles as originating from the superficial layer of the M. rec. abd. medialis and attaching to the skin. Camp interprets these collectively as a distinct muscle, the M. rec. abd. superficialis medianus, and assigns the name M. rec. abd. profundus to the deeper fibers which do not connect with the skin and are not segmented. However, Maurer (1896: 229) emphatically points out that these parts are not separable as distinct muscles: "...die aber kontinuierlich zusammenhängen," ("however they are continuously adherent to each other," my translation).

Camp may also have been easily confused by Maurer's (p. 206) description of *Lacerta*, which I translate as follows:

In *Lacerta* the M. rectus abdominis extends from the pelvis to the neck region as the M. rec. abd. superficialis. It is comprised of two distinguishable parts which I have referred to as the M. rec. abd. medialis and lateralis. The M. rec. abd. medialis extends from the pubic symphysis to the posterior end of the sternum. Superficial fibers attach this layer to the integument, inserting along the posterior edges of the scales. Along its entire lateral margin it meets the M. pectoralis anteriorly... and the M. obliquus abdominis externus profundus posteriorly. The superficial M. rec. abd. medialis is the only muscle present in *Sphenodon*, which is also connected with the M. pectoralis and M. obliquus abd. ext. profundus. The superficial M. rec. abd. lateralis is absent in *Sphenodon*. In *Lacerta* it is attached directly to the M. rec. abd. medialis along its border, and extends from pelvis to throat as a similarly wide band, which arcs laterally along its middle part. It extends from the Ligamentum ileo-pubicum anteriorly and laterally, overlies (superficially) the M. obliquus abd. ext. profundus, and is overlain by the M. obliquus abd. ext. superficialis along its lateral border. It continues forward to overlay the ventral surface of the M. pectoralis, nearly approaches the midline of the M. pectoralis, and terminates in an aponeurosis which attaches to the integument along the border of the anterior most large transverse ventral scutes. Consequently, this M. rec. abd. lateralis is intimately connected with the integument, especially the anterior part which overlies

the M. pectoralis, and is entirely a cutaneous muscle. Superficial fibers of the posterior part of the M. rec. abd. lateralis insert into the skin but deeper fibers comprise a smooth (continuous) muscle belly, without tendinous inscriptions, and is rather unsegmented as Gadow and others have already demonstrated.

Maurer discussed the M. rec. abd. medialis and lateralis collectively as the superficial M. rectus abdominis but not as a formally named M. rec. abd. superficialis. Additionally, he did not describe a M. rec. abd. profundus, which would have been necessary as a contrasting member of the pair of muscles. Apparently, he considered the M. rec. abd. pyramidalis (Camp's internus) as the deep muscle in contrast to his collective superficial M. rectus abdominis.

It is additionally evident that Maurer (p. 219, my translation) intended to recognize only three parts of the M. rectus abdominis from his description of the scincid *Tiliqua* sp:

One recognizes namely... that three proper, separable M. rectus abdominis parts lie near one another, that the M. rectus medialis is divided once again into medial and lateral parts. Directly before the pelvis, a triangular muscle is observed...

Note that the division of the M. rectus medialis is not into formally named muscles but only two anatomical regions which are defined in relationship to the position of the medial edges of the M. rec. abd. lateralis and M. obliquus abd. ext. profundus.

Camp (p. 378) also described the M. rec. abd. lateralis as the lateral continuation of the M. rec. abd. medianus (believed by him to lie superficial to the M. rec. abd. profundus). Referring to the M. rec. abd. lateralis, he wrote: "... according to Maurer (1898) it arises in the embryo of *Lacerta* as the lateral portion of the superficial layer of the M. rectus abdominis." Camp may have based his conclusions on the text figures in Maurer

Figure 2. Anterolateral view of the trunk and thoracic musculature of *Ameiva ameiva* (Teiidae). Windows have been cut in the M. obliquus abdominis externus and the M. intercostalis externus to show deeper layers.

Abbreviations: cos ext, M. intercostalis externus; cos int, M. intercostalis internus; erec sp, M. erector spinae; latis, M. latissimus dorsi, and rib; and see Figure 1.

rather than the text. In adult *Lacerta* spp. the M. rec. abd. lateralis is anatomically adjacent to the superficial fibers of the M. rec. abd. medialis, but Camp misinterpreted both the anatomical positions of the three named parts of the M. rectus abdominis (see above) and the embryonic origin of these parts. A modified and semischematic illustration of the late developmental stages of the trunk musculature of *Lacerta agilis* (Maurer, 1898: Fig. 23) is presented in Figure 3.

In his review of the embryonic development of the ventrolateral trunk muscles of lacertilians, Maurer (1898: 55–56, my translation) observed:

Figure 3. Transverse section of a 31 day old embryo of *Lacerta agilis* (Lacertidae). Illustration is semischematic and modified from Maurer (1898: Fig. 23).

Abbreviations: abd lat, M. rectus abdominis lateralis; abd med, M. rectus abdominis medialis; abd trn, M. abdominis transversus; cos ext brev, M. intercostalis externus brevis; cos ext long, M. intercostalis externus longus; cos int brev, M. intercostalis internus brevis; cos int long, M. intercostalis internus longus; ext obl pro, M. obliquus abdominis profundus; ext obl sup, M. obliquus abdominis externus superficialis; int obl, M. obliquus abdominis internus; perit, Peritoneum, rib, and skin.

The M. rectus abdominis differentiates from a uniform Anlage but in a different manner than in the Urodeles. I distinguish a segmented M. rec. abd. medialis that develops from the caudal border of the sternum to the pubic symphysis. The M. rec. abd. lateralis is unsegmented and extends anteriorly from the pubis to the throat region, where it extends over the ventral surface of the M. pectoralis major muscle and ends in the integument. The M. pyramidalis differentiates from the M. rec. abd. medialis and arises from the pubic symphysis. It is not similar to the identically named mammalian muscle, but is located instead deep and adjacent to the peritoneum. The M. rec. abd. medialis extends laterally to join with the ventral margins of the M. intercostalis externus and internus because these parts are embryonically related.

I summarize the nomenclatural history of the M. rectus abdominis and its parts in Table 1 and suggest that this muscle be recognized as a tripartite muscle. I further suggest that the terms M. rec. abd. medialis and M. rec. abd. lateralis be recognized and used rather than the M. rec. abd. profundus and M. rec. abd. superficialis. This follows either older or more frequently used terminology, more accurately depicts the anatomical positions of the two muscle parts, and avoids confusion with the incorrect muscle divisions recognized and named by Camp (1923). The M. pyramidalis is considered to be embryonically and functionally a part of the M. rectus abdominis.

HISTORICAL REVIEW

Morphological variations of the pectoral girdle musculature (Fürbringer, 1900; Sukhanov, 1961 and 1976) and the ventrolateral trunk musculature (Maurer, 1896; Gadow, 1882; Camp, 1923) have provided several characters frequently used for hypothesizing the phylogenetic relationships of lizard families. In Camp's (1923) "Classification of the Lizards" three of the 34 character complexes reviewed were of the musculature. English speaking herpetologists rely heavily on this monograph as the standard reference for lacertilian com-

TABLE 1. NOMENCLATURAL HISTORY OF THE M. RECTUS ABDOMINIS COMPLEX.

M. RECTUS ABDOMINIS MEDIALIS	(Maurer, 1896, 1898)
= "ventralis"	Gadow, 1882; Hoffman, 1890
= "abdominis"	Mivart, 1867, 1870; Sanders, 1870, 1872, 1874
= "profundus"	Camp, 1923
= "externus"	Byerly, 1925
= "superficialis medianus"	Camp, 1923
M. RECTUS ABDOMINIS LATERALIS	(Gadow, 1882; Hoffman, 1890; Maurer, 1896, 1898; Fürbringer, 1900)
= "suprapectoralis"	Rüdinger, 1868; Fürbringer, 1869, 1887
= "superficialis lateralis"	Camp, 1923
M. RECTUS ABDOMINIS PYRAMIDALIS	(Gorski, 1852; Stannius, 1856; Mivart, 1867, 1870; Sanders, 1870, 1872, 1874; Maurer, 1896, 1898)
= "internus"	Gadow, 1882; Hoffman, 1890; Camp, 1923; Byerly, 1925
= "triangularis"	Maurer, 1896

parative myology, primarily because Camp summarized earlier German works.

Subsequent to Camp (1923) there have been several comparative and functional studies of the limb and girdle musculature of lizards possessing snakelike locomotory adaptations, but comparative and phylogenetic studies of the musculature of nonslinking tetrapod lizards have been few in number. This is unfortunate because these lizards (especially the Iguania) are morphologically more similar to the fossil ancestral lacertilians of the Permian and Triassic (Robinson, 1967) and are presumably more primitive than other extant lizard families (Sukhanov, 1976; Estes, 1983; Moody, in preparation). Cladistic analysis of the character states derived from the present restudy of the M. rectus abdominis supports this view (see discussion section).

Fürbringer (1869) named the cutaneous muscle lying superficial to the M. pectoralis as the M. suprapectoralis. Although he studied several autarchoglossan lizards with a fully developed M. rec. abd. lateralis, he was probably strongly influenced by the condition of this muscle in *Uromastix*, which he had primarily described. As will be discussed later, Fürbringer was correct in believing this muscle to be part of the M. pectoralis.

Mivart (1867, 1870) pioneered the study of lizard musculature with superbly illustrated and described dissections of *Iguana tuberculata* (= *I. iguana*) (Iguanidae) and *Chamaeleo parsonii* (Chamaeleonidae). Sanders (1870, 1872, 1874) published three detailed studies of *Platydictylus japonicus* (= *Gecko japonicus*) (Gekkonidae), *Leiolepis belliana* (Uromastycidae), and *Phrynosoma coronatum* (Iguanidae). DeVis (1884) published a dissection of *Chlamydosaurus kingi* (Agamidae). Shufeldt (1890) thoroughly described the musculature of *Heloderma suspectum* (Helodermatidae). All but one of these studies, which were published in English, were concerned with four lizard families which belong to Camp's defined suborder Ascalabota. This taxon is characterized by simple ventrolateral trunk musculature. The German comparative myological publications, which appeared during the same time period, included primarily representatives of the lizard families belonging to Camp's suborder Autarchoglossa, the sister taxon of Ascalabota. These lizards possess a more complex ventrolateral musculature.

The earliest German study was that of Gorski (1852) who concentrated on the pelvic girdle. Stannius (1856) reviewed lizard anatomy and Rüdinger (1868) produced a comparative study of the fore-

limb musculature of reptiles and birds. Three European lizards *Anguis fragilis* (Anguidae), *Chalcides chalcides* (Scincidae), and *Ophisaurus apodus* (Anguidae) were included in these works. Fürbringer (1869, 1875, 1900) produced a classic series of papers describing and illustrating the skeleton, nerves, and muscles of the lacertilian pectoral girdle, especially that of *Uromastix aegyptius* (Uromastycidae). Gadow (1882) compared the ventrolateral musculature of 16 lizard species representing the families Varanidae, Teiidae, Lacertidae, and Scincidae of the Autarchoglossa, and the families Iguanidae, Gekkonidae, and Chamaeleonidae of the Ascalabota. Hoffman (1890) reviewed these investigations, added original observations, and listed synonymous names for different muscles.

Maurer (1896) grossly compared the abdominal musculature of *Lacerta agilis*, *L. viridis*, *Podarcis muralis* (Lacertidae), *Tiliqua* sp. (Scincidae), *Chamaeleo* sp. (Chamaeleonidae), and *Sphenodon punctatus* (Sphenodonidae). This was followed in 1898 by a study of the embryonic origin of the ventrolateral trunk musculature of *Lacerta agilis*. As I discussed earlier, this study developed accurate terminology and determined the homology of the parts of the M. rectus abdominis.

The phylogenetic significance of the different anatomical arrangements of the M. rectus abdominis in lacertilians was not appreciated until Camp (1923) summarized the literature, made original observations, and hypothesized a phylogenetic classification of the lizard families. Presence or absence of his M. rectus abdominis "superficialis" (including his named parts medianus and lateralis) was selected (pp. 297–298, 355, 385–415) as the major character in creating his phylogenetic classification. He cladistically divided the Lacertilia into two taxa and assigned to them the subordinal names Autarchoglossa and Ascalabota. The former is characterized by presence of

the M. rectus abdominis "superficialis"; the latter by its absence. He also noted the presence of a vestigial M. rectus abdominis "superficialis" in *Leiolepis belliana*, and *Uromastix hardwicki* (Uromastycidae), and *Physignathus lesueurii* (Agamidae).

Prior to Camp's monumental work, the M. rec. abd. "superficialis" had been described and figured (but unfortunately not labeled) for only one agamid, *Uromastix spinipes* (= *U. aegyptius*) (Fürbringer, 1875: 715, Figs. 63, 73). Moody (1980) placed *Leiolepis* and *Uromastix* in the separate family Uromastycidae. I have translated Fürbringer (p. 715) as follows:

Compared with each other (*Uromastix aegyptius* and the scincids *Macrosцинus coctei*, *Scelotes* sp., *Chalcides chalcides*), the lateral part of the superficial layer of the M. pectoralis will be overlain by a small slip of muscle (M. suprapectoralis) which originates either from the posterolateral area of the M. pectoralis itself or alternatively from the M. obliquus abd. externus and extends forwards to just posterior to the axilla (*Uromastix*) or the area of the clavicle (scincids) where it inserts as an aponeurosis into the skin.

Camp (p. 380) incorrectly cited Fürbringer as having examined *Uromastix hardwicki*.

Camp (p. 380) cites Sanders (1872) as having described the M. rec. abd. "superficialis" in the uromastycid *Leiolepis belliana*. However, Sanders never mentioned this muscle slip, although the adjacent M. rec. abd. medialis (Camp's "profundus") was described, and the M. rec. abd. "superficialis" was unambiguously figured, though not labeled (Sanders: Fig. 2). Evidently Camp personally observed the M. rec. abd. "superficialis" in *Physignathus lesueurii* as an accompanying literature citation is lacking. Whether he also examined specimens of *Uromastix hardwicki* and *Leiolepis belliana* cannot be proven because they were neither figured nor identified by museum catalogue numbers.

STATEMENT OF PROBLEM

There has been considerable disagreement among herpetologists, both in literature and especially in private discussions, concerning whether Camp believed the Ascalabota to be primitive or derived relative to the Autarchoglossa and whether the Ascalabota (Gekkota and Iguania) is a monophyletic or polyphyletic suborder. The source of the disagreements is a contradiction between Camp's cladogram (p. 333) and numerous statements in his text. The cladogram suggests that the Iguania is primitive and the sister group to all remaining lacertilians. However in his classification (pp. 297-298) and summary (pp. 416-417) Camp divides the Lacertilia into the currently widely used suborders Ascalabota and Autarchoglossa. In this scheme Camp states that the Gekkota and Iguania are sister taxa and uses the absence of the *M. rec. abd. "superficialis"* as their primary definitive character.

Concerning the direction of evolution of the one or two-layered *M. rectus abdominis*, it is quite clear what Camp believed. He wrote (p. 385) that:

It is apparent . . . that the *Rectus superficialis* is a morphologically widespread and embryologically precocious muscle. Its absence in Ascalabota cannot well be due to primitiveness of that group but more probably to the fact that all forms in the group except certain primitive agamids have *lost* the muscle, while it remains and develops in the Autarchoglossa. This would seem to imply that the pro-Sauria may have carried the body close to the ground and that this mode of locomotion has been correlated with the development of "slinking" musculature in the belly-wall.

Lizards today which retain this mode of locomotion retain the musculature, while the group that seems to have early adopted arboreal life has apparently lost the musculature.

In his summary, Camp wrote (p. 420) that:

The superficial layer of the rectus is regarded as a primitive crawling muscle. Its presence in certain primitive agamids tends to show this and its ab-

sence in all other Ascalabota is considered secondary . . .

Within the Ascalabota, Camp believed that the Agamidae (and Uromastycidae) was evolutionarily derived from within (not as sister taxa) the Iguanidae (p. 333). However, this assumption is contradicted by his observation that *Leiolepis belliana*, *Uromastyx hardwicki*, and *Phrygnathus lesueurii* possess the *M. rec. abd. "superficialis"* because these species are supposedly representative of primitive iguanians which have retained vestigially the *M. rec. abd. "superficialis"* from their autarchoglossan-like ancestor. Camp's argument that this muscle is a plesiomorphic character for the lacertilians is not congruent with the presence of the derived character states such as acrodont dentition, simplified dermocranium and pectoral girdle, lack of inscriptional abdominal ribs in both the Agamidae and Uromastycidae and presence of the epiotic foramen in the Agamidae (Moody, 1980).

Also inexplicable is Camp's defense of the dichotomy of the Autarchoglossa and Ascalabota when the above three species, presumed by him (pp. 416-417) to represent the ancestral clade of the Agamidae, exhibit the "wrong" character state.

Camp dissected the musculature of only a few species and since his work, the *M. rectus abdominis* has been studied phylogenetically only in the Gekkota (Kluge, 1976). Camp examined six agamid genera from a total of 52 recognized genera, two of two uromastycid genera, 19 of 57 iguanid genera, and one of four chamaeleonid genera. I informally recognize the Madagascan genera *Oplurus* and *Chalarodon* as the subfamily Oplurinae in the Iguanidae on the basis of the ulnaris nerve pathway, abdominal inscriptional ribs, dermocranium, hemipenes, scalation and vicariant zoogeography. Camp examined one of two oplurine genera. The two iguanids *Hoplocercus* and *Morunasaurus* considered by Etheridge (personal com-

munication) to be primitive were not examined. If additional agamid and uromastycid species had been examined by Camp, then he would have discovered a character state distribution of the *M. rectus abdominis* which would not have been congruent with the genera *Uromastyx* and *Physignathus* as recognized at the time of his work (see later discussion).

If the Agamidae, Chamaeleonidae, and Uromastycidae were evolutionarily derived from within the Iguanidae as Camp suggested, then at least some of the primitive iguanids should be expected to also possess the *M. rec. abd.* "superficialis." Of course, independent loss of the muscle after the phyletic evolution of the above three families may explain this character state distribution. Subsequently, I surveyed all four families of the Iguania exhaustively, but many more species per genus for the Agamidae were examined because the *M. rec. abd.* "superficialis" was known to be already present and because the family was in sore need of taxonomic revision.

The morphological nature of the *M. rec. abd.* "superficialis" in *Uromastyx*, *Leiolepis* and *Physignathus* is substantially different than the equivalent muscle in the Autarchoglossa. Camp believed that the muscle found in these genera was a vestigial remnant. A goal of this study is to determine whether this homology is correct.

ANATOMICAL OBSERVATIONS

For the Agamidae, I examined 177 of the circa 300 species which represent 48 of the 52 recognized genera and subgenera. The taxonomic nomenclature follows Moody (1980). The four genera not examined (*Harpesaurus*, *Mictopholis*, *Thaumatorhynchus*, undescribed gen. nov.) are known only from type specimens. *Harpesaurus* is closely related to *Pseudocalotes*, and a new genus is being described for "*Cophotis*" *sumatrana*

(Moody and Böhme, in preparation). Both genera were examined. *Mictopholis austeniana* is related to *Salea* according to Smith (1935) and I examined this genus. *Thaumatorhynchus brooksi* is closely related to *Aphaniotis* which I examined. The undescribed gen. nov. (Arnold, in preparation) is closely related to *Phrynocephalus* which I sampled.

For the Uromastycidae, I examined 19 of the 20 species representing the only genera, *Uromastyx* and *Leiolepis*. Only *Uromastyx dispar* was not examined because the single known specimen, the holotype, is only a skin and skull.

For the Chamaeleonidae, I examined 19 of the circa 100 species representing the four genera *Bradypodion*, *Brookesia*, *Chamaeleo*, and *Rhampholeon*. Six species representing four of the nine species groups of *Chamaeleo* inhabiting Madagascar and seven species of the four groups inhabiting Africa were examined. My nomenclature follows Hillenius (1959), Raw (1976), and Klaver and Böhme (in preparation).

For the Iguanidae excluding Oplurinae, I examined 86 of the circa 630 species which represent all 53 currently recognized genera (Etheridge, in preparation). Two recently described monotypic genera from South America, *Vilcunia* Donoso-Barros and Cei (1971) and *Pelusaurus* Donoso-Barros (1973), were not available for study. Moreover, their status as valid genera is in dispute. The original authors hypothesized relationships to *Liolaemus* and *Proctotretus*, but Etheridge (personal communication) has inferred instead from the descriptions, relationship to the *Liolaemus-Ctenoblepharis* complex. In either case these three related genera were examined. Most of the species groups of the two large genera *Anolis* and *Sceloporus*, following the respective arrangements by Etheridge (1959) and Smith (1939) were sampled. I also examined 14 of the 19 species of the three primitive iguanid genera *Morunasaurus*, *Enyaliodes*, and *Hoplocercus*. For the Oplurinae, I ex-

amined six of the seven species belonging to the two genera *Oplurus* and *Chalarodon* of Madagascar.

Camp's conclusion that within the infraorder Iguania, only three genera possess the M. rec. abd. lateralis was confirmed. All five species of *Leiolepis*, both species of *Physignathus* and all species of *Uromastix*, except *U. asmussi* and *U. loricatus*, possess this muscle. In addition, one specimen of *Hydrosaurus pustulatus* (personal collection), a close relative of *Physignathus*, has a tiny muscle slip on only one side. Six other examined specimens representing all three species of *Hydrosaurus* totally lacked the muscle.

All remaining examined agamids, chamaeleonids, iguanids, and oplurines completely lack a lateral or superficial second layer of the M. rectus abdominis or a vestigial muscle slip overlying the M. pectoralis.

Presence or absence of the muscle slip in question was determined by making a paraventral incision through the skin along the entire length of the pectoral girdle and sternum and by reflecting the skin laterally. This muscle is a thin ribbonlike slip lying superficial and lateral to the large fanshaped M. pectoralis. This muscle slip is oriented obliquely with its insertion ventromedial to the axilla and its origin the dorsolateral corner of the thoracic region.

The insertion of this muscle anteriorly is a broad tough sheet of fibrous connective tissue which firmly attaches the muscle to the skin medial to the forearm, which can be detached only by tearing or cutting. The posterior origin of this muscle is not attached to the skin. Instead, the fibers merge imperceptively with fibers of three muscles (M. obliquus abd. externus, M. pectoralis, and M. rec. abd. medialis) at the area of common intersection which overlies the point of articulation of the last xiphisternal rib with the last thoracic rib. The fibers of these four merging muscles have a subparallel orientation in this common area.

This muscle is unsegmented, unlike the M. rec. abd. medialis which is variably segmented in agamids and uromastycids with cartilaginous myocommata. The muscle is thin but distinctly several fibers thick, the thickness in adults being 10 to 20% of the width of the muscle. The width is one third to one half of the distance measured from mid-sternum to the glenohumeral articulation. It is approximately equal in width throughout its length. Anteriorly the fibers are oriented at approximately a 40 degree angle with the adjacent fibers of the M. pectoralis. This muscle in *Physignathus cocincinus*, together with the M. pectoralis and M. rectus abdominis medialis, is illustrated in Figure 4. The configuration of this muscle slip is nearly identical in *Uromastix*, *Leiolepis*, and *Hydrosaurus*.

The skin overlying the M. pectoralis and M. rec. abd. medialis of all iguanians is attached to the musculature only by loose connective tissue and can be peeled away easily. This is in contrast with most autarchoglossan lizards in which the skin is firmly attached segmentally by both fibrous connective tissue and multiple muscular slips. In *Uromastix*, *Leiolepis*, *Hydrosaurus*, and *Physignathus*, only the anterior end of the muscle slip is firmly attached to the skin.

The M. rec. abd. medialis is wide and moderately thick in agamids, uromastycids, iguanids, and oplurines, but considerably reduced in the compressed chamaeleonids. It extends from the anterior margin of the pubis and extends broadly to the xiphisternum. It is comprised of longitudinal fibers in a single muscle layer which may be variously segmented by transverse cartilaginous myocommata in the agamids and uromastycids and by both myocommata and inscriptional ribs (sometimes but not always congruent) in the iguanids, oplurines, and chamaeleonids. Muscle slips do not insert into the overlying ventral skin at regular intervals. Laterally this

Figure 4. Anteroventral view of the superficial trunk and thoracic musculature of *Physignathus cocincinus* (Agamidae). The M. pectoralis pars cutaneous was left adherent to the reflected skin on the right side. This muscle slip was dissected from the skin on the left side and oriented in correct anatomical position alongside the M. pectoralis pars humeralis.

Abbreviations: abd med, M. rectus abdominis medialis; pect cut, M. pectoralis pars cutaneous; pect hum, M. pectoralis pars humeralis.

muscle abuts the M. obliquus abdominis externus within the same fascial layer. The M. obliquus abdominis externus does not have separate superficial and deep layers, except that some fibers overlap superficial and some deep to the abutting M. rectus abdominis medialis.

The only resemblances between the muscle slip overlying the M. pectoralis in *Uromastix*, *Leiolepis*, *Physignathus*, and *Hydrosaurus* and the M. rec. abd. lateralis of autarchoglossan lizards are: 1) inserts into the skin; 2) overlies the M. pectoralis; and 3) some fibers are continuous with the M. rec. abd. medialis. However, these resemblances must be qualified and the differences enumerated.

First, only the anterior end terminates

in an anastomosis with the skin. This muscle does not have segmentally arranged slips of fibers inserting into the skin as in Autarchoglossa. Second, this muscle overlies only the M. pectoralis and does not overlie any part of the M. rec. abd. medialis or M. obliquus abd. externus as in the Autarchoglossa. Third, although it is true that some fibers of this muscle are continuous with the anterolateralmost fibers of the M. rec. abd. medialis, most are continuous with the posterolateralmost fibers of the M. pectoralis. Moreover, in both the Ascalabota and Autarchoglossa the M. pectoralis, M. rec. abd. medialis, and M. obliquus abd. externus imperceptively merge at their common intersection. Fourth, in these primitive iguanians, this muscle is narrow and located laterally. In the Autarchoglossa, it is usually very wide and completely covers the M. pectoralis, and may extend to the throat.

Structurally, this muscle appears to be simply a small lateral slip, surrounded with epimysium, that has detached from the main part of the M. pectoralis and has inserted into the skin just ventromedial to the glenohumeral joint. This muscle is lacking in three of six *Uromastix hardwicki* specimens examined, and presumably develops late into the adult condition. *Uromastix asmussi* and *U. loricatus* usually lack this muscle, but in *U. asmussi* (ZFMK 7925) a thickened band of connective tissue with a few detectable muscle fibers was found. *U. loricatus* (CAS 86468) has a small slip of muscle identical to that found in other species, but six other specimens lacked the muscle. In one specimen of *U. acanthinurus* (personal collection) an interesting developmental variation was discovered (Fig. 5). Approximately one quarter of the central fibers of the M. pectoralis inserted into the skin at the same location where a lateral slip of muscle would insert, and the lateral slip was missing. These structural variations suggest that this muscle slip is derived from the M. pectoralis.

Figure 5. Anteroventral view of the superficial trunk and thoracic musculature of *Uromastix acanthinurus* (Uromastycidae). Only in this specimen was the M. pectoralis pars cutaneous found to arise from the middle of the body of M. pectoralis pars humeralis, rather than existing as a separate slip lateral to the main muscle body.

Abbreviations: see Figure 4.

Finally, several specimens were dissected in order to trace the innervation of the muscle slip in question. The nerve was found to be a branch of a large nerve trunk of the brachial plexus in the axilla, and the large nerve entered the M. pectoralis, in *Leiolepis belliana* (ZFMK 13156), *Uromastix ornatus* (ZFMK 8576), and *Physignathus cocininus* (ZFMK 21459). The latter specimen demonstrated best the ramification of the nerve into the muscle. Because the M. rectus abdominis is innervated by thoracic intercostal and lumbar nerves, innervation of this muscle by a branch of the nerve to the M. pectoralis from the brachial plexus clearly demonstrates correct homology with the M. pectoralis. I have named this muscle slip simply as the M. pectoralis pars cutaneous in contrast with

the principal muscle body M. pectoralis pars humeralis.

DISCUSSION

Camp's (1923) suborders (Autarchoglossa, M. rec. abd. lateralis present; Ascalabota, M. rec. abd. lateralis absent) have remained relatively unchanged since their establishment (Underwood, 1971) even though the Xantusiidae and Pygopodidae were respectively transferred from the Scincomorpha and Anguimorpha to the Gekkota (Kluge 1967, 1976). Recently, the suborder Ascalabota has been challenged as a polyphyletic taxon because of numerous convergent adaptations of the Iguania and Gekkota (Sukhanov 1961, 1976; Northcutt, 1978; Estes, 1983; Moody, in preparation). Even Camp indicated that the Gekkota were more closely related with the Autarchoglossa with his dendrogram (p. 333) but argued for the Ascalabota in his classification and character synopsis (pp. 296–298). A cladistic analysis of Camp's characters has demonstrated that the data available to him would support his dendrogram, not his classification (Moody, in preparation).

Sukhanov (1961, 1976) was the first to rigorously point out the inconsistencies between classification and dendrogram in Camp (1923). He also argued that the Gekkota and Scinco-anguimorpha are sister taxa on the basis of both traditional morphological characters and his observations on the musculature of the pectoral girdle and forelimb. Similar phylogenetic hypotheses have been suggested by Estes (1983) and Presch (personal communication). Northcutt (1978) also derives the Gekkota from the ancestral lineage of Scincomorpha and Anguimorpha and does not hypothesize a close relationship with the Iguania.

Camp (p. 420) argued that the Autarchoglossa was primitive and that the M. rec. abd. lateralis would be present in primitive lizards because of their life style of

terrestrial and burrowing habits, i.e., creeping, crawling, and slinking. He then argued that the Ascalabota were advanced lizards because of their life style of arboreality and saxicolity, i.e., running, jumping, and climbing. I would suggest that the prejudice of progressive evolution may have been involved unwittingly in Camp's evaluation, i.e., advanced lizards would resemble primates. Support for this suggestion comes from the fact that if one examines Camp's predictions for direction of character state evolution separately for his many characters, in a majority of cases he hypothesizes that the state present in the Ascalabota is primitive. But when he summarizes the overall evolution of the lizards, they become advanced (Moody, in preparation). According to Camp, the Gekkota and Iguania shared an ascalabotan ancestor which had lost the M. rec. abd. lateralis; however, the muscle was somehow retained in *Uromastyx*, *Leiolepis*, and *Physignathus*. Camp used the presence or absence of this muscle as the primary diagnostic character for classifying lizards, yet he ranked the muscle 22nd out of 34 evaluated characters for its value as a conservatively predictive phylogenetic character, or in his terminology, its "paleotelic value." Perhaps this inconsistency gave rise to the contradiction between his classification and his dendrogram.

If Camp's classification is followed, then the M. rec. abd. lateralis must be lost at least six times. Once in the Gekkota because the primitive Xantusiidae possesses the muscle (Kluge, 1976). Because primitive agamids and uromastycids possess the muscle, this requires a second loss for the Iguanidae, a third loss for the Chamaeleonidae and a fourth loss for the advanced Agamidae. Because two species of *Uromastyx* lack the muscle, a fifth loss can be argued. Finally, a sixth loss is for *Hydrosaurus*. That the losses in the Iguanidae, Chamaeleonidae and Agamidae are all independent is clear because Camp (pp. 333, 417) believed

that the Agamidae (and Uromastycidae) evolved from within the necessarily paraphyletic Iguanidae.

The discovery with this paper that the observed muscle in *Uromastyx*, *Leiolepis*, *Physignathus*, and *Hydrosaurus* is not homologous with the M. rec. abd. lateralis of the Autarchoglossa and Xantusiidae solves the problem of too many evolutionary losses. If the Gekkota are more closely related to the Scincomorpha and Anguimorpha because Xantusiidae possesses the M. rec. abd. lateralis, then only one loss must be hypothesized. If the Iguania is a relatively primitive lizard group, then they and ancestral lacertilians lacked the M. rec. abd. lateralis. In the common ancestor of uromastycids and primitive agamids, the M. pectoralis pars cutaneous evolved, but was once again lost in the advanced agamids, two species of *Uromastyx*, and *Hydrosaurus*.

This reanalysis of the M. rectus abdominis complex in the Lacertilia, corrections of its terminology and homology, and brief review of alternative phylogenetic hypothesis differing from Camp's classification, should encourage systematic herpetologists to abandon his classification, but accept his dendrogram except for minor shifts of families already discussed above. However, the barrier of tradition and expert opinion is often difficult to overcome as Rosen *et al.* (1981) have so eloquently demonstrated in their presentation of evidence that the lungfishes are the sister group of the tetrapods.

It is interesting that Camp (1923) discovered correctly the three iguanian genera possessing the M. pectoralis pars cutaneous, his vestigial M. rec. abd. "superficialis," not only because so few genera were examined (a probability of circa 1% using the hypergeometric distribution with nonreplacement sampling) but also because two of these genera, *Uromastyx* and *Physignathus*, as then recognized, were not monophyletic.

At the time of Camp's work, *Phy-*

signathus included three species from Australia (*gilberti*, *longirostris*, *temporalis*) together with the originally described *P. lesueurii* and *P. cocincinus*. However, the former three are unrelated to *Physignathus* and lack the compressed swimming tail, *M. pectoralis* pars cutaneous, large lacrimal, and cross-shaped interclavicle which *Physignathus* possesses. The scalation and body habitus of these three species relate them with the *Amphibolurus muricatus* species group and *Chlamydosaurus*. Cogger (1975) resurrected the genus *Lophognathus* for these species and Moody (1980) discussed numerous characters separating these genera.

Also at the time of Camp's work, *Aporoscelis* (Boulenger, 1885) included two species, *princeps* and *batilliferus*, and was considered a sister taxon of *Uromastix*. Anderson (1896: 34) recognized that: "*Uromastix batilliferus* Vaill. from its dentition and the form of the body, is unquestionably a member of the genus *Agama*, but with the tail of an *Uromastix*." His conclusions were not accepted until much later, when Parker (1942) reassigned *princeps* to *Uromastix* and placed *batilliferus* in the new genus *Xenagama*, though without discussing the morphological criteria for the reassignments. Parker's decision is substantiated because *U. princeps* possesses the specialized cranial, skeletal, dentitional, and scalational features characteristic for *Uromastix*, (Moody, 1980), as well as having the *M. pectoralis* pars cutaneous. *Xenagama batillifera* lacks the specialized skull and dentition and the cutaneous muscle. On the basis of scalation and skeleton, *Xenagama* and *Stellio* are sister genera and related to *Agama*. Although the tails of *Xenagama* have very large spiny tubercles, they are not similar to the tails of *Uromastix*.

The *M. pectoralis* pars cutaneous of the four iguanian genera is amazingly similar in structure and presumably in function to the thoracic-axillary portion of the *M. panniculus carnosus*, which is also de-

velopmentally related to the *M. pectoralis*, in mammals. Various cutaneous muscles are known in a few amphibians and reptiles (Romer, 1970). In humans, a slip of the *M. pectoralis* (embryonically part of the *M. panniculus carnosus*) fuses with the adjacent *M. rectus abdominis* and *M. obliquus abdominis externus*. It may insert in the brachial fascia independent of the main tendon of the *M. pectoralis* (Anson, 1966). In the rhesus monkey, the *M. panniculus carnosus* of the lateral thorax is remarkably similar to the *M. pectoralis* pars cutaneous, definitely effects movement of skin over the glenohumeral joint and helps the *M. pectoralis* in depressing the forelimb from an elevated position (Hartman and Straus, 1933).

ACKNOWLEDGMENTS

I appreciate the permission granted to me to make small incisions in specimens loaned to me by the following curators and collections: A. G. C. Grandison (BMNH), A. Leviton (CAS), J. Wright (LACM), H. Marx (FMNH), E. E. Williams (MCZ), A. E. Greer (MVZ), B. Lanza (MZUF), O. Rieppel (NHMB), T. Fritts (SDNHM), H. Wermuth (SMNS), A. G. Kluge (UMMZ), G. Zug (USNM), G. Storr (WAM), and W. Böhme (ZFMK). I especially thank Alan Leviton for rechecking my observations on *Uromastix loricatus* and Joseph Eastman for confirming other observations and for providing comparative information on mammals. Joseph Eastman, Arnold Kluge, and Wolfgang Böhme constructively criticized the manuscript. I thank Paris Mihely for preparing the figures.

Travel money to visit the USA herpetological collections was provided by a Rackham Graduate School Block Grant, University of Michigan. A fellowship from the Deutscher Akademischer Austauschdienst of Bonn, Germany, allowed me to visit all major herpetological collections in Europe. The College of

Osteopathic Medicine, Ohio University, provided funds for the final morphological comparisons and for the illustrations.

APPENDIX: SPECIMENS EXAMINED

Specimens examined are listed below in alphabetical order within each family. The museum collection acronyms follow the published list by the Joint Herpetological Resources Committee (Herpetological Review, 11(4), 1980: 93–102).

IGUANIDAE: *Amblyrhynchus cristatus*, FMNH 19752, UMMZ 50122; *Anisolepis undulatus*, MCZ 123812; *Anolis angusticeps*, UMMZ 115617; *A. bimaculatus*, UMMZ 60215; *A. carolinensis*, UMMZ 86466; *A. chrysolepis planiceps*, UMMZ 57704; *A. coelestinus*, UMMZ 110211; *A. cristatellus*, UMMZ 73703; *A. equestris*, UMMZ 58531; *A. fuscoauratus*, UMMZ 123913; *A. grahami*, UMMZ 85883; *A. latifrons*, UMMZ 48408; *A. onca*, UMMZ 54806; *A. petersi*, UMMZ 91202; *A. sagrei*, UMMZ 85888; *A. valencienni*, UMMZ 85913; *Aperoprists catamarcensis*, MCZ 96710; *Aptycholaemus longicauda*, BMNH 1946.8.9.2; *Basiliscus basiliscus*, UMMZ 78493; *B. galeritus*, UMMZ 51276; *Brachylophus fasciatus*, UMMZ 55772; *Callisaurus draconoides*, UMMZ 134018; *Chamaelinorops barbouri*, MCZ 146951; *Chamaeleolis chamaeleontides*, MCZ 38419; *C. porcus*, UMMZ 98014; *Conolophus subcristatus*, UMMZ 50121; *Cophosaurus texanus*, UMMZ 125303; *Corythophanes cristatus*, UMMZ 46446; *Crotaphytus collaris*, UMMZ 81931; *Ctenoblepharis adpersus*, MCZ 145040; *Ctenosaura pectinata*, UMMZ 118975; *Cyclura cornuta stejnegeri*, UMMZ 73899; *Diplolaemus bibroni*, MCZ 14918; *Dipsosaurus dorsalis*, UMMZ 105886; *Enyaliodes heterolepis*, MCZ 112347, MCZ 6983; *E. l. laticeps*, MCZ 12438; *E. laticeps festae*, MCZ 43752; *E. oshaughnessyi*, MCZ 29297; *E. palpebralis*, MCZ 84035; *Enyaliosaurus quinquecarinatus*, UMMZ 82376; *Enyalius bilineatus*, UMMZ 65946; *E. brasiliensis*, UMMZ 108627; *E. iheringii*, UMMZ 108629; *Gambelia wislizenii*, UMMZ 61997; *Holbrookia maculata*, UMMZ 122308; *Hoplacercus spinosus*, UMMZ 108259; *Iguana iguana*, UMMZ 98389; *Laemactus serratus*, UMMZ 110779; *Leiocephalus carinatus*, UMMZ 85927; *Leiosaurus belli*, UMMZ 89655; *Liolaemus alticolor walkeri*, UMMZ 133307; *Morunasaurus annularis*, USNM 6275; *M. groi*, MCZ 34875; *Ophryoesoides ornatus*, UMMZ 90802; *Petrosaurus mearnsi*, MCZ 100068; *Phenacosaurus richteri*, UMMZ 65203; *Phrynosoma cornutum*, UMMZ 136151; *Phymaturus palluma*, MCZ 2033; *Platynotus semi-taeniatus*, MCZ 126647; *Plica plica*, UMMZ 84744;

Polychrus acutirostris, UMMZ 68076; *P. peruvianus*, UMMZ 107615; *Pristidactylus scapulatus*, MCZ 33584; *Proctotretus azureus*, MCZ 133257; *P. pectinatus*, MCZ 66989; *Sator angustus*, UMMZ 124060; *Sauromalus obesus*, UMMZ 90706; *Sceloporus formosus*, UMMZ 118798; *S. graciosus*, UMMZ 70585; *S. grammicus disparilis*, UMMZ 79442; *S. horridus albiventris*, UMMZ 101975; *S. poinsetti*, UMMZ 118930; *S. scalaris*, UMMZ 71451; *S. undulatus*, UMMZ 97528; *S. utiformis*, UMMZ 115571; *S. variabilis*, UMMZ 105424; *Stenocercus guentheri*, MCZ 146926; *Strobilurus torquatus*, UMMZ 115646; *Tropidurus peruvianus*, UMMZ 83102; *Uma notata*, UMMZ 67367; *Uracentron flaviceps*, MCZ 37270; *Uranoscodon supercilialis*, MCZ 123748; *Urosaurus bicarinatus*, UMMZ 115037; *Urostrophus torquatus*, MCZ 123750; *U. vautieri*, MCZ 132256; *Uta stansburiana*, UMMZ 76480. OPLURINAE: *Chalarodon madagascariensis*, MCZ 11507, 11538; *Oplurus cyclurus*, MCZ 11718, 37189; *O. fierinensis*, MCZ 11627; *O. grandidieri*, MCZ 7735; *O. quadrimaculatus*, MCZ 67945; *O. saxicola*, MCZ 49659; CHAMAELEONIDAE: *Bradypodion pumilus*, UMMZ 60476; *B. ventralis*, UMMZ 61531; *Brookesia stumpffi*, UMMZ 65554; *B. supercilialis*, UMMZ 65555; *Chamaeleo africanus*, UMMZ 113491; *C. bitaeniatus*, UMMZ 70271; *C. brevicornis*, UMMZ 59763; *C. chamaeleon*, UMMZ 113493–4; *C. fallax*, UMMZ 65571; *C. gracilis*, UMMZ 116897; *C. jacksonii*, UMMZ 61184; *C. lateralis*, UMMZ 71611; *C. namaquensis*, UMMZ 65566; *C. oustaleti*, UMMZ 59764; *C. owenii*, UMMZ 60477; *C. tigris*, UMMZ 65733; *C. verrucosus*, UMMZ 65746; *Rhampholeon boulengeri*, UMMZ 60479; *R. spectrum*, UMMZ 35619.

AGAMIDAE: *Acanthosaura armata*, CAS 73721; *A. crucigera*, MVZ 79065; *A. lepidogaster*, MVZ 23431; *Agama (Agama) agama*, MVZ 97473; *A. (A.) anchietae*, MVZ 75527; *A. (A.) atra*, MVZ 69241; *A. (A.) caudospinosa*, MCZ 34957; *A. (A.) doriae*, MCZ 15829; *A. (A.) hartmanni*, MCZ 8809; *A. (A.) hispida*, MVZ 69252–53; *A. (A.) kirkii*, MCZ 67813; *A. (A.) mossambica*, MCZ 50192; *A. (A.) planiceps*, MVZ 9810; *A. (A.) rueppelli*, MCZ 49120; *A. (A.) sankaranica*, MVZ 75522; *A. (A.) weidholzi*, MCZ 44311; *Amphibolurus barbatus*, CAS 77596; *A. caudicinctus*, LACM 55105, 55117; *A. clayi*, LACM 55023, 55046; *A. cristatus*, LACM 55179, 55182; *A. diemensis*, MVZ 81662; *A. femoralis*, MCZ 86572; *A. fionii*, MCZ 10187; *A. fordii*, MVZ 81669; *A. inermis*, CAS 84102; *A. isolepis*, MVZ 81666; *A. maculatus*, MVZ 77631; *A. microlepidotus*, CAS 94360; *A. minimus*, MCZ 32969; *A. minor*, CAS 16214; *A. muricatus*, CAS 34024, 77645; *A. nobbi*, CAS 16215; *A. ornatus*, LACM 55030; *A. pictus*, CAS 16215; *A. reticulatus*, MVZ 78129; *A. rufescens*, MCZ 38595; *A. scutulatus*, LACM 55011, 55013; *Aphaniotis acutirostris*, CAS 64262, *A. fusca*, MCZ 38973; *Brachysaura minor*, MCZ 7211; *Bronchocela cristatellus*, CAS 8447, 8470, 60546; *B. jubatus*, CAS 55125–26; *B. marmoratus*, CAS

- 15457; *Caimanops amphiboluroides*, LACM 55285; *Calotes calotes*, MCZ 39828; *C. elliotti*, MCZ 6207; *C. emma*, MCZ 45268; *C. jerdoni*, MCZ 7204; *C. maria*, MCZ 7203; *C. mystaceus*, CAS 73730; *C. nigrilabris*, MCZ 4119; *C. rouxii*, CAS 104298; *C. versicolor*, CAS 18084, 98885; *Ceratophora stoddartii*, SDNHM 25450, MCZ 116378; *C. tennentii*, MCZ 4126; *Chelosania brunnea*, WAM 11808, 11810; *Chlamydosaurus kingii*, MVZ 77635; *Cophotis ceylanica*, BMNH 72.1.26.7, MCZ 118596; *Corytophyllax subcristatus*, MCZ 27094; *Dendragama boulengeri*, FMNH 109794; *Diporiphora australis*, MVZ 74732; *D. bilineata*, MVZ 81663; *D. winneckei*, LACM 55329, 55331; *Draco blanfordii*, MCZ 39089; *D. fimbriatus*, MCZ 39095; *D. formosus*, MVZ 145268; *D. lineatus beccarii*, MCZ 25354; *D. maculatus*, MVZ 23401; *D. maximus*, MCZ 39100; *D. quinquefasciatus*, MCZ 39126; *D. spilopterus*, MCZ 44141; *D. taeniopterus*, MVZ 15980; *D. volans*, MCZ 79697; *Gonocephalus bellii*, MCZ 15176; *G. beyschlagi*, MCZ 25911; *G. chamaeleontinus*, MCZ 27008; *G. grandis*, MVZ 111826; *G. interruptus*, CAS 25858; *G. liogaster*, MVZ 111835; *G. miotympanum*, MCZ 11265; *G. robinsonii*, MCZ 39172; *G. semperi*, CAS 23551; *G. sophiae*, CAS 19870; *Hydrosaurus amboinensis*, SDNHM 25983, MCZ 9329; *H. pustulatus*, CAS 24607, 28171; *H. weberi*, ZFMK 13993, MCZ 4574; *Hylagama borneensis*, FMNH 131540, 138196; *Hypsilurus (Hypsilurus) binotatus*, MCZ 130351, 137283; *H. (H.) boydii*, MVZ 77648; *H. (H.) bruijnii*, MCZ 126436; *H. (H.) dilophus*, CAS 64251; *H. (H.) godeffroyi*, MVZ 74903; CAS 54718, MCZ 4706; *H. (H.) nigricularis*, CAS 135207; *H. (H.) papuensis*, MCZ 44182; *H. (H.) spinipes*, CAS 84031; *Hypsilurus (Arua) auritus*, MCZ 96203; *H. (Arua) modestus*, CAS 13552; *Japalura brevipes*, MVZ 111835; *J. dymondi*, CAS 63926; *J. flaviceps*, MCZ 12469; *J. planidorsata*, MCZ 7197; *J. polygonata*, CAS 16262; *J. splendida*, MCZ 20700; *J. swinhonis*, MVZ 23391; *J. tricarinata*, MCZ 58288; *J. variegata*, MCZ 7196; *J. yunnanensis*, CAS 31435; *Lophocalotes ludekingi*, MCZ 38974; *Lophognathus gilberti*, CAS 77637-8; *L. longirostris*, CAS 94365, MCZ 35196; *L. temporalis*, MVZ 81675, MCZ 22906, 137294; *Lyriocephalus scutatus*, MCZ 7155; *Moloch horridus*, MVZ 82832, MCZ 6980; *Oriocalotes paulus*, BMNH 70.11.29.43; *Otocryptis wiegmanni*, CAS 16872; *Phoxophrys borneensis*, MCZ 43487; *P. nigrilabris*, MCZ 9036; *Phrynocephalus arabicus*, MCZ 44759; *P. axillaris*, MCZ 7226; *P. clarkorum*, CAS 120236; *P. forsythii*, MCZ 22174; *P. guttatus*, CAS 99719; *P. helioscopus*, CAS 105806; *P. interscapularis*, CAS 120107; *P. luteoguttatus*, CAS 120004; *P. maculatus*, MCZ 7225; *P. mystaceus*, CAS 99716-7; *P. ornatus*, CAS 96273; *P. przewalskii*, MCZ 22177; *P. reticulatus*, CAS 99722; *P. rossikowi*, CAS 94490; *P. scutellatus*, CAS 97978; *P. versicolor*, CAS 99721; *P. vlangalii*, MCZ 22181; *Physignathus cocincinus*, MCZ 20285, 39180, 39182; ZFMK 21459; *P. lesueurii*, UMMZ 65490, CAS 77644, 77647, MCZ 10176, 35212; *Psammophilus blanfordianus*, CAS 94353, 104104; *P. dorsalis*, MCZ 7209; *Pseudocalotes floweri*, MCZ 39068; *P. tympanistriga*, MCZ 7647, 22160; *Ptycolaelamus gularis*, BMNH 1940.6.3.71-2; *Salea anamallayana*, CAS 104246-7; *Salea horsfieldii*, CAS 94351, MCZ 7198; *S. kakhienensis*, MCZ 43055; *Sitana ponticeriana*, MCZ 7153; *Stellio agororensis*, MCZ 7217; *S. atricollis*, MCZ 18590; *S. caucasica*, CAS 120747; *S. cyanogaster*, MCZ 50551; *S. erythrogaster*, CAS 111930; *S. himalayana*, CAS 115934; *S. lehmanni*, CAS 94489; *S. melanura*, MCZ 50192; *S. nupta*, CAS 86332; *S. stellio*, MVZ 97851, 97853; *S. stoliczkan*, CAS 99713; *S. tarimensis*, MCZ 33577; *S. tuberculata*, CAS 90711; *Trapelus (Trapelus) agilis*, MCZ 45627; *T. (T.) flavimaculata*, CAS 97587; *T. (T.) impalearis*, MCZ 29957; *T. (T.) mutabilis*, MVZ 97477, 97480; *T. (T.) persicus*, CAS 84611; *T. (T.) rubrigularis*, MCZ 7213; *T. (T.) rudrata*, MCZ 9810; *T. (T.) savignii*, MVZ 97482-83; *Trapelus (Pseudotrapelus) sinaita*, MCZ 103840; *Tympanocryptis lineata*, MVZ 81663; *T. tetraporophora*, UMMZ 132278; *Xenagama batillifera*, MCZ 49121; *X. taylori*, MCZ 49215-6. Also a new genus is being described for "*Cophotis*" *sumatrana*, ZFMK 20790.
- UROMASTYCIDAE: *Leiolepis b. belliana*, ZFMK 13156, MCZ 122289; *L. belliana ocellata*, MCZ 43088; *L. guttata*, MCZ 16656; *L. peguensis*, BMNH 74.4.29.624; *L. r. reevesii*, MCZ 25185; *L. reevesii rubritaeniata*, MCZ 74102; *L. triploida*, MCZ 103696-7; *Uromastix a. acanthinurus*, CAS 123485, 123489; *U. acanthinurus werneri*, MCZ 27384; *U. aegyptius*, MVZ 97471, ZFMK 2704; *U. asmussi*, NHMB 14352, BMNH 74.11.23.9, ZFMK 7925; *U. benti*, BMNH 99.12.13.51, 1938.2.1.47; *U. geyri*, BMNH 1932.9.10.15, CAS 103072; *U. hardwickii*, CAS 99787, ZFMK 8616-7; *U. loricatus*, NHMB 3491, BMNH 1933.4.1.25, SMNS 607, 609, CAS 86468, 86470, 102480; *U. macfadyeni*, BMNH 1937.12.5.117, BMNH 1937.12.5.120, 1946.8.14.54; *U. microlepis*, ZFMK 20267, BMNH 1920.3.20.2, 1971.7.49; *U. ocellatus*, BMNH 97.10.28.210, 1937.8.38, SMF 10402; *U. ornatus*, SMF 10404, ZFMK 8576, MVZ 97472; *U. philbyi*, MCZ 44760, BMNH 1953.1.8.52; *U. p. princeps*, MCZ 70621, BMNH 1931.7.20.276; *U. princeps scortecci*, MZUF 113; *U. thomasi*, BMNH 1956.1.16.8, 1971.1355.

LITERATURE CITED

- ANDERSON, J. 1896. *Herpetology of Arabia*. London, 122 pp.
- ANSON, B. J. 1966. Editor, Morris' Human Anatomy. 12th edition. McGraw-Hill Co.
- BOULENGER, G. A. 1885. *Catalogue of the lizards in the British Museum (Natural History)*, Second Edition, Volume I. London, 436 pp.
- BYERLY, T. C. 1925. *The myology of Spheonodon*

- punctatum*. Univ. Iowa Studies Natural History, 11(6): 1-51, pl. i-iv.
- CAMP, C. L. 1923. Classification of the lizards. Bull. Amer. Mus. Nat. Hist., 48: 289-481, fig. A-L and 1-112.
- COCGER, H. G. 1975. Reptiles and Amphibians of Australia. Reed, London, 608 pp.
- DEVIS, C. W. 1884. Myology of *Chlamydosaurus kingi*. Proc. Linn. Soc. New South Wales, 8: 300-320.
- DONOSO-BARROS, R. 1973. Un nuevo saurio de Bolivia. Neotropica, 19(60): 132-134.
- DONOSO-BARROS, R., AND J. M. CEI. 1971. New lizards from the Volcanic Patagonian Plateau of Argentina. J. Herpetol., 5: 89-95.
- ESTES, R. 1983. The fossil record and early distribution of lizards, pp. 365-398. In Anders G. J. Rhodin and Kenneth Miyata (eds), Advances in Herpetology and Evolutionary Biology: Essays on Honor of Ernest E. Williams. Cambridge, the Museum of Comparative Zoology, 1983.
- ETHERIDGE, R. 1959. The relationships of the Anoles (Reptilia: Sauria: Iguanidae): An interpretation based on skeletal morphology. Doctoral Dissertation. University of Michigan at Ann Arbor, 236 pp.
- FÜRBRINGER, M. 1869. Die Knochen und Muskeln der Extremitäten bei den schlangenähnlichen Sauriern. Inaug. Dissert. Berlin. Wilh. Engelmann, Leipzig, pp. 1-58.
- . 1875. Zur vergleichenden Anatomie der Schultermuskeln. Vol. III, Chap. IV, Saurier und Crocodile. Morphol. Jahrb., 1: 636-816, pls. xxiii-xxvii.
- . 1900. Zur vergleichenden Anatomie des Brustschulterapparates und der Schultermuskeln. Jena. Zeitschr., 34: 215-718, pls. xiii-xvii.
- GADOW, H. 1882. Untersuchungen über die Bauchmuskeln der Krododile, Eidechsen und Schildkröten. Morphol. Jahrb., 7: 57-100, pl. i.
- GORSKI, C. 1852. Über das Becken der Saurier. Inaug. Dissert. Dorpat.
- HARTMAN, C. G., AND W. L. STRAUS, JR. 1933. The anatomy of the Rhesus Monkey. New York, Hafner Co.
- HILLENIUS, D. 1959. The differentiation within the genus *Chamaeleo* Laurenti, 1768. Beaufortia, 8(89): 1-92.
- HOFFMAN, C. K. 1890. Dr. H. G. Bronn's Klassen und Ordnungen des Thierreichs, wissenschaftlich dargestellt in Wort und Bild. Band VI, Abt. III. Reptilien (Eidechsen II). Leipzig, 443 and 1399 pp.
- KLUGE, A. G. 1967. Higher taxonomic categories of gekkonid lizards and their evolution. Bull. Amer. Mus. Nat. Hist., 135: 1-60.
- . 1976. A reinvestigation of the abdominal musculature of Gekkonoid lizards and its bearing on their phylogenetic relationships. Herpetologica, 32: 295-298.
- MAURER, F. 1896. Die ventrale Rumpfmuskulatur einiger Reptilien. Eine vergleichend-anatomische Untersuchung. Festschrift für Gegenbaur, 1: 183-256, pls. i-iv.
- . 1898. Die Entwicklung der ventralen Rumpfmuskulatur bei Reptilien. Morphol. Jahrb., 26: 1-60, pls. i-iii.
- MIVART, ST. G. 1867. Notes on the myology of *Iguana tuberculata*. Proc. Zool. Soc. Lon.; 1867: 766-797.
- . 1870. On the myology of *Chamaeleo parsonii*. Proc. Zool. Soc. London, 1870: 850-890.
- MOODY, S. M. 1980. Phylogenetic and historical biogeographical relationships of the genera in the family Agamidae (Reptilia: Lacertilia). Doctoral Dissertation, University of Michigan at Ann Arbor, xv + 373 pp.
- NORTHCUTT, R. G. 1978. Forebrain and midbrain organization in lizards and its phylogenetic significance. In, N. Greenberg and P. D. MacLean (eds.) Behavior and Neurology of Lizards. NIMH, Rockville, Maryland.
- PARKER, H. W. 1942. The lizards of British Somaliland. Bull. Mus. Comp. Zool., 91: 1-101.
- RAW, L. R. G. 1976. A survey of the dwarf chameleons of Natal, South Africa, with descriptions of three new species (Sauria: Chamaeleonidae) Durban Mus. Novit., 11: 139-161.
- ROBINSON, P. L. 1967. The evolution of the Lacertilia. Colloques int. Cent. natn. Rech. Scient. No. 163, pp. 395-407.
- ROMER, A. S. 1970. The vertebrate body, fourth ed. Philadelphia, Saunders Co.
- ROSEN, D., P. FOREY, B. GARDINER, AND C. PATTERSON. 1981. Lungfishes, Tetrapods, Paleontology, and Plesiomorphy. Bull. Amer. Mus. Nat. Hist., 167: 159-276.
- RÜDINGER. 1968. Die Muskeln der verdern Extremitäten der Reptilien und Vögel. Hollandsche. Maatsch. Wetensch. Haarlem, 4: 1-187.
- SANDERS, A. 1870. Notes on the myology of *Platy-dactylus japonicus*. Proc. Zool. Soc. London, 1870: 413-426.
- . 1872. Notes on the myology of *Liolepis belli*. Proc. Zool. Soc. London, 1872: 154-183.
- . 1874. Notes on the myology of *Phrynosoma coronatum*. Proc. Zool. Soc., London, 1874: 71-89.
- SHUFELDT, R. W. 1890. Contributions to the study of *Heloderma suspectum*. Proc. Zool. Soc. London, 1890: 148-244.
- SMITH, H. M. 1939. The Mexican and Central American lizards of the genus *Sceloporus*. Fieldiana, Zoology, 26: 1-397, 31 pls.
- SMITH, M. A. 1935. The Fauna of British India, including Ceylon and Burma. Reptilia and Amphibia. Vol. II. Sauria. London, Taylor and Francis, Ltd., 440 pp.
- STANNIUS, H. 1856. Handbuch der Anatomie der Wirbelthiere. In Siebold und Stannius, Handbuch der Zoötomie, Part II, Die Wirbelthiere, 2nd ed. Berlin, 270 pp.

- SUKHANOV, V. B. 1961. Nekotorye voprosy filogenii i sistemy Lacertilia (sue Sauria). Zool. Zhurnal 60(1): 73-83. (In Russian).
- . 1976. Some problems of the phylogeny and systematics of the Lacertilia. Translation of the Russian original. S. Moody and G. Zug, eds. Smithsonian Herpetological Information Service No. 38, pp. 1-15.
- UNDERWOOD, G. 1971. A modern appreciation of Camp's "Classification of the lizards," pp. vii-xvii. In Camp's classification of lizards. Soc. Stud. Amphib. Rept. Fascimilae Reprint Series.
- WERMUTH, H. 1966. Liste der rezenten Amphibien und Reptilien: Chamaeleonidae. Das Tierreich (Berlin), 83: i-x, 1-37.
- . 1967. Liste der rezenten Amphibien und Reptilien: Agamidae. Das Tierreich (Berlin), 86: i-xiv, 1-127.

On the Adaptive Significance of the Reptilian Spectacle: the Evidence from Scincid, Teiid, And Lacertid Lizards

ALLEN E. GREER¹

ABSTRACT. In scincid, teiid, and lacertid lizards the spectacle, a permanent transparent scale covering the eye, is associated with species of small size and diurnal habits inhabiting relatively dry habitats. Presumably the spectacle reduces evaporative water loss from the eye. A similar morphology and ecology may have been associated with the evolution of the spectacle in other squamates (lizards and snakes). The morphology of the squamate eyelid may be tracking, evolutionarily, the interplay between body size and habitat.

INTRODUCTION

Many squamate reptiles (lizards and snakes) have a clear, disc-shaped scale permanently covering the eye. This structure, known as a spectacle or brille, appears to have evolved at least eight times in scincids (see below) and at least once in each of the following six taxa—teiids (*Gymnophthalmus* and *Micrablepharus*), lacertids (*Ophisops*), gekkonids and pygopodids, xantusiids, amphisbaenians, and snakes. The spectacle has always been assumed to have great systematic importance in squamates and in the past was the hallmark for vast polyphyletic assemblages, e.g., the subfamily “Ophiophthalmes” (Duméril and Bibron, 1836) and the genus *Ablepharus* (Boulenger, 1887).

Comparative morphological (Bellairs and Boyd, 1948; Fuhn, 1969; Greer,

1974; Smith, 1935) and embryological (Schwartz-Karsten, 1933; Neher, 1935) studies suggest that the spectacle probably evolved in the following sequence: 1) the lower eyelid moveable and entirely covered by small scales, 2) the eyelid moveable but with either a few central transparent scales or “windows” or only a single large scale or “window” that fills most of the eyelid, and 3) the large, clear eyelid fused in the raised position to give the spectacle² (Fig. 1).

Four hypotheses have been advanced to explain the adaptive significance of the spectacle or its precursor, the windowed eyelid. 1) The spectacle has no adaptive significance on the evidence of its having no discernable ecological correlates (Smith, 1935, 1939; Bellairs, 1970). 2) The spectacle protects the eye from physical damage by foreign bodies on the evidence of a supposed correlation with deserticolous (wind-blown sand), subterranean (soil particles) or nocturnal (reduced visual acuity) habits (Walls, 1934, 1942). 3) The windowed eyelid provides facultative protection from

¹The Australian Museum, Sydney, New South Wales, 2000 Australia.

²A fourth condition which need not concern us here may occur in extreme burrowers: the eye is vestigial and the “eyelid” is a covering scale similar to the surrounding head scales. Examples are most amphisbaenians, dibamids, a few scincids and scolecophidians. Little work has been done on the precursors of this condition, but my own experience, mainly with scincids, suggests that it can evolve from any of the other three conditions.

Figure 1. Close-up photographs of the heads of three different scincid lizards to show the three main structural stages in the evolution of the spectacle in lizards and snakes. Bottom: lower eyelid moveable (shown here raised) and scaly (*Sphenomorphus tympanum*); middle: lower eyelid moveable (shown here raised) but with a clear "window" (*Carlia amax*), and top: the large clear spectacle, permanently covering the eye (*Cryptoblepharus plagiocephalus*).

lids (as opposed to scaly eyelids)—and also with some pigment in the eyelid—are suspected of having crepuscular, cave-dwelling habits and are inferred to protect their presumably light sensitive eyes by raising their eyelids in daylight (Williams and Hecht, 1955). 4) Both the windowed eyelid and the spectacle reduce evaporative water loss from the eye (Arnold, 1973) on the evidence of a) an experimental observation that corneal transpiration may account for over 20% of total transpirational loss in a small lizard at normal activity temperatures (Reichling, 1957; also see Mautz, 1980) and b) anecdotal but informed observation that both eyelid types are widespread in small, sun-dwelling lizards (Arnold, 1973).

The purpose of this paper is fourfold. First, to test a prediction of the evaporative water loss hypothesis using the three squamate families in which the spectacle occurs in lineages of low taxonomic rank and in which, therefore, its original adaptive associations may be especially clear. Second, to show that the evaporative water loss hypothesis either contradicts, or explains more parsimoniously, the observations advanced in support of the other hypotheses. Third, to suggest that the evaporative water loss hypothesis has heuristic value in thinking about the evolution of those groups in which the spectacle characterizes lineages of high taxonomic rank and in which, therefore, its original adaptive significance may be obscure. And fourth, to present a more general hypothesis about the evolution of all three major eyelid types in squamates.

RESULTS

The evaporative water loss hypothesis predicts that taxa with a spectacle will be as small or smaller than their nearest relatives with moveable eyelids and that they will occur in habitats that are as dry or drier. This prediction arises in part from the classic allometric relationship between eye size and body size: in a

strong solar radiation (Plate, 1924; Mertens, 1954) on the evidence that a) certain windowed-eyed lizards—and as it happens species that also have some pigmentation in the eyelid—have been observed to raise their eyelids in strong sunlight (Mertens, 1954) and b) the only two anoline lizards with windowed eye-

series of similarly shaped but different sized animals smaller forms have relatively larger eyes (Walls, 1942; Rensch, 1959; Gould, 1966). This allometric relationship has been verified for scincids. Regression analysis of a log-log plot of eye surface area versus body weight in 20 diurnal, terrestrial scincids represented by single adult individuals with complete tails resulted in a regression coefficient (0.27; 95% confidence interval = ± 0.09) well below even isometric conditions (0.68) under which, of course, eye surface area would still not increase as fast as body volume. Regression analysis was by Bartlett's method as described by Simpson, Roe, and Lewontin (1960).

The predicted morphological and ecological associations of the spectacle are especially clear in the eight scincid and single teiid and lacertid lineages in which this eyelid type has evolved (Fig. 2). These lineages are described briefly below.

1) The Australian scincid genera *Morethia* (8 species) and *Proablepharus* (3 species) both have a spectacle which they probably inherited from a common ancestor (Greer, 1980). Both genera are diurnal and terrestrial. *Morethia* occurs in open habitats throughout Australia except for the mesic southeast (Smyth, 1972; Storr, 1972; Greer, 1980), and *Proablepharus* occurs in open habitats in the arid and semi-arid interior and the seasonally dry north (Storr, 1975; Greer, 1980). The taxon most closely related to these two genera together is the *trilineatum* species group of *Leiopisma* (3 species). This group has windowed eyelids, diurnal and terrestrial habits and a disjunct distribution in the mesic southwest and southeast corners of Australia (Greer, 1980). *Morethia* ranges in size from 33 to 58 mm, *Proablepharus* from 32 to 45 mm and the *trilineatum* species group from 69 to 80 mm.¹ The two spec-

tacted genera are clearly both significantly smaller ($P < 0.001$) and more xeric adapted than their window-eyed relatives.

2) The Australian scincid genus *Lerista* (37 species) occurs throughout most of the continent but is centered over the arid and semi-arid interior and the seasonally dry north (Greer, 1967, 1979b; Storr, 1971, 1976c). The genus is cryptozoic and semifossorial. Nine species have a spectacle, and 27 have windowed eyelids (one species has vestigial eyes and is therefore excluded from the analysis). No attempt has been made to detect a habitat difference between the two groups, but in size the spectacled species are significantly smaller (30–61 mm) than their window-eyed relatives (51–104 mm) ($P < 0.001$). Storr (1971) clearly understood the relationship between the spectacle (his "ablepharine eye") and body size in *Lerista* when he said: "The smallest species . . . [of the *elegans* species group of *Lerista*] have acquired an ablepharine eye (i.e., the lower eyelid is immovable and entirely transparent), a condition for which there is evidently strong selection pressure in litter inhabiting skinks whose snout-vent length averages less than 40 mm."

3) The spectacled scincid genus *Cryptoblepharus* (37 species) occurs in two disjunct areas: the southwest Pacific including Australia and the western Indian Ocean (Mascarene Islands, Madagascar, and associated islands and the east African coast). In most areas it occurs only along open coasts (a dry habitat in terms of potable water) (Mertens, 1931, 1958, 1964; Brongersma, 1942; Canaris and Murphy, 1965), but in Australia it occurs in a variety of habitats throughout most of the arid, semi-arid, and seasonally dry

¹The measure of species size employed throughout is the maximum known snout-vent (SVL) length. Tables giving maximum SVL, sample size,

and sources of information for each species in every taxon discussed are available from the author and the Department of Herpetology, Museum of Comparative Zoology upon request. The distribution of species sizes between taxa were compared using the Mann-Whitney U Test as described by Siegel (1956). The level of significance throughout is 0.05.

Figure 2. Ranges and means of maximum snout-vent lengths for the species in the scincid (1–8), teiid (9), and lacertid (10) lineages discussed in the text. In each lineage taxa with a permanent spectacle covering the eye are indicated by a dashed line and their closest relatives with a moveable eyelid by a solid line. Numbers in parentheses are sample sizes. Numbers to the far right are P values (Mann-Whitney U Test) associated with the size differences between the species with the two different eyelid types in each lineage. Data for individual species are available from the author upon request.

areas (Storr, 1976a; Covacevich and Ingram, 1978). *Cryptoblepharus* is most closely related to the *cyanura* species group of the genus *Emoia* (7 species) which is widespread in the southwest Pacific but extends only to the tip of Cape York Peninsula in Australia (Greer, 1974). It is diurnal, terrestrial to semi-arboreal, and occurs in habitats ranging from moist, closed forest to open coasts (Webster, 1969; McCoy, 1980). *Cryptoblepharus* is significantly smaller than the *cyanura* species group of *Emoia* (34–54 mm versus 45–61 mm; $P < < 0.001$) and appears to occur in habitats that are as dry (e.g., open coasts) or drier (e.g., the interior of Australia).

4) The spectacled, Australian scincid genus *Notoscincus* (2 species) is diurnal

to crepuscular, terrestrial, and distributed over the “less humid parts of northern Australia” (Storr, 1974). The genus is a member of the largest of the three skink lineages in Australia (136 species out of a total of 242), but otherwise its relationships are obscure (Greer, 1979a). It is, however, the smallest taxon in its lineage in Australia (36–41 mm).

5) The *novaeguineae* species group of the scincid genus *Carlia* (5 species) is diurnal and ranges from southern New Guinea and extreme eastern Australia as far south as Sydney. The largest species (42 mm) occurs in moist closed forest to seasonally dry woodland and has a windowed eyelid. The other four species, all Australian, occur in seasonally dry open forest to open woodland. The second

Figure 3. Frequency distributions of maximum snout-vent lengths for all skink species with one of the three basic eyelid types: scaly, windowed, and spectacled.

largest of these (38 mm) has a windowed eyelid; the others have a spectacle. The *novaeaguineae* species group's nearest relative appears to be the diurnal, spectacled Australian genus *Menetia* (5 species) (Greer, 1974; Ingram, 1977) which occurs throughout the arid and semi-arid interior of Australia and the seasonally dry north (Storr, 1976b, 1978; Rankin, 1979) in shrubland and open forest. No *Menetia* exceeds 38 mm. Taking these two groups together, it is evident that there is no significant size difference between the spectacled and window-eyed species ($P < 0.07$) but the former occur in habitats that are generally drier.

6) The subsaharan African scincid subgenus *Panaspis* (6 species) is diurnal and

terrestrial. Five species have windowed eyelids; four of these occur in moist closed forest and the fifth occurs in wooded savanna (Fuhn, 1972; Perret, 1973, 1975). A sixth species has a spectacle and inhabits dense humid forest, mangrove forest, and savanna at the edge of forest (Fuhn, 1970). This latter species is structurally similar to the hypothetical ancestor of the spectacled, subsaharan African subgenus *Afroablepharus* (6 species) (Fuhn, 1970) which is diurnal and terrestrial and associated with savanna (Perret, 1975). The seven spectacle-eyed species in these two groups together are significantly smaller than the five window-eyed species (24–55 mm versus 43–68 mm; $P < 0.01$) and appear to

occur in generally drier habitats (savanna versus forest).

7) The spectacled, south African scincid genus *Typhlacontias* (4 species) is fossorial and occurs in the aeolian sands of the northern Namib Desert (3 species) and in a relatively moist area of the northern Kalahari (1 species) (D. G. Broadley, *in litt.*). *Typhlacontias* is most closely related to the south African genus *Scelotes* (15 species) (Greer, 1970) which has a moveable lower eyelid that is either completely scaled or with a window. The genus is cryptozoic to fossorial and is restricted to the more mesic southern and eastern parts of south Africa (D. G. Broadley, *in litt.*). The genera do not differ significantly in size (83–130 mm for *Typhlacontias* versus 36–120 mm for *Scelotes*; $P \gg 0.05$), but *Typhlacontias* appears to occur in generally drier habitats.

8) The spectacled scincid genus *Ablepharus* (5 species) is diurnal and terrestrial and distributed over southeast Europe and southwest Asia in Mediterranean, prairie, steppe, and desert biomes. *Ablepharus* is most closely related to the window-eyed genus *Scincella* (Greer, 1974) which is diurnal and terrestrial and distributed disjunctly in central and south Asia and southeast North America and Mexico in various forest biomes. In size *Ablepharus* does not differ significantly from neighboring central Asian *Scincella* (7 species) (36–60 mm versus 47–65 mm; $P > 0.05$), but on the basis of the biomes occupied, it appears to occur in generally drier habitats.

9) The only spectacled taxa among teiids are the tropical American genera *Gymnophthalmus* (6 species) and *Micrablepharus* (1 species). The relationships of *Micrablepharus* are poorly known but are thought to lie close to *Gymnophthalmus* (W. Presch, personal communication). The closest relatives of these two genera are the window-eyed, tropical American genera *Anotosaura* (3 species), *Bachia* (15 species), *Heterodactylus* (2 species), *Iphisa* (1 species),

and *Tretioscincus* (2 species) (Dixon, 1973, 1974; Presch, 1980). Both groups occur in a variety of habitats, but in size the spectacled taxa are significantly smaller (41–48 mm) than their window-eyed relatives (39–114 mm but only one species less than 50 mm) ($P < 0.001$).

10) The only spectacled taxon of lacerids is the diurnal, terrestrial genus *Ophisops* which ranges from west India through southwest Asia and southeast Europe to north Africa. *Ophisops* appears to be most closely related to the diurnal, terrestrial genus *Cabrita* which occurs in central and eastern India (Smith, 1935). The two genera do not differ significantly in size (36–65 mm for *Ophisops* versus 44–57 mm for *Cabrita*; $P \gg 0.05$), but it is significant that both genera are small and that the spectacled *Ophisops* ranges over more generally arid country.

DISCUSSION

Before commenting on the spectacle in other squamates, I critically review the three other hypotheses that have been advanced to explain the adaptive significance of the spectacle and, where appropriate, compare them to the evaporative water loss hypothesis. The hypothesis that the spectacle has no adaptive significance can be rejected, because, assertions to the contrary notwithstanding (Smith, 1935, 1939), there are clear ecological associations with the spectacle. The hypothesis that the spectacle protects the eye from physical damage (Walls, 1934, 1942) is unsatisfactory for two reasons. First, it offers three unrelated associations (burrowing, nocturnality, and life in desert environments) to explain the adaptive significance of the spectacle, whereas the evaporative water loss hypothesis suggests only two related associations. Second, two of its associations (burrowing and nocturnality) are not convincing as there are many exceptions (e.g., many burrowing and nocturnal skinks have no spectacles), and the

third association (life in desert environments) is accounted for by the evaporative water loss hypothesis. The hypothesis that the windowed eyelid protects the eye from solar radiation (Plate, 1924; Mertens, 1954; Williams and Hecht, 1955) can be criticized on the grounds that it confounds a modification that reduces the amount of solar radiation entering the eye, i.e., the eyelid pigment¹ with a modification that permits vision while the eyelid is temporarily closed to reduce evaporative water loss, i.e., the spectacle.² The two modifications are often associated but their roles are quite distinct.

Unfortunately, the morphological and ecological associations of the spectacle are not as apparent in other squamates as they are in scincids, teiids, and lacertids, perhaps because subsequent evolution has obscured them. The evidence on the evolution of the spectacle is so telling in the three lizard families discussed here, however, that it may be useful to begin using it in assessing the origin of the spectacle in other groups. This may be especially useful in those groups such as non-eublepharine geckos and snakes where the spectacle probably attended the evolution of the group and where there are several conflicting hypotheses as to the origin of the group. In such considerations, the two most important conclusions that can be drawn from the evidence provided by scincids, teiids, and lacertids are 1) the spectacle has evolved many different times and 2) it has evolved in relatively small forms (gener-

ally 65 mm or less in the lizards discussed here) living in relatively dry habitats.

Finally, it may be useful for the purposes of further testing to state the evaporative water loss hypothesis in its logically broadest terms: the morphology of the squamate eyelid (scaly, windowed, or spectacled), through the intermediate parameter of potential evaporative water loss from the surface of the eye, may be tracking, in an evolutionary sense, the interplay between body size (as it affects the eye surface area/body volume ratio) and habitat (both temporal and spatial). Hence, just as spectacles are generally associated with small size and xeric habitats, scaly eyelids should be associated with large size and mesic habitats, and windowed eyelids associated with intermediate size and intermediate (in terms of moisture) habitats. Indeed, that size alone accounts for much of the variation in eyelid type is evident from the fact that in skinks, the largest and most diverse family of lizards, the size distributions of the species with the three different eyelid types are in the order scaly > windowed > spectacled ($P < 0.001$ for each pairwise comparison; Kolmogorov-Smirnov Two Sample Test as described by Siegel [1956]; Fig. 3).

ACKNOWLEDGMENTS

I thank the following people for providing me with summaries of the ecology and/or size of groups about which I know relatively little: J. P. Bacon, D. G. Broadley, W. C. Brown, R. I. Crombie, J. R. Dixon, W. D. Haacke, M. S. Hoogmoed, A. G. Kluge, W. P. Maclean, J. L. Perret, W. Presch, J. P. Rosado, R. Sage, R. Thomas, P. E. Vanzolini, L. J. Vitt, and A. Schwartz. I also thank C. Gans, S. J. Gould, G. Underwood and E. E. Williams for critical readings of various drafts and P. Greer for preparing the graphs. Work for this report was supported in part by the Australian—

¹A study of eyelid pigmentation in lizards would be interesting in its own right. With regard to its possible function as a sun screen, I note here that window-eyed skinks living in alpine habitats in southeast Australia, e.g., *Leiopisma greeni* and *L. pretiosum*, have intensely pigmented eyelids.

²Note that small, window-eyed skinks almost invariably close their eyelids after a brief period when placed in an environment of relatively dry air but subdued light intensity as, for example, on an open desk in a dimly lit room.

American Educational Foundation,
Canberra, A.C.T. (Fulbright Program).

LITERATURE CITED

- ARNOLD, E. N. 1973. Relationships of the palaearctic lizards assigned to the genera *Lacerta*, *Algroides* and *Psammodromus* (Reptilia: Lacertidae). Bull. Br. Mus. nat. Hist., **25**(8): 289-366.
- BELLAIRS, A. D'A. 1970. The Life of Reptiles. New York, Universe Books, 2 vols, 590 pp.
- BELLAIRS, A. D'A., AND J. D. BOYD. 1948. The lacrimal apparatus in lizards and snakes.—I. The brille, the orbital glands, lachrymal canaliculi and origin of the lachrymal duct. Proc. zool. Soc. Lond., **117**: 81-108.
- BOULENGER, G. A. 1887. Catalogue of the Lizards in The British Museum (Natural History). Vol. 3. London, British Museum (Natural History), 575 pp.
- BRONGERSMA, L. D. 1942. Notes on scincid lizards. Zoöl. Meded. Leiden., **24**(1-2): 125-152.
- CANARIS, A. G., AND D. G. MURPHY. 1965. A scincid reptile feeding primarily on marine crustacea, with a note on its parasites. J. E. Africa nat. Hist. Soc., **25**(2): 129-130.
- COVACEVICH, J., AND G. J. INGRAM. 1978. An undescribed species of rock dwelling *Cryptoblepharus* (Lacertilia: Scincidae). Mem. Qd Mus., **18**(2): 151-154.
- DIXON, J. R. 1973. A systematic review of the teiid lizards, genus *Bachia*, with remarks on *Heterodactylus* and *Anotosaura*. Univ. Kans. Mus. Nat. Hist. Misc. Publ. No. 57, 47 pp.
- . 1974. Systematic review of the lizard genus *Anotosaura* (Teiidae). Herpetologica, **30**(1): 13-18.
- DUMÉRIL, A. M. C., AND G. BIBRON. 1836. *Erpétologie Générale ou Histoire Naturelle Complète des Reptiles*. Vol. 5. Paris, 854 pp.
- FUHN, I. E. 1969. The "polyphyletic" origin of the genus *Ablepharus* (Reptilia, Scincidae): a case of parallel evolution. Z. f. zool. Systematik u. Evolutionsforschung., **7**(1): 67-76.
- . 1970. Contribution à la systématique des lygosomines Africains (Reptilia, Scincidae). I. Les espèces attribuées au genre *Ablepharus*. Rev. Roum. Biol. -Zool., **15**(16): 379-393.
- . 1972. Révision du phylum forestier du genre *Panaspis* Cope (Reptilia, Scincidae, Lygosominae). Rev. Roum. Biol. -Zool. **17**(4): 257-271.
- GOULD, S. J. 1966. Allometry and size in ontogeny and phylogeny. Biol. Rev., **41**: 587-640.
- GREER, A. E. 1967. A new generic arrangement for some Australian scincid lizards. Breviora Mus. Comp. Zool. No. 267, 19 pp.
- . 1970. The systematics and evolution of the subsaharan Africa, Seychelles, and Mauritius scincine scincid lizards. Bull. Mus. comp. Zool., **140**(1): 1-23.
- . 1974. The generic relationships of the scincid lizard genus *Leiopisma* and its relatives. Aust. J. Zool., Suppl. Ser. No. 31, 67 pp.
- . 1979a. A phylogenetic subdivision of Australian skinks. Rec. Aust. Mus., **32**(8): 339-371.
- . 1979b. A new species of *Lerista* (Lacertilia: Scincidae) from northern Queensland, with remarks on the origin of the genus. Rec. Aust. Mus., **32**(10): 383-388.
- . 1980. A new species of *Morethia* (Lacertilia: Scincidae) from northern Australia, with comments on the biology and relationships of the genus. Rec. Aust. Mus., **33**(2): 89-122.
- INGRAM, G. J. 1977. Three species of small lizards—two of them new. Genus *Menetia* (Lacertilia: Scincidae) in Queensland. Victorian Nat., **94**(5): 184-187.
- MAUTZ, W. J. 1980. Factors influencing evaporative water loss in lizards. Comp. Biochem. Physiol., **67A**: 429-437.
- MCCOY, M. 1980. Reptiles of the Solomon Islands. Wau Ecology Inst. Handb. No. 7, Wau, Papua New Guinea, 80 pp.
- MERTENS, R. 1931. *Ablepharus boutonii* (Desjardin) und seine geographische Variation. Zool. Jb. (Syst.), **61**(1/2): 61-210.
- . 1954. Biologische Fahrten in Südwest-Afrika. 3. Im Gebiete des Brandberg. Natur u. Volk, **84**(6): 184-195.
- . 1958. Neue Eidechsen aus Australien. Senck. biol., **39**(1/2): 51-56.
- . 1964. Weitere Mitteilungen über die Rassen von *Ablepharus boutonii*, III. Zool. Anz., **173**(2): 99-110.
- NEHER, E. M. 1935. The origin of the brille in *Crotalus confluentus lutosus* (Great Basin Rattlesnake). Trans. Am. ophthal. Soc. **33**: 533-545.
- PERRET, J. L. 1973. Contribution à l'étude des *Panaspis* (Reptilia, Scincidae) d'Afrique occidentale avec la description de deux espèces nouvelles. Rev. Suisse Zool., **80**(2): 595-630.
- . 1975. La différenciation dans le genre *Panaspis* Cope (Reptilia, Scincidae). Bull. Soc. Neuchâtel. Sci. Natur. **98**: 5-16.
- PLATE, L. 1934. Allgemeine Zoologie und Abstammungslehre. Vol. 2. Jena. (Not seen by me; cited by Williams and Hecht, 1955.)
- PRESCH, W. 1980. Evolutionary history of the South American microteiid lizards (Teiidae: Gymnophthalminae). Copeia, **1980**(1): 36-56.
- RANKIN, P. R. 1979. A taxonomic revision of the genus *Menetia* (Lacertilia, Scincidae) in the Northern Territory. Rec. Aust. Mus., **32**(14): 491-499.
- REICHLING, H. 1957. Transpiration und Vorzugstemperatur mitteleuropäischer Reptilien und Amphibien. Zool. Jb. (Zool.), **67**: 1-64.
- RENSCH, B. 1959. Evolution above the Species Level. London, Methuen, 419 pp.
- SCHWARTZ-KARSTEN, H. 1933. Über Entwicklung und Bau der Brille bei Ophidiern und Lacertiliern und die Anatomie ihrer Tränenwege. Morph. Jb., **72**: 499-540.

- SIEGEL, S. 1956. Nonparametric Statistics for the Behavioral Sciences. New York, McGraw-Hill Book Co., 312 pp.
- SIMPSON, G. G., A. ROE, AND R. C. LEWONTIN. 1960. Quantitative Zoology. 2nd ed. New York; Harcourt, Brace and World, Inc., 440 pp.
- SMITH, M. A. 1935. The Fauna of British India. Reptilia and Amphibia. Vol. II.—Sauria. London, Taylor and Francis, 440 pp.
- . 1939. Evolutionary changes in the eye coverings of certain lizards. Proc. Linn. Soc. Lond., 1939: 190–191.
- SMYTH, M. 1972. The genus *Morethia* (Lacertilia, Scincidae) in South Australia. Rec. S. Aust. Mus., 16(12): 1–14.
- STORR, G. M. 1971. The genus *Lerista* (Lacertilia, Scincidae) in Western Australia. J. Proc. Roy. Soc. West. Aust., 54(3): 59–75.
- . 1972. The genus *Morethia* (Lacertilia, Scincidae) in Western Australia. J. Proc. Roy. Soc. West. Aust., 55(3): 73–79.
- . 1974. The genus *Notoscincus* (Lacertilia, Scincidae) in Western Australia and Northern Territory. Rec. West. Aust. Mus., 3(2): 111–114.
- . 1975. The genus *Proablepharus* (Scincidae, Lacertilia) in Western Australia. Rec. West. Aust. Mus., 3(4): 335–338.
- . 1976a. The genus *Cryptoblepharus* (Lacertilia, Scincidae) in Western Australia. Rec. West. Aust. Mus., 4(1): 53–63.
- . 1976b. The genus *Menetia* (Lacertilia, Scincidae) in Western Australia. Rec. West. Aust. Mus., 4(2): 189–200.
- . 1976c. Revisionary notes on the *Lerista* (Lacertilia, Scincidae) of Western Australia. Rec. West. Aust. Mus., 4(3): 241–256.
- . 1978. Taxonomic notes on the reptiles of the Shark Bay region, Western Australia. Rec. West. Aust. Mus., 6(3): 303–318.
- WALLS, G. L. 1934. The significance of the reptilian "spectacle." Amer. J. Ophthal., 17: 1045–1047.
- . 1942. The Vertebrate Eye and Its Adaptive Radiation. Bloomfield Hills, Michigan; Cranbrook Inst. of Science, 785 pp.
- WEBSTER, T. P. 1969. Aspects of the morphological and ecological variation in the *cyanura* group of the lizard genus *Emoia* (Sauria: Scincidae) in the Solomon Islands. Unpubl. B.A. Honors Thesis, Harvard University, 159 pp.
- WILLIAMS, E. E., AND M. K. HECHT. 1955. "Sunglasses" in two anoline lizards from Cuba. Science, 122: 691–692.

Adaptation and Function of Cranial Kinesis In Reptiles: A Time-Motion Analysis of Feeding in Alligator Lizards

T. H. FRAZZETTA¹

ABSTRACT. Cranial kinesis is widespread among a great many vertebrates, and should never be neglected in any study of the morphological changes in vertebrate skulls. Its adaptive role is not related to an increase in gape. Lizards possess amphikinetism (two kinetic joints) skulls, and thus differ from other kinetic forms. Feeding movements of a number of alligator lizards (*Gerrhonotus multicarinatus*) were studied by means of high speed cinematography. The attack upon prey may be divided into three sequential phases. Almost always the mandibular tip is accurately positioned very near the prey, while the upper jaw is less precisely placed at varying distances to it. Once the mandibles are located properly, relative to the prey, the upper jaw closes down to engage the quarry from above, very shortly after the mandibles engage it. Limitations in the ability to precisely position both upper and lower jaws at the same time enable kinesis to have an adaptive role. In *Gerrhonotus*, probably in many other lizards, and possibly in several other tetrapod groups, cranial kinesis may be important in allowing upper jaw engagement with prey without the need for an extremely large change in momentum of the attacker's head and forebody.

INTRODUCTION

KINETIC SKULLS

"Cranial kinesis" is a term introduced by Versluys in 1910 to designate the condition of many vertebrate skulls that possess a functional, transverse hinge across the skull roof. The effect of this, generally, is to permit the upper jaw

elements to move upward and downward relative to the braincase.

The placement of the transverse hinge varies among vertebrate groups. In many early tetrapods the location was posterior, where the upper, rear corners of the dermatocranium movably join the endocranium. When the kinetic axis is thusly placed, the condition is termed "metakinetic." In lizards especially, there exists a second kinetic axis across the middle of the skull—just above the posterior margin of the orbit—occupying a "mesokinetic" position. Apparently most lizards have both kinetic axes, and can be described as "amphikinetism." These three kinetic designations were provided by Versluys in his early papers (1910, 1912).

Some vertebrates, such as snakes and birds, possess a single kinetic joint located dorsally just anterior to the orbit. This condition is "prokinesis."

Kinetic skulls ordinarily depend upon a sliding joint between braincase and palate. Very often we can see, in a wide variety of kinetic skulls, that this joint is a diarthrosis connecting the pterygoids and a pair of prominently developed basiptyergoid (basitrabecular) processes of the braincase. A notable exception to this particular structural configuration is found in birds where, instead of an involvement with knoblike basiptyergoid processes, the pterygoid and palatine bones are movably coupled and, together, form a slider which can traverse a

¹Department of Ecology, Ethology and Evolution, University of Illinois, Urbana, Illinois 61801, U.S.A.

longitudinal, bony track on the braincase's ventral surface.

Not all moveable upper jaws are properly termed "kinetic." The complex, protrusible upper jaws of teleosts, described by many workers (e.g., Schaeffer and Rosen, 1961; Alexander, 1967; Liem, 1970) do not involve a transverse hinge (nor a functionally complementing braincase-palate modification). Some ganoid fishes (e.g., *Lepisosteus*, Arnold Brody, personal communication) can produce displacements of upper jaw elements which, again, are not accurately described as kinetic movements. Sharks can shift their jaws forward and downward relative to the braincase.

The earliest jawed vertebrates, the acanthodians and the placoderms, seem not to have had kinetic skulls, although the arthrodire placoderms were specially modified to lift the entire head (hence tilt the upper jaws) relative to the shoulder girdle (Heintz, 1932).

Moveable upper jaws, especially kinetic ones, were hardly exceptional throughout vertebrate evolution. From our mammalian vantage point, it perhaps seems "proper" that a skull should be constructed with a rigid association between neurocranium and upper jaws, allowing mandibular displacements alone to perform jaw movements. It is, however, the late therapsids and mammals that were, and are, largely exceptional.

There is an extensive literature on the presence of kinesis in many living and fossil vertebrates. A good summary is provided by Iordansky (1971a) in Russian. Less synoptic references documenting kinesis—but in English—include Moy-Thomas and Miles (1971) and Thomson (1966) for crossopterygian and paleoniscoid fishes; Jarvik (1954) for the structure of the skull in Ichthyostegalia; Panchen (1964), Pfannenstiel (1932), and Romer and Witter (1942) for various labyrinthodonts; Romer (1956) for the anatomy of reptile skulls; and Bock (1964) for birds.

A kinetic skull seems to have been a

widespread, primitive feature of tetrapods that was lost in a number of evolutionary lineages, while retained and often embellished in others. Thus many of the main, evolutionary avenues, leading to major groups, are sequentia of kinetic forms. A number of comparative anatomists still write about kinesis as though it were a fascinating morphological quirk which appeared here and there in vertebrate history. On the contrary, kinetic capabilities were of great importance. The compelling adaptive advantage they conferred often must have proscribed new cranial modifications that would have disrupted the integration of anatomical pieces that comprise kinetic machinery. Hence, investigations on the changes in cranial form are incomplete when the matter of kinesis is neglected.

ADAPTATION IN THE KINESIS OF LIZARDS

Because the present paper deals with adaptation, I should state that I do not believe that all evolutionary change is directly adaptive, nor that diverse animal groups are usually very different in fundamental, epigenetic ways, nor that the evolution of the relatively few fundamental differences which have appeared have come about necessarily in accord with the common statistical models. But I reaffirm that selective advantage, represented tangibly as adaptive form, is the driving and preserving agent in evolution, even though much inadaptive baggage may be carried along in a successful evolutionary line, and even though superior adaptations may compare temporarily to sloppily engineered devices (see Frazzetta, 1975).

The kinetic mechanism of lizards is interestingly complicated, involving several bony links, check ligaments, and muscles. It is more intricate, apparently, than the simpler, solely metakinetic skulls of the great majority of predatory, kinetic tetrapods.

Today, the living tetrapod whose skull most closely represents this primitive, metakinetic condition may not, ironical-

ly, be kinetic at all. *Sphenodon punctatus*, the sole surviving rhynchocephalian possesses a pair of well-developed pterygoid-basipterygoid joints, a firm suggestion of a metakinetic hinge structure, and the levator pterygoid and protractor pterygoid muscles (Fig. 1). Functionally, these muscles can serve no other purpose than to activate the kinetic apparatus. But they are sparsely represented in *Sphenodon*, consisting only of a few muscle fibers.

Versluys (1910) described this condition, and it was noted again by Poglayen-Neuwall (1953) and by Ostrom (1962). I have seen it in an adult specimen which I dissected. The adult skull is, however, functionally akinetic because bones are shaped in a manner to crowd into spaces necessary for kinetic movements. Both Versluys and Ostrom surmise that juvenile specimens are kinetic, but lose the capacity (possibly due to allometric bone growth) toward adulthood. Assuredly this is conjecture. In any case, the general cranial form of *Sphenodon* can represent a good model of the primitive, metakinetic configuration.

Figure 2 shows a primitive type based roughly on *Sphenodon*. Figure 2A represents the skull "at rest," while 2B depicts the major part of the dermal skull including the upper jaws rotated upward (protracted) about the metakinetic hinge. The drawing shows an exaggerated displacement to clarify the action. During protraction the pterygoids slide on the basipterygoids in an arc. The arc's radius is the distance, projected upon any parasagittal plane, from the metakinetic axis to the pterygoid-basipterygoid joint.

Protraction occurs by the contraction of the levator pterygoid and protractor pterygoid muscles, which draw the pterygoid bones forward and upward relative to the braincase. Because the pterygoids are firmly associated with the dermal palate and ectocranium, the upper jaws are lifted.

Retraction, the pulling back of the

dermal cranium relative to the neurocranium, restores the rest position. It is accomplished by any or all jaw closing muscles that originate on the dermal skull, and by extrinsic head muscles (not represented in Fig. 1) that run from the posterior mandibular surfaces to the neck region.

In lizards there are usually two kinetic joints, an amphikenetic condition that is presumably derived from a purely metakinetic skull (although Robinson [1967] makes the surprising statement that lizards arose from akinetic reptiles, and that the metakinetic hinge was a saurian invention). One of the two joints occupies the primitive, metakinetic location. The other is mesokinetic. Figure 3 shows a *Gerrhonotus multicarinatus* skull, with musculature represented. Here, again, there is a pterygoid-basipterygoid joint, but its conformation is specialized. The double-jointed kinesis demands that the pterygoids slip longitudinally, not along an arc's curve. We see in amphikinetic lizard skulls a basic four-bar linkage, one corner of which is "pinned" to the metakinetic axis (Frazzetta, 1962). Figure 4 shows a *Gerrhonotus* skull in outline, with the cranial elements reduced to mechanical links.

The anterior link, the muzzle unit, is formed of a rigid, triangular frame, the posterior strut of which is a typical bar in the four-bar mechanism. The braincase, shown as a rigid triradiate structure, joins the four-bar frame by a rotational joint whose axis is collinear with the metakinetic hinge. Elsewhere, the four-bar linkage is tied to the braincase by strong, elastic ligaments. The epipterygoid does not change the kinematic movement patterns of the linkage, but acts as a limiter and stabilizer. I have discussed its role, briefly, in 1962, and shall provide a future paper on its function.

In amphikinetic lizards, protraction involves the slight *depression* of the dermal cranium about the metakinetic hinge (the opposite of what occurs in more primitive types), and an elevation of the anterior

Figure 1. Skull of *Sphenodon punctatus*. Dorsal, ventral, and lateral views A, B, C. D, lateral view with temporal arcs removed to show placement of intrinsic jaw muscles.

Abbreviations: ar, articular; bo, basioccipital; bp, basipterygoid process; bs, basisphenoid; co, coronoid; de, dentary; ec, ectopterygoid; eo, exoccipital; ep, epipterygoid; fr, frontal; j, jugal; l, lacrimal; max, maxilla; na, nasal; pa, parietal; pal, palatine; pf, prefrontal; pm, premaxilla; po, postorbital; pof, postfrontal; pop, paroccipital process; pr, prootic; pt, pterygoid; q, quadrate; s, stapes; sm, septomaxilla; soc, supraoccipital; sq, squamosal; st, supra-temporal; sur, surangular; vo, vomer.

Figure 2. Diagram of kinetic movements in an hypothetical metakinetic (monokinetic) reptile. A. At rest or retracted. B. Protracted with mandibular depression. Shaded area represents endocranium lying within ectocranium (unshaded). Metakinetic pivot indicated by circled cross.

dermal skull (the muzzle unit) by upward rotation about the mesokinetic axis. The overall effect is nonetheless, a raising of the upper jaw relative to the neurocranium (Fig. 5).

In 1966 Iordansky published an excellent study on the kinetics of lizard skulls. He argued convincingly that amphikinesis, because it permitted a reverse angulation between muzzle and basal units during powerful retraction, could place forces on the prey between upper and lower jaws whose opposing vectors were more nearly collinear. In this manner, fewer unbalanced forces would exist, and the lizard would tend not to lose its meal by jaw forces that wedged the prey forward, inadvertently driving it from the mouth.

Although Iordansky accepts the basic four-bar linkage plan for most lizards, he raises a point concerning *Varanus*. He is accurate when he indicates that the lower strut of the triangular muzzle unit is, itself, interrupted by at least one, rotational turning-pair joint. However, this joint resists flexing by strong connective tissue attachments. As the skull is protracted, the joint allows very little angular displacement. When forcibly protracted, near the anterior limit of kinetic displacement, the intramuzzle

joint will flex a little. The effect is a protracting movement, based mainly on the initial four-bar linkage, which if forced further, will be added to slightly by intramuzzle movement. This small amount of additional movement does not alter significantly the fundamental, four-bar kinematics in such forms as *Varanus*. Nevertheless, the possibility of an adaptive role should not be dismissed.

I do, however, disagree with Iordansky in his claim (1966, 1970) that the pterygoideus muscle cannot act as a kinetic retractor. So long as the lower jaw cannot transmit a potentially nullifying force on the upper jaw, either directly or through prey held in the mouth, the tension produced by an activated pterygoideus muscle will result in an unbalanced force on the muzzle unit. It tends to pull the muzzle backward and downward for precisely similar mechanical reasons that it tends to pull the mandible forward and upward.

This functional point, concerning the pterygoideus muscle, leads to a complicating issue unknown to either Iordansky or myself in the 1960's or early 1970's. Throckmorton's work (1978) on an herbivorous lizard, and Smith's (1980) on a carnivorous species show that a portion of the pterygoideus actually aids in mandi-

Figure 3. Skull of *Gerrhonotus multicarinatus*. (Modified from McDowell and Bogert, 1954, and from several skulls at hand). Conventions as in Figure 1.

Figure 4. Linkage diagram of skull in *Gerrhonotus*. Braincase is represented as triradiate construction. Units of the four-bar linkage are shown, as is the epipterygoid bone.

Figure 5. Diagram of *Gerrhonotus* skull showing A, skull at rest (portion of mandible broken out in A to expose basal unit); B, skull protracted with mandibles depressed; C, skull fully retracted with jaws clamping prey. Compare Figure 2.

bular depression; another portion, a major one, apparently works as a mandibular elevator.

I concluded in 1962 that kinesis could be explained by its provision in allowing the upper jaws to close downward as the mandibles move upward, to insure a nearly simultaneous contact with the prey. This would avoid premature collision by the lower jaw alone, before the prey was secured, and prevent the quarry from being deflected out of seizing range by the very apparatus designed to capture it.

Iordansky (1966) took issue with this by raising the reasonable and challenging question: why is a moveable upper

jaw necessary when, in fact, movement of the entire head will position the upper jaw relative to prey during its capture? Thus, Iordansky rejects the proposal that kinesis is a device specifically evolved because of an advantage in moving the upper jaws downward to match the upward movement of the mandible when prey is seized.

The challenge is meaningful because, obviously, whole head movements can, indeed, position the upper jaws. Yet, I note the intuition of an earlier student of cranial kinesis (in birds), William Beecher who, in 1951, made the observation that a skull without a kinetic upper jaw would compare with a hand without

moveable fingers, but having a moveable thumb. The present paper concentrates on this particular question, by means of a detailed time and motion analysis of lizard feeding.

Since Iordansky's 1966 paper, a number of papers relating to saurian kinesis have appeared. These include Callison (1967), Carroll (1977), Gomes (1974), Gomes and Gasc (1973), Gow (1975), Haas (1973), Iordansky (1970, 1971a, 1971b), MacLean (1974), Rieppel (1978, 1980), Robinson (1967, 1973), Russell (1964), Smith (1980), and Throckmorton (1976, 1978, 1980). Many of these attempt an adaptive explanation for kinesis.

Additionally, in the past decade and one half, a number of general works on vertebrates allude to cranial kinesis in lizards or in other tetrapods. Often these allusions include a brief, adaptive comment. Although it may be difficult to be very confident about what the adaptive significance of kinesis is, in some cases a stronger confidence can be experienced in determining what the kinetic role is not.

Authors will state that cranial kinesis increases the gape, a view rejected by myself (1962), by Iordansky (1966), and by Rieppel (1978). It should be obvious why, in a purely metakinetic skull, there can be no effect on the gape whatsoever. In amphikinetic lizards, mandibular depression through rotation about the quadratomandibular joint can proceed only so far until the tympanum is so tightly sandwiched between the quadrate and the mandible's retroarticular process that the middle ear is at risk. Several ligaments in the lizard skull limit depression of the mandibles relative to the basal and muzzle units; hence, their affect is to allow the quadrate-retroarticular angle to diminish only so far. Because of that, the quadrate and mandible can be modelled as a single rigid unit solidly joined at this limiting angle. As protraction takes place, the muzzle is lifted, but so also is the mandible. Measurements reveal that the gape is actually reduced in protraction.

Perhaps it can be seen intuitively that in protraction of the four-bar mechanism, the shorter quadrate will rotate forward at a faster rate than will the longer, orbital strut of the muzzle unit (see Frazzetta, 1962).

There may be some kinetic skulls, in some vertebrates, that are constructed to increase gape. Many bird skulls may be so constructed (Bock, 1964). However, where gape increase can be shown to exist, caution should be respected in the direction of the simplistic conclusion that gape enlargement is, itself, the adaptive role. Some explanations based on gape can be shown to lead to the unconvincing conclusion that an animal is adapted to seize prey larger than it can swallow.

Suggestions that kinetic jaw structure provides for shock absorption are probably erroneous in all published cases (see Frazzetta, 1962). A future paper will examine the issue in further detail.

Several papers (e.g., Carroll, 1977; Robinson, 1967, 1973) have suggested that kinesis is explainable as a mechanism for swallowing. They propose that kinetic jaw movements move captured prey back into the mouth by alternating protraction and retraction. It is hard to visualize this for the purely metakinetic skulls of lizard ancestors. Amphikinetic lizard skulls have the mechanical capacity to draw the upper jaw's tip slightly backward relative to the mandible. Observations on swallowing in several lizards, however, show that prey is moved backward into the gullet by means of the tongue (Frazzetta, 1962), and/or by inertial feeding (Gans, 1961, 1969). The enhancement of swallowing ability provided by amphikinesis, either in tongue or inertial methods, seems negligible.

MATERIALS AND METHODS

Approximately two dozen *Gerrhonotus multicarinatus* were studied by means of high speed cinematography. Several

other lizard species of the genera *Lacerta*, *Eumeces*, *Sceloporus*, and *Gekko* were filmed as a means to compare similarities and differences between *Gerrhonotus* and several other forms.

A Hycam, high speed camera, equipped with its standard electronic speed control gave excellent service. It was used in conjunction with a Milimite timing light generator, and an Angenieux 12-120 mm zoom lens with a viewing tube. Occasionally a +1 or +2 diopter close-up lens was applied to the zoom lens.

Illumination was provided by four Colortan, spot-flood movie lamps to which were affixed heat absorbing glass filters to protect the animals. Light levels were measured with a Honeywell spot-meter.

Approximately twenty, one-hundred foot rolls of 16 mm Kodak film, of the types Ektachrome B, Tri-X reversal, and Four-X reversal, were exposed at speeds ranging from 250 to 500 fps. Of this, about 18 separate feeding acts were useful. Of the 18, only 8 were analyzable in fine detail.

All of the useful film was studied by means of a stop-motion, variable-speed, remote-controlled LW projector. Sequential tracings were made of the filmed feedings from projections reflected from a front-surfaced mirror angled at 45° to a table top. Overlays of consecutive tracings permitted determination of skull movements during the lizards' acquisition of prey. Eight of the filmed sequences were appropriate for more detailed study by means of a Vanguard Analyzer. This machine permits analysis of point displacements, from one frame to the next, relative to two coordinate axes in measurements of inches $\times 10^{-3}$. It also allows measurement of angular displacements between consecutive frames.

Continuously locatable points were selected on the lizard heads to permit point displacement analysis, or to define cranial axes whose angular changes could be measured. Despite the considerable

technological excellence of the Vanguard Analyzer, its actual use in this study proved difficult. Slight head twistings by the lizards as they lunged, an occasional blurred frame, an unexpected, momentary highlight which masks the sought-after point, were among a number of unwelcome factors. Hence, each point and each angle was measured ten times for all eighteen filmed sequences. In those cases where there was a lack of strong consistency in a set of ten readings, the sequence was rejected from this aspect of the study.

The prey offered was either a mealworm or a cricket, restrained by a fine, cotton thread which was snipped after the capture. Occasionally prey would be suspended above the substrate, to simulate a slow-flying quarry (which I have seen *Gerrhonotus* capture), or instances where the prey is on a leaf, twig or grass stem above the ground.

Dissections were performed on three preserved *Sphenodon punctatus*, and fresh *Gerrhonotus* which died during captivity.

THE PATTERN OF ATTACK ON THE PREY

There are three discernable phases to the attack on prey by a lizard. The first is the rush, or the chase, where the attacker dashes up to the prey—or pursues it—to within lunging distance.

The second phase is the lunge, or what I prefer to call the delivery because the capturing apparatus, the jaws, are delivered to within seizing distance. Then the seizure or grasp is the third and final phase in the capture.

The second and third phases were studied, and are represented in the tracings and the Vanguard analysis. Figure 6 shows a typical delivery and grasp pattern when prey is on a flat substrate shared by the lizard.

With elevated forebody, the neck and head arch above the ground a short dis-

Figure 6. Sequence of tracings of *Gerrhonotus* specimen III, film 2, from projections. "T" designations show elapsed time in seconds from the first frame shown.

tance from the prey toward the lizard. The head is turned downward, its axis roughly in line with the prey.

The delivery phase begins with the mouth closed, and occurs by coordinated movements of the front limbs, forebody, neck and cervicocranial joint. As the delivery progresses, the mandibles are gradually depressed so that the degree of mouth opening increases more or less steadily as the jaws approach their target.

Soon after mandibular depression begins, kinetic protraction becomes progressively conspicuous. The muzzle is seen to lift relative to the more posterior

regions of the head, and the leading, vertical edge of the eardrum—which corresponds to the longitudinal axis of the quadrate, to which it is fastened—becomes gradually angled forward from a point just above it.

The tip of the lower jaw is almost always placed in light contact with the ground, as close to the prey as possible without actually touching it. The precision of this placement is impressive.

At the time the lower jaw tip is placed, the upper jaw is clear of prey contact by a substantial margin of distance, and the kinetic mechanism is usually protracted

to its maximum degree. Often, immediately following this mandibular placement on the substrate, the entire head is turned further downward to close some of the distance between the upper jaw and prey surface. Almost always, as soon as mandibular contact with the ground is established, the head undergoes rapid, kinetic retraction.

At this moment, the muzzle begins to descend, relative to the more posterior region of the head, and the tympanum's leading edge starts its backward swing. Frequently the eye is seen to sink deep into the orbit, presumably a result of contraction of the levator bulbi, a weak kinetic retractor (Frazzetta, 1962) which in *Gerrhonotus* inserts on the pyriform membrane of the palate and on tissues surrounding the eye (Gomes, 1974).

During these movements of the upper jaw, the mandibular tip moves toward the prey along the substrate. This movement is caused only partially by rotation of the mandible about the quadratomandibular joint. It is also effected by pushing the entire head forward as it tilts downward while the neck arches higher upward. The lizard now has positioned itself further over the prey. The lower jaw touches the prey first. The upper jaw, which has been descending toward the prey, may quicken its movement after the lower jaw has made contact with the prey. The upper jaw engages the prey a tiny fraction of a second after the mandibles' contact. Significantly, the prey itself is hardly moved from its position by the seizing action. At the grasp, or just after it, retraction is usually strong enough to carry the muzzle downward past its rest position, to thus hold the prey as Iordansky has postulated (1966).

When prey is taken above the substrate's surface, the success probability of capture appears reduced. Sometimes, this was caused by artificial circumstances, such as the suspending thread being touched by the lizard's jaws which swung the prey out of seizing range. Perhaps, in other instances, there might have

been greater movement of the prey when suspended, despite our efforts to hold the thread still; the lizards' aim might thusly have been less accurate. But beyond these factors, the impression is that the required upward angle of thrust toward the prey, and the prey's position closer to the upper than to the lower jaw, were awkward circumstances that the captor had to overcome.

Figure 7 shows a successful, above-ground capture. In Figure 8 the upper jaw of the lizard bumps the suspending thread and deflects the prey. Because above-ground attacks have a greater variety of factors that the lizard must deal with, they tend themselves to be more diverse, and difficult to typify according to a single, general pattern.

The delivery phase begins with mouth closed and, as before, mandibular depression grows progressively as the prey is neared. Soon the kinetic apparatus begins a steady protraction, reaching a maximum shortly before the jaws are to close on the prey. In both Figure 7 and Figure 8, the tips of the mandibles approach closer to the target than does the upper jaw. This is the usual case in above-ground attacks, as it is when the prey is located on the substrate, as in Figure 6.

When the jaws close, the head may turn downward on the neck (Fig. 8D-E) or it may not (Fig. 7). Jaw closure is produced by elevating the mandible while, at this time, the skull undergoes rapid kinetic retraction. Retraction occurs in Figure 7, but it is but slightly noticeable there. It is conspicuous in Figure 8, and begins (Fig. 8D-E) as soon as the thread is touched. In this sequence the lizard longitudinally twists the head, possibly in a vain attempt to avoid the thread.

VANGUARD ANALYSIS

Capturing prey on the ground is different enough from taking it in the air that

Figure 7. Sequence of tracings of *Gerrhonotus* specimen II, film 1, from projections. "T" designations show elapsed time in seconds from the first frame shown.

convenience enlarges if the two types of attack are regarded separately. The same points and axes, however, are studied for both capture modes.

The points and axes are shown in Figure 9. The purpose in choosing such points as the upper and lower jaw tips (*uj* and *lj*) is clear. Ear top (*et*) is a point that corresponds closely to the joint connecting quadrate, parietal unit, and braincase, and lies close to the cervicocranial articulation. It is a useful point to keep track of.

Figure 10 shows the Vanguard analysis of the same sequence represented in Figure 7. In Figure 10, however, data points are shown for *every* frame, from

the moment the mouth begins to open, early in the delivery phase, to the final seizing of the prey. On any of the twelve curves, each dot is the relevant data point for that frame. Hence, the abscissa is time divided into increments of the reciprocal of camera frame rate, in this case $1/287.0$ of a second, progressing from left to right.

The ordinate units show displacement differences. For data points whose displacements are measured as linear changes projected upon a set of vertical and horizontal axes on the Vanguard machine (not on the axes of Figure 10), the horizontal (X) and vertical (Y) displacements are each graphed separately

Figure 8. Sequence of tracings of *Gerrhonotus* specimen I, film E, from projections. "T" designations show elapsed time in seconds from first frame shown.

here, against the time-related abscissa. A scale of linear displacements is shown in the figure.

Point data (upper jaw point x and y, ujx , ujy ; lower jaw point x and y, ljx , $l jy$; ear top point x and y, etx , ety) are each arranged so that an increase in either horizontal or vertical displacement is indicated by an ascension of sequential data points relative to the ordinate axis of the figure.

Angular data is shown for parietal top axis, pt , and is graphed so that a tilting upward of the head is indicated by a rise in the points shown. Angles between

pairs of several axes (snout top axis with parietal top axis, $\overline{snl-pt}$; muzzle unit upper axis with cranial axis, $\overline{mzu-cr}$; muzzle unit lower axis with cheek axis, $\overline{mzl-ch}$; ear axis with cranial, $\overline{e-cr}$) reveal kinetic flexing of the skull. The graphed points rise when protracting movements are indicated, and fall in retraction.

The mandibular axis' angle with the ear axis, $\overline{m-e}$, shows the turning of the mandible upward or downward in its rotation about its joint with the quadrate.

In several of the eight Vanguard analyses (Figs. 10-13) there are occasional inconsistencies. For example, the

Figure 9. Head of *Gerrhonotus* showing points and axes used in Vanguard analysis.

$\overline{mzl-ch}$ angle sometimes corresponds poorly with other kinetic indicators.

The sequences all start at time-zero when the mouth first starts to open. Hence, the delivery phase has already begun shortly before the graphs begin.

PREY CAPTURED ON THE SUBSTRATE

Typically the ear top point shows a rather smooth, continuous path in both x and y displacements. This is seen in Figure 10. The jaw tips, too, run a fairly continuous course up to the time when the mandibular tip is placed upon the substrate. From that time on, there is little or no vertical change in the lower jaw tip (ljy in Figs. 10, 11A,B,C) because it slides—or nearly does—along the substrate toward the prey. In the cases shown in Figures 10 and 11B, the horizontal motion of the lower jaw tip (ljx) is erratic as the moment of prey capture is approached. In two other cases of capture on the substrate (Fig. 11A,C) the lower jaw tip moves in a smoother manner. During this period, the upper jaw movements—both horizontal and vertical (ujx

and y)—are reasonably smooth in three of the four substrate captures; the exception is the sequence shown in Figure 10.

Tilting of the head, shown by angular changes in the parietal top axis ($\overline{pt}$), is different in each of the four substrate captures. In Figure 10 the head tilts sharply downward just before the mandibular tip touches the substrate, and then rotates further downward at varying rates thereafter. In Figure 11A the downward tilt begins after the mandibles come into contact with the prey and, as before, proceeds at a discontinuous rate. However, in the captures shown in Figure 11B and C, the head begins to turn upward from the time of mandible-substrate contact, and then turns downward later.

Lower jaw depression, relative to the quadrature ($\overline{m-e}$), is mostly a smooth, continuous motion except in Figure 11B. It reaches its greatest relative depression just before mandibular contact with the substrate (except in Figure 11C). Elevation may be smooth or not.

The four kinetic angle differences are the most difficult to measure on the Vanguard Analyzer. Consideration of the

Figure 10. Vanguard analysis of *Gerrhonotus* specimen III, film 2. In this figure, and in Figures 11, 12 and 13, Roman numeral gives specimen number, Arabic number or letter gives film number, fraction (here, 1/287.0) is time interval between successive frames (hence, dots), and mm measurement is image size of head on analyzer screen, measured from eartop point to upper jaw tip point. Vertical line with asterisk is scale: 2.54 mm (image size) for linear data, 2.5° for angular data.

Abbreviations in graph title: AGC, above ground capture; AGC*, above ground capture but prey very nearly missed; AGM, above ground attempt but prey actually missed; GC, capture on ground.

Abbreviations at graph's bottom: lj-g, lower jaw touches ground; lj-p, lower jaw touches prey; uj-p, upper jaw touches prey; es, eye sinks into orbit; ps, prey securely seized by both upper and lower jaws. Other abbreviations and explanations in text.

sequences illustrating substrate-based capture, indicates that kinetic protraction reaches its maximum at, or shortly before, the moment of mandibular contact with the ground. Most times (Figs. 10, 11A,B) protraction seems to sharpen suddenly just before retraction. This is corroborated in the several traced sequences unsuitable for Vanguard analysis (see explanation above).

Prey seizure occurs after a retraction of the kinetic apparatus. As already mentioned, the muzzle-cheek angle ($\overline{mzl-ch}$) corresponds poorly with other kinetic indicators (especially in Figure 11A). I cannot offer an explanation for this. There are at least two possibilities. Accurate measurement of the angle may be more difficult than that in the other kinetic determinations. Or, small adjustments of elements at the joints during their rotation may show up conspicuously in this measurement. If so, the effect could frustrate application of a model which is simplified to regard the moveable articulations between cranial units as mechanical, turning-pair, planar joints.

PREY CAPTURED ABOVE THE SUBSTRATE

It is a little harder to find common denominators in head and jaw movement patterns when prey is taken above the substrate, than when taken upon the substrate. However, in broader aspects, some rough paradigms do exist.

As before, the ear top point (*et*) moves continuously without noticeably sharp breaks in rate or direction. The tips of both upper and lower jaws move only slightly less smoothly than the ear top (Figs. 12, 13). The parietal top axis (*pt*), which reveals the tilting of the head, in all four cases tilts upward at first, and then turns downward. The reversal occurs just before, or when, the lower jaw touches the prey. When prey is taken on the substrate, the mandibles typically are placed close to the prey, and then touch it a tenth of a second or less before the

Figure 11. Vanguard analyses of three film sequences involving captures on the ground. See Figure 10 for explanation.

Figure 12. Vanguard analysis of *Gerrhonotus* specimen II, film 1. See Figure 10 for explanation.

upper jaw makes prey contact. Prey taken above the substrate is, as in the substrate-based captures, approached and touched by the lower jaw before the upper.

The angular change between mandible and ear (quadrate) axis is much like those seen in the previous set of examples. Here, however, in two cases (Fig. 13B, C) the angle begins to close well before contact is made between mandibles and prey.

In Figure 12, retraction begins immediately after mandible-prey contact. The other three cases (Fig. 13) appear to reveal retraction beginning at various times before the prey is touched by the lower jaw.

DISCUSSION

The delivery phase involves a smooth, continuous approach of the neck and neck-cranial joint. The jaw tips themselves are also reasonably flowing in their motions; their smooth movements may become broken as the prey is closely approached. Angular changes in the tilt of the head vary from case to case: sometimes they are continuous, while at other times they seem quite erratic.

Apparently the combination of movements involving the neck-cranial joint, the neck and forebody and front limbs, the angulation of the head, the depression of the lower jaw relative to the quadrate, and kinetic rotations can often control the jaw tip displacements to negotiate an even, unbroken motion. Unquestionably, Iordansky (1966) is correct in that tiltings of the entire head can bring the jaws close to coincidence with the prey. There is, however, more. The mandibles are always first to come near the prey, and they are first to make contact with it. Only afterward does the upper jaw become involved. While the lower jaw is rapidly being positioned near the prey, the margin of distance between prey and upper jaw is usually moderate to fairly great.

Great precision in the aiming of the mandibular tip, relative to the prey's location, is seen in most cases, especially those where the prey is taken on the substrate. At the moment of this accurate mandibular placement, the upper jaw—separated as it is from the prey by a considerable distance—seems, in contrast, imprecisely or but roughly positioned. Very soon after the mandibles have been accurately placed near the target, the upper jaw moves quickly downward to cooperate with the mandibles in making the grasp. This downward motion of the upper jaw may be effected by a variety of factors (head tilting, etc.), but always includes kinetic retraction.

Prey is seized only when the skull has undergone much retraction, and we thus

Figure 13. Vanguard analysis of three film sequences involving attempts to capture prey above ground. See Figure 10 for explanation.

can note on behavioral grounds—as we had, previously, on mechanical ones—that the adaptive meaning of kinesis cannot be the increase of gape for the seizing of larger prey.

It appears that the lizards in this study do not act to set both upper and lower jaws relative to the prey with equal accuracy. There is a strong, perhaps irresistible temptation to interpret the films as revealing an *inability* of the lizards to carefully guide both upper and lower jaws simultaneously. Hence, the lizard must pay attention to the jaw position sequentially, “choosing” to regard the lower jaw first for the best of mechanical and practical reasons.

Lizards live in our world of significant, gravitationally induced forces. Much of their food will lie *upon* a surface, regardless how irregular or complex this surface may be. An obvious disaster in prey capture would be the ramming of the mandibular tip into the substrate. Potential injury would, I think, be less important than the sudden, unplanned-for jolt that would be suffered by the head and jaws, and the subsequent disruption of the complexly controlled delivery phase of prey capture. Equally unwelcome would be the effect of misplacing the mandibular tip behind a tiny hillock of earth, a pebble, a twig, anything that would form a barrier between a rapid, simple movement of the lower jaw to the prey.

Because of these considerations, if the lizard cannot concentrate equally on both upper and lower jaws at the same time, it should devote its first attention to the mandibles. It follows that, because a lizard so often deals with quarry upon a substrate, its morphological and behavioral systems should be adjusted for the sequence: lower jaw, then upper. As a result, even prey taken off the ground, where the sequence is less important, will be approached by the mandibles first.

The fact that the lizards seemed not to aim with both upper and lower jaws simultaneously has a possible parallel in

the grasping operation of a human hand. Beecher's (1951) comparison between the hand and the kinetic bird skull illuminates the hand as a natural model of the amphikinetic lizard skull. In this context the wrist may be seen as representative of the lizard's head-neck joint; the hand, from wrist to the distal ends of metacarpals II through V, compares with the posterior skull half; digits II through V together form the “upper jaw,” with the “mesokinetic hinge” at the metacarpal-phalangeal joints; and the thumb simulates the mandibles.

Among several important discrepancies between the hand and the amphikinetic lizard skull is that, of course, the guidance system (eyes, mainly) travels with the jaws when prey is grasped. Nonetheless, some of the similarities between hand and lizard skull suggests that the hand is a more accurate mechanical analogy than would be a shop-crafted structural model.

When grasping small objects on a desk top, my intuitive impression is that I exercise little conscious direction to the details of placing thumb and fingers relative to the target. But I also notice that while my thumb tip becomes set very close to the subject, my fingers are quite far away, but descend rapidly to complete the grasp.

In an attempt to slightly objectify the hand model of the kinetic skull, I enlisted the aid of two laboratory assistants who were told only to capture a ping pong ball, thrown bouncingly toward them along a table top, by thumb below and fingers above. Those exercises were performed before the high speed camera, set at an average speed of 390 fps. Two one-hundred foot rolls of Tri-X film were thusly exposed. Not all capture attempts were successful, and a number of misses were recorded.

Figure 14 shows a typical (successful) capture. The hand and fingers are positioned by movements of forearm and hand rotations about the wrist. The “gape” between thumb and fingers in-

creases slightly as the target comes nearer. The thumb seems carefully placed in close proximity to the ball. Very soon afterwards the upper fingers—and the hand—move downward. By the time first contact is made by the thumb, the upper fingers are so close that they are a tiny fraction of a second away from reaching the target themselves. Notice that because of the thumb's first contact, the ball

Figure 14. Sequence of tracings from projections of filmed capture of ping pong ball by hand. "T" designations show elapsed time in seconds from first frame shown. Explanation in text.

begins to bounce off of it, but that the proximity of the upper fingers is sufficient to arrest this momentary deflection, and to contain the ball. During the capture, although the upper fingers are brought close to the target, they are not so close that they may accidentally touch—thus deflect and possibly lose—the object during the time that attention is being focused elsewhere (the thumb).

Kinetic lizards may deal with similar problems in prey capture. Ideally, *both* upper and lower jaws should be positioned precisely only a very small distance away from the prey so seizing can occur immediately at the instant that the jaws begin to close. No creature, however, is so accurate. In the lizards studied, this inaccuracy seems much greater in upper jaw than in lower jaw placement. Its deficiency can partially be compensated by certain kinds of adaptive mechanisms which can act to dilute the failure that inaccuracy may cause.

The kinetic jaw machinery can function as such a compensator. But so, presumably, could an akinetic upper jaw (Iordansky, 1966). Hence, in assessing the possible adaptive role of kinesis, it becomes necessary to compare kinetic with akinetic skulls.

Figure 15 illustrates a lizardlike reptile taking prey from the ground. The animal's position is close to the end of the delivery phase, and near the beginning of the seizing phase. When delivery first began the head and jaws were located at some distance "behind" the prey. The delivery brings the jaws rapidly to either the place where the prey was located when this phase began, or to the estimated new location of an active prey as gauged by its movement when delivery was initiated.

Possibly some changes in aim can be made during delivery, as a means of "tracking" elusively moving prey. Whether or not this occurs, the movements during this phase must bring the jaws to a seizing position. In Figure 15, the jaws are positioned for seizing. A

vector is shown representing the momentum of head and forebody, produced by the thrusting effects of limb, trunk and cervicocranial musculature. The synergistic actions of this musculature result in a smoothly, coordinated delivery of jaws to prey.

Seizing of the prey requires an upper jaw movement which can be shown (as in Figure 15) vectorially as the required change in momentum of the head. Data on the lizards have shown, already, that head tiltings can occur suddenly, during the delivery phase, thus changing the head's momentum. This change is most probably caused by cervicocranial musculature.

But a change of momentum at the instant of seizing the prey has different implications than such a change occurring earlier, in the delivery phase. This is because successful seizing depends upon a reasonably precise placement of the jaws upon the prey's surface, with the correct magnitude and development of mouth-closing forces. The momentum change in the seizing phase has to be rapid, complete, and its scheduling well timed. In an akinetic skull, it requires a break from the functional integration and coordinated, sequential action of the considerable musculature which functionally is carrying out the delivery.

A deleting of some muscle group from the control of the delivery, to enlist it suddenly in the seizing phase, all within hundredths of a second to effect a precision grasp of elusive—or potentially elusive—prey, could, despite these ticklish factors, result in a successful capture. But it would seem that much greater assurance of success can be had through kinetic movement. The capacity of any type of kinetic system, enabling the upper jaw to snap down on the prey without usurpation of the delivery musculature, can close the distance between prey and upper jaw established during mandibular placement, without the impairment of other facets of the attack procedure.

In the lizard examples observed, the

consistent, carefully timed retraction just as prey is seized, and the relative inconsistency of head tilt scheduling, suggests that retraction lacks a very constrained, operational connection with any specific kind of head or neck displacements. Kinesis frees the performance of the upper jaw from a rigid coupling with movements and attitudes of other portions of the head and of the forebody.

Perhaps the range of kinetic movement, from the position of the upper jaw in protraction, to that when it is retracted at the moment of seizing prey, is a measure of the inexactness of upper jaw placement. If a lizard, while concentrating on the placing of the mandibles, inaccurately positions the upper jaw, through head and neck movements, farther away from the prey—by a distance s —than its upper jaw can readily traverse

Figure 15. Diagram of generalized reptile capturing prey upon ground. Vector R shows momentum of head and neck immediately prior to capture. Vector S shows direction upper jaw must take to engage prey surface. Magnitude of S is adjusted to make vector C as short as possible. Vector C is the minimum change in momentum required to capture prey if the skull were akinetic. Further explanation in text.

by means of those same sources of movement, the kinetic range should be, approximately, s .

The parameter s need not be very great. Kinetic ranges in some lizards are small. Crompton (1955) in a study of kinesis in some mammal-like reptiles, decided that their kinesis had no adaptive implications because the kinetic range was so restricted. However, even a very small kinetic range can spare the head a very large momentum change (Fig. 15).

In many akinetic predators, head movements are a large part of both the delivery phase and the seizing phase. Kinetic predators, such as *Gerrhonotus*, can reduce the importance of head movements in the seizing function. In these forms, head movements, if they occur beyond the delivery phase, act as an operational link between delivery and seizing, because the final grasp belongs to the functionally independent kinetic jaws.

ACKNOWLEDGMENTS

I am grateful to many colleagues who assisted me throughout this study. Prominent among them are Ernest E. Williams of Harvard's Museum of Comparative Zoology, who loaned specimens, including two of the preserved *Sphenodon*, and discussed this work during its preliminary phases; George Zug provided study skulls from the collections of the U.S. National Museum, and Hyman Marx loaned material from Chicago's Field Museum; Richard Estes offered critical comments on this study in its early days; Dennis W. Nyberg, Timothy C. Moermond, Kenneth V. Kardong, Carlos F. Pinkham, F. William Burley, and the late Sandra Harness provided discussion and laboratory assistance. The work was carried out with the support of National Science Foundation Grant GB 5831.

LITERATURE CITED

- ALEXANDER, R. MCN. 1967. The functions and mechanisms of the protrusible upper jaws of some acanthopterygian fish. *J. Zool., Lond.*, **151**: 43-64.
- BEECHER, W. J. 1951. Adaptations for food getting in the American blackbirds. *Auk*, **68**: 411-440.
- BOCK, W. J. 1964. Kinetics of the avian skull. *J. Morphol.*, **114**: 1-42.
- CALLISON, G. 1967. Intracranial mobility in Kansas mosasaurs. *U. Kans. Paleont. Contr.*, (26): 1-15.
- CARROLL, R. L. 1977. The origin of lizards, pp. 359-396. In S. M. Andrews *et al.* (eds.), *Problems in vertebrate evolution*. Linn. Soc. Symp. Ser.
- CROMPTON, A. W. 1955. A revision of the Sceloposauridae with special reference to kinetism in this family. *Navors. Nas. Mus.*, **1**: 149-183.
- FRAZZETTA, T. H. 1962. A functional consideration of cranial kinesis in lizards. *J. Morphol.*, **111**: 287-320.
- . 1975. *Complex adaptations in evolving populations*. Sinauer Assoc., Sunderland, Mass.
- GANS, C. 1961. The feeding mechanism of snakes and its possible evolution. *Am. Zool.*, **1**: 217-227.
- . 1969. Comments on inertial feeding. *Copeia*, **4**: 855.
- GOMES, N. 1974. Anatomie comparée de la musculature trigéminal des lacertiens. *Mém. Mus. National d'Hist. Natur.*, ser. A, **Zool.**, **90**: 1-107.
- GOMES, N. M. B., AND J. P. GASC. 1973. Etude biomécanique du mouvement de fermeture de la mandibule chez *Ophisaurus apodus* (Sauria, Anguillidae). *Papeis Avulsos d. Zool.*, **27**: 1-25.
- GOW, C. E. 1975. The morphology and relationships of *Youngina capensis* Broom and *Prolacerta broomi* Parrington. *Paleont. afr.*, **18**: 89-131.
- HAAS, G. 1973. Muscles of the jaws and associated structures in the Rhynchocephalia and Squamata, pp. 285-490. In G. Gans and T. S. Parsons (ed.), *Biology of the Reptilia*, **4**, NY, Acad. Press.
- HEINTZ, A. 1932. The structure of *Dinichthys*: A contribution to our knowledge of the Arthrodira. *Am. Mus. Nat. Hist.*, Bashford Dean Mem. Vol., **4**: 115-224.
- IORDANSKY, N. N. 1966. Cranial kinetism in lizards. Contribution to the problem of the adaptive significance of the skull kinetism. *Zool. Zhurn.*, **45**: 1398-1410. (In Russian).
- . 1970. Structure and biomechanical analysis of function of the jaw muscles in the lizards. *Anat. Anz.*, **127**: 383-413.
- . 1971a. Cranial kinesis in the evolution of reptiles. *General Biol.*, **32**: 348-365. (In Russian).
- . 1971b. A contribution to the functional analysis of the skull of lizards (Lacertilia). Peculiarities of the construction of the skull related to

- cranial kinesis. *Zool. Zhurn.*, **50**: 724-733. (In Russian).
- JARVIK, E. 1954. On the visceral skeleton in *Eusthenopteron* with a discussion of the parasphenoid and palatoquadrate in fishes. *K. Svenska. Vetens.-Akad. Handl.*, (4)**5**: 1-104.
- LIEM, K. F. 1970. Comparative functional anatomy of the Nandidae (Pisces: Teleostei). *Fieldiana, Zool.*, **56**: 1-166.
- MACLEAN, W. P. 1974. Feeding and locomotion mechanisms of teiid lizards: functional morphology and evolution. *Papeis Avulsos d. Zool.*, **27**: 179-213.
- MCDOWELL, S. B., AND C. M. BOGERT. 1954. The systematic position of *Lanthanotus* and the affinities of the anguimorph lizards. *Bull. Am. Mus. Nat. Hist.*, **105**: 1-142.
- MOY-THOMAS, J. A., AND R. S. MILES. 1971. Paleozoic fishes. Philadelphia, Saunders Co.
- OSTROM, J. H. 1962. On the constrictor dorsalis muscles of *Sphenodon*. *Copeia*, **1962**: 732-735.
- PANCHEN, A. L. 1964. The cranial anatomy of two Coal Measure anthracosaurs. *Phil. Trans. Roy. Soc. Lond., Ser. B.*, **247**: 593-637.
- PFANNENSTIEL, M. 1932. Der kinetische Schädel karbonischer Stegocephalen. *Fortsch. Geol. U. Palaeont.* (Deecke-Festschrift), **11**: 278-305.
- POGLAYEN-NEUWALL, I. 1953. Untersuchungen über die Trigemini-muskulatur von *Hatteria*. *Zeitschr. Wiss. Zool.*, **157**(1-2): 57-76.
- RIEPEL, O. 1978. Streptostyly and muscle function in lizards. *Experientia*, **34**: 776-777.
- . 1980. The trigeminal jaw adductor musculature of *Tupinambis*, with comments on the phylogenetic relationships of the Teiidae (Reptilia, Lacertilia). *Zool. J. Linn. Soc.*, **69**: 1-29.
- ROBINSON, P. L. 1967. The evolution of the Lacertilia. *Probl. Actuels de Paleont. (Evol. d. Vertebr.) Colloq. Intern. D. Centr. Nat'l. de la Recherche Scient.*, **163**: 395-407.
- . 1973. A problematic reptile from the British Upper Trias. *J. Geol. Soc. Lond.*, **129**: 457-479.
- ROMER, A. S. 1956. Osteology of the reptiles. University of Chicago.
- ROMER, A. S., AND R. V. WITTER. 1942. *Edops*, a primitive rhachitomous amphibian from the Texas Red Beds. *J. Geol.*, **50**: 925-960.
- RUSSELL, D. A. 1964. Intracranial mobility in mosasaurs. *Postilla*, **86**: 1-19.
- SCHAEFFER, B., AND D. E. ROSEN. 1961. Major adaptive levels in the evolution of the actinopterygian feeding mechanism. *Am. Zool.*, **1**: 187-204.
- SMITH, K. K. 1980. Mechanical significance of streptostyly in lizards. *Nature*, **283**: 778-779.
- THOMSON, K. S. 1966. Intracranial mobility in the coelacanth. *Science*, **153**: 999-1000.
- THROCKMORTON, G. S. 1976. Oral food processing in two herbivorous lizards, *Iguana iguana* (Iguanidae) and *Uromastix aegyptius* (Agamidae). *J. Morphol.*, **148**: 363-390.
- . 1978. Action of the pterygoideus muscle during feeding in the lizard *Uromastix aegyptius* (Agamidae). *Anat. Rec.*, **190**: 217-222.
- . 1980. The chewing cycle in the herbivorous lizard *Uromastix aegyptius* (Agamidae). *Arch. Oral Biol.*, **25**: 225-233.
- VERSLUYS, J. 1910. Streptostylie bei Dinosauriern. *Zool. Jahrb. Abt. f. Anat.*, **30**: 177-258.
- . 1912. Das Streptostylie-Problem und die bewegung im Schädel bei Sauropsida. *Zool. Jahrb., Suppl.* **15**, **2**: 545-716.

The Evolution of the Subdigital Pad in *Anolis*.

I. Comparisons Among the Anoline Genera

JANE A. PETERSON¹

ABSTRACT. The morphology of the subdigital pad in each of the three anoline genera, *Chamaeleolis*, *Chamaelinorops*, and *Phenacosaurus*, is compared to that of *Anolis* species with similar body size and where possible similar structural habitat. *Chamaeleolis* and *Phenacosaurus* exhibit gross pad and seta morphologies that are very similar to those of *Anolis*. The gross morphology of the *Chamaelinorops* pad resembles that of several derived and more terrestrial anole species, particularly *A. auratus*, but the structure of the setae and spines is dramatically different from that of any other anoline examined. It is not clear whether the *Chamaelinorops* morphology represents a highly specialized derivative of an *Anolis*-like pad complex or an independent evolutionary experiment in subdigital pads. The comparisons among *Phenacosaurus*, *Chamaeleolis*, and a number of the more primitive anole species suggest that the major elements of the pad complex—the morphotypic series in fine structure, including the setae, and lamellar scales forming a pad—are homologous among these anolines and that the pad complex appeared quite early in the anoline radiation, before *Chamaeleolis* diverged.

INTRODUCTION

Anolis and its close relatives, *Chamaeleolis*, *Chamaelinorops*, and *Phenacosaurus*, constitute a highly successful radiation of arboreal iguanid lizards. The four genera are unique among iguanids, a group which includes many arboreal forms, in possessing a subdigital adhesive pad. The pad consists of laterally expanded flexible scales, the lamellae. The external surface of the lamellae is

covered with microscopic hairs or setae (the presence of setae in *Chamaelinorops*, *Chamaeleolis*, and *Phenacosaurus* has previously been reported by Maderson, 1970). The precise mechanism of grip is unknown, but the setae appear to be the major agent (see Hiller, 1968 and Peterson and Williams, 1981 for discussion of setal function in *Anolis*).

The pad complex appears to be a key innovation (*sensu* Liem, 1973) in the successful radiation of the group. Compared to animals that must rely on their claws and toe position for grip, anoles appear to have greater range of useable locomotor substrates. They can grip or adhere to a perch of almost any diameter (or radius of curvature), spatial orientation, or surface texture (e.g., surfaces as smooth as glass or as rough as bark). It may become possible to state the relationship between the pad complex and physical aspects of the habitat more precisely in the future, but the adhesive pad seems to be among the morphological adaptations that permit anoles to adopt highly acrobatic locomotor behavior and to exploit the spatial heterogeneity of arboreal niches.

In large part through my association with Ernest E. Williams I have become interested in the origin and adaptive diversity of the pad complex. In this study *Chamaeleolis*, *Phenacosaurus*, and *Chamaelinorops* are compared with *Anolis*. Each of the non-*Anolis* genera combines primitive osteological features with unique specializations (Williams, 1977), and two or possibly all three of the

¹Department of Biology, University of California, Los Angeles, Los Angeles, California 90024, U.S.A.

genera diverged from the *Anolis* stock relatively early (Fig. 1). *Chamaeleolis* and *Chamaelinorops* may represent a much earlier invasion of the Greater Antilles than does *Anolis* itself. If they are relics of an early radiation in which pad morphology, as seen in *Anolis*, was not fully established, some features of the pad complex, such as setal stalk diameter and the density of setae on the lamellae, may have evolved in parallel in two or three lineages. Differences in pad morphology among *Chamaelinorops*, *Chamaeleolis*, and *Anolis* could reflect alternate pathways in pad evolution. *Phenacosaurus* is distinct from *Anolis*, but it may be derived from very early South American anoles (Lazell, 1969; Etheridge, 1960). It could represent yet a

third or a fourth lineage in which the pad complex was perfected.

The comparisons are designed to determine whether each of the genera can be distinguished from *Anolis*. In order to minimize the effect of differences in body size or habitat preference on the comparisons, each genus is compared with *Anolis* species which are similar in habitat preference and body size. Ideally the comparisons would also minimize the phylogenetic distance between the *Anolis* species and the anoline genus, but in practice there are too few data to identify nearest neighbors within *Anolis*. *Chamaeleolis*, a large, canopy-dwelling form, is compared with a crown-giant anole ecomorph, *A. cuvieri*. *Chamaelinorops* is a small and

Figure 1. The relationships among the anoline genera according to A) Williams (1977) and B) Etheridge [after Paull, Williams and Hall (1975)]. The osteological and external characters on which the phylogeny is based are included in A; B is based on osteological and chromosomal data.

relatively slow-moving (personal observation) animal which is often found in leaf litter. There is no completely suitable comparison within *Anolis*, but in size, color, body shape, and cryptic behavior, *Chamaelinorops* most closely resembles some of the twig-dwarf anole species, and it is compared with *A. sheplani* and *A. occultus*. These species are also of interest because they are thought to be among the most primitive forms in the West Indian anole radiation (Schwartz, 1974; Williams, 1976). Additional comparative data are included for *A. chrysolepis* and a number of other more terrestrial anole species. *Phenacosaurus* is compared with *A. valencienni*. The two forms are similar in body size and shape and in their preference for slow movement on small diameter perches. *A. valencienni*, however, may be a relatively derived anole, and it is not closely related to the mainland species thought to be close to the ancestors of *Phenacosaurus*. The comparison is supplemented with data from the more primitive West Indian species.

The morphological comparisons between *Anolis* species and each of the genera concentrate on several specific issues: 1) phalangeal proportions and the distribution of scales relative to the phalanges; 2) external morphology of the lamellae and generalized subdigital scales; and 3) the dimensions, spacing, and distribution of setae. 4) Lastly, setae appear to be derived from the spinules which characterize the epidermis of many iguanids (Ruibal and Ernst, 1965; Ruibal, 1968; Maderson, 1970). The two morphologies can be linked through a morphotypic series (Olson, 1975) of transitional forms which actually exist in the basal or proximal portion of the lamellae (Peterson and Williams, 1981; Fig. 2; see discussion). Intergeneric differences in the character of the series could provide some of the most interesting insights into the patterns of evolution associated with the origin of the pad complex.

MATERIALS AND METHODS

Table 1 lists the specimens examined along with their body size (measured as snout-vent length) and digital measurements.

Scales and phalanges of the fourth digit are numbered from distal to proximal following Boulenger (1885) and systematic convention rather than anatomical convention (e.g., phalanx i in our terminology is phalanx iv in anatomical nomenclature). The claw base is included in phalanx i scale counts and measurements. Following Peterson and Williams (1981), lamellar scales are defined as those having a thin, frayed free margin and a width to length ratio greater than one.

The distribution of scales relative to the phalanges was determined by flexing and extending the metatarsal-phalangeal and interphalangeal joints. A scale adjacent to the iii/iv interphalangeal joint was counted as a phalanx iii scale if it moved with phalanx iii. This definition may shift the phalangeal boundaries slightly proximal to the transverse plane through the joint, but it decreases ambiguity in the count and emphasizes functional associations. The i/ii interphalangeal joint is synonymous with the distal border of the pad in most anoles. The distal pad scales are often closely packed and project out over the proximal phalanx i region. These pad scales are included with the phalanx ii scales if their bases are attached to phalanx ii or to the interphalangeal joint. In most anoles the pad of the fourth digit lies under phalanges ii and iii, and the scale count for these phalanges is just one scale greater than the total number of lamellae (the most distal pad scale is non-lamellar; Table 1). There are, however, significant differences in the distribution of lamellae relative to the phalanges. These are described in the text, and a separate count of lamellar scales is included in Table 1 for species in which lamellae do not

TABLE 1. SPECIMENS EXAMINED AND PHALANGEAL DIMENSIONS.

Species Specimen	Snout-vent length	Length 4th digit	Total subdigital scales on 4th digit	Length phalanges ii and iii (length of pad)	Subdigital scales ii and iii (pad scales)	Scales ii & iii		Length ii & iii	
						Digit Length	Total scales (pad scales total scales)	Digit Length	Digit Length (pad length snout-vent length)
<i>Chamaeleolis chamaeleontides</i>									
MCZ 57933*	101 mm	12.5 mm	59	5.1 mm	32	0.12	0.54	0.41 (0.05)	
MCZ 59331	155 mm	20.5 mm	60	8.7 mm	31	0.13	0.52	0.42 (0.06)	
<i>Chamaeleolis porcus</i>									
MCZ 4859	156 mm	19.8 mm	57	8.0 mm	31	0.13	0.54	0.40 (0.05)	
<i>Anolis cuvieri</i>									
MCZ 35971	137 mm	27.8 mm	61	10.6 mm	33	0.20	0.54	0.38 (0.08)	
MCZ 35970	122 mm	24.1 mm	58	9.9 mm	31	0.20	0.53	0.41 (0.08)	
MCZ 35976	124 mm	22.8 mm	58	9.3 mm	31	0.18	0.53	0.41 (0.08)	
Additional specimens: MCZ 35975 126 mm; MCZ 61875 120 mm; MCZ 35971 125 mm; MCZ 35978 125 mm; MCZ 35977 131 mm; MCZ 35974 131 mm.									
<i>Chamaelinorops barbouri</i>									
MCZ 155851	31.0 mm	6.4 mm	25	1.0 mm (1.2 mm)	8 (7)	0.21	0.32 (0.28)	0.34 (0.04)	
MCZ 126708	26.5 mm	4.5 mm	28	1.8 mm (0.9 mm)	8 (6)	0.17	0.32 (0.21)	0.42 (0.03)	
MCZ 74594	22.5 mm	4.8 mm	28	1.4 mm (1.0 mm)	8 (7)	0.21	0.39 (0.25)	0.46 (0.04)	
MCZ 57729	18.5 mm	3.4 mm	27	1.3 mm (0.8 mm)	9 (7)	0.18	0.33 (0.26)	0.41 (0.04)	
MCZ 156923	36.5 mm	6.7 mm	26	2.7 mm (1.5 mm)	9 (6)	0.18	0.38 (0.23)	0.42 (0.04)	
MCZ 57728	30.5 mm	5.6 mm	26	1.7 mm (1.0 mm)	9 (7)	0.18	0.35 (0.27)	0.32 (0.03)	
MCZ 156919	31.0 mm	5.5 mm	25	1.8 mm (1.3 mm)	7 (6)	0.18	0.28 (0.24)	0.40 (0.04)	
Additional specimen: MCZ 98691 43 mm.									
<i>Anolis sheplani</i>									
MCZ 125641	40.8 mm	0.3 mm	28	2.0 mm	16	0.07	0.57	0.67 (0.05)	
MCZ 140021	39.5 mm	3.6 mm	28	2.2 mm	16	0.10	0.57	0.61 (0.06)	
<i>Anolis occultus</i>									
MCZ 156968	37.0 mm	4.0 mm	30	2.0 mm	17	0.11	0.57	0.50 (0.05)	
Additional specimens: Po 322 34.5 mm; P 310 37 mm; P 323 39.0 mm; MCZ 101808 36 mm.									
<i>Phenacosaurus heterodermus</i>									
MCZ 104409	79.0 mm	9.3 mm	36	3.3 mm (7.6 mm)	16 (31)	0.12	0.44 (0.86)	0.35 (0.10)	

MCZ 139566	65.0 mm	8.1 mm	32	3.2 mm (6.6 mm)	14 (27)	0.12	0.44 (0.84)	0.40 (0.10)
MCZ 139570	64.5 mm	8.3 mm	30	2.8 mm (7.6 mm)	14 (25)	0.13	0.47 (0.83)	0.32 (0.10)
MCZ 139567	70.0 mm	10.3 mm	37	3.8 mm (8.4 mm)	14 (32)	0.15	0.39 (0.86)	0.37 (0.12)
P 300	56.0 mm	6.9 mm	34	2.2 mm (5.8 mm)	14 (28)	0.12	0.38 (0.84)	0.32 (0.10)
<i>Anolis valencienni</i>								
P 269	61.0 mm	7.8 mm	45	3.8 mm	26	0.13	0.58	0.49 (0.06)
P 329	58.5 mm	7.6 mm	43	3.4 mm	26	0.14	0.60	0.45 (0.06)
<i>Anolis chrysolepis</i> (narrow pad; generalized beta)								
MCZ 152114	59.0 mm	12.5 mm	30	4.6 mm (2.1 mm)	14 (9)	0.21	0.47 (0.30)	0.37 (0.04)
MCZ 152115	47.5 mm	10.3 mm	31	4.1 mm (2.1 mm)	14 (9)	0.22	0.45 (0.29)	0.40 (0.04)
Additional specimen: LACMNH Δ 74485 62 mm.								
<i>Anolis pentapirion</i> (generalized beta)								
MCZ 126154	55.0 mm	9.0 mm	43	3.6 mm	22	0.16	0.54	0.40 (0.07)
<i>Anolis aequatorialis</i> (narrow pad; Norops condition; alpha)								
MCZ 146979	73.0 mm	17.7 mm	54	7.4 mm (8.6 mm)	23 (26)	0.24	0.43 (0.48)	0.42 (0.12)
<i>Anolis tropidonotus tropidonotus</i> (narrow pad; beta)								
MCZ 38798	41.0 mm	8.7 mm	35	3.8 mm (1.8 mm)	16 (10)	0.21	0.46 (0.29)	0.44 (0.04)
<i>Anolis tropidonotus spilorhipis</i> (narrow pad; beta)								
MCZ 53855	41 mm	10.7 mm	34	4.6 mm (2.3 mm)	16 (11)	0.26	0.47 (0.22)	0.43 (0.06)
<i>Anolis barkeri</i> (narrow pad; Norops condition; beta)								
LACMNH 34567	64 mm	12.8 mm	42	5.5 mm	18	0.20	0.43	0.43 (0.09)
<i>Anolis auratus</i> (narrow pad; Norops condition; beta)								
LACMNH 72781	54 mm	8.7 mm	34	3.7 mm (2.1 mm)	14 (9)	0.16	0.41 (0.29)	0.43 (0.04)
Additional <i>Anolis</i> specimens: A. sp. n. near <i>eulaemus</i> LACMNH 42144.								
<i>Laemactis serratus</i> (basiliscine group of Iguanidae)								
MCZ 28745	110 mm	27.3 mm	34	11.0 mm	14	0.25	0.41	0.40
<i>Laemactis longipes</i> (basiliscine group of Iguanidae)								
MCZ 32034	114 mm	31.3 mm	35	12.8 mm	15	0.27	0.43	0.41
<i>Morunasaurus groi</i> (basal or morunasaurine group of Iguanidae)								
MCZ 37876	88 mm	18.5	24	7.8 mm	11	0.21	0.46	0.42
<i>Morunasaurus annularis</i> (basal or morunasaurine group of Iguanidae)								
MCA 146376	106 mm	20.1 mm	26	8.6 mm	11	0.20	0.42	0.41

* Museum of Comparative Zoology, Harvard University.
 o Specimens in the author's collection.
 Δ Los Angeles County Natural History Museum.

occur throughout the phalanx ii and iii regions. In *Chamaeleolis* and some *Anolis* specimens there are not one, but several series of subdigital scales in the phalanx iv region. Many anole specimens also have double, side-by-side scales in the positions just distal to the pad. In these instances only one of the series is counted so that the scale count reported refers to "scale places" rather than the total number of subdigital scales.

Gross measurements were taken with dial calipers (reading to 0.02 mm) or an ocular micrometer. Phalangeal lengths were taken with the ocular micrometer to the nearest 0.1 mm. The accuracy of the measurement partially depends on the condition of the specimen and error estimates vary. The standard deviation of six replicate measurements of the length of phalanges ii and iii was 0.1 mm (4–5% of the mean value) in five error trials (specimens of *Phenacosaurus*, *A. occultus*, *A. sheplani* and *Chamaelinorops*) and 0.2 mm (11% of the mean value) in one trial (specimen of *Chamaelinorops*). Phalangeal length measurements were taken along the subdigital scale row. Actual lengths were defined by the scale borders rather than the joint axes. For example, the length of phalanx iv was defined as the distance from the base of the most proximal phalanx iv scale to the distal margin of the most distal phalanx iv scale. The length of phalanges ii and iii is in most cases equivalent to the length of the pad region (Table 1). A separate measurement of the length of the pad (= the length of the series of lamellar scales) is included for species in which this is not the case.

The digits were prepared for scanning electron microscopy as described in Peterson and Williams (1981). Terminology for the fine structure follows that of Peterson and Williams (1981; summarized in Fig. 2), Hiller (1968), Maderson (1970) and Ruibal and Ernst (1965). Measurements of fine structure were taken from photographs and are reported

as the mean of 12 to 30 individual measurements with the 95% confidence interval (for some spinule measurements $n=6$). Magnification was adjusted so that measurement with a dial caliper to the nearest 0.1 mm resulted in measurement to the nearest 0.01 μm for stalk diameter, seta tip dimensions and spinule dimensions or to the nearest 0.1 μm for stalk height. The measurement techniques for fine structure are judged to be adequate based on F-tests of the variance among individual setae compared to the variance among replicate measures (F value in each test significant at the 0.001 level).

The major difficulty in characterizing and comparing species in terms of setal dimensions is to obtain a realistic sample of measurements. Measurements taken within about a 30 μm radius on one specimen are quite consistent and confidence intervals are small. Measurements from one or even two such areas underestimate regional variation, particularly in setal stalk height which increases and then decreases again along a proximodistal transect on each lamella (e.g., in *A. cuvieri* [MCZ 61875] mean stalk height for the central and proximal regions of a mid-pad lamella are $16.6 \pm 1.8 \mu\text{m}$ and $24.4 \pm 0.4 \mu\text{m}$, respectively). Because sampling from a large number of areas is prohibitive in a comparative survey, the interspecific and intraspecific comparisons are made for setae from homologous regions of mid-pad lamellae. The mean value for an individual specimen is based on measurements taken from two to four of these regions.

The measurements of stalk dimensions are taken from the center of a mid-pad lamella on setae adjacent to the junction between the lamellar core and the free margin (see Ruibal and Ernst, 1965, or Lillywhite and Maderson, 1968, for further definition of this region). These are usually the tallest setae on the lamella (Ruibal and Ernst, 1965). Variation in stalk diameter is smaller and appears to be less well correlated with

Figure 2. The morphotypic spinule-spine (left) and spine-seta (right) series of *Anolis* (from Peterson and Williams, 1981).

lamellar region so that the thickness of the setal stalk at the lamellar/free margin junction bears no consistent relationship to the range of dimensions (Peterson and Williams, in preparation). Stalk height is defined as the distance between the stalk base on the lamellar surface and the spatulate seta tip. Stalk diameter is measured at a point between one third and one half of the distance between the stalk base and the spatulate tip. The stalk tapers little in this region, and some flexibility in the position of the diameter measurement is necessary because of the presence of debris and the crossing of stalks.

Measurements of the tip dimensions are subject to a) regional variation (e.g., in *A. cuvieri* [MCZ 61875] tip area at the distal margin is 0.165 ± 0.023 sq μm while that midway between the free margin and the base of the lamella is 0.248 ± 0.039 sq μm) and b) variation apparently related to wear or age of the Oberhautchen (= functional epidermal layer; Maderson, 1970). In an *A. occultus* specimen (MCZ 156968) the spatulate tips over the central portion of the lamellae are extremely frayed and small (tip area measures 0.175 ± 0.014 sq μm) while those near the lateral margins of the lamella are much less ragged and larger (tip area measures 0.237 ± 0.020 sq μm). Similar medio-lateral variation in tip area does not occur in *Anolis* specimens in which the central setae appear relatively "unworn." Wherever possible the setal tip measurements have been taken from the central portion of the pad, but in several cases the area was shifted laterally to minimize what appears to be individual variation related to wear or age of the Oberhautchen. In all cases the tip dimensions are taken in regions midway along the proximodistal axis of the lamella. Tip area is calculated from measurements of the width of the seta margin and the height (= distance between the midpoint of the margin and the junction between the flattened tip and the oval distal region of the stalk). In

most anoles the tip is triangular, and area can be estimated as $0.5 (\text{width} \times \text{height})$. In *Chamaelinorops* the tip is square or rectangular, and area is estimated as $\text{width} \times \text{height}$.

The spacing or density of setae is determined by counting the setae over 250 to 750 sq μm . The entire lamella is usually folded along its proximo-distal axis, and in most cases the measurement is based on a series of 120 to 250 sq μm photographs taken midway up a gentle slope where the surface is normal to the viewing plane. This corresponds to a position midway between the free margin of the lamella and its base.

The dimensions and density of spinules were measured on phalanx I scales and/or on a lateral digital scale immediately adjacent to the subdigital scales, both areas were examined. Spinule height is total elevation above the scale surface. Spinules which are less than 0.5 μm tall are nubbinlike, and diameter is measured at their midpoint. Density is determined by counting the spinules over 64 to 100 sq μm .

RESULTS OF COMPARISONS

COMPARISONS BETWEEN *CHAMAELEOLIS* AND *ANOLIS CUVIERI*

Gross Structure and Pad Proportions

The phalangeal dimensions and scale distribution of *Chamaeleolis* are not different from those in *Anolis*. The fourth toe is shorter in *Chamaeleolis* than in *A. cuvieri* (Table 1), but a number of *Anolis* species, including *A. valencienni*, *A. occultus*, and *A. sheplani* are not different from *Chamaeleolis* in this respect. The phalanges that underlie the pad (ii and iii) are absolutely shorter in *Chamaeleolis*, but the proportion of pad length/toe length is similar in *Chamaeleolis* and *A. cuvieri* (Table 1). *Chamaeleolis* has a shorter pad in relation to its body length than *A. cuvieri*, but several other anoles

(e.g., *A. sheplani* and *A. valencienni*) exhibit proportions similar to those of *Chamaeleolis*. The total number of subdigital scales and the number of pad scales are similar in *Chamaeleolis* and *A. cuvieri* (Table 1).

The subdigital scales in both forms exhibit three different morphologies corresponding to a) the pad region (phalanges ii and iii); b) the phalanx i region; and c) the phalanx iv region. In *Chamaeleolis* and *A. cuvieri* all but one (see below) of the scales in the pad region are lamellae, and lamellae are absent in the phalanx i and iv regions. The morphology of the lamellar scales is indistinguishable in the two forms (*Chamaeleolis* lamellae are shown in Fig. 3A). In most specimens one of the small scales that is inserted between the phalanx i and phalanx ii series to "raise" the pad is exposed on the subdigital surface and forms the most distal pad scale. This scale does not have the characteristic free margin of a lamella (it also bears spikes rather than setae). Generalized subdigital scales occur in the phalanx i and iv regions. Those in the phalanx i region wrap around the phalanx more tightly giving them a more convex cross section. The phalanx iv scales in *Chamaeleolis* are very irregular in shape, and a distinct subdigital scale row is not present (Fig. 3B).

Fine Structure

Setae cover the lamellae in both *A. cuvieri* and *Chamaeleolis* (Table 2; Figs. 3C, 3D, 4A, B). The setae stalks are slightly shorter in *Chamaeleolis*, but stalk diameter, tip shape and the density of setae are not different. With the exception of *Chamaeleolis* MCZ 59331, tip area is also comparable in the two forms. *Chamaeleolis* MCZ 59331 has a much larger tip (Table 2). Measurements taken proximally and distally on the pad and over the width of four of the mid-pad lamellae indicate that the enlarged tip is not confined to a small region. It appears to be an individual variation.

In all the specimens the complete spine-seta subdigital series shown in Figure 2 is present in the proximal portion of the lamella.

While the morphology of the pad region is very similar in *Chamaeleolis* and *A. cuvieri*, the phalanx i and iv regions exhibit some differences. The phalanx i scales are grossly similar in the two forms, but their fine structure is different. In *Chamaeleolis* the regions adjacent to the distal margin of the scale are covered by densely packed, small spines 1–2 μm tall and 0.3–0.4 μm in diameter). Spine height decreases toward the base of the scale. The proximal one-half to two-thirds of the scale is covered by spinules (Figs. 3E, F). This area closely resembles the spinulate Oberhautchen (Fig. 2 far left). There are two slightly different populations of spinules (Fig. 3F). The taller ones are $0.79 \pm 0.08 \mu\text{m}$ tall and $0.22 \pm 0.02 \mu\text{m}$ in diameter. The shorter, nubbinlike spinules are $0.34 \pm 0.02 \mu\text{m}$ tall and $0.22 \pm 0.02 \mu\text{m}$ in diameter. Together they occur in a density of 7.0 spinules/sq μm . This morphology is present on all the phalanx i scales; there is no apparent regional differentiation in fine structure. The spinulate phalanx i region is much more conservative in terms of the morphotypic series (Fig. 2) than the lateral digital scales adjacent to the pad. The lateral scales are covered with spikes, prongs, spines, and the seta-prong intermediate morphotype (Figs. 2, 4F). The dimensions and density of these forms are comparable to those of the lamellar series in *Chamaeleolis* and to those described for the generalized anole subdigital series in Figure 2.

In *A. cuvieri* the phalanx i scales are largely covered with spikes and prongs ($10.4 \pm 1.04 \mu\text{m}$ tall and $0.53 \pm 0.07 \mu\text{m}$ in diameter; Fig. 4E). Small spines (Fig. 2), similar to those in *Chamaeleolis*, do occur in *A. cuvieri*, but only in extremely proximal areas of the scales. Spinulate areas are most extensive on the scales just distal to the pad over the phalanx i region. Regions of small spines also occur

Figure 3. The subdigital morphology of *Chamaelinorops*. In this and all subsequent photographs the small arrow adjacent to the figure scale points in the direction of the claw. A. Lamellae numbers 3, 4 and 5 in MCZ 57933. The depth of the frays along the distal margin coincides with the limit of the free margin. B. The phalanx iv subdigital scales in MCZ 59331. C. Oblique view of the setae tips in a mid-pad region of MCZ 57933. D. Setae stalks on the free margin of a lamella in MCZ 57933. E. The surface of phalanx i scale 11 in MCZ 8459. The borders of the epidermal cells are raised, spine-covered ridges. This is the only specimen of an anoline in which this morphology rather than a bare trackway (Ruibal, 1968) has been found. The morphology does occur in a number of non-anoline iguanids. F. Spinules on the proximal portion of phalanx i scale 8. Note the presence of spinelike and nubbinlike spinules.

TABLE 2. COMPARISON OF SETA DIMENSIONS.

Species/Specimen	Seta stalk		Seta tip	
	diameter	height	area	density
<i>Chamaeleolis chameleontides</i>				
MCZ 57933	$0.56 \pm 0.03 \mu\text{m}$	$18.4 \pm 0.8 \mu\text{m}$	$0.206 \pm 0.021 \text{ sq } \mu\text{m}$	$1.0/\text{sq } \mu\text{m}$
MCZ 59331	$0.58 \pm 0.03 \mu\text{m}$	$18.6 \pm 0.3 \mu\text{m}$	$0.377 \pm 0.039 \text{ sq } \mu\text{m}$	$0.7/\text{sq } \mu\text{m}$
<i>Chamaeleolis porcus</i>				
MCZ 8459	$0.53 \pm 0.02 \mu\text{m}$	$22.3 \pm 0.7 \mu\text{m}$	$0.264 \pm 0.019 \text{ sq } \mu\text{m}$	$1.0/\text{sq } \mu\text{m}$
<i>Anolis cuvieri</i>				
MCZ 61875	$0.51 \pm 0.04 \mu\text{m}$	$22.4 \pm 0.5 \mu\text{m}$	$0.229 \pm 0.025 \text{ sq } \mu\text{m}$	$1.0/\text{sq } \mu\text{m}$
MCZ 35975	$0.60 \pm 0.02 \mu\text{m}$	$23.9 \pm 0.5 \mu\text{m}$	$0.232 \pm 0.023 \text{ sq } \mu\text{m}$	$0.9/\text{sq } \mu\text{m}$
MCZ 35971	$0.65 \pm 0.04 \mu\text{m}$	$27.2 \pm 1.4 \mu\text{m}$	$0.253 \pm 0.017 \text{ sq } \mu\text{m}$	$0.9/\text{sq } \mu\text{m}$
MCZ 35977	$0.51 \pm 0.06 \mu\text{m}$	n.a.*	$0.184 \pm 0.020 \text{ sq } \mu\text{m}$	$1.4/\text{sq } \mu\text{m}$
<i>Chamaelinorops barbouri</i>				
MCZ 155851	$0.52 \pm 0.05 \mu\text{m}$	$6.9 \pm 0.7 \mu\text{m}$	$1.014 \pm 0.089 \text{ sq } \mu\text{m}$	$0.5/\text{sq } \mu\text{m}$
MCZ 68691	$0.55 \pm 0.05 \mu\text{m}$	n.a.	$0.649 \pm 0.060 \text{ sq } \mu\text{m}$	$0.5/\text{sq } \mu\text{m}$
MCZ 156919	$0.54 \pm 0.05 \mu\text{m}$	$8.4 \pm 0.4 \mu\text{m}$	$1.472 \pm 0.149 \text{ sq } \mu\text{m}$	$0.6/\text{sq } \mu\text{m}$
MCZ 126708	$0.56 \pm 0.06 \mu\text{m}$	$7.0 \pm 0.5 \mu\text{m}$	$0.759 \pm 0.072 \text{ sq } \mu\text{m}$	$0.5/\text{sq } \mu\text{m}$
MCZ 74594	$0.47 \pm 0.03 \mu\text{m}^\Delta$	$5.0 \pm 0.3 \mu\text{m}^\Delta$	$0.600 \pm 0.047 \text{ sq } \mu\text{m}$	$0.5/\text{sq } \mu\text{m}$
<i>Anolis sheplani</i>				
MCZ 140021	$0.39 \pm 0.02 \mu\text{m}$	$12.6 \pm 0.3 \mu\text{m}$	$0.220 \pm 0.020 \text{ sq } \mu\text{m}$	$1.1/\text{sq } \mu\text{m}$
MCZ 125641	$0.41 \pm 0.03 \mu\text{m}$	$11.8 \pm 0.5 \mu\text{m}$	$0.279 \pm 0.023 \text{ sq } \mu\text{m}$	$1.2/\text{sq } \mu\text{m}$
<i>Anolis occultus</i>				
MCZ 156968	$0.49 \pm 0.04 \mu\text{m}$	$11.0 \pm 0.7 \mu\text{m}$	$0.237 \pm 0.020 \text{ sq } \mu\text{m}$ [or $0.175 \pm 0.024 \text{ sq } \mu\text{m}$ in "worn" regions]	$1.4/\text{sq } \mu\text{m}$
<i>Anolis</i> sp. n. near <i>eulaemus</i>				
LACMNH 72741	$0.57 \pm 0.06 \mu\text{m}$	$20.4 \pm 0.3 \mu\text{m}$	$0.593 \pm 0.045 \text{ sq } \mu\text{m}$	$1.0/\text{sq } \mu\text{m}$
<i>Phenacosaurus heterodermus</i>				
MCZ 35970	$0.41 \pm 0.02 \mu\text{m}$	$14.6 \pm 0.2 \mu\text{m}$	$0.226 \pm 0.028 \text{ sq } \mu$	$1.2/\text{sq } \mu\text{m}$
MCZ 139566	$0.46 \pm 0.02 \mu\text{m}$	$14.8 \pm 0.3 \mu\text{m}$	$0.214 \pm 0.021 \text{ sq } \mu$	$1./2\text{sq } \mu\text{m}$
P 300	$0.45 \pm 0.09 \mu\text{m}$	$13.7 \pm 0.4 \mu\text{m}$	$0.308 \pm 0.025 \text{ sq } \mu$	$1.2/\text{sq } \mu\text{m}$
<i>Anolis valencienni</i>				
P 329	$0.47 \pm 0.03 \mu\text{m}$	$17.2 \pm 0.6 \mu\text{m}$	$0.209 \pm 0.016 \text{ sq } \mu\text{m}$	$1.1/\text{sq } \mu\text{m}$
P 269	$0.40 \pm 0.02 \mu\text{m}$	$15.3 \pm 0.4 \mu\text{m}$	$0.171 \pm 0.018 \text{ sq } \mu\text{m}$	$1.4/\text{sq } \mu\text{m}$

* Measurement not available.

 Δ Measurements taken on the third digit pad.

on the lateral digital scales where they form a landscapelike mosaic with small bare areas (Fig. 4C, D). The dimensions of the small spines ($0.71 \pm 0.08 \mu\text{m}$ tall and $0.35 \pm 0.05 \mu\text{m}$ in diameter) are similar to those of the small spines or larger spinules in *Chamaeleolis*.

The major difference between the spinate areas in *A. cuvieri* and the spinulate regions in *Chamaeleolis* is that in the spinate areas the understory of nubbin-like spinules is very reduced and the density of spines-spinules is about half

that in *Chamaeleolis* (3.6 spinules/ $\text{sq } \mu\text{m}$ compared to 7.0 spinules/ $\text{sq } \mu\text{m}$). In terms of the morphotypic series, the most conservative subdigital scale surface in *A. cuvieri* is less conservative than the surface of the phalanx i scales in *Chamaeleolis*.

Although the phalanx iv scales in *Chamaeleolis* are grossly very different from lamellae, they are covered with setae. The seta stalk height is less than that of lamellar setae ($11.4 \pm 0.3 \mu\text{m}$ tall; cf. Table 2), but the diameter (0.58 ± 0.02

Figure 4. The subdigital morphology of *A. cuvieri* (A-E) and *Chamaeleolis* (F). A. Lateral view of the setae on a mid-pad lamella in MCZ 35971. B. The setae tips in MCZ 35971. C. The surface of a lateral digital scale in MCZ 35975. Smooth bare area similar to that on the crest of a keel occurs to the lower right (distal portion of the scale). The remaining surface is covered with fields of spines. The fine irregular lines separating the fields appear to be the borders of the epidermal cells. D. Detail of the surface of a lateral digital scale showing portions of two small bare areas and a variety of spine morphologies (MCZ 35975). E. Seta-prong intermediate tips in the phalanx iv region of MCZ 35971. F. Seta-prong intermediate tips on the lateral digital scale of *Chamaeleolis* MCZ 8459. Tips which are turned (lower center and upper right) show compression of the tip (cf. setae Fig. 3C).

μm and tip size shape are similar to those of lamellar setae.

The phalanx iv region in *A. cuvieri* consists of well-defined series of generalized subdigital scales. The distal scales bear setae, but within the distal third of the phalanx (between scales 46 and 49), the triangular seta tip is replaced by the flattened, rectangular tip of the seta-prong intermediate morphotype (Figs. 4E, and 2). The width of the tip is about $0.61 \mu\text{m}$ compared to $0.79 \mu\text{m}$ for lamellar setae. At the metatarsal-phalangeal joint, the seta-prong intermediates are $14.5 \pm 1.0 \mu\text{m}$ tall and $.52 \pm .14 \mu\text{m}$ in diameter.

COMPARISONS AMONG *CHAMAELINOROPS*,
A. SHEPLANI AND *A. OCCULTUS*,
AND COMPARISONS AMONG
CHAMAELINOROPS, *A. CHRYSOLEPIS*,
A. AURATUS, AND *A. SP. N. NEAR EULAEMUS*

Gross Morphology of the Digit
and Scales

The gross morphology of the digit in *Chamaelinorops* is significantly different from that in *A. sheplani* and *A. occultus*, although it resembles that of *A. chrysolepis* and *A. auratus* in most respects. The comparison with *A. sheplani* and *A. occultus* is described first.

A. sheplani and *A. occultus* are very similar. A pad is formed by lamellae under phalanges ii and iii (scales 7 through 22 in Fig. 5B; scale 6 is a non-lamellar pad scale). The pad is broad and occupies at least half the length of the toe (Table 1). More than half the subdigital scales are lamellae (Table 1), and generalized subdigital scales occur in the phalanx i (scales 1–5) and phalanx iv (scales 23–30) regions.

In *Chamaelinorops* the pad has a different position relative to the phalanges: it is reduced in size, and less than a quarter of the subdigital scales are pad scales (Table 1; Fig. 5A). a) In *Chamaelinorops* the pad lies under the first and second phalanges rather than the second and third phalanges. The distal 2 to 4 pad

scales lie under phalanx i (scales 8–10 in Fig. 5A), and the proximal 3 to 4 scales lie under phalanx ii (scales 11–13 in Fig. 5A). All the phalanx ii scales are lamellae, but none of the phalanx iii scales have the shape or marginal features of a lamella (scales 14–17 in Fig. 5A correspond to the phalanx iii region). b) The pad is smaller in *Chamaelinorops*. Pad area estimated from the dimensions of the individual lamellae is 0.41 sq mm in *Chamaelinorops* (MCZ 156919; SVL 31 mm) and 0.69 sq mm in *A. sheplani* (MCZ 125641; SVL 40.8 mm). A scaling equation relating pad area and body size (measured as snout-vent length [trunk height]²) which was developed for six Puerto Rican anole ecomorphs (including *A. occultus* and *A. cuvieri*) would predict a pad area of 0.82 sq mm in the *Chamaelinorops* specimen (200% of the observed value) and 0.90 sq mm in the *A. sheplani* specimen (130% of the observed value) (the equation is: $\text{pad area} = \text{constant} [\text{body size}]^{0.656 \pm 0.063}$ [$r=0.963$ for 36 specimens in 6 species] [Peterson *et al.*, 1982, and Peterson, unpublished data]). While both specimens have small pads for their size compared to the Puerto Rican ecomorphs, the discrepancy is much greater for *Chamaelinorops*. c) Differences in phalangeal length and lamellar morphology also contribute to the smaller pad in *Chamaelinorops*. The “pad” phalanges ii and iii are absolutely shorter in *Chamaelinorops*, although the whole digit is almost twice as long as that of *A. occultus* and *A. sheplani* (Table 1). Pad width is also lower in *Chamaelinorops* because the lamellae are much narrower (see below). d) The smaller number of pad scales (Table 1) is partially correlated with the shorter pad, but the shape of the lamellae also contributes to the difference. In *A. sheplani* and *A. occultus*, and to a greater extent in *A. cuvieri* and *A. valencienni* (Fig. 9B), the distal phalanx ii lamellae are very short (proximodistal dimension) so that many lamellae are packed into a short phalangeal length (Figs. 5B, 9). It is the shape of

Figure 5. The subdigital surface of the fourth digit in A) *Chamaelinorops* (MCZ 156919) and B) *A. occultus* (MCZ 156968). In A only the distal 23 of 25 scales are shown. The pad corresponds to scales 8–13. The phalangeal scale series are: 1–10 phalanx i, 11–13 phalanx ii, 14–17 phalanx iii, and 18–25 phalanx iv. In *A. occultus* (B) the regional scale series are: 1–5 phalanx i, 6–22 phalanges ii and iii, and 23–30 phalanx iv. Scale 6 is a non-lamellar pad scale. The morphology of scale 22 is transitional between that of lamellae and generalized scales.

the scales, rather than particularly long pad phalanges, that account for the high proportion of pad to total subdigital scales in most *Anolis* (cf. phalangeal and scale ratios for non-anole iguanids and *A. cuvieri* in Table 1). In *Chamaelinorops* the lamellae are not much shorter than the generalized subdigital scales and only three or four pad scales occur in the phalanx ii region. The width/length proportion of *Chamaelinorops* lamellae is lower than that of *A. sheplani* and *A. occultus* lamellae (mean width/length for the *Chamaelinorops* [MCZ 156919] lamellae is 1.9 while that for *A. sheplani* [MCZ 125641] is 5.0; the maximum for an individual lamella in *Chamaelinorops*, 2.4, is much less than the minimum for *A. sheplani*, 4.0). e) Lamellae are also distinguished from generalized subdigital scales by the presence of a free margin (=that portion of the scale which is distal to the dermal core and consists largely of the outer and inner Oberhautchen layers; Lillywhite and Maderson, 1968). The free margin is poorly developed in *Chamaelinorops*. In *A. sheplani* and *A. occultus* the free margin on phalanx ii lamellae is 20–40 μm long and one-quarter or more of the exposed length of the scale is actually free margin (the limit of the free margin is often indicated by the deeper frays along the lamellar edge; Figs. 4A, 5, 8, 9A, 10A). In *Chamaelinorops* a free margin is present only over the central portion of the scale. It extends for 10 μm proximal to the edge and accounts for less than 5% of the lamellar length. Scales 8 and 13 barely have a free margin; the two layers of Oberhautchen meet at an angle between 45 and 90° and are not adpressed to form the typical bilayer free margin. In both shape and marginal characteristics, *Chamaelinorops* lamellae are convergent on generalized subdigital scales.

The poor differentiation of lamellae from generalized subdigital scales is just one of the factors that contributes to poor differentiation of regional scale series in *Chamaelinorops*. In *A. sheplani* and *A.*

occultus the distal border of the pad is clearly defined by an abrupt change in scale shape and by a series of small scales set deep between the phalanx i and ii scales which "raise" the pad. In *Chamaelinorops* these deep scales are absent and the pad is coplanar with the distal phalanx i scales (=the "Norops" condition). Similarly there is no abrupt change in scale morphology. The distal pad scale (Number 8 in Fig. 5A) is intermediate in shape between the adjacent lamella and the phalanx i scale 7. The scales distal to the pad have a "velvety" texture over their central area similar to that of lamellae, and scales 3 to 5 have very small free margin (Fig. 7A). The central portion of the phalanx i scales is convergent on the lamellar scales. The lateral aspect of the distal phalanx i scales is covered with hillocks (Peterson and Williams, 1981) and low keels. The scales under the third phalanx (numbers 14–17 in Fig. 5A) and those under the proximal portion of the fourth phalanx (numbers 21–25) are very similar. They are so narrow that the lateral digital scales encroach on the subdigital surface. The distal phalanx iv scales (numbers 18–21) are wider and resemble the generalized subdigital scales in the phalanx iv region of *A. sheplani* and *A. occultus*.

While there are few points of similarity between *Chamaelinorops* and *A. sheplani* or *A. occultus*, the gross morphology of the *Chamaelinorops* pad is not entirely unusual among anoles. a) The "unraised" pad occurs in a variety of species groups (e.g., *A. aequatorialis*, *A. barkeri*, and *A. auratus* in Table 1). *A. chrysolepis* exhibits an intermediate condition. b) Many anole and non-anole species share the phalangeal proportions of *Chamaelinorops* (e.g., *A. chrysolepis*, *A. cuvieri* and *Morunasaurus*; Table 1). Indeed *A. sheplani* and *A. occultus* have unusually short toes with relatively long second and third phalanges compared to *A. cuvieri* and *A. chrysolepis*; it is *Chamaelinorops* that appears to exhibit the more generalized phalangeal and

Figure 6. The morphology of lamellae and setae in *Chamaelinorops*. A. Lamellae MCZ 156919. Note that the lamellar margin appears thick and has very shallow frays (cf. Fig. 7A). B. Typical lamellar setae in MCZ 155851 (viewed from the sole of the foot looking toward the claw). Note the large rectangular tip and the absence of an inflection in the stalk. C. Setae on generalized subdigital scale number 25 in MCZ 155851. The setae (also shown in E and F) are unusual because they display a consistent tip flexion. The increase in stalk diameter which occurs about midway along the stalk length marks the beginning of the oval stalk region. A few setae which are turned (e.g., lower center) show the abrupt compression which forms the tip. D. Lamellar setae in MCZ 155851. These setae exhibit considerable wear of the tip, but also show the irregular grooves in the stalk and relatively thicker tip of *Chamaelinorops*. E. Lateral view of the setae on or scale 25 in MCZ 155851. Setae in profile show the irregular stalk, thicker tip, and the abrupt compression of the oval region which forms the tip. Note that the inflection occurs within the flattened tip region and that the distal margin of the tip is often flexed ventrally. F. Normal view of the setae tips on scale 25 in MCZ 155851. Note the rectangular tip shape, reduced density and thick, "rolled" distal margin.

Figure 7. Subdigital morphology in *Chamaelinorops*. A. Phalanx i scales 2 through 4 in MCZ 155851. The small free margin on scale 3 is evident toward the lower portion of the figure. B. The distal margin of scale 4 at the transition between the subdigital and lateral portions of the scale. The "seta" tips are small and the margin itself is studded with large flattened spines. Note how adjacent spines are fused (MCZ 155851). C. The lateral portion of a lamella and the adjacent lateral digital scale in MCZ 98691. The lateral digital scale is contoured by hillocks and several small keels. The distal portion has many large irregular spines. D. Detail of the lateral digital scale showing moderately large spines (lower left) and large diameter spines (top) on adjacent hillocks. Note the irregular, apparently fused spine at the top right (MCZ 98691). E. Proximal and lateral portion of a lamella in MCZ 155851. Spines to the right have a $0.5/\text{sq mm}$ spacing and appear to form the base of the *Chamaelinorops* spine-seta series. Spines to the top left and lower center have a similar spacing, but a much larger diameter. They may form the base of the large spine series. Spines similar to those shown in the lower left have a $1/\text{sq mm}$ density and resemble *Anolis* spines; they may form the "top" of the spinule-spine series in *Chamaelinorops*. F. Adjacent surfaces of a phalanx i and lateral digital scale in MCZ 155851. The cleft between the scales and on a hillock just above the figure scale are covered with spinules and small spines. The adjacent epidermal cells exhibit a variety of large spines. Note in the upper left the large, irregular, apparently fused spines.

digital proportions. c) The *Chamaelinorops* pattern of lamellar distribution is approached by a number of anole species. In *A. chrysolepis* and *A. tropidonotus* lamellae are absent in the phalanx iii region. In *A. auratus* only the distal two phalanx iii scales are lamellae (the West Indian grass anole, *A. pulchellus*, exhibits a less extreme reduction in lamellae). d) Pad area in proportion to body size is similar in *A. auratus* and *Chamaelinorops*. For *A. auratus* (LACMNH 72741; SVL 54 mm) pad area is 1.2 sq mm. The scaling equation described above would predict 2.0 mm from the body dimensions (167% of the observed value). *A. chrysolepis* and *A. tropidonotus* have only slightly larger pads in proportion to their size. e) A similar set of conditions contribute to the small pad of *A. auratus* and *Chamaelinorops*. Pad length is reduced because of the restricted distribution of lamellae, and the lamellae themselves are longer and narrower than in *A. occultus* and *A. sheplani*. The mean width/length proportion of the *A. auratus* lamellae is 2.5 with a maximum value of 3.3 (cf. *A. sheplani* above). Pad length relative to digital length or body length is similar among *Chamaelinorops*, *A. auratus*, *A. chrysolepis*, and *A. tropidonotus* (Table 1). f) The proportion of lamellar to total subdigital scales is similar among *A. auratus*, *A. chrysolepis*, *A. tropidonotus*, and *Chamaelinorops* (Table 1). The restricted distribution of lamellae as well as the greater length of the individual lamellar scales contributes to the condition in each case. g) *A. auratus* has relatively poor regional differentiation of the scale series (Peterson and Williams, 1981). Lamellae grade into generalized subdigital scales in the phalanx iii and iv regions. The distal phalanx i scales are remarkably similar to those of *Chamaelinorops*. Low keels occur over the lateral portion of the scale while the central portion has a "velvet-like" texture (Peterson and Williams, 1981).

Fine Structure

Seta Shape and Dimensions

The "setae" in *Chamaelinorops* are not setae as defined by *A. sheplani*, *A. occultus* (Fig. 8B, C) or any other anole examined to date. The seta tip is rectangular (Fig. 6), as is the anole seta-prong morphotype (Figs. 2, 4E, F), but the shape and dimensions of the *Chamaelinorops* setae are quite distinct from those of the anole seta-prong or seta.

a) The subdigital spine-seta morphotypic series in *Anolis* appears to be based on a fairly constant stalk density—slightly greater than 1/sq μm (values for individual specimens in this study range from 0.9–1.4 stalks/sq μm ; Table 2; Peterson *et al.*, 1982 report that there is no significant difference in stalk density among 6 Puerto Rican ecomorphs including *A. occultus* and *A. cuvieri*). Setal stalk density in *Chamaelinorops* also appears to be relatively constant, but at 0.5 stalks/sq μm , it is roughly half the value for the anole species or *Chamaeleolis* (Table 2).

b) The diameter of the *Chamaelinorops* seta stalks is not different from that of *A. cuvieri* or *A. occultus*, but stalk height is less than that of any anole setae examined to date (Table 2; Peterson and Williams, 1981). The stalks also have a noticeably irregular surface (Fig. 6B, D, E).

c) *A. sheplani* and *A. occultus* are typical of anoles in the shape and size of the tip and in the presence of well-defined stalk and tip regions (Fig. 8B, C and Table 2). In *Chamaelinorops* the stalk and tip regions are less well defined by shape change or by flexion of the stalk, and the tip itself forms in a different fashion.

In *Chamaelinorops* and in anoles such as *A. occultus*, *A. sheplani*, and *A. cuvieri*, the flattened tip lies just distal to a portion of the stalk that is oval in cross section. In *Chamaelinorops* this oval section is 2.0 to 2.8 μm long (roughly one-

Figure 8. Subdigital morphology in *A. sheplani* and *A. occultus*. A. The surface of midpad lamellae (numbers 5 and 6) in *A. sheplani* (MCZ 125641). B. Lateral view of the setae on the free margin of lamella number 5 in *A. sheplani* (MCZ 125641). C. Setae tips at the proximal portion of the lamellar free margin in *A. sheplani* (MCZ 125641). D. Setae-prongs on the most proximal digital scale (number 30) in *A. occultus* (MCZ 156968). E. Oblique view of the spines adjacent to the distal margin of phalanx i scale 4 in *A. occultus* (MCZ 156968). F. Spinules over the proximal portion of phalanx i scale 2 in *A. occultus* (MCZ 156968). Note the presence of nubbinlike and spinelike spinules.

third of the stalk length); in the anoles it is 1.0 to 1.5 μm long (or 5–10% of the stalk length). The major axis of the oval cross section corresponds to the proximo-distal axis of the digit, and this orientation appears to establish the preferred axis for flexion of the tip on the stalk. In the anoles the junction between the round and oval stalk sections is frequently the site of a small inflection which tilts the tip toward the metatarsal—phalangeal joint. The distalmost section of the oval stalk forms the apex of the triangular tip and is the site of a second more frequent and larger inflection that tilts the seta tip 30 to 70° and points the distal margin proximally toward the metatarsal-phalangeal joint (stalks in the upper right of Figure 8B are relatively undisturbed and show this orientation clearly). In *Chamaelinorops* the stalk, including the oval region, is usually straight and normal to the scale surface (Fig. 6B, D). Most of the *Chamaelinorops* setae have no inflection of the tip on the stalk; the flat tip of most lamellar setae is oriented normal to the scale surface and locomotor substrate. In a few regions where there is a consistent tip flexion, the point of flexion lies within the tip region itself, just distal to where the tip differentiates from the stalk (Fig. 6). The inflection tilts the tip toward the metatarsal-phalangeal joint as in *Anolis* or *Chamaeleolis*. There is some taper or reduction in the cross sectional area of the stalk between the round and oval regions and over the length of the oval region in *A. cuvieri*. In *Chamaelinorops* the stalk shape changes slightly, but it is not tapered proximal to the tip.

In the anole species the triangular tip appears to be formed as the oval stalk gradually flattens over the 0.6 μm length of the tip. Relatively gradual compression of the oval cross section produces a triangular surface 0.6–0.8 μm wide (2–3X wider than the distalmost section of the oval). Often it appears that compression of the tip begins in the center and spreads laterally so that in the proximal

portion of the tip the lateral margin is thicker.

In *Chamaelinorops* a tip forms by a more abrupt decrease in the minor axis of the oval stalk (from 0.5–0.6 μm to 0.2–0.3 μm) coupled with a variable increase in the major axis (from 0.6–0.7 μm to 1.0–0.7 μm) (Fig. 6D, E). The amount of material in a cross section through the proximal portion of the tip is very close to that present in any cross section of the stalk. More distally the tip tapers to 0.15 to 0.25 μm thickness (the minor axis of the oval), and in three of the five specimens the tip narrows to 0.5 to 0.6 μm (or even slightly less at the very distal margin; see the paragraphs at the end of this section for discussion of the individual variation in *Chamaelinorops* specimens). The length of the tip is subject to obvious wear variation, but mean length for the five specimens ranges from 1.0 to 1.5 μm . The *Chamaelinorops* seta tip is approximately twice as long and 2 to 3 times thicker than that in the anole species examined. The average tip width is greater than that in anoles, but the width at the distal margin is less. The distal margin of the seta in *Chamaelinorops* is formed by a chipped, uneven edge (Fig. 6B, D) or by a ventrally curled “lip” (Fig. 6C, E, F).

There are a number of very clear differences between the seta tip in generalized anoles and *Chamaelinorops*. In *Chamaelinorops*, 1) there is no taper or reduction in the cross sectional area of the stalk proximal to the tip; 2) the tip appears to form by a more abrupt compression of the oval stalk; 3) this produces a rectangular or a trapezoidal tip surface (rather than a triangular one), 4) which is 3 to 6 times larger than that of the anoles; 5) the tip is substantially wider, longer and thicker, but the distal margin is narrower; 6) the tip is usually oriented parallel to the seta stalk, but may have an orientation similar to that of anoles; 7) if an inflection is present, it occurs within the tip length not in the stalk region; and 8) lastly, the tip and oval stalk section make up about 50% of the

seta height in *Chamaelinorops* (e.g., 4 μm in MCZ 156919) and no more than 10% of the height of lamellar setae in anoles.

d) The distinction between *Chamaelinorops* setae and anole setae (as defined by *A. sheplani*, *A. occultus*, and *A. cuvieri*) is not significantly altered if *Chamaelinorops* is compared to *A. auratus* and *A. tropidonotus*. *A. chrysolepis* has a somewhat wider seta tip and a larger stalk diameter than other anoles examined, but it does not resemble *Chamaelinorops* (e.g., the larger tip is distinctly triangular; Peterson and Williams, 1981).

The only *Anolis* species examined to date which has setae that remotely resemble those in *Chamaelinorops* is an undescribed South American form in the alpha section of the genus (*A. sp. n.* near *eulaemus* LACMNH 72741). The animal is reported to be riparian and semi-aquatic (Williams, personal communication). The gross morphology of the pad, the fine structure of the setae stalks, and the position of the tip inflection are similar to those in other anole species (Table 2). However, the setae tips are rectangular or oval in outline and have areas almost as large as those in *Chamaelinorops* (Table 2; Fig. 10C, D illustrate the rectangular tip). In both *A. sp. n.* near *eulaemus* and *Chamaelinorops* the specialized size and shape of the setae are correlated with increased tip length (e.g., $0.88 \pm 0.04 \mu\text{m}$ in *A. sp. n.* near *eulaemus*), and with the morphology of the tip/stalk junction. In both forms the distalmost stalk section (apex) is much wider than in the anoles ($0.60 \pm 0.03 \mu\text{m}$ wide in *A. sp. n.* near *eulaemus* and $0.6\text{--}0.7 \mu\text{m}$ wide in *Chamaelinorops* compared to $0.2\text{--}4 \mu\text{m}$ in most anoles). Subsequent flattening of the apex to form the tip produces a thick tip ($0.1\text{--}0.2 \mu\text{m}$) which is only slightly wider ($0.75 \mu\text{m}$) than the distal stalk ($0.6 \mu\text{m}$). Compression of a wider, less tapered stalk appears to give the tip its rectangular shape and greater average width. Setae on the pad in *A. sp. n.* near

eulaemus exhibit a gradual flattening to form the tip as in other anoles, but the seta (or seta-prong) tips on the phalanx iv scales are formed from the stalk more abruptly and exhibit the rectangular shape of *Chamaelinorops* setae (Fig. 10C, D). These setae or setae-prongs differ from *Chamaelinorops* setae in tip length ($0.72 \pm 0.06 \mu\text{m}$ compared to $1.1\text{--}1.5 \mu\text{m}$ in *Chamaelinorops*), tip area ($0.488 \pm 0.055 \text{ sq } \mu\text{m}$ compared to $0.6\text{--}1.4 \text{ sq } \mu\text{m}$ in *Chamaelinorops*), the position of the tip flexion, stalk dimensions, and stalk spacing (the latter features are not different from those of the lamellar setae).

e) The anole seta and seta-prong morphotypes generally differ in some of the same features as *Chamaelinorops* setae and anole setae, and it is possible that the *Chamaelinorops* setae should be compared with anole setae-prongs. The major points of similarity include: 1) the seta-prong tip surface usually appears to be thicker and formed by a more abrupt flattening of the stalk, and 2) the apex of the stalk is wider than that of the seta. These differences appear to be responsible for the rectangular tip shape. Further parallels include reduced stalk height and a less well-defined point of flexion of the tip on the stalk (a tip flexion can also be absent; Fig. 8D). However, the tip dimensions and stalk spacing of setae-prongs in any of the anole species examined are clearly different from those of *Chamaelinorops* "setae." For example, in *A. occultus* the poorly differentiated setae (or setae-prongs, the distinction is not clear in this area) at the metatarsal-phalangeal joint (Fig. 8D) are $4.4 \pm 0.3 \mu\text{m}$ tall and $0.44 \pm 0.03 \mu\text{m}$ in diameter with $1.2 \text{ stalks/sq } \mu\text{m}$, and the tips have an area of $0.285 \pm 0.025 \text{ sq } \mu\text{m}$. The tip dimensions are $0.63 \pm 0.06 \mu\text{m}$ long by $0.37 \pm 0.03 \mu\text{m}$ wide proximally and $0.45 \pm 0.02 \mu\text{m}$ wide distally. In *A. cuvieri* the seta-prong tips in the phalanx iv region (Fig. 4E) have somewhat larger dimensions than those in *A. occultus*, but the proportions are the same ($0.77 \pm 0.04 \mu\text{m}$ long by $0.60 \pm 0.03 \mu\text{m}$

wide distally and $0.48 \pm 0.03 \mu\text{m}$ wide proximally). In *Chamaeleolis* the seta-prongs shown in Figure 4F on the lateral digital scales adjacent to the pad are almost perfectly rectangular ($0.60 \pm 0.06 \mu\text{m}$ long by $0.44 \pm 0.03 \mu\text{m}$ wide distally and $0.40 \pm 0.03 \mu\text{m}$ wide proximally.)

Chamaelinorops appears to have a unique setal morphology. Comparisons beyond those with *A. occultus*, *A. sheplani*, and *A. cuvieri* reveal no parallels for the reduced stalk spacing in *Chamaelinorops* nor for the combination of shape and size features. Setae prongs and poorly differentiated setae in a number of anolines approach the tip shape of *Chamaelinorops* setae, but most of the tip dimensions are significantly smaller than those of *Chamaelinorops*. The setae of *A. sp. n.* near *eulaemus* approach the tip dimensions and some of the shape features of *Chamaelinorops* tips, but the resemblance is far from compelling. The setae of *A. sp. n.* near *eulaemus* have more in common with other anole setae than with *Chamaelinorops* setae.

f) There is significant individual variation in tip area among the five *Chamaelinorops* specimens (Table 2; $F_{\text{tip area}} = 82.1$; $p < 0.005$). The variation among the specimens is much greater than the level of apparent "wear" variation in *A. occultus* (Table 2).

The specimen with the greatest tip area (MCZ 156919) does have what appear to be "unworn" setae, but the tips are also about twice as wide as the tips in three of the other specimens ($0.97 \pm 0.05 \mu\text{m}$ in MCZ 156919 compared to $0.58 \pm 0.03 \mu\text{m}$ MCZ 126708, $0.55 \pm 0.03 \mu\text{m}$ MCZ 74594, and $0.52 \pm 0.03 \mu\text{m}$ MCZ 68691). The fourth specimen (MCZ 155851) exhibits tips of an intermediate width ($0.71 \pm 0.04 \mu\text{m}$). The wider tip in MCZ 155851 and MCZ 156919 occurs all over the lamellar surface; there is no suggestion that it is a regional or local variant. Similarly, there is no indication that the narrower tips in the other three specimens are "worn" versions of the larger ones. There are very few scars,

gouges or broken surfaces along the lateral edge of the tip in any specimen. MCZ 126708 exhibits a few of the wide tips scattered among the narrow ones. The width of the seta tip appears to be subject to significant individual variation ($F = 149.8$; $p < 0.005$).

Variation among the *Chamaelinorops* specimens in tip length is also significant ($F = 10.5$; $p < 0.005$), but the range is comparable to that of probable wear variation in *A. occultus* (Table 2). *Chamaelinorops* tip length varies from $1.09 \pm 0.04 \mu\text{m}$ in MCZ 74594 to $1.52 \pm 0.03 \mu\text{m}$ in MCZ 156919 with intermediate values of $1.44 \pm 0.13 \mu\text{m}$ in MCZ 156851, $1.31 \pm 0.10 \mu\text{m}$ in MCZ 126708 and $1.25 \pm 0.08 \mu\text{m}$ in MCZ 68691. The specimen with the shortest setae tips (MCZ 74594) exhibits considerable debris and variation in the outline of the distal edge, probably due to chipping and breakage. Some tips in all the areas are obviously broken off the stalks (these were not measured, but they suggest that the Oberhautchen is old or highly worn). MCZ 68691 with the second shortest tips also has more tips with ragged distal edges than the remaining three specimens. There is no clear evidence of a difference in wear among the remaining specimens, and variation in tip length among them is marginally significant ($F = 4.0$; $0.05 > p > 0.025$).

The Distribution of Setae

In *A. sheplani* setae cover all the lamellar scales and extend as far proximally as the metatarsal-phalangeal joint. In *A. occultus* setae cover the lamellae and some of the phalanx iv scales. The phalanx iv scales exhibit sequential steps in the morphotypic series (Fig. 2). Setae occur on the distal scales (numbers 22 through 26). The seta tip becomes progressively narrower on each more proximal scale, and a progressively smaller central area of the scale bears setae. The surrounding areas are covered by seta-prong and spike morphotypes. Setae are

absent on scales 27 through 30, and setae-prongs occur over the central region of the scales (Fig. 8D). Prongs and spikes occur more laterally on these scales.

In *Chamaelinorops* setae occur on all the subdigital scales. The setae which cover the generalized subdigital scales throughout the phalanx iii and iv regions are somewhat shorter, but otherwise not different from lamellar setae. For example, the setae on the most proximal subdigital scale (number 25) in MCZ 155851 are $5.6 \pm 0.3 \mu\text{m}$ tall and the stalks are $0.58 \pm 0.06 \mu\text{m}$ in diameter (cf. Table 2; Fig. 6C, E, F). The tips are flexed within the flattened tip region so that the entire tip area can not be measured accurately, but the tips are about $1.5 \mu\text{m}$ long and $0.7 \mu\text{m}$ wide.

Shorter setae similar to those on the generalized subdigital scales also occur over the central portion of the distal phalanx i scales (e.g., on scale 4 in MCZ 155851 the stalks are $5.6 \pm 0.7 \mu\text{m}$ tall). Laterally the setae grade into other members of the morphotypic series (see phalanx i description following).

Although setal distribution in *Chamaelinorops* differs from that of *A. sheplani* or *A. occultus*, *A. auratus* exhibits the *Chamaelinorops* pattern. Setae occur on all the subdigital scales, and those on the phalanx i scales occur only in the central portion of the scale (Peterson and Williams, 1981).

The Phalanx i Region

In *A. occultus* and *A. sheplani* the five small phalanx i scales are covered with spines and spinules. Distally the surface is covered with spines which have sharp, curved tips (Fig. 8E). The proximal half of the scale is spinulate (Fig. 8F). Scale 5 is entirely covered with spinules. There are two populations of spinules: short, nubbinlike spinules $0.19 \pm 0.02 \mu\text{m}$ tall and $0.16 \pm 0.03 \mu\text{m}$ in diameter, and taller, spinelike spinules $0.52 \pm 0.14 \mu\text{m}$ tall and $0.12 \pm 0.03 \mu\text{m}$ in diameter (actual dimensions for *A. sheplani* MCZ

125641). Together they occur in a density of 7.4 spinules/sq μm . *A. sheplani* exhibits its similar spinules in a density of 8.4 spinules/sq μm .

In *Chamaelinorops* the distal phalanx i scales (numbers 1–7 in Fig. 5A) exhibit a wide range of fine structure. The central region of the scale is covered with setae. Setae extend distally onto a very small free margin on several of the scales (e.g., scales 2 and 3 in Fig. 7A). The seta-bearing area corresponds to the functionally subdigital portion of the scale. The lateral portions of the scales face laterally, not ventrally, and the morphology of the surface is very similar to that of the adjacent lateral digital scales. In the transition from the subdigital to the lateral surfaces, the distal margin of the subdigital scale becomes studded with large diameter spines (Fig. 7B), and the seta tips become smaller. More laterally the scale surface becomes contoured into “hillocks”—the central portion of each epidermal cell is elevated while the intercellular junction is depressed (Peterson and Williams, 1981; Fig. 7C, D, F). The hillocks create a surface roughness of intermediate magnitude. The surface of each hillock is covered with spines which are fairly similar in size, but adjacent hillocks have spines of very different dimensions (Fig. 7D and text following). True keels occur along the lateral portion of several of the subdigital scales (e.g., scales 2, 5, 6, and 7 of MCZ 156919) and on the lateral digital scales (Fig. 7C).

Chamaelinorops exhibits extraordinary diversity in the morphology of spines on the phalanx i subdigital scales and the lateral digital scales. The diversity in spines is based on variation in diameter and density which, in the extreme, results in the fusion of spines from adjacent epidermal cells. Figure 7 illustrates some of this diversity. Small spines with an understory of spinules occur toward the sides of the scales and in the deep folds between the scales (Fig. 7F lower left and center). The small spines

are $0.88 \pm 0.09 \mu\text{m}$ tall and $0.21 \pm 0.08 \mu\text{m}$ in diameter near their base; the spinules are $0.30 \pm 0.09 \mu\text{m}$ tall and $0.17 \pm 0.05 \mu\text{m}$ in diameter. Together they occur in a density of 3–4 spines and spinules/sq μm . The medium-sized spines shown in Figure 7D bottom are similar to those in *Anolis* ($1.51 \pm 0.31 \mu\text{m}$ tall and $0.48 \pm 0.03 \mu\text{m}$ in diameter with about 1.0–1.5 spines/sq μm). Large pyramidal spines are the most common (Fig. 7D top left and Fig. 7F center; those in Fig. 7F are $2.12 \pm 0.20 \mu\text{m}$ tall and $1.66 \pm 0.20 \mu\text{m}$ in diameter and occur in a density of about 0.2 spines/sq μm). Even larger, irregular spines (upper right Fig. 7B, C and upper left of Fig. 7F) appear to represent a fusion of several single pyramidal spines. Some fused or conglomerate spines assume incredible proportions and cover more than 100 sq μm of the scale surface. Partially fused large spines form serrate ridges stretching for more than 50 μm parallel to the keels on the lateral scales. The spacing of the conglomerate spines is highly irregular and extensive aspinate, bare areas occur around an individual spine (density less than 0.01 conglomerate spines/sq μm). The very large conglomerate spines occur on the distolateral corners of the subdigital scales in the phalanx i region and on the lateral scale surfaces distally and adjacent to the keels (Fig. 7C).

A. sheplani and *A. occultus* exhibit no counterpart for the single and conglomerate spines in *Chamaelinorops*. Spines in the anoles have a maximum diameter of about $0.5 \mu\text{m}$, one-third or less of the maximum spine diameter in *Chamaelinorops*, and the anole spines occur in a density of about 1/sq μm . The *Chamaelinorops* spine series begins and ends at lower densities (3–4/sq μm to 0.01/sq μm) than the spinule-to-spine series in the anoles (e.g., 7.4/sq μm to 1/sq μm in *A. sheplani*). No other anoline examined to date has fused, conglomerate spines. But the mosaic landscape formed by different spine types and the variation in single spine diameter and density which

is exhibited by *Chamaelinorops* does have some parallel in a number of *Anolis* species, including *A. cuvieri* and *A. auratus*. These parallel morphologies occur only on the lateral digital scales. In *A. auratus* there are scattered single epidermal cells covered with large pyramidal spines (about $4 \mu\text{m}$ tall; $2.4 \mu\text{m}$ in diameter near the base and 0.1 spine/sq μm ; cf. $1.9 \mu\text{m}$ tall, $1.8 \mu\text{m}$ in diameter and 0.2 spines/sq μm in *Chamaelinorops*). The lateral digital scales in *A. cuvieri* exhibit a mosaic arrangement of spine types (Fig. 4C, D). The extremes in diameter and spacing are: spines $1.53 \pm 0.15 \mu\text{m}$ tall, $0.75 \pm 0.04 \mu\text{m}$ in diameter in a density of 0.23 spines/sq μm and spines $0.72 \pm 0.04 \mu\text{m}$ tall, $0.32 \pm 0.02 \mu\text{m}$ in diameter in a density of 3.6–5.0 spines/sq μm . The variation in diameter is more conservative than that in *Chamaelinorops*, and there is no contouring of the surface by hillocks and keels.

The Morphotypic Series in Fine Structure

The varieties of fine structure in *A. occultus* and *A. sheplani* are adequately described by Figure 2. Stalk diameter may be somewhat smaller in *A. sheplani*, and the stalk height of spines, prongs, setae-prongs and setae is toward the low end of the ranges given in Figure 2 for both species. All the morphotypes shown in Figure 2, except small spines and spinules which occur on the phalanx i scales, occur in the proximal area of a lamella, and there are no novel morphotypes.

Chamaelinorops does not exhibit the anole spine-seta morphotypic series. There is, however, a parallel, alternative series. Along a transect from distal to proximal on a lamella the *Chamaelinorops* setae become shorter and the tip becomes less compressed leaving a pronglike form. The prong is generally succeeded by a tall spine ($2.5\text{--}3 \mu\text{m}$ tall) with a tapered, hook tip (Fig. 7E right). The series is similar to the spine-seta

series in *Anolis* in that diameter and density remain constant as tip shape and stalk height change. The two series differ in stalk density (about $1/\text{sq } \mu\text{m}$ in *Anolis* and $0.5/\text{sq } \mu\text{m}$ in *Chamaelinorops*). The absence of an *Anolis*-like seta-prong and spike may be only a formalism. The *Chamaelinorops* seta itself is similar to an anole seta-prong, and variation in stalk height is much reduced in *Chamaelinorops* so that the transition to a hooklike tip simply occurs below the $5 \mu\text{m}$ limit of the spike. The *Chamaelinorops* spine-seta series exhibits continuous variation in stalk height and tip shape. It is a simpler series than that of *Anolis*, but there are no "gaps" or "missing" morphotypes. Even more proximally on the lamella, density increases and a small band or scattered patches of anole-like spines can be distinguished ($0.4\text{--}0.5 \mu\text{m}$ diameter and 0.9 spines/ $\text{sq } \mu\text{m}$; Fig. 7E lower left). These grade into regions of small spines about $1 \mu\text{m}$ tall) with an understory of very small, nubbinlike spinules (together in a density of $2.4/\text{sq } \mu\text{m}$). The fields of small spines and spinules are linked to the *Anolis*-type spine by continuous variation in diameter and density. Fields with larger "small spines"—those approaching $1 \mu\text{m}$ height and $0.5 \mu\text{m}$ diameter—have a very reduced understory of small spinules. The 1 spine/ $\text{sq } \mu\text{m}$ density and the *Anolis*-like spine appear to be achieved as the last spinules disappear from the understory (e.g., the area cited above exhibits 1.2 small spines/ $\text{sq } \mu\text{m}$; the spinules effectively double the density in this region). The fields of small spines and spinules in *Chamaelinorops* occur in slightly lower densities than those of most anole species (e.g., the maximum density in *Chamaelinorops* is $4/\text{sq } \mu\text{m}$, while that in *A. sheplani* is $7.4/\text{sq } \mu\text{m}$), but the shape and size of the spines and spinules are similar. The morphotypic series formed between the spinule and *Anolis*-like spine is very similar to that present in the *Anolis* species.

A third, larger spine variety occurs in

patches along the proximodistal borders of the lamella (Fig. 7E left). The dimensions ($2.62 \pm 0.42 \mu\text{m}$ tall and $1.46 \pm 0.09 \mu\text{m}$ diameter) are similar to those of the large single spines on the phalanx i and lateral digital scales, but the lamellar spines are much more closely packed ($0.5\text{--}0.6$ spines/ $\text{sq } \mu\text{m}$) with almost no bare surface between the spines. These large spines resemble the *Chamaelinorops* spine that forms the base of the spine-seta series in density and in height, but they differ in diameter.

The three different spine types—a) the *Anolis*-like spine which is continuous with the spinulate series; b) the large diameter spine which may be linked to the large spines in the phalanx i region and on the lateral digital scales; and c) the $0.5 \mu\text{m}$ diameter- $0.5/\text{sq } \mu\text{m}$ spine which forms the base of the *Chamaelinorops* spine-seta morphotypic series—are topographically contiguous (Fig. 7E). There are no epidermal cells bearing spines with intermediate diameters or densities.

COMPARISONS BETWEEN *PHENACOSAURUS HETERODERMUS* AND *ANOLIS VALENCIENNI*

Pad and Scale Morphology

The major differences between the two forms are in the position of the pad and the shape of the lamellae.

In *A. valencienni* lamellae occur under phalanges ii and iii. The phalanx iv scales have a lamellar shape in some cases, but a free margin is absent. The 26 lamellae (58–60% of the total scales) are packed into 3.4 to $3.8 \mu\text{m}$ of the digit (49–45% of the digital length; Table 1).

In *Phenacosaurus* phalanges ii and iii are shorter relative to the length of the body or digit, and lamellae in this region account for only about 42% of the subdigital scales (Table 1). The distribution of lamellae is expanded, however, to include the entire phalanx iv region (scales 27–34 in Fig. 9A) and the proximal pha-

lanx i region (scales 6–12 in Fig. 9A; Lazell, 1969, describes phalanx i lamellae in all *Phenacosaurus* species). Figure 9C shows the “free” lamellar margin of the scales just distal to the metatarsal-phalangeal joint. This expansion of lamellar scales more than doubles the length of the pad and doubles the number of lamellae (better than 80% of the subdigital scales are lamellae; Table 1). Although the pad is roughly twice as long as that in *A. valencienni*, the number of lamellar scales and pad area are not markedly greater in *Phenacosaurus*. There are 25 actual lamellae in *A. valencienni* and 24 to 31 in *Phenacosaurus* (the first pad scale is non-lamellar). Estimated pad area is 4.6 sq mm in *A. valencienni* (P 329; 58.5 mm SV length) and 4.3 sq mm in *Phenacosaurus* (P 300; 56 mm SV length). The similarity in these values in spite of the differences in pad length and lamellar distribution is related to the difference in lamellar shape. In *A. valencienni* the phalanx ii lamellae are very wide and short (mean width/length = 11.3). Approximately two-thirds of the pad area (2.9 sq mm) and lamellar scales (17 lamellae) are concentrated under the second phalanx. In *Phenacosaurus* there is less variation in lamellar proportions, and the lamellae are narrower and longer than those of *A. valencienni* (the mean width/length ratio for all lamellae is 5.5 in *Phenacosaurus* and 9.0 in *A. valencienni*; the standard deviations are 2.0 and 4.7, respectively). As a consequence of the difference in lamellar shape and distribution, pad area in *Phenacosaurus* is more evenly distributed among the phalanges (approximately 15% under i, 30% under ii, 20% under iii, and 35% under iv).

Expansion of lamellae into the phalanx iv region is unusual among West Indian anoles, but does occur in *Anolis*. *A. aequatorialis*, a South American alpha anole with a narrow pad, has lamellae extending through the proximal half or two-thirds of the phalanx iv region. This is the most proximal distribution of

lamellae among the anole species examined to date, and *Phenacosaurus* may be unique in the presence of lamellae throughout the phalanx iv region.

The expansion of lamellae into the phalanx i region is unique among anole species with a raised pad, but unlike the proximal expansion of the pad, the phalanx i lamellae do not increase the area of the pad. In *A. valencienni*, as in many anoles with a well developed, wide pad, the distal end of the pad is a flap or shelf which is attached to phalanx ii and the i/ii interphalangeal joint, but projects far distally to lie under phalanx i. In *A. valencienni* the distal 7 to 9 lamellae and an estimated 0.65 sq mm of pad area (14% of the total area) lie on this shelf (measurements for P 329). In figure 9B the i/ii interphalangeal joint is flexed, and the portion of the pad which lies under phalanx i, scales 9 to 16, is tilted toward the viewer. The phalanx i pad area in *Phenacosaurus* (0.64 sq mm; 15% of the total area; measurements for P 300) is no greater than that of the pad “shelf” in *A. valencienni*. The pad is simply attached more distally in *Phenacosaurus*.

Fine Structure

There are no major differences in the fine structures of *Phenacosaurus* and *A. valencienni*.

The shape and dimensions of the setae are not different (Table 2; Fig. 10A, B). In *A. valencienni*, setae occur on the lamellae under the ii and iii phalanges and cover the generalized subdigital scales in the phalanx iv region. In *Phenacosaurus* setae occur on all the lamellae (throughout the ii, iii, iv and proximal i phalangeal regions).

The phalanx i scales are covered with spinules and small spines in *A. valencienni*. In *Phenacosaurus* the distal quarter of the scale surface is covered by prongs and spikes which grade into spines and finally, small spines ($1.27 \pm 0.12 \mu\text{m}$ tall, $0.28 \pm 0.03 \mu\text{m}$ diameter) and spinules ($0.33 \pm 0.10 \mu\text{m}$ tall, $.21 \pm$

Figure 9. The subdigital surface of the fourth digit in A) *Phenacosaurus* (P 300) and B) *A. valencienni* (P 369). In *Phenacosaurus* scales 7 through 12 are lamellae attached to phalanx i. Lamellae 13–26 occur under the second and third phalanges, and lamellae 27–34 occur under the fourth phalanx. The free margin of the most proximal subdigital scale (number 34) is shown in C). In *A. valencienni* scales 17–34 are lamellae attached to phalanges ii and iii. Scales 9 (or 10) through 16 are lamellae which are attached to the second phalanx or the i/ii interphalangeal joint region, but lie under phalanx i. The i/ii interphalangeal joint (between scales 16 and 17) is flexed toward the viewer. The ii/iii interphalangeal joint (under scales 27 and 26) is hyperextended.

Figure 10. The subdigital fine structures of *Phenacosaurus* and *A. sp. n. near eulaemus*. A. Lateral view of the setae on the lamella free margin in *Phenacosaurus* (P 300). B. Setae tips of *Phenacosaurus* (MCZ 139566). C. The setae-prongs of *A. sp. n. near eulaemus* (LACMNH 42144) on phalanx iv scale 41. D. Detail of the seta-prong tips in *A. sp. n. near eulaemus*.

0.05 μm diameter; density of small spines and spinules = 2.9/sq μm) over the proximal three-quarters of the scale surface. The prongs, spikes and spines are similar

to those which occur proximally on lamellae in *A. valencienni* and *A. cuvieri*. The spine-seta morphotypic series in *Phenacosaurus* is in no way different

from that of *A. valencienni*, *A. cuvieri*, or most of the generalized anole species examined to date.

DISCUSSION

SUMMARY AND PRELIMINARY EVALUATION OF THE COMPARISONS

Chamaeleolis and *A. cuvieri*

Chamaeleolis and *A. cuvieri* share all the major features of the pad complex. The morphology of the setae and that of other fine structural features of the scales are very similar in the two forms, and the gross morphology of the pad is virtually identical.

The features that distinguish *Chamaeleolis* from *A. cuvieri* are: a) the phalanx i scales have spinules and small spines in *Chamaeleolis* and typical subdigital spines in *A. cuvieri*; b) setae occur on all the phalanx iv scales in *Chamaeleolis*; c) the seta stalk is taller in *C. chamaeleontides*, but not in *C. porcus*; d) a distinct subdigital scale row is not well established in the phalanx iv region; and e) the toe and pad phalanges are absolutely shorter in *Chamaeleolis*.

The difference in stalk height between *C. chamaeleontides* and *A. cuvieri* may be functionally significant, but a number of anole species (e.g., *A. stratulus* and *A. evermanni* among the Puerto Rican ecomorphs; Peterson and Williams, unpublished data) exhibit seta stalk proportions similar to those of *C. chamaeleontides*. The difference does not distinguish *Chamaeleolis* from *Anolis*, and it probably is not indicative of a phylogenetically distinct seta type.

Similarly, the presence of setae on all the phalanx iv scales is a common difference among anole species. Even ecomorph pairs may differ in this feature (e.g., *A. occultus* and *A. sheplani*, and *A. pulchellus* and *A. auratus*; Peterson and Williams, 1981 and unpublished data). The expanded distribution of setae in

Chamaeleolis may be correlated with the short digit and pad phalanges. As one might expect, the number of setae is highly correlated with body size (essentially with the load presented to the setae; Peterson *et al.*, 1982). One would further expect *Chamaeleolis* and *A. cuvieri* to have comparable numbers of setae. Given that seta density and pad width are similar in the two forms, the shorter pad in *Chamaeleolis* would result in fewer setae than in *A. cuvieri*, but for the expansion of setae over the proximal phalanx iv scales. The seta-bearing area is not precisely equal in the two forms (cf. body size in Table 1), but both are within 10% of the area predicted for their body size by a scaling relationship developed for the Puerto Rican ecomorphs (Peterson *et al.*, 1982).

The presence of extensive spinulate areas on the subdigital surface and the absence of a distinct subdigital scale row in the phalanx iv region do not distinguish *Chamaeleolis* from *Anolis*, but these features may be truly primitive characters. Multiple small scales on the proximal subdigital surface occur in many of the preanoline iguanids (e.g., *Diplolaemus bibroni* and *Pristodactylus achalensis*; Peterson, unpublished data) and in several *Anolis* species that are considered relatively primitive by other measures (e.g., *A. pentaprion* and some specimens of *A. sheplani* and *A. occultus*; Schwartz, 1974; Williams, personal communication). This condition has not been observed in more derived anole species, such as *A. cristatellus* or *A. tropidonotus*. Similarly, small, high density spines or spinules occur on the subdigital scales of preanoline iguanids such as *Diplolaemus bibroni* and *Pristodactylus achalensis*, and on the phalanx i scales of some primitive anole species, such as *A. occultus* and *A. sheplani*. More derived anole species, such as *A. cristatellus* and *A. tropidonotus*, exhibit spines, spikes or some combination of morphotypes within the spine-seta portion of the morphotypic series. The spinule morphology

in *Chamaeleolis* is not significantly different from that in *A. sheplani* or *A. occultus*. There is no basis for suggesting it represents an alternative or more primitive condition than those of the relatively primitive anole species.

Chamaeleolis can not be distinguished from *Anolis* in the structure of the pad complex. Setae, lamellae, and other basic features of the pad appear to be homologous in the two genera.

Chamaelinorops and *A. sheplani*, *A. occultus*, *A. chrysolepis*, *A. auratus*, and *A. sp. n. near eulaemus*

Chamaelinorops differs from *A. sheplani* and *A. occultus* in every aspect of the gross and fine structure of the subdigital pad. In spite of their similarity in body size and habitus, *Chamaelinorops* shares little beyond the presence of a pad with these two anole species. The *Chamaelinorops* pad is very small and different in shape and position on the digit. The *Chamaelinorops* setae are distinct in shape, dimensions and distribution.

In its gross morphology, *Chamaelinorops* resembles a variety of the more terrestrial, derived anole species, particularly *A. auratus*. Common features include: the Norops condition, a narrow pad, the absence of lamellae under the third phalanx, expansion of lamellae into the phalanx i region, and the relatively conservative width/length proportions of the lamellae. *Chamaelinorops* differs from *A. auratus* in the extent to which these features are developed. In almost every case the trend is more extreme in *Chamaelinorops*—pad area is lower relative to body size, the lamellar proportions are closer to those of generalized subdigital scales, the lamellar free margin is severely reduced, and lamellae have expanded further into the phalanx i region. The *Chamaelinorops* pad is even less well developed than that of *A. annectens*, a highly derived anole related to the padless *A. onca* (Peterson and Williams, 1981).

In fine structure, *Chamaelinorops* differs from all other anolines examined to date in the reduced density and large size of the seta tip. An undescribed South American species, *A. sp. n. near eulaemus*, has setae whose tips are somewhat larger and approach the rectangular shape of *Chamaelinorops* setae. *A. auratus* exhibits seta distribution similar to that of *Chamaelinorops*, but even these highly derived anole species exhibit setae which are more similar to those of generalized anoles than to those of *Chamaelinorops*.

The distinction between the anole species and *Chamaelinorops* in fine structure is not confined to the seta. The entire *Chamaelinorops* spine-seta series is based on a lower stalk density (0.5/sq m) and reduced variation in stalk height. *Chamaelinorops* also exhibits a parallel series of spines which is based on varying diameter and stalk density. Ultimately, increased diameter and decreased density produce huge keel-like spines. Comparable spine morphology does not occur on the subdigital scales of anoles, although large, pyramidal, single spines and a more conservative spine series are found on the lateral digital scales of some anoles (e.g., *A. auratus*). Possibly the most primitive portion of the fine structural suite, the spinule-spine series, is comparable to that found in the *Anolis* species.

In fine structure, as in gross pad morphology, *Chamaelinorops* resembles derived anole species more than the relatively primitive West Indian forms, *A. occultus* and *A. sheplani*. But in fine structure, the similarity between *Chamaelinorops* and any anole is extremely weak, and it is not at all clear that the *Chamaelinorops* spine-seta series is derived from an *Anolis*-like series.

Phenacosaurus and *A. valencienni*

The fine structure and most aspects of the gross morphology of the pad are very similar in *Phenacosaurus* and *A. valencienni*. The only differences are in the

gross structure of the pad. a) The *Phenacosaurus* lamellae are less variable in shape and have more conservative proportions (the scale width/length ratios are 5.5 compared to 9.0 in *A. valencienni*). This feature does not, however, distinguish *Phenacosaurus* from the dwarf twig ecomorphs, *A. occultus* and *A. sheplani* (lamellar width/length ratio is 5.0). b) The lamellae also have an expanded distribution in *Phenacosaurus*. They extended throughout the phalanx iv region and over the proximal phalanx i region. Both phalanx i and phalanx iv lamellae occur in *Anolis*; the condition in *Phenacosaurus* is an extreme, but not a novel pad position.

The case for an independent origin of the pad complex in *Phenacosaurus* rests solely on the difference in pad position. Among gecko lineages differences in pad position do indicate alternative pathways in pad evolution (Russell, 1975, 1976, 1979), but these cases involve topographic alternatives (e.g., a terminal versus a mid-digit pad) and often major differences in the internal structure of the pad, not simply relative degrees of expansion (see Discussion). Evidence from gross pad morphology alone is insufficient to rule out an independent pad origin, but the fine structural comparisons would argue against it, and other interpretations of the difference in pad position seem at least as plausible. *A. valencienni* and *Phenacosaurus* have similar structural habitats, but extreme pad expansion in *Phenacosaurus* might be correlated with subtle differences in the structural habitat and in foot placement. The pad area in *Phenacosaurus* is distributed so that if the interphalangeal joint mobility is comparable to that in *A. valencienni*, the pad can be made to conform to a smaller radius of perch curvature. Alternatively the difference might be correlated with the adaptive history of the two lineages. The two twig giants might arrive at the same body size and pad area from different ancestral sizes and pad shapes. *Phenacosaurus* could be a "scaled up" twig dwarf with negative

allometry of the ii and iii phalanges forcing expansion of the pad. There is sufficient variation in pad position among anoles to suggest that the *Phenacosaurus* morphology represents an extreme within the context of an *Anolis*-like pad complex rather than an alternative, non-homologous pad.

THE EVOLUTION OF THE SUBDIGITAL PAD COMPLEX AMONG ANOLINES: A PRELIMINARY VIEW FROM INSIDE THE RADIATION

Anolis itself poses the greatest challenge in comprehending the evolutionary diversity of the pad complex. To date only about 15% of the 200 to 250 species have been examined, but this sample includes a variety of West Indian, Central and South American species groups, an arboreal ecomorph series, and terrestrial and semi-aquatic species. The sample is sufficiently large and diverse to make possible a very tentative assessment of some aspects of pad evolution within the genus.

Anolis species exhibit two basic similarities in the pad complex—lamellar scales forming the pad and the spine-seta morphotypic series. Lamellae are wide, flexible, flaplike scales. Anole lamellae vary in shape, size and position on the phalanges. The varieties of anole lamellae probably do not represent independent acquisitions of the anole pad, but at least the external morphology of the pad scales does not provide enough information to examine the issue carefully. Scales that are much wider than long and have a free margin, occur in some geckos (Ruibal and Ernst, 1965) and in the arboreal skink, *Prasinohaema virens* (Williams and Peterson, 1982). These lamellae are certainly derived independently of those in *Anolis* and yet the external morphology provides no clear indication of this. The internal connective tissue and musculoskeletal framework of the digit provide much more information and probably would be a better tool for investigating the macroevolu-

tion of the gross pad (see Russell, 1975, 1976, and 1979 for an analysis of these features in the Gekkonidae). While lamellar scales are a major feature of the pad complex in *Anolis* and appear to be homologous among anoles, their external morphology is not a very satisfactory basis for studying macroevolution among anoles or anolines.

The spine-seta morphotypic series is characterized by a $0.5\ \mu\text{m}$ stalk diameter, variable stalk height, 1 stalk/sq μm spacing, and two basic tip shapes—a hook and a spatula (a flattened triangle joined to the stalk at the apex; Fig. 2). The two tip shapes are linked by intermediate shapes. This series occurs in all the *Anolis* species examined to date. The extent to which it discriminates anoles from other iguanids is under investigation.

There is relatively little novelty in the morphotypes within *Anolis*. None of the West Indian species examined thus far (and these include such unusual forms as *A. eugenegrahami*) exhibit significant differences in the morphotypes in spite of very significant differences in body size, habitat, species group, and gross pad morphology. Instead, adaptation in fine structure appears to be correlated with differences in the absolute numbers and distribution of the morphotypes. Extreme specialization in the complex is, in some cases, associated with a "retrograde" shift in the series and the absence of terminal morphotypes, such as the seta. But it is still possible to identify the series by stalk density, stalk diameter and usually by the remaining tip shapes. For example, *A. oxylophus* and *A. macrolepis*, two Central and South American semi-aquatic species, have a well-developed pad, but setae are replaced by typical spike and prong morphotypes (Peterson and Williams, unpublished data). *A. onca* has neither pad nor setae, but the subdigital spines have the shape and density of anole spines (Peterson and Williams, 1981). There is only one case in which setae are replaced by a novel mor-

photype. In *A. barkeri* the lamellae are covered by a branched prong, but the diameter of the stalk, the density of stalks and the presence of typically anole spikes, prongs, and spines clearly label the branched prong as a morphotype derived from the anole series. There are only three to four species in which the dimensions of any of the morphotypes are unusual or novel. For example, *A. chrysolepis* exhibits larger setae tips and a larger stalk diameter, but the seta density and tip shape clearly place the morphology within the morphotypic series. *A. sp. n.* near *eulaemus* discussed previously is a second example.

The morphotypic series also discriminates the *Anolis* pad from that of skinks and perhaps from that of geckos. The arboreal skink, *Prasinohaema virens*, has setae which are remarkably similar to those of *Anolis* in shape, but the stalk diameter and tip dimensions are an order of magnitude greater, and the stalk density is two orders of magnitude lower (Williams and Peterson, 1982). The spine—seta-prong portion of the morphotypic series is also absent (Williams and Peterson, 1982). The setae of some coleopteran species (Stork, 1980) also have a shape that is remarkably similar to that of *Anolis* setae, but the dimensions appear to be quite different. Gecko pads are too diverse and their epidermal fine structure too poorly known to be certain that the *Anolis* morphotypic series is truly unique. However, the setae of a variety of geckos appear to represent an entirely different design for adhesion. The setae exhibit large diameter branched stalks in low density and many triangular terminal spatulae (Ruibal and Ernst, 1965; Hiller, 1968), but it remains possible that the anole morphotypic series is to some extent paralleled in geckos. Russell (1976) describes a morphotypic series of *Cyrtodactylus* species which involves increasing spine height (with suggested variation in diameter) that leads through an *Anolis*-like seta tip to the branched seta with multi-

ple spatulae. Stalk density and the dimensions of the morphotypes, apart from stalk height are not included and forms which are intermediate between the spine and the anole-like seta are not described. These features might discriminate the *Anolis* and *Cyrtodactylus* series.

The morphotypic series in *Anolis* is a heuristic comparative device based on the existing subdigital morphology in *Anolis*. The morphotypes (e.g., seta or spike) summarize the variety in subdigital fine structure, and the order or arrangement of morphotypes reflects the pattern of topographic variation (e.g., fields of spikes grade into spines or prongs, and are rarely adjacent to spinules). But the arrangement of the morphotypes is also consistent with the patterns of evolutionary change observed in the *A. annectens-onca* lineage (Peterson and Williams, 1981) and in the "retrograde" shift and substitution of spikes or prongs for setae in Central and South American semiaquatic anoles (Peterson and Williams, unpublished data). The morphotypic series may prove to describe a number of phylogenetic sequences, but it might become even more useful in integrating developmental mechanisms and functional change with phylogenetic sequences. In some instances the differences between adjacent morphotypes moving from spine to seta correspond to sequentially later stages in the developmental sequence (e.g., density is determined before stalk height; suggested by Maderson [1970] and Peterson, unpublished data), and the spine-seta series which occurs over the proximal portion of the lamella may be a relic representation of setal development (Peterson and Williams, 1981, see Fig. 23). Most of the differences between morphotypes and thus between species with different morphotypes in homologous areas can be interpreted in terms of the ontogenetic/phylogenetic framework of Alberch, *et al.* (1979). Developmental interpretations of the series are

still very speculative, although considerable information on the shedding cycle and late stages in seta development are available through the studies of Maderson (1970, for example).

Some aspects of the series can already be interpreted functionally. Three of the dimensions of the spine-seta series—stalk diameter, tip size, and stalk density—are relatively constant among *Anolis* species (Peterson, *et al.*, 1982; and unpublished data). The tip, stalk, and epidermal scale surface represent sequential links in the transmission of tensile force to the dermal core of the scale and ultimately to the musculoskeletal system of the digit. The consistency in these dimensions implies that stress is constant over a range of body sizes. Scaling for different body sizes occurs in the absolute number of setae and thus in pad area (Peterson, *et al.*, 1982). In contrast, variation in stalk height has no effect on tensile stress, and stalk height may vary independently of the other dimensions over the surface of a given lamella or between species.

The application and disengagement of setae, spines, etc. form a second set of functional constraints on the system. Height and diameter determine the stalk stiffness and the ability of the tip to "find" the substrate. The size and density of tips could also influence how the tips behave. Increased density, essentially crowding of the tips, could reduce the ability of individual tips to contact the substrate unless diameter and tip size are appropriately reduced. Conversely, extreme reduction in density would give the tips freedom to contact the substrate, but could result in too few tips to support the body mass, unless tip size and stalk diameter were appropriately increased (this would in effect reduce the distance between setae and re-introduce the problems associated with crowding).

The consistency among anoles in stalk density, stalk diameter and seta tip size appear to reflect functional constraints, but there is no reason to expect the

particular dimensions of the anole morphotypic series to be universal. Indeed, there appear to be a variety of setal dimensions and densities (e.g., anoles, *Prasinohaema virens*, geckos and even Coleoptera). The dimensions of *Anolis* morphotypic series may reflect only one of the solutions which might arise in the evolutionary sequence from small spines and spinules to the spine-seta series. Any differences in the character of the series, particularly in the "constants," among the anolines and non-anoline iguanids may indicate an independent acquisition of the pad fine structure.

The existing comparative data suggest that the spine-seta morphotypic series is homologous at least among *Anolis*, *Phenacosaurus* and *Chamaeleolis*. The morphology of the entire pad complex in *Phenacosaurus* is quite consistent with the view that it diverged from an *Anolis* stock in which setae and a gross pad were already well developed.

Chamaeleolis and *Chamaelinorops* are thought to have diverged from the anoline lineage much earlier, perhaps prior to the differentiation of *Anolis* (Fig. 1). It is not clear whether the *Chamaeleolis* lineage diverged earlier or later than *Chamaelinorops* (cf. Fig. 1A, B). The pad complex in *Chamaeleolis* appears to be homologous and virtually identical with that in relatively generalized *Anolis* species. The *Chamaelinorops* morphology is not easily interpreted.

The gross pad morphology of *Chamaelinorops* resembles that of highly derived, more terrestrial anoles, e.g., *A. auratus*. The resemblance in gross morphology involves a number of characters that at least within *Anolis* vary independently (e.g., the Norops condition is not always associated with the absence of phalanx iii lamellae; Table 1). The similarity between *A. auratus* and *Chamaelinorops* is not readily dismissed in favor of the dissimilarity in fine structure, but neither is it easy to accept the

verdict that the *Chamaelinorops* morphology is derived from an *Anolis*-like pad. The resemblance between *A. auratus* and *Chamaelinorops* is based on features which could characterize an early, nascent pad (e.g., the Norops condition and a very few lamellae that are not well differentiated from generalized subdigital scales). The fine structure of *A. auratus* clearly labels it a derived anole (Peterson and Williams, 1981), but the gross morphology might be convergent on the early stages in pad evolution. The *Chamaelinorops* pad morphology might be homologous with that of *Anolis* and represent a) a highly derived pad condition or b) a relic, very early stage in pad evolution, substantially before the well-defined pads of *Chamaeleolis* and *A. sheplani*. The gross morphology also permits a third interpretation because none of the gross characters—lamellar morphology, pad shape and position, or phalangeal proportions—are very incisive criteria for pad homology at this level. The *Chamaelinorops* pad might also represent an independent, parallel acquisition of the pad within the anolines.

The fine structure of the subdigital surface is unique among the anolines examined. It is unlikely that the morphology represents a relic, primitive condition in the sequence leading to *Chamaeleolis* and *Anolis*. a) The spine-seta morphotypic series appears to arise from spinules through a decrease in spinule density and an increase in stalk height and diameter. The trend would have to be reversed to shift from the 0.5 stalks/sq μm series spacing in *Chamaelinorops* to the 1 stalk/sq μm series spacing in *Anolis*. b) The fine structure in *Chamaeleolis* and primitive *Anolis* species exhibits regional differentiation. In these forms the phalanx i scales are covered by spinules or spines which are more conservative and presumably more primitive members of the morphotypic series. If the *Chamaelinorops* morphology represented an intermediate between the

spinate/spinulate scales of preanoline iguanids and those of *Anolis*, it seems unlikely that the phalanx i scales would be covered with setae. c) If the *Chamaelelinorops* morphology represented an intermediate in the sequence of anole pad evolution, one would expect to find at least some evidence of the large spine morphotypic series in more primitive *Anolis* species and in *Chamaeleolis*, not simply in derived *Anolis* species. The fine structure suggests that the *Chamaeleolis* pad is derived, but it does not unequivocally suggest whether it is derived independently or from an *Anolis*-like pad morphology.

There are three morphotypic series in *Chamaelinorops*: a) a "spinule to *Anolis*-like spine" series which is characterized by variation in stalk density, diameter and height; b) a large, pyramidal spine series that is characterized by varying spine diameter (minimum 1.5 μm), varying spine density (minimum 0.5/sq μm) and relatively little variation in spine height; and c) a spine-seta series which is characterized by relatively constant stalk diameter (0.5 μm) and stalk density (0.5/sq μm) and by varying stalk height and tip shape. The three series appear to be linked at the spine morphotype. The development of individual spines has not yet been described, but extensions of the processes implicated in the morphotypic series provide tentative hypotheses for their evolutionary relationships. The spine that forms the base of the *Chamaelinorops* spine-seta series (0.5 μm diameter; 0.5 stalks/sq μm) has the same spacing and height as the smallest and simplest member of the large spine series. A discrete change in the developmental program, perhaps an acceleration (*sensu* Alberch *et al.*, 1979) in the growth in diameter at a stage after the position and density of the spine stalks were set, would convert the 0.5 μm diameter spine into the 1.5 μm diameter spine. Local variation in the developmental program for density and diameter could yield the rest of the large spine se-

ries. The link between the *Anolis*-type spine and the *Chamaelinorops* spine involves merely a change in density or spacing from about 1/sq μm to 0.5/sq μm . This change might also be termed an acceleration (Alberch *et al.*, 1979) in the reduction of spinule density relative to the changes in the spine diameter and height.

These two shifts in the developmental program—an acceleration in spine density and in spine diameter—which could yield the two distinctive *Chamaelinorops* series are more likely to have occurred early than late in the evolution of the pad complex. If a) the *Anolis* spine-seta series was already present, and b) a shift in the developmental program affected the entire subdigital Oberhautchen, all members of the morphotypic series, including the spinules and small spines, would be shifted to a lower density. While the density of the spinules in *Chamaelinorops* is lower than that in *Anolis*, the *Anolis*-type spine with a 0.5 μm diameter and a 1.0/sq μm density would not occur given this scenario. An alternative possibility, that the shift in density occurred only over the fully exposed portions of the scale, would affect the density of the seta morphotype, but not that of spines and the intermediate morphotypes at the base of the scale. Neither scenario will produce the *Chamaelinorops* morphology. Among anoles which exhibit any variation in the series, it is only the seta which is different. If the *Chamaelinorops* fine structures were derived from that of *Anolis*, the comparative data would suggest that only the seta should be unique and not the prong and the spine. The most plausible interpretation of the character of the three series and their linkage through the spine morphotype is that the shift to a 0.5/sq μm density took place before setae differentiated, when the scale surface was spinate. The *Chamaeleolis*-*Phenacosaurus* spine-seta series and the *Chamaelinorops* spine-seta series would then have evolved in parallel (see Fig. 1). If this is the case, the

pre-anoline genera, *Enyalius*, and the para-anoline genera might exhibit comparable variation in spine and/or series density (Fig. 1B). The 0.5 stalks/sq μm spacing of *Chamaelinorops* does occur in *Enyalius*, but the issue needs to be examined in light of comparative data from a number of related iguanid genera (Fig. 1B). A variety of other features, particularly the unique shape and dimensions of the seta tip, would support the argument that the *Chamaelinorops* spine-seta series evolved in parallel with that of *Anolis*.

The opposite argument—that the fine structure of *Chamaelinorops* is derived from that of *Anolis*—can be made based on the similarity between the *Chamaelinorops* and *Anolis* series in stalk diameter and on the parallels between derived anoles and *Chamaelinorops* in seta distribution, in the presence of large, pyramidal spines, and on the weak parallels between *Chamaelinorops* setae and those of *A. sp. n. near eulaemus*. The unique setal shape and dimensions of *Chamaelinorops* might arise through an early arrest in seta development or paedomorphosis. Early cessation of growth in stalk height could leave a larger, untapered “end” of the stalk, and a partial differentiation of this “end” into a tip might produce a thick, square tip. The lower density of the spine-seta series is almost certainly an independent change in the developmental sequence and presumably represents the modification of a much earlier stage in the program. It is more difficult to interpret the series density as a derivative of the *Anolis* condition (see above). The argument that the fine structure is derived from the *Anolis* condition is supported by albumin immunological distance data which suggest that *Chamaelinorops* is close to a relatively generalized Hispaniolan anole, *A. cybotes* (Wyles and Gorman, 1980).

The relative consistency of spine and seta shape and density among *Anolis*, *Chamaeleolis*, and *Phenacosaurus* sug-

gests broad similarity among the forms in the prehensile and adhesive functions of the fine structure (Hiller, 1968; Peterson and Williams, 1981). Behavioral and ecological interpretations of the *Chamaelinorops* morphology are not yet possible; it is not even known if *Chamaelinorops* can maintain grip without its claws. Two specimens kept in captivity for two and four weeks were never observed to voluntarily adhere to glass or leaf surfaces and could not be encouraged to do so. The physical correlates of the leaf litter habitat are also poorly known. The presence of moist films and debris could profoundly affect the options for an adhesive or a prehensile grip based on the fine structure. There also appear to be significant differences in the morphological design of the system. The seta tip in *Chamaelinorops* has an area 3 to 6 times larger than that in *Anolis*, but a similar stalk cross-sectional area. If the setae work by adhesion and the entire flattened area of the tip is involved in an adhesive contact, the stress on the stalk would be 3 to 6 times greater in *Chamaelinorops*, and the stress created in the epidermal surface surrounding the stalk would be approximately 1.5 to 3 times greater. The *Chamaelinorops* setae stalks are shorter and therefore likely to be much stiffer than anole seta stalks. The oval stalk and tip are also thicker and presumably less flexible than those in *Anolis*. Both the increased stiffness of the seta and the absence of any reasonably consistent tip flexion which would orient the tip parallel to the substrate suggest that an individual *Chamaelinorops* seta would be substantially less effective in “finding” and making contact with the locomotor substrate than the anole seta. However, it is possible that the morphological “strategy” for adhesion is simply an alternative to that in *Anolis*, *Phenacosaurus*, and *Chamaeleolis*. The *Chamaelinorops* “strategy” is a lower density of setae that are distributed over the entire subdigital surface; each seta having a lower prob-

ability of establishing contact, but able to support 3 to 6 times more load if it does. This might be as effective a solution as that of the other anolines, but only if the functional tip area is very great, and the non-lamellar setae are as effective as those on the pad. Estimates of the setae numbers and tip areas illustrate this point. Comparing *A. sheplani* (MCZ 125641; 40.8 μm SV length) and *Chamaelinorops* (MCZ 156919; 31 mm SV length), pad area is much greater in *A. sheplani* (0.69 sq mm compared to 0.41 sq mm), but because of the longer toe and the presence of phalanx I setae, the actual seta bearing area is much greater in *Chamaelinorops* (1.24 sq mm compared to 0.98 sq mm). Because of the difference in density the total number of setae is twice as great in *A. sheplani* (1.2×10^6 setae compared to 0.62×10^6 setae; the number of setae is estimated from seta bearing area and seta density), but the total surface area of the setae tips is greater in *Chamaelinorops* (90.0×10^4 sq μm compared to 25.8×10^4 sq μm ; total tip area is estimated as mean tip area [from Table 1] $\times$ the number of setae). *Chamaelinorops* MCZ 156919 has unusually large seta tips (Table 1). If the tip area of MCZ 126708 is substituted for the actual value the sense of the comparison is unchanged, but the difference is reduced to the point where cumulative error in the estimate makes inference less certain (47.0×10^4 sq μm for *Chamaelinorops* and 25.8×10^4 sq μm for *A. sheplani*; an *Anolis* species with the body dimensions of *Chamaelinorops* would have an estimated tip area of 18.1×10^4 sq μm based on the scaling relationships developed for Puerto Rican anoles; Peterson *et al.*, 1982, and Peterson, unpublished data). For more typical *Chamaelinorops* tip areas the advantage in total tip area is less than a factor of 3 while the number of setae or possible points of contact is reduced by half. The *Chamaelinorops* and *Anolis* morphological adaptations are equal, alternative designs for adhesion only if a) all of the

flattened tip area in *Chamaelinorops* can make effective contact with the substrate, and b) the short and presumably stiffer *Chamaelinorops* setae are equally likely to "find" the substrate.

Some of the same arguments apply if the *Chamaelinorops* setae work like Velcro in micro-prehension. The greater stiffness and lower seta density would presumably reduce the number of potential contacts and the ability of the tips to "find" surface irregularities. The tip shape may also be less effective in prehension than that of *Anolis*. The anole seta retains a hook shape and has a wide, thin distal border which might catch microscopic irregularities as the seta is brushed along an environmental surface at the beginning of the propulsive phase (Peterson and Williams, 1981). In *Chamaelinorops*, the tip narrows distally, and it may have a rolled, thick distal edge. These could reduce the probability that the margin will engage an irregularity. Also, without some initial tip flexion toward the metatarsal-phalangeal joint, the distal margin of the tip may not form the leading edge as the digit makes contact with the substrate. Without this orientation the seta can not act like a hook and engage surface irregularities.

It is quite possible that neither of the seta mechanisms presumed in *Anolis* are relevant to *Chamaelinorops*. The distinctive features of the *Chamaelinorops* seta—the absence of a consistent tip flexion, reduced stalk density and increased stiffness—could be advantageous if the setae functioned to increase the frictional coefficient of the surface. If the leaf litter substrate is "slippery" because of moist films or debris, traction might become more important than adhesion or prehension. While it is far from clear how and in what context the setae of *Chamaelinorops* function, there are a number of reasons to suspect that their behavior and adaptive role is different from that of *Anolis* setae.

The large spine series of *Chamaelinorops* differs from most anole spine series

in that a hook tip is consistently absent. It is difficult to imagine how the large spines could act like hooks and form a prehensile, gripping scale surface. The pyramidal spines probably function like miniature keels and increase the roughness or frictional coefficient of the scale surface. The association of large spines with hillocks (which increase surface roughness in a larger dimensional range) and scale regions which have the most relief (e.g., the distolateral corners of the phalanx i scales) supports this interpretation. The series of fused spines converge on the dimensions and appearance of the keels.

Consideration of the possible functional significance of the differences between the *Chamaelinorops* and *Anolis* fine structure is highly speculative, but even interpretations based on alternative assumptions about the seta mechanism corroborate the purely morphological arguments that the *Chamaelinorops* morphology is significantly different from that in *Anolis*, *Chamaeleolis* and *Phenacosaurus*. Functional interpretations can not at this point resolve whether the *Chamaelinorops* morphology represents secondary terrestrial adaptation of an *Anolis*-like fine structure or an independent, perhaps even a less successful "experiment" in seta evolution.

CONCLUSIONS

The gross and fine structures of the *Chamaeleolis* and *Phenacosaurus* pads are similar and appear to be homologous with those of *Anolis*. The pad complex, including the spine-seta morphotypic series and lamellae, probably appeared relatively early in the anoline radiation before the differentiation of *Chamaeleolis*.

With the exception of the expanded pad position in *Phenacosaurus*, the differences between the genera and individual *Anolis* species are no greater than

the differences between even closely related *Anolis* species. If one views *Phenacosaurus*, *Chamaeleolis*, and such *Anolis* species as *cuvieri*, *sheplani*, *auratus*, *chrysolepis*, and *valencienni* as representatives of the taxonomic radiation of anolines, the pad complex exhibits radical differences in the number, distribution and shape of the lamellae, in phalangeal proportions, and in the distribution of setae and spines. The complex appears to be basically conservative in the spacing and diameter of spines and setae, the setae tip shape and dimensions, and the presence of a morphotypic series linking the spine and the seta.

Chamaelinorops has a unique pad fine structure, but resembles derived, more terrestrial anole species in gross morphology. The *Chamaelinorops* morphology may represent a highly specialized version of the *Anolis*-*Chamaeleolis*-*Phenacosaurus* subdigital pad or an independently derived pad complex.

ACKNOWLEDGMENTS

I am very grateful to Ernest E. Williams for discussions of this problem and for permission to examine material in his care and to E. E. Williams, D. B. Wake, A. P. Russell, and D. G. Buth for comments on the manuscript.

I thank J. Berliner and S. Beydler of the Department of Pathology, UCLA, for providing access to an excellent SEM facility. L. Meszoly, M. Kowalczyk and H. Kabe prepared the plates. The study was supported by a UCLA University Research Grant.

LITERATURE CITED

- ALBERCH, P., S. J. GOULD, G. F. OSTER, AND D. B. WAKE. 1979. Size and shape in ontogeny and phylogeny. *Paleobiology*, 5: 296-318.
BOULENGER, G. 1885. Catalogue of the lizards in the British Museum (Natural History), 2nd edition, 2. London, Taylor and Francis.

- ETHERIDGE, R. F. 1960. The Relationships of the Anoles (Reptilia: Sauria: Iguanidae): An Interpretation based on Skeletal Morphology. PhD. Thesis, Univ. Michigan, University Microfilms, Inc., Ann Arbor, Michigan, xiv + 236 pp.
- HILLER, U. 1968. Untersuchungen zum Feinbau und zur Funktion der Haftborsten von Reptilien. *Z. Morphol. Tiere*, **62**: 307–362.
- LAZELL, J. D. 1969. The Genus *Phenacosaurus* (Sauria: Iguanidae). *Breviora Mus. Comp. Zool. No. 325*, pp. 1–24.
- LIEM, K. F. 1973. Evolutionary Strategies and Morphological Innovations: Cichlid Pharyngeal Jaws. *Syst. Zool.*, **22**: 425–441.
- LILLYWHITE, H. B., AND P. F. A. MADERSON. 1968. Historical Changes in the Epidermis of the Subdigital Lamellae of *Anolis carolinensis* during the Shedding Cycle. *J. Morphol.*, **125**: 379–402.
- MADERSON, P. F. A. 1970. Lizard glands and lizard hands: Models for evolutionary study. *Forma et Functio*, **3**: 179–204.
- OLSON, E. C. 1975. Permo-carboniferous paleoecology and morphotypic series. *Am. Zool.*, **15**: 371–391.
- PAULL, D., E. E. WILLIAMS, AND W. P. HALL. 1975. Lizard karyotypes from the Galapagos Islands: Chromosomes in phylogeny and evolution. *Breviora Mus. Comp. Zool.*, No. 441, pp. 1–31.
- PETERSON, J., S. BENSON, J. MORIN, M. NGAI, AND C. OW. 1982. Scaling in tensile “skeletons”: I. Structures with scale independent length dimensions. *Science*, **217**: 1267–1270.
- PETERSON, J. A., AND E. E. WILLIAMS. 1981. A Case Study in Retrograde Evolution; The *Onca* Lineage in Anoline Lizards. II., Subdigital Fine Structure. *Bull. Mus. Comp. Zool.*, **149**: 215–268.
- RUIBAL, R. 1968. The ultrastructure of the surface of lizard scales. *Copeia*, **1968**: 698–704.
- RUIBAL, R., AND V. ERNST. 1965. The structure of the digital setae of lizards. *J. Morphol.*, **117**: 271–294.
- RUSSELL, A. 1975. A contribution to the functional analysis of the foot of the tokay, *Gekko gekko* (Reptilia: Gekkonidae). *J. Zool. Lond.*, **176**: 437–476.
- . 1976. Some comments concerning interrelationships among gekkonine geckos, pp. 217–244. In A. d'A. Bellairs and C. B. Cox (eds.), *Morphology and Biology of Reptiles*. Linnean Society Symposium, Series 3, London, Academic Press.
- . 1979. Parallelism and integrated design in the foot structure of gekkonine and diplodactyline geckoes. *Copeia*, **1979**: 1–22.
- SCHWARTZ, A. 1974. A new species of primitive *Anolis* (Sauria, Iguanidae) from the Sierra de Baoruco, Hispaniola. *Breviora Mus. Comp. Zool. No. 423*, pp. 1–19.
- STORK, N. E. 1980. A scanning electron microscope study of tarsal adhesive setae in the Coleoptera. *Zool. J. Linn. Soc.*, **68**: 173–306.
- WILLIAMS, E. E. 1976. West Indian anoles: A taxonomic and evolutionary summary. 1. Introduction and a species list. *Breviora Mus. Comp. Zool. No. 440*, pp. 1–21.
- . 1977. 3. The macrosystematics of the anoles, pp. 122–131. In E. E. Williams (ed.), the third *Anolis* Newsletter. Cambridge, Mass., Museum of Comparative Zoology.
- WILLIAMS, E. E., AND J. A. PETERSON. 1982. Convergent and alternative designs in the adhesive pad of scincid lizards. *Science*, **215**: 1509–1511.
- WYLES, J. S., AND G. C. GORMAN. 1980. The Classification of *Anolis*: Conflict between Genetic and Osteological Interpretation as Exemplified by *Anolis cybotes*. *J. Herpetol.*, **14**: 149–153.

The Axial Skeleton of *Chamaelinorops*

KARL FORSGAARD¹

ABSTRACT. The axial skeletons of the beta-anoles *Chamaelinorops barbouri* and *Anolis garmani* are described in detail. The caudal transverse processes of *Chamaelinorops* are found to be completely unlike those of other beta-anoles, and thus the close relationship that Etheridge (1960) proposed seems dubious. The zygapophysial plates and other vertebral specializations of *Chamaelinorops* are hypothesized to have evolved to allow an exceptionally strong contraction of the long epaxial musculature, possibly to aid in jumping or in the motions of the head during display behavior.

INTRODUCTION

Etheridge (1960) divided the species of anoline lizards into two groups, alpha and beta. This entire assemblage of small New World iguanids then comprised eleven genera, with the genus *Anolis* containing over 200 species and the other genera being monotypic or containing only a few species. Etheridge (1960) synonymized six of these genera with *Anolis*: *Audantia* (2 species), *Mariguana* (1 sp.), *Diaphoranolis* (1 sp.), *Xiphocercus* (2 sp.), *Deiropyx* (2 sp.), and *Norops* (4 sp.). Williams (1974) additionally synonymized *Tropidodactylus* (1 sp.) with *Anolis*. Currently, only four genera of anoles are recognized: *Anolis*, *Chamaeleolis*, *Chamaelinorops*, and *Phenacosaurus*. Etheridge (1960) classified *Chamaeleolis*, *Phenacosaurus*, and most *Anolis* as the alpha section of anolines; *Chamaelinorops* and the rest of *Anolis* as the beta section.

The alpha-anoles are characterized by the absence of transverse processes on caudal vertebrae posterior to the point where autotomy arises; beta-anoles have such processes. Etheridge (1960) defended the validity of this scheme by noting that caudovertebral structure is useful in distinguishing other lizard families as well as non-anoline iguanids. His phyletic hypotheses, however, have an element of uncertainty, and he admitted the possibility that caudovertebral transverse processes could have been independently lost in several lines of *Anolis*.

Perhaps the most questionable assignment in his phylogeny is the position of *Chamaelinorops* as a beta-anole. Its vertebrae are quite unlike those of other anoles, and, by Etheridge's own admission, are even significantly different from those of all other lizards. He noted (1960) that "each of the vertebrae of the neck and body has a pair of broad, flat, laterally projecting winglike processes arising just below the base of the neural arch and extending between the pre- and postzygapophyses." In the tail region he observed (1960) "the presence of heavily ossified structures" and added, "I am unable to determine just what these structures are from radiographs. They do not appear to be processes of the same sort as are found in the [trunk] vertebrae, yet they seem to be closely associated with, and possibly processes of, the caudal vertebrae." That Etheridge did not directly examine the axial skeleton of *Chamaelinorops* creates considerable doubt about the validity of the position of the genus within his an-

¹University of Washington School of Law, Seattle, Washington 98105, U.S.A. (Present address: 1111 Third Ave., Suite 1500, Seattle, Washington 98101, U.S.A.)

oline phylogeny, particularly in light of the singular importance of caudovertebral structure to that phylogeny.

Relevant skeletal data on *Chamaelinorops* are not available in the literature. Previous works on the external features and ecology of the genus are to be found in Schmidt (1919), Cochran (1928, 1941), Etheridge (1960), Thomas (1966), Schwartz and Thomas (1975), and Schwartz and Inchaustegui (1980). Hoffstetter and Gasc (1969) noted the lack of caudal autotomy, but did not elucidate any other skeletal characteristics. *Chamaelinorops barbouri* is the sole extant representative of the genus, and is restricted to the mountainous areas of Hispaniola. In this paper I provide a detailed osteological description of the axial skeleton of *Chamaelinorops barbouri* (Fig. 1) and compare it to that of the beta-anole *Anolis garmani* (Fig. 2).

DESCRIPTION OF THE AXIAL SKELETON OF *CHAMAELINOROPS BARBOURI*

The vertebral column of *Chamaelinorops* consists of approximately 68 procoelous vertebrae: 8 cervical, 11 thoracic, 5 lumbar, and 2 sacral vertebrae comprise the trunk vertebral series, and approximately 42 comprise the caudal series. The centra are constricted midway between their anterior and posterior ends.

Cervical Vertebrae. There are eight cervical vertebrae in *Chamaelinorops*, the posterior four of which articulate with ribs. The anterior four have hypapophyses, as do the anterior four of all anoles examined.

The atlas has its anterior face ringlike and tilted so that the dorsal border is posterior to the ventral border. It is typical of the "Category B" atlas described by Hoffstetter and Gasc (1969). The ring is formed ventrally by the intercentrum, which projects ventrally as a hypapophysis, and dorsolaterally by the neural arch, which is composed of two flat, oblong

plates, compressed posteriorly, which are separated in the dorsal midline and are overlapped by a forward projecting process on the neural spine of the axis. The atlas of *Anolis garmani* is longer and more massive, and the lateral processes of its neural arch do not extend as far laterally, nor does its hypapophysis extend as far ventrally.

The axis is comprised of neural arch and centrum, intercentra nos. 2 and 3, and the centrum of the atlas. Sutures between these elements are clearly seen in *A. garmani*, but were not evident in any of the *Chamaelinorops* specimens examined. The neural arch of the axis in *Chamaelinorops* has a large, laterally compressed neural spine which projects anteriorly over the neural arch of the atlas as well as posteriorly, like the spines of the remaining trunk vertebrae. The dorsal crest of this spine is a rounded ridge which slants so that its posterior end is dorsal to its anterior end. Also arising from the neural arch dorsolaterally are the zygapophyses. The prezygapophyses project forward and underlie the postzygapophysial area formed by the dorsomedial surface of the atlas neural-arch plates. The postzygapophyses of the axis project laterally as flat plates so that their rounded tips are at a distance from each other about four times the diameter of the axis condyle. The axis of *A. garmani* is similar to that of *Chamaelinorops*, except that its hypapophysis is ventrally flattened into a hypapophysial plate, its prezygapophyses extend farther anteriorly, its postzygapophyses extend laterally rather than dorsolaterally, and its postzygapophysial width is not as great. The hypapophysial plate, similar to the neural-spine splate of *Chamaelinorops*, was also noted in the alpha-anole *A. balearicus* and in the beta-anoles *A. biporcatus*, *A. fuscoauratus*, *A. chrysolepis*, *A. nebulosus*, and *A. onca*.

In *Chamaelinorops*, the axis postzygapophyses articulate with the prezygapophyses of the third cervical vertebra, the postzygapophyses of which extend even

Figure 1. Representative elements of axial skeleton of *Chamaelinorops* (A–B: MCZ 146960, C–F: MCZ 146958). A. Atlas. B. Axis. C. Thoracic vertebra no. 17. D. Sacrum. E. Caudal vertebra no. 5, minus the caudal chevron and one of the transverse processes. F. Caudal vertebra no. 11. For each element, the five views represent lateral, dorsal, ventral, anterior, and posterior views from top to bottom, left to right.

Abbreviations: atc, atlanto-centrum process of the axis; atp, atlantal plate; as, autotomy spetum; bar, base of the atlas ring; cch, caudal chevron; cd, condyle; f, foramen; g, groove; gld, groove-like depression; hp, hypapophysis; hppl, hypapophysial plate; m, medial edge of the zygapophysial plate; ns, neural spine; nsp, neural-spine plate; pp, parapophysis; prz, prezygapophysis; ptz, postzygapophysis; pzp, prezygapophysial plate; s, spine; tp, transverse process; tpp, transverse-process plate; zp, zygapophysial plate.

Figure 2. Representative elements of axial skeleton of *Anolis garmani* (MCZ 140048). A. Atlas. B. Axis (composite view including MCZ 7630). C. Thoracic vertebra no. 16. D. Sacrum. E. Caudal vertebra no. 5. F. Caudal vertebra no. 11 (note lack of terminal bifurcation of transverse process). Views identical to Figure 1.

Abbreviations: see Figure 1.

further apart laterally. This increase in lateral extension of the zygapophyses continues in *Chamaelinorops* so that at cervical vertebra no. 8, the vertebral width is about five times the diameter of the condyle of the eighth vertebra (a diameter that is considerably greater than the condyle diameter of the axis) (see Fig. 3). In *A. garmani*, the reverse occurs: the ratio of the zygapophysial width to the condyle diameter *decreases* from the axis to the middle thoracic vertebrae. In all species examined other than *Chamaelinorops*, the zygapophysial width of virtually all of the presacral vertebrae was approximately that of the postzygapophyses of the axis, and none attained the zygapophysial width of *Chamaelinorops*, and the width of the vertebral body was greatest in the cervical region and just anterior to the sacrum, with the thoracic vertebrae having slightly thinner bodies.

On vertebrae nos. 2 to 6 in *Chamaelinorops*, the pre- and postzygapophyses extend laterally so that they remain distinct from one another in a dorsal view. But beginning with vertebra no. 7, a zygapophysial plate extends laterally between the pre- and postzygapophyses and includes them, so that the prezygapophyses comprise the anterolateral corners of the plate, while the postzygapophyses comprise the posterolateral corners of the plate. No such zygapophysial plate occurs in the vertebrae of *A. garmani* or in those of any other member of the genera *Anolis*, *Chamaeleolis*, and *Phenacosaurus* that were examined in this study.

The neural spines of *Chamaelinorops* vertebrae nos. 3 to 5 are smaller than that of the axis. There is relatively little variation in neural-spine height throughout the rest of the vertebral column. Among other anoles there is considerable variation in the relative height of the neural spines. These spines are low ridges on the midline of the neural arch in *Chamaeleolis*, while in *A. bimaculatus* the neural spines constitute two-thirds of

the vertebral height. In *A. valencienni*, the neural spines are highest in the cervical region; in *A. ophiolepis*, *A. sagrei*, *A. equestris*, and *A. coelestinus* the neural spines are highest in the anterior caudals; in the other anoles examined, as in *Chamaelinorops* and *A. garmani*, the height of the neural spines is fairly constant throughout the column, except for diminution in the posterior caudals.

On *Chamaelinorops* vertebra no. 6 there is a flattening of the posterodorsal area of the neural spine, and on vertebra no. 7 this flattened area widens to form a plate; this neural-spine plate is found on all vertebrae posterior to no. 7, except for the posterior caudals. The anterior region of each neural-spine plate is usually, but not always, narrower than the posterior region of each plate. The dorsal surface of each plate is usually rugose. Neural-spine plates were found in most of the anoline species examined, but only *A. cybotes* had plates nearly as large as those of *Chamaelinorops*. Small neural-spine plates occur in *Phenacosaurus*, *A. latifrons*, *A. equestris*, *A. coelestinus*, *A. carolinensis*, *A. baleatus*, *A. bimaculatus*, *A. fuscoauratus*, *A. chrysolepis*, *A. nebulosus*, and *A. sagrei*. Knobs occur on the crests of the neural spines of *A. angusticeps*, *A. onca*, *A. garmani*, *A. valencienni*, and *A. ophiolepis*, and there is no buildup of the neural spine at all in *Chamaeleolis*. It is perhaps significant that *A. angusticeps* and *A. onca* are at the end points of two of the four main lineages in the Etheridge (1960) phylogeny, and that *A. garmani*, *A. valencienni*, and *A. ophiolepis* are in advanced positions on a third lineage; retention of the neural-spine plate appears from this limited data to be a sign of primitiveness.

All the cervical vertebrae posterior to the axis in *Chamaelinorops* bear parapophyses projecting ventrolaterally from the base of the neural arch; the parapophyses of vertebrae nos. 5 to 7, but not no. 8, are considerably larger than those of any other presacral vertebrae (see Fig. 1B). Unlike *Chamaelinorops*, the other

Figure 3. Trunk vertebrae of *Chamaelinorops* (MCZ 154450). A. Dorsal view of the presacral vertebrae, minus the atlas. B. Ventral view of the presacral vertebrae, minus the atlas. C. Dorsal view of the sacral and anterior caudal area.

anoles examined have the parapophyses of cervical vertebra no. 8 enlarged like those of nos. 5 to 7. In the parapophysial region of the axis, in both *Chamaelinorops* and *A. garmani*, are broad, round, anterolaterally projecting processes which make contact with the short, caudolaterally projecting parapophyses of the atlas; above and between these processes the floor of the neural canal of the axis projects forward as a process which is composed of the centrum of the atlas. The base of the atlas ring surrounds this process, much like the mammalian odontoid.

Thoracic and Lumbar Vertebrae. There are 16 thoracic and lumbar vertebrae in *Chamaelinorops* (nos. 9–24). The lumbar vertebrae are the posterior five of these, lacking ribs and parapophyses on or near

the centrum (they all have small, rounded projections from the portion of the neural arch below the base of the zygapophysial plate). The thoracic vertebrae are the anterior 11; the first ten of these articulate with movable ribs, while the ribs of the last are fused with the parapophyses (see Fig. 5). In *Chamaelinorops* these thoracic and lumbar vertebrae are essentially similar in non-parapophysial structure.

The zygapophysial plates of the thoracic and lumbar vertebrae have centrally located foramina which seem to indicate the point where the pre- and postzygapophyses fuse to form a solid plate. The postzygapophysis comprises the ventral surface of the posterior border of each plate in *Chamaelinorops*, which overlaps the dorsal surface of the anterior border

of the plate of the adjacent posterior vertebra. In addition, the postzygapophysis is composed of the thickened lateral margin of each plate, the posterior portion of which extends, as a spinelike process, underneath the anterolateral border of the plate of the adjacent posterior vertebra. This spine articulates along its length with a groovelike depression in the ventral surface of the plate of the posterior vertebra. Thus the zygapophysial articulation of thoracic and lumbar vertebrae is an interlocking one, but not at all similar to the zygosphenzygantrum articulation found in some lepidosaurs and noted by Romer (1956). In *Chamaelinorops* this spine-groove mode of articulation occurs on the postzygapophyses of vertebrae nos. 8 to 22, but not in all individuals examined. No other anole examined has this mode of articulation.

Additional articulation between the thoracic vertebrae is provided by grooves alongside the anterior face of the neural arch. These are dorsally directed grooves extending for a short distance posteriorly from the anterior face of the vertebra, along the juncture of the arch and the zygapophysial plate. Each groove articulates with the ventrally directed medial edge of the zygapophysial plate of the adjacent anterior vertebra; this latter plate extends posteriorly from the posterior face of the neural arch, so that a right angle is formed with the posterior edge of the plate. This plate-groove mode of articulation occurs on the anterior face of vertebra no. 3 through the posterior caudals. Romer (1956) does not mention the plate-groove mode of intervertebral articulation, but it is found in all the anoles examined, including *A. garmani*. However, in the anoles other than *Chamaelinorops*, the articulation seems to be the result only of the close proximity of the postzygapophyses to the neural arch of the adjacent posterior vertebra. Only in *Chamaelinorops* is the groove deep and the articulation well developed.

Of the five lumbar vertebrae, the two

that are immediately anterior to the sacrum are shorter than the anterior lumbar vertebrae, and hence have shorter zygapophysial plates. These plates also do not extend laterally as far as those of the anterior vertebrae, and the pre- and postzygapophyses of each vertebra are distinct from one another, as they are in the anterior cervical vertebrae (see Fig. 3C). Although five lumbar vertebrae have been recognized in *Chamaelinorops*, Etheridge (1960) notes that the number of lumbar vertebrae in anoles is usually three or four but may be as high as seven. He also cites intraspecific variation in this number, and concludes that "the number of lumbar vertebrae is not a particularly useful taxonomic character." The exact number of true lumbar vertebrae is determined by counting the number of presacral vertebrae without rib articulations. This is usually only possible in alcoholic specimens. Only one specimen of *Chamaelinorops* and none of the other anoles in this study were examined for this feature. However, the two or three vertebrae just anterior to the sacrum were examined for all species in this study and the other anoles were found to differ from *Chamaelinorops* in that the width of the vertebral body just anterior to the sacrum is greater, rather than less, than the width of the body of the thoracic vertebrae.

Sacrum. The sacrum in *Chamaelinorops* is comprised of two vertebrae. The posterior centrum is slightly longer than the anterior centrum. Their neural spines are of the same height as the adjacent anterior and posterior vertebrae. However, the neural spine of the posterior sacral vertebra is twice as long as that of the anterior vertebra, and thus the neural-spine plate of the posterior vertebra has a greater surface area than that of the anterior vertebra. In *A. garmani*, the neural spine of the posterior sacral vertebra is somewhat higher than that of the anterior one. The zygapophyses of the sacral vertebrae in *Chamaelinorops* do not extend as far laterally as those of the lumbar vertebrae, and those of the pos-

terior sacral vertebra are narrower than those of the anterior one. Their parapophyses are massive and extend laterally so that their tips are about as far from each other as are the tips of the widest zygapophysial plates of the thoracic vertebrae. The ends of these parapophyses appear to fuse distally, providing a broad articulating surface for the pelvic girdle. The sacral parapophyses of *A. garmani*, in addition to the other anoles examined, are less massive than those of *Chamaelinorops* and are unfused.

Caudal Vertebrae. There are approximately 42 caudal vertebrae in *Chamaelinorops*. They are not autotomic. In *A. garmani* autotomy occurs and autotomy septa are plainly visible. In *Chamaelinorops* and in *A. garmani*, chevrons occur on both anterior and posterior caudal vertebrae. In *Chamaelinorops* these chevrons articulate with the ventrolateral surface of the centrum just anterior to the condyle, and extend caudoventrally so that their tips are ventral to the anterior portion of the centrum of the posterior vertebra. The anterior caudal chevrons in *A. garmani* are much longer than in *Chamaelinorops*, and consequently their tips are ventral to the posterior portion of the centrum of the posterior vertebra. Caudal chevrons of posterior caudal vertebrae in *A. garmani* are considerably shorter.

The centrum length of caudal vertebrae in *Chamaelinorops* increases progressively posteriorly from the sacrum, so that the length of the centrum of caudal vertebra no. 11 is about three times the length of the centrum of the anterior sacral vertebra. The centrum lengths of caudal vertebra no. 11 and subsequent posterior vertebrae are similar, so that the posterior caudal centra, while having diameters substantially less than that of caudal no. 5, are only slightly shorter in length (see Fig. 4). In *A. garmani*, the centrum length of caudal vertebrae increases posteriorly to only about twice the length of the anterior sacral vertebra.

The parapophyses (transverse processes) of caudal vertebra no. 1 in

Chamaelinorops extend laterally, slightly posteriorly, and slightly ventrally. They are as thick as those of the sacral vertebrae, and slightly greater in length. Those of caudal vertebra no. 2 are dorsoventrally compressed, and their lateral ends are thickened and extend both posteriorly and anteriorly (see Fig. 4). Posterior to caudal vertebra no. 5, these thickened ends of the transverse processes become flattened into transverse-process plates in a plane perpendicular to that of the neural-spine plates and parallel to the dorsal midline. The lateral extension of the transverse processes decreases posteriorly. At about caudal vertebra no. 12 they are approximately three-quarters the length of the centrum. At caudal no. 18 they are half the length of the centrum and considerably less massive, and at caudal no. 20 they are

Figure 4. Dorsal view of caudal vertebrae nos. 1-21 of *Chamaelinorops* (MCZ 146957). Caudal vertebrae posterior to no. 21 are essentially similar to it. Prezygapophysial plates are not well developed in this specimen.

Abbreviations: see Figure 1.

merely longitudinal ridges on the ventrolateral surface of the centrum (see Fig. 4). In contrast, none of the other anoles examined had any expansions on their caudal transverse processes. The transverse processes on the autotomic caudal vertebrae of *A. garmani* are all thin and anterolaterally directed, unlike the large laterally directed processes in *Chamaelinorops*. Furthermore, while Etheridge (1967) asserted that the anterolaterally directed caudal transverse processes of beta-anoles like *A. garmani* are "unusual in being terminally bifurcate in the vertical plane," no such bifurcation was found in the two specimens of *A. garmani* examined.

The neural spines of the caudal vertebrae in *Chamaelinorops* continue the thoracic and lumbar trend of bearing flat plates. Those neural-spine plates increase in size between caudals no. 1 and no. 3, in accordance with the increase in centrum length. This large size is maintained further posteriorly than is the size of the transverse processes. The neural spines of the caudal vertebrae of *A. garmani* do not bear neural-spine plates and are much higher, in relation to the centrum diameter, than those of *Chamaelinorops*, and are also much more posteriorly directed.

The caudal zygapophyses in *Chamaelinorops* remain large but, with the increase in centrum length, lose their platelike structural contiguity as the pre- and postzygapophyses become distinct from one another. The prezygapophyses of caudal vertebra no. 2 extend further laterally than do the postzygapophyses with which they articulate, and the lateral portion of the prezygapophysis that is not overlapped is upturned around the lateral margin of the postzygapophysis. Posteriorly from caudal no. 2, it is upturned more, so that by caudal no. 4, the prezygapophyses are lengthened and their lateral edges, which are upturned, support long flat plates in a plane parallel to and below that of the neural-spine plate. These prezygapophysial plates are

abruptly lost posterior to caudal vertebra no. 8 in specimen MCZ 146958. This trait, however, is subject to intraspecific variation, as the lateral edges of the prezygapophyses of the anterior caudal vertebrae in specimen MCZ 146957 (see Fig. 4) are thickened and upturned but do not bear such plates. A similar prezygapophysial plate does not occur in the anterior caudal vertebrae of *A. garmani* nor the other anoles examined.

In his analysis of the caudovertebral structure of all lizards, Etheridge (1967) makes almost no mention of neural spines or chevrons. However, his evaluation of transverse processes is taxonomically useful and thorough, except that he lumps *Chamaelinorops* in a group with beta-anoles like *A. garmani*, characterized by long transverse processes whose orientation changes sequentially from posterolateral "to lateral then acutely anterolateral." Such a description of the caudovertebral structure of *Chamaelinorops* ignores its major structural uniqueness as compared with other anoles and thus is not adequate to allow acceptance of his association of *Chamaelinorops* with typical beta-anoles.

Ribs. There are 15 pairs of ribs in *Chamaelinorops*. Each pair of ribs, except the most posterior pair, consists of two parts: the curved dorsal bony part and the ventral cartilaginous section, which in the ribs of vertebrae nos. 9 to 18 forms a "parasternum." The fused ribs of vertebra no. 19 have no cartilaginous parts associated with them. There are four pairs of cervical ribs, the first pair being about three-quarters the length of the second pair, which are about half the length of the third pair. The fourth pair is about the same length as the third, and the distal ends of each of these pairs appears to be in close association with the scapulocoracoid. The anterior three pairs of cervical ribs articulate with the enlarged parapophyses of vertebrae nos. 5 to 7, and the capitulum of these ribs is correspondingly enlarged, as is each rib's shaft diameter. The posterior pair of cervical

ribs articulate with the parapophyses of vertebra no. 8, which are similar to those of the thoracic vertebrae, and consequently the capitulum size and average shaft diameter of each of the posterior pair of cervical ribs are much closer to that of the thoracic ribs than to that of the anterior cervical ribs. Each of the anterior two pairs of cervical ribs bears a small, flat, triangular process on its lateral surface, with the point of the triangle directed posterolaterally (see Fig. 5).

The cartilaginous tips of the anterior two pairs of thoracic ribs in *Chamaelinorops* articulate directly with the sternum. The third, fourth, and fifth pairs of thoracic ribs articulate indirectly with the sternum via a cartilaginous xiphisternum. This sternal: xiphisternal formula of 2:3 in the one specimen examined is counter to the claim of Etheridge (1960) that all anolines except *Phenacosaurus* (2:2) have a formula of 3:2. The sixth, seventh, eighth, and ninth pairs of thoracic ribs are

attached inscriptional ribs, with each pair meeting at the midventral line to form an anteriorly directed cartilaginous chevron. Posterior to the cartilaginous chevron of the ninth thoracic rib pair is one free inscriptional rib chevron, which does not extend to the ventral ends of the cartilaginous tips of the bony elements of the tenth pair of thoracic ribs, to which it corresponds. This fixed-chevron to floating-chevron ratio of 4:1 in the one specimen examined differs from the ratios of 4:2 and 3:3 reported for *Chamaelinorops* by Etheridge (1965), indicating that intra-specific variation in that ratio may not be as conservative as he asserted. The ribs of the eleventh thoracic pair are the shortest ribs in the *Chamaelinorops* column, being about the same length as the parapophyses to which they are fused (see Fig. 5).

DISCUSSION

The axial skeleton of *Chamaelinorops* is radically different from that of *Anolis garmani* and the other anoles examined. The caudovertebral transverse processes of beta-anoles like *A. garmani* are slender and are not expanded at their distal ends. Those of *Chamaelinorops*, in contrast, are thick, and those of the anterior caudals posterior to caudal no. 5 bear distal expansions comprising transverse-process plates. These differences indicate oversimplification in the morphological and taxonomic analysis of Etheridge (1960, 1967) insofar as he describes a single type of caudovertebral structure purportedly shared by *Chamaelinorops* and beta-anoles. The caudal vertebrae of *Chamaelinorops* also differ from those of all other anoles in the large size of their neural-spine plates and in the presence of prezygapophysial plates in some specimens of *Chamaelinorops*.

The precaudal vertebrae of *Chamaelinorops* are also unique in several respects. Specializations of the precaudal vertebrae of *Chamaelinorops* include: 1)

Figure 5. Ribs and inscriptional ribs of *Chamaelinorops* (schematic) (MCZ 146960).

presence of "wing-like" zygapophysial plates; 2) greater zygapophysial width; 3) presence of spine-groove intervertebral articulation; 4) more highly developed plate-groove intervertebral articulation; 5) larger neural-spine plates; and 6) apparent fusion of the distal ends of the sacral parapophyses.

These specializations raise questions of function which remain unanswered. The author's review of other reptilian vertebrae has been far from exhaustive, but zygapophysial plates similar to those of *Chamaelinorops* have previously been noted in heavy-bodied arboreal snakes such as *Synopsis bicolor* (Johnson, 1955). He speculated that such plates strengthen the long epaxial musculature through greater area for muscular insertion and assist such snakes in "the habit of extending the unsupported body considerable distances in climbing from hold to hold." Therefore, it is possible that *Chamae-*

linorops might use its vertebral specializations to assist in supporting and lifting the forward portions of its body in such activities as jumping. However, this has not been functionally studied. Moermond (1974) found that certain morphological parameters (ratio of tail length to snout-vent length and ratio of humerus length to femur length) correlated with the percentage of jumps in total movements of seven sympatric species of *Anolis*. Measurements of these parameters in preserved specimens of *Chamaelinorops* (Table 1) predict that *Chamaelinorops* might be a "jumper."

Limited field and laboratory observations of *Chamaelinorops* do not support the hypothesis that it might be a specialized "jumper." However, adequate data are still lacking.

Of interest are observations by E. E. Williams and S. Case (personal communication) regarding aggressive male-

TABLE 1. MEASUREMENTS OF *CHAMAELINOROPS*.

Species		snout-vent length (mm)	tail length (mm)	femur length (mm)	humerus length (mm)	Length Ratio tail: sn-vent	Length Ratio humerus: femur	% jumps of total movements in field
<i>Chamaelinorops</i>	MCZ #							
	146951	41.28	78.41	13.34	7.30	1.899	0.547	—
	56143	39.60	85.75	11.48	6.35	2.165	0.553	—
	146956	37.91	83.59	11.87	6.34	2.205	0.534	—
	146953	36.44	80.44	10.94	6.46	2.209	0.590	—
	146952	32.66	56.71	8.74	4.42	1.736	0.506	—
	57728	31.79	67.70	9.42	6.60	2.130	0.701	—
	126708	26.59	61.85	7.81	4.75	2.326	0.608	—
	146955	26.70	53.68	7.56	4.14	2.049	0.548	—
	146954	24.74	47.75	7.18	4.44	1.930	0.618	—
	74594	22.36	43.12	5.98	3.31	1.928	0.554	—
	38253	22.00	46.38	6.49	3.23	2.108	0.498	—
<i>Chamaelinorops</i> (mean values)		—	—	—	—	2.062	0.569	?
* <i>A. semilineatus</i>		—	—	—	—	2.4	0.6	60
* <i>A. koopmani</i>		—	—	—	—	2.05	0.57	59
* <i>A. monticola</i>		—	—	—	—	2.3	0.56	57
* <i>A. hendersoni</i> (adult)		—	—	—	—	2.55	0.6	46
* <i>A. hendersoni</i> (juv.)		—	—	—	—	2.55	0.6	43
* <i>A. distichus</i> (in trees)		—	—	—	—	1.35	0.8	37
* <i>A. coelestinus</i>		—	—	—	—	2.05	0.8	26
* <i>A. distichus</i> (on rocks)		—	—	—	—	1.35	0.8	16
* <i>A. cybotes</i>		—	—	—	—	1.8	0.7	9

*Data from Moermond (1974).

male interactions of *Chamaelinorops*. These occurred in restrictive habitats in captivity. Williams noted that the "head is carried far backward past the vertical to display the small bicolor dewlap. The head is then waved slowly several times. No push-ups or head bobs" were noted, and "the dewlap is held extended throughout." Case noted that the head was "raised at least 90° from the horizontal axis" and that very slow head-bobs accompanied dewlap extension. This extreme displacement of the head during dewlapping behavior, possibly the greatest vertical displacement of the head among anoles, may require a firmly anchored epaxial musculature and thus may help explain the function of the vertebral specializations of *Chamaelinorops*, but tests of this hypothesis have yet to be made.

ACKNOWLEDGMENTS

I am indebted primarily to my undergraduate thesis advisor, Ernest E. Williams, for loan of specimens and library material and for financial assistance, instruction, and inspiration throughout the course of this study. The author is also grateful to Benjamin Shreve, Catherine McGeary, Patricia Haneline, José Rosado, Susan Rhodin, Anders Rhodin, A. W. Crompton, William Fink, James Lazell, Christopher Kelley, Mark Plotkin, Ross Johnston, Charles Gougeon, William Amaral, Russell Mittermeier, Anthony Russell, Franklin Ross, Susan M. Case, George Lauder, Lisa Moeller, and Timothy Moermond. Illustrations were prepared by Laszlo Meszoly and the author.

APPENDIX: SPECIMENS EXAMINED

Four complete adult skeletons of *Chamaelinorops barbouri* were examined for this study. All had snout-vent lengths of approximately 36 mm.

Alpha-anoles examined: *Phenacosaurus heterodermus*: MCZ 145323; *Chamaeleolis porcus*: MCZ 131918; *Anolis latifrons*: MCZ 77410; *A. equestris*: MCZ 131609; *A. coelestinus*: MCZ

144795; *A. carolinensis*: MCZ 153516; *A. angusticeps*: MCZ 59245; *A. baleatus*: MCZ 7831; *A. bimaculatus*: MCZ 28717; *A. cybotes*: MCZ 134020.

Beta-anoles examined: *Chamaelinorops barbouri*: MCZ 38253, 56143, 57728, 74594, 126708, 146951-60, 154450; *Anolis biporcatus*: MCZ 24396; *A. fuscoauratus*: MCZ 140041; *A. chrysolepis planiceps*: MCZ 43859; *A. nebulosus*: MCZ 133994; *A. onca*: MCZ 139352; *A. garmani*: MCZ 7630, 140048; *A. valencienni*: MCZ 145321; *A. ophiolepis*: MCZ 11181; *A. sagrei*: MCZ 10498.

LITERATURE CITED

- COCHRAN, D. M. 1928. A new species of *Chamaelinorops* from Haiti. *Proc. Biol. Soc. Wash.*, **41**: 45-47.
- . 1941. The herpetology of Hispaniola. *Bull. U.S. Nat. Mus.*, (177): vii + 398 pp.
- ETHERIDGE, R. E. 1960. The relationships of the anoles (reptilia: sauria: Iguanidae): an interpretation based on skeletal morphology. *Ann Arbor, Michigan, University Microfilms, Inc.*, 236 pp.
- . 1965. The abdominal skeleton of lizards in the family Iguanidae. *Herpetologica*, **21**: 161-168.
- . 1967. Lizard caudal vertebrae. *Copeia*, **1967**: 699-721.
- HOFFSTETTER, R., AND J. P. GASC. 1969. Vertebrae and ribs of modern reptiles, pp. 201-310. *In* C. Gans, A. d'A. Bellairs, and T. S. Parsons (eds.), *Biology of the Reptilia*, Vol. I. London, Academic Press.
- JOHNSON, R. G. 1955. The adaptive and phylogenetic significance of vertebral form in snakes. *Evolution*, **9**: 367-388.
- MOERMOND, T. C. 1974. Patterns of habitat utilization in *Anolis* lizards. PhD thesis, Harvard University. Cambridge, Massachusetts, 150 pp.
- ROMER, A. S. 1956. *Osteology of the Reptiles*. Chicago, University of Chicago Press, 772 pp.
- SCHMIDT, K. P. 1919. Descriptions of new amphibians and reptiles. *Bull. Am. Mus. Nat. Hist.*, **41**: 523-524.
- SCHWARTZ, A., AND S. J. INCHAUSTEGUI. 1980. The endemic Hispaniolan lizard genus *Chamaelinorops* Schmidt. *J. Herpetol.*, **14**: 51-56.
- SCHWARTZ, A., AND R. THOMAS. 1975. A Check-list of West Indian Amphibians and Reptiles. Pittsburgh, Carnegie Museum of Natural History, 216 pp.
- THOMAS, R. 1966. A reassessment of the herpetofauna of Navassa Island. *J. Ohio Herpetol. Soc.*, **5**: 73-89.
- WILLIAMS, E. E. 1974. A case history in retrograde evolution: the *onca* lineage in anoline lizards. I. *Anolis annectens* new species, intermediate between the genera *Anolis* and *Tropidodactylus*. *Breviora Mus. Comp. Zool. No.* 421 pp. 1-21.

Variation in the Left Lung and Bronchus Of *Thamnophis sirtalis parietalis*

THOMAS S. PARSONS¹

LUDMELA DJATSCHENKO²

ABSTRACT. In a sample of 55 *Thamnophis sirtalis*, the tracheal attachment of the left lung varied greatly. A distinct bronchus was present in 12 and absent in 13. Another 21 showed bronchial structure ventrally but not dorsally. Nine were damaged and unclassified. We were unable always to recognize cartilaginous bronchial rings until the material was stained with toluidine blue. The size of the left lung varied significantly with the sex of the snake, and the relative size of the left lung is not a useful measure. Taxonomic conclusions should not be based on the examination of respiratory structures in unstained material or in only one or a few specimens.

INTRODUCTION

The usefulness of the structure of the respiratory system in the classification of snakes was suggested by Cope (1900) and more recently has been championed by Underwood (1967). However, there are few detailed studies on the lungs of snakes, and many workers simply note whether a left lung is present or absent and, if present, its length. Beddard (1906 and other papers), Brongersma (1957), Butler (1895), and Cope (1894 and 1900) generally provided more information and they studied many species; in most cases, however, only one or two specimens of each were studied. Bergman (1952, 1958, and other papers) usually studied a large number of specimens of each species, but noted almost nothing about the lungs. We

decided to investigate variation in the left lung of the garter snake, *Thamnophis sirtalis*, to test the validity of conclusions based on observation of a single specimen.

Earlier reports on the presence or absence of a separate left bronchus are contradictory, and therefore we concentrated on this feature. Cope (1900: 695-696) stated that in the Colubroidea (roughly equivalent to the Colubroidea of most authors or to the Caenophidia, excluding the Viperidae, of Underwood, 1967), the left lung is usually present, though very small, and is connected to the right by a "foramen, which perforates the tracheal cartilage It is sometimes connected to the dorsal [= right] lung by a short tube, in which cartilaginous half rings are seen in but two of the genera examined, viz *Heterodon* and *Conopsis*." Thus he concluded that a left bronchus was absent in most forms, including *Thamnophis* (= *Eutaenia* of Cope) *sirtalis parietalis*. Brongersma (1957: 303), on the other hand, says of the Colubridae (equivalent to the Dipsadidae, Homalopsidae, Natricidae, and Colubridae of Underwood, 1967) that "although the left bronchus is very short, it is usually distinct." He also studied *Thamnophis sirtalis parietalis* and did not note it as exceptional in this regard.

Part of the problem here may be in the definition of a bronchus, usually considered to be the single tube connecting a lung to the trachea. In snakes the divi-

^{1,2}Department of Zoology, University of Toronto, Toronto, Ontario, Canada M5S 1A1.

sion between the trachea, the right bronchus, and the lung is obscured by the frequent occurrence of a tracheal lung, i.e., pulmonary tissue in the trachea. We have adopted a fairly simple definition: a left bronchus is any more or less tubular connection between the left lung and the trachea, right lung, or right bronchus that contains cartilaginous rings and no alveolar tissue. Admittedly this is not a perfectly clear-cut distinction, but the presence or absence of cartilaginous rings and of alveolar tissue are determinable features and ones in which we found considerable variation.

We found it very difficult to determine the structure simply by inspection with a dissecting microscope. The cartilaginous rings are small and could not always be recognized without staining. Therefore, we stained all our specimens except for a few that were badly damaged during the initial dissection.

MATERIALS AND METHODS

This study is based on 55 specimens of the red-sided garter snake, *Thamnophis sirtalis parietalis*, 37 males and 18 females, ranging from 300 to 500 mm in snout-vent length. All were obtained from Nasco Educational Materials, Guelph, Ontario, and preserved in Nasco-Guard; some were later transferred to 70 percent ethanol. Specimens were dissected and studied under a stereoscopic microscope.

To determine the distribution of cartilaginous rings in the bronchi, we dissected free and removed the section of the respiratory tract for approximately 15 mm on either side of the attachment of the left lung. This section was then stained with toluidine blue, following the method given by Hildebrand (1968). The washing was done in tap water, and the specimens remained in the stain for six days. Destaining, as outlined by Hildebrand, was accomplished in two to three days. This process left the cartilage

a bright dark blue and all other tissues pale gray. Destaining continued very slowly in a mixture of absolute alcohol and benzene or in the 70% alcohol in which the specimens were stored.

OBSERVATIONS

All of the snakes studied had two lungs, a functional right and a vestigial left one (Fig. 1). The right lung begins dorsal and just anterior to the posterior tip of the heart, with the right lateral portion of the lung as the most anterior point. The anterior surface of the right lung slopes posteriorly for several millimeters to the anterior end of the left lateral border. The trachea runs along the ventral aspect of the right lung, midway between its right and left lateral borders, for approximately 10 mm, making the anterior portion of the lung appear somewhat bilobed. The trachea then disappears into the substance of the lung, but in some cases its dorsal wall contains alveolar tissue (Fig. 2). The functional respiratory tissue of the lung gradually becomes more attenuated posteriorly, leaving a delicate membranous air sac which ends in the vicinity of the right gonad.

The vestigial left lung is located maximally 3 mm anterior to or at the level of entrance of the trachea¹ into the substance of the right lung. It can be overlooked upon initial examination because of its small size and close adherence to the left side of the posterior vena cava. The shape of the left lung varies considerably. Triangular, oval, and circular lungs predominate and are encountered with about equal frequency; vermiform, pouchlike, and irregularly shaped lungs occur less frequently. The locations and lengths of both right and left lungs,

¹The trachea, as used here and in most descriptions of snakes, includes, posterior to the entrance of the left lung, the right bronchus.

expressed as percentages of snout-vent length, are given in Table 1.

The greatest variation in the anatomy of the respiratory organs among specimens involves the attachment of the left lung to the trachea or right lung. Due to the delicate consistency of the respiratory tissue and its rather colorless translucent appearance, in addition to the presence of mesenteries and blood vessels in the area, observations on individual snakes are somewhat subjective. However, in unstained specimens, there appeared to be four distinct patterns of attachment of the left lung. As shown below, these appearances are often deceiving.

In 23 of the 54 snakes the left lung seemed to open directly off the trachea; the alveolar tissue of the lung appeared to begin immediately adjacent to the trachea (Fig. 3).

In eight snakes there appeared to be an intervening tube of membrane between the actual opening into the trachea and the start of the alveolar tissue of the lung (Fig. 4). The membrane ranged from delicate and mesenterylike to stout and suggestive of the tissue found between the cartilaginous tracheal rings. The length of this membrane varied from 0.1 to almost 1.0 mm.

In 12 snakes the attachment appeared different in the ventral and dorsal aspects

TABLE 1. SIZES AND POSITIONS OF LUNGS.*

		Males	Females	Total	d and P values for differences between males and females
Number of specimens		37	18	55	
Snout-vent length (in mm)	min.	300	362	300	
	mean	394 $\pm$ 30.4	447 $\pm$ 36.6	411 $\pm$ 42.8	5.01 (.001)
	max.	476	500	500	
Position of anterior end of right lung	min.	15.1	14.3	14.3	
	mean	17.0 $\pm$ 0.81	16.2 $\pm$ 1.02	16.8 $\pm$ 0.78	2.85 (.01)
	max.	18.9	17.6	18.9	
Position of posterior end of right lung (including air sac)	min.	57.1	54.9	54.9	
	mean	62.3 $\pm$ 2.21	58.9 $\pm$ 2.20	61.2 $\pm$ 2.21	5.39 (.001)
	max.	66.2	63.5	66.2	
Relative length of right lung (including air sac) as a % of snout-vent length	min.	39.5	38.6	38.6	
	mean	45.3 $\pm$ 2.44	42.7 $\pm$ 3.61	44.4 $\pm$ 2.88	2.76 (.01)
	max.	49.8	47.2	49.8	
Position of anterior end of left lung	min.	16.1	15.3	15.3	
	mean	18.3 $\pm$ 0.81	17.3 $\pm$ 0.95	18.0 $\pm$ 0.86	3.30 (.001)
	max.	20.0	18.9	20.0	
Position of posterior end of left lung	min.	16.7	15.9	15.9	
	mean	19.0 $\pm$ 0.79	17.8 $\pm$ 0.79	18.6 $\pm$ 0.78	5.00 (.001)
	max.	20.5	19.4	20.5	
Relative length of left lung as a % of snout-vent length	min.	0.4	0.3	0.3	
	mean	0.7 $\pm$ 0.22	0.5 $\pm$ 0.21	0.7 $\pm$ 0.22	3.64 (.001)
	max.	1.3	0.9	1.3	
Absolute length of left lung (in mm)	min.	0.2	0.1	0.1	1.55 (.10)
	mean	0.28 $\pm$ 0.09	0.25 $\pm$ 0.07	0.27 $\pm$ 0.08	
	max.	0.5	0.4	0.5	

* The positions of lungs are given as percentages of the snout-vent length, with 0 the snout and 100 the vent.

Figures 1-4. Dissections showing the left lung of *Thamnophis* before staining with toluidine blue. Anterior is toward top of page. 1. General view to show the relationships between the two lungs and the trachea. (X 2.8). 2. Specimen showing alveolar tissue in the trachea (tracheal lung). (X 7). 3. Specimen in which the pulmonary tissue of the left lung appears immediately adjacent to the trachea; there is no left bronchus. (X 15). 4. Specimen in which the left lung appears to have a membranous tubular attachment to the trachea; subsequent staining with toluidine blue showed this attachment to actually be a left bronchus with cartilaginous rings. (X 17).

Abbreviations: A, Alveolar tissue of tracheal lung; L, left lung; R, right lung; T, trachea.

of the left lung. In ventral view, nine of these 12 snakes seemed to have a bronchus with cartilaginous rings joining the left lung to the trachea. The remaining three appeared to have a heavy membranous tube joining the two. The length of this bronchus or membrane ranged from about 0.3 to 1.0 mm. However, when the left lung was reflected so that the attachment of its dorsal aspect to the trachea was visible, the distance between the trachea and the start of the alveolar tissue of the lung appeared noticeably smaller due to the decrease in the length of the bronchus or membrane as it swung around to the dorsal aspect of the lung. In four cases the intervening bronchus or membrane disappeared completely, leaving the trachea and the alveolar tissue of the lung apposed.

Eleven of the snakes appeared to have a left bronchus (under our definition of a bronchus) joining the left lung to the trachea. The bronchi ranged in length from 0.3 to 1.0 mm. In most cases the separation between the functional respiratory tissue of the left lung and the bronchus was abrupt. In a few, the line of demarcation between the two structures was less obvious. The left lung of the fifty-fifth snake was destroyed during dissection, and is not considered here.

Due to the difficulty of detecting cartilaginous rings in unstained material, the left bronchi of 47 specimens (in eight this area was destroyed during the initial investigation) were stained and again studied without reference to the earlier observations. The two sets of observations were then compared. The most common pattern of attachment of the left lung was now shown to be the one that displayed a difference in the structure on the ventral and dorsal aspects of the left lung (Table 2; Figs. 5, 6). In all the snakes the tracheal cartilages were approximately 0.2 mm wide and about 0.5 to 0.8 mm apart when measured from the center of one to the center of the adjacent one. In the 21 snakes (11 males and 10 females) which displayed this pat-

tern of attachment, the distance between tracheal rings decreased slightly as the left lung was approached and the rings became less regular in their arrangement. There appeared to be a distinct left bronchus branching off the trachea and running parallel to and in close contact with the underlying trachea. However, when the left lung was viewed from its dorsal aspect, the amount of cartilage visible in the area of attachment was greatly reduced, and often none could be seen. The actual structure in this area was, therefore, an elevation of a portion of the trachea, which resembled the top of a tube sliced lengthwise and which joined the anterior border of the opening in the trachea to the ventral surface of the left lung. The dorsal surface of the left lung was in more direct contact with the posterior edge of the opening into the trachea. This type of left tracheopulmonary junction was found in eight of the snakes which did not, before staining, appear to have any bronchus, two which seemed to have a membranous attachment, and two which appeared to have a definite bronchus. Nine of the twelve snakes originally assigned to this category were confirmed really to have this pattern after staining.

The remaining 25 snakes fell into two categories—those in which the alveolar tissue of the lamella started immediately adjacent to the trachea and those in which there was a definite bronchus. In 13 snakes (11 males and 2 females) the left lung opened directly into a hole in the ventral aspect of the trachea, so that there was only minor disarrangement of the cartilaginous tracheal rings immediately adjacent to the opening (Fig. 7). The alveolar tissue of the left lung extended all the way to the trachea. In 12 snakes (8 males and 4 females) a distinct left bronchus joined the left lung to the trachea (Fig. 8). Again there was a minimal effect of this structure on the rather uniform arrangement of the cartilaginous tracheal rings. The cartilages of the bronchus did not appear to be con-

TABLE 2. COMPARISONS OF DIFFERENT MODES OF ATTACHMENT OF THE LEFT LUNG.

Mode of attachment (for more detailed descriptions see text)	Estimation from inspection of dissection			Observed in specimens stained with toluidine blue		
	males	females	total	males	females	total
Attached directly to trachea	16	7	23	11	2	13
Membranous, duct like attachment	8	0	8	0	0	0
Mode of attachment different dorsally and ventrally	3	9	12	11	10	21
Left bronchus present	9	2	11	8	4	12
Damaged and omitted from sample	1	0	1	7	2	9

tinuous with those of the trachea and usually showed a more irregular pattern. If there were rings, they were always thinner and much more closely spaced than those in the trachea. Often they were connected by longitudinal bars of cartilage. In several instances they formed complete rings around the circumference of the bronchus, unlike those of the trachea. In other cases there were no clear rings, but rather a networklike arrangement of cartilage. The length of the bronchus was also less than originally estimated, 0.8 mm being the maximum.

In no snake was there a purely membranous tube between the trachea and the left lung as was originally described in unstained specimens (Table 2).

Two further variations were noted in the structure of the respiratory tract in this area. In addition to its attachment to the trachea, the left lung appeared, in ten of the snakes (8 males and 2 females), to share common alveolar tissue with the right lung. This characteristic is correlated with the type of attachment of the left lung to the trachea. In half of the cases (4 males and 1 female) the left lung opened directly from the trachea, and in four instances (3 male and 1 female) there was a difference in the method of attachment dorsally and ventrally. In the tenth specimen (male) the attachment was

destroyed before staining so that definite confirmation of the structure was impossible. Before staining, however, the snake appeared to possess a very short bronchus.

The common alveolar tissue was always located posterior to, and usually to the left of, the tracheal attachment of the left lung and always involved the dorsal surface of the left lung. Such common tissue usually formed only a minor portion of the total alveolar tissue of the left lung, but was enough to make dissection of the left lung from the right lung difficult. When the two lungs were separated, the area where they had been fused was always evident on both lungs. In one case the left lung, on its ventral aspect, opened directly into the trachea via a tubelike elevation resembling half a bronchus cut lengthwise. Posterior to this on the dorsal aspect of the left lung, the left lung opened directly into the substance of the right lung rather than into the trachea, which by this point had also entered the substance of the right lung. Thus, posteriorly the left lung appeared to be a direct extension or lobe of the right lung.

Well-developed tracheal lungs were noted in only two specimens, one male and one female (Fig. 2). In both cases the respiratory tissue of the tracheal lung was

Figures 5–8. Dissections of the left lung of *Thamnophis* after staining with toluidine blue. Anterior is toward top of page. 5. Specimen in which dorsal and ventral sides of attachment vary. This ventral view shows a short, bronchus-like connection. (X 12). 6. Same specimen as in Fig. 5, with the left lung folded anteriorly to show the dorsal side of its attachment to the trachea. The attachment appears to be direct with no intervening bronchus. (X 12). 7. Specimen in which pulmonary tissue of left lung is directly adjacent to the trachea; there is no left bronchus. (X 7). 8. Specimen with a distinct left bronchus. (X 12).

Abbreviations: B, Left bronchus; L, left lung; R, right lung; T, trachea.

an anterior extension of the alveolar tissue of the right lung. This tissue was located in the area normally occupied by the membrane separating the ends of the discontinuous tracheal rings, i.e., on the dorsal aspect of the trachea. The tracheal lung tissue tapered out 7 mm (female) or 8 mm (male) anterior to the point at which the trachea entered the substance of the right lung. In the male the left lung opened directly into the ventral surface of the trachea; in the female there was a hint of bronchial structure on the ventral surface of the attachment of the left lung, but none dorsally.

DISCUSSION

Contrary to the studies of Cope (1874, 1900) on the same snake, *Thamnophis sirtalis parietalis*, we found cartilaginous rings in bronchus-like structures in most specimens. Cope, however, does not mention any staining procedure, and apparently simply inspected unstained dissections. The same is true of the observations by Beddard (1906), Brongersma (1957), and Butler (1985). We question the accuracy of the earlier reports on the presence or absence of cartilaginous bronchial rings in snakes, at least in small ones such as *Thamnophis*, since we were often unable to tell if such rings were present without toluidine blue staining (Table 2). Our material was almost certainly at least as well preserved for our purposes as was that used by earlier workers who studied museum specimens fixed in alcohol. Certainly, if the presence or absence of a bronchus is to be used as a taxonomic character, then staining for cartilages appears important and probably essential.

In *Thamnophis* there is considerable variation in the mode of attachment of the left lung. Whether such variation is common in other species is not known. Clearly, however, broad surveys like Brongersma's (1957), which are based on only one or two specimens of most

species, are open to considerable question. Series of specimens of each taxon considered must thus be studied before reliable conclusions can be reached.

Table 1 compares various measurements of the size and position of the lungs in male and female specimens. In every case the difference between the sexes is significant, in all but one case at the 0.01 level or better (all based on the *d* test; Bailey, 1959). The differences in the lengths and positions of the lungs between the sexes are not simply reflections of the difference in lengths of the specimens, as the left lungs of females are both relatively and absolutely smaller than those of males. Thus the sex of the specimens must be checked in any study comparing the size and position of the lungs of different species of snakes.

Another problem is that most previous workers have used some relative measure of the size of the left lung such as a percent of the total body length (Brongersma, 1957; Thorpe, 1975). Unfortunately, our data reveal no correlation between snout-vent length and length of the left lung (correlation coefficient for males = 0.108 and for females = 0.29; the coefficients for relative lengths of the left lung are no better—0.09 for males and 0.10 for females). Thus in our sample, the absolute, not the relative, size should be the more meaningful figure, but obviously it cannot be used in comparing the lungs of snakes of very different sizes.

The problems noted above do not mean that the structure of the lungs cannot be used as a taxonomic character. Some of the cases in which it has been used involve such major differences between taxa that intraspecific variation is probably not critical, e.g., the example of *Gonyosoma* discussed by Underwood (1967). In most cases, however, detailed studies of many specimens of each form are required. For example, Cope (1900) and Underwood (1967) reported the general or at least usual absence of a left lung in the Viperinae, while its presence

was noted by Butler (1895) in *Cerastes cornutus* and by Brongersma (cited by Underwood, 1967) in *Causus*. The character may well be variable in many members of the subfamily.

Our results do not mean that the lungs are highly variable in all snakes. Thorpe (1975) found no significant variation according to size, sex, or geographic origin in the length of the left lung of *Natrix natrix*. He used a different system of measurement, using scales instead of absolute measurement, but this should not change his results greatly. Different species of snakes, even of natricids, must show considerably different degrees of variability.

ACKNOWLEDGMENTS

We wish to thank M. A. Adair, M. N. Loucks, I. H. McCorquodale, and S. T. Richter for technical help and typing. The work was begun by a former student, J. E. Wilkinson, whose report was very useful. The photographs were taken by N. R. Hatton. C. S. Churcher and M. C. Parsons critically read the manuscript and made helpful suggestions as did two anonymous reviewers. The work was supported by Grant A1724 from the Natural Sciences and Engineering Research Council Canada.

LITERATURE CITED

- BAILEY, N. T. J. 1959. Statistical methods in biology. New York, John Wiley and Sons, x + 200 pp.
- BEDDARD, F. E. 1906. Contributions to the knowledge of the vascular and respiratory systems in the Ophidia, and to the anatomy of the genera *Boa* and *Corallus*. Proc. Zool. Soc. London, 1906: 499-532.
- BERGMAN, R. A. M. 1952. L'anatomie du genre *Ptyas* à Java, Riv. Biol. Coloniale, 12: 5-42.
- . 1958. The anatomy of the Acrochordinae. Proc. Koninklijke Ned. Akad. Wetensch., C, 61: 145-184.
- BRONGERSMA, L. D. 1957. Notes upon the trachea, the lungs, and the pulmonary artery in snakes. I-II. Proc. Koninklijke Ned. Akad. Wetensch., C, 60: 299-313.
- BUTLER, G. W. 1895. On the complete or partial suppression of the right lung in the Amphisbaenidae and of the left lung in snakes and snake-like lizards and amphibians. Proc. Zool. Soc. London, 1895: 691-712.
- COPE, E. D. 1894. On the lungs of the Ophidia. Proc. Amer. Phil. Soc., 33: 217-224.
- . 1900. The crocodilians, lizards, and snakes of North America. Rep. U.S. Nat. Mus., 1898: 153-1270.
- HILDEBRAND, M. 1968. Anatomical preparations. Berkeley and Los Angeles, Univ. California Press, viii + 100 pp.
- THORPE, R. S. 1975. Quantitative handling of characters useful in snake systematics with particular reference to intraspecific variation in the ringed snake *Natrix natrix* L. Biol. J. Linn. Soc., London, 7: 27-43.
- UNDERWOOD, G. 1967. A contribution to the classification of snakes. Brit. Mus. (Nat. Hist.), London, x + 179 pp.

On the Dorsal Armor of the Crocodilia

FRANKLIN D. ROSS¹

GREGORY C. MAYER²

ABSTRACT. A new method for counting the scales of the dorsal armor of crocodilians is introduced. Based on the one-to-one correspondence between vertebrae and transverse scale rows, it allows comparisons between species on the basis of homologous scale rows. Scale rows are counted away from the evolutionarily conservative sacro-caudal juncture in precaudal and caudal series, approaching the more variable anterior and posterior ends of the dorsal armor from a fixed central point. Primitively, crocodilian dorsal armor consists of 27 transverse precaudal rows of two scutes each, as is found in protosuchians. Changes occurring in the evolution of the modern crocodilians are 1) broadening of the trunk armor; 2) breakup of the ancestral scute pair into many elements per transverse row; 3) loss of the anteriormost transverse rows; and 4) narrowing the armor at the cervico-thoracic juncture. Descriptions of the dorsal armor and its variation in all living species are given. The most frequent kinds of variation in the living crocodilians, both among and within species, are the deletion of transverse rows and the compounding of transverse rows in the cervical region. *Gavialis* is the most primitive living species, departing least from the mesosuchian type of armor, while *Crocodylus acutus* and *C. porosus* are the most derived, showing great reduction from the primitive eusuchian pattern in both trunk and cervical regions.

INTRODUCTION

Nothing is more fugitive than the forms of crocodiles.
—Geoffroy, in Cuvier, 1831.

Our methodological and conceptual approach to the dorsal armor of the living Crocodilia differs from that of most previous authors (Cuvier, 1807; Duméril and Bibron, 1836; Boulenger, 1889;

Deraniyagala, 1939; Wermuth, 1953; Brazaitis, 1973), but follows directly from insights available in the morphological and paleontological literature (Huxley, 1859; Colbert and Mook, 1951; Nash, 1975). Since the latter part of the eighteenth century, crocodile taxonomists have described dorsal armor from the anterior end first. The traditional methodology involves counting scales (scutes) from occiput to tail tip in terms of postoccipitals, nuchals, dorsals and caudals. The difficulty with counting in this way is that the greatest and most interesting variation occurs in the form of missing, enlarged and fused transverse scale rows in the cervical region. This being the case, the most outstanding variation is encountered initially, resulting in unreliable and oftentimes false homologies. By not establishing the relationships among the sacrum, hind legs, and dorsal armor, various authors have counted the dorsals and caudals beginning or ending with any of several scale rows. We find it interesting and heartening, though, that our conclusions generally agree with those of the traditional method, and that we have little evidence to suggest that the current classification of the living Crocodilia is incorrect.

Our method considers the transverse scale rows on the body and tail in relation to the underlying vertebrae, with which there is primitively a one-to-one relationship. There is always an unarmored space immediately behind the head in living species, and there may be one or two more such spaces near the anterior

^{1,2}Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, U.S.A.

end of the series; but, invariably, there is a continuous series of transverse rows from the region of the shoulders, over the sacrum, and on to the tip of the tail. These transverse rows of bone and scale correspond one-to-one with immediately underlying vertebral elements. Huxley (1859) first observed the one-to-one correspondence in *Caiman crocodilus*, and explained that, "the anterior part of the inner surface of each of the two middle scutes is connected by a ligament with the extremity of the spinous process of a vertebra; at least this is the case in the dorsal, lumbar, sacral, and anterior caudal regions." Our counts are made from the evolutionarily conservative sacro-caudal juncture. We approach the more variable anterior and posterior ends of the dorsal armor from a fixed central point, which is the same for all living and fossil species.

Each transverse row is intervertebral in position, the median elements of a row being attached to the neural spine of the anterior vertebra, and extending posteriorly to overlap the one behind. The dorsal armor thus exhibits the primary segmentation which exists prior to resegmentation in the development of the vertebral column (Williams, 1959). Seidel (1979) observed that the blood supply to the transverse rows of dorsal armor on the body of *Alligator mississippiensis* is by means of a series of unpaired arteries, the dorsal median arteries, which emerge through the deep fascia between successive neural spines, and which also supply the neural spine of the anterior vertebra. All osteoderms in a transverse row and the neural spine of the vertebra anterior to the row receive blood from a common source. Transverse dorsal osteoderm rows are thus associated with the anterior of the two vertebrae they overlap.

Huxley (1859) restricted his observations to the scutes on the body and anterior part of the tail, and Seidel (1979) noted that an interruption in the series of dorsal median arteries occurred in the

shoulder region, although it continued anterior to this. Gadow (1901) also noted the difference between trunk and cervical armor in the Crocodilia, stating that, "the armor of the recent forms consists, so far as the large scutes are concerned, of a considerable number of scutes, which are arranged in transverse rows, each row corresponding with one skeletal segment of the trunk proper. Mostly there is a detached cluster of scutes on the neck." Indeed, there is sometimes a detached cluster of scutes on the neck, particularly in *Crocodylus*, but other forms (e.g., *Gavialis*, *Tomistoma*, *Melanosuchus*) possess continuous thoracic and cervical armor.

As illustrated in Figures 1 and 2, radiographs and dissection show the one-to-one relation between vertebrae and dorsal armor, and that in certain cases some scutes on the neck overlap two vertebrae. Dissection confirms Huxley's observation of association between neural spines and osteoderms. There are normally 26 precaudal vertebrae (24 presacral + 2 sacral) plus a proatlas in all living and fossil forms examined. For further discussion of vertebral morphology, see Higgins (1923), Hoffstetter and Gasc (1969), and Romer (1956).

MATERIALS AND METHODS

Our taxonomy of the living Crocodilia follows Wermuth and Mertens (1961) at the generic and specific level. We do not consider subspecies. We have followed Boulenger (1889) in ordering the genera as follows: *Gavialis*, *Tomistoma*, *Crocodylus*, *Osteolaemus*, and finally *Alligator* and the caimans.

Determinations were made on the basis of cranial characters, coloration, and geographic provenance. We examined skins, mounted, and fluid preserved specimens of all living species. The Appendix is a list of specimens examined and reported on in this paper. In addition, we incorporate some data from the

Figure 1. Tracings of radiographs showing endo- and exoskeletal systems in: A. *Crocodylus siamensis*, MCZ 3716, lateral view of the neck and trunk region. B. *Osteolaemus tetraspis*, MCZ 3589, dorsal view of the lumbar, sacral, and anteriormost caudal regions. C. *Osteolaemus tetraspis*, MCZ 2017, lateral view of the anterior thoracic and all of the cervical series.

literature; and, in the case of *Crocodylus intermedius*, we have relied in part on photographs and a personal communication regarding the type. The type of *Protosuchus richardsoni* (AMNH 3024) was also examined.

Specimens of *Osteolaemus tetraspis*, *Tomistoma schlegelii*, *Melanosuchus niger*, *Paleosuchus trigonatus*, *P. palpebrosus*, *Caiman crocodilus*, *C. latirostris*, *Alligator sinensis*, *A. mississippiensis*, and nine species of *Crocodylus* were radiographed to determine the relationship between the dorsal armor and the skeleton, and for vertebral counts. Additional vertebral counts were performed on skeletons.

METHODOLOGY

Although the tradition has been to count both dorsal armor and vertebrae from the cranium, we count away from the sacro-caudal juncture in both precaudal (PC) and caudal (C) series (Figs. 2, 4). PC 1, the posteriormost transverse precaudal scale row, hangs posteriorly from the tip of the neural spine of the posterior sacral vertebra. PC 1 can be identified unambiguously in fluid preserved specimens by a simple dissection. An incision is made along the medial, posterior, and lateral borders of three transverse rows on one side of the body: the two rows nearest the posterior edge of the iliac crest and the one anterior to them. The skin and osteoderms are then reflected anteriorly, and the musculature dorsal to the level of the sacral ribs removed, thus revealing the relationship between the vertebrae and the osteoderm rows. PC 1 is the transverse row extending posteriorly from the neural spine of the posterior sacral, and dorsal to the postzygapophysis of the same vertebra. The reflected skin and osteoderms may be folded back, leaving the specimen externally unchanged, and the musculature of the other side of the body intact for anatomical studies. Radiographs from dorsal view also allow

unambiguous identification of PC 1, provided the osteoderms are sufficiently opaque to X-rays (Fig. 1B).

Short of dissection or X-ray, the following criteria have been found useful in the identification of PC 1. First, PC 1 lies between the posterior blades of the ilia. The posterior edge of the iliac blades are overlain by PC 1 or the anterior portion of the first caudal row (C 1). The location of the blades can be determined by palpation. Secondly, when held perpendicular to the body, the long axis of the femurs passes through PC 2 or the border between PC 2 and PC 1. Third, in many species, PC 1 is the first transverse row to become broader after monotonically decreasing in breadth from mid-body towards the sacrum (see Figs. 2B, 3C). Finally, the posterior edge of PC 1 is often at the posterior edge of the hind limb (the traditional point of demarcation between dorsals and caudals), but this is not always the case. When these criteria are applied simultaneously, PC 1 can usually be identified unambiguously. Reference to radiographs and dissected specimens is always helpful. PC 1 has been identified by these criteria in most specimens examined in this study.

Skins and stuffed specimens present special problems. Determination of the relationship of the transverse rows to the variously granulated scales around the base of the hind limb in intact specimens, and criterion three above are the best guides to the identification of PC 1 in such specimens. Comparison with intact specimens is helpful.

Posterior to PC 1, the caudal series consists first of double crest caudals and then single crest caudals. Double crest caudals are associated with vertebrae with transverse processes, and have two to four keels across. Single crest caudals are associated with vertebrae having reduced or absent transverse processes, and have a single median keel. The keel of the single crest series and the outer keels of the posterior double crest series form a Y-shaped swimming keel.

Figure 2. Semidiagrammatic illustration of the one-to-one relationship between vertebrae and transverse rows of dorsal osteoderms as revealed by dissection of *Melanosuchus niger*, MCZ 17726. Odd numbered elements stippled in the pre-caudal (PC) series, and even numbered elements stippled in the caudal (C) series. A. PC 26-C 5 in lateral view. B. PC 26-C 2 in dorsal view. C. C 3 in cross section. D. C 4-C 15 in dorsal view. E. C 16-C 28 in lateral view. F. C 29-C 38 in lateral view.

Figure 3. Dorsal armor of three selected crocodilians. A. *Protosuchus richardsoni*, AMNH 3024, type, after Colbert and Mook, 1951 (PC 27-C 39 shown). B. *Gavialis gangeticus*, BMS no #, from the head to the sacrocaudal juncture; C: *Caiman crocodilus*, MCZ 3394, full body.

On the lateral and ventral surfaces of the base of the tail, scales may occur in various configurations which break up the regular (in the sense of showing plane symmetry; see Breder, 1947) disposition of scales in this area. These are collectively termed basicaudal irregularities. Four kinds are recognized: 1) a thin horizontal strip of granular skin separating the dorsal from the lateral scalation; 2) one or more thick strips of granular skin on the dorsolateral surface (Fig. 11); 3) lateral intrusions, in the form of small vertical arrays of scales, two-thirds or more of a normal scale in height, on the postcloacal lateral surface, sometimes present on one side only (Fig. 4; Ross and Ross, 1974: Fig. 1a); and 4) large ventrolateral intrusions composed of vertical arrays of scales on the postcloacal ventral and ventrolateral surfaces (Fig. 4; Ross and Ross, 1974: Fig. 1b). This last increases the number of midventral transverse rows relative to the dorsals.

Anterior to and including PC 1 is the continuous and contiguous precaudal armor (Fig. 4). The *continuous* armor is the series of transverse rows beginning with PC 1 and extending to the row which is bordered in front by a missing transverse row. Soft skin may intrude between two rows without a row being absent. This makes the armor appear discontinuous, although all rows are present. The *Crocodylus moreletii* in Fig. 4 has 16 continuous precaudal rows, PC 17 (and others) being absent. *Contiguous* scutes are scutes in a series crossing (or at least touching) the midline in which adjacent members of the series are in contact. Beginning with PC 1, the number of contiguous scutes per transverse row in Figure 4 is 4, 4, 4, 4, 6, 4, 4, 4, 6, 3, 3, 6, 5, 2, 4, 2.

Additional characteristics of the trunk armor are the presence of detached lateral scutes, sometimes forming a longitudinal row (Fig. 4), the alignment and extent of development of keels on the scutes, the uniformity of scute size, and the presence of left/right asymmetries.

Note that even development and alignment of keels, uniformity of scute size, and absence of asymmetry impart the quality of "regularity" to the armor.

The cervical shield is the cluster of scutes on the neck. The term does not refer to particular precaudal rows, as the rows comprising it and its continuity with the thoracic armor vary among species. The nape region is the area between the occiput and the cervical shield. Vertebral correspondence in the cervical region is determined in two ways. In species with a continuous thoracic and cervical armor, correspondence can be determined directly by counting, with observation of individual variation revealing compound rows. In species with discontinuous cervical and thoracic armor, comparison with related species is necessary. For example, in *Crocodylus* all species have two enlarged rows in the cervical shield. In *C. cataphractus* (Fig. 5D), and frequently in *C. johnstoni* (Fig. 6A), the cervical and thoracic armor is continuous, the last row before the two enlarged rows being PC 19. Thus the enlarged rows begin with PC 20. Individual variation in *C. niloticus* (Fig. 6B-D) shows that these enlarged rows are each compound. Assuming that these rows are homologous among members of the genus allows extrapolation to other species. Thus the *C. moreletii* in Fig. 4 has two scutes in PC 20 + 21 and four in PC 22 + 23. Uncertain homologies might be resolved by careful dissection of arteries and nerves.

Analysis of the cervical region requires comparison of many specimens simultaneously. We found photographs useful in comparing specimens from different institutions. A marker with specimen number was placed on PC 12, and both marker and PC 13 forward photographed from directly above. Properly marked photographs proved to be as good as actual specimens for comparative purposes. We would encourage the collection of photographs of this kind, as it allows obtaining sizable samples without damaging populations of the world's

Figure 4. Some terms used in this paper indicated on *Crocodylus moreletii*, USNM 71955, topotype. From an original drawing by Aleta Karstad.

dwindling supply of crocodilians. The photographs can be stored as a record of the specimens examined in natural populations without removing individuals from those populations.

Meristic characters in reptiles are generally believed not to change in postembryonic life (Hecht, 1952; Kerfoot, 1970; Fox, 1975). Our observations support this. Although we have not studied large series, examination of embryos of various ages from a variety of species indicates that there is little or no ontogenetic change in embryonic life as well. The scutes of unpigmented embryos are somewhat more closely spaced, and thus may appear contiguous or continuous, whereas in hatchlings and adults they would not be in contact. Thus available ontogenetic data does not help to elucidate homology of transverse rows. Medem (1958a) noted that *Crocodylus intermedius* embryos did not differ from adults in scale counts.

There are two exceptions to the general rule of no ontogenetic change. First, the number of single crest caudals decreases due to injury and wear on the tail. Counts of this character are thus possible only on undamaged individuals. Second, and more interesting, the number of scutes per transverse row in *Gavialis* decreases from six to four as an animal ages. This is discussed in the species accounts. Deraniyagala (1939) also noted some ontogenetic changes in *Crocodylus porosus* embryos, involving a decrease in the number of small scales interspersed among the scutes of the trunk armor.

SPECIES ACCOUNTS

The following accounts of the species provide brief descriptions of the dorsal armor of each species, with more detailed treatment of the anterior region, where the greatest variation within and among species occurs. Table 1 gives counts of the contiguous scutes per transverse row for PC 1 to PC 18 for all species. In Table

2, the number of double and single crest caudals is given for each species. All living crocodilians have a Y-shaped caudal swimming keel, consisting of the single crest caudals and the posterior part of the double crest series. The ways in which the Y-shaped keel is formed from the keel rows of the trunk armor as it passes over the pelvis in different taxa are given in Table 3. Table 4 records the occurrence of basicaudal irregularities.

Gavialis gangeticus

In *Gavialis* (Fig. 3B), the continuous and contiguous precaudal armor is remarkably uniform in scute size and shape, and keel row alignment and development. There are six scutes per transverse row in PC 1 to PC 18 in hatchlings and young. In older specimens, there are only four scutes across, apparently because the lateral D-shaped scales of the very young fail to ossify with the four more medial scutes. This is unique in the Crocodilia. There are no detached lateral scutes on the flanks, and the thoracic and cervical armor are continuous. There are 22 continuous rows of precaudal armor, narrowing anteriorly to two scutes in PC 22. The nape is naked except for two prominent scutes and occasional traces of others.

Tomistoma schlegelii

In *Tomistoma* (Fig. 5A-C), the continuous and contiguous precaudal armor is quite regular, with five to eight, usually six, scutes across at midbody (PC 9-PC 12). Detached lateral elements form an additional row on each flank. The thoracic and cervical armor are continuous forward to PC 23, and narrowest at PC 19 or PC 20. There are two scutes in PC 18, 19, 20, and 21. Typically, the terminal row is PC 22+23 compound with two elements. In some specimens PC 23 is present as a distinct row or asymmetrically. Fig. 5A-C shows the range of variation in this region. Scutes on the nape are ir-

Figure 5. Anterior end of the dorsal armor, PC 13 forward, with selected scale rows numbered. A. *Tomistoma schlegelii* FMNH 206755. B. *T. schlegelii*, USNM 145587. C. *T. schlegelii*, USNM 58136; D. *Crocodylus cataphractus*, MCZ 54650.

regularly placed, although sometimes appearing as three transverse rows, presumably PC 24 to PC 26. This latter condition is clearly seen in Figure 5A.

Crocodylus cataphractus

In *Crocodylus cataphractus* (Fig. 5D), the continuous and contiguous precaudal armor is regular in scute size and keel row alignment. There are four to eight, usually six, scutes across at midbody. The median scute pair of each row tends to be broader than the laterals and have lower keels. Detached lateral elements are present and sometimes form an additional keeled row on each side. PC 19 is bordered front and back by short spaces of soft skin. PC 20 + 21 and PC 22 + 23 are compound rows of two elements each. The compound nature of these rows is demonstrated in other species of *Crocodylus*. The nape rows are variably developed, but for the most part are

vestigial. The largest elements appear to be from PC 26.

Crocodylus johnstoni

In *Crocodylus johnstoni* (Fig. 6A), the continuous and contiguous precaudal armor is regular and has six to seven, usually six, scutes across at midbody. Detached lateral scutes form one or two additional well-developed keeled rows on each flank. The thoracic and cervical armor are continuous, but small spaces of soft skin occur between the transverse rows at the cervico-thoracic juncture. Sometimes there is but a single scute in PC 18 or PC 19 (Fig. 6A). PC 20+21 has two large scutes, and PC 22+23 has four, for a total of six fairly closely juxtaposed scutes forming an ovate shield. The two lateral elements of PC 22+23 are often displaced posteriorly, their medial edge bordering the juncture between PC 20+21 and PC 22+23. The nape has

Figure 6. Anterior end of the dorsal armor, PC 13 forward, with selected precaudal scale rows numbered on: A. *Crocodylus johnstoni*, MCZ 35006. B. *C. niloticus*, USNM 195783. C. *C. niloticus*, MCZ 12552. D. *C. niloticus*, USNM 63592.

three to six scutes in PC 26. PC 25 appears lost or perhaps coalesced with PC 26. PC 24 has zero to four minor elements.

Crocodylus niloticus

In *Crocodylus niloticus* (Fig. 6B-D) the continuous and contiguous precaudal armor is moderately regular, particularly in the sacral and lumbar regions and along the midline. There are 16 or 17 continuous rows of precaudal armor, with four to nine, usually six, scutes per transverse row at midbody. The median scute pair of the anteriormost thoracic row is often enlarged. Laterally, detached elements may form up to two additional keeled rows. In some individuals, especially from Madagascar, a row of two scutes intervenes between the thoracic and cervical armor. We interpret this as PC 19 on the basis of it being closer to the cervical than the thoracic armor, but this identification is not certain. PC 20+21 has two scutes, and PC 22+23 has four. These two transverse compounds tend to be closely adjacent, forming an ovate cervical shield. The compound nature of

these rows is demonstrated by occasional variants exhibiting one or the other row in an uncompound state (Fig. 6B, D). The nape has three to six scutes in PC 26, and zero to four small elements in PC 24. PC 25 is absent, or perhaps fused with PC 26.

Crocodylus palustris

In *Crocodylus palustris* (Fig. 7D), the continuous and contiguous precaudal armor is fairly regular. At the anterior end of the thoracic armor, small scales sometimes occur between transverse rows and between the scutes within a transverse row, leading to a loss of contiguity. There are four to eight contiguous scutes per transverse row at midbody, with no outstanding modal number. Detached lateral scutes form one or two additional keeled rows, the more medial particularly well developed. The thoracic and cervical armor are distinctly separated by a space of granular skin. PC 17 has zero to four scutes, and vestiges of PC 18 are rarely present. PC 19 is absent. PC 20+21 has two scutes, and PC 22+23 has four. Sometimes an additional pair of scutes is in contact laterally, giving a total

Figure 7. Anterior end of the dorsal armor, PC 13 forward, with selected precaudal scale rows numbered on: A. *Crocodylus porosus*, USNM 72732. B. *C. novaeguineae*, from New Guinea, CAS 119186. C. *C. novaeguineae*, from the Philippines, FMNH 52362. D. *C. palustris*, FMNH 63739.

of eight scutes in the cervical shield. Traces of PC 24 are sometimes evident. PC 25 is absent or perhaps combined with the normally four scutes in PC 26.

Crocodylus siamensis

In *Crocodylus siamensis* (Fig. 8A), the continuous and contiguous precaudal armor is regular, though often with small triangular scales interspersed between the scutes along the posterior border of a transverse row, as is also found in *C. porosus*. There are four to seven, usually six, scutes across at midbody. Detached scutes form one or two additional rows on the flanks. PC 18 is usually absent, and PC 19 always is. There is a cervical shield of two scutes in PC 20+21 and four scutes in PC 22+23, and sometimes remnants of PC 24. PC 25 is absent or perhaps combined with the two to usually four scutes in PC 26. *C. siamensis* is reported to hybridize with *C. porosus* in captivity, producing phenotypically intermediate and allegedly heterotic off-

spring (Youngprapakorn, 1976). The fertility of the hybrids is not reported.

Crocodylus porosus

In *Crocodylus porosus* (Fig. 7A), the continuous and contiguous precaudal armor is regular in keel row alignment and with six to eight, usually six, narrow elements per transverse row at midbody. Small triangular scales occur interspersed between scutes along the posterior edges of transverse rows. Anteriorly, these small elements also occur along the anterior borders of scute rows, and may contact the posterior triangles, breaking contiguity. Lateral detached scutes are quite small and indistinct. The thoracic and cervical armor are completely discontinuous. PC 17 is usually absent. PC 18 is sometimes represented by vestiges. PC 19 is absent entirely. PC 20 + 21 has two scutes, and PC 22 + 23 has four. The nape is essentially naked. In specimens examined by us PC 24 to PC 26 are usually absent, though rarely two distinct

Figure 8. Anterior end of the dorsal armor, PC 13 forward, with selected precaudal scale rows numbered on: A. *Crocodylus siamensis*, MCZ 3716. B. *C. rhombifer*, MCZ 12081. C. *C. intermedius*, FMNH 75658. D. *C. acutus*, FMNH 23147. E. *C. acutus*, MCZ 10920.

scutes may occur in PC 26. Deraniyagala (1939) reports specimens with up to four.

Crocodylus novaeguineae

In *Crocodylus novaeguineae* (Fig. 7B, C), the continuous and contiguous pre-caudal armor is regular in scute dimensions and keel row alignment. As in *C. porosus* and *C. siamensis*, interscute triangles occur. There are 16 or 17 continuous precaudal rows, with seven to twelve, usually eight, contiguous scutes at midbody. Detached flank scutes, though reduced, sometimes form an additional keeled row on each side. The thoracic and cervical armors are separated by spaces of skin. PC 18 is represented by vestiges or elements separated on the midline. PC 19 is absent, or, in a single specimen, rudimentary. PC 20+21 has two scutes, and PC 22+23 has two large median elements, and may have one or two smaller elements on either side. In many individuals, smooth or fine granular skin separates the two cervical rows (PC 20+21 and PC 22+23) or the left and right halves of one or both (Fig. 7B). Sometimes PC 22 and PC 23 are not compound. All or any combination of cervical scutes may be noncontiguous. There are

four to six scutes in PC 26. PC 24 and PC 25 are sometimes not evident, but are usually present as bluntly keeled scales.

Previous authors have differed on whether the freshwater crocodiles of the Philippines should be recognized as a species (*mindorensis*) distinct from *C. novaeguineae*, or merely as a subspecies of the latter (Wermuth, 1953; Schmidt, 1956; Wermuth and Mertens, 1961). Wermuth and Mertens, in their most recent checklist (1977), have elevated *mindorensis* to full species status. Tables 1 and 2 include with *C. novaeguineae* data for a few *mindorensis*. The preceding account of the cervical armor, however, applies only to *C. novaeguineae* proper. *C. mindorensis* (Fig. 7C) differs from *C. novaeguineae* in that PC 19 is better developed, being present as one or two elements in three of five individuals; PC 22+23 has four elements; the scutes of the cervical shield are little or not at all separated by skin; and the nape scutes between the cervical shield and the prominent nape row (PC 26) are better developed, sometimes in two distinct rows. The extremal count of twelve scutes at midbody is from an individual from the Philippines. In general, *mindorensis* has a more well-developed

armor than *novaeguineae*, though this conclusion is based on a small sample and should be considered tentative.

Crocodylus rhombifer

In *Crocodylus rhombifer* (Fig. 8B), the continuous and contiguous precaudal armor is often remarkably even and regular. There are six or seven, usually six, scutes across at midbody, and detached lateral elements form one or two additional keeled rows. Some individuals have continuous cervical and thoracic armor, others do not. PC 18 has one or two scutes. Sometimes there is a space of soft skin between PC 18 and PC 19, and invariably between PC 19 and PC 20+21. PC 19 is variously absent or present as two scutes. PC 20+21 has two scutes and PC 22+23 has four. Anterior to the cervical shield, scutes are poorly developed, except for a row of four prominent scutes, presumably PC 26. *C. rhombifer* is easily distinguished from *C. acutus* by having six or more scutes in all four transverse rows at midbody. The two species are reported to hybridize in the wild and in semi-captivity (Varona, 1966; Anonymous, 1969). A presumptive wild hybrid, AMNH 82943, has the black and white flecked coloration of *C. rhombifer*, and regularity of the dorsal armor like *rhombifer*, but has fewer contiguous scutes at midbody than *rhombifer*, is lacking PC 17 entirely, and has a head with *C. acutus* proportions and snout character.

Crocodylus moreletii

In *Crocodylus moreletii* (Fig. 4), the continuous and contiguous precaudal armor is fairly irregular and asymmetric, with the contiguous scutes at midbody often reduced to three or four. There are two to six, usually four, contiguous scutes across at midbody, and detached elements on the flank are always present. PC 17 and PC 18 are absent, and PC 19 has zero to

two scutes. PC 20+21 has two scutes, and PC 22+23 usually has four. PC 24 is variably developed, sometimes absent. PC 25 is absent or perhaps compounded with the four to six scutes in PC 26.

Natural hybrids have been reported between *Paleosuchus trigonatus* and *P. palpebrosus* (Medem, 1970) and *Crocodylus rhombifer* and *C. acutus*. In both cases, the presumed hybrids closely resembled one of the parent species, though possessing certain minor, yet characteristic, traits of the other parent. It is probable that hybridization has also occurred between *C. acutus* and *C. moreletii*. Ross and Ross (1974) used a combination of several cranial characters and body color to identify *C. acutus* from throughout its range, and found traces of a *C. moreletii* character (lateral intrusions on the basicaudal surface; see Fig. 4) on *C. acutus* only where they are sympatric with *C. moreletii* (Belize through Chiapas), or where the population could have been influenced by hybrid stock (west coast of Mexico; it is interesting that fossil *C. moreletii* have been reported from Baja [Miller, 1980]). The dorsal armor of presumptive hybrids resembles both parent species in scale counts, but in this case both parent species have remarkably similar dorsal armor.

Crocodylus acutus

In *Crocodylus acutus* (Fig. 8D, E), the continuous and contiguous precaudal armor is highly irregular in scute size, scute shape, and disposition. The median scute pairs of each row are fairly even in length, except anteriorly, where midline irregularities and gross asymmetry may occur. There are two to six, usually four, scutes across at midbody, always with asymmetry in at least some rows. There are always detached scutes on the flank. There are 14 to 17, usually 16, continuous precaudal rows. There is a tendency for the anteriormost median scute pair of the continuous armor to be large and flanked

by noncontiguous elements. PC 18 is always absent. PC 19 is almost always absent. PC 20 – 21 has two elements, and PC 22 + 23 has four, forming a cervical shield of six elements, though variations by deletion are common. PC 24 is vestigial, or more often absent, and PC 25 is absent or perhaps fused with the two to four elements of PC 26.

Crocodylus intermedius

In *Crocodylus intermedius* (Fig. 8C), the continuous and contiguous precaudal armor is more or less regular in scute dimensions and keel row alignment. There are five or six contiguous scutes per transverse row at midbody, with a variable development of detached flank scutes sometimes forming an additional keeled row. There are 16 continuous precaudal rows. One of our specimens has a “normal” *Crocodylus* cervical shield and nape region: PC 20+21 with two elements, PC 22+23 with four elements, and PC 26 with four. Our other specimen has three transverse rows in the cervical shield, with two scutes in each of the two posteriormost, and four in the anteriormost. The nape is “normal” with five scutes. Our sample of *C. intermedius* is small, consisting of a single museum specimen, and photographs of a living individual in the collection of F. Medem. C. A. Ross (personal communication) describes the type (MNHN Paris 7512) as having heavy thoracic armor of six, sometimes even seven contiguous scutes across. There are no detached lateral scutes. The neck has six scutes in a shield, and there are five scutes in the nape row.

Osteolaemus tetraspis

In *Osteolaemus tetraspis* (Fig. 9A), the continuous and contiguous precaudal armor is more or less regular, the median scute pairs the broadest, and with keels poorly developed or lacking. There are

four to eight, usually six, scutes per transverse row at midbody, and detached scutes on the flanks. PC 18 usually consists of two elements contiguous on the midline, often with detached lateral elements with very small, vestigial scales between them. Anterior to PC 18 there is a fold of skin, a row of two small elements, another fold of skin, and then two rows of two large elements each. The anteriormost of these large rows is usually the longest of the two. In a single specimen (Fig. 9A), only one row of two small elements occurs between PC 17 and the two large rows of the cervical shield. There is usually one well-developed row of four scutes on the nape. Some specimens show an additional pair of scutes. Radiographs (Fig. 1C) show that the large neck scutes overlies vertebrae PC 21 and PC 22, and PC 23 and PC 24. However, vertebral correspondences of all cervical scutes and the nape are uncertain.

Alligator sinensis

In *Alligator sinensis* (Fig. 9C), the continuous and contiguous precaudal armor has six to eight, usually six, scutes at midbody, and few or no lateral detached scutes on the flanks. The thoracic and cervical armors are separated by at least a small space of skin. PC 17 is sometimes absent, but usually present with up to five elements. PC 18 is usually absent. Anterior to this there is a strip of skin followed by three transverse rows of two scutes each. The size of these scutes increases anteriorly. The nape has a row of four to six prominent scutes, and sometimes an additional row of two between the prominent row and the cervical shield. The vertebral correspondences of the cervical shield and the nape are uncertain due to the discontinuity at the cervico-thoracic juncture and the relatively slight variation among individuals. The prominent nape row is likely PC 25, as it is in *A. mississippiensis*, leaving PC

Figure 9. Anterior end of the dorsal armor, with selected pre-caudal scale rows numbered, PC 13 forward on: A. *Osteolaemus tetraspis*, USNM 193289. B. *Alligator mississippiensis*, MCZ 17723. C. *A. sinensis*, FMNH 38234. PC 17 forward on: D. *Caiman crocodilus*, MCZ 15334. PC 13 forward on: E. *C. latirostris*, MCZ 3588.

19 to PC 24 to be accounted for by the three rows of the cervical shield and the smaller nape row. The possibility of compounding is indicated by the large size of the anteriormost of the three rows of the cervical shield.

Alligator mississippiensis

In *Alligator mississippiensis* (Fig. 9B), the continuous and contiguous pre-caudal armor is five to eight, usually eight, scutes across at midbody, and there are no detached scutes on the flanks. The thoracic and cervical armors are barely continuous, having reduced elements in PC 18 to PC 19, and possibly PC 20, and two folds of skin in the cervico-thoracic juncture. PC 18 is ordinarily present as two small scutes. There are ordinarily lateral scutes which may be from PC 18 or traces of PC 19 (we don't know which). If PC 19 is absent on the midline, there are two scutes in PC 20, two very long elements in PC 21+22, two in PC 23, and then the nape scales. If the posteriormost neck scute is PC 19, PC 20+21 has two very long scutes, and PC 22+23 has two shorter scutes, and then there are the

nape scutes in two or three rows separated on the midline. The most prominent nape row is PC 25, ordinarily with two, but with up to six ossified elements. PC 25 is the postoccipital of Ross and Roberts (1979); in *Crocodylus* this term refers to PC 26.

Caiman latirostris

In *Caiman latirostris* (Fig. 9E) the continuous and contiguous pre-caudal armor varies from regular to somewhat irregular in the alignment of the keels. There are six to nine, usually six, scutes across at midbody, with detached lateral scutes tending to form additional rows on the flanks. The thoracic and cervical armor are usually discontinuous. PC 18 has two to four elements. PC 19 is normally absent, but may be present in rudimentary fashion (Fig. 9E). There are three transverse rows anterior to PC 19 (or PC 18 when PC 19 is absent) in the cervical shield, with two scutes in the two posteriormost, and four in the anteriormost row. The vertebral correspondences of these rows are uncertain, but are probably PC 20, PC 21, and PC 22+23 com-

pound. In a single specimen (MCZ 17716), PC 24 is well developed and continuous with the cervical shield. The nape consists usually of PC 24 to PC 26, with PC 26 most strongly developed.

Caiman crocodilus

In *Caiman crocodilus* (Figs. 3C, 9D) the continuous and contiguous precaudal armor tends to have low keels in longitudinal alignment, and the detached scutes on the flanks tend to be aligned in additional rows. There are six to twelve, usually eight, scutes across at midbody. The thoracic and cervical armors are sometimes not truly continuous, though they appear so with only narrow strips of skin where PC 18 or PC 19 are lost. There are usually four scutes across somewhere in the PC 21 to PC 23 region, often in PC 23. PC 24 is variously absent, vestigial, or present and rarely (Fig. 3C) continuous with the cervical shield and contiguous across the midline. The nape appears to be PC 24 to PC 26, sometimes PC 25 to PC 26 only. Some specimens exhibit 26 transverse precaudal rows without loss or compounding, but about half of the specimens examined show left-right asymmetry and fewer than 26 transverse precaudal rows. In these specimens it is difficult to determine which rows have been lost or fused. Figure 9D is a probable interpretation.

Melanosuchus niger

In *Melanosuchus niger* (Fig. 2) the continuous and contiguous precaudal armor is regular, with nine to twelve, usually ten scutes across at midbody, and there are no detached lateral scutes. There are four to six scutes in PC 18, two to four in PC 19, two to four in PC 20, two to four in PC 21, and four in PC 22 and PC 23. Remnants of PC 24 to PC 26 in approximately three transverse rows give the nape an overall granular and studded appearance. There are invariably 23 continuous precaudal rows in our sample.

Paleosuchus palpebrosus

In *Paleosuchus palpebrosus* (Fig. 10D) the thoracic armor is broad with more or less regular keel row alignment and scute dimensions. There are four scutes in PC 2 to PC 5, and six to nine, usually eight, contiguous scutes at midbody. The pre-caudal armor is continuous to PC 18. PC 18 has three to six elements. PC 19 is often bordered anteriorly and posteriorly by soft skin, and is reduced in size. There are two contiguous elements in PC 19, PC 20, PC 23, and PC 24. PC 21 and PC 22 may have four or, usually, three elements. PC 24 may be continuous with or separated from the cervical shield. PC 25, as here interpreted, consists of lateral elements and, in one specimen (Fig. 10D), a single element adjacent to the midline. The most prominent row on the nape is PC 26 with four to six elements. *P. palpebrosus* is much more ossified than its congener. This is seen in the greater breadth of its dorsal armor, the continuity between the thoracic and cervical armor, and the greater ossification of the skull and the ventral armor (Medem, 1958b).

Paleosuchus trigonatus

In *Paleosuchus trigonatus* (Fig. 10A-C) the dorsal armor is four to eight, usually six, scutes wide at midbody and often highly irregular in keel row alignment and scute size and shape. These irregular elements form a broad carapace with some detached scutes on the flanks. Some *P. trigonatus* lose the median pair of scutes in PC 2 to PC 5 in various combinations. This character is unique among the living Crocodilia. Our interpretation of *P. trigonatus* is based on the fact that it is less ossified than *P. palpebrosus*. In *P. palpebrosus* PC 18 usually has four elements, while in *P. trigonatus* there are usually two. PC 19 is reduced and has two elements in *P. palpebrosus*. In *P. trigonatus* a single transverse row of the cervical shield is lacking; and, on the

Figure 10. Anterior end of the dorsal armor, PC 13 forward, with selected precaudal scale rows numbered on: A. *Paleosuchus trigonatus*, FMNH 69878. B. *P. trigonatus*, FMNH 69882. C. *P. trigonatus*, FMNH 69877. D. *P. palpebrosus*, FMNH 134989.

basis of its small size in *P. palpebrosus*, we believe the missing row to be PC 19. PC 20 to PC 23 normally have two elements. In some individuals PC 21 and PC 22 appear to be compound. The nape has one well-developed row, PC 26, and small scales lie between PC 26 and PC 23, arranged in one or more additional rows. The larger of these latter small rows is probably PC 24, because this row is better developed in *P. palpebrosus*. Some *P. trigonatus* and *P. palpebrosus* combine median scute pairs in the neck to form median unpaired single or double keeled elements. This again is unique among the living Crocodylia.

DISCUSSION

THE FOSSIL RECORD

The presence of middorsal armor is widespread in the early archosaurs and their descendants, although it is lacking or poorly developed in the most primitive thecodonts, the proterosuchians (Reig, 1970; Romer 1972). It is most highly developed in parasuchians (phyto-

saur), aetosaurs, certain ornithischians, and, of course, crocodilians. In some of these, such as parasuchians (Chatterjee, 1978) and *Stagonolepis* (Walker, 1961) there is also a one-to-one correspondence between vertebrae and transverse scute rows. The wide occurrence of dorsal armor suggests that it is primitive for archosaurs; its general absence in proterosuchians, however, urges caution in accepting this hypothesis.

Fossil crocodilians with dorsal armor in place are rare, but several good examples exist. The earliest known true crocodiles, *Protosuchus* (Colbert and Mook, 1951), from the Upper Triassic or earliest Jurassic, and *Orthosuchus* (Nash, 1975), from the Upper Triassic, have essentially identical dorsal armor. In the neck and trunk region paired rectangular scutes occur in one-to-one correspondence with the vertebrae. At their lateral edges the scutes are sharply downturned to form a lateral face. Behind the occiput an extra small scute or scute pair (corresponding to the proatlas?) occurs. In the more completely known *Protosuchus* this armor extends onto the tail to its tip. The armor forms a long, essentially parallel

sided, rigid bony adjunct to the vertebral column (Fig. 3A). The armor of another early crocodilian, *Stegomosuchus*, seems to be much the same (Walker, 1968). Protosuchian dorsal armor did not form a carapace, though the scutes did enclose the epaxial musculature, and as such were protective.

Romer (1974) derived crocodilians from creatures of amphibious habits because strong hind limbs and tails function well in aquatic environments. Nash (1975) followed Romer, supposing that *Orthosuchus* spent much of its time in the water. However, the completeness of the dorsal armor of protosuchian crocodiles would have prohibited tossing the head upwards and backwards for swallowing prey while keeping the gullet above water in modern crocodilian fashion. Protosuchians could not feed in deep water, and also lacked a compressed tail with a median keel (see Manter, 1940, on the importance of the tail in swimming). We suggest that the group was terrestrial.

The transition from protosuchian to mesosuchian crocodiles involved reduction or loss of the anteriormost transverse scute rows and lateral compression of the posterior half of the tail to a vertical keel. The development of a caudal keel and loss of the anteriormost transverse rows (thus gaining the ability to toss the head backwards to swallow) indicate aquatic habits.

Some mesosuchians retained a parallel sided armor much like that of protosuchians (atoposaurs, Wellnhofer, 1971), while others broadened the scutes in the trunk region (e.g., *Teleosaurus*, Kálin, 1955). Many mesosuchians had peg and socket articulations on the lateral edges of their scutes. This is an elaboration of peglike processes seen in *Protosuchus* and *Orthosuchus*. Most mesosuchians still had only a single pair of scutes per transverse row (Romer, 1956). The pelagic marine thalattosuchians lacked armor altogether (Troxell, 1925). (It is interesting that the most reduced dorsal

armor in modern crocodilians occurs in the two most marine species, *Crocodylus porosus* and *C. acutus*.)

The transition from mesosuchian to eusuchian crocodiles involved further caudal compression, and, in some cases, further loss of anterior transverse scute rows. Eusuchians broadened and fractured the ancestral scute pairs into paired sets of two to six elements each (Fig. 3B, C), lost the lateral peg and socket articulations and consequent rigidity, and covered the trunk and base of the tail with a pavement of bony plates forming a defensive carapace. Interruption or narrowing of the armor in the shoulder region is also typical of eusuchians (Fig. 3C, 4). In general, eusuchian armor is broader and more flexible than that of their mesosuchian predecessors, with a tendency towards interruption of the dorsal armor in the shoulder region. *Procaimanoidea* (*Hassiacosuchus*) (Mook, 1941), *Leidyosuchus* (Erickson, 1976), and *Bernissartia* (Kálin, 1955)¹ are fossil eusuchians whose armor is known, and all are of essentially modern form. Among the living Eusuchia, the dorsal armor of *Gavialis* is most reminiscent of mesosuchian armor.

VARIATION AND EVOLUTION

Primitively, crocodilian dorsal armor consists of 27 transverse precaudal rows of two scutes each, as is found in protosuchians. The preceding review of historical changes in the primitive armor, through mesosuchian and eusuchian stages, allows us to make some general statements on the direction of evolution of some characters. Advanced characteristics include loss of rows, a greater number of scutes per row, broadening of the trunk armor, and narrowing of the armor

¹Buffetaut (1975) has questioned whether *Bernissartia* is a eusuchian on the basis of palatal and vertebral structure; its armor, however, is unquestionably advanced in character.

at the cervico-thoracic juncture. With this in mind, we can look at evolutionary steps and directions in the living taxa. The possibility of reversal cannot be ignored; for example, the low number of scutes per row in *Crocodylus acutus* is probably a secondary loss.

The variation in dorsal armor among the living crocodilians shows that a variety of evolutionary changes have occurred within the group. The two kinds of changes which account for most of this variation are deletion of rows, and the compounding of rows in the cervical region.

Deletion of at least the anteriormost precaudal row (PC 27) occurs in all modern taxa. In most taxa more are missing. In the genus *Crocodylus* a series of stages (not necessarily a phyletic series) in the reduction of armor at the cervico-thoracic juncture can be seen. *C. cataphractus* has continuous precaudal armor through PC 22+23 (Fig. 5D). In *C. johnstoni*, PC 18 and PC 19 may be reduced to a single scute (Fig. 6A). In *C. niloticus* reduction continues, with one or both of PC 18 and PC 19 entirely absent, leaving the precaudal armor continuous to PC 17 and the cervical and thoracic armor discontinuous (Fig. 6B-D). Further rows may be deleted, so that in *C. acutus*, for example, some individuals have continuous armor only to PC 14. Deletion may also occur in the cervical shield, and the presence of rows usually absent also occasionally occurs as an individual variation.

The compounding of rows, which entails a loss of one-to-one correspondence of vertebrae and transverse rows, is clearly derived with respect to separate rows being present. Compounding is indicated by the occurrence of individuals showing asymmetrical fusion. This is clearly seen in the series of *Tomistoma* in Fig. 5: 5A has the enlarged compound, 5B is compound only on the left, and 5C has both PC 22 and PC 23 entirely separate. Though greatly enlarged scutes, and scutes which overlie two vertebrae suggest compounding, the

recognition of compounding depends on individual variation. Some species may be phenotypically fixed for compounds, which would make the compound row difficult to detect.

Overall development of the dorsal armor is reflected in the number of rows present and the number of scutes per row. There is considerable variation among species in the extent to which the armor is developed. In the genus *Paleosuchus*, for example, *palpebrosus*, with more well-developed armor, has 24 continuous precaudal rows and a mode of eight scutes per row at midbody, while the less armored *trigonatus* lacks PC 19, and has modally six scutes per row. Similarly, *Crocodylus rhombifer* has at least 17 continuous precaudal rows and six scutes per row, while *C. acutus* usually has 16 or fewer continuous precaudal rows, and four scutes per row. Overall reduction of armor is sometimes accompanied by marked left/right asymmetries, as in *C. acutus* and *C. moreletii*.

The variation within species is of the same sort as that which distinguishes species. Thus, species differ in number of transverse rows, number of scutes per transverse row, and the presence of compounds, and all of these characters vary within species. Many individual variations are reversions to a more primitive condition, such as the breakup of compounds in *Crocodylus niloticus*, or the presence of PC 17 in some *C. acutus*. Other variations are derived and characteristic of more advanced species, such as the occasional reduction of PC 18 and PC 19 in the relatively primitive *C. johnstoni*. Some individuals show phenotypes more derived than those characteristic of even the most advanced species, such as a *C. niloticus* with a very reduced cervical shield, or a *C. acutus* with only 14 continuous precaudal rows.

ACKNOWLEDGMENTS

About fifteen years ago, one of us (FDR) brought a young crocodilian to the Museum of Comparative Zoology for

Figure 11. Lateral views of the anterior ends of the tails of: A. *Tomistoma schlegelii*, MCZ 9327, showing absence of basicaudal irregularity. B. *Osteolaemus tetraspis*, MCZ 2107, showing two thick granular strips separating the dorsal from the ventrolateral scalation.

identification. The beast had been sold as *Crocodylus acutus*, but appeared to differ from that taxon by the number of osseous scutes on the neck. Ernest E. Williams ducked the question, suggesting that crocodiles were not yet understood, and that work should be done on the problem. We thank him for inspiring and facilitating the bulk of this work.

We report primarily on specimens in the American Museum of Natural History (AMNH), the Field Museum of Natural History (FMNH), the Museum of Comparative Zoology (MCZ), and the National Museum of Natural History (USNM). We have also examined specimens in the Boston Museum of Science (BMS), the California Academy of Sciences (CAS), the Museo Nacional de Historia Natural Santo Domingo, (MNHNSD), the Instituto de Historia Natural (Tuxtla Gutierrez, Chiapas), the National Museum of Canada, and the Yale Peabody Museum. We are indebted to the curators and staffs of these institu-

tions, especially Ernest E. Williams, Hymen Marx, Miguel Alvarez del Toro, George Zug, Charles A. Ross, and the late James A. Peters. We are also indebted to Aleta Karstad for Figure 4, Jim Lovesek for photographs of *Crocodylus intermedius*, and E. E. Williams and Don McAllister for radiographs. In addition, we thank A. S. and S. B. Avery, J. M. Clark, F. R. Cook, P. Elias, K. Miyata, A. Rhodin, J. P. Rosado, J. P. Ross, F. W. and A. K. Schueler, and R., R., and F. Whiston. F.D.R. was supported in part by a grant from the Ella Lyman Cabot Trust, and G.C.M. was supported by an NSF graduate fellowship and NRM.

APPENDIX: SPECIMENS EXAMINED

Museum abbreviations are in the Acknowledgments. Numbers in parentheses are the number of individuals in lots designated by a single museum number.

TABLE 1. NUMBER OF CONTIGUOUS SCUTES PER TRANSVERSE ROW IN PC 1-PC 18.

Number given is mode (or modes) with range underneath in parentheses. Species are identified by the first four letters of their generic and specific names. N = sample size.

Species	N	PC 1	PC 2	PC 3	PC 4	PC 5	PC 6	PC 7	PC 8	PC 9
Gavi gang	8	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)
Tomi schl	12	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4-6)	6 (6)	6 (5-6)	6 (6-7)
Croc cata	11	4 (4)	4 (4)	4 (4)	4 (4-5)	6 (4-6)	6 (4-6)	6 (4-6)	6 (4-8)	6 (4-7)
Croc john	11	4 (4)	4 (4)	4 (4)	4 (3-5)	4,6 (4-6)	6 (6)	6 (6)	6 (6)	6 (6)
Croc nilo	19	4 (4)	4 (4)	4 (4-6)	6 (4-6)	6 (4-6)	6 (4-6)	6 (4-8)	6 (6-8)	6 (6-8)
Croc palu	12	4 (4)	4 (4)	4 (4-5)	4 (4-6)	6 (4-6)	6 (5-6)	6 (6-7)	6 (4-7)	4,5 (4-8)
Croc siam	6	4 (4)	4 (4)	4 (4)	4 (4-6)	6 (4-7)	6 (6-7)	6 (5-8)	5 (4-8)	6 (5-6)
Croc poro	12	4 (4)	4 (4)	4 (4-6)	6 (4-6)	6 (4-6)	6 (6-8)	8 (6-8)	6 (6-8)	6 (6-8)
Croc nova	14	4 (4)	4 ¹ (4)	4 ^k (4-5)	6 (4-6)	6 (6)	6 (6-8)	8 ¹ (6-8)	8 (8)	8 ⁱ (6-11)
Croc rhom	6	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	6 (5-6)	6 (6)	6 (6)	6 (6-7)
Croc more	8	4 (3-5)	4 (4)	4 (3-4)	4 (3-5)	4 (3-5)	4 (2-6)	3 (3-6)	6 (2-6)	6 (4-6)
Croc acut	37	4 (3-5)	4 (2-4)	4 (2-4)	4 (2-4)	4 (2-5)	3,4 (2-6)	4 (2-6)	3 (2-6)	4 (2-6)
Croc inte ^a	2	4,5	4	4	4	5,6	5	5,6	6	5
Oste tetr	18	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4-6)	4 (4-6)	6 (4-6)	6 (4-7)
Alli sine	14	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4)	4 (4-6)	6 (4-6)	6 (6-8)
Alli miss	17	4 (4)	4 (4)	4 (4)	4 (4-6)	4 (4-6)	6 (4-6)	6 (5-7)	6 ^j (6-8)	6 (6-8)
Caim lati	12	4 (4-6)	4 ¹ (4)	4 ¹ (4)	4 (4)	6 (5-6)	6 (6)	6 ¹ (6-8)	8 (4-9)	8 (5-9)
Caim croc	24	4 ^o (4-6)	4 ^o (4-6)	4 ^o (4-6)	6 ^o (4-8)	6 ^o (6-10)	8 ^o (6-11)	8 (7-12)	8 (8-12)	8 (8-12)
Mela nige	15	6 (6)	6 (5-6)	6 (6)	6 (6)	8 (6-8)	8 (8-10)	8 (8-10)	10 (8-12)	10 (9-11)
Pale palp	5	4 (4)	4 ^c (4)	4 (4)	4 (4)	4 (4)	4,5 (4-6)	6 ^c (6-8)	7 (6-7)	8 ^b (6-9)
Pale trig	25	4 (4)	4 ^o (2-4)	2 (2-4)	2 (2-4)	4 (4-5)	4 (4-5)	4 ^o (3-6)	5 (4-6)	6 (4-7)

TABLE 1. CONTINUED.

Species	N	PC 10	PC 11	PC 12	PC 13	PC 14	PC 15	PC 16	PC 17	PC 18
Gavi gang	8	4 (4)	4 (4)	4 (4)	4 ^d (4)	4 ^d (4)	4 ^d (4)	4 ^d (4)	4 ^d (4)	4 ^c (4)
Tomi schl	12	6 (6-8)	6 (6-8)	6 (5-7)	6 ^g (5-6)	4 ^g (4-6)	4 ^g (4-6)	4 ^g (4)	4 ^g (4)	2 ^g (2-4)
Croc cata	11	6 (4-6)	6 (5-8)	6 (4-7)	6 ^f (5-7)	6 ^f (6)	6 ^f (6)	6 ^f (2-6)	4 (2-4)	2 ^f (1-4)
Croc john	11	6 (6)	6 (6-7)	6 (6-7)	6 (6-7)	6 (6)	6 (6)	6 (4-6)	4 (2-4)	2 (1-2)
Croc nilo	19	6 (4-8)	6 (5-8)	6 (6-9)	6 (6-8)	6 (4-8)	6 (3-7)	6 (0-6)	2 (0-4)	0 (0)
Croc palu	12	4,5,6 (4-6)	5 (4-6)	6 (4-6)	6 (4-7)	6 (6-7)	6 (2-6)	4 (3-5)	2 (0-4)	0 (0)
Croc siam	6	6 (4-6)	6 (5-7)	6 (6-7)	6 (5-7)	6 (6)	6 (6)	4 (4-6)	2 (0-4)	0 (0-2)
Croc poro	12	6 (6-8)	6 (6-8)	6 (6-7)	6 ^j (6-8)	6 ^j (4-6)	6 ^j (0-6)	2 ^j (0-4)	0 ^j (0-4)	0 ^j (0)
Croc nova	14	8 ⁱ (8-12)	8 ⁱ (7-11)	8 (8-9)	8 ^m (6-8)	8 ^k (6-8)	6 ^k (6)	6 ^k (4-6)	4 ^k (0-5)	0 ^k (0)
Croc rhom	6	6 (6)	6 (6)	6 (6)	6 (6)	6 (6)	6 (6)	6 (6)	4 (2-6)	0,1,2 (0-2)
Croc more	8	4 (2-6)	4 (4-6)	4 (3-6)	2 (2-6)	6 (2-6)	4 (2-6)	2 (2-4)	0 (0)	0 (0)
Croc acut	37	3 (2-5)	4 (2-4)	4 (2-5)	3 (0-4)	2 (2-5)	2 (0-6)	2 (0-4)	0 (0-2)	0 (0)
Croc inte ^a	2	5,6	5,6	6	6	6	6	4	0	0
Oste tetr	18	6 (4-8)	6 (4-7)	6 (4-8)	6 (6)	6 (5-7)	6 (6)	6 (4-6)	4 (2-6)	2 (0-4)
Alli sine	14	6 (6-8)	6 (6-8)	6,7 (6-8)	6 (6-8)	6 (4-6)	6 (4-6)	6 (4-6)	4 (0-5)	0 (0-2)
Alli miss	17	6 (6-8)	8 (5-8)	8 (6-8)	8 (6-8)	6 (6-8)	6 (4-6)	6 (4-6)	6 (4-6)	2 (2-6)
Caim lati	12	6 (6-9)	6 (5-8)	6 (6-8)	6 ⁱ (6-7)	6 (5-8)	6 (4-6)	6 (4-6)	4 (2-6)	2 ^c (2-4)
Caim croc	24	8 (8-12)	8 (6-12)	8,9 (6-11)	8 (6-11)	8 (6-10)	6 (5-8)	6 (6-8)	4 (2-6)	2 ⁿ (0-6)
Mela nige	15	10 (10-11)	10 ^h (9-12)	10 ^h (9-11)	10 ^h (9-11)	10 (9-11)	10 (8-10)	8 (7-10)	6 (6-8)	6 (4-6)
Pale palp	5	8 (6-8)	7,8 (6-8)	8 ^c (6-8)	6 (6-8)	6 (6-7)	6 (6)	6 (5-7)	6 (4-6)	4 (3-6)
Pale trig	21	6 (4-6)	6 (4-7)	5,6 ^o (4-8)	6 ^o (4-7)	5 ^o (4-6)	4 ^o (4-6)	4 ^o (2-6)	4 ^o (0-4)	2 ^o (2-4)

^aRange not given since N=2. ^bN=6 ^cN=9 ^dN=10 ^eN=11 ^fN=12 ^gN=13 ^hN=14 ⁱN=15 ^jN=16 ^kN=18 ^lN=19
^mN=21 ⁿN=23 ^oN=25

TABLE 2. NUMBER OF DOUBLE AND SINGLE CREST CAUDALS IN UNDAMAGED SPECIMENS OF LIVING CROCODILIANS. Species are identified by the first four letters of their generic and specific names.

Species	Number of double crest caudals															Number of single crest caudals										
	9	10	11	12	13	14	15	16	17	18	19	20	21	16	17	18	19	20	21	22	23	24	25			
Gavi gang										3	2								1	1						
Tomi schl									1	7	3						2	6	2							
Croc cata								5	7					2	1	2	1	1	3	1						
Croc john									10	2						2	1	3								
Croc nilo								1	6	9	8		1		1				6	6						
Croc palu									2	8	2	1		1	5	2	1									
Croc siam										1	3	2				1		1	2							
Croc poro										3	7	9					2	1	3	3						
Croc nova										7	11	2				6	6	3								
Croc rhom										3	4			1	1	1										
Croc more										3	4	1				1	2	1	1							
Croc acut								1	7	8	15	5		1	20	17	2									
Croc inte ^a									1	9	1					3	4	2								
Oste tetr		1	4	9	3	1										2	4	5	2							
Alli sine							2	8	3											4	3	4				
Alli miss										6	11					1			6	2	3					
Caim lati					3	9	2		1											1	1	4	3			
Caim croc			1	3	2	13	4	1											3	7	5	3				
Mela nige								1	8	6										1	3	3	2			
Pale palp	1	5	3																1							
Pale trig	2	21	2											2	4	2	1									

^a Data from Medem, 1958a, Table 1. The single specimen of this species available to us was also examined by Medem. Medem's last dorsal corresponds to PC 1 in this specimen; thus, his counts correspond to the method used in this paper.

TABLE 3. FORMATION OF Y-SHAPED CAUDAL KEEL FROM PELVIC KEEL ROWS.

Median pelvic keel rows merge with lateral pelvic keel row to become Y-shaped caudal keel	Median pelvic keel rows do not contribute to the Y-shaped caudal keel	
	Median pelvic keel rows become an unpaired median keel row diminishing posteriorly	Median pelvic keel rows maintain their paired nature until diminishing posteriorly
<i>Paleosuchus</i> , <i>Alligator sinensis</i>	<i>Caiman</i> , <i>Melanosuchus</i>	<i>Gavialis</i> , <i>Tomistoma</i> , <i>Crocodylus</i> , <i>Osteolaemus</i> , <i>Alligator mississippiensis</i>

TABLE 4. OCCURRENCE OF FOUR KINDS OF BASICAUDAL IRREGULARITIES IN EXAMINED SPECIMENS OF LIVING CROCODYLIANS.

The irregularities are further described in the Methods. + = present; - = absent; * = thinner than in other species. Species are identified by the first four letters of their generic and specific names.

Species	One thin granular strip	One or more thick strips	Lateral intrusions	Ventrolateral intrusions
Gavi gang	-	-	-	-
Tomi schl	-	-	-	-
Croc cata	+/-	-	-	-
Croc john	+/-	-	-	-
Croc nilo	*/-	-	-	-
Croc palu	+/-	+/-	-	-
Croc siam	+/-	+/-	-	-
Croc poro	+/-	+/-	+/-	-
Croc nova	+/-	-	+/-	-
Croc rhom	-	-	-	-
Croc more	*/-	-	+/-	+
Croc acut	+/-	*/-	-	-
Croc inte	-	-	+/- ^a	-
Oste tetr	-	+/-	+/-	-
Alli sine	-	-	+/-	-
Alli miss	-	-	+/-	+/-
Caim lati	-	+/-	+/-	-
Mela nige	*/-	-	+	-
Pale palp	-	-	+/-	-
Pale trig	-	+/-	-	-

^aC. A. Ross, personal communication.

Gavialis gangeticus. INDIA: AMNH 81802, MCZ 5263, 161013, USNM 136615; NO LOCALITY: AMNH 48500, BMS no #, FMNH 82681, no # (2), USNM 211524.

Tomistoma schlegelii. MALAYSIA: USNM 84247, 85106; SUMATRA: FMNH 11084-5, MCZ 9327, USNM 58136, 65509-10, 145586-8; NO LOCALITY: AMNH 66380, 101422, FMNH 134997, 206755, USNM no #.

Crocodylus cataphractus. ANGOLA: MCZ 29360; TANZANIA: MCZ 54650, 54751; ZAIRE: AMNH 10076, 45863, MCZ 16732; NO LOCALITY: AMNH 2206, 6584, 7293, 8877, FMNH 37213, MCZ 5456, USNM 60578.

Crocodylus johnstoni. AUSTRALIA: AMNH 82535, 86536-7, 86539, MCZ 35001-4, 35006, 111978, USNM 64090-1; NO LOCALITY: AMNH 93743.

Crocodylus niloticus. BOTSWANA: USNM 195448, 195782-3; CAMEROON: FMNH 190827-9; CHAD: MCZ 27070-1; ETHIOPIA: USNM 42860, 136621; GHANA: MCZ 55153; KENYA: MCZ 40001, USNM 63592; MADAGASCAR: MCZ 2114, 5461, 12552, 16729-31, 16868; MALAWI: MCZ 30000; MALI: FMNH 20799; SIERRA LEONE: MCZ 54120; SOUTH AFRICA: FMNH 17155; TANZANIA: MCZ 48000.

Crocodylus palustris. INDIA: AMNH 2413-8, FMNH 31538, 63739; SRI LANKA: MCZ 42176, 42180; NO LOCALITY: AMNH 45099, FMNH 95995, MCZ 6753, 17595-7.

Crocodylus siamensis. THAILAND: MCZ 3716, 20301, USNM 79476; NO LOCALITY: AMNH 75035, USNM 3906, 8893.

Crocodylus porosus. AUSTRALIA: AMNH 86541; INDONESIA (IRIAN JAYA): AMNH 62041-2; INDONESIA (SUMATRA): MCZ 43742; PAPUA NEW GUINEA: MCZ 110450-1, 142950, USNM 211310; PHILIPPINES: FMNH 14346, 52363-4, USNM 38081; SANTA CRUZ IDS.: AMNH 42162; SOLOMON IDS.: MCZ 72936; THAILAND: USNM 72730-6; NO LOCALITY: AMNH 66388.

Crocodylus novaeguineae. PAPUA NEW GUINEA: CAS 119186-7, 121997-8; MCZ 65284-5, 101338, 120335, 140805-7, USNM 211297, 211305-6, 082826-7 (field numbers); PHILIPPINES: FMNH 52357-60, 52362; NO LOCALITY: USNM 083141 (field number).

Crocodylus rhombifer. CUBA: FMNH 34677, MCZ 1249, 1397, 12081, 33394, USNM 13576; NO LOCALITY: AMNH 77595.

Crocodylus moreletii. BELIZE: FMNH 4430,

49369, MCZ 18856; GUATEMALA: USNM 71954, 71957-8; MEXICO: MCZ 8047, 53860, USNM 115357; NO LOCALITY: MCZ 17583.

Crocodylus acutus. COLOMBIA: MCZ 17610; CUBA: MCZ 1250, 12278, 17718; DOMINICAN REPUBLIC: MNHNSD no # (8); HAITI: MCZ 1339, 3596(3), USNM 80892; JAMAICA: MCZ 6093(7), 42981, 44853; MEXICO: FMNH 23147; NICARAGUA: MCZ 5454, USNM 15230; PANAMA: MCZ 1010, 1265, 2003, 2720, 7318, 18927-9, 28051; UNITED STATES: MCZ 13428-30, 13432.

Crocodylus intermedius. COLOMBIA: FMNH 75658.

Osteolaemus tetraspis. CAMEROON: MCZ 29348-9, 33594; CONGO: USNM 62226; GHANA: USNM 207060; LIBERIA: AMNH 82166-7, 116354, MCZ 22913, USNM 58273, 193289; NIGERIA: MCZ 2017; ZAIRE: AMNH 29889, MCZ 12458, 42855-6; NO LOCALITY: AMNH 7766, 24740, MCZ no #.

Alligator sinensis. CHINA: AMNH 28689, 28691-3, FMNH 38234, 38236, MCZ 16706-8, 17545-6, 17548-50, 42982, 161014, USNM 72846.

Alligator mississippiensis. UNITED STATES: MCZ 969, 3767, 13433-4, 16715, 17723, 26949, 159136, 159146-7; NO LOCALITY: MCZ 5458, 17592-3, 159149-52.

Caiman latirostris. ARGENTINA: FMNH 9707-8; BRAZIL: FMNH 5777, MCZ 1151, 1455, 2854, 3588-9, 3769, 17600, 17716, 29099, 159154, USNM 98780; PARAGUAY: MCZ 34226.

Caiman crocodilus. BOLIVIA: AMNH 97310, 97315-23, FMNH 81564-5; BRAZIL: AMNH 87939, MCZ 1241, 2606, 3394, 3972, 3975; COSTA RICA: MCZ 15333-4, 54997; GUYANA: FMNH 26682; MEXICO: MCZ 5040, 17581; PANAMA: FMNH 13402, MCZ 2035 2721, 20566, 22332-3, 29760; SURINAM: MCZ 154922; NO LOCALITY: AMNH 66393.

Melanosuchus niger. BOLIVIA: AMNH 97324; BRAZIL: MCZ 1011, 2547-9, 3590, 3592, 15134, 17598, 17726, 51851, 160240, no #; NO LOCALITY: AMNH no # (2).

Paleosuchus palpebrosus. BOLIVIA: AMNH 97328; BRAZIL: AMNH 16574, 97326-7, MCZ 2928, USNM 149134; COLOMBIA: FMNH 69867, 69869, 69875, 134989; NO LOCALITY: MCZ 160000.

Paleosuchus trigonatus. BRAZIL: AMNH 101056, MCZ 2957, 3785; COLOMBIA: FMNH 69877-8, 69881-2; ECUADOR: AMNH 61548-51, 72470, 113635, MCZ 84029; GUYANA: AMNH 16048, 61513, 61516-8, 64823-5; NO LOCALITY: AMNH 46281-2, 159384.

HYBRID: presumed *Crocodylus acutus* × *C. rhombifer* CUBA: AMNH 82943. Counts from this animal or any other presumed hybrids mentioned in the text are not included in Tables 1-4.

LITERATURE CITED

- ANONYMOUS. 1969. Cuban crocodile. *Animal Kingdom*, **72**(5): 29.
- BOULENGER, G. A. 1889. Catalogue of the chelonians, rhynchocephalians, and crocodiles in the British Museum (Natural History). London, Taylor and Francis.
- BRAZAITEIS, P. 1973 (1974). The identification of living crocodilians. *Zoologica*, **58**: 59-101.
- BREDER, C. M., JR. 1947. An analysis of the geometry of symmetry with especial reference to the squamation of fishes. *Bull. Amer. Mus. Nat. Hist.*, **88**: 321-412.
- BUFFETAUT, E. 1975. Sur l'anatomie et la position systématique de *Bernissartia fagesii* Dollo, L., 1883, crocodilien du Wealdien de Bernissart, Belgique. *Bull. Inst. r. Sci. Nat. Belg.*, **51**: 1-20.
- CHATTERJEE, S. 1978. A primitive parasuchid (phytosaur) from the Upper Triassic Maleri Formation of India. *Palaeontology*, **21**: 83-127.
- COLBERT, E. H., AND C. C. MOOK. 1951. The ancestral crocodilian *Protosuchus*. *Bull. Amer. Mus. Nat. Hist.*, **97**: 143-182.
- CUVIER, G. 1807. Sur les différentes espèces de crocodiles vivans et sur leurs caractères distinctifs. *Ann. Mus. Hist. Nat. Paris*, **10**: 8-66.
- . 1831. The animal kingdom arranged in conformity with its organization. Vol. 9. The class Reptilia. London, Whittaker, Treacher.
- DERANIYAGALA, P. E. P. 1939. The tetrapod reptiles of Ceylon. Vol. 1. Testudinates and crocodilians. *Ceylon J. Sci. Colombo Mus. Nat. Hist. Ser.*
- DUMÉRIL, A. M. C., AND G. BIBRON. 1836. *Erpétologie générale ou histoire naturelle complète des reptiles*. Vol. 3. Paris.
- ERICKSON, B. R. 1976. Osteology of the early eusuchian crocodile *Leidyosuchus formidabilis*, sp. nov. *Monogr. Sci. Mus. Minnesota, Paleontology*, **2**: 1-61.
- FOX, S. F. 1975. Natural selection on morphological phenotypes of the lizard *Uta stansburiana*. *Evolution*, **29**: 95-107.
- GADOW, H. F. 1901. *Amphibia and reptiles*. The Cambridge Natural History Vol. 8. London, Macmillan.
- HECHT, M. K. 1952. Natural selection in the lizard genus *Aristelliger*. *Evolution*, **6**: 112-124.
- HIGGINS, G. M. 1923. Development of the primitive reptilian vertebral column, as shown by a study of *Alligator mississippiensis*. *Amer. J. Anat.*, **31**: 373-407.
- HOFFSTETTER, R., AND J.-P. GASC. 1969. Vertebrae and ribs of modern reptiles, pp. 201-310. In C. Gans, A. d'A. Bellairs and T. S. Parsons (eds.), *Biology of the Reptilia*, Vol. 1. London, Academic Press.
- HUXLEY, T. H. 1859. On the dermal armour of *Jacare* and *Caiman*, with notes on the specific and generic characters of recent Crocodilia. *J. Proc. Linn. Soc. Zool. Lond.*, **4**: 1-28.

- KÄLIN, J. 1955. Crocodilia, pp. 695–784, In J. Piveteau (ed.), *Traité de Paléontologie*, Vol. 5. Paris, Masson.
- KERFOOT, W. C. 1970. The effect of functional changes upon the variability of lizard and snake body scale numbers. *Copeia*, 1970: 252–260.
- MANTER, J. T. 1940. The mechanics of swimming in the alligator. *J. Exp. Zool.*, 83: 345–358.
- MEDEM, F. J. 1958a. Informe sobre reptiles Colombianos, III. Investigaciones sobre la anatomía craneal, distribución geográfica y ecología de *Crocodylus intermedius* (Graves) en Colombia. *Caldasia*, 8: 175–215.
- . 1958b. The crocodilian genus *Paleosuchus*. *Fieldiana, Zoology*, 39: 227–247.
- . 1970. Sobre un híbrido inter-específico del género *Paleosuchus* (Crocodylia, Alligatoridae). *Rev. Acad. Colombiana de Cien, Exactas Fis. y Nat.*, 13: 467–479.
- MILLER, W. E. 1980. The late Pliocene Las Tunas local fauna from southernmost Baja California, Mexico. *J. Paleontology*, 54: 762–805.
- MOOK, C. C. 1941. A new crocodilian, *Hassiacosuchus kayi*, from the Bridger Eocene Beds of Wyoming. *Ann. Carnegie Mus.*, 28: 207–221.
- NASH, D. S. 1975. The morphology and relationships of a crocodilian, *Orthosuchus stormbergi*, from the Upper Triassic of Lesotho. *Ann. S. Afr. Mus.*, 67: 227–329.
- REIG, O. A. 1970. The Proterosuchia and the early evolution of the archosaurs; an essay about the origin of a major taxon. *Bull. Mus. Comp. Zool.*, 139: 229–292.
- ROMER, A. S. 1956. *Osteology of the reptiles*. Chicago, Univ. Chicago Press.
- . 1972. The Chañares (Argentina) Triassic reptile fauna. XII. The postcranial skeleton of the thecodont *Chanaresuchus*. *Breviora Mus. Comp. Zool.* No. 385, pp. 1–21.
- . 1974. Aquatic adaptation in reptiles—primary or secondary? *Ann. S. Afr. Mus.*, 64: 221–230.
- ROSS, C. A., AND C. D. ROBERTS. 1979. Scalation of the American alligator. U.S. Dept. Int. Fish and Wildl. Serv. Spec. Sci. Rep.—Wildl. No. 225.
- ROSS, C. A., AND F. D. ROSS. 1974. Caudal scalation of Central American *Crocodylus*. *Proc. Bio. Soc. Wash.*, 87: 231–233.
- SCHMIDT, K. P. 1956. On the status and relations of *Crocodylus mindorensis*. *Fieldiana, Zoology*, 33: 535–539.
- SEIDEL, M. R. 1979. The osteoderms of the American alligator and their functional significance. *Herpetologica*, 35: 375–380.
- TROXELL, E. L. 1925. *Hyposaurus*, a marine crocodilian. *Amer. J. Sci.*, 9: 489–514.
- VARONA, L. S. 1966. Notas sobre los crocodilidos de Cuba y descripción de una nueva especie del Pleistoceno. *Poeyana Ser. A*, 16: 1–34.
- WALKER, A. D. 1961. Triassic reptiles from the Elgin area: *Stagonolepis*, *Dasygnathus* and their allies. *Phil. Trans. Roy. Soc. Lond. Ser. B*, 244: 103–204.
- . 1968. *Protosuchus*, *Proterochampsia*, and the origin of phytosaurs and crocodiles. *Geol. Mag.*, 105: 1–14.
- WELLNHOFFER, P. 1971. Die Atoposauridae (Crocodylia, Mesosuchia) der Oberjura—plattenkalke Bayerns. *Palaeontographica Abt. A*, 138: 133–165.
- WERMUTH, H. 1953. Systematik der rezenten krokodile. *Mitteil. Zool. Mus. Berlin*, 29: 375–514.
- WERMUTH, H., AND R. MERTENS. 1961. Schildkroten, krokodile, bruckenechsen. Gustav Fischer Verlag, Jena.
- . AND ———. 1977. Liste der rezenten amphibien und reptilien. Testudines, Crocodylia, Rhynchocephalia. *Das Tierreich* 100.
- WILLIAMS, E. E. 1959. Gadow's arcualia and the development of tetrapod vertebrae. *Quart. Rev. Bio.*, 34: 1–32.
- YOUNGPRAPAKORN, U. 1976. Samutprakan, Thailand. Borpit, Thailand.

The Tarsus and Metatarsus of *Protosuchus* And Its Phyletic Implications

MAX K. HECHT¹

SAMUEL F. TARSITANO²

ABSTRACT. The tarsus and metatarsus of the Triassic crocodilian *Protosuchus* is compared with that of modern *Alligator*. The two types of tarsi are very similar, indicating that the present crocodilian tarsus is a primitive feature of the Order. Important differences occur only in the tibial-astragalar articulation, the position and development of the distal roller of the astragalus, and degree of development of the fifth metatarsal.

INTRODUCTION

The Order Crocodilia is of great interest to paleobiology because of its longevity as an amniote taxon and because it is one of the two survivors of the archosaurian adaptive radiation. The earliest known crocodilian genera are of Triassic age and generally placed in the suborder Protosuchia (Bonaparte, 1978; Colbert and Mook, 1951; Crompton and Smith, 1980; Nash, 1975; Steel, 1973). It is not a prerequisite axiom, but it is generally expected that the primitive state of a morphological transformation series is to be found among the earliest fossil representatives of the lineage. One of the features which has been used in determining lineages among early reptiles has been the tarsus and metatarsus. The tarsal region has undergone rapid change among the early Archosauria but unfor-

tunately has not been well preserved in most fossils. It is for this reason that a description of the preserved tarsal and metatarsal elements of *Protosuchus* is required.

The description of the type material of *Protosuchus* indicated the presence of an almost complete tarsus and metatarsus (Colbert and Mook, 1951). Further preparation of the material has now made it possible to describe in detail the earliest known crocodilian tarsus and to make further anatomical comparisons. In order to properly evaluate these new data a detailed osteological study of a living eusuchian crocodilian is required. Until now only general descriptions were available for the various crocodilian tarsal elements (Schaeffer, 1941; Cruickshank, 1979; Brinkman, 1980b). Schaeffer (1941) produced the pioneer study on the amphibian and reptilian tarsus that illustrated the functional and phylogenetic problems within the Reptilia. Krebs (1963) discussed the tarsus as an archosaurian character. Cruickshank (1979) discussed the tarsus as an important character and indicator of lineages and functional trends within the early Archosauria. He also described the basic morphology of the archosaurian tarsus and clearly demonstrated the complex interrelationships among the component parts. Cruickshank stated that early in the history of the Archosauria two types of tarsal morphology and function appeared

^{1,2}Department of Biology, Queens College of the City University of New York, Flushing, New York 11367, U.S.A.

in the groups above the proterosuchian level. Chatterjee (1978) and Cruickshank (1979) called these two types, crocodile-normal and crocodile-reversed. Cruickshank postulated that these two types were derived from a primitive archosaurian condition, the proterosuchian condition, which had a relatively mobile relationship between the astragalus and calcaneum. The crocodile-normal condition was represented by the astragalus and its peg articulating with a calcaneum and its socket. In this study Cruickshank (1979) developed a morphological terminology to aid in the description of these complex structures and articulations. He used the genus *Crocodylus* as the typical eusuchian crocodile and the basis of his comparison with other Archosauria. The variation within the living Eusuchia was not considered nor that of other crocodylians. Thulborn (1980) has also discussed the tarsus of early archosaurians but primarily at the proterosuchian level. Brinkman (1980b) described the mechanics of the eusuchian tarsus and metatarsus using *Caiman* as a model. The preservation of tarsal elements among fossil crocodylians is not frequent and is unknown in *Hemiprotosuchus* (Bonaparte, 1971).

The tarsus, metatarsus, and phalanges of *Protosuchus* are almost completely preserved in the type material of *P. richardsoni* (AMNH 3024). It is the intent of this study to compare the tarsus and metatarsus of *Protosuchus* with the typical crocodylian tarsus as represented by members of Eusuchia. In order to accomplish this comparison it is necessary to describe in detail those components of the eusuchian tarsus which are preserved in *Protosuchus*. Schaeffer (1941) and Cruickshank (1979) have provided us with a brief description and terminology based primarily on *Alligator* and *Crocodylus*. Brinkman (1980b), not knowing of Cruickshank's work, described the tarsus using a different terminology. All of the preceding authors considered the functional aspects of the tarsal elements.

THE MORPHOLOGY OF THE EUSUCHIAN TARSUS AND METATARSUS

As an example of a typical crocodylian we are utilizing the tarsus of the genus *Alligator*. The illustrations are of a large *Alligator mississippiensis* in our own collection (R1200) which is believed to have had a total length of eleven feet. The tarsus and metatarsus was well ossified and preserved and all facets and condyles well defined. The morphological terminology used shall be that of Cruickshank (1979). Certain modifications and additions will be required in this terminology. Morphological variation in the Eusuchia will be noted.

ASTRAGALUS

The astragalus of eusuchian crocodyles is firmly attached although not fused to the tibia (Brinkman, 1980b). The proximal surface of the astragalus (Figs. 1, 2) presents a U-shaped tibial facet. The arms of the U are directed laterally while the belly of the U faces medially. The anterior arm of the U is much longer than the posterior arm and culminates in a ridge. This ridge marks the end of the articular surface of the tibial facet and the beginning of the fibular facet lying perpendicular to it. Between the arms of the U-shaped tibial facet lies a moderately large fossa, the astragalar fossa. It is filled in life with connective tissue which conceals many nutritive foramina to the astragalus and helps bind it to the tibia. The fibular facet is a slightly concave uneven rectangular surface which articulates with the medio-distal surface of the fibula through an interposed meniscal cartilage.

The distal surface of the astragalus (Figs. 3, 4) is slightly complex in that it consists of two separate articular surfaces adjacent to one another but separated by a groove. The larger of the two articular surfaces is the anteriorly directed distal roller (Cruickshank, 1979) which articulates with metatarsals 1 and 2. The distal roller extends from the tibial facet prox-

imally and anterior hollow laterally and continues around the anterior (distal) surface of the astragalus to its posterior surface. The second articular surface is a complex low bent scroll-shaped mound henceforth referred to as the astragalar trochlea. The lateral surface of the trochlea forms a prominent feature, the astragalar peg. The surface of the trochlea articulates with a concave shelf of the calcaneum (Cruickshank, 1979) or part of the secondary articulation of Schaeffer (1941). The astragalar peg forms the primary articulation between the socket of the calcaneum and astragalus (ball and socket of Schaeffer, 1941). This peg is formed by the lateral most edge of the astragalar trochlea and articulates with the socket of the calcaneum (concave facet of the calcaneum, Cruickshank, 1979).

The posterior surface (Fig. 4) of the astragalus contains the trochlea described above, part of the distal roller and a large groove. This groove is hereby designated the articulating channel. It is bounded dorsally by the fibular facet, ventrally by the trochlea and peg, and extends medioventrally to become continuous with the lesser groove lying between the distal roller and the medial eminence of the trochlea. The articulating channel also extends to the lateral surface of the astragalus and receives the leading edge of the calcaneal tongue as the calcaneum rotates about the astragalus.

It should be noted that we are not in agreement with some of Cruickshank's terminology because of a contradiction between his labels and orientation of the astragalus (Cruickshank, 1979: Fig. 12). His labeled "groove" in Figures 12b and 12c appear to be different structures and are therefore confusing.

CALCANEUM

Proximally the calcaneum has a fibular condyle, which produces a single moveable articulating surface with the fibula. The fibular condyle (the fibular facet of

Cruickshank, 1979) is rounded proximally but becomes abruptly flattened distally for its articulation with distal tarsal 4, indicating that the movement at this latter joint is restricted.

Medially the calcaneum (Fig. 5) bears two articulating surfaces for the astragalus. The most anterior of the two is a socket (Schaeffer, 1941; or the concave facet of Cruickshank, 1979), which receives the peg of the astragalus. Posterior to this socket is a medially directed tongue of bone (Cruickshank, 1979) which is concave anteriorly and articulates with the rounded trochlea of the astragalus. The dorsal lip of the tongue articulates with the astragalar groove.

Posterolaterally the calcaneum bears a large tuber (Figs. 5, 6) which is distinct from the fibular condyle anteriorly and the tongue anteromedially. On its posterior surface the calcaneal tuber bears a deep groove for the tendons of the long pedal flexors (Brinkman, 1980b), henceforth referred to as the deep groove (Fig. 6). It should be noted that most of the *M. gastrocnemius* passes over the groove or medial to the calcaneal tuber. Most of the long pedal flexor muscles insert on the fifth metatarsal, the tuber changing the application of the force of *M. ambiens*, *M. flexor tibialis externus*, and the fibular head of the *M. gastrocnemius*. One muscle, *M. fibulo-calcaneum*, does insert onto the dorsal surface of the tuber (Fig. 5). Below this point lies a centrally directed channel (Fig. 5) which terminates at the bridge. On the lateral side of the bridge there lies a lateral channel (Fig. 2).

FOURTH DISTAL TARSA

Distal tarsal 4 (Fig. 12) is a triangular bone oriented so that its length is longer than its width. Although the bone is small, it has many articular surfaces. Proximally there is a dorsoventrally broadened flat area which articulates through a meniscus with the distal tarsal 4 facet of the calcaneum. Ventral and distal to the above facet is a nonarticular

concavity perforated with many nutritive foramina. Anterior to this concavity a slightly concave facet extends in a postero-dorsoventral direction to articulate with the fifth metatarsal. The distal surface of the bone is imperfectly concave with an indentation in the middle of its surface. Above the indentation a large eminence bears the articular surfaces for metatarsal 3 (medial surface) and metatarsal 4 (lateral surface). Viewed medially the bone is heart shaped, and its entire surface articulates with distal tarsal 3.

FIFTH METATARSAL

The fifth metatarsal of all crocodilians is reduced and bears no phalanges. This bone viewed dorsally is roughly triangular in shape and is thickened proximally, where there is an elongate convex area for the articulation with distal tarsal 4. The anterolateral surface bears a conical shaped projection which comprises one-third of the total length of the bone. Dorsoproximally there is a prominence which tapers posterolaterally and bears a scarred surface for the insertion of part of *M. gastrocnemius*, other pedal flexor muscles and the cartilage which lies between the calcaneum and fourth distal tarsal (Schaeffer, 1941; Brinkman, 1980b).

OTHER METATARSALS

The metatarsals of crocodilians are modified for a sprawling gait (Brinkman, 1980b) but differently from the sprawling gait of lizards (Brinkman, 1980a). In the Crocodilia this adaptation manifests itself in the torsion and compression of the proximal portion of all of the metatarsals except 5. The first metatarsal is the most robust and the least modified of the four. It is dorsoventrally flattened, and its lateroproximal surface lies on top of the dorsomedial surface of metatarsal 2. Metatarsal 2 is slightly proximally twisted and compressed so that its broad surface faces dorsomedially and underlies the proximal surface of metatarsal 1. Metatarsal 3 is similar in form to meta-

tarsal 2, and its proximal portion lies under metatarsal 2. Metatarsal 4 is the most slender of the five metatarsals similar to the preceding metatarsals in form. The proximolateral surface of metatarsal 4 is covered by metatarsal 3. Metatarsal 1 articulates proximally with the distal roller of the astragalus. Metatarsal 2 articulates proximomedially with the distal roller and proximolaterally with distal tarsal 3. Metatarsal 3 articulates proximomedially with distal tarsal 4. Metatarsal 4 articulates proximally with the dorsodistal surface of tarsal 4, whereas metatarsal 5 articulates proximally with the ventrolateral surface of tarsal 4.

VARIATION AMONG LIVING EUSUCHIA

The preceding description was based on the genus *Alligator*, of a particularly large individual. Some of the osteological features are not as prominent in smaller specimens or in smaller species. Therefore there appears to be an allometric component, which is primarily reflected in slight changes in proportion and degree of ossification. Comparisons with other genera indicate a similar degree of variation as exemplified by the figures of Brinkman (1980b) of *Caiman*, and Cruickshank (1979) and Thulborn (1980) of different species of *Crocodylus*. An examination of *Gavialis*, the most divergent of the living forms, demonstrates similar minor alterations but the basic crocodilian tarsus and metatarsus is still present. Compared to *Alligator*, *Gavialis* has a reduced deep groove of the calcaneal tuber, slightly less spheroidal distal roller and slightly more robust metatarsals.

THE MORPHOLOGY OF THE PROTOSUCHIAN TARSUS AND METATARSUS

ASTRAGALUS

The astragalus of *Protosuchus* (Figs. 7-9) differs from that of modern crocodilians in some proportional and minor

morphological features. The proximal surface of the astragalus differs from eusuchian crocodilians by the presence of a deep channel which receives the end of the tibia, thus locking it into place. The tibial facet is not U-shaped as in the Eusuchia, though the fibular facet is almost perpendicular to the tibial facet as it is in modern crocodilians.

The distal roller is prominent, as in modern crocodilians, but the orientation is more medial and vertical in relation to the remainder of the astragalus. This different orientation causes the tibia be held at a more acute angle in order to enable metatarsal 1 and 2 to contact the ground (compare Figs. 8 and 12).

The anterior hollow adjacent to the distal roller exists in *Protosuchus* but is not as depressed (Fig. 8) as in the modern Crocodilia.

The medial surface of the astragalus bears the partially destroyed peg and a shallow articulating channel (Fig. 9). This channel intersects the lesser groove which is not as well developed as in the eusuchians. There is no astragalar fossa in *Protosuchus* because of the absence of the posterior arm of the tibial facet. As in modern forms the astragalar trochlea is directed postero-laterally to receive the calcaneal tongue.

CALCANEUM

The calcaneum of *Protosuchus* bears a striking resemblance to the typical eusuchian despite the latero-medial compression in the preserved specimen. This severely compressed portion of the calcaneum includes the fibular condyle, distal tarsal 4 facet, and part of the socket. In the lateral view the proximal surface bears a rounded fibular condyle. As in all Eusuchia, there is a distal flattened surface below the fibular condyle for the articulation with distal tarsal 4 (Fig. 10).

The tuber of the calcaneum is a prominent structure distinct from the remainder of the calcaneum. Although the posterior surface of the tuber is distorted

and damaged in the specimen there are clear indications of the deep groove of the tuber (Fig. 11).

On the medial surface there is a medially directed tongue bearing an anteriorly facing concave surface which articulates with the astragalar trochlea. The medial surface of the calcaneum has the identical structures as the Eusuchia, the most prominent of which is the socket.

FOURTH DISTAL TARSAL

Distal tarsal 4 is preserved in the natural position but is clearly medio-laterally compressed. As a result the bone is distorted into a half moon shaped disc. Laterally the articulating surface for metatarsal 5 has been exposed by the rotation of metatarsal 5. The posterior surface is flattened for the articulation with the distal tarsal 4 facet of the calcaneum. The anterior or distal surface of this bone is rounded for articulation with metatarsal 4 and 3. The medial surface is flattened, but this may be an artifact of preservation.

FIFTH METATARSAL

The fifth metatarsal is reduced and as in modern Crocodilia bears no phalanges. Proximally the fifth metatarsal is bulbous and tapers distally into a long fingerlike projection. The entire structure is one third the length of metatarsal 2.

The position of metatarsal 5 in the specimen appears to have been altered from its natural position. The bone has been rotated approximately 90° exposing the articular surface of distal tarsal 4. Thus the lateral view as exposed in the fossil represents the dorsal bone surface and the bottom view, the lateral surface. It is possible that the entire fifth metatarsal may be medio-laterally compressed. The true dorsal surface bears a fossa for probable muscle insertions, possibly *M. peroneus anterior* and *posterior*. The true lateral surface bears a deep

groove or elongate pocket. If this is a natural feature, it would be for the passage and insertion of part of the gastrocnemius muscle.

OTHER METATARSALS AND PHALANGES

The metatarsals of *Protosuchus* have the same relationships to each other in morphology, proportion, lengths, and articulations as those of modern eusuchians (Colbert and Mook, 1951).

The number of phalanges is similar to most modern Eusuchia having a phalangeal formula of 2-3-4-4-0 (Colbert and Mook, 1951).

CONCLUSIONS

THE PROTOSUCHIAN AND EUSUCHIAN TARSUS AND METATARSUS COMPARED

1) The crocodilian "normal" tarsus and its metatarsus were completely developed by the earliest known Crocodilia.

2) Between the protosuchian level and the eusuchian level the tarsus developed greater internal mobility by the reduction of the channel between the astragalus and the tibia and the development of the ligamentous support system described by Brinkman (1980b).

3) The position of the distal roller in *Protosuchus* indicates a more primitive sprawling limb position (Brinkman 1980a, b; Rewcastle, 1980) than is found in modern Crocodilia and the probable inability of the protosuchian crocodile to develop some of the advanced gaits described by Brinkman (1980b).

4) The longer and perhaps more complex fifth metatarsal of *Protosuchus* suggests a more primitive condition but only slightly different from the eusuchian condition. It is evident from the muscle attachments that the vestige of the fifth digit, the fifth metatarsal, in crocodilians cannot be reduced for reasons outlined by Russell and Rewcastle (1979) for the lizard *Sitana*.

The functions of the hooked fifth metatarsal and its distribution among reptilian taxa have been described by Robinson (1975). The hooking of the fifth metatarsal is apparently primitive as indicated by its presence in some early eosuchians (Gow, 1975; Carroll, 1975). The crocodilian condition must be reduced as far as the loss of the distinct hooking pattern.

THE CROCODYLIAN AND PRIMITIVE ARCHOSAURIAN TARSUS

Thulborn's analysis (1980) of the ankles of archosaurian reptiles demonstrates that previous investigators had oversimplified the primitive proterosuchian level of tarsal development (Charig, 1972; Cruickshank, 1979; Hughes, 1968; Krebs, 1963; Reig, 1970). It is obvious that the complexity of the morphological transformation series has been underestimated and the polarities of the tarsal complex have not yet been determined. Thulborn's analysis implies that he has removed reversals and convergences from the tarsal morphocline, but instead he has created paraphyletic and polyphyletic taxa. From our viewpoint he has not removed the contradictions in the systematics of the group (Charig *et al.*, 1976; Krebs, 1974; Reig, 1970; Romer, 1972; Sill, 1967, 1974; Steel, 1973; Tarsitano and Hecht, 1980; Walker, 1972; Whetstone and Martin, 1979), but he has pointed out a serious conflict in the morphological transformation series (Carroll, 1976; Cruickshank, 1975; Ewer, 1965; Krebs, 1963). At the present time the most important question is whether the "crocodile-normal" tarsus is an indicator of a monophyletic origin of taxa or whether it has evolved independently several times (Thulborn, 1980). An examination of the figures of Chatterjee (1978) indicates such similarity in the basic structure of the phytosaurian tarsus as to indicate that such groups must share a common ancestor with crocodilians. The crocodile-normal and crocodile-reversed tarsi are such complex struc-

tures as to require monophyletic origins for the lineages they characterize.

ACKNOWLEDGMENTS

The authors wish to express their thanks to E. Gaffney of the American Museum of Natural History for permission to prepare the material and for his encouragement in making the study; the PSC/BHE faculty research grants (No. 13317) of the City University of New York for the financial support for this study; Matthew Hecht for aid in preparation of the photographs; Otto Simonis for his skill in preparation; and Robin Eisner for one diagram.

The first author would like to express his appreciation to Ernest E. Williams for the many hours of discussion, begun years ago, that has made the study of Evolutionary Biology such a great game.

APPENDIX

The following terminology and their abbreviations have been used in the figures for structures on the astragalus and calcaneum:

ASTRAGALUS

Ac, articulating channel; Ah, anterior hollow; Af, astragalar fossa; At, astragalar trochlea; Atc, astragalus-tibial contact; Dr, distal roller; Ff, fibular facet; Lg, lesser groove; P, peg; Tf, tibial facet.

CALCANEUM

B, bridge of tuber calcaneum; Dg, deep groove; Dt4f, facet of distal tarsal 4; Fc, fibular condyle; Lc, lateral channel; Mc, medial channel; S, socket; T, tongue of the calcaneum; Tc, tuber calcaneum.

LITERATURE CITED

- BONAPARTE, J. F. 1971. Los tetrapodos del sector superior de la Formacion Los Colorados, La Rioja, Argentina (Triasico sup.) Opera lilloana, 22.
- . 1978. El Mesozoico de America del Sur y sus Tetrapodos. Opera lilloana, 26: 1-307.
- BRINKMAN, D. 1980a. Structural correlates of tarsal and metatarsal functioning in Iguana (Lacertilia; Iguanidae) and other lizards. Can. J. Zool., (2)58: 277-289.
- . 1980b. The hind limb step cycle of *Caiman sclerops* and the mechanics of the crocodile tarsus and metatarsus. Can. J. Zool., (12)58: 2187-2200.
- CARROLL, R. L. 1975. Permo-triassic "lizards" from the Karroo. Palaeont. Africana, 18: 133-135.
- . 1976. *Noteosuchus*-The Oldest Known Rhynchosaur. Ann. S. Afr. Mus., 72: 37-57.
- CHARIG, A. J. 1972. The evolution of the archosaur pelvis and hindlimb: an explanation in functional terms, pp. 121-155. In K. A. Joysey and T. S. Kemp, (eds.), Studies in Vertebrate Evolution. Oliver and Boyd.
- CHARIG, A. J., B. KREBS, H. D. SUESS, AND F. WESTPHAL. 1976. Thecodontia. Handbuch der Palaoherpetologie G. Fischer Verlag, Stuttgart.
- CHATTERJEE, S. 1978. A primitive parasuchid (Phytosaur) reptile from the Upper Triassic Maleri formation of India. Palaeontology, (1)21: 83-127.
- COLBERT, E. H., AND C. C. MOOK. 1951. The ancestral crocodilian *Protosuchus*. Bull. Amer. Mus. Nat. Hist., 97: 147-192.
- CROMPTON, A. W., AND K. K. SMITH. 1980. A new genus and species of crocodilian from the Kayenta formation (Late Triassic) of Northern Arizona, pp. 193-217. In L. L. Jacob, (ed.), Aspects of Vertebrate History. Museum of Northern Arizona Press.
- CRUICKSHANK, A. R. I. 1975. The affinities of *Proterochampsia barrionuevei* Reig. Palaeont. Africana, 18: 133-135.
- . 1979. The Ankle Joint in Some Early Archosaurs. South Afric. J. Sci., 75: 168-178.
- EWER, R. F. 1965. The anatomy of the Thecodont Reptile *Euparkeria capensis* Broom Phil Trans. Roy. Soc. Lond. (B), 248: 379-435.
- GOW, C. E. The morphology and relationships of *Youngina capensis* Broom and *Prolacerta broomi* Parrington. Palaeont. Africana, 18: 89-131.
- HUGHES, B. 1968. The tarsus of rhynchocephalian reptiles. J. Zool. Lond., 156: 457-481.
- KREBS, B. 1963. Bau und Funktion des Tarsus eines Pseudosuchiers aus der Trias des Monte San Giorgio (Kanton Tessin, Schweiz). Palaont. Zeit., 57: 88-95.
- . 1974. Die Archosaurier. Naturwiss., 61: 17-24.
- NASH, D. S. 1975. The morphology and relationships of a crocodilian, *Orthosuchus stormbergi*, from the Upper Triassic of Lestho. Ann. S. Afr. Mus., 67(7): 227-329.
- REIG, O. A. 1970. The Proterosuchia and the early evolution of the archosaurs; an essay about the origin of a major taxon. Bull. Mus. Comp. Zool., 139: 229-292.
- REWCASTLE, S. C. 1980. Form and function in lacertilian knee and mesotarsal joints; a contribution to the analysis of sprawling locomotion. J. Zool., 191: 171-192.
- ROBINSON, P. L. 1975. The functions of the hooked

- metatarsal in lepidosaurian reptiles. Colloque in "Problèmes actuels de Paléontologie-Evolution des Vertébrés." Colloque International C.N.R.S. No. 218, pp. 464-483.
- ROMER, A. S. 1972. The Chanares (Argentina) Triassic Reptile Fauna. XVI Thecodont Classification. *Breviora Mus. Com. Zool.* No. 395, pp. 1-24.
- RUSSELL, A. P., AND S. C. NEWCASTLE. 1979. Digital reduction in *Sitana* (Reptilia: Agamidae) and the dual roles of the fifth metatarsal in lizards. *Can. J. Zool.*, **57**: 1129-1135.
- SCHAEFFER, B. 1941. The Morphological and Functional Evolution of the Tarsus in Amphibians and Reptiles. *Bull. Amer. Mus. Nat. Hist.*, (VI)LXXVIII: 395-472.
- SILL, W. D. 1967. *Proterochampsia barrionuevoi* and the early evolution of the Crocodylia. *Bull. Mus. Comp. Zool.*, **135**: 415-446.
- . 1974. The anatomy of *Saurosuchus galilei* and the relationships of the rauisuchid thecodonts. *Bull. Mus. Comp. Zool.*, **146**: 317-362.
- STEEL, R. 1973. Crocodylia, pp. 1-116. In O. Kuhn (ed.), *Handbuch der Paläoherpetologie*: Part 16. G. Fischer Verlag.
- TARSITANO, S., AND M. K. HECHT. 1980. A reconsideration of the reptilian relationships of *Archaeopteryx*. *Zool. J. Linn. Soc.*, **69**: 149-182.
- THULBORN, R. A. 1980. The ankle joints of archosaurs. *Alcheringia*, **4**: 241-261.
- WALKER, A. D. 1972. New Light on the origin of birds and crocodiles. *Nature*, London, **237**: 257-263.
- WHETSTONE, K. N., AND L. D. MARTIN. 1979. New look at the origin of birds and crocodiles. *Nature*, London, **279**: 234-236.

Figure 1. Proximal surface of right astragalus of *Alligator* and explanatory diagram. For symbols see text.

Figure 2. Proximal surface of articulated calcaneum and astragalus of the left side of *Alligator* and explanatory diagram. For symbols see text.

Figure 3. Distal view of the left astragalus of *Alligator* and explanatory diagram. For symbols see text.

Figure 4. Posterior view of the right astragalus of *Alligator* and explanatory diagram. For symbols see text.

Figure 5. Medial view of the right calcaneum of *Alligator* and explanatory diagram. For symbols see text.

Figure 6. Plantar view of right calcaneum of *Alligator* and explanatory diagram. For symbols see text.

Figure 7. Proximal view of right astragalus of *Protosuchus* and explanatory diagram. For symbols see text.

Figure 8. Anterior view of the tibia articulating with astragalus of the left side of *Protosuchus* and explanatory diagram. For symbols see text.

Figure 9. Poster view of the tibia articulating with astragalus of the left side of *Protosuchus* and explanatory diagram. For symbols see text.

Figure 10. Medial view of left calcaneum of *Protosuchus* and explanatory diagram. See text for symbols.

Figure 11. Lateral view of left calcaneum of *Protosuchus* and explanatory diagram. See text for symbols.

Figure 12. Anterior view of the articulated tarsal skeleton of *Alligator* illustrating its relationships to tibia, fibula, and metatarsals.

About the Herpetofauna of Central African Montane Forest

R. F. LAURENT¹

ABSTRACT. The central African montane forest exists as a series of discontinuous habitats where incipient speciation may be occurring. The herpetofauna of the area is examined in regard to microhabitat preference and geographic differentiation.

INTRODUCTION

Exploration of the central African region of the Great Lakes of the Western Graben has focused on Mt. Ruwenzori and the Kivu and Rwanda volcanoes. Other parts of the Nile-Zaire dividing montane range as well as the mountains west of Lake Kivu, of the Ruzizi and of Lake Tanganyika have been poorly explored. The collections made by the author in these little known regions during the 1950's are interesting from the point of view of the differentiation of isolated faunas. Indeed, the montane forest is a highly discontinuous area in central Africa. We can actually count no less than eleven isolated areas: 1) Mt. Ruwenzori; 2) the highlands west of Lake Edward with Mt. Tshiaberimu and scattered mountain forest islands (Butembo, Lubero) in a largely deforested zone; 3) the lakes Mokoto region; 4) the Virunga volcanoes; 5) the highlands west of Lake Kivu with Mt. Kahusi and Mt. Biega; 6) the Idjwi island forests; 7) the Itombwe highlands where the ordinary montane forest surrounds a large grassy central area, the fauna of which is largely

endemic (Laurent, 1964); 8) the highlands between Fizi and Kalemie with Lubitshako and Mt. Kabobo; 9) the Lutsiro highlands of western Rwanda between Kisenyi and Kibuye; 10) the Rugege forest between Bukavu and Butare; 11) the Bururi forest (see Fig. 1). Other practically unknown isolated areas are in the Ituri (Nioka and Blukwa region) and in western Tanzania (Kungwe Mountains).

These forests were interconnected and occupied a larger united area during the last pluvial period, 18 to 25 thousand years ago (Moreau, 1966). Therefore, their flora and fauna is overall rather uniform. It was, however, an attractive problem to investigate if some differentiation had taken place between the isolated parts of this peculiar "milieu" during this relatively short span of time. It must be noted that in some cases the isolation must have been more recent than in others. For instance, the Ituri highlands, the Ruwenzori, the Kungwe Mountains and, to a lesser extent, the Kabobo area have been isolated longer² than the Kahusi and Itombwe range or the forest of the western Rwanda and Burundi where human factors have been at work during a very recent past, as suggested by the high density of population in these countries.

¹Fundación Lillo, Miguel Lillo 205, 4.000 Tucumán, Argentina.

² Evidence for this comes from a) a comparison of the fauna (and flora) on the regions under study; b) the distance; and, c) the depth of the depression between them.

Figure 1. Map of central Africa (eastern Congo, southwestern Uganda, Rwanda, Burundi, and northwestern Tanzania). Striped areas show highlands mostly covered with montane forest. Some include other formations generally at altitudes above 3,000 m (Ruwenzori, Virunga Volcanoes, Itombwe Highlands).

Another feature must be emphasized. On the slopes of the higher mountains which were first explored (Ruwenzori and Kivu volcanoes), the flora is rather distinctly stratified: regular montane forest with *Podocarpus*, Bamboos, *Hagenias* and alpine meadows. Else-

where, in larger areas between 2,000 m and 3,000 m, the first three facies are more or less interwoven and mixed.

Another consequence of the larger surface of the pertinent regions outside the Ruwenzori range and the Kivu volcano is that larger swamps can generally

be found, for instance, in the vicinity of Mt. Biega and Kahusi near Bukavu. Table 1 shows the habitats of the species encountered in the region. This scheme is not a rigid one. Many species, namely the commonest which are also generally the most adaptable, can be found in several kinds of biotopes besides their preferred habitat.

SPECIES OF THE REGION

In the large swamps, the commonest amphibian species of the region are found in great numbers. This is indeed a very striking feature of the amphibian fauna in the montane forest; specimens can easily be collected during the day, while at lower altitudes they hide by day and begin to be active at twilight.

The swamp species are *Xenopus laevis victorianus* Ahl and some polyploids such as the recently described *X. vestitus* Laurent, *X. ruwenzoriensis* Tymowska and Fischberg, *X. wittei* Tinsley and others still undescribed, *Bufo kisoloensis* Loveridge, *Phrynobatrachus versicolor* Ahl, *Phrynobatrachus bequaerti* (Barbour and Loveridge), *Leptopelis kivuensis* Ahl, and *Hyperolius castaneus* Ahl. All of them also inhabit the small marshes which are scattered in the forest and overshadowed by trees. They can also be found in the wells dug by mining prospectors. Except for *Xenopus laevis*, they are often found rather far away from water, but *Phrynobatrachus versicolor* is generally in closer contact with water than the other forms. Some species live in the shadowy puddles of the thick transition forest: *Phrynobatrachus pedropede-*

TABLE 1. ECOLOGICAL DISTRIBUTION OF THE HERPETOFAUNA OF CENTRAL AFRICA MONTANE FOREST.

Small shaded ponds	Bushes	Ground (above and below)
<i>Xenopus</i> sp.	<i>Leptopelis kivuensis</i>	<i>Arthroleptis adolfifriederici</i>
<i>Bufo kisoloensis</i>	<i>Afrixalus laevis</i>	
<i>Phrynobatrachus</i> <i>petropedetoides</i>	<i>Hyperolius castaneus</i>	<i>Cardioglossa cyaneospila</i>
<i>Ph. bequaerti</i>	<i>Cnemaspis</i> <i>quattuorseriatus</i>	<i>Schoutedenella</i> <i>hematogaster</i>
<i>Hyperolius castaneus</i>	<i>Chamaeleo rudis</i>	<i>Lacerta jacksoni</i>
<i>H. frontalis</i>	<i>Ch. adolfifriederici</i>	<i>Algyroides vauereselli</i>
<i>H. chrysogaster</i>	<i>Ch. johnstoni</i>	<i>Leptosiaphos</i> <i>blochmanni</i>
<i>H. leucotaenius</i>	<i>Philothamnus</i> <i>heterodermus</i>	<i>L. meleagris</i>
	<i>Dipsadoboa unicolor</i>	<i>L. graueri</i>
	<i>Atheris nitschei</i>	<i>Typhlops angolensis</i>
Large swamps	Streams	Grassy clearings
<i>Xenopus</i> sp.	<i>Rana angolensis</i>	<i>Phrynobatrachus</i> <i>graueri</i>
<i>Bufo kisoloensis</i>	<i>Rana ruwenzorica</i>	<i>Mabuya megalura</i>
<i>Phrynobatrachus</i> <i>versicolor</i>	<i>Phrynobatrachus</i> <i>acutirostris</i>	<i>Duberria lutrix</i>
<i>Ph. bequaerti</i>	<i>Ph. asper</i>	<i>Psammophylax</i> <i>variabilis</i>
<i>Rana angolensis</i>	<i>Hyperolius frontalis</i>	
<i>chapini</i>	<i>H. discodactylus</i>	
<i>Hyperolius castaneus</i>		
	Bamboos	
	<i>Callixalus pictus</i>	
	<i>Afrixalus orophilus</i>	

toides Ahl, whose digital dilations allow it to climb on leaves, the golden tree frog *Afrixalus laevis* (Ahl), and the white-snouted *Hyperolius frontalis* Laurent. However, in the forest of Rwanda and Burundi the second of these tree frogs is rare, while the third is apparently absent. *Hyperolius discodactylus* Ahl, a green montane tree frog, which was once considered rare, is actually common along streams but rarely encountered in large swamps. *Hyperolius leucotaeniatus* Laurent inhabits the same habitats but its range is restricted to the southern Itombwe highlands and northern Kabobo range, while its northern vicariant *H. chrysogaster* Laurent lives in the transition forest between the latitude of northern Itombwe and Ruwenzori Mt.

Two other montane tree frogs have a limited distribution. *Afrixalus orophilus* (Laurent), described from the Kivu volcano region, is common in Rwanda and Burundi but is unknown in the Ruwenzori and the southern Kivu. The species is often found in rather open places, in contrast with *A. laevis* which likes shade, but in Muhokole (southern Rugege forest) it is common in the broken Bamboos.

The Itombwe Bambu tree frog, *Callixalus pictus* Laurent, has a very peculiar range. West of the Graben it exists in the Itombwe and Kabobo mountains but has never been collected in the Kahusi range nor in the north Kivu, despite careful searching in the bamboo forests. East of the Graben it has been found in the Rugege forest and the Lutsiro highlands south of Kisenyi, but is apparently absent from the volcanoes. This distribution is similar to the range of *Gorilla beringei* (*sensu* Kälin, personal communication), which is limited to the eastern side of a hypothetical ancient Nile coming from upper Katanga, which was captured by the Zaire. *Callixalus pictus* is also limited to the upper levels; rare at 2,300 m, it becomes more and more frequent at higher altitudes and is rather common at 2,600 to 2,700 m. An interesting ecological shift has been observed in the Mt.

Kabobo region where its customary biotopes—the bamboos—is lacking. Here they take refuge beneath the large layers of moss which cover the trunks of big trees.

Besides the green tree frogs like *Hyperolius discodactylus* and *H. frontalis*, some ranids dwell in the streams: giant *Phrynobatrachus* species like *acutirostris* Nieden and *P. asper* Laurent, an Itombwe endemic which is sympatric with *acutirostris* around 2,500 m and the *Ranas*. According to recent investigations (Laurent, 1972), two sibling species exist in the montane forest: a larger form with more extensive webbing, which appears to be *Rana angolensis chapini* Noble (replacing the nominate form in the Great Lakes region) and *R. ruwenzorica* Laurent, a smaller form with reduced webbing. At least one undescribed species seems to live in the Itombwe highlands, increasing further the already high rate of endemism in this interesting region.

In the Itombwe grasslands (see Laurent, 1964), we noticed that *Phrynobatrachus bequaerti*, which belongs to the group of small-sized species previously classified in the genus *Arthroleptis* because of the poorly developed webbing, is more terrestrial than other species. The same species behave similarly in the montane forest, but there is also a closely related species which is even more terrestrial: *P. graueri* Nieden. Like all the species of the genus, it is also aquatic but, whereas *P. bequaerti* prefers still waters, *P. graueri* is more often seen in running waters, i.e., very small brooks. The two species are morphologically very much alike but differ in their coloration: *P. bequaerti* has the thighs and the rear part of the belly orange-colored, while *P. graueri* has the belly and the thighs pinkish.

Completely terrestrial frogs are also found in the montane forest: *Arthroleptis adolfifriederici* Nieden, the most common, exists everywhere, while the more secretive and strangely blue-colored

Cardioglossa cyaneospila Laurent, is mainly found near tree ferns at the lower levels. Both have a largely disjunctive distribution because the forest is interrupted in many places, namely in southern Kivu, Rwanda, and Burundi, but they are also frequent in the transition forest which is the main niche of the blue *Cardioglossa*. The dwarfed arthroleptines of the genus *Schoutedenella* are not known in the northern montane forests except on the Ruwenzori where *S. schubotzi* (Nieden) is not uncommon. In this species, the remarkable sexual dimorphism associated with the arthroleptines, the hypertrophy of the third finger, is very striking, almost as much as in *S. globosa* Witte, since it can be almost as long as the body. In the western Itombwe and the Kabobo massif is *S. hematogaster* Laurent, remarkable for its bright red belly, and in the southern Itombwe and northern Kabobo range the relatively big *S. pyrrhoscelis* (Laurent) with a red network on the belly.

Reptiles are not diverse in these forested highlands. The arboreal species include the secretive gekko, *Cnemaspis quattuorseriatus* (Sternfeld) and three chamaeleons are present everywhere. The robust *Chamaeleo johnstoni* Boulenger is well known by its three-horned males and their spectacular fights. The most common species, however, is *C. rudis* Boulenger, which Stanley Rand (1963) has demonstrated to be a species distinct from *C. bitaeniatus ellioti* Günther, living at lower levels but in the savanna highlands. The strange long-tailed *C. adolfifriederici* Sternfeld is a rare species, but its disjunct range extends from Ituri to the Kahusi Mountains and the Rugege forest. Two still stranger leaf-nosed chamaeleons, *C. xenorhinus* Boulenger and *C. carpenteri* Parker, are Ruwenzori endemics; as they are very rare, Loveridge (1942) supposes that they are canopy dwellers.

Two lacertids very common throughout the montane forests bordering the western Graben are *Lacerta jacksoni*

Boulenger and *Algyroides vauereselli* (Tornier).

The semifossorial skinks of the genus *Leptosiaphos*¹ are also common and live among and under forest floor debris as well as under and within logs: they are *L. graueri* (Sternfeld), *L. meleagris* (Boulenger), *L. hackarsi* (Witte) and *L. blochmanni* (Tornier), which have respectively five fingers and five toes, four fingers and four toes (two species), and three fingers and three toes.

Another skink characteristic of the highland savannas, *Mabuya megalura* (Peters), lives in the Rugege forest (Rwanda and Burundi) in the clearings and *Hagenia* patches which are not dark and where an often grassy soil provides it with its customary niche.

Snakes are infrequent in the montane forests. As always in the forest regions, *Typhlops angolensis* is represented by light-bellied populations: *T. angolensis polylepis* Laurent, in the Itombwe slopes and *T. angolensis irsaci* Laurent, near the Kahusi and the Rugege forest. The only colubrids still common at this level are *Philothamnus heterodermus ruandae* Loveridge, an active frog eater, and *Dasyplectis ater* Sternfeld. *Psammophylax variabilis* Günther is ecologically a highland savanna form like *Mabuya megalura*, but it has been collected in clearings within the montane forest. The beautiful green tree-viper, *Atheris n. nitschei* Tornier is very common in the montane forest. As is generally the case with venomous snakes, its diet is not specialized and comprises frogs as well as lizards and rodents.

DISCUSSION

Following this survey, the main question which arises is whether there is any proof or any hint of a differentiation between the montane forest populations

¹This genus has been recently merged into *Panaspis*.

isolated by distributional gaps created by a strip of highland savanna or of lowland savanna or forest.

An exhaustive analysis of the pertinent material has at present only been made for snakes (Laurent 1956b), but we also have some indications concerning the tree frogs and the chamaeleons.

Let us consider first the case of the snake *Dipsadoboa unicolor viridiventris* Laurent, which is more an inhabitant of the transition forest than of the montane forest, although three specimens have been collected at 2,100 m in the Virunga National Park. There is an apparently homogeneous group of populations living on the Ruwenzori slopes, the forests west of Lake Edward and around Lake Kivu, on the eastern as well as the western side. Four specimens are quite outside of the range of variation of this group and might belong to taxonomically recognizable populations (Fig. 2).¹ Apparently one such distinct population occurs on the Blue Mountains of the Ituri, which have been isolated from other montane forests for a very long time; another is on the Kabobo range, west of Lake Tanganyika and south of the Itombwe, and still another in the Bururi forest (Burundi), which is severed from the Rugege forest by a large deforested area. The fourth specimen comes from the Rwindi plain, which is an unexpected environment for the species.

The viper *Atheris nitschei nitschei* (Tornier) also demonstrates a regional variation, as shown on the diagram (Fig. 3). It can be seen that two samples are especially differentiated: the one on Idjwi Island which is the most completely isolated montane forest of all, and the East Itombwe sample. The eastern slopes of Itombwe are very steep and the inhabited zone consequently narrowed; moreover, the forest appears to be discontinuous, so that some populations can

at the same time be small and isolated, which suggests a Sewall Wright effect (Wright, 1931). Other striking differences appear between Kahusi and Itombwe ranges as well as between the Itombwe range and Upper Lubitshako. In each case, although the two regions are not far away from each other, a low and uninhabited zone intervenes and the gene flow is consequently interrupted. Such a contrast has not been seen for the Bururi sample, which can be explained by the fact that *Atheris nitschei* is well able to withstand the destruction of the forest provided there remain some patches of forestlike undergrowth not too distant from each other. It appears that the Bururi forest is still or has recently been connected to the Rugege forest by a chain of uncultivated areas where *Lobelias* and other montane forest plants form an environment where *Atheris nitschei* can live. If, in the same conditions, *Dipsadoboa unicolor* underwent more differentiation, it is probably because this species is more decidedly a forest dweller and lives at lower altitudes where cultivation has more completely destroyed the suitable habitats. Thus it appears that greater degrees of isolation can be created by human factors at low altitudes than by solely climatic changes at higher altitudes. The apparent reason for this is that the human population is denser and therefore more destructive at moderate heights than on mountain tops.

Chamaeleo rudis schoutedeni Laurent, *Cardioglossa cyaneospila inornata* Laurent, *Hyperolius castaneus submarginatus* Laurent, and *Hyperolius leucotaenitus allogynus* Laurent demonstrate that the populations of the Kabobo range do show an appreciable degree of differentiation. This is a feature which was to be expected when one considers the lowness of the Fizi "bridge," which connects the Kabobo range with the southern Itombwe.

Hyperolius discodactylus appears somewhat differentiated in the Bururi forest. In the Ruwenzori range the

¹They have not been named because these are single specimens.

Figure 2. Correlation between the numbers of subcaudals (ordinate) and ventrals (abscissa) in *Dipsadoboa unicolor viridiventris*. The specimens from Katanda (Rwinda region), Nioka (Ituri), Lubitshako (southern Kivu), and Bururi (Burundi) are outside of the range of variation of the subspecies.

Figure 3. Geographical variation of the numbers of ventrals and subcaudals in *Atheris n. nitschei*. The numbers of specimens in each sample are between parenthesis. The means are connected by a dotted line.

Circled figures correspond to different parts of the montane forest: ① Lubero region; ② Mokoto Lakes region; ③ Virunga volcanoes region; ④ Idjwi Island; ⑤ Kahusi range; ⑥ North Itombwe; ⑦ East Itombwe; ⑧ South Itombwe; ⑨ Lubitshako region; ⑩ Kabobo range; ⑪ Rugege forest; ⑫ Bururi forest.

square-snouted *Hyperolius xenorhinus* has recently been described (Laurent, 1972). Both are extreme areas where such differentiation effects or even endemism (Ruwenzori) were to be expected. Nothing conspicuous has so far been demonstrated elsewhere, but a detailed analysis is likely to prove some incipient divergences wherever the montane forest is discontinuous.

ACKNOWLEDGMENTS

It is a pleasure to dedicate this small essay to Ernest E. Williams, who several years ago revised the text and provided some valuable suggestions to the author. The figures have been executed by Otilia Brizuela.

LITERATURE CITED

- BARBOUR, T., AND A. LOVERIDGE. 1930. Reptiles and Amphibians from the Central African Lakes Region, pp. 764-785. In Strong's African Republic of Liberia and the Belgian Congo. Cambridge, Mass.
- GANS, C. 1964. Further comments on the forms of the African snake genus *Dasypeltis* (Reptilia: Serpentes). *Rev. Zool. Bot. Afr.*, **69**: 279-295.
- LAURENT, R. F. 1950. Diagnoses préliminaires de treize Batraciens nouveaux d'Afrique Centrale. *Rev. Zool. Bot. Afr.*, **44**: 210-212.
- . 1951. Deux Reptiles et onze Batraciens nouveaux d'Afrique Centrale. *Rev. Zool. Bot. Afr.*, **44**: 360-381.
- . 1952a. Reptiles et Batraciens nouveaux du massif du mont Kabobo et du plateau des Marungu. *Rev. Zool. Bot. Afr.*, **46**: 18-34.
- . 1952b. Quelques données nouvelles sur la Systématique et l'Ecologie du genre *Hyperolius*. *Rapp. Ann. Soc. roy. Zool. Belg.*, **82**: 324-334.
- . 1952c. Reptiles et Batraciens nouveaux de la région des grands lacs africains. *Rev. Zool. Bot. Afr.*, **46**: 270-279.
- . 1954a. Remarques sur le genre *Schoutedenella* Witte. *Ann. Mus. Congo. Nov. ser.*, in **4**, **1**, *Miscellanea Zoologica* H. Schouteden: 34-40.
- . 1954b. Aperçu de la Biogéographie des Batraciens et des Reptiles de la région des Grands Lacs. *Bull. Soc. Zool. France*, **79**: 290-310.
- . 1956a. Esquisse d'une faune herpétologique du Ruanda-Urundi. *Les Naturalistes Belges*, **37**: 280-287.
- . 1956b. Contribution à l'Herpétologie de la Région des Grands Lacs de l'Afrique Centrale. I. Généralités II. Chéloniens III. Ophidiens. *Ann. Mus. Roy. Congo Belge*, sér. in **8**, *Sci. Zool.*, **48**: 1-190, pls. I-XXI.
- . 1960a. Notes complémentaires sur les Chéloniens et les Ophidiens du Congo Oriental. *Ann. Mus. Roy. Congo Belge*, sér. in **8**, *Sci. Zool.*, **94**: 1-88.
- . 1960b. Les Serpents du Kivu. *Les Naturalistes Belges*, **41**: 437-452, figs. 1-4.
- . 1964. Adaptive modifications in frogs of an isolated highland fauna in central Africa. *Evolution*, **18**: 458-467.
- . 1971. Caractères biométriques de trois genres monotypiques de la famille des Hyperoliidae. *Bull. Séances Acad. Sc. Outremer. Bruxelles*, **1971-2**: 256-275, 18 figs.
- . 1972. Amphibiens. *Explor. Parc Nat. Virunga*, (2)**22**: 1-124, figs. 1-29, cartes 1-3, pls. I-XI.
- . 1973. Le genre *Leptopelis* Günther (Salientia) au Zaïre. *Ann. Mus. roy. Afr. Centr.* sér. in **8**, *Zool.*, **202**: 1-62.
- . in press. Le genre *Afrixalus* en Afrique Centrale. *Ann. Mus. roy. Afr. Centr.*, sér. in **8**, *Zool.*
- LOVERIDGE, A. 1942. Scientific results of a fourth expedition to forested areas in East and Central Africa IV. Reptiles. *Bull. Mus. Comp. Zool.*, **91**: 237-373, pls. I-VI.
- MOREAU, R. E. 1966. The Bird Faunas of Africa and its islands. New York, Academic Press, 424 pp.
- NIEDEN F. 1912. Amphibia. *Wiss. Ergebn. Deutsch. Zentral Afrika Exped.*, **4**: 165-195, 1 taf.
- RAND, A. S. 1963. Notes on the *Chamaeleo bitaenatus* complex. *Bull. Mus. Comp. Zool.*, **130**: 3-29.
- ROUX-ESTEVE, R. 1974. Revision systématique des Typhlopidae d'Afrique. (Reptilia: Serpentes). *Mem. Mus. Nat. Hist. Nat.*, (n.s.), *A. Zool.*, **87**: 1-313.
- STERNFELD, R. 1912. Reptilia. *Wiss. Ergebn. Deutsch. Zentral Afrika Exped.*, **4**: 197-279, 4 Taf.
- TINSLEY, R. C., H. R. KOBEL, AND M. FISCHBERG. 1979. The biology and systematics of a new species of *Xenopus* (Anura: Pipidae) from the highlands of Central Africa. *J. Zool. London*, **188**: 69-102.
- WITTE, H. G. DE. 1933. Batraciens et Reptiles recueillis par M. L. Burgeon au Ruwenzori, au Kivu et au Tanganyika. *Rev. Zool. Bot. Afr.*, **24**: 97-103.
- . 1941. Batraciens et Reptiles. *Explor. Parc Nat. Albert*, **33**: I-XVII, 1-621, figs. 1-54, pls. I-LXXVI.
- WRIGHT, S. 1931. Evolution in Mendelian populations. *Genetics*, **16**: 97-159.

Zoogeography of the Skinks (Sauria: Scincidae) Of Arno Atoll, Marshall Islands

A. ROSS KIESTER¹

ABSTRACT. The distribution of four species of scincid lizards (*Emoia cyanura*, *E. arnoensis*, *E. boettgeri*, and *Lamprolepis smaragdinum*) on Arno Atoll, Marshall Islands, reveals the following patterns: 1) very small islands are inhabited only by *E. cyanura*, the smallest species; 2) somewhat larger and vegetationally more complex islands usually have *E. cyanura* and *Lamprolepis*, the largest species; 3) as reported by Brown and Marshall (1953), most larger islands have these two plus either *E. arnoensis* or *E. boettgeri*, which are of similar size and intermediate between the first two; 4) two islands were found to have all four species, but these were not the largest or most complex islands. These patterns are discussed in terms of Williams's (1969) generalizations about the ecology of colonization in the *Anolis* of the West Indies.

INTRODUCTION

With the publication of the work of MacArthur and Wilson (1963, 1967), a new methodology, the equilibrium analysis of colonization and extinction rates, revitalized one of the central topics of evolutionary biology. As always, issues involving methodology have corresponding theoretical issues. In biogeography the key issue is often seen as the relative importance of the role of history as opposed to any of those processes which may produce repeated patterns in community structure. This is a general problem in evolutionary ecology (an introduction to it is given in Kiester, 1980). In island biogeography the contrast is nicely exemplified by two books which take

opposite viewpoints for the most part. MacArthur (1972) attempts to develop a generally nonhistorical approach to biogeography, while Lack (1976) argues that the role of history is always paramount.

Some case studies in island biogeography may help to frame the outlines of this problem. In this paper I discuss the pattern of distribution of four species of skinks which occur on Arno Atoll in the Marshall Islands of Micronesia in the hope that analysis of these patterns may contribute to a further understanding of the problem of the role of history.

BACKGROUND AND RESULTS

Arno Atoll in the Marshall Islands, Micronesia, is in most respects a typical coral atoll. Located at about 7° 7' N and 171° 41' E, it consists of 133 islands of various sizes comprising a total of about five sq. miles of dry land. Many of the islets are no larger than emergent coral boulders while some are several miles long, and, in the case of Arno Island nearly one half mile wide. Arno has been described by Agassiz (1903) and by several articles in the Atoll Research Bulletin (Marshall, 1951; Stone, 1951; Wells, 1951; and Mason, 1952).

Six species of skinks occur on Arno Atoll. Two of them will not be considered further here for lack of data. One of these, *Lipinia noctua*, is nocturnal and the other, *Eugongylus albofasciolatus*, is cryptic, living under piles of debris and

¹Department of Biology, Tulane University, New Orleans, Louisiana 70118, U.S.A.

in houses. During my visit I saw no *Lipinia* and only one *Eugongylus*.

The other four species of skinks found on Arno are diurnal and are *Emoia cyanura*, *E. arnoensis*, *E. boettgeri orientalis*, and *Lamprolepis smaragdinum* (placed in the genus *Dasia* until recently). These four species form three distinct size classes with *E. cyanura* the smallest (40 mm snout-vent length [SVL]), *E. arnoensis* and *boettgeri* similar and of intermediate size (70–80 mm SVL), and *Lamprolepis* the largest (110 mm SVL).

As a result of a visit made to Arno in 1950 by J. T. Marshall and I. LaRivers, Brown and Marshall (1953) detailed an unusual distributional pattern for these four species. They found *E. cyanura* present on all islands visited. *Lamprolepis* and the two larger *Emoia* were only found on those islands large enough to support a substantial number of coconut palms. But of the 12 islands which contained either *E. arnoensis* or *E. boettgeri*, none had both species. This remarkable lack of overlap formed no recognizable geographical pattern within the atoll. Further, islets which were inhabited by different species were sometimes only several hundred meters apart. Brown and Marshall (1953) concluded that competition was the probable mechanism which kept these two sibling species allopatric.

In the summer of 1968 I visited Arno for the express purpose of obtaining more distributional data about the skinks, especially the two allopatric species. I censused skink species on 33 islands (about 25% of all of the islands) including 11 of the islands reported on by Brown and Marshall. Each island was searched for at least one rain-free hour. Because of the activity and conspicuousness of these diurnal species, it was usually possible to determine which species were present in a short time. Specimens were collected on seven of these islands and placed in the Museum of Comparative Zoology, Harvard University.

Figure 1 shows the distribution of *E. arnoensis* and *E. boettgeri* throughout the atoll. Table 1 gives the summary data for the four common diurnal species. Of 23 islands which had either *E. arnoensis* or *E. boettgeri*, 13 had *E. arnoensis*, 8 had *E. boettgeri*, and 2 had both species. Thus the basic pattern found by Brown and Marshall still holds, but the allopatry is not perfect; there are two exceptions. All of the islands which had been previously investigated were found to have the same species as before. Thus the pattern was stable for at least 18 years in these cases.

Table 1 gives the distribution of the four species over the 33 islands and the area in hectares of each island. Areas were estimated from the U.S. Navy Chart H.O. 6004 and are approximate. Overall, larger islands have more species. The Spearman rank correlation coefficient for area against number of species is 0.54, and the probability that this correlation does not differ from zero is $p < 0.002$.

Consideration of islands possessing one, two, or three species as detailed in Table 1 shows another pattern: islands which have only one species have *E. cyanura*, the smallest species. Four of the islands which have two species have *E. cyanura* and *Lamprolepis* (the smallest and the largest species) while a single two-species island had *E. cyanura* and *E. arnoensis*. There are then 20 islands with three species and two islands with four species. This pattern (with the one exception) of the smallest species in the simplest communities followed by the largest and then the medium-sized species in the successively more complex communities is reminiscent of some of the "assembly rules" of Diamond (1975).

Casual observations on the ecology of the four species revealed that *E. cyanura* was commonly active on the ground or on piles of dead vegetation. *Lamprolepis* was usually observed on the trunks of coconut palms, but would move onto the ground to forage. The other species of *Emoia* were also usually associated with

Figure 1. Distribution of *Emoia arnoensis* (A) and *E. boettgeri* (B) on Arno Atoll, Marshall Islands.

coconuts, but seemed to spend somewhat more time on the ground.

Two islands, Lojor Ngak and Bokale-Laj, which had only *E. cyanura*, were devoid of coconut palms. Their dominant vegetation consisted of two shrub species, *Scaevola* and *Messerschmidia*. These two genera are usually the earliest shrubs to invade a bare island (Stone, 1951).

Outside of Arno Atoll these four species are widely distributed throughout the Pacific (Brown, 1956). *E. cyanura* is found throughout the Caroline and Marshall Islands in Micronesia, and in much of Melanesia and almost all of

Polynesia, even reaching Clipperton Island in the western Pacific. *Lamprolepis smaragdinum* is found over all of Micronesia and Melanesia, and also New Guinea, Indonesia, and the Philippines. *E. boettgeri* is apparently found throughout the Marshall Islands and the eastern Caroline Islands. The distribution of *E. arnoensis* is less well known. Specimens in the United States National Museum are listed from Kusaie and one specimen in the California Academy of Science is reported from Truk. On either of these islands *arnoensis* would be sympatric with *boettgeri*. I have seen *boettgeri* common on several of the larger islands

TABLE 1. DISTRIBUTION OF THE FOUR SPECIES OF COMMON DIURNAL SKINKS ON THE ISLANDS OF ARNO ATOLL, MARSHALL ISLANDS.

Islands are listed in counterclockwise order from the northwest horn of Arno Atoll. Place names are from Mason (1952); those given in brackets are the names on U.S. Navy Chart H.O. 6004.* = an island also censused by Brown and Marshall (1953).

ISLAND	Number of Species	Previously Censused	Area (hectares)	<i>Emoia cyanura</i>	<i>Emoia arnoensis</i>	<i>Emoia boettgeri</i>	<i>Lamprolepis</i>
NAMWI [NAMI]	3		21.66	C	A		L
LOJOR NGAK	1		2.43	C			
MWET-DRIK [METORIKKU]	3		6.78	C	A		L
BIKAREIJ [PIKAAREJI]	3	*	104.34	C	A		L
BIKONELE	2		3.13	C			L
LIBEIJ-EN'	2		0.97	C			L
BOKALE-LAJ	1		0.63	C			
RAKIJEDR	2		0.80	C			L
BAR-BAR-EN'	2		5.46	C			L
ENEAITOK	2		4.31	C	A		
BOKANBIT	3		1.88	C	A		L
KIDRENEONG	1		2.09	C			
BOKANKORA	3		2.43	C		B	L
JAR-KWIL [CHARUKURU]	3		13.98	C		B	L
ENELIK, ENEAR [Now fused into one]	1		10.26	C			
TUTU [DODO]	3	*	29.17	C		B	L
KIJ-BWE [KEJIBOI]	4		24.58	C	A	B	L
JIL'ANG [CHIRAN]	3		16.44	C		B	L
ENIDRIK [ENIRIKKU]	4		12.27	C	A	B	L
TAK-LIB [TAGELIB]	3		11.65	C		B	L
TAK-LIB-EJ [EAST TAGELIB]	3	*	5.88	C	A		L
AK-TAL	3		1.63	C	A		L
NAJAJ	1		4.66	C			
TAE	3		4.66	C	A		L
L'ANGAR [RAKAARU]	3	*	88.00	C	A		L
KILANGE [KIRAGE]	3	*	69.33	C	A		L
MALEL [MARERI]	3	*	22.84	C	A		L
4th from East in L'OM'ARIN							
MALEL group	1		5.39	C			
IJOEN' [IJOEN]	3	*	80.14	C	A		L
ALJALTUEN' MATOL-EN'							
[CHITTAKINMATOROEN]	3	*	24.93	C	A		L
AOT-LE [AUTORE]	3	*	2.40	C		B	L
INE [INE]	3	*	373.26	C		B	L
ARNO [ARNO]	3	*	190.42	C		B	L

of the Truk group and on Ponape, but saw no *arnoensis* at either locality. *E. arnoensis* also apparently does not occur on Majuro Atoll in the Marshalls, the

closest island to Arno. I did not see it during my visit of three days. It is known from other atolls in the Marshalls, however.

DISCUSSION

Williams (1969) presented a list of ten preliminary generalizations on the ecology of colonization of small islands by lizards of the genus *Anolis*. Several of these claims are open to both historical and equilibrium interpretations and highlight the contrast between the different views of the role of history mentioned above. Although the data are not extensive, the case of the skinks of Arno Atoll may add to an understanding of the generality of some of the claims.

With regard to island size, Williams (1969) claims that "there appears to be some critical island size below which a colonizing species may exclude every other congener." On tiny islands *E. cyanura* does occur alone, but whether it actively excludes the other *Emoia* species and *Lamprolepis* is not known. *E. cyanura* may usually succeed in colonizing small islands first and then be able to prevent the successful colonization of other species. An alternative possibility is that other species are not found on the smallest islets because of a lack of suitable vegetation for perching and perhaps other ecological factors such as food supply and predation. On this view, history plays a dominant role in the sense that there may not have been enough time for suitable vegetation to appear. Were this the case, a species other than *E. cyanura* arriving at such an island would not succeed in colonizing whether *E. cyanura* was there or not.

However, colonization may proceed to the point where Williams (1969) claims that "even on small islands the phenomena of exclusion and coexistence appear to be relatively stable, once achieved." Data from Arno Atoll indicate stability over a period of at least 18 years for those islands which have either *E. arnoensis* or *E. boettgeri*, but not both. The two islands which possessed both species were not larger or more complex than those islands which had only one of the two. I do not know whether this co-

existence represents an equilibrium for these islands. Comparison with the other islands indicates that it is not. This generalization would predict that once one species excluded the other, then stability would occur.

Direct data on the colonization and extinction rates for the lizards of Micronesia is needed to definitely assess the validity of these two claims. This data has not been collected, but some aspects of the natural history of the area suggest that these processes may be both episodic and time dependent between episodes. The destructive typhoons which strike most atolls an average of two or three times a century (Wiens, 1962; see also Brown and Marshall, 1953, for discussion of the case of Arno Atoll) often remove almost all plant and animal life from an atoll island. Obviously this creates very episodic extinction. It may also tend to produce episodic colonization since typhoons may be the primary mechanism by which individual lizards are removed from islands to become rafters and possibly colonists. This pattern of extinction, far from contributing to the creation of an equilibrium species number, moves an island away from any equilibrium. Successful colonization by skinks cannot occur until the islands have been revegetated to some degree. The evidence presented here indicates that *E. cyanura* can move onto islands which have only the earliest plant colonizers (the shrubs *Scaevola* and *Messerschmidia*) whereas the other species may not be able to invade until the coconuts return. Thus the ontogeny of the vegetation of an island affects which species may invade at a given time. The question is: is there enough other colonization and extinction to create an equilibrium between episodes of typhoon-induced extinction and colonization? If the exclusion of *E. arnoensis* and *E. boettgeri* is due to some form of competition, then the possibility of an equilibrium can be considered.

Williams (1969) made two generalizations about coexistence which are rele-

vant to the zoogeography of these Pacific skinks. First, "Coexistence of colonists is possible if the colonizing species have been preadapted in sympatry on more complex source islands..." Islands on which coexistence occurs are not necessarily the source areas for islands with fewer species. The two small islands within Arno Atoll are not necessarily source islands even for the rest of the atoll, just as large islands such as Kusaie and Ponape are not necessarily source islands for the rest of the Marshall and Caroline Islands. For Micronesian skinks, it is not at all clear that there is any source area in the conventional sense. These skinks form a group which is more or less endemic to the Pacific islands (with the exception of *Lamprolepis*), and it is likely that almost all colonization events took place from one island to another rather than from an outside source area to the islands.

Second, "Coexistence may be evolved in situ in the special circumstances of complex geography..." This is the likely pattern for the coexistence (if the reports are accurate) of *E. arnoensis* and *E. boettgeri* on Kusaie and Truk which are large, complex volcanic islands. These islands certainly have enough ecological diversity to support a much larger lizard fauna than Arno Atoll.

In summary, Williams (1969) observes that "exclusion is an extremely common phenomenon but one of low visibility. Coexistence is a less frequent but highly visible phenomenon." Only in special circumstances where the kind of geographical arrangement found in an atoll which essentially constitutes multiple replications of a natural experiment can one expect to see the low visibility phenomenon of exclusion highlighted so well. It is the combination of the replication in space and continued censuses through time of the lizards of Arno Atoll and other islands of Micronesia which will enable us to determine the relative importance of equilibrium and historical factors in island lizard zoogeography.

ACKNOWLEDGMENTS

I wish to thank the Evolutionary Biology Committee of Harvard University, the Society of Sigma Xi, and Ernest E. Williams for support of the field work. I am grateful to A. E. Greer, E. E. Williams, and the late T. P. Webster for discussion on the zoogeography of Pacific lizards. I thank R. A. Lillehei for comments on the manuscript.

LITERATURE CITED

- AGASSIZ, L. 1903. Reports of the scientific results of the expedition to the tropical Pacific... IV, The coral reefs of the tropical Pacific. Mem. Mus. Comp. Zool., 28, xxxiii + 410.
- BROWN, W. 1956. The distribution of terrestrial reptiles in the islands of the Pacific basin. Proc. Eighth Pac. Sci. Cong., 3A: 1479-1491.
- BROWN, W., AND J. T. MARSHALL, JR. 1953. New scincoid lizards from the Marshall Islands, with notes on their distribution. Copeia, 1953: 201-207.
- DIAMOND, J. 1975. Assembly of species communities, pp. 342-444. In M. L. Cody and J. Diamond (eds.), Ecology and the Evolution of Communities. Cambridge, Harvard Univ. Press.
- KIESTER, A. R. 1980. Natural kinds, natural history and ecology. Synthese, 43: 331-342.
- LACK, D. 1976. Island Biology. Berkeley, Univ. of Calif. Press, xvi + 445 pp.
- MACARTHUR, R. 1972. Geographical Ecology. New York, Harper & Row, Publ., xviii + 269 pp.
- MACARTHUR, R. AND E. O. WILSON. 1963. An equilibrium theory of insular zoogeography. Evolution, 17: 373-387.
- AND ———. 1967. The Theory of Island Biogeography. Princeton, N.J., Princeton Univ. Press, 203 pp.
- MARSHALL, J. T., JR. 1951. Vertebrate ecology of Arno Atoll, Marshall Islands. Atoll Res. Bull. No. 3, pp. 1-38.
- MASON, L. 1952. Anthropology-geography study of Arno Atoll, Marshall Islands, Atoll Res. Bull. No. 10, pp. 1-21.
- STONE, E. L., JR. 1951. The soils, of Arno Atoll, Marshall Islands. Atoll Res. Bull. No. 5, pp. 1-56.
- WELLS, J. W. 1951. The coral reefs of Arno Atoll, Marshall Islands. Atoll Res. Bull. No. 9, pp. 1-14.
- WIENS, H. J. 1962. Atoll Environment and Ecology. New Haven, Yale Univ. Press, xxii + 532 pp.
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. Quart. Rev. Biol., 44: 345-389.

The Fossil Record and Early Distribution of Lizards

RICHARD ESTES¹

ABSTRACT. The fossil record of lizards is scanty, but some hypotheses about the early evolution and distribution of the group can be made. Four phases in lizard history may be identified. The first or eolacertilian phase is essentially Triassic, with the group progressively declining during the later Mesozoic. The second phase, during the Jurassic, involved the initial radiation of true lizards, or lacertilians, which probably underwent a primary dichotomy in the latter part of the Early Jurassic when Laurasia and Gondwana began to separate. This vicariance produced a Gondwanan iguanian group and a Laurasian group ancestral to all other lizards; each major group further diversified, both through plate separation and the influence of epicontinental seas or other intracontinental barriers. Some primitive living families may have evolved during this phase although as yet no fossil record documents their presence. In the third, or Cretaceous phase, the remaining Holocene lacertilian families appeared. Plate separation continued to be important in lizard diversification, and was instrumental in partitioning iguanids and agamids into subgroups, perhaps also formed gekkonid and scincid subgroups, and caused the agamid-chamaeleonid dichotomy. During the fourth or Cenozoic phase, expansion of family ranges took place during Paleogene peak tropicality. Cooler and drier climates in the Neogene caused restriction of lizard groups to their present distributions.

An evaluation of Renous's (1979) study on lizard biogeography shows that her cladogram, based essentially on limb nerve character states, is discordant with those based on more diverse data. Owen's (1976) continental reconstruction provides a better background for biogeographic movements of agamids, gekkonids, scincids, and varanids than does the conventional arrangement that includes a widely open eastern Tethys Sea and separation of southeast Asia from Gondwana.

INTRODUCTION

Lizards, as a group, have always provided outstanding opportunities for many kinds of studies. Many anatomical, physiological, ecological, and population studies utilizing these animals have been significant in formulation of new ideas regarding structure and function in reptiles. Lizard taxonomy has always been a popular field, since the diversity of lizards and the extensive modifications of their skeleton and external surface offer a natural laboratory for development of hypotheses of interrelationships and evolution of taxa, as well as providing data for use in biogeographical analysis.

One of the most outstanding examples of such integrated studies in lizards is to be found in the studies of Ernest E. Williams and his colleagues on the genus *Anolis*. Although I have not participated in this project directly myself, Professor Williams facilitated my own work on the lizard fossil record through his vivid interest in anatomy and the history of reptiles, and he has provided an important source of stimulation, information, and constructive criticism directly applicable to my work. It is a pleasure to acknowledge his help and to have the opportunity to contribute to this volume.

In this paper I discuss some aspects of my recent compilation of the lizard fossil record that appears in the *Handbuch der Paläoherpertologie* (Estes, 1982), a summary that I have often informally and somewhat ruefully called "The Telephone Book of Fossil Lizards." While

¹Department of Zoology, San Diego State University, San Diego, California 92182, U.S.A.

this work has provided both the tedious aspects of compilation as well as the stimulating advantages of a comprehensive view, it is the latter that I hope to provide here. In the *Handbuch*, each described fossil species is discussed, diagnoses are given for each genus and species if possible, and a provisional list of character states for each family and other higher categories is provided. The length of the *Handbuch* makes it desirable to give a summary of some aspects of it that will help paleontologists and herpetologists who are interested in what is known of lizard evolution and the fossil record, and who may then use the *Handbuch* itself for documentation.

The format I have adopted here is a narrative one; there is no opportunity to include more than a few of the more important data and bibliographic citations, which are given more fully in the *Handbuch*. Eldredge (1979) has argued that a scenario, to be classified as "good science", should be explicitly based on a well-established genealogy, and Rosen (1978) demonstrated that such a genealogy is essential for biogeographic studies. The cladogram of lizard relationships offered here (Fig. 1) is based on data given in the *Handbuch*, and for the most part is in accord with that generally accepted by herpetologists, at least as far as the primitive-derived branching sequence Iguania—Gekkota—Scincomorpha—Anguimorpha is concerned. However, there are still many differences of opinion regarding the branching sequence of families (or within families), and while this paper demonstrates some agreement between genealogy and vicariant events, it fails in other areas because a robust cladogram is not yet available for lizards. Estes and de Queiroz (in preparation) are providing this information based on more than 100 characters from both the skeletal system and soft anatomy.

THE FOSSIL RECORD OF LIZARD GROUPS

The most important features of the lizard fossil record are summarized below, preliminary to discussion of paleobiogeography of the groups. Marine varanoids have been omitted, as have Ophidia and Amphisbaenia. The latter two are for me separate Orders; the former may actually be convergent on

Figure 1. Suggested interrelationships of lizards and times of divergence. Thick black lines indicate extent of the known fossil record of each group. Dotted lines indicate extinction of the group. Cumulative m.y.b.p. given in column at left, which is in proportion to actual time involved.

varanoids (J. Gauthier, personal communication 1982). The lizards as a whole may be placed in a monophyletic Order Sauria of the Superorder Squamata and divided into two major groups, the probably paraphyletic Suborder Eolacertilia (eolacertilians) and the monophyletic Suborder Lacertilia (lacertilians or true lizards).

EARLY SAURIANS

The Eolacertilia first emerge in the Upper Permian and Lower Triassic of South Africa. These include small insectivorous diapsids that have been referred to the family Paliguanidae by Carroll (1975, 1977), including the genera *Paliguana*, *Palaeagama*, and *Saurosternon*. *Paliguana* itself appears to be an authentic saurian in having an external conch on the quadrate; the other two generally appear more primitive although they are probably lepidosaurian. In *Paliguana*, the first stages of temporal region modifications that are associated in living forms with a streptostylic quadrate appear. In the Upper Triassic, early saurians are found only in England and North America; these later forms, the Kuehneosauridae, are gliding animals (*Kuehneosaurus*, *Kuehneosuchus*, *Icarosaurus*) although other lizards in the English localities were apparently of more normal build (P. Robinson, personal communication). The kuehneosaurs are authentically saurian and had a more progressive streptostylic temporal region than the paliguanids (although the loose quadrate-pterygoid contact necessary for functional streptostyly was not yet present). Another Upper Triassic (possibly Lower Jurassic) form that may be related to the eolacertilians is *Fulengia* (Fulengidae) from southern Asia. It has leaf-shaped, polycusate teeth like those of *Iguana*, and if actually a member of the Sauria, suggests the development of

herbivory in the squamates even at this early date.

Evans (1980) has denied the saurian relationship of eolacertilians, proposing them as merely another group of eosuchians that lost the lower temporal bar, paralleling lizards, some spheodontids, and prolacertilians. Parsimony suggests, however, that this single character cannot be used to deny a relationship, in view of a number of other derived character states that kuehneosaurs, and at least *Paliguana*, share with squamates.

TRUE LIZARDS

The fossil record of the true lizards begins in the Late Jurassic and is thus separated from that of the Triassic eolacertilians by more than 50 million years, approximately the length of the entire Cenozoic. Extrapolation across the gap is thus not possible, and the lack of a fossil record of the primitive precursors of true lizards is unfortunate.

Iguania

The most primitive true lizards appear to be the Iguanidae, a group whose fossil record does not extend prior to the Late Cretaceous, presumably owing to the paucity of Gondwana localities. *Pristiguana*, from the late Cretaceous of Brazil, seems more primitive than the most primitive living iguanids (morunasaurines), although some similarities to teiids were also suggested (Estes and Price, 1973). The Late Paleocene of Brazil has produced a number of interesting undescribed fossil iguanids (Estes, in progress), which include both primitive and derived taxa, one of the latter reminiscent of the tropidurines. This suggests that all major extant iguanid groups were diversified by the beginning of the Cenozoic at least. North American records include several Eocene genera, among which *Parasauromalus* is a primi-

tive form with possible iguanine relationships, but there is no special similarity to *Sauromalus* as the name implies. *Aciprion*, from the Oligocene of North America, is a primitive iguanid that seems to be allied to the basiliscines. An undescribed, related North American Oligocene genus has a basilisklike occipital projection on the skull, helping to corroborate the basiliscine relationship of the group. In Mexico the Paleogene (Oligocene) *Paradipsosaurus* is another misnomer, since this primitive morunasaurine-like iguanid is unrelated to the living *Dipsosaurus*, although resembling it somewhat in having an inflated and shortened snout. Other Oligocene iguanids are known from Canada and represent sceloporines and possibly tropidurines.

North American fossil iguanids from Miocene through Pleistocene time for the most part represent living genera and occur generally in or near existing ranges. An interesting exception is the presence in the Miocene of Florida and Nebraska of the living genus *Leiocephalus*, now restricted to the West Indies.

Only a single taxon suggests the presence of iguanids in Europe. *Geiseltaliellus*, from the Eocene of Central Europe, seems to have been a small agile iguanid perhaps related to *Aciprion*, but the specimens need further study. Presence of iguanid-like forms in Asia is indicated by the Eocene *Arretosauridae* from Central Asia. *Arretosaurus* was a large lizard with a sculptured skull and mandibular character states like those of iguanids, but again further study of these specimens is necessary.

Turning to other iguanian families, the extinct *Euposauridae* from the Upper Jurassic of France seems to be iguanian but its position is still in question. There is a strong derived agamid similarity but euposaurs share derived states lacking in the most primitive living agamids.

Many agamid fossils occur within the

general limits of the family today: a primitive Late Cretaceous agamid *Mimeosaurus* was present in Central Asia; Eocene records referred to the primitive agamid *Tinosaurus* occur in southern Asia; *Agama* (or *Stellio*) is recorded from the Paleogene on into the Holocene in Europe. Other fossils indicate a wider Early Cenozoic dispersal of some agamids; *Tinosaurus* has also been recorded in the Eocene of North America and Europe, while *Uromastix* appears in the Paleogene of Europe.

The Chamaeleonidae have a very limited fossil record, but *Chamaeleo* is present in Central Europe during the Miocene. An unusual Miocene record of *Chamaeleo* from East Africa is known from a cast, in calcite, of the head region.

Gekkota

Some of the earliest true lizard fossils appear to be gekkotans. These animals are represented by complete skeletons and have been placed in the families *Ardeosauridae* and *Bavarisauridae* by Hoffstetter (1964). *Ardeosaurs* occur in the Upper Jurassic lithographic limestones of Germany (*Ardeosaurus*, *Eichstaettisaurus*) as well as in the Upper Jurassic of eastern Asia (*Yabeinosaurus*). *Ardeosaurus* is known from complete specimens and is quite gekkonid in general appearance. It and its close relative *Eichstaettisaurus* (represented by an exquisitely preserved skeleton) have procoelous vertebrae and complete supratemporal arches, the latter showing a derived, "hockey-stick" squamosal that lacks the dorsal process found in *Iguania* and the majority of the *Teiidae*. The bavarisaur *Bavarisaurus* and *Palaeolacerta* are less well preserved, but also appear to be gekkotan; they have derived amphicoelous vertebrae. Other gekkotan features of these families appear in many details of the skull, the fine, numerous teeth, and the configuration of the transverse processes of the first six or seven

caudal vertebrae. The two families appear to be early members of the Gekkota, with the same heterogeneity of vertebral type that occurs in living Gekkonidae. The skull roof of *Ardeosaurus* shows, however, a distinctive pattern of regular, enlarged epidermal scutes. This pattern may be primitive for lacertilians and lost in iguanians and Gekkonidae, or could be interpreted as a derived state of all lizards above the iguanian level but lost in gekkonids. Since pygopodids have such enlarged scales, and they may appear in some iguanids, as well as being present in amphisbaenians and snakes, I believe the first alternative is more parsimonious, indicating that granular scales are derived for both Iguanidae and Gekkonidae. Robinson (1967) suggested that presence of an "hockey-stick" squamosal in ardeosaurus and bavarisaurus is evidence of scincomorphan rather than gekkotan relationships, but the few living gekkonids that have retained remnants of both squamosal and supratemporal show the former to lie closely pressed against the latter and the parietal and appearing to be of "hockey-stick" type. Thus gekkotans may have lost the dorsal process independently of other lizards, if they arose before the scincomorphs as I suppose, since the dorsal process is still present in the vast majority of teiids. Alternatively, if gekkotans are scincomorph derivatives, their possession of an "hockey-stick" squamosal is to be expected.

Described fossils of the family Gekkonidae are all of Cenozoic age. An undescribed gekkonid is known in the Late Cretaceous of Central Asia. Fragmentary New World records occur in the Upper Paleocene of Brazil and the Upper Eocene of California. Cenozoic records are known from Europe (Eocene-Pliocene) and Morocco (Miocene), but these fossils are fragmentary and not distinctive. Cenozoic gekkonids indicate that the family was more or less cosmopolitan at the beginning of the Cenozoic and that

geckos occurred north of their present latitudinal limits in Europe.

Scincomorpha

The primitive scincomorphs (Lacertoidea), have a rather good fossil record that for the teiids goes back to the Late Cretaceous of North America and Central Asia, and Paleocene of South America; lacertids first appear in the Paleocene of Europe. Teiids are already represented by both teiine (*Meniscognathus*, *Penteteius*) and tupinambine (*Chamops*) tribes in the Late Cretaceous of North America. *Polyglyphanodon* is also present there at that time, and is the most highly derived member of the most primitive subfamily of teiids, the Polyglyphanodontinae (see Estes, 1982). The polyglyphanodontines are better represented in the Upper Cretaceous of Central Asia, where they had a brief radiation into a number of rather closely related forms, becoming extinct at the end of the Cretaceous. Rather large (*Iguana*-like) in general, most of them had enlarged molariform or polycusate-spatulate teeth, indicating that both omnivorous and herbivorous adaptations had evolved. No Cretaceous teiids are known in South America as yet (few localities are known), but they must have been there since the Late Paleocene Itaboraí deposits in Brazil have produced well differentiated members of both teiine and tupinambine lines that were similar to modern genera (Estes, in progress). Teiids disappear from northern latitudes in North America at the end of the Cretaceous with appearance of *Cnemidophorus* there in the Miocene. In South America, post-Paleocene records of teiids are essentially limited to the tupinambines *Tupinambis* and *Callopistes*. No fossils are yet known of the Gymnophthalmidae ("microteiids", see Estes, 1982).

The record of lacertids in the Late Paleocene of Europe is of a genus closely allied to *Lacerta* (*Plesirolacerta*), and in

the Middle Eocene *Eolacerta* appears; the latter is a large primitive lacertid with an open supratemporal fenestra. Other records of lacertids are primarily European and referable to *Lacerta* itself or to closely related forms. Lacertids may have occurred in the Late Jurassic of Europe (see below).

The skinks (Scincoidea) have a very poor fossil record, somewhat surprising in view of their present cosmopolitan distribution and great diversity. Some of the Late Jurassic cordylid lizards that I have included in the family Paramacellodidae (Estes, 1982, *Saurillodon*) show hints of scincoid relationship and one apparently had reduced limbs. There is, however, no compelling evidence at present to include them in the Scincoidea. Two Late Cretaceous genera from North America (*Sauriscus* and *Contogenys*) have tentatively been included in the Scincidae. The former is possibly a cordylid, but *Contogenys* is more definitively scincoid and has a *Eumeces*-like aspect, as well as showing some derived dental and mandibular similarities to primitive xantusiids (*Palaeoxantusia*). In the North American Oligocene, *Eumeces* itself appears. South America has produced an undescribed Paleocene form resembling *Contogenys* (Estes, in progress). No skink fossils have been found in Europe although a Miocene record of *Eumeces* from Morocco is known. No fossil skinks have been found in Asia.

The Cordyloidea as constituted here include a primitive Late Jurassic family Paramacellodidae and the living families Cordylidae and Xantusiidae. The paramacellodids occurred in England, Portugal and North America and include *Paramacellodus* and several related genera (Hoffstetter, 1967; Estes, 1982). These taxa show a number of cordylid features of the pelvis, caudal vertebrae and mandible, but according to Hoffstetter (1967) *Paramacellodus* had a primitive open supratemporal fenestra rather than the closed one of the living families. In my opinion, the supratempo-

ral region is not preserved in the specimens of this genus and the fenestral condition cannot be demonstrated. *Paramacellodus* had, however, a derived cordylidlike expanded coracoid region and well developed compound osteoscutes of cordylid type. It is possible that *Sauriscus*, a Late Cretaceous form mentioned above under the Scincoidea, is in fact a cordylid, but definitive character states are lacking.

True cordylid fossils are few, and are limited to the Late Eocene or Early Oligocene of France (*Pseudolacerta*) and the Miocene of Africa (*Gerrhosaurus*). The Xantusiidae, contrary to what is usually stated, has no demonstrable relationship with gekkotans (see Estes, 1982 for further discussion). There are a number of derived character states of the skull and mandible that suggest cordylid relationships, but a scincoid relationship is not falsifiable at present. Xantusiids are represented by fossils now referred to *Palaeoxantusia* from the Eocene of California, Wyoming, and Montana, and the Oligocene of Canada. Some of these fossils appear referable to *Xantusia*, and the only known skull shows some similarity to that of the Central American *Lepidophyma* and perhaps was near the ancestry of the latter.

Anguimorpha

The fossil record of anguimorphs is the best for any group of lizards, primarily because of their extensive, old, Northern Hemisphere distribution. The Xenosauridae has a North American fossil record extending from the Late Cretaceous into the Oligocene. If the Holocene Asian *Shinisaurus* is actually a xenosaur, the group originally had a wider distribution. The Dorsetisauridae is an Upper Jurassic family found in North America and Europe. It seems more derived than the xenosaurs in its predaceous dentition and anguid-like braincase, while its skull roof scutellation is derivable from the primitive scinco-

morph type. If *Dorsetisaurus* is authentically anguimorph, it suggests a split between the lines leading to Anguidae and Xenosauridae prior to the uppermost Jurassic. The Anguidae first appears in the Late Cretaceous of North America, as the common genus *Odaxosaurus*. This genus appears to have been near the ancestry of the glyptosauroids, which like the polyglyphanodontine teiids had enlarged teeth and were probably omnivorous. The group had an extensive Early Cenozoic North American radiation, forming two tribes, the primitive Melanosaurini (*Melanosaurus*, *Xestops*, *Peltosaurus*) and more derived Glyptosaurini (*Glyptosaurus*, *Paraglyptosaurus*, *Eoglyptosaurus*, *Heloderma*). The glyptosaurinid *Placosaurus* occurs in the Eocene and Oligocene of Europe; the melanosaurinid *Xestops* also occurs there in the Eocene. Fragmentary far northern records of both *Odaxosaurus*-like and glyptosauroid-like taxa occur in the Eocene of the Canadian Arctic Archipelago (Estes and Hutchison, 1980). Anguids of modern aspect are also present in the Late Cretaceous and Early Cenozoic of North America, where probable gerrhonotine fossils have been found, and the Diploglossinae, now West Indian, Mexican, Central and South American, appears in the Early Eocene of North America (Gauthier, 1982). The Anguinae is best represented by fossils in Europe, where *Ophisaurus* occurs from the Eocene on and *Anguis* appears in the Oligocene; *Ophisaurus* does not appear in North America until the Oligocene. The Anniellinae is now considered a subfamily of Anguidae (Bellairs, 1970) and is represented by the North American Eocene *Apodosauriscus* and Miocene-Holocene *Anniella* (Gauthier, 1980, 1982).

The Necrosauridae is an extinct family of primitive varanoid lizards. This group has a number of derived varanoid character states (most completely known for the Early Cenozoic European genus *Necrosaurus*), while retaining a more primitive

skull roof. A number of undescribed forms that may belong here occur in the Late Cretaceous of central Asia (Borsuk-Bialynicka, 1982). *Eosaniwa*, from the Eocene of central Europe, may also belong to this group; it is a large lizard with an elongated fish-catching snout unlike that of other necrosaurids. North American necrosaurids include *Parasaniwa* (Cretaceous-Eocene), and perhaps *Provaranosaurus* (Paleocene) and *Colpodontosaurus* (Late Cretaceous). Necrosaurids all had a predaceous dentition and so far as known lacked retraction of the naris; they appear, however, to have had at least a partially developed intramandibular jaw hinge like that of *Heloderma*.

The Helodermatidae may be present in the North American Cretaceous (*Paraderma*) and without doubt appears in North America by the Oligocene and Miocene (*Heloderma*); an helodermatid has also been found in the Late Eocene or Early Oligocene of France (*Eurheloderma*).

The Varanidae had a wide distribution in the Late Cretaceous and Early Cenozoic. Varanids are known from the Late Cretaceous of Central Asia (Borsuk-Bialynicka, 1982) and the North American varanoid *Palaeosaniwa* may be a varanid; *Varanus*-like forms occur in the Eocene of North America and Europe (*Saniwa*). A definitive Early Eocene record of varanids occurs in the Canadian Arctic Archipelago (Estes and Hutchison, 1980). *Saniwa* extends into the Oligocene of North America, and other varanids persist until the Pleistocene in Europe (*Iberosvaranus* and *Varanus*). A Late Cretaceous relative of the extant *Lanthanotus* occurs in central Asia (Borsuk-Bialynicka, 1982). The relationship of *Lanthanotus* to the marine varanoids suggested by McDowell and Bogert (1954) has not been generally accepted (Rieppel, 1980a), and both Estes (1982) and Gauthier (personal communication 1981) include *Lanthanotus* in the Varanidae. Rieppel (1980b) maintained a

separate family for the genus, but his comparisons included only *Lanthanotus*, *Heloderma*, and *Varanus*. Since Rieppel lacked a more extensive generic series, and in view of the many derived characters shared by *Lanthanotus* and *Varanus*, I believe that it is reasonable to place *Lanthanotus* in the Varanidae as a subfamily.

LIZARD PALEOBIOGEOGRAPHY

The summary of the lizard fossil record presented above and derived from Estes (1982) now makes it possible to offer some preliminary suggestions on the biogeographic history of the different families of lizards. Lillegraven *et al.* (1979) have emphasized the tenuous data on which mammalian Mesozoic biogeography is based, and have observed that this lack of data has not hampered the flow of supposition; indeed, as A. S. Romer often observed, "increase in data leads to triumphant loss of clarity." I am well aware that the present paper is based on data equally tenuous as those for mammals so far as the Mesozoic is concerned, and much more so for the Cenozoic. Nevertheless, since the *Handbuch* (Estes, 1982) represents the first codification of data regarding all fossil lizards, some discussion of what it suggests is desirable. No previous detailed discussion of this sort has been based on the fossil record, although a recent study of lizard biogeography by Renou (1979) has made use of continental plate reconstructions. The zoogeographic models discussed here are consistent with the generally accepted movements of continental plates and also provide some corroboration for the position of the Southeast Asian region suggested by Owen (1976, 1981, 1982).

The diagram of continental plates used here in Figures 3–13 is adapted from the Early Cretaceous map in Smith and Bryden (1977). Continental masses are highly diagrammatic, and I have modified the map to reduce the polar distor-

tion of continents, but the relative distance between points of continental plate contact or proximity are kept approximately the same as in Smith and Bryden. The map has utility, however, principally as a base map to be used in the context of this paper; the maps of Smith and Bryden (1977), Owen (1976), Lillegraven *et al.* (1979) and Howarth (1981) should also be consulted. I have chosen the Early Cretaceous map as a base because continental separation first becomes marked at that time, yet the original Pangaeian configuration is still apparent. The Early Cretaceous was in addition an important time in lizard evolution and distribution. The known Jurassic record consists entirely of extinct families, although they adumbrate extant groups to one degree or another. Since most extant families were in existence by at least the Late Cretaceous, the importance of Early Cretaceous time is clear. It should be emphasized, however, that data from times other than the early Cretaceous appear on the maps, and the figure legends and text should be consulted for explanation.

The account of lizard fossils above demonstrates how scanty our record is, as well as how few fossils we have from Gondwana. It is not possible to develop an adequate interpretation of lizard biogeography without this southern record, and a variety of equally plausible hypotheses can be constructed with the existing data. I have chosen the interpretation that seems most parsimonious to me when the cladogram of lizard relationships, the fossil record, and the present distribution of lizards is confronted with data on plate movements, but my account will have to be modified as more information is obtained from the South American and African Jurassic and Cretaceous, the Cretaceous of Europe, and of course any Mesozoic and Cenozoic data from Asia.

PERMIAN-TRIASSIC

The Permian and Triassic record shows only that up to the Early Triassic,

the Squamata was Gondwanan in distribution. The group is not yet recorded from South America but was presumably present there as well. By the Late Triassic the more derived kuehneosaurids appear in Laurasia, and an herbivore, *Fulengia*, occurs in eastern Asia (Fig. 2). Although only a few localities are known, the variety of lizards in the English Triassic fissures (mostly still undescribed) suggests that by the end of the Triassic there was some diversity in Laurasia as well. Cox (1974) has shown that a "world fauna" existed during the Triassic since there were few barriers to animal movement during that time. These data, poor as they are, suggest that lacertilians were Pangaeian in the Triassic, and that the various groups originated as vicariants following the breakup of Pangaea, either by plate separation or through separation by intracontinental barriers.

JURASSIC

The single Lower Jurassic fossil that has been referred to the lizards is the poorly preserved *Protolacerta*, from Argentina. It is not demonstrably a lizard; part of the specimen is from a fish and part from an indeterminate reptile (J. Bonaparte, personal communication 1981).

In the Late Jurassic, the presence of *Cteniogenys* in North America and Europe indicates a significant extension of the geological range of the eolacertilians, already known from Laurasia in the Late Triassic. The most significant aspect of Jurassic fossil lizards, however, is that all the major groups of extant true lizards (Superfamilies of Romer, 1956; Infraorders of Estes, 1982) are represented in the Late Jurassic of a relatively small area—western Europe—and that the groups show very little mixture in localities (Table 1, Fig. 3). Greatest similarity occurs among the English, Wyoming, and Portuguese localities, all of which contain Paramacellodidae and Dorset-

isauridae. *Cteniogenys*, the eolacertilian, is lacking in the English localities, but occurs in Portugal as well as in Wyoming. Primitive scincomorphs (lacertoids?) occur at Lerida, Spain, and the presumed lacertoid *Durotrigia* overlaps with the paramacellodids in England. This series of localities has thus yielded *Cteniogenys*, scincomorphs, and primitive anguimorphs. In the Cerin lithographic limestone in France, only the iguanian euposaurs occur, and in the Solnhofen lithographic limestone in Germany only the ardeosaurs and bavarisaurids are found; an ardeosaur also occurs at the same time in Eastern Asia. The Chinese and German ardeosaurs suggest that the gekkotan diversification was Eurasian. The paramacellodid—dorsetisaur—eolacertilian assemblage suggests that some scincomorph and anguimorph groups were originally Euro-North American. Although ecological and/or sedimentary environment differences may be involved, some barriers to movement seem to have been present (see Howarth, 1981), and there is some geological evidence for this hypothesis. The major extent of European epicontinental seas at the approximate time of these Late Jurassic localities was in Central Europe, extending south through France (Ager, 1975; Hallam and Sellwood, 1976). There was an emergent area in northern Europe, north of the Solnhofen portion of this seaway, where the ardeosaurs and bavarisaurids must have been living, their remains being incorporated into the back-reef lagoons that later became the lithographic limestone (Hallam, 1975). A similar situation occurred at the Cerin locality in France, with an emergent region to the southeast (Delfaud, 1980) being a source for the euposaurs. The Purbeck and Portuguese localities, on the other hand, were associated with an emergent region to the west, more closely linked with North America and Greenland, as indicated both by the plate tectonic evidence as well as that of the fossils. Details of the oscillations of these

Figure 2. Known distribution of Triassic Sauria. Base map for Pangaea in the Late Triassic (200 m.y.b.p.) from Smith and Bryden (1977).

Abbreviations: AF = Africa; AN = Antarctica; AU = Australia; EA = Eurasia; IN = India; NA = North America; NZ = New Zealand; SEA = Southeast Asia; Greenland and Madagascar not marked.

epicontinental seas are not entirely clear, but the archipelagic area formed in western Europe could have permitted vicariance of these and perhaps other lizard groups. For the most part these seas disappeared in the latest part of the Jurassic, when most of Europe (and many other parts of the world) became emergent and provided at least a brief connection with North America before Cretaceous epicontinental seas isolated the two areas by inundating most of Europe once again (Prothero and Estes, 1980; Hallam, 1981).

With the Triassic and Jurassic record in mind, we may now proceed to discussion of the biogeography of Cretaceous and

later lizards and their relationship to the Jurassic forms.

IGUANIAN DISTRIBUTION AND THE MODEL OF RENOUS

A recent review of lizard biogeography by Renous (1979) was based mainly on structures and distributions of living groups. Although she did consider past continental movements, her discussion of lizard fossils ignored many recent contributions in this area. Renous emphasized the Iguania in her discussion, and formulated hypotheses of iguanian (and other lizard) relationships and distribution, based principally on her extensive

Figure 3. Known distribution of Jurassic Lacertilia. Base map for Early Cretaceous (120 m.y.b.p.) from Smith and Bryden (1977). ▲ = paramacellodids, dorsetisaurs, and possible lacertoid *Durotrigia* (England); ● = scincos-morphs (possible lacertoids) *Meyasaurus* and *Ilaerdesaurus* (Spain); ★ = paramacellodids, dorsetisaurs and possible scincoid *Saurillodon* (Portugal); △ = *Paramacellodus*, *Dorsetisaurus* (Wyoming); □ = *Euposaurus* (France); ⊙ = ardeosaurs *Ardeosaurus* and *Eichstaettisaurus*, bavarisaurs *Bavarisaurus* and *Palaeolacerta* (Germany); ⊙ = ardeosaur *Yabeinosaurus* (People's Republic of China). Continents identified in Figure 1. Note large eastern Tethys Sea; see text and compare continental reconstruction in Figure 14.

studies of the limb innervation (Renous, 1978, 1979, 1981; the latter contains little new information). In the lizard forelimb, the course of the ulnar nerve is superficial to the muscles as in most tetrapods (primitive or *lacertide* condition of Renous = L). In some lizards, however, the nerve takes a deep course (derived or *varanide* condition = V). In the hind limb, the dorsal muscles are generally innervated in tetrapods by the peroneal nerve (primitive condition = A), but uniquely in some lizards the peroneal nerve is absent and its function taken over by the interosseous nerve (derived condition = B). The polarity of these

character states seems well established. Derived conditions are as follows: B occurs in most iguanids, all agamids, some cordylids, and *Heloderma*; V occurs in varanids and Afroasian lacertids; V and B together are found in Malagasy iguanids, chamaeleonids, teiids and gymnophthalmids. All other groups have the primitive states L and A. From the developmental standpoint the difference between the *lacertide* and *varanide* conditions of the forelimb does not appear to be fundamental. Since nerves grow, in development, toward the muscle blastemata, slight differences in timing or conditions in the mesenchyme could lead to adop-

TABLE 1. DISTRIBUTION OF KNOWN LATE JURASSIC LIZARDS BY LOCALITY, SHOWING THAT VERY LITTLE OVERLAP IN INFRAORDINAL GROUPINGS OCCURS.

	<i>Ctenogenys</i>	paramacellodids	dorsetisaur	<i>Durotrigia</i>	<i>Meyasaurus</i>	<i>Illaerdesaurus</i>	euposaur	ardeosaur	bavarisaur
Purbeck, England		X	X	X					
Guimarota, Portugal	X	X	X						
Quarry 9, Wyoming	X	X	X						
Lérida, Spain					X	X			
Cerin, France							X		
Solnhofen, Germany								X	X
People's Republic of China								X	

tion of a different pathway; such characters are readily subject to both parallelism and reversal. The difference between the A and B states is less ambiguous since a loss is involved; here again, however, the phylogenetic significance is doubtful since multiple loss can occur. Reversal is possible if the "loss" were to be the result of epigenetic processes not accompanied by loss of potentiality (see Alberch *et al.*, 1979).

Renous (1979) also discussed character states of the dentition, forearm and shoulder musculature, scapulocoracoid fenestrations, humeral foramina, form of the carpus, cervical vertebrae, and femoral-preanal pores. Most of these character states show extensive parallelism; some are correlated, and they will not be discussed further here owing to space restrictions. In any case, Renous' cladistic models for iguanian relationships were based only on the nerve pathways and the presence of an acrodont dentition. She then used these results in biogeographic analysis, an analysis that is

generally unsuccessful because homoplasy (which occurs in most characters) has a greater effect on a cladogram when only a few characters are used; her cladistic models are discordant, in general, with those based on more extensive data. Her view (1979:395) is stated unambiguously in the original as well as in my translation:

Within the classification of lizards, all these characters [i.e., the nerve pathways] do not appear to us to be distributed by chance, but to obey a certain logic, a certain rationality that cannot be other than the phylogenetic history of the group. The distribution of these characters reflects phylogenetic relationships.

It is possible to accept that in some cases the distribution of the nerve pathways indicates phylogenetic relationships, but if all possible cladistic arrangements of these characters are diagrammed it is clear that homoplasy must have occurred, since all possible combinations of the two character states occur in living lizards (i.e., LA, LB, VA, VB). Since B is a state defined by loss, it seems less likely that A has been obtained by reversal, although as we shall see, this possibility has to be considered in some situations.

Several separate origins of V are demonstrated by its occurrence in some iguanians, Afroasian lacertids, teiids and gymnophthalmids, and the varanids, occurrences discordant with the extensive data that indicates lack of relationship between the iguanians, the varanids, and the other groups. The situation with B is more complicated. It is widespread in iguanids, but A is present in iguanines, while the sceloporines and tropidurines may be either A or B. Since these three groups are quite homogenous in most other derived character states, it is possible that the absence of the peroneal nerve in iguanids is the result of genetic suppression without loss of potentiality. Alternatively one must argue for numerous independent losses of the nerve within iguanids, and this is possible yet much less parsimonious. Jullien and

Renous (1972) gave data for morunasaur and *Crotaphytus*, and these primitive iguanids also have both L and B (confirmed by K. de Queiroz, personal communication 1980), suggesting that reversal in iguanines, some of the tropidurines, and some of the sceloporines is the most parsimonious alternative. Separate developments of B have occurred in *Heloderma*, and in some cordylids as well (in the latter group Renous noted, [1979: 394] that the conditions of B are not "perfectly attained"), and in both the teiids and gymnophthalmids. In the latter two groups the presence of both derived conditions V and B may be phylogenetically significant.

Renous (1978, 1979) was particularly concerned with the relationship between the Malagasy iguanids, the agamids, and the chamaeleonids, and emphasized in this regard the common presence of derived conditions V and B in the oplurines and chamaeleonids, but as noted above this combination occurs also in the teiids and gymnophthalmids. She suggested that the agamids, in retaining the primitive condition L with the derived condition B, are more closely related to the New World iguanids, while oplurines gave rise to the chamaeleonids, the latter two groups later invading Madagascar. In her view, the agamids underwent vicariance on the Indian block, only secondarily invading Africa. This hypothesis required her to postulate a parallel origin of acrodonty in both agamids and chamaeleonids. On the other hand, Moody (1980), who has reevaluated the relationships and biogeography of agamids, indicated that determination of relationships among the iguanids, chamaeleonids, and agamids requires further study owing to the fact that chamaeleonids share approximately the same number of derived states with both agamids and Malagasy iguanids (oplurines), and he hypothesized that various agamid groups had centers of origin on Asian, Southeast Asian, and Australian plates, following the continental recon-

struction of Owen (1976). Moody's detailed analysis is in agreement with Renous' conclusion that Africa secondarily received its agamids, although he brings them from Asia rather than from India. Moody also suggested that Australia and Southeast Asia received agamids simultaneously; he emphasized the closer relationship between Southeast Asia and Australia required for this interpretation that can be found in Owen's (1976) reconstruction but not in the more usual continental arrangements (e.g., Smith and Bryden, 1977, and see Fig. 14). Owen's interpretation will be discussed further below.

My own view of early lacertilian evolution is that a Jurassic ancestral morunasaur-like population of iguanoids (essentially at iguanid level) occupied at least the southern portions of the South American-African region of Gondwana (Fig. 4). This iguanoid group was the result of vicariance following separation of Pangaea into two more or less distinct areas, Gondwana and Laurasia; this basal lacertilian dichotomy left the Laurasian lizards to develop into the more derived groups that will be treated in the following section. The iguanid group would have undergone further vicariance when South America and Africa separated (a process initiated in the very latest Jurassic, primarily accomplished in the Early Cretaceous and completed by about Middle Cretaceous, see Tarling, 1980), separating the South American and African forms (Fig. 5). Within the African region, some of the ancestral morunasaur-like forms (which were surely in the Madagascar region before separation of the latter from Africa) became oplurines, and developed condition V in parallel with chamaeleonids (recall that this is simply a different, deeper path for the ulnar nerve, and that V has arisen independently in other lizard groups).

In the South American region, it is possible that *Pristiguana* from the Late Cretaceous of Brazil was a representative of the early morunasaur group that di-

Figure 4. Suggested areas of origin (stippled) in primitive Iguania; see Figure 3 for map data and see text for further explanation. Only initial separation into groups shown; subgroup differentiation shown in Figure 5.

versified into the non-oplurine iguanids. North American iguanids seem to have been the product of rather limited dispersals from South America (Savage, 1966), probably beginning in the Late Cretaceous and extending on into the Early Cenozoic. These movements did not involve vicariance but resulted from chance dispersal across a series of volcanic islands that arose during the Late Cretaceous between North and South America owing to collision of a Pacific aseismic ridge with this island arc system (Schmidt-Effing, 1979), or perhaps transient upwarps produced by heat of plate movements (H. Owen personal communication 1980). Rage (1978, 1981) has suggested that a number of snake genera dispersed across this region, teiids also took this path (see below), as did some hadrosaurine dinosaurs

(Casamiquela, 1980) and mammals (Ferrusquia-Villafranca, 1978; Lillegraven *et al.*, 1979; Hallam, 1981). The North American Late Cretaceous record, however, shows no definitive record of iguanids, which are first recorded by *Paradipsosaurus* in the Paleogene of Mexico and *Parasaurodon* in the Early Eocene of Wyoming; both genera appear to be relatively primitive morunasaur-like forms. The Oligocene *Aciprion* and an undescribed crested relative (see above) apparently represent more northern representatives of basiliscines. The sceloporines probably came from limited dispersal of members of a common ancestor with tropidurines. While the latter diversified in South America and the West Indies, sceloporines exploited the more open, arid habitats in southwestern North America, and only a

single tropidurine, the presently West Indian genus *Leiocephalus*, ever reached North America (Miocene). Iguanines may have been derived from primitive iguanids that also reached Central America from the south. Finally, a single iguanid appears to have reached Europe; the Middle Eocene *Geiseltaliellus* dispersed across the North Atlantic route at the same time as the agamid *Tinosaurus*, glyptosaurine anguids, the varanid *Saniwa*, and a number of mammals and other lower vertebrates during a period of peak tropicality in the Eocene.

Part of the ancestral morunasaurlike iguanid group in the African region may have been separated by some ecological or physical barrier in the northern or northeastern portion, developed acrodonty, and thus became an ancestral group for the agamids and the chamaeleonids. This group would probably have had nerve conditions L and B as do iguanids and agamids today. Following Moody (1980), agamids then originated from portions of this population that underwent vicariance on the Asian, Southeast Asian, Indian, and Australian plates following separation of Africa from eastern Gondwana in the Early Cretaceous (Fig. 5). Southeast Asia and eastern Asia have localities that yield a Triassic *Glossopteris*—*Lystrosaurus* biota (Colbert, 1973), suggesting an association of these areas with Gondwana, an association corroborated by the continental reconstruction of Owen (1976, 1981, 1982, see below). The primitive agamid stock is represented in the Late Cretaceous of Central Asia by *Mimeosaurus*, and later, in the Eocene, by the primitive agamid *Tinosaurus*, which has been recorded from the Paleocene and Eocene of eastern Asia and the Eocene of Europe and North America as well. All records of *Tinosaurus* are fragmentary, and while the Asian forms are primitive agamids the generic reference is in doubt. The North American and European specimens are very similar, however, and appear to be congeneric; corroboration for this is the

presence of Eocene lizard, turtle, champsosaur, crocodilian, and mammal genera common to both these areas, suggesting dispersal across the North Atlantic corridor during Eocene peak tropicality. While the Asian fossil record is very scanty, no data suggest that this dispersal was across the Bering Route, although this is a possibility (Fig. 5).

Portions of the ancestral L-B-acrodont population of iguanoids remaining in Africa after the vicariance of agamids developed nerve condition V and gave rise to the chamaeleonids, which by the Late Cretaceous invaded Madagascar (or perhaps had been there prior to its separation) and in the Cenozoic reached Eurasia.

The Late Jurassic Euposauridae in France may have been a more highly derived branch of the common acrodont ancestor of agamids and chamaeleonids, a group that moved northward, briefly, into Laurasia when agamid ancestors spread to Asia and Southeast Asia. This would explain their general similarity to agamids, their retention of certain primitive conditions such as paired bones of the skull roof, and their development of some derived conditions not found in the most primitive living agamids.

In brief summary, then, a southwestern Gondwanan (South America—Africa—Madagascar) primitive iguanian group diversified during Middle-Late Jurassic time after separation from Laurasian lizard groups. Neotropical iguanid groups and oplurines developed by vicariance as South America and Africa diverged (a process completed by the Middle Cretaceous). A more northern Gondwanan acrodont iguanian group underwent vicariance as the Asia-Southeast Asia-Australia-India blocks separated from Africa. The agamids and chamaeleonids resulted from this plate separation. Much later, following their Mesozoic subgroup differentiation, primitive agamids spread widely in the Eocene, occurring in North America, Europe and Asia; sympatry with iguanids in North America, at least,

Figure 5. Suggested areas of origin (stippled) and later dispersals (arrows) in the families of iguanians. Iguanid subgroups formed after vicariant separation of South America and Africa; oplurines relict in Madagascar; Late Cretaceous or Early Cenozoic dispersal of iguanids to North America and in the Eocene, apparently, to Europe. Agamid subgroups formed by vicariance following separation of Southeast Asia, Australia and India; dispersal of agamids into Europe and North America in the Eocene (North American route chosen here; possible Bering route shown as dotted line). Chamaeleonids disperse to Europe in Miocene; to India after collision with Asia (not shown). Map data as in Figure 3. Dates in or at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. See text for further explanation.

confirms this as dispersal. Some Asian agamids also invaded Africa and India in the Cenozoic in a limited way (*Agama*, *Uromastix*, *Leiolepis*). *Agama* and *Uromastix* reached Europe no later than the Late Eocene or Early Oligocene, either from Asia Minor or Africa, with *Agama* persisting in Europe until the Early Pleistocene (it still occurs there peripherally). It is likely that the chamaeleonids reached India as a result of range extension in the Early or Middle Cenozoic, after collision of India with Asia. While there is no present record of such an extension during Eocene tropicality, the group had reached Central Europe by

the Miocene return of tropical conditions (see e.g., Cracraft, 1973a). Cracraft (1973b, 1975) suggested that agamids and chamaeleonids diverged after separation of South America and Africa in the Middle Cretaceous; it is likely to have been earlier than that time.

A major unanswered question in iguanid distribution is the absence of iguanids in Africa today. If a few have survived in Madagascar and the major evolution of the group was South American, what happened to the intervening African populations? There is no obvious answer, but it is possible that iguanids may have been reduced by

shifting climatic belts (as discussed by Robinson, 1973), perhaps in combination with other, as yet unknown factors.

POST-IGUANIAN LIZARD DISTRIBUTIONS

Gekkota. Renous (1979) suggested Europe as a center of origin for gekkotans. While the apparently gekkotan ardeosaurs and bavarisaurus occur in the Upper Jurassic of northern Europe, the presence of an ardeosaur in eastern Asia indicates a wider distribution (Fig. 3). Since the localities with ardeosaurs contain no trace of the contemporary radiation of paramacellodids, lacertoids and euposaurs, gekkotans may have originated on a portion of Europe and Asia that was at that time not connected with southwestern Europe. During the Late Jurassic, Eurasia began to be bisected by the so-called Obik Sea, which separated the European region and isolated Siberia and eastern Asia from the rest of the world (Cox, 1980). Although the Obik Sea (or Turgai Straits) separated Europe from Asia completely by Cretaceous time, the initiation of this seaway from the north during the Late Jurassic would have effectively separated the northern European ardeosaurs and bavarisaurus from the Asian ones that are more likely to have been the actual gekkonid ancestors from the geographic standpoint; an Asian origin of gekkonids is supported by their present distribution. Kluge (1967) suggested that eublepharine gekkonids gave rise to the now wholly Australian diplodactylines and the gekkonines in Asia, with the latter spreading westward to India, Africa, and Madagascar. South American gekkonines are low in diversity; they and the sphaerodactylines appear to be derivatives of highly dispersable cosmopolitan genera (see Bons and Pasteur, 1977). Whether they reached the New World by Cretaceous predrift distribution or postdrift rafting is unknown, but their low present diversity suggests that rafting is the more likely possibility. Tarling (1980) suggested that South

Atlantic oceanic currents would have fostered dispersal from Africa to South America but not the reverse, and further pointed out that a series of large oceanic islands may have been available along the Ceará-Sierra Leone Rises up to the end of the Eocene. The only South American fossil gekkos are from the Paleocene of Brazil; they have not yet been identified to subgroup, but give a minimum date for emplacement of gekkonids on that continent. Eublepharines are likely to have reached North America from either Asia or Europe in the Early Cenozoic, rather than in the Late Cretaceous when the Bering connection first became available, since there are no gekko fossils in the relatively extensive Late Cretaceous deposits of North America and they were present in Central Asia at the time (unpublished material being studied in Warsaw). An Eocene gekko occurred in California but cannot be identified to subgroup; this animal as well as the Eocene-Pliocene European gekkos probably dispersed during Eocene and Miocene peaks of tropicality (see e.g., Cracraft, 1973a).

The distribution pattern for gekkos is thus very similar to that for agamids as described by Moody (1980), with primitive Southeast Asian forms, a well-defined Australian radiation, and several endemic genera in India, dispersing later into Africa and Madagascar (Fig. 6). Both gekkos and agamids are found in the Late Cretaceous of Central Asia and the Eocene of North America and Europe. Again, however, lack of fossils from many areas of present gekko distribution makes analysis of past movements difficult. The pygopodids lack a fossil record, but if derived from diplodactylines in Australia as Kluge supposes (1981, personal communication), must have lost the type C double visual cells present in all gekkonid subfamilies. Alternatively as G. Underwood believes (personal communication 1981) they may have branched off prior to gekkonid acquisition of these cells; this would fit with their retention of the large,

Figure 6. Suggested area of origin (stippled) and later dispersals (arrows) in geckos. Map data as in Figure 3. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. Possible Bering route to North America for Eocene geckos shown in dotted line, see text. Note vicariance of gekkonid subgroups in Southeast Asia, Australia, and India. Only major paths shown; island or rafting dispersal routes not included. See text for further explanation.

regular dorsal head scales. Dibamids (including anelytropsids) seem to be related to the gekkotans (Underwood, 1971) and while they also lack a fossil record, again point to an originally Laurasian deployment of the Gekkota.

Scincomorpha. While the iguanoids were radiating in western Gondwana, scincomorphs were diversifying in Laurasia, as shown by the presence of cordyloids and perhaps scincoids in the Late Jurassic of Europe. Definitive lacertoids (the most primitive of scincomorphs) are not known at this time, although the presence of the more advanced cordyloid group indicates that lacertoids must have already evolved, and ancestral lacertoids may have split off from a common ancestor with iguan-

ians in Eurasia (Fig. 7). *Meyasaurus* and *Ilaerdesaurus* from the Late Jurassic of Spain, and *Durotrigia* from the Late Jurassic of England, may represent early lacertoids (Fig. 3).

The unity of living lacertoids seems to be demonstrated by a number of derived character states (Hoffstetter, 1962; Rieppel, 1980c; Estes, 1982; Gauthier, in preparation), although Presch (1981, personal communication) believes lacertids and teiids to be widely separated. Presch also allies the gymnophthalmids with the lacertids, but teiids and gymnophthalmids share derived character states of the hemipenial musculature (N. Arnold, 1982, personal communication), and the presence of derived nerve conditions V and B; I thus continue to ally

teiids and gymnophthalmids while awaiting the results of the studies of Presch and Gauthier.

Cordyloids (and perhaps scincoids) are present in the Late Jurassic, and as noted above, lacertoids were present in western Laurasia at least; one possible hypothesis is that the teiid-lacertid dichotomy arose by separation of a part of the ancestral population on one of the large continental islands in the European region during the late Jurassic. This vicariance could then have produced the lacertids, while the teiids developed in North America, where they are extensively represented by the Late Cretaceous, including teiine, tupinambine, and polyglyphanodontine groups. A relatively restricted radiation of polyglyphanodontines was also present in eastern Asia at this time. The polyglyphanodontines were a group that capitalized on relatively large size and herbivorous-omnivorous habits, and while they are a primitive group of teiids their divergent dietary specializations preclude them from ancestry to teiids as a whole. This hypothesis presumes that teiids reached South America from the north. While this is against the ocean current pattern that Tarling (1980) supposes would have fostered only south to north movement, such groups as hadrosaurian dinosaurs clearly must also have come from the north.

So far as lacertids are concerned, much of Europe was submerged for the major part of Cretaceous time, so that extensive habitat for lizard groups was not present. Eurasian lacertids appear to be the most primitive group, and the more derived African forms (with some later reentrants into Eurasia) may have developed from isolation on northwest Africa in the Late Cretaceous, when that region was separate from both Europe and the rest of Africa.

An alternative hypothesis for the lacertid-teiid dichotomy would be to postulate the presence in western Gondwanaland of an ancestral lacertoid population (of

which *Durotrigia* would then, like the euposaurs, be a Laurasian extension) that underwent vicariance during the Early-Middle Cretaceous separation of Africa and South America, producing lacertids in the former, teiids in the latter. Since teiines and tupinambines are known from the Late Paleocene of Brazil, they must have thus had a long history in South America, although since the Cretaceous record in South America is poor we do not know if they were present there at that time. Estes (1970) favored this hypothesis.

In this second hypothesis, the lacertids would be primarily African, but separation of the more primitive, now Eurasian forms from the more derived African forms need not be much different than in the first hypothesis.

Neither of the above alternatives prohibits the gymnophthalmids from having been endemic to the South American region as postulated by Presch (1980). In the first alternative the gymnophthalmid stock could have been the result of extension of a teiid stock into South America from the north during the (Late?) Cretaceous, with divergence possibly from intracontinental factors; in the second hypothesis the source would have been from Africa. When the affinities of the gymnophthalmids are established more clearly perhaps a choice of these alternatives will be easier to make.

In Figure 7, I have chosen the first of these hypotheses, principally on the basis that the known fossil record is earlier in North America and Asia and that *Durotrigia*, the possible ancestral lacertoid, occurs in Laurasia. Once again, however, the record is poor, and I do not think the second hypothesis can be falsified at present.

Beyond the lacertoid level of organization lie the more derived scincoid-cordyloid groups, together forming the sister group of the lacertoids. Origin of these groups must have been no later than the Upper Jurassic, since primitive lacertoids seem already to have been

Figure 7. Suggested area of origin (stippled) and later dispersals in lacertoids. Map data as in Figure 3. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. Alternative hypothesis discussed in text. + = Late Jurassic ?lacertoids; o = Late Cretaceous polyglyphanodontine teiids.

evolved, and primitive cordylids occur in the Late Jurassic of England, Portugal, and Wyoming. Here in particular the lack of a better African and Asian record makes interpretation difficult. At this stage of knowledge the scincoid radiation seems to have taken place in a different area from that of cordylids. Perhaps the simplest explanation is to presume that a primitive scincomorph stock had developed in Laurasia before the latest Jurassic. By then, the Tethys Sea was relatively open from the Pacific to at least as far as southern Asia, effectively isolating much of Laurasia and Gondwana. The fluctuating epicontinental seas of the European region could have first caused the lacertid-teiid dichotomy, and then fostered the divergence of western Laurasian populations that became the

cordylids, and eastern populations that evolved into scincoids.

The scincoids (Scincidae) are now so diverse and cosmopolitan a group that it is difficult to determine their biogeographic history. Greer (1970, 1974, 1977) has given some discussion of the matter, but analysis of the primitive eumecoid skinks is still in progress. Moreover, many skinks are well known for their ability to cross water barriers (see e.g., Darlington, 1957) and they have colonized islands extensively. The most primitive scincines are placed in the subfamily Scincinae, which includes *Eumeces*, probably the most primitive living scincid. This group is most relevant to the present discussion of early diversification of lizard families; the more derived lygosomine and other

scincid groups will not be considered here (Fig. 8). The scincines seem to have spread widely in southern Asia, and other scincid groups reached Australia, Africa, and Madagascar. Since Neotropical scincids are very limited in diversity, being represented (like the geckos) by probable raftings of cosmopolitan genera, the group probably did not arrive in Africa before the Late Cretaceous, after separation of Africa from South America. At that time Africa approached Laurasia more closely as a result of northeastward movement and rotation engendered by separation from South America; this would have facilitated entrance of a number of groups (e.g., geckos, agamids) into Africa. Another factor that may have operated to keep scincids (and perhaps

geckos and cordylids as well) from reaching South America (except by rafting) was the fact that in the Late Cretaceous, much of northwestern Africa was separated into several continental islands by epicontinental seas (see Howarth 1981: Fig. 13.18).

A primitive, probable scincine genus *Contogenys* occurred in the Late Cretaceous and Paleocene of North America, together with the more problematic and perhaps cordyloid genus *Sauriscus*. In the Paleocene of Brazil an apparent relative of *Contogenys* is found. At present there are no fossil scincids known in Europe (Cretaceous localities are essentially unknown, owing to extensive submergence, and no scincids occur in the relatively rich Cenozoic localities there),

Figure 8. Suggested area of origin (stippled) and later dispersals (arrows) in primitive Scincoidea (Scincinae). Entrance of *Contogenys*-like scincine into South America can have been either Late Cretaceous or earliest Paleocene. Map data as in Figure 3. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. See text for further explanation.

or in the large and diverse samples of fossil lizards in the Late Cretaceous of Central Asia; their direction of entrance into North America is thus difficult to predict. As noted above, entrance through South America is unlikely. It is possible that scincids were actually present in Central Asia during the Cretaceous, but in areas or environments different from those that have provided the known fossils. If so, they could have reached North America over the Bering connection, which was utilized by other lizards and mammals in the Late Cretaceous, finally reaching South America briefly in the Paleocene. This may be the simplest explanation at present, since the living South American scincids include only a few African-derived (and probably rafted) lygosomines. *Eumeces* itself occurs in North America in the Oligocene and is very likely to have been a later Bering migrant during the Early Cenozoic. Greer (1974) suggested this route for *Scincella*, an Asian lygosomine that also occurs in the southeastern United States. Origin of the endemic North American *Neoseps* was perhaps through a similar migration path.

The cordylids may have been initially a Laurasian group as indicated by their past distribution in Wyoming, England and Portugal in the Upper Jurassic (Fig. 9, paramacellodids). This wide distribution of a primitive cordylid group provides a means by which xantusiids can have been North American vicariant derivatives of a primitive cordylid stock, while the Cordylidae itself followed a southern direction, eventually becoming limited to subsaharan Africa and Madagascar after the development of the Sahara Desert. Cordylids were present in the Early Cenozoic of Europe, and it is possible that they dispersed northward from an original African center, perhaps during peak Eocene tropicality when other groups also extended their ranges (iguanids, agamids, gekkonids), and varanids). This is perhaps the best choice given that the European cordylid record at present is limited to a single species

and because much of Europe was submerged during the Cretaceous (Hallam, 1981). Given present data, I suggest that an as yet unknown Cretaceous paramacellodid derivative produced widely distributed populations that were eventually limited to North and Central America, becoming the Xantusiidae, and to Africa and Madagascar, becoming the Cordylidae.

Anguimorpha. As Renous (1979) stated, origin and distribution of anguimorphs was not simple. Anguimorpha and Scincomorpha appear to be sister groups (see e.g., Camp, 1923) and some similarities between some of the Jurassic scincomorphs (paramacellodids) and the earliest anguimorphs (dorsetisaurids) support this conclusion. Again we are faced with the great hiatus in the record between the Late Jurassic and the Late Cretaceous, but by the end of Mesozoic time the anguimorphs had diversified widely. In North America all known families (including most marine forms, if actually related) were present by the Late Cretaceous, and at the same time in Central Asia necrosaurids and varanids are known, suggesting a long Late Mesozoic radiation in Laurasia (Cracraft, 1973b also described anguimorphs as Laurasian). Both the Late Jurassic dorsetisaurids and the Cretaceous-Holocene xenosaurs appear to represent a primitive level of anguimorph organization, and the only evidence that more derived levels were present in the Jurassic is the presence of the presumed aigialosaur *Proaigialosaurus* from the Solnhofen limestones in Germany (Hoffstetter, 1964), if that group is in fact varanoid. Both from the standpoint of the fossil record and the distribution of *Xenosaurus* today, the xenosaurs seem to have originated in North America (Fig. 10), although if the Holocene *Shinisaurus* from eastern Asia is actually related to *Xenosaurus*, a possible wider area of origin could be considered (Asiameica of Cox, 1974 and Rage, 1981). Xenosaurs had a limited radiation in the Late

Figure 9. Suggested area of origin (stippled) and later dispersals in cordylids. Movement of cordylids into Europe would have been during Eocene tropicality, unless they had had a previous wider distribution there dating from the Mesozoic. Map data as in Figure 3. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. See text for further explanation.

Cretaceous and Cenozoic in North America (and perhaps Asia) and were eventually limited to Mexico and Central America (and perhaps eastern Asia). They were not present in the Cenozoic of Europe, corroborating a North American or Asiamerican origin; their distribution is thus similar to that for xantusiids, except for the possible Asian connection.

The anguids themselves also seem to have had a North American origin. They occur there in some diversity from the Late Cretaceous on into the present, and the Early Cenozoic anguids from Europe seem to have been a more restricted group derived from Early Cenozoic North American forms, rather than being part of a broad Laurasian distribution (Fig. 11). Lack of a European Cretaceous record again makes this difficult to cor-

roborate, but in the excellent sample of fossil lizards from the Late Cretaceous of Central Asia no anguids are present, suggesting that they were either limited to North America at that point, or to the latter and western Europe. Primitive (melanosaurinid) anguids in the Early Cenozoic of North America include *Xestops* and *Melanosaurus*; at least the first of these genera also occurs in the Eocene of Europe. The more derived (glyptosaurinid) genera in North America seem to be more closely related to each other than to *Placosaurus*, an Eocene glyptosaur from Europe and Asia. The occurrence of *Placosaurus* in Asia indicates that the Obik Sea, which separated Europe and Asia at that time, was not a barrier to at least some lizards. For the above reasons, I have chosen the North

Figure 10. Suggested area of origin (stippled) and later dispersals (arrows) in primitive anguoids. Map data as in Figure 3. Dashed line indicates possible dispersal of xenosaurs to Central Asia. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in that area. See text for further explanation.

Atlantic route for glyptosauroid entrance into Europe (Fig. 11), rather than the Bering connection. An apparent glyptosauroid from the Eocene of the Canadian Arctic Archipelago (Estes and Hutchinson, 1980) indicates that dispersal could have been through the North Atlantic connection to Europe, which was maintained at least until the end of the Early Eocene.

In North America the gerrhonotine and diploglossine groups appeared early, the former perhaps as early as the Late Cretaceous, the latter in the Eocene (Gauthier, 1982). Diploglossine emplacement on the West Indies was probably from North America; how and when they reached South America is less easily determined.

While gerrhonotines and diploglossines appear to have originated in North

America, the case of the anguines is less clear. Ophisauroids appear earlier in Europe (Eocene) than in North America (Oligocene), suggesting that they were Eurasian derivatives from an anguoid stock. The fossil record for these times, from both areas, is good enough to allow some confidence in this hypothesis. Several possibilities for origin of anguines exist that can be tested by future work. One is that the predominantly "grass-swimming" ophisauroids arose in the early Cenozoic in association with the distribution of grassland and savannah environments, finally achieving their present extensive Laurasian distribution. Another possibility, which I consider less likely, is that the Old and New World ophisauroids are parallel developments from primitive anguoid ancestors (see e.g., Klembara, 1979). In either case, ophisauroids appear to

Figure 11. Suggested area of origin (stippled) and later dispersals (arrows) in Anguinae. Glyptosaur movement was in the Eocene. Date of diploglossine dispersal to South America not known. Ophisaurids not shown. Map data as in Figure 3. Dates at end of arrows represent earliest known pre-Pleistocene fossils of the group in the area. See text for further explanation.

have originated subsequent to the early diversification of anguinae in the Cretaceous. The annellines are North American endemics, first appearing in the Early Eocene, and again suggest a North American origin for Anguinae.

The extinct varanoid necrosaurids occurred with true varanids in the Late Cretaceous of North America and Central Asia (different genera appear to be involved and the Asian forms have just recently been described by Borsuk-Bialynicka, 1982). In the Early Cenozoic there are two derived forms in Europe (Fig. 12). Necrosaurids may have originated in Asia, from varanoid ancestors, and dispersed across the Bering connection together with the true varanids during the Late Cretaceous, but it is not possible to falsify an origin in North America

or Asiamerica, perhaps from a common stem with the helodermatids. The Eocene European necrosaurids are derived with respect to the North American Late Cretaceous genus *Parasaniwa*, and probably reached Europe after its emergence in the very early Cenozoic.

The helodermatids (Fig. 12), now isolated in southwestern North America, were once more widely distributed; they occur in the Late Paleogene of Nebraska and Colorado (*Heloderma*), and had undoubtedly members in the Paleogene of France (*Eurheloderma*). Possible helodermatids occur in the Late Cretaceous of Wyoming (*Paraderma*), and if this record is correctly referred, the group had a long North American history. Renous (1979) supposed that helodermatids had a Gondwanan origin, since they

Figure 12. Suggested area of origin (stippled) and later dispersals (arrows) of primitive varanoids. Helodermatid and necrosaur dispersal to Europe was in the Early Cenozoic. Map data as in Figure 3. Dates at end of arrows indicate earliest pre-Pleistocene fossils of the group in that area. See text for further explanation.

have the derived nerve condition B that she believes appeared in the southern hemisphere. There is, however, no reason to suggest an origin outside of Laurasia, and if the early evolution of anguimorphs was also Laurasian, helodermatids could have been derived from a primitive varanoid stock, perhaps in North America, during the Late Cretaceous. Presence of *Eurheloderma* in France was probably a later dispersal during peak tropicality in the Eocene, as for varanids, gekkonids and agamids.

Among the anguimorphs, only the varanids achieved a wide southern continent distribution so far as known. They may have formed subgroups in Asia, Southeast Asia, India and Australia, as suggested for agamids, geckos, and skinks (Fig. 13). Varanids have a high level of activity compared with most

other lizards, and I believe them to have evolved originally in Laurasia as predators on small mammals of the Late Mesozoic, prior to the early evolution of snakes, which was probably a complementary Gondwanan radiation and (at least so far as the colubroid and more derived groups are concerned) primarily a Cenozoic phenomenon. There would have been no problem for varanids to reach Africa, which maintained connections of various sorts with Laurasia. Varanids also spread beyond the Arctic Circle in the Eocene, as did other lizard groups discussed above. Like agamids, they are found in the Eocene of both North America and Europe, and lingered in the former region until the Oligocene and until the Late Pleistocene in Europe. I have chosen the Bering route for Early Cenozoic entrance of *Saniwa* into North

Figure 13. Suggested area of origin (stippled) and later dispersals (arrows) of varanids. X = position of *Lanthanotus*. Varanid dispersal from Asia into North America and Europe was in the Early Cenozoic. Map data as in Figure 3. Dates at end of arrows indicate earliest pre-Pleistocene fossils of the group in that area. See text for further explanation.

America (Fig. 13), but the paucity of the Cretaceous record in Europe and Paleogene in Asia makes a clear-cut decision difficult to make.

OWEN'S HYPOTHESIS AND THE POSITION OF SOUTHEAST ASIA

The continental reconstructions of Owen (1976, 1981, 1982) are of interest here since they provide a much more extensive and intimate connection between eastern Laurasia and Gondwana during the Mesozoic, in contrast to the reconstructions commonly used (e.g., Smith and Bryden, 1977) in which a large Eastern Tethys seaway separates these two regions widely. In Figure 14, I have used Owen's reconstruction for the Early Cretaceous as a base map. In brief,

Owen's view is that this eastern Tethys is a geometric artifact produced by assuming an Earth of constant dimensions. He suggested that the Early Mesozoic earth was about 20% smaller in diameter than at present, and that it has expanded exponentially throughout Phanerozoic time. The assumption of an equal amount of subduction for each area of oceanic crust production by sea floor spreading is unwarranted in this view. Owen has given additional support for his hypothesis in later papers (1981, 1982), and the reviews by Sunderman and Brosche (1978) and Scrutton (1978) are also of interest in this regard. Beyond distortions that may be the result of an expanding Earth, the choice of a map base may further distort this region. A case in point is the Lambert equal-area projection used

Figure 14. Suggested areas of origin of lizard families and suggested dispersals between areas, given on continental reconstruction of Owen (1976) for Early Cretaceous time (120 m.y.b.p.). Family names occur approximately on presumed area of origin. Where names span more than one continent, origin is presumed to have been on the conjoined area. Continents identified in Figure 2. Number of lines connecting regions indicates numbers of families presumed to be involved in dispersals (as opposed to vicariances) during Jurassic, Cretaceous, and Cenozoic time. The number code identifies the families involved as follows: 1) iguanids, teiids, scincids, anguoids; 2) iguanids, glyptosaurs, helodermatids, necrosaurs; 3) agamids, scincids, varanids; 4) agamids, gekkonids, scincids, varanids; 5) gekkonids, scincids; 6) chamaeleonids, lacertids.

by Lillegraven *et al.* (1979). In this outstanding summary of paleogeographic data, the map distortion makes the actual close faunal relationships of Southeast Asia and Gondwana biogeographically inexplicable. In Owen's reconstruction the enormous Tethys seaway in the eastern regions—at least as an entity floored by simatic (oceanic) crust—disappears. Owen stated, however, that this does not rule out a narrower epicontinental Tethys in this region that could have been an effective barrier to animal movement in some cases. The absence of a large Tethys ocean also reduces the northward movement of India to about 1,600 km, thus reducing also the high rate

of movement usually predicted for this region to a figure more like that of rates of plate movement in other areas. In Owen's reconstruction greater India remained close to Southeast Asia throughout Mesozoic time, although its principal movements were the same as in other reconstructions: a counterclockwise rotation that eventually oriented its eastern margin toward the north, and continued northward movement that finally brought its present-day northern margin up against Asia in the Late Cretaceous, when the first phase of Himalayan orogeny began. Based on the conventional reconstruction of India as a free-floating plate in the developing

Indian Ocean, Lillegraven *et al.* (1979: 293) stated that "significant barriers to the exchange of land vertebrates from any continent with India existed from the Early Cretaceous through the remainder of the Mesozoic." The presence of endemic genera of agamids, geckos, lacertids, and skinks in India, however, attests to an ancient connection, almost certainly of Late Mesozoic age, with Southeast Asia. Colbert (1975) demonstrated that India had a fauna of cosmopolitan dinosaurs during the Cretaceous, which corroborates this view; see also Hallam (1981). Based on the double thickness of sialic (continental) crust under the Himalayas and Tibetan Plateau, Owen (1976) suggested that there was originally a more extensive, continental portion of the Indo-Tibetan plate forming a connection to Laurasia, permitting at least a limited amount of exchange to take place. The implications of this connection for the Agamidae have been examined by Moody (1980, see above) and they probably apply to the geckos, skinks, and varanids as well.

Movements northward of India and Australia during the Cenozoic, associated with extensive production of new sea floor in the Wharton Basin west of Australia, deformed Southeast Asia and pressed it to the east, twisting it and bringing Indonesia and the Celebes against Australia-New Guinea (Owen, 1976; Crostella, 1977). This juncture is Weber's Line (Darlington, 1957), an imaginary line of "faunal balance" separating an essentially Oriental fauna to the north from the Australian-New Guinea fauna. Owen's reconstruction thus gives Weber's Line a tectonic rationale.

The relevance of Owen's work here lies in the explanatory power it provides for the distribution of agamids, geckos, skinks, and varanids. In all these cases the most primitive forms of these families are not those found in the Australian region, but are in general Asian (Mertens, 1942; Kluge, 1967; Greer, 1970; Moody,

1980), a situation suggesting that subgroups were formed by vicariance of these areas and India. Cracraft (1975) suggested that Antarctic routes might account for the Australian and New Zealand distribution of some amphibian and reptile groups, although lizards were not among the ones he cited. The West Antarctic connection may have been available to (e.g.) spheonodontid reptiles and leiopelmatid frogs to New Zealand (Estes and Reig, 1973; Cracraft, 1975), and the East Antarctic route to Australia appears to have been open through the Mesozoic (Owen, 1976; Tarling, 1980). Relationships of the lizard groups mentioned above, evidence from fossils, and inferences from recent distributions do not indicate, however, that Antarctica was ever occupied by lizards or used as a migration route, although climatically (at worst cool temperate) and physically it seems to have been available. Perhaps this simply indicates once again that agamids, geckos, skinks, and varanids arose more to the north, or towards Laurasia, and did not have access to this route; alternatively the temperatures may simply have been marginal or prohibitive for lizards.

The above discussion strongly suggests that Owen's continental reconstruction, or some other arrangement that permits closer contact of Southeast Asia and Australia, is preferable zoogeographically to those of Smith and Bryden (1977) or Lillegraven *et al.* (1979), which separate these areas widely. Owen's arrangement in addition helps to explain the Gondwanan biota in some Triassic localities in Indonesia and eastern Asia, provides a tectonic rationale for Weber's Line of faunal balance in the Holocene fauna, and reduces the high relative speed of the Indian plate required by conventional reconstructions to one more in accord with other estimates of plate movement. The additional support for this hypothesis (Owen, 1981, 1982) makes it very difficult to falsify although his expanding Earth model has not yet

been widely accepted; in any case any alternative arrangement should explain adequately the above factors in order to be zoogeographically as well as geologically acceptable.

SUMMARY AND CONCLUSIONS

It seems clear from the above that much of early lizard evolution took place on an essentially Pangaeian continental arrangement, and that most of the crucial early stages of lizard development occurred during the Jurassic and Cretaceous. Some eolacertilian group, as yet unknown, gave rise to generalized lacerilians, perhaps as early as the Late Triassic or Early Jurassic. This generalized group became widely distributed in Pangaea, and a primary dichotomy took place as Laurasia separated from Gondwana in the Middle Jurassic, the Gondwana region retaining the primitive iguanians, the Laurasian populations forming a gekkotan—scincomorph—anguimorph ancestral group, the sister group of iguanians. This latter group appears to have been fragmented by epicontinental seas in the European Jurassic as well as by separation of Africa from North America in Western Gondwana. This led to the development of Late Jurassic groups that already showed the "basic patents" of some living families. Aside from the known Late Jurassic groups, it is probable that some of the more primitive (particularly iguanian) living families were also present in Gondwana, although not yet documented. In this view, the primitive iguanids (at a morunasaurine level) already occupied southern parts, at least, of both Africa and South America (western Gondwana). The origin of the Iguanidae seems to have been the result of plate separation of Laurasia and Gondwana, while the development of an ancestral acrodont agamid-chamaeleonid ancestor in northwestern Gondwana would have been through intracontinental factors. In

the earliest Cretaceous, the initiation of separation of South America and Africa led to the divergence, in isolation, of the South American morunasaurines and other Neotropical iguanids. Loss of African iguanids resulted in the present relict distribution of oplurines on Madagascar. Movement of iguanids from South America into North America and Europe was, as discussed above, essentially a dispersal phenomenon across preexisting barriers and probably did not occur earlier than the latest Cretaceous or the beginning of the Cenozoic.

Late Jurassic and Cretaceous seas extending south along the east coast of Africa, and the separation of Africa from eastern Gondwana (India—Madagascar—Antarctica—Australia) that began in the Early Cretaceous, caused division of the ancestral acrodont population of iguanians. This vicariance resulted in the formation of the Agamidae in the Southeast Asian region and the Chamaeleonidae from the remaining African population.

In Laurasia, intracontinental epicontinental seas fragmented the ancestral group of non-iguanian lizards. The gekkotans became restricted to Asia as the Obik Sea spread southward, perhaps as early as the beginning of the Cretaceous or the end of the Jurassic, resulting in the evolution of gekkonids in Asia (or Southeast Asia), India, and Australia. Diversification within the Gekkonidae probably followed from the separation of the Australia-Southeast Asia-India plates during the Cretaceous. Thus gekkonids vicariated by the action of intracontinental barriers (Obik Sea) but diversified through plate separation, as in the scenario given for agamids by Moody (1980). Since Australia was bisected during the Early Cretaceous by an enormous epicontinental sea, this may have played a part in the evolution of pygopodids from the early gekkonid stock.

Primitive scincomorphs must have arisen from marginal southern Laurasian populations of the ancestral stock of non-

iguanian lizards. Such primitive laceratoids could thus have deployed in North America as teiids, while lacertids developed from the Old World populations, separated by the Jurassic and Cretaceous epicontinental seas that covered, variably, much of what is now Europe. Origin of the gymnophthalmids from teiids in South America was the result of intracontinental factors as yet unknown.

Cordylids and scincids must have diverged as a result of fragmentation of the remaining primitive scincomorph groups in Laurasia by Late Jurassic epicontinental seas. Cordylids appear to have evolved on the North America—Greenland—Great Britain—Iberian Peninsula plate association while scincids may have been evolving on the other side of the European Jurassic seas, in Asia. Scincids subsequently dispersed through Laurasia and into Australia, South America, and Africa. Scincid evolution into subgroups may have followed that of agamids and geckos by vicariance in the Southeast Asian—Australian—Indian regions. Xantusiids and cordylids resulted from restriction of widespread primitive cordylids to areas in North America and Africa, probably through separation by Cretaceous epicontinental seas.

This fragmentation of the Laurasian ancestral group of non-iguanian lizards must have taken place within a relatively short time (Middle to Late Jurassic). In simplified form, I suggest that gekkonoids diverged to the north, lacertoids to the east and west, followed by an east-west dichotomy of cordylids and scincids, following the cladistic relationships given here (Fig. 1).

The origin of anguimorphs brings further complexity. These are the most derived of the major lizard groups, and probably diverged from a scincomorph or pre-scincomorph stem group after the latter had separated from a common ancestor with gekkotans. It is likely that the divergence of xenosaurs, dorsetisaurus and anguids was in the European-North

American region, with intracontinental factors rather than plate separation involved. Necrosaurids and varanids evolved in Asia, being separated from Europe by the Late Jurassic European epicontinental seas and the Cretaceous Obik Sea. Limited dispersal of primitive varanoids into North America in the Late Cretaceous, over the Bering connection, resulted in the North American evolution of helodermatids; a dispersal path for necrosaurids was thus also available.

It is interesting that in the present model, there is a decrease in Gondwanan influence in lizard evolution as more derived groups are considered. The Iguania thus appears wholly Gondwanan, gekkotans seem to have been originally Laurasian, and among scincomorphs only the gymnophthalmids, the pygopodids, and possibly the cordylids may have had a Gondwanan origin. There is no evidence for Gondwanan origin of any anguimorph group.

Several phases in saurian evolution may thus be proposed. The first is a Triassic phase, in which a number of different (perhaps unrelated) pre-lacertilian groups exploited the insectivorous habitus, with one group assuming herbivorous habits. Only a few aspects of this phase are known, the primitive paliguanids in Africa, with normal body type, followed by the gliding kuehneosaurs. After the Triassic, the eolacertilians dwindled as the more progressive lacertilians evolved.

After the climatic extremes of the Triassic, a Jurassic phase of lacertilian evolution resulted in the presence of all living suborders of modern lacertilians by the end of the Jurassic, indicating a long previous history of which we have no record. Since the continents up to the Middle Jurassic were still closely associated in a Pangaeian arrangement, Early Jurassic diversification must have been primarily the result of vicariant separation of populations within a widespread primitive lacertilian group by epicontinental seas or other intracontinental fac-

tors. By the Middle Jurassic, however, separation of Laurasia and Gondwana seems to have been the catalyzing factor in separating the Gondwana iguanians from the more derived Laurasian lizard groups, and if these iguanians were then essentially at iguanid level it could be said that that family arose by this vicariance. At the end of the Jurassic, widespread emergence must have permitted dispersal of the groups earlier formed by isolation in various regions of Laurasia. Very likely this dispersal heightened the competition among the groups already formed, and led to the canalization of various adaptive types within lizards.

In the third or Cretaceous phase of lizard evolution the present family groups evolved. Some of the more primitive living families must have already been in existence in the Late Jurassic although no record of them has yet been found. The Early Cretaceous was a time in which vicariance by plate separation was very important, and resulted in the formation of some iguanid and agamid subgroups and the Chamaeleonidae, as South America and Africa separated and Africa itself diverged from eastern Gondwana. Other than these relatively clear plate movements, epicontinental seas were extensive in the Cretaceous and must have separated populations of various lizard groups that resulted in the evolution of other living families.

The Cenozoic may be thought of as a fourth phase of lizard evolution, in which expansion of ranges in the Paleogene occurred, during periods of peak tropicality, and progressive restriction in the cooler and drier Neogene gradually formed the boundaries of the present distributions. Perhaps as Darlington (1957) has suggested, the rise of colubroid snakes, which seems to have been mainly a Cenozoic phenomenon, played a part in the progressive reduction of diversity and restriction of range seen in lizards.

As shown in Figure 14, there is no major center of lizard evolution in Southeast Asia or even the Old World Tropics,

as was suggested by Darlington (1957) for many vertebrate groups, although this region was important in the evolution of agamids, gekkotans, scincids, and varanids. Such tropical areas were widespread in the past (Lillegraven *et al.*, 1979), and origin of lizard families took place in all parts of the world, except, so far as known, Antarctica. The widespread tropical and temperate conditions of the past permitted lizard dispersal during the later Mesozoic and (particularly) Cenozoic, where connections existed.

ACKNOWLEDGMENTS

The data that permitted this paper to be written were gathered with the help of many colleagues and friends too numerous to mention here; a partial list is given by Estes (1982). For help in specific ways related to this paper I thank E. N. Arnold, Jose Bonaparte, Richard Etheridge, Jacques Gauthier, Max Hecht, Scott Moody, Michael Novacek, Hugh Owen, William Presch, Kevin de Queiroz, and Garth Underwood, although they do not necessarily accept all of my conclusions.

LITERATURE CITED

- AGER, D. 1975. The Jurassic world ocean. In K. Finstad and R. Selley (eds.), *Jurassic Northern Sea Symposium Proceedings*, Norwegian Petrol. Soc., Stavanger.
- ALBERCH, P., S. GOULD, G. OSTER, AND D. WAKE. 1979. Size and shape in ontogeny and phylogeny. *Paleobiology*, 5: 296-317.
- BELLAIRS, A. D'A. 1970. *The life of reptiles*. London, Weidenfeld and Nicholson, 2 vols., xi + 589 pp.
- BONS, J., AND G. PASTEUR. 1977. Solution histologique à un problème de taxonomie herpétologique intéressant les rapports paléobiologiques de l'Amerique du Sud et de l'Afrique. *C. R. Acad. Sci. Paris*, 284: 2547-2550.
- BORSUK-BIAŁYNICKA, M. 1982. Anguimorphan and related lizards from the late Cretaceous of the Gobi Desert, Central Asia. *Paleont. Polonica*, in press.
- CAMP, C. 1923. Classification of the lizards. *Bull. Amer. Mus. Nat. Hist.*, 48: 289-481.
- CARROLL, R. 1975. Permo-Triassic "lizards" from the Karroo. *Paleont. Afr.*, 18: 17-87.

- . 1977. The origin of lizards, pp. 359–396. In S. Andrews, R. Miles and A. Walker (eds.), *Problems in Vertebrate Evolution*. Linn. Soc. London, Symp. Ser. 4.
- CASAMIQUELA, R. 1980. Considerations écologiques et zoogéographiques sur les Vertébrés de la zone littorale de la mer du Maestrichtien dans le Nord de la Patagonie. *Mem. Soc. géol. France*, **139**: 53–55.
- COLBERT, N. 1973. Continental drift and the distribution of fossil reptiles, pp. 395–412. In D. Tarling and S. Runcorn (eds.), *Implications of continental drift to the earth sciences*. New York, Academic Press, NATO Adv. Study Inst., Vol. 1.
- . 1975. Mesozoic tetrapods and the northward migration of India. *J. Paleont. Soc. India* (Orlov Memorial Volume), **20**: 138–145.
- COX, B. 1974. Vertebrate palaeodistributional patterns and continental drift. *J. Biogeog.*, **1**: 75–94.
- . 1980. An outline of the biogeography of the Mesozoic world. *Mem. Soc. géol. France*, **139**: 75–79.
- CRACRAFT, J. 1973a. Continental drift, paleoclimatology and the evolution and biogeography of birds. *J. Zool.*, **169**: 455–545.
- . 1973b. Vertebrate evolution and biogeography in the Old World tropics: Implications of continental drift and paleoclimatology, pp. 373–393. In D. Tarling and S. Runcorn (eds.), *Implications of continental drift to the earth sciences*. New York, Academic Press, NATO Adv. Study Inst., Vol. 1.
- . 1975. Mesozoic dispersal of terrestrial faunas around the southern end of the world. 17th Int. Zool. Congr. 1972, Monaco. *Mem. Mus. Nat. Hist. Nat. Paris*, **88**: 29–54.
- CROSTELLA, A. 1977. Geosynclines and plate tectonics in Banda Arcs, Eastern Indonesia. *Bull. Amer. Assoc. Petrol. Geol.*, **61**: 2063–2081.
- DARLINGTON, P. 1957. *Zoogeography: the geographical distribution of animals*. New York, John Wiley and Sons, xi + 675 pp.
- DELFAUD, J. 1980. Les formations sédimentaires des bassins et plates-formes mésozoïques et cénozoïques. In: A. Autran and J. Dercourt, eds., *Évolutions Géologiques de la France*. 26^e Congr. géol. int. colloq. C7, *Mem. B.R.G.M. France*, **107**: 260–265.
- ELDREDGE, N. 1979. Cladism and common sense, pp. 165–198. In: J. Cracraft and N. Eldredge, eds., *Phylogenetic analysis and paleontology*. New York, Columbia University Press.
- ESTES, R. 1970. Origin of the Recent North American lower vertebrate fauna: an inquiry into the fossil record. *Forma et Functio*, **3**: 139–163.
- . 1982. *Handbuch der Paläoherpetologie*, vol. 10, *Sauria terrestres*. Gustav Fischer Verlag, Stuttgart.
- ESTES, R., AND J. HUTCHISON. 1980. Eocene lower vertebrates from Ellesmere Island, Canadian Arctic Archipelago. *Palaeogeog., Palaeoclimatol., Palaeoecol.*, **30**: 325–347.
- ESTES, R., AND L. PRICE. 1973. Iguanid lizard from the Upper Cretaceous of Brazil. *Science*, **180**: 748–751.
- ESTES, R., AND O. REIG. 1973. The early fossil record of frogs: a review of the evidence, pp. 11–63. In J. Vial (ed.), *Evolutionary Biology of the Anurans*. Columbia, Univ. Missouri Press.
- EVANS, S. 1980. The skull of a new eosuchian reptile from the Lower Jurassic of South Wales. *Zool. Journ. Linn. Soc.*, **70**: 203–264.
- FERRUSQUA-VILLAFRANCA, I. 1978. Distribution of Cenozoic vertebrate faunas in Middle America and problems of migration between North and South America. *Univ. Nac. Auton. Mexico, Inst. Geol. Bol.*, **101**: 182–321.
- GAUTHIER, J. 1980. *Anniella* (Sauria; Anguinae) from the Miocene of California. *PaleoBios*, **31**: 1–7.
- . 1982. Fossil Xenosauridae and Anguinae from the Lower Eocene Wasatch Formation, south central Wyoming, and a revision of the Anguinae. *Univ. Wyoming Contr. Geol.*, in press.
- GREER, A. 1970. A subfamilial classification of scincid lizards. *Bull. Mus. Comp. Zool.*, **139**: 151–184.
- . 1974. The generic relationships of the scincid lizard genus *Leiopisma* and its relatives. *Austr. J. Sci. (Suppl. Ser.)*, **31**: 1–67.
- . 1977. The systematics and evolutionary relationships of the scincid lizard genus *Lygosoma*. *J. nat. Hist.*, **11**: 515–540.
- HALLAM, A. 1975. *Jurassic environments*. Cambridge Earth Sci. Ser.; Cambridge Univ. Press, ix + 269 pp.
- . 1981. Biogeographic relations between the northern and southern continents during the Mesozoic and Cenozoic. *Geol. Rundschau*, **70**: 583–595.
- HALLAM, A., AND B. SELLWOOD. 1976. Middle Mesozoic sedimentation in relation to tectonics in the British area. *J. Geol.*, **84**: 301–321.
- HOFFSTETTER, R. 1962. Revue des récentes acquisitions concernant l'histoire et la systématique des squamates. *Colloq. Intern. Prob. Pal. C.N.R.S.*, **104**: 243–279.
- . 1964. Les Sauria du Jurassique supérieur et spécialement les Gekkota de Bavière et de Mandchourie. *Senck. Biol.*, **45**: 281–324.
- . 1967. Coup d'oeil sur les sauriens (lacertilien) des couches de Purbeck (Jurassique supérieur d'Angleterre: résumé d'un Mémoire). *Colloq. Intern. Prob. Pal. C.N.R.S.*, **163**: 349–371.
- HOWARTH, M. 1981. Palaeogeography of the Mesozoic, pp. 197–220. In Cocks (ed.), *The evolving Earth*. British Mus. (Nat. Hist.) and Cambridge University Press.
- JULLIEN, R. AND S. RENOUS. 1972. Variations du trajet du nerf ulnaire et de l'innervation des muscles dorsaux de la jambe chez les lacertil-

- iens. Bull. Mus. Nat. Hist. Nat. Paris, **29**: 247-252.
- KLEMBARA, J. 1979. Neue funde der Gattungen *Ophisaurus* und *Anguis* (Squamata, Reptilia) aus dem Untermiozän Westböhmens (CSSR). Vest. Ustred. Ust. geol., **54**: 163-169.
- KLUGE, A. 1967. Higher taxonomic categories of gekkonid lizards and their evolution. Bull. Amer. Mus. Nat. Hist., **135**: 1-59.
- LILLEGRAVEN, J., M. KRAUS, AND T. BOWN. 1979. Paleogeography of the world of the Mesozoic. In J. Lillegraven, Z. Kielan-Jaworowska and W. Clemens (eds.), Mesozoic mammals: the first two-thirds of mammalian history. Berkeley, Univ. Calif. Press, 321 pp.
- MCDOWELL, S. AND C. BOGERT. 1954. The systematic position of *Lanthanotus* and the affinities of the anguimorph lizards. Bull. Amer. Mus. Nat. Hist., **105**: 1-142.
- MERTENS, R. 1942. Die Familie der Varanidae. Abh. Senck. Nat. Ges., **462**: 1-391; **465**: 117-234; **466**: 235-391.
- MOODY, S. 1980. Phylogenetic and historical biogeographical relationships of the genera in the Agamidae (Reptilia: Lacertilia). Ph.D. thesis, Univ. Michigan, xv + 373 pp.
- OWEN, H. 1976. Continental displacement and expansion of the earth during the Mesozoic and Cenozoic. Phil. Trans. Roy. Soc. London, ser. A, **281**: 223-291.
- . 1981. Constant dimensions or an expanding Earth?, pp. 179-192. In Cocks (ed.), The evolving Earth. British Mus. (Nat. Hist.) and Cambridge University Press.
- . 1982. Ocean floor spreading evidence of global expansion. Tectonophysics, in press.
- PRESCH, W. 1980. Evolutionary history of the South American microteiid lizards (Teiidae: Gymnophthalminae). Copeia, **1980**(1): 36-56.
- PROTHERO, D., AND R. ESTES. 1980. Late Jurassic lizards from Como Bluff, Wyoming, and their palaeobiogeographic significance. Nature, **286**: 484-486.
- RAGE, J.-C. 1978. Une connexion continentale entre Amérique du Nord et Amérique du Sud au Crétacé supérieur? L' exemple des vertébrés continentaux. C. R. Somm. Géol. France, **1978**: 281-285.
- . 1981. Les continents peri-atlantiques au Cretace supérieur: migrations des faunes continentales et problèmes paléogéographiques. Cret. Res., **2**: 65-84.
- RENOUS, S. 1978. Confrontation entre certaines données morphologiques et la repartition géographique des formes actuelles des Sauriens: interpretation phylogénétique et hypothèse paléobiogéographique. Bull. Soc. Zool., **103**: 219-228.
- . 1979. Application des principes cladistiques a la phylogénèse et la biogéographie des Lacertiliens. Gegenbaurs Morph. Jahrb., **125**: 376-432.
- . 1981. Developpement de l'aspect historique de la biogéographie par la superposition de deux thèses: proposition d'une hypothèse phylogénétique batie selon les principes Hennigiens et theorie de la dérive des continents. C. R. Soc. Biogeogr., **57**: 81-102.
- RIEPEL, O. 1980a. The phylogeny of anguimorph lizards. Denkschr. Schweiz. Naturf. Ges., **94**: 1-86.
- . 1980b. The postcranial skeleton of *Lanthanotus borneensis* (Reptilia, Lacertilia). Amphibia-Reptilia, **1**: 95-112.
- . 1980c. The trigeminal jaw adductor musculature of *Tupinambis* with comments on the phylogenetic relationships of the Teiidae (Reptilia, Lacertilia). Zool. Linnean Soc., **69**: 1-29.
- ROBINSON, P. 1967. The evolution of the Lacertilia. Colloq. Intern. Prob. Pal., C.N.R.S., **163**: 395-407.
- . 1973. Paleoclimatology and continental drift, pp. 451-476. In D. Tarling and S. Runcom (eds.), Implications of continental drift to the earth sciences. New York, Academic Press, NATO Adv. Study Inst., Vol. 1.
- ROMER, A. 1956. Osteology of the reptiles. Univ. Chicago Press, xxi + 772 pp.
- ROSEN, D. 1978. Vicariant patterns and historical explanation in biogeography. Syst. Zool., **27**: 159-188.
- SAVAGE, J. 1968. The origins and history of the Central American herpetofauna. Copeia, **1966**: 719-766.
- SCHMIDT-EFFING, R. 1980. Geodynamic history of oceanic crust in southern central America. Intern. Alfred-Wegener-Symp., Berliner Geowiss. Abh., **A19**: 201-202.
- SCRUTTON, C. 1978. Periodic growth features in fossil organisms and the length of the day and the month, 154-196. In P. Brosche and J. Sundermann (eds.), Tidal friction and the earth's rotation. Berlin, Springer Verlag.
- SMITH, A., AND J. BRYDEN. 1977. Mesozoic and Cenozoic paleocontinental maps. Cambridge Univ. Press, 70 pp.
- SUNDERMANN, J., AND P. BROSCHE. 1978. Numerical computation of tidal friction for present and ancient oceans, pp. 125-144. In P. Brosche and J. Sundermann (eds.), Tidal friction and the earth's rotation. Berlin, Springer Verlag.
- TARLING, D. 1980. The geologic evolution of South America with special reference to the last 200 million years, pp. 1-41. In R. Chiochon and A. Chiarelli (eds.), Evolutionary Biology of New World Monkeys and Continental Drift. New York and London, Plenum Press.
- UNDERWOOD, G. 1971. A modern appreciation of Camp's "Classification of the Lizards." Introduction to facsimile reprint, Soc. Stud. Amphib. Rept., pp. vii-xvii.

Commentary on a Frog and Lizard Newly Recorded from Central Durango, Mexico

ROGER CONANT¹

ABSTRACT. Herpetological and hydrographic features of the Río Nazas, a stream of internal drainage in Durango, Mexico, are discussed. The cliff frog, *Syrrophus guttillatus*, is newly reported from the Nazas valley, and the earless lizard, *Callisaurus (Cophosaurus) texanus scitulus*, apparently invaded a new habitat created near the source of the river as a result of a major flood that occurred in 1968.

INTRODUCTION

The Río Nazas, one of the streams of interior drainage of the *altiplano* of northern Mexico, has long been of interest to zoogeographers, largely because its fauna includes a number of aquatic and semiaquatic organisms that show close affinities with those of the Río Grande which forms the southern boundary of Texas. Meek, as long ago as 1904, called attention to the similarities exhibited by the fish faunas of the two rivers, even though they are now separated by broad areas of arid terrain. A former connection must be assumed, probably during pluvial periods of the Pleistocene. I briefly reviewed the subject (Conant, 1963) and published a map of the general area in a paper that was chiefly concerned with natricine snakes of the Río Nazas and certain other still-existing but disjunct streams that obviously were part of the old Nazas system.

While I was assembling data for that study and a later one (Conant, 1969), I

made a number of visits (during 1960, 1961, 1962, and 1965) to the area at the source of the Nazas near El Palmito, Durango, during which a number of amphibians and reptiles were obtained, most of which were deposited in the collection of the American Museum of Natural History.

The Río Nazas is formed by the union of the Río de Ramos and the Río del Oro, but its water is impounded at its source by a large dam, the Presa Cárdenas, and is controlled by the opening or closing of tunnels at the base of the structure. The Nazas, which flows chiefly eastward across the Chihuahuan Desert, formerly emptied into a large interior basin in southwestern Coahuila. Goldman (1951: Pl. 44) portrayed a sump in the now arid basin where some of the runoff from the river accumulated during earlier years. The water of the Nazas is currently used for agriculture and other purposes in the Laguna District that includes the cities of Torreón, Gómez Palacio, and Ciudad Lerdo. (For information on the hydrography of the Nazas and other streams of the region see Tamayo and West, 1964).

I was able to return to the locality near El Palmito on 20 and 21 July 1976 in company with the late Dr. James D. Anderson, Keith A. Hawthorne, Salome Litwin, and Anderson's daughter, Susan, as part of a field trip through the Chihuahuan Desert as far south as central Zacatecas. Many changes were noted in the relatively broad floodplain of the Nazas immediately below the Presa

¹Department of Biology, University of New Mexico, Albuquerque, New Mexico 87131, U.S.A.

Cárdenas, which formerly had been occupied in large part by arid mesquite scrub. That vegetation had largely disappeared but was being replaced by clumps of miscellaneous growth. An exception was the intermittent gallery forest of very large Mexican cypresses (*Taxodium*) along the edge of the river.

A concrete spillway, which had been constructed near the village of El Palmito to serve as a relief for the main dam during periods of high water, had vanished, and the ravine below it had been steeply eroded to a depth of 10 m or more as was clearly indicated by a comparison with photographs taken at the site about 15 years earlier. Water from the spillway had evidently poured across the Nazas floodplain, sweeping everything before it except the larger rocks, the giant cypresses, and a few much smaller trees along the perimeter or at other places outside the main path of the torrent. Presumably little else remained except rocks, sand, and gravel after the water subsided.

The cause of the flood was Hurricane "Naomi" that struck the Mazatlán area of Sinaloa in September, 1968 with a deluge of rainfall that spilled eastward over the Sierra Madre Occidental and blanketed the watershed feeding the Nazas. The rainy cyclone period lasted from 8 through 13 September during which period 213 mm of rain fell at El Salto and 99 mm at El Palmito-Presa Cárdenas (Dr. Robert H. Schmidt, Jr., *in litt.*, through Dr. Robert G. Webb). Schmidt (1976) reported that the same storm resulted in 320 mm of rain at Siqueros, Sinaloa, in a single 24-hour period.

During the almost eight years that had passed between the time of the deluge and our 1976 visit, part of the herpetological fauna had returned, although we saw no snakes. Turtles (*Kinosternon hirtipes murrayi* and *Chrysemys scripta* ssp.) were abundant in the river and were also present in several small ponds in rocky areas below the former spillway that had been scoured out by the rush of

water and its burden of debris. A large female of the latter species measured 308 mm in carapace length. Two species of spiny lizards (*Sceloporus*) were observed and two *S. undulatus consobrinus* were collected. *Hyla arenicolor* was heard calling and specimens of *Bufo punctatus* and members of the *Rana pipiens* complex were seen. Two taxa of unusual interest, *Syrrophus guttilatus* and *Callisaurus (Cophosaurus) texanus scitulus*, both of which are discussed in detail below, also were found.

Syrrophus guttilatus (Cope)

Five of these frogs (AMNH 107551-5) were obtained, three during the evening of 20 July 1976 and two the following evening. Anderson heard a large number calling from a small cliff face near the northeastern edge of the Nazas floodplain and from talus slides and rock piles at the base of the cliff. He described three distinct calls in his field notes as follows: "A short chirp of low intensity; a louder, more distinct and more metallic chirp; and a soft trill very much like the encounter call of *Hyla crucifer*." A tape was made of the calls of three different frogs, but because their brief, high-pitched sounds are difficult to record well (and to graph), even under the best conditions, only one (from AMNH 107551) was suitable for the production of an audio-spectrogram (Fig. 1).

Some of the frogs responded to crude imitations of their notes, but the majority fell silent when approached. Most seemed to be calling from crevices in the cliff face or from under the piles of rocks. Six were seen, one of which was wedged, head upward, in a vertical opening, between two adjacent rocks each about a half meter high; the frog was about 3 cm inward from the exposed face of the rocks and about 6 cm above their base. One was calling from a small burrow beneath a rock, and another was heard among small rocks at the base of a dead bush.

Figure 1. Audio-spectrogram of *Syrrhophus guttilatus*; same note displayed with 50 Hz (left) and 300 Hz (right) filters. Based on a tape recording made 20 July 1976 (from AMNH 107551) at the edge of the Río Nazas floodplain near El Palmito, Durango, Mexico, temperature 17.8° C.

On 21 July there was a light rain off and on during the late afternoon and early evening, and many more *Syrrhophus* were calling than on the previous night. The cliff apparently was high enough to escape inundation during the flood; at least there were no high water marks on it such as were visible on the cypresses and some of the rocks along the edge of the floodplain. I had not examined it during my earlier visits to the area, nor had I heard any chirps, chiefly because my evenings were fully occupied searching for natricine snakes along the river and in the pools near it (Conant, 1969).

In Anderson's field notes the frogs were described in life as being, "Pale greenish yellow with dark markings. The eye seemed dark although up close it appeared to have some light markings." My field notes were slightly different, viz.: "Pale dorsal areas olive green. Dark dorsal markings black or dark gray. Dorsal and concealed surfaces of hind legs yellowish olive; dark markings match back."

Two of the specimens were sent to Dr. John D. Lynch, an authority on leptodactylid frogs, for a determination as to species. His response is quoted, virtually

in full, as follows: "Both specimens are adult males agreeing with *S. gutttilatus* in the morphology of the hands and feet, size, proportions, and skin texture, but differing in having a more acuminate snout (compare *S. gutttilatus* and *S. marnockii* in my 1970 paper [Figure 9]; these resemble *marnockii* in snout shape), more intense marbling of the dorsum (rather than flecks and thin vermiculations), and in having less distinct tympana. The tympana are large enough for *gutttilatus* (59.1% eye length in the 26.7 mm specimen [AMNH 107554] and 55.2% in the 25.0 mm specimen [AMNH 107553]). The coloration in life as recorded on the specimen tag [Conant notes] also is at variance with what is known of *S. gutttilatus*.

"The frogs either represent somewhat variable *S. gutttilatus* or another weakly differentiated allopatric population. My gut reaction is to call them *S. gutttilatus* and attribute the differences to geographic variation. This decision is based on the lack of difference in hand and foot morphology, the knowledge that techniques of preservation and fluid concentrations affect distinctness of tympana, . . . and a sense that pattern difference is not significant enough to warrant species description. The most vexing feature is the difference in the snout shape, but on the whole it does not compel me to argue that the beast is new."

The three specimens not seen by Lynch agree in having more acuminate snouts and a more intense dorsal marbling by dark pigment (the latter particularly noticeable in AMNH 107551) than in the *gutttilatus* depicted by Lynch (*loc. cit.*). All five specimens have an interorbital bar (slightly interrupted in AMNH 107552) that is both much broader and darker than shown in Lynch's figure.

The discovery of *Syrrhophus gutttilatus* at the source of the Río Nazas in central Durango constitutes a major range extension both for the species and the genus. According to Lynch, in his taxo-

nomic revision of *Syrrhophus* (1970: Fig. 10), the nearest localities for *gutttilatus* were in the Big Bend National Park, Texas, extreme southeastern Coahuila, extreme southwestern San Luis Potosí, and Guanajuato. He has since commented (*in litt.*) that every other specimen of *gutttilatus* he has seen since 1969, when his manuscript was submitted for publication, has been from the area he mapped for this frog. Lynch's postulated range for the genus *Syrrhophus* as a whole (1970: Fig. 21), shows a large blank space surrounding the Nazas River locality.

It is likely that *gutttilatus* will eventually be found at additional stations along the Nazas or its principal tributaries. The upper portions of the river valley abound with rocky cliffs, some of them at the edges of the stream where it passes through gorges. Because of the constant flow of water on a year round basis, the humidity should be high enough in many places to support such small anurans as *Syrrhophus*.

Callisaurus (Cophosaurus) texanus scitulus (Peters)

Four lizards of this taxon (AMNH 119225-8) were collected 21 July 1976 on the floodplain of the Río Nazas where sand and gravel had replaced the many dense clumps of mesquite and other vegetation that were present prior to the 1968 immersion and scouring. Several other *scitulus* were also seen actively prowling during late morning of the same day. Most permitted us to approach closely, but they were watchful and wary. When we moved too near they raced for cover but seldom entered it, stopping short instead and curling their tails and waving them from side to side in characteristic fashion. When we retreated they promptly resumed prowling. Anderson recorded the temperatures of three immediately after their capture at 36.8°, 39.2°, and 40.4° C.

This lizard was not found during my previous trips to the area even though I had walked repeatedly through virtually all open parts of the floodplain during favorable weather at various times in the months of June through September, collectively. Presumably *scitulus* entered the newly created habitat after the 1968 flood, perhaps as soon as the spotty vegetation had grown large enough to provide temporary shelter from the sun and such predators as men, dogs, and birds. It was almost certainly an invader, and in that capacity it would be of interest to Dr. Ernest E. Williams, whose studies on such phenomena are exemplified by his scholarly "The Ecology of Colonization as Seen in the Zoogeography of Anoline Lizards on Small Islands" (1969). It is likely, however, that barring additional severe inundations, the Nazas floodplain eventually will revert, at least in part, to a habitat less favorable for this species.

Among the specimens obtained are two adult males (snout-vent lengths 54 and 59 mm) and two adult females (snout-vent lengths 56 and 58 mm), all of which were found on gravelly or rocky substrata. These lizards agree meristically with Peter's (1951) description of the southwestern race of *texanus* as follows: a) ventral scales in midline, 78 to 82 (mean 81.0); b) dorsal scales along midline, counted for a distance equal to the length of the head measured from tip of snout to posterior edge of interparietal, 36 to 39 (mean 37.8); and c) femoral pores (the sum of the counts for both legs), 27 to 31 (mean 29.5).

No color notes were made in the field, but the color patterns of the four Nazas lizards were compared with six specimens of *scitulus* from Chihuahua and six from Grant County, New Mexico, in the collection of the Museum of Southwestern Biology of the Department of Biology of the University of New Mexico. The patterns were all basically similar. The paired dark spots, one on each side of the midline of the back, were less conspicuous on the Nazas specimens than on

those from the more northern localities, but the dark, wavy crossbands on the tail and the longitudinal, light-bordered dark stripe on the rear of the thigh were about equally intense in the entire sample from the three areas. The dark ventrolateral stripes did not penetrate quite so far anterodorsally in the two Nazas males as they did in the males from Chihuahua and New Mexico. The venters in all were comparable, both in regard to the pale ground color and the intensity and position of the dark pigmentation of the males. Local demes of *scitulus* vary in coloration and pattern and, with this fact in mind, I could detect no significant differences among the several population samples.

The Nazas lizards represent a range extension west of the localities plotted by Peters (1951: Map 1) who listed geographical data for the 1,254 specimens of the *texanus* complex then present in museums in the United States. His nearest localities were approximately 190 km downstream in the Nazas valley between Lerdo and La Goma and "14 miles north of Pedricana" (*sic.* = Pedriceña).

There are widely conflicting opinions about the proper generic name for this lizard. The principal ones, arranged in chronological order, are summarized as follows: 1) Axtell (1958) considered *texana* to belong to the genus *Callisaurus*, and he therefore did not include it in his proposed monographic revision of the genus *Holbrookia*. 2) Earle (1961), based on a study of the middle ear, thought that *texana* represented an "earless" *Callisaurus*. 3) Clarke (1965), as a result of observations on its push-up pattern, assigned *texana* to monotypic status, resurrecting Troschel's (1852) name of *Cophosaurus* for it. 4) Guttman (1970), using electrophoretic evidence from hemoglobin, showed that all the members of the group have a close relationship, and he doubted the validity of *Cophosaurus* as a separate entity. Guttman also cited papers by Carpenter (1962, 1963) on the display action of two

species of *Urosaurus* and three of *Uma* that indicated sufficient intrageneric variation to suggest that the splitting off of *Cophosaurus* on the basis of its behavior was untenable. 5) Cox and Tanner (1977), in a study on the osteology and myology of the head and neck regions, advocated the retention of *Callisaurus*, *Holbrookia*, and *Uma* as separate genera and the return of *Cophosaurus* to the synonymy of *Holbrookia*. 6) Adest (1978), in his electrophoretic study, proposed uniting all under the single genus *Callisaurus*.

Cox and Tanner (1977) may be consulted for a detailed review of the taxonomic peregrinations of the group, the so-called sand lizards, up to the time they submitted their manuscript. Obviously there is wide disagreement about the generic classification, and the group is in need of thorough study, preferably by an investigator who can approach the problem objectively, consider all its facets, and not be overly influenced by his own personal methods of research, however important they may be. In the meantime, the approach I personally favor is to consider *Callisaurus*, *Cophosaurus*, *Holbrookia*, and *Uma* as subgenera of the genus *Callisaurus* Blainville (1835). Such an arrangement would reflect most of the diverse opinions as well as to provide zoologists in general, many of whom are not especially familiar with these iguanids, with a quick and easily envisioned concept of their relationships. Herpetologists who, with a few notable exceptions, have resisted using the subgenus, might do well to emulate practitioners of other disciplines who have found the subgenus to be a useful and convenient category. As Simpson remarked long ago (1961), the aversion to subgenera is mostly psychological.

ACKNOWLEDGMENTS

The several persons who accompanied me on the field trip through the

Chihuahuan Desert to Durango during 1976 were helpful in many ways, especially the late James D. Anderson whose contagious enthusiasm and broad knowledge of the amphibians and reptiles of the region inspired us all. I am grateful to John D. Lynch who provided his opinion on the taxonomic status of the *Syrrhophus* reported in this paper, and I am indebted to Richard G. Zweifel for preparing the audio-spectrogram of the call of one of these frogs. John S. Applegarth, Ralph W. Axtell, Charles M. Bogert, Charles J. Cole, William G. Degenhardt, John D. Lynch, Samuel B. McDowell, Jr., Hobart M. Smith, Robert G. Webb, and Daniel C. Wilhoft undertook a critical reading of the manuscript or were helpful in a variety of other ways or both. The Directors General de la Fauna Silvestre, Departamento de Conservación, in Mexico City very kindly issued collecting permits to me over a period of many years. My field work in Mexico was supported in large part by the National Science Foundation Grants G-9040, G-22657, and GB-2177. Also I must mention my deep obligation to my late wife, Isabelle Hunt Conant, who accompanied me and was my staunch companion in the field during all of my trips to Mexico from 1949 through 1967, but who was too ill to participate with us during 1976.

LITERATURE CITED

- ADEST, G. A. 1978. The relations of the sand lizards *Uma*, *Callisaurus* and *Holbrookia* (Sauria: Iguanidae): an electrophoretic study. Ph.D. dissertation, Univ. California, Los Angeles, 131 pp. Diss. Abstr. Int. B 39: 36.
- AXTELL, R. W. 1958. A monographic revision of the iguanid genus *Holbrookia*. Diss. Abstr., 19: 1476-1477.
- BLAINVILLE, H. M. D. DE. 1835. Description de quelques especes de reptiles de la Californie précédée de l'analyse d'un système général d'herpétologie et d'amphibiologie. Nouv. Ann. Mus. d'Hist. Nat. (Paris), 4:(3) 232-296.
- CARPENTER, C. C. 1962. A comparison of the patterns of display of *Urosaurus*, *Uta*, and *Streptosaurus*. *Herpetologica*, 18: 145-152.

- . 1963. Patterns of behavior in three forms of the fringe-toed lizards (*Uma*-Iguanidae). *Copeia*, 1963: 406-412.
- CLARKE, R. F. 1965. An ethological study of the iguanid lizard genera *Callisaurus*, *Cophosaurus*, and *Holbrookia*. *Emporia State Res. Stud.*, 13: 1-66.
- CONANT, R. 1963. Semiaquatic snakes of the genus *Thamnophis* from the isolated drainage system of the Río Nazas and adjacent areas in Mexico. *Copeia*, 1963: 473-499.
- . 1969. A review of the water snakes of the genus *Natrix* in Mexico. *Bull. Amer. Mus. Nat. Hist.*, 142: 1-140.
- COX, D. C., AND W. W. TANNER. 1977. Osteology and myology of the head and neck regions of *Callisaurus*, *Cophosaurus*, *Holbrookia*, and *Uma* (Reptilia: Iguanidae). *Great Basin Nat.*, 37: 35-56.
- EARLE, A. M. 1961. The middle ear of *Holbrookia* and *Callisaurus*. *Copeia*, 1961: 405-410.
- GOLDMAN, E. A. 1951. Biological investigations in México. *Smithsonian Misc. Coll.*, 115: xiii + 476 pp.
- GUTTMAN, S. I. 1970. An electrophoretic study of the hemoglobins of the sand lizards, *Callisaurus*, *Cophosaurus*, *Holbrookia*, and *Uma*. *Comp. Biochem. Physiol.*, 34: 569-574.
- LYNCH, J. D. 1970. A taxonomic revision of the leptodactylid frog genus *Syrhophus* Cope. *Univ. Kansas Publ., Mus. Nat. Hist.*, 20: 1-45.
- MEEK, S. E. 1904. The fresh-water fishes of Mexico north of the Isthmus of Tehuantepec. *Field Columbian Mus., Zool. Ser.*, 5: lxiii + 252 pp.
- PETERS, J. A. 1951. Studies on the lizard *Holbrookia texana* (Troschel) with descriptions of two new subspecies. *Occ. Papers Mus. Zool., Univ. Michigan*, 537: 1-20.
- SCHMIDT, R. H., JR. 1976. A geographical survey of Sinaloa. *Southwestern Stud., Texas Western Press, Mono.* 50: 1-77.
- SIMPSON, G. G. 1961. Principles of animal taxonomy. New York, Columbia Univ. Press, xiv + 247 pp.
- TAMAYO, J. L., AND R. C. WEST. 1964. The Hydrography of Middle America, pp. 84-121. R. Wauchope and R. C. West (eds.), *Handbook Middle American Indians*, Vol. 1. Austin, Univ. Texas Press. 84-121.
- TROSCHER, F. H. 1852. *Cophosaurus texanus*, neue eidechsen-gattung aus Texas. *Archiv. Naturg. (Berlin)*, 16: 388-394.
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, 44: 345-389.

List of Peruvian *Anolis* with Distributional Data (*Sauria: Iguanidae*)

NELLY CARRILLO DE ESPINOZA¹

ABSTRACT. Species of the iguanid lizard *Anolis* known to occur in Peru are: *A. boettgeri*, *A. bombiceps*, *A. chrysolepis scypheus*, *A. dissimilis*, *A. f. fuscoauratus*, *A. laevis*, *A. orton*i, *A. punctatus boulengeri*, *A. p. punctatus* (new record for Peru), *A. trachyderma*, and *A. transversalis*. Distributional maps for these species in Peru are provided.

INTRODUCTION

This checklist of Peruvian species of the iguanid lizard *Anolis* that I have prepared is based primarily on data contained in the works published by Williams, Vanzolini, Dixon and Soini (see Literature Cited), as well as on the collections of the Section of Herpetology of the Museo de Historia Natural "Javier Prado" of the Universidad Nacional Mayor de San Marcos in Lima, Peru. Maps have been compiled from the available distributional data (Figs. 1, 2, 3).

It is my desire that this contribution will prove useful to all students interested in the Peruvian herpetofauna. I dedicate the paper to the distinguished herpetologist, Dr. Ernest E. Williams, in honor of his recent retirement as Curator of Herpetology at the Museum of Comparative Zoology.

Abbreviations. MCZ, Museum of Comparative Zoology; TCWC, Texas Cooperative Wildlife Collection, Texas A&M University; MHNJP, Museo de Historia Natural "Javier Prado"; FMNH, Field Museum of Natural History, Chicago; ANSP, Academy of Natural Science, Philadelphia; AMNH, American Museum of Natural History.

CHECKLIST

Anolis boettgeri Boulenger 1911

Type Locality. Huancabamba, Peru, 1,000 m, restricted to Oxapampa, Peru by Barbour (1934).

Material. AMNH: Huancabamba, Piura; MCZ: Oxapampa, Pasco.

Distribution. Known only from Huancabamba and Oxapampa of the Cis-Andean region.

Anolis bombiceps Cope 1876

Type Locality. Nauta, Peru.

Material. MHNJP: Saur.0131, San Regis, Río Marañón, Pv. Loreto, Loreto; Saur.0208, Santa María de Nieva, Pv. Bagua, Amazonas; TCWC: Centro Union: Mishana; Moropon (Dixon and Soini, 1975). Additional records: Estirón; Río Ampiyacu; Río Itaya; Río Nanay; San Regis (Vanzolini and Williams, 1970).

Distribution. Amazonian region of Peru, Ecuador, and Brazil plus the Orinoco basin and the Guianas.

¹Section Herpetology, Museo de Historia Natural "Javier Prado", Universidad Nacional Mayor de San Marcos, Lima, Peru.

Anolis chrysolepis scypheus Dumeril and Bibrón 1837

Type Locality. French Guiana and Surinam.

Material. TCWC: Centro Unión (Dixon and Soini, 1975); Pebas (Burt and Myers, 1942). Additional records: Barranca; mouth of Río Santiago; Río Itaya; Iquitos; Yarinacocha; Pedrera; Pampa Hermosa; Río Huallaga, Huánuco (Vanzolini and Williams, 1970).

Distribution. Amazonian region of Colombia, Ecuador, Peru, and Brazil plus the Orinoco basin, the Guianas and central Brazil with adjacent Paraguay.

Anolis dissimilis Williams 1965

Type Locality. Itahuania, Upper Río Madre de Dios, Peru.

Material. FMNH: Itahuania, Upper Río Madre de Dios.

Distribution. Known only from the type locality.

Anolis fuscoauratus fuscoauratus D'Orbigny 1837

Type Locality. "Chile," corrected by D'Orbigny (1847) to Río Mamore between Loreto and Río Sara, Bolivia, restricted to Provincia Moxas, Bolivia, by Bocourt (1873).

Material. MHNJP: Saur.0009, Chanchamayo, Pv. Tarma, Junín; Saur.0013, Pucallpa, Pv. Crnel. Portillo, Ucayali; Saur.0127, La Merced, Pv. Tarma, Junín; Saur.0354, Bellavista, Pv. Jaén, Cajamarca; Saur.0440, Yurinaqui Alto, Pv. Tarma, Junín; Saur.0841–0870, Río La Torre, Río Tambopata, Pv. Tambopata, Madre de Dios; Saur.0899, Tingo Maria, Pv. Huánuco, Huánuco; TCWC: Centro Unión; 5 km NNE Iquitos; Mishana; Moropon; Yanamono. Additional records: Napo and upper Río Marañón (Cope, 1868); Pebas (Cope, 1869); Nauta (Cope, 1876).

Distribution. Amazonian regions of Ecuador, Bolivia, Peru, and Brazil, the

northern tier of NE Colombia and NW Venezuela, the Orinoco and the Guianas, and the Brazilian coastal forest.

Anolis laevis (Cope 1876)

Type Locality. Between Moyobamba and Puerto Balsa, Río Huallaga, eastern Peru.

Material. ANSP: Between Moyobamba and Puerto Balsa, Río Huallaga, eastern Peru.

Distribution. Known only from the type locality.

Anolis ortonii Cope 1868

Type Locality. Río Napo or Upper Río Marañón, Ecuador or Peru.

Material. MHNJP: Saur.0426, Pucallpa, Pv. Crnel. Portillo, Ucayali; TCWC: Iquitos; Moropon; Centro Unión; Mishana. Additional records: Napo or Upper Río Marañón (Cope, 1868); Nauta (Cope, 1876); Pebas (Boulenger, 1885).

Distribution. Amazon and Orinoco basins and the Guianas and the Brazilian coastal forest.

Anolis punctatus punctatus Daudin 1802

Type Locality. "South America."

Material. MHNJP: Saur.0201, Pucallpa, Pv. Crnel. Portillo, Ucayali; Saur.0898, Tingo Maria, Pv. Huánuco, Huánuco (first records from Peru).

Distribution. Amazonian Brazil and the Orinoco basin plus the Pucallpa and Tingo Maria regions of Amazonian Peru.

Anolis punctatus boulengeri O'Shaughnessy 1881

Type Locality. Canelos, Ecuador.

Material. TCWC: Centro Unión; Mishana; Moropon. Additional record: "Pebas" (Cope, 1871).

Distribution. Amazonian regions of Peru, Ecuador, Colombia, and Brazil, the Orinoco and the Guianas and the Brazilian coastal forest.

Anolis trachyderma Cope 1876

Type Locality. Nauta, Peru.

Material. TCWC: Centro Unión; 5 km NNE Iquitos; Indiana; Mishana; Moropon; Río Momon. Additional records: Pebas (Cope, 1885); Iquitos (Williams and Vanzolini, 1966).

Distribution. Known from Amazonian Peru between Nauta and Pebas plus the Orinoco basin and the Guianas.

Anolis transversalis Dumeril 1851

Type Locality. "South America," restricted to Sarayacu, Peru, by Williams and Vanzolini (1966).

Material. TCWC: Centro Unión; Mishana; Moropon. Additional records: Nauta (Cope, 1876); Pebas (Cope, 1885); Achinamisa; Río Huallaga; Bombo, Río Tapiche; Cashiboya; Río Ucayali; Cerro Azul; Iquitos; Pampa Hermosa, near mouth of Río Cushabatay; Río Ucayali valley; Sarayacu (Williams and Vanzolini, 1966).

Distribution. Western Amazonian region, from Loreto in Peru to the Rio Solimões in Brazil, including Ecuador and Colombia. In Ecuador it extends up the Andean foothills for an indeterminate distance.

Comment. The generalized statements on distribution of these species outside Peru are based on works by Peters and Donoso-Barros (1970) and Williams (1965, 1974, 1976) and are not necessarily supported by documented specimens.

LITERATURE CITED

- BARBOUR, T. 1934. The anoles II. The mainland species from Mexico southward. *Bull. Mus. Comp. Zool.*, 77: 121-155.
- BOCOURT, M. F. 1873. Mission scientifique au Mexique. *Rept. Livr.* 1.
- BOULENGER, G. A. 1885. Catalogue of the Lizards in the British Museum (Natural History), London, 2: 11-95.
- BURT, C. E., AND G. S. MYERS. 1942. Neotropical lizards in the collection of the Natural History Museum of Stanford University Stanford Univ. Publ., Univ., Ser., Biol. Sci., 8(2): 1-52.
- COPE, E. D. 1868. An examination of the Reptilia and Batrachia obtained by the Orton Expedition to Ecuador and the Upper Amazon, with notes on other species. *Proc. Acad. Nat. Sci. Philadelphia*, pp. 96-119.
- . 1869. Seventh contribution to the herpetology of Tropical America. *Proc. Amer. Phil. Soc.*, 11: 147-169.
- . 1871. Ninth contribution to the herpetology of Tropical America. *Proc. Acad. Nat. Sci. Philadelphia*, pp. 200-224.
- . 1876. Report on the reptiles brought by Professor James Orton from the middle and upper Amazon, and western Peru. *J. Acad. Nat. Sci. Philadelphia*, 8(2): 159-188.
- . 1885. Catalogue of the species of batrachians and reptiles contained in a collection made at Pebas, Upper Amazon, by John Hauxwell. *Proc. Amer. Phil. Soc.*, 23: 94-103.
- DIXON, J., AND P. SOINI. 1975. The Reptiles of the Upper Amazon Basin, Iquitos Region, Peru. I. Lizards and Amphisbaenians. Milwaukee Public Museum, No. 4, pp. 1-58.
- D'ORBIGNY, A. 1847. Voyage dans l'Amerique meridionale . . . pendant les annees 1826 . . . 1833. Tome cinquieme lere partie: Reptiles. 12 pp. 7, pls. Paris: P. Bertrand Strasbourg: V. Levrault (1835-1847).
- PETERS, J., AND R. DONOSO-BARROS. 1970. Catalogue of the Neotropical Squamata. Part II. Lizards and Amphisbaenians U.S. Nat. Mus., Bull. No. 297, pp. 43-69.
- VANZOLINI, P., AND E. WILLIAMS. 1970. South American Anoles: The Geographic Differentiation and Evolution of the *Anolis chrysoplepis* Species Group (Sauria: Iguanidae). *Arq. Zool. S. Paulo*, 19: 1-298.
- WILLIAMS, E. 1965. South American *Anolis* (Sauria, Iguanidae): two new species of the *Punctatus* Group. *Breviora, Mus. Comp. Zool.* No. 233, pp. 1-15.
- . 1974. South American *Anolis*: Three new Species related to *Anolis nigrolineatus* and *A. dissimilis*. *Breviora Mus. Comp. Zool.* No. 422, pp. 1-15.
- . 1976. South American Anoles: The species Groups. *Pap. Avuls. Dept. Zool. Sao Paulo*, 29(26): 259-268.
- WILLIAMS, E., AND P. VANZOLINI. 1966. Studies on South American Anoles. *Anolis transversalis* A. Duméril. *Pap. Avuls., Dept. Zool. Sao Paulo* 19, Art. 17: 197-204.

Figure 1. Distribution of *Anolis* in Peru.

Figure 2. Distribution of *Anolis* in Peru.

Figure 3. Distribution of *Anolis* in Peru.

Two General Problems Involved in Systematics and Zoogeography of Bats

KARL F. KOOPMAN¹

ABSTRACT. It is difficult to determine true sympatry (at time of mating) in species which are migratory and at the same time have an extended gestation period. The basis for the ecological peculiarity of species which are confined to mountainous regions, but not to high altitudes, is not readily apparent.

INTRODUCTION

My first contact with bat taxonomy and biogeography resulted from an identification problem brought to my attention by Ernest E. Williams, which eventually resulted in a joint paper (Koopman and Williams, 1951). My first field experience with bats was on the same collecting trip to Jamaica, in 1950, that was Ernest's first field experience with *Anolis*. Since then, I have studied the systematics and distribution of bats in many parts of the world. In the course of these studies, two general problems have emerged which, to my knowledge, have not been generally recognized. I would like to discuss these on a more or less theoretical level since I think they have some general implications for evolutionary studies.

HOW DOES ONE DETERMINE SYMPATRY IN MIGRATORY BATS?

Bats, having the ability to fly, are able to cope with the problem of annual fluc-

tuations in climate (winter vs. summer, wet vs. dry seasons) by changing their ranges. In this they resemble migratory birds, though as far as we know, no bats migrate as far as some birds do. However, unlike birds, bats are viviparous mammals with a fairly long gestation period. Moreover, a number of bats have various mechanisms for sperm storage, delayed implantation, or delayed development (Orr, 1970; Wilson, 1979). This more or less extended gestation period may mean that some months will elapse between mating and the birth of young. During this period, the males, females, or both, may fly hundreds of miles.

In birds, it is common to find two or more subspecies occurring sympatrically in the winter range. (Many examples may be found in the American Ornithologists' Union's *Checklist of North American Birds* [1957]. A good example is found among the subspecies of *Sturnella magna*.) However, what is important is that these subspecies are allopatric in the breeding season. In birds, the range during the breeding season is quite simple to determine once nests are found. It can then be safely inferred that mating took place nearby since birds are oviparous and sperm storage is unknown (W. E. Lanyon, personal communication). However, as indicated above, no such inference may be drawn in the case of bats if there is any significant migration. Two taxa may be sympatric during part of the year, allopatric during another part; however, without direct evidence of

¹Department of Mammalogy, American Museum of Natural History, New York, New York 10024, U.S.A.

mating, it is impossible to determine whether or not these two taxa have an opportunity to interbreed. Obviously, the presence of reproductive isolation, the crux of the biological species concept, can only be determined if there is an unrealized opportunity to interbreed.

A few specific cases may point up the problems. *Lasiurus borealis borealis* is a common migratory bat in eastern North America. In the southeastern United States there is another form, *seminolus*, differing chiefly in color. Originally described as a subspecies of *L. borealis*, *seminolus* was considered a distinct species by Barkalow (1948) based on the fact that pregnant females and females with newborn young of both forms could be found sympatrically in Alabama. Also, they apparently both occur throughout the year in Louisiana (see Lowery, 1974: 114–121). The taxonomic status of these two forms is far from clear (see Koopman *et al.*, 1957), and it is quite possible that they are separate species, but the point to be made here is that sympatry at time of parturition is not evidence of sympatry during mating in migratory bats and therefore is not relevant to the question of whether or not separate species are involved.

Tadarida brasiliensis is a widespread New World bat some of whose northernmost populations in the southern half of the United States migrate and others do not (see Cockrum, 1969; Spenrath and LaVal, 1974, and references therein). Some of the authorities cited have considered that more than one species is involved, but the discussion is clouded by uncertainty as to where mating actually takes place. To my knowledge, no one has demonstrated two different populations of the *Tadarida brasiliensis* group mating in the same place at the same time, and it is only this kind of evidence that could establish that more than one species is involved.

In the genus *Leptonycteris*, two species are currently recognized in North America (Davis and Carter, 1962; Baker

and Cockrum, 1966), *nivalis*, and a taxon variously known as *sanborni* or *yerbabuena*. These two forms were originally considered as subspecies, and they are allopatric at the northern ends of their ranges (southwestern Texas and southern Arizona plus southwestern New Mexico respectively). Each occurs at its northern end in summer (where the young may be born) but is confined to more southern regions, chiefly in Mexico, during the winter. In Mexico, their winter ranges certainly overlap, at least macrogeographically, though judging by published records, the two species have not been collected at the same place at the same time. The authors of both the papers mentioned above concluded that the macrogeographical sympatry, which they documented, was evidence that species rather than subspecies were involved. I am inclined to agree with their taxonomic conclusions, but would point out that no one, to my knowledge, has demonstrated that *nivalis* and *sanborni* (or *yerbabuena*) mate at the same time and place and that therefore no true biological sympatry has been demonstrated.

The demonstration of true biological sympatry between closely related taxa in many bats will not be easy. Numerous species living in areas with marked seasonal fluctuations are known or suspected to be migratory to a greater or lesser degree. However, in most cases we have no clear picture concerning the distribution of a species at different times of the year. The best documentation we have for any species of bat with an extensive migration is that of Findley and Jones (1964) for *Lasiurus cinereus* in both North and South America. Cockrum (1969) has summarized what is known about migration (or its absence) in various populations of one subspecies of *Tadarida brasiliensis*. Tuttle (1976) has summarized the movement patterns of a species (*Myotis grisescens*) whose migrations are limited to movements between summer roosts (including maternity colonies) and winter hibernacula.

Hoffmeister (1970) has recognized the problem of summer vs. winter ranges of temperate North American bats and collected all data available to him for these seasonal ranges in the state of Arizona. There have been other studies as well, but these are the principal ones.

Assuming that the ranges of all migratory species can be determined for various times of the year, there is still the problem of when and where mating occurs. Most bats are small, nocturnal, and difficult to observe under natural conditions, so direct observation of mating is rarely possible, though this has been done in some cases (see Thomas *et al.*, 1979). Where a particular species is physiologically capable of mating only during a limited part of the year, indirect evidence such as presence of sperm in the epididymes of males and of active ovulation and fresh vaginal plugs in females, may fix the time and place of mating quite accurately. This kind of evidence is unfortunately available for very few species of migratory bats. The determination of sympatry in a true biological sense will probably entail work by many investigators operating in different places for a long period of time.

THE PROBLEM OF "PARA-MONTANE" DISTRIBUTIONS

During the course of determining geographical and altitudinal ranges of Peruvian bats (Koopman, 1978), I found a rather peculiar distributional pattern emerging in several cases. Though some 75 species were widespread in the Amazon basin, ten had essentially Andean distributions, ranging considerable distances along this mountain chain but not evidently extending far to the east of it. Five of these (*Sturnira erythromos*, *S. bidens*, *Vampyrops vittatus*, *Vampyressa melissa*, *Myotis keaysi*) are not known below 1,000 m (the semiarbitrary upper limit for the lowlands) in Peru, but five others (*Lonchophylla robusta*, *Sturnira*

bogotensis (= "*ludovici*"), *Vampyrops dorsalis*, *Artibeus harti*, *Myotis oxyotus*), although in most cases occurring in the highlands, are also known below 1,000 m. Even among the species apparently confined to the highlands in Peru, two species (*V. vittatus* and *M. keaysi*) are known from lower elevations in Venezuela (Handley, 1976). What keeps these five (or perhaps seven) Andean lowland species from extending their ranges across the Amazon basin as so many other eastern Peruvian lowland species have done? One possibility might be reduced rainfall farther east as the climatic influence of the Andes lessens. However, judging by the rainfall map that Haffer (1974: 17) gives, such is not the case. The only other explanation I can think of is that those species that are restricted to the vicinity of the Andes require caves or rock crevices for roosting and are not able to use trees for this purpose as the species that spread across the Amazon basin obviously can. Unfortunately, most collections of these Andean species have been made by netting and little or no data on roosts are available. It is, of course, possible that some presently unknown factor affecting forest composition and therefore food or roosting sites might explain this anomalous distribution pattern. However, this seems unlikely since species which feed on nectar, on fruit, and on insects and belong to two quite distinct families fall into this pattern. Another point to be emphasized is that bats actually live in two quite different environments, that of the roost and that of the foraging area. These may be separated by many miles and require quite different adaptations. Thus, different species may have similar roosting habits but forage under quite different conditions and vice versa. Where a number of species occur in the same general area, as in tropical America, it is difficult to determine which of these two general factors is more important in restricting numbers and distribution of a given species.

Unlike the problem of determining

true biological sympatry, that of paramontane distributions is not, of course, peculiar to bats. For any group of animals for which reasonably good distributional data are available, it would be interesting to look for species with fairly extensive distributions along a range (or ranges) of mountains that occur in adjacent lowlands but do not occur far from the mountains involved. Explanations for this kind of distribution could well be different in different cases. However, it would certainly be interesting to see how widespread this phenomenon is.

ACKNOWLEDGMENTS

I would like to thank Wesley E. Lanyon of the Department of Ornithology, American Museum of Natural History, for valuable information on birds, and Guy Musser, my colleague in the Department of Mammalogy, for critical reading of the manuscript.

LITERATURE CITED

- AMERICAN ORNITHOLOGISTS' UNION. 1957. Checklist of North American Birds, Edition 5. Baltimore, Maryland.
- BAKER, R. J., AND E. L. COCKRUM. 1966. Geographic and Ecological range of the long-nosed bats, *Leptonycteris*. J. Mammal., 47: 329-331.
- BARKALOW, F. S., JR. 1948. The status of the Seminole bat, *Lasiurus seminolus* (Rhoads). J. Mammal., 29: 415-416.
- COCKRUM, E. L. 1969. Migration of the guano bat, *Tadarida brasiliensis*. Univ. Kansas Mus. Nat. Hist. Misc. Publ., 51: 303-336.
- DAVIS, W. B., AND D. C. CARTER. 1962. Review of the genus *Leptonycteris*. Proc. Biol. Soc. Washington, 75: 193-198.
- FINDLEY, J. S., AND C. JONES. 1964. Seasonal Distribution of the Hoary Bat. J. Mammal., 45: 461-470.
- HAFFER, J. 1974. Avian speciation in tropical South America. Publ. Nuttall Ornith. Club, 14: 1-390.
- HANDLEY, C. O., JR. 1976. Mammals of the Smithsonian Venezuelan Project. Brigham Young Univ. Sci. Bull., Biol. Ser., 20(5): 1-91.
- HOFFMEISTER, D. F. 1970. The Seasonal Distribution of Bats in Arizona: a case for improving Mammalian Range Maps. Southwestern Nat., 15: 11-22.
- KOOPMAN, K. F. 1978. Zoogeography of Peruvian Bats with Special Emphasis on the Role of the Andes. Amer. Mus. Novitates, 2651: 1-33.
- KOOPMAN, K. F., M. K. HECHT, AND E. LEDECKY-JANECEK. 1957. Notes on the Mammals of the Bahamas with Special Reference to the bats. J. Mammal., 38: 164-174.
- KOOPMAN, K. F., AND E. E. WILLIAMS. 1951. Fossil Chiroptera collected by H. E. Anthony in Jamaica, 1919-1920. Amer. Mus. Novitates, 1519: 1-29.
- LOWERY, G. H. 1974. The Mammals of Louisiana and its Adjacent Waters. Baton Rouge, Louisiana State University Press, xxiii + 565 pp.
- ORR, R. T. 1970. Development: Prenatal and Postnatal. Biology of Bats, Vol. 1, 217-231. New York, Academic Press.
- SPENRATH, C. A., AND R. K. LAVAL. 1974. An Ecological Study of a Resident Population of *Tadarida brasiliensis* in Eastern Texas. Occ. Pap. Mus. Texas Tech. Univ., 21: 1-14.
- THOMAS, D. W., M. B. FENTON, AND R. M. R. BARCLAY. 1979. Social Behavior of the Little Brown Bat, *Myotis lucifugus*. I. Mating Behavior. Behav. Ecol. Sociobiol., 6: 129-136.
- TUTTLE, M. D. 1976. Population Ecology of the Gray Bat (*Myotis grisescens*): Philopatry, Timing and Patterns of Movement, Weight Loss during Migration, and Seasonal Adaptive Strategies. Occ. Pap. Mus. Nat. Hist. Univ. Kans., 54: 1-38.
- WILSON, D. E. 1979. Reproductive Patterns. Spec. Publ. Mus. Texas Tech. Univ., 16: 317-378.

Modes of Reproduction of Philippine Anurans

WALTER C. BROWN¹

ANGEL C. ALCALA²

ABSTRACT. Sufficiently complete data on characters of eggs, larvae, and egg deposition sites are available for 28 of the 65 species of anurans known from the Philippines to assign to them a specific mode of reproduction. Less complete but adequate data are available for 28 more species which enable us to assign the General Mode of Reproduction (either "aquatic" or "direct development") with a high degree of probability. Data on reproduction for the remaining nine species are so limited that even the General Mode must be considered as unknown. The diversity of modes of reproduction exhibited by Philippine anurans is discussed in terms of systematics. This diversity is compared to that of the tropical anuran community of 81 species in the area of Santa Cecilia, Ecuador as reported by Crump (1974).

INTRODUCTION

As currently understood, the Philippine anuran fauna is assigned to 65 species distributed in six families and 19 genera. Until relatively recently little was known about their life histories. Taylor (1920) and Inger (1954) published brief notes on eggs, larvae, or both for 14 species. Detailed life histories, however, were known for only two of these, *Polypedates leucomystax* (Villadolid and Rosario, 1930) and *Kaloula picta* (Cendaña and Fermin, 1940).

Since 1954, field and laboratory studies by the present authors have provided detailed life history data on 25% of the

species and data on eggs, egg deposition, larvae, or both for most of the remaining species. Only about half of this data has been published (Alcala, 1955, 1962; Alcala and Brown, 1956, 1957). It therefore seems appropriate at this time to summarize the available information on modes of reproduction of the Philippine anurans, and to compare the diversity with that exhibited by an anuran fauna of about the same size in a similar, tropical forest habitat. The anuran fauna of Santa Cecilia, Ecuador, reported by Crump (1974) best meets these criteria. The evolution and adaptive value of amphibian modes of reproduction are not considered in this paper. These topics have been treated in some detail in a number of early and recent papers (Lutz, 1947, 1948; Noble, 1927; Jameson, 1957; Orton, 1949, 1957; Salthe and Duellman, 1973; Salthe and Mecham, 1974; Crump, 1974; Wassersug, 1975; Lamotte and Lescure, 1977; Wake, 1978, 1980).

CLASSIFICATION OF REPRODUCTIVE MODES

As Salthe and Mecham (1974: 380) point out, the modes of reproduction of anurans are so varied and their evolution so poorly understood that no formal classification has been proposed, although they regarded that of Jameson (1957) to be the most complete up to that time. Examination of the general papers cited in the introduction illustrates this lack of any consensus as to a system of classifica-

¹1070 Fremont Street, Menlo Park, California 94025, U.S.A.

²Silliman University, Dumaquete City, Philippines.

tion. Salthe and Mecham themselves (1974: 380) recognize two primary categories, "Aquatic" and "Terrestrial," with four subcategories and seven sub-subcategories under "Aquatic" and two subcategories and six sub-subcategories under "Terrestrial." Salthe and Duellman (1973: 243) list nine anuran modes without subdivisions. Two of these involve "direct" development. Crump (1974) lists three primary categories based on egg deposition and larval habitat. She lists 10 specific modes exhibited by the Santa Cecilia frogs with three or four of the modes in each primary category. Six of the modes have aquatic larvae. Lamotte and Lescure (1977) classify anuran reproductive modes in six primary categories and 18 modes. The primary categories are largely based on egg deposition sites, larval habitats, and degree of parental care.

In the following scheme we use the two primary categories (General Modes) of Salthe and Mecham but recognize a greater number of Specific Modes (18). The Specific Modes are based on size and pigmentation of eggs, egg deposition, reduction or modification of larval stages and time of retention in egg capsules, larval nourishment, and type and degree of care by parents.

GENERAL MODE I—HAVE AN AQUATIC LARVAL STAGE

SPECIFIC MODE

- 1) Large to moderate number of small, pigmented eggs deposited in clusters in water. Larvae feed.
- 2) Moderate number of moderate-sized, pigmented eggs deposited in small groups attached to rocks on stream bottom or sides. Larvae feed.
- 3) Small to medium-sized eggs deposited in foam nests in water. Larvae feed.
- 4) Moderate number of small to medium-sized, pigmented eggs

deposited on rocks, sand, moss, leaves in close proximity to a body of water to which larvae find their way. Larvae feed.

- 5) Moderate to large number of small to medium-sized, nonpigmented eggs deposited in foam nests on objects overhanging water to which advanced tadpoles find their way. Larvae feed.
- 6) Small number of relatively large, nonpigmented eggs attached to wood above water in tree holes. Larvae fall into water where they develop and feed.
- 7) Small to moderate number of eggs deposited in water-filled leaf axils, tree holes, or other small volumes of water. Larvae feed.
- 8) Small number of relatively large eggs deposited in waterfilled leaf axils, pools or other small volumes of water. Larvae non-feeding.
- 9) Eggs deposited in small water-filled basins constructed in sand or gravel. Larvae feed.
- 10) Eggs deposited on land. Larvae carried to water. Larvae feed.
- 11) Small to moderate number of eggs and larvae carried in some manner by one of the parents until larvae enter water. Larvae feed.
- 12) Small to moderate number of eggs and/or larvae carried in some manner by one of the parents until larvae enter water. Larvae nonfeeding.

GENERAL MODE II—LACK AN AQUATIC LARVAL STAGE AND HAVE DIRECT DEVELOPMENT

- 13) Small to moderate number of large unpigmented eggs deposited in nest in terrestrial site. Tadpoles hatch in nest but never have an aquatic stage.
- 14) Small number of large unpigmented eggs deposited on land or

in leaf axils of trees, shrubs, epiphytic ferns. Froglets hatch directly from eggs.

- 15) Small number of large eggs carried externally in same way by one or the other of the parents until the froglets are released fully metamorphosed.
- 16) Gastric brooding through metamorphosis of small number of large eggs.
- 17) Small number of large eggs retained within the oviducts of the females, with ovoviparous development.
- 18) Small number of small to moderate-sized eggs retained within the oviducts of the female, with viviparous development.

RESULTS

REPRODUCTIVE MODES OF PHILIPPINE ANURANS

We have fairly complete life history data on 16 species of Philippine anurans, part published (Alcala and Brown, 1955; Alcala, 1962) and part in manuscript. We also have sufficiently extensive data on eggs and larvae of 12 additional species, including three for which the information is limited to populations from northern Borneo (Inger, 1956, 1966), to assign all 28 to a specific Reproductive Mode as recognized in the above classification. We know that another 10 species (including two Bornean populations reported by Inger) have aquatic larvae but we do not have data on the site of egg deposition and therefore cannot determine the Specific Mode. The information for these 38 species is summarized in Table 1.

Based on the habitat preference, clutch number, size, and pigmentation of approximately mature eggs dissected from gravid females as well as the general mode of reproduction of close relatives, we can reasonably assume the general mode for 18 more species of Philippine

frogs. Thus we assume direct development within the egg capsule for 11 additional species of *Philautus*, *Oreophryne*, and *Platymantis* (Table 2). The eggs of these species are very large, nonpigmented, and few in number, very similar to those of the species for which direct development is well established. Also the adults of these 11 species occupy similar terrestrial (usually arboreal), humid, forest habitats. Similarly, in terms of characters of the eggs and habitats of adults, we assume aquatic development for seven species of *Pelophryne*, *Rhacophorus*, *Rana*, and *Ooeidozyga*. Thus, of the 65 species of Philippine anurans only nine (*Barbourula busuangensis*, *Rana micrixalus*, *Platymantis cornutus*, *P. subterrestris*, *P. polilloensis*, *Micrixalus mariae*, *Philautus alticola*, *P. williamsi*, and *Rhacophorus hecticus*) are completely lacking in evidence as to their general reproductive mode. With the exception of *B. busuangensis*, these species are known from only one or two specimens and do not include gravid females. However, based on the uniformity of "mode" known or deduced for other Philippine members of the genera *Philautus*, *Rana*, *Platymantis* and *Rhacophorus*, as well as the habitats of the adults, we predict that seven of the nine species listed above will eventually be shown to exhibit a mode which is consistent with that of known Philippine species of the same genera. We make no prediction concerning *B. busuangensis* or *M. mariae*.

When we consider only the 38 species for which we have fairly complete evidence on the mode of reproduction, 31 (81.6%) have aquatic larvae and seven (18.4%) have direct development. When we consider the 65 known species, 39 (60.0%) very probably have aquatic larvae, 17 (26.0%) direct development and nine (14%) are unknown. The discrepancy in the percentages in the two comparisons is attributable to the much greater difficulty in obtaining reasonably

complete evidence of direct development. It is much easier to find aquatic larvae than it is to find eggs which are deposited in hidden sites not in close proximity to some body of water.

In terms of the two general modes, comparable data is available for the new world tropical anurans in the Santa Cecilia area, Ecuador (Crump, 1974). Of the 81 anurans recorded for that area, Crump reported 61 (75%) as having aquatic larvae, 17 (21%) as being direct developers, and 3 (4%) as unknown. These percentages for aquatic and direct development are close to those we found for the Philippine anurans.

At the level of the specific modes, I(1), I(2), etc., we find that we have sufficient data to enable us to so classify 28 of the 65 species (Table 1). Six specific modes are exhibited under aquatic development and one under direct development. Of the 28 species, 10 (35.7%) are type I(1); two (7.1%) I(2); two (7.1%) I(4); three (10.7%) I(5); one (3.5%), I(6); one (3.5%) I(7); and two (7.1%) I(8). Seven species (25%) are known to have direct development, all of type II(14).

Crump (1974) recognized 10 specific modes for the Santa Cecilia frogs as compared to seven for the Philippine anurans. Based on Crump's system, modes three (eggs and larvae in a constructed basin), six (eggs carried by parents, larvae aquatic), seven (eggs and larvae in terrestrial nest), and nine and 10 (eggs and young carried in various ways on dorsum of adults) are not represented among the known Philippine anurans. The correspondence of the modes of Crump to specific modes in our system and the proportions of species exhibiting each of the specific modes are shown in Table 3.

Even though the proportion of the Philippine anurans for which the specific mode is known is much less than that for the Santa Cecilia frogs, the evidence suggests similar trends in the diversity of reproductive modes exhibited by these two tropical anuran communities.

SYSTEMATICS AND MODE

Data on egg deposition sites, larval habitat, and general reproductive mode are summarized by family in Table 4.

Discoglossidae. Adults of *Barbourula busuangensis* are completely aquatic. However, the low number of large, unpigmented eggs in the ovaries of gravid females (Table 2) suggests a specialized mode of reproduction, possibly lacking an aquatic larval stage.

Pelobatidae. Only a single species in each of two genera, *Leptobrachium* and *Megophrys*, occurs in the Philippines. Both species have aquatic, feeding larvae. These are found in ponds or slow streams or in pools adjacent to them. Based on the fact that both species have eggs pigmented on one hemisphere, it is probable that they are deposited in the water or possibly in nearby locations from which the larvae can reach the water on hatching. Thus these two may be any of three specific modes, I(1), I(2), or I(3).

Bufo *idae*. Definitely five and probably all seven Philippine species have an aquatic larval stage. *Bufo marinus* and *B. biporcatus* use rain filled pools or slow moving streams for egg deposition. Larvae have also been collected from such pools (Inger, 1966: 63). The specific mode is I(1).

The larvae of *Ansonia* have been found in rapid mountain streams (Inger, 1954: 240). The mature eggs removed from gravid females are nonpigmented, but the egg deposition site is unknown. The specific mode is probably I(2) or I(4).

Details of the life history of the diminutive bufonid *Pelophryne brevipes* are in manuscript and verify Inger's observation (1960: 416) that the larvae are nonfeeding. The large, nonpigmented eggs and the small clutch size (8–12) of the other two Philippine species (*P. albotaeniata* and *P. lighti*) suggest that they may well have similar specific reproductive modes. This is mode I(8).

Ranidae. Twelve and most probably all

TABLE 1. PHILIPPINE ANURANS FOR WHICH GENERAL OR SPECIFIC REPRODUCTIVE MODE IS KNOWN.

Species	Site of egg deposition	Number of eggs	Eggs with dark pigment	Eggs without dark pigment	Aquatic larvae			Habitat of adults
					Direct development	Feeding	Reproductive Mode	
<i>Megophrys monticola</i>		400-500		X	X	I		semi-terrestrial
<i>Leptobrachium hasselti</i>		400-500	light	X		X	I	semi-terrestrial
<i>Bufo biporcatus</i>	In water	1000-2000	X		X	X	I(1)	semi-terrestrial
<i>Bufo marinus</i>	In water	thousands	X		X	X	I(1)	semi-terrestrial
<i>Pelophryne brevipes</i>	In water in leaf axils	5-20		X		X	I(8)	terrestrial-arboreal
<i>Ansonia mulleri</i>		100-200		X	X	X	I	semi-terrestrial
<i>Ansonia megregori</i>		400-500		X	X	X	I	semi-terrestrial
<i>Ooeidozyga laevis</i>	In water	100-150			X	X	I(1)	aquatic
<i>Rana cancrivora</i>	In water	2000-2500	X		X	X	I(1)	aquatic
<i>Rana erythraea</i>	In water	1200-1900	X		X	X	I(1)	aquatic
<i>Rana everetti</i>	On rocks							
<i>Rana leytensis</i>	in water	800-100	X			X	I(2)	semi-aquatic
	On leaves, rocks etc., near water	40-60	X			X	I(4)	semi-aquatic
<i>Rana limnocharis</i>	In water	1000-2000	X			X	I(1)	aquatic
<i>Rana magna</i>	On moss, rocks sand near water	140-200	X			X	I(4)	semi-aquatic
<i>Rana microdisca parva</i>		10-20	X			X	I	semi-aquatic
<i>Rana nicobarensis</i>		200-400	X			X	I	semi-aquatic
<i>Rana sanguinea</i>		180±	X			X	I	semi-aquatic
<i>Rana signata</i>	On rocks in water	about 300	X			X	I(2)	semi-aquatic
<i>Staurois natator</i>		200-300	X			X	I	semi-aquatic
<i>Platymantis dorsalis</i>	Soil crevices, epiphytic ferns	31-40		X	X		II(14)	terrestrial
<i>Platymantis guentheri</i>	Epiphytic ferns	8-47		X	X		II(14)	arboreal
<i>Platymantis hazelae</i>	Epiphytic ferns, leaf axils	5-9		X	X		II(14)	arboreal
<i>Polypedates leucomystax</i>	Foam nest	150-225		X		X	I(5)	semi-arboreal
<i>Polypedates macrotis</i>	Foam nest	300-400		X		X	I(5)	semi-arboreal

TABLE 1. CONTINUED

<i>Edwardtayloria picta</i>	On wall of tree hole above water	8 (one clutch)	X (gray)		X	I(6)	arboreal
<i>Edwardtayloria spinosa</i>	Probably as for <i>E. picta</i> (?)	20±	X (brown)		X	I	arboreal
<i>Philautus schmackeri</i>	Epiphytic ferns	6-15		X	X	II(14)	arboreal
<i>Philautus lissobranchius</i>	Epiphytic ferns and leaf axils of <i>Pandanus</i>	7-30		X	X	II(14)	arboreal
<i>Philautus sp.*</i>	Leaf axils of <i>Pandanus</i>	5-13		X	X	II(14)	arboreal
<i>Rhacophorus appendiculatus</i>		300-400		X	X	I	arboreal
<i>Rhacophorus pardalis</i>	Foam nest	44-50		X	X	I(5)	arboreal
<i>Chaparina fusca</i>	In water (small pools)	50-100	X		X	I(7)	semi-aquatic
<i>Kalophrynus pleurostigma</i>	In water (small pools)	500-1000		X		X	I(8)
<i>Kaloula baleata</i>	In water	several hundred	X		X	I(1)	terrestrial-arboreal
<i>Kaloula conjuncta</i>	In water	about 1000	X		X	I(1)	terrestrial-arboreal
<i>Kaloula picta</i>	In water	500-1000	X		X	I(1)	terrestrial
<i>Kaloula rigida</i>	In water	500-1000	X		X	I(1)	terrestrial
<i>Oreophryne annulata</i>	Under moss on trees	3-9		X	X	II(14)	arboreal

*This unidentified species also appears in Table 4 as one of the species for which larvae have not been identified.

16 species of the genus *Rana* have an aquatic larval stage. The known Philippine species exhibit three specific modes of aquatic development. *Rana leytenensis* and *R. magna* exhibit mode I(3), *R. everetti* and *R. signata* mode I(2), and *R. cancrivora* and *R. limnocharis* mode I(1) (Table 3).

Ooeidozyga laevis larvae have been collected from temporary ponds and mountain streams, and males and females congregate around such ponds during the rainy season. Alcala (1962: 686) records observing a pair in amplexus in a shallow ditch. Although eggs have not been seen in the turbid ponds (possibly because they are deposited at the bottom), dis-

section of gravid females reveals a clutch size of about 100 to 150 pigmented eggs. The specific mode is I(1) or I(2).

Inger (1966: 244) assigned three mountain stream larvae, two from Borneo and one from Mindanao, to *Staurois natator*. The deposition site of eggs has not been observed, but eggs from two gravid females are brownish pigmented and number 200-300. The specific mode is not known.

Definitely three (very probably four) and probably all 11 species of the genus *Platymantis* have direct development. The fourth species indicated as very probable is *P. corrugatus*. During field work on Negros in 1970, egg capsules

TABLE 2. PHILIPPINE ANURANS FOR WHICH GENERAL REPRODUCTIVE MODE IS UNKNOWN OR ONLY LISTED AS PROBABLE.

Species	Number of eggs*	Eggs with dark pigment	Eggs without dark pigment	General reproductive mode (probable)	Habitat of adults
<i>Barbourula busuangensis</i>	50-75		X	unknown	aquatic
<i>Pelophryne albotaeniata</i>	8-12		X	I	terrestrial-arboreal
<i>Pelophryne lighti</i>	9-12		X	I	terrestrial-arboreal
<i>Platymantis cornutus</i>	unknown			unknown	arboreal?
<i>Platymantis corrugatus</i>	44±		X	II	terrestrial
<i>Platymantis ingeri</i>	21-24		X	II	terrestrial-arboreal
<i>Platymantis insulatus</i>	12±		X	II	terrestrial
<i>Platymantis lawtoni</i>	20±†		X	II	arboreal
<i>Platymantis laevigatus</i>	9±		X	II	semi-terrestrial
<i>Platymantis polilloensis</i>	unknown			unknown	arboreal
<i>Platymantis subterrestris</i>	unknown			unknown	arboreal?
<i>Ooeidozyga diminutiva</i>	15±	X		I	aquatic
<i>Micrixalus mariae</i>	unknown			unknown	unknown
<i>Rana diwata</i>	100±†	X		I	semi-aquatic
<i>Rana melanomenta</i>	probably‡ hundreds	X		I	semi-aquatic
<i>Rana micrixalus</i>	unknown			unknown	semi-aquatic
<i>Rana woodworthi</i>	200-300	X		unknown	semi-aquatic
<i>Philautus acutirostris</i>	16-20		X	II	terrestrial-arboreal
<i>Philautus alticola</i>	unknown			unknown	unknown
<i>Philautus aurifasciatus</i>	12-15		X	II	terrestrial-arboreal
<i>Philautus emembranatus</i>	25±		X	II	arboreal
<i>Philautus leitensis</i>	10±		X	II	terrestrial-arboreal
<i>Philautus surdus</i>	10±		X	II	arboreal
<i>Philautus williamsi</i>	unknown			unknown	unknown
<i>Rhacophorus everetti</i>	30±§		X	I	arboreal
<i>Rhacophorus hecticus</i>	unknown			unknown	unknown
<i>Rhacophorus zamboangensis</i>	30-40		X	I	arboreal
<i>Oreophryne nana</i>	5±		X	II	arboreal

*Based on counts or approximations (over 100) of enlarged or mature ovarian or oviducal eggs.

†This is a rough estimate of eggs in the ovaries of the types of these species. The ovaries were not removed from the types.

‡The eggs in the only female available are in an early stage of maturation. We are assuming that they will all mature at one time.

§There were 15 large eggs in the ovaries and about 30 small to moderate. Whether or not all the latter will mature for the same clutch as the large ones is not known.

with early embryos were collected from a soil crevice. Though these were tentatively identified as *P. corrugatus*, the material was lost before this could be

verified. Eggs of the three well-documented species, (*P. dorsalis*, *P. guentheri*, and *P. hazelae*.) are deposited in such sites as leaf axils of *Pandanus*,

TABLE 3. COMPARISON OF SPECIFIC MODES OF PHILIPPINE AND SANTA CECILIAN FROGS.

	Specific Mode in our Scheme	Number of Philippine species (out of 28)		Corresponding mode in Crump's scheme	Number of Ecuadorian species (out of 78)	
Small, pigmented eggs in water. Larvae feed.	I(1)	10	(36%)	(1)	34	(44%)
Moderate sized pigmented eggs attached to objects in water. Larvae feed.	I(2)	2	(7%)	(4)	14	(18%)
Moderate-sized, pigmented eggs on leaves, mosses, etc. near water. Larvae feed.	I(4)	3	(11%)	(5)	6	(8%)
Relatively large eggs above water in tree holes. Larvae feed.	I(6)	1	(3+%)	not included	0	+
Moderate to large eggs in water in leaf axils, tree holes, etc. Larvae feed.	I(7)	2	(7%)	(2)	1	(1+%)
Relatively large eggs in water in leaf axils, tree holes, etc. Larvae nonfeeding.	I(8)	1	(3+%)	not included		+
Eggs in water-filled basins in sand or gravel. Larva feed.	I(9)	0		(3)	1	(1+%)
Eggs on land. Larvae carried to water by parents. Larva feed.	I(10)	0		(6)	5	(6%)
Eggs and larvae carried by parents till larvae enter water. Larvae feed.	I(11)	0		(9)	1	(1+%)
Large, unpigmented eggs in terrestrial nest. Tadpoles in nest but not aquatic.	II(13)	0		(7)	1	(1+%)
Large, unpigmented eggs in soil, leaf axils, ferns, etc. Direct development.	II(14)	7	(25%)	(8)	14	(18%)
Large eggs carried externally in same way by one of the parents. Direct development.	II(15)	0		(10)	1	(1+%)

epiphytic ferns and soil crevices. Four of the remaining seven species are known to have a small clutch of large, unpigmented eggs and the adults of all 11 spe-

cies occupy similar terrestrial habitats. An interesting morphological feature of the directly developing larvae of the known species is the use of expanded

TABLE 4. EGG DEPOSITION SITE AND LARVAL HABITS OF PHILIPPINE FROGS.

Family	No. of Species	Number undergoing Direct Development	Number with Aquatic Larvae	Unknown (lack evidence of eggs or larvae)	Deposition Sites for eggs (aquatic development)	Deposition Sites for eggs (direct development)	Larval habitats (aquatic development)
Discoglossidae	1			1	Unknown		
Pelobatidae	2	0	2		Unknown (probably in water)		2 species in mountain streams and pools
Bufonidae	7	0	4 feeding, 1 (+2 probable)* non-feeding		2 species in ponds 1 species in waterfilled leaf axils of Pandanus		2 species in ponds 2 species in mountain streams 1 species in water in leaf axils of Pandanus
Ranidae	29	3 (5 probable)	12 (5 probable)	5	2 species on mossy rocks	2 species in leaf axils of	4 species in temporary

	or banks, near water 5 species in water				ferns of Pan- danus and other plants 2 species in leaf axils or moist soil	or perma- nent ponds 8 species in streams
Rhacophoridae	18	3 (4 probable)	6 (2 probable)	3	3 species in foam nests on leaves of trees over water, or grassy banks	2 species in water in tree holes 3 species in stream pools
Microhylidae	8	1 (1 probable)	6		3 or 4 species in leaf axils of ferns of Pandanus 1 species at- tached to wood above water in tree hole	6 species in ponds or rain puddles
Totals	65	7 (18 with probables)	31 (39 with probables)	9	27	

*These probabilities are extrapolations based primarily on the egg number for gravid females, and habitats of adults as well as the mode exhibited by their close relatives occupying similar habitats.

abdominal sacs in respiration (Alcala, 1962). The specific mode is II(14) and is probably exhibited by all 11 species.

Rhacophoridae. Seven, probably eight species in three genera have aquatic larvae. *Polypedates leucomystax*, *P. macrotis*, and *Rhacophorus pardalis* are known to lay nonpigmented eggs in foam nests on leaves of trees or shrubs or grasses at the edge of forest streams or ponds. The hatching larvae find their way to the water where they complete development. These are examples of Mode I(5). Aquatic larvae, but not the site of egg deposition, are also known for *Rhacophorus appendiculatus* and *R. zamboangensis*. The lack of data on larvae and the lower number of relatively large unpigmented eggs in a gravid female of *Rhacophorus everetti* leaves the general mode of reproduction for this species in some doubt, although we list it as aquatic with a question mark (Table 2).

Edwardtayloria picta eggs have been found attached singly to the sides of water filled tree holes just above the level of the water (Inger, 1966: 352). Since larvae of the closely related *E. spinosa* in southeastern Philippines have been found in identical tree-hole microhabitats, it is very probable that the similar eggs (Table 3) are also deposited on the walls. This is mode I(6).

Three species in the genus *Philautus* have direct development, specific mode II(14), and probably at least four other species have the same mode based on characters of eggs and habitats of adults (Table 2). Detailed life histories of the three species noted above are in manuscript. Only *P. alticola* and *P. williamsi*, based on single specimens, are without even the indirect evidence of the type of egg or clutch size. The eggs of the three species reported at this time (Table 1) are deposited in leaf axils of epiphytic ferns or *Pandanus*. Like *Platymantis* in the Ranidae, the larvae do not show any evidence of external gills, oral disks, or coiled guts. They differ from *Platymantis*, however, in possessing highly vas-

cularized, broad tail fins which presumably function in respiration (Alcala and Brown, in manuscript).

Microhylidae. Six of the Philippine species have aquatic larvae. We have found eggs and larvae of *Chaparina fusca* and *Kalophrynus pleurostigma* in small temporary bodies of water in the forest (see also Inger, 1956: 397, 1966: 135). The larvae of *Chaparina* feed, mode I(7), but those of *Kalophrynus* do not, mode I(8).

Apparently all the Philippine species of *Kaloula* deposit eggs in ponds or similar bodies of water such as ditches. Life histories of *K. conjuncta* and *K. picta* have been well studied (Alcala, 1962: 715; Cendaña and Fermin, 1940). *Kaloula rigida* and *K. baleata* use small temporary pools for breeding (Taylor, 1922: 177; Inger, 1966: 125). Alcala (1962: 719) reported that larvae of *K. conjuncta* feed on algae. This is mode I(1).

In the genus *Oreophryne*, one species (*O. annulata*) is known to undergo direct development (Inger, 1954: 447). The highly vascularized tail is the respiratory organ. Most probably the second Philippine species *O. nana* has a similar reproductive mode, since the habitat of the adults and the mature eggs from a gravid female (Table 2) are so similar. The specific mode is II(14).

DISCUSSION

The evidence so far available for the Philippine anurans indicates there is little or no intrageneric diversity of mode of reproduction except for the genus *Rana*. The latter exhibits three specific modes. Otherwise, evidence suggests that a specific mode of reproduction is probably generically uniform and therefore useful in classification. For example, all known species of *Rhacophorus* and the two species of *Polypedates* exhibit mode I(5); all *Philautus* (*Rhacophoridae*) mode II(14), with the larvae using a vascularized tail for respiration; all *Platymantis*

(Ranidae) mode II(14), with the larvae using abdominal sacs for respiration. Also, in view of habitats of the adults and characteristics of the eggs and larvae (where the latter are known), it is very probable that *Oreophryne* and *Edwardtayloria* have uniform modes.

As shown in Table 4, aquatic larvae are known for 31 species and are assumed to be most probable for eight more species. These are distributed in five of the six families of Philippine anurans. Direct development is known for seven species (3 in Ranidae, 3 in Rhacophoridae, and 1 in Microhylidae) and is most probable for 11 more (5 each in Ranidae and Rhacophoridae and 1 more in Microhylidae), see Tables 1 and 2. The family with unknown mode is the Discoglossidae represented by the unique species *Barbourula busuangensis*.

The Ranidae with 29 species exhibits four specific modes. The Rhacophoridae (18 species), Bufonidae (7 species), and Microhylidae (8 species) each exhibits three specific modes. The Discoglossidae (1 species) and Pelobatidae (2 species) probably have only one specific mode.

In the Santa Cecilia fauna 61 species in seven of the eight families of anurans are known to have an aquatic larval stage. The tadpoles of one species in the Leptodactylidae develop through metamorphosis within a foam nest. Sixteen species (14 Leptodactylidae, 1 Pipidae, and 1 Hylidae) undergo direct development. The Hylidae with 37 species exhibits five specific modes, and the Leptodactylidae (25 species) four specific modes. The other families (Pipidae, 1 species; Bufonidae, 4; Dendrobatidae, 5; Centrolenidae, 3; Microhylidae, 5; Ranidae, 1) each exhibit only one specific mode.

In both areas the two families with the greatest diversity of species exhibit the greatest diversity of specific modes of reproduction. Also, with the exception of the Pipidae, direct development is exhibited only by families with relatively

high species diversity. These families, however, are different for the old and the new worlds. It should also be noted that direct development is limited to one specific mode in the Philippine fauna but three specific modes for the Santa Cecilia fauna.

ACKNOWLEDGMENTS

We would like to thank Lawton Alcala, Teodulo Luchavez, Crescencio Lumbod, and Braulio Gargar for assistance in the field. We also thank Marvalee H. Wake and William Duellman for critical suggestions on the manuscript and Robert F. Inger for data on Bornean populations of some of the Philippine species.

Our studies of Philippine amphibians and reptiles were supported by Grants from the National Science Foundation (the most recent, GB 41947).

LITERATURE CITED

- ALCALA, A. C. 1955. Observations on the life history and ecology of *Rana erythrae* Schlegel on Negros Island, Philippines. *Silliman J.*, 2: 175-192.
- . 1962. Breeding behavior and early development of frogs of Negros, Philippine Islands. *Copeia*, 1962: 679-726.
- ALCALA, A. C., AND W. C. BROWN. 1956. Early life history of two Philippine frogs with notes on deposition of eggs. *Herpetologica*, 12: 241-246.
- AND ———. 1957. Discovery of the frog *Cornufer guentheri* on Negros Island, Philippines, with observations on its life history. *Herpetologica*, 13: 182-184.
- CENDAÑA, S. M., AND D. V. FERMIN. 1940. Biology of *Kaloula picta* (Dumeril and Bibron) with special reference to its development and breeding habits. *Philip. Agriculturist*, 28: 626-655.
- CRUMP, M. L. 1974. Reproductive strategies in a tropical anuran community. *Univ. of Kansas Mus. Nat. Hist. Misc. Publ.* 61: 1-68.
- INGER, R. F. 1954. Systematics and zoogeography of Philippine amphibians. *Fieldiana, Zool.*, 33: 183-531.
- . 1956. Some amphibians from the lowlands of north Borneo. *Fieldiana, Zool.*, 34: 389-424.
- . 1960a. Notes on Toads of the genus *Pelophryne*, *Fieldiana, Zool.*, 39: 415-418.

- . 1960b. A review of the oriental toads of the genus *Ansonia* Stoliczka. *Fieldiana, Zool.*, **39**: 473–503.
- . 1966. The systematics and zoogeography of the amphibia of Borneo. *Fieldiana, Zool.*, **52**: 1–402.
- JAMESON, D. L. 1957. Life history and Phylogeny. *Syst. Zool.*, **6**: 75–80.
- LAMÔTTE, M., AND J. LESCURE. 1977. Tendances adaptives a l'affranchissement du milieu aquatique chez les amphibiens anoures. *Terre et la Vie*, **2**: 225–312.
- LUTZ, B. 1947. Trends toward non-aquatic and direct development in frogs. *Copeia*, **1947**: 242–252.
- . 1948. Ontogenetic evolution in frogs. *Evolution*, **2**: 29–39.
- NOBLE, G. K. 1927. The value of life history data in the study of the evolution of amphibians. *Ann. N.Y. Acad. Sci.*, **39**: 31–128.
- ORTON, G. L. 1949. Larval development *Nectophrynoides tornieri* (Roux) with comments on direct development in frogs. *Ann. Carnegie Mus.*, **31**: 257–268.
- . 1957. The bearing of Larval evolution in some problems in frog classification. *Syst. Zool.*, **6**: 79–86.
- SALTHE, S. N., AND W. E. DUELLMAN, 1973. Quantitative constraints associated with reproductive mode in anurans, pp. 229–249. *In* J. L. Vial (ed.), *Evolutionary Biology of the Anurans*. Columbia, Missouri, Univ. Miss. Press.
- SALTHE, S. N. AND J. S. MECHAM. 1974. Reproductive and courtship patterns in Physiology of the Amphibia, B. Lofts, (ed.), New York, Academic Press, **2**: 309–521.
- TAYLOR, A. C., AND J. J. KOLLROS. 1946. Stages in the normal development of *Rana pipiens* larvae. *Anat. Rec.*, **94**: 7–23.
- TAYLOR, E. H. 1920. Philippine Amphibia. *Philip. J. Sci.*, **16**: 213–359.
- . 1922. Additions to the herpetological fauna of the Philippine Islands, I. *Philip. J. Sci.*, **21**: 161–206.
- VILLADOLID, D. V., AND N. DEL ROSARIO. 1930. Studies on the development and feeding habits of *Polypedates leucomyax* (Gravenhorst), with a consideration of the ecology of the more common frogs of Los Banos and vicinity. *Philip. Agriculturist*, **18**: 475–503.
- WAKE, M. H. 1978. The reproductive biology of *Eleutherodactylus jasperi* (Amphibia, Anura, Leptodactylidae), with comments on the evolution of live-bearing systems. *J. Herpetol.*, **12**: 121–133.
- . 1980. The reproductive biology of *Nectophrynoids malcolmi* (Amphibia: Bufonidae), with comments on the evolution of reproductive modes in the genus *Nectophrynoides*. *Copeia*, **1980**: 193–209.
- WASSERSUG, R. J. 1975. The adaptive significance of the tadpole stage with comments on the maintenance of complex life cycles in anurans. *Amer. Zool.*, **15**: 405–417.

Reproductive Data on *Platemys platycephala* (Testudines: Chelidae) in Colombia

FEDERICO MEDEM¹

ABSTRACT. Mating in *Platemys platycephala* occurs mainly towards the end of the wet season. Egg deposition is primarily in the early dry season. A single enormous egg is laid. Incubation lasts approximately 150 days. Hatchlings are large, and subsequent growth appears to be slow. Maximum adult size may be around 160 mm.

INTRODUCTION

Platemys platycephala (Schneider 1792) is a flat-shelled, small chelid turtle, whose habitat comprises marshes, ponds, and creeks within the tropical rain forest of northern South America. Its range in Colombia includes the Colombian Amazon from Leticia to the Rio Atacuari, the upper Putumayo region (Rio Guamués), the upper Caquetá area (Florencia, Rio Ortegua), and the middle course of the Rio Apaporis which belongs to the Caquetá system. It has not yet been recorded from the Rio Vaupés drainage.

Almost nothing is known about the reproductive parameters of *P. platycephala*. According to information given by natives, Medem (1960) stated that it lays between one and six eggs. Dixon and Soini (1977) reported that two females killed for food by natives each contained one well-developed egg. They also reported that on July 31 copulation was observed in shallow water of a small forest stream at Centro Unión, near Iquitos,

Peru. Espinoza (1970) and Moll (1979) reported the clutch size to be four to six and five to seven eggs respectively, though without documentation. Ewert (1979) stated that an average egg measures 51×28 mm and that the hatchling carapace length averages 56.5 mm. No other reproductive information is available on this species.

This paper now provides observations on courtship and reproductive behavior, as well as new data on clutch size, egg size, incubation period, hatchling size, and subsequent growth rate. All observations were carried out in outdoor enclosures at the Instituto "Roberto Franco" in Villavicencio, Colombia.

COURTSHIP AND COPULATION

Courtship takes place during the rainy season (late March/April to November/December), day or night, principally in shallow water (5 cm) but also observed in water up to 50 cm in depth. Copulation occurs infrequently on the bank close to the edge of the water. Courtship is stimulated in captivity by running water, and the animals react immediately if the tank is cleaned by means of a hose. Reproductive behavior was observed on a total of 56 occasions. These occurred in the following frequency: in March, April, and May once each month; June twice; July seven times; August none; September twice; October eight times; November twenty-two times; and December twelve times.

¹Estación de Biología Tropical "Roberto Franco," Apartado aereo 22-61, Villavicencio (Meta), Colombia.

The male initiates courtship by persecuting the female until she is cornered, at which time he touches her face with the tip of his snout. He then climbs upon her carapace from behind, folding his tail under her shell. His forelegs continue to scratch the female's carapace dorso-laterally, until he can secure a grip on the sides of the carapace at the level of marginals 3 to 4. The male then hits and pushes the female's head until she is forced to seek relief by withdrawing her head within her shell, remaining motionless. In this position the male begins trembling motions of the head and neck. Copulation appears to take place at this time, but the exact duration is difficult to ascertain. The total time spent for courtship, mounting and copulation can last from four or five minutes to four and a half hours. The sequence can be interrupted briefly by both sexes in order to lift their heads out of the water to breathe. Successful intromission was specifically noted to occur on October 5 and 8 between the hours of 08:30–08:34 and 09:24–09:29.

EGGS AND HATCHLINGS

During egg deposition the female does not excavate a nest, but rather makes a shallow groove or lays her egg directly on the ground. The nesting site is always to be found under rotten leaves, and the egg is partially covered by sand or earth or not at all.

Between 1968 and 1975 ten episodes of egg deposition were observed during the months of August through February (Table 1). On each occasion only a single, hard-shelled, white oblong egg was laid. In comparison with the size of the female, the egg is enormous, and one wonders how it can pass the orifice of the cloaca. A female with a carapace length of 148 mm was observed to lay an egg measuring 57×28 mm, and an average egg measures 54.8×27.6 mm and weighs 27.0 gm. It is possible that more than one

egg may not have sufficient space within the reproductive tract. In addition, the body cavity volume of this flat-shelled turtle species is quite small, and a second egg of the same proportions could probably not be accommodated within the space available.

The ten eggs obtained were each incubated in the laboratory under rotten leaves at a temperature of 28 to 30°C, which corresponds to the ambient temperature. For unknown reasons, only four of the eggs hatched. The six others were found to be decomposed; two with small dead embryos, and four without embryos, possibly unfertilized. The incubation period lasts from 124 to 177 days (average 150 days). The four viable hatchlings had an average carapace length of 47.8 mm and a weight of 16.2 gm (Table 1).

An egg caruncle was present in two hatchlings only, and fell off after four and 35 days, respectively. The hatchlings were first kept isolated in the laboratory for about six months, and later put in a large enclosure with hatchlings of other species (*Phrynops*, *Rhinoclemmys*, *Kinosternon* spp.). One has survived for five years, the other three died after one or two years. Death was probably due to environmental stress, which was also observed in other chelid turtles (*Phrynops gibbus*, *P. nasutus*) when maintained under crowded conditions.

OTHER COMMENTS

Platemys platycephala apparently grows very slowly (Fig. 1), and evidently to a considerable age. A male (IRF 43—carapace length 146 mm) and a female (IRF 45—carapace length 148 mm) were kept in captivity for 22 and 15 years respectively. Bowler (1977) has recorded a specimen still alive in captivity at the age of nearly 20 years, and Mertens (1954) discussed a 29 year old individual.

The maximum size attainable by *P. platycephala* is unclear. Goeldi (1906)

Figure 1. Growth in captivity of 12 specimens of *Platemys platycephala* followed for one year or greater (average length 5.8 years, longest 22 years). Positions of individual curves interpolated on the graph to mimic a continuous growth curve. Carapace length (mm) plotted against elapsed time (years). Numbers on the graphs represent catalogue numbers from the IRF collection.

stated that the largest specimens were "never greater than 250 mm," and Ewert (1979) gave 200 mm as the size of a "large adult." The largest measured specimens, however, are a 163 mm male (Niceforo Maria, 1952) and a 161 mm male (Duellman, 1978). The largest individual examined in this series of 22 animals was a male measuring 161 mm (Table 2). Reproductively active females usually measure from 140 to 155 mm carapace length.

This species, known in Leticia as "matamata" (as are *Chelus fimbriatus* and *Phrynops nasutus*), can be considered amphibious, since it is frequently found walking on the ground far from water. In captivity it can remain for

several days under rotten leaves, and during the dry season it undergoes a prolonged period of estivation even if water is present. Both sexes are voracious, eating daily, and aggressively bite the necks, hind legs, and tails of cage mates.

It should be noted that whereas each clutch deposited by *P. platycephala* contains but a single egg, the possibility of multiple clutches being laid per season cannot be excluded, especially in view of the broad range of egg-laying dates.

ACKNOWLEDGMENTS

It gives me great pleasure to make a contribution to the collection of essays in

TABLE 1. REPRODUCTIVE DATA ON *PLATEMYS PLATYCEPHALA*.

Egg No.	Date of Deposition	Female Carapace Length (mm)	Female Weight (gm)	Egg Size (mm)	Egg Weight (gm)	Date of Hatching	Incubation Period (days)	Hatching Carapace Length (mm)	Hatching Weight (gm)
1	15 Feb 68	—	—	53 × 28	28.2	—	—	—	—
2	15 Nov 68	—	—	57 × 29	30.7	20 Apr	156	48	17.3
3	1 Feb 69	—	—	55 × 27	27.0	—	—	—	—
4	13 Feb 69	148	300	57 × 28	26.3	—	—	—	—
5	14 Dec 69	—	—	61 × 27	28.8	8 Jun	177	53	18.8
6	24 Nov 72	—	—	56 × 28	27.6	—	—	—	—
7	15 Jan 73	154	325	50 × 26	21.5	—	—	—	—
8	27 Nov 73	—	—	52 × 27	25.1	19 Apr	143	43	12.0
9	20 Jan 75	—	—	51 × 28	26.0	24 May	124	47	16.5
10	14 Aug 75	148	350	56 × 28	28.5	—	—	—	—

Figure 2. Egg and hatchling of *Platemys platycephala*.

TABLE 2. SIZE OF *PLATEMYS PLATYCEPHALA**.

IRF No.	Sex	Carapace Length (mm)	Plastron Length (mm)	Width (mm)	Height (mm)	Weight (gm)
653	m	161	144	99	46	277
238	m	161	—	98	40	—
769	f	158	143	106	46	392
44	f	155	142	103	43	350
555	m	152	131	97	43	325
239	f	150	—	96	39	—
764	m	149	131	93	41	263
45	f	148	134	103	45	350
690	m	148	128	92	43	258
43	m	146	130	95	42	350
350	f	141	—	93	42	—
42	m	138	—	93	38	—
809	f	134	124	91	41	263
810	m	125	113	83	36	222
366	f	115	101	83	33	150
43	m	110	—	75	35	—
814	f	105	95	74	30	129
770	j	98	86	63	26	82
860	j	85	77	63	24	68
815	j	79	72	61	22	76
870	j	75	70	57	21	46
2	j	61	52	45	17	22
2	h	53	45	36	17	18.8
870	j	52	45	38	15	13.5
1	h	48	40	32	17	17.3
860	h	47	39	39	18	16.5
815	h	43	36	29	16	12.0

*Carapace and plastron length measured in midline. Width is maximum value. Height is maximum value along midline.

honor of Ernest E. Williams, as I consider him one of the last of the great classical herpetologists and anatomists, as well as a good friend and mentor. Our association, which began at the MCZ in 1962, has always been intellectually stimulating and rewarding for me, and I owe much of my knowledge of reptile anatomy to him, especially as regards turtles. I owe him a special debt of gratitude for the constant generosity he has shown me in providing classic and recent bibliographies without which I could never have written some of my papers on turtles and crocodilians.

I also wish to express my gratitude to the following persons: Olga Castano Mora, Biologist at the Instituto "Roberto Franco," Alan Lieberman, Herpetologist

and Peace Corps Volunteer, for the preparation of the Tables and the correction of the text respectively, and to Anders and Susan Rhodin for the editing of the manuscript.

APPENDIX: SPECIMENS EXAMINED

IRF = Instituto "Roberto Franco," Villavicencio, Colombia. IRF 42-3, 653, 764, 769, 809-10, 814, 870, Leticia, Amazonas, Colombia; IRF 238, Larandia, upper Rio Orteguzaza, Caquetá, Colombia; IRF 44-5, Santa Rosa de los Kofanes, Rio Guamués, Putumayo, Colombia; IRF 555, Centro Unión, Ancaya, nr. Iquitos, Loreto, Peru; IRF 239, 350, Puerto Nariño, Amazonas, Colombia; IRF 690, Rio Caillaru, Loreto, Peru, ca. 40 km West Leticia, Colombia; IRF 366, Xeruiní, Rio Negro, Brazil; IRF 770, Uru-mutú creek, NE of Leticia, Amazonas, Colombia; IRF 1-2, 815, 860, born in captivity.

LITERATURE CITED

- BOWLER, J. K. 1977. Longevity of reptiles and amphibians in North American collections. *Soc. Stud. Amph. Rept. Herp. Circ.*, **6**: 1-32.
- DIXON, J. R., AND P. SOINI. 1977. The reptiles of the upper Amazon basin, Iquitos region, Peru. II. Crocodilians, turtles and snakes. *Milwaukee Publ. Mus. Contr. Biol. Geol.*, **12**: 1-91.
- DUELLMAN, W. E. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. *Univ. Kansas Mus. Nat. Hist. Misc. Publ.*, **65**: 1-352.
- ESPINOZA, N. C. DE. 1970. Contribucion al conocimiento de los reptiles del Peru (Squamata, Crocodylia, Testudinata: Reptilia). *Publ. Mus. Hist. Nat. "Javier Prado" Zool.*, **22**: 1-64.
- EWERT, M. A. 1979. The embryo and its egg: development and natural history. *In* M. Harless, and H. Morlock (eds.), *Turtles: perspectives and research*. New York, Wiley, pp. 333-413.
- GOELDI, E. A. 1906. Chelonios do Brazil (Jabotys—Kagados—Tartarugas). *Bol. Mus. Goeldi*, **1906**: 699-756.
- MEDEM, F. 1960. Datos zoo-geograficos y ecologicos sobre los Crocodylia y Testudinata de los rios Amazonas, Putumayo y Caqueta. *Caldasia* **8**(38): 341-351.
- MERTENS, R. 1954. Bemerkenswerte Schildkroten aus Sud- und Zentralamerika. *Aquar. Terr. Zeitsch.*, **7**: 239-242.
- MOLL, E. O. 1979. Reproductive cycles and adaptations. *In* M. Harless and H. Morlock (eds.), *Turtles: perspectives and research*. New York, Wiley, pp. 305-331.
- NICEFORO MARIA, H. 1952. Testudineos del suborden Pleurodira en el Museo de La Salle. *Bol. Inst. La Salle*, **39**: 14-21.

The Prehensile-Tailed Skink (*Corucia zebrata*) on Bougainville Island, Papua New Guinea

FRED PARKER¹

ABSTRACT. Little information has been recorded on *Corucia zebrata*. Four years of part-time field work and collecting on Bougainville Island has yielded data on habitat, live color, behavior, feeding, reproduction, predators and vernacular names. Reports indicate that *C. zebrata* may undergo periods of communal torpidity and coprophagy, but this has yet to be verified.

INTRODUCTION

From 1960 to 1963 and for a part of 1966, the author collected reptiles and amphibians on Bougainville Island, which, together with Buka Island and some offshore islets, comprises the North Solomons Province of Papua New Guinea. During that period, and through the following years while the author continued collecting in other parts of Papua New Guinea, Ernest E. Williams provided all the assistance, information, and copies of published material that a part-time field worker could ask for. I am grateful for his patience and perseverance, without which the value of my field collections would have been considerably less. The major part of my herpetological collections from Papua New Guinea and Australia are deposited in the Museum of Comparative Zoology, including the specimens referred to in this paper.

Corucia zebrata is a large and unusual skink (Fig. 1), but since its description by Gray in 1885, very little has been recorded about the species in the wild. Guppy (1887) reports that the species was said to live in the foliage of higher trees; Kinghorn (1928) repeats information from the village people about feeding and behavior; Schmidt (1932) comments on the suitability of the tail and claws for arboreal life; and Loveridge (1946) notes that a specimen was taken from the stomach of a crocodile.

Some specimens have reached zoos, and Honegger (1975) provides information on captive breeding, food, and infections.

DISTRIBUTION AND HABITAT

On Bougainville, *C. zebrata* is fairly common in the lowlands and foothills below 300 m elevation. In the Guava area, west of Kieta, it is reported by villagers to occur on the plateau at about 900 m. It is found in primary forest, including swamp and littoral habitat characterized by *Casuarina* spp. Preferred trees are those with dense foliage and extensive epiphytic growth. Since these are generally the oldest trees, *C. zebrata* does not occur in secondary forest where large trees have been removed as part of traditional gardening practices, or in areas selectively logged for timber.

¹717 Ross River Road, Kirwan Qld 4814, Australia.

Figure 1. *Corucia zebrata*.

Villagers from Buka Island report that *Corucia* is also to be found there, but it is restricted to the undisturbed forests on the low mountain ranges along the southwest coast.

Corucia zebrata is also found on all the larger islands in the Solomon Islands, from Choiseul to San Cristobal (McCoy, 1980).

DESCRIPTION

Corucia zebrata is a large skink, with a large and distinct head, small eyes, well-developed limbs, and digits with long curved claws. The tail is longer than the body, cylindrical and prehensile, and with a blunt tip.

Two gravid females in my series measured 31.5 cm snout-vent length plus

35.5 cm tail length (MCZ 78292 [part]) and 28.5 and 29.7 cm (MCZ 78393). Kinghorn (1928) records a larger specimen 70.5 cm total length and Schmidt (1932) one of 31.7 and 39.8 cm. The snout-vent length comprises 47 and 49% of the total length of my specimens, and 44.3% of Schmidt's. New born specimens measuring 17.8 and 17.5 cm (50.4%) (MCZ 78292 [part]) and 15.5 and 18.2 cm (46%) (Honegger, 1975) extend the range of relative body length, and there may be geographical variation or sexual dimorphism in this feature.

The normal color pattern for *C. zebrata* on Bougainville is pale to medium olive green on head and face with the dorsum olive-brown to brown with narrow irregular pale green to yellow-green bars, usually discontinuous at the vertebral line. The tail lacks pale bars but has

scattered paler scales. The dorsal pattern continues onto the flanks, where the pale bars widen and merge into the uniform pale olive or green ventral coloration. The limbs may have pale bars dorsally and laterally, or on the lateral surfaces only.

A number of specimens have scattered wholly or partly black scales on the post-cephalic dorsal and lateral body surfaces (see illustration in Parker, 1970). Loveridge (1946), Burt and Burt (1932), and Kinghorn (1928) also refer to this apparently common variation.

The iris is bright yellow. The pupil is black and, when contracted, wider above than below, with a small inferior "tail" which points postero-ventrally. This small "tail" is also present when the pupil dilates.

Hatchlings (e.g., MCZ 78292 [part] and 72919) are distinct in having the dorsum pale brown, almost translucent, with the head and venter paler, longitudinal stripes on the nape and transverse bars on the dorsum. There is no green pigment present at birth. I have not observed the ontogenetic change to adult color.

A mechanism for color change exists in this species. On 11 July 1962, at about midday, village people found two sub-adult skinks (MCZ 72916–17) in dense foliage at the extremity of a branch 15 m above the ground in a large tree in primary swamp forest. The branch was cut off and lowered by vines to the ground where I noted: "... the color of the head and the pale dorsal bars as well as ventral surfaces is dull pale blue ... the rest of dorsum and tail are normal brown." By the following morning: "... the bluish color is still present but blotches of normal green are appearing on the heads of both (skinks)—the blotches are irregular in shape and appear to be expanding from the center of each scale." By the evening of July 13, the heads, ventral surfaces and dorsal bars of both lizards were pale dull green. On the 15th, when both lizards were feeding, the same areas

were "... normal dull green but some of the dorsal head scales are darker green in the center, paler at the edges."

This method of color change is similar to that observed in captive sub-adult green tree pythons, *Chondropython viridis*, where the bright yellow, orange, or red juveniles change to green in 2 to 3 days, through a mechanism where a green speck first appears in the center of each scale, then slowly expands to cover the scale.

The pale transverse bars on the dorsum of *C. zebrata* may also become indistinct. An adult female in captivity (MCZ 78292 [part]) became uniformly dull brown dorsally over a period of two months. Honegger (1975) illustrates a strongly marked specimen newly arrived at Zurich Zoo and states the pale dorsal bars faded in captivity.

BEHAVIOR

This skink is arboreal, nocturnal, and according to village informants from a number of different parts of Bougainville, wholly vegetarian. By day it rests in an upright position in dense foliage, grasping twigs and branches with limbs and tail. It avoids sun and light and may shift its position if sunlight does reach it. McCoy (1980) states that it shelters in hollows in the larger forest trees.

When active, its movements are slow and deliberate, with the claws gripping climbing surfaces firmly. The tail is braced against trunks and larger branches, and coiled around smaller branches. Skinks sometimes hang suspended by the tail while feeding or moving. The tongue is used constantly to investigate the surroundings, and more frequently if the skink is disturbed. Its defensive behavior is very similar to that of the larger species of Australian and New Guinean *Tiliqua*—mouth wide open, tongue extended, body compressed to present maximum area to the "intruder," and rapid inhaling and exhal-

ing causing loud hissing. The bite is very powerful and prolonged. Hatchlings display this behavior more readily than adults.

By day these skinks are usually found alone or in twos. Some villagers report that they occasionally find three to five skinks of different sizes and sexes together. Groups of two taken like this were MCZ 72916–17, 93727–28, and 93729–30.

Village informants were consistent in asserting that *Corucia* will not willingly descend to the ground. During my many nocturnal collecting trips, not once did I find one of these skinks on the ground. McCoy (1980), however, states that "... it sometimes forages on the ground at night." On Bougainville, villagers say it will use aerial vines and branches in contact for movement between trees. They associate this skink with the possum, often finding the two animals in close proximity in foliage and in hollows in trees, by day and night. Kinghorn (1928) comments on this association.

Villagers report that this species eats the leaves, flowers, and fruits of various trees and epiphytes. Food is neatly nipped off, fruits in portions and leaves in small circular pieces, and each item swallowed. Kinghorn (1928) notes the stomach contents of one specimen as a mass of circular leaf particles.

The preferred food items on Bougainville are the leaves and fruit of an epiphytic *Scindapsus* vine with elongated leaves, known as *bakuna* in the Nasioi language in the Kieta area. The vine is illustrated in Parker (1970). The ripe fruits are red and sweet. The vine itself, however, has a white milky juice "with numerous needle-like crystals of calcium oxalate which should irritate mucous membranes" (Womersley, *in litt.*). A second common food plant is a wild pepper vine, *Piper* sp., known as *konangkong* in the Nasioi language. Again, leaves and fruit are eaten. Paradoxically,

the presence of the alkaloid piperine, especially in the fruits, should make it unpalatable (Womersley, *in litt.*). The arboreal agamid lizard also found on Bougainville, *Gonocephalus godeffroyi*, feeds readily on this *Piper*.

In captivity on Bougainville, these skinks remained vegetarian, rejecting meat and a variety of insects even if left without food for up to two weeks. Domestic fruits—tomatoes, bananas (skin and pulp) pawpaw, and pineapple—were taken, but all were neglected if either of the preferred food plants were placed in the cage. Villagers report that *Corucia* is not known to enter gardens in search of food. This may be because subsistence gardening as practiced on the island requires felling large forest trees and clearing undergrowth. Dietary preference may be another reason the skinks do not enter gardens. Honegger (1975) reports that the skinks at Zurich Zoo accepted apple, shoots of *Monstera* sp., and *Scindapsus aureus* (the preferred food of all those offered), but rejected bananas, peaches, vegetables, *Tradescantia*, *Philodendron scandens*, and *Codiaeum*. Honegger's specimens are also said to have taken cockroaches, maggots, mealworms and insects, but to have rejected fish. Specimens in the San Diego Zoo took insects and small mice and birds (Anonymous, 1978).

The captive skinks on Bougainville showed a preference for leaves covered in water droplets, and would not take water from containers in the cages. However, Honegger (1975) states that some of his specimens took water from containers at night.

Corucia defecates frequently and the feces have a characteristic odor. Honegger (1975) confirms this. Leaf fiber and portions of undigested leaves are visible in feces of newly captured specimens. Honegger (1975) observed specimens eating feces and portions of skin slough. He also noted that the feces of all

specimens had a high *Salmonella* content, and that dead skinks had high *Salmonella* and amoeba infestations.

COMMUNAL TORPIDITY

Corucia zebrata may exhibit communal torpidity at times. Kinghorn (1928) reports field information obtained by N. S. Heffernan on Ysabel Island:

Western natives assert that if there is more than one of these lizards in the same hollow tree, the topmost one only comes out to feed, the other underneath remaining and feeding on the excreta of the top one, which is always fat and healthy, the second one being thinner, and the third very thin.

People of six major language groups in areas where I collected on Bougainville also report this behavior with more details and embellishments. The reports seem to indicate that four to ten of these lizards, of all sizes and sexes, are occasionally found together in a hollow high in a forest tree. Sometimes there are also possums in the same hollow. The lizards at the top are well fed and healthy, while those below are progressively thinner.

As elsewhere in Papua New Guinea, the people traditionally attach anthropomorphic significance to animal behavior, and this apparent communal torpidity is explained by the people in the following fashion. When the lizards cannot find enough food on adjacent trees, and they cannot reach trees farther away to feed, they enter a tree hollow to sleep and wait for fresh growth of food plants. The lizards nearest the entrance to the hole commiserate with the others, saying how unfortunate it is that they are all without food. They offer to go out at night to see if there is any food available. They may find small quantities of food, and eat this, but when they return to the others they say they found nothing at all. Because they have had something to eat, they defecate, and remain fairly healthy.

The lizards below them in the hollow eat the feces, as there is nothing else for them.

Nothing in these stories indicates that there is a seasonal or even regular pattern to this behavior. During nearly four years of field work on Bougainville, such a group of lizards was never found, nor did village hunters report finding one.

Thus the actual occurrence of communal torpidity remains unconfirmed. From my experience on Bougainville and elsewhere in Papua New Guinea, traditional interpretations of animal behavior are almost invariably based on actual observations. Should such reports be obtained from different language groups within the range of the species concerned, it can be accepted that they are based on fact.

The reports on *Corucia* from six of the more than twenty different language groups of Bougainville and Buka Islands, and from Ysabel Island to the southeast in the Solomon Islands would have a high probability of being based on observed, but apparently infrequent, behavior. The coprophagy part of the reports has now been confirmed by Honegger (1975).

REPRODUCTION

One, or occasionally, two live young are born at a time, covered in a membranous sac from which they soon emerge and are immediately active and able to feed and fend for themselves. Breeding is reported by villagers to occur at any time of the year. One specimen born on 11 March 1961 measured a total of 33 cm, while another born on 17 April 1963 was 17.8 cm snout-vent, the tail 17.5 cm. Honegger (1975) reports a newly-born specimen of 15.5 and 18.2 cm length. The birth of twins is also recorded (Anonymous, 1978).

PREDATORS

On Bougainville, man appears to be the major predator of *Corucia*. Villagers do not specifically hunt these skinks, but frequently find them while searching trees for possums, a favorite subsistence food. When a hunter finds one of these skinks, he grasps it by the tail, and, avoiding its claws, swings its head against a tree to kill it. The body is cooked whole in the ashes of a fire. The skin is removed after cooking. If the lizard is to be kept alive, the hunter ties a length of vine to the end of a stick, makes a noose at the end of the vine, and places the noose around the skink's neck. He then lifts the lizard off the ground until it grasps the stick, and then binds the body, limbs, and tail to the stick with another vine.

Loveridge (1946) records an 18 inch *Corucia* taken from the stomach of an 11 foot crocodile (*Crocodylus porosus*) in the Shortland group, south of Bougainville.

VERNACULAR NAMES

In the areas where data collection was carried out, this was one of the better known species of reptile. This may be because of its large size, vegetarian diet, association with possums, use as food, or because it is the only viviparous lizard on the island. No information indicates that this species is a clan totem. Clan membership is inherited matrilineally on Bougainville, and members may not eat or touch their clan totem. The totem name is often used as the clan name. Other large reptiles, including the crocodile (*Crocodylus porosus*) and some of the marine turtles (*Dermochelys coriacea* and *Eretmochelys imbricata*) are clan totems in parts of Bougainville. *Corucia* appears to be absent from creation and

spirit mythology on Bougainville, whereas the crocodile and the ground boa (*Candoia carinata*) figure prominently.

TABLE 1. NAMES USED FOR *CORUCIA ZEBRATA* IN SOME OF THE LANGUAGES OF BOUGAINVILLE.

Kieta District	Nasioi Language	Katau
	Guava	Katau
	Eivo	Katakau
Buin	Buin	Katau, Kama
	Banoni	Takorai
Kunua	Kunua	Bokapuo
Tinputz	Tinputz	Kokorot

ACKNOWLEDGMENTS

I am grateful to J. Womersley, then Chief of the Division of Botany in Lae, for identification of food plants.

LITERATURE CITED

- ANONYMOUS. 1978. First time at the zoo. ZoonooZ (San Diego Zoo) February, 1978.
- BURT, C. E., AND M. D. BURT. 1932. Herpetological results of the Whitney South Sea Expedition. VI. Bull. Amer. Mus. Nat. Hist., 63: 461-597.
- GUPPY, H. B. 1887. The Solomon Islands and their Natives. London, Swan Sonnenschein, Lowrey & Co., i-xvi + 384 pp.
- KINGHORN, J. R. 1928. Herpetology of the Solomon Islands. Rec. Aust. Mus., 16(3): 123-178.
- HONEGGER, R. E. 1975. Beitrag zur Kenntnis des Wickelskinkes *Corucia zebrata*. Salamandra, 11: 27-32.
- LOVERIDGE, A. 1946. Reptiles of the Pacific World. New York, MacMillan, i-xii + 259 pp.
- MCCOY, M. 1980. Reptiles of the Solomon Islands. Wau Ecology Institute, i-vi + 80 pp.
- PARKER, F. 1970. Collecting reptiles and amphibians in New Guinea. Aust. Nat. Hist. (Australian Museum) March 1970, pp. 309-314.
- SCHMIDT, K. P. 1932. Reptiles and amphibians from the Solomon Islands. Field Mus. Nat. Hist. Zool. Ser., 18: 175-190.

Seasonal and Spatial Variation in the Annual Cycle of a Tropical Lizard

ROBIN M. ANDREWS¹

A. STANLEY RAND²

STELLA GUERRERO³

ABSTRACT. The population ecology of the small tropical lizard, *Anolis limifrons*, was studied over a two year period on two sites at Barro Colorado Island in Panama. Population density was the highest late in the wet season following recruitment of juveniles and the lowest early in the wet season prior to the hatching of eggs. Seasonal fluctuation in density is the result of low egg production during the dry season and annual population turnover. Seasonal changes in egg production appear to be the result of food or water stress, or both. Individual growth rates, physical condition of all individuals, and food intake by juveniles and adult females were lower in the dry than the wet season, paralleling seasonal lows in arthropod abundance. Variation in annual recruitment on Lutz site over four years was not related to food intake by adult females during the period when egg production is maximal. Population density varied spatially with lizards about one-half as abundant on Lutz as on AVA site. The high population density on AVA site was associated with relatively low female reproductivity, relatively poor physical condition, and low food intake. Egg mortality was higher on Lutz than on AVA site resulting in a lower net reproductive rate on Lutz than AVA. The discordance between population density and measures of lizard fitness on the two sites was the result of independence of egg and lizard fitness components.

INTRODUCTION

Population density of *Anolis limifrons* on Barro Colorado Island (BCI) in Panama fluctuates seasonally and from year to year. Low egg production during the dry season and annual population turnover produce the seasonal fluctuations (Sexton *et al.*, 1963, 1971; Andrews and Rand, 1974, 1982). As a result, population density is the highest late in the wet season following the major influx of juveniles into the population and lowest early in the wet season prior to the hatching of eggs.

Population density also fluctuates as much as sixfold from year to year. Andrews and Rand (1982) found a negative correlation between density late in the wet season and the length of the preceding dry season for their ten year study. Thus, both seasonal and year to year fluctuations have been associated with variation in rainfall. Two explanations have been proposed for this association.

The first explanation is that there is insufficient soil moisture for eggs to complete development successfully during the dry season, or desiccation is a problem for hatchlings, or both (Sexton, 1967; Sexton *et al.*, 1971). Thus, egg production is reduced during the dry season and recruitment is inversely related to the length and severity of the dry season.

¹Department of Biology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U.S.A.

²Smithsonian Tropical Research Institute, P.O. Box 2072, Balboa, Panama.

³Department of Entomology, University of California, Berkeley, California 94720, U.S.A.

Evidence to support this hypothesis is indirect, specifically that populations of *A. limifrons* are not food limited during the dry season (Sexton *et al.*, 1972). Although food is less abundant during dry season months, Sexton *et al.* (1972) found that the proportion of adult lizards with empty stomachs, the mean number of prey per lizard stomach, and prey size were similar in the dry and the wet season. Thus, Sexton *et al.* (1972) argued that because individuals get as much to eat in the dry season when food is scarce as in the wet season when it is abundant that food is not limiting.

A second explanation argued by Wright (1979) is that *A. limifrons* populations are food limited during the dry season when food is least abundant. Moreover, since variation in insect abundance is associated with variation in rainfall (Wolda, 1978a), the magnitude of annual recruitment should also be associated with food availability. Wright (1979) further pointed out that the food habit data collected by Sexton *et al.* (1972) was from a dry season that was unusually wet and suggested that even if food were not limiting when Sexton *et al.* (1972) sampled, it might be in most years when the dry season is more severe.

In this paper we describe seasonal changes in *A. limifrons* populations at two sites on BCI, one with a more dense lizard population than the other. The site of relatively low population density is the one on which density has been monitored for ten years (Andrews and Rand, 1982). Comparative data on population density, survival, recruitment, individual growth rate, reproductive condition of females, body size, and food intake were collected over a period of two years. Arthropod abundance was not measured. Since the plasticity of reptilian growth and egg production rates in response to changes in arthropod abundance is well documented (Andrews, 1976; Ballinger, 1977; Dunham, 1978; Smith, 1981), we assume that food intake and growth rates provide direct assessments of food availability.

We propose to evaluate the role of food as a factor regulating population density of *A. limifrons*. We will do so by determining the relationship between food availability and seasonal changes in egg production, year to year changes in annual recruitment, and spatial variation in density. The data analyzed here was collected as part of a study of the effects of malaria on *A. limifrons* (Rand, Guerrero, and Andrews, this volume.)

BARRO COLORADO ISLAND

The vegetation on BCI is classified as tropical moist forest (Holdridge and Budowski, 1956). Seasonality in rainfall is marked. The dry season (months with less than 100 mm of rain, Janzen, 1967) generally includes January through April (Rand and Rand, 1979). The dry season months are characterized by sharply decreased rainfall, decreased relative humidity, increased wind speed, and increased light in the understory as a result of partial leaf drop by canopy trees (Leigh *et al.*, 1982). The wet season includes May through December. We have arbitrarily divided the wet season into an early wet season (May-August) and a late wet season (September-December). The wet season often begins abruptly, and its onset is associated with such seasonal phenomena as leaf flushing of forest trees, reproductive activity of plants and animals, and peak insect abundance (Leigh, 1975; Wolda, 1978a, b; Robinson and Robinson, 1970; Rubinoff, 1974; Toft and Levings, 1979).

One of our study sites, Lutz, is an 890 m² area located in the Lutz watershed (Rubinoff, 1974). The other site, AVA, is a 990 m² area located at the 2,500 meter marker of Armour trail. These sites were established in November 1971 and May 1976, respectively, and they were located about 3 km distant from one another. AVA site was structurally similar to Lutz site. Although in most places the understory was sparse enough to walk through easily, both sites included several

clumps of dense and tangled vegetation resulting from tree falls.

METHODS

Thirteen censuses were conducted at each site (Fig. 1). Four minor censuses consisted of one or two consecutive days of collecting. Nine major censuses consisted of two consecutive days of collecting followed by two additional days of collecting six to seven days later. Lizards were collected in the morning (0930–1200 h). During each census, the sites were searched by two or three people and about 95% of all lizards seen were captured. We moved through the sites slowly, systematically examining all possible perches from tiny stems to large tree trunks from ground level to 2 to 2.5 m. Lizards were captured by hand and individually placed in plastic bags and transported to a laboratory where size, sex, and individual identification number were recorded. Snout-vent length (SVL) was measured to the nearest mm and weight (W) to one-hundredth of a gram. Individuals were identified by clipping the terminal phalanges of toes in unique combinations that required only one toe per foot. In addition, blood smears were taken from each female to assess reproductive condition. Color characteristics of Giemsa stained smears indicates individuals that are depositing yolk (Rand, Guerrero, and Andrews, this volume). Lizards were held in plastic bags for 24 h and then released at the place of capture. Scats produced during this period were dried at 60°C, weighed to one-tenth of a mg, and used to estimate food intake (see below).

Lizard classes were based on age, size, and reproductive attributes. Males and females on BCI both reach a maximum size of 49 to 50 mm SVL. Individuals 40 mm SVL or more were considered adult because most are sexually mature at these sizes (Sexton *et al.*, 1971). For some analyses, the adult class was partitioned into young adults (40–43 mm SVL) and

old adults (> 43 mm SVL). These adult classes were recognized because some individuals cease linear growth at 44 mm SVL and can no longer be aged by size. The juvenile I class consisted of individuals 29 mm SVL or less, and the juvenile II class consisted of individuals 30 to 39 mm SVL.

Total population size for the 1976–1977 censuses was estimated by the method of Jolly (1965). Major censuses provided two estimates of population size, and minor censuses provided one estimate. Bailey's modification of the Lincoln index (Poole, 1974) was used to estimate population size for the 1978 censuses.

Survivorship was determined as

$$S = 1 - (\ln N_0 - \ln N_t)/t$$

where S equals the daily survivorship rate, N_0 is the number of individuals present or known to be alive in an initial census, N_t is the number of individuals included in the initial census present or known to be alive in a following census, and t is the interval between censuses in days. Survivorship was expressed as S^{28} , the survivorship over a 28 day period. Although males and females were initially distinguished, preliminary analyses of survivorship did not detect differences related to sex. Therefore, survivorships were determined for two classes, adults and juveniles. Because home ranges of both juveniles and adults are quite stable (Andrews and Rand, 1983), we assume that few individuals disappearing from the censuses had emigrated. Jolly (1965) estimates for survivorship were not used because they do not distinguish age classes.

Survivorship curves were constructed for the cohorts hatched in the 1976 wet season. Because the high attrition of hatchlings precluded following one set of individuals through time, the curves were based on two age groups. Survivorship to 90 days of age was determined for all individuals 29 mm SVL or less at first capture. Survivorship from 90 days onward was determined for all juveniles

that reached at least 90 days of age. Juvenile ages were determined from growth rates of recaptured individuals.

Growth data from census records were fitted to the logistic growth model using the interval equation derived by Schoener and Schoener (1978). This equation is

$$L_2 = L_1 * a / (L_1 + (a - L_1) * e^{-rD})$$

where the estimated parameters are the rate constant (r) of the logistic growth equation and the asymptote (a). The data provided by the censuses were the size at first capture, L_1 , the size at second capture, L_2 , and the interval between captures in days, D . Estimates of r , a , and the 95% confidence intervals for both parameters were determined by non-linear regression procedures (Barr *et al.*, 1976). Snout-vent length was used as an index of size because weight gave poorer fits to the logistic model as judged by relative widths of the 95% confidence intervals.

The physical condition of adult *A. limifrons* was estimated as

$$IC = (W^{0.333}/SVL) * 100$$

where IC is the index of condition. For *A. limifrons*, the exponent of the power function relating W to SVL is 3 (Andrews, unpublished data). Therefore, to obtain an estimate of IC that was independent of size, W was raised to the 1/3 power.

Changes in the ratio between body weight and body length of adult individuals should reflect changes in their physical condition. High values of IC would be expected when females were producing eggs since eggs and maturing ova may comprise as much as 10% of female body weight (Andrews and Rand, 1974). High values would also be expected during periods of high food intake. Although levels of hydration have not been measured at different times of the year, it is possible that some differences in IC could be attributable to this source of variation.

Food intake was estimated from scat production as

$$I = SCTG * 2.94 * 22,773$$

where I is food intake in $J \cdot g^{-1} \cdot d^{-1}$, $SCTG$ is dry scat weight in g divided by lizard live weight in g , 2.94 is the correction factor which converts scat production into food intake (see below), and 22,773 is the energy value of food items in $J \cdot g^{-1}$. The 2.94 correction factor was determined from energy budgets of 23 adult female *A. limifrons* whose food intake and scat production were monitored daily over a 35 day period (Andrews, 1979). During this period, the mean ratio of dry scat weight to the dry weight of food eaten was $0.34 \pm 0.009(\bar{X} \pm SE)$. In order to determine if individuals captured during censuses would have exhibited similar ratios, 20 females were taken from their cages in late morning after they were fed and placed individually in plastic bags. They were then treated similarly to individuals captured in the field census. After 24 h scats were collected and the females returned to their cages. Each female was used as its own control in this experiment, using as the control value the ratio determined for the day prior to the experiment when the lizards were not disturbed. "Captured" females had ratios of 0.33 ± 0.042 , and the control females had ratios of 0.32 ± 0.024 ($p > 0.05$, paired comparison t-test). A conspicuous difference between the "captured" and the control females was that seven of 20 "captured" females did not defecate during the 24 h confinement, but only two of 20 control females did not defecate during 24 h ($p = 0.0637$, Fisher Exact Test). The low frequency of defecations by "captured" individuals was apparently the result of stress and not of lower food intake. "Captured" individuals that did not defecate had eaten similar quantities in the previous 24 h as those that did defecate (17.6 vs. 18.7 mg, respectively, $p > 0.05$, t-test). For this reason, individuals that did not defecate were not included in the an-

alyses of the census data. We assume that their food intake was the same as that of individuals that produced scats during their 24 h confinement.

Because *A. limifrons* eat a wide variety of small arthropods year round (Sexton, unpublished data; Andrews, unpublished data), it is probably appropriate to use a single energy value for their prey. Because the scat weights collected were not corrected for ash, the value of $22,773 \pm 556 \text{ J.g}^{-1}$ (det. by RMA) for adult crickets was used to represent prey items. The crickets had an ash corrected energy value of $24,577 \text{ J.g}^{-1}$ which was similar to the $24,171 \text{ J.g}^{-1}$ that Reichle *et al.* (1971) considered representative of 15 species of litter arthropods.

All mathematical and statistical manipulations of the SCTG data sets were conducted after natural log transformations. This procedure reduced the significantly ($p < 0.05$, Sokal and Rohlf 1969) skewed data sets (lizard class by census) from 39 to 15%.

RESULTS

POPULATION DENSITIES

During the 1976–1978 study period we observed seasonal and spatial differences in population density of *A. limifrons*. In 1976, population densities reached maximum levels in the late wet season following large increases in population density due to recruitment of juveniles (Fig. 1). The proportion of small juveniles on Lutz was the highest in July to August (Fig. 2). The low proportion of this class during the dry and the early wet season suggest most recruitment occurs within a relatively few months in the middle of the wet season. Proportional changes in class representation were similar on AVA site.

Recruitment was apparently higher in 1976 than in 1977, judging by the numbers of individuals present in the December 1976 to January 1977 and the January to February 1978 censuses, re-

spectively (Fig. 1). As pointed out by Andrews and Rand (1982), the high recruitment of 1976 was associated with a short preceding dry season (January to mid-April) and the low recruitment of 1977 was associated with a long preceding dry season (mid-December 1976 to mid-May 1977).

Spatial differences were pronounced. The density of the *A. limifrons* population on Lutz site was about half that on AVA site (Fig. 1). Densities reached respective maximum values of 1,150 and 1,860 individuals/ha in the late wet season of 1976 and respective minimum values of 260 and 660 individuals/ha in the early wet season of 1977. The maximum density observed on Lutz site during the study period was similar to densities observed in the late wet seasons of 1971 and 1978; 1,150 individuals/ha thus represents the maximal density attained on this site over ten years of censusing (Andrews and Rand, 1982).

SURVIVORSHIP

Survivorship of lizards on Lutz and on AVA site was similar overall ($p > 0.05$, Mann-Whitney U-test, Table 1). Adult survivorship on Lutz site averaged 0.725 per 28 days and did not vary appreciably through time ($p > 0.05$, 2×2 chi-squared tests for all pairwise comparisons of the 1976 to 1977 data). Adult survivorship on AVA site averaged 0.756 per 28 days, but in contrast to Lutz site, survivorship did vary temporally. Survivorship during the last two intervals of 1977 on AVA site was significantly lower than during the preceding four intervals ($p > 0.05$, 2×2 chi-squared tests for all pairwise comparisons). This change in survivorship had no obvious cause.

The similarity of adult and juvenile survivorship during each census interval (Table 1) was an artifact caused by lumping all juvenile survivorship data. The analysis of survivorship using cohort data indicated that the survivorship of individuals of 90 days of age or less was lower than the survivorship of individuals more than 90 days of age on both sites (Fig. 3,

Figure 1. The total size of *Anolis limifrons* populations on Lutz and AVA sites is shown in the top two graphs. Estimates are presented plus and minus their standard errors (see text for estimation procedures). Weekly rainfall (mm) is shown in the bottom graph. Incomplete censuses on AVA precluded population estimates for June 1976 and June 1978.

$p < 0.01$, 2×2 chi-squared tests). The low survivorship of the smallest juveniles was indicated by a significant ($r = 0.74$, $p < 0.05$) increase in survivorship with mean size (Table 1).

Survivorship curves of the 1976 cohort on Lutz and AVA sites were similar in their general appearance (Fig. 3). Survivorship to 90 days of age was 0.636 and 0.539 per 28 days, respectively. The largest juveniles and adults had survivorships on Lutz and AVA sites of 0.797 and 0.845 per 28 days, respectively. The probability of death was constant until about 210 days of age and then dropped

to 0.616 and 0.658 per 28 days, respectively. Although the cohort analysis was based on relatively few individuals, it confirmed the general results of the survivorship analyses based on census intervals (Table 1).

FEMALE REPRODUCTION

Old and young adult females had very different patterns of reproductive activity. Old females exhibited continuous oogenesis on both sites with 88–97% judged reproductively active (Table 2). Fewer young females were judged

Figure 2. Proportional representation of the five lizard classes on Lutz site during the period of most intensive censusing. The darkened bar on the time axis indicates dry season months.

Figure 3. Survivorship curves for *Anolis limifrons* populations on Lutz (●) and on AVA (○) sites. Sample sizes are shown for the initial cohorts at about 30 and 90 days of age. The ordinate indicates the natural log (ln) of the number of survivors with $N=1,000$ at $t=0$.

TABLE 1. SURVIVORSHIP OF ADULT (≥ 40 MM SVL) AND JUVENILE (< 40 MM SVL) *ANOLIS LIMIFRONS* ON LUTZ AND AVA SITES.

Daily survivorship has been converted to survivorship over a 28 day period (S^{28}). The total number of individuals counted in the initial census follows survivorship in parentheses. Survivorship was determined only when the initial number of individuals was 20 or more. Mean juvenile snout-vent length (SVL) shown is for the initial census.

Census Intervals LUTZ SITE			Survivorship		Juvenile Mean SVL
		(days)	Adults	Juveniles	
May-July	1976	(82)	.798 (27)	— (4)	—
July-Oct.	1976	(75)	.761 (29)	.670 (29)	25.3
Oct.-Dec.	1976	(49)	.658 (31)	.671 (24)	30.4
Dec.-Feb.	1977	(70)	.743 (44)	.786 (20)	34.7
Feb.-Apr.	1977	(73)	.736 (42)	— (14)	—
Apr.-June	1977	(70)	.613 (37)	— (11)	—
June-Jan.	1978	(204)	.745 (25)	— (6)	—
Jan.-May	1978	(119)	.752 (30)	.630 (21)	26.8
			$\bar{X} = .725$	.689	
AVA SITE					Juvenile Mean SVL
		(days)	Adults	Juveniles	
June-Sept.	1976	(80)	.838 (53)	— (2)	—
Sept.-Oct.	1976	(56)	.836 (50)	.655 (51)	29.7
Oct.-Jan.	1977	(85)	.842 (64)	.789 (45)	32.0
Jan.-Mar.	1977	(71)	.809 (116)	.858 (25)	33.7
Mar.-June	1977	(63)	.678 (100)	— (14)	—
June-Aug.	1977	(64)	.577 (66)	— (1)	—
Aug.-Feb.	1978	(183)	— (14)	— (10)	—
Feb.-June	1978	(140)	.723 (30)	.736 (23)	35.6
			$\bar{X} = .757$	.760	

TABLE 2. PERCENT OF ADULT FEMALE *ANOLIS LIMIFRONS* DEPOSITING YOLK DURING DRY SEASON (JAN. TO APRIL) AND WET SEASON (MAY TO DECEMBER) CENSUSES (SEE TEXT FOR DETAILS).

Class	LUTZ SITE				AVA SITE			
	Dry		Wet		Dry		Wet	
	%	N	%	N	%	N	%	N
Young adults (40–43 mm SVL)	59	(27)	78	(18)	36	(45)	74	(19)
Old adults (44 mm SVL +)	97	(38)	95	(60)	90	(68)	88	(94)

oogenic than old adults. This difference was significant in the dry season on both sites ($p < 0.05$, 2×2 chi-squared tests). More young adults on both sites were reproductively active in the wet season than in the dry season, but seasonal differences were significant only on AVA site ($p < 0.01$, 2×2 chi-squared tests) where 74% were judged oogenic in the wet season in comparison with 36% in the dry season.

The blood staining technique is largely insensitive to rate of egg production. It distinguishes reproductive individuals as those that have high levels of circulating lipoprotein associated with yolk deposition (Ayala and Spain, 1975). Old (large) females thus exhibited high levels of oogenic activity year round despite more than a twofold difference in egg laying rate between the wet and the dry season (Andrews and Rand, 1974). However, since young (small) adult females have an energy commitment to both growth and reproduction, oogenic events may not be continuous. Thus, the seasonal and site differences in the proportion of oogenic individuals may reflect either differences in egg laying rates, or age (size) of sexual maturity, or both.

GROWTH

Individual growth records, partitioned by sex, site, and season, were fit to a logistic growth model. Preliminary analyses showed that males and females did not differ either in their asymptotic size or their rate constants ($p > 0.05$, two-tailed t-tests). Therefore, only the results of site and season comparisons are pre-

sented (Fig. 4, Table 3). The mean maximum SVL (asymptote) attained was about 47 mm for all the fitted growth curves. In contrast, the rate (r) at which asymptotic size was reached varied considerably. The rate constant was greater in the wet season than the dry season on both sites ($p < 0.05$, two-tailed t-test). The magnitude of the difference was such that an individual hatched in the wet season reached sexual maturity (40 mm SVL) 30 days sooner than an individual hatched in the dry season.

Growth constants (r) were higher on Lutz site than on AVA site during both the wet and the dry season ($p < 0.05$, two-tailed t-tests). For example, during the wet season on Lutz site sexual maturity was reached at 93 days and on AVA site at 125 days. During the dry season, the difference was also about 30 days.

PHYSICAL CONDITION

Index of condition (IC) varied as a function of site, sex, and season (Fig. 5). A three-way analysis of variance using only the first record for each individual for a census period ($N = 917$) was used to test these main effects and all interaction terms. Season, sex, and site all had a significant effect on IC ($p < 0.05$). The interaction term between sex and time was also significant ($p < 0.001$). In general, the early wet season had the highest values of IC, the late wet season intermediate values, and the dry season had the lowest values. Thus, both males and females were in the poorest condition in the dry season and the best condition in the early wet season. Females generally

TABLE 3. FITTED PARAMETERS OF THE LOGISTIC GROWTH EQUATION OF *ANOLIS LIMIFRONS* POPULATIONS ON LUTZ AND AVA SITES.

The rate (r) constants and asymptotes (a) are shown followed by their 95% confidence intervals in parentheses.

	Season	N	r	a
LUTZ SITE	Wet	38	.0228 (.0194-.0262)	46.8 (46.3-47.6)
	Dry	62	.0177 (.0156-.0198)	47.6 (46.7-48.4)
AVA SITE	Wet	74	.0176 (.0156-.0196)	47.7 (47.2-48.3)
	Dry	144	.0135 (.0119-.0150)	47.3 (46.7-48.0)

Figure 4. Fitted logistic growth curves for *A. limifrons* individuals on Lutz and AVA sites during the 1976 wet and the 1977 dry season. Parameter estimates are given in Table 3.

Figure 5. Index of condition for *Anolis limifrons* on Lutz (●) and AVA (○) sites in different seasons and years. Seasonal means with the same letter are not significantly different ($p > 0.05$, Duncan's multiple range test). See text for details.

had higher values of IC than males on each site during all times of the year. Finally, values of IC on Lutz site were generally higher than those of AVA site for within sex comparisons.

FOOD INTAKE

The scat production method of estimating food intake was supported by an independent determination of food intake. In May and June 1977, adult females feeding *ad libitum* in captivity ate $284 \text{ J} \cdot \text{g}^{-1} \cdot \text{d}^{-1}$ (Andrews, 1979). In comparison, adult females censused on Lutz site in May and June to July 1977 had an estimated food intake of 265 and $304 \text{ J} \cdot \text{g}^{-1} \cdot \text{d}^{-1}$ (Fig. 6). Since anoles were probably not food limited on Lutz site during this period at least (see Discussion), the correspondence between the two sets of measurements is corroborative.

In order to evaluate the differences in food intake between the sites, lizard classes, and seasons, a three-way analysis of variance was used to test these main effects and all of the interaction terms. Censuses were combined into six consecutive seasons: early wet 1976, late wet 1976, dry 1977, early wet 1977, dry 1978, early wet 1978. In this analysis all main effects were significant, as was the interaction between season and class ($p < 0.001$). Differences between means were then tested with Duncan's multiple range tests. These results are discussed below.

Food intake was significantly higher on Lutz than on AVA site with respective

Figure 6. Mean food intake ($\text{J.g}^{-1}.\text{d}^{-1}$) for juvenile, adult female, and adult male *Anolis limifrons* individuals on Lutz (●) and AVA (○) sites. Sample sizes for each census are shown next to the means. Diamonds represent mean food intake for the previous year (data from Andrews and Asato, 1977). Food abundance was determined as the number of litter arthropods 2.5 to 23.5 mm in length collected in weekly 9.25 m^2 samples in Lutz watershed (S. Levings and D. M. Windsor, 1982, unpublished data). The values presented here are bimonthly means representing two (rarely three) collections.

means of 263 and 220 $\text{J.g}^{-1}.\text{d}^{-1}$ ($p < 0.001$). Juveniles, adult females, and adult males differed in their food intake with respective means across sites of 296, 261, and 194 $\text{J.g}^{-1}.\text{d}^{-1}$ ($p < 0.05$). The high food intake of juveniles was probably related to their weight specific metabolic rate (Bennett and Dawson, 1976; Andrews and Asato, 1977) and the energy costs of growth processes (Warren, 1971). The wide fluctuation in intake by juveniles may be the result, in part, of

seasonal changes in mean juvenile size (Fig. 2). The food intake of adult females was more similar to that of juveniles than to that of adult males. This suggests that the energy cost of egg production is high relative to the cost of reproduction by males. A comparison of class means for each site further illustrates both site and class differences. Respective means were 330, 293, and 209 $\text{J.g}^{-1}.\text{d}^{-1}$ for Lutz site and 272, 243, and 184 $\text{J.g}^{-1}.\text{d}^{-1}$ for AVA site. Only the comparisons of adult males

vs. adult females and juveniles vs. adult males differed significantly ($p < 0.05$, Duncan's multiple range tests).

Although the effect of season on food intake was highly significant in the complete model, only the early and the late wet seasons of 1976 could be distinguished ($p < 0.05$, Duncan's multiple range test). Inspection of food intake through time (Fig. 6) shows that the highest values for food intake for all classes were observed in that period. The pattern of temporal change in food intake differed between the three classes as indicated by a significant ($p < 0.001$) interaction between time and class.

DISCUSSION

RESPONSE TO SEASONALITY IN FOOD AVAILABILITY

Seasonal variation in arthropod abundance on Barro Colorado Island is documented. In general, abundance is lowest during the dry season and highest in the early wet season in association with the advent of the rainy season (Fig. 6; Robinson and Robinson, 1970; Rubinoff, 1974; Willis, 1976; Wolda, 1978b; Toft and Levings, 1979; Levings and Windsor, 1982). Depending on the arthropod taxon, abundance may gradually decline during the late wet season to dry season levels or abundance may reach seasonal lows as early as August (Rubinoff, 1974).

Wright's (1979) hypothesis that food is limiting during the dry season was supported by our data. Individual growth rates were lower during the dry than the wet season on both sites (Fig. 4). Both males and females had lower weight to length ratios during the dry than the wet season (Fig. 5). Food intake by adult females and juveniles was lower in the dry than in the wet season (Fig. 6). This variation in growth, physical condition, and food intake does parallel known seasonal changes in arthropod abundance on BCI. However, the conclusion that populations are limited by food shortage dur-

ing the dry season must be qualified by two additional considerations.

First, changes in food availability may have behavioral as well as physiological consequences. For example, food intake by adult males was similar year round. If food were limiting during the dry season, males should have increased their food intake during wet season months since females clearly were able to do so. On the other hand, dry season food shortage for males may have increased the time allocated to foraging as has been observed for *Anolis cupreus* (Fleming and Hooker, 1975). Food limitation thus might not be manifested in food intake *per se* but by other costs. For example, increased foraging activity may expose individuals to higher risk of being eaten by predators or may reduce effectiveness of territorial defense. Since females spend most of their activity period searching for prey year round (Andrews, 1971; Fleming and Hooker, 1975), seasonal changes in food intake by females should be more reflective of changes in prey availability.

Second, food may not be the limiting nutrient. Stamps and Tanaka (1981) have demonstrated that water availability overrides food as the limiting factor for growth of juvenile *Anolis aeneus* during dry season months. Presumably, other physiological processes such as oogenesis could be affected detrimentally by water shortage as well. Thus, a more appropriate conclusion for our study is that conditions during the dry season are sufficiently stressful to reduce growth rates and food intake of young individuals and to reduce rates of egg production and food intake of adult females.

RESPONSE TO YEAR TO YEAR VARIATION IN FOOD AVAILABILITY

In order to evaluate the effect of year to year variation in arthropod abundance (Rubinoff, 1974; Wolda, 1978b), on annual recruitment of *A. limifrons* we can compare food intake of adult females during the early wet season when oviposition rates are probably maximal (An-

drews and Rand, 1982) with annual recruitment as measured by population density at the end of the wet season. For 1975–1978, respective mean values for food intake by adult females on Lutz site in the early wet season were 471, 401, 287, and 305 $\text{J}\cdot\text{g}^{-1}\cdot\text{d}^{-1}$ (Fig. 6). The corresponding values for annual recruitment were 23, 100, 66, and 112 (1975 and 1978 population density data are from Andrews and Rand, 1982). Thus, year to year changes in food intake by females were not translated into numbers of surviving offspring.

The independence of annual recruitment and food intake by adult females suggests that annual recruitment may be more influenced by the survivorship of eggs (and possibly very small hatchlings) than by the rate at which they are produced. This argument is supported by the observation that survivorship of lizards did not vary appreciably over 10 years (Andrews and Rand, 1982) and that malaria parasites do not affect survivorship, growth, or fecundity of lizards (Rand, Guerrero, and Andrews, this volume). Moreover, observations on the water relations of eggs of *A. limifrons* suggest that moisture may not be limiting during the dry season in most years (Andrews and Sexton, 1981).

SPATIAL DIFFERENCES

Comparisons between our two sites suggest that the population at AVA site was food limited relative to the population at Lutz site. Individuals on AVA site grew more slowly, matured later, had lower food intake, and had a lower index of condition than individuals on Lutz site. Reproductive activity of young adult females was lower during the dry season and egg production rates of young and old females were presumably lower at AVA than at Lutz site.

The comparisons of food intake, growth, and index of condition suggest that lizards on AVA site were laying fewer eggs than lizards on Lutz site. This creates an apparent paradox. Lizard den-

sities were greater on AVA than on Lutz site despite poorer individual performance. Given a presumed lower reproductive rate and similar mortality schedules (Table 1, Fig. 3), population density on AVA should have declined relative to Lutz site.

However, this result may not be as paradoxical as first appears. For *A. limifrons*, lizard performance is transformed into population density through survivorship of eggs. Selection pressures in the egg environment (soil-leaf litter interface) and the lizard environment may be quite different. Moreover, optimal habitats for eggs and lizards may be found in different places. This model is supported by measurements of mortality rates of eggs (Andrews, 1982). During May and June 1981, eggs on AVA and Lutz sites had mortality rates of 56 and 82%, respectively, measured for a 40 day incubation period. A life table analysis showed that lower mortality during the egg stage accounted for the consistently higher population density on AVA than on Lutz site despite the apparent low fitness of lizards in the AVA population. We therefore suggest that spatial and temporal variation in egg mortality may produce the observed variation in lizard density from place to place and from year to year.

ACKNOWLEDGMENTS

This study was funded by National Science Foundation (NSF) Grants BO19801X and GB37731X to E. E. Williams and NSF Grant 76-05758 to A. S. Rand and S. Ayala. We thank N. Fetcher, S. J. Wright, and two anonymous reviewers for their comments on the manuscript, J. Birch of the Statistical Consulting Service of VPI&SU for statistical help, and S. Levings for allowing us to use unpublished litter arthropod data. The Fortran program used to determine the Jolly estimates of population size was written by G. G. Lawrence and R. H. Giles of the Department of Fisheries and Wild-

life at Virginia Polytechnic Institute and State University.

LITERATURE CITED

- ANDREWS, R. M. 1971. Structural habitat and time budget of a tropical *Anolis* lizard. *Ecology*, **52**: 262-270.
- . 1976. Growth rate in island and mainland anoline lizards. *Copeia*, **1976**: 477-482.
- . 1979. Reproductive effort of female *Anolis limifrons* (Sauria: Iguanidae). *Copeia*, **1979**: 620-626.
- . 1982. Spatial variation in egg mortality of *Anolis limifrons* (Iguanidae). *Herpetol.* **38**: 165-171.
- ANDREWS, R. M., AND T. ASATO. 1977. Energy utilization of a tropical lizard. *Comp. Biochem. Physiol.*, **58A**: 57-62.
- ANDREWS, R. M., AND A. S. RAND. 1974. Reproductive effort in anoline lizards. *Ecology*, **55**: 1317-1327.
- AND ———. 1982, in press. Long term changes in population density of the lizard *Anolis limifrons*, pp. 000-000. In E. G. Leigh, Jr., A. S. Rand, and D. M. Windsor (eds.), The ecology of a tropical forest: seasonal rhythms and long-term changes. Smithsonian Institution Press.
- AND ———. 1983. Limited dispersal of juvenile *Anolis limifrons*. *Copeia* (in press).
- ANDREWS, R. M., AND O. J. SEXTON. 1981. Water relations of eggs of *Anolis auratus* and *Anolis limifrons*. *Ecology*, **62**: 556-662.
- AYALA, S. C., AND J. L. SPAIN. 1975. Annual oogenesis in the lizard *Anolis auratus* determined by a blood smear technique. *Copeia*, **1975**: 138-141.
- BALLINGER, R. E. 1977. Reproductive strategies: food availability as a source of proximal variation in a lizard. *Ecology*, **58**: 628-635.
- BARR, A. J., J. H. GOODNIGHT, J. P. SAIL, AND J. T. HELWIG. 1976. A user's guide to SAS.76. SAS Institute Inc., Sparks Press, Raleigh, North Carolina, USA.
- BENNETT, A. F., AND W. R. DAWSON. 1976. Metabolism, pp. 127-224. In C. Gans (ed.), *Biology of the reptilia*, volume 5 (Physiology A). New York, Academic Press.
- DUNHAM, A. E. 1978. Food availability as a proximate factor influencing individual growth rates in the iguanid lizard *Sceloporus merriami*. *Ecology*, **59**: 770-778.
- FLEMING, T. H., AND R. S. HOOKER. 1975. *Anolis cupreus*: the response of a lizard to tropical seasonality. *Ecology*, **56**: 1243-1261.
- HOLDRIDGE, L. R., AND G. BUDOWSKI. 1956. Report of an ecological survey of the Republic of Panama. *Caribbean Forester*, **17**: 92-110.
- JANZEN, D. H. 1967. Synchronization of sexual reproduction of trees within the dry season in Central America. *Evolution*, **21**: 620-637.
- JOLLY, G. M. 1965. Explicit estimates from capture-recapture data with both death and dilution—Stochastic model. *Biometrika*, **52**: 225-247.
- LEIGH, E. G., JR. 1975. Structure and climate in tropical rain forest. *Ann. Rev. Ecol. and System.* **6**: 67-86.
- LEIGH, E. G., JR., A. S. RAND, AND D. M. WINDSOR (EDS.). 1982. The ecology of a tropical forest: seasonal rhythms and long-term changes. Smithsonian Institution Press, in press.
- LEVINGS, S., AND D. M. WINDSOR. 1982. Seasonal and annual variation in litter arthropod populations. In E. G. Leigh, Jr., A. S. Rand, and D. M. Windsor (eds.), The ecology of a tropical forest: seasonal and long-term changes. Smithsonian Institution Press.
- POOLE, R. W. 1974. An introduction to quantitative ecology. McGraw-Hill Series in Population Biology.
- RAND, A. S., S. GUERRERO, AND R. M. ANDREWS. 1983. The ecological effects of malaria on populations of the lizard *Anolis limifrons* on Barro Colorado Island, Panama. This volume.
- RAND, W. M., AND A. S. RAND. 1979. The rainfall of Barro Colorado Island, Panama, pp. 31-49. *Actas del IV Simposium Internacional de Ecologia Tropical*, Panama, Panama.
- REICHLER, D. E. 1971. Energy and nutrient metabolism of soil and litter invertebrates, pp. 465-477. In P. Duvingeaud (ed.), *Productivity of forest ecosystems*. Paris UNESCO.
- ROBINSON, M. H., AND B. ROBINSON. 1970. Prey caught by a sample population of the spider *Argiope argentata* (Araneae: Araneidae) in Panama: a year's census data. *Zool. J. Linn. Soc.*, **49**: 345-357.
- RUBINOFF, R. W. 1974. Environmental monitoring and baseline data (1973). Smithsonian Institution Environmental Sciences Program, Tropical Studies. Washington, D.C., Smithsonian Institution.
- SCHOENER, T. W., AND A. SCHOENER. 1978. Estimating and interpreting body-size growth in some *Anolis* lizards. *Copeia*, **1978**: 390-405.
- SEXTON, O. J. 1967. Population changes in a tropical lizard *Anolis limifrons* on Barro Colorado Island, Panama Canal Zone. *Copeia*, **1967**: 219-222.
- SEXTON, O. J., H. HEATWOLE, AND E. MESETH. 1963. Seasonal population changes in the lizard, *Anolis limifrons*, in Panama. *Am. Midland Nat.*, **69**: 482-491.
- SEXTON, O. J., E. P. ORTLEB, L. M. HATHAWAY, R. E. BALLINGER, AND P. LICHT. 1971. Reproductive cycles of three species of anoline lizards from the Isthmus of Panama. *Ecology*, **52**: 201-215.
- SEXTON, O. J., J. BAUMAN, AND E. ORTLEB. 1972. Seasonal food habits of *Anolis limifrons*. *Ecology*, **53**: 182-186.

- SMITH, D. C. 1981. Competitive interactions of the striped plateau lizard (*Sceloporus virgatus*) and the tree lizard (*Urosaurus ornatus*). *Ecology*, **62**: 679-687.
- SOKAL, R. R., AND F. J. ROHLF. 1969. *Biometry: the principles and practice of statistics in biological research*. San Francisco, California, W. H. Freeman and Co.
- STAMPS, J. A., AND S. K. TANAKA. 1981. The influence of food and water on growth rates in a tropical lizard (*Anolis aeneus*). *Ecology*, **62**: 33-40.
- TOFT, C. A., AND S. C. LEVINGS. 1979. Tendencias estacionales relacionadas con las poblaciones de arthropodos en la hojarasca, pp. 559-576. *Actas del IV Simposium Internacial de Ecologia Tropical*, Panama, Panama.
- WARREN, C. E. 1971. Biology and water pollution control (Bioenergetics and growth, Chapter 11). Philadelphia, Pennsylvania, W. B. Saunders Co.
- WILLIS, E. O. 1976. Seasonal changes in the invertebrate fauna on Barro Colorado Island, Panama. *Rev. Brasil. de Biologia*, **36**: 643-657.
- WOLDA, H. 1978a. Fluctuations in abundance of tropical insects. *Am. Nat.*, **112**: 1017-1045.
- . 1978b. Seasonal fluctuations in rainfall, food and abundance of tropical insects. *Journal of Animal Ecology* **47**: 369-381.
- WRIGHT, S. J. 1979. Competition between insectivorous lizards and birds in Central Panama. *Am. Zool.*, **19**: 1145-1156.

The Ecological Effects of Malaria on Populations of the Lizard *Anolis limifrons* on Barro Colorado Island, Panama

A. S. RAND¹

STELLA GUERRERO²

ROBIN M. ANDREWS³

ABSTRACT. Prevalence of malaria varied between two sites, with season, and was different in lizards of different sizes. High levels of parasitemia produced anemia, though without measurable effect on growth, reproduction, and survival of the lizards. *Anolis limifrons* and the *Plasmodium* species that infect it may exhibit a coevolved host-parasite relationship.

INTRODUCTION

This project was undertaken as part of a long term study to investigate the factors that influence population levels in a tropical organism. The results demonstrate an apparently well-adjusted relationship between a disease and its vertebrate host. Population fluctuations in the small, forest-understory, iguanid lizard *Anolis limifrons* (Fig. 1) on Barro Colorado Island (BCI), Panama have been reported and discussed several times over the past 14 years (Sexton, 1967; Sexton *et al.*, 1972; Wright, 1979; Andrews and Rand, in press). Attention has focused largely on the importance of food and water as controlling or limiting factors. When Guerrero *et al.* (1977) dis-

covered that these lizards had a high prevalence of malaria and that this varied from place to place on BCI, it seemed possible that this disease was influencing, if not controlling, *A. limifrons* population levels.

We set out to investigate this possibility by examining lizards at roughly monthly intervals at two small study sites on Barro Colorado Island. One site had a high lizard density and lower malaria incidence, the other had fewer lizards and more malaria. *Anolis limifrons* at the latter site had been censused for the past

Figure 1. *Anolis limifrons* from Barro Colorado Island, Panama.

¹Smithsonian Tropical Research Institute, Balboa, Panama.

²Department of Entomology and Parasitology, University of California, Berkeley, California 94720, U.S.A.

³Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U.S.A.

eight years (Andrews and Rand, in press). The ecological results of this study are reported in the preceding paper (Andrews, Rand and Guerrero, this volume); those dealing with malaria are described here.

Few studies have been carried out on the effect of parasites in wild populations (Anderson, 1976; Bennett *et al.*, 1976). Studies on the effect of helminths in populations of wild moose and deer have shown the different effects that a parasite can have in two sympatric host populations, the deer acting as an unaffected reservoir host and the moose being more susceptible and their population density limited by the parasite (Anderson, 1976). Other host-specific parasite infections affect only a single host species. There are only a few reports on the effect of protozoans on populations of birds, such as Warner's (1968) study on the effects of malaria on the Hawaiian honeycreepers.

Malaria in reptiles represents a good model for studies on the evolution and epidemiology of natural infections (Ayala, 1977). Reptiles make good subjects for such studies because they are sedentary and easy to capture so that the same animals can be examined several times. Following the first report that plasmodia occurred in New World lizards (Aragao and Neiva, 1909), many papers have been written on reptilian malaria (156 contributions from 1909-1975: Ayala, 1978). Lizards and snakes are the only reported hosts; of 87 species of New World lizards reported to be infected with malaria 34 are anoles (Ayala, 1977). Most reports have very little information on the ecology of the host and parasite, and there are no studies of the impact of malaria on lizard populations (Ayala, 1977). Ayala and Spain (1976) worked in western Colombia with *Anolis aeneus* infected with *Plasmodium colombiense*, finding fluctuations in the prevalence of malaria through a 12-month period. Scorza (1971) in a detailed study in Venezuela, on *Tropidurus torquatus* infected with *Plasmodium tropiduri*,

reported changes in hematology and in behavior; captive animals with high infections were less active and maintained lower body temperatures. Telford (1974, 1977) described the species of *Plasmodium* infecting anoline lizards at different localities in Panama. He showed fluctuations of natural populations of *Plasmodium* and suggested a relation to the ecology of the vector population. Jordan (1964) also reported fluctuations through the year and changes from year to year in *Plasmodium floridense* infecting *Anolis carolinensis* in Georgia.

Malaria may be an old disease of lizards, possibly much older than malaria in mammals (Ayala, 1977). Our findings suggest that the specific relationships of *Anolis limifrons* with the *Plasmodium* species that infect it are long-standing and the species closely coevolved. As parasitologists have predicted in such situations the disease seems to have little effect on its natural host. In our study, the prevalence of malaria varied between the two sites, with season, and was different in lizards of different sizes. Levels of parasitemia affected the blood picture and high levels produced anemia. However, in the variables which should be relevant to the population dynamics, such as growth, reproduction, and survival, we found little measurable effect of malaria, at least during the chronic phase of the infection.

THE TYPICAL COURSE OF A MALARIA INFECTION

To provide a background for the descriptions which follow, we present a brief review, largely from Ayala (1977), of the typical course of a malaria infection in a lizard. A detailed description of the *Plasmodium* infecting *Anolis limifrons*, and their development and behavior, (Guerrero and Pickering, in preparation) will be published elsewhere. Some information on the course of the infection in captive *A. limifrons* has been presented in Pickering (1980).

Malarial infections in reptiles follow the same general course as in mammals and birds: a rise, peak, decline, and chronic period, sometimes followed by a relapse. The vectors of malaria in reptiles are largely unknown. Ayala (1971) showed that the sandfly *Lutzomyia vextrix* was possibly the vector transmitting *Plasmodium mexicanum* to the iguanid lizard *Sceloporus occidentalis* in California.

The parasite has two phases, a sexual phase in the invertebrate vector and an asexual phase in the vertebrate host. *Plasmodium* gametocytes are picked up by the vector when it takes a blood meal from an infected host. Following fertilization in the vector the zygote may produce up to a thousand sporozoites (Ayala, 1977) which invade the hemocoel of the insect, some going to the salivary glands. When the vector bites a lizard the parasites are transferred when saliva is injected and the lizard becomes infected.

In the lizard, the sporozoites enter red blood cells (RBC's) and become trophozoites which grow, becoming either sexual forms (gametocytes) or asexual forms. Sexual forms remain quiescent until ingested by a vector. Asexual forms become schizonts and then segmentors, dividing several times to produce many merozoites which emerge, destroying the host cell, and infecting new RBC's.

During the first stage of the infection in the lizard the number of parasites in the blood cells is increasing and asexual forms are predominant. This phase of the infection is longer in reptiles than in mammals and birds and takes from one to two months to reach the peak of parasitemia (about 3–4 weeks in *A. limifrons*). The level of parasitemia at this point depends upon both the host and the species of *Plasmodium* (Ayala, 1977). (The highest we have seen in *A. limifrons* under natural conditions is 100 to 170 per 1,000 RBC's.) The peak of the infection is followed by a crisis period when the parasitemia can still be high and the host is responding to it by mas-

sive destruction of old RBC's and replacement by new, immature RBC's. Sometimes animals with high parasitemia die because of the severe anemia (Ayala, 1977). (This apparently happened to two of our captive *A. limifrons*.) The post-crisis period shows a decrease in the level of the infection with sexual forms becoming more common than asexual forms, indicating a low rate of parasite reproduction. After the crisis the infection enters a chronic period.

The chronic period is the longest one, and animals once infected may remain so for the rest of their life time (Ayala, 1977). Gametocytes are the common forms found in the RBC's and blood pictures are very similar to those of healthy individuals. During this period, the infection may be at so low a level that it is not detected in the blood smears. Also the final stages of schizogamy could possibly occur in the hemopoietic sinuses with subsequent reinvasion of the blood (Ayala and Spain, 1976). (Most of the *A. limifrons* caught in the wild with malaria showed chronic infections.)

A lizard with chronic malaria may suffer a relapse during which the parasites rapidly reproduce asexually and levels of parasitemia rise. Another crisis may then occur with high levels of immature RBC's and, if the lizard survives, a return of the infection to the chronic level.

MATERIALS AND METHODS

Intensive censusing, during which lizards were marked, bled, measured, and released, of two study areas on Barro Colorado Island, Panama, continued for about one year and less intensively for an additional year. During the study, 735 *A. limifrons* and 1,287 blood smears were examined. Many anoles were recaptured several times.

The study areas, census periods and measurement techniques have been described in detail in the preceding paper (Andrews, Rand, and Guerrero, this volume). In this paper we describe in

detail only those techniques specific to the malarial study.

Sites. Two sites were used: Lutz, characterized by a lower population density of anoles and higher incidence of malaria; and AVA, characterized by more anoles and less malaria. Both were in tropical forest.

Census. Lizard malaria information was collected during 13 censuses (Table 1). For several analyses censuses were grouped into wet and dry seasons. In some cases, wet and dry season data from different years were combined.

Class. For most analyses, five classes of lizards were recognized: juveniles (SVL < 40 mm), small males, small females (39 mm < SVL < 44 mm), large males, and large females (43 mm < SVL).

Blood smears. Blood was obtained from the tip of a toe when the terminal phalanx was clipped as part of the marking procedure. No more than one blood sample was taken from an animal during a census. It was difficult to bleed young animals without injuring them, so only in cases when the blood was easily obtained were blood smears made from animals less than 30 mm in SV length. Standard thin blood smears were fixed in methanol and stained for 40 minutes with 5% Giemsa solution pH 7.0–7.2 (Ayala and Spain, 1976). Slides were searched, at 1,000 times magnification, long enough to examine at least 20,000 RBC's; often the whole slide was searched before recording it as negative. Once a *Plasmodium*-infected cell was encountered, the microscope field was shifted randomly and 1,000 RBC's were examined, shifting the field systematically. The numbers of *Plasmodium balli* and *P. "tropiciduri"* present in these 1,000 RBC's were recorded. Following Telford (1977), parasitemia levels below 10 per 1,000 were considered low, probably indicating chronic infections.

Next, 100 RBC's were examined, and the number which were immature was recorded. The cytoplasm of an immature RBC is darker than that of a mature RBC's and the nucleus is bigger. Arbi-

trarily, immature RBC counts were considered high when they were equal to or greater than 10 per 100.

The blood smears made from female *Anolis* producing eggs show a very distinctive reddish coloration surrounding the red blood cells. This, as described by Ayala and Spain (1975), is due to the high concentration of lipoprotein that appears in the blood of reptiles at the end of the hydration state of follicle growth and is maintained during yolk deposition.

Species of Plasmodium. Four species of *Plasmodium* have been described from *A. limifrons* in Panama (Telford, 1974). Because controversy exists about the number and nomenclature of the species present (Ayala, 1976; Ayala and Spain, 1976; Guerrero *et al.*, 1977; Telford, 1977), we distinguished two morphospecies: *Plasmodium balli* belonging to the *Sauromoeba* group and *P. "tropiciduri,"* of the *Tropiciduri* group. In this latter category we included parasites that might have been classified as *Plasmodium floridense*, *P. tropiciduri* or *P. minasense* following Telford, 1974 and 1977 or more recently as a single undescribed species, *Plasmodium* sp. (Telford, 1979).

Plasmodium balli

This parasite is bigger than a red blood cell nuclei; it does not show malarial pigment in the cytoplasm and has a vacuole. It causes marked distortion to the host RBC or its nucleus (see Telford, 1969, for descriptions, and 1974, for figure, hosts and geographical distribution).

Plasmodium "tropiciduri"

This parasite is approximately the size of a RBC nucleus. It is oval or elongate in shape with malarial pigment in the cytoplasm, and sometimes a vacuole. These parasites do not visibly distort the host cell or its nucleus (see Telford, 1974, for descriptions, figures, hosts and geographical distributions of *P. floridense*, *P. tropiciduri*, and *P. minasense* and Telford, 1979 for a description of *P. sp.*). We be-

lieve that most, if not all, of our "*tropiduri*" infections involve a single species.

The insect vectors of these parasites has not yet been positively identified. The similarities in distribution of the two parasites seen in this study suggests that they share vectors, at least on Barro Colorado Island. The most likely candidate for vector is the phlebotomine fly, *Luzomya trinidadensis*, that feeds on lizards and is common in the study areas (Kimsey, in preparation).

Statistical analysis of the data was largely done at the Virginia Polytechnic Institute and State University using the Statistical Analysis System programs (Helwig and Council, 1979). Because of the unbalanced design, SAS general linear model procedures were used for the analysis of variance where the effects of several variables were to be considered together.

RESULTS AND DISCUSSION

The results are divided into two sections. First, we consider the prevalence of malaria, levels of parasitemia, and the species of *Plasmodium* as they varied with site, times of sampling, and lizard class as determined by lizard size and sex. Second, we examine the effects of malaria on the lizards considering its prevalence, the species of malaria, and the level of parasitemia on: 1) blood picture as measured by the proportion of immature red blood cells; 2) physical condition as indicated by the ratio of body weight to the cube of length; 3) food intake as estimated by scat weight; 4) time of activity as measured by hour of capture; 5) female reproductive activity as measured by presence of lipoproteins in the blood indicating oogenesis; 6) growth as measured by increase in snout vent length in recaptured animals; 7) survival as measured by recapture; 8) intensity of predation as estimated by tail breakage; and, finally, 9) differential mortality among the size classes as indicated by the size structure of the population.

THE EFFECTS OF SITE, CENSUS, AND CLASS ON MALARIA

Overall. The prevalence of malaria in *Anolis limifrons* on Barro Colorado Island was quite high. Plasmodia were found in 28% of the 1,287 blood smears examined. Of those with malaria, the majority (67%) had low parasitemias ($<10/1,000$), presumably indicating chronic malaria; the remainder had higher parasitemias, presumably indicating acute phase malaria. The prevalence of malaria is probably higher than these figures suggest. Cage experiments suggest that malaria is never lost but may be present at such a very low level that it is undetected in a blood smear (Guerrero and Pickering, in preparation). Some captive *A. limifrons* tested first negative and then positive for malaria even though the screened cages in which they were kept made the possibility of a new infection very unlikely. It is possible that the parasite may be absent from the circulating blood but present in other tissues. Even at levels below which we would reliably detect blood parasites, and we examined at least 20,000 RBC's before declaring a slide negative, a lizard could still be infective to a vector taking a blood meal. A phlebotomine fly typically takes between 0.3 and 0.5 mg of blood, a mosquito 2.0 mg (B. Chaniotis, personal communication) and a mg of *A. limifrons* blood contains about 76,000 RBC's so that a phlebotomine probably ingests at least 22,800 RBC's.

If malaria is never lost, we can estimate the number of undetected infections from the proportion of animals that tested positive for malaria on one capture that tested negative the next time they were examined. At both Lutz and AVA this is about 50%, suggesting that there are about as many undetected malaria infections among the lizards that we examined as there are infections that we detect. This estimate is supported in another way. Half of the 309 blood smears made from large adult *A. limifrons* at Lutz were positive for malaria; 79% of the 75 indi-

viduals that were tested two or more times as large (43 mm < SVL) adults were positive for malaria at least once; 95% of the 36 tested three or more times; and all of the 14 tested four or more times. Probably almost all of the large anoles at Lutz and almost half of those at AVA had malaria at some time during their lives and probably continued to harbor the parasite throughout their lives.

The two morphospecies of malaria, *P. balli* and *P. "tropiciduri"*, were found alone and in combination. Overall (N = 370), mixed infections were the most commonly seen (54%), pure *P. balli* the next most common (34%), and pure *P. "tropiciduri"* the least (13%).

The effects of site, census and class were considered separately, and three aspects of malaria were analyzed: the prevalence of malaria (Table 1), the frequency of high levels of parasitemia among infected animals (Table 2), and the relative abundance of the two species of *Plasmodium* (Table 3).

Site. The overall prevalence of malaria was consistently higher at the Lutz site (43%) than at AVA (19%) (N = 1287, $X^2_1 = 87.3$, $p < 0.0001$). The proportion of infections with high parasitemia were similar at both sites though somewhat higher at Lutz (36%) than at AVA (27%) (N = 370, $X^2_1 = 3.34$, $p = 0.067$). Mixed species infections were relatively more common at AVA (63%) than at Lutz (48%) (N = 370, $X^2_1 = 5.42$, $p = 0.034$). The two species of *Plasmodium* occurred in about the same proportions at the two sites (1 *P. "tropiciduri"* to 1.35 *P. balli* at Lutz, 1 to 1.29 at AVA).

Lutz had about half as many lizards as did AVA (Andrews *et al.*, this volume). This lower lizard density might be expected to reduce the chances of a vector transmitting the disease from one *Anolis limifrons* to another and so reduce the prevalence of the disease. This did not happen; at least 43% of the anoles at Lutz and 19% at AVA had malaria. We initially expected to find that the lower anole population levels at Lutz were the result of higher levels of malaria there. As will

be discussed below, we can not support this idea on the data at hand. Even if it were true, it would not explain why malaria was more common at Lutz than at AVA. If it is not an historical accident, possibly the result of sampling a multi-year disease cycle at different points at the two sites, we can only suggest that it might be due to differences in the two host populations in their susceptibility to the malaria or the result of differences in vector populations. The Lutz site is on a slope close to a small stream; AVA is on level ground much further from water. It is difficult to evaluate these differences until we identify the vector.

Census. For analysis by contingency tables censuses made during the same wet or dry season were combined (see Table 1 for details of the times of censusing and of seasons).

Prevalence of malaria changed over time from census to census in the course of the study (Table 1), with no correlation between sites (N = 12, $r = 0.12$). No clear seasonal pattern is evident, but our late wet season samples were high at both sites. Lutz showed a significant change among seasons censused (N = 517, $X^2_5 = 23.5$, $p = 0.015$) because of a very high prevalence in a single census; this did not occur at AVA (N = 769, $X^2_5 = 13.6$, $p = 0.33$).

The proportion of the infections with high (> 9/1,000) or very high (> 19/1,000) parasitemia changed over time. There is a cross correlation between sites in these changes (N = 12, $r = 0.80$). High parasitemia was more common during the early wet season and became progressively less so throughout the rest of the wet season and into the dry season (Table 1). Looking only at the year of intensive censusing, the differences between seasons are significant at both sites (Lutz N = 178, $X^2_3 = 37.5$, $p < 0.001$; AVA N = 128, $X^2_3 = 26.0$, $p < 0.001$). The three successive early wet seasons sampled show successively lower proportions of high parasitemia (Table 1).

The high frequencies of high parasitemia presumably reflect high frequen-

TABLE 1. PREVALENCE OF MALARIA (THE PERCENTAGE OF LIZARDS EXAMINED THAT HAD MALARIA) AND LEVEL OF PARASITEMIA (THE PERCENTAGE OF INFECTIONS WITH HIGH PARASITEMIA IE. > 9 INFECTIONS PER 1,000 RBC'S) DURING SUCCESSIVE CENSUSES.

				<i>A. limifrons</i> with Malaria					Infections with High Parasitemia			
				Lutz		AVA		Lutz		AVA		
CENSUS				SEASON	N	%	N	%	N	%	N	%
6	May-14	Jun	76	Early Wet	36	41.7	31	25.8	15	93.3	8	62.5
30	Jun-6	Aug	76	"	33	45.5	45	17.8	15	80.0	8	62.5
1	Sep-21	Oct	76	Late Wet	43	46.5	55	21.8	20	40.0	12	25.0
27	Oct-19	Nov	76	"			89	27.0			24	12.5
	1-17	Dec	76	"	59	69.5	35	20.0	41	31.7	7	14.3
	12-27	Jan	77	Dry	33	42.4	135	22.2	14	28.6	30	20.0
	9-23	Feb	77	"	55	40.0	39	20.5	22	27.3	8	37.5
16	Mar-7	Apr	77	"	10	40.0	98	15.3	4	25.0	15	13.3
20	Apr-12	May	77	"	59	42.4	48	8.3	25	20.0	4	25.0
25	May-10	Jun	77	Early Wet	24	33.3	79	11.4	8	37.5	9	55.6
29	Jun-4	Aug	77	"	36	38.9	14	21.4	14	14.3	3	0
18	Jan-9	Feb	78	Dry	65	40.0	59	16.9	26	30.8	10	20.0
17	May-29	Jun	78	Early Wet	64	29.7	42	21.4	19	26.3	9	44.4

N = number examined.

cies of new infections or relapses. These coincide with periods of general insect abundance (see Andrews *et al.*, this volume) and may reflect times when the malaria vector is particularly abundant, rather than times when lizards are particularly susceptible to malaria. It is also a time when food is most abundant and when reproductive and social activity are probably at their highest and the anoles might be the most susceptible to malaria. As one might expect, the peak in new infections precedes the peak in total prevalence of malaria.

The year to year changes make the reality of the apparent seasonal pattern hard to assess and are themselves hard to interpret.

Class. The prevalence of malaria varies with size and sex class (Table 2). It is higher in males than in females, though significantly so only at AVA ($N = 629$, $X^2_1 = 19.1$, $p < 0.001$; at Lutz $N = 410$, $X^2_1 = 3.49$, $p = 0.06$), and higher in larger adults than in small ones (Lutz $N = 517$, $X^2_1 = 89.8$, $p = 0.001$; AVA $N = 770$, $X^2_1 = 36.7$, $p < 0.001$). At both sites large males are more likely to have malaria than is any other size class. Small adults have

lower incidences of malaria. At both sites large females are intermediate; at AVA they are more like large males and at Lutz more like small adults. At both sites juveniles are less likely to have malaria than are adults (Lutz $N = 517$, $X^2_1 = 62.7$, $p < 0.001$; AVA $N = 770$, $X^2_1 = 30.0$).

The differences between size classes are due, at least in part, to differences in age because once a lizard has been infected with malaria, it probably rarely, if ever, loses the parasite. It is not clear why males have a higher prevalence of malaria than do females. Perhaps males, because they tend to use higher and more exposed perches than do females (Talbot, 1979), come into more frequent contact with vectors.

Levels of parasitemia also vary between classes (Table 2). These differences are not significant if the sites are considered separately. However, at both sites the pattern is the same and if they are combined the difference between the small and large adults is marginally significant ($N = 353$, $X^2_1 = 5.24$, $p = 0.02$). Juveniles have the smallest proportion of high parasitemia, small adults the highest proportion, and large adults are inter-

TABLE 2. PREVALENCE OF MALARIA (THE PERCENTAGE OF LIZARDS EXAMINED THAT HAD MALARIA) AND LEVELS OF PARASITEMIA (THE PERCENTAGE OF INFECTIONS WITH HIGH PARASITEMIA IE. > 9 INFECTIONS PER 1,000 RBC'S) AT DIFFERENT SITES AND IN DIFFERENT LIZARD CLASSES.

	<i>A. limifrons</i> with Malaria		Infections with High Parasitemia	
	N	%	N	%
Lutz				
subadults (SVL<40 mm)	107	9	10	20
small females (39<SVL<44)	54	28	15	47
small males (39<SVL<44)	47	32	15	53
large females (43<SVL)	116	55	64	37
large males (43<SVL)	193	62	119	34
Total	517	43	223	36
AVA				
subadults (SVL<40 mm)	141	5	7	14
small females (39<SVL<44)	76	7	5	40
small males (39<SVL<44)	77	17	11	46
large females (43<SVL)	201	17	34	29
large males (43<SVL)	275	32	88	24
Total	770	19	147	27
Grand Total	1287	29	370	33

N = number examined.

mediate. If one compares the frequency of high parasitemias among all smears (not just those with malaria) juveniles have lower frequencies than do adults (Lutz $N = 517$, $X^2_1 = 19.4$, $p < 0.01$; AVA $N = 770$, $X^2_1 = 5.24$, $p < 0.05$). This suggests either that juveniles contract malaria less frequently than do adults or that the early stages of the infection are more frequently fatal for juveniles. Males and females are very similar in the proportion of cases of malaria in which the level of parasitemia is high. Though, as would be expected, where there is a difference males have more high parasitemias.

Both sexes and all three size classes are essentially identical in the species of malaria infecting them.

THE EFFECTS OF MALARIA ON THE ANOLES

Blood Picture. Malaria, at least when parasitemia levels are high, is associated with an abnormally high percentage of immature RBC's. This usually occurs during the crisis phase of an infection

and immediately following it, apparently due to a large scale destruction and replacement of RBC's by the lizard in response to a high level of parasitemia (Ayala, 1977). In our study the number of these immature RBC's per 100 RBC's was routinely counted in all slides with *Plasmodium*. Figure 2 shows a positive correlation between the level of parasitemia and the proportion of the RBC's that are immature (Lutz $N = 220$, $r = 0.45$, $p < 0.001$, AVA $N = 146$, $r = 0.42$, $p < 0.001$).

Testing simultaneously for the effects of the two species of malaria and three levels of parasitemia, we found (Table 4) that *Plasmodium* species either alone, or in interaction with level of parasitemia, made no difference in the proportion of IRBC's. It is surprising that *P. balli* which visibly distorts the nucleus of an RBC apparently has no more effect than does a *P. "tropiduri"* that looks to be much less harmful to the host cell.

Testing simultaneously the effects of the three levels of parasitemia, site, census, class, and their possible interactions

Figure 2. Percentage of red blood cells which are immature, related to number of plasmodia found per 1,000 red blood cells in *Anolis limifrons* on Barro Colorado Island.

on the proportion of immature RBC's, we found (Table 5) that the level of parasitemia has as significant an effect as do census, class, and the interaction between census and level of parasitemia.

Acute malaria produces an effect in lizards which reduces the hemoglobin levels in blood (Scorza, 1971). This anemia presumably occurs in *A. limifrons* in cases where we find large percentages of IRBC's. The effect of high parasitemias on IRBC's and presumably

on anemia is more severe in smaller anoles than it is in larger ones. Any reduction in hemoglobin must mean a reduction in oxygen carrying capacity of the blood and therefore, if nothing else, of the lizard's ability to repay an oxygen debt quickly. Iguanid lizards are well known for their ability to sustain activity anaerobically (Bennett and Licht, 1972), so anemia might not affect a lizard's ability to catch prey, to flee a predator or to interact socially, but it should reduce its

TABLE 3. SPECIES OF MALARIA.

	P. balli		P. balli & P. "trop- iduri"		P. "trop- iduri"	
	N	%	N	%	N	%
Lutz	83	37	106	48	34	15
AVA	41	27	92	63	14	10

ability to recover quickly from such activity. As the following pages will show, we did not find this disability reflected in such things as growth, survival, or reproduction.

Condition. Lizards with malaria were as heavy as those without malaria. Condition, the weight of a lizard divided by its

snout-vent length cubed, was tested (Table 6) for the effects of malaria species and three levels of parasitemia, size, and sex class of lizard and the interactions of these variables. Class and season and their interaction did produce effects at the two sites but at neither site did malaria show any effect. As we have described in the previous paper, males were generally lighter than females of comparable length. Part, at least, of the greater weight of the females was due to the egg or eggs which they were producing. That females with malaria were no lighter than those without is evidence that they are equally reproductive.

Food intake. Anoles with malaria were not getting measurably less food than

TABLE 4. STATISTICAL ANALYSIS SYSTEM. GENERAL LINEAR MODELS PROCEDURE—TYPE IV SUMS OF SQUARES.

Source	Degrees of Freedom	Sum of Squares	F Value	Pr. F
Dependent Variable: Immature Red Blood Cells				
Species of Plasmodium	2	106.40	1.29	0.28
Level of Parasitemia	1	1215.06	29.54	0.0001
Species * Parasitemia	2	3.74	0.05	0.96
Model	5	2383.00	11.59	0.0001
Error	363	14930.59		

* = interaction between variables.

TABLE 5. STATISTICAL ANALYSIS SYSTEM. GENERAL LINEAR MODELS PROCEDURE—TYPE IV SUMS OF SQUARES.

Source	Degrees of Freedom	Sum of Squares	F Value	Pr. F
Dependent Variable: Immature Red Blood Cells				
Site	1	105.89	3.49	0.06
Census	12	809.47	2.22	0.01
Size & Sex Class	4	779.25	6.42	0.0001
Level of Parasitemia	2	387.45	6.38	0.002
Site * Parasitemia	1	0.50	0.02	0.90
Census * Parasitemia	12	813.71	2.23	0.01
Class * Parasitemia	4	218.52	1.80	0.13
Census * Class * Prsta	50	2194.46	1.45	0.03
Model	86	8791.77	3.37	0.0001
Error	283	8593.17		

* = interaction between variables.

TABLE 6. STATISTICAL ANALYSIS SYSTEM. GENERAL LINEAR MODELS PROCEDURE—TYPE IV SUMS OF SQUARES.

Source	Degrees of Freedom	Sum of Squares	F Value	Pr. F
Dependent Variable: Condition (Weight/SVL ³) at Lutz				
Size & Sex Class	4	0.0005	7.03	0.0001
Season	5	0.002	28.10	0.0001
Malaria [†]	6	0.00008	0.74	0.62
Class * Season	19	0.0007	1.99	0.008
Class * Malaria [†]	19	0.0001	0.36	0.99
Season * Malaria [†]	26	0.0003	0.75	0.81
Clss * Ssn * Malaria [†]	21	0.0003	0.90	0.59
Model	100	0.0097	5.48	0.0001
Error	412	0.007		
Dependent Variable: Condition (Weight/SVL ³) at AVA				
Size & Sex class	4	0.0004	6.03	0.0001
Season	5	0.001	15.67	0.0001
Malaria [†]	6	0.00002	0.21	0.97
Class * Malaria [†]	18	0.0007	2.31	0.002
Class * Malaria [†]	13	0.0002	0.86	0.59
Season * Malaria [†]	17	0.0001	0.48	0.96
Clss * Ssn * Malaria [†]	16	0.0003	1.00	0.46
Model	79	0.014	10.39	
Error	686	0.012		

[†] Seven categories used: no malaria and low (< 10/1,000 RBC's) and high (> 9/1,000 RBC's) levels of parasitemia for *Plasmodium balli*, *P. tropiduri*, and both species of *Plasmodium*.

* = interaction between variables.

those without. Food intake was estimated by the weight of scat per gram of lizard produced by a lizard over the 24 hours following capture. A log transformation of the data makes the samples more nearly normally distributed. A test (Table 7) on the transformed data shows that only lizard class had a significant effect on food intake. At Lutz but not at AVA, juveniles and adult females produced a relatively greater weight of feces than did adult males.

Time of activity. Malaria did not seem to affect the time of day at which the lizards were active. Each census was divided into three periods: before 9 am, between 9 and 10 am, and after 10 am. Contingency tables were constructed to compare the times at which lizards were caught. At neither site and in neither the wet nor the dry season did the presence of malaria (Wet season: Lutz N = 293, X₂²

= 0.33, p = 0.85, AVA N = 381, X₁² = 0.05, p = 0.98; Dry season: Lutz N = 221, X₂² = 0.74, p = 0.69, AVA N = 378, X₂² = 282, p = 0.24) or the level of parasitemia (Wet season: Lutz N = 293, X₄² = 7.85, p = 0.097, AVA N = 381, X₄² = 1.90, p = 0.75; Dry season: Lutz N = 221, X₄² = 3.72, p = 0.45, AVA N = 378, X₄² = 3.84, p = 0.43) show a significant effect. Lizards with malaria were not compensating for it by being active earlier nor were they debilitated by it and becoming active later.

Reproduction. Reproduction did not seem to be depressed in those females showing malaria. The proportion of females which tested positive for oogenesis varied with size (Table 8). Those tested below a SVL of 35 mm were never positive; those few above 48 mm always were; at intermediate sizes the proportion that were oogenic varied, increasing

TABLE 7. STATISTICAL ANALYSIS SYSTEM. GENERAL LINEAR MODELS PROCEDURE—TYPE IV SUMS OF SQUARES.

Source	Degrees of Freedom	Sum of Squares	F Value	Pr. F.
Dependent Variable: Food Intake (Log of Scat Weight) at Lutz				
Size & Sex Class	4	8.19	6.21	0.0001
Season	5	2.78	1.69	0.14
Malaria [†]	6	1.26	0.64	0.70
Class * Season	19	8.28	1.32	0.17
Class * Malaria [†]	19	3.61	0.61	0.89
Season * Malaria [†]	25	7.84	0.95	0.53
Clss * Ssn * Malaria [†]	13	5.36	1.25	0.24
Model	91	41.40	1.38	0.02
Error	279	91.98		
Dependent Variable: Food Intake (Log of Scat Weight) at AVA				
Size & Sex Class	4	3.26	1.96	0.10
Season	4	2.04	1.22	0.30
Malaria [†]	6	0.73	0.29	0.94
Class * Malaria [†]	14	5.70	0.98	0.47
Class * Malaria [†]	11	4.78	1.04	0.41
Season * Malaria [†]	12	3.76	0.75	0.70
Clss * Ssn * Malaria [†]	7	3.69	1.27	0.26
Model	58	181.79	1.68	0.0023
Error	437	222.27		

[†]Seven categories used: no malaria and low (< 10/1,000 RBC's) and high (> 9/1,000 RBC's) levels of parasitemia for *Plasmodium balli*, *P. tropiduri*, and both species of *Plasmodium*.

* = interaction between variables.

as SVL increased. Once an *Anolis limifrons* begins to yolk eggs she usually continues to do so. Only 13% of the 93 females tested again after once testing positive tested negative on the second occasion.

A two by two contingency table shows a significant effect of the presence of malaria on the frequency of oogenesis in potentially reproductive (36 mm SVL) females ($N = 504$, $X^2_1 = 25.29$, $p = 0.0001$). Surprisingly the effect is positive, a higher proportion of the females with malaria are oogenic than of those without malaria.

A tabulation (Table 4) of oogenic females with and without malaria shows no negative effect of malaria on oogenesis and suggests that the apparent positive effect is possibly a spurious side effect of the fact that both the prevalence of ma-

laria and the frequency of oogenesis increase with size. A similar lack of effect of malaria on oogenesis was seen in *Anolis auratus* infected with *Plasmodium colombiense* in Colombia (Ayala and Spain, 1976).

Looking at the effect of malaria in another way, we can consider only those adult females that had a very high (> 19 per 1,000 RBC's) parasitemia or a high (> 9 per 100) percentage of IRBC's. Of these 24 anoles, all but one tested positive for oogenesis. Therefore even the acute phase of malaria does not inhibit oogenesis to a measurable extent. Possibly the changes in the blood associated with egg production favors asexual reproduction in the *Plasmodium* so that parasites become more common in oogenic anoles.

Our evidence of reproductive activity

TABLE 8. THE EFFECTS OF MALARIA ON OOGENESIS.

		Female <i>Anolis limifrons</i>			
		Without Malaria		With Malaria	
Snout Vent Length (mm)		N	% Oogenic	N	% Oogenic
21		1	0		
22		1	0		
23					
24					
25	1		0		
26					
27	1		0		
28					
29	1		0		
30	2		0		
31	2		0		
32	6		0	1	0
33	5		0	1	0
34	11		0		
35	5		20.0		
36	9		0		
37	12		16.7	1	0
38	10		20.0	4	50.0
39	10		50.0	1	0
40	22		45.5	2	50.0
41	20		30.0	6	66.7
42	28		64.3	2	100.0
43	28		60.7	7	100.0
44	28		89.3	8	100.0
45	49		89.8	20	95.0
46	53		90.6	24	87.5
47	38		84.2	18	94.4
48	14		85.7	11	100.0
49	7		100.0	3	100.0
50	5		100.0	2	100.0
51	1		100.0	1	100.0
52				2	100.0

eggs per female per week; Mean weight = 0.14 g, N = 48) (Andrews, unpublished data).

Growth. Anoles with malaria seem to grow as rapidly as those without. A test (Table 9) of the effects, on growth rate, of size and the presence of malaria (at the beginning, or end, or both of the growth interval) showed a strong effect of size ($p = 0.0001$) but none of malaria, alone or in interaction with size. The lower growth rates in malarial animals, which one sees when one groups animals without regard for size, seems to be due to a higher prevalence of malaria among larger animals (as noted above) and that larger animals grow more slowly than do smaller ones (Andrews *et al.*, this volume).

Survivorship. Malaria did not appear to decrease the survivorship of infected anoles. To determine if malaria affected the likelihood that a lizard would disappear from the population, contingency tables were prepared for each site comparing the numbers of lizards caught in each census with and without malaria, with our records as to whether they had been recaptured during the same census, recaptured at some subsequent census or never captured again and had probably died or emigrated. At neither site was there a significant difference in the likelihood of recapture during the same or during a subsequent census due to having malaria (Lutz N = 517, $X^2_2 = 1.80$, $p = 0.41$, AVA N = 770, $X^2_2 = 1.72$, $p = 0.42$), or to level of parasitemia (Lutz N = 517, $X^2_2 = 3.12$, $p = 0.54$, AVA N = 770, $X^2_4 = 1.96$, $p = 0.74$). Malaria did not seem to affect either conspicuousness or survivorship.

Predator pressure. Anoles with malaria seemed to be no more at risk from predators than those without malaria. To estimate intensity of predation, frequencies of regenerated tails were tabulated for most captured anoles. Only the condition of the tail on the last occasion when a lizard was captured was scored to avoid counting the same break twice. Combining sites, size, and sex classes shows a

in these anoles in the field is whether their blood smears stained reddish indicating that they were circulating lipoproteins and therefore presumably yolking eggs. One can be reasonably sure that females that are not yolking eggs are not actively reproductive. However, there is no way of knowing from our field data if two oogenic females are producing eggs at the same rate. There is some evidence from females maintained in captivity. Seven captive females with malaria laid as many eggs (Mean = 0.79 per female per week) of the same size (Mean weight = 0.14 g, N = 27) as did 12 similar sized females without malaria (Mean = 0.81

TABLE 9. STATISTICAL ANALYSIS SYSTEM. GENERAL LINEAR MODELS PROCEDURE—TYPE IV SUMS OF SQUARES.

Source	DF	Sum of Squares	F Value	Pr. F
Dependent Variable: Growth Rate				
Lutz, Wet Season				
Snout Vent Length	1	0.87	72.41	0.0001
Presence of Malaria	1	0.001	0.89	0.35
SVL * Malaria	1	0.001	1.01	0.32
Model	3	0.12	32.27	0.0001
Error	34	0.04		
Lutz, Dry Season				
Snout Vent Length	1	0.31	20.95	0.0001
Presence of Malaria	1	0.00002	0.01	0.92
SVL * Malaria	1	0.00001	0.01	0.93
Model	3	0.04	8.71	0.0004
Error	24	0.04		
AVA, Wet Season				
Snout Vent Length	1	0.26	10.98	0.0017
Presence of Malaria	1	0.0000004	0.00	0.99
SVL * Malaria	1	0.0000002	0.00	0.99
Model	3	0.04	5.53	0.002
Error	49	0.11		
AVA, Dry Season				
Snout Vent Length	1	0.004	1.71	0.20
Presence of Malaria	1	0.0005	0.22	0.64
SVL * Malaria	1	0.0005	0.24	0.63
Model	3	0.06	9.06	0.0001
Error	63	0.14		

DF = Degrees of Freedom

* = interaction between variables

significant effect of malaria on frequency of tail breaks (overall % broken = 19.2, $N = 640$, $X^2_1 = 8.35$, $p = 0.004$). This is probably spurious because, as one might expect, at both sites, larger animals are more likely to have their tails broken than are smaller ones (Lutz $N = 272$, $X^2_2 = 8.67$, $p = 0.013$; AVA $N = 368$, $X^2_2 = 10.56$, $p = 0.005$). Larger lizards should more frequently have broken tails if only because they are generally older than small ones and the scar left by a broken tail never disappears.

For all lizard classes where the sample size is large enough to allow testing (large males at Lutz (% broken = 26.3, N

= 99, $X^2_1 = 1.16$, $p = 0.28$), and large males (% broken = 22.0, $N = 132$, $X^2_1 = 0.002$, $p = 0.97$) and females at AVA (% broken = 31.0, $N = 100$, $X^2_1 = 0.06$, $p = 0.81$)), the malarial state when last captured had no significant effect on the frequency of tail breakage.

Among large adults, males have their tails broken more often than do females but the difference is small and not significant (Lutz $N = 159$, $X^2_1 = 2.76$, $p = 0.097$; AVA $N = 232$, $X^2_1 = 2.42$, $p = 0.12$). Because males fight more than do females, if aggressive encounters between conspecifics were an important cause of tail breakage one would expect a

substantially higher percentage of tail breaks among males than among females. We do not find this and so aggressive encounters are probably not important. Most tail breaks probably come from failed predation attempts. It is certainly not necessary that there be a close correlation between failed and successful attempts, but it seems likely that this would be true except where the sample includes differing proportions of naive predators or of predators with different capture techniques. In our situation there could possibly be a seasonal difference in predators but there is unlikely to be a site difference and we can think of no way in which either of these would bias the picture either for or against anoles with malaria.

Differential mortality. Malaria did not have a detectable differential effect on the mortality of the different size and sex classes of anoles. One possible effect of malaria which we have not yet considered is that it might kill a significant number of lizards during the early stages of the infection, leaving the others to recover essentially unaffected even though still circulating the parasite. Two of the several score of naturally infected anoles which we kept in captivity died during the crisis stage of their malaria infection. The malaria-induced anemia may well have been at least a contributing factor.

We can not test for a general effect of this sort but we might detect an effect if the malaria was affecting different lizard classes differently. One might expect that if malaria was sometimes fatal it might kill a higher percentage of young anoles than older ones, or kill more of the egg producing females than males. If this is so one would expect that the classes most strongly affected should be proportionately rarer at Lutz than at AVA where malaria is much less common. A comparison of size and sex classes at the two sites shows no significant difference between them during either dry ($N = 601$, X^2_4 , $p = 0.53$) or wet ($N = 686$, X^2_4 , $p = 0.40$) seasons.

There appears to be no statistically significant difference of the effect of malaria on the two sexes, or on the size classes. However, this approach would not detect an effect in which malaria was killing proportionally more animals in the smallest size class. This early mortality would affect all succeeding size classes as well and so appear as a general lowering of the population level without changing the size structure, unless it was compensated for by reduced mortality at later stages. Since we do not find a difference between the two sites except that the site with more malaria has fewer lizards, we can not exclude the possibility that malaria has a significant negative effect on the survival of an infected anole. We can say that if malaria does have such an effect, it occurs during the early stages of the infection, and that if the effect is different on different age classes it must be more severe on young animals.

We found lower frequencies of high levels of parasitemia (new infections) among juveniles than among adults. This is consistent with the hypothesis that malaria is more serious for younger than for older animals but it is also consistent with the hypothesis that small lizards are less attractive to the vectors and so are infected less frequently.

SUMMARY AND CONCLUSIONS

Malaria was extremely common among *Anolis limifrons* during our study. In one population 95% of the large adults captured three or more times had a patent malaria infection at least once. Two morphologically different types of *Plasmodium* were involved which were similar in their distribution in time, space and class of *A. limifrons* and both were frequently present together in the same individual.

The disease was present at all times of year, and there did not seem to be marked seasonality in either overall

prevalence or frequency of high levels of parasitemia indicative of new infections, though the latter were most common in early rainy season.

Malaria was much more common among large individuals than among small ones. Individuals seemed, once infected, to continue to circulate the parasite for a long time, perhaps throughout their lives, usually at a low level, though relapses with high levels of parasitemia were observed in captive animals. The higher prevalence in large animals is in part the result of this persistence of the parasite. If the probability of being infected remained above zero throughout life then the probability of ever being infected would increase with age and older (larger) animals would have, as we find that they do, a higher prevalence of infection.

Since the frequency of high levels of parasitemia was higher among small adults than large ones, the probability of contracting the infection is as high, or higher, for a small adult as it is for a larger adult. The prevalence of malaria and the occurrence of high levels of parasitemia was low in the juveniles. Either the probability of contracting malaria is low for these small animals, or malaria when contracted is more likely to be fatal to a small animal than it is to a large animal.

Chronic malaria with low levels of parasitemia produced little effect on the blood picture. Higher levels of parasitemia correlated with high levels of immature RBC's and this in turn presumably reflected a high level of anemia. This effect was stronger for small animals than for larger ones.

We examined a variety of other aspects of the anoles from the two study sites including: weight, food intake, time of activity, female reproductive condition, growth, survivorship, and tail breakage. Despite large sample sizes we found no factor, except IRBC's, on which malaria had a statistically significant effect on the lizards in which we detected it, whether we look at all malaria infections or only

the relatively small number of cases with high parasitemias.

Apparently malaria does not have a marked effect on the anoles that it infects, at the very least this is true for adults with chronic malaria. This is what one would expect of a well adapted parasite in a closely coevolved host-parasite relationship. It is to the parasite's advantage, as well as to the advantage of the host, for the host to live for a long time and leave many offspring. However, we can not conclude, with certainty, that malaria has no ecologically significant effect on these populations, because we can not exclude the possibility that the initial stages of a malaria infection, which are rarely represented in our sample, may be fatal, at least to young animals. Still, at least in its chronic phase, malaria in *Anolis limifrons* seems to be behaving as a prudent parasite.

ACKNOWLEDGMENTS

We thank Steve Ayala for his advice and counsel on lizard malaria during the planning stages as well as in the course of the project. A large number of people helped in the field, among them: John Pickering, Bonifacio deLeon, Gordon Burghardt, Lauri McHargue, Sharon Strauss, Hal Hertzog, Marina Guerrero; others assisted in the lab: Humberto and Fanny Carvajal and Luis Carlos Zamora; Don Windsor at STRI and Ned Fetcher, Biology Department and W. R. Price, the Statistical Consulting Service of VPI&SU advised on statistical analysis.

We are indebted to STRI for logistic and financial support, to National Science Foundation Binational Grant No. INT 7605758 to Rand and Ayala, National Institute of Health Grant 2 RO1 AI12511 to Ayala, and a Smithsonian Institution Research award to Rand for financial support.

We thank John Pickering, Ernest E. Williams, Joe Wright, and Bob Kimsey for stimulating discussions and valuable

suggestions on the problems of lizard malaria and anole ecology, and those who read and commented on the manuscript: John Pickering, Joe Wright, Pat Rand, Bob Kimsey, Mike Ryan, Steve Ayala, and Joe Schall.

LITERATURE CITED

- ANDERSON, R. C. 1976. Helminths, pp. 35-43. In L. A. Page (ed.), *Wildlife Diseases. Section II. Impact of parasitic diseases on wildlife populations*. New York, Plenum Press.
- ANDREWS, R. A., AND A. S. RAND. 1982. Long-term changes in population density of the lizard *Anolis limifrons*. In E. G. Leigh, Jr., A. S. Rand, and D. M. Windsor (eds.), *The Ecology of a Tropical Forest: Seasonal Rhythms and Long-term Changes*. Washington, D.C., Smithsonian Institution Press. (in press).
- ARAGAO, H. DE B., AND NEIVA, A. 1909. A contribution to the study of the intraglobular parasites of the lizards. Two new species of *Plasmodium*, *Pl. diploglossi* n. sp. and *Pl. tropiduri* n. sp. Mem. Inst. Oswaldo Cruz, 1: 44-50.
- AYALA, S. C. 1971. Sporogony and experimental transmission of *Plasmodium mexicanum*. J. Parasitol., 57: 598-602.
- . 1977. Plasmodia of reptiles, pp. 267-309. In J. P. Kreier (ed.), *Parasitic Protozoa. Vol. III. Gregarines, Haemogregarines, Coccidia, Plasmodia and Haemoproteids*. New York, Academic Press.
- . 1978. Checklist, host index, and annotated bibliography of *Plasmodium* from reptiles. J. Protozool., 25: 87-100.
- AYALA, S. C., AND J. L. SPAIN. 1975. Annual oogenesis in the lizard *Anolis auratus* determined by a blood smear technique. Copeia, 1975: 138-141.
- , AND ———. 1976. A population of *Plasmodium colombiense* sp. n. in the iguanid lizard, *Anolis auratus*. J. Parasitology., 62: 177-189.
- BENNETT, A., AND P. LICHT. 1972. Anaerobic metabolism during activity in lizards. J. Comp. Physiol., 81: 277-288.
- BENNETT, G. F., E. C. GREINER, AND W. THRELFALL. 1976. Protozoans, pp. 25-33. In L. A. Page (ed.), *Wildlife Diseases. Section II. Impact of parasitic diseases on wildlife populations*. New York, Plenum Press.
- HELVIG, J. T., AND K. A. COUNCIL. 1979. SAS User's Guide, 1979 Edition. SAS Institute Inc., Raleigh.
- JORDAN, H. B. 1964. Lizard malaria in Georgia. J. Protozool., 11: 566-572.
- GUERRERO, S., C. RODRIGUEZ, AND S. C. AYALA. 1977. Prevalencia de hemoparasitos en lagartijas de la Isla Barro Colorado, Panama. Biotropica, 9: 118-123.
- MARKUS, M. B. 1974. Arthropod-borne disease as a possible factor limiting the distribution of birds. Intern. J. Parasit., 4: 609-612.
- PICKERING, J. 1980. Sex ratio, social behavior and ecology in *Polistes* (Hymenoptera, Vespidae), *Paschysomoides* (Hymenoptera, Ichneumonidae) and *Plasmodium* (Protozoa, Haemosporida). Ph.D. Thesis. Harvard University, Cambridge, Mass.
- SEXTON, O. J. 1967. Population changes in a tropical lizard *Anolis limifrons* on Barro Colorado Island, Panama Canal Zone. Copeia, 1967: 219-222.
- SEXTON, O. J., J. BAUMAN, AND E. ORTLEB. 1972. Seasonal food habits of *Anolis limifrons*. Ecology, 53: 182-186.
- SEXTON, O. J., J. HEATWOLE, AND E. MESETH. 1963. Seasonal population changes in the lizard, *Anolis limifrons*, in Panama. Am. Midland Natur., 69: 482-491.
- SCORZA, J. W. 1971. Anaemia in lizard malaria infections. Parasitologia, 13: 391-405.
- TELFORD, S. 1969. A new saurian malarial parasite *Plasmodium balli* from Panama. J. Protozool., 16: 437-441.
- . 1974. The malarial parasites of *Anolis* species (Sauria: Iguanidae) in Panama. Intern. J. Parasitol., 4: 91-102.
- . 1977. The distribution, incidence and general ecology of saurian malaria in Middle America. Intern. J. Parasitol., 7: 299-314.
- . 1979. A taxonomic reconsideration of some *Plasmodium* species from iguanid lizards. Ann. Parasitol. (Paris), 54: 129-144.
- WARNER, R. E. 1968. The role of introduced diseases in the extinction of the endemic Hawaiian avifauna. Condor, 70: 101-120.
- WRIGHT, S. J. 1979. Competition between insectivorous lizards and birds in central Panama. Amer. Zool., 19: 1145-1156.

Eurythermy and Niche Breadth in West Indian *Anolis* Lizards: A Reappraisal

PAUL E. HERTZ¹

ABSTRACT. A comparison of the thermal biologies of the solitary species *Anolis roquet* on Martinique and three species from the complex fauna of Puerto Rico, *A. cristatellus*, *A. evermanni*, and *A. gundlachi*, demonstrates no consistent differences in thermal niche breadth that can be attributed to *A. roquet*'s occupation of an island in the absence of congeneric competitors. Three behavioral-ecological "phenotypes" with respect to thermal biology are described: 1) thermal passivity; 2) obligate thermoregulation; and 3) thermoregulatory versatility. Anoles from the complex faunas exhibit any of the three phenotypes, but solitary anoles all appear to be versatile, at least on thermally complex islands. Eurythermy and thermoregulatory versatility probably represent an ancestral "colonizing phenotype" in West Indian anoles. The thermal biology of solitary anoles may represent a retention of this ancestral phenotype rather than a product of ecological release.

INTRODUCTION

The concept of "ecological release" was developed by ecologists to explain the apparent increase in the range of resources utilized by island populations relative to that used by conspecifics on the mainland (Wilson, 1961). In theory, "solitary" populations are relatively free to exploit a wider variety of resources because of the absence (or near absence) of closely related competitors in depauperate island faunas.

During the last decade, researchers studying the ecology of *Anolis* lizards in the West Indies have extrapolated from the original formulation of the concept to compare patterns of resource utilization among the island-dwelling members of the genus. These researchers have concluded that solitary anoles generally exhibit expanded structural habitat (i.e., perch site) and/or thermal niches relative to those of species from the complex faunas of the Greater Antilles (e.g., Lister, 1976; Rand and Rand, 1967; Ruibal and Philibosian, 1970). Proof of ecological release in the strictest sense (Grant, 1967) is usually lacking, however, in the absence of ecological data on the ancestors of the solitary populations (Huey and Webster, 1976; Ruibal and Philibosian, 1970).

The presumed expansions of structural habitat and thermal niches of the solitary species are thought to be reflections of a single phenomenon: solitary species occupy a greater variety of habitats and microhabitats. In the complex *Anolis* faunas of the Greater Antilles, interspecific resource partitioning of structural and climatic habitats suggests that competitors restrict the range of thermal environments that some species may occupy (Rand, 1964; Schoener and Schoener, 1971a; Rand and Williams, 1969). Indeed, Ruibal and Philibosian (1970) explicitly stated that the relatively broad thermal niche of the solitary spe-

¹Department of Biological Sciences, Barnard College, Columbia University, New York, New York 10027, U.S.A.

cies *A. oculatus* on Dominica resulted from its occupation of "an array of habitats which on Cuba or Puerto Rico... are occupied by different species." One could logically argue that the occupation of a large array of microhabitats would also result in an expansion of the thermal niche.

The broad thermal niche of a species could be produced by either of two phenomena (Huey and Webster, 1975). On the one hand, each population of the species might be eurythermal (i.e., having a broad thermal niche), exhibiting a niche breadth nearly as great as that shown by all populations of the species considered together. On the other hand, a broad species thermal niche could be produced by population differentiation among the niche breadths of several stenothermal (i.e., having a narrow thermal niche) populations living under, but not behaviorally compensating for, distinct climatic regimes (Hertz, 1977). The distinction of "within-population" and "among-population" components of a broad species niche is analogous to Roughgarden's (1972) "within-phenotype" and "between-phenotype" components of a population's niche breadth.

Although previous researchers have considered population stenothermy and eurythermy in their comparisons of thermal niche breadths among *Anolis* species (Huey and Webster, 1975, 1976; Ruibal and Philibosian, 1970), none has made and tested explicit predictions about both within- and among-population niche breadths in solitary versus sympatric species (but see Lister, 1976). A systematic knowledge of these phenomena is essential to our understanding of the evolution and biogeography of the huge West Indian radiation of this genus (Williams, 1969, 1972). The purpose of this paper is to make and test two such predictions, one at the population level and another at the species level.

1) If it is true that populations of solitary anoles are more eurythermal and

occupy broader arrays of microhabitats than do populations in the complex faunas, then we should observe that the within-population niche breadths of the solitary species are at least equal to the combined population niche breadths of two or more sympatric species in the complex fauna.

2) If it is true that the solitary species occupy broader arrays of habitats than do species in the complex fauna, then we should observe that the among-population niche breadth (hereafter, the species niche breadth) of the solitary species is at least equal to the combined species niche breadths (among populations and among species) of those species that replace each other over an equivalent range of habitats in the complex fauna.

To test these predictions, I systematically compared the structural habitats and field thermal biology of a solitary *Anolis* of the Lesser Antilles with those of other *Anolis* species that are, at least superficially, its ecological equivalents in the complex fauna of Puerto Rico. On Martinique, *roquet* is the dominant terrestrial vertebrate in many habitats from sea level to approximately 750 m elevation (Lazell, 1972). On Puerto Rico, *crisatellus* is the dominant trunk-ground anole of the lowlands, and *gundlachi* its upland replacement in montane forests up to approximately 1,000 m elevation. Two other species, *stratulus* and *evermanni*, also commonly occupy tree trunks in the Puerto Rican lowlands and uplands, respectively (Rand, 1964); however, *stratulus* was rare at the lowland *crisatellus* study site, and I did not sample it there.

I sampled along altitudinal transects on both islands. Although such a sampling regime does not ensure the representation of every habitat type present on an island, altitude does provide predictable gradients in ambient temperature, cloud cover, moisture, vegetation, and, on Puerto Rico, community composition.

If the predictions made above are correct, I would expect the following rela-

tionships to emerge from the data on structural habitat and thermal niche breadths. 1) The niche breadths of individual *roquet* populations should be greater than those of individual populations of *crisatellus*, *gundlachi*, and *evermanni* and at least equal to the combined population niche breadths of sympatric *evermanni* and *gundlachi* populations. 2) The overall species niche breadth of *roquet* (among populations sampled along the transect) should be at least equal to the combined species niche breadths of *crisatellus* and *gundlachi* sampled over an equivalent altitudinal range.

MATERIALS AND METHODS

FIELD STUDIES

I sampled three populations of *roquet* (at 20 m, 350 m, and 650 m elevations) in northeastern Martinique during late June and early July 1975. In Puerto Rico, I studied a lowland population of *crisatellus* on the western outskirts of Luquillo (5 m) and four populations (at 270 m, 480 m, 675 m, and 850 m) of its upland replacement, *gundlachi*, in the Luquillo Experimental Forest during July, August, and September 1975. In addition, I simultaneously sampled *evermanni*, sympatric to *gundlachi*, at the two intermediate upland sites. The locations of the study sites, brief descriptions of the vegetation at each, and the subspecific identifications of the *roquet* populations are provided elsewhere (Hertz, 1977, 1981).

I obtained data records from adults and subadults captured with standard techniques and precautions (Hertz, 1977) from dawn to dusk at each locality. For each individual captured, I recorded the following data: sex, snout-to-vent length, body (cloacal) and air (shaded bulb, 1 cm above perch site) temperatures with a quick reading Schultheis thermometer, perch height, and time of capture.

Samples varied from 132 to 180 records per population and were uniformly distributed throughout the animals' full periods of activity at each site. Because there were neither size- nor sex-dependent trends in any population, I have lumped all records within each population in this analysis. A treatment of other aspects of the field data can be found elsewhere (Hertz, 1977, 1981).

ESTIMATION OF NICHE BREADTHS

The estimation of niche breadth for the structural habitat of each population is straightforward. I follow Lister (1976) in using perch height diversity as an index of niche breadth along this resource dimension, using the formula $B = 1/\sum p_i^2$ (Levins, 1968), in which p_i is the frequency with which lizards perched in each one-foot height interval.

Calculation of thermal niche breadth is much more complex, largely because environmental heterogeneity can strongly affect the variability of body temperatures in a population (Soulé, 1963; see also Huey and Slatkin, 1976; Huey, 1982). Hence, one must consider both environmental and body temperatures to accurately describe thermal specialization. For the data presented here, I follow Ruibal and Philibosian (1970) and Huey and Webster (1975, 1976) in using the range of hourly mean body temperatures to estimate thermal niche breadth. Similarly, the range of hourly mean air temperatures indexes the environmental heterogeneity (hereafter, the thermal habitat breadth) experienced by each population. However, the range of hourly mean air temperatures is an imperfect measure of environmental heterogeneity because air temperatures only approximate operative environmental temperatures (Bakken, 1976). Furthermore, the data underestimate the available environmental heterogeneity because these air temperatures are not independent of the animals under study (i.e., I did not measure air temperature in microen-

vvironments that the lizards did not use). Nevertheless, these statistics have the advantage of minimizing the importance of extreme temperatures in any data set. They also provide convenient summaries which are directly comparable among studies in which lizards were sampled over a range of time periods.

RESULTS

POPULATION NICHE BREADTHS: TESTS OF PREDICTION 1

Perch height diversities for populations of *roquet* (maximum = 6.62) on Martinique (Fig. 1) were similar to those for each of the three species on Puerto Rico (Fig. 2). These data indicate that *roquet*, in the absence of congeneric competitors, does not use a wider variety of perch heights than does any of the common species in the complex fauna. Indeed, perch height diversities for two combined samples of sympatric *gundlachi* and *evermanni* were higher (8.50 at 480 m and 7.12 at 675 m) than those for

roquet. These data fail to confirm the first prediction. (The data on perch heights may be biased insofar as *evermanni* perched very high in a tree are less likely to be seen than are those perched lower on the trunk (Rand, 1964). However, such a bias is conservative because it would have fostered an underestimate of *evermanni*'s niche breadth.) Schoener (1975) has shown that the structural habitat breadth of some *Anolis* populations is dependent upon the availability of perch sites in various habitats. Although I did not quantitatively measure perch height availabilities, I do not believe that inter-locality differences in the height of the vegetation influenced these data. Lizards in all populations failed to use the full range of perch heights that were available.

Estimates of the heterogeneity in the thermal environment (thermal habitat breadths, Table 1) show that *roquet* populations experienced more variable environments than did most populations of *gundlachi* and *evermanni*, but the difference is neither consistent nor of great magnitude. The lowland sample of *cristatellus* inhabited the most heterogeneous thermal environment of all the populations surveyed. Moreover, the combined thermal habitat breadths of sympatric *gundlachi* and *evermanni* (3.4 C at 480 m and 2.6 C at 675 m) were indistinguishable from those of the *roquet* populations. Some of these data confirm the first prediction.

Thermal niche breadths (Table 1) of *roquet* populations were consistently greater than those of *gundlachi* and *evermanni* populations, but the niche breadth of the *cristatellus* population was nearly as large as that of the most eurythermal *roquet* population. Thermal niche breadths of each of the *roquet* populations was also greater than the combined niche breadth of sympatric *gundlachi* and *evermanni* populations (3.4° C at each site). These data also provide a partial confirmation of the first prediction.

Figure 1. Sex-weighted perch height frequency distributions for *Anolis roquet* on Martinique. Mean perch heights (cm, $\bar{X} \pm \text{s.e.}$) and perch height diversity ($B = 1/\sum p_i^2$) indicated for each population.

Figure 2. Sex-weighted perch height frequency distributions for three *Anolis* species on Puerto Rico. Mean perch heights and perch height diversities as in Figure 1.

SPECIES NICHE BREADTHS: TESTS OF PREDICTION 2

The environmental heterogeneity experienced by all *roquet* populations combined (6.0°C) was not notably larger than that experienced by the four *gundlachi* populations (5.9°C), but the former was larger than that evident among the two *evermanni* populations (2.6°C). (These data may, however, underestimate species breadths for *evermanni* because this species occupies a larger elevational range than is evident in my samples [Rand, 1964; Schoener and Schoener, 1971b].) My data on one *crist-*

atellus population are insufficient as an estimator for the entire species, but Huey and Webster (1976), using identical techniques in summer, found a species thermal habitat breadth of 8.8°C for 27 hourly samples of *cristatellus* from diverse localities. The species thermal habitat breadth for *roquet* (above) was slightly less than that of the combined thermal habitat breadth of *cristatellus* and *gundlachi* sampled over an equivalent range of elevations (6.7°C , when the 850 m *gundlachi* sample is excluded). The above data, in combination with Huey and Webster's (1976) observation of a species thermal habitat breadth of

TABLE 1. MINIMUM AND MAXIMUM HOURLY MEAN TEMPERATURE STATISTICS, THERMAL HABITAT BREADTHS (THB), AND THERMAL NICHE BREADTHS (TNB) (ALL °C) IN 10 SAMPLES OF WEST INDIAN ANOLIS LIZARDS.

		N ⁺	Air Temperatures			Body Temperatures		
			minimum X̄ ± s.e.	maximum X̄ ± s.e.	THB	minimum X̄ ± s.e.	maximum X̄ ± s.e.	TNB
roquet:	20 m	12	25.1 ± 0.3	*27.7 ± 0.4	2.6	26.4 ± 0.2	30.0 ± 0.2	3.6
	350 m	12	23.7 ± 0.1	27.0 ± 0.3	3.3	24.7 ± 0.2	*31.1 ± 0.2	6.4
	650 m	10	*21.7 ± 0.3	24.8 ± 0.4	3.1	*22.2 ± 0.1	26.7 ± 0.2	4.5
cristatellus:	5 m	12	24.2 ± 0.1	28.3 ± 0.2	4.1	24.3 ± 0.1	30.4 ± 0.2	6.1
gundlachi:	270 m	12	23.7 ± 0.1	*26.5 ± 0.1	2.8	24.0 ± 0.1	*27.1 ± 0.1	3.1
	480 m	12	22.1 ± 0.1	24.3 ± 0.1	2.2	22.3 ± 0.1	25.1 ± 0.1	2.8
	675 m	12	21.6 ± 0.1	23.9 ± 0.2	2.3	21.7 ± 0.1	24.5 ± 0.2	2.8
	850 m	11	*20.6 ± 0.1	22.2 ± 0.1	1.6	*20.8 ± 0.1	22.6 ± 0.2	1.8
evermanni:	480 m	11	22.7 ± 0.1	*24.5 ± 0.2	1.8	23.2 ± 0.2	*25.7 ± 0.2	2.5
	675 m	10	*21.9 ± 0.1	24.2 ± 0.2	2.3	*22.2 ± 0.1	25.1 ± 0.2	2.9

⁺N = number of hourly samples collected at each locality.

*These values used for calculations of species breadths; see text for explanation.

9.5° C for 19 hourly samples of *gundlachi*, collectively refute the second prediction.

The species thermal niche breadth among *roquet* populations (8.9° C) was substantially greater than that among *gundlachi* and *evermanni* populations (6.3° C and 3.5° C, respectively). In addition, the species thermal niche breadth for *roquet* was remarkably similar to that of the combined species thermal niche breadths of *cristatellus* and *gundlachi* sampled over an equivalent altitudinal range (8.7° C, when the 850 m *gundlachi* sample is excluded). These data clearly support the second prediction. However, Huey and Webster (1976) provide contradictory evidence. They found species thermal niche breadths of 9.6° C and 8.7° C for *cristatellus* and *gundlachi*, respectively; these values more nearly approximate those for *roquet* and therefore refute the second prediction.

DISCUSSION

EURYTHERMY AND SPECIES NICHE BREADTHS

These new data and those of Huey and Webster (1976) only partially confirm the

two predictions made in the Introduction. Populations of *roquet* do not exhibit broader structural habitat or thermal habitat niches than do species in the complex Puerto Rican fauna. However, populations of the former species do exhibit broader thermal niches than do some populations in the complex fauna. The comparisons of species breadths indicate no consistent differences between this solitary species and those from the complex fauna of Puerto Rico.

How general are these results? In the two decades since Ruibal's (1961) pioneering study of the Cuban anoles, researchers have studied the thermal biology of numerous West Indian *Anolis* species. Unfortunately, many of the data are not useful for comparison because they were collected over unspecified time periods and are not divisible into temporally restricted samples (e.g., Brooks, 1968, Heatwole *et al.*, 1969; Rand, 1967). However, I have been able to compile data on the thermal habitat and thermal niche breadths for 56 populations of 15 species (Table 2). These data may have limited generality because no researchers have examined seasonal variation in the thermal biology of West Indian anoles; however, the data set is

TABLE 2. SPECIES THERMAL HABITAT AND THERMAL NICHE BREADTHS (THB AND TNB, °C) FOR 15 SPECIES OF WEST INDIAN ANOLIS LIZARDS.

Species	Island	P [§]	S [†]	THB	TNB	Reference
Solitary Anoles:						
<i>acutus</i> ⁺	St. Croix	1	14	6.9	7.0	McManus and Nellis, 1973
<i>marmoratus</i>	Guadeloupe	9	29	9.0	9.9	Huey and Webster, 1975
<i>oculatus</i> [*]	Dominica	4	7	7.7	7.3	Ruibal and Philibosian, 1970
<i>roquet</i>	Martinique	3	32	6.0	8.9	Hertz, 1977, 1981, this paper
<i>sagrei</i> ⁺	Abaco	2	19	?	10.6	Lister, 1976
Anoles from Complex Faunas:						
<i>allisoni</i> [*]	Cuba	2	4	4.7	2.3	Ruibal, 1961; Ruibal and Philibosian, 1970
<i>allogus</i> [*]	Cuba	2	3	2.2	1.7	Ruibal, 1961; Ruibal and Philibosian, 1970
<i>homolechis</i> [*]	Cuba	3	6	4.1	2.6	Ruibal, 1961; Ruibal and Philibosian, 1970
<i>lucius</i> [*]	Cuba	1	3	1.9	2.1	Ruibal, 1961; Ruibal and Philibosian, 1970
<i>sagrei</i> [*]	Cuba	1	4	3.7	3.0	Ruibal, 1961; Ruibal and Philibosian, 1970
<i>sagrei</i> ⁺	Exuma	3	19	?	6.8	Lister, 1976
<i>cybotes</i>	Hispaniola	3	29	9.6	8.8	Hertz, 1977; Hertz and Huey, 1981
<i>shrevei</i>	Hispaniola	1	5	1.7	3.1	Hertz, 1977; Hertz and Huey, 1981
<i>cooki</i>	Puerto Rico	3	19	5.7	4.8	Huey and Webster, 1976
<i>crisatellus</i>	Puerto Rico	8	29	8.8	10.1	Hertz, 1977; Huey and Webster, 1976
<i>evermanni</i>	Puerto Rico	2	21	2.6	3.5	Hertz, 1977; this paper
<i>gundlachi</i>	Puerto Rico	9	65	9.5	9.6	Hertz, 1977, 1981; Huey and Webster, 1976

[§]P = number of populations sampled.

[†]S = number of temporally restricted samples represented. See text for explanation.

^{*}THB calculated from midpoints of air temperatures ranges; mean air temperatures not available.

⁺Data read from figure.

internally consistent because nearly all studies were conducted during summer months.

Three general trends are apparent in Table 2. 1) All of the solitary species exhibit broad thermal habitat and thermal niches. 2) Some species from the complex faunas of Hispaniola and Puerto Rico are as broad-niched as are the solitary species. 3) There is a strong association between species thermal habitat breadth and species thermal niche breadth ($r_s = 0.829$, $P < 0.01$). These conclusions contradict Ruibal and Philibosian's (1970) hypothesis that solitary anoles may generally have broader niches than do species from the complex faunas (see also Huey and Webster, 1976).

Despite the clarity of the above conclusions, the data in Table 2 largely reflect differences in the sampling effort

devoted to different species. Among the 15 species, thermal habitat breadth is significantly rank correlated with number of populations sampled and with the number of hourly samples collected ($r_s = 0.765$ and 0.775 , respectively, P 's < 0.01). Similarly, species thermal niche breadth is also rank correlated with these variables ($r_s = 0.606$ with number of populations and $r_s = 0.835$ with number of hourly samples, P 's < 0.01).

We can limit the effect of this sampling bias by examining only those species in which three or more populations were sampled. (This restriction unfortunately eliminates all but one Cuban species from consideration.) Four species in the complex faunas (*crisatellus*, *cybotes*, *gundlachi*, and *sagrei*) exhibit niche breadths comparable in magnitude to those of three solitary anoles (*marmoratus*, *oculatus*, and *roquet*). Two other

well-sampled species from complex faunas (*cooki* and *homolechis*) exhibit narrow thermal niches, but the small breadth of the latter may be attributable to the limited number of temporally restricted samples available.

EURYTOPY AND THERMOREGULATORY VERSATILITY

The interspecific differences in thermal niche breadths listed in Table 2 reflect interactions between the diversity of ambient thermal regimes which populations occupy and the degree to which populations can alter the precision with which they thermoregulate in response to varying environmental conditions (Hertz, 1977, 1981; Hertz and Huey, 1981). The precision of thermoregulation in a population of small arboreal ectotherms such as *Anolis* can be estimated from the slope of the regression of body temperatures on air temperatures (Huey, 1981; Huey and Slatkin, 1976). A slope near zero indicates the independence of body and air temperatures (perfect thermoregulation), whereas a slope near one indicates a strong dependence of body temperature on air temperature (thermal passivity, Hertz, 1974). The thermoregulatory versatility of a species can therefore be estimated by the range of regression slopes among populations living in diverse thermal habitats (Hertz, 1977). Theoretically, this index of thermoregulatory versatility can vary between 0.0 and 1.0. Although this statistic can describe the range of effects of thermoregulatory behaviors, behavioral observations are still necessary to describe the nature and extent of thermoregulation (Hertz and Huey, 1981; Huey, 1982).

The niche breadths of forest-dwelling *Anolis* species in complex faunas may be determined by their stenotopy (i.e., occupation of a narrow range of habitats) and lack of thermoregulatory versatility. The best studied example is *gundlachi*, the species that is restricted to low

perches in the relatively dark montane rainforests of Puerto Rico. Populations of *gundlachi* are stenothermal, and, because of their thermal passivity at all sites (range of regression slopes = 0.05), body temperatures track air temperatures throughout its broad altitudinal range (Hertz, 1981). However, because populations of *gundlachi* experience narrow and distinct ranges of air temperatures at different altitudes, the species thermal habitat and thermal niche breadths are large. Ruibal (1961) has shown that populations of *allogus* and *lucius* are also stenothermal and thermally passive in the shaded forests of Cuba. However, because these species were not sampled in geographically and thermally diverse localities, one cannot estimate the magnitude of either their species breadths or their thermoregulatory versatility.

Several other stenotopic anoles in complex faunas also show low thermoregulatory versatility. However, because they are restricted to open habitats where the cost of thermoregulation is low (Huey, 1974, Huey and Slatkin, 1976), they thermoregulate precisely, and, as a result, their thermal niche breadths are apparently narrow. For example, *cooki* always regulates body temperature within narrow limits (low regression slopes in all populations, R. B. Huey, personal communication), probably because the potential for overheating is high in the deserts of Puerto Rico (Huey and Webster, 1976). In the montane savanna of Hispaniola, *shrevei* always thermoregulates carefully under sunny skies, presumably because of the benefit accrued from having a high body temperature in a habitat where air temperatures are usually low (Hertz and Huey, 1981).

In contrast to the low versatility shown by the thermoconformers and obligate thermoregulators described above, eurytopic species in complex faunas (e.g., *crisatellus*, *cybotes*, and *sagrei*) exhibit high thermoregulatory versatility. Populations of *crisatellus* living in shaded versus open habitats exhibit a range of

regression slopes (0.80) that virtually span the entire theoretical range from thermal passivity to perfect thermoregulation (Huey, 1974; Huey and Slatkin, 1976). Lister (1976) described an analogous situation for *sagrei* on Exuma in which the precision of thermoregulation varied with the amount of direct solar radiation that penetrated the vegetation in different habitats. Similarly, in *cybotes* sampled along an altitudinal gradient of 1,150 m, the precision of thermoregulation varied (range of slopes = 0.60) among populations. In each of these versatile species, animals predictably shift habitat and/or microhabitat, alter basking frequency, and vary times of activity with geographical changes in air temperature and/or the energetic costs of basking in different habitats. Thermoregulatory versatility in these species therefore arises from their ability to shift regulatory behavior with the occupation of different habitats.

It is difficult to assess the thermoregulatory versatility of the solitary species with great accuracy. Several researchers have noted that they show thermal passivity in many habitats (Brooks 1968 on *oculatus*; McMannus and Nellis 1973 on *acutus*) but exhibit opportunistic thermoregulatory behavior under some circumstances (Lazell, 1972; Ruibal and Philibosian, 1970 on *oculatus*; Huey and Webster 1975 on *marmoratus*). However, my data on three populations of *roquet* suggest at least a moderate degree of versatility (range of regression slopes = 0.59); precision of thermoregulation varied in response to altitudinal changes in air temperatures and basking opportunities (Hertz, 1981) in a fashion analogous to that seen in *cybotes* on Hispaniola (Hertz and Huey, 1981). Presumably, my estimate of versatility in *roquet* would be enlarged by the inclusion of a sample of *A. r. zebrilus* or *A. r. salinei*, the subspecies that occupy the most arid and xeric habitats on Martinique (Lazell, 1972).

The above analysis suggests that West Indian *Anolis* species have adopted one

of three behavioral-ecological "phenotypes" with respect to their thermal biology: 1) thermal passivity under a broad range of ambient conditions; 2) careful thermoregulation whenever possible; and 3) thermoregulatory versatility as a response to varying conditions. (We must, of course, use caution in specifying the phenotype of any species because estimates of thermoregulatory versatility will, like estimates of niche breadth, be sensitive to the number of populations sampled as well as the seasonal schedule of sampling.) *Anolis* species from the complex faunas exhibit any one of the three phenotypes; examples of intermediate phenotypes undoubtedly will be found with additional research. The solitary anoles, however, all exhibit the versatile phenotype.

THERMOREGULATORY VERSATILITY AND COLONIZING POTENTIAL

Anoles are thought to have evolved in the moist tropical forest of the New World as eurythermal and passive species with relatively low activity temperatures (Huey and Webster, 1975). However, because these characteristics might be grossly maladaptive for the transoceanic dispersal of a small animal, thermal passivity and low activity temperatures were probably not ancestral within the West Indian members of this genus. Rather, the colonizing species from which the West Indian *Anolis* radiation is derived almost certainly exhibited thermoregulatory versatility as well as a tolerance of high temperatures (Heatwole *et al.*, 1969; Ruibal, 1967) and desiccation stress (Williams, 1969). These characteristics would enable a small ectotherm not only to survive transoceanic rafting but also to exploit the wide array of habitats available on the newly inhabited island. Thermoregulatory versatility and eurythermy can therefore be considered an ancestral "colonizing phenotype" in most, if not all, island colonizations by members of this genus. Williams (1969: 374-375) has constructed

a similar portrait of colonizing anoles in a more general context.

Thermoregulatory versatility and eurythermy in solitary anoles may not represent "ecological release" or "niche expansion" at all. Rather, these species may have simply retained the widely adaptive colonizing phenotype while exploiting diverse habitats in the absence of congeneric competitors. Morphological differentiation within the solitary anoles of the Lesser Antilles appears not to be related to their thermal ecology (Hertz, 1977; Ruibal and Philibosian, 1970), and these species exhibit relatively little physiological differentiation among populations (Hertz, 1977, 1980, 1981; Hillman *et al.*, 1979). Indeed, Lister's (1976) observations on *sagrei*, and, to some extent, those on *monensis* (but cf. Gorman and Stamm, 1975: 201) are the only reliable demonstration of ecological release of thermal biology among West Indian anoles; the physiological and genetic bases of this phenomenon have not been explored in these species.

Some species in the complex faunas of the Greater Antilles (e.g., *crisatellus*, *cybotes*, and *sagrei*) exhibit phenotypes that are similar to the hypothetical colonizing phenotype, and their occupation of disturbed habitats in the Greater Antilles provides further evidence of their colonizing abilities (e.g., Turner and Gist, 1970). Williams (1969) has, in fact, listed two of these species (*crisatellus* and *sagrei*) among the six best colonizers within West Indian anoles. If Williams's (1972) phylogeny for the Puerto Rican radiation is correct, however, we must assume that, for *crisatellus* at least, thermoregulatory versatility represents convergence to the colonizing phenotype, rather than retention of it.

CONCLUSIONS AND SUGGESTIONS FOR FUTURE RESEARCH

The above analysis does not in any way support the thesis that solitary anoles ex-

perience "ecological release" of their thermal biology. Nonetheless, most *Anolis* ecologists would probably agree that the field biology of solitary anoles is markedly different from that of species in the more complex Greater Antillean faunas. This statement is supported by the intuition of numerous field workers and stems from the belief that competition will almost inevitably constrain a species' ecological flexibility (see, e.g., MacArthur, 1972).

The failure of *Anolis* biologists to demonstrate the "ecological release" of thermal biology is perhaps a methodological, rather than a conceptual, failure. The data gathered during the last two decades apparently provide poor resolution of the biological realities that Rand (1964) and Williams (1972) have described as the "climatic habitat" of each species. Researchers should now adopt more sophisticated methodologies to test the hypotheses generated by theoretical ecologists. The simple measurement of lizard body temperatures and air temperatures, though useful for acquiring baseline information about the field biology of many species, is insufficient for testing general theory about the effects of competition.

The demonstration of ecological release in the thermal biology of these animals will require simultaneous analysis of three factors that interact to produce a species' "thermal biology": 1) the range of habitats and microhabitats available to a population; 2) the populations' use of the available environmental heterogeneity; and 3) the breadth of physiological capacity that has evolved in different populations. The use of biophysical models and micrometeorological measurements (Porter and Gates, 1969), in combination with observations of animals in a diversity of habitats, might adequately address the first two of these factors. Indeed, Roughgarden, *et al.* (1981) have successfully applied these techniques in a study of resource partitioning by anoles on Grenada and St. Kitts. An adequate analysis of the physi-

ological breadth of a population will require detailed studies of the thermal dependence of whole-animal physiological functions (e.g., locomotion, Huey and Stevenson, 1979; paper by Huey in this volume) in addition to the now typical measurements of rates of water loss and critical and preferred temperatures.

Finally, future research on this topic should be directed towards those species that Williams (1969) has identified as the most successful invaders of isolated islands in the Greater Antillean subregion of the Caribbean: *angusticeps*, *carolinensis*, *cristatellus*, *distichus*, *grahami*, and *sagrei*. These species, and their close relatives, will provide the best opportunities for the demonstration of ecological release. Lister's (1976) study of *sagrei* provides a good starting point for additional work: in the absence of detailed information on source island (Cuba) populations, our understanding is far from complete.

ACKNOWLEDGMENTS

I thank A. Garcia, M. Perez, and D. Viera for field assistance in Puerto Rico, and W. Hall and F. Wadsworth for the use of facilities at their disposal. R. B. Huey offered valuable criticism and discussion. W. H. Bossert, K. I. Miyata, A. S. Rand, R. Ruibal, T. W. Schoener, E. E. Williams, B. Wu, and E. Zouros all had something to say about previous drafts.

Financial support of field work was generously provided by grants from the Richmond and Anderson Funds of Harvard University, Sigma Xi, the Explorers Club, and the National Science Foundation Grants DEB 75-16334 and GB 37731X. The manuscript was prepared with the assistance of the Research and Travel Fund of Barnard College.

This paper is dedicated to the Principle of Unsympathetic Magic.

LITERATURE CITED

- BAKKEN, G. S. 1976. A heat transfer analysis of animals: unifying concepts and the application of metabolism chamber data to field ecology. *J. Thermal Biology*, 6: 337-384.
- BROOKS, C. R. 1968. Body temperatures of three lizards from Dominica, West Indies. *Herpetol.*, 24: 209-214.
- COWLES, R. B., AND C. M. BOGERT. 1944. A preliminary study of the thermal requirements of desert reptiles. *Bull. Amer. Mus. Nat. Hist.*, 83: 263-296.
- GORMAN, G. C., AND B. STAMM. 1975. The *Anolis* lizards of Mona, Redondo, and La Blanquilla: Chromosomes, relationships, and natural history notes. *J. Herpetol.*, 9: 197-205.
- GRANT, P. R. 1972. Convergent and divergent character displacement. *Biol. J. Linn. Soc.*, 4: 39-68.
- HEATWOLE, H., T. H. LIN, E. VILLALÓN, A. MUNIZ, AND A. MATTÁ. 1969. Some aspects of the thermal ecology of Puerto Rican anoline lizards. *J. Herpetol.*, 3: 65-77.
- HERTZ, P. E. 1974. Thermal passivity of a tropical forest lizard, *Anolis polylepsis*. *J. Herpetol.*, 8: 323-327.
- . 1977. Altitudinal variation in thermoregulatory strategies, physiological ecology, and morphology of some West Indian anoles. Doctoral Dissertation, Harvard University.
- . 1980. Responses to dehydration in *Anolis* lizards sampled along altitudinal transects. *Copeia*, 1980: 440-446.
- . 1981. Adaptation to altitude in two West Indian anoles. I. Field thermal biology and physiological ecology. *J. Zool., London*, 195: 25-37.
- HERTZ, P. E., AND R. B. HUEY. 1981. Compensation for altitudinal changes in the thermal environment by some *Anolis* lizards on Hispaniola. *Ecology*, 62: 515-521.
- HILLMAN, S., G. C. GORMAN, AND R. THOMAS. 1979. Water loss in *Anolis* lizards: evidence for acclimation and intraspecific differences along a habitat gradient. *Comp. Biochem. Physiol.*, 62A: 491-494.
- HUEY, R. B. 1974. Behavioral thermoregulation in lizards: importance of associated costs. *Science*, 184: 1001-1003.
- . 1982. Temperature, physiology, and the ecology of reptiles, pp. 25-91. In C. Gans and F. H. Pough (eds.), *Biology of the Reptilia*, Vol. 12, in press. New York, Academic Press.
- HUEY, R. B., AND M. SLATKIN. 1976. Cost and benefits of lizard thermoregulation. *Quart. Rev. Biol.*, 51: 363-384.
- HUEY, R. B., AND R. D. STEVENSON. 1979. Integrating thermal physiology and ecology of ecto-

- therms: A discussion of approaches. *Amer. Zool.*, **19**: 357-366.
- HUEY, R. B., AND T. P. WEBSTER. 1975. Thermal biology of a solitary lizard: *Anolis marmoratus* of Guadeloupe, Lesser Antilles. *Ecology*, **56**: 445-452.
- AND —. 1976. Thermal biology of *Anolis* lizards in a complex fauna: the *cristatellus* group on Puerto Rico. *Ecology*, **57**: 985-994.
- LAZELL, J. D., JR. 1972. The anoles (Sauria, Iguanidae) of the Lesser Antilles. *Bull. Mus. Comp. Zool.*, **143**: 1-115.
- LEVINS, R. 1968. *Evolution in Changing Environments*. Princeton, N.J., Princeton University Press.
- LISTER, B. C. 1976. The nature of niche expansion in West Indian *Anolis* lizards. I. Ecological consequences of reduced competition. *Evolution*, **30**: 659-676.
- MACARTHUR, R. H. 1972. *Geographical Ecology*. New York, Harper and Row.
- MCMANUS, J. J., AND D. W. NELLIS. 1973. Temperature and metabolism of a tropical lizard, *Anolis acutus*. *Comp. Biochem. Physiol.*, **45A**: 403-410.
- PORTER, W. P., AND D. M. GATES. 1969. Thermodynamic equilibria of animals with environment. *Ecol. Monogr.*, **39**: 227-244.
- RAND, A. S. 1964. Ecological distribution in anoline lizards of Puerto Rico. *Ecology*, **45**: 745-752.
- . 1967. The ecological distribution of the anoline lizards around Kingston, Jamaica. *Breviora Mus. Comp. Zool.* No. 272, pp. 1-18.
- RAND, A. S., AND P. J. RAND. 1967. Field notes on *Anolis lineatus* in Curacao. *Stud. Fauna Curacao Other Carib. Islds.*, **93**: 112-117.
- RAND, A. S., AND E. E. WILLIAMS. 1969. The anoles of La Palma: aspects of their ecological relationships. *Breviora Mus. Comp. Zool.* No. 327, pp. 1-18.
- ROUGHGARDEN, J. 1972. Evolution of niche width. *Amer. Nat.*, **106**: 683-718.
- ROUGHGARDEN, J., W. PORTER, AND D. HECKEL. 1981. Resource partitioning of space and its relationship to body temperature in *Anolis* lizard populations. *Oecologia*, **50**: 256-264.
- RUIBAL, R. 1961. Thermal relations of five species of tropical lizards. *Evolution*, **15**: 98-111.
- . 1967. Evolution and behavior in West Indian anoles, pp. 116-156. In W. Milstead (ed.), *Lizard ecology: a symposium*. Kansas City, Missouri, University of Missouri Press.
- RUIBAL, R., AND R. PHILIBOSIAN. 1970. Eurythermy and niche expansion in lizards. *Copeia*, **1970**: 645-653.
- SCHOENER, T. W. 1975. Presence and absence of habitat shift in some widespread lizard species. *Ecol. Monogr.*, **45**: 233-258.
- SCHOENER, T. W., AND A. SCHOENER. 1971a. Structural habitats of West Indian *Anolis* lizards. I. Lowland Jamaica. *Breviora Mus. Comp. Zool.* No. 368, pp. 1-53.
- AND —. 1971b. Structural habitats of West Indian *Anolis* lizards. II. Puerto Rican uplands. *Breviora Mus. Comp. Zool.* No. 375, pp. 1-39.
- SOULÉ, M. 1963. Aspects of thermoregulation in nine species of lizards from Baja California. *Copeia*, **1963**: 107-115.
- TURNER, F. B., AND C. S. GIST. 1970. Observations of lizards and frogs in an irradiated Puerto Rican forest, Chapter E-2, pp. E25-E49. In H. T. Odum and R. F. Pigeon (eds.), *A tropical rain forest. A study of irradiation and ecology*. Div. Tech. Inf., U.S. A.E.C. Publ. TID-24270 (PRNC-138).
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, **44**: 345-389.
- . 1972. The origin of faunas. Evolution of lizard congeners in a complex island fauna: a trial analysis, Vol. 16, pp. 47-88. In Th. Dobzhansky, M. K. Hecht, and W. C. Steere (eds.), *Evolutionary Biology*. New York, Appleton-Century-Crofts.
- WILSON, E. O. 1961. The nature of the taxon cycle in the Melanesian ant fauna. *Amer. Nat.*, **95**: 169-193.

Natural Variation in Body Temperature and Physiological Performance in a Lizard (*Anolis cristatellus*)

RAYMOND B. HUEY¹

ABSTRACT. Field data on changes in body temperature (with habitat, time of day, and thermoregulatory behavior) of the Puerto Rican lizard *Anolis cristatellus* are integrated with laboratory data on the effect of body temperature on relative sprint speed. Lizards that thermoregulate in an open park in the early morning should be able to run slightly faster than thermoconforming lizards in a nearby forest. At midday, however, ambient temperatures in the open are high, and the T_b of lizards in the open is elevated above thermal preferences; lizards in the forest may then have a performance advantage. The results suggest that the temporal and spatial variation in physiological performance of lizards in nature can be dynamic.

INTRODUCTION

Body temperatures of some active lizards vary in time and in space (Avery, 1982). The extent to which such variation influences physiological and thus ecological performance is a fundamental, but understudied problem in physiological ecology (Crowder and Magnuson, 1982; Huey, 1982). This problem can be approached by determining how the physical environment influences body temperature (Porter *et al.*, 1973), by investigating the thermal sensitivities of physiological systems that have direct ecological relevance (Bartholomew, 1958; Brett, 1971; Huey and Stevenson, 1979; Bennett, 1980), and then synthesizing ecological and physiological data

(Huey and Slatkin, 1976; Waldschmidt, 1978; Christian and Tracy, 1981; Muth, 1980; Huey, 1982).

I first became interested in integrating physiology and ecology while on a field trip with Ernest E. Williams in the Caribbean and soon decided to pursue this topic for my dissertation under his direction. During this period, I grew to appreciate the relevance of whole-animal physiological performance such as locomotion to ecological analyses, an approach that Ernest supported. Consequently, it is my pleasure to dedicate this paper, which is a direct extension of my thesis research, to Professor Ernest E. Williams for this Festschrift in his honor.

This paper reports a preliminary investigation of the effects of temporal and spatial variation in body temperature on physiological performance in the Puerto Rican lizard *Anolis cristatellus*. Laboratory data on the thermal sensitivity of sprint locomotion are integrated with a previous analysis of body temperatures in nature (Huey, 1974).

This integration of physiological performance and ecology focuses on sprint speed as a single, but ecologically important, metric of physiological performance. Sprint speed can strongly affect the efficiency of predation, escape from predators, and social interactions (Rand, 1964; Webb, 1976; Elliott *et al.*, 1977; Bennett, 1980; Christian and Tracy, 1981; Huey and Hertz, 1982). Ultimately, however, this approach must be gen-

¹Department of Zoology NJ-15, University of Washington, Seattle, WA 98195, U.S.A.

eralized to incorporate additional physiological (e.g., digestion, growth, reproduction) and ecological (e.g., environmental productivity, risks of predation) information (Huey and Slatkin, 1976).

BACKGROUND ECOLOGICAL DATA

The body temperature (T_b) of the trunk-ground lizard *Anolis cristatellus* depends upon habitat, time of day, and thermoregulatory behavior. In open habitats where basking sites are frequently available on lower tree trunks, these lizards bask early and late in the day and behaviorally control body temperatures at relatively high and constant levels (Fig. 1, from Huey, 1974). However, in a nearby forest where basking sites are few and distant, these lizards rarely bask, and thus are thermoconformers. Consequently, average body temperatures in the forest are relatively low and variable (Fig. 1). (Males and females did not differ significantly in T_b in any census.) Comparable patterns have been documented for other *Anolis* (Lister, 1976; Lee, 1980; Hertz and Huey, 1981).

In laboratory thermal gradients, lizards from both forest and open habitats prefer

an average body temperature of 29.6°C (Huey and Webster, 1976). In nature, lizards in the open habitat reach these temperatures very early in the day by basking (Fig. 1). However, the high environmental temperatures in this habitat at midday apparently force these lizards to be active at temperatures a few degrees above preferred levels (Note: this forcing should be verified with a biophysical analysis.) In the shaded forest, however, lizards are active at T_b below preferred levels until late morning. Thereafter, "forest" lizards are active at near-preferred T_b for the rest of the day.

MATERIALS AND METHODS

Six adult male *Anolis cristatellus* used in these experiments were collected along the northern coast of Puerto Rico near the site of the field study (Pta. Salinas) in the summer of 1979 (July), and acclimated for 10 days to a 13:11 L:D cycle and a temperature of 28°C. Crickets (dusted with Vionate® vitamin-mineral supplement) were provided every other day.

Lizards were stimulated to sprint on a horizontal 2.4 × 0.2 m racetrack (rubberized substrate). The track contained 12 photocell stations at set distances (2 m maximum separation). When a lizard broke the first photocell beam, a multi-channel timer was activated, and the time when each subsequent beam was broken was automatically measured and stored in memory (Huey *et al.*, 1981). The time for each 0.25 m interval was determined, and the fastest time converted to speed (to the nearest 0.1 m s⁻¹).

Each lizard was raced daily (6 trials/temperature, 1 temperature/day). The sequence of temperatures (Fig. 2) was random, except that the trials at 32°C were run last to minimize the possibility of injury resulting from exposure to high T_b .

The day after the sprint trials were completed, lizards were transferred to 15°C for 1 h and then cooled (~½°C

Figure 1. Body temperatures of *Anolis cristatellus* in an open and in a forest habitat in lowland Puerto Rico (redrawn from Huey, 1974). Vertical lines represent range, horizontal lines indicate means, and boxes enclose 95% confidence limits of the means.

Figure 2. Normalized sprint speeds of *Anolis cristatellus* (mean $\pm$ 95% confidence limits) as a function of body temperature. The arrow indicates the average preferred body temperature of lizards in a laboratory thermal gradient (data from Huey and Webster, 1976).

min⁻¹) until they lost the righting response (Critical Thermal Minimum). Two days later lizards were transferred to 30°C for 1 h and then heated until they reached the Critical Thermal Maximum, following procedures of Huey and Webster (1976).

Sprint speeds and critical temperatures for each lizard were fitted to a product-exponential curve (Huey and Stevenson, 1979). These equations were then solved to determine the "optimal" T_b for sprint performance (T_b where speed maximized) and the "thermal performance breadth" range of T_b over which sprint speed is greater than some arbitrary level, see Huey and Stevenson (1979). Tolerance ranges (the difference between the Critical Thermal limits) were determined directly.

RESULTS

THERMAL SENSITIVITY OF SPRINT SPEED

Normalized sprint speeds of *Anolis cristatellus* are diagrammed in Fig. 2, and several statistical summaries are presented in Table 1. The lower tem-

perature threshold for locomotion is about 9°C, and the upper threshold is about 37°C. Sprint speed increases from the lower threshold T_b until 30°C, which approximates the average optimal temperature for sprint speed. At higher temperatures, sprint speeds drop precipitously. Interestingly, the optimal $\bar{T}_b$ for sprint speed is nearly identical to the preferred $\bar{T}_b$ of these lizards in laboratory thermal gradients (Table 1).

Thermal performance breadth measures the extent to which physiology is independent of temperature at ecologically meaningful T_b (Huey and Stevenson, 1979). The range of T_b over which these lizards are able to run at 95% and 80% of maximal speed is 6.5° and 12.7°C, respectively.

INTEGRATING SPRINT DATA WITH FIELD TEMPERATURES

The average T_b of lizards from each habitat and time (Fig. 1) are used as an index of temporal and spatial variation in T_b of *Anolis cristatellus*. Of course, this measure ignores within-sample variability in T_b , which is sometimes large in the open habitat (Fig. 1). To summarize the average effect of T_b on sprint velocity, I used the product-exponential equations to calculate relative performance of each individual at each of the $\bar{T}_b$'s measured in the field (Fig. 1) and then averaged performance among individuals. This measure ignores within and between-individual variability (Table 1), but nevertheless serves as a useful first-approximation of how the average relative performance of *cristatellus* might change as a function of habitat, time, and behavior.

Activity begins near sunrise in natural populations. Body temperatures at this time are cool ($\sim$ 24.6°C), but lizards should be able to sprint at about 92% of maximum speed. In less than an hour, lizards in the open elevate their temperatures to about 30.6°C, a temperature at which sprint speed is near maximal (Figs.

TABLE 1. STATISTICS DESCRIBING THERMAL SENSITIVITY OF SPRINT SPEED FOR *ANOLIS CRISTATELLUS* FROM LOW-LAND PUERTO RICO (N = 6).

Character	Temperature °C	
	$\bar{X} \pm \text{s.e.}$	range
Critical Thermal Minimum	8.92 ± 0.50	7.0–10.2
"Optimal" T_b	29.53 ± 0.85	27.0–32.1
Preferred T_b	29.6 ± 0.59	
Critical Thermal Maximum	37.00 ± 0.20	36.6–38.0
Tolerance Range	28.12 ± 0.48	26.8–29.8
Thermal Performance Breadth		
95% of maximum	6.48 ± 0.39	5.4–7.7
80% of maximum	12.73 ± 0.73	10.4–15.2

Tolerance range = CTMax–CTMin.

Preferred T_b from Huey and Webster (1976).

2, 3). In contrast, body temperatures and sprint speeds of the "forest" lizards, which do not bask, remain at lower levels until 1000 or 1100 h (Fig. 3). Thus, lizards in the forest may have slightly submaximal locomotor abilities in the early morning.

By midday, ambient temperatures are hot. Lizards in the forest are finally active at T_b near preferred levels (Fig. 1), and thus their sprint speeds are high and remain high for the remainder of the day (Fig. 3). In contrast, T_b 's of lizards in the open are several degrees above preferred levels (Fig. 1), and sprint speeds appear slightly reduced (Fig. 3). The apparent, slight hyperthermia of lizards in the open should also increase metabolic rate and evaporative water loss, thereby further increasing the possible physiological disadvantage of the open habitat at midday.

COMPARISONS WITH OTHER SPECIES

Data on thermal sensitivity of sprint speeds are now available for several lizards. The close correspondence between the preferred T_b and the optimal T_b for sprint locomotion of *A. cristatellus*, a species with a low thermal preferendum (Table 1), is also found in several thermophilic species (*Cnemidophorus*, *Dipsosaurus*, *Sceloporus*; Tracy, 1979;

Bennett, 1980; Huey, unpublished data). Curiously, however, two species with low thermal preferenda do not show this correspondence (*Eumeces*, *Gerrhonotus*; Bennett, 1980). Further studies are required to determine whether there is a general pattern (or patterns) among lizards.

Data on thermal niche breadth (Table 1) support the proposition that sprint abilities of lizards are somewhat temperature independent (Bennett, 1980): *A. cristatellus* can sprint at 95% of maximum speed over a 6.5°C range of T_b .

Figure 3. Predicted relative sprint speeds of *Anolis cristatellus* in open and forest habitats as a function of time of day. Average values only are plotted.

Nevertheless, the actual diurnal variation in T_b in nature (Fig. 1) seems to be sufficiently large to affect sprint abilities of *A. cristatellus*. Whether this pattern is valid for other species and for other physiological activities of *A. cristatellus* remains to be determined.

Data on thermal niche breadth are also relevant to an hypothesis (Tracy, 1979) concerning the relative degree of physiological specialization for temperature in frogs versus lizards. Because high rates of evaporative water loss hinder the ability of many frogs to control T_b by basking (Tracy, 1976), selection might have favored a relatively temperature-independent physiology in frogs. Indeed, data on the thermal performance breadths (80% level) for *Rana clamitans* (Huey and Stevenson, 1979) and for *Anolis cristatellus* (21.7° and 12.7°C, respectively), are in accord with Tracy's (1979) prediction.

Interestingly, even though the thermal performance breadths of *Rana* and *Anolis* differ strikingly (1.7X), the tolerance ranges for these two species are similar (*Rana* = 30.0°C, *Anolis* = 28.8°C). This observation supports the argument that selection for tolerance range may be independent of that for thermal performance breadth (Huey and Stevenson, 1979; Huey, 1982).

DISCUSSION

If the preferred temperatures of lizards reflect underlying physiological performance (Dawson, 1975), then *A. cristatellus* in open habitats might have a physiological advantage over "forest" lizards in the morning; but "forest" lizards might have an advantage at midday. On the other hand, if some aspects of physiology are relatively independent of T_b near the thermal preferendum (see Bennett, 1980), then the observed variations in T_b (Fig. 1) of *Anolis cristatellus* might not be of sufficient magnitude to impair physiological performance.

Data available to evaluate these alternatives are preliminary rather than definitive: 1) the sample sizes are small (only 6 individuals, only 6 trials/ T_b), 2) the temperature intervals are too broad for detailed analysis (see Fig. 2), 3) the acclimation was at a constant temperature rather than at natural cycles of T_b ; and 4) the performance of all "forest" and of "open" lizards at a given T_b is assumed identical. Keeping these limitations in mind, the data suggest that the observed natural variation in T_b (Fig. 1) is of sufficient magnitude to influence one aspect of the physiological performance of *Anolis cristatellus* in lowland Puerto Rico. In particular, habitat, time, and behavior appear to affect sprint performance. For example, "open" lizards, which bask frequently, may have a locomotor advantage for a few hours in the early morning, but perhaps not at midday. Interestingly, because temperature levels in the forest are warm at midday, "forest" lizards are then active at T_b near optimal for sprinting even without ex-

Figure 4. Body temperature affects maximum speed, mean acceleration, and strike success of gopher snakes (*Pituophis melanoleucus*) striking at mice (data from Greenwald, 1974). However, neither relative speed or acceleration has a 1:1 relationship with relative success (slope of each relationship is plotted).

hibiting thermoregulatory behaviors (Huey, 1974).

The dynamics apparent in these patterns encourage further investigation. In particular, it would be interesting to measure the effect of T_b on digestion, metabolism, and water balance and to learn whether populations in the forest and open differ in food availability or in average growth rates. Moreover, by repeating such studies during winter, when the T_b differential between "forest" and "open" lizards should be exaggerated (see Lister, 1976), the importance of season could be elucidated as well.

Because of the importance of speed to the ecology of lizards (Bennett, 1980; Christian and Tracy, 1981; Huey and Hertz, 1982), the described physiological patterns should have considerable ecological implications. However, predicting the magnitude of these ecological effects is risky from sprint speed alone. Acceleration and reaction times are sometimes very important in predator-prey encounters (Webb, 1976; Elliott *et al.*, 1977; Huey and Hertz, in preparation). Moreover, the actual decrement in rate of food capture that results from a 5% decrement in sprint performance depends in part upon the biology of the prey as well as that of the predator (Huey and Stevenson, 1979). For example, body temperature affects the relative acceleration, velocity, and percent success of gopher snakes (*Pituophis melanoleucus*) striking at mice (Greenwald, 1974). Nevertheless, the two laboratory measures of sprint performance (acceleration, velocity) are only fair predictors of actual strike success (Fig. 4). This problem of scaling physiological to ecological performance is fundamental but presently elusive (Huey, 1982).

ACKNOWLEDGMENTS

This research was supported by the National Science Foundation under

Grants DEB 78-12024 and GB 37731X, the Graduate School Research Fund of the University of Washington, the Miller Institute for Basic Research in Science, and the Museum of Vertebrate Zoology (University of California, Berkeley). I thank C. R. Tracy and P. E. Hertz for comments on the manuscript. Finally, I thank the editors for the opportunity to contribute to this Festschrift.

LITERATURE CITED

- AVERY, R. A. 1982. Field studies of reptilian thermoregulation, pp. 93-166. In C. Gans and F. H. Pough (eds.), *Biology of the Reptilia* Vol. 12, Physiology (B). New York, Academic Press.
- BARTHOLOMEW, G. A. 1958. The role of physiology in the distribution of terrestrial vertebrates, pp. 81-95. C. L. Hubbs (ed.), *Zoogeography*. AAAS, Washington, D.C.
- BENNETT, A. F. 1980. The thermal dependence of behavioural performance of small lizards. *Anim. Behav.*, **28**: 752-762.
- BRETT, J. R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Onchorhynchus nerka*). *Amer. Zool.*, **11**: 99-113.
- CHRISTIAN, K. R., AND C. R. TRACY. 1981. The effect of the thermal environment on the ability of hatchling Galapagos land iguanas to avoid predation during dispersal. *Oecologia*, **49**: 218-223.
- CROWDER, L. B., AND J. J. MAGNUSON. 1982. Cost-benefit analysis of temperature and food resource use: a synthesis with examples from the fishes. In W. P. Aspey and S. I. Lustick (eds.), *Behavioral Energetics: Vertebrate Costs of Survival*. Columbus, Ohio State Univ. Press, in press.
- DAWSON, W. R. 1975. On the physiological significance of the preferred body temperatures of reptiles, pp. 443-473. In D. M. Gates and R. Schmerl (eds.), *Perspectives of Biophysical Ecology*. New York, Springer-Verlag.
- ELLIOTT, J. P., I. MCT. COWAN, AND C. S. HOLLING. 1977. Prey capture by the African lion. *Can. J. Zool.*, **55**: 1811-1828.
- GREENWALD, O. E. 1974. Thermal dependence of striking and prey capture by gopher snakes. *Copeia*, **1971**: 98-106.
- HERTZ, P. E., AND R. B. HUEY. 1981. Compensation for altitudinal changes in the thermal environment by some *Anolis* lizards on Hispaniola. *Ecology*, **62**: 515-521.
- HUEY, R. B. 1974. Behavioral thermoregulation in

- lizards: importance of associated costs. *Science*, **184**: 1001-1003.
- . 1982. Temperature, physiology, and the ecology of reptiles, pp. 25-91. In C. Gans and F. H. Pough (eds.), *Biology of the Reptilia* Vol. 12, Physiology (B). New York, Academic Press, in press.
- HUEY, R. B., AND P. E. HERTZ. 1982. Effects of body size and slope on sprint speed of a lizard *Stellio (Agama) stellio*. *J. exp. Biol.*, **97**: 401-409.
- HUEY, R. B., W. SCHNEIDER, G. L. ERIE, AND R. D. STEVENSON. 1981. A field-portable racetrack for measuring acceleration and velocity of small cursorial animals. *Experientia*, **37**: 1356-1357.
- HUEY, R. B., AND M. SLATKIN. 1976. Costs and benefits of lizard thermoregulation. *Quart. Rev. Biol.*, **51**: 363-384.
- HUEY, R. B., AND R. D. STEVENSON. 1979. Integrating thermal physiology and ecology of ectotherms: a discussion of approaches. *Amer. Zool.*, **19**: 357-366.
- HUEY, R. B., AND T. P. WEBSTER. 1976. Thermal biology of *Anolis* lizards in a complex fauna: the *cristatellus* group on Puerto Rico. *Ecology*, **57**: 985-994.
- LEE, J. C. 1980. Comparative thermal ecology of two lizards. *Oecologia*, **44**: 171-176.
- LISTER, B. C. 1976. The nature of niche expansion in West Indian *Anolis* lizards I: ecological consequences of reduced competition. *Evolution*, **30**: 659-676.
- MUTH, A. 1980. Physiological ecology of desert iguana (*Dipsosaurus dorsalis*) eggs: temperature and water relations. *Ecology*, **61**: 1335-1343.
- PORTER, W. P., J. W. MITCHELL, W. A. BECKMAN, AND C. B. DEWITT. 1973. Behavioral implications of mechanistic ecology. *Oecologia*, **13**: 1-54.
- RAND, A. S. 1964. Inverse relationship between temperature and shyness in the lizard *Anolis lineatopus*. *Ecology*, **45**: 863-864.
- TRACY, C. R. 1976. A model of the dynamic exchanges of water and energy between a terrestrial amphibian and its environment. *Ecol. Monogr.*, **46**: 293-326.
- . 1979. Further thoughts on anuran thermoregulation. In E. H. Burt (ed.), *The Behavioral Significance of Color*. New York, Garland Publishers.
- WALDSCHMIDT, S. 1978. Monthly variation in thermoregulatory behaviors and space utilization in the lizards *Uta stansburiana* and *Sceloporus undulatus*. Unpublished M.S. Thesis, Colorado State University, Fort Collins.
- WEBB, P. W. 1976. The effect of size on the fast-start performance of rainbow trout *Salmo gairdneri*, and a consideration of piscivorous predator-prey interactions. *J. exp. Biol.*, **65**: 157-177.

On the Voluntary Departure of Lizards from Very Small Islands

THOMAS W. SCHOENER¹

AMY SCHOENER²

ABSTRACT. Individuals of the iguanid lizard *Anolis sagrei* were placed on extremely small fragments of natural islands and their subsequent behavior recorded. Between a third and half of the 54 trials resulted in the lizard leaping into the sea and, when allowed, swimming and floating to shore. Lizards that departed showed more exploratory activity than those not jumping; lizards departed more during sunny than cloudy weather. Results support the hypothesis that lizards will leave islands of their own volition if those islands are inhospitable enough.

INTRODUCTION

A significant portion of Ernest E. Williams's scientific contribution has been to the field of island biogeography. Through an examination of distributional patterns in the Caribbean, he inferred the particular characteristics of *Anolis* lizards that predisposed them for island colonization (Williams, 1969). He concluded that the very abundant species of open vegetation, *A. carolinensis* and *A. sagrei*, were especially apt colonizers. Williams's work in this area to a large degree inspired our research on the lizards of very small islands, research begun in 1974 and still underway.

The first stages of this research involved several expeditions by boat, during which we visited approximately 600

islands spanning most of the Bahamas. Among many other characteristics, we recorded the diurnal lizard species present on these islands, including *A. sagrei* and *A. carolinensis*. Our studies revealed that an enormous number of small islands in the Bahamas had at least one species of lizard: single-species islands range down to 0.1 to 2.0 acres (1 acre $\approx$ 0.4 ha), depending on the general locality.

Surprising though these results were to us, we could still ask the question: why do lizards not occur on yet smaller islands? What causes extinction on such islands, assuming the lizards reach them occasionally, or at least were there before the islands became islands, that is before the final post-Pleistocene submergence?

To investigate this problem, in 1977 we initiated a series of experiments, seeding 30 islands smaller than the smallest inhabited-island size with propagules of *A. sagrei* or the iguanid *Leiocephalus carinatus*. A propagule comprised 5 to 10 adult individuals, in the ratio 3 females: 2 males. The details of these introductions will be reported elsewhere. Here the following facts are relevant. First, of the 25 islands seeded with *A. sagrei*, three lost their entire complement of lizards in four to five days; of the five islands seeded with *L. carinatus*, one became empty in eight days. One of the *sagei* introductions even involved 10 individuals—all disappeared without trace. These four islands, measured in terms of vegetated area,

¹Department of Zoology, University of California, Davis, California 95616, U.S.A.

²Department of Oceanography, University of Washington, Seattle, Washington 98195, U.S.A.

were the smallest representatives from their respective localities: vegetated areas for *sagrei* were 6.3 to 39 ft² (total areas 3,020–4,750 ft²). Second, somewhat larger islands lost their individuals more gradually. Even now, however, four years after the introductions, islands down to three orders of magnitude smaller than the smallest islands naturally having *sagrei* still have populations, many thriving, that descended from our propagules. As we will report elsewhere, this second result plus some others suggests to us that periodic catastrophes, in this case hurricanes, set the naturally observed single-species island size well above that inhabitable by lizards most of the time.

But what about the first result? What happened to the lizard propagules on our smallest, most miserable islands? They clearly did not starve to death or desiccate, because initially healthy *sagrei* can live without food or water for longer than four to five days, and because the scalloped, pock-marked surfaces of the islands provide much shade for small animals. Weather was good during this time: lizards could not have been washed away by waves. Predators are more likely, but still unlikely. Terrestrial lizard predators, such as certain hawks, cannot inhabit islands as tiny as those in question. Transient marine birds—herons, terns, gulls, and oystercatchers—are a possibility; all were commonly seen in the general area and a few were seen on the islands in question. However, none of these birds is primarily adapted for catching quick and clever lizards, and it is hard to imagine that they would have found it profitable, much less possible, so totally to wipe out our propagules.

We are, however, left with another alternative. Perhaps the lizards, after sizing up their desperate plight, simply decided to leave the islands of their own volition. Better islands lie nearby and are perhaps visible to lizards, and the future fitness of individuals not leaving would with certainty be zero. This hypothesis is

made less outrageous by an additional set of observations. Some lizards, and *sagrei* in particular, are excellent swimmers and especially floaters. We discovered this in the field, and later in the laboratory we floated lizards in wave-tanks with seawater for varying periods of time (Schoener and Schoener, in preparation). Results show that *sagrei* commonly can survive, floating without aid, for 12 hours, and some can float for at least 24 hours with no visible harm. Hence a lizard, jumping into the relatively quiet waters of our study sites, could quite conceivably reach a larger island, were the currents pulling favorably.

Despite this rather optimistic assessment, we would be more convinced that lizards departed islands voluntarily were we actually to witness it. We could repeat exactly the seeding of the same islands used in 1977, but the likelihood is that we would have to spend days on the islands before seeing a departure. We therefore decided to perform a series of experiments to encourage early departures, making certain variables as extreme as possible, much as physiologists sometimes do. We placed individual lizards on even smaller, more miserable islands than those of the 1977 introductions and watched their behavior. Would they leave voluntarily, and if so, how would they go about it? These experiments are the subject of the present paper.

SUBJECTS AND METHODS

Our experimental procedure was to place a single lizard on a rock of cl-2' (1' = 0.305 m) diameter and record its subsequent behavior. If the lizard jumped from the rock immediately upon release, the trial was not counted. Rocks were placed on the average 5 to 10' from shore, although tidal flux caused considerable variation in this distance. The rock on which the lizard was placed was set upon

one or more additional rocks such that at low tide the lizard was still on an island. Lizards were watched until 1) they departed from the rock, or 2) if they did not depart, for varying lengths of time in half-hour segments up to 6 hours. In the second case, of course, we followed no stopping rules, since we had no idea how long a lizard might remain on a rock before departing, if it departed at all. However, we discovered empirically that no lizard departed a rock once it performed no major movement (body turn or crawl; see below) for a half-hour. As it turned out, we stopped observing four subjects less than a half-hour since their last major movement, and because these individuals might well have departed their rocks, we excluded them from certain comparisons (see below).

For each lizard we recorded details of its behavior while on the rock, including head turns, body turns, number of crawls, and total distance crawled. We also recorded the fates of those individuals departing the rocks.

Experiments were performed April 19–24, 1979, and April 14–25, 1980, at Staniel Cay, Exuma Islands, Bahamas. To avoid wind as much as possible, we rotated trials between three sites with varying degrees of wind exposure. All sites were in fairly protected salt-water “creeks” with little or no current. Experiments were mostly performed between 930 and 1600 hrs (Fig. 1), during April the most favorable time of day for lizard activity. Total observation time for all trials combined was 5,796 minutes.

Lizard subjects were all *Anolis sagrei*, collected from Staniel Cay zero to three days before the experiments. This is the same source locality that supplied colonists for our original 1977 introduction. No individual was used more than twice; and with two exceptions, we did not re-use the same individual during the same day. With one exception, only adults were used, and we attempted to standardize size as much as possible by selecting the largest individuals.

RESULTS AND DISCUSSION

GENERAL OUTCOME OF THE EXPERIMENTS

We ran 54 trials, 29 with males (28 adults, 1 subadult) and 25 with females (all adult). Experiments resulted in three possible outcomes: I) the lizard leapt from the island; II) the lizard remained on the island until the experiment was stopped; and III) the island was nearly flooded by high tide, and then either a) the lizard swam off or b) the experiment was stopped. Among males, 12 jumped from the island and 17 did not; of the latter, 11 were in Class II and 6 were in Class III. Among females, 8 jumped from the island and 17 did not; of the latter, 13 were in Class II and 4 were in Class III.

To test for sexual differences in the tendency to depart from islands, we need to modify these numbers slightly. Because no lizard jumped from an island after remaining motionless for at least a half-hour (see above and below), we deleted the four cases in which the experiment was terminated less than a half-hour after the last major movement. We also deleted the one subadult-male trial. The modified data show that 12 of 26 males jumped, and 8 of 23 females did. In other words, about half the males jumped as compared to a third of the females. The sexual difference, however, is not statistically significant ($\chi^2 = 0.663$).

TEMPORAL AND CLIMATIC CORRELATES OF DEPARTURES

Although the primary purpose of our experiments was to determine whether or not lizards depart from islands a substantial proportion of the time, we can in retrospect search for associations between departure behavior and environmental characteristics.

Most lizards that departed did so close to midday. However, most of our observations were centered around midday also. As Figure 1 shows, males tended to jump from their rocks somewhat earlier

Figure 1. Times of day during which males and females jumped from islands. Top histogram is the percent observation time in various diel intervals for observations of males only. Bottom histogram is the same for females. ○ = times of jumps.

than females, but curiously, our observations also tended to be distributed over earlier portions of the day for males than females. To test whether males departed earlier in the day than would be expected from our observational bias, we performed a Kolmogorov-Smirnov 1-sample test. In this test the expected distribution gives the relative amount of time we spent observing lizards for each diel interval, and the observed distribution gives the proportion of jumps during each of those intervals. Even though males tended to jump earlier than expected from our observational bias, the difference was not statistically significant (maximum difference in cumulative frequency, $D = 0.205$). And, while females tended to jump later than expected, this difference was also not statistically significant ($D = 0.139$). Hence in the absence of more data, we must conclude that departure times for both

sexes were scattered randomly throughout the day.

Considerable variation in wind speed occurred over the course of our experiments. We therefore tested whether or not animals tended to depart more frequently during times of light wind. Wind speeds were divided into two categories: high and moderate-to-slight (We did not measure wind speed, but during the times of our experiments the categories were totally discrete and obvious!). Four departures took place during high winds, and 16 took place during relatively windless times. During high winds 10 lizards did not jump and during light winds 24 did not jump. While a slightly greater fraction of lizards jumped during relatively still periods, the association is not statistically significant ($\chi^2 = 0.489$).

During 1980 we recorded the fraction of the observation time that was cloudy and the fraction that was sunny. Do liz-

ards jump more during sunny periods? Because air temperatures ranged between 24.8 and 28.0°C (in 1979; in 1980, temperatures were slightly higher), we might expect this to be so. Most of our observation periods were fairly sunny: only 13% had greater than 50% cloud cover. Therefore, to divide the data, we had to use a relatively low cutoff for "percent cloudy." Of the lizards that jumped, 15 of 16 did so during weather that was cloudy less than 15% of the time. Of the lizards that did not jump, 13 of 22 were watched during weather that was cloudy less than 15% of the time. This difference is statistically significant ($\chi^2 = 5.739$). If 5% clouds rather than 15% is used, a similar result is obtained ($\chi^2 = 4.968$). Thus lizards tended to depart from islands more on sunny than cloudy days.

MOVEMENT CHARACTERISTICS OF DEPARTING VS. NONDEPARTING LIZARDS

Table 1 gives movement statistics, measured per-unit-time, for four types of data: number of crawls, total inches (1 inch = 2.54 cm) crawled, number of body turns and number of head turns. The table shows that lizards eventually departing (Class I) had higher movement rates in nearly all categories than did lizards that stayed on the island (Class II). Lizards whose island was nearly covered by high tide (Class III) had rates intermediate between those of Classes I and II. Thus lizards that eventually jumped were relatively exploratory, crawling more and farther, reorienting more, and looking around more. For comparisons between Classes I and II, differences are significant for 1) mean crawls/min ($df = 19$, $t = 2.152$, one- or two-tailed $P < 0.05$), 2) mean inches crawled/min ($df = 19$, $t = 2.084$, one-tailed $P < 0.05$), and 3) mean body turns/min ($df = 25$, $t = 1.731$, one-tailed $P < 0.05$), but not for mean head turns/min ($df = 16$, $t = 1.703$; all tests corrected for

inequality of variances according to Bailey, 1959). Lizards that were (or possibly were) disturbed by water splashing over their rocks moved more than those not so disturbed, but not as much as lizards that actually jumped. Of Class III lizards, several were even partially covered by water before they swam off, though in no case did water cover the lizard's nostrils or mouth.

Table I also shows that males tended to move more than did females. Differences, however, are not usually as great as those between classes of outcomes. Indeed only in Class I were they consistent and high: males crawled about twice as far as females, reoriented about twice as often, and looked around about ten times as much! Of the several comparisons possible, only one is statistically significant, mean head turns/min: Class I only ($df = 8$, $t = 2.110$, one-tailed $P < 0.05$), all classes combined ($df = 21$, $t = 2.144$, one- or two-tailed $P < 0.05$). The greater exploratory behavior of males is a natural correlate of their tendency to leap more frequently from islands. Moreover, as shown elsewhere (Schoener and Schoener, 1980), males of *sagrei* on large islands are more likely to be mobile than are females.

DEPARTURE CHARACTERISTICS

For animals of Class I, the elapsed time since the start of the experiment before jumping into the water varied enormously, from 5 seconds to 208 minutes; the mean is 56.3 minutes. Males jumped on average slightly sooner than females (50.3 vs. 65.4 minutes), but the difference is not statistically significant. For all data combined, time before jumping (Fig. 2), and the number of individuals not jumping by a certain time, are approximately exponential. Similarly, Moermond (1979) found that times between movements roughly resembled exponential distributions in several species of Haitian anoles. Such distributions would result from a

TABLE 1. MOVEMENT CHARACTERISTICS FOR LIZARDS PLACED ON VERY SMALL ISLANDS.

Type of movement	Sex	I lizard eventually jumped*			II lizard stayed on island			III island nearly covered with water by end		
		N	$\bar{x}$	s	N	$\bar{x}$	s	N	$\bar{x}$	s
crawls/min	♂	9	.103	.104	11	.036	.037	6	.044	.039
	♀	7	.071	.064	13	.042	.051	4	.047	.025
inches crawled/min	♂	9	.225	.250	11	.052	.064	6	.061	.061
	♀	7	.111	.123	13	.069	.105	4	.058	.038
body turns/min	♂	9	.056	.056	11	.009	.008	6	.016	.018
	♀	7	.022	.030	13	.023	.053	4	.033	.023
head turns/min	♂	9	.315	.389	11	.074	.089	6	.101	.086
	♀	7	.039	.045	13	.038	.081	4	.076	.063

*Only cases where the lizard stayed on the island for more than five seconds were included.

constant per-unit-time probability of jumping or movement.¹

While Class I outcomes cannot be distinguished from those of Class II on the basis of total duration of the experiment, they are distinguishable on the basis of time since the last major movement (crawl or body turn). All animals that jumped did so within a half-hour of their last major movement, and most did so considerably sooner than this: only three jumped more than 5 minutes after their last major movement. Those animals not jumping were watched up to a total of 240 minutes after their last major movement, and 10 were watched more than an hour from this time (or from the start of the experiment, in cases of no movement at all).

All lizards that jumped did so with a decisive leap into the water. Jumping

distances varied from approximately 0.5 to 3.5'. Jumping was sometimes, but not always, oriented toward the nearest land. Some lizards that jumped were allowed to swim and float in the water until they reached shore, a process lasting up to five

Figure 2. Distribution of the total elapsed experimental time before jumping for those lizards (Class I) that jumped from islands.

¹This is well known in probabilists' circles but is given here for the interested. Let $k\Delta t$ be the probability of movement (e.g., jumping) in a small amount of time Δt . In G time units, the probability of not moving $P_{O,T} = (1 - k\Delta t)^G \approx e^{-kT}$, where $\Delta tG = T$. The exponential function $P_{O,T}$ is the same as the fraction of individuals not moving by time T . The fraction moving between times 0 and 1 is then $P_{M,0} = 1 - P_{O,1} = 1 - e^{-k}$, and between times t and $t + 1$ is $P_{M,t} = P_{O,t} - P_{O,t+1} = e^{-kt} - e^{-k(t+1)} = (1 - e^{-k})(e^{-kt}) = P_{M,0} P_{O,t}$. Thus the fraction moving in each of a set of unit time intervals is exponential.

minutes. Often they actively swam more than 4', and three individuals swam more than 10' (Fig. 3). Lizards would commonly change direction while floating, in several cases orienting toward the nearest above-water object, such as a rock or mangrove shoot. In no case did a lizard drown, and most individuals were kept for one to three days after the experiment with no apparent ill effects (indeed, some were reused in a second trial).

CONCLUSIONS

The major objective of our experiments, to show that lizards commonly leave very small islands on their own volition, has been accomplished: between a third and half the trials resulted in this outcome. Our results during sunny versus partly-to-entirely cloudy periods suggest that, were weather conditions sunnier (and perhaps warmer), the fraction departing would have been greater. Many of the lizards not jumping were cold to the touch when removed, and a number had curled up in rock crevices and become motionless. Thus it ap-

peared that sunny periods were generally those in which activity was possible (or optimal), rather than those in which activity was engendered by thermal stress. That thermal stress was not the primary motivation for departures is additionally indicated by the presence of a departure on at least one entirely cloudy day and by the availability of some shade on most of our rocks during most of the day. Moreover, lizards staying in the sun for long periods of time and never jumping showed no apparently ill effects. Nonetheless, because of uncontrolled climatic conditions, we do not wish to ascribe any significance to the actual fraction of individuals that jumped during our experiments.

Because our experiments were run at the same time of year and under the same general climatic conditions as the original 1977 introduction experiments, we do feel that we have adequately simulated most conditions of the latter. One might argue that the greater island size in the 1977 introductions makes the two sets of experiments not very comparable. Our guess is that exploratory behavior would have lasted longer on the larger islands, but that the eventual outcome would have been much the same. But of course we have not proven that lizards voluntarily depart from the same-sized island used in the 1977 experiments. Even if we had shown this with experiments of the present type, however, we could not have proven that lizards actually jumped from the islands in 1977. We have documented, however, an exploratory behavior on the part of lizards placed on very small islands, and a rather remarkable tendency for those lizards to leap into the sea rather than passively await death.

ACKNOWLEDGMENTS

Chris Bayer performed some of the experiments during 1980. Research was supported by National Science Foundation Grant DEB-22798.

Figure 3. Distribution of the distance swum (not including floating) for those lizards that jumped from islands (1 ft = 0.305 m).

LITERATURE CITED

- BAILEY, N. J. T. 1959. *Statistical Methods in Biology*. London, English Universities Press.
- MOERMOND, T. C. 1979. The influence of habitat structure on *Anolis* foraging behavior. *Behavior*, **70**: 147-167.
- SCHOENER, A., AND T. W. SCHOENER. in preparation. Experimental zoogeography: floatation of insular *Anolis* lizards.
- SCHOENER, T. W., AND A. SCHOENER. 1980. Densities, sex ratios and population structure in four species of Bahamian *Anolis* lizards. *J. Anim. Ecol.*, **49**: 19-53.
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, **4**: 345-389.

Experimental Evidence of Strong Present-Day Competition Between the *Anolis* Populations Of the Anguilla Bank—A Preliminary Report

JONATHAN ROUGHGARDEN¹

JOHN RUMMEL²

STEPHEN PACALA³

ABSTRACT. Preliminary experimental results indicate strong present-day competition between the *Anolis* populations native to the Anguilla Bank. *A. wattsi* is absent from xeric habitat in which we show it can successfully survive and reproduce. Introductions of *A. wattsi* to such habitat have low initial survival if *A. gingivinus* is present and higher initial survival if *A. gingivinus* is partially removed prior to the introduction. Furthermore, if *A. wattsi* is introduced in the presence of *A. gingivinus* then the introduced animals are forced into marginal territories. Introduced animals exhibit great site fidelity. The panting temperature of *A. wattsi* is lower than that of *A. gingivinus*. The apparent competitive exclusion of *A. wattsi* by *A. gingivinus* from xeric habitat seems to be caused by a carrying capacity disadvantage of *A. wattsi* relative to *A. gingivinus* in xeric sites, together with a high competition coefficient from *A. gingivinus* to *A. wattsi*. In addition, a four-year survey of three sites on St. Maarten reveals no change in the rank order of abundance of all the populations even though the absolute abundance of every population has fluctuated. The abundance of lizards on an off-shore cay fluctuated less than those on the St. Maarten sites during the same period. Hurricane Frederick in August 1979 caused no reduction in the abundance of the lizard populations at these sites, including the off-shore cay.

INTRODUCTION

Studies on the *Anolis* lizard populations on the island of St. Maarten have

from the beginning revealed exceptional results. Lazell (1972) in his taxonomic monograph on the anoles of the Lesser Antilles reported that the smaller species from St. Maarten, *Anolis wattsi*, was in danger of extinction. He reported also that *A. wattsi* from nearby Anguilla had already become extinct, a consequence of the destruction of its preferred woody habitat. Lazell felt also that *A. wattsi* was destined for extinction on St. Maarten because he believed the species to be concentrated in small, mesic wooded ravines. In contrast, the native populations of *A. wattsi* from the other banks in the Northern Lesser Antilles are found to be distributed throughout all habitats on the island, and were thought to be in no danger of extinction.

In 1972, Williams highlighted the island of St. Maarten as an exception to the biogeographic rule of body size that pertains to islands with two anole species in the Lesser Antilles. The basis for these rules comes from Schoener (1969). Only the island of St. Maarten, among all eight islands with two anole species, does not have a species whose body size in adult males exceeds 100 mm (SVL). In fact, *A. gingivinus* has a body size among adult males of approximately 60 mm (SVL), the size that is typical of endemic species who are the solitary residents of other islands. Since *A. wattsi* is only slightly

^{1,2,3}Department of Biological Sciences, Stanford University, Stanford, California 94305, U.S.A.

smaller than this "solitary size," there is great overlap in body size between the two species on St. Maarten.

Recently, Roughgarden *et al.* (1982) have shown that *A. watsi* is not restricted to small mesic wooded pockets on the island of St. Maarten, but is, in fact, distributed throughout the hills in the center of the island. The maximum elevation of these hills 424 m. The woods on these hills extend down into the xeric habitat at sea level along the walls of ravines. Lazell's observation that *A. watsi* was restricted only to ravines results from his observing *A. watsi* at the margin of its distribution where it approaches sea level elevations through these wooded ravines.

Roughgarden *et al.* (1982) also presented four lines of evidence that suggested the existence of strong present-day competition between the populations of St. Maarten. Moreover, they suggested that the absence of *A. watsi* from sea level habitat is caused by competitive exclusion, in the sense of the Lotka-Volterra equations, by *A. gingivinus*. Thus, they suggest not only that there is bona fide present-day ecological competition between these two populations, but also that it is actually strong enough to result in the competitive exclusion of one population by the other in a certain habitat. The experiments we report here are designed to provide an experimental test of this hypothesis. We report preliminary results which indicate that strong present-day competition is present; our data are not sufficient to establish that this competition is strong enough to explain the actual absence of *A. watsi* from sea level habitat on the island of St. Maarten. Also, we present results of censuses of three sites in St. Maarten which we have been monitoring since 1977. These sites serve to indicate whether there is any long term variation in population sizes on the Anguilla Bank.

RESULTS

Almost all the small cays of the Anguilla Bank contain populations of *A. gingivinus*. The vegetation on these cays is scrub with occasional or no trees at all. This vegetation is similar in aspect to, but more xeric, than the habitat at sea level on St. Maarten, from which *A. watsi* is absent. Most of the cays are formed from Pleistocene coral. For this experiment, we selected a cay known as Anguillita which is due west of the island of Anguilla. We have measured its maximum elevation above sea level as 6 m. This cay is easily accessible by boat from St. Maarten and Anguilla, but is rarely visited. Figure 1 presents a map of this cay. The map also shows the location of two of the study sites that we have established there. The censusing on the cays was conducted by a team of four people. Each site was censused for three days. The lizards were marked with a different color each day. The mark-recapture data were analyzed by finding the simplest log-linear statistical model consistent with the data and then projecting the total population size based on this statistical model. This census technique is presented in detail in Heckel and Roughgarden (1979).

EXPERIMENTAL INTRODUCTIONS TO ANGUILLITA

The first site (SK) was established as a pilot study in August 1979. This site is an isolated grove (clone) of Sea Grape trees (*Coccoloba uvifera* L.). This grove has a profile that has been sculptured by the prevailing wind and exhibits a classic Krummholtz shape. The maximum height of the trees is 4 m towards the leeward side and grades down to approximately 0.5 m towards the windward side of the site. The site is approximately circular with an area of 558.6 m².

In August 1979, we censused the resi-

dent population of *A. gingivinus* and estimated its size at approximately 196 lizards. Then we collected 103 *A. wattsi* from the hills of St. Maarten. Approximately half were collected from the Columbian Ravine at the western margin of the distribution of *A. wattsi*, near sea level. The other *A. wattsi* were collected from Pic du Paradis and are slightly greener in body color than the lizards near the species' border. Hence, it was possible that geographical variation existed in physiological ability to withstand drought stress (Hertz, 1981). Both collections had approximately a 50/50 sex ratio. The lizards were toe-clipped according to the location from which they were collected and were then released into the

center of the SK site. The resident *A. gingivinus* were left undisturbed.

Table 1 presents census data for this site. The population size of the resident *A. gingivinus* was essentially unchanged in March 1980. However, only 15 (approximately) remained of the 103 *A. wattsi* that had been introduced; eight distinct animals were seen. Furthermore, these *A. wattsi* were found solely at the periphery of this site where the vegetation was very low. No *A. wattsi* had assumed territories in the center of the grove. The survivors were generally paired, we observed approximately a 50/50 sex ratio, and individuals from both collection locales were equally represented among the survivors.

In October 1980, and March 1981 the populations of *A. gingivinus* at SK had continued to number approximately 200. The *A. wattsi* that had survived the first six months continued to survive in their territories at the periphery of the site. By March 1981 their number had dropped to about 10. This experiment demonstrates the ability of *A. wattsi* of both sexes and from two collection locales to survive in the environment of the cay for more than one year.

Our next study site (MK) was established in March 1980. It, too, is a Krummholtz of trees (Manzanilla or Manchineel—*Hippomane mancinella* L., and Sea Grape—*Coccoloba uvifera* L.). The area of this site is 262.6 m², approximately one half as large as our first site. The resident population of *A. gingivinus* was estimated at 116, also about one half that of SK. We removed 41 residents, leaving a population of 75 *A. gingivinus*, and we then introduced 55 *A. wattsi* from St. Maarten into the center of the site.

When we returned in October 1980, we discovered that the population of *A. gingivinus* had returned to its original size (116 animals). We attributed this

ANGUILLITA (B.W.I.)

Figure 1. Map of Anguillita.

TABLE 1. CENSUS DATA—ANGUILLITA.

Site	Date		number <i>A. gingivinus</i>	number <i>A. wattsi</i>	density <i>A. gingivinus</i>	density <i>A. wattsi</i>
SK area = 558.6 sq. m.	Aug. 79		196.3 (7.2)	103 intro	35.1 (1.3)	18.4 intro
	May 80		204.6 (9.3)	14.5 (7.0) (8 seen)	36.6 (1.6)	2.6 (1.3)
	Oct. 80		180.0 (6.6)	15.0 (2.1)	32.2 (1.2)	2.7 (0.4)
	Mar. 81		194.1 (14.5)	10.3 (2.6)	34.7 (2.6)	1.8 (0.5)
MK area = 262.6 sq. m.	Mar. 80		115.6 (8.9) 41 removed	55 intro	44.0 (3.4) 28.4 (3.4) after removal	20.9 intro
	Oct. 80		116.4 (5.4)	15.5 (1.7)	44.3 (2.0)	5.9 (0.6)
	Mar. 81		106.0 (5.7)	8.2 (0.6)	40.4 (2.2)	3.1 (0.2)

Note: Standard error of estimate given in parentheses. Density is in units of number of animals per 100 sq. m.

increase to recruitment from outside the site, as indicated by the large body sizes of the new residents. Of the 55 *A. wattsi* that had been introduced, only 16 remained. This initial survivorship is twice that observed in the first experiment in which no residents had been removed, as discussed further below. Furthermore, some of the territories of *A. wattsi* at MK were located in the center of the site where they had been released and where residents had been nosed-out. Finally, we observed one hatchling of *A. wattsi* at this time. By March 1981 the number of *A. gingivinus* had dropped to about 106 and of *A. wattsi* to about eight.

We know that the *A. wattsi* that we have introduced at each of these sites have not merely migrated out of the sites to set up territories elsewhere because we have searched extensively for this possibility.

The principal difference between the introductions at MK, where the resident *A. gingivinus* were partially removed, and SK where the residents were undisturbed, is that the initial survivorship of the introduced *A. wattsi* is higher in MK than SK. But the survivorship of *A. wattsi* within MK for subsequent periods is about the same or possibly even lower than in SK. By the end of the first period the *A. gingivinus* density within MK had returned to its original value due to re-

cruitment from outside the site. Here in the subsequent periods the MK site no longer serves as an experiment to be contrasted with SK but instead as a replicate of SK for the assay of the longevity of *A. wattsi* in the environment provided by the cay.

The length and season of the initial periods for MK and SK are different. The monthly survival probability, l , can be calculated based on the following formula,

$$N(\Delta t) = l^{\Delta t} N_0$$

where $N(\Delta t)$ is the number remaining after Δt months, N_0 is the number introduced, Δt is the number of months between census and l is the monthly survival probability. Upon rearranging we have

$$l = \frac{N(\Delta t)}{N_0}^{\frac{1}{\Delta t}}$$

The number of months for the SK introduction is eight and for the MK introduction is seven yielding

$$\begin{aligned} l_{SK} &= 0.782 \\ l_{MK} &= 0.838. \end{aligned}$$

These numbers indicate that after a standard length of time, say six months, an introduction with the SK monthly survival would have 23% survivors and with the MK value would have a 35% survival, or almost twice as much.

We suspect the mortality differences between SK and MK mostly occur within the first month or two of the introduction and, if so, the values calculated based on the entire census interval underestimate the difference between the sites. Also, the introduction to MK occurred during the dry season, which Stamps and Tanaka (1981) have shown is generally the period of greatest environmental stress for *Anolis* lizards. It is possible that the MK introduction would have succeeded even better if it had been made in the wet season, as was the introduction to SK.

We are repeating these experiments, and further results will be reported as they become available.

LONG TERM CENSUS ON ST. MAARTEN

Table 2 presents census results from three sites on St. Maarten begun in 1977. These results provide the beginning of a longitudinal study of the population sizes of lizards on a tropical island. One site, Naked Boy Mountain, on the windward side of the island, contains members of *A. gingivinus* only. This site is xeric and the vegetation is scrubby. Maximum height of the trees here is about 5 m.

Another site, called Boundary, is on the leeward side of the island, near the species border of *A. watsi*. It is a xeric site, but not as xeric as Naked Boy Mountain above. The maximum tree height is approximately 6 m and cacti are abundant. From July 1977 through March 1980 it contained only *A. gingivinus*, but in November 1980 some *A. watsi* were found there as well.

The third site is located at the highest

TABLE 2. CENSUS DATA—ST. MAARTEN.

Site	Date		number <i>A. gingivinus</i>	number <i>A. watsi</i>	density <i>A. gingivinus</i>	density <i>A. watsi</i>
Naked	July	77	93.2 (7.3)	-0-	43.2 (3.4)	-0-
Boy	July	78	107.9 (11.6)	-0-	50.1 (5.4)	-0-
Mt.	July	79	69.6 (9.0)	-0-	32.3 (4.2)	-0-
area =	Mar.	80	193.6 (14.3)	-0-	89.8 (6.6)	-0-
215.6	Nov.	80	167.0 (6.7)	-0-	77.5 (3.1)	-0-
sq. m.	Mar.	81	98.8 (7.6)	-0-	45.8 (3.5)	-0-
Bound-	July	77	100.4 (4.4)	-0-	50.1 (2.2)	-0-
ary	July	78	130.3 (6.0)	-0-	65.0 (3.0)	-0-
area =	July	79	130.0 (5.7)	-0-	64.8 (2.8)	-0-
200.6	Mar.	80	260.4 (11.1)	-0-	129.8 (5.5)	-0-
sq. m.	Nov.	80	191.4 (6.0)	8.8 (3.7)	95.4 (3.0)	4.4 (1.8)
	Mar.	81	144.7 (6.6)	4.0 (2.0)	72.1 (3.3)	2.0 (1.0)
Pic du	July	77	33.3 (10.1)	41.4 (2.6)	8.4 (2.6)	10.5 (0.7)
Paradis	July	78	56.1 (21.7)	98.7 (12.9)	14.2 (5.5)	24.9 (3.3)
area =	July	79	29.9 (24.5)	83.8 (8.8)	7.6 (6.2)	21.2 (2.2)
396.0	Mar.	80	57.9 (12.3)	174.0 (20.8)	14.6 (3.1)	43.9 (5.2)
sq. m.	Nov.	80	157.4 (18.6)	225.0 (10.8)	39.8 (4.7)	56.8 (2.7)
	Mar.	81	52.8 (11.5)	156.1 (9.7)	13.3 (2.9)	39.4 (2.5)

Note: Standard error of estimate given in parentheses. Density is in units of number of animals per 100 sq. m.

point on St. Maarten, on Pic du Paradis (424 m). This site contains both species and is in the center of the distribution of *A. wattsi*. Pic du Paradis is the most mesic of our sites, and the vegetation consists of woods whose maximum height is about 8 m.

All three sites showed an increase in population size in 1980 for both species. This broad trend appears to be correlated with a dry season that was unusually wet in 1979. In 1980 many more juveniles were observed than in previous years. One of the most interesting points about these data is that they reveal greater fluctuations in population sizes on St. Maarten than on the cay, Anguillita.

DISCUSSION

The results of our study so far establish that *A. wattsi* can survive for over a year and can produce offspring in habitats equivalent to the xeric, scrubby habitat that they do not occupy on St. Maarten. Furthermore, the results show that the initial survivorship of *A. wattsi* upon introduction into this habitat is higher when resident *A. gingivinus* are partially removed prior to the introduction. These findings support the hypothesis that the absence of *A. wattsi* from sea level locations around St. Maarten is caused by competitive exclusion by *A. gingivinus* in that habitat.

Our observations suggest that there is a great site fidelity among introduced animals. T. Schoener (personal communication) has reported a similar high site fidelity in experimental introductions of anoles to small cays in the Bahamas, provided that the cay already contains a resident anole. The observation of high site fidelity also agrees with anecdotal observations on accidental introductions throughout the Caribbean. For example, a population of *A. porcatius* at the site of a former World Trades Fair in Santa Domingo in the Dominican Republic has remained restricted to approximately three city blocks over many

years. Similarly, a population of *A. cristatellus* has remained in a very small area in the town of La Romana in the Dominican Republic since the 1930's. These propagules simply do not spread in the presence of a resident. Still other examples have been compiled by Williams (personal communication). We term a propagule that is not spreading in this sense an "enclave."

We are fortunate to be able to report on the effect of a large hurricane on a population of *A. gingivinus* on the Anguillita cay. Our census in August 1979 preceded hurricane David and another large hurricane (Frederick) that came two weeks later. Hurricane David is the hurricane that demolished much of Dominica. Hurricane Frederick came very close to the Anguilla Bank. Surprisingly enough, this hurricane produced little structural damage to the cay. The hurricane threw chunks of coral up onto the cay, but caused no apparent damage to the trees. Our census of March 1980 reveals that there was essentially no change in the population size of *A. gingivinus* in our study site. This observation suggests that, at least on small islands, there is little mortality to *Anolis* lizards caused by hurricanes, unless there is severe structural damage to the vegetation in which they live.

Finally, we conjecture on the mechanisms that cause the apparent competitive exclusion of *A. wattsi* by *A. gingivinus* from xeric sea level habitats. The phenomenon of competitive exclusion involves two considerations. The first pertains to the disadvantage in carrying capacity (K) that one species has relative to another. The other pertains to the existence of a competition coefficient (α) between the two species. These distinct considerations arise upon inspecting the condition for one species to invade a habitat in the presence of another species according to the Lotka-Volterra competition equations. Here, we are interested in whether *A. wattsi* can invade low-land habitat in the presence of *A. gingivinus*.

For *A. wattsi* to be prevented from invading the habitat, it is necessary that:

$$\alpha_{w,g} > \frac{K_w}{K_g},$$

or in words, that the competition coefficient of *A. gingivinus* against *A. wattsi* exceed the ratio of the carrying capacity of the habitat for *A. wattsi* relative to the carrying capacity of *A. gingivinus*. Obviously, for *A. wattsi* to be prevented from invading the lowland habitat in the presence of *A. gingivinus*, both the competition coefficient $\alpha_{w,g}$ should be of moderate magnitude and the ratio of the carrying capacity for *A. wattsi* relative to *A. gingivinus* should be low.

We speculate that the principal mechanism contributing to a carrying capacity disadvantage to *A. wattsi* in a xeric habitat relative to *A. gingivinus* is that *A. wattsi* is not as physiologically well adapted to xeric conditions as is *A. gingivinus*. Preliminary data obtained at Stanford in collaboration with Hal Mooney and at the University of Wisconsin in collaboration with Warren Porter show that *A. wattsi* has a higher rate of water loss per unit of surface area than *A. gingivinus*. Also, in Figure 2 we show that *A. wattsi* on St. Maarten has a lower panting temperature than *A. gingivinus*. This result parallels our earlier finding that *A. wattsi* on St. Eustatius has a lower panting temperature than *A. bimaculatus* on that island (Roughgarden *et al.*, 1982). *A. bimaculatus* is closely related to *A. gingivinus*, and this lineage generally appears to have more xeric tolerance than the *A. wattsi* lineage. As a consequence, we suspect that *A. wattsi* may find less of the microhabitat suitable as territories and may have a more narrow window in its time budget during which it can forage than *A. gingivinus*. Both factors would translate into lowering the carrying capacity ratio of *A. wattsi* relative to *A. gingivinus* in xeric habitat.

We should add that panting temperature data notwithstanding, individuals of *A. wattsi* and *A. gingivinus* typically have the same body temperature in the field at sites where they co-occur, and there is no evidence that *A. gingivinus* prefers hotter microsites than *A. wattsi* under field conditions.

There are three mechanisms that cause a competitive effect of individuals of *A. gingivinus* against individuals of *A. wattsi* (i.e., that lead to a high $\alpha_{w,g}$). First, members of *A. gingivinus* use insect prey that *A. wattsi* would otherwise eat.

Second, *A. wattsi* and *A. gingivinus* are interspecifically territorial. The space used for territories is the most obvious resource for which these species are competing. The mechanism of competition for this space is interspecific territorial behavior, including both aggression and defense.

Third, interspecific predation may be important as well. Large male *A. gingivinus* have been observed capturing and eating adult as well as juvenile *A. wattsi*. Similarly, *A. wattsi* are capable of capturing and eating at least juvenile *A. gingivinus*. Analysis of stomach contents and direct observation establish that there is both inter- and intra-specific predation among *Anolis* lizards, but the magnitude of the effect of this form of interference competition on population dynamics has

Figure 2. Distribution of panting temperatures for lizards from St. Maarten.

not yet been determined. It is possible, for example, that the readily observed association of *A. watsi* with rock piles is not an adaption whose function is to provide a humid and cool microclimate, but is a trait whose function is to provide an escape from predation by *A. gingivinus* as well as other predators.

In xeric habitat the magnitude of these competitive interactions of *A. gingivinus* against *A. watsi* probably increases. We may summarize these speculations by imagining a transect from the somewhat mesic habitat of the hills on St. Maarten to the xeric habitat of sea level. Along the transect the competition coefficient of *A. gingivinus* against *A. watsi* rises while the carrying capacity ratio of *A. watsi* to *A. gingivinus* falls. The species border of *A. watsi* lies approximately at the point where the competition coefficient equals the carrying capacity ratio. Below this point only *A. gingivinus* occurs.

Over the years Ernest E. Williams has introduced many to the *Anolis* system. He championed the view that this system might someday prove as important to evolution, ecology, and biogeography as *Drosophila* has in genetics. Although it is premature to say with certainty, there is a real possibility that this vision will be realized. Our preliminary efforts with field experiments using *Anolis* populations have provided us with great optimism that *Anolis* will play a leading role as experimental terrestrial community ecology develops during the coming years.

ACKNOWLEDGMENTS

We thank James Bryan of Bryansville, West End, Anguilla for the permission to

conduct these studies on Anguillita and the Fleming family of St. Maarten for their permission to census the population on their property. We also thank Carl and Vikki Leisegang of the VIKKI TOO for their seamanship and hospitality and Stephen Adolph for his able assistance during all phases of the field work. We thank Paul Hertz and an anonymous reviewer for exceptionally helpful comments on the manuscript. We gratefully acknowledge the support of the National Science Foundation.

LITERATURE CITED

- HECKEL, D., AND J. ROUGHGARDEN. 1979. A technique for estimating the size of lizard populations. *Ecology*, **60**: 966-975.
- HERTZ, P. 1981. Compensation for altitudinal changes in the thermal environment by some *Anolis* lizards on Hispaniola. *Ecology*, **62**: 515-521.
- LAZELL, J. 1972. The Anoles (Sauria, Iguanidae) of the Lesser Antilles. *Bull. Mus. Comp. Zool.*, **143**: 1-115.
- ROUGHGARDEN, J., D. HECKEL, AND E. FUENTES. 1982. How coevolutionary theory explains the biogeography and community structure in *Anolis* lizard communities in the Lesser Antilles. To be published in the proceedings of a symposium on the Ecology of Lizards, held at the ASZ meeting in Seattle, December 1980; to be edited by R. Huey, E. Pianka, and T. Schoener.
- SCHOENER, T. 1969. Size patterns in West Indian *Anolis* lizards I. Size and species diversity. *Syst. Zool.*, **18**: 386-401.
- STAMPS, J., AND S. TANAKA. 1981. The influence of food and water on growth rates in a tropical lizard (*Anolis aeneus*). *Ecology*, **62**: 33-40.
- WILLIAMS, E. E. 1972. Origin of faunas. Evolution of lizard congeners in a complex island fauna—a trial analysis. *Evol. Biol.*, **6**: 47-89.

Competition Between *Anolis* and Birds: A Reassessment

TIMOTHY C. MOERMOND¹

ABSTRACT. Birds and anoles show complementary distributions in the West Indies and Central America. The possible effect of these different taxonomic groups on each other is assessed in terms of competition and predation. Birds which are likely competitors and those which are *Anolis* predators are poorly represented in the West Indies. The principal likely competitors of *Anolis* in the West Indies are migratory parulid warblers. The warblers primarily use upper levels of vegetation while anoles predominantly use the lower levels near the ground. Within the lower levels lizards may exist at high population levels and scan many surfaces continuously, possibly depressing arthropod abundance and/or activity. Their hunting patterns may exclude insectivorous birds from lower portions of habitats when lizard densities are high. The ability of lizards to outcompete birds in particular portions of habitat is based on the mechanistic view of habitat use derived from the concept of the structural niche. *Anolis* populations are also sensitive to predation by birds, with the reduced populations in the Central American mainland apparently due to the high level of avian predation, not to competition.

INTRODUCTION

Anolis lizards are a conspicuous part of the fauna of the West Indies both in species richness and abundance (Williams, 1969, 1976; Schoener, 1969) and may be the dominant arboreal insectivores. In contrast, on the Central and South American mainland, *Anolis* lizards are much less abundant (Rand and Humphrey, 1968; Andrews, 1979). Conversely, birds are far more diverse on the

mainland and notably depauperate in the West Indies (Terborgh, 1973; Cox and Ricklefs, 1977; Terborgh and Faaborg, 1980b).

The possibility of competitive interactions between anoles and foliage-gleaning birds is compelling and has been suggested by a number of workers (Schoener, 1969, 1970, 1972; Schoener and Schoener, 1978b; Moermond, 1974, 1976; Morse, 1975; Lister, 1976; Andrews, 1979; Wright, 1979, 1981). Competition is suggested by the many parallels between communities of *Anolis* lizards and of birds, particularly of the small foliage-gleaners: both include generalized insectivores (Schoener and Gorman, 1968; Hespenheide, 1975; Andrews, 1979; Wright, 1979), both are highly arboreal, visual, diurnal foragers and both are equipped with color vision (Forbes *et al.*, 1964; Sexton, 1964; Sillman, 1973).

The demonstration of competition requires evidence 1) that both organisms are using the same resources and 2) that this resource use results in adverse effects to one or both organisms (cf. Schoener, 1972; Pianka, 1978). In the case of anoles and small insectivorous birds, we can infer use of similar resources by similarity in size and taxa of invertebrate prey (e.g., Wright, 1979) or, less certainly, by similarity of hunting locations (i.e., prey taken from same substrates) by animals with similar-sized trophic appendages (i.e., mouth and bill lengths). The latter criterion is probably justified since it is likely that most anoles

¹Department of Zoology, University of Wisconsin, Madison, Wisconsin 53706, U.S.A.

and small avian insectivores are relatively catholic in taxa choice within similar size ranges (cf. Schoener and Gorman, 1968; Hespeneide, 1975; Andrews, 1979).

Adverse effects of one organism on another due to competition are difficult to document. The usual approach is to look for complementary distribution, i.e. distributions in which the density of one organism is low where the other is high (e.g., Kiester, 1971). Complementarity between birds and lizards has been demonstrated within several island groups. For example, Wright (1979) has shown higher lizard densities and lower bird densities on small than on large islands in Gatun Lake, Panama. Lister (1976) refers to another example in the northern Bahamas where Grand Bahama and Abaco, which are unusually rich in small insectivorous birds, support only one species of *Anolis* in contrast to equivalently sized islands farther south. Schoener and Schoener (1978b) carefully document an inverse relationship in bird species richness and *Anolis* density between Andros and Bimini in the Bahamas. Wright (1981) has also shown for a selected series of West Indian islands that smaller islands have higher densities of *Anolis* lizards but lower densities of birds than do larger islands.

The difficulty in using such complementary patterns is in determining the causal factors. Are the patterns primarily results of competition or is predation involved? If the densities of each group are affected by the presence of the other, what factors control the pattern? The concept of the structural niche of *Anolis* developed by Rand and Williams (Rand, 1964; Rand and Williams, 1969; Williams, 1972) offers a mechanistic approach with which to analyze the patterns of habitat use and hence resource use by *Anolis* lizards (see Moermond, 1974, 1979a,b). I shall show here that the same concept can be usefully employed to analyze the interactions between lizards and birds and that such an appli-

cation leads to a more precise definition of the problem. This in turn allows one to distinguish among competing hypotheses.

HYPOTHESIS: COMPETITION OR PREDATION?

The following hypotheses have been offered to account for complementary distributions of lizards and birds in mainland and West Indian habitats:

1) Schoener (1972) suggested that, "...competition with endotherms, the metabolic and therefore home-range requirements of which are too great to allow them to exist on small islands, could reduce the populations of lizards in tropical mainlands."

2) Wright (1979, 1981) argued that lizards opportunistically harvest resources which are not harvested by birds. Higher extinction rates for birds on islands may leave larger portions of resources available for anoles.

3) Terborgh (quoted in Lister, 1976) suggested that *Anolis* lizards, by virtue of their high abundance, tight packing, and low extinction rates, out-compete birds on these islands (i.e., the West Indies).

4) Andrews (1979) suggested that, "...island anoles, with few predators, may be food limited relative to heavily preyed-upon mainland species."

These statements, although not mutually exclusive clearly do not agree on the outcome of competition between lizards and birds. The foregoing suggestions can be combined as follows to form two separate hypotheses:

1) At high species densities birds are capable of out-competing lizards, but there are fewer species of birds in the West Indies due to higher extinction rates of birds over anoles. This presumably allows higher densities of lizards on these islands.

2) At high numerical densities, lizards are capable of out-competing birds, but lizards are sensitive to predation by birds. Thus the higher densities of West

Indian anoles are presumably due to much lower numbers of avian predators compared to the mainland.

The question is not only whether the birds are responsible for the low densities of mainland anoles (Andrews, 1979; Wright, 1979; see also Schoener and Schoener, 1978b) or whether the anoles are responsible for the paucity of bird species in the West Indies (Moermond, 1974; Terborgh and Faaborg, 1980b) but whether the birds are implicated as competitors or as predators. The answer will be sought through a series of comparisons.

MATERIALS AND METHODS

The observations and data presented here were gathered on many trips to the West Indies and Costa Rica during which I have observed over 50 *Anolis* species. Most emphasis here is placed on data collected in Haiti (Nov. 1971 and July-Aug. 1972) and the Dominican Republic (March 1979). Data on lizards were derived mainly from detailed chronologies of the behavior of individuals in the field (described in Moermond, 1974, 1979a). Data on birds were distilled from daily lists and observations on foraging behavior taken at the West Indies sites concurrently with *Anolis* studies as well as from extensive observations of understory birds at the La Selva Biological Station in Costa Rica (Aug.-Dec. 1979, June-Aug. 1980, and Dec. 1980-Jan. 1981).

AVIAN PREDATORS AND COMPETITORS

A comparison of mainland and West Indian habitats shows that, although total bird densities are not appreciably lower in the West Indies, there is a striking decrease in total species diversity and in the population densities of certain foraging types—potential lizard predators, bark-gleaning insectivores and foliage-gleaning insectivores (Cox and

Ricklefs, 1977; Terborgh and Faaborg, 1980b). (Islands such as Trinidad with geologically recent connections to South America are included with the mainland.) If one includes migrants, the totals for species and foliage-gleaning insectivores on the Greater Antilles increase substantially but still remain below corresponding values for mainland habitat (Terborgh and Faaborg, 1980a).

Numerous authors have noted that several families which contribute substantially to the three foraging types listed above are entirely absent from the West Indies (Bond, 1974): Momotidae, Bucconidae, Dendrocolaptidae, Furnariidae, and Formicariidae. Geographical distributions for these species suggest that the members of all five families are poor over-water colonists (MacArthur, *et al.*, 1972; Terborgh, 1973); their absence from the West Indies may be independent of the presence of high lizard densities. At a montane site in southern Haiti (Les Platons) I recorded 38 resident species of birds and 11 migrants (Moermond, 1974). Of the resident species, only six appear as potential competitors of *Anolis*: *Icterus dominicensis*, *Vireo altiloquus*, *Coereba flaveola*, *Anthracothonax dominicus*, *Todus subulatus*, and *T. angustirostris*. Only the first two are usually considered as foliage-gleaners. Closer examination suggests that competition between these two species and the anoles is probably not intense. The *Icterus* is much larger than the anoles, and both *Icterus* and *Vireo* may eat an appreciable amount of fruit. The diet of *V. altiloquus* in Puerto Rico was reported by Wetmore as 57% fruit (Wetmore, 1916). Both *Coereba* and *Anthracothonax* are heavily nectarivorous, and even though both take small insects around flowers, neither is likely to be a substantial competitor of *Anolis*. The two todies take most of their prey on the wing by sallying and hovering, but the majority of these are taken from the surface of foliage at the forest edge (Kepler, 1977; Moermond, unpublished observation).

The todies are thus potential competitors for these insects, but they are also occasionally predators on small juvenile *Anolis* (Wetmore, 1916). Of the six species, only the nectarivores were seen daily at the Les Platons site.

In contrast, the 11 migrant parulids I recorded at Les Platons are all potential competitors of *Anolis*. Six species were seen daily, and three were abundant. Likewise in the area around Constanza, Dominican Republic, I recorded 10 species of parulids in two weeks in 1979; of these, four were abundant.

This predominance of migrant parulid warblers among the foliage-gleaning insectivores in the West Indies is well documented (Lack and Lack, 1972; Lack 1976; Terborgh and Faaborg, 1980a; Keast, 1980). These migrants may remain on their wintering grounds in the West Indies for upwards of seven months (Keast, 1980) and thus are potentially important competitors. Pianka (1975) has stressed the possible influence of migrant birds in Africa in depressing the Kalahari lizard fauna.

Distinction of birds as anole competitors from birds as anole predators is confounded by the dual role of many bird species. In addition to large birds of several families, e.g., Momotidae, Bucconidae, and Cuculidae, that contain known or potential predators on anoles, many smaller species of birds are known to prey occasionally on juvenile anoles. For example, in the rainforest at La Selva field station in Costa Rica, at least half of nearly two dozen understory insectivores are known or likely opportunistic predators on juvenile anoles (personal observation). They include the antbird *Thamnophilus punctatus*, which was one of the principal *Anolis* competitors considered by Wright (1979). Clearly many of these species are likely to have dual roles in their interactions with *Anolis*. It is important to emphasize that the major sources of insectivore competitors with *Anolis* in the West Indies, the parulid warblers, are *not* likely to be significant predators on anoles due to their small size (Table 1).

ARTHROPOD DENSITIES AND FOOD LIMITATION

Wright (1979) argued that birds, as competitors, lower the abundance of arthropod prey used by anoles. Sampling a series of small islands in Gatun Lake, Panama, he demonstrated a negative correlation between density of insectivorous birds (mist net samples) and arthropod abundance (sweep net samples) and a negative correlation between bird and lizard densities. Note that the animal communities on the small islands Wright sampled are reduced subsets of the adjacent mainland communities and are not analogous to the West Indian-mainland comparison. In the latter comparison the correlation between bird and arthropod abundance is *opposite* that found by Wright. Three separate studies of insect abundance at mainland sites in Costa Rica and at sites on Puerto Rico and Dominica have shown a significantly lower abundance of arthropods on West Indian islands (Allan *et al.*, 1973; Janzen, 1973; Andrews, 1979) and significantly fewer large insects in West Indian samples (Andrews, 1979).

With respect to insectivores Morse (1975) reports that Odum *et al.* (1970) found the biomass of reptiles and amphibians in a Puerto Rican forest to be nearly 30 times that of carnivorous birds (including insectivores). Despite the higher metabolic rates of the birds, Morse suggests from preliminary calculations of Lavigne [no reference given] that the birds may account for only 1 to 2% of the insects consumed by both groups.

Andrews's (1976, 1979) experimental studies clearly suggest that island anoles are food limited but that mainland anoles are far less so. Andrews has shown that mainland anoles, when compared with island species, show faster growth rates, earlier maturation, lower adult survivorship, smaller adult size and lower population densities. Andrews (1979) suggests that the most likely cause for this difference is a higher predation rate on the mainland anoles. Experiments with food

TABLE 1. OVERLAP IN SIZE OF TROPHIC STRUCTURE BETWEEN ANOLIS LIZARDS AND FOLIAGE-GLEANING BIRDS AT TWO SITES IN HISPANIOLA.

Trophic size in birds is given as length of exposed culmen. Data from Cruz, 1980; Lack, 1976; and Wetmore and Swales, 1931. Trophic size in *Anolis* is given as head length from the anterior edge of the ear to the tip of the snout. Data from Schoener, 1969. Scale is nonlinear to accommodate the large number of entries at small sizes.

LES PLATONS, HAITI		CONSTANZA, DOMINICAN REPUBLIC	
Bird	Lizard	Bird	
Bill length (mm)	Head Length (mm)	Bill length (mm)	
22	22	22	
<i>Icterus dominicensis</i>			
21 ♂ <i>Todus subulatus</i>	<i>A. cybotes</i> ♂	21 <i>A. cybotes</i> ♂	<i>Todus subulatus</i> ♂
♀ <i>Todus subulatus</i>			<i>Todus subulatus</i> ♀
			<i>Phaenicophilus palmarum</i>
20	<i>A. coelestinus</i> ♂	20 <i>A. chlorocyanus</i> ♂	20
19		19	19
18 <i>Vireo altiloquus</i>		18	18
17		17	17
16	<i>A. hendersoni</i> ♂	16 <i>A. aliniger</i> ♂	16
	<i>A. coelestinus</i> ♀	<i>A. chlorocyanus</i>	
15		15	15
	<i>A. cybotes</i> ♀	<i>A. cybotes</i> ♀	
14		14	14
	<i>A. distichus</i> ♂	<i>A. christophei</i> ♂	
		<i>A. distichus</i> ♂	
13	<i>A. monticola</i> ♂	13	13
		<i>A. cochranæ</i> ♂	<i>Seiurus novaeborealis</i>
	<i>A. semilineatus</i> ♂		
12	<i>A. hendersoni</i> ♀	12	12
<i>Seiurus aurocapillus</i>		<i>A. christophei</i> ♀	<i>Seiurus aurocapillus</i>
<i>Mniotilta varia</i>		<i>A. distichus</i> ♂	<i>Mniotilta varia</i>
		<i>A. etheridgei</i> ♂	
	<i>A. distichus</i> ♂		
11	<i>A. koopmani</i> ♂	11	
<i>Dendroica petechia</i>	<i>A. monticola</i> ♀		
<i>Geothlypis trichis</i>			
10 <i>Dendroica coronata</i>	<i>A. semilineatus</i> ♀	10 <i>A. cochranæ</i> ♀	<i>Dendroica coronata</i>
<i>Parula americana</i>		<i>A. etheridgei</i> ♀	<i>Dendroica tigrina</i>
<i>Dendroica palmarum</i>			<i>Dendroica caerulescens</i>
<i>Dendroica tigrina</i>			
<i>Dendroica caerulescens</i>			
9 <i>Dendroica discolor</i>	<i>A. koopmani</i> ♀	9	9
<i>Setophaga ruticilla</i>			<i>Setophaga ruticilla</i>
8 <i>Coereba flaveola</i>		8	8
			<i>Coereba flaveola</i>

supplementation in enclosures showed no significant increase in growth rates for *A. limifrons* on the mainland but a significant increase in growth for an island species, bringing it into line with the rate for *limifrons* (Andrews, 1976; see also Licht, 1974). Her data clearly suggest that island anoles are food limited while mainland forest anoles are far less so (see also Stamps, 1977; Schoener and Schoener, 1978a).

COMPARISON OF EXPLOITATION PATTERNS

OVERLAP IN SIZES OF TROPHIC STRUCTURES

We can roughly evaluate the overlap in prey size of birds and anoles foraging in similar habitats by examining the relative sizes of their trophic structures. Bill size in birds and head size in *Anolis* have

been shown to correlate well with size of prey items taken (Schoener, 1965, 1967, 1968; Schoener and Gorman, 1968; Hespeneheide, 1975; Howe, *et al.*, in preparation). The bill lengths for insectivorous gleaners and *Todus* species for the two Hispaniolan sites mentioned above are compared to head lengths of the *Anolis* species present at each site in Table 1. Because the head length measure of lizards overestimates their mouth length by a couple of millimeters (varying from species to species), I would expect an *Anolis* with a head length equal to the bill length of a bird to take somewhat smaller prey on average than the bird. Therefore to use Table 1 to compare average prey size taken by lizards and birds, the lizard columns should be shifted downward by a couple of millimeters relative to the birds. The sizes presented for the lizards are for large adults of each sex. All the species presented hatch out as active insectivores but with body and head sizes below that listed for the adults of *A. koopmani*, the smallest species in Table 1. Therefore the ontogenetic series for each species listed in Table 1 overlaps the trophic sizes of all smaller species. In contrast birds fledge and begin foraging with adult-sized bills. Given these two considerations the parulids clearly fall well within the range of trophic sizes encompassed by the lizards.

FORAGING HEIGHT OVERLAP

Within-habitat comparisons of foraging locations reveal important differences between *Anolis* lizards and small foliage-gleaning birds in the West Indies. There is little overlap between the foraging heights of parulid warblers and those of *Anolis* lizards in forest habitats. Data on within-habitat distribution of *Anolis* on Jamaica (Schoener and Schoener, 1971a) and Puerto Rico (Schoener and Schoener, 1971b) (Table 2) and on Hispaniola (Fig. 1) show that the anoles are active pre-

dominantly within 3 m of the ground. In contrast, parulids and other insectivorous foliage-gleaners forage predominantly above 3 m (Lack and Lack, 1972; Moermond, 1974; Cruz, 1980; Terborgh and Faaborg, 1980a; Bennett, 1980). The separation is greater if foraging height distributions are examined in more detail. Perch height distributions for seven Haitian *Anolis* (Fig. 1) show strongly skewed distributions with the mode usually within a few centimeters of the ground. The modes of similar distributions for the birds are usually above three meters, and there is a sharp decline in observations within a meter of the ground (see Diamond, 1973; Cruz, 1980; references in Keast and Morton, 1980).

The height overlap distributions suggest different zones of activity for the two types of organisms. Although it is tempting to interpret these apparently reciprocal distributions as niche shifts, this supposition is difficult to test. Comparisons of foraging height distributions in the West Indies, in mainland wintering grounds, and in breeding and migrating areas over a range of habitats only emphasize the variability of such distributions. As expected, the height of foraging activity is strongly influenced by vegetation structure and, obviously, but less easily documented, by local distribution of available prey (Parnell, 1969; Morse, 1979; and references in Keast and Morton, 1980). Nevertheless, almost all recorded parulid foraging distributions cited in Keast and Morton (1980) show modes at middle or upper ranges of the vegetation, nearly always above three meters. Although *Todus* species may feed below 3 m (Kepler, 1977), these birds are primarily taking prey on the wing from leaf tips.

In contrast, at La Selva, a Costa Rican lowland rainforest, there are at least a dozen species of ground or foliage-gleaning insectivorous birds which, I estimate, spend more than half their foraging time below three meters. A similar pattern is suggested by a careful

TABLE 2. FREQUENCY OF ANOLIS LIZARDS ABOVE TEN FEET (~3 M) ON TWO WEST INDIAN ISLANDS.

Site	No. species in census	No. individuals in census	Per cent above 10 feet (~3 m)	Source
Jamaica				
Mona	4	2,458	7	Schoener & Schoener, 1971a
Port Antonio open	4	1,097	5	Schoener & Schoener, 1971a
Port Antonio closed	3	749	4	Schoener & Schoener, 1971a
Whitehouse	4	1,656	1	Schoener & Schoener, 1971a
Puerto Rico				
Maricao, 20 road edge	3	199	20	Schoener & Schoener, 1971b
Maricao, open forest	3	317	9	Schoener & Schoener, 1971b
Maricao, with cleared understory	3	70	6	Schoener & Schoener, 1971b
Maricao, forest intermediate	3	371	7	Schoener & Schoener, 1971b
El Verde	3	394	4	Schoener & Schoener, 1971b

Figure 1. Foraging height distributions for seven species of *Anolis* at Les Platons, Haiti (after Moermond, 1979a). N = the total number of perch heights recorded for each species. © the Ecological Society of America.

survey of data presented by Karr (1971) and Orians (1969). Pearson (1977) also observed species of foliage-gleaning antbirds feeding predominantly below 3 meters in rainforests in South America. The primary candidates for *Anolis* competitors listed in these mainland studies are suboscines (especially antbirds) and

wrens which are largely absent from the West Indies.

INFLUENCE OF HABITAT STRUCTURE

The differences in foraging heights between parulids and *Anolis* suggest that the habitat constrains the two groups dif-

ferently. For our purpose the forest habitat can be divided into two zones: canopy and understory. Canopy includes the upper and outer branches, twigs and leaves of trees. The understory includes inner limbs and boles of trees as well as shrubs and ground cover. It is not surprising to find that foraging in these different zones requires different adaptations. Below I will briefly discuss the types of constraints that apply to each group of foragers.

FOLIAGE-GLEANING BIRDS

The ability to fly plus the sizes and hence maneuverability of small foliage-gleaners would seem to make all height levels readily exploitable. Although numerous studies show height separations among closely-related birds (Lack, 1971; Lack and Lack, 1972; Cruz, 1980), species often expand their height range of habitat use on islands where total species diversity is lower (MacArthur *et al.*, 1966; MacArthur *et al.*, 1972; Terborgh *et al.*, 1978; Terborgh and Faaborg, 1980a). Nevertheless, studies of niche expansion among colonizing birds on islands (Keast, 1970; Diamond, 1970; Ricklefs and Cox, 1972, 1977; Terborgh and Faaborg, 1973; Terborgh *et al.*, 1978) suggest that not all portions of the habitat are equally available.

Small differences in habitat use near ground levels are sometimes associated with changes in wing shape and especially foot and leg structure (Dilger, 1956; Pearson, 1977; Fitzpatrick, 1978; Moermond, *et al.*, in preparation). The most specialized morphological adaptations appear to be associated with extensive use of trunks or ground (Bock and Miller, 1959; Osterhaus, 1962; Diamond, 1970; Feduccia, 1973). Thus, while foliage-gleaners may readily use much of the canopy zone on islands, they do not appear to readily expand to the trunk or ground in the West Indies even in the absence of the usual specialists on these substrates.

One might, however, expect foliage-gleaners to expand into the understory shrubs. Nevertheless, those warblers, such as *Dendroica palmarum* and *D. caerulescens*, which heavily use low shrubs, primarily do so in the open or at the edges of forest, not usually under the canopy. Here again the mainland groups using this zone, e.g., Formicariidae and Troglodytidae, are absent or rare in the West Indies.

ANOLIS LIZARDS

Although anoles are noted for their arboreal habits and can be found from ground to canopy, their foraging activity is heavily weighted towards the ground. My own detailed observations on the movements of over thirty species have shown that anoles are constrained to move on those surfaces which are either connected to each other or separated by no more than half a meter (Moermond, 1979a,b, and unpublished observations). Anoles use the habitat as if it were a "jungle-gym" with struts (=perches) of varying diameters and spacings. A number of studies have shown the importance of perch diameter to different species of anoles (Rand, 1964; Schoener, 1968; Moermond, 1979a). Any assemblage of four or more species is likely to include species with complementary adaptations such that perch diameters of 2 mm thick twigs to large trunks are used (see comments on ecomorphs in Rand and Williams, 1969; Williams, 1972; and Moermond, 1974).

Different spacings of the habitat "jungle-gyms" also favor different species types, but here we find that combinations of small perches and wide spacings appear difficult for any anole to use effectively (Moermond, 1979a). Such combinations occur predominantly in the canopy zone. Inter-perch distances in the canopies of most trees appear to be so great that most foraging movements are constrained to connected surfaces. Since these surfaces are also narrow, the area

which can be scanned at one time by a hunting lizard is small. As a result, there are frequent changes in scanning locations and probably low prey capture rates (Moermond, 1979b). Consequently anoles cannot easily use much of the canopy and observations within the canopy suggest that the few individuals foraging there usually penetrate the canopy only on the outer tips of large branches which act as "highways" back to other lower locations (personal observation).

Although there are a few species of anoles, such as *A. valencienni* of Jamaica and the giant anoles (Williams, 1962), specialized for movement in the crowns of trees, these species are generally rarer in any given habitat than other anole species (see Schoener and Schoener, 1971a). The "giant" forms may have diverged into different trophic niches. (Schoener [1970] has found that at least one, *A. equestris*, is frugivorous.) I have also observed a large male *A. leachi* on Barbuda picking and eating berries. Other more numerous "canopy" forms such as the trunk-crown or "green" anoles (Williams, 1965; Rand and Williams, 1969) have been observed foraging in canopy (Andrews, personal communication; Moermond, unpublished observation) but still heavily use foraging sites below three meters (see data in references cited in Table 2 and Fig. 1).

Clearly birds have a decided advantage in exploiting the canopy, although I doubt that the scarcity of anoles there is due to competitive pressure from birds. It is more likely due to the inherent difficulties of the nonflying anoles to forage in that zone. Lack of overlap of the two groups in the lower zone in the West Indies is more difficult to explain.

CONSEQUENCES OF HUNTING PATTERNS

In the lower parts of trees, in shrubs and on other surfaces near the ground, easy movement from one surface to another allows anoles (as well as birds) access to most prey. This fact, coupled

with the high lizard densities in these areas (Odum *et al.*, 1970; Schoener and Schoener, 1980; Andrews, 1979), suggests that many surfaces may be under virtually continuous surveillance for much of the day (e.g., Rand, 1967; Andrews, 1971; Fleming and Hooker, 1975; Moermond, 1979b and unpublished observations).

Because *Anolis* hunt by scanning the surfaces around them for prey, particularly moving prey (Curio, 1970; Moermond, 1981), a dense population of anoles may depress prey abundance and activity levels in lower habitat zones (cf. Schoener, 1969; Odum *et al.*, 1970; Schoener and Schoener, 1978b; see also Pough, 1980). This may constitute significant resource depression for active, visual foragers such as birds (cf. Charnov *et al.*, 1976). Schoener (1969) has also pointed out that, "*Anolis*, with its several age-size classes overlapping spatially, is more likely to uniformly reduce food than . . . birds."

Although both *Anolis* lizards and foliage-gleaning birds find prey principally by visually scanning branches, leaves or ground from the same or nearby surfaces, two differences are particularly noticeable: 1) the times spent scanning from each perch, and 2) the rates of prey capture. By either measure birds appear to be much more active foragers. Although few data on rates of movement or prey capture are available for foliage-gleaning birds in the West Indies (Moermond, 1974; Moermond and Denslow, in preparation), the data available suggest a rate of prey capture more than an order of magnitude greater than that reported for *Anolis* in the West Indies (Moermond, 1974; Andrews, 1976).

Several relevant aspects of both birds and anoles are summarized in Table 3. The comparison draws from temperate data on vireos (Williamson, 1971) because these data are quite complete and were taken in a similar fashion to those on parulids and because similar vireos occur among the resident foliage-

TABLE 3. A COMPARISON OF METABOLIC RATES, FEEDING RATES, AND RATE OF MOVEMENT OF ANOLIS LIZARDS WITH SELECTED FOLIAGE-CLEANING BIRDS.

	length of trophic struc. (mm)	weight (g)	cal./day	prey/hr.	moves/hr.
<i>Vireo</i> spp.	10 - 15 *	16 - 20 *	6500 - 9000 †	60 - 120 ‡	810 - 1380 ‡
<i>Parulid</i> spp.	7.5 - 14*	7 - 14 *	3600 - 6000 †	—	900 - 1800 **
<i>Anolis</i> spp. (40-70 mm) SVL)	10 - 17§	1.5 - 7.5	65 - 300 ¶	0.5 - 6 #	5 - 25 ††
RATIOS	<i>Vireo/Anolis</i> <i>Parulid/Anolis</i>	2.5 - 10 2 - 4.5	30 - 100 20 - 55	20 - 120 —	55 - 160 70 - 180

* Values from Roberts, 1955.

† Estimates from Lasiewski and Dawson, 1967, Kcal./day = 129 W^{.724}

‡ Calculated from data in Williamson, 1971.

§ Data from specimens listed in Appendix II of Moermond, 1974.

|| Estimated from weights for similar-sized lizards given in Jenssen, 1973.

¶ Estimated from Licht and Jones, 1967, 40-44 cal/gram body wt./day.

Values from Moermond, 1974.

** Moermond, unpublished data on wintering warblers at Les Platons, Haiti.

†† Values calculated from mean intervals between moves from data in Moermond, 1974.

gleaners in the West Indies (e.g., Cruz, 1980).

The correspondence between movement rate and prey capture rate is consistent for both groups; in general, an individual foliage-gleaning bird must find and capture prey at a much greater pace than an *Anolis* to survive. The obvious implication is that a population of anoles could successfully survive on a much lower prey base than their avian competitors (Pough, 1980) and may also contribute substantially to depressing that base below the minimum needed for the birds. If one is to look for significant competition between birds and lizards it is here.

The apparent advantage of *Anolis* in lower vegetation zones depends on 1) their manner of hunting which is well defined and characteristic of the genus, and 2) their population density. Factors, such as predation, which consistently leave more resources available for birds to exploit—a contrasting argument to that proposed by Wright (1979).

DISCUSSION

The hypotheses presented above can now be reassessed. I find little evidence that birds are capable of outcompeting lizards although the possibility cannot be ruled out. At least lizards are more likely to be constrained in their resource use by the habitat rather than by the species of birds present.

On the mainland, evidence that *Anolis* lizards are generally below carrying capacity is persuasive; predation, not competition from birds, appears to be the most likely cause (Andrews, 1979; Schoener and Schoener, 1978b). In the West Indies, the high densities of lizards and paucity of bird species are unlikely to be due solely to extinction rates of birds as argued by Wright (1981). Schoener (1976) has shown that the high per-species immigration rate for birds can compensate for their relatively small population sizes on islands. Lack of particular species and types of species of birds in the West Indies is likely due to a combination of poor colonization ability

and diffuse competition from other birds (Terborgh and Faaborg, 1980b) and possibly from *Anolis* (*op cit*, Moermond, 1974).

Instead, I suggest that where there are few predators, *Anolis* may be capable of excluding insectivorous birds from particular portions of the habitat via resource depression by the *Anolis* (Moermond, 1974, 1976). The paucity of predators on *Anolis* in the West Indies would not seem to be due to the *Anolis* since they are ubiquitous. Instead many predators on *Anolis*, such as the *Saunderothra* and *Coccyzus* cuckoos or *Tyrannus* flycatchers probably depend heavily on large insects which are relatively scarce in the West Indies (Andrews, 1979). These and probably all avian predators on *Anolis* take lizards opportunistically as a minor portion of a typically insectivorous diet. Therefore if anoles are partly responsible for the depression of insect densities in the West Indies, it is possible this "depression" affects their "predators" also. At least the presence of many species of "large-headed drop-to-the-ground birds" in forests in Costa Rica does appear to be associated with high insect densities which include higher numbers of large insects (Orians, 1969; Schoener, 1971).

I suggest in summary 1) that *Anolis* populations are sensitive to predation by birds which are capable of depressing the lizard populations below carrying capacity; 2) that anoles may effectively exclude small insectivorous birds from the lower, more connected portions of forest habitats if the anoles are near carrying capacity; and 3) that in forest the small insectivorous birds only have a competitive advantage in the outer canopy, although the potential carrying capacity of this portion for lizards may not be high regardless. The latter two points are based primarily on a close view of behavioral mechanisms of habitat and resource use.

The validity of the latter two assessments depends on the accurate identifi-

cation of the site of overlap in resource use between anoles and the relevant bird species. It is at this same level that accurate identification of site and pattern of foraging overlap between groups has led to demonstration of competitive interaction between rodents and ants (Brown and Davidson, 1977) and between fish and golden-eye ducks (Erickson, 1979). In the present case the structural view of the habitat vis-a-vis the concept of the structural niche developed by Rand and Williams (Rand, 1964; Rand and Williams, 1969) provided the key for a mechanistic assessment of habitat use both within the genus *Anolis* (Williams, 1972; Schoener, 1975; Moermond, 1979a,b) and among various groups of arboreal insectivores (Moermond, 1974).

ACKNOWLEDGMENTS

Numerous people have contributed in many ways to this paper over the past nine years. T. W. Schoener offered observations, encouragement, and insights throughout. R. M. Andrews, J. S. Denslow, S. A. Foster, and T. W. Schoener read the manuscript and offered numerous useful suggestions. J. S. Denslow assisted in the Dominican Republic. Field work was supported by National Science Foundation (NSF) grants to E. E. Williams and an NSF grant to the author. Observations from Costa Rica were supported by an NSF grant to the author and J. S. Denslow. The largest debt is to Ernest E. Williams, whose understanding of anoles and people provided the forum for this and other ideas.

LITERATURE CITED

- ALLAN, J. D., L. W. BARNHOUSE, R. A. PRESTBYE, AND D. R. STRONG. 1973. On foliage arthropod communities of Puerto Rican second growth vegetation. *Ecology*, **54**: 628-632.
- ANDREWS, R. M. 1971. Structural habitat and time budget of a tropical *Anolis* lizard. *Ecology*, **52**: 262-270.

- . 1976. Growth rate in island and mainland anoline lizards. *Copeia*, **1976**: 477–482.
- . 1979. Evolution of life histories: a comparison of *Anolis* lizards from matched island and mainland habitats. *Breviora Mus. Comp. Zool.* No. 454, pp. 1–51.
- BENNETT, S. E. 1980. Interspecific competition and the niche of the American redstart (*Setophaga ruticilla*) in wintering and breeding communities, pp. 319–335. In A. Keast and E. S. Morton (eds.), *Migrant Birds in the Neotropics*. Smithsonian, Washington, D.C.
- BOCK, W. J., AND W. DEW. MILLER. 1959. The scansorial foot of the woodpeckers, with comments on the evolution of perching and climbing feet in birds. *Amer. Mus. Nov.* No. 1931, pp. 1–45.
- BOND, J. 1974. *Birds of the West Indies*. 4th Ed. London, Collins.
- BROWN, J. H., AND D. W. DAVIDSON. 1977. Competition between seed-eating rodents and ants in desert ecosystems. *Science*, **196**: 880–882.
- CHARNOV, E. L., G. H. ORIAN, AND K. HYATT. 1976. Ecological implications of resource depression. *Amer. Nat.*, **110**: 247–259.
- COX, G. W., AND R. E. RICKLEFS. 1977. Species diversity and ecological release in Caribbean land bird faunas. *Oikos*, **28**: 113–122.
- CRUZ, A. 1980. Feeding ecology of the black-whiskered vireo and associated gleaner birds in Jamaica. *Wilson Bull.*, **92**: 40–52.
- CURIO, E. 1970. Die Selektion drier Raupenformen eines Schwarmes (Lepidopt., Sphingidae) durch einen *Anolis* (Rept., Iguanidae). *Z. Tierpsychol.*, **27**: 889–914.
- DIAMOND, A. W. 1973. Habitats and feeding assemblages of St. Lucia forest birds. *Ibis*, **115**: 313–329.
- DIAMOND, J. M. 1970. Ecological consequences of island colonization by Southwest Pacific birds. I. Types of niche shifts. *Proc. Nat. Acad. Sci.*, **67**: 529–536.
- DILGER, W. C. 1956. Adaptive modifications and ecological isolating mechanisms in the thrush genera *Catharus* and *Hylocichla*. *Wilson Bull.*, **68**: 170–199.
- ERIKSSON, M. O. G. 1979. Competition between freshwater fish and goldeneyes, *Bucephala clangula* (L.), for common prey. *Oecologia*, **41**: 99–107.
- FEDUCCIA, A. 1973. Evolutionary trends in the neotropical ovenbirds and woodhewers. *Ornith. Monogr.* No. 13, pp. 1–69.
- FITZPATRICK, J. W. 1978. Foraging behavior and adaptive radiation in the avian family Tyrannidae. Ph.D. thesis. Princeton University. University Microfilms International Order No. 78–18333. Ann Arbor, Michigan.
- FLEMING, T. H., AND R. S. HOOKER. 1975. *Anolis cupreus*: the response of a lizard to tropical seasonality. *Ecology*, **56**: 1243–1261.
- FORBES, A., S. FOX, E. MCCARTHY, AND E. YAMASHITA. 1964. Quantitative response to color shift in diurnal lizards. *Proc. Nat. Acad. Sci.*, **52**: 667–672.
- HESPENHEIDE, H. A. 1975. Prey characteristics and predator niche width, pp. 158–180. In M. L. Cody and J. M. Diamond (eds.), *Ecology and Evolution of Communities*. Cambridge, Harvard Univ. Press.
- HOWE, R. W., J. W. BRUSKEWITZ, T. C. MOERMOND, AND K. A. RUSTERHOLZ. In preparation. Prey handling efficiency by captive warblers. II. Interspecific comparisons.
- JANZEN, D. H. 1973. Sweep sample of tropical foliage insects: effects of seasons, vegetation types, elevation, time of day, and insularity. *Ecology*, **54**: 687–702.
- JENSSEN, T. A. 1973. Shift in the structural habitat of *Anolis opalinus* due to congeneric competition. *Ecology*, **54**: 863–869.
- KARR, J. R. 1971. Structure of avian communities in selected Panama and Illinois habitats. *Ecol. Monogr.*, **41**: 207–233.
- KEAST, A. 1970. Adaptive evolution and shifts in niche occupation in island birds, pp. 39–53. In W. L. Stern (ed.), *Adaptive Aspects of Insular Evolution*. Pullman, Washington State Press.
- . 1980. Spatial relationships between migratory Parulid warblers and their ecological counterparts in the Neotropics, pp. 109–130. In A. Keast and E. S. Morton (eds.), *Migrant Birds in the Neotropics*, Smithsonian, Washington, D.C.
- KEAST, A. AND E. S. MORTON. 1980. Migrant birds in the neotropics: ecology, behavior, distribution and conservation. *Symp. Nat. Zool. Park, Smithsonian Inst., Washington, D.C.*, 576 pp.
- KEPLER, A. K. 1977. Comparative study of Todies (Todidae): with emphasis on the Puerto Rican tody, *Todus mexicanus*. *Publ. Nuttall Ornith. Club* No. 16, pp. 1–190.
- KIESTER, A. R. 1971. Species density of North American amphibians and reptiles. *Syst. Zool.*, **20**: 127–137.
- LACK, D. 1971. *Ecological Isolation in Birds*. Cambridge, Harvard Univ. Press.
- . 1976. *Island Biology Illustrated by the Land Birds of Jamaica*. Studies in Ecology, Vol. 3, Berkeley, Univ. Calif. Press.
- LACK, D., AND P. LACK. 1972. Wintering warblers in Jamaica. *Living Bird*, **11**: 129–153.
- LASIEWSKI, R. C., AND W. R. DAWSON. 1967. A re-examination of the relation between standard metabolic rate and body weight in birds. *Condor*, **69**: 13–23.
- LICHT, P. 1974. Response of *Anolis* lizards to food supplementation in nature. *Copeia*, **1974**: 215–221.
- LICHT, P., AND R. E. JONES. 1967. Effects of exogenous prolactin on reproduction and growth in adult males of the lizard *Anolis carolinensis*. *Gen. Comp. Endocrinol.*, **8**: 228–244.
- LISTER, B. C. 1976. The nature of niche expansion

- in West Indian *Anolis* lizards. I. Ecological consequences of reduced competition. *Evolution*, **30**: 659–676.
- MACARTHUR, R. H., J. M. DIAMOND, AND J. R. KARR. 1972. Density compensation in island faunas. *Ecology*, **53**: 330–342.
- MACARTHUR, R. H., H. RECHER, AND M. CODY. 1966. On the relation between habitat selection and species diversity. *Amer. Nat.*, **100**: 319–332.
- MOERMOND, T. C. 1974. Patterns of habitat utilization in *Anolis* lizards. Ph.D. thesis, Harvard University.
- . 1976. The basis for resource partitioning in birds. Abstract. Symp. on Behavioral Allocation of Resources in Vertebrates. Midwest Anim. Behav. Conf. Chicago, Illinois.
- . 1979a. Habitat constraints on the behavior, morphology, and community structure of *Anolis* lizards. *Ecology*, **60**: 152–164.
- . 1979b. The influence of habitat structure on *Anolis* foraging behavior. *Behaviour*, **70**: 147–167.
- . 1981. Prey-attack behavior of *Anolis* lizards. *Z. Tierpsychol.* **56**: 128–136.
- MOERMOND, T. C., AND J. S. DENSLOW. In preparation. A comparison of the foraging patterns of migrant Parulid warblers and *Microlegia palustris* in montane Dominican Republic.
- MOERMOND, T. C., R. W. HOWE, J. W. BRUSKEWITZ, AND K. A. RUSTERHOLZ. In preparation. Experiments on perch use in warblers: interspecific differences due to morphological and behavioral attributes.
- MORSE, D. H. 1975. Ecological aspects of adaptive radiation in birds. *Biol. Rev.*, **50**: 167–214.
- . 1979. Habitat use by the blackpoll warbler. *Wilson Bull.*, **91**: 234–243.
- ODUM, H. T., W. ABBOTT, R. K. SELANDER, F. B. GOLBY, AND R. F. WILSON. 1970. Estimates of chlorophyll and biomass of the tabonuco forest of Puerto Rico, pp. 3–19. H. T. Odum (ed.), *A Tropical Rain Forest*, Section I. Oak Ridge, U.S. Atomic Energy Comm.
- ORIAN, G. 1969. The numbers of bird species in some tropical forests. *Ecology*, **50**: 783–801.
- OSTERHAUS, M. B. 1962. Adaptive modifications in the leg structure of some North American warblers. *Amer. Midl. Nat.*, **68**: 475–486.
- PARNELL, J. F. 1969. Habitat relations of the Parulidae during spring migration. *Auk*, **86**: 505–521.
- PEARSON, D. L. 1977. Ecological relationships of small antbirds in Amazonian bird communities. *Auk*, **94**: 283–292.
- PIANKA, E. R. 1975. Niche relations of desert lizards, pp. 292–314. In M. L. Cody and J. M. Diamond (eds.), *Ecology and Evolution of Communities*. Cambridge, Harvard Univ. Press.
- . 1978. *Evolutionary Ecology*, 2nd Edition. Harper & Rowe, New York.
- POUGH, F. H. 1980. The advantages of ectothermy for tetrapods. *Amer. Nat.*, **115**: 92–112.
- RAND, A. S. 1964. Ecological distribution in anoline lizards of Puerto Rico. *Ecology*, **45**: 745–752.
- . 1967. Ecology and social organization in the Iguanid lizard, *Anolis lineatopus*. *Proc. U.S. Nat. Mus.*, **122**: 1–79.
- RAND, A. S., AND S. S. HUMPHREY. 1968. Interspecific competition in the tropical rain forest: ecological distribution among lizards at Belem, Para. *Proc. U.S. Nat. Mus.*, **125**: 1–17.
- RAND, A. S., AND E. E. WILLIAMS. 1969. The anoles of La Palma: aspects of their ecological relationships. *Breviora Mus. Comp. Zool.* No. 327, pp. 1–17.
- RICKLEFS, R. E., AND G. W. COX. 1972. Taxon cycles in the West Indian avifauna. *Amer. Nat.*, **106**: 195–219.
- . AND ———. 1977. Morphological similarity and ecological overlap among passerine birds on St. Kitts; British West Indies. *Oikos*, **29**: 20–66.
- ROBERTS, T. S. 1955. *A manual for the identification of the birds of Minnesota and neighboring states*. Minneapolis, Univ. Minnesota Press.
- SCHOENER, T. W. 1965. The evolution of bill size differences among sympatric congeneric species of birds. *Evolution*, **19**: 189–213.
- . 1967. The ecological significance of sexual dimorphism in size in the lizard *Anolis conspersus*. *Science*, **155**: 474–477.
- . 1968. The *Anolis* lizards of Bimini: resource partitioning in a complex fauna. *Ecology*, **49**: 704–726.
- . 1969. Size patterns in West Indian *Anolis* lizards. I. Size and species diversity. *Syst. Zool.*, **18**: 386–401.
- . 1970. Size patterns in West Indian *Anolis* lizards. II. Correlations with the size of particular sympatric species—displacement and convergence. *Amer. Nat.*, **104**: 155–174.
- . 1971. Large-billed insectivorous birds: a precipitous diversity gradient. *Condor*, **73**: 154–161.
- . 1972. Competition and the niche, pp. 35–136. In C. Gans and D. W. Tinkle (eds.), *Biology of the Reptilia*, Volume 7, Ecology and Behavior. New York, Academic Press.
- . 1975. Presence and absence of habitat shift in some widespread lizard species. *Ecol. Monogr.*, **45**: 233–258.
- . 1976. The species-area relations within archipelagos: models and evidence from island land birds. *Proc. 16th Internat'l. Ornith. Cong.*, Canberra, pp. 629–642.
- SCHOENER, T. W., AND G. C. GORMAN. 1968. Some niche differences in three lesser Antillean lizards of the genus *Anolis*. *Ecology*, **49**: 819–830.
- SCHOENER, T. W., AND A. SCHOENER. 1971a. Structural habitats of West Indian *Anolis* lizards. I. Lowland Jamaica. *Breviora Mus. Comp. Zool.* No. 368, pp. 1–53.
- . AND ———. 1971b. Structural habitats of West Indian *Anolis* lizards. II. Puerto Rican Uplands.

- Breviora Mus. Comp. Zool. No. 375, pp. 1-39.
- _____. AND _____. 1978a. Estimating and interpreting body-size growth in some *Anolis* lizards. *Copeia*, 1978: 390-405.
- _____. AND _____. 1978b. Inverse relation of survival of lizards with island size and avifaunal richness. *Nature*, 274: 685-687.
- _____. AND _____. 1980. Densities, sex ratios and population structure in four species of Bahamian *Anolis* lizards. *J. Anim. Ecol.*, 49: 19-53.
- SEXTON, O. J. 1964. Differential predation by the lizard *Anolis carolinensis* upon unicoloured and polycoloured insects after an interval of no contact. *Anim. Behav.*, 12: 101-110.
- SILLMAN, A. J. 1973. Avian vision, pp. 349-387. In D. S. Farner and J. R. King (eds.), *Avian Biology*, Volume III. New York, Academic Press.
- STAMPS, J. 1977. Rainfall, moisture and dry season growth rates in *Anolis aeneus*. *Copeia*, 1977: 415-419.
- TERBORGH, J. W. 1973. Chance, habitat and dispersal in the distribution of birds in the West Indies. *Evolution*, 27: 338-349.
- TERBORGH, J. W., AND J. R. FAABORG. 1973. Turnover and ecological release in the avifauna of Mona Island, Puerto Rico. *Auk*, 90: 759-779.
- _____. AND _____. 1980a. Factors affecting the distribution and abundance of North American migrants in the Eastern Caribbean region, pp. 145-155. In A. Keast and E. S. Morton (eds.), *Migrant Birds in the Neotropics*. Smithsonian, Washington, D.C.
- _____. AND _____. 1980b. Saturation of bird communities in the West Indies. *Amer. Nat.*, 116: 178-195.
- TERBORGH, J. W., J. R. FAABORG, AND H. J. BROCKMAN. 1978. Island colonization by lesser Antillean birds. *Auk*, 95: 59-72.
- WETMORE, A. 1916. Birds of Puerto Rico. *Bull. U.S. Dept. Agr.*, 326: 1-140.
- WETMORE, A., AND B. SWALES. 1931. The birds of Haiti and the Dominican Republic. *Bull. U.S. Nat. Mus.*, 155: 1-483.
- WILLIAMS, E. E. 1962. Notes on Hispaniolan herpetology. G. The giant anoles. *Breviora Mus. Comp. Zool.* No. 155, pp. 1-15.
- _____. 1965. The species of Hispaniolan green anoles (Sauria, Iguanidae). *Breviora Mus. Comp. Zool.* No. 227, pp. 1-16.
- _____. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, 44: 345-389.
- _____. 1972. The origin of faunas. Evolution of lizard congeners in a complex island fauna: a trial analysis. *Evol. Bio.*, 6: 47-89.
- _____. 1976. West Indian anoles: a taxonomic and evolutionary summary. I. Introduction and a species list. *Breviora Mus. Comp. Zool.* No. 440, pp. 1-21.
- WILLIAMSON, P. 1971. Feeding ecology of the red-eyed vireo (*Vireo olivaceus*) and associated foliage-gleaning birds. *Ecol. Monogr.*, 41: 129-152.
- WRIGHT, J. S. 1979. Competition between insectivorous lizards and birds in central Panama. *Amer. Zool.*, 19: 1145-1156.
- _____. 1981. Extinction-mediated competition: the *Anolis* lizards and insectivorous birds of the West Indies. *Amer. Nat.*, 117: 181-192.

A Synecological Study of Surinam Monkeys

RUSSELL A. MITTERMEIER¹

MARC G. M. VAN ROOSMALEN²

ABSTRACT. Surinam's eight monkey species were studied in the Raleighvallen-Voltzberg Nature Reserve to determine how these species partition available habitat and food resources. *Cebus apella*, *Saguinus midas midas*, and *Saimiri sciureus* were the most adaptable species and regularly occurred in all five forest types in the study area. *Ateles paniscus paniscus*, *Chiropotes satanas chiropotes*, *Pithecia pithecia*, and *Cebus nigrivittatus* were far more restricted in choice of habitat, and *Alouatta seniculus*, though it entered all five forest types, had a strong preference for one. Seven of the eight species were found most often in high forest, but *Saimiri* was most frequently seen in liane forest and occupied lower levels of the forest than any other species. *Saguinus* was found in edge habitats more often than the other species, but *Saimiri* and *Cebus apella* regularly used edges as well. *Saguinus*, *Saimiri*, *Cebus apella*, and *Cebus nigrivittatus* were omnivorous, with *Saimiri* apparently eating the highest percentage of arthropods. The four other species were not seen to feed on insects or other animal life. All eight species ate a high proportion of fruit, with flowers playing only a very small role. The two largest species (*Alouatta* and *Ateles*) ate the most leaves, whereas the three smallest (*Saguinus*, *Saimiri*, *Pithecia*) ate no leaves at all. *Chiropotes* was found to be an important seed predator, and *Ateles* and *Alouatta* were major seed dispersers.

INTRODUCTION

Although a number of studies (e.g., Gautier and Gautier-Hion, 1969;

Charles-Dominique, 1971; Gartlan and Struhsaker, 1972; Rodman, 1973a, b, 1978; Dunbar and Dunbar, 1974; Quris, 1976; Moreno-Black and Maples, 1977; Struhsaker and Leland, 1979) have investigated certain aspects of habitat utilization and niche separation in a number of Old World primate communities, very little in the way of quantitative synecological data has been collected on rain forest-dwelling Neotropical primates. Hladik and Hladik's (1969, 1971, 1975) studies provided quantitative data only on diet, and other investigations of New World monkey synecology (e.g., Klein and Klein, 1975, 1976, 1977; Izawa, 1975, 1976) have been mainly qualitative.

As a graduate student in the Department of Anthropology and a long-term squatter in the Herpetology Department of the Museum of Comparative Zoology, R.A.M. was much impressed with and inspired by the rigorous quantitative analyses of ecological relationships in various Caribbean *Anolis* communities carried out by Ernest E. Williams and many of his students. Although Neotropical monkeys are not quite the cooperative subjects that *Anolis* lizards are, it seemed possible to apply at least some of the methods being used to learn more of monkeys in South America. Consequently, when we found a suitable study site in the form of a diverse and undisturbed primate community in the interior of Surinam, we decided to make a preliminary attempt to define quantitatively the niches occupied by sympatric New

¹ World Wildlife Fund-U.S. Primate Program, 1601 Connecticut Avenue, N.W., Washington, D.C. 20009, U.S.A. and Department of Anatomical Sciences, SUNY, Stony Brook, New York 11794, U.S.A.

² Rijksinstituut voor Natuurbeheer, Kasteel Broekhuizen 2, P.O. Box 46, 3956ZR Leersum, Netherlands.

Figure 1. Map of Surinam showing the location of the Raleighvallen-Voltzberg Nature Reserve and the other protected areas in the country.

World monkeys and to determine how they divide up available forest resources.

There are eight primate species in Surinam, and they include seven members of the family Cebidae and one species of Callitrichidae. The callitrichid is the golden-handed tamarin (*Saguinus midas midas*), which weighs about 500 g and lives in groups that average six individuals. The cebids range in size from the squirrel monkey (*Saimiri sciureus*), which weighs about 700 g and travels in huge groups that can exceed 50 individuals, to the black spider monkey (*Ateles paniscus paniscus*), with males and females averaging about 7.8 kg. The social structure of spider monkeys is unique among the Surinam species in that the animals apparently live in temporary "parties" or "subgroups" within larger, loosely-organized communities. Other cebids include the white-faced saki (*Pithecia pithecia*) and the bearded saki (*Chiropotes satanas chiropotes*), both of them members of the unusual

cebid subfamily, Pitheciinae. The white-faced saki lives in pairs or small family groups, weighs about 1.5 to 2.0 kg, and is the only Surinam species that is strongly sexually dimorphic in color. The bearded saki is larger, weighing about 3.0 kg, lives in group of eight to more than 30 individuals, and is not dimorphic. Two species of capuchin monkey also occur in Surinam. The tufted capuchin (*Cebus apella apella*) is found throughout the country and is probably the most adaptable monkey. Males average about 3.9 kg and females 3.0 kg, and the animals live in groups of six to 20 or more individuals. The weeper capuchin (*Cebus nigrivittatus*), in contrast to its relative, is probably the rarest Surinam monkey, but its weight and group size are similar to that of *C. apella*. And finally, there is the red howler monkey (*Alouatta seniculus*), a large species that lives in one male groups averaging four to five animals and is sexually dimorphic in size, with males weighing about 8.5 kg and females about 6.0 kg (Mittermeier, 1977).

METHODS

The study discussed here was conducted in the Raleighvallen-Voltzberg Nature Reserve, a 56,000 ha protected area established in 1961 and currently managed by STINASU, the Surinam Nature Conservation Foundation. In early 1976, we established a 300 ha study site at the foot of the 240 m Voltzberg dome, one of a number of huge granitic *inselbergs* found in the reserve.

The study area has four major forest types and a fifth transition type, all of which are briefly described in the Appendix. High forest is the most common formation, followed by liane forest and mountain savanna forest, and *pina* swamp forest grows along the small creeks flowing through the reserve. The Voltzberg area apparently has a greater diversity of forest types and edges than would be usual in a 300 ha tract of forest in the interior of Surinam (J. Lindeman,

personal communication), probably because of the many granite formations that do not always provide adequate support for high forest trees, but do allow the growth of relatively rare forest types like mountain savanna forest and liane forest. As a result of this unusual diversity, it was possible to study the habitat preferences of all eight Surinam monkey species in a much more restricted area than would have been possible in most other parts of the country.

The study discussed here was conducted on 147 field days between March, 1976 and February, 1977. Data were gathered during 12 census walks of the whole study area, and also while searching for "target species" selected on a given day. Every time a group was encountered, the following kinds of information were recorded: time, location in the study area, forest type, edge or non-edge situation, activity (resting, traveling, feeding on plant food, foraging for/feeding on insects), height and stratum in the forest, how located, path animal and observer animal distances (for census purposes), group size and composition (if possible), and direction of travel.

A sighting was recorded as an edge if the first animal encountered was within 20 m of a clearing or another forest type. Feeding on plant food was considered the activity if the first animal observed was eating fruits, flowers, leaves, or other vegetable matter. Foraging for/feeding on insects was recorded as the activity if the first animal seen was eating, catching, or actively searching for insects or other animal life. For stratification observations, the forest was divided into six sections: shrub layer (0–3 m), understory (3–15 m), lower part canopy (15–20 m), middle part canopy (20–25 m), upper part canopy (25–30 m), and emergents ($\geq$ 30 m). In order to avoid possible bias because of the presence of the observer, only first sighting data are used for the histograms of utilization of forest types, edges, vertical stratification, and plant vs. insect eating. Diet histograms, which are

unlikely to be affected by observer presence, are based on all feeding observations.

RESULTS

HABITAT UTILIZATION

Forest type utilization. High forest was the most frequently used forest type for seven of the Surinam species, whereas the eighth, *Saimiri sciureus*) was most often in liane forest (Fig. 4 a–h). *Ateles* was found almost exclusively in high forest; *Cebus nigrivittatus* was very dependent on it as well; and *Chiropotes* was found only in high forest and mountain savanna forest. *Pithecia* was similar to *Chiropotes*, but also entered liane forest on occasion. *Saguinus*, *Saimiri*, *Cebus apella*, and *Alouatta* spent at least some time in all five formations, but *Alouatta* was more dependent on high forest than the three smaller species.

Edge utilization. Three of the eight species studied were regularly seen in edge situations, with *Saimiri* and *C. apella* being found in edges about half the time and *Saguinus* more often than in non-edges. By contrast, *Ateles* was almost never found in edge habitats, and *Pithecia*, *Chiropotes*, *C. nigrivittatus* and *Alouatta* only entered them on occasion (Fig. 5a–h).

Vertical stratification. *Ateles* and *Chiropotes* were found primarily in the two highest strata of the forest; *Alouatta* was most often in the middle to upper strata; and *Saguinus*, *Pithecia*, and the two *Cebus* were mainly in the understory and lower to middle strata. *Saimiri*, in keeping with its preference for liane forest, was in the understory stratum more often than anywhere else. None of the eight species in the Voltzberg area was seen on the ground during the present study, and only three (*Saguinus*, *Saimiri*, *C. apella*) entered the shrub layer. Two species (*Pithecia*, *C. apella*) were never seen in emergents. The understory, and

Figure 2. a. Adult golden-handed tamarin (*Saguinus midas midas*). b. Young adult male squirrel monkey (*Saimiri sciureus*). c. Adult male white-faced saki (*Pithecia pithecia*). d. Adult female white-faced saki (*Pithecia pithecia*). e. Adult bearded saki (*Chiropotes satanas chiropotes*) (photo courtesy of the Köln Zoological Society).

Figure 3. a. Juvenile tufted capuchin monkey (*Cebus apella apella*). b. Young adult male weeper capuchin monkey (*Cebus nigrivittatus*). c. Adult female (above) and subadult (below) red howler monkey (*Alouatta seniculus*). d. Juvenile black spider monkey (*Ateles paniscus paniscus*).

middle, lower, and upper levels of the canopy were utilized to varying extent by all eight species (Figs. 6a-h).

DIET

Herbivory vs. omnivory. Sharp distinctions can be made in diet since half the species are partly insectivorous whereas the other four appear to be entirely herbivorous. *Saguinus*, *Saimiri*, and the two *Cebus* species regularly include arthropods in their diet (Fig. 7a-d), while *Pithecia*, *Chiropotes*, *Alouatta*, and *Ateles* were never seen eating insects or other animals.

Fruits and seeds: seed predation vs. seed dispersal and seed dropping. A further distinction can be made between species that are heavily dependent on seeds as a food source and those that rarely or never actually digest seeds. Here it is necessary to distinguish between ingestion of seeds and digestion of seeds. All species swallow at least some seeds along with the exocarp, mesocarp and/or aril that they eat, but most species simply swallow the seeds intact without mastication and pass them through the digestive system without destroying their powers of germination. The result is referred to as *endozoochory* and is an important method of *seed dispersal* for forest trees. In addition, seeds are often dropped, especially by the smaller monkey species, after the softer, edible parts are removed with the hands or mouth. *Seed dropping* may also be a form of seed dispersal if the monkey carries the seeds to another tree before dropping them (*exozoochory*). However, in most cases, it appears that the seeds are simply dropped out of the tree from which they are taken and do not travel any further than if they had fallen by themselves. This kind of seed dropping is not really a

form of seed dispersal, but we have included it together with endo- and exozoochory in Figure 8 because none of these feeding behaviors destroys the seeds ability to germinate and also because it is at times difficult to distinguish between exozoochory and simple seed dropping.

Contrasting with these behaviors is *seed predation* in which the seeds of certain families (e.g., *Leyceidaceae*) are actually eaten by the monkeys. The seeds are removed from the fruit, chewed into a pulp and digested, obviously losing their powers of germination.

Although all eight monkeys species sometimes perform endozoochory with very small seeds of genera like *Ficus* (*Moraceae*), *Coussapoa* (*Moraceae*), *Cecropia* (*Moraceae*), *Bagassa* (*Moraceae*), *Passiflora* (*Passifloraceae*), *Duroia* (*Rubiaceae*), *Bellucia* (*Melastomataceae*), and *Hylocereus* (*Cactaceae*), the major primate seed dispersers are *Ateles* and *Alouatta*. These monkeys ingest many large and small seeds every day, and several hundred seeds of up to five species are at times found in single fecal samples. *Ateles* is probably a more important dispersal agent because of its much larger range and higher activity level, but *Alouatta* also disperses seeds hundreds of meters from where they were eaten. All eight species also perform exozoochory on occasion, but primates apparently play only a very minor role in this form of dispersal in Surinam.

Seed predation appears to be of major importance for *Chiropotes* (Fig. 8d), and quite possibly for *Pithecia* (Fig. 8c) and *C. nigrivittatus* (Fig. 8f) as well, although the samples for these two species are quite small. *Saguinus*, *C. apella*, and *Ateles* sometimes eat seeds, but much less than *Chiropotes* (Fig. 8a,e,h), where-

Figure 4. Utilization of the five forest types in the Voltzberg study area. a. *Saguinus midas midas*. b. *Saimiri sciureus*. c. *Pithecia pithecia*. d. *Chiropotes satanas chiropotes*. e. *Cebus apella apella*. f. *Cebus nigrivittatus*. g. *Alouatta seniculus*. h. *Ateles paniscus paniscus*.

Figure 5. Utilization of edges in the Voltzberg study area. a. *Saguinus midas midas*. b. *Saimiri sciureus*. c. *Pithecia pithecia*. d. *Chiropotes satanas*. e. *Cebus apella apella*. f. *Cebus nigritatus*. g. *Alouatta seniculus*. h. *Ateles paniscus paniscus*.

Figure 6. Utilization of forest strata in the Voltzberg study area. a. *Saguinus midas midas*. b. *Saimiri sciureus*. c. *Pithecia pithecia*. d. *Chiropotes satanas*. e. *Cebus apella apella*. f. *Cebus nigrivittatus*. g. *Alouatta seniculus*. h. *Ateles paniscus paniscus*.

Figure 7. Feeding on plant food vs. foraging for/feeding on insects and other animals in the Voltzberg study area. Only four species were seen feeding on insects or other animals. a. *Saguinus midas midas*. b. *Saimiri sciureus*. c. *Cebus apella apella*. d. *Cebus nigrivittatus*.

as *Saimiri* and *Alouatta* were never seen acting as seed predators (Fig. 8b,g).

Fruits vs. flowers and leaves. Fruits made up the majority of the plant diet for all species, with flowers and leaves usually contributing only a small portion. Fruits (including seeds) accounted for over 90% of the plant foods eaten by *Saguinus*, *Saimiri*, *Pithecia*, *Chiropotes*, and *C. apella*, and over 80% of those eaten by *C. nigrivittatus* and *Ateles* (Fig. 8a-f,h). *Alouatta* was the only species

that ate a relatively high percentage of leaves, and even it appears to be highly dependent on fruit in the wet season (Fig. 8g). Of the five species eating leaves, *Alouatta* was the only one that included mature leaves in its diet. *Chiropotes*, *C. apella*, *C. nigrivittatus*, and *Ateles* were only seen eating flush leaves, and the three smallest species (*Saguinus*, *Saimiri*, *Pithecia*) did not eat leaves at all (Fig. 8a-h).

Differential utilization of fruit families and fruit species. More detailed analysis of the plant foods eaten by the eight monkeys indicates that they were feeding on 151 identified and 30 unidentified species of at least 54 families during this study. Only one family, the Moraceae, was utilized by all eight monkeys, and no single plant species was eaten by more than six of them. Seven monkey species were observed to eat members of the Leguminosae Mimosaaceae, six fed on various Sapotaceae, Rubiaceae, and Palmae, and five ate some Burseraceae, Lecythidaceae, Myrtaceae, and Passifloraceae. Several families like the Capparaceae and Meliaceae were very important seasonally for some monkeys, but were not eaten by others. The majority of plant families were only occasionally used as dietary supplements for a few monkey species. Comparison of species rather than families reveals even less overlap, since most plant species are infrequently eaten by only one or two monkey species (Mittermeier, 1977; van Roosmalen, *et al.*, in preparation).

Several very interesting examples of niche separation were evident in utilization of the same food species at different stages of growth. For example, *Chiropotes* broke open unripe *Brosimum parinarioides* (Moraceae) and ate the seeds, thus acting as a seed predator, whereas *C. apella* and *Ateles* swallowed the mesocarp and seeds of ripe fruits and later excreted the seeds, thus acting as dispersers. A similar situation was observed with *Ecclinusa guianensis* (Sapotaceae).

Figure 8. Utilization of plant foods in the Voltzberg study area. The "seeds dispersed" category includes actual seed dispersal (endozoochory and exozoochory) and also simple seed dropping. The "p" category includes those plant species for which we were unable to determine if seed predation was taking place. Percentages are based on the total number of feeding observations. a. *Saguinus midas midas*. b. *Saimiri sciureus*. c. *Pithecia pithecia*. d. *Chiropotes satanas chiropotes*. e. *Cebus apella apella*. f. *Cebus nigrivittatus*. g. *Alouatta seniculus*. h. *Ateles paniscus paniscus*.

Chiropotes fed on the young seeds of this species, while *Ateles* swallowed the ripe fruits and excreted the seeds. Competition of this kind was seen in the same tree at the same time, with *Chiropotes* feeding on the young fruits to such an extent that few had a chance to ripen to a stage at which *Ateles* would find them palatable.

Finally, it is necessary to point out that the sample sizes for diet in *Pithecia* and *C. nigrivittatus* are very small (Fig. 8c,f) and give only an approximate indication of the foods eaten by these animals. However, the data on the other six species are a good representation, though by no means complete, of the kinds of plant foods eaten by these monkeys. More in-depth coverage of this same study is provided in Mittermeier (1977) and Mittermeier and van Roosmalen (1981), and an additional two years of data are now in the process of being analyzed (van Roosmalen, *et al.*, in preparation) and should add greatly to our knowledge of these monkeys.

DISCUSSION

COMMENTS ON NICHE SEPARATION

It should be clear from the above data that there is overlap in almost all the variables considered and that the monkeys are at least in competition for many of the fruits of the major edible-fruit producing families like the Moraceae. However, it should be equally clear that the eight species are avoiding competition and are dividing up the habitat in a number of important ways. Some species (e.g., *Ateles*, *Chiropotes*) are spending most of their time in the upper parts of high forest, others (e.g., *Saimiri*) are in the lower strata of formation like liane forest; half the species investigated (*Saguinus*, *Saimiri*, *Cebus apella*, *Cebus nigrivittatus*) are eating arthropods and other animals; others (e.g., *Ateles*, *Alouatta*) are not. Three species

(*Saguinus*, *Saimiri*, *Cebus apella*) frequent edges, perhaps to take advantage of higher insect abundance in such ecotone areas, whereas others (especially the highly frugivorous species, e.g., *Ateles*) are rarely found in edges. Some species act as seed predators (e.g., *Chiropotes*), whereas others (e.g., *Saimiri*, *Alouatta*) rarely or never eat seeds; and each monkey is feeding on a particular set of food items that overlaps partly, but never entirely, with other species.

IMPORTANCE OF THE SYNECOLOGICAL STUDIES FOR CONSERVATION

Wild populations of nonhuman primates are diminishing rapidly in many parts of the world, making the need for adequate protected areas ever more urgent. New national parks and reserves must be created in important areas in which they do not yet exist, and established protected areas must be assessed to determine if they are of long term conservation value. When selecting sites for new reserves and assessing those already in existence, synecological data of the kind presented here can be a very useful tool. Such data can be used to determine if habitats in a particular area are likely to support populations of the primates found in a given country or region and, if not, what kinds of forest should be added for a more complete representation. If one or two primate species are of major concern, a reserve can be selected with their particular ecological requirements in mind. Data from an undisturbed site like the Voltzberg area in Surinam can also be used to assess disturbed areas to determine the extent of damage and the potential for recovery as useful primate habitat.

Synecological data can also be of great help in determining priorities in primate conservation. For instance, the study covered in this paper indicates that *Saguinus midas*, *Saimiri sciureus*, and *Cebus apella* are very adaptable species occurring in a wide variety of habitats,

whereas *Ateles* and *Chiropotes* are restricted to undisturbed formations and are therefore far more vulnerable. Information of this kind, used together with data on hunting and habitat destruction, makes it possible to formulate an overall picture of primate status in a particular country or region, to indicate which species should be the focus of conservation attention, and to evaluate potential reserves with sound ecological principles. Finally, synecological studies can be used to monitor primate populations in established protected areas like the Raleighvallen-Voltzberg Nature Reserve in Surinam, and to ensure that these areas continue to serve their essential conservation functions.

ACKNOWLEDGMENTS

We would like to express our thanks to J. P. Schulz, H. Reichart, and F. Baal of STINASU (Surinam Nature Conservation Foundation) for permission to work in the Raleighvallen-Voltzberg Nature Reserve, and to J. G. Fleagle for reading and commenting on earlier drafts of this manuscript. R.A.M. would also like to express special thanks to Ernest E. Williams for continual support, intellectual stimulation, ever-expanding office space, and a most congenial atmosphere in which to work during his years at Harvard.

R.A.M. was partly supported by the New York Zoological Society, partly by the World Wildlife Fund-U.S., and partly by a Graduate Fellowship, a Doctoral Dissertation Enrichment Grant, and Grant BNS 7724921 from the National Science Foundation. M.G.M.v.R. was supported by WOTRO, the Netherlands Foundation for Tropical Research.

Figures were prepared by L. Meszoly.

APPENDIX

Forest types in the Voltzberg study area (modified from Lindeman and

Moolenaar, 1959 and P. Teunissen, personal communication).

HIGH RAIN FOREST

High rain forest is the typical *terra firme* tropical forest that is not affected by seasonal fluctuations in river level. It can usually be divided into three to four storeys. The upper storey consists of emergents which reach 40–45 m and, on rare occasions, as high as 60 m. Below the emergents is the canopy, which ranges from 15–30 m and can itself be subdivided into several sub-categories. The understory consists of slender trees up to about 15 m. Saplings and undergrowth species make up the bottom 3 m. High rain forest is very rich in tree species and has far more edible fruit species than any other formation. In the Voltzberg area, the understory is dominated by *boegroemaka* palms (*Astrocaryum sciophilum*).

LOW RAIN FOREST

We use the term low forest to designate a sub-category of high rain forest that usually does not exceed 20 m in height, is far richer in lianes than neighboring high forest, and has far fewer *boegroemaka* palms. It is usually a transition between high forest rich in *boegroemaka* and liane forest, or sometimes between high forest and mountain savanna forest.

MOUNTAIN SAVANNA FOREST

Mountain savanna forest is a rare formation that occurs on bauxite hills and low mountains in the coastal region of Surinam and in parts of the interior (e.g., the Voltzberg area) where only a thin layer of soil covers the underlying rock. It is similar to the better known white and sand savanna forest in xeromorphy, thin-stemmed aspect, coriaceous structure of leaves and lack of differentiation into storeys, but differs in floristic composition. In striking contrast to high forest, *boegroemaka* palms are absent from the understory. The term savanna forest in general covers several forest types that grow in areas with very permeable soil that has poor water retention capacity.

LIANE FOREST

Liane forest is a formation found in areas with stony lateritic soils that provide bad rooting conditions and poor footholds for trees. It is especially noteworthy for the absence of storeys. Tall trees do occur, but they are so widely separated from one another that no true canopy exists. The space between the trees is filled with dense tangles of lianes and vines that grow in abundance because of unrestricted exposure to sunlight. Although occasional tall trees may reach 30 m or more, the liane tangle itself rarely exceeds 10 to 15 m.

PINA SWAMP FOREST

The term swamp forest is used for a number of two-storeyed forest types that are inundated to some extent throughout the year. In the Voltzberg area, this forest type is represented by what we call *pina* swamp forest, a formation dominated by the pina palm (*Euterpe oleracea*) and growing in a narrow belt along the small creeks flowing through the rain forest.

LITERATURE CITED

- CHARLES-DOMINIQUE, P. 1971. Éco-éthologie des prosimiens du Gabon. *Biol. Gabon.*, 7(2): 121-228.
- DUNBAR, R. I. M., AND E. P. DUNBAR. 1974. Ecological relations and niche separation between sympatric terrestrial primates in Ethiopia. *Folia primatol.*, 21(1): 36-60.
- GARTLAN, J., AND T. T. STRUHSACKER. 1972. Polyspecific associations and niche separation of rain-forest anthropoides in Cameroon, West Africa. *J. Zool.*, 168: 221-266.
- GAUTIER, J. P., AND A. GAUTIER-HION. 1969. Les associations polyspécifiques chez les Cercopithecidae du Gabon. *Terre et Vie*, 116: 164-201.
- HLADIK, A., AND C. M. HLADIK. 1969. Rapport trophiques entre végétation et primates dans la forêt de Barro Colorado (Panama). *Terre et Vie*, 23: 25-117.
- HLADIK, C. M. 1975. Ecology, diet, and social pattern in Old and New World primates, pp. 3-35. In R. Tuttle (ed.), *Socioecology and Psychology of Primates*.
- HLADIK, C. M., A. HLADIK, J. BROSSET, P. VALDEBOUSE, G. VIROBEN, AND J. DELORT-LAVAL. 1971. Le régime alimentaire des primates de l'île de Barro Colorado (Panama): résultats des analyses quantitatives. *Folia primatol.*, 16: 85-122.
- IZAWA, K. 1975. Foods and feeding behavior of monkeys in the upper Amazon basin. *Primates*, 17: 295-316.
- . 1976. Group sizes and compositions of monkeys in the upper Amazon basin. *Primates*, 17: 367-399.
- KLEIN, L. L., AND D. J. KLEIN. 1975. Social and ecological contrasts between four taxa of Neotropical primates, pp. 59-85. In R. Tuttle (ed.), *Socioecology and Psychology of Primates*. The Hague, Mouton.
- , AND ———. 1976. Neotropical primates: aspects of habitat usage, population density, and regional distribution in La Macarena, Colombia, pp. 70-78. In R. W. Thorington, Jr. and P. G. Heltne (eds.), *Neotropical Primates: Field Studies and Conservation*. National Academy of Sciences, Washington, D.C.
- , AND ———. 1977. Feeding behavior of the Colombian spider monkey, *Ateles belzebuth*, pp. 153-181. In T. Clutton-Brock (ed.), *Primate Ecology: Studies of Feeding and Ranging Behavior in Lemurs, Monkeys and Apes*, New York, Academic Press.
- LINDEMAN, J. C., AND S. P. MOOLENAAR. 1959. Preliminary survey of the vegetation types of northern Suriname. *The Vegetation of Suriname*, Vol. 1, Part 2, pp. 1-45.
- MITTERMEIER, R. A. 1977. Distribution, synecology and conservation of Surinam monkeys. Unpublished Doctoral Dissertation, Harvard Univ.
- MITTERMEIER, R. A., AND M. G. M. VAN ROOSMALEN. 1981. Preliminary observations on habitat utilization and diet in eight Surinam monkeys. *Folia primatol.* (in press).
- MORENO-BLACK, G., AND W. R. MAPLES. 1977. Differential habitat utilization of four Cercopithecidae in a Kenyan forest. *Folia primatol.*, 27: 85-107.
- QURIS, R. 1976. Données comparative sur la socio-écologie de huit espèces de Cercopithecidae vivant dans une même zone de forêt primitive périodiquement inondée (Nord-est du Gabon). *Terre et Vie*, 30: 193-209.
- RODMAN, P. S. 1973a. Synecology of Bornean primates, with special reference to the behavior and ecology of orang-utans. Unpublished Doctoral Dissertation, Harvard Univ.
- . 1973b. Synecology of Bornean primates. 1. A test for interspecific interactions in spatial distribution of five species. *Amer. J. Phys. Anthropol.*, 38(2): 655-659.
- . 1978. Diets, densities and distributions of Bornean primates. In G. G. Montgomery (ed.), *Ecology of Arboreal Folivores*. Smithsonian Institution, Washington, D.C.
- STRUHSACKER, T. T., AND L. LELAND. 1979. Socioecology of five sympatric monkey species in the Kibale Forest, Uganda, pp. 159-227. In *Advances in the Study of Behavior*, 9. New York, Academic Press.

Courtship Behavior in *Varanus bengalensis* (Sauria: Varanidae)

WALTER AUFFENBERG¹

ABSTRACT. *Varanus bengalensis* utilizes 19 behavioral acts during courtship. These are classified into contextural and functional classes. Extensive analysis is carried out on the sequence of adjacent acts both in two-step and three-step interactions. Courtship behavior appears to select the most powerful males for breeding. Though courtship is initiated by males, most subsequent male acts are determined solely by the female's immediately previous act.

INTRODUCTION

The only published information on varanid courtship behavior is that of *Varanus komodoensis* (Auffenberg, 1981a). The distantly related *Varanus bengalensis* is generally placed in a separate subgenus (*Indovaranus*, *fide* Mertens, 1942), and behavioral comparisons would be useful. Individuals of *Varanus bengalensis* maintained in captive facilities in Gainesville, Florida, afforded an opportunity to study the sexual behavior of adult *Varanus bengalensis*. Other contributions deal with sexual differences in nonsexual contexts (Auffenberg, 1979); agonistic behavior in adolescents (Auffenberg and Ganci, in press); and adult combative behavior (Auffenberg, 1981b).

The primary purposes of this contribution are: 1) to describe courtship in this species in respect to its component behavioral acts; 2) to determine the degree of behavioral act complexity; 3) to deter-

mine the size of the act repertoire; and 4) to describe how these factors are related to the reproductive strategy, through courtship, of this species.

MATERIALS AND METHODS

The data comprising the basis of this study were obtained by review and analysis of closed circuit TV tapes. The equipment included one remote camera appropriately mounted in each of two greenhouses (56 m² each). The cameras ran continuously for eight hours each day (0800–1600 hrs) for a total of 720 days, commencing in June 1977 and continuing to the end of July 1979. Each camera was connected to a time lapse VTR automatic recorder set to advance the tape at 0.3 sec intervals. This allowed for rapid review when set at a normal replay speed, yet preserved sufficient detail of the recorded social interactions. In addition to this analysis system, motion pictures (Super-8) were made whenever practical, often within a 2 m range, so that details of the behavioral acts were clearly evident when reviewed by a stop-motion film projector, using a projection system similar to that described by Jenssen (1977).

Each of the two greenhouses usually contained two adult males and four adult females. During the reproductive season the males were sometimes shifted from one greenhouse to another to encourage courtship interactions with the "resi-

¹Florida State Museum, University of Florida, Gainesville, Florida 32611, U.S.A.

dent" females. The four adult males had an average total length of 149 cm, ($\bar{X}$ SVL $\times$ 58.0 cm; $\bar{X}$ weight = 2743 g), and the eight mature females had an average length of 119.3 cm ($\bar{X}$ SVL = 46.3 cm; $\bar{X}$ weight 1452 g).

Of 351 sexual interactions recorded, only 217 proceeded through the mounting phase, including 21 in which coition was completed. These 217 completed courtship sequences form the basis of this report.

Behavioral Acts. Courtship is defined as comprising those acts leading to, but excluding, intromission itself. In this paper the term "act" is used to designate a particular behavior pattern or display performed; the term "step" denotes the place in the sequence in which an act occurred.

Analyses. Many recent investigators of animal communication have used transition matrices to describe their data. Among reptile behaviorists only Cooper (1977) and Gillingham (1979) have analyzed their data by this method so far. Transition matrices were constructed by enumeration of all observed one-step transitions. This method assumes that both the transition probabilities and the unconditional probabilities of single acts are constant over the length of the interaction; that is, the probabilities exhibit "stationarity."

The analysis technique follows that outlined by Oden (1977), in which stationarity is not assumed, and which allows assessments of the significance of dependence between acts separated by an arbitrary number of steps.

In this type of study it is important to determine whether the observed number of adjacent lower order interactions (one-step transitions; i.e., step 1 followed by step 2, step 2 followed by 3, etc.) is significantly different from expected. Expected frequencies for each of the possible act pairs in adjacent steps were calculated by the formula, $\text{Expected} = \frac{(A_1)(B_2)}{N}$, where A represents one act

of the pair and B the other and the subscripts 1 and 2 the adjacent step numbers.

Expected frequencies of higher order interactions (three steps at a time) were calculated by use of the formula:

$$\text{Expected } A_1 B_2 C_3 = \frac{\text{observed } A_1 B_2 (\text{observed } B_2 C_3)}{\text{observed } B_2},$$

where A, B, C are left, and subscripts 1, 2, 3 are the step numbers. Rejection of the assumption means that dependence of C_3 on A_1 is not transmitted through B_2 .

RESULTS

There are 19 behavioral acts (= motor patterns of some behaviorists, including Gillingham, 1979) that are common components of the courtship of *Varanus bengalensis*. These are defined below (letters in parentheses are abbreviations used). More elaborate descriptions and figures of most of these acts are found in Auffenberg (1981b).

Following an earlier paper (Auffenberg, 1981b), behavioral acts included in courtship are categorized into several contextual and functional classes. These are weaponry use, warning, and investigatory tactics and signals used in social (dominance-subordination) or reproductive contexts.

WEAPONRY CLASS

Biting (Bi). In an agonistic context during the early phases of courtship biting is used rarely by females. Bites administered by females are indiscriminately administered on the head, body, neck, or legs. Males mounted on females also bite, but only in this context and only in a specific place: females are bitten on the dorsal skin of the neck, where the scales tend to be largest. It appears to be an immobilization technique and is common in many lizard groups (for review see Carpenter and Ferguson, 1978).

Tail Slap (TS). A rapid lateral swing of

the tail, often, but not always, following tail coiling (see below). The tail slap usually terminates the interaction. The blow is swift and smartly delivered, usually striking the other individual on the side of its body, or less often the head. Tail slapping in combat has been reported in *V. niloticus*, *varius*, *spenceri*, *mertensi*, and *salvadori* (for review see Auffenberg, 1981a).

WARNING CLASS

Tail Coil (TC). Partially or completely coiling the tail in the horizontal plane, usually preceding and leading to a tail slap. Reported in *V. komodoensis* (Auffenberg, 1981a).

Hiss (Hi). An auditory cue, used during courtship only in a defensive context by females when approached by males.

DOMINANT CLASS

Topping (Tp). One individual (usually the female in a courtship context) puts one of its feet on the back of the male or even climbs upon him (= *riding* of Carpenter and Ferguson, 1978). It only occurs during or after courtship and combat.

SUBORDINANT CLASS

Walks Away (W). One individual walks (or runs) away from the other; may be a neutral signal in some contexts, but clearly subordinate in most. Also reported in a subordinate context in *V. komodoensis* (Auffenberg, 1978b, 1981a) and *V. salvator* (Vogel, 1979).

The following three acts may be functionally identical, i.e., they simply serve to move the head away from the tongue of the initiator (avoidance = "turning away" of Vogel, 1979).

Head Down (HD). Head tilted downward and held lower than the body axis. Occurs in investigative context.

Head Away (HA). Head moved lateral to central body axis and held there. Occurs in investigative context.

Head Up (HU). Head lifted above the trunk and held there. Occurs in several contexts, but most common in investigative context. In an agonistic context signifies social dominance (Auffenberg, 1981b; Greenberg, 1977; Weaver, 1970).

INVESTIGATIVE CLASS

Tonguing (To). Touching or extending the tongue toward another individual. It occurs prior to combat or courtship. Concomitant pressing of the snout (= *touch* of Gillingham, 1979, in snakes) against the body is particularly common in *Varanus komodoensis* (Auffenberg, 1978b), but rare in *V. bengalensis*.

Following (F). Pursuit of one individual by another (= *chase* of Gillingham, 1979; Vogel, 1979); chiefly in courtship. It is frequent in the courtship of *V. komodoensis* (Auffenberg, 1981a) and in an agonistic context in *V. salvator* (Vogel, 1979).

STRESS REACTION CLASS

Gular Extension (G). Enlarging the throat by both lowering the hyoid apparatus and inflating it with air, often accompanying a hiss (= *inflated throat* of Carpenter and Ferguson, 1978; Vogel, 1979). Common in most varanid species (Auffenberg, 1981a).

REPRODUCTIVE CLASS

Mount (M). One individual (in this context always the male) assumes a superior position on the other (female), with two or all four legs positioned on the opposite sides of the lower one (female).

Scratching (S). The male (in courtship context) scratches feebly with one foot, usually on the shoulder or middle of the back of the female (Fig. 1A), often when the latter is walking away. The identical acts occurs in *Varanus komodoensis* (Auffenberg, 1981a).

Chin Rubbing (C). Male in mounted position jerkily rubs his chin on the dorsal surface of the neck and/or

head of the female (Fig. 1B). This may be functionally related to the chin adpression of male snakes to the head or body of females (see Gillingham, 1979). It also occurs in *Varanus komodoensis* (Auffenberg, 1981a).

Head Twitch (HT). When mounted by a male, the female often twitches the head laterally in a jerky pattern. It is also common during the courtship of *V. komodoensis*. It may function as a sexual rejection signal, for (1) the male often follows this with vigorous chin rubbing, and (2) head twitching by the female after a long courtship sequence usually leads to the male walking away, or doing nothing (see below). Not observed in *Varanus komodoensis*. The "head jerk" described by Vogel (1979) in agonistic encounters of *V. salvator* is apparently not the same.

Tail Lift (TL). A mounted male tries to lift the tail base of the female with one of his hind feet by stroking upward with the claws (Fig. 1B). The identical act occurs in *Varanus komodoensis*.

MISCELLANEOUS

Do Nothing (N). While this act is intuitively neutral in effect, I agree with Hazlett and Bossert (1965) that for a respondent individual to "do nothing" may be a very meaningful act in many contexts. As will be shown below, it is apparently important in the courtship of *Varanus bengalensis*, for the signal of a female making no response to a courting male seems to encourage further courtship acts by the male.

Brachial Embrace (Br). In the agonistic context, this act (see Auffenberg, 1981b) is used by bipedal males in ritualistic combat, whereby both combatants clasp one another with their front legs. Its major function is to hold both individuals in a ventrally adpressed bipedal position during the wrestling phase. Brachial embrace of the female by the mounted male is also a conspicuous part of the later phases of courtship prior to copulation (Fig. 1C). As in the agon-

istic context, the primary function seems to be to immobilize the outer individual, in this case by pressing the front legs of the female to her sides and making it difficult for her to move forward out from under the mounted male. In a sense, it provides the same function as the partially immobilizing xiphiplastral ramming in land tortoises (e.g., *Geochelone*, Auffenberg, 1964, 1978a), the biting of female lizards on the neck by mounted males (Carpenter and Ferguson, 1978), and the laterally draped body of the male over that of the female during snake courtship (Carpenter, 1977).

Table 1 shows that the most intensive activity in captivity in Florida begins in early April and ends in late May, with peak activity during the middle of this period. At the same time, it is clear that the number of behavioral acts from courtship initiation to mounting decreases steadily from six to one as the season progresses, so that courtship is both more common and shorter later in the season. Coition was noted from as early as June 11 to as late as August 30. Eggs were laid from August 16 to October 3, with most during the height of the Florida rainy season (late August through September).

In Florida, captive-reared females laid their first eggs during their fourth year of life, though the males engaged in some courtship behavior (unsuccessfully) during the preceding year. Among mature adults, unsuccessful courtship was apparently due to the female slipping out from under the mounted male, caused mainly by the fact that the females were not being held tightly enough during the brachial embrace. Thus, the success in courtship is at least partially dependent on the size (= strength) of the males. Smaller males not only had difficulty immobilizing females with the brachial embrace, but their hind feet were too far forward to lift the tail base of the females prior to coition. In general, the larger the male in respect to the female, the better chance for successful courtship to have taken place.

Figure 1. Major behavioral acts comprising the reproductive class in *Varanus bengalensis*. (A) *Scratching* of the back of the female by the male, usually as she *walks away*. (B) When *mounted* the male *rubs chin* horizontally over neck and head of female as she *twitches head* back and forth. (C) The female is grasped in a *brachial embrace* by which her front legs are pinioned to the sides of her body by the front legs of the male. At the same time the male uses one of his hind feet to stimulate the female to lift the base of her tail by raking upwards with the claws.

TABLE 1. DATES AND NUMBER OF MALE BEHAVIORAL ACTS FROM INITIATION OF COURTSHIP TO THE FIRST MOUNTING ATTEMPT IN CAPTIVE *VARANUS BENGALENSIS*.

Inclusive Dates	Average Number Male Behavioral Acts	Number Observations
March 1-14	6.2 ± 3.1	21
March 15-30	5.1 ± 5.6	37
April 1-14	4.2 ± 2.5	59
April 15-30	2.7 ± 1.8	111
May 1-14	2.4 ± 2.0	63
May 15-30	1.7 ± 0.8	35
June 1-14	1.3 ± 1.2	16
June 15-30	1.0 ± 0.8	5
July 1-14	1.0 ± 0.0	1
July 15-30	0.0	0
August 1-14	1.1 ± 0.6	2
August 15-30	1.0 ± 0.0	1

Of 351 early courtship steps observed, 79.9% were initiated by the male when the female was lying still; 16.1% when the female was walking; 0.0% when she was running; and 4.0% when the activity of the female was nonclassifiable. Of the courtships observed, 3.6% resulted when a female walked in front of, or next to, the winning male of a ritual combat only moments before. Females often seemed to watch these contests and showed signs of increasing interest during the phase when one male performed a number of dominant acts against the other, usually moving closer to the interacting males at that time.

The length of time from courtship initiation to the first copulation attempts in 121 timed sequences varied from 3.2 to 44.4 sec ($\bar{X} = 25.5 \pm 22.2$ sec). Complete time *in copula* varied from 10.3 to 87.4 sec, and total courtship plus intromission varied from 20.8 to 123.1 sec ($\bar{X} = 101.8 \pm 60.3$ sec).

The following brief description is typical of courtship behavior in the early part of the season; later, as mentioned above, it tends to include fewer acts. In this case the courtship pattern passed through nine steps (in parentheses).

A basking female is approached from behind by an adult male. When approximately even with her shoulder, the male (1) flicks his tongue onto her occiput one time, her neck two times, and her tympanum region three times, as she (2) turns her head away. The male (3) places his chin on the frontal area of the female, pushing downward as he jerks his head from side to side (see Fig. 1). The female (4) slips away and the male (5) follows, (6) mounts her, (7) applies a strong brachial embrace, and (8) engages in a vigorous tail lift, to which she responds by (9) lifting her tail. Intromission occurs.

The modal number of steps in successful courtship is 10 (range = 3–16), with the longest sequences tending to be more unsuccessful—probably because it becomes apparent to the female that the male is not an appropriate partner. Short unsuccessful courtships are usually terminated by a female that becomes aggressive and/or runs away. Many unsuccessful

courtship sequences contain a series of repetitive acts occurring during the middle or near the end of the sequence, most often following step 5. On this basis I assume the most important discriminatory communication interchange in this species takes place during steps 1–5. It is for this reason that the following analysis concentrates on the acts of the first five steps. Because the male always initiates courtship behavior, acts 1, 3, and 5 are always those of the male, and acts 2 and 4 are those of the respondent female.

To determine the question of stationarity during the first five behavioral steps, the frequencies of each of the 20 acts noted in courtship were considered first. Table 2 shows that the probability of any act occurring is probably dependent on what had occurred previously, for within both male and female sequences certain patterns are more common in the earlier steps and others in the later ones. As examples of temporal dependence, male chinning and tonguing are notably reduced from steps 1 to 5, whereas walking away is increased. Sex is also obviously significant, for mounting, chinning, and scratching are only found in the male, whereas others (usually agonistic), such as indiscriminate biting, topping, head away, and head twitching are restricted to the female. While the remainder are used by both interactants, some are more common in males (brachial embrace, tail lifting, or tonguing), and others in females (head up, do nothing, or walking away). As pointed out under methods above, it is for this reason that the probabilities of specific step acts occurring cannot be stationary, and this important fact is taken into consideration in the analysis.

ADJACENT LOWER ORDER INTERACTIONS

The unconditional probabilities of the acts characteristic of one sex or the other are not constant with time during the courtship bout (Table 2).

Steps 1–2. In this initial phase of

TABLE 2. THE ACTS OCCURRING IN STEPS 1-5 OF 217 COURTSHIP BOUTS IN *VARANUS BENGALENSIS*.

Acts*																			
Steps	Bi	Br	C	F	G	HA	HD	Hi	HT	HU	M	N	S	TC	TL	To	Tp	Ts	W
1			38		1						47		8	1		122	1		
2	1	1			1	8	2		20	8		59		2		4		1	110
3		15	52	16							36		9		7	80	4	1	1
4	2				5	19		6	25	15		42		1	2	6			100
5			4							1	18	84	8	1	2	37	4		56

Note: Even-numbered steps were performed by respondent females and odd-numbered steps by initiator-males.

*Bi = Biting; Br = Brachial Embrace; C = Chin Rubbing; F = Following; G = Gular Extension; HA = Head Away; HD = Head Down; Hi = Hiss; HT = Head Twitch; HU = Head Up; M = Mount; N = Do Nothing; S = Scratching; TC = Tail Coil; TL = Tail Lift; To = Tonguing; Tp = Topping; Ts = Tail Slap; W = Walks Away.

courtship the male initiates the interaction in step 1 and the female responds in step 2. In general, the acts of steps 1-2 show much less variability than steps later in the courtship sequence. Of a total of 21 expressed act pairs in steps 1-2, only 12 are significantly more common than expected; however, in steps 2-3 there are 19 of 30 act pairs that are significantly more frequent than expected, in steps 3-4 there are 27 of 39 act pairs that are significantly higher than expected, and in steps 4-5 there are 19 of 27 act pairs that are significantly higher than expected. Thus, in the early phase of courtship there is low variability in the adjacent acts, which then increases during the middle phase and becomes lower again as the interaction becomes degraded prior to separation of the interacting individuals.

About one half of the significantly high step 1 male acts of the pair belong in the investigatory class (tonguing), and the remainder include only acts in the reproductive class (chinning, mounting, scratching). Male acts that are pair components not significantly different than expected include some of the reproductive class (mounting), some usually used in an aggressive context (gular extension, tail coil), or are investigatory (tonguing).

Respondent female acts (step 2) of act pairs significantly higher than expected

are "do nothing" acts usually used in an aggressive context (tail coil), or of importance socially (head away, head up, walking away), or (?) sexual acts (head twitch). Female acts included in act pairs not significantly different from expected are usually either used in an aggressive context (gular extension, biting, topping, brachial embrace), in a social one (head down), in a reproductive context (head twitch), or "do nothing."

Of those act pairs that occur significantly more frequently than expected, the most common (highest significance) include reproductive acts (chinning, mounting, scratching); the less common (lowest significance) include an investigatory component (tonguing).

Within the act pairs of significantly higher occurrence, there are only four different male acts—almost always in the reproductive class (chinning, mounting, scratching). However, there are seven different female respondent acts—almost twice the level of act variability than in the male acts in this initial stage of courtship. The usual female respondent act involves one of the head movement in the social class, thus apparently reflecting an emphasis by the female on agonistic rather than reproductive motivation at this early stage of courtship.

Steps 2-3. In this act pair the male's response (step 3) to step 2 of the female is

analyzed. First, the most obvious difference when compared with steps 1–2 is that the variety of acts included in steps 2–3 is increased by approximately 50% (30, as opposed to 21, acts significantly higher than expected). Second, the levels of significance of steps 2–3 are generally higher than those of steps 1–2, suggesting a closer association between the component acts of the act pairs than in steps 1–2. This is largely due to the apparently more specific response of the male to the often agonistic acts of the female. Third, the component female acts of the significantly higher act pairs are largely social, involving head movements (head down, away, up, twitch), “do nothing,” walking away, or aggressive or investigatory acts. Of the respondent male acts in the significantly high act pairs, approximately one half are investigatory (tonguing) and the other half are either in the reproductive class (chinning, brachial embrace, mounting, tail lift, scratching) or following. Thus, though the courtship tactic of the male has not changed appreciably from his initial step 1 phase, this, his first response to the female’s initial act, shows greater association with it.

The highest departures from expected values are found in act pairs, including female head movements, suggesting the highest association occurs between male responsive acts (often reproductive or investigatory) and female social indicators. The least significant act pairs (= lowest association between adjacent acts) comprise those in which the male is responding to the female walking away. Mounting becomes common, but is higher than expected only when the female has “done nothing,” though almost all female “do nothing” acts are followed by some form of male reproductive act.

Included within the significantly associated acts in steps 2–3 are eight different component acts and seven component responses. Thus, male act variation becomes almost doubled from step 1.

There is no obvious pattern in the act

pairs exhibiting less than expected frequencies, except that many contain agonistic acts. This suggests that these probably reduce the frequency of associative acts and that their expression tends to reduce the possibility of further communicative sequencing toward successful coition.

Steps 3–4. In this set of adjacent acts the female (step 4) is responding to the previous male act (step 3). It constitutes her second act in the courtship sequence. In steps 3–4, 39 act pairs occur in significantly higher frequencies than expected, so that variability in potentially communicative adjacent acts is again increased by about 30% from steps 2–3. I believe this part (steps 3–4 and nearby) to be the most important of the courtship sequence because (1) there is a gradual reduction in the number of significantly

Figure 2. Common behavioral acts in the first six steps of courtship, with acts significant at the 5% level shown as thick arrows and those significant at the 2% level shown as thin arrows. Steps 1, 3, and 5 are male acts; steps 2, 4, and 6 are female acts. Heavily bordered area includes those acts most commonly associated with successful courtship; lighter bordered area represents less commonly successful sequences.

associated acts in later steps, and (2) remaining steps are often repetitive.

Within the significantly high associations, about one half of the male component acts are reproductive, a third are investigatory, and a few are of the "following" type. Common associated female responding acts are walking away, social acts involving head movements, and aggressive types (biting, gular extension, topping, hissing). Female reproductive acts (head twitch) make their first significant appearance at this stage, but they are not associated with a broad range of male acts.

Unlike the two earlier steps, there is no clear grouping of act pairs into those that are very closely associated and others that are much less associated. All associations tend to be high, with the highest between the tail lifting act of the male and the corresponding respondent tail lift of the female prior to coition.

The female response to tonguing (investigation by the male), though variable, is usually a social act. She may also walk away, but the tonguing-walk away act pair has one of the lowest associations in the entire step 3-4 phase. Of probably high communicative importance is that the female rarely "does nothing" after being scratched, but walks away much more commonly than expected.

There are 7 different component male and 12 different component female acts in the highly significant act pairs. Thus variation in component acts of significantly high act pairs increased from the first to the second communicative act in both males (steps 1 and 3) and females (steps 2 and 4). The result is that the range of probably important communicative acts is greater at this stage than any preceding or succeeding ones.

Steps 4-5. In this set of adjacent acts the male acts for the third time (step 5) and is responding to a female act for the second time. Interestingly, there is a significant reduction in the number of associative acts from 39 in steps 2-3 to only 19. Thus communication potential, as

suggested by the breadth of associative acts, is greatly reduced, to the level of that of steps 2-3, illustrating a general deterioration of the interaction at this stage if mounting has not yet taken place.

The female acts of those significantly associated remain about one half social (head away, head up, walk away) or neutral (do nothing) (total 11), with about one quarter reproductive (tail lift, head twitch) and one investigative (tonguing). Female reproductive acts appear in significantly high frequency for the first time in the courtship sequence. Yet the male acts have not changed greatly, for about one half of the dominant elements among the significantly associated acts continue to be reproductive (mounting, scratch, chinning, tail lift), while investigatory acts (tonguing) are still common. However, a significantly larger proportion of socially "neutral" acts (do nothing, walking) is apparent, often associated with preceding similar or socially subdominant acts by the female. These high associations may reflect increasing loss of interest on the part of the male.

The highest level of association is found in the precopulatory tail lift act by both individuals (as in steps 2-3). Female aggressive acts remain much higher than expected, often followed by rather neutral acts by the male (walk away, tonguing, do nothing). The weakest associations are again seen in act pairs in which the female walked away, suggesting the male may become confused or disinterested. However, in general, the level of association between male and female acts remains high. During this phase there are 10 different female acts included in the significant associations and 8 in the following male acts.

HIGHER ORDER INTERACTIONS

I have shown in the preceding section that certain initiation and respondent acts are often associated in higher frequencies than expected on the basis of chance alone, and that the probabilities of specif-

ic associations change within a temporal and sexual framework, dependent to a large measure on the immediately preceding act. In this section I offer the results of analyses concerned with the probability of a certain act being expressed following a specific prior act of the same individual, or following a specific act pair of preceding acts by the same individual and its interacting partner. Analyses of higher order interactions consider three rather than two steps at a time.

Binomial tests for each of the 3-step sequences show that many of them differ significantly at the $\leq 0.05\%$ level (Table 3). Of the 81 possible three combinations in steps 1–3, only 20 occur significantly more frequently ($P = \leq 0.05$, binomial test) than expected. Of the remainder, there is a much lower association than expected in the 3-step acts $To_1N_2M_3$, $To_1W_2C_3$.

The fact that a complete list of observed and expected frequencies is available for all the steps in the courtship sequence permits the use of a chi-squared test to estimate the significance of the deviation from expected for each 3-step act. Pearson's chi-squared approximation was used to perform the chi-squared test (degrees of freedom $r-1 + c-1$).

The highly associated acts of steps 1–3 include female step 2 acts of walk away (W_2), do nothing (N_2), head twitch (HT_2), tonguing (To_2), and head away (HA_2). Of these, only the first three occurred with sufficiently varied male acts 1 and 3 to make chi-squared tests useful estimations of association. W_2 occurred 110 times, and when compared with the preceding (No. 1) and succeeding (No. 3) male acts, the analysis provided a χ^2 of 0.99 ($df = 10$). Thus the male act 3 occurs as part of the W_2 n-step sequence (C_3 , S_3 , F_3 , Br_3 , To_3 , and M_3) and is *not* dependent on W_2 , or on W_2 preceded by male acts C_1 , M_1 , or To_1 . The same results were obtained for C_3 and To_3 following C_1 Ht_2 ($df = 1$, $\chi^2 = 0.34$), and for C_3 , M_3 ,

TL_3 , and To_3 following MIN_2 , S_1N_2 , and To_1N_2 ($df = 6$, $\chi^2 = 5.45$). The same conclusions can be drawn for the transitions in steps 2–4: HA_3 , W_3 , To_3 , and N_3 are *not* dependent on To_2 or HT_1To_2 , HU_1To_2 , N_1To_2 , N_1To_2 , or To_1To_2 ($df = 9$, $\chi^2 = 1.71$); or HT_3 , W_3 , or N_3 after C_2 , Ht_1C_2 , N_1C_2 , or W_1C_2 ($df = 4$, $\chi^2 = 1.54$); or steps 4–5 here.

Thus, the courtship interaction data do *not* require more than 2-step transition probabilities to describe them; i.e., no more than secondary nonstationary Markov processes need to be postulated. The behavioral acts during courtship are evidently largely dependent on the immediately preceding acts. This conclusion is additionally strengthened by a series of G-tests (likelihood ratio test, see Oden, 1977), which are more conservative than χ^2 tests based on Pearson's method. The results for all significantly associated acts in all steps are shown in Figure 2.

DISCUSSION

Figures 3 and 4 diagram the conclusions of the analyses conducted on the courtship of *Varanus bengalensis*. Figure 3 diagrams the highly significant associations among the adjacent acts comprising the 10 most highly significant act pairs listed in Table 3. The number 10 is arbitrary, but includes what appear to be the most significant adjacent acts in each step to intromission. In that part of the diagram covering steps 1–2 it is obvious that the first act of the male in the courtship sequence is usually a reproductive type—usually scratching, mounting, or chinning—and that these are significantly associated with “do nothing” or walk away by the female. The male acts often lead to a rejection act (head twitch) by the female or other head movements, usually denoting social subordination (head away, head up). The latter are probably most important in the successful completion of the courtship sequence, for in the

Figure 3. Sequence diagram arbitrarily based on the ten highest 2-tuple act behavioral patterns. Circles represent male acts, squares female acts. Arrow widths represent proportion (in %) of associated act pairs in the steps shown.

TABLE 3. UNUSUALLY FREQUENT N-TUPLES FOR 217 LENGTH 5 COURTSIPS OF *VARANUS BENGALENSIS*.

Columns show starting step number, step acts (n-tuple), expected and observed frequencies, z-values, and one-tailed significance (binomial distribution). Only $P \leq 0.05$ deviations from expected are shown. See p. 541 for act abbreviations Step acts are arranged from most to least significant within each step number category.

Step		Frequency		z-value	Significance
No.	Act	Expected	Observed		
1	S-N	0.30	8	15.16	0.0066
	M-HA	0.65	3	14.16	0.0071
	M-W	6.28	29	13.77	0.0073
	C-HT	2.98	17	12.53	0.0080
	C-W	3.68	21	12.46	0.0080
	M-N	2.38	11	12.11	0.0083
	To-TC	1.12	2	9.67	0.0103
	To-HT	1.12	2	9.67	0.0103
	To-To	2.25	5	9.05	0.0111
	To-HA	3.37	6	7.07	0.0141
	To-HU	4.50	8	6.89	0.0145
	To-N	20.25	36	6.54	0.0153
	To-W	33.73	60	6.25	0.0150
2	HD-To	0.02	2	20.63	0.0049
	TC-To	0.02	2	20.63	0.0049
	HA-C	0.07	2	20.10	0.0050
	HU-C	0.07	2	20.10	0.0050
	To-To	0.07	4	16.72	0.0060
	HT-Br	0.28	3	16.39	0.0061
	HU-To	0.22	6	15.54	0.0064
	HT-C	0.65	7	14.43	0.0069
	HT-To	0.74	8	13.64	0.0073
	N-TL	1.60	6	11.73	0.0085
	N-To	3.80	14	11.14	0.0090
	N-M	4.62	17	11.05	0.0090
	N-C	5.71	21	10.99	0.0091
	HA-To	0.18	5	10.53	0.0095
	W-S	4.56	9	7.71	0.0130
	W-Br	4.56	9	7.71	0.0130
	W-F	3.11	16	7.50	0.0133
3	W-M	8.61	17	7.49	0.0134
	W-To	19.26	38	7.36	0.0136
	TL-TL	0.07	2	20.10	0.0050
	F-N	0.15	2	19.27	0.0052
	M-G	0.33	1	17.40	0.0058
	M-Hi	0.33	1	17.40	0.0058
	Br-N	0.21	3	16.57	0.0060
	S-W	0.17	4	16.30	0.0061
	TL-N	0.13	4	16.04	0.0062
	Br-W	0.27	4	15.87	0.0063
	C-Tp	0.48	2	15.83	0.0063
	C-HU	0.48	2	15.83	0.0063
	Br-HT	0.55	8	14.67	0.0068
	F-W	0.88	12	14.26	0.0070
	M-N	1.00	0	13.44	0.0074
	To-HT	0.74	2	13.13	0.0076
	To-Tp	0.74	2	13.13	0.0076
	To-HU	0.74	2	13.13	0.0076
	To-Bi	0.74	2	13.13	0.0076
	To-Hi	0.74	2	13.13	0.0076
	M-W	4.15	25	12.56	0.0080
	C-HT	3.35	14	11.63	0.0086

TABLE 3. CONTINUED.

Step		Frequency		z-value	Significance
No.	Act	Expected	Observed		
4	C-W	5.75	24	11.44	0.0087
	C-N	1.44	6	11.34	0.0088
	To-To	1.84	5	10.40	0.0096
	To-HA	6.64	18	9.57	0.0104
	To-N	6.64	18	9.57	0.0104
	To-W	16.32	28	6.27	0.0160
	S-N	0.12	3	5.36	0.0187
	TC-TL	0.02	2	20.63	0.0049
	Bi-N	0.02	2	20.63	0.0049
	Tp-W	0.06	3	17.71	0.0056
	G-M	0.07	3	17.65	0.0057
	HA-N	0.11	4	16.55	0.0060
	HA-W	0.26	3	16.51	0.0061
	HT-N	0.35	3	15.96	0.0063
	HT-W	0.35	3	15.96	0.0063
	Hi-To	0.03	10	15.58	0.0064
	To-To	0.11	4	15.55	0.0064
	HU-N	0.35	5	15.30	0.0065
	HT-S	0.46	4	15.06	0.0066
	HT-C	0.46	4	15.06	0.0066
	HA-M	6.44	5	15.00	0.0067
	HU-To	0.69	10	14.46	0.0069
	HA-To	0.88	10	14.26	0.0070
	HT-M	1.27	11	13.67	0.0073
	Hi-W	0.03	3	12.91	0.0078
	N-N	7.55	39	12.05	0.0083
	N-M	0.39	2	11.84	0.0084
	W-S	1.38	3	9.76	0.0102
	W-To	5.53	12	8.31	0.0120
	W-N	17.05	37	8.05	0.0124
	W-W	20.28	44	8.03	0.0126
1	To-W-F	0.64	8	14.47	0.0069
	S-N-M	2.30	6	9.95	0.0101
	M-N-C	3.61	9	9.39	0.0107
	M-W-C	3.83	8	8.27	0.0121
	C-W-F	2.93	4	7.66	0.0131
	C-W-M	3.22	6	7.57	0.0107
	M-W-F	4.50	8	6.89	0.0145
	M-W-M	4.51	8	6.89	0.0145
	M-W-To	10.01	7	6.82	0.0546
	To-N-TL	8.51	6	6.72	0.0149
	To-W-Br	4.93	8	6.10	0.0164
	To-HA-To	3.81	6	5.91	0.0169
	To-N-C	7.34	11	5.20	0.0192
	To-N-To	8.52	12	4.49	0.0223
	To-W-S	4.94	4	3.83	0.0261
	M-W-S	2.41	3	3.61	0.0278
	C-W-S	1.70	2	3.13	0.0319
	C-HT-C	3.41	4	2.50	0.0400
	C-HT-To	6.82	8	2.35	0.0426
	To-W-To	20.70	24	2.07	0.0483
2	To-To-To	0.33	4	15.74	0.0063
	To-To-HA	0.92	2	11.46	0.0087
	HT-To-HA	1.10	3	11.45	0.0087
	HT-C-HT	1.64	4	10.21	0.0098
	N-To-HA	3.23	8	9.45	0.0106

TABLE 3. CONTINUED.

Step		Frequency		z-value	Significance
No.	Act	Expected	Observed		
3	W-C-HT	4.30	3	7.83	0.0128
	W-To-HA	7.41	13	7.78	0.0129
	W-M-N	2.68	5	7.57	0.0132
	N-M-W	7.19	12	4.87	0.0205
	N-C-HT	4.88	7	4.77	0.0210
	HT-C-W	2.69	2	3.33	0.0300
	W-To-HA	6.75	6	2.02	0.0495
	W-To-W	10.50	12	1.92	0.0521
	W-M-W	8.91	8	1.77	0.0565
	Br-HT-M	1.00	5	13.33	0.0075
	G-W-N	2.39	17	13.05	0.0077
	To-W-N	1.77	11	13.00	0.0077
	C-HT-M	2.04	6	10.70	0.0093
	F-W-W	6.23	9	9.86	0.0101
	To-W-N	2.79	6	8.68	0.0115
	M-W-W	4.53	10	8.61	0.0116
	To-W-To	2.05	4	8.13	0.0123
	To-N-N	2.57	5	8.10	0.0123
	To-HA-To	4.89	10	7.98	0.0125
	C-W-W	3.41	6	7.00	0.0143
	M-W-To	4.51	9	2.57	0.0389
	Br-HT-S	0.40	2	2.39	0.0418

diagram representing steps 2-3 these same acts are highly significantly associated with further reproductive acts by the male in step 3. Note that "do nothing" on the part of the female is also significantly associated with further male reproductive interest. Investigatory, social, warning, and sexual rejection frequently lead to further investigation by the male. However, rejection by the female is often ignored, with the male pressing the courtship further. During steps 3-4 success is often achieved via mutual tail lifts in both interactants. However, tail lifts by the male more frequently lead to "do nothing" or social acts, including topping which is clearly a dominant act. It also frequently elicits warning behavior and rejection in the form of the female walking away. In steps 4-5 we see that female warning acts are common, though these seem largely ignored by the male, for tail lifting and mounting are closely associated with

them. On the other hand, female weaponry use leads to "do nothing" by the male. Sexual rejection by the female also leads to a breakdown of the sequence at this stage (though not in earlier steps). The male reacts similarly to female social acts—exactly the opposite of his reaction in steps 2-3.

Figure 4 is a diagrammatic conclusion of the more conservative G-tests. Again, the significant male step 1 acts are chin-ning, mounting, or tonguing. These are variously, but significantly, associated with the respondent female acts of head twitch, walking away, or doing nothing. If the female tongues the male at this stage, or turns her head away, there is a very strong association with a responding tonguing act by the male in step 3. There is also a significant association between his tonguing act and her preceding head twitch, and her possibly doing nothing. His chin-ning in step 3 is also closely associated with her doing nothing in step 2.

Figure 4. Behavioral act sequence summary based on G-test analysis.

Tail lifts and mounting by the male at this stage are also closely associated with her doing nothing. Tonguing by the female in step 4 is associated only with similar behavior on the part of the male. Head twitching by the female is associated with his chinning. Her tail lift is closely associated with his previous similar activity. There is a very strong association between her walking away in this step and the male's previous mounting attempt. According to G-test analysis, successful courtship, in that it leads to coition, is most probable in the sequence $M_1 N_2 Tl_3 Tl_4$, or the same sequence be-

ginning one step later. Thus, the social head movements, which seemed so important on the basis of highly significantly associations in Pearson's χ^2 estimates, are here indicated to be less significant in the successful completion of courtship, though undoubtedly of considerable importance in establishing rapport between the interacting individuals.

Because of the data requirements, the G-tests included more courtship sequences during the late part of the courtship season than the data shown in Figure 3. This may help explain the more simplified act system suggested, as well

as the short, direct sequence leading to successful coition. These short, usually more successful, sequences occur later in the same season, after attainment of familiarity between individuals seems to have been accomplished during earlier courtship sequences of the type illustrated in Figure 3. Specific acts frequently deleted in these later phases are those that fall into the agonistic and social classes.

Most female acts in unsuccessful sequences are agonistic and may include actual weaponry use (biting, tail slap). Continued association with particularly large, powerful males able to restrain the aggressive females brings about a reduction in female agonistic behavior. As I pointed out in my earlier study of *Varanus komodoensis* (Auffenberg, 1981a), the large size of this species, plus the aggressive nature of the females and their capability of inflicting serious wounds and of even killing smaller individuals, have undoubtedly contributed to the development of a reproductive behavioral strategy that stresses courtship throughout the year among the mature male and female pairs who share the greater part of the same foraging area (Auffenberg, 1978b). As a result, familiarity is assured among individuals potentially dangerous to one another, thereby reducing stress during the actual breeding period itself. This system apparently tends to develop a long-term pair bond, for individuals of *Varanus komodoensis* seem to inhabit the same area for many years (Auffenberg, 1978b, 1981a). The probability for courtship success is thus enhanced by what amounts to an almost year-round courtship with otherwise very aggressive and powerful females. The same strategy in its basic format is seen in the present study of *Varanus bengalensis*. Though it is not known what role, if any, pair-bonding plays in this species in the wild, it is obvious that a long association (several months in captivity) effected through repeated courtship by the male is usually necessary for successful mating.

The entire social system of *Varanus bengalensis* seems to select the most powerful males and females for ultimate breeding. I base this on the fact that visual displays intimating social rank are of much less importance in varanids than in agamids or iguanids. On the other hand, warning and actual weaponry use are common components of even adolescent interactions (Auffenberg and Ganci, in press). Secondly, unlike lizards in other families, female varanids are much more apt to engage in agonistic displays and use their teeth and tail more often than do males of the same species (Auffenberg, 1981b). This means that male *Varanus bengalensis* interested in copulation must be able to so completely restrain the females that they remain uninjured after breeding. The females show an inclination to breed with the males that are capable of doing this. Third, male varanids engage in a test of strength, the evident purpose of which is to determine the one that has the greater ability to physically manipulate but not harm the other. This is exactly what is required in courtship within the same species (harmless manipulation), for at this time it is the females that must be restrained. Finally, the females themselves often select by overt behavior those males most able to restrain and otherwise manipulate the other; i.e., those males most probably capable of restraining the females themselves. The entire system is based on proven performance, not on bluff, and seems analogous to that in many ungulates (see Wilson, 1975). I believe it is the basis for much of what we know occurs in male-male combat and male-female courtship of snakes (for review see Carpenter and Ferguson, 1978).

The emphasis of the female *Varanus bengalensis* on agonistic behavior during the early part of the courtship season and much of each courtship bout may help explain the fact that the male acts are associated only with the preceding female acts and in no way dependent on his previous acts, either separately or in combination with those of the female. If

my interpretation is correct, the female is less predictable in her response and certainly potentially dangerous in view of her tendency to become aggressive. The latter is clear, while the former is suggested by the greater variability of her responses and the lack of significant association of most of her acts with any specific male initiating act. In view of this, the male probably determines most of what he does during courtship on the basis of the female's immediately previous act alone. His courtship behavior is not rigidly programmed, but remains flexible to contend with more varied reciprocal acts of the female.

ACKNOWLEDGMENTS

This work was made possible through the support of the Florida State Museum, which purchased most of the equipment used; the New York Zoological Society, which provided the greenhouse facilities and funds for the field work in 1974 when some of the individuals were captured; and the United States Customs Office, which permitted use of individuals confiscated as the result of illegal shipments.

LITERATURE CITED

- AUFFENBERG, W. 1964. Notes on the courtship of *Geochelone travancorica* (Boulenger). J. Bombay Nat. Hist. Soc., **61**: 247-253.
- . 1978a. Courtship and breeding behavior in *Geochelone radiata* (Testudines: Testudinidae). Herpetologica, **34**: 277-287.
- . 1978b. Social and feeding behavior in *Varanus komodoensis*. pp. 301-331. In N. Greenberg and P. MacLean (eds.), Behavior and Neurology of Lizards, Natl. Inst. Mental Health.
- . 1979. Intersexual differences in behavior of captive *Varanus bengalensis* (Reptilia, Lacertilia, Varanidae). J. Herpetol., **13**: 313-315.
- . 1981a. Behavioral Ecology of the Komodo Dragon (*Varanus komodoensis*). Univ. Presses of Florida, Gainesville, pp. 1-406.
- . 1981b. Combat behaviour in *Varanus bengalensis* (Sauria: Varanidae). J. Bombay Nat. Hist. Soc., **78**: 54-72.
- AUFFENBERG, W., AND S. GANCI. In press. Behavioral cues and displays in adolescent *Varanus bengalensis*. J. Herpetol.
- CARPENTER, C. C. 1977. Communication and displays of snakes. Amer. Zool., **17**: 217-223.
- CARPENTER, C. C., AND G. FERGUSON. 1978. Variation and evolution of stereotyped behavior in reptiles, pp. 138-308. In C. Gans and D. W. Tinkle (eds.), Biology of the Reptilia. Vol. 7, Ecology and Behavior. London, Academic Press.
- COOPER, W. E. 1977. Information analysis of agonistic behavioral sequences in male iguanid lizards, *Anolis carolinensis*. Copeia, **1977**: 721-735.
- GILLINGHAM, J. C. 1979. Reproductive behavior of the rat snakes of eastern North America, Genus *Elaphe*. Copeia, **1979**: 319-331.
- GREENBERG, N. 1977. An ethogram of the blue spiny lizard, *Sceloporus cyanogenys* (Reptilia, Lacertilia, Iguanidae). J. Herpetol., **11**(2): 177-195.
- HAZLETT, B. A., AND W. H. BOSSERT. 1965. A statistical analysis of the aggressive communication systems of some hermit crabs. Anim. Behav., **13**: 357-373.
- JENSSEN, T. A. 1977. Display diversity and anoline lizards and problems of interpretation, pp. 269-285. In N. Greenberg and P. MacLean (eds.), Behavior and Neurology of Lizards, Natl. Inst. Mental Health.
- MERTENS, R. 1942. Die Familie der Warane (Varanidae). Abh. Senckenb. Naturforsch. Ges., Pts. 1-3: 1-391.
- ODEN, N. 1977. Partitioning dependence in nonstationary behavioral sequences, pp. 203-220. In B. A. Hazlett (ed.), Quantitative Methods in the Study of Animal Behavior. New York, Academic Press.
- VOGEL, P. 1979. Zur biologie der Binseswarens (*Varanus salvator*) im Westjavanischen Naturschutzgebiet Ujung Kulon. Ph.D. dissertation, Univ. Basel, pp. 1-139.
- WEAVER, W. G., JR. 1970. Courtship and combat behavior in *Gopherus berlandieri*. Bull. Florida State Mus., Biol. Sci., **15**(1): 1-43.
- WILSON, E. O. 1975. Sociobiology: The new synthesis. Cambridge, Harvard Univ. Press, pp. 1-607.

Display Behavior of Two Haitian Lizards, *Anolis cybotes* and *Anolis distichus*

THOMAS A. JENSSEN¹

ABSTRACT. The display repertoires of two common and frequently sympatric congeners, *Anolis cybotes* and *A. distichus*, were quantitatively analyzed. The displays for each species were moderately stereotyped, being consistent within a population as well as shared by conspecifics between populations. Comparisons between the two species showed completely different signature display patterns, and a striking difference in the size of their respective repertoires. *Anolis cybotes* has only a single species-unique signature display. In contrast, *A. distichus* has six display types, some with unusual characteristics, and the largest repertoire yet described for an anoline species. An evolutionary scheme is provided, giving the hypothetical steps required to derive the display-rich *distichus* repertoire, and some socio-ecological factors which may be selective for this enlarged repertoire.

INTRODUCTION

Hispaniola supports a complex *Anolis* fauna of at least 34 species (Williams, 1976), with 20 of those species having been described since Cochran's (1941) checklist appeared. Many of the newly recognized taxa are cryptic species, being distinguished primarily by their electrophoretic profiles, collection localities, and a few subtle external characteristics, while others have restricted distributions and were discovered through new and more numerous sampling sites.

With the proliferation of anoline forms, the cybotoid and distichoid species have had their share of revision. *Anolis*

cybotes has recently yielded a cryptic species, *Anolis marcanoi* (Williams, 1975), which has been established to a large measure by electrophoretic criteria (Webster, 1975). Another cybotoid, *Anolis whitemani*, was separated by Williams (1963) from its sibling species *A. cybotes* with whom it shares almost identical albumin immunological characteristics (Wyles and Gorman, 1980); now based on more collection data, *A. whitemani* has been subdivided into three subspecies (Schwartz, 1980). The Hispaniolan *Anolis distichus* represents as many as 12 subspecies (Schwartz, 1968), with the intraspecific nature of some of the forms still in question (Williams, 1977a). *Anolis brevirostris*, a distichoid species, has been recently divided into four cryptic species (Arnold, 1980; Williams, 1976), chiefly through the electrophoretic studies of Webster (1977) and Webster and Burns (1973).

The continued discoveries of sibling species, cryptic species, and extensive variation within species have elicited an exciting and in some ways perplexing perception. *Anolis* systematics of the Hispaniolan fauna seems to become less stable with increased information, and the evolutionary events which shaped the fauna become even more enigmatic with the improved resolution of the taxonomic picture. To achieve any measure of sense to the emerging scene, more morphological, karyotypic, electrophoretic, ecological, and behavioral data will be needed. As Williams (1977b: 135)

¹Biology Department, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24060, U.S.A.

points out, even the most simple question of how many anoline species there are on the island defies an answer; too frequently one is not sure where one species ends and another begins.

In this paper, I present the display repertoires of important representatives of the cybotoid and distichoid groups, *A. cybotes* and *A. distichus dominicensis*. The display repertoires of these two sympatric species are quite different in size, with one being the smallest possible and the other the largest yet described for *Anolis*. The signature displays of *A. cybotes* and *A. distichus* are quite different, while the displays of both species show some remarkable and unusual qualities. These display descriptions serve as a baseline study for future comparative behavioral investigations of Hispaniolan anoles. The presented display data should aid in taxonomic decisions as well as possibly assisting evolutionary interpretations of the genus.

MATERIALS AND METHODS

CYBOTES SAMPLE

Sixty-two displays were filmed in the field. Five of these displays were from a male in Petion-ville, and 57 were by seven males from a population 2.5 km south of Montrouis on Route 100 (Fig. 1). Films were taken of males during undisturbed social activities and during elicited aggressive encounters. The latter were created by: 1) releasing an unrestrained intruder in a resident's territory and 2) placing a male confined within a transparent cubicle into an occupied territory.

DISTICHUS SAMPLE

One hundred ninety-three displays were filmed. Of these, 29 were filmed in the field; 23 were from three males in Petion-ville and six from males in Montrouis. These displays were given

within naturally occurring social contexts.

Five males from Montrouis had 164 displays recorded in my lab. The lizards were housed separately in large enclosures ($1.3 \times 0.7 \times 0.3$ m) containing simulated natural habitat and one to two females. The enclosures were lit with fluorescent lights (12L:12D photoperiod) and maintained at 23 to 25 C. At the time of filming, a subject was transferred to a filming enclosure identical to the holding enclosures. After a day or more of residency, the subject's activities were filmed from a blind with the added illumination from two quartz iodine floodlights. Each subject initially displayed after perch changes (usually non-directed) and when interacting with resident females. Later they displayed during encounters with a released male intruder.

TECHNIQUES

All films were taken with a Nizo S80 super 8 movie camera, using a film advance setting of 18 fps. Only those displays known to be complete and of good resolution were used in the analysis. The resulting sample was analyzed frame-by-frame with a Kodak MFS-8 Ektagraphic projector. Details of the analysis methods are given by Jenssen and Hover (1976).

To analyze the complex bob patterns, I arbitrarily divided each kind of stereotyped head bob pattern (display type) into units. Odd-numbered units were chosen to give the duration of a bob and even-numbered units delineated the duration of an inter-bob pause. For some odd-numbered units, a single bob was delineated by a unit. For the other odd-numbered units, a series of uninterrupted bobs comprised the unit; for such odd-numbered units, a bob count was also made. In addition, counts were made of the number of dewlap extension-retractions (pulses) which followed most of the displays.

Quantification of the data was aided by

Figure 1. Collecting areas in Haiti, indicated by asterisks.

computer analysis. All data of a display were punched on an IBM card; these included: 1) displayer's location; 2) displayer's species; 3) displayer's individual identification if known; 4) display type; 5) social context if known; 6) frames of film for each unit; 7) number of bobs for units depicting a bob series; 8) number of dewlap pulses if present; and 9) whether filmed in the field or lab. An IBM 370 computer then: 1) converted fps to real time for each unit; 2) computed descriptive statistics (mean, standard deviation of the mean, standard error of the mean, and minimum and maximum values) for each display variable; 3) calculated the coefficient of variation (standard deviation $\times$ 100/mean) for each variable; and 4) performed tests of significance.

DAP GRAPHS

The visual summaries resulting from the display analysis appear as display-action-pattern graphs (DAP graphs). These graphs show head amplitude movement along the y-axis and the passage of time along the x-axis. Thus a quick up-and-down head bob will look like a "spike" when graphed, and a bob which is held before the head is lowered will have a "plateaued" appearance. If the dewlap is extended during the display, it appears on the DAP graph under the head bobs; the dewlap extension is shown as a downward vector along the y-axis, dewlap retraction as an upward vector along the y-axis, and passage of time follows the x-axis.

STATISTICAL TABLES

Tables enumerating the descriptive statistics and statistical test results are available from the author and the Department of Herpetology, Museum of Comparative Zoology upon request.

RESULTS

SIGNATURE DISPLAY PATTERN-CYBOTES

Anolis cybotes has but a single stereotyped head bob display in its repertoire. This display pattern is used during non-directed displaying while a male patrols his territory; during male-female courtship, copulatory, and postcopulatory displaying; and during male-male agonistic disputes. The head bob pattern is produced by action of the forelimb and neck muscles (Fig. 2). The bobs of the signature display are rapidly sequenced and of relatively low amplitude. Without benefit of frame-by-frame analysis, the display does not appear to have much pattern to it. However, once the pattern is known, it is readily recognized and quite predictable.

The *cybotes* signature display is composed of two sequences of bobbing (the A and B Elements), with the second sequence repeated during some performances (Fig. 3). The A Element begins with three large bobs (Units 1, 3, and 5). Occasionally a fourth was included (Fig. 4B). This variation was performed by five of the eight subjects and involved 23% of the displays. The A Element concludes with a variable number (3–9, $\bar{x}$ = 7) of small rapid bobs (Unit 7). If a male began a signature display, it always stopped at the end of an element. Usually the display concluded after a B Element, but 12 displays (18%) finished after the initial A Element (Fig. 5).

The B Element is similar to the A, but tends to have one less large bob and fewer smaller rapid bobs (Fig. 3). The B Element is initiated with two large bobs (Units 9 and 11); occasionally two lizards

of my sample performed three (16% displays with a B Element) (Fig. 4C). The concluding series of small bobs (Unit 13) were variable in number (2–7) and averaged four bobs.

A large number of the displays (48%) ended with the first B Element. Some displays, however, concluded after the B Element was repeated two, three, or four times (Fig. 5). Although based on anecdotal observation, it seems that the more aroused a male is, the more likely he will repeat the B Element within a given display. There was no statistical difference in unit durations or number of bobs when comparing between B_1 - B_4 Elements.

The Petion-ville signature displays possessed the same pattern as those by males outside Montrouis in the north. The Wilcoxon Two-Sample test (Sokal and Rohlf, 1969) showed no significant difference between any of the comparable display variables from the two localities. Of interest is the display pattern of two *A. cybotes* males (Fig. 4) filmed during a 1970 trip to Etang Saumatre in the Cul-de-Sac area east of Port-au-Prince (Fig. 1). Their pattern also falls within the signature display characteristics of the males from the Montrouis and Petion-ville regions. This suggests that the *A. cybotes* in at least central Haiti have a fairly stable display pattern across geographic localities.

Figure 2. Minimum (stippled) and maximum (solid) amplitude during head bob displaying by *Anolis distichus* and *Anolis cybotes*.

Figure 3. Summary DAP graph of *Anolis cybotes* signature display. Upper edge of upper solid block shows head amplitude change (y-axis) through time (x-axis), upper edge of stippled block shows head amplitude changes of optional introductory movements, and lower edge of lower solid block shows dewlap extension through time.

Figure 4. Examples of different *Anolis cybotes* head bob behavior. A. Head dipping sequence with intervening quiescent periods. B. Stippled area indicating where a major bob of the A Element was performed as two. C. Stippled area indicating where the usual two major bobs of the B Element are performed as three. D. DAP graph of the signature display from near Etang Saumatre.

Figure 5. Frequency of filmed *Anolis cybotes* signature displays which ended after a particular element.

DISPLAY STEREOTYPY

The first step in a quantified analysis of ritualized behavior is to ask to what degree the behavior is stereotyped. To answer this, a coefficient of variation (CV) is computed for each of the behavior's variables. The CV is an index of variability, where CV values increase as variability increases. Therefore, it can also be used to gauge relative stereotypy, where the smaller the CV, the greater the stereotypy. Most quantified ritualized behaviors which Barlow (1977: 103) found reported in the literature have CV values within a range of 15 to 35%. From this Barlow suggested that behavioral components having CV values less than 15% should be considered highly stereotyped. Test results in this study show that *cybotes* unit durations and number of bobs within Units 7 and 13 fall primarily

within the 15 to 35% range.¹ Therefore, the signature display is classified as stereotyped, but certainly not devoid of variability.

The next question is how this variability is partitioned within the sample. To do this a one-way Analysis of Variance (ANOVA) (Sokal and Rolf, 1969) was used to compute total variance and then to partition the proportion of variance attributed to comparisons between lizards (among component) and that found between displays of the same lizard (within component). Test results show that most of the variance appears in the within component. This means the lizards of the sample tended to share a common range of variability for each unit duration and number of bobs. There was little indication of highly stereotyped individual-unique display characteristics.

The ANOVA (F value) did find significant differences for several variables when comparing between lizards. However, when Duncan's Multiple Range test (Steele and Torrie, 1960) was applied to the display variables, the results showed broadly overlapping variances between lizards, with no lizard or group of lizards being significantly different from the rest of the lizards for any given variable. Again, there was no indication of individual-unique display characteristics.

OTHER BOB MOVEMENTS

When a male was strongly aroused, as during territorial defense, he sometimes prefaced his signature display with an introductory "head roll." This head movement was not particularly stereotyped and consisted of a moderate to slow raising and lowering action of the head (Fig. 3, stippled area adjacent to A Ele-

¹ Tables of descriptive statistics and test results of the display analyses are available upon request from the author and the Department of Herpetology, Museum of Comparative Zoology.

ment). This is a common introductory movement to stereotyped displays of many anoles (Jenssen, 1979).

Another bob movement was also seen to occasionally precede the signature display. This was a quick inverted bob, best described as a "head dip" (Fig. 3 and 4A). Head dipping was also performed independently of the signature display. Solitary males were frequently seen head dipping as they took a new perch or when they appeared to be surveying their surroundings. The dips came in long series, with each separated by an extremely variable time interval (2–30 sec). If this behavior has a social significance, it very likely is not completely emancipated from its original nonritualized source behavior. I am guessing the noncommunicatory function of the head dip behavior is to enhance visual discrimination of distant objects. Head dipping is not performed by the other *Anolis* species I have studied, which leads me to suspect the behavior is unique to *A. cybotes* and possibly its close relatives.

DISPLAY REPERTOIRE-DISTICHUS

In contrast to the single display repertoire of *A. cybotes*, *A. distichus* exhibits a set of six display types (*a*, *A*, *B*, *C*, *D*, and *E*). This is the largest repertoire yet reported for an anoline species. *Anolis limifrons* had the largest previously known repertoire which contained five different display patterns (Hover and Jenssen, 1976). Below is the display-by-display description of the *distichus* repertoire.

TYPE *a* DISPLAY

Type *a* display is not represented by a single display pattern, but by three (Fig. 6). It is likely that the list could be expanded with the observation of more lizards. The criteria used to lump the patterns under the Type *a* category were: 1) extremely low bob amplitude (approx. $\frac{1}{3}$ that of other display types); 2) very short duration (< 2 s); and 3) a general similarity between the patterns which in

no way resembled the other display types.

The *a* displays are simplistic in comparison to the other five display types, giving me the impression that they represent display fragments. These related patterns are composed of 6 to 7 quick, simple bobs which are really more like nods. The low amplitude nods associated with this display type were produced solely by the neck muscles. Dewlap pulses following head bobbing ranged from one to seven, with 12% of the displays having no pulses. On occasion, the *a* displays were not completed, especially the a_2 and a_3 displays.

Stereotypy was fairly strong for most of the *a* units, especially when considering how short the unit durations are. The coefficient of variation values for many of the units fall close to or within Barlow's

Figure 6. Summary DAP graphs of three versions of the Type *a* display by *Anolis distichus*. Numbered boxes immediately under each DAP graph indicate duration and identity of bob and inter-bob units.

(1977) 0–15% range for highly stereotyped behavior.

All but three *a* displays were filmed in the lab. The context was usually associated with exploration; mostly solitary males recently released into a new enclosure performed the *a* displays and, in this situation, the displays were non-directed. The other context for the *a* displays was then two males were in close proximity to each other (< 10 cm). At this time five such displays (11% of the sample) were recorded. They were performed by only one member of the pair. It is very likely the *a* display carries a connotation of uncertainty or subordination.

TYPE A DISPLAY

Type A display is the species' signature display and is represented by a single consistent pattern (Fig. 7A). The pattern is composed of 11 units. Three units delineate single bobs (Units 1, 5, and 9) and three units contain multi-bob series (Units 3, 7, and 11). Separating the bob units are inter-bob pauses. Thus all odd-numbered units depict bobs and even-numbered units are pauses (Fig. 7A). In this display type and in display types B, C, and D, one sees a construct of single bob units alternating with multi-bob units.

All the bobs of the Type A display are quick up and down movements which graph as "spikes." Head amplitude movement of these bobs is created by simultaneous use of neck and limb muscles. In over half the A displays all four limbs came into play (Fig. 2). The use of four legs rather than just forelimbs seems to relate to increased arousal of the displayer. Although anecdotal for *A. distichus*, this correlation has been more firmly established in *Anolis opalinus* (Jenssen, 1979). Four-legged bobs were even more frequently incorporated into the *distichus* Types B, C, D, and E displays.

As is true of all of the *distichus* display types, dewlap pulsing following the head

bob sequence was the usual case, the number of pulses being variable. For 82% of the A displays, concluding dewlap pulses were performed; when present, they ranged from 1–8 pulses.

Display stereotypy was moderate, with 9 to 11 of the 14 display variables (excluding dewlap pulses) having their CV values within the 0 to 35% range for stereotyped behavior.

Comparisons of the Type A display pattern were made between the Montrouis and Petion-ville samples. The two summary DAP graphs are very similar (Fig. 7A_n and 7A_s). When comparing homologous display variables for the two localities, Wilcoxon's Two-Sample tests (Sokal and Rolf, 1969) indicated statistically significant differences for many unit durations and bob numbers within multi-bob units. Thus there are inter-population differences, but these divergences are minor on a qualitative level. They have not produced two different display patterns since bob cadence and the shape of the bobs are shared by the two populations. An A display by any of the sampled lizards fits the summary pattern.

Besides comparing between populations, I also compared the A displays between individuals within a population by using the 42 A displays of five lab-held males from Montrouis. One-way ANOVA tests showed significant differences ($P < 0.05$) for all but three display variables. To see if one or more lizards had display characteristics distinctly different from the rest of the sample, a Duncan's Multiple Range test was also used to compare among lizards. Test results showed there were significantly different groups of subjects for all even-numbered units and Unit 3. This same generalization is reflected in the partitioned variance, where those variables which reflected individual-unique differences (e.g., even-numbered units and Unit 3) would also tend to have most ($\geq 60\%$) of the partitioned variance in the among group category.

Figure 7. Summary DAP graphs of *Anolis distichus* Type A and B displays. A_n and B_n are the patterns from Montrouis in north and A_s and B_s are from Petion-ville in the south. Numbered boxes immediately under each DAP graph indicate duration and identity of bob and inter-bob units.

Examples of interindividual display differences are given in Figure 8; these DAP graphs are summaries for three males for which I had the most displays. The display durations for the three males were made comparable by computing unit durations as a percentage of the total display duration. One sees that the position of homologous bobs between lizards' displays is somewhat variable and the average number of bobs performed for the multi-bob units also differs between lizards. Yet it is evident that these DAP graphs are all Type A displays and cannot be confused with any of the other *distichus* display patterns (e.g., the Type B displays also depicted in Fig. 8).

The Type A display is considered the species' signature display by previously defined criteria (Jenssen, 1978). First, it is a well-defined stereotyped pattern shared by the population. Second, it was the most frequently performed display type; from the film record there were 60 A, 43 a, 29 B, 28 E, 18 C, and 14 D displays. Last, it occurred during non-directed displaying associated with territorial patrol. The Type A display, as expected of the signature display, appeared in all other major social contexts: during courtship, copulation, and the immediate post-copulatory period, and during aggressive interactions between males.

TYPE B DISPLAY

Type *B* display pattern was quite consistent and closely resembled the Type *A* display, the primary difference being the shape and duration of the single bobs of Unit 1 and Unit 5 (Fig. 7B). There was also a trend for more bobs in the multi-bob Units 3 and 11 and fewer in Unit 7. Dewlap pulses usually followed the head bobbing (23 of 29 displays), and when present, ranged from 1–7.

Display stereotypy was moderate, with 11 to 12 of the 14 display variables having CV values within the 0 to 35% range for stereotyped behavior.

When comparing between Montrouis and Petion-ville Type *B* displays, Wilcoxon's Two-Sample tests resulted in nonsignificance for all but four display variables. As with the Type *A* display, there were only minor quantitative differences between the *B* displays of the two populations. In essence, they shared the same general pattern.

The primary context in which the *B* displays were observed was that of male-male aggression. However, on several occasions males were filmed giving *B* displays to females during courtship.

TYPE C DISPLAY

The *C* display type is also related to the Type *A* display, and is composed of the alternation of single and multi-bob units (Fig. 9C). However, there are some large departures from the *A* pattern, much more so than seen in the *B* pattern. The major differences are: 1) a greater separation of bob Units 1 and 3; 2) a change in the shape of the head bobs of Units 1, 3, and 7; 3) the appearance of the dewlap during the head bobbing sequence (rarely during Unit 4, always during Unit 6); and 4) marked qualitative variability in the display pattern. Like the Type *A* display, dewlap pulses usually followed the head bobs; 79% of the *C* displays contained post head bob dewlap pulses, and when present they ranged from 1 to 9. Major amplitude movement was fre-

quently produced by four-legged push-ups.

The *C* displays were qualitatively variable, but tended to be of two versions which I have designated as C_1 and C_2 (Fig. 9C). Individual lizards performed one of the two versions. My sample size was too small, however, to confirm whether there was a polymorphism in the population for these two *C* sub-patterns, or if lizards simply had individual-typical versions of the Type *C* display. A *C* display was classified as one of the two versions on the basis of the bob shapes in Units 1 and 3 (see Fig. 9C). Besides C_1 and C_2 , no other pattern variants were detected in my sample.

Contexts for the *C* displays included courtship as well as male-male disputes. The film record contains fights where paired male contestants were very aroused (i.e., displaying many postural modifiers such as mouth open and orbled eyes) and exchanging *C* displays at close range. These were followed by one of the males lunging forward and locking jaws or biting his adversary on the body.

Figure 8. Summary DAP graph of Type *A* and *B* displays for three male *Anolis distichus* from Montrouis. Display duration is expressed as a percentage.

Figure 9. Summary DAP graphs of *Anolis distichus* Type C and D displays. C_1 and C_2 are two versions of the C display type. Stippled dewlap extension indicates the pulse was not always present. Numbered boxes indicate duration and identity of bob and inter-bob units.

TYPE D DISPLAY

As is true of display Types A-C, the Type D pattern is also based on single bob units alternating with multi-bob units (Fig. 9D). However, the D pattern least resembles the signature display, having several unique characteristics. First, the D pattern is "indeterminate" because it has no fixed end point. In this sense it is similar to the *cybotes* signature display. Second, the multi-bob units 7, 11, and 15 tend to have only two bobs per unit, rather than a wide range of possible bob numbers as seen in the multi-bob units of the A-C patterns. Third, dewlap pulses occur throughout the head bob sequence.

The large number of units for the Type D display reflects the repetition of the single and multi-bob units (essentially the bobs of Units 5 and 7). Most D displays were performed through Unit 11, but some ceased as early as Unit 7, some as late as Unit 15. There was a weak tendency to stop after a multi-bob unit.

With the repetition of the single and multi-bob units, the "plateau-spike" shaped bobs of the D display tend to grade into "spike-like" bobs toward the end of a display (Fig. 9D). Dewlap pulses always appeared after the head bob sequence. In the 58% of the D displays which concluded with dewlap pulses, the number of pulses ranged from 1 to 9.

The context for the Type D display was primarily male-male interactions, but two displays were filmed during male to female displaying. The D display is obviously an agonistic signal as it is accompanied by display modifiers (e.g., tongue out) commonly associated with aggressive behavior (Jenssen, 1979). How the D displays function in male to female relationships is uncertain.

TYPE E DISPLAY

Type E display does not share the construct of alternating single and multi-bob units as do the A-D display types. After Unit 3 there is an indeterminate

series of single bobs (Fig. 10E). This display type appears to be a simplified version of the Type *D* pattern.

As in the *C* and *D* displays, the dewlap is pulsed during head bobbing (i.e., internal dewlap pulses). For most of the *E* displays, there was a single internal dewlap pulse just before the last head bob, and in some performances there were multiple internal dewlap pulses (Fig. 10E, stippled outline of dewlap movement). The dewlap was also pulsed after the last head bob in all *E* displays, ranging from 1 to 7 extensions per display. All *E* displays were performed through Unit 7, but many contained additional single bobs. Covarying with this indeterminate bob series were changes in bob shape (Fig. 10E_{ex}) as seen to a lesser degree in *D* displays. The addition of single bobs during repeated *E* displays by a male, covaried with the bob of Unit 1 changing from a "spike" to a "plateau-spike." The number of bobs in Unit 3 also increased, and the single bobs tended to metamorphose into the plateau-spike configuration. This transformation of the *E* pattern

during repeated performances is unique among previously studied lizard species. The closest resemblance to this phenomenon is a shifting pattern within display performances by the banded iguana, *Brachylophus fasciatus* (Greenberg and Jenssen, 1983).

Because the Type *E* display pattern was not qualitatively stable, unit durations and bob shape of given bob units experienced a wide range of variation. Therefore, the DAP graph of the Type *E* display (Fig. 10E) is not a good generalization for all *E* displays. It simply reflects mean unit durations and intermediate bob shapes. That is why DAP graphs for four consecutive *E* displays were provided in Figure 10.

The context for all *E* displays was resident-intruder interaction in enclosures. No *E* displays were filmed in the field. Their occurrence in the field may require intense territorial defense which is a rare event. In enclosures, high levels of agonistic behavior are more easily induced since there are no potential escape routes available. The progressive transformation of the *E* pattern may indicate the displayer's shift to a more intense level of agonistic intent.

DISCUSSION

Anolis cybotes and *A. distichus dominicensis* are two common species on Hispaniola; the first belongs to the *cybotes* species group and the second to the *stratulus* species group (Williams, 1976). *A. cybotes* is a medium-sized anole (males ~ 65 mm SVL), while *A. distichus* is smaller (males ~ 50 mm SVL). The two species are sympatric in many areas. Their structural habitat preferences are quite similar. Adults of both species utilize broad perch substrates, primarily tree trunks and rock outcroppings. They are usually found from ground level up to about 3 m for *A. cybotes* and up to 8 m for *A. distichus* (Moermond, 1979).

Figure 10. Summary DAP graph of *Anolis distichus* Type *E* display (E), with an example of four sequenced displays by a single male (E_{ex}). Stippled dewlap extension indicates the pulse was not always present. Numbered boxes indicate duration and identity of units.

There was no *a priori* suspicion that these two sympatric congeners would have such radically different display repertoire sizes. *Anolis cybotes* has only a single species-unique head bob display type, its signature display. Other social behaviors observed, but not included in the repertoire count, were display modifiers (see Jenssen, 1978, 1979), some weakly stereotyped head dipping, and "rapid head bobs," sometimes called "jiggling" (see Jenssen, 1977). This last behavior has been reported by almost all studies of iguanid social behavior; it is most frequently performed by a courting male on the move. In contrast, *A. distichus* has six species-unique display types in its repertoire, excluding its own display modifiers and rapid head bobbing.

DISPLAY EVOLUTION

For the display-rich *A. distichus*, it is tempting to speculate on the evolution of its contemporary set of display types. As previously proposed for the evolution of anoline displays (Jenssen, 1977), it is parsimonious to assume that the primitive *distichus* repertoire had but one display type, the signature display. This is a situation not unlike that of present day *A. cybotes*. Since the A-D display types of *A. distichus* are based on alternating single and multi-bob units, the ancestral signature display probably shared this theme. The ancestral signature display is depicted as three series of alternating single and double bobs (Fig. 11). I am curious about the bob cadence of the other *A. distichus* subspecies. It will be instructive to see if their signature displays can be derived from my hypothetical signature DAP, or if some other ancestral pattern will be more appropriate.

The current Type A display was derived from the ancestral signature DAP. The Type B display evolved next (Figs. 11 and 12). The B pattern is very similar to that of the Type A display. One need only visualize the lengthening of

the bobs in Units 1 and 5 to derive the B display from the A (Fig. 11). It is not unusual for anoles and for some other iguanids to have a two display type repertoire in which the second display pattern is very similar to the species' signature display (Jenssen, 1975, 1981; Ortiz, 1979; Rothblum and Jenssen, 1978); this is also true for anoles with more than two display types in their repertoires (Hover and Jenssen, 1976; Jenssen and Rothblum, 1977).

After the Type B evolved, the Type A display also provided the basis for the Type C display. This took place with the additions of an internal dewlap pulse (i.e., a pulse during the head bobbing sequence) and the re-shaping of the multiple bob series of Units 3 and 7 (Figs. 11 and 12). Like the A and B displays, the Type C display is determinate, having no bobs after Unit 11. With the evolution of its internal pulses, the Type C display was most likely the source for the Type D display which also has internal pulses, but is indeterminate. It is most probable that the D pattern was derived from the C in two steps; the bob of Unit 1 was deleted and the D pattern became indeterminate by indefinitely repeating Units 5 and 7 (Figs. 11 and 12). The E display type came from the D through a process of reduction; the Type E pattern has fewer internal pulses and has lost the alternating single and multi-bob units (Fig. 11 and 12). The Type E displays continue the indeterminate construct of the D display type.

The last display type of the repertoire is composed of the *a* displays. These seem to be fragments of either the ancestral signature display or of the current signature pattern. Because of their very low amplitude and simplicity of pattern, the *a* displays do not fit into the A-E flow of the above hypothetical scheme.

I am left with the question of why *A. distichus* has such a large repertoire of social signals, while *A. cybotes*, another quite successful congener, suffices with a simple repertoire. To correlate large rep-

Figure 11. Hypothetical scheme suggesting the evolutionary steps required to derive the *Anolis distichus* display repertoire.

Figure 12. Diagram showing the relationships between *Anolis distichus* display types and the hypothesized evolutionary events producing the multiple display repertoire (refer also to Fig. 11).

ertoire anoles with socio-ecological factors not shared with congeners of limited repertoires is still premature, since there are still so few anoles known to have repertoires of three or more display types. Nevertheless, using my experience with *A. cybotes* and *A. distichus*, some tentative factors can be suggested.

DISTICHUS HYPOTHESIS

All of the *distichus* displays are used in agonistic encounters (Table 1). How these six display types function in aggressive interactions is not well known. However, in *A. limifrons*, the displays of its large repertoire are used sequentially (Hover and Jenssen, 1976). As the distance decreases between contesting males, their exchanged displays progressively shift through the inventory of display types in a hierarchical manner. This phenomenon is an elaborate ritualized aggressive system which seems to allow males to assess each other's willingness to fight; it actually decreases the likeli-

hood of fighting by extending the period for withdrawal without risk of injury. I suspect the *distichus* repertoire is used in the same way.

Possible factors favoring a more elaborate bluff system are: 1) shortlived, high turnover species active throughout the year; 2) relatively high density per unit area with numerous male neighbors; 3) relatively small body size with concomitant small territory size; and 4) structurally complex microhabitat with widely and continuously available perch substrates.

Under this set of conditions, a male would experience frequent contacts with its neighboring males, some being new to the group of associates. A complex microhabitat, such as a large fig tree, would provide a substrate for many territories with no natural boundaries between territories. A single male could not control the entire microhabitat complex, and would need to continuously patrol to maintain his residential boundaries, especially against intrusions by new neighbors. Conversely, the same exten-

TABLE 1. GENERAL ASSOCIATION OF DISPLAY WITH GROSS CONTEXTUAL SITUATIONS. An "X" indicates a strong association, an "x" a weak association, and a "?" an uncertain association.

Species	Display Type	General Context		
		Non-directed	Courtship	Aggression
<i>cybotes</i> <i>distichus</i>	Signature	X	X	X
	A (Signature)	X	X	X
	a	X		x
	B		x	X
	C		x	X
	D		?	X
	E			X

sive habitat complex would provide living space for many lizards (i.e., not be a significant limiting resource), although it would certainly vary in preferred quality. The male population within this patch of habitat could easily reflect a shifting strategy between strict territoriality (specific defended area) and a social hierarchy. A few dominant males would have fairly stable territories, while less dominant males would inhabit less preferred parts of the substrate complex and have unstable or floating territories.

Since many male-male contacts would be experienced throughout the day, enforcement of territorial borders would be best served by graded ritualized aggression, modulated by the potential threat of the intruder. If all members of the population were quick to fight, the incidence of injury would be great; this is not likely to be selected for as indicated below.

Applying game theory to animal conflicts, Maynard Smith and Price (1973) concluded that there should be no stable pure strategy in aggressive contests; a mixed strategy of at times "hawk" and at times "mouse" would be stable (evolutionary stable strategy-ESS). An ESS requires that contestants respond to escalated aggressive behavior by escalating in return, and that individuals vary their behavior from contest to contest depending on the opponent's fighting power and willingness to fight (Dawkins and Krebs, 1978). Where fighting is avoided because

the cost of injury outweighs the benefits (e.g., being injured while maintaining a portion of a relatively abundant resource), one would expect an extensive and graded repertoire of agonistic signals by which to assess an opponent's fighting ability. Thus the ESS becomes one of "assessor/retaliator," using conventional or ritualized displaying; escalation to actual fighting should occur only between evenly matched contestants (Parker, 1974).

How well *A. distichus* fits my suggested scheme requires still ungathered data on its behavioral ecology. However, collected males showed no fight marks, no naturally occurring fights were observed, and only in enclosures (where living space was truly limited) was fighting induced.

CYBOTES HYPOTHESES

The simple repertoire of *A. cybotes* suggests a different approach to social interactions than that of *A. distichus*. Factors likely to maintain a more primitive repertoire are: 1) a relatively low density per unit area, with few male neighbors; 2) a relatively low turnover rate, with a stable set of nearest neighbors; 3) relatively large body size, with concomitant large territory size; and 4) territories which encompass entire or nearly entire patches of preferred microhabitat.

Under these conditions a territorial male would not be frequently challenged by its neighbors. Strips of unsuitable habitat forming an ecological "no man's land" would frequently separate territories. Territories could be rigidly defended and controlled because natural breaks in the microhabitat would delineate territories and decrease casual contacts. Intrusion into a territory would represent a serious threat to a resident, since eviction would mean a significant search time for a new territory by the ex-resident. If resident and intruder were moderately matched for fighting ability (i.e., similar body size), a rapid escalation to fighting is predicted because of the high benefit to holding or acquiring the territory by a contestant (suitable habitat may be a more limited resource than in *A. distichus*). A large repertoire of ritualized signals would be unnecessary because of the relatively rare event of male-male contact and a greater tendency to risk injury to protect a more limited resource.

My observations of *A. cybotes* indicate that in comparison with *A. distichus* they are very aggressive and prone to injurious fighting. *Anolis cybotes* were seen engaged in naturally occurring fights. Many (~ 20%) of the observed and collected males bore scars and fresh head and jaw wounds. Several males had very badly damaged jaws which were obviously broken.

ACKNOWLEDGMENTS

I am indebted to a number of people: Chris Pague and Lars Jenssen assisted me in the field; Teri Brentnall and Mike Menjak conducted initial film analyses on parts of the film record; Ned Fetcher helped with the computer programming; and Pam Pettry typed the manuscript. Financial assistance came from a National Science Foundation Grant DEB 74-20143.

LITERATURE CITED

- ARNOLD, D. L. 1980. Geographic variation in *Anolis brevirostris* (Sauria: Iguanidae) in Hispaniola. *Breviora Mus. Comp. Zool.* No. 461, pp. 1-31.
- BARLOW, G. W. 1977. Modal action patterns, pp. 98-134. In T. A. Sebeck (ed.), *How animals communicate*. Bloomington, Univ. Indiana Press.
- COCHRAN, D. M. 1941. The herpetology of Hispaniola. *Bull. U.S. Nat. Mus.*, 177: 1-398.
- DAWKINS, R., AND J. R. KREBS. 1978. Animal signals: Information or manipulation?, pp. 282-309. In J. R. Krebs and N. B. Davies (eds.), *Behavioural ecology, an evolutionary approach*. Sunderland, Massachusetts, Sinauer Associates.
- GREENBERG, N., AND T. A. JENSSEN. 1983. Social behavior of captive banded iguanas, *Brachylophus fasciatus*. In G. M. Burghardt and A. S. Rand (eds.), *Iguanine lizards of the world: Behavior, ecology, and conservation*. New Jersey, Noyes Publ. In Press.
- HOVER, E. L., AND T. A. JENSSEN. 1976. Descriptive analysis and social correlates of agonistic displays of *Anolis limifrons* (Sauria, Iguanidae). *Behaviour* 58: 173-191.
- JENSSEN, T. A. 1975. Display repertoire of a male *Phenacosaurus heterodermus* (Sauria: Iguanidae). *Herpetologica* 31: 48-55.
- . 1977. Evolution of anoline lizard display behavior. *Amer. Zool.*, 17: 203-215.
- . 1978. Display diversity in anoline lizards and problems of interpretation, pp. 269-285. In N. Greenberg and P. MacLean (eds.), *Behavior and neurology of lizards: An interdisciplinary conference*. N.I.M.H., Rockville, Maryland.
- . 1979. Display modifiers of *Anolis opalinus* (Sauria, Iguanidae). *Herpetologica*, 35: 21-30.
- . 1981. Unusual display behavior by *Anolis grahami* from western Jamaica. *Copeia*, 1981: 728-733.
- JENSSEN, T. A., AND E. L. HOVER. 1976. Display analysis of the signature display of *Anolis limifrons* (Sauria, Iguanidae). *Behavior*, 57: 227-240.
- JENSSEN, T. A., AND L. ROTHBLUM. 1977. Display repertoire analysis of *Anolis townsendi* (Sauria, Iguanidae) from Cocos Island. *Copeia*, 1977: 103-109.
- MAYNARD SMITH, J., AND G. R. PRICE. 1973. The logic of animal conflict. *Nature*, 246: 15-18.
- MOERMOND, T. C. 1979. Habitat constraints on the behavior, morphology, and community structures of *Anolis* lizards. *Ecology*, 60: 152-164.
- ORTIZ, P. 1979. Behavioral comparison of anoline competitors: *Anolis cooki* and *Anolis cristatellus* (Reptilia: Sauria: Iguanidae). M.S. Thesis. Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- PARKER, G. A. 1974. Assessment strategy and the

- evolution of fighting behaviour. *J. Theor. Biol.*, **47**: 223–243.
- ROTHBLUM, L., AND A. JENSSEN. 1978. Display repertoire analysis of *Sceloporus undulatus hyacinthinus* (Sauria, Iguanidae) from south-western Virginia. *Anim. Behav.*, **26**: 130–137.
- SCHWARTZ A. 1968. Geographic variation in *Anolis distichus* Cope (Lacertilia, Iguanidae) in the Bahama Islands and Hispaniola. *Bull. Mus. Comp. Zool.*, **137**: 255–310.
- . 1980. Variation in Hispaniolan *Anolis whitemani* Williams. *J. Herpetol.*, **14**: 399–406.
- SOKAL, R. R., AND F. J. ROHLF. 1969. *Biometry*. San Francisco, W. H. Freeman and Co.
- STEEL, R. G. D., AND J. H. TORRIE. 1960. *Principles and procedures in statistics*. New York, McGraw-Hill.
- WEBSTER, T. P. 1975. An electrophoretic comparison of the Hispaniolan lizards *Anolis cybotes* and *A. marcanoi*. *Bull. Mus. Comp. Zool.*, **431**: 1–8.
- . 1977. Geographic variation in “*Anolis brevirostris*”: Evidence from proteins and a consideration of dewlap color, pp. 153–165. *In* E. E. Williams (ed.), *The Third Anolis newsletter*. Mus. Comp. Zool., Harvard Univ., Cambridge, Massachusetts.
- WEBSTER, T. P., AND J. BURNS. 1973. Dewlap color variation and electrophoretically detected sibling species in a Haitian lizard *Anolis brevirostris*. *Evolution*, **27**: 368–377.
- WILLIAMS, E. E. 1963. *Anolis whitemani*, new species from Hispaniola (Sauria, Iguanidae). *Breviora Mus. Comp. Zool.* No. 197, pp. 1–8.
- . 1975. *Anolis marcanoi* new species: sibling to *Anolis cybotes*: Description and field evidence. *Breviora Mus. Comp. Zool.* No. 430, pp. 1–9.
- . 1976. West Indian anoles: A taxonomic and evolutionary summary. I. Introduction and a species list. *Breviora Mus. Comp. Zool.* No. 440, pp. 1–21.
- . 1977a. Additional notes on distichoids, pp. 174–182. *In* E. E. Williams (ed.), *The Third Anolis newsletter*. Mus. Comp. Zool., Harvard Univ., Cambridge, Massachusetts.
- . 1977b. Species problems, pp. 132–151. *In* E. E. Williams (ed.), *The Third Anolis newsletter*. Mus. Comp. Zool., Harvard Univ., Cambridge, Massachusetts.
- WYLES, J. S., AND G. C. GORMAN. 1980. The classification of *Anolis*: Conflict between genetic and osteological interpretation as exemplified by *Anolis cybotes*. *J. Herpetol.*, **14**: 149–153.

The Social Behavior of *Anolis valencienni*

ROBERT A. HICKS¹

ROBERT L. TRIVERS²

ABSTRACT. *Anolis valencienni* is highly cryptic and displays many features atypical of other *Anolis*. Adult females are completely nonterritorial and nonaggressive, frequently passing within centimeters of each other. Adult females overlap with each other extensively in their daily movements. Sometimes as many as 40 females may occupy a common feeding space. Males are aggressive to each other but show broad overlap in the space occupied. It is not uncommon for five adult males or more to share much of the same space. Consistent with their lack of territories, adult females wander slightly greater distances than do adult males, even though they are smaller. Males appear to suffer differential mortality, and the adult sex ratio is biased toward females. Both sexes feed by searching slowly over the substrate of trees and bushes for insects, many of which are cryptic and at rest. Because many adults of both sexes encounter each other daily, there are unusual opportunities for female choice. Females copulate often during the summer, sometimes, and probably often, more than once a day. Over a period of six weeks a female may copulate with five or more males. Experiments were conducted to test the criteria females use in choosing males, but these were not successful. Field data show that both adult males and adult females copulate more often as they increase in size, but for males this relationship appears to be weaker than in the more size-dimorphic *Anolis garmani*. It is proposed that the many unusual features of *A. valencienni* result from the primary feeding adaptation of searching for stationary, cryptic prey.

INTRODUCTION

In appearance, *Anolis valencienni* is the most cryptic of Jamaica's seven species of anoline lizards. It is also the

slowest moving, crawling deliberately and very slowly over the trees it inhabits. Both in size and in secondary sexual structures such as the dewlap, it is likewise unique in that females are never aggressive toward each other and tolerate the presence of other females at distances of centimeters. More than a year of field work on this species suggest to us that the above attributes of *valencienni* are all interrelated and ultimately explained by *valencienni*'s unique feeding specialization: individuals search actively for and feed upon cryptic insects, many of which may be diurnally immobile. We describe here the results of this field work.

METHODS

Anolis valencienni was studied in two different localities and at several different times in each locality.

Study site 1 (Maryfield) was located in a suburban area of Kingston and consisted of the three acre grounds of the Maryfield Guest House. These, in turn, consisted of scattered large trees (usually over 10 m high) typical of Jamaica, mostly mangos (*Mangifera indica*) but also *Persea americana*, *Guaiacum officinale*, *Delonix regia*, *Terminalia catappa*, *Haematoxylum campechianum*, *Swietenia mahagoni*, *Gliricida sepium*, and *Brya ebenus*. During 12 days in September 1968, feeding behavior was studied by following (through binoculars) unmarked lizards. From June 15 to September 14, 1969, the social behavior of these

¹ P.O. Box 420, Ipswich, Massachusetts 01938, U.S.A.

² Applied Sciences, University of California, Santa Cruz, California 95064, U.S.A.

lizards was studied for a total of eleven weeks by capturing, sexing, measuring, marking, releasing and later following 265 lizards in the study area. Brief visits to this study area (to recapture marked individuals) were made in January 1970, and again in April 1970. All observations in study site 1 were made by Trivers, and no assistants were employed. *A. valencienni* were abundant in this study site, but because the site consisted of many large trees, the lizards were difficult to resight (and recapture) and most of our data on social behavior come from the second study site.

Study site 2 (Southfield) was located in rural St. Elizabeth and was chosen because *valencienni* were abundant and because the study site consisted of many bushes and short trees (most were under 6 m in height). Typical tree species, in addition to mangos, were *Pimenta officinalis*, *Cocos nucifera*, *Manilkara zapotilla*, *Blighia sapida*, *Chrysophyllum cainito*, and *Persea americana*. Social behavior was studied for two weeks in June 1971, by Trivers with the help of field assistants, and for seven weeks in July and August 1973, by Hicks with help of field assistants. All resightings of individuals, including copulations and aggressive encounters, were confirmed by Trivers or Hicks even if first made by one of the assistants. The second site was visited by Trivers for eleven days in December 1973 to January 1974 in order to recapture marked *valencienni*. In addition, Trivers studied *Anolis garmani* for a year in Southfield, where *valencienni* were abundant, and he made numerous casual observations during this year. Finally, five months of field work in the spring of 1976 and in July 1977, concentrated on measuring female choice.

Social behavior was studied as in *A. garmani* (Trivers, 1976); that is, an effort was made to capture, sex, measure, and mark all adults in the study area. Individuals were given a permanent mark by clipping two or three toes in a pattern unique for each lizard. Likewise, a num-

ber was painted on each lizard's back, permitting identification in the field without recapture. As molting takes place about every three weeks in *valencienni*, individuals were often recaptured and repainted. Such recapture work permitted some systematic data on growth rates, and these are analyzed below as a function of size, sex, and rate of dispersal. When an effort was made to gather copulation data (see below), the study area was systematically searched for copulating lizards. If these were marked, numbers were recorded. If one or both were unmarked, the unmarked individuals were captured. It was not possible to estimate precisely to what extent the sample was a biased sample of those copulations actually occurring, but some biases were apparent. These, and other methodological problems are discussed, where relevant, below. Resightings were recorded while searching for unmarked lizards and for copulating pairs. The data from resightings are analyzed below. In addition, observations of aggressive, sexual, and neutral encounters were made in the process of the work described.

RELATIVE CRYPTICITY

Anolis valencienni is unusually cryptic (Lynn and Grant, 1940; Underwood and Williams, 1959; Rand, 1967a; T. W. Schoener and A. Schoener personal communication), and experienced Jamaicans routinely assert that *valencienni* is the most cryptic anole on the island. Several lines of evidence support this opinion. In what follows, we emphasize the comparison between *valencienni* on the one hand and *A. lineatopus* and *A. grahami* on the other; because the latter two species are similar in size to *valencienni*, they often occupy similar perches, and they both occur in our two study areas.

Sightings. 1) On a typical day in either of the study sites, individual *lineatopus* and *grahami* were sighted first, although we were searching for *valencienni*. 2) Marked *grahami* and *lineatopus* are

much more frequently resighted than marked *valencienni*. (This is partly due to the greater crypticity of *valencienni* and partly due to its less predictable choice of perches.) 3) In both study sites, *valencienni* initially appeared to be rare, but marking revealed *valencienni* to be more common than *grahami* in both study sites, more common than *lineatopus* in the second study site, and only slightly less common than *lineatopus* in the first study site. 4) Rand's (1967a) census study of Jamaican anoles found *valencienni* to be less common than any other species except *garmani*. This is expected since lizards were not marked and each study area was traversed only once per day. Underwood and Williams (1959) likewise point out that *valencienni* is much more common than the number of collected specimens would suggest "for it is very clever at concealing itself." 5) It is a common experience to be attracted to a tree because one spots one or more *lineatopus* or *grahami* and after watching these for several minutes to notice an overlooked *valencienni*, often because it finally moves.

Morphology. 1) *A. valencienni* is usually grey in color (sometimes brown) with irregular dark markings which tend to obliterate body outline. This contrasts with the solid blue (or dark brown) color of *grahami* and the regular, reticulated pattern of *lineatopus*. 2) In contrast to *grahami* and *lineatopus*, the splotched, irregular pattern of *valencienni* covers the entire body, including the belly, the face and lower jaw. 3) All other Jamaican anoles have a bright, yellow eye-ring. This eye-ring is obliterated when the lizards are frightened in social encounters and their entire bodies darken. *A. valencienni* lacks the eye-ring entirely.

Behavior. Individual *valencienni* move very close to the surface they are climbing on. When they do move, they move very slowly, so that they appear to glide quietly and nearly invisibly from place to place.

SEXUAL DIMORPHISM

Both in size and in secondary sexual characteristics, *valencienni* is, along with *A. reconditus* (Hicks, 1973), the least dimorphic Jamaican anole.

Data on adult size dimorphism from different visits to the two study areas give consistent results. Since the data from the summer of 1973 (Southfield study site) are the most complete, they are presented here. The snout-vent length of the largest five males averages 85.2 mm while that for the largest five females averages 73.8 mm. The smallest adult male seen copulating was 54 mm in size (= snout-vent length), while the smallest female seen copulating was 50 mm in size. Using these sizes as size of sexual maturity, the average adult male size ($N = 93$) is 72.8 mm, while the average adult female size ($N = 98$) is 64.8 mm. At all sizes for which data are available, males grow faster than females of similar size (Fig. 1). Like other anoles, male *valencienni* have larger jaws than females of similar snout-vent length.

It is of particular interest to know whether the size dimorphism in *valencienni* results entirely from differential growth rate or whether males hatch out at a larger size than females. A sample of 55 *valencienni* hatched from eggs (and then preserved) is available at the Museum of Comparative Zoology. These were measured and then sexed. Forty-nine of these could be sexed unambiguously (by the enlarged post-anal scales of the males), but six individuals had post-anal scales intermediate in size. Although these are probably males, they have been treated separately. No sex difference in size at hatching is apparent. Mean size of known males ($N = 22$) is 23.8 mm, while the mean size of known females ($N = 27$) is 23.7 mm. If the six uncertain individuals are indeed males then the mean size of males ($N = 28$) is 24.0.

There are no pronounced secondary sexual structures in *valencienni* except the dewlap, which is very large and

Figure 1. Growth Rate (mm/100 days) is plotted as a function of initial size (snout-vent length in mm) for males and females between summer 1973 and winter 1973–74 in Southfield. On average each individual of both sexes was recaptured six months after initial capture.

bright purple in the male and smaller and less brightly colored in the female, often showing many white scales.

ADULT SEX RATIO

In all visits to both study areas more adult females were captured than adult males. We divide the data by study area, however, because at Maryfield females were easier to capture (Table 1) while males were easier at Southfield (Table 2). In order to make inferences about differential mortality by sex, we present not only the number of adult males captured (≥ 55 mm), but also the number of males ≥ 50 mm captured, since the latter category certainly contains individuals younger than the youngest adult females (see below).

During the summer of 1969 at Maryfield 198 adult females (≥ 50 mm) were captured compared to 69 adult males (≥ 50 or 55 mm). These data are certainly biased, because males were much less often sighted and were less often captured when sighted (since they stayed higher in the trees). As evidence of this bias, for example, the first 29 individuals captured in the study area were females.

TABLE 1. NUMBERS OF MALES AND FEMALES CAPTURED AT MARYFIELD.

	females (≥ 50 mm)	males (≥ 55 mm or ≥ 50 mm)
Summer 1969	199	62
Winter 1969–1970	26*	9†

*Of these 26 females, 20 had been marked in the summer of 1969.

†Of these 9 males, 3 had been marked in the summer of 1969.

Likewise, of the 121 adult females captured on or before August 6, 48% (58) were, at one time or another, recaptured, but of the 41 adult males marked on or before August 6, only 31.7% (13) were ever recaptured. Thirty-seven females marked on three large connected trees on or before August 6 (see "Female Movements") were resighted a total of 19 times after August 6 for an average of 0.51 resighting per female, while 12 males marked on or before August 6 were resighted thereafter a total of only once, for an average of 0.08 resighting per male.

Recapture data permit a crude estimate of the real adult sex ratio at Maryfield. During seven days in December 1969 to January 1970 and during three days in April 1970, a total of 35 adults were captured of whom 23 had been marked in the summer of 1969. Of the 26 females captured 88.5% (23) were marked the preceding summer, while of the nine adult males captured only 33% (3) had been marked. It is unlikely that many lizards migrated in and out of the study area, which was very large and bounded on three sides by roads. A very crude estimate of the real adult sex ratio at Maryfield (assuming no differential migration) is 186 males and 259 females.

At Southfield males appeared to be slightly easier to spot than females and, once sighted, very few individuals of either sex eluded capture. The 92 adult males captured were resighted a total of 158 times for an average of 1.72 resight-

ing per male while the 97 adult females were resighted a total of 151 times for an average of 1.56 resighting per female. Yet each visit to Southfield showed more adult females than adult males (Table 2). None of the differences are significant; nor, when combined, are the totals significantly different.

Recapture data appear to confirm that more females than males remained uncaptured during the summer of 1973. Of the 31 females captured during the winter of 1973 to 1974, 67% (21) had been marked during the preceding summer, while of the 24 males captured 75% (18) had been marked in the preceding summer. These data may partly reflect greater female mobility, as data from the summer of 1973 show a slightly greater distance between successive sightings of females than of males (see below). Assuming no difference in mobility, a crude estimate of the real adult sex ratio in the summer of 1973 is 92 males and 123 females. The male : female sex ratio for animals of all sizes at Southfield (1973) was 110 : 121.

Males grow faster than females at all sizes for which we have data. If, as in *A. grahami* (Trivers, 1976) and *A. lineatopus* (Rand, unpublished data), the sex difference in growth rates extends to small individuals (as appears likely) then a female 50 mm in length is in fact, older than a male of the same size. Likewise, a newly adult female is almost certainly older than a newly adult male. Thus, the

sex ratio for adults of the same age is expected to be more strongly biased toward females than our data on all adults (and on all individuals ≥ 50 mm) show. If the sex ratio of hatchlings is 50/50 (as expected, since there is no size difference between the sexes at hatching), then our data suggest differential male mortality. What data we have suggest that the sex ratio at hatching may, indeed, be 50/50 but the data are few and our ability to recognize males, imperfect. Of the 24 individuals < 35 mm caught at Southfield during the summer of 1973 and winter 1973 to 1974, 11 were males and 13, females.

FEEDING BEHAVIOR

Individual *valencienni* capture insect prey from the ground at the base of trees to the outer edges of leaves forty feet above the ground. They do so by searching slowly and carefully throughout the tree (or bush) in which they live. When a lizard first spots a prey item, the item is usually within 8 cm of the lizard's face and is usually captured once spotted. These findings emerge from 84 hours of systematic observation of unmarked *valencienni* in 1968 and are supported by numerous incidental observations made during the summers of 1969 and 1973.

In September, 1968, 36 feeding attempts were observed. An attempt at feeding was assumed to occur if an individual moved quickly for several inches and snapped its jaws. The feeding attempt was gauged to be a success if the jaw snap was followed by chewing (i.e., by jaw movements). In the majority of unsuccessful attempts, a small prey item (assumed to be an insect) fluttered away. Likewise, in the majority of successful attempts, an insect was seen in the lizard's mouth before being consumed.

The distance at which the lizard first spotted the prey item was recorded as the distance from which the lizard began its lunge. There is no evidence that individual *valencienni* ever creep up on prey.

TABLE 2. NUMBERS OF MALES AND FEMALES CAPTURED AT SOUTHFIELD.

	females (≥ 50 mm)	males (≥ 55 mm or ≥ 50 mm)
June 1971	44	35
Summer 1973	97	92
Winter 1973-1974	31*	24†

* Of these 31 females, 21 had been marked in the summer of 1973.

† Of these 24 males, 18 had been marked in the summer of 1973.

Usually, the lizard is moving slowly over an area, constantly searching it visually, when it interrupts the searching to lunge and snap, often at something hidden in the bark under its nose. Of the 36 feeding attempts, 27 were successful. Of the 34 for which the distance the prey item was spotted was recorded, all but three times, the prey item was less than 12 cm from the lizard when first spotted. The furthest a prey item appeared to be spotted was 45 cm. The feeding attempts ranged from ground level to 13 m up. Twelve took place 6 m above the ground or higher, but these data are strongly biased toward observations close to the ground, since lizards were more easily spotted and followed at low heights. Because *valencienni* takes prey at short distances with high success, once prey is spotted, the species is, in the terminology of Schoener (1971) a *searcher* (as opposed to a *percher*, such as *lineatopus*). The finding that *valencienni* takes food from throughout the tree is consistent with the data of Rand (1967a) and Schoener and Schoener (1971), namely, that *valencienni* has the most variable "perch" height of a Jamaican anole, i.e., it is sighted from the ground to the tree-tops.

In behavior, *valencienni* is clearly a searching species, in the sense that most individuals can be observed most of the time searching for prey. They do so in five different places: 1) on the ground; 2) on leaves; 3) on the trunk of a tree and its limbs; 4) in holes; and 5) on epiphytes growing on trees, epiphytes such as *Tillandsia recurvata*. The searching behavior itself consists of moving very slowly while frequently cocking the head so as to look at the substrate. The lizards are especially likely to search irregular places in the bark, large holes in a tree and places where limbs have broken off. Likewise, they search out all epiphytes very carefully and on broad-leaved trees such as mangos will go out onto leaves to search their surface.

Typical searching behavior of two dif-

ferent individuals are presented here, condensed from Trivers' field notebook (measurements converted to metric equivalent).

While drawing close to a tree to observe two *lineatopus* at 8:30 am on September 16, 1968, Trivers almost stepped on a *valencienni* (assumed to be a female) who was on the ground a foot from the tree. Trivers withdrew and watched through binoculars:

8:33 lizard jumps onto tree and climbs straight up to 5m.

8:35 stops at series of bromeliads (*Tillandsia recurvata*)

8:38 moves in among them, searching

8:40 carefully, slowly climbing through a bromeliad

8:43 has moved a foot higher since entering bromeliads

8:46 stealthily moves another three inches

8:49 another three inches, then suddenly moves 45 cm very quickly, as if it has spotted something. It then darts, snaps something off a bromeliad and eats.

8:50 moves several inches through bromeliad and stops

8:55 now 6 m up, turns to move out a horizontal branch which is about 7.6 cm thick

9:05 upside down in a large bromeliad, searching

9:10 same place

9:18 still upside down, searching, sometimes on a bromeliad, sometimes on the branch itself

9:25 upside down on a bromeliad when it seems to spot something on a neighboring bromeliad, also upside down. I too spot something on the second bromeliad. Starts to dart the 5 cm to the neighboring bromeliad but—as if forgetting it is upside down—it steps into thin air and falls 6 m to the ground. It appears to be uninjured.

In the second case, an individual was watched continuously for three hours and forty minutes. The individual was almost certainly a female as judged by her appearance, the size and coloration of her dewlap, and by the nonaggressive way she interacted with an adult male who appeared to be courting her. During the time she was observed, she moved from the foot of the tree to the foliage 11 m up and back down to the ground. She fed three times (once at ground level, twice at 11 m). All three prey items were spotted at distances of 15 cm or less. She

used her dewlap on six different occasions, on three of which the adult male was within a couple of meters of her. Observations on this animal are as follows:

At 9:40 on September 6, 1968 an adult *valencienni* was spotted upside down on a horizontal branch 5 m above the ground. Assumed to be a female, the lizard was searching an irregular area in the bark. She moved on to the trunk, and in the next ten minutes moved down to 1 meter above the ground. At 9:56 she moved down to 0.5 m, then down to 7.5 cm. She paused, dashed about 15 cm out onto the ground, caught a prey item, returned to 15 cm up the tree and ate what appeared to be a small, white moth.

10:13 a cat appears, circles the tree rapidly, and scares the female up to 5 m where she stops to search a warty, cracked area of the bark

10:10 1 m higher; a male who was also scared up by the cat is 1 m higher than the female, facing away from her and dewlapping repeatedly

10:14 male 1.5 m higher than female and still dewlapping; she jerks her head as if about to dewlap but no dewlap

10:23 she is now 10 m up going out a 2.5 cm thick branch

10:27 11 m up at end of branch, looking over a whorl of leaves

10:35 goes out another twig which ends in a whorl of over 20 leaves. Pulls itself into leaves and searches visually for over a minute, suddenly darts straight up 7.5 cm and catches a large insect, which she chews

10:40-10:57 searches other leaves in area

11:00 facing down on a branch 0.3 m below the leaves; male is 1 m above her dewlapping

11:04 female just dewlapped (small dewlap, many scales)

11:08 both male and female are 1.5 m lower. Male is dewlapping. Male moves rapidly down the branch, past where the female is and further down

11:13 female is 5 m up slowly searching the bark; for the next five minutes she does not move

11:28 she is 0.3 m lower, facing down, head raised, often looking around (at what?)

11:30 still looking around, moves sideways and dewlaps again, then moves to another branch

11:35 5.4 m up, slowly moving up

11:38 6 m up, moving up. At 11:42 she dewlaps several times (again, small dewlap, much white showing)

For the next 45 minutes she spent most of her time searching several clumps of

leaves (without success). At one point, a large wasp landed within 8 cm of her without eliciting any response. At 11:56 she dewlapped twice; no other lizard was visible near her. At 12:20 she fed on a small insect which was about 2 cm back of her head when she spotted it (10 m above ground on a twig). At 12:26 she dewlapped again and within a minute the large male was spotted about one meter above her, dewlapping. He moved to within a few cm of her and dewlapped repeatedly. Then he ran by her. For the next twenty minutes she and he descended to 5 m up, and the entire time he was within a meter of her, often dewlapping. She dewlapped at 12:53, and he moved rapidly up the tree away from her. For the next half hour she descended slowly to a height of less than a meter. Observations were discontinued at 1:25.

FEMALE-FEMALE INTERACTIONS

Two or more marked adult females were seen within centimeters of each other over 100 times during all visits to the two study areas, and no aggressive interaction was ever observed. Marked females were never seen to display to each other, nor to make any physical contact nor to show any clear avoidance behavior. On four occasions unmarked adults who appeared to be females displayed toward other adults assumed to be females. In three cases, a presumptive female dewlapped at a second presumptive female which was several centimeters away. There was no response nor any follow-up behavior. It is very likely that the dewlapping individual in these three cases was a female, but it is not certain that the recipient was also a female. In the fourth case a presumptive female bit at her own reflection in a mirror. (All other females ignored their mirror images.)

In virtually all interactions between marked females, the relations appeared to be neutral, but in certain situations, described below, females may have been

positively attracted to each other (or to some common stimulus). The neutral relations between adult female *valencienni* contrast strongly with the aggressive interactions that characterize female-female relations in *A. lineatopus* (Rand, 1967b), *A. garmani* (Trivers, 1976), *A. opalinus* (Jenssen, 1973) and *A. grahami* (Trivers, unpublished data). To illustrate the nonaggressive nature of *valencienni* female relations, we give here two examples.

1) On July 22, 1969, three adult females (snout-vent lengths: 60, 65, and 66 mm) were seen within centimeters of each other at a hole about 5 m up a mango tree. The three arrived at the hole within fifteen minutes of each other from three different directions, each stayed several minutes and left independently of the others. Two other adult *valencienni*, which were unmarked and assumed to be females, passed within centimeters of two of the marked females. At no time did any of the individuals clearly alter behavior in response to the others. The hole which attracted the females was one near which females were often seen and one which contained *valencienni* eggs. It is not known whether the sight of a female near such a hole attracts other females.

2) Females (like males) are reluctant to leave bushes or trees for open ground. They appear to be much less reluctant if another individual has preceded them. On August 6, 1969, at 2:15 pm an unmarked adult *valencienni* left a mango tree to climb among some rocks at the base of the tree. Within about 20 minutes, four other unmarked adults left the tree to search among the same rocks. At 2:45, all were caught. Two were then 4 m from the base of the tree but within centimeters of each other. Both were 66 mm females. Two others closer to the tree and within one meter of each other were a 64 mm female and a 62 mm female. And, finally, the individual nearest the tree was a 62 mm male. All appeared to be searching for food, and at least one fed successfully while among the rocks. No

interactions were apparent. The presence of an adult male is not typical, but other such cases have been observed: one male within a meter of one or more females, with no obvious interaction. For example, at 10:30 am July 6, 1969, four adults were caught while searching for food on the trunk of a royal poincianna tree, all within 1 m of the ground and of each other. Three were females (59, 60, and 65 mm), and one was a male (68 mm).

The above example is only atypical in that more than two individuals were involved. In the majority of female-female interactions, one female passes close to another while both are searching for food. It would be interesting to know if females tend to avoid searching through areas recently searched by others, but no good evidence exists one way or the other. Certainly females do not show strong avoidance, and we have often seen one female search the general area recently searched by another. Females may be actively attracted to each other 1) at holes containing eggs and 2) at the base of trees, but this is merely an impression based on the frequency with which we observed more than one female together at such places.

DAILY FEMALE MOVEMENTS AND DISPERSAL

Since females do not interact aggressively and since there is no evidence they avoid each other, it is of interest to know how extensively their home ranges overlap. The best evidence for degree of overlap comes from three large contiguous trees whose occupants were watched during the summer of 1969. (Data from other trees give similar, but much less detailed, results.) The trees were two mangos connected together by a royal poincianna. Each tree stood about 13 m tall and had extensive foliage. The three trees were connected to each other in the foliage but to no other trees. It was possible, then, for a female to wander

throughout the three trees. Distances between the trees and the number of adult females captured on each are given in Table 3.

Between June 16 and September 6, 1969, 43 adult females were caught and marked on the three trees. The great majority of these (37) were marked by August 6. Since the six females caught after August 6 had little opportunity to be resighted (only one of them was, in fact, resighted), the analysis here is limited to the 37 marked by August 6. Of these 37, 29 were resighted at least once for a total of 69 resightings, or almost two resightings per individual marked on or before August 6. Of these 69 resightings, 64 found the lizard on one of the three connected trees. (The other five resightings involved nearby trees and they are discussed below.)

The 64 resightings are analyzed according to each lizard's tree of origin in Table 3. Just as more females were captured on mango-1 than on the other two trees, more females were resighted on that tree as well. This may partly indicate a preference by the lizards for mango-1, but it may result entirely from the greater ease with which lizards could be resighted and caught on mango-1. The tree had a large hole containing over 50 *valencienni* eggs located only one meter above the ground, and females were often sighted near this hole or near a second egg hole 4 m above ground. (The poincianna was not known to contain an egg

hole and mango-2 was only known to have one, 6 m above ground.)

Of the 21 lizards caught on mango-1, 18 were resighted and 82% of the resightings found the females on that same mango. Of the five females caught on the poincianna, all were resighted, but only 22% of the resightings found the lizards on the poincianna. Of the eleven females caught on mango-2, only six were resighted and only 27% of these resightings occurred on that mango. By treating each resighting as an independent event, it is possible to test for heterogeneity in the data. The data almost show a significant tendency for lizards to be resighted more often on the tree of their capture than expected by chance ($\chi^2 = 8.13$; $0.05 < p < 0.10$). What is clear is that the females wander very widely. Only one female (60 mm) showed a marked tendency to remain on the tree of initial capture (mango-1): during a three month period, she was resighted seven times, all of them on mango-1. If data on her are removed from Table 3, no trend toward localized movement is apparent.

Movement from one tree to another can be rapid, taking place in less than two hours. Four cases of females moving between two of the three trees within a day are presented here. 1) On July 8 a 60 mm female was captured and released on mango-2. An hour and a half later, she was spotted at the base of the poincianna. The following day she was seen again on the poincianna, but two days later than this she was seen on mango-1. On July 14 she was again seen on mango-1. 2) On July 18 a 64 mm female was caught at 10:00 am on mango-1 and was resighted that afternoon at 4:15 at the base of mango-2, 29 m away. She presumably reached mango-2 by way of the poincianna. 3) On July 21 a 65 mm female was caught on the poincianna. The following day she was seen at the base of mango-1. 4) On July 24 a 61 mm female was caught on the poincianna and a day later was seen on mango-2.

It might be supposed that these rapid,

TABLE 3. THE NUMBER OF TIMES 37 FEMALES WERE RESIGHTED ON THREE TREES AS A FUNCTION OF THE TREE ON WHICH EACH WAS INITIALLY SIGHTED.

		M-2	POINCI	M-1	TOTAL
TREES					
FIRST	M-1	5	3	36	44
CAUGHT	(21)	2			
ON (No.	Poinci	2	2	5	9
in paren-	(5)				
theses)	M-2	3	2	6	11
	(11)				
	TOTAL	10	9	47	

long-range movements are a response to the trauma of capture, but we doubt this for several reasons. 1) Similar movements have been recorded between resightings (where no capture was involved). For example, a 62 mm female was resighted on August 5 on the poincianna and two days later was resighted again on mango-1 (where it had originally been captured). Likewise, in the data above, the 60 mm female moved to a new tree on the day of capture, where she remained for at least a day, before moving on to the third tree. 2) By following individual unmarked females for several hours through binoculars, two such long-range movements were observed on the same trees in 1968, although no lizards were captured during that visit. 3) In the Maryfield study site, from which these data on movements come, lizards were caught only at ground level, so that a lizard could easily evade capture by remaining above 4 m yet the movements described involve moving to a height of 10 or more meters, crossing to a neighboring tree and sometimes returning to ground level. Within a half hour of capture, a female's behavior could not be distinguished from that of an uncaptured female; that is, a female recently captured rapidly returned to the slow, methodical search for food characteristic of the species. 4) Such long movements in response to capture have not been observed in highly territorial species (*A. lineatopus*, Rand, 1967b; *A. garmani*, Trivers, 1976; *A. grahamsi*, Trivers and Hicks, unpublished data).

In addition to the 64 resightings recorded in Table 3, five additional resightings were recorded in trees near (but not connected in the foliage to) mango-1. In one case a female was captured on mango-2 and resighted only once, nearly two months later, on a mango 30 m from mango-1. This may have been a permanent move. In the second case, a female moved a short distance (about 12 m) across open ground to a neighboring tree and returned to her tree of capture (mango-1) all within two weeks. The

third case involved a 66 mm female who was caught at the base of mango-2 on August 6. Ten days later she was resighted 56 m away on an avocado tree near mango-1. Thirteen days after this she was again seen on mango-2, but twelve days later (on September 10) she was seen back on the avocado tree (as she was on September 13). The female presumably moved from mango-2 onto the poincianna, from there onto mango-1 and then moved 10 m across open ground to some bushes which connected with the avocado tree. Although it is much more likely that the female crossed by way of the poincianna and mango-1 than that she moved 55 m across open ground, such inferential data have not been included in Table 3. (Similarly, one can safely infer presence on the poincianna when a lizard is sighted on mango-1 and then on mango-2, or vice-versa, but five such inferences were not included in Table 3).

Even these examples do not give an adequate picture of *valencienni* female movements. Observations of female movements within a given tree indicate that all females spotted low on a tree soon move high into the foliage. Observations of the same female on successive days reveal no tendency for a female to emerge from the same part of the foliage nor to return to the same part. Although it is likely that more detailed data will reveal some localization, it is clear that the overlap of female home ranges within a tree, or within contiguous trees, is enormous.

The three trees also contained at least 12 adult males. This is certainly a serious underestimate, since many unmarked adult males were seen but never captured, primarily because males stayed considerably higher in the trees than did the females. Based on several lines of evidence reviewed in the "Sex Ratio" section, we estimate that the trees contained at least 25 adult males. Of the 12 actually caught, eight were captured on or before August 6, but only one of these was resighted, so that almost nothing is

known about these males. It is clear, however, that even if males defended nonoverlapping territories, no female would live exclusively within one male's territory and most females would wander (in the space of about two weeks) through virtually every male territory. In the space of a single day a female would be expected to encounter at least five adult males.

In other Jamaican *Anolis*, the larger size of the adult male is associated with a larger territory or home range than seen in the adult female. In *A. valencienni*, we may expect that this relationship will not hold. Females are not territorial, while males, as we shall see, are repelled by each other. Thus, females may wander more widely than do males.

In the summer of 1973, we had a sufficient number of resightings to test this possibility. Lizards were watched for six weeks. The maximum distance between resightings was calculated for each individual who was observed more than one time. Forty-seven adult females were resighted an average of 3.1 times each, while 30 adult males were resighted an average of 4.7 times. Although males were sighted more often than were females, their average maximum distance between resightings, 26.8 m, was slightly shorter than that for females, 28.7 m. If we imagine that each resighting contributed equally to the maximum distance (which is unlikely), then for each resighting, males moved 4.7 m, while females moved 7 m. What seems clear is that adult females move somewhat greater distances during six weeks time than do adult males. These data certainly include some examples of dispersal to new areas, but these instances are few. Most of the data described movements typically made during a period of two or three days wandering.

Individual *A. valencienni* take very unwillingly to the ground. They are slower on the ground than any other Jamaican anole and are found there less often. Like other Jamaican anoles, they descend to

the ground to feed, but we have never seen an individual move more than 3 m from the base of its tree in search of food. We have only twice seen individuals on the ground who were not feeding. One had fallen accidentally out of its tree, and the other was in the process of dispersing 22 m to a new tree. Data on 257 marked individuals indicates that some individuals do move as much as 35 m over ground to reach a new tree. Of the 257 marked individuals, 49 were recaptured and, of these, eight had dispersed across ground to new trees.

Comparing those who dispersed with those who did not shows that those who did not disperse grew more than those who did. Of the 41 individuals who did not disperse, 34 showed measurable increases in size (2 mm or more) in periods ranging from one half month to two months, whereas only two of the nine who dispersed showed significant increases in size in similar time periods ($p = 0.01$). The average growth rate for nondispersers was 2.34 mm/30 days and for dispersers it was 1.06 mm/30 days ($p = 0.001$).

When growth rate is plotted as a function of female size, it is seen that dispersing individuals tend to have lower growth rates for two reasons. Dispersing females tend to be larger and to show smaller growth rates for a given size. Neither effect is in itself significant.

Each effect could be explained as follows. Larger females may be less vulnerable to predation and able to traverse more quickly the open space. Females with low growth rates may be selected to take a greater risk in finding a new feeding area. Of course, females who have low growth rates and disperse, may do so because dispersal causes a subsequent low growth rate in a new and unfamiliar environment. But several lines of evidence support the former interpretation. 1) On August 18, 1969, Trivers predicted that either no. 76 or no. 134 or both would soon disperse, as neither had grown in more than a month. No. 134 was never

found again, but no. 76 was next seen 11 days later on a tree 24 m across open ground from its original one. 2) Four of the eight dispersals were off of one tree which supported a large population of which those remaining behind showed a lower than average growth rate (1.67 mm/30 days). Six dispersals went to a mango, which had been almost empty until dispersing individuals reached it. These facts suggest that the dispersers may be leaving overcrowded trees for less populated ones. 3) The only disperser caught at least two weeks after dispersing showed no significant growth while dispersing, but a 3 mm growth during the six weeks after dispersal.

COPULATORY BEHAVIOR

Female and male *valencienni* appear to copulate wherever they find themselves. We cannot think of a single place (other than inside holes) where we have seen *valencienni* without also seeing them copulate there. They copulate along the thin outer branches of a bush, anywhere along the trunk of a tree, facing up or down, one centimeter off the ground to 10 m up a tree. We have even seen them copulating on the outer leaves of a mango tree 6 m off the ground. There is no evidence that females exert choice over place of copulation, as they appear to in *A. garmani* in which individuals almost invariably copulate face down on the exposed trunk of a tree (Trivers, 1976).

The most striking feature of female copulatory behavior is that females will copulate with more than one male on the same day and with the same male (and different ones) on successive days. Since it is unlikely that females are producing eggs at a rate of two or more per day, sperm competition is expected to be an important determinant of male reproductive success in *valencienni*. The key fact—that females appear to copulate more than once per egg fertilized—is supported by a number of observations, summarized here.

In the summer of 1969 only four copulations were observed involving marked females. Two of the copulations were performed on the same day, by the same female, with two different males. The female was no. 85 (58 mm). She was watched continuously from 10:35 am to 1:15 pm on July 5. For two hours she searched for food at the base of her tree, often in the company of two other marked females. At 12:25 male no. 66 (71 mm) rushed down from a branch and caught no. 85 on a small branch, where they mated. Shortly thereafter, male no. 72 (75 mm) appeared. He dewlapped at the smaller male, no. 66, who jumped on to a neighboring branch and moved up into the foliage. Although male no. 72 and female no. 85 began moving toward each other, no. 72 suddenly jumped on to a nearby branch and moved rapidly up the tree. Within 15 minutes he returned to within 30 cm of no. 85, but at 1:05 male no. 95 (81 mm) rushed down the tree and caught no. 85. Male no. 72 crouched and dewlapped at no. 95 but then jumped out of the way. No. 95 and no. 85 copulated, for her the second copulation in 40 minutes.

During the July-August, 1973 visit to the Southfield study site a special effort was made to observe copulations, and 10 females were seen to copulate more than once, usually twice within a week and sometimes twice within 24 hours. 1) Female no. 171 (60 mm) copulated on July 27 first with male no. 78 (83 mm) and then with male no. 134 (84 mm) and on the following day again with male no. 134. The first copulation took place on one tree, and the next two on a pimento (*Pimenta officianalis*) separated from the first tree by open space. 2) Female no. 24 (57 mm) copulated with male no. 45 on July 27 and on the following day with male no. 165 (73 mm). The copulations took place on contiguous, small trees. 3) Female no. 151 (60 mm) copulated on July 21 with male no. 88 (76 mm) and the following day copulated again with the same male on the same tree. 4) Female

no. LF (68 mm) copulated three days apart on the same tree with the same male (no. 134; 84 mm). 5) Two females copulated four days apart with different males on nearby trees. 6) Three more females accounted for eight more copulations, each separated by between six and nine days from a neighboring copulation. 7) Finally, one 69 mm female copulated 14 days apart on different trees with different males. In all, there were 23 multiple copulations out of a sample of 50 copulations (in which the female was marked).

How is the size of the female associated with her tendency to recopulate? The frequency with which females seen to copulate once copulated additional times was plotted as a function of size of female. Above 59 mm females tended to recopulate *less* the larger they became. In addition, the larger a female becomes the greater the span of time separating recopulations. For example, the four recopulations performed within 24 hours of an earlier copulation were all performed by females 60 mm in size or smaller. Since it is very unlikely that a female matures an egg more often than once a week, these additional copulations should be unnecessary to the female. It is tempting to suggest that the smaller a female is the less control she has over whether she copulates. Data on the percentage of females who copulate at least once (as a function of size) make it unlikely that smaller females are maturing eggs faster than large females (see Figs. 2 and 3). If these data are correct, then sperm competition is expected to be more intense in small females while female choice is expected to be a more important determinant of male reproductive success for large females.

We have timed copulations in nature and in our female choice experiments (see below), and these show remarkable consistency, averaging about 2 minutes, rarely less than 1.5 minutes or more than 2.5 minutes. There is no relationship between size of the lizards and length of copulations.

Figure 2. Frequency of copulation as a function of size (snout-vent length in mm) for females and males. These data were gathered during two weeks in June, 1971. Each fraction gives the number of copulations observed by members of the size class, divided by the number of individuals in that class.

Copulations sometimes appear to be somewhat aggressive. Males often bite females on the skinfold on the top of the neck. This may be done at the time of intromission, and the grip may be maintained during the entire copulation. Sometimes copulations are preceded by a very active stage, in which the male and female run a third of a meter or more while he attempts to cover her. These have the appearance of aggressive chases, in which the male adds an element of force to the persuasion.

Beyond this, in August 1973, Hicks observed an unusual pattern in one small adult male, which he entered in his notes under the title, "The bizarre behavior of Male no. 115, or rape and attempted forcible entry among the lower animals." No. 115 was a 57 mm male. On the 9th of August he copulated with a 50 mm female while they locked jaws. They con-

Figure 3. Frequency of copulation as a function of size (snout-vent length in mm) in females and males. Data for females include all copulations seen except those in June 1971. Data for males include all copulations except those seen in June 1971 and those seen while individual males were observed for three-hour periods.

tinued to lock jaws and fight after copulation was complete. Three days later, Hicks observed courting-fighting between the same male and a 66 mm female. No. 115 dewlapped at the female, but when he approached her closely, he would snap at her with his jaws. She chased him, he chased her. Chasing and dewlapping occurred around the trunk and up and down the tree. Finally, she half hid in a hole at the top of a dead branch. He approached within centimeters several times, once coming up right behind her. She did not move, and he seemed unable to do anything. Finally, he departed.

FEMALE CHOICE

In the highly territorial *A. garmani*, male courtship followed by female rejection has almost never been observed (Trivers, 1976). In *garmani*, females appear to choose the place to copulate and signal their accessibility by going to this place. In addition, the territorial system strongly reduces the opportunities for female choice. By contrast, female *valencienni* are exposed to many adult males each day and we have often observed male courtship and female rejection. In such a situation, a dewlapping male will typically approach a female who will move away from him. If he pursues her, she may run away from him, or she may dewlap back at him and attempt to keep her rear oriented away from him.

In the summer of 1973, we observed eleven cases of courtship and rejection involving marked *valencienni*. In four cases, the female who rejected the male was observed to copulate with that same male at some other time. In one case, she copulated with a male a day after rejecting him. In a second case, she copulated with a male two days after rejecting him. In a third case, a male courted the same female he had copulated with an hour and a half earlier, and she appeared disinterested in his second advance. Finally, a female ran from a courting male who had 16 days earlier copulated with her. These cases suggest that the female may be rejecting the time or the place as much as the male himself or that the female may succeed in rejecting some but not all of the advances of a particular male.

It seems certain that some cases of female rejection resulted because we were watching the lizards at very close quarters, but unambiguous cases of female rejection were also observed at considerable distances.

The most dramatic case of female avoidance took place on July 27, 1973. An 81 mm male was observed from 9:30 until 12:30 during which time he courted two females, both of whom appeared to reject

him. The male first courted a 61 mm female who ran from him. He did not pursue her. He then courted a 65 mm female who also ran from him. He chased the second female who continued to run from him. Finally, forced out on the end of a branch, she leapt from a height of 6 m to the ground in order to escape his attentions. He did not leap after her.

Our data on female rejection are too scanty to show whether there is any relationship between female size and tendency to reject, or between male size and tendency to be rejected, or between relative size of male and female and female tendency to reject.

To find out what criteria females use in choosing males, a series of experiments were run during the summer of 1976. At issue was whether females preferred males who differed by size or recent rate of growth. To ascertain the latter, nearly 300 lizards were captured between February and May. These individuals were then recaptured in June and July and used in choice experiments. Thus, for all individuals used size was known, and for many of these individuals, recent rate of growth to achieve that size was also known.

Initially the plan was to house each adult male in a 40-liter fish tank. All surfaces of the fish tank were painted opaque except one of the two large surfaces. The two tanks were placed next to each other, so that the males could not see each other, but so that a single female in a neighboring 80-liter fish tank could see each of the two males. The plan was then to measure the relative amount of time the female spent near each male. In actual fact, the fish tanks appeared far too confining on the individuals of both sexes. Most females spent the first hour trying to escape from the tank, moving from corner-to-corner. When they realized that they were not going to escape, they settled down in one corner and typically stayed there for dozens of minutes on end, sometimes for as long as two hours. It was difficult to convince

ourselves that females retained any interest in males under these conditions of confinement.

Two other experiments were attempted which were equally unsuccessful. In the first, females were released from a tin can onto a 3 m stick which separated at its end onto two sticks moving apart at a 90° angle. On each of these two sticks were tethered an adult male. Most females moved very slowly along the first 3 m, then chose which turn to make and ran very rapidly underneath the tethered male. Females often seemed to choose the smaller or less active of the two males, as if considering the male only as an obstacle to her own escape.

In the other kind of experiment, a series of eight small bushes were arranged in a row so that the vegetation of each bush connected its neighbors. On several of the bushes we tethered a single adult male per bush. In this experiment, females were provided a more natural and safe environment within which to choose a male. None of the females chose any of the tethered males. Most easily avoided them, and the males were rarely able to act in a natural way. Most moved as far as the tether would permit and strained against it. Some lost their grip on the substratum and hung from their tether. These had to be placed back on the bushes. In short, the females were exposed to the novel opportunity of meeting a series of restrained males, but this was of no use in discerning female choice.

In order to gain some value from our investment in capturing and recapturing the lizards, 108 experiments were run of the following sort. A female was released onto a bush which was placed in a large cleared area. Five minutes later an adult male was released onto the same bush. The two were permitted to remain together for an hour, or until they copulated, whichever came first. These experiments produced 36 copulations. It was clear from the behavior of the lizards that these experiments did not necessarily

measure a female choice alone. Since the bush on which the lizards found themselves was isolated from neighboring vegetation, the female was reluctant to leave the bush, and the male had, in effect, a partly imprisoned female to court. Females often resisted the advances of males by moving away from them, by squirreling behind a limb so as to be out of sight, and by running rapidly when pursued, but females only rarely left the bushes under male duress. Larger males copulated significantly more often than smaller males, and larger females copulated more often than smaller females. Indeed, frequency of copulation for individuals of both sex varied as a function of size in the same way as observed in nature (Figs. 2, 3). The data relating size to frequency of copulation are as follows. For males ≤ 64 mm, 0.17; for successively larger 5 mm categories: 0.25, 0.31, 0.36, 0.38, and 0.42. For females ≤ 59 mm, 0.24; for successively longer 5 mm categories: 0.30, 0.37, 0.38, and 0.40.

Most copulations took place in the second half-hour of the experiment, but there was considerable variation. This allowed us to see whether size of the male, size of the female, or relative size of the two had any influence on the time elapsed until copulation. Relative size had no effect, but larger males copulated significantly quicker and larger females copulated quicker, though not significantly so (see Fig. 4). These data do not support the possibility that relatively larger males were forcing copulations. Instead, larger females appear more willing to copulate and do so more often, while the same is true of males.

For those wishing to pursue the problem, two suggestions are made. One is to build a large natural enclosure into which are released a number of adults of both sexes. A cage 5 m tall, 4 m wide, and 2 m deep enclosing small trees and bushes might permit a population of 10 or 20 adults to be observed simultaneously. Measurements could be taken of behaviors that correlate with female rejection

in order to see whether male characteristics can be correlated with these attempts at rejections. This kind of arrangement would have the virtue of enclosing the adults in a natural setting, but would have the disadvantage that females are not exposed to clear, binary choices. Thus it would be preferable to house adult females in naturalistic cages until their behavior had settled down and then to place two contiguous male cages next to the female cage for a period of time such as an hour, to see if the female will actively demonstrate a preference for remaining near one of the two males. Her behavior can be monitored to see whether this choice represents avoidance of the one male or choice of the other.

Figure 4. Time until copulations (minutes) as a function of size (snout-vent length in mm) for males and females during the mating experiments of 1976. The regression line fitting the male points is $y = 97.02 - 0.754x$ which yields an r of 0.397 with 35 degrees of freedom ($p < 0.05$). The regression fitting the female points is $y = 94.12 - 0.8454x$, $r = 0.302$ with 35 degrees of freedom ($0.05 < p < 0.10$).

THE USE OF THE FEMALE DEWLAP

It seems incongruous that female *valencienni* should possess a dewlap. The dewlap is normally found only on the more aggressive sex, i.e., on males, yet female *valencienni* are far less aggressive than female *grahami*, *garmani*, and *lineatopus*, all of whom lack the dewlap. Furthermore, female *valencienni* rarely use their dewlaps (in contrast to males of their own and other species). To resolve this anomaly, a special effort was made in the summer of 1969 to record all cases in which females dewlapped.

In addition to the case described under "Feeding Behavior," we twice observed a female dewlapping back at a male *valencienni* who approached the female closely. Both times the female was as large or larger than the male, and in both cases the male moved off without copulating. Except for three possible cases of a female dewlapping at another female (see "Female-Female Interactions"), all other observations show females dewlapping at members of other species. Females were observed three times to dewlap at adult male *A. grahami*, two of whom appeared to be courting the females; that is, the male *grahami* were dewlapping while approaching the female *valencienni*. In addition, female *valencienni* were observed to dewlap at 1) an adult male *lineatopus* who was about 30 cm away; at 2) a small female *grahami* who was centimeters away and who leapt over the dewlapping *valencienni* and ran away; and at 3) one of us when he approached the female to within a distance of one meter to photograph her.

These scanty observations do not permit firm conclusions, but it appears that females primarily employ their dewlap to discourage courting males, of their own and closely related species.

MALE-MALE INTERACTIONS

A male is rarely seen in the presence of another male, but when two are seen to-

gether, they are always fighting, displaying aggressively, or one is trying to hide from the other. In contrast to fights in *A. lineatopus* (Rand, 1967; personal observation), *A. grahami* (personal observation) and *A. garmani* (Trivers, 1976), *valencienni* fights appear tame. Males commonly dewlap toward each other and then one male darkens in color and attempts to hide. Movements are slow and chasing is rare. Of nine fights observed in the summer of 1973 (all involving marked males), only one involved actual contact, namely jaw-locking. Of the approximately ten male-male encounters observed in previous visits, none involved any body contact.

Data from eight fights observed in 1973 were reviewed. In only one of the six fights in which the two males were within 4 mm of each other in size did the larger of the two win. In four fights, the smaller appeared victorious (as judged by which male darkened and attempted to hide), and in one the outcome was unclear. Rand (1967) has shown in *A. lineatopus* that when the smaller of two lizards wins a fight it is usually because the smaller is the territory holder and the larger is an intruder.

Although male no. 92 was 4 mm larger than male no. 45, he lost to no. 45 on two separate occasions (separated by a week) (see Fig. 5). Both fights took place within the home range of no. 45, while no. 92 commonly occupied a neighboring home range. Indeed, the only two days no. 92 was spotted within no. 45's home range were the days he was seen fighting with no. 45 (see Fig. 8). It is worth emphasizing, however, that no. 92 copulated in no. 45's home range *after* one of the fights (see Fig. 5, black dot almost touching triangle). (No. 92 was only seen to copulate once in his own home range.) This is consistent with other observations of male-male encounters: a fight does not usually result in the ouster of one of the two males; instead, one male retreats and hides, remaining in the same area to feed and, sometimes, to copulate.

Male no. 45 also appeared to vanquish

Figure 5. Male no. 45's territory during July and August, 1973. Pictured are the locations of fights (▲) and copulations of Male no. 45 (○), and copulations of other males within no. 45's territory (●).

another male who was larger than he was, no. 114 (79 mm). No. 114 was first seen in mid-July nearly 45 m from no. 45's home range; when sighted, no. 114 was copulating. No. 114 was next seen on August 10 in male no. 45's home range; he was seen there August 11 and 12 also but not seen anywhere afterwards. On August 11 he appeared to lose a long fight, involving jaw-locking, to male no. 45. Note that all three of no. 45's fights occurred at the edges of his home range (Fig. 5).

For the two other fights in which a smaller male appeared to vanquish a larger one, the relevant data are even less detailed. In both cases, the loser was seen only once, namely on the day of the fight. By contrast, one winner (male no. 172) was seen on seven different days on a set of eight neighboring trees. The second winner (male no. 107) was seen on eleven different days on two sets of trees. In both cases, the evidence suggests that the loser was either an outsider or sufficiently subordinate to be rarely sighted. Given the frequency with which many small males are resighted, we lean toward the former interpretation.

MALE HOME RANGE OVERLAP

In sharp contrast to more sexually dimorphic Jamaican anoles (*A. garmani*, Trivers, 1976; *A. lineatopus*, Rand, 1976; *A. grahami*, Trivers and Hicks, unpublished data), male home range overlap in *A. valencienni* is substantial. No male in either study area was known to occupy an exclusive area and several males were commonly seen within any one male's home range. To illustrate the extent of the overlap, we present detailed data on the home ranges of two males, one from each study area.

The Home Range of Male No. 72 (74 mm). Between June 25 and July 12 (1969) male no. 72 was seen eleven separate times on one of two connected small trees in the Maryfield study site. These two trees were part of a clump of four small connected trees. On six days, no. 72 was watched for at least an hour and seen to travel through a portion of one, or both, trees. On June 26 he copulated within his home range with female no. 46 (57 mm). Yet during this period of 17 days, five other adult males were seen within no. 72's home range and two of these males copulated with females resident within no. 72's home range.

Between July 1 and July 12 male no. 63 (73 mm) was seen seven times, each time on one of the two trees occupied also by male no. 72. On two of these occasions, no. 63 was watched for over two hours and each time he wandered throughout the two trees much as no. 72 did. On July 10 male no. 63 and male no. 72 encountered each other and no. 72 (although larger) appeared to be subordinate. Male no. 63 was about 4 m up on one of the trees when he spotted no. 72 on a stick 60 cm from the base of the tree. No. 63 dewlapped toward no. 72, then rushed down to a height of one meter and stood for several minutes looking in the direction of no. 72, who, meanwhile, had squirreled around on the stick so that the stick was between him and no. 63. No. 63 slowly started back up the tree dewlapping repeatedly. Fifteen minutes after

spotting no. 72, no. 63 had moved into the canopy out of sight, at which point no. 72 righted himself on the stick.

No. 63 was observed to court a female but was not seen to copulate before July 12. He alone of the males was seen after July 12 (although the trees were searched virtually daily). In the next two months no. 63 was seen six times on the same two trees, the last time of which he was copulating with an unmarked female.

Male no. 66 (71 mm) appeared to occupy one of the two trees contiguous with male no. 72's home range. That is, three of the four times male no. 66 was sighted, he was on this neighboring tree. The fourth time, he was seen on one of male no. 72's trees where he copulated with female no. 85 (58 mm) (see "Female Copulatory Behavior"). After the copulation, he walked onto a branch male no. 72 was sitting on. No. 72 dewlapped at him and no. 66 jumped to a neighboring branch and moved rapidly away.

Two males (63 mm and 81 mm) were seen only once each, and both were seen on one of the two trees that comprised no. 72's home range. The larger male copulated with female no. 85 on July 5 and appeared to frighten male no. 72 shortly afterwards (see "Copulatory Behavior").

In summary, for at least 16 days male no. 72 occupied a home range which he shared with a second male (no. 63) who, although smaller, appeared to be dominant over him. In addition, at least three other adult males were observed within this home range and two copulated within it. All interactions observed between the males were characterized by display and avoidance. Unfortunately, the difficulty of spotting *valencienni* in this study area does not permit us to infer anything from the absence of sightings after July 12. It is possible, for example, that after July 12 only male no. 63 remained within the two trees and that the high overlap observed prior to that day was merely temporary. Detailed data from the summer of 1973 render that an unlikely assumption, however. Throughout the

summer of 1973 the same degree of overlap was observed in all cases for which there were sufficient observations to plot home ranges. An example is the home range of male no. 45.

The Home Range of Male No. 45. Data on male no. 45 give a detailed picture of the overlap in male home ranges and the partly correlated overlap in access to females. Male no. 45 (77 mm) occupied a large home range containing nine small trees and many bushes and covering an area of about 25 square m (see Fig. 5, 6). First captured on July 7, 1973, he was watched throughout the summer visit to the Southfield study area and was last seen on August 18. He was seen on 16 different days and on five of these days his movements were followed for about three hours each day. On three of these three-hour watches, he wandered throughout about a third of his home range. On the other two three-hour watches, he restricted his movements to a corner of the home range.

Twenty-six adult females were seen at one time or another within no. 45's home

Figure 6. Male no. 45's movements in his territory during five days in which he was observed continuously each day. ● = the trees in his territory. Note that in three days he travelled widely, while in two he moved little.

range. These females were sighted a total of 51 times within no. 45's home range and a total of 42 times outside his home range. Sixteen of the 26 females were seen to copulate, and they were seen to copulate a total of 24 times (only seven of which took place outside of no. 45's home range). Male no. 45 accounted for eight of these copulations, each with a different female (see Fig. 5). He twice copulated with two females on the same day. He copulated on five of the seven days in which his movements were followed for three hours. Indeed, six of his eight copulations were seen during these 21 hours of observation.

Five of the females with whom no. 45 copulated also copulated with other males. The five copulated seven other times with a total of five different males. Of the 16 copulations with males other than no. 45, nine took place inside no. 45's home range. A comparison of the location of these nine copulations with the location of no. 45's eight copulations shows broad overlap (Fig. 5).

The degree to which females seen in a male's territory will also be seen outside his territory is shown in Figure 7. The home ranges of six females, who were seen at least once in male no. 45's territory and were sighted at least five times in total, are shown in Figure 7. As we can see, even if male no. 45 maintained exclusive sexual access to these females when they were in his territory, which he does not, they would still wander widely outside of his territory.

The variety of copulations a male may enjoy in a short period of time are suggested by observations of male no. 134 (see Table 4).

Male no. 45 shared his home range with at least ten other adult males. The overlap was considerable. For example, on August 10 four adult males (in addition to no. 45 himself) were seen within no. 45's home range. The day before, three females copulated within no. 45's home range, only one of them with no. 45. The males overlapping with male no.

45 can be divided into four categories. Three were only sighted once or twice (if twice, both sightings were within no. 45's home range). Of those sighted more than twice, two lived entirely within no. 45's home range, two lived more within no. 45's home range than outside it, and three lived primarily outside no. 45's home range (see Fig. 8).

Males completely within no. 45's home range: 1) No. 96 (72 mm) was spotted on four different days on six separate trees within no. 45's home range. No. 96 copulated with three females (on different days) only one of whom was seen to copulate with no. 45. 2) No. 115 (57 mm) was spotted on ten different days and on four different trees, all within no. 45's home range. No. 115 copulated once with a female, and she was not seen to copulate with no. 45.

Males primarily within no. 45's home range: 1) No. 164 (73 mm) was seen eight different times, of which five sightings were within no. 45's home range and three were at its edge. He copulated three times—all of them in no. 45's home range. Only one of the females was also seen to copulate with no. 45. He fought once with male no. 115 (57 mm). 2) Male no. 114 (79 mm) was seen on three successive days in no. 45's home range (August 10–12). He was not observed to copulate there. The only other time he

TABLE 4. HISTORY OF COPULATIONS FOR MALE NO. 134 (84 MM) IN SUMMER, 1973.*

	DATE	FEMALE	SIZE OF FEMALE (mm)
1.	7/22	LF	68
2.	7/25	LF	68
3.	7/26	157	70
4.	7/27	171†	60
5.	7/27	97	67
6.	7/28	171	60
7.	7/29	182	70
8.	8/5	194	62

*All took place in the same pimienta tree.

†No. 171 copulated earlier that day with male no. 78 on a nearby achee tree.

Figure 7. The overlap of Male no. 45's territory is pictured with the home ranges of six adult females observed at least once in his territory, for each of which there were at least five sightings. The mean number of sightings per female was 8.2.

was seen was nearly a month earlier, 43 m away, at which time he was seen copulating.

In summary, it is clear that adult males overlap with each other tremendously, both in space occupied (Fig. 8) and in access to members of the opposite sex (Figs. 5, 7). Commonly several males copulate with the same female and several males copulate with different females in the same space. Fights are short, mild and lead to withdrawal and hiding. Some males, at least, are highly variable in their movements, during some periods moving considerable dis-

tances, during others remaining mostly in one place (Fig. 6). Male-male competition must take an interesting and complex form in *A. valencienni* that would repay careful study.

FEMALE AGGRESSIVENESS IN A SEARCHING SPECIES

Why are female *valencienni* not aggressive? Imagine the contrary. Imagine that each female defends a territory within which to feed. Assume a female completes the search of her territory for food

Figure 8. The overlap of Male no. 45's territory with those of seven other adult males seen at least once in his territory and sighted altogether more than two times. The mean number of sightings for these seven males is 6.7.

in less than a day. She must now decide whether to research her own territory or to move into a neighbor's. To predict her behavior we should know the answers to three questions. 1) At what rate do resources within her territory renew themselves? 2) Have her neighbors completed

the search of their own territories? 3) Is she more likely to find items her neighbors missed or items she herself missed? What is the long-term variance in goodness of territory?

If resources renew themselves slowly (for example, overnight) then there will

be no strong incentive for a female to research her own territory. Indeed, under such conditions, there should be a stronger incentive to search her neighbors' territories, since 1) she knows she has completely searched her own territory and she does not know that her neighbors' territories have also been completely searched and since 2) it is more likely that she will find items her neighbors have overlooked than that she will find items she herself has overlooked (on the assumption that females show individual variability in searching behavior, e.g., in specific search images). Since the same arguments apply to other females, there is, under these assumptions, no incentive for females to guard territories once they have been searched.

FEEDING BEHAVIOR AND CRYPTICITY

For a series of reasons that stem from its feeding adaptation *valencienni* has (we believe) been strongly selected for crypticity. The fact that it is a searcher, means that individuals must constantly be on the move, thereby becoming more visible to predators. At the same time, however, they have not been selected for speed once they sight prey, so they are more vulnerable once sighted by a predator. In addition, they search by scanning very closely the surface centimeters in front of them. This narrows their field of vision considerably compared to perching lizards, making it less likely they will sight a predator early.

One could also argue that differential visibility toward potential prey may have selected for some of *valencienni*'s crypticity. If true, this should only apply to the face and neck and not to the dorsal surface of legs. It would also require that potential prey be able to respond to *valencienni* at distances of centimeters and this is not unlikely. The use of crypticity in social encounters is discussed below.

Our only actual data on predation are four observations of small *valencienni*

being eaten by other anoles. Two were hatchlings at Mayfield in August 1968, eaten apparently within an hour of hatching, one by an adult male *A. lineatopus*, the second by an adult male *A. grahami*. The others were eaten at Southfield 1) on July 21, 1973 by an 82 mm male *valencienni* and 2) on August 8, 1973 by a 56 mm male *valencienni*. In the first case of cannibalism the eaten lizard had been marked and was probably a 43 mm male. It was eaten head first. The second was unmarked and was eaten tail first, the head being bitten off without being swallowed.

COMMUNAL EGGLAYING, SEASONAL REPRODUCTION, AND PREDATION ON YOUNG

Anolis valencienni females lay eggs communally (Rand, 1976a). In Maryfield we found five communal egg deposits and in Southfield, one. These were located in holes in tree trunks from ground level to 6 m up. The number of eggs ranged from only two to well over 30. The largest was in a large hole one meter up the trunk of a mango tree. Although females often frequented this hole, and a mirror was arranged to view their activities inside the hole, egg laying was never observed. (Feeding attempts within the hole were observed, but never directed against the eggs.)

Anolis valencienni appears to breed seasonally. Although all five egg deposits contained eggs during the summertime, none contained eggs in late December and early January. Since the only egg hatched in captivity required 53 days from laying to hatching (Underwood and Williams, 1959), we conclude that no eggs are laid during at least November, December, and early January, but the periods of observation have been short enough so that copulations could have been overlooked by chance. (Except for seven days in December 1969, and eleven days in December 1973 to Jan-

uary 1974, *valencienni* has only been studied in the summertime). However, in two different summers of work at Southfield (June 1971, and July and August 1973) copulations were observed at a rate of slightly more than one per day, so that, at the very least, frequency of copulation must be strongly reduced in wintertime.

The size distribution of lizards caught strongly suggests that there is a peak in lizards hatching in the late summer and fall and that very few hatch between January and March. Of 231 lizards caught in July and August of 1973 (at Southfield) only thirteen (5.7%) were under 35 mm in size, while of 72 lizards caught in the same study site in December 1973 to January 1974, eleven (15.3%) were under 35 mm. The small lizards caught in the summer were probably laid as eggs as early as the beginning of April. A. S. Rand (personal communication) collected over 50 *valencienni* eggs during November and early December, and all of these hatched by mid-December (they are all the hatchlings on which Figure 1 is based).

Casual observations suggest that predation on hatchling *valencienni* is heavy and that hatching may be synchronous (perhaps to swamp predators). During a one hour and 40 minutes period on August 5, 1969, four hatchlings emerged from the large egg hole three feet up a mango tree at Maryfield. The first emerged at 1:50 pm, was captured and died while being measured (male = 25 mm). The second was eaten by an adult male *A. grahami* who had just entered the hole seconds before. It took the male two and a half minutes to swallow the hatchling. The third emerged at 3:10 and slowly (in short bursts of running) made its way up to 3 m up the trunk. During its last burst of running it was spotted by an adult male *A. lineatopus*, who seized it and ate it head first, requiring again two and a half minutes to consume the lizard. The fourth emerged from the hole, rested for ten minutes, then passed by an adult female *grahami* (47 mm). She saw the

hatchling but ignored it. In the next 20 minutes the hatchling made its way up to about 5 m, after which it could no longer be seen. When last seen, it was higher on the tree than where the male *lineatopus* was perched. (The male *grahami* was out of sight high in the tree.)

The synchrony of emergence (and presumably hatching) is striking. Although the hole was searched daily (before and after August 5), no other hatchlings were seen. Between August 4 and August 6, about 20 eggs appear to have hatched in this one hole. On August 6 there were no more eggs stuck to the wall but the bottom of the hole was littered with opened eggs.

One other instance of predation was seen at Maryfield. On August 9 in a different part of the study area a 62 mm adult male *grahami* caught and ate head first a very small *valencienni* who was 1.3 m above the ground on the tip of a one centimeter thick twig. One case of cannibalism was observed at Southfield. On July 21, 1973, an 82 mm male *valencienni* ate a small marked *valencienni* head first (probably no. 95, a 43 mm male).

THE EVOLUTION OF SOCIAL BEHAVIOR AND SEXUAL DIMORPHISM IN *A. VALENCIENNI*

In summary, we argue that the unusual features of *A. valencienni* probably evolved in the following fashion. At some point, individuals switched from perching to searching as a means of acquiring food. Since this new source of food was only slowly renewable within the day, females were no longer selected to maintain exclusive feeding territories, but were instead selected to wander widely and to interact nonaggressively with other females. This new style of movement rendered individuals more vulnerable to predation, since they were now continuously on the go and preoccupied with searching the substratum. Hence they became unusually cryptic.

Once a system of female territoriality had broken down, males were selected to tolerate much greater degrees of territorial overlap. This is because 1) in their search for food, males were also selected to wander widely; 2) excluding all males from a territory no longer gave exclusive access to the females residing in that territory; 3) the individuals of both sexes were more difficult to detect, thus aggressive control more difficult to impose; and 4) females, in their choice of mating partners, may have placed less emphasis on large male size than when they were a perching species.

We must confess, however, that we are ignorant concerning the role that female choice plays in generating male behavior and male size. Certainly *A. valencienni* seems, among lizard species, unusually suited to intense female choice, but we have been unable to design the proper experiment for measuring female preference. What evidence exists suggests that females may prefer larger than average males, but not strongly so. We do not know why sexual dimorphism is reduced in *A. valencienni*. In particular, we cannot discern the relative roles of male-male competition and female choice. We have been unable to penetrate the system of male-male competition. Do males move frequently, relatively large distances? Or do they patrol key crossing points, as if to waylay caravans?

ACKNOWLEDGMENTS

For help spotting and catching lizards, we thank Raymond Simpson, Glenroy Ramsey, Leeman Wynter, Veron Elliott and Paul Bartels. For hospitality in Southfield, we are grateful to Dorel Staple. For financial support we thank the American Philosophical Association,

Ernest E. Williams, and the Milton Fund (Harvard).

DEDICATION (TRIVERS)

Ernest E. Williams took me to Jamaica in 1968. He was on a collecting expedition; I was the driver. We arrived in Kingston in the evening and drove in a blinding rainstorm to the Maryfield Guest House, a decaying English great house set on three acres. The beautiful old trees and well-tended garden attracted a big population of lizards.

We began our field work over breakfast on the veranda watching *Anolis lineatopus*. The males warmed themselves in the sun and then engaged in display and aggressive encounters as they reoccupied their territories. These bright, active, little lizards reminded one almost of puppies or kids, enjoying a little social play in the early morning hours.

Ernest soon drew my attention to a more sinister species, which seemed to hide in the background. This was *Anolis valencienni*, which moved in a very distinctive fashion. Structurally, it was sufficiently aberrant to have appeared in earlier classifications as a separate genus *Xiphocircus* (Underwood and Williams, 1959). Individuals of this species seemed unusually abundant at the Guest House, and since an *Anolis* of this type had not been studied, I soon concentrated on figuring out its social system.

Ernest impressed me very warmly on that trip to Jamaica. He traveled in a very calm, quiet, unpretentious style. When we filled out our immigration cards and were asked to state our occupation, my natural impulse was to jack up the description as high as I could. I expected Ernest to do likewise. Nothing less than "Alexander Agassiz Professor of Zoology at Harvard University" seemed appropriate to the occasion. Instead he wrote, "Teacher." Simple, unpretentious, and

when need be, nonspecific or ambiguous. He also seemed fearless in his travels through Jamaica, and I much respected him for this. He neither swaggered nor scraped, but held himself calmly at all times.

In taking me to Jamaica, Ernest introduced me to more than *A. valencienni*. I have now lived nearly five years in Jamaica. I am married to a Jamaican and have four children by her. I own an acre of land and have planted 250 trees, mostly fruit. I have studied the butterflies of Southfield and collected data on an interesting symbiosis between a carpenter ant and a membracid. The trees I have planted are now beginning to attract anoles, including *A. valencienni*, and I look forward to many more years of watching.

I dedicate this paper to Ernest E. Williams with affection and respect. He was my graduate advisor and well loved at the time. Those were good years, now warmly remembered. What he introduced me to has become an enduring part of my life.

LITERATURE CITED

- HICKS, R., AND JENSSEN, T. A. 1973. New Studies on a montane lizard of Jamaica, *Anolis reconditus*. Breviora Mus. Comp. Zool. No. 404, pp. 1-23.
- LYNN, W. G., AND C. GRANT. 1940. The herpetology of Jamaica. Bulletin of the Institute of Jamaica. Science Series, No. 1.
- RAND, A. S. 1967a. The ecological distribution of the anoline lizards around Kingston, Jamaica. Breviora Mus. Comp. Zool. No. 272, pp. 1-18.
- . 1967b. Ecology and social organization in the iguanid lizard *Anolis lineatopus*. Proceedings of the United States National Museum 122 No. 3595: 1-79.
- SCHOENER, T. W. 1970. Size patterns in West Indian *Anolis* lizards. III. Sexual dimorphism in relation to differential resource utilization and sexual selection. Manuscript pp. 1-113.
- . 1971. Theory of feeding strategies. Ann. Rev. Ecol. Syst., 2: 369-404.
- SCHOENER, T. W., AND A. SCHOENER. 1971. Structural habitats of West Indian *Anolis* lizards. I. Lowland Jamaica. Breviora Mus. Comp. Zool. No. 368, pp. 1-24.
- TRIVERS, R. L. 1976. Sexual selection and resource-accurring abilities in *Anolis garmani*. Evolution, 30: 253-269.
- UNDERWOOD, G., AND E. E. WILLIAMS. 1959. The anoline lizards of Jamaica. Kingston, The Institute of Jamaica.

Conditional Relatedness, Recombination, and The Chromosome Numbers of Insects

JON SEGER¹

ABSTRACT. If two polymorphic loci are out of phase equilibrium, a homozygote at one of these loci is more highly related to its kin, at the other locus, than is an equivalent heterozygote. As a result, selection can favor (1) phenotypic responses to relative heterozygosity, and (2) increased recombination between the loci inducing these responses. Selection is expected to have these consequences only to the extent that kin strongly affect each other's fitnesses. The chromosome numbers of social insects appear to be higher, on average, than those of allied solitary species, which is consistent with this model on the assumption that chromosome numbers are selected in part for their effects on recombination.

INTRODUCTION

Coefficients of relatedness are always conditional on something. In the usual formulations they are conditional on structural relatedness, that is, on the pedigree connections between the individuals whose relatedness is to be evaluated. Full siblings are related by one half, aunt and nephew by one fourth, first cousins by one eighth, and so on, in randomly mating diploid populations of infinite size. At first it seems strange to speak of structural relatedness as something that *conditions* genetic relatedness, since we usually treat it as that which *defines* genetic relatedness. Structural relatedness is certainly one of the most important determinants of genetic

relatedness, but it is not the only one. For example, phenotypic similarity is treated as the conditioner of genetic relatedness in so-called "spotter gene" or "green beard" models. Genetic relatedness at a given locus may be conditional on both structural relatedness and gene frequency if the fitnesses of the related individuals are determined according to one of a number of nonadditive schemes (Seger, 1981, and references therein). Genetic relatedness may also be conditional on heterozygosity, in species with certain kinds of population structures (Seger, 1976). Here I show that where relatedness is conditional on heterozygosity, relatively high rates of recombination may be favored by natural selection. This idea survives a comparative test in which chromosome number is used as a proxy for the average rate of recombination.

Ernest E. Williams became my thesis adviser some time after I first saw the outlines of this argument, but some time before I first attempted to publish it. This is nearly the twentieth version of a paper that has had a long and erratic development. Ernest encouraged me to stay with it on many occasions when I found myself stalled and confused. Several important changes resulted directly from his criticisms. From time to time he asked me whether I thought the model might help to explain lizard microchromosomes. I always said that I supposed it might, but that I was still unable to see how. In fact, I found the question

¹Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, U.S.A.

extremely aggravating. Micros obviously posed a beautiful problem, closely related to all my other interests, and yet I had no idea what to do with them. Couldn't he see that? Why then did he persist in bringing them up? I now suspect that Ernest sets out quite deliberately to tie knots like this in the minds of all of his students. Those who manage to get rid of the things, by solving them, will advance science. The rest will suffer from recurring and unusually vivid flashes of ignorance, no matter what progress they may make on other problems. In either case, Ernest succeeds as a teacher. We may not have been exploited as students, but if this suspicion is correct, most of us were thoroughly outwitted.

RELATEDNESS CONDITIONAL ON HETEROZYGOSITY

Consider two loci each with two alleles (A_1 and A_0 at A, B_1 and B_0 at B). Let p_1 be the frequency of A_1 at A, and let p_2 be the frequency of B_1 at B. Let x_1 , x_2 , x_3 , and x_4 be the frequencies of the gametes A_1B_1 , A_1B_0 , A_0B_1 , and A_0B_0 , respectively. Then we can express the gametic frequencies as

$$\begin{aligned}x_1 &= p_1 p_2 + D, \\x_2 &= p_1 (1 - p_2) - D, \\x_3 &= (1 - p_1) p_2 - D, \\x_4 &= (1 - p_1) (1 - p_2) + D,\end{aligned}$$

by letting

$$D = x_1 x_4 - x_2 x_3.$$

It follows from the definition of conditional probability that

$$P(A_i = A_1 | B_j = B_1) = p_1 + D/p_2 \quad [1]$$

and

$$P(A_i = A_1 | B_j = B_0) = p_1 - D/(1 - p_2). \quad [2]$$

Thus the presence or absence of allele B_1 affects the probability that A_1 occurs on the same gamete if D is anything other than zero.

One of the classic results of population genetics theory states that D decays to zero from any initial value, in a closed randomly mating population, if all four alleles are neutral and there is any recombination between the A and B loci (Robbins, 1981; Malécot, 1948). D is traditionally referred to as the "coefficient of linkage disequilibrium," because the rate of decay depends on the tightness with which the loci are linked. But D can take nonzero values for unlinked loci, even under random mating, and so it is now increasingly referred to as the "coefficient of gametic phase disequilibrium."

Kimura (1956) and many others have studied the ways in which epistatic fitness interactions can generate nonzero values of D, at gene frequency equilibrium, in an infinite randomly mating population. Hill and Robertson (1968) were the first to show that phase disequilibrium is also generated at gene frequency equilibrium in the absence of epistasis, even between neutral loci, if the population is finite in size. The process is difficult to model, but easy to grasp intuitively. Gametes are sampled out of the parental generation, and therefore some are inevitably chosen in proportions that do not perfectly reflect the frequencies of their constituent alleles. On average there will be either an excess of "coupling" gametes (D positive) or an excess of "repulsion" gametes (D negative). These random perturbations of the gametic frequencies tend to cancel, giving an expectation of D equal to zero. But since D is very seldom *exactly* zero, the expectation of D^2 is greater than zero.

The expected magnitude of D depends strongly on the gene frequencies p_1 and p_2 . It can be shown that

$$D(p_1, p_2) = [\sigma_d^2 p_1 p_2 (1 - p_1) (1 - p_2)]^{1/2} \quad [3a]$$

gives a reasonable approximation to the expected absolute value of D if selection is weak (Seger, 1980). In equation [3a]

$$\sigma_d^2 = 1/[3+4N(c+k)-2/(2.5+N(c+2k))], \quad [3b]$$

where N is the effective population size, k is the sum of the four forward and backward mutation rates, and c is the recombination fraction (Ohta and Kimura, 1969). Since k is usually much smaller than c , mutation rates have little effect on σ_d^2 (and thus on $D(p_1, p_2)$), except for very tightly linked genes. Selection also has little effect unless it is very strong (Ohta and Kimura, 1969). Linkage increases the expected magnitude of D , but even for unlinked genes D takes values approximately 0.02, 0.10, and 0.20 times its possible maxima (given p_1 and p_2) in populations of sizes $N=1,000$, $N=50$, and $N=10$, respectively. Such population sizes appear to be typical (Jain, 1976; Lande, 1979; Levin and Kerster, 1969; Loukas *et al.*, 1979; Schaal, 1980; Schaal and Levin, 1978; Wright, 1978). Thus, although the sign of D is unpredictable unless there are epistatic fitness interactions between the A and B loci, the expected magnitude of D may be substantial, even when A and B are unlinked.

Given the conditional probabilities [1] and [2], it is easy to find the correlation between an individual's A-locus alleles, conditional on its B-locus genotype. Assign the arbitrary genic values 1 and 0 to alleles A_1 and A_0 . The joint genic distribution of the gametes that united to form the individual can be written

$$\begin{array}{cc} a_{11} & a_{10} & M_1 \\ a_{01} & a_{00} & M_0, \end{array} \quad [4]$$

where a_{11} is the probability that the individual received A_1 from both parents, $a_{10} = a_{01}$ is the probability that it received A_1 from one parent and A_0 from the other, and a_{00} is the probability that it received A_0 from both. Letting X denote the genic

value of the individual's maternal gamete, and Y that of the paternal, it follows immediately that

$$E(XY) = a_{11}$$

and

$$E(X) = E(X^2) = E(Y) = E(Y^2) = a_{11} + a_{10} = M_1.$$

Suppose the individual is known to be a heterozygote at the B locus. The probability of this event is $k=2p_2(1-p_2)$. The probability that the individual is both B_1B_0 and A_1A_1 is $2x_1x_2$. Thus the conditional probability that it is A_1A_1 , given that it is B_1B_0 , can be written as

$$\begin{aligned} E(XY) = a_{11} &= 2x_1x_2/k \\ &= 2(p_1p_2+D)[p_1(1-p_1)-D]/k \\ &= p_1^2-2D[D+p_1(2p_2-1)]/k. \end{aligned}$$

By similar reasoning

$$\begin{aligned} E(X) = M_1 &= (x_1+x_2)/2k \\ &= p_1-D(2p_2-1)/k. \end{aligned}$$

Thus

$$\text{Cov}(X, Y) = -[D^2/k][2+(2p_2-1)^2/k], \quad [5]$$

and

$$\begin{aligned} \text{Var}(X) &= p_1(1-p_1)+[D(2p_1-1)/k] \\ &[2p_1-1-D(2p_2-1)/k]. \end{aligned} \quad [6]$$

Given any intermediate values of p_1 and p_2 , $\text{Cov}(X, Y)$ is negative or zero. Thus since we expect a nonzero value of D in a finite population, we expect a negative correlation between the A-locus gametes of a B-locus heterozygote. The correlation between an individual's gametes is the same as the coancestry of its parents, so the parents of a B-locus heterozygote have a negative coancestry and are negatively related at the A locus.

Where mating is at random, there can be no overall correlation between parents at the A locus or at any other. This

clearly implies that the parents of B-locus homozygotes are positively related at the A locus, by an amount sufficient to cancel the negative relatedness between parents of B-locus heterozygotes. By means of an argument exactly like that leading to equations [5] and [6] it can be shown that

$$\text{Cov}(X,Y) = [D^2/k][2-(2p_2-1)^2/k] \quad [7]$$

and

$$\text{Var}(X) = p_1(1-p_1) + [D(2p_2-1)/k] [1-2p_1-D(2p_2-1)/k], \quad [8]$$

where $k=1-2p_2(1-p_2)$ is the overall probability of being a B-locus homozygote. In this case $\text{Cov}(X,Y)$ is always zero or positive. Thus the parents of a B-locus homozygote are positively related at the A locus, in a finite population.

This implies, but does not prove, that a B_1B_0 heterozygote will be related to its full sibling by less than one half, across the A locus, while a B_1B_1 or B_0B_0 homozygote will be related to its sibling by more than one half. The explicit demonstration is tedious but straightforward (Seger, 1980). The conclusion can be summarized by saying that A-locus relatedness is *conditional* on B-locus genotype, for full siblings in a finite population. Figure 1 shows these conditional coefficients of relatedness as a function of population size, for the case in which $p_2=0.5$ and the A and B loci are unlinked. Linkage increases the expected magnitude of D , but decreases the extent to which the conditional coefficients diverge from one half. This is illustrated in Figure 2, for populations of effective size 10, 50, and 250.

Suppose the phenotype affected by A-locus genotype has fitness effects on both ego and its sibling. Then ordinary kin selection will favor A-locus alleles that maximize ego's inclusive fitness, given the siblings' nominal relatedness of one half, and given any well defined constraint on their possible joint fitnesses.

Figure 1. Conditional relatedness as a function of population size.

The upper curve shows R^+ , a B-locus homozygote's A-locus relatedness to its sibling, as a function of N , the effective population size. The lower curve shows R^- , the corresponding relatedness of a B-locus heterozygote. The upper horizontal axis shows values of D' , the standardized coefficient of gametic phase disequilibrium, corresponding to the values of N on the lower axis. Loci A and B are unlinked, and $p_2 = 0.5$.

But we know that ego's A-locus relatedness to its sibling is actually *conditional* on ego's B-locus zygosity. Thus it seems to follow that an A-locus allele that made its phenotypic effect appropriately conditional on ego's B-locus genotype could displace all alleles with unconditional phenotypic effects. This conjecture can be shown to hold under plausible assumptions regarding the constraints on joint fitness (Seger, 1980), and can be summarized by saying that an unconditional A-locus phenotype is *not* in general an ESS against a phenotype that is conditioned on the zygotic states of other polymorphic loci (Seger, 1976).

From the point of view of the A locus, the zygotic state of B is a signal that conveys information about the A-locus re-

Figure 2. Conditional relatedness as a function of linkage between A and B.

Increasing the linkage between loci A and B increases the expected magnitude of D' , but decreases the difference between R^+ and R^- . Here $p_2 = 0.5$, and c varies between 0 (no recombination) and 0.5 (free recombination). Three different population sizes are shown. Upper curves are R^+ , lower curves are R^- , as explained in the legend to Figure 1.

latedness of ego and its sibling; genotypes B_1B_1 and B_0B_0 indicate an inflated A-locus relatedness, and B_1B_0 indicates a deflated relatedness, relative to the unconditional relatedness of one half. Thus we can refer to ego's membership in distinct *relatedness sets* which are defined by its B-locus genotype. In the example considered so far, in which there is only one signalling locus, there are only two conditional relatedness sets. But in principle there could be a large array of signalling loci and several distinct relatedness sets corresponding to different configurations of homozygosity and heterozygosity across the array of signalling loci.

Figure 2 shows that linkage between A and B reduces the distinctness of A-locus relatedness sets defined on B. Thus a gene that increased the rate of recombination between A and B would increase the distinctness of relatedness sets, and would thereby promote the spread of conditionally responding alleles at the A locus. But would selection also favor the gene for increased recombination?

SELECTION FOR INCREASED RECOMBINATION BETWEEN SIGNALLING LOCI

Consider a locus C, unlinked to either A or B, with alleles C_1 and C_0 . C_1 promotes recombination between A and B, while C_0 suppresses it. Formally, B is a signalling locus from the point of view of C, just as it is from the point of view of A, despite the fact that C is unable to bring about any phenotypic response to the zygotic state of B. If A were unlinked to B, as C is, it and C would belong to the same relatedness sets defined on B and would therefore have the same inclusive fitness interests. This implies that C_1 , which reduces linkage, might indeed be favored over C_0 . Consider a slightly different model in which there are many polymorphic B loci, all linked to each other but unlinked to A and C. In the limit of complete linkage to each other they constitute a supergene with many pseudoalleles of different degrees of complementarity. It is clear that the average homozygosity over this set of B loci carries less information about the relatedness of an individual's parents when the loci are tightly linked than it does when they recombine freely, which implies that a gene promoting recombination between the B loci could be favored.

Like the fitnesses of sex ratio phenotypes, the fitnesses of recombination phenotypes do not manifest themselves until one or more generations following

their expression. Thus there seems to be no simple way to apply genotypic covariance methods (Price, 1970; Seger, 1980, 1981) to the evolution of recombination modifiers. But positive selection for increased recombination can be demonstrated by simulation of a particular model with many signalling loci. One such simulation is described below.

The model species is a diploid hermaphrodite in which self-fertilization alternates with random mating. The population is infinitely large, but the effective population size is finite, owing to the high frequency of selfing. Offspring are produced in pairs. Each pair is given one unit of resource (x), and an individual's fitness (W) depends on the amount of resource it consumes, according to the function

$$W(x) = 1.0 - e^{-x}.$$

Given this fitness function and total shareable resource, the fitness set for sibling pairs is slightly convex (Fig. 3). One sibling, called "left," divides the unit of resource between itself and the other sibling, called "right."

Locus A controls the left sibling's division of the unit of resource, and locus C controls the rate of recombination at B, a large array of signalling loci to which neither A nor C is linked. In the simulation, A and C are treated explicitly, but B is treated in the aggregate. Typical loci in B are imagined to be highly polymorphic, such that outbred individuals are always more heterozygous than inbred individuals, if recombination between B loci is free. C_1 causes free recombination between all B loci, and C_0 suppresses recombination between them. In some runs C_1 was dominant to C_0 , and in others it was recessive.

The behavior of this genetic system can be described as follows. All of the selfed progeny of recombining parents are moderately homozygous at B, having received the same allele in egg and sperm at approximately half of their loci.

Figure 3. Fitness set for siblings.

The fitness of each sibling is determined by its consumption of resource (x), according to the function $W = 1.0 - e^{-x}$. This function is shown in (a). Given that two siblings share a total of one unit of resource, the set of their possible joint fitnesses is as shown in (b). W_L is the fitness of the "left" sibling, and W_R is the fitness of the "right" sibling, as explained in the text.

Half of the selfed progeny of nonrecombining parents are entirely homozygous, having received an intact copy of the same entire strand in egg and sperm, but half of them are only slightly homozygous, having received the parent's maternal strand in one gamete, and its paternal strand in the other. A_1 causes the left sibling to respond to its zygosity at B and is codominant with A_0 , which when homozygous causes a fixed division of the resource. Left siblings of genotypes A_0A_0 keep 0.8 units of resource and give the remaining 0.2 units to their right siblings. This is the ESS nonresponding phenotype, which was found by playing

A_0 variants against each other in the absence of A_1 . Left siblings of genotypes A_1A_0 and A_1A_1 keep relatively more for themselves when heterozygous at B (0.85 and 0.9 units, respectively), but keep less (0.75 and 0.7) when moderately or entirely homozygous at B. This level of response is derived from a simple inclusive fitness model, based on the formal relatedness of selfed siblings.

The simulation evolves as expected, when allowed to run freely for many generations. A_1 spreads from low frequencies, even in populations fixed for C_0 , the gene that blocks recombination at B. This happens because the zygosity of B contains some information about the siblings' relatedness even when there is no recombination. Consider an outbred carrier of A_1 , the responding gene. Such an individual is heterozygous at B, and therefore gives relatively little of the resource to its right sibling. This is appropriate, owing to the relatively low relatedness of outbred siblings. Now consider the inbred (selfed) carriers of A_1 . Half of them are heterozygous at B, because there is no recombination, and mistakenly act as if they were outbred, but half of them are homozygous and appropriately give a relatively large share of the resource to their right siblings. Thus carriers of A_1 respond in a way that is appropriate to their actual relatedness three quarters of the time, and this is sufficient to give them a higher average inclusive fitness than that realized by the nonresponding A_0A_0 homozygotes, in this particular model.

When C_1 and A_1 are introduced together they both spread, and their progress appears to be mutually reinforcing. They go to fixation whether C_1 is recessive or dominant and whether the C and A loci are linked or unlinked (Fig. 4). This happens because the inbred progeny of recombining parents usually carry C_1 , and those who also carry A_1 never make the inappropriate response made by half of the nonrecombinant inbred responders. C_1 increases the infor-

mation about relatedness that is conveyed by the zygosity of the signalling B loci, and thereby improves the accuracy of the response to zygosity. This benefits both C_1 and A_1 because the siblings are equally related across all loci.

This model differs in several respects from previous models for the evolution of recombination. It does not require high fecundity, environmental variation, or epistatic fitness interactions. It does not require that the recombination modifier be linked to the sites it affects, or even that there be any average fitness differences between alleles at the affected loci. But this model does require that the affected loci (the B array in the realization discussed here) exhibit the form of overdominance that is brought about by the spread of genes that promote responses to the zygosity of the affected loci.

SOCIALITY, RECOMBINATION, AND CHROMOSOME NUMBERS

Selection for phenotypic responses to zygosity should be much stronger in some species than in others. Selection should be weak where relatives seldom compete or where there is never any significant inbreeding or gametic phase disequilibrium. Both of these conditions are probably found in many species with very large effective population sizes, for example, those with planktonic larvae. Selection should be strong where relatives often compete and where conditional relatedness sets are well differentiated. These conditions are probably found in a variety of species with small effective population sizes, for example, at all stages of the life cycle in many social species, during the larval stage in species with gregarious development, and during the embryonic stage in many self-compatible hermaphrodites. The fact that predominantly self-pollinating angiosperms tend to have higher chiasma frequencies than do their predominantly cross-pollinating relatives (Lewis and

Figure 4. Gene frequency trajectories at loci controlling recombination and sibling behavior in a deterministic model.

A_1 promotes responses to zygoty, and C_1 promotes recombination. Figure 4.1a: C_1 is recessive to C_0 . Trajectories terminate at generation 1350. Recombination fractions between A and C are zero (complete linkage), 0.005, and 0.5 (free recombination) in the longest, intermediate, and shortest trajectories, respectively. Figure 4.1b: C_1 is dominant to C_0 . Trajectories terminate at generation 800. Recombination fractions are as in Figure 4.1a.

Both A_1 and C_1 spread more rapidly when linked to each other than when not linked, and both spread more rapidly when C_1 is dominant than when it is recessive. Rates of gene frequency change along the trajectories suggest that the marginal fitnesses of A_1 and C_1 change in complex ways during the course of selection. Apparent selective advantages for A_1 range between 0.003 and 0.012. Those for C_1 range between 0.003 and 0.025. C_1 benefits more from the advance of A_1 than A_1 does from the advance of C_1 , which is expected because recombination affects only the responding phenotypes (those carrying A_1).

John, 1963; Murray, 1976; Ved Brat, 1965; Zarchi *et al.*, 1972) is consistent with this expectation, but it is also consistent with other models for the evolution of recombination under selfing (Charlesworth *et al.*, 1977; Maynard Smith, 1978).

In almost all social insect species there are aspects of larval development and of adult behavior that appear to reflect the distribution of coefficients of relatedness within colonies (Hamilton, 1972; Trivers and Hare, 1976). In some species there is also intense competition between colonies (Wilson, 1971; Michener, 1974). Many species of social insects appear to have populations small enough to generate significant gametic phase disequilibrium

even under local random mating (Talbot, 1965; Wilson, 1958, 1963, 1971; Michener, 1974; Haskins and Wheldon, 1965; Haskins, 1970; Taylor, 1978). If so, Isoptera (termites) and social Hymenoptera (ants, bees, and wasps) should have rates of recombination higher on average than those of their closest solitary relatives.

Unfortunately, there are no comparative data on the recombination of markers, or even chiasma frequencies, for social insects and their relatives. But chromosome number is a major determinant of the average linkage within a genome, and chromosome numbers are known from many species of insects. Loci at the opposite ends of a long chromo-

some may recombine almost freely, if several crossovers usually occur between them. Nonetheless, they remain somewhat linked. A fission occurring between them reduces their linkage all the way to zero. More importantly, other pairs of loci at map positions between them but on opposite sides of the point of fission, which were formerly significantly linked, also become unlinked. Thus given an approximately constant number of crossovers per genome, the average linkage between loci is a strongly decreasing function of the chromosome number. This implies that chromosome number can be used as a proxy for the rate of recombination in comparative studies, as long as large numbers of species are compared.

As expected, the average chromosome numbers of eusocial taxa are consistently higher than those of the most closely related nonsocial taxa for which chromosome data exist (Sherman, 1979). Formicidae, Apidae, and Vespidae represent independent derivations of eusociality within the aculeate Hymenoptera (Wilson, 1971). Although only a few solitary aculeates have been karyotyped, their chromosome numbers are similar to those of the entirely solitary Symphyta and Parasitica (Table 1). This suggests that high numbers have evolved independently in the social families. The Isoptera are entirely eusocial, so no close comparisons are possible in their case. But their average chromosome number exceeds those of the other orthopteroid orders Blattodea, Orthoptera, Mantodea, Dermaptera, and Embioptera (Table 1).

It should be emphasized that the postulated selection in favor of increased recombination has nothing to do with sociality as such. Sociality is merely being used as an indicator of situations in which interactions are expected to be strongly conditioned on relatedness. In such situations natural selection should favor phenotypic responses to individual zygosity, if at the same time the population structure is such as to create signifi-

cant gametic phase disequilibrium. Selection in favor of increased recombination is thus an expected secondary consequence of the establishment of systems of responding loci. It is therefore expected in some solitary species, for example, ones in which extended juvenile dependency leads to parent-offspring conflict over the allocation of resources (Trivers, 1974), and it is not expected in some social species, for example, those with very large effective population sizes. The range of haploid numbers in ants (3–46) exceeds that in all other Hymenoptera. If the argument made here is correct, some of this variation should be associated with differences of population structure. Unusually high chromosome numbers occur in *Rhytidoponera*, *Myrmecia*, and *Nothomyrmecia*. Aptery, brachyptery, and patchy distributions are common in these genera (Haskins and Wheldon, 1965; Haskins, 1970; Taylor, 1978), suggesting that many of their species typically experience small effective population sizes.

The average chromosome numbers of six species of blattid roaches believed to exhibit larval aggregation are 60% higher than the chromosome numbers of ten species believed not to exhibit larval aggregation (Sherman, 1979).

It is difficult to assign levels of statistical significance to the observed patterns of chromosome number variation, because it cannot be assumed that the chromosome numbers of related species are statistically independent. Thus it is not legitimate to perform an analysis of variance on the chromosome numbers of species, as was done by Sherman (1979). The analysis given here treats genus averages as primary observations. This improves matters somewhat, but since genera tend to be correlated within families, as do species within genera, genera cannot properly be treated as independent points in a comparison between families. Thus the significance levels given in the legend to Table 1 should be taken only as indicators of the relative strengths

TABLE 1. AVERAGE HAPLOID CHROMOSOME NUMBERS OF SOCIAL INSECTS AND THEIR ALLIES.*

	N	$\bar{n}$	s.d.
Orthopteroids			
DERMAPTERA	20	13.38	5.92
EMBIOPTERA	4	10.83	0.41
ORTHOPTERA	33	10.84	2.72
PHASMATODEA	40	20.14	4.45
MANTODEA	69	13.20	2.44
BLATTODEA	64	19.05	6.62
Blattidae	7	18.03	5.90
ISOPTERA*	27	20.71	1.95
Hymenoptera			
SYMPHYTA	27	9.41	3.96
APOCRITA			
PARASITICA	19	8.51	2.38
ACULEATA			
Vespoidea			
Formicidae*	69	15.76	6.82
Eumenidae	2	7.63	0.53
Vespidae*	2	14.30	2.40
Sphecoidea			
Halictidae	4	11.50	5.26
Anthophoridae	3	13.67	4.04
Megachilidae	1	16.00	—
Apidae*	18	16.23	2.86

*Sources of data are listed in Literature Cited B.

Social taxa are indicated in the table by an asterisk. Most of the analysis was conducted using genus averages rather than species counts, to increase the independence and representativeness of sample points, and to increase the normality of the resulting distributions. Thus N is the number of genera in a sample, and $\bar{n}$ is the mean of the genus mean haploid chromosome numbers. The average for Orthoptera, however, represents modal numbers for 28 families and subfamilies of Caelifera (based on hundreds of studied species [White, 1973]), and average numbers for five families of Ensifera (based on about 70 genera [Makino, 1951]). Data for Isoptera include nine unpublished species counts in five genera of Kalotermitidae (P. Luykx, personal communication). One-tailed U-tests were applied to comparisons of interest (t-tests with d.f. adjusted for unequal variances were also applied and gave similar results).

Isoptera exceed the blattid roaches, to whom they are believed to be most closely related (McKittrick, 1965), but the difference is not significant. The comparison with Blattodea as a whole is significant ($p < 0.005$), as are those with Orthoptera, Mantodea, Dermaptera ($p < 0.0005$), and Embioptera ($p < 0.001$). The comparison with Phasmatodea is almost significant ($p < 0.1$).

Formicidae (ants) and Apidae (eusocial bees) each exceed both Parasitica (chalcids, ichneumons, gall wasps) and Symphyta (sawflies, horntails) ($p < 0.0005$). The small sample of halictid, antho-

phorid, and megachilid bees consists of five species with haploid numbers of 16, and three species with haploid numbers of 6, 8, and 9. The species with low numbers are communal to quasisocial (Michener, 1974), and three of the species with high numbers are solitary, contrary to expectation. Owing to the complexity of social evolution in the bees, a much larger and more representative sample of chromosome numbers is required to determine whether in fact they exhibit a pattern different from the one seen elsewhere in the Hymenoptera.

The most appropriate comparison is that between the Vespidae (eusocial paper wasps) and the closely related Eumenidae (solitary potter wasps). Their average numbers diverge in the expected direction, but too few genera have been studied to allow a statistical test of the generic averages. If all the published species counts (7 and 5, respectively) are used, the difference is significant ($U=5$, $p=0.024$, $t=2.8$, 8 adjusted d.f., $p < 0.025$). Formicidae exceed Eumenidae ($p < 0.005$) but not Vespidae, using species counts for the wasps.

of different associations, not as statistical tests.

A conservative test of the overall pattern can be carried out by comparing the signs of the differences between groups at the same taxonomic levels. There are at least six relevant comparisons. Apidae exceed all other bees. Vespidae exceed Eumenidae. Aculeata as a whole, who are mainly social in this sample, exceed Parasitica. Isoptera exceed Blattodea. *Rhytidoponera*, *Myrmecia*, and *Nothomyrmecia* exceed other ants. Gregarious blattids exceed nongregarious blattids. The comparison between Formicidae and Eumenidae could be added to this list, as could separate comparisons between Apidae and the other families of bees, but these will be omitted in the interests of conservatism. The probability that six independent comparisons all fall in the same unspecified direction is 0.03125, and the probability that they all fall in the same specified direction is 0.015625. Thus it appears that an association does exist between high chromosome number and circumstances favorable to the evolution of responses to zygoty.

DO BEES HAVE GREEN SETAE?

Sherman (1979) was the first to note the association between eusociality and high chromosome numbers. His explanation for the association, which differs from the one given here, can be summarized as follows. Parent and offspring are always related by exactly one half, because the parent transmits to the offspring exactly one member of each of its pairs of homologous chromosomes. Full siblings also expect to share half of their chromosomes identical by descent through the parents, but owing to the uncertainties of meiosis this expectation has a finite variance. Some siblings have more than half of their DNA in common, identical by descent, and others have less than half. Thus some siblings are related by more than one half, and others by less. If an individual could discriminate between the two kinds of siblings, favoring those to which it was related by more than one half, it could increase its inclusive fitness. In a social insect colony such discrimination is likely to work against the reproductive interests of the queen and her mate. The scope for such discrimination is a function of the variance of actual relatedness between siblings, and the variance of relatedness is an inverse function of the number of chromosome pairs. It is in the queen's interest to reduce the variance of relatedness as far as possible, so as to create a situation in which workers are selected to minister to their reproductive siblings "on the basis of their need rather than kinship" (Sherman, 1979). Increasing the chromosome number is probably one of the simplest and most effective ways to reduce the variance of relatedness. Thus chromosome numbers are expected to increase in eusocial lineages and in others where siblings interact intensely.

This argument turns on two critical assumptions. First, the variance of the proportion of DNA held in common identical by descent corresponds to a variance of relatedness on which kin selection

could act. Second, although an increased variance of relatedness would be in the offspring's interest, a reduced variance will evolve because that is in the queen's interest. This second assumption is problematic, but it will not be considered here in any detail. The first assumption is almost certainly incorrect, at least as stated by Sherman (Dawkins, 1979). The idea seems to arise naturally from the identity-by-descent view of relatedness, which emphasizes amounts rather than covariances. The problem with it is that identity at one locus does not predict identity at another. As Dawkins puts it, "If I want to guess whether your hand of cards contains the ace of spades I would be quite wrong to say: I already know you have the 2, 3, 5, 6, 7, 9, 10, Jack and King of spades; therefore you have a strong hand in spades; therefore you probably have the ace!" A rational gene that controlled discrimination would gain no information regarding its presence in a sibling of its bearer by knowing that the sibling, like the gene's own bearer, had a green beard. If the discrimination gene were tightly linked to the green beard gene the argument might hold, except that selection at all unlinked loci would oppose the favorable discrimination. Few theoretical arguments are affected by the fiction of the single omnipotent gene, but this appears to be one that is. Any act as complex as a comparison of particular details of one's own and one's sibling's phenotypes must depend directly on scores or hundreds of genes, and indirectly on thousands. It is asking a lot to expect them all to be firmly linked to the same phenotypic marker and to be protected from interference on the part of genes elsewhere in the genome.

It was pointed out above that correlations of allelic state arise through drift in finite populations, and that these correlations can drive the evolution of phenotypic responses to zygosity. These same correlations can be shown to underlie a consistent version of Sherman's argument for the evolution of discrimination

based on phenotypic similarity. But as will be seen, this version of the argument may have implications somewhat different from those suggested by Sherman.

Consider a genetic system consisting of two unlinked loci, each with two alleles. Locus B determines a visible phenotypic attribute and has two distinct, fully penetrant, codominant alleles, B_1 and B_0 . Locus A determines a response to the B-locus similarity of its bearer and another individual taken at random from the local population. With an adjustment for structural relatedness the argument can easily be adapted to the case of siblings, but the case of unrelated individuals is both simpler and more general than that of structurally related individuals. Locus A can be taken to represent every locus in the genome that is unlinked to B. We seek the A-locus relatedness of two individuals, conditioned on their B-locus similarity.

Let p_1 be the frequency of A_1 , and let p_2 be the frequency of B_1 . Then from equations [1] and [2] we have that

$$q_1 \equiv P(A_1=A_1|B_j=B_1) = p_1 + D/p_2$$

and

$$q_2 \equiv P(A_1=A_1|B_j=B_0) = p_1 - D/(1-p_2),$$

where D is the coefficient of gametic phase disequilibrium. Define as "alike" those pairs of individuals in which both are B_1B_1 or both are B_0B_0 . Assuming Hardy-Weinberg proportions at both loci, and assuming that individuals meet at random, the probability that two individuals are both B_1B_1 is clearly

$$f_1 = p_2^4,$$

while the probability that both are B_0B_0 is

$$f_0 = (1-p_2)^4.$$

Consider the A-locus genotypes of the B_1B_1 pair. The probability of finding A_1 on either chromosome at the A locus in

either individual is just q_1 . Thus the probability that both individuals are A_1A_1 is q_1^4 . Similarly, if both individuals are B_0B_0 the probability that they are also both A_1A_1 is q_0^4 . Thus if two individuals are known to be "alike" at B, the probability that they are both A_1A_1 is

$$a_{22} = (f_1 q_1^4 + f_0 q_0^4)/k,$$

where $k=f_1+f_0$. The probability a_{22} is the first element of the joint genotypic distribution

$$\begin{array}{cccc} a_{22} & a_{21} & a_{20} & M_2 \\ a_{12} & a_{11} & a_{10} & M_1 \\ a_{02} & a_{01} & a_{00} & M_0. \end{array} \quad [9]$$

Proceeding in this manner it is easy to show that the other elements of the joint genotypic distribution can be represented as

$$\begin{aligned} a_{21} &= a_{12} = 2(f_1 c_1 q_1^2 + f_0 c_0 q_0^2)/k, \\ a_{11} &= 4(f_1 c_1^2 + f_0 c_0^2)/k, \\ M_2 &= (f_1 q_1^2 + f_0 q_0^2)/k, \\ M_1 &= 2(f_1 c_1 + f_0 c_0)/k, \end{aligned}$$

where $c_1=q_1(1-q_1)$ and $c_0=q_0(1-q_0)$. M_0 and the remaining a_{ij} are easily found by subtraction. Substitution of the key elements of [9] into

$$R = [a_{22} + a_{21} + \frac{1}{4}a_{11} - (M_1 + \frac{1}{2}M_2)^2] / [M_1 M_0 + \frac{1}{4}M_2(1-M_2)]$$

(Seger, 1980) gives R^+ , the A-locus relatedness of two randomly chosen individuals who are "alike" at B. If both loci are polymorphic and if D is nonzero then R^+ is positive (Table 2).

Define as "unlike" those pairs of individuals in which one is B_1B_1 and the other is B_0B_0 . Define as "somewhat" those pairs in which at least one individual is the heterozygote B_1B_0 . Then proceeding as above it is easy to find R^- , the relatedness of "unlike" pairs, and R^s , the relatedness of "somewhat" pairs. These are both negative.

Like the conditional coefficients dis-

TABLE 2. CONDITIONAL RELATEDNESS IN A SPOTTER GENE MODEL.*

D'=.05		P ₂		
		.1	.3	.5
P ₁	.1	.00001	.00076	.00499
		-.00271	-.00038	0
		-.01430	-.00596	-.00499
	.3	.00001	.00075	.00499
		-.00269	-.00038	0
		-.01390	-.00594	-.00499
	.5	.00001	.00075	.00499
		-.00269	-.00038	0
		-.01385	-.00594	-.00499
D'=.1		P ₂		
		.1	.3	.5
P ₁	.1	.00003	.00309	.01980
		-.01115	-.00154	0
		-.06279	-.02398	-.01980
	.3	.00003	.00302	.01980
		-.01089	-.00153	0
		-.05575	-.02361	-.01980
	.5	.00003	.00301	.01980
		-.01086	-.00153	0
		-.05501	-.02357	-.01980

*The upper number in each set of three is the A-locus relatedness of two randomly chosen individuals that are "alike" at the B locus. The lower number is the relatedness of "unlike" pairs, and the middle number is the relatedness of "somewhat" pairs. Conditional relatedness is shown for three different frequencies of A₁ and B₁, at two different values of D', the standardized coefficient of gametic phase disequilibrium. Each distribution is symmetrical about both p₁=0.5 and p₂=0.5.

cussed in the introduction, R⁺ and R⁻ diverge as D increases (Table 2). But R⁺ and R⁻ also diverge as the A and B loci become more tightly linked, unlike the conditional coefficients discussed in the introduction, which converge on the unconditional coefficient as linkage tightens. This implies that conflicts might arise between A-loci closely linked to B, and A-loci unlinked to B, because the former will tend to be farther out of phase equilibrium with B than will the

latter. In view of this it is hard to see whether a high rate of recombination looks more like a way to frustrate discrimination than it does like a way to promote it. By reducing the expected gametic phase disequilibrium between B and a typical A locus, recombination reduces the difference between R⁺ and R⁻, thereby reducing the potential advantage of discrimination. But recombination also increases the fraction of the genome that is effectively unlinked to any given B locus. By increasing the similarity of the different values of R⁺ evaluated at different loci, recombination might well increase the ease with which mechanisms inducing discrimination could evolve.

Phenotypic discrimination mechanisms of the form envisioned by Sherman and others require that an individual be able to assess its *own* similarity to another individual. Recent experimental work on discrimination by means of individual odor differences in *Lasioglossum* demonstrates that these bees compare strangers to the aggregate of their nest-mates, and that they apparently do not perceive their own odors (Greenberg, 1980; C. D. Michener, personal communication).

CONCLUSIONS

It is possible that high chromosome numbers are caused by some feature of sociality that has nothing to do with kinship. For example, the genomes of social insects might be larger than those of solitary insects, and there might be an optimum distribution of chromosome sizes, at least within particular lineages. If this were true, chromosome number and genome size would correlate positively, but several lines of evidence suggest that they do not. Chromosome numbers and absolute DNA values for 45 species of insects representing six orders do not correlate either within orders or between them (Fig. 5). Imai, Crozier, and Taylor

Figure 5. Relationship between genome size and chromosome number in six orders of insects.

Genome size is expressed as picograms of DNA per haploid genome. The orders are Orthoptera (●), Hemiptera (▼), Coleoptera (◇), Lepidoptera (△), Diptera (○), and Hymenoptera (□). Spearman rank correlations are nonsignificant within Orthoptera (0.097), Diptera (0.006), and Hymenoptera (zero). The entire sample is significantly correlated ($r_s=0.34$, $p<0.05$) because it is dominated by Orthoptera and Diptera, whose ranges do not overlap on either variable. Using mean values for each of the six orders, the overall correlation is negative and nonsignificant ($r_s=-0.086$).

There is a large average difference in genome size between the hemimetabolous and holometabolous insects in this sample (represented by filled and by open symbols, respectively). The only social species represented (*Apis mellifera* and *A. cerana*, $n=16$, 0.19 pg) have genomes that appear to be small even for Hymenoptera. Other Hymenoptera are *Megachile rotundata* ($n=16$, 0.3 pg), *Habrobracon juglandis* and *H. serinopae* ($n=10$, 0.16 pg), and *Mormoniella vitripennis* ($n=5$, 0.33 pg).

Sources of data are listed in Literature Cited B.

(1977) optically measured chromosome lengths in 16 species of myrmecine and ponerine ants with haploid numbers of four to 42. They found that average chromosome length is inversely proportional to chromosome number, and concluded that the diversity of chromosome number in these subfamilies does not reflect any large or systematic differences

of genome size. Data for angiosperms (Bennett and Smith, 1976) and for vertebrates (White, 1973) also fail to suggest an inherent relationship between genome size and chromosome number.

The first argument outlined here implies that selection will favor relatively high overall rates of recombination in species where it also favors phenotypic responses to heterozygosity. Such species are characterized by certain behavioral, ecological, and demographic parameters. To the extent that sociality indicates the required parameters, and to the extent that chromosome number indicates the rate of recombination, social insects are expected to have elevated chromosome numbers, as they do on average. The second argument, that of Sherman (1979), also leads to the expectation that social insects will have elevated rates of recombination. The two arguments invoke nearly identical sets of causal parameters and thus make similar comparative predictions. There seems to be no way to distinguish between them on the basis of available evidence.

ACKNOWLEDGMENTS

I thank R. L. Trivers for encouragement and for many stimulating discussions during the early stages of this work. I thank S. H. Bartz, M. E. Etter, W. D. Hamilton, P. H. Harvey, R. C. Lewontin, J. Maynard Smith, J. W. Stubblefield, E. E. Williams, and the anonymous reviewer of a previous manuscript for helpful criticisms of various kinds. I thank P. Luykx and R. M. Syren for permission to use their unpublished termite chromosome numbers. This work was supported in part by National Institutes of Health Grant GM07620-02 to Harvard University.

NOTE ADDED IN PROOF

Tooby (J. theor. Biol., 97: 557-576 [1982]) has recently argued that eusocial

insects are expected to have high rates of recombination (and thus high chromosome numbers) because they live at high population densities and should therefore be subject to particularly intense selection for resistance to rapidly evolving pathogens.

LITERATURE CITED A

(TEXT)

- BENNETT, M. D., AND J. B. SMITH. 1976. Nuclear DNA amounts in angiosperms. *Phil. Trans. Roy. Soc. Lond. B*, **274**: 227-274.
- CHARLESWORTH, D., B. CHARLESWORTH, AND C. STROBECK. 1977. Effects of selfing on selection for recombination. *Genetics*, **86**: 213-226.
- DAWKINS, R. 1979. Twelve misunderstandings of kin selection. *Z. Tierpsychol.*, **51**: 184-200.
- HAMILTON, W. D. 1972. Altruism and related phenomena, mainly in social insects. *Ann. Rev. Ecol. Syst.*, **3**: 193-232.
- HASKINS, C. P. 1970. Researches in the biology and social behavior of primitive ants. In L. R. Aronson, *et al.* (eds.), *Development and Evolution of Behavior*. Freeman.
- HASKINS, C. P., AND R. M. WHELDEN. 1965. "Queenlessness," worker sibship, and colony versus population structure in the formicid genus *Rhytidoponera*. *Psyche*, **72**: 87-112.
- HILL, W. G., AND A. ROBERTSON. 1968. Linkage disequilibrium in finite populations. *Theoret. Appl. Genet.*, **38**: 226-231.
- IMAI, H. T., R. H. CROZIER, AND R. W. TAYLOR. 1977. Karyotype evolution in Australian ants. *Chromosoma*, **59**: 341-393.
- JAIN, S. K. 1976. Patterns of survival and microevolution in plant populations. In S. Karlin and E. Nevo (eds.), *Population Genetics and Ecology*. Academic Press.
- KIMURA, M. 1956. A model of a genetic system which leads to closer linkage by natural selection. *Evolution*, **10**: 278-287.
- LANDE, R. 1979. Effective deme sizes during long-term evolution estimated from rates of chromosomal rearrangement. *Evolution*, **33**: 234-251.
- LEVIN, D. A., AND H. KERSTER. 1969. Density-dependent gene dispersal in *Liatris*. *Am. Nat.*, **103**: 61-73.
- LEWIS, K. R., AND B. JOHN. 1963. *Chromosome Marker*. Little, Brown.
- LOUKAS, M., C. B. KRIMBAS, AND Y. VERGINI. 1979. The genetics of *Drosophila subobscura* populations. IX. Studies on linkage disequilibrium in four natural populations. *Genetics*, **93**: 497-523.
- MALÉCOT, G. 1948. *Les Mathématiques de l'Hérédité*. Masson et Cie.
- MAYNARD SMITH, J. 1978. *The Evolution of Sex*. Cambridge University Press.
- MICHENER, C. D. 1974. *The Social Behavior of Bees*. Harvard University Press.
- MURRAY, B. G. 1976. The cytology of the genus *Briza* L. (Gramineae) II. Chiasma frequency, polyploidy and interchange heterozygosity. *Chromosoma*, **57**: 81-93.
- OHTA, T., AND M. KIMURA. 1969. Linkage disequilibrium at steady state determined by random genetic drift and recurrent mutation. *Genetics*, **63**: 229-238.
- PRICE, G. R. 1970. Selection and covariance. *Nature*, **227**: 520-521.
- ROBBINS, R. B. 1918. Some applications of mathematics to breeding problems. II. *Genetics*, **3**: 73-92.
- SCHAAL, B. A. 1980. Measurement of gene flow in *Lupinus texensis*. *Nature*, **284**: 450-451.
- SCHAAL, B. A., AND D. A. LEVIN. 1978. Morphological differentiation and neighborhood size in *Liatris cylindracea*. *Amer. J. Bot.*, **65**: 923-928.
- SEGER, J. 1976. Evolution of responses to relative homozygosity. *Nature*, **262**: 578-580.
- _____. 1980. Ph.D. thesis. Department of Biology, Harvard University.
- _____. 1981. Kinship and covariance. *J. Theor. Biol.*, **91**: 191-123.
- SHERMAN, P. W. 1979. Insect chromosome numbers and eusociality. *Am. Nat.*, **113**: 925-935.
- TALBOT, M. 1965. Populations of ants in a low field. *Insectes Sociaux*, **12**: 19-48.
- TAYLOR, R. W. 1978. *Nothomyrmecia macrops*: a living-fossil ant. *Science*, **210**: 979-985.
- TRIVERS, R. L. 1974. Parent-offspring conflict. *Amer. Zool.*, **14**: 249-264.
- TRIVERS, R. L., AND H. HARE. 1976. Haplodiploidy and the evolution of the social insects. *Science*, **191**: 249-263.
- VED BRAT, S. 1965. Genetic systems in *Allium* III. Meiosis and breeding systems. *Heredity*, **20**: 325-339.
- WHITE, M. J. D. 1973. *Animal Cytology and Evolution*, 3rd edn. Cambridge University Press.
- WILSON, E. O. 1958. Patchy distributions of ant species in New Guinea rain forests. *Psyche*, **65**: 26-38.
- _____. 1963. Social modifications related to rareness in ant species. *Evolution*, **17**: 249-253.
- _____. 1971. *The Insect Societies*. Harvard University Press.
- WRIGHT, S. 1978. *Evolution and the Genetics of Populations*, Vol. 4. University of Chicago Press.
- ZARCHI, Y., G. SIMCHEN, J. HILLEL, AND T. SCHAAP. 1972. Chiasmata and the breeding system in wild populations of diploid wheats. *Chromosoma*, **38**: 77-94.

LITERATURE CITED B

(FIGURE 5 AND TABLE 1)

- COHEN, S., AND L. M. ROTH. 1970. Chromosome numbers of the Blattaria. *Ann. Ent. Soc. Am.*, **63**: 1520-1547.
- CRAIN, W. R., E. H. DAVIDSON, AND R. J. BRITTEN. 1976. Contrasting patterns of DNA sequence arrangement in *Apis mellifera* (Honeybee) and *Musca domestica* (Housefly). *Chromosoma*, **59**: 1-12.
- CROZIER, R. H. 1975. Animal Cytogenetics 3: Insecta 7: Hymenoptera. Gebrüder Borntraeger.
- DANEHOLT, B., AND J.-E. EDSTROM. 1967. The content of deoxyribonucleic acid in individual polytene chromosomes of *Chironomus tentans*. *Cytogenetics*, **6**: 350-356.
- EFSTRATIADIS, A., W. R. CRAIN, R. J. BRITTEN, E. H. DAVIDSON, AND F. C. KAFATOS. 1976. DNA sequence organization in the lepidopteran *Antheraea pernyi*. *Proc. Natl. Acad. Sci. USA*, **73**: 2289-2293.
- FAHRENHORST, H. 1977. Nachweis übereinstimmender Chromosomen-zahlen (N=16) bei allen 4 *Apis*-arten. *Apidologie*, **8**: 89-100.
- FONTANA, F., AND M. AMORELLI. 1975. Chromosome complement of *Kaloterme flavicollis* Fabr. (Isoptera: Kalotermitidae). *Boll. Zool.*, **42**: 99-100.
- FOX, D. P. 1970. A non-doubling DNA series in somatic tissues of the locusts *Schistocerca gregaria* (Forskål) and *Locusta migratoria* (Linn.). *Chromosoma*, **29**: 446-461.
- GALL, J. G., H. C., MACGREGOR, AND M. E. KIDSTROM. 1969. Gene amplification in the oocytes of dytiscid water beetles. *Chromosoma*, **26**: 169-187.
- GLANCEY, B. M., M. K. ST. ROMAIN, AND R. H. CROZIER. 1976. Chromosome numbers of the red and black imported fire ants, *Solenopsis invicta* and *S. richteri*. *Ann. Ent. Soc. Am.*, **69**: 469-470.
- GOODPASTURE, C. 1974. Karyology and taxonomy of some species of eumenid wasps. *J. Kansas Ent. Soc.*, **47**: 364-372.
- GOODPASTURE, C., AND E. E. GRISSSELL. 1975. A karyological study of nine species of *Torymus* (Hymenoptera: Torymidae). *Can. J. Genet. Cytol.*, **17**: 413-422.
- HINDS, T. R., AND R. W. NEWBURGH. 1972. The DNA of the thoracic muscle of *Manduca sexta*. *Biochim. Biophys. Acta*, **281**: 514-521.
- HUNTER, K. W., AND A. C. BARTLETT. 1975. Chromosome number of the parasitic encyrtid *Copidosma truncatellum* (Dalman). *Ann. Ent. Soc. Am.*, **68**: 61-62.
- IMAI, H. T., R. H. CROZIER, AND R. W. TAYLOR. 1977. Karyotype evolution in Australian ants. *Chromosoma*, **59**: 341-393.
- JOHN, B., AND G. M. HEWITT. 1966. Karyotype stability and DNA variability in the Acrididae. *Chromosoma*, **20**: 155-172.
- JORDAN, R. A., AND R. W. BROSEMER. 1974. Characterization of DNA from three bee species. *J. Insect Physiol.*, **20**: 2513-2520.
- JOST, E., AND M. MAMELI. 1972. DNA content in nine species of Nematocera with special reference to the sibling species of the *Anopheles maculipennis* group and the *Culex pipiens* group. *Chromosoma*, **37**: 201-208.
- LAGOWSKI, J. M., M.-Y., WANG YU, H. S. FORREST, AND C. D. LAIRD. 1973. Dispersity of repeat DNA sequences in *Oncopeltus fasciatus*, an organism with diffuse centromeres. *Chromosoma*, **43**: 349-373.
- LAIRD, C. D., AND B. J. MCCARTHY. 1969. Molecular characterization of the *Drosophila* genome. *Genetics*, **63**: 865-882.
- LIMA-DE-FARIA, A., T. GUSTAFSSON, AND H. JAWORSKA. 1972. Amplification of ribosomal DNA in *Acheta*. II. The number of nucleotide pairs of the chromosomes and chromomeres involved in amplification. *Hereditas*, **73**: 119-142.
- MAKINO, S. 1951. An Atlas of the Chromosome Numbers in Animals, 2nd edn. Iowa State University Press.
- MAXWELL, D. E. 1958. Sawfly cytology with emphasis upon the Diprionidae (Hymenoptera: Symphyta). *Proc. Tenth Int. Cong. Ent.*, **2**: 961-978.
- McKITTRICK, F. A. 1965. A contribution to the understanding of cockroach-termite affinities. *Ann. Ent. Soc. Am.*, **58**: 18-22.
- RASCH, E. M. 1974. The DNA content of sperm and hemocyte nuclei of the silkworm, *Bombyx mori* L. *Chromosoma*, **45**: 1-26.
- RASCH, E. M., H. J. BARR, AND R. W. RASCH. 1971. The DNA content of sperm of *Drosophila melanogaster*. *Chromosoma*, **33**: 1-18.
- RASCH, E. M., J. D. CASSIDY, AND R. C. KING. 1977. Evidence for dosage compensation in parthenogenic Hymenoptera. *Chromosoma*, **59**: 323-340.
- ROBERTS, B., AND S. ROBERTS. 1972. The DNA content of testes and malpighian tubule nuclei of *Tricholioprocta impatiens* (Sarcophagidae, Diptera). *Chromosoma*, **39**: 83-91.
- SMITH, S. G., AND N. VIRKKI. 1978. Animal Cytogenetics 3: Insecta 5: Coleoptera. Gebrüder Borntraeger.
- SWIFT, H., AND R. KLEINFELD. 1953. DNA in grasshopper spermatogenesis, oogenesis, and cleavage. *Physiol. Zool.*, **26**: 301-311.
- SYREN, R. M., AND P. LUYKX. 1977. Permanent segmental interchange complex in the termite *Incisitermes schwarzi*. *Nature*, **266**: 167-168.
- VINKE, P. P., AND J. P. TILQUIN. 1978. A sex-linked

- ring quadrivalent in Termitidae (Isoptera). *Chromosoma*, **67**: 151-156.
- WHITE, M. J. D. 1973. *Animal Cytology and Evolution*, 3rd edn. Cambridge University Press.
- . 1976. *Animal Cytogenetics* 3: Insecta 2: Blattodea, Montodea, Isoptera, Grylloblattodea, Phasmatodea, Dermaptera and Embioptera. Gebruder Borntraeger.
- WHITE, M. J. D., AND G. C. WEBB. 1968. Origin and evolution of parthenogenetic reproduction in the grasshopper *Moraba virgo* (Eumasticidae: Morabinae). *Aust. J. Zool.*, **16**: 647-671.
- WILMORE, P. J., AND A. K. BROWN. 1975. Molecular properties of orthopteran DNA. *Chromosoma*, **51**: 337-345.

Is *Sphenodon punctatus* a Maladapted Relict?

CARL GANS¹

ABSTRACT. The tuatara, *Sphenodon punctatus*, is the only Recent member of the Rhynchocephalia. Many aspects of its morphology suggest that it is primitive among the reptiles, and the fossil record indicates that its morphology has hardly changed since the Triassic. Consequently, its peculiar physiology (low metabolic rate correlated with low activity temperature), the relatively long period before sexual maturity, and its longevity have sometimes been interpreted as relicts of past conditions, presumably characterizing the ancestral group. Review of recent work suggests that many of these specializations represent obvious adaptations to current conditions. The tremendous diversity of adaptive patterns seen in Recent lepidosaurians makes it risky to use any small cohort of these animals to infer conditions in various extinct forms.

INTRODUCTION

The tuatara, *Sphenodon punctatus*, sole surviving member of the Rhynchocephalia, is the only living non-squamate lepidosaurian. Textbooks and the semi-popular literature emphasize this point and often note that it is a "living fossil," a "remnant" or "relict"; some even suggest that it (i.e., its skeleton) has remained unchanged since the Triassic. Other comments give one the impression that the occasional specimens of *Sphenodon* seen in zoos were sampled from remnant populations of slow and somehow inadequate relict reptiles, clinging for dear life to a few cold and barren, windswept rocks off the New Zealand coast.

It seems to be tempting to speculate that the physiology of such relict animals

might in some way have retained an ancestral condition rather than being clearly matched to present circumstances. For instance, Bogert (1953b) notes that the very low body temperature at which tuataras are active may have implications for the thermal pattern that existed in extinct members of the group. More recently, Bakker (1971) suggests that the thermal pattern probably reflects a primitively low thermal preference. Stebbins (1958) wonders whether parietal eye function in the tuatara might have a bearing on this function in "ancient reptiles." In contrast, Dawbin (1962) notes that the physiology of *Sphenodon* might well have developed quite independently from its skeleton.

I here suggest that the astonishing ecological and physiological traits of *Sphenodon* are neither maladaptive nor necessarily primitive, but are clearly matched to current conditions; they need not have implications for the specializations exhibited by the ancestral populations. It is my pleasure to dedicate this essay to Professor Ernest E. Williams, for personal reasons and because of his efforts at fostering an increased understanding of the biology of the lower tetrapods.

MORPHOLOGICAL ASPECTS

Sphenodon was early recognized as morphologically primitive, because its temporal region was diapsid and its quadrate fixed (Günther, 1867); this pattern, as well as the acrodont jaws and the

¹Division of Biological Sciences, The University of Michigan, Ann Arbor, Michigan 48109, U.S.A.

absence of male intromittent organs, was then considered to have been that of the earliest reptiles. Comparison showed that the Recent form closely resembles the various Triassic fossils (Romer, 1956), not only in the architecture of the skull, but also in aspects of the postcranial skeleton, such as number and proportions of skeletal elements. This level of similarity between Triassic and Recent forms has held up as more rhynchocephalian fossils became available (Carroll, 1977; Romer, 1956, 1966). By the Triassic, spenodontids were recognizably distinct from lizards (Robinson, 1973), which were already recorded in the Upper Permian (Carroll, 1977).

Various diapsids were initially and inappropriately assigned to the Rhynchocephalia (Romer, 1956). However, the early rhynchocephalian forms *Palacrodon* (South Africa, Lower Triassic), *Poly-sphenodon* (Europe, Upper Triassic), and *Opisthias* (Wyoming, Upper Jurassic; Simpson, 1926) are remarkably similar to *Sphenodon*, as are the excellently preserved *Homoeosaurus* (late Jurassic of Europe; cf. Hoffstetter, 1955; Romer, 1966) and *Toxolophosaurus* (Lower Cretaceous, Montana; Throckmorton *et al.*, 1981). Indeed, the various rhynchosauroids and several levels of advanced forms within the group differ in their structure both from the early rhynchocephalians and the modern form. Recent work suggests that the rhynchosauroids may be quite distinct from the Rhynchocephalia *sensu stricto* (Carroll, 1977). However, the Rhynchocephalia (including *Sphenodon*) retain their assignment separate from the lizards, which has been based on such cranial characteristics as the diapsid skull, the quadratojugal, the rigid attachment of the quadrate, and the acrodont dentition.

Perhaps the most important point in the present context is that even the Triassic spenodontids were functionally specialized, particularly in terms of their feeding system. Their dentition consists of three rows on each side, with the

dentary teeth closing between the maxillary and palatine rows (Robinson, 1976). Functional analysis of their masticatory system indicates that the power stroke or slow-closing component of each masticatory cycle, is divided into two phases. An initial vertical movement is followed by a secondary anterior sliding movement of the mandible (Gorniak *et al.*, 1982; Rosenberg and Gans, 1977). As the upper and lower surfaces of the food object are each held by the tooth cusps, the bolus is bent or sheared as the jaws slide. The triangular shape of the cusps permits the teeth to dig into the prey; thus one sees mice being decapitated (in captive animals) and obtains an understanding of how these animals decapitate seabirds (Crook, 1975). One further discovery is the remarkably limited degree of freedom of the spenodontid mandible. There is almost no opportunity for lateral movements, even when the mouth is partially open.

The pattern of jaw action emphasizes the extreme specialization of the mandibular system. No lizard shows such cutting ability. The most that has been noted in the Sauria is a crushing or piercing action. One superficially parallel action is the leaf-cropping of agamid and iguanid lizards (Throckmorton, 1976); however, in that case the very small and multi-lobed teeth serve as "pinking shears." Furthermore, the pattern in rhynchosauroids appears to have shifted to grinding the food; the various later forms are broad-headed and often show multiple-rowed and widened sets of teeth in the palatal arrays (Sill, 1971).

ECOLOGY AND PHYSIOLOGY

Some years of laboratory observation and a brief visit to the spectacular colony on Stephens Island, New Zealand, indicate that, whatever the physiology of their remote ancestors, tuataras are currently successful animals, well adapted to the circumstances of their current

environment. On Stephens Island, one of some dozen sites where viable populations now survive, *Sphenodon* occurs at an extremely high density (estimates range from 20,000 to 100,000 specimens on the 150 hectares, higher by orders of magnitude than densities elsewhere [D. G. Newman *in litt.*]). They are clearly the top nonavian predators, feeding on ground-dwelling arthropods, amphibians, reptiles and birds (Crook, 1975; Dawbin, 1962; Newman, 1977; Newman *et al.*, 1979; Walls, 1978). They crush and dismember hard prey, crunching large arthropods and carving up sea birds. Their hunting pattern is more than a passive sit-and-wait strategy. Individuals were observed at night inspecting crevices containing tree weta (*Hemideina crassidens*, a 10+ cm long, armored orthopteran); tuataras in such positions immediately moved to attack other insects when these were dropped close to them.

The capacity of *Sphenodon* to operate at low temperatures has often been stressed, particularly after it was shown that many other reptiles can raise their body temperatures to levels from 25° to 42°C and that certain physiological processes of other forms may not even respond below 25°C (Dawson, 1975). There can be little question that *Sphenodon* are active nocturnally at body temperatures between 6.2° and 16.0°C (Bogert, 1953a, personal observation); the hunting tuataras mentioned in the preceding paragraph had body (cloacal) temperatures near 14°C (and force-plate measured activity in the laboratory at the University of Illinois was found to be highest at dusk and dawn; J. Heath and R. G. Northcutt, personal communication). However, it has recently been confirmed that *Sphenodon* will bask, both in the field and in the laboratory (Dawbin, 1949; Stebbins, 1958; Werner and Whitaker, 1978; Barwick, *in litt.*; Gans and Regal, *in litt.*), and that basking animals raise their body temperature to 28°C or more. Still, the energy require-

ments during activity appear to be very low (Milligan, 1923; Wilson and Lee, 1970). Though tuataras have low body temperatures, they are hardly lethargic, thus Günther's (1867: after Smith) comment that they are "slow and sluggish" should be set aside.

In the densest populations, burrows are packed at 2 to 3-meter spacings; this apparently involves a complex social interaction (Newman, 1978). Tuataras produce vocalizations within their best hearing range (Gans and Wever, 1976), suggesting that these may be involved in communication. The burrows that birds, such as the fairy prion, *Pachyptila turtur*, and the diving petrel, *Pelecanoides urinatrix*, construct during the nesting season, are shared by *Sphenodon*. At lower densities, numbers of petrels and tuataras are correlated; however, the relative density of tuataras first becomes constant and then drops as the concentration of petrels and burrows increases further ($> 0.3/m^2$; Crook, 1974, 1975). While this may indicate interaction, it may also reflect different requirements. The animals interact within their colonies. Thus resident tuataras would sometimes emerge from tunnels or other sites to attack and chase other tuataras that had been displaced (being released away from their homesite during the night). Body temperatures of such animals range between 14° and 16°C. Recent laboratory observations document that males bite each other when establishing dominance, and bite females during mating (Gillingham *et al.*, ms.).

Many reports comment that tuataras rarely bask in open areas during the day, whether such areas are due to bird nesting activity or cattle grazing. They do bask in the open shade of the low forest, but do not seem to move far from the mouth of their burrows. They are cryptically colored and may show some color polymorphism yielding green and occasional orange specimens (the latter coloration seemingly matching the patterns of some local fungus *Favolaschia caloce-*

rata, that may have been recently introduced). This suggests that sight-hunting predators, such as hawks (Saint Girons *et al.*, 1981), may be significant. Even adult tuataras are small enough (< 1 kg), so that predation remains a potential problem.

The present range of the tuatara is clearly less than that of even a thousand years ago, when the form was widespread on the main islands of New Zealand (Robb, 1973, 1977). The arrival of man and of the various rats that he introduced changed the situation. Today tuataras have been exterminated on the mainland and on many of the coastal islands they once occupied (Crook, 1973a, b). It is uncertain how far inland the distribution of *Sphenodon* originally extended. There are numerous inland records, but most of these derive from Quaternary (Rich, 1974, 1975) kitchen middens (Crook, 1975). Clearly the Polynesian invasion affected these animals in three ways. The first is that *Sphenodon* and other local vertebrates were hunted for food (Davidson, 1972; Scott, 1973). The second is that the invaders produced significant vegetational and associated ecological change (Jones, 1975; McGlone, 1978). The third is that the invaders introduced *Rattus exulans*, which competes for the same food objects and also preys on the tuatara, particularly on its eggs and most juvenile stages (Whitaker, 1978). On some islands, these rats are obviously limiting the reproductive capacity, as the resident populations of *Sphenodon* consists mainly of adults and may no longer be self-sustaining (Crook, 1973a; Whitaker, 1978). Nevertheless, tuataras seem to do very well except in zones invaded by the rats.

Another set of characteristics about which there is substantial interest and which may be associated with the low body temperature is the longevity of the form. The values of twenty years to achieve sexual maturity and fifty years during which growth continues (Crook, 1970; Dawbin, 1962) rank *Sphenodon* among those reptiles with the slowest

development, though not necessarily among those with the longest lifespan (Case, 1978). In any case, the structure of populations seems curious for a lepidosaurian and is more like that noted in island tortoises (or in trees). The large number of first-year animals, noted in some reptilian populations (Richmond, 1965), does not seem to be in evidence, even on islands that lack rats (Crook, 1970, 1973a). Though growth is slow, not all of their physiological processes are slowed down, as *Sphenodon* are clearly capable of rapid and protracted movement.

INFERENCE FROM SPHENODON

The above set of circumstances permits us to pose two obvious questions. 1) Is *Sphenodon* well adapted to its current environment (i.e., that which it occupied before the disturbance induced by the arrival of man)? 2) Are any of the physiological specializations of this animal likely to have been retained from its ancestral forms, and might they permit conclusions about adaptation or climates in the Triassic?

The first question revolves around the low body temperature of these animals. The suggestion that this represents an ancestral condition in a relict species is probably unjustified for various reasons. Recently, we have had to depart from the earlier concept that all reptiles "prefer" and are primarily active at body temperatures above 30°C. Increasing numbers of reptiles have been shown to be active at body temperatures in the high 10's and low 20's (cf. Gans, 1976; Avery, 1982); other forms, such as geckos, feed nightly at low temperatures and bask under cover during the day. Also, certain populations of lizards selectively occupy environments in which they can never achieve the basking temperatures preferred by or available to populations occupying adjacent sites (Huey and Slatkin, 1976). In such environments, the bio-

topes either lack accessible sources of radiant energy or such radiant energy is only available at times and at locations where its cost, perhaps in energy required to reach it or exposure to predators while doing so, may be higher than the benefit obtained.

What are the factors that restrict *Sphenodon* to a low temperature regime? The first factor appears to be the occurrence of predation. The hawks and other birds of prey are generalist predators that do not depend upon *Sphenodon* as a food resource; thus the population size of these predators is established, in part, by other factors. However, these birds represent a continuing threat to any tuatara and hence limit the net gain, in terms of increased digestive or other physiological rate, achievable by basking. Reduction of the "preferred" body temperature reduces the need to bask.

Another possible set of reasons for a reduced body temperature reflects the adaptive zone occupied by *Sphenodon*. On Stephens Island, tuataras harvest the fabulous energy resource reflected by the enormous number of seabirds, the prions and petrels that nest on coastal rocks and islands. A direct harvest occurs by predation on the birds themselves; thus tuataras take significant quantities of their eggs and offspring (Newman, 1978). Adult birds are more vulnerable at night than in the daytime (Walls, 1978). An indirect harvest occurs, as the tuataras feed mostly on the large arthropods that are a by-product of the nesting colony. The annual reworking of the soil during the formation of nesting tunnels and the deposit of masses of guano, food scraps and dead chicks provide resources for arthropod scavengers, as well as fertilizer for the island vegetation that supports crickets and beetles. The various prey species are also nocturnal and must be hunted when ambient temperatures are low. Thus, *Sphenodon* appears to be adapted for nocturnal predation.

Perhaps the longevity of *Sphenodon* also represents an adaptation to an envi-

ronment in which all of the resources are at the mercy of occasional shifts in the patterns of oceanic currents. Very low energy requirements, not only for metabolism but also for growth, could reduce the effect of unpredictable "bad seasons," when changes in ocean currents cause a crash of the prey populations on which the seabirds subsist.¹ (On steep, high islands, such as Stephens, the vast majority of individual *Sphenodon* would lack access to the seashore, or to alternative food sources, during periods when fewer or no birds breed. At the least, a shift of populations toward the shore would increase their exposure to predators. While the availability of varied herbivorous insects might tend to buffer short-term effects, all of the biota must reflect the annual influx of birds.) However, prolonged immaturity would also slow the recovery of populations depleted during "bad seasons," would prolong the time that the sensitive eggs and juveniles are exposed to predation and may simply be a "cold island" strategy, possible only in the face of a most limited suite of predators.

Whatever the basis of some of the specializations now seen, the tuatara was clearly a well-adapted form at the time man first arrived in New Zealand and remains so now in a more restricted range.

¹ There seem to be relatively few data on long term stability of the breeding success for most oceanic birds (Murphy, 1936). Dunnet *et al.* (1979) document gradual changes (over 28 years) in an Orkney fulmar population, with the hatching success in good years being more than twice that in bad ones. On Peruvian islands (Hutchinson, 1950) breeding may be interrupted and the changes in the numbers of birds may be even more marked. On the islands in question, the birds "always re-appear, although their breeding success may vary from place to place and year to year" (J. Warham, also J. A. Bartle, both *in litt.*; for similar comments on other species, see Richdale, 1963, 1965). Comments suggest that New Zealand's rather drastic geological history (Fleming, 1975a, b) may have led to past interruption of breeding on coastal islands.

The second question is whether the pattern discerned may be deemed a retention from an ancestral condition. Such a retention would imply that *Sphenodon* derives from a "low temperature" stock that survived only in the cool climate of these islands. Two kinds of evidence regarding this point exist. First, there is the documentation that New Zealand underwent considerable climatic variation during the Quaternary and earlier (Fleming, 1979). Secondly, there is the record of the unique lizard fauna of New Zealand which is replete with exceptional life histories. For instance, an unpublished study by R. Barwick, reports that the New Zealand gecko, *Hoplodactylus duvauceli*, also has delayed sexual maturity (7 years) and an extremely slow growth rate, with growth continuing for more than 19 years (D. G. Newman, *in litt.*) while another New Zealand gecko, *H. maculatus*, lived more than 30 years in captivity (D. G. Newman *in litt.*). In these aspects, these animals differ significantly from other geckos for which such data are available (Bowler, 1977). Similar curiosities abound in other New Zealand lizards. Thus Towns (1975a, b) notes that the small nocturnal skink, *Leiolopisma suteri*, has one of the longest developmental times in its family, and the local geckos are unique in including viviparous species (Robb, 1973). It is of further interest that the tree weta are also long-lived (D. G. Newman, *in litt.*), but so are some other members of this primitive orthopteran group (T. H. Hubbell, personal communication).

Neither of these sets of evidence would eliminate the concept that the tuatara had retained its cold-temperature physiology from its ancestral stock, or that the Triassic or Cretaceous sphenodontids were cool-temperature animals. On the other hand, they do not offer any support for these hypotheses; indeed the modifications of the local lizard fauna make it unlikely that *Sphenodon* only

survived in New Zealand because of its temperate climate (Bakker, 1971). Instead it suggests that these specializations most likely reflect current adaptations (to a cool climate with both short-term and long-term environmental fluctuations). The specializations of *Sphenodon* and local lizards also document the extreme diversity seen in Recent Lepidosauria. If *Hoplodactylus* were the "lone survivor," it would provide a most inadequate indication of the physiological responses of other geckos. This example emphasizes the cautions necessary in utilizing growth data, population structure, and predator to prey ratios pertinent to any one set of lepidosaurians as a basis for inferring a uniform situation in major groupings of reptiles now extinct.

ACKNOWLEDGMENTS

I am delighted to thank the New Zealand Wildlife Service, in particular Donald G. Newman, for arranging a trip to Stephens Island and for his friendly assistance and advice. J. Warham shared his unpublished data on the breeding success of sea birds. Many others participated in the discussions that helped formulate and refine these ideas. C. M. Bogert, R. L. Carroll, G. C. Gorniak, Allen Greer, J. A. Hopson, R. W. Marlow, D. G. Newman, J. Ostrom, R. Payne, F. H. Pough, P. Pridmore, R. Shine, and E. E. Williams read and commented on the draft manuscript, though the responsibility for its final formulation is clearly my own. These travels were carried out while a participant in a U.S./Australia Cooperative Science Project; the travel to Stephens Island was supported by a John Simon Guggenheim Memorial Fellowship and preparation of this manuscript by National Science Foundation Grant DEB 77-02605 and DHEW G-IROIDE 05112.

LITERATURE CITED

- AVERY, R. A. 1982. Field studies of body temperatures and thermoregulation. In C. Gans and H. F. Pough (eds.), "Biology of the Reptilia," 12(4): 93-166. London, Academic Press.
- BAKKER, R. T. 1971. Dinosaur physiology and the origin of mammals. *Evolution*, 25(4): 636-658.
- BOGERT, C. M. 1953a. Body temperatures of the tuatara under natural conditions. *Zoologica*, 38(4): 63-64.
- . 1953b. The tuatara: Why is it a lone survivor? *Sci. Mo.*, 76: 163-170.
- BOWLER, J. K. 1977. Longevity of reptiles and amphibians in North American collections as of 1 November 1975. *Soc. Stud. Amph. Rept. Herp. Circ.*, (6): i-iv + 32.
- CARROLL, R. L. 1977. The origin of lizards. *Linn. Soc. Symp. Ser.*, (4): 359-396.
- CASE, T. J. 1978. On the evolution and adaptive significance of postnatal growth rates in the terrestrial vertebrates. *Quart. Rev. Biol.*, 53(3): 243-282.
- CROOK, I. G. 1970. Tuatara. *Wildlife* (Wellington), 2: 16-19.
- . 1973a. The tuatara, *Sphenodon punctatus* Gray, on islands with and without populations of the Polynesian rat, *Rattus exulans* (Peale). *Proc. N. Z. Ecol. Soc.*, 20: 115-120.
- . 1973b. Tuatara and the Polynesian rat. *Wildlife* (Wellington), 4: 47-48.
- . 1974. Are tuataras dependent on petrels? *Wildlife* (Wellington), 5: 43-46.
- . 1975. The tuatara, pp. 331-352. In G. Kuschel (ed.), "Biogeography and Ecology of New Zealand." The Hague, W. Junk, B. V.
- DAVIDSON, J. M. 1972. Archaeological investigations on Motutapu island, New Zealand. *Rec. Auckland Inst. Mus.*, 9: 1-14.
- DAWBIN, W. H. 1949. The tuatara. *Tuatara*, 2: 91-96.
- . 1962. The tuatara in its natural habitat. *Endeavour*, 21: 16-24.
- DAWSON, W. R. 1975. On the physiological significance of the preferred body temperatures of reptiles, pp. 443-473. In D. M. Gates and R. B. Schmerl (eds.), "Perspectives in Biophysical Ecology." New York, Springer Verlag.
- DUNNET, G. M., J. C. OLLASON, AND A. ANDERSON. 1979. A 28-year study of breeding fulmars *Fulmarus glacialis* in Orkney. *Ibis*, 121(3): 293-300.
- FLEMING, C. A. 1975a. The geological history of New Zealand and its biota, pp. 190. In G. Kuschel (ed.), "Biogeography and Ecology in New Zealand." The Hague, Junk.
- . 1975b. The Quaternary record of New Zealand and Australia, pp. 1-20. In R. P. Suggate and M. M. Cresswell (eds.), "Quaternary Studies." Wellington, Roy. Soc. New Zealand.
- . 1979. "The Geological History of New Zealand and its Life." New Zealand, Auckland Univ. Press.
- GANS, C. 1976. Aspects of the biology of uropeltid snakes, pp. 191-204. In A. d'A. Bellairs and C. Barry Cox (eds.), "Morphology and Biology of Reptiles." London, Linn. Soc. Symp. Ser. (3).
- GANS, C., AND E. G. WEVER. 1976. The ear and hearing in *Sphenodon punctatus*. *Proc. Nat. Acad. Sci.*, 73(11): 4244-4246.
- GILLINGHAM, J. C., C. GANS, AND D. L. CLARK. Ms. Courtship, mating and male combat in tuatara, *Sphenodon punctatus*. *J. Herpetol.*, submitted.
- GORNIK, G. C., H. ROSENBERG, AND C. GANS. 1982. Muscle activity during mastication in the tuatara, *Sphenodon punctatus*. *J. Morphol.*,
- GÜNTHER, A. C. L. G. 1867. Contribution to the anatomy of *Hatteria* (*Rhynchocephalus*, Owen). *Phil. Trans. Roy. Soc. London.*, 157(2): 595-629.
- HOFFSTETTER, R. 1955. Rhynchocephalia, pp. 556-576. In J. Piveteau, (ed.), "Traité de Paléontologie. Tome V. Amphibiens, Reptiles, Oiseaux." Paris, Masson et Cie.
- HUEY, R. B., AND M. SLATKIN. 1976. Costs and benefits of lizard thermoregulation. *Q. Rev. Biol.*, 51: 363-381.
- HUTCHINSON, B. E. 1950. Survey of contemporary knowledge of biogeochemistry. 3. The biogeochemistry of vertebrate excretion. *Bull. Amer. Mus. Nat. Hist.*, 96: xviii + 554.
- JONES, R. 1975. The Neolithic, Paleolithic and the hunting gardeners: Man and land in the Antipodes, pp. 21-34. In R. P. Suggate and M. M. Cresswell (eds.), "Quaternary Studies." Wellington, Roy. Soc. New Zealand.
- MCGLONE, M. S. 1978. Forest destruction by early Polynesians, Lake Poukawa, Hawkes Bay, New Zealand. *J. Roy. Soc. New Zealand*, 8(3): 275-281.
- MILLIGAN, R. R. D. 1923. The respiration and metabolism of the tuatara. *Austr. Assn. Adv. Sci. Rept.*, 16: 404-406.
- MURPHY, R. C. 1936. Oceanic Birds of South America. 2 Vols. New York, Amer. Mus. Nat. Hist.
- NEWMAN, D. G. 1977. Some evidence of the predation of Hamilton's frog (*Leiopelma hamiltoni* (McCulloch)) by tuatara (*Sphenodon punctatus* (Grey)) on Stephens Island. *Proc. N. Z. Ecol. Soc.*, 24: 43-47.
- . 1978. Tuataras and petrels. *Wildlife* (Wellington), (9): 16-23.
- NEWMAN, D. G., I. G. CROOK, AND L. R. MORAN. 1979. Some recommendations on the captive maintenance of tuataras *Sphenodon punctatus* based on observations in the field. *Int. Zoo Yearb.*, 68-74.
- RICH, T. H. 1974. *Soc. Vert. Paleontol. News Bull.*, (102): 43-44.
- . 1975. *Soc. Vert. Paleontol. News Bull.*, (105): 35-36.

- RICHDALE, L. E. 1963. Biology of the sooty shearwater, *Puffinus griseus*. Proc. Zool. Soc. Lond., 141: 1-117.
- . 1965. Biology of the birds of Whero Island, New Zealand with special reference to the diving petrel and the broad-billed prion on Whero Island, New Zealand. Trans. Zool. Soc. Lond., 31: 87-155.
- RICHMOND, N. D. 1965. Perhaps juvenile dinosaurs were always scarce. J. Paleontol., 39(3): 503-505.
- ROBB, J. 1973. Reptiles and Amphibia, pp. 285-303. In G. R. Williams (ed.), "The Natural History of New Zealand." Wellington, Reed.
- . 1977. "The Tuatara." Durham, Meadowfield.
- ROBINSON, P. L. 1967. The evolution of the Lacertilia, pp. 395-407. In "Problèmes actuels de Paléontologie." (Evolution des Vertébrés). Col. Int. Cent. Natl. Rech. Sci.
- . 1973. A problematic reptile from the British Upper Trias. J. Geol. Soc. London., 129: 457-459.
- . 1976. How *Sphenodon* and *Uromastyx* grow their teeth and use them, pp. 43-64. In A. d'A. Bellairs and C. Barry Cox (eds.), "Morphology and Biology of Reptiles." Linn Soc. Symp. Ser. (3), London.
- ROMER, A. S. 1956. "Osteology of the Reptilia." Chicago, Univ. Chicago Press.
- . 1966. "Vertebrate Paleontology. Third Edition." Chicago, Univ. Chicago Press.
- ROSENBERG, H. I. AND C. GANS. 1977. Preliminary analysis of mastication in *Sphenodon punctatus*. (Abstract). Amer. Zool., 17(4): 871.
- SAINT GIRONS, H., B. D. BELL AND D. G. NEWMAN. 1981. Brief observations on the activity and thermoregulation of the tuatara, *Sphenodon punctatus* (Reptilia: Rhynchocephalia), on Stephens Island, Cook Strait. New Zealand J. Zool.,
- SCOTT, S. D. 1973. Excavations on the "Sunde Site" N38/24, Motutapu island, New Zealand. Rec. Auckland Inst. Mus., 7: 13-30.
- SILL, W. D. 1971. Functional morphology of the rhynchosaur skull. Forma et functio, 4: 303-318.
- SIMPSON, G. G. 1926. American terrestrial rhynchocephalia. Amer. J. Sci., 12(5): 12-16.
- STEBBINS, R. 1958. An experimental study of the "third eye" in the tuatara. Copeia, 1958(3): 183-190.
- THROCKMORTON, G. S. 1976. Oral food processing in two herbivorous lizards, *Iguana iguana* (Iguanidae) and *Uromastix aegyptius* (Agamidae). J. Morphol. 148: 365-390.
- THROCKMORTON, G. S., J. A. HOPSON, AND P. PARKS. 1981. A redescription of *Toxolophosaurus claudi* Olson, a lower Cretaceous herbivorous sphenodontid reptile. J. Paleontol. (in press).
- TOWNS, D. R. 1975a. Ecology of the black shore skink *Leiopisma suteri* (Lacertilia: Scincidae), in boulder beach habitats. New Zealand J. Zool., 2(4): 389-407.
- . 1975b. Reproduction and growth of the black shore skink *Leiopisma suteri* (Lacertilia: Scincidae), in north-eastern New Zealand. New Zealand J. Zool. 2(4): 409-423.
- WALLS, G. Y. 1978. The influence of the tuatara on fairy prion breeding on Stephens island, Cook Strait. New Zealand J. Ecol., 1: 91-98.
- WERNER, Y. L., AND A. H. WHITAKER. 1978. Observations and comments on the body temperatures of some New Zealand reptiles. New Zealand J. Zool., 5: 375-393.
- WHITAKER A. H. 1978. The effects of rodents on reptiles and amphibians, pp. 75-88. In "The Ecology and Control of Rodents in New Zealand Nature Reserves." Wellington Dept. of Lands and Survey Information, ser. 4.
- WILSON, K. J. AND A. K. LEE. 1970. Changes in oxygen consumption and heart-rate with activity and body temperature in the tuatara *Sphenodon punctatum*. Comp. Biochem. Physiol., 33: 311-322.

Chromosomal C-Banding in *Anolis grahami*

J. A. BLAKE¹

ABSTRACT. The technique of C-banding is applied to chromosomes of *Anolis grahami*, an iguanine lizard showing large amounts of chromosomal variation within and between populations. Differential staining of one chromosome pair distinguished one sample from others. Variation both in diploid number and in amount and distribution of heterochromatin is noted among the three populations sampled.

INTRODUCTION

Although banding techniques have been used extensively in chromosomal analyses of mammals since 1969, they have not been readily adaptable to other groups. Within the reptiles, banding procedures have been utilized to examine sex chromosomes in snakes (Becak, 1972), and have revealed cryptic sex chromosomal differentiation of lizards (Bull, 1978). Comparative karyology of snakes (Stock and Mengden, 1975; Mengden and Stock, 1980) and turtles (Bickham and Baker, 1976; Sites *et al.*, 1979) have been discussed in light of C-, G-, and AgNOR chromosomal banding studies. The identification of chromosome arms (linkage groups) greatly enhances their usefulness as taxonomic characters revealing information about the evolution of the genome. C-banding, or centromeric banding, allows determination of the location and amount of heterochromatin present (Arrighi and Hsu, 1971; Chen and Ruddle, 1971).

In this paper, C-banding is used to identify chromosome groups in the iguanine lizard, *Anolis grahami*, a beta-anole native to Jamaica. Two subspecies are recognized on the basis of distinct color and morphological differences (Lynn and Grant, 1940; Underwood and Williams, 1959). *Anolis g. grahami* is distributed throughout most of Jamaica, excluding the eastern coast. *Anolis g. aquarum* is restricted to the moist north-eastern and eastern areas of Jamaica. There are complex interactions in areas of sympatry (Underwood and Williams, 1959). Chromosomal variation within *Anolis grahami* was first described by Hall (1974), who reported extensive polymorphism reflecting a 'pattern of considerable complexity' involving both occasional accessory chromosomes and fission of up to three pairs of macrochromosomes. My work has confirmed and extended the observations of Hall and his student Espinoza revealing chromosome numbers ranging from $2n = 30$ to $2n = 37$ (Blake, 1981). The purpose of this comparison of C-banding is to examine the types of information available from C-banding studies in anoles, and to initiate discussion of the heterochromatic variation which is apparent in *Anolis grahami*. This report of the use of banding techniques to study chromosomal variation in anoles is a logical extension of the great amount of work concerning chromosomal diversity and its contribution to the study of systematic relationships in lizards which has emanated from the laboratory of Ernest E. Williams (e.g., Gorman *et al.*, 1968; Gorman and Atkins, 1966, 1968;

¹ Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, U.S.A.

Webster *et al.*, 1972; Hall and Selander, 1973; Paull *et al.*, 1976).

MATERIALS AND METHODS

Specimens of *Anolis grahami* were collected on Jamaica from the vicinity of Anchovy (N = 6, St. James Parish, *A. g. grahami*), Mandeville (N = 5, St. Thomas Parish, *A. g. grahami*), and Bath (N = 3, St. Thomas Parish, *A. g. aquarum*) during March 1980 (Fig. 1). All animals used in this study are deposited in the MCZ. One male from each population was C-banded (MCZ 158586, 158594, 158643).

Chromosome samples were prepared in the field according to procedures described by Hall and Selander (1973). Animals were injected with colchicine 12 to 24 hours before dissection. Bone marrow, testes, and spleen were removed and macerated in 2.2% NaCitrate. Suspensions were centrifuged and hypotonic solution (1.8% NaCitrate) added to each pellet, which was then suspended and held for 12 minutes before centrifugation. After centrifugation (700 rpm for 7 minutes in an IEC tabletop centrifuge), fresh Carnoy's fixative was added drop by drop to the pellet. This preparation can be stored in 2 mls of fixative at 4°C for at least 6 months. A drop of suspension in fresh fixative was deposited on an acid-cleansed slide which had been drained after being held in ice-cold 50% methanol. The slide was then tilted and air dried.

Diploid number was determined from a minimum of 20 spreads in each animal.

C-banding technique is modified from that of Sumner (1972) and Schmid (1978). Four- to seven-day-old slides are treated in 0.2N HCl for 45 minutes and rinsed in distilled deionized water (d.d. H₂O). A series of slides is subjected to 5% Ba(OH)₂ at 37°C for times ranging from 1 to 5 minutes at one minute increments. After rinsing in d.d. H₂O, the slides are incubated in 2XSSC (1X SSC is 0.15M NaCl, 0.015M NaCitrate) pH = 7.0 at

Figure 1. Map of Jamaica showing localities where animals used in this study were collected.

60°C for 1 hour. The slides are finally rinsed and stained for 10 minutes in Giemasa/citric acid stain. Selected spreads were photographed on a Leitz compound microscope using Kodak Tech-Pan film.

RESULTS

Diploid Number. Five individuals from Anchovy showed a diploid number of 36 with 20 macrochromosomes and 16 microchromosomes. One individual from Anchovy had a diploid number of 37. Individuals from Mandeville showed diploid numbers of 30(2), 31(2), and 32(1) (Table 1). Of the three individuals from Bath, one had a diploid number of 30; the other two had 31. All individuals have 16 microchromosomes. The differences appear in the number of macrochromosomes. As the number of macrochromosomes increases, so does the number of telocentrics.

Centromeric Heterochromatin. All metacentrics showed centromeric banding indicative of centromeric heterochromatin (Figs. 2, 3, 4). In the Anchovy individual, the largest pair of metacentrics shows a difference in intensity of staining in the centromeric region (Fig. 2). One chromosome is heavily stained at the centromere. The other is less intensely stained, and the centromere and blocks of heterochromatin on either side of the centromere are distinguishable from each other. This heteromorphism was observed in five different C-banded prints, as well as in C-band spreads of

TABLE 1. GENERAL KARYOTYPE DESCRIPTIONS FOR *ANOLIS GRAHAMI* FROM THREE JAMAICAN LOCALITIES.

Locality	N	2n	V = metacentrics
			sV = small metacentrics
			I = telocentrics
			A = acrocentrics
			m = microchromosomes
Anchovy	6	36(5)	4V + 4sV + 12I + 16m
		37(1)	3V + 4sV + 14I + 16m
Mandeville	5	30(2)	10V + 4sV + 16m
		31(2)	9V + 4sV + 2I + 16m
		32(1)	8V + 4sV + 4I + 16m
Bath	3	30(1)	10V + 4sV + 16m
		31(2)	1A + 8V + 4sV + 2I + 16m

Figure 2. C-Banded karyotype of a ♂ *Anolis grahami* from Anchovy. Arrows show differential centromeric staining of large metacentric pair. $2n = 36$.

individuals of another *Anolis g. grahami* population at Laughlands, St. Ann Parish not reported here. The differential staining is not observed in the specimens from Mandeville (Fig. 3) or Bath (Fig. 4). The telocentrics also show centromeric staining with two prominent blocks of heterochromatin distinguishable in the centromeric region (Fig. 2).

Telomeric Heterochromatin. All telomers are stained, although some are stained more intensely than others. This can be diagnostic, as in the case of the largest metacentrics of the Anchovy karyotype (Fig. 2). Here the largest pair is distinguishable from other metacentrics not only on the basis of size, but also on

Figure 3. C-Banded karyotype of a ♂ *Anolis grahami* from Mandeville. Arrows show absence of differential centromeric staining in large metacentric pair. $2n = 30$. Figure 4. C-Banded karyotype of a ♂ *Anolis grahami* from Bath. Only 15 microchromosomes are observed, the 16th is presumed to be covered by other chromosomes. The arrow indicates the heterochromatic acrocentric arm. $2n = 31$.

the basis of telomeric staining. One telomeric region of the chromosomes is heavily stained, the other is not. This is repeatedly observed, and is apparent also in the individuals from Mandeville and Bath. In Mandeville and Bath such distinction is more important in identifying

chromosomes as there are three pairs of metacentrics of approximately the same size in these populations.

Whole-Arm Heterochromatin. In all cases, the two pairs of small metacentrics are distinguishable on the basis of whole-arm staining (Figs. 2, 3, 4). One pair is heavily stained, the other is not. In many spreads, differences in staining between members of the darkly staining pair was often observed. One member of the pair stained heavily over both arms, the other seemed to stain only over one arm. This distinction may be important as it is these small metacentrics which are involved in sex chromosome arrangements in other *grahami* series species.

One of the macrochromosomes in the Bath spread appears to have a heterochromatic arm. The presence of an acrocentric chromosome with a heterochromatic short arm in *A. g. aquarum* populations is under further investigation.

Microchromosomes. The microchromosomes are stained in all spreads. The heterochromatic nature of the microchromosomes is especially apparent in meiotic metaphase I spreads (Blake, 1981). Microchromosomes are an important part of sex chromosome systems in *Anolis* when these have been observed (Gorman *et al.*, 1968; Gorman and Atkins, 1969), and the lack of distinction among the microchromosomes is therefore of special interest.

DISCUSSION

Anolis grahami presents a complex case of chromosomal polymorphism. Subspecies designation does not clarify the variation observed. *Anolis g. grahami* diploid numbers range from 30 to 32 at Mandeville, to 37 at Anchovy. Populations with intermediate numbers of chromosomes exist, and some have been extensively studied (Blake, 1981). Whether the polymorphism evident at Mandeville is balanced and stable can not be decided here. The presence of

such polymorphism, both at Mandeville, and in the *Anolis g. aquarum* at Bath, is indicative both of the widespread tendency for karyotypic change in this species, and of the presence of chromosomal polymorphisms throughout its range.

Such conditions of chromosomal polymorphism are not unknown in *Anolis*. *Anolis monticola* provides a parallel example of fissioning of the karyotype and presence of polymorphisms (Webster *et al.*, 1972). In that case, the primitive karyotype of $12M + 24m$ is altered through fission events to $48M + 24m$. Other *monticola* series species also appear to have undergone karyotypic changes through fission events (Webster *et al.*, 1972).

Anolis grahami is a member of the *grahami* series, a group of closely related beta-anoles inhabiting Jamaica. Within this series, *A. lineatopus*, *A. opalinus*, and *A. garmani* have diploid numbers of 30, $14M + 16m$, although some polymorphism in *A. opalinus* is suspected (Gorman and Atkins, 1968a; Gorman, 1969). *Anolis conspersus*, another member of this series from nearby Grand Cayman Island, shows sex chromosomal heteromorphism, with the males having a small acrocentric and metacentric in contrast to two small metacentrics in the female (Gorman and Atkins, 1968b). Most of these determinations were based on small sample sizes and, in light of current information concerning the variable *Anolis grahami* karyotype, warrant further investigation.

The presence of differential staining of the centromere of the large metacentric of the Anchovy individual is strongly reminiscent of the kind of variation reported in the lizard *Cnemidophorus tigris* by Bull (1978). In that case, Bull demonstrated a sex difference in the staining pattern and designated the chromosomes as cryptic sex chromosomes, an intermediate stage in sex chromosomal differentiation. The observation that such differential staining is

not observed in the three males studied here makes such an argument unlikely in this case. In addition, the C-banding pattern in female *Anolis grahami* is not known. The absence of repeatability of such differential staining in different populations is warning that the karyotypic description of a species must be based on a fair sampling of the population.

ACKNOWLEDGMENTS

I thank S. Abmeyr for her assistance in micro-photographic procedures and S. M. Case for critical reading of the manuscript. This work was supported by National Science Foundation Grant DEB 79-05837 to E. E. Williams, and by the Anderson Fund.

LITERATURE CITED

- ARRIGHI, F. E., AND T. C. HSU. 1971. Localization of heterochromatin in human chromosomes. *Cytogenetics*, **10**: 81-86.
- BECAK, W. 1972. W-sex chromatin fluorescence in snakes. *Experientia*, **28**: 228-229.
- BICKHAM, J. W., AND R. J. BAKER. 1976. Chromosome homology and evolution of emydid turtles. *Chromosoma*, **54**: 201-219.
- BLAKE, J. A. 1981. Chromosomal variation in the Jamaican lizard *Anolis grahami*. Ph.D. Thesis, Dept. of Biol., Harvard Univ., Cambridge.
- BULL, J. 1978. Sex chromosome differentiation: An intermediate stage in a lizard. *Can. J. Genet. Cytol.*, **20**: 205-209.
- CHEN, T. R., AND F. H. RUDDLE. 1971. Karyotype analysis utilizing differentially stained constitutive heterochromatin of human and murine chromosomes. *Chromosoma*, **34**: 51-72.
- GORMAN, G. C. 1969. Mammalian Chromosome Newsletter. Vol. 10, pp. 222-224.
- GORMAN, G. C., AND L. ATKINS. 1966. Chromosomal heteromorphism in some male lizards of the genus *Anolis*. *Am. Natur.* **100**: 579-583.
- . 1968a. New karyotypic data for 16 species of *Anolis* from Cuba, Jamaica, and the Cayman Islands. *Herpetologica*, **24**: 13-20.
- . 1968b. Confirmation of an X-Y sex determining mechanism in lizards (*Anolis*). *Copeia*, **1**: 159-160.
- . 1969. The Zoogeography of Lesser Antillean *Anolis* lizards. In *Analysis Based upon Chromosomes and Lactic Dehydrogenases*. *Bull. Mus. Comp. Zool.*, **138**: 53-80.
- GORMAN, G. C., R. THOMAS, AND L. ATKINS. 1968. Intra- and interspecific chromosome variation in the lizard *Anolis cristatellus* and its closest relatives. *Breviora Mus. Comp. Zool. No.* 293, pp. 1-13.
- HALL, W. P. 1974. 2nd *Anolis* Newsletter. E. E. Williams (ed.), 2 pp.
- HALL, W. P., AND R. K. SELANDER. 1973. Hybridization of karyotypically differentiated populations in the *Sceloporus grammicus* complex (Iguanidae). *Evolution*, **27**: 226-242.
- LYNN, W. G., AND C. GRANT. 1940. The Herpetology of Jamaica. *Bull. Inst. Jamaica Sci. Ser. No.* 1, pp. 1-148.
- MENGDON, G. A., AND A. D. STOCK. 1980. Chromosomal evolution in serpentes: A comparison of G and C chromosome banding patterns of some colubrid and boid genera. *Chromosome*, **79**: 53-64.
- PAULL, D., E. E. WILLIAMS, AND W. P. HALL. 1976. Lizard karyotypes from the Galapagos Island: (Chromosomes in phylogeny and evolution.) *Breviora, Mus. Comp. Zool. No.* 441, pp. 1-31.
- SCHMID, M. 1978. Chromosome banding in Amphibia I. Constitutive heterochromatin and nucleolus organizer regions in *Bufo* and *Hyla*. *Chromosoma*, **66**: 361-388.
- SITES, J. W., J. W. BICKHAM, M. W. HAIDUK, AND J. B. IBERSON. 1979. Banded karyotypes of six taxa of kinosternid turtles. *Copeia*, **4**: 692-698.
- STOCK, A. D., AND G. A. MENGDON. 1975. Chromosome banding pattern conservatism in birds and non-homology of chromosome banding patterns between birds, turtles, snakes and amphibians. *Chromosoma*, **50**: 69-77.
- SUMNER, A. J. 1972. A simple technique for demonstrating centromeric heterochromatin. *Exp. Cell Res.*, **75**, 304-306.
- SUMNER, A. T., H. J. EVANS, AND R. A. BUCKLAND. 1971. A new technique for distinguishing between human chromosomes. *Nature N. B.*, **232**: 31-32.
- UNDERWOOD, G., AND E. E. WILLIAMS. 1959. The Anoline Lizards of Jamaica. *Bull. Inst. Jamaica Sci. Ser. No.* 9, pp. 1-48.
- WEBSTER, T. P., W. P. HALL, AND E. E. WILLIAMS. 1972. Fission in the evolution of lizard karyotype. *Science*, **177**: 611-613.

The Relationships of the Puerto Rican *Anolis*: Electrophoretic and Karyotypic Studies

GEORGE C. GORMAN¹

DONALD BUTH²

MICHAEL SOULÉ³

S. Y. YANG⁴

ABSTRACT. Relationships of the Puerto Rican anoles were investigated through both phenetic and cladistic treatments of two independently generated multi-locus allozyme data sets. These findings are compared with a phylogeny developed from previously reported karyotypic data. Phylogenetic treatment of the data supports a rather conventional view of the relationships of the Puerto Rican anoline radiation. There is electrophoretic support for the recognition of three major karyotypically limited lineages: 1) *Anolis evermanni* and *A. stratulus*; 2) *A. cooki* and *A. cristatellus*; and 3) *A. gundlachi*, *A. poncensis*, *A. pulchellus*, and *A. krugi*, although *A. gundlachi* clusters with the latter group based on a karyotypic rather than any electrophoretic character. Within the third lineage, *A. poncensis* is the sister-taxon of a group comprised of *A. krugi* and *A. pulchellus*, and these three "grass-bush" anoles are in a sister group relationship to *A. gundlachi*.

INTRODUCTION

There are ten species of *Anolis* lizards on Puerto Rico, all endemic to the island bank. This adaptive radiation has received considerable attention from

evolutionary biologists in the past two decades. Rand (1964) studied ecological niche partitioning among the species, and Schoener and Schoener (1971) further elaborated upon this.

In 1972, Ernest E. Williams published a stimulating paper on the anoline radiation of Puerto Rico. In that paper he presented a model based largely upon ecological principles that described how the Puerto Rican anoline fauna may have evolved *in place* following an initial colonization event. Subsequent analyses tend to support much of the model (e.g., Gorman *et al.*, 1980a). It now appears that there is a primary radiation accounting for eight of the species. Two additional species (*Anolis cuvieri* and *A. occultus*) that appear to have more primitive relatives living on the island of Hispaniola may have independently colonized Puerto Rico.

The morphology of most Greater Antillean *Anolis* is highly correlated with their ecology. Williams (1972) assigned the *Anolis* of Puerto Rico to various "ecomorph" categories. *Anolis cuvieri* is a "crown-giant" and *A. occultus* is a "twig-dwarf" species. The other ecomorph categories each have several species which replace one another by climatic niche. The three "trunk-ground" species, *A. gundlachi*, *A. cristatellus*, and *A. cooki*, are distributed in the rainforest,

^{1,2}Department of Biology, University of California (UCLA), Los Angeles, California 90024, U.S.A.

³Department of Biology, University of California (UCSD), La Jolla, California 92037, U.S.A. Present Address: Institute for Transcultural Studies, 905 South Normandie Avenue, Los Angeles, California, 90006, U.S.A.

⁴Department of Biology, University of California (UCSD), La Jolla, California 92037, U.S.A. Present Address: Department of Biology, Inha University, Incheon, Korea.

lowland mesic areas, and the xeric southwestern part of the island, respectively. The three "grass-bush" species, *A. krugi*, *A. pulchellus*, and *A. poncensis*, replace one another similarly. Finally, there is a pair of "trunk-crown" species: *Anolis evermanni* is the typical upland species, and *A. stratulus* is widespread in the lowlands.

Only the three widespread lowland (mesic) species occur on the islands east of Puerto Rico that form part of the Puerto Rico Bank. (Politically, some of these islands are part of Puerto Rico, the remainder being the United States and British Virgin Islands. The island of St. Croix, of the U.S.V.I., is not part of the Puerto Rico Bank, and has an endemic anole.)

In 1976 Williams published a formal classification of West Indian *Anolis*. The classification closely follows the osteological interpretations of Etheridge (1960). In this classification, the genus is divided into two sections (alpha and beta); the alpha section is divided into two subsections (*carolinensis* and *punctatus*). All of the Puerto Rican anoles belong to the alpha section, and all but one belong to the *punctatus* subsection. The diminutive *Anolis occultus* is the only Puerto Rican member of the *carolinensis* subsection.

The remaining nine species were placed in three different series. *Anolis cuvieri* belongs to the *cuvieri* series, along with several Hispaniolan giant anoles, and the possibly extinct giant anole *A. roosevelti* of Culebra (a small island that is part of the Puerto Rico Bank). The "trunk-ground" and "grass-bush" anoles were placed in the *cristatellus* series. This series was divided into two subseries—the *cristatellus* subseries and the *cybotes* subseries. The former contains only the six Puerto Rican species plus two extralimital forms that are closely allied with *cristatellus* (Gorman *et al.*, 1980b). The *cybotes* subseries is a Hispaniolan radiation of a single super-species. The "trunk-crown" anoles of

Puerto Rico were placed in the *bimaculatus* series. This series, as defined by Williams (1976), is represented throughout the northern part of the Lesser Antilles, the island of St. Croix, and on Hispaniola where there is a single radiation of closely related species (the distichoids). In addition, the Hispaniolan *A. distichus* has colonized some of the Bahamas and Florida.

Interpretation of data obtained from albumin immunology, as well as interpretation of karyotype data, has raised some questions about the classification proposed by Williams (1976). Shochat (1976) and Shochat and Dessauer (1981) questioned the alpha-beta dichotomy; Wyles and Gorman (1980) raised some questions about the division of the alpha section into two subsections. The latter study revealed that *A. cybotes* is not closest in albumin immunological distance or karyotype to the *cristatellus* radiation, but that its closest affinities appears to be with other Hispaniolan anoles, some of which were placed by Williams in the *carolinensis* subsection.

Gorman *et al.* (1980a) presented a cladistic analysis of eastern Caribbean *Anolis* based upon albumin immunological data. This analysis did not deal with the alpha-beta dichotomy, or the *carolinensis-punctatus* dichotomy; these investigators chose to restrict the analysis exclusively to the *punctatus* subsection of the alpha section as defined by Williams (1976). The analysis of immunological data led to a proposed classification that differs somewhat from the one proposed by Williams (1976). Specifically, the cybotoids were removed from the *cristatellus* series and placed in their own series. The *cristatellus* series was then divided into two subseries, designated *cristatellus* and *bimaculatus*. The latter is restricted to the Lesser Antilles; the former now includes the Greater Antillean (and St. Croix) forms that Williams placed in both the *bimaculatus* and *cristatellus* series. Thus the "trunk-crown" anoles of Puerto Rico are placed

in the same series and subseries as the "trunk-ground" and "grass-bush" radiations. The revised classification is thus completely in accord with Williams' (1972) idea of a "radiation in place."

The purpose of the present paper is to utilize previously reported karyotypic data combined with new data obtained from multi-locus electrophoretic studies to help us further analyze the relationships of the Puerto Rican anoles. Our primary concern is with lower-level patterns of relationships, not with the classification of the genus; however, we have chosen certain "outside" forms for comparison specifically to examine some of the controversial questions in anole systematics.

There are two independent allozyme data sets utilized in this paper. Almost ten years ago, three of us (Yang, Soulé, Gorman) collaborated on a 19 locus electrophoretic study of all species of Puerto Rican *Anolis*, plus several selected outside representatives. The original study failed to resolve clearly some of the patterns of relationship among the eight Puerto Rican species assigned by Gorman *et al.* (1980a) to the *cristatellus* subseries. Thus a second data set was obtained, based on fresh material, for the eight species. The number of loci examined was expanded to 35 and different outside reference species were utilized in the quest for clearer patterns to emerge. Nei and Roychoudhury (1974) pointed out that genetic distance estimates are far more sensitive to the number of loci examined than to the number of specimens examined. This has been confirmed empirically for *Anolis* by Gorman and Renzi (1979).

MATERIALS AND METHODS

ALLOZYME DATA SETS

The first electrophoretic study involved 22 population samples representing 15 species of *Anolis*, including all ten

Puerto Rican species, which were examined for gene products of 19 loci. The electrophoretic methods have been previously presented (Yang *et al.*, 1974; Gorman and Kim, 1976) as have been the allelic data for the populations of *A. scriptus*, *A. cooki*, *A. monensis*, and the representative population of *A. cristatellus* (#8 of Gorman *et al.*, 1980b), which are also compared in the present study. Sample size and locality data for all populations in this first study are listed in Appendix I.

The second electrophoretic study involved 12 population samples representing 11 species of *Anolis*, including eight species from Puerto Rico, which were examined for gene products of 35 loci. The electrophoretic methods for the second study are essentially those of Buth *et al.* (1980) with the exceptions that adenosine deaminase Ada-A, esterase Est-2, peptidase Pept-3, superoxide dismutase Sod-2, and xanthine dehydrogenase Xdh-A of the latter study have been replaced by creatine kinase (Ck-C; EC 2.7.3.2), dihydrolipoamide reductase (Dlr-1; EC 1.6.4.3), fumarate hydratase (Fum-A; EC 4.2.1.2), L-iditol (sorbitol) dehydrogenase (Iddh-A; EC 1.1.1.14), and a second "malic enzyme" (Me-2; EC 1.1.1.40). Staining procedures for these five systems were modified from Shaw and Prasad (1970), Brewer (1970), Lin *et al.* (1969), Kaplan and Beutler (1967), and Ayala *et al.* (1972), respectively. Sample size and locality data for all populations in this second study are listed in Appendix II.

TREATMENT OF ALLOZYME DATA

Phenetics. Differentiation among populations was quantified for both of the allozyme data bases by computing Nei's (1972) genetic distance (D) and similarity (I) coefficients between each pair of populations. The distance coefficients were then clustered using the unweighted pair group method with

arithmetic averages (Sneath and Sokal, 1973).

Cladistics. Phylogenetic relationships within the *crisatellus* subseries (Gorman *et al.*, 1980) and the Puerto Rican representatives of this subseries were estimated using the Wagner procedure (Farris, 1970) as applied to both of the allozyme data bases. Allelic products were encoded as "present" or "absent" following Mickevich and Johnson (1976). While the latter study employed a frequency minimum of 0.05 for allelic "presence," our treatment for the two data sets employed a much higher frequency requirement (0.25) to minimize the resolution limitations set by the gross sample size differences between the two studies. Most of the uncommon alleles omitted in this coding procedure were population-specific and contributed little to the estimation of relationships between taxa. Initially estimated unrooted Wagner networks (Lundberg, 1972) were rooted by including species presumed to be external to the phyletic radiation in question (see Farris, 1972). Thus, the first electrophoretic study employed *A. cuvieri*, *A. cybotes*, *A. occultus*, and *A. oculatus* for rooting purposes whereas the second study employed *A. gadovi*, *A. grahami*, and *A. carolinensis* for this purpose. The first two members of the rooting array used in the second study are members of the beta section; the third species is a North American form of Cuban affinity (Buth *et al.*, 1980) that represents the other (i.e., *carolinensis*) subsection of alpha *Anolis*.

RESULTS

The allozyme data bases for the first and second electrophoretic studies are presented in Tables 1 and 2, respectively.

Phenetic relationships derived from both allozyme data sets are depicted in Figures 1 and 2. In all cases where more than a single population was examined

from a given species, the intraspecific genetic distances are uniformly low and the populations of a species cluster together (e.g., *A. acutus*, *A. crisatellus*, *A. krugi*, *A. poncensis*, *A. pulchellus*, *A. stratulus*). Among the eight Puerto Rican species examined in common between the two electrophoretic studies, the average genetic distance value (D) for pairwise interspecific comparisons is similar. The average absolute difference between the studies is $D=0.07$. Despite these small average differences, there are certain trends in distances of the second study that provide a different perspective regarding interspecific similarities. The most striking discrepancy is in the relationship of *A. evermanni* and *A. stratulus*. In the first study, the mean genetic distance between this pair was approximately $D = 0.5$, and the two species were not closely associated (Fig. 1). In the second study, $D = 0.27$ between these forms and this pair clusters together (Fig. 2).

The cladistic analyses were limited to the members of the *crisatellus* subseries of Gorman *et al.* (1980a). The cladistic treatment of both allozyme data sets yielded Wagner trees with levels of homoplasy that might be considered high. Consistency indices (Farris, 1969; Mickevich and Johnson, 1976) of $c = 0.64$ and $c = 0.61$ were computed for Wagner trees based on the first and second electrophoretic studies, respectively. Thus, a large number of equally parsimonious minimum-length Wagner trees may be constructed. Evaluations of several such trees revealed repeated clustering of certain groups of species in both cases. These conservatively estimated phylogenetic relationships are depicted in Figures 3 and 4.

The phylogenetic relationships of the eight Puerto Rican species in the second electrophoretic study were examined with reference to a comparison of karyotypic evolution previously reported by Gorman *et al.* (1968) and Gorman and Atkins (1969). These karyotype descrip-

TABLE 1. ALLELIC VARIABILITY WITHIN AND AMONG SAMPLES OF 22 POPULATIONS OF 15 SPECIES OF *ANOLIS*.

Allelic products are lettered corresponding to increasing anodal electrophoretic mobility and are monomorphic within samples unless otherwise noted. Allelic notation is equivalent to the data base of Gorman *et al.* (1980b) which includes additional populations of *A. cristatellus*; however, allele letters used here do not necessarily correspond to those in other tables in the present study. Populations are numbered in the order they are listed in Appendix I. * = no expression of the locus (null allele?). ** = equivalent to creatine kinase (Ck-A).

Locus	1	2	3	4	5	6	7	8	9	10	11
1. Adh-A	f	f	f	c(0.17) f(0.83)	c(0.02) f(0.98)	f(0.92) h(0.08)	f(0.98) g(0.02)	f(0.98) h(0.02)	f(0.83) h(0.17)	c(0.05) f(0.77) h(0.18)	f
2. S-Aat-A	d(0.97) g(0.03)	d	d	d(0.94) e(0.02) f(0.04)	b(0.94) d(0.06)	c(0.13) d(0.87)	d	c(0.02) d(0.98)	c(0.02) d(0.98)	d	d(0.96) g(0.04)
3. Est-1	c(0.07) d(0.03) f(0.10) h(0.73) j(0.07)	b(0.04) f(0.96)	j	k(0.04) m(0.52) p(0.46)	i	e(0.07) f(0.04) h(0.05) j(0.74) j(0.84)	c(0.02) e(0.02) j(0.74) l(0.22)	e(0.02) f(0.12) j(0.72) l(0.14)	b(0.02) e(0.98)	a(0.12) b(0.02) e(0.86)	*
4. Fum-A	d	d	d(0.87) f(0.13)	d	a(0.02) d(0.98)	d(0.75) e(0.25)	d	d	d	d	d
5. Gp-1	d(0.07) e(0.93)	e	c(0.20) e(0.80)	e	f(0.02) h(0.98)	d(0.98) f(0.02)	d(0.74) e(0.08) f(0.18)	c(0.02) e(0.96) g(0.02)	e	c(0.02) e(0.98)	f
6. Gp-2	b	b	b	b	b	b	b	b	b	b	b
7. Gp-3 ^{**}	a	a	c	c	a	a	a	a	a	a	a
8. Gp-4	b	b	b	b	b	b	b	b	b	b	b
9. Gpi-A	f	f	a(0.07) c(0.93)	c	b(0.04) c(0.02) f(0.94)	c(0.05) e(0.92) f(0.03)	e	d(0.12) e(0.88)	e	c(0.08) e(0.90) g(0.02)	c
10. G-3-pdh-A	e	e	b(0.02) e(0.98)	e	a(0.04) e(0.96)	e(0.98) f(0.02)	e	e(0.98) f(0.02)	e	e	e
11. M-lcdh-A	c(0.97) e(0.03)	c	c(0.73) e(0.27)	c	a(0.71) c(0.29)	b(0.90) c(0.10)	b(0.85) e(0.15)	b	c	c	c
12. S-lcdh-A	c	c	a(0.15) c(0.85)	c	c	c	c	c	c	c	c
13. Ldh-A	c	c	c	c	c	c(0.90) f(0.10)	c	a(0.02) c(0.98)	c	c	c
14. Ldh-B	c(0.67) e(0.33)	c(0.98) h(0.02)	f	e(0.98) f(0.02)	f	a(0.57) f(0.43)	a(0.70) f(0.30)	a(0.62) f(0.38)	a	a(0.95) f(0.05)	f
15. M-Mdh-A	d	d	d	d	a(0.12) e(0.88)	d	d	d	d	d	d
16. S-Mdh-A	d	d	d	d	d(0.94) e(0.06)	d	d(0.95) e(0.05)	d	d(0.63) e(0.37)	d(0.25) e(0.75)	d
17. Pept-1	b(0.10) d(0.90)	d(0.98) f(0.02)	d	d	b(0.96) e(0.04)	d	d	d	d	d	d
18. Pgm-A	d(0.64) g(0.33) h(0.03)	g	d	d	d	a(0.02) d(0.87) f(0.08) h(0.03)	a(0.10) d(0.81) d(0.86) e(0.02) g(0.07)	a(0.06) d(0.88) g(0.08)	a(0.04) d(0.79) f(0.17)	a(0.17) d(0.83)	d(0.50) g(0.50)
19. Pgdh-A	a(0.07) d(0.83) e(0.03) f(0.07)	d	d	a(0.02) d(0.87) e(0.11)	d(0.02) i(0.98)	d	d(0.80) e(0.20)	a(0.02) b(0.02) d(0.88) e(0.02) f(0.06)	c(0.20) d(0.04) e(0.76)	c(0.17) e(0.83)	d

TABLE 1. CONTINUED.

Locus	12	13	14	15	16	17	18	19	20	21	22
1. Adh-A	f	c(0.02) f(0.98)	f	f	f	f	c(0.06) f(0.94)	f	f	d	a(0.03) b(0.90) e(0.07)
2. S-Aat-A	d(0.93) g(0.05)	c(0.98) d(0.02)	c	b	b	b	c(0.96) d(0.04)	b	a	d	c
3. Est-1	*	l(0.04) n(0.96)	n	p	k(0.11) p(0.89)	p(0.95) q(0.05)	p(0.98) q(0.02)	i	i	g(0.03) i(0.78) l(0.19)	*
4. Fum-A	d	b(0.02) d(0.76) f(0.22)	b(0.03) d(0.64) f(0.33)	d	d	d	d	d	c	d	d
5. Gp-1	f	e	e	e	e(0.78) f(0.22)	c(0.10) e(0.90)	a(0.17) g(0.83)	c(0.19) f(0.81)	g	d	e
6. Gp-2	b	b	b	b	b	b	b	b	a	b	b
7. Gp-3 ^{††}	a	a	a	a	a	a	a	a	b	d	c
8. Gp-4	b	b	b	b	b	b	b	b	b	c	b
9. Gpi-A	a(0.11) c(0.89)	a(0.02) c(0.98)	a(0.03) c(0.97)	c	c	c(0.95) f(0.05)	c(0.04) f(0.96)	a(0.04) c(0.96)	c(0.95) f(0.05)	f	c(0.13) e(0.84) g(0.03)
10. G-3-pdh-A	e	e	e	b	b	b	c(0.04) e(0.96)	e	e	d(0.97) e(0.03)	b
11. M-Icdh-A	c(0.89) e(0.11)	c	c	d	d	d	c	c(0.60) e(0.40)	c	c	a(0.03) c(0.94) e(0.03)
12. S-Icdh-A	c	c	c	c	c	c	b(0.23) c(0.77)	c	d	c	c(0.86) e(0.14)
13. Ldh-A	c	c(0.98) e(0.02)	c	b(0.04) c(0.96)	c	c	a(0.01) c(0.98) d(0.01)	c	d	c	f
14. Ldh-B	f	f	f	d	d	d	g	c	f	c	f
15. M-Mdh-A	d	d	d	c	a(0.05) c(0.95)	c	c	c	b	a	d
16. S-Mdh-A	d	d	d	b	b	b	d	c(0.24) d(0.76)	c	a	c
17. Pept-1	d	d	d	d	d	d	d	c	d	b(0.50) d(0.47) f(0.03)	a
18. Pgm-A	d(0.44) g(0.56)	a(0.04) d(0.94) f(0.02)	d(0.92) f(0.08)	d(0.98) g(0.02)	d	d	d	d	d	d	c(0.03) e(0.03) f(0.94)
19. Pgdh-A	d	e	c(0.06) e(0.94)	d	d(0.98) f(0.02)	d	a(0.04) d(0.29) e(0.67)	f(0.88) h(0.12)	d(0.77) g(0.23)	j	e(0.07) g(0.46) i(0.47)

tions are summarized in Table 3, and the karyotype phylogeny is shown in Figure 5. Unfortunately, there have been no further chromosome studies on these anoles

incorporating modern banding techniques. Thus, the karyotype phylogeny is based upon several assumptions. The presence of six pairs of metacentric

TABLE 2. ALLELIC VARIABILITY WITHIN AND AMONG SAMPLES OF 12 POPULATIONS OF 11 SPECIES OF ANOLIS. Allelic products are lettered corresponding to increasing anodal electrophoretic mobility and are monomorphic within samples unless otherwise noted. Allelic notation is relative to this data base alone, i.e., allele letters used here do not necessarily correspond to those in other tables in the present study. Populations are numbered in the order they are listed in Appendix II.

Locus	1	2	3	4	5	6	7	8	9	10	11	12
1. M-Acon-A	c	c	b(0.95) c(0.05)	d	c	b	a	b	b	c	c	c
2. S-Acon-A	a(0.50) b(0.17) c(0.33)	b(0.33) c(0.67)	c	b(0.63) c(0.37)	b	b(0.88) c(0.12)	b	b(0.25) c(0.75)	a(0.50) b(0.50)	b(0.50) c(0.38) e(0.12)	d	a
3. Adh-A	a	a	a	a	a	a	a	a	a	a	a	b
4. Ap-A	b	b	b	b	b	b	b	b	b	b	c	a
5. S-Aat-A	b	b	b	e	a	a	a	b	b	c(0.88) d(0.12)	f	g
6. Cbp-1	a	a	a	a	a	a	a	a	a	a	a	a
7. Cat-A	a	a	a	a	a	a	a	a	a	b	b	a
8. Ck-A	a	a	b	a	a	a	a	a	a	a	a	a
9. Ck-C	b	b	b	b	b	b	b	b	b	c	a	b
10. Dir-1	a	a	a	a	a	a	a	a	a	a	a	a
11. Est-1	a	a	a	a	a	a	a	a	a	a	a	a
12. Fum-A	a	a	a	a	a	a	a	a	a	a	a	b
13. Gp-1	a	a	a	a	a	a	a	a	a	a	a	a
14. Gp-2	a	a	a	a	a	a	a	a	a	a	b	c
15. Gp-3	c	c	c	e	c	c	c	c	c	d	b	a
16. Gpi-A	a(0.17) d(0.83)	a(0.17) d(0.83)	a	d(0.88) e(0.12)	c(0.93) e(0.07)	c(0.88) e(0.12)	a	b	d	d	d	d
17. Gdh-A	a	a	a	a	a	a	a	b	b	a	a	a

18. G-3-pdh-A	d	d	a(0.50) d(0.50)	d	b(0.07) d(0.93)	d	d	b	d	b(0.75) e(0.25)	f	c
19. Hadh-A	c(0.17) d(0.50) g(0.33)	d	d(0.88) h(0.12)	c	d(0.72) h(0.07) i(0.07) j(0.07) k(0.07)	c	c	d(0.50) h(0.50)	d	b(0.63) f(0.37)	e	a
20. Iddh-A	a	a	a	a	a	a	a	a	a	c	a	b
21. M-Icdh-A	b	b	b	b	b	b	b	d	a(0.50) b(0.50)	b	c	b
22. S-Icdh-A	e	e	e	c	d(0.93) e(0.07)	e	e	f	e	a	b	e
23. Ldh-A	b	b	b	b	b	b	b	b	b	b	a	c
24. Ldh-B	a(0.83) e(0.17)	a(0.83) e(0.17)	e	e	c(0.64) e(0.36)	c	e	b	f	d	d	g
25. M-Mdh-A	a	a	e	d	e	e	e	b	b	c	a	a
26. S-Mdh-A	c	c	c	c	c	c	c	a	c	b	b	d
27. Me-1	b	b	f	a	f	c	f	e	e	f	d	d
28. Me-2	f	f	c	f	d(0.07) e(0.28) f(0.65)	e	f	f	f	b(0.37) e(0.63)	b	a
29. Mpi-A	e	e	f	e	d	e	e	d	b(0.17) d(0.83)	c	a	d
30. Pept-1	c	c	c	a	c	c	c	c	c	d(0.63) e(0.37)	b	c
31. Pept-2	f	f	d	f	f	f	b	c	c	e	f	a
32. Pgm-A	a(0.83) e(0.17)	a(0.50) e(0.50)	a	a	a b(0.12)	a(0.88) b(0.12)	a(0.25) d(0.75)	a	a	a c(0.50)	a(0.50) c(0.50)	a
33. Pgdh-A	b	b	b	j	a(0.07) b(0.93)	a(0.63) e(0.37)	b	b(0.76) c(0.12) g(0.83) i(0.12)	b(0.17) g(0.83)	d(0.12) h(0.88)	i	f
34. Pnp-A	b	b	b	d	d	d	d	d	d	a	c	d
35. Sod-1	e	e(0.83) f(0.17)	f	b(0.88) d(0.12)	e(0.07) f(0.93)	c	e	c	e	g	a	h

Figure 1. Phenogram of genetic relationships among populations of ten species of Puerto Rican *Anolis* (solid lines) plus five non-Puerto Rican species (dashed lines). Relationships are derived from calculations of Nei's (1972) coefficient of genetic distance (D) clustered using the unweighted pair-group method with arithmetic averages (UPGMA: Sneath and Sokal, 1973: 230). Calculations are based on the analysis of 121 allelic products encoded by 19 presumptive loci. Populations are numbered as listed in Appendix I.

Figure 2. Phenogram of genetic relationships among nine populations of Puerto Rican *Anolis* (solid lines) plus *A. carolinensis*, *A. gadovi*, and *A. grahami* (dashed lines). Relationships are derived from calculations of Nei's (1972) coefficient of genetic distance (D) clustered using the unweighted pair-group method with arithmetic averages (UPGMA: Sneath and Sokal, 1973: 230). Calculations are based on the analysis of 154 allelic products encoded by 35 presumptive loci.

Figure 3. Cladogram of the species of the *cristatellus* subseries based on the allozyme data of the first electrophoretic study (see Table 1). A number of minimum-length Wagner trees were evaluated to conservatively estimate the relationships depicted here although many relationships remain unresolved.

macrochromosomes, 24 microchromosomes and the absence of sexual heteromorphism is considered primitive (Gorman, 1973; Paull et al., 1976). Reduction in diploid number and the evolution of X_1X_2Y heteromorphism are derived conditions. Major macrochromo-

Figure 4. Cladogram of eight Puerto Rican species of *Anolis* based on the allozyme data of the second electrophoretic study (see Table 2). A number of minimum-length Wagner trees were evaluated to conservatively estimate the relationships depicted here although many relationships remain unresolved.

somal rearrangement affects only the Lesser Antillean *bimaculatus* subseries (group III). *Anolis distichus* has the highest diploid number in the *cristatellus* series but also is the only member of the series with possible loss of macrochromosomes; thus its position (IIG) in Figure 5 is open to other interpretations. *Anolis evermanni* (IIF in Figure 5) has the lowest diploid number in the *cristatellus* series (and in the genus) and also has XY rather than X_1X_2Y heteromorphism. Except for the sex chromosomes and the reduction of a pair of microchromosomes, its karyotype is identical to *A. stratulus* (IIB); it is thus considered derived from the IIB state even though its sex chromosomes appear simpler. States IIB, C, and D all have one fewer pair of microchromosomes than IIA. They differ in relative size of smaller chromosomes. It is not certain that they represent three independent lineages, but it is impossible to suggest a single sequence of derivation based simply on

TABLE 3. DESCRIPTIVE SUMMARY OF KARYOTYPES OF PUERTO RICAN AND OTHER EASTERN CARIBBEAN *ANOLIS*.*

Group I.

Primitive karyotype, $2n=36$; 6 pairs of metacentric macrochromosomes; 12 pairs of distinctly smaller microchromosomes. No obvious sex chromosome heteromorphism.

Included species:

<i>A. occultus</i>	Puerto Rico
<i>A. cuvieri</i>	Puerto Rico
<i>A. cybotes</i>	Hispaniola

Group II.

Derived karyotype, usually involving only the microchromosomes which are reduced in number. Sex chromosome heteromorphism is obvious.

A. $2n=31$, $2n=32$

Autosome pairs 1-6 metacentric, large

Autosome pair 7 metacentric, distinctly intermediate in size

Seven pairs of microchromosomes

X_1X_2Y male heteromorphism. Y chromosome is metacentric, approximately same size as autosome 7.

Included species:

<i>A. acutus</i>	St. Croix
------------------	-----------

B. $2n=29$, $2n=30$

Same as II-A except six pairs of microchromosomes

Included species:

<i>A. stratulus</i>	Puerto Rico
---------------------	-------------

C. $2n=29$, $2n=30$

Same as II-B except that there are two pairs of intermediate sized metacentric autosomes (pairs 7 and 8), and thus only five pairs of microchromosomes

Included species:

<i>A. gundlachi</i>	Puerto Rico
<i>A. krugi</i>	Puerto Rico
<i>A. pulchellus</i>	Puerto Rico
<i>A. poncensis</i>	Puerto Rico

D. $2n=29$, $2n=30$

Same as II-C except no distinct size break between intermediate autosome pair 8, and largest of microchromosomes, pair 9

Included species:

<i>A. cooki</i>	Puerto Rico
<i>A. monensis</i>	Mona

E. $2n=27$, $2n=28$

Same as II-D except one fewer pair of microchromosomes

Included species:

<i>A. cristatellus</i>	Puerto Rico
<i>A. scriptus</i>	Southern Bahamas

F. $2n=26$, $2n=26$

Autosome pairs 1-7 as in II-A and II-B

Five pairs of microchromosomes

X-Y male heteromorphism (the only species in the group that is not X_1X_2Y).

The Y chromosome is acrocentric, slightly smaller than autosome 7.

Included species:

<i>A. evermanni</i>	Puerto Rico
---------------------	-------------

TABLE 3. CONTINUED.

G.	2n=33,	2n=34
Macrochromosomes differ from all of the above in that there are essentially 7 pairs, with a sharp break in size between pairs 5 and 6 (alternatively we might say there are 5 pairs of metacentric macrochromosomes, and 2 intermediate sized pairs of metacentric autosomes).		
Eight pairs of autosomal microchromosomes		
X_1X_2Y sex chromosome heteromorphism. Y chromosome is metacentric, just smaller than autosome 7.		
<i>Included species:</i>		
	<i>A. distichus</i>	Hispaniola
Group III.		
Derived karyotype. Microchromosomes and sex chromosomes (X_1X_2Y) as in group II, but macrochromosomes show major rearrangement.		
<i>Included species:</i>		
	<i>A. oculatus</i>	Dominica
	All northern Lesser Antillean anoles from the Anguilla Bank to Dominica	

*For figures and detailed descriptions of karyotypes, see Gorman *et al.* (1968) and Gorman and Atkins (1969).

karyotype morphology. State IIE appears to be derived directly from IID by a loss of a pair of microchromosomes.

DISCUSSION

A phenetic treatment of the first allozyme data set (including all ten Puerto Rican species, plus five additional species of *Anolis*) reveals that two of the extralimital species, *A. scriptus* and *A. monensis*, are very similar to *A. cristatellus* and *A. cooki*, respectively (Fig. 1). These relationships have been previously discussed (Gorman *et al.*, 1980b); they will receive no further comment. Two Puerto Rican species, *A. occultus* and *A. cuvieri*, and the Hispaniolan *A. cybotes* are not similar to one another or to any of the remaining species that were considered. This lack of similarity is further support for our decision (Gorman *et al.*, 1980a) to remove the Hispaniolan cybotoids from the *cristatellus* series as defined by Williams (1976), and also supports the idea that neither *A. cuvieri* nor *A. occultus* is intimately related to the remainder of the Puerto Rican anoline

radiation. Further discussion of the Puerto Rican anoline radiation will exclude consideration of *A. occultus* and *A. cuvieri*.

Five species from Puerto Rico, *A. cristatellus*, *A. cooki*, *A. pulchellus*, *A. krugi*, and *A. poncensis*, are very similar and all phenetically cluster at about $D = 0.2$. The three remaining species, *A. gundlachi*, *A. evermanni*, and *A. stratus*, are phenetically distinctive, clustering with the previously mentioned species at about $D = 0.5$. The electrophoretic distinction of *A. gundlachi* is unexpected given that its karyotype is indistinguishable from that of the "grass-bush" species (Gorman *et al.*, 1968), and that its albumin immunological distances to *A. cristatellus* and the "grass-bush" species are low and indistinguishable, respectively (Shochat, 1976). The apparent lack of congruence among these phenetically-treated data sets may be the result of independent rates of relative evolutionary change; allozymes may be evolving faster in *A. gundlachi* than albumin or chromosome structure. *Anolis gundlachi* phenetically clusters with *A. oculatus* of Dominica, Lesser Antilles (Fig. 1), but

Figure 5. Phyletic relationships of karyotypic character states described in Table 3. Species of the *cristatellus* series expressing each state are indicated.

this similarity is not necessarily indicative of an evolutionary relationship between these two taxa. Further studies may elucidate the relationship of *A. oculatus* to the Puerto Rican assemblage.

Anolis acutus of St. Croix exhibits greater similarity to the main body of the Puerto Rican radiation than do the electrophoretically distinctive Puerto Rican forms—*A. gundlachi*, *A. evermanni*, and *A. stratulus*. The phenetic relationship of *A. acutus* to these forms differs from the cladistic relationship discussed below.

Anolis grahami was chosen for comparison in the second electrophoretic study specifically because Shochat (1976) suggested that the beta anoles of the Jamaican *grahami* series were intimately

associated with the eastern Caribbean *cristatellus*-*bimaculatus* radiation. The genetic distances between *A. grahami* and the species in the *cristatellus* subseries prove to be quite large, but *A. grahami* does exhibit a slightly greater similarity to these species than do *A. gadovi* or *A. carolinensis* (Fig. 2).

By combining the implications of both Wagner trees (Figs. 3, 4), we can attempt to assess some of the patterns of evolutionary relationships. In Figure 3, the species *A. krugi* and *A. pulchellus* form a sister group, the third "grass-bush" species, *A. poncensis*, arises from a common node with the previous pair plus *A. cristatellus* and *A. cooki*. However, in Figure 4 the three "grass-bush" anoles are a unit

forming an unresolved trichotomy. Thus we believe *A. krugi* and *A. pulchellus* to be a sister group, and *A. poncensis* as the sister group of this pair.

Anolis gundlachi is one of the most problematic species in the Puerto Rican radiation. One reason is that it has many autapomorphic alleles. Although Figures 3 and 4 show unresolved polychotomies, *A. gundlachi* could be the sister group of the "trunk-ground" and "grass-bush" ecomorphs. This is concordant with the thoughts of Etheridge (1960), Williams (1972) and Gorman et al. (1968) that *A. gundlachi* represents the descendent of the lineage that later radiated into the other "trunk-ground" and "grass-bush" species.

In both Figures 3 and 4, *A. evermanni* and *A. stratulus* form a pair, and their lineage is not cladistically associated with any of the remaining Puerto Rican species. Some of our equally parsimonious Wagner trees associated *A. acutus* with *A. evermanni* and *A. stratulus* while other trees show no relationship of *A. acutus* to the other Puerto Rican anoles. We believe there is some relationship among these three species but the relationship is not clear.

SUMMARY CLADOGRAM AND CLASSIFICATION

In Figure 6 we present our best estimate of a summary cladogram. Following the proposed terminology of Williams (1976), and our recent proposals (Gorman et al., 1980a), we are discussing the *cristatellus* subseries of the *cristatellus* series. Within the subseries, we recognize two species groups, the *acutus* species group and the *cristatellus* species group. The groups may be distinguished by an osteological character; there are 23 presacral vertebrae in the *cristatellus* species group and 24 in the *acutus* species group. The *acutus* species group is not further divided into subgroups because, despite the fact that *A.*

Figure 6. Summary cladogram of relationships in the *cristatellus* subseries. Allelic synapomorphies are indicated by number: (1) Pept-2 allele c, (2) Me-1 allele e, (3) Gdh-A allele b, (4) Pnp-A allele b, (5) S-Aat-A allele a, (6) Me-2 allele e, (7) Ldh-B allele c. Allele letters are those used in Table 2. One karyotypic synapomorphy is indicated as IIC, corresponding to the state designations of Table 3 and Figure 5. The six species of the *cristatellus* species group share an osteological synapomorphy (23 presacral vertebrae).

evermanni and *A. stratulus* appear to be a natural subgroup, we cannot presently place *A. acutus* and *A. distichus* in this radiation.

The *cristatellus* species group is divided into two subgroups. The *cristatellus* subgroup contains the *cristatellus* superspecies (*A. cristatellus* and *A. scriptus*) and the *monensis* superspecies (*A. cooki* and *A. monensis*). Thus what Williams (1976) called the *cristatellus* species group is now reduced one rank to a subgroup. The second unit is the *gundlachi* subgroup. It contains *A. gundlachi* and the "grass-bush" anoles. Since the Williams classification (1976) includes no divisions between subgroup and superspecies, and since the "grass-bush" anoles are not a superspecies by conventional definition, there are no further available categories to separate *A. gundlachi* from the "grass-bush" anoles. Although it is hard to imagine that anyone will accept yet another division in this hierarchical classification, we could define a *gundlachi* infragroup and a *pulchellus* infragroup. This would unite the "grass-bush" anoles and distinguish

TABLE 4. PROPOSED CLASSIFICATION OF THE *CRISTATELLUS* SUBSERIES.

<i>cristatellus</i> subseries
<i>acutus</i> species group
<i>A. evermanni</i>
<i>A. stratulus</i>
<i>A. distichus</i> superspecies
<i>A. acutus</i>
<i>cristatellus</i> species group
<i>cristatellus</i> subgroup
<i>A. cristatellus</i> superspecies
<i>A. monensis</i> superspecies
<i>gundlachi</i> subgroup
<i>A. gundlachi</i>
<i>A. poncensis</i>
<i>A. pulchellus</i>
<i>A. krugi</i>

them as a lineage separate from *A. gundlachi*. Table 4 summarizes the proposed classification of the *cristatellus* subseries.

ACKNOWLEDGMENTS

This study was supported by National Science Foundation Grants GB-27358 (M. Soulé), B-019801 (E. E. Williams), and DEB 77-03259 (G. C. Gorman). Field and logistic assistance of W. P. Maclean, III of the College of the Virgin Islands, and Jose A. Colon and the supporting staff of the Puerto Rican Nuclear Center greatly facilitated our work. C. S. Lieb, R. D. Orton, and T. L. Vance assisted in the laboratory. Genetic distances and similarities were calculated using a program supplied by E. G. Zimmerman. Wagner trees were estimated with the HENNIG MEMORIAL PROGRAM written by J. S. Farris. We also thank P. Fuerst, M. Nei, and W. J. Rainboth, Jr. for additional data analysis and discussion. Finally, we are indebted to E. E. Williams and Susan Case for thoughtful comments on an earlier draft of this paper.

APPENDIX I

Specimens and localities for the first electrophoretic study. No voucher specimens were deposited. Sample sizes are indicated in parentheses.

Population 1. *Anolis cristatellus* PUERTO RICO: Rio Piedras (15). 2. *A. scriptus* TURKS AND CAICOS ISLANDS: Grand Turk (25). 3. *A. cooki* PUERTO RICO: Hwy. 333 between km posts 2.7 and 3.1 near Guanica (20). 4. *A. monensis* ISLA MONA (23). 5. *A. gundlachi* PUERTO RICO: Luquillo National Forest, near La Mina recreation area (30). 6. *A. pulchellus* PUERTO RICO: Hwy. 186, km 22 near El Verde (30). 7. *A. pulchellus* PUERTO RICO: Hacienda Roses near Utuado (30). 8. *A. pulchellus* U.S.V.I.: St. Thomas: Estate Tutu (24). 9. *A. krugi* PUERTO RICO: Hacienda Roses near Utuado (30). 11. *A. poncensis* PUERTO RICO: 2.3 miles W of La Parguera (9). 12. *A. poncensis* PUERTO RICO: La Parguera (27). 13. *A. acutus* U.S.V.I.: St. Croix (23). 14. *A. acutus* U.S.V.I.: Green Cay near St. Croix (18). 15. *A. stratulus* B.V.I.: Virgin Gorda (26). 16. *A. stratulus* PUERTO RICO: Rio Piedras (27). 17. *A. stratulus* PUERTO RICO: Hacienda Roses near Utuado (10). 18. *A. evermanni* PUERTO RICO: Luquillo National Forest, near La Mina recreation area (30). 19. *A. oculatus* DOMINICA: Roseau (26). 20. *A. cuvieri* PUERTO RICO: Hwy. 186, near km 19 [El Verde Field Station] (11). 21. *A. occultus* PUERTO RICO: Luquillo National Forest, base of El Toro Trail (16). 22. *A. cybotes* HAITI: 9 km W of Marmelade (15).

APPENDIX II

Specimens and localities for the second electrophoretic study. Vouchers are deposited with the Natural History Museum of Los Angeles County.

Population 1. *Anolis cristatellus* PUERTO RICO: Hwy. 186, km 22 (3, LACM 130688-90). 2. *A. cristatellus* PUERTO RICO: Jct. Hwys. 191 and 988 (3, LACM 130691-93). 3. *A. cooki* PUERTO RICO: Salt Plant near Pta. Jaguey (5, LACM 130683-87). 4. *A. gundlachi* PUERTO RICO: Jct. Hwys. 191 and 988 (4, LACM 130694-97). 5. *A. pulchellus* PUERTO RICO: Hwy. 3, NE edge of Luquillo (7, LACM 130676-82). 6. *A. krugi* PUERTO RICO: Jct. Hwys. 191 and 988 (4, LACM 130698-701). 7. *A. poncensis* PUERTO RICO: Lighthouse, Pta. Jaguey (2, LACM 130667-68). 8. *A. stratulus* PUERTO RICO: Hwy. 186, km 22 (4, LACM 130669-72). 9. *A. evermanni* PUERTO RICO: Jct. Hwys. 191 and 988 (3, LACM 130673-75). 10. *A. grahami* JAMAICA: Discovery Bay (6, LACM 130704-09). 11. *A. gadovi* MEXICO: Guerrero: Tierra Colorado (1, LACM

129728). 12. *A. carolinensis* [purchased, no locality data] (2, no vouchers).

NOTE ADDED IN PROOF

Mickevich and Mitter (1981) have noted that the "independent alleles model" for encoding allozyme data, as used in this study, usually yields lower consistency indices compared to other methods such as Transition Series Analysis. The independent alleles model may also be biologically unrealistic in some situations as it does not prohibit the formation of a hypothetical intermediate lacking alleles at a locus. Readers should be aware of the potential limitations of our allozyme analyses.

LITERATURE CITED

- AYALA, F. J., J. R. POWELL, M. L. TRACEY, C. A. MOURAO, AND S. PEREZ-SALAS. 1972. Enzyme variability in the *Drosophila willistoni* group. IV. Genic variation in natural populations of *Drosophila willistoni*. *Genetics*, **70**: 113-139.
- BREWER, G. J. 1970. An Introduction to Isozyme Techniques. New York, Academic Press.
- BUTH, D. G., G. C. GORMAN, AND C. S. LIEB. 1980. Genetic divergence between *Anolis carolinensis* and its Cuban progenitor, *Anolis porcatus*. *J. Herpetol.*, **14**: 279-284.
- ETHERIDGE, R. E. 1960. The relationships of the anoles (Reptilia: Sauria: Iguanidae): an interpretation based on skeletal morphology. Ph.D. thesis, University of Michigan. University Microfilms, Inc., Ann Arbor, Michigan, xiv + 236 pp.
- FARRIS, J. S. 1969. A successive approximations approach to character weighting. *Syst. Zool.*, **18**: 374-385.
- . 1970. Methods for computing Wagner trees. *Syst. Zool.*, **19**: 83-92.
- . 1972. Estimating phylogenetic trees from distance matrices. *Amer. Natur.*, **106**: 645-668.
- FILDES, R. A., AND H. HARRIS. 1966. Genetically determined variation of adenylate kinase in man. *Nature*, **209**: 261-263.
- GORMAN, G. C. 1973. The chromosomes of the Reptilia, a cytotoxic interpretation, pp. 349-424. In A. B. Chiarelli and E. Capenna (eds.), *Cytotaxonomy and Vertebrate Evolution*. London and New York, Academic Press.
- GORMAN, G. C., AND L. ATKINS. 1969. The zoogeography of Lesser Antillean *Anolis* lizards—an analysis based upon chromosomes and lactic dehydrogenases. *Bull. Mus. Comp. Zool.*, **138**: 53-80.
- GORMAN, G. C., AND Y. S. KIM. 1976. *Anolis* lizards of the eastern Caribbean: A case study in evolution. II. Genetic relationships and genetic variation of the *bimaculatus* group. *Syst. Zool.*, **20**: 167-185.
- GORMAN, G. C., AND J. R. RENZI, JR. 1979. Genetic distance and heterozygosity estimates in electrophoretic studies: Effects of sample size. *Copeia*, **1979**: 242-249.
- GORMAN, G. C., R. THOMAS, AND L. ATKINS. 1968. Intra- and interspecific chromosome variation in *Anolis cristatellus* and its closest relatives. *Breviora Mus. Comp. Zool.* No. 293, pp. 1-12.
- GORMAN, G. C., D. G. BUTH, AND J. S. WYLES. 1980a. *Anolis* lizards of the eastern Caribbean: A case study in evolution. III. A cladistic analysis of albumin immunological data, and the definition of species groups. *Syst. Zool.*, **29**: 143-158.
- GORMAN, G. C., D. G. BUTH, M. SOULÉ, AND S. Y. YANG. 1980b. The relationship of the *Anolis cristatellus* species group: Electrophoretic analysis. *J. Herpetol.*, **14**: 269-278.
- KAPLAN, J. C., AND E. BEUTLER. 1967. Electrophoresis of red cell NADH- and NADPH-diaphorases in normal subjects and patients with congenital methemoglobinemia. *Biochem. Biophys. Res. Commun.*, **29**: 605-610.
- LIN, C. C., G. SCHIPMANN, W. A. KITTELL, AND S. OHNO. 1969. The predominance of heterozygotes found in wild goldfish of Lake Erie at the gene locus for sorbitol dehydrogenase. *Biochem. Genet.*, **3**: 603-607.
- LUNDBERG, J. G. 1972. Wagner networks and ancestors. *Syst. Zool.*, **21**: 398-413.
- MICKEVICH, M. F., AND M. S. JOHNSON. 1976. Congruence between morphological and allozyme data in evolutionary inference and character evolution. *Syst. Zool.*, **25**: 260-270.
- MICKEVICH, M. F., AND C. MITTER. 1981. Treating polymorphic characters in systematics: A phylogenetic treatment of electrophoretic data, pp. 45-58. In V. A. Funk and D. R. Brooks (eds.), *Advances in Cladistics*. Bronx, N.Y., The New York Botanical Garden.
- NEI, M. 1972. Genetic distance between populations. *Amer. Natur.*, **106**: 283-292.
- NEI, M., AND A. K. ROYCHOUDHURY. 1974. Sampling variances of heterozygosity and genetic distance. *Genetics*, **76**: 379-390.
- PAULL, D., E. E. WILLIAMS, AND W. P. HALL. 1976. Lizard karyotypes from the Galapagos Islands: chromosomes in phylogeny and evolution. *Breviora Mus. Comp. Zool.* No. 441, pp. 1-31.
- RAND, A. S. 1964. Ecological distribution in anoline lizards of Puerto Rico. *Ecology*, **45**: 745-752.
- SCHOENER, T., AND A. SCHOENER. 1971. Structural habitats of West Indian *Anolis* lizards. II. Puerto Rican uplands. *Breviora Mus. Comp. Zool.* No. 375, pp. 1-39.
- SHAW, C. R., AND R. PRASAD. 1970. Starch gel elec-

- trophoresis of enzymes—A compilation of recipes. *Biochem. Genet.*, **4**: 297–320.
- SHOCHAT, D. 1976. Comparative immunological study of albumins of *Anolis* lizards of Caribbean Islands. Ph.D. Thesis, Louisiana State Univ., School of Medicine, New Orleans, x + 129 pp.
- SHOCHAT, D., AND H. C. DESSAUER. 1981. Comparative immunological study of albumins of *Anolis* lizards of the Caribbean islands. *Comp. Biochem. Physiol.*, **68A**: 67–73.
- SNEATH, P. H. A., AND R. R. SOKAL. 1973. Numerical Taxonomy. San Francisco, W. H. Freeman and Co.
- WILLIAMS, E. E. 1972. The origin of faunas: Evolution of lizard congeners in a complex island fauna: A trial analysis. *Evol. Biol.*, **6**: 47–89.
- . 1976. West Indian anoles: A taxonomic and evolutionary summary. I. Introduction and a species list. *Breviora Mus. Comp. Zool.* No. 440, pp. 1–21.
- WYLES, J. S., AND G. C. GORMAN. 1980. The classification of *Anolis*: Conflict between genetic and osteological interpretation as exemplified by *Anolis cybotes*. *J. Herpetol.*, **14**: 149–53.
- YANG, S. Y., M. SOULÉ, AND G. C. GORMAN. 1974. *Anolis* lizards of the eastern Caribbean: A case study in evolution. I. Genetic relationships, phylogeny, and colonization sequence of the *roquet* group. *Syst. Zool.*, **23**: 387–399.

Modes of Speciation and Evolution in the Sceloporine Iguanid Lizards.

I. Epistemology of the Comparative Approach and Introduction to the Problem

WILLIAM P. HALL¹

ABSTRACT. Comparative studies of speciation mechanisms and their evolutionary consequences in the radiation of ± 120 sceloporine lizard species are introduced. The epistemological foundations of the comparative approach are explained. Illustrations from the sceloporine study show how the approach guided the work and aided interpretation. A foundation problem in evolutionary biology over methods for generating and testing explanatory models traces from one in the philosophy of science over the values of different "logics" of discovery and heuristics for "verification." Popper's model of scientific discovery and deductive falsification is inappropriate for processes like speciation—which are not universal, and where it cannot be assumed either that one set of causes is responsible or that they are deterministic. The worth of the comparative approach is evaluated and its heuristic structure is outlined. In a collection of case histories of a phenomenon, the methodology finds modes of correlation between observations on the cases and markers identifying the phenomenon which may be obscured by randomly unrelated features. If the number of different mechanisms causing the phenomenon is small relative to the case histories, each mode should result from one causal mechanism. Models explaining the observed correlations are realistic to the degree that they logically explain the observed correlations. Their realism may be increased or diminished by further testing, but their "truth" or "falsity" can never be proved.

INTRODUCTION

Mayr (1942, 1947, 1954, 1963, 1970) has argued that essentially all animal speciation results from genetic changes which can evolve between populations of a species only when these populations become allopatrically isolated from one another by geographic barriers to gene flow. Bush (1975b), Bush *et al.* (1977), Wilson *et al.* (1975), and White (1978), among others, using modes of argument which differ from Mayr's, concluded that the majority of animal speciation does not require geographic isolation. They also suggested that much of this non-allopatric speciation involves fixation of chromosomal modifications in initially small, inbred, but not geographically isolated local populations. Bush and Wilson, particularly, also observed that rapid morphological (but not biochemical) evolution is associated with rapid chromosomal evolution, and suggested Goldschmidt's (1955) idea of macromutation as a possible explanation for the putative relationship, resurrecting a controversy laid to rest 20 years previously.

Aside from proposing obvious and non-controversial models for speciation by parthenogenesis and polyploidy, other workers, many following still other and less readily definable modes of argument, have proposed a multitude of models for non-allopatric speciation (e.g.,

¹Queen's College and Department of Genetics, University of Melbourne, Parkville, Vic. 3052, Australia.

Bush, 1969, 1974, 1975; Ehrlich and Raven, 1969; Endler, 1973, 1977; Goldschmidt, 1955; John and Lewis, 1965, 1966; Lewis, 1966, 1973; Lewis and Raven, 1958; Matthey, 1964, 1966; Murray, 1972; Spurway, 1953; Spurway and Callan, 1960; Todd, 1970; Wallace, 1959, etc.). Thus, evolutionists face a Babel of different and frequently incommensurable attempts to explain speciation and to account for its significance in evolution. This situation resembles the crisis which Kuhn (1962) believes precedes a revolutionary change in the paradigm of a scientific community.

The present work will approach the problem of speciation in still another, and clearer way—one which should help reduce the confusing Babel. This series of papers will report systematically controlled comparative studies of mechanisms of speciation and evolution in the sceloporine branch of the Iguanidae (Savage, 1958; Etheridge, 1964; Presch, 1969). The sceloporines include some 120 species (Hall, part II), and enough evidence now exists to allow most speciation events in their history to be reconstructed with some degree of plausibility. From its inception, this research program has had three major goals: 1) to determine how many qualitatively different modes of speciation have occurred and how frequent each was in the proliferation of the sceloporines; 2) to develop and critically test models for the mechanism of each mode, where this has not already been done; and 3) to develop and critically test models to predict the evolutionary potentials for each of the modes. The results then provide a solid foundation from which to extrapolate to explain speciation patterns in more distantly related organisms.

In this, the introductory paper to the series, the characteristics of the sceloporines which led them to be studied are described, and I explain my plan of attack for the study. The organization of this plan is relatively novel and uses unusual but powerful forms of logical argument.

These must be explained and justified scientifically and epistemologically. To do so involves confronting major paradigmatic crises both in evolutionary biology and in the philosophy of science. The second paper (Hall, part II) will present an overview of sceloporine evolution and cytosystematics which is needed to orient the more detailed reports to follow.

THE SPECIES PROBLEM: A CRISIS IN EVOLUTIONARY BIOLOGY

Ghiselin (1974) and others have observed that evolutionary biology now faces a crisis (Kuhn, 1962) in its historical development similar to that before Darwin published his "On the Origin of Species" in 1859, or before Dobzhansky (1937), Huxley (1940, 1942), Mayr (1942), and Simpson (1944) achieved the consensus of ideas known as the "synthetic theory of evolution". These past crises were both concerned primarily (but not exclusively) with major revisions in understandings of the mechanisms by which species evolved through time (e.g., anagenesis; White, 1978). The present crisis, as indicated by the confusion of papers cited in the introduction, relates primarily to the species problem (e.g., cladogenesis—White, 1978). To what does the species concept actually refer (e.g., Hull, 1976)? How are these species formed? What effects, if any, does the mode of species formation have on the evolution of the speciating lineage? The philosophical problems which underlie these questions and make them so difficult to answer also involve the whole of evolutionary biology (e.g., Løvtrup, 1975; Peters, 1976; Platnick and Gaffney, 1978). The array of ideas on species and speciation already proposed, and the obvious problems in answering questions they raise, suggest that something more than a new paradigm in Kuhn's (1962) rather loose sense is needed. Rather, more efficient heuristic

schemata are needed for rejecting unrealistic or patently unscientific proposals.

Kuhn (1962, 1970a) publicized the importance of "paradigms" in scientific research and communication. At the same time, he admits to having hopelessly confused usage of the word paradigm (Kuhn, 1969, 1970b; Masterman, 1970), a confusion not entirely resolved by his introduction of the substitute terms "disciplinary matrix" and "exemplar" (Kuhn, 1969). Kuhn's ideas most relevant here concern scientific revolutions, incommensurability, and communication within and between scientific communities following different paradigms. These ideas also seem to apply to a more precisely defined concept, the "heuristic schemata" of a research program. Based on usage of the two words separately in Suppe [ed.] (1977), the term "heuristic schemata" will be used here to refer to a particular framework of initial metaphysical assumptions, and the pattern of logical argument which is followed within those assumptions, either to test or to discover a theory or putative knowledge. This is not to be confused with specific observations of nature which form the basis of the knowledge, the mechanical apparatus used to collect the observations, or with the theory or knowledge which is discovered or tested by following the heuristic schemata.

Reviewers of my earlier attempts to describe the comparative studies of sceloporine speciation and evolution (e.g., Hall, 1977)¹ have had conspicuous

difficulties in following what I thought were clearly organized arguments used to discover, develop, and test hypotheses. Conversely, I have also found it exceptionally difficult to either review other work intelligibly or to follow it as a model for writing these arguments. From trying to find the source of these long standing communication difficulties, I have concluded that the heuristic schemata used in the sceloporine research, which I collectively referred to as the comparative approach (Hall, 1977), are unusual and poorly understood by many evolutionists. Once I extracted them from my own work I have been unable to find sources where these schemata either are described or are laid out clearly in forms which could usefully serve as exemplars for the approach I have followed. Most importantly, more detailed, but still incomplete studies to understand the problems raised by this apparent incommensurability (Kuhn, 1962) of heuristic schemata suggest that the approach followed here should be more efficacious than others applied to the species problem.

However, although I have modified and/or used some in new contexts, the schemata used in the sceloporine research are not entirely original with me. For instance, as his logical methodology is described by Ghiselin (1969) and Hull (1973), Darwin probably used similar approaches to guide the development and testing of his many revolutionary

¹Reviewers were uniformly impressed by the models proposed in Hall (1977) but seemed unable to follow my arguments. Only when one well-intentioned reviewer took the extraordinary effort to completely rewrite my awkward prose did it become unavoidably apparent that he and I were seeing the same data from incommensurable vantages. At this point I withdrew the manuscript and attempted to understand the problem. The present series of papers is intended to cover the same ground, but with this understanding in hand. Un-

fortunately, because of the wide interest in its conclusions, the 1977 manuscript has already been effectively published by unauthorized but widespread photocopying from copies I circulated for informal reviews. Given the situation, I have no choice but to declare this manuscript to be in the public domain. I hereby grant permission to anyone concerned to either make further copies or to cite the paper. I request that the following citation style be used: "Hall, W. P. 1977. Cascading chromosomal speciation and the paradoxical role of contact hybridization as a barrier to gene flow. Informally published, 91 pp. [cited by author's permission, see Hall, (= full citation to the present paper)]."

theories. Unfortunately, Darwin did not explain or justify his schemata clearly enough for his critics to understand or accept his arguments (Ghiselin, 1969; Hull, 1973) although many of Darwin's most revolutionary theories have been fully substantiated with time. Ghiselin also asserts that many present workers *still* do not understand Darwin's heuristic schemata. I would add here, to emphasize the problems involved in extracting such schemata from the interstices of the work which they organize, that if I have used a Darwinian approach in the sceloporine studies, and if I understand my own approach, then even Ghiselin (1969, 1974) fails to recognize all of the roles or significances of all of Darwin's schemata. This is especially true for Ghiselin's understanding of the approach to originating hypotheses, as compared to practices for testing them, which are described in more detail. Hull (1973) makes the distinction between the two functions of Darwin's research program more clearly than Ghiselin does, but Hull describes neither set of schemata in useful detail.

More recently, Bush (1975); Bush, *et al.* (1977); Wilson *et al.* (1975), and perhaps White (1978) have used comparative approaches which have some similarities with the ones here; but in none of their works are the heuristic procedures adequately explained, justified, or used with full epistemic power. Bush especially can also be criticized for logical and syntactical errors in his probably intuitive usage of these schemata.

Thus, given the present crisis in evolutionary biology over the species problem, the history of my own previous attempts to report my sceloporine findings, and the variety of communication problems which result from the implicit use of different heuristic schemata (Kuhn, 1962), it is desirable to take the unusual step to explain and justify philosophically the *a priori* assumptions and logical structures

of the schemata which I have followed. I illustrate this discussion with examples from the historical development of my sceloporine studies to demonstrate how the schemata have been used.

Because of the problems dealt with, I will unavoidably raise many issues in this paper which demand more complete treatment than I can provide here. For example, the heuristic schemata I present should be axiomatized and tested for logical consistency. Their historical origins should be traced in detail, and they should be compared with schemata used by other evolutionists. However, these issues are beyond the scope of the present work. Such projects would minimally take several years' work, and the present paper is primarily intended: 1) to explain why and how I have organized my sceloporine studies as I have; 2) to explain the unusual logical organization of the arguments I use to reach and defend the conclusions of these studies; and 3) to provide enough of a philosophical justification for this approach to show that it is scientifically valid.

To successfully meet these intentions, certain pitfalls must be avoided. To illustrate some of the traps I will begin with some (probably oversimplified) comments on relevant trends and problems in the discipline of the philosophy of science. Most of my references will be to Popper (1972a,b,c); and to the information on recent developments in the philosophy of science in Suppe [ed.] (1977).

At the outset, despite my harsh comments, I subscribe to most of Popper's philosophy—as philosophic ideals. However, the self-limitations of these ideals should be examined very critically before any attempt is made to apply them to a discipline as complex as evolutionary biology. At least some of the confusion already rampant in the field is due to oversimplistic attempts to use Popperian ideals.

A CRISIS IN THE PHILOSOPHY OF SCIENCE

PROBLEMS WITH INDUCTION AND SEPARATING SCIENCE FROM FANTASY

Epistemology is concerned with the problem: How and how much do we truly know of that which we think we know? The question is central to all science, and it still lacks a generally satisfactory answer. Logic is concerned with developing rules of argument, which if properly followed, will yield a true or at least a probabilistically true output from a true input. The rules of logical argument are central to the program of gaining scientific knowledge, but even here, there are important and unsolved problems. This is particularly true of inductive reasoning.

Induction, as used in science, is the process of inferring conclusions about the existence and details of natural laws from specific observations of the consequences of these laws. In other words, it is the logic of making generalisations from "facts." Popper (e.g., 1972a, b, c) and many others have unquestionably shown that conclusions reached by inductive methods can never be proven to be absolutely true. Yet, how does science gain "knowledge" of natural laws except through the specific observations it makes? This paradox is known as the problem of induction, or Hume's problem (Popper, 1934). A related, and even more important epistemic question is: how does one distinguish factual, rational, or scientific knowledge from fantasies? Popper calls this the "problem of demarcation".

Many philosophers assert that because one cannot argue logically from specific facts to certain truths about natural laws, then no method or schema of inductive discovery can be justified philosophically as a reliable procedure to produce probably, or certainly true conclusions

(Suppe, 1973). For example (Popper, 1972a: 31, 32): "... The act of conceiving or inventing a theory, seems to be neither to call for logical analysis nor be susceptible of it There is no such thing as a logical method of having new ideas, or a logical reconstruction of this process . . ." Or, similarly (Hempel, 1965: 7): "What determines the soundness of a hypothesis is not the way it is arrived at (it may even have been suggested by a dream or hallucination), but the way it stands up when tested, i.e., when confronted with the relevant observational data." Thus, many of these workers have explicitly washed their hands of discovery as being "irrational", and have left it to "psychologists, historians, and sociologists" to explain how scientists find new ideas (Hanson, 1958; Suppe, 1973; and others in Suppe [ed.], 1977).

Somewhat contradictorily, given their position on discovery, many philosophers have argued that repeatedly successful predictions can statistically (or inductively) verify the *truth* of a hypothesis: specific and observable consequences can be deduced with logical certainty from the hypothesis and stated initial conditions. If these predictions are confirmed consistently by a variety of tests, then it can be assumed that the theory is probably true (Hempel, 1965). However, Popper (1972a, b, c) shows in well-justified arguments that no finite number of specific facts can ever absolutely prove the universal theory or generalization to be true. It makes no difference whether these facts come before or after the statement of theory. Arguing from metaphysical assumptions (uniformitarianism for the future is not accepted), Popper goes even further to assert that repeated substantiations do not even show a theory to be "probably" true. This is the problem of induction in another form.

Popper (1972c: chapter 1) claims to have solved the problem of induction in

1933 by accepting that all knowledge remains theoretical, and that although some of this knowledge may be true, its truth can never be proved, even to a degree of probability. This solution makes distinguishing science from fantasy all the more important. How then do we account for the obvious practical successes of science versus fantasy in providing useful understandings of how the world works? Popper proposes an "evolutionary" theory of knowledge, which also answers his problem of demarcation.

According to him, the only logically defensible test of knowledge is that of falsification. If knowledge can be stated in the form of a theory, then specific predictions can be deduced logically from it and given initial conditions. (I pass over the critical and unresolved problem of what a theory is—cf. Suppe [ed.], 1977.) With a proper test, the disconfirmation of a single prediction logically proves the theory to be false, but no number of confirmatory tests will ever prove its truth. Popper then answers his problem of demarcation by concluding that no idea is scientific or rational if it cannot make potentially falsifiable predictions about the empirical world. Thus defined, scientific understanding grows only through proposing bold hypotheses—hypotheses which cover or include the "known" phenomena, but which are "improbable" (or information rich) because they deductively predict heretofore unexpected relationships or situations. These new predictions are then subjected to empirical tests, or in Popper's words "ingenious and severe attempts to refute them." As bold hypotheses are proposed and tested ever more stringently against reality, those which contain high contents of untruth are falsified and selectively eliminated. Thus, a bold hypothesis which survives criticism contains additional information which is not demonstrably false, even though its absolute truth cannot be proved. The net result of Popper's pro-

gram to "discover" scientific knowledge is then to "evolve" selectively an increasing and empirically realistic understanding of the world (Kuhn, 1965; Popper, 1970, 1972a, b, c).

Caveat emptor. As an ideal, Popper's program has much to recommend it, but as a scientist following Popper's advice to critically test generalizations against empirical reality, I find it difficult to apply. There are several traps in the philosophical ideals abstracted above, but a digression into metaphysics and semantics is needed before I can adequately discuss them.

My most basic metaphysical assumptions for my practice as a scientist are uniformitarianism, empiricism or realism, and nondeterminism. I see no reason to assume that anything has happened in the past, or will happen in the future, which cannot at least in principle happen in the present. Nor do I see any justification for assuming the existence behind the empirical world of an ultimate reality of deterministic causality and essences. Many aspects of the empirical world are usefully and pragmatically explained by the theoretical world of nondeterministic quantum physics. For those unfamiliar with the physical evidence for the existence of this world, I recommend Eisberg and Resnick (1974) as a nontrivial introduction to the effects of quantum level uncertainty on more familiar levels of matter. Although fairly rigorous, the nonphysicist should still find this readable. If this quantum physical world is accepted as the basis for reality, the deduced conclusion is that no causal relationship involving matter is deterministic, except to a (sometimes nearly exact) approximation. Quantum level uncertainty leads to uncertainty in radioactive decay and Brownian motion. In turn, these and other nondeterministic effects on molecules directly influence such macroscopically important phenomena as neuron firing; fertilization; and the mutation, recombination, and assortment of chromosomes. Thus, nondeterministic

processes significantly affect the predictability of the development and behavior of individual organisms.

This world view has profound implications for the natural philosophy of evolutionary processes. However, one thing not implied is that these processes must be completely unpredictable. All that is indicated is that they cannot be predicted exactly. Both theory and practice suggest that stochastic or probabilistic predictions are still possible (e.g., Monod, 1971; Morowitz, 1968).

To avoid misunderstanding, either here or in later discussions, the terms deterministic, nondeterministic, stochastic, random, and cause should be defined as I use them. A deterministic relationship between two events or things signifies that the occurrence of one specific event or thing requires that the other also occur with exact and unchanging qualities and properties in comparison to the first. The deterministic relationship is assumed to apply universally. Nondeterminism simply implies that the relationship between two events or things is no longer exactly deterministic. Nondeterministic relationships may be either stochastic or random. A stochastic relationship is where one specified event or thing has a definite but not deterministic effect on the probability of the occurrence, qualities, or properties of the second. In general this relationship can be described by some probability distribution function which may reflect the actions or characteristics of some natural law. A random relationship is where one event or thing has absolutely no effect on the probability of occurrence, qualities, or properties of the second. A causal relationship between two events or things bearing a temporal relationship to one another assumes that the antecedent event influences the occurrence, qualities, or properties of the subsequent event or thing through the action of natural law. This influence may be either stochastic or deterministic, depending on the law. The antecedent event or thing is

the cause, and the influenced characteristic(s) of the subsequent event or thing is the effect.

Given this groundwork, the scientist wishing to use the ideals of the philosophers of science should beware of the following difficulties:

1) From Plato and Aristotle to at least the 1960's most philosophers have been searching for a logical methodology to reveal absolute truth in an ideal world of universal and deterministic laws—a world which science has convincingly demonstrated simply does not exist in empirical reality. By 1969 the philosophers were just getting this message (Suppe [ed.], 1977).

2) Many situations which scientists are concerned with are not universal by any definition of the term (e.g., Kitts, 1977). Many logical arguments philosophers use to support their theses depend crucially on the assumption of universal determinism. For instance, although Popper accepts nondeterminism in his world view (Popper, 1972c), his argument that a generalization can be absolutely falsified by a single disconfirmation is logically invalid if the prediction to be falsified is nondeterministic. For stochastic predictions, no number of disconfirmations will ever absolutely prove falsity. All arguments, whether inductive or deductive, must be criticized on their initial assumptions and statistical merits, not by some ideal of absolute truth or falsity. (But note that the logic of statistical inference also has critical and unresolved foundation problems—Salmon, 1966, 1967.)

However, that hypotheses should be empirically realistic is still the principle which makes science different from fantasy. The idea of a *test* assumes that the generalization predicts empirically observable conditions which may in principle be distinguished from imaginable alternative conditions. If such predictions are confirmed, they support the belief that the generalization is realistic. If they fail, the generalization is probably

unrealistic. However, there are also other ways to eliminate unrealistic generalizations which will be discussed in more detail below.

3) Many philosophers of science, including Popper, may be criticized because they have chosen to ignore processes used to discover generalizations. Although Popper gave his main thesis the title, "The Logic of Scientific Discovery," his program of "discovery" is no more than his evolutionary theory of knowledge as I have outlined it above. His "logic" is simply to be "bold"—to include as many untried predictions as possible under a single covering generalization, and to test as many different bold generalizations as possible—to entail the maximum information content. The logical result of this boldness should be that some increase in realistic information will survive testing. Thus, despite Popper's lip service to the logic of making generalizations, he dismisses the process of discovery as irrational and not worth considering because it involves induction.

My conclusions amplify those of Suppe (1977) and others in Suppe [ed.] (1977): at least until recently, most philosophers of "science" have paid scant attention to how science actually works. Aside from their quixotic search for truth in a non-existent ideal world, they have assumed that one set of heuristic schemata (e.g., the Hypothetico-Deductive schema) is adequate for all science. Most have also ignored the logic and epistemic significance of making generalizations (except for example, Hanson, 1958). Yet real science inescapably involves both making generalizations and assessing whether they are realistic. Both components yield information on the empirical realism of the putative knowledge (Salmon, 1966, 1967; Suppe, 1977), but philosophers have not even fully clarified how realistic generalizations are tested, let alone generated. Nickles (1973) and Suppe (1977) indicate the path to resolving this situation. It is to accept that each

scientific discipline tends to have its own armory of heuristic schemata, which evolve as new practices prove better than older ones in producing empirically realistic knowledge. Philosophers of science should then develop a taxonomy of the various kinds of problems scientists wish to solve, identify the specialized schemata used to solve them, and ascertain the logical and epistemic values of these schemata.¹ Then, perhaps philosophers can innovate and justify still better logics for problem solving. This work is only beginning (Suppe, 1977).

My study of heuristic schemata used in evolutionary biology is incomplete. However, I have isolated those used in my own work from the data and hypotheses within which they are embedded. Once isolated, their functions can be described and their logical and epistemic values for solving particular kinds of problems assessed. The schemata are surprisingly general: they should apply with little modification to a wide range of problems with similar logical structures in a variety of fields beyond biology. My discussion of the schemata is broken into four sections: in the first, speciation represents a class of problems which share many characteristics. Given these characteristics the possible forms of a solution and the possible ways this solution can be reached are constrained. Solving a problem of this type is equivalent to locating and then understanding an unknown but repeated signal hidden

¹Chorley and Haggett [eds.] (1967) and Harvey (1969) have done almost exactly this for geography, which studies some problems that are structurally very similar to those of evolutionary biology. Harvey (1969) reviews and develops the relevant epistemological theory much more completely than I have. These works are commendable exemplars for what could be done with the heuristic methods of evolutionary biology. Also interesting is that they seem to result from a major revolution in the logical methodology of problem identification and solving in geography which is at least as fundamental as that evolutionists now seem to face (see also footnote page 660, below).

in the random fluctuations of a noisy communication channel. The second section describes schemata used to locate those features in the noise which should be included in an explanatory model to solve the problem. This may be called a program for discovery. The third section then discusses how models may be constructed once the components of the potential solution have been located. The final section then discusses several heuristic schemata which may be used to further test the empirical realism of the hypothesis. Each of these will be illustrated by examples of how I have used them in the sceloporine program.

THE PROGRAM FOR DISCOVERY BY COMPARISON

THE PROBLEM TO BE SOLVED: TO EXPLAIN COMPLEX STOCHASTIC BUT ITERATED HISTORICAL PROCESSES

The nature of the problem of speciation. Speciation is typical of problems many evolutionists study. Most evolutionists probably would accept the following generalizations about speciation: the formation of two or more present species from one past species results from historical processes comprised of individually unique and unrepeatable combinations of events which bear spatial, temporal, and perhaps causal relationships to one another. When the question of how one past species became two present species is considered, the only evidence usually available will be observable effects of the past history of events on the present. The effects of some, or many of these past events, may be partially or completely obscured by the effects of other historical events having nothing to do with the case of speciation being considered (i.e., are randomly related to it). Speciation is also a phenomenon where most causal relationships can be assumed to be nondeterministic (i.e., involving mutation, recombina-

tion, migration, selection, and sampling errors of reproduction and death—all of which trace stochastic properties from quantum physics). Although some evolutionists might disagree, there is also no reason to believe *a priori* that there is only one set of qualitatively similar processes which form new species. The desired solution to the problem of speciation is to find the most realistic understanding of the laws of causation governing processes which result in speciation.

As defined here with respect to historical processes, a law is a fundamental property of nature. Although the scientist can never prove that the law is known exactly, it may at least be understood to an empirically testable approximation. The most useful understandings of laws are those which prove to have the greatest explanatory or predictive power. Explanations should logically account for observations already made. Predictions should specify observations to be expected in unexamined sets of the phenomenon in question. Predictions may concern either unexamined properties of the initial cases studied, or they may apply to the same types of properties originally studied but in cases not studied initially. Obviously, if the explanation is based on unrealistic assumptions, if it is not logical, or if its predictions are not substantiated, the understanding is poor and should be revised or replaced.

Generalized statement of the problem. In its most general sense, the problem of speciation is to identify those *kinds* of past events which result in the formation of two or more descendent species from one ancestral species and to explain how this result is achieved. Thus it belongs to a general class of fundamentally similar problems characterized by at least most of the following assumptions about properties of the phenomenon to be explained, "P", and of the world in which P occurs:

- 1) P results from complex historical processes which are evolutionary

(i.e., they involve a series of related events and changes spread through time) and which produce the observable consequence(s) by which P is recognized.

- 2) Each process producing P may be resolved into a finite number of unit events which are unique for that particular example in detail, number, and spatio-temporal relationships (i.e., the process is "unrepeatable").
- 3) Causal relationships among the unit events producing P are governed by stochastic laws. These laws derive from fundamental properties of matter, space, and time and may be explained by them.
- 4) Uniformitarianism applies: nothing has happened in the past which is in principle not observable in the present.
- 5) Several causes may affect one event, and one cause may affect several events. Such multiple causes or causations will obey rational laws of interaction as in 3 and 4, above.
- 6) The only evidence of past events in processes which result in P is their empirically observable consequences in the present.
- 7) These consequences may be obscured to some unknown extent by the effects (or "noise") of events unrelated to P.
- 8) Although processes producing P are unique because of their differences in detail, P's are iterated, and similar consequences produced by similar processes may be examined independently.
- 9) Processes producing P's may be found in various stages of completion.
- 10) The desired solution is to understand how the consequences by which P is recognized are produced and to determine the

range of initial conditions which result in P. The understanding should be as realistic as possible.

Besides their frequency in evolutionary biology, problems with these characteristics (or simpler relatives) are found in disciplines as diverse as the history of science, geology, economics, and psychology, as recognized by Ghiselin (1969, 1974). I would add political science, ecology, geography, and cosmology to this list. The comparative approach should be efficacious for achieving understandings in all of these fields. It involves the successive and/or parallel use of various heuristic schemata and also includes several stages of corrective feedback from nature which help the evolving understanding become more realistic in its explanations of nature. Figure 1 provides a flow chart to indicate the informational relationships among the various components of the entire program of the approach.

SIGNAL AVERAGING SCHEMAS

Overview. The major source of inductive power in the comparative approach is a logical methodology which facilitates generalizations about causal relationships among present variables. It works with sets of facts obtained from observations of the present characteristics of multiple examples of P. The core of the method is a statistical signal averaging or cross correlation procedure: if multiple cases of P exist and can be studied, and if the causes of these P's are iterated,¹ then their effects projected onto present features should be detectable as modes of correlation with one another and with the features that identify P. The "noise" due to random features should not correlate significantly. The added information

¹ If P results from more than one set of causes, at least the number of these sets should be small with respect to the number of cases available for analysis, and there should be several cases of each set.

Figure 1. Flow chart tracing the informational relationships of the heuristic schemata used to discover and test understandings about the causal processes involved in the evolutionary development of classes of historical phenomena.

from correlated features should limit “guessing” for an explanation, since the explanation should be able to account deductively for the coexistence of the newly identified features along with those used to identify P. In other words, the cross correlation procedure provides a logical basis for developing bold hypotheses in Popper’s sense, which will contain an information content *already* shown to be realistic.

More power is gained if stages in the development of P can be correlated across similar developmental stages.

Presumably, initial events will be less obscured in the earlier stages by noise and subsequent events than they will be in later ones. Dated stages should also provide evidence on the temporal evolution of the processes involved.

It is even better if a group of genetically or otherwise related cases can be found which resemble the P cases, but which lack the features of P itself. These may be considered as “controls” for the “experimental” cases which do exhibit features of P. If the experimental or P cases show correlated features *not* found

among the selected controls, this provides strong evidence that these are causally related to P.

The signal averaging principle will be amplified and exemplified, as I describe and discuss the individual steps in the discovery program (Fig. 1). These examples serve to introduce the sceloporine studies.

STEP 1: PROBLEM IDENTIFICATION AND INITIAL SPECULATIONS

General principle. Contrasted with mere data gathering, any scientific research should begin with a problem. The problem will generally first be seen by the scientist as a situation or phenomenon which is inadequately explained by his knowledge of the world. The more clearly the problem can be demarcated and questions relating to it formulated, the easier it will be to develop specific methodologies to find the desired knowledge.

Background information on the sceloporines. While I was an undergraduate in biology at San Diego State University, from 1961 through 1964, I became interested in lizard biology and studied some simple field ecology problems with local species under the supervision of Don Hunsaker II. From my courses (including Richard Etheridge's comparative anatomy of the vertebrates) and informal contact with Etheridge and Hunsaker, both experts on sceloporines, I became familiar with these animals and the problem they posed.

According to Smith and Taylor (1950), the lizard fauna of continental North America included 54 species of *Sceloporus* and perhaps 50 *Anolis* (also iguanids). The next most speciose genera in North American representation were *Eumeces* (Scincidae) with 22 species, and *Cnemidophorus* (Teiidae) with about 17 species. No other genus contained more than about 15 species by this taxonomy. Aside from being remarkably numerous, *Sceloporus* species are also ecologically

remarkably diverse. They are found in almost every habitat available to lizards; over a range from the Canadian border to Panama, from below sea level in desert basins to above timberlines in high mountains, and from extreme tropical rainforests to extreme temperate deserts. In many areas three to seven species occur sympatrically. No other North American genus approaches this ecological diversity or species density,¹ although both *Eumeces* and *Cnemidophorus* have wider geographic ranges due to their occurrence on other continents.

Savage (1958) grouped *Sceloporus* with eight other genera (*Phrynosoma*, *Uta*, *Urosaurus*, *Petrosaurus*, *Callisaurus*, *Holbrookia*, *Uma*, and *Sator*) to form the sceloporine branch of the family. Etheridge (1964) excluded *Phrynosoma* from this assemblage, but acknowledged its close relationship. Etheridge's (1964) osteological observations indicated that *Sceloporus* are not primitive sceloporines. *Sceloporus*, *Uta*, *Urosaurus*, and *Sator* were grouped in a relatively derived position within the sceloporines (see Fig. 2, from Presch, 1969, based primarily on Etheridge, 1964). Although osteology does not differentiate among these four genera, external morphology (development and imbrication of body scales, and the loss of the gular fold) suggests that *Sceloporus* are probably derived even with respect to *Uta* and *Urosaurus*. Also, a comparatively recent derivation is supported by the fact that morphological variation among the 54+ species of *Sceloporus* does not exceed that found in other iguanid genera with only 10 to 15 species. A thesis that *Scel-*

¹*Cnemidophorus*, as it is presently known, may have up to 5 or 6 species occurring sympatrically in areas of the Rio Grande Valley of New Mexico. However, in these cases, all but one or two of the "species" prove to be diploid or triploid unisexual parthenospecies, which clearly are the products of instantaneous speciation. Most of these are proven to be of hybrid origin (Cole, 1975; Parker and Selander, 1976).

Figure 2. Phylogenetic relationships and numbers of species of the sceloporine genera. Phylogeny based on Presch (1969: 286), after Etheridge (1964). Primitive character states designated by capital letters. Small letters indicate derived character states. Species numbers are from Smith and Taylor (1950). By 1966 taxonomic revisions changed *Holbrookia* to 3 species, *Petrosaurus* to 3, *Phrynosoma* to 14, and *Sceloporus* to 57.

Sceloporus had more opportunities for speciation simply because it is an older radiation cannot reasonably explain the striking species diversity. The conclusion from these data showed that the species diversity of *Sceloporus* was extraordinary compared to other North American genera, and particularly to their close relatives.

Hunsaker continually reminded me that this anomaly presented a fascinating

problem to be explained. However, no opportunities in biogeography or specializations in ecology or morphology explained why *Sceloporus*, instead of some combination of the related genera, should have achieved such a striking proliferation of species. The anomalous species diversity most likely was a result rather than a cause.

For a cytogenetics course project (Hall, 1964) also suggested by Hunsaker, I col-

lected all information then available on iguanid lizard cytogenetics (Painter, 1921; Matthey, 1931; Cavazos, 1951; Hunsaker, personal communication; Schroeder, personal communication; Zeff, 1962). Data were found for 11 species (Table 1): five were from genera distantly related to *Sceloporus* and to one another. All were reported to have $2n=36$ karyotypes, which I argued (Hall, 1964) were primitive in the Iguanidae (see Paull *et al.*, 1976, for a discussion of the logic followed). The remaining six species, all *Sceloporus* or their closest relatives (Fig. 2), were reported to have $2n$'s from 22 to 30 or more. Many of these data proved to be wrong, but the concentration of chromosomal diversity in *Sceloporus* has been fully validated (Paull *et al.*, 1976).

First statement of the sceloporine problem. As I finished my undergraduate degree in January 1964, I planned to specialize in community ecology, but my ecological interests did not suggest practical master's thesis projects. With the question in mind—"what explains the extraordinary species diversity of *Sceloporus*?"—I saw that the correlation of chromosomal diversity and species diversity might lead towards this explanation. I did not expect that the problem could actually be solved, but that the

attempt would help me to understand better the species concept as a useful background for studying species interactions in ecological communities. The comparative approach provided the ideal tool for attacking the problem.

Even though $2n$'s were available for only 11 of more than 500 species in the family, because I could see no other obvious correlation with prolific speciation, I thought that the chromosomal diversity might offer a clue (Hall, 1964). Chromosomal differentiation might allow some form of non-allopatric "chromosomal speciation," while more conservatively evolving genera could form species only by slower allopatric speciation (Mayr, 1963). Thus, chromosomally variable lineages might form species besides those formed allopatrically. These extra species would provide added opportunities to evolve a wider variety of ecological specializations. *Sceloporus* was particularly suitable for intensive study, not only because of the possible association between species diversity and chromosomal diversity, but also because its many closely related species offered the possibility of finding relatively early stages in the process of chromosomal differentiation and the possibly associated speciation (i.e., where morphospecies were still "polymorphic"). The compara-

TABLE 1. IGUANID KARYOTYPE FORMULAS AVAILABLE IN 1964.

species	$2n$:formula*	source
<i>Anolis carolinensis</i>	36:12MM,24m	Painter, 1921; Matthey, 1931
<i>Dipsosaurus dorsalis</i>	36:12MM,24m	Zeff, 1962
<i>Crotaphytus collaris</i>	36:12MM,24m	Painter, 1921; Zeff, 1962
<i>Phrynosoma cornutum</i>	36:12MM,24m [†]	Cavazos, 1951
<i>Holbrookia texana</i>	34-36:12MM,22-24m [†]	Painter, 1921
<i>Urosaurus ornatus</i>	30:12MM,18m [†]	Painter, 1921
<i>Sator angustus</i>	~28:12MM,16m [‡]	Hunsaker, pers. comm.
<i>Sceloporus graciosus</i>	30-36:12MM,18-24m [†]	Schroeder and Hunsaker, pers. comm.
<i>Sceloporus undulatus</i>	30:12MM,18m [§]	Painter, 1921
<i>Sceloporus occidentalis</i>	22:12MM,10m	Schroeder and Hunsaker, pers. comm.
<i>Sceloporus olivaceus</i>	22:12MM,10m	Painter, 1921

*MM = Metacentric Macrochromosome, m = microchromosome.

[†]Actual formulas = 34:12MM,22m.

[‡]Actual formula = 30:12MM,18m.

[§]Actual formula = 22:12MM,10m.

tive approach might reveal details that would show how chromosomal speciation differed from other kinds. I also mentioned but did not discuss in detail that variation should be examined in other genera, such as *Anolis* (not closely related to *Sceloporus*) to test further the reality of the correlation between chromosomal differentiation and speciation.

STEP 2: SELECT APPROPRIATE "EXPERIMENTAL" AND "CONTROL" CASES TO ILLUSTRATE THE PROBLEM

General principle. The principle of "controlling" a comparative study is analogous to that of controlling a laboratory experiment. In the laboratory, the experimenter uses an apparatus where an independent variable can be manipulated to learn its effects on dependent variables whose variations are presumed to be causally connected to those of the independent variable. To exclude effects in the output of the experimental apparatus from unknown or extraneous inputs not under the experimenter's control, the experiment is controlled by an apparatus which is as similar to the experimental apparatus as possible, except that the independent variable is held constant. Any variation in the output of the control apparatus is assumed to be experimental artifact, and subtracted from the output of the experimental apparatus. The variation in the dependent variable remaining after subtracting the artifact should result from the causal influences of the parameter manipulated by the experimenter.

The principle in a comparative study is the same, except that "nature" is the experimenter. The investigator selects a set of natural experiments, or cases, Cx_i , from nature which exhibit the diagnostic variables, V^* , which identify the phenomenon of interest, P ; and a set of similar control cases, Co_j , that do not show the V^* of P . Many features, V_c , will be constant, or correlate strongly across both sets of cases. These presumably are

not causally related to V^* but result from the selection procedure. Other features, V_x , will show correlations within the set of Cx_i , but not among the Co_j . These V_x are assumed to be causally related to V^* in the production of P .

Comparative studies differ from the laboratory experiment in that experimental cases may be selected either because they exhibit a presumed "cause" or independent variable, e.g., the fixation of a certain kind of chromosomal rearrangement, or because they exhibit the presumed "effect" or dependent variable, e.g., speciation. Also, the heuristic schema requires no hypothesis to account for the relationship of the variables, V^* and V_x , or even an idea of what the V_x should be. The relationship is demonstrated as a logical consequence of the causal connections between the variables, and not by preconceived beliefs of the investigator. In practice, there may be a working hypothesis to explain P , which can help to select potentially important variables to study. This kind of selection is useful, if not too restrictive, since it is impractical to study all aspects of the cases (this is the difficulty with Baconian induction). On the other hand, preconceptions should not influence selection of *which* experimental and control cases to study. This should be determined arbitrarily by the way the problem is defined. In biological systems, the most obvious and probably least biased controls would be lineages which ecologically parallel the experimental lineages, and which are phylogenetically closely related to them. If such controls are not practical, depending on the problem, controls should be selected which meet at least one of these criteria.

Sceloporine experiments and controls. The research question formulated to guide the selection of cases for the comparative study of sceloporine speciation was based on the anomalous species diversity of *Sceloporus* relative to the other North American lizard genera: is

this anomaly explained by the fact that *Sceloporus* formed "extra" species by nonallopatric speciation mechanisms not available to conservatively speciating genera? Does this involve chromosomal differentiation, and if so, what other feature(s) result from this speciation mechanism (or mechanisms), and how is this constellation of features produced? The obvious experimental cases were speciation events in *Sceloporus*, and the obvious controls were speciation events in the other eight sceloporine genera.

Although there was no evidence, initially, to determine which *Sceloporus* species may have resulted from the supposed non-allopatric speciation mechanism(s), it was assumed that *Sceloporus* probably included many cases of non-allopatric speciation in the evolution of its 54+ species. However, some *Sceloporus* species were presumably also formed allopatrically. Although later in the study these would provide the best controls for nonallopatric speciation, initially, the cases of allopatric speciation would represent noise or artifact among the experimental cases. Thus appropriate control genera should be studied to identify features that correlate only with allopatric speciation, so they could be subtracted from the *Sceloporus* data base, to leave primarily those cases likely to have resulted from nonallopatric speciation.

The other sceloporine genera serve as controls for identifying the allopatrically speciating *Sceloporus*. These are *Petrosaurus* (2 species), *Phrynosoma* (14 species), sand lizards—*Callisaurus*, *Holbrookia*, and *Uma* (~10 species), *Uta* (6 species), *Urosaurus* (~10 species), and *Sator* (2 species). According to Savage (1958), Etheridge (1964), and Presch (1969), all share a close common ancestry with *Sceloporus*. The biogeographical opportunities for speciation appear to have been similar for all of the genera except the insular *Sator*: all have distributions centering on the North American deserts, and belong to the "New North-

ern Faunal Element" of the North American herpetofauna (Savage, 1960, 1966) which evolved *in situ* along with the development of the North American deserts (Axelrod, 1950, 1958; Axtell, 1958; Norris, 1958; see also Morafka, 1977). Within this desert environment, the sceloporine genera show three main ecological specializations: *Phrynosoma*, which are specialized nomadic anteaters; the sand lizards, which are cursorial insectivores that run after their prey, and which normally perch directly on the ground or small stones; and the *Uta-Sceloporus* assemblage, which normally perch off the ground (on rocks, trees, bushes, etc.), and wait for prey to come within easy striking distance. At least *Uta* and *Urosaurus* should ecologically parallel *Sceloporus*. Other controls are needed to distinguish speciation of *any* kind from non-speciation. These are provided by studying different populations included *within* operationally defined species.

Extending the idea of the natural experiment, if the correlation between chromosomal diversity and species diversity seen in the sceloporines results from generally applicable natural laws, rather than from some unique specialization of *Sceloporus*, then the "experiments" should be repeatable: similar patterns of correlation should be seen when other prolifically speciose genera such as *Anolis* and *Liolaemus* are compared with their appropriate controls.

STEP 3: INITIAL DATA COLLECTING

General principle. Once appropriate experimental cases and their suitable controls have been selected, then as much information as practical relating to each should be collected: a "history" should be developed for each case. The inductive power of the methodology depends on *not* limiting observations to those parameters which would support the preconceived speculations. Every attempt should be made to survey a

variety of parameters, some of which might potentially relate to P, and some of which should not.

Data collected or available from the sceloporines. In my studies of the sceloporines, besides collecting information on a variety of cytogenetic parameters from as many species as practical, I have observed reproductive biology, anatomy, population biology, and ecology. Also, I have tried to keep track of all published work dealing with any aspect of the biology of the genera being studied. These data will be presented in the more detailed papers to come later in the present series.

STEP 4: CROSS CORRELATION ANALYSES

General principles: Signal averaging schemata. The major inductive power in the comparative approach comes from heuristic schemata which facilitate generalizing from specific observations to assumptions about causal relationships among the variables studied from the case histories. At the same time the methodology keeps these generalizations in touch with reality. Modes of inter-correlated variables which are probably causally involved with P can be identified independently from any *a priori* speculations. The schemata are developed by analogy from an inductive statistical procedure known as signal averaging.

The concept of signal averaging is well known in neurophysiology (Glaser and Ruchkin, 1976) where a frequently encountered problem is to extract iterated, but otherwise unknown signals, which are individually completely hidden within random noise in an input channel. This is the simple signal averaging schema.

Assume that a noisy input includes a signal in the form of a voltage fluctuation which has a fixed time relationship to some identifiable marker event. A computer or tape storage can then be triggered by the marker event to record

the voltage fluctuations during the time sequence believed to contain the unknown signal. The marker triggered recording sequence is repeated many times with the same time delay relative to the marker event. Each recorded sequence is placed in register with the triggering events and added algebraically to sum the recordings. As many recorded cycles are added, voltage deviations which are randomly related to the marker will tend to an average or null value, since positive deviations for a given time delay after the marker will tend to be as frequent for negative deviations at the same delay. However, any repeated signal embedded in the noise which shows a constant voltage fluctuation relative to the marker, will always show the same slight positive or negative deviation added to the noise. With enough repetitions, a generalized signal will eventually emerge from the masking noise as a statistical average of the repeated signals. Assuming unlimited repetitions and a constant signal, the signal-to-noise-ratio can be increased to any desired value by continued averaging, even though the signal is undetectable in any one repetition.

A slightly different schema—an autocorrection procedure—can be used to extract unmarked signals from a noise channel if they are repeated at a constant interval, even if nothing is known about the signal's periodicity or other characteristics. Here, one long recording from the input channel is chopped into many short segments of a given duration, and the segments are then added together in register relative to the chopping points. Any recurring signal in the chopped recording which has a simple harmonic relationship relative to the period determined by the duration of the chopped segment will add algebraically. With the summing of enough segments, the signal will emerge from the masking noise as in the simple signal averaging schema. With a computer, a long recording from the channel can be systematically chopped

and added this way for a variety of different duration periods until a signal with the highest signal-to-noise ratio is found for a constant number of additions. This presumably occurs when the length of a single chopped segment exactly coincides with the actual repetition period of the original signal. Other procedures for extracting signals from noise have also been developed, but the basic principles of signal averaging illustrated above are all that are needed for the analogies to be developed here.

Cross-correlation schemata. Analysis of data from complex, nondeterministic, but iterated historical processes involves extracting signals produced by causal events in the past. These signals are transmitted from the past through the noisy communication channel of time onto the event surface of the present, and need to be extracted from the variety of "noise" generated by random historical events. Simple signal averaging from a communication channel involves making essentially longitudinal correlations of input voltage deviations stored along the linear axis of a recording tape. By analogy, in a comparative study, when case histories are made of the present states of parameters of the selected experiments and controls, information on the variables is stored on various parameter axes (one for each parameter examined) of a data matrix for each case. The stored information in these data matrices or case histories can then be averaged to search for correlations among their parameters. This is exactly analogous to what is done with the one dimensional data matrices of the signal averaging schema. Any variables bearing relatively constant relationships among the data sets of the case histories should stand out as strong additive correlations against the "noise" due to unrelated events. I call this the *n*-dimensional cross-correlation schema.¹

To clarify the analogy, in signal averaging the input triggered by a marker event is recorded as a voltage deviation along the linear time axis established by the uniform movement of a tape past a recording head, or by the sequential filling of adjacent "bins" in a computer memory. If records associated with triggering events are stored individually for later off-line analysis, each recorded time deviation sequence is associated with a unique address for each specific triggering event. Thus, a one dimensional cross-correlation matrix is formed. For each specific triggering event, the input deviations are recorded along the one dimensional time axis. Note that the filled data matrix is already three dimensional in a physical sense: one dimension is established by a sequence of addresses corresponding to the sequence of individual triggering events. A second is established by the time axis of the recorded sequence for each triggering event. And the third is required to hold the specific deviations for each given instant of delay along this

cently begun to think and work with multi-dimensional "data matrices" very similar to the idea I propose here. Matrix manipulations and multivariate statistical procedures (many of them from numerical taxonomy!) are applied to identify modes of correlation and relationships among the various parameters of these matrices (e.g., Berry, 1964; Haggett and Chorley, 1967; Harvey, 1969; King, 1969). To quote from Harvey (1969: 347-348):

We are concerned to find general classes and general relationships among attributes, we are concerned with finding comparable underlying structures in complex data matrices, and, above all, we are concerned with identifying a theory about structures which can command our confidence as an analytic, retrodictive, or predictive device. *Quantitative techniques for classification bode well as search procedures. They can lead us to new ideas, new frameworks for analysis, and so on.* [My italics]

Although all of these points are touched on by Harvey (1969), the geographers have not focused on the ideas of selecting natural experiments and their controls, the epistemic contributions of the process of discovery to the realism of explanations derived from the process, or the ways the empirical content of the explanation can be increased through different kinds of attempts to refute it.

¹As part of their revolution in heuristic methodology (see footnote p. 650), geographers have re-

time axis. The average signal is then extracted from the matrix by averaging all of the individual input voltage $\times$ time histories to form the single 2-dimensional plot of the iterated signal.

When the signal averaging schema is modified to extract information from the effects of past events on the present, the method of selecting the cases to be examined is exactly analogous to the triggering event in the one dimensional signal averaging paradigm. Clearly, just as in triggering a recording from a noisy communication channel, the success of the n -dimensional procedure will depend on how well selection has limited the cases to be studied to one, or at most a few, underlying causes relative to the number of cases to be studied. In the sceloporine lizards, by present count (Hall, part II) there are about 57 species which are known to be, or probably are, chromosomally derived. Each of these derived species may then be the reflection on the event surface of the present of one or more instances of a past event of "chromosomal speciation". The remaining approximately 65 species in the radiation all have, or are suspected to have, identical, presumably conservative $2n=34$ karyotypes. These almost certainly do not reflect past chromosomal speciation. The radiation includes approximately 120 species in total. Each pair of species will include one or more speciation events in its derivation from a common ancestor. When averaged, the cases should form at least two modes of correlation—one resulting from allopatric speciation and the other(s) from non-allopatric speciation. Differences between these pairs can also be compared with differences between paired populations belonging to a single species. Once cases have been selected, an n -dimensional data matrix is established for each case history. A different axis of the standard data matrix is used to store observations for each kind or parameter being studied across the selected cases.

If features of the selected cases reflect

iterated but still unknown underlying mechanism(s), and if all cases have similar ages relative to P (i.e., if the triggering event bears a relatively precise time relationship to the signal), then when a given axis is averaged from the data matrices from the selected cases, states of a parameter having a specific causal (either caused or causing) relationship to the process should frequently coincide. Conversely, states of a parameter only randomly associated with the process should be randomly distributed and should average to some neutral value. In some cases, effects due to unrelated processes may also add to the value of a causally related parameter, but presumably the deviations from these other effects would also be random with respect to the value causally related to P. In other words, the effects of random events should be exactly analogous to the noise in a communication channel. For causally unrelated parameters, observations would include only noise, and should average null as the number of record matrices examined is increased. For a causally related parameter, the observations will include a signal embedded in the noise, which should give increasingly stronger correlations as the sample size is increased. Similarly, causal relationships between different parameters will show up as cross correlations across the parameter axes involved.

Here the importance of having "controls" for experimental cases should be obvious. The controls establish a null or baseline value for a parameter which contains only noise relative to experimental cases. Strongly intercorrelated values from experimental cases which differ from this null are fairly conclusively related causally to the factor(s) which led these cases to be designated as "experiments".

However, actual historical processes will rarely be of the same age or at the same degree of completion, and values of important parameters may vary causally according to the stage of completion of

the process. This presents a problem which makes the schema more complex, but its resolution contributes importantly to the inductive power of the approach. If available information (e.g., taxonomy and phylogenetic reconstruction in the sceloporine example) allows the cases to be segregated into subsets that can be ranked according to approximately equivalent stages in the process, then it is probably most efficient to proceed by doing so. Each subset of case histories resulting from this segregation is then averaged to identify correlated parameters associated with particular stages of the process. A tentative model generated from such ranked subsets may suggest better criteria for re-segregating and re-ranking; which may, in turn, suggest revisions in the model. This negative feedback process should lead through a series of successive approximations to a stable relationship between the arrangement of the data and the theoretical generalizations drawn from the data. If the process does not converge to a stable relationship, the basic model is probably fundamentally unrealistic and other, completely different models should be considered.

In many instances it may not be obvious how to segregate the cases, or the attempt to do so may have given no useful results. An auto-correlation schema may then be used to randomly segregate the cases into variously sized subsets. Each possible combination of cases is then independently checked for cross correlation of information. The procedure is analogous to auto-correlating from a communication channel where both the periodicity and characteristics of the signal are originally known. As applied to historical problems, the schema may be termed auto-correlation by random association.

If many cases are available for study, and data have been collected for many parameters of each, auto-correlation by random association would be extremely laborious if performed manually. But

data involved in such historical studies may not lend themselves to computer coding and analysis either. It is intriguing to speculate that human memories may be organized like the n -dimensional data matrices described for the cross-correlation and auto-correlation schemas (Harvey, 1969). Possibly human memories automatically correlate these matrices by random association as part of the unconscious filing system. I would even suggest that this process might account for the powerful inductive "intuition" of some comparative biologists. However, the auto-correlation process has enough logical rigor so computer processing could be used where observations can be suitably quantified, and it is also one which can be performed manually by hand sorting where few enough recording matrices (e.g., data entered in a standard format on a page) are involved to make it feasible.

In summation, signal averaging schemata provide logical methodologies to reveal correlations among variables from a given number of empirical observations of real-world situations. These correlations provide objective statistical evidence to support the inductive generalization that the correlated variables are causally related. Similarly, the methodology will also provide objective statistical evidence on the absence of causal relationships among other variables which do not show correlation. These generalizations are based directly on the evidence, completely independently from any preconception which may have been held before the observations were made. The problem of model building then becomes a greatly simplified one of plausibly explaining causal relationships which are corroborated by already existing statistical evidence. This aspect of the comparative methodology may be sufficient to explain "the truly amazing feature of Darwin's intellect" which accounted for "the frequency with which he was able to 'guess' correctly, even though he lacked the requisite data

and anything like an adequate theory governing the phenomena" (Hull, 1973: 77).

CONSTRUCTING EXPLANATORY MODELS

STEP 5: MODEL GENERATION

General principles. When certain variables correlate across cases selected because they demonstrate some particular P, this correlation provides evidence to suggest that the variables represent effects of past events in the causation of P. Model building then involves developing a covering explanation that requires the observed correlations as logical consequences. This resembles Hanson's (1958, 1961) program of retrodiction (Achinstein, 1971). Obviously, explanations can require conditions other than those already observed. An explanation may depend on various assumed initial conditions, and unanticipated additional consequences may follow logically from these assumed initial conditions. These dependencies and requirements beyond the original observations provide opportunities for testing the realism of the explanation. However, at the very least, any realistic model should account logically and simply for those conditions already shown to correlate with P. Similarly, a realistic model should not require conditions already observed not to exist.

Developing a realistic explanation does not test its realism, yet a program which discovers information helping to explain P contributes epistemically to the realism of any explanation which logically requires this information. Examining the alternative, to have no discovery program, demonstrates this: on probabilistic grounds, explanations which logically require consequences already observed to correlate with P should entail more empirically realistic information than would any single attempt to explain P by a random guess, no matter how "bold" the guess. Irrationally gener-

ating and then testing many different bold hypotheses might eventually produce an empirically realistic explanation for P which entailed as much information as an explanation yielded by the discovery program. However, for such explanations with similar contents, it should make no epistemological difference whether observations corroborating the required circumstances are made before or after developing the explanation. Note that this argument assumes that the information content of the explanations derives from examining the same number of variables in the same number of cases (cf. Popper, 1972c). In practice, guessing for fruitful explanations is difficult when few data limit the guesses. It is more efficient (and more justifiable epistemologically) to collect a series of related case histories in advance of guessing, and scan them for correlations. Large domains of possible guesses may then be avoided because observations already made contradict some conditions demanded by the domains. Thus, attention for guessing is focused logically on what is frequently a very much smaller domain of possible explanations not already contradicted by observations.

Models to explain modes of sceloporine speciation. I will show in detail later in this series that features of the cases of sceloporine speciation group into two particularly distinct modes. The characteristics of the two modes will be abstracted here from Hall (1973, 1977) to provide background for illustrating the logic followed to explain them. In one mode, pairs of closely related species show a sibling type relationship and belong to conservatively evolving radiations containing few species. The paired species frequently have an obvious history of allopatric isolation from one another and do not differ chromosomally. Usually, where species are presently in geographic contact they are so similar ecologically that close sympatry is prevented. The allopatric speciation model (Mayr, 1963, etc.) easily explains all of

these features, so this mode will not be considered further here.

In the second mode, pairs of closely related species have ancestor-descendant relationships and belong to predominantly linear sequences of phylogenetic derivation. These lineages frequently contain many species. Paired species show little or no evidence of past allopatric isolation from one another, and they frequently differ enough ecologically to coexist in extensive sympatry or syntopy. Paired species in lineages frequently differ by chromosomal rearrangements which can potentially cause meiotic malassortment, thereby reducing fitness of heterozygotes (i.e., heterozygotes show negative heterosis). Chromosomally primitive species near the origins of sequences of chromosomal derivation tend to be ecologically conservative, while highly derived species towards the ends of sequences either have extreme ecological specializations or show impressive ecological dominance. Most sequences of chromosomal derivation involve only one type of chromosomal rearrangement (e.g., all fissions or all fusions), and it seems that rates of speciation accelerate towards the terminations of the sequences of derivation. Terminal species at the ends of sequences of derivation frequently have either exhausted the karyotypic substrate for the particular type of chromosomal rearrangement involved in the sequence, or show polymorphisms for that kind of rearrangement. The most closely related populations known to differ chromosomally, form narrow hybrid zones where the populations meet geographically (e.g., Hall and Selander, 1973). Paradoxically, although hybrids in hybrid zones are fertile and backcross, there seems to be a complete block to gene flow between the chromosomally different populations. No sceloporine species outside of *Sceloporus* shows any evidence for chromosomal differentiation; and even within *Sceloporus*, speciation events between close relatives that differ

chromosomally are considerably less frequent than allopatric speciation events not involving chromosomal differentiation. Yet, a greatly disproportionate number of *Sceloporus* species have a background of recent chromosomal derivation in their phylogenetic histories.

As observations forming this second mode accumulated, I developed a chromosomal speciation model to account logically for the correlations that emerged (Hall, 1973, 1977). The complete explanatory model resolves into three major components, each of which explains some of the features listed above. One part of the model describes how negatively heterotic chromosomal rearrangements can become fixed in populations and initiate speciation. Another explains how hybrid zones between chromosomally differentiated populations function to block gene flow. The third part of the complete chromosomal speciation model explains: 1) the disproportionate evolutionary successes of chromosomally derived species in comparison to species formed allopatrically, 2) the predominantly linear nature of sequences of chromosomal derivation, and 3) details of patterns of chromosomal variation within the lineages. These partial explanations were developed in the order listed, to account for particular correlations in the growing body of information on sceloporine speciation. Without attempting to detail their actual historical development, I abstract the reasoning followed to reach these explanations, to illustrate the logic of model generation.

The chance fixation model. The first and most obvious correlation seen with the prolific speciation of *Sceloporus* related the fixation of chromosomal differences between species to prolific speciation. Species differ by Robertsonian mutations, which, at least theoretically, are negatively heterotic because they cause meiotic malassortment (White, 1973, 1978). Thus, heterozygous hybrids and backcrosses between

chromosomally different populations should be less fit than either homozygous parental type. Two consequences follow directly from this assumption: 1) Reduced hybrid fitness should serve as a partial barrier to reduce gene flow between the chromosomally different homozygous populations. This will make it easier for them to evolve independently. 2) Reduced hybrid fitness should also selectively favor individual genotypes which avoid hybridization. Thus, the chromosomally differentiated populations should rapidly evolve isolating mechanisms to prevent hybridization. Both consequences would work in the absence of allopatric isolation to speed the evolution of a completed barrier to gene flow between the chromosomally different populations. Neither consequence would occur if the populations were geographically isolated from one another.

Thus, in theory, negatively heterotic chromosomal differences can aid speciation; but how are the differences first established? Any mutation which has a strong enough negative heterosis to favor the evolution of a barrier between incipient species could not become fixed in a large, randomly breeding population. This is because the Hardy-Weinberg equilibrium frequency equals the frequency of the new (rare) chromosome, while the frequency of the neutral or advantageous homozygous mutant genotype equals the square of the frequency of the new arrangement. In a large population, essentially all of the chromosomes carrying a new rearrangement will occur in the selectively disadvantageous heterozygous state. Hence selection will quickly eliminate it, even if it potentially has a strong selective advantage in the homozygous state.

However, if the negatively heterotic mutations should occur in a very small population, statistical sampling errors in mating (= genetic drift) become evolutionarily significant. Here, a negatively heterotic mutation has an evolutionarily

interesting possibility of becoming fixed by chance (about one chance in 10^{-3} for a population size of 10 and a 50% reduction in heterozygote fitness [Wright, 1941]). Also, selection against negatively heterotic mutants constantly works to eliminate whichever chromosome arrangement is rarest in a population. If a new mutation once passes 50% frequency by drift, selection will work to push it to complete fixation, even against some immigration of individuals carrying the ancestral chromosome. Hence, as long as the small population remains predominantly inbred, small amounts of outcrossing per generation (say up to 10 or 15%) will not greatly affect the probability of fixing a new mutation. Given reasonable conditions of population structure, fixation of the chromosomal rearrangement may occur entirely without allopatric isolation of the populations in the classical sense.

In this partial model, initiation of chromosomal speciation requires certain antecedent conditions: 1) Chromosomal heterozygotes are substantially less fit than either homozygote. 2) Negatively heterotic mutations occur with appreciable frequency. 3) These negatively heterotic mutations occur in populations which include many small inbred demes. It follows from these initial conditions and the chance fixation model that: 1) Chromosomally derived species will be "founded" by initially very small demes within which the chromosomal rearrangements first become fixed. 2) Such founder populations occur within the geographical range of the chromosomally more primitive ancestral stock. 3) Competitive interactions between the founder population of the chromosomally derived incipient species and the much more massive ancestral stock should force the derived species to differentiate ecologically away from the niche of the more massive ancestral stock. Thus, given only the correlation between chromosomal diversity and species diversity, other observed correlations follow logically from the most straight-

forward guess to explain the first correlation: species will be formed in an ancestor-descendent relationship, and chromosomally derived species will be ecologically derived relative to their ancestral species.

However, predictions of this simplistic chance fixation model seem to be contrary to the evidence in at least one respect: selection against parents which hybridize to form the negatively heterotic chromosomal heterozygotes should lead to the rapid evolution of premating isolating mechanisms to prevent this hybridization. Evidence from the narrow hybrid zones in *Sceloporus grammicus* (Hall, 1973; Hall and Selander, 1973), and similar studies in other organisms (e.g., Nevo and Bar-El, 1976; Szymura, 1976a, b; Szymura *et al.*, in preparation; White *et al.*, 1964, 1967, 1969), suggests that selection in these hybrid zones does not lead to the evolution of premating isolation. Paradoxically, although the failure to evolve premating barriers allows frequent hybridization and backcrossing rather than just reducing gene flow, the hybrid zones appear to completely block it. The second partial model was developed to explain this paradox.

The hybrid sink model. If populations differ by a negatively heterotic mutation and hybridize, their hybrid zone will include a relatively high frequency of less fit heterozygotes. Because of the negative heterosis of the hybrids, populations in the center of the zone will have a reduced productivity compared with pure homozygous populations outside of it. Thus, hybrid populations will put less pressure on the carrying capacity of their environment than will the pure populations. Consequently, there should be a net migration or diffusion of individuals from pure populations towards the reduced pressure of the hybrid zone. Migrating individuals will carry their genes towards the "sink" for gene flow, represented by the chromosomally unbalanced (and therefore genetically

lethal) gametes of the chromosomal heterozygotes and the balanced gametes that may combine with them to form lethally unbalanced zygotes. It seems reasonable that the diffusion gradient towards the hybrid zone may become steep enough to prevent any gene from leaving the vicinity once it enters the zone, and that this may happen well before a complete sterility barrier evolves. Hence, the hybrid zone may completely block gene flow between the pure populations even though hybrids and backcrosses retain appreciable fertility. It also follows that the residence time of any gene in the hybrid zone should be relatively short, as long as individuals carrying this gene cannot discriminate absolutely against mismating with chromosomally different individuals. Thus, the sink would probably prevent the evolution of premating isolation in the hybrid zone, even though selection favors this. No single gene mutation would be likely to confer its carrier with a complete discriminatory ability to avoid mismating, and any gene conferring only partial discriminatory ability would almost certainly be lost in the sink before other mutations allowing partial discrimination could combine with it to produce a perfected isolating mechanism.

Many details of the fully developed hybrid sink model have been suggested by field observations of the sceloporine hybrid zone. These will be discussed in later papers of this series. However, of concern here is that hybrid sinks should exhibit surface tension (Key, 1968, 1974). Populations on the concave side of a curved hybrid zone will have a shorter contact front than will those on the convex (or outer) side of the curvature. If other circumstances are equal, the population on the concave side cannot feed as many immigrants into the sink as can the other population. Consequently, the larger number of immigrants from the convex side will push the center of the zone towards the concave side. Where

populations meet on a broad front, this surface tension effect will tend to straighten out any kinks in the contact. But where a newly differentiated founder population occupies only a very small area within the range of its parental stocks, the surface tension will press inward around this entire circumference. Many newly differentiated populations must be quickly swamped by this effect, thus raising the question: could any chromosomally differentiated populations survive swamping in this critical early state? Attempts to answer this question and to account for the impressive ecological dominance of the products of one chain of derivation in *Sceloporus* led to guessing for the third part of the chromosomal speciation model.

The cascading or chain speciation model. Three sets of assumed or demonstrated circumstances provide the basis for this explanation, which also accounts for the remaining features correlated with rapid speciation in *Sceloporus*: 1) Chromosomally differentiated species originate as very small founder populations. 2) The probability that chromosomal differentiation will occur at all is profoundly affected by: a) rates of chromosomal mutation, b) behavior of the chromosomes in meiosis and the genetic consequences of any meiotic errors, c) details of the species' mating system, and d) details of its population structure. 3) The aspects of a species' genetic system listed above are genetically controlled. These circumstances allow evolution by two unusual mechanisms, a group selection effect and a positive feedback or deviation amplifying process (Szarski, 1971). These may counteract natural selection working at the level of single individuals.

The genetic system parameters listed above can vary considerably and still have minimal effects on individual fitness. Hence, if loci determining them are polymorphic, demes small enough for fixing chromosomal rearrangements may

also show substantial variation in these parameters due to genetic drift. Over the range of a species which has an appropriate population structure, through chance some demes will have a much more favorable genetic background for chromosomal speciation than will others. It follows that chromosomal speciation will more likely begin in these "favorable" demes than in others. Founder populations that survive as new species will tend to perpetuate the favorable genetic backgrounds. Thus, chromosomally derived species will, on the average, offer more favorable circumstances of further chromosomal speciation than will ancestral species. This positive feedback amplification process may work in each chromosomal speciation event in a sequence to produce a predominantly linear and increasingly rapid chain of chromosomal speciation. (see Hall, 1973, 1977, and later papers in this series). Three circumstances (2 intrinsic and 1 extrinsic) can terminate such a chain of speciation.

1) If chromosomal fissions and fusions are qualitatively different kinds of mutations, different genes should affect their mutation rates. Similarly, different genes may control their meiotic behavior. Thus, if a speciation event involved centric fissions, then positive feedback effects would result in more favorable conditions for further speciation involving fissions. The chain of derivation may then proceed at an accelerating rate until all metacentric chromosomes in the karyotype were fissioned. At this point, further chromosomal speciation involving fissions would be blocked, and the genetic system might not be conditioned to favor speciation involving other kinds of chromosomal mutations.

2) One variable which has a directly proportional effect on the probability that chromosomal speciation will occur is the rate of chromosome mutation for a given kind of rearrangement. This rate could easily be raised by the positive feedback process to such a high level that many

individuals in a population would suffer from the negative heterotic effects of chromosomal heterozygosity. If this happened, individual selection would favor evolving modifications in the mechanics of meiotic assortment to prevent malassortment in the large number of heterozygotes carrying new mutations. Chromosomal heterozygosity would no longer show negative heterosis and could no longer produce the hybrid sink required for successful speciation. Consequently, the species forced into this adaptation could form no more species chromosomally, but would be left with a high rate of chromosome mutation, and should be polymorphic for the kinds of rearrangements involved in the chain of derivation in its ancestry.

3) The third type of termination involves extrinsic circumstances. If the chain speciation process works as I have suggested, a fairly large number of species may form in a geologically short time under circumstances where they are not geographically isolated from one another. If species need an ecological niche of some minimum width, and if their basic adaptations allow them to use only a certain subdivision of the environment, then as the lineage proliferates new species, they would be limited to progressively more restricted niches or geographic areas. Eventually a point would be reached where further speciation either would be impossible or could not occur without extinction of other forms. If further speciation was blocked for long enough, individual selection would eventually force genetic system parameters to revert to their "unamplified" original conditions, which would not favor chromosomal speciation even with new ecological opportunities.

Group selection in this model is associated with the positive feedback mechanism, but its effects show primarily in the ecological consequences of chromosomal speciation. The major "selective" force involved is the surface tension produced by the hybrid zone sur-

rounding newly differentiated founder populations. According to the competitive exclusion principle, two species cannot coexist in the same ecological niche. If a nascent species is formed within the range of its ancestral stock, the ancestral and derived forms will inevitably compete ecologically. Three outcomes are possible: 1) One of the species becomes extinct—most probably this will be the nascent species. 2) The two species remain in similar niches but displace one another geographically along some environmental gradient (i.e., geographical exclusion). 3) The two species coexist geographically but displace one another along one or more resource axis of the environment (i.e., character displacement). As mentioned above in the discussion of the random fixation model, in character displacement the ancestral population will almost always displace the derived population, thereby forcing it into a new and probably more specialized niche. However, in geographic exclusion, exclusion of the initially very small founder population will be equivalent to extinction.

When a lineage invades a new environment or adaptive range, and its species have few competitors, character displacement is probably easy (e.g., Williams, 1972). The ready availability of vacant sympatric niches and the initial selection against hybridization in the random fixation model may allow the rapid evolution of premating isolation. One can also imagine that establishment of a hybrid sink situation along some resource axis of the environment would prevent gene flow along the axis so that disruptive selection would favor the rapid evolution of sympatric species. However, once an environment becomes fairly saturated with related species (as in the current situation for the *Sceloporus grammicus* complex), most chromosomal speciation probably involves a long period of geographic exclusion. The consequences logically required by the group selection model are interesting.

Because of the surface tension of its hybrid sink, which occurs independently of any environmental circumstances, the nascent chromosomally differentiated species will be at an immediate disadvantage. If the risk of hybridization cannot be reduced quickly by character displacement and expansion into a sympatric niche (or an uncontested geographic range) the founder population will probably be consumed by its own hybrid sink. Where the founder population cannot escape hybridization its survival is likely only if the parental stock can be pushed back to increase the hybrid zone's radius of curvature enough to reduce the surface tension. Two kinds of chance circumstances allow the founder population to achieve this initial expansion. The first depends entirely on extrinsic events, such as a catastrophe, to eliminate an adjacent ancestral population so the nascent species can spread into the depopulated area. The other circumstance depends on intrinsic aspects of the genetic system.

The founder population required for the chance fixation of a negatively heterotic chromosomal rearrangement is ideal for other stochastic phenomena such as the random fixation or drift of alleles at a variety of polymorphic gene loci. Also, founder populations will be exposed to specific local circumstances which may result in selection pressures that differ considerably from the average selective environment of the ancestral species. The ancestral species must maintain an adaptation to some average habitat occupied by populations exchanging genes with one another, while the initial barrier provided by chromosomal differentiation allows the founder population to adapt to its own specific local habitat. Thus, although most incipient chromosomally differentiated species will be quickly consumed by their own hybrid sinks, one may rarely achieve a fortuitous genotype which is sufficiently superior in its local environment so that it can feed enough migrants into the hybrid

sink to counteract its surface tension. If the founder population once manages to push the sink back to a greater radius of curvature, the pressure will be reduced and the founder population's superior fitness may then allow it to displace the ancestral stock from a wide geographic range. Chromosomal differentiation events may become frequent towards the end of a chain of derivation, but the environment will probably be too saturated to allow easy character displacement. The vast majority of these incipient species will be immediately consumed by their hybrid zones. Barring the lucky environmental circumstances, the *only* populations to survive will be those which achieve an especially superior adaption which allows them to overwhelm the surface tension of their hybrid zone.

Note that the positive feedback in chain speciation will also work here. Each founder population must be fit enough to be able to displace its immediate ancestor against the added impediment or pressure of the hybrid sink's surface tension. Thus, species at the ends of chains of geographic displacement should be especially effective competitors against species formed early in the chain. Chromosomally conservative species should survive competition with their derivatives, because they presumably held extensive geographic ranges before chromosomal speciation began. However, intermediate species in a chain may easily be excluded into extinction by their competitively superior derivatives, because they may have little chance to expand geographically before producing further derivatives.

In sum, the discovery program of the comparative approach has identified in the sceloporine radiation, independently of models, a variety of circumstances of cytogenetics, ecology, and phylogenetic relationships which appear to correlate with prolific speciation in *Sceloporus*. These are not adequately explained by the allopatric speciation of Mayr (1963,

etc.). The "guessing" to explain the totality of these relationships has been very straightforward and has led in easy steps to a three-part model which fully explains all of the phenomena revealed so far by the discovery program. By contrast, none of the guesses made in advance of collecting these data make predictions which approach the richness or realism of the circumstances logically required by the complete model and reasonable initial conditions. The complete model is, of course, fully corroborated by the information content of the conditions it has been developed to explain. The question remains—can the realism of these explanations be tested further? The answer provided by the next section is yes.

TESTING EXPLANATORY MODELS

General principles. The logic of testing the realism of explanatory models for historical processes is not difficult. However, it is my impression that many workers who fail to understand how the models are generated, fail also to understand all of the tests which can be used to further corroborate them (e.g., Peters, 1976). Under any circumstances it is logically impossible to prove any universal explanation to be absolutely true; although if uniformitarianism is accepted, the explanation can be demonstrated to be realistic to some degree of statistical confidence. If the world is fundamentally nondeterministic, falsity cannot be absolutely proven either. However, the idea of empirically testing and criticizing the logically derived statements a theory makes about nature still must form the basis that makes science different from fantasy.

If the program of discovery described above is followed it will lead to generalizations which already contain a large quantity of empirically realistic (or pre-tested) information. However, the pro-

gram of science is always to extend the frontiers of knowledge, and Popper's evolutionary theory of knowledge suggests how this should be done, except rather than beginning by testing wild guesses, the program begins with testing generalizations which already include a large body of pretested and empirically realistic information. Following Popper the content of realistic information entailed by the explanation can be increased by attempting to refute it.

There are at least three ways to refute an explanation (most workers think only of testing "predictions", and assume that an explanation which cannot be attacked in this way must be metaphysical, e.g., Peters, 1976): 1) An explanation may be refuted if its predictions are consistently unrealistic. 2) If an explanation depends logically on certain initial conditions or assumptions and it can be shown that these are unrealistic, then the explanation is refuted (Ghiselin, 1969). 3) Also, if it can be shown that supposed final conditions actually do not follow logically from supposed initial conditions, the explanation is obviously faulty. Full corroboration of an explanation should require that it survive all of these attempts to refute it. Each of these tests will be explained in more detail and illustrated with examples from my research program.

STEP 6: TEST LOGIC

Principle. In principle, testing the logic of an explanation requires no additional data gathering or processing. This should be the first and most obvious type of test to apply, but judging by many of the published attempts to explain non-allopatric speciation, it is frequently overlooked. The critical question to be asked in this kind of test is, do the supposed consequences the model proposes to explain actually follow logically from its supposed initial conditions. It should

be possible to reduce deterministic models to mathematical or logical symbols to see if the consequences do follow. For stochastic models, some of which can become quite complex, it should be possible to test their logic by computer simulation to see if the supposed consequences follow from the initial conditions and the explanation.

Application. Several aspects of the chromosomal speciation model are susceptible to testing via computer simulation. The most critical points to attack are:

1) Fixation of negatively heterotic mutations in small populations: the probability of fixing a negatively heterotic mutation should be calculated or determined by simulations for a variety of population sizes, migration rates, and relative fitnesses of the three possible genotypes (two homozygous conditions versus heterozygotes).

2) The hybrid sink effect: for various conditions of population structure and density, vagility, and degrees of negative heterosis, how effective is the hybrid sink as a block to the penetration of unlinked or linked alleles? What relative fitnesses of the two populations are required to maintain a curved hybrid sink in equilibrium and how do these vary as a function of the radius of curvature of the center of the sink? What conditions would be required to form a hybrid sink along an environmental resource axis rather than a geographical axis?

3) Chain speciation and positive feedback amplification: how reasonable is the group selection argument? For a plausible random fixation model, what mutation rate would be required to generate enough incipient species to provide a plausible frequency of species able to displace their ancestors against the initially high surface tension pressure? What changes in the frequency of alleles in controlling genetic system parameters are plausible in chromosomal speciation and how rapidly could selection at the

individual level change these frequencies? Many other aspects could also be modeled, but the tests listed here would seem to have the greatest power to corroborate the proposed explanations.

STEP 7: TEST ASSUMPTIONS

Principle. Any demonstration that a necessary *a priori* assumption of the explanation is unrealistic, immediately makes the model unrealistic to the extent that it depends on the assumption, and to the extent that the assumption is shown to be realistic. There are many ways to provide such demonstrations. Any model which assumes a "fact" of nature is falsified by demonstrating that the assumed "fact" is in reality not true.

Applications. The most critical assumption in the chromosomal speciation model is that the chromosomal rearrangements fixed between species were negatively heterotic when they originally occurred as new mutations. Presumably, where they are involved in narrow hybrid zones which appear to function as sinks, such as found in the *Sceloporus grammicus* complex, they should still be negatively heterotic. Thus, examining meiotic assortment in heterozygotes from these hybrid zones should test this important assumption. Such tests have already been done for a similar situation of rapid chromosomal differentiation in European mice of the genus *Mus* (Cattanach and Moseley, 1973), where single metacentric chromosomes have been backcrossed for the test into essentially pure *Mus musculus* genotypes to isolate them from possible genic problems in the hybrids. It should also be possible to do the same with *Sceloporus grammicus*. Another critical assumption concerns population subdivision. In sceloporines the most recent chromosomal speciation appears to have been in the *Sceloporus grammicus* complex. Superficially it appears to have an ideally

subdivided population. However, the assumption can be tested further by making detailed observations on population structure, mating system and vagility by using mark-observation and recapture methods.

STEPS 8 AND 9: TESTING PREDICTIONS

Principles. If an explanation requires potentially observable circumstances (conditions which should be causally related to P) which have not been examined, then attempts to demonstrate these circumstances will test the explanation. For stochastic models, confirmation or disconfirmation of these predictions will support or reject the explanation with some degree of statistical confidence. Such tests can take at least two qualitatively different forms: testing predictions in a relatively strict meaning of the term, and testing the ability of the model to reconstruct case histories realistically.

Testable predictions can be subdivided into three categories: 1) The model may predict already observed classes of phenomena in new cases (i.e., the natural experiments should be repeatable). 2) The model may predict new classes of phenomena in cases already observed. 3) The model may predict new classes of phenomena in new cases. These are listed here in the order of the epistemic values of the tests from least to most (cf. Popper, 1972a, b, c). However, it should be recognized that an efficient discovery program may already have found all of the classes of phenomena causally related to P. If the explanation covers these, the failure of the model to predict "new classes of testable phenomena" does not necessarily suggest that the model is unscientific as Popper might claim. If both the logic and assumptions of the model have been tested, the failure to predict new phenomena suggests that the discovery program has already provided the model

with nearly the maximum content of information available in the domain it covers. That is, the model has been proven to be realistic within the limitations of the real world. This would correspond to Kuhn's (1962, 1970a, b) "normal science".

Historical explanations can be used to make "predictions" of still another kind. Models will usually be expressed in the form of generalizations about assumed initial conditions and the *kinds* of effects expected to follow from them. Aside from predicting general classes of phenomena, as discussed above, the explanations should also allow the past histories of specific cases to be reconstructed in detail. If evidence not already included in the model by the discovery program allows an independent reconstruction of the past histories of cases, then the realism of the model is tested to the degree that the independent reconstructions are realistic and coincide with the reconstructions provided by the model.

Applications. Except for requiring certain initial conditions which can be tested empirically, the chromosomal speciation model predicts no phenomena not already suggested by the discovery program. However, these "predictions" can be tested over a wider range of sceloporine species than have been examined to date. Also, the predicted patterns of chromosomal variability and phylogenetic relationships should be found in other speciose genera of the Iguanidae (i.e., in *Anolis* and *Liolaemus*) and they should *not* occur in other small genera (Paull *et al.*, 1976 have already begun this test). Also, can these patterns be found in other lizard families, other classes of vertebrates, and other phyla? The most critical tests available are those of the accuracy of historical explanations. The chromosomal speciation model greatly constrains possible phylogenies which can be reconstructed from karyotypic evidence. Do these karyotypic phylogenies correspond to phylogenies

which can be reconstructed from evidence provided by independent character sets?

OTHER CONSIDERATIONS

Corrective feedback and the growth of knowledge. Tests provide new empirical observations which tend to falsify, support, or suggest extending the tested model. Whatever the "favorable" or "unfavorable" import of the observations for the model being considered, these new data increase the information content of the domain of the explanation which includes the observations by giving a more complete view of the world than existed previously (Popper, 1970, 1972c). A methodology which does not take advantage of this input is less than fully effective. Where some assumption of a model has been proven false, the obvious response is to find another explanation for the available data which does not depend on this assumption. If this cannot be done, perhaps the problem is poorly defined and the new data may suggest a clearer formulation of the research question, further data collecting, and/or looking at the available case histories in new combinations. Predictive or reconstructive tests automatically provide new data which can be entered in the cross correlation analyses to improve their inductive precision, and the evidence provided by the new data should clearly indicate the strengths and weaknesses of the tested model. Assuming that a model has at least some contact with reality, this feedback should lead through successive approximations of data gathering, testing, and revisions to ever more realistic understandings of the fundamental laws governing the processes modeled (Popper, 1972).¹

Semantics and syntax. The final epistemological difficulties of a methodology for the comparative study of evolutionary problems concern the precision of its tools. Since the comparative approach seeks to understand processes which cannot be manipulated readily in the laboratory, information and speculations concerning them must be abstracted through several levels of language before they can be manipulated logically in building and testing explanatory models. The logical sequences of abstracting inductively from reality to formulating a model, and reflecting deductively back to reality for testing the model, both depend on the precision of the symbols used for these manipulations. The conclusion of any logical argument of this nature can be no stronger than the weakest or foggiest definitions used in the chain of reasoning. (Note that this is a very different problem from determining the "essential" meaning of a word; cf. Popper, 1976.) Many proposed explanations of non-allopatric speciation are unrealistic because of logical problems introduced by using common words with semantically or syntactically faulty definitions. These faults are easily made by using as descriptors of nature (i.e., "facts") terms which were originally defined on the basis of some unsubstantiated model of nature. As Popper (e.g., 1970, 1976) and others (e.g., Kuhn, 1962; and others in Suppe [ed.], 1977) have shown, all words are theory laden, so the problem is unavoidable to some degree; but the utmost should be done to minimize it, and it is certainly unwise for an investigator to forget that it exists. Another reason for the confusion of proposals cited in my introduction results from inconsistent use of terms in discussing speciation phenomena between the different

¹See Lakatos and Musgrave [eds.] (1970) for a debate between Popper, Kuhn, and their various followers and critics about their respective ideas concerning how scientific "knowledge" grows and

can be defined through its contacts with empirical reality.

"scientific communities" (Kuhn, 1970: 176) concerned with the problems of species and speciation.

Thus, although many before me have discussed these definitions, I think it essential for the following papers in this series to redefine much of the vocabulary concerned with geographic and genetic relationships of populations and species, and with isolating mechanisms and their failures. A glossary of these definitions will be provided later in this series.

CONCLUSIONS

Solving the problem of speciation. To understand realistically how new species form is probably the most difficult task evolutionists have attempted to achieve. Besides confronting the unresolved problem of what species are, these attempts confront the fundamental epistemological problem of determining how realistic the explanations are, when speciation cannot be directly manipulated or observed. I propose that the comparative methodology described above answers the epistemological problem and provides an efficacious approach towards explaining how species are formed, and thus also, what they are. Many evolutionists claim to use comparative methodologies but few appear to understand clearly the epistemic power of the approach; so far as I know, none have adequately formulated or justified the logical methodology. From this, and my own experience with the approach, I conclude that it is sufficiently novel to be explained in detail and justified philosophically. Therefore, in the present paper I have presented and justified epistemologically the logical schemata I have followed in my own attempts to solve the problem of how species are formed. These schemata offer seemingly novel tools, both for discovering causally related effects of historical processes hidden among the noise of unrelated events, and for empirically testing the

realism of any models proposed to explain the process and its causally related effects. In justifying these schemata I confront and tentatively answer two major problems in the philosophical foundations of scientific methodology: 1) Is there an inductive logic which can be used to infer realistic generalizations about nature from specific observations of nature, and if so what is the epistemic value of this logic? 2) How can one test the empirical realism of generalizations about historical processes, when the processes cannot be observed or manipulated experimentally? The answers to these questions provide a methodology which appears to be effective for understanding speciation, and a surprisingly wide diversity of other historical problems ranging from cosmology through economics and possibly even to fundamental particle physics. The approaches to answering these questions also seem to indicate the way to solutions of some of the foundation problems of the philosophy of science.

The inductive power of the comparative approach is most effective if it can be applied to a compact phylogenetic radiation which can be subdivided into a series of genetically and ecologically related cases, some of which serve as "natural experiments" to demonstrate the phenomenon to be explained, and others which serve as "controls" because they are similar to the "experiments" in most respects, but do not show the experimental phenomenon. The 120+ species of the sceloporine radiation of the lizard family Iguanidae provide an ideal radiation for this comparative approach. The radiation involves at least two qualitatively different modes of speciation which may be compared and contrasted by the comparative methods to help isolate specific features that help to explain the differences. I abstract my research program on the comparative cytogenetics, speciation, and evolution of the sceloporine lizards to illustrate how the comparative methodology has actually

been used in my own research program. Thus, the present paper serves to introduce the series of papers to follow, which will present in detail the results of the research program on sceloporine speciation. It also serves as a methodology section for this program.

Modes of speciation: where we stand. To illustrate and recapitulate the epistemological problems the present paper faces and attempts to solve, I offer the following quotations as a necessary background to the attempt to understand sceloporine speciation:

It is rather discouraging to read this perennial controversy because the same old arguments are cited again and again in favor of sympatric speciation In the last analysis, all the various schemes make arbitrary postulates that at once endow the speciating individuals with the attributes of a full species. They attempt thus to bypass the real problem of speciation. One would think that it should no longer be necessary to devote so much time to this topic, but past experience permits one to predict that the issue will be raised again at regular intervals. Sympatric speciation is like the Lernaean Hydra which grew two new heads whenever one of its old heads was cut off. There is only one way in which final agreement can be reached and that is to clarify the whole relevant complex of questions to such an extent that disagreement is no longer possible. (Mayr, 1963: 451)

False facts are highly injurious to the progress of science, for they often endure long; but false views, if supported by some evidence, do little harm, for everyone takes a salutary pleasure in proving their falseness; and when this is done, one path toward error is closed and the road to truth is often at the same time opened. (Darwin, 1889: 606, as quoted in Ghiselin, 1969—emphasizing the importance of distinguishing syntactically between observations and explanations)

For evolutionary biology . . . history . . . provides the needed key to scientific knowledge. Our approach to evolutionary and population problems should involve historical reconstruction as well as more traditional comparative and experimental techniques. We must account for what actually has occurred: what might take place under *ideal* conditions will never do. *Otherwise we shall justify our conceptions of things from our ideas, rather than letting the way things are determine how we shall conceive of them.* [Italics mine] (Ghiselin, 1974: 27, stressing the impor-

tance of not begging the conclusion by working logically from unrealistic initial assumptions [i.e., to avoid the error of *petitio principii*])

Two alternatives seem open to us at this juncture: either revert to a hypothetic-inductive model of science or argue that evolutionary theory after a century is still inadequately formulated and that in a more finished form will conform to the H-D model. The problem with the first alternative is that there is no H-I model. Thus, this alternative reduces to the admission that there is no reconstruction of science appropriate to evolutionary theory as it now stands. Biologists are currently working on the second alternative. Some are attempting to reformulate macro-evolutionary theory more rigorously so that deductive confirmation or disconfirmation is possible. Others find evolutionary theory in terms of organisms and their interactions too crude to permit an adequate formulation of evolutionary theory. Instead, they call for a molecular version of evolutionary theory, hoping in this manner to fulfill the requirements of the H-D model. . . . (Hull, 1973: 34–35)

I would claim that the comparative approach outlined in this paper is as logically rigorous both for making inductive generalizations and for deductively testing them as is possible, given a fundamentally stochastic reality. It remains to demonstrate the utility of the methodology, as will be attempted in the following papers.

ACKNOWLEDGMENTS

The heuristic schemata trace from diverse exemplars of methodology, beginning with my father's geology background. My formal training started with physics—to statistical mechanics and atomic physics. I changed to biology at UCLA in 1960 with Raymond Cowles's natural history, which taught "population thinking" and anti-essentialism, and with William Beckwith's comparative ethology, which taught the "natural experiment." G. A. Misrahy (Sensory and Developmental Physiology Lab, Children's Hospital, Los Angeles—NIH Training Grant 2B-5264) taught "black-box" systems analysis, controlling ex-

periments, and how to use signal averaging computers.

The sceloporine study owes its genesis to Don Hunsaker, II and Richard Etheridge, who taught me at San Diego State University from 1961–1965. (Etheridge also demonstrated the comparative approach in his teaching.) Discussions and/or manuscript reviews by Don Gartside, George Gorman, Max Hecht, Bernard John, Max King, Ernst Mayr, Jeffry Mitton, Hobart Smith, Jacek Szymura, Henryk Szarski, Bruce Wallace, Michael White, Max Whitten and many students, among others, helped greatly. Special thanks go to Ralph Axtell and Ernest E. Williams, at Southern Illinois University-Edwardsville and Harvard, where I did most of the field work, and to Scott Moody, who identified communication problems in my 1977 paper. Roslyn Dunn (now my wife) typed the manuscript and eliminated some of my worst prose. Hall (1977) was completed at the University of Colorado, Boulder, and the University of Melbourne Research Fellowship supported the present work. Field work was supported by grants from the Society of the Sigma Xi (1966 and 1968), Evolutionary Biology Committee Training Grants (Harvard, 1968 and 1969), and National Geographic Society Grants 864 and 972 (1970–1971).

LITERATURE CITED

- ACHINSTEIN, P. 1971. Law and Explanation: An Essay in the Philosophy of Science. London, Oxford University Press, 168 pp.
- AXELROD, D. I. 1950. Evolution of desert vegetation in western North America, pp. 215–306. In D. I. Axelrod (ed.), *Studies in Late Tertiary Paleobotany*. Contrib. Paleont., Carnegie Inst. Washington, Publ. No. 590.
- . 1958. Evolution of the Madro-Tertiary Geoflora. *Bot. Rev.*, **24**: 433–509.
- AXTELL, R. W. 1958. A monographic revision of the iguanid genus *Holbrookia*. Ph.D. Thesis, University of Texas, Austin, 222 pp.
- BERRY, B. J. L. 1964. Approaches to regional analysis: A synthesis. *Ann. Assoc. Amer. Geogr.*, **54**: 2–11.
- BUSH, G. L. 1969. Sympatric host race formation and speciation in frugivorous flies of the genus *Rhagoletis* (Diptera, Tephritidae). *Evolution*, **23**: 237–251.
- . 1974. The mechanism of sympatric host race formation in the true fruit flies (Tephritidae), pp. 3–23. In M. J. D. White (ed.), *Genetic Mechanisms of Speciation in Insects*. Sydney, Australia and New Zealand Book Co.
- . 1975a. Sympatric speciation in phytophagous parasitic insects, pp. 187–206. In P. W. Price (ed.), *Evolutionary Strategies of Parasitic Insects*. London, Plenum.
- . 1975b. Modes of animal speciation. *Ann. Rev. Ecol. Syst.*, **6**: 339–364.
- BUSH, G. L., S. M. CASE, A. C. WILSON AND J. L. PATTON. 1977. Rapid speciation and chromosomal evolution in mammals. *Proc. Nat. Acad. Sci. USA* **74**: 3942–3946.
- CAIN, A. J. 1953. Geography, ecology and coexistence in relation to the biological definition of the species. *Evolution*, **7**: 76–83.
- CATTANACH, B. M., AND H. MOSELEY. 1973. Non-disjunction and reduced fertility caused by the tobacco mouse metacentric chromosomes. *Cytogenet. Cell Genet.*, **12**: 264–287.
- CAVAZOS, L. F. 1951. Spermatogenesis of the horned lizard *Phrynosoma cornutum*. *Amer. Nat.*, **75**: 373–379.
- CHORLEY, R. J., AND P. HAGGETT (EDS.). 1967. *Models in Geography*. London, Methuen & Co., 816 pp.
- COLE, C. J. 1975. Evolution of parthenogenetic species of reptiles, pp. 340–355. In R. Reinboth (ed.), *Intersexuality in the Animal Kingdom*. New York, Springer-Verlag.
- DARWIN, C. 1889. *The Descent of Man, and Selection in Relation to Sex*, 2nd. Ed. New York, D. Appleton.
- DOBZHANSKY, T. 1937. *Genetics and the Origin of Species* [1st. Ed.]. New York, Columbia University Press.
- . 1970. *Genetics of the Evolutionary Process*. New York, Columbia University Press, 505 pp.
- EHRRLICH, P. R., AND P. H. RAVEN. 1969. Differentiation of populations. *Science*, **165**: 1228–1232.
- EISBERG, R., AND R. RESNICK. 1974. *Quantum Physics, Atoms, Molecules, Solids, Nuclei, and Particles*. New York, John Wiley & Sons, 713 pp. + Appendices and Index.
- ENDLER, J. A. 1973. Gene flow and population differentiation. *Science*, **179**: 243–250.
- . 1977. *Geographic Variation, Speciation, and Clines*. Monogr. Pop. Biol. no. 10, Princeton, Princeton University Press, 246 pp.
- ETHERIDGE, R. 1964. The skeletal morphology and systematic relationships of sceloporine lizards. *Copeia*, **1964**: 610–631.

- GHISELIN, M. T. 1969. *The Triumph of the Darwinian Method*. Berkeley, Univ. of California Press, 287 pp.
- . 1974. *The Economy of Nature and the Evolution of Sex*. Berkeley, Univ. of California Press, 346 pp.
- GLASER, E. M., AND D. S. RUCHKIN. 1976. *Principles of Neurobiological Signal Analysis*. New York, Academic Press, 471 pp.
- GOLDSCHMIDT, R. 1955. *Theoretical Genetics*. Berkeley, Univ. of California Press.
- HAGGETT, P., AND R. J. CHORLEY. 1967. Models, paradigms and the new geography, pp. 19–41. In Chorley and Haggett (eds.).
- HALL, W. P. 1964. Cytogenetic studies in the family Iguanidae. Unpublished manuscript. San Diego State University Dept. Zoology, San Diego.
- . 1973. Comparative population cytogenetics, speciation, and evolution of the iguanid lizard genus *Sceloporus*. Ph.D. Thesis, Harvard Univ., Cambridge, Massachusetts, 193 pp. + addenda.
- . 1977. Cascading chromosomal speciation and the paradoxical role of contact hybridization as a barrier to gene flow. Informally Published [see footnote 1, p. xxx, this paper], 91 pp.
- . . Modes of speciation and evolution in the sceloporine iguanid lizards. II. An overview of chromosomal variation and phylogeny. In Preparation.
- HALL, W. P., AND R. K. SELANDER. 1973. Hybridization of karyotypically differentiated populations in the *Sceloporus grammicus* complex (Iguanidae). *Evolution*, 27: 226–242.
- HANSON, N. R. 1958. *Patterns of Discovery, An Inquiry into the Conceptual Foundations of Science*. London, Cambridge University Press, 241 pp.
- . 1961. Is there a logic of discovery?, pp. 20–42. In H. Feigl and G. Maxwell (eds.), *Current Issues in the Philosophy of Science*. New York, Holt, Reinhart and Winston.
- HARVEY, D. 1969. *Explanation in Geography*. London, Edward Arnold, 521 pp.
- HEMPEL, C. G. 1965. *Aspects of Scientific Discovery and Other Essays in the Philosophy of Science*. New York, The Free Press, 504 pp.
- HULL, D. L. 1973. Introduction, pp. 1–77. In D. L. Hull (ed.), *Darwin and His Critics, The Reception of Darwin's Theory of Evolution by the Scientific Community*. Cambridge, Harvard Univ. Press.
- . 1976. Are species really individuals? *Syst. Zool.*, 25: 174–191.
- HUNT, W. G., AND R. K. SELANDER. 1973. Biochemical genetics of hybridization in European house mice. *Heredity*, 31: 11–33.
- HUXLEY, J. S., (ED.). 1940. *The New Systematics*. Oxford, Clarendon Press.
- . 1942. *Evolution, the Modern Synthesis*. London, Allen and Unwin.
- JOHN, B., AND K. R. LEWIS. 1965. Genetic speciation in the grasshopper *Eyprepocnemis plorans*. *Chromosoma*, 16: 308–344.
- . 1966. Chromosome variability and geographic distribution in insects. *Science*, 152: 711–721.
- KEY, K. H. L. 1968. The concept of stasipatric speciation. *Syst. Zool.*, 17: 14–22.
- . 1974. Speciation in the Australian morabine grasshoppers—taxonomy and ecology, pp. 43–56. In M. J. D. White (ed.), *Genetic Mechanisms of Speciation in Insects*. Sydney, Australia & New Zealand Book Co.
- KING, L. J. 1969. *Statistical Analysis in Geography*. Englewood Cliffs, Prentice-Hall, 288 pp.
- KITTS, D. B. 1977. Karl Popper, verifiability, and systematic zoology. *Syst. Zool.*, 26: 185–194.
- KOESTLER, A. 1967. *The Ghost in the Machine*. London, Pan Books, Ltd., 384 pp.
- KUHN, T. S. 1962. *The Structure of Scientific Revolutions*. Chicago, University of Chicago Press, 173 pp.
- . [1965.] Logic of discovery or psychology of research, pp. 1–23. In Lakatos and Musgrave (eds.), 1970.
- . [1969.] Second thoughts on paradigms [Session VI of Symposium, March 26 to 29, 1969], pp. 459–519 [incl. commentary]. In F. Suppe (ed.), 1977.
- . 1970a. Postscript, 1969, pp. 174–210. In T. S. Kuhn. *The Structure of Scientific Revolutions*, 2nd Ed., Enlarged. Chicago, University of Chicago Press.
- . 1970b. Reflections on my critics, pp. 231–278. In Lakatos and Musgrave (eds.), 1970.
- LAKATOS, I., AND A. MUSGRAVE (EDS.). 1970. *Criticism and the Growth of Knowledge*. Proceedings of the International Colloquium in the Philosophy of Science, London, 1965, Vol. 4. London, Cambridge Univ. Press, 282 pp.
- LEWIS, H. 1966. Speciation in flowering plants. *Science*, 152: 167–172.
- . 1973. The origin of diploid neospecies in *Clarkia*. *Am. Nat.*, 107: 161–170.
- LEWIS, H., AND P. H. RAVEN. 1958. Rapid evolution in *Clarkia*. *Evolution*, 12: 319–336.
- LOVETRUP, S. 1975. Letter. *Syst. Zool.*, 24: 507–511.
- MASTERMAN, M. 1970. The nature of a paradigm, pp. 59–89. In Lakatos and Musgrave (eds.), 1970.
- MATTHEY, R. 1931. Chromosomes de reptiles—sauriens, ophiidiens, cheloniens. L'évolution de la formule chromosomiale chez les sauriens. *Rev. Suisse Zool.*, 38: 117–186 + 8 pl.
- . 1964. La signification des mutations chromosomiques dans les processus de speciation: Etude cytologique du sous-genre *Leggada* Gray (Mammalia—Muridae). *Arch. Biol.*, 75: 169–206.
- . 1966. Genetic mechanisms and speciations of

- mammals, pp. 1-11. In G. Gergamian (ed.), *The Chromosomes, Structural and Functional Aspects*. In Vitro, Publ. Tissue Culture Association.
- MAYR, E. 1942. *Systematics and the Origin of Species*. New York, Columbia University Press, 334 pp.
- . 1947. Ecological factors in speciation. *Evolution*, 1: 263-288.
- . 1954. Change of genetic environment and evolution, pp. 157-180. In A. C. Hardy and E. B. Ford (eds.), *Evolution as a Process*. London, Allen and Unwin.
- . 1963. *Animal Species and Evolution*. Cambridge, Harvard Belknap Press, 797 pp.
- . 1970. *Populations, Species and Evolution*. Cambridge, Harvard Belknap Press, 453 pp.
- MONOD, J. 1971. *Chance and Necessity: An Essay on the Natural Philosophy of Modern Biology* [Trans. from the French by A. Wainhouse]. New York, Vintage Books, 199 pp.
- MORAFKA, D. J. 1977. A Biogeographic Analysis of the Chihuahuan Desert through its Herpetofauna. *Biogeographica* Vol. 9. The Hague, Dr. W. Junk B. V., Publ., 313 pp.
- MOROWITZ, H. J. 1968. *Energy Flow in Biology: Biological Organization as a Problem in Thermal Physics*. New York, Academic Press, 179 pp.
- MURRAY, J. 1972. *Genetic Diversity and Natural Selection*. Edinburgh, Oliver and Boyd, 128 pp.
- NEVO, E., AND H. BAR-EL. 1976. Hybridization and speciation in fossorial mole rats. *Evolution*, 30: 831-840.
- NEVO, E. Y., J. KIM, C. R. SHAW, AND C. S. THAELE JR. 1974. Genetic variation, selection and speciation in *Thomomys talpoides* pocket gophers. *Evolution*, 28: 1-23.
- NICKLES, T. [1973.] Heuristics and justification in scientific research: Comments on Shapere, pp. 571-589. In Suppe (ed.), 1977.
- NORRIS, K. S. 1958. The evolution and systematics of the iguanid genus *Uma* and its relation to the evolution of the other North American desert reptiles. *Bull. Amer. Mus. Nat. Hist.*, 114: 247-326.
- PAINTER, T. S. 1921. Studies in reptilian spermatogenesis. I. The spermatogenesis of lizards. *J. Exptl. Zool.*, 34: 281-327.
- PARKER, E. D., JR., AND R. K. SELANDER. 1976. The organization of genetic diversity in the parthenogenetic lizard, *Cnemidophorus tesselatus*. *Genetics*, 84: 791-805.
- PATTON, J. L., R. K. SELANDER, AND M. H. SMITH. 1972. Genic variation in hybridizing populations of gophers (genus *Thomomys*). *Syst. Zool.*, 21: 263-270.
- PAULL, D., E. E. WILLIAMS, AND W. P. HALL. 1976. Lizard karyotypes from the Galapagos Islands: Chromosomes in phylogeny and evolution. *Breviora Mus. Comp. Zool. No.* 441, 31 pp.
- PETERS, R. H. 1976. Tautology in evolution and ecology. *Am. Nat.*, 110: 1-12.
- PLATNICK, N. I., AND E. S. GAFFNEY. 1978. Evolutionary biology: A Popperian perspective. *Syst. Zool.*, 27: 137-141.
- POPPE, K. R. 1934. *Logik der Forschung*. Vienna. Julius Springer Verlag.
- . 1970. Normal science and its dangers, pp. 51-58. In Lakatos and Musgrave (eds.), 1970.
- . 1972a. *The Logic of Scientific Discovery, Sixth Impression (Revised)* [First Impression, 1959]. London, Hutchinson & Co., Ltd., 480 pp.
- . 1972b. *Conjectures and Refutations: The Growth of Scientific Knowledge*, 4th Ed. (Revised) [1st Ed., 1963]. London, Routledge and Kegan Paul, 431 pp.
- . 1972c. *Objective Knowledge: An Evolutionary Approach*. London, Oxford Univ. Press, 380 pp.
- . 1976. *Unended Quest: An Intellectual Autobiography*, Revised Ed. [first publ. 1974]. La Salle, Illinois. Open Court Publ. Co., 255 pp.
- PRESCH, W. 1969. Evolutionary osteology and relationships of the horned lizard genus *Phrynosoma* (family Iguanidae). *Copeia*, 1969: 250-275.
- RIVAS, L. R. 1964. A reinterpretation of the concepts "sympatric" and "allopatric" with proposal of the additional terms "syntopic" and "allotopic." *Syst. Zool.*, 13: 42-43.
- SALMON, W. C. 1966, 1967. *The Foundations of Scientific Inference*. Pittsburg, Univ. of Pittsburg Press, 157 pp.
- SAVAGE, J. M. 1958. The iguanid lizard genera *Urosaurus* and *Uta* with remarks on related groups. *Zoologica, Sci. Contrib. New York Zool. Soc.*, 43: 41-54.
- . 1960. Evolution of a peninsular herpetofauna. *Syst. Zool.*, 9: 184-212.
- . 1966. The origins and history of the Central American herpetofauna, *Copeia*, 1966: 719-766.
- SIMPSON, G. G. 1944. *Tempo and Mode in Evolution*. New York, Columbia Univ. Press.
- . 1961. *Principles of Animal Taxonomy*. New York, Columbia Univ. Press, 247 pp.
- SMITH, H. M., AND E. H. TAYLOR. 1950. An annotated checklist and key to the reptiles of Mexico, exclusive of the snakes. *Bull. U.S. Nat. Mus.*, 199: 1-253.
- SMITH, S. G. 1966. Natural hybridization of the coccinellid genus *Chilocorus*. *Chromosoma*, 18: 380-406.
- SPURWAY, H. 1953. Genetics of specific and sub-specific differences in European newts. *Symp. Soc. Exptl. Biol.*, 7: 200-237.
- SPURWAY, H., AND H. G. CALLAN. 1960. The vigor and male sterility of hybrids between the species *Triturus vulgaris* and *T. helveticus*. *J. Genet.*, 57: 84-118.
- SUPPE, F. [1973.] *The search for philosophic under-*

- standing of scientific theories, pp. 1-241. In Suppe (ed.), 1977.
- . 1977. Afterword, pp. 615-730. In Suppe (ed.), 1977.
- . (ed.) 1977. *The Structure of Scientific Theories*, 2nd Ed. Urbana, Univ. of Illinois Press, 818 pp.
- SZARSKI, H. 1971. The importance of deviation amplifying circuits for the understanding of the course of evolution. *Acta Biotheor.*, **20**: 159-170.
- SZYMURA, J. M. 1976a. Hybridization between discoglossid toads *Bombina bombina* and *Bombina variegata* in southern Poland as revealed by the electrophoretic technique. *Z. f. Zool. Syst. Evol.-Forsch.*, **14**: 227-236.
- . 1976b. New data on the hybrid zone between *Bombina bombina* and *Bombina variegata* (Anura, Discoglossidae). *Bull. Acad. Polon. Sci. C1 2*, **34**: 355-363.
- THAELER, C. S., JR. 1974. Four contacts between ranges of different chromosome forms of the *Thomomys talpoides* complex (Rodentia: Geomyidae). *Syst. Zool.*, **23**: 343-354.
- TODD, N. B. 1970. Karyotypic fissioning and canid phylogeny. *J. Theor. Biol.*, **26**: 445-480.
- WALLACE, B. 1959. Influence of genetic systems on geographic distribution. Cold Spring Harbor Symp. Quant. Biol., **24**: 193-204.
- WHITE, M. J. D. 1973. *Animal Cytology and Evolution*, 3rd. Ed. London, Cambridge University Press., 961 pp.
- . 1978. *Modes of Speciation*. San Francisco, W. H. Freeman and Co., 455 pp.
- WHITE, M. J. D., R. E. BLACKITH, R. M. BLACKITH, AND J. CHENEY. 1967. Cytogenetics of the *viatica* group of morabine grasshoppers. I. The "coastal" species. *Aust. J. Zool.*, **15**: 263-302.
- WHITE, M. J. D., H. L. CARSON, AND J. CHENEY. 1964. Chromosomal races in the Australian grasshopper, *Moraba viatica* in a zone of geographic overlap. *Evolution*, **19**: 417-429.
- WHITE, M. J. D., K. H. L. KEY, M. ANDRE, AND J. CHENEY. 1969. Cytogenetics of the *viatica* group of morabine grasshoppers. II. Kangaroo Island populations. *Aust. J. Zool.*, **17**: 313-328.
- WILLIAMS, E. E. 1972. The origin of faunas. Evolution of lizard congeners in a complex island fauna: A trial analysis. *Evol. Biol.*, **6**: 47-89.
- WILSON, A. C., G. L. BUSH, S. M. CASE, AND M.-C. KING. 1975. Social structuring of mammalian populations and rate of chromosomal evolution. *Proc. Nat. Acad. Sci. USA*, **72**: 5061-5065.
- WRIGHT, S. 1941. On the probability of fixation of reciprocal translocations. *Am. Nat.*, **75**: 513-522.
- WOODRUFF, D. S. 1973. Natural hybridization and hybrid zones. *Syst. Zool.*, **22**: 213-218.
- ZASLEVSKY, V. A. 1963. Hybrid sterility as a limiting factor in the distribution of allopatric species. *Dokl. Akad. Nauk SSSR*, **149**: 470-471.
- ZEFF, E. W. 1962. A technique for delineation of chromosomal constitution of reptilian leucocytes grown in culture. Unpublished manuscript, UCLA Dept. Zool., Los Angeles.

Immigration and the Dynamics of Peripheral Populations

ROBERT D. HOLT¹

ABSTRACT. The influence of immigration upon the abundance and dynamics of peripheral populations should reflect the degree of genetic differentiation between residents and immigrants. The interplay of immigration, local dynamics, and natural selection in models of population growth is examined. Given that no genetic variation exists upon which local selection may act, and that a population is numerically stable, immigration should increase population size. In populations with discrete-time dynamics, immigration may either destroy or create local stability. If a population is unstable, an increase in the rate of immigration may reduce its average density. The introduction of haploid genetic variation dramatically changes the character of the relation between immigration rate and density. If fitnesses are density-dependent, in a stable, polymorphic population total density is independent of the rate of immigration. Decreasing the fitness of immigrants relative to residents may enhance the stabilizing influence of immigration. It is briefly argued that frequency-dependent fitnesses or diploid genetic variation with density-dependent fitnesses can produce an inverse relation between the rate of immigration and population size. These theoretical results suggest that the consequences of dispersal for population dynamics may be strongly influenced by the degree of local genetic adaptation that exists within a species.

INTRODUCTION

Patterns of dispersal are central to both island and continental biogeography, for the range ultimately occupied by a newly formed species depends upon that species' success as a colonizer (Williams, 1969). The importance of dispersal in

other phenomena of ecology and evolution is not as well understood and has been the subject of considerable debate. Over the past two decades, evolutionists and ecologists have moved in curiously opposite directions in their views on the significance of dispersal in natural populations. In 1963, Ernst Mayr in *Animal Species and Evolution* strongly argued that "gene flow is the main factor responsible for genetic cohesion among the populations of a species . . . [and] one of the principal reasons for the slow rate of evolution of common widespread species." This view of gene flow as a significant homogenizing force was accepted evolutionary doctrine for many years but has recently come under increasing attack. The empirical and theoretical studies of Antonovics (1968), Endler (1977), Levin and Kerster (1974) and Slatkin (1978) (to note just a few prominent landmarks) have demonstrated that gene flow does not as a general rule effectively counter spatially varying selection. Many evolutionary biologists are now persuaded that even though gene flow can provide an important source of variation upon which selection might act, in most natural populations rates of gene flow are usually not great enough to prevent the evolution of local adaptation.

Until the early 1970's theoretical ecology consisted largely of glosses on the elegant mathematical edifice erected by Vito Volterra and Alfred Lotka fifty years earlier. This body of theory, designed to analyze the dynamics of spatially homogeneous populations, neglect-

¹ Museum of Natural History and the Department of Systematics and Ecology, University of Kansas, Lawrence, Kansas 66045, U.S.A.

ted the possible significance of movement patterns in population processes. But in recent years the role of dispersal in local dynamics and interspecific interactions has become the focus of much research activity by theoretical ecologists (e.g., Levin, 1978; Okubo, 1980). Empirical evidence is steadily mounting that movement patterns affect both the mean density of populations and the pattern of fluctuations around that mean (Gaines and McClenaghan, 1980; Taylor and Taylor, 1977; Tamarin, 1977). We are left in the slightly awkward position of appearing to believe that dispersal is important in local ecological dynamics, but not in local evolutionary dynamics!

There is one circumstance for which all seem to agree that dispersal can be an important conservative evolutionary force. Peripheral populations in an ecotone often exist in low densities adjacent to populations at much higher densities. In such peripheral populations, natural selection favoring locally adapted alleles can be readily swamped by rates of immigration which are low in absolute numbers of individuals per unit time, but high in proportion to the peripheral population's low abundance (Antonovics, 1976). Mayr (1963) suggested that such regular disruption of local adaptations by immigration should be expected wherever a border population exists at low densities, and he argued that this process could produce evolutionary stasis at a species' border.

It is premature to judge the empirical adequacy of Mayr's model, as too little is known about the causes and dynamics of species' borders. However, the scenario is consistent with analytic studies of the countervailing influence exerted by gene flow upon local selection in low-density populations (e.g., Nagylaki, 1978). Hence, it is possible that gene flow neither accounts for the genetic cohesion of a species over its entire range, nor as a rule prevents local adaptation, yet does confer a kind of evolutionary stability upon that species' border.

In this paper I explore an ecological

question suggested by Mayr's evolutionary hypothesis. What is the relationship between the rate of immigration into a population and that population's size? I will argue that the answer to this question should depend upon the existence and character of genetic variation in the species. Given that residents and immigrants are genetically identical, it seems intuitively reasonable to expect that as the number of immigrants increases, so should population size. Conversely, if immigrants and residents differ genetically, the average immigrant should be less fit in the local environment than is the average resident. The more immigrants there are, the lower the mean fitness of the population should be. We may reasonably conjecture that a sufficient reduction in mean fitness will decrease the size of the population.

Using verbal arguments alone it is difficult to gauge the relative magnitude of these two opposing consequences of immigration. I have therefore explored the effect of immigration upon the dynamics of peripheral populations in three classes of population models. In order of increasing complexity, these are continuous-time and discrete-time ecological models: 1) without genetics, 2) with haploid genetic variation, and 3) with diploid genetic variation. This paper discusses the first two classes of models. My studies of the third class of models—by far the most complicated—will be presented elsewhere, along with certain technical details of the models described below.

PERIPHERAL POPULATION MODELS: GENETICALLY HOMOGENEOUS POPULATIONS

The two general growth models to be discussed are

$$\frac{dN}{dt} = F(N) + I \quad (1)$$

and $N(t + 1) = G(N(t)) + I$, or, equivalently,

$$\Delta N = G'(N(t)) + I \quad (2)$$

where N is population density, $F(N)$ and $G(N)$ are growth functions, $G'(N) \equiv G(N) - N$, and I denotes immigration. Throughout this paper the rate of immigration is assumed to be temporally invariant, and the growth functions F and G are assumed to be continuous. A continuous-time differential equation such as (1) is strictly appropriate as a growth model only for populations with completely overlapping generations, negligible time lags and no variation in survivorship or fecundity with age. The quantity I is an instantaneous rate of immigration. The discrete-time difference equation (2) best describes populations in which age-classes or growth stages are temporally segregated, so that only one class is present at any time. If dispersal occurs during a single stage in the life history of the organism, and the population is censused immediately following each pulse of dispersal and immigration, equation (2) is an appropriate model for studying the influence of immigration upon population dynamics. Both models assume that there are no qualitative differences, including genetic differences, between residents and immigrants.

The three standard steps in the analysis of growth models such as (1) and (2) are as follows: 1) solve for point equilibria, N^* ; 2) determine the local stability character of each equilibrium; 3) for model (2), attempt to understand the often complex dynamical behavior that may exist when populations do not have a stable point equilibrium. For any biologically reasonable model, in the absence of immigration there will be an equilibrium N^* of maximal size—the carrying capacity, K —above which the population has a negative growth rate. The number of equilibria found at densities below K is determined by the specific details of particular models.

IMMIGRATION AND POPULATION EQUILIBRIA

In model (1), equilibria are found by solving $F(N^*) + I = 0$. The local stability of each equilibrium is determined by the slope of the growth curve,

$$\lambda \equiv \frac{d}{dN} \left(\frac{dN}{dt} \right) = \frac{dF}{dN}$$

where the derivative is evaluated at $N = N^*$. If $\lambda < 0$, the equilibrium is locally stable, and if $\lambda > 0$ it is unstable. A population nudged away from an unstable equilibrium will converge asymptotically to a stable equilibrium. How does immigration shift the position of each equilibrium? By applying the chain rule of differentiation to (1) we have

$$\frac{dN^*}{dI} = -\left(\frac{dF}{dN}\right)^{-1} = -\lambda^{-1}, \quad \left(\frac{dF}{dN} = 0\right).$$

Hence, the slope λ of the growth curve at an equilibrium determines both the stability of the equilibrium population and the way in which immigration shifts its density. Immigration increases the density of populations at locally stable equilibria and decreases the densities at which unstable equilibria occur. The quantity $|\lambda|$ measures the strength of density-dependence in the population; increasing $|\lambda|$ reduces the sensitivity of population size to changes in immigration rate. Regardless of the detailed form of the growth model, however, if a peripheral population is at its carrying capacity, $\lambda < 0$ and immigration will always increase that population's size.

A similar relation between the rate of immigration and equilibrial density holds in the discrete-time growth model. We find the equilibria by solving $G'(N^*) + I = 0$. By linearizing around N^* in the usual way it can be shown that the local stability properties of each equilibrium are set by the sign and magnitude of

$$\lambda' = \frac{dG'}{dN},$$

where the derivative is evaluated at N^* . In general, three classes of equilibria are possible: 1) $\lambda' > 0$. A population deviating by a small amount from N^* will monotonically diverge away from N^* . 2) $-2 < \lambda' < 0$. The equilibrium is locally stable. 3) $\lambda' < -2$. The negative density-dependence at the equilibrium is so severe that the population repeatedly overshoots the equilibrium then crashes to low densities. The population ultimately settles into sustained oscillations around the locally unstable equilibrium; the character of these fluctuations depends upon the specific details of the full, nonlinear model. (May, 1976).

Applying the chain rule of differentiation again, we have

$$\frac{dN^*}{dI} = -(\frac{dG'}{dN})^{-1} = -(\lambda')^{-1}$$

where $\frac{dG'}{dN}$ is evaluated at N^* . Hence, if $\lambda' > 0$, immigration reduces the density at which equilibrium occurs, and if $\lambda' < 0$, population size increases with immigration.

A simple graphical illustration of this conclusion is shown in Figure 1. The solid line marks a growth curve $G(N)$ of an isolated peripheral population. There are two stable nontrivial equilibria (points 1 and 3 in the figure) and one unstable equilibrium (point 2). In a comparable but less isolated population with I immigrants entering per generation, the growth curve is $G(N) + I$. Adding a constant rate of immigration I to the growth equation is geometrically equivalent to a rigid, vertical translation of the growth curve graph by I units. The dashed line in the figure portrays the growth curve of the less isolated population. It can be seen that the influx of immigrants has pushed the population well above its

carrying capacity and has caused the paired equilibria at lower densities to disappear entirely. As the rate of immigration is increased, populations are less likely to be trapped at such low-density equilibria.

IMMIGRATION AND POPULATION STABILITY IN DISCRETE-TIME GROWTH MODELS

Given that a population is locally stable, we have seen that immigration should increase population size. But if negative density-dependence is too severe at high densities, populations may exist in a permanent state of fluctuation around an unstable point equilibrium. This suggests the following two questions about the impact of immigration upon a peripheral population. Can immigration destabilize an otherwise stable population, or, conversely, can immigration impose stability onto an unstable, fluctuating population? Given that a

Figure 1. The effect of immigration upon equilibrium density in a discrete-time model. Equilibria occur where $G(N)$, the curved line, crosses the straight, 45° line. Adding a constant amount of immigration raises the growth curve uniformly and equilibrium 3 increases, whereas the other equilibria vanish.

population is varying cyclically or chaotically, how does immigration modify the distribution of densities shown by the population over time?

As immigration increases, the slope of the growth curve at an equilibrium where $\lambda' < 0$ changes according to

$$\frac{\partial \lambda'}{\partial I} = \left(\frac{d^2 G}{dN^2} \right) \left(\frac{dN^*}{dI} \right).$$

Given a growth curve that is concave downward,

$$\frac{d^2 G}{dN^2} < 0,$$

immigration will push the population toward levels of increasingly severe density-dependence and may even destabilize an intrinsically stable peripheral population. Conversely, immigration may stabilize a population whose growth curve is concave upward—

$$\frac{d^2 G}{dN^2} > 0.$$

Therefore the effect of immigration upon population stability is determined by the concavity of the growth curve $G(N)$.

Standard discrete-time models provide ready examples of both effects. One model for which immigration is destabilizing is the discrete logistic equation

$$N(t+1) = N(t) \left(1 + r - \frac{r}{K} N(t) \right) + I \quad (3)$$

where r is the intrinsic growth rate. The stability properties of this model without immigration are well-understood (May and Oster, 1976; Roughgarden, 1979). In the Appendix, I outline the stability character of the discrete logistic with immigration. Figure 2 depicts the stability domains of this model. (The dashed and dotted lines are explained in the Appendix.) The overall impression from this figure is that immigration reduces or eliminates stability in the peripheral population.

By contrast, in other models immigration may stabilize an intrinsically unstable population. Figure 3 shows the results of a local stability analysis of the following model:

$$N(t+1) = N(t) \exp[r(1 - N(t)/K)] + I \quad (4)$$

With no immigration and $r < 3$, the dynamics of this model resemble those of (3) May (1976). As immigration increases, the domain of unstable behavior diminishes. The contrast between Figures 2 and 3 is striking. In a more detailed analysis it can be shown that the opposing consequences of immigration for the two models are due to the downward concavity of (3) and the upward concavity of (4) at high values of r and N .

Numerical studies of a number of discrete-time models were carried out in order to ascertain how immigration modifies the temporal distribution of densities in cyclic or chaotic populations. In such populations, a reasonable measure of abundance is the arithmetic time-average of densities

$$\langle N \rangle = \frac{1}{T} \sum_{t=1}^T N(t).$$

Figure 2. Stability regions for the discrete logistic (3). s = monotonic return to equilibrium. os = oscillatory return to equilibrium. u = locally unstable (stable cycles or chaos). The solid line is the largest value of I allowing non-negative numbers for $G(N)$. The dashed line is the outer bound of the unstable region for the modified logistic model discussed in the Appendix.

Figure 3. Stability regions for the exponential logistic (equation (6)). s = monotonic stability. o = oscillatory stability. u = unstable. The central line in the o region separates parameter choices where an increase in I decreases the rate of return to an equilibrium (to the left) from those parameters where I effects a faster return (to the right).

In cyclic populations the appropriate value for T is one cycle length, whereas in a chaotic population a large number of generations per run starting from a number of initial conditions may be required to fully characterize $\langle N \rangle$.

What is the relationship between average density and the rate of immigration? An example of a pattern that emerged repeatedly in the simulations is depicted in Figure 4. The model used for this figure has been extensively exploited by insect ecologists (e.g., Hassell, 1975). With an added immigration term the growth model is

$$N(t+1) = N(t)e^r (1 + dN(t))^{-b} + I. \quad (5)$$

The four curves in the figure correspond to four values for the intrinsic growth rate, r . At $I=0$ and high r , populations obeying equation (5) fluctuate, sometimes greatly, around an unstable point equilibrium. In these unstable populations, $\langle N \rangle$ does not increase monotonically with I . The influx of a few immigrants per generation may dramatically increase $\langle N \rangle$, and a yet greater rate of immigration may actually decrease $\langle N \rangle$.

This nonmonotonic relation between $\langle N \rangle$ and I has a simple explanation. In

discrete-time growth models such as (5), populations with high intrinsic growth rates tend to exhibit chaotic behavior (May, 1976). Time-series of populations in chaos typically show overshoots of K followed by precipitous declines in abundance. Following each population crash, several generations may elapse before population numbers are sufficiently large to produce a high total growth rate, culminating in another explosive overshoot and crash. Even a slow trickle of immigrants can greatly reduce the number of generations between successive overshoots. In model (5), population growth is essentially geometric at densities well below K . If $N(0)$ is the number in the population immediately following a crash, the population size t generations later is approximately

Figure 4. Time-averaged densities for model (7). The numbers labelling the curves are the intrinsic growth rates. The other parameters are $d = 1$, and $b = r$. For populations exhibiting chaotic behavior, the lines represent the average over many runs of 5,000 generations length, starting over a range of initial densities.

$$N(t) \cong N(0)e^{rt} + I \sum_{s=0}^t e^{rs}$$

given that numbers remain low enough for negative density-dependence to be negligible. From this expression it is clear that the time required to reach a given $\bar{N}$ (e.g., an $\bar{N}$ such that $G(\bar{N}) > K$) from $N(0)$ may be substantially shortened by regular immigration. This reduction in the amount of time needed for recovery from a population crash increases the number of generations at high densities, and therefore increases $\langle N \rangle$. At higher rates of immigration, density-dependence among the immigrants themselves tends to diminish the magnitude of the overshoot. In model (5), this diminution leads to a decrease in $\langle N \rangle$.

We can summarize the above results as follows: 1) Given no genetic differentiation between immigrants and residents, immigration should increase the density of stable peripheral populations. 2) The concavity of the growth curve determines the relation between immigration rate and stability. And finally, 3) in fluctuating peripheral populations, an increase in the rate of immigration may decrease the average size of the population.

PERIPHERAL POPULATION MODELS: THE EFFECT OF HAPLOID GENETIC VARIATION

The conjecture to be examined is that genetic differentiation between immigrants and residents alters the functional relationship between the rate of immigration and population size. In the remainder of this paper I explore the properties of models which incorporate a particularly simple kind of genetics—haploid variation with two alternative alleles at a single locus. One allele—type 1—is fixed within the species' main range, and so all immigrants are type 1, whereas the other allele—type 2—is selectively favored in the peripheral population. To explore how the balance

between selection and immigration affects both the gene frequency and population size, we modify models (1) and (2) as follows.

Counting *gene numbers*, the appropriate generalization of model (1) is

$$\begin{aligned} \frac{dN_1}{dt} &= N_1 \phi_1(N_1, N_2) + I \\ \frac{dN_2}{dt} &= N_2 \phi_2(N_1, N_2) \end{aligned} \quad (6)$$

where N_i is the density of type i ($i=1,2$), ϕ_i is the *per capita* instantaneous growth rate or the absolute fitness of type i , and I is the immigration rate of type 1. The total population size is defined as $N = N_1 + N_2$, the gene frequency of allele 1 is $p = N_1/N$, and the gene frequency of allele 2 is $q = 1 - p$. The mean fitness of the population is $\bar{\phi} = p\phi_1 + q\phi_2$. An alternative representation of system (6) is obtained by differentiating N and p with respect to t

$$\frac{dN}{dt} = N\bar{\phi} + I \quad (6')$$

$$\frac{dp}{dt} = pq(\phi_1 - \phi_2) + q\frac{I}{N}.$$

In like manner, we may embody haploid genetic variation within the framework of the discrete-time model (2) as follows:

$$\begin{aligned} N_1(t+1) &= N_1(t)g_1(N_1, N_2) + I \\ N_2(t+1) &= N_2(t)g_2(N_1, N_2) \end{aligned} \quad (7)$$

or, equivalently,

$$\Delta N = N(\bar{g} - 1) + I \quad (7')$$

$$\Delta p = \frac{pq(g_1 - g_2) + q\frac{I}{N}}{\bar{g} + \frac{I}{N}}$$

where g_i is the per-capita growth rate or absolute fitness of type i and $\bar{g} \equiv pg_1 +$

qg_2 is the mean fitness of the population. The analysis of these two models proceeds through the usual steps of first solving for equilibria (N^*, p^*) , determining the local stability of each equilibrium, and then examining the behavior of the model away from equilibria. Models (6)' and (7)' explicitly display the genetic character implicit in models (6) and (7).

DENSITY-DEPENDENT SELECTION

It is difficult to make much headway without specifying in more detail the fitness functions ϕ_i and g_i . A selective regime that has received a great deal of attention from population biologists is density-dependent selection, for which ϕ_i and g_i may be written as functions of total density N , $\phi_i(N)$ and $g_i(N)$ (Roughgarden 1971, 1976; Charlesworth 1971, 1980; Asmussen, 1979). A significant finding of these theoretical studies is that if fitnesses are strictly density-dependent, and the population is stable, natural selection adjusts gene frequencies in such a way that population density is locally maximized at the joint demographic and genetic equilibrium.

First consider the continuous-time model (6). To simplify the analysis, I assume that each allele has a carrying capacity K_i such that $\phi_i(K_i) = 0$, and that there is negative density-dependence at all densities:

$$\frac{d\phi_i}{dN_i} < 0.$$

In the absence of immigration, it is straightforward to show that the allele with higher K excludes the alternate allele. Since type 2 is assumed to be locally favored, $K_2 > K_1$ and the population equilibrates at K_2 .

As the rate of immigration increases, so does the frequency of type 1. However, given that the population is polymorphic for both alleles, population size is *independent* of the rate of immigration. For the population to be at its equilibrium

(N^*, p^*) both growth rates in (6) must equal 0, so $N_2^* \phi_2(N^*) = 0$. As the population is assumed to be polymorphic, $N_2^* > 0$. Thus $\phi_2(N^*) = 0$. But by assumption, $\phi_2 = 0$ only when $N^* = K_2$. Therefore, the total population size remains at K_2 as the rate of immigration increases. Immigration does not change the total population size, but instead shifts the relative proportions of the two types. From (6') we find that

$$p^* = \frac{-I}{K_2 \phi_1(K_2)}$$

(since $K_2 > K_1$, $\phi_1(K_2) < 0$). Hence, a polymorphic equilibrium exists only if $I < |K_2 \phi_1(K_2)|$. In the Appendix it is shown that the equilibrium

$$(N^*, p^*) = (K_2, \frac{-I}{K_2 \phi_1(K_2)})$$

is both locally and globally stable.

Thus we have completely characterized the interplay of immigration and selection in a general, haploid model of density-dependent selection. As long as $I < |K_2 \phi_1(K_2)|$, immigration merely shifts the genetic composition of the peripheral population without changing its total density at all. The opposing ecological and genetic effects of immigration discussed in the introduction exactly cancel each other out.

By a parallel argument it can be shown that the comparative statistics of the discrete-generation model (7) are identical to those of the continuous-generation model. Given fitnesses that are density-dependent, equilibrium densities are independent of the rate of immigration in polymorphic populations. As with the purely ecological models discussed earlier, the principal difference between models (6) and (7) is that the latter may exhibit sustained oscillatory behavior. The point equilibrium of (7) is locally stable only if the magnitude of the real part of the dominant eigenvalue λ is less than unity. Let

$$a_{ij} \equiv \frac{\partial}{\partial N_j} (N_i g_i).$$

Then the two eigenvalues are

$$\lambda = [a_{11} + a_{22} \pm ((a_{11} - a_{22})^2 + 4a_{12}a_{21})^{1/2}]/2.$$

As both a_{12} and a_{21} are negative, the discriminant is positive and both eigenvalues are real numbers. We wish to understand how immigration alters the stability properties of a peripheral population, and how immigration perturbs average abundance in unstable populations. Substituting explicit fitness functions such as (3) or (4) leads to cumbersome masses of algebra from which it is difficult to extract necessary and sufficient conditions for local stability. However, it is plausible that if immigration reduces the frequency of a locally favored allele, the peripheral population could be made more stable. In discrete-time population models, the intrinsic growth rate r typically "tunes" the dynamic behavior of the model—the magnitude and period of oscillations usually increase with r . As immigrants are assumed to be less fit than residents, the values of r , or K , or both, for the immigrant should be lower. Diluting a population at high r with an admixture of immigrants with lower r should reduce the average r of the population as a whole, and therefore tend to stabilize an unstable peripheral population.

In like manner, a lower tolerance of crowding in immigrants ($K_1 < K_2$) may enhance the stabilizing effect of immigration. As a particular example, consider a population with discrete generations in which the fitnesses of each type are described by $g_i(N) = \exp[r(1 - N/K_i)]$. The two alleles share the same r but differ in K . If the rate of immigration is too great, the polymorphism will not persist. As depicted in Figure 5, the maximum I consonant with polymorphism varies with both r and K_1 . The local stability

Figure 5. Maximum rate of immigration I permitting polymorphism in model (9) with equation (6) as a fitness function. $K_2 = 10$. $r_1 = r_2$. K_1 is denoted by the number marking each curve.

Figure 6. Stability regions of model (9) with fitnesses given by (6). s = monotonically stable. os = damped oscillations. u = unstable. The solid lines mark the edges of stability regions for $K_1 = 1$, and the dashed lines for $K_1 = 5$. The dotted lines are from Figure 3 and depict the stability character of a population in which the immigrants are type 2.

properties of this model are displayed in Figure 6. We have already observed that immigration is weakly stabilizing in this model when immigrants are of the same genetic type as the residents; the dotted lines, lifted from Figure 4, demarcate transitions from monotonic to oscillatory stability (left line) and from locally stable to unstable point equilibria (right line). The solid lines demarcate the edges of

the corresponding stability regions for immigrants with $K_1 = 1$, and in like manner the dashed lines for $K_1 = 5$. Decreasing the carrying capacity of the immigrants heightens the stabilizing influence of immigration.

In discrete-time haploid models such as (6), as in the purely ecological model (2) discussed above, instability may complicate the relation between immigration and population size. In numerical studies I have found that polymorphisms are difficult to maintain if the demographic attributes of each type lead to severe oscillations in monotypic populations. In a stable population, a low immigration rate produces a low frequency of the immigrant type. The same rate of immigration can lead to a much higher gene frequency and even fixation in an unstable population, for the simple reason that following a population crash a few immigrants may comprise a sizeable fraction of the total population. During these crashes the locally adapted type can go extinct because of genetic drift, while immigration steadily replenishes the less fit type. Gene flow should hamper local adaptation more readily in unstable than in stable peripheral populations.

If the polymorphism does persist, over the course of population fluctuations high densities are correlated with low frequencies of the immigrant type. In the particular model used for Figures 5 and 6, the time-average of density, $\langle N \rangle$, seems to be independent of I , but in other haploid models $\langle N \rangle$ may decrease with increasing I .

From this analysis of density-dependent selection in haploid populations, we may conclude that 1) the rate of immigration does not affect the total density of stable, polymorphic populations; 2) decreasing the fitness of immigrants relative to residents strengthens the stabilizing effect of immigration upon population dynamics; and 3) the main-

tenance of local adaptation in the face of gene flow is unlikely in severely fluctuating populations.

OTHER GENETIC MODELS

Two obvious generalizations of this model are to employ more general fitness functions in (5) and (6) and to develop comparable diploid models. Space limitations preclude a full treatment of these extensions here, so I will merely outline a few salient changes in the results produced by these modifications in the models. If the absolute fitness functions in models (5) or (6) are functions of the separate densities of the two types rather than of their summed density, the selective regime is a mixture of density-dependent and frequency-dependent selection. Alternative stable states may exist (Fenchel, 1975). For the continuous-time model (6), if the population is at a stable equilibrium, it can be shown that

$$\frac{dN^*}{dI} = \frac{dN_1^*}{dI} \left(1 - \frac{\partial \phi_2 / \partial N_1}{\partial \phi_2 / \partial N_2} \right)$$

where the partial derivatives are evaluated at (N_1^*, N_2^*) . Since immigration increases the density of N_1 , the overall effect of immigration upon the total population hinges upon the relative magnitudes of the derivatives $\partial \phi_2 / \partial N_1$ and $\partial \phi_2 / \partial N_2$. If absolute fitnesses are determined solely by the total density of the population,

$$\frac{\partial \phi_2}{\partial N_2} = \frac{\partial \phi_2}{\partial N_1},$$

and N^* is independent of I . Immigrants that exert a disproportionate effect on the fitness of residents, compared with the effect of residents upon themselves,

$$\frac{\partial \phi_2}{\partial N_1} > \frac{\partial \phi_2}{\partial N_2},$$

decrease the total size of the population.

Even without immigration the diploid equivalents of (6)' and (7)' can manifest a rich panoply of behaviors. Asmussen (1979) has analyzed a discrete-generation model of density-dependent selection in which (4) served as a fitness function. She demonstrated the existence of regular and chaotic cycling in both population size and gene frequency. In contrast to classical selection models, heterozygote superiority in crowding tolerance is not required for the maintenance of genetic variation in these cyclic populations. All these features are retained in the models with immigration. An additional feature of the diploid models is that the relationship between immigration and population size depends upon the dominance relations of alternative alleles in the peripheral population. For instance, consider an allele that is simultaneously genetically dominant yet locally unfit. An increase in the rate of immigration of individuals bearing that allele has three distinct effects, two of which were discussed above. First, there is the ecological effect of increasing density by increasing the influx of individuals into the population; this corresponds to a purely ecological model such as (1). Second, if fitnesses are density-dependent, this increase in density lowers the fitness of the locally favored genotype, the numbers of which will decline until the overall density of the population is unchanged; this accounts for the uncoupling of immigration rates and equilibrium densities in the haploid models (6) and (7). Third, since in a Mendelian population the average fitness of the offspring produced by a genotype is partially determined by the array of genotypes available for mating, the more immigrants there are, the more the fitness of the resident's offspring will be diluted by cross-matings. The number of residents

declines even further to compensate for this mating effect. The final pattern is an inverse relation between immigration rate and population density. By contrast, if the immigrant allele is recessive the mating effect is greatly diminished, and, just as in the haploid model, population size may be essentially independent of the rate of immigration.

A third theoretical approach, and one that ultimately might be the most profitable, is to extend quantitative genetic models of phenotypic evolution (Lande, 1976) in order to examine how immigration by individuals with nonoptimal phenotypes in one or more character states influences population size in density-regulated populations. Antonovics (1976) and Slatkin (1978) suggest some promising lines of development for such a theory.

DISCUSSION AND CONCLUSIONS

I have argued that two characteristics of a peripheral population should mold the functional relation between immigration rate and population size—its dynamic stability, and the degree of its genetic adaptation to the local environment. To begin understanding the dynamics of a peripheral population we must first isolate those factors responsible for its low population density. It is widely believed that density-independent factors act more severely in peripheral than in central populations. For instance, Mayr (1963) asserts that "the border region is a place in the area of a species where density-dependent factors are of minor importance." Alternatively, the peripheral population may be rare because of an increase in the intensity of density dependence. In populations with delayed density-dependence, these two explanations for rarity at a species' border have fundamentally different implications. If rarity results from a high rate of density-independent mortality the intrinsic growth rate of the population will be low

and the population will stably persist at its carrying capacity, at least in temporally constant environments. By contrast, if r is high but K is low, the population may exhibit fluctuations of considerable magnitude around K , sometimes to the point of extinction. As the demographic and genetic consequences of immigration depend upon the stability character of the peripheral population, the first step in gauging the role of immigration should be to try to understand the causal basis for population fluctuations at the species' border.

Immigration may not change, or may even decrease, the density of a population that has adapted to its local environment. Scant data exists for testing this idea, in part because the requisite evidence is technically difficult to obtain. As noted above, there recently has been a de-emphasis of gene flow as a constraint on the evolution of local adaptation. To the extent that this view is valid, we should expect the predicted phenomenon to be rarely observed. An additional reason for the rarity of relevant data, however, is that many ecologists simply assume that organisms are well adapted to their environment. Although usually reasonable, in peripheral or ecologically marginal populations this assumption may well be false. I know of two possible examples of local maladaptation in peripheral populations resulting from gene flow. Camin and Ehrlich (1958) argued that on islands in Lake Erie a balance between migration and selection maintained a polymorphism in banding patterns of water snakes (*Natrix sipedon*). A substantial fraction of the island populations had locally unfit banding patterns. Stearns and Sage (1980) have suggested that the life-history traits of one freshwater population of the mosquito-fish (*Gambusia affinis*) might be best interpreted as maladaptation to the freshwater environment. Gene flow from a nearby brackish-water population may account for the apparent lack of local adaptation in this population. In neither

instance is anything known about the effect of the apparent maladaptation upon population density. Comparable situations might well exist in many organisms with passive dispersal or territoriality residing in locally heterogeneous environments. For instance, were the habitat of a species of *Anolis* a mosaic of high and low quality patches, territorial behavior could lead to a steady flow of individuals into low density patches. This influx could reduce the efficacy of selection for better adaptation to the low-quality patch type.

A critical and very difficult empirical problem is to gauge the relative magnitudes of spatial scales associated with dispersal, selection, and nonselective determinants of density. Slatkin (1973) has suggested that a useful quantity for understanding evolution in spatially structured populations is the ratio between the average dispersal distance and a measure of the gradient in selection. If this ratio is small, a species should closely track spatial variation in selection (Roughgarden, 1979); since immigrants into a population are drawn primarily from nearby populations, and nearby populations experience similar selective regimes, immigrants should genetically resemble residents. In this case the purely ecological models discussed above are appropriate. By contrast, if the ratio is large, many immigrants will be from relatively distant populations with different selection pressures, and immigrants may differ greatly from the locally favored type. Understanding the role of immigration in population dynamics in this case requires a blending of ecology with genetics.

For the rate of immigration to be large enough, relative to carrying capacity, to qualitatively perturb the abundance or dynamics of a peripheral population, there usually must be a second, more abundant population in reasonable proximity to the first—a sharp spatial gradient in densities should exist. This density gradient need not correspond to a

comparable selection gradient, since many factors that affect density do so indirectly. For instance, a consumer such as a detritivore may be able to alter the abundance, R , but not the rate of renewal of its resource. Individual consumers are food-limited, and there is no direct interference among them, we can write the *per capita* consumer growth rate as some increasing function of resource abundance, $Y(R)$. The consumer population should grow until its collective demand has reduced resource availability to some level, C , at which each consumer just replaces itself with a single descendent. If the parameters determining the form of Y and the quantity C are spatially invariant, selection will act uniformly on these parameters throughout the consumer's range. However, the number of consumers present at equilibrium is determined indirectly by the renewal rate of the resource, which might well vary greatly over space. Hence, a consumer's density may be highly variable in space even though selection is not; dispersal could then be of more consequence for the ecology than for the evolution of the species. In other circumstances, of course, density gradients may closely match selection gradients.

The degree to which ecological phenomena are influenced by intraspecific genetic variation is a problem of great current interest. I have here argued that the ecological consequences of dispersal into a population depend upon whether or not immigrants differ genetically from residents. In the study of species' borders, in particular, it should be realized that ecological and genetical phenomena may be inextricably entangled.

ACKNOWLEDGMENTS

Ernest E. Williams has long emphasized, both in his own work and in his guidance of graduate students, the potential value of species' borders as loci for the study of significant ecological and

evolutionary problems. My own work on models of peripheral populations, some of which is presented in this paper, received its initial stimulation from conversations we had on the question of the ecological and evolutionary stability of species borders.

I would like to express my gratitude to Ernest and to William Bossert for patiently listening to my ruminations about species' borders while I was a graduate student at Harvard. I thank Steve Hubbell, Phil Hedrick, Jim Hamrick, Mike Gaines, Norman Slade, and Gregory Glass for comments on the work reported here, and Russell Lande and Juli Armstrong for valuable comments on the manuscript. My graduate assistant Juli Armstrong kindly assisted with the numerical simulations, and the University of Kansas supplied the requisite computer funds. I thank Jan Elder, Coletta Spencer, and my wife Lynne for assisting with manuscript preparation.

APPENDIX

THE DISCRETE LOGISTIC MODEL

Without immigration, the single non-trivial equilibrium is $N^* = K$, and the stability-setting eigenvalue is $\lambda = 1 - r$. N converges asymptotically to K if $r < 2$. Stable limit cycles occur when $2 < r < 2.828$, whereas for $2.828 < r \leq 3$ the population exists in a chaotic regime in which the long-term behavior of any particular population depends upon its initial conditions; the behavior may be either periodic or aperiodic (May and Oster, 1976). Given $r > 3$, the model can predict negative densities; to preclude this biological absurdity we require that $r \leq 3$. In like manner, as I increases we would like to ensure that population production $G(N)$ is never negative. For the discrete logistic, a population minimum occurs during the generation immediately following a maximum. The maximum value of $N(t + 1)$ occurs at the critical point $(N(t) = (K(1 + r)/2 \text{ of } (3),$

so $\max N(t + 1) = \frac{K(1 + r)^2}{4r} + I$. If we require that $G(\max N) \geq 0$, we find that the ratio I/N is bound by the following expression:

$$\frac{I}{K} \leq \frac{1}{2} + \frac{3}{4r} - \frac{r}{4}.$$

This bound on I/K is the dotted line of Figure 2. We can avoid the annoying possibility of negative densities by simply setting $N(t + 1) = I$ whenever $G(N)$ is a negative quantity, which occurs whenever

$$N(t) > K \left(\frac{1+r}{r} \right).$$

For this modified discrete logistic, the dashed line in the Figure is an *upper* bound on the parameter values that allow fluctuating densities.

Let $Y = (1 + 4I/rK)^{1/2}$. Then $N^* = K(1 + Y)/2$, and the eigenvalue is $\lambda = 1 - rY$.

As N^* increases with increasing I , λ decreases. The maximum r consistent with local stability is

$$r = 2((I^2/K^2 + 1)^{1/2} - I/K).$$

THE STABILITY OF THE CONTINUOUS-TIME HAPLOID MODEL (6)

The two eigenvalues are

$$\lambda = [a \pm (a^2 - 4\phi_1 N_2 \frac{d\phi_2}{dN})^{1/2}]/2,$$

where

$$a = \phi_1 + N_1 \frac{d\phi_1}{dN} + N_2 \frac{d\phi_2}{dN}.$$

Since the quantities

$$\phi_1, \frac{d\phi_1}{dN}, \text{ and } \frac{d\phi_2}{dN}$$

are all negative at $N = K_2$, $a < 0$. This implies that $\text{Re}(\lambda) < 0$. Moreover, straightforward manipulations show that the separatrix

$$a^2 - 4\phi_1 N_2 \frac{d\phi_2}{dN}$$

is positive, so (N^*, p^*) is a stable node. The per capita growth rates of type 1 and 2 were assumed to be negatively density-dependent at all densities. Applying Theorem (2) of Hastings (1978) we find that the equilibrium is globally stable. The critical assumption required for this result is that the fitness of both types depends only upon the total density of the population.

LITERATURE CITED

- ANTONOVICS, J. 1968. Evolution in closely adjacent plant populations. VI. Manifest effects of gene flow. *Heredity*, **23**: 507-524.
- . 1976. The nature of limits to natural selection. *Ann. Missouri Bot. Gard.*, **63**: 224-247.
- ASMUSSEN, M. A. 1979. Regular and chaotic cycling in models of ecological genetics. *Theor. Pop. Biol.*, **16**: 172-190.
- CAMIN, J. H., AND P. R. EHRLICH. 1958. Natural selection in water snakes (*Natrix sipedon* L.) on islands in Lake Erie. *Evolution*, **12**: 504-511.
- CHARLESWORTH, B. 1971. Selection in density-regulated populations. *Ecology*, **52**: 469-474.
- . 1980. *Evolution in age-structured populations*. Cambridge, Cambridge University Press.
- ENDLER, J. A. 1977. *Geographic variation, speciation, and clines*. Princeton, Princeton Univ. Press.
- FENCHEL, T. 1975. Factors determining the distribution patterns of mud snails (Hydrobiidae). *Oecologia*, **20**: 1-17.
- GAINES, M. S., AND L. R. MCCLENAGHAN, JR. 1980. Dispersal in small mammals. *Ann. Rev. Ecol. Syst.*, **11**: 163-195.
- HASSELL, M. P. 1975. Density dependence in single-species populations. *J. Anim. Ecol.*, **44**: 283-295.
- HASTINGS, A. 1978. Global stability of two-species systems. *J. Math. Biol.*, **5**: 399-403.
- LANDE, R. 1976. Natural selection and random

- genetic drift in phenotypic evolution. *Evolution*, **30**: 314-334.
- LEVIN, D. A., AND H. W. KERSTER. 1974. Gene flow in seed plants. *Evol. Biol.*, **7**: 139-220.
- LEVIN, S. A. 1978. Population models and community structure in heterogeneous environments, pp. 439-476. *In* S. A. Levin (ed.), *Studies in Mathematical Biology Vol. 2: Populations and Communities*. Washington: Math. Assoc. Amer.
- MAY, R. M. 1976. Simple mathematical models with very complicated dynamics. *Nature*, **261**: 459-467.
- MAY, R. M. AND G. F. OSTER. 1976. Bifurcations and dynamic complexity in simple ecological models. *Amer. Nat.*, **110**: 573-599.
- MAYR, E. 1963. *Animal species and evolution*. Cambridge, Harvard Univ. Press.
- NAGYLAKI, T. 1978. Clines with asymmetric migration. *Genetics*, **88**: 813-827.
- OKUBO, A. 1980. *Diffusion and ecological problems: mathematical models*. Berlin, Springer-Verlag.
- ROUGHGARDEN, J. 1971. Density-dependent natural selection. *Ecology*, **52**: 453-468.
- . 1976. Resource partitioning among competing species—A coevolutionary approach. *Theor. Pop. Biol.*, **9**: 388-424.
- . 1979. *Theory of population genetics and evolutionary ecology: An introduction*. New York, MacMillan.
- SLATKIN, M. 1973. Gene flow and selection in a cline. *Genetics*, **75**: 733-756.
- . 1978. Spatial patterns in the distributions of polygenic characters. *J. Theor. Biol.*, **70**: 213-228.
- SPIETH, P. T. 1979. Environmental heterogeneity: A problem of contradictory selection pressures, gene flow and local polymorphism. *Amer. Nat.*, **113**: 247-260.
- STEARNS, S. C., AND R. D. SAGE. 1980. Maladaptation in a marginal population of the mosquito fish, *Gambusia affinis*. *Evolution*, **34**: 65-75.
- TAMARIN, R. H. 1977. Dispersal in island and mainland voles. *Ecology*, **58**: 1044-1054.
- TAYLOR, L. R., AND R. A. J. TAYLOR. 1977. Aggregation, migration, and population mechanics. *Nature*, **265**: 415-421.
- WILLIAMS, E. E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. *Quart. Rev. Biol.*, **44**: 345-389.

Distribution and Conservation of the Black Caiman (*Melanosuchus niger*)

MARK J. PLOTKIN¹

FEDERICO MEDEM²

RUSSELL A. MITTERMEIER³

ISABEL D. CONSTABLE⁴

ABSTRACT. The black caiman (*Melanosuchus niger*) once ranged throughout much of Amazonia, but its populations have decreased sharply due to hunting for the skin trade and for food, incompatibility with man and his domestic animals, and habitat destruction. Viable populations of black caiman apparently still exist in Ecuador, French Guiana, and Peru, but only in Peru do they occur within protected areas. Establishment of reserves for protection of the Ecuador and the French Guiana populations is recommended as well as better enforcement of existing legislation in all of the countries in which the black caiman occurs.

INTRODUCTION

Most of the world's living crocodilian species are rapidly disappearing, mainly as a result of hunting for the skin trade, hunting for food, incompatibility with man and his domestic animals, and, in some areas, habitat destruction or modification. Five crocodilian species are found in the Amazonian region of northern South America (including the Amazon and Orinoco drainages and the

Guianas). The three smaller species (*Caiman crocodilus*, *Paleosuchus trigonatus*, *Paleosuchus palpebrosus*), are still relatively abundant over much of their range and are probably among the world's least endangered crocodilians. In striking contrast, the two larger species, *Crocodylus intermedius* and *Melanosuchus niger*, are both considered endangered and are already extinct over much of their former range. Here we review the conservation status of the black caiman, *Melanosuchus niger* and discuss several key areas important for its continued survival.

PAST DISTRIBUTION

The black caiman was once widespread in Amazonia. It has been recorded from Brazil, Bolivia, Peru, Colombia, Ecuador, Guyana, French Guiana, and even Paraguay. In fact, the only Amazonian countries from which there are no authenticated records are Surinam and Venezuela. The historical distribution of the species, based on museum localities, literature records, and sightings by the authors or other reliable researchers, is shown in Figure 1.

Within this vast area, the black caiman

¹Harvard Botanical Museum, Cambridge, Massachusetts 02138, U.S.A.

²Universidad Nacional de Colombia, Bogota, Colombia.

³World Wildlife Fund, 1601 Connecticut Avenue, NW, Washington, D.C. 20009, U.S.A.

⁴Brown University, Providence, Rhode Island 02912, U.S.A.

Figure 1. Historical distribution of *Melanosuchus niger*. ▲ = populations currently considered to be viable.

appears to prefer quiet waters to the currents of the larger rivers. In large rivers like the Amazon it is found in quieter water around bends and especially in the lakes and lagoons beside the mainstream. It is also fond of the creeks and swamps associated with the Rio Branco and the Amazon delta islands of Marajó, Mexiana, and Caviana. In the dry months of November and December in lower Amazonia, large individuals sometimes estivate until the beginning of the wet season in January.

The black caiman was once incredibly abundant in many areas. Bates (1863) recorded their high density in the upper Rio Solimões, and enormous numbers were also to be found around the delta islands of Mexiana, Marajó, and Caviana (Wallace, 1853; Goeldi, 1898; Hagmann, 1902).

Today, however, the black caiman is either extinct or very rare in areas where it once was common. It is no longer possible to see them along the mainstream of the Rio Solimões where Bates was so impressed by their numbers, and they have all but disappeared from the delta islands. They are now rare or absent in most other parts of Amazonia, and are regularly encountered only in the more remote lake and tributary rivers—and never in numbers approaching those of the past century.

REASONS FOR DECLINE

The main reasons for the decline of the black caiman are large scale hide-hunting and incompatibility with man and his domestic animals, with hunting as a food source playing a role in some areas. Habitat destruction does not appear to have been a major factor in the past, although it could well become one in the near future.

Hide-hunting was responsible for tremendous slaughter during this century, and the black caiman was long the species of choice among the caimans because of its size and abundance. Commercial hide-hunting apparently began in the late 1930's and 1940's, and by the 1950's enormous numbers of caiman skins were being exported from Amazonia. For instance, Fittkau (1973) estimated that up to 5,000,000 caiman skins were removed from Brazilian Amazonia each year in the early 1950's, and that one or two additional animals were killed for each of the officially reported skins. Between 1960 and 1970, the total number of skins was still between 200,000 and 700,000 per year. However, many of these skins were of the smaller *Caiman crocodilus* which became increasingly important as *Melanosuchus niger* populations declined in Brazil. Export figures vary somewhat (e.g., Melo Carvalho, 1967; Medem, 1972), but it is clear that many millions of black caimans were

killed for their hides in the middle decades of this century.

Incompatibility with man and his domestic animals was another problem. Though not nearly as dangerous as some of the larger Asian, African, and South American crocodiles (e.g., *C. porosus*, *C. niloticus*, *C. intermedius*), the black caiman occasionally attacks man (Bates, 1863; P. Soini, personal communication). As a result of their potential danger to man, black caiman have often been wiped out in the vicinity of human settlements—even if their hides aren't necessarily used.

Black caiman were considered a major threat to cattle ranching activities, especially in the seasonally flooded campos or grasslands like those on Marajó or Mexiana. Ranchers once held great caiman drives in the dry season to exterminate as many of the crocodilians as possible. Hagmann (1902) reported a drive in which about 800 animals between 1 and 4.2 m were killed in two days. Similar drives were also held in other parts of Amazonia, the most recent we know of in the 1940's in the Brazilian state of Amapá (R. Renau Ferrer, personal communication).

On top of this, the black caiman was and sometimes still is hunted for its meat or fat in parts of its range (Wallace, 1853). Black caiman meat, especially that of the tail, is still eaten in some parts of the animal's range (e.g., the Kaw region of French Guiana, the Río Tapiche in Peru, and several tributaries south of Belem, Pará).

CURRENT STATUS

The tremendous slaughter of black caimans over the past century resulted in a steep population decline, and this species is now either extinct or rare in most of its former range. Only a few areas apparently harbor substantial populations, although none of these approach those observed by the nineteenth-century naturalists.

The black caiman is officially protected in most of countries in which it occurs and appears in the Red Data Book as an endangered species and on Appendix 1 of the Convention on International Trade in Endangered Species, and the U.S. Endangered Species List. The hide trade has clearly diminished from the levels of the 1950's, but this is probably as much due to large scale decline of populations as to protective legislation. Although export is now illegal, black caimans continue to be killed, and enforcement in the remote areas where they still occur is virtually nonexistent. Although the vastness of the species' range will probably prevent it from disappearing completely before the turn of the century, it is certainly not exaggerating to say that the present population is probably less than 1% of what existed a century ago, and the decline still continues.

In this section, we review the current situation and existing protective legislation on a country by country basis.

BOLIVIA

The black caiman once occurred in much of northern and eastern Bolivia, apparently as far south as Laguna Caceres on the Brazilian border near Corumbá, and Puerto Grether, further inland on the Río Mamoré.

Hide-hunting was particularly heavy during the period 1942–1960, and continues today. In 1973, Medem found over 2,000 skins (including 120 over 3 m in length) in tanneries in Cochabamba and Trinidad. He also saw a shipment of 500 black caiman skins in Riberalta in June 1973, all of them from the middle course of the Beni. Some 70,000 *Melanosuchus* skins were exported between 1967 and 1973 and Campbell (1977) reported 4580 skins exported in 1976.

The hide industry in Bolivia has developed sophisticated methods for exploiting caimans. Small planes are used to locate isolated lakes and ponds in the forest, the position is fixed on aerial

photographs, and well-equipped hunting crews then cut trails or travel by canoe to the area and exterminate the caiman population. Indications are that helicopters may be used in the future if enough caiman are present to make it commercially feasible.

Since the tanning industry in Bolivia is second only to mining and oil and gas in export and since nature legislation is largely ineffective, the black caiman does not seem to have much of a future in this country and has already been wiped out in areas in which it once occurred. It is clearly a highly endangered animal within Bolivia.

BRAZIL

Once widespread in Brazilian Amazonia, the black caiman is now rare or locally extinct over most of its historical range. A small population occurs in the Amazonas National Park (Magnusson, 1979) and this may also be true in the other large parks that have been established in Brazilian Amazonia in the past few years, but survey work is needed to see if they are large enough to be viable over the long term.

Illegal hunting continues and Magnusson (1980) reports that most river passenger/freight boats carried caiman (*Melanosuchus* and *Caiman*) hides in 1979 and 1980. In 1973, Mittermeier found skulls of black caiman that had been eaten in small villages along the Rio Negro and the Rio Solimões.

COLOMBIA

The black caiman was once abundant in the Colombian Amazon, from Leticia to the Río Atacuari at the Peruvian border, and along the Río Putumayo and the Río Caquetá, major Colombian tributaries of the Amazon. Along the upper Putumayo, its range was limited by the falls and rapids of Araracuara; on the Río Yari, a tributary of the Caquetá below Araracuara, it was limited by the rapids of La Gamitana; and on the Río Mesai, a

tributary of the Yari, it was limited by the falls known as El Bufo. It occurred in the lower Río Apaporis, largest tributary of the Río Caquetá, but its spread up this river was prevented by the rapids of La Libertad as well as two other sets of rapids.

Although larger falls and rapids proved a barrier to *Melanosuchus* distribution in Colombian Amazonia, smaller ones did not. The falls known as Chorro de Cordoba on the Caquetá, above the village of La Pedrera, did not limit its spread into the Río Mirití-Paraná and the Río Cahuinari, and later up to the Serranía de Araracuara (Medem, 1963).

In the 1940's some attempts were made to introduce black caiman into the upper Río Caquetá, above Araracuara (Medem, 1960, 1963). For instance, in 1943, a sergeant of the Colombian navy released 25 juvenile *M. niger* into the Caquetá near the village of Tres Troncos. These animals apparently did not reproduce. However, some may have survived for at least a while since an individual approximately 300 cm in length was killed along the Río Sencella, a tributary of the upper Caquetá, and it may have been one of the last of these 25 juveniles.

Commercial hide-hunting began along the Río Amazonas before 1945 and on the lower Río Caquetá and the Río Putumayo in 1950 and resulted in a rapid decline of *Melanosuchus* populations throughout the region. By the late 1950's, the species was already rare. Medem, for instance, did not observe any specimens during expeditions to the La Pedrera region and the lower Río Apaporis. He only saw six juveniles in Lago Tarapoto and the Río Loreto-yacu above Puerto Nariño (October, 1958) and saw none at La Concepción on the Putumayo (November, 1958). In 1968, Medem also did not see any black caiman during a three month expedition on the Río Caquetá, Mirití-Paraná, and Cahuinari, although local Indians claimed that a few widely-scattered and shy individuals still existed.

Few black caiman were seen in Colombia in the 1970's. Roger Foote observed black caiman in the wild only once between 1973 and 1975. This was a sighting of four individuals under 120 cm in length in a shallow lake about one hour from Leticia and 30 minutes from the town of Benjamin Constant (R. Foote, *in litt.* to Medem, July 5, 1975). In 1977, two Colombian biologists, Hernando Chirivi Callego and Jorge Morales, made a two-month survey of *M. niger* populations along the Amazon and the Putumayo. They found only one specimen, a 62 cm juvenile kept by a settler at the mouth of the Río Caucaya. Finally, in 1977 a female about 350 cm in length was reported killed by a settler on the Río Caquetá, close to La Pedrera (P. von Hildebrand, personal communication). No more recent records exist.

Despite the passage of laws designed to protect the black caiman, hunting continued at least until 1972. Statistics obtained from INDERENA and from the Instituto de Comercio Exterior (INCOMEX) reveal that a total of 66,369 hides and live animals were illegally exported from Leticia, Bogotá, Barranquilla, and Cali in 1970 (Medem, 1971) and 1972 (Table 1). After 1972, official export statistics were no longer available.

It is clear that hide hunting has all but exterminated the black caiman in Colombia, and that this animal is highly endangered having been reduced to small numbers of widely dispersed individuals. A major conservation program would be necessary to restore it to even a fraction of its former abundance, and it may already be too late for such a program to be undertaken.

ECUADOR

In Ecuador, the black caiman is found in the drainage of the Río Napo, extending east almost to the foothills of the Andes. It is especially fond of lakes and lagoons and sometimes occurs in large rivers at places where the current is weak. It has

been recorded from several tributaries of the Curaray, the Conambo, and the Pindo, themselves tributaries of the Marañon (upper Río Amazonas). Black caiman occur at relatively high densities in Limon Cocha (K. Miyata, personal communication) and may also occur in established conservation units in Amazonian Ecuador such as Yasuní National Park, Cuyabeno Faunal Production Area, and Lagartococha Ecological Reserve (C. Freese, personal communication).

There appears to be little active hide-hunting in Ecuador at this time. However, there was once a very active hide industry for the American crocodile (*Crocodylus acutus*) in Ecuador (C. Freese, personal communication), but we were unable to determine if the black caiman was also included.

FRENCH GUIANA

The black caiman occurs only in the extreme northeastern corner of French Guiana. In this region, it can still be seen in the seasonally flooded grasslands (*les Marais de Kaw*) of the remote Kaw River and in the largely inaccessible swampy area known as *la Savanne Angelique*, which is bordered by the Kaw River, the Montagnes de Kaw, and the Montagne de Gabrielle. Black caiman are also sometimes found to the east of Kaw between the lower Approuague River and the Ounary River (J. Lescure, personal communication), at a few sites along the lower Approuague River (particularly at the mouth of the Kaw Canal and by the Mantouni and Aipoto Islands, J. Lescure, personal communication), and in the small Ouapou Creek to the south of the Montagnes de Kaw. They once occurred in the Gabrielle Creek, the Savanne Gabrielle, and the Mahury River to the west of Kaw, but are now extinct there (J. Lescure, personal communication). They also were once common along the lower Oyapock and its tributaries, the Gabaret and the Ounary River, but Brazilian

TABLE 1. EXPORT DATA FOR *MELANOSUCHUS NIGER* SKINS FROM COLOMBIA.

Year	Hides/Live caimans	Site	Source
1970	30,105 (H)	Leticia	Medem, 1971
1970	436 (H)	Tarapaca	INDERENA, March, 1971
1970	7,633 (H)	Bogota, Baranquilla	Medem, 1971
1970	259 (L)	Bogota, Baranquilla	Mendal Bros., Lao, INDERENA
1970	43,417 (H&L)	(Total, 1970)	Medem, 1971
1972	22,254 (H)	Bogota, Cali	INCOMEX
1972	698 (H)	Leticia	Scheuerman and Foote, <i>in litt.</i>
1972	22,952 (H)	(Total, 1972)	
Total for 1970 and 1972: 66,369 (H & L)			

poachers have almost completely wiped out the animal along the Oyapock (although occasional individuals can still be seen there—e.g., Condamin, 1975; R. Renau Ferrer, personal communication). Stragglers even appear from time to time on the beaches of Cayenne and nearby Montjoly. A 3 m specimen was killed by a gendarme in 1918, and two others were shot on the Montjoly beach in June 1972 and January 1975. A distribution map for the species in French Guiana is given in Condamin (1976).

Laws forbidding the hunting of black caiman have not been effective as evidenced by the fact that a local businessman requested permission to export 500 black caiman hides in March 1975 and apparently was not prosecuted. French officials started to seize in-transit shipments, however, and there have not been any more since 1977. Nonetheless, there is a small-scale trade carried out by local hunters, and some animals are killed by Brazilian hunters or by tourists from Cayenne (J. Lescure, personal communication).

Fair numbers of *Caiman crocodilus* hides are shipped from French Guiana to tanneries in France, and the tanners are trying to open a trade in *Melanosuchus* as well. Such shipment is not international

since French Guiana is a department of France rather than a separate country. France has taken a reservation on *Melanosuchus* under CITES so there is every reason to believe that French tanners will continue to push for the French Guiana hides. They are already supplying French Guiana papers on shipments of *Caiman crocodilus yacare* hides originating from Paraguay and U.S. officials are accepting the papers for U.S. imports (F. Wayne King, personal communication).

Stuffed juvenile spectacled caiman are sold in fairly large numbers in the markets and tourist shops of Cayenne, but protective legislation has at least been effective enough to exclude the black caiman from this trade (although the remoteness of the animal's habitat has undoubtedly played a role as well). In three surveys conducted in 1975, 1976, and 1979, we saw hundreds of spectacled caiman but not a single black caiman for sale in the shops in Cayenne.

Although the Kaw population may be one of the largest enclaves of black caiman left in South America, it is a relatively small area and still not a park or reserve. We therefore believe that the black caiman should be considered an endangered species in French Guiana.

GUYANA

In Guyana, the black caiman is found in the Rupununi, the Essequibo, and the Berbice Rivers. The black caiman was apparently once abundant in the Rupununi District, but hide-hunting was a major cause of decline. People from Guyana would apply for permits to hunt, and would then arrange for Brazilian skin dealers in Boa Vista to cross the border and organize local Amerindians to undertake the actual hunting and skinning. A handful of skins were declared to customs officials situated at Lethem, and much larger numbers were sent across the Tacutú River at uncontrolled points. According to a manager of the Balata Company (personal communication to F. Medem), the commercial hunting resulted in such a wholesale slaughter "that the lower reaches of the river at Apoteri stank for weeks, and a large number of piranhas suddenly appeared to feed on the numerous black caiman carcasses." It was impossible to estimate the number of skins shipped out to Brazil, but Medem learned that one hunter obtained 5,000 skins on his own and a second got about 700. As a result of the slaughter, the Guyana government in 1965 declared a total ban on caiman hunting for a period of five years.

We do not have any information on further protection in Guyana, but it is clear that the species has declined drastically in that country and must be considered endangered there.

PARAGUAY

During his 1973 survey in Paraguay, Medem obtained unexpected information suggesting that the black caiman might occur in that country. Although he did not himself see live individuals or hides, two reliable informants told him that they had seen or shot several specimens 3 to 5 m in length, which, by their descriptions (e.g., white ventral surface, size), are unlikely to have been *Caiman crocodilus yacare* or *Caiman latirostris*,

the other caiman species in Paraguay. The localities in question included a stretch between the Ríos Tebicuary and Tebicuary-mi and the village of Villa del Pilar (1950), and the Laguna Ypoá (1965), both within the drainage of the Río Paraguay.

The black caiman is not mentioned by Bertoni (1913) in his list of Paraguayan reptiles, but this is not surprising given what would be a very low black caiman population density in a little-known area. Populations of black caiman exist (or existed) nearby in the upper Río Paraguay of Brazil's Mato Grosso (e.g., São Luis de Cáceres, Corumbá, the town of Mato Grosso), so it would not have been difficult for the animal to enter Paraguay from these areas.

The black caiman would be protected under the general decree (Decreto No. 18.796 of 1975) prohibiting hunting, commercialization, and export of Paraguayan wildlife. Since this law was passed, numerous warehouses of animals have been shut down in the capital city of Asunción. However, one can still buy novelty items such as stuffed *Caiman latirostris* in tourist shops located along Asunción's Calle Colon (Jody Stallings, personal communication). In addition, illegal shipments of caimans from Paraguay still enter the U.S. and Europe (F. Wayne King, personal communication), although it is not known whether *Melanosuchus* has been included among these.

Recently, the Servicio Forestal Nacional has initiated a national biological survey in Paraguay. This project will add to our knowledge of caiman distribution within Paraguay. In any case, if the black caiman does occur, it would definitely have to be considered highly endangered.

PERU

The black caiman was once abundant throughout the upper Amazon drainage in Peru. However, commercial hide-hun-

ting, which started around 1950 took a tremendous toll with at least 26,463 still being exported as recently as 1970 (FAO statistics, quoted in Medem, 1971).

This crocodilian is now on the verge of extinction in Peru, with good populations existing only in the Manú National Park and possibly in the large swampy area on the lower Río Madre de Dios. A population also exists in the Pacaya-Samiria Reserve, largest of Peru's protected areas, and its status is currently being investigated by Pekka Soini.

IMPORTANCE OF THE BLACK CAIMAN IN THE AMAZONIAN ECOSYSTEM

The black caiman is not only important as a wildlife heritage. It also occupied an important position in the Amazonian ecosystem and its elimination in many areas has resulted in a number of ecological changes, some of them affecting man.

Fittkau (1970, 1973), reported that a decline in caiman populations has brought about an accompanying decline of fish populations, apparently due to interruption of a delicate nutrient chain. Caiman excrement in the lakes and swamps apparently provides a key food source for zoo- and phytoplankton, which in turn serve as food for fish fry of the many species that use these lakes as their major egg-laying sites. When the caiman is eliminated, its excrement no longer forms the basis of the nutrient chain, and the fish species dependent on it drop significantly. This effect is accentuated in the nutrient poor "black water" areas where living animals constitute a major portion of the biomass and are responsible for bringing allochthonous nutrients into the partially closed system.

Overexploitation of caimans therefore results not only in the loss of the valuable resource that these animals themselves represent, but also can have a significant effect on fish populations, which are the major protein source for man in Amazonia. (This effect, however, may at

least be partially offset by the increase in the population of spectacled caiman [*C. crocodilus*] that has apparently accompanied the decline of the black caiman [Magnusson, 1980].)

Black caiman also play an important role in control of animals harmful to man's agricultural activities in Amazonia. In Bolivia and part of Brazilian Amazonia, for instance, it is reported that the widespread extermination of the black caiman has resulted in an increase of their herbivore prey species, notably capybara and other smaller rodents, and these have become serious agricultural pests. In some parts of Bolivia, an increase in the piranha (*Serrasalmus* spp.) population was noted, to the extent that these fish frequently attacked cattle crossing the flooded grasslands.

The message here is that black caiman conservation is not only of interest to preservationists. It is also an essential part of habitat management and helps assure the continued usefulness of the Amazonian riverine ecosystem for man.

THE FUTURE OF THE BLACK CAIMAN—KEY CONSERVATION AREAS

Although small populations or isolated individuals undoubtedly hang on in many remote corners of Amazonia (e.g., Magnusson, 1979), few substantial concentrations of black caiman remain. We know with certainty of only three, and briefly discuss them here.

LIMON COCHA, ORIENTE PROVINCE, ECUADOR

Limon Cocha is a lake in Amazonian Ecuador, which harbors a large population of black caiman. During the dry season (December–February), it is possible to see over 100 individuals in several hours of night searching (Ken Miyata, personal communication). Limon Cocha was proposed for ecological reserve status in the late 1970's, but we were un-

able to learn whether any official action has been taken.

MANÚ NATIONAL PARK, DEPARTMENT OF MADRE DE DIOS, PERU

One of the largest, most important and most remote protected areas in Amazonia, the 1,500,000 ha Manú National Park has a large and undisturbed population of black caiman that was estimated to number some 150 adult individuals in the early 1970's (K. C. Otte, personal communication to Medem). In the Manú River, the animals can still be seen during the day, a habit that remaining individuals in heavily hunted areas have abandoned long ago. The Manú *Melanosuchus* were studied by a German ecologist, Dr. K.-C. Otte, but thus far only a brief report has appeared on this work (Otte, 1974).

Otte (1974) also surveyed a number of other rivers in Peru, among them the Sotileja, the Piedras, the Heath, and the Pariamanú. Apart from the upper Río de las Piedras, he did not encounter black caiman in any of these. Based on reports from hunters and skin dealers and from local government statistics, he concluded that the black caiman existed in economic numbers only on the upper courses of the Río Tambopata, Manú, Piedras, and Amigo. Viable populations have recently been seen in lagoons along the Río Tambopata (K. Miyata, personal communication), and Plotkin found evidence of continued hunting of black caiman along the eastern border of Manú Park in July of 1980. It is clear that continued protection of the important Manú National Park population of black caiman is essential to the survival of this animal in Peru.

THE KAW REGION, FRENCH GUIANA

The Kaw region is evidently the last major black caiman concentration in the Guianas, and perhaps even in all of lower Amazonia. The vast swamps to the west of the Kaw River and north of the Monta-

gnes de Kaw are inaccessible, and the Kaw River itself has a very small human population (not exceeding 100 individuals) in the village of Kaw and on two nearby cattle ranches.

A former animal dealer familiar with Kaw estimates that there may still be 1,500 to 2,000 large black caiman in the whole region. However, the French herpetologist J. Lescure (personal communication) believes that 1,000 should be considered a maximum for all of French Guiana. In any case, the Kaw region certainly still harbors some of the 5 m giants common in the past. Towards the end of the dry season, when most of the swamps are without water, a large pond known locally as the "Mare aux Caimans" forms in the uninhabited area to the west and it is said that individuals 6 to 7 m in length can still be seen.

Small individuals are commonly seen at night in the Kaw River. During a brief expedition to this river in July, 1979, Plotkin, Mittermeier, and Constable saw a number of small black caiman one meter in length and under, and a single large adult of about 5 m. Since it was the wet season, most of the larger caiman were dispersed throughout the flooded grasslands, but local informants told us that many concentrated in the river during the drier months.

Black caiman are occasionally hunted for food by the inhabitants of the region, but this is rare and other mammal and avian game are much preferred. A greater threat is the seasonal burning of the dried-out swamps around Kaw, a practice that probably results in the destruction of many caiman nests. Furthermore, the region's inaccessibility will soon end with the construction of a road to Kaw (J. Lescure, personal communication).

The area should obviously be protected, and this was already suggested by M. Condamin, a French zoologist working in the Cayenne branch of ORSTOM, in 1975. In a report entitled "Projets de Reserves Naturelles sur le Littoral Guyanais" (Condamine, 1975, see also

Condamin, 1976), he suggested four areas for reserves along the French Guiana coast. These include Mana (22,000 ha) for sea turtles, Sinnamary (15,000 ha) for scarlet ibis and other coastal birds, Le Grand Connetable (1 ha), a tiny island important for nesting marine birds, and Kaw (49,000 ha) for black caiman and hoatzin. The first three were established, but Kaw still has not because of lack of funds. The Society for Nature Protection in French Guiana (SEPANGUY) and its director, Dr. Leon Sanite, are very concerned that Kaw be fully protected as soon as possible, and we are recommending to the French government that Condamin's plan for Kaw be implemented as soon as possible.

ACKNOWLEDGMENTS

We would like to thank the following people for the information that they provided: J. Dorst, R. Ferrer, R. Foote, C. Freese, P. von Hildebrand, F. W. King, J. Lescure, K. Miyata, P. Soini, and J. Stallings.

APPENDIX

South American legislation concerning *Melanosuchus niger*:

BOLIVIA - Officially protected by Decreto Supremo No. 08063 of August 16, 1967 (Article 20) which imposes a total ban on hunting from July 31 to January 1.

BRAZIL - The 1967 Fauna Protection Law (Lei No. 5.197/67) officially protects all forms of Brazilian wildlife. The black caiman does not appear on the Brazilian List of Endangered Species.

COLOMBIA - Officially protected by a total ban on hunting (Resolution No. 411, July 6, 1969) passed by the Division de los Recursos Naturales of the Ministerio de Agricultura. In 1969, the Instituto de Recursos Naturales Renovables (INDERENA) also declared a ban on black

caiman hunting and egg collection (Resolution No. 573, July 24, 1969).

ECUADOR - The black caiman was not included in the wildlife protection Resolution No. 818 of 1970, but was covered by the total ban on wildlife export declared in 1972.

FRENCH GUIANA - First protected in French Guiana in 1968 (Arrete Prefectoral No. 68-719 ID/2B - January 31, 1975). *Melanosuchus* received stronger protection in 1975 (Arrete Prefectoral No. 172 ID/2B - January 31, 1975).

GUYANA - The government passed a total ban on caiman hunting in 1968 for a period of five years.

PARAGUAY - All wildlife is protected by general decree (Decreto No. 18.796 of 1975) which prohibits hunting, commercialization, and export of Paraguayan fauna.

PERU - The black caiman is officially protected by law No. 21147 which protects all forms of wildlife other than those of importance to local people as food sources.

LITERATURE CITED

- BATES, H. W. 1863. The Naturalist on the River Amazons. London, John Murray.
- BERTONI, A. W. 1913. Fauna paraguaya. Catálogos sistemáticas de los vertebrados del Paraguay. In: Descripción Física y Económica del Paraguay. Asuncion, Moises S. Bertoni.
- CAMPBELL, H. 1977. SSS/IUCN Croc. Gp. Newsl. 12.
- CONDAMIN, M. 1975. Projects de reserves naturelles sur le littoral Guyanais. Unpublished report.
- . 1976. Faunes originales (plaines côtières). Atlas des Départements d'Outre Mer. IV. La Guyane, 1-2. Edit. CNRS. ORSTOM.
- FITTKAU, E. J. 1970. The role of caimans in the nutrient regime of mouth-lakes of Amazon affluents. Biotropica, 2(2): 138-142.
- . 1973. Crocodiles and the nutrient metabolism of Amazonian waters. Amazoniana, IV(1): 103-133.
- GOELDI, E.A. 1898. Die Eier von 13 brasilianischen Reptilien, nebst Bemerkungen ueber Lebens- und Fortpflanzungsweise letzterer. Zool. Jahrb. (Syst.), 10: 640-676.
- HAGMANN, G. 1902. Die Eier von *Caiman niger*. Zool. Jahrb. (Syst.), 16: 405-410.
- MAGNUSSON, W. E. 1979. The distribution of caiman within the Parque Nacional da Amazonia

- (Tapajós): Unpublished report to the National Amazon Research Institute, Manaus.
- . 1980. Report on the status of Amazonian crocodilians. Unpublished report to the IUCN/SSC Crocodile Specialist Group.
- MEDEM, F. 1960. Datos zoo-geograficos y ecológicos sobre los Crocodylia y Testudinata de los ríos Amazonas, Putumayo, y Caquetá. *Caldasia*, 8(38): 341–351.
- . 1963. Osteologia craneal, distribución geográfica, y ecología de *Melanosuchus niger* (Spix), (Crocodylia, Alligatoridae). *Rev. Acad. Colomb. Ci. Exact., Fis., Nat.*, 12(45): 5–19.
- . 1970. El estado actual respecto a la exterminación de los crocodilídeos en la hoya Orinoco Colombiano. Unpublished report.
- . 1971. Situation report on crocodilians from three South American countries. In: Crocodiles. Proc. First Work. Meeting IUCN/SSC Croc. Spec. Group. IUCN Publ. (32): 54–71.
- . 1972. Situation report on South American crocodilians. Unpublished report.
- . 1973. Summary of the surveys of the status of crocodilian species in South America. In: Proc. Sec. Work. Meeting IUCN/SSC Croc. Spec. Group. IUCN Publ. (41): 33–36.
- MELO CARVALHO, J. C. 1967. A conservação da natureza e recursos naturais na Amazônia Brasileira. Atas Simposio Biota Amazônica, Rio de Janeiro, (7): 32–33.
- OTTE, K.-C. 1974. Research programme *Melanosuchus niger* in the Manú National Park. *World Wildlife Yearbook* 1973–1974: 257–260.
- WALLACE, A. W. 1853. A narrative of travels on the Amazon and Rio Negro. London, Reeve and Co.

The Leatherback Sea Turtle, *Dermochelys coriacea*, Nesting in the Dominican Republic

JAMES PERRAN ROSS¹

JOSE ALBERTO OTTENWALDER²

ABSTRACT. An estimated 300 female *Dermochelys* nest each year in the Dominican Republic, and 37% of nestings occur on four beaches. Nearly 100% of nesting females and eggs are taken by local people for food. Favored beaches have no offshore fringing reef and few human inhabitants nearby. Conservation measures need implementation and enhancement.

INTRODUCTION

The leatherback sea turtle *Dermochelys coriacea* is the largest of the sea turtles and is considered to be an endangered species by both the United States Department of the Interior and the International Union for the Conservation of Nature. Leatherbacks nest widely on mainland shores of the tropics, but large concentrations are only reported from Trengganu, Malaysia, French Guiana and Surinam, and Pacific Mexico. Few nesting populations have been adequately surveyed or receive protection (Pritchard, 1971; Ross, 1982).

In the Caribbean there are small populations of leatherbacks nesting in Florida, Vieques, Culebra, the United States Virgin Islands, Trinidad, and Costa Rica. Only the Costa Rican population is thought to number more than a few dozen individuals. Some of these small

populations have been proposed for protection (Dodd, 1978).

Leatherbacks are known to migrate into temperate waters from their tropical breeding grounds. They are commonly reported in Massachusetts waters (Lazell, 1980) and as many as several hundred sightings per year have been made in recent years. (C. R. Shoop, personal communication)

The origin of these leatherbacks is unknown, but there appear to be too few adults in the reported nesting areas to account for the sightings made in temperate waters. It appears likely that there are unreported nesting sites in the Caribbean area.

Recently, a previously unreported nesting site for leatherbacks was reported in the northeastern Dominican Republic (D.R.) on the island of Hispaniola (A. Carr; T. Carr, personal communication). The aim of the present survey was to accurately locate the nesting beaches in the D.R., to estimate how many leatherbacks nest there; to evaluate the activities of people taking eggs and killing adults for food; to initiate contact with the D.R. authorities; and to assess the population for conservation action.

METHODS

To achieve these aims the senior author visited the Dominican Republic between March 24th and April 13th, 1980.

¹ Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, U.S.A.

² Museo Nacional de Historia Natural, Plaza de la Cultura, Santo Domingo, Dominican Republic.

On March 27th, the authors flew in a chartered aircraft around the eastern part of the D.R. to search for beaches and signs of nesting. In the next three weeks we traveled by car to many coastal locations. At each place we talked with local people and examined the beaches for signs of recent nesting and skeletal material.

AERIAL SURVEY

We flew around the coast from Miches to La Romana, maintaining a height of 31 m and a speed of 160 km/hr., positioned slightly offshore of the beach. We noted beaches that appeared suitable for

nesting and looked for tracks, body pits, and carcasses on the beaches. Locations were established using time of flight, a 1:600,000 map, and prominent landmarks.

GROUND SURVEY

Areas surveyed in detail on the ground are shown in Figure 1 and Table 1.

INTERVIEWS

Interviews were conducted in Spanish through a fluent translator. No formal interview sequence was used but conversation led into asking what types of sea

Figure 1. Nesting of *Dermochelys coriacea* in the Dominican Republic. P = Playa; P.d. = Playa del; C. = Cabo (Cape).

turtles occur nearby? Do they nest? Where? How many nest in one night? Are the eggs and meat good to eat? How many are taken? Are turtles more or less abundant than in previous years? Care was taken to maintain neutrality to answers and avoid leading people into statements meant to please us. A total of twenty-four interviews were used as a data base after eliminating those containing clearly erroneous or inaccurate information. The distribution of these informants by area is given in Table 1.

RESULTS

The local name for *Dermochelys* is "Tinglar," similar to names used in Pacific Mexico, "Tinglada" and Peru, "Tinglado."

NESTING SEASON

Only one recent nest was seen, on 30 March at Playa del Muerto in Altagracia province. Local informants generally agreed that most *Dermochelys* nesting takes place from mid-April to mid-June.

DISTRIBUTION OF NESTING

Most coastal people recognized the local name for *Dermochelys*, "Tinglar," and say that they are occasionally encountered. However, in most locations people agreed that *Dermochelys* were only seen from once/year to once/lifetime—in other words, they are uncommon. *Dermochelys* were reported to nest occasionally in very low densities throughout the D.R. wherever suitable sandy beaches occur. Three sections of coastline support more regular nesting although the individuals are widely dispersed. Within these regions four locations were identified that support modest concentrations of nesting *Dermochelys* (see Fig. 1 for locations).

TABLE 1. AREAS SURVEYED AND INTERVIEWS WITH LOCAL INFORMANTS.

Punta Rucia -Luperon	(20 km) — 3 interviews
Puerto Plata - Sosua	(20 km) — 1 interview
Nagua - Las Terrenas	(37 km) — 5 interviews
Miches - Boca Maimón	(64 km) — 6 interviews
Macao	(5 km) — 1 interview
Barahona -Enriquillo	(140 km) — 1 interview
Cabo San Luis	(13 km) — 2 interviews
Cabo Falso - Cabo Rojo	(30 km) — 5 interviews

LOW DENSITY NESTING CONCENTRATIONS OF *DERMOCHELYS*

1. Nagua - Cabo Samaná (North side of Peninsula) (88 km).

There are many fine beaches along the north coast of the Samaná Peninsula. These are difficult to approach by car, but there are many small settlements with almost continuous cultivation of coconuts. The beaches at Las Terrenas and Coson are narrow and backed by coconut palms. Nesting is said to occur throughout the area in low density. Local people estimate from one to two per year up to one per night on various sections of the beach. Particular locations are Matancitas, El Pozo, Las Terrenas, Boca Río Limón (Puerto Escondido), and Galeras (Bahía del Rincón).

2. Miches - Cabo Engaño (94 km).

A series of narrow beaches backed almost entirely by coconut plantations supports low density nesting throughout its length. The beach is accessible by a sandy track through the coconuts. Estimates of local informants range from one to two per year up to 10 to 12 per year of various sections. Included within this section are two concentrations of nesting that are discussed below.

3. Enriquillo - Pedernales (140 km).

This dry area with relatively low density of human population has extensive beaches lying around the tip of the Barahona Peninsula. Two concentrations are reported below. It is highly likely that

there is more *Dermochelys* nesting in this remote area, particularly between Cabo Falso and Punta San Luis.

Other occasional sightings of nesting *Dermochelys* were reported as occurring at Isla Beata, Bahía de Ocoa, and Isla Saona.

HIGH DENSITY NESTING CONCENTRATIONS OF *DERMOCHELYS*

1. *Playa del Muerto* (- de los Muertos) Altagracia province.

Approximately 4 km of excellent nesting beach is located 10 km south of the mouth of the Río Nisibón. The beach is accessible by motor vehicle from the main road at Las Vacamas. The beach is 40 m wide with a wide, well-defined beach platform stabilized by *Ipomea* and grasses. There is no fringing reef, and consequently there is heavy surf on the beach. Behind the beach is a strip of coconut plantation 200 m wide, and behind this is a mangrove swamp. Local informants including two nearby residents agree that this beach is the major concentration of *Dermochelys* between Río Nisibón and Río Maimón. This location corresponds to the location near Nisibón reported by T. Carr (personal communication) who visited it in 1978. Estimates vary from two to six turtles nesting each night in April and May. Knowledgeable informants were properly cautious about estimating annual populations, citing the difficulties of multiple nesting and incomplete observations. When pressed, informants guessed from 40 to 60 turtles nesting each year.

Informants readily admitted to taking all the eggs they found and killing the females whenever they were encountered. Three separate informants agreed that between 12 and 15 *Dermochelys* were killed here in 1979. Examination of the beach revealed several old nest pits and bones from two separate individuals. A freshly killed carcass, 4 to 10 days old,

was found on the beach with plastron, limb bones, and meat removed.

Nesting at lower densities is reported on the narrow beaches between Río Nisibón and Río Maimón (approx. 32 km).

2. *Playa Macao* Altagracia province. Approximately 5 km of excellent nesting beach lies south of Punta Macao. The track to Cabo Engaño lies 250 m inland of the beach separated from it by low dunes and scrub. The beach is 40 m to 120 m wide with a high beach platform stabilized by dune vegetation. Local informants report that 3 to 4 *Dermochelys* can be seen each night during April and May and that an experienced collector working on foot from El Macao can collect 500 eggs in one night. Local people claim to only occasionally kill females for meat. Examination of 2.5 km of the beach did not reveal any recent nesting or turtle specimens.

Dermochelys nesting is said to occur in lesser density on the narrower beaches southward to Cabo Engaño (approx. 20 km).

3. *Playa San Luis* Pedernales province. Approximately 15 km of nesting beach lie between Cabo San Luis and Cabo Inglesa. There is a narrow strip of palm scrub behind the beach, then a large saline lagoon that effectively isolates the beach. Access is difficult, on foot or across the lagoon by small boat from the main road at Oviedo Viejo. This location is the site of a new tagging and conservation program of the Museo Nacional. Under the direction of Ottenwalder, a local resident patrols about 5 km of the beach each night and is instructed to tag and measure all *Dermochelys*. This person estimates that between 1 and 3 *Dermochelys* come ashore most nights in April and May. In 1979 he observed 15 nesting turtles. Recommendations for enhancing the effectiveness of this program are given below.

4. *Playa Aguilas* Pedernales province. About 5 km of remote beach is located 9 km south of Cabo Rojo in Bahía de las Aguilas. Another small beach (*Playa Majer* or *Playa Caliente*) is said to be further south. The beach lies in a sparsely inhabited region and is accessible by a very rough track from Cabo Rojo or by boat from the nearby fishing village of Cueva. Fishermen at Cueva (population about 80) report that one to two *Dermochelys* can be seen any night in April–June. I found parts of two old carcasses butchered for meat, and fishermen say they regularly take the eggs. The beach is rather narrow (5–15 m) but slopes steeply up to a stabilized dune platform.

Topographical details for these areas can be seen on U.S. Army Topographic maps: Dominican Republic 1:50,000 Series E 733 Ed. 2. Sheets: 64721, 647211, 6572111, and Ed. 3. Sheets: 58691, 586911, 5969111.

PREDATION BY PEOPLE

Predation by people on the eggs and nesting females of *Dermochelys* is common throughout the Dominican Republic. A striking feature of most of the coastline is the high density of the human population. Every single beach visited showed signs of at least weekly visitation, and most beaches are patrolled daily by people who live nearby. As a result a very high proportion of eggs of all turtle species are found and taken. The eggs are both for home consumption and for sale.

People mark the location of nests and return between 10 and 30 days later in an attempt to intercept the female on a later nesting visit. The females are slaughtered without regard for whether they lay or not, and the meat is transported to nearby villages for sale. *Dermochelys* eggs retail for 30¢ (U.S.) each and the meat for \$1/lb. and thus represent a sizeable income for the vendor.

Coastal dwellers frequently claimed to kill all the *Dermochelys* they encounter-

ed. All the *Dermochelys* specimens found in this study (five individuals) appeared to have been killed for meat. The vigor of persecution seems to be most intense on the east coast in the region of Nisibíson and more casual at Macao, in the Samaná Peninsula and in the southwest. However, exact data are lacking. The taking of eggs and the killing of nesting females constitute the most serious threat to the survival of *Dermochelys* in the Dominican Republic.

Skin divers, who prey heavily on hawksbill turtles (*Eretmochelys imbricata*), claim to rarely encounter *Dermochelys* and do not spear them because they are too large to handle.

Accidental capture in fisheries aimed at other species was not reported and appears to be negligible.

POPULATION ESTIMATES

Population numbers of sea turtles are notoriously difficult to assess as several of our informants were aware. The following estimates are included to give 'order of magnitude' understanding of the importance of D.R. nesting beaches and to show the relative importance of different areas. In the absence of nightly data on nesting at the height of the season, these estimates are based on information from informants and are consequently of low reliability. We believe most informants overestimate numbers of turtles because they remember occasional nights of heavy nesting and do not calculate in many nights when few turtles nest. On the other hand, individuals' estimates are usually limited to short sections of coast that they know well and thus estimates from distant informants can be summed. The highly dispersed low density of nesting in much of the D.R. makes population assessment even more difficult.

Population numbers are estimated in two ways. We take the estimated number of turtles nesting each night and, assuming a season 60 days long, and that each

turtle nests three times, extrapolate to an annual nesting population. We also give the informants' estimates of annual nesting. These data are shown in Table 2.

The estimates show that each year perhaps 300 *Dermochelys* nest in the Dominican Republic. The estimate is highly tentative and needs refining by further research (see recommendations).

A considerable proportion of the estimated nesting is concentrated on four beaches.

TURTLE CONSERVATION IN THE DOMINICAN REPUBLIC

PROBLEMS

The major problem for the continued existence of *Dermochelys* nesting populations in the D.R. is the uncontrolled taking of eggs and killing of nesting females for food by local people. The problem is rendered complex because the rural population is very poor, widely dispersed, and uniformly dense. Turtles, including *Dermochelys*, constitute an important cash source for some members of coastal communities.

A secondary problem is the development of isolated beaches as tourist resorts

with a consequent increase in back lighting and disturbance.

Laws pertaining to wildlife and fisheries conservation are contained in several Presidential decrees of the Dominican Republic. These are given in detail in Ross (1980). In summary, there are provisions for closed seasons and lower size limits for sea turtles. However, this body of legislation has some deficiencies and is inadequate for protection of all sea turtles, particularly *Dermochelys*. Nomenclature in the legislation is unclear, and there is little biological justification for the size limit and closed season. All of our informants agreed that there is no policing of these regulations.

This situation is currently under review and revised protective legislation is pending implementation.

RESEARCH IN PROGRESS

There is a small but active group of researchers, headed by Ottenwalder, that is working to improve conservation in the D.R. and is collecting data. The Museo Nacional de Historia Natural (MNHN) is designing a series of aerial surveys to evaluate sea turtle nesting. MNHN has begun a small tagging and protection project on *Dermochelys* at Playa San Luis. A

TABLE 2. FIRST APPROXIMATION NESTING POPULATION ESTIMATES. FEMALE *DERMOCHELYS CORIACEA* IN THE DOMINICAN REPUBLIC.

Location	Informants		Estimated annual
	nightly	annual*	
Playa del Muerto	2-6	40-60	40-120
Playa Macao	3-4	100	60-80
Playa San Luis	1-3	45	20-60
Playa de Aguilas	1-2	—	20-40
Miches - Cabo Engaño (excluding concentrations above)	—	2-12	20
Nagua - Cabo Samaná	4	less than 100	80
Dispersed other	—	—	240
			60 (?)
			300

*The range of values estimated by local informants is given. See Table 1 for distribution of informants.

local man patrols the beach nightly, tagging turtles and discouraging people from taking eggs or killing turtles. These studies are greatly handicapped by an almost total lack of funding and suffer because the few trained and interested personnel in the D.R. need to spread their activities over several conservation projects in addition to their routine work. Nevertheless there are promising beginnings.

CONSERVATION PLANNED

A pan-Caribbean turtle conservation symposium is planned for 1983 in San Jose, Costa Rica sponsored by IOCARIBE and NOAA. It is hoped that this conference will stimulate participating nations to establish an expanded data base on sea turtles.

DISCUSSION

The documentation of a large number of *Dermochelys* nesting in the Dominican Republic is a useful addition to our knowledge of the distribution of this species, although further work is necessary to improve the estimate of numbers. It is likely that some of the leatherbacks nesting in the D.R. migrate up the U.S. coast to temperate waters. Tagging programs in the D.R. will provide a means of verifying this postulated pathway.

Within the three general areas where *Dermochelys* are more common, there are four beaches where concentrations of nesting occur. These beaches share two factors that appear to make them more suitable for *Dermochelys*. All are backed by infertile land with reduced human population density. Playa del Muerto and Playa Macao are backed by dunes and mangrove swamp, Playa San Luis is backed by a saline lagoon and Playa Aguilas by barren limestone karst. This may serve to reduce the regularity of disturbance and predation by local people.

Each of the beaches where *Dermochelys* nest in greater density lacks an off-

shore fringing reef and has heavy surf. As a result the beaches are wide, rise steeply to stable beach platforms, and provide unimpeded access to the sea. In contrast, much of the remainder of the D.R. coastline has a fringing reef 100 to 500 m offshore, with beaches that are uniformly narrow (5–10 m) with coastal vegetation forming a dense barrier at the top. Much of this coastline supports a strip of coconut palm plantations and the dense root masses of the coconut trees form a barrier just above the high tide line.

Hawksbill turtles (*Eretmochelys imbricata*) nest intermittently along the narrow beaches, although their numbers have been greatly reduced in recent years. However, the narrow low profile beaches are apparently poorly suited to the nesting requirements of *Dermochelys*. Sand quality seems to be relatively unimportant to nesting turtles within broad limits (Stancyk and Ross, 1978). However, the extremely large *Dermochelys* does require more open, unimpeded beaches than the smaller, more agile *Eretmochelys*, a conclusion similar to that of Hendrickson and Balasingam (1966). Unfortunately, remote beaches of the sort favored by sea turtles are in great demand for resort developments. Two such developments were noted in the aerial survey. Care should be taken that similar development does not occur on the very few beaches where concentrated *Dermochelys* nesting occurs.

The widely dispersed nature of *Dermochelys* nesting in much of the D.R. and the complex socio-economic problems surrounding the use of turtles by people will make conservation of *Dermochelys* difficult. However, the small number of nesting concentrations provides an opportunity for effective action. We suggest that conservation activities be focused on the four important nesting beaches: Playa del Muerto, Playa Macao, Playa San Luis, and Playa Aguilas. We also recommend the present MNHN survey and tagging program should be enhanced to accurately assess the numbers and distri-

bution of nesting turtles. The basic requirement is money to pay salaries and provide very basic equipment. A limited amount of expert advice on the expansion of the tagging program will ensure that useful biological data are accumulated.

Aerial surveys would be of limited use in D.R. In addition to the usual problems of ground truth calibration and observer error, the narrow, littered and overgrown beaches make accurate observations of nesting difficult.

A large scale public relations and educational program on endangered sea turtles needs developing in rural areas. Something like the WWF mobile audio visual center would be ideal, as most coastal villages are accessible by road, have electric power, and lack competing entertainment on most evenings. Such a program should be developed by Dominican personnel, perhaps as an extension service of the National Zoo which has already assumed a burden of public education on natural resources.

As a further conservation measure, it may be possible to buy fresh *Dermochelys* eggs collected from areas of diffuse nesting and hatch them from natural beach hatcheries at a central protected location.

ACKNOWLEDGMENTS

Ernest E. Williams's early published work concerned the taxonomy and systematics of turtles, including the *Dermochelyidae* (Williams, 1950). Turtles continued to attract his interest, resulting in over 20 papers and articles spanning 17 years. In reference to the extinct tortoises of the West Indies he wrote, "... much remains to be learned about the West Indian fauna" (Williams, 1952), and he has continued to show a lively interest in

turtles of the Caribbean region. His concerns extend beyond systematics to the problem of conservation of the present fauna, and in early 1980 he encouraged the senior author to visit the Dominican Republic to assess the status of populations of *Dermochelys coriacea* reported to be nesting there.

This work was carried out under contract to the U.S. Fish and Wildlife Service. Additional support was received from Ocean Research and Education Society, Chelonia Institute, Fauna Preservation Society and Ernest E. Williams. We thank K. Musik for assistance with translation in the field.

LITERATURE CITED

- DODD, C. K. 1978. Terrestrial critical habitat and marine turtles. *Bull. Maryland Herpetol. Soc.*, 14(4): 223-240.
- HENDRICKSON, J. R., AND E. BALASINGAM. 1966. Nesting beach preferences of Malayan sea turtles. *Bull. Nat. Hist. Mus. Singapore*, 33: 69-76.
- LAZELL, J. D., JR. 1980. New England waters: critical habitat for marine turtles. *Copeia*, 1980: 290-295.
- PRITCHARD, P. C. H. 1971. The leatherback or leathery turtle. IUCN Publications, monogr. 1: 39 pp. Switzerland, Morges.
- ROSS, J. P. 1980. Report on the leatherback sea turtle (*Dermochelys coriacea*) nesting in the Dominican Republic. U.S. Fish and Wildlife Service, mimeo report, 17 pp.
- . 1982. Historical decline of loggerhead, ridley and leatherback turtles, pp. 189-195. In Bjørndal, K. A. (ed). *Biology and Conservation of Sea Turtles*. Washington, D.C., Smithsonian Institution Press.
- STANCYK, S. E., AND J. P. ROSS. 1978. An analysis of sand from green turtle nesting beaches on Ascension Island. *Copeia*, 1978(1): 93-99.
- WILLIAMS, E. E. 1950. Variation and selection in the cervical central articulations of living turtles. *Bull. Am. Mus. Nat. Hist.*, 94(9): 509-561.
- . 1952. A new fossil tortoise from Mona Island, West Indies and a tentative arrangement of tortoises of the world. *Bull. Am. Mus. Nat. Hist.*, 99(9): 545-559.

SYSTEMATIC INDEX

PHYLUM PROTOZOA

- Plasmodium balli* 458, 460
- colombiense* 456, 466
- floridense* 456, 458
- mexicanum* 457
- minasense* 458
- sp. 458
- tropiduri* 456, 458, 460

PHYLUM PLATYHELMINTHES

- helminths 456

PHYLUM ARTHROPODA

- Aculeata 605
- Anthophoridae 605
- ants 517, 603
- Apidae 604
- Apis cerana* 609
- mellifera* 609
- arthropods 445, 451, 510
- Blattidae 605
- Blattodea 604
- Coleoptera 278, 609
- crickets 445
- Dermaptera 604
- Diptera 609
- Drosophila* 506
- Embioptera 604
- Eumenidae 605
- Formicidae 605
- Habrobracon juglandis* 609
- serinopae* 609
- Halictidae 605
- Hemideina crassidens* 615
- Hemiptera 609
- Hymenoptera 604, 609
- insects 596
- Isoptera 604
- Kalotermitidae 605
- Lasioglossum* 608
- Lepidoptera 609
- Lutzomyia vextrix* 457
- Luzomya trinidadensis* 459
- Mantodea 604
- Megachile rotundata* 609
- Megachilidae 605
- Mormoniella vitripennis* 609
- Myrmecia* 604
- Nothomyrmecia* 604
- Orthoptera 604, 609
- Parasitica 604
- Phasmatodea 605
- Rhytidoponera* 604

- roaches 604
- Symphyta 604
- Vespidae 604

PHYLUM CHORDATA

SUBPHYLUM VERTEBRATA

SUPERCLASS AGNATHA

- acanthodians 223
- placoderms 223

SUPERCLASS GNATHOSTOMATA

SERIES PISCES

- crossopterygians 223
- fish 517
- Gambusia affinis* 691
- Ichthyostegalia 223
- labyrinthodonts 223
- Lepisosteus* 223
- paleoniscoids 223
- Serrasalmus* 702
- sharks 223
- teleosts 223

SERIES TETRAPODA

CLASS AMPHIBIA

ORDER GYMNOPHIONA

- Gymnophiona 186

ORDER URODELA

FAMILY PLETHODONTIDAE

- Aneides lugubris* 9
- Batrachoseps* 11
- Batrachoseps campi* 11
- wrighti* 11
- Bolitoglossa* 2, 5, 8
- Bolitoglossa cuchumatana* 5
- hartwegi* 5
- sp. 5
- Chirotroterotriton* 2
- Hydromantes* 11
- Lineatriton* 2, 9
- Nyctanolis*, **gen. nov.** 2, 11
- Nyctanolis pernix*, **sp. nov.** 2
- Oedipina* 2, 9
- Parvimolge* 2
- Pseudoeurycea* 2
- Thorius* 2

ORDER ANURA

FAMILY BUFONIDAE

- Ansonia* 419
- Ansonia megregori* 420
- mulleri* 420
- Bufo biporcatus* 419

- kisoloensis* 352
lemur 104
marinus 419
punctatus 400
Bufonidae 426
Pelophryne albotaeniata 419
brevipes 419
lighti 419
FAMILY CENTROLENIDAE
 Centrolenidae 426
FAMILY DENDROBATIDAE
 Dendrobatidae 426
FAMILY DISCOGLOSSIDAE
Barbourula busuangensis 418, 427
FAMILY HYLIDAE
Hyla arenicolor 400
bogerti 35
carnifex 35
charlesbogerti 35
columbiana 33, 40, 46
crucifer 400
labialis 44, 46
leucophyllata 46
marmorata 46
microcephala 46
minuta 36, 46
parviceps 36, 46
praestans, **sp. nov.** 43
variabilis 40
 Hylidae 426
Smilisca 48
FAMILY LEOPELMATIDAE
 Leiopelmatidae 393
FAMILY LEPTODACTYLIDAE
Eleutherodactylus 33, 110
Eleutherodactylus anolirex, **sp. nov.** 52
antillensis 104
briceni 53, 56
cochranae 104
devillei 56
schwartzi 104, 111
supernatis 52, 56
unistrigatus 54
vertebralis 52, 56
 Leptodactylidae 426
Leptodactylus albilabris 104
Syrrophus guttilatus 400
marnockii 402
FAMILY MICROHYLIDAE
Chaparina fusca 426
Kalophrynus pleurostigma 426
Kaloula baleata 426
conjuncta 426
picta 416, 426
rigida 426
 Microhylidae 426
Oreophryne annulata 426
nana 426
FAMILY PELOBATIDAE
Leptobranchella mjobergi 27
Leptobranchium 419
Leptobranchium gracilis 23
hasselti 16, 20, 420
hendricksoni 19
minimum 26
montanum 16
nigrops 18
oshanense 26
pelodytoides 26
pullum 16, 21
sp. 21
Leptolalax 13, 31
Megophrys 30, 419
Megophrys minor 30
monticola 420
Scaphiopus bombifrons 30
FAMILY PIPIDAE
 Pipidae 426
Xenopus laevis victorianus 352
ruwenzoriensis 352
vestitus 352
wittei 352
FAMILY RANIDAE
Arthroleptis adolfifriederici 353
Cardioglossa cyaneospila 354
cyaneospila inornata 355
Micrixalus mariae 418, 422
Ooeidozyga diminutiva 422
laevis 421
Phrynobatrachus acutirostris 353
asper 353
bequaerti 352, 353
graueri 353
pedropedetoides 352
versicolor 352
Platymantis 425
Platymantis cornutus 418, 422
corrugatus 421
dorsalis 422
guentheri 422
hazela 422
ingeri 422
insulatus 422
laevigatus 422
lawtoni 422
polilloensis 418, 422
subterrestris 418, 422
Rana angolensis chapini 353
cancrivora 421
clamitans 488
diwata 422
erythraea 420
everetti 421
leytensis 421
limnocharis 421
magna 421
melanomenta 422
micrixalus 418, 422
microdisca parva 420
nicobarensis 420
piapiens 400
ruwenzorica 353
sanguinea 420
signata 421

- woodworthi* 422
 Ranidae 427
Schoutedenella globosa 354
 hematogaster 354
 pyrrhoscelis 354
 schubotzi 354
Staurois natator 421
FAMILY RHACOPHORIDAE
Afraxalus laevis 353
 orophilus 353
Callixalus pictus 353
Edwardtayloria picta 426
 spinosa 426
Hyperolius castaneus 352
 castaneus submarginatus 355
 chrysogaster 353
 discodactylus 353, 355
 frontalis 353
 leucotaenius 353
 leucotaenius allogynus 355
 xenorhinus 358
Leptopelis kivuensis 352
Philautus 426
Philautus acutirostris 422
 alticola 418, 426
 aurifasciatus 422
 emembranatus 422
 leitensis 422
 lissobranchius 421
 schmackeri 421
 sp. 421
 surdus 422
 williamsi 418, 426
Polypedates leucomystax 416, 426
 macrotis 426
Rhacophorus appendiculatus 426
 everetti 426
 hecticus 418, 422
 pardalis 426
 zamboangensis 426
CLASS REPTILIA
SUBCLASS ANAPSIDA
ORDER COTYLOSAURIA
Captorhinus 191
Eocaptorhinus 192
ORDER TESTUDINES
Cryptodira 179, 191
Pleurodira 179, 180, 191
FAMILY CHELIDAE
 Chelidae 58
Chelodina 160, 170, 179, 181
Chelodina expansa 179
 longicollis 180
 novaequinae 174, 179
 oblonga 160, 180
 rugosa 179
 steindachneri 180
Chelus 59, 180, 184
Chelus fimbriatus 179, 431
Elseya 66, 160, 177
Elseya dentata 169
 latisternum 170
 novaequinae 170
Emydura 66, 169, 186
Emydura australis 171
 dentata 170
 krefftii 171
 latisternum 177, 179
 macquaria 170, 178
 subglobosa 170, 176
Hydraspis geoffroyana 59, 70
Hydromedusa 59, 180
Phrynos 59, 186
Phrynos dahl 59, 70
 geoffroyana 60
 geoffroyana geoffroyana 59, 70
 geoffroyanus 58, 179, 184
 geoffroyanus geoffroyanus 59
 geoffroyanus ssp. 60
 geoffroyana 59
 gibbus 59, 65, 70, 179, 181, 184, 430
 hilarii 59, 65, 69
 hoge 59, 65, 70
 nasutus 59, 70, 160, 430
 rufipes 59, 65, 70
 tuberculatus 59, 70
 tuberosus 59, 70
 vanderhaegei 59, 65, 70
 wermuthi 59, 70
 williamsi, **sp. nov.** 59, 65, 70
Platemys 59, 179, 181
Platemys geoffreyana 59, 70
 geoffroyana 59
 platycephala 160, 179, 184, 429
Pseudemydura 66, 179
Pseudemydura umbrina 169, 178
FAMILY CHELONIIDAE
Caretta caretta caretta 150
Chelonia mydas 186
Eretmochelys imbricata 440, 710, 712
FAMILY DERMOCHELYIDAE
Dermochelys coriacea 440, 706
FAMILY EMYDIDAE
Chrysemys picta 150
 scripta ssp. 400
Clemmys guttata 186
Ocadia sinensis 186
Pseudemys dorbignyi 69
 scripta elegans 150
FAMILY GLYPTOPSIDAE
Glyptops 192
Mesochelys 192
FAMILY KINOSTERNIDAE
Kinosternon hirtipes 186
 hirtipes murrayi 400
FAMILY PELOMEDUSIDAE
Dacquemys 75
Erymnochelys 76, 160
Erymnochelys madagascariensis 75
Eusarkia rotundiformis 160
Kenyemys, **gen. nov.** 74
Kenyemys williamsi, **sp. nov.** 75
Latisternon microsulca 79

Pelomedusa 76, 160, 181
Pelomedusa subrufa 161, 179
Pelusios 78, 159, 179, 181, 184
Pelusios adansonii 161
 bechuanicus 159, 163
 bechuanicus bechuanicus 161
 upembae 162
 blanckenhorni 163
 carinatus 162
 castaneus 159, 166
 castaneus castaneus 159, 162
 castanoides 162
 chapini 159
 derbianus 159
 kapika 159, 162
 lutescens 162
 rhodesianus 159
 williamsi 162
 derbianus 179
 dewitzianus 163
 gabonensis 161
 nanus 161
 niger 161
 rhodesianus 162
 rudolphi 161, 163
 rusingae 161
 sinuatus 160, 162
 sinuatus leptus 160
 zuluensis 160
 subniger 161
 williamsi lutescens 159
 williamsi 159
Podocnemis 69, 160, 179, 181, 184
Podocnemis aegyptiaca 79
 antiqua 75
 blanckenhorni 75
 bramlyi 77
 congolensis 75
 fajumensis 75, 79
 madagascariensis 75
 podocnemoides 79
 sextuberculata 65, 69, 78
 unifilis 78, 179
 venezuelensis 75, 160
Shweboemys antiqua 75
Stereogenys cromeri 80
 libyca 79
Sternothaerus derbianus 160
Taphrosphys congolensis 75
FAMILY PLATYSTERNIDAE
Platysternon megacephalum 186
FAMILY PROGANOCHELYIDAE
Proganochelys quenstedti 190
Stegochelys dux 190
Triassochelys 191
FAMILY TESTUDINIDAE
Geochelone 538
Geochelone carbonaria 101, 111
FAMILY TRIONYCHIDAE
Trionyx 181, 186
Trionyx ater 186

SUBCLASS LEPIDOSAURIA
ORDER RHYNCHOCEPHALIA
FAMILY SPHENODONTIDAE

Homoeosaurus 614
Meyasaurus 376, 382
Opistias 614
Palacrodon 614
Polysphenodon 614
Sphenodon 186, 196
Sphenodon punctatus 201, 224, 613
Sphenodontidae 393
Toxolophosaurus 614

ORDER SQUAMATA
SUBORDER EOLACERTILIA

Cteniogenys 373, 376
Fulengia 367
Icarosaurus 367
Kuehneosaurus 367
Kuehneosuchus 367
Palaeagama 367
Paliguana 367
Saurosternon 367

SUBORDER LACERTILIA
FAMILY AGAMIDAE

Acanthosaura armata 209
 crucigera 209
 lepidogaster 209
Agama 208, 368, 380
Agama agama 209
 anchietae 209
 atra 209
 caudospinosa 209
 doriae 209
 hartmanni 209
 hispidia 209
 kirkii 209
 mossambica 209
 planiceps 209
 rueppelli 209
 sankaranica 209
 weidholzi 209
Amphibolurus barbatus 209
 caudicinctus 209
 clayi 209
 cristatus 209
 diemensis 209
 femoralis 209
 fionni 209
 fordi 209
 inermis 209
 isolepis 209
 maculatus 209
 microlepidotus 209
 minimus 209
 minor 209
 muricatus 208, 209
 nobbi 209
 ornatus 209
 pictus 209
 reticulatus 209
 rufescens 209

- scutulatus* 209
Aphaniotis 203
Aphaniotis acutirostris 209
fusca 209
Aporoscelis 208
Brachysaura minor 209
Bronchocela cristatellus 209
jubatus 209
marmoratus 209
Caimanops amphiboluiroides 210
Calotes calotes 210
elliotti 210
emma 210
jerdoni 210
maria 210
mystaceus 210
nigrilabris 210
rouxii 210
versicolor 210
Ceratophora stoddartii 210
tennentii 210
Chelosania brunnea 210
Chlamydosaurus 208
Chlamydosaurus kingii 200, 210
Cophotis ceylanica 210
sumatrana 203, 210
Coryophophylax subcristatus 210
Dendragama boulengeri 210
Diporiphora australia 210
bilineata 210
winneckei 210
Draco blanfordii 210
fimbriatus 210
formosus 210
lineatus beccarii 210
maculatus 210
maximus 210
quinquefasciatus 210
spilopterus 210
taeniopterus 210
volans 210
Gonocephalus bellii 210
beyschlagi 210
chamaeleontinus 210
godeffroyi 438
grandis 210
interruptus 210
liogaster 210
miotympanum 210
robinsonii 210
semperi 210
sophiae 210
Harpesaurus 203
Hydrosaurus 195, 204
Hydrosaurus amboinensis 210
pustulatus 204, 210
weberi 210
Hylagama borneensis 210
Hypsilurus (Arua) auritus 210
modestus 210
(Hypsilurus) binotatus 210
boydii 210
bruijni 210
dilophus 210
godeffroyi 210
nigrigularis 210
papuensis 210
spinipes 210
Japalura brevipes 210
dymondi 210
flaviceps 210
planidorsata 210
polygonata 210
splendida 210
swinhonis 210
tricarinata 210
variegata 210
yunnanensis 210
Lophocalotes ludekingi 210
Lophognathus 208
Lophognathus gilberti 210
longirostris 210
temporalis 210
Lyriocephalus scutatus 210
Mictopholis austeniana 203
Mimeosaurus 368, 379
Moloch horridus 210
Oriocalotes paulus 210
Otocryptis wiegmanni 210
Phoxophrys borneensis 210
nigrilabris 210
Phrynocephalus 203
Phrynocephalus arabicus 210
axillaris 210
clarkorum 210
forsythii 210
guttatus 210
helioscopus 210
interscapularis 210
luteoguttatus 210
maculatus 210
mystaceus 210
ornatus 210
przewalskii 210
reticulatus 210
rossikowi 210
scutellatus 210
versicolor 210
vlangalii 210
Physignathus 195, 203
Physignathus cocincinus 204, 206, 210
gilberti 208
lesueurii 201, 210
longirostris 208
temporalis 208
Psammophilus blanfordianus 210
dorsalis 210
Pseudocalotes 203
Pseudocalotes floweri 210
tympanistriga 210
Ptyctolaemus gularis 210
Salea 203
Salea anamallayana 210
horsfieldii 210

kakhiensis 210
Sitana 337
Sitana ponticeriana 210
Stellio 208, 368
Stellio agorensis 210
 atricollis 210
 caucasica 210
 cyanogaster 210
 erythrogaster 210
 himalayana 210
 lehmanni 210
 melanura 210
 nupta 210
 stellio 210
 stoliczkana 210
 tarimensis 210
 tuberculata 210
Thaumatorhynchus brooksi 203
Tinosaurus 368, 379
Trapelus (Pseudotrapelus) sinaita 210
 (*Trapelus*) *agilis* 210
 flavimaculata 210
 impalearis 210
 mutabilis 210
 persicus 210
 rubrigularis 210
 runderata 210
 savignii 210

Tympanocryptis lineata 210
tetraporophora 210
Xenagama batillifera 208, 210
taylori 210

FAMILY AIGIALOSAURIDAE

Proaigialosaurus 386

FAMILY AMPHISBAENIDAE

Amphisbaena fenestrata 111

FAMILY ANGUIDAE

Anguidae 371, 389

Anguis 371

Anguis fragilis 201

Anniella 371

Barisia 5

Eoglyptosaurus 371

Gerrhonotus 487

Gerrhonotus multicarinatus 222

Glyptosaurus 371

Helodermoides 371

Melanosaurus 371, 387

Odaxosaurus 371

Ophisaurus 371

Ophisaurus apodus 201

Paraglyptosaurus 371

Peltosaurus 371

Placosaurus 371, 387

Xestops 371, 387

FAMILY ANGUOIDEA

Ilaerdesaurus 376

FAMILY ARDEOSAURIDAE

Ardeosauridae 376

Ardeosaurus 368

Eichstaettisaurus 368

Yabeinosaurus 368

FAMILY ARRETOSAURIDAE

Arretosaurus 368

FAMILY BAVARISAURIDAE

Bavarisauridae 376

Bavarisaurus 368

Palaeolacerta 368

FAMILY CHAMAELEONIDAE

Bradypodion 203

Bradypodion pumilus 209

ventralis 209

Brookesia 203

Brookesia stumpffi 209

superciliaris 209

Chamaeleo 201, 203, 368

Chamaeleo adolfifriederici 354

africanus 209

bitaeniatus 209

bitaeniatus ellioti 354

brevicornis 209

carpenteri 354

chamaeleon 209

fallax 209

gracilis 209

jacksonii 209

johnstoni 354

lateralis 209

namaquensis 209

oustaleti 209

owenii 209

parsonii 200

rudis 354

rudis schoutedeni 355

tigris 209

verrucosus 209

xenorhinus 354

Rhampholeon 203

Rhampholeon boulengeri 209

spectrum 209

FAMILY CORDYLIDAE

Cordylidae 386

Cordylus giganteus 196

Gerrhosaurus 370

Pseudolacerta 370

FAMILY DIBAMIDAE

Dibamidae 382

FAMILY DORSETISAURIDAE

Dorsetisauridae 370

FAMILY EUPOSAURIDAE

Euposauridae 368, 379

FAMILY GEKKONIDAE

Cnemaspis quattuorseriatus 354

Cyrtodactylus 276

Gecko japonicus 200

Gekko 230

Gekkonidae 369, 381

Hemidactylus mabouia 100, 111, 115

Hoplodactylus duvauceli 618

maculatus 618

Platydactylus japonicus 200

Sphaerodactylus altavelensis 86

altavelensis altavelensis 89

brevirostratus 93

- enriquilloensis* 91
lucioi, **ssp. nov.** 94
asterulus 95, 97
brevirostratus 89
brevirostratus brevisrostratus 93
enriquilloensis 91
cinereus 95, 97
cryptus 86
difficilis 86, 96, 97
difficilis randi 89
elegans 97
macrolepis 110
macrolepis macrolepis 115
notatus 86
nycteropus 86
ocoae 86
parthenopion 111
randi 86, 89
shrevei 97
williamsi, **sp. nov.** 96
zygaena 86
Thecadactylus rapicaudus 111
FAMILY GYMNOPTHALMIDAE
 Gymnophthalmidae 369, 383
FAMILY HELODERMATIDAE
Eurheloderma 371, 389
Heloderma 371, 375, 389
Heloderma suspectum 200
Paraderma 371, 389
FAMILY IGUANIDAE
Aciprion 368, 378
Agama marmorata 119
Amblyrhynchus cristatus 209
Anisolepis grillii 127
undulatus 120, 209
Anolis 5, 33, 203, 245, 284, 363, 365, 406, 472, 499,
 507, 514, 626, 654, 672, 691
Anolis acutus 480, 629, 638
aeneus 451
aequatorialis 259, 270
aliniger 511
allisoni 478
allogus 479
angusticeps 209, 288, 482
annectens 274, 277
auratus 257, 274, 278, 456, 466
baleatus 285, 288
barkeri 259, 276
bimaculatus 209, 288, 505, 627, 635
biporcatus 285
boettgeri 406
bombiceps 406
brevirostris 552
carolinensis 209, 288, 456, 482, 491, 627, 638, 656
chlorocyanus 511
christophei 511
chrysolepis 247, 257, 274, 276, 285, 288
chrysolepis planiceps 209, 295
scypheus 407
cochranae 511
coelestinus 209, 288, 511, 513
conspersus 624
cooki 479, 626, 637
cristatellus 102, 109, 209, 273, 473, 484, 504, 626,
 629
cristatellus cristatellus 107
wileyae 101, 107, 115
cupreus 451
cuvieri 101, 246, 252, 273, 626, 637
cybotes 280, 288, 478, 511, 513, 552, 627, 637
desechensis 108, 109
dissimilis 407
distichus 294, 482, 511, 513, 552, 627, 635
distichus dominicensis 553, 563
equestris 209, 288, 515
ernestwilliamsi, **sp. nov.** 102
etheridgei 511
eugengrahami 276
evermanni 273, 474, 627, 629
fuscoauratus 209, 285, 288
fuscoauratus fuscoauratus 407
gadovi 629, 638
garmani 285, 571, 624
gingivinus 499
grahami 209, 482, 571, 621, 629, 638
grahami aquarum 621
grahami 621
gundlachi 474, 626, 637
hendersoni 294, 511, 513
homolechis 479
koopmani 294, 512
krugi 627, 629
laevis 407
latifrons 209, 288
leachi 515
limifrons 441, 455, 511, 558, 566
lineatopus 571, 624
lucius 479
macrolepis 276
marcanoi 552
marmoratus 478
meridionalis 127
monensis 108, 109, 481, 628, 637
monticola 294, 511, 513, 624
nebulosus 285, 288
occultus 247, 252, 257, 274, 626, 637
oculatus 473, 478, 629, 637
onca 209, 274, 276, 285, 288
opalinus 559, 577, 624
ophiolepis 288
ortoni 407
oxylophus 276
pentaprion 273
petersi 209
poncensis 627, 629
porcatus 504
pulchellus 110, 262, 273, 627, 629
punctatus 627
punctatus boulengeri 407
punctatus 407
reconditus 572
roosevelti 627
roquet 474
roquet salinei 480

- zebrilus* 480
sagrei 209, 288, 478, 491
scriptus 108, 628, 637
semilineatus 294, 511, 513
sheplani 247, 252, 257, 274
shrevei 479
sp. n. near eulaemus 257, 265, 274, 280
stratulus 115, 273, 473, 627, 629
trachyderma 408
transversalis 408
tropidonotus 262, 273
tropidonotus spilorhipis 249
tropidonotus 249
valencienni 209, 247, 252, 257, 269, 274, 288, 515, 570
wattsi 499
whitemani 552
Aperoprists catamarcensis 209
Aptycholaemus longicauda 209
Audantia 284
Basiliscus basiliscus 209
galeritus 209
Brachylophus 110
Brachylophus fasciatus 209, 563
Callisaurus 654, 658
Callisaurus (Cophosaurus) texanus scitulus 402
draconoides 209
Chalarodon 202
Chalarodon madagascariensis 209
Chamaeleolis 245, 284, 288
Chamaeleolis chamaeleontides 209, 273
porcus 209, 273, 295
Chamaelinorops 245, 284
Chamaelinorops barbouri 209, 248, 285
Conolophus subcristatus 209
Cophosaurus 403
Cophosaurus texanus 209
Corythophanes cristatus 209
Crotaphytus 377
Crotaphytus collaris 209, 656
Ctenoblepharis 203
Ctenoblepharis adpersus 209
Ctenosaura pectinata 209
Cyclura 110
Cyclura cornuta stejnegeri 209
Deiroptyx 284
Diaphoranolis 284
Diplolaemus bibroni 209, 273
Dipsosaurus 368, 487
Dipsosaurus dorsalis 209, 656
Ecphymotes acutirostris 120
pictus 120
plica 120
undulatus 120
Enyaliodes 203
Enyaliodes heterolepis 209
laticeps festae 209
laticeps 209
oshaughnessyi 209
palpebralis 209
Enyaliosaurus quinquecarinatus 209
Enyalius 280
Enyalius bilineatus 209
brasiliensis 209
iheringii 209
pictus 120
Gambelia wislizenni 209
Geiseltaliellus 368, 379
Holbrookia 403, 654, 658
Holbrookia maculata 209
texana 656
Hoplocercus 203
Hoplocercus spinosus 209
Iguana 110, 367
Iguana iguana 101, 200, 209
pinguis 111
tuberculata 200
Lacerta marmorata 118, 122
Laemactus longipes 249
serratus 209, 249
Leiocephalus 368, 379
Leiocephalus carinatus 209, 491
vinculum altavelensis 89
Leirolepis auduboni 120, 122
Leiosaurus belli 209
Liolaemus 203, 658, 672
Liolaemus alticolor walkeri 209
Mariguana 284
Morunasaurus 203, 259
Morunasaurus annularis 209, 249
groi 209, 249
Norops 284
Ophryoessoides ornatus 209
Oplurus 202
Oplurus cyclurus 209
fierinensis 209
grandidieri 209
quadrinaculatus 209
saxicola 209
Paradipsosaurus 368, 378
Parasauromalus 367, 378
Pelusaurus 203
Petrosaurus 654, 658
Petrosaurus mearnsi 209
Phenacosaurus 245, 274, 284, 288, 293
Phenacosaurus heterodermus 269, 295
richteri 209
Phrynosoma 654, 658
Phrynosoma cornutum 209, 656
coronatum 200
Phymaturus palluma 209
Platynotus semitaeniatus 209
Plica plica 120, 209
Polychroides peruvianus 121
Polychrus acutirostris 118, 209
angustirostris 121
anomalus 120, 122
fasciatus 120, 122
femoralis 121, 128
geometricus 120
gutturosus 120
liogaster 121
marmoratus 118
marmoratus acutirostris 121

- femoralis* 121
liogaster 121
marmoratus 121
spurrelli 121
neovidanus 120, 122
peruvianus 121, 128, 209
spurrelli 121
strigiventris 120, 122
virescens 119, 122
Pristidactylus scapulatus 209
Pristiguana 367, 377
Pristodactylus achalensis 273
Proctotretus 203
Proctotretus azureus 209
pectinatus 209
Psilocercus 119
Psilocercus marmoratus 122
Sator 654, 658
Sator angustus 209, 656
Sauromalus 368
Sauromalus obesus 209
Sceloporus 5, 203, 230, 487, 654, 656, 664
Sceloporus formosus 209
graciosus 209, 656
grammicus 666, 668, 671
grammicus disparilis 209
horridus albiventris 209
occidentalis 457, 656
olivaceus 656
poinsetti 209
scalaris 209
undulatus 209, 656
undulatus consobrinus 400
uniformis 209
variabilis 209
Sphaerops anomalus 120, 122
Stenocercus guentheri 209
Streptosaurus 655
Strobilurus torquatus 209
Tropidodactylus 284
Tropidurus 128
Tropidurus peruvianus 209
torquatus 456
Uma 404, 654, 658
Uma notata 209
Uracentron flaviceps 209
Uranoscodon superciliaris 209
Urosaurus 404, 654, 658
Urosaurus bicarinatus 209
ornatus 656
Urostrophus torquatus 209
vautieri 209
Uta 654, 658
Uta stansburiana 209
Vilcunia 203
Xiphocercus 284
FAMILY LACERTIDAE
Algyroides vauereselli 354
Cabrita 218
Durotrigia 373, 382
Eolacerta 370
Lacerta 196, 230, 370
Lacerta agilis 150, 196, 199
jacksoni 354
lepida 196
viridis 196, 201
Ophisops 218
Plesiolacerta 369
Podarcis muralis 196, 201
FAMILY NECROSAURIDAE
Colpodontosaurus 371
Eosaniwa 371
Necrosauridae 389
Necrosaurus 371
Parasaniwa 371, 389
Provaranosaurus 371
FAMILY PARAMACELLODIDAE
Paramacellodidae 376
Paramacellodus 370
Saurillodon 370
FAMILY PYGOPODIIDAE
Pygopodiidae 213, 381
FAMILY SCINCIDAE
Ablepharus 218
Afroablepharus 217
Carlia amax 214
novaeguineae 216
Chalcides chalcides 201
Contogenys 370, 385
Corucia zebrata 435
Cryptoblepharus 215
Cryptoblepharus plagiocephalus 214
Dasia smaragdinum 360
Emoia arnoensis 360
boettgeri 360
boettgeri orientalis 360
cyanura 216, 360
Eugongylus albofasciolatus 359
Eumeces 230, 370, 384, 386, 487, 654
Lamprolepis smaragdinum 360
Leiopisma greeni 219
pretiosum 219
suteri 618
trilineatum 215
Leptosiphos blochmanni 354
graueri 354
hackarsi 354
meleagris 354
Lerista 215
Lerista elegans 215
Lipinia noctua 359
Mabuya megalura 354
sloanei 110, 115
Macroscincus coctei 201
Menetia 217
Morethia 215
Neoseps 386
Notoscincus 216
Panaspis 217, 354
Panaspis cabindae 216
Prasinohaema virens 275, 276
Proablepharus 215
Sauriscus 370, 385
Scelotes 201, 218

Scincella 5, 218, 386

Scincidae 384

Sphenomorphus tympanum 214

Tiliqua 198, 201, 437

Tiliqua scincoides 196

Typhlacontias 218

FAMILY TEIIDAE

Ameiva ameiva 197

exsul 104, 115

Anotosaura 218

Bachia 218

Callopistes 369

Chamops 369

Cnemidophorus 369, 487, 654

Cnemidophorus tigris 624

Gymnophthalmus 218

Heterodactylus 218

Iphisa 218

Meniscognathus 369

Micrablepharus 218

Peneteius 369

Polyglyphanodon 369

Teiidae 383

Tretioscincus 218

Tupinambis 369

FAMILY UROMASTYCIDAE

Leiolepis 195, 380

Leiolepis belliana 200, 202, 206

belliana belliana 210

ocellata 210

guttata 210

peguensis 210

reevesii reevesii 210

rubritaeniata 210

triploida 210

Uromastyx 195, 368, 380

Uromastyx acanthinurus 205

acanthinurus acanthinurus 210

wernerii 210

aegyptius 201, 210

asmussi 204, 205, 210

benti 210

dispar 203

geyeri 210

hardwickii 201, 205, 210

loricatus 204, 205, 210

macfadyeni 210

microlepis 210

ocellatus 210

ornatus 206, 210

philbyi 210

princeps 208

princeps princeps 210

scortecci 210

spinipes 201

thomasi 210

FAMILY VARANIDAE

Iberovaranus 371

Indovaranus 535

Lanthanotus 371

Palaeosaniwa 371

Saniwa 371, 379, 390

Varanidae 390, 535

Varanus 226, 371

Varanus bengalensis 535, 544

komodoensis 535, 537, 550

mertensi 537

niloticus 537

salvadori 537

salvator 537

spenceri 537

varius 537

FAMILY XANTUSIIDAE

Lepidophyma 5, 370

Palaeoxantusia 370

Xantusiidae 207, 386

FAMILY XENOSAURIDAE

Shinisaurus 370, 386

Xenosaurus 386

SUBORDER OPHIDIA

FAMILY BOIDAE

Candoia carinata 440

Chondropython viridis 437

Epicrates monensis 104

FAMILY COLUBRIDAE

Alsophis 110

Arrhyton 110

Conophis 296

Dasypeltis ater 354

Dipsadoboa unicolor viridiventris 355

Dromicus 110

Duberria lutrix 352

Gonyosoma 303

Heterodon 296

Leimadophis epinephelus ecuadorensis 138

Leptodeira 5

Liophis 110

Liophis alticolus 132

bimaculatus 132

bimaculatus lamonae 132

bipraeocularis 132

cobella alticolus 132

epinephelus 132

epinephelus albiventris 138

bimaculatus 138

ecuadorensis 132

epinephelus 137

fraseri 138

juvenalis 139

lamonae 139

opisthotaenia 139

pseudocobella 139

exiguus 104

fraseri 132

opisthotaenia 132

portoricensis 104, 112

pseudocobella 132

reginae albiventris 132

quadrilineata 132

taeniurus juvenalis 132

williamsi 146

Natrix natrix 304

sipedon 691

Opheomorphus alticolus 138
Philothamnus heterodermus ruandae 354
Pituophis melanoleucus 489
Psammophylax variabilis 354
Synophis bicolor 294
Thamnophis sirtalis 296
 sirtalis parietalis 296

Zamensis ater 132

FAMILY CROTALIDAE

Crotalus 128

FAMILY TYPHLOPIDAE

Typhlops angolensis irsaci 354
 polylepis 354

richardi 104, 112

FAMILY VIPERIDAE

Atheris nitschei nitschei 354

Causus 304

Cerastes cornutus 304

SUBCLASS ARCHOSAURIA

ORDER THECODONTIA

aetosaurs 322

parasuchians 322

phytosaurs 322

proterosuchians 322

Stagonolepis 322

Stegomosuchus 323

ORDER CROCODILIA

Crocodylia 186, 305, 332

SUBORDER PROTOSUCHIA

Hemiprotosuchus 333

Orthosuchus 322

Protosuchus 322, 332

Protosuchus richardsoni 308, 310, 333

SUBORDER MESOSUCHIA

Bernissartia 323

mesosuchians 323

Teleosaurus 323

SUBORDER THALATTOSUCHIA

thalattosuchians 323

SUBORDER EUSUCHIA

FAMILY CROCODYLIDAE

Alligator 306, 333

Alligator mississippiensis 306, 320, 333
 sinensis 319

Caiman 333, 335

Caiman crocodilus 306, 321, 695, 700, 702

crocodilus yacare 700, 701

latirostris 320, 701

Crocodylus 306, 333, 335

Crocodylus acutus 318, 699

cataphractus 314

intermedius 319, 695, 697

johnstoni 314

mindorensis 317

moreletii 318

niloticus 315, 697

novaeaguinae 317

palustris 315

porosus 316, 440, 697

rhombifer 318

siamensis 316

Gavialis 306, 335

Gavialis gangeticus 313

Hassiacosuchus 323

Leidyosuchus 323

Melanosuchus niger 321, 695

Osteolaemus 306

Osteolaemus tetraspis 319

Paleosuchus palpebrosus 318, 321, 695

trigonatus 318, 321, 695

Procaimanoidea 323

Tomistoma 306

Tomistoma schlegelii 313

ORDER SAURISCHIA

Plateosaurus 191

ORDER ORNITHISCHIA

ornithischians 322

SUBCLASS SYNAPSIDA

ORDER THERAPSIDA

Glossopteris 379

Lystrosaurus 379

therapsids 223

REPTILIA INCERTAE SEDIS

Protolacerta 373

CLASS AVES

Apterygidae 186

Anatidae 186

Anthracothonax dominicus 509

birds 186, 223, 412, 507

Bucconidae 509

Burhinidae 186

Cariama 128

Coccyzus 517

Coereba flaveola 509, 511

Cracidae 186

Cuculidae 510

Dendrocolaptidae 509

Dendroica caerulescens 511, 514

coronata 511

discolor 511

palmarum 511, 514

petechia 511

tigrina 511

Dromaeus 186

ducks 517

Falco sparverius 110

Formicariidae 509, 514

Furnariidae 509

Geothlypis trichis 511

hoatzin 704

honeycreepers 456

ibis 704

Icterus dominicensis 509, 511

Mniotilta varia 511

Momotidae 509

Otididae 186

Pachyptila turtur 615

Parula americana 511

parulids 510, 512, 516

Pelecanoides urinatrix 615

Phaenicophilus palmarum 511
Phaethon aethureus 110
 Phoenicopteridae 186
 Rheidae 186
Saurothera 517
Seiurus aurocapillus 511
 novaeboealis 511
Setophaga ruticilla 511
 Struthionidae 186
Sturnella magna 412
 suboscines 513
Thamnophilus punctatus 510
 Threskiornithidae 186
 Tinamidae 186
Todus 512
Todus angustirostris 509
 subulatus 509, 511
 Troglodytidae 514
Tyrannus 517
Vireo 516
Vireo altiloquus 509, 511
 wrens 513
CLASS MAMMALIA
Alouatta seniculus 522
Artibeus harti 414
Ateles paniscus paniscus 522
 bats 412
 capybara 702
Cebus apella apella 522
 nigrivittatus 522
Chiropterus satanas chiropterus 522

deer 456
Felis catus 112
Gorilla beringei 353
Lasiurus borealis borealis 413
 cinereus 413
 seminolus 413
Leptonycteris nivalis 413
 sanborni 413
 yerbabuenae 413
Lonchophylla robusta 414
 monkeys 521
 moose 456
Mus musculus 671
Myotis grisescens 413
 keaysi 414
 oxyotus 414
Pithecia pithecia 522
 possum 438
Rattus exulans 616
 rhesus monkey 208
 rodents 517
Saguinus midas midas 522
Saimiri sciureus 522
Sturnira bidens 414
 bogotensis 414
 erythromos 414
 ludovici 414
Tadarida brasiliensis 413
Vampyressa melissa 414
Vampyrops dorsalis 414
 vittatus 414

3 2044 118 660 331

ISBN 0-910999-00-7