QK
191
T55
1907
Bot^{H.C.}~~any~~
~~Hitch-~~
~~Cock -~~
~~Chase~~

Banister's Catalogue
of
Virginia Plants

WASHINGTON
1907

HITCHCOCK AND CHASE LIBRARY
Smithsonian Institution

In Memory of John Banister
Pioneer of Virginia Botany

QK

191

TSS

1907

Botany

Hitchcock - *Passiflora*

Chase

PREFACE

The object of this little volume is to bring within reach of those who have not access to the earlier works on our flora, the most important facts relating to the botany of Maryland and Virginia. This is perhaps best accomplished by the reprinting of old records, as the reader will thus obtain an objective view of the works of the pioneers.

Jacques Philippe Cornut's *Historia Plantarum Canadensium*, published in Paris, 1635, is undoubtedly one of the earliest records of North American plants. This work contains descriptions and illustrations of many species the geographical ranges of which include not only *New France* but also *New England* and the territory southward to *Virginia* and the *Carolinas*. Of the numerous species described whose identity has been established beyond any possible doubt, we may cite *Filix bac-cifera* (*F. bulbifera*), found occasionally in our region, and *Adiantum americanum* (*A. pedatum*) one of our characteristic ferns of low, deciduous woods.

In 1640, Parkinson published his *Theatrum Botanicum* wherein he records many plants from Virginia. We may cite again *Adiantum pedatum* under the name of *Adiantum fruticosum americanum*. The weird, but beautiful *Juniperus virginiana*, the characteristic ornament of the lower Potomac region and elsewhere, is mentioned by Parkinson.

In 1686, Ray began the publishing of his *Historia Plantarum* [vol. 1, 1686; 2, 1688; 3, 1704], in which numerous species from Maryland and Virginia are described. Banister's Catalogue of Virginian Plants was published here. This document is of special interest since it is without much doubt the earliest record, dealing exclusively with Virginian plants. Many of the species are recognized at once by any one familiar with the plants of our region, and in the notes to follow the reprint of *Banister's Catalogue* attempts will be made to determine the species and give their history as fully as possible.

The works of Morison, Hermann, Plukenet, Bobart and others will be referred to in the account of the species.

The task of identifying an old species from the descriptions only is a difficult one, for it requires a careful study of the prominent authors of the period in which a species was published. The method of description of two centuries ago presupposes also an intimate knowledge of the older masters and their terminology. This is readily seen if we examine Banister's second species: [*Alsine Spargula latifolia reptans*] *Becabungae folio*. The expression serves both as a name and as a description, wherein a knowledge of another plant, *Veronica beccabunga*, is prerequisite.

I am indebted to Messrs. *Frederick V. Coville* and *P. L. Ricker* for their kind aid in the correction of the manuscripts and proofs.

Washington, D. C.

Ivar Tidestrom

BANISTER'S CATALOGUE
OF
VIRGINIAN PLANTS

(From RAY, HIST. PLANT. 2: 1926-1928. 1688.)

E Catalogo huc transmisso *Anno* 1680. quem composuit eruditissimus Vir & consummatissimus Botanicus D. *Johannes Banister* Plantarum a seipso in Virginia observatarum.

A

Alsine *Spergula latifolia reptans.*

Becabungae folio.

Althaea lutea Pimpinellae majoris folio, floribus parvis, seminibus rostratis. Folia hujus plantae pediculis insident.

Althaea magna Aceris folio, cortice Cannabino, floribus parvis semina rotatim in summitate caulium, singula singulis cuticulis rostratis co-operta ferens.

Althaea magna quinquecapsularis, cortice Cannabino, foliis integris subtus albicantibus, floribus magnis ex fundo saturate rubro albis.

Alth. magna quinquecapsularis, cortice Cannabino, foliis Malvarum modo divisis, subtus viridibus.

Ambrosia inodora foliis non divisis.

gigantea inodora foliis asper is trifidis.

Anchusa lutea minor, quam Indi Paccoon vocant seipsos ea pingentes.

Anemone latifolia sylvestris alba.

Apocynum erectum non ramosum folio subrotundo, umbellis florum rubris.

Apoc. erect. non ramos. latiore folio, umbellis florum albicantibus.

Apoc. erect. minus, umbella florum candida.

1. *Stellaria* *thunbergii*

2

3

4 *Napaea* *hexa*

7 *ambrosia* *trifida* *antegrifolia*

8

9 *ambrosia* *virginiana* L.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Emolobus laevis

9 *Alisma terrefolium*

10 " *pusillum*

11 " *dracunculium*

12 *Aster virginicus*

13 *Crataegus cuneata*

14

15 *Urtica virginiana* L.

16 *Castanea pumila*

1 Apoc. erect. non ram. Asclepiadis folio, umbellis
florum rubentibus.

2 Apoc. minus non lactescens, caule & foliis hirsu-
tis, floribus saturate luteis.

3 Apoc. erect. non ram. Roris marini foliis umbel-
lis florum candidis.

4 Apoc. petraeum ramosum Salicis folio.

5 Apoc. scandens, capsulis brevibus spinis asperis.

6 Apoc. scand. capsulis alatis.

7 Apoc. scand. capsulis planis.

Haec omnia siliquas ferunt tumentes.

8 Apoc. erect. ramosum, caule rubente, foliis ob-
longis parvis, siliquis [ex flosculis albis] tenu-
issimis, binatim ad extremitates conjunctis.

9 Arisarum triphyllum, pene viridi.

10 Aris. triph. minus, pene atro-rubente.

11 Aris. Dracontii foliis pene longo acuminato.

12 Arum aquaticum, foliis in acumen desinentibus,
fructu viridi.

13 Arum fluitans, pene nudo.

C

14 Carduus Jaceoides purp. foliis subtus incanis, ca-
pite viscoso.

15 Caryophyllata flore semper albo.

16 Castanea pumila racemoso fructu parvo, in sin-
gulis capsulis echinatis unico, The Chinqua-
pin. Autor descriptionis Carolinae ex hac
nuce Chocolatam fieri refert non multo inferi-
orem ei quae ex Cacao fit.

Centaurium minus caule quadrato alato, flore
carneo amplo, umbilico luteo.

Centaurium luteum Ascyroides.

Clematis purpurea repens petalis florum coriaceis.

Clem. erecta, humilis non ramosa, foliis subro-
tundis, flore unico ochroleuco.

Cochlearia flore majori. In locis udis a salsis
procul remotis.

Conyza coerulea acris Americana.

Cucumis fructu minimo viridi, ad maturitatem
perducto nigricante. Fructus Bryoniae albae
bacca non multo major est, cujus primo aspectu
speciem esse putaveram.

D

Dens caninus flore luteo.

Digitalis flore pallido transparenti, foliis & caule
molli hirsutiae imbutis.

Digit. rubra minor, labiis florum patulis, foliis
parvis angustis.

Digit. lutea elatior Jaceae nigrae foliis.
lutea altera, foliis tenuius dissectis thecis
florum foliaceis.

parva comis coccineis.

E

Eryngium campestre Yuccae foliis, spinis tenellis
hinc inde marginibus appositis.

Euonymus capsulis eleganter bullatis.

F

Filix mas foliis integris auriculatis.

mas rachi seu nervo medio alato.

1 *Sabbatia difformis* L.

2

3 *Clematis viorna* L.

4 *Clematis ochroleuca*

5 *Androsace virginica*

6 *Epistemon coeruleum*

7 *Melothrin pendula* L.

8 *Gaythronium dens canis* L.

9 *Pentstemon hispidus*

10 *Gerardia purpurea*

11 *Dacystema flavum*

12 *Gerardia pedicularis*

13

14 *Caryophyllus aquaticum*

15 *Cissampelos bullata*

16 *Polystichum acrostichoides*

17 *Phryopteris hexagonoptera*

8 *Passiflora cucurbitaria*

9 *Rhus toxicodendron*

10 *Lycopodium pubescens*

11 *Polygonum sarothamum*

12 *Chenopodium verrucosum*

13

1 [Filix] foemina foliis per margines pulverulentis,
seminibus fimbriatis.

2 Fumaria siliquosa lutea.

3 Siliquosa altera grumosa radice, floribus
gemellis ad labia conjunctis.

✓ Fungus (ex stercore equino) capillaceus capitulo
rorido, nigro punctulo in summitate notato.

Ex recenti fimo noctu exoritur cauliculis erectis, vix digitum longis capillorum instar tenuibus nec minus densis seu confertis. Singuli Cauliculi parvulo globulo aqueo coronantur, qui in summa sui parte macula parva nigra Limacis oculo simili insignitur.

G

5 Gentianae affinis foliis glabris serratis, floribus Ran-
nae referentibus.

6 Gladiolus coeruleus hexapetalos, caule etiam gladiato.

7 Gratiola foliis latioribus serratis.

H

8 Hedera trifolia Canadensis foliis sinuatis.

9 Helleborine flore rotundo luteo, purpureis venis
striato. The Mockasine flower.

10 Helxine late scandens seminibus majoribus.

11 Helxine frutescens Bryoniae nigrae foliis, capsulis
triquetris amplis Pergamenis.

12 Hieracium fruticosum latifolium foliis punctulis
& venis sanguineis notatis.

13 Hyacinthus Occidentalis flore pallide coeruleo.

Hypericum parvum caule quadrato seu Ascyron
minimum.

Hyper. pumilum semper virens caule compresso
ligneo; ad bina latera alato, flore luteo tetra-
petalo, seu Crux S. Andreae.

Hyper. frutescens luteum Phillyrrheae foliis.

I

Jacea non ramosa tuberosa radice, foliis plurimis
rigidis perangustis, flores ferens multos parvos
rubentes acaules in spica ad caulem sessiles.

Jac. non ram. tub. rad. foliis latioribus flores fe-
rens pauciores, majores, squamis hiantibus ar-
matis & pediculis curtis insidentes.

Jacobaea lanata foliis brevibus subrotundis, lanata
altera foliis longis angustis.

Jasminum arboreum foliis amplis, oblongis, su-
perne virentibus, subtus leni canitie pubescen-
tibus, flores albos in quatuor lacinias longas an-
gustas ad umbilicum usque partitos racematim
ferens.

Iris aculeata baccifera arborea minus ferax.

Iris coerulea latifolia & angustifolia.

Chamae-Iris verna odoratissima, latifolia coerulea
repens.

L

Pseudo-Lathyrus luteus glaber, siliquis tumentibus,
duplicem seminum seriem continentibus.

Pseudo-Lath. lut. hirsutus siliq. tument. continenti-
bus duplicem seriem seminum.

Laurus Tinus floribus albidis eleganter bullatis.

1

2

3

4 *Lacinaria stricta*

5 " *serotina*

6 *Senecio tomentosus*

7 *Chionanthus virginicus*

8

9 *Iris varicolor* - *I. prismatica*

10

11 *Baptisia tinctoria*

12 " *villosa*

13 *Thalictrum latifolium* L.

- 1. *Amargosa* *abundans* L.
- 2. *Lilium* *superbum* L.
- 3. *Asclepias* *virginiana* L.

4. *Agave* *virginica* L.

5. *Helianthus* *pyramidalis* L.

6.

7. *Cruciferae* *stolonifera* L.

Flos nondum apertus pyxidi S.M. Magdalenae
(ut nonnunquam pictam vidi) similis est.

Lilio-narcissus humilis albus.

Lilium S. Martagon floribus reflexis ex luteo rubentibus, purpureis maculis eleganter notatis.

Lilium S. Martagon pusillum florib. minutissimis herbaceis. Caulem habet vix dodrantalem, verticillo foliorum unico cinctum, cujus summitas quatuor floribus reflexis Solani lignosi floribus magnitudine haud aequalibus, pediculis parvis insidentibus coronantur.

Lilium Squillae foliis, denticellis parvis ad margines serratis. Caule est alto, nudo, ad cujus summitatem prodeunt flores in spica sessiles, petalis paene ad imum (ut loqui amant faciales) quasi erasis; staminibus sex (si non male memini) purpuro-coeruleis, mole sua aliquid tantulum depressis. Flos quamvis aspectu non admodum pulcher sit pergratum habet odorem. Radicem habet imbricatam instar Lili, folia crassa admodum & succulenta, non tamen Sempervivi species est.

Lonchitis maxima foliis planis *i. e.* non dentatis, nec pulverulentis maculis notatis, uno in cespite foliorum unicum protrudens caulem foliis angustioribus, pulverem seu semina in membranulis quasi in capsulis ferentibus compositum.

Lonchitis major Polypodii facie. Haec atque etiam vulgaris simili modo florida est.

Lychnis plum aria alba, foliis ad geniculum quatuor cruciatim positis, thecis florum tumentibus.

Lysimachia lutea minor, foliis & floribus purpureo punctatis.

Lysimachia siliquosa lutea minor.

M

Melissa elatior foliis magnis dentatis glabris, ad geniculum binis : flores odoratos luteos patulos stamina bina quasi cornua protrudenti[a] in summitate caulium racematim ferens.

Mercurialis tricoccus hermaphroditica, s. ad foliorum juncturas ex foliolis cristatis julifera simul ac fructum ferens.

Muscus erectus dense complicatus *Cupressi* foliis major & minor. In rupe quadam prope Sabinas foliis *Cupressi*.

Myrrhis minor procumbens, seu potius *Cerefolium*.

N

Nux vesicaria Virginiana Park.

O

Orchis palmata elegans lutea cum longis calcaribus luteis, palmata lutea minor nullis calcaribus. Hermaphroditica, flore minore, calcare longiore.

Origanum floribus amplis luteis, purpureo maculatis, cujus caulis sub quovis verticillo decem vel duodecimo foliis est circumcinctus.

Orig. cujus ramorum summitates floribus dilute rubris in verticillos congestis coronantur.

Orig. foliis ad summitatem caulium canis, floribus multis pallide coeruleis in cymis ramorum dense stipatis.

3 *Collinsia scutellaria* L.

4 *Scutellaria virginica* L.

5 *Lupinus* L.

6 *Chorizanthe maculata* L.

7 *Phacelia trifoliata* L.

8 *Thlaspi virginicum* L.

9 *Thlaspi virginicum* L.

10 *Thlaspi virginicum* L.

11 *Thlaspi virginicum* L.

1 *Agavea erecta* L.
2 *Monarda mollis* L.

3

4

5

6 *Thalictrum chrysothellum*

7

8 *Alchemilla filomatensis* L.

9 *Collyria lutea*

10 *Androsace*

11 *Androsace*

12 *Androsace*

13 *Androsace*

14 *Androsace*

15 *Androsace* ?

16

Ornithogalum luteum parvum foliis gramineis hirsutis.

Orobanche radice dentata caule & flore albo.

Flos ejus quem unicum in uno caule Goodyeri Orobanches similis est sed major. Conceptaculi seminalis venter seu pars protuberans non rotunda est canaliculata.

P

Pepo fructu parvo compresso.

Phalangium ramosum floribus albis, ad fundum viridibus.

Phalangium album non ramosum floribus albis ad caulem spicatum sessilibus.

Phyllitis parva saxatilis per summitates folii proli-fera.

Pisum spontaneum purpureum.

Plantago aquatica latiore folio.

Polygala seu Flos ambarvalis floribus luteis in caput oblongum congestis.

Polyg. rubra spica parva compacta.

Polygala spicata rubra major foliis & caulibus coerulescentibus.

Pol. quadrifolia S. cruciata floribus ex viridi rubentibus, in globum compactis.

Pol. quadrif. minor spica parva rubente.

Polygonatum ramosum capsula prismatica, ramosum perfoliatum flore ochroleuco capsula trigona.

Polypodium parvum foliis minutim serratis.

Polypodium minus alterum Scolopendriae facie.

Potamogiton Virginianum.

Q

Quercus variae species, 1. Pumila. 2. Alba. 3. Rubra. 4. Hispanica. 5. Castaneae folio. 6. Lini aut Salicis foliis. 7. Fruticosa. Harum primam in Historia, quintam in Appendice meminimus.

R

Ranunculus Thalictri folio, radice grumosa; Anemone sylv.

Rapuntum minimum glabrum. In uliginosis minus glabrum flore pallidiore.

Rhamnus Prunifolius fructu nigro, ossiculo compresso. The black Haw.

Rhus ramis ex stipe pullulantibus glabris. Hujus truncus carpi crassitiem nunquam superat, pollicis raro excedit. Baccae sapore sunt subsalso cum pungenti illo acore qui in Tamarindis sentitur mixto.

Rubia parva foliolis ad geniculum unumquodque binis, flore coeruleo fistuloso.

Rub. parva latifolia foliis ad genic. binis, flore rubente.

S

Sanicula seu Auricula ursi Cyclamini flore.

Saxifraga petraea Alsinefolia.

Sedum saxatile parvum, caule gracili aphylo, floribus rubentibus.

Solanum verticillatum latifolium molle, floribus obsolete rubris, baccis luteis.

Rumex *penula*, *alba*, *rubra*,
galeata, *crucis*, *phellon*,
primoides.

2 *Synedra* *Chakobsky*

3 *Sabunum* *penifolium*

4 *Hosobus* *coccinea* &
5 *H* " *purpurea*

- 3 *Tristemon coccineus*
- 2
- 1 *Peltandra caroliniana*
- 5
- 4 *Solanum carolinense*
- 7 *Sorbus arbutifolia*
- 8 *Lilium pudica*
- 10 *Urtica dioica*

1 *Macchia carolinensis*

10 *Campanula medium*

11

12

Sol. verticill. angustiore folio, flore ochroleuco.

Sol. triphyllum flo. tripetalo, atro-purpureo, in foliorum sinu absque pediculo sessili.

Staphis agria fol. dilute viridibus.

Stramonium fundo floris coeruleo, pomis longioribus spinis armatis.

T

Trichomanes major foliis longis auriculatis.

U

Valeriana Graeca seu Valeriana coerulea minor.

Veronica pratensis Serpyllifolia.

Viola tricolor nudo caule, foliis tenuius dissectis.

Viola alabastrites pentaphyllea, Cochleariae sapore, Nasturtii species.

Urtica urena major seminibus rotundis, compressis loculamentis viridibus inclusis & in caulis summitate racematim dispositis.

Inter semina ad me e Virginia transmissa, Anno 1687. ab eodem *D. Banister*, nonnulla invenio quorum nomina *in hoc* Catalogo non habentur, v. g.

Erigeron (frutex marit.) Halimi folio: Senecio arborescens Atriplicis folio.

Eryngium Plinii Portulacae foliis a *D. Spragge* acceptum.

Euonymus (ni male memini) Pyracanthae foliis.

Lysimachia lutea corniculata maritima.

Cynoglossum coeruleum Buglossi foliis.

Senae spuriae tres species quarum una siliquis est hirsutis a D. *Spragge* acceptae. Duae a me satae germinarunt & plantae produxerunt, quae duae species Paiomiriobae a Pisone descriptae esse videbantur.

Pistacia nigra Coryli folio D. *Spragge*.

Ulmus fructu Lupulino.

Ricinus parvus Urticae folio.

Convolvulus bicapsularis seminibus pappo alatis.

Phalangium spicatum flosculo Arbuteo bullato aureo.

Lithospermum floribus rostratis.

Ricinus frutescens Fici foliis.

Gramen marinum echinatum. Hujus semina etiam a D. *Spragge* accepi, quae hac aestate in horto sata germinarunt, sed nondum ad frugem pervenerunt.

GK
191
T55
1907
Bot H.C.

Banister's Catalogue
of
Virginia Plants

WASHINGTON
1907

HITCHCOCK AND CHASE LIBRARY
Smithsonian Institution

In Memory of John Banister
Pioneer of Virginia Botany

OK
191
755
1909
Botany
Hitchcock
Chace

PREFACE

The object of this little volume is to bring within reach of those who have not access to the earlier works on our flora, the most important facts relating to the botany of Maryland and Virginia. This is perhaps best accomplished by the reprinting of old records, as the reader will thus obtain an objective view of the works of the pioneers.

Jacques Philippe Cornut's *Historia Plantarum Canadensium*, published in Paris, 1635, is undoubtedly one of the earliest records of North American plants. This work contains descriptions and illustrations of many species the geographical ranges of which include not only *New France* but also *New England* and the territory southward to *Virginia* and the *Carolinas*. Of the numerous species described whose identity has been established beyond any possible doubt, we may cite *Filix baccifera* (*F. bulbifera*), found occasionally in our region, and *Adiantum americanum* (*A. pedatum*) one of our characteristic ferns of low, deciduous woods.

In 1640, Parkinson published his *Theatrum Botanicum* wherein he records many plants from Virginia. We may cite again *Adiantum pedatum* under the name of *Adiantum fruticosum americanum*. The weird, but beautiful *Juniperus virginiana*, the characteristic ornament of the lower Potomac region and elsewhere, is mentioned by Parkinson.

In 1686. Ray began the publishing of his *Historia Plantarum* [vol. 1, 1686; 2, 1688; 3, 1704], in which numerous species from Maryland and Virginia are described. Banister's Catalogue of Virginian Plants was published here. This document is of special interest since it is without much doubt the earliest record, dealing exclusively with Virginian plants. Many of the species are recognized at once by any one familiar with the plants of our region, and in the notes to follow the reprint of *Banister's Catalogue* attempts will be made to determine the species and give their history as fully as possible.

The works of Morison, Hermann, Plukenet, Bobart and others will be referred to in the account of the species.

The task of identifying an old species from the descriptions only is a difficult one, for it requires a careful study of the prominent authors of the period in which a species was published. The method of description of two centuries ago presupposes also an intimate knowledge of the older masters and their terminology. This is readily seen if we examine Banister's second species: [*Alsine Spergula latifolia reptans*] *Becabungae folio*. The expression serves both as a name and as a description, wherein a knowledge of another plant, *Veronica becabunga*, is prerequisite.

I am indebted to Messrs. *Frederick V. Coville* and *P. L. Ricker* for their kind aid in the correction of the manuscripts and proofs.

Washington, D. C.

Ivar Tidestrom

BANISTER'S CATALOGUE
OF
VIRGINIAN PLANTS

(From RAY, HIST. PLANT. 2 : 1926-1928. 1688.)

E Catalogo huc transmisso Anno 1680. quem composuit eruditissimus Vir & consummatissimus Botanicus D. *Johannes Banister* Plantarum a seipso in Virginia observatarum.

A

Alsine *Spergula latifolia reptans.*

Becabungae folio.

Althaea latea Pimpinellae majoris folio, floribus parvis, seminibus rostratis. Folia hujus plantae pediculis insident.

Althaea magna Aceris folio, cortice Cannabino, floribus parvis semina rotatim in summitate caulium, singula singulis cuticulis rostratis cooperta ferens.

Althaea magna quinquecapsularis, cortice Cannabino, foliis integris subtus albicantibus, floribus magnis ex fundo saturate rubro albis.

Alth. magna quinquecapsularis, cortice Cannabino, foliis Malvarum modo divisis, subtus viridibus.

Ambrosia inodora foliis non divisis.

gigantea inodora foliis asperis trifidis.

Anchusa lutea minor, quam Indi Paccoon vocant seipsos ea pingentes.

Anemone latifolia sylvestris alba.

Apocynum erectum non ramosum folio subrotundo, umbellis florum rubris.

Apoc. erect. non ramos. latiore folio, umbellis florum albicantibus.

Apoc. erect. minus, umbella florum candida.

Apoc. erect. non ram. Asclepiadis folio, umbellis
florum rubentibus.

Apoc. minus non lactescens, caule & foliis hirsu-
tis, floribus saturate luteis.

Apoc. erect. non ram. Roris marini foliis umbel-
lis florum candidis.

Apoc. petraeum ramosum Salicis folio.

Apoc. scandens, capsulis brevibus spinis asperis.

Apoc. scand. capsulis alatis.

Apoc. scand. capsulis planis.

Haec omnia siliquas ferunt tumentes.

Apoc. erect. ramosum, caule rubente, foliis ob-
longis parvis, siliquis [ex flosculis albis] tenu-
issimis, binatim ad extremitates conjunctis.

Arisarum triphyllum, pene viridi.

Aris. triph. minus, pene atro-rubente.

Aris. Dracontii foliis pene longo acuminato.

Arum aquaticum, foliis in acumen desinentibus,
fructu viridi.

Arum fluitans, pene nudo.

C

Carduus Jaceoides purp. foliis subtus incanis, ca-
pite viscoso.

Caryophyllata flore semper albo.

Castanea pumila racemoso fructu parvo, in sin-
gulis capsulis echinatis unico, The Chinqua-
pin. Autor descriptionis Carolinae ex hac
nuce Chocolatam fieri refert non multo inferi-
orem ei quae ex Cacao fit.

Centaureum minus caule quadrato alato, flore
carneo amplo, umbilico luteo.

Centaureum luteum Ascyroides.

Clematis purpurea repens petalis florum coriaceis.

Clem. erecta, humilis non ramosa, foliis subro-
tundis, flore unico ochroleuco.

Cochlearia flore majori. In locis udis a salsis
procul remotis.

Conyza coerulea acris Americana.

Cucumis fructu minimo viridi, ad maturitatem
perducto nigricante. Fructus Bryoniae albae
bacca non multo major est, cujus primo aspectu
speciem esse putaveram.

D

Dens caninus flore luteo.

Digitalis flore pallido transparenti, foliis & caule
molli hirsutiae imbutis.

Digit. rubra minor, labiis florum patulis, foliis
parvis angustis.

Digit. lutea elatior Jaceae nigrae foliis.
lutea altera, foliis tenuius dissectis thecis
florum foliaceis.
parva comis coccineis.

E

Eryngium campestre Yuccae foliis, spinis tenellis
hinc inde marginibus apposis.

Euonymus capsulis eleganter bullatis.

F

Filix mas foliis integris auriculatis.

mas rachi seu nervo medio alato.

[Filix] foemina foliis per margines pulverulentis,
seminibus fimbriatis.

Fumaria siliquosa lutea.

Siliquosa altera grumosa radice, floribus
gemellis ad labia conjunctis.

Fungus (ex stercore equino) capillaceus capitulo
rorido, nigro punctulo in summitate notato.

Ex recenti fimo noctu exoritur cauliculis erectis,
vix digitum longis capillorum instar tenuibus nec minus densis seu confertis. Singuli
Cauliculi parvulo globulo aqueo coronantur,
qui in summa sui parte macula parva nigra Limacis oculo simili insignitur.

G

Gentianae affinis foliis glabris serratis, floribus Ranæ referentibus.

Gladiolus coeruleus hexapetalos, caule etiam gladiato.

Gratiola foliis latioribus serratis.

H

Hedera trifolia Canadensis foliis sinuatis.

Helleborine flore rotundo luteo, purpureis venis striato. The Mockasine flower.

Helxine late scandens seminibus majoribus.

Helxine frutescens Bryoniae nigrae foliis, capsulis triquetris amplis Pergamenis.

Hieracium fruticosum latifolium foliis punctulis & venis sanguineis notatis.

Hyacinthus Occidentalis flore pallide coeruleo.

Hypericum parvum caule quadrato seu *Ascyron minimum*.

Hyper. pumilum semper virens caule compresso ligneo; ad bina latera alato, flore luteo tetrapetalo, seu *Crux S. Andreae*.

Hyper. frutescens luteum Phillyrrheae foliis.

I

Jacea non ramosa tuberosa radice, foliis plurimis rigidis perangustis, flores ferens multos parvos rubentes acaules in spica ad caulem sessiles.

Jac. non ram. tub. rad. foliis latioribus flores ferens pauciores, majores, squamis hiantibus armatos & pediculis curtis insidentes.

Jacobaea lanata foliis brevibus subrotundis, lanata altera foliis longis angustis.

Jasminum arboreum foliis amplis, oblongis, superne virentibus, subtus leni canitie pubescentibus, flores albos in quatuor lacinias longas angustas ad umbilicum usque partitos racematim ferens.

Iris aculeata baccifera arborea minus ferax.

Iris coerulea latifolia & angustifolia.

Chamae-Iris verna odoratissima, latifolia coerulea repens.

L

Pseudo-Lathyrus luteus glaber, siliquis tumentibus, duplicem seminum seriem continentibus.

Pseudo-Lath. lut. hirsutus siliq. tument. continentibus duplicem seriem seminum.

Laurus Tinus floribus albidis eleganter bullatis.

Flos nondum apertus pyxidi S.M. Magdalenae
(ut nonnunquam pictam vidi) similis est.

Lilio-narcissus humilis albus.

Lilium S. Martagon floribus reflexis ex luteo rubentibus, purpureis maculis eleganter notatis.

Lilium S. Martagon pusillum florib. minutissimis herbaceis. Caulem habet vix dodrantalem, verticillo foliorum unico cinctum, cujus summitas quatuor floribus reflexis Solani lignosi floribus magnitudine haud aequalibus, pediculis parvis insidentibus coronantur.

Lilium Squillae foliis, denticellis parvis ad margines serratis. Caule est alto, nudo, ad cujus summitatem prodeunt flores in spica sessiles, petalis paene ad inum (ut loqui amant faciales) quasi erasis; staminibus sex (si non male memini) purpuro-coeruleis, mole sua aliquantulum depressis. Flos quamvis aspectu non admodum pulcher sit pergratum habet odorem. Radicem habet imbricatam instar Lili, folia crassa admodum & succulenta, non tamen Sempervivi species est.

Lonchitis maxima foliis planis *i. e.* non dentatis, nec pulverulentis maculis notatis, uno in cespite foliorum unicum protrudens caulem foliis angustioribus, pulverem seu semina in membranulis quasi in capsulis ferentibus compositum.

Lonchitis major Polypodii facie. Haec atque etiam vulgaris simili modo florida est.

Lychnis plum aria alba, foliis ad geniculum quatuor cruciatim positis, thecis florum tumentibus.

Lysimachia lutea minor, foliis & floribus purpureo punctatis.

Lysimachia siliquosa lutea minor.

M

Melissa elatior foliis magnis dentatis glabris, ad geniculum binis : flores odoratos luteos patulos stamina bina quasi cornua protrudenti[a] in summitate caulium racematim ferens.

Mercurialis tricoccus hermaphroditica, s. ad foliorum juncturas ex foliolis cristatis julifera simul ac fructum ferens.

Muscus erectus dense complicatus *Cupressi* foliis major & minor. In rupe quadam prope Sabinas foliis *Cupressi*.

Myrrhis minor procumbens, seu potius *Cerefolium*.

N

Nux vesicaria Virginiana Park.

O

Orchis palmata elegans lutea cum longis calcaribus luteis, *palmata lutea minor* nullis calcaribus. *Hermaphroditica*, flore minore, calcare longiore.

Origanum floribus amplis luteis, purpureo maculatis, cujus caulis sub quovis verticillo decem vel duodecimo foliis est circumcinctus.

Orig. cujus ramorum summitates floribus dilute rubris in verticillos congestis coronantur.

Orig. foliis ad summitatem caulium canis, floribus multis pallide coeruleis in cymis ramorum dense stipatis.

1. *Polygala lutea* L.
2. *Polygala fr. nigra* Willd. ?
 fr. angustata L.
3. *Polygala minor* L.
4. *Polygala crassata* L.
5. *Polygala multiflora* Willd.

Ornithogalum luteum parvum foliis gramineis
hirsutis.

Orobanche radice dentata caule & flore albo.

Flos ejus quem unicum in uno caule Goodyeri
Orobanches similis est sed major. Conceptacu-
li seminalis venter seu pars protuberans non
rotunda est canaliculata.

P

Pepo fructu parvo compresso.

Phalangium ramosum floribus albis, ad fundum
viridibus.

Phalangium album non ramosum floribus albis
ad caulem spicatum sessilibus.

Phyllitis parva saxatilis per summitates folii proli-
fera.

Pisum spontaneum purpureum.

Plantago aquatica latiore folio.

/ Polygala seu Flos ambarvalis floribus luteis in ca-
put oblongum congestis.

2 Polyg. rubra spica parva compacta.

3 Polygala spicata rubra major foliis & caulibus
coerulescentibus.

4 Pol. quadrifolia S. cruciata floribus ex viridi ru-
bentibus, in globum compactis.

5 Pol. quadrif. minor spica parva rubente.

Polygonatum ramosum capsula prismatica, ra-
mosum perfoliatum flore ochroleuco capsula
trigona.

Polypodium parvum foliis minutim serratis.

Polypodium minus alterum Scolopendriae facie.

Potamogiton Virginianum.

Q

Quercus variae species, 1. Pumila. 2. Alba. 3. Rubra. 4. Hispanica. 5. Castaneae folio. 6. Lini aut Salicis foliis. 7. Fruticosa. Harum primam in Historia, quintam in Appendice meminimus.

R

Ranunculus Thalictri folio, radice grumosa; Anemone sylv.

Rapuntum minimum glabrum. In uliginosis minus glabrum flore pallidiore.

Rhamnus Prunifolius fructu nigro, ossiculo compresso. The black Haw.

Rhus ramis ex stipite pullulantibus glabris. Hujus truncus carpi crassitiem nunquam superat, pollicis raro excedit. Baccae sapore sunt subsalso cum pungenti illo acore qui in Tamarindis sentitur mixto.

Rubia parva foliolis ad geniculum unumquodque binis, flore coeruleo fistuloso.

Rub. parva latifolia foliis ad genic. binis, flore rubente.

S

Sanicula seu Auricula ursi Cyclamini flore.

Saxifraga petraea Alsinefolia.

Sedum saxatile parvum, caule gracili aphylo, floribus rubentibus.

Solanum verticillatum latifolium molle, floribus obsolete rubris, baccis luteis.

Sol. verticill. angustiore folio, flore ochroleuco.
Sol. triphyllum flo. tripetalo, atro-purpureo, in
foliorum sinu absque pediculo sessili.
Staphis agria fol. dilute viridibus.
Stramonium fundo floris coeruleo, pomis longio-
ribus spinis armatis.

T

Trichomanes major foliis longis auriculatis.

U

Valeriana Graeca seu Valeriana coerulea minor.
Veronica pratensis Serpyllifolia.
Viola tricolor nudo caule, foliis tenuius dissectis.
Viola alabastrites pentaphyllea, Cochleariae sapo-
re, Nasturtii species.
Urtica urena major seminibus rotundis, compres-
sis loculamentis viridibus inclusis & in caulis
summitate racematim dispositis.

Inter semina ad me e Virginia transmissa, *Anno*
1687. ab eodem *D. Banister*, nonnulla invenio
quorum nomina *in hoc* Catalogo non habentur,
v. g.

Erigeron (frutex marit.) Halimi folio: Senecio
arborescens Atriplicis folio.

Eryngium Plinii Portulacae foliis a *D. Spragge* ac-
ceptum.

Euonymus (ni male memini) Pyracanthae foliis.

Lysimachia lutea corniculata maritima.

Cynoglossum coeruleum Buglossi foliis.

Senae spuriae tres species quarum una siliquis est
hirsutis a D. *Spragge* acceptae. Duae a me satae
germinarunt & plantae prodixerunt, quae duae
species Paiomiriobae a Pisone descriptae esse vi-
debantur.

Pistacia nigra Coryli folio D. *Spragge*.

Ulmus fructu Lupulino.

Ricinus parvus Urticae folio.

Convolvulus bicapsularis seminibus pappo alatis.

Phalangium spicatum flosculo Arbuteo bullato
aureo.

Lithospermum floribus rostratis.

Ricinus frutescens Fici foliis.

Gramen marinum echinatum. Hujus semina eti-
am a D. *Spragge* accepi, quae hac aestate in
horto sata germinarunt, sed nondum ad frugem
pervenerunt.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00088 4056