ALBERT R. MANN
LIBRARY
CORNELL UNIV.

Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

<http://www.archive.org/details/cu31924000615397>

SYSTEMATIC CENSUS
OF
AUSTRALIAN PLANTS,

WITH

Chronologic, Literary and Geographic Annotations;

BY

BARON FERDINAND VON MUELLER,

K.C.M.G., M.D., PH.D., F.R.S.,

F.I.S., F.G.S., F.R.G.S., C.M.Z.S., H.F.R.H.S. &c.

GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

^{* h}
"LEVATE OCULOS IN EXCELSUM ET VIDETE, QUIS CREAVIT HÆC OMNIA, QUI REDUCIT MULTITUDINEM FORUM."—*Isaiah* xl. 26.

MELBOURNE:

PRINTED FOR THE VICTORIAN GOVERNMENT

By M'CARRON, BIRD & Co., 37 FLINDERS LANE WEST.*

1882.

QK 431
M 94.

To

GEORGE BENTHAM, Esq., F.R.S., V.P.L.S.,

TO

SIR JOSEPH HOOKER, K.C.S.I., C.B., F.R.S., V.P.L.S.

AND TO

M. ALPHONSE DE CANDOLLE, F.M.R.S., F.M.L.S.,

WHO AS HEIRS OF GREAT NAMES WORTHILY SUSTAIN WORLD-WIDE ANCESTRAL FAME,

AND WHO AS LEADERS IN PHYTOGRAPHY DURING MORE THAN A GENERATION'S TIME

WILL BE PRE-EMINENT IN BIOMORPHIC SCIENCE THROUGH ALL AGES,

THESE PAGES

ARE OFFERED AS AN APPRECIATIVE TRIBUTE

FROM A YOUNG COLONY IN ANTIPODAL REMOTENESS.

Q
QK 431
M94

@ 38730

THIS work needs hardly to be introduced by prefatory observations to the notice of professed naturalists, who will recognise its scope and object at a glance. To those however, who may be less conversant in phytographic science, and who may use these pages, some explanatory preface would be needful, in order that they may fully understand the views, which guided the author through the details of this enumerative essay on Australian indigenous plants. Again, to all recipients it may be of some interest, to be informed of the causes, which gave rise to the present literary issue. Of the seven volumes of the "*Flora Australiensis*," elaborated by Mr. Bentham, mainly through the aid of the Botanic Departments of Kew and Melbourne, the earlier ones were written at a time, when a large portion of Australia remained yet untraversed by any explorers. Since then the territorial occupation as a whole has become nearly doubled, while new lines of geographic research have opened up vast additional tracts of lands for future settlements. Thus, the *Flora* of our island-continent became also more revealed, with the result of about 850 species of vascular plants having been added already to those described in the "*Flora Australiensis*"; but, irrespective of this, we have learned in later years much more concerning the regional distribution of the species formerly recorded. All these additional data, accumulated gradually during the last two decennia, though mainly kept together in the volumes of the "*Fragmenta Phytographiae Australiae*," became necessarily much disconnected and dispersed during the progress of discoveries and the early records connected therewith, so much so, that much difficulty was experienced of late, to command a clear and easy view over these scattered literary fragments. To collect and rearrange them into one or more volumes, conformous with those of the "*Flora Australiensis*," and supplementary to that work, proved beyond the range of operations of the venerable savant, who spent much of his precious time during fully sixteen years for grandly systematising throughout the native vegetation of these great parts of the British Dominions,—because his unrivalled experience and scientific valour were called more pressingly into channels of research, by which not only the realm of plants of Australia, but indeed that of the whole globe, became drawn both comprehensively and connectedly into the cycle of his observations. Perhaps it may fall yet to the share of the writer of the present pages, to furnish supplements to the "*Flora Australiensis*"; indeed, he has it under contemplation, to add one volume, solely to be devoted to a descriptive enumeration of all the additional species, and to be arranged in consonance with Bentham's views, whereas another volume would be rendered exclusively the vehicle of record of all new localities, from whence any species of the "*Flora Australiensis*" were obtained, when perhaps also such alterations, as largely augmented material may suggest, could be made to the descriptive text. But to complete the "*Flora*" in this manner, so far as our present state of knowledge will admit of it, would involve the thorough revision of the whole vast collections now stored in our local Botanical Museum; therefore, the connected re-examination of at least 80,000 specimens, from which the pages of the "*Flora*" largely, and those of the "*Fragmenta*" solely, have emanated. Such a task, to be performed with conscientious care, would need continued and close application throughout several years. Under these circumstances it seemed best, to pave the way for subsequent purposes through the issue of a preliminary publication, by which anyhow the use of both works, above referred to, would be facilitated. A postponement of the work for supplementary volumes, thus arising, would bring with it one advantage also, that of gaining time for carrying the augmentation of the *Flora* still further, as we would meanwhile profit from phytologic searches of travellers and settlers in wide regions of Australia, as yet in reference to vegetation but scantily investigated or altogether unexplored.

By deferring therefore the elaboration of supplemental volumes, we could aim more at future exhaustiveness, whereas the work as left by its illustrious author will meet mainly all present wants, and must remain the foundation for all future systematic study of the vegetation peculiar to this part of the globe. In offering now a separate statistic volume, the writer has availed himself of the opportunity, to place on record independently his views on preferable systematic sequences of orders, wherever occasionally his opinions differed from those enunciated by other phytographers. He furthermore could draw the limits of some genera and many species in accordance with original observations of his own; besides he was able to apply more rigorously the rules for systematic naming in strict application of chronologic data. But to avoid misapprehensions, the author at the very outset endeavours to anticipate objections, which may perhaps be raised against some portion of the nomenclature adopted in these pages. If, for instance, Haller is regarded as the earliest indicator of *Cyperaceae*, it must be conceded, that his wording of the order was "*Cyper*" in 1742, as well shown in Pfeiffer's extensive work. If however in accordance with the more exact limitation of this ordinal group of plants we ascribe it under the name of

Cyperaceae to De Candolle, as defined in 1805, or to St. Hilaire as circumscribed in the same year, or to A. L. De Jussieu, who already in 1789 adopted the same order under the appellation Cyperoideae, then we evidently act unjustly in not recognising prior claims; indeed, so far as wording is concerned, we lose the substance for the shadow, especially as no necessity exists, to strive for etymologic uniformity at the sacrifice of fair priority. To overcome difficulties or ambiguities of these kinds, we could use the preposition "from" (to be rendered *nach* in German and *d'après* in French) for indicating the originator of an order, genus or species, after the quotation of any later accepted authority. Indeed this principle was adopted already for a list of Australian plant-genera, published by the Royal Society of New South Wales in 1881; and had the scope assigned to the present pages admitted of it, the same rule would have been extended to the species so far as applicable.

To accomplish the object of rendering this publication, as a commentary index, one for easy reference, the schematic form appeared the most eligible. But to bring it within the means available for issue, all specific synonymy and any secondary notations had to be abandoned. Indeed for these, in quoting the "Flora" and the "Fragmenta" throughout, all needful indications are afforded. The geographic notes had to be limited to the initial letters, indicative of our main colonial territories, in which respect a plan was followed, precisely the same as that adopted in 1866 for a geographic list of nearly one thousand species of trees, then already known to constitute the very varied arboreous vegetation of Australia (see volume of the Intercolonial Exhibition of 1866, and of the second Paris "Exposition," 1867.) On that occasion also the terms Choripetaleae and Synpetaleae were already used, to distinguish the chief divisions of the Dicotyledoneae, while the sequence of orders, effected then, was also mainly the same as that followed in the present pages, the Apetaleae of Jussieu or Monochlamydeae of De Candolle having been merged into the petaliferous divisions. But the system, built up by these two great masters in botanic science, is in its main features so genuinely natural, that no subsequent research could bring about very material changes, except in the one particular direction above indicated, a design more or less happily carried out from the time of the earlier writings of Brogniart to that of the latest essays of Jean Mueller, because it was felt, that so long as the Monochlamydeae remained isolated and associated with the Gymnospermeae, so long would we have an imperfect natural system. To proceed in chief alterations further than this, would give a less acceptable system. Even among the Thalamiflorae and Calyciflorae, represented in Australia, as placed together by Bentham, we had already not less than 58 genera, which are entirely apetalous or contain species, in which the corolla remains undeveloped, leaving out of consideration numerous absolutely extra-Australian genera wholly or partly apetalous. Again in Haloragis, Myriophyllum, Cotula, Soliva and some other genera the corolla is absent in the non-antheriferous flowers, while even the very commencement of the Candollean arrangement is made with the apetalous genera Clematis, Thalictrum and Anemone, soon followed by the equally well-known Caltha. Furthermore the Euphorbiaceae, which in the train of their alliances must carry with them always the Urticeae, count among their numerous genera more than one third provided with petals. Besides, in Proteaceae the floral envelope may be regarded as homologous to that of the closely allied Lorantheae, with an absence of a calyx, comparable to the suppression of that organ in Diplolaena, Asterolasia and few other thus far exceptional genera. Again, a firm thread runs uninterruptedly through all the orders of Curvembryonatae or Amyliferae, whether placed in Thalamiflorae or Calyciflorae or Monochlamydeae or even as regards Plumbagineae into Corolliflorae; nor is there any real difficulty of finding for the rest of the Monochlamydeae proper places among naturally allied orders of supposed higher organisation. Nevertheless affinity is variously radial, not altogether uniserial, as beautifully demonstrated already by Linné in his map of ordinal alliances of plants, published by Fabricius and Giseke, or as lucidly exhibited as long ago as the middle of the last century by Bernard de Jussieu in the Royal Garden of Trianon, through arranging the plants in a class-ground, a method adopted in our Botanic Garden here also already in 1857. In using the term Calyceae for the three main-divisions of the Monocotyledoneae, the author wishes it to be understood, that it remains an open question, whether the flowers of two of these divisions are to be regarded as mono- or di-chlamydeous. But if in the latter case a petalaceous development is attributed to the flowers, we are then bound by homology to carry the term of petals also to Junceae even and Restiaceae, and a change in the naming of the three great sections would be needful accordingly. In suppressing throughout the epigynous divisions for unison with the perigynous, into which they often pass so gradually, simplification is aimed at, without causing any disarrangement, or without attempting alterations in the great received systems except nominally. Furthermore we should always regard the apocarpic orders as those of the highest development, because the fruit—the final effort of all growth—when multiplying into separate and complete carpel elements, exhibits the most advanced vegetable organisation.

Much could it have been wished, to allot in this systematic census also columns of citation to those of the larger works, which are exclusively devoted to the Australian Flora. But Robert Brown's ever-memorable "Prodromus" records only vascular Acotyledoneae, Monocotyledoneae, some Apetaleae

and a portion of the Synpetaleae, as far as known in 1810. The two important volumes, issued by Lehmann and his coadjutors as "*Plantae Preissianae*," refer solely to South-West Australian species. Sir Joseph Hooker's grand "*Flora Tasmaniae*" became necessarily limited to the plants of that island, although prefaced by an extensive general essay.

A future re-edition of this census from augmented material, in ampler paginal form, might afford space also for noting these works. The "*Flora Tasmaniae*" will replace however the "*Flora Australiensis*" in the quotation-column of the second part of this census, in which all the "*Evasculares*" of Australia are to be enumerated, the total of them having risen already to about 3750 recorded species. For, be it understood, that the genial and learned Director of the Royal Botanic Garden of Kew has laid a firm foundation for the system of evascular Acotyledoneae of All-Australia in his *Tasmanian Flora*, that work being the first, in which any extensive and connectedly elaborated account of Mosses, Lichens, Fungs and Algs of Australia has been presented.

It was beyond the scope of these pages, to extend the statistics here given; otherwise the writer would have gladly assigned geographic columns also to Europe, Asia, Africa, North and South America for indicating the respective range of many Australian species of plants into other regions of the globe. Whoever may be interested in such phytogeographic subjects is referred to an able article, written by Dr. Engler, Director of the Botanic Garden of Kiel, for the volume of 1881 of his "*Jahrbücher*," or to the print of a discourse, delivered a few months ago before the School of Mines at Ballarat. To sum up some of the results of this statistic volume, tables are appended, setting forth the number of the species in each order as well as the grand totals arrived at, clearly immigrated plants being excluded from these pages throughout. Future phytological explorations are not likely, to add beyond several hundred specific forms, considered in conservative limitations, to those now adduced; but extensive additions are sure to be made yet to the records of regional distribution. Indeed even while this first part of the work went through the press, so many new localities became known, as to render an appendix now already necessary. For these addenda I am largely indebted either in notations or material to the Rev. Dr. Woolls, Professor Tate, the Rev. B. Scortechini and Mr. F. M. Bailey, while the supplemental notes concerning West Australian plants emanated chiefly from gatherings, which we owe to the Hon. J. Forrest, C.M.G. It should however be observed, that the whole of the vast additional collections, which accumulated in our Botanic Museum since the volumes of the "*Flora Australiensis*" successively appeared, could not receive in all instances timely close and critical attention for the elaboration of the present work, though its author had the advantage of departmental help from Mr. G. Luehmann in preliminary sorting and comparing of much of the supplemental material; and here it is an apt place, to recognise likewise the aid, afforded by Mr. Léon Henry, in writing out primarily the requisite notes from the "*Flora*" and the "*Fragmenta*." Nearly one hundred species more might have been taken into account already, for estimating the number of species in various orders; but the samples hitherto secured did not admit of obtaining accurate specific data, though the ordinal position could be recognised. Thus we know, that already some access to Anonaceae, Menispermaceae, Lauraceae, Euphorbiaceae, Rutaceae and a few other orders could be recorded.

The chronologic references for species, here more extensively carried out, than ever before, amount to nearly 10,000, including those for genera, in this first part of the present publication. The bibliographic works of Pritzels, Pfeiffer and Jackson have done excellent services in these particular inquiries. The position, given to several of the orders on this occasion, is to some extent assailable; thus—as well-known—reasons exist for moving Piperaceae to Nymphaeaceae, Aristolochiaceae and Nepenthaceae to Sarracenaceae, Viniferae to Araliaceae, Thymeleae to Rhamnaceae, Balanophoreae and Podostemoneae to Haloragaceae, Droseraceae to Saxifrageae, Rubiaceae to Loganiaceae, Fluviales to Hydrocharideae, Casuarineae to Coniferae, although the two latter are neither anatomically nor morphologically approaching each other, notwithstanding some deceptive external resemblance, in which however Exocarpus also shares, not to speak of a few other genera. But in considerations like these we should not insist in a dogmatic spirit on the full acceptance of hitherto recognised systematic arrangements, particularly as not yet all the forms of the world's vegetation are known, and as at any moment the discovery of a new plant may turn the scale in weighing the affinity of its allies.

In incurring the responsibility of restoring some generic names, which had sunk almost into oblivion, the writer has been guided by the impartial rules of strict priority, feeling assured, that phytographers will become quite as quickly accustomed to changes from *Stylidium* to *Candollea*, or *Ionidium* to *Hybanthus*, or *Chenolea* to *Bassia*, as they became reconciled to the long forgotten appellations of *Trema*, *Hevea*, *Centipeda*, *Galeola*, *Floriscopa*, *Nunnezharia* and others, thereby discarding names, sanctioned by lengthened use and familiar to all of us, with a readiness as surprising as universal. To alterations or restitutions of specific names shall only be alluded here with a few words. Why Lamarek for an Eleusine selected designedly the appellation *E. cruciata*, or why Schleiden called a *Wolffia* purposely *W. Michellii*, can be as easily perceived as the advisability of withdrawing the specific name *antarctica* from a *Vitis*, *Cymodocea* or *Dicksonia*. Where authors bestowed several names

at the same time on a species, it should be free to those, who effect the reductions, to choose a collective designation for the consolidated species; cases in point are afforded by *Euphrasia Brownii*, *Fimbristylis communis*, *Pappophorum commune*, *Danthonia penicillata*.

As to the circumscription of species themselves, it will require yet much of assiduous research of future generations for arriving at a full understanding of the diagnostic value, which has to be attached respectively to the thousands of specific and therefore original forms in the Australian and any other vegetation. Of the plants, enumerated as species in the present pages, a considerable number may prove, that they owe their particularity to hybridism. In such cases they should occupy a subordinate position, and should also here, as elsewhere, receive their double parental name;—thus *Lasiopetalum Tepperi* would require to have its appellation changed into *Lasiopetalum Baueri* × *discolor*; but as the instances of ascertained hybridism among our native plants are as yet so very few, it was deemed best to admit at present the very limited number of known bastards under ordinary specific rank.

The geographical limitations in this work coincide with the political boundaries of the colonial territories, except that the tropic of Capricorn eastward to the 138th degree separates what is here called Northern Australia (N.A.), from the South- and West-Australian extratropic possessions. Such geographic segregations are necessarily quite arbitrary, though they serve our present purpose of assigning to each of the colonial divisions of Australia its number of specified plants; the limitation is the same as that adopted in the “*Flora Australiensis*,” and as regards abbreviations also identical with the method of indications, chosen for the list of Australian trees in 1866. Here may it further be remarked, that the plants of the small and isolated insular spot, called Norfolk-Island, have been counted with those of N.S.W., to whose dominions this isle as well as Lord Howe’s Island politically belongs. To draw the species into physiographic and regional complexes must be the work of future periods, when climatic and geologic circumstances throughout Australia shall have become more extensively known. The geographic columns in these pages indicate simply the occurrence of plants within any of the colonial areas, but have been extended even to such species, which merely may pass boundary-lines. In this way are noted for W.A. and S.A. such tropical plants, as reach not beyond Shark-Bay or the vicinity of the MacDonnell-Range; in the same manner Victoria is credited with those plants of New South Wales, which barely advance into East-Gippsland, while South-Australia obtains credit for several Tasmanian plants, which are confined to very limited stretches of country, chiefly in the extreme south-east of that province; and again a few of the New England plants are recorded for Queensland, although they pass merely on to the nearest adjacent ranges across the boundary. At the whole however it can be foreseen, that many a species, still accepted as genuine for these lists, will in the course of further and more facilitated research be eliminated from any future editions of this work, though still larger arrays, containing many a novelty, will have to be mustered for systematic rolls.

The lines of demarkation between truly indigenous and more recently immigrated plants can no longer in all cases be drawn with precision; but whereas *Alchemilla vulgaris* and *Veronica serpyllifolia* were found along with several European Carices in untrodden parts of the Australian Alps during the author’s earliest explorations, *Alchemilla arvensis* and *Veronica peregrina* were at first only noticed near settlements. The occurrence of *Arabis glabra*, *Geum urbanum*, *Agrimonia Eupatoria*, *Eupatorium cannabinum*, *Carpesium cernuum* and some others may therefore readily be disputed as indigenous, and some questions concerning the nativity of various of our plants will probably remain for ever involved in doubts. While concluding these introductory remarks, it is incumbent on the writer, to acknowledge in grateful terms the consideration, shown him by the Hon. James Macpherson Grant, M.L.A., the Ministerial Chief of the Department, in allowing this work to be issued under the auspices of the Victorian Government and at departmental expenditure. Encouraged by generous support, the author will cheerfully continue through such time, as may still be allotted for his worldly career, to devote his strength and resources also in future mainly for the furtherance of Australian phytography, on which much of his efforts were concentrated through more than a third of a century; and as these enquiries are largely carried on without worldly gain, he feels free, to claim the co-operation of any educated and high-minded colonists in these great dominions of the British Crown for promoting studies, which cannot fail to be of great industrial advantage at all future time, which are calculated to advance continuously the cause of education in this part of the world also, which must far and wide contribute to intellectual enjoyments, and above all should lead to religious reverence of that Supreme Godly power, so gloriously revealed in nature’s wondrous works.

DICOTYLEDONEAE.

Ray, Method. Plant. nova 2 (1682).

CHORIPETALEAE HYPOGYNAE.

F. v. Mueller, native plants of Victoria I, 1 (1879).

RANUNCULACEAE.

A. L. de Jussieu, Rec. de l'Acad. des sc. (1773) from B. de Jussieu (1759).

CLEMATIS, Linné, gen. pl. 163 (1737) from l'Ecluse (1576).

C. aristata, R. Brown in De Candolle, syst. veg. I, 147 (1818) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,6	M.fr.X,2.
C. pubescens, Huegel, enum. pl. Nov. Holl. austr. occ. I (1837) ...	W.A.	—	—	—	—	—	—	B.f.I,6	
C. glycinoides, De Candolle, syst. veg. I, 145 (1818) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I,7	M.fr.X,2.
C. microphylla, De Candolle, syst. veg. I, 147 (1818) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,7	M.fr.X,2; XI,20,27.
C. Fawcettii, F. v. Mueller, fragm. phytogr. Austr. X, 1 (1876) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.X,1.

ANEMONE, Tournefort, inst. rei herb. 275, t. 147 (1700) from Hippocrates, Theophrastus and Dioscorides.

A. crassifolia, Hooker, icones pl. t. 257 (1840) ...	—	—	—	—	—	—	—	B.f.I,8	M.fr.X,1.
--	---	---	---	---	---	---	---	---------	-----------

MYOSURUS, Dillenius, catal. pl. Gie-s. 106, t. 4 (1719) from Tabernaemontanus (1588). (Myosuros).

M. minimus, Linné, sp. pl. 234 (1753) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I,8	M.fr.X,2; XI,27.
---	---	------	---	----	--------	---	---	---------	------------------

RANUNCULUS, Tournefort, inst. 285, t. 149 (1700) from Bock (1552).

R. aquatilis, Dodonæus stirp. hist. pempt. 387 (1583) ...	—	S.A.	T.	V.	—	—	—	B.f.I,10	
R. Millani, F. v. Mueller in Hook. Kew misc. VII, 358 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.f.I,10	M.fr.X,2.
R. anemoneus, F. v. Mueller in Trans. Phil. Soc. Vict. I, 97 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.f.I,11	
R. Gunnianus, Hooker, Journ. of Bot. I, 245, t. 233 (1834) ...	—	—	T.	V.	N.S.W.	—	—	B.f.I,11	
R. lappaceus, Smith in Rees's Cyclopædia, XXIX (1815) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,12	M.fr.X,2.
R. Muelleri, Benth. Austr. I, 13 (1863) ...	—	—	—	V.	N.S.W.	—	—	B.f.I,13	
R. rivularis, Bks. & Sol. in G. Forst. fl. ins. austr. prodr. 90 (1786) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,13	M.fr.X,2.
R. hirtus, Bks. & Sol. in G. Forst. fl. ins. austr. prodr. 90 (1786) ...	W.A.	—	—	V.	N.S.W.	Q.	—	B.f.I,13	M.fr.X,2.
R. parviflorus, Linné, sp. pl. sec. ed. 780 (1762) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,14	M.fr.X,2; XI,27.

CALTHA, Ruppius, flora Jenensis 119 (1718) from Bock (1552). (Trollius).

C. introloba, F. v. Mueller in Trans. Phil. Soc. Vict. I, 98 (1855) ...	—	—	T.	V.	N.S.W.	—	—	B.f.I,15	M.fr.X,2; XI,136.
---	---	---	----	----	--------	---	---	----------	-------------------

NYMPHAEACEAE.

Salisbury in Koenig & Sims's Ann. of Bot. II, 70 (1805).

CABOMBA, Aublet, Hist. pl. Guian. I, 321, t. 124 (1775). (Brasenia, Hydropeltis).

C. peltata, F. v. Mueller, native pl. of Vict. 15 (1874) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.I,60	M.fr.X,77.
--	---	---	---	----	--------	----	---	----------	------------

NYMPHAEA, Tournefort, inst. 260, t. 137 8 (1700) from Theophrastus and Dioscorides.

N. stellata, Willdenow, spec. pl. II, 1153 (1799) ...	—	—	—	—	—	—	N.A.	—	M.fr.II,142; X,77.
N. gigantea, Hooker, Bot. Mag. t. 4647 (1852) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I,61	M.fr.II,141; X,78.

NELUMBO, Tournefort, inst. 261 (1700). (Nelumbium).

N. nucifera, Gaertner, de fruct. I, 73 (1788) ...	—	—	—	—	—	—	Q.	N.A.	B.f.I,62	M.fr.X,77.
---	---	---	---	---	---	---	----	------	----------	------------

DILLENIACEAE.

Salisbury, Parad. Lond. I, 73 (1801).

WORMIA, Rottboell, Nye Saml. Vid. Selsk. Skrivt. II, 532 (1783).

W. alata, R. Brown in De Cand. syst. veg. I, 434 (1818) ...	—	—	—	—	—	Q.	—	B.f.I,16	M.fr.VIII,124.
---	---	---	---	---	---	----	---	----------	----------------

HIBBERTIA, Andrews, Bot. Reposit. t. 126 (1800). (Candollea, Pleurandra, Hemistemma, Adrastaea, Ochrolasia, Hemistephus, Huttia.)

H. Banksii, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	—	—	—	—	—	Q.	—	B.f.I,21	M.fr.II,1; XI,92.
H. Brownii, Benth. Austr. I, 21 (1863) ...	—	—	—	—	—	Q.	—	B.f.I,21	M.fr.II,1.
H. dealbata, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	—	—	—	—	—	—	N.A.	B.f.I,21	M.fr.II,1.
H. candicans, Benth. Austr. I, 21 (1863) ...	—	—	—	—	—	Q.	N.A.	B.f.I,21	
H. Benthami, H. angustifolia, Benth. fl. Austr. I, 21 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I,21	M.fr.II,1.
H. Muelleri, Benth. Austr. I, 21 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I,21	
H. ledifolia, Benth. Austr. I, 22 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I,22	
H. verrucosa, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	W.A.	—	—	—	—	—	—	B.f.I,22	M.fr.XI,92.
H. spicata, F. v. Mueller, fragm. II, 1 (1860) ...	W.A.	—	—	—	—	—	—	B.f.I,22	M.fr.II,1; XI,20,92.
H. polystachya, Benth. Austr. I, 22 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I,22	M.fr.XI,92.
H. rhadinopoda, F. v. Mueller, fragm. XI, 91 (1880) ...	W.A.	—	—	—	—	—	—	—	M.fr.XI,91.
H. furfuracea, Benth. Austr. I, 23 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I,23	M.fr.I,161; XI,92.
H. hypericoides, F. v. Mueller, pl. of Vict. I, 13, implied (1860) ...	W.A.	—	—	—	—	—	—	B.f.I,23	
H. microphylla, Steudel in Lehmann pl. Preiss. I, 273 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I,23	M.fr.I,217.
H. recurvifolia, Benth. Austr. I, 24 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I,24	
H. lineata, Steudel in Lehmann pl. Preiss. I, 272 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I,24	M.fr.XI,92.
H. acerosa, Benth. Austr. I, 24 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I,24	M.fr.XI,92.
H. aurea, Steudel in Lehmann pl. Preiss. I, 272 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I,25	M.fr.XI,92.
H. crassifolia, Benth. Austr. I, 25 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I,25	
H. nitida, F. v. Mueller, pl. of Vict. I, 15 (1860) ...	—	—	—	—	N.S.W.	—	—	B.f.I,25	
H. bracteata, Benth. Austr. I, 26 (1863) ...	—	—	—	—	N.S.W.	—	—	B.f.I,25	
H. densiflora, F. v. Mueller, pl. of Vict. I, 15 (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.I,26	M.fr.VII,125; XI,92.
H. hirsuta, Benth. Austr. 26 (1863) ...	—	—	—	T.	—	—	—	B.f.I,26	
H. synandra, F. v. Mueller, fragm. IV, 151 (1864) ...	—	—	—	—	—	Q.	—	—	M.fr.IV,151,
H. stricta, R. Brown in De Candolle, syst. veg. I, 422 (1818) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I,27	
H. humifusa, F. v. Mueller, pl. of Vict. I, 16, t. Suppl. I (1860) ...	—	—	—	V.	—	—	—	B.f.I,27	

H. Billardieri, F. v. Mueller, pl. of Vict. I, 14 (1860) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 28	M.fr. IV, 116.
H. gracilipes, Benth. Fl. Austr. I, 28 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 28	M.fr. XI, 92.
H. acicularis, F. v. Mueller, pl. of Vict. I, 17 (1860) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 29	M.fr. XI, 93.
H. mucronata, F. v. Mueller, pl. of Vict. I, 17 (1860) ...	W.A.	—	—	—	—	—	—	B.f.I, 29	M.fr. XI, 92.
H. hermannifolia, De Candolle, syst. veg. I, 431 (1818) ...	—	—	—	—	N.S.W.	—	—	B.f.I, 30	—
H. velutina, R. Brown in Benth. Fl. Austr. I, 30 (1863) ...	—	—	—	—	—	Q.	—	B.f.I, 30	—
H. oblongata, R. Brown in De Candolle, syst. veg. I, 431 (1818) ...	—	—	—	—	—	—	N.A.	B.f.I, 30	M.fr. XI, 94.
H. melhanoides, F. v. Mueller, fragm. IV, 116 (1864) ...	—	—	—	—	—	Q.	—	—	M.fr. IV, 116.
H. tomentosa, R. Brown in De Candolle, syst. veg. I, 432 (1818) ...	—	—	—	—	—	—	N.A.	B.f.I, 30	—
H. cistifolia, R. Brown in De Candolle, syst. veg. I, 431 (1818) ...	—	—	—	—	—	—	N.A.	B.f.I, 30	—
H. echiifolia, R. Brown in Benth. Fl. Austr. I, 31 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 31	—
H. scabra, R. Brown in Benth. Fl. Austr. I, 31 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 31	—
H. lepidota, R. Brown in De Candolle, syst. veg. I, 432 (1818) ...	—	—	—	—	—	—	N.A.	B.f.I, 31	—
H. vestita, Cunningham in Benth. Fl. Austr. I, 31 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 31	—
H. serpyllifolia, R. Brown in De Cand. syst. veg. I, 430 (1818) ...	—	—	—	T.	V.	N.S.W.	—	B.f.I, 32	—
H. pedunculata, R. Brown in De Cand. syst. veg. I, 430 (1818) ...	—	—	—	—	V.	N.S.W.	—	B.f.I, 32	—
H. ochrolasia, Benth. Fl. Austr. I, 32 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 32	M.fr. VII, 125; XI, 93.
H. angustifolia, Salisbury, parad. Lond. 73 (1807) ...	—	—	—	T.	V.	—	—	B.f.I, 33	—
H. fasciculata, R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 33	—
H. virgata, R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 34	M.fr. XI, 94.
H. inclusa, Benth. Fl. Austr. I, 34 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 34	—
H. rostellata, Turcz. in Bull. Soc. Mosc. XXII, part III, 8 (1849) ...	W.A.	—	—	—	—	—	—	B.f.I, 34	M.fr. XI, 93.
H. glomerata, Benth. Fl. Austr. I, 34 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 34	—
H. argentea, Steudel in Lehmann, pl. Preiss. I, 268 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 35	—
H. pilosa, Steudel in Lehmann, pl. Preiss. I, 272 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 35	—
H. montana, Steudel in Lehmann, pl. Preiss. I, 270 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 35	M.fr. XI, 94.
H. linearis, R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.I, 36	—
H. diffusa, R. Brown in De Candolle, syst. veg. I, 429 (1818) ...	—	—	—	—	V.	N.S.W.	—	B.f.I, 36	—
H. saligna, R. Brown in De Candolle, syst. veg. I, 427 (1818) ...	—	—	—	—	—	N.S.W.	—	B.f.I, 37	M.fr. XI, 94.
H. volubilis, Andrews in Bot. Reposit. t. 126 (1800) ...	—	—	—	—	—	N.S.W.	Q.	B.f.I, 37	M.fr. IV, 116; VII, 125; XI, 94.
H. grossularifolia, Salisbury, parad. Lond. t. 73 (1806) ...	W.A.	—	—	—	—	—	—	B.f.I, 37	M.fr. XI, 94.
H. dentata, R. Brown in De Candolle, syst. veg. I, 426 (1818) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.I, 38	M.fr. VII, 125; XI, 94.
H. Cunninghamii, Aiton in Hook. Bot. Mag. t. 318 (1832) ...	W.A.	—	—	—	—	—	—	B.f.I, 39	M.fr. XI, 93.
H. perfoliata, Huegel, enum. pl. Nov. Holl. austr. occ. 3 (1837) ...	W.A.	—	—	—	—	—	—	B.f.I, 38	M.fr. XI, 93.
H. bracteosa, Turczaninow, in Bull. Soc. Mosc. XXV, 140 (1852) ...	W.A.	—	—	—	—	—	—	B.f.I, 38	M.fr. XI, 94.
H. longifolia, F. v. Mueller, fragm. IV, 115 (1864) ...	—	—	—	—	—	Q.	—	—	M.fr. IV, 115.
H. oenotheroides, F. v. Mueller, fragm. VII, 37 (1870) ...	—	—	—	—	—	Q.	—	—	M.fr. VII, 37.
H. glaberrima, F. v. Mueller, fragm. III, 1 (1862) ...	—	S.A.	—	—	—	—	—	B.f.I, 39	M.fr. III, 1; XI, 94.
H. Mylnei, Benth. Fl. Austr. I, 39 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 39	—
H. lasiopus, Benth. Fl. Austr. I, 40 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 40	M.fr. XI, 94.
H. potentilliflora, F. v. Mueller in Benth. Fl. Austr. I, 40 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 40	M.fr. XI, 94.
H. pungens, Benth. Fl. Austr. I, 40 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 40	—
H. nutans, Benth. Fl. Austr. I, 40 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 40	—
H. leptopus, Benth. Fl. Austr. I, 41 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 41	—
H. stellaris, Endlicher in Hueg. enum. 3 (1837) ...	W.A.	—	—	—	—	—	—	B.f.I, 41	M.fr. XI, 93.
H. obcuneata, Salisbury, parad. Lond. t. 73 (1807) ...	W.A.	—	—	—	—	—	—	B.f.I, 42	M.fr. XI, 94.
H. glomerosa, F. v. M.; Candollea glomerosa, Benth. Fl. Austr. I, 43 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 43	—
H. teretifolia, F. v. Mueller, fragm. IV, 117 (1864) ...	W.A.	—	—	—	—	—	—	B.f.I, 43	M.fr. IV, 117; XI, 95.
H. desmophylla, F. v. Mueller, fragm. XI, 95 (1880) ...	W.A.	—	—	—	—	—	—	B.f.I, 43	M.fr. XI, 95.
H. helianthemoides, F. v. M.; Candollea helianthemoides, Turcz. in Bull. Mosc. XXII, part III, 8 (1849) ...	W.A.	—	—	—	—	—	—	B.f.I, 43	—
H. depressa, Steudel in Lehm. pl. Preiss. I, 268 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 44	M.fr. XI, 95.
H. Huegelii, F. v. Mueller, fragm. XI, 95 (1880) ...	W.A.	—	—	—	—	—	—	B.f.I, 44	M.fr. XI, 95.
H. subvaginata, F. v. Mueller, fragm. XI, 95 (1880) ...	W.A.	—	—	—	—	—	—	B.f.I, 45	M.fr. XI, 95.
H. vaginata, F. v. Mueller, fragm. XI, 96 (1880) ...	W.A.	—	—	—	—	—	—	B.f.I, 45	M.fr. XI, 96.
H. subexcisa, Steudel in Lehmann, pl. Preiss. I, 269 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 46	M.fr. XI, 95.
H. squamosa, Turczaninow in Bull. Soc. Mosc. XXII, part III, 9 (1849) ...	W.A.	—	—	—	—	—	—	B.f.I, 46	—
H. uncinata, F. v. M.; Candollea uncinata, Benth. Fl. Austr. I, 46 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 46	—
H. salicifolia, F. v. Mueller, fragm. I, 161 (1859) ...	—	—	—	—	—	N.S.W.	Q.	B.f.I, 46	M.fr. I, 161; VII, 124; XI, 94.
H. Huttii, F. v. Mueller, fragm. VII, 123 (1870) ...	W.A.	—	—	—	—	—	—	B.f.I, 47	M.fr. VII, 123; XI, 20.
H. Goyderi, F. v. Mueller, fragm. VII, 123 (1870) ...	—	—	—	—	—	—	N.A.	—	M.fr. VII, 123.
TETRACERA, Linné, gen. pl. 345 (1737).									
T. Nordiana, F. v. Mueller, fragm. V, 1 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr. V, 1.
T. Daemeliana, F. v. Mueller, fragm. V, 191 (1866) ...	—	—	—	—	—	Q.	—	—	M.fr. V, 191.
T. Wuthiana, F. v. Mueller, fragm. X, 49 (1876) ...	—	—	—	—	—	Q.	—	—	M.fr. X, 49.
DILLENIA, Linné, gen. pl. 162 (1737).									
D. Andreaana, F. v. Mueller, fragm. V, 175 (1866) ...	—	—	—	—	—	Q.	—	—	M.fr. V, 175.
PACHYNEMA, R. Brown in De Candolle, syst. veg. I, 412 (1818).									
P. junceum, Benth. Fl. Austr. I, 47 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 47	M.fr. X, 124.
P. complanatum, R. Brown in De Cand., syst. veg. I, 412 (1818) ...	—	—	—	—	—	—	N.A.	B.f.I, 48	M.fr. X, 124.
P. dilatatum, Benth. Fl. Austr. I, 48 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 48	M.fr. X, 124.

MAGNOLIACEAE.

J. de St. Hilaire, Expos. fam. nat. II, 74, t. 83-84 (1805).

DRIMYS, R. & G. Forster, char. gen. 83, t. 42 (1776). (Tasmania).

D. aromatica, F. v. Mueller, pl. of Vict. I, 20 (1860) ...	—	—	—	T.	V.	N.S.W.	—	B.f.I, 49	M.fr. VII, 18.
D. membranacea, F. v. Mueller, fragm. V, 175 (1866) ...	—	—	—	—	—	—	Q.	—	M.fr. V, 175; VII, 18.
D. dipetala, F. v. Mueller, pl. of Vict. I, 21 (1860) ...	—	—	—	—	—	N.S.W.	Q.	B.f.I, 49	M.fr. VII, 18.
D. Howeana, F. v. Mueller, fragm. VII, 17 (1870) ...	—	—	—	—	—	N.S.W.	—	—	M.fr. VII, 17.

ANONACEAE.

A. L. de Jussieu, Gen. 283 (1739) from B. de Jussieu (1759).

UVARIA, Linné, Amoen. acad. 404 (1747).															
U. membranacea,	Bentham,	Fl. Austr. I,	51	(1863)	...	...	—	—	—	—	Q.	—	B.fl.I,51		
U. Goezeana,	F. v. Mueller,	fragm. VII,	125	(1871)	...	...	—	—	—	—	Q.	—	—	M.fr.VII,125.	
FITZALANIA, F. v. Mueller, fragm. IV, 33 (1863).															
F. heteropetala,	F. v. Mueller,	fragm. IV,	33	(1863)	...	...	—	—	—	—	Q.	—	B.fl.I,51	M.fr.III,1;VII,126.	
CANANGA, Rumphius, herb. Amboin. II, 195, t. 65 (1741).															
C. odorata,	J. Hooker and Thomson,	fl. Indic. 130	(1855)	...	...	—	—	—	—	—	Q.	—	—	M.fr.VI,1.	
ANCANA, F. v. Mueller, fragm. V, 27 (1865).															
A. stenopetala,	F. v. Mueller,	fragm. V,	27	(1865)	...	...	—	—	—	N.S.W.	Q.	--	—	M.fr.V,27,189;VII,121.	
POLYALTHIA, Blume, Fl. Jav. Anon. 68 (1829).															
P. nitidissima,	Bentham,	Fl. Austr. I,	51	(1863)	...	...	—	—	—	N.S.W.	Q.	—	B.fl.I,51	M.fr.III,2;VII,126.	
POPOWIA, Endlicher, Gen. 831 (1839).															
P. australis,	Bentham,	Fl. Austr. I,	52	(1863)	...	...	—	—	—	—	—	N.A.	B.fl.I,52		
MELODORUM, Loureiro, Fl. Cochinch. I, 351 (1790).															
M. Leichhardtii,	Bentham,	Fl. Austr. I,	52	(1863)	...	...	—	—	—	N.S.W.	Q.	—	B.fl.I,52	M.fr.III,41,VI,2;VII,126.	
M. Uchrii,	F. v. Mueller,	fragm. VI,	2	(1867)	...	...	—	—	—	—	Q.	—	—	M.fr.VI,2.	
M. Maccraei,	F. v. Mueller,	fragm. VI,	176	(1868)	...	...	—	—	—	—	Q.	—	—	M.fr.VI,176,250.	
MILIUSIA, Leschenault in Mém. Genève. V, 213 (1832).															
Species undetermined	...	...	...	...	...	...	—	—	—	—	Q.	—	—		
SACCOPETALUM, Bennett in Horfs. pl. Jav. rar. 115, t. 35 (1838).															
S. Bidwilli,	Bentham,	Fl. Austr. I,	53	(1863)	...	...	—	—	—	—	Q.	—	B.fl.I,53		
S. Brahei,	F. v. Mueller,	fragm. VIII,	159	(1874)	...	...	—	—	—	—	Q.	—	—	M.fr.VIII,159.	
EUPOMATIA, R. Brown in App. Flind. voy. II, 597, t. 2 (1814).															
E. Bennettii,	F. v. Mueller,	fragm. I,	45	(1859)	...	...	—	—	—	N.S.W.	Q.	—	B.fl.I,54	M.fr.I,45;VI,177.	
E. laurina,	R. Brown in App. Flind. voy. II,	597, t. 2	(1814)	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,54	M.fr.I,45;VI,2;VII,126.	

MONIMIEAE.

A. L. de Jussieu in *Ann. du Mus.* XIV, 30 (1809).

DORYPHORA, Endlicher, Gen. 3015 (1837) (Learosa).									
D. sassafras, Endlicher, iconogr. gen. pl., t 10 (1838)	...	...	—	—	—	—	N.S.W.	—	B.fl.V, 283
ATHEROSPERMA, Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806).									
A. moschatum, Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806)	—	—	—	—	—	—	T. V. N.S.W.	—	B.fl.V, 284 M.fr.VIII, 142; X, 106.
DAPHNANDRA, Bentharn, Fl. Austr. V, 285 (1870).									
D. micrantha, Bentharn, Fl. Austr. V, 285 (1870)...	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 285
D. repandula, F. v. Mueller, fragm. X, 105 (1877)	...	...	—	—	—	—	—	Q.	M.fr.X, 105.
MOLLINEDIA, Ruiz & Pavon, prodr. 83, t. 15 (1794). (Kibara, Wilkiea).									
M. Huegeliana, Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855)	—	—	—	—	—	—	N.S.W.	Q.	B.fl.V, 286
M. Wardellii, F. v. Mueller, fragm. V, 153 (1866)	...	...	—	—	—	—	—	Q.	B.fl.V, 287 M.fr.V, 155; VI, 252.
M. loxocarya, Bentharn, Fl. Austr. V, 257 (1870)...	...	...	—	—	—	—	—	Q.	B.fl.V, 287
M. acuminata, F. v. Mueller, fragm. V, 155 (1866)	...	...	—	—	—	—	—	Q.	B.fl.V, 287 M.fr.V, 155.
M. macrophylla, Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855)	—	—	—	—	—	—	N.S.W.	Q.	B.fl.V, 288
M. longipes, Bentharn, Fl. Austr. V, 289 (1870)	...	...	—	—	—	—	—	Q.	B.fl.V, 289
M. laxiflora, Bentharn, Fl. Austr. V, 289 (1870)	...	...	—	—	—	—	—	Q.	B.fl.V, 289
M. pubescens, Bentharn, Fl. Austr. V, 290 (1870)	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 290
HEDYCARYA, R. & G. Forster, char. gen. 127, t. 64 (1776).									
H. Cunninghamii, Tulasne in Archiv. du Mus. hist. nat. VIII, 408 (1855)	—	—	—	—	—	—	V. N.S.W.	Q.	B.fl.V, 291
PALMERIA, F. v. Mueller, fragm. IV, 151 (1864).									
P. scandens, F. v. Mueller, fragm. IV, 152 (1864)	...	...	—	—	—	—	N.S.W.	Q.	B.fl.V, 291 M.fr.IV, 152.
PIPTOCALYX, Oliver in Bentharn's Fl. Austr. V, 292 (1870).									
P. Moorei, Oliver in Bentharn's Fl. Austr. V, 292 (1870)...	...	...	—	—	—	—	N.S.W.	—	B.fl.V, 292 M.fr.X, 106.

MYRISTICAE.

R. Brown, prodr. 399 (1810) from A. L. de Jussieu (1806).

MYRISTICA, Linné, gen. pl. ed. II, 524 (1742).
M. insipida, R. Brown, prodr. 400 (1810) — — — — — Q. N.A. B.f.V, 281

LAURACEAE.

Ventenat, Tabl. I, 245 (1799).

CRYPTOCARYA, R. Brown, prodr. 402 (1810). (Caryodaphne).												
C. Murrayi, F. v. Mueller, fragm. V, 170 (1866) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.V, 295	M.fr.V, 170; X, 253.
C. Mackinnoniana, F. v. Mueller, fragm. V, 169 (1866) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.V, 296	M.fr.V, 169.
C. patentinervis, F. v. Mueller in DeCandolle, prodr. XV, 508 (1864) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 296	M.fr.V, 166.
C. obovata, R. Brown, prodr. 402 (1810) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 296	M.fr.V, 170.
C. glaucescens, R. Brown, prodr. 402 (1810) ...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V, 297	M.fr.V, 165, 166.
C. triplinervis, R. Brown, prodr. 402 (1810) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 297	M.fr.V, 166.
C. cinnamomifolia, Benth. Fl. Austr. V, 298 (1870) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.V, 298	
C. Meissneri, F. v. Mueller, fragm. V, 170 (1866) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 298	M.fr.V, 170.
C. australis, Benth. Fl. Austr. V, 299 (1870) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 299	M.fr.V, 167.
*BEILSCHMIEDIA, Nees in Wallich, pl. Asiat. rarior II, 69 (1831). (Nesodaphne).												
B. obtusifolia, Benth. in B. & H. gen. III, 1521 (1880) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 299	M.fr.V, 166.

ENDIANDRA, R. Brown, prodr. 402 (1810)

E. glauca, R. Brown, prodr. 402 (1810)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 300	
E. hypotephra, F. v. Mueller, fragm. V, 166 (1866)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 301	M.fr.V, 166.
E. discolor, Benth, Fl. Austr. V, 301 (1870)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 301	
E. Sieberi, Nees, Syst. Laurin. 194 (1836)	...	...	...	—	—	—	N.S.W.	—	—	B.fl.V, 301	M.fr.V, 166.
E. virens, F. v. Mueller, fragm. II, 90 (1860)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 302	M.fr.V, 166.
E. Muelleri, Meissner in De Candolle, prodr. XV, 509 (1864)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 302	
E. pubens, Meissner in De Candolle, prodr. XV, 509 (1864)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 302	

CINNAMOMUM, Burman, fl. zeil. 62 (1737).

C. Tamala, C. G. & T. F. Nees, Syst. Laurin. 56 (1836)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 303	M.fr.V, 165.
--	-----	-----	-----	---	---	---	---	----	---	-------------	--------------

LITSEA, Lamarck, Diction. III, 574 (1789). (Glabraria 1771, Tetranthera, Cylicodaphne, Litsaea).

L. chinensis, Lamarck, Diction. III, 574 (1789)	...	...	...	—	—	—	—	Q.	N.A.	B.fl.V, 305	
L. Bindoniana, F. v. Mueller, Tetranthera, fragm. V, 167 (1866)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 305	M.fr.V, 167.
L. Hexanthus, A. L. de Jussieu in Annal. du Mus. VI, 212 (1805)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 305	
L. reticulata, Benth in B. & J. H. gen. pl. III, part I, 161 (1880)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 306	
L. Zeylanica, C. G. & T. F. Nees, in Amoën. Bonn. I, 58t. 5 (1823)	...	...	...	—	—	—	—	Q.	—	B.fl.V, 307	
L. dealbata, Nees, Syst. Laurin. 630 (1836)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 307	

CASSYTHA, Osbeck in Linné, sp. pl. 35 (1753).

C. nodiflora, Meissner in De Candolle, prodr. XV, 252 (1864)	...	...	...	W.A.	—	—	—	—	—	B.fl.V, 309	
C. glabella, R. Brown, prodr. 404 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V, 309
C. flava, Nees in Lehmann, pl. Preiss. I, 620 (1845)	...	...	...	W.A.	—	—	—	—	—	B.fl.V, 310	
C. pubescens, R. Brown, prodr. 404 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V, 310
C. phaeolasia, F. v. Mueller, fragm. V, 167 (1866)	...	...	...	—	—	—	V.	N.S.W.	—	B.fl.V, 310	
C. paniculata, R. Brown, prodr. 404 (1810)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 311
C. filiformis, Linné, sp. pl. 35 (1753)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V, 311
C. melantha, R. Brown, prodr. 404 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.fl.V, 311	M.fr.V, 167.
C. micrantha, Meissner in De Candolle, prodr. XV, 256 (1864)	...	...	...	W.A.	—	—	—	—	—	B.fl.V, 312	
C. racemosa, Nees in Lehmann, pl. Preiss. I, 621 (1845)	...	...	...	W.A.	—	—	—	—	Q.	B.fl.V, 312	
C. pomiformis, Nees in Lehmann, pl. Preiss. I, 620 (1845)	...	...	...	W.A.	—	—	—	—	—	B.fl.V, 313	

HERNANDIA, Plumier, Gen. 6, t. 40 (1703).

H. peltata, Meissner in De Candolle, prodr. XV, 263 (1864)	...	...	...	—	—	—	—	—	Q.	—	B.fl.V, 314
H. bivalvis, Benth, Fl. Austr. V, 314 (1870)	...	...	...	—	—	—	—	—	Q.	—	B.fl.V, 314

MENISPERMEAE.

A. L. de Jussieu, Gen. 284 (1789).

TINOSPORA, Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 35 (1851).

T. smilacina, Benth in Journ. Linn. Soc. V, suppl. II, 2 (1861)	...	...	...	—	—	—	—	Q.	N.A.	B.fl.I, 55	M.fr.IX, 82.
T. Walcottii, F. v. Mueller in Benth's Fl. Austr. I, 56 (1863)	...	...	...	—	—	—	—	—	N.A.	B.fl.I, 56	

FAWCETTIA, F. v. Mueller, fragm. X, 93 (1877).

F. tinoporoides, F. v. Mueller, fragm. X, 93 (1877)	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.X, 93; XI, 136.
---	-----	-----	-----	---	---	---	--------	---	---	---	----------------------

COCCULUS, De Candolle, sys. veg. I, 515 (1818) from C. Bauhin (1623). (Pericampylus, Legnephora).

C. Moorei, F. v. Mueller, fragm. I, 162 (1859)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 56
--	-----	-----	-----	---	---	---	---	--------	----	---	------------

HYPSPERPA, Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 36 (1851). (Selwynia).

H. Selwyni, F. v. Mueller, fragm. IX, 82 (1875)	...	...	...	—	—	—	—	—	Q.	—	M.fr.IV, 153.
---	-----	-----	-----	---	---	---	---	---	----	---	---------------

TRISTICHOCALYX, F. v. Mueller, fragm. IV, 27 (1863).

T. pubescens, F. v. Mueller, fragm. IV, 27 (1863)	...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 58
T. diffusus, Miers, contrib. to Bot. III, 286 (1871)	...	...	...	—	—	—	—	—	Q.	—	M.fr.IV, 27.

SARCOPETALUM, F. v. Mueller, pl. of Vict. I, 26 (1860).

S. Harveyanum, F. v. Mueller, pl. of Vict. 27 and 221 (1860)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I, 57
--	-----	-----	-----	---	---	---	----	--------	----	---	------------

LEICHHARDTIA, F. v. Mueller, fragm. X, 67 (1876).

L. clamboides, F. v. Mueller, fragm. X, 68 (1876)	...	...	...	—	—	—	—	—	Q.	—	M.fr.X, 68.
---	-----	-----	-----	---	---	---	---	---	----	---	-------------

STEPHANIA, Loureiro, Fl. Cochinch. II, 608 (1790).

S. hernandifolia, Walpers, report bot. syst. I, 96 (1842)	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.I, 57
---	-----	-----	-----	---	---	---	----	--------	----	------	------------

PACHYGONE, Miers in Ann. and Mag. nat. hist. sec. ser. VII, 37 (1851).

P. Hullsii, F. v. Mueller, fragm. IX, 81 (1875)	...	...	...	—	—	—	—	—	Q.	N.A.	—
---	-----	-----	-----	---	---	---	---	---	----	------	---

PLEOGYNE, Miers in Taylor's Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851). (Microclisia).

P. Cunninghami, Miers in Ann. & Mag. of nat. hist. sec. ser. VII, 37 (1851)	...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 59
---	-----	-----	-----	---	---	---	---	---	----	---	------------

CARRONIA, F. v. Mueller, fragm. IX, 84 (1875).

C. multiseptata, F. v. Mueller, fragm. IX, 171 (1875)	...	...	...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 84.
---	-----	-----	-----	---	---	---	---	--------	---	---	--------------

ADELIOPSIS, Benth & J. Hooker, Gen. pl. I, 436 (1862).

A. decumbens, Benth, Fl. Austr. I, 59 (1863)	...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 59
--	-----	-----	-----	---	---	---	---	---	----	---	------------

PAPAVERACEAE.

A. L. de Jussieu, Gen. 235 (1789) from B. de Jussieu (1759).

PAPAVER, Tournefort, inst. 237, t. 119 (1700).

P. aculeatum, Thunberg, Fl. Capensis 431 (1813)	...	...	...	—	—	—	S.A.	T.	V.	N.S.W.	Q.
---	-----	-----	-----	---	---	---	------	----	----	--------	----

CAPPARIDEAE.

Ventenat, Tabl. III, 118 (1799).

CLEOME, Linné, syst. nat. 9 (1735); Linné, gen. 200 (1737). (Polanisia).

C. oxalidea, F. v. Mueller, fragm. I, 69 (1858)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 90
C. tetrandra, Banks in De Candolle prodr. I, 240 (1824)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 90
C. viscosa, Linné, sp. pl. 672 (1753)	...	...	...	—	—	—	S.A.	—	—	N.S.W.	Q.

- ROEPERLA, F. v. Mueller in Hooker's Kew Misc. IX, 15 (1857). (Tetratelia).
 R. clemoides, F. v. Mueller in Hooker's Kew Misc. IX, 15 (1857) — — — — — Q. N.A. B.f.I, 91 M.fr. V, 105; IX, 174.
 EMBLINGIA, F. v. Mueller, fragm. II, 2 (1860).
 E. calceoliflora, F. v. Mueller, fragm. II, 3 (1860) ... W.A. — — — — — B.f.I, 92 M.fr. II, 3; XI, 20.
 CADABA, Forskael, Fl. Aeg. Arab. 67 (1775).
 C. capparoides, De Candolle, prodr. I, 244 (1824) ... — — — — — N.A. B.f.I, 92
 CAPPARIS, Tournefort, inst. 261, t. 139 (1700) from Theophrastos, Dioscorides and Plinius. (Busbeckea, Busbeckea).
 C. umbellata, R. Brown in De Candolle, prodr. I, 247 (1824) ... — — — — — Q. N.A. B.f.I, 93 M.fr. V, 104; IX, 172.
 C. lasiantha, R. Brown in De Candolle, prodr. I, 247 (1824) ... — S.A. — — — — — N.S.W. Q. N.A. B.f.I, 94 M.fr. IX, 173.
 C. quiniflora, De Candolle, prodr. I, 247 (1824) ... — — — — — Q. N.A. B.f.I, 94 M.fr. IX, 173.
 C. spinosa, Linné, sp. pl. 503 (1753) ... — — — — — W.A. — — — — — N.A. B.f.I, 94 M.fr. I, 143; V, 104; IX, 173.
 C. sarmentosa, Cunningham in Benth. Fl. Austr. I, 95 (1863) ... — — — — — Q. — B.f.I, 95 M.fr. V, 104; IX, 173.
 C. ornans, F. v. Mueller in Benth. Fl. Austr. I, 95 (1863) ... — — — — — Q. — B.f.I, 95 M.fr. IX, 173.
 C. nobilis, F. v. Mueller in Benth. Fl. Austr. I, 95 (1863) ... — — — — — N.S.W. Q. — B.f.I, 95 M.fr. I, 163; V, 104.
 C. canescens, Banks in De Candolle, prodr. I, 246 (1824) ... — — — — — Q. — B.f.I, 96 M.fr. IX, 173.
 C. lucida, R. Brown in Benth. Fl. Austr. I, 96 (1863) ... — — — — — Q. N.A. B.f.I, 96 M.fr. I, 163; V, 104; IX, 173.
 C. Mitchellii, Lindley in Mitchell's Three Exped. I, 315 (1838) ... — S.A. — — — — — V. N.S.W. Q. N.A. B.f.I, 96 M.fr. IX, 173.
 C. loranthifolia, Lindley in Mitchell's Trop. Austr. 220 (1848) ... — — — — — N.S.W. Q. — B.f.I, 97
 C. umbonata, Lindley in Mitchell's Trop. Austr. 257 (1848) ... — — — — — Q. N.A. B.f.I, 97 M.fr. IX, 173.
 C. uberiflora, F. v. Mueller, fragm. IX, 172 (1875) ... — — — — — Q. — — M.fr. IX, 172.
 C. Shanesiana, F. v. Mueller, fragm. X, 94 (1877) ... — — — — — Q. — — M.fr. X, 94.
 C. humistrata, F. v. Mueller, fragm. V, 156 (1866) ... — — — — — Q. — — M.fr. V, 156.
 C. Thozetiana, F. v. Mueller, fragm. V, 104 (1866) ... — — — — — Q. — — M.fr. V, 156.
 APOPHYLLUM, F. v. Mueller in Hooker's Kew Misc. IX, 307 (1857).
 A. anomalum, F. v. Mueller in Hooker's Kew Misc. IX, 307 (1857) — — — — — N.S.W. Q. N.A. B.f.I, 97 M.fr. IX, 173; XI, 27.

CRUCIFERAE.

A. L. de Jussieu, Gen. 237 (1789) from B. de Jussieu (1759).

- NASTURTIUM, R. Brown in W. T. Aiton's hortus Kewensis IV, 110 (1812) from Linné, syst. nat. 9 (1735).
 N. terrestre, R. Brown in Aiton's hort. Kewens. IV, 110 (1812) — S.A. T. V. N.S.W. Q. — B.f.I, 66 M.fr. XI, 60.
 BARBARAEA, Beckmann, Lex. botan. 33 (1801). (Barbarea).
 B. vulgaris, R. Brown in De Candolle, prodr. I, 140 (1824) ... — — — — — T. V. N.S.W. — — B.f.I, 66 M.fr. VIII, 142; XI, 6, 60.
 ARABIS, Linné, gen. pl. 198 (1737) from Dalechamps (1554). (Turritis).
 A. glabra, Crantz, stirp. Austriac. 36 (1767) ... — — — — — V. N.S.W. — — B.f.I, 67
 CARDAMINE, Tournefort, inst. 224, t. 109 (1700) from l'Ecluse (1576).
 C. stylosa, De Candolle, regn. veg. syst. nat. II, 24 (1821) ... — — — — — T. V. N.S.W. Q. — B.f.I, 68 M.fr. VII, 19; XI, 60.
 C. dictyosperma, Hooker, Journ. of Bot. I, 246 (1834) ... — — — — — T. V. N.S.W. — — B.f.I, 68 M.fr. VII, 19; X, 6, 60.
 C. radicata, J. Hooker in icones pl. t. 882 (1852) ... — — — — — T. — — — — — B.f.I, 69 M.fr. XI, 6.
 C. laciniata, F. v. Mueller in Transact. Phil. Soc. Vict. I, 34 (1854) ... — S.A. — — — — — V. N.S.W. — — B.f.I, 69 M.fr. XI, 27, 60.
 C. hirsuta, Linné, sp. pl. 655 (1753) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.f.I, 70 M.fr. VII, 19; XI, 60.
 C. eustylis, F. v. Mueller in Transact. Vict. Inst. I, 114 (1854) ... — S.A. — — — — — V. N.S.W. Q. N.A. B.f.I, 71 M.fr. VII, 19; XI, 60.
 ALYSSUM, Tournefort, inst. 217 (1700) from l'Ecluse (1576). (Meniocus, Alysson).
 A. minimum, Pallas, Reise Prov. Russ. Reich. III, 746 (1776) ... W.A. S.A. — — — — — V. N.S.W. Q. N.A. B.f.I, 71 M.fr. XI, 27, 60.
 WILKIA, Scopoli, introd. hist. nat. 317 (1777). (Malcolmia).
 W. Africana, F. v. Mueller, native plants of Vict. I, 33 (1879) ... — — — — — V. N.S.W. — — — — — M.fr. VII, 18.
 SISYMBRIUM, Tournefort, inst. 225, t. 109 (1700). (Blennodia).
 S. filifolium, F. v. Mueller in Transact. Phil. Soc. Vict. I, 34 (1854) ... — S.A. — — — — — N.S.W. — — B.f.I, 73
 S. trisetum, F. v. Mueller in Transact. Vict. Inst. I, 114 (1854) ... W.A. S.A. — — — — — V. N.S.W. Q. — B.f.I, 74 M.fr. VII, 20; XI, 60.
 S. nasturtioides, F. v. Mueller in Transact. Vict. Inst. I, 115 (1854) ... — S.A. — — — — — V. N.S.W. — — B.f.I, 74
 S. Richardsii, F. v. Mueller, fragm. X, 105 (1877) ... — — — — — W.A. S.A. — — — — — — — M.fr. X, 105.
 S. eremigenum, F. v. Mueller, fragm. II, 143 (1861) ... — — — — — — — N.S.W. Q. — B.f.I, 74 M.fr. II, 143.
 S. Lucae, F. v. Mueller, fragm. XI, 59 (1879) ... — — — — — — — V. N.S.W. — — — — — M.fr. XI, 59.
 S. cardaminoides, F. v. Mueller in Benth. Fl. Austr. I, 75 (1863) ... — S.A. — — — — — V. N.S.W. — — B.f.I, 75 M.fr. XI, 27, 59.
 ERYSIMUM, Linné, gen. pl. 198 (1737). (Blennodia).
 E. curvipes, F. v. Mueller in Linnaea XXV, 368 (1852) ... — — — — — S.A. — — — — — V. N.S.W. — — B.f.I, 75 M.fr. VII, 20.
 E. brevipes, F. v. Mueller in Linnaea XXV, 367 (1852) ... — — — — — W.A. S.A. — — — — — V. N.S.W. — — B.f.I, 75 M.fr. VII, 20; XI, 6.
 E. blennodioides, F. v. Mueller in Linnaea XXV, 367 (1852) ... — — — — — S.A. — — — — — V. N.S.W. Q. — B.f.I, 76 M.fr. VII, 20; XI, 27, 60.
 E. Cunninghamii, Benth. Fl. Austr. I, 76 (1863) ... — — — — — — — N.S.W. Q. — B.f.I, 76
 E. Blennodia, F. v. Mueller, fragm. X, 78 (1877) ... — — — — — S.A. — — — — — N.S.W. — — B.f.I, 76 M.fr. VII, 20; X, 78.
 E. capsellinum, F. v. Mueller, native pl. of Vict. I, 35 (1879) ... — — — — — — — V. N.S.W. — — — — — B.f.I, 77
 STENOPETALUM, R. Brown in De Candolle, Mém. du Mus. VII, 239 (1821).
 S. velutinum, F. v. Mueller, pl. Vict. I, 49 (1860) ... — — — — — S.A. — — — — — V. N.S.W. Q. — B.f.I, 78 M.fr. VII, 20; XI, 27.
 S. lineare, R. Brown in De Candolle, syst. veg. I, 513 (1821) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.f.I, 78 M.fr. VII, 20; XI, 6.
 S. sphaerocarpum, F. v. Mueller in Trans. Ph. Soc. Vict. I, 35 (1854) ... W.A. S.A. — — — — — V. N.S.W. — — B.f.I, 78 M.fr. VII, 20.
 S. nutans, F. v. Mueller, fragm. III, 27 (1862) ... — — — — — — — S.A. — — — — — N.S.W. Q. — B.f.I, 79 M.fr. III, 27; VII, 20; XI, 6.
 S. robustum, Endlicher in Huegel enum. 4 (1837) ... — — — — — W.A. S.A. — — — — — — — B.f.I, 79 M.fr. VII, 20; XI, 5, 60.
 S. pedicellare, F. v. Mueller in Benth. Fl. Austr. I, 79 (1863) ... — — — — — W.A. — — — — — — — B.f.I, 79 M.fr. VII, 20.
 S. croceum, Bunge in Lehmann, pl. Preiss. I, 258 (1844) ... — — — — — W.A. — — — — — — — — — M.fr. XI, 6.
 GEOCOCCUS, Drummond and Harvey in Hook. Kew Misc. VII, 52 (1855).
 G. pusillus, Drumm. & Harv. in Hook. Kew Misc. VII, 52 (1855) ... W.A. S.A. — — — — — V. N.S.W. — — B.f.I, 80 M.fr. VII, 19; XI, 6, 26.
 MENKEA, Lehmann, Ind. sem. hort. bot. Hamb. 8 (1843).
 M. australis, Lehmann in Ind. sem. hort. bot. Hamb. 8 (1843) ... — — — — — W.A. S.A. — — — — — V. N.S.W. — — B.f.I, 80 M.fr. II, 142.
 M. draboides, J. Hooker in Benth. Fl. Austr. I, 80 (1863) ... — — — — — W.A. — — — — — — — B.f.I, 80 M.fr. II, 142; XII, 19.
 M. sphaerocarpa, F. v. Mueller, fragm. VIII, 223 (1874) ... — — — — — S.A. — — — — — — — — — M.fr. VIII, 223.

CAPSELLA, Medicus, Pflanzen-Gattungen 85 (1792). (Hutchinsia partly, Thlaspi partly, Microlepidium).									
C. elliptica, C. A. Meyer, Verz. der Pfl. v. Caucas. 194 (1831) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 81	M.fr. VII, 19; XI, 60.
C. antipoda, F. v. Mueller, pl. of Vict. I, 44 (1860) ...	—	—	T.	V.	—	—	—	B.f.I, 81	
C. pilosula, F. v. Mueller, pl. of Vict. I, 44 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 82	M.fr. XI, 60.
C. Tasmanica, F. v. Mueller, fragm. XI, 26 (1878) ...	—	—	T.	—	—	—	—	B.f.I, 87	
C. cochlearina, F. v. Mueller, pl. of Vict. I, 51 (1860) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.I, 88	M.fr. XI, 26.
C. Drummondii, F. v. Mueller, fragm. XI, 26 (1878) ...	W.A.	S.A.	—	—	—	—	—	B.f.I, 88	M.fr. VII, 90; XI, 26.
C. humistrata, F. v. Mueller, fragm. XI, 25 (1878) ...	—	S.A.	—	—	N.S.W.	—	—	—	M.fr. XI, 25.

SENEBIERA, De Candolle in Mém. soc. hist. nat. Par. I, 140 (1799).									
S. integrifolia, De Candolle in Mém. soc. his. nat. Par. I, 144 (1799) ...	—	—	—	—	—	—	Q.	—	B.f.I, 82
LEPIDIDIUM, Tournefort, inst. 215, t. 103 (1700) from Diosc. and Plinius. (Iberis partly, Lepia, Monoploca).									
L. stronglylophyllum, F. v. Mueller in Benth. Fl. Austr. I, 84 (1863) ...	—	S.A.	—	—	—	—	Q.	N.A.	B.f.I, 84 M.fr. XI, 6.
L. linifolium, Benth. Fl. Austr. I, 84 (1863) ...	W.A.	—	—	—	—	—	—	—	B.f.I, 84 M.fr. VII, 20; XI, 6.
L. leptopetalum, F. v. Mueller, pl. of Vict. I, 48 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	—	B.f.I, 84
L. rotundum, De Candolle, syst. veg. II, 537 (1818) ...	W.A.	—	—	—	—	—	—	—	B.f.I, 85 M.fr. VII, 20; XI, 6.
L. phlebotopetalum, F. v. Mueller, pl. of Vict. I, 47 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	—	B.f.I, 85 M.fr. VII, 19; XI, 60.
L. pedicellum, F. v. Mueller, fragm. XI, 27 (1878) ...	—	—	—	—	—	—	—	N.A.	— M.fr. XI, 27.
L. monoplocoides, F. v. Mueller in Trans. Phil. Soc. Vict. 35 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	—	B.f.I, 85 M.fr. XI, 27.
L. papillosum, F. v. Mueller, in Linnaea XXV, 370 (1852) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	—	B.f.I, 86 M.fr. VII, 19; XI, 6, 28.
L. foliosum, Desvaux, Journ. de Bot. III, 164 et 180 (1813) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	—	B.f.I, 86 M.fr. VII, 19.
L. ruderales, Linné, sp. pl. 645 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.I, 86	M.fr. VII, 19; XI, 27, 61.
CAKILE, Tournefort, coroll. 49, t. 483 (1700).									
C. maritima, Scopoli, Fl. Carniolica, 344 (1760) ...	—	S.A.	T.	V.	N.S.W.	—	—	—	M.fr. VII, 19; XI, 6, 28, 60.

VIOLACEAE.

De Candolle, Fl. Franc. IV, 801 (1805).

VIOLA, Tournefort, inst. 419, t. 236 (1700) from Plinius.									
V. betonicifolia, Smith in Rees's Cyclopaedia XXXVII (1817) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 99	M.fr. X, 82.
V. hederacea, Labillardière, Nov. Holl. pl. spec. I, 66, t. 91 (1804) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 99	M.fr. X, 82.
V. Cunninghamii, J. Hooker, Fl. N. Zel. I, 16 (1853) ...	—	—	T.	—	—	—	—	B.f.I, 100	
V. Caleyana, G. Don, gen. syst. dichlam. pl. I, 329 (1831) ...	—	—	T.	V.	N.S.W.	—	—	B.f.I, 100	M.fr. X, 82.
HYBANTHUS, Jacquin, Stirp. Amer. hist. 77, t. 75 (1763). (Ionidium, Pigea, Vlamingia).									
H. floribundus, F. v. Mueller, nat. pl. of Vict. I, 45 (1879) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.I, 112	M.fr. X, 81.
H. Vernonii, F. v. Mueller, nat. pl. of Vict. I, 45 (1879) ...	—	—	—	V.	N.S.W.	—	—	B.f.I, 103	M.fr. X, 81.
H. filiformis, F. v. Mueller, native pl. of Vict. I, 44 (1879) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.I, 103	M.fr. X, 81.
H. Tatei, F. v. Mueller in Trans. R. Socy. of S. Aust. IV, 102 (1882) ...	—	S.A.	—	—	—	—	—	—	
H. debilissimus, F. v. Mueller, fragm. XI, 4 (1878) ...	W.A.	—	—	—	—	—	—	—	M.fr. XI, 4.
H. calycinus, F. v. Mueller, fragm. X, 81 (1877) ...	W.A.	—	—	—	—	—	—	B.f.I, 104	M.fr. XI, 4.
H. encaspermus, F. v. Mueller, fragm. X, 81 (1877) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 101	M.fr. X, 81.
HYMENANTHERA, R. Brown in Tuck. Cong. 442 (1818).									
H. Banksii, F. v. Mueller, first. gen. report 9 (1853) ...	—	—	T.	V.	N.S.W.	—	—	B.f.I, 104	M.fr. X, 82.

FLACOURTIEAE.

Cl. Richard in Mém. du Mus. I, 366 (1815).

COCHLOSPERMUM, Kunth, Malvac. 6 (1822).									
C. Fraseri, Planchon in Hook. Lond. Journ. VI, 307 (1847) ...	—	—	—	—	—	—	—	N.A.	B.f.I, 106
C. heteronemum, F. v. Mueller in Hook. Kew. Misc. IX, 15 (1857) ...	—	—	—	—	—	—	—	N.A.	B.f.I, 106 M.fr. IX, 61.
C. Gillivraei, Benth. Fl. Austr. I, 106 (1863) ...	—	—	—	—	—	—	—	Q.	B.f.I, 106
C. Gregorii, F. v. Mueller, fragm. I, 71 (1858) ...	—	—	—	—	—	—	—	Q.	N.A. B.f.I, 106 M.fr. IX, 61.
SCOLOPIA, Schreber, Gen. 335 (1789). (Phoberos).									
S. Brownii, F. v. Mueller, fragm. III, 11 (1862) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 107	M.fr. III, 11; IX, 61.
XYLOSMA, G. Forster, prodr. 380 (1786).									
X. ovata, Benth. Fl. Austr. I, 108 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 108	M.fr. IX, 60.
STREPTOTHAMNUS, F. v. Mueller, fragm. III, 28 (1862).									
S. Moorei, F. v. Mueller, fragm. III, 28 (1862) ...	—	—	—	—	N.S.W.	—	—	B.f.I, 108	M.fr. III, 28.
S. Beckleri, F. v. Mueller, fragm. III, 28 (1862) ...	—	—	—	—	N.S.W.	—	—	B.f.I, 108	M.fr. IX, 61.

SAMYDACEAE.

J. Gaertner, de Fruct. III, 328 (1805).

CASEARIA, Jacquin, Stirp. Amer. hist. 132, t. 85 (1763).									
C. esculenta, Roxburgh, Fl. Ind. II, 422 (1832) ...	—	—	—	—	—	—	Q.	—	B.f. VI, 309
C. Dallachii, F. v. Mueller, fragm. V, 107 (1866) ...	—	—	—	—	—	—	Q.	—	B.f. III, 309 M.fr. VI, 252.
HOMALIUM, Jacquin, Stirp. Amer. hist. 170, t. 183 (1763). (Blackwellia).									
H. Vitiense, Benth. in Journ. Linn. Soc. IV, 36 (1860) ...	—	—	—	—	—	—	Q.	—	B.f. III, 310
H. brachybotrys, F. v. Mueller, fragm. II, 127 (1861) ...	—	—	—	—	—	—	Q.	—	B.f. III, 310 M.fr. II, 127.

PITTOSPOREAE.

R. Brown in Flinders' voy. 542 (1814).

PITTOSPORUM, Banks in Gaertner, de Fruct. I, 286, t. 59 (1788).									
P. rhombifolium, Cunningham in Hook. icon. pl. t. 621 (1844) ...	—	—	—	—	N.S.W.	Q.	—	B.f. I, 110	M.fr. XII, 4.
P. melanosperrum, F. v. Mueller, fragm. I, 70 (1858) ...	—	—	—	—	—	—	—	N.A.	B.f. I, 111 M.fr. VI, 167; XII, 4.
P. undulatum, Andrews, Bot. reposit. 383 (1800) ...	—	—	—	—	T.	V.	N.S.W.	Q.	B.f. I, 111 M.fr. VI, 167; XII, 4.
P. revolutum, Aiton, Hort. Kew. sec. ed. II, 27 (1811) ...	—	—	—	—	V.	N.S.W.	Q.	—	B.f. I, 111 M.fr. VI, 167 & 186; VII, 140.
P. venulosum, F. v. Mueller, fragm. VI, 186 (1868) ...	—	—	—	—	—	—	—	Q.	— M.fr. VII, 140; XII, 4.
P. ferrugineum, Aiton, Hort. Kew. sec. ed. II, 27 (1811) ...	—	—	—	—	—	—	Q.	—	B.f. I, 112 M.fr. II, 78; III, 165; VI, 167

<i>P. rubiginosum</i> , Cunningham in Ann. of Nat. Hist. IV, 108 (1842)	—	—	—	—	Q.	—	B.f.I, 112	M. fr. VI, 167; VII, 140; IX, 190.
<i>P. phillyroides</i> , De Candolle, prodr. I, 347 (1824)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.I, 112
<i>P. bicolor</i> , Hooker, Journ. Bot. I, 249 (1834)	—	—	—	T.	V.	N.S.W.	—	B.f.I, 113
<i>P. erioluma</i> , C. Moore et F. v. Mueller, fragm. VII, 139 (1871)	—	—	—	—	N.S.W.	—	—	M. fr. VII, 139.
HYMENOSPORUM, R. Brown in F. v. Mueller, fragm. II, 77 (1860).								
<i>H. flavum</i> , F. v. Mueller, fragm. II, 77 (1860)	—	—	—	—	N.S.W.	Q.	—	B.f.I, 114
BURSARIA, Cavanilles, icon. et descr. pl. IV, 30, t. 350 (1797).								
<i>B. spinosa</i> , Cavanilles, Icon. pl. IV, 30, t. 350 (1797)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.I, 115
MARIANTHUS, Huegel, enum. pl. Nov. Holl. 8 (1837). (Oncosporum, Rhytidosporum, Calopetalon).								
<i>M. procumbens</i> , Benth. Fl. Austr. I, 117 (1863)	—	—	—	T.	V.	N.S.W.	Q.	—
<i>M. microphyllus</i> , Benth. Fl. Austr. I, 117 (1863)	W.A.	—	—	—	—	—	—	B.f.I, 117
<i>M. villosus</i> , Benth. Fl. Austr. I, 117 (1863)	W.A.	—	—	—	—	—	—	B.f.I, 117
<i>M. parviflorus</i> , F. v. Mueller, fragm. II, 144 (1861)	W.A.	—	—	—	—	—	—	B.f.I, 118
<i>M. bignoniaceus</i> , F. v. Mueller in Trans. Phil. Soc. Vict. 6 (1854)	—	S.A.	—	V.	—	—	—	B.f.I, 118
<i>M. Drummondianus</i> , Benth. Fl. Austr. I, 119 (1863)	W.A.	—	—	—	—	—	—	B.f.I, 119
<i>M. coeruleo-punctatus</i> , Klotzsch in Link and Otto, icon. pl. 28, t. 12 (1841)	W.A.	—	—	—	—	—	—	B.f.I, 119
<i>M. candidus</i> , Huegel, enum. pl. Nov. Holl. 8 (1837)	W.A.	—	—	—	—	—	—	B.f.I, 120
<i>M. erubescens</i> , Putterlick in Endl. nov. stirp. dec. 60 (1839)	W.A.	—	—	—	—	—	—	B.f.I, 120
<i>M. ringens</i> , F. v. Mueller, fragm. I, 218 (1859)	W.A.	—	—	—	—	—	—	B.f.I, 120
<i>M. pictus</i> , Lindley in Bot. Regist. XXV, App. XXV (1839)	W.A.	—	—	—	—	—	—	B.f.I, 121
CITRIOBATUS, Cunningham, in Loudon's hort. Brit. supp. I, 585 (1832). (Ixiosporum).								
<i>C. multiflorus</i> , Cunningham in Loud. hort. Brit. I, 585 (1832)	—	—	—	—	N.S.W.	Q.	—	B.f.I, 121
<i>C. pauciflorus</i> , Cunningham in Loud. hort. Brit. I, 585 (1832)	—	—	—	—	—	Q.	N.A.	B.f.I, 122
BILLARDIERA, Smith, Sp. Bot. New Holl. I, t. 1 (1793). (Pronaya, Campylanthera).								
<i>B. longiflora</i> , Labillardiere, Nov. Holl. plant. spec. I, 64, t. 89 (1804)	—	—	—	T.	V.	N.S.W.	—	B.f.I, 123
<i>B. scandens</i> , Smith, Spec. bot. New Holl. I, t. 1 (1793)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 123
<i>B. coriacea</i> , Benth. Fl. Austr. I, 124 (1863)	W.A.	—	—	—	—	—	—	B.f.I, 124
<i>B. cymosa</i> , F. v. Mueller in Transact. Vict. Inst. I, 29 (1855)	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 124
<i>B. elegans</i> , F. v. Mueller, pl. of Vict. I, 78 (1860)	W.A.	—	—	—	—	—	—	B.f.I, 125
<i>B. variifolia</i> , De Candolle, prodr. I, 346 (1824)	W.A.	—	—	—	—	—	—	B.f.I, 125
<i>B. Lehmanniana</i> , F. v. Mueller, pl. of Vict. I, 78 (1860)	W.A.	—	—	—	—	—	—	B.f.I, 125
<i>B. floribunda</i> , F. v. Mueller in South Sc. Record II, 1 (1882)	W.A.	—	—	—	—	—	—	B.f.I, 120
SOLLYA, Lindley, Bot. Regist. XXII, t. 1460 (1831). (Xerosollya.)								
<i>S. heterophylla</i> , Lindley, Bot. Regist. XVII, t. 1466 (1831)	W.A.	—	—	—	—	—	—	B.f.I, 126
<i>S. parviflora</i> , Turcz. in Bull. Soc. Mosc. XXVII, 361 (1854)	W.A.	—	—	—	—	—	—	B.f.I, 126
CHEIRANTHERA, Cunningham, in Brongn. Bot. voy. Coquille, t. 77 (1829).								
<i>C. linearis</i> , Cunningham in Bot. Regist. 1719 (1834)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.I, 127
<i>C. filifolia</i> , Turczaninow in Bull. Soc. Mosc. XXVII, 364 (1854)	W.A.	—	—	—	—	—	—	B.f.I, 127
<i>C. parviflora</i> , Benth. Fl. Austr. I, 128 (1863)	W.A.	—	—	—	—	—	—	B.f.I, 128

DROSERACEAE.

Salisbury, Parad. Lond. 95 (1809).

ALDROVANDA, Monti in Act. Bonon. II, 404, t. 12 (1747).								
<i>A. vesiculosa</i> , Linné, sp. pl. 281 (1753)	—	—	—	—	—	Q.	—	M. fr. VI, 104; X, 79.
DROSERA, Linné, gen. pl. 89 (1737). (Londera).								
<i>D. Indica</i> , Linné, sp. pl. 282 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.II, 456
<i>D. Adelaë</i> , F. v. Mueller, fragm. IV, 154 (1864)	—	—	—	—	—	Q.	—	M. fr. X, 80.
<i>D. Arcturi</i> , Hooker, Journ. of Bot. I, 247 (1834)	—	—	—	T.	V.	N.S.W.	—	B.f.II, 456
<i>D. glanduligera</i> , Lehmann, pugill. VIII, 37 (1844)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.II, 457
<i>D. pygmaea</i> , De Candolle, prodr. I, 317 (1824)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 457
<i>D. platystigma</i> , Lehmann, pugill. VIII, 37 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 457
<i>D. pulchella</i> , Lehmann, pugill. VIII, 38 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 458
<i>D. leucoblasta</i> , Benth. Fl. Austr. II, 458 (1864)	W.A.	—	—	—	—	—	—	B.f.II, 458
<i>D. nitidula</i> , Planchon in Ann. des sc. nat. IX, 285 (1848)	W.A.	—	—	—	—	—	—	B.f.II, 458
<i>D. paleacea</i> , R. Brown in De Candolle, prodr. I, 318 (1824)	W.A.	—	—	—	—	—	—	B.f.II, 458
<i>D. parvula</i> , Planchon in Ann. des sc. nat. IX, 287 (1848)	W.A.	—	—	—	—	—	—	B.f.II, 459
<i>D. Burmanni</i> , Vahl, Symbol. III, 50 (1794)	—	S.A.	—	—	—	Q.	N.A.	B.f.II, 459
<i>D. spatulata</i> , Labillardiere, Nov. Holl. pl. spec. I, 79, t. 106 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 459
<i>D. Drummondii</i> , Lehmann, pl. Preiss. II, 235 (1847)	W.A.	—	—	—	—	—	—	B.f.II, 460
<i>D. scorpioides</i> , Planchon in Ann. des sc. nat. IX, 288 (1848)	W.A.	—	—	—	—	—	—	B.f.II, 460
<i>D. petiolaris</i> , R. Brown in De Candolle, prodr. I, 318 (1824)	—	—	—	—	—	Q.	N.A.	B.f.II, 460
<i>D. binata</i> , Labillardiere, Nov. Holl. pl. spec. I, 78, t. 105 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 461
<i>D. zonaria</i> , Planchon in Ann. des sc. nat. IX, 303 (1848)	W.A.	—	—	—	—	—	—	B.f.II, 462
<i>D. bulbosa</i> , Hooker, icon. pl. t. 375 (1841)	W.A.	—	—	—	—	—	—	B.f.II, 462
<i>D. rosulata</i> , Lehmann, pugill. VIII, 36 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 462
<i>D. Whittakerii</i> , Planchon in Ann. des sc. nat. IX, 302 (1848)	—	S.A.	—	V.	—	—	—	B.f.II, 462
<i>D. macrophylla</i> , Lindley, Bot. Regist. XXV, App. XX, (1839)	W.A.	—	—	—	—	—	—	B.f.II, 463
<i>D. squamosa</i> , Benth. Fl. Austr. II, 463 (1864)	W.A.	—	—	—	—	—	—	B.f.II, 463
<i>D. erythrorrhiza</i> , Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.f.II, 463
<i>D. stolonifera</i> , Endlicher in Hueg. enum. 5 (1837)	W.A.	—	—	—	—	—	—	B.f.II, 463
<i>D. humilis</i> , Planchon in Ann. des sc. nat. IX, 300 (1848)	W.A.	—	—	—	—	—	—	B.f.II, 464
<i>D. ramellosa</i> , Lehmann, pugill. VIII, 40 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 464
<i>D. platypoda</i> , Turczaninow in Bull. del'Ac. de St. Petersburg. 343 (1852)	W.A.	—	—	—	—	—	—	B.f.II, 464
<i>D. auriculata</i> , Backhouse in Ann. des sc. nat. IX, 295 (1848)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 465
<i>D. lunata</i> , Buchanan in De Cand. prodr. I, 319 (1824)	—	—	—	—	—	Q.	—	—
<i>D. peltata</i> , Smith in Willd. spec. pl. I, 1546 (1797)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 465

D. Neesii, Lehmann, pugill. VIII, 42 (1844)	W.A.	—	—	—	—	—	B.f.II, 466	
D. gigantea, Lindley, Bot. Regist. XXV, App. XX (1839) ...	W.A.	—	—	—	—	—	B.f.II, 466	
D. myriantha, Planchon in Ann. des sc. nat. IX, 291 (1848) ...	W.A.	—	—	—	—	—	B.f.II, 466	
D. pallida, Lindley, Bot. Regist. XXV, App. XX (1839) ...	W.A.	—	—	—	—	—	B.f.II, 467	
D. penicillaris, Benth. Fl. Austr. II, 467 (1864)	W.A.	—	—	—	—	—	B.f.II, 467	
D. filicaulis, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	B.f.II, 467	
D. Huegelii, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	B.f.II, 467	M.fr. VIII, 185; X, 81.
D. macrantha, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	B.f.II, 468	
D. Menziesii, R. Brown in De Candolle, prodr. I, 319 (1824) ...	W.A. S.A. T. V. N.S.W.	—	—	—	—	—	B.f.II, 468	
D. calycina, Planchon in Ann. des sc. nat. IX, 299 (1848) ...	W.A.	—	—	—	—	—	B.f.II, 468	
D. heterophylla, Lindley, Bot. Regist. XXV, App. XX (1839) ...	W.A.	—	—	—	—	—	B.f.II, 469	
D. Banksii, R. Brown in De Candolle, prodr. I, 319 (1824) ...	—	—	—	—	—	Q.	B.f.II, 469	
BYBLIS, Salisbury, Parad. Lond. t. 95 (1808).								
B. liniflora, Salisbury, Parad. Lond. t. 95 (1808)	—	—	—	—	—	Q. N.A.	B.f.II, 470	M.fr. X, 81.
B. gigantea, Lindley Bot. Regist. XXV, App. XXI (1839) ...	W.A.	—	—	—	—	—	B.f.II, 470	

ELATINEAE.

Cambessèdes in Mém. du Mus. XVIII, 225 (1829).

ELATINE, Linné, gen. pl. 118 (1737).								
E. Americana, Arnott in Edinb. journ. nat. sc. I, 431 (1830) ...	W.A. S.A. T. V. N.S.W. Q.	—	—	—	—	—	B.f.I, 178	M.fr. XI, 27.
BERGIA, Linné, mantiss. II, 152 (1771).								
B. ammannioides, Roth, nov. pl. spec. 219 (1821)	—	—	—	—	V. N.S.W. Q. N.A.	—	B.f.I, 180	M.fr. II, 147.
B. pedicellaris, F. v. Mueller in Benth. Fl. Austr. I, 180 (1863) ...	—	—	—	—	—	N.A.	B.f.I, 180	M.fr. II, 145.
B. perennis, F. v. Mueller in Benth. Fl. Austr. I, 180 (1863) ...	—	—	—	—	—	N.A.	B.f.I, 181	M.fr. II, 146.

HYPERICINAE.

J. de St. Hilaire, Expos. fam. II, 23, t. 75 (1805).

HYPERICUM, Tournefort, inst. 254, t. 131 (1700), from Dioscorides and Plinius.								
H. Japonicum, Thunberg, Fl. Jap. 295, t. 31 (1784)	W.A. S.A. T. V. N.S.W. Q. N.A.	—	—	—	—	—	B.f.I, 182	

GUTTIFERAE.

A. L. de Jussieu, Gen. 243 (1789).

CALOPHYLLUM, Linné, gen. 154 (1737).								
C. inophyllum, Linné, sp. pl. 513 (1753)	—	—	—	—	—	Q.	—	B.f.I, 183 M.fr. IX, 175.
C. tomentosum, Wight, Illustrat. Pl. Ind. Orient. I, 128 (1840) ...	—	—	—	—	—	Q.	—	M.fr. IX, 174.

POLYGALEAE.

A. L. de Jussieu, in Ann. du Mus. XIV, 386 (1809).

SALOMONIA, Loureiro, Fl. Cochinch. I, 14 (1790).								
S. oblongifolia, De Candolle, prodr. I, 334 (1824)	—	—	—	—	N.S.W. Q.	—	B.f.I, 138	M.fr. XI, 4.
POLYGALA, Tournefort, inst. 174, t. 79 (1700).								
P. Sibirica, Linné, sp. pl. 702 (1753)	—	—	—	—	V. N.S.W. Q.	—	B.f.I, 139	M.fr. XI, 4.
P. leptalea, De Candolle, prodr. I, 325 (1824)	—	—	—	—	—	Q. N.A.	B.f.I, 139	M.fr. XI, 4.
P. eriocephala, F. v. Mueller in Benth. Fl. Austr. I, 139 (1863) ...	—	—	—	—	—	N.A.	B.f.I, 139	
P. persicarifolia, De Candolle, prodr. I, 326 (1818)	—	—	—	—	—	—	N.A.	M.fr. XI, 4.
P. Chinensis, Linné, sp. pl. 704 (1753)	—	—	—	—	—	Q. N.A.	B.f.I, 140	M.fr. XI, 4.
P. stenoclada, Benth. Fl. Austr. I, 141 (1863)	—	—	—	—	—	N.A.	B.f.I, 141	
COMESPERMA, Labillardière, Nov. Holl. pl. spec. II, 21 (1806).								
C. sphaerocarpaceum, Steetz in Lehm. pl. Preiss. II, 314 (1847) ...	—	—	—	—	N.S.W. Q.	—	B.f.I, 143	M.fr. XI, 3.
C. scoparium, Steetz in Lehm. pl. Preiss. II, 309 (1847) ...	W.A. S.A.	—	—	V. N.S.W.	—	—	B.f.I, 143	
C. aphyllum, R. Brown in Benth. Fl. Austr. I, 143 (1863) ...	—	—	—	—	—	N.A.	B.f.I, 143	
C. spinosum, F. v. Mueller, fragm. I, 144 (1859)	W.A.	—	—	—	—	—	B.f.I, 144	M.fr. I, 144.
C. volubile, Labillardière, Nov. Holl. pl. spec. II, 24, t. 163 (1806) ...	W.A. S.A. T. V. N.S.W. Q.	—	—	—	—	—	B.f.I, 144	M.fr. XI, 3.
C. ciliatum, Steetz in Lehm. pl. Preiss. II, 304 (1847) ...	W.A.	—	—	—	—	—	B.f.I, 144	M.fr. X, 3.
C. integerimum, Endlicher in Hueg. enum. (1837)	W.A.	—	—	—	—	—	B.f.I, 145	M.fr. XI, 3.
C. secundum, Banks in De Candolle, prodr. I, 334 (1824) ...	—	—	—	—	—	Q. N.A.	B.f.I, 145	
C. Drummondii, Steetz in Lehm. pl. Preiss. II, 301 (1847) ...	W.A.	—	—	—	—	—	B.f.I, 145	M.fr. XI, 2.
C. praecelsum, F. v. Mueller, fragm. XI, 2 (1878)	—	—	—	—	—	Q.	—	M.fr. XI, 2.
C. retusum, Labillardière, Nov. Holl. pl. spec. II, 22, t. 160 (1806) ...	—	—	—	—	T. V. N.S.W. Q.	—	B.f.I, 145	M.fr. XI, 2.
C. sylvestre, Lindley in Mitch. Trop. Austr. 342 (1848) ...	—	—	—	—	—	Q.	B.f.I, 146	M.fr. I, 49; XI, 2.
C. acerosum, Steetz in Lehm. pl. Preiss. II, 299 (1847) ...	W.A.	—	—	—	—	—	B.f.I, 146	M.fr. XI, 3.
C. ericinum, De Candolle, prodr. I, 334 (1824)	—	—	—	—	T. V. N.S.W. Q.	—	B.f.I, 146	M.fr. I, 48; XI, 2.
C. viscidulum, F. v. Mueller, fragm. X, 4 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 4; XI, 2.
C. confertum, Labillardière, Nov. Holl. pl. spec. II, 23, t. 161 (1806) ...	W.A.	—	—	—	—	—	B.f.I, 147	M.fr. XI, 3.
C. flavum, De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	B.f.I, 147	M.fr. XI, 3.
C. calymega, Labillardière, Nov. Holl. pl. spec. II, 23, t. 162 (1806) ...	W.A. S.A. T. V. N.S.W.	—	—	—	—	—	B.f.I, 147	M.fr. XI, 3.
C. rhadinocarpaceum, F. v. Mueller, fragm. XI, 1 (1878) ...	W.A.	—	—	—	—	—	—	M.fr. XI, 1.
C. lanceolatum, R. Brown in Benth. Fl. Austr. I, 148 (1863) ...	W.A.	—	—	—	—	—	B.f.I, 148	M.fr. XI, 1.
C. defoliatum, F. v. Mueller, pl. of Vict. I, 189 (1862) ...	—	—	—	—	T. V. N.S.W. Q.	—	B.f.I, 148	M.fr. XI, 3.
C. nudiusculum, De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	B.f.I, 148	M.fr. XI, 4.
C. virgatum, Labillardière, Nov. Holl. pl. spec. II, 21 t. 159 (1806) ...	W.A.	—	—	—	—	—	B.f.I, 149	M.fr. XI, 2.
C. polygaloides, F. v. Mueller in Trans. Phil. Soc. Vict. I, 7 (1854) ...	W.A. S.A.	—	—	V.	—	—	B.f.I, 149	M.fr. XI, 4.
XANTHOPHYLLUM, Roxburgh, Pl. Corom. III, 81, t. 284 (1819).								
X. Macintyrii, F. v. Mueller, fragm. V, 57 (1865)	—	—	—	—	—	Q.	—	M.fr. V, 57; XI, 4.

TREMANDREAE.

R. Brown in Flind. voy. II, 554 App. (1814).

PLATYTHECA, Steetz in Lehmann, pl. Preiss. I, 220 (1845).											
P. galioides, Steetz in Lehm. pl. Preiss. I, 220 (1845)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 136	M.fr. XII.
TETRATHECA, Smith, Spec. Bot. New Holl. I, t. 2 (1793).											
T. Harperi, F. v. Mueller, fragm. V, 49 (1865)	...	...	W.A.	—	—	—	—	—	—	—	M.fr. V, 180.
T. ciliata, Lindley in Mitch. three Exped. II, 206 (1838)	...	...	—	S.A.	T. V.	—	—	—	—	B.f.I, 130	M.fr. XII.
T. efoliata, F. v. Mueller, fragm. X, 3 (1876)	...	...	W.A.	—	—	—	—	—	—	—	M.fr. X, 3.
T. aphylla, F. v. M. in South. Science Record, vol. II, May (1852)	...	...	W.A.	—	—	—	—	—	—	—	M.fr. XII.
T. ericifolia, Smith, Exot. Bot. I, 37, t. 20 (1804)	...	...	—	S.A.	T. V.	N.S.W.	Q.	—	—	B.f.I, 131	M.fr. XII.
T. juncea, Smith, specim. of Bot. of New Holl. 5, t. 2 (1793)	...	...	—	—	—	—	N.S.W.	—	—	B.f.I, 132	M.fr. XII.
T. affinis, Endlicher in Hueg. enum. 7 (1837)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 132	M.fr. XII.
T. nuda, Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 133	M.fr. XII.
T. virgata, Steetz in Lehm. pl. Preiss. I, 212 (1845)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 133	M.fr. XII.
T. confertifolia, Steetz in Lehm. pl. Preiss. I, 214 (1845)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 133	M.fr. XII.
T. setigera, Endlicher in Hueg. enum. 8 (1837)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 133	M.fr. XII.
T. hirsuta, Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 134	M.fr. XII.
T. pilifera, Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 135	M.fr. XII.
T. filiformis, Benth. Fl. Austr. I, 135 (1863)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 135	M.fr. XII.
TREMANDRA, R. Brown in App. Flind. voy. II, 544 (1814).											
T. stelligera, R. Brown in De Candolle, prodr. I, 344 (1824)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 136	M.fr. XII.
T. diffusa, R. Brown in De Candolle, prodr. I, 344 (1824)	...	...	W.A.	—	—	—	—	—	—	B.f.I, 137	M.fr. XII.

MELIACEAE.

Ventenat, Tabl. III, 159 (1799).

HEDRAIANTHERA, F. v. Mueller, fragm. V, 58 (1865).											
H. porphyropetala, F. v. Mueller, fragm. V, 58 (1865)	...	...	—	—	—	—	—	Q.	—	—	M.fr. V, 58.
TURRAEA, Linné, mantiss. II, 150 (1771).											
T. pubescens, Hellenius in K. Sv. Vent. Acad. Handl. 26, t. 10 (1788)	—	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 379	M.fr. V, 144.	
MELIA, Linné, gen. pl. 127 (1737).											
M. Azedarach, Linné, sp. pl. 384 (1753)	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 380	M.fr. V, 144; IX, 134; XI, 135.	
DYSOXYLUM, Blume, Bijdrag. 172 (1825). (Dysoxylon, Epicharis, Hartigsheia).											
D. oppositifolium, F. v. Mueller, fragm. V, 144 (1866)	...	...	—	—	—	—	Q.	—	—	M.fr. V, 144; IX, 61.	
D. latifolium, Benth. Fl. Austr. I, 381 (1863)	...	...	—	—	—	—	Q.	—	B.f.I, 381	M.fr. IX, 61.	
D. Klanderi, F. v. Mueller, fragm. V, 176 (1866)	...	...	—	—	—	—	Q.	—	—	M.fr. V, 176; IX, 61, 134.	
D. Fraserianum, Benth. Fl. Austr. I, 381 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 381	M.fr. IX, 61.	
D. arborescens, Miquel in Annal. Mus. Bot. Lugd. IV, 24 (1868)	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 134.	
D. Muelleri, Benth. Fl. Austr. I, 381 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 381	M.fr. V, 145.	
D. Schiffneri, F. v. Mueller in Melb. Chemist, 53 (1881)	...	...	—	—	—	—	Q.	—	—	M.fr. XI, 138.	
D. Becklerianum, C. de Candolle in Monogr. phan. I, 509 (1878)	...	...	—	—	—	N.S.W.	Q.	—	—	M.fr. X, 138.	
D. Lessertianum, Benth. Fl. Austr. I, 382 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 382	M.fr. V, 145; IX, 61.	
D. Schultzii, C. de Candolle in Monogr. phanerogr. I, 509 (1878)	...	...	—	—	—	—	—	N.A.	—	M.fr. XI, 138.	
D. rufum, Benth. Fl. Austr. I, 382 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 382	M.fr. V, 145.	
D. Nernstii, F. v. Mueller, fragm. V, 176 (1866)	...	...	—	—	—	—	Q.	—	—	M.fr. V, 176.	
AMOORA, Roxburgh, pl. Corom. III, 54, t. 258 (1819).											
A. nitidula, Benth. Fl. Austr. I, 383 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 383	—	
SYNOUM, Adr. de Jussieu in Mém. du Mus. XIX, 226, t. 15 (1830).											
A. glandulosum, de Jussieu in Mém. du Mus. XIX, 227, t. 15 (1830)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 384	M.fr. V, 145; IX, 134.	
A. Muelleri, C. de Candolle in Monogr. phanerogr. I, 593 (1878)	...	...	—	—	—	—	Q.	—	—	M.fr. V, 145; XI, 138.	
OWENIA, F. v. Mueller in Hook. Kew Misc. IX, 303 (1857).											
O. acidula, F. v. Mueller in Hook. Kew Misc. IX, 304 (1857)	...	...	—	S.A.	—	N.S.W.	Q.	—	B.f.I, 385	M.fr. III, 14; IX, 134.	
O. vernicosa, F. v. Mueller, fragm. III, 15 (1862)	...	...	—	—	—	—	—	N.A.	B.f.I, 385	M.fr. III, 15; IX, 134.	
O. venosa, F. v. Mueller in Hook. Kew Misc. IX, 304 (1857)	...	...	—	—	—	—	Q.	—	B.f.I, 386	M.fr. III, 14; IX, 134.	
O. cepiodora, F. v. Mueller, fragm. XI, 82 (1880)	...	...	—	—	—	N.S.W.	—	—	—	M.fr. XI, 82.	
O. reticulata, F. v. Mueller in Hook. Kew Misc. IX, 305 (1857)	...	...	W.A.	—	—	—	—	N.A.	B.f.I, 386	M.fr. III, 13; IX, 134.	
AGLAI A, Loureiro, Fl. Cochinch. I, 173 (1790), non Allemand (1770). (Milnea, Nemedra).											
A. elaeagnoides, Benth. Fl. Austr. I, 383 (1863)	...	...	—	—	—	—	—	N.A.	B.f.I, 383	M.fr. V, 145; IX, 134.	
HEARNIA, F. v. Mueller, fragm. V, 55 (1865). (Aglaiopsis).											
H. sapindina, F. v. Mueller, fragm. V, 55 (1865)	...	...	—	—	—	—	Q.	—	—	M.fr. V, 55.	
CARAPA, Aublet, Hist. pl. Guian. II, Suppl. 32, t. 357 (1775). (Xylocarpus).											
C. Moluccensis, Lamarck, Encycl. meth. Bot. I, 621 (1783)	...	...	—	—	—	—	Q.	N.A.	B.f.I, 387	M.fr. V, 145; IX, 134.	
CEDRELA, P. Browne, Nat. hist. of Jamaica 158 (1756).											
C. australis, F. v. Mueller, fragm. I, 4 (1858)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 387	M.fr. IX, 133; XI, 82.	
FLINDERSIA, R. Brown in Flind. voy. II, 595 (1814). (Oxleya, Strzeleckia).											
F. australis, R. Brown in Flind. voy. II, 595, t. 1 (1814)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 388	M.fr. I, 65; II, 174; IX, 133.	
F. Bourjotiana, F. v. Mueller, fragm. IX, 133 (1875)	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 133.	
F. Schottiana, F. v. Mueller, fragm. III, 25 (1862)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 388	M.fr. III, 25; VI, 143; IX, 133.	
F. Oxleyana, F. v. Mueller, fragm. I, 65 (1858)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 389	M.fr. I, 65; III, 25; IX, 133.	
F. Pimenteliana, F. v. Mueller, fragm. IX, 132 (1875)	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 132.	
F. Bennettiana, F. v. Mueller in Benth. Fl. Austr. I, 389 (1863)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 389	M.fr. IX, 131; XI, 135.	
F. Inflaiana, F. v. Mueller, fragm. X, 94 (1877)	...	...	—	—	—	—	Q.	—	—	M.fr. X, 94.	
F. Brayleyana, F. v. Mueller, fragm. V, 143 (1866)	...	...	—	—	—	—	Q.	—	—	M.fr. I, 143; VI, 252.	
F. Strzeleckiana, F. v. Mueller, fragm. I, 65 (1858)	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 389	M.fr. IX, 133.	

OCHNACEAE.

De Candolle in Ann. du Mus. XVII, 398 (1811).

BRACKENRIDGEA, Asa Gray, Bot. Wilk. Expl. Exped. 361, t. 42 (1854).

B. Australiana, F. v. Mueller, fragm. V, 29 (1865) ... — — — — — Q. — — M.fr. V, 29.

RUTACEAE.

A. L. de Jussieu, Gen. 296 (1789).

ZIERIA, Smith in Transact. Linn. Soc. IV, 216 (1798).

Z. laevigata, Smith in Rees's Cyclopaedia, XXXIX (1819) ... — — — — V. N.S.W. Q. — B.f.I, 304 M.fr. IX, 116.
 Z. pilosa, Rudge in Transact. Linn. Soc. X, 293, t. 17 (1809) ... — — — — N.S.W. — — B.f.I, 305 M.fr. I, 100; IX, 116.
 Z. obcordata, Cunningham in Field. N.S. Wales, 330 (1825) ... — — — — N.S.W. Q. — B.f.I, 305 M.fr. I, 101; IX, 116.
 Z. cytisoides, Smith in Rees's Cyclopaedia, XXXIX (1819) ... — — — — T. V. N.S.W. Q. — B.f.I, 306 M.fr. IX, 116.
 Z. Smithii, Andrews' Bot. Rep. t. 606 (1810) ... — — — — T. V. N.S.W. Q. — B.f.I, 306 M.fr. I, 100; IX, 116.
 Z. granulata, C. Moore in Benth. Fl. Austr. I, 307 (1863) ... — — — — N.S.W. Q. — B.f.I, 307 M.fr. IX, 116.
 Z. veronica, F. v. Mueller in Trans. phil. Soc. Vict. I, 11 (1854) — — S.A. T. V. — — B.f.I, 305 M.fr. IX, 116.

BORONIA, Smith, Tracts relat. nat. hist. 285 (1798). (Cyanothamnus).

B. grandisepala, F. v. Mueller, fragm. I, 66 (1858) ... — — — — — N.A. B.f.I, 311 M.fr. I, 66; II, 179.
 B. artemisifolia, F. v. Mueller, fragm. I, 66 (1858) ... — — — — — N.A. B.f.I, 311 M.fr. I, 66; IX, 111.
 B. affinis, R. Brown in Benth. Fl. Austr. I, 311 (1863) ... — — — — — N.A. B.f.I, 311
 B. filicifolia, Cunningham in Benth. Fl. Austr. I, 311 (1863) ... — — — — — N.A. B.f.I, 311
 B. alata, Smith in Transact. Linn. Soc. VIII, 283 (1807) ... W.A. — — — — — B.f.I, 312 M.fr. IX, 111.
 B. algida, F. v. Mueller in Transact. phil. Soc. Vict. I, 100 (1855) ... — — — — V. N.S.W. — — B.f.I, 312 M.fr. IX, 111.
 B. Edwardsii, Benth. Fl. Austr. I, 312 (1863) ... — — — — S.A. — — — — B.f.I, 312
 B. ternata, Endlicher, nov. stirp. dec. 6 (1839) ... — — — — W.A. — — — — B.f.I, 312 M.fr. IX, 111.
 B. ericifolia, Benth. Fl. Austr. I, 313 (1863) ... — — — — W.A. — — — — B.f.I, 313 M.fr. IX, 111.
 B. inconspicua, Benth. Fl. Austr. I, 313 (1863) ... — — — — W.A. — — — — B.f.I, 313
 B. eriantha, Lindley in Mitch. Trop. Austr. 298 (1848) ... — — — — — Q. — — B.f.I, 313
 B. alulata, Solander in Benth. Fl. Austr. I, 313 (1863) ... — — — — — Q. — — B.f.I, 313
 B. ledifolia, J. Gay, monogr. des Lasiopetalées 20 (1821) ... — — — — N.S.W. Q. — B.f.I, 314 M.fr. IX, 111.
 B. lanceolata, F. v. Mueller, fragm. I, 66 (1858) ... — — — — — Q. N.A. B.f.I, 314 M.fr. I, 66; VII, 38; IX, 112; XI, 97.
 B. Bowmanii, F. v. Mueller, fragm. IV, 135 (1864) ... — — — — — Q. — — M.fr. VII, 38.
 B. Fraseri, Hooker, Bot. Mag. t. 4052 (1844) ... — — — — N.S.W. — — B.f.I, 315 M.fr. IX, 111.
 B. mollis, Cunningham in Lindl. Bot. Reg. 47 (1841) ... — — — — N.S.W. — — B.f.I, 315 M.fr. IX, 112.
 B. megastigma, Nees in Lehm. pl. Preiss. II, 227 (1847) ... — — — — W.A. — — — — B.f.I, 315 M.fr. II, 97; IV, 112; IX, 112.
 B. heterophylla, F. v. Mueller, fragm. II, 98 (1861) ... — — — — W.A. — — — — B.f.I, 315 M.fr. II, 98; IX, 112.
 B. elatior, Bartling in Lehm. pl. Preiss. I, 170 (1844) ... — — — — W.A. — — — — B.f.I, 316 M.fr. II, 98; IX, 122; XI, 97.
 B. tetrandra, Labillardière, Nov. Holl. pl. spec. I, 98, t. 125 (1804) ... — — — — W.A. — — — — B.f.I, 316
 B. crassifolia, Bartling in Lehm. pl. Preiss. I, 169 (1844) ... — — — — W.A. — — — — B.f.I, 316 M.fr. IX, 112.
 B. albiflora, R. Brown in Benth. Fl. Austr. I, 317 (1863) ... — — — — W.A. — — — — B.f.I, 317 M.fr. IX, 114.
 B. lanuginosa, Endlicher in Hueg. enum. 16 (1837) ... — — — — W.A. — — — — B.f.I, 317 M.fr. IX, 114.
 B. pulchella, Turczaninow in Bull. Soc. Mosc. XXV, 162 (1852) ... — — — — W.A. — — — — B.f.I, 318 M.fr. IX, 114.
 B. gracilipes, F. v. Mueller, fragm. II, 99 (1861) ... — — — — W.A. — — — — B.f.I, 318 M.fr. II, 99; IX, 114.
 B. microphylla, Sieber in Spreng. syst. cur. post. 148 (1827) ... — — — — N.S.W. Q. — B.f.I, 318
 B. pinnata, Smith, Tracts relat. nat. hist. 290, t. 4 (1798) ... — — — — S.A. T. V. N.S.W. Q. — B.f.I, 318 M.fr. IX, 114.
 B. pilosa, Labillardière, Nov. Holl. pl. spec. I, 97, t. 124 (1804) ... — — — — T. V. — — B.f.I, 319
 B. spinescens, Benth. Fl. Austr. I, 319 (1863) ... — — — — W.A. — — — — B.f.I, 319
 B. baeckeaceae, F. v. Mueller, fragm. IX, 112 (1875) ... — — — — W.A. — — — — — M.fr. IX, 112.
 B. coerulescens, F. v. Mueller in Trans. phil. Soc. Vict. II (1854) ... — — — — W.A. S.A. — — V. N.S.W. — — B.f.I, 320 M.fr. IX, 112.
 B. defoliata, F. v. Mueller, fragm. IX, 113 (1875) ... — — — — W.A. — — — — — M.fr. IX, 113.
 B. busselliana, F. v. Mueller, fragm. IX, 113 (1875) ... — — — — W.A. — — — — — M.fr. IX, 113.
 B. subcoerulea, F. v. Mueller, fragm. II, 100 (1860) ... — — — — W.A. — — — — — B.f.I, 320 M.fr. II, 100; IX, 112 & 198.
 B. polygalifolia, Smith, Tracts relat. nat. hist. 297, t. 7 (1798) ... — — — — S.A. T. V. N.S.W. Q. — B.f.I, 320 M.fr. IX, 114.
 B. falcifolia, Cunningham in Lindl. Bot. Reg. 47 (1841) ... — — — — N.S.W. Q. — B.f.I, 322 M.fr. IX, 114.
 B. penicillata, Benth. Fl. Austr. I, 322 (1863) ... — — — — W.A. — — — — B.f.I, 322
 B. crassipes, Bartling in Lehm. pl. Preiss. I, 168 (1844) ... — — — — W.A. — — — — B.f.I, 322 M.fr. IX, 116.
 B. subsessilis, Benth. Fl. Austr. I, 322 (1863) ... — — — — W.A. — — — — B.f.I, 322
 B. capitata, Benth. Fl. Austr. I, 323 (1863) ... — — — — W.A. — — — — B.f.I, 323 M.fr. IX, 116.
 B. nematophylla, F. v. Mueller, fragm. II, 100 (1861) ... — — — — W.A. — — — — B.f.I, 323 M.fr. II, 100.
 B. crenulata, Smith in Transact. Linn. Soc. VIII, 284 (1807) ... — — — — W.A. — — — — B.f.I, 323 M.fr. XI, 115 & 198.
 B. serrulata, Smith, Tracts relat. nat. hist. 292 (1798) ... — — — — N.S.W. — — B.f.I, 323 M.fr. IX, 115.
 B. rhomboidea, Hooker icon. pl. t. 722 (1848) ... — — — — T. — — — — B.f.I, 324
 B. parviflora, Smith, Tracts relat. nat. hist. 295, t. 5 (1798) ... — — — — S.A. T. V. N.S.W. Q. — B.f.I, 324 M.fr. IX, 115.
 B. vinea, Lindley, Bot. Regist. XXV, App. XVII (1839) ... — — — — W.A. — — — — B.f.I, 324
 B. filifolia, F. v. Mueller, fragm. I, 3 (1858) ... — — — — S.A. — — V. — — B.f.I, 325 M.fr. I, 31; IX, 114.
 B. clavellifolia, F. v. Mueller in Trans. phil. Soc. Vict. I, 12 (1854) ... — — — — W.A. S.A. — — V. N.S.W. — — B.f.I, 325 M.fr. I, 99; IX, 114 & 198.
 B. scabra, Lindley, Bot. Regist. XXV, App. XVII (1839) ... — — — — W.A. — — — — B.f.I, 326
 B. thymifolia, Turcz. in Bull. Soc. Mosc. XXV, part II, 165 (1852) ... — — — — W.A. — — — — B.f.I, 326 M.fr. I, 99; II, 99; IX, 116.
 B. ovata, Lindley, Bot. Regist. 47 (1841) ... — — — — W.A. — — — — B.f.I, 326 M.fr. IX, 116.
 B. Barkeriana, F. v. Mueller, fragm. XI, 96 (1880) ... — — — — N.S.W. — — — — M.fr. XI, 96.
 B. denticulata, Smith in Transact. Linn. Soc. VIII, 284 (1807) ... — — — — W.A. — — — — B.f.I, 327 M.fr. IX, 115 & 198.
 B. machardiana, F. v. Mueller, fragm. IX, 115 (1875) ... — — — — W.A. — — — — — M.fr. IX, 115.
 B. spathulata, Lindley, Bot. Regist. XXV, App. XVII (1839) ... — — — — W.A. — — — — B.f.I, 327 M.fr. IX, 115.
 B. juncea, Bartling in Lehm. pl. Preiss. I, 166 (1844) ... — — — — W.A. — — — — B.f.I, 327 M.fr. IX, 115.
 B. cymosa, Endlicher in Hueg. enum. 16 (1837) ... — — — — W.A. — — — — B.f.I, 328 M.fr. II, 101; IX, 116.

ERIOSTEMON, Smith in Transact. Linn. Soc. IX, 221 (1798). (Crowea, Phebalium, Asterolasia, Microcybe, Pleurandropsia, Geleznowia, Urocarpus, Sandfordia, Actinostigma).

E. virgatus, Cunningham in Hooker's Journ. of Bot. II, 417 (1840) ... — — — — T. — — — — B.f.I, 332 M.fr. IX, 111.
 E. nodiflorus, Lindley, Bot. Regist. XXV, App. XVII (1839) ... — — — — W.A. — — — — B.f.I, 336 M.fr. IX, 110.

<i>E. spicatus</i> , A. Richard, sert. Astrolab. 76, t. 27 (1833) ...	W.A.	—	—	—	—	—	B.f.I, 336	M.fr.IX, 110.
<i>E. pungens</i> , Lindley in Mitch. Three Exped. II, 156 (1838) ...	—	S.A.	—	V.	N.S.W.	—	B.f.I, 338	M.fr.IX, 110.
<i>E. montanus</i> , F. v. M., pl. of Vict. I, 129 (1860) ...	—	—	T.	—	—	—	B.f.I, 338	M.fr.IX, 109.
<i>E. phyllicifolius</i> , F. v. M., fragm. I, 105 (1859) ...	—	—	—	V.	—	—	B.f.I, 339	M.fr.I, 105.
<i>E. umbellatus</i> , Turcz. in Bull. Soc. Mosc. XXII, part III, 15 (1849) ...	—	—	—	V.	N.S.W.	—	B.f.I, 339	M.fr.I, 104; IX, 109.
<i>E. Ralstoni</i> , F. v. M., fragm. II, 101, t. 14 (1861) ...	—	—	—	V.	N.S.W.	—	B.f.I, 339	M.fr.III, 165; IX, 109.
<i>E. Hillebrandii</i> , F. v. M. in Transact. phil. Soc. Vict. I, 10 (1854) ...	—	S.A.	T.	V.	—	—	B.f.I, 340	M.fr.I, 4; IX, 109.
<i>E. lamprophyllus</i> , F. v. M. in Journ. pharm. Soc. Vict. II, 43 (1858) ...	—	—	—	V.	—	—	B.f.I, 340	M.fr.IX, 109.
<i>E. elatior</i> , F. v. M., fragm. I, 181 (1859) ...	—	—	—	—	N.S.W.	—	B.f.I, 340	M.fr.IX, 109.
<i>E. ambiens</i> , F. v. M., fragm. VI, 166 (1868) ...	—	—	—	—	N.S.W.	—	—	M.fr.VI, 166; IX, 110.
<i>E. Oldfieldii</i> , F. v. M., fragm. I, 3 (1858) ...	—	—	—	T.	—	—	B.f.I, 340	M.fr.IX, 109.
<i>E. rotundifolius</i> , Cunningham in Hueg. enum. 15 (1837) ...	—	—	—	—	N.S.W.	Q.	B.f.I, 341	—
<i>E. microphyllus</i> , F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854) ...	—	S.A.	—	—	—	—	B.f.I, 341	M.fr.IX, 109.
<i>E. phyllicoides</i> , F. v. M., fragm. I, 107 (1859) ...	—	—	—	V.	N.S.W.	—	B.f.I, 341	M.fr.IX, 107.
<i>E. ozothamnoides</i> , F. v. M., fragm. I, 103 (1859) ...	—	—	—	V.	N.S.W.	—	B.f.I, 342	M.fr.IX, 107.
<i>E. Mortonii</i> , F. v. M., fragm. IX, 108 (1875) ...	—	—	—	—	N.S.W.	—	B.f.I, 342	M.fr.IX, 108.
<i>E. sediflorus</i> , F. v. M., fragm. I, 102 (1859) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.I, 342	M.fr.I, 102.
<i>E. Nottii</i> , F. v. M., fragm. VI, 22 (1867) ...	—	—	—	—	—	Q.	—	M.fr.VI, 22.
<i>E. lepidotus</i> , Sprengel, syst. II, 322 (1825) ...	—	S.A.	—	V.	N.S.W.	Q.	B.f.I, 343	M.fr.I, 104; IX, 107.
<i>E. alpinus</i> , F. v. M., fragm. I, 103 (1859) ...	—	—	—	V.	N.S.W.	—	—	M.fr.I, 103.
<i>E. tuberculosus</i> , F. v. M., pl. of Vict. I, 130 (1862) ...	W.A.	—	—	—	—	—	B.f.I, 343	M.fr.IX, 108.
<i>E. Benthami</i> , F. v. M., fragm. IX, 108 (1875) ...	W.A.	—	—	—	—	—	B.f.I, 343	M.fr.IX, 108.
<i>E. Maxwellii</i> , F. v. M., fragm. IX, 108 (1875) ...	W.A.	—	—	—	—	—	—	M.fr.IX, 103.
<i>E. filifolius</i> , F. v. M., fragm. IX, 108 (1875) ...	W.A.	—	—	—	—	—	B.f.I, 344	M.fr.IX, 108.
<i>E. squameus</i> , Labillardière, Nov. Holl. pl. spec. I, 111, t. 141 (1804) ...	—	—	T.	V.	N.S.W.	Q.	B.f.I, 344	M.fr.I, 104; IX, 107.
<i>E. anceps</i> , Sprengel, syst. II, 322 (1825) ...	W.A.	—	—	—	—	—	B.f.I, 345	M.fr.I, 103; IX, 108.
<i>E. ovatifolius</i> , F. v. M., fragm. I, 103 (1859) ...	—	—	—	V.	N.S.W.	—	B.f.I, 345	M.fr.I, 103.
<i>E. bilobus</i> , F. v. M., fragm. I, 102 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 345	M.fr.IX, 108.
<i>E. Beckleri</i> , F. v. M., fragm. IX, 109 (1875) ...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 108.
<i>E. capitatus</i> , F. v. M., fragm. I, 116 (1859) ...	W.A.	S.A.	—	—	—	—	B.f.I, 346	M.fr.I, 106; IX, 107.
<i>E. Geleznovii</i> , F. v. M., fragm. I, 107 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 347	M.fr.IX, 107.
<i>E. correfolius</i> , F. v. M., fragm. I, 105 (1859) ...	—	—	—	V.	N.S.W.	—	B.f.I, 350	M.fr.IX, 107.
<i>E. Cunninghamii</i> , F. v. M., fragm. IX, 107 (1875) ...	—	—	—	—	N.S.W.	—	B.f.I, 351	M.fr.IX, 107.
<i>E. mollis</i> , Asterolasia mollis, Benth. Fl. Austr. I, 351 (1863) ...	—	—	—	—	N.S.W.	—	B.f.I, 349	—
<i>E. pleurandroides</i> , F. v. M., fragm. I, 106 (1859) ...	—	—	—	V.	—	—	B.f.I, 349	M.fr.IX, 107.
<i>E. trymaloides</i> , F. v. M., fragm. I, 106 (1859) ...	—	—	—	V.	N.S.W.	—	B.f.I, 351	M.fr.I, 106.
<i>E. Hookeri</i> , F. v. M., fragm. I, 104 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 352	M.fr.VII, 22; IX, 107.
<i>E. pallidus</i> , Asterolasia pallida, Benth. Fl. Austr. I, 352 (1863) ...	W.A.	—	—	—	—	—	B.f.I, 352	—
<i>E. Drummondii</i> , F. v. M., fragm. I, 105 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 352	M.fr.VII, 22.
<i>E. grandiflorus</i> , F. v. M., fragm. I, 105 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 353	M.fr.VI, 21.
<i>E. Crowei</i> , F. v. Mueller, pl. of Vict. I, 119 (1860) ...	—	—	—	V.	N.S.W.	Q.	B.f.I, 329	M.fr.IX, 106.
<i>E. Turczaninowii</i> , F. v. Mueller, pl. of Vict. I, 120 (1860) ...	W.A.	—	—	—	—	—	B.f.I, 330	M.fr.IX, 106.
<i>E. lanceolatus</i> , K. F. Gaertner, de fruct. III, 154, t. 210 (1807) ...	—	—	—	V.	N.S.W.	—	B.f.I, 331	M.fr.IX, 110.
<i>E. Banksii</i> , Cunningham in Hueg. enum. 15 (1837) ...	—	—	—	—	—	Q.	B.f.I, 332	—
<i>E. trachyphyllus</i> , F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854) ...	—	—	—	V.	N.S.W.	—	B.f.I, 332	M.fr.IX, 110.
<i>E. myoporoides</i> , De Candolle, prodr. I, 720 (1824) ...	—	—	—	V.	N.S.W.	Q.	B.f.I, 333	M.fr.IX, 110.
<i>E. hispidulus</i> , Sieber in Spreng. syst. cur. post. 164 (1827) ...	—	—	—	—	N.S.W.	—	B.f.I, 333	M.fr.IX, 110.
<i>E. buxifolius</i> , Smith in Rees's Cyclop. XIII, 2 (1809) ...	—	—	—	—	N.S.W.	—	B.f.I, 333	M.fr.IX, 111.
<i>E. obovatis</i> , Cunningham in Field's N. S. Wales, 331 (1825) ...	—	S.A.	T.	V.	N.S.W.	—	B.f.I, 334	M.fr.IX, 110.
<i>E. scaber</i> , Paxton, Mag. of Bot. XIII, 127 (1846) ...	—	—	—	—	N.S.W.	Q.	B.f.I, 334	M.fr.IX, 111.
<i>E. Brucei</i> , F. v. M., fragm. VII, 38 (1869) ...	W.A.	—	—	—	—	—	—	M.fr.IX, 110.
<i>E. linearis</i> , Cunningham in Hueg. enum. 16 (1837) ...	—	S.A.	—	—	N.S.W.	—	B.f.I, 334	—
<i>E. difformis</i> , Cunningham in Hueg. enum. 15 (1837) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	B.f.I, 335	M.fr.IX, 110.
<i>E. parvifolius</i> , R. Brown in Benth. Fl. Austr. I, 335 (1863) ...	—	—	—	—	—	Q.	B.f.I, 335	—
<i>E. ericifolius</i> , Cunningham in Benth. Fl. Austr. I, 335 (1863) ...	—	—	—	—	N.S.W.	—	B.f.I, 335	—
PHILOTHECA, Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815). (Drummondita).								
<i>P. australis</i> , Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815) ...	—	—	—	—	N.S.W.	Q.	B.f.I, 348	M.fr.VII, 20, 141.
<i>P. ericoides</i> , F. v. M., fragm. I, 107 (1859) ...	W.A.	—	—	—	—	—	B.f.I, 349	M.fr.VII, 21.
<i>P. calida</i> , F. v. M., fragm. VII, 21 (1869) ...	—	—	—	—	—	Q.	—	M.fr.VII, 38.
CORREA, Smith in Transact. Linn. Soc. IV, 219 (1798). (Didymeria).								
<i>C. aemula</i> , F. v. M., fragm. I, 3 (1859) ...	—	S.A.	—	V.	—	—	B.f.I, 353	M.fr.IX, 117.
<i>C. alba</i> , Andrews, Bot. Reposit. t. 18 (1797) ...	—	S.A.	T.	V.	N.S.W.	—	B.f.I, 354	M.fr.IX, 117.
<i>C. speciosa</i> , Andrews, Bot. Reposit. t. 653 (1811) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.I, 354	M.fr.IX, 117.
<i>C. Lawrenciana</i> , Hooker, Journ. of Bot. I, 254 (1834) ...	—	—	T.	V.	N.S.W.	—	B.f.I, 355	M.fr.VIII, 142; IX, 117.
<i>C. decumbens</i> , F. v. M. in Transact. phil. Soc. Vict. I, 30 (1854) ...	—	S.A.	—	—	—	—	B.f.I, 356	—
NEMATOLEPIS, Turczaninow in Bull. Soc. Mosc. XXV, part III, 158 (1852). (Symphyopetalum).								
<i>N. phlebaloides</i> , Turczaninow in Bull. Soc. Mosc. part III, 158 (1852) ...	W.A.	—	—	—	—	—	B.f.I, 356	M.fr.IX, 106.
<i>N. Euphemiae</i> , F. v. M., fragm. III, 149 (1863) ...	W.A.	—	—	—	—	—	B.f.I, 345	M.fr.IX, 106.
CHORILAENA, Endlicher in Hueg. enum. 17 (1837).								
<i>C. quercifolia</i> , Endlicher in Hueg. enum. 17 (1837) ...	W.A.	—	—	—	—	—	B.f.I, 357	M.fr.IX, 106.
<i>C. hirsuta</i> , Benth. Fl. Austr. I, 357 (1863) ...	W.A.	—	—	—	—	—	B.f.I, 357	M.fr.IX, 106.
DIPLOLAENA, R. Brown in Append. Flind. voy. II, 546 (1814).								
<i>D. Dampieri</i> , Desfontaines in Mém. du Mus. 452, t. 20 (1817) ...	W.A.	—	—	—	—	—	B.f.I, 358	M.fr.IX, 106.
BOSISTOA, F. v. M. in Benth. Fl. Austr. I, 359 (1863).								
<i>B. sapindiformis</i> , F. v. M. in Benth. Fl. Austr. I, 359 (1863) ...	—	—	—	—	N.S.W.	Q.	B.f.I, 359	M.fr.IX, 103.
<i>B. euodiformis</i> , F. v. M., fragm. IX, 174 (1875) ...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 174.

ACRADENIA, Kippist in Transact. Linn. Soc. XXI, 207 (1855).									
A. Frankliniae, Kippist in Transact. Linn. Soc. XXI, 207 (1855)	—	T.	—	—	—	—	B.fl.I, 323	M.fr.VI, 167; IX, 103.	
EUODIA, R. et G. Forster, char. gen. 13, t. 7 (1776). (Melicope, Medicosma).									
E. pentacocca, F. v. M., fragm. III, 41 (1862)	...	...	...	...	N.S.W.	Q.	—	M.fr.III, 168; IV, 175.	
E. Cunninghami, F. v. M., fragm. III, 2 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 362	M.fr.I, 27; IX, 102.	
E. Fareana, F. v. M., fragm. IX, 101 (1875)	...	...	...	...	—	Q.	—	M.fr.IX, 101.	
E. erythrocoeca, F. v. M., fragm. I, 26 (1858)	...	...	...	...	N.S.W.	Q.	B.fl.I, 360	M.fr.II, 103, 178; IX, 102.	
E. contermina, Moore & F. v. M., fragm. VII, 144 (1871)	...	...	...	...	N.S.W.	—	—	M.fr.VII, 144.	
E. octandra, F. v. M., fragm. II, 102 (1861)	...	...	...	...	N.S.W.	—	B.fl.I, 360	M.fr.II, 102; IX, 102.	
E. microcoeca, F. v. M., fragm. I, 144 (1859)	...	...	...	...	N.S.W.	Q.	B.fl.I, 361	M.fr.II, 103, 180; IX, 102.	
E. xanthoxyloides, F. v. M., fragm. IV, 155 (1864)	...	...	...	...	—	Q.	—	M.fr.IV, 155.	
E. Bonwickii, F. v. M., fragm. V, 56 (1865)	...	...	...	...	—	Q.	—	M.fr.V, 179.	
E. alata, F. v. M., fragm. VII, 142 (1871)	...	...	...	...	—	Q.	—	M.fr.VII, 142.	
E. acedens, Blume, Bijdr. 246 (1825)	...	...	...	...	N.S.W.	Q.	—	M.fr.V, 456; VII, 22; IX, 102.	
E. littoralis, Endlicher, prodr. Fl. Norfolk. 86 (1833)	...	...	...	...	N.S.W.	—	—	M.fr.IX, 102.	
E. polybotrya, Moore & F. v. M., fragm. VII, 143 (1871)	...	...	...	...	N.S.W.	—	—	M.fr.IX, 193.	
E. vitiflora, F. v. M., fragm. VII, 144 (1871)	...	...	...	...	—	Q.	—	M.fr.VII, 144.	
BROMBYA, F. v. M., fragm. V, 4 (1865).									
B. platynema, F. v. M., fragm. V, 4 (1865)	...	...	...	...	—	Q.	—	M.fr.V, 56; IX, 102.	
PAGETIA, F. v. M., fragm. V, 178 (1866).									
P. medicinalis, F. v. M., fragm. V, 178 (1866)	...	...	...	...	—	Q.	—	M.fr.V, 178; IX, 103.	
BOUCHARDATIA, Baillon in Adans. VII, 350, t. 10 (1867).									
B. neurocoeca, Baillon in Adans. IX, 110 (1869)	...	...	...	...	N.S.W.	Q.	B.fl.I, 360	M.fr.I, 28; II, 163, 178; IX, 103.	
XANTHOXYLUM, Catesby in Linné, hort. Cliff. 487 (1737). (Zanthoxylum, Blackburnia).									
X. brachyacanthum, F. v. M., pl. of Vict. I, 108 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 363	M.fr.VII, 141; IX, 104.	
X. Blackburnia, Benth. Fl. Austr. I, 363 (1863)	...	...	...	...	N.S.W.	—	B.fl.I, 363	—	
X. parviflorum, Benth. Fl. Austr. I, 363 (1863)	...	...	...	...	—	N.A.	B.fl.I, 363	M.fr.VII, 141; IX, 104.	
X. torum, F. v. M., fragm. VII, 140 (1871)	...	...	...	...	—	Q.	—	M.fr.VII, 140; IX, 104.	
GELJERA, Schott, Rutac. 4, t. 7 (1834). (Coatesia).									
G. Muelleri, Benth. Fl. Austr. I, 364 (1863)	...	...	...	...	—	Q.	B.fl.I, 364	M.fr.I, 26; IX, 104.	
G. salicifolia, Schott, Rutac. t. 4 (1834)	...	...	...	...	N.S.W.	Q.	B.fl.I, 364	M.fr.VII, 22; IX, 105; XI, 134.	
G. parviflora, Lindley in Mitch. Trop. Austr. 102 (1848)	...	W.A.	S.A.	—	V.	N.S.W.	Q.	B.fl.I, 364	M.fr.VII, 22; XI, 105; XI, 139.
PLEIOCOCCA, F. v. Mueller, fragm. IX, 117 (1875).									
P. Wilcoxiana, F. v. M., fragm. IX, 117 (1875)	...	...	...	...	N.S.W.	—	—	M.fr.IX, 117.	
ACRONYCHIA, R. & G. Forster, char. gen. 53, t. 27 (1776). (Cyminosma).									
A. Baueri, Schott, Rutac. t. 3 (1834)	...	...	...	...	N.S.W.	Q.	B.fl.I, 366	M.fr.I, 26; IX, 103.	
A. laevis, R. & G. Forster, char. gen. 53, t. 27 (1776)	...	...	...	...	V.	N.S.W.	Q.	B.fl.I, 366	M.fr.I, 27; IX, 104.
A. imperfecta, F. v. M., fragm. I, 26 (1858)	...	...	...	...	—	Q.	B.fl.I, 367	M.fr.I, 26; IX, 104.	
A. melicopoides, F. v. M., fragm. V, 3 (1865)	...	...	...	...	N.S.W.	Q.	—	M.fr.IV, 117; VII, 145.	
A. acidula, F. v. M., fragm. IV, 154 (1864)	...	...	...	...	—	Q.	—	M.fr.IV, 154.	
A. vestita, F. v. M., fragm. IV, 155 (1864)	...	...	...	...	—	Q.	—	M.fr.IV, 155; IX, 104.	
A. Endlicheri, Schott, fragm. bot. III, t. 2	...	...	...	...	N.S.W.	—	—	M.fr.IX, 103.	
A. tetrandra, F. v. M., fragm. IX, 104 (1875)	...	...	...	...	—	Q.	—	M.fr.V, 179; IX, 104.	
HALFORDIA, F. v. Mueller, fragm. V, 43 (1865).									
H. drupifera, F. v. M., fragm. V, 43 (1865)	...	...	...	...	N.S.W.	Q.	—	M.fr.V, 5, 189; IX, 103.	
H. scleroxyla, F. v. M., fragm. VII, 142 (1871)	...	...	...	...	—	Q.	—	M.fr.VII, 142; IX, 103.	
GLYCOSMIS, Correa in Ann. du Mus. VI, 384 (1805).									
G. pentaphylla, Correa in Ann. du Mus. VI, 384 (1805)	...	...	...	...	—	Q.	B.fl.I, 367	M.fr.IX, 105.	
MICROMELUM, Blume, Bijdrag. 137 (1825).									
M. pubescens, Blume, Bijdrag. 137 (1825)	...	...	...	...	—	Q.	N.A.	B.fl.I, 368	M.fr.I, 25; IX, 105.
MURRAYA, Koenig in Linné, mant. alt. 563 (1771). (Chalcas, 1767).									
M. exotica, Koenig in Linné, mant. alt. 563 (1771)	...	...	...	...	—	Q.	B.fl.I, 369	M.fr.IX, 106.	
M. crenulata, Oliver in Journ. Linn. Soc. V, suppl. II, 29 (1861)	...	...	...	...	—	Q.	B.fl.I, 369	—	
CLAUSENA, Burman, Fl. Indica index & t. 29 (1768).									
C. brevistyla, Oliver in Journ. Linn. Soc. V, suppl. II, 31 (1861)	...	...	...	...	—	Q.	B.fl.I, 369	M.fr.IX, 106.	
ATALANTIA, Correa in Ann. du Mus. VI, 383 (1805).									
A. glauca, J. Hooker in B. & H. Gen. pl. I, 305 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 370	M.fr.XI, 105.	
A. recurva, Benth. Fl. Austr. I, 370 (1863)	...	...	...	...	—	N.A.	B.fl.I, 370	M.fr.VII, 142.	
CITRUS, Linné, gen. pl. 230 (1737), from Plinius.									
C. Planchonii, F. v. M., Report Intercol. Exhib. 23 (1867)	...	...	...	...	N.S.W.	Q.	B.fl.I, 371	M.fr.IX, 105.	[IX, 105.]
C. Australasica, F. v. M., fragm. I, 26 (1858)	...	...	...	...	N.S.W.	Q.	B.fl.I, 371	M.fr.I, 26; II, 178; III, 167;	
PENTACERAS, J. Hooker in B. & H. Gen. pl. I, 298 (1862).									
P. australis, J. Hooker in B. & H. Gen. pl. I, 298 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 365	M.fr.IX, 104.	

SIMARUBEAE.

De Candolle in Ann. du Mus. XVII, 323 (1811).

AILANTUS, Desfontaines, in Act. Acad. Paris, 265 (1786).									
A. imberbiflora, F. v. M., fragm. III, 42 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 373	M.fr.III, 42; VI, 166.	
BRUCEA, J. S. Miller, icon. plantar. t. 25 (1780).									
B. Sumatrana, Roxburgh, Fl. Indica I, 469 (1820)	...	...	...	...	—	N.A.	B.fl.I, 373	M.fr.IX, 166.	
HYPTIANDRA, J. Hooker in B. & H. Gen. pl. 294 (1862).									
H. Bidwilli, J. Hooker in B. & H. Gen. pl. 294 (1862)	...	...	...	...	N.S.W.	Q.	B.fl.I, 374	M.fr.VI, 165.	

- CADELLIA, F. v. Mueller, fragm. II, 25 (1860).
 C. pentastylis, F. v. M., fragm. II, 26, t. 12 (1860) ... — — — — N.S.W. — — B.fl.I, 374 M.fr.II, 25 & 171.
 C. monostylis, Benth. Fl. Austr. I, 375 (1863) ... — — — — N.S.W. Q. — — B.fl.I, 375 M.fr.IV, 166; VIII, 34.
 SURIANA, Plumier, nov. plant. Amer. gen. 37, t. 40 (1703).
 S. maritima, Linné, sp. pl. 284 (1753) ... — — — — — Q. — — B.fl.I, 375
 HARRISONIA, R. Brown in Mém. du Mus. XII, 517 (1825). (Ebelingia).
 H. Brownii, A. L. de Juss. in Mém. du Mus. Par. XII, 530, t. 28 (1825) — — — — — N.A. B.fl.I, 376

ZYGOPHYLLEAE.

R. Brown in App. Flind. voy. II, 545 (1814).

- NITRARIA, Linné, syst. nat. X, 1044 (1759).
 N. Schoberi, Linné, sp. pl. ed. sec. 638 (1762) ... — — — — W.A. S.A. — — V. N.S.W. — — B.fl.I, 291 M.fr.XI, 30.
 ZYGOPHYLLUM, Linné, syst. nat. 8 (1735), Linné, gen. 126 (1737). (Sarcozygium).
 Z. apiculatum, F. v. M. in Linnaea XXV, 373 (1852) ... — — — — S.A. T. V. N.S.W. Q. — — B.fl.I, 292 M.fr.XI, 29.
 Z. glaucescens, F. v. M. pl. of Vict. I, 228 (1862) ... — — — — S.A. — — V. N.S.W. — — B.fl.I, 293 M.fr.XI, 29.
 Z. iodicarpum, F. v. M. in Linnaea XXV, 372 (1852) ... — — — — W.A. S.A. — — V. N.S.W. — — B.fl.I, 293 M.fr.XI, 29.
 Z. prismatothecum, F. v. M. in Linnaea XXV, 375 (1852) ... — — — — S.A. — — — — — B.fl.I, 293
 Z. ammophilum, F. v. M. in Linnaea XXV, 376 (1852) ... — — — — W.A. S.A. — — V. N.S.W. — — — — M.fr.XI, 28.
 Z. Billardieri, de Candolle, prodr. I, 705 (1824) ... — — — — W.A. S.A. T. V. N.S.W. — — — — B.fl.I, 293 M.fr.XI, 29.
 Z. fruticosum, de Candolle, prodr. I, 705 (1824) ... — — — — W.A. S.A. — — V. N.S.W. Q. — — B.fl.I, 294 M.fr.XI, 29.
 Z. Howittii, F. v. M., fragm. III, 150 (1863) ... — — — — S.A. — — — — — M.fr.III, 150; XI, 29.
 TRIBULUS, L'Obel, Icon. II, 84 (1581), from Theophr. Diosc. & Plinius. (Tribulopsis).
 T. terrestris, L'Obel, stirp. icones II, 84 (1581) ... — — — — W.A. S.A. — — V. N.S.W. Q. N.A. B.fl.I, 288 M.fr.XI, 30.
 T. ranunculiflorus, F. v. M., fragm. I, 48 (1858) ... — — — — — — — — — — N.A. B.fl.I, 288 M.fr.I, 48.
 T. hystrix, R. Brown in App. Sturt. Exped. 6 (1849) ... — — — — S.A. — — — — — N.A. B.fl.I, 289 M.fr.XI, 30.
 T. macrocarpum, F. v. M. in Benth. Fl. Austr. I, 289 (1863) ... — — — — — — — — — — N.A. B.fl.I, 289
 T. platypterus, Benth. Fl. Austr. I, 289 (1863) ... — — — — — — — — — — N.A. B.fl.I, 289
 T. hirsutus, Benth. Fl. Austr. I, 289 (1863) ... — — — — — — — — — — N.A. B.fl.I, 289
 T. minutus, Leichhardt in Benth. Fl. Austr. I, 291 (1863) ... — — — — — — — — — — Q. — — B.fl.I, 291
 T. Brownii, F. v. M., pl. of Vict. I, 99 (1860) ... — — — — — — — — — — Q. N.A. B.fl.I, 290 M.fr.I, 48.
 T. bicolor, F. v. M., pl. of Vict. I, 99 (1860) ... — — — — — — — — — — N.A. B.fl.I, 290 M.fr.I, 47.
 T. Solandri, F. v. M., pl. of Vict. I, 99 (1860) ... — — — — — — — — — — Q. N.A. B.fl.I, 290 M.fr.I, 47; XI, 30.

LINEAE.

De Candolle. Théor. élém. bot. 214 (1813).

- HUGONIA, Linné, gen. pl. 134 (1737).
 H. Jenkensis, F. v. M., fragm. V, 7 (1865) ... — — — — — — — — — — Q. — — — — M.fr.IX, 190.
 ERYTHROXYLUM, P. Brown, Hist. pl. of Jamaica 278 (1756), from Theophr. Diosc. & Plinius. (Erythroxyton).
 E. australe, F. v. M. in Transact. Vict. Inst. III, 22 (1855) ... — — — — — — — — — — Q. — — B.fl.I, 284
 E. ellipticum, R. Brown in Benth. Fl. Austr. I, 284 (1863) ... — — — — — — — — — — N.A. B.fl.I, 284
 LINUM, Tournefort, inst. 339, t. 176, (1700), from Theophr. Diosc. & Plinius.
 L. marginale, Cunningh. in Hook. Lond. Jour. Bot. VII, 169 (1848) W.A. S.A. T. V. N.S.W. Q. — — B.fl.I, 283

GERANIACEAE.

A. L. de Jussieu, Gen. 268 (1780), from B. de Jussieu (1759).

- GERANIUM, Tournefort, inst. 266, t. 142 (1700), from Diosc. & Plinius.
 G. Carolinianum, Linné, sp. pl. 682 (1753) ... — — — — — — — — — — W.A. S.A. T. V. N.S.W. Q. — — B.fl.I, 296
 G. sessiliflorum, Cavanilles, Diss. 198, t. 77 (1790) ... — — — — — — — — — — T. V. N.S.W. — — — — B.fl.I, 297
 ERODIUM, L. Héritier, Geraniologia, t. 1 (1787).
 E. cygnotum, Nees in Lehm. pl. Preiss. I, 162 (1844) ... — — — — — — — — — — W.A. S.A. — — V. N.S.W. Q. N.A. B.fl.I, 297 M.fr.XI, 20.
 PELARGONIUM, L'Héritier, Geraniologia, t. 7 (1787).
 P. australe, Willdenow, sp. pl. III, 675 (1800) ... — — — — — — — — — — W.A. S.A. T. V. N.S.W. Q. — — B.fl.I, 298
 P. Rodneyanum, Mitchell, three Exped. II, 144 (1838) ... — — — — — — — — — — W.A. S.A. — — V. N.S.W. — — — — B.fl.I, 299
 OXALIS, Linné, gen. pl. 134 (1737), from Plinius.
 O. Magellanica, G. Forster in Comm. Goetting. IX, 33 (1789) ... — — — — — — — — — — T. V. — — — — B.fl.I, 300
 O. corniculata, Linné, sp. pl. 435 (1753) ... — — — — — — — — — — W.A. S.A. T. V. N.S.W. Q. — — B.fl.I, 301

MALVACEAE.

Adanson in Mém. Acad. fr. 224 (1761).

- LAVATERA, Tournefort in Act. Acad. Par. 86, t. 3 (1706).
 L. plebeja, Sims in Bot. Mag. t. 2269 (1821) ... — — — — — — — — — — W.A. S.A. T. V. N.S.W. — — — — B.fl.I, 185 M.fr.IX, 130.
 MALVASTRUM, Asa Gray in Mém. Amer. Acad. IV, 21 (1849). (Malva partly).
 M. spicatum, A. Gray in Mém. Amer. Acad. IV, 21 (1849) ... — — — — — — — — — — S.A. — — — — N.S.W. Q. N.A. B.fl.I, 187 M.fr.VI, 170; IX, 130.
 M. tricuspidatum, A. Gray in Mém. Amer. Acad. IV, 21 (1849) ... — — — — — — — — — — — — — — — — N.S.W. Q. — — B.fl.I, 187
 PLAGIANTHUS, R. & G. Forster, char. gen. 85, t. 43 (1776). (Asterotrichon, Blepharanthemum, Lawrencia, Halothamnus).
 P. sidoides, Hooker, Bot. Mag. t. 3396 (1835) ... — — — — — — — — — — — — — — — — T. — — — — B.fl.I, 188 M.fr.VI, 169.
 P. pulchellus, A. Gray, Bot. Amer. Expl. Exped. I, 181 (1854) ... — — — — — — — — — — — — — — — — T. V. N.S.W. — — — — B.fl.I, 189 M.fr.IX, 130.
 P. spicatus, Benth. in proceed. Linn. Soc. VI, 103 (1862) ... — — — — — — — — — — W.A. S.A. T. V. N.S.W. — — — — B.fl.I, 189 M.fr.VI, 170; IX, 130.
 P. glomeratus, Benth. in proceed. Linn. Soc. VI, 103 (1862) ... — — — — — — — — — — W.A. S.A. — — — — N.S.W. Q. — — B.fl.I, 190 M.fr.IX, 130.
 P. diffusus, Benth. Fl. Austr. I, 190 (1863) ... — — — — — — — — — — W.A. — — — — — — — — — — N.A. B.fl.I, 190
 P. microphyllus, F. v. M., fragm. I, 29 (1858) ... — — — — — — — — — — W.A. S.A. — — V. N.S.W. — — — — B.fl.I, 190 M.fr.VII, 170; IX, 13.
 P. Berthae, F. v. M., fragm. V, 103 (1866) ... — — — — — — — — — — W.A. S.A. — — — — — — — — — — — — M.fr.V, 103.

SIDA, Linné, gen. pl. 205 (1737), from Theophrastus.

S. corrugata, Lindley in Mitch. Three Exped. II, 13 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.I, 192	
S. intricata, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1860)...	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.f.I, 193	
S. macropoda, F. v. M. in Benth. Fl. Austr. I, 193 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 193	
S. virgata, Hooker in Mitch. Trop. Austr. 361 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 194	
S. echinocarpa, F. v. M., fragm. XI, 62 (1879) ...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 62.
S. cryphiopetala, F. v. M., fragm. III, 4 (1862) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.f.I, 194	M.fr. III, 4.
S. petrophila, F. v. M. in Linnaea XXV, 381 (1852) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 194	
S. calychymenia, J. Gay in de Candolle, prodr. I, 462 (1824) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f.I, 194	
S. physocalyx, F. v. M., fragm. III, 3 (1862) ...	—	—	—	—	—	—	N.A.	B.f.I, 195	M.fr. III, 3.
S. subspicata, F. v. M. in Benth. Fl. Austr. I, 195 (1863) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 195	
S. pleiantha, F. v. M. in Benth. Fl. Austr. I, 195 (1863) ...	—	—	—	—	—	Q.	—	B.f.I, 195	M.fr. VI, 169.
S. spinosa, Linné, sp. pl. 683 (1753) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 196	
S. rhombifolia, Linné, sp. pl. 684 (1753) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 196	
S. cordifolia, Linné, sp. pl. 684 (1753) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 196	
S. platycalyx, F. v. M. in Benth. Fl. Austr. I, 197 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 197	
S. cleisocalyx, F. v. M., fragm. X, 73 (1876) ...	—	—	—	—	—	—	N.A.	—	M.fr. X, 73.
S. inclusa, Benth. Fl. Austr. I, 197 (1863) ...	—	S.A.	—	—	—	—	N.A.	B.f.I, 197	M.fr. IX, 181; XI, 32.
S. cardiophylla, F. v. M., fragm. VIII, 242 (1874) ...	W.A.	S.A.	—	—	—	—	N.A.	—	M.fr. VIII, 242.
S. Hookeriana, Miquel in Lehm. pl. Preiss. I, 242 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 197	
S. lepidia, F. v. M., fragm. VI, 168 (1868) ...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 32.

ABUTILON, Tournefort, inst. 99, t. 25 (1700), from Camerarius (1586).

A. tubulosum, Hooker in Mitch. Trop. Austr. 390 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.I, 200	M.fr. IX, 131.
A. amplum, Benth. Fl. Austr. I, 200 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 200	
A. leucopetalum, F. v. M. in Benth. Fl. Austr. I, 200 (1863) ...	—	S.A.	—	—	N.S.W.	—	N.A.	B.f.I, 200	M.fr. II, 12.
A. Mitchellii, Benth. Fl. Austr. I, 201 (1863) ...	—	S.A.	—	—	—	Q.	—	B.f.I, 201	
A. micropetalum, F. v. M. in Benth. Fl. Austr. I, 201 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 201	M.fr. VI, 169 S.; IX, 131; XI,
A. cryptopetalum, F. v. M. in Benth. Fl. Austr. I, 201 (1863) ...	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.f.I, 201	M.fr. II, 11; IX, 13. [32.
A. geranioides, Benth. Fl. Austr. I, 202 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 202	
A. otocarpum, F. v. M. in Transact. phil. Soc. Vict. I, 13 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.I, 202	M.fr. XI, 32.
A. longilobum, F. v. M., fragm. IX, 130 (1875) ...	W.A.	—	—	—	—	—	—	—	M.fr. IX, 130.
A. subviscosum, Benth. Fl. Austr. I, 202 (1863) ...	—	—	—	—	—	Q.	—	B.f.I, 202	
A. Indicum, G. Don, gen. syst. I, 504 (1831) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 202	
A. auritum, G. Don, gen. syst. I, 500 (1831) ...	—	—	—	—	—	Q.	—	B.f.I, 203	M.fr. XI, 32.
A. Avicennae, Gaertner, de fruct. II, 251, t. 135 (1791) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 203	
A. graveolens, Wight & Arnott, Prodr. Fl. Pen. Ind. or. I, 56 (1840) ...	—	—	—	—	—	Q.	—	B.f.I, 204	
A. oxycarpum, F. v. M. in Benth. Fl. Austr. I, 204 (1863) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 204	M.fr. II, 12; IX, 131; XI, 32.
A. muticum, G. Don, gen. syst. I, 502 (1831) ...	—	—	—	—	—	Q.	—	B.f.I, 204	
A. Cunninghamii, Benth. Fl. Austr. I, 205 (1863) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 205	
A. exonemum, F. v. M., fragm. XI, 63 (1879) ...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 63.
A. Fraseri, Hooker in Walpers, annal. II, 158 (1851) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 205	
A. halophilum, F. v. M. in Linnaea XXV, 381 (1852) ...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.I, 206	M.fr. XI, 27.
A. macrum, F. v. M., fragm. IX, 59 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 59.
A. crispum, G. Don, gen. syst. I, 502 (1831) ...	—	—	—	—	—	—	N.A.	B.f.I, 206	M.fr. VI, 169.

URENA, Dillenius, Hort. Eltham. 430, t. 319 (1732).

U. lobata, Linné, sp. pl. 692 (1753) ...	—	—	—	—	—	Q.	—	B.f.I, 206	M.fr. VI, 169; IX, 130.
U. Armitiana, F. v. M., fragm. X, 78 (1877) ...	—	—	—	—	—	Q.	—	—	M.fr. X, 78.

PAVONIA, Cavanilles, monadelph. dissert. II, App. II (1786). (Greevesia).

P. hastata, Cavanilles, dissert. III, 138, t. 47 (1786) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 207	M.fr. IX, 130.
---	---	---	---	---	--------	----	---	------------	----------------

HOWITTIA, F. v. Mueller in Transact. Vict. Inst. I, 116 (1855).

H. trilocolata, F. v. M. in Transact. Vict. Inst. I, 116 (1855) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 198	M.fr. VI, 169.
---	---	------	---	----	--------	---	---	------------	----------------

HIBISCUS, Linné, syst. nat. (1735); Linné, gen. pl. 207 (1737). (Fugosia partly, Abelmoschus, Paritium).

H. ficulneus, Linné, sp. pl. 605 (1753) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 209	M.fr. I, 67.
H. Abelmoschus, Linné, sp. pl. 696 (1753) ...	—	—	—	—	—	—	N.A.	—	M.fr. VI, 169.
H. rhodopetalus, F. v. M. in Benth. Fl. Austr. I, 209 (1863) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 209	M.fr. IX, 229; XI, 31.
H. Manihot, Linné, sp. pl. 696 (1753) ...	—	—	—	—	—	Q.	—	B.f.I, 210	
H. Notho-Manihot, F. v. M., fragm. V, 57 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr. IX, 130.
H. Trionum, Linné, sp. pl. 697 (1753) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 210	M.fr. II, 115; IX, 129; XI, 31.
H. brachysiphonius, F. v. M., fragm. I, 67 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.I, 210	M.fr. I, 67; VI, 169.
H. Drummondii, Turczan. in Bull. Soc. Mosc. XXXI, 195 (1858) ...	W.A.	—	—	—	—	—	—	B.f.I, 211	M.fr. I, 220; IX, 129; XI, 31.
H. microlaenus, F. v. M., fragm. II, 116 (1861) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f.I, 211	M.fr. III, 5; IX, 129.
H. Pinonius, Gaudichaud in Freyc. voy. Bot. 476, t. 100 (1826) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f.I, 211	M.fr. II, 116. [IX, 129; XI, 31.
H. cannabinus, Linné, sp. pl. ed. sec., 979 (1763) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 212	M.fr. II, 117; III, 166; VI, 269;
H. divaricatus, Graham in Edinb. phil. Journ. (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 212	M.fr. II, 118; IX, 129; XI, 31.
H. Fitzgeraldii, F. v. M., fragm. VIII, 242 (1874) ...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 242.
H. Elsworthii, F. v. M., fragm. VIII, 241 (1874) ...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 241.
H. heterophyllus, Ventenat, jard. de la Malm., t. 103 (1804) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 212	M.fr. VI, 170; XI, 31.
H. diversifolius, N. Jacquin, icon. pl. rar. 551 (1789) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 213	M.fr. II, 117; VI, 169.
H. splendens, Fraser in Edinb. phil. Journ. (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 213	M.fr. VI, 170; XI, 31 & 123.
H. zonatus, F. v. M., fragm. I, 221 (1859) ...	—	—	—	—	—	—	N.A.	B.f.I, 213	M.fr. VI, 170.
H. Coatesii, F. v. M., fragm. III, 5 (1862) ...	—	—	—	—	—	—	N.A.	B.f.I, 214	M.fr. III, 5.
H. leptocladius, Benth. Fl. Austr. I, 214 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 214	M.fr. VI, 169; IX, 129.
H. setulosus, F. v. M., fragm. I, 221 (1859) ...	—	—	—	—	—	—	N.A.	B.f.I, 214	M.fr. I, 221; XI, 31.
H. Goldsworthii, F. v. M., fragm. XI, 30 (1878) ...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 30.
H. pentaphyllus, F. v. M., fragm. II, 13 (1860) ...	—	—	—	—	—	—	N.A.	B.f.I, 214	M.fr. VI, 169; XI, 32.
H. geranioides, Cunningham in Benth. Fl. Austr. I, 215 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 215	
H. vitifolius, Linné, sp. pl. 696 (1753) ...	—	—	—	—	—	Q.	—	B.f.I, 215	M.fr. II, 114; VI, 169.
H. panduriformis, N. Burman, Fl. Ind. I, 151, t. 47 (1768) ...	—	—	—	—	—	—	N.A.	B.f.I, 215	M.fr. II, 115; VI, 169; IX, 129.
H. Normani, F. v. M., fragm. III, 4 (1862) ...	—	—	—	—	—	Q.	—	B.f.I, 216	M.fr. III, 4; VI, 169.

H. Krichauffii, F. v. M., Rep. Babb. Exped. 7 (1858) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I, 216	M.fr. VI, 169; IX, 129.
H. Farragei, F. v. M., fragm. VIII, 241 (1874) ...	...	...	W.A.	S.A.	—	—	N.S.W.	—	—	—	M.fr. IX, 129 & 196; XI, 32.
H. Sturtii, Hooker in Mitch. trop. Austr. 363 (1848) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I, 216	M.fr. II, 13.
H. phyllochaenus, F. v. M., fragm. IX, 128 (1875) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. IX, 128.
H. Huegelii, Endlicher in Hueg. enum. 10 (1837) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.I, 217	M.fr. IX, 128.
H. Wrayae, Lindley, bot. Regist. t. 69 (1840) ...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr. IX, 128.
H. tiliaceus, Linné, sp. pl. 694 (1753) ...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I, 218	M.fr. VI, 170; IX, 129; XI, 32.
H. cuneiformis, de Candolle, prodr. I, 454 (1824) ...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl.I, 219	M.fr. XI, 32.
H. hakeaeifolius, Giordano, Mém. di nuov. sp. d'Ibisco (1833) ...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.I, 220	M.fr. IX, 128.
LAGUNARIA, G. Don, gen. syst. dichlam. pl. I, 485 (1831).											
L. Patersoni, G. Don, gen. syst. I, 485 (1831) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 218	M.fr. IX, 130; XI, 32.
THESPIA, Solander in Ann. du Mus. IX, 290 (1807).											
T. populnea, Solander in Ann. du Mus. IX, 290, t. 8 (1807) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 221	M.fr. VII, 170; IX, 130; XI, 32.
GOSSYPIUM, Linné, gen. pl. 206 (1737), from Camerarius (1586). (Fugosia partly, Sturtia). [IX, 127; XI, 63.											
G. Sturtii, F. v. M., fragm. III, 6 (1862) ...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.I, 222	M.fr. III, 174; VI, 169 & 251;
G. australe, F. v. M., fragm. I, 46 (1858) ...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.I, 220	M.fr. III, 6; IX, 127, XI, 32 & 63
G. thespesioides, F. v. M., fragm. IX, 127 (1875) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 220	M.fr. V, 177; VI, 169; XI, 32 & 63
G. Cunninghamii, Todaro, Monogr. Gen. Gossyp. 7 (1878) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 220	M.fr. VI, 169; IX, 129; XI, 32 & 63
G. costulatum, Todaro, Monogr. Gen. Gossyp. 7 (1878) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 221	M.fr. IX, 127; XI, 32 & 63. [63.
G. flaviflorum, F. v. M., fragm. IX, 127 (1875) ...	...	...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 63.
G. populifolium, F. v. M., fragm. IX, 127 (1875) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 221	M.fr. XI, 32 & 63.
G. Robinsonii, F. v. M., fragm. IX, 126 (1875) ...	...	...	W.A.	—	—	—	—	—	N.A.	—	M.fr. XI, 63.
ADANSONIA, Linné, sp. pl. I, 1190 (1753).											
A. Gregorii, F. v. M. in Hook. Kew misc. IX, 14 (1857) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 223	M.fr. VII, 40; XI, 32.
CAMPTOSTEMON, Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876).											
C. Schultzei, Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876) ...	...	...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 32.
BOMBAX, Linné, sp. pl. I, 511 (1753).											
B. Malabarica, de Candolle, prodr. I, 479 (1824) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 223	M.fr. XI, 32.
STERCULIACEAE.											
Ventenat, Jard. de la Malm. II, 91 (1804).											
STERCULIA, Linné, Fl. Zeyl. 166 (1747).											
S. quadrifida, R. Brown in Benn. pl. Jav. rar. 233 (1852) ...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I, 227	M.fr. VI, 173.
S. laurifolia, F. v. M., fragm. VI, 172 (1868) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VI, 172.
BRACHYCHITON, Schott & Endlicher, Melat. bot. 34 (1832). (Trichosiphon, Poecilodermis, Delabechea).											
B. paradoxum, Schott, Melat. 34 (1832) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 227	—
B. Bidwilli, Hooker, Bot. Mag. t. 5133 (1859) ...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 228	—
B. discolor, F. v. M., fragm. I, 1 (1858) ...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I, 228	M.fr. I, 242; II, 177.
B. incanum, R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 228	—
B. luridum, C. Moore in F. v. M., fragm. I, 1 (1858) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 228	M.fr. I, 1; II, 177.
B. platanoides, R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 229	M.fr. VI, 173.
B. acerifolium, F. v. M., fragm. I, 1 (1858) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 229	M.fr. I, 1; VI, 173.
B. diversifolium, R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	W.A.	—	—	—	—	—	N.A.	B.fl.I, 230	—
B. populneum, R. Brown in Benn. pl. Jav. rar. 234 (1852) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I, 229	—
B. Delabechii, F. v. M., pl. of Viet. I, 157 (1860) ...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 230	—
B. Gregorii, F. v. M. in Hook. Kew misc. IX, 199 (1857) ...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr. IX, 137.
TARRIETIA, Blume, Bijdr. 227 (1825). (Argyrodendron).											
T. argyrodendron, Benth. Austr. I, 230 (1863) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I, 230	M.fr. VI, 173; IX, 42.
T. trifoliolata, F. v. M., fragm. IX, 43 (1875) ...	...	...	—	—	—	—	—	N.S.W.	Q.	—	M.fr. IX, 43.
HERITIERA, Dryander in Aiton, Hort. Kew III, 546 (1789).											
H. littoralis, Dryander in Aiton, Hort. Kew III, 546 (1789) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 231	M.fr. VI, 173.
UNGERIA, Schott & Endlicher, Melat. 27—31 (1832).											
U. floribunda, Schott & Endlicher, Melat. 27, t. 4 (1832) ...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 169.
HELICTERES, Plukenet Phytogr. 181, t. 245 (1692).											
H. Isora, Linné, sp. pl. 963 (1753) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 232	M.fr. VI, 173.
METHORIUM, Schott & Endlicher, Melat. 29—30 (1832).											
M. spicatum, Helicteris, Colebrooke in G. Don, syst. I, 507 (1831) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 39.
M. canum, Schott, Melat. 29, t. 5 (1832) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 232	—
M. dentatum, F. v. M. in Benth. Fl. Austr. I, 232 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 232	—
M. semiglabrum, F. v. M., fragm. V, 43 (1865) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. V, 43.
MELHANIA, Forskael, Fl. Aeg. Arab. 64 (1775).											
M. incana, Heyne in Wight & Arn. prodr. 68 (1834) ...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I, 234	M.fr. I, 69; VI, 174.
MELOCHIA, Dillenius, Hort. Eltham. 221, t. 176 (1732). (Riedleya).											
M. pyramidata, Linné, sp. pl. 674 (1753) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 234	M.fr. VI, 173.
M. corchorifolia, Linné, sp. pl. 675 (1753) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I, 235	M.fr. VI, 173.
HERMANNIA, Tournefort, inst. 656, t. 432 (1700). (Gilesia, Mahernia partly).											
H. Gilesii, F. v. M., fragm. IX, 42 (1875) ...	...	...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 42.
DICARPIDIUM, F. v. Mueller in Hook. Kew misc. IX, 302 (1857).											
D. monicum, F. v. M., in Hook. Kew misc. IX, 302 (1857) ...	...	...	—	—	—	—	—	—	N.A.	B.fl.I, 235	—
WALTHERIA, Linné, gen. pl. 203 (1737).											
W. Indica, Linné, sp. pl. 637 (1753) ...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.I, 236	M.fr. VI, 173.
ABROMA, N. Jacquin, Hort. Vindob. III, t. 1 (1776).											
A. fastuosa, Salisbury, Parad. Lond. 102 (1807) ...	...	...	—	—	—	—	—	Q.	—	B.fl.I, 236	M.fr. VI, 173.

L. oppositifolium, F. v. M., fragm. II, 5 (1860) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 262	M.fr. XI, 112.
L. micranthum, J. Hooker, Fl. Tasman. I, 51 (1860) ...	...	...	—	—	T.	—	—	—	B.f.I, 262	
L. macrophyllum, Graham in Bot. Mag. t. 3098 (1844) ...	...	...	—	—	—	N.S.W.	—	—	B.f.I, 262	
L. Baueri, Steetz in Lehm. pl. Preiss. II, 339 (1847) ...	...	...	—	S.A.	—	V.	N.S.W.	—	B.f.I, 263	M.fr. VI, 174; XI, 110.
L. Tepperi, F. v. M., fragm. XI, 109 (1881) ...	...	...	—	S.A.	—	—	—	—	—	M.fr. XI, 109.
L. rufum, R. Brown in Benth. Fl. Austr. I, 263 (1863) ...	...	...	—	—	—	N.S.W.	—	—	B.f.I, 263	
L. ferrugineum, Smith in Bot. Reposit. t. 308 (1799) ...	...	...	—	—	—	V.	N.S.W.	—	B.f.I, 263	
L. acutiflorum, Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 264	M.fr. XI, 109.
L. Oldfieldii, F. v. M., fragm. II, 6 (1860) ...	...	...	W.A.	—	—	—	—	—	—	M.fr. II, 6.
L. quinquenervium, Turcz. in Bull. Soc. Mosc. XXV, 146 (1852) ...	...	...	W.A.	—	—	—	—	—	—	M.fr. XI, 109.
L. Drummondii, Benth. Fl. Austr. I, 264 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 264	M.fr. VI, 174; XI, 112.
L. rosmarinifolium, Benth. Fl. Austr. I, 264 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 264	M.fr. XI, 112.
L. cordifolium, Endlicher in Hueg. enum. 10 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 265	M.fr. VI, 174; XI, 112.
L. Schulzenii, F. v. M., pl. of Vict. I, 145 implied (1861) ...	...	...	—	S.A.	—	V.	—	—	B.f.I, 265	
L. floribundum, Benth. Fl. Austr. I, 265 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 265	M.fr. XI, 112.
L. molle, Benth. Fl. Austr. I, 265 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 265	M.fr. XI, 111.
L. membranaceum, Benth. Fl. Austr. I, 266 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 266	M.fr. XI, 112.
L. bracteatum, Benth. Fl. Austr. I, 266 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 266	

LYSIOSEPALUM, F. v. Mueller, fragm. I, 143 (1859).

L. Barryanum, F. v. M., fragm. I, 143 (1859) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 267	M.fr. VI, 174.
L. rugosum, Benth. Fl. Austr. I, 267 (1863) ...	...	...	W.A.	—	—	—	—	—	B.f.I, 267	

TILIACEAE.

A. L. de Jussieu, Gen. 289 (1789).

BERRYA, Roxburgh, Pl. Coromand. III, 60, t. 264 (1819).

B. ammonilla, Roxburgh, pl. Coromand. III, 60, t. 264 (1819) ...	—	—	—	—	—	Q.	—	—	B.f.I, 268	
--	---	---	---	---	---	----	---	---	------------	--

GREWIA, Linné, syst. nat. 9 (1753); Linné, gen. pl. 276 (1737).

G. orientalis, Linné, sp. pl. 964 (1753) ...	...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 270
G. multiflora, A. L. de Jussieu in Ann. du Mus. IV, 89, t. 47 (1804) ...	—	—	—	—	—	—	—	Q.	N.A.	B.f.I, 270
G. breviflora, Benth. Fl. Austr. I, 270 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 270
G. latifolia, F. v. M. in Benth. Fl. Austr. I, 271 (1863) ...	...	...	—	—	—	N.S.W.	—	Q.	—	B.f.I, 271
G. polygama, Roxburgh, Fl. Ind. II, 588 (1832) ...	...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 271
G. xanthopetala, F. v. M. in Benth. Fl. Austr. I, 271 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 271
G. scabrella, Benth. Fl. Austr. I, 272 (1863) ...	...	...	—	—	—	—	—	Q.	—	B.f.I, 272
G. orbifolia, F. v. M. in Benth. Fl. Austr. I, 272 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 272
G. pleiostigma, F. v. M., fragm. VIII, 4 (1872) ...	...	...	—	—	—	—	—	Q.	—	—

TRIUMFETTA, Plumier, nov. gen. 40, t. 8 (1703).

T. procumbens, G. Forster, prodr. fl. insul. Austr. 35 (1786) ...	—	—	—	—	—	—	—	Q.	—	B.f.I, 273
T. appendiculata, F. v. M., fragm. III, 7 (1862) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 273
T. chactocarpa, F. v. M., fragm. XI, 61 (1879) ...	...	...	—	—	—	—	—	—	N.A.	—
T. glaucescens, R. Brown in Benth. Fl. Austr. I, 273 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 273
T. denticulata, R. Brown in Benth. Fl. Austr. I, 273 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 274
T. micracantha, F. v. M., fragm. III, 7 (1862) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 274
T. leptacantha, F. v. M., fragm. XI, 62 (1879) ...	...	...	—	—	—	—	—	—	N.A.	—
T. plumigera, F. v. M., fragm. I, 69 (1858) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 274
T. pilosa, Roth, nov. pl. spec. 223 (1821) ...	...	...	—	—	—	—	—	Q.	—	—
T. parviflora, Benth. Fl. Austr. I, 274 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 274

NETTOA, Baillon in Adans. VI, 238, t. 7 (1866).

N. chrozophorifolia, Baillon in Adans. VI, 238, 242 (1866) ...	—	—	—	—	—	—	—	—	N.A.	—
--	---	---	---	---	---	---	---	---	------	---

CORCHORUS, Tournefort, inst. 259, t. 135 (1700).

C. echinatus, Benth. Fl. Austr. I, 276 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 276
C. hygrophilus, Cunningham in Benth. Fl. Austr. I, 276 (1863) ...	...	...	—	—	—	—	—	—	Q.	—
C. Cunninghamii, F. v. M., fragm. III, 8 (1862) ...	...	...	—	—	—	N.S.W.	—	Q.	—	B.f.I, 276
C. olitorius, Linné, sp. pl. 529 (1753) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 276
C. tridens, Linné, mantiss. alter. 566 (1771) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 276
C. acutangulus, Lamarck, encycl. meth. II, 104 (1786) ...	...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 277
C. fascicularis, Lamarck, encycl. meth. II, 104 (1786) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 277
C. pumilio, Benth. Fl. Austr. I, 277 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 277
C. vermicularis, F. v. M., fragm. III, 10 (1862) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 277
C. tomentellus, F. v. M., fragm. III, 10 (1862) ...	...	...	—	—	—	—	—	Q.	—	B.f.I, 278
C. sidoides, F. v. M., fragm. III, 9 (1862) ...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.f.I, 278
C. leptocarpus, Cunningham in Benth. Fl. Austr. I, 278 (1863) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 278
C. elachocarpus, F. v. M., fragm. VIII, 6 (1872) ...	...	...	—	—	—	—	—	—	N.A.	—
C. Walcottii, F. v. M., fragm. III, 9 (1862) ...	...	...	—	—	—	—	—	—	N.A.	B.f.I, 278
C. trilocularis, Linné, mantiss. 77 (1771) ...	...	...	—	—	—	—	—	Q.	—	—

SLOANEA, Linné, hort. Cliffort. 210 (1737). (Echinocarpus.)

S. Australis, F. v. M., fragm. IV, 91 (1864) ...	...	...	—	—	—	N.S.W.	Q.	—	B.f.I, 279	M.fr. VI, 172; VIII, 2.
S. Langii, F. v. M., fragm. V, 28 (1865) ...	...	...	—	—	—	—	Q.	—	—	M.fr. VIII, 2.
S. Macbrydei, F. v. M., fragm. VI, 170 (1868) ...	...	...	—	—	—	—	Q.	—	—	M.fr. VI, 170.
S. Woollsii, F. v. M., fragm. VI, 171 (1868) ...	...	...	—	—	—	N.S.W.	Q.	—	—	M.fr. VIII, 2.

ARISTOTELIA, L'Héritier, stirp. II, 21, t. 16 (1784). (Friesia.)

A. peduncularis, J. Hooker, Fl. Nov. Fel. I, 33 (1853) ...	...	...	—	—	T.	—	—	—	B.f.I, 280	M.fr. VIII, 2.
A. Australasica, F. v. M., fragm. II, 79 (1860) ...	...	...	—	—	—	N.S.W.	—	—	B.f.I, 280	M.fr. VIII, 2.
A. megalosperma, F. v. M., fragm. IX, 84 (1875) ...	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 84.

ELAEOCARPUS, Burman, thesaur. Zeil. 39, t. 40 (1737). (Elaeocarpos, Dicera.)

E. holopetalus, F. v. M., fragm. II, 143 (1861) ...	...	...	—	—	—	V.	N.S.W.	—	B.f.I, 281	M.fr. IV, 173; VI, 172;
E. Arnheimicus, F. v. M., Docum. intercol. Exhib. 24 (1867) ...	...	...	—	—	—	—	—	—	N.A.	M.fr. X, 4.

[VIII.]

<i>E. obovatus</i> , G. Don, gen. syst. dichl. pl. I, 599 (1831) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 281	M.fr. II, 80; VI, 172.
<i>E. cyaneus</i> , Aiton, Epit. hort. Kew, addend. 367 (1814) ...	—	—	—	T. V.	N.S.W.	Q.	—	B.f.I, 281	M.fr. VI, 172; VIII, 2. [12.
<i>E. grandis</i> , F. v. M., fragm. II, 81 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 281	M.fr. II, 175; VI, 172; VIII,
<i>E. foveolatus</i> , F. v. M., fragm. V, 157 (1866) ...	—	—	—	—	—	Q.	—	—	M.fr. VI, 172; VIII, 2.
<i>E. ruminatus</i> , F. v. M., fragm. VIII, 1 (1872) ...	—	—	—	—	—	Q.	—	—	M.fr. X, 4.
<i>E. Grahami</i> , F. v. M., fragm. X, 3 (1876) ...	—	—	—	—	—	Q.	—	—	M.fr. X, 3.
<i>E. sericopetalus</i> , F. v. M., fragm. VI, 171 (1868) ...	—	—	—	—	—	Q.	—	—	M.fr. VI, 171.

EUPHORBIACEAE.

A. L. de Jussieu, Gen. 384 (1789) from B. de Jussieu (1759).

CALYCOPEPLUS, Planchon in Bull. de la Soc. Bot. de Fr. VIII, 31 (1861).

<i>C. ephedroides</i> , Planchon in Bull. Soc. Bot. Fr. VIII, 31 (1861) ...	W.A.	—	—	—	—	—	—	B.f. VI, 53	
<i>C. marginatus</i> , Benth. in Fl. Austr. VI, 53 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 53	

EUPHORBIA, Linné, gen. pl. 152 (1737) from Plinius.

<i>E. Atoto</i> , G. Forster, florul. ins. Austr. prodr. 36 (1786) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 46	
<i>E. Sparmanni</i> , Boissier, Cent. Euphorb. 5 (1860) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 46	
<i>E. Mitchelliana</i> , Boissier in De Candolle, prodr. XV, 25 (1866) ...	W.A.	—	—	—	—	Q.	N.A.	B.f. VI, 47	M.fr. VI, 182.
<i>E. schizolepis</i> , F. v. M. in Benth. Fl. Austr. VI, 47 (1873) ...	—	—	—	—	—	—	N.A.	B.f. VI, 47	
<i>E. Schultzii</i> , Benth. in Fl. Austr. VI, 47 (1873) ...	—	—	—	—	—	—	N.A.	B.f. VI, 47	
<i>E. Armstrongiana</i> , Boissier in De Candolle, prodr. XV, 47 (1866) ...	—	—	—	—	—	—	N.A.	B.f. VI, 48	
<i>E. erythrantha</i> , F. v. M., fragm. II, 152 (1861) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. VI, 48	M.fr. XI, 64.
<i>E. Muelleri</i> , Boissier in De Candolle, prodr. XV, 27 (1866) ...	—	—	—	—	—	—	N.A.	B.f. VI, 48	
<i>E. Drummondii</i> , Boissier, Cent. Euphorb. 14 (1860) ...	W.A.	S.A.	T. V.	N.S.W.	Q.	N.A.	B.f. VI, 49	B.f. VI, 49	M.fr. VI, 182.
<i>E. alsiniflora</i> , Baillon, Adans. VI, 288 (1866) ...	—	—	—	—	—	—	N.A.	B.f. VI, 49	
<i>E. Wheeleri</i> , Baillon, Adans. VI, 286 (1866) ...	—	S.A.	—	—	—	—	N.A.	B.f. VI, 49	
<i>E. myrtoidea</i> , Boissier in De Candolle, prodr. XV, 15 (1866) ...	W.A.	—	—	—	—	—	N.A.	B.f. VI, 50	
<i>E. micradenia</i> , Boissier in De Candolle, prodr. XV, 27 (1866) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 50	
<i>E. Macgillivrayi</i> , Boissier in De Candolle, prodr. XV, 26 (1866) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 50	
<i>E. serrulata</i> , Reinwardt in Blum. Bijdrag. 635 (1826) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 51	
<i>E. filipes</i> , Benth. in Fl. Austr. VI, 51 (1873) ...	—	—	—	—	—	—	N.A.	B.f. VI, 51	
<i>E. Careyi</i> , F. v. M., fragm. XI, 64 (1879) ...	—	—	—	—	—	—	N.A.	—	M.fr. XI, 64.
<i>E. pilulifera</i> , Linné, sp. pl. 454 (1753) ...	—	—	—	—	—	Q.	—	B.f. VI, 51	
<i>E. Norfolkiana</i> , Boissier in De Cand., prodr. XV, 206 (1866) ...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 169.
<i>E. eremophila</i> , Cunningham in Mitch. Trop. Austr. 348 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VI, 52	

MONOTAXIS, Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829). (Hippocrepandra.)

<i>M. macrophylla</i> , Benth. in Fl. Austr. VI, 79 (1873) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 79	
<i>M. linifolia</i> , Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 79	
<i>M. occidentalis</i> , Endlicher in Hueg. enum. 19 (1837) ...	W.A.	—	—	—	—	—	—	B.f. VI, 79	
<i>M. lurida</i> , Benth. in Fl. Austr. VI, 80 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 80	
<i>M. megacarpa</i> , F. v. M., fragm. IV, 143 (1865) ...	W.A.	—	—	—	—	—	—	B.f. VI, 80	M.fr. IV, 143.
<i>M. gracilis</i> , Baillon, Adans. VI, 293 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 80	
<i>M. grandiflora</i> , Endlicher in Hueg. enum. 19 (1837) ...	W.A.	—	—	—	—	—	—	B.f. VI, 81	
<i>M. luteiflora</i> , F. v. M., fragm. X, 51 (1877) ...	W.A.	—	—	—	—	—	—	—	M.fr. X, 51.

PORANTHERA, Rudge in Transact. Linn. Soc. X, 302, t. 22 (1811).

<i>P. ericifolia</i> , Rudge in transact. Linn. Soc. X, 302, t. 22 (1811) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 55	
<i>P. ericoides</i> , Klotzsch in Lehm. pl. Preiss. II, 239 (1847) ...	W.A.	S.A.	—	V.	—	—	—	B.f. VI, 55	
<i>P. Huegelii</i> , Klotzsch in Lehm. pl. Preiss. II, 231 (1847) ...	W.A.	—	—	—	—	—	—	B.f. VI, 55	
<i>P. corymbosa</i> , Brongniart in Duperr. Voy. Cog. Bot. 219, t. 56 (1826) ...	—	—	—	V.	N.S.W.	—	—	B.f. VI, 56	M.fr. VI, 183.
<i>P. microphylla</i> , Brongniart in Duperr. Voy. Cog. Bot. 218, t. 50 (1826) ...	W.A.	S.A.	T. V.	N.S.W.	Q.	N.A.	—	B.f. VI, 56	M.fr. XI, 20.

MICRANTHEUM, Desfontaines in Mém. du Mus. hist. nat. IV, 253 (1818). (Caletia.)

<i>M. ericoides</i> , Desfontaines in Mém. du Mus. IV, 253, t. 14 (1818) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 57	
<i>M. hexandrum</i> , J. Hooker in Lond. Journ. VI, 283 (1847) ...	—	S.A.	T. V.	N.S.W.	—	—	—	B.f. VI, 57	

PSEUDANTHUS, Sieber in Sprengel, syst. veg. cur. post. 25 (1827). (Stachystemon.)

<i>P. pimeloides</i> , Sieber in Sprengel, syst. veg. cur. post. 25 (1827) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 59	
<i>P. ovalifolius</i> , F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857) ...	—	—	T. V.	N.S.W.	—	—	—	B.f. VI, 59	
<i>P. micranthus</i> , Benth. in Fl. Austr. VI, 59 (1873) ...	—	S.A.	—	—	—	—	—	B.f. VI, 59	
<i>P. divaricatissimus</i> , Benth. in Fl. Austr. VI, 60 (1873) ...	—	—	—	V.	N.S.W.	—	—	B.f. VI, 60	
<i>P. orientalis</i> , F. v. M., fragm. II, 14 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 60	M.fr. II, 14.
<i>P. occidentalis</i> , F. v. M., fragm. I, 107 (1859) ...	W.A.	—	—	—	—	—	—	B.f. VI, 60	M.fr. I, 107.
<i>P. nematophorus</i> , F. v. M., fragm. II, 14 (1860) ...	W.A.	—	—	—	—	—	—	B.f. VI, 60	M. II, 14.
<i>P. polyandrus</i> , F. v. M., fragm. II, 153 (1861) ...	W.A.	—	—	—	—	—	—	B.f. VI, 62	M.fr. II, 153; IV, 173
<i>P. brachyphyllus</i> , F. v. M. in Transact. R. Soc. N.S. Wales 11 (1881) ...	W.A.	—	—	—	—	—	—	B.f. VI, 62	
<i>P. vermicularis</i> , F. v. M. in Transact. R. Soc. N.S.W. 11 (1881) ...	W.A.	—	—	—	—	—	—	B.f. VI, 62	

BEYERIA, Miquel in Ann. sc. nat. sér. trois. I, 350 (1844). (Calypotrostigma, Beyerlopsis.)

<i>B. viscosa</i> , Miquel in Ann. des. sc. nat. sér. trois. I, 350, t. 15 (1844) ...	W.A.	S.A.	T. V.	N.S.W.	Q.	—	—	B.f. VI, 64	M.fr. I, 230.
<i>B. lasiocarpa</i> , F. v. M. in Benth. Fl. Austr. VI, 65 (1873) ...	—	—	—	V.	N.S.W.	—	—	B.f. VI, 65	M.fr. VI, 182.
<i>B. opaca</i> , F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854) ...	—	S.A.	T. V.	N.S.W.	—	—	—	B.f. VI, 65	M.fr. VI, 182.
<i>B. uncinata</i> , F. v. M. in Benth. Fl. Austr. VI, 65 (1873) ...	—	S.A.	—	—	—	—	—	B.f. VI, 65	
<i>B. latifolia</i> , Baillon, Adans. VI, 304 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 66	
<i>B. cygnorum</i> , Baillon, Adans. VI, 309 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 66	
<i>B. cinerea</i> , Baillon, Adans. VI, 309 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 66	
<i>B. cyanescens</i> , Benth. in Fl. Austr. VI, 66 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 66	
<i>B. lepidopetala</i> , F. v. M., fragm. I, 230 (1859) ...	W.A.	—	—	—	—	—	—	B.f. VI, 67	M.fr. I, 230; III, 164.
<i>B. similis</i> , Baillon, Adans. VI, 309 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 67	
<i>B. brevifolia</i> , Baillon, Adans. VI, 309 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VI, 67	
<i>B. Drummondii</i> , J. Mueller in Linnaea XXXIV, 58 (1865) ...	W.A.	—	—	—	—	—	—	B.f. VI, 68	
<i>B. tristigma</i> , F. v. M., fragm. VI, 181 (1868) ...	—	—	—	—	—	Q.	—	B.f. VI, 68	M.fr. VI, 181.

RICINOCARPUS, Desfontaines in Mém. du Mus. III, 459, t. 22 (1817). (Roeperia.)									
R. pinifolius, Desfontaines in Mém. du Mus. III, 459, t. 22 (1817)	...	—	—	T. V. N.S.W.	Q.	—	B.f. VI, 70	M.fr. VI, 182.	
R. tuberculatus, J. Mueller in Linnaea XXXIV, 60 (1865)	...	W.A.	—	—	—	—	B.f. VI, 70		
R. cyanescens, J. Mueller in Linnaea XXXIV, 60 (1865)	...	W.A.	—	—	—	—	B.f. VI, 70		
R. psilocladus, Benth. Fl. Austr. VI, 71 (1873)	...	W.A.	—	—	—	—	B.f. VI, 70		
R. glaucus, Endlicher in Hug. enum. 18 (1837)	...	W.A.	—	—	—	—	B.f. VI, 71	M.fr. VI, 182.	
R. major, J. Mueller in Linnaea XXXIV, 59 (1865)	...	—	—	T.	—	—	B.f. VI, 72		
R. Bowmanii, F. v. M., fragm. I, 181 (1859)	...	—	—	—	N.S.W.	Q.	B.f. VI, 72	M.fr. I, 181.	
R. ledifolius, F. v. M., fragm. I, 76 (1858)	...	—	—	—	—	Q.	B.f. VI, 72	M.fr. I, 76.	
R. rosmarinifolius, Benth. Fl. Austr. VI, 72 (1873)	...	—	—	—	—	N.A.	B.f. VI, 72		
R. marginatus, Benth. Fl. Austr. VI, 73 (1873)	...	—	—	—	—	N.A.	B.f. VI, 73		
R. speciosus, J. Mueller in De Candolle, prodr. XV, 204 (1863)	...	—	—	—	N.S.W.	—	B.f. VI, 73		
R. trichophorus, J. Mueller in Linnaea XXXIV, 60 (1865)	...	W.A.	—	—	—	—	B.f. VI, 73		
R. muricatus, J. Mueller in Linnaea XXXIV, 61 (1865)	...	W.A.	—	—	—	—	B.f. VI, 73		
R. velutinus, F. v. M., fragm. IX, 2 (1875)	...	W.A.	—	—	—	—	—	M.fr. IX, 2.	
BERTYA, Planchon in Hook. Lond. Journ. Bot. IV, 472, t. 16 (1845).									
B. gummifera, Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	—	—	—	N.S.W.	—	B.f. VI, 75		
B. pinifolia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	—	—	—	N.S.W.	Q.	B.f. VI, 75		
B. quadrise-pala, F. v. M., fragm. X, 52 (1876)	...	W.A.	—	—	—	—	—	M.fr. X, 52.	
B. Cunninghamii, Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	—	—	V.	N.S.W.	—	B.f. VI, 75	M.fr. X, 52.	
B. dimerostigma, F. v. M. in Wing's S. Science Journ. II (1882)	...	W.A.	—	—	—	—	—	M.fr. XII.	
B. rosmarinifolia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	—	—	T.	N.S.W.	—	B.f. VI, 76		
B. Mitchellii, J. Mueller in Linnaea XXXIV, 63 (1865)	...	—	—	S.A.	V.	N.S.W.	B.f. VI, 76		
B. oppositifolia, F. v. M. & O'Shanesyn in S. Sc. Rec. II, May (1882)	...	—	—	—	—	Q.	—	M.fr. XII.	
B. oleae-folia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	...	—	—	—	N.S.W.	Q.	B.f. VI, 76		
B. rotundifolia, F. v. M., fragm. IV, 34 (1864)	...	—	—	S.A.	—	—	B.f. VI, 77	M.fr. IV, 34.	
B. pedicellata, F. v. M., fragm. IV, 143 (1865)	...	—	—	—	—	Q.	B.f. VI, 77	M.fr. IV, 143.	
B. pomaderrioides, F. v. M., fragm. IV, 34 (1864)	...	—	—	—	N.S.W.	—	B.f. VI, 77	M.fr. IV, 34.	
B. Bendlayi, F. v. M., fragm. VIII, 141 (1874)	...	—	—	—	V.	N.S.W.	—	M.fr. VIII, 141.	
AMPEREA, Adr. de Jussieu, Euphorb. gen. 35, t. 10 (1824).									
A. protensa, Nees in Lehmann, pl. Preiss. II, 229 (1847)	...	W.A.	—	—	—	—	B.f. VI, 82		
A. volubilis, F. v. M. in Benth. Fl. Austr. VI, 82 (1873)	...	W.A.	—	—	—	—	B.f. VI, 82	M.fr. VI, 182.	
A. micrantha, Benth. Fl. Austr. VI, 83 (1873)	...	W.A.	—	—	—	—	B.f. VI, 83		
A. conferta, Benth. Fl. Austr. VI, 83 (1873)	...	W.A.	—	—	—	—	B.f. VI, 83		
A. ericoides, Adr. de Jussieu, Tent. Euphorb. 112, t. 10 (1824)	...	W.A.	—	—	—	—	B.f. VI, 83		
A. spartioides, Brongniart in Duperr. Voy. Cog. 221, t. 49 (1826)	...	S.A.	T. V.	N.S.W.	Q.	—	B.f. VI, 84		
ANTIDESMA, Linné, Fl. Zeyl. 169 (1747).									
A. Ghaesembilla, Gaertner, de Fruct. I, 189, t. 39 (1788)	...	—	—	—	—	N.A.	B.f. VI, 85	M.fr. VI, 183.	
A. Dallachyanum, Baillon, Adans. VI, 337 (1866)	...	—	—	—	—	Q.	N.A.	B.f. VI, 85	M.fr. VI, 183.
A. Bunius, Sprengel, syst. veg. I, 826 (1825)	...	—	—	—	—	Q.	—	B.f. VI, 86	
A. parviflorum, Thwaites & F. v. M., fragm. IV, 86 (1864)	...	—	—	—	—	Q.	—	B.f. VI, 86	M.fr. IV, 86; VI, 183.
A. Schultzii, Benth. Fl. Austr. VI, 86 (1873)	...	—	—	—	—	N.A.	B.f. VI, 86		
A. erostre, F. v. M. in Benth. Fl. Austr. VI, 87 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 87	
A. sinuatum, Benth. Fl. Austr. VI, 87 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 87	
ANDRACHNE, Linné, syst. nat. 9 (1735); Linné, gen. pl. 287 (1737).									
A. Decaisnei, Benth. Fl. Austr. VI, 88 (1873)	...	W.A.	—	—	—	Q.	N.A.	B.f. VI, 88	M.fr. VI, 183.
ACTEPHILA, Blume, Bijdr. XII, 581 (1825).									
A. grandifolia, Baillon, Adans. VI, 330 & 360, t. 10 (1866)	...	—	—	—	N.S.W.	Q.	—	B.f. VI, 89	
A. Mooreana, Baillon, Adans. VI, 330 & 336 (1866)	...	—	—	—	N.S.W.	Q.	—	B.f. VI, 89	
A. latifolia, Benth. Fl. Austr. VI, 89 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 89	
A. petiolaris, Benth. Fl. Austr. VI, 89 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 89	
A. sessilifolia, Benth. Fl. Austr. VI, 90 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 90	
DISSILIARIA, F. v. M. in Baillon, Adans. VII, 356, pl. 1 (1867). (Choriceras.)									
D. baloghoides, F. v. M. in Baillon, Adans. VII, 356 (1867)	...	—	—	—	—	Q.	—	B.f. VI, 90	
D. Muelleri, Baillon, Adans. VII, 359, t. 1 (1867)	...	—	—	—	—	Q.	—	B.f. VI, 91	
D. tricornis, Benth. Fl. Austr. VI, 91 (1873)	...	—	—	—	—	Q.	N.A.	B.f. VI, 91	
PETALOSTIGMA, F. v. M. in Hook. Kew Misc. IX, 17 (1857). (Hylococcus.)									
P. quadriloculare, F. v. M. in Hook. Kew Misc. IX, 17 (1857)	...	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 92	M.fr. VI, 182.
PHYLLANTHUS, Linné, gen. pl. 282 (1737), from J. Commelyn. (Glochidion, Synostemon, Bradleia.)									
P. Ferdinandi, J. Mueller in Regensb. Flora, 379 (1865)	...	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 96	M.fr. VI, 183.
P. lobocarpus, Benth. Fl. Austr. VI, 97 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 97	
P. ditassoides, J. Mueller in Regensb. Flora, 487 (1864)	...	—	—	—	—	—	N.A.	B.f. VI, 97	
P. Adami, J. Mueller in De Candolle, prodr. XV, 327 (1866)	...	—	—	—	—	—	N.A.	B.f. VI, 97	
P. thesioides, Benth. Fl. Austr. VI, 98 (1873)	...	—	—	—	N.S.W.	Q.	—	B.f. VI, 98	
P. hirtellus, J. Mueller in De Candolle, prodr. XV, 326 (1866)	...	—	—	—	—	Q.	—	B.f. VI, 98	M.fr. III, 89.
P. rigens, J. Mueller in Regensb. Flora, 513 (1864)	...	—	—	—	N.S.W.	Q.	—	B.f. VI, 99	
P. ochrophyllus, Benth. Fl. Austr. VI, 99 (1873)	...	—	—	—	—	—	N.A.	B.f. VI, 99	
P. rigidulus, F. v. M. in Linnaea XXXIV, 72 (1865)	...	—	—	—	—	—	N.A.	B.f. VI, 99	
P. ramosissimus, J. Mueller in Linnaea XXXIV, 70 (1865)	...	—	—	—	N.S.W.	Q.	—	B.f. VI, 100	
P. Tatei, F. v. M. in Wing's South. Science Record II, 55 (1882)	...	—	S.A.	—	—	—	—	—	
P. rhytidosperrnus, F. v. M. in Linnaea XXXIV, 70 (1865)	...	—	—	—	—	—	N.A.	B.f. VI, 100	
P. albiflorus, F. v. M. in Linnaea XXXIV, 70 (1865)	...	—	—	—	—	Q.	—	B.f. VI, 100	
P. crassifolius, J. Mueller in Regensb. Flora, 513 (1864)	...	W.A.	—	—	—	—	—	B.f. VI, 100	
P. elachophyllus, F. v. M. in Benth. Fl. Austr. VI, 101 (1873)	...	—	—	—	—	Q.	—	B.f. VI, 101	
P. uberiflorus, F. v. M. in Adansonia, 343 (1866)	...	—	—	—	—	Q.	—	B.f. VI, 101	
P. reticulatus, Poiret, encycl. meth. V, 298 (1804)	...	—	—	—	—	—	N.A.	B.f. VI, 101	
P. baccatus, F. v. M. in Benth. Fl. Austr. VI, 102 (1873)	...	—	—	—	—	—	N.A.	B.f. VI, 102	

P. Urinaria, Linné, sp. pl. 982 (1753)	...	...	...	...	...	...	...	N.A.	B.f. VI, 102	
P. trachygynne, Benth. Fl. Austr. VI, 103 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 103	
P. Maderaspatanus, Linné, sp. pl. 982 (1753)	...	...	...	...	...	...	...	N.A.	B.f. VI, 103	
P. Mitchellii, Benth. Fl. Austr. VI, 103 (1873)	...	...	...	...	...	...	...	Q.	B.f. VI, 103	
P. Gastroemii, J. Mueller in De Cand., prodr. XV, 358 (1866)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 104	
P. Dallachyanus, Benth. Fl. Austr. VI, 104 (1873)	...	...	...	...	...	...	Q.	—	B.f. VI, 104	
P. subcrenulatus, F. v. M., fragm. I, 108 (1859)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 104	M.fr. I, 108.
P. calycinus, Labillardiere, Nov. Holl. pl. spec. II, 75, t. 225 (1806)	W.A.	S.A.	—	—	—	—	—	—	B.f. VI, 105	
P. flagellaris, Benth. Fl. Austr. VI, 106 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 106	
P. similis, J. Mueller in Linnaea XXXIV, 71 (1865)	...	...	...	...	...	...	...	Q.	B.f. VI, 106	
P. microcladus, J. Mueller in Linnaea XXXIV, 71 (1865)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 106	
P. grandisepalus, F. v. M. in Linnaea XXXIV, 72 (1863)	...	...	...	...	...	...	...	N.A.	B.f. VI, 106	
P. Carpentariae, J. Mueller in Linnaea XXXIV, 72 (1863)	...	...	...	...	...	...	...	N.A.	B.f. VI, 107	
P. Fuernrohrii, F. v. M. in Transact. phil. Soc. Vict. I, 15 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	—	B.f. VI, 107	
P. hebecarpus, Benth. Fl. Austr. VI, 108 (1873)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 108
P. lacunarius, F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854)	—	S.A.	—	V.	N.S.W.	—	—	—	B.f. VI, 108	
P. trachyspermus, F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854)	—	S.A.	—	V.	N.S.W.	—	—	—	B.f. VI, 108	
P. australis, J. Hooker in Lond. Journ. VI, 284 (1847)	...	...	...	...	...	T.	N.S.W.	—	B.f. VI, 108	
P. thymoides, Sieber in Linnaea XXVIII, 566 (1866)	...	...	...	...	...	S.A.	V.	N.S.W.	—	B.f. VI, 109
P. scaber, Klotzsch in Lehm. pl. Preiss. I, 179 (1844)	...	...	...	...	...	...	...	—	B.f. VI, 109	
P. indigoferoides, Benth. Fl. Austr. VI, 110 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 110	
P. aridus, Benth. Fl. Austr. VI, 110 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 110	
P. Gunnii, J. Hooker in Lond. Journ. VI, 284 (1847)	...	...	...	...	...	S.A.	T.	V.	N.S.W.	M.fr. VI, 183.
P. simplex, Retzius, observ. V, 29 (1789)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 111
P. filiculis, Benth. Fl. Austr. VI, 111 (1873)	...	...	...	...	...	...	...	N.S.W.	—	B.f. VI, 111
P. minutiflorus, F. v. M. in Linnaea XXXIV, 75 (1863)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 112
P. Armstrongii, Benth. Fl. Austr. VI, 112 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 112	
P. buxifolius, Reinwardt in Blume, catal. plant. Buitenz. 106 (1823)	...	...	...	...	...	...	...	Q.	—	M.fr. X, 121.
(Melanthesa, Melanthesiopsis.)										
B. cernua, J. Mueller in De Candolle, prodr. XV, 439 (1866)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 113
B. oblongifolia, J. Mueller in De Candolle, prodr. XV, 110 (1866)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 114	
B. stipitata, J. Mueller in De Candolle, prodr. XV, 442 (1866)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 114
B. rhynchocarpa, Benth. Fl. Austr. VI, 114 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 114	
SECURINEGA, A. L. de Jussieu, Gen. 388 (1789). (Flueggea.)										
S. Abyssinica, A. Richard, tentam. fl. Abyss. II, 256 (1851)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 116
S. Leucopyrus, J. Mueller in De Cand., prodr. XV, 451 (1866)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 116
NEOROEPEA, F. & J. Mueller in De Candolle, prodr. XV, 489 (1866).										
N. buxifolia, F. & J. Mueller in De Cand., prodr. XV, 489 (1866)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 116
N. Banksii, Benth. Fl. Austr. VI, 117 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 117
BISCHOFFIA, Blume, Bijdr. 1168 (1825).										
B. Javanica, Blume, Bijdr. 1168 (1825)	...	...	...	...	...	...	...	Q.	—	M.fr. VIII, 141.
HEMICYCLIA, Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833).										
H. sepiaria, Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833)	...	...	...	...	...	...	...	N.A.	B.f. VI, 117	M.fr. IV, 119; VI, 182.
H. Australasica, J. Mueller in De Candolle, prodr. XV, 487 (1866)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 118	
H. lasiogyne, F. v. M., fragm. IV, 119 (1864)	...	...	...	...	...	...	...	N.A.	B.f. VI, 118	M.fr. IV, 119.
APOROSA, Blume, Bijdr. 514 (1824).										
A. Australiana, F. v. M., fragm. XII ined.	...	...	...	...	...	...	...	Q.	—	M.fr. VIII, 141; XII.
BRIDELIA, Willdenow, Spec. IV, 978 (1805).										
B. exaltata, F. v. M., fragm. III, 32 (1862)	...	...	...	...	...	N.S.W.	—	—	B.f. VI, 119	M.fr. III, 32.
B. ovata, Decaisne in nouv. ann. du Mus. V, 484 (1835)	...	...	...	...	...	...	...	N.A.	B.f. VI, 120	
B. tomentosa, Blume, Bijdr. 597 (1825)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 120
B. faginea, F. v. M. in Benth. Fl. Austr. VI, 120 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 120
CLEISTANTHUS, J. Hooker, icon. pl. t. 779 (1847). (Lebediera, Amanoa.)										
C. Cunninghamii, J. Mueller in De Cand., prodr. XV, 506 (1866)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 122	
C. apodus, Benth. Fl. Austr. VI, 122 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 122
C. Dallachyanus, Baillon in Benth. Fl. Austr. VI, 122 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 122
C. semiopacus, F. v. M. in Benth. Fl. Austr. VI, 123 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 123
CROTON, Linné, gen. pl. 288 (1737).										
C. Schultzii, Benth. Fl. Austr. VI, 124 (1873)	...	...	...	...	...	...	...	N.A.	B.f. VI, 124	
C. insularis, Baillon, Adans. II, 217 (1862)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 124	
C. phleballoides, F. v. M. in Benth. Fl. Austr. VI, 125 (1873)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 125	M.fr. IV, 140.
C. opponens, F. v. M. in Benth. Fl. Austr. VI, 125 (1873)	...	...	...	...	...	...	...	Q.	B.f. VI, 125	
C. tomentellus, F. v. M., fragm. IV, 141 (1864)	...	...	...	...	...	...	...	N.A.	B.f. VI, 126	M.fr. IV, 141.
C. Verreauxii, Baillon, Etud. Euph. 357 (1858)	...	...	...	...	...	N.S.W.	Q.	N.A.	B.f. VI, 126	M.fr. IV, 141; VI, 186.
C. acronychioides, F. v. M., fragm. IV, 142 (1864)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 127
C. triacros, F. v. M., fragm. VI, 185 (1867)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 127
C. Arnemicus, F. v. M. in Linnaea XXXIV, 112 (1865)	...	...	...	...	...	...	...	Q.	N.A.	B.f. VI, 127
ALEURITES, R. & G. Forster, char. gen. 111, t. 56 (1776).										
A. triloba, R. & G. Forster, char. gen. 111, t. 56 (1776)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 128
CLAOSYXON, Adr. de Jussieu, Euphorb. gen. 43, t. 14 (1824).										
C. angustifolium, J. Mueller in Linnaea XXXIV, 165 (1865)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 129
C. tenerifolium, F. v. M. in Baillon, Adans. VI, 323 (1866)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 130
C. australe, Baillon, Etud. Euphorb. 493 (1858)	...	...	...	...	...	V.	N.S.W.	Q.	—	B.f. VI, 130
C. Hillii, Benth. Fl. Austr. VI, 131 (1873)	...	...	...	...	...	...	...	Q.	—	B.f. VI, 131
ACALYPHA, Royen in Linné, hort. Cliffort. 495 (1737).										
A. nemorum, F. v. M. in Linnaea XXXIV, 38 (1865)	...	...	...	...	...	N.S.W.	Q.	—	B.f. VI, 132	

A. eremorum, F. v. M. in Regensb. Flora 440 (1864) ...	—	—	—	—	Q.	—	B.f. VI, 132	
A. capillipes, F. v. M. in Linnaea XXXIV, 40 (1865) ...	—	—	—	—	N.S.W.	—	B.f. VI, 133	
ADRIANA, Gaudichaud in Ann. sc. nat. V, 223 (1825). (Trachycaryon.)								
A. tomentosa, Gaudichaud in Ann. sc. nat. V, 223 (1825) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VI, 134 M.fr. IX, 2.
A. quadrupartita, Gaudichaud in Freyc. voy. Bot. 489 (1826) ...	W.A.	S.A.	—	V.	—	—	—	B.f. VI, 135
ALCHORNEA, Solander in Swartz, prodr. 698 (1788). (Coelebobogyne, Cladodes.)								
A. ilicifolia, J. Mueller in Linnaea XXXIV, 170 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 136
A. Thozetiana, F. v. M. in Benth. Fl. Austr. VI, 137 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 137
TRAGIA, Plumier, gen. 14, t. 12 (1703).								
T. Novae Hollandiae, J. Mueller in Linnaea XXXIV, 180 (1865) ...	—	—	—	—	—	Q.	—	B.f. VI, 137 M.fr. VI, 183.
MALLOTUS, Loureiro, Fl. Cochinch. II, 635 (1790). (Echinus, Rottlera, Echinocroton.)								
M. ricinoides, J. Mueller in Linnaea XXXIV, 187 (1865) ...	—	—	—	—	—	Q.	—	B.f. VI, 140 M.fr. IV, 138; VI, 184.
M. Chinensis, Loureiro, Fl. Cochinch. II, 635 (1770) ...	—	—	—	—	—	Q.	—	B.f. VI, 140 M.fr. VI, 184.
M. claoxyloides, J. Mueller in Linnaea XXXIV, 192 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 140 M.fr. VI, 184.
M. Philippinensis, J. Mueller in Linnaea XXXIV, 196 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 141 M.fr. VI, 183.
M. angustifolius, Benth. in Fl. Austr. VI, 141 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 142
M. polyadenos, F. v. M., fragm. VI, 184 (1867) ...	—	—	—	—	—	Q.	—	B.f. VI, 142 M.fr. VI, 184.
M. repandus, J. Mueller in Linnaea XXXIV, 197 (1865) ...	—	—	—	—	—	Q.	—	B.f. VI, 142 M.fr. VI, 185.
M. nesophilus, F. v. M. in Linnaea XXXIV, 196 (1865) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 143 M.fr. VI, 184.
M. discolor, F. v. M. in Benth. Fl. Austr. VI, 143 (1873) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 143 M.fr. IV, 139.
MACARANGA, Petit-Thouars, Gen. Madag. n. 88 (1809). (Mappa.)								
M. Dallachyi, F. v. M. in Benth. Fl. Austr. VI, 144 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 144 M.fr. VI, 184.
M. subdentata, Benth. in Fl. Austr. VI, 145 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 145
M. inamoena, F. v. M. in Benth. Fl. Austr. VI, 145 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 145
M. involutifolia, Baillon, Etud. Euphorb. 432 (1858) ...	—	—	—	—	—	Q.	—	B.f. VI, 146 M.fr. IV, 139; VI, 183.
M. Tanaria, J. Mueller in De Candolle, prodr. XV, 997 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 146 M.fr. VI, 183.
CODIAEUM, Rumphius, herbar. Amboin. IV, 65, t. 25 (1743).								
C. variegatum, Blume, Bijdrag. 605 (1825) ...	—	—	—	—	—	Q.	—	B.f. VI, 147 M.fr. VI, 182.
BALOGHIA, Endlicher, prodr. fl. Norfolk. 84 (1833).								
B. lucida, Endlicher, prodr. fl. Norfolk. 84 (1833) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 148 M.fr. VI, 182.
FONTAINEA, Heckel, these inaugurale, Montpell. (1870).								
F. Pancheri, Heckel, these inaugurale (1870) ...	—	—	—	—	—	Q.	—	B.f. VI, 149
OMALANTHUS, Adr. de Jussieu, Euphorb. gen. 50, t. 16 (1824). (Carumbium, Wartmannia.)								
O. populifolius, Graham in Bot. Mag. 2780 (1827) ...	—	—	—	—	V.	N.S.W.	Q.	B.f. VI, 150 M.fr. I, 32.
O. stilligifolius, F. v. M., fragm. I, 32 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 150 M.fr. I, 32.
SEBASTIANA, Sprengel, Neue Entd. II, 118 (1821). (Microstachys, Elachocroton.)								
S. chamelaeae, J. Mueller in De Cand., prodr. XV, 1175 (1866) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 151
EXCAECARIA, Linné, syst. veg. ed. decim. 1288 (1759).								
E. Agallocha, Linné, syst. ed. decim. 1288 (1759) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 152 M.fr. XI, 64.
E. Dallachyana, Baillon, Adans. VI, 324 (1866) ...	—	—	—	—	—	Q.	—	B.f. VI, 153
E. parvifolia, J. Mueller in Flora, 433 (1864) ...	—	—	—	—	—	—	N.A.	B.f. VI, 153

URTICACEAE.

Ventenat, Table III, 524 (1799).

CELTIS, Tournefort, inst. 612, t. 383 (1700), from Camerarius (1586). (Solenostigma.)								
C. Philippinensis, Blanco, Fl. de Filipinas. 197 (1837) ...	—	—	—	—	Q.	N.A.	B.f. VI, 156	
C. paniculata, Planchon in Ann. sc. nat. sér. trois. X, 305 (1848) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 156 M.fr. IV, 88.
C. amblyphylla, F. v. M., fragm. IX, 76 (1875) ...	—	—	—	—	N.S.W.	—	—	M.fr. IX, 76.
ULMUS, Tournefort, inst. 601, t. 372 (1700), from Dodoens (1583).								
U. parviflora, N. Jacquin, pl. rar. hort. Schoenbr. descr. et icon. III, t. 269 (1798) ...	—	—	—	—	—	Q.	—	M.fr. VI, 101; XI, 133.
TREMA, Loureiro, Fl. Cochinch. II, 562 (1790). (Sponia.)								
T. cannabina, Loureiro, Fl. Cochinch. II, 689 (1790) ...	—	—	—	—	V.	N.S.W.	Q.	N.A. B.f. VI, 158 M.fr. VIII, 247.
APHANANTHE, Planchon in Ann. sc. nat. X, 265 (1848).								
A. Philippinensis, Planchon in Ann. sc. nat. X, 337 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 160
FICUS, Tournefort, inst. 662 (1700), from Plinius. (Urostigma, Covellia.)								
F. colosseae, F. v. M. in Benth. Fl. Austr. VI, 163 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 163
F. pilosa, Reinwardt in Blume, Bijdr. 446 (1825) ...	—	—	—	—	—	Q.	—	B.f. VI, 164 M.fr. VI, 195.
F. nesophila, F. v. M., Docum. Interc. Exhib. 26 (1866) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 164 M.fr. VI, 195.
F. Cunninghamii, Miquel in Ann. Mus. Lugd. Bot. III, 286 (1867) ...	—	—	—	—	—	Q.	—	B.f. VI, 165 M.fr. VI, 195; VIII, 246.
F. Henneana, Miquel in Ann. Mus. Lugd. Bot. III, 216 (1867) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 165
F. validinervis, F. v. M. in Benth. Fl. Austr. VI, 166 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 166
F. retusa, Linné, mantissa plant. 129 (1767) ...	—	—	—	—	—	Q.	N.A.	B.f. VI, 166 M.fr. VI, 195.
F. eugenioides, F. v. M., Docum. Interc. Exhib. 26 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VI, 166 M.fr. VI, 195.
F. Benjaminea, Linné, mantissa plant. 129 (1767) ...	—	—	—	—	—	Q.	—	B.f. VI, 167 M.fr. VI, 195.
F. Muellieri, Miquel in Ann. Mus. Lugd. Bot. III, 287 (1867) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 167 M.fr. VI, 195; VIII, 246.
F. leucotricha, Miquel in Ann. Mus. Lugd. Bot. III, 285 (1867) ...	—	—	—	—	—	N.A.	—	B.f. VI, 167 M.fr. VI, 195.
F. rubiginosa, Desfontaines, catal. pl. hort. Paris. 209 (1804) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 168 M.fr. VI, 194; VIII, 247.
F. columnaris, F. v. M. & Moore in proc. Accl. Soc. Vict. III, 71 (1874) ...	—	—	—	—	N.S.W.	—	—	M.fr. VIII, 247.
F. puberula, Cunningham in Hook. Lond. Journ. VI, 562 (1847) ...	—	—	—	—	—	N.A.	—	B.f. VI, 169
F. platypoda, Cunningham in Hook. Lond. Journ. VI, 561 (1847) ...	—	S.A.	—	—	—	Q.	N.A.	B.f. VI, 169 M.fr. VI, 194; VIII, 247.
F. dictyophleba, F. v. M. in Benth. Fl. Austr. VI, 170 (1873) ...	—	—	—	—	—	Q.	—	B.f. VI, 170

F. macrophylla, Desfontaines, catal. pl. hort. Paris. 209 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 170	M.fr. VI, 194; VIII, 247.
F. magnifolia, F. v. M., fragm. IV, 50 (1864)	...	...	...	...	—	Q.	—	B.fl. VI, 171	M.fr. IV, 50&177; VI, 196.
F. ehretioides, F. v. M. in Benth. Fl. Austr. VI, 171 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 171	
F. pumila, Linné, spec. plant. 1060 (1753)	...	...	...	...	—	N.S.W.	Q.	—	M.fr. VI, 196. [246.
F. coronulata, F. v. M. in Journ. Bot. Neerland 241 (1861)	...	...	...	...	—	—	N.A.	B.fl. VI, 172	M.fr. IV, 49; VI, 196; VIII,
F. leptoclada, Benth. Fl. Austr. VI, 172 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 172	
F. depressa, Benth. Fl. Austr. VI, 172 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 172	
F. Philippinensis, Miquel in Hooker, Lond. Journ. VII, 435 (1848)	...	...	...	...	—	Q.	—	B.fl. VI, 173	
F. mollior, F. v. M. in Benth. Fl. Austr. VI, 173 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 174	M.fr. IX, 152. [114.
F. stenocarpa, F. v. M. in Benth. Fl. Austr. VI, 174 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 174	M.fr. VI, 196; IX, 152; X,
F. scabra, G. Forster, florul. ins. Austr. prodr. 76 (1786)	...	...	...	...	—	V. N.S.W.	Q.	—	M.fr. VI, 196.
F. orbicularis, Cunningham in Hook. Lond. Journ. VII, 426 (1848)	...	S.A.	—	—	—	—	N.A.	B.fl. VI, 175	
F. aculeata, Cunningham in Hook. Lond. Journ. VII, 426 (1848)	...	—	—	—	—	—	N.A.	B.fl. VI, 175	
F. opposita, Miquel in Hook. Lond. Journ. VII, 426 (1848)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 176	M.fr. VI, 196; VIII, 246.
F. scobina, Benth. Fl. Austr. VI, 176 (1873)	...	...	...	...	—	—	N.A.	B.fl. VI, 176	
F. hispida, Linné, fil. suppl. 442 (1781)	...	...	...	...	—	Q.	N.A.	B.fl. VI, 176	
F. fasciculata, F. v. M. in Benth. Fl. Austr. VI, 177 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 177	
F. casearia, F. v. M. in Benth. Fl. Austr. VI, 177 (1873)	...	...	...	...	—	Q.	—	B.fl. VI, 177	
F. glomerata, Willdenow, spec. pl. IV, 1148 (1806)	...	...	...	...	—	Q.	N.A.	B.fl. VI, 178	M.fr. VI, 195.
F. pleurocarpa, F. v. M., fragm. VIII, 246 (1874)	...	...	...	...	—	Q.	—	—	M.fr. VIII, 246.
CUDRANIA, Trécul in Ann. des. sc. nat. VIII, 122 (1847).									
C. Javanensis, Trécul in Ann. des. sc. nat. VIII, 123 (1847)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 179	M.fr. VIII, 247.
ANTLIARIS, Leschenault in Ann. du Mus. XVI, 476 (1810).									
A. macrophylla, R. Brown in Flind. voy. II, 602, t. 5 (1814)	...	—	—	—	—	—	N.A.	B.fl. VI, 179	M.fr. VI, 194.
MALAISIA, Blanco, Fl. Filip. 789 (1837).									
M. tortuosa, Blanco, Fl. Filip. 789 (1837)	...	...	...	...	—	—	N.S.W.	Q. N.A.	B.fl. VI, 180 M.fr. VI, 193&255; VIII, [247.
PSEUDOMORUS, Bureau in Ann. des. sc. nat. sér. cinq. XI, 372 (1872).									
P. Brunoniana, Bureau in Ann. des. sc. nat. sér. cinq. XI, 372 (1872)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 181	
FATOUA, Gaudichaud in Freyc. voy. Bot. 509 (1826).									
F. pilosa, Gaudichaud in Freyc. voy. Bot. 509 (1826)	...	...	...	...	—	—	—	N.A.	B.fl. VI, 182
ELATOSTEMMA, R. & G. Forster, char. gen. 105, t. 53 (1776).									
E. reticulatum, Weddell in Ann. des. sc. nat. sér. 4, I, 188 (1854)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 183	
E. stipitatum, Weddell in Ann. des. sc. nat. sér. 4, I, 190 (1854)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 184	
PROCRIIS, Commerçon in A. L. de Jussieu, gen. 403 (1789).									
P. montana, Steudel, nomencl. bot. II, 398 (1841)	...	...	...	...	—	—	—	—	M.fr. IX, 169.
BOEHMERIA, N. J. Jacquin, stirp. Amer. hist. 246, t. 157 (1763).									
B. calophleba, Moore & Mueller, fragm. VIII, 11 (1870)	...	...	...	...	—	—	—	N.S.W.	— B.fl. VI, 184 M.fr. VIII, 11.
B. australis, Endlicher, prodr. Fl. Norfolk, 38 (1833)	...	...	...	...	—	—	—	N.S.W.	— M.fr. IX, 169.
PIPTURUS, Weddell in Ann. des. sc. nat. sér. quatr. I, 196 (1854).									
P. propinquus, Weddell in Archiv. du Mus. 447, t. 15 (1856)	...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 185	M.fr. VI, 197.
POUZOLZIA, Gaudichaud in Freyc. voy. Bot. 503 (1826). (Memorialis, Hyrtanandra.)									
P. Indica, Gaudichaud in Freyc. voy. Bot. 503 (1826)	...	...	...	...	—	Q.	N.A.	B.fl. VI, 186	M.fr. IV, 87.
P. quinquenervis, Bennett, Pl. Jav. rar. 66 (1852)	...	...	...	...	—	Q.	—	B.fl. VI, 187	
PARIETARIA, Tournefort, inst. 509, t. 289 (1700), from C. Bauhin (1623). (Freirea.)									
P. debilis, G. Forster, florul. ins. Austr. prodr. 73 (1786)	...	W.A.	S.A.	T. V.	N.S.W.	Q.	N.A.	B.fl. VI, 188	M.fr. XI, 20.
AUSTRALINA, Gaudichaud in Freyc. voy. Bot. 497 (1826).									
A. pusilla, Gaudichaud in Freyc. voy. Bot. 505 (1826)	...	...	...	...	—	—	T. V.	N.S.W.	— B.fl. VI, 189 M.fr. VIII, 247.
FLEURYA, Gaudichaud in Freyc. voy. Bot. 497 (1826).									
F. interrupta, Gaudichaud in Freyc. voy. Bot. 497 (1826)	...	...	...	...	—	—	—	Q.	— M.fr. X, 114.
URTICA, Tournefort, inst. 534, t. 308 (1700), from Plinius.									
U. incisa, Poir. Encycl. meth. suppl. IV, 223 (1817)	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q. — B.fl. VI, 190
LAPORTEA, Gaudichaud in Freyc. voy. Bot. 498 (1826).									
L. gigas, Weddell in Arch. du Mus. IX, 129, t. 3 & 4 (1856)	...	...	...	...	—	—	—	N.S.W.	Q. — B.fl. VI, 192
L. photiniphylla, Weddell in Arch. du Mus. IX, 138 (1856)	...	...	...	...	—	—	—	N.S.W.	Q. — B.fl. VI, 192
L. moroides, Weddell in Arch. du Mus. IX, 142 (1856)	...	...	...	...	—	—	—	Q.	— B.fl. VI, 192 M.fr. VIII, 248.

CUPULIFERAE.

L. Cl. Richard, Anal. de fr. 32 & 92 (1808).

FAGUS, Tournefort, inst. 584, t. 351 (1700), from Camerarius (1586).

F. Gunnii, J. Hooker, icon. plant., t. 881 (1852)	...	...	...	...	—	T.	—	—	— B.fl. VI, 210
F. Cunninghamii, Hooker, Journ. of Bot. II, 152, t. 7 (1840)	...	...	...	...	—	T. V.	—	—	— B.fl. VI, 210
F. Moorei, F. v. M., fragm. V, 109 (1866)	...	...	...	...	—	—	—	N.S.W.	— B.fl. VI, 211 M.fr. V, 109.
BALANOPS, Baillon, Hist. des pl. VI, 237 & 258 (1876).									
B. Australiana, F. v. M., fragm. X, 114 (1877)	...	...	...	...	—	—	—	Q.	— M.fr. X, 114.

CASUARINEAE.

Mirbel in Ann. du Mus. XVI, 451 (1810).

CASUARINA, Rumphius, herb. Amboin. III, 87, t. 58 (1743).

C. quadrivalvis, Labill., nov. Holl. pl. spec. II, 67, t. 218 (1806)	...	S.A.	T. V.	N.S.W.	—	—	—	— B.fl. VI, 195	M.fr. VI, 18&19; X, 62.
C. trichodon, Miquel in Lehm. pl. Preiss. I, 641 (1845)	...	W.A.	—	—	—	—	—	— B.fl. VI, 196	M.fr. VI, 17.
C. lepidophloia, F. v. M., fragm. X, 115 (1877)	...	...	...	...	—	S.A.	—	N.S.W.	— M.fr. X, 115.
C. glauca, Sieber in Sprengel, syst. III, 803 (1826)	...	...	...	...	—	S.A.	—	V. N.S.W.	Q. — B.fl. VI, 196 M.fr. VI, 18.

C. Huegeliana, Miquel in Lehm. pl. Preiss. I, 640 (1845) ...	W.A.	—	—	—	—	—	B.fl. VI, 196	M.fr. VI, 19; X, 62.
C. equisetifolia, R. & G. Forster, char. gen. 103, t. 52 (1776) ...	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 197	M.fr. VI, 17.
C. suberosa, Otto & Dietrich, allgem. Gartenzeit. 155 (1841) ...	—	—	T. V.	N.S.W.	Q.	—	B.fl. VI, 197	M.fr. VI, 20 & 234.
C. Cunninghamiana, Miquel, Revis. Casuarin. 56, t. 6 (1848) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 198	—
C. distyla, Ventenat, pl. dans le jard. de Cels. t. 62 (1800) ...	W.A.	S.A.	T. V.	N.S.W.	—	—	B.fl. VI, 198	M.fr. VI, 20.
C. inophloia, F. v. M. & Bailey in Melb. Chemist, April (1882) ...	—	—	—	—	Q.	—	—	M.fr. XII.
C. Fraseriana, Miquel, Revis. Casuarin. 50, t. 6 (1848) ...	W.A.	—	—	—	—	—	B.fl. VI, 199	M.fr. VI, 19.
C. nana, Sieber in Sprengel, syst. III, 804 (1826) ...	—	—	—	N.S.W.	—	—	B.fl. VI, 199	—
C. humilis, Otto & Dietrich, allgem. Gartenzeit. 163 (1841) ...	W.A.	—	—	—	—	—	B.fl. VI, 200	M.fr. VI, 18.
C. torulosa, Aiton, hort. Kew. III, 320 (1789) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 200	M.fr. VI, 17.
C. decussata, Benth. Fl. Austr. VI, 200 (1873) ...	W.A.	—	—	—	—	—	B.fl. VI, 200	—
C. Decaisneana, F. v. M., fragm. I, 61 (1858) ...	—	S.A.	—	—	—	N.A.	B.fl. VI, 201	M.fr. I, 61; X, 62.
C. Drummondiana, Miquel, Revis. Casuar. 26, t. 1 (1848) ...	W.A.	—	—	—	—	—	B.fl. VI, 201	—
C. microstachya, Miquel in Lehm. pl. Preiss. I, 642 (1845) ...	W.A.	—	—	—	—	—	B.fl. VI, 201	M.fr. VI, 17; X, 62.
C. acutivalvis, F. v. M., fragm. X, 61 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 61.
C. bicuspidata, Benth. Fl. Austr. VI, 202 (1873) ...	W.A.	S.A.	T.	—	—	—	B.fl. VI, 202	—
C. thuyoides, Miquel in Lehm. pl. Preiss. I, 641 (1845) ...	W.A.	—	—	—	—	—	B.fl. VI, 202	M.fr. VI, 17.
C. corniculata, F. v. M., fragm. X, 62 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 62.
C. acunaria, F. v. M., fragm. VI, 16 (1868) ...	W.A.	—	—	—	—	—	B.fl. VI, 202	M.fr. VI, 16 & 250.

PIPERACEAE.

L. Cl. Richard in Humb. Bonpl. & Kunth, Nov. gen. I, 46 (1815).

PIPER, Linné, gen. pl. 333 (1737), from C. Bauhin (1623). (Macropiper.)								
P. subpeltatum, Willdenow, spec. plant. V, 166 (1810) ...	—	—	—	—	Q.	—	B.fl. VI, 204	—
P. excelsum, G. Forster, florul. ins. Austr. prodr. 5 (1786) ...	—	—	—	N.S.W.	—	—	B.fl. VI, 204	—
P. Novae Hollandiae, Miquel in Medd. Akad. Vetensk. Amsterd. sec. ser. II, 8 (1866) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 204	—
P. Banksii, Miquel in Medd. Akad. Amst. sec. ser. II, 9 (1866) ...	—	—	—	—	Q.	—	B.fl. VI, 205	—
P. triandrum, F. v. M., fragm. V, 197 (1866) ...	—	—	—	—	Q.	—	B.fl. VI, 205	M.fr. V, 197.
P. hederaceum, Cunningham in De Cand. prodr. XVI, 366 (1869) ...	—	—	—	N.S.W.	—	—	B.fl. VI, 205	—

PEPEROMIA, Ruiz & Pavon, fl. Peruv. & Chil. prodr. 8, t. 2 (1794).

P. leptostachya, Hooker & Arnott, Bot. Beech. 96 (1841) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 206	M.fr. IX, 76.
P. reflexa, A. Dietrich, spec. plant. I, 180 (1831) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 206	—
P. Urvilleana, Ach. Richard in d'Urville, voy. Bot. 356 (1832) ...	—	—	—	—	—	—	—	M.fr. IX, 76.

PODOSTEMONACEAE.

L. Cl. Richard in Humb. Bonpl. & Kunth, Nov. gen. I, 246 (1815).

Genus and species not yet determined ...	—	—	—	—	Q.	—	—	—
--	---	---	---	---	----	---	---	---

NEPENTHACEAE.

Reichenbach, Conspectus regni vegetabilis 45 (1828).

NEPENTHES, Linné, syst. nat. 9 (1735); Linné, gen. pl. 273 (1737).

N. Kennedyana, F. v. M., fragm. V, 154 (1866) ...	—	—	—	—	Q.	—	B.fl. VI, 40	M.fr. V, 154.
---	---	---	---	---	----	---	--------------	---------------

ARISTOLOCHIEAE.

A. L. de Jussieu, in Ann. du Mus. V, 221 (1804).

ARISTOLOCHIA, Tournefort, inst. 162, t. 71 (1700), from Hippocrates, Theophrastus & Dioscorides.

A. deltantha, F. v. M., fragm. VI, 179 (1869) ...	—	—	—	—	Q.	—	B.fl. VI, 208	M.fr. VI, 179 & 255.
A. praevenosa, F. v. M., fragm. II, 166 (1861) ...	—	—	—	N.S.W.	—	—	B.fl. VI, 208	M.fr. VI, 180.
A. pubera, R. Brown, prodr. 349 (1810) ...	—	—	—	N.S.W.	Q.	—	B.fl. VI, 208	—
A. Thozetii, F. v. M., fragm. II, 167 (1861) ...	—	—	—	—	Q.	—	B.fl. VI, 208	M.fr. II, 167.
A. Indica, Linné, spec. plant. 960 (1753) ...	—	—	—	—	Q.	—	B.fl. VI, 209	—

BALONOPHOREAE.

L. Cl. Richard in Mém. du Mus. VIII, 404 & 429 (1822).

BALONOPHORA, R. & G. Forster, char. gen. 99, t. 50 (1776).

B. fungosa, R. & G. Forster, char. gen. 99, t. 50 (1776) ...	—	—	—	—	Q.	—	B.fl. VI, 232	—
--	---	---	---	---	----	---	---------------	---

VINIFERAE.

J. de St. Hilaire, Exp. fam. II, 48, t. 79 (1805).

VITIS, Tournefort, inst. 613, t. 384 (1700), e Latinis. (Cissus.)

V. Baudiniana, F. v. M., fragm. IV, 136 (1864) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl. I, 447	M.fr. V, 212; VI, 178.
V. oblonga, Benth. Fl. Austr. I, 447 (1863) ...	—	—	—	—	—	Q.	—	B.fl. I, 447	M.fr. VI, 178; IX, 125.
V. cordata, Wallich in Benth. fl. Hongk. 54 (1861) ...	—	—	—	—	—	Q.	N.A.	B.fl. I, 447	M.fr. II, 73; VI, 178; IX, 126.
V. adnata, Wallich in Wight & Arnott, prodr. 126 (1834) ...	—	—	—	—	—	—	N.A.	B.fl. I, 448	M.fr. VI, 178.
V. nitens, F. v. M., fragm. II, 73 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. I, 448	M.fr. VI, 178.
V. brachypoda, F. v. M., fr. IX, 125 (1875) ...	—	—	—	—	—	Q.	—	—	M.fr. IX, 125.
V. saponaria, Seemann, Bonplandia, 254 (1859) ...	—	—	—	—	—	Q.	—	B.fl. I, 448	M.fr. VI, 178.
V. acris, F. v. M., fragm. II, 75 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. I, 449	M.fr. VI, 178.
V. trifolia, Linné, spec. plant. 203 (1753) ...	—	—	—	—	—	Q.	N.A.	B.fl. I, 449	M.fr. II, 75; VI, 177; IX, 126.
V. clematidea, F. v. M., fragm. II, 74 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. I, 449	M.fr. VI, 177; IX, 126.
V. acetosa, F. v. M., plant. of Vict. I, 94 (1860) ...	—	—	—	—	—	—	N.A.	B.fl. I, 449	M.fr. VI, 177; IX, 126.
V. hypoglauca, F. v. M., pl. of Vict. I, 95 (1860) ...	—	—	—	—	V.	N.S.W.	Q.	B.fl. I, 450	M.fr. VI, 177.
V. sterculiifolia, F. v. M. in Benth. Fl. Austr. I, 450 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. I, 450	M.fr. IX, 126.
V. penninervis, F. v. M., fr. VI, 177 (1868) ...	—	—	—	—	—	Q.	—	—	M.fr. VI, 177.

- V. opaca, F. v. M. in Benth. Fl. Austr. I, 450 (1863) ... — — — — N.S.W. Q. — B.fl.I,450 M.fr.VI,177;IX,125.
 V. augustissima, F. v. M., fragm. I, 141 (1859) ... — — — — W.A. — — — — — B.fl.I,450 M.fr.I,244;II,180;IX,126.
 LEEA, Linné, mantiss. 17 (1767).
 L. staphylea, Roxburgh, hort. Beng. 18 (1814) ... — — — — — Q. — B.fl.I,415 M.fr.VI,178.
 L. aculeata, Blume, Bijdrag. 197 (1825) ... — — — — — Q. N.A. — M.fr.VI,178.

SAPINDACEAE.

A. L. de Jussieu, Gen. 246 (1789).

- CARDIOSPERMUM, Linné, syst. nat. (1735); Linné, gen. pl. 117 (1737).
 C. Halicacabum, Linné, sp. pl. 366 (1753) ... — — — — — Q. N.A. B.fl.I,453
 GANOPHYLLUM, Blume, Mus. Bot. Lugd. I, 230 (1850).
 G. falcatum, Blume, Mus. Bot. Lugd. I, 230 (1850) ... — — — — — Q. — — M.fr.VII,24;IX,193.
 ATALAYA, Blume, Rumphia III, 186 (1847). (Pseudatalaya, Sapindus partly.)
 A. coriacea, Radlkofer in Sitzungsber. d. Akad. zu Münch. 326 (1878) ... — — — — — N.S.W. — — —
 A. salicifolia, Blume, Rumphia III, 186 (1847) ... — — — — — — N.A. B.fl.I,463
 A. australis, Radlk. in Sitzungsber. der Akad. zu Münch. 327 (1878) ... — — — — — Q. — B.fl.I,464
 A. multiflora, Benth. Fl. Austr. I, 463 (1863) ... — — — — — N.S.W. Q. — B.fl.I,463 M.fr.IX,90.
 A. hemiglaucula, F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — — S.A. — — N.S.W. Q. N.A. B.fl.I,463 M.fr.IX,90.
 A. variifolia, F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — — — — — — N.A. B.fl.I,463 M.fr.I,45.
 DIPLOGLOTTIS, J. Hooker in H. & B. gen. pl. 395 (1862).
 D. Cunninghamii, J. Hooker in H. & B. gen. pl. 395 (1862) ... — — — — — N.S.W. Q. — B.fl.I,454 M.fr.V,145;IX,90.
 ERIOGLOSSUM, Blume, Bijdr. 229 (1829). (Pancovia partly.)
 E. edule, Blume, Bijdr. 229 (1829) ... — — — — — — N.A. B.fl.I,454 M.fr.IX,100.
 CASTANOSPORA, F. v. Mueller, fragm. IX, 92 (1875).
 C. Alphonandii, F. v. M., fragm. IX, 92 (1875) ... — — — — — Q. — — M.fr.IV,158;IX,92,197.
 ALLOPHYLUS, Linné, amoen. acad. 124 (1747). (Schmidelia.)
 A. ternatus, Loureiro, Fl. Cochinch. ed. Willd. 286 (1790) ... — — — — — Q. N.A. B.fl.I,455 M.fr.IX,90.
 CUPANIA, Plumier, Gen. 45, t. 19 (1703). (Guioa, Ratonia, Aryteria, Elattostachys, Mischocarpus, Euphoria partly.)
 C. Wadsworthii, Harpullia, F. v. M., fragm. IV, 1, t. XXVI (1863) ... — — — — — Q. — — M.fr.IV,1;IX,89,197.
 C. anacardioides, A. Richard, sert. Astrol. 33, t. 13 (1833) ... — — — — — N.S.W. Q. N.A. B.fl.I,458 M.fr.IX,91.
 C. Robertsonii, F. v. M., fragm. V, 146 (1866) ... — — — — — Q. — — M.fr.V,146;IX,94.
 C. serrata, F. v. M., fragm. III, 43 (1862) ... — — — — — N.S.W. Q. — B.fl.I,458 M.fr.III,43;IX,94.
 C. foveolata, F. v. M., fragm. IX, 95 (1875) ... — — — — — Q. — — M.fr.IX,95.
 C. tomentella, F. v. M. in Benth. Fl. Austr. I, 458 (1863) ... — — — — — Q. — B.fl.I,458
 C. pseudorhus, A. Richard, sert. Astrol. 34, t. 14 (1833) ... — — — — — N.S.W. Q. — B.fl.I,459 M.fr.IX,92.
 C. xylocarpa, Cunningham in Benth. Fl. Austr. I, 459 (1863) ... — — — — — N.S.W. Q. — B.fl.I,459 M.fr.IX,95.
 C. erythrocarpa, F. v. M., fragm. V, 7 (1865) ... — — — — — Q. — — M.fr.V,7;IX,91.
 C. Mortoniana, F. v. M., fragm. V, 177 (1866) ... — — — — — Q. — — M.fr.V,177;IX,94.
 C. lachnocarpa, F. v. M., fragm. IV, 157 (1864) ... — — — — — Q. — — M.fr.IV,156;IX,91.
 C. grandissima, F. v. M., fragm. IV, 156 (1864) ... — — — — — Q. — — M.fr.IV,156;IX,91.
 C. pyriformis, F. v. M., App. Rep. Intercol. Exhib. 25 (1867) ... — — — — — N.S.W. Q. — B.fl.I,461 M.fr.I,2;II,76;IX,90.
 C. anodonta, F. v. M., fragm. II, 76 (1860) ... — — — — — Q. — B.fl.I,461 M.fr.I,2;II,76;IX,90.
 C. stipitata, F. v. M., fragm. II, 75 (1860) ... — — — — — N.S.W. — — B.fl.I,461 M.fr.II,75;IX,91.
 C. punctulata, F. v. M., fragm. III, 12 (1862) ... — — — — — Q. — B.fl.I,458 M.fr.III,12;IX,91.
 C. exangulata, F. v. M., fragm. IV, 156 (1864) ... — — — — — Q. — — M.fr.IV,156;IX,91.
 C. O'Shanesiana, F. v. M., fragm. IX, 96 (1875) ... — — — — — Q. — — M.fr.IX,96.
 C. Daemelianana, F. v. M., fragm. IX, 96 (1875) ... — — — — — Q. — — M.fr.IX,96.
 C. Martiana, F. v. M., fragm. V, 6 (1865) ... — — — — — Q. — — M.fr.V,6;IX,94.
 C. Cordieri, F. v. M., fragm. IX, 93 (1875) ... — — — — — Q. — — M.fr.IX,93.
 C. tenax, Cunningham in Benth. Fl. Austr. I, 461 (1863) ... — — — — — N.S.W. Q. — B.fl.I,461 M.fr.IX,94.
 C. semiglaucula, F. v. M. in Benth. Fl. Austr. I, 457 (1863) ... — — — — — N.S.W. Q. — B.fl.I,457 M.fr.IX,97.
 NEPHELIUM, Linné, mantiss. 18 (1767). (Euphoria partly, Alectryon, Spanoghea.)
 N. connatum, F. v. M. in Benth. Fl. Austr. I, 465 (1863) ... — — — — — N.S.W. Q. — B.fl.I,465 M.fr.IX,98.
 N. semicinerum, F. v. M., fragm. IV, 158 (1864) ... — — — — — Q. — — M.fr.IV,158.
 N. subdentatum, F. v. M. in Benth. Fl. Austr. I, 465 (1863) ... — — — — — N.S.W. — — B.fl.I,465 M.fr.IX,99.
 N. tomentosum, F. v. M. in Transact. Vict. Inst. II, 64 (1858) ... — — — — — N.S.W. Q. — B.fl.I,466 M.fr.IX,99.
 N. coriaceum, Benth. Fl. Austr. I, 466 (1863) ... — — — — — Q. — B.fl.I,466
 N. foveolatum, F. v. M. in Benth. Fl. Austr. I, 466 (1863) ... — — — — — Q. — B.fl.I,466 M.fr.IX,99.
 N. leiocarpum, F. v. M. in Transact. phil. Inst. Vict. III, 25 (1858) ... — — — — — V. N.S.W. Q. — B.fl.I,467 M.fr.IX,98.
 N. Beckleri, Benth. Fl. Austr. I, 467 (1863) ... — — — — — N.S.W. — — B.fl.I,467
 N. divaricatum, F. v. M. in Benth. Fl. Austr. I, 467 (1863) ... — — — — — Q. — B.fl.I,467 M.fr.IX,98.
 N. microphyllum, Benth. Fl. Austr. I, 468 (1863) ... — — — — — Q. — B.fl.I,468
 N. Leichhardtii, F. v. M., App. Rep. Intercol. Exhib. 25 (1867) ... — — — — — Q. — B.fl.I,468 M.fr.IX,99.
 N. distyle, F. v. M., fragm. IX, 99 (1875) ... — — — — — Q. — B.fl.I,462 M.fr.IX,99.
 HETERODENDRON, Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818).
 H. oleaeifolium, Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818) ... — — — — — V. N.S.W. Q. N.A. B.fl.I,469 M.fr.IX,90;X,82.
 H. diversifolium, F. v. M., fragm. I, 46 (1858) ... — — — — — N.S.W. Q. — B.fl.I,469 M.fr.I,46;IX,90;X,82.
 HARPULLIA, Roxburgh, Fl. Ind. ed. Carey II, 44 (1826).
 H. alata, F. v. M., fragm. II, 103 (1861) ... — — — — — N.S.W. Q. — B.fl.I,470 M.fr.II,103;IX,89.
 H. Hillii, F. v. M. in Transact. Vict. Inst. III, 26 (1859) ... — — — — — N.S.W. Q. — B.fl.I,470
 H. Leichhardtii, F. v. M. in Benth. Fl. Austr. I, 470 (1863) ... — — — — — N.A. B.fl.I,470
 H. pendula, Planchon in Transact. Vict. Inst. III, 26 (1859) ... — — — — — N.S.W. Q. — B.fl.I,471 M.fr.IX,89&187.
 AKANIA, J. Hooker in B. & H. Gen. pl. 409 (1862).
 A. Hillii, J. Hooker in B. & H. Gen. pl. 409 (1862) ... — — — — — N.S.W. Q. — B.fl.I,471 M.fr.III,44;IX,89&197.

DIPLOPELTIS, Endlicher in Huegel, enum. 13 (1837).

D. Huegelii, Endlicher in Huegel, enum. 13 (1837)...	...	W.A.	—	—	—	—	N.A.	B.f.I, 456	M.fr. III, 12; IX, 89.
D. Stuartii, F. v. M., fragm. III, 12 (1862)...	...	—	S.A.	—	—	—	N.A.	B.f.I, 456	M.fr. III, 12 & 167; IX, 89.

DODONAEA, Linné, gen. pl. 341 (1737). (Empleurosmia).

D. triquetra, Wendland, Bot. Beobacht. p. 44 (1798)...	...	—	—	—	V.	N.S.W.	Q.	—	B.f.I, 474	M.fr. I, 73; IX, 88.
D. lanceolata, F. v. M., fragm. I, 73 (1858)...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.I, 475	M.fr. I, 73.
D. petiolaris, F. v. M., fragm. III, 13 (1862)...	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.I, 475	M.fr. IX, 89.
D. viscosa, Linné, mantissa X, altera 149 & 228 (1771)...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.I, 475	M.fr. IX, 80.
D. peduncularis, Lindley in Mitch. Trop. Austr. 361 (1848)...	...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 478	M.fr. IX, 87.
D. procumbens, F. v. M. in Transact. Vict. Inst. I, 3 (1854)...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 478	—
D. ericifolia, G. Don, gen. syst. I, 674 (1831)...	...	—	—	T.	—	—	—	—	B.f.I, 478	M.fr. IX, 87.
D. filifolia, Hooker in Mitch. Trop. Austr. 241 (1848)...	...	—	—	—	—	—	Q.	—	B.f.I, 478	M.fr. I, 71.
D. lobulata, F. v. M. in Linnaea XXV, 372 (1852)...	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.I, 479	M.fr. IX, 87.
D. ptarmicifolia, Turcz. in Bull. Soc. Mosc. XXV, part III, 165 (1852)...	...	W.A.	—	—	—	—	—	—	B.f.I, 479	M.fr. I, 97; IX, 87.
D. truncatiales, F. v. M., fragm. II, 143 (1861)...	...	—	—	—	V.	N.S.W.	Q.	—	B.f.I, 479	M.fr. IX, 89.
D. platyptera, F. v. M., fragm. I, 73 (1858)...	...	—	—	—	—	—	—	N.A.	B.f.I, 480	M.fr. IX, 87.
D. stenophylla, F. v. M., fragm. I, 72 (1858)...	...	—	—	—	—	—	Q.	—	B.f.I, 480	M.fr. IX, 87.
D. pinifolia, Miquel in Lehm. pl. Preiss. I, 227 (1844)...	...	W.A.	—	—	—	—	—	—	B.f.I, 480	M.fr. IX, 88.
D. ceratocarpa, Endlicher in Huegel, enum. 13 (1837)...	...	W.A.	—	—	—	—	—	—	B.f.I, 481	M.fr. X, 88.
D. divaricata, Benth. Fl. Austr. I, 481 (1863)...	...	W.A.	—	—	—	—	—	—	B.f.I, 481	M.fr. IX, 87.
D. triangularis, Lindley in Mitch. Trop. Austr. 219 (1848)...	...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 481	M.fr. IX, 87.
D. aptera, Miquel in Lehm. pl. Preiss. I, 225 (1844)...	...	W.A.	—	—	—	—	—	—	B.f.I, 481	M.fr. IX, 88.
D. bursarifolia, Behr & F. v. M. in Transact. Vict. Inst. I, 8 (1855)...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.I, 482	M.fr. IX, 87.
D. Baueri, Endlicher in Huegel, enum. 13 (1837)...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 482	M.fr. IX, 87.
D. trifida, F. v. M., fragm. IX, 88 (1875)...	...	W.A.	—	—	—	—	—	—	—	M.fr. IX, 88.
D. humifusa, Miquel in Lehm. pl. Preiss. I, 266 (1844)...	...	W.A.	—	—	—	—	—	—	B.f.I, 482	M.fr. IX, 88.
D. hexandra, F. v. M. in Transact. Vict. Inst. I, 117 (1855)...	...	—	S.A.	—	—	—	—	—	B.f.I, 483	—
D. ericoides, Miquel in Lehm. pl. Preiss. I, 227 (1844)...	...	W.A.	—	—	—	—	—	—	B.f.I, 483	M.fr. IX, 88.
D. polyzyga, F. v. M. in fragm. I, 74 (1858)...	...	—	—	—	—	—	—	N.A.	B.f.I, 483	M.fr. I, 74.
D. megazyga, F. v. M. in Benth. Fl. Austr. I, 483 (1863)...	...	—	—	—	—	N.S.W.	—	—	B.f.I, 483	M.fr. IX, 86.
D. macrozyga, F. v. M., fragm. IV, 135 (1864)...	...	—	—	—	—	—	Q.	—	—	M.fr. IX, 86.
D. physocarpa, F. v. M., fragm. I, 74 (1858)...	...	—	—	—	—	—	—	N.A.	B.f.I, 484	M.fr. I, 74; IX, 86.
D. vestita, Hooker in Mitch. Trop. Austr. 265 (1848)...	...	—	—	—	—	—	Q.	—	B.f.I, 484	M.fr. IX, 86.
D. pinnata, Smith in Rees' Cycl. XII (1809)...	...	—	—	—	—	N.S.W.	—	—	B.f.I, 484	—
D. oxyptera, F. v. M., fragm. I, 74 (1858)...	...	—	—	—	—	—	—	N.A.	B.f.I, 484	M.fr. I, 74.
D. humilis, Endlicher, nov. stirp. dec. 26 (1839)...	...	—	S.A.	—	—	—	—	N.A.	B.f.I, 485	M.fr. IX, 86.
D. Macrossani, F. v. M. & Scortechini in Melb. Chemist Jan. 1882	...	—	—	—	—	—	Q.	—	—	M.fr. XII.
D. microzyga, F. v. M. annual Rep. bot. Gard. 12 (1863)...	...	—	—	—	—	—	—	—	—	M.fr. IX, 86.
D. boronifolia, G. Don, gen. syst. I, 674 (1831)...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.I, 485	M.fr. IX, 86.
D. multijuga, G. Don, gen. syst. I, 674 (1831)...	...	—	—	—	—	N.S.W.	—	—	B.f.I, 485	M.fr. I, 219; II, 182; IX, 86.
D. larraeoides, Turcz. in Bull. Soc. Mosc. XXXI, 408 (1858)...	...	W.A.	—	—	—	—	—	—	B.f.I, 486	—
D. inaequifolia, Turcz. in Bull. Soc. Mosc. XXXI, 408 (1858)...	...	W.A.	—	—	—	—	—	—	B.f.I, 486	M.fr. I, 219; IX, 89.
D. adenophora, Miquel in Linnaea XVIII, 95 (1845)...	...	W.A.	—	—	—	N.S.W.	Q.	—	B.f.I, 486	M.fr. I, 98.
D. tenuifolia, Lindley in Mitch. Trop. Austr. 248 (1848)...	...	—	—	—	—	—	—	—	—	M.fr. IX, 85.
D. stenozyga, F. v. M., fragm. I, 98 (1859)...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.I, 486	M.fr. IX, 87.
D. concinna, Benth. Fl. Austr. I, 487 (1863)...	...	W.A.	—	—	—	—	—	—	B.f.I, 487	M.fr. IX, 87.

DISTICHOSTEMON, F. v. Mueller in Hook. Kew misc. IX, 306 (1857).

D. phyllopterus, F. v. M. in Hook. Kew misc. IX, 306 (1857)...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 487	M.fr. IX, 89.
--	-----	---	---	---	---	---	----	------	------------	---------------

BLEPHAROCARYA, F. v. Mueller, fragm. XI, 15 (1878).

B. involucrigera, F. v. M., fragm. XI, 16 (1878)...	...	—	—	—	—	—	Q.	—	—	M.fr. XI, 16 & 137
---	-----	---	---	---	---	---	----	---	---	--------------------

MALPIGHIACEAE.

Ventenat, Tabl. III, 131 (1799).

RYSSOPTERYS, Blume in Delessert ic. sel. III, 21, t. 35 (1837).

R. Timorensis, Blume in Delessert ic. sel. III, 21, t. 35 (1837)...	...	—	—	—	—	—	Q.	—	B.f.I, 285	—
---	-----	---	---	---	---	---	----	---	------------	---

TRISTELLATEIA, Petit-Thouars in Roemer. Collect. bot. 206 (1809).

T. Australasica, A. Richard, Sert. Astrol. 38, t. 15 (1833)...	...	—	—	—	—	—	Q.	—	B.f.I, 286	—
--	-----	---	---	---	---	---	----	---	------------	---

BUESERACEAE.

Kunth in Ann. des sc. nat. II, 346 (1824).

GARUGA, Roxburgh, Pl. of Coromandel III, 5, t. 208 (1819).

G. floribunda, Descaisne, in nouv. annal. du Mus. III, 477 (1834)...	...	—	—	—	—	—	—	N.A.	B.f.I, 377	—
--	-----	---	---	---	---	---	---	------	------------	---

CANARIUM, Rumphius, herb. Amboin. II, 145, t. 47, 49, 54 (1741). (Sonzaya.)

C. Australianum, F. v. M., fragm. III, 15 (1860)...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 378	M.fr. VII, 25.
---	-----	---	---	---	---	---	----	------	------------	----------------

ANACARDIACEAE.

R. Brown in Tuckey's Expedit. Congo 431 (1818).

RHUS, Tournefort, Inst. 611, t. 381 (1700), from Hippocr. Theophr. Diosc. & Plin. (Rhodospaera.)

R. rhodanthema, F. v. M. in Journ. pharm. soc. Vict. pl. I, 43 (1858)...	...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 489	M.fr. VII, 23.
R. panaciformis, F. v. M., fragm. VII, 22 (1869)...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 22.

EUROSCHINUS, J. Hooker in B. & H. Gen. pl. I, 422 (1862).

E. falcatus, J. Hooker in B. & H. Gen. pl. I, 422 (1862)...	...	—	—	—	—	N.S.W.	Q.	—	B.f.I, 490	M.fr. VII, 24.
---	-----	---	---	---	---	--------	----	---	------------	----------------

BUCHANANIA, Sprengel in Schrader's Journ. II, 234 (1800).

B. angustifolia, Roxburgh, Pl. Corom. III, 68, t. 262 (1819)...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 490	M.fr. VII, 23.
B. mangoides, F. v. M., fragm. VII, 23 (1869)...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 23.
B. arborescens, Blume, Mus. Bot. Lugd. Batav. I, 183 (1850)...	...	—	—	—	—	—	Q.	—	—	M.fr. VII, 23.

SEMECARPUS, Linné fl., suppl. 25 (1781).									
S. Anacardium, Linné fl., suppl. 182 (1781)	...	...	...	—	—	—	—	Q.	N.A. B.fl.I, 491 M.fr.VII, 23.
SPONDIAS, Linné, gen. pl. 365 (1737).									
S. Solandri, Benth., Fl. Austr. I, 492 (1863)	...	...	...	—	—	—	—	Q.	— B.fl.I, 492
S. pleiogyna, F. v. M., fragm. IV, 78 (1864)	...	...	...	—	—	—	—	Q.	— M.fr.III, 15.

CELASTRINEAE.

R. Brown in App. Flind. Voy. 554 (1814).

EUONYMUS, Tournefort, inst. 617, t. 388 (1700), from Theophr. & Plinius. (Evonymus.)									
E. Australianus, F. v. M., fragm. IV, 118 (1864)...	...	...	...	—	—	—	—	Q.	— M.fr.IV 118; VI, 202.
CELASTRUS, Linné, gen. pl. 59 (1737).									
C. Australis, Harvey & F. v. M. in Trans. phil. Soc. Vict. I, 41 (1854)	—	—	—	—	—	V.	N.S.W.	Q.	— B.fl.I, 398 M.fr.III, 93.
C. Muellieri, Benth., Fl. Austr. I, 399 (1863)	...	...	...	—	—	—	—	N.A.	B.fl.I, 399
C. dispermus, F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	—	—	—	—	—	—	—	Q.	— B.fl.I, 399 M.fr.VI, 203.
C. bilocularis, F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	—	—	—	—	—	—	—	Q.	— B.fl.I, 399 M.fr.VI, 204.
C. Cunninghamii, F. v. M. in Trans. phil. Inst. Vict. III, 30 (1859)	—	—	—	—	—	—	N.S.W.	Q.	N.A. B.fl.I, 399 M.fr.VI, 203.
GYMNOSPORIA, Wight & Arnott, prodr. fl. paenius. Ind. I, 159 (1834.)									
G. montana, Wight & Arnott, prodr. I, 159 (1834)	...	...	...	—	—	—	—	Q.	— B.fl.I, 400
LEUCOCARPUM, A. Richard, sert. Astrolabe 46 (1834). (Denhamia, Leucocarpon.)									
L. Oleaster, F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881)	—	—	—	—	—	—	—	Q.	— B.fl.I, 401 M.fr.VI, 203.
L. obscurum, A. Richard, sert. Astrol. 46, t. 18 (1834)	...	...	...	—	—	—	—	Q.	N.A. B.fl.I, 401 M.fr.VI, 203.
L. pittosporoides, F. v. M. in Trans. Roy. Soc. of N.S.W. 15 (1881)	—	—	—	—	—	—	N.S.W.	Q.	— B.fl.I, 402 M.fr.VI, 203.
ELAEODENDRON, J. F. Jacquin in Nov. Act. Helv. I, 36 (1787).									
E. Australe, Ventenat, Jard. Malm. t. 117 (1804)	...	...	...	—	—	—	—	N.S.W.	Q. N.A. B.fl.I, 402 M.fr.III, 61; IV, 175; VI, 204.
CARYOSPERMUM, Blume, Mus. bot. Lugd. I, 175 (1850).									
C. arborescens, F. v. M., fragm. VI, 202 (1868)	...	...	...	—	—	—	—	Q.	— M.fr.VI, 202.
SIPHONODON, Griffith in Calc. journ. of Nat. Hist. IV, 150 (1843).									
S. Australe, Benth., Fl. Austr. I, 403 (1863)	...	...	...	—	—	—	—	N.S.W.	Q. — B.fl.I, 403 M.fr.VI, 204.
HIPPOCRATEA, Linné, gen. pl. 363 (1737).									
H. obtusifolia, Roxburgh, flor. Ind. edit. Carey, I, 170 (1820)	...	...	...	—	—	—	—	N.S.W.	Q. — B.fl.I, 404
SALACIA, Linné, mantiss. II, 159 (1771).									
S. prinoides, De Candolle, prodr. I, 571 (1824)	...	...	...	—	—	—	—	Q.	— M.fr.VI, 202.

STACKHOUSIEAE.

R. Brown in App. to Flind. Voy. 555 (1814).

STACKHOUSIA, Smith in Transact. Linn. Soc. IV, 218 (1798). (Tripterococcus, Plokiostigma.)									
S. pulvinaris, F. v. M. in Transact. phil. Soc. Vict. I, 101 (1855)	—	—	T. V.	N.S.W.	—	—	—	B.fl.I, 405	M.fr.III, 88.
S. linarifolia, Cunningham in Field's N.S. Wales 365 (1825)	...	...	S.A.	T. V.	N.S.W.	Q.	—	B.fl.I, 406	M.fr.III, 87; XI, 27.
S. pubescens, A. Richard, sert. Astrol. 89, t. 33 (1834)	...	...	W.A.	—	—	—	—	B.fl.I, 407	
S. Huegelii, Endlicher in Hueg. enum. 17 (1837)	...	...	W.A.	—	—	—	—	B.fl.I, 407	
S. flava, Hooker, icon. pl. t. 269 (1840)	...	...	W.A.	S.A.	T. V.	—	—	B.fl.I, 407	
S. muricata, Lindley, Bot. Reg. XXII, 1917 (1837)	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.I, 408 M.fr.VIII, 36.
S. viminea, Smith in Rees' Cycl. XXXIII (1816)	...	...	—	—	—	V.	N.S.W.	Q.	N.A. B.fl.I, 408 M.fr.VIII, 36.
S. elata, F. v. M., fragm. III, 86 (1862)	...	...	W.A.	—	—	—	—	B.fl.I, 408	M.fr.III, 86.
S. scoparia, Benth., Fl. Austr. I, 409 (1863)	...	...	W.A.	—	—	—	—	B.fl.I, 409	
S. spathulata, Sieber in Spreng. syst. cur. post. 124 (1827)	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.fl.I, 406	M.fr.III, 86.
S. megaloptera, F. v. M., fragm. VIII, 35 (1873)	...	...	W.A.	S.A.	—	—	—	—	M.fr.X, 119.
S. Brunonis, Benth., Fl. Austr. I, 409 (1863)	...	...	W.A.	—	—	—	—	N.A.	B.fl.I, 409 M.fr.VIII, 36.
MACGREGORIA, F. v. Mueller in Caruel, Giorn. Bot. Italian. 128 (1873).									
M. racemigera, F. v. M. in Caruel, Giorn. Bot. Ital. 128 (1873)	—	—	S.A.	—	—	—	—	—	M.fr.VIII, 161; XI, 134.

FRANKENIACEAE.

A. de St. Hilaire in Bull. de la Soc. Philom. 22 (1815).

FRANKENIA, Linné, gen. pl. 92 (1737).									
F. laevis, Linné, spec. plant. 331 (1753)	...	...	W.A.	S.A.	T. V.	N.S.W.	Q.	N.A.	B.fl.I, 151
F. bracteata, Turczan. in Bull. Soc. Mosc. XXVII, 367 (1854)	...	...	W.A.	—	—	—	—	—	B.fl.I, 150
F. glomerata, Turczan. in Bull. Soc. Mosc. XXVII, 368 (1854)	...	...	W.A.	—	—	—	—	—	B.fl.I, 151
F. parvula, Turczan. in Bull. Soc. Mosc. XXVII, 368 (1854)	...	...	W.A.	—	—	—	—	—	B.fl.I, 152
F. Drummondii, Benth., Fl. Austr. I, 152 (1863)	...	...	W.A.	—	—	—	—	—	B.fl.I, 152
F. tetrapetala, Labillard., nov. Holl. pl. spec. I, 88, t. 114 (1804)	...	...	W.A.	—	—	—	—	—	B.fl.I, 152
F. punctata, Turczan. in Bull. Soc. Mosc. XXVII, 367 (1854)	...	...	W.A.	—	—	—	—	—	B.fl.I, 153

PLUMBAGINEAE.

A. L. de Jussieu, gen. plant. 92 (1789).

PLUMBAGO, Tournefort, inst. 140, t. 58 (1700).									
P. Zeilanica, Linné, sp. pl. 151 (1753)	...	...	—	—	—	—	N.S.W.	Q.	N.A. B.fl.IV, 267 M.fr.XI, 7.
STATICE, Linné, gen. pl. 88 (1737).									
S. Taxantheme, Roemer & Schultes, syst. veg. VI, 798 (1820)	...	...	—	—	—	—	T. V.	N.S.W.	Q. — B.fl.IV, 267 M.fr.XI, 7.
S. salicorniacea, F. v. M., fragm. XI, 7 (1878)	...	...	W.A.	—	—	—	—	—	N.A. — M.fr.XI, 7.
AEGIALITIS, R. Brown, prodr. 426 (1810). (Aegialinites).									
A. annulata, R. Brown, prodr. 426 (1810)	...	...	—	—	—	—	—	Q.	— B.fl.IV, 266

PORTULACEAE.

A. L. de Jussieu, Gen. 312 (1789).

PORTULACA, Tournefort, inst. 236, t. 118 (1700), from 1 Obel (1581).

P. oleracea, Linné, spec. plant. 445 (1753) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.I, 169	M.fr. X, 71.
P. napiformis, F. v. M. in Benth. Fl. Austr. I, 169 (1863) ...	—	—	—	—	—	—	Q.	N.A. B.f.I, 169	M.fr. X, 72.
P. australis, Endlicher, Atakta 7, t. 6 (1833) ...	—	—	—	—	—	—	Q.	N.A. B.f.I, 169	M.fr. X, 72.
P. filifolia, F. v. M., fragm. I, 169 (1859) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.I, 169	M.fr. III, 164; X, 72.
P. digyna, F. v. M., fragm. I, 170 (1859) ...	—	—	—	—	—	—	N.A.	B.f.I, 170	M.fr. I, 170.
P. oligosperma, F. v. M., fragm. I, 170 (1859) ...	—	—	—	—	—	—	Q.	N.A. B.f.I, 170	M.fr. III, 164; X, 72.
P. bicolor, F. v. M., fragm. I, 171 (1859) ...	—	—	—	—	—	—	Q.	N.A. B.f.I, 170	M.fr. III, 164; X, 72.
P. Armitii, F. v. M., fragm. X, 97 (1877) ...	—	—	—	—	—	—	Q.	N.A.	M.fr. X, 97.

CLAYTONIA, Gronovius in Linné, gen. pl. 339 (1737). (Calandrinia, Talinum.)

C. spergularina; Calandrinia, F. v. M., fragm. I, 175 (1859) ...	—	S.A.	—	—	—	Q.	N.A.	B.f.I, 176	M.fr. I, 175; X, 71.
C. ptychosperma; Calandrinia, F. v. M., fragm. IV, 137 (1864) ...	—	S.A.	—	—	—	Q.	—	—	M.fr. IV, 137; X, 70
C. strophilata, F. v. M., fragm. XI, 82 (1880) ...	W.A.	—	—	—	—	—	N.A.	—	M.fr. XI, 82.
C. Lehmanni, F. v. M.; Calandr., Endl. in Lehm. pl. Preiss. II, 235 (1847) ...	W.A.	—	—	—	—	—	—	B.f.I, 172	M.fr. X, 71.
C. uniflora, F. v. M.; Calandr. in Trans. Inst. Vict. III, 41 (1859) ...	—	—	—	—	—	Q.	N.A.	B.f.I, 172	M.fr. X, 71.
C. Balonnensis, F. v. M.; Calandrinia, Lindley in Mitch. Trop. Austr. 148 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.I, 172	M.fr. X, 71.
C. polyandra, F. v. M.; Calandr., Benth. Fl. Austr. I, 172 (1863) ...	W.A.	S.A.	—	—	N.S.W.	—	—	B.f.I, 172	M.fr. X, 71.
C. pleiopetala; Calandrinia, F. v. M., fragm. X, 70 (1877) ...	—	S.A.	—	—	—	Q.	—	—	M.fr. X, 70.
C. quadrivalvis; Calandrinia, F. v. M., fragm. I, 176 (1859) ...	—	—	—	—	—	—	N.A.	B.f.I, 173	M.fr. I, 176.
C. liniflora, F. v. M.; Calandr., Fenzl. in Hueg. enum. 52 (1837) ...	W.A.	—	—	—	—	—	—	B.f.I, 173	M.fr. X, 71.
C. gracilis, F. v. M.; Calandrinia, Benth. Fl. Austr. I, 173 (1863) ...	—	—	—	—	—	—	N.A.	B.f.I, 173	—
C. polypetala, F. v. M.; Calandr., Fenzl. in Hueg. enum. 51 (1837) ...	W.A.	—	—	—	—	—	—	B.f.I, 174	—
C. Pickeringi, F. v. M.; Calandr., A. Gray, Bot. Wilkes' Exped. 144 (1854) ...	—	—	—	—	N.S.W.	—	—	—	M.fr. X, 69; XI, 136.
C. volubilis, F. v. M.; Calandr., Benth. Fl. Austr. I, 174 (1863) ...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.f.I, 174	M.fr. X, 68; XI, 27.
C. calyptrata, F. v. M., fragm. III, 89 (1862) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 174	M.fr. IV, 175; X, 68.
C. pumila; Calandrinia, F. v. M., fragm. X, 68 (1877) ...	W.A.	S.A.	—	—	—	Q.	—	—	M.fr. X, 68.
C. composita, F. v. M.; Calandrinia, Nees in Lehm. pl. Preiss. I, 247 (1844) ...	W.A.	—	—	—	—	—	—	B.f.I, 175	M.fr. X, 69.
C. corrigioloides; Calandr., F. v. M., native pl. of Vict. I, 135 (1879) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.I, 175	M.fr. X, 69.
C. brevipedata; Calandr., F. v. M., fragm. X, 69 (1877) ...	W.A.	S.A.	—	—	—	—	—	—	M.fr. X, 69; XI, 136.
C. granulifera, F. v. M.; Calandr., Benth. Fl. Austr. I, 176 (1863) ...	W.A.	—	—	—	—	—	—	B.f.I, 176	—
C. poganophora; Calandrinia, F. v. M., fragm. X, 69 (1877) ...	—	—	—	—	—	Q.	—	—	M.fr. X, 69.
C. pygmaea, F. v. M., fragm. III, 89 (1862) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.I, 176	M.fr. I, 175; X, 71.
C. Australasica, J. Hooker, icon. plant. t. 293 (1840) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 177	M.fr. III, 89; X, 71.

MONTIA, Micheli, nov. pl. gen. 17, t. 13 (1729).

M. fontana, Linné, spec. plant. 87 (1753) ...	—	—	T.	—	—	—	—	B.f.I, 177	—
---	---	---	----	---	---	---	---	------------	---

CARYOPHYLLEAE.

Linné, philosoph. bot. 31 (1751).

STELLARIA, Linné, spec. plant. 421 (1753).

S. pungens, Brongniart, Voy. sur la Coquille t. 78 (1826) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 157	—
S. glauca, Withering, Arr. veg. in Great Brit. I, 420 (1776) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 158	—
S. flaccida, Hooker, Comp. Bot. Mag. I, 275 (1835) ...	—	—	T.	V.	N.S.W.	—	—	B.f.I, 158	—
S. multiflora, Hooker, Comp. Bot. Mag. I, 275 (1835) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 159	M.fr. XI, 27.

SAGINA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 118 (1737).

S. procumbens, Linné, spec. plant. 128 (1753) ...	—	—	—	V.	N.S.W.	—	—	B.f.I, 160	—
S. apetala, Linné, mantiss. 559 (1771) ...	—	S.A.	T.	V.	—	—	—	B.f.I, 160	—

COLOBANTHUS, Bartling, Reliq. Haenk. II, 13, t. 49 (1830).

C. Benthianus, Fenzl in Ann. Wien. Mus. I, 49 (1838) ...	—	—	—	V.	N.S.W.	—	—	B.f.I, 160	—
C. Billardieri, Fenzl in Ann. des Wien. Mus. I, 48 (1838) ...	—	S.A.	T.	V.	—	—	—	B.f.I, 161	—

SCLERANTHUS, Linné, gen. plant. 130 (1737). (Mniarum.)

S. pungens, R. Brown, prodr. 412 (1810) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 260	—
S. diander, R. Brown, prodr. 412 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f.V, 260	—
S. biflorus, J. Hooker, Fl. N. Zeal. I, 74 (1853) ...	—	—	T.	V.	N.S.W.	Q.	—	B.f.V, 259	—
S. mniaroides, F. v. M., pl. of Vict. I, 215 (1862) ...	—	—	—	—	V.	N.S.W.	—	B.f.V, 259	—

SAPONARIA, Linné, gen. plant. 130 (1737), from Camerarius (1586). (Gypsophila.)

S. tubulosa, F. v. M., native pl. of Vict. I, 136 (1879) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.I, 155	—
--	---	------	---	----	--------	---	---	------------	---

SPERGULARIA, Persoon, synops. pl. I, 504 (1805). (Lepigonum, Arenaria partly.)

S. rubra, Cambessèdes in St. Hilaire, Fl. Brazil. II, 179 (1829) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.I, 162	M.fr. XI, 27
--	------	------	----	----	--------	---	---	------------	--------------

DRYMARIA, Willdenow in Roemer & Schultes, syst. V, 406 (1819).

D. filiformis, Bentham, Fl. Austr. I, 162 (1863) ...	W.A.	S.A.	—	—	—	—	—	B.f.I, 162	—
--	------	------	---	---	---	---	---	------------	---

POLYCARPON, Loeffing, iter Hispanic. 7 (1758).

P. tetraphyllum, Loeffing in Linné, syst. nat. ed. X, 881 (1759) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.I, 163	—
--	------	------	----	----	--------	----	---	------------	---

POLYCARPAEA, Lamarck in Journ. d'hist. nat. II, 8, t. 25 (1792). (Aylmeyera.)

P. longiflora, F. v. M., Rep. of Babb. Exped. 8 (1858) ...	—	—	—	—	—	—	N.A.	B.f.I, 164	—
P. spirostylis, F. v. M., Rep. of Babb. Exped. 8 (1858) ...	—	—	—	—	—	—	Q.	N.A. B.f.I, 165	—
P. synandra, F. v. M., Rep. of Babb. Exped. 8 (1858) ...	—	S.A.	—	—	—	—	Q.	N.A. B.f.I, 165	—
P. violacea, Bentham, Fl. Austr. I, 165 (1863) ...	—	—	—	—	—	—	—	N.A. B.f.I, 165	—
P. Indica, Lamarck in Journ. d'hist. nat. II, 8 (1792) ...	—	S.A.	—	—	—	—	Q.	N.A. B.f.I, 166	—

P. breviflora, F. v. M., Rep. of Babb. Exped. 9 (1858) ...	...	—	—	—	—	—	Q.	N.A.	B.f.I, 166
P. spicata, Arnott in Ann. of nat. hist. III, 91 (1839) ...	...	—	—	—	—	—	—	N.A.	B.f.I, 167
P. involucrata, F. v. M., Rep. of Babb. Exped. 9 (1858) ...	...	—	—	—	—	—	—	N.A.	B.f.I, 167

AMARANTACEAE.

A. L. de Jussieu, Gen. 87 (1789).

GOMPHRENA, Linné, gen. plant. 69 (1737). (Philoxerus.)

G. canescens, R. Brown, prodr. 416 (1810) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 253	
G. flaccida, R. Brown, prodr. 416 (1810) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 254	M.fr.III, 123.
G. affinis, F. v. M. in Benth. Fl. Austr. V, 254 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 254	
G. humilis, R. Brown, prodr. 416 (1810) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 254	
G. Brownii, Moquin in De Cand. prodr. XIII, 2, 397 (1849) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 255	
G. brachystylis, F. v. M., fragm. III, 124 (1862) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 255	M.fr.III, 124.
G. leptoclada, Benth. Fl. Austr. V, 255 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 255	
G. Maitlandii, F. v. M., fragm. III, 124, t. 23 (1862) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 256	M.fr.VI, 234.
G. pusilla, Benth. Fl. Austr. V, 256 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 256	
G. tenella, Benth. Fl. Austr. V, 256 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 256	
G. conica, Sprengel, syst. I, 824 (1825) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 256	M.fr.III, 125.
G. conferta, Benth. Fl. Austr. V, 257 (1870) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 257	
G. diffusa, Sprengel, syst. I, 824 (1825) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 257	
G. parviflora, Benth. Fl. Austr. V, 257 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 257	

ALTERNANTHERA, Forskael, flor. Aegypt. Arab. 28 (1775). (Telanthera.)

A. triandra, Lamarck, encycl. meth. I, 95 (1783) ...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.V, 249	
A. nana, R. Brown, prodr. 417 (1810) ...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.V, 250	
A. angustifolia, R. Brown, prodr. 417 (1810) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 250	
A. decipiens, Benth. Fl. Austr. V, 251 (1870) ...	...	—	—	—	—	—	Q.	—	B.f.V, 251	
A. polyccephala, Benth. Fl. Austr. V, 251 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 251	
A. leptophylla, Benth. Fl. Austr. V, 251 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 251	
A. longipes, Benth. Fl. Austr. V, 252 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 252	

ACHYRANTHES, Linné, gen. plant. 34 (1737).

A. aspera, Linné, spec. plant. 204 (1753) ...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.V, 246	
A. arborescens, R. Brown, prodr. 417 (1810) ...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX, 169.

NYSSANTHES, R. Brown, prodr. 418 (1810).

N. erecta, R. Brown, prodr. 418 (1810) ...	...	—	—	—	—	N.S.W.	Q.	—	B.f.V, 247	
N. diffusa, R. Brown, prodr. 418 (1810) ...	...	—	—	—	—	N.S.W.	Q.	—	B.f.V, 247	

PTILOTUS, R. Brown, prodr. 415 (1810). (Trichinium, Psilotrichum, Goniotriche, Hemisteirus, Arthrotrichum.)

P. conicus, R. Brown, prodr. 415 (1810) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 242	
P. corymbosus, R. Brown, prodr. 415 (1810) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 243	M.fr.III, 125.
P. grandiflorus, F. v. M., fragm. I, 237 (1859) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 243	M.fr.I, 237.
P. spicatus, F. v. M. in Benth. Fl. Austr. V, 243 (1870) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 243	
P. Murrayi, F. v. M., fragm. III, 145 (1863) ...	...	—	S.A.	—	—	—	—	N.A.	B.f.V, 243	M.fr.III, 145.
P. gomphrenoides, F. v. M., fragm. VI, 233 (1868) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 244	
P. latifolius, R. Brown in App. Sturt's Exped. 25 (1849) ...	...	—	S.A.	—	—	—	—	N.A.	B.f.V, 244	M.fr.VI, 232.
P. macrotrichus, F. v. M., fragm. IV, 90 (1864) ...	...	W.A.	—	—	—	—	—	N.A.	B.f.V, 244	M.fr.VI, 232.
P. villosiflorus, F. v. M., fragm. III, 125 (1863) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 245	M.fr.III, 125.
P. humilis, F. v. M., fragm. VI, 229 (1868) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 245	M.fr.III, 161; VI, 229.
P. obovatus, F. v. M., fragm. VI, 228 (1868) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 220	M.fr.XI, 27.
P. incanus, Poiret, Encycl. méth. suppl. IV, 620 (1817) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 221	M.fr.VI, 228.
P. parviflorus, Trich., Lindley in Mitch. three Exp. II, 13 (1838) ...	...	—	—	—	—	N.S.W.	Q.	—	B.f.V, 222	
P. astrolasius, F. v. M., fragm. VI, 233 (1869) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 222	M.fr.VI, 233.
P. rotundifolius, F. v. M., fragm. VI, 230 (1868) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 223	M.fr.III, 122.
P. dissitiflorus, F. v. M., fragm. IV, 89 (1864) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 223	M.fr.IV, 89.
P. distans, Poiret, Encycl. suppl. IV, 620 (1817) ...	...	—	—	—	—	—	Q.	N.A.	B.f.V, 223	M.fr.VI, 228.
P. alopecuroides, F. v. M., fragm. VI, 227 (1868) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 224	M.fr.VI, 227.
P. nobilis, F. v. M., fragm. VI, 227 (1868) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 224	
P. macrocephalus, Poiret, Encycl. suppl. IV, 620 (1817) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 225	M.fr.VI, 228; XI, 27.
P. hemisteirus, F. v. M., fragm. IV, 90 (1864) ...	...	W.A.	S.A.	—	—	—	—	N.A.	B.f.V, 226	M.fr.VI, 231.
P. parvifolius, F. v. M., fragm. VI, 229 (1869) ...	...	—	S.A.	—	—	N.S.W.	—	—	B.f.V, 241	M.fr.VI, 229.
P. exaltatus, Nees in Lehm. pl. Preiss. I, 630 (1845) ...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 227	
P. Manglesii, F. v. M., fragm. VI, 230 (1869) ...	...	W.A.	—	—	—	—	—	N.A.	B.f.V, 228	M.fr.VI, 230.
P. Beckeri, F. v. M., fragm. VI, 233 (1869) ...	...	—	S.A.	—	—	—	—	—	B.f.V, 228	
P. gomphrenoides, Moquin in De Cand., prodr. XIII, 2, 287 (1849) ...	...	—	S.A.	—	—	—	—	—	B.f.V, 229	
P. esquamatus, Trichinium, Benth. Fl. Austr. V, 229 (1870) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 229	
P. declinatus, Nees in Lehm. pl. Preiss. I, 631 (1845) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 230	
P. erubescens, Schlechtendal in Linnaea XX, 575 (1847) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 230	M.fr.VI, 229.
P. divaricatus, F. v. M., fragm. VI, 229 (1868) ...	...	W.A.	—	—	—	—	—	N.A.	B.f.V, 231	M.fr.VI, 229.
P. helipteroides, F. v. M., fragm. VI, 231 (1868) ...	...	—	S.A.	—	—	—	—	N.A.	B.f.V, 231	M.fr.III, 122&161.
P. Stirlingii, Trichinium, Lindley in Bot. Reg. under n. 28 (1839) ...	...	W.A.	—	—	—	—	—	N.A.	B.f.V, 232	
P. laxus, Trichinium, Benth. Fl. Austr. V, 232 (1870) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 232	
P. axillaris, Trichinium, F. v. M. in Benth. Fl. Austr. V, 232 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 232	
P. striatus, Trichinium, Moquin in Benth. Fl. Austr. V, 233 (1870) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 233	
P. auriculifolius, Trichinium, Cunningham in De Cand., prodr. XIII, part II, 287 (1849) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 233	
P. sericostachyus, F. v. M., fragm. VI, 230 (1868) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 234	M.fr.VI, 230
P. roseus, Trichin., Moq. in De Cand., prodr. XIII, p. II, 284 (1849) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 234	
P. fusiformis, Poiret, Encycl. suppl. IV, 619 (1817) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 234	
P. gracilis, Poiret, Encycl. suppl. IV, 620 (1817) ...	...	—	—	—	—	—	—	N.A.	B.f.V, 235	M.fr.VI, 228.
P. Drummondii, F. v. M., fragm. VI, 229 (1868) ...	...	W.A.	—	—	—	—	—	—	B.f.V, 235	M.fr.VI, 229.

P. calostachyus, F. v. M., fragm. VI, 231 (1868) ...	—	—	—	—	—	N.A.	B.f.V. 236	M.fr. VI, 231.
P. Fraseri; Trichinium, Cunningham in De Cand., prodr. XIII, part II, 295 (1849) ...	—	—	—	—	N.S.W.	—	B.f.V. 236	
P. spathulatus, Poirer, Encycl. suppl. IV, 620 (1817) ...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.V. 236	M.fr. VI, 228.
P. pyramidatus, F. v. M., fragm. VI, 230 (1868) ...	W.A.	—	—	—	—	—	B.f.V. 237	M.fr. VI, 230.
P. holosericeus, F. v. M., fragm. VI, 229 (1868) ...	W.A.	—	—	—	—	—	B.f.V. 237	M.fr. VI, 229.
P. lanatus, Cunningham in De Cand., prodr. XIII, p. II, 281 (1849) ...	—	—	—	—	—	N.A.	B.f.V. 238	
P. leucocoma; Trichinium, Moquin in De Cand., prodr. XIII, part II, 292 (1849) ...	—	S.A.	—	—	—	—	B.f.V. 238	
P. brachyanthus; Trichin., F. v. M. in Benth. Fl. Austr. V, 639 (1870) ...	—	—	—	—	—	N.A.	B.f.V. 239	
P. arthrolasius, F. v. M., fragm. VI, 232 (1868) ...	—	—	—	—	—	N.A.	B.f.V. 239	M.fr. VI, 232.
P. Forrestii, F. v. M., pl. of N. W. Austr. 7 (1881) ...	—	—	—	—	—	N.A.	—	
P. aervoides, F. v. M., fragm. VI, 231 (1868) ...	—	—	—	—	Q.	N.A.	B.f.V. 240	M.fr. III, 123; VI, 231.
P. Roei; Trichinium, F. v. M. in Benth. Fl. Austr. V, 240 (1870) ...	W.A.	—	—	—	—	—	B.f.V. 240	
P. Hoodii, F. v. M., fragm. VIII, 232 (1874) ...	—	S.A.	—	—	—	—	—	M.fr. VIII, 232.
P. caespitosus, F. v. M., fragm. VI, 232 (1868) ...	W.A.	—	—	—	—	—	B.f.V. 240	M.fr. VI, 232.
P. helichrysoides, F. v. M., fragm. VI, 231 (1868) ...	W.A.	—	—	—	—	—	B.f.V. 241	M.fr. VI, 231.
P. psilotrichoides, F. v. M., fragm. VI, 231 (1869) ...	W.A.	—	—	—	—	—	B.f.V. 209	M.fr. XI, 97.

EUXOLUS, Rafinesque, Fl. Tell. 42 (1836).

E. Mitchellii, F. v. M. in Giles, geogr. trav. Centr. Austr. 214 (1875) ...	S.A.	—	—	N.S.W.	Q.	—	B.f.V. 214	
E. interruptus, Moquin in De Cand., prodr. XIII, part II, 275 (1849) ...	—	—	—	—	Q.	N.A.	B.f.V. 215	
E. viridis, Moquin in De Cand., prodr. XIII, part II, 273 (1849) ...	W.A.	—	—	N.S.W.	Q.	—	B.f.V. 215	
E. macrocarpus; Amarantus, Benth. Fl. Austr. V, 216 (1870) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.V. 216

AMARANTUS, Dodoens, stirp. hist. pempt. 185 (1583).

A. leptostachyus, Benth., Fl. Austr. V, 214 (1870) ...	—	—	—	—	Q.	N.A.	B.f.V. 213	
A. pallidiflorus, F. v. M., fragm. I, 140 (1859) ...	W.A.	—	—	—	Q.	N.A.	B.f.V. 214	M.fr. I, 140.

POLYCNEMON, Linné, gen. pl. ed. sec. 21 (1742). (Hemichroa.)

P. pentandrum, F. v. M., native pl. of Vict. I, 162 (1879) ...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.V. 211	M.fr. VI, 234.
P. diandrum, F. v. M., native pl. of Vict. I, 163 (1879) ...	W.A.	S.A.	—	V.	N.S.W.	N.A.	B.f.V. 211	M.fr. VI, 234.
P. mesembrianthemum, F. v. M., fragm. VIII, 38 (1873) ...	—	S.A.	—	—	—	—	—	M.fr. VIII, 38.

DEERINGIA, R. Brown, prodr. 413 (1810). (Lestibudesia partly, Celosia partly, Lagrezia.)

D. celosioides, R. Brown, prodr. 413 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V. 210	M.fr. II, 92.
D. altissima, F. v. M., fragm. II, 92 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V. 210	M.fr. VI, 251.

SALSOLACEAE.

Linné, class. pl. 507 (1738).

RHAGODIA, R. Brown, prodr. 408 (1810).

R. Billardieri, R. Brown, prodr. 408 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.V. 152	M.fr. VII, 10.
R. parabolica, R. Brown, prodr. 408 (1810) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.V. 153	M.fr. VII, 10.
R. dioica, Nees in Lehm. pl. Preiss. I, 636 (1845) ...	W.A.	—	—	—	—	—	—	B.f.V. 154	
R. Gaudichaudiana, Moquin, Chenop. monogr. enum. 11 (1840) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V. 154	M.fr. VII, 10.
R. crassifolia, R. Brown, prodr. 408 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.f.V. 154	
R. Preissii, Moquin in De Cand. prodr. XIII, part II, 49 (1849) ...	W.A.	—	—	—	—	—	—	B.f.V. 155	M.fr. VII, 10.
R. obovata, Moquin, Chenop. monogr. enum. 10 (1840) ...	W.A.	—	—	—	—	—	—	B.f.V. 155	
R. spinescens, R. Brown, prodr. 408 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V. 155	M.fr. VII, 10; XI, 20.
R. hastata, R. Brown, prodr. 408 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.V. 156	M.fr. VII, 10.
R. nutans, R. Brown, prodr. 408 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.V. 156	M.fr. VII, 10.
R. linifolia, R. Brown, prodr. 408 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V. 157	M.fr. VII, 10.

CHENOPODIUM, Tournefort, inst. 506, t. 288 (1700). (Blitum, Ambrin.)

C. nitrariaceum, F. v. M. in Benth. Fl. Austr. V, 158 (1870) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V. 158	M.fr. XI, 20&27.
C. auricomum, Lindley in Mitch. Trop. Austr. 94 (1848) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V. 159	M.fr. VII, 11.
C. triangulare, R. Brown, prodr. 407 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V. 160	M.fr. VII, 10.
C. microphyllum, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V. 161	M.fr. VII, 11.
C. rhadinostachyum, F. v. M. in Wing's S. Science Rec. May (1880) ...	—	S.A.	—	—	—	—	—	—	M.fr. XII.
C. carinatum, R. Brown, prodr. 407 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V. 162	M.fr. VII, 11.
C. cristatum, F. v. M., fragm. VII, 11 (1869) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V. 163	M.fr. XI, 27.
C. atriplicinum, F. v. M., fragm. VII, 11 (1869) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V. 163	M.fr. I, 140; XI, 27.

DYSPHANIA, R. Brown, prodr. 411 (1810).

D. plantaginella, F. v. M., fragm. I, 61 (1858) ...	—	S.A.	—	—	—	—	N.A.	B.f.V. 164	M.fr. I, 61.
D. litoralis, R. Brown, prodr. 411 (1810) ...	—	S.A.	—	—	—	Q.	N.A.	B.f.V. 164	M.fr. VII, 11.
D. myrioccephala, Benth., Fl. Austr. V, 165 (1870) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V. 165	

ATRIPLEX, Tournefort, inst. 506, t. 288 (1700), from l'Obel (1581). (Obione, Theleophyton.)

A. Moquinianum, Webb in Cand. prodr. XIII, part. II, 97 (1849) ...	W.A.	—	—	—	—	—	N.A.	B.f.V. 169	
A. stipitatum, Benth., Fl. Austr. V, 168 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V. 168	M.fr. VII, 9.
A. paludosum, R. Brown, prodr. 406 (1810) ...	W.A.	S.A.	T.	V.	—	—	—	B.f.V. 169	M.fr. VII, 9.
A. Drummondii, Moquin in Cand. prodr. XIII, part. II, 102 (1849) ...	W.A.	—	—	—	—	—	—	B.f.V. 170	
A. isatideum, Moquin, Chenop. monogr. enum. 63 (1840) ...	W.A.	—	—	—	—	—	—	B.f.V. 170	
A. nummularium, Lindley in Mitch. Trop. Austr. 64 (1848) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V. 170	M.fr. VII, 10; XI, 27.
A. cinereum, Poirer, Encycl. suppl. I, 471 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.V. 171	M.fr. VII, 9.
A. rhagodioides, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V. 172	
A. incrassatum, F. v. M., Rep. Babb. Exped. 20 (1858) ...	—	S.A.	—	—	—	—	—	B.f.V. 172	
A. vesicarium, Heward in Benth. Fl. Austr. V, 172 (1870) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V. 172	
A. hymenothecum, Moquin in Cand. prodr. XIII, part. II, 101 (1849) ...	W.A.	—	—	—	—	—	—	B.f.V. 173	M.fr. VII, 10.
A. humile, F. v. M., fragm. IV, 48 (1866) ...	—	—	—	—	—	—	N.A.	B.f.V. 174	M.fr. IV, 48.
A. fissivalve, F. v. M., fragm. IX, 123 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 123.
A. velutinellum, F. v. M., Rep. Babb. Exped. 20 (1858) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.V. 174	M.fr. VII, 9.
A. angulatum, Benth., Fl. Austr. V, 174 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V. 174	

A. semibaccatum, R. Brown, prodr. 406 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 175	M.fr. VII, 9.
A. exilifolium, F. v. M., fragm. VII, 9 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 175	M.fr. VII, 9.
A. Muelleri, Benth. Fl. Austr. V, 175 (1870) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 175	M.fr. VII, 9.
A. elachophyllum, F. v. M., fragm. VII, 8 (1870) ...	—	—	—	—	—	—	N.A.	B.f.V, 176	M.fr. VII, 8.
A. microcarpum, Benth. Fl. Austr. V, 176 (1870) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.V, 176	—
A. prostratum, R. Brown, prodr. 406 (1810) ...	—	S.A.	—	—	—	—	—	B.f.V, 176	—
A. pumilio, R. Brown, prodr. 406 (1810) ...	—	S.A.	—	—	—	—	—	B.f.V, 176	—
A. campanulatum, Benth. Fl. Austr. V, 177 (1870) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.V, 177	—
A. leptocarpum, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 178	M.fr. VII, 9; XI, 27
A. limbatum, Benth. Fl. Austr. V, 178 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 178	—
A. halimoides, Lindley in Mitch. Three Exped. I, 232 (1838) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 179	M.fr. VII, 10; XI, 2
A. holocarpum, F. v. M., Rep. Babb. Exped. 19 (1858) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 179	—
A. spongiosum, F. v. M. in Trans. phil. Inst. II, 74 (1857) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 179	—
A. crystallinum, J. Hooker in Lond. journ. of Bot. VI, 279 (1847) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.V, 180	M.fr. VII, 9.
KOCHIA, Roth in Schrader's Journ. Bot. I, 303 (1799). (Maireana, Sclerochlamys.)									
K. fimbriolata, F. v. M., fragm. IX, 75 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 75.
K. lobiflora, F. v. M. in Benth. Fl. Austr. V, 184 (1870) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.V, 184	—
K. lanosa, Lindley in Mitch. Trop. Austr. 88 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 184	M.fr. VII, 13.
K. triptera, Benth. Fl. Austr. V, 185 (1870) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V, 185	—
K. oppositifolia, F. v. M. in Transact. Vict. Inst. 134 (1855) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V, 185	M.fr. VII, 12.
K. brevifolia, R. Brown, prodr. 409 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 185	M.fr. VII, 13.
K. pyramidata, Benth. Fl. Austr. V, 186 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 186	—
K. eriantha, F. v. M., Rep. Babb. Exped. 20 (1858) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.V, 186	M.fr. VII, 13.
K. villosa, Lindley in Mitch. Trop. Austr. 91 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 186	M.fr. VII, 12.
K. planifolia, F. v. M., fragm. I, 213 (1859) ...	W.A.	—	—	—	—	—	—	B.f.V, 187	M.fr. I, 213.
K. sedifolia, F. v. M. in Transact. Vict. Inst. I, 134 (1855) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 187	M.fr. VII, 12; VIII, 37.
K. humillima, F. v. M., fragm. IX, 168 (1875) ...	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr. IX, 168.
K. microphylla, F. v. M., fragm. VIII, 148 (1874) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V, 181	M.fr. VIII, 37.
K. dichoptera, F. v. M., fragm. VIII, 37 (1873) ...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 37, 163.
K. decaptera, F. v. M., fragm. IX, 75 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 75.
K. ciliata, F. v. M., Rep. of Babb. Exped. 20 (1858) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 188	M.fr. VII, 12.
K. brachyptera, F. v. M., second gen. Rep. 15 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 189	M.fr. VII, 13.
K. stelligera, F. v. M., fragm. VII, 13 (1869) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 189	M.fr. VII, 13.
DIDYMANTHUS, Endlicher, nov. stirp. decad. 7 (1839).									
D. Roei, Endlicher, nov. stirp. decad. 8 (1839) ...	W.A.	—	—	—	—	—	—	B.f.V, 193	M.fr. VII, 11.
BABBAGIA, F. v. M., Rep. on pl. of Babb. Exped. 21 (1858).									
B. dipterocharpa, F. v. M., Rep. of Babb. Exped. 21 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.V, 192	M.fr. VII, 11.
BASSIA, Allioni in Misc. Taurin. III, 117, t. 4 (1766). (Chenolea, Sclerolaena, Anisacantha, Echinopsilon, Kentropsis, Dissocarpum, Eriochiton, Osteocarpum, Coilocarpus.)									
B. carnosa; Chenolea, Benth. Fl. Austr. V, 190 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 190	M.fr. VII, 13.
B. dallachyana; Chenolea, Benth. Fl. Austr. V, 191 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 191	—
B. tricornis; Chenolea, Benth. Fl. Austr. V, 191 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 191	—
B. eurotioides; Chenolea, F. v. M. in Benth. Fl. Austr. V, 191 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 191	M.fr. VII, 13.
B. Muelleri; Chenolea, Benth. Fl. Austr. V, 191 (1870) ...	—	—	—	—	—	—	N.A.	B.f.V, 191	—
B. sclerolaenoides; Echinopsilon, F. v. M. in Tr. phil. Inst. Vict. II, 75 (1857) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V, 192	M.fr. VII, 13.
B. uniflora; Sclerolaena, R. Brown, prodr. 410 (1810) ...	W.A.	S.A.	—	—	—	—	—	B.f.V, 194	—
B. diacantha; Anisacantha, Nees in Lehm. pl. Preiss. I, 635 (1844) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 194	M.fr. VII, 14; X, 91.
B. lanicuspis; Anisacantha, F. v. M., fragm. II, 170 (1861) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.V, 195	M.fr. II, 140, 170; VII, 14.
B. bicornis; Sclerolaena, Lindl. in Mitch. Three Exp. II, 47 (1838) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 195	M.fr. VII, 14.
B. biflora; Sclerolaena, R. Brown, prodr. 410 (1810) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 196	M.fr. VII, 11; X, 91.
B. paradoxa; Sclerolaena, R. Brown, prodr. 410 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.V, 196	M.fr. VII, 11; X, 91.
B. quinquecuspis; Anisac., F. v. M. in Tr. Vict. Inst. I, 134 (1855) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 199	M.fr. VII, 14; X, 91.
B. Drummondii; Anisacantha, Benth. Fl. Austr. V, 199 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 199	—
B. Birchii; Anisacantha, F. v. M., fragm. VIII, 163 (1874) ...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 163.
B. divaricata; Anisacantha, R. Brown, prodr. 410 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 200	M.fr. VII, 14; X, 92.
B. bicuspis; Anisac., F. v. M. in Trans. Vict. Inst. I, 133 (1855) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.V, 200	M.fr. VII, 13.
B. glabra; Kentropsis, F. v. M., fragm. I, 139 (1859) ...	—	—	—	—	—	—	N.A.	B.f.V, 200	—
B. echinopsila; Anisacantha, F. v. M., fragm. VII, 14 (1870) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.V, 201	M.fr. I, 139; X, 92.
B. enchylaenoides; Chenolea, F. v. M., fragm. X, 92 (1876) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 182	M.fr. X, 92.
B. micrantha; Enchylaena, Benth. Fl. Austr. V, 181 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 181	—
B. breviscupis; Anisacantha, F. v. M., fragm. IV, 150 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.f.V, 198	M.fr. VII, 14; VIII, 38.
B. salsuginosa; Osteocarpum, F. v. M., second gen. Rep. 15 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 197	M.fr. VII, 12; X, 92.
ENCHYLAENA, R. Brown, prodr. 408 (1810).									
E. tomentosa, R. Brown, prodr. 408 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.V, 181	M.fr. VII, 12; XI, 27.
THRELKELDIA, R. Brown, prodr. 409 (1810).									
T. diffusa, R. Brown, prodr. 410 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.V, 197	M.fr. VII, 12.
T. procerriflora, F. v. M., fragm. VIII, 38 (1873) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr. VIII, 38.
SALICORNIA, Tournefort, coroll. 51 (1703). (Halocnemum, Arthrocnemum, Tecticornia, Pachycornia).									
S. robusta, F. v. M., fragm. VI, 251 (1868) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.V, 202	M.fr. I, 139; VII, 14.
S. arbuscula, R. Brown, prodr. 411 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.V, 203	M.fr. VII, 15.
S. cinerea, F. v. M., fragm. VI, 251 (1869) ...	—	—	—	—	—	—	—	B.f.V, 203	M.fr. VII, 15.
S. leiostachya, Benth. Fl. Austr. V, 203 (1870) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.V, 203	—
S. bidens, Benth. Fl. Austr. V, 204 (1870) ...	W.A.	—	—	—	—	—	—	B.f.V, 204	—
S. tenuis, Benth. Fl. Austr. V, 204 (1870) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.V, 204	—
S. australis, Solander in G. Forster, prodr. 88 (1786) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.V, 205	M.fr. VII, 15; XI, 20, 27.
SUAEDA, Forskael, Fl. Aegypt. Arab. 69 (1779). (Chenopodium partly, Schoberia, Chenopodina.)									
S. maritima, Dumortier, Fl. Belg. 22 (1827) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.V, 206	M.fr. XI, 20.

SALSOLA, Linné, gen. pl. 67 (1737), from Cisalpinii.
S. Kali, Linné, spec. plant. 222 (1753) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,207 M.fr.XI,20,27.

FICOIDEAE.

A. L. de Jussieu, Gen. 315 (1789).

MESEMBRIANTHEMUM, Breyne, prodr. fasc. rar. pl. II, 67 (1689).

M. aequilaterale, Haworth, misc. nat. 77 (1803) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.XI,20,27.
M. australe, Solander in G. Forster, prodr. 90 (1786) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158;X,83;XI,20,27

TETRAGONIA, Linné, syst. nat. 9 (1735); Linné, gen. pl. 144 (1737). (Tetragonella.)

T. expansa, Murray in Comm. Goetting. VI, 13, t. 5 (1783) ... S.A. T. V. N.S.W. Q. — B.fl.III,325 M.fr.VII,129;X,84.
T. diptera, F. v. M., fragm. XI, 8 (1878) ... W.A. — — — — N.A. — — M.fr.XI,8.
T. implexicoma, J. Hooker, Fl. Tasman. I, 148 (1859) ... W.A. S.A. T. V. N.S.W. — N.A. B.fl.III,326 M.fr.X,84;XI,8.

AIZOON, Linné, gen. pl. 161 (1737). (Aizoon.)

A. quadrifidum, F. v. M., fragm. II, 148 (1861) ... W.A. S.A. — — N.S.W. — — B.fl.III,327 M.fr.VII,129;X,83.
A. zygophylloides, F. v. M., fragm. VII, 129 (1871) ... W.A. S.A. — — — — — M.fr.X,83.

GUNNIA, F. v. M., Rep. on pl. of Babb. Exped. 9 (1858).

G. septifraga, F. v. M., Rep. of Babb. Exped. 9 (1858) ... — S.A. — — — — — B.fl.III,327
G. Drummondii, Benth. Fl. Austr. III, 327 (1866) ... W.A. — — — — — B.fl.III,327

SESUVIUM, Linné, syst. ed. decim. 1058 (1759).

S. portulacastrum, Linné, syst. ed. dec. 1058 (1759) ... — — — — N.S.W. Q. N.A. B.fl.III,328 M.fr.VII,129;X,83.

ZALEYA, N. L. Burmann, Fl. Indic. 110, t. 31 (1768).

Z. decandra, N. L. Burmann, Fl. Ind. 110, t. 31 (1768) ... — S.A. — — N.S.W. Q. N.A. B.fl.III,329 M.fr.I,172;X,83.

TRIANTHEMA, Sauvage, Méth. fol. 127 (1751). (Ancistrostigma.)

T. turgidifolia, F. v. M., fragm. X, 83 (1876) ... — — — — — N.A. — — M.fr.X,83.
T. crystallina, Vahl, symbol. I, 32 (1790) ... — S.A. — — — — — Q. N.A. B.fl.III,330 M.fr.I,171-2.
T. glaucifolia, F. v. M., fragm. I, 172 (1859) ... — — — — — Q. — — M.fr.I,172.
T. pilosa, F. v. M., fragm. I, 174 (1859) ... — S.A. — — — — — N.A. B.fl.III,330 M.fr.X,83.
T. oxycalyptra, F. v. M., fragm. I, 173 (1859) ... — — — — — N.A. B.fl.III,330 M.fr.I,173.
T. rhynchocalyptra, F. v. M., fragm. I, 174 (1859) ... — — — — — N.A. B.fl.III,330 M.fr.I,174.
T. cypseloides, Benth. Fl. Austr. III, 331 (1866) ... — — — — — N.S.W. Q. N.A. B.fl.III,331 M.fr.X,83.

POMATOTHECA, F. v. M., fragm. X, 72 (1876).

P. humillima, F. v. M., fragm. X, 72 (1876) ... — S.A. — — N.S.W. — — — M.fr.X,72.

MACARTHURIA, Huegel, enum. pl. Nov. Holl. Austr. occ. 11 (1837).

M. apetalia, Harvey in Hook. Kew. misc. VII, 55 (1855) ... W.A. — — — — — B.fl.III,332 M.fr.XI,20.
M. australis, Huegel, enum. pl. 11 (1837) ... W.A. — — — — — B.fl.III,332
M. Neo-Cambrica, F. v. M., fragm. V, 28 (1866) ... — — — — — N.S.W. Q. — B.fl.III,332 M.fr.V,28.

MOLLUGO, Linné, gen. plant. 336 (1737). (Glinus, Trigastrotheca.)

M. Ginus, A. Richard, tentam. fl. Abyss. I, 48 (1847) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,333
M. orygioides, F. v. M. in Benth. Fl. Austr. III, 333 (1866) ... — S.A. — — N.S.W. Q. — B.fl.III,333
M. Spergula, Linné, spec. pl. ed. alt. 131 (1762) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,334
M. trigastrotheca, F. v. M., pl. of Vict. I, 201 (1860) ... — — — — — N.A. B.fl.III,334
M. Cerviana, Seringe in De Cand., prodr. I, 392 (1824) ... — S.A. — V. N.S.W. — — B.fl.III,334 M.fr.II,148.

POLYGONACEAE.

A. L. de Jussieu, gen. pl. 82 (1789), from B. de Jussieu (1759).

EMEX, Necker, elem. bot. II, 214 (1790). (Perhaps immigrated.)

E. australis, Steinheil in Ann. des sc. nat. ser. 2, IX, 195, t. 7 (1839) ... W.A. S.A. — — — — — B.fl.V,262

RUMEX, Linné, gen. pl. 105 (1737), from Plinius.

R. Brownii, Campdera, Monogr. Rum. 81 (1819) ... — S.A. T. V. N.S.W. Q. — B.fl.V,263
R. flexuosus, Solander in G. Forst., fl. ins. Austr. prodr. 90 (1786) ... — S.A. — — N.S.W. — — B.fl.V,264
R. crystallinus, Lange, ind. sem. hort. Hafn. 28 (1861) ... — S.A. — V. N.S.W. Q. N.A. B.fl.V,265 M.fr.IV,48.
R. bidens, R. Brown, prodr. 421 (1810) ... — S.A. T. V. N.S.W. Q. — B.fl.V,265

POLYGONUM, L'Obel, stirp. hist. 228 (1576), from Dioscorides and Plinius.

P. plebejum, R. Brown, prodr. 420 (1810) ... — S.A. — V. N.S.W. Q. N.A. B.fl.V,267
P. strigosum, R. Brown, prodr. 420 (1810) ... — — T. V. N.S.W. Q. — B.fl.V,268
P. prostratum, R. Brown, prodr. 419 (1810) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.V,268
P. hydropiper, Linné, spec. plant. 361 (1753) ... — S.A. — V. N.S.W. Q. — B.fl.V,269
P. minus, Hudson, Fl. Anglica I, 148 (1762) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.V,269
P. subsessile, R. Brown, prodr. 419 (1810) ... — — T. V. N.S.W. Q. — B.fl.V,269
P. barbatum, Linné, spec. plant. 362 (1753) ... — — — — N.S.W. Q. — B.fl.V,270
P. articulatum, R. Brown, prodr. 420 (1810) ... — — — — — Q. — B.fl.V,270
P. lapathifolium, Linné, spec. plant. 360 (1753) ... — S.A. T. V. N.S.W. Q. — B.fl.V,270
P. orientale, Linné, spec. plant. 362 (1753) ... — — — — N.S.W. Q. — B.fl.V,271
P. attenuatum, R. Brown, prodr. 420 (1810) ... — S.A. — — N.S.W. Q. N.A. B.fl.V,272

MUEHLENBECKIA, Meissner, gen. pl. vasc. 316 (1840).

M. adpressa, Meissner, gen. pl. comm. 227 (1843) ... W.A. S.A. T. V. N.S.W. — — B.fl.V,273 M.fr.XI,20.
M. australis, Meissner, gen. pl. comm. 227 (1843) ... — — — — N.S.W. — — M.fr.IX,169.
M. gracillima, Meissner in De Cand., prodr. XIV, 145 (1856) ... — — — — N.S.W. Q. — B.fl.V,274
M. rhyticarya, F. v. M., fragm. V, 92 (1866) ... — — — — N.S.W. Q. — B.fl.V,274
M. axillaris, J. Hooker in Lond. Journ. VI, 278 (1847) ... — — T. V. N.S.W. — — B.fl.V,275
M. polybotrya, Meissner in Lehm. pl. Preiss. I, 623 (1845) ... W.A. — — — — — B.fl.V,475 M.fr.IV,130.
M. polygonoides, F. v. M., fragm. V, 73 (1865) ... — S.A. — V. N.S.W. — — B.fl.V,275 M.fr.V,73.
M. stenophylla, F. v. M., fragm. I, 138 (1859) ... — — — — V. — — B.fl.V,275 M.fr.I,138.
M. Cunninghamii, F. v. M., fragm. V, 91 (1865) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,275 M.fr.V,95.

MONOCOCCUS, F. v. M., fragm. I, 47 (1858).									
M. echinophorus, F. v. M., fragm. I, 47 (1858)	...	...	...	—	—	—	—	N.S.W. Q.	— B.f.V, 144 M.fr.I, 47.
DIDYMOTHECA, J. Hooker in Lond. Journ. Bot. VI, 278 (1847). (Cyclothea.)									
D. thesioides, J. Hooker in Lond. Journ. V, 279 (1847)	...	...	...	W.A.	S.A.	T.	—	—	— B.f.V, 145
D. pleiococca, F. v. M., fragm. I, 202 (1859)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	— — B.f.V, 146 M.fr.I, 202.
GYROSTEMON, Desfontaines in Mém. du Mus. VI, 16, t. 6 (1820).									
G. brachystigma, F. v. M. in Benth Fl. Austr. V, 146 (1870)	...	...	...	W.A.	—	—	—	—	— B.f.V, 146
G. ramulosus, Desfontaines in Mém. du Mus. V, 17, t. 6 (1820)	...	...	...	W.A.	S.A.	—	—	—	— N.A. B.f.V, 147
CODONOCARPUS, Cunningham in Hook. Bot. Misc. I, 244 (1830). (Hymenotheca.)									
C. pyramidalis, F. v. M., plants of Vict. I, 201 (1862)	...	...	...	S.A.	—	—	—	—	— B.f.V, 148
C. australis, Cunningham in Hook. bot. misc. I, 244 (1830)	...	...	...	—	—	—	N.S.W.	Q.	— B.f.V, 148
C. cotinifolius, F. v. M., pl. of Vict. I, 200 (1862)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	— N.A. B.f.V, 149
TERSONIA, Moquin in De Cand., prodr. XIII, part sec. 40 (1849).									
T. brevipes, Moquin in De Cand., prod. XIII, part II, 40 (1849)	...	...	...	W.A.	—	—	—	—	— B.f.V, 149
T. subvolubilis, Benth, Fl. Austr. V, 150 (1870)	...	...	...	W.A.	—	—	—	—	— B.f.V, 150
CYPSELOCARPUS, F. v. M., fragm. VIII, 36 (1873).									
C. haloragoides, F. v. M., fragm. VIII, 36 (1873)	...	...	...	W.A.	—	—	—	—	— B.f.V, 198 M.fr. VIII, 36.

BOERHAAVIA, Vaillant, serm. de struct. fl. 50 (1718).									
B. diffusa, Linné, spec. fl. Zeylan. 4 (1747)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A. B.fl.V, 277
B. repanda, Willdenow, sp. pl. I, 22 (1797)	...	...	...	W.A.	S.A.	—	—	—	Q. N.A. B.fl.V, 278
PISONIA, Plumier, nov. pl. Amer. gen. 7, t. 11 (1703).									
P. aculeata, Linné, sp. pl. 1026 (1753)	...	...	...	—	—	—	—	N.S.W.	Q. — B.fl.V, 279 M.fr.VI, 197.
P. inermis, G. Forster, prodr. 75 (1786)	...	...	...	—	—	—	—	—	Q. N.A. B.fl.V, 280 M.fr.VI, 197.
P. Brunonianana, Endlicher, prodr. fl. Norf. 43 (1833)	...	...	...	—	—	—	—	N.S.W.	Q. — B.fl.V, 280 M.fr.I, 20; IV, 169.

ROUREA, Aublet, Hist. des pl. de la Guian. I, 467, t. 187 (1775).												
R. brachyandra, F. v. M., fragm. VIII, 6 (1872) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 6.	
TRICHOLOBUS, Blume, Mus. Bot. Lugd. I, 237 (1850).												
T. connaroides, F. v. M., fragm. VIII, 224 (1874) ...	...	...	—	—	—	—	—	Q.	—	—	M.fr. V, 224; X, 120.	

JANSONIA, Kippist in Gard. Chron. 19 (1847). (Cryptosema.)									
J. formosa, Kippist in Transact. Linn. Soc. XX, 384, t. 16 (1851)	W.A.	—	—	—	—	—	—	—	B.f. II, 8
BRACHYSEMA, R. Brown in Ait. hort. Kew. sec. edit. III, 10 (1811). (Leptosema, Kaleniczenkia, Burgessia.)									
B. praemorsum, Meissner in Lehm. pl. Preiss. I, 25 (1844)	...	W.A.	—	—	—	—	—	—	B.f. II, 10
B. lanceolatum, Meissner in Lehm. pl. Preiss. I, 24 (1844)	...	W.A.	—	—	—	—	—	—	B.f. II, 10
B. latifolium, R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	...	W.A.	—	—	—	—	—	—	B.f. II, 10
B. undulatum, Ker in Bot. Regist. t. 642 (1822)	...	W.A.	—	—	—	—	—	—	B.f. II, 11
B. subcordatum, Benth. Fl. Austr. II, 11 (1864)	...	W.A.	—	—	—	—	—	—	B.f. II, 11
B. bracteolusum, F. v. M., fragm. IV, 10 (1864)	...	W.A.	—	—	—	—	—	—	B.f. II, 11
B. bossiaoides, Benth. Fl. Austr. II, 12 (1864)	...	—	—	—	—	—	N.A.	B.f. II, 12	M.fr. IV, 10.
B. oxylobioides, Benth. Fl. Austr. II, 12 (1864)	...	—	—	—	—	—	Q. N.A.	B.f. II, 12	
B. uniflorum, R. Brown in Benth., Fl. Austr. II, 12 (1864)	...	—	—	—	—	—	N.A.	B.f. II, 12	
B. aphyllum, Hooker, Bot. Mag. t. 4481 (1849)	...	W.A.	—	—	—	—	—	B.f. II, 12	M.fr. I, 222.
B. macrocarpum, Benth. Fl. Austr. II, 13 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 13	
B. tomentosum, Benth. Fl. Austr. II, 13 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 13	
B. Chambersii, F. v. M. in Benth., Fl. Austr. II, 13 (1864)	...	—	S.A.	—	—	—	N.A.	B.f. II, 13	
B. daviesioides, Benth. Fl. Austr. II, 13 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 13	
OXYLOBIUM, Andrews, Bot. Reposit. t. 492 (1809). (Chorizematis subgenus, Callistachys, Podolobium.)									
O. Callistachys, Benth. Fl. Austr. II, 16 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 16	M.fr. IV, 18.
O. ellipticum, R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	...	—	—	T. V.	N.S.W.	Q.	—	B.f. II, 16	
O. alpestre, F. v. M. in Transact. phil. Soc. Vict. I, 38 (1854)	...	—	—	—	V.	N.S.W.	—	B.f. II, 17	
O. lineare, Benth. Fl. Austr. II, 17 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 17	M.fr. IV, 17.
O. carinatum, Benth. Fl. Austr. II, 18 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 18	
O. spatulatum, Benth. Fl. Austr. II, 18 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 18	
O. obtusifolium, Sweet, Fl. Austr. t. 5 (1827)	...	W.A.	—	—	—	—	—	B.f. II, 18	
O. cordifolium, Andrews, Bot. Regist. t. 492 (1809)	...	—	—	—	—	N.S.W.	—	B.f. II, 19	M.fr. IV, 17.
O. microphyllum, Benth. Fl. Austr. II, 19 (1864)	...	W.A.	—	—	—	—	—	B.f. II, 19	
O. Pulteneae, De Candolle, prodr. II, 104 (1825)	...	—	—	—	—	N.S.W.	—	B.f. II, 19	M.fr. IV, 19.
O. hamulosum, Benth. in A. Gray, Bot. Wilk. Exped. t. 379	...	—	—	—	—	N.S.W.	—	B.f. II, 20	
O. scandens, Benth. in Ann. Wien. Mus. II, 70 (1839)	...	—	—	—	—	N.S.W.	Q.	B.f. II, 20	
O. procumbens, F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)	...	—	—	—	—	V.	N.S.W.	B.f. II, 20	
O. truncipidatum, Meissner in Lehm. pl. Preiss. I, 30 (1844)	...	W.A.	—	—	—	—	—	B.f. II, 21	

O. spectabile, Endlicher, nov. stirp. dec. 2 (1839)	W.A.	—	—	—	—	—	—	B.f.II,21	
O. atropurpureum, Turczanin. in Bull. Soc. Mosc. XXVI, 250 (1853)	W.A.	—	—	—	—	—	—	B.f.II,22	
O. retusum, R. Brown in Bot. Regist. t. 913 (1825)	W.A.	—	—	—	—	—	—	B.f.II,22	
O. virgatum, Benth. in Fl. Austr. II, 22 (1864)	W.A.	—	—	—	—	—	—	B.f.II,22	
O. reticulatum, Meissner in Lehm. pl. Preiss. I, 29 (1844)	W.A.	—	—	—	—	—	—	B.f.II,23	
O. capitatum, Benth. in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.f.II,23	
O. cuneatum, Benth. in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.f.II,23	
O. acutum, Benth. in Fl. Austr. II, 24 (1864)	W.A.	—	—	—	—	—	—	B.f.II,24	
O. parviflorum, Benth. in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.f.II,24	
O. heterophyllum, Benth. in Fl. Austr. II, 25 (1864)	W.A.	—	—	—	—	—	—	B.f.II,25	
O. aciculiferum, F. v. M., fragm. I, 75, implied (1859)	—	—	—	—	—	—	Q.	B.f.II,25	M.fr.I,75.
O. trilobatum, F. v. M., annual report 23 (1858)	—	—	—	—	N.S.W.	Q.	—	B.f.II,25	M.fr.IV,19.

CHORIZEMA, Labillardière, Voy. I, 405 (1799). (Orthotropis.)

C. Dicksonii, Graham in Maund's Botanist, t. 106 (1841)	W.A.	—	—	—	—	—	—	B.f.II,27	
C. nervosum, Th. Moore in Gard. Companion (1852)	W.A.	—	—	—	—	—	—	B.f.II,27	
C. varium, Benth. in Bot. Regist. XXV, t. 49 (1839)	W.A.	—	—	—	—	—	—	B.f.II,28	
C. cordatum, Lindley, Bot. Regist. t. 10 (1838)	W.A.	—	—	—	—	—	—	B.f.II,28	
C. ilicifolium, Labillardière, Voy. I, 405, t. 21 (1799)	W.A.	—	—	—	—	—	—	B.f.II,28	
C. rhombeum, R. Brown in Ait. hort. Kew. sec. ed. III, 9 (1811)	W.A.	—	—	—	—	—	—	B.f.II,28	
C. diversifolium, A. de Candolle, Pl. rar. jard. Genève. 44, t. 8 (1836)	W.A.	—	—	—	—	—	—	B.f.II,29	
C. angustifolium, Benth. in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.f.II,29	
C. reticulatum, Meissner in Lehm. pl. Preiss. I, 34 (1844)	W.A.	—	—	—	—	—	—	B.f.II,30	
C. trigonum, Turcz. in Bull. de la Soc. Mosc. XXVI, 254 (1853)	W.A.	—	—	—	—	—	—	B.f.II,30	
C. humile, Turcz. in Bull. de la Soc. Mosc. XXVI, 254 (1853)	W.A.	—	—	—	—	—	—	B.f.II,30	
C. parviflorum, Benth. in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	N.S.W.	Q.	—	B.f.II,30	
C. cytisoides, Turczaninow in Bull. Soc. Mosc. XXVI, 256 (1853)	W.A.	—	—	—	—	—	—	B.f.II,31	
C. Henchmanni, R. Brown in Bot. Regist. t. 986 (1826)	W.A.	—	—	—	—	—	—	B.f.II,31	
C. ericifolium, Meissner in Lehm. pl. Preiss. II, 209 (1847)	W.A.	—	—	—	—	—	—	B.f.II,32	

GASTROLOBIUM, R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811).

G. pyramidale, Th. Moore in Gard. Comp. I, 81 (1852)	W.A.	—	—	—	—	—	—	B.f.II,98	
G. Lehmanni, Meissner in Lehm. pl. Preiss. I, 70 (1844)	W.A.	—	—	—	—	—	—	B.f.II,98	M.fr.X,35.
G. pulchellum, Turcz. in Bull. de la Soc. Mosc. XXVI, 274 (1853)	W.A.	—	—	—	—	—	—	B.f.II,98	
G. stipulare, Meissner in Lehm. pl. Preiss. II, 218 (1847)	W.A.	—	—	—	—	—	—	B.f.II,99	
G. Brownii, Meissner in Lehm. pl. Preiss. I, 71 (1844)	W.A.	—	—	—	—	—	—	B.f.II,99	
G. reticulatum, Benth. in Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.f.II,99	M.fr.X,35.
G. elachistum, F. v. M., fragm. IX, 67 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,67.
G. truncatum, Benth. in Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.f.II,99	
G. spathulatum, Benth. in Bot. Regist. XX, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.f.II,100	
G. plicatum, Turczanin. in Bull. Soc. Mosc. XXVI, 274 (1853)	W.A.	—	—	—	—	—	—	B.f.II,100	
G. tricuspidatum, Meissner in Lehm. pl. Preiss. I, 66 (1844)	W.A.	—	—	—	—	—	—	B.f.II,100	
G. obovatum, Benth. in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.f.II,101	M.fr.X,35.
G. epacridioides, Meissner in Lehm. pl. Preiss. I, 72 (1844)	W.A.	—	—	—	—	—	—	B.f.II,102	
G. trilobum, Benth. in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.f.II,101	
G. ilicifolium, Meissner in Lehm. pl. Preiss. I, 67 (1844)	W.A.	—	—	—	—	—	—	B.f.II,101	
G. villosum, Benth. in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.f.II,102	
G. polystachyum, Meissner in Lehm. pl. Preiss. II, 217 (1847)	W.A.	—	—	—	—	—	—	B.f.II,102	
G. ovalifolium, Henfrey in Gard. Comp. I, 41 (1852)	W.A.	—	—	—	—	—	—	B.f.II,102	
G. grandiflorum, F. v. M., fragm. III, 17 (1862)	—	—	—	—	—	Q.	N.A.	B.f.II,103	M.fr.III,17;IV,174;X,35.
G. pycnostachyum, Benth. in Fl. Austr. II, 103 (1864)	W.A.	—	—	—	—	—	—	B.f.II,103	
G. spinosum, Benth. in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.f.II,103	M.fr.X,35.
G. rotundifolium, Meissner in Lehm. pl. Preiss. II, 216 (1847)	W.A.	—	—	—	—	—	—	B.f.II,104	
G. microcarpum, Meissner in Lehm. pl. Preiss. I, 70 (1844)	W.A.	—	—	—	—	—	—	B.f.II,104	
G. oxylobioides, Benth. in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.f.II,104	
G. calycinum, Benth. in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.f.II,104	M.fr.X,35.
G. Callistachys, Meissner in Lehm. pl. Preiss. II, 216 (1846)	W.A.	—	—	—	—	—	—	B.f.II,105	
G. stenophyllum, Turczanin. in Bull. Soc. Mosc. XXVI, 275 (1853)	W.A.	—	—	—	—	—	—	B.f.II,105	
G. crassifolium, Benth. in Fl. Austr. II, 105 (1864)	W.A.	—	—	—	—	—	—	B.f.II,105	
G. parvifolium, Benth. in Fl. Austr. II, 106 (1864)	W.A.	—	—	—	—	—	—	B.f.II,106	
G. hamulosum, Meissner in Lehm. pl. Preiss. II, 218 (1847)	W.A.	—	—	—	—	—	—	B.f.II,106	
G. velutinum, Lindley & Paxton, Flower Gard. III, 76 (1853)	W.A.	—	—	—	—	—	—	B.f.II,106	M.fr.X,35.
G. bidens, Meissner in Regensb. Bot. Zeit. 29 (1855)	W.A.	—	—	—	—	—	—	B.f.II,106	
G. bilobum, R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.f.II,107	M.fr.X,35.
G. seorsifolium, F. v. M., fragm. X, 35 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,35.

ISOTROPIS, Benth. in Hueg. enum. 28 (1837).

I. striata, Benth. in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.f.II,39	M.fr.III,16.
I. Drummondii, Meissner in Lehm. pl. Preiss. I, 31 (1844)	W.A.	—	—	—	—	—	—	B.f.II,39	
I. juncea, Turczanin. in Bull. Soc. Mosc. XXVI, 251 (1853)	W.A.	—	—	—	—	—	—	B.f.II,39	M.fr.X,51.
I. canescens, F. v. M., fragm. X, 51 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,51.
I. atropurpurea, F. v. M., fragm. III, 16 (1862)	—	S.A.	—	—	—	—	N.A.	B.f.II,40	M.fr.III,16;X,51.
I. filicaulis, Benth. in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	—	—	Q.	B.f.II,40	M.fr.IV,20.
I. parviflora, Benth. in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	—	—	N.A.	B.f.II,40	
I. Wheeleri, F. v. M. in Benth. Fl. Austr. II, 40 (1864)	—	S.A.	—	—	N.S.W.	—	—	B.f.II,40	

MIRBELIA, Smith in Koenig & Sims, Ann. of Bot. I, 511 (1805). (Dichosema, Oxycladium.)

M. dilatata, R. Brown in Ait. hort. Kew. sec. ed. III, 21 (1811)	W.A.	—	—	—	—	—	—	B.f.II,33	
M. racemosa, Turczanin. in Bull. Soc. Mosc. XXVI, 282 (1853)	W.A.	—	—	—	—	—	—	B.f.II,33	M.fr.IV,18.
M. grandiflora, Aiton in Bot. Mag. t. 2771 (1827)	—	—	—	—	N.S.W.	—	—	B.f.II,34	
M. subcordata, Turczanin. in Bull. Soc. Mosc. XXVI, 282 (1853)	W.A.	—	—	—	—	—	—	B.f.II,34	
M. ovata, Meissner in Lehm. pl. Preiss. I, 77 (1844)	W.A.	—	—	—	—	—	—	B.f.II,34	

J. purpurascens, F. v. M., fragm. IV, 161 (1864)...	...	...	—	—	—	—	Q.	—	B.f.II,62	M.fr.IV,161.
J. piptomeris, Benth. in Fl. Austr. II, 62 (1864)...	...	...	W.A.	—	—	—	—	—	—	—
SPHAEROLOBUM, Smith in Koenig & Sims, Ann. of Bot. I, 509 (1805). (Roea.)										
S. linophyllum, F. v. M., fragm. I, 167 (1859) ...	...	...	W.A.	—	—	—	—	—	B.f.II,64	M.fr.I,167.
S. foliosum, F. v. M., fragm. I, 166 (1859)...	...	...	W.A.	—	—	—	—	—	B.f.II,64	M.fr.I,166.
S. gracile, Benth. in Fl. Austr. II, 64 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,64	—
S. racemosum, Benth. in Fl. Austr. II, 64 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,65	—
S. alatum, Benth. in Hueg. enum. 32 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.II,65	—
S. vimineum, Smith in Koen. & Sims, Ann. of Bot. I, 509 (1805)...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II,65
S. grandiflorum, R. Brown in Benth. Fl. Austr. II, 66 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,66	—
S. fornicatum, Benth. in Hueg. enum. 32 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.II,66	—
S. medium, R. Brown in Ait. hort. Kew. sec. ed. III, 14 (1811) ...	...	...	W.A.	—	—	—	—	—	B.f.II,66	—
S. scabriusculum, Meissner in Lehm. pl. Preiss. II, 214 (1849)...	...	...	W.A.	—	—	—	—	—	B.f.II,66	—
S. macranthum, Meissner in Lehm. pl. Preiss. II, 213 (1847) ...	...	...	W.A.	—	—	—	—	—	B.f.II,67	—
S. daviesioides, Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853) ...	...	...	W.A.	—	—	—	—	—	B.f.II,67	—
VIMINARIA, Smith, Exotic Botany 51, t. 27 (1804).										
V. denudata, Smith, Exot. Bot. 51, t. 27 (1804) ...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II,68
DAVIESIA, Smith in Transact. Linn. Soc. IV, 222 (1798).										
D. cordata, Smith in Transact. Linn. Soc. IX, 259 (1808) ...	...	...	W.A.	—	—	—	—	—	B.f.II,72	—
D. ovata, Benth. in Fl. Austr. II, 72 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,72	—
D. crenulata, Turczaninow in Bull. Soc. de Mosc. XXVI, 265 (1853) ...	...	...	W.A.	—	—	—	—	—	B.f.II,72	M.fr.IV,16.
D. oppositifolia, Endl. in den. Ann. des Wien. Mus. II, 199 (1838) ...	...	...	W.A.	—	—	—	—	—	B.f.II,73	—
D. alternifolia, Endlicher in Ann. Wien. Mus. II, 199 (1838) ...	...	...	W.A.	—	—	—	—	—	B.f.II,73	—
D. elongata, Benth. in Fl. Austr. II, 74 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,74	—
D. pedunculata, Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	B.f.II,74	—
D. arthropoda, F. v. M., fragm. VIII, 225 (1874)...	...	...	W.A.	S.A.	—	—	—	—	—	M.fr.VIII,225.
D. mollis, Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853)...	...	...	W.A.	—	—	—	—	—	B.f.II,74	—
D. concinna, R. Brown in Benth. Fl. Austr. II, 75 (1864) ...	...	...	—	—	—	—	N.S.W.	Q.	B.f.II,75	—
D. umbellulata, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	—	—	—	N.S.W.	Q.	B.f.II,75	—
D. Wyattii, Bailey in pap. Gardn. Assoc. of S. Austr. (1880) ...	...	...	—	—	—	—	V.	N.S.W.	Q.	—
D. latifolia, R. Brown in Ait. hort. Kew. sec. ed. III, 20 (1811) ...	...	...	—	—	—	—	T.	V.	N.S.W.	—
D. corymbosa, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	S.A.	—	—	V.	N.S.W.	Q.	B.f.II,76
D. horrida, Meissner in Lehm. pl. Preiss. I, 54 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,77	—
D. reclinata, Cunningham in Benth. Fl. Austr. II, 77 (1864) ...	...	...	—	—	—	—	—	N.A.	B.f.II,77	—
D. obtusifolia, F. v. M., fragm. II, 104 (1861) ...	...	...	W.A.	—	—	—	—	—	B.f.II,77	M.fr.II,104.
D. obovata, Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853) ...	...	...	W.A.	—	—	—	—	—	B.f.II,78	—
D. longifolia, Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	B.f.II,78	—
D. chordophylla, Meissner in Lehm. plant. Preiss. I, 48 (1844)...	...	...	W.A.	—	—	—	—	—	B.f.II,78	—
D. nematophylla, F. v. M. in Benth. Fl. Austr. II, 78 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,78	—
D. daphnoides, Meissner in Lehm. plant. Preiss. I, 54 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,79	—
D. nudiflora, Meissner in Lehm. plant. Preiss. I, 53 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,79	—
D. rhombifolia, Meissner in Lehm. plant. Preiss. I, 56 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,79	—
D. Drummondii, Meissner in Lehm. plant. Preiss. I, 53 (1844)...	...	...	W.A.	—	—	—	—	—	B.f.II,80	—
D. filipes, Benth. in Mitch. Trop. Austr. 363 (1848) ...	...	...	—	—	—	—	N.S.W.	Q.	B.f.II,80	—
D. cardiophylla, F. v. M., fragm. II, 105 (1861) ...	...	...	W.A.	—	—	—	—	—	B.f.II,79	M.fr.II,105.
D. squarrosa, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805) ...	...	...	—	—	—	—	N.S.W.	Q.	B.f.II,80	—
D. ulicina, Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805) ...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.II,81	—
D. acicularis, Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805) ...	...	...	—	—	—	—	N.S.W.	—	B.f.II,81	—
D. pachyphylla, F. v. M., fragm. IV, 15 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,82	M.fr.IV,15.
D. teretifolia, R. Brown in Benth. Fl. Austr. II, 82 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,82	—
D. genistifolia, Cunningham in Annal. Wien. Mus. II, 75 (1838) ...	...	...	—	S.A.	—	—	V.	N.S.W.	Q.	B.f.II,82
D. hakeoides, Meissner in Lehm. plant. Preiss. I, 47 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,82	—
D. colletioides, Meissner in Lehm. plant. Preiss. I, 48 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,83	—
D. reversifolia, F. v. M., fragm. I, 145 (1859) ...	...	...	W.A.	—	—	—	—	—	B.f.II,83	M.fr.I,145.
D. incrassata, Smith in Transact. Linn. Soc. IX, 255 (1808) ...	...	...	W.A.	S.A.	—	—	—	—	B.f.II,83	—
D. brevifolia, Lindley in Mitch. Three Exped. II, 201 (1838) ...	...	...	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.II,84
D. acanthoclona, F. v. M., fragm. X, 32 (1876) ...	...	...	W.A.	—	—	—	—	—	—	M.fr.X,32.
D. Freissii, Meissner in Lehm. plant. Preiss. I, 50 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,84	—
D. spinosissima, Meissner in Lehm. plant. Preiss. I, 51 (1844)...	...	...	W.A.	—	—	—	—	—	B.f.II,85	—
D. pachylina, Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853) ...	...	...	W.A.	—	—	—	—	—	B.f.II,85	—
D. quadrilatera, Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	B.f.II,85	—
D. striata, Turczaninow in Bull. Soc. Mosc. XXVI, 264 (1853) ...	...	...	W.A.	—	—	—	—	—	B.f.II,85	M.fr.II,105.
D. polyphylla, Benth. in Bot. Regist. XXV, App. XIV (1839) ...	...	...	W.A.	—	—	—	—	—	B.f.II,86	—
D. microphylla, Benth. in Fl. Austr. II, 86 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,86	—
D. flexuosa, Benth. in Hueg. enum. 32 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.II,86	—
D. pectinata, Lindley in Mitch. Three Exped. II, 151 (1838) ...	...	...	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.II,87
D. trigonophylla, Meissner in Lehm. plant. Preiss. II, 213 (1847) ...	...	...	W.A.	—	—	—	—	—	B.f.II,87	—
D. epiphylla, Meissner in Regensb. Bot. Zeit. 27 (1855) ...	...	...	W.A.	—	—	—	—	—	B.f.II,87	—
D. euphorbioides, Benth. in Fl. Austr. II, 88 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,88	—
D. divaricata, Benth. in Hueg. enum. 31 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.II,88	—
D. paniculata, Benth. in Hueg. enum. 31 (1837) ...	...	...	W.A.	—	—	—	—	—	B.f.II,88	—
D. aphylla, F. v. M. in Benth. Fl. Austr. II, 88 (1864) ...	...	...	W.A.	—	—	—	—	—	B.f.II,88	—
D. juncea, Smith in Transact. Linn. Soc. IX, 260 (1808) ...	...	...	W.A.	—	—	—	—	—	B.f.II,88	—
D. alata, Smith in Transact. Linn. Soc. IX, 259 (1808) ...	...	...	—	—	—	—	N.S.W.	—	B.f.II,89	—
D. anceps, Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853)...	...	...	W.A.	—	—	—	—	—	B.f.II,89	—
AOTUS, Smith in Koenig & Sims, Ann. of Bot. I, 504 (1805).										
A. gracillima, Meissner in Lehm. pl. Preiss. I, 59 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.II,90	—
A. villosa, Smith in Koen. & Sims, Ann. of Bot. I, 504 (1805) ...	...	...	—	—	—	—	T.	V.	N.S.W.	Q.
A. mollis, Benth. in Mitch. Trop. Austr. 236 (1848) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.II,91

A. Preissii, Meissner in Lehm. plant. Preiss. II, 214 (1846) ...	W.A.	—	—	—	—	—	—	B.f.II, 91	
A. physicoides, F. v. M. in Benth. Fl. Austr. II, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 92	
A. lanigera, Cunningham in Annal. des Wien. Mus. II, 78 (1838) ...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 92	
A. genistoides, Turczaninow in Bull. Soc. Mosc. XXVI, 268 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 92	
A. carinata, Meissner in Lehm. plant. Preiss. II, 215 (1849) ...	W.A.	—	—	—	—	—	—	B.f.II, 93	
A. passerinoides, Meissner in Lehm. plant. Preiss. I, 61 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 93	
A. Tietkensis, F. v. M., fragm. X, 33 (1876) ...	W.A.	—	—	—	—	—	—	—	M.fr.X, 33.
A. cordifolia, Benth in Hueg. enum. 33 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 93	
PHYLLOTA, De Candolle, prodr. II, 113 (1825). (Urodon.)									
P. barbata, Benth in Hueg. enum. 33 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 94	
P. gracilis, Turczaninow in Bull. Soc. Mosc. XXVI, 267 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 94	
P. sturtii, Benth in Fl. Austr. II, 95 (1864) ...	—	S.A.	—	—	—	—	—	B.f.II, 95	
P. phyllicifolia, Benth in Ann. des Wien. Mus. II, 77 (1838) ...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 95	M.fr.X, 33.
P. humifusa, Benth in Fl. Austr. II, 95 (1864) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 95	
P. diffusa, F. v. M., fragm. I, 8 (1858) ...	—	—	—	—	T.	—	—	B.f.II, 119	M.fr.I, 8.
P. pleurandroides, F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854) ...	—	S.A.	—	—	V.	—	—	B.f.II, 96	M.fr.X, 33.
P. Luehmanni, F. v. M., fragm. X, 33 (1876) ...	W.A.	—	—	—	—	—	—	—	M.fr.X, 33.
P. Urodon, F. v. M. in papers of the Roy. Soc. of Tasm. 120 (1877) ...	W.A.	—	—	—	—	—	—	B.f.II, 124	
PULTENAEA, Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793). (Euchilus, Spadostyles, Bartlingia.)									
P. daphnoides, Wendland, Bot. Beob. 49 (1798) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 112	
P. stricta, Sims, Bot. Mag. 1588 (1813) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 112	
P. retusa, Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.II, 113	
P. Benthamii, F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854) ...	—	—	—	—	V.	—	—	B.f.II, 113	
P. pycnocephala, F. v. M. in Benth. Fl. Austr. II, 114 (1864) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 114	
P. myrtoides, Cunningham in Annal. des Wien. Mus. II, 81 (1838) ...	—	—	—	—	—	—	Q.	B.f.II, 114	
P. mucronata, F. v. M., fragm. I, 8 (1858) ...	—	S.A.	—	—	V.	—	—	B.f.II, 114	M.fr.I, 8.
P. polifolia, Cunningham in Field's N. S. Wales 346 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 115	
P. petiolaris, Cunningham in Ann. des Wien. Mus. II, 82 (1838) ...	—	—	—	—	—	—	Q.	B.f.II, 115	
P. paleacea, Willdenow, spec. plant. II, 506 (1799) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 115	
P. Gunni, Benth in Ann. des Wien. Mus. II, 82 (1839) ...	—	—	—	—	T.	V.	N.S.W.	B.f.II, 116	
P. scabra, R. Brown in Ait. hort. Kew. sec. ed. III, 18 (1811) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 116	
P. Hartmanni, F. v. M., fragm. VIII, 166 (1874) ...	—	—	—	—	—	N.S.W.	Q.	—	M.fr.VIII, 166.
P. microphylla, Sieber in De Cand. prodr. II, 11 (1825) ...	—	—	—	—	—	N.S.W.	Q.	B.f.II, 117	
P. Skinneri, F. v. M., fragm. VIII, 166 (1874) ...	W.A.	—	—	—	—	—	—	—	M.fr.VIII, 166.
P. Drummondii, Meissner in Lehm. plant. Preiss. II, 219 (1847) ...	W.A.	—	—	—	—	—	—	B.f.II, 117	
P. pinifolia, Meissner in Lehm. plant. Preiss. II, 220 (1847) ...	W.A.	—	—	—	—	—	—	B.f.II, 118	
P. pedunculata, Hooker, Bot. Magaz. t. 2859 (1828) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 118	
P. conferta, Benth in Fl. Austr. II, 118 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 118	
P. aciphylla, Benth in Hueg. enum. 35 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 119	
P. ochreate, Meissner in Lehm. plant. Preiss. I, 73 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 120	
P. aspalathoides, Meissner in Lehm. plant. Preiss. I, 73 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 120	
P. obcordata, Benth in Fl. Austr. II, 120 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 120	
P. rotundifolia, Benth in Fl. Austr. II, 121 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 121	M.fr.I, 145.
P. calycina, Benth in Fl. Austr. II, 121 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 121	
P. spinulosa, Benth in Fl. Austr. II, 121 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 121	
P. tenella, Benth in Fl. Austr. II, 122 (1864) ...	—	—	—	—	V.	—	—	B.f.II, 122	
P. ternata, F. v. M., fragm. I, 8 (1858) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.II, 122	M.fr.I, 8; IV, 10.
P. styphelioides, Cunningham in G. Don, gen. syst. II, 124 (1832) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 122	
P. altissima, F. v. M. in Benth. Fl. Austr. II, 123 (1864) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 123	
P. obovata, Benth in Fl. Austr. II, 123 (1864) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 123	
P. incurvata, Cunningham in Field's N. S. Wales 346 (1825) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 124	
P. subumbellata, Hooker, Bot. Mag. t. 3254 (1833) ...	—	—	—	—	T.	V.	N.S.W.	B.f.II, 124	
P. stipularis, Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 125	
P. glabra, Benth in Fl. Austr. II, 124 (1864) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 125	
P. dentata, Labillardière, Nov. Holl. pl. spec. I, 103, t. 131 (1804) ...	—	—	—	—	T.	V.	N.S.W.	B.f.II, 125	
P. aristata, Sieber in De Cand. prodr. II, 112 (1825) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 126	
P. plumosa, Sieber in De Cand. prodr. II, 111 (1825) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 126	
P. viscosa, R. Brown in Benth. Fl. Austr. II, 127 (1864) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 127	
P. echinula, Sieber in De Cand. prodr. II, 112 (1825) ...	—	—	—	—	—	N.S.W.	Q.	B.f.II, 127	
P. hibernioides, J. Hooker, Fl. Tasm. I, 89 (1860) ...	—	—	—	—	T.	V.	N.S.W.	B.f.II, 127	
P. rosea, F. v. M., fragm. II, 15 (1860) ...	—	—	—	—	V.	—	—	B.f.II, 128	M.fr.II, 15.
P. mollis, Lindley in Mitch. Three Exped. II, 260 (1838) ...	—	S.A.	—	—	V.	—	—	B.f.II, 128	
P. strobilifera, Meissner in Lehm. pl. Preiss. I, 75 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 128	
P. ericifolia, Benth in Bot. Regist. XXV, App. XIII (1839) ...	W.A.	—	—	—	—	—	—	B.f.II, 129	
P. verruculosa, Turcz. in Bull. Soc. Mosc. XXVI, 278 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 129	
P. empetrifolia, Meissner in Lehm. plant. Preiss. I, 76 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 129	
P. adunca, Turczaninow in Bull. Soc. Mosc. XXVI, 279 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 130	
P. neurocalyx, Turcz. in Bull. Soc. Mosc. XXVI, 281 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 130	
P. rigida, R. Brown in Benth. Fl. Austr. II, 130 (1864) ...	—	S.A.	—	—	—	—	—	B.f.II, 130	
P. juniperina, Labillardière, Nov. Holl. pl. spec. I, 102, t. 130 (1804) ...	—	—	—	—	T.	V.	—	B.f.II, 131	
P. acerosa, R. Brown in Benth. Fl. Austr. II, 131 (1864) ...	—	S.A.	—	—	—	—	—	B.f.II, 131	
P. humilis, Benth in J. Hook. Fl. Tasm. I, 91 (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 131	
P. parviflora, Sieber in De Cand. prodr. II, 111 (1825) ...	—	—	—	—	V.	N.S.W.	—	B.f.II, 132	
P. setulosa, Benth in Fl. Austr. II, 132 (1864) ...	—	—	—	—	—	—	Q.	B.f.II, 132	
P. vestita, R. Brown in Ait. hort. Kew. sec. ed. III, 19 (1811) ...	W.A.	S.A.	—	—	—	—	—	B.f.II, 132	
P. procumbens, Cunningham in Field's N. S. Wales 347 (1838) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 133	
P. hispidula, R. Brown in Benth. Fl. Austr. II, 133 (1864) ...	—	—	—	—	—	N.S.W.	—	B.f.II, 133	
P. laxiflora, Benth in Fl. Austr. I, 133 (1864) ...	—	S.A.	—	—	V.	—	—	B.f.II, 133	
P. largiflorens, F. v. M. in Benth. Fl. Austr. II, 134 (1864) ...	—	S.A.	—	—	V.	—	—	B.f.II, 134	

<i>P. villosa</i> , Willdenow, spec. plant. II, 507 (1799) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.II, 134	
<i>P. foliolosa</i> , Cunningham in Ann. des Wien. Mus. II, 83 (1838) ...	—	—	—	V.	N.S.W.	—	—	B.f.II, 135	
<i>P. flexilis</i> , Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 135	
<i>P. euchila</i> , De Candolle, prodr. II, 112 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 135	
<i>P. selaginoides</i> , J. Hooker, Fl. Tasm. I, 87 (1860) ...	—	—	—	T.	—	—	—	B.f.II, 136	
<i>P. densifolia</i> , F. v. M. in Transact. Vict. Inst. 119 (1854) ...	—	S.A.	—	—	—	—	—	B.f.II, 136	
<i>P. elliptica</i> , Smith in Transact. Linn. Soc. IX, 246 (1808) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 136	
<i>P. subpicata</i> , Benth. Fl. Austr. II, 137 (1864) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 137	
<i>P. villifera</i> , Sieber in De Cand. prodr. II, 111 (1825) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 137	
<i>P. involucreta</i> , Benth. Fl. Austr. II, 138 (1864) ...	—	S.A.	—	—	—	—	—	B.f.II, 138	
<i>P. Muelleri</i> , Benth. Fl. Austr. II, 138 (1864) ...	—	—	—	V.	—	—	—	B.f.II, 138	
<i>P. prostrata</i> , Benth. in J. Hook. Fl. Tasm. I, 89 (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 138	
<i>P. canaliculata</i> , F. v. M. in Transact. Vict. Inst. 119 (1854) ...	—	S.A.	—	V.	—	—	—	B.f.II, 139	
<i>P. fasciculata</i> , Benth. in Ann. des Wien. Mus. II, 82 (1838) ...	—	—	T.	V.	N.S.W.	—	—	B.f.II, 139	
<i>P. tenuifolia</i> , R. Brown in Bot. Mag. t. 2086 (1819) ...	W.A.	S.A.	T.	V.	—	—	—	B.f.II, 139	M.fr.I, 9.
LATROBEA , Meissner in Lehm. plant. Preiss. II, 219 (1847). (Leptocytisus.)									
<i>L. pungens</i> , Benth. Fl. Austr. II, 140 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 140	M.fr.II, 106.
<i>L. genistoides</i> , Meissner in Lehm. plant. Preiss. II, 219 (1847) ...	W.A.	—	—	—	—	—	—	B.f.II, 141	
<i>L. Brunonis</i> , Meissner in Lehm. plant. Preiss. II, 219 (1847) ...	W.A.	—	—	—	—	—	—	B.f.II, 141	
<i>L. tenella</i> , Benth. Fl. Austr. II, 141 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 141	
<i>L. hirtella</i> , Benth. Fl. Austr. II, 142 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 142	
<i>L. diosmifolia</i> , Benth. Fl. Austr. II, 142 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 142	
EUTAXIA , R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811). (Sclerotheramnus.)									
<i>E. cuneata</i> , Meissner in Lehm. plant. Preiss. I, 65 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 143	
<i>E. myrtifolia</i> , R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811) ...	W.A.	—	—	—	—	—	—	B.f.II, 144	
<i>E. epacridoides</i> , Meissner in Lehm. plant. Preiss. I, 64 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 144	
<i>E. virgata</i> , Benth. in Hueg. enum. 34 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 144	
<i>E. densifolia</i> , Turczaninow in Bull. Soc. Mosc. XXVI, 271 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 145	
<i>E. dillwynioides</i> , Meissner in Lehm. plant. Preiss. I, 63 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 145	
<i>E. parvifolia</i> , Benth. in Hueg. enum. 34 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 145	
<i>E. empetrifolia</i> , Schlechtendal, Linnaea XX, 667 (1847) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.II, 145	M.fr.I, 7.
DILLWYNIA , Smith in Koenig & Sims, Ann. of Bot. I, 510 (1805).									
<i>D. hispida</i> , Lindley in Mitch. Three Exped. II, 251 (1838) ...	—	S.A.	—	V.	—	—	—	B.f.II, 147	
<i>D. ericifolia</i> , Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 147	
<i>D. floribunda</i> , Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 149	
<i>D. Preissii</i> , Benth. Fl. Austr. II, 149 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 149	
<i>D. juniperoides</i> , Meissner in Lehm. pl. Preiss. I, 62 (1844) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 150	
<i>D. bruniperina</i> , Sieber in Hueg. enum. pl. N. Holl. occ. 33 (1837) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.II, 150	
<i>D. pungens</i> , Mackay in den Annal. des Wien. Mus. II, 79 (1838) ...	W.A.	—	—	—	—	—	—	B.f.II, 150	
<i>D. cinerascens</i> , R. Brown in Bot. Mag. t. 2247 (1821) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 151	
<i>D. divaricata</i> , Benth. Fl. Austr. II, 151 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 151	
<i>D. patula</i> , F. v. M., fragm. IV, 16 (1864) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.II, 151	M.fr.IV, 16.
EUCHILOPSIS , F. v. M. in Melbourne Chemist, June (1882).									
<i>E. linearis</i> , F. v. M. in Melb. Chem., June (1882) ...	W.A.	—	—	—	—	—	—	B.f.II, 67	
PLATYLOBIUM , Smith in Transact. Linn. Soc. II, 350 (1794). (Cheilococca.)									
<i>P. formosum</i> , Smith in Transact. Linn. Soc. II, 350 (1794) ...	—	—	T.	V.	N.S.W.	Q.	—	B.f.II, 153	M.fr.IX, 157.
<i>P. obtusangulum</i> , Hooker, Bot. Mag. t. 3258 (1833) ...	—	S.A.	T.	V.	—	—	—	B.f.II, 153	
<i>P. triangulare</i> , R. Brown in Ait. hort. Kew. sec. ed. III, 266 (1811) ...	—	S.A.	T.	V.	—	—	—	B.f.II, 152	M.fr.IX, 158.
BOSSIAEA , Ventenat, Jardin de Cels I, 7, t. 7 (1800). (Lalage, Scottia.)									
<i>B. dentata</i> , Benth. Fl. Austr. II, 156 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 156	
<i>B. Aquifolium</i> , Benth. Fl. Austr. II, 157 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 157	
<i>B. strigillosa</i> , Benth. Fl. Austr. II, 157 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 157	
<i>B. cordigera</i> , Benth. in J. Hook. Fl. Tasm. I, 95, t. 16 (1860) ...	W.A.	—	T.	V.	—	—	—	B.f.II, 157	M.fr.III, 100; IX, 45.
<i>B. lenticularis</i> , Sieber in De Cand. prodr. II, 117 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 157	
<i>B. Kiamensis</i> , Benth. Fl. Austr. II, 158 (1864) ...	—	—	—	—	N.S.W.	—	—	B.f.II, 158	
<i>B. ornata</i> , F. v. M., fragm. IV, 13 (1863) ...	W.A.	—	—	—	—	—	—	B.f.II, 158	M.fr.IX, 45.
<i>B. eriocarpa</i> , Benth. in Hueg. enum. 36 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 159	M.fr.IX, 45.
<i>B. divaricata</i> , Turczaninow in Bull. Soc. Mosc. XXVI, 285 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 159	M.fr.IV, 12.
<i>B. calycina</i> , Benth. Fl. Austr. II, 159 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 159	
<i>B. foliosa</i> , Cunningham in Field's N. S. Wales, 347 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.f.II, 160	M.fr.III, 100; IX, 45.
<i>B. cinerea</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 160	M.fr.III, 98.
<i>B. biloba</i> , Benth. in Hueg. enum. 36 (1837) ...	W.A.	—	—	—	—	—	—	B.f.II, 160	M.fr.IX, 45.
<i>B. Preissii</i> , Meissner in Lehm. pl. Preiss. I, 82 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 161	M.fr.IX, 45.
<i>B. concinna</i> , Benth. Fl. Austr. II, 161 (1864) ...	W.A.	—	—	—	—	—	—	B.f.II, 161	
<i>B. carinalis</i> , Benth. in Mitch. Trop. Austr. 290 (1848) ...	—	—	—	—	—	Q.	—	B.f.II, 161	M.fr.IX, 45.
<i>B. rupicola</i> , Cunningham in Benth. Fl. Austr. II, 162 (1864) ...	—	—	—	—	—	Q.	—	B.f.II, 162	
<i>B. linophylla</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	W.A.	—	—	—	—	—	—	B.f.II, 162	M.fr.IX, 45.
<i>B. disticha</i> , Lindley, Bot. Regist. t. 55 (1841) ...	W.A.	—	—	—	—	—	—	B.f.II, 162	
<i>B. prostrata</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 162	M.fr.III, 97.
<i>B. Neo-Anglica</i> , F. v. M., fragm. V, 106 (1866) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.V, 106.
<i>B. buxifolia</i> , Cunningham in Field's N. S. Wales, 348 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.f.II, 163	M.fr.I, 9; IX, 45.
<i>B. Brownii</i> , Benth. Fl. Austr. II, 163 (1864) ...	—	—	—	—	—	Q.	—	B.f.II, 163	
<i>B. rhombifolia</i> , Sieber in De Cand., prodr. II, 117 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 164	M.fr.IX, 45.
<i>B. pulchella</i> , Meissner in Lehm. pl. Preiss. I, 84 (1844) ...	W.A.	—	—	—	—	—	—	B.f.II, 164	
<i>B. microphylla</i> , Smith in Transact. Linn. Soc. IX, 303 (1808) ...	—	—	—	V.	N.S.W.	—	—	B.f.II, 164	M.fr.III, 99.
<i>B. peduncularis</i> , Turczaninow in Bull. Soc. Mosc. XXVI, 287 (1853) ...	W.A.	—	—	—	—	—	—	B.f.II, 165	
<i>B. heterophylla</i> , Ventenat, Jardin de Cels t. 7 (1800) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.II, 163	M.fr.III, 97.
<i>B. rufa</i> , R. Brown in Ait. hort. Kew. sec. ed. III, 267 (1812) ...	W.A.	—	—	—	—	—	—	B.f.II, 165	M.fr.IX, 45.

B. bracteosa, F. v. M. in Benth. Fl. Austr. II, 166 (1864) ...	—	—	—	V.	—	—	B.f. II, 166	
B. riparia, Cunningham in Benth. Fl. Austr. II, 166 (1864) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. II, 166	
B. ensata, Sieber in De Cand. prodr. II, 117 (1825) ...	—	—	—	V.	N.S.W.	Q.	B.f. II, 167	M. fr. XI, 27 & 123.
B. scolopendria, Smith in Transact. Linn. Soc. IX, 307 (1808) ...	—	—	—	—	N.S.W.	—	B.f. II, 167	M. fr. III, 95.
B. Walkeri, F. v. M., fragm. II, 120 (1861) ...	—	S.A.	—	—	N.S.W.	—	B.f. II, 167	M. fr. II, 120; IX, 45; XI, 27.
B. Armitii, F. v. M., fragm. IX, 44 (1875) ...	—	—	—	—	—	Q.	—	M. fr. IX, 44.
B. phylloclada, F. v. M. in Transact. phil. Inst. Vict. III, 52 (1859) ...	—	—	—	—	—	—	N.A.	B.f. II, 168
TEMPLETONIA, R. Brown in Ait. hort. Kew. sec. ed. 269 (1812).								
T. retusa, R. Brown in Ait. hort. Kew. sec. ed. IV, 269 (1872) ...	W.A.	S.A.	—	—	—	—	B.f. II, 169	M. fr. IX, 158.
T. Drummondii, Benth. Fl. Austr. II, 169 (1864) ...	W.A.	—	—	—	—	—	B.f. II, 169	
T. Muelleri, Benth. Fl. Austr. II, 169 (1864) ...	—	—	—	V.	N.S.W.	Q.	B.f. II, 169	M. fr. I, 9; II, 178; IX, 158.
T. aculeata, Benth. Fl. Austr. II, 170 (1864) ...	W.A.	—	—	—	—	—	B.f. II, 170	M. fr. II, 120.
T. egena, Benth. Fl. Austr. II, 170 (1864) ...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.f. II, 170
T. sulcata, Benth. Fl. Austr. II, 171 (1864) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. II, 171
HOVEA, R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812). (Poiretia, Plagiolobium, Platychilum.)								
H. linearis, R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812) ...	—	—	—	—	N.S.W.	—	B.f. II, 172	
H. heterophylla, Cunningham in J. Hook. Fl. Tasm. I, 93, t. 15 (1860) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. II, 172
H. longifolia, R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812) ...	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f. II, 172
H. acutifolia, Cunningham in G. Don, gen. syst. II, 126 (1832) ...	—	—	—	—	N.S.W.	Q.	—	B.f. II, 174
H. longipes, Benth. in Hueg. enum. 37 (1837) ...	—	—	—	—	N.S.W.	Q.	—	B.f. II, 174
H. acanthoclada, F. v. M., fragm. IV, 15 (1864) ...	W.A.	—	—	—	—	—	—	B.f. II, 174
H. chorizemifolia, De Candolle, prodr. II, 116 (1825) ...	W.A.	—	—	—	—	—	—	B.f. II, 174
H. Celsi, Bonpland, plant. cult. à Malmaison. t. 51 (1813) ...	W.A.	—	—	—	—	—	—	B.f. II, 175
H. trisperma, Benth. in Hueg. enum. 37 (1837) ...	W.A.	—	—	—	—	—	—	B.f. II, 176
H. stricta, Meissner in Lehm. pl. Preiss. I, 79 (1844) ...	W.A.	—	—	—	—	—	—	B.f. II, 176
H. pungens, Benth. in Hueg. enum. 37 (1837) ...	W.A.	—	—	—	—	—	—	B.f. II, 176
NEMATOPHYLLUM, F. v. M. in Hook. Kew. miscell. IX, 20 (1857).								
N. Hookeri, F. v. M. in Hook. Kew. misc. IX, 20 (1857) ...	—	—	—	—	—	—	N.A.	B.f. II, 170
GOODIA, Salisbury, paradis. London. t. 41 (1806).								
G. lotifolia, Salisbury, parad. Lond. t. 41 (1806) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. II, 177
G. medicaginea, F. v. M., Second Gen. Rep. II (1854) ...	W.A.	S.A.	—	V.	—	—	—	—
CROTALARIA, Hermann, hort. acad. Lugd. Bat. 196—202, t. 197—203 (1687). (Pentadynamis.)								
C. verrucosa, Linné, spec. plant. 715 (1753) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 179
C. crispata, F. v. M. in Benth. Fl. Austr. II, 179 (1864) ...	—	—	—	—	—	—	N.A.	B.f. II, 179
C. juncea, Linné, spec. plant. 714 (1753) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 179
C. linifolia, Linné, fl. suppl. 322 (1781) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. II, 180
C. humifusa, Graham in Wall. numeric. list 5421 (1828) ...	—	—	—	—	—	Q.	—	—
C. calycina, Schrank, pl. rar. hort. Monac. t. 12 (1819) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 180
C. retusa, Linné, spec. plant. 715 (1753) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. II, 181
C. Mitchelli, Benth. in Mitch. Trop. Austr. 120 (1848) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. II, 181
C. Novae-Hollandiae, De Candolle, prodr. II, 127 (1825) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 181
C. crassipes, Hooker, icon. plant. t. 830 (1852) ...	—	—	—	—	—	—	N.A.	B.f. II, 182
C. Cunninghamii, B. Brown, App. to Sturt's Centr. Austr. 8 (1849) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f. II, 182
C. medicaginea, Lamarck, encycl. meth. II, 201 (1786) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 183
C. incana, Linné, spec. plant. 716 (1753) ...	—	—	—	—	N.S.W.	Q.	—	B.f. II, 183
C. dissitiflora, Benth. in Mitch. Trop. Austr. 386 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. II, 184
C. laburnifolia, Linné, spec. plant. 715 (1753) ...	—	—	—	—	—	Q.	—	B.f. II, 184
C. quinquefolia, Linné, spec. plant. 716 (1753) ...	—	—	—	—	—	Q.	—	B.f. II, 184
C. alata, Hamilton in D. Don, prodr. Fl. Nepal. 241 (1825) ...	—	—	—	—	—	—	N.A.	—
WESTONIA, Sprengel, syst. veg. III, 152 (1826).								
W. trifoliata, Sprengel, syst. veg. III, 230 (1826) ...	—	—	—	—	—	—	N.A.	B.f. II, 185
TRIGONELLA, Linné, gen. pl. 351 (1737).								
T. suavissima, Lindley in Mitch. Three Exped. I, 255 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. II, 187
LOTUS, Tournefort, inst. 402, t. 227 (1700), from C. Bauhin (1623).								
L. corniculatus, Linné, spec. plant. 775 (1753) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. II, 188
L. australis, Andrews, Botanist's Reposit. t. 624 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f. II, 188
PSORALEA, Linné, gen. ed. sec. 358 (1742).								
P. badocana, Benth. Fl. Austr. II, 190 (1864) ...	—	—	—	—	—	Q.	N.A.	B.f. II, 190
P. cephalantha, F. v. M., fragm. IV, 35 (1863) ...	—	—	—	—	—	Q.	—	—
P. Martini, F. v. M., fragm. V, 11 (1865) ...	—	—	—	—	—	—	N.A.	—
P. Archeri, F. v. M., fragm. IV, 21 (1863) ...	—	—	—	—	—	—	N.A.	B.f. II, 190
P. balsamica, F. v. M. in Transact. Vict. Inst. III, 55 (1858) ...	—	—	—	—	—	—	N.A.	B.f. II, 191
P. plumosa, F. v. M., fragm. IV, 22 (1864) ...	—	—	—	—	—	—	N.A.	B.f. II, 191
P. pustulata, F. v. M. in Transact. Vict. Inst. III, 54 (1858) ...	—	—	—	—	—	—	N.A.	B.f. II, 191
P. lachnostachys, F. v. M., fragm. III, 105 (1863) ...	—	—	—	—	—	—	N.A.	B.f. II, 191
P. eriantha, Benth. in Mitch. Trop. Austr. 131 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f. II, 192
P. patens, Lindley in Mitch. Three Exped. II, 9 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. II, 192
P. cinerea, Lindley in Mitch. Three Exped. II, 65 (1838) ...	W.A.	—	—	—	N.S.W.	—	N.A.	B.f. II, 192
P. leucantha, F. v. M. in Transact. Vict. Inst. III, 54 (1859) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.f. II, 193
P. parva, F. v. M. in Transact. Vict. Inst. I, 40 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f. II, 194
P. tenax, Lindley in Mitch. Three Exped. II, 10 (1838) ...	—	—	—	—	N.S.W.	Q.	—	B.f. II, 193
P. adscendens, F. v. M. in Transact. Vict. Inst. I, 40 (1854) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. II, 193
P. Schultzii, F. v. M., fragm. IX, 155 (1875) ...	—	—	—	—	—	—	—	—
P. Testariae, F. v. M., fragm. V, 45 (1865) ...	—	—	—	—	—	Q.	—	—

INDIGOFERA, Royen, pl. hort. Lugd. 372 (1740), from Linné (1737).

I. linifolia, Retzius, observ. bot. IV, 29 (1786) ...	...	...	...	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II, 195	M.fr.III, 101; IX, 44.
I. cordifolia, Heyne in Roth, nov. pl. spec. 357 (1821) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 196	
I. enneaphylla, Linné, mantissa altera 272 (1771) ...	...	...	...	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II, 196	M.fr.III, 102; IX, 44. [44.
I. glandulosa, Willdenow, spec. plant. III, 1227 (1800) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 196	M.fr.IV, 22; VII, 106; IX,
I. haplophylla, F. v. M., fragm. III, 102 (1863) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 196	M.fr.IX, 44.
I. trifoliata, Linné, amoen. acad. IV, 327 (1759) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 197	M.fr.III, 104.
I. trita, Linné fil., suppl. plant. 335 (1781) ...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.II, 197	M.fr.III, 103.
I. parviflora, Heyne in Wall. numeric. list 5457 (1828) ..	...	...	...	—	—	—	—	Q.	N.A.	B.f.II, 197	M.fr.III, 103; IX, 44.
I. viscosa, Lamarck, Encycl. méth. III, 247 (1789) ...	...	...	...	S.A.	—	—	—	Q.	N.A.	B.f.II, 198	M.fr.III, 104; IX, 44.
I. hirsuta, Linné, spec. plant. 751 (1753) ...	...	...	...	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II, 198	M.fr.III, 105; IX, 44.
I. pratensis, F. v. M., essay on plants of Burdek. Exped. 10 (1860)	...	...	...	—	—	—	—	Q.	—	B.f.II, 198	
I. Schultzeana, F. v. M., fragm. VII, 105 (1870) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.VII, 105. [44.
I. monophylla, De Candolle, prodr. II, 222 (1825) ...	...	...	...	S.A.	—	—	—	—	N.A.	B.f.II, 199	M.fr.III, 44; VII, 106; IX,
I. rugosa, Benth. Fl. Austr. II, 199 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 199	
I. saxicola, F. v. M. in Benth. Fl. Austr. II, 199 (1864)...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 199	
I. australis, Willdenow, spec. plant. III, 1235 (1800) ...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 199
I. brevidens, Benth. in Mitch. Trop. Austr. 385 (1848) ...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.II, 200
I. Baileyi, F. v. M., fragm. IX, 43 (1875) ...	...	...	...	—	—	—	—	Q.	—	—	
I. coronillifolia, Cunningham in Benth. Fl. Austr. II, 201 (1864)	...	...	...	S.A.	—	—	N.S.W.	—	—	B.f.II, 201	M.fr.IX, 43.
PTYCHOSEMA, Benth. in Lindl. Bot. Regist. XXV, App. XVI (1839).											
P. pusillum, Benth. in Lindl. Bot. Regist. XXV, App. XVI (1839) W.A.	...	...	...	—	—	—	—	—	—	B.f.II, 201	
P. anomalum, F. v. M., fragm. IX, 62 (1875) ...	...	...	...	—	S.A.	—	—	—	—	—	M.fr.IX, 62.
P. trifoliolatum, F. v. M. in Wing's S. Science Record II, 72 (1882) W.A.	...	...	...	—	—	—	—	—	—	—	
LAMPROLOBIUM, Benth. Fl. Austr. II, 202 (1864).											
L. fruticosum, Benth. Fl. Austr. II, 202 (1864) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 202	M.fr.IX, 68.
L. megalophyllum, F. v. M., fragm. IX, 67 (1875) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.IX, 67.
TEPHROSIA, Persoon, synops. pl. II, 328 (1807).											
T. coriacea, Benth. Fl. Austr. II, 204 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 204	M.fr.IX, 65&66.
T. lamprolobioides, F. v. M., fragm. IX, 64 (1875) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.IX, 64.
T. Forrestiana, F. v. M., fragm. XI, 98 (1880) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.XI, 98.
T. flammea, F. v. M. in Benth. Fl. Austr. II, 204 (1864) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 204	
T. reticulata, Benth. Fl. Austr. II, 205 (1864) ...	...	...	...	—	—	—	—	Q.	N.A.	B.f.II, 205	M.fr.IX, 64.
T. crocea, Benth. Fl. Austr. II, 205 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 205	
T. oblongata, Benth. Fl. Austr. II, 205 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 205	
T. porrecta, Benth. Fl. Austr. II, 206 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 206	
T. polyzyga, F. v. M. in Benth. Fl. Austr. II, 206 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 206	
T. graminifolia, F. v. M. in Benth. Fl. Austr. II, 206 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 206	M.fr.IX, 66.
T. simplicifolia, F. v. M. in Benth. Fl. Austr. II, 206 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 206	
T. nematophylla, F. v. M., fragm. IX, 63 (1879) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.IX, 63.
T. leptoclada, Benth. Fl. Austr. II, 207 (1864) ...	...	...	...	—	—	—	—	Q.	N.A.	B.f.II, 207	
T. brachycarpa, F. v. M. in Benth. Fl. Austr. II, 207 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 207	
T. uniovulata, F. v. M., fragm. XI, 70 (1879) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.XI, 70.
T. Stuartii, Benth. Fl. Austr. II, 207 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 207	
T. conferta, F. v. M., fragm. IX, 65 (1879) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.IX, 75.
T. eriocarpa, Benth. Fl. Austr. II, 207 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 207	
T. phaeosperma, F. v. M. in Benth. Fl. Austr. II, 208 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 208	
T. astragaloides, Benth. Fl. Austr. II, 208 (1864) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 208	
T. juncea, Benth. Fl. Austr. II, 208 (1864) ...	...	...	...	—	—	—	—	Q.	—	B.f.II, 208	M.fr.IX, 64.
T. filipes, Benth. Fl. Austr. II, 208 (1864) ...	...	...	...	—	—	—	—	Q.	N.A.	B.f.II, 208	
T. remotiflora, F. v. M. in Benth. Fl. Austr. II, 209 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 209	
T. oligophylla, Benth. Fl. Austr. II, 209 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 209	
T. macrocarpa, Benth. Fl. Austr. II, 209 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 209	
T. purpurea, Persoon, synops. plant. II, 329 (1807) ...	...	...	...	—	S.A.	—	N.S.W.	Q.	N.A.	B.f.II, 209	
T. Bidwilli, Benth. Fl. Austr. II, 210 (1864) ...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f.II, 210	
T. rosea, F. v. M. in Benth. Fl. Austr. II, 211 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 211	
WISTARIA, Nuttall, Gen. of N. Amer. pl. II, 115 (1818). (Milletim.)											
W. megasperma, F. v. M., fragm. I, ad hist. natur. 10 (1858) ...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f.II, 211	M.fr.I, 10.
SESBANIA, Scopoli, Introd. 308 (1777). (Agati.)											
S. grandiflora, Persoon, synops. pl. II, 316 (1807) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 212	M.fr.II, 88, XI, 33.
S. Aegyptiaca, Persoon, synops. pl. II, 316 (1807) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 212	M.fr.XI, 33.
S. aculeata, Persoon, synops. pl. II, 316 (1807) ...	...	...	...	—	S.A.	—	N.S.W.	Q.	N.A.	B.f.II, 213	M.fr.XI, 33.
S. simpliciuscula, F. v. M. in Benth. Fl. Austr. II, 213 (1864) ...	...	...	...	—	—	—	—	—	N.A.	B.f.II, 213	
S. brachycarpa, F. v. M., fragm. XI, 32 (1878) ...	...	...	...	—	—	—	—	—	N.A.	—	M.fr.IX, 32.
CARMICHAELIA, R. Brown in Bot. Regist. XI, 912 (1825).											
C. exsul, F. v. M., fragm. VII, 126 (1871) ...	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.VII, 126, 193.
CLIANthus, Banks & Solander in G. Don, dichlam. pl. II, 463; indicative (1832). (Donia.)											
C. Dampieri, Cuning. in Trans. hort. soc. Lond. sec. ser. I, 522 (1835) W.A.	...	...	...	S.A.	—	—	N.S.W.	—	N.A.	B.f.II, 214	
STREBLORRHIZA, Endlicher, prodr. fl. Norfolk. 92 (1833).											
S. speciosa, Endlicher, prodr. fl. Norf. 93 (1833) ...	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr.IX, 169.
SWAINSONIA, Salisbury, parad. Londin. t. 28 (1806). (Cyclogyne, Diplolobium.)											
S. Greyana, Lindley, Bot. Regist. t. 66 (1846) ...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.II, 216
S. galegifolia, R. Brown in Ait. hort. Kew. sec. ed. III, 327 (1811) ...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f.II, 217	
S. coronillifolia, Salisbury, parad. Lond. t. 28 (1806) ...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f.II, 217	M.fr.XI, 70.
S. brachycarpa, Benth. Fl. Austr. II, 217 (1864) ...	...	...	...	—	—	—	N.S.W.	Q.	—	B.f.II, 217	M.fr.IX, 155.
S. phacoides, Benth. in Mitch. Trop. Austr. 363 (1848) ...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.II, 217
S. Burkittii, F. v. M. in Benth. Fl. Austr. II, 218 (1864) ...	...	...	...	—	S.A.	—	N.S.W.	—	—	B.f.II, 218	

- GLYCINE, Linné, gen. pl. 349 (1737). (Leptolobium, Leptocyanus.)
- G. falcata, Benth., Fl. Austr. II, 243 (1864) — S.A. — — N.S.W. Q. N.A. B.f.II, 243
- G. clandestina, Wendland, Bot. Reob. 54 (1798) W.A. S.A. T. V. N.S.W. Q. — B.f.II, 243
- G. Latrobeana, Benth., Fl. Austr. II, 244 (1864) — S.A. T. V. — — — B.f.II, 244
- G. tabacina, Benth., Fl. Austr. II, 244 (1864) W.A. S.A. — V. N.S.W. Q. — B.f.II, 244
- G. sericea, Benth., Fl. Austr. II, 245 (1864) — S.A. — V. N.S.W. — — B.f.II, 245
- G. tomentosa, Benth., Fl. Austr. II, 245 (1864) — S.A. — — N.S.W. Q. N.A. B.f.II, 245
- KENNEDYA, Ventenat, Jardin de la Malmaison II, t. 104, 106 (1804). (Caulinia, Moench 1802, Hardenbergia, Physolobium, Zichya, Amphodus.)
- K. nigricans, Lindley, Bot. Regist. t. 1715 (1835) W.A. — — — — — B.f.II, 249 M.fr. VII, 127.
- K. Beckxiana, F. v. M., fragm. XI, 98 (1880) W.A. — — — — — — M.fr. XI, 98.
- K. rubicunda, Ventenat, Jard. de la Malm. t. 104 (1804) — — — V. N.S.W. Q. — B.f.II, 249 M.fr. VII, 127.
- K. procurrens, Benth. in Mitch. trop. Austr. 365 (1848) — — — N.S.W. Q. — B.f.II, 249
- K. prorepens, F. v. M., fragm. VIII, 225 (1874) — S.A. — — — — — M.fr. VIII, 225.
- K. prostrata, R. Brown in Ait. hort. Kew. sec. ed. IV, 299 (1812) W.A. S.A. T. V. N.S.W. Q. — B.f.II, 250 M.fr. VII, 128.
- K. bracteata, Gaudichaud in Freyc. voy. 286, t. 113 (1826) W.A. — — — — — B.f.II, 250
- K. eximia, Lindley in Paxt. Mag. XVI, 35 W.A. — — — — — B.f.II, 250
- K. microphylla, Meissner in Lehm. pl. Preiss. I, 91 (1844) W.A. — — — — — B.f.II, 251
- K. parviflora, Meissner in Lehm. pl. Preiss. I, 91 (1844) W.A. — — — — — B.f.II, 251
- K. Stirlingii, Lindley, Bot. Regist. t. 1845 (1837) W.A. — — — — — B.f.II, 252
- K. glabrata, Lindley, Bot. Regist. t. 1838 (1837) W.A. — — — — — B.f.II, 252 M.fr. VII, 128.
- K. macrophylla, F. v. M., fragm. IV, 79 (1864) W.A. — — — — — B.f.II, 252 M.fr. IV, 79.
- K. coccinea, Ventenat, Jard. de la Malm. t. 105 (1804) W.A. — — — — — B.f.II, 250 M.fr. VII, 128.
- K. monophylla, Ventenat, Jard. de la Malm. t. 106 (1804) — S.A. T. V. N.S.W. Q. — B.f.II, 246 M.fr. VII, 128.
- K. Comptoniana, Link, enum. pl. hort. bot. Berol. II, 235 (1822) W.A. — — — — — B.f.II, 247 M.fr. VII, 128.
- K. retusa, F. v. M., fragm. V, 106 (1866) — — — — — Q. — B.f.II, 247 M.fr. VII, 128.
- ERYTHRINA, Linné, gen. pl. 216 (1737).
- E. vespertilio, Benth. in Mitch. Trop. Austr. 218 (1848) — S.A. — — — — — Q. N.A. B.f.II, 253
- E. Indica, Lamarck, Encyclop. méthod. II, 391 (1786) — — — — — Q. N.A. B.f.II, 253
- STRONGYLODON, T. Vogel in Linnaea X, 585 (1836).
- S. ruber, T. Vogel in Linnaea X, 585 (1836) — — — — — Q. — —
- MUCUNA, Maregraf, hist. nat. Brazil. 18 (1648).
- M. gigantea, De Candolle, prodr. II, 405 (1825) — — — — — N.S.W. Q. N.A. B.f.II, 254
- GALACTIA, P. Browne, Civil & Nat. Hist. of Jamaica 298 (1756).
- G. tenuiflora, Wight & Arnott, prodr. fl. penins. Ind. 206 (1834) — S.A. — — — — — N.S.W. Q. N.A. B.f.II, 255
- G. Muelleri, Benth., Fl. Austr. II, 255 (1864) — — — — — — N.A. B.f.II, 255
- CANAVALIA, De Candolle, prodr. II, 403 (1825). (Canavali.)
- C. obtusifolia, De Candolle, prodr. II, 404 (1825) — — — — — N.S.W. Q. N.A. B.f.II, 256
- PHASEOLUS, Tournefort, inst. 412, t. 232 (1700), from Dioscorides & Columella.
- P. vulgaris, de l'Obel, stirp. icon. 59 (1581) — — — — — — N.A. B.f.II, 257
- P. Truxillensis, Humb. Bonpl. & Kunth, nov. gen. Amer. VI, 451 (1823) — — — — — Q. N.A. B.f.II, 257
- P. Mungo, Linné, mantissa plantar. 101 (1767) — — — — — Q. N.A. B.f.II, 257
- VIGNA, Savi, Dissert. 6 (1824). (Callicystus.)
- V. vexillata, Benth. in Mart. Fl. Bras. Papil. 193, t. 50 (1862) — — — — — N.S.W. Q. N.A. B.f.II, 258
- V. lutea, A. Gray, Bot. Wilk. Exped. I, 454 (1854) — — — — — N.S.W. Q. N.A. B.f.II, 259
- V. luteola, Benth. in Mart. Fl. Bras. Papil. 194, t. 50 (1862) — — — — — N.S.W. Q. — B.f.II, 260
- V. lanceolata, Benth. in Mitch. trop. Austr. 350 (1848) — S.A. — — — — — N.S.W. Q. N.A. B.f.II, 260
- DOLICHOS, Linné, gen. plant. 222 (1737).
- D. biflorus, Linné, spec. plant. 727 (1753) — — — — — — Q. N.A. B.f.II, 261
- DUNBARIA, Wight & Arnott, prodr. I, 258 (1834).
- D. conspersa, Benth. in plant. Junghuhn. (1851) — — — — — — Q. — B.f.II, 262
- CAJANUS, De Candolle, Catal. hort. Monsp. 85 (1813). (Cajan, Atylosia.)
- C. marmoratus, F. v. M.; Atylosia, Benth., Fl. Austr. II, 263 (1864) — — — — — Q. N.A. B.f.II, 263
- C. scarabaeoides, F. v. M.; Atyl., Benth. in pl. Jungh. I, 242 (1851) — — — — — Q. — B.f.II, 263
- C. reticulatus, F. v. M.; Atylosia, Benth., Fl. Austr. II, 263 (1864) — — — — — Q. N.A. B.f.II, 263
- C. grandifolius, F. v. M., Essay on pl. Burdekin Exped. 9 (1860) — — — — — Q. N.A. B.f.II, 264
- C. confertiflorus, F. v. M., Essay on pl. Burdekin Exped. 9 (1860) — — — — — Q. — B.f.II, 264
- C. cinereus; Atylosia, F. v. M. in Benth. Fl. Austr. II, 264 (1864) — — — — — — N.A. B.f.II, 264
- RHYNCHOSIA, Loureiro, Fl. Cochinch. II, 460 (1790).
- R. rhomboidea, F. v. M. in Benth. Fl. Austr. II, 265 (1864) — — — — — — N.A. B.f.II, 265
- R. acutifolia, Benth., Fl. Austr. II, 266 (1864) — — — — — — N.A. B.f.II, 266
- R. rostrata, Benth., Fl. Austr. II, 266 (1864) — — — — — — N.A. B.f.II, 266
- R. Cunninghamii, Benth., Fl. Austr. II, 266 (1864) — — — — — Q. — B.f.II, 266
- R. minima, De Candolle, prodr. II, 385 (1825) — — — — — N.S.W. Q. N.A. B.f.II, 267
- ERIOSEMA, De Candolle, prodr. II, 388 (1825).
- E. Chinense, T. Vogel in Meyen's Beitr. zur Bot. 31 (1843) — — — — — Q. N.A. B.f.II, 268
- FLEMINGIA, Roxburgh in Ait. hort. Kew. sec. ed. IV, 349 (1812).
- F. lineata, Roxburgh in Ait. hort. Kew. sec. ed. IV, 350 (1812) — — — — — — Q. N.A. B.f.II, 268
- F. pauciflora, Benth., Fl. Austr. II, 269 (1864) — — — — — — N.A. B.f.II, 269
- F. parviflora, Benth., Fl. Austr. II, 269 (1864) — — — — — Q. — B.f.II, 269
- F. involucreta, Benth. in pl. Jungh. I, 246 (1851) — — — — — Q. — B.f.II, 269
- ABRUS, Rauwolf in Dalechamps, hist. general. plant. append. 193 (1587).
- A. precatorius, Linné, syst. natur. ed. XII, 533 (1767) — — — — — Q. N.A. B.f.II, 270

- DALBERGIA, Linné, fil. suppl. 52 (1781).
D. densa, Benth in Hook. Lond. Journ. II, 217 (1843) ... — — — — — Q. — B.f.II, 271
- LONCHOCARPUS, Humboldt, Bonpland & Kunth, nov. gen. VI, 383 (1823).
L. Blackii, Benth in Fl. Austr. II, 271 (1864) ... — — — — — N.S.W. Q. — B.f.II, 272
- DERRIS, Loureiro, Fl. Cochinch. II, 432 (1790). (Brachypterum.)
D. scandens, Benth in Journ. Linn. soc. IV, suppl. 103 (1860) ... — — — — — N.S.W. Q. — B.f.II, 272
D. uliginosa, Benth in pl. Jungh. I, 252 (1851) ... — — — — — Q. N.A. B.f.II, 272
- PONGAMIA, Lamarck in Ventenat, Jardin de la Malmaison, t. 28 (1803).
P. glabra, Ventenat, Jard. Malm. t. 28 (1803) ... — — — — — Q. N.A. B.f.II, 273
- SOPHORA, Linné, gen. plant. 125 (1737).
S. tomentosa, Linné, spec. plant. 373 (1753) ... — — — — — N.S.W. Q. N.A. B.f.II, 274 M.fr. VII, 27.
S. Fraseri, Benth in Fl. Austr. II, 274 (1864) ... — — — — — N.S.W. Q. — B.f.II, 274 M.fr. V, 31; VII, 27.
S. tetraptera, J. Miller, icon. plant. t. 1 (1780) ... — — — — — N.S.W. — — — M.fr. VII, 26.
- PODOPETALUM, F. v. M. in Melbourne Chemist, June (1882).
P. Ormondi, F. v. M., fragm. XII (inedit.) ... — — — — — Q. — — —
- CASTANOSPERMUM, Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830).
C. Australe, Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830) ... — — — — — N.S.W. Q. — B.f.II, 275 M.fr. VII, 27.
- BARKLYA, F. v. M., fragm. I, 109 (1859).
B. syringifolia, F. v. M. in Journ. Linn. Soc. III, 158 (1859) ... — — — — — N.S.W. Q. — B.f.II, 275 M.fr. I, 109; III, 163; VIII, 27.
- CAESALPINA, Plumier, nov. pl. Amer. gen. 28, t. 9 (1703), from Ammand. (Guilandina, Caesalpinia.)
C. Bonducella, Fleming in Asiat. Research XI, 159 (1801) ... — — — — — N.S.W. Q. N.A. B.f.II, 276
C. Nuga, W. T. Aiton, hort. Kew., sec. ed. III, 32 (1811) ... — — — — — Q. — B.f.II, 277 M.fr. X, 7.
C. sepiaria, Roxburgh, Fl. Indic. II, 360 (1832) ... — — — — — Q. — B.f.II, 277 M.fr. X, 7.
- MEZONEURON, Desfontaines in Mém. du Mus. IV, 245, t. 10 et 11 (1818).
M. brachycarpum, Benth in Fl. Austr. II, 278 (1864) ... — — — — — N.S.W. Q. — B.f.II, 278 M.fr. X, 7.
M. Scortechinii, F. v. M. in Wing's South. Sc. Record, II, 73 (1882) ... — — — — — N.S.W. Q. — — —
- PTEROLOBIUM, R. Brown in Salt. Abyssin. 65 (1814).
P. nitens, F. v. M. in Benth in Fl. Austr. II, 279 (1864) ... — — — — — Q. — B.f.II, 279 M.fr. X, 7.
- PELTOPHORUM, Vogel in Linnaea XI, 406 (1837).
P. ferrugineum, Benth in Fl. Austr. II, 279 (1864) ... — — — — — N.A. B.f.II, 279 M.fr. X, 7.
- CASSIA, Tournefort, inst. 619, t. 392 (1700), from Plumier (1693). (Cathartocarpus.)
C. Brewsteri, F. v. M., fourth ann. Rep. 17 (1858) ... — — — — — N.S.W. Q. — B.f.II, 282 M.fr. X, 8.
C. laevigata, Willdenow, enum. pl. hort. bot. Berol. I, 441 (1809) ... — — — — — N.S.W. Q. — B.f.II, 282 M.fr. IV, 14; X, 8.
C. Sophora, Linné, spec. plant. 379 (1853) ... — — — — — S.A. — — — N.S.W. Q. — B.f.II, 283 M.fr. IV, 14; X, 9.
C. laxiflora, Benth in Fl. Austr. II, 283 (1864) ... — — — — — — — — — N.A. B.f.II, 283
C. magnifolia, F. v. M., fragm. I, 166 (1859) ... — — — — — — — — — Q. N.A. B.f.II, 283 M.fr. I, 166; X, 9.
C. venusta, F. v. M., fragm. I, 165 (1859) ... — — — — — S.A. — — — — — Q. N.A. B.f.II, 284 M.fr. I, 165; X, 9.
C. notabilis, F. v. M., fragm. III, 28 (1862) ... — — — — — W.A. S.A. — — — — — N.A. B.f.II, 284 M.fr. III, 28; X, 9.
C. pleurocarpa, F. v. M., fragm. I, 223 (1859) ... — — — — — W.A. S.A. — — — — — N.S.W. Q. — B.f.II, 284 M.fr. I, 223; X, 9.
C. glauca, Lamarck, Encycl. méth. I, 647 (1783) ... — — — — — — — — — N.S.W. Q. N.A. B.f.II, 285 M.fr. IV, 13; X, 9.
C. retusa, Solander in Linnaea XI, 72 (1842) ... — — — — — — — — — Q. — B.f.II, 285
C. australis, Sims in Bot. Mag. t. 2676 (1826) ... — — — — — V. N.S.W. Q. N.A. B.f.II, 285 M.fr. X, 9.
C. Chatelainiana, Gaudichaud in Freyc. voy. 485, t. 111 (1826) ... — — — — — W.A. — — — — — N.A. B.f.II, 286
C. glutinosa, De Candolle, prodr. II, 495 (1825) ... — — — — — S.A. — — — — — N.A. B.f.II, 286
C. pruinosa, F. v. M., fragm. III, 48 (1862) ... — — — — — S.A. — — — — — Q. N.A. B.f.II, 286 M.fr. III, 48.
C. circinata, Benth in Mitch. trop. Austr. 384 (1848) ... — — — — — S.A. — — — — — N.S.W. Q. — B.f.II, 286 M.fr. I, 165.
C. phyllodinea, R. Brown, App. to Sturt's Centr. Austr. 15 (1849) ... — — — — — S.A. — — — — — V. N.S.W. — — — B.f.II, 287
C. eremophila, Cunningham, in T. Vogel, gen. Cass. synops. 47 (1837) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. — B.f.II, 287
C. artemisioides, Gaudichaud in De Cand. prodr. II, 495 (1825) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. — B.f.II, 288
C. Sturtii, R. Brown, App. to Sturt's Centr. Austr. 14 (1849) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. — B.f.II, 288
C. desolata, F. v. M. in Linnaea XXV, 387 (1852) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. N.A. B.f.II, 289
C. oligophylla, F. v. M., fragm. III, 49 (1862) ... — — — — — — — — — — — N.A. B.f.II, 289 M.fr. III, 49.
C. cardiosperma, F. v. M., fragm. X, 50 (1877) ... — — — — — W.A. — — — — — — — — — — — M.fr. X, 50.
C. oligoclada, F. v. M., fragm. III, 49 (1862) ... — — — — — — — — — — — N.A. B.f.II, 289 M.fr. III, 49; X, 9.
C. heptanthera, F. v. M., fragm. X, 8 (1876) ... — — — — — — — — — — — N.A. — — — M.fr. X, 8.
C. leptoclada, Benth in Fl. Austr. II, 290 (1864) ... — — — — — — — — — — — N.A. B.f.II, 290
C. Absus, Linné, spec. plant. 376 (1753) ... — — — — — — — — — — — Q. N.A. B.f.II, 290 M.fr. III, 50; X, 9.
C. pumila, Lamarck, Encycl. méth. I, 620 (1783) ... — — — — — — — — — — — Q. — B.f.II, 290 M.fr. III, 47; X, 10.
C. concinna, Benth in Fl. Austr. II, 291 (1864) ... — — — — — — — — — — — N.S.W. Q. N.A. B.f.II, 291 M.fr. X, 10.
C. mimosoides, Linné, spec. plant. 379 (1753) ... — — — — — — — — — — — N.S.W. Q. N.A. B.f.II, 291 M.fr. III, 48; X, 10.
- PETALOSTYLIS, R. Brown, App. to Sturt's Centr. Austr. 79 (1849).
P. labichoides, R. Brown, App. to Sturt's Centr. Austr. 17 (1849) ... — — — — — W.A. S.A. — — — — — N.S.W. Q. N.A. B.f.II, 292 M.fr. X, 7.
- LABICHEA, Gaudichaud in De Cand. prodr. II, 507 (1825).
L. cassioides, Gaudichaud in De Cand. prodr. II, 507 (1825) ... — — — — — W.A. — — — — — — — — — B.f.II, 292
L. nitida, Benth in Fl. Austr. II, 293 (1864) ... — — — — — — — — — — — Q. N.A. B.f.II, 293 M.fr. X, 7.
L. lanceolata, Benth in Hueg. enum. 41 (1837) ... — — — — — W.A. — — — — — — — — — — — B.f.II, 293 M.fr. X, 7.
L. rupestris, Benth in Mitch. Trop. Austr. 342 (1848) ... — — — — — — — — — — — Q. — B.f.II, 293 M.fr. X, 7.
L. punctata, Benth in Bot. Regist. XXV, App. XV (1839) ... — — — — — W.A. — — — — — — — — — — — B.f.II, 294 M.fr. X, 8.
L. Buettneriana, F. v. M. in Melb. Chemist, June (1882) ... — — — — — — — — — — — Q. — — —
- TAMARINDUS, Tournefort, inst. 660, t. 445. App. (1700) from C. Bauhin (1623).
T. Indicus, Linné, spec. plant. 34 (1853) ... — — — — — — — — — — — N.A. B.f.II, 294
- BAUHINIA, Plumier, nov. gen. pl. Amer. 22, t. 13 (1703).
B. Leichhardtii, F. v. M. in Transact. Vict. Inst. III, 50 (1858) ... — — — — — S.A. — — — — — — — — — N.A. B.f.II, 295 M.fr. X, 8.
B. Carronii, F. v. M. in Transact. Vict. Inst. III, 49 (1858) ... — — — — — — — — — — — N.S.W. Q. — B.f.II, 295 M.fr. X, 8.
B. Hookeri, F. v. M. in Transact. Vict. Inst. III, 51 (1858) ... — — — — — — — — — — — Q. N.A. B.f.II, 295 M.fr. X, 8.

B. Gilesii, F. v. M. and Bailey in Wing's S. Sc. Record, Jul. (1882)

—

—

—

—

N.A.

—

AFZELIA, Smith in Transact. Linn. Soc. IV, 221 (1798).

A. bijuga, A. Gray, Bot. Wilk. Expl. Exped. 467, t. 51 (1854)...

—

—

—

—

Q.

—

—

CYNOMETRA, Linné in Act. Soc. Ups. 78 (1741).

C. ramiflora, Linné, spec. plant. 382 (1753)

...

...

...

—

—

Q.

—

B.f.II, 296

ERYTHROPHLAEUM, Afzelius in Tuckey's Congo 430 (1818).

(Laboucheria.)

E. Laboucherii, F. v. M. in Benth. Fl. Austr. II, 297 (1864)

...

—

—

—

—

Q.

N.A.

B.f.II, 297

ENTADA, Adanson, Familles des plantes II, 318 (1763).

E. Pursaetha, De Cand., memoir. sur la fam. des Légum. 12 (1825)

—

—

—

—

—

Q.

—

B.f.II, 298

ADENANTHERA, Royen in Linné, coroll. 7 (1737).

A. pavonina, Linné, spec. plant. 384 (1753)

...

...

...

—

—

Q.

—

—

M.fr.V, 30.

A. abrosperma, F. v. M., fragm. V, 30 (1865)

...

...

...

—

—

Q.

—

B.f.II, 298

M.fr.V, 31.

NEPTUNIA, Loureiro, Fl. Cochinch. II, 653 (1790).

(Dichrostachys.)

N. cinerea, F.v. M.; Dichrostachys, Wight & Arnott, prodr. 271 (1834)

—

—

—

—

—

N.A.

B.f.II, 299

N. spicata, F. v. M., fragm. III, 151 (1862)

...

...

...

—

—

N.A.

B.f.II, 299

M.fr.III, 151.

N. gracilis, Benth in Hook. journ. of Bot. IV, 355 (1842)

...

—

—

—

—

N.S.W.

Q.

N.A.

B.f.II, 300

N. monosperma, F. v. M. in Benth. Fl. Austr. II, 299 (1864)

...

—

—

—

—

Q.

N.A.

B.f.II, 300

ACACIA, Tournefort, inst. 605, t. 375 (1700) from Dioscorides & Plinius.

(Vachellia, Chithonanthus, Tetracheilus, Mimosa partly.)

I. ALATAE.

A. glaucoptera, Benth in Schlecht. Linnaea XXVI, 604 (1853)

W.A.

—

—

—

—

—

—

B.f.II, 320

A. alata, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813)

...

W.A.

—

—

—

—

—

—

B.f.II, 320

A. diptera, Lindley, Bot. Regist. XXV, App. XV (1839)

...

W.A.

—

—

—

—

—

—

B.f.II, 321

A. stenoptera, Benth in Hook. Lond. Journ. I, 325 (1842)

...

W.A.

—

—

—

—

—

—

B.f.II, 321

II. CONTINUAE.

A. incurva, Benth in Hook. Lond. Journ. I, 325 (1842)

...

W.A.

—

—

—

—

—

—

B.f.II, 322

A. trigonophylla, Meissner in Lehm. pl. Preiss. II, 199 (1846)

...

W.A.

—

—

—

—

—

—

B.f.II, 322

M.fr.IV, 3.

A. continua, Benth in Fl. Austr. II, 322 (1864)

...

...

...

—

S.A.

—

V.

N.S.W.

—

—

B.f.II, 322

A. Peuce, F. v. M., fragm. III, 151 (1863)...

...

...

...

—

S.A.

—

—

—

Q.

—

B.f.II, 323

M.fr.III, 151.

III. PUNGENTES.

A. spinescens, Benth in Hook. Lond. Journ. I, 323 (1842)

...

—

S.A.

—

V.

N.S.W.

—

—

—

B.f.II, 323

A. cochlearis, H. L. Wendland, comm. de Acac. aphyll. 15 (1820)

W.A.

S.A.

—

—

—

—

—

—

B.f.II, 324

A. lanigera, Cunningham in Field's N.S. Wales, 345 (1825)

...

—

—

—

V.

N.S.W.

—

—

—

B.f.II, 324

A. phlebocarpa, F. v. M. in Journ. Linn. Soc. III, 119 (1858)

...

—

—

—

—

—

—

N.A.

—

B.f.II, 325

A. trinervata, Sieber in De Cand. prodr. II, 451 (1825)

...

—

—

—

—

N.S.W.

—

—

—

B.f.II, 325

A. colletioides, Cunningham in Hook. Lond. Journ. I, 336 (1842)

...

—

S.A.

—

V.

N.S.W.

—

—

—

B.f.II, 325

M.fr.IV, 4.

A. striatula, Benth in Hook. Lond. Journ. I, 336 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 326

A. campylophylla, Benth in Schlecht. Linnaea XXVI, 605 (1853)

W.A.

—

—

—

—

—

—

—

B.f.II, 326

A. teretifolia, Benth in Hook. Lond. Journ. I, 326 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 326

A. sulcata, R. Brown in Ait. hort. Kew. sec. ed. V, 460 (1813)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 327

A. costata, Benth in Hook. Lond. Journ. I, 339 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 327

A. barbinervis, Benth in Hook. Lond. Journ. I, 326 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 327

A. ataxiphylla, Benth in Schlecht. Linnaea XXVI, 605 (1853)

W.A.

—

—

—

—

—

—

—

B.f.II, 328

A. Baxteri, Benth in Hook. Lond. Journ. I, 327 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 328

A. aureonitens, Lindley, Bot. Regist. XXV, App. XV (1839)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 328

A. quadrisulcata, F. v. M., fragm. III, 127 (1863)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 328

M.f.III, 127.

A. erioclada, Benth in Schlecht. Linnaea XXVI, 606 (1853)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 329

A. siculiformis, Cunningham in Hook. Lond. Journ. I, 337 (1842)

...

—

T.

V.

N.S.W.

—

—

—

B.f.II, 329

A. patens, F. v. M. in Journ. Linn. Soc. III, 120 (1858)...

...

—

—

—

—

—

—

N.A.

—

B.f.II, 329

M.fr.III, 46.

A. laricina, Meissner in Lehm. pl. Preiss. I, 6 (1844)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 330

A. tetragonophylla, F. v. M. in Journ. Linn. Soc. III, 121 (1858)

...

—

S.A.

—

—

N.S.W.

—

—

—

B.f.II, 330

M.fr.IV, 3.

A. genistoides, Cunningham in Benth. Fl. Austr. II, 330 (1864)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 330

A. spacelata, Benth in Hook. Lond. Journ. I, 338 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 331

A. ingrata, Benth in Fl. Austr. II, 331 (1864)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 331

A. juniperina, Willdenow, spec. plant. IV, 1049 (1805)

...

—

—

T.

V.

N.S.W.

Q.

—

—

B.f.II, 331

A. asparagoides, Cunningham in Field's N.S. Wales 343 (1825)...

...

—

—

—

—

N.S.W.

—

—

—

B.f.II, 332

A. tenuifolia, F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)

...

—

—

—

V.

—

—

—

B.f.II, 332

A. diffusa, Edwards, Bot. Regist. t. 634 (1822)

...

—

—

T.

V.

N.S.W.

—

—

—

B.f.II, 332

A. rupicola, F. v. M. in Schlecht. Linnaea XXVI, 610 (1853)

...

—

S.A.

—

V.

—

—

—

B.f.II, 333

IV. CALAMIFORMES.

A. volubilis, F. v. M., fragm. X, 98 (1877)...

...

W.A.

—

—

—

—

—

—

—

M.fr.X, 98.

A. tetragonocarpa, Meissner in Lehm. pl. Preiss. I, 4 (1844)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 336

A. restiacea, Benth in Hook. Lond. Journ. I, 3 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 336

A. squamata, Lindley, Bot. Regist. XXV, App. XV (1839)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 336

A. brachyphylla, Benth in Schlecht. Linnaea XXVI, 615 (1853)

W.A.

—

—

—

—

—

—

—

B.f.II, 337

A. Bynoeana, Benth in Schlecht. Linnaea XXVI, 614 (1853)

...

—

—

—

—

—

—

N.A.

—

B.f.II, 337

M.fr.IV, 9.

A. triptycha, F. v. M. in Benth. Fl. Austr. II, 337 (1864)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 337

A. leptoneura, Benth in Hook. Lond. Journ. I, 341 (1842)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 337

A. rigens, Cunningham in Loudon, hort. Brit. II, 406 (1830)

...

—

S.A.

—

V.

N.S.W.

—

—

—

B.f.II, 337

A. papyrocarpa, Benth in Fl. Austr. II, 338 (1864)

...

W.A.

S.A.

—

—

—

—

—

—

B.f.II, 338

A. Gilesiana, F. v. M. in Melb. Chemist, July (1882)

...

...

...

—

S.A.

—

—

—

—

—

A. sessiliceps, F. v. M. in Melb. Chemist, July (1882)

...

...

...

—

S.A.

—

—

—

—

—

A. pugioniformis, Wendl., comm. de Acac. aphyll. 38, t. 9 (1820)

...

—

—

—

—

N.S.W.

Q.

—

B.f.II, 338

A. juncifolia, Benth in Hook. Lond. Journ. I, 341 (1842)

...

—

—

—

—

N.S.W.

Q.

N.A.

—

B.f.II, 339

A. calamifolia, Sweet in Bot. Regist. t. 839 (1824)

...

...

...

—

S.A.

—

V.

N.S.W.

—

—

B.f.II, 339

A. scirpifolia, Meissner in Bot. Zeitung 10 (1855)...

...

...

W.A.

—

—

—

—

—

—

B.f.II, 339

A. extensa, Lindley, Bot. Regist. XXV, App. XV (1839)

...

W.A.

—

—

—

—

—

—

—

B.f.II, 340

<i>A. gonophylla</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 613 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 340	
<i>A. ericifolia</i> , Benth in Hook. <i>Lond. Journ.</i> I, 345 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 340	
<i>A. uncinella</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 613 (1853)...	W.A.	—	—	—	—	—	—	B.f.II, 341	
<i>A. oxyclada</i> , F. v. M. in Benth. <i>Fl. Austr.</i> II, 341 (1864)	W.A.	—	—	—	—	—	—	B.f.II, 341	
V. BRUNIOIDAE.									
<i>A. cedroides</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 615 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 341	
<i>A. lycopodiifolia</i> , Cunningham in Hook. <i>icon. plant.</i> t. 172 (1837)	—	S.A.	—	—	—	Q.	N.A.	B.f.II, 342	M.fr. 243.
<i>A. spondylophylla</i> , F. v. M., <i>fragm.</i> VIII, 243 (1874)	—	S.A.	—	—	—	—	N.A.	—	
<i>A. galioides</i> , Benth in Hook. <i>Lond. Journ.</i> I, 344 (1842)	—	—	—	—	—	Q.	N.A.	B.f.II, 342	
<i>A. Baueri</i> , Benth in Hook. <i>Lond. Journ.</i> I, 344 (1842)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 342	M.fr. XI, 33.
<i>A. subternata</i> , F. v. M. in <i>Journ. Linn. Soc.</i> III, 124 (1858)	—	—	—	—	—	—	N.A.	B.f.II, 343	
<i>A. minutifolia</i> , F. v. M., <i>fragm.</i> VIII, 243 (1874)...	—	S.A.	—	—	—	—	—	—	M.fr. X, 243.
<i>A. bruniioides</i> , Cunningham in G. Don, <i>gen. syst.</i> II, 404 (1832)	—	—	—	—	—	Q.	—	B.f.II, 343	
<i>A. conferta</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 345 (1842)...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 343	
VI. UNINERVES.									
<i>A. scabra</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 605 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 344	
<i>A. nodiflora</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 621 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 344	
<i>A. spinosissima</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 621 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 344	
<i>A. ulicina</i> , Meissner in <i>Lehm. pl. Preiss.</i> II, 202 (1846)	W.A.	—	—	—	—	—	—	B.f.II, 345	
<i>A. erinacea</i> , Benth in Hook. <i>Lond. Journ.</i> I, 360 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 345	
<i>A. Huegelii</i> , Benth in <i>Hueg. enum.</i> 42 (1837)	W.A.	—	—	—	—	—	—	B.f.II, 345	
<i>A. nervosa</i> , De Candolle, <i>Mem. sur la fam. des Légum.</i> 444 (1825)	W.A.	—	—	—	—	—	—	B.f.II, 346	
<i>A. obovata</i> , Benth in Hook. <i>Lond. Journ.</i> I, 329 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 346	
<i>A. congesta</i> , Benth in Hook. <i>Lond. Journ.</i> I, 327 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 346	
<i>A. dermatophylla</i> , Benth in <i>Fl. Austr.</i> II, 346 (1864)	W.A.	—	—	—	—	—	—	B.f.II, 346	
<i>A. aspera</i> , Lindley in <i>Mitch. Three Exped.</i> II, 139 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 347	
<i>A. armata</i> , R. Brown in <i>Ait. hort. Kew. sec. ed.</i> V, 463 (1813)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.II, 347	
<i>A. idiomorpha</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 329 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 348	
<i>A. Shuttleworthii</i> , Meissner in <i>Lehm. pl. Preiss.</i> I, 7 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 348	
<i>A. Gregorii</i> , F. v. M., <i>fragm.</i> III, 47 (1862)	—	—	—	—	—	—	N.A.	B.f.II, 348	M.fr. III, 47.
<i>A. pilosa</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 607 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 348	
<i>A. crispula</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 607 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 349	
<i>A. crassistipulea</i> , Benth in Hook. <i>Lond. Journ.</i> I, 326 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 349	
<i>A. hastulata</i> , Smith in <i>Rees' Cyclopaed.</i> XXXIX, Suppl. (1819)	W.A.	—	—	—	—	—	—	B.f.II, 349	
<i>A. horridula</i> , Meissner in <i>Lehm. pl. Preiss.</i> I, 9 (1844)	W.A.	—	—	—	—	—	—	B.f.II, 350	
<i>A. divergens</i> , Benth in Hook. <i>Lond. Journ.</i> I, 331 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 350	
<i>A. vomeriformis</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 332 (1842)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 350	
<i>A. plagiophylla</i> , F. v. M. in <i>Journ. Linn. Soc.</i> III, 131 (1858)	—	—	—	—	N.S.W.	Q.	—	—	
<i>A. biflora</i> , R. Brown in <i>Ait. hort. Kew. sec. ed.</i> V, 463 (1813)	W.A.	—	—	—	—	—	—	B.f.II, 351	
<i>A. decipiens</i> , R. Brown in <i>Ait. hort. Kew. sec. ed.</i> V, 463 (1813)	W.A.	—	—	—	—	—	—	B.f.II, 351	
<i>A. cuneata</i> , Benth in <i>Hueg. enum.</i> 42 (1837)...	W.A.	—	—	—	—	—	—	B.f.II, 351	
<i>A. dilatata</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 608 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 352	
<i>A. bidentata</i> , Benth in Hook. <i>Lond. Journ.</i> I, 333 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 352	
<i>A. acanthoclada</i> , F. v. M., <i>fragm.</i> III, 127 (1863)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.II, 352	M.fr. III, 127.
<i>A. obliqua</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 334 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 353	
<i>A. acinacea</i> , Lindley in <i>Mitch. Three Exped.</i> II, 267 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 353	
<i>A. lineata</i> , Cunningham in G. Don, <i>gen. syst.</i> II, 403 (1832)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 353	M.fr. I, 5; II, 177.
<i>A. lachnophylla</i> , F. v. M. in <i>Wing's S. Sc. Record</i> , July (1882)	W.A.	—	—	—	—	—	—	—	
<i>A. triquetra</i> , Benth in Hook. <i>Lond. Journ.</i> I, 358 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 354	
<i>A. ligustrina</i> , Meissner in <i>Lehm. pl. Preiss.</i> II, 203 (1846)	W.A.	—	—	—	—	—	—	B.f.II, 354	
<i>A. Meissneri</i> , Lehmann, <i>delect. sem. hort. bot. Hamb.</i> (1842)	W.A.	—	—	—	—	—	—	B.f.II, 354	
<i>A. anceps</i> , De Candolle, <i>Mem. sur la fam. des Légum.</i> 446 (1825)	W.A.	S.A.	—	—	—	—	—	B.f.II, 355	
<i>A. hispidula</i> , Willdenow, <i>spec. plant.</i> IV, 1054 (1805)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 355	
<i>A. undulifolia</i> , Fraser in <i>Loddiges, Bot. Cabinet</i> XVI, 1544 (1829)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 355	
<i>A. flexifolia</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 359 (1842)...	—	—	—	—	N.S.W.	—	—	B.f.II, 356	
<i>A. dura</i> , Benth in Schlecht. <i>Linnaea</i> XXVI, 622 (1853)	W.A.	—	—	—	—	—	—	B.f.II, 356	
<i>A. spathulata</i> , F. v. M. in Benth. <i>Fl. Austr.</i> II, 356 (1864)	W.A.	—	—	—	—	—	N.A.	B.f.II, 356	
<i>A. microcarpa</i> , F. v. M., <i>second gen. Report</i> 7 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 357	M.fr. I, 6.
<i>A. montana</i> , Benth in Hook. <i>Lond. Journ.</i> I, 360 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 357	
<i>A. verniciflua</i> , Cunningham in <i>Field's N.S. Wales</i> 344 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 358	
<i>A. leprosa</i> , Sieber in <i>De Cand. prodr.</i> II, 450 (1825)	—	—	—	V.	N.S.W.	—	—	B.f.II, 358	
<i>A. stricta</i> , Willdenow, <i>spec. plant.</i> IV, 1052 (1805)	—	—	T.	V.	N.S.W.	—	—	B.f.II, 358	
<i>A. dodonaeifolia</i> , Willdenow, <i>enum. pl. hort. Berol. suppl.</i> 68 (1813)	—	S.A.	—	—	—	—	—	B.f.II, 359	
<i>A. Gnidium</i> , Benth in <i>Fl. Austr.</i> II, 359 (1864)	—	—	—	—	—	Q.	—	B.f.II, 359	
<i>A. ramosissima</i> , Benth in Hook. <i>Lond. Journ.</i> I, 356 (1842)...	W.A.	—	—	—	—	—	—	B.f.II, 360	
<i>A. Dempsteri</i> , F. v. M., <i>fragm.</i> XI, 65 (1879)	W.A.	—	—	—	—	—	—	—	M.fr. XI, 65.
<i>A. Sentis</i> , F. v. M., <i>second gen. Report</i> 11 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.II, 360	M.fr. XI, 65.
<i>A. dentifera</i> , Benth in Maund, <i>Botanist</i> IV, t. 179 (1839)	W.A.	—	—	—	—	—	—	B.f.II, 361	
<i>A. sclerosperma</i> , F. v. M. in <i>Wing's South. Sc. Rec.</i> , July (1882)	W.A.	—	—	—	—	—	N.A.	—	
<i>A. fasciculifera</i> , F. v. M. in Benth. <i>Fl. Austr.</i> II, 361 (1864)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 361	
<i>A. falcata</i> , Willdenow, <i>spec. plant.</i> IV, 1053 (1805)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 361	
<i>A. macradenia</i> , Benth in <i>Mitch. Trop. Austr.</i> 360 (1848)	—	—	—	—	—	Q.	—	B.f.II, 362	
<i>A. penninervis</i> , Sieber in <i>De Cand. prodr.</i> II, 452 (1825)	—	—	T.	V.	N.S.W.	Q.	—	B.f.II, 362	
<i>A. amblyphylla</i> , F. v. M. in <i>Wing's S. Sc. Record</i> , July (1882)	W.A.	—	—	—	—	—	—	—	
<i>A. retinodes</i> , Schlechtendal, <i>Linnaea</i> XX, 664 (1847)	—	S.A.	—	V.	—	—	—	B.f.II, 362	
<i>A. nerifolia</i> , Cunningham in Hook. <i>Lond. Journ.</i> I, 357 (1842)	—	—	—	—	N.S.W.	Q.	—	B.f.II, 363	
<i>A. microbotrya</i> , Benth in Hook. <i>Lond. Journ.</i> I, 353 (1842)...	W.A.	—	—	—	—	—	—	B.f.II, 363	
<i>A. leiophylla</i> , Benth in Hook. <i>Lond. Journ.</i> I, 351 (1842)	W.A.	—	—	—	—	—	—	B.f.II, 364	
<i>A. cyanophylla</i> , Lindley, <i>Bot. Regist. Misc.</i> 49 (1835)	W.A.	—	—	—	—	—	—	B.f.II, 364	
<i>A. pycnantha</i> , Benth in Hook. <i>Lond. Journ.</i> I, 351 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II, 365	

<i>A. notabilis</i> , F. v. M., fragm. I, 6 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.f.II,365	M.fr.I,6.
<i>A. gladiiformis</i> , Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	—	—	—	N.S.W.	—	—	B.f.II,365	
<i>A. obtusata</i> , Sieber in De Cand. prodr. II, 453 (1825)	—	—	—	—	N.S.W.	—	—	B.f.II,366	
<i>A. rubida</i> , Cunningham in Field's N.S.Wales, 344 (1825)	—	—	—	—	N.S.W.	—	—	B.f.II,366	
<i>A. amoena</i> , H. L. Wendland, comm. de Acac. aphyll. 10, t. 4 (1820)	—	—	—	—	V. N.S.W.	—	—	B.f.II,366	
<i>A. hakeoides</i> , Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	S.A.	—	—	V. N.S.W.	—	—	B.f.II,367	
<i>A. salicina</i> , Lindley in Mitch. Three Exped. II, 20 (1838)	W.A.	S.A.	—	—	V. N.S.W.	Q.	N.A.	B.f.II,367	
<i>A. rostellifera</i> , Benth. in Hook. Lond. Journ. I, 356 (1842)	W.A.	—	—	—	—	—	—	B.f.II,368	
<i>A. pycnophylla</i> , Benth. in Fl. Austr. II, 368 (1864)	W.A.	—	—	—	—	—	—	B.f.II,368	
<i>A. Harveyi</i> , Benth. in Fl. Austr. II, 368 (1864)	W.A.	—	—	—	—	—	—	B.f.II,368	
<i>A. iteaphylla</i> , F. v. M. in Schlecht. Linnaea XXVI, 617 (1853)	—	—	—	—	—	—	—	—	
<i>A. suaveolens</i> , Willdenow, spec. plant. IV, 1050 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II,369	
<i>A. subcoerulea</i> , Lindley, Bot. Regist. t. 1075 (1827)	W.A.	—	—	—	—	—	—	B.f.II,369	
<i>A. Lindleyi</i> , Meissner in Lehm. pl. Preiss. I, 14 (1844)	W.A.	—	—	—	—	—	—	B.f.II,370	
<i>A. leptopetala</i> , Benth. in Schlecht. Linnaea XXVI, 619 (1853)	W.A.	—	—	—	—	—	—	B.f.II,370	
<i>A. Dietrichiana</i> , F. v. M. in Melb. Chemist, July (1882)	—	—	—	—	—	—	—	—	
<i>A. Murrayana</i> , F. v. M. in Benth. Fl. Austr. II, 370 (1864)	—	S.A.	—	—	—	Q.	—	B.f.II,370	
<i>A. subulata</i> , Bonpland, pl. rar. cult. à Malm. 110 t. 45 (1813)	—	—	—	—	N.S.W.	—	—	B.f.II,370	
<i>A. linifolia</i> , Willdenow, spec. plant. IV, 1051 (1805)	—	—	—	—	N.S.W.	Q.	—	B.f.II,371	
<i>A. Leichhardtii</i> , Benth. in Fl. Austr. II, 372 (1864)	—	—	—	—	—	Q.	—	B.f.II,372	
<i>A. crassiuscula</i> , H. L. Wendland, comm. de Acac. 31, t. 8 (1820)	—	—	T.	—	N.S.W.	Q.	—	B.f.II,372	
<i>A. buxifolia</i> , Cunningham in Field's N.S.Wales, 344 (1825)	—	—	—	—	N.S.W.	Q.	—	B.f.II,372	
<i>A. lunata</i> , Sieber in De Cand. prodr. II, 452 (1825)	—	—	—	—	V. N.S.W.	Q.	—	B.f.II,373	
<i>A. brachybotrya</i> , Benth. in Hook. Lond. Journ. I, 347 (1842)	—	S.A.	—	—	V. N.S.W.	—	—	B.f.II,373	
<i>A. Wattiana</i> , F. v. M. in Benth. Fl. Austr. II, 374 (1864)	—	S.A.	—	—	—	—	—	B.f.II,374	
<i>A. podalyrifolia</i> , Cunningham in Loudon, hort. Brit. 707 (1830)	—	—	—	—	N.S.W.	Q.	—	B.f.II,374	
<i>A. uncinifera</i> , Benth. in Mitch. Trop. Austr. 341 (1848)	—	—	—	—	—	Q.	—	B.f.II,374	
<i>A. vestita</i> , Ker in Bot. Regist. t. 698 (1823)	—	—	—	—	N.S.W.	—	—	B.f.II,375	
<i>A. cultiriformis</i> , Cunningham in G. Don. gen. syst. II, 406 (1832)	—	—	—	—	N.S.W.	—	—	B.f.II,375	M.fr.XI,123.
<i>A. pravissima</i> , F. v. M., first gen. report 12 (1853)	—	—	—	—	V. N.S.W.	—	—	B.f.II,375	M.fr.I,5.
<i>A. strongylophylla</i> , F. v. M., fragm. VIII, 226 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,226
<i>A. pyrifolia</i> , De Candolle, Mém. sur la fam. des Légum. 447 (1825)	W.A.	—	—	—	—	—	N.A.	B.f.II,376	M.fr.III,17.
<i>A. myrtifolia</i> , Willdenow, spec. plant. IV, 1054 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II,376	
VII. PLURINERVES.									
<i>A. scalpelliformis</i> , Meissner in Lehm. pl. Preiss. II, 200 (1847)	W.A.	—	—	—	—	—	—	B.f.II,377	
<i>A. urophylla</i> , Benth. in Bot. Reg. Misc. 24 (1841)	W.A.	—	—	—	—	—	—	B.f.II,377	
<i>A. Luehmanni</i> , F. v. M., fragm. XI, 116 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.XI,116.
<i>A. sublanata</i> , Benth. in Hueg. enum. 42 (1837)	—	S.A.	—	—	—	—	—	B.f.II,378	M.fr.XI,117.
<i>A. pravifolia</i> , F. v. M., fragm. I, 4 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.I,4.
<i>A. ambylogona</i> , Cunningham in Hook. Lond. Journ. I, 332 (1842)	—	—	—	—	N.S.W.	Q.	—	B.f.II,378	M.fr.IV,3.
<i>A. adnata</i> , F. v. M. in Melb. Chemist, July (1882)	W.A.	—	—	—	—	—	—	—	
<i>A. deltoidea</i> , Cunningham in G. Don. gen. syst. II, 401 (1832)	—	—	—	—	—	—	N.A.	B.f.II,378	
<i>A. stipulosa</i> , F. v. M. in Journ. Linn. Soc. III, 119 (1858)	W.A.	—	—	—	—	—	N.A.	B.f.II,379	M.fr.XI,117.
<i>A. loxophylla</i> , Benth. in Schlecht. Linnaea XXVI, 625 (1853)	W.A.	—	—	—	—	—	—	B.f.II,379	
<i>A. setulifera</i> , Benth. in Schlecht. Linnaea XXVI, 625 (1853)	—	—	—	—	—	—	N.A.	B.f.II,379	
<i>A. translucens</i> , Cunningham in Hook. icon. plant. t. 160 (1837)	—	—	—	—	—	—	N.A.	B.f.II,379	
<i>A. impressa</i> , F. v. M. in Journ. Linn. Soc. III, 133 (1858)	—	—	—	—	—	—	N.A.	B.f.II,380	
<i>A. bivenosa</i> , De Candolle, prodr. II, 452 (1825)	W.A.	—	—	—	—	—	N.A.	B.f.II,380	
<i>A. trineura</i> , F. v. M., pl. of Vict. II, 25 (1860)	—	S.A.	—	V.	—	—	N.A.	B.f.II,381	M.fr.IV,5.
<i>A. nitidula</i> , Benth. in Fl. Austr. II, 381 (1864)	W.A.	—	—	—	—	—	—	B.f.II,381	
<i>A. estropholata</i> , F. v. M. in Wing's South. Sc. Rec., July (1882)	—	S.A.	—	—	—	—	—	—	
<i>A. heteroclita</i> , Meissner in Lehm. pl. Preiss. I, 318 (1845)	W.A.	—	—	—	—	—	—	B.f.II,381	M.fr.IV,6.
<i>A. elongata</i> , Sieber in De Cand. prodr. II, 451 (1825)	—	—	—	—	V. N.S.W.	Q.	—	B.f.II,381	
<i>A. subporosa</i> , F. v. M., pl. of Vict. II, 24 (1860)	—	—	—	—	V. N.S.W.	—	—	B.f.II,382	M.fr.IV,5.
<i>A. Simsii</i> , Cunningham in Hook. Lond. Journ. I, 368 (1842)	—	—	—	—	—	Q.	N.A.	B.f.II,382	
<i>A. dissonera</i> , F. v. M. in Wing's S. Sc. Record, July (1882)	—	—	—	—	—	—	—	—	
<i>A. leptospermoides</i> , Benth. in Schlecht. Linnaea XXVI, 626 (1853)	W.A.	—	—	—	—	—	—	B.f.II,383	
<i>A. homalophylla</i> , Cunningham in Hook. Lond. Journ. I, 365 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II,383	
<i>A. pendula</i> , Cunningham in Loudon, hort. Brit. 490 (1830)	—	—	—	—	N.S.W.	Q.	—	B.f.II,383	
<i>A. Osswaldi</i> , F. v. M., pl. of Vict. II, 27 (1860)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.II,384	M.fr.IV,5.
<i>A. lineolata</i> , Benth. in Schlecht. Linnaea XXVI, 626 (1853)	W.A.	—	—	—	—	—	—	B.f.II,384	
<i>A. quadrimarginea</i> , F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,31.
<i>A. coriacea</i> , De Candolle, Mém. sur la fam. des Légum. 446 (1825)	W.A.	—	—	—	—	—	N.A.	B.f.II,385	
<i>A. stenophylla</i> , Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.II,385	M.fr.XI,67
<i>A. hemignosta</i> , F. v. M. in Journ. Linn. Soc. III, 134 (1858)	—	—	—	—	—	—	N.A.	B.f.II,385	
<i>A. sclerophylla</i> , Lindley in Mitch. Three Exped. II, 138 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II,386	
<i>A. farinosa</i> , Lindley in Mitch. Three Exped. II, 145 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.f.II,386	
<i>A. Whanii</i> , F. v. M. in Benth. Fl. Austr. II, 386 (1864)	—	—	—	V.	—	—	—	B.f.II,386	
<i>A. heteroneura</i> , Benth. in Schlecht. Linnaea XXVI, 624 (1853)	W.A.	—	—	—	—	—	—	B.f.II,387	
<i>A. viscidula</i> , Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	—	—	—	N.S.W.	Q.	—	B.f.II,387	
<i>A. ixiophylla</i> , Benth. in Hook. Lond. Journ. I, 364 (1842)	W.A.	—	—	—	N.S.W.	Q.	—	B.f.II,387	M.fr.IV,6.
<i>A. dictyophleba</i> , F. v. M., fragm. III, 128 (1863)	—	S.A.	—	—	—	Q.	N.A.	B.f.II,388	M.fr.III,128.
<i>A. venulosa</i> , Benth. in Hook. Lond. Journ. I, 366 (1842)	—	—	—	—	N.S.W.	Q.	—	B.f.II,388	
<i>A. cyclops</i> , Cunningham in Loudon, hort. Brit. 407 (1830)	W.A.	S.A.	—	—	—	—	—	B.f.II,388	
<i>A. melanoxylon</i> , R. Brown in Ait. hort. Kew. sec. ed. V, 462 (1813)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II,388	
<i>A. oraria</i> , F. v. M., fragm. XI, 66 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,66.
<i>A. implexa</i> , Benth. in Hook. Lond. Journ. I, 368 (1842)	—	—	—	—	V. N.S.W.	Q.	—	B.f.II,389	
<i>A. harpophylla</i> , F. v. M. in Benth. Fl. Austr. II, 389 (1864)	—	—	—	—	—	Q.	—	B.f.II,389	
<i>A. excelsa</i> , Benth. in Mitch. Trop. Austr. 225 (1848)	—	—	—	—	—	—	—	B.f.II,390	M.fr.IV,6.
<i>A. homaloclada</i> , F. v. M., fragm. XI, 34 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,34.

- A. humifusa, A. Cunningham in Lond. Journ. I, 382 (1842) ... — — — — — Q. N.A. B.f.II,412
 A. denticulosa, F. v. M., fragm. X, 32 (1876) ... W.A. — — — — — — — M.fr.X,32

IX. BIPINNATAE.

- A. Drummondii, Lindley, Bot. Regist. XXV, App. XV (1839)... W.A. — — — — — B.f.II,419
 A. pulchella, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) W.A. — — — — — B.f.II,416
 A. Mitchelli, Benth in Hook. Journ. I, 387 (1842) ... — S.A. — V. — — — B.f.II,417
 A. pentadenia, Lindley, Bot. Regist. t. 1521 (1832) ... W.A. — — — — — B.f.II,417
 A. Gilberti, Meissner in Lehm. pl. Preiss. II, 204 (1847) ... W.A. — — — — — B.f.II,417
 A. nigricans, R. Brown in Ait. hort. Kew. sec. ed. V, 465 (1813) W.A. — — — — — B.f.II,418
 A. Tayloriana, F. v. M. in Wing's S. Sc. Record, July (1882) ... W.A. — — — — — — —
 A. strigosa, Link, enum. pl. hort. Berol. 444 (1822) ... W.A. — — — — — B.f.II,418
 A. elata, Cunningham in Hook. Lond. Journ. I, 383 (1842) ... — — — — N.S.W. — — B.f.II,413
 A. pruinosa, Cunningham in Hook. Lond. Journ. I, 383 (1842)... — — — — N.S.W. — — B.f.II,413
 A. spectabilis, Cunningham in Hook. Lond. Journ. I, 383 (1842) — — — — N.S.W. Q. — B.f.II,413
 A. polybotrya, Benth in Hook. Lond. Journ. I, 384 (1842) ... — — — — N.S.W. Q. — B.f.II,414
 A. discolor, Willdenow, spec. plant. IV, 1068 (1805) ... — — T. V. N.S.W. — — B.f.II,414
 A. decurrens, Willdenow, spec. plant. IV, 1872 (1805) ... — S.A. T. V. N.S.W. Q. — B.f.II,414
 A. dealbata, Link, enum. pl. hort. Berol. 445 (1822) ... — S.A. T. V. N.S.W. — — B.f.II,415
 A. cardiophylla, Cunningham in Hook. Lond. Journ. I, 385 (1842) — — — — N.S.W. — — B.f.II,415
 A. leptoclada, Cunningham in Hook. Lond. Journ. I, 385 (1842) — — — — N.S.W. Q. — B.f.II,416
 A. pubescens, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) — — — — N.S.W. — — B.f.II,416
 A. Farnesiana, Willdenow, spec. plant. IV, 1083 (1805)... — S.A. — — N.S.W. Q. N.A. B.f.II,419
 A. suberosa, Cunningham in Hook. Lond. Journ. I, 499 (1842)... — — — — — N.A. B.f.II,420 M fr.XI,36.
 A. Bidwilli, Benth in Schlecht. Linnaea XXVI, 629 (1853) ... — — — — — Q. N.A. B.f.II,420
 A. pallida, F. v. M. in Journ. Linn. Soc. III, 14 (1858) ... — — — — — N.A. B.f.II,421

ALBIZZIA, Durazzini in Magazzino Toscano III, part IV, 13 (1772). (Zygia 1756, Pithecolobium, Cathormion, Calliandra Enterolobium, Serianthes.)

- A. lophantha, Benth in Hook. Lond. Journ. III, 86 (1844) ... W.A. — — — — — B.f.II,421
 A. Sutherlandi, F. v. M., fragm. VI, 22 (1867) ... — — — — — N.A. — — M.fr.IX,179.
 A. basaltica, Benth, Fl. Austr. II, 422 (1864) ... — — — — — Q. — B.f.II,422
 A. Thozetiana, F. v. M. in Benth. Fl. Austr. II, 422 (1864) ... — — — — — Q. — B.f.II,422 M.fr.IV,9&176
 A. amoenissima, F. v. M., fragm. VIII, 165 (1874) ... — — — — V. N.S.W. Q. — — M.fr.IX,179.
 A. procera, Benth in Hook. Lond. Journ. III, 88 (1844) ... — — — — — N.A. B.f.II,422
 A. pruinosa, F. v. M. in Trimen's Journ. of Bot. 9 (1872) ... — — — — N.S.W. Q. — B.f.II,423 M.fr.IX,179.
 A. canescens, Benth, Fl. Austr. II, 423 (1864)... — — — — — Q. — B.f.II,423 M.fr.IX,179.
 A. Tozeri, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ... — — — — N.S.W. Q. — B.f.II,424 M.fr.IX,179.
 A. Hendersoni, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ... — — — — N.S.W. Q. — — M.fr.X,178.
 A. monilifera, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ... — — — — — N.A. B.f.II,424
 A. ramiflora, F. v. M., fragm. IX, 178 (1875) ... — — — — — Q. — — M.fr.V,10;IX,178.

ARCHIDENDRON, F. v. M., fragm. V, 59 (1865).

- A. Vaillantii, F. v. M., fragm. V, 60 (1865) ... — — — — — Q. — — M.fr.V,9;IX,178.
 A. Lucyi, F. v. M., fragm. VI, 201 (1868)... — — — — — Q. — — M.fr.IX,178.

ROSACEAE.

A. L. de Jussieu, gen. pl. 334 (1789), from B. de Jussieu (1759).

PARINARIUM, A. L. de Jussieu, gen. pl. 342 (1789).

- P. Nonda, F. v. M. in Benth. Fl. Austr. II, 426 (1864) ... — — — — — Q. N.A. B.f.II,426
 P. Griffithianum, Benth in Hook. Niger-Flora 334 (1849) ... — — — — — Q. N.A. B.f.II,426 M.fr.VIII,34

STYLOBASIUM, Desfontaines in Mém. du Mus. V, 37, t. 2 (1819). (Macrostigma.)

- S. spatulatum, Desfontaines in Mém. du Mus. V, 37, t. 2 (1819) W.A. S.A. — — — — — N.A. B.f.II,427
 S. lineare, Nees in Lehm. pl. Preiss. I, 95 (1844) ... W.A. — — — — — — — B.f.II,427

GEUM, Linné, gen. plant. 148 (1737) from Plinius. (Sieversia.)

- G. urbanum, Linné, spec. plant. 501 (1753) ... — S.A. T. V. N.S.W. — — B.f.II,428
 G. renifolium, F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857) — — T. — — — — B.f.II,428

POTENTILLA, Linné, gen. plant. 147 (1737) from Camerarius (1586).

- P. anserina, Linné, spec. plant. 495 (1753) ... — — S.A. T. V. — — — B.f.II,429

RUBUS, Tournefort, Inst. 614, t. 385 (1700) from Plinius.

- R. Gunnianus, Hooker, icon. plant. t. 291 (1840) ... — — T. — — — — B.f.II,430
 R. parvifolius, Linné, spec. plant. 1197 (1753) ... — S.A. T. V. N.S.W. Q. — — B.f.II,430 M.fr.IV,30.
 R. rosifolius, Smith, plantar. icon. ex herb. Linn. t. 60 (1790) ... — — V. N.S.W. Q. — — B.f.II,431 M.fr.IV,32;VIII,34.
 R. Moluccanus, Linné, spec. plant. 1197 (1753) ... — — V. N.S.W. Q. N.A. B.f.II,430 M.fr.IV,31;VIII,34.
 R. Moorei, F. v. M. in Transact. phil. Inst. Vict. II, 67 (1857)... — — — — N.S.W. Q. — — B.f.II,431 M.fr.IV,29&176;VIII,34.

AGRIMONIA, Tournefort, inst. 301, t. 155 (1700) from C. Bauhin (1623).

- A. Eupatoria, Linné, spec. plant. 448 (1753) ... — — — — N.S.W. — — — M.fr.VIII,34.

ALCHEMILLA, Linné, gen. pl. 30 (1737) from C. Bauhin (1623). (Alchimilla.)

- A. vulgaris, C. Bauhin, pinax 319 (1623) ... — — — — V. N.S.W. — — — B.f.II,432

ACAENA, Mutis in Linné, mantiss. II, 200 (1771). (Ancistrum.)

- A. ovina, Cunningham in Field's N.S. Wales, 358 (1825) ... W.A. S.A. T. V. N.S.W. Q. — — B.f.II,433
 A. sanguisorbae, Vahl, enum. plant. I, 294 (1804) ... — S.A. T. V. N.S.W. Q. — — B.f.II,434
 A. montana, J. Hooker in Hook. Lond. Journ. VI, 276 (1847) ... — — T. — — — — B.f.II,434

SAXIFRAGEAE.

Ventenat, Tabl. III, 277 (1799).

ARGOPHYLLUM, R. et G. Forster, charact. gen. 29, t. 15 (1776).

- A. Lejoudanii, F. v. M., fragm. IV, 33 (1864) ... — — — — N.S.W. Q. — — B.f.II,437 M.fr.VI,188.

ABROPHYLLUM, J. Hooker in Benth. fl. Austr. II, 437 (1864). (Brachynema.)									
A. ornans, J. Hooker in Benth. pl. Austr. II, 437 (1864)	...	...	...	...	...	N.S.W.	Q.	—	B.fl.II, 437 M.fr. VI, 189; VII, 150
CUTTSIA, F. v. M., fragm. V, 47 (1865).									
C. viburnea, F. v. M., fragm. V, 47 (1865)...	...	...	...	...	...	N.S.W.	Q.	—	M.fr. V, 47 & 189; VI, 189
COLMEIROA, F. v. M., fragm. VII, 149 (1871).									
C. carpodetoides, F. v. M., fragm. VII, 149 (1871)	...	...	...	...	...	N.S.W.	—	—	M.fr. VII, 149.
QUINTINIA, A. de Candolle, Mon. Campan. 92 (1830).									
Q. Sieberi, A. De Candolle, Monogr. Camp. 92 (1830)	...	...	...	...	...	N.S.W.	Q.	—	M.fr. III, 126; VI, 189.
Q. Verdonii, F. v. M., fragm. II, 225 (1861)	...	...	...	...	...	N.S.W.	Q.	—	M.fr. VI, 189.
Q. Fawkeri, F. v. M., fragm. VI, 92 (1867)	...	...	...	...	...	—	Q.	—	M.fr. VII, 150.
POLYOSMA, Blume, Bijdr. 658 (1825).									
P. Cunninghamii, J. J. Bennett, pl. Jav. rar. 196 (1840)	...	...	...	...	...	N.S.W.	—	—	M.fr. VI, 189.
P. alangiacea, F. v. M., fragm. VIII, 8 (1872)	...	...	...	...	...	—	Q.	—	M.fr. VIII, 8.
ANOPTERUS, Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804.)									
A. glandulosus, Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804)	—	—	—	—	—	T.	—	—	B.fl.II, 439
A. Macleayanus, F. v. M. in Journ. Pharm. Soc. Vict. 43 (1858)	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II, 439 M.fr. VI, 188; VII, 150.
CALLICOMA, Andrews, Bot. Reposit. t. 566 (1809). (Calycomis partly.)									
C. serratifolia, Andrews, Bot. Reposit. t. 566 (1809)	...	...	...	...	...	N.S.W.	Q.	—	M.fr. V, 32.
C. Stutzeri, F. v. M., fragm. V, 34 (1865)	...	...	...	...	...	—	Q.	—	M.fr. VI, 188 & 252; VII, 150.
ANODOPETALUM, Cunningham in Endl. gen. 818 (1839).									
A. biglandulosum, Cunningham in J. Hook. Fl. Tasm. I, 148 (1860)	—	—	—	—	—	T.	—	—	B.fl.II, 440 M.fr. VI, 189.
APHANOPETALUM, Endlicher, nov. stirp. decad. I, 34 (1839). (Platyptelea.)									
A. resinosum, Endlicher, nov. stirp. dec. I, 35 (1839)	...	...	...	...	...	V.	N.S.W.	Q.	— B.fl.II, 441 M.fr. I, 228.
A. occidentale, F. v. M., fragm. I, 228 (1858)	...	...	...	...	...	W.A.	—	—	B.fl.II, 441 M.fr. XI, 20.
CERATOPETALUM, Smith, specim. of the Bot. of New Holland, 9 (1793).									
C. gummiferum, Smith, specim. of the Bot. of New Holl. t. 3 (1793)	—	—	—	—	—	N.S.W.	—	—	B.fl.II, 442
C. apetalum, D. Don in Edinb. new phil. journ. (1830)	...	...	...	...	...	N.S.W.	—	—	B.fl.II, 442 M.fr. VI, 189; VII, 150.
SCHIZOMERIA, D. Don in Edinb. new phil. journ. IX, 94 (1830).									
S. ovata, D. Don in Edinb. new phil. journ. IX, 94 (1830)	...	...	...	...	...	N.S.W.	Q.	—	B.fl.II, 443 M.fr. VI, 189.
ACROPHYLLUM, Benth. in Maund & Henslow, Botanist II, 95 (1840). (Calycomis partly.)									
A. venosum, Benth. in Maund & Henslow, Bot. II, 95 (1840)	—	—	—	—	—	N.S.W.	—	—	B.fl.II, 443 M.fr. VI, 189; VII, 150.
WEINMANNIA, Linné, syst. ed. X, 1005 (1759). (Ackama.)									
W. paniculosa, F. v. M., fragm. II, 175 (1861)	...	...	...	...	...	N.S.W.	Q.	—	B.fl.II, 444 M.fr. II, 83; VI, 188; VII, 150.
W. Biagiana, F. v. M., fragm. V, 16 (1865)	...	...	...	...	...	—	Q.	—	M.fr. VII, 150. [150.]
W. lachnocarpa, F. v. M., fragm. VIII, 7 (1872)...	...	...	...	...	...	N.S.W.	Q.	—	M.fr. VIII, 7.
W. rubifolia, Benth. in Fl. Aust. II, 445 (1864)	...	...	...	...	...	N.S.W.	Q.	—	B.fl.II, 445 M.fr. II, 82 & 175; VI, 181.
W. Benth. in F. v. M., Docum. intercol. Exhibit. 31 (1866)	...	...	...	...	...	—	—	—	B.fl.II, 446 M.fr. V, 180; VI, 188.
GILLBEEA, F. v. M., fragm. V, 17 (1865).									
G. adenopetala, F. v. M., fragm. V, 17 (1865)	...	...	...	...	...	—	Q.	—	M.fr. V, 17; VI, 188.
DAVIDSONIA, F. v. M., fragm. VI, 4 (1867).									
D. pruriens, F. v. M., fragm. VI, 4 (1867)...	...	...	...	...	...	—	Q.	—	M.fr. VI, 249.
TETRACARPAEA, J. Hooker, icon. plant. t. 264 (1840).									
T. Tasmanica, J. Hooker, icon. plant. t. 264 (1840)	...	...	...	...	...	T.	—	—	B.fl.II, 445
EUCRYPHIA, Cavanilles, icon. IV, 48, t. 372 (1797). (Carpodontos.)									
E. Billardieri, Spach, veg. phan. V, 344 (1836)	...	...	...	...	...	T.	—	—	B.fl.II, 446 M.fr. VI, 189; VII, 150.
E. Moorei, F. v. M., fragm. IV, 2 (1864)	...	...	...	...	...	V.	N.S.W.	—	B.fl.II, 447 M.fr. IV, 176; VI, 189; VII, 150 [150]
BAUERA, Banks & Kennedy in Andrews, Bot. Rep. t. 198 (1793).									
B. rubioides, Andrews, Bot. Reposit. t. 198 (1793)	...	...	...	...	...	S.A.	T. V.	N.S.W.	— B.fl.II, 447 M.fr. IV, 23.
B. capitata, Seringe in De Cand. prodr. IV, 13 (1830)	...	...	...	...	...	—	—	N.S.W.	Q. — B.fl.II, 448 M.fr. IV, 24.
B. sessiliflora, F. v. M. in Transact. phil. Soc. Vict. I, 41 (1854)	...	...	...	...	...	—	—	V.	— B.fl.II, 448 M.fr. IV, 24; VII, 150
EREMOSYNE, Endlicher in Hueg. enum. pl. Nov. Holl. austr. occ. 53 (1837).									
E. pectinata, Endlicher in Hueg. enum. pl. 53 (1837)	...	...	...	...	...	W.A.	—	—	— B.fl.II, 449 M.fr. VI, 189.
CEPHALOTUS, Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806).									
C. follicularis, Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806) W.A.	—	—	—	—	—	—	—	—	B.fl.II, 449 M.fr. VI, 189.

CRASSULACEAE.

De Candolle, Bull. de la soc. philom. n. 49 (1801).

TILLAEA, Micheli, nov. gen. 22, t. 20 (1729). (Bulliarda, Crassulaceae subgenus.)									
T. verticillaris, De Candolle, prodr. III, 382 (1828)	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W. Q. — B.fl.II, 451 M.fr. XI, 118.
T. purpurata, J. Hooker in Lond. Journ. of Bot. VI, 472 (1847)	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W. — B.fl.II, 451 M.fr. XI, 118.
T. macrantha, J. Hooker in Hook. icon. plant. t. 310 (1841)	...	...	...	...	...	—	S.A.	T. V.	N.S.W. — B.fl.II, 452 M.fr. XI, 117.
T. pedicellosa, F. v. M., fragm. XI, 118 (1881)	...	...	...	...	...	W.A.	—	—	— M.fr. XI, 118.
T. intricata, Nees in Lehm. pl. Preiss. I, 278 (1844)	...	...	...	...	...	W.A.	—	—	— M.fr. XI, 117.
T. recurva, J. Hooker, Fl. Tasm. I, 146 (1860)	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W. Q. — B.fl.II, 452 M.fr. XI, 118.

HAMAMELIDEAE.

R. Brown in Abel, Narr. journ. Chin. 374 (1818).

Genus and species undetermined	...	...	...	...	...	—	—	—	— Q. — —
--------------------------------	-----	-----	-----	-----	-----	---	---	---	----------

ONAGREAE.

Adanson, Familles des plantes II, 81 (1763), from B. de Jussieu (1759).

OENOTHERA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 112 (1737).									
O. Tasmanica, J. Hooker, Fl. Tasm. I, 119 (1860)	...	...	—	—	T.	—	—	—	B.f.III, 303
EPILOBIUM, Dillenius in Linné, syst. nat. 8 (1735), from Gesner (1542).									
E. tetragonum, Linné, spec. plant. 348 (1753)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W. Q. — B.f.III, 305
JUSSIEUA, Linné, gen. plant. 126 (1737). (Jussiaea.)									
J. repens, Linné, spec. plant. 338 (1753)	...	...	...	...	—	S.A.	—	V.	N.S.W. Q. — B.f.III, 306
J. suffruticosa, Linné, spec. plant. 338 (1753)	...	...	...	...	—	—	—	N.S.W. Q.	N.A. B.f.III, 307 M.fr.III, 13.
LUDWIGIA, Linné, coroll. 3 (1737).									
L. parviflora, Roxburgh, Fl. Ind. edit. Carey. I, 440 (1820)	—	—	—	—	—	—	—	Q.	N.A. B.f.III, 307 M.fr.III, 129; IV, 175.

SALICARIEAE.

Adanson, Familles des plantes II, 232 (1763), from B. de Jussieu (1759).

ROOTALA, Linné, mantissa altera 143 (1771).									
R. verticillaris, Linné, mantissa altera 143 (1771)	...	...	—	—	—	—	—	N.A.	B.f.III, 295 M.fr.III, 108.
R. Roxburghiana, Wight, Icon. t. 260 (1840)	...	...	—	—	—	—	—	Q.	N.A. B.f.III, 296 M.fr.XI, 118.
R. diandra, F. v. M. in Transact. Roy. Soc. of N.S. Wales 89 (1880)	—	—	—	—	—	—	—	N.A.	B.f.III, 296
AMMANNIA, Houston in Linné, gen. plant. 337 (1737).									
A. crinipes, F. v. M. in Transact. phil. Soc. Vict. III, 49 (1858)	—	—	—	—	—	—	—	N.A.	B.f.III, 296
A. triflora, R. Brown in Benth. Fl. Austr. III, 297 (1866)	—	—	—	—	—	—	—	N.A.	B.f.III, 297
A. baccifera, Linné, spec. plant. ed. sec. 175 (1762)	...	...	—	—	S.A.	—	—	Q.	N.A. B.f.III, 297
A. auriculata, Willdenow, hort. Berol. t. 7 (1806)	...	...	—	—	—	—	—	Q.	N.A. B.f.III, 297
A. multiflora, Roxburgh, Fl. Ind. edit. Carey. I, 447 (1820)	...	...	—	—	S.A.	—	V.	N.S.W. Q.	N.A. B.f.III, 298
LAGERSTROEMIA, Linné, syst. ed. X, 1076 (1759).									
L. Indica, Linné, spec. plant. 734 (1753)	...	...	—	—	—	—	—	Q.	— M.fr.VIII, 35.
L. Flos Reginae, Retzius, observ. botan. V, 25 (1789)	...	...	—	—	—	—	—	Q.	— M.fr.XII.
LYTHRUM, Linné, syst. nat. 8 (1735); Linné, gen. plant. 136 (1737).									
L. Salicaria, Linné, spec. plant. 446 (1753)	...	...	—	—	S.A.	T.	V.	N.S.W. Q.	— B.f.III, 298 M.fr.VII, 147.
L. Hyssopifolia, Linné, spec. plant. 447 (1753)	...	...	—	—	S.A.	T.	V.	N.S.W. Q.	— B.f.III, 299 M.fr.VII, 147.
NESAEA, Commerçon in A. L. de Jussieu, gen. plant. 332 (1789).									
N. Robertsii, F. v. M., fragm. VII, 145 (1871)	...	...	—	—	—	—	—	Q.	— M.fr.VII, 145.
N. Arnheimica, F. v. M., fragm. VII, 146 (1871)	...	...	—	—	—	—	—	N.A.	B.f.III, 299 M.fr.III, 109; VII, 146.
PEMPHIS, R. et G. Forster, charact. gen. 67, t. 34 (1776).									
P. acidula, R. & G. Forster, charact. gen. 67, t. 34 (1776)	...	...	—	—	—	—	—	Q.	N.A. B.f.III, 300 M.fr.VII, 147.
LAWSONIA, Linné, gen. plant. 111 (1737).									
L. alba, Lamarck, Encycl. méth. III, 106 (1789)	...	...	—	—	—	—	—	—	N.A. B.f.III, 301

HALORAGAEAE.

R. Brown in Flind. voy. II, 549 (1814).

LOUDONIA, Lindley, Bot. Regist. XXV, App. XLII (1839). (Glischrocaryon.)									
L. aurea, Lindley, Bot. Regist. XXV, App. XLII (1839)	...	...	W.A.	S.A.	—	—	—	—	B.f.II, 472 M.fr.VIII, 162; XI, 20.
L. Behrii, Schlechtendal, Linnaea XX, 648 (1847)	...	...	—	S.A.	—	V.	N.S.W.	—	B.f.II, 472 M.fr.VIII, 162.
L. Roei, Schlechtendal, Linnaea XX, 648 (1847)	...	...	W.A.	—	—	—	—	—	B.f.II, 472 M.fr.VIII, 162.
HALORAGIS, R. et G. Forster, charact. gen. 61, t. 31 (1776). (Cercodia, Goniocarpus.)									
H. Gossei, F. v. M., fragm. VIII, 161 (1874)	...	...	—	S.A.	—	—	—	N.A.	— M.fr.XI, 134.
H. trigonocarpa, F. v. M., fragm. X, 84 (1876)	...	...	W.A.	S.A.	—	—	—	—	— M.fr.X, 84.
H. digyna, Labillardière, Nov. Holl. pl. spec. I, 101, t. 129 (1804)	...	...	W.A.	—	—	—	—	—	B.f.II, 475
H. mucronata, Benth. Fl. Austr. II, 475 (1864)	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f.II, 475
H. pithyoides, Benth. Fl. Austr. II, 476 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 476
H. cordigera, Fenzl in Hueg. enum. 45 (1837)	...	...	W.A.	—	—	—	—	—	B.f.II, 476
H. elata, Cunningham in Hueg. enum. 45 (1837)	...	...	—	S.A.	—	V.	N.S.W. Q.	—	B.f.II, 476 M.fr.VIII, 162.
H. tenuifolia, Benth. Fl. Austr. II, 477 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 477
H. scoparia, Fenzl in Hueg. enum. 45 (1837)	...	...	W.A.	—	—	—	—	—	B.f.II, 477 M.fr.X, 54.
H. aculeolata, Benth. Fl. Austr. II, 477 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 477 M.fr.X, 54.
H. foliosa, Benth. Fl. Austr. II, 477 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 477
H. confertifolia, F. v. M., fragm. X, 53 (1876)	...	...	W.A.	—	—	—	—	—	— M.fr.X, 54.
H. platycarpa, Benth. Fl. Austr. II, 478 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 478
H. ceratophylla, Zahlbruckner in Endl. Atakta, 16, t. 15 (1833)	—	—	S.A.	T.	V.	N.S.W. Q.	N.A.	—	B.f.II, 478
H. acutangula, F. v. M. in Transact. Vict. Inst. 125 (1854)	...	...	—	S.A.	—	—	—	—	B.f.II, 478
H. hexandra, F. v. M., fragm. III, 31 (1862)	...	...	W.A.	—	—	—	—	—	B.f.II, 478 M.fr.III, 31.
H. odontocarpa, F. v. M., fragm. I, 108 (1859)	...	...	—	S.A.	—	V.	N.S.W.	—	B.f.II, 479 M.fr.I, 108; X, 54.
H. serra, Brongniart in Duperr. Voy. sur la Coquill. t. 69 (1829)	—	—	—	—	—	N.S.W.	—	—	B.f.II, 479
H. glauca, Lindley in Mitch. Trop. Austr. 91 (1848)	...	...	—	—	—	N.S.W.	—	—	B.f.II, 479
H. alata, N. J. Jacquin, icon. plant. rar. I, 7, t. 69 (1786)	...	...	—	—	—	V.	N.S.W.	—	B.f.II, 479
H. racemosa, Labillardière, Nov. Holl. pl. spec. I, 100, t. 128 (1804)	...	...	W.A.	—	—	—	—	—	B.f.II, 480 M.fr.VIII, 162.
H. rotundifolia, Benth. Fl. Austr. II, 480 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 480 M.fr.VIII, 162.
H. rudis, Benth. Fl. Austr. II, 480 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 480
H. paniculata, R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 481
H. pusilla, R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 481
H. intricata, Benth. Fl. Austr. II, 481 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 481
H. nodulosa, Walpers, repert. V, 672 (1846)	...	...	W.A.	—	—	—	—	—	B.f.II, 481
H. trichostachya, Benth. Fl. Austr. II, 481 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 481
H. lanceolata, R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	W.A.	—	—	—	—	—	B.f.II, 482
H. micrantha, R. Brown in Flind. Voy. II, 550 (1814)	...	...	—	S.A.	T.	V.	N.S.W. Q.	—	B.f.II, 482 M.fr.VIII, 162; X, 54.

H. stricta, R. Brown in Benth. Fl. Austr. II, 482 (1864)	...	—	—	—	—	Q.	—	B.f.II, 482	
H. heterophylla, Brongniart in Duperr. Voy. t. 68 (1829)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.II, 483
H. pinnatifida, A. Gray, Bot. Wilk. Expl. Exped. I, 627 (1854)	...	—	—	—	—	N.S.W.	—	—	B.f.II, 483
H. leptothoca, F. v. M., fragm. III, 32 (1862)	...	—	—	—	—	—	N.A.	—	B.f.II, 483
H. tetragyna, R. Brown in Flind. Voy. II, 550 (1814)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 484
H. teucroides, A. Gray, Bot. Wilk. Expl. Exped. I, 625 (1854)	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 484
H. scordioides, Benth. Fl. Austr. II, 485 (1864)	...	W.A.	—	—	—	—	—	—	B.f.II, 485
H. depressa, Walpers, repertor. II, 99 (1843)	...	—	—	T.	V.	—	—	—	B.f.II, 485
H. salsoloides, Benth. Fl. Austr. II, 485 (1864)	...	—	—	—	—	N.S.W.	—	—	B.f.II, 485

M.fr. VIII, 162.
M.fr. IV, 26.

MEIONECTES, R. Brown in Flind. Voy. II, 550 (1814).									
M. Brownii, J. Hooker in Hook. icon. plant. t. 306 (1841)	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 486

M.fr. VIII, 163.

MYRIOPHYLLUM, l'Ecluse, rar. stirp. hist. II, 252 (1583) from Dioscorides. (Pelonastes).									
M. variifolium, J. Hooker in Hook. icon. plant. t. 239 (1840)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 487
M. elatinoides, Gaudichaud in Ann. des sc. nat. V, 105 (1825)	...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 487
M. verrucosum, Lindley in Mitch. Trop. Austr. 384 (1848)	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.II, 488
M. latifolium, F. v. M., fragm. II, 87, (1860)	...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 488
M. Muelleri, Sonder in Linnaea, XXVIII, 233 (1855)	...	W.A.	S.A.	—	V.	—	—	—	B.f.II, 488
M. amphibium, Labillardiere, Nov. Holl. pl. spec. II, 70, t. 220 (1806)	...	—	S.A.	T.	V.	—	—	—	B.f.II, 489
M. pedunculatum, J. Hooker in Lond. Journ. VI, 474 (1847)	...	W.A.	S.A.	T.	V.	—	—	—	B.f.II, 489
M. dicoccum, F. v. M. in Transact. phil. Inst. Vict. III, 41 (1858)	...	—	—	—	—	—	N.A.	—	B.f.II, 489
M. trachycarpum, F. v. M., fragm. II, 87 (1860)	...	—	—	—	—	—	N.A.	—	B.f.II, 489
M. gracile, Benth. Fl. Austr. II, 489 (1864)	...	—	—	—	—	N.S.W.	Q.	—	B.f.II, 489
M. filiforme, Benth. Fl. Austr. II, 489 (1864)	...	—	—	—	—	—	N.A.	—	B.f.II, 489
M. integrifolium, J. Hooker, Fl. Tasm. I, 123, t. 23 (1860)	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.II, 490
M. Drummondii, J. Hooker, Fl. Tasm. I, 123, implied (1860)	...	W.A.	—	—	—	—	—	—	B.f.II, 490

M.fr. VIII, 163.
M.fr. II, 87; IV, 172; X, 54.
M.fr. VIII, 162.
M.fr. VIII, 163.
M.fr. VIII, 162.
M.fr. VIII, 162.
M.fr. II, 87.

GUNNERA, Linné, mantiss. plant. 16 (1767). (Milligania.)									
G. cordifolia, J. Hooker, Fl. Tasm. I, 125 (1860)	...	—	—	T.	—	—	—	—	B.f.II, 491

CERATOPHYLLUM, Linné, syst. nat. 9 (1735); Linné, gen. plant. 290 (1737).									
C. demersum, Linné, spec. plant. 992 (1753)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.II, 491

M.fr. VIII, 163.

CALLITRICHE, Linné, syst. nat. ed. sext. 82 (1748).									
C. verna, Linné, flor. Suec. edit. sec. (1755)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.II, 492
C. Muelleri, Sonder in Linnaea XXVIII, 229 (1855)	...	—	—	—	V.	N.S.W.	—	—	—

M.fr. VIII, 163.

RHIZOPHOREAE.

R. Brown in Flind. Voy. II, 549 (1814).

RHIZOPHORA, Linné, gen. plant. 137 (1737).									
R. mucronata, Lamarck, Encycl. méth. VI, 169 (1804)	...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.II, 493
CERIOPS, Arnott in Jardine, Ann. of nat. hist. I, 363 (1838).									
C. Candolleana, Arnott in Ann. of nat. hist. I, 364 (1838)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 494
BRUGUIERA, Lamarck, Encycl. méth. IV, 696, t. 397 (1796).									
B. Rhéedii, Blume, enum. pl. Jav. 92 (1827)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 494
B. caryophylloides, Blume, enum. pl. Jav. 93 (1827)	...	—	—	—	—	—	—	—	—
B. gymnorhiza, Lamarck, Encycl. méth. IV, 696, t. 397 (1796)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 495
CARALLIA, Roxburgh, in Flind. Voy. II, 549 (1814).									
C. integririma, De Candolle, prodr. III, 33 (1823)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 495

M.fr. IX, 159.
M.fr. IX, 159.
M.fr. IX, 159.
M.fr. IX, 159.
M.fr. IX, 159.

COMBRETACEAE.

R. Brown, prodr. 351 (1810).

TERMINALIA, Linné, mantiss. plant. 21 (1767). (Chuncoa.)									
T. platyptera, F. v. M., fragm. II, 151 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 498	M.fr. II, 151; IX, 160.
T. volucris, R. Brown in Benth. Fl. Austr. II, 498 (1864)	...	—	—	—	—	—	N.A.	B.f.II, 498	
T. oblongata, F. v. M., fragm. II, 152 (1861)	...	—	—	—	—	—	Q.	—	B.f.II, 499
T. bursarina, F. v. M., fragm. II, 149 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 499	M.fr. III, 167; IX, 160.
T. circumalata, F. v. M., fragm. III, 91 (1862)	...	—	—	—	—	—	N.A.	B.f.II, 499	M.fr. II, 149.
T. pterocarya, F. v. M., fragm. II, 152 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 500	M.fr. III, 91.
T. Thozetii, Benth. Fl. Austr. II, 500 (1864)	...	—	—	—	—	—	Q.	—	B.f.II, 500
T. Catappa, Linné, mantiss. pl. 128 (1767)	...	—	—	—	—	—	Q.	—	—
T. melanocarpa, F. v. M., fragm. III, 92 (1862)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 500
T. Muelleri, Benth. Fl. Austr. II, 500 (1864)	...	—	—	—	—	—	Q.	—	B.f.II, 500
T. latipes, Benth. Fl. Austr. II, 501 (1864)	...	—	—	—	—	—	N.A.	B.f.II, 501	M.fr. III, 92; IX, 160.
T. edulis, F. v. M., fragm. II, 151 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 501	M.fr. IX, 160.
T. discolor, F. v. M., fragm. III, 92 (1862)	...	—	—	—	—	—	N.A.	B.f.II, 501	M.fr. II, 151; IX, 160.
T. porphyrocarpa, F. v. M. in Benth. Fl. Austr. II, 501 (1864)	...	—	—	—	—	—	Q.	—	B.f.II, 501
T. platyphylla, F. v. M., fragm. II, 150 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 502	M.fr. III, 92.
T. sericocarpa, F. v. M., fragm. IX, 159 (1875)	...	—	—	—	—	—	Q.	—	—
T. microcarpa, Decaisne in Ann. du Mus. III (1834)	...	—	—	—	—	—	N.A.	B.f.II, 502	M.fr. II, 150.
T. petiolaris, Cunningham in Benth. Fl. Austr. II, 502 (1864)	...	—	—	—	—	—	N.A.	B.f.II, 502	M.fr. IX, 159.
T. erythrocarpa, F. v. M., fragm. II, 150 (1861)	...	—	—	—	—	—	N.A.	B.f.II, 503	
T. grandiflora, Benth. Fl. Austr. II, 503 (1864)	...	—	—	—	—	—	N.A.	B.f.II, 503	M.fr. II, 150.

M.fr. IX, 160.

LUMNITZERA, Willdenow, in den Verh. der Ges. Nat. Freunde zu Berl. IV, 186 (1803).									
L. coccinea, Wight & Arnott, prodr. 316 (1834)	...	—	—	—	—	—	Q.	—	B.f.II, 503
L. racemosa, Willd. in Verh. der Nat. Freunde zu Berl. IV, 186 (1803)	...	—	—	—	—	—	Q.	N.A.	B.f.II, 504

M.fr. IX, 160.
M.fr. IX, 160.

MACROPTERANTHES, F. v. M., fragm. III, 91 (1862).

M. montana, F. v. M., fragm. III, 91 (1862)	—	—	—	—	Q.	—	B. fl. II, 504	M. fr. II, 149; III, 91 & 166.
M. Kekwickii, F. v. M., fragm. III, 151 (1863)	—	—	—	—	—	N.A.	B. fl. II, 504	M. fr. III, 151.
M. Leichhardtii, F. v. M., fragm. III, 91 (1862)	—	—	—	—	Q.	—	B. fl. II, 505	M. fr. III, 91.
M. Fitzalani, F. v. M., fragm. VIII, 160 (1874)	—	—	—	—	Q.	—	—	M. fr. VIII, 160.

GYROCARPUS, N. Jacquin, select. stirp. Amer. hist. 282 (1763).

G. Americanus, N. Jacquin, select. stirp. Amer. hist. 282, t. 178 (1763)	—	—	—	—	Q.	N.A.	B. fl. II, 505	M. fr. IX, 160.
---	---	---	---	---	----	------	----------------	-----------------

MYRTACEAE.

Adanson, Familles des plantes II, 86 (1763), from B. de Jussieu (1759).

ACTINODIUM, Schauer in Schlecht. Linnæa X, 311 (1835). (Triphelia.)

A. Cunninghamii, Schauer in Lindl. introd. to the nat. syst., sec. ed. 440 (1835)	—	—	—	—	—	—	B. fl. III, 5	
--	---	---	---	---	---	---	---------------	--

DARWINIA, Rudge in Transact. Linn. Soc. XI, 299 (1813). (Genetyllis, Chamaelaucium, Homoranthus, Hedaroma, Polyzone, Francisia, Decalophium, Schuermannia, Cryptostemon.)

D. macrostegia, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 8	
D. Hookeriana, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 9	M. fr. VIII, 182.
D. fimbriata, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 9	M. fr. VIII, 182; IX, 177.
D. speciosa, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 9	M. fr. XI, 10.
D. Meissneri, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 9	M. fr. VIII, 182; IX, 177.
D. helichrysoides, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 10	
D. Neildiana, F. v. M., fragm. IX, 177 (1875)	W.A.	—	—	—	—	—	—	M. fr. IX, 177.
D. oederoides, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 10	M. fr. IX, 177; XI, 9.
D. virescens, Benth. in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	B. fl. III, 10	M. fr. XI, 10.
D. Oldfieldii, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 10	
D. purpurea, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 11	M. fr. XI, 9. [177; XI, 9.
D. citriodora, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 11	M. fr. I, 169; IV, 174; IX,
D. thymoides, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 11	M. fr. VIII, 182; IX, 176.
D. taxifolia, Cunningham in Field's N.S. Wales 352 (1825)	—	—	—	V.	N.S.W.	—	B. fl. III, 12	M. fr. IX, 176.
D. vestita, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 12	M. fr. VIII, 182; IX, 176.
D. pauciflora, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 12	M. fr. XI, 9.
D. diosmoides, Benth. in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	B. fl. III, 13	M. fr. IX, 176.
D. fascicularis, Rudge in Transact. Linn. Soc. XI, 299, t. 22 (1813)	—	—	—	—	N.S.W.	—	B. fl. III, 13	M. fr. IX, 176.
D. pinifolia, Benth. in Journ. Linn. Soc. IX, 181 (1865)	W.A.	—	—	—	—	—	B. fl. III, 14	
D. rhadinophylla, F. v. M., fragm. IX, 175 (1875)	W.A.	—	—	—	—	—	—	M. fr. IX, 175.
D. sanguinea, Benth. in Journ. Linn. Soc. IX, 181 (1865)	W.A.	—	—	—	—	—	B. fl. III, 14	M. fr. IX, 176.
D. micropetala, Benth. in Journ. Linn. Soc. IX, 181 (1865)	—	S.A.	—	—	—	—	B. fl. III, 14	M. fr. I, 12; IX, 176.
D. Schuermanni, Benth. in Journ. Linn. Soc. IX, 181 (1865)	—	S.A.	—	—	—	—	B. fl. III, 14	M. fr. I, 12.
D. verticordina, Benth. in Journ. Linn. Soc. IX, 181 (1865)	W.A.	—	—	—	—	—	B. fl. III, 15	M. fr. IV, 57.
D. Thomasii, Benth. in Journ. Linn. Soc. IX, 181 (1865)	—	—	—	—	Q.	—	B. fl. III, 15	M. fr. IV, 137.
D. virgata, F. v. M., fragm. IX, 176 (1875)	—	—	—	—	N.S.W.	Q.	B. fl. III, 16	M. fr. IX, 176.
D. ciliata, F. v. M.; Chamaelaucium, Desfontaines in Mém. du Mus. Par. V, 40, t. 3 (1821)	W.A.	—	—	—	—	—	B. fl. III, 36	M. fr. VIII, 182.
D. gracilis, Chamaelaucium, F. v. M., fragm. IV, 62 (1864)	W.A.	—	—	—	—	—	B. fl. III, 36	M. fr. IV, 62; VIII, 182.
D. Forrestii, F. v. M., fragm. XI, 9 (1878)	W.A.	—	—	—	—	—	—	M. fr. XI, 9.
D. heterandra, F. v. M.; Cham., Benth., Fl. Austr. III, 36 (1866)	W.A.	—	—	—	—	—	B. fl. III, 36	
D. Drummondii, F. v. M.; Chamaelaucium, Meissner in Journ. Linn. Soc. I, 44 (1857)	W.A.	—	—	—	—	—	B. fl. III, 37	
D. Endlicheri, F. v. M.; Chamaelaucium virgatum, Endlicher in Ann. Wien. Mus. II, 193 (1838)	W.A.	—	—	—	—	—	B. fl. III, 37	
D. brevifolia, F. v. M.; Cham., Benth., Fl. Austr. III, 37 (1866)	W.A.	—	—	—	—	—	B. fl. III, 37	
D. uncinata, F. v. M.; Cham., Schauer in Pl. Preiss. I, 97 (1844)	W.A.	—	—	—	—	—	B. fl. III, 37	
D. megalopetala, Cham., F. v. M. in Benth. Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	B. fl. III, 38	M. fr. VIII, 182.
D. Turczaninowii, F. v. M.; Chamaelaucium pauciflorum, Benth., Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	B. fl. III, 38	M. fr. IX, 176; XI, 9.
D. axillaris, Cham., F. v. M. in Benth. Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	B. fl. III, 38	

VERTICORDIA, De Candolle in Dict. class. XI, 400 (1826). (Chrysorrhoea.)

V. Wilhelmii, F. v. M. in Transact. Vict. Inst. 122 (1855)	—	S.A.	—	—	—	—	B. fl. III, 19	M. fr. X, 28.
V. densiflora, Lindley in Bot. Regist. XXV, App. VI (1839)	W.A.	—	—	—	—	—	B. fl. III, 20	
V. stelluligera, Meissner in Journ. Linn. Soc. I, 38 (1857)	W.A.	—	—	—	—	—	B. fl. III, 20	
V. minutiflora, F. v. M., fragm. IV, 58 (1864)	W.A.	—	—	—	—	—	B. fl. III, 20	M. fr. IV, 38.
V. Fontainesii, De Candolle, prodr. III, 209 (1828)	W.A.	—	—	—	—	—	B. fl. III, 21	M. fr. VIII, 182; X, 28.
V. helichrysanthra, F. v. M. in Benth. Fl. Austr. III, 21 (1866)	W.A.	—	—	—	—	—	B. fl. III, 21	
V. Brownii, De Candolle, prodr. III, 209 (1828)	W.A.	—	—	—	—	—	B. fl. III, 21	
V. fastigiata, Turcz. in Bullet. de l'Ac. de St. Petersb. 402 (1852)	W.A.	—	—	—	—	—	B. fl. III, 22	M. fr. VIII, 122; X, 28.
V. Harveyi, Benth. in Journ. Linn. Soc. III, 22 (1864)	W.A.	—	—	—	—	—	B. fl. III, 22	
V. ambrilepis, Turczaninow in Bull. Mosc. XX, 158 (1847)	W.A.	—	—	—	—	—	B. fl. III, 22	M. fr. X, 28.
V. serrata, Schauer in nov. act. acad. Caes. XIX, suppl. II, 70 (1841)	W.A.	—	—	—	—	—	B. fl. III, 23	
V. nitens, Schauer in nov. act. acad. Caes. XIX, suppl. II, 71, t. 4 (1841)	W.A.	—	—	—	—	—	B. fl. III, 23	M. fr. X, 28.
V. grandiflora, Endlicher in Ann. Wien. Mus. II, 195 (1838)	W.A.	—	—	—	—	—	B. fl. III, 23	
V. chrysanthra, Endlicher in Ann. Wien. Mus. II, 195 (1838)	W.A.	—	—	—	—	—	B. fl. III, 24	M. fr. X, 28.
V. Preissii, Schauer in Lehm. pl. Preiss. I, 101 (1844)	W.A.	—	—	—	—	—	B. fl. III, 24	M. fr. VIII, 182.
V. acerosa, Lindley in Bot. Regist. XXV, App. VI (1839)	W.A.	—	—	—	—	—	B. fl. III, 25	
V. polytricha, Benth. in Journ. Linn. Soc. III, 25 (1866)	W.A.	—	—	—	—	—	B. fl. III, 25	
V. oxylepis, Turcz. in Bullet. de l'Ac. de St. Petersb. 402 (1852)	W.A.	—	—	—	—	—	B. fl. III, 25	M. fr. VIII, 182.
V. humilis, Benth. in Journ. Linn. Soc. III, 26 (1866)	W.A.	—	—	—	—	—	B. fl. III, 26	
V. penicillaris, F. v. M., fragm. I, 226 (1859)	W.A.	—	—	—	—	—	B. fl. III, 26	M. fr. I, 226.
V. multiflora, Turczaninow in Bull. Mosc. XX, 159 (1847)	W.A.	—	—	—	—	—	B. fl. III, 26	M. fr. X, 28.
V. Huegelii, Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	B. fl. III, 27	

V. insignis, Endlicher in Hueg. enum. 47 (1837) ...	W.A.	—	—	—	—	—	B.f. III, 27	M. fr. X, 23.
V. habrantha, Schauer in Lehm. pl. Preiss. I, 100 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 28	M. fr. I, 164.
V. monadelpha, Turczaninow in Bull. Mosc. XX, 158 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 28	M. fr. X, 28.
V. Lehmanni, Schauer in Lehm. pl. Preiss. I, 99 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 29	
V. Cunninghamii, Schauer in nov. act. acad. Caes. XIX, suppl. II, 55 (1841) ...	—	—	—	—	—	N.A.	B.f. III, 29	M. fr. X, 28.
V. picta, Endlicher in Ann. des Wien. Mus. II, 194 (1838) ...	W.A.	—	—	—	—	—	B.f. III, 30	M. fr. X, 28.
V. pennigera, Endlicher in Hueg. enum. 46 (1837) ...	W.A.	—	—	—	—	—	B.f. III, 30	M. fr. X, 28.
V. Drummondii, Schauer in nov. act. acad. Caes. XIX, suppl. II, 56 (1841) ...	W.A.	—	—	—	—	—	B.f. III, 31	
V. Hughani, F. v. M., fragm. XI, 10 (1878) ...	W.A.	—	—	—	—	—	—	M. fr. XI, 10.
V. pholidophylla, F. v. M., fragm. I, 227 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 31	M. fr. I, 227.
V. spicata, F. v. M., fragm. I, 226 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 32	M. fr. I, 226.
V. lepidophylla, F. v. M., fragm. I, 228 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 32	M. fr. I, 228.
V. ovalifolia, Meissner in Journ. Linn. Soc. I, 40 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 32	M. fr. X, 28.
V. chrysostachya, Meissner in Journ. Linn. Soc. I, 41 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 33	M. fr. X, 28.
V. oculata, Meissner in Journ. Linn. Soc. I, 41 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 33	
V. grandis, Drummond in Hook. Kew Misc. V, 119 (1853) ...	W.A.	—	—	—	—	—	B.f. III, 33	M. fr. X, 28; XI, 10.
PILEANTHUS, Labillardière, Nov. Holl. pl. specim. II, 11 (1806).								
P. peduncularis, Endl. in Ann. des Wien. Mus. II, 196 (1838) ...	W.A.	—	—	—	—	—	B.f. III, 34	M. fr. I, 225.
P. Limacis, Labillardière, Nov. Holl. pl. specim. II, 11, t. 149 (1806) ...	W.A.	—	—	—	—	—	B.f. III, 34	
P. filifolius, Meissner in Journ. Linn. Soc. I, 45 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 35	
CALYCOTHRIX, Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806). (Calytrix, Calythrix.)								
C. aurea, Lindley, Bot. Regist. XXV, App. V, t. 3 (1839) ...	W.A.	—	—	—	—	—	B.f. III, 41	
C. puberula, Meissner in Journ. Linn. Soc. I, 48 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 42	
C. flavescens, Cunningham in Bot. Mag. 3323 (1834) ...	W.A.	—	—	—	—	—	B.f. III, 42	
C. asperula, Schauer in Lehm. pl. Preiss. I, 106 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 42	
C. sapphirina, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	B.f. III, 43	M. fr. I, 224.
C. breviseta, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	B.f. III, 43	M. fr. VIII, 182.
C. simplex, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	B.f. III, 43	
C. empetroides, Schauer in nov. act. acad. Caes. XIX, suppl. II, 102 (1841) ...	W.A.	—	—	—	—	—	B.f. III, 44	
C. variabilis, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	B.f. III, 44	
C. muricata, F. v. M., fragm. I, 224 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 44	M. fr. I, 224.
C. gracilis, Benth. Fl. Austr. III, 45 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 45	
C. brevifolia, Meissner in Journ. Linn. Soc. I, 46 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 45	
C. brachyphylla, Turczaninow in Bull. Mosc. XX, 161 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 45	
C. Leschenaultii, Schauer in Lehm. pl. Preiss. I, 104 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 46	
C. Oldfieldii, Benth. Fl. Austr. III, 46 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 46	
C. glutinosa, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	B.f. III, 46	
C. angulata, Lindley, Bot. Regist. XXV, App. VI (1839) ...	W.A.	—	—	—	—	—	B.f. III, 47	
C. depressa, Turczaninow in Bull. Soc. Mosc. XX, 162 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 47	
C. tenuifolia, Meissner in Journ. Linn. Soc. I, 46 (1857) ...	W.A.	—	—	—	—	—	B.f. III, 47	
C. strigosa, Cunningham in Bot. Mag. 3323 (1834) ...	W.A.	—	—	—	—	—	B.f. III, 47	
C. plumulosa, F. v. M., fragm. X, 27 (1876) ...	W.A.	—	—	—	—	—	—	M. fr. X, 27.
C. decandra, R. Brown in De Cand. prodr. III, 208 (1828) ...	W.A.	—	—	—	—	—	B.f. III, 48	M. fr. I, 146.
C. Birdii, F. v. M., fragm. X, 26 (1876) ...	W.A.	—	—	—	—	—	—	M. fr. X, 26.
C. tenuiramea, Turczaninow in Bull. Mosc. XXII, 20 (1849) ...	W.A.	—	—	—	—	—	B.f. III, 48	
C. Fraseri, Cunningham in Bot. Mag. 3323 (1834) ...	W.A.	—	—	—	—	—	B.f. III, 48	
C. granulosa, Benth. Fl. Austr. III, 49 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 49	
C. Creswellii, F. v. M., fragm. X, 27 (1876) ...	W.A.	—	—	—	—	—	—	M. fr. X, 27.
C. microphylla, Cunningham in Bot. Mag. 3323 (1834) ...	—	—	—	—	—	N.A.	B.f. III, 49	
C. longiflora, F. v. M., fragm. I, 12 (1858) ...	—	S.A.	—	—	—	Q.	B.f. III, 49	M. fr. I, 12.
C. leptophylla, Benth. Fl. Austr. III, 50 (1866) ...	—	—	—	—	—	Q.	B.f. III, 50	
C. megaphylla, F. v. M., fragm. I, 146 (1859) ...	—	—	—	—	—	N.A.	B.f. III, 50	M. fr. I, 146.
C. tetragona, Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f. III, 50	M. fr. IV, 36; VIII, 183.
C. Sullivanii, F. v. M., fragm. IX, 1 (1875) ...	—	—	—	V.	—	—	—	M. fr. IX, 1.
C. conferta, Cunningham in Bot. Mag. 3323 (1834) ...	—	—	—	—	—	N.A.	B.f. III, 51	
C. arborescens, F. v. M. in Transact. phil. Inst. Vict. III, 42 (1858) ...	—	—	—	—	—	N.A.	B.f. III, 51	
C. brachychaeta, F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858) ...	—	—	—	—	—	N.A.	B.f. III, 52	
C. achaeta, F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858) ...	—	—	—	—	—	N.A.	B.f. III, 52	
C. laricina, R. Brown in Benth. Fl. Austr. III, 52 (1866) ...	—	—	—	—	—	N.A.	B.f. III, 52	
LHOTZKYA, Schauer in Linnaea X, 309 (1835).								
L. glaberrima, F. v. M., fragm. I, 13 (1858) ...	—	S.A.	—	—	—	—	B.f. III, 53	M. fr. I, 13.
L. genetylloides, F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854) ...	—	S.A.	—	V.	—	—	B.f. III, 54	
L. violacea, Lindley, Bot. Regist. XXV, App. VII (1839) ...	W.A.	—	—	—	—	—	B.f. III, 54	M. fr. XI, 8.
L. ciliata, F. v. M. in Benth. Fl. Austr. III, 54 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 54	
L. brevifolia, Schauer in Lehm. pl. Preiss. I, 103 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 54	
L. purpurea, F. v. M., fragm. I, 224 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 55	M. fr. I, 224.
L. ericoides, Schauer in Lindl. Introd. sec. ed. 439 (1835) ...	W.A.	—	—	—	—	—	B.f. III, 55	M. fr. XI, 9.
L. acutifolia, Lindley, Bot. Regist. XXV, App. VII (1839) ...	W.A.	—	—	—	—	—	B.f. III, 55	M. fr. XI, 8.
L. Harvestiana, F. v. M., fragm. XI, 8 (1878) ...	W.A.	—	—	—	—	—	—	
THRYPTOMENE, Endlicher in den Ann. des Wien. Mus. II, 192 (1838). (Homalocalyx, Paryphantha, Astraea, Eremopyxis, Micromyrtus.)								
T. mucronulata, Turczaninow in Bull. Mosc. XX, 156 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 58	
T. australis, Endlicher in den Ann. des Wien. Mus. II, 192 (1838) ...	W.A.	—	—	—	—	—	B.f. III, 58	
T. tenella, Benth. Fl. Austr. III, 59 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 59	M. fr. VIII, 182; X, 24.
T. prolifera, Turczaninow in Bull. Mosc. XXXV, 324 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 59	

T. saxicola, Schauer in Lehm. pl. Preiss. I, 102 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 59	M.fr. IV, 75.
T. Johnsonii, F. v. M., fragm. IV, 77 (1864) ...	W.A.	—	—	—	—	—	B.f. III, 60	M.fr. IV, 77.
T. racemulosa, Turczaninow in Bull. Mosc. XX, 156 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 60	
T. denticulata, Benth. in Fl. Austr. III, 60 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 60	M.fr. IV, 75.
T. baeckeacea, F. v. M., fragm. IV, 65 (1864) ...	W.A.	—	—	—	—	—	B.f. III, 60	M.fr. IV, 65.
T. strongylophylla, F. v. M. in Benth. Fl. Austr. III, 61 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 61	
T. hyporhysis, Turczaninow in Bull. Mosc. XXXV, 324 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 61	
T. Maisonneuvii, F. v. M., fragm. IV, 64 (1864) ...	—	S.A.	—	—	—	N.A.	B.f. III, 61	M.fr. IV, 64.
T. urceolaris, F. v. M., fragm. X, 25 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 25.
T. Mitchelliana, F. v. M., fragm. I, 11 (1858) ...	—	S.A.	—	V.	—	—	B.f. III, 61	M.fr. I, 11; IX, 62.
T. Miqueliana, F. v. M., fragm. I, 11 (1858) ...	—	S.A.	—	—	N.S.W.	—	B.f. III, 62	M.fr. I, 11.
T. micrantha, J. Hooker in Hook. Kew Misc. V, 299, t. 8 (1853) ...	—	—	T.	—	—	—	B.f. III, 62	
T. stenocalyx, F. v. M., fragm. X, 23 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 23.
T. ericaea, F. v. M., fragm. I, 12 (1858) ...	—	S.A.	—	—	—	—	B.f. III, 62	M.fr. I, 12.
T. oligandra, F. v. M., fragm. I, 11 (1858) ...	W.A.	—	—	—	—	Q.	N.A.	B.f. III, 63
T. elobata, F. v. M., fragm. IV, 63 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 64
T. racemosa, F. v. M. in Benth. Fl. Austr. III, 64 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 64
T. trachycalyx, F. v. M., fragm. X, 25 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 25.
T. imbricata, F. v. M.; Micromyrtus, Benth., Fl. Aus. III, 64 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 64
T. flaviflora, F. v. M., fragm. VIII, 13 (1873) ...	W.A.	S.A.	—	—	—	—	—	M.fr. X, 25.
T. Drummondii, F. v. M.; Microm., Benth., Fl. Aus. III, 64 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 64
T. Elliotii, F. v. M., fragm. IX, 62 (1875) ...	—	S.A.	—	—	—	—	—	M.fr. VIII, 183.
T. auriculata, F. v. M., fragm. X, 24 (1876) ...	—	S.A.	—	—	—	—	—	M.fr. IX, 62.
T. hymenonema, F. v. M., fragm. X, 26 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 24.
T. ciliata, F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880) ...	—	S.A.	—	V.	N.S.W.	Q.	—	M.fr. I, 30 & 242; IV, 63.
T. minutiflora, F. v. M. in Woolls, pl. of the neighb. of Syd. 23 (1880) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 65
T. leptocalyx, F. v. M.; Microm., Benth., Fl. Austr. III, 65 (1866) ...	—	—	—	—	—	Q.	—	B.f. III, 65
T. homalocalyx, F. v. M., fragm. IV, 63 (1864) ...	—	—	—	—	—	—	N.A.	B.f. III, 66
T. polyandra, F. v. M., fragm. IV, 77 (1864) ...	—	—	—	—	—	—	N.A.	B.f. III, 56
WEHLIA, F. v. M., fragm. X, 22 (1876).								
W. thryptomenoides, F. v. M., fragm. X, 22 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 22.
W. coarctata, F. v. M., fragm. X, 23 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 23.
BAECKEA, Linné, spec. plant. 353 (1753). (Jungia, Imbricaria, Scholtzia, Schidiomyrtus, Rinzia, Euryomyrtus, Camphoromyrtus, Tetrapora, Harmogia, Oxymyrrhine, Babingtonia, Ericomyrtus, Piptandra, Anticoryne.)								
B. platystemonea, Benth. in Fl. Austr. III, 74 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 74
B. Fumana, F. v. M., fragm. IV, 68 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 74
B. dimorphandra, F. v. M. in Benth. Fl. Austr. III, 74 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 74
B. schollerifolia, Lehmann in Lehm. pl. Preiss. II, 369 (1847) ...	W.A.	—	—	—	—	—	—	B.f. III, 75
B. oxycoccoides, Benth. in Fl. Austr. III, 75 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 75
B. Drummondii, Benth. in Fl. Austr. III, 75 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 75
B. diffusa, Sieber in De Cand. prodr. III, 230 (1828) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. III, 76
B. crassifolia, Lindley in Mitch. Three Exped. II, 115 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. III, 76
B. cryptandroides, F. v. M., fragm. X, 29 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. IV, 66.
B. tetragona, F. v. M. in Benth. Fl. Austr. III, 77 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 77
B. ericaea, F. v. M., fragm. I, 31 (1858) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f. III, 77
B. polystemonea, F. v. M., fragm. II, 124 (1861) ...	—	S.A.	—	—	—	—	—	B.f. III, 77
B. crenulata, R. Brown in Flind. voy. 548 (1814) ...	—	—	—	—	N.S.W.	Q.	—	B.f. III, 78
B. brevifolia, De Candolle, prodr. III, 230 (1828) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 78
B. Gunniana, Schauer in Walp. Rep. II, 920 (1843) ...	—	—	T.	V.	N.S.W.	—	—	B.f. III, 78
B. diosmifolia, Rudge in Transact. Linn. Soc. VIII, 298, t. 13 (1807) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 79
B. leptocaulis, J. Hooker, icon. pl. t. 298 (1840) ...	—	—	T.	—	—	—	—	B.f. III, 79
B. arbuscula, R. Brown in Benth. Fl. Austr. III, 79 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 79
B. astarteoides, Benth. in Fl. Austr. III, 80 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 80
B. linifolia, Rudge in Transact. Linn. Soc. VIII, 297, t. 12 (1807) ...	—	—	—	V.	N.S.W.	—	—	B.f. III, 80
B. stenophylla, F. v. M., fragm. I, 13 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.f. III, 80
B. camphorata, R. Brown in Bot. Mag. t. 2694 (1826) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 81
B. virgata, Andrews, Bot. Reposit. t. 598 (1810) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. III, 81
B. crenatifolia, F. v. M., fragm. IV, 70 (1864) ...	—	—	—	V.	—	—	—	B.f. III, 82
B. Cunninghamii, Benth. in Fl. Austr. III, 82 (1866) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 82
B. densifolia, Smith in Transact. Linn. Soc. III, 260 (1797) ...	—	—	—	—	N.S.W.	—	—	B.f. III, 82
B. Behrii, F. v. M., fragm. IV, 68 (1864) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. III, 83
B. uncinella, Benth. in Fl. Austr. III, 84 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 84
B. polyandra, F. v. M., fragm. IV, 12 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 84
B. corynophylla, F. v. M., fragm. IV, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 84
B. pachyphylla, Benth. in Fl. Austr. III, 85 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 85
B. crispiflora, F. v. M., fragm. IV, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 85
B. Blackettii, F. v. M., fragm. VIII, 181 (1874) ...	W.A.	—	—	—	—	—	—	M.fr. II, 31; IV, 72.
B. camphorosmae, Endlicher in Hueg. enum. 51 (1837) ...	W.A.	—	—	—	—	—	—	M.fr. VIII, 181.
B. pulchella, De Candolle, prodr. III, 230 (1828) ...	W.A.	—	—	—	—	—	—	—
B. ochropetala, F. v. M., fragm. X, 29 (1876) ...	W.A.	—	—	—	—	—	—	—
B. pygmaea, R. Brown in Benth. Fl. Austr. III, 86 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 86
B. corymbulosa, Benth. in Fl. Austr. III, 87 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 87
B. floribunda, Benth. in Fl. Austr. III, 87 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 87
B. pentandra, F. v. M., fragm. II, 31; IV, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 87
B. pentagonantha, F. v. M., fragm. IV, 73 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 87
B. robusta, F. v. M., fragm. IV, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 88
B. ovalifolia, F. v. M., fragm. IV, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 88
B. subcuneata, F. v. M., fragm. IV, 73 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 88
B. grandiflora, Benth. in Fl. Austr. III, 89 (1866) ...	W.A.	—	—	—	—	—	—	B.f. III, 89
B. uberiflora; Scholtzia, F. v. M., fragm. IV, 74 (1864) ...	W.A.	—	—	—	—	—	—	B.f. III, 67

[183.

M.fr. II, 31; IV, 72; VIII,

M.fr. IV, 73.

M.fr. IV, 72.

M.fr. II, 32; IV, 72; VIII,

M.fr. IV, 73.

M.fr. IV, 74.

B. involucreta, Endlicher in Hueg. enum. 51 (1837) ...	W.A.	—	—	—	—	—	B.f.III,68	M.fr.IV,74.
B. spathulata, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 68 (1866) ...	W.A.	—	—	—	—	—	B.f.III,68	
B. ciliata; Scholtzia, F. v. M., fragm. IV, 76 (1864) ...	W.A.	—	—	—	—	—	B.f.III,68	M.fr.IV,76.
B. capitata; Scholtzia, F. v. M. in Benth. Fl. Austr. III, 69 (1866) ...	W.A.	—	—	—	—	—	B.f.III,69	
B. umbellifera; Scholtzia, F. v. M., fragm. IV, 75 (1864) ...	W.A.	—	—	—	—	—	B.f.III,69	M.fr.IV,75.
B. laxiflora, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 69 (1866) ...	W.A.	—	—	—	—	—	B.f.III,69	
B. leptantha, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 69 (1866) ...	W.A.	—	—	—	—	—	B.f.III,69	
B. parviflora; Scholtzia, F. v. M., fragm. IV, 76 (1864) ...	W.A.	—	—	—	—	—	B.f.III,70	M.fr.IV,76.
B. oligandra; Scholtzia, F. v. M. in Benth. Fl. Austr. III, 70 (1866) ...	W.A.	—	—	—	—	—	B.f.III,70	
B. serpillifolia, F. v. M., fragm. X, 30 (1876) ...	W.A.	—	—	—	—	—	B.f.III,70	M.fr.VIII,183.
B. teretifolia, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 70 (1866) ...	W.A.	—	—	—	—	—	B.f.III,70	
ASTARTEA, De Candolle in Dict. class. XI, 400 (1826).								
A. ambigua, F. v. M., fragm. II, 32 (1860) ...	W.A.	—	—	—	—	—	B.f.III,89	M.fr.II,32.
A. fascicularis, De Candolle, prodr. III, 210 (1828) ...	W.A.	—	—	—	—	—	B.f.III,90	M.fr.VIII,183.
A. intratropica, F. v. M., fragm. I, 83 (1858) ...	—	—	—	—	—	N.A.	B.f.III,90	M.fr.I,83.
HYPOCALYMYMA, Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 51 (1837).								
H. xanthopetalum, F. v. M., fragm. II, 29 (1860) ...	W.A.	—	—	—	—	—	B.f.III,92	M.fr.II,29.
H. robustum, Endlicher in Hueg. enum. 50 (1837) ...	W.A.	—	—	—	—	—	B.f.III,92	
H. longifolium, F. v. M., fragm. II, 23 (1860) ...	W.A.	—	—	—	—	—	B.f.III,92	M.fr.II,28.
H. strictum, Schauer in Lehm. pl. Preiss. I, 111 (1844) ...	W.A.	—	—	—	—	—	B.f.III,93	
H. tetrapterum, Turczaninow in Bull. Mosc. XXXV, 325 (1862) ...	W.A.	—	—	—	—	—	B.f.III,93	
H. linifolium, Turczaninow in Bull. Mosc. XXXV, 325 (1862) ...	W.A.	—	—	—	—	—	B.f.III,93	
H. angustifolium, Endlicher in Hueg. enum. 50 (1837) ...	W.A.	—	—	—	—	—	B.f.III,94	
H. ericifolium, Benth., Fl. Austr. III, 94 (1866) ...	W.A.	—	—	—	—	—	B.f.III,94	
H. cordifolium, Lehmann, plant. Preiss. I, 112 (1844) ...	W.A.	—	—	—	—	—	B.f.III,94	
H. speciosum, Turcz. in Bull. de l'Acad. de St. Pet. 409 (1852) ...	W.A.	—	—	—	—	—	B.f.III,95	M.fr.VIII,183.
H. Phillipsii, Harvey in nat. hist. Rev. V, 296, t. 22 (1865) ...	W.A.	—	—	—	—	—	B.f.III,95	
H. myrtifolium, Turcz. in Bullet. de St. Petersb. 410 (1852) ...	W.A.	—	—	—	—	—	B.f.III,95	M.fr.VIII,183.
BALAUSTION, Hooker, icones plantarum 852 (1852). (Punicella, Cheynia.)								
B. pulcherrimum, Hooker, icon. plant. 852 (1852) ...	W.A.	—	—	—	—	—	B.f.III,96	M.fr.VIII,183.
AGONIS, Lindley, Bot. Regist. XXV, p. X (1839). (Billiotia, Billotia.)								
A. spathulata, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	B.f.III,97	M.fr.XI,120.
A. floribunda, Turczaninow in Bull. Mosc. II, 20 (1849) ...	W.A.	—	—	—	—	—	B.f.III,97	
A. marginata, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	B.f.III,98	M.fr.XI,120.
A. linearifolia, Schauer in Lehm. pl. Preiss. I, 118 (1844) ...	W.A.	—	—	—	—	—	B.f.III,98	M.fr.VIII,183;XI,120.
A. juperina, Schauer in Lehm. pl. Preiss. I, 118 (1844) ...	W.A.	—	—	—	—	—	B.f.III,98	M.fr.XI,120.
A. parviceps, Schauer in Lehm. pl. Preiss. I, 119 (1844) ...	W.A.	—	—	—	—	—	B.f.III,99	M.fr.XI,120.
A. obtusissima, F. v. M., fragm. XI, 119 (1881) ...	W.A.	—	—	—	—	—	—	M.fr.XI,119.
A. flexuosa, Schauer in Lehm. pl. Preiss. I, 116 (1844) ...	W.A.	—	—	—	—	—	B.f.III,99	M.fr.XI,120.
A. undulata, Benth., Fl. Austr. III, 100 (1866) ...	W.A.	—	—	—	—	—	B.f.III,100	
A. hypericifolia, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	B.f.III,100	M.fr.XI,120.
A. Scortechiniana, F. v. M., fragm. XI, 118 (1881) ...	—	—	—	—	—	Q.	—	M.fr.XI,118.
A. grandiflora, Benth., Fl. Austr. III, 100 (1866) ...	W.A.	—	—	—	—	—	B.f.III,100	M.fr.XI,120.
LEPTOSPERMUM, R. et G. Forster, char. gen. 71, t. 36 (1776). (Fabricia, Homalospermum, Pericalymma.)								
L. Fabricia, Benth., Fl. Austr. III, 102 (1866) ...	—	—	—	—	—	Q.	B.f.III,102	
L. laevigatum, F. v. M., Annual Report 22 (1858) ...	—	S.A.	T.	V.	N.S.W.	—	B.f.III,103	M.fr.IV,60.
L. firmum, Benth., Fl. Austr. III, 104 (1866) ...	W.A.	—	—	—	—	—	B.f.III,104	
L. flavescens, Smith in Transact. Linn. Soc. III, 262 (1797) ...	—	—	T.	V.	N.S.W.	Q.	B.f.III,104	
L. scoparium, R. & G. Forster, char. gen. 48 (1776) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.III,105	
L. arachnoideum, Smith in Transact. Linn. Soc. III, 263 (1797) ...	—	—	—	—	N.S.W.	—	B.f.III,105	
L. spinescens, Endlicher in Hueg. enum. 51 (1837) ...	W.A.	—	—	—	—	—	B.f.III,106	
L. lanigerum, Smith in Transact. Linn. Soc. III, 263 (1797) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.III,106	
L. parvifolium, Smith in Transact. Linn. Soc. III, 263 (1797) ...	—	—	—	—	N.S.W.	—	B.f.III,107	
L. stellatum, Cavanilles, icon. pl. IV, 16, t. 330 ...	—	—	—	—	N.S.W.	Q.	B.f.III,107	
L. attenuatum, Smith in Transact. Linn. Soc. III, 262 (1797) ...	—	—	—	V.	N.S.W.	Q.	B.f.III,108	
L. myrtifolium, Sieber in De Cand., prodr. III, 238 (1828) ...	—	—	T.	V.	N.S.W.	Q.	B.f.III,108	
L. rupestre, J. Hooker in Hook. icon. pl. t. 308 (1841) ...	—	—	T.	—	—	—	B.f.III,108	
L. myrsinoides, Schlechtendal in Linnaea XX, 653 (1847) ...	—	S.A.	—	V.	N.S.W.	—	B.f.III,109	
L. erubescens, Schauer in Lehm. pl. Preiss. I, 121 (1844) ...	W.A.	—	—	—	—	—	B.f.III,109	M.fr.VIII,183.
L. abnorme, F. v. M. in Benth. Fl. Austr. III, 109 (1866) ...	—	—	—	—	N.S.W.	Q.	B.f.III,109	M.fr.II,27.
L. Roei, Benth., Fl. Austr. III, 110 (1866) ...	W.A.	—	—	—	—	—	B.f.III,110	
L. floridum, Benth., Fl. Austr. III, 110 (1866) ...	W.A.	—	—	—	—	—	B.f.III,110	
L. ellipticum, Endlicher in Hueg. enum. 51 (1837) ...	W.A.	—	—	—	—	—	B.f.III,110	
L. crassipes, Lehmann, ind. sem. hort. Hamb. (1842) ...	W.A.	—	—	—	—	—	B.f.III,110	
KUNZEA, Reichenbach, conspect. regn. veget. 175 (1828). (Salisia, Pentagonaster.)								
K. micrantha, Schauer in Lehm. pl. Preiss. I, 125 (1844) ...	W.A.	—	—	—	—	—	B.f.III,112	M.fr.VIII,183.
K. eriocalyx, F. v. M., fragm. II, 28 (1860) ...	W.A.	—	—	—	—	—	B.f.III,112	M.fr.II,28.
K. Muelleri, Benth., Fl. Austr. III, 113 (1866) ...	—	—	—	V.	N.S.W.	—	B.f.III,113	
K. ericifolia, Reichenbach, consp. 175 (1828) ...	W.A.	—	—	—	—	—	B.f.III,113	
K. Preissiana, Schauer in Lehm. pl. Preiss. I, 125 (1844) ...	W.A.	—	—	—	—	—	B.f.III,113	
K. recurva, Schauer in Lehm. pl. Preiss. I, 125 (1844) ...	W.A.	—	—	—	—	—	B.f.III,114	M.fr.VIII,183.
K. micromera, Schauer in Lehm. pl. Preiss. III, 223 (1847) ...	W.A.	—	—	—	—	—	B.f.III,114	
K. pauciflora, Schauer in Lehm. pl. Preiss. I, 124 (1844) ...	W.A.	—	—	—	—	—	B.f.III,114	M.fr.VIII,183.
K. calida, F. v. M., fragm. VI, 23 (1867) ...	—	—	—	—	—	Q.	—	M.fr.VI,23.
K. parvifolia, Schauer in Lehm. pl. Preiss. I, 124 (1844) ...	—	—	—	—	N.S.W.	—	B.f.III,115	M.fr.VIII,13.
K. peduncularis, F. v. M. in Transact. Vict. Inst. 144 (1855) ...	—	—	—	V.	N.S.W.	—	B.f.III,115	M.fr.X,55.
K. corifolia, Reichenbach, consp. 175 (1828) ...	—	—	—	T.	V.	N.S.W.	B.f.III,115	
K. capitata, Reichenbach, consp. 175 (1828) ...	—	—	—	—	N.S.W.	—	B.f.III,116	
K. pomifera, F. v. M. in Transact. Vict. Inst. 124 (1855) ...	—	S.A.	—	V.	—	—	B.f.III,116	

K. sericea, Turczaninow in Bull. Mosc. XX, 162 (1847)...	W.A.	—	—	—	—	—	B.f. III, 117	
K. Baxteri, Schauer in Lehm. pl. Preiss. I, 123 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 117	M.fr. III, 153.
K. opposita, F. v. M., fragm. VI, 24 (1867) ...	—	—	—	N.S.W.	—	—	—	M.fr. VI, 24.
CALLISTEMON, R. Brown in Flind. voy. II, 547 (1814).								
C. speciosus, De Candolle, prodr. III, 224 (1828)...	W.A.	—	—	—	—	—	B.f. III, 119	M.fr. I, 14.
C. phoeniceus, Lindley, Bot. Regist. XXV, App. X (1839) ...	W.A.	—	—	—	—	—	B.f. III, 119	M.fr. IV, 53.
C. lanceolatus, De Candolle, prodr. III, 224 (1828) ...	—	—	—	V. N.S.W.	Q.	—	B.f. III, 120	M.fr. IV, 53.
C. coccineus, F. v. M., fragm. I, 13 (1858)...	—	—	—	S.A.	V. N.S.W.	—	B.f. III, 120	M.fr. I, 13.
C. salignus, De Candolle, prodr. III, 223 (1828) ...	—	—	—	S.A.	T. V. N.S.W.	Q.	B.f. III, 120	M.fr. IV, 54.
C. paludosus, F. v. M., fragm. I, 14 (1858) ...	—	—	—	—	—	Q.	—	M.fr. I, 14.
C. rigidus, R. Brown in Bot. Regist. t. 393 (1819) ...	—	—	—	—	N.S.W.	—	B.f. III, 121	
C. linearis, De Candolle, prodr. III, 223 (1828) ...	—	—	—	—	N.S.W.	—	B.f. III, 122	
C. pinifolius, De Candolle, prodr. III, 223 (1828)...	—	—	—	—	N.S.W.	—	B.f. III, 122	
C. teretifolius, F. v. M. in Schlecht. Linnaea XXV, 387 (1852) ...	—	S.A.	—	—	—	—	B.f. III, 122	
C. brachyandrus, Lindley in Journ. Hort. Soc. IV, 112 (1849) ...	—	S.A.	—	V. N.S.W.	—	—	B.f. III, 122	M.fr. IV, 52.
MELALEUCA, Linné, mantiss. plant. 14 (1767). (Gymnagathis, Asteromyrtus.)								
M. macronychia, Turcz. in Bullet. de St. Petersb. 420 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 129	M.fr. VIII, 183.
M. lateritia, Otto in allem. Gart. Zeit. II, 257 (1834) ...	W.A.	—	—	—	—	—	B.f. III, 130	
M. calothamnoides, F. v. M., fragm. III, 114 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 130	M.fr. III, 114.
M. blairiaefolia, Turczaninow in Bull. Mosc. XX, 165 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 130	
M. diosmifolia, Andrews, Bot. Reposit. t. 476 (1808) ...	W.A.	—	—	—	—	—	B.f. III, 130	
M. elliptica, Labillardière, Nov. Holl. pl. spec. II, 31, t. 173 (1806) ...	W.A.	—	—	—	—	—	B.f. III, 131	M.fr. VIII, 183.
M. hypericifolia, Smith in Transact. Linn. Soc. III, 279 (1797)...	—	—	—	N.S.W.	—	—	B.f. III, 131	
M. fulgens, R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812) ...	W.A.	—	—	—	—	—	B.f. III, 131	
M. acuminata, F. v. M., fragm. I, 15 (1858) ...	W.A.	S.A.	—	V. N.S.W.	—	—	B.f. III, 132	M.fr. I, 15.
M. leptoclada, Benth. Fl. Austr. III, 132 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 132	
M. basiculiflora, Benth. Fl. Austr. III, 133 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 133	
M. gibbosa, Labillardière, Nov. Holl. pl. spec. II, 30, t. 172 (1806) ...	—	S.A.	T. V.	—	—	—	B.f. III, 133	
M. decussata, R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812) ...	—	S.A.	—	V.	—	—	B.f. III, 133	
M. Wilsonii, F. v. M., fragm. II, 124, t. 15 (1861) ...	—	S.A.	—	V.	—	—	B.f. III, 134	M.fr. II, 124.
M. thymifolia, Smith in Transact. Linn. Soc. III, 278 (1797) ...	—	—	—	N.S.W.	Q.	—	B.f. III, 134	M.fr. X, 55.
M. violacea, Lindley, Bot. Regist. XXV, App. VIII (1839) ...	W.A.	—	—	—	—	—	B.f. III, 135	
M. cardiophylla, F. v. M., fragm. I, 225 (1859) ...	W.A.	—	—	—	—	—	B.f. III, 135	M.fr. I, 225; III, 164.
M. undulata, Benth. Fl. Austr. III, 135 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 135	
M. depauperata, Turcz. in Bullet. de St. Petersb. 424 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 136	M.fr. III, 120; VIII, 183.
M. lateriflora, Benth. Fl. Austr. III, 136 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 136	
M. exarata, F. v. M., fragm. III, 114 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 136	M.fr. III, 114; VIII, 184.
M. fasciculiflora, Benth. Fl. Austr. III, 137 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 137	M.fr. VIII, 184.
M. teretifolia, Endlicher in Hueg. enum. 49 (1837) ...	W.A.	—	—	—	—	—	B.f. III, 137	M.fr. III, 117.
M. alsophila, Cunningham in Benth. Fl. Austr. III, 137 (1866)...	—	—	—	—	—	N.A.	B.f. III, 137	
M. acacioides, F. v. M., fragm. III, 116 (1862) ...	—	—	—	—	—	N.A.	B.f. III, 138	M.fr. III, 116.
M. Baxteri, Benth. Fl. Austr. III, 138 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 138	
M. symphyocarpa, F. v. M. in Transact. phil. Inst. Vict. III, 44 (1858) ...	—	—	—	—	—	N.A.	B.f. III, 138	
M. angustifolia, Gaertner de fruct. I, 172, t. 35 (1788) ...	—	—	—	—	—	Q.	B.f. III, 139	
M. pauciflora, Turczaninow in Bull. Mosc. XX, 166 (1847) ...	—	—	—	N.S.W.	—	—	B.f. III, 139	
M. squarrosa, Donn, Hort. Cantabr. ed. sec. 101 (1800) ...	—	S.A.	T. V.	N.S.W.	—	—	B.f. III, 139	M.fr. X, 55.
M. adnata, Turcz. in Bullet. de St. Petersb. 425 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 140	M.fr. III, 117; VIII, 184.
M. linariifolia, Smith in Transact. Linn. Soc. III, 278 (1797) ...	—	—	—	N.S.W.	Q.	—	B.f. III, 140	M.fr. X, 55.
M. trichostachya, Lindley in Mitch. Trop. Austr. 277 (1848) ...	—	S.A.	—	—	Q.	N.A.	—	
M. radula, Lindley, Bot. Regist. XXV, App. VIII (1839) ...	W.A.	—	—	—	—	—	B.f. III, 141	M.fr. X, 55.
M. pulchella, R. Brown in Ait. hort. Kew. sec. ed. IV, 414 ...	W.A.	—	—	—	—	—	B.f. III, 141	
M. conferta, Benth. Fl. Austr. III, 142 (1866)...	W.A.	—	—	—	—	—	B.f. III, 142	
M. Leucadendra, Linné, mantiss. 105 (1767) ...	W.A.	—	—	N.S.W.	Q.	N.A.	B.f. III, 142	
M. lasiandra, F. v. M., fragm. III, 115 (1862) ...	—	—	—	—	—	N.A.	B.f. III, 143	M.fr. III, 115.
M. genistifolia, Smith in Transact. Linn. Soc. III, 277 (1797) ...	—	—	—	N.S.W.	Q.	N.A.	B.f. III, 143	M.fr. I, 15; X, 55.
M. styphelioides, Smith in Transact. Linn. Soc. III, 275 (1797) ...	—	—	—	N.S.W.	Q.	—	B.f. III, 144	
M. Huegelii, Endlicher in Hueg. enum. 48 (1837) ...	W.A.	—	—	—	—	—	B.f. III, 144	
M. dissitiflora, F. v. M., fragm. III, 153 (1863) ...	—	—	—	—	—	N.A.	B.f. III, 144	M.fr. III, 153.
M. linophylla, F. v. M., fragm. III, 115 (1863) ...	—	—	—	—	—	N.A.	B.f. III, 145	M.fr. III, 115.
M. parviflora, Lindley, Bot. Regist. XXV, p. VIII (1839) ...	W.A.	S.A.	—	V. N.S.W.	Q.	—	B.f. III, 145	
M. laxiflora, Turczaninow in Bullet. de St. Petersb. 421 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 145	M.fr. VIII, 184; X, 55.
M. armillaris, Smith in Transact. Linn. Soc. III, 277 (1797) ...	—	—	—	V. N.S.W.	—	—	B.f. III, 146	
M. hamulosa, Turczaninow in Bull. Mosc. XX, 165 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 146	
M. subfalcata, Turczaninow in Bullet. de St. Petersb. 422 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 146	M.fr. III, 119; VIII, 184.
M. glaberrima, F. v. M., fragm. III, 119 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 147	M.fr. III, 119.
M. rhaphiophylla, Schauer in Lehm. pl. Preiss. I, 143 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 147	
M. cymbifolia, Benth. Fl. Austr. III, 148 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 148	
M. cuticularis, Labillard., Nov. Holl. pl. spec. II, 30, t. 171 (1806) ...	W.A.	—	—	—	—	—	B.f. III, 148	
M. sparsiflora, Turczaninow in Bull. Mosc. XX, 167 (1847) ...	W.A.	—	—	—	—	—	B.f. III, 148	
M. calycina, R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812) ...	W.A.	—	—	—	—	—	B.f. III, 148	M.fr. VIII, 184.
M. cordata, Turcz. in Bullet. de St. Petersb. 418 (1852)...	W.A.	—	—	—	—	—	B.f. III, 149	M.fr. VIII, 184.
M. globifera, R. Brown in Ait. hort. Kew. sec. ed. IV, 411 (1812) ...	W.A.	—	—	—	—	—	B.f. III, 149	
M. megacephala, F. v. M., fragm. III, 117 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 149	M.fr. III, 117.
M. nesophila, F. v. M., fragm. III, 113 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 150	M.fr. III, 150.
M. Oldfieldii, F. v. M., fragm. III, 118 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 150	
M. uncinata, R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812) ...	W.A.	S.A.	—	V. N.S.W.	—	—	B.f. III, 150	
M. concreta, F. v. M., fragm. III, 118 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 151	M.fr. III, 118.
M. filifolia, F. v. M., fragm. III, 119 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 151	M.fr. III, 119.
M. hakeoides, F. v. M. in Benth. Fl. Austr. III, 151 (1866) ...	—	S.A.	—	—	N.S.W.	—	B.f. III, 151	
M. glomerata, F. v. M., Rep. Babb. Exped. 8 (1858) ...	W.A.	S.A.	—	—	—	N.A.	B.f. III, 151	

M. pentagona, Labillardière, Nov. Holl. pl. spec. II, 27, t. 166 (1806) W.A.	—	—	—	—	—	—	B.f. III, 152	M.fr. VIII, 184.
M. ciliosa, Turczaninow in Bull. Soc. Mosc. XXXV, 326 (1862) W.A.	—	—	—	—	—	—	B.f. III, 152	
M. polycephala, Benth. Fl. Austr. III, 152 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 152	M.fr. VIII, 184.
M. spatulata, Schauer in Lehm. pl. Preiss. I, 134 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 153	M.fr. VIII, 184.
M. eriantha, Benth. Fl. Austr. III, 153 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 153	
M. subtrigona, Schauer in Lehm. pl. Preiss. I, 139 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 153	
M. seriata, Lindley, Bot. Regist. XXV, App. VIII (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 154	
M. scabra, R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812) W.A.	—	—	—	—	—	—	B.f. III, 154	M.fr. VIII, 184.
M. urceolaris, F. v. M. in Benth. Fl. Austr. III, 154 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 154	
M. trichophylla, Lindley, Bot. Regist. XXV, App. VIII (1839) W.A.	—	—	—	—	—	—	B.f. III, 155	M.fr. III, 114.
M. holosericea, Schauer in Lehm. pl. Preiss. I, 139 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 155	
M. squamea, Labillardière, Nov. Holl. pl. spec. II, 23, t. 168 (1806) — S.A. T. V. N.S.W.	—	—	—	—	—	—	B.f. III, 155	
M. densa, R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812) ... W.A.	—	—	—	—	—	—	B.f. III, 156	M.fr. VIII, 184.
M. thymoides, Labillardière, Nov. Holl. pl. spec. II, 27, t. 167 (1806) W.A.	—	—	—	—	—	—	B.f. III, 156	
M. striata, Labillardière, Nov. Holl. pl. spec. II, 26, t. 165 (1806) W.A.	—	—	—	—	—	—	B.f. III, 157	
M. polygaloides, Schauer in Lehm. pl. Preiss. I, 142 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 157	
M. incana, R. Brown in Bot. Regist. t. 410 (1819) ... W.A.	—	—	—	—	—	—	B.f. III, 157	
M. nodosa, Smith in Transact. Linn. Soc. III, 276 (1797) ... — N.S.W. Q.	—	—	—	—	—	—	B.f. III, 158	M.fr. VIII, 184.
M. pungens, Schauer in Lehm. pl. Preiss. I, 138 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 158	
M. ericifolia, Smith in Transact. Linn. Soc. III, 276 (1797) ... — S.A. T. V. N.S.W. Q.	—	—	—	—	—	—	B.f. III, 159	
M. pustulata, J. Hooker in Lond. Journ. VI, 476 (1844) ... — S.A. T. V. N.S.W.	—	—	—	—	—	—	B.f. III, 160	
M. viminea, Lindley, Bot. Regist. XXV, App. VIII (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 159	M.fr. III, 116.
M. microphylla, Smith in Rees, Cyclop. XXIII (1812) ... W.A.	—	—	—	—	—	—	B.f. III, 159	
M. tenella, Benth. Fl. Austr. III, 160 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 160	
M. leiocarpa, F. v. M., fragm. X, 55 (1876) ... W.A.	—	—	—	—	—	—	B.f. III, 160	M.fr. X, 55.
M. leiopyxis, F. v. M. in Benth. Fl. Austr. III, 160 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 160	
M. acerosa, Schauer in Lehm. pl. Preiss. I, 137 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 161	M.fr. III, 118.
M. pauperiflora, F. v. M., fragm. III, 116 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 161	M.fr. III, 116.
M. aspalathoides, Schauer in Lehm. pl. Preiss. I, 140 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 161	
M. cucullata, Turczaninow in Bullet. de St. Petersb. 424 (1852) W.A.	—	—	—	—	—	—	B.f. III, 161	M.fr. VIII, 184.
M. minutifolia, F. v. M. in Transact. phil. Soc. Vict. III, 45 (1858) — Q. N.A.	—	—	—	—	—	—	B.f. III, 162	
M. foliolosa, Cunningham in Benth. Fl. Austr. III, 162 (1866) ... — Q.	—	—	—	—	—	—	B.f. III, 162	
M. micromera, Schauer in Lehm. pl. Preiss. I, 146 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 162	
M. thyoides, Turczaninow in Bull. Soc. Mosc. XX, 167 (1847) ... W.A.	—	—	—	—	—	—	B.f. III, 162	M.fr. III, 114.
M. tamariscina, Hooker in Mitch. Trop. Austr. 262 (1848) ... — Q.	—	—	—	—	—	—	B.f. III, 163	
CONOTHAMNUS, Lindley, Bot. Regist. XXV, App. IX (1839). (Trichobasis.)								
C. trinervis, Lindley, Bot. Regist. XXV, App. IX (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 163	
C. divaricatus, Benth. Fl. Austr. III, 164 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 164	M.fr. VIII, 184.
BEAUFORTIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812). (Schizopleura.)								
B. sparsa, R. Brown in Ait. hort. Kew. sec. ed. IV, 419 (1812) W.A.	—	—	—	—	—	—	B.f. III, 165	
B. decussata, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812) W.A.	—	—	—	—	—	—	B.f. III, 166	
B. squarrosa, Schauer in nov. act. acad. Caes. XXI, 11 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 166	
B. orbifolia, F. v. M., fragm. III, 110 (1863) ... W.A.	—	—	—	—	—	—	B.f. III, 166	M.fr. III, 110.
B. anisandra, Schauer in nov. act. acad. Caes. XXI, 13, t. I, (1844) W.A.	—	—	—	—	—	—	B.f. III, 167	
B. macrostemon, Lindley, Bot. Regist. XXV, App. X (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 167	
B. heterophylla, Turczaninow in Bullet. de St. Petersb. 427 (1852) W.A.	—	—	—	—	—	—	B.f. III, 167	M.fr. VIII, 184.
B. Schaueri, Preiss in nov. act. acad. Caes. XXI, 18 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 168	
B. purpurea, Lindley, Bot. Regist. XXV, App. X, t. 3 (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 168	
B. Dampieri, Cunningham in Bot. Mag. t. 3272 (1833) ... W.A.	—	—	—	—	—	—	B.f. III, 168	
B. elegans, Schauer in nov. act. acad. Caes. XXI, 20 (1844) ... W.A.	—	—	—	—	—	N.A.	B.f. III, 169	
B. interstans, F. v. M., fragm. X, 30 (1876) ... W.A.	—	—	—	—	—	—	—	M.fr. X, 30.
B. micrantha, Schauer in nov. act. acad. Caes. XXI, 22 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 169	M.fr. VIII, 184.
REGELIA, Schauer in Linnaea XVII, 243 (1843).								
R. grandiflora, Benth. Fl. Austr. III, 170 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 170	M.fr. VIII, 184; IX, 46; XI, [11.
R. ciliata, Schauer in nov. act. acad. Caes. XXI, 11 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 171	M.fr. XI, 11.
R. inops, Schauer in Lehm. pl. Preiss. II, 224 (1847) ... W.A.	—	—	—	—	—	—	B.f. III, 171	M.fr. XI, 11.
PHYMATOCARPUS, F. v. Mueller, fragm. III, 120 (1862).								
P. porphyrocephalus, F. v. M., fragm. III, 121 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 172	M.fr. III, 121.
P. Maxwellii, F. v. M., fragm. IX, 45 (1875) ... W.A.	—	—	—	—	—	—	—	M.fr. IX, 45.
CALOTHAMNUS, Labillardière, Nov. Holl. pl. specim. II, 25 (1806). (Billotia.)								
C. pachystachyus, Benth. Fl. Austr. III, 173 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 173	
C. longissimus, F. v. M., fragm. III, 112 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 174	M.fr. III, 112.
C. blepharanthus, F. v. M., fragm. III, 112 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 174	M.fr. III, 112; X, 31 & 55.
C. sanguineus, Labillardière, Nov. Holl. pl. spec. II, 25, t. 164 (1806) W.A.	—	—	—	—	—	—	B.f. III, 174	M.fr. X, 31.
C. torulosus, Schauer in nov. act. acad. Caes. XXI, 25 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 175	M.fr. X, 31.
C. gibbosus, Benth. Fl. Austr. III, 175 (1866) ... W.A.	—	—	—	—	—	—	B.f. III, 175	
C. gracilis, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812) W.A.	—	—	—	—	—	—	B.f. III, 175	
C. blepharosperrmus, F. v. M., fragm. III, 111 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 175	M.fr. III, 111.
C. Gilesii, F. v. M., fragm. X, 31 (1876) ... W.A.	—	—	—	—	—	—	—	M.fr. X, 31.
C. chrysanthus, F. v. M., fragm. III, 112 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 176	M.fr. III, 112.
C. Oldfieldii, F. v. M., fragm. III, 113 (1862) ... W.A.	—	—	—	—	—	—	B.f. III, 176	M.fr. III, 113.
C. planifolius, Lehmann, delect. sem. hort. bot. Hamb. 7 (1842) W.A.	—	—	—	—	—	—	B.f. III, 176	M.fr. X, 31.
C. lateralis, Lindley, Bot. Regist. XXV, App. IX (1839) ... W.A.	—	—	—	—	—	—	B.f. III, 177	
C. affinis, Turczaninow in Bullet. de St. Petersb. 428 (1852) ... W.A.	—	—	—	—	—	—	B.f. III, 177	M.fr. III, 113; VIII, 184.
C. Preissii, Schauer in nov. act. acad. Caes. XXI, 31 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 177	
C. Schaueri, Lehmann, delect. sem. hort. bot. Hamb. 7 (1842) ... W.A.	—	—	—	—	—	—	B.f. III, 178	
C. Lehmanni, Schauer in nov. act. acad. Caes. XXI, 31 (1844) ... W.A.	—	—	—	—	—	—	B.f. III, 178	
C. villosus, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812) W.A.	—	—	—	—	—	—	B.f. III, 178	
C. pinifolius, F. v. M., fragm. III, 153 (1863) ... W.A.	—	—	—	—	—	—	B.f. III, 179	M.fr. III, 153.

<i>C. rupestris</i> , Schauer in nov. act. acad. Caes. XXI, 26 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 179	
<i>C. quadrifidus</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812) ...	W.A.	—	—	—	—	—	B.f. III, 179	M.fr. X, 31.
<i>C. asper</i> , Turczaninow in Bull. Mosc. XXII, 25 (1849) ...	W.A.	—	—	—	—	—	B.f. III, 180	M.fr. X, 31.
<i>C. homalophyllus</i> , F. v. M., fragm. III, 111 (1862) ...	W.A.	—	—	—	—	—	B.f. III, 180	M.fr. X, 31.
LAMARCHEA, Gaudichaud in Freyc. Voy. Bot. 483, t. 110 (1826).								
<i>L. hakeaeifolia</i> , Gaudichaud in Freyc. Voy. 484, t. 110 (1826) ...	W.A.	—	—	—	—	—	B.f. III, 123	
EREMAEA, Lindley, Bot. Regist. XXV, App. XI (1839).								
<i>E. ambriata</i> , Lindley, Bot. Regist. XXV, App. XI (1839) ...	W.A.	—	—	—	—	—	B.f. III, 181	M.fr. XI, 11.
<i>E. acutifolia</i> , F. v. M., fragm. II, 30 (1860) ...	W.A.	—	—	—	—	—	B.f. III, 181	M.fr. II, 30.
<i>E. violacea</i> , F. v. M., fragm. XI, 10 (1878) ...	W.A.	—	—	—	—	—	—	N.fr. XI, 10.
<i>E. pilosa</i> , Lindley, Bot. Regist. XXV, App. XI (1839) ...	W.A.	—	—	—	—	—	B.f. III, 182	M.fr. XI, 11.
<i>E. ebracteata</i> , F. v. M., fragm. II, 29 (1860) ...	W.A.	—	—	—	—	—	B.f. III, 182	M.fr. XI, 11.
<i>E. beaufortoides</i> , Benth. Fl. Austr. III, 182 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 182	M.fr. XI, 11.
ANGOPHORA, Cavanilles, icon. IV, 21, t. 331 (1797).								
<i>A. cordifolia</i> , Cavanilles, icon. IV, 21, t. 338 (1797) ...	—	—	—	—	N.S.W.	—	B.f. III, 183	
<i>A. subvelutina</i> , F. v. M., fragm. I, 31 (1858) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 184	M.fr. I, 31.
<i>A. intermedia</i> , De Candolle, prodr. III, 222 (1828) ...	—	—	—	—	V.	N.S.W.	B.f. III, 184	
<i>A. lanceolata</i> , Cavanilles, icon. IV, 22, t. 339 (1797) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 184	
EUCALYPTUS, L'Heritier, sertum Anglicum 18 (1788). (Eudesmia, Symphyomyrtus.)								
I. RENANTHERAE.								
<i>E. stellulata</i> , Sieber in De Cand., prodr. III, 217 (1828) ...	—	—	—	V.	N.S.W.	—	B.f. III, 200	M.fr. II, 45; IX, 172.
<i>E. pauciflora</i> , Sieber in Sprengel, cur. poster. 195 (1827) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. III, 201	M.fr. III, 52.
<i>E. regnans</i> , F. v. M. in Report Acclim. Soc. Vict. 20 (1870) ...	—	—	T.	V.	N.S.W.	—	—	
<i>E. amygdalina</i> , Labill., Nov. Holl. pl. spec. II, 14, t. 154 (1806) ...	—	—	T.	V.	N.S.W.	—	B.f. III, 202	M.fr. II, 53; VII, 42&44.
<i>E. coccifera</i> , J. Hooker in Lond. Journ. of Bot. VI, 477 (1847) ...	—	—	T.	—	—	—	B.f. III, 204	
<i>E. obliqua</i> , L'Heritier, sertum Anglicum 18, t. 20 (1788) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. III, 204	M.fr. II, 44&171; XI, 33.
<i>E. stricta</i> , Sieber in De Candolle, prodr. III, 218 (1828) ...	—	—	—	—	N.S.W.	—	B.f. III, 217	M.fr. XI, 38.
<i>E. planchoniana</i> , F. v. M., fragm. XI, 43 (1878) ...	—	—	—	—	—	Q.	—	M.fr. XI, 43.
<i>E. Baileyana</i> , F. v. M., fragm. XI, 37 (1878) ...	—	—	—	—	—	Q.	—	M.fr. XI, 37.
<i>E. sepulchralis</i> , F. v. M., Eucalyptographia eight dec. (1882) ...	W.A.	—	—	—	—	—	—	
<i>E. bupestium</i> , F. v. M., fragm. III, 57 (1862) ...	W.A.	—	—	—	—	—	—	
<i>E. marginata</i> , Smith in Transact. Linn. Soc. VI, 302 (1802) ...	W.A.	—	—	—	—	—	B.f. III, 205	M.fr. III, 57. [XI, 38.
<i>E. santalifolia</i> , F. v. M. in Transact. Vict. Inst. I, 35 (1854) ...	W.A.	S.A.	—	—	—	—	B.f. III, 209	M.fr. II, 40&174; VII, 43;
<i>E. macrorrhyncha</i> , F. v. M., First general Report, 12 (1853) ...	—	—	—	—	—	—	B.f. III, 206	
<i>E. capitellata</i> , Smith in White, Journ. Voy. N.S.W. 216 (1790) ...	—	S.A.	—	V.	N.S.W.	—	B.f. III, 207	M.fr. XI, 45.
<i>E. eugenoides</i> , Sieber in Spreng. cur. poster. 195 (1827) ...	—	—	—	V.	N.S.W.	—	B.f. III, 206	M.fr. II, 173; XI, 38.
<i>E. piperita</i> , Smith in White, Journ. Voy. N.S.Wales 226 (1790) ...	—	—	—	V.	N.S.W.	—	—	
<i>E. pilularis</i> , Smith in Transact. Linn. Soc. III, 284 (1797) ...	—	—	—	V.	N.S.W.	Q.	B.f. III, 207	M.fr. II, 173; VII, 44; XI, 38
<i>E. triantha</i> , Link, enum. pl. hort. bot. Berol. II, 20 (1822) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 208	M.fr. II, 61&172; VII, 44;
<i>E. haemastoma</i> , Smith in Transact. Linn. Soc. III, 285 (1797) ...	—	—	T.	V.	N.S.W.	Q.	—	[XI, 38.
<i>E. Sieberiana</i> , F. v. M., Eucalyptogr. sec. decade (1879) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. III, 212	M.fr. II, 51&175; VII, 43;
<i>E. microcorys</i> , F. v. M., fragm. II, 50 (1860) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 202	M.fr. XI, 37. [XI, 38.
—	—	—	—	—	N.S.W.	Q.	B.f. III, 212	M.fr. II, 50.
II. PORANTHERAE.								
<i>E. paniculata</i> , Smith in Transact. Linn. Soc. III, 287 (1797) ...	—	S.A.	—	V.	N.S.W.	—	B.f. III, 211	M.fr. II, 174; VII, 43.
<i>E. Leucoxydon</i> , F. v. M. in Transact. Vict. Inst. I, 33 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	B.f. III, 209	M.fr. II, 60&175.
<i>E. melliodora</i> , Cunningham, in Walp. rep. bot. syst. II, 924 (1843) ...	—	—	—	V.	N.S.W.	—	B.f. III, 210	M.fr. VII, 43.
<i>E. polyanthema</i> , Schauer in Walpers, rep. bot. syst. II, 924 (1843) ...	—	—	—	V.	N.S.W.	—	B.f. III, 213	M.fr. VII, 44.
<i>E. populifolia</i> , Hooker, icon. plant. 279 (1852) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 214&243	
<i>E. ochrophloia</i> , F. v. M., fragm. XI, 36 (1878) ...	—	—	—	—	N.S.W.	Q.	—	M.fr. XI, 36. [XI, 36.
<i>E. gracilis</i> , F. v. M. in Transact. Vict. Inst. I, 35 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	B.f. III, 222	M.fr. VII, 43; VIII, 184;
<i>E. uncinata</i> , Turczaninow in Bull. de Mosc. XXII, 23 (1849) ...	W.A.	S.A.	—	V.	N.S.W.	—	B.f. III, 216	
<i>E. odorata</i> , Behr in Schlecht. Linnaea XX, 657 (1847) ...	—	S.A.	—	—	—	—	B.f. III, 215	M.fr. II, 66.
<i>E. largiflorens</i> , F. v. M. in Transact. Vict. Inst. I, 34 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	B.f. III, 215	M.fr. VII, 43.
<i>E. Behriana</i> , F. v. M. in Transact. Vict. Inst. I, 34 (1854) ...	—	S.A.	—	V.	—	—	B.f. III, 214	
<i>E. hemiphloia</i> , F. v. M., fragm. II, 62 (1860) ...	—	S.A.	—	V.	N.S.W.	Q.	B.f. III, 216	M.fr. II, 62.
<i>E. pruinosa</i> , Schauer in Walp. rep. bot. syst. II, 926 (1843) ...	—	—	—	—	—	N.A.	B.f. III, 213	M.fr. III, 132.
<i>E. melanophloia</i> , F. v. M. in Journ. Linn. Soc. III, 93 (1858) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 220	
III. PARALLELANTHERAE.								
<i>E. Cloeziana</i> , F. v. M., fragm. XI, 44 (1878) ...	—	—	—	—	—	Q.	—	M.fr. XI, 44.
<i>E. Howittiana</i> , F. v. M. in Wing's S. Sc. Record Aug. (1882) ...	—	—	—	—	—	Q.	—	
<i>E. drepanophylla</i> , F. v. M. in Benth. Fl. Austr. III, 221 (1866) ...	—	—	—	—	—	Q.	B.f. III, 221	
<i>E. crebra</i> , F. v. M. in Journ. Linn. Soc. III, 87 (1858) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. III, 221
<i>E. decipiens</i> , Endlicher in Hueg. enum. 49 (1837) ...	W.A.	—	—	—	—	—	B.f. III, 218	
<i>E. concolor</i> , Schauer in Lehm. pl. Preiss. I, 129 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 247	
<i>E. siderophloia</i> , Benth. Fl. Austr. III, 220 (1866) ...	—	—	—	—	N.S.W.	Q.	B.f. III, 220	M.fr. II, 61; VII, 44.
<i>E. microtheca</i> , F. v. M. in Journ. Linn. Soc. III, 87 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	B.f. III, 223	
<i>E. Raveretiana</i> , F. v. M., fragm. X, 99 (1877) ...	—	—	—	—	—	Q.	—	M.fr. X, 99.
<i>E. brachyandra</i> , F. v. M. in Journ. Linn. Soc. III, 97 (1858) ...	—	—	—	—	—	N.A.	B.f. III, 223	
<i>E. pachyphylla</i> , F. v. M. in Journ. Linn. Soc. III, 98 (1858) ...	—	S.A.	—	—	—	N.A.	B.f. III, 237	
<i>E. pyriformis</i> , Turczaninow in Bull. Mosc. XXII, 22 (1849) ...	W.A.	S.A.	—	—	—	—	B.f. III, 226	M.fr. II, 32; X, 5&84.
<i>E. macrocarpa</i> , Hooker, icones plant. t. 405-407 (1842) ...	W.A.	—	—	—	—	—	B.f. III, 224	M.fr. II, 41; XI, 38.
<i>E. Preissiana</i> , Schauer in Lehm. pl. Preiss. I, 131 (1844) ...	W.A.	—	—	—	—	—	B.f. III, 232	M.fr. II, 38; VII, 42.
<i>E. alpina</i> , Lindley in Mitch. Three Exped. II, 175 (1838) ...	—	—	—	V.	—	—	B.f. III, 225	M.fr. II, 68; VII, 42.
<i>E. Globulus</i> , Labillardiere, Voy. I, 153, t. 13 (1799) ...	—	—	T.	V.	N.S.W.	—	B.f. III, 225	M.fr. II, 68.
<i>E. megacarpa</i> , F. v. M., fragm. II, 70 (1860) ...	W.A.	—	—	—	—	—	B.f. III, 232	M.fr. VII, 43.
<i>E. cosmophylla</i> , F. v. M. in Transact. Vict. Inst. 32 (1854) ...	—	S.A.	—	—	—	—	B.f. III, 225	M.fr. VII, 43.
<i>E. cordata</i> , Labillardiere, Nov. Holl. pl. spec. II, 13, t. 152 (1806) ...	—	—	T.	—	—	—	B.f. III, 224	
<i>E. urnigera</i> , J. Hooker in Lond. Journ. of Bot. VI, 477 (1847) ...	—	—	T.	—	—	—	B.f. III, 227	

<i>E. longifolia</i> , Link, enum. pl. hort. bot. Berol. II, 29 (1822) ...	—	—	—	V.	N.S.W.	—	—	B. f. III, 226	M. fr. II, 50; VII, 43.
<i>E. erythronema</i> , Turcz. in Bull. de l'acad. de St. Petersb. 415 (1852) ...	W.A.	—	—	—	—	—	—	B. f. III, 227	M. fr. VIII, 184.
<i>E. caesia</i> , Benth, Fl. Austr. III, 227 (1866) ...	W.A.	—	—	—	—	—	—	B. f. III, 227	M. fr. XI, 40. [184.
<i>E. tetraptera</i> , Turczaninow in Bull. de Mosc. XXII, 22 (1849) ...	W.A.	—	—	—	—	—	—	B. f. III, 228	M. fr. II, 34; IV, 172; V, III,
<i>E. miniata</i> , A. Cunningham in Walp. rep. bot. syst. II, 925 (1843) ...	—	—	—	—	—	Q.	N.A.	B. f. III, 228	M. fr. XI, 38 & 42.
<i>E. phoenicea</i> , F. v. M. in Journ. Linn. Soc. III, 91 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 251	
<i>E. robusta</i> , Smith, Specim. Bot. New Holl. 40, t. 13 (1793) ...	—	—	—	—	N.S.W.	Q.	—	B. f. III, 228	M. fr. II, 43 & 171; VII, 44.
<i>E. botryoides</i> , Smith in Transact. Linn. Soc. III, 286 (1797) ...	—	—	—	—	V.	N.S.W.	Q.	B. f. III, 229	M. fr. II, 48 & 175; VII, 43.
<i>E. gonocalyx</i> , F. v. M., fragm. II, 48 (1860) ...	—	—	S.A.	—	V.	N.S.W.	—	B. f. III, 229	M. fr. IV, 52.
<i>E. pallidifolia</i> , F. v. M., fragm. III, 131 (1863) ...	—	—	—	—	—	—	N.A.	B. f. III, 236	M. fr. III, 131.
<i>E. incrassata</i> , Labillard., Nov. Holl. pl. spec. II, 12, t. 150 (1806) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B. f. III, 231	
<i>E. oleosa</i> , F. v. M. in Nederl. Kruidk. Arch. IV, 127 (1859) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B. f. III, 248	M. fr. II, 56; XI, 14.
<i>E. goniantha</i> , Turczaninow in Bullet. de Mosc. XX, 163 (1847) ...	W.A.	—	—	—	—	—	—	B. f. III, 248	
<i>E. falcata</i> , Turczaninow in Bullet. de Mosc. XX, 163 (1847) ...	W.A.	—	—	—	—	—	—	B. f. III, 248	
<i>E. salmonophloia</i> , F. v. M., fragm. XI, 11 (1878) ...	—	—	—	—	—	—	—	—	M. fr. XI, 11.
<i>E. leptopoda</i> , Benth, Fl. Austr. III, 238 (1866) ...	—	—	—	—	—	—	—	B. f. III, 238	M. fr. XI, 11 & 13.
<i>E. salubris</i> , F. v. M., fragm. X, 54 (1876) ...	—	—	—	—	—	—	—	—	M. fr. XI, 12.
<i>E. angustissima</i> , F. v. M., fragm. IV, 25 (1863) ...	—	—	S.A.	—	—	—	—	B. f. III, 238	M. fr. IV, 25.
<i>E. Doratoxylon</i> , F. v. M., fragm. II, 55 (1860) ...	—	—	—	—	—	—	—	B. f. III, 249	M. fr. II, 55.
<i>E. decurva</i> , F. v. M., fragm. III, 130 (1863) ...	—	—	—	—	—	—	—	B. f. III, 249	M. fr. III, 130.
<i>E. Cooperiana</i> , F. v. M., fragm. XI, 83 (1880) ...	—	—	—	—	—	—	—	—	M. fr. XI, 83.
<i>E. corynocalyx</i> , F. v. M., fragm. II, 43 (1860) ...	—	—	S.A.	—	V.	—	—	B. f. III, 218	M. fr. VII, 43.
<i>E. gomphocephala</i> , De Candolle, prodr. III, 220 (1828) ...	—	—	—	—	—	—	—	B. f. III, 231	M. fr. II, 36.
<i>E. Oldfieldii</i> , F. v. M., fragm. II, 37 (1860) ...	—	—	—	—	—	—	—	B. f. III, 237	M. fr. II, 37.
<i>E. orbifolia</i> , F. v. M., fragm. V, 50 (1865) ...	—	—	—	—	—	—	—	B. f. III, 238	M. fr. V, 50.
<i>E. diversicolor</i> , F. v. M., fragm. III, 131 (1863) ...	—	—	—	—	—	—	—	B. f. III, 251	M. fr. III, 131.
<i>E. patellaris</i> , F. v. M. in Journ. Linn. Soc. III, 84 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 244	
<i>E. tessellaris</i> , F. v. M. in Journ. Linn. Soc. III, 88 (1858) ...	—	—	—	—	—	Q.	N.A.	B. f. III, 251	
<i>E. clavigera</i> , A. Cunningham in Walp. rep. bot. syst. II, 926 (1843) ...	—	—	—	—	—	—	N.A.	B. f. III, 250	
<i>E. ferruginea</i> , Schauer in Walp. repert. II, 926 (1843) ...	—	—	—	—	—	—	N.A.	B. f. III, 254	
<i>E. grandifolia</i> , R. Brown in Benth. Fl. Austr. III, 250 (1866) ...	—	—	—	—	—	—	N.A.	B. f. III, 250	
<i>E. aspera</i> , F. v. M. in Journ. Linn. Soc. III, 95 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 250	
<i>E. patens</i> , Benth, Fl. Austr. III, 247 (1866) ...	W.A.	—	—	—	—	—	—	B. f. III, 247	
<i>E. Todtiana</i> , F. v. M., in Wing's S. Sc. Record Aug. (1882) ...	W.A.	—	—	—	—	—	—	—	
<i>E. vernicosa</i> , J. Hooker in Lond. Journ. of Bot. VI, 478 (1847) ...	—	—	T.	—	—	—	—	B. f. III, 232	
<i>E. Gunnii</i> , J. Hooker in Lond. Journ. Bot. III, 499 (1844) ...	—	S.A.	T.	V.	N.S.W.	—	—	B. f. III, 246	M. fr. II, 62; VII, 43; XI, 38.
<i>E. resinifera</i> , Smith in White's Journ. Voy. N.S. Wales 231 (1790) ...	—	—	—	—	N.S.W.	Q.	—	B. f. III, 245	M. fr. II, 172; VII, 43.
<i>E. saligna</i> , Smith in Transact. Linn. Soc. III, 285 (1797) ...	—	—	—	—	N.S.W.	—	—	B. f. III, 245	M. fr. II, 173; VII, 44.
<i>E. punctata</i> , De Candolle, prodr. III, 217 (1828) ...	—	—	—	—	N.S.W.	—	—	B. f. III, 242	M. fr. VII, 43.
<i>E. oligantha</i> , Schauer in Walp. rep. II, 926 (1843) ...	—	—	—	—	—	—	N.A.	B. f. III, 213	
<i>E. alba</i> , Reinwardt in Blume, Bijdr. 1101 (1826) ...	—	—	—	—	—	Q.	N.A.	B. f. III, 243	
<i>E. pulverulenta</i> , Sims, Bot. Mag. t. 2089 (1819) ...	—	—	—	V.	N.S.W.	—	—	B. f. III, 224	M. fr. II, 70.
<i>E. Stuartiana</i> , F. v. M. in Benth. Fl. Austr. III, 243 (1866) ...	—	S.A.	T.	V.	N.S.W.	—	—	B. f. III, 243	M. fr. II, 62.
<i>E. viminalis</i> , Labillardière, Nov. Holl. pl. spec. II, 12, t. 151 (1806) ...	—	S.A.	T.	V.	N.S.W.	—	—	B. f. III, 239	M. fr. II, 64; VII, 44.
<i>E. rostrata</i> , Schlechtendal, Linnaea XX, 655 (1847) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B. f. III, 240	M. fr. VII, 44.
<i>E. tereticoxis</i> , Smith, Spec. Bot. New Holl. 41 (1793) ...	—	—	—	V.	N.S.W.	Q.	—	B. f. III, 241	
<i>E. rudis</i> , Endlicher in Hueg. enum. 49 (1837) ...	W.A.	—	—	—	—	—	—	B. f. III, 244	M. fr. XI, 14.
<i>E. foecunda</i> , Schauer in Lehm. pl. Preiss. I, 130 (1844) ...	W.A.	—	—	—	—	—	—	B. f. III, 252	
<i>E. redunca</i> , Schauer in Lehm. pl. Preiss. I, 127 (1844) ...	W.A.	—	—	—	—	—	—	B. f. III, 253	M. fr. XI, 15.
<i>E. grossa</i> , F. v. M. in Benth. Fl. Austr. III, 232 (1866) ...	W.A.	—	—	—	—	—	—	B. f. III, 232	M. fr. VII, 41.
<i>E. obovata</i> , Turczaninow in Bull. Acad. Petersb. 416 (1852) ...	W.A.	—	—	—	—	—	—	B. f. III, 234	M. fr. VII, 44; VIII, 184.
<i>E. occidentalis</i> , Endlicher in Hueg. enum. 49 (1837) ...	W.A.	—	—	—	—	—	—	B. f. III, 235	M. fr. II, 39; VII, 43.
<i>E. cornuta</i> , Labillardière, Voy. I, 403, t. 20 (1799) ...	W.A.	—	—	—	—	—	—	B. f. III, 234	M. fr. II, 39; XI, 38.
<i>E. gamophylla</i> , F. v. M., fragm. XI, 40 (1878) ...	—	S.A.	—	—	—	—	N.A.	—	M. fr. XI, 40.
<i>E. perfoliata</i> , R. Brown in Benth. Fl. Austr. III, 253 (1866) ...	—	—	—	—	—	—	N.A.	B. f. III, 253	
<i>E. setosa</i> , Schauer in Walpers, rep. bot. syst. II, 926 (1843) ...	—	—	—	—	—	Q.	N.A.	B. f. III, 254	M. fr. III, 123; X, 106.
<i>E. Torelliana</i> , F. v. M., fragm. X, 106 (1877) ...	—	—	—	—	—	Q.	—	—	M. fr. X, 106.
<i>E. peltata</i> , Benth, Fl. Austr. III, 254 (1866) ...	—	—	—	—	—	Q.	—	B. f. III, 254	
<i>E. latifolia</i> , F. v. M. in Journ. Linn. Soc. III, 94 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 255	
<i>E. ptychocarpa</i> , F. v. M. in Journ. Linn. Soc. III, 90 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 255	M. fr. VII, 44.
<i>E. Abergiana</i> , F. v. M., fragm. XI, 41 (1878) ...	—	—	—	—	—	Q.	—	—	M. fr. XI, 41.
<i>E. calophylla</i> , R. Brown in Journ. Geogr. Soc. 20 (1831) ...	W.A.	—	—	—	—	—	—	B. f. III, 255	M. fr. II, 35 & 171. [XI, 15.
<i>E. ficifolia</i> , F. v. M., fragm. II, 85 (1860) ...	W.A.	—	—	—	—	—	—	B. f. III, 256	M. fr. II, 85 & 176; VI, 25;
<i>E. corymbosa</i> , Smith, Specim. Bot. New Holl. 43 (1793) ...	—	—	—	N.S.W.	Q.	N.A.	—	B. f. III, 256	M. fr. II, 46 & 174.
<i>E. terminalis</i> , F. v. M. in Journ. Linn. Soc. III, 89 (1858) ...	—	S.A.	—	—	—	Q.	N.A.	B. f. III, 257	
<i>E. dichromophloia</i> , F. v. M. in Journ. Linn. Soc. III, 89 (1858) ...	—	—	—	—	—	—	N.A.	B. f. III, 257	
<i>E. trachyphloia</i> , F. v. M. in Journ. Linn. Soc. III, 90 (1858) ...	—	—	—	—	—	Q.	—	B. f. III, 221	
<i>E. eximia</i> , Schauer in Walpers, rep. bot. syst. II, 925 (1843) ...	—	—	—	—	N.S.W.	—	—	B. f. III, 258	
<i>E. maculata</i> , Hooker, icones plant. t. 619 (1844) ...	—	—	—	—	N.S.W.	Q.	—	B. f. III, 258	M. fr. II, 47; VII, 43.
<i>E. Watsoniana</i> , F. v. M., fragm. X, 98 (1877) ...	—	—	—	—	—	Q.	—	—	M. fr. X, 98.
<i>E. odontocarpa</i> , F. v. M. in Journ. Linn. Soc. III, 98 (1858) ...	W.A.	—	—	—	—	—	—	B. f. III, 260	
<i>E. tetradonta</i> , F. v. M. in Journ. Linn. Soc. III, 97 (1858) ...	—	—	—	—	—	Q.	N.A.	B. f. III, 260	
<i>E. erythrocorys</i> , F. v. M., fragm. II, 33 (1860) ...	W.A.	—	—	—	—	—	—	B. f. III, 258	M. fr. II, 33.
<i>E. tetragona</i> , F. v. M., fragm. VI, 51 (1864) ...	W.A.	—	—	—	—	—	—	B. f. III, 259	M. fr. II, 37.
<i>E. eudesmioides</i> , F. v. M., fragm. II, 35 (1860) ...	W.A.	—	—	—	—	—	—	B. f. III, 260	M. fr. II, 35.

TRISTANIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812). (Lophostemon, Tristaniopsis.)

<i>T. neriifolia</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812) ...	—	—	—	N.S.W.	—	—	B. f. III, 261	M. fr. IV, 56.
<i>T. suaveolens</i> , Smith in Rees, Cycl. XXXVI (1817) ...	—	—	—	N.S.W.	Q.	N.A.	B. f. III, 261	M. fr. I, 81.
<i>T. conferta</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812) ...	—	—	—	N.S.W.	Q.	N.A.	B. f. III, 263	M. fr. I, 82.
<i>T. lactiflora</i> , F. v. M., fragm. I, 82 (1858) ...	—	—	—	—	—	N.A.	B. f. III, 263	M. fr. I, 82.

<i>T. exiliflora</i> , F. v. M., fragm. V, 11 (1865)...	—	—	—	—	Q.	—	B. fl. III, 264	M. fr. IX, 190.
<i>T. laurina</i> , R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	—	—	—	V.	N. S. W.	Q.	—	B. fl. III, 264
<i>T. psidioides</i> , Cunningham in Bot. Regist. XXII, n. 1839 (1837)	—	—	—	—	—	—	N. A.	B. fl. III, 264
<i>T. umbrosa</i> , Cunningham in Bot. Regist. XXII, n. 1839 (1837)...	—	—	—	—	—	—	N. A.	B. fl. III, 265
METROSIDEROS, Banks in Gaertner, de fruct. I, 170, t. 34 (1788). (Syncarpia, Kamptzia, Lysicarpus, Xanthostemon, Nania, Fremya, Cloezia, Spermalepis.)								
<i>M. glomulifera</i> , Smith in Transact. Linn. Soc. III, 269 (1797) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 265
<i>M. leptopetala</i> , F. v. M., Docum. Intercol. Exhib. 30 (1867) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 266
<i>M. ternifolia</i> , F. v. M., Docum. Intercol. Exhib. 30 (1867) ...	—	—	—	—	—	Q.	—	B. fl. III, 267
<i>M. eucalyptoides</i> , F. v. M., fragm. I, 243 (1858) ...	—	—	—	—	—	—	N. A.	B. fl. III, 267
<i>M. polymorpha</i> , Gaudichaud, Freyc. Voy. Bot. 99 t. 108 (1826)	—	—	—	—	N. S. W.	—	—	M. fr. IX, 14.
<i>M. nervulosa</i> , C. Moore & F. v. M., fragm. VIII, 15 (1873) ...	—	—	—	—	N. S. W.	—	—	M. fr. I, 80.
<i>M. chrysanthia</i> , F. v. M., fragm. IV, 159 (1864) ...	—	—	—	—	—	Q.	—	B. fl. III, 268
<i>M. paradoxa</i> , F. v. M., fragm. I, 243 (1859) ...	—	—	—	—	—	—	N. A.	B. fl. III, 269
<i>M. tetrapetala</i> , F. v. M., fragm. VII, 41 (1870) ...	—	—	—	—	—	—	N. A.	—
BACKHOUSIA, Hooker and Harvey in Bot. Mag. t. 4133 (1845).								
<i>B. myrtifolia</i> , Hooker & Harvey in Bot. Mag. t. 4133 (1845) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 269
<i>B. angustifolia</i> , F. v. M., fragm. I, 79 (1858) ...	—	—	—	—	—	Q.	—	B. fl. III, 270
<i>B. sciadophora</i> , F. v. M., fragm. II, 26 (1860) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 270
<i>B. citriodora</i> , F. v. M., fragm. I, 78 (1858) ...	—	—	—	—	—	Q.	—	B. fl. III, 270
OSBORNIA, F. v. Mueller, fragm. III, 30 (1862).								
<i>O. octodonta</i> , F. v. M., fragm. III, 31 (1862) ...	—	—	—	—	—	Q.	N. A.	B. fl. III, 271
RHODOMYRTUS, De Candolle, prodr. III, 240 (1828).								
<i>R. psidioides</i> , Benth. Fl. Austr. III, 272 (1866) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 272
<i>R. trineura</i> , F. v. M. in Benth. Fl. Austr. III, 272 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 272
<i>R. cymiflora</i> , F. v. M. in Benth. Fl. Austr. III, 273 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 273
<i>R. macrocarpa</i> , Benth. Fl. Austr. III, 273 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 273
MYRTUS, Tournefort, inst. 640, t. 409 (1700), from Plinius.								
<i>M. rhytisperma</i> , F. v. M., fragm. I, 77 (1858) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 274
<i>M. tenuifolia</i> , Smith in Transact. Linn. Soc. III, 280 (1797) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 274
<i>M. gonolada</i> , F. v. M. in Benth. Fl. Austr. III, 275 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 275
<i>M. lasioclada</i> , F. v. M., fragm. IX, 148 (1875) ...	—	—	—	—	—	Q.	—	—
<i>M. Hillii</i> , Benth. Fl. Austr. III, 275 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 275
<i>M. Becklerii</i> , F. v. M., fragm. II, 85 (1861) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 275
<i>M. Bidwillii</i> , Benth. Fl. Austr. III, 275 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 275
<i>M. racemulosa</i> , Benth. Fl. Austr. III, 276 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 276
<i>M. acmenoides</i> , F. v. M., fragm. I, 77 (1858) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 276
<i>M. fragrantissima</i> , F. v. M. in Benth. Fl. Austr. III, 277 (1866) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 277
<i>M. Shepherdi</i> , F. v. M., fragm. IX, 148 (1875) ...	—	—	—	—	—	Q.	—	—
RHODAMNIA, Jack in Hook. Comp. to the Bot. Mag. I, 153 (1835).								
<i>R. sessiliflora</i> , Benth. Fl. Austr. III, 275 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 277
<i>R. trinervia</i> , Blume, Mus. Bot. Lugd. I, 79 (1849) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 278
<i>R. argentea</i> , Benth. Fl. Austr. III, 276 (1866) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 278
<i>R. Blairiana</i> , F. v. M., fragm. IX, 141 (1875) ...	—	—	—	—	—	Q.	—	—
FENZLIA, Endlicher, Atacta 19, t. 17 & 18 (1833).								
<i>F. obtusa</i> , Endlicher, Atacta 19, t. 17 (1833) ...	—	—	—	—	—	Q.	—	B. fl. III, 279
<i>F. retusa</i> , Endlicher, Atacta 20, t. 18 (1833) ...	—	—	—	—	—	—	N. A.	B. fl. III, 279
DECASPERMUM, R. & G. Forster, char. gen. 73, t. 37 (1776). (Nelitris.)								
<i>D. paniculatum</i> , Baillon, hist. des plant. 350 (1871) ...	—	—	—	—	—	Q.	—	B. fl. III, 279
EUGENIA, Micheli, nov. pl. gen. 226, t. 108 (1729). (Plinia, Plumier 1703, Aemene, Jambosa, Syzygium.)								
<i>E. rariflora</i> , Benth. in Hook. Lond. Journ. II, 221 (1843) ...	—	—	—	—	—	Q.	—	B. fl. III, 282
<i>E. Smithii</i> , Poir. Encycl. method. Suppl. III, 126 (1813) ...	—	—	—	—	V.	N. S. W.	Q.	N. A.
<i>E. hemilampra</i> , F. v. M., fragm. IX, 145 (1875) ...	—	—	—	—	—	N. S. W.	Q.	—
<i>E. Ventenatii</i> , Benth. Fl. Austr. III, 283 (1866) ...	—	—	—	—	—	N. S. W.	Q.	—
<i>E. leptantha</i> , Wight, Illustr. II, 15 (1850) ...	—	—	—	—	—	Q.	—	B. fl. III, 283
<i>E. Moorei</i> , F. v. M., fragm. V, 33 (1865) ...	—	—	—	—	—	N. S. W.	Q.	—
<i>E. corynantha</i> , F. v. M., fragm. IX, 144 (1875) ...	—	—	—	—	—	N. S. W.	Q.	—
<i>E. corniflora</i> , F. v. M., fragm. V, 32 (1865) ...	—	—	—	—	—	Q.	—	B. fl. III, 284
<i>E. Hodgkinsoniae</i> , F. v. M., fragm. IX, 145 (1875) ...	—	—	—	—	—	N. S. W.	Q.	—
<i>E. Tierneyana</i> , F. v. M., fragm. V, 14 (1865) ...	—	—	—	—	—	Q.	—	B. fl. III, 284
<i>E. grandis</i> , Wight, Illustr. II, 17 (1850) ...	—	—	—	—	—	Q.	—	B. fl. III, 285
<i>E. suborbicularis</i> , Benth. Fl. Austr. III, 285 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 285
<i>E. Wilsonii</i> , F. v. M., fragm. V, 12 (1865) ...	—	—	—	—	—	Q.	—	B. fl. III, 285
<i>E. eucalyptoides</i> , F. v. M., fragm. IV, 55 (1863) ...	—	—	—	—	—	—	N. A.	B. fl. III, 285
<i>E. myrtifolia</i> , Sims, Bot. Mag. t. 2230 (1821) ...	—	—	—	—	N. S. W.	Q.	—	B. fl. III, 286
<i>E. angophoroides</i> , F. v. M., fragm. V, 33 (1865) ...	—	—	—	—	—	Q.	—	B. fl. III, 286
<i>E. Armstrongii</i> , Benth. Fl. Austr. III, 286 (1866) ...	—	—	—	—	—	—	N. A.	B. fl. III, 286
<i>E. oleosa</i> , F. v. M., fragm. V, 15 (1865) ...	—	—	—	—	—	Q.	—	B. fl. III, 287
<i>E. Dallachiana</i> , F. v. M. in Benth. Fl. Austr. III, 287 (1866) ...	—	—	—	—	—	Q.	—	B. fl. III, 287
<i>E. cryptophlebia</i> , F. v. M., fragm. IX, 144 (1875) ...	—	—	—	—	—	Q.	—	—
ACICALYPTUS, A. Gray, New Gen. Un. St. Exped. 9 (1853). (Calyptanthus partly.)								
<i>A. Fullageri</i> , F. v. M., fragm. VIII, 15 (1873) ...	—	—	—	—	N. S. W.	—	—	—
BARRINGTONIA, R. & G. Forster, char. gen. 75, t. 38 (1776). (Stravadium.)								
<i>B. speciosa</i> , R. & G. Forster, char. gen. 76, t. 38 (1776) ...	—	—	—	—	—	Q.	—	B. fl. III, 288
<i>B. racemosa</i> , Gaudichaud in Freyc. Voy. Bot. 483, t. 107 (1826) ...	—	—	—	—	—	Q.	—	—
<i>B. acutangula</i> , Gaertner, de fruct. II, 97, t. 101 (1791) ...	—	—	—	—	—	—	N. A.	B. fl. III, 288

- CAREYA, Roxburgh, pl. Corom. III, 13, t. 217 (1816).
 C. australis, F. v. M., fragm. V, 183 (1866) ... — — — — — Q. N.A. B.f.III,289 M.fr.V,183.
 SONNERATIA, Linné fl., suppl. plant. 38 (1781).
 S. acida, Linné fl., suppl. plant. 38 (1781)... — — — — — — N.A. B.f.III,301

MELASTOMACEAE.

A. L. de Jussieu, gen. plant. 328 (1789).

- MEMECYLON, Linné, Fl. Zeyl. app. 9 (1747).
 M. umbellatum, N. L. Burmann, Fl. Ind. 87 (1768) ... — — — — — Q. N.A. B.f.III,293
 OSBECKIA, Linné, spec. plant. I, 345 (1753).
 O. Chinensis, Linné, spec. plant. 345 (1753) ... — — — — — Q. — B.f.III,291 M.fr.IV,160.
 O. Australiana, Naudin in Ann. sc. nat. trois. sér. XIV, 59 (1850) ... — — — — — — N.A. B.f.III,291
 O. perangusta, F. v. M., fragm. V, 181 (1866) ... — — — — — — N.A. — M.fr.V,181.
 OTANTHERA, Blume in Flora, Regensb. IV, 488 (1831).
 O. bracteata, Korthals in Verh. Nat. Gesch. der Ned. Bes. 235, t. 51 (1841) ... — — — — — Q. — B.f.III,292
 MELASTOMA, J. Burmann, thesaur. Zeyl. 72 (1737).
 M. Malabathricum, Linné, spec. plant. 390 (1753) ... — — — — — N.S.W. Q. N.A. B.f.III,292

RHAMNACEAE.

A. L. de Jussieu, gen. plant. 376 (1789), from B. de Jussieu (1759).

- VENTILAGO, Gaertner, de fruct. I, 223, t. 49 (1788).
 V. viminalis, Hooker in Mitch. Trop. Austr. 369 (1848) ... — S.A. — — — N.S.W. Q. N.A. B.f.I,411 M.fr.IX,140.
 ZIZYPHUS, Tournefort, inst. 627, t. 403 (1700), from Plinius (Ziziphus).
 . Oenoplia, Miller, Gardeners Dictionary n. 3 (1731) ... — — — — — — N.A. B.f.I,412
 . Jujuba, Lamarck, Encycl. méth. III, 318 (1789) ... — — — — — — Q. — B.f.I,412
 . quadrilocularis, F. v. M., fragm. III, 57 (1862) ... — — — — — — N.A. B.f.I,412 M.fr.III,57.
 DALLACHYA, F. v. M., fragm. IX, 140 (1875). (Rhamnus partly.)
 D. Vitiensis, F. v. M., fragm. IX, 140 (1875) ... — — — — — Q. — — M.fr.XI,134.
 BERCHEMIA, Necker, elem. bot. II, 122 (1790).
 B. corollata, F. v. M., fragm. IX, 141 (1875) ... — — — — — Q. — — M.fr.IX,141. ~~~~~
 COLUBRINA, L. C. Richard in Ann. des sc. nat. X, 368 (1827).
 C. Asiatica, Brogniart in Ann. des sc. nat. X, 369 (1827) ... — — — — — Q. N.A. B.f.I,413 M.fr.IX,139.
 ALPHITONIA, Reisseck in Endl. gen. plant. 1098 (1840).
 A. excelsa, Reisseck in Endl. gen. plant. 1098 (1840) ... — — — — — N.S.W. Q. N.A. B.f.I,414 M.fr.IX,140.
 EMMENOSPERMUM, F. v. M., fragm. III, 63 (1862).
 E. alphitonioides, F. v. M., fragm. III, 63 (1862)... — — — — — N.S.W. Q. — B.f.I,415 M.fr.IX,140.
 E. Cunninghamii, Benth, Fl. Austr. I, 415 (1863) ... — — — — — — N.A. B.f.I,415
 POMADERRIS, Labillardière, Nov. Holl. pl. specim. I, 61, t. 86 (1804).
 P. lanigera, Sims, Bot. Mag. t. 1823 (1816) ... — — — — — T. V. N.S.W. — — B.f.I,416
 P. elliptica, Labillardière, Nov. Holl. pl. spec. I, 61, t. 86 (1804) ... — — — — — T. V. N.S.W. — — B.f.I,417 M.fr.IX,139.
 P. phillyroides, Sieber in De Cand., prodr. II, 33 (1825)... — — — — — N.S.W. Q. — B.f.I,418
 P. vacciniifolia, Reisseck & F. v. M. in Linnaea XXIX, 266 (1856) ... — — — — — V. N.S.W. — — B.f.I,418 M.fr.III,71.
 P. myrtilloides, Fenzl in Hueg. enum. 22 (1837) ... — — — — — W.A. — — — — — B.f.I,419 M.fr.IX,138.
 P. grandis, F. v. M., fragm. III, 68 (1862)... — — — — — W.A. — — — — — B.f.I,417 M.fr.IX,139.
 P. Calvertiana, F. v. M., fragm. IX, 138 (1875) ... — — — — — — N.S.W. — — — — — M.fr.IX,138.
 P. ledifolia, Cunningham in Field, N.S. Wales 351 (1825) ... — — — — — V. N.S.W. — — B.f.I,419 M.fr.III,69.
 P. apetala, Labillardière, Nov. Holl. pl. spec. I, 62, t. 87 (1804) ... — S.A. T. V. N.S.W. — — B.f.I,419 M.fr.III,69&73;IX,139.
 P. cinerea, Benth, Fl. Austr. I, 420 (1863) ... — — — — — V. N.S.W. — — B.f.I,420
 P. prunifolia, Cunningham in Hueg. enum. 22 (1837) ... — — — — — V. N.S.W. — — B.f.I,420 M.fr.III,75;IX,139.
 P. ligustrina, Sieber in De Cand., prodr. II, 34 (1825) ... — — — — — N.S.W. — — B.f.I,420 M.fr.III,71;IX,139.
 P. betulina, Cunningham in Bot. Mag. t. 3212 (1833) ... — — — — — V. N.S.W. — — B.f.I,421 M.fr.III,76;IX,139.
 P. Forrestiana, F. v. M., fragm. IX, 139 (1875) ... — — — — — W.A. — — — — — M.fr.IX,139.
 P. obcordata, Fenzl in Hueg. enum. 23 (1837) ... — — — — — W.A. S.A. — — — — — B.f.I,421 M.fr.III,73.
 P. racemosa, Hooker, Journ. of Bot. I, 256 (1834) ... — — — — — S.A. T. V. N.S.W. — — B.f.I,421 M.fr.III,75&168;IX,139.
 P. subrepanda, Reisseck & F. v. M. in Linnaea XXIX, 267 (1856) ... — — — — — V. — — — — — B.f.I,422 M.fr.III,74.
 P. elachophylla, F. v. M., fragm. II, 131 (1861) ... — — — — — V. — — — — — B.f.I,422 M.fr.II,166;IX,139.
 P. phyllicifolia, Loddiges, Bot. Cabinet t. 120 (1818) ... — — — — — T. V. — — — — — B.f.I,422 M.fr.III,73.
 P. intangenda, F. v. M., fragm. X, 52 (1876) ... — — — — — W.A. — — — — — — M.fr.X,53.
 CRYPTANDRA, Smith in Transact. Linn. Soc. IV, 217 (1798). (Trymalium, Spyridium, Stenanthemum, Wichurea, Stenodiscus.)
 C. albicans, F. v. M.; Trym., Reisseck in pl. Preiss. II, 280 (1847) W.A. — — — — — — B.f.I,423
 C. Billardieri, F. v. M.; Trym., Fenzl in Hueg. enum. 25 (1837) W.A. — — — — — — B.f.I,423
 C. ledifolia, F. v. M.; Trymalium, Fenzl in Hueg. enum. 24 (1837) W.A. — — — — — — B.f.I,424
 C. Daltoni; Trymalium, F. v. M., fragm. IX, 135 (1875) ... — — — — — V. — — — — — M.fr.IX,135.
 C. angustifolia, F. v. M.; Trym., Reisseck in pl. Preiss. II, 284 (1847) W.A. — — — — — — B.f.I,424 M.fr.IX,135.
 C. Wichurea, F. v. M.; Trym., Nees in Lehm. pl. Preiss. II, 281 (1847) W.A. — — — — — — B.f.I,425 M.fr.IX,135.
 C. ericifolia, Smith in Transact. Linn. Soc. X, 294, t. 18 (1811)... — — — — — N.S.W. Q. — B.f.I,438 M.fr.IX,137.
 C. propinqua, Cunningham in Hueg. enum. 23 (1837) ... — — — — — S.A. — — — — — N.S.W. — — B.f.I,442 M.fr.IX,137.
 C. hispidula, Reisseck & F. v. M. in Linnaea XXIX, 294 (1856) ... — — — — — S.A. — — — — — — B.f.I,439
 C. spyridioides, F. v. M., fragm. III, 68 (1862) ... — — — — — W.A. — — — — — — B.f.I,439 M.fr.IX,138.
 C. scoparia, Reisseck in Lehm. pl. Preiss. II, 283 (1847) W.A. — — — — — — B.f.I,439
 C. spinescens, Sieber in De Cand., prodr. II, 38 (1825) ... — — — — — N.S.W. — — — — — B.f.I,439
 C. amara, Smith in Transact. Linn. Soc. X, 295, t. 18 (1811) ... — — — — — S.A. T. V. N.S.W. Q. — B.f.I,440 M.fr.III,66&67;IX,137.
 C. lanosiflora, F. v. M., fragm. III, 65 (1862) ... — — — — — — N.S.W. — — B.f.I,440 M.fr.III,65.
 C. tomentosa, Lindley in Mitch. Three Exped. II, 178 (1838) ... — — — — — S.A. — — — — — V. N.S.W. — — B.f.I,441 M.fr.III,67;IX,138.
 C. nutans, Steudel in Lehm. pl. Preiss. I, 186 (1844) ... — — — — — W.A. — — — — — — B.f.I,441 M.fr.IX,138.

C. glabriflora, Benth. Fl. Austr. I, 441 (1863) ...	...	W.A.	—	—	—	—	—	B.f.I, 441		
C. alpina, J. Hooker, Fl. Tasm. I, 75, t. 12 (1860) ...	...	—	—	T.	—	—	—	B.f.I, 441		
C. leucopogon, Meissner in Lehm. pl. Preiss. II, 287 (1847) ...	...	W.A.	—	—	—	—	—	B.f.I, 442		
C. parvifolia, Turczaninow in Bull. de Mosc. I, 459 (1858) ...	...	W.A.	—	—	—	—	—	B.f.I, 442		
C. buxifolia, Fenzl in Hueg. enum. 23 (1837) ...	...	—	—	—	—	N.S.W.	—	B.f.I, 442	M.fr. IX, 138.	
C. pungens, Steudel in Lehm. pl. Preiss. I, 187 (1844) ...	...	W.A.	—	—	—	—	—	B.f.I, 443	M.fr. IX, 138.	
C. mutila, Nees in Lehm. pl. Preiss. II, 289 (1847) ...	...	W.A.	—	—	—	—	—	B.f.I, 443		
C. longistaminea, F. v. M., fragm. III, 64 (1862) ...	...	—	—	—	—	N.S.W.	—	B.f.I, 443	M.fr. IX, 138.	
C. arbutiflora, Fenzl in Hueg. enum. 26 (1837) ...	...	W.A.	—	—	—	—	—	B.f.I, 444	M.fr. IX, 138.	
C. miliaris, Reisseck in Lehm. pl. Preiss. II, 288 (1847) ...	...	W.A.	—	—	—	—	—	B.f.I, 444		
C. nudiflora, F. v. M., fragm. III, 64 (1862) ...	...	W.A.	—	—	—	—	—	B.f.I, 444	M.fr. III, 64.	
C. ulicina, Hooker, Journ. of Bot. I, 257 (1834) ...	...	—	—	T.	—	—	—	B.f.I, 434	M.fr. IX, 137.	
C. pomaderoides, Reisseck in Endl. nov. stirp. dec. 29 (1839) ...	...	W.A.	—	—	—	—	—	B.f.I, 435	M.fr. IX, 137.	
C. pumila; Spyridium, F. v. M., fragm. IX, 137 (1875) ...	...	W.A.	—	—	—	—	—	B.f.I, 435	M.fr. IX, 137.	
C. leucophracta, Schlechtendal, Linnaea XX, 640 (1847) ...	...	—	S.A.	T.	V.	—	—	B.f.I, 435	M.fr. III, 77; IX, 136.	
C. coronata, Reisseck in Lehm. pl. Preiss. II, 288 (1847) ...	...	W.A.	—	—	—	—	—	B.f.I, 436	M.fr. IX, 137.	
C. humilis, F. v. M.; Stenanthemum, Benth., Fl. Austr. I, 436 (1863) ...	...	W.A.	—	—	—	—	—	B.f.I, 436		
C. Waterhousii; Spyridium, F. v. M., fragm. III, 83 (1862) ...	...	—	S.A.	—	—	—	—	B.f.I, 436	M.fr. III, 83.	
C. tridentata, Steudel in Lehm. pl. Preiss. I, 186 (1844) ...	...	W.A.	—	—	—	—	—	B.f.I, 427	M.fr. IX, 137.	
C. divaricata, F. v. M.; Spyridium, Benth. Fl. Austr. I, 427 (1863) ...	...	W.A.	—	—	—	—	—	B.f.I, 427		
C. obcordata, J. Hooker, Fl. Tasm. I, 71 (1860) ...	...	—	—	T.	V.	—	—	B.f.I, 427	M.fr. III, 80.	
C. Lawrencii, J. Hooker, Fl. Tasm. I, 72 (1860) ...	...	—	—	T.	—	—	—	B.f.I, 430	M.fr. IX, 136.	
C. Hookeri, F. v. M.; C. parvifolia, J. Hooker, Fl. Tasm. I, 73 (1860) ...	...	—	S.A.	T.	V.	N.S.W.	—	B.f.I, 428	M.fr. III, 79; IX, 136.	
C. spadicea, F. v. M.; Trymalium, Fenzl in Hueg. enum. 26 (1837) ...	...	W.A.	—	—	—	—	—	B.f.I, 428	M.fr. IX, 137.	
C. globulosa, F. v. M.; Pomad., G. Don in Loud. hort. Brit. 84 (1830) ...	...	W.A.	—	—	—	—	—	B.f.I, 429	M.fr. IX, 137.	
C. obovata, J. Hooker, Fl. Tasm. I, 74 (1860) ...	...	—	S.A.	T.	—	—	—	B.f.I, 429	M.fr. IX, 136.	
C. spathulata; Trymalium, F. v. M. in Trans. Vict. Inst. I, 122 (1855) ...	...	—	S.A.	—	—	—	—	B.f.I, 430	M.fr. IX, 136.	
C. cordata, Turczaninow in Bull. de Mosc. XXXI, 459 (1858) ...	...	W.A.	—	—	—	—	—	B.f.I, 430	M.fr. IV, 25; IX, 137.	
C. rotundifolia; Spyridium, F. v. M., fragm. IV, 25 (1863) ...	...	W.A.	—	—	—	—	—	B.f.I, 430	M.fr. IV, 25.	
C. phlebophylla; Trymalium, F. v. M. in Linnaea XXIX, 272 (1856) ...	...	—	S.A.	—	—	—	—	B.f.I, 430		
C. coactilifolia, F. v. M.; Spyr., Reiss. in Linnaea XXIX, 291 (1856) ...	...	—	S.A.	—	—	—	—	B.f.I, 431	M.fr. IX, 136.	
C. complicata; Spyridium, F. v. M., fragm. III, 78 (1862) ...	...	W.A.	—	—	—	—	—	B.f.I, 431	M.fr. III, 78.	
C. westringifolia, F. v. M.; Trym., Reiss. in pl. Preiss. II, 284 (1847) ...	...	W.A.	—	—	—	—	—	B.f.I, 431		
C. villosa, Turczaninow in Bull. de Mosc. XXXI, 458 (1858) ...	...	W.A.	—	—	—	—	—	B.f.I, 432		
C. pauciflora, Turczaninow in Bull. de Mosc. XXXI, 458 (1858) ...	...	W.A.	—	—	—	—	—	B.f.I, 432		
C. halmaturina; Trym., F. v. M. in Linnaea XXIX, 283 (1856) ...	...	—	S.A.	—	—	—	—	B.f.I, 432		
C. bifida; Trymalium, F. v. M. in Linnaea XXIX, 282 (1856) ...	...	—	S.A.	—	—	—	—	B.f.I, 432	M.fr. IX, 136.	
C. subochreata; Trym., F. v. M. in Trans. Vict. Inst. I, 122 (1855) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f.I, 432	M.fr. III, 82; IX, 136.	
C. vexillifera, Hooker, Journ. Bot. I, 257 (1834) ...	...	—	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.I, 433	M.fr. III, 81; IX, 136.
COLLETTIA, Commerçon in A. L. de Jussieu, gen. 380 (1789). (Discaria, Tetrapasma.)										
C. pubescens, Brogniart in Annal. des sc. nat. X, 366 (1827) ...	...	—	—	T.	V.	N.S.W.	—	B.f.I, 445	M.fr. III, 83; IV, 175.	
GOUANIA, N. J. Jacquin, stirp. Amer. hist. 263, t. 179 (1763).										
G. Hillii, F. v. M., fragm. IX, 163 (1875) ...	...	—	—	—	—	—	Q.	—	M.fr. IX, 163.	
G. Australiana, F. v. M., fragm. IV, 144 (1864) ...	...	—	—	—	—	—	Q.	—	M.fr. IV, 144.	

ARALIACEAE.

Ventenat, Tabl. III (1799). (Umbelliferarum subordo.)

ASTROTTRICHA, De Candolle, Mémoire sur la fam. des Ombellifères 25 (1829).								
A. pterocarpa, Benth. Fl. Austr. III, 379 (1866) ...	—	—	—	—	—	Q.	—	B.f.III, 379 M.fr.VII, 148.
A. floccosa, De Candolle, Mém. des Ombell. 30, t. 5 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 380
A. longifolia, Benth. in Hueg. enum. 55 (1837) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 380 M.fr.VII, 148.
A. ledifolia, De Candolle, Mém. des Ombell. 30, t. 6 (1829) ...	—	—	—	—	V.	N.S.W.	—	B.f.III, 380 M.fr.VII, 148.
A. Hamptoni, F. v. M., fragm. VI, 125 (1868) ...	W.A.	—	—	—	—	—	—	M.fr.VI, 125&250
POROSPERMUM, F. v. M., fragm. VII, 94 (1870).								
P. Michianum, F. v. M., fragm. VII, 95 (1870) ...	—	—	—	—	—	Q.	—	M.fr.VII, 95.
PANAX, Linné, spec. plant. edit. sec. 1513 (1763). (Nothopanax.)								
P. Gunnii, J. Hooker in Lond. Journ. of Bot. VI, 466 (1847) ...	—	—	T.	—	—	—	—	B.f.III, 381
P. Murrayi, F. v. M., fragm. II, 106 (1861) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 381 M.fr.VII, 96.
P. mollis, Benth. Fl. Austr. III, 382 (1866) ...	—	—	—	—	—	Q.	—	B.f.III, 382 M.fr.VII, 96.
P. Macgillivraei, Benth. Fl. Austr. III, 382 (1866) ...	—	—	—	—	—	Q.	—	B.f.III, 382
P. sambucifolius, Sieber in De Cand. prodr. III, 255 (1825) ...	—	—	T.	V.	N.S.W.	Q.	—	B.f.III, 382 M.fr.VII, 95.
P. cephalobotrys, F. v. M., fragm. II, 83 (1861) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 382 M.fr.II, 83; VII, 95.
P. elegans, C. Moore & F. v. M. in Trans. phil. Inst. Vict. II, 68 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 383
P. Cissodendron, C. Moore & F. v. M., fragm. VII, 96 (1870) ...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 96.
MOTHERWELLIA, F. v. M., fragm. VII, 107 (1871).								
M. haplosciadea, F. v. M., fragm. VII, 107 (1871) ...	—	—	—	—	—	Q.	—	M.fr.VII, 107.
CISSODENDRON, Seemann, Journ. of Bot. II, 303 (1864). (Hedera partly, Irvingia, Kissodendron.)								
C. Australicum, Seemann, Journ. of Bot. II, 303 (1864) ...	—	—	—	—	—	Q.	—	B.f.III, 384 M.fr.IV, 120; V, 18.
HEPTOPLEURUM, Gaertner, de fruct. II, 472, t. 178 (1791). (Paratropia.)								
H. venulosum, Seemann, Journ. of Bot. III, 80 (1865) ...	—	—	—	—	—	Q.	—	B.f.III, 384
BRASSAIA, Endlicher, nov. stirp. decad. I, 89 (1839).								
B. actinophylla, Endlicher, nov. stirp. dec. I, 89 (1839) ...	—	—	—	—	—	Q.	—	B.f.III, 385
MERVTA, R. & G. Forster, char. gen. 119, t. 60 (1776). (Bothryodendron.)								
M. latifolia, Seemann in Bonplandia, 295 (1862) ...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 169.
M. angustifolia, Seemann in Bonplandia, 296 (1862) ...	—	—	—	—	N.S.W.	—	—	M.fr.IX, 169.
MACKINLAYA, F. v. M., fragm. IV, 120 (1864).								
M. macrosciadea, F. v. M., fragm. IV, 120 (1864) ...	—	—	—	—	—	Q.	—	B.f.III, 383 M.fr.II, 108&176; IV, 120.

HYDROCOTYLE, Tournefort, inst. 328, t. 173 (1700).

H. vulgaris, Linné, spec. plant. 234 (1753)...	...	...	...	...	S.A.	—	V.	N.S.W.	Q.	—	B.f. III, 339
H. peduncularis, R. Brown in annal. des sc. phys. VI, 62, t. 61 (1820)	—	—	T.	—	—	—	—	—	—	—	B.f. III, 339
H. hirta, R. Brown in annal. des sc. phys. VI, 64 (1820)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f. III, 339
H. laxiflora, De Candolle, prodr. IV, 61 (1830)	...	...	...	...	...	S.A.	—	V.	N.S.W.	Q.	B.f. III, 340
H. rhombifolia, F. v. M., fragm. VII, 147 (1871)...	...	...	...	...	—	—	—	—	N.S.W.	—	—
H. pedicellosa, F. v. M., fragm. IV, 182 (1864)	...	...	...	...	—	—	—	—	N.S.W.	Q.	B.f. III, 341
H. tripartita, R. Brown in annal. des sc. phys. VI, 69, t. 61 (1820)	—	...	...	...	S.A.	T.	V.	N.S.W.	Q.	—	B.f. III, 341
H. plebeia, R. Brown in annal. des sc. phys. VI, 46, t. 60 (1820)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 341
H. pterocarpa, F. v. M. in Transact. Vict. Inst. 126 (1855)	...	...	...	...	—	—	T.	V.	—	—	B.f. III, 342
H. geraniifolia, F. v. M. in Transact. Vict. Inst. 126 (1855)	...	...	...	...	—	—	—	V.	N.S.W.	—	B.f. III, 342
H. medicaginoides, Turczaninow in Bull. Mosc. XXII, 27 (1849)	...	...	...	...	W.A.	S.A.	—	V.	—	—	B.f. III, 342
H. muriculata, Turczaninow in Bull. Mosc. 28 (1849)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 342
H. callicarpa, Bunge in Lehm. pl. Preiss. I, 283 (1844)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f. III, 343
H. scutellifera, Benthani, Fl. Austr. III, 343 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 343
H. hispida, Bunge in Lehm. pl. Preiss. I, 283 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 343
H. trachycarpa, F. v. M. in Schlecht. Linnaea XXV, 394 (1852)	—	...	...	...	—	S.A.	—	—	N.S.W.	—	B.f. III, 343
H. pilifera, Turczaninow in Bull. Mosc. XXII, 26 (1849)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 344
H. capillaris, F. v. M., fragm. IV, 178 (1864)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f. III, 344
H. rugulosa, Turczaninow in Bull. Soc. Mosc. XXII, 27 (1849)...	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 344
H. grammotocarpa, F. v. M., fragm. II, 128 (1861)	...	...	...	...	—	—	—	—	—	N.A.	B.f. III, 344
H. diantha, De Candolle, prodr. IV, 63 (1830)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 345
H. lemnoides, Benthani, Fl. Austr. III, 345 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 345
H. alata, R. Brown in annal. des sc. phys. VI, 75, t. 61 (1820)...	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 345
H. tetragonocarpa, Bunge in Lehm. pl. Preiss. I, 284 (1844)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 345
H. glochidiata, Benthani, Fl. Austr. III, 346 (1866)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 346
H. verticillata, Turczaninow in Bull. Mosc. XXII, 28 (1849)	...	...	...	...	W.A.	—	—	—	—	—	B.f. III, 346
H. Asiatica, Linné, spec. plant. 234 (1753)...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f. III, 346

DIDISCUS, De Candolle in Bot. Mag. t. 2857 (1828).

(Dimetopia, Pritzelia, Huegelia, Cesatia, Hemicarpus.)

D. pusillus, F. v. M., fragm. IX, 47 (1875)	W.A. S.A.	—	V. N.S.W.	—	B. fl. III, 348	M. fr. IX, 47.
D. cyanopetalus, F. v. M., fragm. IX, 46 (1875)	W.A. S.A.	—	V. N.S.W.	—	B. fl. III, 348	M. fr. I, 231.
D. eriocarpus, F. v. M., fragm. IX, 46 (1875)	W.A. S.A.	—	—	—	B. fl. III, 348	M. fr. IX, 46.
D. villosus, F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)	—	—	—	—	N.A.	B. fl. III, 349
D. coeruleus, De Candolle in Hook. Bot. Mag. 2875 (1828)	W.A.	—	—	—	B. fl. III, 349	M. fr. IX, 47.
D. pilosus, Benthani in Hueg. enum. 54 (1837)	W.A. S.A.	T.	V.	N.S.W.	Q.	B. fl. III, 349
D. glaucifolius, F. v. M. in Schlecht. Linnaea XXV, 395 (1852)	—	S.A.	—	N.S.W.	—	B. fl. III, 350
D. glandulosus, F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)	—	—	—	—	N.A.	B. fl. III, 350
D. incisus, Hooker, Bot. Magaz. 2875 (1828)	—	—	—	N.S.W.	Q.	B. fl. III, 350
D. procumbens, F. v. M. in Proc. Roy. Soc. Tasm. III, 237 (1857)	—	—	—	N.S.W.	Q.	B. fl. III, 350
D. hemicarpus, F. v. M. in Trans. Bot. Soc. Edinb. VII, 491 (1857)	—	—	—	—	N.A.	B. fl. III, 351
D. humilis, J. Hooker in icon. pl. t. 304 (1841)	—	—	T.	V.	N.S.W.	B. fl. III, 351

TRACHYMENE, Rudge in Transact. Linn. Soc. X, 300 (1810).

(Siebera, Fischera, Platysace, Platycarpidium.)

T. compressa, Rudge in Transact. Linn. Soc. X, 300 (1810)	...	W.A.	—	—	—	—	B.f.III, 352
T. juncea, Bunge in Lehm. pl. Preiss. I, 286 (1844)	...	W.A.	—	—	—	—	B.f.III, 353
T. haplosciadea, F. v. M.; Siebera, Benth., Fl. Austr. III, 353 (1866)	...	W.A.	—	—	—	—	B.f.III, 353
T. cirrosa, F. v. M.; Platysace, Bunge in pl. Preiss. I, 285 (1844)	—	S.A.	—	V.	—	—	B.f.III, 354
T. heterophylla, F. v. M., First Gen. Rep. 13 (1853)	...	W.A.	S.A.	—	—	—	B.f.III, 354
T. tenuissima, F. v. M.; Siebera, Benth., Fl. Austr. III, 354 (1866)	...	W.A.	—	—	—	—	B.f.III, 354
T. dissecta, F. v. M.; Siebera, Benth., Fl. Austr. III, 354 (1866)	...	W.A.	—	—	—	—	B.f.III, 354
T. commutata, Turczaninow in Bull. Mosc. XXII, 30 (1849)	...	W.A.	—	—	—	—	B.f.III, 355
T. effusa, Turczaninow in Bull. Mosc. XXII, 31 (1849)	...	W.A.	—	—	—	—	B.f.III, 355
T. deflexa, Turczaninow in Bull. Mosc. XXII, 31 (1849)	...	W.A.	—	—	—	—	B.f.III, 355
T. valida; Platycarpidium, F. v. M. in Kew. Misc. IX, 310 (1857)	—	—	—	—	—	Q.	B.f.III, 355
T. ericoides, Sieber in De Candolle, prodr. IV, 73 (1830)	...	—	—	V.	N.S.W.	Q.	B.f.III, 356
T. linearis, Sprengel, spec. umbellif. 7 (1818)	...	—	—	—	N.S.W.	—	B.f.III, 356
T. Billardieri, F. v. M.; Siebera, Benth., Fl. Austr. III, 356 (1866)	...	—	—	V.	N.S.W.	—	B.f.III, 356
T. Stephensonii, Turczaninow in Bull. de Mosc. XX, 170 (1847)	—	—	—	—	N.S.W.	—	B.f.III, 357

XANTHOSIA, Rudge in Transact. Linn. Soc. X, 361 (1810).

(Leucolaena, Schoenolaena, Pentapeltis.)

X. juncea, Benth., Fl. Austr. III, 359 (1866) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 359	
X. tenuior, Benth., Fl. Austr. III, 359 (1866) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 359	
X. peltigera, Steudel, nomencl. bot. 791 (1841) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 359	
X. hederifolia, Benth., Fl. Austr. III, 359 (1866) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 359	
X. candida, Steudel, nomencl. bot. 790 (1841) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 259	
X. tridentata, De Candolle, prodr. IV, 75 (1830) ...	...	...	...	—	—	—	V.	N.S.W.	—	B.fl. III, 260	
X. singuliflora, F. v. M., fragm. IV, 184 (1864) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 260	
X. pilosa, Rudge in Transact. Linn. Soc. X. 301, t. 22 (1811) ...	...	...	...	—	—	T.	V.	N.S.W.	Q.	B.fl. III, 260	M.fr. IV, 184.
X. ciliata, Hooker, icon. pl. t. 726 (1848) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 261	
X. pusilla, Bunge in Lehm. pl. Preiss. I, 291 (1844) ...	...	...	...	W.A.	S.A.	T.	V.	—	—	B.fl. III, 261	
X. fruticulosa, Benth., Fl. Austr. III, 361 (1866) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 261	
X. Huegelii, Steudel, nomenclat. bot. 791 (1841) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 362	
X. dissecta, J. Hooker in icon. plant. t. 302 (1841) ...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	B.fl. III, 362	
X. peduncularis, Benth., Fl. Austr. III, 362 (1866) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 362	
X. vestita, Benth., Fl. Austr. III, 363 (1866) ...	...	...	...	—	—	—	—	N.S.W.	—	B.fl. III, 363	
X. Atkinsoniana, F. v. M., fragm. II, 127 (1861) ...	...	...	...	W.A.	—	—	—	N.S.W.	—	B.fl. III, 363	M.fr. II, 127.
X. rotundifolia, De Candolle, prodr. IV, 75 (1830) ...	...	...	...	W.A.	—	—	—	—	—	B.fl. III, 363	

- AZORELLA**, Lamarck, Encycl. méthodiq. I, 344, t. 189 (1783). (Fragosa, Pozoa, Microsciadium, Schizcilema, Oschatzia, Centella partly, Dichopetalum.)
- A. Muellieri, Benth., Fl. Austr. III, 364 (1866) ... — — — V. — — — B.f. III, 364
 A. cuneifolia, F. v. M. in Transact. phil. Inst. Vict. I, 103 (1855) ... — — — V. — — — B.f. III, 365
 A. saxifraga, Benth., Fl. Austr. III, 365 (1866) ... — — — T. — — — B.f. III, 365
 A. dichopetala, Benth., Fl. Austr. III, 365 (1866) ... — — — T. V. — — — B.f. III, 365
- HUANACA**, Cavanilles, icon. VI, 18, t. 528 (1801). (Diplaspis, Pozoopsis.)
- H. hydrocotylea, Benth. & J. Hooker, gen. pl. I, 877 (1867) ... — — — T. V. — — — B.f. III, 366
 H. cordifolia, Benth. & J. Hooker, gen. pl. I, 877 (1867) ... — — — T. — — — B.f. III, 366
- ACTINOTUS**, Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804). (Hemiphues, Eriocalia, Holotome.)
- A. Helianthi, Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804) ... — — — V. N.S.W. Q. — — — B.f. III, 367
 A. leucocephalus, Benth. in Hueg. enum. 56 (1837) ... — — — W.A. — — — — — B.f. III, 368
 A. minor, De Candolle, prodr. IV, 83 (1830) ... — — — — — N.S.W. — — — B.f. III, 368
 A. Gibbonsii, F. v. M., fragm. VI, 23 (1867) ... — — — — — N.S.W. — — — M.fr. VII, 23.
 A. omnifertilis, Benth., Fl. Austr. III, 368 (1866) ... — — — W.A. — — — — — B.f. III, 368
 A. rhomboideus, Benth., Fl. Austr. III, 368 (1866) ... — — — W.A. — — — — — B.f. III, 368
 A. bellidioides, Benth., Fl. Austr. III, 369 (1866) ... — — — T. — — — B.f. III, 369
 A. glomeratus, Benth., Fl. Austr. III, 369 (1866) ... — — — W.A. — — — — — B.f. III, 369
- ERYNGIUM**, Tournefort, inst. 327, t. 173 (1700) from Theophrastus and Dioscorides.
- E. rostratum, Cavanilles, icon. pl. VI, 35, t. 552 (1801) ... — — — W.A. S.A. — — — V. N.S.W. Q. — — — B.f. III, 370
 E. vesiculosum, Labillardière, Nov. Holl. pl. spec. I, 73, t. 98 (1804) ... — — — S.A. T. V. N.S.W. Q. — — — B.f. III, 370 M.fr. VII, 148.
 E. plantagineum, F. v. M. in Pap. Roy. Soc. Tasm. III, 235 (1857) ... — — — S.A. — — — N.S.W. Q. — — — B.f. III, 371 M.fr. VII, 148.
 E. expansum, F. v. M. in Pap. Roy. Soc. Tasm. III, 236 (1857) ... — — — — — N.S.W. Q. — — — B.f. III, 371 M.fr. VII, 148.
- APIUM**, Tournefort, inst. 305, t. 160 (1700).
- A. prostratum, Labillardière, relat. du voy. I, 141 (1799) ... — — — W.A. S.A. T. V. N.S.W. Q. — — — B.f. III, 372 M.fr. VII, 148.
 A. leptophyllum, F. v. M. in Benth. Fl. Austr. III, 372 (1866) ... — — — — — V. N.S.W. Q. — — — B.f. III, 372 M.fr. VII, 148.
- SIUM**, Tournefort, inst. 308, t. 162 (1700), from C. Bauhin (1623).
- S. latifolium, C. Bauhin, pinax 154 (1623) ... — — — — — S.A. — — — V. N.S.W. Q. — — — —
 S. angustifolium, Linné, spec. plant. ed. sec. 1672 (1763) ... — — — — — N.S.W. — — — — —
- SESELI**, Rivinus in Rupp. Fl. Jen. 267 (1718).
- S. Harveyanum, F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855) ... — — — — — V. — — — — — B.f. III, 373
 S. algens, F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855) ... — — — — — V. — — — — — B.f. III, 373
- CRANTZIA**, Nuttall, Gen. of North Amer. plants I, 177 (1818). (Non Scopoli, 1777, Crantziola, F. v. M.)
- C. lineata, Nuttall, Gen. of North Amer. pl. I, 178 (1818) ... — — — — — S.A. T. V. N.S.W. Q. — — — B.f. III, 374
- OENANTHE**, Tournefort, inst. 312, t. 166 (1700).
- O. Javanica, De Candolle, prodr. IV, 138 (1830) ... — — — — — — — — — Q. — — — M.fr. V, 182.
- ACIPHYLLA**, R. & G. Forster, char. gen. 135, t. 68 (1776). (Anisotome, Calosciadium, Gingidium partly, Ligustici subgenus.)
- A. simplicifolia, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ... — — — — — V. N.S.W. — — — B.f. III, 375
 A. glacialis, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ... — — — — — V. N.S.W. — — — B.f. III, 375
 A. procumbens, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ... — — — — — T. — — — — — B.f. III, 375
- DAUCUS**, Tournefort, inst. 307, t. 161 (1700), from l'Ecluse (1576).
- D. brachiatus, Sieber in De Cand., prodr. IV, 214 (1830) ... — — — W.A. S.A. T. V. N.S.W. Q. — — — B.f. III, 376
- OREOMYRRHIS**, Endlicher, gen. plantar. 787 (1839).
- O. andicola, Endlicher, gen. plantar. 787 (1839) ... — — — — — S.A. T. V. N.S.W. — — — B.f. III, 377 M.fr. VIII, 185.
 O. pulvinifica, F. v. M., fragm. VIII, 185 (1874) ... — — — — — V. N.S.W. — — — — — M.fr. VIII, 185.

SYNPETALEAE PERIGYNÆ.

F. v. M. in Woolls's plants of the neighb. of Sydney, 27 (1880).

AQUIFOLIACEAE.

De Candolle, Théor. élém. I, 217 (1813).

- BYRONIA**, Endlicher in den Ann. des Wien. Mus. I, 184 (1836).
- B. Arnhemensis, F. v. M., fragm. II, 119 (1861) ... — — — — — — — — — N.A. B.f. I, 397 M.fr. II, 119.
- ILEX**, Linné, gen. plant. 33 (1737), from C. Bauhin (1623).
- I. peduncularis, F. v. M., fragm. VII, 105 (1871) ... — — — — — — — — — Q. — — — M.fr. VII, 105.

OLACINEAE.

Mirbel in Bull. de la Soc. philomat. 377 (1813).

- XIMENIA**, Plumier, nova plant. Amer. genera 6, t. 21 (1703).
- X. Americana, Linné, spec. plant. 1193 (1753) ... — — — — — — — — — Q. N.A. B.f. I, 391 M.fr. VI, 3; IX, 150.
- OLAX**, Linné, amoen. acad. 387 (1747). (Spermaxyrum, Lopadocalyx.)
- O. phyllanthi, R. Brown, prodr. 358 (1810) ... — — — — — W.A. — — — — — B.f. I, 392 M.fr. IX, 151.
 O. retusa, F. v. M. in Benth. Fl. Austr. I, 392 (1863) ... — — — — — N.S.W. Q. — — — B.f. I, 392 M.fr. IX, 151.
 O. stricta, R. Brown, prodr. 358 (1810) ... — — — — — N.S.W. Q. — — — B.f. I, 392 M.fr. IX, 151.
 O. Benthamiana, Miquel in Lehmann, pl. Preiss. I, 228 (1844) ... — — — — — W.A. S.A. — — — — — N.A. B.f. I, 393 M.fr. IX, 151.
 O. aphylla, R. Brown, prodr. 358 (1810) ... — — — — — — — — — N.A. B.f. I, 393 M.fr. IX, 151.
- CANSJERA**, A. L. de Jussieu, gen. pl. 448 (1789).
- C. leptostachya, Benth. in Hook. Lond. Journ. of Bot. II, 231 (1843) ... — — — — — — — — — Q. — — — B.f. I, 394 M.fr. VI, 3.
- OPILIA**, Roxburgh, pl. Corom. II, 31, t. 158 (1799).
- O. amentacea, Roxburgh, pl. Corom. II, 31, t. 158 (1799) ... — — — — — — — — — Q. N.A. B.f. I, 394 M.fr. VI, 4; IX, 150.
- PHLEBOCALYMMMA**, Griffith in Benth. & J. Hook. gen. pl. I, 333 (1862). (Phlebocalymma).
- P. lobospora, F. v. M., fragm. IX, 151 (1875) ... — — — — — — — — — Q. — — — M.fr. IX, 151.

- PENNANTIA, R. & G. Forster, char. gen. 133, t. 67 (1776).
 P. Cunninghamii, Miens in Ann. of nat. hist. sec. ser. IX, 491 (1852) — — — — N.S.W. Q. — B.f.I, 395 M.fr. VI, 4; IX, 150.
 P. Endlicheri, Reisseck in Schlecht. Linnaea XVI, 341, t. 13 (1843) — — — — N.S.W. — — — — M.fr. IX, 150.
 APODYTES, E. Meyer in Hook. Journ. of Bot. III, 155 (1841).
 A. brachystylis, F. v. M., fragm. IX, 149 (1875) ... — — — — — Q. — — — — M.fr. IX, 149.
 VILLAREZIA, Ruiz & Pavon, flor. Peruv. et Chil. prodr. 35 (1794).
 V. Moorei, F. v. M. in Benth. Fl. Austr. I, 396 (1863) ... — — — — N.S.W. — — — — B.f.I, 395
 V. Smythii, F. v. M., fragm. V, 156 (1866) ... — — — — — Q. — — — — M.fr. IX, 150.
 GOMPHANDRA, Wallich, Catal. n. 3718 & 7204 (1832).
 G. Australiana, F. v. M., fragm. VI, 3 (1877) ... — — — — — Q. — — — — M.fr. VI, 3&253; IX, 150.

ELAEAGNEAE.

R. Brown, prodr. I, 350 (1810.)

- ELAEAGNUS, Tournefort, coroll. 53, t. 459 (1703) from Camerarius (1586).
 E. latifolia, Linné, spec. plant. 121 (1753) ... — — — — — Q. — — — — M.fr. VI, 39 M.fr. IX, 190.

SANTALACEAE.

R. Brown, prodr. I, 350 (1810).

- THESIUM, Linné, gen. pl. 60 (1737).
 T. australe, R. Brown, prodr. 353 (1810) ... — — — — — T. V. N.S.W. Q. — — — — M.fr. VI, 212 M.fr. IX, 3.
 SANTALUM, Linné, gen. ed. sec., 165 (1742), from C. Bauhin (1623). (Fusanus, Eucarya).
 S. lanceolatum, R. Brown, prodr. 356 (1810) ... — — — — — W.A. S.A. — — — — — N.S.W. Q. N.A. B.f. VI, 214 M.fr. I, 85; VIII, 11; IX, 3.
 S. ovatum, R. Brown, prodr. 355 (1810) ... — — — — — — — — — — N.A. B.f. VI, 214
 S. obtusifolium, R. Brown, prodr. 356 (1810) ... — — — — — V. N.S.W. — — — — — B.f. VI, 215 M.fr. VIII, 11.
 S. acuminatum, A. de Candolle, prodr. XIV, 684 (1856) ... — — — — — W.A. S.A. — — — — — V. N.S.W. — — — — — B.f. VI, 216 M.fr. IX, 169; XI, 20.
 S. persicarium, F. v. M. in Transact. Vict. Inst. 41 (1854) ... — — — — — W.A. S.A. — — — — — V. N.S.W. — — — — — B.f. VI, 216 M.fr. I, 86; IX, 3; XI, 20.
 S. cygnorum, Miquel in Lehm. pl. Preiss. I, 615 (1845) ... — — — — — W.A. — — — — — — — — — — B.f. VI, 217 M.fr. VIII, 11.
 S. crassifolium, A. de Candolle, prodr. XIV, 685 (1856) ... — — — — — — — — — — N.S.W. — — — — — B.f. VI, 217
 CHORETRUM, R. Brown, prodr. 354 (1810).
 C. glomeratum, R. Brown, prodr. 354 (1810) ... — — — — — W.A. S.A. — — — — — V. N.S.W. — — — — — B.f. VI, 218
 C. chrysanthum, F. v. M. in Transact. phil. Soc. Vict. I, 23 (1854) ... — — — — — S.A. — — — — — V. N.S.W. — — — — — — — — — — M.fr. VIII, 11; XI, 138.
 C. spicatum, F. v. M., fragm. I, 21 (1858) ... — — — — — S.A. — — — — — V. N.S.W. — — — — — B.f. VI, 218 M.fr. VIII, 11.
 C. lateriflorum, R. Brown, prodr. 354 (1810) ... — — — — — — — — — — V. N.S.W. — — — — — B.f. VI, 219
 C. Candollei, F. v. M. in Bentham's Fl. Austr. VII, 219 (1873) ... — — — — — — — — — — N.S.W. Q. — — — — — B.f. VI, 219
 LEPTOMERIA, R. Brown, prodr. 353 (1810).
 L. spinosa, A. de Candolle, prodr. XIV, 678 (1856) ... — — — — — W.A. — — — — — — — — — — B.f. VI, 220 M.fr. VIII, 10.
 L. Preissiana, A. de Candolle, prodr. XIV, 678 (1856) ... — — — — — W.A. — — — — — — — — — — B.f. VI, 221 M.fr. VIII, 10.
 L. pauciflora, R. Brown, prodr. 354 (1810) ... — — — — — W.A. — — — — — — — — — — B.f. VI, 221 M.fr. VIII, 10.
 L. scrobiculata, R. Brown, prodr. 354 (1810) ... — — — — — W.A. — — — — — — — — — — B.f. VI, 221 M.fr. VIII, 10.
 L. acida, R. Brown, prodr. 353 (1810) ... — — — — — — — — — — V. N.S.W. Q. — — — — — B.f. VI, 222 M.fr. VIII, 10.
 L. Billardieri, R. Brown, prodr. 354 (1810) ... — — — — — — — — — — T. — — — — — N.S.W. Q. — — — — — B.f. VI, 222 M.fr. VIII, 10.
 L. aphylla, R. Brown, prodr. 354 (1810) ... — — — — — — — — — — S.A. — — — — — V. N.S.W. — — — — — B.f. VI, 222 M.fr. VIII, 10.
 L. glomerata, F. v. M. in J. Hook. Tasm. II, 370 (1860) ... — — — — — — — — — — T. — — — — — — — — — — B.f. VI, 223 M.fr. VIII, 10.
 L. squarrola, R. Brown, prodr. 354 (1810) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 223 M.fr. VIII, 11.
 L. Cunninghamii, Miquel in Lehm. pl. Preiss. I, 611 (1845) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 223 M.fr. VIII, 10.
 L. empetrifolius, Miquel in Lehm. pl. Preiss. I, 610 (1845) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 224
 L. axillaris, R. Brown, prodr. 354 (1810) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 224
 L. laxa, Miquel in Lehm. pl. Preiss. I, 612 (1845) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 224
 L. obovata, Miquel in Lehm. pl. Preiss. I, 613 (1845) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 224
 OMPHACOMERIA, Endlicher, gen. pl. 326 (1838).
 O. acerba, A. de Candolle, prodr. XIV, 681 (1856) ... — — — — — — — — — — V. N.S.W. — — — — — B.f. VI, 224
 ANTHOBOLUS, R. Brown, prodr. 357 (1810).
 A. filifolius, R. Brown, prodr. 357 (1810) ... — — — — — — — — — — — — — — — — N.A. B.f. VI, 226 M.fr. IX, 3.
 A. triquetus, R. Brown, prodr. 357 (1810) ... — — — — — — — — — — — — — — — — Q. — — — — — B.f. VI, 226
 A. foveolatus, F. v. M., fragm. I, 212 (1859) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 226 M.fr. II, 181; IX, 3; XI, 20.
 A. leptomerioides, F. v. M., fragm. I, 21 (1858) ... — — — — — — — — — — — — — — — — Q. — — — — — B.f. VI, 227 M.fr. IX, 3.
 A. exocarpoides, F. v. M., fragm. IX, 3 (1875) ... — — — — — S.A. — M.fr. IX, 3.
 EXOCARPOS, Labillardière, Relat. du voy. à la rech. de La Pérouse I, 155 (1798). (Exocarpus.)
 E. latifolia, R. Brown, prodr. 356 (1810) ... — — — — — — — — — — — — — — — — N.S.W. Q. N.A. B.f. VI, 228 M.fr. VIII, 10.
 E. odorata, A. de Candolle, prodr. XIV, 689 (1856) ... — — — — — W.A. — — — — — — — — — — — — — — — — B.f. VI, 228 M.fr. VIII, 10.
 E. cupressiformis, Labillardière, voy. I, 155, t. 14 (1798) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — — — — — B.f. VI, 229
 E. spartea, R. Brown, prodr. 356 (1810) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. — — — — — B.f. VI, 229 M.fr. VIII, 10; XI, 20. [20.
 E. aphylla, R. Brown, prodr. 357 (1810) ... — — — — — W.A. S.A. — — — — — V. N.S.W. Q. — — — — — B.f. VI, 230 M.fr. VIII, 10; IX, 169; XI,
 E. homaloclada, C. Moore & F. v. M., fragm. VIII, 9 (1871) ... — — — — — — — — — — — — — — — — N.S.W. — — — — — B.f. VI, 230 M.fr. VIII, 9.
 E. phyllanthoides, Endlicher, prodr. ins. Norf. 46 (1833) ... — — — — — — — — — — — — — — — — N.S.W. — — — — — — — — — — M.fr. IX, 169.
 E. stricta, R. Brown, prodr. 357 (1810) ... — — — — — S.A. T. V. N.S.W. — — — — — B.f. VI, 230 M.fr. IX, 169.
 E. humifusa, R. Brown, prodr. 356 (1810) ... — — — — — — — — — — T. — — — — — — — — — — B.f. VI, 231
 E. nana, J. Hooker in Lond. Journ. of Bot. VI, 281 (1847) ... — — — — — T. V. — — — — — — — — — — B.f. VI, 230

LORANTHACEAE.

A. L. de Jussieu, in Ann. du Mus. XII, 292 (1808).

- VISCUM, Tournefort, inst. 609, t. 380 (1700), from Camerarius (1586).
 V. orientale, Willdenow, spec. plant. IV, 737 (1805) ... — — — — — — — — — — Q. — — — — — B.f. III, 396
 V. angulatum, Heyne in De Cand., prodr. IV, 283 (1830) ... — — — — — — — — — — Q. — — — — — B.f. III, 396
 V. articulatum, N. L. Burmann, Fl. Ind. 311 (1768) ... — — — — — — — — — — N.S.W. Q. — — — — — B.f. III, 396

NOTOTHIXOS, Oliver in Journ. Linn. Soc. VII, 104 (1865).									
N. incanus, Oliver in Journ. Linn. Soc. VII, 104 (1865)...	—	—	—	—	N.S.W.	Q.	—	B.f.III,397	M.fr.II,109,176;IV,173.
LORANTHUS, Linné, syst. nat. ed. sec. 22 (1740).									
L. celastroides, Sieber in Roem. & Schult. syst. veg. VII, 163 (1829)	—	—	—	V.	N.S.W.	Q.	—	B.f.III,389	M.fr.VI,252.
L. Bidwillii, Benth. Fl. Austr. III, 390 (1866)...	—	—	—	—	—	Q.	—	B.f.III,390	
L. myrtifolius, Cunningham in Benth. Fl. Austr. III, 390 (1866)	—	—	—	—	N.S.W.	Q.	—	B.f.III,390	
L. longiflorus, Desrousseaux in Lam. Encycl. méth. IV, 598 (1796)	—	—	—	—	N.S.W.	Q.	N.A.	B.f.III,390	
L. angustifolius, R. Brown in Benth. Fl. Austr. III, 390 (1866)	—	—	—	—	—	—	—	B.f.III,390	
L. dictyophlebus, F. v. M., Rep. Burdek. Exped. 14 (1858)	—	—	—	—	N.S.W.	Q.	—	B.f.III,391	
L. alyxifolius, F. v. M. in Benth. Fl. Austr. III, 391 (1866)	—	—	—	—	N.S.W.	Q.	—	B.f.III,391	
L. odontocalyx, F. v. M. in Benth. Fl. Austr. III, 391 (1866)	—	—	—	—	—	—	Q.	N.A.	B.f.III,391
L. linearifolius, Hooker in Mitchell, Trop. Austr. 102 (1848)	—	—	—	—	—	—	Q.	—	B.f.III,391
L. Exocarpi, Behr in Schlecht. Linnaea XX, 624 (1847)	—	—	—	—	W.A.	S.A.	—	—	—
L. acacioides, Cunningham in Benth. Fl. Austr. III, 392 (1866)	—	—	—	—	—	—	—	N.A.	B.f.III,392
L. signatus, F. v. M. in Benth. Fl. Austr. III, 392 (1866)	—	—	—	—	—	—	Q.	N.A.	B.f.III,392
L. maytenifolius, A. Gray, Bot. Wilk. Expl. Exped. I, 739, t. 99 (1854)	—	—	—	—	N.S.W.	—	—	B.f.III,393	
L. sanguineus, F. v. M., fragm. I, 177 (1859)	—	—	—	—	—	—	Q.	N.A.	B.f.III,393
L. bifurcatus, Benth. Fl. Austr. III, 393 (1866)	—	—	—	—	—	—	—	N.A.	B.f.III,393
L. linophyllus, Fenzl in Huegel. enum. 56 (1837)...	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.
L. pendulus, Sieber in De Candolle, prodr. IV, 294 (1830)	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.
L. Quandang, Lindley in Mitch. Three Exped. II, 69 (1838)	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.
L. grandibracteus, F. v. M., Rep. Burdek. Exped. 14 (1858)	—	—	—	—	—	S.A.	—	—	Q.
ATKINSONIA, F. v. Mueller, fragm. V, 34 (1865).									
A. ligustrina, F. v. M., fragm. V, 34 (1865)	—	—	—	—	N.S.W.	—	—	B.f.III,388	M.fr.II,130;VI,252.
NUYTSIA, R. Brown in Journ. of the geogr. Soc. I, 17 (1831).									
N. floribunda, R. Brown in Journ. geogr. Soc. I, 17 (1831)	—	—	—	—	W.A.	—	—	B.f.III,387	M.fr.VI,252.

PROTEACEAE.

A. L. de Jussieu, gen. pl. 78 (1789).

PETROPHILA, R. Brown in Transact. Linn. Soc. X, 67 (1809).									
P. teretifolia, R. Brown in Transact. Linn. Soc. X, 63 (1809)	—	—	—	—	—	—	—	B.f.V,321	M.fr.VI,245.
P. longifolia, R. Brown, prot. nov. 5 (1830)	—	—	—	—	—	—	—	B.f.V,322	M.fr.VI,255.
P. media, R. Brown, prot. nov. 5 (1830)	—	—	—	—	—	—	—	B.f.V,322	M.fr.VI,255.
P. acicularis, R. Brown in Transact. Linn. Soc. X, 69 (1809)	—	—	—	—	—	—	—	B.f.V,323	M.fr.VI,245.
P. megalostegia, F. v. M., fragm. X, 61 (1876)	—	—	—	—	—	—	—	—	M.fr.X,61.
P. linearis, R. Brown, prot. nov. 6 (1830)	—	—	—	—	—	—	—	B.f.V,323	M.fr.VI,243.
P. anceps, R. Brown, prot. nov. 5 (1830)	—	—	—	—	—	—	—	—	M.fr.VI,243;X,47.
P. heterophylla, Lindley, Bot. Regist. XXV, App. XXXV (1839)	—	—	—	—	—	—	—	B.f.V,324	M.fr.VI,244.
P. biloba, R. Brown, prot. nov. 7 (1830)	—	—	—	—	—	—	—	B.f.V,324	M.fr.VI,255.
P. propinqua, R. Brown, prot. nov. 7 (1830)	—	—	—	—	—	—	—	B.f.V,325	
P. squamata, R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	—	—	—	B.f.V,325	M.fr.VI,244.
P. colorata, Meissner in Lehm. pl. Preiss. II, 246 (1847)	—	—	—	—	—	—	—	B.f.V,326	
P. striata, R. Brown, prot. nov. 6 (1830)	—	—	—	—	—	—	—	B.f.V,326	M.fr.VI,244.
P. divaricata, R. Brown, prot. nov. 7 (1830)	—	—	—	—	—	—	—	B.f.V,326	M.fr.VI,344.
P. serruriae, R. Brown, prot. nov. 6 (1830)	—	—	—	—	—	—	—	B.f.V,327	M.fr.VI,243.
P. inconspicua, Meissner in Hook. Kew Misc. VII, 68 (1855)	—	—	—	—	—	—	—	B.f.V,327	M.fr.VI,243.
P. trifida, R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	—	—	—	B.f.V,328	M.fr.VI,244.
P. carduacea, Meissner in Hook. Kew Misc. IV, 182 (1852)	—	—	—	—	—	—	—	B.f.V,328	M.fr.VI,244.
P. Shuttleworthiana, Meissner in Lehm. pl. Preiss. II, 246 (1847)	—	—	—	—	—	—	—	B.f.V,329	M.fr.VI,245.
P. macrostachya, R. Brown, prot. nov. 7 (1830)	—	—	—	—	—	—	—	B.f.V,329	M.fr.VI,245.
P. diversifolia, R. Brown in Transact. Linn. Soc. X, 70 (1809)...	—	—	—	—	—	—	—	B.f.V,329	M.fr.VI,244.
P. biternata, Meissner in Hook. Kew Misc. VII, 69 (1855)	—	—	—	—	—	—	—	B.f.V,330	M.fr.VI,245.
P. plumosa, Meissner in Hook. Kew Misc. VII, 69 (1855)	—	—	—	—	—	—	—	B.f.V,330	M.fr.VI,244.
P. ericifolia, R. Brown, prot. nov. 5 (1830)	—	—	—	—	—	—	—	B.f.V,331	M.fr.VI,243.
P. chrysanthra, Meissner in Hook. Kew Misc. VII, 68 (1855)	—	—	—	—	—	—	—	B.f.V,331	M.fr.VI,243.
P. pedunculata, R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	N.S.W.	Q.	—	B.f.V,332	M.fr.VI,242.
P. pulchella, R. Brown in Transact. Linn. Soc. X, 69 (1809)	—	—	—	—	N.S.W.	—	—	B.f.V,332	M.fr.VI,242.
P. sessilis, Sieber in Roem. & Schult. mant. 262 (1822)	—	—	—	—	N.S.W.	Q.	—	B.f.V,332	
P. fastigiata, R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	—	—	—	B.f.V,333	M.fr.VI,245.
P. seminuda, Lindley, Bot. Regist. XXV, App. XXXIV (1839)	—	—	—	—	—	—	—	B.f.V,333	M.fr.VI,243.
P. circinata, Kippist in Hook. Kew Misc. VII, 67 (1855)	—	—	—	—	—	—	—	B.f.V,333	M.fr.VI,244.
P. Drummondii, Meissner in Lehm. pl. Preiss. I, 496 (1845)	—	—	—	—	—	—	—	B.f.V,334	M.fr.VI,243.
P. crispata, R. Brown, prot. nov. 6 (1830)...	—	—	—	—	—	—	—	B.f.V,334	
P. rigida, R. Brown in Transact. Linn. Soc. X, 69 (1809)	—	—	—	—	—	—	—	B.f.V,334	M.fr.VI,245.
P. multiseeta, F. v. M., fragm. VI, 242 (1868)	—	—	—	—	S.A.	—	—	B.f.V,335	M.fr.VI,242.
P. conifera, Meissner in Hook. Kew Misc. VII, 67 (1855)	—	—	—	—	—	—	—	B.f.V,335	M.fr.VI,245.
P. semifurcata, F. v. M. in Benth. Fl. Austr. V, 335 (1870)	—	—	—	—	—	—	—	B.f.V,335	M.fr.X,47.
ISOPOGON, R. Brown in Transact. Linn. Soc. X, 70 (1809).									
I. latifolius, R. Brown, prot. nov. 8 (1830)...	—	—	—	—	—	—	—	B.f.V,338	M.fr.VI,237.
I. cuneatus, R. Brown in Transact. Linn. Soc. X, 73 (1809)	—	—	—	—	—	—	—	B.f.V,339	M.fr.VI,238.
I. linearis, Meissner in Hook. Kew Misc. VII, 69 (1855)	—	—	—	—	—	—	—	B.f.V,339	M.fr.VI,236.
I. polyccephalus, R. Brown in Transact. Linn. Soc. X, 73 (1809)	—	—	—	—	—	—	—	B.f.V,339	M.fr.VI,236.
I. attenuatus, R. Brown in Transact. Linn. Soc. X, 73 (1809)	—	—	—	—	—	—	—	B.f.V,340	M.fr.VI,237.
I. sphaerocephalus, Lindl., Bot. Regist. XXV, App. XXXIV (1839)	—	—	—	—	—	—	—	B.f.V,340	M.fr.VI,236.
I. uncinatus, R. Brown, prot. nov. 8 (1830)	—	—	—	—	—	—	—	B.f.V,341	M.fr.VI,237.
I. buxifolius, R. Brown in Transact. Linn. Soc. X, 74 (1809)	—	—	—	—	—	—	—	B.f.V,341	M.fr.VI,237.
I. axillaris, R. Brown in Transact. Linn. Soc. X, 74 (1809)	—	—	—	—	—	—	—	B.f.V,341	M.fr.VI,237.
I. tridens, F. v. M., fragm. VI, 239 (1868)...	—	—	—	—	—	—	—	B.f.V,342	M.fr.VI,239.

I. Baxteri, R. Brown, prot. nov. 9 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 342	M.fr. VI, 240.
I. roseus, Lindley, Bot. Regist. Misc. n. 37 (1842) ...	W.A.	—	—	—	—	—	B.f.V, 343	M.fr. VI, 240
I. adenanthoides, Meissner in Hook. Kew Misc. VII, 69 (1855) ...	W.A.	—	—	—	—	—	B.f.V, 343	M.fr. VI, 241.
I. trilobus, R. Brown in Transact. Linn. Soc. X, 72 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 343	M.fr. VI, 239.
I. tripartitus, R. Brown, prot. nov. 8 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 344	
I. longifolius, R. Brown in Transact. Linn. Soc. X, 73 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 344	M.fr. VI, 237.
I. Drummondii, Benth, Fl. Austr. V, 344 (1870) ...	W.A.	—	—	—	—	—	B.f.V, 344	M.fr. VI, 241.
I. heterophyllus, Meissner in Lehm. pl. Preiss. I, 504 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 345	M.fr. VI, 241.
I. villosus, Meissner in De Candolle, prodr. XIV, 277 () ...	—	—	—	—	—	—	—	M.fr. VI, 241.
I. teretifolius, R. Brown in Transact. Linn. Soc. X, 71 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 345	M.fr. VI, 241.
I. anethifolius, R. Brown in J. Knight, proteac. 94 (1809) ...	W.A.	—	—	N.S.W.	—	—	B.f.V, 346	M.fr. VI, 238.
I. petiolaris, Cunningham in R. Br., prot. nov. 8 (1830) ...	—	—	—	N.S.W.	—	—	B.f.V, 346	
I. anemonifolius, R. Brown in J. Knight, prot. 93 (1809) ...	—	—	—	N.S.W.	Q.	—	B.f.V, 347	M.fr. VI, 238; X, 90.
I. ceratophyllus, R. Brown in Transact. Linn. Soc. X, 72 (1809) ...	—	S.A.	T.	N.S.W.	—	—	B.f.V, 347	M.fr. VI, 238.
I. asper, R. Brown, prot. nov. 8 (1830) ...	W.A.	S.A.	T.	V.	—	—	B.f.V, 348	M.fr. VI, 240.
I. crithmifolius, F. v. M., fragm. VI, 239 (1868) ...	W.A.	—	—	—	—	—	B.f.V, 348	M.fr. VI, 239.
I. formosus, R. Brown in Transact. Linn. Soc. X, 72 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 349	M.fr. VI, 240.
I. divergens, R. Brown, prot. nov. 7 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 349	M.fr. VI, 241.
I. scabriusculus, Meissner in Hook. Kew Misc. IV, 182 (1852) ...	W.A.	—	—	—	—	—	B.f.V, 349	M.fr. VI, 240.
ADENANTHOS, Labillardière, Nov. Holl. pl. specim. I, 28, t. 36 (1804).								
A. Detmoldi, F. v. M., fragm. VIII, 149 (1874) ...	W.A.	—	—	—	—	—	—	M.fr. VIII, 149.
A. barbigerus, Lindley, Bot. Regist. XXV, App. XXXVI (1839) ...	W.A.	—	—	—	—	—	B.f.V, 351	M.fr. VI, 205; VIII, 149
A. obovatus, Labillardière, Nov. Holl. pl. spec. I, 29, t. 37 (1804) ...	W.A.	—	—	—	—	—	B.f.V, 352	
A. cuneatus, Labillardière, Nov. Holl. pl. spec. I, 28, t. 36 (1804) ...	W.A.	—	—	—	—	—	B.f.V, 352	
A. Cunninghami, Meissner in Lehm. pl. Preiss. I, 513 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 352	
A. pungens, Meissner in Lehm. pl. Preiss. I, 515 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 353	
A. venosus, Meissner in Hook. Kew Misc. IV, 183 (1852) ...	W.A.	—	—	—	—	—	B.f.V, 353	
A. Dobsoni, F. v. M., fragm. VI, 204 (1868) ...	W.A.	—	—	—	—	—	B.f.V, 353	M.fr. VI, 204.
A. linearis, Meissner in Hook. Kew Misc. IV, 183 (1852) ...	W.A.	—	—	—	—	—	B.f.V, 354	
A. sericeus, Labillardière, Nov. Holl. pl. specim. I, 29, t. 38 (1804) ...	W.A.	S.A.	—	—	—	—	B.f.V, 354	
A. Meissneri, Lehmann, pl. Preiss. I, 515 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 354	
A. filifolius, Benth, Fl. Austr. V, 355 (1873) ...	W.A.	—	—	—	—	—	B.f.V, 355	
A. terminalis, R. Brown in Transact. Linn. Soc. X, 152 (1809) ...	—	S.A.	—	V.	—	—	B.f.V, 355	
A. flavidiflorus, F. v. M., fragm. I, 157 (1859) ...	W.A.	—	—	—	—	—	B.f.V, 355	M.fr. I, 157.
A. apiculatus, R. Brown, prot. nov. 9 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 356	
SIMSIA, R. Brown in Transact. Linn. Soc. X, 152 (1809). (Stirlingia.)								
S. simplex, F. v. M.; Stirl., Lindl. Bot. Reg. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	B.f.V, 357	
S. abrotanoides, F. v. M.; Stirlingia, Meissn. in pl. Preiss. I, 517 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 357	
S. teretifolia, F. v. M.; Stirlingia, Meissn. in pl. Preiss. 515 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 357	
S. tenuifolia, R. Brown in Transact. Linn. Soc. X, 152 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 358	
S. latifolia, R. Brown, prot. nov. 9 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 358	
SYNAPHEA, R. Brown in Transact. Linn. Soc. X, 155 (1809).								
S. polymorpha, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 360	
S. dilatata, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 360	
S. favosa, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 360	
S. Preissii, Meissner in Lehm. pl. Preiss. I, 529 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 361	
S. acutiloba, Meissner in Lehm. pl. Preiss. I, 528 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 361	
S. petiolaris, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 361	
S. decorticans, Lindley, Bot. Regist. XXV, App. XXXII (1839) ...	W.A.	—	—	—	—	—	B.f.V, 362	
S. pinnata, Lindley, Bot. Regist. XXV, App. XXXII (1839) ...	W.A.	—	—	—	—	—	B.f.V, 362	
CONOSPERMUM, Smith in Transact. Linn. Soc. IV, 213 (1798).								
C. capitatum, R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 365	
C. petiolare, R. Brown, prot. nov. 11 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 365	
C. teretifolium, R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 365	M.fr. VI, 224.
C. flexuosum, R. Brown, prot. nov. 11 (1830) ...	W.A.	—	—	—	—	—	B.f.V, 366	M.fr. VI, 224.
C. acerosum, Lindley, Bot. Regist. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	B.f.V, 366	
C. amoenum, Meissner in Lehm. pl. Preiss. I, 522 (1845) ...	W.A.	—	—	—	—	—	B.f.V, 366	M.fr. I, 157; VI, 224.
C. nervosum, Meissner in Hook. Kew Misc. VII, 71 (1855) ...	W.A.	—	—	—	—	—	B.f.V, 367	M.fr. VI, 224.
C. diffusum, Benth, Fl. Austr. V, 367 (1870) ...	W.A.	—	—	—	—	—	B.f.V, 367	
C. glumaceum, Lindley, Bot. Regist. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	B.f.V, 367	
C. ephedroides, Kippist in Hook. Kew Misc. VII, 70 (1855) ...	W.A.	—	—	—	—	—	B.f.V, 368	
C. Toddii, F. v. M., fragm. X, 20 (1876) ...	W.A.	—	—	—	—	—	—	
C. polyccephalum, Meissner in Lehm. pl. Preiss. II, 249 (1847) ...	W.A.	—	—	—	—	—	B.f.V, 368	M.fr. X, 20, 90.
C. coeruleum, R. Brown in Transact. Linn. Soc. X, 154 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 368	M.fr. VI, 224.
C. debile, Kippist in Hook. Kew Misc. VII, 70 (1855) ...	W.A.	—	—	—	—	—	B.f.V, 369	
C. scaposum, Benth, Fl. Austr. V, 369 (1870) ...	W.A.	—	—	—	—	—	B.f.V, 369	
C. Huegelii, R. Brown in Endlicher, nov. stirp. dec. 58 (1839) ...	W.A.	—	—	—	—	—	B.f.V, 369	
C. densiflorum, Lindley, Bot. Regist. XXV, App. XXXII (1839) ...	W.A.	—	—	—	—	—	B.f.V, 369	
C. Brownii, Meissner in Lehm. pl. Preiss. II, 248 (1847) ...	W.A.	—	—	—	—	—	B.f.V, 370	
C. longifolium, Smith, Exot. Bot. II, 45, t. 82 (1805) ...	—	—	—	N.S.W.	—	—	B.f.V, 370	
C. tenuifolium, R. Brown in Transact. Linn. Soc. X, 154 (1809) ...	—	—	—	N.S.W.	—	—	B.f.V, 370	M.fr. VI, 223.
C. Mitchellii, Meissner in De Candolle, prodr. XIV, 320 (1856) ...	—	—	—	V.	—	—	B.f.V, 371	
C. sphacelatum, Hooker in Mitchell, Trop. Austr. 342 (1848) ...	—	—	—	—	Q.	N.A.	B.f.V, 371	M.fr. VI, 223.
C. patens, Schlechtendal, Linnaea XX, 587 (1847) ...	—	S.A.	—	V.	N.S.W.	—	B.f.V, 371	M.fr. VI, 223.
C. taxifolium, Smith in Rees's Cycl. IX (1808) ...	—	—	—	T.	V.	N.S.W.	Q.	B.f.V, 372
C. ericifolium, Smith in Rees's Cycl. IX (1808) ...	—	—	—	—	N.S.W.	—	B.f.V, 372	M.fr. VI, 224.
C. ellipticum, Smith in Rees's Cycl. IX (1808) ...	—	—	—	—	N.S.W.	—	B.f.V, 372	
C. distichum, R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	W.A.	—	—	—	—	—	B.f.V, 373	M.fr. VI, 224.
C. floribundum, Benth, Fl. Austr. V, 373 (1870) ...	W.A.	—	—	—	—	—	B.f.V, 373	

<i>C. incurvum</i> , Lindley, Bot. Regist. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	B.f. V, 373	
<i>C. brachyphyllum</i> , Lindley, Bot. Regist. XXV, App. XXXI (1839) ...	W.A.	—	—	—	—	—	B.f. V, 374	
<i>C. stoechadis</i> , Endlicher in Ann. Wien. Mus. II, 208 (1838) ...	W.A.	—	—	—	N.S.W.	—	B.f. V, 374	M.fr. XI, 20.
<i>C. triplinervium</i> , R. Brown, prot. nov. 11 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 375	
<i>C. bracteosum</i> , Meissner in Lehm. pl. Preiss. I, 518 (1845) ...	W.A.	—	—	—	—	—	B.f. V, 375	M.fr. VI, 224.
<i>C. crassinervium</i> , Meissner in Hook. Kew Misc. IV, 184 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 375	
FRANKLANDIA, R. Brown in Transact. Linn. Soc. X, 157 (1809).								
<i>F. fucifolia</i> , R. Brown in Transact. Linn. Soc. X, 157 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 376	M.fr. VI, 223; VII, 133.
<i>F. triaristata</i> , Benth. Fl. Austr. V, 377 (1870) ...	W.A.	—	—	—	—	—	B.f. V, 377	
SYMPHYONEMA, R. Brown in Transact. Linn. Soc. X, 157 (1809). (Symphonema.)								
<i>S. montanum</i> , R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 377	M.fr. VI, 223.
<i>S. paludosum</i> , R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 378	M.fr. VI, 223.
BELLENDENA, R. Brown in Transact. Linn. Soc. X, 156 (1809).								
<i>B. montana</i> , R. Brown in Transact. Linn. Soc. X, 166 (1809) ...	—	—	T.	—	—	—	B.f. V, 378	
AGASTACHYS, R. Brown in Transact. Linn. Soc. X, 158 (1809).								
<i>A. odorata</i> , R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	—	—	T.	—	—	—	B.f. V, 379	M.fr. X, 90.
CENARRHENES, Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804).								
<i>C. nitida</i> , Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804) ...	—	—	T.	—	—	—	B.f. V, 380	
PERSOONIA, Smith in Transact. Linn. Soc. IV, 215 (1798). (Linkia, 1797, Pentadactylon.)								
<i>P. hakeiformis</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 383	M.fr. VI, 223.
<i>P. teretifolia</i> , R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 383	M.fr. VI, 223.
<i>P. saccata</i> , R. Brown, prot. nov. 12 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 384	M.fr. VI, 223.
<i>P. Saundersiana</i> , Kippist in Hook. Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 384	
<i>P. diadema</i> , F. v. M., fragm. X, 46 (1876) ...	W.A.	—	—	—	—	—	—	M.fr. X, 46.
<i>P. comata</i> , Meissner in Hooker's Kew Misc. VII, 71 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 385	
<i>P. brachystylis</i> , F. v. M., fragm. VI, 221 (1868) ...	W.A.	—	—	—	—	—	B.f. V, 385	
<i>P. falcata</i> , R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	—	—	—	—	Q.	N.A.	B.f. V, 385	M.fr. VI, 222.
<i>P. trinervis</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 386	
<i>P. tortifolia</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 386	
<i>P. angustiflora</i> , Benth. Fl. Austr. V, 386 (1870) ...	W.A.	—	—	—	—	—	B.f. V, 386	
<i>P. rudis</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 387	
<i>P. microcarpa</i> , R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 387	M.fr. VI, 222.
<i>P. sulcata</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 387	
<i>P. acicularis</i> , F. v. M., fragm. VI, 220 (1868) ...	W.A.	—	—	—	—	—	B.f. V, 388	M.fr. VI, 220.
<i>P. scabrella</i> , Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 388	
<i>P. dillwynioides</i> , Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 388	M.fr. VI, 222.
<i>P. striata</i> , R. Brown, prot. nov. 13 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 388	M.fr. VI, 223.
<i>P. quinquenervis</i> , Hooker, icon. plant. t. 425 (1842) ...	W.A.	—	—	—	—	—	B.f. V, 389	
<i>P. rufiflora</i> , Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 389	
<i>P. scabra</i> , R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 389	
<i>P. graminea</i> , R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 390	M.fr. VI, 222.
<i>P. longifolia</i> , R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 390	M.fr. VI, 222.
<i>P. articulata</i> , R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 390	
<i>P. elliptica</i> , R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 391	M.fr. VI, 222.
<i>P. ferruginea</i> , Smith, Exot. Bot. II, 47, t. 83 (1805) ...	—	—	—	—	N.S.W.	—	B.f. V, 391	
<i>P. media</i> , R. Brown, prot. nov. 16 (1830) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 391	
<i>P. cornifolia</i> , Cunningham in R. Br., prot. nov. 16 (1830) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 392	
<i>P. marginata</i> , R. Brown, prot. nov. 16 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 392	
<i>P. sericea</i> , Cunningham in R. Br., prot. nov. 14 (1830) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 393	
<i>P. Mitchellii</i> , Meissner in Hook. Kew Misc. VII, 73 (1809) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 393	
<i>P. fastigiata</i> , R. Brown, prot. nov. 13 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 393	
<i>P. hirsuta</i> , Persoon, synops. plant. I, 118 (1805) ...	—	—	—	—	N.S.W.	—	B.f. V, 394	M.fr. VI, 222.
<i>P. Chamaeepithys</i> , Cunningham in Field, N.S. Wales, 329 (1825) ...	—	—	—	—	N.S.W.	—	B.f. V, 394	M.fr. VI, 223.
<i>P. arborea</i> , F. v. M., fragm. V, 37 (1865) ...	—	—	—	—	V.	—	B.f. V, 394	M.fr. V, 37; VI, 221.
<i>P. salicina</i> , Persoon, synops. plant. I, 118 (1805) ...	—	—	—	—	V.	N.S.W.	B.f. V, 395	
<i>P. prostrata</i> , R. Brown in Transact. Linn. Soc. X, 163 (1809) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 395	
<i>P. lanceolata</i> , Andrews, Bot. Reposit. t. 74 (1800) ...	—	—	—	—	N.S.W.	—	B.f. V, 395	M.fr. VI, 222.
<i>P. confertiflora</i> , Benth. Fl. Austr. V, 396 (1870) ...	—	—	—	—	V.	—	B.f. V, 396	
<i>P. lucida</i> , R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 396	
<i>P. linearis</i> , Andrews, Bot. Reposit. t. 77 (1800) ...	—	—	—	—	V.	N.S.W.	B.f. V, 397	M.fr. VI, 222.
<i>P. pinifolia</i> , R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 397	M.fr. VI, 222; X, 47.
<i>P. Caleyi</i> , R. Brown, prot. nov. 13 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 398	
<i>P. ledifolia</i> , Cunningham in De Cand. prodr. XIV, 339 (1856) ...	—	—	—	—	N.S.W.	—	B.f. V, 398	
<i>P. revoluta</i> , Sieber in Roem. & Schult. mant. 272 (1822) ...	—	—	—	—	N.S.W.	—	B.f. V, 398	M.fr. VI, 221.
<i>P. Gunnii</i> , J. Hooker in Lond. Journ. Bot. VI, 283 (1847) ...	—	—	T.	—	—	—	B.f. V, 399	
<i>P. mollis</i> , R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 399	M.fr. VI, 221. [47.
<i>P. rigida</i> , R. Brown, prot. nov. 14 (1830) ...	—	—	—	—	V.	N.S.W.	B.f. V, 399	M.fr. VI, 221; VIII, 149, X,
<i>P. curvifolia</i> , R. Brown, prot. nov. 13 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 400	
<i>P. oblongata</i> , Cunningham in R. Br. prot. nov. 14 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 400	
<i>P. Cunninghamii</i> , R. Brown, prot. nov. 15 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 400	
<i>P. myrtilloides</i> , Sieber in Roem. & Schult. mant. 272 (1822) ...	—	—	—	—	V.	N.S.W.	B.f. V, 401	M.fr. VI, 221.
<i>P. oxycooides</i> , Sieber in Roem. & Schult. mant. 270 (1822) ...	—	—	—	—	N.S.W.	—	B.f. V, 401	M.fr. VI, 221.
<i>P. nutans</i> , R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	—	—	—	—	N.S.W.	—	B.f. V, 401	M.fr. VI, 222.
<i>P. angulata</i> , R. Brown, prot. nov. 14 (1830) ...	—	—	—	—	N.S.W.	—	B.f. V, 402	
<i>P. virgata</i> , R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 402	
<i>P. Chamaepeuce</i> , Lhotsky in De Cand. prodr. XIV, 336 (1856) ...	—	—	—	—	V.	N.S.W.	B.f. V, 402	M.fr. VIII, 149; X, 47.
<i>P. juniperina</i> , Labillardière, Nov. Holl. pl. spec. I, 33, t. 45 (1804) ...	S.A.	T.	—	—	N.S.W.	—	B.f. V, 403	
<i>P. tenuifolia</i> , R. Brown, prot. nov. 12 (1830) ...	—	—	—	—	N.S.W.	Q.	B.f. V, 403	
<i>P. acerosa</i> , Sieber in Roem. & Schult. mant. 269 (1822) ...	—	—	—	—	N.S.W.	—	B.f. V, 403	M.fr. VI, 222.

MACADAMIA, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857). (Panopsis partly.)									
M. ternifolia, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.V, 406	M.fr. II, 91; VI, 191, 224,
HELICIA, Loureiro, Fl. Cochinch. I, 83 (1790).									
H. Youngiana, C. Moore & F. v. M., fragm. IV, 84 (1864)	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 406	M.fr. V, 186.
H. praealta, F. v. M., fragm. III, 37 (1862)	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 404	M.fr. IV, 174; VI, 224.
H. Australasica, F. v. M. in Hook. Kew Misc. IX, 27 (1857)	...	—	—	—	—	—	N.A.	B.fl.V, 405	M.fr. II, 91.
H. glabriflora, F. v. M., fragm. II, 91 (1861)	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 405	M.fr. V, 38, 186.
H. ferruginea, F. v. M., fragm. III, 37 (1862)	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 405	M.fr. IV, 174; V, 186; VI,
ROUPALA, Aublet, Hist. des pl. de la Guian. I, 83, t. 32 (1775). (Rhopala, Bleasdalea, Adenostephanus partly.)									
R. Bleasdalei, F. v. M. in proceed. Roy. Soc. N.S. Wales, 28 (1881)	—	—	—	—	—	Q.	—	B.fl.V, 417	M.fr. V, 90.
XYLOMELUM, Smith in Transact. Linn. Soc. IV, 214 (1798).									
X. pyriforme, Smith in J. Knight, prot. 105 (1809)	...	—	—	—	N.S.W.	—	—	B.fl.V, 408	M.fr. VI, 220.
X. occidentale, R. Brown, prot. nov. 31 (1830)	...	W.A.	—	—	—	—	—	B.fl.V, 408	—
X. salicinum, Cunningham in R. Br. prot. 31 (1830)	...	—	—	—	—	Q.	—	B.fl.V, 408	M.fr. IV, 107, 174; VI, 220.
X. angustifolium, Kippist in Hook. Kew Misc. IV, 209 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 409	M.fr. VI, 220.
LAMBERTIA, Smith in Transact. Linn. Soc. IV, 214 (1798).									
L. uniflora, R. Brown in Transact. Linn. Soc. X, 188 (1809)	...	W.A.	—	—	—	—	—	B.fl.V, 413	M.fr. VI, 248.
L. rariflora, Meissner in Lehm. pl. Preiss. II, 263 (1847)	...	W.A.	—	—	—	—	—	B.fl.V, 414	—
L. inermis, R. Brown in Transact. Linn. Soc. X, 188 (1809)	...	W.A.	—	—	—	—	—	B.fl.V, 414	M.fr. VI, 248.
L. ericifolia, R. Brown, prot. nov. 30 (1830)	...	W.A.	—	—	—	—	—	B.fl.V, 414	M.fr. VI, 248.
L. multiflora, Lindley, Bot. Regist. XXV, App. XXXII (1839)	...	W.A.	—	—	—	—	—	B.fl.V, 415	—
L. formosa, Smith in Transact. Linn. Soc. IV, 214, t. 20 (1798)	...	W.A.	—	—	N.S.W.	—	—	B.fl.V, 415	M.fr. VII, 133.
L. echinata, R. Brown in Transact. Linn. Soc. X, 189 (1809)	...	W.A.	—	—	—	—	—	B.fl.V, 416	M.fr. VI, 248.
L. ilicifolia, Hooker, icon. plant. t. 553 (1843)	...	W.A.	—	—	—	—	—	B.fl.V, 416	—
ORTITES, R. Brown in Transact. Linn. Soc. X, 189 (1809). (Oritina.)									
O. excelsa, R. Brown, prot. nov. 32 (1832)	...	—	—	—	N.S.W.	—	—	B.fl.V, 411	M.fr. V, 153; VI, 223.
O. diversifolia, R. Brown in Transact. Linn. Soc. X, 190 (1809)	...	—	—	T.	—	—	—	B.fl.V, 411	M.fr. V, 153.
O. Milligani, Meissner in Hook. Kew Misc. IV, 209 (1852)	...	—	—	T.	—	—	—	B.fl.V, 411	—
O. lancifolia, F. v. M. in Transact. phil. Soc. Vict. I, 108 (1855)	...	—	—	—	V.	—	—	B.fl.V, 412	M.fr. V, 153; VI, 223; VII,
O. revoluta, R. Brown in Transact. Linn. Soc. X, 19 (1809)	...	—	—	T.	—	—	—	B.fl.V, 412	[133.]
O. acicularis, R. Brown, prot. nov. 32 (1830)	...	—	—	T.	—	—	—	B.fl.V, 412	—
STRANGEA, Meissner in Hooker's Kew Misc. VII, 66 (1855). (Molloya.)									
S. linearis, Meissner in Hook. Kew Misc. VII, 66 (1855)	...	—	—	—	N.S.W.	Q.	—	B.fl.V, 453	M.fr. VI, 223; VII, 132.
S. cynanchocarpa, F. v. M., fragm. VII, 132 (1871)	...	W.A.	—	—	—	—	—	—	M.fr. VII, 132.
GREVILLEA, R. Brown in Transact. Linn. Soc. X, 168 (1809). (Anadenia, Lyssanthus, Stylurus, Manglesia.)									
G. Pinaster, Meissner in Hook. Kew Misc. VII, 76 (1855)	...	W.A.	—	—	—	—	—	B.fl.V, 427	M.fr. VI, 212.
G. obtusifolia, Meissner in Hook. Kew Misc. IV, 187 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 427	—
G. sparsiflora, F. v. M., fragm. VI, 206 (1868)	...	W.A.	—	—	—	—	—	B.fl.V, 428	M.fr. VI, 206.
G. macrostylis, F. v. M., fragm. I, 137 (1859)	...	W.A.	—	—	—	—	—	B.fl.V, 428	M.fr. VI, 211.
G. tripartita, Meissner in Hook. Kew Misc. IV, 186 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 428	M.fr. VI, 213.
G. platypoda, F. v. M., fragm. VI, 205 (1868)	...	W.A.	—	—	—	—	—	B.fl.V, 428	M.fr. VI, 205.
G. patentiloba, F. v. M., fragm. I, 137 (1859)	...	W.A.	—	—	—	—	—	B.fl.V, 429	M.fr. I, 137.
G. pectinata, R. Brown, prot. nov. 23 (1830)	...	W.A.	—	—	—	—	—	B.fl.V, 429	M.fr. VI, 212.
G. pluri-juga, F. v. M., fragm. IV, 84 (1864)	...	W.A.	—	—	—	—	—	B.fl.V, 430	M.fr. VI, 212.
G. nudiflora, Meissner in Hook. Kew Misc. IV, 186 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 430	M.fr. I, 35; VI, 212.
G. stenomera, F. v. M., fragm. IV, 85 (1864)	...	W.A.	—	—	—	—	—	B.fl.V, 430	M.fr. IV, 85.
G. Thelemanniana, Endlicher, nov. stirp. dec. 6 (1839)	...	W.A.	—	—	—	—	—	B.fl.V, 431	M.fr. VI, 212.
G. concinna, R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	W.A.	—	—	—	—	—	B.fl.V, 431	M.fr. VI, 212.
G. Hookeriana, Meissner in Lehm. pl. Preiss. I, 546 (1845)	...	W.A.	—	—	—	—	—	B.fl.V, 432	M.fr. VI, 212.
G. Baxteri, R. Brown, prot. nov. 22 (1830)	...	W.A.	—	—	—	—	—	B.fl.V, 432	—
G. eriobotrya, F. v. M., fragm. X, 44 (1876)	...	W.A.	—	—	—	—	—	—	B.fl. X, 44.
G. pterosperma, F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	...	W.A.	S.A.	—	V.	N.S.W.	—	B.fl.V, 432	M.fr. VI, 210.
G. stenobotrya, F. v. M., fragm. IX, 3 (1875)	...	—	S.A.	—	—	—	N.A.	—	M.fr. IX, 3.
G. eriostachya, Lindley, Bot. Regist. XXV, App. XXXVI (1839)	...	—	—	—	—	—	N.A.	B.fl.V, 433	M.fr. VI, 208 & 213.
G. thyrsoidea, Meissner in Hooker's Kew Misc. VII, 77 (1855)	...	W.A.	—	—	—	—	—	B.fl.V, 433	—
G. Chrysodendron, R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	—	—	—	—	Q.	N.A.	B.fl.V, 434	M.fr. VI, 210.
G. Banksii, R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	—	—	—	—	Q.	—	B.fl.V, 434	M.fr. VI, 210; VIII, 150; X,
G. Caleyi, R. Brown, prot. nov. 22 (1830)	...	—	—	—	N.S.W.	—	—	B.fl.V, 435	—
G. asplenifolia, R. Brown in J. Knight, prot. 120 (1869)	...	—	—	—	N.S.W.	—	—	B.fl.V, 435	M.fr. VI, 212.
G. cirsiifolia, Meissner in Lehm. pl. Preiss. II, 253 (1847)	...	W.A.	—	—	—	—	—	B.fl.V, 436	—
G. laurifolia, Sieber in Roem. & Schult. mant. 279 (1822)	...	—	—	—	N.S.W.	—	—	B.fl.V, 436	[X, 46.]
G. Barklyana, F. v. M., gen. report 14 & 18 (1861)	...	—	—	—	V.	—	—	B.fl.V, 436	M.fr. VII, 133; VIII, 150;
G. repens, F. v. M. in Schlecht. Linnaea XXVI, 355 (1853)	...	—	—	—	V.	—	—	B.fl.V, 437	—
G. Aquifolium, Lindley in Mitchell, Three Exped. II, 178 (1838)	...	—	—	—	V.	—	—	B.fl.V, 437	—
G. ilicifolia, R. Brown, prot. nov. 21 (1830)	...	—	S.A.	—	V.	—	—	B.fl.V, 437	M.fr. VI, 212.
G. Gaudichaudii, R. Brown in Freye. Voy. Bot. 443, t. 46 (1826)	...	—	—	—	N.S.W.	—	—	B.fl.V, 438	—
G. acanthifolia, Cunningham in Field, N.S. Wales, 328 (1825)	...	—	—	—	N.S.W.	—	—	B.fl.V, 438	M.fr. VI, 212; VIII, 150.
G. bipinnatifida, R. Brown, prot. nov. 23 (1830)	...	W.A.	—	—	—	—	—	B.fl.V, 439	M.fr. VI, 212.
G. armigera, Meissner in Hook. Kew Misc. IV, 186 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 439	—
G. asparagoides, Meissner in Hook. Kew Misc. IV, 186 (1852)	...	W.A.	—	—	—	—	—	B.fl.V, 439	M.fr. IX, 123.
G. Treueriana, F. v. M., fragm. IX, 123 (1875)	...	W.A.	S.A.	—	—	—	—	—	M.fr. IX, 123.
G. floribunda, R. Brown, prot. nov. 19 (1830)	...	—	—	—	V.	N.S.W.	Q.	B.fl.V, 440	—
G. cinerea, R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	—	—	—	N.S.W.	—	—	B.fl.V, 440	—
G. alpina, Lindley in Mitchell, Three Exped. II, 179 (1838)	...	—	—	—	V.	—	—	B.fl.V, 441	—
G. montana, R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	—	—	—	N.S.W.	—	—	B.fl.V, 441	—
G. obtusiflora, R. Brown, prot. nov. 19 (1830)	...	—	—	—	N.S.W.	—	—	B.fl.V, 442	—
G. arenaria, R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	—	—	—	N.S.W.	—	—	B.fl.V, 442	—
G. mucronulata, R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	—	—	—	N.S.W.	—	—	B.fl.V, 443	—

G. Baueri, R. Brown in Transact. Linn. Soc. X, 173 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 443	
G. lanigera, Cunningham in R. Br., prot. nov. 20 (1830) ...	—	—	—	V. N.S.W.	—	B.f. V, 444	
G. rosmarinifolia, Cunningham in Field, N.S. Wales, 328 (1825) ...	—	—	—	V. N.S.W.	—	B.f. V, 445	
G. Goodii, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	—	Q. N.A.	B.f. V, 446	M.fr. VI, 211.
G. venusta, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	Q. —	B.f. V, 446	
G. longistyla, Hooker in Mitchell, Trop. Austr. 343 (1848) ...	—	—	—	—	Q. N.A.	B.f. V, 446	M.fr. VI, 211.
G. juncifolia, Hooker in Mitchell, Trop. Austr. 341 (1848) ...	—	S.A.	—	N.S.W.	Q. —	B.f. V, 447	M.fr. VI, 209; VIII, 150.
G. Wilsoni, Cunningham in Wilson's Voy. 273 (1835) ...	W.A.	—	—	—	—	B.f. V, 447	M.fr. VI, 212; X, 45.
G. erectiloba, F. v. M., fragm. X, 44 (1876) ...	W.A.	—	—	—	—	—	M.fr. X, 44.
G. lavandulacea, Schlechtendal, Linnaea XX, 586 (1847) ...	—	S.A.	—	V. N.S.W.	—	B.f. V, 448	
G. insignis, Kippist in De Cand., prodr. XIV, 379 (1856) ...	W.A.	—	—	—	—	B.f. V, 448	M.fr. VIII, 150; X, 45.
G. Brownii, Meissner in Lehm. pl. Preiss. I, 537 (1845) ...	W.A.	—	—	—	—	B.f. V, 449	M.fr. VI, 213.
G. fasciculata, R. Brown, prot. nov. 20 (1830) ...	W.A.	—	—	—	—	B.f. V, 449	
G. aspera, R. Brown in Transact. Linn. Soc. X, 172 (1809) ...	W.A.	S.A.	—	—	—	B.f. V, 450	M.fr. VI, 211.
G. brachystylis, Meissner in Lehm. pl. Preiss. I, 538 (1845) ...	W.A.	—	—	—	—	B.f. V, 450	
G. saccata, Benth., Fl. Austr. V, 450 (1870) ...	W.A.	—	—	—	—	B.f. V, 450	
G. Drummondii, Meissner in Lehm. pl. Preiss. I, 536 (1845) ...	W.A.	—	—	—	—	B.f. V, 451	
G. disjuncta, F. v. M., fragm. VI, 206 (1868) ...	W.A.	—	—	—	—	B.f. V, 451	M.fr. VI, 206.
G. haplantha, F. v. M. in Benth. Fl. Austr. V, 451 (1870) ...	W.A.	—	—	—	—	B.f. V, 452	
G. pinifolia, Meissner in Hook. Kew Misc. 186 (1852) ...	W.A.	—	—	—	—	B.f. V, 452	
G. acuaria, F. v. M. in Benth. Fl. Austr. V, 452 (1870) ...	W.A.	—	—	—	—	B.f. V, 452	
G. singuliflora, F. v. M., fragm. VI, 92 (1868) ...	—	—	—	—	Q. —	B.f. V, 452	M.fr. VI, 92.
G. pauciflora, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	W.A.	S.A.	—	—	—	B.f. V, 453	M.fr. I, 135; VI, 206, 208.
G. quercifolia, R. Brown, prot. nov. 23 (1830) ...	W.A.	—	—	—	—	B.f. V, 454	M.fr. VI, 213.
G. angulata, R. Brown, prot. nov. 24 (1830) ...	—	S.A.	—	—	—	B.f. V, 455	
G. Wickhami, Meissner in Hook. Kew Misc. 187 (1852) ...	—	S.A.	—	—	—	B.f. V, 455	
G. agrifolia, Cunningham in R. Br., prot. nov. 24 (1830) ...	—	—	—	—	—	B.f. V, 455	
G. Cunninghamii, R. Brown, prot. nov. 23 (1830) ...	—	—	—	—	—	B.f. V, 456	
G. pungens, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	—	B.f. V, 456	
G. Huegelii, Meissner in Lehm. pl. Preiss. I, 543 (1845) ...	W.A.	S.A.	—	V. N.S.W.	—	B.f. V, 457	M.fr. VI, 212; X, 46.
G. dimidiata, F. v. M., fragm. III, 146 (1863) ...	—	—	—	—	—	B.f. V, 457	M.fr. III, 146.
G. heliosperma, R. Brown in Transact. Linn. Soc. X, 176 (1809) ...	—	—	—	—	—	B.f. V, 457	M.fr. VI, 210.
G. refracta, R. Brown in Transact. Linn. Soc. X, 176 (1809) ...	—	—	—	—	—	B.f. V, 458	
G. Dryandri, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	Q. N.A.	B.f. V, 458	M.fr. VI, 209.
G. polystachya, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	—	—	—	Q. N.A.	B.f. V, 459	M.fr. VI, 210.
G. robusta, Cunningham in R. Br., prot. nov. 24 (1830) ...	—	—	—	N.S.W.	Q. —	B.f. V, 459	M.fr. VI, 210; VII, 133; XI, 20.
G. annulifera, F. v. M., fragm. IV, 85 (1864) ...	W.A.	—	—	—	—	B.f. V, 460	M.fr. IV, 85; VI, 212; VII, 132;
G. leucopteris, Meissner in Hook. Kew Misc. VII, 76 (1855) ...	W.A.	—	—	—	—	B.f. V, 460	M.fr. III, 145; IV, 85, 176;
G. Leucadendron, Cunningham in R. Br., prot. nov. 25 (1830) ...	—	—	—	—	—	B.f. V, 461	M.fr. I, 136-7. [XI, 20]
G. pyramidalis, Cunningham in R. Br., prot. nov. 25 (1830) ...	—	—	—	—	—	B.f. V, 462	
G. striata, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	S.A.	—	N.S.W.	Q. N.A.	B.f. V, 462	
G. mimosoides, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	—	—	—	—	B.f. V, 462	
G. Hilliana, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857) ...	—	—	—	N.S.W.	Q. —	B.f. V, 463	
G. gibbosa, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	—	—	—	Q. N.A.	B.f. V, 463	M.fr. VI, 210.
G. buxifolia, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 464	
G. phyllicoides, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 464	M.fr. VI, 213.
G. sphacelata, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 464	
G. occidentalis, R. Brown in Transact. Linn. Soc. X, 173 (1809) ...	W.A.	—	—	—	—	B.f. V, 465	
G. acerosa, F. v. M., fragm. I, 136 (1859) ...	W.A.	—	—	—	—	B.f. V, 465	M.fr. I, 136.
G. umbellulata, Meissner in Lehm. pl. Preiss. II, 252 (1847) ...	W.A.	—	—	—	—	B.f. V, 465	
G. oxystigma, Meissner in Lehm. pl. Preiss. I, 540 (1845) ...	W.A.	—	—	—	—	B.f. V, 466	
G. Candolleana, Meissner in Lehm. pl. Preiss. I, 541 (1845) ...	W.A.	—	—	—	—	B.f. V, 466	M.fr. VI, 213.
G. scabra, Meissner in Lehm. pl. Preiss. I, 541 (1845) ...	W.A.	—	—	—	—	B.f. V, 466	
G. Miqueliana, F. v. M. in Transact. Vict. Inst. 132 (1855) ...	—	—	—	V. —	—	B.f. V, 467	
G. Victoriae, F. v. M. in Transact. phil. Soc. Vict. I, 107 (1865) ...	—	—	—	V. N.S.W.	—	B.f. V, 468	M.fr. VIII, 142, 150.
G. punicea, R. Brown in Transact. Linn. Soc. X, 169 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 468	M.fr. XI, 123.
G. oleoides, Sieber in Roem. & Schult. mant. 277 (1822) ...	—	—	—	V. N.S.W.	—	B.f. V, 468	
G. trinervis, R. Brown, prot. nov. 18 (1830) ...	—	—	—	N.S.W.	—	B.f. V, 469	M.fr. VI, 211.
G. juniperina, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 469	
G. sericea, R. Brown in Transact. Linn. Soc. X, 170 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 470	
G. capitellata, Meissner in Hook. Kew Misc. IV, 187 (1852) ...	—	—	—	N.S.W.	—	B.f. V, 470	
G. leiophylla, F. v. M. in Benth. Fl. Austr. V, 471 (1870) ...	—	—	—	N.S.W.	Q. —	B.f. V, 471	
G. linearis, R. Brown in Transact. Linn. Soc. X, 170 (1809) ...	—	—	—	N.S.W.	—	B.f. V, 471	
G. confertifolia, F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854) ...	—	—	—	V. —	—	B.f. V, 471	M.fr. VI, 211.
G. parviflora, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	—	S.A.	—	V. N.S.W.	—	B.f. V, 472	
G. australis, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	—	—	T. V.	N.S.W.	—	B.f. V, 472	M.fr. VII, 133.
G. commutata, F. v. M., fragm. VI, 207 (1868) ...	W.A.	—	—	—	—	B.f. V, 473	M.fr. VII, 207.
G. pinnatisecta, F. v. M. in Benth. Fl. Austr. V, 473 (1870) ...	W.A.	—	—	—	—	B.f. V, 473	
G. argyrophylla, Meissner in Hook. Kew Misc. VII, 75 (1855) ...	W.A.	—	—	—	—	B.f. V, 474	M.fr. VI, 411.
G. brachystachys, Meissner in Lehm. pl. Preiss. II, 254 (1847) ...	W.A.	—	—	—	—	B.f. V, 474	M.fr. VI, 211.
G. Endlicheriana, Meissner in Lehm. pl. Preiss. I, 546 (1845) ...	W.A.	—	—	—	—	B.f. V, 474	M.fr. VI, 211.
G. manglesoides, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	B.f. V, 475	
G. diversifolia, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	B.f. V, 475	
G. filifolia, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	B.f. V, 475	
G. hakeoides, Meissner in Lehm. pl. Preiss. II, 252 (1847) ...	W.A.	—	—	—	—	B.f. V, 476	
G. teretifolia, Meissner in Lehm. pl. Preiss. II, 255 (1847) ...	W.A.	—	—	—	—	B.f. V, 476	
G. eryngioides, Benth., Fl. Austr. V, 476 (1870) ...	W.A.	—	—	—	—	B.f. V, 476	M.fr. X, 46.
G. bracteosa, Meissner in Lehm. pl. Preiss. II, 254 (1847) ...	W.A.	—	—	—	—	B.f. V, 477	
G. crithmifolia, R. Brown, prot. nov. 23 (1830) ...	W.A.	—	—	—	—	B.f. V, 477	
G. trachytheca, F. v. M., fragm. VI, 207 (1868) ...	W.A.	—	—	—	—	B.f. V, 477	M.fr. VI, 207.
G. triternata, R. Brown, prot. nov. 21 (1830) ...	—	—	—	V. N.S.W.	—	B.f. V, 478	

G. ramosissima, Meissner in De Candolle, prodr. XIV, 388	...	—	—	—	N.S.W.	—	B.f. V, 478	M.fr. VIII, 150; X, 46.
G. monticola, Meissner in Lehm. pl. Preiss. II, 259 (1847)	...	W.A.	—	—	—	—	B.f. V, 478	M.fr. VI, 209.
G. Muelleri, Benth. Fl. Austr. V, 479 (1870)	...	W.A.	—	—	—	—	B.f. V, 479	
G. trifida, Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	W.A.	—	—	—	—	B.f. V, 479	M.fr. VI, 209.
G. Synapheae, R. Brown, prot. nov. 23 (1830)	...	W.A.	—	—	—	—	B.f. V, 480	
G. flexuosa, Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	W.A.	—	—	—	—	B.f. V, 480	
G. leptobotrya, Meissner in Lehm. pl. Preiss. II, 256 (1847)	...	W.A.	—	—	—	—	B.f. V, 480	M.fr. VI, 209.
G. brevicuspis, Meissner in Lehm. pl. Preiss. II, 256 (1847)	...	W.A.	—	—	—	—	B.f. V, 481	
G. intricata, Meissner in Hook. Kew Misc. VII, 74 (1855)	...	W.A.	—	—	—	—	B.f. V, 481	
G. didymobotrya, Meissner in Hook. Kew Misc. IV, 186 (1852)	...	W.A.	—	—	—	—	B.f. V, 481	M.fr. I, 136; VI, 208.
G. polybotrya, Meissner in Hook. Kew Misc. IV, 185 (1852)	...	W.A.	—	—	—	N.A.	B.f. V, 482	M.fr. IV, 129; VI, 208.
G. nematophylla, F. v. M., fragm. I, 136 (1859)	...	—	S.A.	—	N.S.W.	—	B.f. V, 482	M.fr. I, 136; VI, 208.
G. anethifolia, R. Brown, prot. nov. 21 (1830)	...	—	—	—	N.S.W.	—	B.f. V, 482	M.fr. VI, 209.
G. paradoxa, F. v. M., fragm. VI, 246 (1868)	...	W.A.	—	—	—	—	B.f. V, 483	M.fr. VI, 246.
G. petrophiloides, Meissner in Lehm. pl. Preiss. II, 257 (1847)	...	W.A.	—	—	—	—	B.f. V, 483	M.fr. VI, 209; X, 46.
G. tenuiflora, Meissner in Lehm. pl. Preiss. I, 554 (1845)	...	W.A.	—	—	—	—	B.f. V, 483	
G. pulchella, Meissner in Lehm. pl. Preiss. I, 553 (1845)	...	W.A.	—	—	—	—	B.f. V, 484	M.fr. VI, 209.
G. rudis, Meissner in Hook. Kew Misc. VII, 73 (1855)	...	W.A.	—	—	—	—	B.f. V, 484	
G. apiculoba, F. v. M., fragm. X, 45 (1876)	...	W.A.	—	—	—	—	—	M.fr. X, 45.
G. Shuttleworthiana, Meissner in Lehm. pl. Preiss. II, 258 (1847)	...	W.A.	—	—	—	—	B.f. V, 484	
G. integrifolia, Meissner in De Candolle, prodr. XIV, 385 (1856)	...	W.A.	—	—	—	—	B.f. V, 485	
G. stenocarpa, F. v. M. in Benth. Fl. Austr. V, 485 (1870)	...	W.A.	—	—	—	—	B.f. V, 485	
G. acrobotrya, Meissner in Hooker's Kew Misc. VII, 74 (1855)	...	W.A.	—	—	—	—	B.f. V, 486	M.fr. VI, 213.
G. glabrata, Meissner in Lehm. pl. Preiss. I, 549 (1845)	...	W.A.	—	—	—	—	B.f. V, 486	
G. ornithopoda, Meissner in Lehm. pl. Preiss. II, 256 (1847)	...	W.A.	—	—	—	—	B.f. V, 486	
G. paniculata, Meissner in Lehm. pl. Preiss. I, 550 (1845)	...	W.A.	—	—	—	—	B.f. V, 487	
G. biternata, Meissner in Lehm. pl. Preiss. I, 549 (1845)	...	W.A.	—	—	—	—	B.f. V, 487	M.fr. VI, 213.
G. triloba, Meissner in Hooker's Kew Misc. VII, 74 (1855)	...	W.A.	—	—	—	—	B.f. V, 487	M.fr. VI, 213.
G. amplexans, F. v. M. in Benth. Fl. Austr. V, 488 (1870)	...	W.A.	—	—	—	—	B.f. V, 488	
G. vestita, Meissner in Lehm. pl. Preiss. I, 548 (1845)	...	W.A.	—	—	—	—	B.f. V, 488	M.fr. VI, 213.
G. tridentifera, Meissner in Lehm. pl. Preiss. I, 547 (1845)	...	W.A.	—	—	—	—	B.f. V, 488	
G. erinacea, Meissner in Hooker's Kew Misc. VII, 74 (1855)	...	W.A.	—	—	—	—	B.f. V, 489	M.fr. X, 45.
HAKEA, Schrader, sert. Hannov. I, fasc. 3, 27, t. 17 (1797). (Conchium.)								
H. chordophylla, F. v. M. in Hooker's Kew Misc. IX, 23 (1857)	...	—	—	—	—	N.A.	B.f. V, 495	M.fr. VI, 190.
H. Cunninghamii, R. Brown, prot. nov. 26 (1830)	...	—	—	—	—	N.A.	B.f. V, 495	M.fr. VI, 190.
H. lorea, R. Brown, prot. nov. 25 (1830)	...	W.A.	S.A.	—	—	Q.	B.f. V, 495	M.fr. VI, 189; VII, 133.
H. Fraseri, R. Brown, prot. nov. 26 (1830)	...	—	—	—	N.S.W.	—	B.f. V, 496	
H. macrocarpa, Cunningham in R. Brown, prot. nov. 30 (1830)	...	—	—	—	—	N.A.	B.f. V, 496	M.fr. VI, 190, 210.
H. arborescens, R. Brown in Transact. Linn. Soc. X, 187 (1809)	...	—	—	—	—	Q.	B.f. V, 497	M.fr. VI, 190.
H. stenophylla, Cunningham in De Cand. prodr. XIV, 417 (1856)	...	—	—	—	—	N.A.	B.f. V, 497	
H. trineura, F. v. M., fragm. III, 146 (1863)	...	—	—	—	—	Q.	B.f. V, 497	M.fr. VI, 216.
H. cyclocarpa, Lindley, Bot. Regist. XXV, App. XXXVI (1839)	...	W.A.	—	—	—	—	B.f. V, 498	
H. crassifolia, Meissner in Lehm. pl. Preiss. I, 570 (1845)	...	W.A.	—	—	—	—	B.f. V, 498	M.fr. VI, 216.
H. pandanocarpa, R. Brown, prot. nov. 29 (1830)	...	W.A.	—	—	—	—	B.f. V, 499	M.fr. VI, 216.
H. Roei, Benth. Fl. Austr. V, 499 (1870)	...	W.A.	—	—	—	—	B.f. V, 499	
H. adnata, R. Brown, prot. nov. 26 (1830)	...	W.A.	—	—	—	—	B.f. V, 499	
H. obliqua, R. Brown in Transact. Linn. Soc. X, 180 (1809)	...	W.A.	—	—	—	—	B.f. V, 500	M.fr. VI, 218.
H. Hookeriana, Meissner in Hook. Kew Misc. IV, 208 (1852)	...	W.A.	—	—	—	—	B.f. V, 500	M.fr. VI, 216.
H. incrassata, R. Brown, prot. nov. 29 (1839)	...	W.A.	—	—	—	—	B.f. V, 500	
H. flabellifolia, Meissner in Hooker's Kew Misc. VII, 116 (1855)	...	W.A.	—	—	—	—	B.f. V, 501	
H. Brownii, Meissner in Lehm. pl. Preiss. I, 569 (1845)	...	W.A.	—	—	—	—	B.f. V, 501	
H. Baxteri, R. Brown, prot. nov. 28 (1830)	...	W.A.	—	—	—	—	B.f. V, 501	
H. ceratophylla, R. Brown in Transact. Linn. Soc. X, 184 (1809)	...	W.A.	—	—	—	—	B.f. V, 501	M.fr. IV, 49; VI, 217.
H. lasiantha, R. Brown, prot. nov. 29 (1830)	...	W.A.	—	—	—	—	B.f. V, 502	M.fr. VI, 215.
H. eriantha, R. Brown, prot. nov. 29 (1830)	...	—	—	—	V. N.S.W.	—	B.f. V, 502	M.fr. VI, 217.
H. megalosperma, Meissner in Hook. Kew Misc. VII, 117 (1855)	...	W.A.	—	—	—	—	B.f. V, 503	
H. clavata, Labillardière, Nov. Holl. pl. spec. I, 31, t. 41 (1804)	...	W.A.	—	—	—	—	B.f. V, 503	M.fr. VI, 216.
H. orthorrhyncha, F. v. M., fragm. V, 214 (1868)	...	W.A.	—	—	—	—	B.f. V, 503	M.fr. VI, 214.
H. Candolleana, Meissner in Lehm. pl. Preiss. II, 262 (1847)	...	W.A.	—	—	—	—	B.f. V, 504	M.fr. VI, 215.
H. trifurcata, R. Brown in Transact. Linn. Soc. X, 183 (1809)	...	W.A.	—	—	—	—	B.f. V, 504	M.fr. VI, 220.
H. erinacea, Meissner in Lehm. pl. Preiss. I, 559 (1845)	...	W.A.	—	—	—	—	B.f. V, 505	M.fr. VI, 220.
H. platysperma, Hooker, icones pl. t. 433 (1842)	...	W.A.	—	—	—	—	B.f. V, 505	
H. brachyptera, Meissner in Hook. Kew Misc. IV, 208 (1852)	...	W.A.	—	—	—	—	B.f. V, 505	M.fr. VI, 217.
H. Kippistiana, Meissner in Hooker's Kew Misc. VII, 115 (1855)	...	W.A.	—	—	—	—	B.f. V, 506	
H. Preissii, Meissner in Lehm. pl. Preiss. I, 557 (1845)	...	W.A.	—	—	—	—	B.f. V, 506	M.fr. VI, 219.
H. pugioniformis, Cavanilles, Annal. hist. nat. I, 213, t. 11 (1800)	...	—	—	T. V.	N.S.W.	—	B.f. V, 506	M.fr. VI, 217.
H. Pampliniana, Kippist in Hook. Kew Misc. VII, 115 (1855)	...	—	S.A.	—	V. N.S.W.	Q.	B.f. V, 507	
H. vittata, R. Brown in Transact. Linn. Soc. X, 182 (1809)	...	—	S.A.	—	—	—	B.f. V, 507	
H. rostrata, F. v. M. in Schlecht. Linnaea XXVI, 259 (1853)	...	—	S.A.	—	V.	—	B.f. V, 508	M.fr. VI, 218; VII, 133.
H. rugosa, R. Brown in Transact. Linn. Soc. X, 179 (1809)	...	—	S.A.	—	V.	—	B.f. V, 508	M.fr. VI, 218.
H. Epiglottis, Labillardière, Nov. Holl. pl. spec. I, 30, t. 40 (1804)	...	—	—	T.	—	—	B.f. V, 508	
H. amplexicaulis, R. Brown in Transact. Linn. Soc. X, 184 (1809)	...	W.A.	—	—	—	—	B.f. V, 509	M.fr. VI, 217; VII, 133.
H. glabella, R. Brown, prot. nov. 28 (1830)	...	W.A.	—	—	—	—	B.f. V, 509	
H. auriculata, Meissner in Hook. Kew Misc. VII, 116 (1855)	...	W.A.	—	—	—	—	B.f. V, 510	M.fr. VI, 217.
H. cristata, R. Brown, prot. nov. 28 (1830)	...	W.A.	—	—	—	—	B.f. V, 510	
H. linearis, R. Brown in Transact. Linn. Soc. X, 183 (1809)	...	W.A.	—	—	—	—	B.f. V, 511	
H. stenocarpoides, F. v. M. in Benth. Fl. V, 511 (1870)	...	W.A.	—	—	—	—	B.f. V, 511	
H. ruscifolia, Labillardière, Nov. Holl. pl. spec. I, 30, t. 39 (1804)	...	W.A.	—	—	—	—	B.f. V, 511	
H. saligna, R. Brown in J. Knight, prot. 108 (1809)	...	—	—	—	N.S.W.	Q.	B.f. V, 512	M.fr. VI, 217.
H. verrucosa, F. v. M., fragm. V, 25 (1865)	...	W.A.	—	—	—	—	B.f. V, 512	M.fr. V, 25; VI, 218.

H. rhombales, F. v. M., fragm. X, 90 (1876) ...	—	—	—	—	—	N.A.	—	M.fr. X, 90.
H. purpurea, Hooker in Mitch. Trop. Austr. 348 (1848)...	—	—	—	—	N.S.W.	Q.	B.fl. V, 513	
H. gibbosa, Cavanilles, Anal. hist. nat. I, 215 (1800) ...	—	—	—	—	N.S.W.	—	B.fl. V, 513	M.fr. VI, 218.
H. propinqua, Cunningham in Field N.S. Wales, 327 (1825) ...	—	—	—	—	N.S.W.	—	B.fl. V, 513	M.fr. V, 26.
H. nodosa, R. Brown in Transact. Linn. Soc. X, 179 (1809) ...	—	—	—	T. V.	—	—	B.fl. V, 514	M.fr. V, 26; VI, 217.
H. acicularis, R. Brown in Transact. Linn. Soc. X, 181 (1809)...	—	—	—	T. V.	N.S.W.	—	B.fl. V, 514	M.fr. VI, 218.
H. leucoptera, R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	—	S.A.	—	V.	N.S.W.	Q.	B.fl. V, 515	M.fr. VI, 219; VII, 133.
H. cycloptera, R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	—	S.A.	—	—	—	—	B.fl. V, 515	M.fr. VI, 217.
H. microcarpa, R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	—	—	—	T. V.	N.S.W.	—	B.fl. V, 516	M.fr. VI, 217.
H. recurva, Meissner in Hook. Kew Misc. IV, 207 (1852) ...	W.A.	—	—	—	—	—	B.fl. V, 516	M.fr. VI, 218.
H. circumalata, Meissner in Hook. Kew Misc. VII, 114 (1855)...	W.A.	—	—	—	—	—	B.fl. V, 516	M.fr. VI, 219.
H. commutata, F. v. M., fragm. V, 26 (1865) ...	W.A.	—	—	—	—	—	B.fl. V, 517	M.fr. VI, 219.
H. strumosa, Meissner in Hook. Kew Misc. IV, 208 (1852) ...	W.A.	—	—	—	—	—	B.fl. V, 517	M.fr. VI, 219.
H. multilineata, Meissner in Lehm. pl. Preiss. II, 261 (1847) ...	W.A.	S.A.	—	—	—	—	B.fl. V, 518	M.fr. VI, 215.
H. laurina, R. Brown, prot. nov. 29 (1830)...	W.A.	—	—	—	—	—	B.fl. V, 518	M.fr. IV, 130, 216.
H. obtusa, Meissner in Hook. Kew Misc. IV, 209 (1852) ...	W.A.	—	—	—	—	—	B.fl. V, 519	
H. cinerea, R. Brown in Transact. Linn. Soc. X, 186 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 519	
H. corymbosa, R. Brown, prot. nov. 28 (1830) ...	W.A.	—	—	—	—	—	B.fl. V, 519	M.fr. VI, 216.
H. undulata, R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 520	M.fr. VI, 216.
H. petiolaris, Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 520	
H. neurophylla, Meissner in Hook. Kew Misc. VII, 117 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 521	
H. loranthifolia, Meissner in Lehm. pl. Preiss. I, 574 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 521	
H. cucullata, R. Brown, prot. nov. 30 (1830) ...	W.A.	—	—	—	—	—	B.fl. V, 521	M.fr. VI, 216.
H. ferruginea, Sweet, Fl. Austral. t. 45 (1828) ...	W.A.	—	—	—	—	—	B.fl. V, 522	M.fr. VI, 216.
H. smilacifolia, Meissner in Lehm. pl. Preiss. I, 567 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 522	M.fr. VI, 217.
H. elliptica, R. Brown in Transact. Linn. Soc. X, 187 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 523	M.fr. VI, 216.
H. ambigua, Meissner in Lehm. pl. Preiss. II, 260 (1847) ...	W.A.	—	—	—	—	—	B.fl. V, 523	
H. plurinervia, F. v. M. in Benth. Fl. Austr. V, 523 (1870) ...	—	—	—	—	—	Q.	B.fl. V, 523	
H. dactyloides, Cavanilles, Anal. hist. nat. I, 215, t. 12 (1800)...	—	—	—	—	N.S.W.	—	B.fl. V, 524	M.fr. VI, 215.
H. ulicina, R. Brown, prot. nov. 29 (1830)...	—	S.A.	T.	V.	N.S.W.	—	B.fl. V, 524	
H. falcata, R. Brown, prot. nov. 29 (1830)...	W.A.	—	—	—	—	—	B.fl. V, 524	
H. pycnoneura, Meissner in Hook. Kew Misc. VII, 117 (1855) ...	W.A.	—	—	—	—	—	B.fl. V, 525	M.fr. VI, 215.
H. stenocarpa, R. Brown, prot. nov. 29 (1830) ...	W.A.	—	—	—	—	—	B.fl. V, 525	
H. marginata, R. Brown in Transact. Linn. Soc. X, 185 (1809)...	W.A.	—	—	—	—	—	B.fl. V, 525	M.fr. VI, 216.
H. myrtoides, Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 526	
H. costata, Meissner in Lehm. pl. Preiss. I, 575 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 526	
H. oleifolia, R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 527	M.fr. VI, 215.
H. florida, R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 527	
H. varia, R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 527	M.fr. VI, 219.
H. sulcata, R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 528	M.fr. VI, 219.
H. Meissneriana, Kippist in Hook. Kew Misc. VII, 114 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 529	M.fr. VI, 219.
H. subsulcata, Meissner in Lehm. pl. Preiss. I, 555 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 529	M.fr. VI, 219.
H. Lehmanniana, Meissner in Lehm. pl. Preiss. I, 557 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 529	M.fr. VI, 218.
H. flexilis, F. v. M. in Schlecht. Linnaea XXVI, 359 (1853) ...	—	S.A.	—	V.	N.S.W.	—	B.fl. V, 530	M.fr. VI, 216.
H. nitida, R. Brown in Transact. Linn. Soc. X, 184 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 530	M.fr. V, 72; VI, 217.
H. Oldfieldii, Benth, Fl. Austr. V, 530 (1870) ...	W.A.	—	—	—	—	—	B.fl. V, 530	
H. suaveolens, R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 531	M.fr. VI, 220.
H. lissocarpa, R. Brown, prot. nov. 27 (1830) ...	W.A.	—	—	—	—	—	B.fl. V, 531	M.fr. VI, 219.
H. bipinnatifida, R. Brown, prot. nov. 28 (1830) ...	W.A.	—	—	—	—	—	B.fl. V, 532	
CANARVONIA, F. v. M., fragm. VI, 81 (1867).								
C. aralifolia, F. v. M., fragm. VI, t. 55 (1867) ...	—	—	—	—	—	Q.	B.fl. V, 410	M.fr. VI, 81, 247; VII, 59.
BUCKINGHAMIA, F. v. M., fragm. VI, 248 (1868).								
B. celsissima, F. v. M., fragm. VI, 248 (1868) ...	—	—	—	—	—	Q.	B.fl. V, 532	M.fr. VI, 248.
DARLINGIA, F. v. M., fragm. V, 152 (1866).								
D. spectatissima, F. v. M., fragm. V, 152 (1866) ...	—	—	—	—	—	Q.	B.fl. V, 533	M.fr. X, 90.
CARDWELLIA, F. v. M., fragm. V, 23 (1865).								
C. sublimis, F. v. M., fragm. V, 24 (1865)...	—	—	—	—	—	Q.	B.fl. V, 538	M.fr. V, 24.
STENOCARPUS, R. Brown in Transact. Linn. Soc. X, 201 (1809). (Cybele, Agnostus.)								
S. sinuatus, Endlicher, gen. pl. suppl. IV, 88 (1847) ...	—	—	—	—	N.S.W.	Q.	B.fl. V, 539	M.fr. VI, 224. [VII, 133.
S. salignus, R. Brown in Transact. Linn. Soc. X, 202 (1809) ...	—	—	—	—	N.S.W.	Q.	B.fl. V, 539	M.fr. I, 135, 244; VI, 224;
S. Cunninghamii, R. Brown, prot. nov. 34 (1830)...	—	—	—	—	—	N.A.	B.fl. V, 540	
LOMATIA, R. Brown in Transact. Linn. Soc. X, 199 (1809). (Tricondylus.)								
L. fraxinifolia, F. v. M. in Benth. Fl. Austr. V, 536 (1870) ...	—	—	—	—	—	Q.	B.fl. V, 536	
L. ilicifolia, R. Brown in Transact. Linn. Soc. X, 200 (1809) ...	—	—	—	V.	N.S.W.	—	B.fl. V, 536	M.fr. VII, 133.
L. longifolia, R. Brown in Transact. Linn. Soc. X, 200 (1809) ...	—	—	—	V.	N.S.W.	—	B.fl. V, 537	
L. silaifolia, R. Brown in Transact. Linn. Soc. X, 199 (1809) ...	—	—	—	—	N.S.W.	Q.	B.fl. V, 537	M.fr. V, 153; VI, 191.
L. tinctoria, R. Brown in Transact. Linn. Soc. X, 199 (1809) ...	—	—	—	T.	—	—	B.fl. V, 537	M.fr. VI, 224.
L. polymorpha, R. Brown in Transact. Linn. Soc. X, 200 (1809) ...	—	—	—	T.	—	—	B.fl. V, 538	
EMBOTHRUM, R. & G. Forster, charact. gen. 15, t. 8 (1776).								
E. Wickhami, F. v. M., fragm. VIII, 164 (1874)...	—	—	—	—	—	Q.	—	M.fr. VIII, 164; IX, 194.
TELOPEA, R. Brown in Transact. Linn. Soc. X, 197 (1809).								
T. speciosissima, R. Brown in Transact. Linn. Soc. X, 198 (1809) ...	—	—	—	—	N.S.W.	—	B.fl. V, 534	M.fr. X, 90.
T. oreades, F. v. M., Annual Report 18 (1861) ...	—	—	—	V.	—	—	B.fl. V, 534	M.fr. II, 170; VII, 133.
T. truncata, R. Brown in Transact. Linn. Soc. X, 198 (1809) ...	—	—	—	T.	—	—	B.fl. V, 535	M.fr. II, 171.
BANKSIA, Linné, fl. suppl. 15 et 126 (1781).								
B. pulchella, R. Brown in Transact. Linn. Soc. X, 202 (1809) ...	W.A.	—	—	—	—	—	B.fl. V, 544	M.fr. VII, 54.
B. Meissneri, Lehmann, plant. Preiss. I, 582 (1845) ...	W.A.	—	—	—	—	—	B.fl. V, 545	

B. nutans, R. Brown in Transact. Linn. Soc. X, 203 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 545	M.fr. IV, 108; VII, 54.
B. sphaerocarpa, R. Brown in Transact. Linn. Soc. X, 203 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 546	M.fr. VII, 54.
B. tricuspid, Meissner in Hook. Kew Misc. VII, 118 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 546	M.fr. VII, 54.
B. occidentalis, R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 546	M.fr. VII, 54.
B. littoralis, R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 547	M.fr. VII, 55.
B. ericifolia, Linné, fil. suppl. 127 (1781) ...	—	—	—	N.S.W.	—	—	B.f. V, 547	M.fr. VII, 54.
B. spinulosa, Smith, Specim. Bot. New Holl. 13, t. 4 (1793) ...	—	—	—	N.S.W.	—	—	B.f. V, 547	
B. collina, R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	—	—	—	V. N.S.W. Q.	—	—	B.f. V, 548	M.fr. VII, 54.
B. verticillata, R. Brown in Transact. Linn. Soc. X, 207 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 548	
B. dryandroides, Baxter in Sweet, Fl. Austral. t. 56 (1828) ...	W.A.	—	—	—	—	—	B.f. V, 549	M.fr. VII, 58.
B. Brownii, Baxter in R. Br. prot. nov. 37 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 549	M.fr. VII, 55.
B. attenuata, R. Brown in Transact. Linn. Soc. X, 209 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 550	M.fr. IV, 109.
B. media, R. Brown, prot. nov. 35 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 550	M.fr. VII, 58.
B. Solandri, R. Brown, prot. nov. 35 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 550	M.fr. VII, 58.
B. Goodii, R. Brown, prot. nov. 36 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 551	M.fr. IV, 109.
B. petiolaris, F. v. M., fragm. IV, 109 (1864) ...	W.A.	—	—	—	—	—	B.f. V, 551	M.fr. IV, 108, 177; VII, 58.
B. repens, Labillardière, Voy. I, 411, t. 23 (1798) ...	W.A.	—	—	—	—	—	B.f. V, 551	M.fr. VII, 57.
B. prostrata, R. Brown, prot. nov. 36 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 552	M.fr. VII, 57.
B. grandis, Willdenow, spec. plant. I, 535 (1797) ...	W.A.	—	—	—	—	—	B.f. V, 552	M.fr. VII, 57.
B. quercifolia, R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 552	M.fr. IV, 107; VII, 56.
B. Baueri, R. Brown, prot. nov. 35 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 553	M.fr. VII, 55, 133; X, 90.
B. marginata, Cavanilles, Anal. hist. nat. I, 227, t. 13 (1800) ...	—	S.A.	T. V.	N.S.W.	—	—	B.f. V, 554	M.fr. VII, 55.
B. integrifolia, Linné, fil. suppl. 127 (1781) ...	—	—	—	V. N.S.W. Q.	—	—	B.f. V, 555	M.fr. VII, 57.
B. dentata, Linné, fil. suppl. 127 (1781) ...	—	—	—	—	Q.	N.A.	B.f. V, 555	M.fr. VII, 56.
B. latifolia, R. Brown in Transact. Linn. Soc. X, 208 (1809) ...	—	—	—	N.S.W. Q.	—	—	B.f. V, 556	M.fr. VII, 56.
B. serrata, Linné, fil. suppl. 126 (1781) ...	—	—	T. V.	N.S.W.	—	—	B.f. V, 556	M.fr. VII, 56.
B. aemula, R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	—	—	—	N.S.W. Q.	—	—	B.f. V, 557	M.fr. VII, 56.
B. ornata, F. v. M. in Schlecht. Linnaea XXVI, 352 (1853) ...	—	S.A.	—	V.	—	—	B.f. V, 557	M.fr. VII, 56.
B. coccinea, R. Brown in Transact. Linn. Soc. X, 207 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 557	M.fr. VII, 56.
B. Sceptum, Meissner in Hook. Kew Misc. VII, 120 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 558	M.fr. VII, 56.
B. Menziesii, R. Brown, prot. nov. 36 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 558	
B. laevigata, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 558	M.fr. VII, 56.
B. Hookeriana, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 558	
B. prionotes, Lindley, Bot. Regist. XXV, App. XXXIV (1839) ...	W.A.	—	—	—	—	—	B.f. V, 559	M.fr. VII, 56.
B. Victoriae, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 559	M.fr. VII, 58.
B. speciosa, R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 559	M.fr. VII, 58.
B. Baxteri, R. Brown, prot. nov. 36 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 560	M.fr. VII, 56.
B. marcescens, R. Brown in Transact. Linn. Soc. X, 208 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 560	M.fr. VII, 57.
B. Lemniana, Meissner in Hook. Kew Misc. 210 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 560	M.fr. VII, 57.
B. Caleyi, R. Brown, prot. nov. 35 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 561	M.fr. VII, 55.
B. Lindleyana, Meissner in Hook. Kew Misc. VII, 120 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 561	
B. elegans, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 561	M.fr. VII, 58.
B. Candolleana, Meissner in Hook. Kew Misc. VII, 118 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 561	M.fr. VII, 58.
B. ilicifolia, R. Brown in Transact. Linn. Soc. X, 211 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 561	
DRYANDRA, R. Brown in Transact. Linn. Soc. X, 211 (1809). (Josephia, Hemicleidia.)								
D. quercifolia, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 566	M.fr. VII, 50.
D. praemorsa, Meissner in Lehm. pl. Preiss. II, 265 (1847) ...	W.A.	—	—	—	—	—	B.f. V, 566	M.fr. VII, 50.
D. cuneata, R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 567	M.fr. VII, 50.
D. falcata, R. Brown in Transact. Linn. Soc. X, 213 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 567	
D. armata, R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 568	M.fr. VII, 52.
D. longifolia, R. Brown in Transact. Linn. Soc. X, 215 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 568	M.fr. VI, 92; VII, 50.
D. Fraseri, R. Brown, prot. nov. 39 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 569	M.fr. VII, 51.
D. floribunda, R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 569	M.fr. VII, 53.
D. carduacea, Lindley, Bot. Regist. XXV, App. XXXIII (1839) ...	W.A.	—	—	—	—	—	B.f. V, 570	M.fr. VII, 52.
D. carlinoides, Meissner in Lehm. pl. Preiss. II, 267 (1847) ...	W.A.	—	—	—	—	—	B.f. V, 571	M.fr. VII, 52.
D. polycephala, Bentham, Fl. Austr. V, 570 (1870) ...	W.A.	—	—	—	—	—	B.f. V, 571	M.fr. VII, 51.
D. Kippistiana, Meissner in Hook. Kew Misc. VII, 122 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 571	M.fr. VII, 52.
D. squarrosa, R. Brown, prot. nov. 38 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 572	M.fr. VII, 51.
D. Serra, R. Brown, prot. nov. 38 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 572	M.fr. VII, 51.
D. concinna, R. Brown, prot. nov. 38 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 572	
D. foliolata, R. Brown, prot. nov. 38 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 573	M.fr. VII, 51.
D. stuppea, Lindley, Bot. Regist. XXV, App. XXXIII (1839) ...	W.A.	—	—	—	—	—	B.f. V, 573	M.fr. VII, 52.
D. nobilis, Lindley, Bot. Regist. XXV, App. XXXIII (1839) ...	W.A.	—	—	—	—	—	B.f. V, 573	
D. mucronulata, R. Brown in Transact. Linn. Soc. X, 213 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 574	M.fr. VII, 52.
D. furcata, R. Brown in Transact. Linn. Soc. X, 213, t. 3 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 574	
D. Baxteri, R. Brown, prot. nov. 38 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 575	M.fr. VII, 52.
D. nivea, R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 575	M.fr. VII, 54.
D. Aretotidis, R. Brown, prot. nov. 39 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 576	M.fr. VII, 52.
D. nana, Meissner in Hook. Kew Misc. VII, 121 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 576	
D. Preissii, Meissner in Lehm. pl. Preiss. I, 599 (1845) ...	W.A.	—	—	—	—	—	B.f. V, 576	M.fr. VII, 51.
D. sclerophylla, Meissner in Hook. Kew Misc. VII, 123 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 577	M.fr. VII, 53.
D. pulchella, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 577	M.fr. VII, 52.
D. plumosa, R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	W.A.	—	—	—	—	—	B.f. V, 577	M.fr. VII, 53.
D. senecionifolia, R. Brown, prot. nov. 39 (1830) ...	W.A.	—	—	—	—	—	B.f. V, 578	
D. vestita, Kippist in Hook. Kew Misc. VII, 121 (1855) ...	W.A.	—	—	—	—	—	B.f. V, 578	
D. cirsioides, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 578	
D. Hewardiana, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	B.f. V, 578	
D. patens, Bentham, Fl. Austr. V, 578 (1870) ...	W.A.	—	—	—	—	—	B.f. V, 579	M.fr. VII, 52.
D. conferta, Bentham, Fl. Austr. V, 578 (1870) ...	W.A.	—	—	—	—	—		
D. horrida, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—		

<i>D. serratuloides</i> , Meissner in Hook. Kew Misc. VII, 123 (1855)	W.A.	—	—	—	—	—	—	B.f.V, 579	M.fr. VII, 52.
<i>D. comosa</i> , Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.f.V, 579	M.fr. VII, 53.
<i>D. Shuttleworthiana</i> , Meissner in Hook. Kew Misc. VII, 122 (1855)	W.A.	—	—	—	—	—	—	B.f.V, 580	M.fr. VII, 52.
<i>D. speciosa</i> , Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.f.V, 580	
<i>D. tridentata</i> , Meissner in Hook. Kew Misc. VII, 120 (1855)	W.A.	—	—	—	—	—	—	B.f.V, 580	M.fr. VII, 53.
<i>D. tenuifolia</i> , R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.f.V, 581	M.fr. VII, 53.
<i>D. proteoides</i> , Lindley, Bot. Regist. XXV, App. XXXIII (1839)	W.A.	—	—	—	—	—	—	B.f.V, 582	
<i>D. runcinata</i> , Meissner in Hook. Kew Misc. IV, 210 (1852)	W.A.	—	—	—	—	—	—	B.f.V, 582	
<i>D. obtusa</i> , R. Brown in Transact. Linn. Soc. X, 214 (1809)	W.A.	—	—	—	—	—	—	B.f.V, 582	
<i>D. bipinnatifida</i> , R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.f.V, 583	M.fr. V, 185.
<i>D. pteridifolia</i> , R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.f.V, 583	M.fr. VII, 54.
<i>D. calophylla</i> , R. Brown, prot. nov. 40 (1830)	W.A.	—	—	—	—	—	—	B.f.V, 583	M.fr. VII, 54.

THYMELEAE.

A. L. de Jussieu, gen. plant. 76 (1789).

PIMELEA, Banks & Solander in Gaert. de fruct. I, 186 (1788). (Thecanthes, Gymnococca, Heterolaena, Calyptrastegia, Macrosteigia;—Banksia, Forst., 1776.)

<i>P. punicea</i> , R. Brown, prodr. 359 (1810)	—	—	—	—	—	N.A.	B.f.VI, 6	M.fr. V, 74; VII, 3.
<i>P. concreta</i> , F. v. M., fragm. V, 73 (1865)	—	—	—	—	—	N.A.	B.f.VI, 6	M.fr. V, 73.
<i>P. cornucopiae</i> , Vahl, enum. plant. I, 305 (1804)	—	—	—	—	—	Q.	B.f.VI, 6	M.fr. VII, 3.
<i>P. sanguinea</i> , F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	B.f.VI, 7	M.fr. VII, 3; XI, 47.
<i>P. alpina</i> , F. v. M., second general report, 14 (1854)	—	—	—	—	V. N.S.W.	—	B.f.VI, 7	M.fr. VII, 4.
<i>P. longifolia</i> , Banks & Solander in Rees, Cycl. XXVI (1814)	—	—	—	—	N.S.W.	—	B.f.VI, 7	M.fr. VIII, 9.
<i>P. cinerea</i> , R. Brown, prodr. 361 (1810)	—	—	—	T.	—	—	B.f.VI, 8	
<i>P. Milligani</i> , Meissner in De Candolle, prodr. XIV, 509 (1856)	—	—	—	T.	—	—	B.f.VI, 8	
<i>P. spectabilis</i> , Lindley, Bot. Regist. t. 33 (1841)	W.A.	—	—	—	—	—	B.f.VI, 9	M.fr. VII, 4.
<i>P. Lehmanniana</i> , Meissner in Lehm. pl. Preiss. I, 603 (1845)	W.A.	—	—	—	—	—	B.f.VI, 9	M.fr. VII, 4.
<i>P. hispida</i> , R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	B.f.VI, 10	
<i>P. rosea</i> , R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	B.f.VI, 10	
<i>P. ferruginea</i> , Labillardière, Nov. Holl. pl. spec. I, 10, t. 5 (1804)	W.A.	—	—	—	—	—	B.f.VI, 10	M.fr. VII, 4.
<i>P. brachyphylla</i> , Benth. Fl. Austr. VI, 11 (1873)	W.A.	—	—	—	—	—	B.f.VI, 11	
<i>P. sylvestris</i> , R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	B.f.VI, 11	M.fr. VII, 2&5.
<i>P. brevifolia</i> , R. Brown, prodr. 359 (1810)	W.A.	—	—	—	—	—	B.f.VI, 12	M.fr. VII, 4.
<i>P. Maxwelli</i> , F. v. M. in Benth. Fl. Austr. VI, 12 (1873)	W.A.	—	—	—	—	—	B.f.VI, 12	
<i>P. angustifolia</i> , R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	B.f.VI, 13	M.fr. VII, 3.
<i>P. nervosa</i> , Meissner in Lehm. pl. Preiss. II, 269 (1847)	W.A.	—	—	—	—	—	B.f.VI, 13	M.fr. VII, 3.
<i>P. sulphurea</i> , Meissner in Mohl & Schlecht. Bot. Zeitung, 396 (1848)	W.A.	—	—	—	—	—	B.f.VI, 14	M.fr. VII, 5.
<i>P. floribunda</i> , Meissner in De Candolle, prodr. XIV, 505 (1856)	W.A.	—	—	—	—	—	B.f.VI, 14	M.fr. VII, 5.
<i>P. suaveolens</i> , Meissner in Lehm. pl. Preiss. I, 603 (1845)	W.A.	—	—	—	—	—	B.f.VI, 14	M.fr. VII, 2, 4, 5.
<i>P. physodes</i> , Hooker, icon. plant. t. 865 (1852)	W.A.	—	—	—	—	—	B.f.VI, 15	M.fr. VII, 4.
<i>P. glauca</i> , R. Brown, prodr. 360 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	B.f.VI, 15	M.fr. VII, 4; XI, 123.
<i>P. colorans</i> , Cunningham in De Cand. prodr. XIV, 499 (1856)	—	—	—	—	N.S.W.	—	B.f.VI, 16	
<i>P. collina</i> , R. Brown, prodr. 359 (1810)	—	—	—	—	V. N.S.W.	Q.	B.f.VI, 16	
<i>P. spatulata</i> , Labillardière, Nov. Holl. pl. spec. I, 9, t. 4 (1804)	—	S.A.	T.	V.	N.S.W.	—	B.f.VI, 17	
<i>P. linifolia</i> , Smith, Specim. of Bot. of N. Holl. 31, t. 11 (1793)	—	—	T.	V.	N.S.W.	Q.	B.f.VI, 17	M.fr. VII, 3.
<i>P. ligustrina</i> , Labillardière, Nov. Holl. pl. spec. I, 9, t. 3 (1804)	—	S.A.	T.	V.	N.S.W.	—	B.f.VI, 18	M.fr. I, 87; IV, 163; VII, 5.
<i>P. humilis</i> , R. Brown, prodr. 361 (1810)	—	S.A.	T.	V.	N.S.W.	—	B.f.VI, 19	M.fr. VII, 4; XI, 47.
<i>P. sericea</i> , R. Brown, prodr. 361 (1810)	—	—	T.	—	—	—	B.f.VI, 19	M.fr. VII, 5.
<i>P. nivea</i> , Labillardière, Nov. Holl. pl. spec. I, 10, t. 6 (1804)	—	—	T.	—	—	—	B.f.VI, 20	M.fr. VII, 5.
<i>P. imbricata</i> , R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	B.f.VI, 20	M.fr. VII, 5.
<i>P. villifera</i> , Meissner in Lehm. pl. Preiss. II, 271 (1847)	W.A.	—	—	—	—	—	B.f.VI, 21	
<i>P. drupacea</i> , Labillardière, Nov. Holl. pl. spec. I, 10, t. 7 (1804)	—	—	T.	V.	—	—	B.f.VI, 21	M.fr. VII, 8.
<i>P. haematostachya</i> , F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	B.f.VI, 22	M.fr. I, 84; VII, 8; XI, 47.
<i>P. spicata</i> , R. Brown, prodr. 362 (1810)	—	—	—	—	N.S.W.	—	B.f.VI, 22	M.fr. VII, 7; VIII, 9.
<i>P. spiculigera</i> , F. v. M. in Benth. Fl. Austr. VI, 23 (1873)	W.A.	—	—	—	—	—	B.f.VI, 23	M.fr. XI, 46.
<i>P. Forrestiana</i> , F. v. M., fragm. XI, 46 (1879)	—	—	—	—	—	N.A.	—	M.fr. XI, 46.
<i>P. filiformis</i> , J. Hooker in Lond. Journ. VI, 280 (1847)	—	—	T.	—	—	—	B.f.VI, 23	M.fr. VII, 7.
<i>P. latifolia</i> , R. Brown, prodr. 362 (1810)	—	—	—	—	—	Q.	B.f.VI, 23	M.fr. IV, 49; VII, 7.
<i>P. simplex</i> , F. v. M. in Linnaea XXV, 443 (1852)	—	S.A.	—	V.	N.S.W.	—	B.f.VI, 23	
<i>P. sericostachya</i> , F. v. M., fragm. IV, 162 (1864)	—	—	—	—	N.S.W.	Q.	B.f.VI, 24	M.fr. XI, 47.
<i>P. trichostachya</i> , Lindley in Mitch. Trop. Austr. 355 (1848)	—	—	—	—	N.S.W.	Q.	B.f.VI, 24	M.fr. VII, 7.
<i>P. leptostachya</i> , Benth. Fl. Austr. VI, 24 (1873)	—	—	—	—	—	Q.	B.f.VI, 24	
<i>P. argentea</i> , R. Brown, prodr. 362 (1810)	W.A.	—	—	—	—	—	B.f.VI, 25	M.fr. VII, 7.
<i>P. clavata</i> , Labillardière, Nov. Holl. pl. spec. I, 11 (1804)	W.A.	—	—	—	—	—	B.f.VI, 25	M.fr. VI, 160; VII, 5.
<i>P. axiflora</i> , F. v. M., First general Report 17 (1853)	—	—	T.	V.	N.S.W.	—	B.f.VI, 26	M.fr. VII, 7.
<i>P. leptospermoides</i> , F. v. M., fragm. VII, 2 (1870)	—	—	—	—	—	Q.	B.f.VI, 27	M.fr. VII, 2.
<i>P. microcephala</i> , R. Brown, prodr. 361 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	B.f.VI, 27	M.fr. XI, 27, 47.
<i>P. pauciflora</i> , R. Brown, prodr. 360 (1810)	—	—	T.	V.	N.S.W.	Q.	B.f.VI, 27	M.fr. VII, 8; XI, 47.
<i>P. elachantha</i> , F. v. M., First general Report 17 (1853)	—	—	—	V.	—	—	B.f.VI, 28	M.fr. VII, 6.
<i>P. pygmaea</i> , F. v. M., in Schlecht. Linnaea XXVI, 346 (1853)	—	—	T.	—	—	—	B.f.VI, 28	
<i>P. serpyllifolia</i> , R. Brown, prodr. 360 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.VI, 29	M.fr. VII, 6.
<i>P. flava</i> , R. Brown, prodr. 361 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f.VI, 29	M.fr. VII, 6.
<i>P. petrophila</i> , F. v. M. in Schlecht. Linnaea XXV, 442 (1852)	—	S.A.	—	—	N.S.W.	—	B.f.VI, 30	
<i>P. Bowmanni</i> , F. v. M. in Benth. Fl. Austr. VI, 30 (1873)	—	—	—	—	—	Q.	B.f.VI, 30	M.fr. VII, 6.
<i>P. ammodharis</i> , F. v. M. in Hook. Kew Misc. IX, 24 (1857)	—	—	—	—	—	N.A.	B.f.VI, 30	M.fr. VII, 5; XI, 47.
<i>P. curviflora</i> , R. Brown, prodr. 362 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	B.f.VI, 31	M.fr. VII, 6; VIII, 9.
<i>P. hirsuta</i> , Meissner in De Cand. prodr. XIV, 513 (1856)	—	—	—	—	N.S.W.	—	B.f.VI, 32	
<i>P. altior</i> , F. v. M., fragm. I, 84 (1858)	—	—	—	—	N.S.W.	Q.	B.f.VI, 32	M.fr. VII, 7.
<i>P. octophylla</i> , R. Brown, prodr. 361 (1810)	—	S.A.	—	V.	N.S.W.	—	B.f.VI, 32	M.fr. VII, 6.

P. petraea, Meissner in Schlecht. Linnaea XXVI, 347 (1853) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.VI,33	
P. phylloides, Meissner in Lehm. pl. Preiss. II, 271 (1847) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,33	M.fr.VII,6.
P. Eyrei, F. v. M., fragm. V, 109 (1865) ...	...	W.A.	—	—	—	—	—	B.fl.VI,34	M.fr.V,109.
P. longiflora, R. Brown, prodr. 361 (1810)...	...	W.A.	—	—	—	—	—	B.fl.VI,34	M.fr.VII,5.
P. stricta, Meissner in Schlecht. Linnaea XXVI, 348 (1853) ...	...	S.A.	T.	V.	—	—	—	B.fl.VI,34	M.fr.VII,4.
P. Preissii, Meissner in Lehm. pl. Preiss. I, 601 (1845) ...	...	W.A.	—	—	—	—	—	B.fl.VI,35	M.fr.VII,3.
P. Holroydi, F. v. M., fragm. VI, 159, t. 59 (1868) ...	...	—	—	—	—	—	N.A.	B.fl.VI,35	M.fr.250.
DRAPETES, Lamarck, Journ. d'hist. nat. I, 119, t. 10 (1792). (Kelleria, Daphnobryon).									
D. Tasmanica, J. Hooker in Kew Misc. V, 299, t. 7 (1853) ...	...	—	T.	V.	N.S.W.	—	—	B.fl.VI,36	M.fr.VII,1.
WICKSTROEMIA, Endlicher, prodr. fl. insul. Norfolk. 47 (1833). (Daphne partly).									
W. Indica, C. A. Meyer in Bull. de l'Acad. de Petersb. I, 357 (1843) ...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,37	M.fr.VII,1.
PHALERIA, Jack in Hook. Comp. to Bot. Mag. I, 156 (1835). (Drymispermum).									
P. Blumei, Benth. Fl. Austr. VI, 38 (1873) ...	...	—	—	—	—	—	Q.	B.fl.VI,38	
P. Neumannii, F. v. M., fragm. VIII, 9 (1873) ...	...	—	—	—	—	—	Q.	B.fl.VI,38	M.fr.V,26.
P. Clerodendron, F. v. M., fragm. VIII, 9 (1873) ...	...	—	—	—	—	—	Q.	B.fl.VI,38	M.fr.VII,1.

CORNACEAE.

Humboldt, Bonpland & Kunth, nov. gen. Americ. III, 430 (1818).

STYLIDIUM, Loureiro, Fl. Cochinch. I, 220 (1790). (Marlea, Rhytidandra, Pseudalangium).

S. Vitiense, F.v.M.; Rhytid., Asa Gray, Bot. Wilk. Ex. I, 303, t. 28 (1854) — — — N.S.W. Q. — B.fl.III, 386

RUBIACEAE.

A. L. de Jussieu, gen. plant. 196 (1789), from B. de Jussieu (1759).

SARCOCEPHALUS, Afzelius in Transact. hort. soc. Lond. V, 422, t. 18 (1824).

S. cordatus, Miquel, Fl. Ind. Batav. II, 133 (1856) ... — — — — — Q. N.A. B.fl.III, 492

OLDENLANDIA, Plumier, nov. plant. Americ. gen. 42, t. 36 (1703). (Hedyotis, Synaptantha).

O. corymbosa, Linné, sp. pl. 119 (1753) ...	...	—	—	—	—	—	Q.	—	M.fr.VII,45.
O. paniculata, Linné, spec. plant. edit. sec. 1667 (1763) ...	...	—	—	—	—	—	Q.	—	M.fr.VIII,47.
O. auricularia, F. v. M.; Hedyotis, Linné, spec. plant. 101 (1753) ...	...	—	—	—	—	—	Q.	B.fl.III,404	
O. polyclada; Hedyotis, F. v. M., fragm. VIII, 146 (1874) ...	...	—	—	—	—	—	Q.	—	M.fr.VIII,146.
O. coerulescens; Hedyotis, F. v. M., fragm. IV, 38 (1863) ...	...	—	—	—	—	—	Q.	B.fl.III,404	M.fr.IV,38.
O. spermacocoides; Hedyotis, F. v. M., fragm. VIII, 146 (1874) ...	...	—	—	—	—	—	N.A.	—	M.fr.VIII,146.
O. Crouchiana; Hedyotis, F. v. M., fragm. X, 85 (1876) ...	...	—	—	—	—	—	N.A.	—	M.fr.X,85.
O. mitrasacmoides; Hedyotis, F. v. M., fragm. IV, 37 (1863) ...	...	—	—	—	—	—	N.A.	B.fl.III,404	M.fr.IV,37.
O. galioides; Hedyotis, F. v. M., fragm. IV, 38 (1863) ...	...	—	—	—	—	—	Q.	N.A. B.fl.III,404	M.fr.IV,38.
O. scleranthoides; Hedyotis, F. v. M., fragm. IV, 39 (1863) ...	...	—	—	—	—	—	N.A.	B.fl.III,405	M.fr.IV,39.
O. elatinoides, F. v. M.; Hedyotis, Benth., Fl. Austr. III, 405 (1866) ...	...	W.A.	—	—	—	—	—	B.fl.III,405	M.fr.VIII,147.
O. tillaeacea; Hedyotis, F. v. M., fragm. IV, 39 (1863) ...	...	—	S.A.	—	N.S.W.	Q.	—	B.fl.III,405	M.fr.IV,39.
O. trachymenioides; Hedyotis, F. v. M., fragm. IV, 40 (1863) ...	...	—	—	—	—	—	Q.	B.fl.III,406	M.fr.IV,40.
O. pteropora; Hedyotis, F. v. M., fragm. IV, 40 (1863) ...	...	—	—	—	—	—	N.A.	B.fl.III,407	M.fr.IV,40,177.

DENTELLA, R. et G. Forster, char. gen. 25, t. 13 (1776). (Lippaya).

D. repens, R. & G. Forster, char. gen. 25, t. 13 (1776) ... — S.A. — — N.S.W. Q. N.A. B.fl.III,407 M.fr.IX,187.

OPHIORRHIZA, Linné, fl. Zeylan. 190 et 239 (1747).

O. Australiana, Benth. Fl. Austr. III, 407 (1866) ... — — — — — Q. — B.fl.III,408 M.fr.IX,187.

ABBOTTIA, F. v. M., fragm. IX, 181 (1875).

A. singularis, F. v. M., fragm. IX, 181 (1875) ... — — — — — Q. — — M.fr.IX,181.

GARDENIA, Ellis in Phil. Transact. LI, 935, t. 23 (1761).

G. edulis, F. v. M., fragm. I, 54 (1858) ...	...	—	—	—	—	—	N.A.	B.fl.III,408	M.fr.I,54.
G. resinosa, F. v. M., fragm. I, 54 (1858) ...	...	—	—	—	—	—	N.A.	B.fl.III,408	M.fr.I,54.
G. pyriformis, Cunningham in Benth. Fl. Austr. III, 409 (1866) ...	...	—	—	—	—	—	N.A.	B.fl.III,409	
G. megasperma, F. v. M., fragm. I, 54 (1858) ...	...	—	—	—	—	—	N.A.	B.fl.III,409	M.fr.I,54.
G. Macgillivraei, Benth. Fl. Austr. III, 409 (1866) ...	...	—	—	—	—	—	Q.	B.fl.III,409	M.fr.VII,46.
G. ochreatea, F. v. M., fragm. I, 55 (1858) ...	...	—	—	—	—	—	Q.	B.fl.III,409	M.fr.155;VII,46.
G. suffruticosa, R. Brown in Benth. Fl. Austr. III, 410 (1866) ...	...	—	—	—	—	—	N.A.	B.fl.III,410	
G. fucata, R. Brown in Benth. Fl. Austr. III, 410 (1866) ...	...	—	—	—	—	—	N.A.	B.fl.III,410	
G. Jardinei, F. v. M. in Benth. Fl. Austr. III, 410 (1866) ...	...	—	—	—	—	—	Q.	B.fl.III,410	M.fr.VII,46.

RANDIA, Houston in Linné, hort. Cliffort. 485 (1737).

R. hirta, F. v. M., fragm. VII, 46 (1869) ...	...	—	—	—	—	—	Q.	—	M.fr.IX,181.
R. chartacea, F. v. M., fragm. IX, 180 (1875) ...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,410	M.fr.VII,47.
R. Benthiana, F. v. M., fragm. IX, 180 (1875) ...	...	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,180.
R. Moorei, F. v. M. in Benth. Fl. Austr. III, 411 (1866) ...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,411	
R. stipularis, F. v. M., Papuan plants I, 69 (1876) ...	...	—	—	—	N.S.W.	—	—	—	M.fr.VII,47;IX,70,181.
R. Fitzalani, F. v. M., Rep. Burdek. Exped. 12 (1860) ...	...	—	—	—	—	—	Q.	B.fl.III,411	M.fr.V,19;IX,180.
R. sessilis, F. v. M., fragm. VII, 47 (1869) ...	...	—	—	—	—	—	Q.	N.A.	M.fr.VII,47. [180.
R. densiflora, Benth. Fl. Hongk. 155 (1861) ...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,412	M.fr.II,132; III, 166; IX,

DIPLOSPORA, De Candolle, prodr. IV, 477 (1830). (Discospermum.)

D. Australis, Benth. Fl. Austr. III, 413 (1866) ...	...	—	—	—	—	—	Q.	—	B.fl.III,413
D. ixoroides, F. v. M., fragm. IX, 182 (1875) ...	...	—	—	—	—	—	Q.	—	B.fl.III,416 M.fr.IX,182.

IXORA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 27 (1737). (Pavetta, Webera, Stylocorne.)

I. Pavetta, Roxburgh, Fl. Ind. I, 395, ed. Carey (1820) ...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,414 M.fr.IX,182.
I. tomentosa, Roxburgh, Fl. Ind. I, 396, ed. Carey (1820) ...	...	—	—	—	—	—	—	N.A.	B.fl.III,414
I. coccinea, Linné, sp. pl. 110 (1753) ...	...	—	—	—	—	—	—	N.A.	B.fl.III,415
I. Timorensis, Decaisne in Nouv. Ann. du Mus. III, 419 (1834) ...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,415 M.fr.V,18;IX,183.
I. Becklerii, Benth. Fl. Austr. III, 415 (1866) ...	...	—	—	—	N.S.W.	Q.	—	B.fl.III,415	M.fr.IX,183.

I. pentamera, Benth., Fl. Austr. III, 416 (1866)	...	...	...	...	...	N.A.	B.f. III, 416	
I. dallachyana, F. v. M., Webera, Benth., Fl. Austr. III, 412 (1866)	...	...	...	...	...	Q.	B.f. III, 412	
I. expandens, Webera, F. v. M., fragm. VI, 25 (1867)	...	...	...	...	...	N.A.	—	M.fr. VI, 25.
COFFEA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 55 (1737), from Ray (1691).								
C. bengalensis, Roxburgh, hort. Bengal. 15 (1814)	...	...	...	...	...	Q.	—	M.fr. VIII, 147.
TIMONIUS, Rumphius, herb. Amboin. III, 216, t. 140 (1743). (Polyphragmon.)								
T. Rumphii, De Candolle, prodr. IV, 461 (1830)	...	...	...	...	...	Q.	N.A.	B.f. III, 417 M.fr. II, 134; IX, 157.
SCYPHIPHORA, K. F. Gaertner, de fruct. III, 91, t. 196 (1805). (Epithinia.)								
S. hydrophylacea, K. F. Gaertner, de fruct. III, 91, t. 196 (1805)	...	...	...	...	...	Q.	N.A.	B.f. III, 415 M.fr. VII, 46; IX, 157.
GUETTARDA, Osbeck in Linné, spec. plant. 991 (1753). (Antirrhoea, Guettardella.)								
G. speciosa, Linné, spec. plant. 991 (1753)	...	...	...	...	...	Q.	N.A.	B.f. III, 419 M.fr. IX, 183.
G. tenuiflora, F. v. M., fragm. VII, 48 (1869)	...	...	...	...	...	Q.	—	B.f. III, 415 M.fr. IX, 183.
G. putaminosa, F. v. M., fragm. IX, 153 (1875)	...	...	...	...	...	Q.	—	B.f. III, 419 M.fr. IV, 92.
G. myrtoides, F. v. M., fragm. IX, 184 (1875)	...	...	...	...	...	Q.	—	M.fr. IX, 184.
HODGKINSONIA, F. v. M., fragm. II, 132 (1861).								
H. ovatiflora, F. v. M., fragm. II, 132 (1861)	...	...	...	...	...	N.S.W.	Q.	— B.f. III, 420 M.fr. III, 166.
CANTHIUM, Lamarck, Encycl. method. I, 602 (1783). (Psydrax, Electronia partly.)								
C. cymosum, Persoon, synops. plant. I, 200 (1805)	...	...	...	...	...	Q.	—	M.fr. IX, 155.
C. latifolium, F. v. M. in Benth. Fl. Austr. III, 421 (1866)	...	W.A.	S.A.	—	N.S.W.	Q.	—	B.f. III, 421
C. attenuatum, R. Brown in Benth. Fl. Austr. III, 421 (1866)	...	—	—	—	—	N.A.	B.f. III, 421	[135.
C. lucidum, Hooker & Arnott, Bot. of Beechey's voy. 65 (1841)	...	—	—	—	N.S.W.	Q.	N.A.	B.f. III, 421 M.fr. II, 133; IX, 185; XI,
C. oleifolium, Hooker in Mitch. Trop. Austr. 397 (1845)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 422
C. buxifolium, Benth., Fl. Austr. III, 422 (1866)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 422
C. vacciniifolium, F. v. M. in Trans. phil. Inst. Vict. III, 47 (1855)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 422 M.fr. II, 134; IX, 186.
C. coprosimoides, F. v. M. in Trans. phil. Inst. Vict. III, 47 (1858)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 422
MORINDA, Vaillant in act. Acad. Par. 202 (1722).								
M. citrifolia, Linné, spec. plant. 176 (1753)	...	...	...	...	...	Q.	N.A.	B.f. III, 422
M. acutifolia, F. v. M., fragm. IX, 179 (1875)	...	...	...	...	...	Q.	—	B.f. III, 421 M.fr. IX, 179.
M. jasminoides, Cunningham in Hook. Bot. Mag. t. 3351 (1834)	...	—	—	—	V. N.S.W.	Q.	—	B.f. III, 423
M. umbellata, Linné, spec. plant. 176 (1753)	...	...	...	...	...	Q.	—	B.f. III, 423
M. reticulata, Benth., Fl. Austr. III, 424 (1866)	...	...	...	...	...	Q.	N.A.	B.f. III, 423 M.fr. IX, 180.
COELOSPERMUM, Blume, Bijdr. 994 (1826). (Pogonolobus.)								
C. paniculatum, F. v. M., fragm. V, 19 (1865)	...	...	...	...	N.S.W.	Q.	—	B.f. III, 425 M.fr. VII, 46; IX, 185.
C. reticulatum, Benth., Fl. Austr. III, 425 (1866)	...	...	...	...	...	Q.	N.A.	B.f. III, 425 M.fr. I, 56; IX, 185.
HYDNOPHYTUM, Jack in Transact. Linn. Soc. XIV, 124 (1823).								
H. formicarum, Jack in Transact. Linn. Soc. XIV, 124 (1823)	...	—	—	—	—	Q.	N.A.	— M.fr. VII, 45.
MYRMECODIA, Jack in Transact. Linn. Soc. XIV, 122 (1823).								
M. echinata, Gaudichaud, Bot. voy. Bonite 472, t. 96 (1844)	...	—	—	—	—	Q.	N.A.	— M.fr. VII, 45.
LASIANTHUS, Jack in Transact. Linn. Soc. XIV, 125 (1823).								
L. strigosus, Wight in Calc. Journ. nat. hist. VI, 512 (1846)	...	—	—	—	—	Q.	—	B.f. III, 426
PSYCHOTRIA, Linné, syst. ed. decim. 929 (1759). (Uragoga, 1737, Grumilia.)								
P. nesophila, F. v. M., fragm. II, 135 (1861)	...	...	...	...	...	Q.	N.A.	B.f. III, 427 M.fr. IX, 184.
P. dallachyana, Benth., Fl. Austr. III, 427 (1866)	...	...	...	...	...	Q.	—	B.f. III, 427
P. nematopoda, F. v. M., fragm. VII, 45 (1869)	...	...	...	...	...	Q.	—	M.fr. IX, 184.
P. loniceroides, Sieber in De Cand., prodr. IV, 523 (1830)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 427 M.fr. IX, 184.
P. daphnoides, Cunningham in Hook. Bot. Mag. 3228 (1833)	...	—	—	—	N.S.W.	Q.	—	B.f. III, 428 M.fr. IX, 184.
P. polioctemma, Benth., Fl. Austr. III, 427 (1866)	...	...	...	...	...	Q.	—	B.f. III, 425
P. Fitzalanii, Benth., Fl. Austr. III, 427 (1866)	...	...	...	...	...	Q.	—	B.f. III, 425
P. Carronis, Moore & F. v. M., fragm. VII, 49 (1869)	...	...	...	...	N.S.W.	—	—	M.fr. IX, 70, 185.
GEOPHILA, D. Don, prodr. fl. Nepal. 136 (1825). (Cephaelis partly.)								
G. reniformis, D. Don, prodr. fl. Nepal. 136 (1825)	...	—	—	—	—	Q.	—	M.fr. VII, 45; VIII, 147.
COPROSMA, R. et G. Forster, char. gen. 137, t. 69 (1776). (Nertera, Marquisia.)								
C. Nertera, F. v. M., fragm. IX, 186 (1875)	...	...	...	...	T. V. N.S.W.	—	—	M.fr. IX, 186.
C. reptans, F. v. M., fragm. IX, 186 (1875)	...	...	...	...	—	—	—	M.fr. IX, 186.
C. pumila, J. Hooker in Lond. Journ. VI, 465 (1847)	...	...	...	...	T. V.	—	—	B.f. III, 430 M.fr. IX, 186.
C. nitida, J. Hooker in Lond. Journ. VI, 465 (1847)	...	...	...	...	T. V.	—	—	B.f. III, 430 M.fr. IX, 186.
C. Billardieri, J. Hooker in Lond. Journ. VI, 465 (1847)	...	...	...	...	T. V. N.S.W.	—	—	B.f. III, 430
C. hirtella, Labillardière, Nov. Holl. pl. spec. I, 70, t. 95 (1804)	...	...	S.A.	T. V. N.S.W.	—	—	—	B.f. III, 429 M.fr. IX, 186.
C. Baueri, Endlicher, iconogr. gen. pl. XI, t. 111 (1835)	...	...	—	—	N.S.W.	—	—	M.fr. IX, 69.
C. putida, C. Moore & F. v. M., fragm. VII, 45 (1869)	...	...	—	—	N.S.W.	—	—	M.fr. IX, 69.
C. lanceolaris, F. v. M., fragm. IX, 70 (1875)	...	...	—	—	N.S.W.	—	—	M.fr. IX, 70.
OPERCULARIA, J. Gaertner, de fruct. I, 111, t. 24 (1758).								
O. vaginata, Labillardière, Nov. Holl. pl. spec. I, 34, t. 46 (1804)	...	W.A.	—	—	—	—	—	B.f. III, 433
O. spermacoea, Labillard., Nov. Holl. pl. spec. I, 35, t. 47 (1804)	...	W.A.	—	—	—	—	—	B.f. III, 433
O. scabrida, Schlechtendal, Linnaea XX, 604 (1847)	...	...	S.A.	—	V. N.S.W.	—	—	B.f. III, 433
O. hirsuta, F. v. M. in Benth. Fl. Austr. III, 434 (1866)	...	W.A.	—	—	—	—	—	B.f. III, 434
O. aspera, Gaertner, de fruct. I, 112, t. 24 (1758)	...	...	—	—	V. N.S.W.	Q.	—	B.f. III, 434
O. hispida, Sprengel, syst. veg. I, 355 (1825)	...	...	—	—	V. N.S.W.	—	—	B.f. III, 434
O. diphylla, Gaertner, de fruct. I, 113 (1758)	...	...	—	—	N.S.W.	Q.	—	B.f. III, 434
O. ovata, J. Hooker in Lond. Journ. VI, 465 (1847)	...	...	S.A.	T. V. N.S.W.	—	—	—	B.f. III, 435
O. varia, J. Hooker in Lond. Journ. VI, 466 (1847)	...	...	S.A.	T. V. N.S.W.	—	—	—	B.f. III, 435
O. rubioides, A. L. de Jussieu in Annal. du Mus. IV, 428 (1804)	...	W.A.	—	—	—	—	—	B.f. III, 435
O. volubilis, R. Brown in Benth. Fl. Austr. III, 435 (1866)	...	W.A.	—	—	—	—	—	B.f. III, 435
O. hispidula, Endlicher in Hueg. enum. 58 (1837)	...	W.A.	—	—	—	—	—	B.f. III, 436
O. echinocephala, Benth., Fl. Austr. III, 436 (1866)	...	W.A.	—	—	—	—	—	B.f. III, 436
O. apiciflora, Labillardière, Nov. Holl. pl. spec. I, 35, t. 48 (1804)	...	W.A.	—	—	—	—	—	B.f. III, 436

POMAX, Solander in Gaertn., de fruct. I, 112 (1788).									
P. umbellata, Solander in Gaertn., de fruct. I, 112 (1788)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III,437 M.fr.IX,187.
ELEUTHRANTHES, F. v. M., fragm. IV, 92 (1864).									
E. opercularina, F. v. M., fragm. IV, 92 (1864)	...	...	W.A.	—	—	—	—	—	B.f.III,437 M.fr.IV,92.
KNOXIA, Linné, Fl. Zeyl. 189 (1747).									
K. corymbosa, Willdenow, spec. pl. I, 582 (1798)	..	...	—	—	—	—	—	Q.	N.A. B.f.III,438 M.fr.IX,187.
SPERMACEAE, Dillenius, hort. Elth. 369, t. 277 (1732).									
S. brachystoma, R. Brown in Benth. Fl. Austr. III, 439 (1866)	—	—	—	—	—	—	Q.	N.A.	B.f.III,439 M.fr.IV,41.
S. pogostoma, Benth. Fl. Austr. III, 440 (1866)	...	...	—	—	—	—	—	N.A.	B.f.III,440
S. leptoloba, Benth. Fl. Austr. III, 440 (1866)	...	...	—	—	—	—	—	N.A.	B.f.III,440
S. marginata, Benth. Fl. Austr. III, 440 (1866)	...	...	—	—	—	—	—	Q.	N.A. B.f.III,440
S. multicaulis, Benth. Fl. Austr. III, 440 (1866)	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.III,440
S. exserta, Benth. Fl. Austr. III, 441 (1866)	...	...	—	—	—	—	—	N.A.	B.f.III,441
S. membranacea, R. Brown in Benth. Fl. Austr. III, 441 (1866)	...	...	—	—	—	—	—	N.A.	B.f.III,441
S. debilis, Benth. Fl. Austr. III, 441 (1866)	...	...	—	—	—	—	—	Q.	— B.f.III,441
S. inaperta, F. v. M., fragm. IV, 43 (1863)	...	...	—	—	—	—	—	—	N.A. B.f.III,441 M.fr.IV,43.
S. stenophylla, F. v. M., fragm. IV, 43 (1863)	...	...	—	—	—	—	—	—	N.A. B.f.III,442 M.fr.IV,43.
S. laevigata, F. v. M., fragm. IV, 41 (1863)	...	...	—	—	—	—	—	—	N.A. B.f.III,442 M.fr.IV,41.
S. breviflora, F. v. M. in Benth. Fl. Austr. III, 442 (1866)	...	...	—	—	—	—	—	—	N.A. B.f.III,442
S. auriculata, F. v. M., fragm. IV, 42 (1863)	...	...	—	—	—	—	—	—	N.A. B.f.III,442 M.fr.IV,42.
S. suffruticosa, R. Brown in Benth. Fl. Austr. III, 443 (1866)	...	...	—	—	—	—	—	—	N.A. B.f.III,443
ASPERULA, Dodoens, pemptad. 355 (1583).									
A. geminifolia, F. v. M., fragm. V, 147 (1865)	...	...	—	S.A.	—	V.	N.S.W.	Q.	— B.f.III,443 M.fr.IX,188;XI,27.
A. oligantha, F. v. M. in Neerl. Kruidk. Arch. IV, 111 et 112 (1859)	—	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,444 M.fr.IX,187;XI,27.
GALIUM, Dodoens, pemptad. 335 (1583), from Dioscorides.									
G. umbrosum, Solander in G. Forst. prodr. 89 (1786)	...	...	—	S.A.	T.	V.	N.S.W.	—	— B.f.III,446 M.fr.IX,188.
G. australe, De Candolle, prodr. IV, 608 (1830)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	— B.f.III,446

CAPRIFOLIACEAE.

Adanson, Familles des plantes II, 133 (1763).

SAMBUCUS, Tournefort, inst. 606, t. 396 (1700), from Dodoens (1583). (Tripetelus.)									
S. xanthocarpa, F. v. M. in Transact. phil. Inst. Vict. I, 42 (1855)	—	—	—	—	V.	N.S.W.	Q.	—	B.f.III,398
S. Gaudichaudiana, De Candolle, prodr. IV, 322 (1830)	...	...	—	S.A.	T.	V.	N.S.W.	Q.	— B.f.III,398

PASSIFLOREAE.

A. L. de Jussieu in Ann. du Mus. VI, 102 (1805).

PASSIFLORA, Plukenet, phytograph. 202 et 282, t. 104, 210, 211, 212 (1692). (Disemma.)									
P. Herbertiana, Lindley, Bot. Regist. t. 737 (1823)	...	...	—	—	—	N.S.W.	Q.	—	B.f.III,311 M.fr.IX,68.
P. cinnabarina, Lindley in Gard. Chron. 724 (1855)	...	...	—	—	V.	N.S.W.	—	—	— M.fr.IX,68.
P. aurantia, G. Forster, fl. ins. Austr. prodr. 62 (1786)	...	...	—	—	—	N.S.W.	Q.	—	B.f.III,312 M.fr.IX,68.
P. brachystephanea, F. v. M. in Benth. Fl. Austr. III, 312 (1866)	...	...	—	—	—	—	Q.	—	B.f.III,312 M.fr.I.56.
MODECCA, Lamarck, Encycl. méthod. IV, 283 (1797).									
M. australis, R. Brown in De Cand. prodr. III, 337 (1828)	...	...	—	—	—	—	Q.	N.A.	B.f.III,313 M.fr.IX,69.

CUCURBITACEAE.

Haller, enum. stirp. Helv. praef. 34 (1742).

TRICHOSANTHES, Linné, gen. plant. 295 (1737).									
T. pentaphylla, F. v. M. in Benth. Fl. Austr. III, 314 (1866)	...	...	—	—	—	—	Q.	—	B.f.III,314
T. cucumerina, Linné, spec. plant. 1008 (1753)	...	...	—	—	—	—	—	N.A.	B.f.III,314
T. palmata, Roxburgh, Fl. Ind. III, 704 (1832)	...	...	—	—	—	N.S.W.	Q.	—	B.f.III,315
T. Hearnii, F. v. M. in Benth. Fl. Austr. III, 315 (1866)	...	...	—	—	—	—	Q.	—	B.f.III,315
LAGENARIA, Seringe in Mém. Soc. Genév. III, 25, t. 2 (1825).									
L. vulgaris, Seringe in De Cand. prodr. III, 279 (1828)	...	...	—	—	—	—	Q.	—	B.f.III,316 M.fr.VII,62.
LUFFA, Tournefort in act. Acad. Paris. 84, t. 2 (1706), from Vesling (1638).									
L. Aegyptiaca, Miller, Gardener's Dictionary (1731)	...	...	—	—	—	—	Q.	—	B.f.III,316 M.fr.III,107.
L. graveolens, Roxburgh, Fl. Ind. III, 716 (1832)	...	...	—	—	—	—	—	N.A.	B.f.III,316 M.fr.III,106;IV,175.
L. foetida, Cavanilles, icon. I, 7, t. 9—10 (1791)	...	...	—	—	—	—	—	N.A.	— M.fr.XI,138.
ZANONIA, Linné, coroll. 19 (1737). (Alsomitra.)									
Z. Capricornica, F. v. M., fragm. VII, 61 (1870)	...	...	—	—	—	—	Q.	—	— M.fr.VII,61.
Z. Hookeri, F. v. M., fragm. VI, 188 (1868)	...	...	—	—	—	—	Q.	—	— M.fr.V,181;VII,62.
Z. Stephensiana, F. v. M., fragm. VIII, 181 (1874)	...	...	—	—	—	—	Q.	—	— M.fr.VIII,181.
CUCUMIS, Tournefort, inst. 104, t. 31 (1700), from Plinius.									
C. trigonus, Roxburgh, Fl. Ind. III, 722 (1832)	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.III,317 M.fr.VI,187.
C. Chate, Linné, syst. nat. edit. decim. 5 (1759)	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.
BENINCASA, Savi in Bibl. Ital. IX, 158 (1818).									
B. vacua; Cucurbita vacua, F. v. M., fragm. VI, 186 (1868)	...	...	—	—	—	—	Q.	—	— M.fr.VI,186.
MOMORDICA, Tournefort, institut. rei herbar. 103, t. 29 et 30 (1700).									
M. Balsamina, Linné, sp. pl. 1009 (1753)	...	...	—	—	—	N.S.W.	Q.	—	B.f.III,318 M.fr.VI,187.
M. Charantia, Linné, sp. pl. 1009 (1753)	...	...	—	—	—	—	Q.	—	—
BRYONOPSIS, Arnott in Hook. Journ. of Bot. III, 274 (1841). (Bryonia partly.)									
B. laciniata, Naudin in Annal. des sc. nat. sér. cinq. VI, 30 (1869)	—	—	—	—	—	N.S.W.	Q.	N.A.	B.f.III,319
B. Pancheri, Naudin in Annal. des sc. nat. sér. cinq. V, 30 (1869)	—	—	—	—	—	N.S.W.	—	—	—

MELOTHRIA, Linné, coroll. 1 (1737). (Zehneria, Mukia.)

M. Cunninghamii, Benth. Fl. Austr. III, 320 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.III, 320	M.fr. VI, 188; VII, 62.
M. Baueriana, F. v. M., fragm. VI, 188 (1868) ...	—	—	—	—	—	—	—	—	M.fr. VI, 188.
M. Muellieri, Benth. Fl. Austr. III, 320 (1866) ...	—	—	—	—	S.A.	—	V.	N.S.W.	—
M. Maderaspata, Cogniaux in De Cand. mon. phan. III, 623 (1881) ...	W.A.	S.A.	—	—	—	—	—	N.S.W.	Q.
SICYOS, Linné, syst. nat. 9 (1735); Linné, gen. plant. 297 (1737).	—	—	—	—	—	—	—	—	—
S. angulata, Linné, spec. plant. 1013 (1753) ...	—	—	—	—	T.	V.	N.S.W.	Q.	—
								B.f.III, 322	M.fr. VI, 187; VII, 62.

COMPOSITAE.

Vaillant in act. Acad. Paris. 143 (1718).

CENTRATHERUM, Cassini in Bull. de la Soc. philom. (1817).

C. muticum, Lessing in Schlecht. Linnaea II, 320 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 460	
---	---	---	---	---	--------	----	---	--------------	--

PLEUROCARPÆA, Benth. Fl. Austr. III, 460 (1866).

P. denticulata, Benth. Fl. Austr. III, 460 (1866) ...	—	—	—	—	—	—	—	N.A.	B.f.III, 460
---	---	---	---	---	---	---	---	------	--------------

VERNONIA, Schreber, gen. plant. II, 541 (1791).

V. cinerea, Lessing in Schlecht. Linnaea II, 291 (1829) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.III, 459	
---	---	---	---	---	--------	----	------	--------------	--

V. vagans, De Candolle, prodr. V, 32 (1843) ...	—	—	—	—	—	—	—	Q.	—
									M.fr. VI, 234.

ELEPHANTOPUS, Vaillant in act. Acad. Par. 309 (1719).

E. scaber, Linné, spec. plant. 814 (1753) ...	—	—	—	—	—	—	—	Q.	N.A.
									B.f.III, 461

ADENOSTEMMA, R. et G. Forster, char. gen. 89, t. 45 (1776).

A. viscosum, R. & G. Forster, char. gen. 89, t. 45 (1776) ...	—	—	—	—	S.A.	—	V.	N.S.W.	Q.
									B.f.III, 462

AGERATUM, Linné, gen. plant. 247 (1737).

A. conyzoides, Linné, spec. plant. 839 (1753) ...	—	—	—	—	—	—	—	Q.	—
									B.f.III, 462

EUPATORIUM, Tournefort, inst. 455, t. 259 (1700), from C. Bauhin (1623).

E. cannabinum, Linné, spec. pl. 838 (1753) ...	—	—	—	—	—	—	—	N.S.W.	Q.
									B.f.III, 462
									M.fr. V, 62.

LAGENOPHORA, Cassini in Bull. de la Soc. philom. 34 (1818). (Ixauchenus, Solenogyne, Emphysopus.)

L. Billardieri, Cassini in Diction. XXV, 111 (1822) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III, 507	
---	------	------	----	----	--------	----	---	--------------	--

L. Huegelii, Benth. in Hueg. enum. 59 (1837) ...	W.A.	S.A.	T.	V.	—	—	—	B.f.III, 507	
--	------	------	----	----	---	---	---	--------------	--

L. Solenogyne, F. v. M., fragm. V, 62 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 508	M.fr. V, 62.
--	---	---	---	---	--------	----	---	--------------	--------------

L. Emphysopus, J. Hooker, Fl. Tasman. I, 189 (1860) ...	—	—	—	—	T.	V.	N.S.W.	—	B.f.III, 508
---	---	---	---	---	----	----	--------	---	--------------

BRACHYCOME, Cassini in Bull. de la Soc. philom. 199 (1816). (Bellidis subgenus, Pacquerina, Brachystephium, Steiroglossa, Silphiospermum.)

B. diversifolia, Fischer & Meyer, ind. sem. hort. Petrop. 31 (1835) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 511	
---	---	------	----	----	--------	---	---	--------------	--

B. segmentosa, Moore & F. v. M., fragm. VIII, 144 (1874) ...	—	—	—	—	N.S.W.	—	—	—	M.fr. V, 144.
--	---	---	---	---	--------	---	---	---	---------------

B. melanocarpa, Sonder & F. v. M. in Schl. Linn. XXV, 476 (1852) ...	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 511	
--	---	------	---	---	----	--------	---	--------------	--

B. radicans, Steetz in Lehm. pl. Preiss. I, 429 (1845) ...	—	—	—	—	T.	V.	N.S.W.	—	B.f.III, 512
--	---	---	---	---	----	----	--------	---	--------------

B. goniocarpa, Sonder & F. v. M. in Schl. Linn. XXV, 474 (1852) ...	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 512	
---	------	------	---	---	----	--------	---	--------------	--

B. pachyptera, Turczaninow in Bull. de Mosc. XXIV, 175 (1851) ...	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 512	
---	------	------	---	---	----	--------	---	--------------	--

B. iberidifolia, Benth. in Hueg. enum. 59 (1837) ...	W.A.	—	—	—	—	—	—	B.f.III, 512	M.fr. XI, 123.
--	------	---	---	---	---	---	---	--------------	----------------

B. pusilla, Steetz in Lehm. pl. Preiss. I, 427 (1845) ...	W.A.	—	—	—	—	—	—	B.f.III, 513	
---	------	---	---	---	---	---	---	--------------	--

B. microcarpa, F. v. M., fragm. I, 50 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 513	M.fr. I, 50.
--	---	---	---	---	--------	----	---	--------------	--------------

B. Stuartii, Benth. Fl. Austr. III, 513 (1866) ...	—	—	—	—	N.S.W.	—	—	B.f.III, 513	
--	---	---	---	---	--------	---	---	--------------	--

B. scapigera, De Candolle, prodr. VII, 277 (1838) ...	—	—	—	—	V.	N.S.W.	—	B.f.III, 513	
---	---	---	---	---	----	--------	---	--------------	--

B. Muellieri, Sonder in Schlecht. Linnaea XXV, 475 (1852) ...	—	S.A.	—	—	—	—	—	B.f.III, 514	
---	---	------	---	---	---	---	---	--------------	--

B. graminea, F. v. M., fragm. I, 49 (1858) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 514	M.fr. VIII, 142.
--	---	------	----	----	--------	---	---	--------------	------------------

B. angustifolia, Cunningham in De Cand., prodr. V, 306 (1836) ...	—	—	—	—	T.	V.	N.S.W.	—	M.fr. I, 49.
---	---	---	---	---	----	----	--------	---	--------------

B. linearifolia, De Candolle, prodr. V, 306 (1836) ...	—	—	—	—	N.S.W.	—	—	B.f.III, 515	
--	---	---	---	---	--------	---	---	--------------	--

B. basaltica, F. v. M., fragm. I, 50 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.III, 515	M.fr. I, 50.
---	---	------	---	---	--------	----	---	--------------	--------------

B. trachycarpa, F. v. M. in Schlecht. Linnaea XXV, 339 (1852) ...	—	S.A.	—	—	V.	N.S.W.	Q.	B.f.III, 515	
---	---	------	---	---	----	--------	----	--------------	--

B. exilis, Sonder in Linnaea XXV, 473 (1852) ...	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 516	
--	---	------	---	---	----	--------	---	--------------	--

B. ptychocarpa, F. v. M. in Transact. phil. Soc. Vict. I, 43 (1854) ...	—	—	—	—	V.	N.S.W.	—	B.f.III, 516	
---	---	---	---	---	----	--------	---	--------------	--

B. debilis, Sonder in Linnaea XXV, 477 (1852) ...	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 516	
---	---	------	---	---	----	--------	---	--------------	--

B. decipiens, J. Hooker in Lond. Journ. VI, 114 (1847) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 516	
--	---	------	----	----	--------	---	---	--------------	--

B. cardiocarpa, F. v. M. in Benth. Fl. Austr. III, 516 (1866) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 517	
---	---	------	----	----	--------	---	---	--------------	--

B. nivalis, F. v. M. in Transact. phil. Soc. Vict. I, 43 (1854) ...	—	—	—	—	V.	N.S.W.	—	B.f.III, 517	
---	---	---	---	---	----	--------	---	--------------	--

B. scapiformis, De Candolle, prodr. V, 306 (1836) ...	—	—	—	—	T.	V.	N.S.W.	—	B.f.III, 517
---	---	---	---	---	----	----	--------	---	--------------

B. stricta, De Candolle, prodr. V, 305 (1836) ...	—	—	—	—	T.	V.	N.S.W.	—	B.f.III, 518
---	---	---	---	---	----	----	--------	---	--------------

B. heterodonta, De Candolle, prodr. V, 305 (1836) ...	—	—	—	—	N.S.W.	—	—	B.f.III, 518	
---	---	---	---	---	--------	---	---	--------------	--

B. Billardieri, Benth. Fl. Austr. III, 518 (1866) ...	W.A.	—	—	—	—	—	—	B.f.III, 518	
---	------	---	---	---	---	---	---	--------------	--

B. cheilocarpa, F. v. M. in Wing's South. Sc. Record, Aug. (1882) ...	W.A.	—	—	—	—	—	—	—	
---	------	---	---	---	---	---	---	---	--

B. ciliaris, Lessing, syn. comp. 172 (1832) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 518	
---	------	------	----	----	--------	---	---	--------------	--

B. calocarpa, F. v. M. in Schlecht. Linnaea XXV, 399 (1852) ...	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 519	M.fr. XI, 27.
---	---	------	---	---	----	--------	---	--------------	---------------

B. chrysoglossa, F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854) ...	—	S.A.	—	—	V.	N.S.W.	—	—	
--	---	------	---	---	----	--------	---	---	--

B. marginata, Benth. in Hueg. enum. 60 (1837) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III, 519	
---	---	---	---	---	--------	----	---	--------------	--

B. Sieberi, De Candolle, prodr. V, 306 (1836) ...	—	—	—	—	N.S.W.	—	—	B.f.III, 520	
---	---	---	---	---	--------	---	---	--------------	--

B. discolor, C. Stuart in Benth. Fl. Austr. III, 520 (1866) ...	—	—	—	—	N.S.W.	—	—	B.f.III, 520	
---	---	---	---	---	--------	---	---	--------------	--

B. multifida, De Candolle, prodr. V, 306 (1836) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.III, 520	M.fr. VIII, 145.
---	---	---	---	---	----	--------	----	--------------	------------------

B. latisquamata, F. v. M., fragm. XI, 16 (1878) ...	W.A.	—	—	—	—	—	—	—	M.fr. XI, 16.
---	------	---	---	---	---	---	---	---	---------------

B. glandulosa, Benth. Fl. Austr. III, 520 (1866) ...	W.A.	—	—	—	—	—	—	B.f.III, 520	
--	------	---	---	---	---	---	---	--------------	--

B. collina, Benth. Fl. Austr. III, 521 (1866) ...	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 521	
---	------	------	---	---	----	--------	---	--------------	--

ERODIOPHYLLUM, F. v. M., fragm. IX, 119 (1875).

E. Elderi, F. v. M., fragm. IX, 120 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr. IX, 119.
--	---	------	---	---	---	---	---	---	----------------

MINURIA, De Candolle, prodr. V, 298 (1836). (Therogeron, Elachothamnus, Kippistia.)

M. leptophylla, De Candolle, prodr. V, 298 (1836) ...	W.A.	S.A.	—	—	V.	N.S.W.	Q.	—	M.fr. IX, 119; X, 56.
---	------	------	---	---	----	--------	----	---	-----------------------

M. Cunninghamii, Benth. Fl. Austr. III, 498 (1866) ...	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 498	
--	---	------	---	---	----	--------	---	--------------	--

M. Candollei, F. v. M., fragm. IX, 119 (1875)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.III, 499	M.fr. X, 56.
M. suaedifolia, F. v. M. in Benth. Fl. Austr. III, 499 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 499	M.fr. X, 56.
CALOTIS, R. Brown in Bot. Regist. t. 504 (1820). (Huenefeldia, Goniopogon, Cheiroloma.)									
C. dentex, R. Brown in Bot. Regist. 504 (1820)	—	—	—	—	N.S.W.	Q.	—	B.f.III, 501	
C. cuneifolia, R. Brown in Bot. Regist. t. 504 (1820)	—	S.A.	—	—	V.	N.S.W.	Q.	B.f.III, 501	
C. glandulosa, F. v. M. in Transact. Vict. Inst. 129 (1855)	—	—	—	—	V.	N.S.W.	—	B.f.III, 502	
C. cymbacantha, F. v. M. in Schlecht. Linn. XXV, 400 (1852)	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 502	
C. erinacea, Steetz in Lehm. pl. Preiss. I, 424 (1845)	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 502	
C. scabiosifolia, Sonder & F. v. M. in Linn. XXV, 471 (1852)	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 503	M.fr. XI, 27.
C. scapigera, Hooker in Mitch. Trop. Austr. 75 (1848)	—	S.A.	—	—	V.	N.S.W.	Q.	N.A.	M.fr. XI, 27.
C. anthemoides, F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854)	—	—	—	—	V.	—	—	B.f.III, 504	
C. lappulacea, Benth. in Hueg. enum. 60 (1837)	W.A.	S.A.	—	—	V.	N.S.W.	Q.	B.f.III, 504	
C. microcephala, Benth. Fl. Austr. III, 504 (1866)	—	S.A.	—	—	V.	N.S.W.	—	B.f.III, 504	
C. brevifolia, Benth. in Hueg. enum. 60 (1837)	—	—	—	—	—	—	N.A.	B.f.III, 505	
C. plumulifera, F. v. M. in Transact. phil. Inst. Vict. III, 57 (1859)	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 505	
C. porphyroglossa, F. v. M. in Benth. Fl. Austr. III, 505 (1866)	—	S.A.	—	—	—	N.S.W.	—	N.A.	B.f.III, 505
C. pterosperma, E. Brown in Benth. Fl. Austr. III, 505 (1866)	—	—	—	—	—	—	N.A.	B.f.III, 505	
C. hispidula, F. v. M. in Transact. Vict. Inst. 130 (1855)	W.A.	S.A.	—	—	V.	N.S.W.	—	B.f.III, 506	
C. Kempei, F. v. M. in Transact. Roy. Soc. S. Austr. IV, 112 (1881)	—	S.A.	—	—	—	—	—	—	M.fr. XI, 139.
ASTER, Tournefort, inst. 481, t. 274 (1700), from Dioscorides. (Olearia, Celmisia, Eurybia, Steetzia.)									
A. megalophyllus, F. v. M., fragm. V, 70 (1865)	—	—	—	—	V.	N.S.W.	—	B.f.III, 468	M.fr. V, 70.
A. chrysophyllus, Cunningham in De Cand. prodr. V, 266 (1836)	—	—	—	—	N.S.W.	—	—	B.f.III, 468	
A. alpicola, F. v. M., fragm. V, 70 (1865)	—	—	—	—	V.	N.S.W.	—	B.f.III, 468	M.fr. V, 70.
A. rosmarinifolius, Cunningham in De Cand. prodr. V, 268 (1836)	—	—	—	—	N.S.W.	—	—	B.f.III, 468	
A. viscosus, Labillardiere, Nov. Holl. pl. spec. II, 53, t. 203 (1806)	—	—	T.	V.	—	—	—	B.f.III, 468	M.fr. V, 71.
A. Sonderi, F. v. M., fragm. V, 83 (1865)	—	S.A.	—	—	—	—	—	B.f.III, 469	M.fr. V, 83.
A. pannosus, F. v. M., fragm. V, 83 (1865)	—	S.A.	—	—	V.	—	—	B.f.III, 469	M.fr. V, 83.
A. oliganthemus; Olearia, F. v. M. in Benth. Fl. Austr. III, 469 (1866)	—	—	—	—	N.S.W.	—	—	B.f.III, 469	
A. argophyllus, Labillard., Nov. Holl. pl. spec. II, 52, t. 201 (1806)	—	—	T.	V.	N.S.W.	—	—	B.f.III, 470	M.fr. V, 68.
A. myrsinifolius, Cunningham in De Cand. prodr. V, 267 (1836)	—	—	—	—	N.S.W.	—	—	B.f.III, 470	
A. cyrsinoides, Labillard., Nov. Holl. pl. spec. II, 53, t. 202 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 471	M.fr. V, 69.
A. personioides, Cunningham in De Cand. prodr. V, 267 (1836)	—	—	T.	—	—	—	—	B.f.III, 471	M.fr. V, 69.
A. obcordatus, F. v. M., fragm. V, 69 (1865)	—	—	T.	—	—	—	—	B.f.III, 471	M.fr. V, 69.
A. pinifolius, F. v. M., fragm. V, 71 (1865)	—	—	T.	—	—	—	—	B.f.III, 472	M.fr. V, 71.
A. ledifolius, Cunningham in De Cand. prodr. V, 269 (1836)	—	—	T.	—	—	—	—	B.f.III, 472	
A. dentatus, Andrews, Bot. Reposit. t. 61 (1799)	—	—	—	—	N.S.W.	—	—	B.f.III, 472	M.fr. V, 82.
A. stellulatus, Labillard., Nov. Holl. pl. spec. II, 50, t. 196 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III, 473	M.fr. V, 81.
A. asterotrichus, F. v. M., fragm. V, 79 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 473	M.fr. I, 111.
A. gravis, F. v. M., fragm. V, 82 (1865)	—	—	—	—	N.S.W.	—	—	B.f.III, 474	M.fr. V, 82.
A. Nernstii, F. v. M., fragm. V, 81 (1865)	—	—	—	—	N.S.W.	Q.	—	B.f.III, 474	M.fr. V, 81.
A. hygrophilus, Cunningham in De Cand. prodr. V, 269 (1836)	—	—	—	—	—	Q.	—	B.f.III, 474	
A. Siemssenii, F. v. M., fragm. V, 71 (1865)	—	—	—	—	N.S.W.	—	—	B.f.III, 475	M.fr. I, 50.
A. tubuliflorus, F. v. M., fragm. V, 65 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 475	M.fr. V, 65.
A. axillaris, F. v. M., fragm. V, 64 (1865)	W.A.	S.A.	T.	V.	N.S.W.	—	N.A.	B.f.III, 475	M.fr. V, 64.
A. revolutus; Olearia, F. v. M. in Benth. Fl. Austr. III, 476 (1866)	W.A.	—	—	—	—	—	—	B.f.III, 476	
A. exilifolius, F. v. M., fragm. V, 69 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 476	M.fr. V, 69.
A. ramulosus, Labillard., Nov. Holl. plant. spec. II, 51, t. 198 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 476	
A. florulentus, F. v. M., fragm. V, 82 (1865)	—	—	T.	V.	N.S.W.	—	—	B.f.III, 477	M.fr. V, 82.
A. microphyllus, Persoon, synops. pl. II, 442 (1807)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 478	M.fr. V, 70.
A. Mitchellii, F. v. M., fragm. V, 78 (1865)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III, 478	M.fr. V, 78.
A. exiguifolius, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 478	M.fr. V, 67.
A. Cassiniae, F. v. M., fragm. V, 68 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 479	M.fr. V, 68.
A. cyanodiscus, F. v. M., fragm. V, 82 (1865)	—	—	—	—	N.S.W.	—	—	B.f.III, 479	M.fr. V, 82.
A. pimeleoides, Cunningham in De Cand. prodr. V, 268 (1836)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 479	
A. iodochromus, F. v. M., fragm. V, 81 (1865)	—	—	—	—	V.	N.S.W.	—	B.f.III, 480	M.fr. II, 110.
A. adenolasius, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 480	M.fr. V, 67.
A. conocephalus, F. v. M., fragm. V, 79 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 480	M.fr. V, 79.
A. calcareus; Olearia, F. v. M. in Benth. Fl. Austr. III, 481 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 481	
A. magniflorus, F. v. M., fragm. V, 80 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 481	M.fr. V, 80.
A. Muelleri; Eurybia, Sonder in Schlecht. Linn. XXV, 459 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.III, 481	
A. Stuartii, F. v. M., fragm. V, 76 (1865)	—	S.A.	—	—	—	—	—	B.f.III, 481	M.fr. I, 202.
A. decurrens, Cunningham in De Cand. prodr. V, 269 (1836)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 481	
A. glutescens, F. v. M., fragm. V, 77 (1865)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 482	M.fr. V, 77.
A. orarius, F. v. M., fragm. V, 78 (1865)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 482	M.fr. V, 78.
A. vernicosus, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 482	M.fr. V, 67.
A. teretifolius, F. v. M., fragm. V, 77 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.f.III, 483	M.fr. V, 77.
A. Turczaninowii, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 483	M.fr. V, 67.
A. ellipticus, Cunningham in De Cand. prodr. V, 271 (1836)	—	—	—	—	N.S.W.	Q.	—	B.f.III, 483	M.fr. I, 16.
A. glandulosus, Labillard., Nov. Holl. pl. spec. II, 50, t. 197 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.III, 484	
A. heleophilus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 484	M.fr. V, 66.
A. muricatus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 484	M.fr. V, 66.
A. Benth. F. v. M.; Olearia stricta, Benth. Fl. Austr. III, 485 (1866)	—	—	—	—	V.	—	—	B.f.III, 485	
A. Steetzii, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 485	M.fr. V, 66.
A. Cunninghamii, F. v. M.; Eurybia, De Cand. prodr. V, 269 (1836)	—	—	—	—	N.S.W.	—	—	B.f.III, 486	
A. paucidentatus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 485	M.fr. V, 66.
A. megalodontus, F. v. M., fragm. VIII, 244 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr. VIII, 244.
A. adenophorus, F. v. M., fragm. V, 78 (1865)	—	—	—	—	V.	N.S.W.	—	B.f.III, 486	M.fr. I, 111.
A. homolepis, F. v. M., fragm. V, 65 (1865)	W.A.	—	—	—	—	—	—	B.f.III, 486	M.fr. V, 65.
A. xerophilus, F. v. M., fragm. V, 76 (1865)	—	—	—	—	—	Q.	—	B.f.III, 486	M.fr. I, 51, 76 & 86.
A. Ferresii, F. v. M., fragm. V, 75 (1865)	—	S.A.	—	—	—	—	N.A.	B.f.III, 487	M.fr. III, 18.

A. exul, Lindley, Bot. Regist. XXV, App. XXIV (1839) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.III,487	M.fr.V,75.
A. argutus, R. Brown in Benth. Fl. Austr. III, 488 (1866) ...	—	—	—	—	—	—	N.A.	B.f.III,488	
A. Huegelii, F. v. M., fragm. V, 79 (1865)... ..	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.III,488	M.fr.V,79.
A. Balli, F. v. M., fragm. VIII, 143 (1874)... ..	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,143.
A. Mooneyi, F. v. M., fragm. VIII, 144 (1874) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.
A. Celmisia, F. v. M., fragm. V, 84 (1865)... ..	—	—	—	T.	V.	N.S.W.	—	B.f.III,489	M.fr.V,84.
VITTADINIA, Ach. Richard, Voy. d'Astrol. bot. 250 (1832). (Eurybiopsis, Microgyne.)									
V. brachycomoides, F. v. M., fragm. V, 86 (1865) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,490
V. australis, A. Richard, Voy. d'Astrol. 251 (1832) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,490	M.fr.VI,86.
V. scabra, De Candolle, prodr. V, 281 (1836) ...	—	—	—	—	N.S.W.	Q.	—	B.f.III,491	M.fr.XI,20.
V. macrorrhiza, A. Gray in Proc. Am. Acad. V, 118 (1863) ...	—	—	—	—	—	—	N.A.	B.f.III,492	
PODOCOMA, Cassini in Bull. de la Soc. philom. 137 (1817). (Ixiochlamys.)									
P. cuneifolia, R. Brown, App. to Sturt, Centr. Austr. 17 (1849) ...	—	S.A.	—	—	N.S.W.	—	N.A.	B.f.III,493	
ERIGERON, Linné, hort. Clifford. 407 (1737). (Haplopappus partly.)									
E. pappochromus, Labillard., Nov. Holl. pl. spec. II, 47, t. 193 (1806) ...	—	—	—	T.	V.	N.S.W.	—	—	B.f.III,494
E. ambiguus, F. v. M. in Transact. phil. Inst. Vict. III, 53 (1858) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,494
E. sessilifolius, F. v. M., fragm. X, 100 (1881) ...	—	—	—	—	—	—	—	N.A.	—
E. minurioides, Benth., Fl. Austr. III, 495 (1866) ...	—	—	—	—	V.	—	—	—	B.f.III,495
E. conyzoides, F. v. M. in Transact. phil. Soc. Vict. I, 105 (1855) ...	—	—	—	—	V.	N.S.W.	—	—	B.f.III,495
CONYZA, Lamarck, Encycl. méthod. II, 87, 89 (1786).									
C. viscidula, Wallich, numer. list 3006 (1828) ...	—	—	—	—	—	N.S.W.	Q.	—	B.f.III,496
C. Aegyptiaca, Aiton, hort. Kew. III, 183 (1789)... ..	—	—	—	—	—	—	Q.	N.A.	B.f.III,497
BLUMEA, De Candolle in Guillemin, Arch. de Bot. II, 514 (1833).									
B. glandulosa, De Candolle in Wight, contrib. 14 (1834) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,525
B. amplexens, De Candolle in Wight, contrib. 13 (1834)... ..	—	—	—	—	—	—	Q.	—	B.f.III,525
B. integrifolia, De Candolle, prodr. V, 433 (1836)... ..	—	—	—	—	—	—	—	N.A.	B.f.III,525
B. diffusa, R. Brown in Benth. Fl. Austr. III, 525 (1866) ...	—	—	—	—	—	—	—	N.A.	B.f.III,525
B. hieracifolia, De Candolle in Wight, contrib. 15 (1834)... ..	—	—	—	—	—	—	Q.	N.A.	B.f.III,526
B. Cunninghamii, De Candolle, prodr. V, 435 (1836) ...	—	—	—	—	—	—	—	N.A.	B.f.III,526
B. acutata, De Cand. in Ann. du Mus. d'hist. nat. Par. III, 409 (1834) ...	—	—	—	—	—	—	Q.	—	M.fr.IX,160.
B. lacera, De Candolle in Wight, contrib. 14 (1834) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,526
PLUCHEA, Cassini in Bull. de la Soc. philom. 31 (1817). (Spiropodium, Eyrea.)									
P. Indica, Lessing in Schlecht. Linnaea VI, 150 (1831) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,527
P. tetranthera, F. v. M., Rep. Babb. Exped. 12 (1860) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,528
P. baccharoides, F. v. M. in Benth. Fl. Austr. III, 528 (1866) ...	—	—	—	—	—	—	Q.	—	B.f.III,528
P. Eyrea, F. v. M., Rep. Babb. Exped. 11, 12 (1860) ...	—	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.III,528
P. squarrosa, Benth., Fl. Austr. III, 529 (1866)... ..	—	W.A.	—	—	—	—	—	—	B.f.III,529
P. dentex, R. Brown in Benth. Fl. Austr. III, 529 (1866) ...	—	—	—	—	—	—	Q.	—	B.f.III,529
PTERIGERON, De Candolle, prodr. V, 293 (1836). (Streptoglossa, Oliganthemum.)									
P. decurrens, De Candolle, prodr. V, 293 (1836) ...	—	—	—	—	—	—	—	N.A.	B.f.III,531
P. liatroides, Benth., Fl. Austr. III, 532 (1866)... ..	—	W.A.	S.A.	—	—	—	—	—	B.f.III,532
P. odoratus, Benth., Fl. Austr. III, 532 (1866) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,532
P. macrocephalus, Benth., Fl. Austr. III, 532 (1866) ...	—	—	—	—	—	—	—	N.A.	B.f.III,532
P. microglossus, Benth., Fl. Austr. III, 532 (1866) ...	—	—	—	—	—	—	—	N.A.	B.f.III,532
P. dentatifolius, F. v. M., fragm. IX, 119 (1875) ...	—	—	S.A.	—	—	—	—	—	M.fr.X,119.
P. adscendens, Benth., Fl. Austr. III, 533 (1866) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,533
P. filifolius, Benth., Fl. Austr. III, 533 (1866) ...	—	—	—	—	—	—	—	N.A.	B.f.III,533
THESPIDIUM, F. v. M. in Journ. of Landsborough's Exped. app. (1862).									
T. basiflorum, F. v. M. in Journ. of Landsb. Exped. App. (1862) ...	—	—	—	—	—	—	—	N.A.	B.f.III,534
COLEOCOMA, F. v. M. in Hook. Kew Misc. IX, 19 (1857).									
C. Centaurea, F. v. M. in Hook. Kew Misc. IX, 19 (1857) ...	—	—	—	—	—	—	—	N.A.	B.f.III,533
EPALTES, Cassini in Bull. de la Soc. philom. 139 (1818). (Sphaeromorphaea partly, Ethuliopsis.)									
E. Cunninghamii, Benth., Fl. Austr. III, 530 (1866) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.III,530	
E. australis, Lessing in Linnaea V, 148 (1831) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.III,530	M.fr.XI,101.
E. pleiochaeta, F. v. M., fragm. X, 100 (1877) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.X,100.
E. Harrisii, F. v. M., fragm. XI, 101 (1881) ...	—	—	—	—	—	—	Q.	—	M.fr.XI,101.
SPHAERANTHOS, Vaillant in act. Acad. Par. 289 (1719). (Sphaeranthus.)									
S. hirtus, Willdenow, spec. plant. III, 2395 (1800) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,521
S. microcephalus, Willdenow, spec. plant. III, 2395 (1800) ...	—	—	—	—	—	—	—	N.A.	B.f.III,522
PTEROCAULON, Elliot, Sketch of the Bot. of S. Carolina and Georgia II, 323 (1824). (Monenteles.)									
P. verbascifolius, Benth. & J. Hooker, gen. pl. II, 294 (1873) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,523
P. Billardieri, F. v. M., Papuan plants, 43 (1876)... ..	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.III,523	M.fr.XI,101.
P. sphacelatus, Benth. & J. Hooker, gen. plant. II, 94 (1873) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.III,523	
P. glandulosus, Benth. & J. Hooker, gen. plant. II, 94 (1873) ...	—	—	—	—	—	—	Q.	N.A.	B.f.III,523
P. sphaeranthoides, Benth. & J. Hooker, gen. plant. II, 94 (1873) ...	—	—	—	—	—	—	—	N.A.	B.f.III,524
STUARTINA, Sonder in Schlechtendal's Linnaea XXV, 521 (1852).									
S. Muellieri, Sonder in Schlecht. Linnaea XXV, 522 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.III,657	M.fr.VIII,145;XI,27.
GNAPHALIUM, Linné, gen. plant. 250 (1737), from J. & C. Bauhin (1619). (Euchiton.)									
G. luteo-album, Linné, spec. plant. 851 (1753) ...	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.III,653
G. Japonicum, Thunberg, Fl. Japon. 311 (1784) ...	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III,653
G. alpinum, F. v. M., second gen. Report 12 (1854) ...	—	—	—	T.	V.	—	—	—	B.f.III,654
G. purpureum, Linné, spec. plant. 854 (1753) ...	—	—	—	—	—	N.S.W.	Q.	—	B.f.III,655
G. indicum, Linné, spec. plant. 852 (1753)... ..	—	—	—	—	—	N.S.W.	Q.	N.A.	B.f.III,655
G. indutum, J. Hooker in Lond. Journ. of Bot. VI, 121 (1847) ...	—	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.III,655
G. Traversii, J. Hooker, Handb. N. Zeal. Fl. 154 ...	—	—	—	—	V.	N.S.W.	—	—	B.f.III,656

ANTENNARIA, Gaertner, de fruct. II, 410, t. 167 (1791). (Raoulia partly.)									
A. Planchoni, F. v. M.; Gnaphalium, Hook. Fl. Tasm. I, 217, t. 62 (1860)	—	—	—	—	—	—	—	—	—
A. uniceps, F. v. M. in Transact. phil. Soc. Vict. t. 105 (1854)...	—	—	—	V.	N.S.W.	—	—	B.f.III, 652	M.fr. VIII, 145.
LEONTOPODIUM, R. Brown in Transact. Linn. Soc. XII, 124 (1817). (Raoulia partly.)									
L. Catipes, F. v. M. in Papers of the Roy. Soc. of Tasm. 44 (1882)	—	—	—	T.	V.	N.S.W.	—	B.f.III, 651	—
L. Mereditiae; Antennaria, F. v. M. in Pap. Roy. Soc. of Tasm. 15 (1870)	—	—	—	T.	—	—	—	—	—
PTERYGOPAPPUS, J. Hooker in Lond. Journ. of Bot. VII, 120 (1847).									
P. Lawrencii, J. Hooker in Lond. Journ. VII, 120 (1847)	...	—	—	T.	—	—	—	B.f.III, 656	—
PODOTHECA, Cassini in Diction. XXIII, 561 (1822). (Podospermum, Phaenopoda, Lophoclinium.)									
P. gnaphalioides, Graham in Bot. Mag. t. 3920 (1842)	...	...	...	W.A.	—	—	—	B.f.III, 601	—
P. angustifolia, Lessing, synops. gen. compos. 273 (1832)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—
P. pygmaea, A. Gray in Hook. Kew Misc. IV, 227 (1852)	...	...	...	W.A.	—	—	—	B.f.III, 602	M.fr. XI, 20.
P. chrysanthia, Benth. Fl. Austr. III, 602 (1866)	...	...	...	W.A.	—	—	—	B.f.III, 602	—
P. fuscescens, Benth. Fl. Austr. III, 602 (1866)	...	...	...	W.A.	—	—	—	B.f.III, 602	—
IXIOLAENA, Benth. in Hueg. enum. 66 (1837).									
I. brevicompta, F. v. M., fragm. I, 53 (1858)	...	...	...	—	—	—	—	N.S.W.	Q.
I. leptolepis, Benth. Fl. Austr. III, 597 (1866)...	...	...	...	—	S.A.	—	V.	N.S.W.	—
I. supina, F. v. M. in Transact. Vict. Inst. 37 (1855)	...	...	...	—	S.A.	T.	—	—	—
I. tomentosa, Sonder & F. v. M. in Linnaea XXV, 504 (1852)	...	...	...	—	S.A.	—	V.	N.S.W.	—
I. viscosa, Benth. in Hueg. enum. 66 (1837)	...	...	...	W.A.	—	—	—	—	—
PODOLEPIS, Labillardière, Nov. Holl. pl. specim. II, 57, t. 28 (1806). (Scalia, Panaetia, Scaliopsis, Siemssenia, Stylolepis, Rutidochlamys.)									
P. rhytidochlamys, F. v. M., fragm. IV, 79 (1864)	...	...	...	—	—	—	—	N.S.W.	Q.
P. longipedata, Cunningham in De Cand. prodr. VI, 163 (1837)...	...	...	...	—	—	—	—	V.	N.S.W.
P. acuminata, R. Brown in Ait. hort. Kew. ed. sec. V, 82 (1813)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.
P. canescens, Cunningham in De Cand. prodr. VI, 163 (1837)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.
P. aristata, Benth. in Hueg. enum. 64 (1837)	...	...	...	W.A.	—	—	—	—	—
P. pallida, Turczaninow in Bull. de Mosc. XXIV, 78 (1851)	...	...	...	W.A.	—	—	—	—	N.A.
P. nutans, Steetz in Lehm. pl. Preiss. I, 464 (1845)	...	...	...	W.A.	—	—	—	—	—
P. gracilis, Graham in Edinb. New phil. journ. V, 379 (1828)	...	...	...	W.A.	—	—	—	—	—
P. rugata, Labillardière, Nov. Holl. pl. spec. II, 57, t. 208 (1806)	...	...	...	W.A.	S.A.	—	V.	—	—
P. Lessoni, Benth. Fl. Austr. III, 606 (1866)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
P. Siemssenia, F. v. M. in Benth. Fl. Austr. III, 607 (1866)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
P. microcephala, Benth. Fl. Austr. III, 607 (1866)...	...	...	...	W.A.	—	—	—	—	—
ATHRIXIA, Ker in Bot. Regist. VIII, 681 (1823). (Asteridea, Chrysodiscus, Trichostegia.)									
A. australis, Steetz in Lehm. pl. Preiss. I, 482 (1845)	...	...	...	W.A.	—	—	—	—	—
A. gracilis, Benth. Fl. Austr. III, 599 (1866)	...	...	...	W.A.	—	—	—	—	—
A. multiceps, Benth. Fl. Austr. III, 599 (1866)	...	...	...	W.A.	—	—	—	—	—
A. stricta, Benth. Fl. Austr. III, 600 (1866)	...	...	...	W.A.	—	—	—	—	—
A. tenella, Benth. Fl. Austr. III, 600 (1866)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
A. chaetopoda, F. v. M., fragm. X, 56 (1876)	...	...	...	W.A.	—	—	—	—	—
LEPTORRHYNCHOS, Lessing, syn. composit. 273 (1832). (Rhytidanthe.)									
L. squamatus, Lessing, syn. comp. 273 (1832)	...	...	...	—	S.A.	T.	V.	N.S.W.	—
L. panaetioides, Benth. Fl. Austr. III, 609 (1866)	...	...	...	—	S.A.	—	V.	N.S.W.	—
L. tenuifolius, F. v. M., fragm. I, 52 (1858)	...	...	...	—	S.A.	—	V.	—	—
L. pulchellus, F. v. M. in Linnaea XXV, 500 (1852)	...	...	...	—	S.A.	—	V.	N.S.W.	—
L. Baileyi, F. v. M., fragm. X, 101 (1877)	...	...	...	—	—	—	—	—	Q.
L. elongatus, De Candolle, prodr. VI, 160 (1837)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—
L. medius, Cunningham in De Cand. prodr. VI, 160 (1837)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
L. Waitzia, Sonder in Schlecht. Linnaea XXV, 501 (1852)	...	...	...	—	S.A.	—	V.	N.S.W.	—
L. nitidulus, De Candolle, prodr. VI, 160 (1837)	...	...	...	—	—	—	T.	V.	N.S.W.
WAITZIA, Wendland, Collect. pl. II, 13, t. 42 (1808). (Viraya, Morna, Pterochaete.)									
W. corymbosa, Wendland, Coll. pl. II, 13, t. 42 (1808)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
W. aurea, Steetz in Lehm. pl. Preiss. I, 452 (1845)	...	...	...	W.A.	—	—	—	—	—
W. nivea, Benth. Fl. Austr. III, 636 (1866)	...	...	...	W.A.	—	—	—	—	—
W. Steetziana, Lehmann, pl. Preiss. I, 454 (1845)	...	...	...	W.A.	—	—	—	—	—
W. podolepis, Steetz in Lehm. pl. Preiss. I, 450 (1845)	...	...	...	W.A.	—	—	—	—	—
W. paniculata, F. v. M. in Benth. Fl. Austr. III, 637 (1866)	...	...	...	W.A.	—	—	—	—	—
HELIPTERUM, De Candolle, prodr. VI, 211 (1837). (Argyrocome, 1822, Pteropogon, Rodanthe, Xyridanthe, Anisolepis, Hyalosperma, Triptilodiscus, Acroclinium, Monencyanthes, Dimorpholepis, Duttonia, Cassiniola.)									
H. Manglesii, F. v. M., fragm. V, 200 (1866)	...	...	...	W.A.	—	—	—	—	—
H. Margaritae, F. v. M., fragm. XI, 48 (1878)	...	...	...	—	—	—	—	N.A.	—
H. roseum, Benth. Fl. Austr. III, 640 (1866)	...	...	...	W.A.	S.A.	—	—	—	—
H. anthemoides, De Candolle, prodr. VI, 216 (1837)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.
H. polygalifolium, De Candolle, prodr. VI, 216 (1837)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.
H. rubellum, Benth. Fl. Austr. III, 641 (1866)	...	...	...	W.A.	—	—	—	—	—
H. chlorocephalum, Benth. Fl. Austr. III, 641 (1866)	...	...	...	W.A.	—	—	—	—	—
H. floribundum, De Candolle, prodr. VI, 217 (1837)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.
H. Pyrethrum, Benth. Fl. Austr. III, 642 (1866)	...	...	...	W.A.	—	—	—	—	—
H. heteranthum, Turczan. in Bull. de Mosc. XXIV, 198 (1851)	...	...	...	W.A.	S.A.	—	—	—	—
H. Kendallii, F. v. M., fragm. VIII, 168 (1874)	...	...	...	W.A.	—	—	—	—	—
H. stipitatum, F. v. M. in Benth. Fl. Austr. III, 643 (1866)	...	...	...	—	S.A.	—	—	—	—
H. incanum, De Candolle, prodr. VI, 215 (1837)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.
H. Cotula, De Candolle, prodr. VI, 215 (1837)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.
H. hyalospermum, F. v. M. in Benth. Fl. Austr. III, 644 (1866)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—
H. condensatum, F. v. M., fragm. III, 136 (1863)	...	...	...	W.A.	—	—	—	—	N.A.
H. polyphyllum, F. v. M., fragm. I, 35 (1858)	...	...	...	—	—	—	—	N.S.W.	Q.

H. Humboldtianum, De Candolle, prodr. VI, 216 (1866)	...	W.A.	—	—	—	—	N.A.	B.f.III, 645	M.fr.III, 135; X, 109.
H. Haigii, F. v. M., fragm. X, 107 (1877) ...	...	W.A.	S.A.	—	—	—	—	—	M.fr. X, 107. [20.
H. tenellum, Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	...	W.A.	S.A.	—	—	—	—	B.f.III, 646	M.fr.III, 135; X, 109; XI,
H. strictum, Benth., Fl. Austr. III, 646 (1866)	...	W.A.	S.A.	—	—	N.S.W.	—	B.f.III, 646	M.fr. X, 109.
H. corymbiflorum, Schlechtendal, Linnaea XXI, 448 (1848)	...	—	S.A.	—	V.	N.S.W.	Q.	B.f.III, 647	M.fr. X, 109.
H. pygmaeum, Benth., Fl. Austr. III, 647 (1866)	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f.III, 647	M.fr. X, 109.
H. spicatum, F. v. M. in Benth., Fl. Austr. III, 647 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 647	M.fr. X, 109.
H. Charsleyae, F. v. M., fragm. VIII, 168 (1874) ...	...	W.A.	S.A.	—	—	—	—	—	M.fr. VIII, 168; XI, 134.
H. moschatum, Benth., Fl. Austr. III, 648 (1866)	...	—	S.A.	—	V.	N.S.W.	Q.	B.f.III, 648	M.fr. X, 107.
H. Tietkensii, F. v. M., fragm. VIII, 227 (1874) ...	...	—	S.A.	—	—	—	—	—	M.fr. VIII, 227.
H. pterochaetum, Benth., Fl. Austr. III, 648 (1866)	...	—	S.A.	—	—	N.S.W.	—	B.f.III, 648	M.fr. X, 109.
H. polycephalum, Benth., Fl. Austr. III, 649 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 649	M.fr.III, 139; X, 109.
H. corymbosum, Benth., Fl. Austr. III, 649 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 649	M.fr. V, 200.
H. laeve, Benth., Fl. Austr. III, 649 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 649	M.fr. V, 200.
H. exiguum, F. v. M. in Transact. Vict. Inst. 39 (1855) ...	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.III, 649	M.fr. X, 109.
H. Dimorpholepis, Benth., Fl. Austr. III, 650 (1866) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f.III, 650	M.fr. X, 109; XI, 27.
HELICHRYSUM, Vaillant in Act. Acad. Par. 290 (1719), from Theophrastus and Dioscorides. (Elichrysium, Schoenia, Petalolepis, Faustula, Ozothamnus, Swammerdamia, Lawrenceella, Argyrophanes, Chrysocephalum, Conanthodium, Xanthochrysium, Argyroglottis, Acanthocladium, Raoulia partly.)									
H. Cassinianum, Gaudichaud in Freyc. Voy. Bot. 466, t. 87 (1826)	...	W.A.	S.A.	—	—	—	—	B.f.III, 616	
H. Lawrenceella, F. v. M. in Benth. Fl. Austr. III, 616 (1866) ...	...	W.A.	S.A.	—	—	—	—	B.f.III, 616	
H. subulifolium, F. v. M., fragm. III, 134 (1863) ...	...	W.A.	—	—	—	—	—	B.f.III, 616	M.fr. III, 134.
H. filifolium, F. v. M., fragm. III, 134 (1863) ...	...	W.A.	—	—	—	—	—	B.f.III, 617	M.fr. III, 134.
H. semifertile, F. v. M., Rep. Babb. Exped. 14 (1858) ...	...	—	S.A.	—	—	N.S.W.	—	B.f.III, 617	
H. Baxteri, Cunningham in De Cand. prodr. VI, 193 (1837)	...	—	S.A.	V.	—	N.S.W.	—	B.f.III, 617	
H. Ayersii, F. v. M., fragm. VIII, 167 (1874) ...	...	—	S.A.	—	—	—	—	—	M.fr. IX, 195.
H. rutidolepis, De Candolle, prodr. VI, 194 (1837) ...	...	—	S.A.	—	V.	N.S.W.	—	B.f.III, 618	
H. scorpioides, Labillardiere, Nov. Holl. pl. spec. II, 45, t. 191 (1806)	...	—	S.A.	T.	V.	N.S.W.	Q.	B.f.III, 618	M.fr. VIII, 167.
H. oligochaetum, F. v. M., fragm. VI, 235 (1868) ...	...	W.A.	—	—	—	—	N.A.	—	M.fr. VI, 235.
H. Tepperi, F. v. M. in Wing's South. Science Record, II, 1 (1882)	...	—	S.A.	—	—	—	—	—	
H. obtusifolium, Sonder & F. v. M. in Linnaea XXV, 515 (1852)	...	W.A.	S.A.	—	V.	N.S.W.	—	B.f.III, 619	
H. Calvertianum, F. v. M., fragm. X, 108 (1877) ...	...	—	—	—	—	N.S.W.	—	—	M.fr. XI, 136.
H. Spicerii, F. v. M., fragm. XI, 47 (1878) ...	...	—	—	T.	—	—	—	—	M.fr. XI, 47.
H. dealbatum, Labillardiere, Nov. Holl. pl. spec. II, 45, t. 190 (1806)	...	—	—	T.	—	—	—	B.f.III, 619	
H. pumilum, J. Hooker, Fl. Tasman. I, 213, t. 60 (1860)	...	—	—	T.	—	—	—	B.f.III, 619	
H. lucidum, Henckel, adumbr. pl. hort. Hal. 5 (1806)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.III, 620
H. elatum, Cunningham in De Cand. prodr. VI, 193 (1837)	...	—	—	—	V.	N.S.W.	Q.	—	M.fr. XI, 48.
H. adenophorum, F. v. M. in Transact. Vict. Inst. 38 (1855)	...	—	S.A.	—	—	—	—	B.f.III, 621	
H. leucopsidium, De Candolle, prodr. VI, 193 (1837)	...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f.III, 622	
H. Blandowskianum, Steetz in Schlecht. Linnaea XXV, 512 (1852)	...	—	S.A.	—	V.	N.S.W.	—	B.f.III, 622	M.fr. VI, 255.
H. oxylepis, F. v. M., fragm. I, 35 (1858) ...	...	—	—	—	—	N.S.W.	Q.	B.f.III, 623	M.fr. I, 35.
H. collinum, De Candolle, prodr. VI, 190 (1837) ...	...	—	—	—	—	N.S.W.	Q.	B.f.III, 623	
H. rupicola, De Candolle, prodr. VI, 190 (1837) ...	...	—	—	—	—	—	Q.	B.f.III, 623	
H. podolepideum, F. v. M., Rep. Babb. Exped. 13 (1858)	...	—	S.A.	—	—	N.S.W.	—	B.f.III, 624	
H. Gilesii, F. v. M., fragm. X, 85 (1876) ...	...	W.A.	—	—	—	—	N.A.	—	M.fr. X, 86.
H. ambiguum, Turczaninow in Bull. de Mosc. 195 (1851)	...	W.A.	S.A.	—	—	N.S.W.	—	B.f.III, 609	
H. apiculatum, De Candolle, prodr. VI, 195 (1837)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.III, 624
H. semipapposum, De Candolle, prodr. VI, 195 (1837)	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.III, 625
H. Dockerii, F. v. M. in Benth. Fl. Austr. III, 626 (1866)	...	—	S.A.	—	—	N.S.W.	—	B.f.III, 626	M.fr. II, 150.
H. Aggyroglottis, Benth., Fl. Austr. III, 626 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 626	
H. ramosum, De Candolle, prodr. VI, 181 (1837) ...	...	W.A.	—	—	—	—	—	B.f.III, 626	
H. cordatum, De Candolle, prodr. VI, 180 (1837) ...	...	W.A.	—	—	—	—	—	B.f.III, 627	
H. obovatum, De Candolle, prodr. VI, 180 (1837) ...	...	—	—	—	—	N.S.W.	—	B.f.III, 627	M.fr. II, 89.
H. Bidwillii, Benth., Fl. Austr. III, 627 (1866) ...	...	—	—	—	—	N.S.W.	Q.	B.f.III, 627	
H. Becklerii, F. v. M. in Benth. Fl. Austr. III, 627 (1866)	...	—	—	—	—	N.S.W.	—	B.f.III, 627	M.fr. I, 183.
H. diotophyllum, F. v. M., fragm. V, 150 (1866) ...	...	—	—	—	—	N.S.W.	Q.	B.f.III, 628	M.fr. VIII, 46.
H. Thomsoni, F. v. M., fragm. VIII, 45 (1873) ...	...	—	S.A.	—	—	—	—	—	M.fr. VIII, 45.
H. diosmifolium, Don in Steud. Nom. Bot. ed. sec. 738 (1841)	...	—	—	—	—	N.S.W.	Q.	B.f.III, 628	
H. retusum, F. v. M., fragm. VIII, 46 (1873) ...	...	—	S.A.	—	V.	N.S.W.	—	—	M.fr. VIII, 46.
H. decurrens, F. v. M., fragm. VIII, 46 (1873) ...	...	—	S.A.	—	V.	N.S.W.	—	—	M.fr. VIII, 46.
H. Cunninghamii, Benth., Fl. Austr. III, 629 (1866)	...	—	—	—	—	N.S.W.	—	B.f.III, 629	
H. reticulatum, Lessing in Steud. Nom. Bot. ed. sec. 740 (1841)	...	—	—	T.	—	—	—	B.f.III, 629	
H. cinereum, F. v. M. in Benth. Fl. Austr. III, 629 (1866)	...	—	S.A.	T.	V.	N.S.W.	—	B.f.III, 629	M.fr. V, 199.
H. bracteolatum, Benth., Fl. Austr. III, 630 (1866)	...	—	—	T.	—	—	—	B.f.III, 630	
H. Kempei, F. v. M. in Melb. Chemist, Jan. (1882) ...	...	—	S.A.	—	—	—	—	—	
H. cassinioides, Benth., Fl. Austr. III, 630 (1866)	...	—	—	—	—	—	Q.	B.f.III, 630	
H. Gunnii, F. v. M. in Benth. Fl. Austr. III, 630 (1866)	...	—	—	T.	—	—	—	B.f.III, 630	M.fr. V, 199.
H. rosmarinifolium, Lessing, synops. gen. comp. 274 (1832)	...	—	—	T.	V.	N.S.W.	—	B.f.III, 631	
H. ledifolium, F. v. M. in Benth. Fl. Austr. III, 630 (1866)	...	—	—	T.	—	—	—	B.f.III, 630	M.fr. V, 199.
H. ferrugineum, Lessing, synops. gen. comp. 274 (1832) ...	...	—	S.A.	T.	V.	N.S.W.	—	B.f.III, 631	
H. antennarium, F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	...	—	—	T.	—	—	—	B.f.III, 632	M.fr. V, 199.
H. oboordatum, F. v. M. in Benth. Fl. Austr. III, 632 (1866)	...	—	—	T.	V.	N.S.W.	—	B.f.III, 632	M.fr. V, 199.
H. Backhousii, F. v. M. in Benth. Fl. Austr. III, 632 (1866)	...	—	—	T.	—	—	—	B.f.III, 632	M.fr. V, 199.
H. cuneifolium, F. v. M. in Benth. Fl. Austr. III, 633 (1866)	...	—	—	—	V.	—	—	B.f.III, 633	
H. baccharoides, F. v. M. in Benth. Fl. Austr. III, 633 (1866)	...	—	—	T.	V.	—	—	B.f.III, 633	M.fr. V, 199.
H. lepidophyllum, F. v. M. in Benth. Fl. Austr. III, 633 (1866)	...	W.A.	—	—	—	—	—	B.f.III, 633	M.fr. V, 199.
H. scutellifolium, Benth., Fl. Austr. III, 633 (1866)	...	—	—	T.	—	—	—	B.f.III, 633	M.fr. V, 199.
H. pholidotum, F. v. M. in Benth. Fl. Austr. III, 634 (1866)	...	—	S.A.	—	V.	N.S.W.	—	B.f.III, 634	M.fr. II, 131; V, 199.
H. lycopodioides, Benth., Fl. Austr. III, 634 (1866)	...	—	—	T.	—	—	—	B.f.III, 634	M.fr. V, 199.
H. selaginoides, F. v. M. in Benth. Fl. Austr. III, 634 (1866)	...	—	—	T.	—	—	—	B.f.III, 634	M.fr. V, 199.

- CASSINIA, R. Brown in Transact. Linn. Soc. XII, 126 (1817). (Non zoologorum, Apolochlamys, Achromolaena, Chromochiton.)
- C. leptoccephala, F. v. M., fragm. III, 138 (1863) ... — — — N.S.W. — — B.f. III, 585 M. fr. III, 138.
- C. compacta, F. v. M., fragm. I, 18 (1858) ... — — — N.S.W. Q. — — B.f. III, 585 M. fr. I, 18.
- C. denticulata, R. Brown in Transact. Linn. Soc. XII, 127 (1817) ... — — — N.S.W. — — B.f. III, 586
- C. longifolia, R. Brown in Transact. Linn. Soc. XII, 127 (1817) ... — — T. V. N.S.W. Q. — — B.f. III, 586
- C. aurea, R. Brown in Transact. Linn. Soc. XII, 127 (1817) ... — — — N.S.W. — — B.f. III, 586
- C. aculeata, R. Brown in Transact. Linn. Soc. XII, 127 (1817) ... — S.A. T. V. N.S.W. — — B.f. III, 586
- C. laevis, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ... — S.A. — — N.S.W. Q. — — B.f. III, 587
- C. tenuifolia, Benth. Fl. Austr. III, 585 (1866) ... — — — N.S.W. — — B.f. III, 585
- C. quinquefaria, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ... — — — N.S.W. Q. — — B.f. III, 587
- C. arcuata, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ... W.A. S.A. — V. N.S.W. — — B.f. III, 587
- C. subtropica, F. v. M., fragm. I, 17 (1858) ... — — — N.S.W. Q. — — B.f. III, 588 M. fr. I, 17.
- C. Theodori, F. v. M., fragm. V, 148 (1866) ... — — — N.S.W. — — B.f. III, 588 M. fr. V, 148.
- C. spectabilis, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ... — S.A. T. V. — — — B.f. III, 588
- HUMEA, Smith, Exot. bot. I, t. 1 (1804). (Haeckeria, Calomeria.)
- H. elegans, Smith, Exot. bot. I, t. 1 (1804) ... — — — V. N.S.W. — — B.f. III, 589 M. fr. I, 17.
- H. cassiniacea, F. v. M., fragm. I, 17 (1858) ... — S.A. — — — — — B.f. III, 589 M. fr. I, 17.
- H. punctulata, F. v. M., fragm. III, 137 (1863) ... — S.A. — — — — — B.f. III, 589 M. fr. III, 137.
- H. ozothamnoides, F. v. M., fragm. I, 17 (1858) ... — — — V. N.S.W. — — B.f. III, 590 M. fr. VIII, 142.
- H. squamata, F. v. M., fragm. XI, 86 (1880) ... — S.A. — V. — — — — — B.f. III, 586
- ACOMIS, F. v. M., fragm. II, 89 (1860).
- A. Rutidosi, F. v. M., fragm. II, 89 (1860) ... — — — — — N.S.W. — — B.f. III, 591 M. fr. II, 89.
- A. macra, F. v. M., fragm. IV, 145 (1864) ... — — — — — Q. — — B.f. III, 591 M. fr. IV, 145.
- RUTIDOSIS, De Candolle, prodr. VI, 158 (1837). (Pumilo, Actinopappus, Lepidocoma.)
- R. leioplepis, F. v. M. in Transact. Vict. Inst. 131 (1855) ... — — — — — V. — — — B.f. III, 593
- R. leptorrhynchoideis, F. v. M., fragm. V, 148 (1866) ... — — — — — V. N.S.W. — — B.f. III, 593 M. fr. V, 148.
- R. Brownii, Benth. Fl. Austr. III, 594 (1866) ... — — — — — Q. N.A. — — B.f. III, 594
- R. helichrysoides, De Candolle, prodr. VI, 159 (1837) ... — S.A. — V. N.S.W. — — B.f. III, 594
- R. leucantha, F. v. M., fragm. I, 35 (1858) ... — — — — — Q. — — B.f. III, 594 M. fr. I, 35.
- R. Murchisonii, F. v. M., fragm. I, 34 (1858) ... — — — — — Q. — — B.f. III, 594 M. fr. I, 34.
- R. Pumilo, Benth. Fl. Austr. III, 595 (1866) ... W.A. S.A. T. V. N.S.W. — — B.f. III, 595
- PITHOCARPA, Lindley, Bot. Regist. XXV, App. XXIII (1839).
- P. corymbosa, Lindley, Bot. Regist. XXV, App. XXIII (1839) W.A. — — — — — — — B.f. III, 590
- AMMOBIUM, R. Brown in Bot. Magaz. t. 2459 (1824).
- A. alatum, R. Brown in Bot. Magaz. t. 2459 (1824) ... — — — — — N.S.W. Q. — — B.f. III, 583
- A. craspedioides, Benth. Fl. Austr. III, 584 (1866) ... — — — — — N.S.W. — — B.f. III, 584
- IXODIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812).
- I. achilleoides, R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812) ... — S.A. — V. — — — — B.f. III, 583
- MILLOTIA, Cassini in Ann. sc. nat. XVII, 416 (1829).
- M. tenuifolia, Cassini in Ann. sc. nat. XVII, 416 (1829) ... W.A. S.A. T. V. N.S.W. — — B.f. III, 596
- M. Greevesii, F. v. M., fragm. III, 18, t. 19 (1863) ... — — S.A. — — — — — N.S.W. — — B.f. III, 596 M. fr. III, 18.
- M. Kempei, F. v. M. in Wing's South. Sc. Record II, 1 (1882) ... — S.A. — — — — — — — —
- TOXANTHUS, Turczaninow in Bull. de Mosc. XXIV, 177 (1851). (Anthocerastes, Scyphocoronis.)
- T. perpusillus, Turczaninow in Bull. Mosc. XXIV, 177 (1851) ... W.A. S.A. — V. N.S.W. — — B.f. III, 592 M. fr. XI, 49.
- T. Muellieri, Benth. Fl. Austr. III, 592 (1866) ... W.A. S.A. — — — — — — — B.f. III, 592 M. fr. XI, 49.
- T. major, Turczaninow in Bull. de Mosc. XXIV, 64 (1851) ... W.A. — — — — — — — B.f. III, 592 M. fr. XI, 49.
- QUINETIA, Cassini in Diction. sc. nat. LX, 579 (1830).
- Q. Urvillei, Cassini in Diction. LX, 579 (1830) ... W.A. S.A. — — — — — — — B.f. III, 595
- PHACELOTHRIX, F. v. M., fragm. XI, 49 (1878).
- P. cladochaeta, F. v. M., fragm. XI, 49 (1878) ... — — — — — — — Q. — — — M. fr. V, 199; XI, 49.
- ERIOCHLAMYS, Sonder and F. v. M. in Schlecht. Linnaea XXV, 488 (1852).
- E. Behrii, Sonder & F. v. M. in Linnaea XXV, 488 (1852) ... — S.A. — V. N.S.W. — — B.f. III, 591
- MYRIOCEPHALUS, Benth. in Hueg. enum. 61 (1837). (Hyalolepis, Anthediosorus, Gilberta, Lamprochlaena, Elachopappus, Polycalymma; some readily to be restituted.)
- M. rhizocephalus, Benth. Fl. Austr. III, 557 (1866) ... W.A. S.A. — V. N.S.W. — — B.f. III, 557 M. fr. III, 155.
- M. nudus, A. Gray in Hook. Kew Misc. III, 174 (1851) ... W.A. — — — — — — — B.f. III, 558 M. fr. III, 157.
- M. appendiculatus, Benth. in Hueg. enum. 61 (1837) ... W.A. — — — — — — — B.f. III, 558 [145]
- M. Rudallii, F. v. M., fragm. III, 157 (1863) ... W.A. S.A. — — — — — — — B.f. III, 559 M. fr. III, 157; VI, 235; VII,
- M. gracilis, Benth. Fl. Austr. III, 559 (1866) ... W.A. — — — — — — — B.f. III, 559 M. fr. VIII, 169.
- M. helichrysoides, A. Gray in Hook. Kew Misc. III, 175 (1851) ... W.A. — — — — — — — B.f. III, 559
- M. suffruticosus, Benth. Fl. Austr. III, 559 (1866) ... W.A. — — — — — — — B.f. III, 559
- M. Stuartii, Benth. Fl. Austr. III, 560 (1866) ... — S.A. — V. N.S.W. — — B.f. III, 560
- M. Gueriniae, F. v. M., fragm. VIII, 169 (1874) ... W.A. — — — — — — — — M. fr. VIII, 169.
- ANGIANTHUS, Wendland, Collect. pl. II, 31 (1809). (Siloxerus, Styloncerus, Ogcerostylus, Cylindrosorus, Phyllocalymma, Skirrophorus, Chrysocoryne, Eriocladium, Pogonolepis, Piptostemma, Epitriche, Gamozygis, Cephalosorus partly, Hyalochlamys, Dithyrostegia, Pleuropappus; some readily to be restituted.)
- A. humifusus, Benth. Fl. Austr. III, 563 (1866) ... W.A. — — — — — — — B.f. III, 563
- A. tomentosus, Wendland, Collect. pl. II, 31, t. 48 (1809) ... W.A. S.A. — V. N.S.W. — N.A. B.f. III, 562
- A. pleuropappus, Benth. Fl. Austr. III, 563 (1866) ... — S.A. — — — — — — — B.f. III, 563
- A. brachypappus, F. v. M. in Trans. phil. Soc. Vict. I, 44 (1854) ... — S.A. — V. N.S.W. — — B.f. III, 563
- A. myosuroides, Benth. Fl. Austr. III, 563 (1866) ... W.A. — — — — — — — B.f. III, 563
- A. tenellus, Benth. Fl. Austr. III, 564 (1866) ... W.A. S.A. — — — — — — — B.f. III, 564
- A. pusillus, Benth. Fl. Austr. III, 564 (1866) ... W.A. S.A. — V. N.S.W. — — B.f. III, 564
- A. Milnei, Benth. Fl. Austr. III, 564 (1866) ... W.A. — — — — — — — B.f. III, 564
- A. Cunninghamii, Benth. Fl. Austr. III, 565 (1866) ... W.A. — — — — — — — B.f. III, 565
- A. phyllocephalus, Benth. Fl. Austr. III, 565 (1866) ... W.A. — — — — — — — B.f. III, 565 M. fr. III, 159.

A. micropoides, Benthām, Fl. Austr. III, 565 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 565	
A. microcephalus, Benthām, Fl. Austr. III, 566 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 566	M.fr. III, 158.
A. platycephalus, Benthām, Fl. Austr. III, 566 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 566	
A. Drummondii, Benthām, Fl. Austr. III, 566 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 566	
A. Preissianus, Benthām, Fl. Austr. III, 566 (1866) ...	W.A. S.A. T. V. N.S.W.	—	—	—	—	—	B.f. III, 566	
A. pygmaeus, Benthām, Fl. Austr. III, 567 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 567	
A. globifer, Benthām, Fl. Austr. III, 567 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 567	
A. demissus, Benthām, Fl. Austr. III, 567 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 567	
A. strictus, Benthām, Fl. Austr. III, 568 (1866) ...	W.A. S.A. — V. N.S.W.	—	—	—	—	—	B.f. III, 568	
A. plumiger, Benthām, Fl. Austr. III, 568 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 568	
A. amplexicaulis, Benthām, Fl. Austr. III, 568 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 568	
GNEPHOSIS, Cassini in Bull. de la Soc. philom. 45 (1820). (Cephalosorus partly, Nematopus, Crossolepis, Leptotriche, Trichanthodium, Cyathopappus; some readily to be restituted.)								
G. Burkittii, Benthām, Fl. Austr. III, 570 (1866) ...	— S.A.	—	—	—	—	—	B.f. III, 570	
G. eriocarpa, Benthām, Fl. Austr. III, 570 (1866) ...	— S.A.	—	—	N.S.W.	—	—	B.f. III, 570	
G. macrocephala, Turcz. in Bull. Soc. Mosc. XXIV, 190 (1851) ...	W.A.	—	—	—	—	—	B.f. III, 570	
G. skirophora, Benthām, Fl. Austr. III, 570 (1866) ...	W.A. S.A. — V. N.S.W.	—	—	—	—	—	B.f. III, 570	
G. cyathopappa, Benthām, Fl. Austr. III, 571 (1866) ...	— S.A.	—	—	N.S.W.	—	N.A.	B.f. III, 571	
G. codonopappa, F. v. M. in Giles, geogr. trav. 217 (1875) ...	— S.A.	—	—	—	—	—	—	M.fr. IX, 2.
G. leptoclada, Benthām, Fl. Austr. III, 571 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 571	M.fr. III, 158.
G. arachnoidea, Turcz. in Bull. Soc. Mosc. XXIV, 189 (1851) ...	W.A. S.A.	—	—	—	—	—	B.f. III, 571	
G. tenuissima, Cassini in Bull. de la Soc. philom. 43 (1820) ...	W.A.	—	—	—	—	—	B.f. III, 572	
G. acicularis, Benthām, Fl. Austr. III, 572 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 572	
G. pygmaea, Benthām, Fl. Austr. III, 572 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 572	
G. brevifolia, Benthām, Fl. Austr. III, 572 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 572	
G. eriocephala, Benthām, Fl. Austr. III, 573 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 573	
DECAZESIA, F. v. M., fragm. XI, 71 (1879).								
D. hecatcephala, F. v. M., fragm. XI, 72 (1879) ...	—	—	—	—	—	N.A.	—	M.fr. XI, 72.
CALOCEPHALUS, R. Brown in Transact. Linn. Soc. XII, 106 (1817). (Leucophyta, Pachysura, Blennospora, Achrysum.)								
C. Drummondii, Benthām, Fl. Austr. III, 574 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 574	
C. Brownii, F. v. M., Rep. Rabb. Exped. 13 (1858) ...	W.A. S.A. T. V. N.S.W.	—	—	—	—	—	B.f. III, 574	
C. Sonderi, F. v. M., Rep. Rabb. Exped. 13 (1858) ...	— S.A. T. V. N.S.W.	—	—	—	—	—	B.f. III, 575	
C. lacteus, Lessing, syn. gen. comp. 271 (1832) ...	W.A. S.A. T. V. N.S.W.	—	—	—	—	—	B.f. III, 575	
C. citreus, Lessing, syn. gen. comp. 271 (1832) ...	— S.A. T. V. N.S.W.	—	—	—	—	—	B.f. III, 575	
C. angianthoides, Benthām, Fl. Austr. III, 575 (1866) ...	W.A. — T.	—	—	—	—	—	B.f. III, 575	
C. Francisii, Benthām, Fl. Austr. III, 576 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 576	
C. platycephalus, Benthām, Fl. Austr. III, 576 (1866) ...	— S.A.	—	—	N.S.W.	—	—	B.f. III, 576	
C. multiflorus, Benthām, Fl. Austr. III, 576 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 576	
C. aeruoides, Benthām, Fl. Austr. III, 576 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 576	
CEPHALIPTERUM, A. Gray in Hook. Kew Misc. IV, 271 (1852).								
C. Drummondii, A. Gray in Hook. Kew Misc. IV, 272 (1852) ...	W.A. S.A.	—	—	—	—	—	B.f. III, 577	
GNAPHALODES, A. Gray in Hook. Kew Misc. IV, 228 (1852).								
G. uliginosus, A. Gray in Hook. Kew Misc. IV, 228 (1852) ...	W.A. S.A. — V. N.S.W. Q.	—	—	—	—	—	B.f. III, 578	M.fr. XI, 27.
G. condensatum, A. Gray in Hook. Kew Misc. IV, 228 (1852) ...	W.A.	—	—	—	—	—	B.f. III, 578	
G. filifolium, Benthām, Fl. Austr. III, 578 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 578	
CRASPEDIA, G. Forster, florul. insul. Austr. prodr. 306 (1786). (Richea, Pycnosorus.)								
C. Richea, Cassini in Diction. XI, 353 (1818) ...	W.A. S.A. T. V. N.S.W. Q.	—	—	—	—	—	B.f. III, 579	
C. pleiocephala, F. v. M. in Schlecht. Linnaea XXV, 404 (1852) ...	— S.A. — V. N.S.W.	—	—	—	—	—	B.f. III, 580	M.fr. XI, 27.
C. chrysanthia, Benthām, Fl. Austr. III, 580 (1866) ...	— S.A. — V. N.S.W. Q.	—	—	—	—	—	B.f. III, 580	
C. globosa, Benthām, Fl. Austr. III, 580 (1866) ...	— S.A. — V. N.S.W.	—	—	—	—	—	B.f. III, 580	
CHTHONOCEPHALUS, Steetz in Lehm. pl. Preiss. I, 444 (1845). (Chamaesphaerion, Gyrostephium, Lachnothalamus.)								
C. tomentellus, Benthām, Fl. Austr. III, 581 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 581	
C. Pseudevax, Steetz in Lehm. pl. Preiss. I, 445 (1845) ...	W.A. S.A. — V. N.S.W.	—	—	—	—	—	B.f. III, 582	
C. pygmaeus, Benthām, Fl. Austr. III, 582 (1866) ...	W.A.	—	—	—	—	—	B.f. III, 582	
NABLONIUM, Cassini in Diction. XXXIV, 101 (1825).								
N. calyceroides, Cassini, Dict. XXXIV, 101 (1825) ...	—	—	T.	—	—	—	B.f. III, 545	
CHRYSOGONUM, Linné, hort. Cliffort. 424 (1737). (Moonia, Pentalepis.)								
C. trichodesmoides, Benthām & J. Hooker, gen. pl. II, 350 (1873) ...	—	—	—	—	—	N.A.	B.f. III, 540	M.fr. VIII, 145.
C. ecliptoides, Benthām & J. Hooker, gen. pl. II, 350 (1873) ...	—	—	—	—	—	N.A.	B.f. III, 540	M.fr. VIII, 145.
C. procumbens, Benthām & J. Hooker, gen. pl. II, 350 (1873) ...	—	—	—	—	—	N.A.	B.f. III, 540	M.fr. VIII, 145.
SIEGESBECKIA, Linné, hort. Cliffort. 412 (1737).								
S. orientalis, Linné, spec. plant. 900 (1753) ...	W.A. S.A. — V. N.S.W. Q.	—	—	—	—	—	B.f. III, 535	
ENHYDRA, Loureiro, Flor. Cochinch. II, 510 (1790). (Enydra.)								
E. paludosa, De Candolle, prodr. V, 637 (1836) ...	—	—	—	—	N.S.W. Q.	—	B.f. III, 546	M.fr. III, 139.
ECLIPTA, Linné, mantissa altera, 157 (1771).								
E. alba, Hasskarl, plant. Javan. rar. 528 (1848) ...	—	—	—	—	N.S.W. Q.	—	B.f. III, 536	
E. latifolia, Linné, fil. suppl. 378 (1781) ...	—	—	—	—	Q.	N.A.	—	M.fr. VIII, 145.
E. platyglossa, F. v. M., fragm. II, 135 (1861) ...	— S.A. — V. N.S.W. Q.	N.A.	—	—	—	—	B.f. III, 536	M.fr. XI, 27.
WEDELIA, Jacquin, stirp. Amer. hist. 217, t. 130 (1763). (Wollastonia.)								
W. calendulacea, Lessing, synops. gen. comp. 222 (1832) ...	—	—	—	—	—	Q.	—	B.f. III, 537
W. urticifolia, De Candolle in Wight, Contrib. Bot. Ind. 18 (1834) ...	—	—	—	—	—	N.A.	B.f. III, 538	
W. spilanthis, F. v. M., fragm. V, 64 (1865) ...	—	—	—	—	N.S.W. Q.	—	B.f. III, 538	M.fr. V, 64.
W. verbesinoides, F. v. M. in Benth. Fl. Austr. III, 538 (1866) ...	—	—	—	—	—	N.A.	B.f. III, 538	
W. biflora, De Candolle in Wight, Contrib. Bot. Ind. 18 (1834) ...	—	—	—	—	N.S.W. Q.	N.A.	B.f. III, 539	
W. asperrima, Benthām, Fl. Austr. III, 539 (1866) ...	—	—	—	—	—	N.A.	B.f. III, 539	

- SPILANTHES**, Jacquin, stirp. Amer. hist. 214, t. 126 (1763).
S. grandiflora, Turczaninow in Bull. de Mosc. XXIV, 185 (1851) — — — — N.S.W. Q. N.A. B.f.III,541 M.fr.V,63.
S. anactina, F. v. M., fragm. V, 63 (1865) ... — — — — N.A. B.f.III,541 M.fr.V,63.
- BIDENS**, Dillenius, hort. Eltham. 51 and 52, t. 43 et 44 (1732), from Cesalpini (1583). (Possibly immigrated.)
B. tripartitus, Linné, spec. plant. 831 (1753) ... — — — — V. N.S.W. — — B.f.III,543
B. pilosus, Linné, spec. plant. 832 (1753) ... — — — — N.S.W. Q. — — B.f.III,543
B. bipinnatus, Linné, spec. plant. 832 (1753) ... — — — — N.S.W. Q. N.A. B.f.III,543
- GLOSSOGYNE**, Cassini in Diction. LI, 475 (1827). (Diodontium.)
G. tenuifolia, Cassini in Dict. LI, 475 (1827) ... — — — — S.A. — — N.S.W. Q. N.A. B.f.III,544 M.fr.I,51.
G. retroflexa, F. v. M., fragm. I, 51 (1858) ... — — — — — Q. — — B.f.III,544
G. filifolia, F. v. M. in Benth. Fl. Austr. III, 544 (1866) ... — — — — — N.A. B.f.III,544
- FLAVERIA**, A. L. de Jussieu, gen. plant. 186 (1789).
F. Australasica, Hooker in Mitch. Trop. Austr. 118 (1848) ... — — — — — Q. N.A. B.f.III,546 M.fr.I,183.
- SOLIVA**, Ruiz et Pavon, prodr. 113, t. 24 (1794). (Gymnostyles; possibly immigrated.)
S. anthemifolia, R. Brown in Transact. Linn. Soc. XII, 102 (1817) — — — — N.S.W. Q. — — B.f.III,552
- COTULA**, Linné, syst. nat. 9 (1753); Linné, gen. pl. 256 (1737). (Gymnogyne, Strongylospermum, Pleiogyne, Symphyomera, Stenosperma, Leptinella.)
C. filifolia, Thunberg, prodr. pl. Capens. 161 (1800) ... W.A. S.A. T. V. N.S.W. — — B.f.III,548 M.fr.VI,114.
C. coronopifolia, Linné, sp. pl. 892 (1753) ... W.A. S.A. T. V. N.S.W. — — B.f.III,549 M.fr.VI,114.
C. Gymnogyne, F. v. M. in Benth. Fl. Austr. III, 549 (1866) ... W.A. — — — — — B.f.III,549
C. australis, J. Hooker, Fl. N. Zeal. I, 128 (1853) ... W.A. S.A. T. V. N.S.W. Q. — — B.f.III,550
C. alpina, J. Hooker, Fl. Tasman. I, 192, t. 51 (1860) ... — — T. V. N.S.W. — — B.f.III,550
C. Drummondii, Benth. Fl. Austr. III, 550 (1866) ... W.A. — — — — — B.f.III,550
C. reptans, Benth. Fl. Austr. III, 551 (1866) ... — — S.A. T. V. N.S.W. — — B.f.III,551
C. Filicula, J. Hooker in Benth. Fl. Austr. III, 551 (1866) ... — — T. V. N.S.W. — — B.f.III,551
- CENTIPEDA**, Loureiro, Fl. Cochinch. II, 492 (1790). (Myriogyne, Sphaeromorphaea partly.)
C. racemosa, F. v. M.; Myriogyne, Hook. in Mitch. Tr. Austr. 353 (1848) — — — — N.S.W. Q. N.A. B.f.III,553
C. orbicularis, Loureiro, Fl. Cochinch. II, 492 (1790) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.f.III,553 M.fr.VIII,142.
C. Cunninghamii, F. v. M., fragm. VIII, 143 (1874) ... W.A. S.A. — — V. N.S.W. — — — —
C. thespidioides, F. v. M., fragm. VIII, 143 (1874) ... — — S.A. — — V. N.S.W. — — — —
- ABROTANELLA**, Cassini in Dict. XXXVI, 27 (1825). (Scleroleima, Trineuron.)
A. fosterioides, J. Hooker, Hand. N. Zeal. Fl. 139 (1864) ... — — — — T. — — — — B.f.III,554
A. nivigena, F. v. M. in Benth. Fl. Austr. III, 554 (1866) ... — — — — V. — — — — B.f.III,554
A. scapigera, F. v. M. in Benth. Fl. Austr. III, 554 (1866) ... — — — — T. — — — — B.f.III,554
- ELACHANTHUS**, F. v. M. in Schlecht. Linnaea XXV, 410 (1852).
E. pusillus, F. v. M. in Schlecht. Linnaea XXV, 411 (1852) ... — — S.A. — — V. N.S.W. — — B.f.III,555
- CERATOGYNE**, Turczaninow in Bull. de Mosc. XXIV, 68 (1851). (Diotosperma.)
C. obionoides, Turczaninow in Bull. de Mosc. XXIV, 68 (1851) W.A. — — — — — — B.f.III,555
- ISOETOPSIS**, Turczaninow in Bull. de Mosc. XXIV, 175 (1851).
I. graminifolia, Turcz. in Bull. de Mosc. XXIV, 175, t. 3 (1851) W.A. S.A. — — V. N.S.W. — — B.f.III,556
- EMILIA**, Cassini in Bulletin de la Soc. philomat. 68 (1817).
E. purpurea, Cassini in Diction. des sc. nat. XXXIV, 393 (1825) — — — — — — Q. — — — — M.fr.XII.
- GYNURA**, Cassini in Diction. des sc. nat. XXXIV, 391 (1825).
G. Pseudo-China, De Candolle, prodr. VI, 299 (1837) ... — — — — — N.S.W. Q. — — B.f.III,661
- SENECIO**, Tournefort, inst. 456, t. 260 (1700), from Plinius. (Bedfordia, Centropappus.)
S. Gregorii, F. v. M., pl. of Greg. Search-Exped. for Leichh. 7 (1859) — — S.A. — — V. N.S.W. Q. — — B.f.III,661
S. platylepis, De Candolle, prodr. VI, 371 (1837) ... — — — — V. N.S.W. Q. — — B.f.III,662 M.fr.XI,27.
S. papillosus, F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857) ... — — T. — — — — — B.f.III,662
S. primulifolius, F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857) ... — — T. — — — — — B.f.III,662
S. pectinatus, De Candolle, prodr. VI, 372 (1837) ... — — — — T. V. N.S.W. — — B.f.III,662
S. spathulatus, A. Richard, sert. Astrol. 125 (1833) ... — — S.A. T. V. N.S.W. — — B.f.III,663
S. megaglossus, F. v. M. in Linnaea XXV, 419 (1852) ... — — S.A. — — — — — B.f.III,663
S. magnificus, F. v. M. in Linnaea XXV, 418 (1852) ... — — S.A. — — V. N.S.W. — — B.f.III,663
S. insularis, Benth. Fl. Austr. III, 666 (1866) ... — — — — — N.S.W. — — B.f.III,664
S. Centropappus, F. v. M., annual report 26 (1858) ... — — — — T. — — — — B.f.III,664
S. macranthus, A. Richard, sert. Astrol. 126 (1833) ... — — — — — N.S.W. — — B.f.III,664
S. Daltoni, F. v. M., fragm. VI, 27 (1867) ... — — — — — N.S.W. Q. — — — — M.fr.VI,27.
S. lautus, Solander in G. Forst. prodr. 91 (1786) ... W.A. S.A. T. V. N.S.W. Q. — — B.f.III,665 M.fr.XI,20.
S. capillifolius, J. Hooker in Lond. Journ. VI, 123 (1847) ... — — — — T. — — — — B.f.III,667
S. vagus, F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854) ... — — — — V. N.S.W. — — B.f.III,667
S. amygdalifolius, F. v. M., fragm. I, 232 (1859) ... — — — — — N.S.W. Q. — — B.f.III,668 M.fr.I,232.
S. velleioides, Cunningham in De Cand. prodr. VI, 374 (1837) ... — — — — T. V. N.S.W. — — B.f.III,668
S. australis, A. Richard, sert. Astrol. 131, t. 39 (1833) ... — — S.A. T. V. N.S.W. — — B.f.III,668
S. Behrianus, Sonder and F. v. M. in Linnaea XXV, 527 (1852) ... — — S.A. — — V. N.S.W. — — B.f.III,669
S. leucoglossus, F. v. M., fragm. II, 15 (1861) ... W.A. — — — — — — B.f.III,669 M.fr.II,15.
S. brachyglossus, F. v. M. in Schlecht. Linnaea XXV, 525 (1852) W.A. S.A. — — V. N.S.W. — — B.f.III,669 M.fr.XI,27.
S. Georgianus, De Candolle, prodr. VI, 371 (1837) ... W.A. S.A. T. V. N.S.W. — — B.f.III,670
S. Gilberti, Turczaninow in Bull. Soc. Mosc. XXIV, 208 (1851) W.A. — — — — — — B.f.III,670
S. ramosissimus, De Candolle, prodr. VI, 371 (1837) ... W.A. — — — — — — B.f.III,671
S. odoratus, Horneman, hort. reg. bot. Hafn. II, 809 (1815) ... — — S.A. T. V. N.S.W. — — B.f.III,671
S. Cunninghamii, De Candolle, prodr. VI, 371 (1837) ... W.A. S.A. — — V. N.S.W. — — B.f.III,671
S. anethifolius, Cunningham in De Cand. prodr. VI, 371 (1837) ... — — S.A. — — — — — N.S.W. Q. — — B.f.III,672
S. Bedfordii, F. v. M. annual report 26 (1858) ... — — — — T. V. N.S.W. — — B.f.III,673
S. Billardieri, F. v. M. annual report 26 (1858) ... — — — — T. — — — — — B.f.III,673

ERECHTITES, Rafinesque, Fl. Ludov. 65 (1817).									
E. prenanthoides, De Candolle, prodr. VI, 296 (1837)	...	...	—	—	T. V. N.S.W.	—	—	B.f.III, 658	M.fr.V, 88.
E. Atkinsoniae, F. v. M., fragm. V, 88 (1866)	...	...	—	—	—	N.S.W. Q.	—	B.f.III, 658	
E. arguta, De Candolle, prodr. VI, 296 (1837)	...	...	W.A.	S.A.	T. V. N.S.W.	Q.	—	B.f.III, 659	
E. mixta, De Candolle, prodr. VI, 297 (1837)	...	...	—	—	—	N.S.W.	—	B.f.III, 659	
E. quadridentata, De Candolle, prodr. VI, 295 (1837)	...	...	W.A.	S.A.	T. V. N.S.W.	Q.	—	B.f.III, 660	
E. hispidula, De Candolle, prodr. VI, 296 (1837)	...	...	W.A.	S.A.	T. V. N.S.W.	—	—	B.f.III, 660	
CYMBONOTUS, Cassini in Diction. XXXV, 397 (1825).									
C. Lawsonianus, Gaudichaud in Freyc. Voy. Bot. 462, t. 86 (1826)	W.A.	S.A.	T. V. N.S.W.	Q.	—	—	—	B.f.III, 674	M.fr.XI, 27.
SAUSSUREA, De Candolle in Ann. du Mus. XVI, 156 (1810). (Aplotaxis, Haplotaxis.)									
S. carthamoides, Benth, Fl. Hongk. 168 (1861)	...	...	—	—	—	N.S.W. Q.	—	B.f.III, 456	
CENTAUREA, Linné, hort. Cliffort. 420 (1737). (Leuzea partly.)									
C. Australis, Benth, Fl. Hongk. 168 (1861)	...	...	—	—	—	V. N.S.W. Q.	—	B.f.III, 457	
TRICHOCLINE, Cassini in Bull. de la Soc. philom. 13 (1817). (Amblyserma.)									
T. scapigera, Benth, Fl. Hongk. 168 (1861)	...	...	W.A.	—	—	—	—	B.f.III, 676	
MICROSERIS, D. Don in Edinb. phil. Mag. XI, 388 (1832). (Phyllopappus, Scozonera partly.)									
M. Forsteri, J. Hooker, Fl. N. Zeal. I, 151 (1853)	...	...	W.A.	S.A.	T. V. N.S.W.	—	—	B.f.III, 676	M.fr.XI, 27.
CREPIS, Linné, gen. pl. 240 (1737). (Youngia.)									
C. Japonica, Benth, Fl. Hongk. 194 (1861)	...	...	—	—	—	N.S.W. Q.	—	B.f.III, 679	

CAMPANULACEAE.

A. L. de Jussieu, gen. plant. 163 (1789), from B. de Jussieu (1759).

LOBELIA, Linné, Fl. Lappon. 227 (1737). (Rapuntium, Pratia, Grammatotheca, Holostigma.)									
L. heterophylla, Labillard., Nov. Holl. pl. spec. I, 52, t. 74 (1804)	W.A.	S.A.	—	—	—	—	—	B.f.IV, 124	M.fr.IV, 184; X, 41.
L. Browniana, Roemer & Schultes, syst. veg. V, 71 (1819)	...	...	—	S.A.	T. V. N.S.W.	Q.	—	—	
L. simplicicaulis, R. Brown, prodr. 564 (1810)	...	...	—	S.A.	T. V. N.S.W.	—	—	B.f.IV, 124	M.fr.IV, 184; X, 41.
L. microsperma, F. v. M., fragm. X, 41 (1876)	...	...	W.A.	S.A.	T. V. N.S.W.	Q.	—	—	M.fr.X, 41.
L. dentata, Cavanilles, icon. pl. VI, 14, t. 522 (1801)	...	...	—	—	—	N.S.W. Q.	—	B.f.IV, 125	M.fr.IV, 183; X, 42.
L. gracilis, Andrews, Bot. Reposit. t. 340 (1805)	...	...	—	—	—	N.S.W. Q.	—	B.f.IV, 125	M.fr.IV, 183; X, 42.
L. rhytidisperma, Benth, Fl. Austr. IV, 126 (1869)	...	...	W.A.	—	—	—	—	B.f.IV, 126	M.fr.X, 40.
L. tenuior, R. Brown, prodr. 564 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 126	M.fr.IV, 183; X, 42.
L. rhombifolia, De Vriese in Lehm. pl. Preiss. I, 397 (1845)	...	...	W.A.	S.A.	—	V.	—	B.f.IV, 126	M.fr.X, 42.
L. parvifolia, R. Brown, prodr. 564 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 127	M.fr.IV, 183.
L. trigonocaulis, F. v. M., fragm. I, 18 (1858)	...	...	—	—	—	N.S.W. Q.	—	B.f.IV, 127	M.fr.I, 18, 242.
L. Bergiana, Chamisso in Schlecht. Linnaea VIII, 217 (1835)	...	...	W.A.	—	—	—	—	B.f.IV, 127	M.fr.II, 20; IV, 171; X, 43.
L. anceps, Thunberg, prodr. plant. cap. 40 (1794)	...	...	W.A.	S.A.	T. V. N.S.W.	Q.	—	B.f.IV, 128	M.fr.IV, 184; VI, 114; X, 43.
L. surrepens, J. Hooker, Fl. Tasman. I, 237, t. 69 (1860)	...	...	—	—	T.	—	—	B.f.IV, 129	M.fr.IV, 183.
L. membranacea, R. Brown, prodr. 563 (1810)	...	...	—	—	—	—	Q.	B.f.IV, 129	M.fr.IV, 171; X, 42.
L. stenophylla, Benth, Fl. Austr. III, 130 (1869)	...	...	—	—	—	—	Q. N.A.	B.f.IV, 130	M.fr.X, 42.
L. humistrata, F. v. M. in Benth. Fl. Austr. III, 130 (1869)	...	...	—	—	—	—	Q. N.A.	B.f.IV, 130	M.fr.IV, 171; X, 42.
L. dioica, R. Brown, prodr. 565 (1810)	...	...	—	—	—	—	Q. N.A.	B.f.IV, 130	M.fr.IV, 183; X, 43.
L. purpurascens, R. Brown, prodr. 563 (1810)	...	...	—	S.A.	—	V. N.S.W. Q.	—	B.f.IV, 131	M.fr.IV, 183; X, 42.
L. pratoides, Benth, Fl. Austr. IV, 131 (1869)	...	...	—	S.A.	T. V. N.S.W.	—	—	B.f.IV, 131	M.fr.X, 43.
L. irrigua, R. Brown, prodr. 563 (1810)	...	...	—	—	T.	—	—	B.f.IV, 132	
L. gelida, F. v. M., fragm. IV, 183 (1864)	...	...	—	—	—	V. N.S.W.	—	B.f.IV, 132	M.fr.IV, 183.
L. platycalyx, F. v. M., fragm. IV, 183 (1864)	...	...	—	—	—	T. V.	—	B.f.IV, 133	M.fr.X, 43.
L. concolor, R. Brown, prodr. 563 (1810)	...	...	—	S.A.	—	V. N.S.W. Q.	—	B.f.IV, 133	M.fr.X, 43.
L. Benthami, F. v. M.; Pratia puberula, Benth. Fl. Austr. IV, 133 (1869)	...	...	—	—	—	V. N.S.W.	—	B.f.IV, 133	
L. pedunculata, R. Brown, prodr. 563 (1810)	...	...	—	S.A.	T. V. N.S.W.	Q.	—	B.f.IV, 133	M.fr.X, 42.
ISOTOMA, R. Brown, prodr. 565 (1810). (Lobelia partly, Enchysia partly, Laurencia partly.)									
I. Brownii, G. Don, gen. syst. III, 716 (1834)	...	...	W.A.	—	—	—	—	B.f.IV, 134	M.fr.X, 40.
I. axillaris, Lindley, Bot. Regist. t. 964 (1826)	...	...	—	—	—	V. N.S.W. Q.	—	B.f.IV, 135	M.fr.X, 39.
I. petraea, F. v. M. in Linnaea XXV, 420 (1852)	...	...	W.A.	S.A.	—	N.S.W. Q.	—	B.f.IV, 135	M.fr.X, 39.
I. pusilla, Benth, Fl. Austr. III, 130 (1869)	...	...	W.A.	—	—	—	—	B.f.IV, 135	
I. Gulliverii, F. v. M., fragm. X, 39 (1876)	...	...	—	—	—	—	N.A.	—	M.fr.X, 39.
I. scapigera, G. Don, gen. syst. III, 716 (1834)	...	...	W.A.	S.A.	—	—	—	B.f.IV, 136	M.fr.IV, 183; X, 39.
I. fluviatilis, F. v. M. in Benth, Fl. Austr. IV, 136 (1869)	...	...	—	S.A.	T. V. N.S.W.	Q.	—	B.f.IV, 136	M.fr.X, 39.
WAHLENBERGIA, Schrader, cat. hort. bot. Goetting. (1814).									
W. gracilis, A. De Candolle, Monogr. comp. 142 (1830)	...	...	W.A.	S.A.	T. V. N.S.W.	Q. N.A.	—	B.f.IV, 137	

CANDOLLEACEAE.

F. v. M., fragm. VIII, 41 (1873). (Stylideae, R. Brown, prodr. 565, 1810.)

CANDOLLEA, Labillardière in Ann. du Mus. VI, 451 (1805). (Stylidium, Ventenata, Forsteropsis.)									
C. (St.) carnosa, Benth, Fl. Austr. III, 130 (1869)	...	...	W.A.	—	—	—	—	B.f.IV, 6	
C. (St.) pilosa, Labillardière, Nov. Holl. pl. spec. II, 63, t. 213 (1806)	W.A.	—	—	—	—	—	—	B.f.IV, 7	
C. (St.) reduplicata, R. Brown, prodr. 568 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 7	
C. (St.) scabrata, Lindley, Bot. Regist. XXV, App. XXVIII (1839)	W.A.	—	—	—	—	—	—	B.f.IV, 8	
C. (St.) hirsuta, R. Brown, prodr. 568 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 8	
C. (St.) crossocephala, F. v. M., fragm. VI, 5 (1866)	...	...	W.A.	—	—	—	—	B.f.IV, 8	M.fr.VI, 5, 254; X, 58.
C. (St.) juncea, R. Brown, prodr. 569 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 9	M.fr.X, 58.
C. (St.) guttata, R. Brown, prodr. 571 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 9	
C. (St.) repens, R. Brown, prodr. 571 (1810)	...	...	W.A.	—	—	—	—	B.f.IV, 9	
C. (St.) graminifolia, Swartz, Mag. der nat. Ges. Berl. 49 (1807)	...	...	—	S.A.	T. V. N.S.W.	Q.	—	B.f.IV, 10	M.fr.VI, 6; X, 58.
C. (St.) linearis, Swartz, Mag. der nat. Ges. Berl. 50 (1807)	...	...	—	—	—	N.S.W.	—	B.f.IV, 11	
C. (St.) elongata, Benth, Fl. Austr. IV, 11 (1869)	...	...	W.A.	—	—	—	—	B.f.IV, 11	M.fr.X, 86.
C. (St.) spinulosa, R. Brown, prodr. 569 (1815)	...	...	W.A.	—	—	—	—	B.f.IV, 11	

C. (St.) limbata, F. v. M., fragm. X, 57 (1876) ...	W.A.	—	—	—	—	—	—	—	M.fr.X,57.
C. (St.) caespitosa, R. Brown, prodr. 569 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,11	M.fr.X,57.
C. (St.) squamellosa, De Candolle, prodr. VII, 782 (1838) ...	W.A.	—	—	—	—	—	—	—	M.fr.X,56.
C. (St.) violacea, R. Brown, prodr. 569 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,12	
C. (St.) lutea, R. Brown, prodr. 570 (1810)...	W.A.	—	—	—	—	—	—	B.fl.IV,12	
C. (St.) filifera, R. Brown, prodr. 569 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,12	M.fr.X,58,86.
C. (St.) ciliata, Lindley, Bot. Regist. XXV, App. XXVIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,13	
C. (St.) sobolifera, F. v. M. in transact. Vict. Inst. 131 (1855) ...	—	—	—	V.	—	—	—	B.fl.IV,13	
C. (St.) Floodii, F. v. M., fragm. I, 149 (1859) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,13	M.fr.VI,6;X,58.
C. (St.) disperma, F. v. M., fragm. IV, 93 (1864)...	W.A.	—	—	—	—	—	—	B.fl.IV,14	M.fr.IV,93.
C. (St.) calcarata, R. Brown, prodr. 570 (1810) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,14	M.fr.VI,6.
C. (St.) perpusilla, J. Hooker in Lond. Journ. of Bot. VI, 266 (1847) ...	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,15	M.fr.VI,78;X,58.
C. (St.) eriorrhiza, R. Brown, prodr. 569 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,15	M.fr.I,147;VI,6;X,58.
C. (St.) debilis, F. v. M., fragm. I, 149 (1859) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,15	M.fr.VI,6;X,58.
C. (St.) floribunda, R. Brown, prodr. 569 (1810) ...	—	S.A.	—	—	—	—	N.A.	B.fl.IV,15	M.fr.I,148;VI,6;X,58.
C. (St.) leptorrhiza, F. v. M., fragm. I, 148 (1859) ...	—	—	—	—	—	—	N.A.	B.fl.IV,16	M.fr.I,148.
C. (St.) assimilis, R. Brown, prodr. 569 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,16	M.fr.X,57.
C. (St.) rupestris, Sonder in Lehm. pl. Preiss. I, 375 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,16	
C. (St.) spatulata, R. Brown, prodr. 569 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) Barleei, F. v. M., fragm. VI, 5, t. 69 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.IV,17	M.fr.VI,5,249.
C. (St.) lineata, Sonder in Lehm. pl. Preiss. I, 376 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) glauca, Labillard., Nov. Holl. pl. spec. II, 64, t. 214 (1806) ...	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) amoena, R. Brown, prodr. 570 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,18	
C. (St.) striata, Lindley, Bot. Regist. XXV, App. XXVIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,18	
C. (St.) diversifolia, R. Brown, prodr. 570 (1810)...	W.A.	—	—	—	—	—	—	B.fl.IV,18	M.fr.X,58.
C. (St.) articulata, R. Brown, prodr. 570 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,19	
C. (St.) Brunoniana, Benth in Hueg. enum. 72 (1837)...	W.A.	—	—	—	—	—	—	B.fl.IV,19	
C. (St.) diuroides, Lindley, Bot. Regist. XXV, App. XXIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,20	
C. (St.) scandens, R. Brown, prodr. 570 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,20	M.fr.VI,78.
C. (St.) verticillata, F. v. M., fragm. IV, 94 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,20	M.fr.IV,94.
C. (St.) glandulosa, Salisbury, Parad. Lond. t. 77 (1807) ...	W.A.	—	—	—	—	—	—	B.fl.IV,20	
C. (St.) loricifolia, Richard in Pers. syn. II, 210 (1807) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,21	
C. (St.) Preissii, F. v. M., fragm. III, 122 (1863)...	W.A.	—	—	—	—	—	—	B.fl.IV,21	M.fr.III,122.
C. (St.) imbricata, Benth in Hueg. enum. 73 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,21	
C. (St.) adpressa, Benth in Fl. Austr. IV, 22 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,22	
C. (St.) despecta, R. Brown, prodr. 571 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,22	M.fr.VI,58.
C. (St.) utricularioides, Benth in Hueg. enum. 73 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) pygmaea, R. Brown, prodr. 571 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) longitubea, Benth in Hueg. enum. 73 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) diffusa, R. Brown, prodr. 571 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,23	
C. (St.) fissiloba, F. v. M., fragm. I, 154 (1859) ...	—	—	—	—	—	N.A.	—	B.fl.IV,24	M.fr.I,154.
C. (St.) trichopoda, F. v. M., fragm. X, 86 (1877) ...	—	—	—	—	—	Q.	—	B.fl.IV,24	M.fr.X,86.
C. (St.) alsinoides, R. Brown, prodr. 572 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,24	M.fr.I,151;VI,6;X,58.
C. (St.) tenerima, F. v. M., fragm. I, 150 (1859)...	—	—	—	—	—	N.A.	—	B.fl.IV,24	M.fr.I,150.
C. (St.) brachyphylla, Sonder in Lehm. pl. Preiss. I, 386 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,24	
C. (St.) capillaris, R. Brown, prodr. 570 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,25	
C. (St.) rotundifolia, R. Brown, prodr. 571 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,25	M.fr.I,151.
C. (St.) schizantha, F. v. M., fragm. I, 152 (1859) ...	—	—	—	—	—	N.A.	—	B.fl.IV,25	M.fr.I,152.
C. (St.) lobuliflora, F. v. M., fragm. I, 153 (1859)...	—	—	—	—	—	N.A.	—	B.fl.IV,25	M.fr.I,153.
C. (St.) uliginosa, Swartz in Mag. nat. Fr. Berl. 52, t. 2 (1807) ...	—	—	—	—	—	Q.	—	B.fl.IV,26	
C. (St.) pulchella, Sonder in Lehm. pl. Preiss. I, 381 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,26	
C. (St.) petiolaris, Sonder in Lehm. pl. Preiss. I, 382 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,26	
C. (St.) emarginata, Sonder in Lehm. pl. Preiss. I, 383 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) corymbosa, R. Brown, prodr. 571 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) lepidia, F. v. M. in Benth. Fl. Austr. IV, 27 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) streptocarpa, Sonder in Lehm. pl. Preiss. I, 385 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) uniflora, Sonder in Lehm. pl. Preiss. I, 381 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,28	M.fr.VI,6.
C. (St.) pedunculata, R. Brown, prodr. 571 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,28	M.fr.VI,91.
C. (St.) pachyrrhiza, F. v. M., fragm. I, 152 (1859) ...	—	—	—	—	—	N.A.	—	B.fl.IV,28	M.fr.I,152.
C. (St.) muscicola, F. v. M., fragm. I, 153 (1859)...	—	—	—	—	—	N.A.	—	B.fl.IV,29	M.fr.I,153.
C. (St.) crassifolia, R. Brown, prodr. 571 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,29	
C. (St.) pycnostachya, Lindley, Bot. Regist. XXV, App. XXIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,29	
C. (St.) pubigera, Sonder in Lehm. pl. Preiss. I, 383 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) canaliculata, Lindley, Bot. Regist. XXV, App. XXIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) leptophylla, De Candolle, prodr. VII, 783 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) dichotoma, De Candolle, prodr. VII, 783 (1838)...	W.A.	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) bulbifera, Benth in Hueg. enum. 73 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,31	
C. (St.) breviscapex, R. Brown, prodr. 572 (1810)...	W.A.	—	—	—	—	—	—	B.fl.IV,31	
C. (St.) eglandulosa, F. v. M., fragm. I, 150 (1859) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,31	M.fr.VI,6;X,58.
C. (St.) fasciculata, R. Brown, prodr. 572 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,32	
C. (St.) falcata, R. Brown, prodr. 572 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,32	
C. (St.) rhynchocarpa, Sonder in Lehm. pl. Preiss. I, 389 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,33	
C. (St.) adnata, R. Brown, prodr. 572 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,33	M.fr.X,58.
LEEWEENHOEKIA, R. Brown, prodr. 572 (1810). (Levenhookia, Coleostylis.)									
L. pusilla, R. Brown, prodr. 573 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,34	M.fr.X,58.
L. dubia, Sonder in Lehm. pl. Preiss. I, 392 (1845) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,34	M.fr.III,121;X,58.
L. Sonderi, F. v. M., fragm. I, 18 (1858) ...	—	—	—	—	V.	—	—	B.fl.IV,35	
L. pauciflora, Benth in Hueg. enum. 74 (1837)...	W.A.	—	—	—	—	—	—	B.fl.IV,35	M.fr.VI,77.
L. leptantha, Benth in Fl. Austr. IV, 35 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,35	
L. stipitata, F. v. M., fragm. IV, 94 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.
L. Preissii, F. v. M., fragm. IV, 94 (1864)...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.

- PHYLLACHNE, R. & G. Forster, char. gen. 115, t. 58 (1776). (Forstera, Helophyllum, Oreostylidium.)
 P. bellidifolia, F. v. M., fragm. VIII, 39 (1873) ... — — — — — T. — — — — — M.fr. X, 58.
 DONATIA, R. et G. Forster, char. gen. 9, t. 5 (1776).
 D. Novae Zealandiae, J. Hooker, Fl. N. Zeal. I, 81, t. 20 (1853) — — — — — T. — — — — — B.fl. II, 450 M.fr. VIII, 46; X, 58.

GOODENIACEAE.

R. Brown, prodr. 573 (1810).

- BRUNONIA, Smith in Transact. Linn. Soc. X, 366 (1809).
 B. australis, Smith in Transact. Linn. Soc. X, 367, t. 28 (1809) W.A. S.A. T. V. N.S.W. Q. — B.fl. IV, 121 M.fr. X, 111.
 DAMPIERA, R. Brown, prodr. 587 (1810). (Linschotenia.)
 D. luteifolia, F. v. M., fragm. X, 11 (1876) ... W.A. — — — — — — — — — — M.fr. X, 11.
 D. Linschotenii, F. v. M., fragm. VI, 28 (1866) ... — — — — — Q. — — — — — B.fl. IV, 108 M.fr. VI, 28.
 D. spicigera, Benth. Fl. Austr. IV, 109 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 109
 D. candicans, F. v. M., fragm. X, 86 (1876) ... W.A. S.A. — — — — — — — — — — M.fr. X, 86.
 D. teres, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 109
 D. trigona, De Vriese in Lehm. pl. Preiss. I, 401 (1845) ... W.A. — — — — — — — — — — B.fl. IV, 109 M.fr. II, 17; VIII, 245; X, 11
 D. prostrata, De Vriese, Goodenov. 83 (1854) ... W.A. — — — — — — — — — — B.fl. IV, 110
 D. alata, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 110 M.fr. X, 11.
 D. coronata, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 110
 D. carinata, Benth. Fl. Austr. IV, 111 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 111
 D. sacculata, F. v. M. in Benth. Fl. Austr. IV, 111 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 111
 D. incana, R. Brown, prodr. 588 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 111 M.fr. II, 17.
 D. hederacea, R. Brown, prodr. 588 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 112 M.fr. X, 12.
 D. ferruginea, R. Brown, prodr. 588 (1810) ... — — — — — — — — — — Q. — — — — — B.fl. IV, 112 M.fr. VI, 29.
 D. Brownii, F. v. M., fragm. VI, 29 (1866) ... — — — — — V. N.S.W. — — — — — B.fl. IV, 112 M.fr. VI, 29.
 D. lanceolata, Cunningham in De Cand., prodr. VII, 503 (1839) ... S.A. — — — — — V. N.S.W. — — — — — B.fl. IV, 113 M.fr. VI, 29.
 D. altissima, F. v. M. in Benth. Fl. Austr. IV, 113 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 113
 D. marifolia, Benth. Fl. Austr. IV, 114 (1869) ... — — — — — S.A. — — — — — V. N.S.W. — — — — — B.fl. IV, 114
 D. rosmarinifolia, Schlechtendal, Linnaea XX, 603 (1847) ... — — — — — S.A. — — — — — V. N.S.W. — — — — — B.fl. IV, 114 M.fr. VI, 29; X, 12.
 D. lavandulacea, Lindley, Bot. Regist. XXV, App. XXVII (1839) W.A. — — — — — — — — — — B.fl. IV, 114
 D. juncea, Benth. Fl. Austr. IV, 115 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 115
 D. oligophylla, Benth. Fl. Austr. IV, 115 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 115
 D. loranthifolia, F. v. M. in Benth. Fl. Austr. IV, 115 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 115
 D. stricta, R. Brown, prodr. 589 (1810) ... — — — — — S.A. T. V. N.S.W. — — — — — B.fl. IV, 116 M.fr. X, 12; XI, 121.
 D. Scottiana, F. v. M., fragm. XI, 120 (1881) ... — — — — — — — — — — N.S.W. — — — — — M.fr. XI, 120.
 D. leptoclada, Benth. Fl. Austr. IV, 116 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 116
 D. fasciculata, R. Brown, prodr. 588 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 116 M.fr. XI, 121.
 D. subspicata, Benth. Fl. Austr. IV, 117 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 117
 D. trilobata, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 117
 D. linearis, R. Brown, prodr. 588 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 117
 D. cuneata, R. Brown, prodr. 588 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 118
 D. sericantha, F. v. M. in Benth. Fl. Austr. IV, 118 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 118
 D. parvifolia, R. Brown, prodr. 589 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 118
 D. glabrescens, Benth. Fl. Austr. IV, 119 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 119
 D. adpressa, Cunningham in De Cand., prodr. VII, 503 (1839) ... — — — — — N.S.W. Q. — — — — — B.fl. IV, 119
 D. diversifolia, De Vriese in Lehm. pl. Preiss. I, 403 (1845) ... W.A. — — — — — — — — — — B.fl. IV, 119
 D. eriocephala, De Vriese, Goodenov. 118, t. 21 (1854) ... W.A. — — — — — — — — — — B.fl. IV, 120 M.fr. VI, 29; X, 12.
 D. Wellsiana, F. v. M., fragm. X, 12 (1876) ... W.A. — — — — — — — — — — — — — — — — M.fr. X, 12.
 DIASPASIS, R. Brown, prodr. 586 (1810).
 D. filifolia, R. Brown, prodr. 587 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 104 M.fr. I, 206; II, 181; VIII, 58
 LESCHENAULTIA, R. Brown, prodr. 581 (1810). (Latourea.)
 L. formosa, R. Brown, prodr. 581 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 40 M.fr. VI, 226.
 L. oblata, Sweet, flor. Australas. 26, 46, t. 46 (1828) ... W.A. — — — — — — — — — — — — — — — — M.fr. VI, 26.
 L. chlorantha, F. v. M., fragm. II, 20 (1860) ... W.A. — — — — — — — — — — B.fl. IV, 40 M.fr. II, 20.
 L. linarioides, De Candolle, prodr. VII, 519 (1818) ... W.A. — — — — — — — — — — B.fl. IV, 40
 L. tubiflora, R. Brown, prodr. 581 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 41
 L. superba, F. v. M., fragm. VI, 10 (1866) ... W.A. — — — — — — — — — — B.fl. IV, 41 M.fr. VI, 10.
 L. striata, F. v. M., fragm. VIII, 245 (1874) ... — — — — — S.A. — — — — — — — — — — M.fr. VIII, 245.
 L. acutiloba, Benth. Fl. Austr. IV, 41 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 41
 L. larinica, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 41
 L. hirsuta, F. v. M., fragm. VI, 9 (1866) ... W.A. — — — — — — — — — — B.fl. IV, 42 M.fr. VI, 9.
 L. longiloba, F. v. M., fragm. VI, 10 (1866) ... W.A. — — — — — — — — — — B.fl. IV, 42 M.fr. VI, 10.
 L. biloba, Lindley, Bot. Regist. XXV, App. XXVII (1839) ... W.A. — — — — — — — — — — B.fl. IV, 42
 L. expansa, R. Brown, prodr. 581 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 42
 L. floribunda, Benth. in Hueg. enum. 70 (1837) ... W.A. — — — — — — — — — — B.fl. IV, 43
 L. heteromera, Benth. Fl. Austr. IV, 43 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 43
 L. divaricata, F. v. M., fragm. III, 33, 167 (1863) ... — — — — — S.A. — — — — — — — — — — B.fl. IV, 43 M.fr. VIII, 246.
 L. filiformis, R. Brown, prodr. 581 (1810) ... — — — — — — — — — — Q. N.A. B.fl. IV, 44 M.fr. VI, 9, 249.
 L. agrostophylla, F. v. M., fragm. VI, 8, pl. XLVII (1866) ... — — — — — — — — — — N.A. B.fl. IV, 44 M.fr. VI, 8; 249.
 ANTHOTIUM, R. Brown, prodr. 582 (1810).
 A. humile, R. Brown, prodr. 582 (1810) ... W.A. — — — — — — — — — — B.fl. IV, 44 M.fr. VIII, 58.
 A. rubriflorum, F. v. M. in Benth. Fl. Austr. IV, 45 (1869) ... W.A. — — — — — — — — — — B.fl. IV, 45
 CATOSPERMA, Benth. in Hooker, icon. pl. t. 1028 (1868).
 C. Muellieri, Benth. in Hook. icon. pl. t. 1028 (1868) ... — — — — — — — — — — N.A. B.fl. IV, 83 M.fr. I, 121; VIII, 58.
 SCAEVOLA, Linné, mantiss. II, 145 (1771). (Pogonetes, Crossotoma, Temminckia, Kamphusia, Molkenboeria, Merkusia, Verreauxia.)
 S. Koenigii, Vahl, symb. bot. III, 36 (1794) ... — — — — — — — — — — Q. N.A. B.fl. IV, 86 M.fr. XI, 76.
 S. enantophylla, F. v. M., fragm. VIII, 58 (1873) ... — — — — — — — — — — Q. — — — — — M.fr. VI, 225.

S. spinescens, R. Brown, prodr. 586 (1810)...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 87	M.fr. XI, 20, 27, 76.
S. Groeneri, F. v. M., fragm. VI, 15 (1866)...	...	...	W.A.	S.A.	—	—	—	—	—	B.f.IV, 88	M.fr. VIII, 58.
S. tomentosa, Gaudichaud in Freyc. Voy. Bot. 460, t. 81 (1826)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 88	
S. atriplicina, F. v. M., fragm. II, 18 (1860)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 88	M.fr. III, 181.
S. striata, R. Brown, prodr. 586 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 89	
S. phlebopetala, F. v. M., fragm. II, 18 (1860)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 89	M.fr. II, VI, 16; XI, 76.
S. pilosa, Benth. in Hueg. enum. 69 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 89	
S. hispida, Cavanilles, icon. et descr. pl. VI, 7, t. 510 (1801)...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.f.IV, 90	M.fr. XI, 76.
S. apterantha, F. v. M., fragm. I, 121 (1859)...	...	...	—	—	—	V.	—	—	—	B.f.IV, 90	M.fr. I, 121.
S. Hookeri, F. v. M. in Benth. Fl. Austr. IV, 90 (1869)...	...	...	—	—	—	T.	V.	N.S.W.	—	B.f.IV, 90	M.fr. XI, 76.
S. parvifolia, F. v. M. in Benth. Fl. Austr. IV, 91 (1869)...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.f.IV, 91	M.fr. VIII, 58; X, 59; XI, 76.
S. oxyclona, F. v. M., fragm. X, 58 (1876)...	...	...	W.A.	—	—	—	—	—	—	—	M.fr. X, 58.
S. restiacea, Benth. Fl. Austr. IV, 91 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 91	
S. depauperata, R. Brown, App. Sturt's Exped. 20 (1849)...	...	...	—	S.A.	—	—	—	—	—	B.f.IV, 91	M.fr. III, 33; VIII, 58; XI, 76.
S. tortuosa, Benth. Fl. Austr. IV, 91 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 91	
S. Cunninghamii, De Candolle, prodr. VII, 508 (1838)...	...	...	—	—	—	—	—	—	N.A.	B.f.IV, 92	
S. collaris, F. v. M., Rep. Babb. Exped. 15 (1858)...	...	...	—	S.A.	—	—	—	—	—	B.f.IV, 92	M.fr. VIII, 58; XI, 76.
S. angulata, R. Brown, prodr. 586 (1810)...	...	...	—	—	—	—	—	—	N.A.	B.f.IV, 92	
S. nitida, R. Brown, prodr. 584 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 93	
S. globuliflora, Labillardière, Nov. Holl. pl. spec. I, 55, t. 78 (1804)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 93	
S. porocarya, F. v. M., fragm. II, 19 (1860)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 94	M.fr. II, 19; XI, 76.
S. attenuata, R. Brown, prodr. 583 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 94	
S. glandulifera, De Candolle, prodr. VII, 510 (1838)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 94	
S. anchusifolia, Benth. in Hueg. enum. 68 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 94	
S. holosericea, De Vriese in Lehm. pl. Preiss. I, 408 (1845)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 95	
S. suaveolens, R. Brown, prodr. 585 (1810)...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.IV, 95	
S. revoluta, R. Brown, prodr. 586 (1810)...	...	...	—	—	—	—	—	—	N.A.	B.f.IV, 96	
S. ovalifolia, R. Brown, prodr. 584 (1810)...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.IV, 96	M.fr. VIII, 58; XI, 76.
S. crassifolia, Labillardière, Nov. Holl. pl. spec. I, 56, t. 79 (1804)...	...	...	W.A.	S.A.	—	V.	—	—	—	B.f.IV, 96	M.fr. VIII, 245; XI, 20, 76.
S. macrostachya, Benth. Fl. Austr. IV, 97 (1869)...	...	...	—	—	—	—	—	—	N.A.	B.f.IV, 97	M.fr. VIII, 58; XI, 76.
S. longifolia, De Vriese in Lehm. pl. Preiss. I, 410 (1845)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 97	
S. lanceolata, Benth. in Hueg. enum. 69 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 97	M.fr. I, 207; XI, 76.
S. thesioides, Benth. in Hueg. enum. 68 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 98	M.fr. XI, 77.
S. macrophylla, Benth. Fl. Austr. IV, 98 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 98	
S. platyphylla, Lindley, Bot. Regist. XXV, App. XXVI (1839)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 98	M.fr. XI, 77.
S. auriculata, Benth. Fl. Austr. IV, 99 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 99	M.fr. XI, 77.
S. aemula, R. Brown, prodr. 584 (1810)...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 99	M.fr. XI, 77.
S. humilis, R. Brown, prodr. 585 (1810)...	...	...	—	S.A.	—	—	—	—	—	B.f.IV, 100	
S. amblyanthera, F. v. M., fragm. I, 121 (1859)...	...	...	—	—	—	—	—	—	N.A.	B.f.IV, 100	M.fr. I, 121.
S. microphylla, Benth. Fl. Austr. IV, 100 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 100	
S. cuneiformis, Labillard., Nov. Holl. pl. spec. I, 56, t. 80 (1804)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 101	M.fr. XI, 77.
S. microcarpa, Cavanilles, icon. et descr. pl. VI, 6, t. 509 (1801)...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 101	M.fr. VI, 16; XI, 77.
S. linearis, R. Brown, prodr. 586 (1810)...	...	...	—	S.A.	—	—	—	—	—	B.f.IV, 102	
S. Oldfieldii, F. v. M., fragm. II, 19 (1860)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 102	M.fr. II, 19; XI, 77.
S. paludosa, R. Brown, prodr. 586 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 102	M.fr. XI, 77.
S. sericophylla, F. v. M. in Benth. Fl. Austr. IV, 102 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 102	M.fr. XI, 77.
S. canescens, Benth. in Hueg. enum. 69 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 103	
S. humifusa, De Vriese in Lehm. pl. Preiss. I, 410 (1845)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 103	M.fr. XI, 77.
S. fasciculata, Benth. in Hueg. enum. 68 (1837)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 104	M.fr. XI, 77.
S. stenophylla, Benth. Fl. Austr. IV, 104 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 104	M.fr. I, 113; XI, 77.
S. Reinwardtii, De Vriese in Lehm. pl. Preiss. I, 409 (1845)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 106	
S. Verreauxii, F. v. M., Bot. Teachings, 65 (1877)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 105	
SELLIERA, Cavanilles in Anal. Cien. Nat. I, 41, t. 5 (1799).											
S. radicans, Cavanilles, icon. et descr. pl. V, 49, t. 474 (1799)...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 82	M.fr. VIII, 58.
S. exigua, F. v. M., fragm. III, 142 (1862)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 82	M.fr. III, 142.
CALOZYNE, R. Brown, prodr. 579 (1810). (Distylis.)											
C. pilosa, R. Brown, prodr. 579 (1810)...	...	...	—	—	—	—	—	Q.	N.A.	B.f.IV, 81	M.fr. VI, 7, 28.
C. Berardiana, F. v. M., fragm. VI, 7 (1867)...	...	...	W.A.	—	—	—	—	—	N.A.	B.f.IV, 81	M.fr. VI, 7; VIII, 58.
C. heteroptera, F. v. Mueller, fragm. X, 43 (1876)...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr. X, 43.
C. purpurea, F. v. M., fragm. VIII, 57 (1873)...	...	...	—	—	—	—	—	—	N.A.	—	M.fr. VIII, 57.
GOODENIA, Smith in Transact. Linn. Soc. II, 347 (1794). (Goodenoughia, Tetraphylax, Picrophyta, Stekhovia, Aillya.)											
G. phylloides, F. v. M., fragm. I, 206 (1859)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 55	M.fr. I, 206.
G. viscida, R. Brown, prodr. 578 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 55	M.fr. I, 120; III, 35.
G. xanthotricha, De Vriese, Gooden. 155 (1854)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 56	M.fr. VI, 13.
G. scapigera, R. Brown, prodr. 578 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 56	M.fr. I, 114; III, 163; XI, 50.
G. quadrilocularis, R. Brown, prodr. 578 (1810)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 57	M.fr. III, 141; IV, 13.
G. Ramelii, F. v. M., fragm. III, 20, t. 17 (1862)...	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.f.IV, 57	M.fr. VIII, 245; X, 13.
G. pinifolia, De Vriese, Gooden. 157, t. 30 (1854)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 58	
G. decurrens, R. Brown, prodr. 575 (1810)...	...	...	—	—	—	—	N.S.W.	—	—	B.f.IV, 58	
G. racemosa, F. v. M., fragm. I, 114 (1859)...	...	...	—	—	—	—	—	Q.	—	B.f.IV, 58	M.fr. I, 114.
G. bellidifolia, Smith in Transact. Linn. Soc. II, 349 (1794)...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 58	M.fr. XI, 50.
G. stelligera, R. Brown, prodr. 575 (1810)...	...	...	—	—	—	—	V.	N.S.W.	Q.	B.f.IV, 59	M.fr. VI, 14.
G. ovata, Smith in Transact. Linn. Soc. II, 347 (1794)...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 59	M.fr. X, 110.
G. amplexans, F. v. M. in Transact. phil. Inst. Vict. II, 70 (1859)...	...	...	—	S.A.	—	V.	—	—	—	B.f.IV, 60	M.fr. I, 15.
G. strophiolata, F. v. M., fragm. I, 119 (1859)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 60	M.fr. I, 119.
G. varia, R. Brown, prodr. 576 (1810)...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 61	M.fr. I, 205.
G. laevis, Benth. Fl. Austr. IV, 61 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 61	
G. barbata, R. Brown, prodr. 576 (1810)...	...	...	—	—	—	T.	V.	N.S.W.	—	B.f.IV, 61	M.fr. II, 110, 176; VI, 15.
G. arthrotricha, F. v. M. in Benth. Fl. Austr. IV, 62 (1869)...	...	...	W.A.	—	—	—	—	—	—	B.f.IV, 62	
G. disperma, F. v. M., fragm. I, 113 (1859)...	...	...	—	—	—	—	—	Q.	—	B.f.IV, 62	M.fr. IV, 145; VI, 15.

G. geniculata, R. Brown, prodr. 577 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV,62	M.fr.VI,14.	
G. hederacea, Smith in Transact. Linn. Soc. II, 349 (1794)	...	...	...	—	—	—	—	V.	N.S.W.	Q.	—	M.fr.VI,15,110.	
G. xanthosperma, F. v. M., fragm. X, 12 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,12.	
G. hirsuta, F. v. M., fragm. III, 35 (1862)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.f.IV,64	M.fr.III,35.	
G. heterophylla, Smith in Transact. Linn. Soc. II, 349 (1794)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,64	—	
G. glabra, R. Brown, prodr. 577 (1810)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,64	M.fr.VI,14.	
G. stronglyphylla, F. v. M., fragm. VI, 12 (1867)	...	...	...	—	—	—	—	—	Q.	—	B.f.IV,65	M.fr.VI,12.	
G. rotundifolia, R. Brown, prodr. 576 (1810)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,65	M.fr.VI,15.	
G. azurea, F. v. M., fragm. I, 117 (1859)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,66	M.fr.I,117.	
G. scaevolina, F. v. M., fragm. I, 118 (1859)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,66	M.fr.I,118.	
G. Stobbsiana, F. v. M., fragm. XI, 49 (1879)	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.XI,49,137.	
G. incana, R. Brown, prodr. 578 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,66	M.fr.I,155;III,141. [50.	
G. leptoclada, Benth. Fl. Austr. IV, 67 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,67	M.fr.VI,227,VIII,186;XI,	
G. Eatoniana, F. v. M., fragm. VIII, 186 (1874)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.XI,51.	
G. coerulea, R. Brown, prodr. 578 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,67	M.fr.III,140;XI,20.	
G. trichophylla, De Vriese in Benth. Fl. Austr. IV, 67 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,67	—	
G. Hassallii, F. v. M., fragm. VI, 10, t. 51 (1867)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,68	M.fr.XI,50.	
G. pterygosperma, R. Brown, prodr. 578 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,68	M.fr.I,155;III,163.	
G. Vilmoriniae, F. v. M., fragm. III, 19, t. 16 (1862)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.f.IV,68	M.fr.III,19;VIII,245.	
G. Bonneyana, F. v. M. in Benth. Fl. Austr. IV, 69 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,69	M.fr.VI,226,250.	
G. calcarata, F. v. M., fragm. VI, 14 (1867)	...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.f.IV,69	M.fr.VIII,245.	
G. albiflora, Schlechtendal, Linnaea XX, 599 (1847)	...	...	...	—	S.A.	—	—	—	—	—	B.f.IV,70	—	
G. Nicholsoni, F. v. M., fragm. I, 203, t. 4 (1859)	...	...	...	—	S.A.	—	—	—	—	—	B.f.IV,69	M.fr.I,203.	
G. Macmillani, F. v. M., fragm. I, 119, t. 5 (1859)	...	...	...	—	—	—	—	V.	—	—	B.f.IV,69	M.fr.I,119.	
G. grandiflora, Sims, Bot. Mag. t. 890 (1806)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,69	M.fr.I,204;VI,14.	
G. Chambersii, F. v. M., fragm. I, 204 (1859)	...	...	...	—	S.A.	—	—	—	—	—	B.f.IV,70	M.fr.I,204.	
G. Strangfordii, F. v. M., fragm. VI, 11, t. 52 (1867)	...	...	...	—	—	—	—	—	Q.	N.A.	B.f.IV,70	—	
G. Mitchellii, Benth. Fl. Austr. IV, 71 (1869)	...	...	...	—	S.A.	—	—	—	Q.	—	B.f.IV,71	M.fr.XI,75.	
G. Mueckeana, F. v. M., fragm. VIII, 56 (1873)	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.IX,194;XI,133.	
G. melanoptera, F. v. M., fragm. I, 115 (1859)	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.I,115.	
G. heterochila, F. v. M., fragm. III, 142 (1863)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.f.IV,71	M.fr.VIII,245.	
G. sepalosa, F. v. M. in Benth. Fl. Austr. IV, 72 (1869)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,72	—	
G. hispida, R. Brown, prodr. 577 (1810)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,72	M.fr.VI,12.	
G. auriculata, Benth. Fl. Austr. IV, 72 (1869)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,72	—	
G. Armstrongiana, De Vriese, Goodenov. 138, t. 24 (1854)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,73	M.fr.I,205;VII,40.	
G. corynocarpa, F. v. M., fragm. II, 16 (1860)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,73	M.fr.II,16;VI,15.	
G. mollissima, F. v. M. in Benth. Fl. Austr. IV, 73 (1869)	...	...	...	—	—	—	—	—	Q.	—	B.f.IV,73	—	
G. cycloptera, R. Brown in App. Sturt's Exped. 20 (1849)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV,74	M.fr.VI,14;XI,51.	
G. tenella, R. Brown, prodr. 577 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,74	M.fr.XI,51. [X,110.	
G. elongata, Labillardiere, Nov. Holl. pl. spec. I, 52, t. 75 (1804)	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.IV,74	M.fr.VI,14;VIII,142,245;	
G. pinnatifida, Schlechtendal, Linnaea XXI, 440 (1848)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV,75	M.fr.VI,14;XI,51.	
G. coronopifolia, R. Brown, prodr. 577 (1810)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,75	—	
G. heteromera, F. v. M., fragm. I, 115 (1859)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV,76	M.fr.VI,14;XI,27.	
G. concinna, Benth. Fl. Austr. IV, 76 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,76	—	
G. glauca, F. v. M. in Transact. Vict. Inst. 40 (1855)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV,76	M.fr.VI,14;XI,27.	
G. filiformis, R. Brown, prodr. 578 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,77	M.fr.VI,14;VIII,245.	
G. Armitiana, F. v. M., fragm. X, 110 (1877)	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.X,110.	
G. microptera, F. v. M., fragm. III, 34 (1862)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.f.IV,77	M.fr.VI,227;XI,75.	
G. bicolor, F. v. M. in Benth. Fl. Austr. IV, 80 (1869)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,80	[245;X,110.	
G. humilis, R. Brown, prodr. 575 (1810)	...	...	...	—	—	—	—	—	—	—	B.f.IV,79	M.fr.VII,227;VIII,142,	
G. paniculata, Smith in Transact. Linn. Soc. II, 348 (1794)	...	...	...	—	—	—	—	V.	N.S.W.	Q.	B.f.IV,78	M.fr.I,116.	
G. purpurascens, R. Brown, prodr. 578 (1810)	...	...	...	—	—	—	—	—	Q.	N.A.	B.f.IV,78	M.fr.I,117.	
G. gracilis, R. Brown, prodr. 575 (1810)	...	...	...	—	—	—	—	V.	N.S.W.	Q.	N.A.	B.f.IV,79	M.fr.XI,51.
G. lamprosperma, F. v. M., fragm. I, 116 (1859)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,79	M.fr.I,116.	
G. minutiflora, F. v. M., fragm. VIII, 244 (1874)	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.VIII,244.	
G. claytoniacea, F. v. M. in Benth. Fl. Austr. IV, 79 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,79	M.fr.II,111.	
G. Pumilio, R. Brown, prodr. 579 (1810)	...	...	...	—	—	—	—	—	Q.	N.A.	B.f.IV,80	M.fr.X,111.	

VELLEYA, Smith in Transact. Linn. Soc. IV, 217 (1798).	(Velleia, Euthales.)											
V. Daviesii, F. v. M., fragm. X, 10 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,10.
V. panduriformis, Cunningham in Benth. Fl. Austr. IV, 46 (1869)	...	...	...	—	—	—	—	—	—	N.A.	B.f.IV,46	M.fr.XI,73.
V. connata, F. v. M. in Transact. phil. Soc. Vict. I, 18 (1855)	...	...	...	—	S.A.	—	V.	N.S.W.	—	N.A.	B.f.IV,46	M.fr.VI,227;VIII,246.
V. perfoliata, R. Brown, prodr. 581 (1810)	...	...	...	—	—	—	—	N.S.W.	—	—	B.f.IV,46	
V. discophora, F. v. M., fragm. X, 10 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,10.
V. trinervis, Labillardiere, Nov. Holl. pl. spec. I, 54, t. 77 (1804)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,47	M.fr.XI,75.
V. macrophylla, Benth. Fl. Austr. IV, 47 (1869)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,47	M.fr.XI,75.
V. paradoxa, R. Brown, prodr. 580 (1810)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV,48	M.fr.VI,11;XI,75.
V. cynopotaamica, F. v. M., fragm. VI, 7 (1867)	...	...	...	W.A.	—	—	—	—	—	—	B.f.IV,48	M.fr.VI,249;X,10.
V. lyrata, R. Brown, prodr. 580 (1810)	...	...	...	—	—	—	—	N.S.W.	—	—	B.f.IV,49	
V. macrocalyx, De Vriese, Goodenov. 176, t. 34 (1854)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,49	M.fr.X,75.
V. spathulata, R. Brown, prodr. 580 (1810)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.f.IV,50	M.fr.XI,75.
V. pubescens, R. Brown, prodr. 581 (1810)	...	...	...	—	—	—	—	—	Q.	—	B.f.IV,49	M.fr.VI,11;XI,75.
V. montana, J. Hooker in Lond. Journ. of Bot. VI, 265 (1847)	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.f.IV,50	

SYNPETALEAE HYPOGYNAE.

F. v. M. in Woolls, pl. of the neighb. of Sydney 34 (1880).

GENTIANEAEE.

Necker in Act. Acad. Theod. Pal. II, 477 (1770), from B. de Jussieu (1759).

LIMNANTHEMUM, Gmelin, Nov. Comm. Acad. Petrop. XIV, 257 (1770). (Villarsia, Liparophyllum.)

L. Indicum, Thwaites, enum. pl. Ceyl. 205 (1862) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV, 378	M.fr. VI, 136.
L. Moonii, Thwaites, enum. pl. Ceyl. 205 (1862) ...	—	—	—	—	—	Q.	—	—	M.fr. IX, 163.
L. minimum, F. v. M., fragm. I, 40 (1858) ...	—	—	—	—	—	—	N.A.	B.f.IV, 379	M.fr. VI, 137. [165]
L. crenatum, F. v. M. in Transact. phil. Soc. Vict. I, 17 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 379	M.fr. IV, 127; VI, 137; IX,
L. geminatum, Grisebach, gen. et spec. Gent. 346 (1839) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 380	M.fr. VI, 137, 254.
L. hydrocharoides, F. v. M. in Benth. Fl. Austr. IV, 380 (1869) ...	—	—	—	—	—	Q.	—	B.f.IV, 380	M.fr. VI, 139; IX, 165.
L. exiliflorum, F. v. M., fragm. V, 46 (1865) ...	—	—	—	—	—	Q.	—	B.f.IV, 381	M.fr. VI, 137.
L. exiguum, F. v. M., fragm. I, 40 (1858) ...	—	—	T.	—	—	—	—	B.f.IV, 381	M.fr. VI, 137.
L. Gunnii, J. Hooker, Fl. Tasman. II, 368 (1860) ...	—	—	T.	—	—	—	—	B.f.IV, 382	M.fr. VI, 137; IX, 164.
L. capitatum, F. v. M., fragm. IX, 164 (1875) ...	W.A.	—	—	—	—	—	—	B.f.IV, 375	M.fr. VI, 142.
L. calthifolium, F. v. M., fragm. IX, 164 (1875) ...	W.A.	—	—	—	—	—	—	B.f.IV, 374	M.fr. VI, 140.
L. congestiflorum; Villarsia, F. v. M., fragm. VI, 141 (1868) ...	W.A.	—	—	—	—	—	—	B.f.IV, 375	M.fr. VI, 141.
L. latifolium, F. v. M., fragm. IX, 164 (1875) ...	W.A.	—	—	—	—	—	—	B.f.IV, 375	M.fr. VI, 141.
L. exaltatum, F. v. M., fragm. IX, 165 (1875) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 376	M.fr. II, 21; VI, 139.
L. parnassifolium, F. v. M., fragm. IX, 165 (1875) ...	W.A.	—	—	—	—	—	—	B.f.IV, 376	M.fr. VI, 140.
L. violifolium; Villarsia, F. v. M., fragm. VI, 138 (1868) ...	W.A.	—	—	—	—	—	—	B.f.IV, 377	M.fr. VI, 138.
L. lasiospermum, F. v. M., fragm. IX, 165 (1875) ...	W.A.	—	—	—	—	—	—	B.f.IV, 377	M.fr. VI, 137.
L. albiflorum; Villarsia, F. v. M., fragm. II, 21 (1860) ...	W.A.	—	—	—	—	—	—	B.f.IV, 377	M.fr. VI, 138.

SEBAEA, Solander in R. Brown, prodr. 451 (1810).

S. ovata, R. Brown, prodr. 452 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 371	M.fr. VII, 136; IX, 165.
S. albidiflora, F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854) ...	—	S.A.	T.	V.	—	—	—	B.f.IV, 371	M.fr. VI, 136.

ERYTHRAEA, Reneaulme, specim. hist. plant. 77, t. 76 (1611).

E. australis, R. Brown, prodr. 451 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.IV, 371	M.fr. VI, 136; IX, 165; XI, [20]
---	------	------	----	----	--------	----	------	-------------	----------------------------------

CANSCORA, Lamarck, Encycl. method. I, 602 (1783). (Orthostemon.)

C. diffusa, R. Brown, prodr. 451 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 372	M.fr. VI, 136; IX, 165.
---	---	---	---	---	---	----	------	-------------	-------------------------

GENTIANA, Tournefort, inst. 80, t. 40 (1700), from Dioscorides.

G. saxosa, Forster in Svensk. Kongl. Vet. Ac. Handl. 183, t. 5 (1777) ...	S.A.	T.	V.	N.S.W.	—	—	—	B.f.IV, 373	M.fr. VI, 136, 255; IX, 165.
---	------	----	----	--------	---	---	---	-------------	------------------------------

LOGANIACEAE.

R. Brown in App. Flind. Voy. 564 (1814).

STRYCHNOS, Linné, syst. nat. 8 (1735); Linné, spec. plant. 189 (1753).

S. lucida, R. Brown, prodr. 469 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 369	—
S. psiloperma, F. v. M., fragm. IV, 44 (1863) ...	—	—	—	—	—	Q.	—	B.f.IV, 369	M.fr. VI, 131.

FAGRAEA, Thunberg in Kongl. Svensk. Vetensk. Acad. Handl. 125 (1782).

F. racemosa, Jack in Roxb. Fl. Ind. ed. Carey. II, 35 (1824) ...	—	—	—	—	—	—	N.A.	B.f.IV, 367	M.fr. II, 137.
F. Muellieri, Benth. Fl. Austr. IV, 368 (1869) ...	—	—	—	—	—	Q.	—	B.f.IV, 368	—

MITREOLA, Linné, hort. Cliffort. 492 (1737).

M. oldenlandioides, Wallich, numeric. list 4350 (1828) ...	—	—	—	—	—	—	N.A.	B.f.IV, 349	M.fr. VI, 131.
--	---	---	---	---	---	---	------	-------------	----------------

MITRASACME, Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804). (Mitragyne.)

M. Archeri, J. Hooker, Fl. Tasman. II, 368 (1860) ...	—	—	T.	—	—	—	—	B.f.IV, 351	—
M. montana, J. Hooker, Fl. Tasman. I, 274, t. 88 (1859) ...	—	—	T.	V.	—	—	—	B.f.IV, 351	M.fr. VI, 131.
M. serpillifolia, R. Brown, prodr. 454 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f.IV, 352	—
M. paludosa, R. Brown, prodr. 453 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 352	M.fr. I, 134.
M. pilosa, Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804) ...	—	—	T.	V.	N.S.W.	—	—	B.f.IV, 352	M.fr. VI, 131.
M. alsinoides, R. Brown, prodr. 453 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 353	—
M. polymorpha, R. Brown, prodr. 452 (1810) ...	—	—	—	—	V.	N.S.W.	Q.	B.f.IV, 353	M.fr. VI, 131.
M. longiflora, F. v. M. in Benth. Fl. Austr. IV, 353 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 353	—
M. elata, R. Brown, prodr. 453 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 354	M.fr. I, 132.
M. exserta, F. v. M., fragm. I, 131 (1859) ...	—	—	—	—	—	—	N.A.	B.f.IV, 354	M.fr. I, 131.
M. tenuiflora, Benth. Fl. Austr. IV, 354 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 354	—
M. ambigua, R. Brown, prodr. 454 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 354	—
M. nudicaulis, Reinwardt in Blume, Bijdr. 849 (1826) ...	—	—	—	—	—	—	N.A.	B.f.IV, 355	—
M. connata, R. Brown, prodr. 454 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 355	M.fr. I, 131.
M. laevis, Benth. in Journ. Linn. Soc. I, 93 (1857) ...	—	—	—	—	—	—	N.A.	B.f.IV, 355	M.fr. I, 133.
M. Indica, Wight, icon. pl. Ind. or. IV, t. 1601 (1850) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV, 356	—
M. stellata, R. Brown, prodr. 454 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 356	—
M. Cunninghamii, Benth. Fl. Austr. IV, 357 (1869) ...	—	—	—	—	—	Q.	—	B.f.IV, 357	—
M. pygmaea, R. Brown, prodr. 453 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 357	—
M. lutea, F. v. M., fragm. I, 133 (1859) ...	—	—	—	—	—	—	N.A.	B.f.IV, 357	M.fr. IV, 170.
M. multicaulis, R. Brown, prodr. 453 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 357	—
M. loricifolia, R. Brown, prodr. 453 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 358	M.fr. VI, 131.
M. prolifera, R. Brown, prodr. 453 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 358	—
M. gentiana, F. v. M., fragm. I, 130 (1859) ...	—	—	—	—	—	—	N.A.	B.f.IV, 358	M.fr. I, 130.
M. phascoides, R. Brown, prodr. 454 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 359	—
M. paradoxa, R. Brown, prodr. 454 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 359	M.fr. VI, 131.
M. distylis, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ...	—	—	T.	V.	—	—	—	B.f.IV, 359	M.fr. VI, 131.

GENIOSTOMA, R. & G. Forster, char. gen. 23, t. 12 (1776).

G. Australianum, F. v. M., fragm. V, 19 (1865) ...	—	—	—	—	—	Q.	—	B.f.IV, 367	M.fr. V, 19.
G. petiolosum, Moore & F. v. M., fragm. VII, 28 (1869) ...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 193.

LOGANIA, R. Brown, prodr. 454 (1810). (Euosma 1808, restorable.)

L. longifolia, R. Brown, prodr. 456 (1810) ...	W.A. S.A.	—	—	—	—	—	B.fl. IV, 361	
L. latifolia, R. Brown, prodr. 455 (1810) ...	W.A.	—	—	—	—	—	B.fl. IV, 361	M.fr. VI, 132.
L. crassifolia, R. Brown, prodr. 455 (1810) ...	W.A. S.A.	—	—	—	—	—	B.fl. IV, 362	
L. ovata, R. Brown, prodr. 455 (1810) ...	W.A. S.A.	—	V.	—	—	—	B.fl. IV, 362	
L. buxifolia, F. v. M., fragm. VI, 132 (1868) ...	W.A.	—	—	—	—	—	B.fl. IV, 362	M.fr. IV, 132.
L. stenophylla, F. v. M., fragm. I, 128 (1859) ...	W.A.	—	—	—	—	—	B.fl. IV, 362	M.fr. I, 128.
L. micrantha, Benth in Journ. Linn. Soc. I, 94 (1857) ...	W.A.	—	—	—	—	—	B.fl. IV, 363	M.fr. VI, 132.
L. linifolia, Schlechtendal, Linnaea XX, 605 (1847) ...	—	S.A.	—	V.	N.S.W.	—	B.fl. IV, 363	M.fr. VI, 131.
L. fasciculata, R. Brown, prodr. 456 (1810) ...	W.A.	—	—	—	—	—	B.fl. IV, 363	M.fr. VI, 131.
L. floribunda, R. Brown, prodr. 456 (1810) ...	—	—	—	V.	N.S.W.	—	B.fl. IV, 364	M.fr. VI, 131.
L. cordifolia, Hooker in Mitch. Trop. Austr. 341 (1848) ...	—	—	—	—	—	Q.	B.fl. IV, 364	
L. nuda, F. v. M., fragm. I, 129 (1859) ...	W.A. S.A.	—	V.	N.S.W.	—	—	B.fl. IV, 365	M.fr. III, 163; VI, 131.
L. spermacoea, F. v. M., fragm. VI, 134 (1868) ...	W.A.	—	—	—	—	—	B.fl. IV, 365	M.fr. VI, 134.
L. callosa, F. v. M., fragm. VI, 134 (1868) ...	W.A.	—	—	—	—	—	B.fl. IV, 365	M.fr. VI, 134.
L. campanulata, R. Brown, prodr. 456 (1810) ...	W.A.	—	—	—	—	—	B.fl. IV, 365	M.fr. VI, 133.
L. serpillifolia, R. Brown, prodr. 456 (1810) ...	W.A.	—	—	—	—	—	B.fl. IV, 366	M.fr. VI, 133.
L. pusilla, R. Brown, prodr. 456 (1810) ...	—	—	—	N.S.W.	Q.	—	B.fl. IV, 366	M.fr. VI, 131.

PLANTAGINEAE.

A. L. de Jussieu, gen. plant. 89 (1789).

PLANTAGO, Tournefort, inst. 126, t. 48 (1700), from l'Ecluse (1576).

P. varia, R. Brown, prodr. 424 (1810) ...	W.A. S.A. T. V. N.S.W. Q.	—	—	—	—	B.fl. V, 139	M.fr. XI, 20.
P. stellaris, F. v. M., fragm. II, 23 (1860) ...	—	—	V.	N.S.W.	—	—	M.fr. VIII, 148.
P. Brownii, Rapin in Mém. de la Soc. Linn. Par. VI, 484 (1827) ...	—	—	T.	—	—	—	M.fr. II, 23.
P. Gunnii, J. Hooker in Lond. Journ. V, 446, t. 13 (1846) ...	—	—	T. V. N.S.W.	—	—	B.fl. V, 142	M.fr. VIII, 148; XI, 134.

PRIMULACEAE.

Ventenat, Tabl. du régn. végét. II, 285 (1799).

ANAGALLIS, Tournefort, inst. 142, t. 59 (1700), from Hippocrates and Dioscorides. (Centunculus, Micropyxia.)

A. pumila, Swartz, Fl. Ind. occ. I, 345 (1797) ...	—	—	—	—	—	Q. N.A.	B.fl. IV, 270	
--	---	---	---	---	---	---------	---------------	--

LYSIMACHIA, Tournefort, inst. 141, t. 59 (1700), from Dioscorides.

L. salicifolia, F. v. M. in Benth. Fl. Austr. IV, 269 (1869) ...	—	—	—	V.	N.S.W.	—	B.fl. IV, 269	
L. Japonica, Thunberg, F. Japon. 83 (1784) ...	—	—	—	—	N.S.W.	—	B.fl. IV, 269	

SAMOLUS, Tournefort, inst. 143, t. 60 (1700). (Sheffieldia.)

S. Valerandi, Linné, spec. plant. 171 (1753) ...	—	—	—	V.	N.S.W. Q.	—	B.fl. IV, 271	
S. repens, Persoon, synops. plant. I, 171 (1805) ...	W.A. S.A. T. V. N.S.W. Q. N.A.	—	—	—	—	—	B.fl. IV, 271	M.fr. XI, 20.
S. platyphyllus, F. v. M. inedit. ...	—	—	S.A.	—	—	—	N.A.	—

MYRSINACEAE.

R. Brown, prodr. 532 (1810). (Primulacearum subordo.)

MAESA, Forskael, Fl. Aegypt. Arab. 66 (1775). (Baeobotrys.)

M. dependens, F. v. M., fragm. V, 107 (1865) ...	—	—	—	—	—	Q.	—	B.fl. IV, 273	M.fr. V, 107.
M. haplobotrys, F. v. M., fragm. V, 161 (1866) ...	—	—	—	—	—	Q.	—	B.fl. IV, 273	M.fr. V, 161.

SAMARA, Linné, mantiss. II, 144 (1771). (Choripetalum.)

S. Australiana, F. v. M., fragm. VI, 164 (1868) ...	—	—	—	—	N.S.W. Q.	—	B.fl. IV, 274	M.fr. IV, 36.
---	---	---	---	---	-----------	---	---------------	---------------

MYRSINE, Linné, syst. nat. 8 (1735); Linné, gen. plant. 54 (1737).

M. urceolata, R. Brown, prodr. 534 (1810)...	...	...	...	—	—	—	—	Q.	—	B.fl. IV, 274	
M. campanulata, F. v. M., fragm. VI, 235 (1868)...	...	...	...	—	—	—	—	Q.	—	—	M.fr. VI, 235.
M. crassifolia, R. Brown, prodr. 534 (1810)	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl. IV, 275	M.fr. IV, 81; VI, 164; VIII, 148.
M. platystigma, F. v. M., fragm. VIII, 48 (1873)	...	...	...	—	—	—	N.S.W.	—	—	—	M.fr. VIII, 48.
M. variabilis, R. Brown, prodr. 534 (1810)...	...	...	...	—	—	—	V. N.S.W.	Q.	—	B.fl. IV, 275	
M. achradifolia, F. v. M., fragm. VI, 164 (1868)	...	...	...	—	—	—	—	Q.	—	B.fl. IV, 275	M.fr. VI, 164.

ARDISIA, Swartz, nov. gen. et spec. 3 et 48 (1788).

A. Pseudo-Jambosa, F. v. M., fragm. IV, 81 (1864) ...	...	—	—	—	N.S.W. Q.	—	B.fl.IV,276	M.fr.IV,81.	
A. brevipedata, F. v. M., fragm. VI, 163 (1868) ...	...	—	—	—	—	Q.	—	B.fl.IV,276	M.fr.VI,163.

AEGICERAS, Gaertner, de fruct. I, 216, t. 46 (1788).

A. majus, Gaertner, de fruct. I, 216, t. 46 (1788) ...	—	—	—	—	N.S.W. Q. N.A.	—	B.fl. IV, 277	
--	---	---	---	---	----------------	---	---------------	--

SAPOTACEAE.

A. L. de Jussieu, gen. 151 (1789), from B. de Jussieu (1759).

NIEMEYERA, F. v. M., fragm. VII, 114 (1870). (Chrysophyllum partly.)

N. prunifera, F. v. M., fragm. VII, 114 (1870) ...	—	—	—	—	N.S.W. Q.	—	B.fl. IV, 278	M.fr. VI, 26.
--	---	---	---	---	-----------	---	---------------	---------------

AMORPHOSPERMUM, F. v. M., fragm. VII, 112 (1870).

A. antilogum, F. v. M., fragm. VII, 113 (1870) ...	—	—	—	—	N.S.W. Q.	—	—	M.fr. X, 119.
--	---	---	---	---	-----------	---	---	---------------

LUCUMA, Molina, Saggio, 186 (1782). (Sersalisia.)

L. sericea, Benth & J. Hooker, gen. pl. II, 654 (1876) ...	—	—	—	—	—	Q. N.A.	B.fl. IV, 279	M.fr. VII, 112.
L. Galactoxylon, Benth & J. Hooker, gen. pl. II, 654 (1876) ...	—	—	—	—	—	Q.	—	B.fl. IV, 279

SIDEROXYLON, Dillenius, hort. Eltham. 357, t. 265 (1732). (Achras partly, Sapota partly, Sersalisia partly.)

S. Arnheimicum, Benth & J. Hooker, gen. pl. II, 655 (1876)...	—	—	—	—	—	N.A.	B.fl. IV, 280		
S. Pohlmanianum, Benth & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	Q.	—	B.fl. IV, 281	M.fr. V, 184.
S. xerocarpum, Benth & J. Hooker, gen. pl. II, 655 (1876) ...	—	—	—	—	—	Q.	—	B.fl. IV, 281	
S. echartaceum, Benth & J. Hooker, gen. pl. II, 655 (1876) ...	—	—	—	—	—	Q.	—	B.fl. IV, 281	

CARISSA, Linné, mantiss. pl. 7 (1767).											
C. laxiflora, Benth. Fl. Austr. IV, 305 (1869) ...	—	—	—	—	—	—	Q.	—	B.fl. IV, 305		
C. Brownii, F. v. M., fragm. IV, 45 (1863) ...	—	—	—	—	S.A.	—	N.S.W.	Q.	N.A.	B.fl. IV, 305	M.fr. IV, 45.
CERBERA, Linné, gen. plant. 62 (1737).											
C. Odollam, Gaertner, de fruct. 193, t. 124 (1791) ...	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 306	M.fr. VI, 117.
ALYXIA, Banks in R. Brown, prodr. 469 (1810). (Gynopogon.)											
A. buxifolia, R. Brown, prodr. 470 (1810) ...	—	—	—	—	W.A.	S.A.	T. V.	N.S.W.	—	B.fl. IV, 307	M.fr. VI, 117.
A. ruscifolia, R. Brown, prodr. 470 (1810) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 308	M.fr. VI, 117.
A. ilicifolia, F. v. M., fragm. IV, 149 (1864) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 308	M.fr. IV, 149.
A. squamulosa, C. Moore & F. v. M., fragm. VIII, 47 (1873) ...	—	—	—	—	—	—	—	N.S.W.	—	—	M.fr. VI, 117; VIII, 47.
A. obtusifolia, R. Brown, prodr. 470 (1810) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 308	
A. Gynopogon, Roemer & Schultes, syst. veg. IV, 440 (1819) ...	—	—	—	—	—	—	—	N.S.W.	—	—	M.fr. VIII, 47.
A. spicata, R. Brown, prodr. 470 (1810) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 308
A. Lindii, F. v. M., fragm. VIII, 46 (1873) ...	—	—	—	—	—	—	—	N.S.W.	—	—	M.fr. VI, 117.
A. thyrsiflora, Benth. Fl. Austr. IV, 309 (1869) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 309	M.fr. VIII, 46.
A. Thozetii, F. v. M., fragm. X, 103 (1877) ...	—	—	—	—	—	—	—	—	Q.	—	M.fr. X, 103.
OCHROSIA, A. L. de Jussieu, gen. 143 (1789). (Lactaria.)											
O. elliptica, Labillardière, sert. Austr. Caled. 27, t. 30 (1824) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 310	M.fr. VI, 118.
O. Moorei, F. v. M. in Benth. Fl. Austr. IV, 310 (1869) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 310	M.fr. VI, 118.
O. Kilneri, F. v. M., fragm. VII, 129 (1871) ...	—	—	—	—	—	—	—	—	Q.	—	M.fr. VIII, 129.
NOTONERIUM, Benth. in B. & J. H. gen. II, 698 (1876).											
N. Gossei, Benth. in B. & J. H. gen. pl. 698 (1876) ...	—	—	—	—	S.A.	—	—	—	—	—	M.fr. X, 103.
TABERNAEMONTANA, Plumier, nov. pl. Amer. gen. 18, t. 30 (1703).											
T. orientalis, R. Brown, prodr. 468 (1810) ...	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 311
T. pubescens, R. Brown, prodr. 468 (1810) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 311
VINCA, Rivinus in Rupp. Fl. Jenens. 27 (1818), from Plinius. (Perhaps immigrated.)											
V. rosea, Linné, spec. plant. ed. sec. 305 (1762) ...	—	—	—	—	—	—	—	—	Q.	—	M.fr. XI, 123.
ALSTONIA, R. Brown in Mem. Wern. Soc. I, 75 (1809).											
A. scholaris, R. Brown in Mem. Wern. Soc. I, 75 (1809) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 312	M.fr. VI, 117.
A. verticillata, F. v. M., fragm. VI, 116 (1868) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 313
A. ophioxylodes, F. v. M., fragm. I, 57 (1858) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 313
A. villosa, Blume, Bijdr. 1038 (1826) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 313	M.fr. VI, 117.
A. linearis, Benth. Fl. Austr. IV, 314 (1869) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 314
A. constricta, F. v. M., fragm. I, 57 (1858) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 314	M.fr. III, 163; IV, 170; VI, 117.
ICHNOCARPUS, R. Brown in Mem. Wern. Soc. I, 75 (1809).											
I. frutescens, R. Brown in Mem. Wern. Soc. I, 75 (1809) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 315
WRIGHTIA, R. Brown in Mem. Wern. Soc. I, 75 (1809). (Balfouria.)											
W. pubescens, R. Brown, prodr. 467 (1810) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 316
W. saligna, F. v. M. in Benth. Fl. Austr. IV, 316 (1869) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 316
W. Cunninghamii, Benth. Fl. Austr. IV, 317 (1869) ...	—	—	—	—	W.A.	—	—	—	—	B.fl. IV, 317	M.fr. V, 161; VI, 118.
PARSONSIA, R. Brown in Mem. Wern. Soc. I, 64 (1809).											
P. lanceolata, R. Brown, prodr. 466 (1810) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 318	M.fr. VI, 126.
P. velutina, R. Brown, prodr. 466 (1810) ...	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 318
P. Leichhardtii, F. v. M., fragm. VI, 128 (1868) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 319	M.fr. VI, 128.
P. ventricosa, F. v. M. in Transact. phil. Inst. Vict. II, 71 (1857) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 319	M.fr. VI, 130.
LYONSIA, R. Brown in Mem. Wern. Soc. I, 66 (1809).											
L. lilacina, F. v. M. in Benth. Fl. Austr. IV, 321 (1869) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 321	M.fr. VI, 127.
L. induplicata, F. v. M. in Benth. Fl. Austr. IV, 321 (1869) ...	—	—	—	—	—	—	—	N.S.W.	—	B.fl. IV, 321	M.fr. VI, 129.
L. straminea, R. Brown, prodr. 466 (1810) ...	—	—	—	—	—	—	T. V.	N.S.W.	—	B.fl. IV, 321	M.fr. VI, 129.
L. reticulata, F. v. M., Rep. Burdek. Exped. 16 (1860) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 321	M.fr. VI, 129.
L. diaphanophleba, F. v. M. in Benth. Fl. Austr. IV, 322 (1869) ...	—	—	—	—	W.A.	—	—	—	—	B.fl. IV, 322	M.fr. II, 158; VI, 130.
L. Langiana, F. v. M., fragm. VI, 128 (1868) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 322	M.fr. VI, 128.
L. largiflorens, F. v. M. in Benth. Fl. Austr. IV, 322 (1869) ...	—	—	—	—	—	—	—	N.S.W.	Q.	B.fl. IV, 322	
L. latifolia, Benth. Fl. Austr. IV, 323 (1869) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 323	
L. oblongifolia, Benth. Fl. Austr. IV, 323 (1869) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 323	
L. eucalyptifolia, F. v. M. in Benth. Fl. Austr. IV, 323 (1869) ...	—	—	—	—	—	—	—	—	Q.	B.fl. IV, 323	M.fr. II, 159; VI, 130

ASCLEPIADEAE.

N. J. Jacquin, Misc. Austr. I, 35 (1778).

GYMNANTHERA, R. Brown in Mem. Wern. Soc. I, 58 (1809).											
G. nitida, R. Brown, prodr. 464 (1810) ...	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl. IV, 326
SECAMONE, R. Brown in Mem. Wern. Soc. I, 55 (1809).											
S. elliptica, R. Brown, prodr. 464 (1810) ...	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 327
S. ovata, R. Brown, prodr. 464 (1810) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 328
VINCETOXICUM, Dodoens, stirp. hist. pemptades, 704 (1583). (Oxystelma partly, Cynoctonum.)											
V. ovatum, Benth. Fl. Austr. IV, 330 (1869) ...	—	—	—	—	—	—	—	—	Q.	—	B.fl. IV, 330
V. elegans, Benth. Fl. Austr. IV, 330 (1869) ...	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 332
V. carnosum, Benth. Fl. Austr. IV, 331 (1869) ...	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. IV, 331
V. leptolepis, Benth. Fl. Austr. IV, 331 (1869) ...	—	—	—	—	—	—	—	—	Q.	—	B.fl. IV, 331
CYNANCHUM, Linné, gen. plant. 63 (1737).											
C. erubescens, R. Brown, prodr. 463 (1810) ...	—	—	—	—	—	—	—	—	Q.	—	B.fl. IV, 332
C. floribundum, R. Brown, prodr. 463 (1810) ...	—	—	—	—	S.A.	—	—	—	Q.	N.A.	B.fl. IV, 332
C. pedunculatum, R. Brown, prodr. 463 (1810) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 333
C. puberulum, F. v. M. in Benth. Fl. Austr. IV, 333 (1869) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl. IV, 333

- SARCOSTEMMA, R. Brown in Mem. Wern. Soc. I, 50 (1809).
 S. Australe, R. Brown, prodr. 463 (1810) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.IV,328 M.fr.V,160;XI,80.
- DAEMIA, R. Brown in Mem. Wern. Soc. I, 50 (1809). (Pentatropis, Rhyncharrhena.)
 D. linearis, F. v. M.; Pentatropis, Decaisne in De Cand., prodr. VIII, 536 (1833) ... W.A. — — — — — B.fl.IV,329
 D. atropurpurea; Rhyncharrh., F. v. M., fr. I, 128 (1859) ... — — — — Q. — B.fl.IV,329 M.fr.I,128.
 D. quinquepartita; Rhyncharrh., F. v. M., fr. I, 128 (1859) ... — — — — V. N.S.W. — — B.fl.IV,329 M.fr.I,128;XI,80.
 D. Kempeana, F. v. M. in Wing's S. Sc. Record, Aug. (1882) ... — S.A. — — — — — —
- GYMNEMA, R. Brown in Mem. Wern. Soc. I, 33 (1809). (Bidaria.)
 G. sylvestre, R. Brown in Mem. Wern. Soc. I, 33 (1809) ... — — — — — N.A. B.fl.IV,343
 G. pleiadenium, F. v. M., fragm. XI, 77 (1879) ... — — — — — Q. — — M.fr.XI,77.
 G. Muellieri, Benth., Fl. Austr. IV, 343 (1869) ... — — — — — N.A. B.fl.IV,343
 G. brevifolium, Benth., Fl. Austr. IV, 343 (1869) ... — — — — — Q. — — B.fl.IV,343
 G. trinerve, R. Brown, prodr. 462 (1810) ... — — — — — N.A. B.fl.IV,343
 G. stenophyllum, A. Gray in Proc. Am. Ac. of Sc. V, 335 (1864) ... — — — — — N.A. B.fl.IV,344
- GONGRONEMA, Decaisne in De Candolle, prodr. VIII, 624 (1844).
 G. micradenia, Benth. & J. Hooker, gen. plant. II, 770 (1876) ... — — — — — Q. — — B.fl.IV,344 M.fr.XI,78.
- TYLOPHORA, R. Brown in Mem. Wern. Soc. I, 28 (1809).
 T. erecta, F. v. M. in Benth. Fl. Austr. IV, 334 (1869) ... — — — — — Q. — — B.fl.IV,334 M.fr.XI,79.
 T. macrophylla, Benth., Fl. Austr. IV, 334 (1869) ... — — — — — N.A. B.fl.IV,334
 T. grandiflora, R. Brown, prodr. 460 (1810) ... — — — — — N.S.W. Q. — — B.fl.IV,334 M.fr.XI,80.
 T. floribunda, Benth., Fl. Austr. IV, 335 (1869) ... — — — — — N.S.W. Q. — — B.fl.IV,335
 T. calcarata, Benth., Fl. Austr. IV, 335 (1869) ... — — — — — Q. — — B.fl.IV,335
 T. barbata, R. Brown, prodr. 460 (1810) ... — — — — — V. N.S.W. — — B.fl.IV,335 M.fr.V,160;XI,79.
 T. Woollsi, Benth., Fl. Austr. IV, 335 (1869) ... — — — — — N.S.W. — — B.fl.IV,335
 T. paniculata, R. Brown, prodr. 460 (1810) ... — — — — — N.S.W. — — B.fl.IV,336
 T. enervis, F. v. M., fragm. IX, 70 (1875) ... — — — — — N.S.W. — — — M.fr.IX,70.
 T. flexuosa, R. Brown, prodr. 460 (1810) ... — — — — — N.A. B.fl.IV,336 M.fr.XI,79.
 T. biglandulosa, A. Gray in Proc. of Am. acad. sc. vol. V (1864) ... — — — — — N.S.W. — — — M.fr.IX,169.
- MARSDENIA, R. Brown in Mem. Wern. Soc. I, 28 (1809). (Leichhardtia.)
 M. cinerascens, R. Brown, prodr. 461 (1810) ... — — — — — N.A. B.fl.IV,337
 M. rhyncholepis, F. v. M., fragm. XI, 78 (1879) ... — — — — — Q. — — M.fr.XI,78.
 M. flavesces, Cunningham in Bot. Mag., t. 3289 (1833) ... — — — — — N.S.W. — — B.fl.IV,337 M.fr.V,160.
 M. cymulosa, Benth., Fl. Austr. IV, 338 (1869) ... — — — — — Q. — — B.fl.IV,338
 M. velutina, R. Brown, prodr. 461 (1810) ... — — — — — N.A. B.fl.IV,338
 M. Hullsi, F. v. M. in Benth. Fl. Austr. IV, 338 (1869) ... — — — — — N.A. B.fl.IV,338
 M. araujacea, F. v. M., fragm. VI, 135 (1868) ... — — — — — Q. — — B.fl.IV,339 M.fr.VI,135.
 M. tubulosa, F. v. M., fragm. IX, 71 (1875) ... — — — — — N.S.W. — — — M.fr.IX,71.
 M. rostrata, R. Brown, prodr. 461 (1810) ... — — — — — V. N.S.W. Q. — — B.fl.IV,339 M.fr.V,160;VI,135.
 M. Fraseri, Benth., Fl. Austr. IV, 339 (1869) ... — — — — — Q. — — B.fl.IV,339
 M. longiloba, Benth., Fl. Austr. IV, 340 (1869) ... — — — — — N.S.W. Q. — — B.fl.IV,340 M.fr.XI,79.
 M. suaveolens, R. Brown, prodr. 461 (1810) ... — — — — — N.S.W. — — B.fl.IV,340
 M. leptophylla, F. v. M. in Benth. Fl. Austr. IV, 340 (1869) ... — — — — — Q. — — B.fl.IV,340
 M. Leichhardtiana, F. v. M., fragm. V, 160 (1866) ... — — — — — W.A. S.A. — — V. N.S.W. Q. N.A. B.fl.IV,341 M.fr.V,160.
 M. viridiflora, R. Brown, prodr. 461 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.IV,341 M.fr.V,160.
 M. coronata, Benth., Fl. Austr. IV, 341 (1810) ... — — — — — Q. — — B.fl.IV,341
 M. microlepis, Benth., Fl. Austr. IV, 342 (1869) ... — — — — — Q. — — B.fl.IV,342
- THOZETIA, F. v. M. in Benth. Fl. Austr. IV, 347 (1869).
 T. racemosa, F. v. M. in Benth. Fl. Austr. IV, 347 (1869) ... — — — — — Q. — — B.fl.IV,347
- HOYA, R. Brown in Mem. Wern. Soc. I, 26 (1809).
 H. carnosa, R. Brown in Mem. Wern. Soc. I, 26 (1809) ... — — — — — Q. — — B.fl.IV,346
 H. australis, R. Brown in Transact. hort. Soc. VII, 28 (1826) ... — — — — — N.S.W. Q. — — B.fl.IV,346 M.fr.XI,80.
 H. Nicholsoniae, F. v. M., fragm. V, 159 (1866) ... — — — — — Q. — — B.fl.IV,347 M.fr.V,159;IX,70,190.
- DISCHIDIA, R. Brown in Mem. Wern. Soc. I, 32 (1809).
 D. nummularia, R. Brown, prodr. 461 (1810) ... — — — — — Q. — — B.fl.IV,345 M.fr.V,161.
 D. Timorensis, Decaisne in nouv. annal. du Mus. III, 377, t. 17 (1835) ... — — — — — Q. N.A. — — —
- MICROSTEMMA, R. Brown in Mem. Wern. Soc. I, 25 (1809).
 M. tuberosum, R. Brown, prodr. 459 (1810) ... — — — — — Q. N.A. B.fl.IV,345 M.fr.I,58.
 M. glabriflorum, F. v. M., fragm. I, 58 (1858) ... — — — — — N.A. B.fl.IV,345 M.fr.I,58.
- CEROPEGIA, Linné, gen. plant. 65 (1737).
 C. Cumingiana, Decaisne in De Cand., prodr. VIII, 643 (1844) ... — — — — — Q. — — B.fl.IV,348 M.fr.V,159.

CONVOLVULACEAE.

A. L. de Jussieu, gen. plant. 132 (1789), from B. de Jussieu (1759).

- ERYCIBE, Roxburgh, pl. Corom. III, 31, t. 159 (1798).
 E. paniculata, Roxburgh, pl. Corom. III, 31, t. 159 (1798) ... — — — — — Q. — — B.fl.IV,411 M.fr.VI,100.
- LEPISTEMON, Blume, Bijdrag. 722 (1825).
 L. urceolatus, F. v. M.; L. Fitzalanii, F. v. M., fragm. X, 111 (1877) ... — — — — — Q. — — B.fl.IV,427 M.fr.X,111.
- IPOMOEA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 47 (1737). (Pharbitis, Batatas, Calonyction, Aniscia, Skinnera.)
 I. paniculata, R. Brown, prodr. 486 (1810) ... — — — — — Q. N.A. B.fl.IV,414 M.fr.VI,98;X,111.
 I. Davenporti, F. v. M., fragm. VI, 97 (1868) ... — — — — — S.A. — — — N.A. B.fl.IV,415 M.fr.VIII,17;X,111.
 I. bona nox, Linné, spec. plant. edit. secund. 228 (1762) ... — — — — — N.S.W. Q. — — — M.fr.IX,74;X,113;X,73.
 I. palmata, Forskael, Fl. Aegypt. Arabic. 43 (1775) ... — — — — — N.S.W. Q. — — B.fl.IV,415 M.fr.VI,98;X,111.
 I. quinata, R. Brown, prodr. 486 (1810) ... — — — — — Q. N.A. B.fl.IV,415 M.fr.VI,98;X,112.
 I. diversifolia, R. Brown, prodr. 487 (1810) ... — — — — — Q. N.A. B.fl.IV,416 M.fr.X,112.

I. dissecta, Willdenow, phytogr. 5, t. 2 (1794) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 416	M.fr. VI, 98; X, 112.
I. sinuata, Ortega, decad. nov. pl. hort. Matrit. VII, 84 (1799) ...	—	—	—	—	—	Q.	—	—	M.fr. VIII, 17; X, 113.
I. hederacea, N. J. Jacquin, Collectan. I, 124 (1786) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 416	M.fr. VI, 99; X, 112.
I. congesta, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 417	M.fr. VI, 99; X, 112.
I. purpurea, Roth, Catalact. bot. I, 36 (1797) ...	—	—	—	—	—	Q.	—	B.f.IV, 417	M.fr. VI, 99; X, 112.
I. peltata, Choisy, Convolvulac. orient. 70 (1834) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 418	—
I. Calobra, Hill & F. v. M., fragm. XI, 73 (1879) ...	—	—	—	—	—	Q.	—	—	M.fr. XI, 73, 137.
I. alata, R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	N.A.	—	B.f.IV, 418	M.fr. X, 112.
I. Turpethum, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 418	M.fr. VI, 98.
I. longiflora, R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 418	M.fr. VI, 98; IX, 74; X, 112.
I. costata, F. v. M. in Benth, Fl. Austr. IV, 419 (1869) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f.IV, 419	M.fr. X, 112.
I. Pes Caprae, Roth, nov. spec. plant. 109 ...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.f.IV, 419	M.fr. VI, 97; X, 112.
I. carnosa, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 420	—
I. reptans, Poiret, Encycl. méth. suppl. III, 460 (1813) ...	W.A.	—	—	—	—	Q.	N.A.	B.f.IV, 420	M.fr. VI, 97.
I. graminea, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 421	—
I. velutina, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 421	—
I. abrupta, R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 421	M.fr. X, 112.
I. denticulata, Choisy in De Cand: prodr. IX, 379 (1845) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 421	M.fr. I, 185; X, 112.
I. gracilis, R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 422	M.fr. I, 185; X, 112.
I. sepiaia, Koenig in Roxb. Fl. Indic. ed. Carey II, 90 (1824) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 422	—
I. Muelleri, Benth, Fl. Austr. IV, 423 (1869) ...	W.A.	—	—	—	—	—	N.A.	B.f.IV, 423	M.fr. X, 112.
I. cymosa, Roemer & Schultes, syst. veg. IV, 241 (1819) ...	—	—	—	—	—	Q.	—	B.f.IV, 423	—
I. linifolia, Blume, Bijdrag. 721 (1825) ...	—	—	—	—	—	Q.	—	B.f.IV, 423	—
I. chryseides, Ker in Bot. Regist. t. 270 (1818) ...	—	—	—	—	—	Q.	—	B.f.IV, 423	M.fr. VI, 98.
I. flava, F. v. M. in Benth. Fl. Austr. IV, 424 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 424	—
I. obscura, Ker in Bot. Regist. t. 239 (1817) ...	—	—	—	—	—	Q.	—	B.f.IV, 424	—
I. incisa, R. Brown, prodr. 486 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 424	M.fr. I, 186.
I. uniflora, Roemer & Schultes, syst. IV, 247 (1819) ...	—	—	—	—	—	Q.	—	B.f.IV, 425	M.fr. VI, 99.
I. angustifolia, N. J. Jacquin, Collect. II, 367 (1788) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 425	M.fr. VI, 97; X, 112.
I. plebeja, R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 426	M.fr. VI, 99; X, 112.
I. eriocarpa, R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 426	M.fr. VI, 98; X, 112.
I. heterophylla, R. Brown, prodr. 487 (1810) ...	—	S.A.	—	—	—	Q.	N.A.	B.f.IV, 426	M.fr. VI, 98; X, 112.
I. erecta, R. Brown, prodr. 487 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 427	M.fr. VI, 98.

CONVOLVULUS, Tournefort, inst. 82, t. 17 (1700). (Calystegia.)									
C. erubescens, Sims, Bot. Magaz. t. 1067 (1808) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 429	M.fr. VI, 99; X, 113.
C. parviflorus, Vahl, symbol. bot. III, 29 (1794) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 430	M.fr. VI, 99; X, 113.
C. marginatus, Poiret, Encycl. méth. suppl. II (1811) ...	—	—	—	—	V.	N.S.W.	Q.	—	B.f.IV, 430
C. sepium, Linné, spec. plant. 153 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 430	M.fr. VI, 99; X, 113.

POLYMERIA, R. Brown, prodr. 488 (1810).									
P. marginata, Benth, Fl. Austr. IV, 432 (1869) ...	—	—	—	—	—	Q.	—	B.f.IV, 432	—
P. longifolia, Lindley in Mitch. Trop. Austr. 398 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.IV, 432	M.fr. X, 114.
P. angusta, F. v. M., fragm. VI, 100 (1868) ...	—	—	—	—	—	—	N.A.	B.f.IV, 432	M.fr. VI, 100.
P. distigma, Benth, Fl. Austr. IV, 433 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 433	—
P. occidentalis, F. v. M., inedit. ...	W.A.	—	—	—	—	—	—	B.f.IV, 433	—
P. calycina, R. Brown, prodr. 488 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 433	M.fr. VI, 100.
P. ambigua, R. Brown, prodr. 488 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 433	—

PORANA, Burmann, Fl. Ind. 51, t. 21 (1768). (Duperreya.)									
P. sericea, F. v. M., fragm. VI, 100 (1868) ...	W.A.	—	—	—	—	—	N.A.	B.f.IV, 434	M.fr. II, 22.

BREWERIA, R. Brown, prodr. 487 (1810).									
B. linearis, R. Brown, prodr. 488 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 435	—
B. media, R. Brown, prodr. 488 (1810) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.IV, 436	M.fr. X, 113.
B. brevifolia, Benth, Fl. Austr. IV, 436 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 436	—
B. rosea, F. v. M., fragm. I, 233 (1858) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f.IV, 436	M.fr. VI, 100; X, 113.
B. pannosa, R. Brown, prodr. 488 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV, 436	M.fr. VI, 100; X, 113.

EVOLVULUS, Linné, spec. plant. ed. sec. 391 (1763).									
E. linifolius, Linné, spec. plant. ed. sec. 392 (1762) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.IV, 437	M.fr. VI, 100; X, 113.

DICHONDRA, R. & G. Forster, char. gen. 39, t. 20 (1776).									
D. repens, R. & G. Forster, char. gen. 39, t. 20 (1776) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.IV, 438	M.fr. VI, 100; X, 113.

CRESSA, Linné, Amoen. Acad. I, 121 (1747).									
C. Cretica, Linné, spec. plant. 223 (1753) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 437	M.fr. VI, 100.

WILSONIA, R. Brown, prodr. 490 (1810).									
W. humilis, R. Brown, prodr. 490 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 439	M.fr. VI, 101; X, 113.
W. rotundifolia, Hooker, icon. plant. t. 410 (1842) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 440	M.fr. VI, 101.
W. Backhousi, J. Hooker in Lond. Journ. VI, 275 (1842) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 440	M.fr. VI, 101; VIII, 17.

CUSCUTA, Tournefort, inst. app. 652, t. 422 (1700), from C. Bauhin (1623).									
C. Chinensis, Lamarck, Encycl. méth. II, 229 (1786) ...	—	—	—	—	—	Q.	—	B.f.IV, 441	M.fr. X, 113.
C. Australis, R. Brown, prodr. 491 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f.IV, 441	M.fr. I, 100; X, 113.
C. Tasmanica, Engelmann in Transact. Acad. St. Louis I, 512 (1860) ...	—	—	T.	V.	—	—	—	B.f.IV, 441	—

SOLANACEAE.

Haller, enum. stirp. Helv. praef. 34 (1742).

PHYSALIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 51 (1737).									
P. minima, Linné, spec. plant. 183 (1753) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV, 466	M.fr. VI, 144.

SOLANUM, Tournefort, inst. 148, t. 62 (1700), from Celsus.									
S. nigrum, Linné, spec. plant. 186 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.IV, 446	M.fr. VI, 145.
S. vescum, F. v. M. in Transact. Vict. Inst. 69 (1855) ...	—	—	T.	V.	N.S.W.	—	—	—	—

S. aviculare, G. Forster, prodr. 18 (1786) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV, 447	M.fr. VI, 144.
S. simile, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 448	M.fr. I, 123; VI, 144.
S. fasciculatum, F. v. M., fragm. I, 123 (1859) ...	W.A.	S.A.	—	—	—	—	—	—	M.fr. VI, 144.
S. Bauerianum, Endlicher, prodr. fl. Norfolk, 54 (1833) ...	—	—	—	—	N.S.W.	—	—	—	M.fr. IX, 74.
S. viride, R. Brown, prodr. 445 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 449	M.fr. VI, 145.
S. tetrandrum, R. Brown, prodr. 445 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 449	—
S. Shanesii, F. v. M., fragm. VI, 144 (1868) ...	—	—	—	—	—	Q.	—	B.f.IV, 448	M.fr. VI, 144.
S. verbascifolium, Linné, spec. plant. 184 (1753) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 449	M.fr. VI, 145.
S. discolor, R. Brown, prodr. 445 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV, 450	M.fr. II, 166; VI, 147.
S. stelligerum, Smith, Exot. Bot. II, 57, t. 88 (1805) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 450	M.fr. VI, 147.
S. parvifolium, R. Brown, prodr. 446 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 451	M.fr. II, 164; VI, 147.
S. ferocissimum, Lindley in Mitch. Three Exped. II, 58 (1838) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f.IV, 451	—
S. sporadotrichum, F. v. M. in Melb. Chemist, Oct. (1882) ...	—	—	—	—	—	Q.	—	—	—
S. defensum, F. v. M., fragm. V, 193 (1866) ...	—	—	—	—	—	Q.	—	B.f.IV, 451	M.fr. V, 193.
S. violaceum, R. Brown, prodr. 445 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 452	—
S. tetrathecum, F. v. M., fragm. II, 165 (1861) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 453	M.fr. II, 165.
S. elachophyllum, F. v. M., fragm. II, 164 (1861) ...	—	—	—	—	—	Q.	—	B.f.IV, 453	M.fr. VI, 145.
S. orbiculatum, Dunal in Poir. Encycl. méth. XIII, 762 (1817) ...	W.A.	S.A.	—	—	—	—	—	B.f.IV, 453	M.fr. VI, 145.
S. oligacanthum, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.IV, 454	M.fr. VI, 145.
S. esuriale, Lindley in Mitch. Three Exped. II, 43 (1838) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 454	—
S. chenopodium, F. v. M., fragm. II, 165 (1861) ...	—	S.A.	—	—	N.S.W.	—	N.A.	B.f.IV, 454	M.fr. VI, 146.
S. Sturtianum, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854) ...	—	S.A.	—	—	N.S.W.	—	N.A.	B.f.IV, 454	M.fr. VI, 146.
S. furfuraceum, R. Brown, prodr. 446 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV, 455	—
S. dianthophorum, Dunal, hist. des Solan. 183 (1813) ...	—	—	—	—	—	Q.	—	B.f.IV, 455	M.fr. VI, 145.
S. Dallachyi, Benth, Fl. Austr. IV, 456 (1869) ...	—	—	—	—	—	Q.	—	B.f.IV, 456	N.fr. VI, 145.
S. densivestitum, F. v. M. in Benth. Fl. Austr. IV, 456 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 456	—
S. nemophilum, F. v. M., fragm. II, 161 (1861) ...	—	—	—	—	—	Q.	—	B.f.IV, 456	M.fr. II, 161.
S. Oldfieldii, F. v. M., fragm. II, 161 (1861) ...	W.A.	—	—	—	—	—	—	B.f.IV, 457	M.fr. VI, 145.
S. semiarmatum, F. v. M., fragm. II, 163 (1861) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 457	M.fr. VI, 146.
S. armatum, R. Brown, prodr. 446 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.IV, 458	—
S. Hystrix, R. Brown, prodr. 446 (1810) ...	—	S.A.	—	—	—	—	—	B.f.IV, 458	—
S. cataphractum, Cunningham in Benth. Fl. Austr. IV, 459 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 459	—
S. pungetium, R. Brown, prodr. 446 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.IV, 459	M.fr. VI, 146.
S. eremophilum, F. v. M. in Schlecht. Linnaea XXV, 432 (1852) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.IV, 459	M.fr. VI, 146.
S. campanulatum, R. Brown, prodr. 446 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 460	M.fr. VI, 146.
S. adenophorum, F. v. M., fragm. II, 162 (1861) ...	—	—	—	—	—	Q.	—	B.f.IV, 460	M.fr. II, 162.
S. cinereum, R. Brown, prodr. 446 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.IV, 460	—
S. lacunarum, F. v. M. in Transact. phil. Soc. Vict. I, 18 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 461	M.fr. VI, 146.
S. petrophilum, F. v. M. in Linnaea XXV, 433 (1852) ...	—	S.A.	—	—	N.S.W.	—	—	B.f.IV, 461	M.fr. VI, 146.
S. diversiflorum, F. v. M., fragm. VI, 146 (1868) ...	—	—	—	—	—	—	N.A.	B.f.IV, 461	M.fr. VI, 146.
S. carduiforme, F. v. M., fragm. II, 163 (1861) ...	—	—	—	—	—	—	N.A.	B.f.IV, 462	M.fr. II, 163.
S. melanospermum, F. v. M., fragm. II, 163 (1861) ...	—	—	—	—	—	—	N.A.	B.f.IV, 462	M.fr. II, 163.
S. horridum, Dunal, Solan. synops. 28 (1816) ...	—	—	—	—	—	—	N.A.	B.f.IV, 463	—
S. echinatum, R. Brown, prodr. 447 (1810) ...	—	—	—	—	—	—	N.A.	B.f.IV, 463	M.fr. VI, 145.
S. lasiophyllum, Dunal in Poir. Encycl. méth. XIII, 764 (1817) ...	W.A.	—	—	—	—	—	N.A.	B.f.IV, 463	M.fr. VI, 145.
S. ellipticum, R. Brown, prodr. 446 (1810) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f.IV, 464	M.fr. VI, 145.
S. quadriloculatum, F. v. M., fragm. II, 161 (1861) ...	—	—	—	—	—	—	N.A.	B.f.IV, 464	M.fr. II, 161.
S. phlomoideus, Cunningham in Benth. Fl. Austr. IV, 464 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 464	—
S. Cunninghamii, Benth, Fl. Austr. IV, 465 (1869) ...	—	—	—	—	—	—	N.A.	B.f.IV, 465	—
LYCIUM, Linné, gen. plant. 57 (1737), from Celsus.									
L. australe, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 467	M.fr. I, 83, 243; II, 179.
DATURA, Linné, gen. plant. 48 (1737).									
D. Leichhardtii, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ...	—	S.A.	—	—	—	Q.	N.A.	B.f.IV, 468	M.fr. VI, 144.
NICOTIANA, Tournefort, inst. 117, t. 41 (1700), from C. Bauhin (1623).									
N. suaveolens, Lehmann, gener. Nicot. hist. 43 (1818) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.IV, 469	M.fr. VI, 144.
ANTHOTROCHE, Endlicher, nov. stirp. mus. Vindob. dec. 6 (1839).									
A. pannosa, Endlicher, nov. stirp. mus. Vind. dec. 7 (1839) ...	W.A.	—	—	—	—	—	—	B.f.IV, 467	M.fr. VI, 142.
A. Blackii, F. v. M., fragm. VIII, 232 (1874) ...	W.A.	S.A.	—	—	—	—	—	—	M.fr. VIII, 232.
A. Walcottii, F. v. M., fragm. I, 123 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 468	M.fr. I, 123.
ANTHOCERCIS, Labillardière, Nov. Holl. pl. spec. II, 19 (1806). (Cyphanthera, Eadesia.)									
A. viscosa, R. Brown, prodr. 448 (1810) ...	W.A.	—	—	—	—	—	—	B.f.IV, 475	M.fr. VI, 143.
A. litorea, Labillard., Nov. Holl. pl. spec. II, 19, t. 158 (1806) ...	W.A.	—	—	—	—	—	—	B.f.IV, 476	M.fr. VI, 143; XI, 123.
A. gracilis, Benth in De Cand., prodr. X, 192 (1846) ...	W.A.	—	—	—	—	—	—	B.f.IV, 476	—
A. genistoides, Miers, Illustr. of plants II, App. 27, t. 83 (1857) ...	W.A.	—	—	—	—	—	—	B.f.IV, 476	M.fr. I, 122; VI, 143.
A. anisantha, Endlicher in Ann. Wien. Mus. II, 201 (1838) ...	W.A.	S.A.	—	—	—	—	—	B.f.IV, 477	M.fr. VI, 143.
A. intricata, F. v. M., fragm. I, 211 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 477	M.fr. I, 211.
A. arborea, F. v. M., fragm. I, 212 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 477	M.fr. I, 212; VI, 143.
A. angustifolia, F. v. M. in Transact. phil. Soc. Vict. I, 21 (1854) ...	—	S.A.	—	—	—	—	—	B.f.IV, 478	—
A. fasciculata, F. v. M., fragm. I, 122 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 478	M.fr. I, 122; VI, 143.
A. microphylla, F. v. M., fragm. I, 179 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 478	M.fr. I, 179; VI, 143.
A. myosotidea, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 478	M.fr. I, 179; VI, 143.
A. scabrella, Benth in De Cand., prodr. X, 192 (1846) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 479	—
A. albicans, Cunningham in Field's N.S.W. 335, t. 2 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 479	M.fr. VI, 143.
A. Odgersii, F. v. M., fragm. X, 19 (1876) ...	—	—	—	—	—	—	N.A.	—	M.fr. X, 19.
A. Tasmanica, J. Hooker, Fl. Tasman. I, 239, t. 92 (1860) ...	—	—	T.	—	—	—	—	B.f.IV, 479	M.fr. VI, 143.
A. Eadesii, F. v. M., fragm. II, 139 (1861) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 480	M.fr. VI, 142.
A. racemosa, F. v. M., fragm. I, 211 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 480	M.fr. I, 211.

- DUBOISIA, R. Brown, prodr. fl. Nov. Holl. 448 (1810).
 D. myoporoides, R. Brown, prodr. 448 (1810) ... — — — — N.S.W. Q. — B.fl. IV, 474 M.fr. VI, 144.
 D. Hopwoodi, F. v. M., fragm. VIII, 232 (1874) ... — — — — N.S.W. Q. — B.fl. IV, 480 M.fr. II, 138; VI, 143; X, 20.
 D. Leichhardtii, F. v. M. in Wing's S. Sc. Record, II, 222 (1882) ... — — — — N.S.W. — — B.fl. IV, 480 M.fr. VI, 142.

SCROPHULARINAE.

Mirbel, Elém. de Botan. II, 879 (1815).

- MIMULUS, Linné in Act. Upsal. 82 (1741). (Uvedalia.)
 M. Uvedaliae, Benth. in De Cand. X, 369 (1846) ... — — — — — Q. N.A. B.fl. IV, 482 M.fr. VI, 103; IX, 167.
 M. debilis, F. v. M. in Transact. phil. Inst. Vict. I, 62 (1857) ... — — — — — N.A. — — M.fr. VI, 103.
 M. gracilis, R. Brown, prodr. 439 (1810) ... — — — — — S.A. — — V. N.S.W. Q. N.A. B.fl. IV, 482 M.fr. VI, 103; IX, 167.
 M. repens, R. Brown, prodr. 439 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl. IV, 482 M.fr. VI, 103; IX, 166.
 M. prostratus, Benth. in De Cand. X, 373 (1846) ... — — — — — S.A. — — V. N.S.W. Q. — B.fl. IV, 483 M.fr. IX, 167.
- MAZUS, Loureiro, Fl. Cochinch. II, 385 (1790).
 M. Pumilio, R. Brown, prodr. 439 (1810) ... — — — — — S.A. T. V. N.S.W. Q. — B.fl. IV, 484 M.fr. VI, 104; IX, 167.
- ADENOSMA, R. Brown, prodr. fl. Nov. Holl. 442 (1810). (Pterostigma.)
 A. coerulea, R. Brown, prodr. 443 (1810) ... — — — — — Q. — — B.fl. IV, 484
 A. Muellieri, Benth. in Fl. Austr. IV, 485 (1869) ... — — — — — — — — N.A. B.fl. IV, 485
- STEMODIA, Linné, syst. ed. dec. 1118 (1759). (Morgania, Limnophila.)
 S. lythrifolia, F. v. M. in Benth. Fl. Austr. IV, 486 (1869) ... — — — — — — — — N.A. B.fl. IV, 486
 S. grossa, Benth. in Fl. Austr. IV, 486 (1869) ... — — — — — — — — N.A. B.fl. IV, 486
 S. viscosa, Roxburgh, pl. of Coromand. II, 33, t. 163 (1798) ... — — — — — W.A. S.A. — — — — — N.A. B.fl. IV, 486
 S. linophylla, F. v. M., fragm. X, 88 (1876) ... — — — — — W.A. — — — — — — — — M.fr. X, 88.
 S. debilis, Benth. in Fl. Austr. IV, 487 (1869) ... — — — — — — — — N.A. B.fl. IV, 487
 S. pedicellaris, F. v. M., fragm. VIII, 231 (1874) ... — — — — — W.A. S.A. — — — — — — — — M.fr. VIII, 231.
 S. Morgania, F. v. M., fragm. X, 89 (1876) ... — — — — — W.A. S.A. — — V. N.S.W. Q. N.A. B.fl. IV, 488 M.fr. VI, 104.
 S. sessiliflora, F. v. M., Limnophila, Blume, Bijdr. 750 (1826) ... — — — — — — — — N.A. — —
 S. gratiolioides, F. v. M., fragm. X, 89 (1876) ... — — — — — — — — Q. N.A. B.fl. IV, 489 M.fr. X, 89.
 S. punctata, F. v. M.; Limnophila, Blume, Bijdr. 750 (1826) ... — — — — — — — — Q. — — B.fl. IV, 490
 S. hirsuta, Heyne in Wallich, numer. list. 3930 (1828) ... — — — — — — — — N.A. B.fl. IV, 490
 S. tenuiflora, Benth. in Scrophul. Indic. 23 (1835) ... — — — — — — — — N.A. B.fl. IV, 490
- BRAMIA, Lamarck, Encycl. méthod. I, 459 (1783). (Herpestis.)
 B. floribunda, F. v. M., fragm. IX, 167 (1875) ... — — — — — — — — Q. N.A. B.fl. IV, 491 M.fr. VI, 104.
 B. Indica, Lamarck, Encycl. méth. I, 459 (1783) ... — — — — — — — — N.S.W. Q. N.A. B.fl. IV, 491 M.fr. IX, 167.
- GRATIOLA, Ruppius, Fl. Jenens. 241 (1718), from Dodoens (1556).
 G. pedunculata, R. Brown, prodr. 435 (1810) ... — — — — — W.A. S.A. — — V. N.S.W. Q. — B.fl. IV, 492 M.fr. VI, 108; IX, 167.
 G. Peruviana, Linné, spec. plant. 17 (1753) ... — — — — — W.A. S.A. T. V. N.S.W. Q. N.A. B.fl. IV, 493 M.fr. VI, 103; IX, 167.
 G. nana, Benth. in De Cand. prodr. X, 404 (1846) ... — — — — — — — — T. V. N.S.W. — — B.fl. IV, 493 M.fr. IX, 167.
- DOPATRIUM, Hamilton in Benth. Scroph. Ind. 4 (1835).
 D. junceum, Hamilton in Benth. Scroph. Ind. 4 (1745) ... — — — — — — — — Q. — — B.fl. IV, 494
- ARTANEMA, D. Don in Sweet, Fl. gard. t. 234 (1829).
 A. fimbriatum, D. Don in Sweet, Fl. Gard. t. 234 (1829) ... — — — — — — — — N.S.W. Q. — B.fl. IV, 495 M.fr. VI, 104; IX, 167.
- LINDERNIA, Allioni, rarior. Pedemont. stirp. specim. t. 5 (1755). (Vandellia, Bonnaya, Ilysanthes, Torenia partly, Tittmannia partly.)
 L. crustacea, F. v. M.; Vandellia, Benth. in Scroph. Indic. 35 (1835) ... — — — — — — — — N.S.W. Q. N.A. — —
 L. pubescens, F. v. M.; Vandellia, Benth. in De Cand. pr. X, 415 (1846) ... — — — — — — — — — — N.A. B.fl. IV, 496 M.fr. VI, 102.
 L. alsinoides, R. Brown, prodr. 441 (1810) ... — — — — — W.A. — — — — — N.S.W. Q. N.A. B.fl. IV, 496
 L. scapigera, R. Brown, prodr. 441 (1810) ... — — — — — — — — — — N.A. B.fl. IV, 497 M.fr. VI, 102.
 L. subulata, R. Brown, prodr. 441 (1810) ... — — — — — — — — — — Q. N.A. B.fl. IV, 497 M.fr. VI, 102.
 L. lobelioides; Vandellia, F. v. M. in Transact. phil. Inst. Vict. III, 61 (1858) ... — — — — — — — — — — N.A. B.fl. IV, 497 M.fr. VI, 102.
 L. clausa, F. v. M., fragm. VI, 102 (1868) ... — — — — — — — — — — N.A. B.fl. IV, 498
 L. veronicifolia, F. v. M., fragm. VI, 101 (1868) ... — — — — — — — — — — Q. — — B.fl. IV, 498 M.fr. VI, 101.
 L. serrata, F. v. M.; Gratiola, Roxburgh. fl. Ind. ed. Car. I, 140 (1820) ... — — — — — — — — — — Q. — —
- HEMIARRHENA, Benth. in Fl. Austr. IV, 518 (1869).
 H. plantaginea, Benth. in Hook. icon. pl. t. 1059 (1869) ... — — — — — — — — — — N.A. B.fl. IV, 518 M.fr. VI, 102.
- PEPLIDIUM, Delile, Fl. Aegypt. 148 (1813).
 P. humifusum, Delile, Fl. Aegypt. 148 (1813) ... — — — — — — — — — — N.S.W. Q. N.A. B.fl. IV, 500 M.fr. VI, 105.
 P. Muellieri, Benth. in Fl. Austr. IV, 500 (1869) ... — — — — — — — — — — N.A. B.fl. IV, 500
- MICROCARPAEA, R. Brown, prodr. 435 (1810).
 M. muscosa, R. Brown, prodr. 435 (1810) ... — — — — — — — — — — Q. N.A. B.fl. IV, 501 M.fr. VI, 104.
- GLOSSOSTIGMA, Arnott in nov. act. acad. Caes. Leop. Carol. XVIII, 355 (1836). (Tricholoma.)
 G. spathulatum, Arnott in nov. act. Leop. XVIII, 355 (1836) ... — — — — — — — — — — Q. — — B.fl. IV, 501
 G. Drummondii, Benth. in De Cand. prodr. X, 426 (1846) ... — — — — — W.A. — — — — — — — — B.fl. IV, 502 M.fr. VI, 104.
 G. elatinoides, Benth. in J. Hook. Fl. New Zeal. I, 189 (1853) ... — — — — — S.A. T. V. N.S.W. — — — — B.fl. IV, 502
- LIMOSELLA, Lindern, opusc. plant. Argentorat. 156, t. 5 (1728).
 L. aquatica, Linné, spec. plant. 631 (1753) ... — — — — — W.A. S.A. T. V. N.S.W. — — — — B.fl. IV, 502 M.fr. VI, 105; IX, 165.
 L. Curdieana, F. v. M., fragm. IX, 166 (1875) ... — — — — — — — — — — S.A. — — V. N.S.W. — — M.fr. IX, 166.
- SCOPARIA, Linné, syst. nat. ed. sext. 87 (1748).
 S. dulcis, Linné, spec. plant. 116 (1753) ... — — — — — — — — — — Q. N.A. B.fl. IV, 504 M.fr. VI, 104.
- OURISIA, Commerçon in A. L. de Jussieu, gen. pl. 100 (1789).
 O. integrifolia, R. Brown, prodr. 439 (1810) ... — — — — — — — — — — T. — — — — B.fl. IV, 512 M.fr. VI, 105; IX, 168.
- VERONICA, Tournefort, inst. 143, t. 60 (1700), from Fuchs (1543). (Pygmaea.)
 V. densifolia, F. v. M., fragm. II, 137, t. 63 (1861) ... — — — — — — — — — — V. N.S.W. — — B.fl. IV, 505 M.fr. VI, 103.
 V. formosa, R. Brown, prodr. 434 (1810) ... — — — — — — — — — — T. — — — — B.fl. IV, 506 M.fr. VI, 102; IX, 167.

V. decorosa, F. v. M. in Schlechtend., <i>Linnaea</i> XXV, 430 (1852)	—	S.A.	—	—	—	—	—	B.f.IV,506	
V. perfoliata, R. Brown, prodr. 434 (1810)	—	—	—	V.	N.S.W.	—	—	B.f.IV,507	M.fr.VI,102;IX,167.
V. Derwentia, Littlejohn in Andr. Bot. Reposit. t. 531 (1808) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV,507	M.fr.VI,102;IX,168.
V. arenaria, Cunningham in De Cand. prodr. X, 463 (1846) ...	—	—	—	—	N.S.W.	—	—	B.f.IV,507	M.fr.VI,102;IX,167.
V. nivea, Lindley, Bot. Regist. Misc. 42 (1842)	—	—	—	T.	V.	N.S.W.	—	B.f.IV,508	M.fr.VI,102;IX,167.
V. gracilis, R. Brown, prodr. 435 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.IV,508	M.fr.VI,103;IX,168.
V. distans, R. Brown, prodr. 435 (1810)	—	W.A.	S.A.	T.	V.	—	—	B.f.IV,509	M.fr.VI,102.
V. calycina, R. Brown, prodr. 435 (1810)	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f.IV,509	M.fr.VI,103.
V. plebeja, R. Brown, prodr. 435 (1810)	—	—	—	T.	V.	N.S.W.	Q.	B.f.IV,510	M.fr.VI,102.
V. notabilis, F. v. M. in Benth. Fl. Austr. IV, 511 (1869)	—	—	—	T.	V.	N.S.W.	Q.	B.f.IV,511	M.fr.IX,168.
V. serpillifolia, Linné, spec. plant. 12 (1753)	—	—	—	—	V.	N.S.W.	—	B.f.IV,511	M.fr.VI,103.
V. peregrina, Linné, spec. plant. 14 (1753)	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f.IV,511	M.fr.VI,103;IX,168.

RAMPHICARPA, Benth. in Hook. Comp. to the Bot. Mag. I, 368 (1835).

R. longiflora, Benth. in Hook. Comp. Bot. Mag. I, 368 (1835)	—	—	—	—	Q.	N.A.	B.f.IV,518	M.fr.VI,104;IX,168.
--	---	---	---	---	----	------	------------	---------------------

CENTRANTHERA, R. Brown, prodr. fl. Nov. Holland. 438 (1810).

C. hispida, R. Brown, prodr. 438 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV,513	M.fr.VI,104;IX,168.
---	---	---	---	---	--------	----	------	------------	---------------------

SOPUBIA, Hamilton in D. Don, prodrom. flor. Nepalens. 88 (1825).

S. trifida, Hamilton in D. Don, prodr. fl. Nep. 88 (1825)	—	—	—	—	—	Q.	—	B.f.IV,513	
---	---	---	---	---	---	----	---	------------	--

BUECHNERA, Linné, hort. Clifford. 501 (1737). (Buchnera, Striga.)

B. tetragona, R. Brown, prodr. 437 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV,514	M.fr.IX,168.
B. urticifolia, R. Brown, prodr. 437 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f.IV,514	M.fr.VI,104.
B. linearis, R. Brown, prodr. 437 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV,515	
B. tenella, R. Brown, prodr. 437 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV,515	
B. gracilis, R. Brown, prodr. 437 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f.IV,515	M.fr.VI,104.
B. ramosissima, R. Brown, prodr. 438 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV,515	
B. coccinea, Benth. in Scrophul. Ind. 40 (1835) ...	—	—	—	—	—	Q.	—	B.f.IV,516	
B. parviflora, R. Brown, prodr. 438 (1810) ...	—	—	—	—	—	Q.	—	B.f.IV,517	M.fr.VI,104.
B. multiflora, F. v. M.; Striga, Benth. in De Cand., prodr. X, 501 (1836) ...	—	—	—	—	—	—	N.A.	B.f.IV,517	
B. curviflora, R. Brown, prodr. 438 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f.IV,517	M.fr.VI,104.

EUPHRASIA, Tournefort, instit. 174, t. 78 (1700), from Matthaeus (1478).

E. Brownii, F. v. M., fragm. V, 88 (1865) ...	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV,521	M.fr.IX,168.
E. scabra, R. Brown, prodr. 437 (1810) ...	—	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.IV,521	M.fr.IX,168.
E. cuspidata, J. Hooker, Fl. Tasman. I, 298 (1860) ...	—	—	—	T.	—	—	—	—	B.f.IV,522	
E. antarctica, Benth. in De Cand., prodr. X, 555 (1846) ...	—	—	—	—	V.	N.S.W.	—	—	B.f.IV,522	

OROBANCHEAE.

A. L. de Jussieu in Ann. Mus. V, 25 (1804).

OROBANCHE, Tournefort, inst. 175, t. 81 (1700), from l'Ecluse (1583).

O. Australiana, F. v. M., inedit. ...	—	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.IV,533	
---------------------------------------	---	------	------	---	----	--------	---	---	------------	--

LENTIBULARINAE.

L. C. Richard in Bulliard, Fl. Paris. ed. sec. I, 26 (1796).

UTRICULARIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 5 (1737).

U. stellaris, Linné, fil. suppl. plant. 86 (1781) ...	...	...	...	...	...	Q.	N.A.	B.f.IV,525	M.fr.VI,161.
U. flexuosa, Vahl, enum. plant. I, 198 (1804) ...	...	...	W.A.	S.A.	T. V.	N.S.W.	Q.	N.A.	B.f.IV,525
U. exoleta, R. Brown, prodr. 430 (1810) ...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.IV,526
U. tubulata, F. v. M., fragm. IX, 48 (1875) ...	...	...	—	—	—	—	Q.	—	—
U. albiflora, R. Brown, prodr. 431 (1810) ...	...	...	—	—	—	—	Q.	—	B.f.IV,526
U. pygmaea, R. Brown, prodr. 432 (1810) ...	...	...	—	—	—	—	Q.	—	B.f.IV,526
U. fulva, F. v. M. in Transact. phil. Inst. Vict. III, 63 (1858)...	...	...	—	—	—	—	—	N.A.	B.f.IV,527
U. chrysanthra, R. Brown, prodr. 432 (1810) ...	...	...	—	—	—	—	Q.	N.A.	B.f.IV,527
U. bifida, Linné, spec. plant. 18 (1753) ...	...	...	—	—	—	—	Q.	—	B.f.IV,527
U. cyanea, R. Brown, prodr. 431 (1810) ...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.f.IV,527
U. lateriflora, R. Brown, prodr. 431 (1810)...	...	...	—	—	T. V.	N.S.W.	—	—	B.f.IV,528
U. simplex, R. Brown, prodr. 431 (1810) ...	...	...	W.A.	—	—	—	—	—	B.f.IV,528
U. monantha, J. Hooker, Fl. Tasman. I, 299 (1860) ...	...	...	—	—	T.	—	—	—	B.f.IV,528
U. dichotoma, Labillard., Nov. Holl. pl. spec. I, 11, t. 8 (1804) ...	...	...	—	S.A.	T. V.	N.S.W.	Q.	—	B.f.IV,529
U. volubilis, R. Brown, prodr. 430 (1810) ...	...	...	W.A.	—	—	—	—	—	B.f.IV,529
U. Hookeri, Lehmann, nov. stirp. pugill. VIII, 47 (1844) ...	...	...	W.A.	—	—	—	—	—	B.f.IV,530
U. violacea, R. Brown, prodr. 431 (1810) ...	...	...	W.A.	—	—	—	—	—	B.f.IV,530
U. Menziesii, R. Brown, prodr. 431 (1810)...	...	...	W.A.	—	—	—	—	—	B.f.IV,530
U. biloba, R. Brown, prodr. 432 (1810) ...	...	...	—	—	—	N.S.W.	—	—	B.f.IV,531
U. limosa, R. Brown, prodr. 432 (1810) ...	...	...	—	—	—	—	Q.	—	B.f.IV,531
U. Baueri, R. Brown, prodr. 431 (1810) ...	...	...	—	—	—	—	Q.	—	B.f.IV,531

POLYPOMPHOLYX, Lehmann in der Bot. Zeitung, Halle, 109 (1844). (Tetralobus.)

P. multifida, F. v. M., fragm. VI, 162 (1868) ...	—	W.A.	—	—	—	—	—	—	B.f.IV,532	M.fr.VI,162.
P. tenella, Lehmann, nov. stirp. pugill. VIII, 50 (1844) ...	—	W.A.	S.A.	T.	V.	—	—	—	B.f.IV,532	M.fr.VI,162.

GESNERIACEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. pl. II, 392 (1817).

FIELDIA, Cunningham in Field's New South Wales 363, t. 2 (1825).

F. australis, Cunningham in Field's N.S.W. 364, t. 2 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.f.IV,535	M.fr.IX,191.
---	---	---	---	----	--------	---	---	------------	--------------

NEGRIA, F. v. M., fragm. VII, 151 (1871). (Rhabdotheramnus partly.)

N. rhabdotheramnoides, F. v. M., fragm. VII, 152 (1871) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,152;XI,133.
---	---	---	---	---	--------	---	---	---	----------------------

BAEA, Commerçon in Lamarck, Encycl. méth. I, 401 (1783). (Streptocarpus.)

B. hygroscopica, F. v. M., fragm. IV, 146 (1864) ...	—	—	—	—	—	—	Q.	—	B.f.IV,535	M.fr.IV,146.
--	---	---	---	---	---	---	----	---	------------	--------------

BIGNONIACEAE.

Ventenat, Tabl. du règn. végét. II, 402 (1799).

TECOMA, A. L. de Jussieu, gen. plant. 139 (1789).

T. australis, R. Brown, prodr. 471 (1810) ...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,537	
T. Hillii, F. v. M., fragm. X, 101 (1877) ...	...	...	...	—	—	—	—	—	—	Q.	—	M.fr.X,101;XI,136.
T. jasminoides, Lindley, Bot. Regist. t. 2002 (1838) ...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,537	

DOLICHANDRONE, Fenzl in Ann. nat. hist. third ser. X, 31 (1862). (Spathodea partly, Dolichandra partly.)

D. heterophylla, F. v. M., fragm. IV, 149 (1864)...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,539	M.fr.IV,149.
D. filiformis, Seemann, Journ. of Bot. I, 226 (1863) ...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.IV,539	

HAUSSMANNIA, F. v. M., fragm. IV, 148 (1864).

H. jucunda, F. v. M., fragm. IV, 148 (1864) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,540	M.fr.IV,148.
---	-----	-----	-----	---	---	---	---	---	----	---	-------------	--------------

DIPLANTHERA, R. Brown, prodr. Fl. Nov. Holl. 449 (1810). (Deplanchea, Bulweria.)

D. tetraphylla, R. Brown, prodr. 449 (1810) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,540	
---	-----	-----	-----	---	---	---	---	---	----	---	-------------	--

ACANTHACEAE.

A. L. de Jussieu, gen. plant. 102 (1789), from B. de Jussieu (1759).

THUNBERGIA, Retzius, Physiogr. Saellsk. Handl. I, 163 (1776).

T. Arnheimica, F. v. M., fragm. IX, 73 (1875) ...	...	...	...	—	—	—	—	—	N.A.	—		M.fr.IX,73.
T. Powellii, F. v. M. in Wing's South. Sc. Rec. II, 34 (1882) ...	...	...	...	—	—	—	—	—	N.A.	—		

NELSONIA, R. Brown, prodr. fl. Nov. Holl. 485 (1810).

N. campestris, R. Brown, prodr. 481 (1810) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,543	M.fr.VI,88;XI,18.
--	-----	-----	-----	---	---	---	---	---	----	------	-------------	-------------------

EBERMAYERA, Nees in Wallich, pl. Asiat. rar. III, 75 (1832).

E. glauca, Nees in De Candolle, prodr. XI, 73 (1847) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,544	M.fr.VI,88;XI,18
--	-----	-----	-----	---	---	---	---	---	----	------	-------------	------------------

HYGROPHILA, R. Brown, prodr. Fl. Nov. Holl. 479 (1810).

H. angustifolia, R. Brown, prodr. 479 (1810) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,544	M.fr.VI,88.
--	-----	-----	-----	---	---	---	---	---	----	------	-------------	-------------

STROBILANTHES, Blume, Bijdr. tot de fl. van Nederl. Ind. 796 (1826).

S. Tatei, F. v. M. in transact. Roy. Soc. S. Austr. V, Aug. (1882) ...	...	...	...	—	—	—	—	—	N.A.	—		
--	-----	-----	-----	---	---	---	---	---	------	---	--	--

RUELLIA, Plumier, nov. gen. 12, t. 2 (1703). (Dipteracanthus.)

R. bracteata, R. Brown, prodr. 479 (1810)...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,546	M.fr.VI,91.
R. primulacea, F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ...	...	...	...	—	S.A.	—	—	—	Q.	—	B.fl.IV,546	M.fr.XI,18.
R. corynotheca, F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,546	M.fr.XI,18.
R. australis, R. Brown, prodr. 479 (1810) ...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,547	M.fr.VI,91.
R. spiciflora, F. v. M. in Benth. Fl. Austr. IV, 547 (1869) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,547	
R. acaulis, R. Brown, prodr. 479 (1810) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,547	

ACANTHUS, Tournefort, inst. 176, t. 81 (1700), from Theophrastos. (Dilivaria.)

A. ilicifolius, Linné, spec. plant. 939 (1753) ...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,548	M.fr.X,18.
--	-----	-----	-----	---	---	---	---	---	----	------	-------------	------------

JUSTICIA, Houston in Linné, gen. plant. 4 (1737). (Rostellaria, Rostellularia.)

J. procumbens, Linné, spec. plant. 15 (1753) ...	...	...	...	...	W.A.	S.A.	—	N.S.W.	Q.	N.A.	B.fl.IV,549	M.fr.VI,91;XI,18.
J. Bonneyana, F. v. M. in Wing's S. Sc. Record II, 74 (1882) ...	...	...	...	...	S.A.	—	—	N.S.W.	—	—	—	M.fr.X,73.
J. hygrophiloides, F. v. M., fragm. VI, 89 (1868)...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,550	M.fr.VI,89.
J. Kempeana, F. v. M., fragm. XI, 101 (1881) ...	...	...	...	...	W.A.	S.A.	—	—	—	N.A.	—	M.fr.XI,101.
J. cavernarum, F. v. M., fragm. VI, 91 (1868) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,550	M.fr.VI,91.
J. eranthemoides, F. v. M., fragm. VI, 90 (1868)...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.IV,551	M.fr.VI,90.

GRAPTOPHYLLUM, Nees in Wall. pl. Asiat. rar. III, 102 (1832). (Earlia, Thyracanthus.)

G. Earlii, F. v. M., fragm. VI, 87 (1868) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,551	M.fr.VI,87.
G. ilicifolium, F. v. M. in Benth. Fl. Austr. IV, 552 (1869) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,552	
G. spinigerum, F. v. M., fragm. XI, 17 (1878) ...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.XI,17.

DICLIPTERA, A. L. de Jussieu in Ann. du Mus. IX, 267 (1807). (Brochosiphon.)

D. glabra, Decaisne in Nouv. Annal. du Mus. III, 388 (1834) ...	—	—	—	—	—	—	—	—	—	N.A.	B.fl.IV,552	M.fr.VI,88;XI,18.	
D. spicata, Decaisne in Nouv. Annal. du Mus. III, 389 (1834) ...	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.IV,553	M.fr.VI,89.
D. Burmanni, Nees in Wallich, pl. Asiat. rar. III, 112 (1832) ...	—	—	—	—	—	—	—	—	—	Q.	—	—	M.fr.VII,62.

HYPOESTES, Solander in R. Brown, prodr. 474 (1810).

H. floribunda, R. Brown, prodr. 474 (1810) ...	...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,554	M.fr.XI,18.
--	-----	-----	-----	-----	---	---	---	--------	----	------	-------------	-------------

ERANTHEMUM, Linné, Fl. Zeyl. 6 (1747).

E. variabile, R. Brown, prodr. 477 (1810) ...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.IV,555	M.fr.VI,91;XI,18.
E. tenellum, Benth, Fl. Austr. IV, 555 (1869) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,555	

HYDROPHYLLEAE.

R. Brown in Bot. Regist. t. 242 (1817).

HYDROLEA, Linné, spec. plant. ed. sec. 328 (1762).

H. Zeilanica, Vahl, symbol. botan. II, 46 (1791) ...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.IV,382	M.fr.V,192.
H. spinosa, Linné, spec. plant. ed. sec. 328 (1762) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,383	M.fr.V,192.

ASPERIFOLIAE.

Haller, enum. stirp. Helv. Praef. 34 (1742).

CORDIA, Plumier, gen. plant. 13, t. 14 (1703).

C. subcordata, Lamarck, Illustr. des genr. I, 421 (1791)...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.IV,385	M.fr.VI,114;IX,122.[122.
C. aspera, G. Forster, florul. ins. austr. prodr. 18 (1786)...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.V,193;VI,114;IX,
C. Myxa, Linné, spec. plant. 190 (1753) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.I,59;VI,114.

EHRETIA, P. Browne, Civ. and Nat. Hist. of Jamaic. 168, t. 16 (1756).

EURELIA, F. Browne, Civ. and Nat. Hist. of Samoa, 103, t. 16 (1930).													
E. acuminata, R. Brown, prodr. 497 (1810)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,387	M.fr.V,21;IX,122;XI,124
E. pilosula, F. v. M., fragm. V, 20 (1865)...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,388	M.fr.V,20.
E. saligna, R. Brown, prodr. 497 (1810)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,388	M.fr.V,21;IX,122.

- E. membranifolia*, R. Brown, prodr. 497 (1810) — — — — — Q. — B.f. IV, 388 M.fr. V, 21; VI, 115; IX, 122
E. laevis, Roxburgh, Pl. Corom. I, 42, t. 56 (1795) — — — — — Q. — B.f. IV, 389
- TOURNEFORTIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 55 (1737.)
- T. argentea*, Linné fil. suppl. plant. 133 (1781) — — — — — Q. N.A. B.f. IV, 389 M.fr. IV, 95; VI, 116.
T. mollis, F. v. M., fragm. I, 59 (1858) — — — — — Q. N.A. B.f. IV, 390 M.fr. VI, 116.
T. sarmentosa, Lamarck, Illustr. des genr. I, 416 (1791) — — — — — Q. — B.f. IV, 390 M.fr. IV, 95; VI, 116.
- COLDENIA, Linné, amoen. acad. ed. prim. 119 (1747). (Lobophyllum.)
- C. procumbens*, Linné, spec. plant. 125 (1753) — — — — — Q. N.A. B.f. IV, 391 M.fr. VI, 115.
- HELIOTROPIUM, Tournefort, inst. 138, t. 57 (1700), from Theophr., Diosc. & Plinius. (Tiaridium, Heliophytum.)
- H. indicum*, Linné, spec. plant. 130 (1753) — — — — — N.A. — M.fr. XI, 124.
H. pleiopterum, F. v. M., fragm. IX, 121 (1875) — S.A. — — — — — M.fr. IX, 121.
H. curassavicum, Linné, spec. plant. 130 (1753) W.A. S.A. — V. N.S.W. Q. N.A. B.f. IV, 393 M.fr. VI, 116; IX, 121.
H. europaeum, Linné, spec. plant. 130 (1753) — S.A. — V. N.S.W. — — B.f. IV, 394 M.fr. VI, 116. [124.
H. undulatum, Vahl, symbol. bot. I, 13 (1795) W.A. S.A. — — — — — N.A. B.f. IV, 394 M.fr. VI, 116; IX, 121; XI,
H. asperrimum, R. Brown, prodr. 493 (1810) W.A. S.A. — V. N.S.W. — — B.f. IV, 394 M.fr. VI, 116; IX, 121.
H. crispatum, F. v. M. in Benth. Fl. Austr. IV, 395 (1869) — — — — — N.A. B.f. IV, 395
H. fasciculatum, R. Brown, prodr. 494 (1810) — — — — — Q. N.A. B.f. IV, 395
H. vestitum, Benth. Fl. Austr. IV, 395 (1869) — — — — — N.S.W. — — B.f. IV, 395
H. brachygyne, Benth. Fl. Austr. IV, 396 (1869) — — — — — Q. — B.f. IV, 396
H. epacrideum, F. v. M. in Benth. Fl. Austr. IV, 396 (1869) — — — — — N.A. B.f. IV, 396
H. ovalifolium, Forskael, Fl. Aegypt. Arab. 47 (1775) — S.A. — — — — — Q. N.A. B.f. IV, 396 M.fr. VI, 115.
H. strigosum, Willdenow, sp. pl. I, 743 (1797) — — — — — Q. N.A. B.f. IV, 397
H. prostratum, R. Brown, prodr. 494 (1810) — — — — — N.A. B.f. IV, 397
H. bracteatum, R. Brown, prodr. 493 (1810) — — — — — Q. N.A. B.f. IV, 397
H. pauciflorum, R. Brown, prodr. 493 (1810) — — — — — Q. N.A. B.f. IV, 398
H. conocarpum, F. v. M. in Benth. Fl. Austr. 398 (1869) — — — — — N.A. B.f. IV, 398
H. filaginoides, Benth. Fl. Austr. IV, 398 (1869) — S.A. — — — — — B.f. IV, 398
H. ventricosum, R. Brown, prodr. 494 (1810) — — — — — Q. N.A. B.f. IV, 399
H. tenuifolium, R. Brown, prodr. 494 (1810) — — — — — Q. N.A. B.f. IV, 399
H. paniculatum, R. Brown, prodr. 494 (1810) — — — — — N.A. B.f. IV, 399
H. Cunninghamii, Benth. Fl. Austr. IV, 400 (1869) — — — — — N.A. B.f. IV, 400
H. diversifolium, F. v. M. in Benth. Fl. Austr. IV, 400 (1869) — — — — — N.A. B.f. IV, 400
- HALGANIA, Gaudichaud in Freyc. in voy. Bot. 448 (1826).
- H. solanacea*, F. v. M. in Hook. Kew Misc. IX, 21 (1857) — — — — — N.A. B.f. IV, 401 M.fr. VI, 115.
H. litoralis, Gaudichaud in Freyc. voy. Bot. 449, t. 59 (1826) ... W.A. — — — — — B.f. IV, 401 M.fr. VI, 115.
H. corymbosa, Lindley, Bot. Regist. XXV, App. XL (1839) ... W.A. — — — — — B.f. IV, 402
H. sericiflora, Benth. Fl. Austr. IV, 402 (1869) W.A. — — — — — B.f. IV, 402
H. cyanea, Lindley, Bot. Regist. XXV, App. XL (1839) ... W.A. S.A. — V. N.S.W. Q. N.A. — M.fr. VI, 115; IX, 122.
H. lavandulacea, Endlicher in Ann. Wien. Mus. II, 205 (1838) W.A. S.A. — V. N.S.W. — — B.f. IV, 402 M.fr. I, 209; VI, 115; XI, 125
H. integriloba, Endlicher in Ann. Wien. Mus. II, 205 (1838) ... W.A. — — — — — B.f. IV, 402
- POLLICHIA, Medikus, Botan. Beobach. 247 (1783). (Trichodesma.)
- P. zeylanica*, F. v. M.; Trichodesma, R. Brown, prodr. 496 (1810) W.A. S.A. — — N.S.W. Q. N.A. B.f. IV, 404 M.fr. XI, 125.
P. latispalea; Trichodesma, F. v. M., fragm. X, 102 (1877) ... — — — — — — — M.fr. X, 121.
- MYOSOTIS, Rupp. Fl. Jenens. 9 (1718), from Dioscorides and Plinius. (Exarrhena.)
- M. australis*, R. Brown, prodr. 495 (1810) W.A. S.A. T. V. N.S.W. — — B.f. IV, 405 M.fr. VI, 116; IX, 122.
M. suaveolens, Poir. Encycl. méth. suppl. IV, 44 (1816) — — T. V. N.S.W. — — B.f. IV, 406 M.fr. VI, 115; IX, 122.
- ERITRICHUM, Schrader in commentat. Goetting. IV, 186 (1820). (Eritrichium.)
- E. Australasicum*, A. De Candolle, prodr. X, 134 (1846) ... W.A. S.A. — V. N.S.W. — — B.f. IV, 406 M.fr. XI, 125.
- LAPPULA, Rivinus in Rupp. Fl. Jenens. (1718), from Dalechamp. (Echinosperrum.)
- L. concava*; Echinosperrum, F. v. M., fragm. II, 139 (1861) ... W.A. S.A. — V. N.S.W. — — B.f. IV, 407 M.fr. VI, 116; IX, 121.
- ROCHELIA, Reichenbach in Fl. Regensb. bot. Zeit. I, 243 (1824).
- R. Maccoya*, F. v. M. in Benth. Fl. Austr. IV, 408 (1869) — S.A. — V. N.S.W. — — B.f. IV, 408 M.fr. I, 127; IX, 121.
- CYNOGLOSSUM, Tournefort, inst. 139, t. 57 (1700), from Dioscorides and Nicandros. [125.
C. latifolium, R. Brown, prodr. 495 (1810) — — T. V. N.S.W. Q. — B.f. IV, 408 M.fr. VI, 115; IX, 122; XI,
C. suaveolens, R. Brown, prodr. 495 (1810) — S.A. T. V. N.S.W. — — B.f. IV, 409 M.fr. VI, 115; IX, 122.
C. australe, R. Brown, prodr. 495 (1810) — S.A. T. V. N.S.W. Q. — B.f. IV, 409 M.fr. VI, 115; IX, 122.
C. Drummondii, Benth. Fl. Austr. IV, 409 (1869) W.A. S.A. — — — — — B.f. IV, 409 M.fr. IX, 122.
- LABIATAE.
- Adanson, Famil. des plant. II, 180 (1763), from B. de Jussieu (1759).
- OCIMUM, Tournefort, inst. 203, t. 96 (1700), from Theophrastus and Dioscorides. (Ocimum.)
- O. sanctum*, Linné, mantiss. plant. 85 (1767) — — — — — Q. N.A. B.f. V, 74 M.fr. IV, 46.
- MOSCHOSMA, Reichenbach, conspect. regn. vegetab. 115 (1828). (Basilicum 1802.)
- M. polystachya*, Benth. in Wall. pl. As. rar. II, 13 (1831) — — — — — N.S.W. Q. N.A. B.f. V, 75 M.fr. VI, 108.
- ORTHOSIPHON, Benth. Labiatar. gen. et spec. 25 (1832).
- O. stamineus*, Benth. in Wall. pl. As. rar. II, 15 (1831) — — — — — Q. N.A. B.f. V, 76 M.fr. VI, 108; IX, 161.
- PLECTRANTHUS, L'Héritier, stirp. nov. I, 85 (1785).
- P. longicornis*, F. v. M., fragm. V, 51 (1865) — — — — — Q. — B.f. V, 77 M.fr. V, 51.
P. parviflorus, Willdenow, hort. Berol. I, t. 65 (1806) — S.A. — V. N.S.W. Q. N.A. B.f. V, 78 M.fr. VI, 108; IX, 161.
P. congestus, R. Brown, prodr. 506 (1810) — — — — — Q. — B.f. V, 78 M.fr. VI, 108.
P. foetidus, Benth. Labiat. 35 (1832) — — — — — Q. — — M.fr. IX, 161; XI, 135.
- COLEUS, Loureiro, Fl. Cochinch. II, 372 (1790).
- C. scutellarioides*, Benth. in Wall. pl. Asiat. rar. II, 16 (1831) — — — — — Q. N.A. B.f. V, 79 M.fr. VI, 108.
- POGOSTEMON, Desfontaines in Mém. du Mus. II, 154, t. 6 (1815). (Dysophylla.)
- P. verticillatus*, Hasskarl, catal. hort. Bogor. 131 (1844) — — — — — Q. N.A. B.f. V, 81 M.fr. V, 200; IX, 161.

MENTHA, Tournefort, inst. 188, t. 89 (1700), from Hippocrates and Theophrastos.									
M. laxiflora, Benth in De Cand. XII, 174 (1848)	...	...	—	—	—	V. N.S.W.	—	B.fl.V,82	M.fr.VI,109;IX,162.
M. grandiflora, Benth in Mitch. Trop. Austr. 362 (1848)	...	...	—	—	—	N.S.W.	Q.	B.fl.V,83	
M. australis, R. Brown, prodr. 505 (1810)...	...	...	—	—	S.A. T. V. N.S.W.	Q.	N.A.	B.fl.V,83	M.fr.VI,109;IX,162.
M. gracilis, R. Brown, prodr. 505 (1810)	...	...	—	—	S.A. T. V. N.S.W.	—	—	B.fl.V,83	M.fr.VI,109.
M. saturejoides, R. Brown, prodr. 505 (1810)	...	...	...	...	W.A. S.A. T. V. N.S.W.	Q.	—	B.fl.V,84	M.fr.VI,109.
LYCOPUS, Tournefort, inst. 180, t. 83 (1700), from Plinius.									
L. australis, R. Brown, prodr. 500 (1810)	...	...	—	—	S.A. T. V. N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109.
SALVIA, Linné, gen. plant. 6 (1737), from Plinius.									
S. plebeja, R. Brown, prodr. 501 (1810)	...	...	—	—	—	V. N.S.W.	Q.	B.fl.V,85	M.fr.VI,109;IX,161.
BRUNELLA, L'Ecluse, rar. stirp. hist. II, 42, 43 (1576), from Brunfels (1536). (Prunella.)									
B. vulgaris, A. de Candolle, prodr. XII, 410 (1848)	...	...	—	—	S.A. T. V. N.S.W.	Q.	—	B.fl.V,87	M.fr.VI,109.
SCUTELLARIA, Hermann, hort. Lugd. bot. cat. (1687), from Cortusi (1591).									
S. mollis, R. Brown, prodr. 507 (1810)	...	...	—	—	—	V. N.S.W.	—	B.fl.V,88	M.fr.VI,109.
S. humilis, R. Brown, prodr. 507 (1810)	...	...	—	—	S.A. T. V. N.S.W.	Q.	—	B.fl.V,88	M.fr.VI,109.
ANISOMELES, R. Brown, prodr. 503 (1810).									
A. salvifolia, R. Brown, prodr. 503 (1810)...	...	...	—	—	—	—	Q. N.A.	B.fl.V,89	M.fr.VI,109;IX,162.
LEUCAS, J. Burmann, thesaur. Zeyl. 140 (1737).									
L. flaccida, R. Brown, prodr. 505 (1810)	...	...	—	—	—	—	Q. N.A.	B.fl.V,90	M.fr.VI,109.
DEPREMESNILIA, F. v. M., fragm. X, 59 (1876).									
D. chrysocalyx, F. v. M., fragm. X, 59 (1876)	...	...	W.A.	—	—	—	—	—	M.fr.X,59.
PROSTANTHERA, Labillardière, Nov. Holl. pl. spec. 18, t. 157 (1806). (Chilodia, Cryphia, Klanderia.)									
P. lasiantha, Labillard., Nov. Holl. pl. spec. II, 18, t. 15 (1806)	...	...	—	S.A. T. V. N.S.W.	Q.	—	—	B.fl.V,93	M.fr.VI,105.
P. prunelloides, B. Brown, prodr. 508 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,94	M.fr.VI,106.
P. coerulea, R. Brown, prodr. 508 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,94	M.fr.VI,106.
P. ovalifolia, R. Brown, prodr. 509 (1810)...	...	...	—	—	—	N.S.W.	Q.	B.fl.V,95	M.fr.VI,106.
P. melissifolia, F. v. M., fragm. I, 19 (1858)	...	...	—	—	—	V.	—	B.fl.V,95	M.fr.I,19,242.
P. incisa, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	Q.	B.fl.V,95	M.fr.VI,107;IX,162.
P. rotundifolia, R. Brown, prodr. 509 (1810)	...	...	—	S.A. T. V. N.S.W.	—	—	—	B.fl.V,96	M.fr.VI,107;IX,162.
P. violacea, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,96	M.fr.VI,107;IX,162.
P. incana, Cunningham in Benth. lab. 455 (1836)...	...	...	—	—	—	N.S.W.	—	B.fl.V,97	M.fr.IX,162.
P. hirtula, F. v. M., first gen. Report 16 (1853)	...	...	—	—	—	V. N.S.W.	—	B.fl.V,97	
P. denticulata, R. Brown, prodr. 509 (1810)	...	...	—	—	—	V. N.S.W.	—	B.fl.V,97	M.fr.VI,105.
P. rugosa, Cunningham in Benth. lab. 456 (1836)...	...	...	—	—	—	N.S.W.	—	B.fl.V,98	
P. marifolia, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,98	M.fr.VI,105.
P. rhombea, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,99	
P. spinosa, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	...	...	—	S.A.	—	V.	—	B.fl.V,99	M.fr.VI,108;IX,162.
P. cuneata, Benth in De Cand., prodr. XII, 560 (1848)	...	...	—	—	T. V. N.S.W.	—	—	B.fl.V,99	M.fr.VI,108.
P. linearis, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	—	B.fl.V,100	M.fr.VI,106.
P. phyllofolia, F. v. M., fragm. I, 19 (1858)	...	...	—	—	—	V. N.S.W.	Q.	B.fl.V,100	M.fr.I,19;IX,162.
P. decussata, F. v. M., fragm. I, 126 (1858)	...	...	—	—	—	V.	—	B.fl.V,100	M.fr.I,126.
P. empetrifolia, Sieber in Spreng. syst. cur. post. 226 (1827)	...	...	—	—	—	N.S.W.	—	B.fl.V,101	
P. lithospermoides, F. v. M., fragm. VI, 107 (1868)	...	...	—	—	—	—	Q.	B.fl.V,101	M.fr.VI,107.
P. Behriana, Schlechtendal, Linnaea XX, 610 (1847)	...	...	—	S.A.	—	V. N.S.W.	—	B.fl.V,102	M.fr.VI,106.
P. Wilkieana, F. v. M., fragm. VIII, 230 (1874)...	...	...	—	S.A.	—	—	—	—	M.fr.VIII,230.
P. Baxteri, Cunningham in Benth. lab. 452 (1836)	...	...	W.A.	S.A.	—	—	—	B.fl.V,102	M.fr.VI,106.
P. canaliculata, F. v. M., fragm. VI, 105 (1868)	...	...	W.A.	—	—	—	—	B.fl.V,102	M.fr.VI,105.
P. nivea, Cunningham in Benth. lab. 452 (1836)	...	...	—	—	—	V. N.S.W.	Q.	B.fl.V,103	M.fr.VI,106.
P. striatiflora, F. v. M. in Schlechtend. Linnaea XXV, 425 (1852)	...	...	—	S.A.	—	N.S.W.	Q. N.A.	B.fl.V,103	M.fr.VI,105;IX,162.
P. saxicola, R. Brown, prodr. 509 (1810)	...	...	—	—	—	N.S.W.	Q.	B.fl.V,104	M.fr.VI,107;IX,162.
P. debilis, F. v. M., fragm. VIII, 147 (1874)	...	...	—	—	—	V.	—	—	M.fr.VIII,147.
P. odoratissima, Benth in Mitch. Trop. Austr. 291 (1848)	...	...	—	—	—	—	Q.	B.fl.V,104	
P. Eckersleyana, F. v. M., fragm. X, 17 (1876)	...	...	W.A.	—	—	—	—	—	M.fr.X,17.
P. euphrasioides, Benth in Mitch. Trop. Austr. 360 (1848)	...	...	—	—	—	N.S.W.	Q.	B.fl.V,104	M.fr.VI,108.
P. staurophylla, F. v. M., fragm. IX, 73 (1875)	...	...	—	—	—	N.S.W.	—	—	M.fr.IX,73.
P. cryptandroides, Cunningham in Benth. lab. 453 (1836)	...	...	—	—	—	N.S.W.	—	B.fl.V,105	
P. eurybioides, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	...	...	—	S.A.	—	—	—	B.fl.V,105	M.fr.VI,105.
P. ringens, Benth in Mitch. Trop. Austr. 363 (1848)	...	...	—	—	—	N.S.W.	Q.	B.fl.V,106	
P. Leichhardtii, Benth in Fl. Austr. V, 106 (1870)	...	...	—	—	—	—	Q.	B.fl.V,106	
P. Walteri, F. v. M., fragm. VII, 108 (1870)	...	...	—	—	—	V. N.S.W.	—	—	M.fr.VII,108.
P. coccinea, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)...	...	...	W.A.	S.A.	—	V. N.S.W.	Q.	B.fl.V,106	M.fr.VI,108;IX,162.
P. chlorantha, F. v. M. in Benth. Fl. Austr. V, 108 (1870)	...	...	—	S.A.	—	V. N.S.W.	—	B.fl.V,108	M.fr.IX,162;X,18.
P. calycina, F. v. M. in Benth. Fl. Austr. V, 107 (1870)	...	...	—	S.A.	—	—	—	B.fl.V,107	
P. Grylloana, F. v. M., fragm. X, 17 (1876)	...	...	W.A.	—	—	—	—	—	M.fr.X,17.
WRIXONIA, F. v. M., fragm. X, 18 (1876).									
W. prostantheroides, F. v. M., fragm. X, 18 (1876)	...	...	W.A.	—	—	—	—	—	M.fr.X,18.
HEMIGENIA, R. Brown, prodr. 502 (1810). (Hemiandra, Colobandra, Atelandra.)									
H. macrantha, F. v. M., fragm. I, 210 (1859)	...	...	W.A.	—	—	—	—	B.fl.V,112	M.fr.I,210;XI,19.
H. rigida, Benth in De Cand. prodr. XII, 563 (1848)...	...	...	W.A.	—	—	—	—	B.fl.V,112	
H. ramosissima, Benth in De Cand. prodr. XII, 565 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,113	
H. microphylla, Benth in De Cand. prodr. XII, 565 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,113	M.fr.VI,112;IX,162.
H. incana, Benth in De Cand. prodr. XII, 566 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,113	
H. canescens, Benth in De Cand. prodr. XII, 566 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,114	
H. podalyrina, F. v. M., fragm. VI, 112 (1868)	...	...	W.A.	—	—	—	—	B.fl.V,114	M.fr.VI,112.
H. platyphylla, Benth in De Cand. prodr. XII, 566 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,115	
H. obovata, F. v. M., fragm. XI, 19 (1878)	...	...	W.A.	—	—	—	—	—	B.fl.XI,19
H. glabrescens, Benth in De Cand. prodr. XII, 566 (1848)	...	...	W.A.	—	—	—	—	B.fl.V,115	

N. chrysotricha, F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 15.
N. hexarrhena, F. v. M., fragm. X, 16 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 16.
N. spodiotricha, F. v. M., fragm. III, 21, t. 21 (1862)	W.A.	S.A.	—	—	—	N.A.	B.fl. V, 40	M.fr. VIII, 230; IX, 4.
N. cephalantha, F. v. M., fragm. IX, 4 (1875)	—	S.A.	—	—	—	—	—	M.fr. IX, 4.
PHYSOPSIS, Turczaninow in Bullet. de la Soc. Imper. des Natur. de Mosc. XXII, 34 (1849).								
P. spicata, Turczaninow in Bull. de Mosc. XXII, 35 (1849)	W.A.	—	—	—	—	—	B.fl. V, 41	M.fr. VI, 154.
MALLOPHORA, Endlicher in Annal. des Wien. Mus. II, 206 (1838). (Lachnocephalus.)								
M. globiflora, Endlicher in Ann. des Wien. Mus. II, 206 (1838)	W.A.	—	—	—	—	—	B.fl. V, 41	M.fr. VI, 154; X, 15.
DICRASTYLIS, Drummond & Harvey in Hook. Kew Misc. VII, 56 (1855). (Dicrastyles.)								
D. Doranii, F. v. M., fragm. VIII, 230 (1874)	—	S.A.	—	—	—	—	—	M.fr. VIII, 230.
D. ochrotricha, F. v. M., fragm. IV, 161 (1864)	—	S.A.	—	—	—	N.A.	B.fl. V, 42	M.fr. I, 60; VI, 155; VIII, 230
D. Gilesii, F. v. M., fragm. VIII, 229 (1874)	—	S.A.	—	—	—	—	—	M.fr. X, 15.
D. fulva, Drummond & Harvey in Kew Misc. VII, 56 (1855)	W.A.	—	—	—	—	—	B.fl. V, 43	M.fr. I, 236; VI, 155.
D. reticulata, Drum. & Harvey in Hook. Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	B.fl. V, 43	M.fr. VI, 155; IX, 4.
D. Nicholasii, F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 15.
D. parvifolia, F. v. M. fragm. II, 16 (1861)	W.A.	—	—	—	—	—	B.fl. V, 43	M.fr. VI, 155; X, 15.
D. Beveridgei, F. v. M., fragm. VIII, 50 (1873)	—	S.A.	—	—	—	—	—	M.fr. IX, 4.
D. Stoechas, Drummond & Harvey in Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	B.fl. V, 44	M.fr. VI, 155.
D. Lewellini, F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	—	—	—	—	M.fr. VIII, 50; XI, 137.
CHLOANTHES, R. Brown, prodr. Flor. Nov. Holl. 513 (1810). (Pithyrodia, Quoya, Dasymalla.)								
C. Stoechadis, R. Brown, prodr. 514 (1810)	—	—	—	—	N.S.W.	—	B.fl. V, 45	M.fr. VI, 156.
C. parviflora, Walpers, repertor. bot. syst. IV, 58 (1848)	—	—	—	—	N.S.W.	Q.	B.fl. V, 46	M.fr. IX, 5.
C. coccinea, Bartling in Lehm. pl. Preiss. I, 352 (1845)	W.A.	—	—	—	—	—	B.fl. V, 48	M.fr. VI, 156; IX, 5.
C. salvifolia, F. v. M.; Pithyrodia, R. Brown, prodr. 513 (1810)	—	—	—	—	—	Q.	B.fl. V, 48	M.fr. VI, 155; IX, 5.
C. hemigenioides, F. v. M., fragm. VI, 156 (1868)	W.A.	—	—	—	—	N.A.	B.fl. V, 48	M.fr. VI, 156.
C. uncinata, Turczaninow in Bull. de Mosc. XXXVI, 194 (1863)	W.A.	—	—	—	—	—	B.fl. V, 48	M.fr. VI, 156.
C. Bartlingii, Lehmann, Ind. sem. hort. Hamb. (1844)	W.A.	—	—	—	—	—	B.fl. V, 49	M.fr. VI, 156.
C. verbascina, F. v. M., fragm. I, 233 (1859)	W.A.	—	—	—	—	—	B.fl. V, 50	M.fr. IV, 80; VI, 157; IX, 5.
C. stachyodes, F. v. M., fragm. V, 50 (1855)	W.A.	—	—	—	—	—	B.fl. V, 50	M.fr. VI, 158.
C. loxocarpa, F. v. M., fragm. II, 22 (1861)	W.A.	—	—	—	—	N.A.	B.fl. V, 51	M.fr. IV, 80; VI, 156.
C. dilatata, F. v. M., fragm. VI, 157 (1868)	W.A.	—	—	—	—	—	B.fl. V, 51	M.fr. VI, 157.
C. cuneata, F. v. M.; Quoya, Gaudich. in Freyc. voy. 454 t. 66 (1826)	W.A.	—	—	—	—	—	B.fl. V, 51	
C. Oldfieldii, F. v. M., fragm. I, 234 (1859)	W.A.	—	—	—	—	—	B.fl. V, 52	M.fr. IV, 80.
C. atriplicina, F. v. M., fragm. I, 235 (1859)	W.A.	—	—	—	—	N.A.	B.fl. V, 52	M.fr. VI, 157; IX, 5.
C. paniculata, F. v. M., fragm. IV, 80 (1864)	W.A.	—	—	—	—	—	B.fl. V, 53	
C. loricata, F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 14.
C. halganicea, F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 14.
HEMIPHORA, F. v. M., fragm. X, 13 (1876).								
H. Elderi, F. v. M., fragm. X, 13 (1876)	W.A.	—	—	—	—	—	—	M.fr. X, 13.
DENISONIA, F. v. M., fragm. I, 124 (1859).								
D. ternifolia, F. v. M., fragm. I, 124, t. 2 (1859)	—	—	—	—	—	N.A.	B.fl. V, 54	M.fr. I, 124.
CYANOSTEGIA, Turczaninow in Bull. Soc. Mosc. XXII, 35 (1849). (Bunnya.)								
C. Turczaninovii, F. v. M., fragm. VI, 154 (1868)	W.A.	—	—	—	—	—	B.fl. V, 54	M.fr. IX, 5.
C. Bunnyana, F. v. M., fragm. V, 36 (1865)	—	—	—	—	—	N.A.	B.fl. V, 54	M.fr. VI, 153.
CALLICARPA, Linné in act. litter. et scienc. Upsal. 80 (1741).								
C. cana, Linné, mantiss. plant. alt. 198 (1771)	—	—	—	—	—	Q.	N.A.	B.fl. V, 56
C. pedunculata, R. Brown, prodr. 513 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 57
C. longifolia, Lamarck, Encycl. méth. I, 562 (1783)	—	—	—	—	—	Q.	—	B.fl. V, 57
PREMNA, Linné, mantiss. plant. alter. 154 (1771).								
P. obtusifolia, R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. V, 58
P. integrifolia, Linné, mant. plant. alt. 154 (1771)	—	—	—	—	—	—	N.A.	B.fl. V, 59
P. limbata, Benth. Fl. Austr. V, 59 (1870)	—	—	—	—	—	—	—	B.fl. V, 59
P. Dallachyana, Benth. Fl. Austr. V, 59 (1870)	—	—	—	—	—	Q.	—	B.fl. V, 59
P. acuminata, R. Brown, prodr. 512 (1810)	—	—	—	—	—	—	N.A.	B.fl. V, 60
CLERODENDRUM, Burmann, thesaur. Zeylan. 66, t. 29 (1737). (Clerodendron.)								
C. hemiderma, F. v. M. in Benth. Fl. Austral. V, 61 (1870)	—	—	—	—	—	Q.	—	B.fl. V, 61
C. inerme, R. Brown, prodr. 511 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. V, 61
C. Tracyanum, F. v. M. in Benth. Fl. Austr. V, 62 (1870)	—	—	—	—	—	Q.	—	B.fl. V, 62
C. tomentosum, R. Brown, prodr. 510 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 62
C. lanceolatum, F. v. M., fragm. III, 145 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl. V, 63
C. floribundum, R. Brown, prodr. 511 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. V, 63
C. Cunninghamii, Benth. Fl. Austr. V, 64 (1870)	—	—	—	—	—	Q.	N.A.	B.fl. V, 64
C. costatum, R. Brown, prodr. 511 (1810)	—	—	—	—	—	Q.	—	B.fl. V, 64
GMELINA, Linné, gen. plant. ed. sec. 526 (1742).								
G. macrophylla, Benth. Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	N.A.	B.fl. V, 65
G. fasciculiflora, Benth. Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	—	B.fl. V, 65
G. Leichhardtii, F. v. M. in Benth. Fl. Austr. V, 66 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 66
VITEX, Tournefort, inst. 603, t. 373 (1700), from Plinius.								
V. trifolia, Linné, spec. plant. 938 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. V, 66
V. Lignum vitae, Cunningham in De Cand. prodr. XI, 692 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl. V, 67
V. acuminata, R. Brown, prodr. 512 (1810)	—	—	—	—	—	—	N.A.	B.fl. V, 67
V. glabrata, R. Brown prodr. 512 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. V, 68
FARADAYA, F. v. M., fragm. phytogr. Austr. V, 21 (1865).								
F. splendida, F. v. M., fragm. V, 21 (1865)	—	—	—	—	—	Q.	—	B.fl. V, 69
AVICENNIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 27 (1737).								
A. officinalis, Linné, spec. plant. 110 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. V, 69

MYOPORUM, Banks & Solander in G. Forster, prodr. 44 (1786). (Bertolonia, Andrewsia, Pogonia, Disoon.)

M. tenuifolium, G. Forster, fl. ins. Austr. prodr. 44 (1786) ...	—	—	—	—	V. N.S.W. Q. N.A.	B.f. V,4	M.fr. VI, 149.
M. montanum, R. Brown, prodr. 515 (1810) ...	W.A.	S.A.	—	V. N.S.W. Q. N.A.	B.f. V,4	M.fr. VI, 149.	
M. deserti, Cunningham in Hueg. enum. 78 (1837) ...	W.A.	S.A.	—	V. N.S.W. Q. —	B.f. V,5	M.fr. VI, 148; VII, 110.	
M. laxiflorum, Benth. Fl. Austr. V, 6 (1870) ...	—	—	—	—	Q. —	B.f. V,6	M.fr. V, 23.
M. insulare, R. Brown, prodr. 515 (1810) ...	W.A.	S.A.	T. V.	N.S.W. —	—	B.f. V,6	M.fr. VI, 149; VII, 110.
M. viscosum, R. Brown, prodr. 516 (1810) ...	—	S.A.	—	V. N.S.W. —	—	B.f. V,5	M.fr. VII, 109.
M. serratum, R. Brown, prodr. 516 (1810) ...	W.A.	—	—	—	—	B.f. V,4	M.fr. VI, 149.
M. oppositifolium, R. Brown, prodr. 516 (1810) ...	W.A.	—	—	—	—	B.f. V,7	M.fr. VI, 149.
M. brevipes, Benth. Fl. Austr. V, 6 (1870) ...	—	S.A.	—	—	—	B.f. V,6	—
M. humile, R. Brown, prodr. 516 (1810) ...	W.A.	S.A.	T. V.	N.S.W. —	—	B.f. V,6	M.fr. VI, 150.
M. debile, R. Brown, prodr. 516 (1810) ...	—	—	—	N.S.W. Q. —	—	B.f. V,8	M.fr. VI, 149.
M. platycarpum, R. Brown, prodr. 516 (1810) ...	W.A.	S.A.	—	V. N.S.W. —	—	B.f. V,7	M.fr. VI, 150.
M. Bateae, F. v. M. in Proceed. Linn. Soc. N.S.W. VI, 792 (1851) ...	—	—	—	N.S.W. —	—	—	M.fr. XI, 139.
M. floribundum, Cunningham in Hueg. enum. 78 (1837) ...	—	—	—	V. N.S.W. —	—	B.f. V,8	M.fr. I, 126; VI, 150.
M. Beckeri, F. v. M. in Benth. Fl. Austr. V, 7 (1870) ...	W.A.	—	—	—	—	B.f. V,7	M.fr. I, 126, 244; VI, 150.
M. salsoloides, Turcz. in Bull. de Mosc. XXXVI, 226 (1863) ...	W.A.	—	—	—	—	B.f. V,8	M.fr. I, 126; VI, 150.
EREMOPHILA, R. Brown, prodr. 518 (1810). (Pholidia, Stenochilus, Eremodendron, Pseudopholidia, Sentis, Duttonia, Pholidiopsis.)							
E. Mackinlayi, F. v. M., fragm. IV, 80 (1864) ...	W.A.	—	—	—	—	B.f. V,17	M.fr. IV, 80.
E. Bowmani, F. v. M., fragm. II, 139 (1863) ...	—	S.A.	—	N.S.W. Q. —	—	B.f. V,18	M.fr. II, 139.
E. stronglyphylla, F. v. M., fragm. X, 87 (1876) ...	W.A.	S.A.	—	—	—	—	M.fr. X, 87.
E. leucophylla, Benth. Fl. Austr. V, 18 (1870) ...	W.A.	—	—	—	—	B.f. V,18	—
E. Turtoni, F. v. M., fragm. X, 87 (1876) ...	W.A.	S.A.	—	—	—	—	M.fr. X, 87.
E. Forrestii, F. v. M., fragm. VII, 49 (1869) ...	W.A.	—	—	—	—	B.f. V,18	M.fr. VII, 49.
E. eriocalyx, F. v. M., fragm. I, 236 (1859) ...	W.A.	—	—	—	—	B.f. V,19	M.fr. I, 236.
E. Maitlandi, F. v. M. in Benth. Fl. Austr. V, 19 (1870) ...	W.A.	—	—	—	—	B.f. V,19	—
E. rotundifolia, F. v. M., fragm. I, 207 (1859) ...	—	S.A.	—	—	—	B.f. V,19	M.fr. I, 207.
E. oppositifolia, R. Brown, prodr. 518 (1810) ...	—	S.A.	—	V. N.S.W. —	—	B.f. V,20	—
E. Mitchelli, Benth. in Mitch. Trop. Austr. 31 (1848) ...	—	—	—	N.S.W. Q. —	—	B.f. V,21	M.fr. VI, 148.
E. Paisleyi, F. v. M., Rep. Babb. Exped. 17 (1858) ...	—	S.A.	—	—	—	B.f. V,20	—
E. Sturtii, R. Brown, App. to Sturt's Exped. 22 (1849) ...	—	S.A.	—	N.S.W. —	—	B.f. V,21	M.fr. VI, 148.
E. exilifolia, F. v. M., fragm. X, 88 (1876) ...	—	S.A.	—	—	—	—	M.fr. X, 88.
E. Dempsteri, F. v. M., fragm. X, 60 (1876) ...	W.A.	—	—	—	—	—	M.fr. X, 60.
E. Gibsoni, F. v. M., fragm. VIII, 227 (1874) ...	—	S.A.	—	—	—	—	M.fr. VIII, 227.
E. Berryi, F. v. M., fragm. VIII, 228 (1874) ...	W.A.	S.A.	—	—	—	—	M.fr. VIII, 228; X, 88.
E. Clarkei, F. v. M., fragm. I, 208 (1859) ...	W.A.	—	—	—	—	B.f. V,21	M.fr. I, 208.
E. Gilesii, F. v. M., fragm. VIII, 49 (1873) ...	—	S.A.	—	—	—	—	M.fr. VIII, 49.
E. Latrobei, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858) ...	W.A.	S.A.	—	N.S.W. Q. N.A.	B.f. V,22	M.fr. I, 125.	
E. graciliflora, F. v. M., fragm. I, 208 (1858) ...	W.A.	—	—	—	—	B.f. V,23	M.fr. I, 208.
E. longifolia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858) ...	W.A.	S.A.	—	V. N.S.W. Q. N.A.	B.f. V,23	M.fr. VI, 148.	
E. Drummondii, F. v. M., fragm. VI, 147 (1868) ...	W.A.	—	—	—	—	B.f. V,24	M.fr. VI, 147.
E. Hughesii, F. v. M., fragm. VIII, 228 (1874) ...	W.A.	S.A.	—	—	—	—	M.fr. VIII, 228.
E. polyclada, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858) ...	—	S.A.	—	V. N.S.W. Q. —	B.f. V,24	—	—
E. bignoniiflora, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858) ...	—	S.A.	—	V. N.S.W. Q. N.A.	B.f. V,25	M.fr. VI, 148.	
E. Freelingii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858) ...	—	S.A.	—	N.S.W. —	B.f. V,25	M.fr. VI, 148.	
E. Fraseri, F. v. M., fragm. XI, 51 (1878) ...	W.A.	—	—	—	N.A. —	—	M.fr. XI, 51.
E. Macdonnelli, F. v. M., Rep. Babb. Exped. 18 (1858) ...	—	S.A.	—	—	—	B.f. V,22	M.fr. VI, 150.
E. Goodwinii, F. v. M., Rep. Babb. Exped. 17 (1858) ...	—	S.A.	—	N.S.W. —	B.f. V,25	M.fr. VI, 148.	
E. Elderi, F. v. M., fragm. VIII, 228 (1874) ...	W.A.	S.A.	—	—	—	—	M.fr. VIII, 228.
E. Willsii, F. v. M., fragm. III, 21, t. 20 (1862) ...	W.A.	S.A.	—	—	—	B.f. V,26	M.fr. III, 21.
E. platycalyx, F. v. M., fragm. V, 109 (1866) ...	W.A.	—	—	—	—	B.f. V,26	M.fr. V, 109.
E. viscida, Endlicher, nov. stirp. dec. 51 (1839) ...	W.A.	—	—	—	—	B.f. V,26	—
E. Brownii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858) ...	W.A.	S.A.	—	V. N.S.W. Q. —	B.f. V,27	M.fr. VI, 148.	
E. subfloccosa, Benth. Fl. Austr. V, 28 (1870) ...	W.A.	—	—	—	—	B.f. V,28	—
E. Oldfieldii, F. v. M., fragm. I, 208 (1859) ...	W.A.	—	—	—	—	B.f. V,28	M.fr. I, 208.
E. Duttonii, F. v. M., Rep. Babb. Exped. 16 (1858) ...	—	S.A.	—	N.S.W. —	B.f. V,28	M.fr. VI, 148.	
E. maculata, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858) ...	W.A.	S.A.	—	V. N.S.W. Q. N.A.	B.f. V,29	—	—
E. denticulata, F. v. M., fragm. I, 125 (1859) ...	W.A.	—	—	—	—	B.f. V,29	M.fr. IV, 170.
E. latifolia, F. v. M. in Schlechtend. Linnaea XXV, 428 (1852) ...	W.A.	S.A.	—	N.S.W. —	B.f. V,30	M.fr. VI, 148.	
E. alternifolia, R. Brown, prodr. 518 (1810) ...	—	S.A.	—	V. N.S.W. —	B.f. V,30	M.fr. VI, 148.	
E. Dalyana, F. v. M., fragm. V, 22 (1865) ...	—	S.A.	—	N.S.W. —	B.f. V,10	M.fr. V, 22.	
E. scoparia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858) ...	—	S.A.	—	V. N.S.W. —	B.f. V,10	M.fr. V, 22; VI, 148.	
E. Delisserii, F. v. M., fragm. V, 108, t. 42 (1865) ...	W.A.	S.A.	—	—	—	B.f. V,11	M.fr. V, 189.
E. crassifolia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858) ...	—	S.A.	—	—	—	B.f. V,11	M.fr. VI, 148.
E. resinosa, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858) ...	W.A.	—	—	—	—	B.f. V,11	—
E. Behrii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858) ...	—	S.A.	—	—	—	B.f. V,12	—
E. Woollsiana, F. v. M., fragm. I, 125, t. 7 (1859) ...	W.A.	—	—	—	—	B.f. V,12	M.fr. VI, 151.
E. Christophori, F. v. M., fragm. IX, 120 (1875) ...	—	S.A.	—	—	—	—	M.fr. IX, 198.
E. brevifolia, F. v. M.; Myoporum, Bartling in pl. Preiss. I, 350 (1845) ...	W.A.	—	—	—	—	B.f. V,12	M.fr. VI, 150.
E. Weldii, F. v. M., fragm. VII, 109 (1870) ...	W.A.	S.A.	—	—	—	—	M.fr. IX, 120.
E. imbricata, F. v. M.; Pholidia, Benth. Fl. Austr. V, 13 (1870) ...	W.A.	—	—	—	—	B.f. V,13	—
E. densifolia, F. v. M., fragm. II, 160 (1861) ...	W.A.	—	—	—	—	B.f. V,13	M.fr. II, 160.
E. gibbosifolia, F. v. M., Rep. Babb. Exped. 18 (1858) ...	—	S.A.	—	V. —	—	B.f. V,13	M.fr. VI, 150.
E. divaricata, F. v. M. in Proceed. Roy. Soc. Tasm. III, 293 (1858) ...	—	S.A.	—	V. N.S.W. —	—	B.f. V,14	M.fr. IV, 48; VI, 150.
E. microtheca, F. v. M. in Benth. Fl. Austr. V, 14 (1870) ...	W.A.	—	—	—	—	B.f. V,14	M.fr. II, 160.
E. adenotricha, F. v. M. in Benth. Fl. Austr. V, 15 (1870) ...	W.A.	—	—	—	—	B.f. V,14	—
E. santalina, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858) ...	—	S.A.	—	—	—	B.f. V,15	—
E. Youngii, F. v. M., fragm. X, 16 (1876) ...	W.A.	—	—	—	—	—	M.fr. X, 16.

PEDALINAE.

R. Brown, prodr. Fl. Nov. Holl. 519 (1810).

JOSEPHINIA, Ventenat, Jardin de la Malmaison, 103 t. 67 (1804).

J. grandiflora, R. Brown, prodr. 520 (1810)	...	...	...	...	Q.	—	B.f.IV, 556	
J. Imperatricis, Ventenat, Jard. de la Malm. t. 67 (1804)	...	...	...	...	—	N.A.	B.f.IV, 557	M.fr. VI, 163.
J. Eugeniae, F. v. M. in Hook. Kew Misc. IX, 370, t. 11 (1857)	W.A.	S.A.	—	—	Q.	N.A.	B.f.IV, 557	M.fr. VI, 163.

ERICACEAE.

A. L. de Jussieu, gen. plant. 159 (1789).

PERNETTYA, Gaudichaud in Ann. des sc. nat. V, 102 (1825).

P. Tasmanica, J. Hooker in Lond. Journ. of Bot. VI, 268 (1847)	—	—	T.	—	—	—	B.f.IV, 140	
--	---	---	----	---	---	---	-------------	--

GAULTIERA, Kalm in Linné, amoen. acad. III, 14, fig. 6 (1751). (Gaultheria.)

G. hispida, R. Brown, prodr. 559 (1810)	...	...	...	...	T.	V.	N.S.W.	—	B.f.IV, 141
G. lanceolata, J. Hooker in Lond. Journ. of Bot. VI, 267 (1847)	—	—	T.	—	—	—	—	—	B.f.IV, 141
G. antipoda, G. Forster, florul. ins. Austr. pr. 34 (1786)...	...	...	—	—	T.	—	—	—	B.f.IV, 142

WITTSTEINIA, F. v. M., fragm. II, 136 (1861).

W. vacciniacea, F. v. M., fragm. II, 136 (1861)	...	...	—	—	—	V.	—	—	B.f.IV, 139	M.fr. III, 166.
---	-----	-----	---	---	---	----	---	---	-------------	-----------------

EPACRIDAE.

R. Brown, prodr. 535 (1810).

STYPHELIA, Solander in G. Forster, fl. ins. Austr. prodr. 13 (1786). (Epacris, Forst. (1776) partly, Stiphelia, Ardisia, Peroa, Perojoa, Ventenatia, Cyathodes, Stenanthra, Astroloma, Leucopogon, Melichrus, Acrotriche, Monotoca, Soleniscia, Stomarrhena, Pentataphrus, Mesotriche, Phanerandra, Froebelia, Pentaptelion, Androstoma, Cyathopsis, Lissanthe partly.)

S. adscendens, R. Brown, prodr. 537 (1810)	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	--

<i>S. gibbosa</i> , F. v. M.; Leucopogon, Stschegleew in Bull. de Mosc. XXXII, 12 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 189	
<i>S. thymifolia</i> , F. v. M.; Leucopogon, Lindley in Benth. Fl. Austr. IV, 189 (1869) ...	—	—	—	V.	—	—	—	B.f.IV, 189	
<i>S. cordata</i> , F. v. M.; Leucop., Sonder in pl. Preiss. I, 313 (1845) ...	W.A.	—	—	—	—	—	—	B.f.IV, 190	
<i>S. Bossiaca</i> , F. v. M., fragm. VI, 47 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 190	M.fr. VI, 47.
<i>S. hirsuta</i> , F. v. M., fragm. VI, 31 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 191	M.fr. VI, 31.
<i>S. collina</i> , Labillardière, Nov. Holl. pl. spec. I, 47, t. 65 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 191	M.fr. VI, 45; VIII, 54.
<i>S. glacialis</i> , F. v. M.; Leucopogon, Lindl. in Mitch. three Exped. II, 127 (1838) ...	—	—	—	V.	—	—	—	B.f.IV, 191	
<i>S. compacta</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 13 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 192	
<i>S. squarrosa</i> , F. v. M., fragm. VI, 31 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 192	M.fr. VI, 31.
<i>S. microphylla</i> , Sprengel, syst. I, 656 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 192	M.fr. VI, 31.
<i>S. tetragona</i> , F. v. M., fragm. VI, 31 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 193	M.fr. VI, 31.
<i>S. phyllostachys</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 193 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 193	M.fr. VI, 32.
<i>S. glabella</i> , Sprengel, syst. I, 655 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 194	M.fr. VI, 32.
<i>S. semiopposita</i> , F. v. M., fragm. VI, 49 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 194	M.fr. VI, 49.
<i>S. florulenta</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 194 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 194	
<i>S. striata</i> , Sprengel, syst. I, 656 (1825) ...	W.A.	S.A.	—	—	—	—	—	B.f.IV, 194	
<i>S. lasiostachya</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 11 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 195	
<i>S. carinata</i> , Sprengel, syst. I, 658 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 195	
<i>S. opponens</i> , F. v. M., fragm. VI, 48 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 196	M.fr. VI, 48.
<i>S. oppositifolia</i> , F. v. M., fragm. VI, 32 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 196	M.fr. VI, 32.
<i>S. tamariscina</i> , Sprengel, syst. I, 656 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 197	
<i>S. bracteolaris</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 197 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 197	
<i>S. blepharophylla</i> , F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 197	M.fr. VI, 34.
<i>S. tenuis</i> , F. v. M.; Leucop., De Candolle prodr. VII, 744 (1833) ...	W.A.	—	—	—	—	—	—	B.f.IV, 197	
<i>S. gnaphalioides</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 14 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 198	
<i>S. concurva</i> , F. v. M., fragm. VI, 36 (1867) ...	—	S.A.	—	—	—	—	—	B.f.IV, 198	M.fr. III, 144.
<i>S. Gilbertii</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXII, 15 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 198	
<i>S. gracilis</i> , Sprengel, syst. I, 658 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 199	
<i>S. acicularis</i> , F. v. M.; Leucop., Benth., fl. Austr. IV, 199 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 199	
<i>S. cryptantha</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 199 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 199	
<i>S. gracillima</i> , F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 200	M.fr. VI, 34.
<i>S. cymbiformis</i> , F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 200	M.fr. VI, 34.
<i>S. apiculata</i> , Sprengel, syst. I, 656 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 201	M.fr. IV, 105; VI, 31.
<i>S. polystachya</i> , Sprengel, syst. I, 659 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 201	M.fr. VI, 31.
<i>S. virgata</i> , Labillardière, Nov. Holl. pl. spec. I, 46, t. 64 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.IV, 201	M.fr. VI, 42.
<i>S. pulchella</i> , F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 202	M.fr. VI, 34.
<i>S. polymorpha</i> , F. v. M., fragm. VI, 31 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 202	M.fr. VI, 31.
<i>S. assimilis</i> , F. v. M.; Leucopogon, R. Brown, prodr. 545 (1810) ...	W.A.	—	—	—	—	—	—	B.f.IV, 202	M.fr. IV, 106; VI, 32.
<i>S. Oldfieldii</i> , F. v. M.; Leucopogon, Benth., fl. Austr. IV, 203 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 203	
<i>S. cucullata</i> , Sprengel, syst. I, 656 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 203	M.fr. VI, 32.
<i>S. sprengelioides</i> , F. v. M.; Leuc., Sonder in pl. Preiss. I, 319 (1845) ...	W.A.	—	—	—	—	—	—	B.f.IV, 204	M.fr. VI, 34.
<i>S. obtusata</i> , F. v. M.; Leuc., Sonder in pl. Preiss. I, 313 (1845) ...	W.A.	—	—	—	—	—	—	B.f.IV, 204	
<i>S. fimbriata</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 17 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 204	
<i>S. ozothamnoides</i> , Leuc., F. v. M. in Benth. fl. Austr. IV, 205 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 204	M.fr. VI, 155.
<i>S. plumuliflora</i> , F. v. M., fragm. VI, 29 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 205	M.fr. XI, 122.
<i>S. unilateralis</i> , F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 19 (1859) ...	W.A.	—	—	—	—	—	—	B.f.IV, 205	
<i>S. montana</i> , F. v. M., fragm. VI, 45 (1867) ...	—	T.	V.	N.S.W.	—	—	—	B.f.IV, 206	M.fr. VI, 45.
<i>S. Macraei</i> , F. v. M., fragm. VI, 46 (1867) ...	—	—	—	V.	N.S.W.	—	—	B.f.IV, 206	M.fr. VI, 46; VIII, 54.
<i>S. pleurandroides</i> , F. v. M., fragm. VI, 32 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 206	M.fr. III, 143.
<i>S. linifolia</i> , F. v. M., fragm. VI, 36 (1867) ...	—	—	—	—	N.S.W. Q.	—	—	B.f.IV, 207	M.fr. VI, 36.
<i>S. pluriloculata</i> , F. v. M., fragm. VI, 32 (1867) ...	—	—	—	—	N.S.W. Q.	—	—	B.f.IV, 207	M.fr. I, 37.
<i>S. pleiosperma</i> , F. v. M., fragm. VI, 41 (1867) ...	—	—	—	—	N.S.W. Q.	—	—	B.f.IV, 207	M.fr. VI, 41.
<i>S. rubicunda</i> , F. v. M., fragm. VI, 31 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 208	M.fr. IV, 99.
<i>S. attenuata</i> , F. v. M.; Leucopogon, Cunningham in Field's N.S.W. 341 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 208	
<i>S. conferta</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 208 (1869) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 208	
<i>S. mutica</i> , F. v. M., fragm. VI, 45 (1867) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 209	M.fr. VI, 45.
<i>S. ericoides</i> , Smith, Specim. of the Bot. of New Holl. 48 (1793) ...	—	S.A.	T.	V.	N.S.W. Q.	—	—	B.f.IV, 209	M.fr. VI, 45.
<i>S. brevicuspis</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 210 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 210	
<i>S. propinqua</i> , Sprengel, syst. I, 658 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 210	M.fr. VI, 34.
<i>S. subulifolia</i> , F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 211	M.fr. IV, 103.
<i>S. Allittii</i> , F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 211	M.fr. IV, 103.
<i>S. racemulosa</i> , F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 211	M.fr. VI, 33.
<i>S. pendula</i> , Sprengel, syst. I, 657 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 212	M.fr. VI, 33.
<i>S. concinna</i> , F. v. M.; Leuc., Benth., fl. Austr. IV, 212 (1869) ...	W.A.	—	—	—	—	—	—	B.f.IV, 212	
<i>S. margarodes</i> , Sprengel, syst. I, 657 (1825) ...	—	—	—	—	N.S.W. Q.	—	—	B.f.IV, 213	M.fr. VI, 36.
<i>S. flavescens</i> , F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 213	M.fr. IV, 100.
<i>S. blepharolepis</i> , F. v. M., fragm. VI, 48 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 213	M.fr. VI, 48.
<i>S. esquamata</i> , Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	—	—	B.f.IV, 214	
<i>S. rotundifolia</i> , Sprengel, syst. I, 655 (1825) ...	W.A.	—	—	—	—	—	—	B.f.IV, 214	
<i>S. cordifolia</i> , F. v. M., fragm. VIII, 54 (1873) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f.IV, 214	M.fr. VI, 45.
<i>S. megacarpa</i> , F. v. M., fragm. VI, 32 (1867) ...	W.A.	—	—	—	—	—	—	B.f.IV, 215	M.fr. IV, 102.

S. ruscifolia, Sprengel, syst. I, 656 (1825) ...	—	—	—	—	Q.	—	B.f. IV, 215	
S. imbricata, Sprengel, syst. I, 656 (1825) ...	—	—	—	—	Q.	—	B.f. IV, 215	
S. cuspidata, Sprengel, syst. I, 657 (1825) ...	—	—	—	—	Q.	—	B.f. IV, 216	
S. leptospermoides, Sprengel, syst. I, 659 (1825) ...	—	—	—	—	Q.	—	B.f. IV, 216	
S. acuminata, Sprengel, syst. I, 659 (1825) ...	—	—	—	—	N.A.	—	B.f. IV, 216	
S. flexifolia, Sprengel, syst. I, 659 (1825) ...	—	—	—	—	Q.	—	B.f. IV, 217	
S. biflora, Sprengel, syst. I, 659 (1825) ...	—	—	—	V.	N.S.W.	Q.	B.f. IV, 217	M.fr. IV, 104.
S. setigera, Sprengel, syst. I, 659 (1825) ...	—	—	—	—	N.S.W.	—	B.f. IV, 217	M.fr. VI, 45.
S. exolasia, F. v. M., fragm. VI, 34 (1867) ...	—	—	—	—	N.S.W.	—	B.f. IV, 217	M.fr. VI, 34.
S. Fraseri, F. v. M.; Leucop., Cunn. in Ann. nat. hist. II, 47 (1840) ...	—	—	—	T. V.	N.S.W.	—	B.f. IV, 218	M.fr. IV, 105; VI, 46.
S. hirtella, F. v. M.; Leucopogon in Benth. Fl. Austr. IV, 218 (1869) ...	S.A.	—	—	—	—	—	B.f. IV, 218	
S. Oldfieldii, F. v. M.; Leuc. ovalifolius, Sond. in pl. Preiss. I, 324 (1845) W.A.	—	—	—	—	—	—	B.f. IV, 218	
S. erubescens, F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 219	M.fr. IV, 102.
S. lissanthoides, F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 219	M.fr. IV, 101.
S. stricta, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 219 (1869) W.A.	—	—	—	—	—	—	B.f. IV, 219	
S. Mitchellii, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 220 (1869) ...	—	—	—	—	Q.	—	B.f. IV, 220	
S. juniperina, Sprengel, syst. I, 658 (1825) ...	—	—	—	V.	N.S.W.	Q.	B.f. IV, 220	M.fr. IV, 104; VI, 46.
S. rufa, F. v. M., fragm. VI, 46 (1867) ...	S.A.	—	V.	N.S.W.	—	—	B.f. IV, 221	M.fr. VIII, 54; IX, 48.
S. conostephioides, F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 221	M.fr. VI, 34.
S. deformis, Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	—	B.f. IV, 221	
S. pogonocalyx; Leucop., F. v. M. in Benth. Fl. Austr. IV, 222 (1869) W.A.	—	—	—	—	—	—	B.f. IV, 222	
S. breviflora; Leucopogon, F. v. M., fragm. IV, 102 (1864) ...	W.A.	—	—	—	—	—	B.f. IV, 222	M.fr. IV, 102.
S. dura, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 222 (1869) W.A.	—	—	—	—	—	—	B.f. IV, 222	
S. multiflora, Sprengel, syst. I, 658 (1825) ...	W.A.	—	—	—	—	—	B.f. IV, 222	M.fr. VI, 32.
S. appressa, Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	—	B.f. IV, 223	
S. Neo-Anglica; Leucop., F. v. M. in Benth. Fl. Austr. IV, 222 (1869) ...	—	—	—	—	N.S.W.	Q.	B.f. IV, 223	
S. obtecta, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 223 (1869) W.A.	—	—	—	—	—	—	B.f. IV, 223	
S. crassiflora, F. v. M., fragm. VI, 40 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 224	M.fr. VI, 40.
S. stronglyphylla, F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 224	M.fr. IV, 101.
S. crassifolia, F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 224	M.fr. VI, 33.
S. corynocarpa, F. v. M.; Leucop., Sonder in pl. Preiss. I, 322 (1845) W.A.	—	—	—	—	—	—	B.f. IV, 224	
S. Woodsii; Leucopogon, F. v. M., fragm. I, 178 (1859) ...	W.A.	S.A.	—	V.	—	—	B.f. IV, 225	M.fr. VI, 33.
S. leptantha, F. v. M.; Leucop., Benth. Fl. Austr. IV, 225 (1869) W.A.	—	—	—	—	—	—	B.f. IV, 225	
S. divaricata, Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	Q.	B.f. IV, 226	M.fr. VI, 44.
S. aggregata, Sprengel, syst. I, 657 (1825) ...	—	—	—	—	N.S.W.	Q.	B.f. IV, 227	M.fr. VI, 44; VIII, 55.
S. serrulata, Labillardiere, Nov. Holl. pl. spec. I, 45, t. 62 (1804) ...	S.A.	T.	V.	N.S.W.	—	—	B.f. IV, 227	M.fr. VI, 44.
S. patula, Sprengel, syst. I, 657 (1825) ...	—	S.A.	—	—	—	—	B.f. IV, 227	M.fr. VI, 44; XI, 122.
S. ovalifolia, Sprengel, syst. I, 656 (1825) ...	W.A.	S.A.	—	V.	—	—	B.f. IV, 228	M.fr. VI, 44.
S. ramiflora, Sprengel, syst. I, 659 (1825) ...	W.A.	—	—	—	—	—	B.f. IV, 228	M.fr.
S. depressa, Sprengel, syst. I, 655 (1825) ...	W.A.	S.A.	—	—	—	—	B.f. IV, 228	M.fr. VI, 44.
S. fasciculiflora, F. v. M., fragm. VIII, 55 (1873) ...	—	S.A.	—	—	—	—	B.f. IV, 229	M.fr. VI, 44.
S. elliptica, Smith, Specim. of the Bot. of New Holl. 49 (1793) ...	—	—	T.	V.	N.S.W.	Q.	B.f. IV, 230	M.fr. VI, 58.
S. glauca, Labillardiere, Nov. Holl. pl. spec. I, 45, t. 61 (1804) ...	—	—	T.	—	—	—	B.f. IV, 230	
S. scoparia, Smith, Specim. of the Bot. of New Holl. 49 (1793) ...	—	—	T.	V.	N.S.W.	Q.	B.f. IV, 231	M.fr. VI, 58; XI, 122.
S. ledifolia, F. v. M.; Monotoca, Cunningham in De Cand., prodr. VII, 756 (1839) ...	—	—	—	—	N.S.W.	—	B.f. IV, 231	
S. empetrifolia, F. v. M. in Papers of R. S. Tasm. 86 (1874) ...	—	—	T.	—	—	—	B.f. IV, 231	
S. oligarrhenoides, F. v. M., fragm. IX, 47 (1875) ...	W.A.	—	—	—	—	—	—	M.fr. IX, 47.
S. minutiflora, F. v. M., fragm. XI, 122 (1881) ...	W.A.	—	—	—	—	—	B.f. IV, 231	M.fr. XI, 122.
OLIGARRHENA, R. Brown, prodr. Fl. Nov. Holl. 549 (1810).								
O. micrantha, R. Brown, prodr. 459 (1810) ...	W.A.	—	—	—	—	—	B.f. IV, 232	
NEEDHAMIA, R. Brown, prodr. Fl. Nov. Holl. 549 (1810).								
N. pumilio, R. Brown, prodr. 549 (1810) ...	W.A.	—	—	—	—	—	B.f. IV, 174	
BRACHYLOMA, Sonder in Lehm. pl. Preiss. I, 304 (1845). (Lobopogon, Lissanthe partly.)								
B. daphnoides, Benth. Fl. Austr. IV, 173 (1869) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f. IV, 173	M.fr. VI, 42; VIII, 55.
B. ciliatum, Benth. Fl. Austr. IV, 173 (1869) ...	—	S.A.	T.	V.	—	—	B.f. IV, 173	M.fr. VI, 42.
B. depressum, Benth. Fl. Austr. IV, 173 (1869) ...	—	—	T.	V.	—	—	B.f. IV, 173	M.fr. I, 36; VI, 42.
B. Scortechinii, F. v. M., fragm. XI, 121 (1881) ...	—	—	—	—	—	Q.	—	M.fr. XI, 121.
B. Preissii, Sonder in Lehm. pl. Preiss. I, 305 (1845) ...	W.A.	—	—	—	—	—	B.f. IV, 172	M.fr. VI, 39; VIII, 55.
B. concolor, F. v. M., fragm. VI, 39 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 172	M.fr. IV, 98.
B. ericoides, Sonder in Schlecht. Linnaea XXVI, 247 (1853) ...	—	S.A.	—	V.	N.S.W.	—	B.f. IV, 172	M.fr. IV, 98; VI, 39; XI, 46.
CONOSTEPHIUM, Benth. in Hueg. enum. 76 (1837). (Conostephiopsis.)								
C. pendulum, Benth. in Hueg. enum. 76 (1837) ...	W.A.	—	—	—	—	—	B.f. IV, 160	M.fr. VI, 40.
C. minus, Lindley, Bot. Regist. XXV, App. XXV (1839) ...	W.A.	—	—	—	—	—	B.f. IV, 160	M.fr. VI, 40.
C. Roei, Benth. Fl. Austr. IV, 160 (1869) ...	W.A.	—	—	—	—	—	B.f. IV, 160	M.fr. VIII, 55.
C. Preissii, Sonder in Lehm. pl. Preiss. I, 304 (1845) ...	W.A.	—	—	—	—	—	B.f. IV, 161	M.fr. VI, 40.
C. planifolium, F. v. M.; fragm. VI, 30 (1867) ...	W.A.	—	—	—	—	—	B.f. IV, 161	M.fr. VI, 30.
COLEANTHERA, Stschegleew in Bull. Soc. Mosc. XXXII, 4 (1859). (Michiea.)								
C. coelophylla, Benth. Fl. Austr. IV, 150 (1869) ...	W.A.	—	—	—	—	—	B.f. IV, 150	
C. myrtoides, Stschegleew in Bull. de Mosc. XXXII, 4 (1859) ...	W.A.	—	—	—	—	—	B.f. IV, 150	M.fr. VI, 80.
C. virgata, Stschegleew in Bull. de Mosc. XXXII, 5 (1859) ...	W.A.	—	—	—	—	—	B.f. IV, 151	
TROCHOCARPA, R. Brown, prodr. 548 (1810). (Decaspora, Pentachondra.)								
T. laurina, R. Brown, prodr. 548 (1810) ...	—	—	—	—	N.S.W.	Q.	B.f. IV, 166	M.fr. VI, 57; IX, 48.
T. disticha, Sprengel, syst. I, 660 (1825) ...	—	—	T.	—	—	—	B.f. IV, 166	M.fr. VI, 57; VIII, 55.
T. thymifolia, Sprengel, syst. I, 660 (1825) ...	—	—	T.	—	—	—	B.f. IV, 166	M.fr. VI, 57; VIII, 55.
T. Clarkei, F. v. M., fragm. VI, 57 (1867) ...	—	—	—	V.	—	—	B.f. IV, 167	M.fr. VI, 57.
T. Gunnii, Benth. Fl. Austr. IV, 167 (1869) ...	—	—	T.	—	—	—	B.f. IV, 167	M.fr. VIII, 55.
T. parviflora, Benth. Fl. Austr. IV, 167 (1869) ...	W.A.	—	—	—	—	—	B.f. IV, 167	

T. involucreta, F. v. M., fragm. VI, 57 (1867)	—	—	—	—	T. — — —	B.fl. IV, 164	M.fr. VIII, 55.
T. pumila, F. v. M., fragm. VI, 57 (1867)	—	—	—	—	T. V. N.S.W. —	B.fl. IV, 164	M.fr. VI, 57.
T. ericifolia, F. v. M. in Papers R. S. Tasm. 86 (1874)	—	—	—	—	T. — — —	B.fl. IV, 164	
T. verticillata, F. v. M. in Papers R. S. Tasm. 86 (1874)	—	—	—	—	T. — — —	B.fl. IV, 164	
EPACRIS, Cavanilles, icon. et descr. pl. IV, 25, t. 344 et 345 (1797). (Archeria.)							
E. micranthera, F. v. M., fragm. VI, 72 (1867)	—	—	—	—	T. — — —	B.fl. VI, 246	M.fr. VIII, 56.
E. hirtella, J. Hooker in Lond. Journ. of Bot. VI, 271 (1847)	—	—	—	—	T. — — —	B.fl. VI, 246	M.fr. VI, 71.
E. longiflora, Cavanilles, icon. et descr. pl. IV, 25, t. 344 (1797)	—	—	—	—	N.S.W. — —	B.fl. IV, 234	M.fr. VI, 70; VIII, 56.
E. reclinata, Cunningham in Benth. Fl. Austr. IV, 234 (1869)	—	—	—	—	N.S.W. — —	B.fl. IV, 234	M.fr. VIII, 56. [122]
E. impressa, Labillardière, Nov. Holl. pl. spec. I, 43, t. 58 (1804)	—	S.A.	T. V.	N.S.W.	— —	B.fl. IV, 235	M.fr. VI, 70; VIII, 56; XI,
E. sparsa, R. Brown, prodr. 551 (1810)	—	—	—	—	N.S.W. — —	B.fl. IV, 235	M.fr. VIII, 53.
E. petrophila, J. Hooker, Fl. Tasman. I, 261 (1860)	—	—	T. V.	N.S.W.	— —	B.fl. IV, 236	
E. rigida, Sieber in Spreng. cur. poster. 64 (1827)	—	—	—	—	N.S.W. — —	B.fl. IV, 236	
E. coriacea, Cunningham in Cand. prodr. VII, 763 (1839)	—	—	—	—	N.S.W. — —	B.fl. IV, 236	
E. crassifolia, R. Brown, prodr. 551 (1810)	—	—	—	—	N.S.W. — —	B.fl. IV, 237	
E. robusta, Benth. Fl. Austr. IV, 237 (1869)	—	—	—	—	V. N.S.W. — —	B.fl. IV, 237	
E. obtusifolia, Smith, Exot. Bot. I, 77, t. 40 (1804)	—	S.A.	T. V.	N.S.W.	Q. —	B.fl. IV, 237	M.fr. VI, 71; VIII, 56.
E. myrtifolia, Labillardière, Nov. Holl. pl. spec. I, 41, t. 55 (1804)	—	—	T. —	—	— —	B.fl. IV, 238	
E. exserta, R. Brown, prodr. 551 (1810)	—	—	T. —	—	— —	B.fl. IV, 238	
E. mucronulata, R. Brown, prodr. 552 (1810)	—	—	T. —	—	— —	B.fl. IV, 238	
E. lanuginosa, Labillardière, Nov. Holl. pl. spec. I, 42, t. 57 (1804)	—	S.A.	T. V.	—	— —	B.fl. IV, 238	M.fr. VIII, 56; XI, 122.
E. paludosa, R. Brown, prodr. 551 (1810)	—	—	—	—	V. N.S.W. — —	B.fl. IV, 239	M.fr. VIII, 56.
E. calvertiana, F. v. M., fragm. VIII, 53 (1874)	—	—	—	—	N.S.W. — —	—	M.fr. VIII, 53.
E. heteronema, Labillardière, Nov. Holl. pl. spec. I, 42, t. 56 (1804)	—	—	T. V.	N.S.W.	— —	B.fl. IV, 239	M.fr. VI, 71; VIII, 56.
E. serpillifolia, R. Brown, prodr. 240 (1810)	—	—	T. V.	N.S.W.	— —	B.fl. IV, 240	
E. microphylla, R. Brown, prodr. 550 (1810)	—	S.A.	T. V.	N.S.W.	Q. —	B.fl. IV, 240	
E. acuminata, Benth. Fl. Austr. IV, 240 (1869)	—	—	T. —	—	— —	B.fl. IV, 240	
E. apiculata, Cunningham in Field, N.S. Wales, 340 (1825)	—	—	—	—	N.S.W. — —	B.fl. IV, 241	
E. pulchella, Cavanilles, icon. IV, 26, t. 345 (1797)	—	—	—	—	N.S.W. Q. —	B.fl. IV, 241	M.fr. VI, 71; XI, 122.
E. purpurascens, R. Brown, prodr. 550 (1810)	—	—	—	—	N.S.W. — —	B.fl. IV, 241	M.fr. IV, 127; VIII, 56.
WOOLLSIA, F. v. M., fragm. VIII, 52 (1872). (Lysinema partly.)							
W. pungens, F. v. M., fragm. VIII, 52 (1872)	—	—	—	—	N.S.W. Q. —	B.fl. IV, 243	M.fr. IV, 126; VI, 70, 88; VIII, 55; IX, 48; XI, 122.
LYSINEMA, R. Brown, prodr. 552 (1810).							
L. lasianthum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	— —	B.fl. IV, 243	M.fr. III, 142; VI, 70; VIII, 55. [122]
L. conspicuum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	— —	B.fl. IV, 243	M.fr. VI, 70; VIII, 55; XI,
L. ciliatum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	— —	B.fl. IV, 243	M.fr. VI, 69; VII, 55.
L. fimbriatum, F. v. M., fragm. IV, 125 (1864)	W.A.	—	—	—	— —	B.fl. IV, 244	M.fr. VI, 70.
L. elegans, Sonder in Lehm. pl. Preiss. I, 327 (1845)	W.A.	—	—	—	— —	B.fl. IV, 244	M.fr. VI, 70.
PRIONOTES, R. Brown, prodr. fl. Nov. Holl. 553 (1810).							
P. cerinthoides, R. Brown, prodr. 553 (1810)	—	—	T. —	—	— —	B.fl. IV, 246	M.fr. VI, 69; XI, 122.
COSMELLA, R. Brown, prodr. fl. Nov. Holl. 553 (1810).							
C. rubra, R. Brown, prodr. 553 (1810)	W.A.	—	—	—	— —	B.fl. IV, 247	M.fr. VI, 64.
PONCELETIA, R. Brown, prodr. fl. Nov. Holl. 554 (1810).							
P. sprengelioides, R. Brown, prodr. 554 (1810)	—	—	—	—	N.S.W. Q. —	B.fl. IV, 248	M.fr. I, 39; VI, 60.
P. monticola, Cunningham in De Cand. prodr. VII, 763 (1843)	—	—	—	—	N.S.W. — —	B.fl. IV, 249	
SPRENGELIA, Smith in Svensk. Vetens. Acad. Handling. 260 (1794). (Poirotia.)							
S. incarnata, Smith in Vetens. Ac. Handl. 260, t. 8 (1794)	—	S.A.	T. V.	N.S.W.	— —	B.fl. IV, 249	M.fr. I, 39; VI, 54; IX, 48; [VIII, 56; XI, 122.]
ANDERSONIA, R. Brown, prodr. 553 (1810). (Atherocephala, Homalostoma, Sphincterostoma.)							
A. colosseae, F. v. M., fragm. VI, 63 (1867)	W.A.	—	—	—	— —	B.fl. IV, 251	M.fr. VI, 63.
A. patricia, F. v. M., fragm. VI, 79 (1868)	W.A.	—	—	—	— —	B.fl. IV, 251	M.fr. VI, 79.
A. grandiflora, Stschegleew in Bull. de Mosc. XXXII, 21 (1859)	W.A.	—	—	—	— —	B.fl. IV, 252	M.fr. VI, 62.
A. setifolia, Benth. Fl. Austr. IV, 252 (1869)	W.A.	—	—	—	— —	B.fl. IV, 252	M.fr. VIII, 51.
A. involucreta, Sonder in Lehm. pl. Preiss. I, 331 (1845)	W.A.	—	—	—	— —	B.fl. IV, 252	M.fr. VI, 62.
A. homalostoma, Benth. Fl. Austr. IV, 253 (1869)	W.A.	—	—	—	— —	B.fl. IV, 253	M.fr. VIII, 56.
A. sprengelioides, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	— —	B.fl. IV, 253	M.fr. VI, 64.
A. latiflora, F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	— —	B.fl. IV, 253	M.fr. VI, 61.
A. gracilis, De Candolle, prodr. VII, 767 (1843)	W.A.	—	—	—	— —	B.fl. IV, 254	
A. aristata, Lindley, Bot. Regist. XXV, App. XXV (1839)	W.A.	—	—	—	— —	B.fl. IV, 254	M.fr. VI, 63.
A. macronema, F. v. M., fragm. VIII, 51 (1873)	W.A.	—	—	—	— —	—	M.fr. VIII, 51.
A. parvifolia, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	— —	B.fl. IV, 254	M.fr. IV, 125; V, 62.
A. depressa, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	— —	B.fl. IV, 255	
A. coerulea, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	— —	B.fl. IV, 255	M.fr. VI, 64.
A. subulata, Benth. Fl. Austr. IV, 256 (1869)	W.A.	—	—	—	— —	B.fl. IV, 256	
A. heterophylla, Sonder in Lehm. pl. Preiss. I, 333 (1845)	W.A.	—	—	—	— —	B.fl. IV, 256	M.fr. IV, 124.
A. brachyanthera, F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	— —	B.fl. IV, 256	M.fr. VI, 61.
A. brevifolia, Sonder in Lehm. pl. Preiss. I, 332 (1845)	W.A.	—	—	—	— —	B.fl. IV, 256	M.fr. VI, 61.
A. variegata, Sonder in Lehm. pl. Preiss. I, 334 (1845)	W.A.	—	—	—	— —	B.fl. IV, 257	
A. micrantha, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	— —	B.fl. IV, 257	M.fr. VI, 61.
RICHEA, R. Brown, prodr. 555 (1810). (Cystenthe, Pilitis.)							
R. sprengelioides, F. v. M., fragm. VI, 63 (1867)	—	—	T. —	—	— —	B.fl. IV, 258	M.fr. I, 38; VI, 68.
R. procera, F. v. M., fragm. VI, 68 (1867)	—	—	T. —	—	— —	B.fl. IV, 259	M.fr. I, 38; VI, 68.
R. acerosa, F. v. M., fragm. VI, 69 (1867)	—	—	T. —	—	— —	B.fl. IV, 259	M.fr. I, 38; VI, 69.
R. Milligani, F. v. M., fragm. VI, 69 (1867)	—	—	T. —	—	— —	B.fl. IV, 259	M.fr. I, 38; VIII, 56.
R. Gunnii, J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T. V.	—	— —	B.fl. IV, 260	M.fr. VI, 67; XI, 122.
R. scoparia, J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T. —	—	— —	B.fl. IV, 260	M.fr. VI, 68.

<i>R. dracophylla</i> , R. Brown, prodr. 555 (1810)	—	—	T.	—	—	—	—	B.f.IV, 260	M.fr.VI, 67.
<i>R. pandanifolia</i> , J. Hooker, Fl. Antarct. I, 50 (1844) ..	—	—	T.	—	—	—	—	B.f.IV, 261	M.fr.VI, 67; IX, 48.
DRACOPHYLLUM, Labillardière, Voy. II, 211, t. 40 (1798). (Epacris, Forst. 1776 partly.)									
<i>D. Milligani</i> , Hooker, icon. t. 845 (1852)	—	—	T.	—	—	—	—	B.f.IV, 262	M.fr.VI, 65.
<i>D. Fitzgeraldi</i> , F. v. M., fragm. VII, 27 (1869)	—	—	—	—	—	N.S.W.	—	—	M.fr.X, 119.
<i>D. secundum</i> , R. Brown, prodr. 556 (1810)	—	—	—	—	—	N.S.W.	—	B.f.IV, 262	M.fr.VI, 65; VIII, 56.
<i>D. minimum</i> , F. v. M., fragm. I, 39 (1858)	—	—	T.	—	—	—	—	B.f.IV, 265	M.fr.VI, 65; VII, 56.
SPHENOTOMA, Sweet, Flora Australasica, t. 44 (1828).									
<i>S. squarrosus</i> , G. Don, gen. syst. III, 785 (1834)	W.A.	—	—	—	—	—	—	B.f.IV, 263	M.fr.VI, 66.
<i>S. Drummondii</i> , F. v. M.; Dracophyllum, Benth. Fl. Austr. IV, 263 (1869)	W.A.	—	—	—	—	—	—	B.f.IV, 263	—
<i>S. dracophylloides</i> , Sonder in Lehm. pl. Preiss. I, 33 (1844) ...	W.A.	—	—	—	—	—	—	B.f.IV, 264	M.fr.VI, 65.
<i>S. capitatum</i> , Lindley, Bot. Regist. t. 1515 (1832)	W.A.	—	—	—	—	—	—	B.f.IV, 264	—
<i>S. gracile</i> , Sweet, Fl. Austral. t. 44 (1828)	W.A.	—	—	—	—	—	—	B.f.IV, 264	M.fr.VI, 66.
<i>S. parviflorum</i> ; Dracophyllum, F. v. M. in Benth. Fl. Austr. IV, 265 (1869)	W.A.	—	—	—	—	—	—	B.f.IV, 265	M.fr.VI, 66.

APETALEAE GYMNOSPERMEAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 40 (1880).

CONIFERAE.

Haller, enum. stirp. Helv. I, 145 (1742).

DAMMARA, Rumphius, herb. Amboin. II, 174, t. 54 (1741). (Agathis.)									
<i>D. robusta</i> , C. Moore in Transact. pharm. Soc. Vict. II, 174 (1860) — — — — —	Q.	—	—	—	—	—	—	B.f.VI, 244	M.fr.XI, 104.
ARAUCARIA, A. L. de Jussieu, gen. plant. 413 (1789). (Colymbea, Columbea, Eutassa, Eutacta, Altingia.)									
<i>A. Cunninghamii</i> , Aiton in Sweet. hort. Brit. 475 (1827)	—	—	—	—	N.S.W.	Q.	—	B.f.VI, 243	—
<i>A. Bidwilli</i> , Hooker, Lond. Journ. of Bot. II, 503, t. 18 (1850) ...	—	—	—	—	—	Q.	—	B.f.VI, 243	—
<i>A. excelsa</i> , R. Brown in Ait. hort. Kew. sec. ed. V, 412 (1813) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX, 169.
ATHROTAXIS, D. Don in Transact. Linn. Soc. XVIII, 171 (1839). (Arthrotaxis, Sequoia.)									
<i>A. cupressoides</i> , D. Don in Trans. Linn. Soc. XVIII, 173, t. 13 (1839) — — — — —	T.	—	—	—	—	—	—	B.f.VI, 242	—
<i>A. laxifolia</i> , Hooker, icon. plant. t. 573 (1843)	—	—	T.	—	—	—	—	B.f.VI, 242	—
<i>A. selaginoides</i> , D. Don in Trans. Linn. Soc. XVIII, 172, t. 14 (1839) — — — — —	T.	—	—	—	—	—	—	B.f.VI, 242	M.fr.XI, 104.
CALLITRIS, Ventenat, decas generum novorum 10 (1808). (Frenela, Actinostrobus, Leichhardtia, Octoclinis.)									
<i>C. Macleayana</i> , F. v. M. in Rep. Burdek. Exped. 17 (1860)	—	—	—	—	N.S.W.	Q.	—	B.f.VI, 235	M.fr.XI, 104.
<i>C. Parlatores</i> , F. v. M. in Seemann Journ. of Bot. 267 (1866)	—	—	—	—	N.S.W.	Q.	—	B.f.VI, 235	M.fr.V, 186; XI, 104.
<i>C. Roei</i> , Bentham & J. Hooker, gen. plant. III, 424 (1880)	W.A.	—	—	—	—	—	—	B.f.VI, 236	—
<i>C. Drummondii</i> , Bentham & J. Hooker, gen. plant. III, 424 (1880) ...	W.A.	—	—	—	—	—	—	B.f.VI, 236	—
<i>C. verrucosa</i> , R. Brown in Mém. du Mus. Par. XIII, 74 (1826) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.VI, 236	M.fr.V, 237.
<i>C. columellaris</i> , F. v. M., fragm. V, 198 (1866)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V, 198.
<i>C. Muelleri</i> , Bentham & J. Hooker, gen. plant. III, 424 (1880) ...	—	—	—	—	N.S.W.	—	—	B.f.VI, 237	—
<i>C. cupressiformis</i> , Ventenat, decas gen. nov. 10 (1808)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VI, 237	—
<i>C. calcarata</i> , R. Brown in Mém. du Mus. Par. XIII, 74 (1826) ...	—	—	—	V.	N.S.W.	Q.	—	B.f.VI, 238	—
<i>C. oblonga</i> , L. Cl. Richard, comment. de conf. 49, t. 18 (1826) ...	—	—	T.	—	—	—	—	B.f.VI, 238	—
<i>C. Actinostrobus</i> , F. v. M., Rep. Burdek. Exped. 19 (1860)	W.A.	—	—	—	—	—	—	B.f.VI, 240	—
<i>C. acuminata</i> , F. v. M.; Actinostrobus, Parlatores, enum. sem. hort. Florent. 25 (1862)	W.A.	—	—	—	—	—	—	B.f.VI, 240	M.fr.XI, 104.
FITZROYA, J. Hooker in Journ. of the hort. Soc. VI, 264 (1851). (Diselma.)									
<i>F. Archeri</i> , Bentham & J. Hooker, gen. pl. III, 426 (1880)	—	—	T.	—	—	—	—	B.f.VI, 240	—
PHEROSPHAERA, Archer in Hooker's Kew Misc. II, 52 (1850). (Dacrydium partly.)									
<i>P. Hookeriana</i> , Archer in Hooker, Kew Misc. II, 52 (1850)	—	—	T.	—	—	—	—	B.f.VI, 245	M.fr.XI, 103.
<i>P. Fitzgeraldi</i> , F. v. M. in J. Hooker, icon. pl. XIV, 64, t. 1383 (1882) — — — — —	—	—	—	—	N.S.W.	—	—	—	M.fr.XI, 102, 138.
MICROCACHRYIS, J. Hooker in Lond. Journ. of Bot. IV, 149 (1845). (Dacrydium partly.)									
<i>M. tetragona</i> , J. Hooker, Fl. Tasman. I, 358, t. 100 (1860)	—	—	T.	—	—	—	—	B.f.VI, 241	—
DACRYDIUM, Solander in G. Forster, pl. escul. insul. ocean. austr. 80 (1786).									
<i>D. Franklinii</i> , J. Hooker in Lond. Journ. of Bot. IV, 152, t. 6 (1845) ...	—	—	T.	—	—	—	—	B.f.VI, 245	—
NAGEIA, Gaertner, de fructib. t. 39 (1788). (Podocarpus.)									
<i>N. elata</i> , F. v. M.; Podoc., R. Brown in Mém. de Mus. XIII, 75 (1826) — — — — —	—	—	—	—	N.S.W.	Q.	—	B.f.VI, 247	—
<i>N. spinulosa</i> , F. v. M.; Podocarpus, R. Brown in Mém. du Mus. XIII, 75 (1826)	—	—	—	—	N.S.W.	—	—	B.f.VI, 247	—
<i>N. Drouyniana</i> ; Podocarpus, F. v. M., fragm. IV, 86 (1864)	W.A.	—	—	—	—	—	—	B.f.VI, 247	M.fr.VI, 251; XI, 104.
<i>N. alpina</i> , F. v. M. in papers Roy. Soc. of Tasm. 23 (1879)	—	—	T.	V.	N.S.W.	—	—	B.f.VI, 248	M.fr.XI, 104.
THALAMIA, Sprengel, Anleit. zur Kenntn. der Gew. zweite Ausg. II, 218 (1817). (Phyllocladus.)									
<i>T. asplenifolia</i> , Sprengel, Anl. zur Kenntn. der Gew. II, 218 (1817) — — — — —	T.	—	—	—	—	—	—	B.f.VI, 246	M.fr.XI, 104.

CYCADEAE.

L. C. Richard in Persoon, synops. 630 (1807).

CYCAS, Linné, hort. Cliffort. 482 (1737).									
<i>C. media</i> , R. Brown, prodr. 348 (1810)	—	—	—	—	—	Q.	N.A.	B.f.VI, 249	M.fr.VIII, 171.
<i>C. Normanbyana</i> , F. v. M., fragm. VIII, 169 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII, 169.
<i>C. Kennedyana</i> , F. v. M. in Melb. Chemist, March (1882)	—	—	—	—	—	Q.	—	—	—
<i>C. Cairnsiana</i> , F. v. M., fragm. X, 63 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X, 63, 121.

- ENCEPHALARTOS, Lehmann, pugill. VI, 9, t. I, et 3 (1834). (Macrozamia, Lepidozamia, Catakidozamia, Zamia partly.)
 E. Pauli Guilielmi, F. v. M. in Trans. Pharm. Soc. Vict. II, 91 (1859) — — — — N.S.W. Q. — B.fl. VI, 251 M.fr. I, 86, 243; IX, 124.
 E. tridentatus, Lehmann, pugill. VI, 13 (1834) ... — — — — N.S.W. — — B.fl. VI, 253 M.fr. III, 38; XI, 125.
 E. spiralis, Lehmann, pugill. VI, 13 (1834) ... — — — — N.S.W. Q. — B.fl. VI, 251 M.fr. I, 41; II, 179; VIII, 173
 E. Douglasii, F. v. M.; Macrozamia, Hill (inedit.) ... — — — — Q. — — — [167; VIII, 173; XI, 126.
 E. Fraseri, Miquel in Versl. K. Akad. Wet. Amst. XV, 368 (1863) W.A. — — — — — B.fl. VI, 252 M.fr. I, 41, 243; II, 179; III,
 E. Moorei, F. v. M., fragm. XI, 125 (1881) ... — — — — Q. — — M.fr. I, 243; XI, 125.
 E. Macdonnelli, F. v. M. in Versl. K. Akad. Wet. Amst. XV, 376 (1863) — S.A. — — — — — M.fr. IX, 124; XI, 126. [124.
 E. Denisonii, F. v. M. in Journ. Pharm. Soc. Vict. II, 90 (1859) — — — — N.S.W. Q. — B.fl. VI, 253 M.fr. I, 41; VIII, 173; IX,
 BOWENIA, Hooker, Bot. Magaz. t. 5398 (1863).
 B. spectabilis, Hooker, Bot. Mag. t. 5398 (1863) ... — — — — — Q. — B.fl. VI, 254 M.fr. V, 171.

MONOCOTYLEDONEAE.

Ray, meth. plant. nov. (1682).

CALYCEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 41 (1880).

ORCHIDEAE.

Haller, enum. stirp. Helv. praefat. 33 (1742).

- PHOLIDOTA, Lindley in Hooker, Exotic Flora II, t. 138 (1825).
 P. imbricata, Lindley in Hook. Exot. Fl. II, t. 138 (1825) ... — — — — — Q. — B.fl. VI, 290 M.fr. IV, 163; V, 105.
 STURMIA, Reichenbach in Moessler's Handb. II, 1552 (1828). (Liparis, Richard 1818 not of Zoologists 1738.)
 S. reflexa, F. v. M., fragm. II, 72 (1861) ... — — — — N.S.W. Q. — B.fl. VI, 273 M.fr. III, 165; V, 102.
 S. angustilabris, F. v. M., fragm. IV, 164 (1864) ... — — — — Q. — B.fl. VI, 273 M.fr. IV, 164.
 S. coelogyneoides, F. v. M., fragm. II, 71 (1861) ... — — — — N.S.W. Q. — B.fl. VI, 273 M.fr. IX, 50.
 S. habenarina, F. v. M., fragm. IV, 131 (1864) ... — — — — Q. — B.fl. VI, 273 M.fr. V, 102; XI, 53.
 OBERONIA, Lindley, gen. and spec. of orchid. pl. 15 (1830). (Titania.)
 O. iridifolia, Lindley, gen. and spec. of orchid. pl. 15 (1830) ... — — — — N.S.W. Q. — B.fl. VI, 274 [88.
 O. palmicola, F. v. M., fragm. II, 24 (1860) ... — — — — N.S.W. Q. — B.fl. VI, 274 M.fr. II, 24; VII, 30; XI, 53,
 PHREATIA, Lindley, gen. and spec. of orchid. pl. 63 (1830). (Plexaure.)
 P. Limenophylax, G. Reichenbach in Seem. Bonpl. 54 (1857) ... — — — — — Q. — B.fl. VI, 290 M.fr. X, 64.
 DENDROBIUM, Swartz in nov. act. Upsal. VI, 82 (1799). (Thelychiton partly, Pedilonum, Coelandria.)
 D. bigibbum, Lindley in Paxt. Flower Gard. III, 25 (1853) ... — — — — — Q. N.A. B.fl. VI, 277 M.fr. VI, 119; XI, 87.
 D. dicuphum, F. v. M., fragm. VIII, 28 (1871) ... — — — — — N.A. B.fl. VI, 277 M.fr. VI, 119.
 D. Sumneri, F. v. M., fragm. VI, 94 (1868) ... — — — — — Q. — B.fl. VI, 278 M.fr. VI, 94.
 D. superbiens, G. Reichenbach in Garden. Chron. 516 (1876) ... — — — — — Q. — — M.fr. XI, 22.
 D. Phalaenopsis, Fitzgerald in Gardeners' Chronicle, July (1880) ... — — — — — Q. — —
 D. undulatum, R. Brown, prodr. 332 (1810) ... — — — — — Q. — B.fl. VI, 278 M.fr. I, 87.
 D. Johannis, G. Reichenbach in Garden. Chron. 890 (1865) ... — — — — — Q. — B.fl. VI, 279
 D. speciosum, Smith, Exot. Botany I, 17, t. 10 (1804) ... — — — — V. N.S.W. Q. — B.fl. VI, 279 M.fr. VIII, 249.
 D. falconirostre, Fitzgerald in Sydney Morn. Herald, Nov. 18 (1876) ... — — — — N.S.W. — — M.fr. XI, 87.
 D. tetragonum, Cunningham in Bot. Regist. XXV, Misc. 33 (1839) ... — — — — N.S.W. Q. — B.fl. VI, 279 M.fr. I, 87; VI, 119.
 D. aemulum, R. Brown, prodr. 333 (1810) ... — — — — N.S.W. Q. — B.fl. VI, 280 M.fr. I, 213; XI, 87.
 D. Kingianum, Bidwill in Bot. Regist. XXV, Misc. 11 (1839) ... — — — — N.S.W. Q. — B.fl. VI, 280 M.fr. III, 60.
 D. brachypus, G. Reichenbach in Garcke, Linnaea XLI, 42 (1876) ... — — — — N.S.W. — —
 D. gracilicaule, F. v. M., fragm. I, 179 (1859) ... — — — — N.S.W. Q. — B.fl. VI, 281 M.fr. IV, 171.
 D. Moorei, F. v. M., fragm. VII, 29 (1870) ... — — — — N.S.W. — — B.fl. VI, 281 M.fr. XI, 54.
 D. agrostophyllum, F. v. M., fragm. VIII, 28 (1871) ... — — — — — Q. — B.fl. VI, 282 M.fr. VIII, 28.
 D. Baileyi, F. v. M., fragm. VIII, 173 (1874) ... — — — — — Q. — — M.fr. VIII, 173.
 D. Smilliae, F. v. M., fragm. VI, 94 (1868) ... — — — — — Q. — B.fl. VI, 282 M.fr. VII, 63; XI, 87.
 D. ophioglossum, G. Reichenbach in Journ. L. Soc. XV, 113 (1876) ... — — — — — Q. — — M.fr. XI, 54.
 D. canaliculatum, R. Brown, prodr. 333 (1810) ... — — — — — Q. — B.fl. VI, 282 M.fr. III, 126.
 D. Foelschei, F. v. M. in Wing's S. Sc. Record, Oct. (1882) ... — — — — — N.A. — —
 D. monophyllum, F. v. M., fragm. I, 189 (1859) ... — — — — N.S.W. Q. — B.fl. VI, 283 M.fr. VII, 64.
 D. hispidum, A. Richard, sert. Astrol. 13, t. 5 (1833) ... — — — — — Q. — B.fl. VI, 283 M.fr. VII, 30.
 D. cucumerinum, MacLeay in Bot. Reg., new ser. V, Misc. 58 (1842) ... — — — — N.S.W. Q. — B.fl. VI, 283 M.fr. III, 59; VIII, 151.
 D. pugioniforme, Cunningham in Bot. Reg. XXV, Misc. 33 (1839) ... — — — — N.S.W. Q. — B.fl. VI, 284 M.fr. I, 189.
 D. rigidum, R. Brown, prodr. 333 (1810) ... — — — — — Q. — B.fl. VI, 284 M.fr. VIII, 248; XI, 87.
 D. linguiforme, Swartz in Svensk. Vetensk. Ac. Handl. 247 (1800) ... — — — — N.S.W. Q. — B.fl. VI, 284
 D. teretifolium, R. Brown, prodr. 333 (1810) ... — — — — N.S.W. Q. — B.fl. VI, 285 M.fr. I, 89.
 D. Fairfaxii, Fitzgerald & F. v. M. in Sydney Mail 360, Nov. (1872) ... — — — — N.S.W. — — B.fl. VI, 285 M.fr. X, 122.
 D. striolatum, G. Reichenbach in Hamburg Gartenz. 313 (1857) ... — — T. V. N.S.W. — — B.fl. VI, 285 M.fr. I, 88; VII, 30.
 D. Mortii, F. v. M., fragm. II, 93 (1860) ... — — — — N.S.W. Q. — B.fl. VI, 286 M.fr. II, 93.
 D. Beckleri, F. v. M., fragm. V, 95 (1865) ... — — — — N.S.W. Q. — B.fl. VI, 286
 BULBOPHYLLUM, Petit-Thouars, Orchid. Afric. t. 95 et 108 (1822). (Bulbophyllum, Thelychiton partly.)
 B. Lichenastrum, F. v. M., fragm. VII, 60 (1869) ... — — — — — Q. — B.fl. VI, 287 M.fr. X, 119.
 B. Baileyi, F. v. M., fragm. IX, 5 (1875) ... — — — — — Q. — — M.fr. IX, 5.
 B. nematopodum, F. v. M., fragm. VIII, 30 (1873) ... — — — — — Q. — B.fl. VI, 287 M.fr. VIII, 130.

B. Shepherdii, F. v. M., fragm. III, 40 (1862)	—	—	—	N.S.W.	Q.	—	B.f. VI, 288	M.fr. I, 190.
B. Taylori, F. v. M., fragm. VIII, 150 (1874)	—	—	—	—	Q.	—	—	M.fr. VIII, 150.
B. aurantiacum, F. v. M., fragm. III, 39 (1862)	—	—	—	N.S.W.	Q.	—	B.f. VI, 288	M.fr. VII, 98.
B. argyropus, G. Reichenbach in Garcke, <i>Linnaea</i> XLI, 42 (1876)	—	—	—	N.S.W.	—	—	—	—
B. exiguum, F. v. M., fragm. II, 72 (1861)	—	—	—	N.S.W.	Q.	—	B.f. VI, 288	M.fr. XI, 22.
B. minutissimum, F. v. M., fragm. XI, 53 (1878)	—	—	—	N.S.W.	Q.	—	—	M.fr. V, 85; XI, 88.
B. Prenticei, F. v. M. in Wing's South. Sc. Record I, 173 (1881)	—	—	—	—	Q.	—	—	B.f. X, 139.
B. Elisae, F. v. M., fragm. VI, 120 (1868)	—	—	—	N.S.W.	Q.	—	B.f. VI, 289	M.fr. VIII, 28.
SARCOCHILUS, R. Brown, prodr. 332 (1810). (<i>Thrixperum</i> , <i>Gunnia</i> , <i>Cleisostoma</i> .)								
S. erectus; <i>Cleisostoma</i> , Fitzgerald, Austr. Orchids part 4 (1877)	—	—	—	N.S.W.	—	—	—	M.fr. XI, 52, 139.
S. tridentatus, G. Reichenbach in Walp. <i>Annal.</i> VI, 500 (1861)	—	—	—	N.S.W.	Q.	—	B.f. VI, 296	M.fr. I, 192; II, 181, VII, 98.
S. Armitii, F. v. M., fragm. IX, 49 (1875)	—	—	—	—	Q.	—	—	M.fr. IX, 49.
S. Beckleri; <i>Cleisostoma</i> , F. v. M. in Benth. <i>Fl. Austr.</i> VI, 296 (1870)	—	—	—	N.S.W.	—	—	B.f. VI, 296	M.fr. VIII, 248.
S. brevibrabris; <i>Cleisostoma</i> , F. v. M., fragm. XI, 87 (1880) ...	—	—	—	—	Q.	—	—	M.fr. VIII, 248; XI, 139.
S. Macphersoni, F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	Q.	—	B.f. VI, 297	M.fr. VII, 96.
S. Hartmanni, F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	Q.	—	—	M.fr. VIII, 248.
S. divitiflorus, F. v. M. in Benth. <i>Fl. Austr.</i> VI, 292 (1870) ...	—	—	—	N.S.W.	Q.	—	B.f. VI, 292	M.fr. IX, 50; XI, 22.
S. falcatus, R. Brown, prodr. 332 (1810)	—	—	—	N.S.W.	Q.	—	B.f. VI, 293	M.fr. I, 89; VII, 87; X, 50.
S. Fitzgeraldi, F. v. M., fragm. VII, 115 (1870)	—	—	—	N.S.W.	Q.	—	B.f. VI, 293	M.fr. VII, 115.
S. olivaceus, Lindley, Bot. <i>Regist.</i> Misc. XXV, 32 (1839) ...	—	—	—	N.S.W.	Q.	—	B.f. VI, 293	M.fr. I, 90, 191; VII, 97.
S. parviflorus, Lindley, Bot. <i>Regist.</i> Misc. XXIV, 34 (1838) ...	—	—	T. V.	N.S.W.	—	—	B.f. VI, 294	M.fr. I, 89; VII, 98.
S. Ceciliae, F. v. M., fragm. V, 42, t. 42 (1865)	—	—	—	—	Q.	—	B.f. VI, 294	M.fr. VII, 98; XI, 22.
S. Hillii, F. v. M., fragm. II, 94 (1861)	—	—	—	N.S.W.	Q.	—	B.f. VI, 295	M.fr. I, 88; III, 39; VII, 98.
S. phyllorrhizus, F. v. M., fragm. V, 201 (1866)	—	—	—	—	Q.	—	B.f. VI, 295	M.fr. VII, 30.
TAENIOPHYLLUM, Blume, <i>Bijdr.</i> 355 (1825).								
T. Muelleri, Lindley in Benth. <i>Fl. Austr.</i> VI, 291 (1870) ...	—	—	—	—	Q.	—	B.f. VI, 291	—
ORNITHOCHILUS, Wallich in Lindley, gen. and spec. of orchid. plants 242 (1833). (<i>Saccolabium</i> partly.)								
O. Hillii, Benth in Journ. Linn. Soc. XVIII, 334 (1880)	—	—	—	N.S.W.	Q.	—	B.f. VI, 298	M.fr. VII, 97.
GEODORUM, Jackson in Andrews, Botan. <i>Reposit.</i> t. 626 (1810).								
G. pictum, Lindley, gen. and spec. of orchid. pl. 175 (1833) ...	—	—	—	—	Q.	N.A.	B.f. VI, 299	M.fr. III, 24; V, 102; VIII, [31.
EULOPHIA, R. Brown in Edwards, Bot. <i>Regist.</i> 686 (1822). (<i>Eulophus</i> 1822.)								
E. venosa, G. Reichenbach in Benth. <i>Fl. Austr.</i> VI, 300 (1870)	—	—	—	—	Q.	N.A.	B.f. VI, 300	M.fr. I, 61.
E. Fitzalani, F. v. M., fragm. VIII, 30 (1872)	—	—	—	—	Q.	—	B.f. VI, 300	M.fr. VII, 30.
DIPODIUM, R. Brown, prodr. 330 (1810).								
D. punctatum, R. Brown, prodr. 331 (1810)	—	S.A.	T. V.	N.S.W.	Q.	N.A.	B.f. VI, 301	M.fr. X, 64.
D. ensifolium, F. v. M., fragm. V, 42 (1865)	—	—	—	—	Q.	—	B.f. VI, 301	M.fr. V, 42.
CYMBIDIUM, Swartz in nov. act. Upsal. VI, 70 (1799).								
C. Hillii, F. v. M. in Regel's <i>Gartenflora</i> , 138 (1879)	—	—	—	—	Q.	—	—	M.fr. XI, 88.
C. canaliculatum, R. Brown, prodr. 331 (1810)	—	S.A.	—	N.S.W.	Q.	N.A.	B.f. VI, 302	M.fr. XI, 88.
C. albuciflorum, F. v. M., fragm. I, 188 (1859)	—	—	—	—	Q.	—	B.f. VI, 303	M.fr. I, 188.
C. suave, R. Brown, prodr. 331 (1810)	—	—	—	N.S.W.	Q.	—	B.f. VI, 303	M.fr. I, 187.
SPATHOGLOTTIS, Blume, <i>Bijdr.</i> 400 (1825).								
S. Paulinae, F. v. M., fragm. VI, 95 (1868)	—	—	—	—	Q.	—	B.f. VI, 304	M.fr. X, 64.
PHAJUS, Loureiro, <i>Fl. Cochinch.</i> II, 529 (1790).								
P. grandifolius, Loureiro, <i>Fl. Cochinch.</i> II, 529 (1790)	—	—	—	N.S.W.	Q.	—	B.f. VI, 304	M.fr. I, 42; IV, 163; X, 6
CALANTHE, R. Brown in Edwards, Bot. <i>Regist.</i> 573 (1821).								
C. veratrifolia, R. Brown in Bot. <i>Regist.</i> 573 (1821)	—	—	—	N.S.W.	Q.	—	B.f. VI, 305	M.fr. VIII, 30; IX, 50.
GALEOLA, Loureiro, <i>Fl. Cochinch.</i> II, 520 (1790). (<i>Erythrorchis</i> , <i>Ledgeria</i> .)								
G. cassythoides, G. Reichenbach, Xen. <i>Orchid.</i> II, 77 (1869) ...	—	—	—	N.S.W.	Q.	—	B.f. VI, 307	M.fr. II, 167; V, 102; VIII, 31; [XI, 105.
G. foliata, F. v. M., fragm. VIII, 31 (1872)	—	—	—	N.S.W.	Q.	—	B.f. VI, 307	M.fr. II, 167; V, 102; XI, 127.
EPIPOGUM, Gmelin, <i>Fl. Sibir.</i> I, 11, t. 2 (1747).								
E. nutans, Lindley in Journ. Linn. Soc. I, 177 (1856)	—	—	—	N.S.W.	Q.	—	B.f. VI, 308	M.fr. IX, 50; XI, 105.
GASTRODIA, R. Brown, prodr. 330 (1810).								
G. sesamoides, R. Brown, prodr. 330 (1810)	—	—	T. V.	N.S.W.	Q.	—	B.f. VI, 309	M.fr. VIII, 151; X, 64.
POGONIA, A. L. de Jussieu, gen. plant. 65 (1789).								
P. uniflora, F. v. M., fragm. V, 201 (1866)	—	—	—	—	Q.	—	B.f. VI, 310	M.fr. V, 201.
P. holochila, F. v. M., fragm. V, 200 (1866)	—	—	—	—	Q.	—	B.f. VI, 310	M.fr. V, 200.
P. Dallachyana, F. v. M. in Benth. <i>Fl. Austr.</i> VI, 310 (1870) ...	—	—	—	—	Q.	—	B.f. VI, 310	—
P. pachystomoides, F. v. M., fragm. VIII, 174 (1874)	—	—	—	—	Q.	—	—	M.fr. VIII, 174.
CORYMBORCHIS, Petit-Thouars, hist. des plant. <i>Orchid.</i> recueill. sur les isles d'Afrique, t. 37-38 (1822). (<i>Corymbis</i> .)								
C. veratrifolia, Blume, <i>Collect. des Orchidées</i> 125, t. 42 & 43 (1858)	—	—	—	—	Q.	—	B.f. VI, 311	M.fr. VIII, 30; X, 64.
ETAERIA, Blume, <i>Bijdr.</i> 409 (1825). (<i>Hetaeria</i> , <i>Ramphidia</i> .)								
E. tenuis, Benth in Journ. Linn. Soc. XVIII, 345 (1880)	—	—	—	—	Q.	—	B.f. VI, 312	—
MICROSTYLIS, Nuttall, <i>Genera of North American plants</i> II, 196 (1818).								
M. Bernaysii, F. v. M., fragm. XI, 21 (1878)	—	—	—	—	Q.	—	—	M.fr. XI, 21.
GOODYERA, R. Brown in Aiton, hort. Kew. sec. ed. V, 197 (1813).								
G. viridiflora, Blume, <i>Coll. des Orchid.</i> 41, t. 9 (1858)	—	—	—	—	Q.	—	B.f. VI, 313	M.fr. VIII, 29. [64.
G. polygonoides, F. v. M., fragm. VIII, 29 (1872)	—	—	—	—	Q.	—	B.f. VI, 313	M.fr. VII, 30; VIII, 29; X,
SPIRANTHES, L. C. Richard in Mém. du Mus. IV, 40 (1818). (<i>Gyrostachis</i> , 1807).								
S. australis, Lindley, Bot. <i>Regist.</i> 823 (1824)	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VI, 314	M.fr. X, 65.
THELYMITRA, R. & G. Forster, char. gen. 97, t. 49 (1776). (<i>Macdonaldia</i> .)								
T. ixioides, Swartz in Kong. Svensk. Acad. <i>Handl.</i> 223, t. 3 (1800) W.A.	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VI, 317	—
T. circumsepta, Fitzgerald, Austral. <i>Orchid.</i> , part 4 (1877) ...	—	—	—	N.S.W.	—	—	—	—

T. crinita, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	B.f. VI, 319	M.fr. VI, 84.
T. aristata, Lindley, gen. and spec. of orchid. pl. 521 (1840) ...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f. VI, 319	
T. longifolia, R. & G. Forster, char. gen. 98, t. 49 (1776) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f. VI, 319	
T. villosa, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	B.f. VI, 320	M.fr. V, 94, 214.
T. tigrina, R. Brown, prodr. 315 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 320	
T. fusco-lutea, R. Brown, prodr. 315 (1810) ...	W.A.	S.A.	—	—	—	—	B.f. VI, 321	M.fr. V, 94.
T. stellata, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	B.f. VI, 321	
T. carnea, R. Brown, prodr. 314 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. VI, 321	M.fr. V, 98.
T. flexusa, Endlicher, nov. stirp. decad. 23 (1839) ...	W.A.	S.A.	T.	V.	—	—	B.f. VI, 322	
T. antennifera, J. Hooker, Fl. Tasman. II, 4, t. 101 (1860) ...	W.A.	S.A.	T.	V.	—	—	B.f. VI, 322	
T. Macmillani, F. v. M., fragm. V, 93 (1865) ...	—	—	—	V.	—	—	B.f. VI, 323	M.fr. V, 93.
T. Mackibbinii, F. v. M. in Melb. Chemist 44 (1881) ...	W.A.	S.A.	—	V.	—	—	—	M.fr. XI, 139.
T. variegata, Lindley in Benth. Fl. Austr. VI, 323 (1870) ...	W.A.	—	—	—	—	—	B.f. VI, 323	M.fr. V, 97, 98.
T. venosa, R. Brown, prodr. 314 (1870) ...	—	—	—	—	N.S.W.	—	B.f. VI, 323	
T. cyanea, Lindley in Benth. Fl. Austr. VI, 323 (1870) ...	—	—	T.	—	—	—	B.f. VI, 323	
EPIBLEMA, R. Brown, prodr. fl. Nov. Holl. 315 (1810).								
E. grandiflorum, R. Brown, prodr. 315 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 324	M.fr. XI, 21.
DIURIS, Smith in Transact. Linn. Soc. IV, 222 (1798).								
D. alba, R. Brown, prodr. 316 (1810) ...	—	—	—	—	N.S.W.	Q.	B.f. VI, 325	M.fr. V, 102.
D. punctata, Smith, Exot. Bot. I, 13, t. 8 (1804) ...	—	S.A.	—	V.	N.S.W.	—	B.f. VI, 326	M.fr. V, 102.
D. secundiflora, Fitzgerald, Austral. Orchids, part 4 (1877) ...	—	—	—	—	N.S.W.	—	—	
D. aurea, Smith, Exot. Bot. 15, t. 9 (1804) ...	—	—	—	—	N.S.W.	Q.	B.f. VI, 327	M.fr. V, 172.
D. palustris, Lindley, gen. and spec. of orchid. pl. 507 (1840) ...	—	S.A.	T.	V.	—	—	B.f. VI, 327	M.fr. V, 102.
D. maculata, Smith, Exot. Bot. I, 57, t. 30 (1804) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f. VI, 327	M.fr. V, 102.
D. pedunculata, R. Brown, prodr. 316 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. VI, 328	M.fr. V, 102, 173.
D. pallens, Benth. Fl. Austr. VI, 329 (1870) ...	—	—	—	—	N.S.W.	—	B.f. VI, 329	
D. abbreviata, F. v. M. in Benth. Fl. Austr. VI, 329 (1870) ...	—	—	—	—	N.S.W.	Q.	B.f. VI, 329	
D. setacea, R. Brown, prodr. 316 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 329	M.fr. V, 101.
D. emarginata, R. Brown, prodr. 316 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 330	
D. sulphurea, R. Brown, prodr. 316 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. VI, 330	M.fr. V, 102, 173.
D. aequalis, F. v. M. in Fitzgerald, Austral. orch. part 2 (1876) ...	—	—	—	—	N.S.W.	—	—	M.fr. X, 65.
D. dendrobioides, Fitzgerald, Austral. Orchids, part 7 (1882) ...	—	—	—	—	N.S.W.	—	—	
D. longifolia, R. Brown, prodr. 316 (1810) ...	W.A.	S.A.	T.	V.	—	—	B.f. VI, 331	M.fr. V, 101, 172.
D. pauciflora, R. Brown, prodr. 316 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 331	
ORTHOCERAS, R. Brown, prodr. fl. Nov. Holl. 317 (1810).								
O. strictum, R. Brown, prodr. 317 (1810) ...	—	S.A.	—	V.	N.S.W.	—	B.f. VI, 332	
CALOCHILUS, R. Brown, prodr. fl. Nov. Holl. 320 (1810).								
C. campestris, R. Brown, prodr. 320 (1810) ...	—	—	T.	V.	N.S.W.	Q.	B.f. VI, 315	M.fr. VIII, 151.
C. Robertsoni, Benth. Fl. Austr. VI, 315 (1870) ...	—	S.A.	—	V.	—	—	B.f. VI, 315	
C. paludosus, R. Brown, prodr. 320 (1810) ...	—	—	—	—	N.S.W.	Q.	B.f. VI, 316	
CRYPTOSTYLIS, R. Brown, prodr. 317 (1810). (Zosterostylis.)								
C. longifolia, R. Brown, prodr. 317 (1810) ...	—	—	T.	V.	N.S.W.	Q.	B.f. VI, 333	M.fr. XI, 21.
C. ovata, R. Brown, prodr. 317 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 334	
C. erecta, R. Brown, prodr. 317 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 334	
C. leptochila, F. v. M. in Benth. Fl. Austr. VI, 324 (1870) ...	—	—	—	—	N.S.W.	—	B.f. VI, 334	
PRASOPHYLLUM, R. Brown, prodr. 317 (1810). (Genoplesium.)								
P. australe, R. Brown, prodr. 318 (1810) ...	—	S.A.	T.	V.	—	—	B.f. VI, 337	
P. flavum, R. Brown, prodr. 318 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 337	
P. elatum, R. Brown, prodr. 318 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	B.f. VI, 337	M.fr. V, 100.
P. brevifolium, J. Hooker, Fl. Tasman II, 11, t. 110 (1860) ...	—	—	T.	V.	N.S.W.	Q.	B.f. VI, 338	M.fr. V, 101.
P. hians, G. Reichenbach, Beitr. zur syst. Pflanzenk. 61 (1871) ...	W.A.	—	—	—	—	—	B.f. VI, 338	
P. patens, R. Brown, prodr. 318 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f. VI, 339	M.fr. V, 101.
P. fuscum, R. Brown, prodr. 318 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	B.f. VI, 339	M.fr. V, 101.
P. cyphochilum, Benth. Fl. Austr. VI, 340 (1870) ...	W.A.	—	—	—	—	—	B.f. VI, 340	
P. ovale, Lindley, Bot. Regist. XXV, App. LIV (1839) ...	W.A.	—	—	—	—	—	B.f. VI, 341	
P. macrostachyum, R. Brown, prodr. 318 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 341	
P. Fimbria, G. Reichenbach, Beitr. zur syst. Pflanzenk. 60 (1871) ...	W.A.	—	—	—	—	—	B.f. VI, 341	
P. striatum, R. Brown, prodr. 318 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 342	M.fr. V, 100.
P. parvifolium, Lindley, Bot. Regist. XXV, App. LIV (1839) ...	W.A.	—	—	—	—	—	B.f. VI, 342	M.fr. V, 101.
P. gibbosum, R. Brown, prodr. 331 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 342	M.fr. V, 100.
P. cucullatum, G. Reichenbach, Beitr. zur syst. Pflanzenk. 59 (1871) ...	W.A.	—	—	—	—	—	B.f. VI, 343	
P. nigricans, R. Brown, prodr. 319 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	B.f. VI, 343	M.fr. IX, 50.
P. rufum, R. Brown, prodr. 319 (1810) ...	—	—	T.	V.	N.S.W.	Q.	B.f. VI, 344	M.fr. V, 100.
P. brachystachyum, Lindley, gen. and spec. of orchid. pl. 513 (1840) ...	—	—	T.	—	—	—	B.f. VI, 344	M.fr. V, 100.
P. despectans, J. Hooker, Fl. Tasman. II, 13, t. 113 (1860) ...	—	S.A.	T.	V.	—	—	B.f. VI, 345	M.fr. V, 100.
P. fimbriatum, R. Brown, prodr. 319 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 345	M.fr. V, 100.
P. Archeri, J. Hooker, Fl. Tasman. II, 14, t. 113 (1860) ...	—	—	T.	V.	—	—	B.f. VI, 345	M.fr. V, 100.
P. intricatum, C. Stuart in Benth. Fl. Austr. VI, 346 (1870) ...	—	—	T.	—	—	—	B.f. VI, 346	
P. Woollii, F. v. M., fragm. V, 100 (1866) ...	—	—	—	—	N.S.W.	—	B.f. VI, 346	M.fr. V, 100.
MICROTIS, R. Brown, prodr. fl. Nov. Holl. 320 (1810).								
M. porrifolia, R. Brown, prodr. 320 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.f. VI, 347	M.fr. X, 65; XI, 21.
M. media, R. Brown, prodr. 321 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 348	
M. alba, R. Brown, prodr. 321 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 348	
M. pulchella, R. Brown, prodr. 321 (1810) ...	W.A.	—	—	—	—	—	B.f. VI, 349	
M. atrata, Lindley, Bot. Regist. XXV, App. LIV (1839) ...	W.A.	S.A.	—	V.	—	—	B.f. VI, 349	M.fr. I, 90, 244; X, 65.
CORYSANTHES, R. Brown, prodr. 328 (1810). (Coryhas, Nematoceras.)								
C. unguiculata, R. Brown, prodr. 328 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 350	
C. pruinosa, R. Cunningham in N.S. Wales, Magaz. n. 1 (1833) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	M.fr. IX, 49.

<i>C. fimbriata</i> , R. Brown, prodr. 328 (1810) ...	—	—	—	—	N.S.W.	—	B.f. VI, 351	M.fr. X, 65.
<i>C. bicalcarata</i> , R. Brown, prodr. 328 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 351
PTEROSTYLIS, R. Brown, prodr. fl. Nov. Holl. 326 (1810).								
<i>P. ophioglossa</i> , R. Brown, prodr. 326 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 354
<i>P. concinna</i> , R. Brown, prodr. 326 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VI, 355
<i>P. curta</i> , R. Brown, prodr. 326 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 355
<i>P. acuminata</i> , R. Brown, prodr. 326 (1810) ...	—	—	—	V.	N.S.W.	—	—	B.f. VI, 355
<i>P. nutans</i> , R. Brown, prodr. 327 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 356
<i>P. pedaloglossa</i> , Fitzgerald, Austral. Orchids, part 3 (1877) ...	—	—	—	—	N.S.W.	—	—	—
<i>P. pedunculata</i> , R. Brown, prodr. 327 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 356
<i>P. nana</i> , R. Brown, prodr. 327 (1810) ...	W.A.	S.A.	T.	V.	—	—	—	B.f. VI, 357
<i>P. cucullata</i> , R. Brown, prodr. 327 (1810) ...	—	S.A.	T.	V.	—	—	—	B.f. VI, 357
<i>P. grandiflora</i> , R. Brown, prodr. 327 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 358
<i>P. truncata</i> , Fitzgerald, Austral. Orchids, part 4 (1878) ...	—	—	—	—	N.S.W.	—	—	—
<i>P. reflexa</i> , R. Brown, prodr. 327 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. VI, 359
<i>P. praecox</i> , Lindley, gen. and spec. of Orchid. pl. 388 (1840) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 359
<i>P. obtusa</i> , R. Brown, prodr. 327 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VI, 360
<i>P. recurva</i> , Benth. Fl. Austr. VI, 360 (1870) ...	W.A.	—	—	—	—	—	—	B.f. VI, 360
<i>P. parviflora</i> , R. Brown, prodr. 327 (1810) ...	—	—	T.	V.	N.S.W.	Q.	—	B.f. VI, 361
<i>P. barbata</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 362
<i>P. turfosa</i> , Endlicher in Lehm. pl. Preiss. II, 5 (1846) ...	W.A.	—	—	—	—	—	—	B.f. VI, 362
<i>P. mutica</i> , R. Brown, prodr. 328 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 362
<i>P. rufa</i> , R. Brown, prodr. 327 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 363
<i>P. daintreana</i> , F. v. M. in Benth. Fl. Austr. VI, 360 (1870) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 360
<i>P. longifolia</i> , R. Brown, prodr. 327 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 364
<i>P. vittata</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	S.A.	—	V.	—	—	—	B.f. VI, 364
CALEYA, R. Brown in Aiton, Hort. Kew. sec. ed. V, 204 (1813). (<i>Calceana</i> , 1810.)								
<i>C. major</i> , R. Brown, prodr. 329 (1810) ...	—	—	T.	V.	N.S.W.	Q.	—	B.f. VI, 366
<i>C. minor</i> , R. Brown, prodr. 329 (1810) ...	—	—	T.	—	N.S.W.	—	—	B.f. VI, 366
<i>C. nigrita</i> , Lindley, Bot. Reg. XXV, App. LIV (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 366
<i>C. Sullivanii</i> , F. v. M. in Melb. Chemist 44 (1882) ...	—	—	—	V.	—	—	—	—
DRAKAEA, Lindley, Bot. Regist. XXV, App. LV (1839). (<i>Spiculaea</i> , <i>Arthrochilus</i> .)								
<i>D. ciliata</i> , G. Reichenbach, Beitr. zur syst. Pflanzenk. 68 (1871) ...	W.A.	—	—	—	—	—	—	B.f. VI, 367
<i>D. irritabilis</i> , G. Reichenbach, Beitr. zur syst. Pflanzenk. 68 (1871) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 368
<i>D. elastica</i> , Lindley, Bot. Regist. XXV, App. LV (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 368
ACIANTHUS, R. Brown, prodr. fl. Nov. Holl. 321 (1810).								
<i>A. caudatus</i> , R. Brown, prodr. 321 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VI, 369
<i>A. fornicatus</i> , R. Brown, prodr. 321 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VI, 370
<i>A. exsertus</i> , R. Brown, prodr. 321 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 370
<i>A. viridis</i> , J. Hooker, Fl. Tasman. II, 372 (1860) ...	—	—	T.	—	—	—	—	B.f. VI, 371
CYRTOSTYLIS, R. Brown, prodr. flor. Nov. Holl. 322 (1810).								
<i>C. reniformis</i> , R. Brown, prodr. 322 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 376
LYPERANTHUS, R. Brown, prodr. 325 (1810). (<i>Burnettia</i> .)								
<i>L. nigriscans</i> , R. Brown, prodr. 325 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 374
<i>L. Burnettii</i> , F. v. M., fragm. V, 96 (1865) ...	—	—	T.	V.	—	—	—	B.f. VI, 375
<i>L. Forrestii</i> , F. v. M. in Wing's S. Sc. Rec. II, 55 (1882) ...	W.A.	—	—	—	—	—	—	—
<i>L. ellipticus</i> , R. Brown, prodr. 325 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f. VI, 374
ERIOCHILUS, R. Brown, prodr. fl. Nov. Holl. 323 (1810).								
<i>E. autumnalis</i> , R. Brown, prodr. 323 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 372
<i>E. scaber</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 372
<i>E. tenuis</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 372
<i>E. dilatatus</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 372
<i>E. multiflorus</i> , Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 373
<i>E. fimbriatus</i> , F. v. M. in Wing's S. Sc. Record II, 152 (1882) ...	W.A.	S.A.	—	V.	—	—	—	B.f. VI, 377
CALADENIA, R. Brown, prodr. 325 (1810). (<i>Leptoceras</i> , <i>Adenochilus</i> .)								
<i>C. Menziesii</i> , R. Brown, prodr. 325 (1810) ...	W.A.	S.A.	T.	V.	—	—	—	B.f. VI, 377
<i>C. Nortoni</i> , Adenochilus, Fitzgerald, Austral. Orch. part 2 (1876) ...	—	—	—	—	N.S.W.	—	—	—
<i>C. Cairnsiana</i> , F. v. M., fragm. VII, 31 (1869) ...	W.A.	—	—	—	—	—	—	B.f. VI, 378
<i>C. multiclavia</i> , G. Reichenbach, Beitr. zur syst. Pflanzenk. 64 (1871) ...	W.A.	—	—	—	—	—	—	B.f. VI, 378
<i>C. discoidea</i> , Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 382
<i>C. Patersonii</i> , R. Brown, prodr. 324 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 382
<i>C. Drummondii</i> , Benth. Fl. Austr. VI, 383 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 383
<i>C. hirta</i> , Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 383
<i>C. Roei</i> , Benth. Fl. Austr. VI, 383 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 383
<i>C. Barbarossae</i> , G. Reichenbach, Beitr. z. syst. Pflanzenk. 64 (1871) ...	W.A.	—	—	—	—	—	—	B.f. VI, 383
<i>C. flava</i> , R. Brown, prodr. 324 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VI, 384
<i>C. latifolia</i> , R. Brown, prodr. 324 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 384
<i>C. reptans</i> , Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 385
<i>C. suaveolens</i> , G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871) ...	—	T.	V.	N.S.W.	—	—	—	B.f. VI, 385
<i>C. serrata</i> , G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871) ...	W.A.	—	—	—	—	—	—	B.f. VI, 385
<i>C. carnea</i> , R. Brown, prodr. 324 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VI, 386
<i>C. congesta</i> , R. Brown, prodr. 324 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VI, 387
<i>C. aphylla</i> , Benth. Fl. Austr. VI, 387 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VI, 387
<i>C. coerulea</i> , R. Brown, prodr. 324 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 388
<i>C. deformis</i> , R. Brown, prodr. 324 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VI, 388
<i>C. sericea</i> , Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 389
<i>C. gemmata</i> , Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VI, 389

C. ixioideis, Lindley, Bot. Regist. XXV, App. LII (1839)	..	W.A.	—	—	—	—	—	—	B.fl. VI, 389	
C. unita, Fitzgerald in Gardeners' Chronicle 461 (1882) ...	...	W.A.	—	—	—	—	—	—	—	
CHILOGLOTTIS, R. Brown, prodr. fl. 322 (1810).										
C. diphylla, R. Brown, prodr. 323 (1810)	...	—	—	T. V.	N.S.W.	Q.	—	—	B.fl. VI, 390	M.fr. X, 116.
C. trapeziformis, Fitzgerald, Austral. Orchids, part 3 (1877)	...	—	—	—	N.S.W.	—	—	—	—	M.fr. X, 116.
C. formicifera, Fitzgerald, Austral. Orchids, part 3 (1877)	...	—	—	—	N.S.W.	—	—	—	—	M.fr. X, 116. [116; XI, 22.
C. Gunnii, Lindley, gen. and spec. Orchid. 387 (1840) ...	...	—	—	T. V.	—	—	—	—	B.fl. VI, 391	M.fr. VIII, 151; IX, 50; X,
GLOSSODIA, R. Brown, prodr. 325 (1810).										
G. major, R. Brown, prodr. 326 (1810)	...	—	S.A.	T. V.	N.S.W.	Q.	—	—	B.fl. VI, 392	
G. minor, R. Brown, prodr. 326 (1810)	...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VI, 392	
G. Brunonis, Endlicher, nov. stirp. dec. 16 (1839) ...	...	W.A.	—	—	—	—	—	—	B.fl. VI, 393	M.fr. VI, 83.
G. intermedia, Fitzgerald in Gardeners' Chron. 462 (1882)	...	W.A.	—	—	—	—	—	—	—	
G. emarginata, Lindley, gen. and spec. of Orchid. pl. 424 (1840)	...	W.A.	—	—	—	—	—	—	B.fl. VI, 393	M.fr. XI, 22.
HABENARIA, Willdenow, spec. plant. IV, 44 (1805).										
H. trinervis, Wight, icon. plant. Ind. or. t. 1701 (1852)...	...	—	—	—	—	—	—	N.A.	B.fl. VI, 394	
H. elongata, R. Brown, prodr. 313 (1810)	...	—	—	—	—	—	—	Q. N.A.	B.fl. VI, 394	M.fr. VII, 15.
H. graminea, Lindley, gen. and spec. of orchid. pl. 318 (1840) ...	...	—	—	—	—	—	—	Q. N.A.	B.fl. VI, 394	M.fr. VII, 16.
H. ochroleuca, R. Brown, prodr. 313 (1810)	...	—	—	—	—	—	—	N.A.	B.fl. VI, 395	M.fr. V, 102.
H. xanthantha, F. v. M., fragm. VII, 16 (1869)	...	—	—	—	—	—	—	Q. N.A.	B.fl. VI, 395	M.fr. VII, 16.
APOSTASIA, Blume, Bijdr. 423 (1825). (Niemeysera.)										
A. stylidioides, G. Reichenbach in Benth, Fl. Austr. VI, 396 (1873)	—	—	—	—	—	—	—	Q.	—	B.fl. VI, 396

SCITAMINEAE.

Linné, philos. botan. 27 (1751).

MUSA, l'Ecluse, <i>exoticorum libri decem</i> 229 (1605).											
M. Banksii, F. v. M., fragm. IV, 132 (1864)	...	...	...	—	—	—	—	Q.	—	B.fl. VI, 261	M.fr. IV, 132.
M. Fitzalanei, F. v. M., fragm. IX, 188 (1875)	...	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 188.
M. Hillii, F. v. M., fragm. IX, 169 (1875)...	...	...	...	—	—	—	—	Q.	—	—	M.fr. IX, 169.
CURCUMA, Linné, <i>gen. plant.</i> 332 (1737), from Hermann (1687).											
C. Australasica, J. Hooker, <i>Bot. Mag.</i> t. 5620 (1867)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 263	M.fr. VIII, 26.
AMOMUM, Linné, <i>gen. plant.</i> 330 (1737).											
A. Dallachyi, F. v. M., fragm. VIII, 25 (1872)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 263	M.fr. VIII, 25.
ELETTARIA, White and Matton in <i>Transact. Linn. Soc.</i> X, 249 (1809), from Rheede (1692).											
E. Scottiana, F. v. M., fragm. VIII, 24 (1872)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 264	M.fr. VIII, 24.
ALPINIA, Linné, <i>gen. plant.</i> 332 (1737). (Hellenia.)											
A. racemigera, F. v. M., fragm. VIII, 27 (1872)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 365	M.fr. VIII, 27.
A. caerulea, Benthani, <i>Fl. Austr.</i> VI, 265 (1873)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 365	M.fr. VIII, 26.
A. arctiflora, F. v. M., fragm. VIII, 25 (1872)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 366	M.fr. VIII, 25.
COSTUS, C. Bauhin, <i>pinax</i> , 36 (1623).											
C. Potierae, F. v. M., <i>fragm. IX</i> , 164 (1864)	...	...	—	—	—	—	—	Q.	—	B.fl. VI, 296	M.fr. IV, 164; VIII, 27.
TAPEINOCHELOS, Miquel in <i>annal. mus. bot. Lugdun.</i> IV, 101, t. 4 (1869).											
T. pungens, Miquel in <i>ann. mus. Lugd.</i> IV, 101, t. 4 (1869)	...	—	—	—	—	—	—	Q.	—	B.fl. VI, 297	M.fr. VIII, 26.

IRIDEAE.

Ventenat, Tableau du règne végétal II, 188 (1799).

DIPLARRHENA, Labillardière, Relation du voy. à la rech. de La Per. I, 275, t. 15 (1799).									
D. Moraea, Labillardière, Voy. I, 157, t. 15 (1799)	...	...	—	—	T. V.	N.S.W.	—	—	B.f. VI, 400 M.fr. VII, 94.
PATERSONIA, R. Brown, in Bot. Magaz. t. 1041 (1807). (Genosiris, 1804.)									
P. glauca, R. Brown, prodr. 304 (1810)	...	...	...	...	S.A.	T. V.	N.S.W.	—	B.f. VI, 402 M.fr. VII, 36.
P. longiscapa, Sweet, Fl. Austral. t. 39 (1828)	...	...	...	...	S.A.	T. V.	—	—	B.f. VI, 402 M.fr. VII, 36.
P. occidentalis, R. Brown, prodr. 304 (1810)	...	...	...	...	W.A.	—	—	—	B.f. VI, 403 M.fr. VII, 31.
P. umbrosa, Endlicher in Lehm. pl. Preiss. II, 31 (1846)	...	...	...	...	W.A.	—	—	—	B.f. VI, 404 M.fr. VII, 32.
P. xanthina, F. v. M., fragm. I, 214 (1859)	...	...	...	...	W.A.	—	—	—	B.f. VI, 404 M.fr. I, 214.
P. limbata, Endlicher in Lehm. pl. Preiss. II, 29 (1846)...	...	...	...	...	W.A.	—	—	—	B.f. VI, 404
P. juncea, Lindley, Bot. Regist. XXV, App. LVIII (1839)	...	...	...	...	W.A.	—	—	—	B.f. VI, 404 M.fr. VII, 33.
P. Maxwellii, F. v. M. in Benth. fl. Austr. VI, 405 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 405 M.fr. VII, 34.
P. pygmaea, Lindley, Bot. Regist. XXV, App. LVIII (1839)	...	...	...	...	W.A.	—	—	—	B.f. VI, 405 M.fr. VII, 33.
P. sericea, R. Brown in Bot. Mag. t. 1041 (1807)...	...	...	...	...	—	V.	N.S.W.	Q.	B.f. VI, 406 M.fr. VII, 35.
P. lanata, R. Brown, prodr. 303 (1810)	...	...	...	...	W.A.	—	—	—	B.f. VI, 406 M.fr. VII, 35.
P. rudis, Endlicher in Lehm. pl. Preiss. II, 29 (1846)	...	...	...	...	W.A.	—	—	—	B.f. VI, 406 M.fr. VII, 35.
P. macrantha, Bentham, Fl. Austr. VI, 407 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 407
P. glabrata, R. Brown, prodr. 304 (1810)	...	...	...	...	—	—	V.	N.S.W.	Q.
P. Drummondii, F. v. M. in Benth. Fl. Austr. VI, 407 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 407 M.fr. VII, 35.
P. inaequalis, Bentham, Fl. Austr. VI, 408 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 408
P. graminea, Bentham, Fl. Austr. VI, 408 (1873)...	...	...	...	...	W.A.	—	—	—	B.f. VI, 408 M.fr. XI, 23.
P. babianoides, Bentham, Fl. Austr. VI, 408 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 408
IRIS, Tournefort, inst. 358, t. 186, 188 (1700), from Hippocrates, Dioscorides and Plinius. (Moraea partly.)									
I. Robinsoniana, F. v. M., fragm. VII, 153 (1870)	...	...	—	—	—	N.S.W.	—	—	B.f. VI, 409 M.fr. VII, 153.
SISYRINCHIUM, Linné, gen. plant. 273 (1737). (Libertia, Orthrosanthus, Renealmia, Nematostigma.)									
S. paniculatum, R. Brown, prodr. 305 (1810)	...	...	...	...	—	—	V.	N.S.W.	—
S. pulchellum, R. Brown, prodr. 305 (1810)	...	...	...	...	—	T. V.	N.S.W.	—	B.f. VI, 413 M.fr. VII, 91.
S. cyaneum, Lindley, Bot. Regist. t. 1090 (1827)...	...	...	...	...	W.A.	S.A.	—	—	B.f. VI, 413 M.fr. VII, 92.
S. polystachyum, F. v. M.; Orthrosanthus, Benth. Fl. Austr. VI, 411 (1873)	...	...	...	...	W.A.	—	—	—	B.f. VI, 410 M.fr. VII, 92.
	...	...	...	...	W.A.	—	—	—	B.f. VI, 411

CAMPYNEMA, Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804).
 C. lineare, Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804) — — T. — — — — B.f. VI, 415 M.fr. VI, 96.

BURMANNIACEAE.

Sprengel, syst. veg. I, 125 (1825).

BURMANNIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 100 (1737).
 B. disticha, Linné, spec. plant. 287 (1753)... — — — — N.S.W. Q. N.A. B.f. VI, 397
 B. juncea, Solander in R. Brown, prodr. 265 (1810) ... — — — — Q. N.A. B.f. VI, 397

TACCACEAE.

Presl, reliq. Haenk. I, 149 (1828).

TACCA, R. et G. Forster, char. gen. 65 (1776).
 T. pinnatifida, R. & G. Forster, char. gen. 65 (1776) ... — — — — Q. N.A. B.f. VI, 458

DIOSCORIDEAE.

Du Mortier, Analyse des familles des plantes 57 (1829), from R. Brown (1810).

DIOSCOREA, Plumier, nov. plant. Americ. gen. 9, t. 26 (1703). (Dioscoridea.)
 D. hastifolia, Nees in Lehm. pl. Preiss. II, 33 (1846) ... — — — — B.f. VI, 461 M.fr. VII, 80; XI, 20.
 D. transversa, R. Brown, prodr. 295 (1810) ... — — — — N.S.W. Q. N.A. B.f. VI, 460 M.fr. VII, 80.
 D. sativa, Linné, spec. plant. 1033 (1753) ... — — — — Q. N.A. B.f. VI, 461 M.fr. VII, 81.
 PETERMANNIA, F. v. M., fragm. II, 92 (1860).
 P. cirrosa, F. v. M., fragm. II, 93 (1860) ... — — — — N.S.W. — — B.f. VI, 462 M.fr. II, 176.

HYDROCHARIDEAE.

Lamarck and De Candolle, Flore Française, III, 265 (1805).

HYDROCHARIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 308 (1737).
 H. morsus ranae, Linné, spec. plant. 1036 (1753)... — — — — Q. — B.f. VI, 256 M.fr. VI, 199.
 ENHALUS, L. C. Richard in Mém. de l'Inst. II, 64 (1811).
 E. Koenigii, Richard in Mém. de l'Inst. II, 64 (1811) ... — — — — Q. — — M.fr. VIII, 219
 HALOPHILA, Du Petit-Thouars, gen. nov. Madag. n. 6 (1806).
 H. ovata, Gaudichaud, Bot. Voy. Freyc. 470, t. 40 (1826) ... — S.A. T. V. N.S.W. Q. — B.f. VII, 182 M.fr. VIII, 219.
 H. spinulosa, Ascherson in Neumayer, Anleit. zu wiss. Beob. 368 (1875) — — — — Q. — B.f. VII, 183 M.fr. VIII, 219.
 OTTELLA, Persoon, synops. plant. I, 400 (1805). (Damasonium partly.)
 O. alismoides, Persoon, synops. plant. I, 400 (1805) ... — — — — Q. N.A. B.f. VI, 257 M.fr. VI, 199.
 O. ovalifolia, L. C. Richard in Mém. de l'Inst. II, 78 (1811) ... W.A. S.A. — V. N.S.W. Q. N.A. B.f. VI, 257
 BLYXA, Noronha in Du Petit-Thouars, gen. nov. Madag. n. 14 (1806).
 B. Roxburghii, L. C. Richard, Mém. de l'Inst. II, 77, t. 5 (1811) — S.A. — — — Q. N.A. B.f. VI, 258
 VALLISNERIA, Micheli, nov. plant. gen. 12, t. 10 (1729).
 V. spiralis, Linné, spec. plant. 1015 (1753)... — — — — S.A. T. V. N.S.W. Q. N.A. B.f. VI, 259
 HYDRILLA, L. C. Richard in Mém. de l'Inst. II, 61 (1811).
 H. verticillata, Caspary in Bot. Zeitung, Dec. (1856) ... — — — — S.A. — V. N.S.W. Q. N.A. B.f. VI, 259

AMARYLLIDEAE.

J. St. Hilaire, Expos. des famill. natur. des pl. I, 134, t. 21 (1805).

HAEMODORUM, Smith in Transact. Linn. Soc. 213 (1798).
 H. distichophyllum, Hooker, icon. plant. t. 866 (1852) ... — — — — T. — — — — B.f. VI, 419 M.fr. VII, 118; IX, 51.
 H. brevicaulis, F. v. M., fragm. I, 64 (1858) ... — — — — — N.A. B.f. VI, 419 M.fr. VII, 118.
 H. sparsiflorum, F. v. M., fragm. VII, 117 (1870) ... — — — — — B.f. VI, 420 M.fr. VII, 117.
 H. spicatum, R. Brown, prodr. 300 (1810) ... — — — — — B.f. VI, 420 M.fr. VII, 118.
 H. brevipesaleum, Bentham, Fl. Austr. VI, 420 (1873) ... — — — — — B.f. VI, 420
 H. paniculatum, Lindley, Bot. Regist. XXV, App. XLIV (1839) W.A. — — — — — B.f. VI, 420
 H. laxum, R. Brown, prodr. 300 (1810) ... — — — — — B.f. VI, 421 M.fr. VII, 118.
 H. simplex, Lindley, Bot. Regist. XXV, App. XLIV (1839) ... W.A. — — — — — B.f. VI, 421
 H. simulans, F. v. M., fragm. VII, 117 (1870) ... — — — — — B.f. VI, 421 M.fr. VII, 117.
 H. planifolium, R. Brown, prodr. 300 (1810) ... — — — — — N.S.W. Q. — B.f. VI, 422 M.fr. VII, 118.
 H. teretifolium, R. Brown, prodr. 300 (1810) ... — — — — — N.S.W. — — B.f. VI, 422 M.fr. VII, 118.
 H. coccineum, R. Brown, prodr. 300 (1810) ... — — — — — Q. N.A. B.f. VI, 422
 H. subvirens, F. v. M., fragm. I, 63 (1858) ... — — — — — N.A. B.f. VI, 423 M.fr. I, 63.
 H. ensifolium, F. v. M., fragm. I, 64 (1858) ... — — — — — Q. N.A. B.f. VI, 423 M.fr. VII, 118.
 H. parviflorum, Bentham, Fl. Austr. VI, 423 (1873) ... — — — — — N.A. B.f. VI, 423
 H. leptostachyum, Bentham, Fl. Austr. VI, 423 (1873) ... — — — — — N.A. B.f. VI, 423
 H. tenuifolium, Cunningham in Benth. Fl. Austr. VI, 425 (1873) — — — — — Q. — B.f. VI, 423
 PHLEBOCARYA, R. Brown, prodr. 301 (1810).
 P. ciliata, R. Brown, prodr. 301 (1810) ... — — — — — W.A. — — — — — B.f. VI, 424 M.fr. VIII, 23.
 P. pilosissima, F. v. M. in Benth. fl. Austr. VI, 425 (1873) ... — — — — — W.A. — — — — — B.f. VI, 425 M.fr. VIII, 23.
 P. filifolia, F. v. M. in Benth. fl. Austr. VI, 425 (1873) ... — — — — — W.A. — — — — — B.f. VI, 425 M.fr. VIII, 23.

TRIBONANTHES, Endlicher, nov. stirp. dec. I, 27 (1839).

T. brachypetala, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,426	M.fr.VIII,23.
T. uniflora, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.
T. australis, Endlicher, nov. stirp. dec. I, 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.
T. variabilis, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	
T. longipetala, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.

CONOSTYLIS, R. Brown, prodr. 300 (1810). (Blancoa, Androstemma.)

C. breviscapa, R. Brown, prodr. 301 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,430	
C. vaginata, Endlicher in Lehm. pl. Preiss. II, 23 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,431	M.fr.VIII,19.
C. petrophiloides, F. v. M. in Benth. Fl. Austr. VI, 431 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,431	
C. setosa, Lindley, Bot. Regist. XXV, App. XLIV, t. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,431	M.fr.VIII,19.
C. aurea, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,19.
C. melanopogon, Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. setigera, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. Psyllium, Endlicher in Lehm. pl. Preiss. II, 21 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. villosa, Benth. Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. Drummondii, Benth. Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. involucreta, Endlicher in Lehm. pl. Preiss. II, 23 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. gladiata, Benth. Fl. Austr. VI, 434 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. seorsiflora, F. v. M., fragm. I, 158 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,19.
C. teretiuscula, F. v. M., fragm. VIII, 18 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,18.
C. stylioides, F. v. M., fragm. VIII, 17 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,17.
C. prolifera, Benth. Fl. Austr. VI, 436 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. racemosa, Benth. Fl. Austr. VI, 436 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. candicans, Endlicher, nov. stirp. dec. 20 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,436	M.fr.VI,255.
C. dealbata, Lindley, Bot. Regist. XXV, App. XLV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. Preissii, Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. bracteata, Endlicher in Lehm. pl. Preiss. II, 16 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. filifolia, F. v. M., fragm. VIII, 18 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,18.
C. spinuligera, F. v. M. in Benth. Fl. Austr. VI, 438 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. bromelioides, Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. aculeata, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,19.
C. laxiflora, Benth. Fl. Austr. VI, 439 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. cymosa, F. v. M. in Benth. Fl. Austr. VI, 439 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. serrulata, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,439	M.fr.VIII,19.
C. caricina, Lindley, Bot. Regist. XXV, App. XLV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,20.
C. Androstemma, F. v. M., fragm. VIII, 19 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,19.
C. Bealiana, F. v. M., fragm. IX, 50 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,50.
C. canescens, F. v. M., fragm. VIII, 19 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,441	M.fr.VIII,19.

AGONIZANTHOS, Labillardière, Relat. du voy. à la rech. de La Perouse I, 411 (1798). (Anigosanthus, Macropodia, Macropodia, Schwaegrichenia)

A. rufa, Labillardière, Voy. I, 411, t. 22 (1798)	W.A.	—	—	—	—	—	—	B.fl.VI,442	M.fr.VIII,21.
A. pulcherrima, Hooker, Bot. Mag. t. 4180 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. flavida, Redouté et De Candolle, les Liliacées t. 176 (1807)	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. Preissii, Endlicher in Lehm. pl. Preiss. II, 26 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,23.
A. humilis, Lindley, Bot. Regist. XXV, App. XLVI (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,21.
A. viridis, Endlicher in Lehm. pl. Preiss. II, 25 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. Manglesii, D. Don in Sweet, Brit. Fl. Gard. sec. ser. t. 265 (1836)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. bicolor, Endlicher in Lehm. pl. Preiss. II, 26 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,446	M.fr.VIII,22.
A. fuliginosus, Hooker, Bot. Mag. t. 4291 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,447	M.fr.VIII,20.

CURCULIGO, Gaertner, de fruct. I, 63, t. 16 (1788).

C. recurvata, Aiton, hort. Kew. sec. ed. II, 253 (1811)	—	—	—	—	—	Q.	—	B.fl.VI,448	M.fr.VIII,27.
C. ensifolia, R. Brown, prodr. 290 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,448	M.fr.VI,96.

HYPOXIS, Linné, syst. ed. decim. 986 (1759).

H. hygrometrica, Labillardière, Nov. Holl. pl. spec. I, 82, t. 108 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,449	M.fr.VI,96.
H. glabella, R. Brown, prodr. 289 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,450	M.fr.VI,96.
H. occidentalis, Benth. Fl. Austr. VI, 451 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,451	
H. marginata, R. Brown, prodr. 289 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,451	

DORYANTHES, Correa in Transact. Linn. Soc. VI, 211 (1802).

D. excelso, Correa in Transact. Linn. Soc. VI, 211, t. 23 (1802)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,452	M.fr.IX,51.
D. Palmeri, W. Hill in Benth. Fl. Austr. VI, 453 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,452	M.fr.IX,51.

CRINUM, Linné, gen. plant. 97 (1737).

C. venosum, R. Brown, prodr. 297 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,453	M.fr.III,22.
C. Asiaticum, Linné, spec. plant. 292 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VI,454	M.fr.III,23.
C. brachyandrum, Herbert in Bot. Mag. t. 2121 (1820)	—	—	—	—	—	Q.	—	—	M.fr.XI,139.
C. angustifolium, R. Brown, prodr. 297 (1810)	—	—	—	—	—	Q.	—	—	M.fr.XI,139.
C. flaccidum, Herbert in Bot. Mag. t. 2133 (1820)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,454	M.fr.III,23.
C. uniflorum, F. v. M., fragm. III, 23 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.VI,454	M.fr.VI,251.
C. pedunculatum, R. Brown, prodr. 297 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,455	M.fr.III,23.

EURYCYLES, Salisbury in Transact. of the Hort. Soc. I, 337 (1812).

E. silvestris, Salisbury in Transact. Hort. Soc. I, 337 (1812)	—	—	—	—	—	Q.	—	B.fl.VI,456	
E. Cunninghamii, Aiton in Loud. Hort. Brit. Suppl. 588 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,456	

CALOSTEMMA, R. Brown, prodr. fl. Nov. Holl. 297 (1810).

C. album, R. Brown, prodr. 298 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,458	
C. purpureum, R. Brown, prodr. 298 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,457	
C. luteum, Sims, Bot. Mag. t. 2101 (1819)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VI,457	

CALYCEAE HYPOGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 44 (1880).

ROXBURGHIAEAE.

Lindley in Wallich, pl. Asiat. rar. III, 50 (1832).

ROXBURGHIA, Jones in Roxburgh, pl. Corom. I, 29, t. 32 (1795). (Stemona 1790.)

R. Javanica, Kunth, enum. V, 288 (1850) ... — — — — — Q. N.A. B.fl.VII,1 M.fr.VII,81

LILIACEAE.

Haller, enum. stirp. Helv. I, 279 (1742).

PHORMIUM, R. & G. Forster, charact. gener. 47, t. 24 (1776). (Chlamydia.)

P. tenax, R. & G. Forster, char. gen. 47, t. 24 (1776) ... — — — — — N.S.W. — — —

SMILAX, Tournefort, inst. 654, t. 421 (1700), from Theophrastos and Dioscorides.

S. glycyphylla, Smith in White, Voy. to N.S. Wales 230 (1790) ... — — — — — N.S.W. Q. — B.fl.VII,7 M.fr.VII,77.

S. australis, R. Brown, prodr. 293 (1810) ... — — — — — V. N.S.W. Q. N.A. B.fl.VII,7 M.fr.VII,78.

RHIPOGONUM, R. & G. Forster, char. gen. 49, t. 25 (1776). (Ripogonum.)

R. album, R. Brown, prodr. 293 (1810) ... — — — — — V. N.S.W. Q. — B.fl.VII,9 M.fr.VII,79.

R. discolor, F. v. M., fragm. VII, 78 (1870) ... — — — — — N.S.W. Q. — B.fl.VII,9 M.fr.VII,78.

R. Fawcettianum, F. v. M. in Benth. Fl. Austr. VII, 9 (1878) ... — — — — — N.S.W. Q. — B.fl.VII,9 M.fr.VII,80.

R. Elseyanum, F. v. M., fragm. I, 44 (1858) ... — — — — — N.S.W. Q. — B.fl.VII,10 M.fr.VII,80.

FLAGELLARIA, Linné, amoen. acad. 396 (1747).

F. Indica, Linné, spec. plant. 333 (1753) ... — — — — — N.S.W. Q. N.A. B.fl.VII,10

DRYMOPHILA, R. Brown, prodr. fl. Nov. Holl. 292 (1810).

D. cyanocarpa, R. Brown, prodr. 292 (1810) ... — — — — — T. V. N.S.W. — — — B.fl.VII,12 M.fr.VII,73;IX,191.

D. Moorei, Baker in Journ. Linn. Soc. XIV, 571 (1875) ... — — — — — N.S.W. — — — B.fl.VII,13 M.fr.IX,190.

DIANELLA, Lamarek, Encyclop. méthod. II, t. 250 (1786).

D. Tasmanica, J. Hooker, fl. Tasman. II, 57, t. 133 (1860) ... — — — — — T. V. N.S.W. — — — B.fl.VII,14 M.fr.VI,121.

D. longifolia, R. Brown, prodr. 280 (1810) ... — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,14 M.fr.VI,122.

D. revoluta, R. Brown, prodr. 280 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,15 M.fr.VI,121;XI,20.

D. caerulea, Sims, Bot. Magaz. t. 505 (1799) ... — — — — — N.S.W. Q. — B.fl.VII,16 M.fr.VI,123.

D. ensifolia, De Candolle & Redouté, Liliac. t. 1 (1802) ... — — — — — Q. N.A. B.fl.VII,16 M.fr.VI,123.

ASPARAGUS, Tournefort, inst. 300, t. 154 (1700), from Theophrastos, Dioscorides and Plinius. (Asparagopsis.)

A. racemosus, Willdenow, spec. plant. II, 152 (1799) ... — — — — — Q. N.A. B.fl.VII,17 M.fr.VII,73.

EUSTREPHUS, R. Brown, prodr. 281 (1810).

E. Brownii, F. v. M., fragm. VII, 73 (1870) ... — — — — — V. N.S.W. Q. — B.fl.VII,18 M.fr.VII,73.

GEITONOPLESIMUM, Cunningham in Bot. Magaz. t. 3131 (1832). (Luzuriago partly.)

G. cymosum, Cunningham in Bot. Mag. t. 3131 (1832) ... — — — — — V. N.S.W. Q. — B.fl.VII,19 M.fr.VII,74.

DRACAENA, Vandelli, dissertatio de Dracaena t. 1 (1762).

D. angustifolia, Roxburgh, fl. Indic. II, 155 (1832) ... — — — — — Q. N.A. B.fl.VII,20 M.fr.V,194;VI,120;IX,190.

CORDYLINE, Commerçon in R. Hedwig, gen. plant. 243 (1806). (Charlwoodia.)

C. terminalis, Kunth in act. acad. Berol. 30 (1820) ... — — — — — N.S.W. Q. — B.fl.VII,21 M.fr.V,196.

C. Murchisoniae, F. v. M., fragm. V, 195 (1866) ... — — — — — Q. — B.fl.VII,22 M.fr.V,195.

C. stricta, Endlicher in Ann. Wien. Mus. I, 162 (1838) ... — — — — — N.S.W. Q. — B.fl.VII,22 M.fr.V,195;VI,121.

C. Baueri, J. Hooker in Gardeners' Chronicle 792 (1860) ... — — — — — N.S.W. — — —

BLANDFORDIA, Smith, Exot. Bot. I, 5, t. 4 (1804).

B. marginata, Herbert in Bot. Regist. Misc. 84 (1842) ... — — — — — T. — — — B.fl.VII,23 M.fr.VII,71.

B. grandiflora, R. Brown, prodr. 296 (1810) ... — — — — — N.S.W. — — — B.fl.VII,23

B. nobilis, Smith, Exot. Bot. I, 5, t. 4 (1804) ... — — — — — N.S.W. — — — B.fl.VII,24 M.fr.VII,70.

B. flammula, Hooker, Bot. Mag. t. 4819 (1854) ... — — — — — N.S.W. Q. — B.fl.VII,24

HEWARDIA, Hooker, icon. plant. 858 (1852).

H. Tasmanica, Hooker, icon. plant. t. 858 (1852) ... — — — — — T. — — — B.fl.VII,25 M.fr.VII,72.

ASTELIA, Banks & Solander in R. Brown, prodr. 291 (1810). (Hamelinia.)

A. alpina, R. Brown, prodr. 291 (1810) ... — — — — — T. V. N.S.W. — — — B.fl.VII,11

MILLIGANIA, J. Hooker in Kew misc. V, 296, t. IX (1853).

M. densiflora, J. Hooker, Fl. Tasman. II, 62 (1860) ... — — — — — T. — — — B.fl.VII,26

M. longifolia, J. Hooker in Kew misc. V, 296, t. IX (1853) ... — — — — — T. — — — B.fl.VII,26

M. Johnstoni, F. v. M. in Benth. Fl. Austr. VII, 26 (1878) ... — — — — — T. — — — B.fl.VII,26

M. stylosa, F. v. M. in papers of the R. S. of Tasm. 11 (1876) ... — — — — — T. — — — B.fl.VII,27

WURMBEA, Thunberg, nov. gen. I, 18 (1781). (Anguillaria, Melanthium partly, Pleea partly.)

W. tubulosa, Bentham, Fl. Austr. VII, 28 (1878) ... — — — — — W.A. — — — B.fl.VII,28 M.fr.VII,75.

W. dioica, F. v. M., fragm. X, 119 (1877) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,30 M.fr.VII,74.

IPHIGENIA, Kunth, enum. plant. omn. hucusq. cognit. IV, 212 (1843).

I. Indica, Kunth, enum. plant. IV, 213 (1843) ... — — — — — Q. N.A. B.fl.VII,31 M.fr.VII,74;IX,193.

SCHELHAMMERA, R. Brown, prodr. 273 (1810).

S. undulata, R. Brown, prodr. 274 (1810) ... — — — — — V. N.S.W. — — — B.fl.VII,31 M.fr.VII,71.

S. multiflora, R. Brown, prodr. 274 (1810) ... — — — — — Q. — — — B.fl.VII,32 M.fr.VII,72.

KREYSSIGIA, Reichenbach, iconogr. bot. exotic. III, 11, t. 229 (1829). (Kreysigia, Tripladenia.)

K. Cunninghami, F. v. M., fragm. VII, 71 (1870) ... — — — — — N.S.W. Q. — B.fl.VII,32 M.fr.VII,71.

BURCHARDIA, R. Brown, prodr. 272 (1810).

B. umbellata, R. Brown, prodr. 272 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,33 M.fr.VII,77.

BULBINE, Linné, hort. Cliffort. 122 (1737). (Anthericum partly, Chrysobactron.)											
B. bulbosa, Haworth, revis. pl. succul. 33 (1821) ...	...	...	...	...	...	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 34 M.fr. VII, 70; XI, 27.
B. semibarbata, Haworth, rev. pl. succ. 33 (1821) ...	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 35 M.fr. VII, 70.
AGROSTOCRINUM, F. v. M., fragm. phytogr. Austr. II, 95 (1860).											
A. stypandroides, F. v. M., fragm. II, 95 (1860) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 36 M.fr. VII, 65.
THYSANOTUS, R. Brown, prodr. 282 (1810). (Chlamyspermum.)											
T. multiflorus, R. Brown, prodr. 285 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 38 M.fr. VII, 69.
T. triandrus, R. Brown, prodr. 284 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 39 M.fr. VII, 69.
T. glaucus, Endlicher in Lehm. pl. Preiss. II, 38 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 39
T. Drummondii, Baker in Journ. Linn. Soc. XV, 341 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 39
T. pauciflorus, R. Brown, prodr. 285 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 39
T. asper, Lindley, Bot. Regist. XXV, App. LVIII (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 39
T. chrysantherus, F. v. M. in Bentham, flor. Hongk. 372 (1861) ...	...	...	...	...	...	—	—	—	—	N.A.	B.f. VII, 40 M.fr. V, 202.
T. exiliflorus, F. v. M. in transact. R. S. of S. Austr. IV, 112 (1881) ...	...	...	...	...	...	S.A.	—	—	—	—	M.fr. XI, 139.
T. isantherus, R. Brown, prodr. 283 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 40
T. tenellus, Endlicher in Lehm. pl. Preiss. II, 37 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 40
T. scaber, Endlicher in Lehm. pl. Preiss. II, 37 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 41
T. tuberosus, R. Brown, prodr. 282 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q. N.A. B.f. VII, 41 M.fr. VII, 69.
T. exasperatus, F. v. M., fragm. I, 21 (1858) ...	...	...	...	...	...	—	S.A.	—	—	—	B.f. VII, 42 M.fr. I, 21.
T. thyrsoides, Baker in Journ. Linn. Soc. XV, 336 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 42
T. Baueri, R. Brown, prodr. 283 (1810) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	— B.f. VII, 42 M.fr. I, 22; VII, 69.
T. Patersoni, R. Brown, prodr. 284 (1810) ...	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W.	—	B.f. VII, 43 M.fr. VII, 69; XI, 20.
T. junceus, R. Brown, prodr. 283 (1810) ...	...	...	...	...	...	—	—	—	N.S.W.	—	B.f. VII, 43 M.fr. VII, 69.
T. dichotomus, R. Brown, prodr. 284 (1810) ...	...	...	...	...	...	W.A.	S.A.	—	—	—	B.f. VII, 44 M.fr. VII, 69.
T. arbuscula, Baker in Journ. Linn. Soc. XV, 339 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 45
T. anceps, Lindley, Bot. Regist. XXV, App. LVIII (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 45
HODGSONIOLA, F. v. M., fragm. II, 176 (1861). (Hodgsonia.)											
H. junciformis, F. v. M., fragm. II, 176 (1861) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 46 M.fr. II, 95, 176.
CAESIA, R. Brown, prodr. fl. Nov. Holl. 277 (1810).											
C. vittata, R. Brown, prodr. 277 (1810) ...	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 47 M.fr. I, 63; VII, 68.
C. chlorantha, F. v. M., fragm. I, 63 (1858) ...	...	...	...	...	...	—	—	—	—	Q. N.A.	— M.fr. I, 63.
C. parviflora, R. Brown, prodr. 277 (1810) ...	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 47 M.fr. VII, 67.
C. rigidifolia, F. v. M., fragm. X, 48 (1876) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 47 M.fr. X, 48.
C. setifera, Baker in Journ. Linn. Soc. XV, 359 (1876) ...	...	...	...	...	...	—	—	—	—	N.A.	B.f. VII, 47
CHAMAESCILLA, F. v. M., fragm. VII, 68 (1870). (Caesia partly.)											
C. corymbosa, F. v. M., fragm. VII, 68 (1870) ...	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W.	—	B.f. VII, 48 M.fr. VII, 68.
C. spiralis, F. v. M., fragm. VII, 68 (1870) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 48 M.fr. VII, 68.
CORYNOTHECA, F. v. M., fragm. VII, 68 (1870). (Caesia partly.)											
C. lateriflora, F. v. M., fragm. VII, 68 (1870) ...	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q. N.A. B.f. VII, 49 M.fr. VII, 68.
C. dichotoma, F. v. M., fragm. VII, 68 (1870) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 50 M.fr. I, 215; XI, 20.
C. acanthoclada, F. v. M., fragm. VII, 68 (1870) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 50 M.fr. I, 215; XI, 20.
TRICHORYNE, R. Brown, prodr. fl. Nov. Holl. 278 (1810).											
T. platyptera, G. Reichenbach, Beitr. zur syst. Pflanzenk. 72 (1871) ...	...	...	...	...	...	—	—	—	—	Q.	— B.f. VII, 51
T. anceps, R. Brown, prodr. 278 (1810) ...	...	...	...	...	...	—	—	—	—	Q.	— B.f. VII, 51 M.fr. VII, 67.
T. muricata, Baker in Journ. Linn. Soc. XV, 363 (1876) ...	...	...	...	...	...	—	—	—	—	Q.	— B.f. VII, 52
T. elatior, R. Brown, prodr. 278 (1810) ...	...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 52 M.fr. VII, 67; XI, 20.
T. simplex, R. Brown, prodr. 278 (1810) ...	...	...	...	...	...	—	—	—	N.S.W.	Q.	— B.f. VII, 52 M.fr. VII, 67.
T. humilis, Endlicher in Lehm. pl. Preiss. II, 36 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 53 M.fr. VII, 67.
STYPANDRA, R. Brown, prodr. flor. Nov. Holl. 278 (1810).											
S. glauca, R. Brown, prodr. 278 (1810) ...	...	...	...	...	...	W.A.	—	—	V.	N.S.W.	Q. — B.f. VII, 53 M.fr. VII, 64.
S. caespitosa, R. Brown, prodr. 279 (1810) ...	...	...	...	...	...	—	—	T. V.	N.S.W.	Q.	— B.f. VII, 54 M.fr. VII, 65.
ARTHROPODIUM, R. Brown, prodr. 276 (1810). (Dichopogon.)											
A. paniculatum, R. Brown, prodr. 276 (1810) ...	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 56 M.fr. VII, 66.
A. minus, R. Brown, prodr. 276 (1810) ...	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	—	B.f. VII, 56 M.fr. VII, 67; XI, 27.
A. capillipes, Endlicher in Lehm. pl. Preiss. II, 34 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 57
A. Preissii, Endlicher in Lehm. pl. Preiss. II, 35 (1846) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 57 M.fr. VII, 66.
A. dianellaceum, F. v. M., fragm. X, 65 (1876) ...	...	...	...	...	...	—	—	—	—	Q.	— B.f. VII, 57 M.fr. X, 65.
A. strictum, R. Brown, prodr. 276 (1810) ...	...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	— B.f. VII, 58 M.fr. VII, 66.
A. laxum, Sieber in Roemer & Schultes, syst. VII, 441 (1830) ...	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	— B.f. VII, 59
CHLOROPHYTUM, Ker in Bot. Mag. t. 1071 (1808). (Chlorophyton, Phalangium partly.)											
C. laxum, R. Brown, prodr. 277 (1810) ...	...	...	...	...	...	—	—	—	—	N.A.	B.f. VII, 60 M.fr. I, 63; VII, 71.
C. alpinum, Baker in Journ. Linn. Soc. XV, 329 (1876) ...	...	...	...	...	...	—	—	T.	—	—	— B.f. VII, 60
HERPOLIRION, J. Hooker, Fl. of New Zeal. I, 258 (1853).											
H. Novae Zelandiae, J. Hooker, Fl. of New Zeal. I, 258 (1853) ...	...	...	...	...	...	—	—	T. V.	N.S.W.	—	— B.f. VII, 61 M.fr. VII, 68.
SOWERBAEA, Smith in Transact. Linn. Soc. V, 159 (1800). (Sowerbya.)											
S. juncea, Smith in Transact. Linn. Soc. V, 160, t. 6 (1800) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q. — B.f. VII, 61 M.fr. VI, 180.
S. laxiflora, Lindley, Botan. Regist. t. 10 (1841) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 62 M.fr. VI, 180; VII, 87.
S. alliacea, F. v. M., fragm. VI, 180 (1868) ...	...	...	...	...	...	—	—	—	—	N.A.	B.f. VII, 62 M.fr. VII, 87.
ALLANIA, Endlicher, gen. plant. 151 (1837). (Alania.)											
A. Endlicheri, Kunth, enum. IV, 644 (1843) ...	...	...	...	...	...	—	—	—	N.S.W.	—	— B.f. VII, 63 M.fr. VI, 180.
BARTLINGIA, F. v. M., fragm. VII, 88 (1870). (Laxmannia.)											
B. grandiflora, F. v. M.; Laxmannia, Lindl. Bot. Reg. XXV, App. LVI (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 64 M.fr. I, 159; VII, 88.
B. squarrosa, F. v. M.; Laxm., Lindl. Bot. Reg. XXV, App. LVI (1839) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 64
B. minor, F. v. M.; Laxmannia, R. Brown, prodr. 286 (1810) ...	...	...	...	...	...	W.A.	—	—	—	—	B.f. VII, 65 M.fr. VII, 88, 89.

B. gracilis, F. v. M., fragm. VII, 88 (1870)	—	—	—	V.	N.S.W.	Q.	—	B.f. VII, 65	M.fr. I, 159.
B. ramosa, F. v. M., fragm. VII, 88 (1870)	W.A.	—	—	—	—	—	—	B.f. VII, 66	M.fr. VI, 255.
B. sessiliflora, F. v. M. in pap. R. S. of Tasm. 116 (1877) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 66	—
B. brachyphylla; Laxm., F. v. M. in Benth. Fl. VII, 67 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 66	M.fr. I, 158.
B. sessilis, F.v.M.; Laxm., Lindl. Bot. Reg. XXV, App. LVI (1839)	W.A.	—	—	—	—	—	—	B.f. VII, 67	—
STAWELLIA, F. v. M., fragm. phytogr. Austr. VII, 85 (1870).									
S. dimorphantha, F. v. M., fragm. VII, 85 (1870)	W.A.	—	—	—	—	—	—	B.f. VII, 67	M.fr. VII, 85.
JOHNSONIA, R. Brown, prodr. fl. Nov. Holl. 287 (1810).									
J. lupulina, R. Brown, prodr. 287 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 68	M.fr. VII, 86.
J. pubescens, Lindley, Bot. Regist. XXV, App. LVII (1839) ...	W.A.	—	—	—	—	—	—	B.f. VII, 68	M.fr. VI, 255.
ARNOCRINUM, Endlicher & Lehmann in pl. Preiss. II, 41 (1846).									
A. Drummondii, Endlicher in Lehm. pl. Preiss. II, 41 (1846) ...	W.A.	—	—	—	—	—	—	B.f. VII, 69	M.fr. VII, 73.
A. Preissii, Lehmann, pl. Preiss. II, 42 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 70	—
BORYA, Labillardiere, Nov. Holl. pl. specim. 81, t. 107 (1804).									
B. nitida, Labillardiere, Nov. Holl. pl. spec. 81, t. 107 (1804) ...	W.A.	—	—	—	—	—	—	B.f. VII, 71	M.fr. VII, 87.
B. septentrionalis, F. v. M., fragm. 41 (1856)	—	—	—	—	—	Q.	—	B.f. VII, 71	M.fr. V, 41.
CALECTASIA, R. Brown, prodr. fl. Nov. Holl. 263 (1810).									
C. cyanea, R. Brown, prodr. 264 (1810)	W.A.	S.A.	—	V.	—	—	—	B.f. VII, 121	—
BAXTERIA, R. Brown in Hook. Lond. Journ. of Bot. II, 492, t. 13-15 (1843).									
B. australis, R. Brown in Hook. Lond. Journ. II, 492 (1843) ...	W.A.	—	—	—	—	—	—	B.f. VII, 120	—
ACANTHOCARPUS, Lehmann, pl. Preiss. II, 274 (1847). (Chamaexeros).									
A. Preissii, Lehmann, pl. Preiss. II, 274 (1847)	W.A.	—	—	—	—	—	—	B.f. VII, 111	—
A. Serra, F. v. M.; Xerotes, Endlicher in pl. Preiss. II, 49 (1847)	W.A.	—	—	—	—	—	—	B.f. VII, 111	M.fr. VIII, 211.
A. fimbriatus; Xerotes, F. v. M., fragm. VIII, 211 (1874) ...	W.A.	—	—	—	—	—	—	B.f. VII, 111	M.fr. VIII, 211.
XEROTES, R. Brown, prodr. 257 (1810). (Lomandra, 1804.)									
X. Banksii, R. Brown, prodr. 263 (1810)	—	—	—	—	—	Q.	—	B.f. VII, 96	M.fr. VIII, 213.
X. dura, F. v. M. in Transact. Vict. Inst. 42 (1854)	—	S.A.	—	—	—	—	—	B.f. VII, 97	M.fr. VIII, 207.
X. longifolia, R. Brown, prodr. 262 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 98	M.fr. VIII, 210.
X. rigida, R. Brown, prodr. 262 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 99	M.fr. VIII, 209.
X. Sonderi, F. v. M., fragm. VIII, 206 (1874)	W.A.	—	—	—	—	—	—	B.f. VII, 99	M.fr.
X. odora, Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 99	—
X. Brownii, F. v. M., fragm. VIII, 206 (1874)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 100	M.fr. VIII, 206.
X. Ordii, F. v. M., fragm. XI, 23 (1878)	W.A.	—	—	—	—	—	—	—	M.fr. XI, 137.
X. sororia, F. v. M., first gen. Report 15 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 100	—
X. Endlicheri, F. v. M., fragm. VIII, 205 (1874)	W.A.	—	—	—	—	—	—	B.f. VII, 101	M.fr. VIII, 205.
X. sericea, Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 101	M.fr. VIII, 207.
X. purpurea, Endlicher in Lehm. pl. Preiss. II, 49 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 101	M.fr. VIII, 213.
X. Preissii, Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 102	—
X. effusa, Lindley in Mitch. Three Exped. II, 151 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. VII, 102	M.fr. VIII, 209.
X. micrantha, Endl. in Lehm. pl. Preiss. II, 49 (1846)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.f. VII, 103	M.fr. VIII, 212.
X. Thunbergii, F. v. M., fragm. VIII, 208 (1874)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 103	M.fr. VIII, 208.
X. caespitosa, Benth. Fl. Austr. VII, 104 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 104	—
X. pauciflora, R. Brown, prodr. 261 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 104	M.fr. 212.
X. flexifolia, R. Brown, prodr. 260 (1810)	—	—	—	—	N.S.W.	—	—	B.f. VII, 105	M.fr. VIII, 207.
X. glauca, R. Brown, prodr. 260 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 105	M.fr. VIII, 208.
X. elongata, Benth. Fl. Austr. VII, 106 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 106	—
X. rupestris, Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 106	—
X. collina, R. Brown, prodr. 260 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 107	—
X. suaveolens, Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 107	M.fr. VIII, 207.
X. turbinata, Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 107	M.fr. VIII, 211.
X. spartea, Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 108	M.fr. VIII, 211.
X. juncea, F. v. M. in Transact. Vict. Inst. 135 (1855)	—	S.A.	—	—	—	—	—	B.f. VII, 108	M.fr. VIII, 210.
X. leucocephala, R. Brown, prodr. 260 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 109	M.fr. VIII, 210.
X. hastilis, R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 110	M.fr. VIII, 210; XI, 54.
DASYPOGON, R. Brown, prodr. fl. Nov. Holl. 263 (1810).									
D. bromelifolius, R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 118	—
D. Hookeri, Drummond in Hook. Lond. Journ. II, 168 (1843) ...	W.A.	—	—	—	—	—	—	B.f. VII, 119	M.fr. II, 112; IV, 173.
XANTHORRHOEA, Smith in Transact. Linn. Soc. IV, 219 (1798).									
X. gracilis, Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 114	M.fr. IV, 112.
X. pumilio, R. Brown, prodr. 288 (1810)	—	—	—	—	—	Q.	—	B.f. VII, 114	—
X. macronema, F. v. M., fragm. IV, 112 (1864)	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 113	—
X. hastilis, R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 115	M.fr. IV, 113.
X. arborea, R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 115	M.fr. IV, 113.
X. semiplana, F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.f. VII, 115	M.fr. IV, 111.
X. bracteata, R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	—	—	B.f. VII, 116	—
X. minor, R. Brown, prodr. 288 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 114	M.fr. IV, 112.
X. australis, R. Brown, prodr. 288 (1810)	—	—	T.	V.	—	—	—	B.f. VII, 116	M.fr. IV, 110.
X. quadrangulata, F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.f. VII, 117	M.fr. IV, 111.
X. Preissii, Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 117	M.fr. IV, 110.
KINGIA, R. Brown in King's Narrat. of survey of Austr. coasts II, 529 (1827).									
K. australis, R. Brown in King's Narr. of surv. II, 535 (1827) ...	W.A.	—	—	—	—	—	—	B.f. VII, 119	—

PALMAE.

Ray, method. pl. emend. 135 (1703).

CALAMUS, Linné, spec. plant. 325 (1753).									
C. australis, Martius, hist. nat. palm. III, 213 (1839)	—	—	—	—	—	Q.	—	B.f. VII, 134	M.fr. V, 48.
C. Muellert, Wendland & Drude in Linnaea XXXIX, 193 (1871) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 135	—
C. caryotoides, A. Cunn. in Martius, hist. nat. palm. III, 212 (1839) ...	—	—	—	—	—	Q.	—	B.f. VII, 135	—

- BACULARIA**, F. v. M., fragm. VII, 103 (1870). (Linospadiæ).
- B. monostachya*, F. v. M., fragm. VII, 103 (1870) ... — — — — N.S.W. Q. — B.f. VII, 136 M.fr. VII, 82, 103; XI, 58.
- KENTIA**, Blume in Bull. Neerl. 64 (1838). (Grisebachia, Hydiarstele, Hedyscepe, Kentiopsis, Veitchia partly).
- K. minor*, F. v. M., fragm. VIII, 235 (1874) ... — — — — Q. — B.f. VII, 137 M.fr. IX, 195; XI, 58, 137.
- K. Wendlandiana*, F. v. M., fragm. VII, 102 (1870) ... — — — — Q. N.A. B.f. VII, 138 M.fr. VII, 102.
- K. Belmoreana*, C. Moore & F. v. M., fragm. VII, 99 (1870) ... — — — — N.S.W. — — B.f. VII, 137 M.fr. VIII, 234.
- K. Forsteriana*, C. Moore & F. v. M., fragm. VII, 100 (1870) ... — — — — N.S.W. — — — —
- K. Baueri*, Seemann, fl. Vitiens. 269 (1869) ... — — — — N.S.W. — — — — M.fr. VII, 102; VIII, 233.
- K. Canterburyana*, C. Moore & F. v. M., fragm. VII, 101 (1870) ... — — — — N.S.W. — — B.f. VII, 138 M.fr. VIII, 234; X, 119.
- K. acuminata*, Wendland & Drude in Linnaea XXXIX, 207 (1871) ... — — — — N.A. B.f. VII, 138
- CLINOSTIGMA**, Herm. Wendland in Seemann's Bonplandia X, 196 (1862). (Cyphokentia). [120.]
- C. Mooreanum*, F. v. M., fragm. VIII, 235 (1874) ... — — — — N.S.W. — — B.f. VII, 139 M.fr. VII, 101; IX, 195; X,
- PTYCHOSPERMA**, Labillardiere in Mém. de l'Inst., ann. 1808, I, 251, pl. X (1809). (Seaforthia, Laccospadix, Archontophoenix).
- P. Laccospadix*, Benth., fl. Austr. VII, 140 (1878) ... — — — — Q. — B.f. VII, 140
- P. Beatricae*, F. v. M. in Melb. Chemist. Febr. (1882) ... — — — — Q. — — — —
- P. Alexandrae*, F. v. M., fragm. V, 47, t. 43 (1865) ... — — — — Q. — B.f. VII, 140 M.fr. V, 47, 190, 213.
- P. Cunninghamii*, Herm. Wendland in Bot. Zeit. 346 (1858) ... — — — — N.S.W. — — B.f. VII, 141 M.fr. VIII, 222.
- P. elegans*, Blume, Rumphia II, 118 (1836) ... — — — — Q. — B.f. VII, 141 M.fr. VIII, 222.
- P. Normanbyi*, F. v. M., fragm. XI, 56 (1878) ... — — — — Q. — B.f. VII, 142 M.fr. VIII, 235.
- ARECA**, Linné, spec. plant. 1189 (1753) from Ray (1688).
- A. Alicaë*, F. v. M. in Regel's Gartenflora, 199-201 (1879) ... — — — — Q. — — — — M.fr. XI, 88.
- COCOS**, Linné, spec. plant. 1188 (1753).
- C. nucifera*, Linné, spec. plant. 1188 (1753) ... — — — — Q. — B.f. VII, 143
- CARYOTA**, Linné, gen. plant. 355 (1737).
- C. Rumphiana*, Martius, hist. nat. palm. III, 195 (1839) ... — — — — Q. — B.f. VII, 144
- LICUALA**, Rumphius, herb. Amboin. I, 44, t. 9 (1741).
- L. Muelleri*, Wendland & Drude in Linnaea XXXIX, 223 (1871) ... — — — — Q. — B.f. VII, 145 M.fr. VIII, 221; X, 120.
- LIVISTONA**, R. Brown, prodr. fl. Nov. Holl. 267 (1810). (Corypha partly).
- L. humilis*, R. Brown, prodr. 268 (1810) ... — — — — — N.A. B.f. VII, 146 M.fr. VIII, 221.
- L. inermis*, R. Brown, prodr. 268 (1810) ... — — — — — N.A. B.f. VII, 146
- L. Mariae*, F. v. M. in Giles's Geog. Trav. in Cntr. Austr. 222 (1875) ... — S.A. — — — — N.A. — — — —
- L. australis*, Martius, hist. nat. palm. III, 241 (1839) ... — — — — — N.A. — — — — M.fr. XI, 54.
- PANDANACEAE.**
- R. Brown, prodr. Nov. Holl. (1810)
- PANDANUS**, Rumphius, herb. Amboin. IV, 139, t. 74 (1750) (Athrodactylis, Fiquetia).
- P. odoratissimus*, Linné, fil. suppl. 424 (1781) ... — — — — — N.A. B.f. VII, 148 M.fr. XI, 55.
- P. pedunculatus*, R. Brown, prodr. 341 (1810) ... — — — — N.S.W. Q. — B.f. VII, 149 M.fr. VIII, 220; IX, 193.
- P. Forsteri*, Moore & F. v. M., fragm. VIII, 220 (1874) ... — — — — N.S.W. — — B.f. VII, 149 M.fr. VIII, 220.
- P. aquaticus*, F. v. M., fragm. V, 40 (1865) ... — — — — Q. N.A. B.f. VII, 148 M.fr. VIII, 220. [193]
- P. monticola*, F. v. M., fragm. V, 40 (1865) ... — — — — Q. — B.f. VII, 150 M.fr. VII, 63; VIII, 220; IX,
- FREYCINETIA**, Gaudichaud in Annal. des scienc. natur. III, 509 (1824).
- F. Gaudichaudii*, Brown & Bennett, pl. Jav. rar. 31, t. 9 (1838) ... — — — — Q. — B.f. VII, 151
- F. excelsa*, F. v. M., fragm. V, 39 (1865) ... — — — — Q. — B.f. VII, 151 M.fr. V, 39.
- F. Baueriana*, Endlicher, prodr. fl. ins. Norf. 25 (1833) ... — — — — N.S.W. — — — —
- NIPA**, Wurm in Verhandl. Batav. Genootsch. I, 349 (1779), from Camellus, Ray & Rumphius.
- N. fruticans*, Wurm in Verhandl. Batav. Genootsch. I, 349 (1779) ... — — — — Q. — — — — M.fr. XI, 128.
- AROIDEAE.**
- A. L. de Jussieu, gen. plant. 23 (1789), from B. de Jussieu (1759).
- TYPHONIUM**, Schott in Wien. Zeitschr. III, 72 (1829). (Arum partly.)
- T. liliifolium*, F. v. M. in Schott, prodr. syst. aroid. 107 (1860) ... — — — — — N.A. B.f. VII, 153 M.fr. VIII, 187.
- T. alismifolium*, F. v. M., fragm. VIII, 186 (1874) ... — — — — Q. — B.f. VII, 153 M.fr. VIII, 186.
- T. Brownii*, Schott, icon. aroid. II, t. 15 (1857) ... — — — — N.S.W. Q. — B.f. VII, 154 M.fr. VIII, 187.
- T. angustilobum*, F. v. M., fragm. X, 66 (1876) ... — — — — Q. — B.f. VII, 154 M.fr. X, 66.
- AMORPHOPHALLUS**, Blume in Nouv. Ann. du Mus. III, 366 (1834). (Brachyspatha).
- A. variabilis*, Blume, Rumphia I, 146, t. 35 (1835) ... — — — — — N.A. B.f. VII, 155 M.fr. VIII, 87.
- COLOCASIA**, Ray, Method. 157 (1682). (Caladium partly.)
- C. antiquorum*, Schott, Meletemat. botan. 18 (1832) ... — — — — Q. — B.f. VII, 155 M.fr. VIII, 87.
- C. macrorrhiza*, Schott, Meletemat. bot. 18 (1832) ... — — — — N.S.W. Q. — B.f. VII, 155 M.fr. VIII, 87.
- RHAPHIDOPHORA**, Hasskarl in Fl. Regensb. Beibl. II, 11 (1842). (Epipremnum partly.)
- R. pertusa*, Schott in Seem. Bonplandia V, 45 (1857) ... — — — — Q. — B.f. VII, 156 M.fr. VIII, 87.
- GYMNOSTACHYS**, R. Brown, prodr. fl. Nov. Holl. 337 (1810).
- G. anceps*, R. Brown, prodr. 337 (1810) ... — — — — N.S.W. Q. — B.f. VII, 157 M.fr. VIII, 87.
- POTHOS**, Linné, amcen. acad. I, 410 (1747).
- P. Loureiri*, Hooker & Arnott, Bot. of Beech. Voy. 220 (1841) ... — — — — N.S.W. Q. — B.f. VII, 158 M.fr. I, 62; VIII, 87.

TYPHACEAE.

A. L. de Jussieu, gener. plant. 25 (1789).

- TYPHA**, Tournefort, inst. 530, t. 301 (1700), from Theophrastus, Dioscorides and Plinius.
- T. angustifolia*, Linné, spec. plant. 971 (1753) ... — — — — W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 159 M.fr. VII, 116.
- SPARGANIUM**, Tournefort, inst. 530, t. 302 (1700).
- S. angustifolium*, R. Brown, prodr. 338 (1810) ... — — — — V. N.S.W. Q. — B.f. VII, 160 M.fr. VII, 117.

LEMNACEAE.

J. E. Gray, Nat. arrang. of Brit. pl. II, 729 (1821).

- LEMNA, Linné, syst. nat. 9 (1735); Linné, Fl. Lapon. 351 (1737). (Telmatophace, Spirodela.)
 L. trisulca, Linné, spec. plant. 970 (1753) ... — S.A. T. V. N.S.W. Q. — B.f. VII, 162 M.fr. VIII, 188.
 L. minor, Linné, spec. plant. 970 (1753) ... — W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 163 M.fr. VIII, 188.
 L. oligorrhiza, Kurz in Journ. Linn. Soc. IX, 287 (1867) ... — S.A. — V. N.S.W. Q. N.A. B.f. VII, 163 M.fr. VIII, 188.
 L. polyrrhiza, Linné, spec. plant. 970 (1753) ... — — V. N.S.W. — — B.f. VII, 164 M.fr. VIII, 188.
 L. gibba, Linné, spec. plant. 970 (1753) ... — W.A. — — — — B.f. VII, 163 M.fr. VIII, 188.
 WOLFFIA, Horkel & Schleiden in Schlecht. Linnaea XIII, 389 (1839). (Wolfia.)
 W. Michellii, Schleiden, Beitr. zur Botan. 233 (1844) ... — — — V. N.S.W. — — B.f. VII, 162 M.fr. VIII, 187.

FLUVIALES.

Ventenat, Tabl. du règn. végét. II, 80, (1799).

- TRIGLOCHIN, Rivinus in Ruppius, fl. Jenens. 54 et 366 (1718), from Dalechamps (1586). (Cycnogeton.)
 T. centrocarpa, Hooker, icon. plant. t. 728 (1848) ... — W.A. S.A. T. V. N.S.W. — — B.f. VII, 167 M.fr. VI, 82.
 T. striata, Ruiz & Pavon, Fl. Per. et Chil. III, 72 (1802) ... — W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 166 M.fr. VI, 83.
 T. mucronata, R. Brown, prodr. 343 (1810) ... — W.A. S.A. — V. — — — B.f. VII, 168 M.fr. VI, 81.
 T. procera, R. Brown, prodr. 343 (1810) ... — W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 168 M.fr. VI, 83.
 T. Maundii, F. v. M., fragm. VI, 83 (1867) ... — — — N.S.W. Q. — B.f. VII, 169

- APONOGETON, Linné fil., suppl. plant. 32 (1781).
 A. monostachyus, Linné fil., suppl. 214 (1781) ... — — — — — Q. N.A. B.f. VII, 188 M.fr. VIII, 215.
 A. elongatus, F. v. M. in Benth. Fl. Austr. VII, 188 (1878) ... — — — — — N.S.W. Q. N.A. B.f. VII, 188 M.fr. VIII, 216.

- POTAMOGETON, Fuchs, histor. stirp. comment. 651 (1542).
 P. natans, Linné, spec. plant. 126 (1753) ... — W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 170 M.fr. VIII, 217.
 P. Javanicus, Hasskarl in Act. Soc. Ind. Neerl. I, 26 (1857) ... — — — — — Q. N.A. B.f. VII, 171 M.fr. VIII, 217.
 P. Drummondii, Benth. Fl. Austr. VII, 171 (1878) ... — W.A. — — — — — — B.f. VII, 171
 P. perfoliatus, Linné, spec. plant. 126 (1753) ... — — — T. V. N.S.W. — — B.f. VII, 172 M.fr. VIII, 217.
 P. crispus, Linné, spec. plant. 126 (1753) ... — — — S.A. — V. N.S.W. Q. N.A. B.f. VII, 172 M.fr. VIII, 217.
 P. obtusifolius, Mertens & Koch, Deutschl. Fl. I, 855 (1823) ... — — — S.A. T. V. N.S.W. Q. — B.f. VII, 172 M.fr. VIII, 216.
 P. acutifolius, Link in Roem. & Schult. syst. veg. III, 513 (1818) ... — — — S.A. — V. N.S.W. — — B.f. VII, 173 M.fr. VIII, 216.
 P. pectinatus, Linné, spec. plant. 127 (1753) ... — — — S.A. T. V. — — — B.f. VII, 173 M.fr. VIII, 217.

- RUPPIA, Linné, syst. nat. 9 (1735); Linné, gen. plant. 277 (1737).
 R. maritima, Linné, spec. plant. 127 (1753) ... — — — W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 174 M.fr. VIII, 217

- POSIDONIA, Koenig, Annals of Botany II, 95, t. 6 (1806). (Caulinia partly, Kernera partly.)
 P. australis, J. Hooker, Fl. Tasman. II, 43 (1860) ... — — — W.A. S.A. T. V. — — — B.f. VII, 175 M.fr. VIII, 218.

- ZOSTERA, Linné, Waesgoeta-Resa, 166-168 et fig. (1747).
 Z. nana, F. K. Mertens in Roth, enum. pl. phaner. Germ. 8 (1827) ... — S.A. T. V. N.S.W. — — — B.f. VII, 176 M.fr. VIII, 218.
 Z. Tasmanica, G. v. Mertens in Garcke, Linnaea XXXV, 68 (1867) ... — S.A. T. V. — — — — B.f. VII, 176 M.fr. VIII, 218.

- CYMODOCEA, C. Koenig, Annals of Botany II, 96, t. 7 (1806). (Amphibolis, Graumuellera, Kernera partly.)
 C. zosterifolia, F. v. M.; Amphibolis, Agardh., spec. algar. II, 477 (1822) W.A. S.A. T. V. — — — — B.f. VII, 177 M.fr. VIII, 218.
 C. ciliata, Ehrenberg in Garcke, Linnaea XXXV, 162 (1867) ... — — — — — Q. N.A. B.f. VII, 178 M.fr. VIII, 218.
 C. rotundata, Ascherson & Schweinfurth in Neumayer, Anleit. zu wissenschaftl. Beob. 361 (1875) ... — — — — — — — N.A. — — M.fr. X, 162.

- C. serrulata, Ascherson & Magnus in Neumayer, Anleit. zu wissenschaftl. Beob. 362 (1875) ... — — — — — — — Q. — — B.f. VII, 178 M.fr. VIII, 218.
 C. isoetifolia, Ascherson in Garcke, Linnaea XXXV, 163 (1867) W.A. — — — — — — — N.A. B.f. VII, 178

- HALODULE, Endlicher, gen. plant. 1368 (1840). (Diplanthera, 1806.)
 H. tridentata, Endlicher, gen. plant. 1368, implied (1840) ... — — — — — — — N.A. — — M.fr. VIII, 218.

- LEPILAENA, Drummond & Harvey in Hook. Kew misc. VII, 57 (1855). (Zannichellia partly, Hexatheca, Altheniae subgenus.)
 L. australis, J. Drummond in Hook. Kew misc. VII, 58 (1855) ... — W.A. S.A. — — — — — — B.f. VII, 179 M.fr. VIII, 217.
 L. Preissii, F. v. M., fragm. VIII, 217 (1874) ... — — — W.A. S.A. T. V. — — — — B.f. VII, 180 M.fr. VIII, 217.

- NAJAS, Linné, syst. nat. 9 (1735); Linné, gen. plant. 278 (1737). (Caulinia partly.)
 N. major, Allioni, Fl. Pedem. II, 221 (1785) ... — — — W.A. S.A. — — — — — N.A. B.f. VII, 181 M.fr. VIII, 218.
 N. tenuifolia, R. Brown, prodr. 345 (1810) ... — — — — — S.A. — V. N.S.W. Q. — B.f. VII, 181 M.fr. VIII, 219.

ALISMACEAE.

Ventenat, Tabl. du règn. végét. II, 157 (1799).

- ALISMA, Rivinus in Ruppius, fl. Jen. 54 (1718), from Dioscorides and Plinius. (Caldesia; Echinodorus partly).
 A. Plantago, Linné, spec. plant. 842 (1753) ... — — — — — V. N.S.W. — — B.f. VII, 185 M.fr. VIII, 215.
 A. acanthocarpum, F. v. M., fragm. I, 23 (1858) ... — — — — — — — Q. N.A. B.f. VII, 185 M.fr. VIII, 214.
 A. oligococcum, F. v. M., fragm. I, 23 (1858) ... — — — — — — — Q. N.A. B.f. VII, 185 M.fr. VI, 169; VIII, 214.
 A. parnassifolium, Bassi in act. Bonon. (1768) ... — — — — — — — Q. — — B.f. VII, 186 M.fr. VIII, 214.

- DAMASONIUM, Tournefort, inst. 256 (1700). (Actinocarpus.)
 D. australe, Salisbury in Transact. hort. Soc. Lond. I, 268 (1812) W.A. S.A. — V. N.S.W. Q. — B.f. VII, 186 M.fr. VIII, 215.

- TENAGOPHARIS, Hochstetter in Regensb. bot. Zeit. June (1841). (Butomopsis.)
 T. Cordofana, Hochstetter in Regensb. bot. Zeit. June (1841) ... — — — — — — — Q. N.A. B.f. VII, 187 M.fr. X, 103.

PONTEDERIACEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. plant. I, 265 (1815).

- MONOCHORIA, Presl, reliq. Haenk. I, 127 (1827). (Limnostachys.)
 M. cyanea, F. v. M., fragm. VIII, 44 (1872) ... — — — — — — — Q. N.A. B.f. VII, 74 M.fr. VIII, 44.

PHILYDREAE.

R. Brown in Flinders' voy. II, 57 (1814).

PHILYDRUM, Banks in Gaertner, de fruct. I, 62 t. 16 (1788). (Philhydrium.)

P. lanuginosum, Banks in Gaertn. de fruct. I, 62 t. 16 (1788) ... — — — V. N.S.W. Q. N.A. B.fl.VII,73 M.fr.V,203.

PRITZELIA, F. v. M., Papuan plants 13 (1875). (Hetaeria, Philydrella.)

P. pygmaea, F. v. M., Papuan plants 13 (1875) ... — — — — — — — B.fl.VII,74 M.fr.V,203.

HELMHOLTZIA, F. v. M., fragm. V, 202 (1866).

H. acorifolia, F. v. M., fragm. V, 203 (1866) ... — — — — N.S.W. Q. — B.fl.VII,75 M.fr.V,203.

COMMELINEAE.

R. Brown, prodr. fl. Nov. Holl. 268 (1810).

ZYGOMENES, Salisbury in Transact. hort. Soc. I, 271 (1812). (Cyanotis).

Z. axillaris, Salisbury in Transact. hort. Soc. I, 271 (1812) ... — — — — — Q. N.A. B.fl.VII,82 M.fr.VIII,62.

COMMELINA, Plumier, nov. pl. Amer. gen. 48, t. 38 (1703). (Commelynia).

C. ensifolia, R. Brown, prodr. 269 (1810) ... — — — — S.A. — — N.S.W. Q. N.A. B.fl.VII,83 M.fr.VIII,60;IX,191.

C. cyanea, R. Brown, prodr. 269 (1810) ... — — — — — — — N.S.W. Q. N.A. B.fl.VII,84 M.fr.VIII,59;IX,191.

C. lanceolata, R. Brown, prodr. 269 (1810) ... — — — — — — — Q. N.A. B.fl.VII,84 M.fr.VIII,59.

C. agrostophylla, F. v. M. in J. Hook., Fl. Tasm. p. XLVII (1860) ... — — — — — — — Q. N.A. — M.fr.VIII,59.

ANEILEMA, R. Brown, prodr. fl. Nov. Holl. 270 (1810).

A. acuminatum, R. Brown, prodr. 270 (1810) ... — — — — — N.S.W. Q. — B.fl.VII,85 M.fr.VIII,61.

A. biflorum, R. Brown, prodr. 270 (1810) ... — — — — — N.S.W. Q. — B.fl.VII,86 M.fr.VIII,61.

A. sclerocarpum, F. v. M., fragm. VIII, 61 (1873) ... — — — — — — — Q. — B.fl.VII,86 M.fr.VIII,61.

A. siliculosum, R. Brown, prodr. 270 (1810) ... — — — — — — — Q. N.A. B.fl.VII,87 M.fr.VIII,61;IX,191.

A. gramineum, R. Brown, prodr. 270 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.VII,87 M.fr.VIII,62.

A. giganteum, R. Brown, prodr. 271 (1810) ... — — — — — — — Q. N.A. B.fl.VII,88 M.fr.VIII,62.

FLORISCOPIA, Loureiro, Fl. Cochinch. I, 192 (1790). (Floscopa, Dithyrocarpus.)

F. scandens, Loureiro, fl. Cochinch. I, 193 (1790) ... — — — — — — — Q. — B.fl.VII,89 M.fr.VIII,63.

POLLIA, Thunberg, nov. gen. plant. I, 11 (1781).

P. cyanococca, F. v. M., fragm. V, 40 (1865) ... — — — — — N.S.W. Q. — B.fl.VII,90 M.fr.VIII,63.

CARTONEMA, R. Brown, prodr. fl. Nov. Holl. 271 (1810).

C. philydroides, F. v. M., fragm. I, 62 (1858) ... — — — — — W.A. — — — B.fl.VII,91 M.fr.VIII,64.

C. spicatum, R. Brown, prodr. 271 (1810) ... — — — — — — — N.A. B.fl.VII,91 M.fr.I,62;VIII,64.

C. parviflorum, Hasskarl in Flora Regensb. 365 (1869) ... — — — — — — — N.A. B.fl.VII,91 M.fr.VIII,64.

C. trigonospermum, Clarke in De Cand., mon. phaner. III, 24 (1881) ... — — — — — — — N.A. — M.fr.XI,139.

C. brachyantherum, Benthams, Fl. Austr. VII, 92 (1878) ... — — — — — — — Q. — B.fl.VII,92 M.fr.VIII,64.

C. tenue, Benthams in De Cand., monogr. phaner. III, 264 (1881) ... — — — — — — — N.A. — M.fr.XI,139.

XYRIDEAE.

Salisbury in Transact. of the Hort. Soc. I, 326 (1812).

XYRIS, Gronovius in Linné, genera plantar. 11 (1737).

X. complanata, R. Brown, prodr. 256 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.VII,77 M.fr.VIII,205.

X. pauciflora, Willdenow, phytogr. 2, t. 1 (1794) ... — — — — — — — Q. N.A. B.fl.VII,77 M.fr.VIII,205.

X. lacera, R. Brown, prodr. 257 (1810) ... — — — — — W.A. — — — B.fl.VII,78 M.fr.VIII,204.

X. flexifolia, R. Brown, prodr. 256 (1810) ... — — — — — W.A. — — — B.fl.VII,78 M.fr.VIII,204.

X. gracilis, R. Brown, prodr. 256 (1810) ... — — — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,79 M.fr.VIII,204.

X. operculata, Labillardière, Nov. Holl. pl. spec. I, 14 t. 10 (1804) ... — — — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,79 M.fr.VIII,204.

X. lanata, R. Brown, prodr. 257 (1810) ... — — — — — W.A. — — — B.fl.VII,80 M.fr.VIII,204.

X. laxiflora, F. v. M., fragm. VIII, 203 (1874) ... — — — — — W.A. — — — B.fl.VII,80 M.fr.VIII,203.

X. gracillima, F. v. M., fragm. VIII, 203 (1874) ... — — — — — W.A. — — — B.fl.VII,80 M.fr.VIII,203.

JUNCEAE.

R. Brown, prodr. fl. Nov. Holl. 257 (1810).

LUZULA, De Candolle, Fl. franc. III, 158 (1805).

L. campestris, De Candolle, Fl. franc. III, 158 (1805) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,123

L. longiflora, Benthams, Fl. Austr. VII, 123 (1878) ... — — — — — — — N.S.W. — — B.fl.VII,123

JUNCUS, Tournefort, inst. 246, t. 127 (1700), from Camerarius, J. & C. Bauhin, Ray & Morison.

J. gracilis, R. Brown, prodr. 259 (1810) ... — — — — — W.A. — — — B.fl.VII,125

J. planifolius, R. Brown, prodr. 259 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. — — B.fl.VII,125

J. caespitosus, E. Meyer in Lehm. pl. Preiss. II, 47 (1846) ... — — — — — W.A. S.A. T. V. N.S.W. — — B.fl.VII,126

J. falcatus, E. Meyer, synops. luzular. 34 (1823) ... — — — — — — — T. V. N.S.W. — — B.fl.VII,126

J. bufonius, Linné, spec. plant. 328 (1753) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,127

J. homalocaulis, F. v. M., first gen. Rep. 19 (1853) ... — — — — — — — S.A. — — B.fl.VII,128

J. Brownii, F. v. M., first gen. Rep. 19 (1853) ... — — — — — — — T. V. N.S.W. — — B.fl.VII,128

J. communis, E. Meyer, synops. Juncor. 12 (1822) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,128

J. vaginatus, R. Brown, prodr. 258 (1810) ... — — — — — — — — — B.fl.VII,129

J. pauciflorus, R. Brown, prodr. 259 (1810) ... — — — — — S.A. T. V. N.S.W. Q. — B.fl.VII,129

J. pallidus, R. Brown, prodr. 258 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,130

J. maritimus, Lamarek, Encycl. méth. III, 264 (1789) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,130

J. prismatocarpus, R. Brown, prodr. 259 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,131

J. pusillus, Buchenau in Abhandl. naturw. Vereins zu Bremen VI, 395 (1879) ... — — — — — — — T. V. N.S.W. — — B.fl.VII,132

ERIOCAULEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. plant. I, 251 (1815).

ERIOCAULON, Linné, gen. plant. ed. sec. 35 (1742). (Electrosperma.)

E. setaceum, Linné, spec. plant. 87 (1753) ...	—	—	—	—	Q.	N.A.	B.f. VII, 191	
E. australe, R. Brown, prodr. 254 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 192
E. quinquangulare, Linné, spec. plant. 87 (1753) ...	—	—	—	—	—	N.A.	B.f. VII, 192	
E. Smithii, R. Brown, prodr. 254 (1810) ...	—	—	—	—	V.	N.S.W.	Q.	B.f. VII, 192
E. nanum, R. Brown, prodr. 254 (1810) ...	—	—	—	—	—	Q.	—	B.f. VII, 193
E. cinereum, R. Brown, prodr. 254 (1810) ...	—	—	—	—	—	N.A.	B.f. VII, 193	M.fr. I, 95.
E. pusillum, R. Brown, prodr. 254 (1810) ...	—	—	—	—	—	Q.	—	B.f. VII, 194
E. pallidum, R. Brown, prodr. 254 (1810) ...	—	—	—	—	—	Q.	—	B.f. VII, 194
E. nigricans, R. Brown, prodr. 254 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 194
E. electrospermum; Electrosperma Australasicum, F. v. M. in Trans. phil. Soc. Vict. I, 24 (1854) ...	—	—	—	V.	N.S.W.	—	—	B.f. VII, 195
E. lividum, F. v. M., fragm. I, 92 (1858) ...	—	—	—	—	—	N.A.	B.f. VII, 195	M.fr. I, 92.
E. concretum, F. v. M., fragm. I, 92 (1858) ...	—	—	—	—	—	N.A.	B.f. VII, 195	M.fr. I, 92.
E. Schultzii, Benth. Fl. Austr. VII, 195 (1878) ...	—	—	—	—	—	N.A.	B.f. VII, 195	
E. tortuosum, F. v. M., fragm. I, 94 (1858) ...	—	—	—	—	—	N.A.	B.f. VII, 196	M.fr. I, 91.
E. monoscapum, F. v. M., fragm. I, 94 (1858) ...	—	—	—	—	—	N.A.	B.f. VII, 196	M.fr. I, 94.
E. spectabile, F. v. M., fragm. I, 95 (1858) ...	—	—	—	—	—	N.A.	B.f. VII, 196	M.fr. I, 95.
E. scarosum, R. Brown, prodr. 255 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 197
E. depressum, R. Brown, prodr. 255 (1810) ...	—	—	—	—	—	N.A.	B.f. VII, 197	M.fr. I, 92.

RESTIACEAE.

R. Brown, prodr. flor. Nov. Holl. 243 (1810).

TRITHURIA, J. Hooker, Fl. Tasman. II, 79, t. 138 (1860). (Juncella, 1854.)

T. submersa, J. Hooker, Fl. Tasman. II, 79, t. 138 (1860) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 199	M.fr. VIII, 237.
---	------	------	----	----	--------	---	---	---------------	------------------

APHELIA, R. Brown, prodr. fl. Nov. Holl. 251 (1810). (Brizula.)

A. cyperoides, R. Brown, prodr. 252 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 200	M.fr. VIII, 236.
A. nutans, J. Hooker in Benth. Fl. Austr. VII, 200 (1878) ...	W.A.	—	—	—	—	—	—	B.f. VII, 200	
A. gracilis, Sonder in Schlecht. Linnaea XXVIII, 227 (1855) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 201	M.fr. VIII, 237.
A. Drummondii, Benth. Fl. Austr. VII, 201 (1878) ...	W.A.	—	—	—	—	—	—	B.f. VII, 201	
A. Pumilio, F. v. M., Schlecht. in Linnaea XXVIII, 226 (1855) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 201	M.fr. VIII, 237.
A. Brizula, F. v. M., fragm. V, 203 (1866) ...	W.A.	—	—	—	—	—	—	B.f. VII, 202	M.fr. VIII, 237.

CENTROLEPIS, Labillardière, Nov. Holl. pl. specim. I, 7 (1804). (Desvauxia, Alepyrum.)

C. humillima, F. v. M. in Benth. Fl. Austr. VII, 203 (1878) ...	W.A.	—	—	—	—	—	—	B.f. VII, 203	
C. polygyna, Hieronymus in Abh. naturf. Ges. Halle XII, 96 (1873) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 203	M.fr. VIII, 237.
C. alepyroides, Hieronymus in Abh. nat. Ges. Halle XII, 96 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 204	M.fr. VIII, 237.
C. mutica, Hieronymus in Abh. nat. Ges. Halle XII, 97 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 204	
C. glabra, F. v. M. in Abh. nat. Ges. Halle XII, 95 (1873) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 204	M.fr. VIII, 237.
C. muscoides, Hieronymus in Abh. nat. Ges. Halle XII, 95 (1873) ...	—	—	T.	—	—	—	—	B.f. VII, 205	
C. monogyna, Benth. Fl. Austr. VII, 205 (1878) ...	—	—	T.	—	—	—	—	B.f. VII, 205	
C. pulvinata, Desvaux in Ann. des sc. nat. XIII, 42, t. 2 (1828) ...	—	—	T.	—	—	—	—	B.f. VII, 205	
C. pusilla, Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	—	B.f. VII, 205	
C. aristata, Roemer & Schultes, syst. veg. I, 44 (1817) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 206	M.fr. VIII, 237.
C. Drummondii, Hieronymus in Abh. nat. Ges. Halle XII, 98 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 206	M.fr. VIII, 237.
C. Banksii, Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 207	M.fr. VIII, 237.
C. fascicularis, Labillardière, Nov. Holl. pl. spec. I, 7, t. 1 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 207	
C. pilosa, Hieronymus in Abh. nat. Ges. Halle 102 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 207	
C. strigosa, Roemer & Schultes, syst. veg. I, 43 (1817) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 207	
C. exserta, Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 208	M.fr. VIII, 237.

LYGINIA, R. Brown, prodr. fl. Nov. Holl. 248 (1810). (Schoenodum partly.)

L. barbata, R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 210	M.fr. VIII, 79.
---	------	---	---	---	---	---	---	---------------	-----------------

ECDEIOCOLEA, F. v. M., fragm. phytogr. Austr. VIII, 236 (1874).

E. monostachya, F. v. M., fragm. VIII, 236 (1874) ...	W.A.	—	—	—	—	—	—	B.f. VII, 211	M.fr. VIII, 236.
---	------	---	---	---	---	---	---	---------------	------------------

ANARTHRIA, R. Brown, prodr. fl. Nov. Holl. 248 (1810).

A. scabra, R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 212	M.fr. VIII, 80.
A. laevis, R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 212	M.fr. VIII, 81.
A. gracilis, R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 213	M.fr. VIII, 72, 81.
A. prolifera, R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 213	M.fr. VIII, 82.
A. polyphylla, Nees in Lehm. pl. Preiss. II, 63 (1846) ...	W.A.	—	—	—	—	—	—	B.f. VII, 214	M.fr. VIII, 83.

LEPYRODIA, R. Brown, prodr. fl. Nov. Holl. 247 (1810).

L. scariosa, R. Brown, prodr. 248 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f. VII, 215	M.fr. VIII, 72.
L. Muelleri, Benth. Fl. Austr. VII, 215 (1878) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 215	
L. anarthria, F. v. M., fragm. VIII, 73 (1873) ...	—	—	—	—	N.S.W.	—	—	B.f. VII, 216	
L. gracilis, R. Brown, prodr. 247 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f. VII, 216	M.fr. VIII, 73.
L. Tasmanica, J. Hooker, Fl. Tasman. II, 72, t. 135 (1860) ...	—	—	T.	V.	—	—	—	B.f. VII, 216	M.fr. VIII, 73, 75; IX, 194.
L. interrupta, F. v. M., fragm. VIII, 74 (1873) ...	—	—	—	—	V.	N.S.W.	Q.	B.f. VII, 217	M.fr. VIII, 74.
L. hermaphrodita, R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 217	M.fr. VIII, 75.
L. monoica, F. v. M., fragm. VIII, 76 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 217	M.fr. VIII, 76.
L. Muirii, F. v. M., fragm. VIII, 78 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 218	M.fr. VIII, 76.
L. stricta, R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.f. VII, 218	M.fr. VIII, 74.
L. macra, Nees in Lehm. pl. Preiss. II, 60 (1846) ...	W.A.	—	—	—	—	—	—	B.f. VII, 218	
L. Drummondiana, Steudel, syn. pl. glum. II, 248 (1855) ...	W.A.	—	—	—	—	—	—	B.f. VII, 219	M.fr. VIII, 75.
L. glauca, F. v. M., fragm. VIII, 77 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 219	M.fr. VIII, 77.
L. anaectocolea, F. v. M., fragm. VIII, 78 (1873) ...	W.A.	—	—	—	—	—	—	B.f. VII, 220	M.fr. VIII, 78.

RESTIO, Linné, syst. nat. edit. duo dec. II, 735 (1767). (Megalotheca.)

R. fastigiatus, R. Brown, prodr. 246 (1810)	...	...	...	...	N.S.W.	—	—	B.f. VII, 222	M.fr. VIII, 67.
R. Megalotheca, F. v. M., fragm. VIII, 99 (1873)	...	...	W.A.	—	—	—	—	B.f. VII, 222	
R. tropicus, R. Brown, prodr. 246 (1810)	...	...	—	—	—	—	N.A.	B.f. VII, 223	
R. applanatus, Sprengel, syst. I, 185 (1825)	...	...	W.A.	—	—	—	—	B.f. VII, 223	M.fr. VIII, 69.
R. monocephalus, R. Brown, prodr. 245 (1810)	...	...	—	—	T.	—	—	B.f. VII, 224	M.fr. VIII, 68.
R. dimorphus, R. Brown, prodr. 246 (1810)	...	...	—	—	—	—	N.S.W. Q.	B.f. VII, 224	M.fr. VIII, 68.
R. conferto-spiciatus, Steudel, syn. glum. II, 256 (1855)	...	...	W.A.	—	—	—	—	B.f. VII, 224	
R. sphacelatus, R. Brown, prodr. 246 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 225	
R. deformis, R. Brown, prodr. 245 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 225	M.fr.
R. crispatus, R. Brown, prodr. 246 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 225	
R. nitens, Nees in Lehm. pl. Preiss. II, 59 (1846)	...	...	W.A.	—	—	—	—	B.f. VII, 226	M.fr. VIII, 70.
R. gracilior, F. v. M. in Benth. Fl. Austr. VII, 226 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 226	
R. chaunocoleus, F. v. M., fragm. VIII, 64 (1873)	...	...	W.A.	—	—	—	—	B.f. VII, 226	M.fr.
R. australis, R. Brown, prodr. 245 (1810)	...	...	—	—	T. V.	N.S.W.	—	B.f. VII, 227	M.fr. VIII, 69.
R. gracilis, R. Brown, prodr. 245 (1810)	...	...	—	—	T. V.	N.S.W. Q.	—	B.f. VII, 227	M.fr. VIII, 69.
R. complanatus, R. Brown, prodr. 245 (1810)	...	...	—	—	S.A.	T. V.	N.S.W. Q.	B.f. VII, 228	M.fr. VIII, 67.
R. tetraphyllus, Labillardière, Nov. Holl. pl. spec. II, 77 (1806)	...	...	—	—	S.A.	T. V.	N.S.W. Q.	B.f. VII, 228	M.fr. VIII, 66.
R. laxus, R. Brown, prodr. 245 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 228	M.fr. VIII, 71.
R. ornatus, Steudel, syn. pl. glum. II, 256 (1855)	...	...	W.A.	—	—	—	—	B.f. VII, 229	M.fr. VIII, 70.
R. leptocarpoides, Benth. Fl. Austr. VII, 229 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 229	M.fr. VIII, 65.
R. amblycoleus, F. v. M., fragm. VIII, 65 (1873)	...	...	W.A.	—	—	—	—	B.f. VII, 230	M.fr. VIII, 65.
R. tremulus, R. Brown, prodr. 230 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 230	M.fr. VIII, 70.

LOXOCARYA, R. Brown, prodr. fl. Nov. Holl. 249 (1810). (Desmocladus.)

L. densa, Benth. Fl. Austr. VII, 241 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 241	
L. Benth. F. v. M.; Hypolaena, Masters in Cand. mon. phan. I, 367 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 242	
L. pubescens, Benth. Fl. Austr. VII, 242 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 242	M.fr. VIII, 100.
L. fasciculata, Benth. Fl. Austr. VII, 242 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 242	M.fr. VIII, 98.
L. flexuosa, Benth. Fl. Austr. VII, 243 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 243	M.fr. VIII, 100.
L. cinerea, R. Brown, prodr. 249 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 243	

CALOSTROPHUS, Labillardière, Nov. Holl. pl. spec. II, 78 (1806). (Calorophus, Hypolaena.)

C. elongatus, F. v. M., fragm. VIII, 86 (1873)	...	...	—	—	T.	—	—	B.f. VII, 238	M.fr. VIII, 86; IX, 94; X, 120
C. lateriflorus, F. v. M., fragm. VIII, 87 (1873)	...	...	—	—	S.A.	T. V.	N.S.W. Q.	B.f. VII, 238	M.fr. VIII, 87.
C. gracillimus, F. v. M., fragm. VIII, 88 (1873)	...	...	W.A.	—	—	—	—	B.f. VII, 239	M.fr. VIII, 88.
C. fastigiatus, F. v. M. in pap. R. Soc. Tasm. 117 (1878)	...	...	—	—	S.A.	T. V.	N.S.W.	B.f. VII, 239	M.fr. VIII, 84.
C. exsulcus, F. v. M.; Hypolaena, R. Brown, prodr. 251 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 240	M.fr. VIII, 86.

LEPTOCARPUS, R. Brown, prodr. 250 (1810). (Schoenodum partly.)

L. scariosus, R. Brown, prodr. 250 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 232	M.fr. VIII, 94.
L. tenax, R. Brown, prodr. 250 (1810)	...	...	W.A.	S.A.	T. V.	N.S.W.	—	B.f. VII, 232	M.fr. VIII, 96.
L. Brownii, J. Hooker, Fl. Tasman. II, 73, t. 136 (1860)	...	...	—	S.A.	T. V.	N.S.W.	—	B.f. VII, 233	M.fr. VIII, 91.
L. canus, Nees in Ann. of nat. hist. VI, 50 (1843)	...	...	W.A.	—	—	—	—	B.f. VII, 234	M.fr. VIII, 95.
L. coangustatus, Nees in Lehm. pl. Preiss. II, 65 (1846)	...	...	W.A.	—	—	—	—	B.f. VII, 234	M.fr. VIII, 94.
L. aristatus, R. Brown, prodr. 250 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 235	M.fr. VIII, 90.
L. erianthus, Benth. Fl. Austr. VII, 235 (1878)	...	...	W.A.	—	—	—	—	B.f. VII, 235	
L. ramosus, R. Brown, prodr. 250 (1810)	...	...	—	—	—	—	Q.	B.f. VII, 236	M.fr. VIII, 92.
L. elatior, R. Brown, prodr. 250 (1810)	...	...	—	—	—	—	Q.	N.A.	B.f. VII, 236
L. spathaceus, R. Brown, prodr. 250 (1810)	...	...	—	—	—	—	—	N.A.	B.f. VII, 236
L. Schultzei, Benth. Fl. Austr. VII, 237 (1878)	...	...	—	—	—	—	—	N.A.	B.f. VII, 237

LEPIDOBOLUS, Nees in Lehm. pl. Preiss. II, 66 (1846).

L. drapetocoleus, F. v. M., fragm. VIII, 84 (1873)	...	...	—	S.A.	—	V.	—	B.f. VII, 244	M.fr. VIII, 84.
L. Preissianus, Nees in Lehm. pl. Preiss. II, 66 (1846)	...	...	W.A.	—	—	—	—	B.f. VII, 245	M.fr. VIII, 83.
L. chaetocephalus, F. v. M., fragm. VIII, 84 (1873)	...	...	W.A.	—	—	—	—	B.f. VII, 245	

CHAETANTHUS, R. Brown, prodr. 251 (1810). (Prionosepalum)

C. leptocarpoides, R. Brown, prodr. 251 (1810)	...	...	W.A.	—	—	—	—	B.f. VII, 246	M.fr. VIII, 97.
--	-----	-----	------	---	---	---	---	---------------	-----------------

ONYCHOSEPALUM, Steudel, syn. pl. glum. II, 249 (1855).

O. laxiflorum, Steudel, syn. pl. glum. II, 249 (1855)	...	...	W.A.	—	—	—	—	B.f. VII, 246	M.fr. IX, 51.
---	-----	-----	------	---	---	---	---	---------------	---------------

ACALYCEAE HYPOGYNEAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 48 (1880).

CYPERACEAE.

Haller, enum. stirp. Helvet. I, 234 (1742).

KYLINGIA, Rottboell, descr. et icon. rar. et nov. plant. 13, t. 4 (1773). (Killingia.)

K. intermedia, R. Brown, prodr. 219 (1810)	...	...	—	S.A.	—	V.	N.S.W. Q.	—	B.f. VII, 251	M.fr. VIII, 27.
K. monocephala, Rottboell, descr. et icon. pl. 13, t. 4 (1773)	...	...	—	S.A.	—	—	N.S.W. Q.	—	B.f. VII, 251	M.fr. VIII, 271; IX, 54.
K. cylindrica, Nees in Wight's Contr. to Bot. of Ind. 91 (1834)	...	...	—	—	—	—	N.S.W. Q.	—	B.f. VII, 252	M.fr. VIII, 271.
K. triceps, Rottboell, descr. et icon. pl. 14, t. 4 (1773)	...	...	—	—	—	—	—	Q.	—	B.f. VII, 252

CYPERUS, Tournefort, inst. 527, t. 299 (1700), from Hippocr., Theophr. and Plinius. (Pycnus, Mariscus, Anosporum, Sorostachys.)

C. pumilus, Linné, amoen. acad. IV, 302 (1759) ...	...	...	—	—	—	—	Q.	—	B. fl. VII, 258	M. fr. VIII, 267.	
C. Eragrostis, Vahl, enum. plant. II, 322 (1806) ...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B. fl. VII, 258	M. fr. VIII, 260; IX, 52.
C. flavescens, Linné, spec. plant. 46 (1753) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B. fl. VII, 259	
C. globosus, Allioni, auctuar. ad fl. Pedem. 49 (1789) ...	...	...	—	—	—	V.	—	Q.	—	B. fl. VII, 260	M. fr. VIII, 260; IX, 52.
C. unioloides, R. Brown, prodr. 216 (1810) ...	...	...	—	—	—	V.	N.S.W.	Q.	—	B. fl. VII, 260	M. fr. VIII, 259; IX, 52.

F. cardiocarpa, F. v. M., fragm. I, 194 (1859) ...	—	—	—	—	—	N.A.	B. fl. VII, 303	M. fr. I, 194.
F. leucocolea, Benth. Fl. Austr. VII, 304 (1878) ...	—	—	—	—	—	N.A.	B. fl. VII, 304	
F. polytrichoides, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 304
F. androgyna, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	N.A.	B. fl. VII, 304	
F. subbulbosa, Benth. Fl. Austr. VII, 305 (1878) ...	—	—	—	—	—	Q.	—	B. fl. VII, 305
F. tetragona, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	N.A.	B. fl. VII, 305	M. fr. I, 194; VIII, 274.
F. trigastrocarya, F. v. M., fragm. I, 194 (1859) ...	—	—	—	—	—	N.A.	B. fl. VII, 305	M. fr. I, 194.
F. monandra, F. v. M., fragm. I, 195 (1859) ...	—	—	—	—	—	N.A.	B. fl. VII, 306	M. fr. I, 195.
F. pterygosperma, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	N.A.	B. fl. VII, 306	M. fr. I, 193; IX, 55.
F. sphaerocephala, Benth. Fl. Austr. VII, 306 (1878) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 306
F. xyridis, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 307
F. oxystachya, F. v. M., fragm. I, 195 (1859) ...	—	—	—	—	—	N.A.	B. fl. VII, 307	M. fr. VIII, 272.
F. macrantha, Boeckeler in Garcke in Linnaea XXXVIII, 388 (1874) ...	—	—	—	—	—	N.A.	B. fl. VII, 307	M. fr. IX, 55.
F. squarrola, F. v. M., fragm. I, 216 (1859) ...	—	—	—	—	—	N.A.	B. fl. VII, 308	M. fr. I, 216.
F. monostachya, Hasskarl, pl. Jav. rar. 61-64 (1848) ...	—	—	—	—	N.S.W.	Q.	N.A.	B. fl. VII, 308
F. Brownii, Benth. Fl. Austr. VII, 308 (1878) ...	—	—	—	—	—	N.A.	B. fl. VII, 308	M. fr. VIII, 273.
F. Dallachyi, F. v. M. in Benth. Fl. Austr. VII, 309 (1878) ...	—	—	—	—	—	Q.	—	B. fl. VII, 309
F. velata, R. Brown, prodr. 227 (1810) ...	—	—	—	S.A.	—	Q.	N.A.	B. fl. VII, 309
F. aestivalis, Vahl, enum. II, 288 (1806) ...	—	—	—	—	V.	N.S.W.	Q.	N.A.
F. dichotoma, Vahl, enum. II, 287 (1806) ...	—	—	—	—	—	N.S.W.	Q.	N.A.
F. depauperata, R. Brown, prodr. 227 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 311
F. spirostachya, F. v. M. in Benth. Fl. Austr. VII, 311 (1878) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 311
F. communis, Kunth, enum. II, 234 (1837) ...	—	—	—	S.A.	—	V.	N.S.W.	Q.
F. ferruginea, Vahl, enum. 291 (1806) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 312
F. denudata, R. Brown, prodr. 227 (1810) ...	—	—	—	—	—	—	Q.	N.A.
F. elata, R. Brown, prodr. 227 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 313
F. caespitosa, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	—	Q.	N.A.
F. spiralis, R. Brown, prodr. 226 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 314
F. subaristata, Benth. Fl. Austr. VII, 314 (1878) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 314
F. leptoclada, Benth. Fl. Austr. VII, 314 (1878) ...	—	—	—	—	—	Q.	—	B. fl. VII, 314
F. debilis, F. v. M., fragm. I, 198 (1859) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 315
F. corynocarya, F. v. M., fragm. I, 197 (1859) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 315
F. solidifolia, F. v. M., fragm. I, 198 (1859) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 315
F. obtusangula, F. v. M., fragm. I, 198 (1859) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 315
F. miliacea, Vahl, enum. II, 287 (1806) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 316
F. rara, R. Brown, prodr. 227 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 316
F. microcarya, F. v. M., fragm. I, 200 (1859) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 316
F. quinqueangularis, Kunth, enum. II, 229 (1837) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 317
F. cyperoides, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	N.S.W.	Q.	N.A.
F. furva, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	Q.	—	B. fl. VII, 318
F. cymosa, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 318
F. multifolia, Boeckeler in Garcke, Linnaea XXXVIII, 397 (1874) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 319
F. sericea, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 319
F. macrostachya, Boeckeler in Garcke, Linn. XXXVIII, 386 (1874) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 319
F. Neilsoni, F. v. M., fragm. IX, 79 (1875) ...	—	—	—	S.A.	—	N.S.W.	—	B. fl. VII, 320
F. capitata, R. Brown, prodr. 228 (1810) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 320
F. Schultzii, Boeckeler in Garcke, Linnaea XXXVIII, 391 (1874) ...	—	—	—	—	—	—	N.A.	B. fl. VII, 321
F. barbata, Benth. Fl. Austr. VII, 321 (1878) ...	—	—	—	S.A.	—	N.S.W.	Q.	N.A.
F. capillaris, A. Gray, Man. of Bot. N.U.S. fifth ed. 567 (1867) ...	—	—	—	W.A.	—	—	Q.	N.A.
SCIRPUS, Tournefort, inst. 528, t. 300 (1700), from Terentius. (Isoplepis, Malacochaete.)								
S. humillimus, Benth. Fl. Austr. VII, 324 (1878) ...	—	—	—	—	—	N.A.	B. fl. VII, 324	M. fr. IX, 7.
S. fluitans, Linné, spec. plant. 48 (1753) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	—
S. arenarius, Benth. Fl. Austr. VII, 325 (1878) ...	—	—	—	W.A.	—	V.	—	B. fl. VII, 325
S. lenticularis, Sprengel, syst. I, 208 (1825) ...	—	—	—	—	—	N.S.W.	—	B. fl. VII, 326
S. crassiusculus, J. Hooker in Benth. Fl. Austr. VII, 326 (1878) ...	—	—	—	—	—	T. V.	N.S.W.	—
S. cyperoides, Sprengel, syst. I, 208 (1825) ...	—	—	—	W.A.	—	—	—	B. fl. VII, 326
S. setaceus, Linné, spec. plant. 49 (1753) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	Q.
S. riparius, Sprengel, syst. I, 208 (1825) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	—
S. cartilagineus, Sprengel, syst. I, 208 (1825) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	—
S. squarrosus, Linné, mant. plant. alt. 181 (1771) ...	—	—	—	—	—	—	Q.	N.A.
S. inundatus, Sprengel, syst. I, 207 (1825) ...	—	—	—	—	S.A.	T. V.	N.S.W.	Q.
S. prolifer, Rottboell, descr. et icon. pl. 55 t. 17 (1773) ...	—	—	—	—	—	—	N.S.W.	—
S. articulatus, Linné, spec. plant. 47 (1753) ...	—	—	—	—	—	—	Q.	N.A.
S. nodosus, Rottboell, descr. et icon. pl. 52, t. 8 (1773) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	Q.
S. supinus, Linné, spec. plant. 49 (1753) ...	—	—	—	—	S.A.	—	V.	N.S.W.
S. debilis, Pursch, flor. Amer. Sept. I, 55 (1814) ...	—	—	—	—	—	—	Q.	N.A.
S. mucronatus, Linné, spec. plant. 50 (1753) ...	—	—	—	—	—	—	N.S.W.	Q.
S. pungens, Vahl, enum. II, 255 (1806) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	—
S. lacustris, Linné, spec. plant. 48 (1753) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	Q.
S. litoralis, Schrader, Fl. Germ. I, 142, t. 5 (1806) ...	—	—	—	—	S.A.	—	—	N.S.W.
S. maritimus, Linné, spec. plant. 51 (1753) ...	—	—	—	W.A.	S.A.	T. V.	N.S.W.	Q.
S. grossus, Linné, fil. suppl. plant. 104 (1781) ...	—	—	—	—	—	—	Q.	—
S. polystachyus, F. v. M. in Transact. phil. Soc. Vict. I, 108 (1854) ...	—	—	—	—	—	V.	N.S.W.	—
LIPOCARPHA, R. Brown in Tuckey's narrat. of an Exped. to Congo, 459 (1818). (Hypaclyptum.)								
L. argentea, R. Brown in Tuck. Congo, 459 (1818) ...	—	—	—	—	—	Q.	—	B. fl. VII, 336
L. microcephala, R. Brown in Tuck. Congo, 459 (1818) ...	—	—	—	W.A.	S.A.	—	V.	N.S.W.
FUIRENA, Rottboell, descr. et icon. pl. illust. 70 (1773).								
F. umbellata, Rottboell, descr. et icon. pl. 70, t. 19 (1773) ...	—	—	—	—	—	Q.	N.A.	B. fl. VII, 337
F. glomerata, Lamarck, Illustr. des genr. I, 150 (1791) ...	—	—	—	—	—	N.S.W.	Q.	N.A.

HYPELYTRUM, L. C. Richard in Persoon, synops. I, 70 (1805). (Hypolytrum.)									
H. latifolium, L. C. Richard in Pers. syn. I, 70 (1805) ...	...	...	...	...	...	...	...	Q.	B.f. VII, 339 M.fr. VIII, 238; IX, 57.
EXOCARYA, Benth in J. Hooker, icon. plant. t. 1206 (1877).									
E. scleroides, Benth in Hook. icon. pl. t. 1206 (1877)...	...	...	...	...	...	...	...	N.S.W. Q.	B.f. VII, 339 M.fr. IX, 12.
MAPANIA, Aublet, Hist. pl. Guian. 47, t. 17 (1775). (Pandanophyllum.)									
M. hypelytroides, F. v. M., fragm. IX, 16 (1875)...	...	...	...	...	...	...	...	Q.	B.f. VII, 341 M.fr. IX, 16.
SCIRPODENDRON, Zippelius in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869).									
S. costatum, Kurz in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869) ...	...	...	...	...	...	...	...	Q.	B.f. VII, 341 M.fr. X, 104.
LEPIRONIA, L. C. Richard in Persoon, synops. I, 70 (1805). (Chondrachne.)									
L. mucronata, L. C. Richard in Pers. syn. I, 70 (1805) ...	...	...	...	...	...	...	...	N.S.W. Q.	B.f. VII, 342 M.fr. IX, 17, 58.
CHORIZANDRA, R. Brown, prodr. 221 (1810). (Chorisandra.)									
C. spaerocephala, R. Brown, prodr. 221 (1810) ...	...	...	...	...	...	...	...	N.S.W. Q.	B.f. VII, 344 M.fr. IX, 18, 58.
C. enodis, Nees in Lehm. pl. Preiss. II, 73 (1846)...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W.	B.f. VII, 344 M.fr. IX, 18.
C. multiarticulata, Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 345 M.fr. IX, 18.
C. cymbaria, R. Brown, prodr. 221 (1810) ...	...	...	...	...	W.A.	...	...	V. N.S.W. Q.	B.f. VII, 345 M.fr. IX, 18, 58
OREOBOLUS, R. Brown, prodr. fl. Nov. Holl. 235 (1810).									
O. Pumilio, R. Brown, prodr. 226 (1810) ...	...	...	...	...	...	...	T. V.	N.S.W.	B.f. VII, 346 M.fr. IX, 20, 58.
REMIREA, Aublet, Hist. des pl. de la Guian. franc. I, 44, t. 16 (1775).									
R. maritima, Aublet, Hist. pl. Guian. I, 45, t. 16 (1775) ...	...	...	...	...	...	...	...	Q.	B.f. VII, 347 M.fr. V, 92; IX, 20, 58.
RHYNCHOSPORA, Vahl, enum. plant. II, 229 (1806). (Morisia.)									
R. aurea, Vahl, enum. II, 229 (1806) ...	...	...	...	...	...	...	...	Q.	B.f. VII, 348 M.fr. IX, 17, 57.
R. glauca, Vahl, enum. II, 229 (1806) ...	...	...	...	...	...	...	...	N.S.W. Q.	B.f. VII, 349 M.fr. IX, 17, 57.
R. Wallichiana, Kunth, enum. II, 289 (1837) ...	...	...	...	...	...	...	...	Q. N.A.	B.f. VII, 349 M.fr. IX, 17, 52.
R. longisetis, R. Brown, prodr. 230 (1810)...	...	...	...	...	...	...	...	Q. N.A.	B.f. VII, 350 M.fr. IX, 17, 58.
R. tenuifolia, Benth in Fl. Austr. VII, 350 (1878) ...	...	...	...	...	...	...	...	Q. N.A.	B.f. VII, 350
CYATHOCHAETE, Nees in Lehm. pl. Preiss. II, 86 (1846). (Tetralepsis.)									
C. clandestina, Benth in Fl. Austr. VII, 351 (1878) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 351 M.fr. IX, 40.
C. avenacea, Benth in Fl. Austr. VII, 351 (1878) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 351 M.fr. IX, 40.
C. diandra, Nees in Lehm. pl. Preiss. II, 86 (1846) ...	...	...	...	...	...	...	V. N.S.W.	...	B.f. VII, 352 M.fr. IX, 39.
CARPHA, Banks & Solander in R. Brown, prodr. 230 (1810).									
C. alpina, R. Brown, prodr. 280 (1810) ...	...	...	...	...	...	...	T. V.	N.S.W.	B.f. VII, 381 M.fr. IX, 39.
SCHOENUS, Linné, coroll. gen. 2 (1737). (Chaetospora, Elynanthus, Tricostularia, Helothrix, Isoschoenus, Gymnochaete, Gymnoschoenus, Mesomelaena, Discopodium.)									
S. cruentus, F. v. M., fragm. IX, 36 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 357 M.fr. IX, 37.
S. Benthani, F. v. M.; S. compressus, Benth. Fl. Austr. VII, 357 (1878) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 357 M.fr. IX, 37.
S. lanatus, Labillard., Nov. Holl. pl. spec. I, 19, t. 20 (1804) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 357 M.fr. IX, 37.
S. curvifolius, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 358 M.fr. IX, 36.
S. capitatus, F. v. M., fragm. IX, 58 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 358 M.fr. IX, 37.
S. setifolius, Benth in Fl. Austr. VII, 359 (1878) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 359
S. Drummondii, Benth in Fl. Austr. VII, 359, non Steud. (1870) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 359 M.fr. IX, 37.
S. turbinatus, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	...	...	N.S.W.	...	B.f. VII, 359 M.fr. IX, 33.
S. barbatus, Boeckeler in Garcke, Linnaea XXXVIII, 277 (1874) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 360 M.fr. IX, 30.
S. flavus, Boeckeler in Garcke, Linnaea XXXVIII, 278 (1874) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 360 M.fr. IX, 30.
S. brevisetis, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 360 M.fr. IX, 30.
S. Armeria, Boeckeler in Garcke, Linnaea XXXVIII, 279 (1874) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 361 M.fr. IX, 30.
S. aphyllus, Boeckeler in Garcke, Linnaea XXXVIII, 280 (1874) ...	...	...	...	...	S.A.	V.	N.S.W.	...	B.f. VII, 361 M.fr. IX, 28.
S. imberbis, R. Brown, prodr. 231 (1810) ...	...	...	...	...	...	...	V. N.S.W.	...	B.f. VII, 361 M.fr. IX, 28.
S. ericetorum, R. Brown, prodr. 231 (1810) ...	...	...	...	...	...	...	N.S.W. Q.	...	B.f. VII, 362 M.fr. IX, 58
S. nitens, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	W.A.	S.A.	T. V.	N.S.W. Q.	B.f. VII, 362 M.fr. IX, 35.
S. cygneus, Nees in Lehmann, pl. Preiss. II, 81 (1846) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 363 M.fr. IX, 80.
S. minutulus, F. v. M., fragm. IX, 32 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 363 M.fr. IX, 32.
S. trachycarpus, F. v. M., fragm. IX, 33 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 363 M.fr. IX, 33.
S. Tepperi, F. v. M., fragm. XI, 106 (1880) ...	...	...	...	...	...	S.A.	...	...	M.fr. XI, 106.
S. nanus, F. v. M., fragm. IX, 36 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 364 M.fr. IX, 28.
S. pleiostomoneus, F. v. M., fragm. IX, 52 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 364 M.fr. IX, 52.
S. breviculmis, Benth in Fl. Austr. VII, 364 (1878) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 364
S. deformis, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	...	S.A.	...	...	B.f. VII, 364 M.fr. IX, 39.
S. unispiculatus, F. v. M. in Benth. Fl. Austr. VII, 365 (1878) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 365
S. obtusifolius, Boeckeler in Garcke, Linnaea XXXVIII, 281 (1874) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 366 M.fr. IX, 31.
S. grammatophyllum, F. v. M., fragm. IX, 31 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 366 M.fr. IX, 31.
S. asperocarpus, F. v. M., fragm. IX, 29 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 366 M.fr. IX, 29.
S. Moorei, Benth in Fl. Austr. VII, 367 (1878) ...	...	...	...	...	...	...	N.S.W.	...	B.f. VII, 367
S. villosus, R. Brown, prodr. 231 (1810) ...	...	...	...	...	...	...	N.S.W.	...	B.f. VII, 367 M.fr. IX, 28.
S. grandiflorus, F. v. M., fragm. IX, 30 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 367 M.fr. IX, 30.
S. distans; Chaetospora, F. v. M., fragm. IX, 35 (1875) ...	...	...	...	...	W.A.	...	...	...	M.fr. IX, 35.
S. calostachyus, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	...	...	N.S.W. Q.	...	B.f. VII, 368 M.fr. IX, 35.
S. scabripes, Benth in Fl. Austr. VII, 368 (1878) ...	...	...	...	...	...	...	...	Q.	B.f. VII, 368
S. multiglumis, Benth in Fl. Austr. VII, 368 (1878) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 368
S. efoliatus, F. v. M., fragm. IX, 32 (1875) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 369 M.fr. IX, 32.
S. acuminatus, R. Brown, prodr. 231 (1810) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 369 M.fr. IX, 80.
S. pedicellatus, Poiret, Encycl. méth. suppl. II, 251 (1811) ...	...	...	...	...	W.A.	...	...	...	B.f. VII, 369
S. fascicularis, Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842) W.A.	...	...	...	...	...	...	...	...	B.f. VII, 370
S. brevifolius, R. Brown, prodr. 231 (1810) ...	...	...	...	...	W.A.	S.A.	V.	N.S.W. Q.	B.f. VII, 370 M.fr. IX, 29, 80.
S. melanostachys, R. Brown, prodr. 231 (1810) ...	...	...	...	...	...	...	V. N.S.W. Q.	...	B.f. VII, 370 M.fr. IX, 29.
S. spartens, R. Brown, prodr. 231 (1810) ...	...	...	...	...	...	...	...	Q. N.A.	B.f. VII, 371
S. vaginatus, F. v. M. in Benth. Fl. Austr. VII, 371 (1878) ...	...	...	...	...	...	...	N.S.W. Q.	...	B.f. VII, 371

<i>S. falcatus</i> , R. Brown, prodr. 232 (1810)	—	—	—	—	—	Q.	N.A.	B.f.VII,372	M.fr.IX,29.
<i>S. punctatus</i> , R. Brown, prodr. 232 (1810)... ..	—	—	—	—	—	—	N.A.	B.f.VII,372	
<i>S. indutus</i> , F. v. M. in Benth. Fl. Austr. VII, 372 (1878) ...	W.A.	—	—	—	—	—	—	B.f.VII,372	
<i>S. bifidus</i> , Boeckeler in Garcke, Linnaea XXXVIII, 282 (1874) ...	W.A.	—	—	—	—	—	—	B.f.VII,373	M.fr.IX,38,80.
<i>S. apogon</i> , Roemer et Schultes, syst. veg. II, 77 (1817) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VII,373	M.fr.IX,56;X,120.
<i>S. odontocarpus</i> , F. v. M., fragm. IX, 32 (1875)	W.A.	—	—	—	—	—	—	B.f.VII,374	M.fr.IX,32.
<i>S. humilis</i> , Benth. Fl. Austr. VII, 374 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,374	
<i>S. sculptus</i> , Boeckeler in Garcke, Linnaea XXXVIII, 286 (1874) ...	W.A.	—	—	—	—	—	—	B.f.VII,374	M.fr.IX,30.
<i>S. axillaris</i> , Poir. encycl. méth. suppl. II, 251 (1811) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,375	M.fr.IX,34.
<i>S. tenellus</i> , Benth. Fl. Austr. VII, 375 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,375	
<i>S. natans</i> , F. v. M., fragm. IX, 36 (1875)	W.A.	—	—	—	—	—	—	B.f.VII,375	M.fr.IX,38.
<i>S. fluitans</i> , J. Hooker, Fl. Tasman. II, 81, t. 141 (1860)... ..	—	—	T.	—	—	—	—	B.f.VII,376	M.fr.IX,28.
<i>S. octandrus</i> , F. v. M., fragm. IX, 31 (1875)	W.A.	—	—	—	—	—	—	B.f.VII,377	M.fr.IX,31.
<i>S. capillaris</i> , Chaetosporea, F. v. M., fragm. IX, 377 (1875) ...	W.A.	S.A.	T.	V.	—	—	—	B.f.VII,377	M.fr.IX,34,80.
<i>S. stygius</i> , Poir. encycl. méth. suppl. II, 251 (1811) ...	W.A.	—	—	—	—	—	—	B.f.VII,378	M.fr.IX,36.
<i>S. deustus</i> , F. v. M.; Carpha deusta, R. Brown, prodr. 230 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,379	M.fr.IX,39.
<i>S. tetragonus</i> , Poir. encycl. méth. suppl. II, 251 (1811) ...	W.A.	—	—	—	—	—	—	B.f.VII,380	M.fr.IX,36.
<i>S. sphaerocephalus</i> , Poir. encycl. méth. suppl. II, 251 (1811)... ..	—	—	T.	V.	N.S.W.	—	—	B.f.VII,380	M.fr.IX,33.
<i>S. anceps</i> , Poir. encycl. méth. suppl. II, 251 (1811) ...	W.A.	—	—	—	—	—	—	B.f.VII,380	M.fr.IX,30.
<i>S. paludosus</i> , Poir. encycl. méth. suppl. II, 251 (1811) ...	—	—	—	—	N.S.W.	Q.	—	B.f.VII,382	M.fr.IX,35.
<i>S. pauciflorus</i> ; Lepidosperma, F. v. M., fragm. IX, 23 (1875) ...	—	—	—	—	V.	—	—	B.f.VII,383	M.fr.IX,23.
<i>S. Tricostularia</i> , F. v. M.; Tricostularia compressa, Nees in pl. Preiss. II, 83 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,383	M.fr.IX,37.
<i>S. Neesii</i> , F. v. M.; Tricostularia, Lehmann, pl. Preiss. II, 83 (1846) ...	W.A.	—	—	—	—	—	—	B.f.VII,383	M.fr.IX,37.
<i>S. fimbriatylloides</i> ; Chaetosporea, F. v. M., fragm. IX, 34 (1875) ...	—	—	—	—	—	—	N.A.	B.f.VII,384	M.fr.IX,34.

LEPIDOSPERMA, F. v. M., fragm. phytograph. Austral. IX, 34 (1875).

<i>L. tenuissima</i> , F. v. M., fragm. IX, 34 (1875)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,385	M.fr.IX,34.
--	---	------	----	----	--------	---	---	-------------	-------------

LEPIDOSPERMA, Labillardière, Nov. Holl. pl. spec. I, 14 (1804).

<i>L. gladiatum</i> , Labillardière, Nov. Holl. pl. spec. I, 15, t. 12 (1804) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,387	
<i>L. effusum</i> , Benth. Fl. Austr. VII, 387 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,387	
<i>L. rupestre</i> , Benth. Fl. Austr. VII, 388 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,388	
<i>L. elatius</i> , Labillardière, Nov. Holl. pl. spec. I, 15, t. 11 (1804) ...	—	S.A.	T.	V.	—	—	—	B.f.VII,388	M.fr.IX,25.
<i>L. tetraquetrum</i> , Nees in Lehm. pl. Preiss. II, 90 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,388	M.fr.IX,24.
<i>L. Oldfieldii</i> , J. Hooker, Fl. Tasman. II, 91, t. 146 (1860)	—	—	T.	—	—	—	—	B.f.VII,389	
<i>L. exaltatum</i> , R. Brown, prodr. 234 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.VII,389	
<i>L. longitudinale</i> , Labillardière, Nov. Holl. pl. spec. I, 16, t. 13 (1804) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,389	M.fr.IX,25.
<i>L. concavum</i> , R. Brown, prodr. 234 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VII,390	M.fr.IX,24.
<i>L. angustatum</i> , R. Brown, prodr. 235 (1810)	W.A.	—	—	—	—	—	—	B.f.VII,391	M.fr.IX,26,80.
<i>L. Drummondii</i> , Benth. Fl. Austr. VII, 391 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,391	
<i>L. Brunonianum</i> , Nees in Lehm. pl. Preiss. II, 92 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,392	
<i>L. tuberculatum</i> , Nees in Lehm. pl. Preiss. II, 90 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,392	M.fr.IX,26.
<i>L. resinosa</i> , F. v. M. in Benth. Fl. Austr. VII, 392 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,392	
<i>L. viscidum</i> , R. Brown, prodr. 234 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.f.VII,393	
<i>L. costale</i> , Nees in Lehm. pl. Preiss. II, 92 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,393	
<i>L. laterale</i> , R. Brown, prodr. 234 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VII,393	
<i>L. congestum</i> , R. Brown, prodr. 234 (1810)	—	S.A.	—	—	—	—	—	B.f.VII,394	
<i>L. globosum</i> , Labillardière, Nov. Holl. pl. spec. I, 16, t. 14 (1804) ...	—	S.A.	T.	V.	—	—	—	B.f.VII,394	
<i>L. lineare</i> , R. Brown, prodr. 235 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,395	M.fr.IX,26.
<i>L. aphyllum</i> , R. Brown, prodr. 235 (1810)	W.A.	—	—	—	—	—	—	B.f.VII,395	
<i>L. gracile</i> , R. Brown, prodr. 235 (1810)	W.A.	—	—	—	—	—	—	B.f.VII,395	
<i>L. semiteres</i> , F. v. M. in Garcke, Linnaea XXXVIII, 327 (1874) ...	—	S.A.	—	V.	—	—	—	B.f.VII,395	
<i>L. canescens</i> , Boeckeler in Garcke, Linnaea XXXVIII, 330 (1874) ...	—	S.A.	—	V.	N.S.W.	—	—	B.f.VII,396	M.fr.IX,24.
<i>L. pubisquamum</i> , Steudel, syn. pl. glum. II, 158 (1855)	W.A.	—	—	—	—	—	—	B.f.VII,396	M.fr.IX,27.
<i>L. scabrum</i> , Nees in Lehm. pl. Preiss. II, 92 (1846)	W.A.	—	—	—	—	—	—	B.f.VII,397	M.fr.IX,27.
<i>L. tenue</i> , Benth. Fl. Austr. VII, 397 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,397	
<i>L. leptostachyum</i> , Benth. Fl. Austr. VII, 397 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,397	
<i>L. leptophyllum</i> , Benth. Fl. Austr. VII, 398 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,398	
<i>L. tortuosum</i> , F. v. M., general Report 9 (1855)	—	—	—	—	V.	—	—	B.f.VII,398	M.fr.IX,23.
<i>L. flexuosum</i> , R. Brown, prodr. 235 (1810)	—	—	—	—	N.S.W.	—	—	B.f.VII,398	
<i>L. filiforme</i> , Labillardière, Nov. Holl. pl. spec. I, 17, t. 15 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,398	M.fr.IX,27.
<i>L. striatum</i> , R. Brown, prodr. 235 (1810)	W.A.	—	—	—	—	—	—	B.f.VII,399	M.fr.IX,26,27.
<i>L. Neesii</i> , Kunth, enum. II, 319 (1837)	—	—	—	—	V.	N.S.W.	—	B.f.VII,399	M.fr.IX,24,27.
<i>L. carphoides</i> , F. v. M. in Benth. Fl. Austr. VII, 400 (1878) ...	W.A.	S.A.	—	V.	—	—	—	B.f.VII,400	

CLADIUM, P. Browne, Civ. and Nat. Hist. of Jamaica. 114 (1756). (Gahnia, Lampocarya, Melachne, Epiandra, Didymonema, Baumea, Chapelliera, Morelotia.)

<i>C. Mariscus</i> , R. Brown, prodr. 236 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f.VII,402	M.fr.IX,14,57.
<i>C. insulare</i> , Benth. Fl. Austr. 403 (1878)	—	—	—	—	N.S.W.	—	—	B.f.VII,403	
<i>C. articulatum</i> , R. Brown, prodr. 237 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f.VII,403	M.fr.IX,14,57.
<i>C. arthrophyllum</i> , F. v. M., fragm. IX, 14 (1875)	W.A.	—	—	—	—	—	—	B.f.VII,403	M.fr.IX,14.
<i>C. glomeratum</i> , R. Brown, prodr. 237 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.VII,404	M.fr.IX,15,57.
<i>C. Preissii</i> , F. v. M. in Benth. Fl. Austr. VII, 405 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,405	M.fr.IX,13,15,56.
<i>C. laxum</i> , Benth. Fl. Austr. VII, 405 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,405	
<i>C. riparium</i> , Benth. Fl. Austr. VII, 405 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,405	
<i>C. teretifolium</i> , R. Brown, prodr. 237 (1810)	—	—	—	—	N.S.W.	Q.	—	B.f.VII,406	M.fr.IX,15,56.
<i>C. tetraquetrum</i> , J. Hooker, Fl. Tasman. II, 95, t. 149 (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,406	
<i>C. schoenoides</i> , R. Brown, prodr. 237 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f.VII,407	
<i>C. Gunnii</i> , J. Hooker, Fl. Tasman. II, 95, t. 148 (1860)	—	S.A.	T.	V.	N.S.W.	—	—	B.f.VII,407	M.fr.IX,15,57.
<i>C. junceum</i> , R. Brown, prodr. 237 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f.VII,408	M.fr.IX,16,57.
<i>C. vaginale</i> , Benth. Fl. Austr. VII, 409 (1878)	W.A.	—	—	—	—	—	—	B.f.VII,408	
<i>C. elynanthoides</i> , F. v. M., fragm. IX, 31 (1875)	W.A.	—	—	—	—	—	—	B.f.VII,409	M.fr.IX,31.

C. Filum, R. Brown, prodr. 237 (1810)	—	S.A.	T. V.	—	—	—	B.f. VII, 409	M.fr. IX, 14, 57.	
C. trifidum, F. v. M. in Pap. R. S. Tasm. 117 (1878)	W.A.	S.A.	T. V.	N.S.W.	—	—	B.f. VII, 413		
C. decompositum, R. Brown, prodr. 237 (1810)	W.A.	—	—	—	—	—	B.f. VII, 417	M.fr. IX, 18.	
C. microstachyum, F. v. M.; Gahnia, Benth. Fl. Austr. VII, 414 (1878)	—	—	V.	N.S.W.	—	—	B.f. VII, 414		
C. polyphyllum, F. v. M.; Gahnia, Benth. Fl. Austr. VII, 415 (1878)	W.A.	—	—	—	—	—	B.f. VII, 415		
C. anicistrophyllosum, F. v. M. in Benth. Fl. Austr. VII, 415 (1878)	W.A.	—	—	—	—	—	B.f. VII, 415		
C. lanigerum, R. Brown, prodr. 237 (1810)	W.A.	S.A.	—	—	—	—	B.f. VII, 415	M.fr. IX, 14, 57.	
C. aristatum, F. v. M. in Benth. Fl. Austr. 416 (1878)	W.A.	—	—	—	—	—	B.f. VII, 416		
C. deustum, R. Brown, prodr. 237 (1810)	W.A.	S.A.	—	—	—	—	B.f. VII, 416	M.fr. IX, 14.	
C. Radula, R. Brown, prodr. 237 (1810)	—	—	T. V.	—	—	—	B.f. VII, 417	M.fr. IX, 13, 56.	
C. Sieberi, F. v. M., fragm. IX, 14 (1875)	—	—	—	N.S.W.	—	—	B.f. VII, 414	M.fr. IX, 14, 57.	
C. tetragonocarpum, F. v. M.; Gahnia, Boeckeler in Linnaea XXXVIII, 347 (1874)	—	—	—	—	—	—	B.f. VII, 418		
C. melanocarpum, F. v. M., fragm. IX, 13 (1875)	—	—	T. V.	N.S.W.	—	—	B.f. VII, 414	M.fr. IX, 13, 56.	
C. psittacorum, F. v. M., fragm. IX, 13 (1875)	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 418	M.fr. IX, 13, 56.	
C. asperum, F. v. M., fragm. IX, 12 (1875)	—	—	—	N.S.W.	Q.	—	B.f. VII, 412	M.fr. IX, 12, 56.	
C. xanthocarpum, F. v. M., fragm. IX, 13 (1875)	—	—	—	N.S.W.	—	—	B.f. VII, 418	M.fr. IX, 13, 56.	
CAUSTIS, R. Brown, prodr. 239 (1810). (Eurostorrhiza.)									
C. pentandra, R. Brown, prodr. 240 (1810)	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 420	M.fr. IX, 19, 58.	
C. flexuosa, R. Brown, prodr. 239 (1810)	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 421	M.fr. IX, 58.	
C. recurvata, Sprengel, syst. cur. post. 26 (1827)	—	—	—	N.S.W.	—	—	B.f. VII, 421	M.fr. IX, 19.	
C. restiacea, F. v. M. in Benth. Fl. Austr. VII, 421 (1878)	—	—	V.	N.S.W.	—	—	B.f. VII, 421	M.fr. IX, 19.	
C. dioica, R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	B.f. VII, 422	M.fr. IX, 19, 58.	
ARTHROSTYLIS, R. Brown, prodr. 229 (1810). (Arthrostyles.)									
A. aphylla, R. Brown, prodr. 229 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 423	M.fr. IX, 9, 55.
REEDIA, F. v. M., fragm. I, 240, t. 10 (1859).									
R. spathacea, F. v. M., fragm. I, 240, t. 10 (1859)	W.A.	—	—	—	—	—	B.f. VII, 423	M.fr. IX, 15.	
EUANDRA, R. Brown, prodr. 239 (1810). (Evandra.)									
E. aristata, R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	B.f. VII, 424	M.fr. IX, 18.	
E. pauciflora, R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	B.f. VII, 425	M.fr. IX, 18.	
SCLERIA, Bergius in Svensk. Vet. Acad. Handl. 149, t. 4 (1765). (Diplacrum, Hypoporum, Sphaeropus.)									
S. caricina, Benth. Fl. Austr. VII (1878)	—	—	—	—	—	Q.	—	B.f. VII, 426	M.fr. IX, 58.
S. pygmaea, R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 427	M.fr. IX, 22, 58.
S. rugosa, R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 428	M.fr. IX, 22, 58.
S. laxa, R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 428	M.fr. IX, 21.
S. Brownii, Kunth, enum. II, 349 (1837)	—	—	—	—	—	Q.	N.A.	B.f. VII, 429	M.fr. IX, 21, 58.
S. lithosperma, Willdenow, spec. plant. IV, 316 (1805)	—	—	—	—	—	Q.	N.A.	B.f. VII, 429	M.fr. IX, 21, 58.
S. tessellata, Willdenow, spec. plant. IV, 315 (1805)	—	—	—	—	—	Q.	—	B.f. VII, 430	
S. margaritifera, Willdenow, spec. plant. IV, 312 (1805)	—	—	—	—	—	Q.	—	B.f. VII, 430	M.fr. IX, 21, 58.
S. Graeffeana, Boeckeler in Regensb. Flora, 121 (1875)	—	—	—	—	—	Q.	N.A.	B.f. VII, 431	
S. hebecarpa, Nees in Wight, Contrib. to Bot. of Ind. 117 (1834)	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 431	M.fr. IX, 21, 58.	
S. Chinensis, Kunth, enum. II, 357 (1837)	—	—	—	—	—	Q.	—	B.f. VII, 431	M.fr. IX, 20, 58.
S. oryzoides, Presl, reliq. Haenk. I, 201 (1830)	—	—	—	—	—	N.A.	B.f. VII, 432		
S. sphacelata, F. v. M., fragm. IX, 20 (1875)	—	—	—	N.S.W.	Q.	—	B.f. VII, 432	M.fr. IX, 20.	
UNCINIA, Persoon, synopsis. plant. I, 534 (1807).									
U. tenella, R. Brown, prodr. 241 (1810)	—	—	T. V.	—	—	—	B.f. VII, 433	M.fr. VIII, 151.	
U. compacta, R. Brown, prodr. 241 (1810)	—	—	T. V.	—	—	—	B.f. VII, 434	M.fr. VIII, 152.	
U. riparia, R. Brown, prodr. 241 (1810)	—	—	T. V.	N.S.W.	—	—	B.f. VII, 434	M.fr. VIII, 152.	
U. debilior, F. v. M., fragm. VIII, 151 (1874)	—	—	—	N.S.W.	—	—	B.f. VII, 435	M.fr. VIII, 151.	
CAREX, Ruppius, Fl. Jenens. 306 (1718).									
C. cephalotes, F. v. M. in Transact. phil. Soc. Vict. I, 110 (1854)	—	—	V.	N.S.W.	—	—	B.f. VII, 437	M.fr. VIII, 251.	
C. acicularis, Boott in J. Hook. Fl. N. Zel. I, 280, t. 63 (1853)	—	—	T. V.	N.S.W.	—	—	B.f. VII, 437	M.fr. VIII, 251.	
C. capillacea, Boott, Illustr. of the gen. Carex I, 44, t. 110 (1858)	—	—	—	N.S.W.	—	—	B.f. VII, 438	M.fr. IX, 191.	
C. inversa, R. Brown, prodr. 242 (1810)	W.A.	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 438	M.fr. VIII, 252.	
C. canescens, Linné, spec. plant. 974 (1753)	—	—	V.	N.S.W.	—	—	B.f. VII, 439	M.fr. VIII, 253.	
C. echinata, Murray, prodr. design. stirp. Goett. 76 (1770)	—	—	V.	N.S.W.	—	—	B.f. VII, 439	M.fr. VIII, 252.	
C. hypandra, F. v. M., fragm. VIII, 259 (1874)	—	—	V.	N.S.W.	—	—	B.f. VII, 439	M.fr. VIII, 259.	
C. chlorantha, R. Brown, prodr. 242 (1810)	—	S.A.	T. V.	N.S.W.	—	—	B.f. VII, 440	M.fr. VIII, 256.	
C. paniculata, Linné, amoen. acad. IV, 294 (1759)	W.A.	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 440	M.fr. VIII, 256.	
C. declinata, Boott, Ill. of the gen. Carex IV, 17, t. 580 (1867)	—	—	V.	N.S.W.	Q.	—	B.f. VII, 441	M.fr. VIII, 257.	
C. tereticaulis, F. v. M., fragm. VIII, 256 (1874)	W.A.	S.A.	T. V.	N.S.W.	—	—	B.f. VII, 441	M.fr. VIII, 256.	
C. fissilis, Boott, Ill. of the gen. Carex II, 86, t. 245 (1860)	—	—	—	—	—	Q.	B.f. VII, 441	M.fr. VIII, 249; IX, 80.	
C. gracilis, R. Brown, prodr. 242 (1810)	—	—	—	N.S.W.	Q.	—	B.f. VII, 442	M.fr. VIII, 250.	
C. contracta, F. v. M., fragm. VIII, 258 (1874)	—	—	—	N.S.W.	—	—	B.f. VII, 442	M.fr. VIII, 258.	
C. Gaudichaudiana, Kunth, enum. plant. II, 417 (1837)	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 442	M.fr. VIII, 257.	
C. acuta, Linné, spec. plant. 978 (1753)	—	—	V.	N.S.W.	Q.	—	B.f. VII, 443	M.fr. VIII, 259.	
C. lobolepis, F. v. M., fragm. VIII, 258 (1874)	—	—	—	N.S.W.	—	—	B.f. VII, 443	M.fr. VIII, 258.	
C. flava, Linné, spec. plant. 975 (1753)	—	—	T.	—	—	—	B.f. VII, 444	M.fr. VIII, 250.	
C. Buxbaumii, Wahlenberg in Act. Holmiens. 139 (1803)	—	—	V.	N.S.W.	—	—	B.f. VII, 444	M.fr. VIII, 252.	
C. pumila, Thunberg, Fl. Japon. 38 (1784)	—	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 445	M.fr. VIII, 251.	
C. breviculmis, R. Brown, prodr. 242 (1810)	—	S.A.	T. V.	N.S.W.	—	—	B.f. VII, 445	M.fr. VIII, 255.	
C. Preissii, Nees in Lehm. pl. Preiss. II, 94 (1846)	W.A.	—	—	—	—	—	B.f. VII, 446	M.fr. VIII, 251.	
C. Gunniana, Boott in Transact. Linn. Soc. XX, 143 (1851)	—	S.A.	T. V.	N.S.W.	—	—	B.f. VII, 446	M.fr. VIII, 251.	
C. Bichenoviana, Boott in J. Hook. Fl. Tasm. II, 101 (1860)	—	—	T.	—	—	—	B.f. VII, 446		
C. maculata, Boott in Transact. Linn. Soc. XX, 128 (1851)	—	—	—	N.S.W.	Q.	—	B.f. VII, 447	M.fr. VIII, 258.	
C. Brownii, Tuckerman, enum. meth. Caric. 21 (1849)	—	—	—	N.S.W.	—	—	B.f. VII, 447	M.fr. IX, 250.	
C. alsophila, F. v. M., fragm. VIII, 257 (1874)	—	—	V.	—	—	—	B.f. VII, 447	M.fr. VIII, 257.	
C. longifolia, R. Brown, prodr. 242 (1810)	—	—	T. V.	N.S.W.	Q.	—	B.f. VII, 448	M.fr. VIII, 250.	
C. Pseudo-Cyperus, Linné, spec. plant. 978 (1753)	W.A.	S.A.	T. V.	N.S.W.	Q.	—	B.f. VII, 448	M.fr. VIII, 249.	

GRAMINEAE.

Haller, enum. stirp. Helv. I, 203 (1742).

ERIOCHLOA, Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 94, t. 30 (1815). (Helopus.)

- E. punctata, Hamilton, prodr. pl. Ind. occ. 5 (1825) ... — — — — V. N.S.W. Q. N.A. B.f. VII, 462 M.fr. VI, 85; VIII, 126
 E. annulata, Kunth, Graminées I, 30 (1832) ... — — — — S.A. — — — — N.S.W. Q. — B.f. VII, 463 M.fr. VI, 85.

PASPALUM, Linné, syst. nat. ed. decim. 855 (1759).

- P. serobiculatum, Linné, mantiss. 29 (1767) ... — — — — N.S.W. Q. N.A. B.f. VII, 360 M.fr. VIII, 156.
 P. distichum, Linné, syst. nat. ed. decim. 855 (1759) ... — — — — W.A. — — — — N.S.W. Q. — B.f. VII, 460 M.fr. VIII, 156.
 P. brevifolium, Fluegge, gram. monogr. 150 (1810) ... — — — — N.S.W. Q. N.A. B.f. VII, 461 M.fr. VIII, 195.
 P. minutiflorum, Steudel, sym. glum. I, 17 (1855) ... — — — — — Q. — B.f. VII, 461

PANICUM, Tournefort, inst. 515, t. 293 (1700), from Plinius. (Isachne, Digitaria, Echinochloa, Coridochloa, Chamaeraphis.)

- P. coenicolum, F. v. M. in Transact. Vict. Inst. 45 (1854) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. — B.f. VII, 467
 P. divaricatissimum, R. Brown, prodr. 192 (1810) ... — — — — S.A. — — — — V. N.S.W. Q. — B.f. VII, 467 M.fr. VIII, 154; X, 76.
 P. macractinum, Benth. Fl. Austr. VII, 468 (1878) ... — — — — — — — — N.S.W. Q. — B.f. VII, 468
 P. papposum, R. Brown, prodr. 192 (1810) ... — — — — — — — — — N.A. B.f. VII, 468 M.fr. VIII, 155, 196.
 P. sanguinale, Linné, spec. plant. 57 (1753) ... — — — — W.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 469 M.fr. VIII, 153.
 P. ctenanthum, F. v. M., fragm. VIII, 153 (1874) ... — — — — — — — — — N.A. B.f. VII, 469 M.fr. VIII, 153.
 P. stenostachyum, Benth. Fl. Austr. VII, 470 (1878) ... — — — — — — — — — N.A. B.f. VII, 470
 P. tenuissimum, Benth. Fl. Austr. VII, 470 (1878) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 470
 P. parviflorum, R. Brown, prodr. 192 (1810) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 470 M.fr. VIII, 196; XI, 129.
 P. Baileyi, Benth. Fl. Austr. VII, 470 (1878) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 471
 P. gibbosum, R. Brown, prodr. 193 (1810) ... — — — — — — — — — Q. N.A. B.f. VII, 471 M.fr. VIII, 155; XI, 129.
 P. leucophaeum, Humb., Bonpl. & Kunth, nov. gen. et spec. I, 97 (1815) ... — — — — S.A. — — — — V. N.S.W. Q. — B.f. VII, 472 M.fr. VIII, 155.
 P. semialatum, R. Brown, prodr. 192 (1810) ... — — — — — — — — — N.S.W. Q. N.A. B.f. VII, 472 M.fr. VIII, 196.
 P. rarum, R. Brown, prodr. 189 (1810) ... — — — — — — — — — — N.A. B.f. VII, 473
 P. argenteum, R. Brown, prodr. 190 (1810) ... — — — — — — — — — — N.A. B.f. VII, 473
 P. holosericeum, R. Brown, prodr. 190 (1810) ... — — — — — — — — — Q. N.A. B.f. VII, 473 M.fr. VIII, 155.
 P. flavidum, Retzius, observ. IV, 15 (1787) ... — — — — — — — — — N.S.W. Q. N.A. B.f. VII, 474 M.fr. VIII, 189.
 P. gracile, R. Brown, prodr. 190 (1810) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 475 M.fr. VIII, 189.
 P. prostratum, Lamarck, illustr. I, 171 (1791) ... — — — — — — — — — Q. N.A. B.f. VII, 476 M.fr. VIII, 191.
 P. helopus, Trinius in Spreng. Neue Entd. II, 84 (1821) ... — — — — S.A. — — — — N.S.W. Q. N.A. B.f. VII, 476
 P. Gilesii, Benth. Fl. Austr. VII, 477 (1878) ... — — — — S.A. — — — — — Q. N.A. B.f. VII, 477 M.fr. XI, 129.
 P. piligerum, F. v. M. in Benth. Fl. Austr. VII, 477 (1878) ... — — — — — — — — — — N.A. B.f. VII, 477
 P. polyphyllum, R. Brown, prodr. 190 (1810) ... — — — — — — — — — — N.A. B.f. VII, 477 M.fr. VIII, 194.
 P. distachyon, Linné, mantiss. alt. pl. 138 (1771) ... — — — — S.A. — — — — N.S.W. Q. — B.f. VII, 478 M.fr. VIII, 194.
 P. reversum, F. v. M., fragm. VIII, 152 (1874) ... — — — — W.A. S.A. — — — — N.S.W. Q. — B.f. VII, 478 M.fr. X, 76.
 P. colonum, Linné, syst. nat. ed. decim. 5 (1758) ... — — — — — — — — — — N.A. B.f. VII, 478
 P. Crus Galli, Linné, spec. plant. 56 (1753) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 479 M.fr. VIII, 198.
 P. myosuroides, R. Brown, prodr. 189 (1810) ... — — — — — — — — — Q. N.A. B.f. VII, 480
 P. Indicum, Linné, mantiss. pl. alt. 184 (1771) ... — — — — — — — — — N.S.W. Q. N.A. B.f. VII, 480 M.fr. VIII, 197; XI, 129.
 P. Myurus, Lamarck, illustr. I, 172 (1791) ... — — — — — — — — — Q. — B.f. VII, 480 M. X, 114.
 P. foliosum, R. Brown, prodr. 191 (1810) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 481 M.fr. VIII, 194.
 P. adpersum, Trinius, spec. gram. t. 169 (1830) ... — — — — S.A. — — — — N.S.W. Q. — B.f. VII, 481
 P. inaequale, F. v. M., fragm. VIII, 189 (1874) ... — — — — — — — — — Q. — B.f. VII, 482 M.fr. XI, 129. [129.
 P. uncinulatum, R. Brown, prodr. 191 (1810) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 482 M.fr. VIII, 195; X, 76; XI,
 P. majusculum, F. v. M. in Benth. Fl. Austr. VII, 482 (1878) ... — — — — — — — — — N.A. B.f. VII, 482
 P. pauciflorum, R. Brown, prodr. 483 (1810) ... — — — — S.A. — — — — — Q. N.A. B.f. VII, 483 M.fr. VIII, 193.
 P. semitonsum, F. v. M. in Benth. Fl. Austr. VII, 483 (1878) ... — — — — — — — — — N.A. B.f. VII, 483
 P. antidotale, Retzius, observ. bot. IV, 17 (1878) ... — — — — — — — — — N.A. B.f. VII, 483
 P. repens, Linné, spec. pl. ed. sec. 87 (1762) ... — — — — — — — — — V. N.S.W. Q. N.A. B.f. VII, 484
 P. Buncei, F. v. M. in Benth. Fl. Austr. VII, 487 (1878) ... — — — — — — — — — Q. — B.f. VII, 487
 P. capillipes, Benth. Fl. Austr. VII, 484 (1878) ... — — — — — — — — — N.A. B.f. VII, 484 M.fr. VIII, 193.
 P. pygmaeum, R. Brown, prodr. 191 (1810) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 484 M.fr. VIII, 193.
 P. brevifolium, Linné, spec. plant. 59 (1753) ... — — — — — — — — — Q. N.A. B.f. VII, 485 M.fr. VIII, 193; XI, 129.
 P. hermaphroditum, Steudel, syn. glum. I, 67 (1855) ... — — — — — — — — — Q. — B.f. VII, 485
 P. marginatum, R. Brown, prodr. 190 (1810) ... — — — — — — — — — V. N.S.W. Q. — B.f. VII, 485 M.fr. VIII, 190.
 P. lachnophyllum, Benth. Fl. Austr. VII, 486 (1878) ... — — — — — — — — — Q. — B.f. VII, 486
 P. obseptum, Trinius, gram. panic. diss. alt. 149 (1826) ... — — — — — — — — — N.S.W. — B.f. VII, 486
 P. bicolor, R. Brown, prodr. 191 (1810) ... — — — — — — — — — N.S.W. Q. — B.f. VII, 487
 P. melananthum, F. v. M. in Transact. Vict. Inst. 47 (1854) ... — — — — — — — — — V. N.S.W. Q. — B.f. VII, 488 M.fr. VIII, 192.
 P. effusum, R. Brown, prodr. 191 (1810) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. — B.f. VII, 488 M.fr. VIII, 191.
 P. Mitchellii, Benth. Fl. Austr. VII, 489 (1878) ... — — — — S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 489 M.fr. X, 76.
 P. decompositum, R. Brown, prodr. 191 (1810) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 489 M.fr. VI, 86; VIII, 191.
 P. trachyrhachis, Benth. Fl. Austr. VII, 490 (1878) ... — — — — — — — — — N.S.W. Q. N.A. B.f. VII, 490 M.fr. VIII, 192; XI, 129.
 P. prolutum, F. v. M. in Transact. Vict. Inst. 46 (1854) ... — — — — S.A. — — — — V. N.S.W. Q. — B.f. VII, 490 M.fr. VIII, 192.
 P. spinosens, R. Brown, prodr. 193 (1810) ... — — — — W.A. S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 499 M.fr. VIII, 197; XI, 129.
 P. paradoxum, R. Brown, prodr. 193 (1810) ... — — — — — — — — — V. N.S.W. — B.f. VII, 499 M.fr. VIII, 197.
 P. Chamaeraphis, Trinius in Mem. acad. Petersb. ser. 6, III, 217 (1835) ... — — — — — — — — — N.A. B.f. VII, 500
 P. atro-virens, Trinius in Spreng. neue Entdeckung. II, 89 (1821) ... — — — — — — — — — V. N.S.W. Q. — B.f. VII, 625 M.fr. VIII, 193; XI, 129.
 P. Myosotis, Steudel, syn. glum. I, 96 (1855) ... — — — — — — — — — Q. — B.f. VII, 625 M.fr. VIII, 193.

OPLISMENUS, Palisot, Flore d'Oware et de Benin. II, 14 (1807). (Orthopogon.)

- O. compositus, Palisot, agrostogr. 54 (1812) ... — — — — — — — — — V. N.S.W. Q. — B.f. VII, 491 M.fr. VIII, 199.

SETARIA, Palisot, Essai d'une nouv. Agrostogr. 51, t. 13 (1812).

- S. glauca, Palisot, Agrostogr. 51, t. 13 (1812) ... — — — — S.A. — — — — V. N.S.W. Q. N.A. B.f. VII, 492 M.fr. VIII, 110.
 S. macrostachya, Humb., Bonpl. & Kunth, nov. gen. et spec. I, 110 (1815) ... — — — — S.A. — — — — N.S.W. Q. N.A. B.f. VII, 493 M.fr. VIII, 110; XI, 129.
 S. viridis, Palisot, Agrostogr. 51, t. 13 (1812) ... — — — — W.A. S.A. — — — — — — — — — N.A. B.f. VII, 494
 S. verticillata, Palisot, Agrostogr. 51 (1812) ... — — — — W.A. — — — — — — — — — N.A. B.f. VII, 494 M.fr. VIII, 110.

- PENNISETUM, L. C. Richard in Persoon, synops. plant. I. 72 (1805). (Gymnothrix, Plagiosetum.)
P. compressum, R. Brown, prodr. 195 (1810) ... — — — — N.S.W. Q. — B.f. VII, 495 M.fr. VIII, 109.
P. Arnhemicum, F. v. M., fragm. VIII, 109 (1874) ... — — — — — N.A. B.f. VII, 496 M.fr. VIII, 109.
P. refractum, F. v. M., fragm. VIII, 109 (1874) ... — S.A. — — — — Q. N.A. B.f. VII, 495 M.fr. III, 147; IX, 194.
- CENCHRUS, Linné, coroll. gen. 20 (1737).
C. australis, R. Brown, prodr. 196 (1810) ... — — — — N.S.W. Q. — B.f. VII, 497 M.fr. VIII, 107; XI, 129.
C. inflexus, R. Brown, prodr. 195 (1810) ... — — — — — N.A. B.f. VII, 497 M.fr. VIII, 107.
C. elymoides, F. v. M., fragm. VIII, 107 (1874) ... — — — — — Q. N.A. B.f. VII, 497 M.fr. VIII, 107.
- XEROCHLOA, R. Brown, prodr. fl. Nov. Holl. 196 (1810).
X. imberbis, R. Brown, prodr. 197 (1810) ... — — — — — N.A. B.f. VII, 501 M.fr. VIII, 117.
X. barbata, R. Brown, prodr. 197 (1810) ... — — — — — N.A. B.f. VII, 501 M.fr. VIII, 117.
X. laniflora, Benth. Fl. Austr. VII, 502 (1878) ... — — — — — N.A. B.f. VII, 502
- THUAREA, Persoon, synops. plant. I, 110 (1805). (Thouarsea.)
T. sarmentosa, Persoon, synops. I, 110 (1805) ... — — — — — Q. N.A. B.f. VII, 502 M.fr. VI, 86.
- SPINIFEX, Linné, syst. nat. ed. XII, 757 (1767).
S. hirsutus, Labillard., Nov. Holl. pl. spec. II, 81, t. 230 (1806) W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 503 M.fr. VIII, 138.
S. longifolius, R. Brown, prodr. 198 (1810) ... — — — — — W.A. — — — — — Q. N.A. B.f. VII, 504 M.fr. VIII, 139. [IX, 195.
S. paradoxus, Benth. in Hook. icon. plant. t. 1243 (1877) ... — S.A. — V. N.S.W. Q. — B.f. VII, 504 M.fr. VI, 86; VIII, 117, 199;
- LEPTASPIS, R. Brown, prodr. fl. Nov. Holl. 211 (1810).
L. Banksii, R. Brown, prodr. 211 (1810) ... — — — — — Q. — B.f. VII, 548 M.fr. VIII, 116.
- PEROTIS, Aiton, hort. Kewens. I, 85 (1789).
P. rara, R. Brown, prodr. 172 (1810) ... — S.A. — — N.S.W. Q. N.A. B.f. VII, 509 M.fr. VIII, 115.
- POLYPOGON, Desfontaines, flor. Atlantic. I, 66 (1798).
P. fugax, Nees in Steudel, syn. glum. I, 184 (1855) ... — — — — — W.A. — — — — — B.f. VII, 547
P. tenellus, R. Brown, prodr. 173 (1810) ... — — — — — W.A. — — — — — B.f. VII, 547 M.fr. VIII, 114.
- ARUNDINELLA, Raddi, agrostograph. Brasil. 37 (1823).
A. Nepalensis, Trinius, spec. gram. t. 268 (1833) ... — — — — — N.S.W. Q. N.A. B.f. VII, 545 M.fr. VIII, 139.
A. Schultzii, Benth. Fl. Austr. VII, 545 (1878) ... — — — — — — N.A. B.f. VII, 545
- TRAGUS, Haller, hist. stirp. Helvet. n. 1413 (1768). (Lappago.)
T. racemosus, Haller, stirp. Helv. n. 1413 (1768) ... — S.A. — V. N.S.W. Q. N.A. B.f. VII, 507 M.fr. VIII, 107.
- NEURACHNE, R. Brown, prodr. fl. Nov. Holl. 196 (1810). [129.
N. alopecuroides, R. Brown, prodr. 196 (1810) ... — — — — — W.A. S.A. — V. N.S.W. — — B.f. VII, 507 M.fr. VI, 86; VIII, 200; XI,
N. Mitchelliana, Nees in Hook. Lond. Journ. of Bot. II, 410 (1843) — S.A. — V. N.S.W. — — B.f. VII, 508 M.fr. VI, 86; VIII, 200.
N. Munroi, F. v. M., fragm. VIII, 200 (1874) ... — S.A. — V. N.S.W. — — B.f. VII, 508 M.fr. V, 204; XI, 129.
- ZOYSIA, Willdenow, Neue Schrift. nat. Freunde zu Berl. III, 440 (1801).
Z. pungens, Willdenow, Neue Schrift. nat. Fr. Berl. III, 440 (1801) — — T. V. N.S.W. Q. — B.f. VII, 506 M.fr. VIII, 116.
- IMPERATA, Cyrillo, plant. rar. Neap. II, t. 11 (1792).
I. arundinacea, Cyrillo, plant. rar. Neap. II, t. 11 (1792) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 536 M.fr. VIII, 126.
- ERIANTHUS, L. C. Richard in Michaux, fl. bor. Amer. I, 54 (1803). (Pollinia, Eulalia, Pogonatherum, Leptatherum, Saccharum partly.)
E. articulatus, F. v. M., fragm. VIII, 118 (1874) ... — — — — — Q. N.A. B.f. VII, 525 M.fr. X, 76.
E. irritans, Kunth, Graminées I, 160 (1832) ... — — — — — Q. N.A. B.f. VII, 526 M.fr. VIII, 118.
E. fulvus, Kunth, Graminées I, 160 (1832) ... — — — — — W.A. S.A. — — N.S.W. Q. N.A. B.f. VII, 526 M.fr. VIII, 118.
E. Roxburghii, F. v. M., fragm. VIII, 117 (1874) ... — — — — — Q. — B.f. VII, 527 M.fr. VIII, 117.
E. Mackinlayi, Pollinia, F. v. M. in Benth. fl. Austr. VII, 527 (1878) — — — — — — N.A. B.f. VII, 527 M.fr. VIII, 118.
- DIMERIA, R. Brown, prodr. fl. Nov. Holl. 204 (1810).
D. acinaciformis, R. Brown, prodr. 204 (1810) ... — — — — — Q. — B.f. VII, 523
D. tenera, Trinius in Mém. de l'Acad. de Petersb. six. sér. II, 225 (1833) ... — — — — — Q. N.A. B.f. VII, 523 M.fr. VII, 104; XI, 129.
- ARTHAXON, Palisot, Agrostograph. 111, t. XI, f. 6 (1812). (Batratherum.)
A. ciliare, Palisot, Agrostogr. 111, t. 11 (1812) ... — — — — — N.S.W. Q. — B.f. VII, 524 M.fr. VIII, 119; X, 76.
- ELIONURUS, Willdenow, spec. plant. IV, 941 (1805).
E. citreus, Munro in Benth. Fl. Austral. VII, 510 (1878) ... — — — — — Q. — B.f. VII, 510 M.fr. XI, 129.
- LEPTURUS, R. Brown, prodr. fl. Nov. Holl. 207 (1810).
L. incurvatus, Trinius, fundam. agrostogr. 123 (1820) ... — S.A. — V. N.S.W. — — B.f. VII, 668 M.fr. VIII, 117.
L. cylindricus, Trinius, fundam. agrostogr. 123 (1820) ... — — — — — W.A. S.A. — V. N.S.W. — — B.f. VII, 668
L. repens, R. Brown, prodr. 207 (1810) ... — — — — — Q. N.A. B.f. VII, 668 M.fr. VI, 86.
- ROTTBOELLIA, Linné, fil. gram. gen. 22 (1779). (Coelorrhachis.)
R. formosa, R. Brown, prodr. 206 (1810) ... — — — — — Q. N.A. B.f. VII, 513 M.fr. VI, 85; XI, 129.
R. exaltata, Linné, fil. suppl. 114 (1781) ... — — — — — Q. N.A. B.f. VII, 513
R. muricata, Retzius, observ. bot. III, 12 (1786) ... — — — — — Q. N.A. B.f. VII, 514
R. ophiuroides, Benth. Fl. Austr. VII, 514 (1878) ... — — — — — Q. N.A. B.f. VII, 514 M.fr. VIII, 123.
- OPHIUROS, K. F. Gaertner, de fruct. III, 3, t. 181 (1805). (Ophiurus.)
O. corymbosus, K. F. Gaertner, de fruct. III, 3, t. 181 (1805) ... — — — — — Q. N.A. B.f. VII, 512 M.fr. VI, 85.
- MANISURIS, Linné, syst. nat. ed. duodec. 762 (1767).
M. granularis, Swartz, I, 186 (1797) ... — — — — — Q. N.A. B.f. VII, 514 M.fr. VII, 116; XI, 129.
- HEMARTHRIA, R. Brown, prodr. fl. Nov. Holl. 207 (1810).
H. compressa, R. Brown, prodr. 207 (1810) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 511 M.fr. XI, 129.
- ISCHAEMUM, Linné, syst. nat. 8 (1735); Linné, gen. pl. ed. sec. 525 (1742). (Hologamium, Spodiopogon partly.)
I. truncatiglume, F. v. M. in Benth. Fl. Austr. VII, 518 (1878) ... — — — — — — N.A. B.f. VII, 518
I. arundinaceum, F. v. M. in Benth. Fl. Austr. VII, 519 (1878) ... — — — — — — N.A. B.f. VII, 519
I. triticeum, R. Brown, prodr. 205 (1810) ... — — — — — N.S.W. Q. — B.f. VII, 519

I. australe, R. Brown, prodr. 205 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 519	
I. muticum, Linné, spec. plant. 1049 (1753) ...	—	—	—	—	—	Q.	—	B.f. VII, 520	M.fr. VIII, 120.
I. ciliare, Retzius, observ. bot. VI, 36 (1791) ...	—	—	—	—	N.S.W.	—	—	B.f. VII, 520	M.fr. VIII, 119.
I. decumbens, Benthām, Fl. Austr. VII, 521 (1878) ...	—	—	—	—	—	—	N.A.	B.f. VII, 521	
I. pectinatum, Trinius in Mem. de l'Acad. de Petersb. six. sér. II, 296 (1833) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 521	M.fr. VIII, 118.
I. fragile, R. Brown, prodr. 205 (1810) ...	—	—	—	—	—	Q.	—	B.f. VII, 522	M.fr. VIII, 123; XI, 130.
I. laxum, R. Brown, prodr. 205 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 522	M.fr. VIII, 119.
ANDROPOGON, Roeyen, prodr. exhib. pl. hort. acad. Lugd. 52 (1740). (Sorghum, Chrysopogon, Heteropogon, Schizachyrium, Trachypogon, Holcus partly, Spodiopogon partly.)									
A. erianthoides, F. v. M., fragm. X, 75 (1876) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 529	M.fr. X, 75.
A. sericeus, R. Brown, prodr. 201 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 529	
A. affinis, R. Brown, prodr. 201 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 530	
A. pertusus, Willdenow, spec. plant. IV, 922 (1805) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 530	M.fr. VIII, 122; X, 76.
A. annulatus, Forskael, Fl. Aegypt. Arab. 173 (1775) ...	—	S.A.	—	—	—	Q.	N.A.	B.f. VII, 531	M.fr. VIII, 123.
A. Ischaemum, Linné, spec. plant. 1047 (1753) ...	W.A.	—	—	—	—	—	—	B.f. VII, 531	M.fr. VIII, 122.
A. intermedius, R. Brown, prodr. 202 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 531	
A. procerus, R. Brown, prodr. 202 (1810) ...	—	—	—	—	—	—	N.A.	B.f. VII, 532	M.fr. VIII, 124.
A. exaltatus, R. Brown, prodr. 202 (1810) ...	W.A.	S.A.	—	—	—	—	N.A.	B.f. VII, 532	M.fr. VIII, 124.
A. bombycinus, R. Brown, prodr. 202 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 533	M.fr. VIII, 124.
A. Schoenanthus, Linné, spec. plant. 1046 (1753) ...	—	—	—	—	—	Q.	—	B.f. VII, 534	M.fr. VIII, 124.
A. refractus, R. Brown, prodr. 202 (1810) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 534	M.fr. VIII, 124; X, 76.
A. lachnatherus, Benthām, Fl. Austr. VII, 534 (1874) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 534	
A. fragilis, R. Brown, prodr. 202 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 535	
A. contortus, Linné, spec. plant. 1045 (1753) ...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 517	M.fr. VIII, 120.
A. triticeus, R. Brown, prodr. 201 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 517	M.fr. VIII, 120.
A. Gryllus, Linné, amoen. acad. IV, 332 (1759) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 537	M.fr. VIII, 121; XI, 130.
A. montanus, Roxburgh, Flor. Indic. edit. Carey. I, 271 (1820) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 538	M.fr. VIII, 122; X, 76; XI, 130.
A. acicularis, Retzius, observ. bot. V, 22 (1789) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 538	M.fr. VIII, 122.
A. elongatus, Sprengel, syst. I, 289 (1825) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 539	M.fr. VIII, 121.
A. Halepensis, Sibthorp & Smith, Fl. Graeca, I, 52, t. 68 (1806) ...	W.A.	—	—	—	N.S.W.	Q.	—	B.f. VII, 540	M.fr. VIII, 119.
A. australis, Sprengel, syst. I, 287 (1825) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 540	
A. tropicus, Sprengel, syst. I, 287 (1825) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 541	
A. intrans; Sorghum, F. v. M. in Benth. Fl. Austr. VII, 541 (1878) ...	—	—	—	—	—	—	N.A.	B.f. VII, 541	
ANTHISTIRIA, Linné, fil. gram. gen. 35 (1779). (Iseilema.)									
A. ciliata, Linné fil., gram. gen. 35 (1779) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.f. VII, 542	M.fr. V, 207; VIII, 139.
A. frondosa, R. Brown, prodr. 200 (1810) ...	—	—	—	—	—	Q.	N.A.	B.f. VII, 542	M.fr. VIII, 139; XI, 130.
A. avenacea, F. v. M., fragm. V, 206 (1866) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 543	M.fr. VI, 253; VIII, 139.
A. membranacea, Lindley in Mitch. Trop. Austr. 88 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. VII, 543	M.fr. V, 207; VIII, 139.
APLUDA, Linné, spec. plant. 82 (1753).									
A. mutica, Linné, spec. plant. 82 (1753) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 544	M.fr. VIII, 140.
CHIONACHNE, R. Brown in Horsfield's pl. Jav. rar. 15 (1838).									
C. barbata, R. Brown in Bennett, pl. Javan. rar. 18 (1838) ...	—	—	—	—	—	Q.	—	B.f. VII, 515	
C. cyathopoda, F. v. M. in Benth. Fl. Austr. VII, 516 (1878) ...	—	—	—	—	—	—	N.A.	B.f. VII, 515	M.fr. VIII, 116.
ALOPECURUS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 18 (1737). (Perhaps immigrated.)									
A. geniculatus, Linné, spec. plant. 60 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 555	M.fr. VIII, 138.
LEERSIA, Solander in Swartz, nov. gen. et spec. plant. 1 et 21 (1788). (Asprella, Asperella.)									
L. hexandra, Swartz, nov. gen. et spec. pl. 21 (1788) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 549	M.fr. VIII, 115.
ORYZA, Tournefort, inst. 513, t. 296 (1700), from Theophrastus & Dioscorides.									
O. sativa, Linné, spec. plant. 333 (1753) ...	—	—	—	—	—	—	N.A.	B.f. VII, 550	M.fr. VIII, 115; XI, 130.
POTAMOPHILA, R. Brown, prodr. fl. Nov. Holl. 211 (1810).									
P. parviflora, R. Brown, prodr. 211 (1810) ...	—	—	—	—	N.S.W.	—	—	B.f. VII, 550	M.fr. VIII, 126.
EHRHARTA, Thunberg in Svensk. Ventsk. Acad. Handl. 216, t. 8 (1779). (Tetrarrhena, Michrolaena, Diplax.)									
E. distichophylla, Labill., Nov. Holl. pl. spec. I, 90, t. 117 (1804) ...	—	—	T.	V.	—	—	—	B.f. VII, 554	M.fr. VII, 90.
E. juncea, Sprengel, syst. II, 114 (1825) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 554	M.fr. VII, 90.
E. laevis, Sprengel, syst. II, 115 (1825) ...	W.A.	—	—	—	—	—	—	B.f. VII, 554	M.fr. VII, 90.
E. acuminata, Sprengel, syst. II, 114 (1825) ...	—	—	T.	V.	—	—	—	B.f. VII, 555	M.fr. VII, 90; XI, 130.
E. stipoides, Labillard., Nov. Holl. pl. spec. I, 91, t. 118 (1804) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 552	M.fr. VII, 90.
E. diarrhena, F. v. M., fragm. VII, 89 (1870) ...	—	—	T.	—	—	—	—	B.f. VII, 553	M.fr. VII, 89.
HIEROCHLOE, J. G. Gmelin, Fl. Sibirie. I, 100 (1747). (Hierochloa, Disarrhenum.)									
H. redolens, R. Brown, prodr. 209 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 558	M.fr. VIII, 137.
H. rariflora, J. Hooker, Fl. Austr. I, 93 (1844) ...	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 559	M.fr. VIII, 138.
ARISTIDA, Linné, spec. plant. 82 (1753). (Chaetaria, Arthratherum.)									
A. hygrometrica, R. Brown, prodr. 174 (1810) ...	—	—	—	—	—	—	N.A.	B.f. VII, 561	
A. stipoides, R. Brown, prodr. 174 (1810) ...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. VII, 561	M.fr. VIII, 111.
A. arenaria, Gaudichaud in Frey, voy. Bot. 407 (1826) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 561	
A. Behriana, F. v. M. in Transact. Vict. Inst. 44 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 562	M.fr. XI, 130.
A. leptopoda, Benthām, Fl. Austr. VII, 562 (1878) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 562	M.fr. XI, 130.
A. vagans, Cavanilles, icon. et descr. pl. V, 45, t. 471 (1799) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 562	M.fr. VIII, 111.
A. ramosa, R. Brown, prodr. 173 (1810) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 563	
A. calycina, R. Brown, prodr. 173 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 563	M.fr. XI, 130.
A. depressa, Retzius, observ. bot. IV, 22 (1787) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 563	
STIPA, Linné, spec. plant. 78 (1737). (Streptachne, Urachne partly.)									
S. elegantissima, Labillardière, Nov. Holl. pl. spec. I, 23, t. 29 (1804) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 565	M.fr. VIII, 103.
S. Tuckeri, F. v. M., fragm. XI, 128 (1881) ...	—	—	—	V.	N.S.W.	—	—	—	M.fr. XI, 129.
S. verticillata, Nees in Spreng. syst. veg. cur. post. 30 (1827) ...	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 566	M.fr. VIII, 105.

S. flavescens, Labillardiere, Nov. Holl. pl. spec. I, 24, t. 30 (1804)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 566	M.fr. VIII, 104.
S. teretifolia, Steudel, syn. glum. I, 128 (1855)	W.A.	—	T.	V.	—	—	—	B.f. VII, 567	M.fr. VIII, 104.
S. compressa, R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 567	—
S. Drummondii, Steudel, syn. glum. I, 128 (1855)	W.A.	—	—	—	—	—	—	B.f. VII, 567	—
S. pycnostachya, Benth. Fl. Austr. VII, 568 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 568	—
S. setacea, R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 568	—
S. semibarbata, R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 568	M.fr. VIII, 104; XI, 129.
S. hemipogon, Benth. Fl. Austr. VII, 569 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 569	M.fr. VIII, 104.
S. pubescens, R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 569	M.fr. VIII, 104.
S. aristiglumis, F. v. M. in Transact. Vict. Inst. 43 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 570	M.fr. VIII, 103.
S. Eriopus, Benth. Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 570	—
S. trichophylla, Benth. Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	—	B.f. VII, 570	—
S. scabra, Lindley in Mitch. Trop. Austr. 31 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 570	—
S. Streptachne, F. v. M. in Journ. R.S. of N.S.W. 237 (1881)	—	—	—	—	—	Q.	—	B.f. VII, 572	—
DICHELACHNE, Endlicher, prodr. fl. Norfolk. 20 (1833). (Muehlenbergia partly.)									
D. crinita, J. Hooker, Fl. N. Zel. I, 293 (1853)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 574	M.fr. VIII, 105.
D. sciurea, J. Hooker, Fl. N. Zel. I, 294 (1853)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 574	M.fr. VIII, 105; XI, 129.
PENTAPOGON, R. Brown, prodr. fl. Nov. Holl. 173 (1810).									
P. Billardieri, R. Brown, prodr. 173 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 572	M.fr. VIII, 106.
ECHINOPOGON, Palisot, Essai d'une nouv. Agrostogr. 42, t. 9, f. 5 (1812). (Cinna partly.)									
E. ovatus, Palisot, Agrostogr. 42, t. 9 (1812)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 599	M.fr. VIII, 106.
DIPLOPOGON, R. Brown, prodr. fl. Nov. Holl. 176 (1810). (Dipogonia.)									
D. setaceus, R. Brown, prodr. 176 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 573	M.fr. VI, 86; VIII, 106.
AMPHIPOGON, R. Brown, prodr. 175 (1810). (Aegopogon partly, Pentacraspedion, Gamelythrum.)									
A. debilis, R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 597	M.fr. VIII, 201.
A. strictus, R. Brown, prodr. 175 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 597	M.fr. VIII, 201.
A. laguroides, R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 598	M.fr. VI, 86; VIII, 201.
A. cygnorum, Nees in Lehm. pl. Preiss. II, 100 (1846)	W.A.	—	—	—	—	—	—	B.f. VII, 599	—
A. turbinatus, R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 599	M.fr. VI, 86; VIII, 201.
PAPPOPHORUM, Schreber, gen. pl. II, 787 (1791). (Euneapogon.)									
P. commune, F. v. M., plants of Gregory's Exp. 10 (1859)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 600	M.fr. VIII, 200.
SPOROBOLUS, R. Brown, prodr. fl. Nov. Holl. 169 (1810). (Vilfa partly.)									
S. Virginicus, Humboldt & Kunth, gram. I, 67 (1832)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.f. VII, 621	M.fr. VI, 84.
S. Indicus, R. Brown, prodr. 170 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 622	M.fr. VI, 84; X, 76.
S. diander, Palisot, Agrostogr. 26 (1812)	—	—	—	—	N.S.W.	Q.	—	B.f. VII, 622	M.fr. VI, 84; VIII, 140.
S. pulchellus, R. Brown, prodr. 170 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.f. VII, 623	M.fr. VIII, 140.
S. Lindleyi, Benth. Fl. Austr. VII, 623 (1878)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.f. VII, 623	M.fr. XI, 130. [130.]
S. actinocladius, F. v. M., fragm. VIII, 140 (1874)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.f. VII, 623	M.fr. VI, 84; IX, 194; XI,
AGROSTIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 19 (1737). (Deyeuxia, Didymochaeta, Trichodium, Bromidium, Lachnagrostis partly.)									
A. Muelleri, Benth. Fl. Austr. VII, 576 (1878)	—	—	—	V.	N.S.W.	—	—	B.f. VII, 576	M.fr. VI, 86.
A. scabra, Willdenow, spec. plant. I, 378 (1797)	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 576	—
A. venusta, Trinius in Mém. de l'Acad. de Petersb. VI, 340 (1850)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 576	—
A. aequata, Nees in Hook. Lond. Journ. of Bot. II, 412 (1843)	—	—	T.	—	—	—	—	B.f. VII, 579	—
A. Solandri, F. v. M., Veget. of the Chath. Isl. 60 (1864)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 579	M.fr. XI, 27, 130.
A. montana, R. Brown, prodr. 171 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 581	—
A. quadriseta, R. Brown, prodr. 171 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.f. VII, 581	—
A. cylindrica, R. Brown, prodr. 171 (1810)	W.A.	—	—	—	—	—	—	B.f. VII, 582	—
A. densa, F. v. M.; Deyeuxia, Benth. Fl. Austr. VII, 582 (1878)	—	S.A.	—	V.	—	—	—	B.f. VII, 582	—
A. frigida, F. v. M. in Benth. Fl. Austr. VII, 583 (1878)	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 583	—
A. scabra, R. Brown, prodr. 172 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.f. VII, 583	M.fr. VI, 85.
A. nivalis, F. v. M. in Transact. Vict. Inst. 43 (1854)	—	—	—	V.	N.S.W.	—	—	B.f. VII, 584	—
A. Gunniana, F. v. M.; Deyeuxia, Benth. Fl. Austr. VII, 584 (1878)	—	—	T.	—	—	—	—	B.f. VII, 584	—
A. breviglumis, F. v. M.; Deyeuxia, Benth. Fl. Austr. VII, 584 (1878)	—	—	—	—	N.S.W.	—	—	B.f. VII, 584	—
COELACHNE, R. Brown, prodr. fl. Nov. Holl. 187 (1810).									
C. pulchella, R. Brown, prodr. 187 (1810)	—	—	—	—	—	Q.	—	B.f. VII, 626	—
MICRAIRA, F. v. M., fragm. V, 208 (1866).									
M. subulifolia, F. v. M., fragm. V, 208 (1866)	—	—	—	—	—	Q.	—	—	M.fr. V, 208.
AIRA, Linné, gen. plant. 335 (1737). (Aera, Deschampsia.)									
A. caespitosa, Linné, spec. plant. 64 (1753)	—	S.A.	T.	V.	N.S.W.	—	—	B.f. VII, 587	M.fr. VIII, 138.
TRisetum, Persoon, synopsis. plant. I, 97 (1805).									
T. subspicatum, Palisot, Agrostogr. 88 (1812)	—	—	T.	V.	N.S.W.	—	—	B.f. VII, 588	M.fr. VIII, 136.
ERIACHNE, R. Brown, prodr. fl. Nov. Holl. 183 (1810).									
E. stipacea, F. v. M., fragm. V, 206 (1866)	—	—	—	—	—	Q.	N.A.	B.f. VII, 626	M.fr. VIII, 137; X, 76.
E. Armitii, F. v. M. in Benth. Fl. Austr. VII, 627 (1878)	—	—	—	—	—	Q.	N.A.	B.f. VII, 627	—
E. squarrosa, R. Brown, prodr. 183 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 628	—
E. glauca, R. Brown, prodr. 184 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 628	—
E. rara, R. Brown, prodr. 183 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 628	—
E. agrostidea, F. v. M., fragm. VII, 82 (1870)	—	—	—	—	—	Q.	N.A.	B.f. VII, 628	M.fr. VIII, 137.
E. ciliata, R. Brown, prodr. 184 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 629	—
E. setacea, Benth. Fl. Austr. VII, 629 (1878)	—	—	—	—	—	Q.	N.A.	B.f. VII, 629	—
E. avenacea, R. Brown, prodr. 184 (1810)	—	—	—	—	—	Q.	N.A.	B.f. VII, 629	—
E. aristidea, F. v. M., fragm. V, 205 (1866)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.f. VII, 629	M.fr. V, 205.
E. pallescens, R. Brown, prodr. 184 (1810)	—	—	—	—	—	Q.	—	B.f. VII, 630	—
E. festucacea, F. v. M., fragm. V, 205 (1866)	—	—	—	—	—	Q.	N.A.	B.f. VII, 630	M.fr. V, 205.

- E. ovata*, Nees in Hook. Lond. Journ. II, 418 (1843) ... W.A. S.A. — — — — N.A. B.fl.VII,630 M.fr.VIII,137.
E. meliacea, F. v. M., fragm. V, 205 (1866) ... — — — — — — N.A. B.fl.VII,631 M.fr.V,205.
E. pallida, F. v. M. in Benth. Fl. Austr. VII, 631 (1878) ... — — — — — — N.A. B.fl.VII,631
E. scleranthoides, F. v. M., fragm. VIII, 233 (1874) ... — — S.A. — — — — — B.fl.VII,631
E. mucronata, R. Brown, prodr. 184 (1810) ... — — — — — — Q. — B.fl.VII,632
E. obtusa, R. Brown, prodr. 184 (1810) ... — — — — — N.S.W. Q. N.A. B.fl.VII,632 M.fr.VIII,137;XI,130.
E. capillaris, R. Brown, prodr. 184 (1810) ... — — — — — — N.A. B.fl.VII,632
- ANISOPOGON**, R. Brown, prodr. fl. Nov. Holl. 176 (1810).
A. avenaceus, R. Brown, prodr. 176 (1810) ... — — — — — N.S.W. Q. — B.fl.VII,590 M.fr.VIII,106.
- DANTHONIA**, De Candolle, Fl. franç. III, 32 (1805). (Amphibromus, Monacather, Plinthacanthus.)
D. paradoxa, R. Brown, prodr. 177 (1810) ... — — — — — N.S.W. — — B.fl.VII,591
D. bipartita, F. v. M., fragm. I, 160 (1859) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,592 M.fr.VIII,136.
D. carphoides, F. v. M. in Benth. Fl. Austr. VII, 592 (1878) ... — S.A. — V. N.S.W. — — B.fl.VII,592 M.fr.XI,130.
D. penicillata, F. v. M., fragm. VIII, 135 (1873) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,592 M.fr.VIII,135.
D. robusta, F. v. M. in Transact. Vict. Inst. 44 (1854) ... — — — — V. N.S.W. — — B.fl.VII,593 M.fr.VIII,136.
D. pauciflora, R. Brown, prodr. 177 (1810) ... — — — — T. V. N.S.W. — — B.fl.VII,596
D. nervosa, J. Hooker, Fl. Tasman. II, 121, t. 163 (1860) ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,589 M.fr.VIII,135.
- ASTREBLA**, F. v. M., fragm. phytogr. Austr. X, 76 (1876).
A. pectinata, F. v. M., fragm. X, 76 (1876) ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,602 M.fr.VIII,134.
A. triticoideus, F. v. M., fragm. X, 76 (1876) ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,602 M.fr.VIII,134.
- MICROCHLOA**, R. Brown, prodr. fl. Nov. Holl. 208 (1810).
M. setacea, R. Brown, prodr. 208 (1810) ... — — — — — — N.A. B.fl.VII,608
- CYNODON**, L. C. Richard in Persoon, synops. I, 85 (1805).
C. Dactylon, L. C. Richard in Persoon, synops. I, 85 (1805) ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,609 M.fr.VIII,113.
C. tenellus, R. Brown, prodr. 187 (1810) ... — — — — — — N.A. B.fl.VII,609 M.fr.VIII,113.
C. convergens, F. v. M., fragm. VIII, 113 (1874) ... — — — — — — Q. N.A. B.fl.VII,610 M.fr.X,76.
C. ciliaris, Benth. Fl. Austr. VII, 610 (1878) ... — S.A. — — — — — B.fl.VII,610
- CHLORIS**, Swartz, nov. gen. et spec. plant. 25 (1788).
C. unispicea, F. v. M., fragm. VII, 118 (1871) ... — — — — — — Q. — B.fl.VII,611 M.fr.
C. pumilio, R. Brown, prodr. 186 (1810) ... — — — — — — Q. N.A. B.fl.VII,611
C. pectinata, Benth. Fl. Austr. VII, 612 (1878) ... — — S.A. — — — — — Q. N.A. B.fl.VII,612
C. divaricata, R. Brown, prodr. 186 (1810) ... — — — — — — Q. N.A. B.fl.VII,612 M.fr.VI,85.
C. acicularis, Lindley in Mitch. Trop. Austr. 33 (1848) ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,612 M.fr.VI,85.
C. truncata, R. Brown, prodr. 186 (1810) ... — — S.A. — V. N.S.W. Q. — B.fl.VII,612 M.fr.VI,85;XI,130.
C. ventricosa, R. Brown, prodr. 186 (1810) ... — — — — — N.S.W. Q. — B.fl.VII,613 M.fr.VIII,132.
C. barbata, Swartz, nov. gen. et spec. plant. 25 (1788) ... — — S.A. — — — — — N.A. B.fl.VII,613 M.fr.VI,85.
C. scariosa, F. v. M., fragm. VI, 85 (1868) ... — — — — — — Q. N.A. B.fl.VII,614 M.fr.XI,130.
- ELEUSINE**, J. Gaertner, de fruct. I, 7 (1788). (Leptochloa, Dactyloctenium, Acrachne).
E. Aegyptiaca, Persoon, syn. I, 82 (1805) ... — — S.A. — V. N.S.W. Q. N.A. B.fl.VII,615 M.fr.VIII,111.
E. Indica, J. Gaertner, de fruct. I, 7 (1788) ... — — — — — N.S.W. Q. — B.fl.VII,615 M.fr.VIII,112.
E. verticillata, Roxburgh, Fl. Ind. ed. Car. & Wall. I, 346 (1820) ... — — — — — — Q. N.A. B.fl.VII,616 M.fr.VIII,112. [130.
E. digitata, Sprengel, syst. cur. post. 36 (1827) ... W.A. S.A. — — — — — N.S.W. Q. N.A. B.fl.VII,617 M.fr.I,216;VIII,112;XI,
E. Chinensis, F. v. M.; Poa, Koenig in Roth, nov. pl. spec. 65 (1821) ... — — — — — N.S.W. Q. — B.fl.VII,617 M.fr.VIII,132.
E. digitata, Sprengel, syst. veg. cur. post. 36 (1827) ... — — — — — — Q. N.A. B.fl.VII,617 M.fr.VIII,113.
- CENTOTHECA**, Desvaux in Palisot, Agrost. 69, t. 14 (1812). [76.
C. lappacea, Desvaux in Palisot, Agrost. 69, t. 14 (1812) ... — — — — — — Q. — B.fl.VII,640 M.fr.VI,85;VIII,116;X,
- POA**, Linné, gen. plant. 20 (1737). (Glyceria partly.)
P. Billardieri, Steudel, syn. glum. I, 262 (1855) ... W.A. S.A. T. V. — — — — N.A. B.fl.VII,651
P. homomalla, Nees in Lehm. pl. Preiss. II, 104 (1846) ... W.A. — — — — — — — B.fl.VII,651
P. caespitosa, G. Forster, fl. ins. Austr. prodr. 89 (1786) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,651 M.fr.VIII,133;XI,13.
P. Maxwellii, Benth. Fl. Austr. VII, 653 (1878) ... W.A. — — — — — — — B.fl.VII,652
P. nodosa, Nees in Lehm. pl. Preiss. II, 105 (1846) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,653 M.fr.VIII,132.
P. saxicola, R. Brown, prodr. 654 (1810) ... — — — — — T. — — B.fl.VII,654
P. lepida, F. v. M., fragm. VIII, 130 (1874) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,654 M.fr.XI,130.
P. Fordeana, F. v. M., fragm. VIII, 130 (1874) ... — — S.A. — V. N.S.W. — — B.fl.VII,657 M.fr.VIII,130.
P. fluitans, Scopoli, flor. Carniol. 106 (1772) ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,657 M.fr.VIII,129.
P. latispicea; Festuca, F. v. M., VIII, 127 (1873) ... — — — — — N.S.W. — — B.fl.VII,658 M.fr.VIII,128.
P. syrtica, F. v. M. in Transact. Vict. Inst. 45 (1854) ... W.A. S.A. T. V. — — — — B.fl.VII,658 M.fr.VIII,130. [130.
P. dives; Festuca, F. v. M., fragm. III, 147 (1863) ... — — — — — V. — — B.fl.VII,659 M.fr.VI,147;VIII,129,XI,
P. ramigera, F. v. M. in Transact. Vict. Inst. 45 (1854) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,639 M.fr.VI,85;VIII,131,XI,
130.
- FESTUCA**, Dillenius, nov. gen. 90, t. 3 (1719). (Glyceria partly, Vulpia partly, Schedonorus, Schenodorus, Brizopyrum.)
F. duriuscula, Linné, spec. plant. 74 (1753) ... — — S.A. T. V. N.S.W. — — B.fl.VII,663 M.fr.VI,85.
F. scirpoidea, F. v. M., fragm. VIII, 129 (1874) ... — — W.A. — — — — — — B.fl.VII,655 M.fr.XI,130.
F. litoralis, Labillardière, Nov. Holl. spec. plant. I, 22, t. 27 (1804) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,656 M.fr.VIII,129.
F. Hookeriana, F. v. M. in J. Hooker, Fl. Tasm. II, 122, t. 115 (1860) ... — — T. V. N.S.W. — — B.fl.VII,656 M.fr.VIII,131.
- DIPLACHNE**, Palisot, Essai d'une nouv. Agrostogr. 9, t. 16 (1812). (Uralapis.)
D. loliiformis, F. v. M. in Benth. fl. Austr. VII, 618 (1878) ... — — S.A. — V. N.S.W. Q. — B.fl.VII,616 M.fr.VIII,128;XI,130.
D. Muelleri, Benth. Fl. Austr. VII, 619 (1878) ... — — S.A. — — — — — N.A. B.fl.VII,619
D. fusca, Palisot, Agrost. 163 (1812) ... — — W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,619 M.fr.VIII,127.
D. parviflora, Benth. Fl. Austr. VII, 620 (1878) ... — — — — — — — N.A. B.fl.VII,620 M.fr.VIII,129,

- TRIODIA, R. Brown, prodr. fl. Nov. Holl. 182 (1810).
- T. Mitchelli, Benth. Fl. Austr. VII, 606 (1878) ... — — — — N.S.W. Q. — B.fl. VII, 606
- T. pungens, R. Brown, prodr. 182 (1810) ... — S.A. — — — — N.A. B.fl. VII, 606 M.fr. VIII, 129.
- T. Cunninghamii, Benth. Fl. Austr. VII, 606 (1878) ... — — — — — Q. N.A. B.fl. VII, 606
- T. irritans, R. Brown, prodr. 182 (1810) ... — — — — — W.A. S.A. — V. N.S.W. Q. — B.fl. VII, 607 M.fr. VIII, 129.
- T. procera, R. Brown, prodr. 182 (1810) ... — — — — — — N.A. B.fl. VII, 607
- T. microstachya, R. Brown, prodr. 182 (1810) ... — — — — — — N.A. B.fl. VII, 607
- T. microdon, F. v. M.; Triraphis, Benth. Fl. Austr. VII, 605 (1878) — — — — — N.S.W. — — B.fl. VII, 605
- DISTICHLIS, Rafinesque in Journ. de Phys. LXXXIX, 104 (1819). (Uniola partly, Brizopyrum partly.)
- D. maritima, Rafinesque in Journ. de Phys. LXXXIX, 104 (1819) — S.A. T. V. N.S.W. — — B.fl. VII, 637 M.fr. VIII, 129.
- BROMUS, Dillenius in Linné, syst. nat. 8 (1735); Linné, gen. plant. 15 (1737).
- B. arenarius, Labillardière, Nov. Holl. pl. spec. I, 23, t. 28 (1804) W.A. S.A. — V. N.S.W. Q. — B.fl. VII, 660 M.fr. VI, 126; XI, 20.
- ERAGROSTIS, Palisot, Essai d'une nouv. Agrostogr. 70, t. 14 (1812). (Poa partly).
- E. tenella, Palisot in Roem. et Schult. syst. veg. II, 576 (1817) — — — — — S.A. — V. N.S.W. Q. N.A. B.fl. VII, 643 M.fr. VIII, 132; XI, 130
- E. nigra, Nees in Steud. nomencl. bot. I, 563 (1841) ... — — — — — N.S.W. — — B.fl. VII, 643
- E. imbecilla, Benth. Fl. Austr. VII, 643 (1878) ... — — — — — — Q. — B.fl. VII, 643
- E. trichophylla, Benth. Fl. Austr. VII, 644 (1878) ... — — — — — S.A. — — — — — B.fl. VII, 644
- E. leptocarpa, Benth. Fl. Austr. VII, 644 (1878) ... — — — — — S.A. — — — — — Q. N.A. B.fl. VII, 644
- E. megalosperma, F. v. M. in Benth. Fl. Austr. VII, 644 (1878) — — — — — N.S.W. Q. — B.fl. VII, 644
- E. pilosa, Palisot, Agrost. 71 (1812) ... — — — — — S.A. — V. N.S.W. Q. — B.fl. VII, 645 M.fr. VIII, 133; XI, 130
- E. leptostachya, Steudel, syn. glum. I, 279 (1855) ... — — — — — N.S.W. Q. — B.fl. VII, 645
- E. Schultzei, Benth. Fl. Austr. VII, 646 (1878) ... — — — — — — — N.A. B.fl. VII, 646
- E. diandra, Steudel, syn. glum. I, 279 (1855) ... — — — — — W.A. S.A. — — N.S.W. Q. N.A. B.fl. VII, 646 M.fr. XI, 130.
- E. Brownii, Nees in Steud. nomencl. bot. I, 562 (1841) ... — — — — — W.A. S.A. — V. N.S.W. Q. N.A. B.fl. VII, 646
- E. concinna, Steudel, syn. glum. I, 279 (1855) ... — — — — — S.A. — — — — — Q. N.A. B.fl. VII, 647
- E. speciosa, Steudel, syn. glum. I, 279 (1855) ... — — — — — S.A. — — — — — Q. N.A. B.fl. VII, 648
- E. laniflora, Benth. Fl. Austr. VII, 648 (1878) ... — — — — — S.A. — — — — — N.S.W. Q. — B.fl. VII, 648
- E. eriopoda, Benth. Fl. Austr. VII, 648 (1878) ... — — — — — S.A. — — — — — N.S.W. — N.A. B.fl. VII, 648
- E. setifolia, Nees in Hook. Lond. Journ. II, 419 (1843) ... — — — — — W.A. S.A. — — N.S.W. Q. N.A. B.fl. VII, 648
- E. lacunaria, F. v. M., first gen. Rep. 20 (1853) ... — — — — — S.A. — V. N.S.W. Q. — B.fl. VII, 649
- E. falcata, Gaudichaud in Voy. Freyc. Bot. 408, t. 25 (1826) ... — — — — — W.A. S.A. — V. N.S.W. Q. N.A. B.fl. VII, 649
- E. stenostachya, Steudel, syn. glum. I, 279 (1855) ... — — — — — — Q. N.A. B.fl. VII, 650
- HETERACHNE, Benth. in J. Hook. icon. plant. XIII, 39 (1877).
- H. Brownii, Benth. in J. Hook. icon. XIII, 40 (1877) ... — — — — — — Q. N.A. B.fl. VII, 635 M.fr. VIII, 132.
- H. Gulliveri, Benth. in J. Hook. icon. XIII, 39, t. 1250 (1877) — — — — — — Q. N.A. B.fl. VII, 635
- ECTROSLA, R. Brown, prodr. fl. Nov. Holl. 185 (1810).
- E. Schultzei, Benth. Fl. Austr. VII, 633 (1878) ... — — — — — — — N.A. B.fl. VII, 633
- E. leporina, R. Brown, prodr. 186 (1810) ... — — — — — N.S.W. Q. N.A. B.fl. VII, 633 M.fr. VIII, 109.
- E. agrostoides, Benth. Fl. Austr. VII, 634 (1878) ... — — — — — — — N.A. B.fl. VII, 634
- E. Gulliveri, F. v. M., fragm. VIII, 201 (1874) ... — — — — — — Q. N.A. B.fl. VII, 634 M.fr. VIII, 201.
- ELYTHROPHORUS, Palisot, Essai d'une nouv. Agrostogr. 67, t. 14 (1812).
- E. articulatus, Palisot, Agrost. 67, t. 14 (1812) ... — — — — — S.A. — V. N.S.W. Q. N.A. B.fl. VII, 638 M.fr. VIII, 109.
- TRIRHAPHIS, R. Brown, prodr. fl. Nov. Holl. 185 (1810). (Triraphis.)
- T. mollis, R. Brown, prodr. 185 (1810) ... — — — — — W.A. S.A. — V. N.S.W. Q. N.A. B.fl. VII, 603 M.fr. VIII, 108.
- T. pungens, R. Brown, prodr. 185 (1810) ... — — — — — — — N.A. B.fl. VII, 604 M.fr. VIII, 108, 125.
- T. bromoides, F. v. M., fragm. VIII, 108 (1874) ... — — — — — W.A. — — — — — B.fl. VII, 604 M.fr. VIII, 108.
- T. danthonioides, F. v. M., fragm. VIII, 125 (1874) ... — — — — — W.A. — — — — — B.fl. VII, 604 M.fr. VIII, 108.
- AGROPYRON, J. Gaertner, nov. comm. Petrop. XIV, 539 (1770). (Agropyrum, Triticum partly, Anthosachne, Vulpia partly.)
- A. scabrum, Palisot, Agrost. 102 (1812) ... — — — — — W.A. S.A. T. V. N.S.W. Q. — B.fl. VII, 665 M.fr. VI, 85; XI, 27, 130.
- A. velutinum, Nees in Hook. Lond. Journ. of Bot. II, 417 (1843) — — — — — T. V. N.S.W. — — B.fl. VII, 665
- A. pectinatum, Palisot, Agrost. 102 (1812) ... — — — — — T. V. N.S.W. — — B.fl. VII, 666 M.fr. VI, 85.
- ARUNDO, Tournefort, inst. 526 (1700), from Varro. (Phragmites.)
- A. Phragmites, Dodoens, stirp. hist. pemptad. 602 (1583) — — — — — S.A. T. V. N.S.W. Q. — B.fl. VII, 637
- A. Roxburghii, F. v. M.; Phragmites, Kunth in nov. act. acad. Caes. XIX, suppl. I, 152 (1843) ... — — — — — — Q. N.A. — M.fr. XI, 139.
- BAMBUSA, Schreber, gener. plantar. I, 236 (1789).
- Species (three, of this or allied genera, not yet found in flower) undetermined ... — — — — — — Q. N.A. — M.fr. VI, 85; XI, 129.

ACOTYLEDONEAE.

A. L. de Jussieu, genera plantarum 1 (1789).

ACOTYLEDONEAE VASCULARES.

Meissner, plantarum vascularium genera, 429 (1843).

RHIZOSPERMAE.

G. Weber, primit. Fl. Holsat. 74 (1780).

- AZOLLA, Lamarek, Encycl. méthod. I, 343 (1783).
 A. pinnata, R. Brown, prodr. 167 (1810) ... — S.A. — V. N.S.W. Q. — B.fl. VII, 679 M.fr. V, 140.
 A. filiculoides, Lamarek, Encycl. méth. I, 343 (1783) ... — S.A. T. V. N.S.W. Q. — B.fl. VII, 680 M.fr. V, 140; XI, 127.
 MARSILEA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 326 (1737). (Marsiglia.)
 M. quadrifolia, Linné, spec. plant. 1099 (1753) ... — W.A. S.A. — V. N.S.W. Q. N.A. B.fl. VII, 683 M.fr. V, 140.
 PILULARIA, Vaillant, Bot. Paris. 159, t. 15 (1727).
 P. globulifera, Linné, spec. plant. 1100 (1753) ... — W.A. S.A. T. V. N.S.W. — — B.fl. VII, 684 M.fr. V, 140.
 ISOETES, Linné, skanska resa 420 (1751).
 I. Drummondii, A. Braun in Berl. Monatsb. 593 (1863) ... — W.A. — — V. — — — B.fl. VII, 672 M.fr. V, 140.
 I. elatior, F. v. M. & A. Braun in Linnaea XXV, 722 (1852) ... — — T. — — — — — M.fr. V, 140; XI, 139.
 I. humilior, F. v. M. & A. Braun in Linnaea XXV, 722 (1852) ... — — T. — — — — — M.fr. V, 140; XI, 139.
 I. Gunnii, A. Braun in Berl. Monatsb. 535 (1868) ... — — T. — — — — — M.fr. XI, 139.
 I. Muelleri, A. Braun in Berl. Monatsb. 535 (1868) ... — — — — — Q. — — M.fr. XI, 139.

LYCOPODINEAE.

Swartz, synops. filic. XV (1806).

- PSILOTUM, Swartz in Schrader, Journ. fuer die Bot. II, 109 (1800.)
 P. triquetrum, Swartz, synops. filic. 187 (1806) ... — — — — N.S.W. Q. — B.fl. VII, 681 M.fr. V, 112; X, 118.
 P. complanatum, Swartz, synops. filic. 187 (1806) ... — — — — — Q. — B.fl. VII, 682 M.fr. V, 112.
 TMESIPTERIS, Bernhardt in Schrader, Journ. fuer die Bot. II, 131 (1800).
 T. Tannensis, Bernhardt in Schrad. Journ. II, 131, t. 2 (1800) ... — — T. V. N.S.W. Q. — B.fl. VII, 680 M.fr. V, 112; X, 118.
 LYCOPODIUM, Ruppius, Fl. Jenens. 32 (1718), from Tabernaemontanus (1588).
 L. Selago, Linné, spec. plant. 1102 (1753) ... — — T. V. N.S.W. — — B.fl. VII, 674 M.fr. V, 111; X, 118.
 L. Phlegmaria, Linné, spec. plant. 1101 (1753) ... — — — — — Q. — B.fl. VII, 674 M.fr. V, 111; X, 118.
 L. clavatum, Linné, spec. plant. 1101 (1753) ... — — T. V. N.S.W. — — B.fl. VII, 675 M.fr. V, 111.
 L. Carolinianum, Linné, spec. plant. 1104 (1753) ... — — W.A. — T. V. N.S.W. Q. — B.fl. VII, 675 M.fr. V, 111; X, 118.
 L. laterale, R. Brown, prodr. 165 (1810) ... — — S.A. T. V. N.S.W. Q. — B.fl. VII, 675 M.fr. V, 111; VI, 124; X, 118.
 L. cernuum, Linné, spec. plant. 1103 (1753) ... — — — — — Q. N.A. B.fl. VII, 676 M.fr. V, 111; X, 118.
 L. densum, Labill., Nov. Holl. plant. spec. II, 104, t. 251 (1806) ... — — S.A. T. V. N.S.W. — — B.fl. VII, 676 M.fr. V, 111; X, 118.
 L. scariosum, G. Forster, florul. ins. austr. prodr. 87 (1786) ... — — T. V. — — — B.fl. VII, 676 M.fr. V, 112.
 L. volubile, G. Forster, florul. ins. austr. prodr. 86 (1786) ... — — — — — N.A. B.fl. VII, 677 M.fr. V, 112.
 SELAGINELLA, Palisot, prodr. des fam. de l'Aethéogamie 101 (1805). (Lycopodium partly.)
 S. Preissiana, Spring in Mém. de l'Ac. de Brux. 61 (1842) ... — — W.A. S.A. T. V. N.S.W. Q. — B.fl. VII, 677 M.fr. V, 112; VIII, 275.
 S. uliginosa, Spring in Mém. de l'Ac. de Brux. II, 60 (1842) ... — — S.A. T. V. N.S.W. Q. — B.fl. VII, 678 M.fr. V, 112; VIII, 275.
 S. flabellata, Spring in Mém. de l'Ac. de Brux. 174 (1849) ... — — — — — Q. — B.fl. VII, 678 M.fr. V, 112; VIII, 275.
 S. concinna, Spring in Mém. de l'Ac. de Brux. 199 (1849) ... — — — — — Q. — B.fl. VII, 678 M.fr. V, 112; VIII, 274.
 S. Belangeri, Spring in Mém. de l'Ac. de Brux. 242 (1849) ... — — — — — N.S.W. Q. N.A. B.fl. VII, 679
 PHYLLOGLOSSUM, Kunze in Mohl & Schlechtendal, Bot. Zeitung, 721 (1843).
 P. Drummondii, Kunze in der Bot. Zeit. 721 (1843) ... — — W.A. — T. V. N.S.W. — — B.fl. VII, 672 M.fr. V, 112; X, 118.

FILICES.

Linné, gen. plant. 322 (1737).

- OPHIOGLOSSUM, Tournefort, inst. rei herb. 548, t. 325 (1700), from Bock (1539). (Ophioderma.)
 O. vulgatum, C. Bauhin, pinax theatr. bot. 354 (1623) ... — — S.A. T. V. N.S.W. Q. N.A. B.fl. VII, 688 M.fr. V, 112; VI, 124.
 O. pendulum, Linné, sp. pl. edit. sec. 1518 (1763) ... — — — — — N.S.W. Q. — B.fl. VII, 689 M.fr. V, 113; VI, 124.
 BOTRYCHIUM, Swartz in Schrader, Journ. fuer die Bot. II, 110 (1800).
 B. Lunaria, Swartz in Schrad. Journ. II, 110 (1800) ... — — T. V. N.S.W. — — B.fl. VII, 690 M.fr. V, 113.
 B. ternatum, Swartz, synops. filic. 172 (1806) ... — — T. V. N.S.W. Q. — B.fl. VII, 690 M.fr. V, 113.
 HELMINTHOSTACHYS, Kaulfuss, enum. filic. Chamiss. 28 (1824).
 H. Zeylanica, Kaulfuss, enum. fil. 28 (1824) ... — — — — — Q. — B.fl. VII, 690 M.fr. V, 113; IX, 190.
 LYGODIUM, Swartz in Schrader, Journ. fuer die Bot. II, 7 & 106 (1800). (Hydroglossum, Lygodietyon.)
 L. scandens, Swartz in Schrad. Journ. II, 106 (1800) ... — — — — — N.S.W. Q. N.A. B.fl. VII, 691 M.fr. V, 113; VII, 84, X, 118.
 L. reticulatum, Schkuhr, kryptog. Gewaechse 139, t. 139 (1809) ... — — — — — Q. — B.fl. VII, 692 M.fr. VII, 83; VIII, 157.
 L. Japonicum, Swartz, syn. filic. 154 (1806) ... — — — — — Q. N.A. B.fl. VII, 692 M.fr. V, 113; VII, 84, 156.
 SCHIZAEA, Smith in Mem. de l'Acad. Turin. V, 149, t. 19 (1791). (Lopidium, Actinostachys.)
 S. rupestris, R. Brown, prodr. 162 (1810) ... — — — — — N.S.W. — — B.fl. VII, 693 M.fr. V, 113.
 S. fistulosa, Labillardière, Nov. Holl. pl. spec. II, 103, t. 250 (1806) ... — — S.A. T. V. — — — B.fl. VII, 693 M.fr. V, 113.
 S. dichotoma, Smith in Mém. de l'Ac. Tur. V, 149 (1791) ... — — S.A. T. V. N.S.W. Q. N.A. B.fl. VII, 694 M.fr. V, 113; X, 118.
 S. Forsteri, Sprengel, Anleit. zur Kenntn. der Gew. III, 157 (1804) ... — — — — — Q. — — M.fr. VIII, 275.
 ANGIOPTERIS, G. Hoffmann in Comment. Goett. XII, 29 (1796). [118.
 A. evecta, G. Hoffmann in Comment. Goett. XII, 29 (1796) ... — — — — — Q. — B.fl. VII, 694 M.fr. V, 114; VII, 120; X,

- MARATTIA, Swartz, nov. gen. et spec. plant. 8 & 128 (1788).
M. fraxinea, Smith, plant. icon. hact. ined. t. 48 (1790)... — — — N.S.W. Q. — B.f. VII, 695 M.fr. V, 114; X, 118.
- TRICHOMANES, Linné, syst. nat. 8 (1835); Linné, hort. Cliffort. 476 (1737).
T. peltatum, Baker in Journ. Linn. Soc. IX, 336, t. 8 (1865) ... — — — — Q. — B.f. VII, 701 M.fr. X, 117.
T. Vitiense, Baker in Journ. Linn. Soc. IX, 338, t. 8 (1865) ... — — — — N.S.W. Q. — B.f. VII, 701 M.fr. VIII, 32; IX, 132.
T. Motleyi, Bosch, Hymenoph. Javan (1861) ... — — — — Q. — — — —
T. cuspidatum, Willdenow, spec. plant. V, 499 (1810) ... — — — — Q. — — — — M.fr. XI, 131 & 139.
T. parvulum, Poiret, Encycl. méth. VIII, 44 (1810) ... — — — — N.S.W. Q. — B.f. VII, 701 M.fr. VII, 155.
T. digitatum, Swartz, synops. filic. 370 & 422 (1806) ... — — — — N.S.W. — — — — B.f. VII, 702
T. venosum, R. Brown, prodr. 159 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 702 M.fr. V, 116; IX, 191.
T. Javanicum, Blume, flor. Jav. filic. 224 (1836) ... — — — — — Q. — — — — B.f. VII, 702
T. rigidum, Swartz, nov. gen. et spec. pl. 137 (1788) ... — — — — N.S.W. Q. — B.f. VII, 702 M.fr. V, 115.
T. bipunctatum, Poiret, Encycl. méth. VIII, 44 (1810) ... — — — — — Q. — — — — B.f. VII, 703 M.fr. X, 117; XI, 132.
T. humile, G. Forster, florul. ins. Austr. prodr. 84 (1786) ... — — — — N.S.W. — — — — —
T. caudatum, Brackenridge, Bot. of Wilk. U.S. Exped. fil. 256, t. 36 (1854) ... — — — — N.S.W. Q. — B.f. VII, 703 M.fr. V, 116; VII, 122.
T. Bauerianum, Endlicher, prodr. fl. Norf. 17 (1833) ... — — — — N.S.W. Q. — B.f. VII, 703 M.fr. VII, 121; XI, 132.
T. parviflorum, Poiret, Encycl. méth. VIII, 44 (1810) ... — — — — — Q. — — — — B.f. VII, 704 M.fr. VII, 122.
- HYMENOPHYLLUM, Smith in Roemer, Archiv. I, 56 (1797).
H. marginatum, Hooker & Greville, icon. filic. I, t. 34 (1829) ... — — — — N.S.W. — — — — B.f. VII, 705
H. nitens, R. Brown, prodr. 159 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 705
H. Javanicum, Sprengel, syst. veg. IV, 132 (1827) ... — — — — T. V. N.S.W. Q. — B.f. VII, 705 M.fr. V, 116.
H. Tunbridgense, Smith in Roemer, Archiv. I, 56 (1797) ... — — — — T. V. N.S.W. Q. — B.f. VII, 706 M.fr. V, 116.
H. multifidum, Swartz, synops. filic. 149 & 378 (1806) ... — — — — N.S.W. — — — — B.f. VII, 707
- CYATHEA, Smith in Mém. de l'Acad. Turin. V, 416 (1791). (Hemitelia, Amphicosmia.)
C. Lindsayana, Hooker, syn. filic. 25 (1865) ... — — — — N.S.W. Q. — B.f. VII, 708 M.fr. VIII, 177.
C. arachnoidea, Hooker, syn. filic. 24 (1866) ... — — — — — Q. — — — — B.f. VII, 708 M.fr. VI, 200.
C. Macarthurii, F. v. M., fragm. VII, 177 (1874) ... — — — — N.S.W. — — — — B.f. VII, 708 M.fr. VIII, 176; X, 120.
C. medullaris, Swartz, synops. filic. 140 (1806) ... — — — — T. V. N.S.W. — — — — B.f. VII, 708 M.fr. V, 116; VIII, 177.
C. Cunninghami, J. Hooker, icon. plant. 985 (1854) ... — — — — T. V. — — — — —
C. brevipinna, Baker in Bentham fl. Austr. VII, 709 (1878) ... — — — — N.S.W. — — — — B.f. VII, 709 M.fr. XI, 139.
C. Moorei, F. v. M.; Hemitelia, Baker in Gardn. Chron. 252 (1872) ... — — — — N.S.W. — — — — B.f. VII, 709 M.fr. VIII, 157.
- CERATOPTERIS, Brogniart in Bull. de la Soc. philom. 186 (1821). (Parkeria.)
C. thalictroides, Brogniart in Bull. Soc. philom. 186 (1821) ... — — — — — Q. N.A. B.f. VII, 695 M.fr. V, 122; VIII, 158.
- GLEICHENIA, Smith in Mém. de l'Acad. Turin. V, 418 (1791). (Mertensia, Platyzoma, Stromatopteris.) [118].
G. platyzoma, F. v. M., fragm. V, 114 (1866) ... — — — — N.S.W. Q. N.A. B.f. VII, 696 M.fr. V, 114; VIII, 157; X,
G. circinata, Swartz, syn. filic. 165 et 394 (1806) ... — — — — S.A. T. V. N.S.W. Q. N.A. B.f. VII, 697 M.fr. V, 115.
G. dicarpa, R. Brown, prodr. 161 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 698 M.fr. V, 115.
G. flabellata, R. Brown, prodr. 161 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 698 M.fr. V, 114; VIII, 157.
G. Hermannii, R. Brown, prodr. 161 (1810) ... — — — — N.S.W. Q. N.A. B.f. VII, 698 M.fr. V, 114; X, 118.
- OSMUNDA, Tournefort, inst. rei herb. 547, t. 324 (1700), from l'Obel (1576). (Todea, Leptopteris.)
O. barbara, Thunberg, prodr. plant. Capens. 171 (1800) ... — — — — S.A. T. V. N.S.W. Q. — B.f. VII, 699 M.fr. V, 114.
O. Fraseri, F. v. M.; Todea, Hooker & Greville, icon. filic. t. 101 (1829) ... — — — — N.S.W. — — — — B.f. VII, 700 M.fr. V, 114; VII, 156.
O. Moorei, F. v. M.; Todea, Baker in Trimen, Journ. of Bot. 16 (1873) ... — — — — N.S.W. — — — — B.f. VII, 700 M.fr. VIII, 157.
- ALSOPHILA, R. Brown, prodr. 158 (1810).
A. Rebecca, F. v. M., fragm. V, 53 (1865) ... — — — — — Q. — — — — B.f. VII, 710 M.fr. V, 117, 215; VIII, 180.
A. Loddigesii, Kunze in Schlecht. Linnaea XXIII, 221 (1850) ... — — — — N.S.W. Q. — B.f. VII, 710 [VIII, 178].
A. australis, R. Brown, prodr. 158 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 710 M.fr. V, 52, 116; VII, 120;
A. excelsa, R. Brown, prodr. 158 (1810) ... — — — — — N.S.W. Q. — — — — —
A. Leichhardtiana, F. v. M., fragm. V, 53 (1865) ... — — — — N.S.W. Q. — B.f. VII, 711 M.fr. V, 117; VI, 200; VII,
A. Robertiana, F. v. M., fragm. V, 54 (1865) ... — — — — — Q. — — — — B.f. VII, 712 M.fr. V, 117, 213. [120].
- DICKSONIA, L'Heritier, sert. Angl. 31 (1788). (Cibotium, Patania, Dennstaedtia, Deparia, Balantium.)
D. Billardieri, F. v. M., fragm. VIII, 175 (1874) ... — — — — S.A. T. V. N.S.W. Q. — B.f. VII, 712 M.fr. V, 117; VI, 199; VII,
D. Youngiae, C. Moore in Hook. & Bak. syn. filic. 51 (1865) ... — — — — N.S.W. Q. — B.f. VII, 713 M.fr. VI, 200. [120].
D. davallioides, R. Brown, prodr. 158 (1810) ... — — — — V. N.S.W. Q. — B.f. VII, 713 M.fr. VIII, 158. [104].
D. nephroides, Deparia, Baker in Gardn. Chron. 253 (1872) ... — — — — N.S.W. — — — — B.f. VII, 714 M.fr. VIII, 157; IX, 78; X,
- DAVALLIA, Smith in Mém. de l'Acad. de Turin. V, 414 (1791). (Microlepia, Humata.)
D. solida, Swartz, synops. filic. 132 & 345 (1806) ... — — — — — Q. — — — — B.f. VII, 715
D. elegans, Swartz, synops. filic. 132 & 347 (1806) ... — — — — — Q. — — — — B.f. VII, 715 M.fr. V, 118; VIII, 157.
D. pyxidata, Cavanilles, descripcion de las plantas n. 694 (1802) ... — — — — V. N.S.W. Q. — B.f. VII, 716 M.fr. V, 118.
D. pedata, Smith in Mém. de l'Ac. de Turin. V, 414 (1791) ... — — — — — Q. — — — — B.f. VII, 716 M.fr. V, 118.
D. dubia, R. Brown, prodr. 157 (1810) ... — — — — T. V. N.S.W. Q. — B.f. VII, 716 M.fr. V, 118.
D. flaccida, R. Brown, prodr. 157 (1810) ... — — — — — Q. — — — — B.f. VII, 717 M.fr. V, 118; VII, 156.
D. tripinnata, F. v. M. in Benth. Fl. Austr. VII, 717 (1878) ... — — — — — Q. — — — — B.f. VII, 717
- VITTARIA, Smith in Mém. de l'Acad. de Turin. V, 414 (1791).
V. elongata, Swartz, synops. filic. 109 & 302 (1806) ... — — — — N.S.W. Q. — B.f. VII, 718 M.fr. V, 122; VII, 121.
- LINDSAYA, Dryander in Mém. de l'Acad. de Turin. V, 413 (1791). (Lindsaea, Isoloma, Synophlebium, Schizoloma.)
L. linearis, Swartz, synops. filic. 118 & 318, t. 3 (1806) ... — — — — W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 719 M.fr. V, 119.
L. dimorpha, Bailey, Queensland Ferns 19 (1874) ... — — — — — Q. — — — — B.f. VII, 719
L. cultrata, Swartz, synops. filic. 119 (1806) ... — — — — — Q. — — — — B.f. VII, 719
L. flabellulata, Dryander in Transact. Linn. Soc. III, 41, t. 8 (1797) ... — — — — — Q. N.A. B.f. VII, 720 M.fr. V, 119.
L. lobata, Poiret, Encycl. méth. XI (1813) ... — — — — — Q. — — — — B.f. VII, 720
L. trichomanoides, Dryander in Trans. Linn. Soc. III, 43, t. 11 (1797) ... — — — — T. — — — — B.f. VII, 720 M.fr. V, 118.
L. microphylla, Swartz, synops. filic. 120 (1806) ... — — — — N.S.W. Q. — B.f. VII, 721 M.fr. V, 119.
L. incisa, Prentice in Trim. Journ. of Bot. 295 (1873) ... — — — — — Q. — — — — B.f. VII, 721
L. Fraseri, Hooker, spec. filic. I, 221, t. 70 (1846) ... — — — — — Q. — — — — B.f. VII, 721 M.fr. V, 118.

- L. ensifolia*, Swartz, synops. filic. 118 (1806) Q. N.A. B.f. VII, 721 M.fr. V, 118.
L. lanuginosa, Wallich, numerical list n. 154 (1828) Q. — B.f. VII, 722 M.fr. V, 118.
- ADIANTUM**, Tournefort, in 543, t. 317 (1700), from Hippocrates, Theophrastus, Dioscorides and Plinius.
A. lunulatum, N. Burmann, Fl. Ind. 235 (1768) Q. N.A. B.f. VII, 723 M.fr. V, 119.
A. Capillus veneris, Linné, spec. plant. 1096 (1753) Q. — B.f. VII, 723
A. Aethiopicum, Linné, syst. nat. edit. decim. n. 15 (1759) ... W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 724 M.fr. V, 119.
A. formosum, R. Brown, prodr. 155 (1810) V. N.S.W. Q. — B.f. VII, 724 M.fr. V, 120; VIII, 157.
A. affine, Willdenow, spec. plant. V, 448 (1810) N.S.W. Q. — B.f. VII, 724 M.fr. V, 119.
A. diaphanum, Blume, flora Javae, filic. 215 (1836) N.S.W. Q. — B.f. VII, 725 M.fr. VII, 121.
A. hispidulum, Swartz, syn. filic. 124 & 321 (1806) V. N.S.W. Q. — B.f. VII, 725 M.fr. V, 120.
- CHEILANTHES**, Swartz, syn. filic. 126, t. 3 (1806). (Notholaena, Nothochlaena.)
C. Prenticei, Luerssen in Uhlw. & Behr. Bot. Centralblatt IX, 440 (1882) N.A. —
C. pumilio, F. v. M.; Notholaena, R. Brown, prodr. 146 (1810) Q. — B.f. VII, 773 M.fr. VIII, 175.
C. fragillima, F. v. M., fragm. V, 123 (1866) N.A. B.f. VII, 774 M.fr. V, 123.
C. vellea, F. v. M., fragm. V, 123 (1866) ... W.A. S.A. — V. N.S.W. Q. N.A. B.f. VII, 773 M.fr. V, 123; VIII, 176.
C. distans, A. Braun, index sem. hort. Berol. (1859) ... W.A. S.A. — V. N.S.W. Q. — B.f. VII, 774 M.fr. V, 122; VIII, 176.
C. tenuifolia, Swartz, syn. filic. 126 (1806) .. W.A. S.A. T. V. N.S.W. Q. N.A. B.f. VII, 726 M.fr. V, 122.
- PTERIS**, Linné, syst. nat. 9 (1735); Linné, gen. plant. 322 (1737.) (Pellaea, Cheiloplecton, Platyloma, Litobrochia.)
P. geraniifolia, Raddi, syn. filic. Brasil. 46 (1819) N.S.W. Q. — B.f. VII, 728 M.fr. V, 124.
P. paradoxa, Baker in Benth. Fl. Austr. VII, 729 (1878) N.S.W. Q. — B.f. VII, 729
P. falcata, R. Brown, prodr. 154 (1810) T. V. N.S.W. Q. — B.f. VII, 729 M.fr. V, 123.
P. rotundifolia, G. Forster, florul. ins. Austr. prodr. 79 (1786) Q. — B.f. VII, 730
P. longifolia, Linné, spec. plant. 1074 (1753) V. N.S.W. Q. — B.f. VII, 730 M.fr. V, 126.
P. ensiformis, N. Burmann, Fl. Ind. 230 (1768) N.S.W. Q. — B.f. VII, 730 M.fr. V, 123.
P. umbrosa, R. Brown, prodr. 154 (1810) V. N.S.W. Q. — B.f. VII, 730 M.fr. V, 126.
P. quadriaurita, Retzius, observ. bot. VI, 38 (1791) Q. — B.f. VII, 731 M.fr. V, 125.
P. arguta, Aiton, hort. Kew. III, 458 (1789) T. V. N.S.W. Q. — B.f. VII, 731 M.fr. V, 126.
P. aquilina, Linné, spec. plant. 1075 (1753) ... W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 731 M.fr. V, 126.
P. incisa, Thunberg, prodr. pl. Capens. 171 (1800) S.A. T. V. N.S.W. Q. — B.f. VII, 732 M.fr. V, 124; VIII, 158.
P. marginata, Bory. voy. dans les îles d'Afr. II, 192 (1804) Q. — B.f. VII, 733 M.fr. V, 125.
P. comans, G. Forster, florul. ins. Austr. prodr. 79 (1786) T. V. N.S.W. Q. — B.f. VII, 733 M.fr. V, 125; VII, 121.
- LOMARIA**, Willdenow in Magaz. der Ges. naturf. Freunde zu Berlin, III, 160 (1809.) (Stegania, Plagiogyria.)
L. Patersoni, Sprengel, syst. veg. IV, 62 (1827) T. V. N.S.W. Q. — B.f. VII, 734 M.fr. V, 122; VII, 158.
L. vulcanica, Blume, fl. Javae, filic. 202 (1836) T. — B.f. VII, 735 M.fr. V, 121.
L. discolor, Willdenow, spec. plant. V, 293 (1810) S.A. T. V. N.S.W. Q. — B.f. VII, 735 M.fr. V, 121.
L. lanceolata, Sprengel, syst. veg. IV, 62 (1827) S.A. T. V. N.S.W. Q. — B.f. VII, 735 M.fr. V, 121.
L. attenuata, Willdenow, spec. plant. V, 290 (1810) N.S.W. — B.f. VII, 736
L. alpina, Sprengel, syst. veg. IV, 62 (1827) T. V. N.S.W. — B.f. VII, 736 M.fr. V, 121.
L. fluviatilis, Sprengel, syst. veg. IV, 65 (1827) T. V. N.S.W. — B.f. VII, 736 M.fr. V, 121.
L. Fullagari, F. v. M., fragm. VIII, 157 (1874) N.S.W. — B.f. VII, 737 M.fr. VIII, 157.
L. Capensis, Willdenow, spec. plant. V, 291 (1810) S.A. T. V. N.S.W. Q. — B.f. VII, 737 M.fr. V, 121.
L. euphlebia, Kunze in Mohl & Schlecht. Bot. Zeit. VI, 521 (1849) Q. — B.f. VII, 738 M.fr. VI, 124.
- BLECHNUM**, Linné, spec. plant. 1077 (1753.)
B. cartilagineum, Swartz, syn. filic. 114 & 312 (1806) V. N.S.W. Q. — B.f. VII, 738 M.fr. V, 120.
B. laevigatum, Cavanilles, description de las plantas 650 (1802) N.S.W. — B.f. VII, 739 M.fr. V, 120.
B. serrulatum, C. Richard in Annal. du Mus. d'Hist. nat. Par. I, 114 (1802) N.S.W. Q. N.A. B.f. VII, 739 M.fr. V, 120.
B. orientale, Linné, syst. nat. ed. decim. (1759) Q. N.A. B.f. VII, 740
- MONOGRAMMA**, Commerçon in Schkuhr's Kryptog. Gew. 82, t. 87 (1809). (Monogramme, Vaginularia, Pleurogramme, Didymopteris.)
M. Junghuhnii, Hooker, spec. filic. V, 123, t. 289 (1864) Q. — B.f. VII, 740 M.fr. VII, 119.
- WOODWARDIA**, Smith in mém. de l'Acad. de Turin. V, 411 (1791). (Doodia.)
W. aspera, Mettenius, fil. hort. bot. Lips. 65 (1856) V. N.S.W. Q. — B.f. VII, 740 M.fr. V, 130.
W. caudata, Cavanilles, description de las pl. n. 653 (1802) T. V. N.S.W. Q. N.A. B.f. VII, 742 M.fr. V, 129; VIII, 158.
- ASPLENIUM**, Linné, gen. plant. 322 (1737) from C. Bauhin (1651.) (Scolopendrium, Allantodea, Diplazium, Callipteris, Anisogonium, Thamnopteris, Neottopteris, Darea, Caenopteris, Athyrium, Diplora, Asplenium.)
A. Nidus, Linné, spec. plant. 1079 (1753) V. N.S.W. Q. — B.f. VII, 744 M.fr. V, 130.
A. simplicifrons, F. v. M., fragm. V, 74 (1865) Q. — B.f. VII, 744 M.fr. V, 74, 130; VIII, 158.
A. attenuatum, R. Brown, prodr. 150 (1810) N.S.W. Q. — B.f. VII, 745 M.fr. V, 130.
A. Trichomanes, Linné, spec. plant. 1080 (1753) T. V. N.S.W. — B.f. VII, 745 M.fr. V, 131.
A. flabellifolium, Cavanilles, description de las pl. n. 636 (1802) W.A. S.A. T. V. N.S.W. Q. — B.f. VII, 745 M.fr. V, 131.
A. paleaceum, R. Brown, prodr. 150 (1810) Q. — B.f. VII, 746 M.fr. V, 131; VIII, 158.
A. falcatum, Lamarck, Encycl. méth. I, 303 (1783) N.S.W. Q. — B.f. VII, 746 M.fr. V, 131; VII, 156.
A. Hookerianum, Colenso in Journ. of Nat. Sc. Tasm. II, 169 (1842) V. N.S.W. — B.f. VII, 747
A. furcatum, Thunberg, prodr. pl. Cap. 172 (1800) ... W.A. S.A. — V. N.S.W. — B.f. VII, 748 M.fr. V, 131.
A. laserpitifolium, Lamarck, Encycl. méth. I, 310 (1783) Q. — B.f. VII, 749 M.fr. V, 131.
A. maritimum, Linné, spec. plant. 1081 (1753) T. V. N.S.W. — B.f. VII, 747 M.fr. V, 132.
A. bulbiferum, G. Forster, floral. ins. Austr. prodr. 433 (1786) ... S.A. T. V. N.S.W. Q. — B.f. VII, 749 M.fr. V, 132.
A. pteridoides, Baker, syn. filic. sec. edit. 483 (1874) N.S.W. — B.f. VII, 749
A. umbrosum, J. Smith in Hook. Lond. Journ. IV, 174 (1842) T. V. N.S.W. — B.f. VII, 749 M.fr. V, 132.
A. pallidum, Blume, fl. Javae, filic. 776 (1836) Q. — M.fr. XI, 131.
A. sylvaticum, Presl, reliq. Haenk. I, 42 (1830) Q. — B.f. VII, 750 M.fr. V, 133.
A. maximum, D. Don, prodr. fl. Nepal. 8 (1825) N.S.W. Q. — B.f. VII, 751
A. polypodioides, Mettenius, fil. hort. Lips. 78 (1856) Q. — B.f. VII, 751 M.fr. V, 132.
A. melanochlamys, Hooker, spec. filic. III, 259 (1860) N.S.W. — B.f. VII, 751
A. decussatum, Swartz, spec. fil. 76 & 260 (1806) Q. — B.f. VII, 751

- CYSTOPTERIS, Bernhardt in Schrader's neuem Journ. II, 26 (1806).
 C. fragilis, Bernhardt in Schrad. Journ. I, 26 (1806) ... — — — — T. — — — — B.f. VII, 752 M.fr. V, 118.
- ASPIDIUM, Swartz in Schrader's Journ. II, 429 (1000). (Nephrodium, Nephrolepis, Polystichum, Lastraea, Sagenia, Oleandra.)
 A. cordifolium, Swartz, syn. filic. 45 (1806) ... — — — — N.S.W. Q. — — B.f. VII, 754 M.fr. V, 136.
 A. exaltatum, Swartz, syn. filic. 45 (1806) ... — — — — Q. N.A. B.f. VII, 754 M.fr. V, 136.
 A. ramosum, Palisot, Fl. d'Oware et de Benin II, 53, t. 91 (1807) ... — — — — N.S.W. Q. — — B.f. VII, 754 M.fr. V, 135.
 A. unitum, Swartz, syn. filic. 47 (1806) ... — — — — W.A. — — — — N.S.W. Q. N.A. B.f. VII, 755 M.fr. V, 137; VIII, 158.
 A. pteroides, Swartz, syn. filic. 49 (1806) ... — — — — — — — — Q. — — B.f. VII, 755
 A. molle, Swartz, syn. filic. 49 (1806) ... — — — — S.A. — — V. N.S.W. Q. N.A. B.f. VII, 756 M.fr. V, 135.
 A. truncatum, Gaudichaud in Voy. Freyc. Bot. 332, t. 10 (1826) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 756 M.fr. V, 135.
 A. confluent, Mettenius in Schlecht. Linnaea XXXVI, 125 (1863) ... — — — — — — — — Q. N.A. B.f. VII, 757 M.fr. V, 133; VIII, 158.
 A. aculeatum, Swartz in Schrad. Journ. II, 37 (1800) ... — — — — S.A. T. V. N.S.W. Q. — — B.f. VII, 757 M.fr. V, 134.
 A. aristatum, Swartz, syn. filic. 53 & 253 (1806) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 757 M.fr. V, 134.
 A. Capense, Willdenow, spec. plant. V, 267 (1810) ... — — — — T. V. N.S.W. — — — — B.f. VII, 758 M.fr. V, 134.
 A. decompositum, Sprengel, syst. IV, 109 (1827) ... — — — — S.A. T. V. N.S.W. Q. — — B.f. VII, 758 M.fr. V, 136.
 A. tenerum, Sprengel, syst. IV, 109 (1827) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 759
 A. uliginosum, Kunze in Schlecht. Linnaea XX, 6 (1847) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 759 M.fr. V, 133.
 A. hispidum, Swartz, syn. filic. 56 (1806) ... — — — — T. V. — — — — — — B.f. VII, 760 M.fr. V, 133.
- POLYPODIUM, Tournefort, inst. 540, t. 316 (1700), from Theophrastos, Dioscorides & Plinius. (Niphoebolus, Goniophlebium, Goniopteris, Phlebodium, Phegopteris, Phymatodes, Pleopeltis, Drynaria, Dictyopteris, Arthropteris, Xiphopteris, Meniscium partly; from Dodoens, l'Ecluse, Bauhin, Morison, Ray, Plumier and particularly Petiver as Polypodium.)
 P. australe, Mettenius, filic. hort. bot. Lips. 36 (1856) ... — — — — T. V. N.S.W. Q. — — B.f. VII, 762 M.fr. V, 127.
 P. Hookeri, Brackenridge, Bot. Wilk. Exped. Filic. 4 (1854) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 763
 P. contiguum, Brackenridge, Bot. Wilk. Exp. Fil. 6 t. (1854) ... — — — — — — — — Q. — — B.f. VII, 763 M.fr. V, 127.
 P. graminifolium, R. Brown, prodr. 147 (1810) ... — — — — T. V. N.S.W. — — — — B.f. VII, 764 M.fr. V, 127.
 P. tenellum, G. Forster, florul. ins. Austr. prodr. 440 (1786) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 764 M.fr. V, 126; VII, 121.
 P. proliferum, Roxburgh in Wall. num. list 11 (1828) ... — — — — — — — — N.S.W. Q. N.A. B.f. VII, 765 M.fr. V, 128.
 P. urophyllum, Wallich, numerical list 11 (1828) ... — — — — — — — — Q. — — B.f. VII, 765 M.fr. IV, 165; V, 137; VIII, [121].
 P. Hillii, Baker in Hook., syn. filic. sec. edit. 505 (1874) ... — — — — — — — — Q. — — B.f. VII, 766
 P. poecilophlebium, Hooker, spec. filic. V, 14 (1863) ... — — — — — — — — Q. — — B.f. VII, 766 M.fr. V, 127.
 P. serpens, G. Forster, florul. insul. austr. prodr. 81 (1786) ... — — — — — — — — V. N.S.W. Q. — — B.f. VII, 767 M.fr. V, 129.
 P. confluent, R. Brown, prodr. 146 (1810) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 767
 P. acrostichoides, G. Forster, florul. insul. austr. prodr. 81 (1786) ... — — — — — — — — Q. — — B.f. VII, 767 M.fr. V, 129.
 P. attenuatum, R. Brown, prodr. 146 (1810) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 768 M.fr. V, 128; V, 121.
 P. simplicissimum, F. v. M. in Hook. & Bak. syn. fil. sec. edit. 513 (1874) ... — — — — — — — — Q. — — B.f. VII, 768 M.fr. VII, 120, 156.
 P. nigrescens, Blume, Fl. Javae, filic. 161, t. 70 (1836) ... — — — — — — — — Q. — — B.f. VII, 769
 P. phymatodes, Linné, mant. pl. alt. 306 (1771) ... — — — — — — — — Q. N.A. B.f. VII, 769
 P. pustulatum, G. Forster, florul. ins. Austr. pr. 81 (1786) ... — — — — T. V. N.S.W. Q. — — B.f. VII, 769 M.fr. V, 127.
 P. scandens, G. Forster, florul. ins. Austr. pr. 81 (1786) ... — — — — — — — — V. N.S.W. Q. — — B.f. VII, 770 M.fr. V, 128.
 P. verrucosum, Wallich, numerical list 11 (1828) ... — — — — — — — — Q. — — B.f. VII, 770
 P. subauriculatum, Blume, Fl. Javae, filic. 177, t. 83 (1836) ... — — — — — — — — Q. — — B.f. VII, 770 M.fr. V, 128.
 P. rigidulum, Swartz, synops. filic. 38 & 320 (1806) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 771
 P. quercifolium, Linné, spec. plant. 1087 (1753) ... — — — — — — — — Q. N.A. B.f. VII, 772 M.fr. V, 127.
 P. irioides, Poiré, Encycl. méth. V, 513 (1804) ... — — — — — — — — Q. — — B.f. VII, 771 M.fr. V, 127.
 P. punctatum, Thunberg, fl. Japonic. 337 (1784) ... — — — — S.A. T. V. N.S.W. Q. — — B.f. VII, 764 M.fr. V, 129.
- HYPOLEPIS, Bernhardt in Schrader, neu. Journ. I, 34 (1806).
 H. tenuifolia, Bernhardt in Schrad. Journ. II, 24 (1800) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 726 M.fr. V, 129.
- GRAMMITIS, Swartz in Schrader, Journ. II, 3 & 17 (1800). (Gymnogramme, Selligenea).
 G. Reynoldsii, F. v. M. in Benth. Fl. Austr. VII, 775 (1878) ... — — — — S.A. — — — — — — B.f. VII, 775 M.fr. VIII, 175.
 G. Muelleri, Hooker in F. v. M. fragm. V, 138 (1866) ... — — — — — — — — Q. — — B.f. VII, 775 M.fr. V, 138; VI, 124; VII, [121].
 G. rutifolia, R. Brown, prodr. 146 (1810) ... — — — — W.A. S.A. T. V. N.S.W. Q. — — B.f. VII, 775 M.fr. V, 137.
 G. leptophylla, Swartz, syn. fil. 23 et 218, t. 1 (1806) ... — — — — W.A. S.A. T. V. N.S.W. — — — — B.f. VII, 776 M.fr. V, 137.
 G. pinnata, F. v. M., fragm. VI, 124 (1868) ... — — — — — — — — Q. — — B.f. VII, 776 M.fr. VI, 124.
 G. ampla, F. v. M., fragm. V, 188 (1866) ... — — — — — — — — Q. — — B.f. VII, 777 M.fr. V, 188.
- ANTROPHYUM, Kaulfuss, enum. filic. 197 (1824).
 A. reticulatum, Kaulfuss, enum. filic. 197 (1824) ... — — — — — — — — Q. — — B.f. VII, 777 M.fr. V, 138.
- ACROSTICHUM, Linné, gen. plant. 322 (1737), indicative. (Elaphoglossum, Stenochlaena, Lomariopsis, Hymenolepis, Gymnopteris, Chrysodium.)
 A. conforme, Swartz, syn. filic. 10 & 192, t. 1 (1836) ... — — — — — — — — Q. — — B.f. VII, 778 M.fr. V, 138.
 A. scandens, J. Smith in Hook. spec. fil. V, 249 (1864) ... — — — — — — — — Q. N.A. B.f. VII, 778 M.fr. VI, 124.
 A. sorbifolium, Linné, spec. plant. 1069 (1753) ... — — — — — — — — Q. — — B.f. VII, 779 M.fr. VII, 119.
 A. neglectum, Bailey in proc. Linn. Soc. of N.S.W., V, 32 (1880) ... — — — — — — — — Q. — — M.fr. XI, 131.
 A. repandum, Blume, Fl. Javae 39, t. 14 & 15 (1836) ... — — — — — — — — Q. — — B.f. VII, 779 M.fr. V, 138.
 A. aureum, Linné, spec. plant. 1069 (1753) ... — — — — — — — — N.S.W. Q. N.A. B.f. VII, 779 M.fr. V, 138; VII, 120.
 A. spicatum, Linné, fil. suppl. pl. 444 (1781) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 780 M.fr. V, 138.
 A. pteroides, R. Brown, prodr. 145 (1810) ... — — — — — — — — Q. N.A. B.f. VII, 780 M.fr. X, 76.
- PLATYTERUM, Desvaux in Mém. de la Soc. Linn. de Paris VI, 213 (1827).
 P. alpicorne, Desvaux in Mém. Soc. Linn. Par. VI, 213 (1827) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 780 M.fr. V, 139.
 P. grande, J. Smith in Hooker, Lond. Journ. III, 402 (1841) ... — — — — — — — — N.S.W. Q. — — B.f. VII, 781 M.fr. V, 139.

ADDITIONAL SPECIES.

Pachynema spheandrum, F. v. M. & Tate in Transact. R. S. of S. Austr. V, 79 (1882)	—	—	—	—	—	—	N.A.	—
Polyalthia Holtzeana, F. v. M. in Wing's S. Sc. Record II, 230 (1882)	—	—	—	—	—	—	N.A.	—
Philotheca Hassellii, F. v. M., fragm. XII, inedit.	W.A.	—	—	—	—	—	—	—
Tribulus astrocarpus, F. v. M., fragm. XII, 4 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII,
Euphorbia obliqua, Bauer in Endl. prodr. fl. Norfolk. 85 (1833)	—	—	—	—	N.S.W.	—	—	—
Phyllanthus Tatei, F. v. M. in Wing's S. Sc. Record II, 55 (1882)	—	S.A.	—	—	N.S.W.	—	—	—
Ficus Pinkiana, F. v. M. in Wing's S. Sc. Record II, 273 (1882)	—	—	—	—	—	Q.	—	—
Peperomia Baueriana, Miquel, syst. piperac. 120 (1843)	—	—	—	—	N.S.W.	—	—	—
Nepenthes Bernaysii, Bailey in proceed. L. S. of N.S.W. 185 (1880)	—	—	—	—	—	Q.	—	—
Ptilotus Polakii, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	W.A.	—	—	—	—	—	—	—
Atriplex Bunburyanum, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	—	—	—	—	—	—	—	—
Kochia prosthocochaeta, F. v. M., fragm. XII, 14 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 14
Kochia melanocoma, F. v. M., fragm. XII, 14 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 14
Bassia astracantha, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 12
Bassia tridens, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 12
Bassia Forrestiana, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 12
Bossiaea Scortechinii, F. v. M. (inedit.)	—	—	—	—	N.S.W.	Q.	—	—
Bossiaea Webbii, F. v. M. in Melb. Chemist, Dec. (1882)	W.A.	—	—	—	—	—	—	—
Isotropis Forrestii, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	W.A.	—	—	—	—	—	—	—
Daviesia arborea, F. v. M. & Scortechinii in proc. L. S. of N.S.W. VII, 221 (1882)	—	—	—	—	N.S.W.	Q.	—	—
Crotalaria striata, De Candolle, prodr. II, 131 (1825)	—	—	—	—	—	Q.	—	—
Indigofera tinctoria, Linné, spec. plant. 751 (1753)	—	—	—	—	—	Q.	—	—
Swainsona Oliverii, F. v. M. in Wing's S. Sc. Rec. II, 152 (1882)	W.A.	S.A.	—	—	—	—	—	—
Callistemon pithyoides, Miquel in Nederl. Kruidk. Arch. IV, 142 (1859)	—	—	—	—	V. N.S.W.	Q.	—	—
Melaleuca cylindracea, R. Brown in Benth. Fl. Austral. III, 146 (1866)	—	S.A.	—	—	—	—	—	—
Eucalyptus Foelscheana, F. v. M. in Melb. Chemist, Nov. (1882)	—	—	—	—	—	N.A.	—	—
Cryptandra Wayii; Trymalium, F. v. M. & Tate in Transact. R. S. of S. Austr. V, 80 (1882)	—	S.A.	—	—	—	—	—	—
Adenanthos Forrestii, F. v. M. in Wing's S. Sc. Record II, 230 (1882)	W.A.	—	—	—	—	—	—	—
Grevillea deflexa, F. v. M. inedit.	W.A.	—	—	—	—	—	—	—
Epaltes Tatei, F. v. M. inedit.	—	S.A.	—	—	—	—	—	—
PODOSPERMA, Labillardière, Nov. Holl. plant. specim. II, 35, t. 177 (1806). (Podotheca.)	—	—	—	—	—	—	—	—
Podosperma Polakii, F. v. M., fragm. XII, 21 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 21
Helipterum sterilesens, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	W.A.	—	—	—	—	—	—	—
Helipterum Forrestii, F. v. M. in Wing's S. Sc. Record II, 273 (1882)	W.A.	—	—	—	—	—	—	—
Velleya macropectra, F. v. M., fragm. XII, 22 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 22
Thunbergia fragrans, J. Koenig in Roxb. pl. of Coromand. I, 44, t. 67 (1795)	—	—	—	—	—	—	N.A.	—
ISANDRA, F. v. M., fragm. phyt. Austr. XII (inedit.)	Near Anthocercis.	—	—	—	—	—	—	—
Isandra Bancroftii, F. v. M., fr. XII (inedit.)	W.A.	—	—	—	—	—	—	—
DICLADANTHERA, F. v. M., fragm. phytogr. Austral. XII, 23 (1882).	—	—	—	—	—	—	—	—
Dicladanthera Forrestii, F. v. M., fragm. XII, 23 (1882)	W.A.	—	—	—	—	—	—	M.fr. XII, 23
Prostanthera Campbelli, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	W.A.	—	—	—	—	—	—	—
Chloanthes lepidota, F. v. M. inedit.	W.A.	—	—	—	—	—	—	—
Myoporum obscurum, Endlicher, prodr. fl. Norfolk. 54 (1833)	—	—	—	—	N.S.W.	—	—	—
Eremophila Pantoni, F. v. M. in Wing's S. Sc. Record II, 251 (1882)	W.A.	—	—	—	—	—	—	—
ERIA, Lindley in Edwards, Bot. Regist. t. 904 (1825).	—	—	—	—	—	—	—	—
Eria Fitzalanii, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	—	—	—	—	—	Q.	—	—
LUISIA, Gaudichaud in Freycin. voy. Bot. 426, t. 37 (1826).	—	—	—	—	—	—	—	—
Luisia teretifolia, Gaudichaud in Freyc. voy. 426, t. 37 (1826)	—	—	—	—	—	Q.	—	—
Prasophyllum Tepperi, F. v. M. in Tepp. pl. about Ardross., Oct. (1880)	—	S.A.	—	—	—	—	—	—
Smilax purpurata, G. Forster, florul. ins. Austr. prodr. 70 (1786)	—	—	—	—	N.S.W.	—	—	—
Pandanus Moorei, F. v. M. inedit.	—	—	—	—	N.S.W.	—	—	—
Carex haematostoma, Nees in Wight, contrib. to the Bot. of Ind. 125 (1834)	—	—	T.	—	—	—	—	—

ADDITIONAL ANNOTATIONS.

Myosurus from l'Obel (1576).
 Myosurus minimus Q.
 Ranunculus from Plinius.
 Ranunculus aquatilis N.S.W.
 Hibbertia hirsuta S.A.
 Doryphora Sassafras Q.
 Endiandra Sieberi Q.
 Tristichocalyx pubescens N.A.
 Papaver from Plinius.
 Cleome viscosa W.A.
 Capparis spinosa S.A.
 Capparis lasiantha W.A.
 Barbarea from l'Obel (1576).
 Barbarea vulgaris S.A.
 Capsella pilosula W.A.
 Senebiera = Carara, Medicus, Pflanzen-Gattung. 34 (1792), from
 Cesalpini (1583). (Coronopus partly).
 Lepidium strongylophyllum W.A.
 Lepidium leptopetalum W.A.
 Lepidium rotundum S.A.
 Cakile from l'Obel (1576).
 Hybanthus enneaspermus W.A.
 Hymenanthra Banksii S.A.
 Pittosporum revolutum M.fr.XII,4.
 Pittosporum ferrugineum M.fr.XII,4.
 Pittosporum melanospermum Q.
 Hymenosporum flavum M.fr.XII,4.
 Marianthus microphyllus M.fr.XII,2.
 Citriobatus pauciflorus N.S.W.
 Billardiera variifolia M.fr.XII,2.
 Cheiranthra linearis M.fr.XII,4.
 Cheiranthra filifolia M.fr.XII,4.
 Bergia perennis S.A.
 Polygala from Dioscorides & Plinius.
 Polygala persicarifolia Q.
 Polygala Chinensis S.A.
 Comesperma sylvestre S.A., N.S.W.
 Comesperma viscidulum S.A.
 Tribulus Hystrix W.A.
 Hedraianthera porphyropetala N.S.W.
 Tribulus macrocarpus W.A.
 Tribulus platypterus W.A.
 Sida virgata W.A.
 Sida petrophila W.A.
 Sida rhombifolia S.A.
 Sida physocalyx W.A.
 Sida lepida W.A., S.A.
 Abutilon tubulosum N.A.
 Abutilon cryptopetalum M.fr.IX,131.
 Abutilon geranioides W.A.
 Abutilon Avicennae, Gerard, gen. hist. of pl. 790 (1597).
 Urena lobata N.A.
 Hibiscus panduriformis W.A., Q.
 Hibiscus Normani M.fr.V,44.
 Hibiscus Sturtii W.A.
 Gossypium australe W.A. M.fr.V,44.
 Gossypium flaviflorum M.fr.V,44.
 Waltheria Indica W.A.
 Commerçon Fraseri Q.
 Euphorbia erythrantha W.A.
 Euphorbia pilulifera N.A.
 Monotaxis luteiflora S.A.
 Poranthera microphylla M.fr.XII,11.
 Pseudanthus micranthus M.fr.XII,11.
 Beyeria viscosa M.fr.XII,10.
 Ricinocarpus pinifolius S.A.
 Bertya rosmarinifolia Q. M.fr.XII,10.
 Bertya rotundifolia M.fr.XII,11.
 Micrantheum hexandrum M.fr.XII,11.
 Dissiliaria tricornis M.fr.XII,11.
 Phyllanthus rhytidospermus S.A. M.fr.XII,11.
 Phyllanthus rigens M.fr.XII,11.

Phyllanthus Gastroemii M.fr.XII,11.
 Phyllanthus australis M.fr.XII,11.
 Securinea Leucopyrus S.A., N.A.
 Adriana tomentosa M.fr.XII,11.
 Adriana quadripartita M.fr.XII,11.
 Tragia Novae Hollandiae M.fr.XII,11.
 Fontainea Pancheri M.fr.XII,11.
 Onalanthus populifolius M.fr.XII,11.
 Ficus platypoda W.A.
 Ficus scabra N.A.
 Ficus stenocarpa = Ficus subglabra, F. v. M.,
 fragm. IX, 152 (1875) N.S.W., Q.
 Pouzolzia quinquenervis N.A.
 Casuarina suberosa S.A.
 Casuarina inophloia N.S.W.
 Peperomia Urvilleana N.S.W.
 Leea aculeata = L. Brunoniana, Clarke in Britten's Journ. of Bot. X,
 105 (1881).
 Castanospora Alphanthi N.S.W.
 Atalaya variifolia Q.
 Dodonaea lanceolata S.A.
 Dodonaea truncatiales = D. calycina, Cunningham in A. Gray, Bot.
 of Wilk. Expl. Exped. I, 262 (1854).
 Dodonaea platyptera W.A.
 Dodonaea microzyga S.A.
 Bueseraceae = Burseraceae.
 Celastrus bilocularis N.S.W.
 Stackhousia muricata S.A.
 Plumbago Zeilanica S.A.
 Aegialitis annulata N.A.
 Portulaca from W. Turner (1538).
 Sagina apetala N.S.W.
 Saponaria from Bock (1539).
 Ptilotus incanus S.A.
 Ptilotus psilotrichoides N.A.
 Ptilotus helipteroides W.A.
 Ptilotus rotundifolius W.A.
 Ptilotus latifolius W.A.
 Rhagodia Billardieri N.A. M.fr.XII,16.
 Rhagodia parabolica W.A., V. M.fr.XII,15.
 Rhagodia crassifolia Q. M.fr.XII,15.
 Rhagodia Preissii S.A. M.fr.XII,15.
 Rhagodia Gaudichaudii M.fr.XII,15.
 Rhagodia spinescens M.fr.XII,15.
 Rhagodia nutans W.A., N.A. M.fr.XII,15.
 Rhagodia linifolia M.fr.XII,15.
 Chenopodium nitrariaceum M.fr.XII,16.
 Chenopodium microphyllum M.fr.XII,16.
 Chenopodium rhadinostachyum (1882).
 Chenopodium cristatum M.fr.XII,17.
 Chenopodium atriplicinum M.fr.XII,17.
 Chenopodium carinatum N.A. M.fr.XII,17.
 Dysphania plantaginella W.A.
 Dysphania litoralis M.fr.XII,17.
 Dysphania myriocephala M.fr.XII,17.
 Atriplex from W. Turner (1538).
 Atriplex prostratum includes A. microcarpum
 and A. Pumilio M.fr.XII,17.
 Atriplex fissivalve M.fr.XII,16.
 Atriplex paludosum M.fr.XII,16.
 Atriplex halimoides W.A., N.A. M.fr.XII,17.
 Atriplex vesicarium W.A. M.fr.XII,17.
 Atriplex holocarpum M.fr.XII,17.
 Atriplex nummularium M.fr.XII,17.
 Atriplex leptocarpum M.fr.XII,17.
 Atriplex crystallinum M.fr.XII,17.
 Kochia sedifolia W.A. M.fr.XII,15.
 Kochia lobiflora M.fr.XII,15.
 Kochia lanosa M.fr.XII,15.
 Kochia triptera M.fr.XII,15.
 Kochia oppositifolia M.fr.XII,15.
 Kochia brevifolia M.fr.XII,15.

<i>Kochia humillima</i>	M.fr. XII, 15.	<i>Melaleuca armillaris</i>	Q.
<i>Kochia planifolia</i>	M.fr. XII, 15.	<i>Eucalyptus amygdalina</i>	S.A.
<i>Kochia microphylla</i>	M.fr. XII, 15.	<i>Eucalyptus macrorrhyncha</i>	S.A.
<i>Kochia brachyptera</i>	M.fr. XII, 15.	<i>Eucalyptus capitellata</i>	Q.
<i>Kochia ciliata</i>	Q. M.fr. XII, 15.	<i>Eucalyptus clavigera</i>	Q.
<i>Kochia stelligera</i>	Q. M.fr. XII, 15.	<i>Eucalyptus tessellaris</i>	S.A.
<i>Babbagia dipterocarpa</i>	M.fr. XII, 15.	<i>Eucalyptus populifolia</i>	N.A.
<i>Bassia tricornis</i>	S.A.	<i>Eucalyptus microtheca</i>	N.A.
<i>Bassia carnosia</i>	M.fr. XII, 13.	<i>Pomaderris prunifolia</i>	Q.
<i>Bassia sclerolaenoides</i>	M.fr. XII, 13.	<i>Pomaderris myrtilloides</i>	S.A.
<i>Bassia bicornis</i>	M.fr. XII, 13.	<i>Astrotricha Hamptoni</i>	N.A.
<i>Bassia paradoxa</i>	M.fr. XII, 13.	<i>Didiscus hemiarpus</i> = <i>D. setulosus</i> , F. v. M. in Proceed. R. S. Tasm	
<i>Bassia salsuginosa</i>	M.fr. XII, 13.	III, 238 (1857).	
<i>Bassia lanicuspis</i>	W.A., N.A. M.fr. XII, 13.	<i>Trachymene cirrosa</i> (not in S.A. and V.) ...	W.A.
<i>Bassia biflora</i>	Q. M.fr. XII, 13.	<i>Trachymene heterophylla</i> (not in W.A.) ...	V.
<i>Bassia echinopsila</i>	W.A. M.fr. XII, 13.	<i>Actinotus Gibbonsii</i>	Q.
<i>Bassia encchylaenoides</i>	Q. M.fr. XII, 13.	<i>Oenanthe</i> from l'Obel (1576).	
<i>Enchylaena tomentosa</i>	M.fr. XII, 15.	<i>Choretum lateriflorum</i>	Q.
<i>Salicornia</i> from Gerard (1597).		<i>Viscum</i> from W. Turner (1538).	
<i>Salicornia robusta</i>	W.A., N.A. M.fr. XII, 18.	<i>Viscum angulatum</i>	N.A.
<i>Salicornia arbuscula</i>	M.fr. XII, 18.	<i>Viscum articulatum</i>	S.A.
<i>Salicornia cinerea</i>	M.fr. XII, 18.	<i>Isopogon ceratophyllus</i>	W.A.
<i>Salicornia australis</i>	M.fr. XII, 18.	<i>Isopogon asper</i> (not in S.A., T., V.) ...	W.A.
<i>Salsola Kali</i>	M.fr. XII, 18.	<i>Persoonia lanceolata</i>	Q.
<i>Tetragonia expansa</i>	W.A.	<i>Grevillea agrifolia</i>	S.A.
<i>Zaleya decandra</i>	W.A.	<i>Grevillea ilicifolia</i>	Q.
<i>Trianthema turgidifolia</i>	W.A.	<i>Grevillea juncifolia</i>	N.A.
<i>Trianthema crystallina</i>	W.A.	<i>Grevillea ramosissima</i>	V.
<i>Trianthema pilosa</i>	W.A.	<i>Grevillea Chrysodendron</i>	W.A.
<i>Macarthuria australis</i>	M.fr. II, 11.	<i>Grevillea stenobotrya</i>	W.A.
<i>Mollugo Glinus</i> = <i>M. Glinus</i> .		<i>Grevillea striata</i>	W.A.
<i>Mollugo Cerviana</i>	W.A., N.A.	<i>Hakea Cunninghamii</i>	W.A.
<i>Polygonum barbatum</i>	N.A.	<i>Hakea saligna</i>	V.
<i>Didymotheca pleiococca</i>	N.A.	<i>Hakea microcarpa</i>	Q.
<i>Gastrolobium grandiflorum</i>	S.A.	<i>Hakea multilineata</i>	N.A.
<i>Mirbelia speciosa</i>	Q.	<i>Banksia serrata</i>	Q.
<i>Mirbelia daviesioides</i> = <i>M. aphylla</i> , F. v. M., fragm. IV, 11 (1863).		<i>Pimelea sanguinea</i>	N.A.
<i>Gompholobium latifolium</i>	V.	<i>Pimelea trichostachya</i>	S.A.
<i>Aotus villosa</i>	S.A.	<i>Pimelea elachantha</i>	S.A.
<i>Crotalaria medicaginea</i>	S.A.	<i>Pimelea amnocharis</i>	S.A.
<i>Psoralea cephalantha</i>	N.A.	<i>Ixora Dallachyana</i>	N.A.
<i>Psoralea balsamica</i>	S.A.	<i>Canthium coprosmodies</i>	N.A.
<i>Psoralea Schultzii</i>	N.A.	<i>Canthium latifolium</i>	W.A.
<i>Indigofera monophylla</i>	W.A.	<i>Oldenlandia</i> (Anotis).	
<i>Tephrosia flammea</i>	N.A.	<i>Oldenlandia scleranthoides</i> , Benth. & J. Hooker, gen. pl. II, 88 (1873).	
<i>Tephrosia juncea</i>	N.A.	<i>Oldenlandia galioides</i>	W.A.
<i>Sesbania aculeata</i>	W.A.	<i>Opercularia varia</i>	Q.
<i>Swainsona coronillifolia</i>	S.A.	<i>Asperula</i> from Dalechamps (1554).	
<i>Swainsona colutooides</i>	W.A.	<i>Galium umbrosum</i>	W.A.
<i>Aeschynomene Indica</i>	S.A., N.S.W. M.fr. XII, 18.	<i>Sambucus</i> from Plinius.	
<i>Aeschynomene falcata</i>	M.fr. XII, 18.	<i>Trichosanthes cucumerina</i>	Q.
<i>Aeschynomene Americana</i>	M.fr. XII, 18.	<i>Cucumis trigonus</i>	W.A.
<i>Uria lagopoides</i>	N.A.	<i>Eupatorium</i> from Gerard (1597).	
<i>Lespedeza lanata</i>	N.A.	<i>Lagenophora Emphysopus</i>	S.A.
<i>Erythrina vespertilio</i>	W.A.	<i>Brachycome Stuartii</i>	Q.
<i>Canavalia obtusifolia</i>	W.A.	<i>Brachycome graminea</i> , F. v. M., first gen.	
<i>Vigna lanceolata</i>	W.A.	Rep. 14 (1853)	Q.
<i>Rhynchosia minima</i>	W.A., S.A.	<i>Brachycome ciliaris</i>	Q.
<i>Caesalpinia Bonducella</i> (1810).		<i>Calotis microcephala</i>	Q.
<i>Labichea Buettneriana</i>	M.fr. XII, 18.	<i>Aster ramulosus</i>	Q.
<i>Labichea punctata</i>	M.fr. XII, 19.	<i>Vittadinia macrorrhiza</i>	Q.
<i>Labichea rupestris</i>	M.fr. XII, 19.	<i>Podocoma cuneifolia</i>	W.A.
<i>Adenantha abrosperma</i>	N.A.	<i>Pterigeron liatroides</i>	N.A.
<i>Acacia iteaphylla</i>	S.A.	<i>Pterigeron microglossus</i>	S.A.
<i>Acacia dissonera</i>	N.A.	<i>Epilates</i> (Gynaphanes).	
<i>Acacia Doratoxylon</i>	N.A.	<i>Pterocaulon sphacelatus</i>	W.A.
<i>Acacia Farnesiana</i>	W.A.	<i>Podotheca fusciscens</i> = <i>Helipterum</i> , Turczaninow	
<i>Acacia triptera</i> , <i>A. rhizophylla</i> , <i>A. Oxycedrus</i> , <i>A. verticillata</i> , <i>A. Riceana</i> typographically transposed from <i>Julifera</i> to <i>Plurineres</i> .		in Bull. de Mosc. 80 (1851)	M.fr. XII, 22.
<i>Albizia Hendersoni</i>	M.fr. V, 191.	<i>Podosperma gnaphaloides</i>	M.fr. XII, 21.
<i>Alchemilla</i> from Bock (1539).		<i>Podosperma chrysanthum</i>	M.fr. XII, 21.
<i>Ammannia crinipes</i> = <i>Koehnea crinipes</i> , F. v. M. (inedit.) = <i>Nesaea</i> , sect. <i>Crinipedium</i> , Koehne in Engler, bot. Jahrb. III, 337 (1882).		<i>Podosperma angustifolium</i>	M.fr. XII, 21.
<i>Lagerstroemia Flos Reginae</i>	M.fr. XII, 20.	<i>Leptorhynchus squamatus</i>	Q.
<i>Ceratophyllum demersum</i>	N.A.	<i>Helipterum Kendallii</i> = <i>Podolepis Kendallii</i> , F. v. M. (inedit.)	
<i>Bruguiera caryophylloides</i>	Q.	<i>Helipterum Tietkensii</i>	W.A., N.S.W.
<i>Thryptomene cricaea</i>	V.	<i>Helichrysum dealbatum</i>	V.
<i>Baeckea densifolia</i>	Q.	<i>Helichrysum baccharoides</i>	N.S.W.
<i>Kunzea peduncularis</i>	Q.	<i>Helichrysum pholidotum</i> = <i>Humea squamata</i> .	
		<i>Eclipta alba</i>	N.A.
		<i>Wedelia verbesinoides</i>	S.A.
		<i>Bidens bipinnatus</i>	S.A.
		<i>Flaveria Australasica</i>	W.A., S.A.

Erechtites mixa	S.A., V.	Wolffia Michellii	S.A.
Erechtites prenanthoides	S.A.	Cyperus Eragrostis	M.fr. XII, 25.
Candollea pilosa, Labillardière in Annal. du Mus. VI., 453, t. 63 (1805).		Cyperus pulchellus	M.fr. XII, 25.
Candollea glauca, Labillardière in Annal. du Mus. VI., 454, t. 64 (1805).		Cyperus gracilis	M.fr. XII, 25.
Wahlenbergia (Strzeleckia).		Cyperus tenellus	M.fr. XII, 25.
Lobelia Benthamii	S.A.	Cyperus squarrosus	M.fr. XII, 25.
Isotoma petraea	N.A.	Cyperus fulvus	W.A.
Dampiera Brownii	Q.	Cyperus rotundus	M.fr. XII, 25.
Dampiera diversifolia	M.fr. I, 120.	Cyperus polystachys	M.fr. XII, 25.
Dampiera incana	N.A.	Cyperus Iria	M.fr. XII, 25.
Leschenaultia hirsuta	M.fr. XII, 23.	Cyperus decompositus	M.fr. XII, 25.
Leschenaultia longiloba	M.fr. XII, 23.	Scirpus grossus	M.fr. XII, 24.
Catosperma Muelleri	S.A.	Scirpus supinus	M.fr. XII, 25.
Scaevola depauperata	N.A.	Scirpus litoralis	M.fr. XII, 25.
Scaevola globuliflora = S. globulifera.		Cyathochaete clandestina	M.fr. XII, 25.
Goodenia Chambersii	N.A.	Cyathochaete diandra	M.fr. XII, 25.
Goodenia Hassallii = G. Hassellii.		Schoenus aphyllus	M.fr. XII, 26.
Mitrasacme pilosa	S.A.	Schoenus ericetorum	S.A.
Logania floribunda	Q.	Schoenus sphaerocephalus	S.A.
Plantago from W. Turner (1538).		Lepidospora tenuissima	M.fr. XII, 26.
Ligustrum from W. Turner (1538).		Lepidosperma gladiatum	M.fr. XII, 26.
Notelaea linearis	Q.	Lepidosperma filiforme	M.fr. XII, 26.
Cynanchum floribundum	W.A.	Cladium junceum	M.fr. XII, 25.
Gymnanthera nitida	W.A.	Cladium schoenoides	M.fr. XII, 26.
Tylophora erecta	N.A.	Cladium trifidum	M.fr. XII, 26.
Ipomoea Turpethum	N.A.	Cladium filum	M.fr. XII, 26.
Ipomoea Muelleri	S.A.	Cladium melanospermum	M.fr. XII, 26.
Convolvulus from W. Turner (1538).		Cladium Mariscus	M.fr. XII, 25.
Polymeria angusta	S.A.	Cladium teretifolium	M.fr. XII, 25.
Cuscuta from Bock (1539).		Cladium psittacorum	M.fr. XII, 25.
Cuscuta australis	W.A., S.A.	Cladium Radula	S.A.
Solanum viride, Solander in G. Forster, prodr. 89 (1786).		Caustis pentandra	M.fr. XII, 26.
Datura Leichhardtii	W.A.	Arthrostylis aphylla	M.fr. XII, 25.
Lindernia, Allioni in misc. Taurin. III, 178 (1766), from Gagnebin (Pyxidaria).		Carex Brownii	Q. M.fr. XII, 26.
Peplidium Muelleri	W.A., S.A.	Carex tereticaulis	M.fr. XII, 26.
Buechnera parviflora	W.A.	Carex declinata	M.fr. XII, 26.
Coldenia procumbens	S.A.	Carex breviculmis	M.fr. XII, 26.
Pollichia latiseipalea	Q., N.A.	Carex inversa	M.fr. XII, 26.
Clerodendrum lanceolatum	W.A.	Carex acuta	M.fr. XII, 26.
Spartothamnus teucriiflorus	W.A.	Carex pumila	M.fr. XII, 26.
Eremophila oppositifolia	W.A.	Tragus racemosus	W.A.
Eremophila Mitchellii	S.A.	Andropogon Gryllus	W.A.
Eremophila exilifolia	W.A.	Andropogon procerus	W.A.
Eremophila Freelingii	W.A.	Anthistiria membranacea	W.A.
Eremophila Macdonnellii	N.A.	Chionachne cyathopoda	Q.
Eremophila Goodwinii	N.A.	Sporobolus Lindleyi	N.A.
Eremophila platycalyx	S.A.	Eriachne obtusa, W.A., S.A.	
Eremophila alternifolia	W.A.	Eleusine Aegyptia = E. cruciata, Lamarck, tabl. encycl. et meth. I, 203 t. 48 (1791)	W.A.
Eremophila scoparia	W.A.	Setaria glauca	W.A.
Prasophyllum Fimbria = P. Reichenbachii, F. v. M. (inedit.)		Setaria macrostachya	W.A.
Chiloglottis Gunnii	N.S.W.	Eragrostis pilosa	W.A.
Astelia (Funkia 1805).		Eragrostis tenella	W.A.
Typhonium Brownii	N.A.	Trirhaphis pungens	Q.
		Arundo Phragmites	W.A.
		Botrychium ternatum	S.A.

SEQUENCE OF ORDERS ACCORDING TO PREDOMINANCE, WITH INDICATIONS OF THEIR NUMBERS OF SPECIES.

Total number of Vasculares=8646. Dicotyledoneae=6897. Monocotyledoneae=1522. Acotyledoneae=227.

Leguminosae ... 1058	Tiliaceae ... 52	Rosaceae ... 17	Ericaceae ... 5
Myrtaceae ... 651	Loganiaceae ... 51	Anonaceae ... 16	Nymphaeaceae ... 4
Proteaceae ... 586	Apocynaceae ... 48	Salicariaceae ... 16	Magnoliaceae ... 4
Compositae ... 529	Droseraceae ... 46	Junceae ... 16	Samydaceae ... 4
Cyperaceae ... 372	Santalaceae ... 43	Monimiceae ... 15	Elatineae ... 4
Gramineae ... 346	Pittosporaceae ... 38	Celastrinae ... 15	Lineae ... 4
Epacridaceae ... 273	Lauraceae ... 37	Olacinae ... 15	Cupuliferae ... 4
Orchideae ... 255	Meliaceae ... 36	Ebenaceae ... 15	Plumbaginaceae ... 4
Euphorbiaceae ... 224	Saxifragaceae ... 35	Menispermaceae ... 14	Plantaginaceae ... 4
Goodeniaceae ... 212	Campanulaceae ... 34	Stackhousiaceae ... 13	Dioscorideae ... 4
Filices ... 200	Polygaleae ... 32	Cycadeae ... 13	Gesneriaceae ... 3
Rutaceae ... 185	Portulacaceae ... 32	Violaceae ... 12	Pedalinae ... 3
Liliaceae ... 161	Acanthaceae ... 30	Myrsinaceae ... 12	Philydreae ... 3
Rubiaceae ... 124	Coniferae ... 29	Phytolaccaeae ... 11	Nepenthaceae ... 2
Labiatae ... 124	Fluviales ... 29	Scitamineae ... 11	Guttiferae ... 2
Sterculiaceae ... 123	Combretaceae ... 27	Piperaceae ... 10	Malpighiaceae ... 2
Salsolaceae ... 112	Ficoideae ... 27	Aroideae ... 10	Burseraceae ... 2
Malvaceae ... 105	Palmae ... 26	Pandaneae ... 10	Connaraceae ... 2
Umbelliferae ... 103	Polygonaceae ... 25	Anacardiaceae ... 9	Caprifoliaceae ... 2
Sapindaceae ... 100	Loranthaceae ... 25	Hydrocharideae ... 9	Aquifoliaceae ... 2
Candolleaceae ... 95	Irideae ... 25	Xyrideae ... 9	Styraceae ... 2
Dilleniaceae ... 95	Caryophylleae ... 24	Rhizospermeae ... 9	Hydrophyllaeae ... 2
Amarantaceae ... 94	Capparidaceae ... 23	Flacourtiaceae ... 8	Burmanniaceae ... 2
Restiaceae ... 93	Casuarineae ... 23	Simarubeae ... 7	Typhaceae ... 2
Rhamnaceae ... 89	Cucurbitaceae ... 23	Geraniaceae ... 7	Myristicaceae ... 1
Amaryllideae ... 86	Gentianeae ... 23	Frankeniaceae ... 7	Papaveraceae ... 1
Solanaceae ... 80	Lentibularinae ... 23	Bignoniaceae ... 7	Hypericinae ... 1
Scrophularinae ... 78	Araliaceae ... 21	Crassulaceae ... 6	Ochnaceae ... 1
Myoporinae ... 78	Zygophylleae ... 20	Rhizophoreae ... 6	Podostemoneae ... 1
Verbenaceae ... 77	Jasminae ... 20	Melastomaceae ... 6	Balonophoreae ... 1
Thymeleae ... 74	Sapotaceae ... 19	Primulaceae ... 6	Hamamelidae ... 1
Convolvulaceae ... 67	Commelineae ... 19	Lemnaceae ... 6	Elaeagneae ... 1
Urticaceae ... 63	Viniferae ... 18	Alismaceae ... 6	Cornaceae ... 1
Haloragaeae ... 60	Eriocaulaeae ... 18	Aristolochiaeae ... 5	Orobanchaeae ... 1
Asclepiadeae ... 60	Lycopodinae ... 18	Nyctagineae ... 5	Taccaceae ... 1
Cruciferae ... 54	Ranunculaceae ... 17	Onagreae ... 5	Roxburghiaceae ... 1
Asperfoliae ... 53	Tremandreae ... 17	Passifloreae ... 5	Pontederiaceae ... 1

Total number of Genera of Vasculares=1355.

INDEX OF ORDERS AND GENERA.

Abbotia 74	Aciphylla 53	Adenanthos 66, 140	Anigozanthos 116
Abelmoschus 14	Ackama 48	Adenochilus 113	Agrimonia 47
Abildgaardia 125	Acmene 59	Adenosma 97	Agropyron 135
Abroma 15	Acomis 82	Adenostemma 77	Agrostis 133
Abrophyllum 48	Acrachne 134	Adenostephanus 68	Agrostocrinum 118
Abrotanella 84	Acradenia 12	Adiantum 137	Ailantus 12
Abrus 41	Acroclinium 80	Adrastaea 1	Aillya 88
Abutilon 14, 141	Acronychia 12	Adriana 21, 141	Aira 133
Acacia 43, 142	Acrophyllum 48	Aegialinites 26	Aizoon 31
Acaena 47	Acrostichum 139	Aegialitis 26, 141	Ajuga 102
Acalypha 20	Acrotriche 105	Aegiceras 91	Akania 24
Acanthaceae 99	Actephila 19	Aegopogon 133	Albizzia 47, 142
Acanthocarpus 119	Actinocarpus 121	Aeschynomene 40, 142	Aldrovanda 7
Acanthocladium 81	Actinodium 51	Azelia 43	Alchemilla 47, 142
Acanthus 99	Actinopappus 82	Agastachys 67	Alchornea 21
Achilleopsis 16	Actinostachys 136	Agathis 109	Alectryon 24
Achras 91	Actinostigma 10	Agati 39	Alepyrum 123
Achromolaena 82	Actinostrobus 109	Ageratum 77	Aleurites 20
Achrysum 83	Actinotus 63, 142	Aglaia 9	Alisma 121
Achyranthes 28	Adansonia 15	Aglaiaopsis 9	Alismaceae 121
Acianthus 113	Adeliopsis 4	Agnostus 71	Allania 118
Acicalyptus 59	Adenantha 43, 142	Agonis 54	Allantodea 138

- Allophylus 24
 Alopecurus 132
 Alphonsonia 60
 Alpinia 114
 Alsomitra 76
 Alsophila 137
 Althenia 121
 Altingia 109
 Alstonia 93
 Alternanthera 28
 Alysicarpus 40
 Alyssum 5
 Alyxia 93
 Amanoia 20
 Amaranthaceae 28
 Amaranthus 29
 Amaryllidaceae 115
 Amblyosperma 85
 Ambrina 29
 Ammannia 49, 142
 Ammobium 82
 Amomum 114
 Amora 9
 Amorphophallus 120
 Amorphospermum 91
 Amperea 19
 Amphibolis 121
 Amphibromus 134
 Amphicosmia 137
 Amphipogon 133
 Amphodus 41
 Anacardiaceae 25
 Anadenia 68
 Anagallis 91
 Anarthria 123
 Ancana 3
 Ancistrostigma 31
 Ancistrum 47
 Andersonia 108
 Andrachne 19
 Andrewsia 104
 Andropogon 132, 143
 Androstemma 116
 Androstoma 105
 Aneilema 122
 Anemone 1
 Angianthus 82
 Angioperis 136
 Angophora 57
 Anguillaria 117
 Anisacantha 30
 Anisandra 102
 Aniseia 94
 Anisogonium 138
 Anisolepis 80
 Anisomeles 101
 Anisopogon 134
 Anisotome 63
 Anonaceae 3
 Anodopetalum 48
 Anopterus 48
 Anosporum 124
 Antennaria 80
 Anthediosorus 82
 Anthericum 118
 Anthistria 132, 143
 Anthobolus 64
 Anthocerales 82
 Anthocercis 96
 Anthotium 87
 Anthotroche 96
 Anthosachne 135
 Antiaris 22
 Anticoryne 53
 Antidesma 19
 Antirrhoea 75
 Antrophyum 139
 Aotus 35, 142
 Aphananthe 21
 Aphelia 123
 Apium 63
 Aploxaxis 85
 Apluda 132
 Apocynaceae 92
 Apodytes 64
 Apolochlamys 82
 Aponogeton 121
 Apophyllum 5
 Aporosa 20
 Apostasia 114
 Aquifoliaceae 63
 Arabis 5
 Araliaceae 61
 Araucaria 109
 Archeria 108
 Archidendron 47
 Archontophoenix 120
 Ardisia 91, 105
 Areca 120
 Arenaria 27
 Argophyllum 47
 Argyrocome 80
 Argyrodermum 15
 Argyroglottis 81
 Argyrophanes 81
 Aristida 131
 Aristolochia, 23
 Aristolochiaceae 23
 Aristotelia 17
 Arnocrinum 119
 Aroideae 120
 Artanema 97
 Arthratherum 132
 Arthraxon 131
 Arthrochilus 113
 Arthrocnemum 30
 Arthropodium 118
 Arthropteris 139
 Arthrostylis 129, 143
 Arthrotrichum 28
 Arum 120
 Arundinella 130
 Arundo 135, 143
 Aryteria 24
 Asclepiadeae 93
 Asparagus 117
 Asparagopsis 117
 Asperella 132
 Asperifoliae 99
 Asperula 76, 142
 Aspidium 138
 Asplenium 138
 Asprella 132
 Astartea 54
 Astelia 117, 143
 Aster 78, 142
 Asteridea 80
 Asterochiton 16
 Asterolasia 10
 Asteromyrtus 55
 Asterotrichon 13
 Astraea 52
 Astrebla 134
 Astroloma 105
 Astrotricha 61, 142
 Atalantia 12
 Atalaya 24, 141
 Atelandra 101
 Atherocephala 108
 Atherosperma 3
 Athrixia 80
 Athrodactylis 120
 Athrotaxis 109
 Athyrium 138
 Atkinsonia 65
 Atriplex 29, 140, 141
 Atylosia 41
 Australina 22
 Avicennia 103
 Aylmeyeria 27
 Azolla 136
 Azorella 63
 Babbagia 30, 141
 Babingtonia 53
 Backhousia 59
 Bacularia 120
 Baeca 98
 Baeckea 53, 142
 Baeobotrys 91
 Balanops 22
 Balantium 137
 Balaustion 54
 Balfouria 93
 Baloghia 21
 Balonophora 23
 Balonophoreae 23
 Bambusa 135
 Banksia 71, 142
 Barbarea 5, 141
 Barklya 42
 Barringtonia 39
 Bartlingia 36, 118
 Basilicum 100
 Bassia 30, 140, 141
 Batatas 94
 Batracherum 131
 Bauera 48
 Bauhinia 42
 Baumea 128
 Baxteria 119
 Beaufortia 56
 Bedfordia 84
 Beilschmiedia 3
 Bellis 77
 Bellendena 67
 Benincasa 76
 Berchemia 60
 Bergia 8, 141
 Berrya 17
 Bertolonia 104
 Bertya 19, 141
 Beyeria 18, 141
 Beyerlopsis 18
 Bidaria 94
 Bidens 84, 142
 Bignoniaceae 99
 Billardiera 7, 141
 Billotia 54
 Billotia 56
 Bischoffia 20
 Blackburnia 12
 Blackwellia 6
 Blancoa 116
 Blandfordia 117
 Bleasdalea 68
 Blechnum 138
 Blennodia 5
 Blennospora 83
 Blepharanthemum 13
 Blepharocarya 25
 Blitum 29
 Blumea 79
 Blyxa 115
 Boehmeria 22
 Boerhaavia 32
 Bombax 15
 Bonnaya 97
 Boronia 10
 Borya 119
 Bosistoa 11
 Bossiaea 37, 140
 Bothryodendron 61
 Botrychium 136, 143
 Bouchardatia 12
 Bowenia 110
 Brachychiton 15
 Brachycome 77, 142
 Brachyloma 107
 Brachynema 48
 Brachypterum 42
 Brachysema 32
 Brachyspatha 120
 Brachystephium 77
 Brackenridgea 10
 Bradlea 19
 Bramia 97
 Brassenia 1
 Brassia 61
 Breweria 95
 Breynia 20
 Bridelia 20
 Brizopyrum 134
 Brizula 123
 Brochosiphon 99
 Brombya 12
 Bromidium 133
 Bromus 135
 Brucea 12
 Bruguiera 50, 142
 Brunella 101
 Brunonia 87
 Bryonia 76
 Bryonopsis 76
 Buchanania 25
 Buckinghamia 71
 Buechnera 98, 143
 Bulbine 118
 Bulbophyllum 110
 Bulliardia 48
 Bulweria 99
 Bunya 103
 Burchardia 117
 Burmannia 115
 Burmanniaceae 115
 Burnettia 113
 Burgesia 32
 Bursaria 7
 Burseraceae 25, 151
 Burtonia 34
 Busbeckea 5
 Butomopsis 121
 Byblis 8
 Byronia 63
 Cabomba 1
 Cadaba 5
 Cadellia 13
 Caesalpinia 42, 142
 Caesia 118
 Cajanus 41
 Cakile 6, 141
 Caladenia 113
 Caladium 120
 Calamus 119
 Calandrinia 27
 Calanthe 111
 Caldesia 121
 Caleana 113
 Calcectasia 119
 Caletia 18
 Caleyia 113
 Calliandra 47
 Callicarpa 103
 Callicoma 48
 Callicystus 41
 Callipteris 138
 Callistachys 32
 Callistemon 55, 140
 Callitriche 50
 Callitris 109
 Calocephalus 83
 Calochilus 112
 Calogyne 88
 Calomeria 82
 Calonyction 94
 Calopetalon 7
 Calophyllum 8
 Calosciadium 63
 Calothamnus 56
 Calorophus 124
 Calostemma 116
 Calostrophus 124
 Calotis 78, 142
 Caltha 1

- Calycoplepus 18
 Calycotrix 52
 Calyptranthes 59
 Calyptrastegia 73
 Calyptrastigma 18
 Calystegia 95
 Campanulaceae 85
 Camphoromyrtus 53
 Camptostemon 15
 Campylanthera 7
 Campynema 115
 Cananga 3
 Canarium 25
 Canarvonia 71
 Canavalia 41,142
 Candollea 1,85,143
 Candolleaceae 85
 Canscora 90
 Cansjera 63
 Canthium 75,142
 Capparideae 4
 Capparis 5,141
 Caprifoliaceae 76
 Capsella 6,141
 Carallia 50
 Carapa 9
 Carara 141
 Cardamine 5
 Cardiospermum 24
 Cardwellia 71
 Carex 129,140,143
 Careya 60
 Cargillia 92
 Carissa 93
 Carmichaelia 39
 Carpha 127
 Carpodontos 48
 Carronia 4
 Cartonema 122
 Carumbium 21
 Caryodaphne 3
 Caryophylleae 27
 Caryospermum 26
 Caryota 120
 Casearia 6
 Cassia 42
 Cassinia 82
 Cassiniola 80
 Cassytha 4
 Castanospermum 42
 Castanospora 24,141
 Casuarina 22,141
 Casuarineae 22
 Catakidozamia 110
 Cathartocarpus 42
 Cathormion 47
 Catosperma 87,142
 Caulinia 41, 121
 Caustis 129
 Cedrela 9
 Celastrineae 26
 Celastrus 26,141
 Celmisia 78
 Celosia 29
 Celtis 21
 Cenarrhenes 67
 Cenchrus 130
 Centaurea 85
 Centella 63
 Centipeda 84
 Centranthera 98
 Centrathium 77
 Centrolepis 123
 Centropappus 84
 Centrotheca 134
 Centunculus 91
 Cephaelis 75
 Cephalipterum 83
 Cephalosorus 82
 Cephalotus 48
 Ceranthes 92
 Ceratogyne 84
 Ceratopetalum 48
 Ceratophyllum 50,142
 Ceratopteris 137
 Cerbera 93
 Cercodia 49
 Ceriops 50
 Ceropogia 94
 Cesatia 62
 Chaetanthus 124
 Chaetaria 132
 Chaetospira 127
 Chalcas 12
 Chamaeraphis 130
 Chamaecilla 118
 Chamaesphaerion 83
 Chamaexeros 119
 Chamaelaucium 51
 Chapelliera 127
 Charlwoodia 117
 Cheilanthes 138
 Cheilococca 37
 Cheiloplecton 138
 Cheiranthra 7,141
 Cheiroloma 78
 Chenolea 30
 Chenopodina 30
 Chenopodium 29,141
 Cheynia 54
 Chilocarpus 92
 Chiodia 101
 Chiloglottis 114,143
 Chionachne 132,143
 Chionanthus 92
 Chithonanthus 43
 Chlamydia 117
 Chlamyspermum 118
 Chloanthes 103,140
 Chloris 134
 Chlorophytum 118
 Chondrachne 127
 Choretrum 64,142
 Choriceras 19
 Chorilaena 11
 Choripetalum 91
 Chorizandra 127
 Chorizema 32,33
 Chromochiton 82
 Chrysobactron 118
 Chrysocephalum 81
 Chrysocoryne 82
 Chrysodium 139
 Chrysodiscus 80
 Chrysogonum 83
 Chrysophyllum 91
 Chrysopogon 132
 Chrysorrhoea 51
 Chionocephalus 83
 Chuncoa 50
 Cibotium 137
 Cinna 133
 Cinnamomum 4
 Cissodendron 61
 Citriobatus 7,141
 Citrus 12
 Cladium 128,143
 Cladodes 21
 Claoxylon 20
 Clausena 12
 Claytonia 27
 Clematis 1
 Cleistanthus 20
 Cleome 4,141
 Clerodendrum 103,143
 Cleisostoma 111
 Clianthus 39
 Clidanthra 40
 Clinostigma 120
 Clitoria 40
 Cloezia 59
 Coatesia 12
 Cocculus 4
 Cochlospermum 6
 Cocos 120
 Codiaeum 21
 Codonocarpus 32
 Coelachne 133
 Coelandria 110
 Coelebogyne 21
 Coelorrhachis 131
 Coelospermum 75
 Coenopteris
 Coffea 75
 Coilocarpus 30
 Coldenia 100,143
 Coleanthera 107
 Coleocoma 79
 Coleostylis 86
 Coleus 100
 Colobandra 101
 Colobanthus 27
 Colletia 61
 Colmeiroa 48
 Colocasia 120
 Colubrina 60
 Columbea 109
 Combretaceae 50
 Comesperma 8,141
 Commelina 122
 Commelineae 122
 Commerconia 16,141
 Compositae 77
 Conanthodium 81
 Conchium 70
 Coniferae 109
 Connaraceae 32
 Conospermum 66
 Conostephiopsis 107
 Conostephium 107
 Conostylis 116
 Conothamnus 56
 Convolvulaceae 94
 Convolvulus 95,143
 Conyza 79
 Coprosma 75
 Corchorus 17
 Cordia 99
 Cordyline 117
 Corethrostylis 16
 Coridochloa 130
 Cornaceae 74
 Coronopus 141
 Correa 11
 Corybas 112
 Corymbis 111
 Corymborchis 111
 Corypha 120
 Corynotheca 118
 Corysanthes 112
 Cosmelia 108
 Costus 114
 Cotula 84
 Covellia 21
 Craspedia 83
 Crassulaceae 48
 Crantzia 63
 Crantziola 63
 Crepis 85
 Cressa 95
 Crinum 116
 Crossolepis 83
 Crossotoma 87
 Crotalaria 38,140,142
 Croton 20
 Crowea 10
 Cruciferae 5
 Cryphia 101
 Cryptandra 60,140
 Cryptocarya 3
 Cryptosema 32
 Cryptostemon 51
 Cryptostylis 112
 Cucumis 76,142
 Cucurbitaceae 76
 Cudrania 22
 Cupania 24
 Cupuliferae 22
 Curculigo 116
 Curcuma 114
 Cuscuta 95,143
 Cutisia 48
 Cyanostegia 103
 Cyanothamnus 10
 Cyanotis 121
 Cyathea 137
 Cyathochaete 127,143
 Cyathodes 105
 Cyathopappus 83
 Cyathopsis 105
 Cybele 71
 Cycadeae 109
 Cycas 109
 Cyclogyne 39
 Cyclothea 32
 Cycnogeton 121
 Cylicodaphne 4
 Cyllindrosorus 82
 Cymbidium 111
 Cymbonotus 85
 Cyminosma 12
 Cymodocea 121
 Cynanchum 93,143
 Cynoctomum 93
 Cynodon 134
 Cynoglossum 100,143
 Cynometra 43
 Cyperaceae 124
 Cyperus 124,143
 Cyphanthera 96
 Cyphokentia 120
 Cypselocarpus 32
 Cyrtostylis 113
 Cystanthe 108
 Cystopteris 138
 Dacrydium 109
 Dactyloctenium 134
 Daemia 94
 Dalbergia 42
 Dallachya 60
 Damasonium 121
 Dammara 109
 Dampiera 87,142
 Danthonia 134
 Daphnandra 3
 Daphne 74
 Daphnobryon 74
 Darea 138
 Darlingia 71
 Dasymalla 103
 Dasypogon 119
 Datura 96,143
 Daucus 63
 Davallia 137
 Davidsonia 48
 Daviesia 35,140
 Darwinia 51
 Decalophium 51
 Decaspermum 59
 Decaspora 107
 Decazesia 83
 Deeringia 29
 Delabechea 15
 Dendrobium 110
 Denhamia 26
 Denisonia 103
 Dennstaedtia 137
 Dentella 74
 Deparia 137
 Deplanchea 99
 Depremesnilia 101
 Derris 42
 Deschampsia 133

- Desmocladus 124
 Desmodium 40
 Desvauxia 123
 Deyeuxia 133
 Diaspasis 87
 Dianella 117
 Dicarpidium 15
 Dicera 17
 Dicerma 40
 Dichelachne 133
 Dichondra 95
 Dichopetalum 63
 Dichopogon 118
 Dichosema 33
 Dichrostachys 43
 Dicksonia 137
 Dieladanthra 140
 Dielidopteris 138
 Dieliptera 99
 Dicrastylis 103
 Dictyopteris 139
 Didiscus 62,142
 Didymanthus 30
 Didymeria 11
 Didymochaete 133
 Didymonema 128
 Didymotheca 32,142
 Digitaria 130
 Dilivaria 99
 Dillenia 2
 Dilleniaceae 1
 Dillwynia 37
 Dimeria 130
 Dimetopia 62
 Dimorpholepis 80
 Diodontium 84
 Dioscorea 115
 Dioscorideae 115
 Diospyros 92
 Diotosperma 84
 Diplachne 134
 Diplacrum 129
 Diplanthera 99,121
 Diplarrhena 114
 Diplaspis 63
 Diplax 132
 Diplazium 138
 Diploglottis 24
 Diplolaena 11
 Diplobolium 39
 Diplopeltis 25
 Diplopogon 133
 Diplora 138
 Diplospora 74
 Dipodium 111
 Dipogonia 133
 Dipteracanthus 99
 Disarrhenum 132
 Discaria 61
 Dischidia 94
 Discopodium 127
 Discospermum 74
 Diselma 109
 Disemma 76
 Disoon 104
 Dissiliaria 19,141
 Dissocarpus 30
 Distichlis 135
 Distichostemon 25
 Distylis 88
 Dithyrocarpus 122
 Dithyrostegia 82
 Diuris 112
 Dodonaea 25,141
 Dolichandra 99
 Dolichandrone 99
 Dolichos 41
 Donatia 87
 Donia 39
 Doodia 138
 Dopatrium 97
- Doryanthes 116
 Doryphora 3,141
 Dracaena 117
 Dracophyllum 109
 Drakaea 113
 Drapetes 74
 Drosera 7
 Droseraceae 7
 Drummondia 11
 Dryandra 72
 Drymaria 27
 Drymophila 117
 Drymis 2
 Drynaria 139
 Duboisia 96
 Dumbaria 41
 Duperreya 95
 Duttonia 80,104
 Dysophylla 100
 Dysoxylum 9
 Dysphania 29,141
- Eadesia 96
 Earlia 99
 Ebelingia 13
 Ebenaceae 92
 Ebermayera 99
 Ecdeiocolea 123
 Echinocarpus 17
 Echinochloa 130
 Echinocroton 21
 Echinodorus 121
 Echinopogon 133
 Echinopsilon 30
 Echinosperrum 100
 Echinus 21
 Eclipta 83,142
 Ectrosia 135
 Ehretia 99
 Ehrharta 132
 Elachanthus 84
 Elachocroton 21
 Elachopappus 82
 Elachothamnus 77
 Elaeagneae 64
 Elaeagnus 64
 Elaeocarpus 17
 Elaeodendron 26
 Elaphoglossum 139
 Elatine 8
 Elatineae 8
 Elatostemma 22
 Elattostachys 24
 Electrosperra 123
 Eleoichiton 125
 Elephantopus 77
 Eletaria 114
 Eleusine 134,143
 Eleutranthes 76
 Elionurus 131
 Elynanthus 127
 Elythrophorus 135
 Emblingia 5
 Embotrium 71
 Emex 31
 Emilia 84
 Emmenospermum 60
 Emphysopus 77
 Empleurosmia 25
 Encephalartos 110
 Enchylaena 30,142
 Enchysia 85
 Endiandra 4,141
 Enhalus 115
 Enhydra 83
 Entada 43
 Enterolobium 47
 Epacrideae 105
 Epacris 108
 Epaltes 79,140,142
 Epiandra 128
- Epiblema 112
 Epicharis 9
 Epilobium 49
 Epipogon 111
 Epipremnum 120
 Epithinia 75
 Epitriche 82
 Eragrostis 135,143
 Eranthemum 99
 Erechites 85,143
 Eremaea 57
 Eremodendron 104
 Eremophila 104,140
 Eremopyxis 52
 Eremosyne 48
 Eria 140
 Eriachne 133,143
 Erianthus 131
 Ericaceae 105
 Ericomyrtus 53
 Erigeron 79
 Eriocalia 63
 Eriocaulae 123
 Eriocaulon 123
 Eriochilus 113
 Eriochiton 30
 Eriochlamys 82
 Eriochloa 130
 Eriocladium 82
 Eriosema 41
 Eriostemon 10
 Eritrichum 100
 Erodium 13
 Erodiphyllum 77
 Erycibe 94
 Eryngium 63
 Erysimum 5
 Erythraea 90
 Erythrina 41,142
 Erythrorchis 111
 Erythrophlaeum 43
 Erythroxyllum 13
 Etaeria 111
 Ethulopsis 79
 Euandra 129
 Eucalyptus 57,141,142
 Eucarya 64
 Euchilopsis 37
 Euchilus 36
 Euchiton 79
 Eucryphia 48
 Eudesmia 57
 Eugenia 59
 Euneapogon 133
 Euodia 12
 Euonymus 26
 Euosma 91
 Eulalia 131
 Eulophia 111
 Eupatorium 77,141
 Euphorbia 18,140,141
 Euphorbiaceae 18
 Euphoria 24
 Eupomatia 3
 Euphrasia 98
 Eurosichinus 25
 Eurostorrhiza 129
 Eurybia 78
 Eurybiopsis 79
 Eurycles 116
 Euryomyrtus 53
 Eustrephus 117
 Eutacta 109
 Eutassa 109
 Eutaxia 32
 Euthales 89
 Euxolus 29
 Exarrhena 100
 Excaecaria 21
 Exocarpos 64
 Exocarya 127
- Evolvulus 95
 Eyrea 79
- Fabricia 54
 Fagus 22
 Fagraea 90
 Faradaya 103
 Fatona 22
 Faustula 81
 Fawcettia 4
 Fenzlia 59
 Festuca 134
 Fieldia 98
 Ficoideae 31
 Ficus 21,140,141
 Filices 136
 Fimbristylis 125
 Fischera 62
 Fisquetia 120
 Fitzalanina 3
 Fitzroya 109
 Flacourtiaceae 6
 Flagellaria 117
 Flaveria 84,142
 Flemingia 41
 Fleurya 22
 Flindersia 9
 Floriscopa 122
 Floscopa 122
 Flueggea 20
 Fluviales 121
 Fontainea 21,141
 Forstera 86
 Forsteropsis 85
 Fragosa 63
 Francisia 51
 Frankenia 26
 Frankeniaceae 26
 Franklandia 67
 Freirea 22
 Freyma 59
 Frenela 109
 Freycinetia 120
 Friesia 17
 Froebelia 105
 Fugosia 15
 Fuirena 126
 Fusanus 64
- Gahnia 128
 Galactia 41
 Galeola 111
 Galium 76,142
 Gamelythrum 133
 Gamozygis 82
 Ganophyllum 24
 Gardenia 74
 Garuga 25
 Gastrodia 111
 Gastrolobium 33,141
 Gaultheria 105
 Gaultiera 105
 Geijera 12
 Geitonoplesium 117
 Geleznovia 10
 Genetyllis 51
 Geniostoma 90
 Genoplesium 112
 Genosiris 114
 Gentiana 90
 Gentianeae 90
 Geococcus 5
 Geodorum 111
 Geophila 75
 Geraniaceae 13
 Geranium 13
 Geum 47
 Gilberta 82
 Gilesia 15
 Gillbeea 48
 Gingidium 63

- Glabaria 4
 Gleichenia 137
 Glinus 31
 Glischocaryon 49
 Glochidion 19
 Glossodia 114
 Glossogyne 84
 Glossostigma 97
 Glyceria 134
 Glycine 41
 Glycosmis 12
 Glycyrrhiza 40
 Gmelina 103
 Gnaphalium 79
 Gnaphalodes 83
 Gnephosis 83
 Gomphandra 64
 Gompholobium 34,142
 Gomphrena 28
 Gongronema 94
 Goniocarpus 49
 Goniophlebium 139
 Goniopogon 78
 Goniopteris 139
 Goniotriche 28
 Goodenia 88,143
 Goodeniaceae 87
 Goodenoughia 88
 Goodia 38
 Goodyera 111
 Gossypium 15,141
 Gouania 61
 Gramineae 131
 Grammatotheca 85
 Grammitis 139
 Graptophyllum 99
 Gratiola 97
 Graumuellera 121
 Greevesia 14
 Grevillea 68,140,142
 Grewia 17
 Grisebachia 120
 Grumilia 75
 Guettarda 75
 Guettardella 75
 Guichenotia 16
 Guilandina 42
 Gunnera 50
 Gunnia 31,111
 Guioia 24
 Guttiferae 8
 Gymnagathis 55
 Gymnanthera 93,143
 Gymnema 94
 Gymnochaete 127
 Gymnoschoenus 127
 Gymnococca 73
 Gymnogramme 139
 Gymnogyne 84
 Gymnopteris 139
 Gymnopogon 93
 Gymnosporia 26
 Gymnostachys 120
 Gymnostylis 84
 Gymnothrix 131
 Gynaphanes 142
 Gynura 84
 Gypsophila 27
 Gyrocarpus 51
 Gyrostachys 111
 Gyrostemon 32
 Gyrostephium 83

 Habenaria 114
 Haekeria 82
 Haemodorum 115
 Hakea 70,142
 Halgania 100
 Halfordia 12
 Halocnemum 30
 Halodule 121

 Halophila 115
 Halorageae 49
 Haloragis 49
 Halothamnus 13
 Hamamelideae 48
 Hamelinia 117
 Hannafordia 16
 Haplopappus 79
 Hardenbergia 41
 Harmogia 53
 Harpulia 24
 Harrisonia 13
 Hartigshea 9
 Haussmannia 99
 Hearnia 9
 Hedera 51
 Hedera 61
 Hedraianthera 9,141
 Hedyocarya 3
 Hedyotis 74
 Hedyoscepe 120
 Heleiocharis 125
 Helichrysum 81,142
 Helicia 68
 Helicteres 15
 Heliophytum 100
 Heliotropium 100
 Helipterum 80,140,142
 Hellenia 114
 Helmholtzia 122
 Helmintostachys 136
 Helophyllum 87
 Helopus 130
 Helostigma 85
 Helothrix 127
 Hemarthria 131
 Hemarrhena 97
 Hemianthera 101
 Hemiphorus 63
 Hemisteirus 28
 Hemistemma 1
 Hemistephus 1
 Hemitelia 137
 Heptopleurum 61
 Heritiera 15
 Hermannia 15
 Hernandia 4
 Herpestis 97
 Herpolirion 118
 Hetaeria 122
 Heterachne 135
 Heterodendron 24
 Heterolaena 73
 Heteropogon 132
 Hewardia 117
 Hexatheca 121
 Hibbertia 1,141
 Hibiscus 14,141
 Hierochloa 132
 Hippocratea 26
 Hippocrepandra 18
 Hodgkinsonia 75
 Hodgsonia 118
 Hodgsoniella 118
 Holcus 132
 Hologamium 131
 Holotome 63
 Homalium 6
 Homalocalyx 52
 Homalospermum 54
 Homalostoma 108
 Hormogyne 92
 Hovea 38
 Howittia 14
 Hoya 94

 Huanaca 63
 Huegelia 62
 Huenefeldia 78
 Hugonia 13
 Humata 137
 Humea 82,142
 Hutchinsia 6
 Huttia 1
 Hyalochlamys 82
 Hyalolepis 82
 Hyalosperma 80
 Hybanthus 6,141
 Hydriastele 120
 Hydrilla 115
 Hydrocharideae 115
 Hydrocharis 115
 Hydrocotyle 62
 Hydroglossum 136
 Hydrolea 99
 Hydropeltis 1
 Hydrophila 99
 Hydnophytum 75
 Hydrophyllaeae 99
 Hylococcus 19
 Hymenanthra 6,141
 Hymenolepis 139
 Hymenophyllum 137
 Hymenopodium 7,141
 Hymenotheca 32
 Hypaelytrum 126
 Hypelytrum 127
 Hypericinae 8
 Hypericum 8
 Hypocalymma 54
 Hypoestes 99
 Hypolepis 139
 Hypoporum 129
 Hypolaena 124
 Hypoxis 116
 Hypserpa 4
 Hyptiandra 12
 Hyrtandra 22

 Iberis 6
 Ichnocarpus 93
 Ilex 63
 Ilysanthes 97
 Imbricaria 53
 Imperata 131
 Indigofera 39,140,142
 Ionidium 6
 Iphigenia 117
 Ipomoea 94,143
 Irideae 114
 Iris 114
 Irwingia 61
 Isachne 130
 Isandra 140
 Ischaemum 131
 Iseilema 132
 Isoetes 136
 Isoetopsis 84
 Isolepis 125
 Isoloma 137
 Isopogon 65,142
 Isoschoenus 127
 Isotoma 85,142
 Isotropis 33,140
 Ixauchenius 77
 Ixioclamys 79
 Ixiolaena 80
 Ixiosporum 7
 Ixodia 82
 Ixora 74,142

 Jacksonia 34
 Jambosa 59
 Jansonia 32
 Jasmineae 92
 Jasminum 92
 Johnsongia 119

 Josephia 72
 Josephinia 105
 Junceae 122
 Juncella 123
 Juncus 122
 Jungia 53
 Jussieuia 49
 Justicia 99

 Kaleniczenkia 32
 Kamphusia 87
 Kamptzia 59
 Kellaria 74
 Kennedyia 41
 Kentia 120
 Kentiopsis 120
 Kentropis 30
 Keraudrenia 16
 Kerneria 121
 Kibara 3
 Kingia 119
 Kippistia 77
 Kissodendron 61
 Klandella 101
 Knoxia 76
 Kochia 30,140,142
 Koehnea 142
 Kreyssigia 117
 Kunzea 54,142
 Kyllingia 124

 Labiatae 100
 Labichea 42,142
 Laboucheria 43
 Laccospadix 120
 Lachnagrostis 133
 Lachnocephalus 103
 Lachnostachys 102
 Lachnothalamus 85
 Lactaria 93
 Lagenaria 76
 Lagenophora 77,142
 Lagerstroemia 49,142
 Lagrezia 29
 Lagunaria 15
 Lalage 37
 Lamarchea 57
 Lambertia 68
 Lampocarya 128
 Lamprochlaena 82
 Lamprolobium 39
 Laportea 22
 Lappago 131
 Lappula 100
 Lasianthus 75
 Lasiopetalum 16
 Lastraea 135
 Latourea 87
 Latrobea 37
 Lauraceae 3
 Laurencia 85
 Lavatera 13
 Lawrencia 13
 Lawrenceella 81
 Lawsonia 49
 Laxmannia 118
 Learosa 3
 Lebediera 20
 Ledgeria 111
 Leea 24,141
 Leersia 132
 Leewenhoekia 86
 Legnophora 4
 Leguminosae 32
 Leichhardtia 4,94,109
 Lemna 121
 Lemnaceae 121
 Lenticularinae 98
 Leontopodium 80
 Lepia 6
 Lepidium 6,141

- Lepidobolus 124
 Lepidocoma 82
 Lepidosperma 128
 Lepidospora 128,143
 Lepidozamia 110
 Lepigonum 27
 Lepironia 127
 Lepistemon 94
 Leptaspis 131
 Leptatherum 131
 Leptinella 84
 Leptocarpus 124
 Leptoceras 113
 Leptochloe 134
 Leptocymnus 41
 Leptocytisus 37
 Leptolobium 41
 Leptomeria 64
 Leptopteris 137
 Leptorrhynchus 80,142
 Leptosema 32
 Leptospermum 54
 Leptotriche 83
 Lepturus 131
 Lepyrodia 123
 Leschenaultia 87,142
 Lespedeza 40,142
 Lestibudesia 29
 Leucas 101
 Leucocarpus 26
 Leucolaena 62
 Leucophyta 83
 Leucopogon 105
 Leucothamnus 16
 Leuzea 85
 Lhotzkia 52
 Libertia 114
 Licuala 120
 Ligusticum 63
 Ligustrum 92,143
 Liliaceae 117
 Limnanthemum 90
 Limnophila 97
 Limnostachys 121
 Limosella 97
 Lindernia 97,143
 Lindsaea 137
 Lindsaya 137
 Lineae 13
 Linkia 67
 Linociera 92
 Linospadix 120
 Linschotenia 87
 Linum 13
 Liparis 110
 Liparophyllum 90
 Lipocarpa 126
 Lippaya 74
 Lippia 102
 Lissanthus 105
 Litobrochia 138
 Litsea 4
 Livistona 120
 Lobelia 85,142
 Lobophyllum 100
 Lopopogon 107
 Logania 91,143
 Loganiaceae 90
 Lomandra 119
 Lomaria 138
 Lomariopsis 139
 Lomatia 71
 Lonchocarpus 42
 Lopadocalyx 63
 Lophoclinium 80
 Lophostemon 58
 Lopidium 136
 Lorantheaceae 64
 Loranthus 65
 Lotus 38
 Loudonia 49
- Lourea 40
 Loxocarya 124
 Lucuma 91
 Ludwigia 49
 Luffa 76
 Luisia 140
 Lumnitzera 50
 Luzula 122
 Luzuriago 117
 Lycium 96
 Lycopodiaceae 136
 Lycopodium 136
 Lycopus 101
 Lyginia 123
 Lygodictyon 136
 Lygodium 136
 Lyonsia 93
 Lyperanthus 113
 Lysanthe 68
 Lysicarpus 59
 Lysimachia 91
 Lysinema 108
 Lysiosepalum 17
 Lythrum 49
- Maba 92
 Macadamia 68
 Macaranga 21
 Macarthuria 31,142
 Macdonaldia 111
 Macgregoria 26
 Mackinlaya 61
 Macropidia 116
 Macropiper 23
 Macropodia 116
 Macropteranthus 51
 Macrostegia 73
 Macrostigma 47
 Macrozamia 110
 Maesa 91
 Magnoliaceae 2
 Mahernia 15
 Maireana 30
 Malacochaete 126
 Malaisia 22
 Malcolmia 5
 Mallophora 103
 Mallotus 21
 Malpighiaceae 25
 Malva 13
 Malvaceae 13
 Malvastrum 13
 Manglesia 68
 Manisuris 131
 Mapania 127
 Mappa 21
 Marattia 136
 Marianthus 7,141
 Mariscus 124
 Marlea 74
 Marsdenia 94
 Marsilea 136
 Marquisia 75
 Mayepea 92
 Mazus 97
 Megalotheca 124
 Meionectes 50
 Medicosma 12
 Melachne 128
 Melaleuca 55,140,142
 Melanthesia 20
 Melanthesiopsis 20
 Melanthium 117
 Melastoma 60
 Melastomaceae 60
 Melhania 15
 Melia 9
 Meliaceae 9
 Melichrus 105
 Melicope 12
 Melochia 15
- Melodinus 92
 Melodorum 3
 Melothria 77
 Memecylon 60
 Memorialis 22
 Menkea 5
 Meniocus 5
 Meniscium 139
 Menispermeae 4
 Mentha 101
 Merkusia 87
 Mertya 61
 Mesembrianthemum 31
 Mesomelaena 127
 Mesotriche 105
 Methorium 15
 Metrosideros 59
 Mezoneurum 42
 Michiea 107
 Micraira 133
 Micranthemum 18,141
 Microcachrys 109
 Microcarpaea 97
 Microchloa 134
 Microcladium 63
 Microclisia 4
 Microcorys 102
 Microcybe 10
 Microgyne 79
 Microlaena 132
 Microlepidium 6
 Microlepis 137
 Micromelum 12
 Micromyrtus 52
 Micropyxis 91
 Microseris 85
 Microstachys 21
 Microstemma 94
 Microtis 112
 Microstylis 111
 Milisia 3
 Milletia 39
 Milligania 50,117
 Millotia 82
 Milnea 9
 Mimosa 43
 Mimulus 97
 Mimusops 92
 Minuria 77
 Mirbelia 33,142
 Mischocarpus 24
 Mitragyne 90
 Mitrasacme 90,142
 Mitreola 90
 Mnium 27
 Modecca 76
 Molkenboeria 87
 Mollinedia 3
 Molloya 68
 Mollugo 31,142
 Momordica 76
 Monacather 134
 Monencyanthes 80
 Monenteles 79
 Monimieae 3
 Monogramma 138
 Monochoria 121
 Monococcus 32
 Monoploca 6
 Monotaxis 18,141
 Monotoca 105
 Montia 27
 Moonia 83
 Moraea 114
 Morelotia 128
 Morgania 97
 Morinda 75
 Morisia 127
 Morna 80
 Moschosma 100
 Motherwellia 61
- Mucuna 41
 Muehlenbeckia 31
 Muehlenbergia 133
 Mukia 77
 Murraya 12
 Musa 114
 Myoporinae 104
 Myoporum 104,140
 Myosotis 100
 Myosurus 1,141
 Myriocephalus 82
 Myriogyne 84
 Myriophyllum 50
 Myristica 3
 Myristiceae 3
 Myrmecodia 75
 Myrsinaceae 91
 Myrsine 91
 Myrtaceae 51
 Myrtus 59
- Nablonium 83
 Nageia 109
 Najas 121
 Nania 59
 Nasturtium 5
 Needhamia 107
 Negria 98
 Nelitris 59
 Nelsonia 99
 Nelumium 1
 Nelumbo 1
 Nematoceras 112
 Nematolepis 11
 Nematophyllum 38
 Nematopus 83
 Nematostigma 114
 Nemedra 9
 Neuroopera 20
 Neotopteris 138
 Nepenthaceae 23
 Nepenthes 23,140
 Nephelium 24
 Nephrodium 138
 Nephrolepis 138
 Neptunia 43
 Nertera 75
 Nesaea 49,142
 Nesodaphne 3
 Nesogenes 102
 Nettoa 17
 Neurachne 131
 Newcastleia 102
 Nicotiana 96
 Niemeyera 91,114
 Nipa 120
 Niphobolus 139
 Nitraria 13
 Notelaea 92,143
 Notholaena 138
 Nothopanax 61
 Notonierium 93
 Notothixos 65
 Nuytsia 65
 Nyctagineae 32
 Nymphaea 1
 Nymphaeaceae 1
 Nyssanthus 28
- Oberonia 110
 Obione 29
 Ochraceae 10
 Ochrolasia 9
 Ochrosia 93
 Ocimum 100
 Octoclinis 109
 Oenanthe 63,142
 Oenothera 49
 Ogcerostylus 82
 Olacinae 63
 Olax 63

- Oldenlandia 74,142
 Olea 92
 Oleandra 138
 Olearia 78
 Oliganthemum 79
 Oligarrhena 107
 Omalanthus 21,141
 Omphacomeria 64
 Onagreae 49
 Oncosporum 7
 Oncostylis 125
 Onychosepalum 124
 Opercularia 75,142
 Ophioderma 136
 Ophioglossum 136
 Ophiorrhiza 74
 Ophiuros 131
 Opilia 63
 Oplismenus 130
 Orchideae 110
 Oreobolus 127
 Oreomyrrhis 63
 Oreostylidium 87
 Orites 68
 Oritina 68
 Ormocarpum 40
 Ornithochilus 111
 Orobanche 98
 Orobancheae 98
 Orthoceras 112
 Orthopogon 130
 Orthostemon 90
 Orthosiphon 100
 Orthotropis 33
 Orthrosanthus 114
 Oryza 132
 Osbeckia 60
 Osbornia 59
 Oschatzia 63
 Osmunda 137
 Osteocarpum 30
 Otanthera 60
 Ottelia 114
 Ourisia 97
 Owenia 9
 Oxalis 13
 Oxleya 9
 Oxycladium 33
 Oxylobium 32
 Oxymyrrhine 53
 Oxytelma 93
 Ozothamnus 81

 Pachycornia 30
 Pachygone 4
 Pachynema 2,140
 Pachysurus 83
 Pacquerina 77
 Pagetia 12
 Palmae 119
 Palmeria 3
 Panax 61
 Pancovia 24
 Panaetia 80
 Pandaneae 120
 Pandanophyllum 127
 Pandanus 120,140
 Panicum 130
 Panopsis 68
 Papaver 4,141
 Papaveraceae 4
 Pappophorum 133
 Paratropia 61
 Parkeria 137
 Parietaria 22
 Parinarium 47
 Paritium 14
 Parsonsia 93
 Paryphantha 52
 Paspalum 130
 Passiflora 76

 Passifloreae 76
 Patania 137
 Patersonia 114
 Pavetta 74
 Pavonia 14
 Pedalinae 105
 Pedilonum 110
 Pelargonium 13
 Pellaea 138
 Pelonastes 50
 Peltophorum 42
 Pemphis 49
 Pennantia 64
 Pennisetum 131
 Pentaceras 12
 Pentachondra 107
 Pentacraspedion 133
 Pentadactylon 67
 Pentadynamis 38
 Pentagonaster 54
 Pentalepis 83
 Pentapeltis 62
 Pentataphrus 105
 Pentapogon 133
 Pentaptelion 105
 Pentatropis 94
 Peperomia 23,140,141
 Peplidium 97,143
 Pericalymma 54
 Pericampylus 4
 Pernettya 105
 Peroa 105
 Perojoa 105
 Perotis 131
 Persoonia 67,142
 Petalolepis 81
 Petalostigma 19
 Petalostylis 42
 Petermannia 115
 Petrophila 65
 Phacelothrix 82
 Phaenopoda 80
 Phajus 111
 Phalangium 118
 Phaleria 74
 Phanerandra 105
 Pharbitis 94
 Phaseolus 41
 Phebalium 10
 Phegopteris 139
 Pherosphaera 109
 Philotheca 11,140
 Philoxerus 28
 Philydreae 122
 Philydrum 122
 Phlebocalymma 63
 Phlebocarya 115
 Phlebodium 138
 Phoberos 6
 Pholidia 104
 Pholidiopsis 104
 Pholidota 110
 Phormium 117
 Phragmites 135
 Phreatia 110
 Phyllachne 87
 Phyllanthus 19,140,141
 Phyllocalymma 82
 Phyllocladus 109
 Phylloodium 40
 Phylloglossum 136
 Phyllopappus 85
 Phyllota 36
 Phymatocarpus 56
 Phymatodes 139
 Physalis 95
 Physolobium 41
 Physopsis 103
 Phytolacceae 32
 Picrophyta 88
 Pigea 6

 Pileanthus 52
 Pilitis 108
 Pilularia 136
 Pimelea 73,142
 Piper 23
 Piperaceae 23
 Piptandra 53
 Piptomeris 34
 Piptostemma 82
 Pisonia 32
 Pithecolobium 47
 Pithocarpa 82
 Pithyrodia 103
 Piptocalyx 3
 Pipturus 22
 Pittosporae 6
 Pittosporum 6,141
 Plagianthus 13
 Plagiogyria 138
 Plagiolobium 38
 Plagiosetum 131
 Plantagineae 91
 Plantago 91,143
 Platycarpidium 62
 Platycerium 139
 Platychilum 38
 Platylobium 37
 Platylooma 138
 Platyptelea 48
 Platysace 62
 Platytheca 9
 Platyzoa 137
 Plectranthus 100
 Plea 117
 Pleiococca 12
 Pleiogyne 84
 Pleogyne 4
 Pleopeltis 139
 Pleurandra 1
 Pleurandropsis 10
 Pleurocarpaea 77
 Pleurogramme 138
 Pleuropappus 82
 Plexaure 110
 Plinia 59
 Plintacanthus 134
 Plokiostigma 26
 Pluchea 79
 Plumbagineae 26
 Plumbago 26,141
 Poa 134
 Podocarpus 109
 Podocoma 79,142
 Podolepis 80,142
 Podolobium 32
 Podopetalum 42
 Podosperma 80,140,142
 Podostemoneae 23
 Podotheca 80,140,142
 Poeciloderms 15
 Pogonia 104,111
 Pogonolepis 82
 Pogonolobus 75
 Pogonatherum 131
 Pogonetes 87
 Pogostemon 100
 Poirertia 38,108
 Polanisia 4
 Pollia 122
 Pollichia 100,143
 Pollinia 131
 Polyalthia 3,140
 Polycalymma 82
 Polycarpaea 27
 Polycarpon 27
 Polyenemon 29
 Polygala 8,141
 Polygaleae 8
 Polygonaceae 31
 Polygonum 31,142
 Polymeria 95,143

 Polyosma 48
 Polyphragmon 75
 Polypodium 139
 Polypogon 131
 Polypompholyx 98
 Polystichum 138
 Polyzone 51
 Pomaderris 60,142
 Pomatotheca 31
 Pomax 76
 Ponceletia 108
 Pongamia 42
 Pontederiaceae 121
 Popowia 3
 Porana 95
 Poranthera 18,141
 Porospermum 61
 Portulaca 27,141
 Portulacaceae 27
 Posidonia 121
 Potamogeton 121
 Potamophila 132
 Potentilla 47
 Pothos 120
 Pouzolzia 22,141
 Pozoa 63
 Pozoopsis 63
 Prasophyllum 112,140,143
 Pratia 85
 Premna 103
 Primulaceae 91
 Prionosepalum 124
 Prionotes 108
 Pritzelia 62,122
 Procris 22
 Pronaya 7
 Prostanthera 101,140
 Proteaceae 65
 Prunella 101
 Pseudalangium 74
 Pseudanthus 18,141
 Pseudatalaya 24
 Pseudomorus 22
 Pseudopholidia 104
 Psilotrichum 28
 Psilotum 136
 Psoralea 38,142
 Psychotria 75
 Psydrax 75
 Pterigeron 79,142
 Pteris 138
 Pterocaulon 79
 Pterochaete 80
 Pterolobium 42
 Pteroloma 40
 Pterostigma 97
 Pterostylis 113
 Pterygopappus 80
 Ptilotus 28,140,141
 Ptychosema 39
 Ptychosperma 120
 Pultenaea 36
 Pumilio 82
 Punicella 54
 Pycnolachne 102
 Pycnosorus 83
 Pycnospora 40
 Pycneus 124
 Pygmaea 97
 Pyxidaria 143

 Quinetia 82
 Quintinia 48
 Quoya 103

 Ramphidia 111
 Randia 74
 Ranunculaceae 1
 Ranunculus 1,141
 Raoulia 80
 Rapuntium 85

- Ratonia 24
 Reedia 129
 Regelia 56
 Remirea 127
 Renealmia 114
 Restiaceae 123
 Restio 124
 Rhabdothermus 98
 Rhagodia 29,141
 Rhamnaceae 60
 Rhamnus 60
 Rhamphicarpa 98
 Rhamphidia 111
 Rhabdiphora 120
 Rhipogonum 117
 Rhizophora 50
 Rhizophoreae 50
 Rhizospermeae 136
 Rhodamnia 59
 Rhodanthe 80
 Rhodomyrtus 59
 Rhodosphaera 25
 Rhopala 68
 Rhus 25
 Rhyncharrhena 94
 Rhynchosia 41,142
 Rhynchospora 127
 Rhychothemon 16
 Rhytidandra 74
 Rhytidanthe 80
 Rhytidosporum 7
 Richea 83,108
 Ricinocarpus 19,141
 Riedleya 15
 Rinzia 53
 Rochelia 100
 Roa 35
 Roeperia 5,19
 Rosaceae 47
 Rostellaria 99
 Rostellularia 99
 Rotala 49
 Rottboellia 131
 Rottlera 21
 Roupala 68
 Rourea 32
 Roxburghia 117
 Roxburghiaceae 117
 Rubiaceae 74
 Rubus 47
 Ruellingia 16
 Ruellia 99
 Rumex 31
 Ruppia 121
 Rutaceae 10
 Rutidochlamys 80
 Rutidosis 82
 Ryssopteris 25

 Saccharum 131
 Saccolabium 111
 Saccopetalum 3
 Sagenia 138
 Sagina 27,141
 Salacia 26
 Salicariaceae 49
 Salicornia 38,142
 Salisia 54
 Salomonina 8
 Salsola 31,142
 Salsolaceae 29
 Salvia 101
 Samara 91
 Sambucus 76,142
 Samolus 91
 Samydaceae 6
 Sandfordia 10
 Santalaceae 64
 Santalum 64
 Sapindaceae 21
 Sapindus 24

 Saponaria 27,141
 Sapota 91
 Sapotaceae 91
 Sarcocephalus 74
 Sarcochilus 111
 Sarcopetalum 4
 Sarcostemma 94
 Sarcozygium 13
 Sarotes 16
 Saussurea 85
 Saxifrageae 47
 Scaevola 87,142
 Scalia 80
 Scaliopsis 80
 Schelhammra 117
 Schenodorus 134
 Schidiomyrtus 53
 Schizaea 136
 Schizachyrium 132
 Schizeilema 63
 Schizolema 137
 Schizomeria 48
 Schizopleura 56
 Schmidelia 24
 Schoberia 30
 Schoenia 81
 Schoenodum 123
 Schoenolaena 62
 Schoenus 127,143
 Scholtzia 53
 Schuermannia 51
 Schwaegrichenia 116
 Scirpidium 125
 Scirpodendron 127
 Scirpus 126,143
 Scitamineae 114
 Scleranthus 27
 Scleria 129
 Sclerochlamys 30
 Sclerolaena 30
 Scleroleima 84
 Sclerothermus 37
 Scolopendrium 138
 Scolopia 6
 Scoparia 97
 Scorzonera 85
 Scottia 37
 Scrophularinae 97
 Scutellaria 101
 Scyphiphora 75
 Scyphocoronis 82
 Seaforthia 120
 Sebaea 90
 Sebastiania 21
 Secamone 93
 Securinea 20,141
 Selaginella 136
 Selliera 88
 Selliguea 139
 Semecarpus 26
 Senebiera 6,141
 Senecio 84
 Sentis 104
 Sequoia 109
 Serianthes 47
 Seringea 16
 Sersalisia 91
 Sesbania 39,142
 Seseli 63
 Sesuvium 31
 Setaria 130,143
 Selwynia 4
 Sheffieldia 91
 Sicyos 77
 Sida 14,141
 Sideroxylon 91
 Siebera 62
 Siegesbeckia 83
 Siemssenia 80
 Sieversia 47
 Siloxerus 82

 Silphiospermum 77
 Simarubeae 12
 Simsia 66
 Siphonodon 26
 Sisymbrium 5
 Sisyrinchium 114
 Sium 63
 Skinnera 94
 Skirrhophorus 82
 Sloanea 17
 Smilax 117,140
 Smithia 40
 Solanaceae 95
 Solanum 95,143
 Soleniscia 105
 Solenogyne 77
 Solenostigma 21
 Soliva 84
 Sollya 7
 Sonneratia 60
 Sonzaya 25
 Sophora 42
 Sopubia 98
 Sorghum 132
 Sorostachys 124
 Sowerbaea 118
 Spadostyles 36
 Spanoghea 24
 Sparganium 120
 Spartothermus 102,143
 Spathoglottis 111
 Spathodea 99
 Spergularia 27
 Spermacoe 76
 Spermalepis 59
 Spermaxyrum 63
 Sphaeranthos 79
 Sphaerolobium 35
 Sphaeromorphaea 79
 Sphaeropus 129
 Sphenotoma 109
 Sphincterostoma 108
 Spiculaea 113
 Spinifex 131
 Spilanthus 84
 Spiranthus 111
 Spirodela 121
 Spiropodium 79
 Spodiopogon 131
 Spondias 26
 Sponia 21
 Sporobolus 133,143
 Sprengelia 108
 Spyridium 60
 Stachystemon 18
 Stackhousia 26,141
 Stackhousiaceae 26
 Statice 26
 Stawellia 119
 Steetzia 78
 Stegania 138
 Steiroglossa 77
 Stekovia 88
 Stellaria 27
 Stemodia 97
 Stenanthemum 60
 Stenanthera 105
 Stenocarpus 71
 Stenochilus 104
 Stenochlaena 138
 Stenodiscus 60
 Stenopetalum 5
 Stenosperma 84
 Stephania 4
 Sterculia 15
 Sterculiaceae 15
 Stipa 132
 Stirlingia 66
 Stomarrhena 105
 Strangea 68
 Stravadium 59

 Streblorrhiza 39
 Streptachne 132
 Streptothamnus 6
 Streptocarpus 98
 Streptoglossa 79
 Strobilanthes 99
 Stomatopteris 137
 Strongyloclon 41
 Strongylospermum 84
 Strychnos 90
 Strzeleckia 9,143
 Stuartina 79
 Sturtia 110
 Sturtia 15
 Stylidium 74, 85
 Stylobasium 47
 Stylocoryne 74
 Stylolepis 80
 Styloncerus 82
 Stylurus 68
 Stypantra 118
 Styphelia 105
 Styraceae 92
 Suaeda 30
 Suriana 13
 Symphyomera 84
 Symphyomyrtus 57
 Symphyonema 67
 Symphyopetalum 11
 Symplocos 92
 Synaphea 66
 Synaptantha 74
 Syncarpia 59
 Synophlebius 137
 Synostemon 19
 Synoum 9
 Syzygium 59
 Swainsona 39,141,142
 Swammerdamia 81

 Tabernaemontana 93
 Tacca 115
 Taccaceae 115
 Taeniophyllum 111
 Talinum 27
 Tamarindus 42
 Tapeinocheilos 114
 Tarrietia 15
 Tasmania 2
 Tecoma 99
 Telanthera 28
 Telmatophace 121
 Telopea 71
 Temminkia 87
 Templetonia 38
 Tenagocharis 121
 Tephrosia 39,142
 Terminalia 50
 Tersonia 32
 Tetracarpaea 48
 Tetracera 2
 Tetracheilus 43
 Tetragonella 31
 Tetragonia 30,142
 Tetralepis 127
 Tetralobus 98
 Tetranthera 4
 Tetrapasma 61
 Tetraphora 53
 Tetraphylax 88
 Tetrarrhena 132
 Tetratelia 5
 Tetrathea 9
 Teucrium 102
 Thalamia 109
 Thamnopteris 138
 Thecanthes 73
 Theleophyton 29
 Thelychiton 110
 Thelymitra 111
 Therogeron 77

- Thesium 64
 Thespesia 15
 Thespidium 79
 Thlaspi 6
 Thomasia 16
 Thouarsea 131
 Thozetia 94
 Threlkeldia 30
 Thrixpermum 111
 Thryptomene 52,142
 Thuarea 131
 Thunbergia 99,140
 Thymelee 73
 Thyrsacanthus 99
 Thysanotus 118
 Tiaridium 100
 Tiliaceae 17
 Tillaea 48
 Tinonius 75
 Tinospora 4
 Titania 110
 Tittmannia 97
 Tmesipteris 136
 Todea 137
 Torenia 97
 Tournefortia 100
 Toxanthus 82
 Trachycaryon 21
 Trachymene 62,142
 Trachypogon 132
 Tragia 21,141
 Tragus 131,143
 Trema 21
 Tremandra 9
 Tremandrea 9
 Trianthema 31,142
 Tribonanthes 116
 Tribulopsis 13
 Tribulus 13,140,141
 Trichanthodium 83
 Trichelostylis 125
 Trichinium 28
 Trichobasis 56
 Trichobolus 32
 Trichocline 85
 Trichodesma 100
 Trichodium 133
 Tricholoma 97
 Trichomanes 137
 Trichosanthes 76,142
 Trichosiphon 15
 Trichostegia 80
 Tricondylus 71
 Tricoryne 118
 Tricostularia 127
 Trigastrotheca 31
 Triglochin 121
 Trigonella 38
 Trineuron 84
 Triodia 135
 Triphelia 51
 Tripladenia 117
 Tripetelus 76
 Tripterococcus 26
 Triptilodiscus 80
 Triphaphis 135,143
 Trisetum 133
 Tristania 58
 Tristaniopsis 58
 Tristellateia 25
 Tristichocalyx 4,141
 Trithuria 123
 Triticum 135
 Triumphetta 17
 Trochocarpa 107
 Trollius 1
 Trymalium 60
 Turraea 9
 Turritis 5
 Tylophora 94,143
 Typha 120
 Typhaceae 120
 Typhonium 120,143
 Ulmus 21
 Umbelliferae 62
 Uncinia 129
 Ungeria 15
 Urachne 132
 Uragoga 75
 Uralepis 134
 Uraria 40,142
 Urena 14,141
 Urocarpus 10
 Urodon 36
 Urostigma 21
 Urtica 22
 Urticaceae 21
 Utricularia 98
 Uvaria 3
 Uvedallia 97
 Vachellia 43
 Vaginularia 138
 Vallisneria 115
 Vandellia 97
 Veitchia 120
 Velleya 89,140
 Ventenata 85
 Ventenatia 105
 Ventilago 60
 Verbena 102
 Verbenaceae 102
 Vernonia 77
 Veronica 97
 Verreauxia 87
 Verticordia 51
 Vigna 41,142
 Vilfa 133
 Villarsia 90
 Villaresia 64
 Viminaria 35
 Vinca 93
 Vincetoxicum 93
 Viniferae 23
 Viola 6
 Violaceae 6
 Viraya 80
 Viscum 64,142
 Vitex 103
 Vitis 23
 Vittaria 137
 Vittadinia 79,142
 Vlamingia 6
 Vulpia 134
 Wahlenbergia 85,143
 Waitzia 80
 Walcottia 102
 Waltheria 15,141
 Wartmannia 21
 Webera 74
 Wedelia 83,142
 Wehlia 53
 Weinmannia 48
 Westonia 38
 Westringia 102
 Wichurea 60
 Wickstroemia 74
 Wilkiea 5
 Wilkiea 3
 Wilsonia 95
 Wistaria 39
 Wittsteinia 105
 Wolffia 121,143
 Wollastonia 83
 Woodwardia 138
 Woollsia 108
 Wormia 1
 Wrightia 93
 Wrixonia 101
 Wurmbea 117
 Xanthochrysum 81
 Xanthophyllum 8
 Xanthorrhoea 119
 Xanthosia 62
 Xanthostemon 59
 Xanthoxylum 12
 Xerochloa 131
 Xerosollya 7
 Xerotes 119
 Ximenia 63
 Xiphopteris 139
 Xyllocarpus 9
 Xylomelum 68
 Xylosma 6
 Xyridanthus 80
 Xyrideae 122
 Xyris 122
 Youngia 85
 Zaleya 31,142
 Zamia 110
 Zannichellia 121
 Zanonina 76
 Zapania 102
 Zehneria 77
 Zichya 41
 Zieria 10
 Zizyphus 60
 Zornia 40
 Zostera 121
 Zosterostylis 112
 Zoysia 131
 Zygia 47
 Zygomenes 122
 Zygophylleae 13
 Zygophyllum 13