QK
603
C62
1909

ALBERT R. MANN LIBRARY
AT
CORNELL UNIVERSITY

All books are subject to recall after two weeks

GAYLORD			PRINTED IN U.S.A.

Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

THE
GENERA OF FUNGI

FREDERIC EDWARD CLEMENTS, PH. D.

*Professor of Botany and Head of the Department of Botany
in the University of Minnesota*

MINNEAPOLIS
THE H. W. WILSON COMPANY
1909

COPYRIGHT 1909
FREDERIC E. CLEMENTS

QK
603
C62
1909

PREFACE.

The present book is an outgrowth of a translation of the keys in the original eight volumes of Saccardo's "Sylloge Fungorum." This translation was mimeographed and bound for the use of classes in mycology. It immediately proved so convenient and usable that the preparation of a complete guide to the fungi was begun the same year. Many things have occurred during the past two years to delay the completion of the guide until this time. In its present form, the book is based upon Saccardo's great work, though in certain groups other authors have been followed, and in some cases, the discomycetes and lichens, the treatment amounts almost to a revision. The arrangement of the orders and families is different in a large measure, and in the distribution of the lichens is original. No attempt has been made to revise the genera, except where the treatment had lagged behind current practice, as is particularly true of the lichens. In some cases, genera have been included in others, but this is done only for the sake of the beginner, when the descriptions reveal no differences, and is by no means intended as a revision.

Questions of nomenclature have necessarily been left largely to one side, but no hesitation has been felt in making certain corrections. These have dealt mostly with mistaken or neglected transliteration, and with faulty composition. A considerable number of sesquipedalian words have been shortened, and the greater number of hybrid names have been corrected. These corrections have been made in such a way as to retain as much of the original name as possible. Corrections are indicated by the sign † with the original form in parenthesis below. New genera are designated by an asterisk, and are listed with their types on a later page.

The genera described in volumes 9-18 of the "Sylloge" have been included in the proper family keys. Genera placed under "incertae sedis" are excluded as a rule, since it is impossible to locate them definitely. A few genera occur more than once when they show the characters of two families, or when superficial and developmental features indicate different positions. An endeavor has been made to make the keys as consistent as possible, and as simple as is profitable. The mycologist must have a fair equipment of technical terms, as well as a Latin vocabulary, and the sooner these are acquired the better. In many cases, definiteness will seem to be lost by the use of such terms as "typically," "usually," etc., but the beginner must quickly learn that the line between families is rarely clear-cut, but often on the contrary most devious. The tyro must constantly be warned that some species belong as naturally in one family as in another, and must consequently be sought in more than one place. The color of a spore, the position of a perithecium, or the texture of a cup does not always

conform with a definite term, and the beginner must be governed accordingly.

While the writer is particularly indebted to Saccardo's "Sylloge Fungorum," he is also indebted to Thaxter's "Monograph of the Laboulbeniaceae," and his "Preliminary Diagnoses of New Species of Laboulbeniaceae," II-VI, for the material for the key to this group. The treatment of the Pezizales is largely that of Rehm's "Discomyceten," modified by the inclusion of the lichens. From Engler and Prantl's "Pflanzenfamilien," material has been drawn in the monographs of the bacteria by Migula, of phycomycetes and other groups by Schröter and Lindau, and especially of the lichens by Zahlbruckner. The writer is also under heavy obligation to Dr. Edith Clements, for the preparation of the Glossary, and for much other work of preparation and of publication. His thanks are also due to Professor Raymond J. Pool for assistance in the original mimeograph copies.

FREDERIC EDWARD CLEMENTS.

The University of Minnesota,

June 1, 1909.

CONTENTS.

PREFACE	
KEY TO ORDERS AND FAMILIES	I
KEY TO GENERA	7
KEY TO SPORE SECTIONS	165
GUIDE TO THE VOLUMES OF SACCARDO'S "SYLLOGE FUNGORUM"	167
INDEX TO FAMILIES IN SACCARDO'S "SYLLOGE FUNGORUM" AND REHM'S "DISCOMYCETEN"	171
LIST OF NEW GENERA AND TYPES	173
GLOSSARY OF LATIN AND ENGLISH TERMS	177
INDEX TO GENERA, SUBFAMILIES, FAMILIES AND ORDERS	207

Key to Orders and Families

- I. Filaments one-celled, rarely septate, typically aquatic or endobiotic; propagation by fission or by conidia, the latter usually in sporangia; sex-cells typically present, uniting to form resting-spores **Phycomycetes** **1**
- II. Filaments septate, typically saprophytic or epibiotic; conidia borne on conidiophores; sex-cells usually absent
 - 1. Spores in a hymenium composed of asci or club-shaped basidia
 - a. Spores in asci **Ascomycetes** **2**
 - b. Spores on more or less club-shaped basidia **Basidiomycetes** **5**
 - 2. Conidia on conidiophores of various form, not in asci or on true basidia **Fungi Imperfecti** **6**

Phycomycetes

- I. True mycelium lacking or rudimentary
 - 1. Threads simple, globose to filamentous, often motile; propagating by fission or by conidia also
 - a. Cells single or in colonies, never forming plasmodium-like masses **Bacteriales** **7**
 - (1) Cells filamentous, not spirally twisted
 - (a) Filaments motile, sheathless **Beggiatoaceae** **7**
 - (b) Filaments non-motile, sheathed **Chlamydobacteriaceae** **7**
 - (2) Cells cylindric to globose, spirally twisted when filamentous
 - (a) Cells more or less spirally twisted **Spirillaceae** **7**
 - (b) Cells not spirally twisted or curved
 - x. Cells oblong to cylindric **Bacteriaceae** **8**
 - y. Cells globose or cuboid **Coccaceae** **8**
 - b. Cells secreting a gelatinous matrix and forming pseudoplasmodia, passing into cysts or spore-masses which are often stalked **Myxobactrales** **8**
 - 2. Threads absent or slightly developed; propagation by sporangia which produce zoogonids; sex-cells rare **Chytridiaceae** **9**
- II. Mycelium present, typically well-developed and branched; propagation by zoogonids or by non-motile conidia borne in sporangia or on conidiophores; sex-cells usually present
 - 1. Aerial fungi propagating by conidia
 - a. Conidia typically in globose to cylindric sporangia; mostly saprophytes; zygosporous **Mucoraceae** **12**

b. Conidia single or in chains on conidiophores		
(1) Typically parasitic on insects; zygosporous	Entomophthoraceae	14
(2) Typically parasitic on leaves and stems; oosporous	Peronosporaceae	17
2. Typically aquatic fungi propagating by zoogonids		
a. Mycelium mostly well-developed		
(1) Antheridial tube touching or penetrating oogone	Saprolegniaceae	15
(2) Antherids producing antherozoids	Monoblepharidaceae	18
b. Mycelium more or less scanty, developing wholly or chiefly into sporangia and sex-organs	Ancylistaceae	16

Ascomycetes

I. Asci completely or partly enclosed in a pericarp		
1. Asci in a perithecium		
a. Perithecia one to many on a receptacle; sex-organs present; typically on insects	Laboulbeniales	18
b. Perithecia not on a receptacle; sex organs very rare; rarely on insects	Sphaeriales	21
(1) Mycelium or subicle typically present; ostiole and paraphyses usually absent		
(a) Subicle white; perithecia usually with appendages; asci one to few, more or less ovoid	Erysibaceae	21
(b) Subicle dark or black; appendages mostly lacking; asci usually numerous, more or less cylindric		
x. Perithecia more or less globose	Perisporiaceae	22
y. Perithecia clavate to cylindric, often branched	Capnodiaceae	25
(2) Subicle usually absent; ostiole and paraphyses typically present		
(a) Perithecia fleshy or waxy, bright colored	Hypocreaceae	42
(b) Perithecia hard, membranous to carbonous, typically brown to black		
x. Perithecia distinct, not reduced to cavities or locules		
(x) Perithecia normally globose, single, clustered or in a stroma		
m. Mycelium not forming a thallus with algae	Sphaeriaceae	25
n. Mycelium forming a thallus	Verrucariaceae	38
(y) Perithecia flattened, dimidiate and radiate	Microthyriaceae	51
(z) Perithecia with a broad and compressed or a funnellform ostiole		
m. Ostiole broad and compressed, cleft; perithecia mostly carbonous	Lophiostomataceae	53
n. Ostiole elongate, then expanded and		

funnel form; perithecia mostly coriaceous	Coryneliaceae	54
y. Perithecia reduced to locules in a stroma		
(x) Thallus absent		
m. Stromata mostly carbonous or membranous, not attached by a stipe-like point	Dothideaceae	48
n. Stromata subcarnose, attached by a stipe-like point	Coccoideaceae	50
(y) Thallus present	Mycoporaceae	50
2. Asci in a hysterothecium, i. e., a perithecium with a cleft-like ostiole, typically oblong to linear, rarely vertical	Hysteriales	54
a. Hysterothecium imperfect, dimidiate-scutate, but the ostiole a cleft	Hemihysteriaceae	54
b. Hysterothecium more or less elongate and rimose, or rounded and stellately cleft		
(1) Hysterothecium elongate, rimose, rarely vertical		
(a) Thallus absent	Hysteriaceae	55
(b) Thallus present	Graphidaceae	58
(2) Hysterothecium round to linear, ostiole more or less stellate or lobed; thallus present or absent	Arthoniae	58
3. Asci in an apothecium	Pezizales	61
a. Apothecia closed at first, then open, disk-shaped to cup-shaped, rarely elongate		
(1) Thallus lacking		
(a) Apothecia sunken, then erumpent, usually opening by lobes, rarely by a cleft		
x. Apothecia opening by stellate or irregular lobes or by a cleft		
(x) Apothecia dark, brown or black		
m. Apothecia mostly carbonous or leathery; hypothecium thin	Phacidiaaceae	61
n. Apothecia mostly membranous or horny; hypothecium thick	Tryblidiaceae	65
(y) Apothecia white or bright colored, typically waxy	Stictidaceae	62
y. Apothecia usually opening circularly, mostly leathery or horny, brown or black	Dermateaceae	65
(b) Apothecia typically superficial and opening circularly, usually waxy or fleshy but often carbonous, gelatinous or leathery		
x. Asci disappearing early; spores and paraphyses forming a mazaedium	Caliciaceae	70
y. Asci persistent; mazaedium lacking		

(x) Apothecia not branched-stipitate at the tips of branches		
m. Apothecia gelatinous	Bulgariaceae	66
n. Apothecia not gelatinous		
(m) Apothecia usually dark or black, carbonous to leathery, rarely waxy	Patellariaceae	68
(n) Apothecia usually bright colored, waxy to fleshy		
r. Apothecia typically waxy, on plant parts		
(r) Exciple brownish, parenchymatic all over or at the base; mostly sessile	Mollisiaceae	84
(s) Exciple concolorous, prosenchymatic; mostly stalked	Helotiaceae	85
s. Apothecia typically fleshy, usually terrestrial, often fimicole		
(r) Apothecia usually terrestrial, medium to large; asci mostly cylindric, not exerted	Pezizaceae	88
(s) Apothecia usually fimicole; asci broad, exerted from disk at maturity	Ascobolaceae	92
(y) Apothecia branched-stipitate at the tips of branches	Cordieritaceae	92
(2) Thallus present		
(a) Asci disappearing early; disk with a mazaedium	Caliciaceae	70
(b) Asci persistent; mazaedium absent		
x. Thallus cottony, cobwebby or spongy; algae yellow-green	Chrysotrichaceae	72
y. Thallus more or less distinctly gelatinous; algae blue-green	Collemataceae	72
z. Thallus firm, layered, neither gelatinous nor cottony		
(x) Thallus of two sorts: one horizontal, the other erect, i. e., a podetium	Cladoniaceae	78
(y) Thallus of one sort only, horizontal or erect		
m. Spores typically 2-celled, with a thickened cross-wall, usually traversed by a narrow canal	Physciaceae	83
n. Spores without thickened cross-wall and intersecting canal		
(m) Apothecia sunken, or grown together with the thallus on the whole underside	Peltophoraceae	75
(n) Apothecia typically superficial when mature, not attached broadly		

r. Apothecia with proper exciple	Lecideaceae	76
s. Apothecia typically with thalline exciple	Parmeliaceae	78
b. Apothecia open from the first, stalked, saddle-shaped, pileate to club-shaped, terrestrial as a rule	Helvellaceae	90
4. Asci in a closed globoid body or ascoma, containing cavities or veins	Tuberales	94
a. Ascomata epigean		
(1) Ascomata fleshy with locules at the margin, forming swellings on branches of living trees	Cyttariaceae	94
(2) Ascomata minute, waxy to subcarbonous, crowded with locules containing a single ascus each	Phymatosphaeriaceae	95
(3) Ascomata fragile, asci evanescent, then powdery within; epizoid	Onygenaceae	96
b. Ascomata hypogean		
(1) Ascomata woody, crustose or carbonous, powdery within	Elaphomycetaceae	96
(2) Ascomata fleshy or waxy, not powdery but veined or lacunose within	Tuberaceae	96
II. Asci exposed, apothecium lacking		
I. Spores free in the ascus	Gymnascales	93
a. Asci parallel and crowded, usually deforming living plant parts	Exascaceae	93
b. Asci solitary or grouped irregularly, saprophytic or when parasitic scarcely deforming the host	Gymnascaceae	93
c. Asci abnormal, rare; mycelium poorly developed, propagating by budding	Saccharomycetaceae	94
2. Spore wall united with ascus wall, or asci disappearing at maturity	Uredinales	98
a. Spores and ascus united; aecidia and uredinia often present	Uredinaceae	98
b. Asci disappearing early, leaving a firm or powdery spore-mass	Ustilaginaceae	101
Basidiomycetes		
I. Hymenium variously modified, exposed at maturity	Agaricales	102
1. Basidia septate crosswise or lengthwise, or furcate; usually gelatinous	Tremellaceae	103
2. Basidia not septate; pileus fleshy, waxy, leathery or woody		
a. Hymenium more or less uniform		
(1) Pileus funnel-form, dimidiate or resupinate	Thelephoraceae	106
(2) Pileus club-shaped, coralloid or filiform	Clavariaceae	105
b. Hymenium modified into teeth, pores or gills		
(1) Hymenium of teeth or granules	Hydnaceae	107
(2) Hymenium of pores or tubes	Polyporaceae	108

(3) Hymenium of gills or gill-like veins	Agaricaceae	110
II. Definite hymenium lacking; spore-mass gelatinous or powdery, typically enclosed in a peridium, or elevated at maturity	Lycoperdales	115
1. Gleba more or less gelatinous, enclosed at first in a volva, then raised on the receptacle	Phallaceae	115
2. Gleba firm or powdery, not gelatinous, enclosed in a peridium		
a. Peridium epigean		
(1) Gleba typically powdery or cellular, enclosed in a more or less globose peridium which opens irregularly or by a definite mouth	Lycoperdaceae	116
(2) Gleba in seed-like sporiangioles which are borne in a more or less cup-shaped peridium	Nidulariaceae	120
b. Peridium hypogean, closed	Hymenogastraceae	119

Fungi Imperfecti

I. Conidia present		
1. Conidia in globoid, cup-shaped or hysteroioid pycnidia	Phomatales	121
a. Pycnidia fleshy or waxy, bright colored	Zythiaceae	128
b. Pycnidia typically membranous to carbonous, dark, brown or black		
(1) Pycnidia more or less globose, rarely cylindric	Phomataceae	121
(2) Pycnidia dimidiate, shield-shaped	Leptostromataceae	130
(3) Pycnidia disciform, cup-shaped or hysteroioid	Excipulaceae	133
2. Conidia not in pycnidia		
a. Hyphae short or obsolete, borne on a matrix or stratum	Melanconiales	135
b. Hyphae not on a matrix, typically well-developed, but sometimes short or even lacking	Moniliales	138
(1) Hyphae in more or less loose cottony masses		
(a) Hyphae and conidia clear or bright colored	Moniliaceae	138
(b) Hyphae and conidia both typically dark or one or the other always dark	Dematiaceae	146
(2) Hyphae compactly united to form a globose to cylindric body which is often stalked		
(a) Hyphal body cylindric to capitate, stalked, i. e., a synnema	Stilbaceae	154
(b) Hyphal body more or less globose, sessile, i. e., a sporodochium	Tuberculariaceae	158
II. Conidia lacking	Sterile Mycelia	164

Key to the Genera

Class 1. SCHIZOMYCETES

Typically one-celled fungi, dividing by fission in 1, 2 or 3 planes, sometimes forming true filaments, but then motile or sheathed, and without true branches; resting cells often developed; sexual reproduction lacking.

Order 1. BACTERIALES

Globose, rod-like or filamentous, single or in colonies, sometimes grouped into a loose mass (zoogloea), but never forming pseudoplasmodia or sporangium-like masses.

Family 1. BEGGIATOACEAE

MIGULA 40

Filaments simple, free, motile, continuous or septate, sheathless, usually filled with shining or yellowish sulphur granules.

A single genus

Beggiatoa 8: 935

Family 2. CHLAMYDOBACTERIACEAE

MIGULA 35

Filaments simple or false-branched, typically attached, non-motile, septate, with a more or less conspicuous sheath; propagation by ciliate, creeping or non-motile conidia.

I. Cells without sulphur granules

1. Filaments simple

a. Fission always in one plane

Nocardia 8: 927

b. Fission in 3 planes during conidia formation

(1) Filaments marine, sheath very thin

Phragmidiothrix 8: 935

(2) Filaments fresh-water, sheath distinct

Crenothrix 8: 925

2. Filaments false-branched

Cladothrix 8: 927

II. Cells with sulphur granules

Thiothrix 8: 934

Family 3. SPIRILLACEAE

MIGULA 30

One-celled, more or less spirally twisted, rod-like or short-filamentous, usually motile by means of one to many flagella.

I. Cells stiff or rigid

I. Flagella lacking	Spirosoma M. 31
2. Flagella present	
a. Flagellum 1, rarely 2-3, polar	Microspira M. 31
b. Flagella clustered, polar	Spirillum 8: 1006
II. Cells flexible	Spirochaete 8: 1006

Family 4. BACTERIACEAE

MIGULA 20

One-celled, cells oblong to cylindric, straight or at least never spirally curved, flagella often present.

I. Flagella lacking	Bacterium 8: 1020
II. Flagella present	
1. Flagella peripheral	Bacillus 8: 943
2. Flagella polar	Pseudomonas M. 29

Family 5. COCCACEAE

MIGULA 15

One-celled, cells globose, usually flattened when grouped in rows or masses, flagella usually absent.

I. Flagella lacking	
1. Fission in one plane, cells in rows	Streptococcus 8: 1054
2. Fission in two planes, cells in plates	Micrococcus 8: 1076
3. Fission in three planes, cells in bundles	Sarcina 8: 1044
II. Flagella present	
1. Fission in two planes	Planococcus M. 19
2. Fission in three planes	Planosarcina M. 20

Order 2. MYXOBACTRALES

Cells rod-like, motile, fission in one plane; cells secreting a gelatinous base and forming pseudoplasmodia, then passing into cysts, or spore-masses which are often stalked (cystophore).

Family 6. MYXOBACTERIACEAE

11: 460, T. 389

Characters of the order.

I. Cells always rod-like, distinct cysts present	
1. Cysts free, usually on a cystophore	Chondromyces 14: 842
2. Cysts one or more in a gelatinous matrix	Myxobacter 14: 844
(Polyangium 7: 47)
II. Cells finally forming rows of globose spores, no definite cysts	
	Myxococcus 14: 843

Class 2. CHLOROPHYCEAE

Typically one-celled or filamentous plants, for the most part chlorophyllous but

each order containing at least one fungous family; propagation by fission and zoogonids; sexual reproduction present in most.

Order 3. PROTOCOCCALES

Typically one-celled algae, usually dividing by fission and producing zoogonids; sexual reproduction often lacking; one fungous family.

Family 7. CHYTRIDIACEAE

7:286, SCHROETER 65

Mycelium lacking or in the form of delicate protoplasmic threads, rarely of hyphae, one-celled; sporangiophore lacking or but slightly developed; sporangia producing zoogonids, thin-walled and ripening quickly, or thick-walled and resting for a time (resting sporangia); sexual reproduction present in a few forms, the sex organs scarcely distinguishable.

Key to the Subfamilies

- I. Resting sporangium asexual, rarely formed by the union of two zoogonids
 - 1. Mycelium completely lacking
 - a. Sporangia separate, one formed from each fruit-mass
Olpidiae
 - b. Sporangia in sori, formed by division of fruit-mass
Synchytriae
 - 2. Mycelium present
 - a. Mycelium of delicate transient strands
 - (1) Mycelium limited to one terminal sporangium
Rhizidiae
 - (2) Mycelium extended, sporangia intercalary and terminal
Cladochytriae
 - b. Mycelium consisting of permanent hyphae
Hyphochytriae
- II. Sexual resting spores formed by union of two sporangia and passing of contents of one into the other
Oochytriae
- III. Sexual spores formed by conjugation
Zygochytriae

Subfamily Olpidiae

SCHROETER 67

Mycelium lacking; fruit-mass endobiotic, globose, elliptic, rarely subclavate, undivided, finally forming a simple zoosporangium or resting sporangium, in which zoospores are formed after a period of rest.

- I. Fruit-body amoeboid before maturity
Reessia 7:304, S. 67
- II. Fruit-body without movement
 - 1. Sporangia free in the host-cell
 - a. Membrane delicate, dissolving to free zoospores
Sphaerita 7:314, S. 67
 - b. Membrane firm, with a definite opening
 - (1) Sporangia globose or elliptic
 - (a) Sporangia with 1, rarely 2, openings

- x. Zoospores 1-ciliate; resting sporangium smooth
Olpidium 7: 310, S. 67
- y. Zoospores 2-ciliate; resting sporangium spiny or warted
Olpidiopsis 7: 299, S. 69
- (b) Zoosporangia with many openings
Pleotrachelus 7: 315, S. 69
- (2) Sporangia elongate or clavate
Ectrogella 7: 315, S. 70
- 2. Wall of sporangium fused with wall of host-cell
Pleolpidium S. 70

Subfamily Synchytriae

SCHROETER 71

Mycelium lacking; fruit-body endobiotic, when mature dividing simultaneously to form zoosporangia grouped in rows or in a sorus; resting sporangia arising directly from the fruit-body or by the division of it.

- I. Zoosporangia arising through direct division of entire plasm of fruit-body, not surrounded by a common membrane
 - 1. Sporangia filling host-cell completely, wall fused with that of host-cell
Rozella 7: 300, S. 71
 - 2. Sporangia free, aggregated
Woronina 7: 301, S. 71
- II. Zoosporangia arising through division of the full-grown fruit-body, surrounded by the common membrane of the mother cell
 - 1. Sporangia formed directly from the full-grown fruit-body
Synchytrium 7: 288, S. 72
 - 2. Sporangia formed from the division of a thin-walled mother-cell which escapes from the fruit body
Pycnochytrium S. 73

Subfamily Rhizidiace

SCHROETER 75

Fruit-body endophytic, epiphytic, or living free between the nutrient media, at base, with a slender (in epiphytic forms sometimes scarcely perceptible) often branched mycelium, distinct for each fruit-body and imbedded in the matrix; zoosporangia globose or oblong, simple, often with a sterile swollen cell at base; zoospores globose, 1-ciliate; resting sporangia formed asexually, usually like the zoosporangia.

- I. Zoosporangia breaking out with an irregular or tube-like mouth, like the resting sporangia, which arise at the same place; mycelium delicate
 - 1. Sporangia without basal cell, arising directly from mycelium
Entophlyctis 14: 443, S. 75
 - a. Sporangia endophytic
 - b. Sporangia epiphytic or free
 - (1) Sporangia epiphytic, seated thickly on host-cell
Rhizophidium 7: 298, S. 76
 - (2) Sporangia free, mycelium only penetrating nutrient medium
 - (a) Zoospores escaping singly
Rhizophlyctis 14: 445, S. 77
 - (b) Zoospores escaping as a ball
Nowakowskia 7: 313, S. 77
 - 2. Sporangia with stalk-like or swollen basal cell
 - a. Sporangia with a stalk-like cell
 - (1) Epiphytic; stalk separated by wall from sporangium

- (a) Sporangium straight, rounded above
Podochytrium S. 77
- (b) Sporangium curved, pointed above
Harpochytrium 11: 249, S. 77
- (2) Saprophytic; stalk not separated from sporangium
Obelidium 7: 299, S. 77
- b. Sporangia with swollen basal cell
 - (1) Sporangium and basal cell endophytic
Diplophlyctis S. 78
 - (2) Sporangium epiphytic or free
 - (a) Sporangium epiphytic
 - x. Zoospores escaping singly
Phlyctochytrium S. 78
 - y. Zoospores escaping in a ball
Rhizidiomyces 7: 316, S. 79
 - (b) Sporangia saprophytic, free
Rhizidium 7: 296, S. 79
- II. Zoosporangia opening by a lid, epiphytic; resting sporangia endophytic, mycelium tubular or saccate
Chytridium 7: 304, S. 80

Subfamily Cladochytriae

SCHROETER 80

Mycelium diffuse, repeatedly branched, saprophytic, intercellular or intracellular, forming many sporangia, delicate, disappearing by the maturity of the spores; sporangia intercalary or terminal, zoospores 1-ciliate; resting sporangia produced asexually.

- I. Resting sporangia alone present
Physoderma 7: 317, S. 81
- II. Zoosporangia alone present
Cladochytrium 7: 295, S. 81
 - 1. Endophytic, intracellular
 - 2. Free, in algal slime
 - a. Sporangia opening by a hole
Amoebochytrium 7: 315, S. 82
 - b. Sporangia opening by a lid
Nowakowskiella 17: 514, S. 82

Subfamily Harpochytriae

SCHROETER 83

Mycelium strongly developed, cylindric, persistent; sporangia alone known, formed asexually.

- I. Mycelium and sporangia in the host-cell
Catenaria 9: 360, S. 83
- II. Sporangia in part at least free
 - 1. Parasitic
 - a. Mycelium endophytic
Harpochytrium 11: 249, S. 84
 - b. Mycelium endozoic
Polyrrhina 7: 314, S. 84
 - 2. Saprophytic
Tetrachytrium 7: 295, S. 84

Subfamily Oochytriae

SCHROETER 84

Mycelium lacking or variously developed; resting sporangium formed by the union of two young fruit-bodies, in which the plasm of one passes into the other which develops as an oogone; zoosporangia present, spherical to elongate.

- I. Mycelium entirely lacking
Diplophysa 7: 302, S. 85
- II. Mycelium present

1. Mycelium producing a single fruit-body **Polyphagus** 7: 302, S. 85
2. Mycelium producing several fruit-bodies **Urophlyctis** 7: 303, S. 86

Subfamily Zygochytriae

SCHROETER 87

Mycelium one-celled, upright, branched, producing zoospores and zygospores; zoosporangia single on ends of the branches, opening by a lid, zoospores one-ciliate; zygospores produced by the fusion of the end-cells of conjugating tubes, growing into a filament upon germination; intermediate between Chytridiaceae and Mucoraceae.

A single genus

Zygochytrium 7: 294, S. 87

Order 4. SPIROGYRALES

Typically one-celled or simple filamentous algae, without zoospores; sexual reproduction by the conjugation of similar gametes; two fungous families.

Family 8. MUCORACEAE

SCHROETER 119, 7: 182, 9: 335, 11: 239, 14: 432, 16: 383, 17: 494

Saprophytes, rarely parasites, with a well-developed branching mycelium in which cross-walls are absent; propagation by spores (conidia) arising within sporangia, the latter apparently reduced to chains of conidia in one family; reproduction by the union of the end-cells or gametes of conjugating tubes.

Key to the Subfamilies

- I. Sporangia always present, conidia sometimes present
 1. Columella present; zygospore naked or with a few appendages
 - a. Wall of the sporangium homogeneous, not cuticularized, diffluent
Mucorae
 - b. Wall cuticularized and persistent above, thin and diffluent below
Pilobolae
 2. Columella absent; zygospore enveloped in a dense covering
Mortierellae
- II. Sporangia rarely present, conidia always present
 1. Conidia solitary; zygospore arising directly from the gametes
 - a. Sporangia present
Choanophorae
 - b. Sporangia lacking
Chaetocladiace
 2. Conidia in chains; zygospore arising from outgrowths of gametes
Syncephalidae

Subfamily Mucorae

7: 184, S. 123

Mycelium similar throughout or consisting of aerial and nutritive parts; sporangia alike or of two sorts, primary and accessory, the former with columella, the latter mostly without one; zygospore naked or with separate appendages arising from the suspensors.

I. Sporangia similar

1. Sporangiphore simple or branched, but not repeatedly dichotomous

a. Suspensors without appendages at maturity

(1) Aerial mycelium lacking

(a) Sporangia single, terminal

Mucor 7: 190, S. 124

(b) Sporangia clustered, lateral

x. Sporangia globose

Circinella 7: 215, S. 125

y. Sporangia long pear-shaped

Pirella 7: 216, S. 125

(2) Aerial mycelium present

(a) Aerial mycelium stoloniferous

Rhizopus 7: 212, S. 125

(b) Aerial mycelium with many short thorn-like branches

Spinellus 7: 205, S. 125

b. Suspensors with thorny appendages at maturity

(1) Appendages spreading

Phycomyces 7: 204, S. 126

(2) Appendages loosely enclosing the zygospore

Absidia 7: 214, S. 126

2. Sporangiphore repeatedly dichotomous

Sporodinia 7: 206, S. 127

II. Sporangia of two sorts, primary and secondary

1. Primary sporangia with, secondary without columella

Thamnidium 7: 211, S. 127

2. Both kinds of sporangia with columella

Dicranophora 11: 240, S. 128**Subfamily Pilobolae**

7: 184, S. 123

Mycelium similar throughout; sporangia alike, with columella, sporangial wall cuticularized and persistent above; zygospores naked.

I. Sporangiphore equal, sporangium not thrown off

Pilaira 7: 188, S. 129

II. Sporangiphore swollen above, sporangium thrown off

Pilobolus 7: 184, S. 129**Subfamily Mortierellae**

7: 184, S. 130

Sporangia similar, terminal, without columella; conidia single, spherical on short lateral branches of the aerial mycelium; zygospore enclosed in a dense mass of hyphae arising from the suspensors.

I. Sporangiphores erect, branches attenuate toward tip

Mortierella 7: 220, S. 130

II. Sporangiphores creeping, branches equal

Herpocladiella 7: 225, S. 130**Subfamily Choanophorae**

9: 339, S. 131

Mycelium parasitic on plant parts; sporangia and conidia both present; conidio-

phores simple or branched, bearing one-celled conidia; sporangiophores simple, sporangia with a small columella.

A single genus

Choanophora 9: 339, S. 131

Subfamily Chaetocladiae

7: 220, S. 131

Mycelium parasitic on species of Mucor; propagation by conidia, sporangia lacking, conidia arising on short side branches; zygosporangia arising directly from the fused gametes.

A single genus

Chaetocladium 7: 220, S. 131

Subfamily Syncephalidae

7: 225, S. 132

Conidia in chains on short basidia borne on the end of the sporophores; zygosporangia arising as an outgrowth from the tips of the suspensors after conjugation.

I. Sporophores not swollen at tip

Piptocephalis 7: 225, S. 132

II. Sporophores swollen into a head at tip

1. Sporophore simple

Syncephalis 7: 227, S. 132

2. Sporophore branched

Syncephalastrum 7: 232, S. 134

Family 9. ENTOMOPHTHORACEAE

SCHROETER 134, 7: 280, 9: 349, 14: 437, 16: 388, 17: 510

Mycelium usually well-developed, tubular or filamentous, mostly parasitic or endozoic, rarely saprophytic, at first one-celled, then septate; propagation by one-celled conidia terminal on one-celled clavate conidiophores; zygosporangia globose.

I. Mycelium endozoic (in insects)

1. Conidia always present

a. Conidiophore simple, zygosporangia unknown, azygosporangia present

(1) Cystidia and holdfasts lacking; azygosporangia lateral

Empusa 7: 281, S. 138

(2) Cystidia and holdfasts present; azygosporangia terminal

Lamia S. 139

b. Conidiophore repeatedly branched, zygosporangia and azygosporangia present

Entomophthora 7: 282, S. 139

2. Azygosporangia alone present

Tarichium 7: 284, S. 140

II. Mycelium endophytic or saprophytic

1. Mycelium little developed, intracellular

Completozia 7: 286, S. 140

2. Mycelium well-developed, not intracellular

a. Parasitic on fungi

Conidiobolus 7: 285, S. 141

b. Saprophytic

- **Basidiobolus** 7: 285, S. 141

Order 5. VAUCHERIALES

Unicellular, multinucleate, saccate or filamentous algae and fungi; propagation by zoospores or conidia; sexual reproduction in the three fungous families by unlike gametes, produced in antherids and oogones.

Family 10. SAPROLEGNIACEAE

SCHROETER 93, 7: 264, 9: 345, 11: 244, 14: 450, 16: 395, 17: 519

Mycelium strongly developed, broadly filamentous, more or less branched; propagation by zoosporangia, producing ciliate, rarely non-motile, zoospores; sexual reproduction by antherids and oogones, their contents fusing by means of a connecting tube.

Key to the Subfamilies

- I. Vegetative mycelium broad, tubular, aquatic; zoosporangia cylindric, of the same width as the mycelium
 - 1. Filaments uniform, not constricted **Saprolegniae**
 - 2. Filaments constricted regularly **Leptomitae**
- II. Vegetative mycelium thin, mostly saprophytic on plant tissues; zoosporangia several times broader than the filaments **Pythiae**

Subfamily Saprolegniae

SCHROETER 96

Nutritive mycelium sunken in the substratum, finely branched, water mycelium tubular, repeatedly branched, cylindric; zoosporangia narrowly cylindric; oogones mostly terminal, globose, 1- to many-spored, antheridia clavate, the tube penetrating the oogone.

- I. Zoospores escaping before germination
 - 1. Zoosporangia cylindric-clavate, zoospores several-rowed
 - a. Zoospores escaping together through a terminal pore
 - (1) Zoospores scattering upon escape
 - (a) Zoosporangia ovate **Pythiopsis S. 97**
 - (b) Zoosporangia cylindric **Saprolegnia 7: 268, S. 97**
 - (2) Zoospores remaining massed about the pore
 - Achlya 7: 274, S. 99**
 - b. Zoospores not escaping through a common opening
 - (1) Each zoospore escaping singly through its own lateral pore
 - Dictyuchus 7: 273, S. 99**
 - (2) Zoospores freed by the falling apart of the whole sporangium
 - Thraustotheca S. 100**
 - 2. Zoosporangia linear, zoospores 1-rowed
 - a. Zoospores scattering upon escape **Leptolegnia S. 100**
 - b. Zoospores remaining in a ball at the pore
 - Aphanomyces 7: 276, S. 100**
- II. Zoospores germinating in the sporangium
 - Aplanes S. 101**

Subfamily Leptomitae

SCHROETER 101

Filaments thin, branched, divided by regular constrictions; zoosporangia cylindric, pear-shaped or elliptic; oogones 1-spored.

- I. Branches similar to the main stem

- I. Zoospores escaping singly from the pore
 - Leptomitus** 7: 265, S. 101
 2. Zoospores remaining in a hollow ball about the pore before swimming
Apodachlya S. 102
- II. Branches different from the main stem
 1. Branches whorled **Naegeliella** S. 163
 2. Branches repeatedly umbellate-ramose **Araeospora** 14: 454
 3. Branches springing from the swollen tip of the main stem
Rhipidium 7: 268, S. 103

Subfamily Pythiae

SCHROETER 104

Vegetative mycelium very narrow, uniform, much-branched; sporangiophores not distinct from mycelium; zoosporangium filamentous, cylindric, ellipsoid or globose, contents escaping in a globose vesicle in which the zoospores arise, zoospores 2-ciliate; oogones globose, terminal, rarely intercalary, 1-spored.

- I. Zoosporangia filamentous **Nematosporangium** S. 104
- II. Zoosporangia globose or lemon-shaped **Pythium** 7: 270, S. 104

Family 11. ANCYLISTACEAE

SCHROETER 89, 7: 278, 9: 348, 14: 450, 16: 395, 17: 516

Mycelium mostly poorly developed and scarcely distinct from the fruit-body, the latter tubular, when mature divided into vegetative cells, sporangia or oogones and antherids; entire contents of antherid passing into oogone, oospore lying free; sporangia always producing zoospores.

Key to the Subfamilies

- I. Filament or fruit-body producing wholly sporangia or sex cells, mycelium entirely lacking **Lagenidiace**
- II. Filament producing vegetative cells also, the latter germinating to form threads **Ancylistae**

Subfamily Lagenidiace

Fruit-body filamentous, tubular, simple or branched, dividing into cells which develop into sporangia or sex cells; antherids on the same or on different fruit bodies; sporangia and oospores always giving rise to zoospores.

- I. In fresh-water algae, rarely in animals
 1. Filament simple
 - a. Zoospores escaping singly from the sporangium
Achlyogeton 7: 277, S. 89
 - b. Sporangial plasm poured out into a vesicle in which the zoospores are formed
Myzocytiium 7: 279, S. 90
 2. Filament with short side-branches **Lagenidium** 7: 278, S. 90
- II. In the root-hairs of plants **Rhizomyxa** 7: 278, S. 91

Subfamily Ancylistae

Fruit-body tubular, mycelium-like, unbranched or with few short side-branches, when mature dividing into a number of chain-like cells, which develop into vegetative

cells, sporangia or sex cells; sporangia producing zoospores; vegetative cells producing a long tube, which penetrates new host-cells; oospores globose or elliptic.

I. Sporangia lacking, vegetative and sex cells alone formed

Ancylistes 7: 280, S. 92

II. Sporangia also present

Reticularia 9: 348, S. 92

Family 12. PERONOSPORACEAE

SCHROETER 110, 7: 233, 9: 340, 11: 242, 14: 457, 16: 396, 17: 519

Mycelium abundant, filamentous, much branched, one-celled, endophytic; propagation by conidia borne on the ends of conidiophores, conidia producing zoospores or a germinating tube; sexual reproduction by means of endophytic antherids and oogones, borne on the ends of lateral branches; oospores single, globose, producing zoospores or a germinating tube.

Key to the Subfamilies

I. Conidia in chains, conidiophores club-shaped

Albuginae

II. Conidia single, conidiophores branched

Peronosporae

Subfamily Albuginae

Mycelium intercellular, haustoria globose; conidiophores densely grouped into a conidial layer beneath the epidermis; conidia globose, ellipsoid or subcylindric, in chains on the ends of the conidiophores, usually producing zoospores, rarely a germinating tube; oospores globose, producing zoospores.

A single genus

Albugo 7: 233, S. 110

Subfamily Peronosporae

Mycelium intercellular, rarely intracellular, haustoria of various form; conidiophores thread-like, above the epidermis, branched, without cross-walls; conidia single on the tips of the branchlets, producing zoospores or a germinating tube; oospores globose, with a well-developed outer wall, germinating by means of a tube.

I. Conidiophores slender, with long and slender branches

1. Conidiophore growing after the formation of the first conidia, producing new joints

Phytophthora 7: 237, S. 113

2. Conidiophore not growing and making new extensions

a. Conidia papillate at the tip

(1) Conidia on stalks arising from irregular disks

Bremia 7: 243, S. 116

(2) Conidia on stalks arising directly from the unchanged ends of the conidiophores

Plasmopara 7: 239

b. Conidia not papillate at the tip

Peronospora 7: 244, S. 117

II. Conidiophores stout, swollen at the tip, or with short thick branches

1. Conidiophore simple up to the enlarged tip, which bears the conidia on slender stalks

Basidiophora S. 114

2. Conidiophore with short thick branches bearing the conidia on flask-like stalks

Sclerospora 7: 238, S. 114

Order 6. CONFERVALES

Typically multicellular filamentous algae, propagating by zoospores, and reproducing by the union of isogametes, or by heterogametes borne in antherids and oogones; one fungous family.

Family 13. MONOBLEPHARIDACEAE

SCHROETER 106, 7: 277, 14: 452, 16: 394

Mycelium filamentous, one-celled or septate, producing zoospores and sex cells; zoospores 1-ciliate arising in terminal sporangia; antherids cylindric producing ciliate antherozoids; oogones globose, terminal, opening by a pore, 1-spored.

I. Zoospores 1-ciliate

1. Mycelial threads equal throughout **Monoblepharis 7: 277, S. 107**
2. Mycelial threads constricted, necklace-like **Gonapodya 14: 452, S. 107**

II. Zoospores two or more ciliate

1. Zoospores 2-ciliate **Diblepharis 16: 395**
2. Zoospores many-ciliate **Myrioblepharis 14: 455**

Class 4. ASCOMYCETES

Fungi usually destitute of a conspicuous mycelium, reproducing by means of a spore-fruit containing asci (perithecium or apothecium), the spore-fruit occasionally reduced to a group of naked asci.

Order 7. LABOULBENIALES

THAXTER 197, LINDAU 491

Family 14. LABOULBENIACEAE

8: 909, 9: 1130, 11: 446, 14: 725, 16: 674, 17: 915

Receptacle consisting of two to many cells in a row, or parenchyma-like, regularly producing from the cells one or more appendages bearing antherids as a rule; antherozoids normally endogenous, borne within flask-like, simple or compound antherids, rarely produced like conidia, i. e., naked or exogenous; perithecia one to many, stalked or sessile, terminal or lateral on the receptacle, resulting from fertilization by means of a trichogyne; asci seriate, mostly 4-spored, spores usually 2-celled.

I. Antherozoids endogenous, i. e., in closed antherids

1. Antheridial cells forming a compound antherid
 - a. Dioecious
 - (1) Perithecia and appendages in pairs to the right and left **Dimorphomyces T. 264, L. 497**
 - (2) Perithecia and appendages in a row **Dimeromyces T. 267, L. 497**

b. Monoecious

- (1) Antherids arising on an appendage
 - (a) Antherids lateral
 - x. On a subbasal cell of the appendage **Cantharomyces T. 271, L. 497**

- y. On short opposite branchlets of the appendage
Stichomyces T. 4: 37
- (b) Antherids terminal
 - x. Antherid with a short spine at the tip
Haplomyces T. 269, L. 497
 - y. Antherid without a spine but with a neck-like canal cell
 - (x) Ascogenic cells at least 36 **Polyascomyces** T. 2: 414
 - (y) Ascogenic cells few
 - m. Stalk of antherid a single cell
 - (m) Antheridial cells obliquely in vertical rows
 - r. Subbasal cell of receptacle with a sterile appendage
Eumonoecomyces T. 4: 21
 - s. Subbasal cell of receptacle without sterile appendage
Eucantharomyces T. 273, L. 497
 - (n) Antherid parenchyma-like, many-celled
 - r. Antheridial cells with three marginal cells
Euhaplomyces T. 4: 25
 - s. Antherial cells without marginal cells
Camptomyces T. 274, L. 498
 - (o) Antherid of several superposed cells bearing single simple antherids directly
 - r. Simple antherids two **Acallomyces** T. 5: 23
 - s. Simple antherids several
Acompsomyces T. 4: 37
 - n. Stalk of two cells placed side by side
Monoecomyces T. 2: 412, 4: 23
 - (2) Antherids arising on the receptacle
 - (a) Perithecia free
 - x. Receptacle of a single row of several to many superposed cells
Enarthromyces T. 276, L. 498
 - y. Receptacle of one or two superposed cells followed by two or three oblique or transverse rows
 - (x) Receptacle with one basal cell
 - m. Basal cell followed by two tiers of cells
Limnaeomyces T. 2: 428
 - n. Basal cell followed by three symmetrical series
Dichomyces T. 282, L. 499
 - (y) Receptacle with two superposed basal cells
Peyritschiella T. 278, L. 499
 - (b) Perithecia grown together with distal portion of receptacle
 - x. Base of receptacle of two superposed cells
Chitonomyces T. 285, L. 499
 - y. Base of three superposed cells
Hydraeomyces T. 293, L. 500
 - 2. Antheridial cells distinct, discharging independently
 - a. Dioecious
 - (1) Perithecium borne by the basal or subbasal cell of receptacle
 - (a) Perithecium on the single basal cell, spores continuous
Amorphomyces T. 295, L. 501

- (b) Perithecium lateral on the subbasal cell, spores obliquely 1-septate
Dioecomyces T. 4: 33
- (2) Two-celled normal receptacle producing secondary receptacles on which the perithecia are borne
Herpomyces T. 5: 11
- b. Monoecious
 - (1) Antherids in definite series on the appendages
 - (a) Arising directly from cells of the appendages
 - x. Appendage one
 - (x) Antherids in 4 vertical series
Helminthophana T. 297, L. 501
 - (y) Antherids in a single vertical series
Stigmatomyces T. 298, L. 501
 - y. Appendages numerous, antherids in 3 vertical series
Idiomyces T. 302, L. 501
 - (b) Borne on branches of the appendages
 - x. Appendage one
 - (x) Appendage with sterile terminal branchlets, antherids in short series near its base
Rhadinomyces T. 305, L. 501
 - (y) Appendage with fertile terminal branchlets bearing antherids laterally
Eucorethromyces T. 2: 433
 - y. Appendages forming a tuft, antherids on lateral branchlets
Corethromyces T. 303, L. 501
 - (2) Antherids not in definite series on the appendages
 - (a) Receptacle 2-celled
 - x. Basal cell with rhizoids
 - (x) A single receptacle from each rhizoid base
Rhizomyces T. 307, L. 502
 - (y) Several receptacles from a common rhizoid base
Moschomyces T. 368, L. 504
 - y. Basal cell not from a rhizoid
 - (x) Appendage single
 - m. Receptacle of 2 superposed cells
 - (m) Basal cell spheric, penetrating by a long filament
Ceraiomycetes T. 3: 410
 - (n) Basal cell elongate
Sphaleromyces T. 365, L. 504
 - n. Receptacle of a series of superposed cells
Ectinomyces T. 5: 26
 - (y) Appendages several to many
 - m. Appendages and perithecium in a whorl
Compsomyces T. 366, L. 504
 - n. Appendages in a row
Clematomyces T. 2: 439
 - (b) Receptacle more than 2-celled
 - x. Receptacle of seriate, regularly superposed cells
 - (x) Plant bilaterally symmetrical
Diplomyces T. 357, L. 503
 - (y) Plant asymmetrical
 - m. Receptacle of two contiguous and united rows
 - (m) A single basal cell
Rhachomyces T. 358, L. 504

(n) Basal and subbasal cell present

Distichomyces T. 6: 308

n. Receptacle of a single row **Chaetomyces** T. 364, L. 504

y. Receptacle more or less parenchyma-like, at most only part of the cells superposed in series

(x) Appendages all on one side **Laboulbenia** T. 308, L. 502

(y) Appendages on two sides **Rickia** 16: 689

(z) Appendages completely surrounding the perithecium

Teratomyces T. 354 L. 502

II. Antherozoids exogenous, i. e., produced terminally or laterally on the appendages as naked cells

1. Receptacle large, very many-celled, parenchyma-like

a. Perithecium with six wall cells in each row

(1) Base of trichogyne persistent as a one-celled appendage

Caenomyces T. 4: 44

(2) Base of trichogyne not persistent as an appendage

Zodiomyces T. 371, L. 504

b. Perithecium with 9-10 wall cells in each row

Euzodiomyces T. 2: 449

2. Receptacle of a series of superposed cells

a. Appendage single

Ceratomyces T. 372, L. 505

b. Appendages several

Coreomyces T. 5: 56

The genus *Misgomyces* T. 2: 443 has not been included in the key owing to the fact that its antherids are unknown; it is very closely related, apparently, to *Laboulbenia*.

Order 8. SPHAERIALES

Mycelium sometimes superficial and abundant, often forming a thallus with algae, but usually scanty and imbedded in the matrix, the threads branched and septate; propagation by means of conidia borne on branches of the mycelium, or by means of pycnidia; reproduction resulting in a globose, flask-shaped or flattened perithecium, with a round mouth or ostiole except in the simpler forms, in which appendages are also often found; asci usually 8-spored and with paraphyses; spores hyaline, yellowish or brown, one to many-celled.

Family 15. ERYSIBACEAE

I: 1, 9: 364, II: 253, 14: 404, 17: 526

Mycelium white, cobwebby, superficial, penetrating the epiderm by means of haustoria; propagation by chains of conidia cut off from upright simple branches; perithecium without mouth, membranous, regularly with simple or modified appendages, often imbedded in the mycelium; ascus one to several, globose to ovoid, 2-8-spored, without paraphyses; spores usually 1-celled, hyaline.

Hyalosporae

Spores 1-celled, hyaline

I. Perithecium with one ascus

1. Appendages simple

Sphaerotheca 1: 3

2. Appendages dichotomously branched

Podosphaera 1: 2

II. Perithecium with several asci

1. Appendages present

a. Appendages simple, thread-like **Erysibe 1: 15**

b. Appendages branched or otherwise modified

(1) Appendages dichotomously branched

Microsphaera 1: 10

(2) Appendages modified but not branched

(a) Appendages stiff and bristle-like

x. Appendages numerous, not swollen at base

Pleochaete 1: 9

y. Appendages few, swollen at base

Phyllactinia 1: 5(b) Appendages coiled at tip **Uncinula 1: 6**

2. Appendages absent; perithecium surrounded by the mycelium

Erysibella 1: 23**Dictyosporae**

Spores usually hyaline, muriform

A single genus

Saccardia 1: 24**Family 16. PERISPORIACEAE**

1: 24, 9: 371, 11: 253, 14: 462, 16: 398, 17: 524

Mycelium superficial, dark, filamentous, sometimes lacking, rarely forming a firm stroma; conidia or pycnidia rarely present; perithecium without a mouth, or opening irregularly, usually globose, membranous or coriaceous, rarely carbonous, appendages usually lacking; asci mostly numerous, clustered, more or less cylindric, mostly 8-spored, paraphyses regularly lacking; spores various.

Hyalosporae

Spores 1-celled, hyaline or yellowish

I. Perithecia bright-colored, yellow or reddish, rarely white

1. Asci 8-spored

a. Perithecia with setae, or hairs

(1) With long rigid setae

Chaetotheca 11: 254

(2) With many hairs, immersed in a dense subicle

Cryptothecium 14: 465

b. Perithecia glabrous

(1) Spores with an unequal samariform appendage

Samarospora 11: 254

(2) Spores not appendaged

(a) Spores verrucose

Anixiopsis 14: 464

(b) Spores smooth

x. Conidiophores branched

Allescheria 14: 464

y. Conidiophores simple, swollen at tip

Eurotium 1: 25**(Kickxella 9: 372)****Pisomyxa 1: 29**

2. Asci many-spored

II. Perithecia dark or black, spores hyaline

1. Asci 2-8-spored

- a. Ascus single Cystotheca 16: 407
- b. Asci several or many
 - (1) Perithecia numerous in setose stroma-like cups Lasiobotrys 1: 29
 - (2) Perithecia not in cups
 - (a) Perithecia globose Meliolopsis 1: 68
 - (b) Perithecia applanate Asterula 1: 47
- 2. Asci many-spored
 - a. Asci many Apiosporium 1: 30
 - b. Ascus single Monascus 9: 373
- III. Perithecia brown, then black, spores yellow Anixia 1: 34

Phaeosporae

Spores 1-celled, dark

- I. Asci capitate on tips of branched hyphae Cephalotheca 1: 36
- II. Asci sessile or on simple stalks
 - 1. Perithecia with appendages
 - a. Spores globose, conglobate
 - (1) Appendages closely spiral, convolute Pleurascus 16: 1123
 - (2) Appendages flexuose-tortuose Arachnomycetes 17: 532
 - b. Spores ellipsoid
 - (1) Appendages several times branched Ascotricha 1: 37
 - (2) Appendages circinate at apex Magnusia 1: 38
 - 2. Perithecia without appendages
 - a. Perithecia hairy or setose Chaetomidium 1: 39
 - b. Perithecia glabrous
 - (1) Perithecia innate upon a radiate subicle Asteronia 1: 47
 - (2) Perithecia not on a radiate subicle
 - (a) Spores at first conglobate Laaseomyces 16: 405
 - (b) Spores free from the first
 - x. Growing on lichen thalli Orbicula 1: 38
 - y. Growing on roots Thielavia 1: 39

Hyalodidymae

Spores 2-celled, (1-septate), hyaline

- I. Asci 8-spored
 - 1. Cells of spore separating easily Neorehmia 17: 536
 - 2. Cells of spore not separating
 - a. Perithecia on a radiate subicle Asterella 1: 42
 - b. Perithecia on a uniform subicle Dimerosporium 1: 51
- II. Asci many-spored Pampolysporium 16: 411

Phaeodidymae

Spores 1-septate, dark when mature, rarely yellowish

- I. Perithecia on a subicle
 - 1. Subicle radiate: perithecia lenticular Asterina 1: 39
 - 2. Subicle uniform, dematium-like; perithecia globose

- a. Perithecia without basal setae
 - (1) Asci several or many **Dimerium** 1: 51, 17: 537
 - (2) Ascus one, rarely two **Balladyna** 16: 411
 - b. Perithecia with basal setae **Kusanobotrys** 17: 881
- II. Perithecia not seated on a subicle
- 1. Perithecia gelatinous when wet, honey-yellow **Englerula** 17: 529
 - 2. Perithecia membranous or carbonous, usually dark
 - a. Spores apiculate-appendaged, very large **Zopfia** 1: 54
 - b. Spores not appendaged, small or medium
 - (1) Spores smooth
 - (a) Spores elongate-oblong, very large **Richonia** 9: 379
 - (b) Spores subtrapeziform, small **Argynna** 14: 470
 - (c) Spores ellipitic, medium **Parodiella** 1: 717, 9: 409
 - (2) Spores spiny or roughened
 - (a) Perithecium irregularly dehiscent; asci not long-stalked **Marchaliella** 11: 257
 - (b) Perithecia regularly areolate-dehiscent; asci long-stalked **Testudina** 9: 378

Hyalophragmiae

Spores with 2 or more cross walls, hyaline

- I. Perithecia on a radiate subicle **Asteridium** 1: 49
- II. Perithecia on a uniform subicle
 - 1. Subicle effuse, dematium-like; perithecium closed **Zukalia** 9: 431
 - 2. Subicle fibrous, subcrustose; perithecium perforate **Perisporiopsis** 17: 544

Phaeophragmiae

Spores 2-several-septate, dark

- I. Perithecia on a radiate subicle **Meliola** 1: 60
(**Limacinia** 14: 474)
- II. Subicle uniform or absent
 - 1. Spores separating at the joints
 - a. Paraphyses lacking **Perisporium** 1: 55
 - b. Paraphyses present **Schenckia** 11: 268
 - 2. Spores not separating **Perisporina** 17: 545

Hyalodictyae

Spores muriform, hyaline

- I. Perithecia on a subicle, closed **Zukaliopsis** 17: 554

Phaeodictyae

Spores muriform, dark

- I. Perithecia globose
 - 1. Spores with an appendage at each end **Ceratocarpia** 14: 474
 - 2. Spores without appendages

- a. Subicle radiate **Pleomeliola** 1: 70, 17: 554
- b. Subicle lacking **Cleistotheca** 11: 270
- II. Perithecia applanate **Cookella** 1: 71

Scolecosporae

Spores filiform, septate or continuous, hyaline or subhyaline

- I. Perithecium opening by a small pore **Saccardomyces** 17: 530
- II. Perithecium without a pore
 - 1. Subicle radiate, paraphyses present **Ophiomeliola** 16: 416
 - 2. Subicle uniform, paraphyses absent **Hyaloderma** 9: 437

Family 17. CAPNODIACEAE

1: 73, 9: 438, 11: 270, 14: 476, 17: 555

Perithecia vertically elongate, clavate or cylindric, obtuse or acute, simple or branched, usually lacinate-dehiscent at the apex, on a thick black mycelium, which is rarely absent.

- I. Subicle crustose
 - 1. Spores 1-celled, globose **Capnodiella** 1: 74
 - 2. Spores 3-4-septate, dark **Capnodaria** 1: 74
 - 3. Spores muriform, dark **Capnodium** 1: 73, 80
- II. Subicle very thick, spongy **Scorias** 1: 83
- III. Subicle sparse or lacking
 - 1. Spores 1-celled, hyaline **Capnodiopsis** 17: 555
 - 2. Spores 2-celled, hyaline; perithecium gelatinous **Seuratia** 17: 558

Family 18. SPHAERIACEAE

1: 88, 2: 1, 9: 4, 11: 271, 14: 478, 16: 417, 17: 560

Mycelium scanty and immersed, or often producing a stroma, rarely a subicle; perithecia typically globoid, often drawn out into a beak, membranous, coriaceous, or carbonous, brown or black, dehiscing by a round pore or ostiole, single, cespitose or composite in a stroma; in the latter case each perithecium is distinct, not merely a locule in the stroma; asci usually numerous, elongate, usually paraphysate; spores various.

Allantosporae

Spores 1-celled, obtuse, curved-oblong, hyaline or olivascens

- I. Perithecia sparse or cespitose
 - 1. Ostiole central, very short
 - a. Asci 8-spored
 - (1) Perithecia covered
 - (a) Perithecia minute, glabrous **Massalongiella** 1: 89
 - (b) Perithecia largish, strigose-pilose **Enchnoa** 1: 89
 - (2) Perithecia subsuperficial
 - (a) Perithecia globose, never collapsing **Bizzozera** A: 24, 9: 445

- (b) Perithecia collapsing, becoming cup-shaped
 - x. Perithecia gregarious **Coelosphaeria** 1: 91
 - y. Perithecia cespitose **Nitschkea** 11: 272
- b. Asci many-spored **Fracchiacea** 1: 93
- 2. Ostiole central, papillate **Neoarcangelia** 16: 419
- 3. Ostiole lateral, conic **Pleurostoma** 1: 95
- II. Perithecia composite, typically in a stroma
 - 1. True stroma lacking; perithecia heaped together between bark and wood
 - a. Asci 8-spored; ostiole short or long **Calosphaeria** 1: 95 (16: 419, 421)
 - b. Asci many-spored; ostiole very short **Coronophora** 1: 103
 - 2. True stroma present; perithecia immersed in bark or wood
 - a. Stroma formed by the changed matrix
 - (1) Stroma valsous, i. e., perithecia in a circle
 - (a) Asci 4-8-spored
 - x. Perithecia usually 4, never more than 6, in each stroma **Quaternaria** 1: 106
 - y. Perithecia many, 8-30, in most stromata at least
 - (x) Perithecia circinate or monostichous, ostiole entire; asci subsessile. **Valsa** 1: 108
 - (y) Perithecia monostichous or polystichous, ostiole not entire; asci stipitate **Eutypella** 1: 145, 17: 569
 - (b) Asci many-spored **Valsella** 1: 158
 - (2) Stroma eutypeous, i. e., broadly and indefinitely effuse
 - (a) Asci 8-spored
 - x. Stroma conspicuous, cortical or woody **Eutypa** 1: 162, 17: 569
 - y. Stroma more or less obsolete
 - (x) Stroma woody; ostiole largish; spores subfuscous **Endoxyla** 1: 181
 - (y) Stroma cortical; ostiole small; spores subhyaline **Cryptosphaeria** 1: 182
 - (b) Asci many-spored
 - x. Stroma manifest, cortical or woody
 - y. Stroma obsolete, cortical **Cryptovalsa** 1: 187
 - b. Stroma different from the substance of the matrix **Cryptosphaerella** 1: 186

Hyalosporae

1: 407, A 58, 9: 577, 11: 289, 14: 515, 16: 452, 17: 573

Spores 1-celled, hyaline or nearly hyaline, ovoid, oblong or fusoid, rarely irregular or stellate, not allantoid.

- I. Perithecia single or separate
 - 1. Perithecia beaked or with a stellate ostiole
 - a. Perithecia subcarbonous

- (1) Spores normal, i. e., not modified
 - (a) Perithecia superficial, glabrous or dark hairy **Ceratostomella** 1: 408
 - (b) Perithecia innate-erumpent, yellow-hairy **Camptosphaeria** 1: 413
- (2) Spores with a ring-like appendage **Rostrella** 17: 609
- b. Perithecia submembranous, usually phyllogenuous
 - (1) Ostiole black, not stellate **Gnomoniella** 1: 413
 - (2) Ostiole white, stellate with black wartlike appendages **Rinia** 17: 591
- 2. Perithecia not beaked
 - a. Perithecia covered
 - (1) Asci 1-2- or 4-8-spored
 - (a) Paraphyses present **Physalospora** 1: 433
(incl. **Stigmatula** 1: 543)
 - (b) Paraphyses lacking
 - x. Spores long-caudate
 - (x) Spores caudate at one end only **Urcospora** 1: 448
 - (y) Spores caudate at both ends **Urosporella** 14: 523
 - y. Spores not caudate
 - (x) Asci 1-2-spored
 - m. Perithecia perforate †**Diplospor** 11: 292
(**Geminispora**)
 - n. Perithecia closed, then splitting irregularly at apex **Spolverinia** 17: 577
 - (y) Asci 4-8-spored
 - m. Perithecia lenticular, perforate **Laestadia** 1: 420
 - n. Perithecia globose, papillate **Phomatospora** 1: 432
 - (2) Asci many-spored
 - (a) Perithecia glabrous **Ditopella** 1: 450
 - (b) Perithecia strigose-pilose **Polytrichia** 1: 451
 - b. Perithecia superficial
 - (1) Perithecia smooth, i. e., glabrous
 - (a) Spores stellate **Inzengaea** 9: 610
 - (b) Spores not stellate
 - x. Perithecia on a dark crustose subicle **Pilgeriella** 16: 464
 - y. Perithecia not on a subicle
 - (x) Perithecia surrounded by dark hyphae at base **Guignardiella** 16: 465
 - (y) Perithecia without dark hyphae at base **Wallrothiella** 1: 455
(incl. **Zignoina** 2: 219)
 - (2) Perithecia hairy
 - (a) Asci 8-spored **Trichosphaeria** 1: 452
 - (b) Asci 16-spored **Trichosphaerella** 9: 604

II. Perithecia upon or within a stroma or subicle

1. Perithecia beaked **Glomerella** 16: 452, 17: 573
2. Perithecia not beaked
 - a. Perithecia immersed in a subicle **Scortechinia** A 68, 9: 604
 - b. Perithecia in or upon a stroma
 - (1) Stroma radiate, phyllogenous **Trabutia** 1: 449
 - (2) Stroma not radiate, usually caulicole
 - (a) Necks of perithecia wanting, stroma disk-like
Botryosphaeria 1: 456
(incl. **Gibellia** A 406, 9: 608 and
Coutinia 17: 589)
 - (b) Necks of perithecia present, stroma valsiform
Cryptosporella 1: 466
(incl. **Diaporthopsis** 9: 610)

Phaeosporae

1: 214, 9: 481, 11: 278, 14: 489, 16: 427, 17: 593

Spores 1-celled, colored, usually yellowish or brown, ovoid, oblong or fusoid

I. Perithecia separate, at least without a stroma

1. Covered, often erumpent
 - a. Asci 1-spored **Haplosporium** A 40, 9: 495
 - b. Asci 4-8-spored
 - (1) Perithecia covered by the blackened adhering epiderm
Anthostomella 1: 278
 - (2) Perithecia erumpent with a stellate volva
Astrocystis 1: 293
 - c. Asci many-spored
 - (1) Spores smooth **Müllerella** A 40, 9: 495
 - (2) Spores verrucose **Mesnieria** 16: 440
2. Superficial or subsuperficial
 - a. Perithecia long-beaked
 - (1) Spores lunulate; fimicole **Micrascus** A 37, 9: 483
 - (2) Spores globose to elliptic; not fimicole
Ceratostoma 1: 215
 - b. Perithecia not beaked
 - (1) Perithecia submembranous
 - (a) Spores with a mucous sheath or tail; usually fimicole
 - x. Asci 4-8-spored
 - (x) Spores with a hyaline tail or cauda
Sordaria 1: 230
 - (y) Spores with a mucous sheath
 - m. Perithecia sparse **Hypocopa** 1: 240
 - n. Perithecia densely aggregate, almost stroma-like
Coprolepa 1: 248
 - y. Asci many-spored, spores usually caudate
Philocopa 1: 249
 - (b) Spores without mucous sheath or tail
 - x. Perithecia with simple setae, asci persistent
Helminthosphaeria 1: 230

- y. Perithecia with branched, hooked or spiral setae; asci diffluent
(x) Spores subglobose to elliptic

Chaetomium 1: 220

- (y) Spores triangular

Bommerella A 38, 9: 486

- (2) Perithecia typically carbonous

Rosellinia 1: 252

(incl. **Pleosporopsis** 14: 501 and

Tyimpanopsis 11: 283

- (3) Perithecia coriaceous, firm, ascending-elongate

Bombardia 1: 277

II. Perithecia in a stroma

1. Stroma immersed, somewhat woody; perithecia membranous

Anthostoma 1: 293

2. Stroma superficial, carbonous or leathery; perithecia carbonous

- a. Stroma terete, fruticose or filiform

- (1) Stroma fimicole

†**Pedisordaria** 14: 494

(**Podosordaria**)

- (2) Stroma not fimicole

- (a) Stroma with a single perithecium at apex

Capnodiella 17: 621

- (b) Stroma containing many perithecia

- x. Perithecia immersed laterally

- (x) Stroma fruticose, clavate or filiform

Xylaria 1: 309

(incl. **Kretschmaria** 9: 565)

- (y) Stroma disk-like or cupulate above

Xylariodiscus 16: 449

- y. Perithecia immersed vertically

- (x) Perithecia immersed annulately about the truncate apex

Camillea 1: 346

- (y) Perithecia crowded beneath an operculate disk

Henningsinia 16: 450

- b. Stroma effuse, globose or cupulate, adnate or substipitate

- (1) Conidia superficial on the young stroma

- (a) Stroma usually fimicole

Poronia 1: 348

- (b) Stroma not fimicole

- x. Stroma concentrically zonate

Daldinia 1: 393

- y. Stroma not concentrically zonate

- (x) Stroma repand-pulvinate, somewhat hollow

Ustilina 1: 351

- (y) Stroma solid

- m. Stroma subglobose, hemispheric or obpiriform

- (m) Stroma not modified with squarrose papery membranes

Penzigia 9: 567

- (n) Stroma modified by squarrose papery membranes

Squamotubera 17: 620

- n. Stroma effuse

- (m) Perithecia immersed, necks rather long

Bolinia 1: 352

(n) Perithecia innate-prominent, necks lacking

Hypoxyllum 1: 352

(2) Conidia arising beneath the upper layer of the disk-like or cupulate stroma

(a) Perithecia flask-shaped

Nummularia 1: 395

(b) Perithecia long-cylindric

Solenoplea 17: 619

Hyalodidymae

1: 475, 9: 611, 11: 295, 14: 525, 16: 468, 17: 635

Spores 1-septate (2-celled), hyaline or subhyaline, ovoid, oblong or fusoid

I. Perithecia separate

1. Perithecia covered or nearly so

a. Perithecia beaked, submembranous

(1) Asci 8-spored

Gnomonia 1: 561

(2) Asci many-spored

Rehmiella 9: 676

b. Perithecia not beaked

(1) Asci 8-spored

(a) Perithecia in a phyllogenous pseudostroma

Hypoaspilina 2: 190

(b) Perithecia not in a phyllogenous pseudostroma

x. Paraphyses lacking

Sphaerella 1: 476

(incl. *Lizoniella* 17: 661)

y. Paraphyses present

(x) Spores surrounded with mucus

Massarinula 14: 536

(y) Spores not surrounded with mucus

m. Spores septate near the base

Apiospora 1: 539

(incl. *Stigmathea* 1: 541)

n. Spores septate near the middle

(m) Perithecia smooth

Didymella 1: 545

(incl. *Stigmathea* 1: 545)

(n) Perithecia long-hairy

Arcangelia 9: 696

(2) Asci 16-24-spored

(a) Asci 16-spored

Mycosphaerella 9: 659

(b) Asci 24-spored

Hariotia 9: 672

2. Perithecia superficial or nearly so

a. Perithecia beaked

(1) Spores expelled in a mucous mass

Spumatoria 16: 1134

(2) Spores not expelled in a mucous mass

Lentomita 1: 584

b. Perithecia not beaked

(1) Perithecia smooth

(a) Asci 8-spored

x. Paraphyses lacking

(x) Perithecia borne in lichen thalli

Pharcidia 9: 676, 17: 635

(incl. *Epicymatia* 1: 570)

(y) Perithecia not in lichen thalli

Bertia 1: 581

- y. Paraphyses present
 - (x) Spores with a mucous layer produced into a spatulate ring
Pteridiospora 14: 539
 - (y) Spores without a mucous layer
 - m. Spores ellipsoid to fusoid **Melanopsamma 1: 575**
 - n. Spores botuliform **Thaxteria 9: 687**
 - (b) Asci 16-spored **Pseudolizonia 9: 682**
 - (2) Perithecia with hairs or bristles
 - (a) Paraphyses lacking
 - x. Perithecia lichenicole **Echinothecium 16: 484**
 - y. Perithecia typically on leaves, rarely on stems
Venturia 1: 586
 - (b) Paraphyses present **Eriosphaeria 1: 597**
- II. Perithecia cespitose **Othiella 1: 739, 17: 662**
- III. Perithecia in, or rarely upon, a stroma
- 1. Stroma scanty
 - a. Perithecia smooth **Gibbera 1: 599**
 - b. Perithecia setose **Cacosphaeria 9: 699**
 - 2. Stroma well-developed
 - a. Stroma white or colored
 - (1) Stroma white and soft **Melchiora 14: 538**
 - (2) Stroma bright yellow **Endothia 1: 601**
 - b. Stroma black, rarely yellowish
 - (1) Perithecia botryose, erumpent, superficial
Myrmaecium 1: 600
 - (2) Perithecia immersed
 - (a) Spores septate near the base **Aplacodina 16: 485**
 - (b) Spores septate near the middle
 - x. Stroma valsa-like
 - (x) Conidial stage **Melanconium**
Melanconis 1: 602
 - (y) Pycnidial stage **Rabenhorstia**
Hercospora 1: 605
 - (z) Pycnidial stage **Phoma**
Diaporthe 1: 606
 - y. Stroma eutype-like or diatrype-like
Euporthe 1: 631, 1: 662

Phaeodidymae

I: 701, 9: 723, II: 312, 14: 551, 16: 498, 17: 675

Spores 1-septate, dark, fuliginous to brown, ovoid, oblong or fusoid

- I. Perithecia separate
 - 1. Perithecia covered
 - a. Paraphyses lacking **Phaeosphaerella 9: 723**
(incl. **Lizonia 1: 574**)
 - b. Paraphyses present
 - (1) Asci 8-spored
 - (a) Spores surrounded by a hyaline sheath
Massariella 1: 716

- (b) Spores without a sheath **Didymosphaeria** 1: 701
 - (2) Asci many-spored **Tichothecium** 17: 676, 9: 723
- 2. Perithecia superficial or immersed at the base
 - a. Subicle present
 - (1) Perithecia beaked
 - (a) Paraphyses lacking **Rhynchomeliola** A. 127, 9: 751
 - (b) Paraphyses present **Gibellina** A: 413, 9: 740, 11: 317
 - (2) Perithecia not beaked
 - (a) Perithecia glabrous **Neopeckia** A: 26, 9: 749
 - (b) Perithecia setose **†Dimerosporis** 17: 686
(**Dimerosporiopsis**)
 - b. Subicle lacking
 - (1) Perithecia beaked
 - (a) Asci paraphysate **Rhynchostoma** 1: 730
 - (b) Asci not paraphysate **†Dysrhynchis** 17: 689
(**Henningsomyces**)
 - (2) Perithecia not beaked
 - (a) Perithecia glabrous
 - x. Perithecia carbonous **Amphisphaeria** 1: 718
 - y. Perithecia membranous or submembranous
 - (x) Asci 8-spored
 - m. Perithecia globose, fimicole **Delitschia** 1: 732
 - n. Perithecia cupulate, not fimicole
 - Gaillardiiella** 14: 559
 - Delitschiella** 17: 688
 - (y) Asci many-spored **Protoventuria** A: 113, 9: 741
- II. Perithecia cespitose or forming a crust, not stromate
 - 1. Perithecia forming an effuse crust **Parodiella** 1: 717
 - 2. Perithecia in groups
 - a. Perithecia foliicole **Pseudothia** 16: 507
 - b. Perithecia lichenicole **Sorothelia** A: 122, 9: 728
 - c. Perithecia ramicole **Othia** 1: 735
- III. Perithecia in a stroma
 - 1. Spore with a mucous covering **Massariovalsa** 9: 755
 - 2. Spore without a mucous covering
 - a. Stroma erect, subterete
 - Xylobotryum** 11: 319, 14: 20
 - (**Trachyxylaria** 16: 510, **Xyloceras** 17: 690)
 - b. Stroma flat, round or cushion-like, immersed or emerging
 - (1) Paraphyses lacking
 - (a) Stroma bearing conidia of Melanconium **Melanconiella** 1: 740
 - (b) Stroma without conidia **Camarops** 1: 753
 - (2) Paraphyses present
 - (a) Stroma phyllogenuous; perithecia superficial **Licopolia** 16: 508

(b) Stroma not phyllogenous

x. Perithecia valsoid

Valsaria 1: 741

y. Perithecia eutypoid

Endoxylina 11: 318

Hyalophragmiae

2: 152, 9: 824, 11: 332, 14: 581, 16: 528, 17: 692

Spores 2-several-septate, hyaline, oblong to cylindric

I. Perithecia separate

1. Perithecia covered or erumpent

a. Perithecia beaked

(1) Perithecia xylogenous, carbonous

Ceratosphaeria 2: 227

(2) Perithecia phyllogenous, submembranous

(a) Spores separating into halves **Cryptoderis** 2: 229

(b) Spores not separating into halves

Gnomoniopsis 17: 716

b. Perithecia not beaked

(1) Spores with a mucous covering **Massarina** 2: 153

(2) Spores without a mucous covering

(a) Perithecia submembranous, pseudostroma lacking

x. Paraphyses lacking

Sphaerulina 2: 186

y. Paraphyses present

(x) Spores muticate

Metasphaeria 2: 156

(incl. **Charrinia** 14: 585)

(y) Spores with a seta or cusp at either end

Ceriosporella 2: 184, 14: 19

(b) Perithecia membranous, in a leafy pseudostroma

Hypospila 2: 189

(c) Perithecia subcarbonous, pseudostroma lacking, spores 20-30-septate

Saccardoella 2: 190

2. Perithecia superficial or subsuperficial

a. Perithecia glabrous

(1) Perithecia stalked, covered with a bright powder

Bombardiastrum 11: 338

(2) Perithecia not stalked, powdery covering lacking

(a) Spores 2-septate

Melomastia 2: 213

(b) Spores typically 3 or more-septate

x. Perithecia carbonous, black

Zignoella 2: 214

(incl. **Bertiella** 17: 708)

y. Perithecia softish, greenish or reddish

Winterina 14: 589

b. Perithecia hairy or byssiside

(1) Perithecia of one color

(a) Spores chain-like, separating into globose joints

Hormosperma 14: 591

(b) Spores not separating into joints

x. Perithecia carbonous, large

(x) Spores cylindric, elongate

Lasiosphaeria 2: 191

- (y) Spores fusoid, somewhat short
Enchnosphaeria 2: 205
- y. Perithecia submembranous, small
Acanthostigma 2: 207
- z. Perithecia fleshy-coriaceous, hairs fascicled on a central disk
Actiniopsis 16: 543
- (2) Perithecia of two colors, usually reddish at vertex
Herpotrichia 2: 211
- II. Perithecia cespitose, erumpent, superficial, membranous
Baumiella 17: 708
- III. Perithecia in a stroma or on a subicle
1. Perithecia on a subicle; asci many-spored, paraphyses lacking
Sydowia 11: 341
2. Perithecia in a stroma
- a. Stroma lichenicole, white, lanose
Dichosporium 16: 542
- b. Stroma not lichenicole, black
- (1) Stroma immersed
Calospora 2: 231
- (2) Stroma superficial
- (a) Stroma lentiform, adnate to the pycnidium
Melanops 2: 231
- (b) Stroma pulvinate or hemispheric
Holstiella 14: 593
- Phaeophragmiae**
- 2: 1, 9: 759, 11: 319, 14: 561, 16: 510, 17: 718
- Spores 2-several-septate, olive, melleous or fuliginous, oblong to cylindric
- I. Perithecia separate
1. Perithecia covered or erumpent
- a. Spores with a mucous covering
Massaria 2: 2
- b. Spores without a mucous covering
- (1) Perithecia depressed beneath a black cortical clypeus
Clypeosphaeria 2: 90
- (2) Perithecia without a stromatic clypeus
- (a) Spores muticate
- x. Paraphyses lacking
Phaeospora 16: 519
- y. Paraphyses present
- (x) Cells of spore concolorous
- m. Perithecia glabrous
- (m) Perithecia rostrate
Rhynchospaeria 16: 524
- (n) Perithecia not beaked
- r. Spores cylindric, connected in pairs in the ascus
Leptosphaeropsis 9: 770, 11: 321
Leptosphaeria 2: 13
 (incl. **Cladosphaeria 11: 321, Chitonospora 9: 797**)
- s. Spores separate
- n. Perithecia setose or hairy
Pocosphaeria 11: 325
- (y) Cells of spore discolorous
Heptameria 2: 88
 (incl. **Passeriniella 11: 326**)
- (b) Spores caudate or cuspidate
- x. Spores caudate at base
Rebentischia 2: 12
- y. Spores cuspidate at both ends
Ceriospora 14: 19, 2: 184

2. *Perithecia* superficial or subsuperficial
 - a. *Perithecia* glabrous
 - (1) *Phytophilous*
 - (a) Spores finally separating into joints
 - x. Joints 1-celled *Ohleriella* 17: 736
 - y. Joints 2-celled *Ohleria* 2: 96 .
 - (b) Spores not separating into joints
 - x. *Perithecia* smooth or nearly so
 - (x) Spores biconic with a mucous covering
Caryospora 2: 122
 - (y) Spores medium, no mucous covering
 - m. Ostiole narrow *Melanomma* 2: 98
 - n. Ostiole widely open *Trematosphaeria* 2: 115
 - y. *Perithecia* verrucose
 - Stuartella* 2: 123
 - (2) *Fimicole* *Sporormia* 2: 123
 - b. *Perithecia* pilose or byssiside
 - (1) *Perithecia* concolorous
 - (a) Spores cylindric, elongate **Lasiosphaeris* 2: 194
 - (b) Spores fusoid, somewhat short *Chaetosphaeria* 2: 92
 - (2) *Perithecia* discolorous at the vertex
 - *Herpothrix* 2: 211
 - II. *Perithecia* cespitose, erumpent *Gibberidea* 2: 132
 - III. *Perithecia* in a stroma
 1. Stroma lichenicole *†Trematosphaeris* 17: 735
(*Trematosphaeriopsis*)
 2. Stroma not lichenicole
 - a. Asci 1-spored *Titania* 9: 823
 - b. Asci 4-8-spored
 - (1) Stroma valsa-like, innate
 - (a) Asci 4-spored *Aglaospora* 2: 133
 - (b) Asci 6-8-spored
 - x. Acervuli covered with a reddish or yellowish bran
Thyridaria 2: 140
 - y. Acervuli not covered with a bran
Pseudovalsa 2: 135
 - (2) Stroma eutype-like, i. e., woody, effuse
 - (a) Paraphyses lacking *Cryptosphaerina* 16: 521
 - (b) Paraphyses present *Kalmusia* 2: 142
 - (3) Stroma pulvinate, emerging *Melogramma* 2: 144
- Hyalodictyae**
2: 238, 11: 349, 9: 872, 14: 611, 16: 554, 17: 743
- Spores transversally and longitudinally septate, usually muriform,
hyaline, oblong to fusoid.
- I. *Perithecia* separate
 1. *Perithecia* covered or erumpent
 - a. Asci 8-spored
 - (1) Paraphyses lacking

- i. Wall of perithecium double
Scleroplea 16: 548
 - (s) Spores compressed, flattened
 - h. Perithecia smooth **Clathrospora 9: 894**
 - i. Perithecia hairy ***Comoclathris**
 - (n) Spores appendaged at both ends
Delacourea 2: 288
 - n. Perithecia setose, especially about ostiole
Pyrenophora 2: 277
 - (2) Perithecia in a phyllogenous pseudostroma
Isothea 2: 290
 - 2. Perithecia superficial
 - a. Phytogenous
 - (1) Perithecia soft, light colored **Winteria 14: 589**
 - (2) Perithecia carbonous, black
 - (a) Perithecia corrugate-tuberculate
Crotonocarpia 2: 306
 - (b) Perithecia not corrugate
 - x. Perithecia glabrous **Tichospora 2: 290**
 - y. Perithecia hairy **Pleosphaeria 2: 304**
 - b. Fimicole; each spore of 3 10-celled chains
Pleophragmia 2: 307
- II. Perithecia cespitose **Cucurbitaria 2: 307**
- III. Perithecia in a stroma
- 1. Spores with a mucous layer **Montagnula 14: 603**
 - 2. Spores without a mucous layer
 - a. Stroma effuse, eutypeous **Thyridium 2: 323**
 - b. Stroma valsous **Fenestella 2: 325**
- Scolecosporae**
- 2: 337, 9: 923, 11: 351, 14: 613, 16: 557, 17: 767
- Spores linear or filiform, continuous or septate, hyaline or yellowish.
- I. Perithecia separate
- 1. Perithecia covered or erumpent
 - a. Perithecia covered by a phyllogenous clypeus
Linospora 2: 354
 - b. Perithecia not covered by a clypeus
 - (1) Perithecia beaked **Ophiognomonina 17: 776**
 - (2) Perithecia not beaked
 - (a) Perithecia glabrous
 - x. Spores muticate
 - (x) Spores in a hyaline sheath **Ophiomassaria 11: 353**
 - (y) Spores not in a hyaline sheath
 - m. Perithecia globose to conoid
Ophiobolus 2: 337
 - n. Perithecia cylindric, truncate
Cylindrina A: 421, 9: 937
 - y. Spores awned at each end

- (x) Perithecia very large, disk-form, corticole
Therrya 2: 358
- (y) Perithecia small, globose, on grasses and palms
Dilophia 2: 357
- (b) Perithecia hairy
Ophiochaete 11: 353
- 2. Perithecia superficial or immersed at base
 - a. Perithecia beaked
Ophiochaete 11: 352
 - b. Perithecia not beaked
Bovilla 2: 360
 - (1) Perithecia fimicole
 - (2) Perithecia not fimicole
 - (a) Perithecia glabrous
 - x. Perithecia globose
 - (x) Perithecia immersed at base
Acerbia 11: 353, 14: 619
 - (y) Perithecia wholly superficial
Leptospora 14: 619
 - y. Perithecia elongate cylindric; ostiole sulcate
Bactrosphaeria 14: 617
 - Acerbiella 17: 768**
 - (b) Perithecia hairy
- II. Perithecia in a stroma
 - 1. Stroma superficial
 - a. Perithecia in an effuse definite stroma
Maurya 14: 620
 - b. Perithecia densely heaped in a thin vanishing stroma
Pseudomeliola 9: 938
 - 2. Stroma immersed or erumpent
 - a. Stroma erumpent, yellow within
Sillia 1: 361
 - b. Stroma immersed, valvous
 - (1) Necks of perithecia short, scarcely converging
Vialaea 14: 619
 - (2) Necks long, converging into a disk
Cryptospora 2: 361

Family 19. VERRUCARIACEAE

Zahlbruckner 51

Mycelium parasitic on bluegreen or yellow green algae, and forming a more or less distinct crustose, foliose or fruticose thallus, the latter usually superficial but sometimes below the surface; perithecia distinct, single or cespitose or united in a stroma, usually globose and ostiolate, membranous, coriaceous or carbonous; asci 1-many-spored; spores various.

- I. Perithecia separate, at least not in a stroma (Cfr. Lichinae, page 74.)
 - 1. Algae bluegreen, Nostoc, Scytonema, Sirospira, or Calothrix
Subfamily Pyrenidiaceae 76
 - a. Asci 4-8-spored
 - (1) Asci 4-spored; spores 3-septate
Pyrenidium 77
 - (2) Asci 6-8-spored
 - (a) Spores spheric, 1-celled: algae Calothrix
Calothricopsis 165
 - (b) Spores fusiform, 1-septate

- x. Algae Sirosiphon or Scytonema
 - Eolichen 76
- y. Algae Nostoc
 - Pyrenocollema 169
- (c) Spores filiform, continuous
 - Hassea 76
- b. Asci many-spored; spores 1-celled
 - Placothelium 77
- 2. Algae yellow green, Pleurococcus, Palmella, Chroolepus, etc.
 - a. Thallus crustose or gelatinous
 - (1) Thallus gelatinous, hyphae loose
 - Epigloea 53
 - (2) Thallus crustose, not gelatinous, hyphae compact
 - (a) Algae Cystococcus, in sheathed colonies
 - Subfamily Moriola 52
 - x. Thallus without pseudoparenchyma
 - Moriola 52
 - y. Thallus with pseudoparenchyma
 - (x) Asci 8-spored
 - m. Spores dark, 1-septate
 - *Dimerisma 52
 - n. Spores dark, 4-8-septate
 - *Phaeomeris 52
 - o. Spores hyaline, 2-4-septate
 - Spheconisca 52
 - (y) Asci many-spored; spores hyaline, 1-celled
 - *Pleophalis 52
- (b) Algae Pleurococcus or Palmella
 - Subfamily Verrucariae 53
- x. Paraphyses lacking, or soon disappearing
 - (x) Asci 1-8-spored
 - m. Algae present within the perithecium; spores muriform
 - (m) Spores hyaline
 - *Phalostauris 57
 - (n) Spores dark
 - Staurothele 56
 - n. Algae lacking in perithecium
 - (m) Spores 1-celled
 - r. Spores globose to elliptic
 - (r) Perithecia more or less superficial
 - h. Spores hyaline
 - Verrucaria 54
 - i. Spores dark
 - *Phaeosporis 55
 - (s) Perithecia immersed
 - *Lithoecis 55
 - s. Spores vermiform, clavate at each end
 - Saccopyrenia 54
 - (n) Spores 2-4-celled, hyaline
 - r. Spores 2-celled
 - Thelidium 56
 - s. Spores 4-celled
 - *Phragmothele 56
 - (o) Spores muriform
 - Polyblastia 56
 - (y) Asci many-spored
 - Trimmatothele 56
- y. Paraphyses persistent
 - (x) Algae present in the perithecium
 - Thelenidia 57
 - (y) Hymenial algae lacking
 - m. Perithecia with normal ostiole

- (m) Spores 1-celled
 - r. Spores hyaline **Thrombium 57**
 - s. Spores dark ***Phaeothrombis 57**
- (n) Spores septate
 - r. Spores elliptic, 3-few-septate **Geisleria 57**
 - s. Spores muriform
 - (r) Spores hyaline **Microglæna 57**
 - (s) Spores dark ***Phaeoglæna 57**
 - t. Spores needle-shaped, many-celled **Gongylia 57**
- n. Ostiole margined by a broad disk
 - (m) Spores transeptate **Aspidopyrenium 58**
 - (n) Spores muriform **Aspidothelium 58**
- (c) Algae Chroolepus
 - x. Perithecia upright, with vertical ostiole **Subfamily Pyrenulæ 62**
- (x) Paraphyses free, simple
 - m. Perithecia smooth
 - (m) Spores 1-celled, colorless **Coccotrema 66**
 - (n) Spores septate
 - r. Asci 4-8-spored
 - (r) Asci persistent
 - h. Spores transeptate
 - (h) Spores hyaline
 - + Spores 1-septate
 - (+) Spore cells separating ***Dichoporis 66**
 - (—) Spore cells not separating ***Diporina 66**
 - Spores 2-many-septate **Porina 66**
 - (i) Spores dark
 - + Spores 1-septate ***Dipyrenis 68**
 - Spores several-septate **Pyrenula 67**
(incl. **Blastodesmia 67**)
 - i. Spores muriform
 - (h) Spores hyaline **Clathroporina 67**
 - (i) Spores brown **Anthracotheicum 68**
 - (s) Asci evanescent; spores acicular, clear **Belonia 67**
 - s. Asci many-spored; spores septate, clear
 - (r) Spores 1-celled ***Holothelis 67**
 - (s) Spores septate
 - h. Spores 1-septate ***Dithelopsis 67**

- i. Spores 2-many-septate
 - Thelopsis** 67
- n. Perithecia with stiff fascicled hairs
 - Stereochlamys** 68
- (y) Paraphyses lacking, or branched and united
 - m. Ostiole round or dot-like
 - (m) Spores hyaline
 - r. Spores 1-septate ***Pyrenyllum** 64
 - s. Spores 2-many-septate
 - (r) Spores oval to oblong
 - Arthropyrenea** 64
 - (incl. **Pseudopyrenula** 65)
 - (s) Spores acicular to filiform
 - Leptorhaphis** 65
 - t. Spores muriform **Polyblastiopsis** 65
 - (n) Spores brown
 - r. Spores 1-septate **Microthelia** 62
 - s. Spores 2-several-septate ***Polythelis** 64
 - n. Ostiole radiate, torn or lobed
 - Asteroporum** 62
- y. Perithecia oblique or horizontal with oblique or lateral ostiole
 - Subfamily Paratheliae** 71
- (x) Spores transeptate
 - m. Spores hyaline
 - (m) Spores 1-septate ***Ditremis** 71
 - (n) Spores several-septate, oblong
 - Pleurotrema** 71
 - (incl. **Plagiotrema** 72)
 - (o) Spores filiform, many-celled
 - *Trichotrema** 71
 - n. Spores brown
 - Parathelium** 72
 - (y) Spores muriform
 - m. Spores hyaline **Campylothelium** 72
 - n. Spores brown **Pleurothelium** 72
- (d) Algae **Phyllactidium** or **Cephaleurus**
 - Subfamily Strigulae** 74
- x. Perithecia smooth
 - (x) Paraphyses simple, free
 - m. Spores transeptate
 - (m) Spores 1-septate ***Phylloporis** 75
 - (n) Spores several-septate
 - r. Thallus uniform **Phylloporina** 75
 - s. Thallus orbicular, lobed at edge
 - Strigula** 76
 - Phyllobathelium** 75
 - n. Spores muriform
 - (y) Paraphyses branched and united
 - m. Spores 1-celled, dark **Haplopyrenula** 74
 - n. Spores 2-4-celled, brown **Microtheliopsis** 75

- y. Perithecia with fascicled nearly horizontal hairs at apex
Trichothelium 75
- b. Thallus foliose or scaly **Subfamily Dermatocarpae 58**
 - (1) Algae Palmella
 - (a) Hymenial algae lacking
 - x. Paraphyses lacking, or fused into a mass
 - (x) Paraphyses lacking; thallus without cortex
Normandina 59
 - (y) Paraphyses fused; thallus corticate
 - m. Spores 1-celled, colorless **Dermatocarpum 60**
 - n. Spores septate
 - (m) Spores colorless **Placidiopsis 60**
 - (n) Spores brown **Heterocarpum 60**
 - y. Paraphyses persistent
 - (x) Spores 1-celled, brown **Anapyrenium 59**
 - (y) Spores muriform, colorless **Psoroglaena 59**
 - (b) Hymenial algae present **Endocarpum 61**
 - (2) Algae Chroolepus; spores colorless, 1-celled
Lepolichen 69
Mastodia 241
 - (3) Algae Prasiola
 - c. Thallus fruticose, branched, with Pleurococcus; spores muriform, brown
Pyrenothamnia 61
- II. Perithecia in a stroma (Cfr. Pertusariae, page 79.)
 - 1. Perithecia upright, with individual pores **Subfamily Trypetheliae 69**
 - a. Spores colorless
 - (1) Spores transeptate
 - (a) Spores oval to fusiform **Trypethelium 70**
 - (b) Spores filiform **Tomasiella 69**
 - (2) Spores muriform **Laurera 71**
 - b. Spores brown
 - (1) Spores transeptate **Melanotheca 70**
 - (2) Spores muriform **Bottaria 71**
 - 2. Perithecia oblique or horizontal, with a common canal or pore **Subfamily Astrotheliae 72**
 - a. Spores transeptate
 - (1) Spores colorless **Astrothelium 73**
(incl. **Lithothelium 73**)
 - (2) Spores brown **Pyrenastrum 73**
 - b. Spores muriform
 - (1) Spores colorless **Heufleria 74**
 - (2) Spores brown **Parmenteria 74**
- III. Perithecia sunken in stroma-like warts; horizontal thallus lacking; asci many-spored; spores 1-celled, clear **Thelocarpum 150**

Family 20. HYPOCREACEAE

2: 447, 9: 941, 11: 354, 14: 621, 16: 559, 17: 777.

Mycelium scanty and immersed or producing a subicle or a stroma; perithecia

globoid, sometimes beaked, fleshy, waxy or waxy-membranous, bright colored, usually reddish, more rarely blue, yellow or whitish, never carbonous, opening by a round pore or ostiole, single, cespitose or composite in a stroma; asci and spores as in Sphaeriaceae.

Allantosporae

17: 778

Spores 1-celled, obtuse, curved-oblong, hyaline or olivascens

One genus

Allantonectria 17: 778

Hyalosporae

2: 447, 9: 941, 11: 354, 14: 621, 16: 559, 17: 778

Spores 1-celled, hyaline

I. Perithecia separate

1. Perithecia covered

a. Asci 8-spored

Hyponectria 2: 455

b. Asci many-spored

Thelocarpum 9: 946

2. Perithecia superficial or nearly so

a. Perithecia beaked; spores ciliate

Eleutheromyces 2: 455

b. Perithecia not beaked

(1) Spores smooth

Nectriella 2: 448

(2) Spores ciliate or spiny

(a) Spores 1-ciliate at each end

Heteronectria 14: 624

(b) Spores spiny, hemispheric

Cleistosoma A: 195, 9: 943

II. Perithecia cespitose

1. Asci 8-spored

Lisiella 9: 945

2. Asci many-spored

Chilonectria 2: 453

III. Perithecia in a subicle or stroma

1. Perithecia in a subicle, i. e., a cobwebby or cottony stroma

a. Paraphyses lacking, fungicole

Peckiella 9: 944

b. Paraphyses numerous, not fungicole

Byssonectria 2: 456

2. Perithecia in a definite stroma

a. Stroma effuse, globose, verruciform or linear

(1) Asci 8-spored

(a) Perithecia circinate, valsiform

Balzania 16: 561

(b) Perithecia not circinate, mostly irregular

x. Spores globose

Battarina 2: 533

y. Spores ovate to oblong

(x) Stroma globose or verruciform

m. Stroma globose, smooth, dark

Pseudotrype 16: 561

n. Stroma verruciform, hairy, red

Selinia 2: 457

(y) Stroma lirelliform, clear

Monographus 2: 457

(z) Stroma effuse, phyllogenous

Polystigma 2: 458

(2) Asci many-spored; phyllogenous

Moelleriella 14: 626

b. Stroma elongate, erect

- (1) Asci 8-spored
 - (a) Stroma capitate, spores smooth **Sphaerostilbella 17: 778**
 - (b) Stroma clavaria-like; spores asperate **Penicilliopsis 9: 945**
- (2) Asci 16-spored; stroma clavate; on insects **Podostroma 11: 355**

Phaeosporae

2: 459, 9: 949, 11: 355, 14: 626, 16: 562, 17: 781

Spores 1-celled, dark

- I. Perithecia separate
 - 1. Perithecia more or less covered **Baculospora 9: 952**
 - 2. Perithecia superficial
 - a. Perithecia not beaked
 - (1) Perithecia smooth
 - (a) Spores globose, verruculose **Neocosmospora 16: 562**
 - (b) Spores oval to elliptic, smooth ***Sphaerodes 2: 460**
 - (2) Perithecia hairy **Erythrocarpum 9: 950**
 - b. Perithecia beaked
 - (1) Asci 8-spored **Melanospora 2: 461**
 - (2) Asci many-spored **Scopinella 9: 953**
- II. Perithecia in a subicle or a stroma
 - 1. Perithecia immersed in a subicle
 - a. Perithecia beaked ***Rhynchomelas 2: 461**
 - b. Perithecia not beaked **Sphaeroderma 2: 459**
 - 2. Perithecia in a stroma
 - a. Spores spheric **Thuemenella 14: 628**
 - b. Spores ovoid
 - (1) Stroma clavate, pendulous **Xylocrea 16: 451**
 - (2) Stroma more or less globose
 - (a) Perithecia in one layer **Entonaema 16: 450**
 - (b) Perithecia in several layers **†Stromne 16: 452**
(Engleromyces)

Hyalodidymae

2: 465, 9: 953, 11: 356, 14: 628, 16: 565, 17: 782.

Spores 2-celled, hyaline

- I. Perithecia separate or cespitose
 - 1. Perithecia immersed; in leaves **Charonectria 2: 466**
 - 2. Perithecia superficial
 - a. Perithecia red, yellow or white
 - (1) Asci of one kind, 8-spored
 - (a) Perithecia beaked **Rhynchonectria 17: 798**
 - (b) Perithecia not beaked
 - x. Spore cells separating **Bresadolella 17: 797**
 - y. Spore cells not separating
 - (x) Perithecia smooth

- m. Perithecia often on a tubercularoid base
Nectria 2: 479
 - n. Perithecia on or with a stilboid base
Sphaerostilbe 2: 511
***Dasyphthora 2: 505**
 - (y) Perithecia hairy
 - (2) Asci of two kinds, 8-spored and many-spored
Aponectria 2: 516
Metanectria 2: 517
 - (3) Asci many-spored
 - b. Perithecia blue or violet
(1) Asci 8-spored
Lisea 2: 517
(2) Asci many-spored
Cyanocephalum 11: 360
 - II. Perithecia in a subicle or stroma
 - 1. Perithecia in a subicle
 - a. Perithecia gibbose-conic, fungicole
Hypomyces 2: 466
 - b. Perithecia scutate-dimidiata, phyllogenous
Puiggariella 2: 478
 - 2. Perithecia in a stroma
 - a. Perithecia adnate to a fruticose stroma
Corallomyces 2: 519
 - b. Perithecia immersed in a clavate, globose, pulvinate or effuse stroma
 - (1) Perithecia long-beaked
Treleasia 14: 640
 - (2) Perithecia not long-beaked
 - (a) Spore divided near base
Lambro 16: 589
 - (b) Spore divided near middle
 - x. Spore cells separating
 - (x) Stroma vertically elongate
Podocrea 17: 799
 - (y) Stroma globose to effuse
 - m. Conidiophore (Stilbum) arising from stroma
Stilbocrea 16: 588
 - n. Conidiophore lacking or not Stilbum
Hypocrea 2: 520
(incl. **Cryphonectria 17: 783**, **My-**
cocitrus 16: 589)
 - y. Spore cells not separating
Hypocreopsis 9: 980
(incl. **Clintoniella 16: 588**)
- Phaeodidymae**
2: 537, 9: 981, 14: 646, 16: 591, 17: 808.
Spores 2-celled, dark
- I. Perithecia separate or cespitose
 - 1. Perithecia immersed
 - a. Perithecia white, ostiole cylindric; on black fungi
Passerinula 2: 537
 - b. Perithecia darkish, ostiole broad, bright; in bark
Spegazzinula 2: 537
 - 2. Perithecia superficial
 - a. Spore cells separating
Neoskofitzia 9: 981
 - b. Spore cells not separating

- (1) Perithecia on or with a stilbum-like base

Calostilbe 16: 591

- (2) Perithecia without stilbum-like base, often with Helminthosporium

Letendreaa 2: 538

(incl. **Phaeonectria** 11: 359)

- II. Perithecia in a stroma

Phaeocreopsis 16: 591

Hyalophragmiae

2: 539, 9: 982, 11: 363, 14: 647, 16: 592, 17: 808

Spores 2-several-septate, hyaline

- I. Perithecia separate or cespitose

1. Perithecia immersed, spores falcate

Cesatiella 2: 557

2. Perithecia superficial

- a. Perithecia red, yellow or white

- (1) Perithecia on or with a stilbum base

Stilbonectria 9: 986

- (2) Perithecia without a stilbum base

- (a) Perithecia astomous

Malmeomyces 16: 592

- (b) Perithecia ostiolate

- x. Spores ciliate at each end

Paranectria 2: 552

(incl. **Debaryella** 17: 809)

- y. Spores muticate

Calonectria 2: 540

- b. Perithecia blue, violet or greenish

- (1) Spores muticate

Gibberella 2: 552

- (2) Spores appendiculate each way

Lecithium 11: 364

- II. Perithecia in a subicle or in a stroma

1. Perithecia in a subicle

Berkelella 9: 989

2. Perithecia in a pulvinate or discoid stroma

Broomella 2: 557

Phaeophragmiae

2: 539, 9: 982, 11: 363, 16: 599

Spores 2-several-septate, dark

- I. Perithecia in a large tuberiform stroma

Peloronectria 16: 599

Hyalodictyae

2: 558, 9: 990, 11: 364, 14: 650, 16: 599, 17: 814

Spores muriform, hyaline

- I. Perithecia separate or cespitose, superficial

1. Perithecia red or yellow to whitish

- a. Perithecia with a stilbum base

Megalonectria 2: 560

- b. Perithecia without a stilbum base

Pleonectria 2: 559

2. Perithecia blue or violet

Pleogibberella 9: 992

- II. Perithecia in a valsoid stroma

Thyronectria 2: 561

Phaeodictyae

2: 558, 9: 990, 11: 364, 16: 600, 17: 815

Spores muriform, dark

I. Perithecia separate or cespitose

1. Perithecia beaked, asci 8-spored **Bivonella** 9: 989
2. Perithecia not beaked, asci many-spored **Feracia** 17: 815

II. Perithecia in a stroma

1. Asci paraphysate
 - a. Stroma conoid, snow-white **Leucocrea** 16: 601
 - b. Stroma tuberiform, rimose **Shiraia** 16: 600
2. Asci not paraphysate
 - a. Stroma pulvinate, disk greenish **Mattirolia** 9: 993
 - b. Stroma subcrustose **Uleomyces** 11: 364

Scolecosporae

2: 562, 9: 993, 11: 365, 14: 651, 17: 815, 16: 601

Hyaloscoleciae

Spores needle-shaped or filiform, hyaline or nearly so

I Perithecia separate or cespitose

1. Perithecia enclosed in a sack **Oomyces** 2: 564
2. Perithecia not in a sack
 - a. Perithecia immersed or erumpent
 - (1) Perithecia many-perforate above **Coscinnaria** 9: 1003
 - (2) Perithecia with a single ostiole **Micronectria** 9: 996
 - b. Perithecia superficial
 - (1) Perithecia globose-conic, papillate, reddish
Ophionectria 2: 563
 - (2) Perithecia vertically oblong, not papillate, white
Tubeufia 14: 652

II. Perithecia in a subicle or in a stroma

1. Perithecia in a subicle or byssoid stroma
 - Torrubiella** 9: 994
(**Helminthascus** 16: 616)
2. Perithecia in a stroma
 - a. Stroma vertical
 - (1) Stroma from a sclerotium or a blackened matrix
Claviceps 2: 564
(incl. **Balansia** 9: 997, **Balansiella** 17: 822)
 - (2) Stroma without sclerotium; on insects or fungi
Cordyceps 2: 566
Dussiella 9: 1004
 - b. Stroma effuse or pulvinate
 - (1) Stroma on a white subicle
Epichloe 2: 578
 - (2) Stroma without a subicle
 - (a) Stroma effuse, encircling culms
Mycomalus 16: 604
 - (b) Stroma pulvinate to globose
 - x. Spore cells separating
 - (x) Perithecia in a definite peripheral zone

(y) Perithecia not arranged in a zone

m. Stroma hard and black **Fleischera 17:819**

n. Stroma fleshy and soft

(m) Stroma fertile over entire surface

Hypocrella 2:579

(n) Stroma fertile above, sterile below

Ascopolyporus 16:605

y. Spore cells not separating

Echinodopsis 17:819

Phaeoscoleciae

Spores filiform, dark

I. Stroma black, perithecia immersed; spores dilabent, brown

Konradia 16:605

Family 21. DOTHIDEACEAE

Mycelium typically producing a stroma, in which the perithecia are more or less completely sunken and reduced to locules; otherwise as in Sphaeriaceae.

Hyalosporae

2:588, A:222, 9:1004, 11:368, 14:663, 16:616, 17:827

Spores 1-celled, hyaline or nearly hyaline, ovoid, oblong or fusoid, rarely globose

I. Asci 8-spored

1. Stroma globose, pulvinate or cup-shaped

a. Stroma cupulate-discoïd, attached at center

Schweinitziella 9:1005

b. Stroma pulvinate or subcylpeate

(1) Stroma pulvinate

(a) Stroma subcoriaceous

Bagnisiella 2:589

(b) Stroma corneous

Kullhemia 2:591

(2) Stroma subcylpeate, often oval to oblong

Mazzantia 2:591

(incl. **Diachora 11:374**)

2. Stroma oblong, linear or effuse

a. Stroma superficial, on flowers

Hyalodothis 11:374

b. Stroma erumpent or superficial

(1) Stroma waxy or fleshy

a. Stroma more or less waxy within, linear, black

Scirrhella 9:1030

b. Stroma fleshy, white

Monographus 2:457

(2) Stroma more or less carbonous, round to effuse

(a) Asci usually shorter than 30 μ

Euryachora 2:625

(b) Asci usually longer than 50 μ

Phyllachora 2:594

II. Asci 3-spored; stroma subglobose, subcorneous

Zimmermanniella 17:827

Phaeosporae

2:626, 9:1031, 11:374, 14:675, 16:625, 17:841

Spores 1-celled, colored, usually yellowish or brown, ovoid, oblong or fusoid

- I. Stroma subhemispheric to effuse; asci 8-spored

Auerswaldia 2: 626

Hyalodidymae

2: 627, 9: 1034, 11: 375, 14: 676, 16: 625, 17: 844

Spores 1-septate (2-celled), hyaline or subhyaline, ovoid, oblong or fusoid

- I. Stroma pulvinate or disciform

1. Stroma pulvinate, erumpent, usually ramicole

- a. Asci 4-8-spored

Plowrightia 2: 635

- b. Asci many-spored

***Pleodothis 11: 376**

2. Stroma disciform, superficial, foliicole

Microcyclus 17: 844

- II. Stroma oblong to linear or effuse

1. Stroma linear

Scirrhia 2: 634

2. Stroma oblong to effuse, sometimes orbicular

- a. Cells of spore very unequal

Munkiella 9: 1034

- b. Cells of spore equal

- (1) Locules immersed in stroma

Dothidella 2: 627

- (2) Locules completely exerted from stroma

Rosenscheldia 9: 1036

Phaeodidymae

2: 639, 9: 1043, 11: 377, 14: 680, 16: 628, 17: 852

Spores 1-septate, dark, fuliginous to brown, ovoid, oblong or fusoid

- I. Stroma superficial, disciform

Maurodothis 17: 856

- II. Stroma erumpent, pulvinate to effuse

1. Stroma usually effuse

Phaeodothis 17: 854

2. Stroma pulvinate

- a. Stroma subcarbonous

Russoella 9: 1044

- b. Stroma subcoriaceous

Dothidea 2: 639

(incl. **Hypoxylopsis 17: 855**)

Hyalophragmiae

2: 646, 9: 1045, 11: 377, 14: 682, 16: 629, 17: 856

Spores 2-several-septate, hyaline, oblong to cylindric

- I. Perithecia or locules exerted from the stroma; spores sometimes colored

Montagnella 2: 646

- II. Perithecia immersed

1. Stroma fleshy or waxy

Dangardiella 14: 683

2. Stroma carbonous

- a. Perithecia disposed in radiate lines

Telimena 16: 631

- b. Perithecia not radiate

Darwiniella 9: 1048

Phaeophragmiae

2: 646, 9: 1045, 11: 377, 14: 682, 16: 629, 17: 857

Spores 2-several-septate, colored, yellowish to brown, oblong to cylindric

- I. Stroma elongate or linear

Rhopographus 2: 647

- II. Stroma subhemispheric

Homostegia 2: 649

Hyalodictyae

8: 847

Spores muriform, hyaline, ovate to oblong

- I. Stroma with a round black receptacle stuffed with locules
Pyrenotheca 8: 847
- II. Stroma disciform or hemispheric
***Discostroma** 11: 379

Phaeodictyae

2: 651, 9: 1051, 11: 378, 14: 684, 16: 632, 17: 858

Spores muriform, dark, ovate to oblong

- I. Stroma disciform or hemispheric
Curreya 2: 651

Scolecosporae

2: 652, 9: 1051, 14: 685, 16: 632, 17: 859

Spores filiform, hyaline, continuous, guttate or septate

- I. Asci 8-spored
 - 1. Spores narrowly filiform, 1-2 μ wide
Ophiodothis 2: 652
 - 2. Spores broadly filiform, 5-8 μ wide
Oxydothis 14: 674
- II. Asci many-spored
Myriogenospora 14: 685

Family 22. MYCOPORACEAE

ZAHLEBRUCKNER 77

Mycelium parasitic on *Palmella* or *Chroolepus*, forming a uniform thallus without a cortex; perithecia reduced to locules in a stroma as in *Dothideaceae*, to which family the genera might well be referred.

- I. Spores transeptate; algae *Chroolepus*
 - 1. Spores 1-septate
 - a. Spores colorless
***Chlorodothis** 78
 - b. Spores brown
***Sciodothis** 78
 - 2. Spores several-septate
 - a. Spores colorless
***Nothostroma** 78
 - b. Spores brown
***Mycoporis** 78
 - 3. Spores needle-shaped
Mycoporellum 78
- II. Spores muriform; algae *Palmella*
Mycoporum 78

Family 23. COCCOIDEACEAE

17: 860 (16: 624)

Stromata with immersed locules, affixed to the matrix by a central stipitiform point, subcarnose when fresh, subcorneous when dry; locules without distinct proper walls.

Hyalosporae

16: 624

Spores 1-celled, hyaline, ellipsoid

- I. Stroma superficial, disciform-pulvinate, subcarbonous
Coccoidea 16: 624
- II. Stroma superficial, cupulate-discoid
Schweinitziella 9: 1005

Phaeosporae

17:860

Spores 1-celled, dark, ovoid

- I. Stroma subcarnose, discoid **Coccodiscus** 17:860

Hyalodidymae

17:860

Spores 1-septate, hyaline, fusoid

- I. Stroma subcarnose or corneous, disciform-pulvinate
Yoshinagaia 17:860

Family 24. MICROTHYRIACEAE

2:658, 9:1053, 11:379, 14:686, 16:633, 17:861

Perithecia separate, or rarely in a stroma, dimidiate, applanate, context usually beautifully radiate, subsuperficial, black, membranous or carbonous, perforate or astomous; asci 4-8-spored, usually short.

Subfamily Microthyriae

Perithecia typically not seated on a subicle

Hyalosporae

2:659, 9:1053, 11:379, 14:686, 16:633, 17:861

• Spores 1-celled, hyaline, ovoid to oblong or fusiform

- I. Spores oblong, curved **Piptostoma** 9:1054

- II. Spores elliptic to fusiform, straight

1. Spores elliptic, short **Myiocoprum** 2:659

2. Spores fusiform, long, sometimes 1-septate
Pemphidium 2:670

Phaeosporae

2:662, 9:1054, 16:634, 17:861

Spores 1-celled, dark, globose to oblong

- I. Spores globose; perithecia on a hyaline subicle
Blasdalea 16:634
- II. Spores oblong; subicle lacking **Vizella** 2:662

Hyalodidymae

2:662, 9:1055, 11:379, 14:687, 16:635, 17:862

Spores 1-septate, hyaline, oblong to fusoid

- I. Asci with paraphyses
1. Perithecia with several ostioles **Polystomella** 9:1063
2. Perithecia astomous **Clypeolum** 2:667
- II. Asci without paraphyses
1. Perithecia smooth
- a. Perithecia more or less mytiliform and confluent
Brefeldiella 9:1063

- b. Perithecia not mytiliform or confluent

Microthyrium 2: 662

2. Perithecia setulose

Chaetothyrium 9: 1061

Phaeodidymae

2: 668, 9: 1064, 11: 381, 14: 689, 16: 639, 17: 865

Spores 1-septate, dark, oblong to fusoid

- I. Perithecia superficial, carbonous, perforate

Seynesia 2: 668

Hyalophragmiae

2: 668, 9: 1068, 11: 381, 14: 690, 16: 642, 17: 868

Spores 2-several-septate, hyaline, fusoid to cylindric

- I. Perithecia separate

1. Perithecia on a fibrous mycelium

Trichopeltis 9: 1068

2. Perithecia without a mycelium

- a. Perithecia smooth

Micropeltis 2: 669

- b. Perithecia margined with rigid appendages

Actiniopsis 17: 871

- II. Perithecia in a dimidiate many-perforate stroma

Gilletiella 14: 691

Phaeophragmiae

2: 668, 9: 1068, 11: 381, 14: 690, 16: 642, 17: 872

Spores 2-several-septate, dark, fusoid, to cylindric

- I. Perithecia membranous, subfibrous; spores conglobate

Phaeoscutella 17: 872

- II. Perithecia carbonous or coriaceous

Scutellum 2: 668

Hyalodictyae

A: 253, 9: 1071, 14: 692, 16: 645

Spores muriform, hyaline, oblong to elliptic

- I. Perithecia membranous, ostiolate

Saccardinula 9: 1071

Phaeodictyae

17: 873

Spores muriform, dark, oblong to elliptic

- I. Perithecia superficial, phyllogenous, subradiate

†**Phaeopeltis** 17: 873

(**Phaeosaccardinula**)

Scolecosporae

9: 1072, 16: 646, 17: 873

Spores acicular, hyaline or colored, continuous or septate

- I. Spores separating into cells

Scolecopeltis 9: 1072

- II. Spores not separating

Ophiopeltis 17: 873

Subfamily Asterinae

14: 692, 16: 646, 17: 875

Perithecia typically seated upon an effuse radiate black subicle

Hyalosporae

14: 692, 16: 646

- I. Spores hyaline, one-celled

Asterula 1: 47, 14: 692**Phaeosporae**

14: 693

- I. Spores dark, one celled

Asteronia 1: 47, 14: 693**Hyalodidymae**

14: 693, 16: 646, 17: 882

- I. Spores hyaline, 1-septate

Asterella 1: 42, 14: 698**Phaeodidymae**

14: 693, 16: 646, 17: 875

- I. Spores dark, 1-septate

Asterina 1: 39, 14: 693
(incl. *Trichothyrium* 9: 1062)**Hyalophragmiae**

14: 699, 16: 650, 17: 884

- I. Spores hyaline, several-septate

Asteridium 1: 49, 14: 699**Phaeophragmiae**

14: 699, 17: 885

- I. Spores dark, several-septate

Asteridiella 14: 701**Family 25. LOPHIOSTOMATACEAE**

2: 672, 9: 1074, 11: 382, 14: 702, 16: 650, 17: 886

Perithecia simple, separate, at first covered, then subsuperficial or insculptate, carbonous, rarely submembranous, black, with a very narrowly rimose, broad and compressed ostiole; asci paraphysate, usually 8-spored; matrix often blackened giving the appearance of a stroma.

Hyalosporae

(Not represented)

Phaeosporae

2: 673, 17: 886

- I. Spores 1-celled, dark

Lophiella 2: 673**Hyalodidymae**

2: 675, 9: 1075, 11: 383, 14: 702, 17: 886

Spores 1-septate, hyaline, oblong to fusoid

- I. Perithecia smooth

Lophiosphaera 2: 675

- II. Perithecia hairy, with wool at base

Lophiotricha 9: 1082**Phaeodidymae**

2: 673, 9: 1074, 11: 382, 14: 702, 16: 650, 17: 887

- I. Spores 1-septate, dark

Schizostoma 2: 673

Hyalophragmiae

2: 678, 9: 1076, 14: 703, 16: 651, 17: 887

- I. Spores hyaline, several-septate **Lophiotrema 2: 678**

Phaeophragmiae

2: 689, 9: 1083, 11: 383, 14: 704, 16: 651, 17: 887

Spores dark, several-septate

- I. Spores caudate **Brigantiella 17: 889**
 II. Spores not caudate **Lophiostoma 2: 689**

Hyalodictyae

9: 1093

- I. Spores hyaline or nearly so, muriform **Lophidiopsis 9: 1093**

Phaeodictyae

2: 710, 9: 1091, 11: 384, 14: 706, 16: 653, 17: 889

- I. Spores dark, muriform **Platystomum 17: 889**
 (**Lophidium 2: 710**)

Scolecosporae

2: 717, 9: 1094

- i. Spores filiform, hyaline or dilutely colored **Lophionema 2: 717**

Family 26. CORYNELIACEAE

9: 1073, 11: 385, 16: 650

Perithecia separate or in a stroma, coriaceous, black, lageniform, with an elongated ostiole, perforate at the apex and then broadly expanded and infundibuliform.

Phaeosporae

9: 1073, 16: 650

- I. Spores dark, 1-celled, spherical **Corynelia 9: 1073**

Phaeophragmiae

11: 385

- I. Spores dark, 3-several-septate **Coryneliella 11: 385**

Phaeodictyae

9: 1073

- I. Spores black, stellate, cells radiating **Tripodora 9: 1073**

Order 9. HYSTERIALES

Perithecia oblong to linear, rarely round, carbonous or membranous, rarely coriaceous, ostiole a cleft or slit; mycelium often forming a thallus with algae.

Family 27. HEMIHYSTERIACEAE

9: 1094, 11: 385, 14: 707, 16: 653, 17: 892

Perithecia simple or aggregated into a stroma, dimidiate-scutate, subicle lacking,

or more or less developed, ostiole hysterium-like; asci 8-spored, spores usually 2-celled, dark.

Phaeosporae

14: 707

- I. Spores dark, 1-celled; subicle lacking **Cyclostomella** 14: 707

Phaeodidymae

9: 1094, 11: 385, 14: 708, 16: 653, 17: 892

Spores dark, 1-septate, elliptic to fusoid

- I. Perithecia on a subicle; stroma lacking **Morenoella** 9: 1094

II. Perithecia in a stroma

1. Asci with paraphyses **Parmularia** 14: 708
(**Schneepia** 9: 1097)
2. Asci without paraphyses **Hysterostomella** 9: 1098

Hyalophragmiae

17: 892

- I. Spores hyaline, 3-several-septate **Parmulariella** 17: 892

Family 28. HYSTERIACEAE

2: 721, 9: 1100, 11: 385, 14: 710, 16: 657, 17: 893

Perithecia simple or very rarely in a stroma, erumpent-superficial, horizontally, rarely vertically oblong or linear, membranous, coriaceous or carbonous, rarely carnosule at first, usually black, opening along the whole surface by a somewhat narrow cleft; asci usually paraphysate, 4-8-spored, rarely many-spored.

Hyalosporae

2: 723, 9: 1100, 11: 385, 14: 710, 16: 657, 17: 893

Spores 1-celled, hyaline, globose to fusoid

- I. Asci 4-spored; spores covered with mucus **Hypodermella** 11: 385

II. Asci 8-spored

1. Perithecia single or at least not coalescing
Schizothyrium 2: 723
(**Henriquesia** 2: 726)
2. Perithecia coalescing in stellate groups of 4-6
Delpinoella 16: 658

Phaeosporae

2: 727, 9: 1100, 14: 710

Spores 1-celled, dark, globose to ovoid

I. Asci 8-spored

1. Perithecia separate; asci paraphysate **Farlowiella** 2: 727, 9: 1100
2. Perithecia stromatic at base; asci aparaphysate **Erikssonia** 14: 710

II. Asci 10-12-spored

Lembosiella 9: 1101

Hyalodidymae

2: 727, 9: 1101, 11: 386, 14: 711, 16: 659, 17: 895

Spores 1-septate, hyaline, ovoid to fusoid

- I. Perithecia membranous
 - 1. Perithecia separate, minute **Aulographum 2: 727**
 - 2. Perithecia in a dimidiate stroma **Cycloschizum 17: 896**
- II. Perithecia carbonous
 - 1. Perithecia separate
 - a. Perithecia simple or scarcely branched
 - (1) Asci 8-spored **Glonium 2: 731**
 - (2) Asci many-spored ***Pleoglonis 9: 1103**
 - b. Perithecia radiately branched, or stellate **Actidium 2: 738**
 - 2. Perithecia connected in orbicular sori **Synglonium 14: 711**
- III. Perithecia at first somewhat fleshy, reddish or yellow **Angelinia 2: 739**

Phaeodidymae

2: 740, 9: 1103, 11: 387, 14: 711, 16: 659, 17: 897

Spores 1-septate, dark, ovoid to oblong

- I. Perithecia on a fibrillose-radiate subicle **Lembosia 2: 741**
- II. Perithecia without a subicle
 - 1. Perithecia coriaceous **Tryblidium 2: 740**
 - 2. Perithecia carbonous
 - a. Perithecia linear; cleft very narrow, straight **Bulliardiella 17: 902**
 - b. Perithecia scutellate; cleft subcircular **Dielsiella 17: 902**

Hyalophragmiae

2: 765, 9: 1112, 11: 388, 14: 715, 16: 664, 17: 903

Spores several-septate, hyaline, oblong to cylindric

- I. Perithecia saprogenous
 - 1. Perithecia carbonous, cleft narrow **Glioniella 2: 765**
 - 2. Perithecia subcoriaceous, cleft wide **Pseudographis 2: 769**
- II. Perithecia biogenous, gregarious in spots
 - 1. Perithecia corticole **Dichaena 2: 771**
 - 2. Perithecia foliicole
 - a. Perithecia merely gregarious **Phragmographium 17: 906**
 - b. Perithecia radiately disposed **Aldona 16: 667**

Phaeophragmiae

2: 743, 9: 1108, 11: 387, 14: 715, 16: 664, 17: 907

Spores several-septate, dark, oblong to cylindric

- I. Edges of cleft somewhat obtuse, then more or less distant
 - 1. Asci 4-8-spored
 - a. Perithecia transversely densely and coarsely sulcate **Rhytidhysterium 2: 759**

b. Perithecia smooth

(1) Perithecia covered by the epidermis

Hypodermopsis 17: 908

(2) Perithecia erumpent or superficial

(a) Perithecia carbonous

Hysterium 2: 743

(b) Perithecia coriaceous

Tryblidiella 2: 757

2. Asci many-spored, perithecia subcoriaceous

Baggea 2: 760

II. Edges of cleft very thin, closely connivent

1. Asci 4-spored; perithecia subcarbonous, striate

Ostreium 2: 765

2. Asci 8-spored; perithecia somewhat membranous, fragile

Mytilidium 2: 760**Hyalodictyae**

2: 772, 9: 1116, 11: 389, 14: 717, 16: 668, 17: 909

Spores muriform, hyaline, ovoid to oblong

I. Perithecia separate

1. Perithecia carbonous, erumpent; spores without mucus

Gloniopsis 2: 772

2. Perithecia membranous, innate; spores with mucus sheath

Hysteropsis 9: 1118

II. Perithecia in a lenticular, radiate stroma

Mendogia 16: 669**Phaeodictyae**

2: 776, 9: 1119, 11: 389, 14: 717, 16: 668, 17: 912

Spores muriform, dark, ovoid to oblong

I. Perithecia carbonous or corneo-carbonous, firm

Hysterographium 2: 776

II. Perithecia membranous, thin

Graphyllum 16: 1145, 17: 913**Scolecosporae**

2: 784, 9: 1123, 11: 389, 14: 719, 16: 669, 17: 913

Spores bacillar to filiform, hyaline or dark

I. Spores 2-5 times shorter than the asci; perithecia membranous

Hypoderma 2: 784

II. Spores filiform, nearly as long as the asci

1. Perithecia horizontally elongate, rarely ampulliform

a. Perithecia elongate

(1) Perithecia membranous, applanate

Lophodermium 2: 791

(2) Perithecia subcarbonous, conchiform

Lophium 2: 799

(3) Perithecia subcoriaceous, depressed

(a) Perithecia subcorneous

Sporomega 2: 801

(b) Perithecia subcarnose

Colpoma 2: 803

b. Perithecia subspheroid or ampulliform

(1) Perithecia depressed spheroid, cleft longitudinal

Ostropa 2: 804

- (2) Perithecia horizontally ampulliform, ostiole roundish
Robergea 2: 806
2. Perithecia vertically elongate, cylindric; cleft obsolete
 a. Spores breaking apart into cells **Microstelium 16: 672**
 b. Spores not breaking apart **Acrospermum 2: 807**
(Schizacrospermum 16: 672)

Family 29. GRAPHIDACEAE

ZAHLEBRUCKNER 87

Mycelium parasitic on yellow green algae, forming a crustose, foliose or fruticose thallus, the latter often immersed, or thallus lacking, and parasitic on lichens or on bark; perithecia single or cespitose or united in a stroma, typically oblong to elongate with a cleft-like opening, more rarely disk-shaped and with an irregular often stellate opening, more or less carbonous.

I. Perithecia separate

1. Thallus lacking, parasitic on lichens or on bark
Subfamily Arthoniae 89, R. 414
- a. Parasitic on lichens
 1. Spores 1-celled **Phacopsis R. 419**
 2. Spores 2-celled **Conida R. 420**
 3. Spores 4-6-celled **Celidium R. 425**
- b. On bark
 1. Spores 2-celled **Lecideopsis R. 432**
 2. Spores 2-several-septate **Arthonia R. 435**
 3. Spores muriform **Arthothelium R. 438**
2. Thallus present, crustose, or uniform
 a. Perithecia without an exciple, i. e., not margined
Subfamily Arthoniae 89
- (1) Algae *Palmella* or *Protococcus*; spores colorless
 (a) Spores 1-septate **Allarthonia 91**
 (b) Spores several-septate ***Plearthonis 91**
 (c) Spores muriform **Allarthothelium 241**
- (2) Algae *Chroolepus*
 (a) Spores transeptate
 x. Spores colorless
 (x) Spores 1-septate ***Diarthonis 91**
 (y) Spores 2-several-septate **Arthonia 89**
 y. Spores brown **Gymnographa 94**
 (b) Spores muriform **Arthothelium 91**
- (3) Algae *Phyllactidium*
 (a) Spores 1-septate ***Merarthonis 91**
 (b) Spores 2-several-septate **Arthoniopsis 91**
- b. Perithecia margined with a distinct proper exciple
Subfamily Graphidae 92
- (1) Thallus without cortex
 (a) Algae *Palmella*
 x. Perithecia with a single hymenium
 (x) Spores colorless
 m. Spores 1-celled

- (m) Hypothecium clear or brownish
Xylographa 93
- (n) Hypothecium black, carbonous
Lithographa 93
- n. Spores transeptate
Aulaxina 94
- (y) Spores dark
 - m. Spores transeptate
Encephalographa 94
 - n. Spores finally muriform
Xyloschistes 94
- y. Perithecia with 2-4 parallel hymenia
 - (x) Spores 1-celled
Ptychographa 94
 - (y) Spores transeptate
Diplogramma 94
- (b) Algae Chroolepus
 - x. Asci many-spored; spores filiform
Spirographa 96
 - y. Asci 1-8-spored
 - (x) Spores clear
 - m. Spores transeptate
 - (m) Paraphyses simple and not united
 - r. Ends of paraphyses little thickened, smooth
 - (r) Spores 1-septate
***Digraphis 98**
 - (s) Spores 2-several-septate
Graphis 96
 - s. Ends clavate and warted or spiny
***Psorographis 102**
 - (n) Paraphyses branched and united
Opegrapha 94
 - n. Spores muriform
 - (m) Paraphyses simple and not united
 - r. Ends of paraphyses not thickened, smooth
Graphina 99
 - s. Ends of paraphyses clavate, warted or spiny
†Acanthothecis 101
(not *Acanthothecium* Speg.)
 - (n) Paraphyses branched and united
Helminthocarpum 102
(incl. *Dictyographa 96*)
 - (y) Spores dark
 - m. Spores 1-septate
Melaspilea 96
 - n. Spores 2-several-septate
Phaeographis 99
 - o. Spores muriform
Phaeographina 100
- (c) Algae Phyllactidium: spores transeptate
 - x. Spores clear; paraphyses branched and united
Opegraphella 102
 - y. Spores dark; paraphyses simple and free
Micrographa 102
- (2) Thallus with a cortex: algae Chroolepus
Subfamily Dirinae 105
- (a) Spores elliptic to fusoid, 4-8-celled, clear
Dirina 106

- (b) Spores similar but brown **Dirinastrum 106**
- 3. Thallus present, fruticose, erect **Subfamily Roccellae 106**
 - a. Hyphae of cortex parallel with thallus surface
 - (1) Perithecia elongate, furrowed; spores clear, 8-9-celled **Ingaderia 107**
 - (2) Perithecia round
 - (a) Hypothecium black; spores clear
 - x. Exciple with algae **Dendrographa 107**
 - y. Exciple without algae **Roccellaria 107**
 - (b) Hypothecium clear; spores brown, spiny **Darbishirella 108**
 - b. Hyphae perpendicular to surface
 - (1) Perithecia elongate, furrowed
 - (a) Perithecia immersed; hypothecium clear **Roccellographa 108**
 - (b) Perithecia superficial; hypothecium black **Reinkella 108**
 - (2) Perithecia round
 - (a) Spores clear; perithecia entire
 - x. Hypothecium black
 - (x) Thallus mostly crustose, slightly fruticose **Roccellina 108**
 - (y) Thallus distinctly fruticose **Roccella 109**
 - y. Hypothecium clear
 - (x) Algae present below the hypothecium **Pentagenella 110**
 - (y) No algae below the hypothecium **Combea 109**
 - (b) Spores brown or brownish; perithecia deeply lobed
 - x. Medulla clear throughout **Schizopelte 110**
 - y. Inner medullary layer black **Simonyella 110**
- II. Perithecia in a stroma, mostly immersed **Subfamily Chiodectae 102**
 - I. Algae *Chroolepus*
 - a. Paraphyses simple and free
 - (1) Spores transeptate
 - (a) Spores clear **Glyphis 103**
 - (b) Spores brown **Sarcographa 103**
 - (2) Spores muriform
 - (a) Spores clear **Enterodictyum 104**
 - (b) Spores brown **Sarcographina 103**
 - b. Paraphyses branched and reticulately united
 - (1) Spores transeptate
 - (a) Spores colorless **Chiodectum 104**
 - (b) Spores brown or dark
 - x. Perithecia margined **Sclerophytum 105**
 - y. Perithecia marginless **Synarthonia 91**
 - (2) Spores muriform
 - (a) Spores clear **Minksia 241**

- (b) Spores brown **Enterostigma** 105
2. Algae **Phyllactidium**
- a. Spores 2-celled; paraphyses simple and free **Pycnographa** 105
- b. Spores many-celled; paraphyses branched and united **Mazosia** 105

Order 10. PEZIZALES

Mycelium various, but typically inconspicuous or invisible; propagaton by conidia, but usually not in evidence; reproductive body or apothecium at first closed and more or less globose, rarely elongate, then opening more or less completely into a cup, saucer or disk, waxy or fleshy, more rarely carbonous, leathery or gelatinous; asci typically 8-spored and paraphysate; spores various.

Family 30. PHACIDIACEAE

REHM 60

Apothecia sunken, more or less erumpent, disk-like or elongate, single or grouped, leathery or carbonous, black, firm, opening by lobes or by a rift; hypothecium poorly developed as a rule.

Hyalosporae

8:705, 11:431, 10:48, 14:813, 16:783, 18:155

Spores hyaline, 1-celled, globose to oblong

I. Apothecia concrete above with the epiderm

1. Apothecia and epiderm splitting radiately

Phacidium 8:709

2. Apothecia and epiderm splitting circumscissilely

Stegia 8:733

3. Apothecia and epiderm splitting irregularly

Cryptomyces 8:707

II. Apothecia and epiderm little or not at all concrete

Pseudophacidium R. 94

Phaeosporae

14:814

Spores dark, 1-celled, oblong

I. Apothecia superficial, membranous, lacinate

Phaeophacidium 14:814

Hyalodidymae

Spores hyaline, 1-septate, elliptic to oblong

I. Apothecia scutellate or oblong, lacinate

Schizothyrium R. 75

(incl. **Rhagadolobium** 14:816)

Phaeodidymae

Spores dark, 1-septate, elliptic to oblong

I. Apothecia in black foliicole spots

Cocconia 8:738

II. Apothecia stellately erumpent through epiderm

Metadothella 18: 162

III. Apothecia and epiderm concrete, laciniate

Keithia 10: 49**Phragmosporae**

8: 740

Spores typically hyaline, 2-several-septate, ovoid to oblong

I. Apothecia and epiderm concrete, laciniate

Sphaeropezia 8: 740, R. 72

II. Apothecia and epiderm not concrete, splitting irregularly

Pseudographis R. 90**Dictyosporae**

8: 764, 16: 790

Spores muriform, typically hyaline, ovoid to oblong

I. Apothecia round to oblong, splitting irregularly; paraphysate

Dothiora 8: 764, R. 108**Scolecosporae**

8: 744, 10: 51, 11: 432, 14: 817, 16: 789, 18: 163

Spores bacillar to filiform, typically hyaline, continuous or septate

I. Apothecia and epiderm concrete

1. Apothecia in black foliicole stroma-like spots

Rhytisma 8: 752, R. 82(incl. **Duplicaria** 8: 764)

2. Apothecia not in stroma-like spots

a. Apothecia and epiderm laciniate

Coccomyces 8: 744, R. 76

b. Apothecia and epiderm operculately circumscissile

Moutoniella 18: 163

II. Apothecia and epiderm not concrete

1. Apothecia round, laciniate

Coccophacidium R. 97

2. Apothecia oblong to elongate, hysterooid

Clithris 18: 165, R. 101**Family 31. STICTIDACEAE**

REHM 112

Apothecia sunken, finally more or less erumpent, round or elongate, single or grouped, typically waxy, rarely membranous or leathery, white or bright-colored, at least never black, splitting the epiderm laciniately or irregularly, hypothecium little developed.

Subfamily Eustictidae

REHM 113

Apothecia waxy, not deeply sunken, finally opening widely, and exposing the hymenium.

Hyalosporae

8:648, 10:44, 11:428, 14:806, 16:776, 18:146

Spores hyaline, 1-celled, globose to oblong

I. Spores globose

- 1. Asci 8-spored Lindaueella 16:777
- 2. Asci many-spored Flaminia 16:777

II. Spores elliptic to oblong

- 1. Paraphyses long-pointed, much longer than the asci Stegia 8:733, R. 155
- 2. Paraphyses blunt, swollen or branched
 - a. Paraphyses thread-shaped or forked
 - (1) Apothecia round
 - (a) Apothecia blackish; ascus pore blue with iodine Trochila 8:728, R. 127
 - (b) Apothecia bright-colored
 - x. Ascus pore blue with iodine
 - (x) Paraphyses forked, enlarged and colored above Ocellaria 8:654, R. 133
 - (y) Paraphyses little if at all enlarged or colored *Habrostictis R. 137
 - y. Ascus pore not blue with iodine Naevia 8:658, R. 145
 - (2) Apothecia oblong or elongate
 - (a) Hymenium blue with iodine Xylographa 8:664, R. 153
 - (b) Hymenium not blue with iodine Briardia 16:776, R. 151
 - b. Paraphyses irregularly branched
 - (1) Asci 8-spored Propolis 8:648, R. 141
 - (2) Asci many-spored Propolina 8:654

PhaeosporaeSpores 1-celled, dark, oblong Stictophacidium R. 1215**Didymosporae**

8:666, 10:45, 11:428, 14:808, 16:778, 18:147

Spores 1-septate, typically hyaline or bright-colored, oblong

- I. Paraphyses lacking Coccopeziza 10:45
- II. Paraphyses present
 - 1. Spores blue or green Ploettnera 16:778
 - 2. Spores hyaline
 - a. Spores with 1-2 cilia at each end; hysteroioid Iridionia 16:788
 - b. Spores muticate
 - (1) Paraphyses filiform or forked
 - (a) Apothecia round
 - x. Asci not blue with iodine *Naeviella R. 164
 - y. Asci blue with iodine
 - (x) Ascus pore alone blue with iodine Diplonaevia 8:666, R. 161
 - (y) Whole hymenium blue with iodine *Diplocryptis R. 158

- (b) Apothecia rounded, with flexuose clefts

Lauterbachella 16: 788

- (2) Paraphyses irregularly branched

- (a) Apothecia round; not blue with iodine

Propolidium 8: 667

- (b) Apothecia elongate; ascus pore blue with iodine

***Xyloglyphis** R. 170

Phragmosporae

8: 669, 10: 46, 11: 429, 14: 808, 16: 778, 18: 148

Spores 2-several-septate, hyaline, rarely darkish, oblong to elongate

- I. Spores somewhat fuscous

Eupropolis 8: 676

(incl. **Janseella** 16: 780)

- II. Spores hyaline

1. Paraphyses filiform or forked

- a. Asci not blue with iodine

***Merostictis** R. 164

- b. Asci blue with iodine

- (1) Ascus pore alone blue with iodine

Phragmonaevia 8: 674, R. 160

- (2) Whole hymenium blue with iodine

Cryptodiscus 8: 669, R. 158

2. Paraphyses branched; apothecia elongate

Xylogramma 8: 677, R. 169

Dictyosporae

8: 704, 11: 431, 14: 812, 16: 782, 18: 151

Spores muriform, typically hyaline, ovoid to oblong

- I. Asci 1-spored

Pleostictis 8: 703

- II. Asci 8-spored

1. Apothecia oblong, hysterioid

Melittiosporium 8: 704, R. 172

2. Apothecia round

- a. Apothecia urceolate

Platysticta 8: 703

- b. Apothecia disk-like

Delpontia 18: 151

Scolecosporae

8: 681, 10: 46, 11: 429, 14: 810, 16: 781, 18: 152

Spores bacillar or filiform, typically hyaline

- I. Asci 8-spored

1. Apothecia pilose

Lasiostictis 8: 696

2. Apothecia not pilose

- a. Spore cells separating

Schizoxylum 8: 697, R. 181

- b. Spore cells not separating

- (1) Paraphyses filiform or nearly so; asci cylindric

Stictis 8: 681, R. 175

(incl. **Karstenia** 8: 702, **Cerion**
18: 154)

- (2) Paraphyses much branched; asci clavate

Naemacyclus 8: 701, R. 173

- II. Asci many-spored

Carestiella 14: 810

Subfamily Ostropae

REHM 185

Apothecia membranous or leathery, deeply sunken, the scarcely opened tip alone erumpent.

- I. Spores 1-celled, elliptic; asci clavate **Laquearia R. 187**
- II. Spores many-celled, filiform; asci cylindric
 - 1. Apothecia cask-shaped, partly erumpent **Ostropa R. 188**
 - 2. Apothecia with only the thick ostiole erumpent **Robergea R. 189**

Family 32. TRYBLIDIACEAE

REHM 191

Apothecia sunken, then erumpent, often lobed, brown or black, membranous or horny; hypothecium well-developed, thick.

- I. Apothecia scattered
 - 1. Spores 1-septate
 - a. Spores with a mucose covering ***Tryblidis R. 194**
 - b. Spores without a mucose covering **Heterosphaeria R. 198**
 - 2. Spores 2-several-septate
 - a. Spores with a mucose covering **Tryblidiopsis R. 193**
 - b. Spores without a mucose covering **Odontotrema R. 204**
 - 3. Spores muriform **Tryblidium R. 196**
 - 4. Spores filiform ***Odontura R. 207**
- II. Apothecia cespitose or stromate; spores bacillar or filiform **Scleroderris R. 208**

Family 33. DERMATEACEAE

REHM 241

Apothecia sunken, then erumpent, cup-shaped to oblong, single or grouped, waxy, leathery or horny, mostly brownish or black; hypothecium more or less developed.

Hyalosporae

8: 547, 10: 36, 11: 422, 14: 794, 16: 762, 18: 121

Spores hyaline, 1-celled, globose to oblong

- I. Apothecia large, usually stalked or radicate at base
 - 1. Apothecia ear-shaped, more or less vertical, leathery
 - a. Spores ovoid to oblong **Midotis 8: 547**
 - b. Spores globose **Midotiopsis 18: 121**
 - 2. Apothecia urceolate or turbinate
 - a. Apothecia stalked; exciple and hypothecium prosenchymatic **Urnula 8: 548**
 - b. Apothecia stalked; exciple and hypothecium parenchymatic **Choriactis 18: 121**
 - c. Apothecia sessile, hairy; exciple parenchymatic, hypothecium prosenchymatic **Scytopezis 18: 122**
- II. Apothecia small, sessile or nearly so

1. Asci 8-spored

- a. Apothecia more or less corky **Dermatea** 8: 550, R. 246
 b. Apothecia coriaceous to subcorneous **Cenangium** 8: 556, R. 219
 (incl. *Ameghiniella* 8: 584, *Ephelina* 8: 585)

2. Asci many-spored, or 8-spored and many-spored

Tympanis 8: 578, R. 264**Phaeosporae**

16: 764, 18: 127

Spores dark, 1-celled, oblong

I. Apothecia coriaceous, erumpent

Phaeangium 16: 764**Hyalodidymae**

8: 587, 10: 37, 11: 424, 14: 798, 18: 127

Spores hyaline, 1-septate, elliptic to oblong

I. Apothecia patellate, coriaceous to corneous

Cenangella 8: 587

II. Apothecia elongate, cleft, subcorneous

Angelinia 18: 129**Phaeodidymae**

18: 128

Spores dark, 1-septate, elliptic to oblong

I. Apothecia patellate, coriaceous

Phaeangella 18: 128**Hyalophragmiae**

8: 594, 16: 765, 18: 129

Spores hyaline, 2-several-septate, elliptic to fusoid

I. Apothecia waxy-membranous, pilose, urceolate

Crumenula 8: 600, R. 235**Phaeophragmiae**

2: 757, R. 233

Spores dark, 2-several-septate, elliptic to fusoid

I. Apothecia hysterioid, cleft, coriaceous

Tryblidiella R. 233**Scolecosporae**

8: 601, 10: 37, 11: 425, 18: 130

Spores filiform, hyaline or subhyaline

I. Apothecia urceolate to cup-shaped, subcoriaceous

Godronia 8: 601, R. 237

II. Apothecia clavate, stipe corneous, disk submucose

Crinula 8: 606Family 34. **BULGARIACEAE**

РЕМ 444

Apothecia mostly superficial, cup-shaped to disk-shaped, usually smooth, gelatinous-fleshy or gelatinous-waxy, horn-like when dry; hypothecium gelatinous, more or less developed.

Hyalosporae

4: 609, 10: 38, 11: 425, 14: 801, 16: 766, 18: 131

Spores hyaline, 1-celled, globose to oblong

I. Spores globose**Pulparia 8: 612****II. Spores elliptic to bacillar****1. Apothecia in a lens-shaped gelatinous stroma****Physmatomyces 16: 770****2. Apothecia not in a stroma****a. Exciple lacking****(1) Asci 8-spored****(a) Apothecia microscopic, margined by changed paraphyses****Gloeopeziza 10: 41****(b) Apothecia larger; paraphyses not modified****Agyrium 8: 634, R. 450*****Agyrina 8: 636****(2) Asci 16-spored****b. Exciple present****(1) Lichenicole****Ahlesia 8: 633****(2) Not lichenicole****(a) Apothecia stipitate****Ombrophila 8: 613, R. 475****(incl. Stammnaria 8: 620, R. 465)****(b) Apothecia sessile****x. Asci 8-spored****(x) Apothecia smooth outside****m. Apothecia with an even disk****Orbilina 8: 621, R. 453****(incl. Bulgariopsis 18: 135)****n. Apothecia with a much folded disk****Haematomyces 8: 633****(y) Apothecia veined or roughened outside****m. Apothecia 1-2 cm. wide****Gloeocalyx 18: 132****n. Apothecia 2-9 cm. wide****Sarcosoma 10: 42, R. 497****y. Asci many-spored*****Myridium 8: 631****Phaeosporae**

8: 636, 10: 41, 14: 804, 16: 770, 18: 140

Spores dark, 1-celled, elliptic to fusoid

I. Apothecia turbinate, substipitate, closed at first, large**Bulgaria 8: 636, R. 494****II. Apothecia disciform, sessile, open at first, smaller****Bulgariella 8: 638****Hyalodidymae**

8: 639, 10: 42, 11: 427, 14: 805, 16: 771, 18: 142

Spores hyaline or subhyaline, 1-septate, elliptic to fusoid

I. Parasitic, urn-shaped; paraphyses forming an epithecium**Paryphedria 10: 43, R. 484****II. Saprophytic, disciform; epithecium lacking****Calloria 8: 639, R. 462**

Phaeodidymae

10: 42, 16: 771, 18: 142

Spores brown, 1-septate, elliptic to fusoid

- I. Apothecia subturbinate, sessile **Sorokinia** 10: 42

Phragmosporae

8: 641, 10: 43, 11: 427, 16: 773, 18: 143

Spores typically hyaline, 2-several-septate, fusoid

- I. Apothecia turbinate to disciform **Coryne** 8: 641, **R.** 485

Hyalodictyae

18: 145

Spores hyaline, muriform, ovoid

- I. Apothecia cupulate to plane **Dictyonina** 18: 144

Phaeodictyae

8: 646, 10: 44, 18: 144

Spores dark, muriform, ovoid to oblong

- I. Hymenium sinuate-gyrose, not margined **Haematomyxa** 8: 646
 II. Hymenium smooth, acute-margined **Sarcomyces** 10: 44

Scolecosporae

8: 646, 14: 805, 16: 775, 18: 145

Spores filiform, typically hyaline

- I. Apothecia without an exciple **Agyriopsis** 14: 805

- II. Exciple present

1. Apothecia dark or black; spores medium

Holwaya 8: 646

2. Apothecia gray or bright-colored; spores very long

Ophiogloea 18: 145**Family 35. PATELLARIACEAE**

REHM 277

Apothecia mostly superficial, cupulate to disk-shaped, more rarely boat-shaped or oblong, usually dark or black, carbonous, leathery, corneous or waxy; hypothecium typically well-developed.

Hyalosporae

8: 769, 10: 52, 11: 433, 14: 818, 16: 791, 18: 165

Spores hyaline, 1-celled, globose to oblong

- I. Asci many-spored

1. Spores globose

Biatorella 8: 469, **R.** 303

2. Spores allantoid

Biatorellina 18: 172

- II. Asci 8-spored

1. Apothecia oblong to elongate, cleft

Placographa **R.** 313

2. Apothecia round

- a. Parasitic on lichen thalli

- (1) Exciple present

Rhymbocarpus 14: 819

- (2) Exciple lacking **Nesolechia** 10: 53, R. 315
- b. Saprophytic
 - (1) Paraphyses branched, forming an epithecium
 - (a) Asci club-shaped
 - x. Subicle absent **Patinella** 8: 769, R. 310
 - y. Subicle present, radiate **Actinoscypha** 8: 774
 - (b) Asci cylindric **Starbaeckia** 10: 53
 - (2) Paraphyses simple; epithecium none **Psilothecium** 18: 168

Phaeosporae

10: 55

Spores dark, 1-celled, globose to elliptic

- I. Apothecia patellate, margined, black **Lagerheimia** 10: 55

Hyalodidymae

8: 779, 10: 56, 11: 434, 14: 820, 16: 792, 18: 173

Spores hyaline, 1-septate, elliptic to fusoid

- I. Parasitic on lichen thalli **Scutula** R. 321
- II. Not lichenicole
 - 1. Apothecia smooth, saprophytic **Patellea** 8: 783, R. 283
 - 2. Apothecia setose, parasitic on leaves **Johansonia** 8: 785

Phaeodidymae

8: 779, 10: 56, 11: 434, 14: 820, 16: 792, 18: 173

Spores dark, 1-septate, elliptic to fusoid

- I. Asci 8-spored
 - 1. Apothecia on a foliicole radiate subicle **Woodiella** 16: 794
 - 2. Apothecia not on a subicle
 - a. Apothecia round
 - (1) Apothecia superficial
 - (a) Saprophytic **Karschia** 8: 779, R. 345
 - (b) Parasitic on lichens ***Epilichen** 18: 177, R. 350
 - (2) Apothecia sunken, then erumpent
 - (a) Parasitic on lichens **Abrothallus** 8: 739, R. 358
 - (b) Saprophytic **Caldesia** R. 289
 - b. Apothecia elliptic to linear
 - (1) Apothecia irregularly elliptic or oblong
 - Melaspilea** 10: 58, R. 362
 - (2) Apothecia boat-shaped to linear **Hysteropatella** R. 367
- II. Asci 16-spored **Ravenelula** 8: 782
- III. Asci many-spored ***Pleospilis** 18: 179

Hyalophragmiae

8: 786, 10: 59, 11: 434, 14: 821, 16: 795, 18: 179

Spores hyaline, 2-several-septate, elliptic to fusoid

- I. Parasitic on lichens **Mycobilimbia** 10: 60, R. 327
- II. Saprophytic

1. Apothecia twisted when dry	Durella 8: 790, R. 286
2. Apothecia not contorted	Patellaria R. 329
(incl. Lecanidion 8: 795)

Phaeophragmiae

8: 786, 10: 59, 11: 434, 14: 821, 16: 795, 18: 179

Spores dark, 2-several-septate, elliptic to fusoid

I. Asci 8-spored

1. Margin of cup involute, densely costate-rugose	Rhytidopeziza 10: 65
2. Margin not costate-rugose	
a. Apothecia erumpent	Pseudotryblidium 10: 65, R. 370
b. Apothecia superficial	
(1) Parasitic typically on lichens	
(a) Apothecia round	Leciographa 10: 61, R. 372
(b) Apothecia elliptic to elongate	*Lecoglyphis R. 380
(2) Saprophytic	*Mycolecis , R. 372, 10: 61

II. Asci many-spored**Dictyosporae**

8: 802, 11: 435, 14: 823, 18: 185

Spores hyaline or subhyaline, muriform, ovoid to oblong

I. Apothecia laciniate, depressed-spheroid	Blitrydium 8: 802
II. Apothecia not laciniate, patellate	Tryblidaria 18: 186

Scolecosporae

8: 807, 10: 65, 11: 435, 14: 823, 16: 798

Spores hyaline or subhyaline, bacillar to filiform

I. Spores separating at the joints	Bactrospora 10: 67, R. 344
II. Spores not separating	
I. Apothecia sessile	
a. Parasitic	Mycobacidia 10: 66, R. 337
b. Saprophytic	Pragmopara R. 339
(incl. Scutularia 8: 807)
2. Apothecia stalked, turbinate	
a. Parasitic	*Parathalle R. 343
b. Saprophytic	Lahmia 10: 65, R. 341

Family 36. CALICIACEAE

REHM 388, ZAHLBRUCKNER 80

Mycelium inconspicuous and saprophytic, or parasitic on algae, forming a powdery, crustose, foliose or fruticose thallus; apothecia sessile or stalked, cup- to top-shaped, opening more or less completely, asci disappearing very early and the disk then covered with a persistent mass of spores and paraphyses, i. e., mazaedium; exciple prosenchymatic, horny, proper or thalline.

- I. Mycelium saprophytic, at least not forming a thallus**
- 1. Spores 1-celled, globose or globoid**

- a. Spores clear or merely yellowish
 - (1) Algae present but not forming a thallus
Farriolla 83
 - (2) Algae lacking
 - (a) Asci long and slender stalked, ovoid above
Caliciopsis R. 388
 - (b) Asci cylindric
Roesleria 8: 826, R. 396
- b. Spores dark
 - (1) Apothecia black, nearly sessile
Sphinctrina 83, R. 389
 - (2) Apothecia bright-colored, with a slender stalk
**Eucyphelis* R. 392
(*Cyphelium* Rehm)
- 2. Spores typically 2-several-celled
 - a. Spores 2-celled
 - (1) Apothecia sessile
Acolium R. 398
 - (2) Apothecia with a slender stalk
Mycocalicium R. 401
 - b. Spores 3-several-celled
Stenocybe 82 R. 413
- II. Mycelium forming a thallus with algae
 - 1. Thallus crustose
 - a. Spores 1-celled, globose or globoid
 - (1) Asci 8-spored
 - (a) Spores dark; disk more or less flat
 - x. Apothecia stalked
Chaenotheca 81
 - y. Apothecia sessile
**Holocyphis* 84
 - (b) Spores clear or yellowish; disk globose
Coniocybe 82
 - (2) Asci many-spored
Tylophorella 85
 - b. Spores 2-several-celled, transeptate or muriform
 - (1) Spores transeptate
 - (a) Spores 2-celled, dark or brown
 - x. Apothecia stalked
 - (x) Apothecia long-stalked
Calicium 81
 - (y) Apothecia with short thick stalk
Pyrgidium 83
 - y. Apothecia sessile
 - (x) Algae *Pleurococcus*
Cyphelium 83
 - (y) Algae *Chroolepus*
 - m. Proper exciple alone present
**Dipyrgis* 84
 - n. Thalline exciple also present
**Ditylis* 84
 - (b) Spores 3-many-celled
 - x. Proper exciple alone present
Pyrgillus 84
 - y. Thalline exciple also present
Tylophorum 84
 - (2) Spores muriform
Pseudacolium 84
 - 2. Thallus foliose
 - a. Thallus of horizontal scales with marginal apothecia
Calycidium 85

- b. Horizontal scales sterile; apothecia on cylindric podetia
Tholurna 85
- 3. Thallus fruticose
 - a. Thallus hollow; apothecia on the under side
Pleurocybe 85
 - b. Thallus with solid medulla; apothecia terminal
 - (1) Apothecia without thalline covering, goblet-like
Acroscyphus 86
 - (2) Apothecia enclosed in a globose thalline exciple, which finally opens irregularly at the top
Sphaerophorus 86

Family 37. CHRYSOTRICHACEAE

ZAHLEBRUCKNER 117, 127

Apothecia disk-form, margined, asci persistent; mazaedium lacking, thallus uniform, cobwebby, cottony or spongy, loose, without layers, algae *Palmella*, *Pleurococcus*, *Chroolepus* or *Cladophora*.

- I. Thallus with *Palmella* or *Pleurococcus*
 - 1. Spores 1-celled
Crocynia 242
 - 2. Spores 2-4-celled
Chrysothrix 117
- II. Thallus with *Chroolepus*; spores clear
 - 1. Spores 1-celled
***Holocoenis 128**
 - 2. Spores 2-celled
Coenogonium 127
- III. Thallus with *Cladophora*; apothecia lacking
Racodium 128

Family 38. COLLEMATACEAE

ZAHLEBRUCKNER 154, 158, 167, 168

Apothecia disk-form or pitcher-form, with persistent asci; thallus more or less gelatinous when moist, mostly without layers, always with blue-green algae, scaly, foliose or fruticose, rarely crustose.

- I. Algae *Gloeocapsa*, *Chroococcus* or *Xanthocapsa*; spores typically 1-celled, colorless
Subfamily Pyrenopsidae 158
- 1. Algae *Gloeocapsa*
 - a. Thallus crustose, scaly or dwarf fruticose
 - (1) Spores 1-celled
 - (a) Asci 8-spored
Pyrenopsis 159
 - (b) Asci 32-spored
***Pleopyrenis 160**
 - (2) Spores 2-celled
Cryptothele 159
 - b. Thallus foliose, of a single leaf; spores clear, 1-celled
Phylliscidium 160
 - c. Thallus fruticose, with rhizoids; spores clear, 1-celled
Synalissa 160
- 2. Algae *Chroococcus*
 - a. Thallus crustose; apothecia more or less open
Pyrenopsidium 160
 - b. Thallus foliose, of one leaf. umbilicate; apothecia closed
Phylliscum 161

3. Algae *Xanthocapsa*

a. Thallus crustose

(1) Spores 1-celled

- (a) Hymenium covered with a mass of algae and hyphae

Gonohymenia 161

- (b) Hymenium without epithelial mass

- x. Thallus pseudoparenchymatic at margin

Forssellia 161

- y. Thallus nowhere pseudoparenchymatic

Psorotichia 161

- (2) Spores 2-celled; apothecia closed
- Collemopsidium 161**

b. Thallus of one leaf, umbilicate, often lobed

- (1) Thallus pseudoparenchymatic

Anema 162

- (2) Thallus not pseudoparenchymatic

- (a) Spores 1-celled

- x. Hyphae loose, net-like at margin

Thyrea 162

- y. Hyphae perpendicular to the margin

Jenmania 162

- (b) Spores 2-celled

Paulia 163

c. Thallus fruticose, branched, upright

- (1) Thallus without layers

- (a) Asci 8-spored

Peccania 163

- (b) Asci 12-many-spored

***Pleoconis 164**

- (2) Thallus layered, with a cortex

Phloeopeccania 164II. Thallus with *Nostoc*; spores clear**Subfamily Collematae 168**

1. Apothecia with proper exciple only, biatorin

a. Spores 1-celled

- (1) Spores globose to fusoid, straight

- (a) Thallus crustose, scarcely gelatinous

Leprocollema 170

- (b) Thallus scaly or dwarf fruticose, gelatinous

Leciophysma 170

- (2) Spores needle-shaped, twisted

Koerberia 173

b. Spores transeptate, 2-many-celled

- (1) Spores 2-celled; thallus without cortex

Homothecium 171

- (2) Spores 4-8-celled; thallus with cortex

Arctomia 173

2. Apothecia with thalline exciple, lecanorin

a. Spores 1-celled

- (1) Thallus scaly or dwarf fruticose; spores thin-walled

- (a) Thallus without cortex

Physma 170

- (b) Thallus with pseudoparenchymatic cortex

Lemmopsis 171

- (2) Thallus large-leaved; spores thick-walled or mucose

Dichodium 171

b. Spores transeptate to muriform

- (1) Thallus without cortex

- (a) Spores 2-celled ***Dicollema 172**
 - (b) Spores transeptate, many-celled **Collema 171**
 - (c) Spores muriform **Blennothallia 172**
 - (2) Thallus with a pseudoparenchymatic cortex on one or both sides or pseudoparenchymatic throughout
 - (a) Spores transeptate, 3-many-celled **Leptogiopsis 175**
 - (b) Spores muriform **Leptogium 174**
- III. Thallus with Scytonema or Stigonema; spores colorless
Subfamily Ephebae 154
- 1. Thallus crustose to scaly
 - a. Thallus uniform, not corticate
 - (1) Spores 1-celled **Pterygiopsis 157**
 - (2) Spores 4-celled **Petractis 124**
 - b. Thallus corticate above **Porocyphus 157**
 - 2. Thallus dwarf fruticose, much branched, dark
 - a. Apothecia sunken in swellings of the thallus
 - (1) Spores 1-celled; paraphyses present **Ephebeia 155**
 - (2) Spores 2-3-celled **Ephebe 155**
 - b. Apothecia superficial
 - (1) Thallus without pseudoparenchymatic cortex or central medulla
 - (a) Paraphyses capitate, septate **Spilonema 154**
 - (b) Paraphyses filiform, not septate **Thermutis 154**
 - (2) Thallus with large-celled pseudoparenchymatic cortex and central medulla
 - (a) Cortex of one row of cells; spores 2-celled **Leptodendriscum 155**
 - (b) Cortex of several rows
 - x. Spores 1-celled **Leptogidium 156**
 - y. Spores 2-celled **Polychidium 156**
- IV. Algae Rivularia; spores clear **Subfamily Lichinae 164**
- 1. Apothecia disk-form; thallus scaly to granular
 - a. Apothecia with proper exciple; algae horizontal **Pterygium 165**
 - b. Apothecia with thalline exciple; algae erect **Steinera 166**
 - 2. Apothecia almost perithecioid; thallus dwarf fruticose
 - a. Algae in the middle of the thallus and parallel with the long axis of the branches **Lichinodium 166**
 - b. Algae absent from the middle but marginal beneath the cortex
 - (1) Algae parallel with the long axis of the branches **Lichina 167**
 - (2) Algae perpendicular to the long axis
 - (a) Paraphyses present **Lichinella 166**
 - (b) Paraphyses absent **Homopsella 167**

Family 39. PELTOPHORACEAE

ZAHLEBRUCKNER 122, 176, 190

Thallus firm, not at all gelatinous, crustose or foliose, more or less lobed and somewhat erect at the margin but never truly fruticose, typically attached to the substratum by rhizoids or by a navel, with a pseudoparenchymatic cortex on one or both sides or pseudoparenchymatic throughout; apothecia typically sunken in the thallus or grown together with it on the whole under side, more or less margined by the thallus, but a proper exciple lacking.

I. Thallus uniform to crustose; algae Protococcus, rarely Pleurococcus

Subfamily Ectolechia 122

1. Spores transeptate, usually 2-3-celled
 - a. Paraphyses not branched
 - (1) Paraphyses free; no algae below the hypothecium
Asterothyrium 123
 - (2) Paraphyses united; algae below the hypothecium
Lecaniella 124
 - b. Paraphyses branched and united
 - (1) Spores 2-celled
Actinoplaca 124
 - (2) Spores many-celled
Tapellaria 243
2. Spores muriform
 - a. Asci 1-spored; hypothecium without algae
 - (1) Paraphyses unbranched, free
Lopadiopsis 123
 - (2) Paraphyses branched, united
 - (a) Epithecium without algae
Sporopodium 123
 - (b) Epithecium with algae
***Gonothecis 123**
 - b. Asci 8-spored; hypothecium with algae below
Arthotheliopsis 124

II. Thallus foliose or foliose scaly, rarely subfruticose; algae typically bluegreen, rarely bright-green

1. Apothecia not marginal; thallus pseudoparenchymatic throughout
Subfamily Heppiae 176
- One genus, parasitic on Scytonema
Heppia 177
2. Apothecia typically marginal or even with the thallus; thallus layered
Subfamily Peltophorae 190

- a. Thallus foliose, usually large-leaved
 - (1) Apothecia on the upper side of the thallus
 - (a) Apothecia marginal on lobes of thallus; lower surface of thallus without cortex
 - x. Algae Nostoc
†Peltophora 194
(Peltigera)
 - y. Algae Palmella (Dactylococcus)
***Chloropeltis 194**
 - (b) Apothecia superficial; lower surface with cortex below the apothecia
 - x. Algae Nostoc
Solorina 192
 - y. Algae Palmella
Solorinina 192
 - (2) Apothecia on the under side of elongate thallus lobes; thallus completely corticate on both sides
 - x. Algae Nostoc
Nephromium 194

- y. Algae *Palmella* **Nephroma 193**
 b. Thallus minute, small triangular scales radiating from the apothecium
 (1) Asci 8-spored; spores brownish, 4-6-celled **Asteristium 191**
 (2) Asci many-spored; spores clear, 2-celled **Solorinella 192**

Family 40. LECIDEACEAE

Zahlbruckner 114, 129, 138, 144

Thallus firm, not gelatinous, crustose, scaly or foliose, exceptionally dwarf fruticose, with rhizoids or a navel in the larger forms, with or without cortex; apothecia superficial or somewhat sunken at first, with a characteristic proper exciple, very rarely lacking, but without a thalline exciple. The absence of the latter distinguishes this family from the Parmeliaceae.

I. Thallus uniform or crustose

1. Algae *Chroolepus* or *Phyllactidium* **Subfamily Lecanactidae 114**
 a. Proper exciple lacking, or rudimentary and lateral
 (1) Spores transeptate; exciple mostly absent **Schismatomma 115**
 (2) Spores muriform; exciple thin, complete **Melampyidium 116**
 b. Proper exciple well-developed, carbonous
 (1) Spores 2-celled **Arthoniactis 115**
 (2) Spores 4-many-celled **Lecanactis 115**
 (3) Spores needle-shaped ***Scoleactis 115**
 2. Algae *Pleurococcus* or *Palmella* **Subfamily Lecideae 129**
 a. Thallus uniform-crustose, loose, without cortex; spores clear, fusoid, 4-celled **Pilocarpum 116**
 b. Thallus typically crustose, firm
 (1) Asci 1-8-spored, rarely 16-32-spored
 (a) Spores 1-celled
 x. Spores clear
 (x) Asci 1-2-spored; spores large, thick-walled **Mycoblastus 133**
 (y) Asci 8-spored
 m. Exciple black, carbonous **Lecidea 130**
 n. Exciple clear or colored, not carbonous **Biatora 132**
 (z) Asci 16-32-spored ***Pleolecis 132**
 y. Spores brown **Orphniospora 133**
 (b) Spores 2-celled
 x. Spores clear
 (x) Paraphyses simple
 m. Spores thick-walled, large **Megalospora 134**
 n. Spores thin-walled, small
 (m) Thallus with cortex ***Diphloeis 136**
 (n) Thallus without cortex

- r. Exciple and hypothecium dark or black
Catillaria 133
 - s. Exciple and hypothecium clear or bright
Biatorina 134
 - (y) Paraphyses branched, in a slimy hymenium
***Diphanis 138**
 - y. Spores brown; paraphyses branched
***Diphaeis 138**
 - (c) Spores 4-many-celled
 - x. Spores elliptic to long-fusoid
 - (x) Thallus not corticate, crustose-uniform
 - m. Spores thin-walled **Bacidia 135**
 - n. Spores thick-walled **Bombyliospora 136**
 - (y) Thallus corticate, warty to scaly
Toninia 136
 - y. Spores needle-shaped or filiform
†Scolecosporis 136
(Scoliciosporum)
 - (d) Spores muriform
 - x. Spores clear
 - (x) Spores with mucus covering; paraphyses branched
***Phalodictyum 138**
 - (y) Spores without mucus cover; paraphyses simple
Lopadium 137
 - y. Spores brown, mucose
Rhizocarpum 137
 - (2) Asci many-spored
 - (a) Exciple bright-colored, soft **Biatorella 151**
 - (b) Exciple dark or black, hard **Sporostatia 152**
- II. Thallus scaly or foliose; algae *Pleurococcus* or *Palmella*
Subfamily Phyllopsorae 138
- I. Thallus scaly, with rhizoids; disk even
 - a. Spores 1-celled
 - (1) Hypothecium pseudoparenchymatic
Phyllopsora 138
 - (2) Hypothecium not pseudoparenchymatic
 - (a) Exciple clear or bright **Psoromaria 183**
 - (b) Exciple dark or black **Psora 132**
 - b. Spores transeptate **Psorella 139**
 - 2. Thallus mostly with one large leaf; disk often furrowed
Subfamily Gyrophorae 147
 - a. Spores 1-celled; disk furrowed in most of the species
Gyrophora 147
 - b. Spores transeptate
 - (1) Spores 2-many-celled, colorless ***Merophora 148**
 - (2) Spores 2-celled, brown **Dermaticum 149**
 - c. Spores muriform, dark **Umbilicaria 149**
- III. Thallus dwarf fruticose, of low erect slightly branched podetia, horizontal
thallus lacking; spores clear, 2-celled **Sphaerophoropsis 133**

Family 41. CLADONIACEAE

ZAHLEBRUCKNER 139

Thallus of two kinds, one horizontal on the substratum, crustose, scaly to foliose, the other consisting of erect clubshaped, cupshaped or filiform, simple or branched podetia; algae typically *Pleurococcus*; apothecia terminal or lateral, mostly convex to globose, with proper exciple only, except in *Chlorocaulum*; spores colorless.

I. Apothecia with proper exciple**1. Podetia short, simple, rarely forked; apothecia terminal****a. Podetia equal, not broadened above****(1) Podetia covering the surface****(a) Hypothecium clear****x. Spores 1-celled*****Baeomyces* 140****y. Spores transeptate****(x) Spores elliptic to rod-shaped****m. Spores 2-celled******Dibaeis* 140****n. Spores 4-celled****(m) Algae bluegreen******Cyanobaeis* 141****(n) Algae yellow-green*****Heteromyces* 141****(y) Spores filiform, many-celled*****Gomphyllus* 141****(b) Hypothecium dark; spores 1-celled*****Pilophorum* 142****(2) Podetia marginal on a foliose thallus*****Gymnoderma* 142****b. Podetia broadened above into lobes or tongues bearing the hymenium on one side****(1) No algae below the hymenium; medulla uniform*****Glossodium* 142****(2) Algae below the hymenium; medulla with thicker strands*****Thysanothecium* 142****2. Podetia funnellform, cupshaped or more or less branched, large****a. Spores 1-celled; podetia hollow*****Cladonia* 143****b. Spores 4-many-celled*****Stereocaulum* 146****c. Spores muriform*****Argopsis* 146****II. Apothecia with thalline exciple******Chlorocaulum* 146****Family 42. PARMELIACEAE**

ZAHLEBRUCKNER 118, 124, 150, 195, 199, 207, 216

Thallus of one kind, podetia lacking, firm, not gelatinous, crustose, scaly, foliose or fruticose, often with rhizoids, typically layered, algae typically yellow green, but bluegreen in two subfamilies; apothecia characterized by a thalline exciple, which is rarely lacking, superficial, rarely immersed

I. Thallus typically crustose, sometimes scaly or lobed at the margin**1. Algae *Pleurococcus* or *Parmella*, rarely *Protococcus*****a. Asci 1-32-spored, mostly 8-spored****(1) Disk conspicuous, not perithecioid****Subfamily *Leanorae* 199****(a) Spores 1-celled**

- x. Asci 1-8-spored
 - (x) Paraphyses simple, free
 - m. Spores straight, elliptic to oblong
 - (m) Thallus bright yellow; pycnoconidia elliptic
Candelariella 207
 - (n) Thallus rarely bright yellow; conidia filiform
 - r. Cortex not pseudoparenchymatic
Lecanora 201
 - s. Cortex pseudoparenchymatic
Psoroma 183
 - n. Spores crescent to falcate
Harpidium 199
 - (y) Paraphyses branched and united
Ochrolechia 203
***Myriolectis** 202
 - y. Asci 12-many-spored
- (b) Spores 2-celled
 - x. Paraphyses simple, free
 - (x) Sterigmata exobasidial
Lecania 204
 - (y) Sterigmata endobasidial
Icmadophila 204
(incl. *Placolecania* 205)
 - y. Paraphyses branched, united
Calenia 205
- (c) Spores 4-many-celled
 - x. Apothecia superficial
 - (x) Asci 1-8-spored
 - m. Thallus with cortex
Haematomma 205
 - n. Thallus without cortex
 - (m) Paraphyses forked; spores moniliform, 30-40-celled
Conotrema 121
 - (n) Paraphyses simple; spores not moniliform, 8-30-celled
***Adermatis** 204
***Dyslecanis** 204
 - (y) Asci 16-32-spored
***Dyslecanis** 204
 - y. Apothecia immersed; thallus without cortex
 - (x) Paraphyses simple, free
Phlyctella 206
 - (y) Paraphyses branched and united
Phlyctidia 206
- (d) Spores muriform
 - x. Spores clear, at least not dark
 - (x) Apothecia superficial, broad
Myxodictyum 206
 - (y) Apothecia immersed, small
Phlyctis 206
 - y. Spores dark
Diploschistes 122
- (2) Disk small, more or less closed and perithecioid; apothecia mostly sunk-
en in warts
Subfamily Pertusariae 195
 - (a) Spores 1-celled
 - x. Paraphyses simple, free
Perforaria 195
 - y. Paraphyses branched and united
Pertusaria 195

- (b) Spores 2-celled; paraphyses branched and united
Varicellaria 198
- b. Asci many-spored; spores 1-celled, more rarely 2-celled
Subfamily Acarosporae 150
 - (1) Apothecia superficial
 - (a) Thallus bright yellow ***Pleochroma 207**
 - (b) Thallus not bright yellow **Maronea 152**
 - (2) Apothecia typically immersed, with mostly narrow disk
Acarospora 152
- 2. Algae *Chroolepus* or *Phyllactidium*; apothecia with thalline exciple, at least when young
Subfamily Gyalectae 124
(incl. *Thelotremae* 118)
 - a. Thalline exciple present and persistent
 - (1) Spores 1-celled **Jonaspis 125**
 - (2) Spores 2-celled ***Ocellis 118**
 - (3) Spores 4-many-celled
 - (a) Spores clear
 - x. Apothecia sprouting repeatedly from the margin, forming erect forked chains of apothecia **Polystroma 121**
 - y. Apothecia not in chains
 - (x) Algae *Chroolepus*
 - m. Exciple and hypothecium clear
Ocellularia 118
 - n. Exciple and hypothecium dark, hard
Sagiolechia 126
 - (y) Algae *Phyllactidium* **Phyllophthalmaria 120**
 - (b) Spores brown **Phaeotrema 119**
 - (4) Spores muriform
 - (a) Spores clear
 - x. Paraphyses simple, free **Thelotrema 119**
 - y. Paraphyses branched and united
***Phanotylum 121**
 - (b) Spores dark or brown
 - x. Paraphyses simple, free **Leptotrema 120**
 - y. Paraphyses branched and united
 - (x) Apothecia sunken in groups in a stroma
Tremotylum 120
 - (y) Apothecia not in a stroma
Gyrostomum 120
 - b. Thalline exciple present at first, then more or less completely disappearing
 - (1) Asci 1-8-spored
 - (a) Spores 2-celled **Microphiale 125**
 - (b) Spores 4-many-celled **Bryophagus 126**
 - (c) Spores muriform **Gyalecta 125**
 - (2) Asci 12-many-spored
 - (a) Spores 2-celled **Ramonia 125**
 - (b) Spores 6-many-celled **Pachyphiale 126**
- II. Thallus typically foliose or fruticose, sometimes small-leaved or scaly; thalline exciple sometimes lacking

- i. Algae *Pleurococcus*, *Protococcus*, *Palmella* or *Cystococcus*
 - a. Asci many-spored; apothecia cespitose on a one-leaved thallus
***Glypholecia* 153**
 - b. Asci 1-32-spored
 - (1) Thallus foliose, horizontal or upright, rarely fruticose, typically dorsiventral
 - (a) Thallus with cyphellae or pseudocyphellae or furnished with well-developed clubshaped cephalodia
 - x. Lower side of thallus with cyphellae or pseudocyphellae
 - (x) Apothecia with thalline exciple
 - m. Spores 2-celled
 - (m) Spores clear ****Diphanosticta* 189**
 - (n) Spores brown ****Diphaeosticta* 189**
 - n. Spores 4-many-celled
 - (m) Spores clear ****Phanosticta* 189**
 - (n) Spores brown ***Sticta* 188**
 - (y) Apothecia with proper exciple only
****Dysticta* 189**
 - y. Lower side of thallus without cyphellae or pseudocyphellae; thallus typically with cephalodia
 - (x) Algae *Protococcus* ***Lobaria* 185**
 - (y) Algae *Cystococcus*, i. e., in mucose colonies
****Cystolobis* 188**
 - (b) Thallus typically without cyphellae, pseudocyphellae, and cephalodia
Subfamily *Parmeliae* 207
 - x. Asci 16-32-spored ***Candelaria* 209**
 - y. Asci 2-8-spored
 - (x) Cortex on both sides of thallus
 - m. Apothecia superficial
 - (m) Lower cortex more or less cellular, usually with rhizoids
***Parmelia* 211**
(incl. *Parmeliopsis* 209)
 - (n) Lower cortex without rhizoids, spongy, of net-like hyphae
***Anzia* 213**
 - n. Apothecia marginal or terminal; thallus often fruticose
 - (m) Disks upright from the beginning
***Cetraria* 214**
 - (n) Disks on the under side of thallus lobes, later upright by the twisting of the lobes
***Nephromopsis* 216**
 - (y) Cortex on the upper side alone
 - m. Apothecia superficial; lower surface without cyphellae
***Physcidia* 209**
 - n. Apothecial terminal; cyphellae on lower side
***Heterodea* 208**
 - (2) Thallus fruticose, erect or hanging, often long and hair-like; radial, rarely dorsiventral in structure **Subfamily *Usneae* 216**
 - (a) Spores 1-celled or unknown

- x. Medulla traversed by varying solid strands

Letharia 218

- y. Medulla uniform without strands

- (x) Cortex formed of hyphae running lengthwise

- m. Spores clear; asci 8-spored

Bryopogon 219

- n. Spores brownish; asci 4-spored

Alectoria 219

- (y) Cortex of hyphae more or less perpendicular to the long axis, pseudoparenchymatic

- m. Medulla of hyphae running lengthwise

- (m) Medulla loose, not horny; apothecia unknown

Thamnolia 225

- (n) Medulla firm, horny

- r. Thallus low, podetium-like; apothecia unknown

Siphula 225

- s. Thallus fruticose, elongate; apothecia known

- (r) Thallus dorsiventral, without fibrous branches; medulla and cortex not separable

Everniopsis 218

- (s) Thallus radial, usually with fibrous branches; medulla and cortex readily separable

Usnea 223

- n. Medulla of hyphae running in all directions

- (m) Thallus more or less hollow

- r. Thallus swollen, tubular

Dactylina 218

- s. Thallus not swollen and tubular

- (r) Thallus fruticose, erect

Dufourea 218

- (s) Thallus podetium-like; apothecia unknown

Endocena 226

- (n) Thallus flattened, not hollow, dorsiventral

Evernia 217

- (b) Spores 2-celled

Ramalina 220

- (c) Spores muriform, brown, large; asci 1-spored

Oropogon 220

2. Algae bluegreen, Scytonema or Nostoc

- a. Thallus large-leaved, with cyphellae, pseudocyphellae or cephalodia

- (1) Lower side of thallus with cyphellae or pseudocyphellae

- (a) Apothecia with thalline exciple

- x. Spores clear, bacillar to acicular, 2-8-celled

***Podostictina 189**

- y. Spores brown

- (x) Spores 2-celled

Stictina 189

- (y) Spores 4-celled

***Merostictina 189**

- (b) Apothecia with proper exciple only

***Dystictina 190**

(2) Cyphellae or pseudocyphellae absent; cephalodia usually present

(a) Apothecia with thalline exciple

***Phycodiscis 188**

(b) Apothecia with proper exciple only

Lobarina 188

b. Thallus scaly to small-leafy, sometimes crustose, exceptionally large-leafy, without cyphellae, etc. **Subfamily Pannariae 178**

(1) Lower surface of thallus scarcely or not at all veined; spores 1-2-celled

(a) Upper cortex well-developed, distinct

x. Upper cortex with hyphae perpendicular to it

(x) Upper cortex hairy or pilose

Erioderma 183

(y) Upper cortex not hairy

m. Apothecia with thalline exciple

(m) Spores 1-celled; algae Nostoc

Pannaria 181

(n) Spores 2-celled; algae Scytonema

Massalongia 183

n. Apothecia with proper exciple only

(m) Spores 1-celled **Parmeliella 181**

(n) Spores 2-many-celled **Placynthium 181**

y. Upper cortex of horizontal hyphae

Coccocarpia 184

(b) Upper cortex indistinct; algae occupying nearly the whole width of the thallus

Lepidocellema 180

(2) Lower surface of thallus with distinct forked veins; spores 4-celled

Hydrothyria 184

Family 43. PHYSICIACEAE

ZAHLBRUCKNER 226-234

Thallus crustose, foliose or fruticose, as in Parmeliaceae; apothecia mostly lecanorin, sometimes with proper exciple alone; spores normally 2-celled, with more or less thickened cross-wall, often traversed by a line-like canal, or exceptionally 1-many-celled or muriform

I. Spores 2-celled

1. Spores clear

a. Thallus without cortex, uniform or crustose

(1) Apothecia with thalline exciple **Caloplaca 227**

(2) Apothecia with proper exciple only

Blastenia 226

b. Thallus with cortex, foliose or fruticose

(1) Thallus foliose, horizontal or ascending, dorsiventral, with rhizoids, cortex pseudoparenchymatic on both sides

Xanthoria 229

(2) Thallus fruticose, erect, radial, cortex of conglutinate longitudinal hyphae

Theloschistes 230

2. Spores dark or brown

a. Thallus without cortex, uniform or crustose

- (1) Apothecia with thalline exciple
 - (a) Asci 8-spored **Rinodina 232**
 - (b) Asci 12-24-spored ***Pleorinis 233**
- (2) Apothecia with proper exciple only **Buellia 231**
- b. Thallus with cortex, foliose or fruticose
 - (1) Upper cortex of perpendicular hyphae, pseudoparenchymatic
 - (a) Apothecia with thalline exciple
 - x. Hypothecium clear **Physcia 234**
 - y. Hypothecium black **Dirinaria 235**
 - (b) Apothecia with proper exciple only **Pyxine 234**
 - (2) Upper cortex of hyphae parallel with the long axis, not pseudoparenchymatic; apothecia with proper exciple **Anaptychia 236**
- II. Spores 3-4-celled
 - 1. Spores clear
 - a. Thallus without cortex, uniform or crustose
 - (1) Apothecia with thalline exciple ***Meroplacis 228**
 - (2) Apothecia with proper exciple only **Xanthocarpia 227**
 - b. Thallus with cortex, fruticose **Niorma 230**
 - 2. Spores brown
 - a. Thallus without cortex, uniform or crustose
 - (1) Apothecia with thalline exciple ***Merorinis 233**
 - (2) Apothecia with proper exciple alone **Diplotomma 232**
 - b. Thallus with cortex, foliose; exciple proper ***Phragmopyxine 234**
- III. Spores muriform, brown
 - 1. Thallus without cortex, uniform or crustose ***Dictyorinis 233**
 - 2. Thallus with cortex, foliose **Hyperphyscia 236**

Family 44. MOLLISIACEAE

REHM 503

Apothecia superficial or erumpent, cupulate to disk-shaped, mostly smooth, rarely with hairs, typically soft-waxy; distinguished from all other families by the typically brownish exciple, which is entirely parenchymatic, or at least about the base.

Subfamily Eumollisiae

Apothecia superficial from the beginning

Hyalosporae

Spores hyaline, 1-celled, globose to elliptic

- I. Apothecia not on a subicle
 - 1. Spores globose **Mollisiella 18:64**
 - 2. Spores elliptic to fusoid **Mollisia R. 511, 8:321**

- II. Apothecia on a subicle **Tapesia** R. 573, 8: 371

Hyalodidymae

Spores hyaline, 1-septate, elliptic to oblong

- I. Apothecia not on a subicle **Niptera** R. 549, 8: 480

- II. Apothecia on a subicle

1. Spores with a mucose covering **Stictoclypeolum** 18: 110

2. Spores not mucose

- a. Spores constricted, large, $50 \times 25 \mu$ **Psorotheciopsis** 16: 746

- b. Spores not constricted, small, $12 \times 5 \mu$ **Linhartia** 16: 744

Hyalophragmiae

Spores hyaline, 2-several-septate, elliptic to fusoid

- I. Apothecia not on a subicle or thallus **Belonidium** R. 561, 8: 496

- II. Apothecia on a subicle or thallus

1. Spores ciliate at each end **Ciliella** 16: 748

2. Spores not ciliate

- a. Apothecia on a subicle of hyphal threads **Trichobelonium** R. 590, 16: 747

- b. Apothecia on a parenchymatic thallus **Pazschkea** 14: 788
(incl. **Psorotheciella** 16: 746)

Hyalodictyae

Spores hyaline, muriform, ovoid to oblong

- I. Subicle present; asci 1-4-spored; spores mucose

†Melittosporis 16: 751
(**Melittosporiopsis**)

Scolecosporae

Spores hyaline, filiform, usually septate

- I. Apothecia gregarious; subicle lacking **Belonopsis** R. 571, 16: 752

Subfamily Pyrenopezizae

Apothecia at first covered, then erumpent and more or less superficial

Hyalosporae

Spores hyaline, 1-celled, globose to oblong

- I. Apothecia bright-colored, on living leaves

Pseudopeziza R. 596, 8: 723

- II. Apothecia dark-brown without, not on living leaves

1. Apothecia with bristles **Pirottaea** R. 636, 8: 386

2. Apothecia without bristles, but sometimes with projecting rows of cells

- a. Subicle lacking **Pyrenopeziza** R. 608, 8: 354

- b. Subicle present ***Spilopezis** R. 620

Phaeosporae

Spores dark or brownish, 1-celled, elliptic to oblong

- I. Apothecia leathery, bright-colored outside

Velutaria R. 645, 8: 488

Hyalodidymae

Spores hyaline, 1-septate, elliptic to fusoid

- I. Apothecia scarcely erumpent, bright colored

Fabraea R. 599, 8: 735

- II. Apothecia nearly superficial, dark-brown without

**Dibelonis* R. 638

Hyalophragmiae

Spores hyaline, 2-several-septate, oblong to fusoid

- I. Apothecia at last superficial, more or less roughened

Beloniella R. 638

Family 45. HELOTIACEAE

REHM 647

Apothecia mostly superficial, rarely erumpent or arising from a sclerotium, typically stalked, sometimes sessile, cupulate to disk-shaped, waxy; distinguished by an exciple which is completely prosenchymatic.

Subfamily Helotiae

Apothecia not hairy

Hyalosporae

Spores hyaline, 1-celled, globose to oblong

- I. Apothecia on a subicle

Eriopeziza R. 693

- II. Apothecia not on a subicle

1. Apothecia arising from a sclerotium, long-stalked

Sclerotinia R. 803, 8: 195

2. Apothecia not arising from a sclerotium

- a. Apothecia green, arising from a green substratum

Chlorosplenium R. 752, 8: 315

- b. Apothecia not on a green substratum

- (1) Apothecia margined by a row of triangular teeth

- (a) Apothecia stalked

Cyathicula R. 740, 8: 304

- (b) Apothecia sessile

**Pezoloma*

- (2) Apothecia without teeth

- (a) Asci many-spored

Comesia 8: 468

- (b) Asci typically 8-spored

- x. Apothecia sessile

Pezizella R. 653, 8: 275

- y. Apothecia stalked

- (x) Ascus pore blue with iodine

Helotium R. 772, 8: 210

(incl. *Ciboria* R. 754, 8: 201)

- (y) Ascus pore not blue with iodine

Phialea R. 708, 8: 251

(incl. *Helotium* in part)

Hyalodidymae

Spores hyaline, 1-septate, elliptic to fusoid

- I. Apothecia typically sessile *Eubelonis R. 685
- II. Apothecia stalked
 - 1. Stalk ridged or folded Lanzia 8: 479
 - 2. Stalk not ridged or folded Hymenoscypha R. 781

Hyalophragmiae

Spores hyaline, 2-several-septate, elliptic to fusoid

- I. Apothecia not toothed at margin
 - 1. Apothecia sessile Belonium R. 685, 8: 492
 - 2. Apothecia stalked
 - a. Subicle lacking
 - (1) Spores muticate Belonioscypha R. 743
 - (a) Paraphyses colorless, epithecium lacking Rutstroemia R. 763
 - (b) Paraphyses colored, forming an epithecium *Belospora R. 744, 8: 488
 - (2) Spores 1-ciliate at each end Massea 18: 99
 - b. Subicle present
- II. Apothecia with a row of triangular teeth at margin
 - 1. Apothecia sessile *Merodontis 18: 102
 - 2. Apothecia stalked Davincia 18: 101

Scolecosporae

Spores typically hyaline, filiform

- I. Apothecia sessile or merely narrowed below
 - 1. Apothecia smooth Gorgoniceps R. 690, 8: 504
 - 2. Apothecia hairy Arachnopeziza R. 698
- II. Apothecia stalked Pocillum R. 747, 8: 605

Subfamily Dasyscyphae

REHM 824

Apothecia hairy

Hyalosporae

Spores hyaline, 1-celled, globose to fusoid

- I. Spores globose Lachnellula R. 862, 8: 390
- II. Spores elliptic to fusoid
 - 1. Paraphyses lance-shaped, pointed
 - a. Apothecia sessile *Dyslachnum R. 868, 888
 - b. Apothecia stalked Lachnum R. 870
 - 2. Paraphyses filiform, blunt
 - a. Apothecia divided above into 3-6 lobes, black Arenaea 18: 75
 - b. Apothecia entire, rarely black

- (1) Apothecia hairy with distinct bristles
 - (a) Hairs shining, clear, non-septate, nearly solid
**Phalothrix* R. 831
 - (b) Hairs dull, usually septate, hollow
 - x. Apothecia sessile **Dasypezis* R. 829, 842
 - y. Apothecia stalked *Dasyscypha* R. 832, 8: 432
- (2) Apothecia villose with projecting hyphae
Hyphoscypha 18: 87

Hyalodidymae

Spores hyaline, 1-septate, elliptic to fusoid

- I. Spores at first 1-celled, but finally 2-celled

Lachnella R. 853, 8: 391
(incl. *Perrotia* 18: 90)

Hyalophragmiae

Spores hyaline, 2-several-septate, oblong to fusoid

- I. Paraphyses lance-shaped, pointed *Erinella* R. 910, 8: 507
- II. Paraphyses bearing conidia at the tips *Diplocarpa* 18: 110

Family 46. PEZIZACEAE

REHM 913

Apothecia typically terrestrial, erumpent or superficial, sessile or stalked, urn-shaped to disciform, smooth or hairy, fleshy or fleshy-waxy, rarely leathery; usually medium to large forms.

Subfamily Pezizae

Apothecia smooth, i. e., without hairs

Hyalosporae

Spores hyaline, 1-celled, globose to fusoid

- 1. Asci not blue with iodine
 - 1. Apothecia cleft on one side, ear-like *Otidea* R. 1023, 8: 94
 - 2. Apothecia not ear-like
 - a. Spores globose
 - (1) Apothecia fleshy or fleshy-waxy
 - (a) Substipitate, parasitic *Pitya* R. 925, 8: 209
 - (b) Sessile, terrestrial *Detonia* R. 927, 1269, 8: 105
(*Barlaea* 8: 111, *Otidella* 8: 99)
 - (2) Apothecia cartilaginous †*Peltophoromyces* 16: 720
(*Peltigeromyces*)
 - b. Spores elliptic to fusoid
 - (1) Apothecia sessile
 - (a) Spores with reticulately thickened wall
Aleuria R. 968
 - (b) Spores smooth or roughened
 - x. Apothecia not on a subicle *Humaria* R. 934, 8: 118

- y. Apothecia on a subicle **Pyronema** R. 962, 8: 107
(incl. **Phycascus** 16: 709)
- (2) Apothecia stalked
 - (a) Stalk narrow, cylindric, mealy-rough, almost hairy
Macropodia R. 984, 8: 158
 - (b) Stalk mostly short and wide, not mealy-rough
 - x. Stalk large and thick, deeply furrowed
Phleboscypus R. 981, 18: 13
(**Acetabula**)
 - y. Stalk even or slightly furrowed
 - (x) Apothecia persistently cup-shaped
Geopyxis R. 971, 8: 63
 - (y) Apothecia finally open and flat
Discina R. 976, 8: 99
- II. Asci blue with iodine
 - 1. Apothecia cleft on one side, ear-like ***Iotidea** R. 1028
 - 2. Apothecia not ear-like
 - a. Spores globose **Plicariella** R. 993
 - b. Spores elliptic to fusoid
 - (1) Apothecia sessile
 - (a) Apothecia with a milky juice **Galactinia** 8: 106
 - (b) Apothecia without milky juice
 - x. Apothecia not on a subicle
 - (x) Apothecia leathery, black **Urnula** R. 999, 8: 548
 - (y) Apothecia fleshy, not black
 - m. Apothecia on the surface of the ground
Plicaria R. 1000
(**Pustularia** in part)
 - n. Apothecia large, sunken, lobed
Peziza R. 1019, 8: 73 and 511
(**Pustularia** in part)
 - y. Apothecia on a subicle **Melachroia** R. 997
 - (2) Apothecia with a long, slender stalk
Tarzettia R. 1021

Phaeosporae

Spores dark, 1-celled, globose to oblong

- I. Spores globose **Phaeopezia** 8: 471, R. 995
- II. Spores elliptic
 - 1. Apothecia sessile **Aleurina** 18: 88
 - 2. Apothecia stalked ***Podaleuris** 18: 88

Subfamily Scutelliniae

Apothecia setose or hairy

Hyalosporae

Spores hyaline, 1-celled, globose to fusoid

- I. Spores globose

- I. Spores smooth
 - a. Cup dark or black, more or less strigose at base
Pseudoplectania R. 1039, 8: 165
 - b. Cup bright-colored, hairy or setose
Sphaerospora R. 1037, 8: 188
- 2. Spores warted or reticulate; cups white-hairy
Pyronemella R. 1038, 8: 194
- II. Spores elliptic to fusoid
 - 1. Spores rostrate at base
Puttemansia 18: 98
 - 2. Spores muticate
 - a. Apothecia sunken in the ground, opening by lobes
Sepultaria R. 1075, 8: 166
 - b. Apothecia superficial
 - (1) Apothecia sessile
 - (a) Apothecia dark-hairy or ciliate
 - x. Apothecia uniformly dark-hairy
Pelodiscus 16: 1147, 18: 35
 - y. Apothecia also with long cilia at the margin
 - (x) Paraphyses clavulate, blunt
Scutellinia R. 1042, 8: 173
(*Lachnea*)
 - (y) Paraphyses equal, brown, pointed
Desmazierella R. 1041, 8: 386
 - (b) Apothecia bright-hairy or ciliate
 - x. Apothecia uniformly bright-hairy
**Leucopezis*
 - y. Apothecia with marginal cilia also
Neottiopezis 8: 190, R. 1068
 - (2) Apothecia stalked
 - (a) Apothecia dark or black
 - x. Stalk long, slender, mealy
Macropodia R. 984, 8: 158
 - y. Stalk short, thick with brown hairs and rhizoids
Plectania 8: 163, R. 1070
 - (b) Apothecia and hairs bright-colored
Sarcoscypha R. 1070, 8: 153
(incl. *Trichoscypha* 8: 160, *Pilocratera* 18: 31)

Phaeosporae

Spores hyaline, 1-celled, globose to fusoid

- I. Apothecia with a cylindric verrucose stalk
Phaeomacropus 16: 740
- II. Apothecia sessile
**Trichaleuris* 18: 89

Family 47. HELVELLACEAE

REHM 1134

Apothecia typically terrestrial, and stalked, sometimes sessile, club-shaped, conical or saddle-shaped, rarely flat, mostly smooth, fleshy, cartilaginous or rarely gelatinous; usually large forms.

Subfamily Rhizinae

Apothecia sessile, flat, arched or irregularly globose

- I. Spores globose Sphaerosoma R. 1140, 8: 56
- II. Spores elliptic or fusoid
 - 1. Spores elliptic, rounded at ends Psilopezia R. 1137, 8: 152
(incl. Peltidium 18: 11)
 - 2. Spores fusoid, pointed at the thickened ends Rhizina R. 1138, 8: 57

Subfamily Helvellae

Apothecia stalked, cap- or saddle-shaped, or columnar

- I. Hymenium ridged in both directions
 - 1. Ridged cap stalked Morchella R. 1200, 8: 8
 - 2. Ridged cap sessile Underwoodia 10: 1
- II. Hymenium smooth, convolute or ridged longitudinally
 - 1. Hymenium saddle-like, more or less lobed Helvella R. 1179, 8: 17
 - 2. Hymenium globoid, convolute Gyromitra R. 1189, 8: 15
 - 3. Hymenium cap- or bell-shaped, smooth or ridged Verpa R. 1195, 8: 29

Subfamily Geoglossae

Apothecia stalked, clavate or capitate

- I. Hymenium distinct from stem, disciform or capitate
 - 1. Spores 1-celled *Haplocybe R. 1168
(incl. Moellerodiscus 18: 8)
 - 2. Spores 2-4-celled
 - a. Apothecia gelatinous Leotia R. 1164, 8: 609
 - b. Apothecia waxy or fleshy-waxy Cudoniella R. 1166, 8: 41
 - 3. Spores filiform or acicular
 - a. Apothecia fleshy, cap-shaped with involute margin Cudonia R. 1169, 8: 527
(Leotiella 16: 700)
 - b. Apothecia waxy, button-shaped, solid Vibrissea R. 1170, 8: 51
- II. Hymenium club-shaped, not distinct from stem or but slightly so
 - 1. Spores hyaline
 - a. Spores 1-celled
 - (1) Spores globose Neolecta 8: 40
 - (2) Spores elliptic Mitrula R. 1146, 8: 32
(Spragueola 14: 742)
 - b. Spores 2-4-celled, fusoid
 - (1) Hymenium covering the whole club Microglossum R. 1151, 8: 39
 - (2) Hymenium on one side only Hemiglossum 10: 2
 - c. Spores more or less filiform Spathularia R. 1158, 8: 48
(incl. Mitruliopsis 18: 10)
 - 2. Spores brown, clavate or cylindric, many-celled Geoglossum R. 1153, 8: 42

Family 48. ASCOBOLACEAE

REHM 1078

Apothecia superficial, typically fimicole, scutellate to disciform, fleshy or waxy or gelatinous; asci mostly broad and clavate, projecting above the hymenium at maturity.

Subfamily Ascophanae

Spores colorless

- I. Hymenium within an exciple
 - 1. Asci 4- or 8-spored
 - a. Spores globose
 - (1) Asci 4-spored **Boudierella 14: 792**
 - (2) Asci 8-spored **Cubonia 8: 527**
 - b. Spores elliptic to fusoid; asci 8-spored
 - (1) Apothecia smooth **Ascophanus R. 1085, 8: 528**
 - (2) Apothecia hairy or setose
 - (a) Spores smooth **Lasiobolus R. 1096, 8: 536**
 - (b) Spores spiny **Aphanascus 10: 35**
 - 2. Asci 16-many-spored
 - a. Asci many
 - (1) Apothecia fimbriate with delicate hairs; asci 32-spored **Streptotheca 10: 34**
 - (2) Apothecia not hairy; asci 16-many-spored
 - Rhyparobius R. 1099**
 - Thelebolus R. 1106**
 - b. Ascus one **1106**
- II. Hymenium without an exciple; asci many-spored
 - Zukalina R. 1108**

Subfamily Ascobolae

Spores colored

- I. Spores globose **Boudiera R. 1113, 8: 512**
- II. Spores elliptic to fusoid
 - 1. Spores in a gelatinous mass in ascus **Saccobolus R. 1115, 8: 524**
 - 2. Spores free in the ascus
 - a. Apothecia smooth
 - (1) Exciple present, normal **Ascobolus R. 1120, 8: 514**
 - (2) Exciple lacking **Ascodesmis 8: 824**
 - b. Apothecia hairy or ciliate **Dasybolus 11: 421**

Family 49. CORDIERITACEAE

8: 810, 16: 803

Apothecia suberose or corneo-carbonous, superficial, ramose-stipitate, arising at the tips of the branches, finally cup-like and open; asci terete-clavate, 6-8-spored; spores 1- or 2-celled, mostly hyaline.

- I. Spores 1-celled, hyaline; stipe much branched above, horny-carbonous **Cordierites 8: 810**

II. Spores 2-celled; stipe fascicled-ramose, suberose

Acroscyphus 8:811

Order 11. GYMNASCALES

Apothecia imperfect, more or less effuse or obsolete, maculiform, byssoid or dot-like, exciple absent; asci mostly free, often single, 1-many-spored, rarely with paraphyses.

Family 50. EXASCACEAE

8:811, 10:67, 11:435, 14:823, 16:803, 18:196

Asci parallel and crowded, sessile or enlarged at base; parasitic in living plants and deforming the part attacked as a rule.

I. Asci few-spored, usually 8-spored

1. Spores 1-celled, more or less globose *Exascus* 8:816

2. Spores 2-3-septate, oblong *Elsinoe* 16:804

II. Asci many-spored

1. Asci more or less globose *Taphridium* 18:203

2. Asci terete-clavate *Taphrina* 8:812

Family 51. GYMNASCACEAE

8:820, 10:70, 11:437, 14:824, 16:805, 18:194

(incl. Ascoidaceae, Ascocortiaceae, Endomycetaceae, Protomycetaceae)

Asci more or less solitary or grouped in masses of mycelium; for the most part saprophytic.

I. Saprogenous

1. Asci 1-2-spored *Bargellinia* 8:823

2. Asci 3-8-spored

a. Spores globose or nearly so

(1) Spores brown or violet *Amaurascus* 11:438

(2) Spores hyaline or golden

(a) Asci 3-5-spored *Conidiascus* 16:807

(b) Asci 8-spored

x. Asci surrounded by serrate spiral hyphae

Ctenomyces 8:824

y. Asci without serrate spiral hyphae

(x) Asci solitary

m. Asci acrogenous *Eremascus* 8:822

n. Asci intercalary *Oleina* 8:822

(y) Asci grouped or congested in masses

Gymnascus 8:823

(incl. *Arachniotus* 11:438)

b. Spores elliptic, hyaline; asci vertical, clavate

Ascocorticium 10:71

3. Asci many-spored

a. Spores globose

(1) Asci elongate, split at base *Dipodascus* 11:439

(2) Asci terete-clavate, simple at base *Ascoidea* 10:71

- b. Spores elliptic †*Ascodes* 16:807
(*Oscarbrefeldia*)

II. Biogenous

1. Asci 4-8-spored
 - a. Asci 4-spored, solitary; on fungi *Endomyces* 8:821
 - b. Asci 8-spored
 - (1) Spores 1-celled
 - (a) Hyphae of palmiform haustoria; on fungi *Podocapsa* 8:820
 - (b) Hyphae filamentous; on animals *Eidamella* 16:805
 - (2) Spores muriform; on leaves *Nostocotheca* 16:806
2. Asci many-spored
 - a. Mycelium present *Eremothecium* 8:821
 - b. Mycelium none
 - (1) Haustoria present; on fungi **Podocapsium* 8:820
 - (2) Haustoria absent; mostly on flowering plants *Protomyces* 7:319

Family 52. SACCHAROMYCETACEAE

8:916, 11:457, 14:828, 16:818, 18:198

True hyphae lacking, unicellular, propagating by buds; asci spurious?, globose to elliptic, mostly 1-4-spored; growing typically in sugary or starchy liquids or materials.

- I. Cells increasing by fission *Schizosaccharomyces* 18:201
- II. Cells increasing by budding
 1. Spores pileiform or limoniform, costate *Willia* 18:198
 2. Spores globose to irregular
 - a. Vegetative cells conjugating *Zygosaccharomyces* 18:198
 - b. Vegetative cells normal *Saccharomyces* 18:198

Order 12. TUBERALES

Ascoma or apothecium typically more or less globose, and indehiscent, with one to many hollows, locules or veins, fleshy, waxy, leathery or even subcarbonous, saprophytic or parasitic, usually subterranean; asci present, 1-many-spored.

Family 53. CYTTARIACEAE

8:4, 16:695, 18:1

Ascomata globose or obovate, firm fleshy, subcorneous when dry, stuffed or hollow, loculiferous at the periphery, producing tubercular swellings on the branches of living trees; locules globose, large, dehiscing by lobes, filled with asci and paraphyses; asci cylindric 8-spored; spores hyaline.

- I. Ascoma globose or obovate; all locules bearing asci *Cyttaria* 8:4

- II. Ascoma turbinate, fenestrate below; asci on a definite disk

Rickiella 18: 1

Family 54. PHYMATOSPHAERIACEAE
(incl. MYRIANGIACEAE)

8: 843, 11: 440, 16: 799, 18: 191

Ascomata verruciform, small, waxy, membranous or subcarbonous, superficial, densely loculiferous within; locules with a single ascus, indehiscent; asci globose or short clavate, 8-spored.

Hyalosporae

Spores hyaline, 1-celled, ovoid to elliptic

- I. Ascomata globose-depressed, membranous *Phillipsiella* 8: 844

Phaeosporae

Spores dark, 1-celled, elliptic to fusoid

- I. Spores angulose, verrucose; fimicole *Guillermondia* 18: 191

Hyalodidymae

Spores hyaline, 1-septate, elliptic to fusoid

- I. Ascomata dark, globose-depressed *Microphyma* 8: 844

- II. Ascomata bright-colored, applanate *Leptophyma* 8: 844

Hyalophragmiae

Spores hyaline, 2-several-septate, oblong to fusoid

- I. Ascomata elongate, rugose *Eurytheca* 8: 846

- II. Ascomata punctiform to obconic

1. Ascomata punctiform or applanate

- a. Ascomata punctiform; asci clavate *Harknessiella* 8: 845

- b. Ascomata applanate-disciform; asci ovoid to globose

Myriangium 16: 800

(incl. *Myriangella* 18: 192)

2. Ascomata hemispheric or obconic; asci globose

Molleriella 8: 845

Phaeophragmiae

Spores dark, 2-several-septate, oblong to fusoid

- I. Ascomata blood-red, membranous-waxy *Kusanoa* 16: 800

Hyalodictyae

Spores hyaline, muriform

- I. Ascomata bright-colored

1. Ascomata on a radiate subicle

Phymatosphaeria 8: 847

2. Ascomata not on a subicle

Ascomycetella 8: 846

- II. Ascomata dark or black

Trichophyma 18: 194

Phaeodictyae

Spores dark, muriform

- I. Ascomata applanate-tuberculiform, black **Cookella 8:846**

Family 55. ONYGENACEAE

8:861, 10:80, 11:440, 16:807

Ascomata subglobose, sessile or stipitate, membranous, fragile, epizoid; gleba waxy, then pulverulent; asci 8-spored, globose, evanescent; spores continuous, sub-hyaline.

A single genus

Onygena 8:861**Family 56. ELAPHOMYCETACEAE**(incl. **CENOCOCCACEAE**)

8:863, 10:80, 11:441

Ascomata hypogaeal, woody, crustose or carbonous, more or less globose, indehiscent, finally producing a powdery spore mass or gleba; asci 1-8-spored, sometimes spurious.

- I. Gleba interwoven with silky threads; asci normal

Elaphomyces 8:863

- II. Gleba without capillitium; asci spurious, cell-shaped

Cenococcum 8:871**Family 57. TUBERACEAE**(incl. **ENDOAGONACEAE, EOTERFEZIACEAE**)

8:872, 10:80, 11:442, 14:826, 16:808, 18:205

Ascomata hypogaeal, rarely epigaeal or parasitic, fleshy or waxy hardened, more or less globose, indehiscent; gleba never becoming a powdery mass, typically veined or lacunose, rarely continuous; asci 1-8-spored, rarely spurious.

Hyalosporae

Spores hyaline, 1-celled, globose to elliptic

- I. Gleba without veins, but with one or more cavities

1. Asci linear or elongate

- a. Spores verrucose or roughened

- (1) Spores globose

Pseudogenea 16:808

- (2) Spores ovoid to elliptic

Genea 8:873

- b. Spores smooth

- (1) Gleba with a single large cavity

Hydnocystis 8:876

- (2) Gleba convolute lacunose

- (a) Densely lanate; canals not produced to surface

Geopora 8:877

- (b) Not lanate; canals produced to surface

Pseudohydnotria 16:808

2. Asci globose to oblong

- a. Spores roughened or alveolate, globose

- (1) Asci 2-4-spored; spores with recurved spines
Terfeziopsis 16:816
 - (2) Asci 8-spored
 - x. Hollows or canals not reaching the surface
 - (x) Gleba with irregular stellate hollows
Myrmecocystis 16:809
 - (y) Microscopic; gleba central, lax
Lilliputia 16:816
 - y. Hollows or canals reaching the surface
Hydnobolites 8:879
 - b. Spores smooth
 - (1) Gleba of numerous locules; epigaeal, parasitic on fungi
Eoterfezia 18:205
 - (2) Hypogaeal
 - (a) Ascoma brown villous
Phaeangium 11:442
 - (b) Ascoma not villous
Balsamia 8:877
- II. Gleba with veins, solid or also lacunose**
- 1. Veins of two colors; spores globose, smooth
Stephensia 8:880
 - 2. Veins all of one color
 - a. Spores globose, roughened
 - (1) Gleba with distinct veins; asci mostly 2-3-spored
Delastria 8:904
 - (2) Gleba marbled with brown spots; asci 3-4-spored
Piersonia 16:812
 - b. Spores ellipsoid, smooth
 - (1) Spores apiculate at each end, limoniform
Leucangium 8:899
 - (2) Spores not apiculate
 - (a) Asci 8-spored, broadly stipitate
Tirmania 11:444
 - (b) Asci 6-8-spored, not stipitate
Picoa 8:899

Phaeosporae

Spores dark, 1-celled

- I. Gleba without veins; typically with hollows or canals**
 - 1. Spores globose, roughened
 - a. Asci linear or cylindric
 - (1) Gleba with one or more hollows
Gyrocratera 16:815
(incl. **Cryptica 10:82**)
 - (2) Gleba homogeneous, lax
Ruhlandiella 17:241
 - b. Asci broad, oblong
Hydnotrya 8:879
 - 2. Spores ovoid, smooth
Genabea 8:878
- II. Gleba with veins**
 - 1. Veins of two colors
 - a. Some veins white
Pachyphloeus 8:881
 - b. No veins white
Tuber 8:882
 - 2. Veins of one color

- a. Asci elongate; gleba not divided into masses

Choeromyces 8:900

- b. Asci ovate to globose; gleba divided into masses

Terfezia 8:902

Order 13. UREDINALES

Apothecia reduced to a mass of persistent or evanescent asci, waxy, leathery, gelatinous or powdery; parasites.

Family 58. UREDINACEAE

7:528, 9:291, 11:174, 14:269, 16:257, 17:244

Parasitic; apothecia reduced to a mass of asci with fixed spore cells, i. e., teleutospores with 1 or more cells; conidia normally present, produced in cluster cups (aecidia, aecia), sori (uredinia), or spermagonia (pycnia); the asci and conidia may occur on the same host or upon different hosts, or one or the other alone may occur; teleutospores producing a promycelium and sporidioles upon germination.

Amerosporae

Teleutospores 1-celled, colored, rarely hyaline, or absent

I. Teleutospores present

1. Teleutospores hyaline

- a. Teleutospores catenate

Monosporidium 9:297

- b. Teleutospores single

Zaghoulania 17:268

2. Teleutospores colored

- a. Spore mass or sorus horizontal

(1) Teleutospores catenate

- (a) Spores in a pseudoperidium

Dietelia 14:291

- (b) Spores not in a pseudoperidium

Clastopsora 17:263

(2) Teleutospores not catenate

- (a) Uredospores not in a pseudoperidium

- x. Spores half smooth, half roughened

Hemileia 7:585

- y. Spore cells alike smooth or rough

- (x) Teleutospores on a stalk

Uromyces 7:531

- (y) Teleutospores not stalked

- m. Teleutospores connate in a lentiform layer

†Uromyces 14:290

(Schroeteriaster)

- n. Teleutospores not connate

Chaonia 14:290

(b) Uredospores in a pseudoperidium

- x. Teleutospore sorus determinate, black or dark-brown

Melampsora 7:586

(incl. Phacopsora 14:289)

- y. Teleutospore sorus indeterminate, pale or reddish

Melampsorella 7:596

(incl. Hyalopsora 17:268)

- b. Spore mass or sorus with a cylindric columella, more or less vertical, globose to cylindric

- (1) Teleutospores mucose; uredospores lacking
Masseella 14: 292
- (2) Teleutospores not mucose; uredospores present
 - (a) Uredospores in a pseudoperidium
Cronartium 7: 597
 - (b) Uredospores not in a pseudoperidium
Skierka 16: 271

II. Teleutospores absent; pycnia, aecia or uredinia only

- I. Spores in a pseudoperidium or cup
 - a. Spores in pycnia
Aecidiolum 7: 773
 - b. Spores in aecia
 - (1) Aecia cup-shaped, usually dentate or crenate at margin
Aecidium 7: 774
 - (2) Aecia cylindric, margin fimbriate
Roestelia 7: 833
 - (3) Aecia irregular, more or less globose
 - (a) Spores catenate; on conifers
Peridermium 7: 835
 - (b) Spores free; not on conifers
Pericladium 7: 838
- 2. Spores not in a pseudoperidium; uredinia
 - a. Spores single
Uredo 7: 838
 - b. Spores catenate
Caeoma 7: 863

Didymosporae

Teleutospores 2-celled, colored or hyaline

- I. Teleutospores absent; aecia alone present
Aecidiella 14: 389
- II. Teleutospores present
 - I. Sori horizontal
 - a. Teleutospores catenate, in a pseudoperidium
†Didymosira 11: 205
(Puccinosira)
 - b. Teleutospores single
 - (1) Teleutospores not in a pseudoperidium
 - (a) Teleutospores subpenicillate at each end
Dasyscypha 9: 313
 - (b) Teleutospores not penicillate
 - x. Pedicel of spore with a hyaline gelatinous sheath
†Coleoma 9: 313
(Coleopuccinia)
 - y. Pedicel without gelatinous sheath
 - (x) Teleutospores longitudinally 1-septate
Diorchidium 7: 736
 - (y) Teleutospores transversely 1-septate
 - m. Teleutospores with a hyaline integument
Uropyxis 7: 735
 - n. Teleutospores without hyaline integument

(m) Spore cells with germination pores

Puccinia 7: 600

(inc. *Trichopsora*, *Chrysopsora*

11: 206, *Gymnoconia* 14: 360)

(n) Spore cells without germination pores

Leptinia 14: 358

(2) Teleutospores in a pseudoperidium

Schizospora 14: 361

2. Sori vertical

a. Teleutospores confluent into a gelatinous stratum

Gymnosporangium 7: 737

b. Teleutospores closely joined in a columella

(1) Spores catenate

Gambleola 16: 314

(2) Spores not catenate

Didymopsora 16: 315

Phragmosporae

Teleutospores 2-several-septate

I. Teleutospores not in a pseudoperidium

1. Teleutospores transversely septate

a. Teleutospores catenate

†**Phragmostele 16: 321**

b. Teleutospores not catenate

(Pucciniostele)

(1) Uredospores not catenate

(a) Teleutospores cylindric; cells separating with difficulty

Phragmidium 7: 742

(incl. *Phragmopyxis* 14: 361, *Rostrupia*, *Barclayella* 9: 316)

(b) Teleutospores moniliform; cells separating easily

Xenodochus 7: 750

(2) Uredospores catenate, at least at first

(a) Wall of teleutospore thick; promycelium simple with a single sporidiale at apex

Coleosporium 7: 751

(incl. *Stichopsora* 16: 318)

(b) Wall of teleutospore thin; promycelium 3-septate, with a sporidiale at each cell

Chrysomyxa 7: 759

2. Teleutospores longitudinally or obliquely septate

a. Teleutospores developed within the host cells

(1) Uredospores in a pseudoperidium; homoeocious

Thecopsora 7: 764

(2) Uredospores lacking; heteroeocious

Calyptospora 7: 766

b. Teleutospores developed outside the host cells

Pucciniastrum 7: 762

II. Teleutospores in a pseudoperidium

1. Teleutospores catenate, verrucose

Endophyllum 7: 767

2. Teleutospores not catenate, echinulate

Milesia 7: 768

(incl. *Uredinopsis* 17: 269)

Dictyosporae

Teleutospores septate in two directions, or muriform

I. Teleutospores more or less radiately 3-septate

Triphragmium 7: 768

(incl. **Hapalophragmium** 16: 1121)

II. Teleutospores radiately 4-many-septate or muriform

Ravenelia 7: 770

(incl. **Sphaerophragmium** 11: 209,
Alveolaria 11: 212, **Hemileiopsis** 16: 269, **Anthomyces** 16: 325,
Pleoravenelia and **Neoravenelia**, 17: 407)

Family 59. USTILAGINACEAE

7: 449, 9: 282, 11: 230, 14: 410, 16: 367, 17: 472

Mycelium growing widely through parts of living plants, chiefly flowers and fruits, finally disappearing, leaving the mass of spores; spores producing upon germination a promycelium upon which sporidioles are borne.

Amerosporae

Spores 1-celled

I. Sori without a fungal involucre

1. Sporidioles typically pleurogenous on the promycelium

a. Spores arising from a compact subgelatinous stroma

Cintractia 7: 480

b. Spores not arising from a compact subgelatinous stroma

Ustilago 7: 451

(incl. **Anthracoidea** 14: 420)

2. Sporidioles many, acrogenous, crowning the promycelium

a. Sori powdery at maturity

(1) Sporidioles many, in a capitulum

Neovossia 16: 375

(2) Sporidioles not in a capitulum

Tilletia 7: 481

b. Sori not powdery at maturity

(1) Spores catenate, then separating

Sirentyloma 14: 425

(2) Spores not catenate

(a) Spores rostrate

Rhamphospora 9: 287

(b) Spores not rostrate

x. In stems and leaves

(x) Sori pustulate, pale or rust-brown

Entyloma 7: 487

(y) Sori explanate, widely expanded, black

Melanotaenium 7: 496

y. In roots

(x) Spores conglobate in spheroid cysts

Oedomycetes 11: 234

(y) Spores not conglobate

Entorrhiza 7: 497

z. In ovaries

†*Ustilaginula* 7: 498
(*Ustilagopsis*)

II. Sori with a fungal involucre

1. Spores in a powdery mass *Sphacelotheca* 7: 499
2. Spores in a hard black crust *Melanopsichium* 17: 484

Didymosporae

Spores united by twos or 2-celled

I. Spore-bearing hyphae tubular, enclosed in a stroma

Mycosyrinx 17: 484

II. Spore-bearing hyphae not in a stroma

1. Spores joined laterally by a narrow isthmus; sporidioles pleurogenous
Schizonella 7: 500
2. Spores joined horizontally and broadly; sporidioles acrogenous
Schroeteria 7: 500

Dictyosporae

Spores closely joined in masses, the latter appearing to be many-celled spores

I. Spores or cells of each mass alike

1. Sporidioles pleurogenous or acrogenous; usually not foliicole
 - a. Promycelium simple *Tolyposporium* 7: 501
 - b. Promycelium branched *Tolyposporella* 14: 427
2. Sporidioles acrogenous, typically foliicole
 - a. Sporidioles numerous
 - (1) Spore masses covered by a layer of sterile cells
Doassansia 7: 502
(incl. *Cornuella*, *Burrillia* 11: 236)
 - (2) Spore masses without a sterile layer
Tubercinia 7: 507
 - b. Sporidioles solitary; sori reddish, usually fructicole
Thecophora 7: 507
3. Sporidioles unknown; sori mostly very black
Sorosporium 7: 511
(incl. *Poecilosporium* 16: 380)

II. Spores or cells of two kinds in each mass, central few large, peripheral many, small

1. Sori of many sacks containing spore masses
Polysaccopsis 16: 381
2. Sori without sacks
Urocystis 7: 515

Class 5. BASIDIOMYCETES

Spores produced on basidia, not inclosed in asci.

Order 14. AGARICALES (HYMENOMYCETES)

Basidia exposed on an even or modified hymenium, the latter usually in the form of gills, pores or teeth.

Family 60. TREMELLACEAE

6: 760, 9: 257, 11: 142, 14: 244, 16: 215, 17: 203

Pileus typically gelatinous and homogeneous, horny when dry, reviving when wet, sometimes waxy or leathery but then with divided basidia; hymenium typically amphigenous or superior, smooth or somewhat convolute; basidia globose to terete, transversely or longitudinally divided, or in one subfamily merely terete-clavate and furcate, 1-4-sterigmate; spores globose to reniform and oblong, continuous or septate, producing sporidioles on germination; conidia often present with the spores. Some gelatinous forms included in the following families on account of the character of the hymenium seem to belong properly in this family.

Subfamily Auriculariae

Basidia transversely septate, elongate or fusoid

- I. Pileus, or at least the hymenium, gelatinous
 - i. Entire pileus gelatinous
 - a. Pileus verruciform or effuse
 - (1) Basidia mixed with paraphyses **Mytilopsis** 14: 246
 - (2) Basidia without paraphyses
 - (a) Spores not producing sporidioles on germination
Platyglora 6: 771
 - (b) Spores producing sporidioles
Helicogloea 11: 145
 - b. Pileus disciform, cupulate or columnar
 - (1) Pileus erect, filiform, columnar **Eucronartium** 17: 211
 - (2) Pileus not columnar, disciform or cupulate
 - (a) Basidia without sterigmata **Auriculariella** 6: 407
 - (b) Basidia with sterigmata
 - x. Basidia 2-sterigmate; pileus applanate
Phlebophora 16: 215
 - y. Basidia 3-4-sterigmate; pileus pezizoid
†Collopezis 16: 216
(**Tjibodasia**)
 2. Pileus coriaceous or membranous, hymenium gelatinous
 - a. Pileus coriaceous; hymenium reticulate-costate
Auricularia 6: 762
 - b. Pileus membranous; hymenium smooth or plicate
Hirneola 6: 764
- II. Pileus waxy, crust-like or byssoid
 - i. Pileus waxy or crust-like
 - a. Pileus very minute, disciform, on a pedicel
Pilacrella 14: 246
 - b. Pileus membranous, incrusting
Jola 14: 245
 2. Pileus byssoid
 - a. Basidia without a sac near the base **Stypinella** 14: 244
 - b. Basidia with a sack near the base **Saccoblastia** 14: 244

Subfamily Tremellae

Basidia longitudinally 4-divided, or cruciate, globose or ovoid

- I. Spores alone present, i. e., homosporous

1. Pileus waxy or byssoid
 - a. Pileus waxy, scarcely gelatinous
 - (1) Pileus effuse **Protomerulius 11: 142**
 - (2) Pileus cupulate or concave **Hirneolina 17: 208**
 - b. Pileus byssoid **Stypella 14: 246**
 2. Pileus gelatinous
 - a. Pileus covered with sterile setae, effuse **Heterochaete 14: 247**
 - b. Pileus without sterile setae
 - (1) Pileus erect, clavate, columnar or spatulate
 - (a) Pileus clavate, simple or branched **Clavariopsis 16: 219**
(incl. **Hyaloria 14: 252**)
Gyrocephalus 6: 795
 - (b) Pileus spatulate, large, simple **Gyrocephalus 6: 795**
 - (2) Pileus effuse, globose, cupulate or pulvinate
 - (a) Spores 1-celled
 - x. Pileus cupulate, radicate **Femsjonia 6: 779**
 - y. Pileus pulvinate or effuse
 - (y) Basidia in chains; hymenium not cerebriiform **Sirobasidium 14: 248**
 - (y) Basidia not in chains; hymenium cerebriiform **Tremella 6: 780**
(inc. **Naematelia 6: 792**)
 - (b) Spores 2-4-celled, at least upon germination, reniform
 - x. Spores 2-4-celled, sporidioles allantoid; pileus truncate-cupulate or effuse **Exidia 6: 772**
 - y. Spores 2-celled, sporidioles straight; pileus pulvinate, gyrose **Ulocolla 6: 777**
- II. Spores and conidia present, i. e., heterosporous
 1. Pileus ascending and dendroid **†Collodendrum 17: 208**
(**Tremellodendron**)
 2. Pileus effuse to pulvinate
 - a. Spores on the disk, conidia on the exciple **Craterocolla 6: 778**
 - b. Conidia and spores usually succeeding each other on the same area
 - (1) Pileus cerebriiform, pulvinate or effuse **Tremella 6: 780**
 - (2) Pileus not cerebriiform, crust-like
 - (a) Spores reniform, conidia ovoid **Sebacina 6: 540**
 - (b) Spores ovoid, conidia hamate **Exidiopsis 14: 248**
- Subfamily Dacryomycetae**
Basidia terete-clavate, furcate above
- I. Pileus effuse, pulvinate or globose, typically sessile
 1. Spores septate, at least upon germination
 - a. Pileus gyrose; spores not horseshoe-shaped **Dacryomyces 6: 796**
 - b. Pileus tuberculiform; spores horseshoe-shaped **Delortia 6: 795**

- 2. Spores not septate
 - a. Spores hyaline; pileus more or less effuse, waxy
 - Arrhytidia* 6: 804
 - (incl. *Ceracea* 6: 805)
 - b. Spores colored; pileus subglobose
 - Seismosarca* 9: 260
- II. Pileus cupulate, clavate or foliose, typically stalked
 - 1. Pileus irregularly cup-shaped, usually stipitate
 - a. Pileus gelatinous or cartilaginous, cupulate
 - Guepinia* 6: 805
 - b. Pileus leathery, hymenium gelatinous, cupulate-disciform
 - Ditiola* 6: 813
 - 2. Pileus erect, foliose-lobed
 - †*Tremellastrum* 17: 193
 - (*Tremellopsis*)
 - 3. Pileus capitate to lanceolate, stipitate
 - a. Pileus capitate, head inflated, corrugate; stipe hollow
 - (1) Homosporous
 - Collyria* 6: 811
 - (2) Heterosporous
 - Dacryopsis* 11: 149
 - b. Pileus clavate, club plicate
 - Dacryomitra* 6: 811
 - c. Pileus lanceolate, hanging
 - Myxomycidium* 16: 220

Family 61. CLAVARIACEAE

6: 690, 9: 247, 11: 134, 14: 235, 16: 203, 18: 193

Hymenium not discrete from the hymenophore, amphigenous; pileus more or less clavate or coralloid, subcarnose or leathery, simple or branched.

- I. Pileus with many crowded, leaf-like branches
 - Sparassis* 6: 690
- II. Branches not leaf-like
 - 1. Pileus fleshy
 - a. Branches fibrous-splitting
 - Acurtis* 6: 691
 - b. Branches not splitting
 - Clavaria* 6: 692
 - (incl. *Phaeoclavulina* 14: 238)
 - 2. Pileus leathery, rarely subgelatinous
 - a. Pileus somewhat gelatinous
 - (1) Pileus capitate; cap hollow, inflated
 - Baumannella* 14: 244
 - (2) Pileus clavate or coralloid
 - Calocera* 6: 732
 - b. Pileus leathery
 - (1) Pileus tomentose
 - Lachnocladium* 6: 738
 - (2) Pileus not tomentose
 - (a) Pileus terete or compressed, dry, cartilaginous
 - Pterula* 6: 740
 - (incl. *Phaeopterula* 17: 201)
 - (b) Pileus simple, filiform or capitate
 - Hirsutella* 11: 140
 - x. Pileus capitate, inflated
 - Physalacria* 6: 759
 - y. Pileus more or less filiform

- (x) Pileus clavulate with filiform stipe
Typhula 6: 743
- (y) Pileus linear or subclavate; stipe short or none
Pistillaria 6: 752

Family 62. THELEPHORACEAE

6: 513, 9: 218, 11: 115, 14: 212, 16: 181, 18: 160

Hymenium inferior or amphigenous, leathery, waxy or membranous, smooth, i. e., without spines, pores, etc., sometimes somewhat ridged, or cracked; spores various.

I. Not parasitic on algae

1. Pileus more or less gelatinous

a. Pileus effuse

- (1) Spores hyaline **Cerocorticium 16: 196**
- (2) Spores olivaceous **Aldridgea 11: 129**

b. Pileus convex to discoid

Discocyphella 16: 202

2. Pileus not gelatinous

a. Hymenium somewhat ridged or roughened

- (1) Hymenium subcarnose, infundibuliform, costate

Craterellus 6: 514

- (2) Hymenium leathery

- (a) Hymenium woody, with radiating ridges, warty-roughened

Cladoderris 6: 547

- (b) Hymenium similar, but with fan-like ridges

Beccariella 6: 550

b. Hymenium smooth, or absent

- (1) Hymenium present, smooth

- (a) Hymenium without cystidia

- x. Pileus urn-shaped, stipitate **Hypolyssus 6: 521**

- y. Pileus typically crateriform to dimidiate

- (x) Pileus with distinct intermediate stratum

Stereum 6: 551

- (y) Pileus homogeneous or nearly so

- m. Pileus vertical, beautifully convolute, mitriform

Skepperia 6: 603

- n. Pileus not convolute

- (m) Basidia not transeptate **Thelephora 6: 521**

(incl. **Friesula 6: 685**)

- (n) Basidia transeptate **Septobasidium 11: 118**

- z. Pileus resupinate, effuse, rarely cupulate

- (x) Pileus not cupulate

- m. Hymenium waxy

- (m) Spores large, citriform **Michenera 6: 652**

- (n) Spores medium, not citriform

Corticium 6: 603

(incl. **Kneiffia 6: 510**)

- n. Hymenium fleshy, spores minute, colored

- (m) Spores smooth **Coniophora 6: 647**
 - (n) Spores angular or aculeate **Prillieuxia 14: 225**
 - (y) Pileus cupulate or cylindric
 - m. Pileus cupulate **Cyphella 6: 667**
 - n. Pileus terete to cylindric **Solenia 6: 424**
 - (b) Hymenium with cystidia
 - x. Cystidia simple
 - (x) Cystidia hyaline **Peniophora 6: 640**
(incl. *Coniophorella* 17: 183)
 - (y) Cystidia colored **Hymenochaete 6: 588**
(incl. *Lloydia* 16: 1116)
 - y. Cystidia septate **Bonia 11: 123**
 - (2) Hymenium absent, or more or less cobwebby
 - (a) Biogenous
 - x. Hymenium endophytic **Endobasidium 17: 190**
 - y. Hymenium erumpent
 - (x) Basidia circinate **Helicobasidium 6: 666**
 - (y) Basidia not circinate
 - m. Spores globose; on galls **Urobasidium 11: 131**
 - n. Spores cylindric; on roots ***Chrysobasidium 11: 131**
(*Aureobasidium*)
 - o. Spores oblong; on leaves **Exobasidium 6: 664**
 - (b) Saprogenous
 - x. Spores septate, fuscous **Heterobasidium 9: 237**
 - y. Spores 1-celled, hyaline
 - (x) Brown stellate hyphae present **Asterostroma 9: 236**
 - (y) Brown stellate hyphae absent
 - m. Basidia 4-spored **Hypochnus 6: 653**
 - n. Basidia 2-spored **Matruchotia 11: 118**
- (Cfr. Tulasnellaceae 14: 234)**

II. Parasitic on algae

- 1. Algae *Chroococcus* **Cora 6: 685**
- 2. Algae *Scytonema* **Rhipidonema 6: 687**

(Zahlbruckner 237)

Family 63. HYDNACEAE

6: 429, 9: 208, 11: 106, 14: 201, 16: 174, 18: 147

Pileus cap-shaped to resupinate, fleshy, gelatinous, woody or leathery; hymenium consisting of spines, teeth, or granules, rarely somewhat pore-like; spores various.

I. Pileus more or less gelatinous

- 1. Gelatinous, stalked or dimidiate; with teeth **Tremellodon 6: 479**
- 2. Waxy-gelatinous, resupinate, with granules **Grandiniella 14: 208**

II. Pileus fleshy, woody or leathery

- 1. Hymenium of more or less subulate teeth or spines

- a. Pileus present
 - (1) Perennial; woody †Hydnophysa 16: 177
(Hydnofomes)
 - (2) Not perennial
 - (a) Pileus clavaria-like Hericum 6: 478
 - (b) Pileus not clavaria-like
 - x. Teeth free; mostly carnose
 - (x) Pileus typically stalked Hydnum 6: 430
(incl. Echinodontium 16: 176)
 - (y) Pileus horizontal Sistotrema 6: 480
 - y. Teeth connected at base; coriaceous
 - (x) Cystidia lacking Irpex 6: 482
 - (y) Cystidia present
 - m. Cystidia subulate Asterodon 11: 111
 - n. Cystidia stellate Hydnochaete 14: 211
- b. Pileus lacking
 - (1) Teeth on a membranous subicle Caldesiella 6: 477
 - (2) Teeth without a subicle Mucronella 6: 512
- 2. Hymenium of granules, warts or folds
 - a. Hymenium of granules or warts
 - (1) Hymenium with penicillate-multifid warts Odontia 6: 506
 - (2) Hymenium with simple granules or warts
 - (a) Hymenium porose-reticulate, granular Grammothele 6: 505
 - (b) Hymenium with difform, obtuse cylindric warts Radulum 6: 493
(incl. Phaeoradulum 16: 179)
 - (c) Hymenium with globose hollowed granules Grandinia 6: 500
 - b. Hymenium with folds or laminae
 - (1) Hymenium with fold-like crests
 - (a) Crests with edge entire Phlebia 6: 497
 - (b) Crests with edge incised Lopharia 6: 500
 - (2) Hymenium with anastomosing radiate laminae Thwaitesiella 11: 112

Family 64. POLYPORACEAE

6: 1, 9: 150, 11: 79, 14: 164, 16: 138, 17: 95

Pileus cap-shaped, shelf-like, or resupinate, very rarely volvate or annulate, fleshy, leathery or woody, rarely gelatinous; hymenium consisting of pores, very rarely somewhat lamellar; spores typically 1-celled, hyaline or colored.

I. Pileus fleshy, putrescent, or gelatinous

1. Pileus fleshy

a. Stipe volvate or annulate

(1) Stipe volvate

†Boletium 14: 164
(Volvoboletus)

- (2) Stipe annulate **Boletopsis 14: 164**
- b. Stipe not volvate or annulate
 - (1) Stipe central, tubes usually not discrete from each other
 - (a) Spores cylindric, minute **†Bactroboletus 16: 142**
(**Filoboletus**)
 - (b) Spores globose to fusoid
 - x. Pileus and stipe beautifully squarrose-scaly
Strobilomyces 6: 49
 - y. Pileus and stipe not squarrose-scaly
 - (x) Layer of tubes separating readily from the hymenophore
Boletus 6: 2
(incl. **Suillus**, **Tylopilus 16: 142**)
 - (y) Layer of tubes not separating readily from the hymenophore
 - m. Tubes not discrete from each other
 - (m) Tubes radiate; hymenophore mucronate
Boletinus 6: 51
 - (n) Tubes sinuose or gyrose; hymenophore smooth
Gyrodon 6: 51
 - n. Tubes discrete from each other
Fistulinella 17: 101
 - (2) Stipe lateral; tubes discrete from each other
Fistulina 6: 54
- 2. Pileus gelatinous
 - a. Stalked; spores brown
 - (1) Pileus single **Rodwaya 16: 172**
 - (2) Pileus many, superimposed on the stipe
Mycodendrum 9: 206
Laschia 6: 404
 - b. Mostly sessile; spores hyaline
- II. Pileus leathery, corky or woody, rarely tough-fleshy
 - 1. Tubes gelatinous **Gloeoporus 6: 403**
 - 2. Tubes not gelatinous
 - a. Hymenium covered by a volva-like membrane
Cryptoporus 17: 125
 - b. Hymenium not volvate
 - (1) Tubes in several layers; perennial, woody
Fomes 6: 150
 - (2) Tubes not stratified in layers
 - (a) Tubes typically pore-like
 - x. Tube layer distinct but not separable from the hymenophore; tough-fleshy to leathery
 - (x) Pileus thick, tough-fleshy, stalked or sessile
Polyporus 6: 55
(incl. **Laccocephalum 11: 87**)
 - (y) Pileus thin, coriaceous or membranous
 - m. Pileus stipitate to dimidiate
 - (m) Tubes not spiny inside **Polystictis 6: 208**
 - (n) Tubes spiny inside **Mucronoporus 9: 188**
 - n. Pileus resupinate **Poria 6: 292**

- y. Tube layer not distinct from hymenophore; tubes often unequally sunken
 - (x) Pileus suberose; typically sessile to resupinate
 - m. Tubes subrotund **Trametes 6: 334**
(incl. *Sclerodopsis* 9: 194)
 - n. Tubes not round, or of two forms
 - (m) Tubes of two forms, one normal, the other loculiform, enclosed **Myriadoporus 6: 384**
 - (n) Tubes alike, superficial
 - r. Tubes hexagonal **Hexagonia 6: 356**
 - s. Tubes sinuose-labyrinthine, elongate **Daedalea 6: 370**
 - (y) Pileus leathery, membranous or waxy; sessile
 - m. Tubes immersed in discrete warts; resupinate **Porothelium 6: 421**
 - n. Tubes not immersed in warts
 - (m) Tubes with a papilla in the center **Theloporus 6: 421**
 - (n) Tubes reticulate-gyrose, not papillate **Merulius 6: 411**
(incl. *Poroptycha* 9: 206)
 - (b) Tubes lamella-like (see *Daedalea* also)
 - x. Tubes of many little laminae **Bresadolina 6: 388**
 - y. Tubes lamellose, in radiating series **Favolus 6: 390**
 - z. Tubes really concentric lamellae **Cyclomyces 6: 389**

Family 65. AGARICACEAE

Pileus typically cap-shaped and stalked, rarely sessile and the hymenium above, fleshy to corky; pileus sometimes enclosed in a cap veil which persists at the base of the stipe as a volva; hymenium consisting of radiating lamellae or gills, often protected by a gill veil which remains on the stipe as a ring; gills covered with basidia, bearing typically 4 sterigmata and spores; spores typically 1-celled, hyaline or colored.

Leucosporae

5:8, 9:1, 11:1, 14:63, 16:1, 18:1

Spores colorless, or very dilutely colored even in spore prints, globose to fusoid, smooth or rough

- I. Edge of the gills entire, not canaliculate or split
 - 1. Fleshy, putrescent, not reviving when wet
 - a. Edge of the gills acute, not fold-like
 - (1) Trama of the pileus not vesiculose; spores typically smooth
 - (a) Gills more or less fleshy, readily separable into two layers
 - x. Stipe central or nearly so
 - (x) Hymenophore discrete from the fleshy stipe
 - m. Stipe volvate

- (m) Stipe annulate **Amanita 5:8**
- (n) Stipe not annulate **Amanitopsis 5:20**
- n. Stipe not volvate
 - (m) Stipe annulate **Lepiota 5:27**
 - (n) Stipe not annulate **Schulzeria 5:72**
- (y) Hymenophore homogeneous and confluent with the fleshy or fibrous-elastic stipe
 - m. Stipe annulate, without a volva **Armillaria 5:73**
 - n. Stipe not annulate or volvate
 - (m) Gills adnate or sinuate, not decurrent **Tricholoma 5:87**
 - (n) Gills typically decurrent **Clitocybe 5:141**
 - (z) Hymenophore confluent with the cartilaginous stipe but heterogeneous from it
 - m. Gills not decurrent
 - (m) Cap very thin, diaphanous **Hiatala 5:305**
 - (n) Cap not diaphanous
 - r. Margin of the young cap turned in **Collybia 5:200**
 - s. Margin of the young cap straight **Mycena 5:251**
(incl. *Eomycenella* 17:21)
 - n. Gills decurrent; cap umbilicate **Omphalia 5:308**
 - y. Stipe excentric or none **Pleurotus 5:339**
 - (b) Gills waxy rather than fleshy, splitting with difficulty **Hygrophorus 5:387**
 - (2) Trama of cap more or less vesiculose; spores globose, spiny
 - (a) Gills with milky, white or bright-colored sap **Lactarius 5:423**
(incl. *Lactariopsis* 17:30)
 - (b) Gills with clear sap, if any **Russula 5:453**
 - b. Edge of gills obtuse or fold-like
 - (1) Gills decurrent, dichotomous, somewhat waxy **Cantharellus 5:482**
 - (2) Gills not decurrent
 - (a) Gills somewhat broad, obtuse **Nyctalis 5:499**
 - (b) Gills thin or obsolete
 - x. Gills thin
 - (x) Gills vein-like, fleshy **Arrhenia 5:498**
(incl. *Campanella* 14:100, *Rim-bachia* 11:32)
 - (y) Gills of two sorts, gelatinous **Stylobates 5:502**

- y. Gills obsolete **Cymatella 16: 49**
2. Fleishy-leathery, leathery, corky or woody, persistent, reviving when wet
- a. Fleishy-leathery or gelatinous-leathery
- (1) Gills distinct
- (a) Stipe discrete from the hymenophore
- x. Cap fleshy and tough or thin and leathery **Marasmius 5: 503**
(incl. **Marasmiopsis 14: 101**)
- y. Cap gelatinous-leathery **Heliomyces 5: 569**
- (b) Stipe and hymenophore continuous
- x. Edge of gills acute
- (x) Edge serrate **Lentinus 5: 571**
(incl. **Lentodium 14: 121**, **Lentodopsis 17: 47**)
- (y) Edge entire **Panus 5: 614**
- y. Edge of gills obtuse, gills dichotomous **Xerotus 5: 630**
- (2) Gills fold-like, edges canaliculate or crisp **Trogia 5: 635**
- b. Corky
- (1) Gills distinct
- (a) Gills tomentose **Tilotus 5: 652**
- (b) Gills smooth **Lenzites 5: 637**
- (2) Gills line-like, parallel, flexuous **Hymenogramme 5: 652**
- II. Edge of gill split or appendiculate
- I. Fleishy
- a. Stipe central; edge of gills split **Oudemansiella 5: 653**
- b. Stipe lateral; edge with appendages **Pterophyllus 5: 654**
2. Membranous or coriaceous
- a. Membranous; stipe central; gills split into flexuous fragments **Rhacophyllus 5: 654**
- b. Coriaceous; stipe none or lateral; edge split and revolute **Schizophyllum 5: 654**
- Rhodosporae**
- 5: 656, 9: 82, 11: 43, 14: 124, 16: 69, 18: 52
- Spores rosy, salmon-colored or rosy-rust-colored in spore prints, paler under the microscope
- I. Stipe central
1. Hymenophore discrete from the stipe
- a. Stipe volvate at base
- (1) Stipe annulate also **Metraria 9: 82**
- (2) Stipe not annulate **Volvaria 5: 656**
- b. Stipe not volvate
- (1) Stipe annulate **Annularia 5: 663**
- (2) Stipe not annulate
- (a) Fleishy; gills free **Pluteus 5: 665**

- (b) Tough; gills adnexed **Schinzinia 11: 44**
- 2. Hymenophore homogeneous and confluent with the stipe
 - a. Gills decurrent
 - (1) Stipe fleshy-fibrous **Clitopilus 5: 698**
 - (2) Stipe cartilaginous **Eccilia 5: 729**
 - b. Gills adnexed, sinuate or free
 - (1) Stipe fleshy-fibrous; gills sinuate **Entoloma 5: 679**
 - (2) Stipe cartilaginous; gills not sinuate
 - (a) Cap convex; margin at first inflexed **Leptonia 5: 706**
 - (b) Cap campanulate; margin straight from the first **Nolanea 5: 716**
- 3. Hymenophore continuous with the cartilaginous stipe, but different from it; volvate **Volvariella 16: 70**
- II. Stipe excentric or none; lignicole **Claudopus 5: 733**

Ochrosporae

5: 735, 9: 90, 11: 48, 14: 131, 16: 83, 18: 62

Spores ochraceous or more or less rust-colored

- I. Gills not separating readily or naturally from hymenophore
 - 1. Gill veil not cobwebby
 - a. Stipe central
 - (1) Stipe volvate or annulate
 - (a) Stipe volvate **Locellina 5: 761**
 - (b) Stipe annulate **Pholidota 5: 736**
(incl. **Pholiotella 9: 90**)
 - (2) Stipe not volvate or annulate
 - (a) Gills not deliquescing
 - x. Stipe fleshy
 - (x) Gills adnate or decurrent **Flammula 5: 809**
 - (y) Gills mostly sinuate
 - m. Cap fibrillose, silky or scaly **Inocybe 5: 762**
 - n. Cap smooth, more or less viscid **Hebeloma 5: 791**
 - y. Stipe cartilaginous
 - (x) Gills decurrent **Tubaria 5: 872**
 - (y) Gills not decurrent
 - m. Margin of cap inflexed at first **Naucoria 5: 828**
 - n. Margin of cap straight
 - (m) Stipe discrete from hymenophore; gills free **Pluteolus 5: 859**
 - (n) Stipe homogeneous with hymenophore **Galera 5: 860**
 - (b) Gills deliquescing **Bolbitius 5: 1073**
 - b. Stipe excentric or none; lignicole **Crepidotus 5: 876**

2. Gill veil cobwebby, hanging curtain-like from the margin, often disappearing completely with age **Cortinarius 5: 889**
- II. Gills separating readily from the hymenophore; margin of cap persistently involute **Paxillus 5: 983**

Melanosporae

5: 991, 9: 136, 11: 69, 14: 149, 16: 112, 18: 82

Spores purple, dark-purple to black

- I. Spores purple or dark-purple
1. Hymenophore discrete from stipe
- a. Stipe volvate at base
- (1) Stipe annulate **Chitoniella 14: 149**
- (2) Stipe not annulate **†Chitonis 5: 992**
(Chitonia, Clarkeinda)
- b. Stipe not volvate
- (1) Stipe annulate **Agaricus 5: 993**
- (2) Stipe not annulate; gills free **Pilosace 5: 1010**
2. Hymenophore continuous with stipe
- a. Stipe annulate **Stropharia 5: 1012**
- b. Stipe not annulate; margin sometimes cortinate
- (1) Margin of cap cortinate; rarely subannulate **Hypholoma 5: 1027**
- (2) Margin not cortinate
- (a) Gills decurrent **Deconica 5: 1058**
- (b) Gills not decurrent
- x. Margin of cap inflexed at first **Psilocybe 5: 1043**
- y. Margin of cap straight **Psathyra 5: 1060**
- II. Spores dark or black, not purple
1. Gills deliquescing **Coprinus 5: 1078**
2. Gills not deliquescing
- a. Gills united above to the hymenophore
- (1) Cap fleshy, fleshy-waxy or membranous
- (a) Gills waxy; spores globose, spiny **Phaeohygrocybe 17: 81**
- (b) Gills not waxy
- x. Margin of cap with a viscid cobwebby cortina **Phaeolimacium 16: 110**
- y. Margin of cap not viscid-cortinate
- (x) Spores globose to elliptic
- m. Stipe annulate; variegated gills exceeding the margin **Anellaria 5: 1125**
- n. Stipe not annulate
- (m) Cap fleshy, not striate; variegated gills exceeding the margin **Panaeolus 5: 1118**
- (n) Cap membranous, striate; uniform gills not exceeding the margin **Psathyrella 5: 1126**

- (y) Spores elongate, fusoid; gills decurrent
Gomphidius 5: 1137
- (2) Cap leathery-horny; spores minute, globose
Anthracoephyllum 5: 1139
- b. Gills free above, not united to the hymenophore; stipe dilated into a lamellar disk above
Montagnites 5: 1140

Order 15. LYCOPERDALES (GASTEROMYCETES)

Typically terrestrial, sometimes lignicole or hypogaeous, fleshy, leathery or membranous; spores borne on basidia, in a receptacle or a peridium, continuous, hyaline or colored.

Family 66. PHALLACEAE

7: 2, 9: 262, 11: 153, 14: 254, 16: 224, 17: 212

Receptacle arising from a volva, bearing outside or inside the sporiferous pulp or gleba, stalk-like, pileiform, or sessile and more or less clathrate

I. Gleba covering the outside of receptacle; receptacle stalk-like, pileate or appendaged

1. Receptacle pileate; gleba on outer surface of pileus
 - a. Stalk with an appendage below the pileus
 - (1) Appendage net-like; volva smooth **Dictyophora 7: 3**
 - (2) Appendage collar-like; volva acuminate
Echinophallus 16: 226
 - b. Stalk without an appendage
 - (1) Upper part of volva remaining with pileus, and enclosing the gleba
Cryptophallus 14: 254
 - (2) Upper part of volva not enclosing gleba at maturity
Ithyphallus 7: 8
(incl. Alboffiella 16: 227)
2. Receptacle without hanging pileus; gleba borne directly on the apex of the stalk-like receptacle
 - a. Receptacle without appendages
 - (1) Receptacle floccose **Floccimutinus 14: 255**
 - (2) Receptacle not floccose **Mutinus 7: 12**
(incl. Aporophallus Itajahya 11: 153, Jansia 16: 226)
 - b. Receptacle or gleba with coralloid processes
Kalchbrennera 7: 14

II. Gleba on the inside of the hollow receptacle, which is clathrate or lobed

1. Receptacle hollow and clathrate, or formed of a few vertical branches joined at the apex
 - a. Receptacle stalked
 - (1) Gleba dimorphous, apex with sterile radiate laminae, lower part with convolute subclathrate lobes **Dictyobole 17: 213**
 - (2) Gleba not dimorphous
 - (a) Receptacle hollow-clathrate, stalked

- x. Openings polygonal **Simblum 7: 16**
 - y. Openings vertically elongate **Colus 7: 21**
 - (b) Receptacle of thin anastomosing branches, stipitiform at base **Clathrella 16: 228**
- b. Receptacle sessile
 - (1) Hollow-clathrate, or of a few united vertical branches **Clathrus 7: 18**
 - (2) Radiately loculate within **Protuberia 11: 155**
- 2. Receptacle divided above into free laciniae or lobes
 - a. Receptacle expanded above into a horizontal border which is laciniate at the margin **Aseroe 7: 25**
 - b. Receptacle divided directly into lobes
 - (1) Lobes distinct from stalk in structure and color
 - (a) Lobes without winged appendages **Lysurus 7: 22**
 - (b) Lobes with membranous winged appendages **Blumenavia 11: 154**
 - (2) Lobes like the stalk in structure and color
 - (a) Receptacle spheric, lobes contiguous **Phallogaster 11: 155**
 - (b) Receptacle elongate or cupulate; lobes more or less spreading
 - x. Lobes sporiferous **Anthurus 7: 23**
 - y. Lobes not sporiferous **Calathiscus 7: 24**

Family 67. LYCOPERDACEAE

7: 48, 9: 266, 11: 157, 14: 257, 16: 230, 17: 217

Epigaeous, rarely hypogaeous or lignicole, peridium usually globose to pyriform, sessile or stipitate, membrano-coriaceous, furnished with a mouth or opening irregularly, enclosing a more or less powdery, often floccose, gleba; spores globose to ellipsoid, hyaline or colored, smooth or rough.

- I. Peridium more or less completely traversed by a continuation of the stipe, i. e., a columella; gleba lamellate or with membranous septa or more or less uniform

Subfamily Podaxae

- 1. Gleba lamellate; capillitium none; peridium turbinate **Gyrophragmium 7: 51**
- 2. Gleba not lamellate, more or less divided by anastomosing septa, or uniform
 - a. Gleba with septa
 - (1) Capillitium none; stipe central, not volvate, short
 - (a) Peridium with broad false radiate lamellae beneath **Elasmomyces 14: 258**
 - (b) Peridium without lamellae beneath **Secotium 7: 51**
 - (2) Capillitium present, filamentous; stipe volvate **Polyplocium 7: 55**
 - b. Gleba without septa or locules; capillitium copious
 - (1) Peridium subsessile; columella free, not touching the apex of the peridium
 - (a) Epigaeous

- x. Columella cup-shaped; exoperidium areolate
Cycloderma 7: 56
- y. Columella obturbinate; exoperidium splitting into lobes
Geasteropsis 17: 229
- (b) Hypogaeous; spores subfusoid **Mesophellia 7: 56**
- (2) Peridium stipitate; columella touching the apex of the peridium
 - (a) Peridium splitting longitudinally, or laterally lacerate
 - x. Peridium opening lengthwise by valves
Chaenoderma 9: 268
 - y. Peridium laterally lacerate **Cauloglossum 7: 57**
 - (b) Peridium opening horizontally or circularly
 - x. Peridium opening around the stipe
Podaxon 7: 58
 - y. Peridium opening circularly around the middle
† **Sphaerocybis 7: 60**
(**Sphaericeps**)
- II. Peridium typically without a columella, with exo- and endoperidium; gleba floccose, rarely septate
Subfamily Geasterae
- I. Peridium stalked
 - a. Inner peridium alone persistent
 - (1) Peridium fixed to stipe, with distinct mouth
Tylostoma 7: 60
 - (2) Peridium easily separable from stipe; mouth none
Queletia 7: 65
 - b. Both peridial layers persistent
 - (1) Exoperidium forming a volva about the stipe
 - (a) Endoperidium convex; spores on upper surface
Battarea 7: 65
 - (b) Endoperidium hemispheric; spores within
† **Podoloma 17: 223**
(**Battareopsis**)
 - (2) Exoperidium not volvate; inner peridium with a mouth
 - (a) Endoperidium with plicate-sulcate mouth; capillitium copious
Husseyia 7: 67
 - (b) Endoperidium suspended free in cavity of exoperidium, mouth with bright-colored scales
Mitromyces 7: 68
- 2. Exoperidium sessile, typically stellate-laciniate, containing 1 or more endoperidia
 - a. Endoperidium one
 - (1) Spores borne on the inside
 - (a) Exoperidium closed **Diploderma 7: 92**
 - (b) Exoperidium opening stellately or circularly
 - x. Exoperidium stellate
 - (x) Endoperidium dehiscent, usually by a mouth; capillitium present
Geaster 7: 70
 - (y) Endoperidium indehiscent; capillitium none
Stella 9: 272

- y. Exoperidium cup-shaped, mouth minute, ciliate
Diplocystis 7: 92
 - (2) Spores borne on the outside of endoperidium; stellate
Trichaster 7: 93
 - b. Endoperidia several
 - (1) Mycelium crust-like; capillitium not hollow
Broomeia 7: 93
 - (2) Mycelium not crust-like; capillitium hollow
Coelomyces 7: 94
- III. Peridium without a columella; exoperidium lacking or consisting of a papery or spiny cortex; gleba floccose **Subfamily Lycoperdace**
- 1. Peridium with a distinct, stalk-like sterile base; exoperidium spiny or warty
Lycoperdon 7: 106
 - 2. Peridium without sterile base; gleba fertile throughout
 - a. Peridium sessile or nearly so
 - (1) Capillitium a dense elastic mass discrete from the peridium
 - (a) Peridium persistent **Lanopila 7: 95**
 - (b) Peridium falling away **Eriosphaera 7: 96**
 - (2) Capillitium not dense elastic and discrete
 - (a) Peridium persistent
 - x. Mouth at apex, or lacking **Povista 7: 96**
 - y. Mouth at base when in the ground
Catastoma 11: 165
 - (b) Peridium entirely falling away **Lycoperdopsis 16: 243**
 - b. Peridium stipitate; exoperidium dehiscing above along undulating folds
Calvatia 7: 105
- IV. Peridium without columella; gleba with cell-like spaces, often containing sporangioles, or powdery **Subfamily Sclerodermatae**
- 1. Gleba without sporangioles, finally powdery
 - a. Peridium none; gleba naked, subcylindric
Gymnoglossum 11: 158
 - b. Peridium present, enclosing the gleba
 - (1) Peridium sessile or nearly so
 - (a) Peridium not dehiscent
 - x. Gleba reticulate-veined, hard **Corditubera 14: 266**
 - y. Gleba not reticulate-veined, somewhat floccose
 - (x) Spores globose **Hippoperdon 7: 133**
 - (y) Spores fusiform **Castoreum 7: 142**
 - (b) Peridium dehiscent stellately or irregularly
Scleroderma 7: 134
(incl. **Caloderma 16: 243**)
 - (2) Peridium stalked
 - (a) Peridium not dehiscent, clavate **†Corynogaster 14: 266**
(**Clavogaster**)
 - (b) Peridium dehiscent

- x. Peridium clavate, splitting above and entirely disappearing
Favillea 7: 146
- y. Peridium globoid, not entirely disappearing
 - (x) Stipe hollow; peridium dehiscing irregularly, or rimose
Phellorina 7: 145
 - (y) Stipe not hollow
 - m. Peridium many-lobed; stipe fibrous-woody
Xylopodium 7: 143
 - n. Peridium reticulately dehiscent; stipe solid
Areolaria 7: 144
- 2. Gleba containing numerous sporangioles
 - a. Sporangioles fleshy or gelatinous
 - (1) Peridium stipitate; stipe with persistent cupulate volva
Dictyocephalus 17: 238
 - (2) Peridium not volvate, sessile or with stipe-like base
 - (a) Parasitic in glumes; peridium not dehiscent
Testicularia 7: 150
 - (b) Terrestrial or parasitic on roots
 - x. Peridium with sterile stipe-like base, mucose-cellular within
Polysaccum 7: 146
 - y. Peridium sessile, fleshy-cellular within
Polygaster 7: 146
 - b. Sporangioles membranous, not fleshy or gelatinous
 - (1) Peridium corky; sporangioles round
Arachnium 7: 150
 - (2) Peridium membranous; sporangioles cylindric, gyrose
Scoleciocarpus 7: 151
 - (3) Peridium hard; sporangioles large, flexuous
Paurocotylis 7: 152

Family 68. HYMENOGASTRACEAE

7: 154, 9: 280, 11: 168, 14: 267, 16: 245, 17: 239

Typically subterranean, very rarely epigaeous, mycelium often persistent; peridium not opening at maturity, wall occasionally lacking, more or less globose; gleba fleshy or gelatinous, putrescent, more or less cellular or loculate, capillitium none.

I. Peridium wall present, distinct

- 1. Peridium easily separating from the gleba
 - a. Peridium volvate
 - (1) Peridium silky, reticulate-sulcate; volva gelatinous
Clathrogaster 16: 250
 - (2) Peridium waxy-gelatinous, not sulcate
Torrendia 17: 241
 - b. Peridium not volvate
 - (1) Peridium vertical, elongate-cylindric; basidia 2-spored
Protoglossum 11: 158
 - (2) Peridium more or less globose

- (a) Endosporium and exosporium separated by a hyaline mucus
Leucogaster 9: 281
- (b) Endosporium and exosporium contiguous
 - x. Spores elliptic to lanceolate, smooth
Hysterangium 7: 155
 - y. Spores globose, rough or spiny
 - (x) Peridium lanate; basidia usually 7-spored
Sclerogaster 11: 169
 - (y) Peridium not lanate; basidia 3-4-spored
 - m. Gleba with a sterile base, radicate
Octaviania 7: 158
 - n. Gleba without a sterile base, not radicate
Martellia 16: 252
- 2. Peridium separating from the gleba with difficulty or not at all
 - a. Peridium covered with thread-like masses of mycelium
 - (1) Spores hyaline
Rhizopogon 7: 161
 - (2) Spores colored
Melanogaster 7: 164
 - b. Peridium without thread-like masses of mycelium
 - (1) Spores spiny
 - (a) Gleba percurrent by a columella
Arcangeliella 16: 255
 - (b) Gleba without a columella
Hydnangium 7: 175
 - (2) Spores not spiny, smooth, verrucose, rugose, etc.
 - (a) Gleba with branching columella and sterile base
Dendrogaster 17: 240
 - (b) Gleba without columella or sterile base
Hymenogaster 7: 168
(incl. **Chamonixia**, **Leucophleps** 16: 251)
- II. Peridium wall lacking
 - I. Hypogaeous
 - a. Spores elliptic, striate-sulcate
Gautiera 7: 177
 - b. Spores globose, spiny or warty
Gymnomycetes 16: 249
 - 2. Epigaeous; spores globose, warty
Macowanites 7: 179

Family 69. NIDULARIACEAE

7: 28, 9: 265, 11: 156, 14: 256, 16: 229, 17: 214

Epigaeous, fimicole or lignicole, funnel-shaped to cup-shaped, leathery, containing one to many lentiform or globoid sporangioles, the latter attached by a cord to the wall of the peridium; spores elliptic, smooth.

I. Peridium single

- 1. Peridium with several to many sporangioles
 - a. Peridium torn at the apex in opening **Nidularia** 7: 28
 - b. Peridium opening by a deciduous membrane
 - (1) Sporangioles attached to wall by a cord
 - (a) Spores mixed with filaments; peridium of three united layers
Cyathus 7: 32

(b) Spores not mixed with filaments; peridium of a single cottony layer

Crucibulum 7: 43

(2) Sporangioles densely crowded in a glutinous substance

Nidula 17: 215

2. Peridium with a single gelatinous sporangiole

Dacryobolus 7: 45

II. Peridium double, outer stellate, inner with a single viscous sporangiole

Sphaerobolus 7: 46

FUNGI IMPERFECTI

Secondary or propagative stages of other fungi, largely Ascomycetes, characterized by the presence of conidia borne in perithecia-like or disk-like structures, on a stroma, or on a mycelial mass. Many of these forms have been connected by means of experiment with the corresponding perfect stage, but the vast majority of them are found alone in nature.

Order 16. PHOMATALES (Sphaeropsideae Sacc. 3: 1)

Conidia borne on simple or branched threads, so-called basidia, in pycnidia; pycnidia globose, conic, elongate, dimidiate, disk-shaped or cup-shaped, membranous, carbonous, coriaceous or somewhat fleshy, usually black, sometimes bright-colored.

Family 70. PHOMATACEAE (Sphaerioidaceae 3: 1)

Pycnidia globose, conic or lens-like, membranous, carbonous or subcoriaceous, black, immersed or superficial, separate or in a stroma; conidia from 1 to many-celled, hyaline or dark.

Hyalosporae

3: 1, 10: 100, 11: 472, 14: 844, 16: 825, 18: 220

Conidia 1-celled, hyaline, globose, ovoid or oblong, often curved

I. Pycnidia separate

1. Pycnidia smooth

a. Pycnidia borne in discolored areas, i. e., maculicole

Phyllosticta 3: 3

b. Pycnidia not maculicole

(1) Conidia single, not in chains

(a) Conidia muticate, not ciliate or trigonous

x. Subicle none

(x) Pycnidia muticate or papillate, not rostrate or cylindric

m. Pycnidia erumpent or immersed

(m) Basidia 1-spored, mostly short

r. Pycnidia papillate

(r) Growing on lichens

Lichenosticta 16: 851

(s) Not lichenicole

h. Basidia hamate

Phomopsis 18: 264

- i. Basidia not hamate
 - (h) Conidia less than 15μ
Phoma 3: 65
 - (i) Conidia 15μ or more long
Macrophoma 10: 189
- s. Pycnidia astomous or irregularly dehiscent
 - (r) Pycnidia subcarnose, sclerotoid
h. Conidia obtuse at both ends
Plenodomus 3: 184
 - i. Conidia acute at both ends
Sclerotiopsis 3: 184
 - (s) Pycnidia carbonous, circumscissile
Piptostomum 3: 183
- (n) Basidia several-spored, branched
Dendrophoma 3: 178
- n. Pycnidia superficial
 - (m) Pycnidia dense in asteroma-like spots
Asteromella 3: 182
 - (n) Pycnidia not in such spots
 - r. Pycnidia globose or nearly so
 - (r) Basidia short, straight
Aposphaeria 3: 169
 - (s) Basidia beautifully circinate
Pyrenotrichum 3: 184
 - (t) Basidia none
Mycogala 3: 185
 - s. Pycnidia turbinate, carnose
Crocicreas 3: 183
- (y) Pycnidia rostrate or cylindric
 - m. Pycnidia globose, rostrate
Sphaeronaema 3: 185
 - n. Pycnidia cylindric
Glutinium 11: 500
- y. Subicle present
 - (x) Subicle white, cobwebby
Cicinnobolus 3: 216
(incl. *Byssocystis* 11: 502)
 - (y) Subicle dark
 - m. Subicle usually radiate
Asteroma 3: 201
 - n. Subicle not radiate
Chaetophoma 3: 199
- (b) Conidia ciliate, forked or angled
 - x. Conidia ciliate at apex
 - (x) Apex 1-ciliate
Strasseria 18: 284
 - (y) Apex several-ciliate
Neottiospora 3: 216
 - y. Conidia forked or angled
 - (x) Conidia Y-like; subicle present
Ypsilonia 3: 215
 - (y) Conidia trigonous
Trigonosporium 16: 892
- (2) Conidia in chains

- (a) Chains of spores simple or nearly so
Sirococcus 3: 217
- (b) Chains of spores connected, often net-like
Peckia 3: 217
- 2. Pycnidia with hairs or bristles
 - a. Bristles stellate; conidia ovoid **Staurochaeta** 3: 218
 - b. Bristles simple
 - (1) Basidia usually simple, conidia fusoid
Vermicularia 3: 221
 - (2) Basidia usually branched, conidia oblong
Pyrenochaeta 3: 219
- II. Pycnidia in a stroma
 - 1. Stroma globose, conic or valsa-like
 - a. Conidia in chains ***Sirodothis**
 - b. Conidia single
 - (1) Stroma globose, conic or pulvinate
 - (a) Stroma more or less globose or pulvinate
 - x. Stroma unilocular **Dothiopsis** 10: 228
 - y. Stroma several- or many-locular
 - (x) Pycnidia distinct
 - m. Pycnidia aggregate in a basal stroma
Dothiorella 3: 235
 - n. Pycnidia more deeply immersed
 - (m) Necks not joined in one ostiole
Lamyella 11: 510
 - (n) Necks joined in a single ostiole
Torsellia 11: 510
 - (y) Pycnidia merely locules in the stroma
 - m. Locules several, not numerous
Rabenhorstia 3: 243
 - n. Locules very numerous
Fuckelia 3: 244
 - (b) Stroma conic-truncate, conidia bacillar
Ceuthospora 3: 277
 - (2) Stroma valsa-like
 - (a) Conidia fusoid or bacillar **Fusicoccum** 3: 247
 - (b) Conidia allantoid **Cytospora** 3: 252
 - (c) Conidia globose or ovoid **Cytosporella** 3: 251
 - 2. Stroma applanate, effuse or linear
 - a. Stroma linear, conidia connate in fours
Gamosporella 10: 238
 - b. Stroma applanate or effuse
 - (1) Growing on leaves and stems **Placosphaeria** 3: 244
 - (2) Growing on fungi **Anthraceroderm** 10: 238

Of Uncertain Position.

Manginia 18: 266. a Phoma with micro- and macropycnidia

Phaeosporae

3: 291, 10: 251, 11: 511, 14: 919, 16: 905, 18: 302

Conidia 1-celled, dark, globose, ovoid or oblong

I. Pycnidia separate**1. Pycnidia without mycelium or subicle****a. Pycnidia smooth, not hairy**

(1) Conidia in chains, globose

Sirothecium 10: 270

(2) Conidia not in chains

(a) Pycnidia sessile, spheroid

x. Pycnidia beaked

Naemosphaera 10: 259

y. Pycnidia not beaked

(x) Pycnidia with a distinct orbicular locule

Hypocenia 3: 320

(y) Pycnidia without such a locule

m. Conidia on long basidia

(m) Pycnidia thin, white-lacerate at top

Harknessia 3: 320

(n) Pycnidia subcarbonous, not lacerate

Sphaeropsis 3: 291

n. Basidia very short or obsolete

Coniothyrium 3: 305

(b) Pycnidia stipitate, clavate

Levieuxia 3: 321**b. Pycnidia hairy or setose****Chaetomella 3: 321****2. Pycnidia with distinct mycelium or subicle**

a. Pycnidia astomous, in a dark subicle

Capnodiastrum 10: 272

b. Pycnidia perforate, with basal hyphae

Cicinnobella 18: 302**II. Pycnidia cespitose or in a stroma****1. Pycnidia in dense erumpent clusters****Haplosporella 3: 323****2. Pycnidia in a definite stroma**

a. Stroma applanate or effuse, foliicole

Discomycetopsis 11: 517

b. Stroma dot-like, discoid or hemispheric

(1) Stroma dot-like, immersed

Melanconiopsis 16: 915

(2) Stroma discoid to hemispheric

(a) Stroma discoid; spores large

Nothopatella 11: 517

(b) Stroma pulvinate; spores minute, catenulate

Cytoplea 3: 325

(c) Stroma hemispheric; pycnidia circinate

†Circinastrum 3: 325**(Weinmannodora)****Hyalodidymae**

3: 384, 10: 295, 11: 522, 14: 942, 16: 925, 18: 335

Conidia hyaline, 1-septate, ovoid, ellipsoid or oblong

I. Pycnidia separate**1. Pycnidia not beaked**

a. Pycnidia in discolored areas, maculicole

- (1) Pycnidia immersed, then erumpent, perforate
 - (a) Conidia muticate **Ascochyta 3:384**
 - (b) Conidia with setae at the apex **Robillardia 3:407**
- (2) Pycnidia superficial, astomous **Puccinospora 10:317**
- b. Pycnidia not maculicole
 - (1) Pycnidia hairy **Didymochaete 14:953**
(**Vermiculariella 16:940**)
 - (2) Pycnidia smooth
 - (a) Conidia with an appendage at each end
 - x. Conidia with 1 or more bristles **Darlucella 3:410**
 - y. Conidia with cap-like appendages **Tiarospora 10:311**
 - (b) Conidia muticate
 - x. Basidia 1-spored
 - (x) Pycnidia on a cobwebby subicle, phyllogenous
Actinonema 3:408
 - (y) Pycnidia without subicle, ramicole
Diplodina 3:411
 - y. Basidia several-many-spored **Cystotricha 3:413**
- 2. Pycnidia beaked **Rhynchophoma 3:414**
- II. Pycnidia in a stroma
 - 1. Stroma effuse
 - a. Stroma consisting of two distinct layers
Thoracella 16:941
 - b. Stroma of a single layer
Placosphaerella 14:948
 - 2. Stroma verruciform
 - a. Stroma superficial
Pazschkella 16:528
 - b. Stroma erumpent
Cytodiplospora 11:942

Phaeodidymae

2:329, 10:275, 11:518, 14:927, 16:915, 18:319

Spores dark, 1-septate, ovoid to oblong

- I. Pycnidia separate
 - 1. Pycnidia beaked
 - a. Pycnidia hairy **Rhynchodiplodia 18:329**
 - b. Pycnidia smooth **Pellioniella 18:329**
 - 2. Pycnidia not beaked
 - a. Pycnidia hairy **Chaetodiplodia 3:374**
 - b. Pycnidia smooth
 - (1) Conidia with a mucous layer, very large
Macrodiplodia 3:374
 - (2) Conidia without a mucous layer
 - (a) Pycnidia erumpent
 - x. Conidia 1-ciliate at apex
***Chaetoconis 10:337**
(**Kellermannia in part**)
 - y. Conidia muticate
 - (x) Conidia less than 15 μ long
Microdiplodia 18:323

- (y) Conidia 15 μ or more long **Diplodia 3:329**
- (b) Pycnidia superficial, lignicole **Diplodiella 3:375**
- II. Pycnidia cespitose or in a stroma **Botryodiplodia 3:377**
 - 1. Pycnidia cespitose
 - 2. Pycnidia in a stroma
 - a. Pycnidia and subicle enclosed in a hemispheric stroma **Lasiodiplodia 14:939**
 - b. Pycnidia without subicle, in a globose stroma **Diplodiopsis 18:335**

Hyalophragmiae

3:418, 10:330, 11:533, 14:962, 16:947, 18:358

Conidia hyaline, 2-several-septate, oblong to fusoid

- I. Pycnidia more or less globose
 - 1. Subicle none
 - a. Conidia appendaged at apex
 - (1) Seta 1 **Kellermannia 10:337**
 - (2) Setae 3 **Bartalinia 16:951**
 - b. Conidia muticate **Stagonospora 3:445**
 - 2. Subicle present, dark, phyllogenous **Asteromidium 10:338**
- II. Pycnidia elongate to cylindric **Mastomyces 3:456**

Phaeophragmiae

3:418, 10:317, 11:528, 14:953, 16:943, 18:362

Conidia hyaline, 2-several-septate, oblong to fusoid

- I. Pycnidia separate
 - 1. Conidia free from each other
 - a. Conidia muticate
 - (1) Pycnidia papillate or subastomous
 - (a) Pycnidia with flattened base **Macrobatis 11:532**
 - (b) Pycnidia globose, without flattened base
 - x. Pycnidia on a stellate subicle, superficial **Couturea 3:442**
 - y. Pycnidia without a subicle, crumpeut
 - (x) Pycnidia hairy **Wojnowicia 14:960**
 - (y) Pycnidia smooth **Hendersonia 3:418**
 - (2) Pycnidia opening widely, with an operculum
 - (a) Pycnidia superficial, dark, hairy **Angiopoma 3:442**
 - (b) Pycnidia immersed, pale, smooth **Lichenopsis 3:442**
 - b. Conidia appendaged
 - (1) Conidia 1-ciliate at each end **Cryptostictis 3:443**
 - (2) Conidia 1-ciliate at base by the basidium
 - †Uroconis 18:368**
 - (Urohendersonia)**
 - (3) Conidia with a round or cup-like appendage at each end **Santiella 16:947**

- 2. Conidia united in groups
 - a. Conidia united into a fascicle Eriosporina 11:532
 - b. Conidia stellately united Prosthemia 3:444
- II. Pycnidia locules in a stroma Hendersonula 3:445

Hyalodictyae

16:955

Conidia hyaline, muriform, ovoid or oblong

- I. Pycnidia erumpent, papillate †Hyalothyris 16:955
(Hyalothyridium)

Phaeodictyae

3:459, 10:338, 11:536, 14:964, 16:951, 18:369

Conidia dark, muriform, oblong to ovoid, rarely radiate or cruciate

- I. Pycnidia separate
 - 1. Conidia not reticulately roughened
 - a. Pycnidia corticole, erumpent Camarosporium 3:459
 - b. Pycnidia xylogenous, subsuperficial Cytosporium 3:470
 - 2. Conidia reticulately roughened Endobotrya 3:470
- II. Pycnidia locules in a stroma Dichomera 3:471

Scolecosporae

3:474, 10:349, 11:538, 14:967, 16:956, 18:376

Conidia hyaline or dilutely colored, elongate-fusoid, bacillar or filiform, continuous or septate.

- I. Pycnidia separate
 - 1. Pycnidia membranous or carbonous
 - a. Pycnidia superficial
 - (1) Pycnidia hairy
 - (a) Conidia single on the basidia Trichocollonema 18:404
 - (b) Conidia ternate on the basidia Gamospora 10:402
 - (2) Pycnidia smooth
 - (a) Pycnidia beaked Cornularia 3:598
 - (b) Pycnidia not beaked
 - x. Conidia usually expelled in a ball Collonema 10:397
 - y. Conidia not expelled in a ball Septorella 14:981
 - b. Pycnidia immersed or erumpent
 - (1) Pycnidia hairy, maculicole Trichoseptoria 11:548
 - (2) Pycnidia smooth
 - (a) Pycnidia beaked Sphaerographium 3:596
 - (b) Pycnidia not beaked
 - x. Pycnidia maculicole, phyllogenous Septoria 3:474
 - y. Pycnidia not maculicole
 - (x) Pycnidia complete at top, usually papillate Rhabdospora 3:578

- (y) Pycnidia more or less incomplete at top
 - m. Pycnidia gaping, showing a gelatinous spore mass
Gelatinosporium 3: 596
 - n. Pycnidia not exposing a gelatinous mass
 - (m) Pycnidia foliicole **Phleospora 3: 577**
 - (n) Pycnidia rami-caulicole **Phlyctaena 3: 593**
- 2. Pycnidia suberose, incomplete, often pale
 - a. Pycnidia cespitose **Micropera 3: 604**
 - b. Pycnidia merely gregarious **Micula 3: 604**
- II. Pycnidia in a stroma
 - 1. Conidia 4-6 fasciculate on a basidium **Eriospora 3: 600**
 - 2. Conidia separate
 - a. Conidia setose-penicillate **Dilophospora 3: 600**
 - b. Conidia muticate
 - (1) Stroma superficial, setose **†Merodothidis 18: 405**
(Septodothideopsis)
 - (2) Stroma erumpent or immersed
 - (a) Pycnidia distinct in the stroma **Cytosporina 3: 601**
 - (b) Pycnidia locules in the stroma **Septosporiella 10: 403**

Family 71. ZYTHIACEAE

(Nectrioidaceae Sacc. 3: 613)

Pycnidia, and stromata when present, fleshy or waxy, light-colored, white, yellow, red or orange, globose, more rarely cup-shaped or hysterioid; conidia various, mostly hyaline.

Subfamily Zythiae

Pycnidia more or less globose

Hyalosporae

3: 613, 10: 404, 11: 552, 14: 988, 16: 983, 18: 407

- I. Pycnidia separate
 - 1. Pycnidia smooth
 - a. Pycnidia beakless
 - (1) Conidia in chains **Sirozythia 18: 410**
 - (2) Conidia not catenulate
 - (a) Pycnidia on creeping hyphae **Eurotiopsis 10: 406** ..
 - (b) Pycnidia without mycelium
 - x. Conidia spiny or ciliate
 - (x) Conidia spiny **Roumegueriella 3: 616**
 - (y) Conidia with several cilia at apex **Ciliospora 18: 410**
 - y. Conidia smooth
 - (x) Pycnidia single-walled
 - m. Pycnidia more or less papillate
Zythia 3: 614

- n. Pycnidia with crateriform ostiole
 - Libertiella** 3:616
 - o. Pycnidia cup-shaped
 - Lemalis** 3:672
 - (y) Pycnidia with outer circumscissile wall
 - Dichlaena** 3:620
 - Sphaeronaemella** 3:617
 - b. Pycnidia beaked
 - 2. Pycnidia hairy or spiny
 - a. Pycnidia densely beset with conoid 1-celled setae
 - Muricularia** 3:218
 - b. Pycnidia with slender bristles or hairs
 - (1) Hairs fasciculate
 - Collocystis** 3:616
 - (2) Hairs separate
 - (a) Hairs everywhere but at the apex
 - Chaetozythia** 10:406
 - (b) Hairs only around the wide ostiole
 - Pseudozythia** 18:409
 - II. Pycnidia caespitose or in a stroma
 - 1. Pycnidia caespitose, beaked; conidia in chains
 - Trelesiella** 14:989
 - 2. Pycnidia in a stroma
 - a. Stroma more or less pulvinate; conidia fusoid
 - Aschersonia** 3:619
 - b. Stroma fruticose branched; conidia bacillar
 - Hypocreodendrum** 14:992
- Phaeosporae**
10:409, 18:416
- Conidia dark, 1-celled, globose to oblong
- I. Pycnidia separate, beaked; basidia obsolete
 - Ampullaria** 18:416
 - II. Pycnidia in a stroma
 - Martinella** 10:409
- Hyalodidymae**
3:621, 10:409, 11:553, 16:986, 18:416
- Conidia hyaline or nearly so, 1-septate, ovoid to oblong
- I. Basidia simple or nearly so
 - Pseudodiplodia** 3:621
 - II. Basidia dendroid branched
 - Diplozythia** 18:417
- Hyalophragmiae**
3:621, 10:410, 18:417
- Conidia hyaline, several-septate, elliptic to fusoid
- I. Conidia oblong-fusoid
 - Stagonopsis** 3:621
 - II. Conidia 4-radiate, with septate radii
 - Chiastospora** 3:621
- Scolecosporae**
3:622, 10:410, 18:418
- Conidia hyaline, bacillar or filiform, continuous or septate
- I. Pycnidia separate

- 1. Pycnidia beakless, almost discoid **Trichocrea 10:410**
- 2. Pycnidia beaked; conidia 1-ciliate **Mycorhynchus 18:418**
- II. Pycnidia in a stroma; conidia hamate **Polystigmina 3:622**

Subfamily Patellinae

Pycnidia cupulate or hysterioid

Hyalosporae

3:622, 10:411, 11:553, 18:419

Conidia hyaline, 1-celled, globose to oblong

I. Pycnidia separate

1. Pycnidia cup-shaped

a. Pycnidia smooth

- (1) Pycnidia carnose; basidia simple, cylindric

Patellina 3:622

- (2) Pycnidia submembranous; basidia branched

Ollula 10:411

b. Pycnidia hairy

- (1) Conidia in chains

***Sirocypis**

- (2) Conidia not in chains

Cyphina 3:623

2. Pycnidia flattened, oblong, cleft

Hysteromyxa 3:622

II. Pycnidia in a stroma

1. Stroma suberose, white

Munkia 10:408

2. Stroma corneous, black

†**Pycnostroma 18:415**
(**Aschersoniopsis**)

Hyalophragmiae

11:553

Conidia hyaline, several-septate, oblong

I. Pycnidia immersed, waxy

Pseudostictis 11:553

Scolecosporae

10:411

Conidia hyaline, filiform, continuous

I. Pycnidia waxy, cup-shaped, on a white subicle

Trichosperma 10:411

Family 72. LEPTOSTROMATACEAE

Pycnidia membranous or carbonous, black, more or less distinctly dimidiate, scutiform, astomous, ostiolate or cleft, erumpent or superficial.

Hyalosporae

3:625, 10:412, 11:553, 14:992, 16:986, 18:419

Conidia hyaline, 1-celled, globose to oblong

I. Pycnidia separate

1. Pycnidia astomous or variously perforate, but not cleft

a. Basidia lacking

- (1) Pycnidia on a subicle
 - (a) Subicle of fumiginous hyphae **Eriothyrium** 10: 418
 - (b) Subicle of broad fibers **†Trichopeltium** 10: 418
(**Trichopeltulum**)
- (2) Pycnidia without subicle
 - (a) Conidia muticate
 - x. Pycnidia stellately divided or cleft **Actinothecium** 3: 638
 - y. Pycnidia depressed-clypeate, not stellate
 - Leptothyrium** 3: 626
 - (**Sacidium** 3: 649)
 - (b) Conidia setulose at each end **Tracyella** 18: 424
 - b. Basidia present, cylindric **Piggotia** 3: 636
- 2. Pycnidia more or less clearly cleft lengthwise
 - a. Pycnidia elongate or lanceolate **Leptostroma** 3: 639
 - b. Pycnidia subcircular **Labrella** 3: 647
- II. Pycnidia in a stroma
 - 1. Stroma phyllogenuous **Melasmia** 3: 637
 - 2. Stroma growing on animal hairs **Trichophila** 10: 423

Phaeosporae

3: 653, 10: 423, 14: 996, 18: 429

Conidia dark, 1-celled, globose to oblong

- I. Pycnidia separate
 - 1. Pycnidia on a dark subicle, radiately dehiscent **Asterostomella** 10: 423
 - 2. Pycnidia not on a subicle
 - a. Conidia conglobate, verrucose **Discomycopsella** 18: 429
 - b. Conidia not conglobate, smooth **Pirostoma** 3: 653
- II. Pycnidia in a stroma
 - 1. Stroma membranous
 - a. Pycnidia distinct, exerted **Peltostroma** 18: 430
 - b. Pycnidia merely locules, immersed **Lasmenia** 10: 425
 - 2. Stroma carbonous; locules many, immersed **Poropeltis** 18: 430

Hyalodidymae

10: 426, 11: 557, 18: 431

Conidia hyaline, 1-septate, oblong to fusoid

- I. Pycnidia separate
 - 1. Pycnidia astomous or variously perforate, not cleft
 - a. Conidia muticate **Leptothyrella** 10: 426
 - b. Conidia cuspidate at apex, falcate **Kabatia** 18: 433
 - 2. Pycnidia cleft lengthwise, elongate **Fioriella** 18: 432
- II. Pycnidia in a stroma, rimose **Pseudomelasmia** 18: 434

Phaeodidymae

10: 426, 18: 431

Conidia dark, 1-septate, oblong to fusoid

- I. Pycnidia separate
 - a. Pycnidia ostiolate **Diplopeltis 10: 426**
 - b. Pycnidia longitudinally cleft **Holcomyces 18: 431**
- II. Pycnidia in a stroma, ostiolate **Seynesiopsis 18: 431**

Hyalophragmiae

3: 653, 10: 426, 11: 557, 14: 996, 16: 992, 18: 434

Conidia hyaline, 2-several-septate, oblong to fusoid

- I. Pycnidia astomous or ostiolate, not cleft
 - 1. Conidia muticate; pycnidia with creeping hyphae **Asterothyrium 18: 434**
 - 2. Conidia ciliate
 - a. Conidia fusoid, 1-ciliate at each end **Discosia 3: 653**
 - b. Conidia cruciate, each arm 1-ciliate **Entomosporium 3: 657**
- II. Pycnidia rimose dehiscent **Cystothyrium 10: 427**

Phaeophragmiae

14: 997, 18: 435

Conidia dark, 1-several-septate, oblong to fusoid

- I. Pycnidia separate, rimose-gaping; conidia 1-ciliate each way **Labridium 14: 997**
- II. Pycnidia in a stroma; conidia muticate, finally black **Phragmopeltis 18: 435**

Scolecosporae

3: 658, 10: 428, 11: 557, 14: 997, 16: 992, 18: 436

Conidia normally hyaline, bacillar or filiform, continuous or septate

- I. Pycnidia astomous or opening variously
 - 1. Pycnidia with a round ostiole; conidia catenate **Crandallia 14: 998**
 - 2. Pycnidia astomous or irregularly dehiscent
 - a. Pycnidia with radiate-fimbriate margin **Actinothyrium 3: 658**
 - b. Pycnidia not radiate-fimbriate
 - (1) Pycnidia of two kinds, small simple and large loculate **Brunchorstia 10: 431**
 - (2) Pycnidia of one kind
 - (a) Conidia muticate
 - x. Pycnidia corrugate, not hairy; conidia not separating **Melophia 3: 658**
 - y. Pycnidia hairy; conidia separating into joints **Chaetopeltis 14: 998**
 - (b) Conidia ciliate-penicillate at apex **Giulia 18: 435**

II. Pycnidia elongate, longitudinally cleft

1. Basidia simple, bacillar **Leptostromella 3: 659**
2. Basidia umbellately branched ***Petasodes 14: 998**

Family 73. EXCIPULACEAE

Pycnidia membranous or carbonous, black, cup-shaped, patellate or hysterioid, at first more or less spheric, but at length widely open, erumpent or superficial, glabrous or hairy.

Hyalosporae

3: 665, 10: 432, 11: 558, 14: 999, 16: 993, 18: 436

Conidia hyaline, 1-celled, globose to oblong

I. Pycnidia pilose or setose

1. Conidia muticate; pycnidia cupulate **Amerosporium 3: 680**
2. Conidia ciliate; pycnidia cupulate
 - a. Conidia several-ciliate at apex **Polynema 3: 687**
 - b. Conidia 1-ciliate at each end **Dinemasporium 3: 683**

II. Pycnidia smooth or nearly so

1. Pycnidia more or less cup-shaped, or disciform
 - a. Pycnidia composed of conglutinate dark hyphae **Godroniella 3: 665**
 - b. Pycnidia with cellular context
 - (1) Pycnidia cup-like when mature, sometimes obconoid
 - (a) Basidia simple **Excipula 3: 665**
 - x. Pycnidia cup-shaped **Catinula 3: 673**
 - y. Pycnidia terete-conic **Heteropatella 3: 670**
 - (b) Basidia branched **Dothichiza 3: 671**
 - (2) Pycnidia subglobose-collabent, disciform or verruciform
 - (a) Pycnidia subglobose, irregularly dehiscent and collabent **Discula 3: 674**
 - (b) Pycnidia disciform, often imperfect and covered by epiderm **Agyriellopsis 18: 438**
 - (c) Pycnidia verruciform; conidia mucose-involute **Psilospora 3: 679**
2. Pycnidia hysterioid or valvately gaping
 - a. Pycnidia widely hysterioid **Sporonema 3: 677**
 - b. Pycnidia valvately gaping
 - (1) Basidia typically branched **Pleococcum 3: 679**
 - (2) Basidia simple or none

Phaeosporae

10: 439, 18: 441

Conidia dark, 1-celled, globose to oblong

- I. Pycnidia patellate, smooth **Phaeodiscula 10: 439**
- II. Pycnidia cupulate, setulose at margin **†Coniothyris 10: 439**
(**Coniothyriella**)

Hyalodidymae

3: 687, 10: 440, 11: 560, 14: 1002, 16: 993, 18: 442

Conidia hyaline, 1-septate, oblong to fusoid

I. Pycnidia discoid or patellate

1. Pycnidia discoid, veiled; basidia simple **Discella 3: 687**2. Pycnidia patellate, subsuperficial; basidia branched
Pseudopatella 3: 688

II. Pycnidia hysterioid or irregularly gaping

1. Pycnidia hysterioid, elongate **Scaphidium 18: 443**2. Pycnidia globose, then irregularly gaping; conidia catenate
Siropatella 18: 443**Hyalophragmiae**

3: 688, 10: 441, 11: 560, 14: 1002, 18: 443

Conidia hyaline, 2-several-septate, oblong to fusoid

I. Pycnidia cupulate or subcupulate

1. Pycnidia smooth; conidia sometimes 1-ciliate
Excipulina 3: 688

2. Pycnidia setulose

a. Conidia fusoid, inner cells somewhat colored
Excipularia 3: 689b. Conidia X-shaped, entirely hyaline **Acanthothecium 10: 442**

II. Pycnidia discoid and unequal, margin lacerate

Pilidium 3: 689**Phaeophragmiae**

10: 443, 18: 444

Conidia dark, 2-several-septate, oblong to fusoid

I. Pycnidia hysterioid; conidia not catenate **Dichaenopsis 18: 444**

II. Pycnidia laciniately dehiscent; conidia catenate

Taeniophora 10: 443**Scolecosporae**

3: 690, 10: 443, 14: 1002, 16: 993, 18: 445

Conidia typically hyaline, bacillar or filiform, continuous or septate

I. Pycnidia separate

1. Conidia separating at the joints **Schizothyrella 3: 690**
(incl. **Pseudocenangium 10: 445**)

2. Conidia not separating

a. Pycnidia discoid, margin lacerate; conidia filiform
Protostegia 3: 690

b. Pycnidia mostly cupulate, not lacerate; conidia hamate

Oncospora 3: 691

II. Pycnidia in a stroma, pezizoid

Ephelis 3: 691

Order 17. MELANCONIALES

Family 74. MELANCONIACEAE

Pycnidia lacking, or reduced to a stratum merely; strata typically bearing basidia of various sorts upon which conidia arise, forming masses or acervuli, which are immersed or erumpent, black, gray or light-colored, waxy, corneous or even submembranous.

Hyalosporae

3: 698, 10: 446, 11: 562, 14: 1004, 16: 995, 18: 447

Conidia hyaline, 1-celled, globose to oblong, rarely dilutely colored

I. Conidia muticate

1. Masses, or acervuli, not setose

a. Conidia not catenate

(1) Masses bright-colored, subtremelloid

Hainesia 3: 698

(2) Masses gray to black, rarely bright-colored, waxy or horny

(a) Masses gray, rarely bright-colored, waxy

x. Growing on leaves or fruits for the most part

Gloeosporium 3: 699

y. Growing usually on twigs of trees or shrubs

Myxosporium 3: 722

(b) Masses black, discoid, horny

Melanostroma 3: 728

b. Conidia in chains

(1) Masses oblong, hysteroïd, dark, hard

Hypodermium 3: 728

(2) Masses discoid, pulvinate or conoid

(a) Masses bright-colored, softish

Myxosporella 3: 729

(b) Masses dark to black

x. Basidia repeatedly branched

(x) Masses discoid; basidia dichotomous

Blennoria 3: 730

(y) Masses depressed-pulvinate; basidia verticillate

Agyriella 3: 731

(z) Masses perithecioid; basidia irregularly branched

***Hormyllum** 3: 733

y. Basidia simple

(x) Masses perithecioid, black

***Thecostroma** 3: 752

(y) Masses scutellate, olive or ashen

Myxormia 3: 734

(z) Masses truncate, black below, pale above

Bloxamia 3: 734

2. Masses setose at margin; basidia short, fasciculate

Colletotrichum 3: 735

II. Conidia aristate with a branched awn at apex

Pestalozziella 3: 737

Phaeosporae

3: 749, 10: 471, 11: 571, 14: 1018, 16: 1008, 18: 469

Conidia dark, 1-celled, globose to oblong or fusoid

I. Conidia solitary on the basidia

- 1. Conidia globose or oblong **Melanconium 3: 749**
- 2. Conidia fusoid, often arcuate
 - a. Basidia not swollen at base **Cryptomela 3: 760**
 - b. Basidia swollen at base **Basiascum 10: 474**

II. Conidia in chains

- 1. Conidial chains separate **Trullula 3: 731**
- 2. Conidial chains in a mucose head **Thyrsidium 3: 761**

Hyalodidymae

3: 766, 10: 475, 11: 572, 14: 1020, 16: 1009, 18: 472

Conidia hyaline or nearly so, 1-septate, ovoid to fusoid

I. Conidia muticate

- 1. Saprogenous, on stems and fruits **Septomyxa 3: 766**
- 2. Biogenous, typically on leaves **Marsonia 3: 767**

II. Conidia 3-4-ciliate at each end

Gloeosporiella 11: 575**Phaeodidymae**

3: 763, 10: 475, 11: 572, 14: 1029, 16: 1009

Conidia dark, 1-septate, ovoid to fusoid

I. Conidia solitary

- 1. Conidia muticate **Didymosporium 3: 763**
- 2. Conidia 1-3-ciliate at apex **Neobarclaya 14: 46, 10: 475**

II. Conidia catenate, connected by hyaline isthmi

Bullaria 3: 766**Hyalophragmiae**

3: 801, 10: 480, 11: 575, 14: 1022, 16: 1012, 18: 474

Conidia hyaline, 2-several-septate, oblong to fusoid or clavate

I. Conidia separate

- 1. Conidia muticate
 - a. Conidia oblong or fusoid, masses usually pale **Septogloeum 3: 801**
 - b. Conidia long-clavate; masses dark **Rhopalidium 3: 801**
- 2. Conidia 1-several-ciliate, usually at the apex **Pestalozzina 11: 580**

II. Conidia united at base into a radiate or stellate group

Prosthemmiella 3: 803
 (incl. *Psammmina* 10: 498)

Phaeophragmiae

3: 771, 10: 480, 11: 575, 14: 1022, 16: 1012, 18: 475

Conidia dark, at least in part, 2-several-septate, oblong to cylindric

I. Conidia muticate

1. Conidia separate, not in chains

a. Conidia oblong or elongate

(1) Conidia curved-attenuate, i. e., hyaline-rostrate

(a) Conidia dark, except the hyaline beak

Scolecosporium 3: 782

(b) Conidia with 2 inner cells opaque, others clear

Toxosporium 14: 1030

(2) Conidia oblong, not rostrate

(a) Conidia cirrhose protruded and atro-inquant

Stilbospora 3: 771

(b) Conidia not protruded and atro-inquant

Coryneum 3: 774

b. Conidia stellate-lobed, lobes several-septate

Asterosporium 3: 782

2. Conidia in chains

a. Conidia connected by filiform isthmi

Siridium 3: 782

b. Conidia chains without isthmi

Siridiella 11: 580(incl. *Septotrullula* 18: 487)

II. Conidia ciliate

1. Conidia ciliate at apex alone

a. Conidia 1-ciliate

Monochaetia 18: 485

b. Conidia several-ciliate

Pestalozzia 3: 784

2. Conidia 1-ciliate at each end

Hyaloceras 3: 783(incl. *Amphichaeta* 18: 486)**Phaeodictyae**

3: 803, 10: 508, 11: 565, 14: 1035, 16: 1022, 18: 488

Conidia dark, muriform, ovoid or oblong

I. Conidia muticate

1. Conidia not catenate

Steganosporium 3: 803

2. Conidia catenate by cylindric isthmi

Phragmotrichum 3: 806

II. Conidia pluriciliate at apex; end cells subhyaline

Morinia 10: 508**Scolecosporae**

3: 737, 10: 498, 11: 582, 14: 1031, 16: 1018, 18: 488

Conidia cylindric, filiform or suballantoid, hyaline, mostly continuous

I. Conidia allantoid

Naemospora 3: 746

II. Conidia bacillar to filiform

1. Conidia fasciculate at the apex of the basidia

Trichodytes 14: 1031

2. Conidia solitary

a. Masses white or pale, foliicole; conidia filiform

Cylindrosporium 3: 737, 18: 491

b. Masses gray or dark, usually ramicole; conidia falcate

Cryptosporium 3: 740

- c. Masses bright-colored, saprophytic; conidia falcate

Libertella 3:744

Staurosporae

18:493

Conidia star-shaped, hyaline

- I. Masses phyllogenous, bright-colored; conidia 4-radiate

Asteroconium 18:493

Order 18. MONILIALES (Hyphomyceteae Sacc. 4:1)

Hyphae more or less developed, cobwebby or more or less compacted, but rarely arising from a definite stratum or stroma, never enclosed in a pycnidium, typically superficial.

Family 75. MONILIACEAE (Mucedineae 4:2)

Hyphae hyaline or bright-colored, more or less fragile, lax, not cohering in fascicles; conidia concolorous, i. e., hyaline or bright-colored.

Hyalosporae

4:2, 10:510, 11:586, 14:1037, 16:1023, 18:495

Conidia hyaline, or bright-colored, 1-celled, globose, ovoid to short-cylindric

Micronemeae

Hyphae very short or obsolete, or little different from the conidia

- I. Conidia not in chains

1. Conidia solitary, at least not capitate

- a. Saprogenous

- (1) Hyphae none

- (a) Conidia separate

Chromosporium 4:6

- (b) Conidia joined in twos or threes, not catenate

Selenotila 11:587

- (2) Hyphae very short, branched, septate

Coccospora 4:9

- b. Entomogenous

Massospora 4:10

(incl. **Sorosporella** 10:512)

- c. Phytogenous

- (1) In fungi

- (a) Conidia ovoid, smooth

Myceliophthora 11:587

- (b) Conidia globose, verrucose

Coccosporella 11:586

- (2) In leaves

- (a) Hyphae paliform, stipate, very short

Microstroma 4:9

- (b) Hyphae vermiform-tortuose; biophilous

Ophiocladium 11:587

2. Conidia capitate; hyphae lacking; biophilous

Glomerularia 4:10

II. Conidia in chains

1. Saprophilous

a. Conidial chains arising in the hyphae

(1) Conidial branches simple, arcuate **Malbranchea 4:11**

(2) Conidial branches dichotomous, not arcuate

Glycophila 4:11

b. Chains arising at the apex of the hyphae

(1) Conidia globose, elliptic or fusiform

(a) Hyphae short, simple or nearly so

x. Conidia globose or suboblong **Oospora 4:11**y. Conidia fusoid, acute each way **Fusidium 4:25**

(b) Hyphae longer, distinctly branched

Monilia 4:31(incl. *Halobysus* 11:588)

(2) Conidia bacillar or cuboid

(a) Hyphae nearly obsolete; conidia bacillar

Cylindrium 4:36

(b) Hyphae distinctly present

x. Conidia bacillar

Polyscytalum 4:38

y. Conidia cuboid

Geotrichum 4:39

2. Biophilous

a. Growing within leaf tissue

Oidiopsis 18:507

b. Growing on leaves or other parts

(1) Conidia ellipsoid, without isthmi **Oidium 4:40**

(2) Conidia globose, connected by isthmi

Paepalopsis 4:47**Macronemeae**

Hyphae elongate and distinct from the conidia

I. Conidia in heads

Cephalosporiae

1. Conidia not catenulate

a. Conidia globose or oblong

(1) Conidia sessile on the head or nearly so

(a) Fertile hyphae inflated at apex

x. Apical vesicle globose-inflated

(x) Conidia sessile, not mucus-covered

m. Vesicle verrucose or muriculate

(m) Fertile hyphae simple **Oedocephalum 4:47**

(n) Fertile hyphae sigmoid, much branched

Sigmoideomyces 10:523

n. Vesicle hexagonally areolate

Rhopalomyces 4:50

(y) Conidia on stalks, mucus-covered

Gliocephalus 16:1031

y. Vesicle clavate or lobed

(x) Vesicle disk-shaped, stellate-lobed

Coronella 4:51

- (y) Vesicle clavate or subpalmate **Buseella 18: 509**
- (b) Fertile hyphae not inflated at apex
 - x. Conidial head covered with mucus
 - (x) Fertile hyphae simple **Hyalopus 4: 51**
 - (y) Fertile hyphae with verticillate branches at tip **Gliobotrys 18: 510**
 - y. Head without mucus
 - (x) Fertile hyphae with one head
 - m. Conidia not separating **Papulospora 4: 58**
 - n. Conidia separating
 - (m) Head elongate **Doratomyces 4: 53**
 - (n) Head globose or slightly clavate
 - r. Sterile hyphae scanty **Haplotrichum 4: 53**
 - s. Sterile hyphae long, decumbent **Cephalosporium 4: 56**
 - (y) Fertile hyphae with 2-several heads
 - m. Conidia upright on verticillate basidia **Coemansiella 4: 55**
 - n. Conidia in more definite heads
 - (m) Fertile hyphae simple, with 3-several heads of conidia on spines **Botryosporium 4: 54**
 - (n) Fertile hyphae several times 2-3-fid **Trichoderma 4: 59**
- (2) Conidia borne on little stalks or sterigmata
 - (a) Fertile hyphae simple **Corethrospis 4: 62**
 - (b) Fertile hyphae verticillate branched **Spicularia 4: 63**
- b. Conidia short cylindric
 - (1) Conidia without mucus **Cylindrocephalum 4: 63**
 - (2) Conidia covered with mucus **Acontium 18: 512**
- 2. Conidia catenulate **Aspergillae**
 - a. Fertile hyphae inflated at apex
 - (1) Fertile hyphae simple or nearly so
 - (a) Sterigmata of apical vesicle none or simple
 - x. Conidia terminal on sterigmata **Aspergillus 4: 64**
 - y. Conidia lateral and terminal on sterigmata **Dimargaris 4: 76**
 - (b) Sterigmata verticillate branched **Sterigmatocystis 4: 71**
(incl. **Alliospora 18: 516**)
 - (2) Fertile hyphae dichotomous, branches curved **Dispira 4: 77**
 - b. Fertile hyphae little or not at all inflated
 - (1) Fertile hyphae verticillately branched at tip
 - (a) Tips equally verticillate; conidia doliiform **Amblyosporium 4: 77**
 - (b) Tips unequally verticillate; conidia globoid

- x. Conidia without mucus **Penicillium 4: 78**
(incl. **Citromyces 11: 593**)
 - y. Conidia enclosed in mucus **Gliocladium 4: 84**
- (2) Fertile hyphae not verticillate at tip
Briarea 4: 85
- II. Conidia borne irregularly on simple or branched but not inflated or verticillate hyphae **Botrytidae**
- I. Conidia smooth or scarcely roughened
 - a. Saprogenous
 - (1) Conidia typically pleurogenous
 - (a) Fertile hyphae 2-several-furcate **Haplaria 4: 85**
 - (b) Fertile hyphae simple or nearly so
 - x. Conidia globose or ellipsoid **Acladium 4: 87**
 - y. Conidia short cylindric **Cylindrotrichum 4: 88**
 - (2) Conidia acrogenous or pleurogenous
 - (a) Some intermediate joints of the hyphae swollen and denticulate conidia-bearing **Physospora 4: 88**
 - (b) Intermediate joints equal
 - x. Conidia-bearing hyphae of two sorts, the upright alone denticulate **Blastomyces 10: 529**
 - y. Conidia-bearing hyphae of one sort
 - (x) Fertile hyphae simple or nearly so
 - m. Hyphae not denticulate; conidia solitary
 - (m) Hyphae forming a crust-like stratum **Hyphoderma 4: 89**
 - (n) Hyphae loose, cobwebby **Acremonium 4: 89**
(incl. **Thermomyces 18: 524**)
 - n. Hyphae denticulate; conidia usually grouped
 - (m) Hyphae everywhere denticulate, bearing conidia only at tip **Xenopus 18: 524**
 - (n) Hyphae denticulate or proliferous at tip alone
 - r. Apex denticulate, many-spored **Rhinotrichum 4: 91**
 - s. Apex inflated-ampulliform, 1-spored **Olpitrichum 11: 594**
 - (y) Fertile hyphae branched
 - m. Conidia globose to ovoid
 - (m) Both sterile and fertile hyphae procumbent
 - r. Sterile hyphae intracellular **Hartigiella 16: 1031**
 - s. Sterile hyphae superficial
 - (r) Fertile hyphae vaguely branched
 - h. Conidia acro-pleurogenous **Sporotrichum 4: 96**
(incl. **Leiosepium 16: 1036**)
 - i. Conidia on a one-sided sympodium **Monopodium 10: 544**

- (s) Fertile hyphae dichotomous; conidia acrogenous on spine-like branches **Langloisula** 10: 535
 - (n) Fertile hyphae erect or ascending
 - r. Conidia solitary acrogenous
 - (r) Fertile hyphae spiny-branched at apex **Plectothrix** 18: 525
 - (s) Fertile hyphae not spiny-branched **Monosporium** 4: 113
(incl. *Allescheriella* 14: 1075)
 - s. Conidia loosely grouped about the apex
 - (r) Conidia not involved in mucus
 - h. Conidia on inflated muriculate apices **Phymatotrichum** 16: 1033
 - i. Apices not muriculate or inflated **Botrytis** 4: 116
 - (s) Conidia involved in mucus **Tolypomyria** 4: 137
 - n. Conidia fusoid to cylindric
 - (m) Fertile hyphae mostly procumbent **Sporotrichella** 10: 534
 - (n) Fertile hyphae erect or ascending
 - r. Conidia fusoid on the upper side of curved branches **Martensella** 4: 138
 - s. Conidia acrogenous
 - (r) Conidia-bearing branches terete **Cylindrophora** 4: 138
 - (s) Conidia-bearing branches ellipsoid **Cylindrodendrum** 4: 139
 - b. Biogenous
 - (1) Conidia smooth, solitary, more rarely subcatenate **Ovularia** 4: 139
(incl. *Ovulariopsis* 16: 1036)
 - (2) Conidia densely spiny **Ramulaspera** 18: 532
 - 2. Conidia muricate or tuberculose-stellate
 - a. Conidia globose
 - (1) Conidia merely muricate
 - (a) Hyphae loose, cobwebby **Sepedonium** 4: 146
 - (b) Hyphae woven into a subgelatinous pellicle **Pellicularia** 4: 149
 - (2) Conidia setose at apex as well as muricate **Chaetoconidium** 10: 544
 - b. Conidia tuberculose-stellate **Asterophora** 4: 148
- III. Conidia acrogenous on verticillate branches **Verticilliae**
- 1. Conidia solitary or loosely grouped, not in chains

- a. Conidia-bearing branches very short, ampulliform
Pachybasium 4: 149
 - b. Conidia-bearing branches terete or longer
 - (1) Conidia globose to ovoid
 - (a) Tips of branches clavate, in twos rectangularly
Verticilliosis 11: 600
 - (b) Tips of branches normal
 - x. Conidia conglutinate into a stratum
Corymbomyces 18: 533
 - y. Conidia not conglutinate
 - (x) Conidia separating readily from the tips
Verticillium 4: 150
 - (y) Conidia separating with difficulty from the tips
Cladobotryum 4: 160
 - (2) Conidia cylindric or elongate
 - (a) Conidia-bearing branches or sporophores 1-spored
 - x. Sporophores straight
Acrocylindrium 4: 161
 - y. Sporophores uncinat
Uncigera 4: 162
 - (b) Sporophores several-spored
 - x. Sporophore inflated verrucose at apex
Calcarisporium 4: 162
 - y. Sporophore incurved, with seriate conidia below
Coemansia 4: 162
 - 2. Conidia capitate or densely spicate, not in chains
 - a. Conidia sessile
 - (1) Conidia capitate, involved in mucus
 - (a) Fertile hyphae smooth
Acrostalagmus 4: 163
(incl. Harziella 16: 1037)
 - (b) Fertile hyphae asperate
Gloeosphaera 18: 535
 - (2) Conidia densely spirally spicate at apices
Clonostachys 4: 165
 - b. Conidia on small stalks
Sceptromyces 4: 166
 - 3. Conidia in chains
Spicaria 4: 166
(incl. Nomuraea 18: 533)
- IV. Joints of the hyphae inflated here and there and bearing pleurogenous conidia
Gonatobotrytae
- 1. Joints smooth
 - a. Conidia catenulate
Gonatorrhodum 4: 169
 - b. Conidia solitary
Nematogonium 4: 170
 - 2. Joints muricate or punctate
 - a. Conidia solitary
Gonatobotrys 4: 168
 - b. Conidia catenulate, forming a spheric head
Gonatorrhodiella 10: 548
- Hyalodidymae**
4: 176, 10: 548, 11: 600, 14: 1057, 16: 1038, 18: 539
Conidia hyaline or bright-colored, 1-septate, ovoid oblong or short fusoid

I. Conidia not in chains

1. Saprophilus

a. Conidia smooth

(1) Fertile hyphae simple or nearly so

(a) Hyphae inflated at apex or joints

x. Hyphae denticulate inflated at apex; conidia fusoid

Diplorhinotrichum 18: 540

y. Hyphae inflated at both apex and joints

Arthrobotrys 4: 181

(b) Hyphae not inflated

x. Conidia spirally pleurogenous

Haplariopsis 18: 539

y. Conidia solitary acrogenous or capitate

(x) Conidia capitate at apex

Cephalothecium 4: 180

(y) Conidia solitary at apex

m. Fertile hyphae long

Trichothecium 4: 178

n. Fertile hyphae very short

Didymopsis 4: 182

(2) Fertile hyphae branched

(a) Fertile hyphae irregularly branched

Diplosporium 4: 178

(b) Fertile hyphae verticillate or dichotomous

x. Fertile hyphae verticillate

Diplocladium 4: 176

y. Fertile hyphae dichotomous; sterigmata subternate

Cylindrocladium 11: 600

b. Conidia echinulate; conidial cells unequal

Mycogone 4: 183

2. Biophilous

a. Conidia obliquely beaked

Rhynchosporium 18: 540

b. Conidia not beaked

(1) Hyphae mostly simple, not spirally twisted

Didymaria 4: 184

(2) Hyphae simple, spirally twisted

Bostrichonema 4: 185

II. Conidia catenulate

1. Fertile hyphae simple, short

Hormiactis 4: 186

2. Fertile hyphae verticillately branched

Didymocladium 4: 186**Hyalophragmiae**

4: 188, 10: 551, 11: 601, 14: 1059, 16: 1041, 18: 544

Conidia hyaline or bright-colored, 2-several-septate, oblong, fusoid or elongate

Micronemeae

Fertile hyphae very short and little different from the conidia

I. Conidia in chains, cylindric or oblong

Septocylindrium 4: 223

II. Conidia not in chains

1. Sporophore 3-celled, upper cell much inflated

Milowia 4: 222

2. Sporophore not inflated, sometimes obsolete

- a. Conidia ciliate at apex and upper septum
Mastigosporium 4: 220
- b. Conidia not ciliate
 - (1) Hyphae lacking; conidia not aggregate
Fusoma 4: 220
 - (2) Hyphae distinct; conidia aggregate
 - (a) Conidia in mucose glomerules
Rotaea 4: 222
 - (b) Conidia in fascicles, not mucose
Paraspora 4: 222

Macronemeae

Fertile hyphae manifest and distinct from the conidia

I. Saprophilous

- 1. Conidia solitary or at least not capitate
 - a. Fertile hyphae simple
 - (1) Sterile hyphae lacking
Dactylella 4: 193
 - (2) Sterile hyphae abundant
Monacrosporium 4: 193
 - b. Fertile hyphae branched
 - (1) Hyphae verticillately branched
Dactylium 4: 188
 - (2) Hyphae irregularly branched
Blastotrichum 4: 191
- 2. Conidia capitate
 - a. Fertile hyphae vesiculose at tip; fimicole
Cephalophora 18: 544
 - b. Fertile hyphae not swollen
 - (1) Hyphae simple; sterile lacking
Dactylaria 4: 194
 - (2) Hyphae verticillate; sterile hyphae present
Mucrosporium 4: 190

II. Biophilous

- 1. Conidia mucose-conglobate, allantoid, often continuous
Allantospora 14: 1043
- 2. Conidia not mucose-conglobate
 - a. Conidia ciliate at apex
***Trichoconis 18: 545**
 - b. Conidia not ciliate
 - (1) Conidia ovate-cylindric or elongate, often catenate
Ramularia 4: 196
 - (2) Conidia obclavate-piriform
Piricularia 4: 217
 - (3) Conidia long vermiform
Cercosporella 4: 218

Hyalodictyae

11: 608, 18: 561

Conidia hyaline, or bright-colored, muriform, ovoid to globose or cubic

- I. Hyphae much branched; conidia elliptic or globose, cells uniform
Stemphyliopsis 18: 561
- II. Hyphae little branched; conidia six-lobed and sarciniform, central cell larger, colored, lobes hyaline
Synthetospora 11: 608

Staurospora

4: 230, 10: 567, 11: 608, 14: 1067, 16: 1049, 18: 559

Conidia hyaline or bright-colored, stellate, radiate or forked, septate or continuous

- I. Hyphae lacking; conidia trident-shaped **Tridentaria 4: 231**
- II. Hyphae present
 - 1. Conidia globose to cylindric, permanently attached to 2-3 divergent sterigmata **Tetracladium 14: 1067**
 - 2. Conidia themselves stellate or radiate
 - a. Conidia bilobate-forked; lobes parallel, contiguous **Pedilospora 18: 559**
 - b. Conidia narrowly digitate **Prismaria 4: 230**
 - c. Conidia 3-4-radiate
 - (1) Conidia ciliate at the apex **Titaea 4: 231**
 - (2) Conidia muticate
 - (a) Conidia 3-radiate **Trinacrium 4: 231**
 - (b) Conidia 4-radiate
 - x. Fertile hyphae very short, simple **Tetracium 18: 560**
 - y. Fertile hyphae branched **Lemonniera 14: 1067**

Helicosporae

4: 233, 10: 568, 11: 608

Conidia hyaline or bright-colored, spirally curved, cylindric

- I. Hyphae very short; conidia spiral **Helicomycetes 4: 233**
- II. Hyphae various; conidia spirally twisted into a conic or ovate tube **Helicoum 11: 609**

Family 76. DEMATIACEAE

Hyphae dark or black, cobwebby, loose, usually rigid, not cohering in definite fascicles; conidia typically dark and concolorous, but sometimes the hyphae are dark and conidia clear, or the conidia dark and the hyphae clear. This family is parallel with the Moniliaceae and certain intermediate forms must be sought in both places.

Amerosporae

2: 235, 10: 569, 11: 610, 14: 1068, 16: 1059, 18: 563

Conidia dark, or sometimes hyaline but the hyphae then dark, 1-celled, globose to oblong.

Micronemeae

Hyphae very short or scarcely different from the conidia.

- I. Conidia not in chains
 - 1. Conidia globose to elliptic
 - a. Sterile hyphae nearly obsolete **Coniosporium 4: 238**
 - b. Sterile hyphae elongate **Cordella 10: 586**
 - 2. Conidia elongate, usually fusoid **Fusella 4: 246**
- II. Conidia in chains
 - 1. Conidia of two sorts, larger catenate, smaller glomerate **Heterobotrys 4: 267**
 - 2. Conidia all alike

a. Hyphae dark

(1) Chains breaking up readily

(a) Conidia globose or ovoid

Torula 4: 247

(b) Conidia clavate

Gongromeriza 4: 263

(2) Chains breaking up with difficulty

(a) Chains curved

Gyroceras 4: 266

(b) Chains straight or nearly so

Hormiscium 4: 263

b. Hyphae hyaline

Torulina 18: 566

III. Conidia in heads or racemes; conidia usually piriform

Echinobotryum 4: 268**Macronemeae**

Hyphae manifest and distinct from the conidia

I. Conidia dark, rarely subhyaline

i. Conidia not in chains

a. Conidia capitate

(1) Fertile hyphae simple, but often with short apical branches

(a) Hyphae with apical branches or basidia

x. Biophilous

Periconiella 4: 275

y. Saprophilous

(x) Apex with heterogeneous basidia

m. Apex swollen; basidia 3-4

Haplobasidium 10: 578

n. Apex not swollen; basidia many

Stachybotrys 4: 269

(y) Apex short-branched, rarely simple

m. Apex short-branched or simple

(m) Apex not swollen

Periconia 4: 270

(n) Apex swollen

Stachybotryella 18: 570

n. Apex capitate-branched; branches 2-3-furcate and spine-bearing

Cephalotrichum 4: 275

(b) Hyphae without apical branches or basidia

x. Conidia globose

Trichobotrys 18: 571

y. Conidia boat-shaped curved; hyphae dark-ringed

Camptoum 4: 276

z. Conidia fusoid, sometimes subhyaline

Acrotheca 4: 276

(2) Fertile hyphae branched below the apex

(a) Hyphae forked below apex; conidia oblong

Synsporium 4: 278

(b) Hyphae repeatedly dichotomous; conidia globose or elliptic

Dicyma 18: 570

b. Conidia verticillate-pleurogenous

(1) Hyphae dark nodose-inflated; conidia ovoid

Gonatobotryum 4: 278

(2) Hyphae hyaline, dark-ringed

(a) Conidia globose-angulose

Goniosporium 4: 280

(b) Conidia fusoid

Arthrimum 4: 279

- c. Conidia inserted irregularly
 - (1) Hyphae loose, typically saprogenous
 - (a) Hyphae vesiculose-inflated here and there
 - x. Conidia-bearing vesicles pleurogenous
Oedemium 4: 297
 - y. Conidia-bearing vesicles acrogenous
Cystophora 4: 298
 - (b) Hyphae not vesiculose-inflated
 - x. Fertile hyphae erect
 - (x) Branches circinate at apex; conidia mesogenous, muricate
Acrospira 4: 282, 14: 1056
 - (y) Branches spirally twisted; conidia exogenous
Streptothrix 4: 282
 - (z) Hyphae simple or with straight branches
Virgaria 4: 280
 - y. All hyphae more or less creeping
 - (x) Branches curved or lash-like
Campsotrichum 4: 295
 - (y) Branches not curved
 - m. Conidia spiny, rarely smooth
Zygodemus 4: 283
 - n. Conidia smooth
 - (m) Conidia sessile
Trichosporium 4: 288
 - (n) Conidia on stalks or basidia
 - r. Conidia on tooth-like sterigmata
Rhinocladium 4: 295
 - s. Conidia on jar-like basidia
Basisporium 18: 533
 - (2) Hyphae forming a crust, biogenous
Glenospora 4: 298
 - d. Conidia solitary, acrogenous
 - (1) Fertile hyphae simple
 - (a) Sterile hyphae lacking
 - x. Fertile hyphae short and fascicled at base
Hadrotrichum 4: 301
 - y. Fertile hyphae longer, separate
Monotospora 4: 299
 - (b) Sterile hyphae present
 - x. Conidia with a loose hyaline membrane
†Phaeoconis 18: 571
(Nigrospora)
 - y. Conidia without a membrane
 - (x) Conidia with a large shining gutta
Sporoglena 14: 1074
 - (y) Conidia without a shining gutta
Acremoniella 4: 302
(incl. Cordella 10: 586)

- (2) Hyphae branched; conidium at first enclosed in a vesicle from which it escapes at the apex **Conioscypha 18: 572**
2. Conidia in chains
- a. Sterile hyphae all creeping or obsolete
- (1) Conidia of two kinds; larger catenulate fuscous, smaller internal catenulate cylindric hyaline **Thielaviopsis 11: 612**
- (2) Conidia all alike
- (a) Conidia produced in the hyphae **Sporendonema 10: 515**
- (b) Conidia produced on the hyphae
- x. Fertile hyphae spirally twisted, forming a head of conidia **Helicocephalum 10: 512**
- y. Fertile hyphae not twisted
- (x) Fertile hyphae simple, not branched at tip
- m. Chains of conidia lateral **Dematium 4: 308**
- n. Chains terminal
- (m) Conidia without isthmi **Catenularia 4: 303**
- (n) Conidia connected by cylindric isthmi **Prophytroma 4: 309**
- (y) Fertile hyphae branched
- m. Hyphae dendroid **Hormodendrum 4: 310**
- n. Hyphae capitate branched at tip **Haplographium 4: 304**
- b. Some sterile hyphae erect and mixed with the fertile **Hormiactella 4: 311**
- II. Conidia hyaline or subhyaline
1. Conidia acrogenous on short heteromorphic basidia at the lower part or at the base of erect hyphae
- a. Conidia capitate glomerate
- (1) Sterile hyphae simple and circinate at apex **Bolacotricha 4: 316**
- (2) Sterile hyphae much branched below **Myxotrichum 4: 317**
- b. Conidia not capitate
- (1) Conidia solitary
- (a) Eruptent; conidia fusoid, usually setose **Ellisiella 4: 315**
- (b) Superficial
- x. Sterile hyphae simple
- (x) Conidia globose **Botryotrichum 4: 313**
- (y) Conidia bacillar
- m. Sterile hyphae tortuous **Sarcopodium 4: 312**
- n. Sterile hyphae circinate at apex **Helicotrichum 4: 313**
- y. Sterile hyphae branched
- (x) Hyphae irregularly branched; basidia verticillate **Costantinella 16: 1054**
- (y) Hyphae repeatedly dichotomous

- m. Branches continuous; basidia terete, basal
Circinotrichum 4: 314
 - n. Branches septate; basidia ampulliform, above base
Ceratocladium 4: 315
 - (2) Conidia loosely catenate; conidia basilar, ovoid
Stirochaete 4: 316
 - 2. Conidia on hyphae of the same kind
 - a. Conidia solitary, neither catenate or capitate
 - (1) Hyphae erect, simple
 - (a) Hyphae with a single lateral basidium near base
Zygosporium 4: 328
 - (b) Hyphae with pleurogenous conidia
Chloridium 4: 320
 - (2) Hyphae branched
 - (a) Hyphae erect, smooth
 - x. Hyphae verticillate branched **Verticicladium 4: 327**
 - y. Hyphae more or less irregularly branched
 - (x) Conidia ovoid **Mesobotrys 4: 324**
 - (y) Conidia cylindric **Chaetopsis 4: 324**
 - (z) Conidia falcate, sometimes ciliate
Menispora 4: 325
 - (b) Hyphae somewhat decumbent, more or less spiny
 - x. Hyphae nodose-spiny here and there
Gonytrichum 4: 329
 - y. Hyphae spiny but not swollen **Cladorrhinum 4: 330**
 - b. Conidia capitate
 - (1) Hyphae simple, with basidia only at the tip
 - (a) Conidia globose
 - x. Basidia verticillate **Fuckelina 4: 330**
 - y. Basidia irregular **Pimina 16: 1054**
 - (b) Conidia ovoid, mucose **Scopularia 4: 330**
 - (2) Hyphae more or less verticillate branched
Stachylidium 4: 331
 - c. Conidia catenate, arising within the hyphae
 - (1) Conidia in simple chains **Chalara 4: 333**
 - (2) Conidia conglutinate into a long curl
Cirromyces 18: 627
- Didymosporae**
- 4: 341, 10: 595, 11: 616, 14: 1077, 16: 1056, 18: 575
- Conidia 1-celled, dark, more rarely hyaline, ovoid to oblong
- Micronemeae**
- Hyphae very short or scarcely different from the conidia.
- I. Conidia not in chains
 - 1. Hyphae lacking **Dicoccum 4: 342**
 - 2. Hyphae present, circinate **Cycloconium 4: 343**
 - II. Conidia in chains **Bispora 4: 343**

Macronemeae

Hyphae distinctly different from the conidia

- I. Conidia smooth, muticate
 - i. Conidia not capitate
 - a. Conidia more or less catenulate at first
 - (1) Hyphae and conidia biform, the latter 1-celled dark or continuous hyaline
Epochnium 4: 375
 - (2) Hyphae and conidia uniform
 - (a) Hyphae here and there inflated **Cladotrichum 4: 370**
 - (b) Hyphae not inflated
 - x. Hyphae erect; conidia long-catenate
Diplococcium 4: 374
 - y. Hyphae somewhat decumbent; conidia short-catenate or finally solitary
Cladosporium 4: 350
 - b. Conidia not catenate
 - (1) Hyphae beautifully flexuose-torulose
Polythrincium 4: 350
 - (2) Hyphae not torulose or flexuose
 - (a) Hyphae inflated at tip, branched
Pseudobeltrania 18: 578
 - (b) Hyphae not inflated, usually short and little branched
 - x. Conidia merely acrogenous
Fusicladium 4: 345
(incl. *Passalora* 4: 344)
 - y. Conidia acro-pleurogenous
Scolecotrichum 4: 347
 2. Conidia capitate
Cordana 4: 376
 - II. Conidia muriculate or ciliate
 1. Conidia muriculate
Trichocladium 4: 376
 2. Conidia ciliate at apex; fertile and sterile hyphae intermixed
Beltrania 4: 377

Phragmosporae

4: 380, 10: 606, 11: 621, 14: 1082, 16: 1060, 18: 581

Conidia 2-several-septate, dark, rarely hyaline, ovoid to cylindric or vermicular

Micronemeae

Fertile hyphae very short or little different from the conidia

- I. Conidia not in chains
 1. Conidia muticate
 - a. Conidia united at base, fasciculate, cylindric
Cryptocoryneum 4: 395
 - b. Conidia separate
 - (1) Conidia ovoid to cylindric
 - (a) Saprogenous
Clasterosporium 4: 382
 - (b) Phyllogenous
Stigmina 4: 394
 - (2) Conidia fusoid-falcate
Fusariella 4: 395
 2. Conidia cuspidate or setose

- a. Hyphae dichotomous and broadened at apex

Urosporium 4: 397

- b. Hyphae not dichotomous or broadened

Ceratophorum 4: 395

II. Conidia in chains

1. Conidia not connected by isthmi

Septonema 4: 397

2. Conidia connected by isthmi

Polydesmus 4: 401

Macronemeae

Fertile hyphae distinctly different from the conidia

I. Conidia solitary or nearly so, acrogenous for the most part

1. Conidia muticate

- a. Conidia echinulate

Heterosporium 4: 480

- b. Conidia smooth

(1) Biophilous

- (a) Hyphae creeping, radiate

Ophiotrichum 10: 617

- (b) Hyphae ascending or erect

- x. Conidia ovoid to oblong

Napicladium 4: 481

(incl. **Cercosporidium 18: 594**)

- y. Conidia filiform or vermicular

Cercospora 4: 431

(2) Saprophilous

- (a) Hyphae rigid; conidia ovoid to elongate

- x. Conidia ovoid

Brachysporium 4: 423

- y. Conidia elongate

Helminthosporium 4: 402

- (b) Hyphae flexuous, pannose

Drepanospora 4: 430

2. Conidia 1-3-ciliate at apex

Camposporium 4: 482

II. Conidia verticillate or capitate

1. Hyphae dark

- a. Conidia acrogenous, forming a head

- (1) Hyphae simple

Acrothecium 4: 483

- (2) Hyphae branched at the apex

Atractina 18: 584

- b. Conidia pleurogenous, somewhat verticillate

- (1) Hyphae rostrate and naked at apex

Rhynchomyces 18: 584

- (2) Hyphae not rostrate at apex

Spondylocadium 4: 482

2. Hyphae hyaline or bright-colored, apex denticulate

Neomichelia 18: 593

III. Conidia catenate as a rule

1. Conidia arising from the interior of the hyphae

Sporoschisma 4: 486

2. Conidia arising from the apex, sometimes solitary

Dendryphium 4: 487

Dictyosporae

4: 496, 10: 665, 11: 632, 14: 1090, 16: 1075, 18: 612

Conidia dark, rarely hyaline, muriform, globose to oblong

Micronemeae

Hyphae very short or scarcely different from the conidia

I. Conidia not in chains

1. Conidia muticate

a. Conidia irregularly muriform or sarciniform

(1) Conidia with a conic point at each side

Oncopodium 18: 616

(2) Conidia muticate

(a) Conidia globose to oblong

x. Conidia ovoid to oblong, loose **Sporodesmium 4: 497**

y. Conidia globose to ovoid, aggregated

Stigmella 4: 507

(b) Conidia sarciniform, often coalescent

Coniothecium 4: 508

b. Conidia as if composed of parallel chains of cells

(1) Chains of conidia never separating

Dictyosporium 4: 513

(2) Chains of conidia separating

Spira 4: 514

2. Conidia corniculate at apex

Tetraploa 4: 516

II. Conidia in chains, often asperate or with isthmi

Sirodesmium 4: 516

Macronemeae

Hyphae distinctly different from the conidia

I. Conidia of the same form

1. Conidia not in chains or capitate

a. Conidia bearing little conidia on their surface

Xenosporium 18: 612

b. Conidia normal

(1) Hyphae alike

(a) Conidia cruciate-divided, verrucose

†**Tetracoccosporis 18: 617**

(**Tetracoccosporium**)

(b) Conidia muriform, typically smooth

x. Hyphae decumbent

Stemphylium 4: 519

y. Hyphae erect or ascending

(x) Conidia globose, pleurogenous

m. Conidia around the apex of the hyphae

Coccosporium 4: 542

n. Conidia conglobate around the base

Trichaegum 4: 542

(y) Conidia ovoid to oblong, mostly acrogenous

Macrosporium 4: 523

(incl. **Mystrosporium 4: 539**)

(2) Hyphae of two kinds, longer sterile, shorter fertile

Septosporium 4: 543

Dactylosporium 4: 545

2. Conidia capitate

- 3. Conidia catenate
 - a. Hyphae velvety, erect, subsimple; conidia caudate
Alternaria 4: 545
 - b. Hyphae crustose, various; conidia 2-celled; conidia-like ganglia sarciniform
Fumago 4: 547
- II. Conidia of two forms, dark sarciniform and subhyaline falcate
Sarcinella 4: 548

Staurosporae

4: 552, 11: 639, 14: 1107, 16: 1181, 18: 625

Conidia forked or stellate, usually dark, septate or continuous

- I. Conidia of two forms, small fusoid hyaline, large lobate many-celled, brown
Desmidiospora 10: 568

II. Conidia alike

- 1. Fertile hyphae present; conidia 3-4-radiate
Tripodsporium 4: 554

2. Fertile hyphae lacking

- a. Conidia on a cellular stroma, 2-4-digitate

Chiromyces 4: 554

b. Cellular stroma lacking

- (1) Conidia 3-several-radiate; xylogenous

Ceratosporium 4: 552

- (2) Conidia 2-radiate; phyllogenous

Hirudinaria 4: 553**Scolecosporae**

Conidia long-filiform or vermicular

- One genus
Cercospora 4: 431, 14: 1099

Helicosporae

4: 557, 10: 680, 11: 638, 14: 1107, 16: 1081, 18: 624

Conidia cylindric, spiral or convolute, typically septate, dark or hyaline

- I. Hyphae obsolete
Helicopsis 10: 680
- II. Hyphae present
 - 1. Conidia septate transversely
Helicosporium 4: 557
 - 2. Conidia muriform
Helicoma 11: 638

Family 77. STILBACEAE

Sterile hyphae creeping, scanty; fertile hyphae collected into stalk-like or stroma-like fascicles bearing conidia at the top, more rarely along the side, pale, bright-colored or dark.

Hyalostilbae

Hyphae and conidia pale or bright-colored, not dark or black

Amerosporae

4: 561, 10: 681, 11: 640, 14: 1107, 16: 1082, 18: 630

Conidia globose, elliptic or oblong, 1-celled, hyaline or pale, or bright-colored

- I. Conidial part distinctly capitate or at least terminal

1. Conidia not in chains
 - a. Head of conidia not gaping or splitting above
 - (1) Head not spiny
 - (a) Conidiophores of head normal
 - x. Conidia covered with mucus
 - (x) Synnema monocephalous
 - m. Conidiophores dendroid-verticillate
 - (m) Without distinct sterigmata

Dendrostilbella 18: 635
 - (n) With obpiriform sterigmata

Pirobasidium 18: 638
 - n. Conidiophores not dendroid-verticillate

Stilbum 4: 564
 - (y) Synnema polycephalous
 - m. Capitula on extremely short branches

Polycephalum 4: 575
 - n. Capitula on spreading subulate branches

Tilachlidium 4: 576
 - o. Capitula on erect branches

Corallodendrum 4: 576
 - y. Conidia without mucus
 - (x) Synnema monocephalous
 - m. Conidiophores spirally twisted

Martindalia 4: 578
 - n. Conidiophores more or less straight
 - (m) Conidia rhombic or biconic

Rhombostilbella 18: 636
 - (n) Conidia globose to fusoid

Ciliciopodium 4: 577
(incl. *Clavularia* 10: 686)
 - (y) Synnema polycephalous
 - m. Terrestrial, large, 1-2 cm.; conidia ovoid

Macrostilbum 16: 1083
 - n. Small, not terrestrial; conidia elongate-ovate

Chondromyces 4: 576
 - (b) Conidiophores conidium-like, septate; monocephalous

Atractiella 4: 578
 - (2) Head spiny with radiating spicules
 - (a) Spicules conic, granulate

Actiniceps 4: 579
 - (b) Spicules with many curved branches at middle

Heterocephalum 18: 642
 - b. Head of conidia persistent below, splitting above

Pilacre 4: 579
2. Conidia in chains
 - a. Synnema with conidia above; conidia without mucus
 - (1) Synnema not pubescent

Coremium 4: 581
(incl. *Pritzeliella* 18: 644)
 - (2) Synnema pubescent

Lasioderma 4: 584
 - b. Synnema with conidia below; conidia with mucus

Microspatha 10: 687

II. Conidial part cylindric or long-clavate

1. Conidia more or less equally scattered

a. Biophilous; sterigmata denticulate branched

Cladosterigma 11: 640

b. Saprophilous; sterigmata none or simple

Isaria 4: 584

2. Conidia in lateral heads or racemes

a. Conidia in racemes; synnema lobate

Peribotryum 4: 595

b. Conidia in heads

(1) Conidiophores with lateral nodes, usually escaping through the stomata

Helostroma 18: 630

(2) Conidiophores without nodes, usually entomophilous

Gibellula 11: 643**Didymosporae**

18: 645

Conidia 2-celled, hyaline, globose to oblong

I. Synnema cylindric, fimbriate at apex; conidia oblong

Didymobotryopsis 18: 645

II. Synnema capitate; conidia fusoid

Didymostilbe 18: 645**Phragmosporae**

4: 598, 10: 691, 14: 1109, 18: 646

Conidia 2-several-septate, hyaline, oblong to bacillar

I. Conidia solitary

1. Conidia bacillar, aristate above, separating at joints

Stilbomyces 14: 1109

2. Conidia not aristate or separating

a. Conidia oblong

Arthrosporium 4: 598

b. Conidia elongate-falcate

Atractium 4: 599

II. Conidia catenate, cylindric

Symphyosira 4: 600**Helicosporae**

18: 658

Conidia filiform, spirally twisted

I. Synnema erect, setose

Helicostilbe 18: 657**Phaeostilbae**

Hyphae and conidia or one or the other dark

Amerosporae

4: 603, 10: 692, 11: 643, 14: 1109, 16: 1086, 18: 648

Conidia 1-celled, dark, globose to elongate

I. Conidia not in chains

1. Synnema setose

Saccardaea 11: 643

2. Synnema naked

a. Conidia asperate, on minute basidia

Basidiella 10: 698

b. Conidia smooth

- (1) Synnema carnose, racemose-branched **Stilbothamnium 14: 1110**
- (2) Synnema fibrous or corneous, not racemose
 - (a) Basidia lageniform **Ceratocladium 18: 649**
 - (b) Basidia lacking, at least not lageniform
 - x. Synnema stalked, fibrous
 - (x) Conidia dark, globose to elliptic **Sporocybe 4: 604**
 - (y) Conidia hyaline
 - m. Conidia ovoid to oblong **Graphium 4: 609**
 - u. Conidia elongate or falcate **Harpographium 4: 619**
 - y. Synnema sessile, corneous **Glutinium 4: 620**

II. Conidia in chains

- 1. Synnema setose **Trichurus 14: 1112**
- 2. Synnema not setose
 - a. Stalk scopulate branched above **Stemmaria 10: 696**
 - b. Stalk simple or nearly so
 - (1) Capitule loose
 - (a) Base of synnema subequal; usually on stems **Stysanus 4: 620**
 - (b) Base of synnema perithecioid; usually on leaves **Graphiothecium 4: 624**
 - (2) Capitule compact
 - (a) Conidia globose
 - x. Conidia echinulate **Harpocephalum 14: 1111**
 - y. Conidia smooth
 - (x) Conidia pleurogenous **Heydenia 4: 625**
 - (y) Conidia acrogenous **Briosia 10: 698**
 - (b) Conidia ovoid to oblong **Antromycopsis 14: 1113**

Didymosporae

4: 626, 10: 699, 18: 654

Conidia 1-septate, dark or hyaline, oblong to cylindric

- I. Conidia muticate **Didymobotryum 4: 626**
- II. Conidia 1-ciliate at apex **Hoehneliella 18: 654**

Phragmosporae

4: 627, 10: 699, 11: 644, 14: 1113, 16: 1089, 18: 655

Conidia 2-several-septate, dark or hyaline, oblong to cylindric

- I. Conidia capitule
 - 1. Synnema simple
 - a. Synnema black; conidia densely capitate **Arthrobotryum 4: 628**
 - b. Synnema fuscous or pale; conidia loosely capitate **Isariopsis 4: 630**
 - 2. Synnema dendroid branched **Xylocladium 16: 1089**

II. Conidia not capitate

1. Conidia catenulate

Dendrographium 11: 644

2. Conidia not catenulate

a. Stalk fibrous

(1) Synnema simple or branched; conidia acro-pleurogenous

Podosporium 4: 627

(2) Synnema branched; conidia acrogenous

Negeriella 14: 1114

b. Stalk parenchyma-like

(1) Conidia pleurogenous, on a disk

Riccoa 18: 656

(2) Conidia acrogenous

Podosporella 11: 644**Dictyosporae**

4: 632

Conidia muriform, dark or hyaline, oblong

I. Synnema stalked, capitate

Sclerographium 4: 632**Staurosporae**

I. Conidia of 4-5-radiate cells, hyaline

Riessia 4: 627**Family 78. TUBERCULARIACEAE**

Hyphae compacted into a globose, discoid or verruciform body or sporodochium; sporodochia typically sessile, waxy or subgelatinous, white, bright-colored or dark to black.

Mucedinae

Hyphae and conidia white or bright-colored

Amerosporae

4: 635, 10: 700, 11: 645, 14: 1115, 16: 1090, 18: 658

Conidia hyaline or bright-colored, 1-celled, globose to fusoid

I. Sporodochia smooth or nearly so

1. Conidiophores normal

a. Conidia muticate

(1) Conidia not covered with mucus

(a) Conidia not acrogenous capitate

x. Sporodochium girt by a heterogeneous cup

Patellina 4: 677

y. Sporodochium without a heterogeneous cup

(x) Conidia not catenate or scarcely so

m. Conidia escaping from interior of hyphae

(m) Conidiophores branched **Endoconidium 10: 708**(n) Conidiophores simple **Trichotheca 10: 714**

n. Conidia arising on outside of hyphae

(m) Conidiophores lacking

r. Conidia large, pellucid

(r) Conidia globose

Sphaerosporium 4: 664

(s) Conidia oval

Diaphanium 4: 672

s. Conidia small, not pellucid

Pactilia 4: 672

- (n) Conidiophores present
 - r. Conidia pleurogenous or acro-pleurogenous
 - (r) Conidia globose **Beniowskia** 16: 1091
 - (s) Conidia ovoid to oblong
 . **Tubercularia** 4: 638
 - (t) Conidia fusoid to cylindric
Fusicolla 4: 664
 - s. Conidia acrogenous
 - (r) Conidiophores verrucose
Dacrymycella 4: 671
 - (s) Conidiophores not verrucose
 - h. Uredinicole **Tuberculina** 4: 653
 - i. Not uredinicole
 - (h) Sporodochia globose
 + Conidia globose; conidiophores short
Aegerita 4: 661
 - Conidia ovoid; conidiophores branched
Granularia 4: 649
 - (i) Sporodochia pulvinate
 + Conidia acicular
Kmetia 16: 1158
 - Conidia terete-oblong
Bactridiopsis 18: 662
 - (j) Sporodochia disk-shaped, or cupulate
 + Sporodochia disk-shaped
Hymenula 4: 667
(Hymenella 16: 1105)
 - Sporodochia cupulate
Hyphostereum 11: 649
 - (k) Sporodochia verruciform or effuse
 + Conidiophores simple
 (+) Conidiophores radiate, united at base
Clinoconidium 16: 1093
 - (—) Conidiophores not united or radiate
Sphacelia 4: 666
 - Conidiophores dendroid branched
Dendrodochium 4: 650
- (y) Conidia in chains
 - m. Conidia covered with mucus
Collodochium 18: 661
 - n. Conidia without mucus
 - (m) Conidia globose
 - r. Conidia hyaline **Sphaerocolla** 11: 648
 - s. Conidia blue **Sporoderma** 4: 676
 - (n) Conidia elliptic to oblong
 - r. Sporodochium disk-shaped, orange-red
Necator 16: 1094
 - s. Sporodochium subglobose, whitish
Patouillardia 4: 677

- (o) Conidia cylindric
 - r. Sporodochium dilated above, stalked
Bizzozzeriella 10: 713
 - s. Sporodochia globose to verruciform
 - (r) Sporodochia gelatinous, sessile
Cylindrocolla 4: 673
 - (s) Sporodochia not gelatinous, short-stalked
Sphaeridium 4: 675
 - (b) Conidia acrogenous capitate; sporodochia turbinate
Cephalodochium 4: 678
 - (2) Conidia covered with mucus
 - (a) Sporodochium globose, hardened
Thecospora 4: 679
 - (b) Sporodochia verruciform or discoid, gelatinous or waxy
 - x. Sporodochia verruciform or subeffuse
Illosporium 4: 656
(incl. **Myxonema** 10: 714)
 - y. Sporodochia discoid
Epidochiopsis 11: 648
 - b. Conidia ciliate
 - (1) Conidia 1-ciliate at base only
Stigmatella 4: 679
 - (2) Conidia ciliate at both ends
 - (a) Conidia 1-ciliate at each end
Thozetia 4: 679
 - (b) Conidia 7-8-ciliate at each end
Chaetospermum 10: 706
 - 2. Conidiophores with internal conidia-bearing areoles
Scoriomyces 4: 680
- II. Sporodochia setulose, ciliate or uniformly woolly
- 1. Sporodochia woolly or setulose
 - a. Sporodochia setulose; conidia catenate
Periola 4: 681
 - b. Sporodochia woolly or velvety; conidia capitate
 - (1) Conidia globose
Dacryodochium 14: 1122
 - (2) Conidia oblong
Lachnodochium 14: 1122
 - 2. Sporodochia ciliate at the margin
 - a. Sporophores none; conidia coacervate
Volutellaria 4: 682
 - b. Sporophores distinct
 - (1) Conidia in chains
Volutina 18: 667
 - (2) Conidia not in chains
 - (a) Conidiophores 6-ciliate above, united below
Guelichia 10: 720
 - (b) Conidiophores not ciliate or united
Volutella 4: 682
- Didymosporae**
4: 690, 10: 721, 18: 668
- Conidia 1-septate, hyaline or bright-colored
- I. Conidia in chains
 - 1. Sporodochia setulose
Endodesmia 4: 691
 - 2. Sporodochia smooth
Gymnodochium 18: 668

II. Conidia not in chains

1. Sporodochia setulose

Leptotrichum 4: 690

2. Sporodochia smooth

a. Conidia verrucose

Cosmariospora 4: 690

b. Conidia smooth

Patouillardia 10: 721**Phragmosporae**

4: 691, 10: 721, 11: 649, 14: 1123, 16: 1097, 18: 669

Conidia 2-several-septate, hyaline or bright-colored, fusoid to falcate (in *Fusarium* sometimes short and simple).

I. Conidia somewhat catenate, cylindric

Discocolla 11: 653

II. Conidia rarely catenate

1. Conidia cruciately 4-celled; sporodochium gelatinous

Sarcinodochium 18: 677

2. Conidia not cruciate

a. Conidiophores short, simple

(1) Conidia very large, terete-oblong

Bactridium 4: 691

(2) Conidia doliiform

Pithomyces 4: 693

b. Conidiophores more or less branched

(1) Conidiophores dichotomous; conidia key-like

Heliscus 4: 693

(2) Conidiophores usually verticillately branched; conidia usually falcate, sometimes oblong

(a) Sporodochium gelatinous

Pionnotes 4: 725

(b) Sporodochium waxy or byssoid

Fusarium 4: 694(incl. *Microcera* 4: 727)**Dictyosporae**

18: 676

Conidia muriform, hyaline, subglobose

I. Sporodochia globose

Sporocystis 18: 676**Staurosporae**

4: 728, 16: 1104, 18: 677

Conidia forked or cruciate, hyaline or bright-colored

I. Conidiophores simple; conidia horseshoe-like

Lituarina 4: 728

II. Conidiophores branched

1. Conidia with short irregular branches or lobes

Aegeritopsis 18: 677

2. Conidia forked or cruciate

a. Conidia 2-forked, septate

Dicranidium 4: 728

b. Conidia 3-forked or subcruciate, continuous

Triglyphium 4: 728**Helicosporae**

4: 729, 10: 732, 11: 653, 18: 678

Conidia spirally convolute

- I. Conidiophores lacking **Everhartia 4: 729**
- II. Conidiophores present
1. Conidia continuous **Troposporium 4: 729**
 2. Conidia septate **Hobsonia 11: 653**

Dematiace

Hyphae olive, to brown or black; conidia concolorous, rarely hyaline

Amerosporae

4: 736, 10: 732, 11: 654, 14: 1129, 16: 1104, 18: 678

Conidia 1-celled, globose to elongate, sometimes unequal

- I. Conidia not in chains
1. Sporodochia not setose
 - a. Conidiophores lacking
 - (1) Lichenicole **Spilomium 18: 678**
 - (2) Not lichenicole
 - (a) Sporodochia gelatinous; conidia globose, vesiculose
Myriophysa 4: 742
 - (b) Sporodochia not gelatinous
 - x. Sporodochia hemispheric, with a stratum of conidia
Spermodermia 4: 742
 - y. Sporodochia disk-like, applanate **Sclerodiscus 10: 735** ..
 - b. Conidiophores present
 - (1) Sporodochia thick, tremelloid **Epidochium 4: 747**
 - (2) Sporodochia not tremelloid
 - (a) Conidiophores with a slender apical appendage; conidia globose
Bonplandiella 10: 732
 - (b) Conidiophores not appendaged
 - x. Conidia globose
 - (x) Sporodochia cellular, uniform
Epicoccum 4: 736
 - (y) Sporodochia of three hyphal layers
Triplicaria 10: 734
 - y. Conidia ovoid to bacillar
 - (x) Conidiophores bacillar; sporodochia subdiscoid
Hymenopsis 4: 744
 - (y) Conidiophores branched
 - m. No brown radiate hyphae at base
Strumella 4: 742
 - n. Brown radiate hyphae at base
Astrodochilum 14: 1117
 2. Sporodochia ciliate or with exserted hypae
 - a. Sporodochia with loose exserted conidiophores, verruciform
Trichostroma 4: 752
 - b. Sporodochia margined with hairs or setae
 - (1) Setae dark **Chaetostroma 4: 749**
 - (2) Setae or hairs white **Myrothecium 4: 750**

II. Conidia in chains

1. Conidiophores lacking **Exosporina** 18: 684
2. Conidiophores present
 - a. Sporodochium tremelloid ***Hormodochis** 4: 749
 - b. Sporodochium not tremelloid
 - (1) Sporodochium ciliate ***Chaetodochis** 4: 750
 - (2) Sporodochium not ciliate
 - (a) Sporodochia globose **Sphaeromyces** 4: 753
 - (b) Sporodochia stellate **Actinomma** 4: 753

Didymosporae

4: 754, 10: 737, 16: 1105, 18: 684

Conidia 1-septate, typically dark, elliptic to fusoid

- I. Sporodochia lichenicole, globose **Sclerococcum** 4: 754
- II. Sporodochia not lichenicole
 1. Sporodochia foliicole
 - a. Sporodochia annuliform asteroid **Hyphaster** 18: 685
 - b. Sporodochia subglobose **Pucciniopsis** 10: 737
 2. Sporodochia lignicole **Epiclinium** 4: 754

Phragmosporae

4: 755, 10: 738, 11: 656, 14: 1131, 16: 1106, 18: 685

Conidia 2-several-septate, usually colored, oblong to cylindric

- I. Conidia in chains; sporodochium discoid **Trimmatostroma** 4: 757
- II. Conidia not in chains
 1. Conidia 1-ciliate at each end **Ciliofusarium** 11: 656
 2. Conidia muticate
 - a. Sporodochium hairy **Excipularia** 18: 688, 3: 689
 - b. Sporodochium smooth
 - (1) Conidia laterally proliferate and joined in bundles **Amallospora** 14: 1131
 - (2) Conidia not proliferate and united
 - (a) Sporodochia convex-pulvinate **Exosporium** 4: 755
 - (b) Sporodochia vertically cylindric or clavate **Listeromyces** 18: 685

Dictyosporae

4: 758, 10: 739, 11: 656, 14: 1131, 16: 1107, 18: 689

Conidia muriform, usually dark

- I. Conidia in chains **Bonordeniella** 18: 689
- II. Conidia not in chains
 1. Sporodochia setulose **Chaetostromella** 11: 656
 2. Sporodochia smooth **Spegazzinia** 4: 758

Scolecosporae

18: 689

Conidia filiform, hyaline

- I. Sporodochia globose, setulose **Schizotrichum** 18: 688

Staurosporae

4: 753

Conidia angulose-stellate, hyaline

- I. Sporodochia scutellate, pilose **Stephanoma 4: 753**

Helicosporae

11: 654

Conidia spirally twisted, smoky

- I. Sporodochia pulvinate **Troposporella 11: 654**

Sterile Mycelia

14: 1138, 16: 1108, 18: 690

Conidia permanently absent so far as known

- I. Parasitic on algae **Lepraria, Pulveraria, etc. Z. 239**

- II. Not parasitic on algae

1. Tubercle-like

- a. Tubercles connected with fibrils

Rhizoctonia 14: 1175**(Coccobotrys 16: 1108)**

- b. Tubercles without fibrils

- (1) Cortex discrete

Acinula 14: 1174

- (2) Cortex not discrete

Sclerotium 14: 1139

2. Maculiform; black stromata in leaves and stems

Ectostroma 14: 1177

3. Root-like

- a. Filaments rigid, broad, terete or depressed, dark, white within

Rhizomorpha 14: 1180

- b. Filaments rigid, capilliform, dark, closely adhering

Capillaria 14: 1184

4. Clavariform; filaments terete, vertical, simple or branched

Anthina 14: 1184

5. Cobwebby or byssoid

- a. Cespitose interwoven, primary hyphae joined in bundles

Ozonium 14: 1187

- b. Cespitose interwoven, hyphae not fasciculate, black

Rhacodium 14: 1189

- c. Cobwebby, soft, fleeting, white or pale

Hypha 14: 1192

- d. Adpressed, creeping, dendritic, white to brownish, not forming a continuous membrane

Himantia 14: 1194

6. Membrane-like; densely interwoven, forming a continuous suberose or coriaceous membrane

Xylostroma 14: 1197

7. Deformed, discolored corky cells of plants

Phloeconis 14: 1197

Key to Spore Sections

- Amerosporae: spores one-celled, not stellate or spiral
 - Allantosporae: spores sausage-shaped, mostly clear
 - Hyalosporae: spores hyaline or clear, globose to oblong
 - Phaeosporae: spores dark, yellow, brown or black, globose to oblong
 - Leucosporae: spores clear, rarely faintly colored
 - Rhodosporae: spores rose-colored
 - Ochrosporae: spores yellow to yellow-brown
 - Melanosporae: spores dark purple to black
- Didymosporae: spores 1-septate or 2-celled
 - Hyalodidymae: spores hyaline, 2-celled
 - Phaeodidymae: spores dark, 2-celled
- Phragmosporae: spores few-many-transeptate, 3-many-celled
 - Hyalophragmiae: spores hyaline, 3-many-celled
 - Phaeophragmiae: spores dark, 3-many-celled
- Dictyosporae: spores septate crosswise and lengthwise, i. e., muriform
 - Hyalodictyae: spores hyaline, muriform
 - Phaeodictyae: spores dark, muriform
- Scolecosporae: spores needle-shaped to filiform, continuous or septate
 - Hyaloscoleciae: spores hyaline, filiform
 - Phaeoscoleciae: spores dark, filiform
- Staurosporae: spores stellate or radiate, hyaline or dark, continuous or septate
- Helicosporae: spores spirally twisted, hyaline or dark, continuous or septate

Guide to the Volumes of Saccardo's "Sylloge Fungorum"

Volume I.

Bibliotheca Mycologica	IX-XIX
Pyrenomycetes: Perisporiaceae, Sphaeriaceae-Phaeodidymae.....	I-754

Volume II.

Pyrenomycetes: Sphaeriaceae-Phaeophragmiae, Hysteriaceae.....	I-809
Addenda to Volume I.....(815)	I-LV
Addenda to Volume II.....	LVI-LXIX
Index to genera in Volumes I and II.....	I-10
Index to species in Volumes I and II.....	II-77

Volume III.

Sphaeropsideae	I-695
Melanconieae	696-807
Index to genera in Volume III.....	813-816
Index to species in Volume III.....	817-860

Volume IV.

Hyphomycetes: Mucedineae-Tuberculariae	I-758
Index to genera in Volume IV.....	763-768
Index to species in Volume IV.....	769-807

Additions to Volumes I-IV.

Pyrenomycetes	I-273
Sphaeropsideae	285-360
Melanconieae	361-371
Hyphomycetes	372-392
Appendix 39	393-445
Index to genera in this volume.....	453-457
Index to species in this volume.....	459-484

Volume V.

Agaricineae	I-II44
Index to genera in Volume V.....	II45

Volume VI.

Hymenomyces: Polyporaceae-Tremellaceae	I-815
Index to genera in Volumes V and VI.....	817-824
Index to species in Volumes V and VI.....	825-928

Volume VII.

Part I:—

Gasteromycetes	I-180
Phycomycetes	180-322
Myxomycetes	323-468

Additions to Gasteromycetes	469-492
Index to genera in Volume VI, part 1.....	I-IV
Index to species in Volume VII, part 1.....	V-XXX

Part 2:—

Ustilagineae	449-524
Uredineae	528-869
Index to genera in Volume VII, part 2.....	XXXI-XXXIII
Index to species in Volume VII, part 2.....	XXXV-LIX

Volume VIII.

Conspectus Systematicus Generalis	XIII-XVI
Discomycetes	3-842
Phymatosphaeriaceae	843-847
Onygenaceae	861-862
Tuberoideae	863-908
Laboulbeniaceae	909-915
Saccharomycetes	916-922
Schizomycetes	923-1087
Index to genera in Volume VIII.....	1089-1095
Index to species in Volume VIII.....	1097-1143

Volume IX.

Hymenomycetes	1-261
Gasteromycetes	262-281
Hypodermiaceae	282-334
Phycomycetes	335-363
Pyrenomycetes	364-1129
Laboulbeniaceae	1130-1134
Index to genera in Volume IX	1135-1141

Volume X.

De nominibus generum	VII-IX
Bibliotheca Mycologica	XI-XXX
Discomycetes	1-79
Onygenaceae	80
Tuberoideae	80-83
Myxomycetes	83-99
Schizomycetes	100
Sphaeropsidae	100-446
Melanconieae	446-509
Hyphomycetes	510-739
Fossil Fungi	745-808
Index Universalis to Volumes I-X	841-869
Index to species in Volumes IX and X.....	871-964

Volume XI.

Hymenomycetes	I-151
Gasteromycetes	152-173
Hypodermieae	174-238
Phycomycetes	239-251
Pyrenomycetes	252-390

Discomycetes	391-439
Phymatosphaeriaceae	440
Onygenaceae	440
Tuberoideae	441-445
Laboulbeniaceae	446-456
Saccharomycetes	457-459
Myxobacteria	460-461
Myxomycetes	462-471
Sphaeropsidaceae	472-561
Melanconieae	562-585
Hyphomycetes	586-656
Fossil Fungi	657-659
Index to species in Volume XI.....	677-718
Index Universalis to Volumes I-XI.....	

Volume XII.

Index Universalis to genera, species and varieties in Volumes I-XI.

Abbreviations	
Fungi in living and dead parts of plants.....	1-858
Fungi in living and dead parts of man and animals.....	859-872
Fungi on dung and artificial substrata (textiles, etc.).....	873-902
Fungi on earth, stones, walls, turf and charcoal.....	903-1039
Fossil Fungi	1042-1053

Volume XIII.

Abbreviations.

Host Index to the close of 1897 (Volumes I-XI).

Volume XIV.

Table of orders and families.....	3-4
Table of spore sections and the genera included in them.....	5-62
Hymenomycetes	63-253
Gasteromycetes	254-268
Uredinaceae	269-409
Ustilaginaceae	410-430
Hyphostomaceae	431
Plycomycetes	432-461
Pyrenomycetes	462-724
Laboulbeniaceae	725-737
Discomycetes	738-825
Tuberoideae	826-828
Saccharomycetes	828
Protomycetes	829-830
Myxomycetes	831-841
Myxobacteria	842-844
Sphaeropsidaceae	844-1003
Melanconiaceae	1004-1036
Hyphomycetes	1037-1132
Appendix containing some omitted and new species.....	1133-1137
Sterile Mycelia	1138-1198

Index to species in Volume XIV.....	1219-1272
Index universalis to Volumes I-XIV.....	1277-1316

Volume XV.

Synonyms of genera, species and varieties in Volumes I-XIV.

Volume XVI.

Hymenomycetes	I-223
Gasteromycetes	224-256
Uredinaceae	257-366
Ustilaginaceae	367-382
Phycomycetes	383-397
Pyrenomycetes	398-673
Laboulbeniaceae	674-694
Discomycetes	695-817
Saccharomycetes	818
Myxomycetes	819-824
Deuteromycetes	825-1107
Sterile Mycelia	1108-1109
Appendix: new fungi	1110-1158
Index to species in Volume XVI.....	1179-1233
Index universalis to Volumes I-XVI.....	1239-1291

Volume XVII.

Bibliotheca Mycologica	IX-CVII
Hymenomycetes	I-211
Gasteromycetes	212-243
Uredinaceae	244-471
Ustilaginaceae	472-493
Phycomycetes	494-523
Pyrenomycetes	524-914
Laboulbeniaceae	915-924
Index to species in Volume XVII.....	941-986
Index to genera in Volume XVII.....	987-991

Volume XVIII.

Mycological diagnosis and nomenclature.....	III-VII
Discomycetes	I-207
Myxomycetes	208-217
Myxobacteria	217-219
Deuteromycetes	219-690
Sterile Mycelia	690-691
Index to species in Volume XVIII.....	707-740
Index universalis to Volumes I-XVIII.....	743-838

Index to Families in Saccardo's "Sylloge Fungorum and Rehm's "Discomyceten"

Acrospormaceae	 R: 53
Agaricaceae	 5: 3, 9: 3, 11: 1, 14: 63, 16: 1, 16: 1110, 17: 1
Ancylistaceae	 16: 395
Arthoniaceae	 10: 74, 16: 798, 18: 186, R. 414
Ascobolaceae	 8: 512, 10: 31, 11: 420, 14: 792, 16: 758, 1149, 18: 116, R. 1078
Ascocorticiaceae	 18: 198
Ascoidaceae	 16: 806
Bulgariaceae	 8: 607, 10: 38, 11: 425, 14: 801, 16: 766, 1150, 18: 131, R. 444
Caliciaceae	 8: 825, 10: 72, 11: 439, 14: 825, 18: 189, R. 388
Cenangiaceae	 R. 213
Cenococcaceae	 8: 871
Chytridiaceae	 7: 286, 9: 357, 11: 246, 14: 437, 16: 389, 1122, 17: 511
Clavariaceae	 6: 670, 9: 247, 11: 134, 14: 235, 16: 1116, 17: 193
Coccoideaceae	 17: 860
Cordieritaceae	 8, 810, 16: 803
Coryneliaceae	 9: 1073, 11: 385, 16: 650
Cyttariaceae	 8: 4, 16: 695, 18: 1
Dematiaceae	 4: 235, 9: 378, 10: 569, 11: 610, 14: 1068, 16: 1050, 1157, 18: 563
Dermateaceae	 8: 545, 10: 36, 11: 422, 14: 794, 16: 762, 1149, 18: 121, R. 241
Dothideaceae	 2: 588, A. 222, 9: 1004, 11: 368, 14: 663, 16: 616, 1144, 17: 827
Elaphomycetaceae	 8: 863, 10: 80, 11: 441
Endogonaceae	 8: 905
Endomycetaceae	 18: 202
Entomophthoraceae	.. 7: 280, 9: 349, 14: 437, 16: 388, 1123, 17: 510
Eotterfeziaceae	 18: 205
Euphacidiaceae	 R. 60
Euphacidiaceae	 R. 913
Eustictaceae	 R. 113
Excipulaceae	 3: 664, A. 358, 10: 432, 11: 558, 14: 999, 16: 993, 18: 436
Exoascaceae	 14: 823, 16: 803, 18: 196
Geoglossaceae	 R. 1142
Gymnoascaceae	 8: 811, 10: 67, 11: 435, 14: 824, 16: 805, 1152
Helotiaceae	 R. 647
Helvellaceae	 8: 7, 10: 1, 11: 391, 14: 738, 16: 695, 1146, 18: 2, R. 1134
Hemihysteriaceae	 A. 260, 9: 1094, 11: 385, 14: 707, 16: 653, 17: 892
Heterosphaeriaceae	... R. 198
Hydnaceae	 6: 429, 9: 208, 11: 106, 14: 201, 16: 174, 1116, 17: 147
Hymenogastraceae	 7: 154, 9: 280, 11: 168, 14: 267, 16: 245, 17: 239
Hypocreaceae	 2: 447, A. 194, 9: 941, 11: 354, 14: 621, 16: 559, 1140, 17: 777
Hypodermaceae	 10: 748, R. 28

- Hysteriaceae 2: 721, A. 263, 9: 1100, 11: 385, 14: 710, 16: 657, 1145, 17: 893, R. 3
 Laboulbeniaceae 8: 909, 9: 1130, 11: 446, 14: 725, 16: 674, 17: 915
 Leptostromaceae 3: 625, A. 356, 10: 412, 11: 553, 14: 992, 16: 986, 18: 419
 Lophiostomaceae 2: 672, A. 254, 9: 1074, 11: 382, 14: 702, 16: 650, 1144, 17: 886
 Lycoperdaceae 7: 48, 9: 266, 11: 157, 14: 257, 16: 230, 1117, 17: 217
 Melanconiaceae 3: 696, A. 361, 10: 446, 11: 562, 14: 1004, 16: 995, 1155, 18: 447
 Microthyriaceae 2: 658, A. 246, 9: 1053, 11: 379, 14: 686, 16: 633, 1141, 17: 861
 Mollisiaceae R. 503
 Monascaceae 14: 825
 Monoblepharidaceae .. 16: 394
 Mucedinaceae 4: 2, A. 372, 10: 510, 11: 586, 14: 1037, 16: 1022, 1156, 18: 495
 Mucoraceae 7: 182, 9: 335, 11: 239, 14: 432, 16: 383, 17: 494
 Myriangiaceae 16: 799, 1151, 18: 191
 Myxobacteriaceae 11: 467, 14: 842, 18: 217
 Nectrioidaceae 3: 613, A. 354, 10: 404, 11: 552, 14: 988, 16: 983, 18: 407
 Nidulariaceae 7: 28, 9: 265, 11: 156, 14: 256, 16: 229, 17: 214
 Onygenaceae 8: 861, 10: 80, 11: 440, 16: 807
 Ostropaceae R. 185
 Patellariaceae 8: 768, 10: 52, 11: 433, 14: 818, 16: 791, 1151, 18: 165, R. 277
 Perisporiaceae 1: 1, A. 1, 9: 364, 11: 252, 14: 462, 16: 398, 1123, 17: 524
 Peronosporaceae 7: 233, 9: 340, 11: 242, 14: 457, 16: 396, 17: 519
 Pezizaceae 8: 53, 10: 3, 11: 393, 14: 744, 16: 701, 1146, 18: 11, R. 913
 Phacidiaceae 8: 705, 10: 48, 11: 431, 14: 813, 16: 783, 1150, 18: 155, R. 60
 Phallaceae 7: 1, 9: 262, 11: 152, 14: 254, 16: 224, 1118, 17: 212
 Phymatosphaeriaceae.. 8: 843, 11: 440, 14: 826
 Polyporaceae 6: 1, 9: 150, 11: 79, 14: 164, 16: 138, 1115, 17: 95
 Protomycetaceae 7: 319, 9: 363, 11: 251, 14: 829, 16: 816, 1153, 18: 202
 Pseudophacidiaceae ... R. 87
 Rhiziniaceae R. 1134
 Saccharomycetaceae .. 8: 916, 11: 457, 14: 828, 16: 818, 1153, 18: 198
 Saprolegniaceae 7: 254, 9: 345, 11: 244, 14: 450, 16: 395, 1123, 17: 518
 Schizomycetaceae 8: 923, 10: 100
 Sphaeriaceae 1: 88, 2: 1, A. 22, 9: 442, 11: 271, 14: 478, 16: 417, 1128, 17: 561
 Sphaerioidaceae 3: 1, A. 285, 10: 100, 11: 472, 14: 844, 16: 825, 1154, 18: 219
 Sphaeropsidaceae 10: 785, 16: 1154
 Stictidaceae 8: 647, 10: 44, 11: 428, 14: 806, 16: 776, 1150, 18: 146, R. 112
 Stilbaceae 4: 563, A. 386, 10: 681, 11: 640, 14: 1107, 16: 1082, 1157, 18: 630
 Thelephoraceae 6: 513, 9: 218, 11: 115, 14: 212, 16: 181, 1116, 17: 160
 Tremellaceae 6: 760, 9: 257, 11: 142, 14: 244, 16: 215, 1117, 17: 203
 Tryblidiaceae R. 191
 Tuberaceae 8: 872, 10: 80, 11: 442, 14: 826, 16: 808, 1146, 18: 205
 Tuberculariaceae 4: 635, A. 389, 10: 700, 11: 645, 14: 1115, 16: 1090, 1158, 18: 658
 Tulasnellaceae 14: 234
 Uredinaceae 7: 528, 9: 291, 11: 174, 14: 269, 16: 257, 1118, 17: 244
 Ustilaginaceae 7: 449, 9: 282, 11: 230, 14: 410, 16: 367, 1122, 17: 472

List of New Genera and Types

Sphaeriaceae

Genus	Character	Type	Page
<i>Lasiosphaeris</i>	<i>Lasiosphaeria phaeophragmia</i>	<i>L. hispida</i> (Tode)	35
<i>Herpothrix</i>	<i>Herpotrichia phaeophragmia</i>	<i>H. calospora</i> (Winter)	35
<i>Comoclathris</i>	<i>Clathrospora pilosa</i>	<i>Comoclathris lanata</i> Clements	37

Verrucariaceae

<i>Dimerisma</i>	<i>Speconisca phaeodidyma</i>	<i>D. tenebrosum</i> (Norm.)	39
<i>Phaeomeris</i>	<i>Speconisca phaeophragmia</i>	<i>Ph. confusa</i> (Norm.)	39
<i>Pleophalis</i>	<i>Speconisca polyspora</i>	<i>P. nova</i> (Norm.)	39
<i>Phalostauris</i>	<i>Staurothele hyalospora</i>	<i>Ph. diffractella</i> (Tuck.)	39
<i>Phaeosporis</i>	<i>Verrucaria phaeospora</i>	<i>Ph. melasperma</i> (Nyl.)	39
<i>Lithoecis</i>	<i>Verrucaria immersa</i>	<i>L. tristis</i> (Kremp.)	39
<i>Phragmothele</i>	<i>Thelidium hyalophragmium</i>	<i>Ph. papularis</i> (Fr.)	39
<i>Phaeothrombis</i>	<i>Thrombium phaeosporum</i>	<i>Ph. melaspermiza</i> (Stnr.)	40
<i>Phaeoglaena</i>	<i>Microglaena phaeodictya</i>		40
<i>Dichoporis</i>	<i>Porina schizospora hyalodid.</i>	<i>D. schizospora</i> (Wain.)	40
<i>Diporina</i>	<i>Porina hyalodidyma</i>	<i>D. subsimplicans</i> (Nyl.)	40
<i>Dipyrenis</i>	<i>Pyrenula phaeodidyma</i>	<i>D. trachysperma</i> (Müll. Arg.)	40
<i>Holothelis</i>	<i>Thelopsis hyalospora</i>	<i>H. flaveola</i> (Arn.)	40
<i>Dithelopsis</i>	<i>Thelopsis hyalodidyma</i>	<i>D. subporinella</i> (Nyl.)	40
<i>Pyrenyllum</i>	<i>Arthropyrenia hyalodidyma</i>	<i>P. analeptum</i> (Ach.)	41
<i>Polythelis</i>	<i>Microthelia phaeophragmia</i>	<i>P. sexlocularis</i> (Müll. Arg.)	41
<i>Ditremis</i>	<i>Pleurotrema hyalodidymum</i>	<i>D. inspersa</i> (Müll. Arg.)	41
<i>Trichotrema</i>	<i>Pleurotrema scolecosporum</i>	<i>T. trichosporum</i> (Müll. Arg.)	41
<i>Phylloporis</i>	<i>Phylloporina hyalodidyma</i>	<i>Ph. phyllogena</i> (Müll. Arg.)	41

Hypocreaceae

<i>Sphaerodes</i>	<i>Sphaeroderma esubiculatum</i>	<i>S. episphaerium</i> (Ph. & Pl.)	44
<i>Rhynchomelas</i>	<i>Melanospora stromatica</i>	<i>Rh. arenaria</i> (Mont.)	44
<i>Dasyphthora</i>	<i>Nectria pilosa</i>	<i>D. lasioderma</i> (Ell.)	45

Dothideaceae

<i>Pleodothis</i>	<i>Plowrightia polyspora</i>	<i>P. polyspora</i> (Bref.)	49
<i>Discostroma</i>	<i>Curreya hyalodictya</i>	<i>D. rehmi</i> (Schnabl)	50

Mycoporaceae

<i>Chlorodothis</i>	<i>Mycoporellum hyalodidymum</i>	<i>Ch. lahmii</i> (Müll. Arg.)	50
<i>Sciodothis</i>	<i>Mycoporellum phaeodidymum</i>	<i>S. leucoplaca</i> (Müll. Arg.)	50
<i>Nothostroma</i>	<i>Mycoporellum hyalophragmium</i>	<i>N. roseolum</i> (Müll. Arg.)	50
<i>Mycoporis</i>	<i>Mycoporellum phaeophragmium</i>	<i>M. perexigua</i> (Müll. Arg.)	50

Hysteriaceae

<i>Pleoglonis</i>	<i>Glonium polysporum</i>	<i>P. strobiligena</i> (Desm.)	56
-------------------	---------------------------	--------------------------------	----

Graphidaceae

Plearthonis	Allarthonia hyalophragmia	P. caesia (Fw.)	58
Diarthonis	Arthonia hyalodidyma	D. lurida (Ach.)	58
Merarthonis	Arthoniopsis hyalodidyma	M. leptosperma (Müll. Arg.)	58
Digraphis	Graphis hyalodidyma	D. turbulenta (Nyl.)	59
Psorographis	Acanthothecis hyalophragmia	Ps. clavuliger (Wain.)	59

Stictidaceae

Habrostictis	Naevia iodata	H. pallida (Fckl.)	63
Naeviella	Naevia didymospora	N. paradoxa (Rehm)	63
Diplocryptis	Diplonaevia iodata	D. foveolaris (Rehm)	63
Xyloglyphis	Xylogramma didymosporum	X. striola (Fr.)	64
Merostictis	Phragmonaevia non-iodata	M. emergens (Karst.)	64

Tryblidiaceae

Tryblidis	Tryblidiopsis didymospora	T. pinastri (Pers.)	65
Odontura	Odontotrema scolecosporum	O. raphidospora (Rehm)	65

Bulgariaceae

Agyrina	Agyrium polysporum	A. sexdecimspora (Fckl.)	67
Myridium	Orbilbia polyspora	M. myriosporum (Ph. & Hark.)	67

Patellariaceae

Epilichen	Karschia lichenicola	E. scabrosus (Ach.)	69
Pleospilis	Melaspilea polyspora	P. vermifera (Leight.)	69
Lecoglyphis	Leciographa hysteroidea	L. centrifuga (Mass.)	70
Mycolecis	Leciographa saprophytica	M. leidecina (Rehm)	70
Parathalle	Lahmia parasitica	P. fuistingii (Körb.)	70

Caliciaceae

Eucyphelis	Sphinctrina laeta stipitata	E. acicularis (Smith)	71
Holocyphis	Cyphelium sphaerosporum	H. bolanderi (Tuck.)	71
Dipyrgis	Pyrgillus didymosporus		71
Ditylis	Tylophorum didymosporum	D. moderata (Nyl.)	71

Chrysotrichaceae

Holocoenis	Coenogonium amerosporum	H. lepieurii (Mont.)	72
------------	-------------------------	----------------------	----

Collemataceae

Pleopyrenis	Pyrenopsis polyspora	P. picina (Nyl.)	72
Pleoconis	Peccania polyspora	P. kansana (Tuck.)	73
Dicollema	Collema didymosporum	D. pycnocarpum (Nyl.)	74

Peltophoraceae

Gonothecis	Sporopodium phycepitheciale	G. phyllocharis (Mont.)	75
Chloropeltis	Peltophora palmellicola	Ch. apthosa (L.)	75
Scoleactis	Lecanactis scolecospora	S. myriadea (Fee)	76
Pleolecis	Lecidea polyspora	P. geophana (Nyl.)	76
Diphloeis	Toninia didymospora	D. candida (Web.)	76
Diphanis	Rhizocarpum hyalodidymum	D. polycarpa (Hepp)	77
Diphaeis	Rhizocarpum phaeodidymum	D. badiatra (Flk.)	77
Phalodictyum	Rhizocarpum hyalodictyum	Ph. obscuratum (Ach.)	77
Merophora	Gyrophora merospora	M. haplocarpa (Nyl.)	77

Cladoniaceae

Dibaeis	Baeomyces didymosporus	D. rosea (Pers.)	78
Cyanobaeis	Baeomyces cyanophyceus	C. paeminosa (Kremp.)	78
Chlorocaulum	Stereocaulum lecanorinum	Ch. salazinum (Bory)	78

Parmeliaceae

Myriolecis	Lecanora polyspora	M. sambuci (Pers.)	79
Adermatis	Lecania phragmospora	A. nylanderiana (Mass.)	79
Dyslecanis	Lecania polyspora	D. syringea (Ach.)	79
Pleochroma	Candelariella polyspora	P. vitellina (Ehrh.)	80
Ocellis	Ocellularia didymospora		80
Phanotylum	Tremotylum hyalodictyum	P. australiense (Müll. Arg.)	80
Diphanosticta	Sticta lecanorina hyalodid.	D. cellulifera (H. & T.)	81
Diphaeosticta	Sticta lecanorina phaeodid.	D. physciospora (Nyl.)	81
Phanosticta	Sticta lecan. hyalophragmia	Ph. freycinetii (Del.)	81
Dysticta	Sticta lecideina	D. sinuosa (Pers.)	81
Cystolobis	Lobaria cysticoccola	C. leucocarpa (Müll. Arg.)	81
Podostictina	Stictina hyalophr. stipitata	P. endochrysoides (Müll. Arg.)	82
Merostictina	Stictina phaeophragmia	M. mougeotiana (Del.)	82
Dystictina	Stictina lecideina	D. tomentosa (Sw.)	82
Phycodiscis	Lobarina lecanorina	Ph. retigera (Bory)	83

Physciaceae

Pleorinis	Rinodina polyspora	P. polyspora (Th. Fr.)	84
Meroplacis	Caloplaca phragmospora	M. brebissonii (Fee)	84
Merorinis	Rinodina phragmospora	M. conradi (Körb.)	84
Phragmopyxine	Pyxine phragmospora	Ph. eschweileri (Tuck.)	84
Dictyorinis	Rinodina dictyospora	D. diplinthia (Nyl.)	84

Mollisiaceae

Spilopezis	Pyrenopeziza subiculata	S. radians (Rob.)	85
Dibelonis	Beloniella hyalodidyma	D. dehnii (Rabh.)	86

Helotiaceae

Pezoloma	Cyathicula sessilis	P. griseum Clements	86
Eubelonis	Belonium hyalodidymum	E. drosodes (Rehm)	87
Belospora	Belonioscypha chaetospora	B. ciliatospora (Fckl.)	87
Merodontis	Davincia sessilis	M. tenella (Penz. & Sacc.)	87
Dyslachnum	Lachnum sessile	D. mollissimum (Lasch)	87
Phalothrix	Dasyscypha phalotrichia	Ph. hyalotricha (Rehm)	88
Dasypezis	Dasyscypha sessilis	D. albolutea (Pers.)	88

Pezizaceae

Iotideia	Otideia iodata	I. pleurota (Phill.)	89
Podaleuris	Aleurina stipitata	P. reperta (Boud.)	89
Leucopezis	Neottiopezis eciliata	L. excipulata Clements	90
Trichaleuris	Aleurina setosa	T. crinita (Bull.)	90

Helvellaceae

Haplocybe	Cudoniella hyalospora	H. aquatica (Lib.)	91
-----------	-----------------------	--------------------	----

Gymnascaceae

Podocapsium	Podocapsa polyspora	P. diffusum (Van Tieg.)	94
-------------	---------------------	-------------------------	----

Phomataceae

Sirodothis	Dothiorella catenata	S. populi Clements	123
Chaetoconis	Kellermannia phaeodidyma	Ch. polygoni (E. & E.)	125

Zythiaceae

Sirocyphis	Cyphina catenata	S. nivea Clements	130
------------	------------------	-------------------	-----

Leptostromataceae

Petasodes	Leptostromella basidiis um- bellatis	P. umbellatum (Vesterg.)	133
-----------	---	--------------------------	-----

Melanconiaceae

Hormyllum	Trullula hyalospora basidiis ramosis	H. populi (Preuss)	135
Thecostroma	Trullula hyalospora	Th. nitidulum (Sacc.)	135

Moniliaceae

Trichoconis	Ramularia trichospora	T. caudata (Ap. & Str.)	145
-------------	-----------------------	-------------------------	-----

Tuberculariaceae

Hormodochis	Epidochium catenatum	H. melanochlora (Desm.)	163
Chaetodochis	Chaetostroma catenatum	Ch. caricis (Fckl.)	163

Glossary of Latin and English Terms

A

a, without (in comp.)

ab, from

abbreviatus, shortened

abeuns, deviating

abhorreo, abhor, differ from

abiegnus, fir

abietinus, fir

abnormis, abnormal

abortivus, abortive

abortus, aborted

abrupte, abruptly

abundans, abundant

abunde, abundantly

ac, and

acaudatus, without a tail

accedo, to approach

accessory, additional

accipio, to accept

acerinus, maple

acervulatus, heaped, massed

acervulus, **i**, **m.**, a little heap

acervus, **i**, **m.**, a heap

achromaticus, without color

achrous, colorless

acicularis, acicular, needle-shaped

acidulus, slightly acid

acies, **ei**, **f.**, edge

acotyledon, **nis**, **m.**, cryptogam

acquirō, to acquire

acrogenus, acrogenous, borne at tip

acropleurogenus, borne at the tip and
on the sides

acris, sharp

aculeatus, spiny, pointed

aculeolatus, spiny, pointed

acuminatus, long-pointed

acus, **us**, **f.**, needle

acutatus, acute

acutiusculus, somewhat acute

acutus, acute

ad, to

adesse, to be present

adhibitus, used, applied

adhuc, as yet, hitherto

adinterim, meanwhile

admiro, to look, wonder at

admodum, at least, fully, very

adnatus, adnate, touching broadly

adparenter, apparently

adproximatus, drawn near

adscendens, ascending

adsociatus, clustered

adspectus, **us**, **m.**, sight, appearance

adultus, fully grown

adustus, burned, blackened

aecidiiformis, aecidium-shaped

aecium, a cluster cup

aegre, poorly, with difficulty

aemulans, rivalling

aemulus, similar

aeneus, brazen, coppery

aequalis, equal

aequans, equalling

aequidistans, equally distant

aerius, aerial

aerobius, growing in the air

aerophilus, aerial

aeruginosus, copper-colored

aeternus, eternal

affectus, affected

affixus, attached

afflatus, swollen

agamicus, asexual

agamus, asexual

ager, **ri**, **m.**, field

agglomeratus, heaped together

aggregatus, grouped together

albicans, whitening

albidus, white

albofarctus, white-stuffed

albolutescens, whitish yellow

albus, white

alcoholicus, alcoholic

alienus, foreign, strange

aliquantisper, for a while

- aliquantulus**, somewhat, a little
alius, another, other
alius,—alius, some—others
allantoid, sausage-shaped, short and curved
allantoideus, a, um, allantoid, sausage-shaped
alliaceus, a, um, of an onion
alpis, mountain
alte, deeply
alternus, a, um, alternate
altitudo, f., height
altus, a, um, high
alutaceus, grayish yellow
alveolatus, a, um, with hollows
amaricans, making bitter, irritating
ambiens, surrounding
ambitus, m., periphery
amentum, n., catkin
amerosporus, a, um, with one-celled spores
amethysteus, a, um, amethyst-colored
amissus, a, um, lost, dismissed
ammoniacalis, e, like ammonia
amnis, is, m., a brook
amoebiformis, e, amoeba-form
amoeboid, amoeba-like
amoeboideus, a, um, amoeba-like
amoene, beautifully
amoenus, a, um, beautiful, pleasant
amoveo, to withdraw
amphibius, a, um, amphibial
amphigenus, a, um, borne on both sides
amplectens, clasping
amplecto, to wind or clasp
amplus, a, um, broad, ample
ampulliformis, ampulliform, cushion-like
amycelicus, without mycelium
amygdalinus, almond-like, pink
analogus, similar
anastomosans, anastomosing, running together
anceps, cipitis, two-headed, double
androgynus, with male and female
angularis, angular
angulosus, angulose, angular
angustatus, narrowed
angustus, narrow
animalcula, ae, f., little animal
annularis, ring-like
annulatum, in a ring
annulatus, annulate, with a ring, ringed
annuliform, ring-like
annulus, i, m., a ring
annuosus, aged, old
anormaliter, abnormally
anserinus, of or pertaining to geese
ante, before
antecedens, preceding
antheridiiformis, antheridium-like
antheridium, ii, m., antherid
antherozoidium, ii, n., antherozoid
antice, in front
aparaphysatus, without paraphyses
apertus, open
aperio, to open, uncover
apex, icis, m., tip
apiculatus, apiculate, with a point
apiculiformis, like a little point
apophysatus, with a supporting cell
apothecium, ii, n., cup or disk containing asci
appendicula, ae, f., little appendage
appendiculatus, appendiculate, appendaged
appendix, icis, f., appendage
applanatus, applanate, flattened
approximatus, close, near
apricus, wild
apud, at
apus, odis, without a stalk
aquaeductus, us, m., aqueduct
aquaticus, aquatic
aquosus, watery
arachnoideus, cobwebby
araneosus, cobwebby
arbor, is, f., tree
arbusculiformis, shrub-like
arcte, closely
arcticus, arctic
arcuatim, bow-like, curved
arcuatus, arcuate, bow-like
area, ae, f., space
areola, ae, f., little space
areolatus, areolate, marked by areas or spaces
arescens, drying
aresco, to become dry

argenteus, silvery
argentinus, silvery
argillaceus, clay-color
aridus, dry
arista, *ae, f.*, awn
aristatus, aristate, awned
arrectus, upright, stiff
arrhizus, without roots
articulatus, jointed
articulus, *i, m.*, joint
asciger, ascus-bearing
ascogenic, producing asci
ascoma, *atis, n.*, spore-fruit, ascus-bearing body
ascophorus, ascus-bearing
ascus, *i, m.*, sac
asiaticus, Asiatic
asper, rough
asperatus, asperate, roughened
aspergo, to scatter, sprinkle
asperulus, slightly roughened
asser, *eris, m.*, branch, beam, post
assurgens, ascending
asterigmaticus, without stalks
asterineus, star-like, radiate
asteroid, star-like, radiate
asteroma-like, with radiate subicle
astomus, mouthless
astromatoideus, without a stroma
asymmetricus, irregular
ater, dark, black
atomatus, with small particles
atomisticus, tiny
atque, also
atrans, blackening
atratu, dark
atro-fuscus, dark
atro-inquinans, blackening
atro-nitidu, black and shining
atropiceus, black as pitch
atropurpureus, dark purple
attenuatus, tapering
attingens, touching
attolens, raising
atypicus, abnormal
auctio, *onis, f.*, growth
auctor, *is, comm.*, author
auctus, enlarged
audeo, to dare
augmentum, *i, n.*, increase, growth

aurantiaceus, orange, golden
aurantinus, orange
auratus, golden
aureus, golden
auriformis, ear-shaped
australis, southern
aut, or
autem, moreover
authenticus, authentic
autumnus, autumn
avulsus, torn off, separated
axicola, growing on the axis
axiformis, axis-like
axilaris, axillary
azonus, without zones
azygospore, a zygospore formed without conjugation

B

bacca, *ae, f.*, berry
baccatus, berry-like
bacillaris, bacillar, rod-shaped
bacteriformis, bacterium-like
bactrosporus, with rod-shaped spores
baculum, *i, n.*, rod
badius, brown
basidiosporus, with spores borne on stalks
basidium, *ii, n.*, rod, basidium
basilaris, basal
basis, *is, f.*, base
bene, plainly, well
benevole, kindly
betulicola, growing on birch
betulinus, birchen
bi-, two, twice
bibulus, absorbing
biclavuligerus, bearing two club-shaped branches
biconic, conic at each end
biconvexus, biconvex
bicornus, with two horns, two-branched
bicorticus, with two barks
bidentatus, two-toothed
bifidu, split into two parts
biformis, or **-us**, of two forms
bifrons, on both sides of the leaf
bifurcatus, two-forked

- biguttulatus**, with two globules or vacuoles
bilabellulatus, two-lipped
bilabiatus, two-lipped
bilobus, two-lobed
bilocularis, two-celled
binatim, by twos
binucleolatus, with two oil-drops
binus, two-fold
biogenus, biogenous, growing on organisms
biophilus, biophilous, growing on organisms
bipunctatus, with two vacuoles
bis, twice
biscocitiformis, biscuit-shaped
biserialis, in two rows
biseriatus, in two rows
bisporus, two-spored
bitunicatus, with two walls
biuncinatus, two-hooked
bombardus, cannon-like
borealis, northern
botryosus, botryose, clustered like grapes
botuliformis, botuliform, sausage-shaped
brachiatus, with arms
bractea, *ae, f.*, bract
brevicollis, short-necked
brevis, short
breviter, shortly
breviusculus, somewhat short
brunneolus, brownish
brunneus, brown
bullula, *ae, f.*, bubble
bullula, *ae, f.*, a little swelling
byssinus, cottony
byssisedus, byssisede, seated on cotton
byssoides, byssoid, cottony
byssus, *i, f.*, cotton
- C**
- caerulescens**, turning blue
caesius, bluish-grey
caespes, *itis, m.*, tuft
caespitosus, caespitose, in tufts
caesus, fallen
calamus, *i, m.*, stem
calcareus, of lime, calcareous
calcariferus, bearing lime
calcifer, bearing lime
calidarium, *ii, n.*, hot-house
callosus, roughened
calvescens, becoming bare
calvitium, *ii, n.*, bald spot
calvus, bare, bald, not pubescent
calx, *calcis, f.*, lime
calyciformis, cup-shaped
calycicola, living on the calyx
calycularis, cup-shaped
calyptra, *ae, f.*, cap
calyx, *ycis, m.*, calyx, cup
campanulatus, bell-shaped
campaniformis, bell-shaped
campylotropus, curved
canaliculatus, canaliculate, channeled
candicans, growing white
cannabinus, of hemp
canus, hoary
capillaris, hair-like
capillatura, *ae, f.*, mass of hair
capilliform, hair-like
capillitium, *ii, n.*, mass of threads
capillus, *i, m.*, hair
capitatus, capitate, in heads
capitulatus, borne in little heads
capitulum, *i, n.*, a little head
capreolus, *i, m.*, goat
caprinus, of or pertaining to goats
capsula, *ae, f.*, capsule
caput, *itis, n.*, head
carbo, *onis, m.*, carbon, charcoal
carbonaceus, like coal
carbonicola, on burned-over ground or on charcoal
carbonous, like coal or carbon
carens, lacking
caries, *ei, f.*, decay
carinatus, keeled
cariosus, decaying
carneus, flesh-colored
carnosus, carnose, fleshy
caro, *carnis, f.*, flesh
carpogenus, living on fruit
carpogonium, *ii, n.*, carpogone
cartilagineus, cartilaginous, tough but pliable
caryopsis, *idis, f.*, grain

castaneus, chestnut brown
catenate, in chains
catenifer, chain-bearing
catenigerus, bearing chains
catenula, *ae, f.*, chain
catenulatus, catenulate, in chains
catenuliformis, chain-like
catenulus, *m., -a, f.*, a small chain
caterva, *ae, f.*, heap, crowd
catervatim, in heaps, in groups
cauda, *ae, f.*, tail
caudatus, caudate, tailed
caudex, *icis, m.*, stalk
caudicula, *ae, f.*, a little stalk
caulicola, growing on stems
caulis, *is, m.*, stem
caulogenus, on stems
caverna, *ae, f.*, a cavern, hollow
cavernosus, with hollows
cavernula, *ae, f.*, a little cavity
cavitas, *atis, f.*, cavity
cavitus, hollow
cavus, *i, m.*, hollow
celans, hiding
cella, *ae, f.*, a cell
celluliformis, cell-shaped
cellulosus, cellular
censeo, to think, estimate
centrifugus, centrifugal
centrum, *i, n.*, the centre
cephalodium ii, n., a globose to club-shaped projection on a lichen thallus
ceraceus, waxy
cerebriformis, brain-like
cereus, waxy
cerno, to perceive, separate
cernuus, nodding, inclined
cerumen, *inis, n.*, wax
cervinus, tawny
cespitose, clustered, crowded
ceterum, remaining
chalybeus, of steel
character, *eris, m.*, character, style
charta, *ae, f.*, paper
chartaceus, papery
chlamydosporicus, with chlamydo-spores
chlorinus, greenish
chlorophyllous, green, with chlorophyll

chorda, *ae, f.*, twine, a cord
cibaria, *ae, f.*, food
cicatrix, *icis, f.*, a scar
ciliatulus, slightly ciliate
ciliatus, ciliate, with long hairs on the margin
ciliolatus, ciliolate, with cilia
cincinnatus, curled
cinctus, surrounded
cinerascens, becoming ashen
cinereus, ashen
cingens, surrounding
cingulatus, surrounded
cingulus, *i, m.*, a little belt
cinnabarinus, orange red
cinnamomeus, cinnamon-colored
circa, near
circinatus, circinate, coiled
circino, to circle
circiter, about
circuitus, *us, m.*, a circuit
circulus, *i, m.*, a circle
circumambiens, encircling
circumdatus, surrounded
circumscissile, splitting circularly
circumscrip-tus, circumscribed
circumtextus, surrounded
circumvallatus, surrounded
cirr-hatus, curled
cirrhosus, curly
citatus, cited
cito, to name, mention
cito, soon, rather
citriformis, citriform, lemon-shaped
citrinus, lemon yellow
cladodium ii, n., a leaf, branch
cladogenus, borne on branches
clathratus, clathrate, latticed
clausus, closed
clava, *ae, f.*, a club
clavaria-like, club-shaped, or coral-like
clavatus, club-shaped
claviformis, club-shaped
clavis, *is, f.*, a key
clavula, *ae, f.*, a little club
clavulatus, club-shaped
clypeatus, shield-like
clypeus, *i, m.*, a shield

- coacervatus**, coacervate, heaped together
coadunatio, onis, f., a summing up
coadunatus, united, collected
coalescens, coalesced, running together
coalitus, joined, running together
coarctatus, crowded
coccineus, bright red
coccus, i, m., round cell, berry
cochleariformis, spoon-shaped
cochleatus, ear-like
coctus, cooked
coenobium, ii, n., a colony
coerulescens, turning blue
coffeatus, coffee-like
coffeicolor, coffee-colored
coffeiformis, coffee-shaped
cognatus, related
cogo, to act, collect
cohabitans, living together
cohaerens, cohering
collabasco, to fall in
collabens, collapsing, crumbling up
collabent, collapsing, falling in
collapsus, collapsed
collariatus, collared, attached to a collar
collectivus, collected
colliculosus, with tiny elevations
collum, i, n., a neck
colonia, ae, f., a colony
color, is, m., color
coloratio, onis, f., coloration, color
coloratus, colored
coloreus, colored
columnella, ae, f., a small pillar, columnella
columnaris, columnar
comatus, shaggy
comestibilis, eatable
commissura, ae, f., commissure, path, cleft
commixtus, commingled
communico, to share, communicate
communis, common
comosus, hairy
compactus, dense
compaginatus, united
complectens, comprising, clasping
complecto(r), to clasp
complexus, complex
compositus, composed, compound
compressus, compressed
concatenatus, in chains
concavus, concave
concentricus, concentric
conceptaculum, i, n., conceptacle
conchiformis, conchiform, shell-shaped
concolor, concolorous, of like color
concrescens, growing together
concretus, united
condensus, condensed
conditio, onis, f., condition
confero, to collect
confertus, crowded
confirmatio, onis, f., confirmation
conflatus, swollen
confluens, running together
confluo, to merge
conformis, all alike, similar
confundo, to mingle, confuse
congestus, crowded
conglobatus, conglobate, heaped together
conglomeratus, heaped
conglutinated, conglutinate, glued together
congregatus, aggregated
congruo, to agree
conicus, conical
conidium, ii, n., an asexual spore
conidial, producing or pertaining to conidia
conidicus, conidial
conidiferus, conidia-bearing
condiophorum, i, n., a hypha bearing conidia, a condiophore
conjugatio, onis, f., conjugation
connatus, connate, joined
connexus, connected
connivens, connivent, approaching
conoideus, conoid, cone-shaped
consortium, ii, n., company
conspersgens, sprinkled
conspersus, scattered
conspicuus, conspicuous
conspurcatus, polluted
constipatio, onis, f., a crowding

- constituens**, constituting
consuetudo, inis, f., a habit
consumptus, destroyed
contemno, to condemn, disparage
contextum, i, n., texture, context
contiguus, close
continens, containing
continuus, continuous, one-celled
contortus, twisted
contra, against
contractus, narrowed
contusus, bruised
conus, i, m., a cone
convergens, coming together
convolutus, convolute, coiled
convolutio, onis, f., a fold
copiosus, abundant
coprophilus, growing on dung
copulans, copulating
coralloid, coral-like
coralloideus, coralloid, like much-branched coral
coriaceus, leathery
corneus, corneous, horn-like
corniculatus, corniculate, horned
corniformis, horn-shaped
cornutus, horned
coronatus, crowned
corpusculum, i, n., a little body
corrugatus, corrugate, ridged
corruptus, corrupted, spoiled
cortex, icis, m., the bark
corticalis, cortical, of bark
corticatus, corticate, with a bark or epiderm
corticola, corticole, growing on bark
cortina, ae, f., veil
cortinate, with a curtain-like veil
corvinus, pertaining to the raven, black
costa, ae, f., ridge
cestatus, costate, ridged
crassities, ei, f., thickness
crassitudo, inis, f., thickness, width
crassiusculus, somewhat broad
crassus, broad
crateriformis, crateriform, crater-shaped
creber, crowded
cremicolor, cream-colored
cribrosus, sieve-like
crinitus, hairy, crested
crispulus, somewhat crisp
crispus, crisp
crista, ae, f., crest
cristatus, crested
crocatius, yellow
croceus, yellow
cruciatim, cruciately, cross-like
cruentatus, bloody
crusta, ae, f., crust
crustaceus, crust-like
crustiformis, crust-shaped
crustose, forming a crust, more or less interrupted
crustula, ae, f., a little crust
cubile, is, n., a bed
cuboideus, cuboid, cubical
cucullatus, hooded
cucumeriformis, cucumber-shaped
culmicola, growing on grass-stems
culmus, i, m., culm, a stalk, stem
cultellus, i, m., a small knife
culter, tri, m., a knife
cultriformis, knife-like
cultus, cultivated
cum, with
cumulatus, heaped up
cuneatus, wedge-shaped
cuneiformis, wedge-shaped
cuniculus, i, m., a rabbit
cupreus, coppery
cuprinus, coppery
cupula, ae, f., a little cup
cupularis, **cupulatus**, **cupuliformis**, cup-shaped
curtus, short
curvatus, curved
cuspidatus, cuspidate, with a tooth
cuticula, ae, f., cuticle
cuticularized, with firm cover or cuticle
cutis, is, f., the skin
cyaneus, blue
cyathiformis, cup-like
cyclus, i, m., a cycle
cylindraceus, **cylindricus**, cylindrical
cymbiformis, boat-shaped
cyphella, ae, f., an opening or hollow

in a thallus, more or less cup-shaped
cystidium, ii, n., cyst
cystophore, the stalk which bears a cell or cyst

D

daedaleus, labyrinthine
dealbatus, whitened
debilis, weak
deciduus, falling
decies, ten times
decorticatus, without bark
decumbens, prostrate
decurrens, decurrent, running down the stem
defectus, lacking
deficiens, lacking
deficio, to lack
definitus, definite
deflexus, deflexed
deformus, deformed
degenero, to degenerate
dehiscens, dehiscent, splitting
dein, then, at length
dejectus, fallen
dejiciens, throwing down
delicatulus, delicate
delineatus, figured
deliquescens, deliquescing, liquefying
delitescens, hiding
delitescio, to conceal, lurk
deltoideus, delta-like, triangular
dematium-like, black and cobwebby
dematius, black and cottony
demonstro, to show
demum, at length
dendritice, dendritically, tree-like
dendriticus, tree-like
dendroideus, dendroid, tree-like
denigratus, blackened
denique, at length
densus, close, dense
dentatus, toothed
denticulatus, denticulate, with little teeth
denudans, denuding
denudatus, denuded
deorsum, downward
dependens, hanging
deplanatus, flattened
depressus, depressed
derumpens, breaking
descendens, descending
desciscens, leaving, deserting
describo, to describe
descriptus, described
desicco, to dry up
desinens, ending, closing
desum, to fail, be absent
destitutus, lacking
destruens, destroying
detergibilis, removable, breakable
deustus, burnt
diametralis, of the diameter
diametrum, i, n., diameter
diaphanus, diaphanous, transparent
diatrype-like, with a stroma different from the tissue of the matrix
dichotomus, dichotomous, two-forked
diclinus, with separate sexes
dictyosporus, spores having cross and longitudinal walls
didymosporus, with two-celled spores
didymus, two-fold or two-celled
differo, to differ
difficilis, difficult
diffluens, diffluent, dissolving
diffractus, broken
difformis, of two forms
digestus, broken up
digitiformis, finger-shaped
digitaliformis, digitate, finger-like
digitatus, digitate, having fingers
dignosco, to differ
dignotus, to distinguish
dilabens, breaking apart
dilatatus, spread out
dilute, dilutely
dilutus, dilute
dimidiatus, dimidiate, two-lobed, halved
dimidius, half
dimorphus, of two forms
dioecious, sex organs on separate plants
directio, onis, f., direction
directus, straight
dirumpens, breaking apart
disciformis, disc-shaped
discolorus, discolorous, discolored

discretus, discrete, separate
discrimen, inis, n., difference
disculus, i, m., little disc
disfractus, broken
disparens, disappearing
dispergens, scattering
dispositus, arranged
disruptus, broken
disseco, to cut up
dissectus, cut up
disseminatus, scattered
dissentio, to disagree
dissepimentum, i, n., partition, wall
distal, distant, further
distans, remote
distichus, distichous, in two rows
distinguo, to distinguish
diu, long
divaricatus, spreading
divergens, diverging
diversimodus, in different ways
diversus, diverse, different
divinans, conjecturing
divisio, onis, f., a division
divisus, divided
doliiformis, doliiform, cask-shaped, jar-shaped
dolium, ii, n., cask, jar
donacinus, of a reed
donatus, furnished
dorsiventral, with two unlike sides
dorsum, i, n., back
dothideaceus, like *Dothidea*, i. e., loculate
dubitantur, doubtfully
dubius, doubtful
duco, to lead
ductus, led
dulcis, sweet
dumetum, i, n., a thicket
duo, two
duodecim, twelve
duplo, twice
duriusculus, somewhat hard
durities, ei, f., hardness
durus, hard

E

eburneus, ivory-white
ecaudatus, without a tail
eccentricus, eccentric, lateral

echinatus, spiny
echinulatus, echinulate, spiny
edulis, edible
effiguratus, shaped, formed
effoetus, worn out
efformatus, formed
effusus, effuse, spread out
egrediens, growing out
elasticus, elastic
elatus, tall
elevatus, raised
ellipticus, elliptical
ellipsoideus, ellipsoid
elongatus, lengthened
emarginatus, without a margin
emergens, emerging
emergeo, to emerge
emersus, emerging
emittens, emitting
emortuus, dead
enatus, arising from
endobasidial, continuous with the basidium
endobiotic, growing within living things
endochroma, atis, n., colored contents
endogenus, endogenous, born within
endoperidium, ii, n., inner peridium
endophytic, growing in plants
endoplasma, atis, n., protoplasm
endoxylus, within wood
endozoic, growing in animals
enim, for
endoparasiticus, internally parasitic
entomogenus, entomogenous, living in insects
epelliculosus, without a covering or pellicle
epidermis, idis, f., epiderm, the surface skin
epigaeus, epigaeal, on the ground
epigenus, borne above
epiphloeodus, on the bark
epiphragma, an upper wall or division
epiphyllus, on the upper side of the leaf
epiphytic, upon plants
epispodium, ii, n., outer wall of spore
epithecium, a layer above the asci, usually formed of the tips of the paraphyses

- epizoic, growing on animals
 equinus, equine, belonging to horses
 erectus, erect
 ergo, therefore
 erostratus, without a beak
 erostris, without a beak
 erraticus, erratic, wandering
 error, is, m., error
 eructatus, thrown up
 erumpens, bursting out
 erysiphoides, like Erysiphe, cobwebby
 esepate, without cross walls
 estriatus, without lines or markings
 etiam, also
 etsi, although
 eumorphus, well-formed
 eutype-like, eutypeous, eutypoid, with an effuse stroma similar to the tissue of the matrix
 evacuans, emptying
 evacuatus, emptied
 evado, to escape
 evaginatus, without a sheath
 evanescens, evanescent, disappearing
 evanidus, vanishing
 evidentius, more clearly
 evolutus, developed
 evolvatus, without a volva
 evolvens, developing
 exacte, exactly
 exalbescens, becoming white
 exalbidus, whitish
 exalbugo, to whiten
 exannulatus, without a ring
 exappendiculatus, not appendaged
 exaridus, dried out
 exasperans, roughened
 exasperatus, roughened
 exaspero, to roughen
 excavatio, onis, f., an excavation, hollowing out
 excavatus, hollowed out
 excedens, exceeding
 excentric, out of the centre, lateral
 exciple, the outer wall or covering of an apothecium
 excipuliformis, cup-shaped
 excipulum, i, n., exciple, margin
 excrescens, growing out
 excutiens, shaking out
 exemplaris, model
 exemplarium, ii, n., specimen, sample
 exemplum, i, n., an example
 exesus, consumed, destroyed
 exhibens, exhibiting
 exigens, scanty
 exiguitas, atis, f., smallness, scantiness
 exiguus, little, small
 exilis, thin, slender
 eximie, exceedingly
 existimo, to estimate
 exitus, us, m., a departure, escape
 exobasidial, separated by a wall from the basidium
 exogenus, arising on the outside
 exoperidium, ii, n., outer peridium
 exoriens, arising
 exosporium, ii, n., exospore, outer wall of the spore
 expallens, becoming pale
 explodens, exploding
 expulsus, expelled
 exquisite, beautifully
 exsertus, exerted, thrust out
 exsiccatio, onis, f., a drying out
 exsiccatus, dried out
 exsiliens, escaping
 exsuccus, without milk or juice
 extensio, onis, f., extension
 externus, external
 extimus, outermost, ultimate
 extra, without, outside
 extrico, to extricate
 extrorsum, toward the edge
 extus, outside

F

- fabiformis, bean-shaped
 fabrica, ae, f., texture
 facies, ei, f., face, form
 facilis, easily
 fagineus, beechen
 falcatus, falcate, scythe-shaped, curved
 falciformis, beak-shaped, scythe-shaped
 familia, ae, f., family
 familiola, ae, f., a little family
 farctus, stuffed

farina, ae, f., meal, flour
farinaceus, mealy
fascia, ae, f., fascicle
fasciatus, grouped
fasciculatus, fasciculate, fascicled, in bundles
fastigiatus, bunched
faticens, disappearing, breaking up
favosus, hollow
femineus, feminine
fenestratus, with windows or openings
fere, almost
fermentatio, onis, f., fermentation
fermentum, i, n., yeast
ferruginascens, turning rust-colored
ferrugineus, rust-colored
ferrumequinum, i, n., a horse-shoe
ferrum, i, n., iron
fibra, ae, f., a fiber, filament
fibrilla, ae, f., little fibril
fibrillula, ae, f., a little fibril
fibrosus, fibrous
fictitius, fictitious
filamentosus, filamentous, thread-like
filia, ae, f., daughter
filiformis, filiform, thread-shaped
filiger, filament-bearing
filum, i, n., thread
fimbria, ae, f., fringe
fimbrians, fringing
fimbriatus, slightly fringed
fimbriatus, fimbriate, fringed
fimicola, fimicole, dwelling on dung
finus, i, m., dung
findo, to cleave, divide
firmulus, somewhat firm
fissilis, cleft, ruptured
fissuratus, fissured, split
fissus, split
fistulosus, hollow
flabelliformis, fan-shaped
flaccidus, weak
flagella, ae, f., lash
flagellatus, bearing long bristles or threads
flagelliformis, lash-like
flamens, flame-colored
flavens, yellowing
flavidus, yellowish

flavus, yellow
flexuosus, flexuous, full of turns or windings
flexus, bent
flocciformis, tuft-like
floccosus, floccose, cottony
floccus, i, m., tuft
floralis, floral
flumen, inis, n., river
fluvius, ii, m., a river
fluxilis, flowing
foedatus, dark, soiled
foetidus, with a bad odor
foliicola, foliicole, living on leaves
foliose, like a leaf in form
folium, ii, n., leaf
foramen, inis, n., a hole
forma, ae, f., form
formans, forming
formo, to form
formosus, beautiful
fornix, icis, m., a vault
forsan, perhaps
forsitan, perhaps
fortasse, perhaps
forte, strongly
fovens, nourishing
fraccidus, soft, mellow
fractus, broken
fragilis, fragile
fragmentum, i, n., a bit, fragment
frequens, frequent
friabilis, falling to pieces
frigidarium, ii, n., a cold place, cold storage
frondosus, leafy
frons, dis, f., a leaf
fruticicola, living on fruits
fructiferus, fructifer, fruit-bearing
fructificans, fruiting
fructificatio, nis, f., fruiting
fructus, us, m., fruit
frustulatus, fragmentary
frustum, i, n., a bit, piece
fruticosus, fruticose, shrub-like
fruticulosus, fruticulose, shrub-like
fucatus, colored
fugans, fleeting
fulciens, supporting
fuliginus, fuliginous, sooty

fuligo, inis, f., soot
fultus, supported
fulvellus, somewhat tawny
fulvescent, becoming tawny
fumagineus, fumaginous, smoky.
fumosus, smoky
fungicola, fungicole, growing on fungi
fungillus, i, m., a little fungus
fungus, i, m., a fungus
funicularis, rope-like
funiculus, i, m., a little rope
uniformis, rope-like
furcatus, furcate, forked
furfur, uris, m., bran
furfuraceus, bran-like
furfurellus, covered with bran
fuscatus, darkened
fuscillus, somewhat dark
fuscescens, darkening
fuscidus, dark
fuscidulus, dark
fuscus, dark, or dark brown
fusiformis, fusiform, spindle-shaped
fusisporus, with spindle-shaped spores
fusoideus, fusoid, spindle-shaped

G

galeiformis, hood-shaped
galeriformis, cap-shaped
gamete, sex-cell
gangliformis, forming knots
gangligerus, bearing knots
gelatina, ae, f., gelatine
geminatus, paired, twinned
gemmiparus, producing buds
generans, generating
genesis, is, f., origin
geniculatus, bent
genuflexus, bent
genuinus, genuine
genus, eris, n., genus
gerens, bearing
germinans, germinating
germinatio, onis, f., germination
gibbosus, swollen
gigastylosporus, with very large stylospores
gignens, producing
gigno, to bear
gilvus, brownish

glaber, smooth
glabrescens, becoming smooth
glacies, ei, f., glacier, ice
glans, glandis, f., a nut,
glaucescens, turning bluish-green
glaucus, sea-green
gleba, ae, f., soil, mass
globosus, globose, rounded
globuliger, bearing a ball
globulus, i, m., a globule
glomerula, ae, f., a little mass
glomerulatum, in heaps
gluten, inis, n., glue
glutinosus, glutinous
gonidium, ii, n., an algal cell
gossypinus, cottony
gracilis, graceful, slender
gradatim, gradually
gradus, us, m., grade, step
gramen, inis, n., grass
gramineus, grassy
graminicola, growing on grass
grandis, large
grandiusculus, somewhat large
granulatus, granular
granulosus, granular
graphidoideus, long and cleft, like

Graphis

graveolens, of unpleasant odor
gregarius, gregarious, in clusters
gregatim, in clusters
grex, gregis, m., a flock
griseolus, grayish
griseus, gray
grossus, thick
grumosus, heaped
grumulus, i, m., a heap
gumosus, gummy
gutta, ae, f., a vacuole
guttatus, with little drops
guttula, ae, f., a drop or vacuole
guttulosus, with drops
gyalectoideus, Gyalecta-like
gypseus, gypsum-like
gyrosus, gyrose, spiral

H

habeo, to have
habitatio, onis, f., habitat
habitus, us, m., habit

hactenus, up to the present time
haerens, adhering
haereo, to hold to
halos, o, f., a halo
hamatus, hamate, hooked
haud, not at all
haustorium, ii, n., a sucker
helicoideus, spiral-like
heliotropicus, heliotropic
helvolus, deep purple
herba, ae, f., a plant
herbicola, dwelling on herbs
heterogamete, one of two unlike sex-cells
heterogeneus, different
heteroicus, on two hosts
heteromorphus, heteromorphic, of different kinds
hexagonus, hexagonal
hexasporus, six-spored
hians, gaping
hiascens, gaping
hibernans, resting
hicillic, here and there
hinc, hence
hirtellus, somewhat shaggy
hodiernus, of today
homogeneous, homogeneous
homoicus, on one host
homomorphus, alike, of one form
horizontalis, horizontal
hornotinus, of this year
hortus, i, m., a garden
hospes, itis, m., a host
hospitalis, of a host
huc, hither, in this direction
humectatus, wet
humectus, moist
humidulus, moist
humilis, low, small
humistratus, moist
humus, i, f., the earth
hyalinulus, somewhat clear
hyalinus, hyaline, clear
hyalosporus, with clear, one-celled spores
hydrophilus, aquatic
hygrometricus, absorbing moisture
hygrophanus, translucent
hymeniferus, membrane-bearing

hymenium, ii, n., fruiting surface, consisting of asci, or of basidia.
hymenophorum, i, n., that which bears the hymenium
hypertrophians, hypertrophying
hypha, ae, f., a fungus filament
hyphasma, atis, n., the mycelium.
hyphoideus, hypha-like
hypohomycetus, mould-like, cobwebby
hypocreaceus, *Hypocrea*-like, fleshy and bright-colored
hypodermicus, under the epiderm
hypogaeus, hypogaeal, underground
hypogenus, on the under side
hypophloeodus, under the bark
hypophyllus, on the under side of leaf
hypostroma, atis, n., lower stroma
hypothallus, i, m., hypothallus
hypothecium, the area just below the layer of asci
hysteriformis, *Hysterium*-like, long and cleft
hysterinus, long and cleft as in *Hysterium*
hysterothecium, an oblong or linear perithecium opening by a cleft

I.

ibi, there, then
icon, onis, f., an image, figure
idem, the same
ideoque, therefore
idoneus, fit
igitur, therefore, accordingly
ignotus, unknown
imbricatus, imbricate
immaculatus, without spots
immarginatus, without a margin
immaturus, young
immediate, directly
immersus, sunken
immutatus, unchanged
impalpabilis, extremely fine and minute
impervius, impervious
implens, filling
implexus, infolded
impolitus, not polished
impositus, imposed

- imprimis*, especially
improbabile, improbably
imus, lowest
inaequilateralis, unequal-sided
inaequaliter, unequally
inaequipolaris, with unequal poles
inanis, empty
inarticulatus, without divisions
incarceratus, hidden
incarnatus, pink
incertus, uncertain
incisio, onis, f., incision, cutting
incisus, cut
inclinatus, bent
inclusus, inclosed
incoctus, not cooked
incolens, dwelling in
incoloratus, without color
inconditus, confused, unformed
incrassatulus, somewhat thickened
incrassatus, broadened, thickened
increasco, to grow in, increase
incumbens, lying upon
incurviusculus, somewhat incurved
incusus, forged, made
indeterminatus, indefinite
indico, to indicate
indigito, to utter, announce
indivisus, undivided
indoles, is, f., nature, natural ability
indumentum, i, n., a covering
induratus, hardened
indurescens, growing hard
indusium, ii, n., indusium
indutus, covered
ineptum, improper
inermis, unarmed
inferior, lower
inferus, below, lower
infestans, infesting
inficiens, infecting
inifimus, lowest
infixus, fastened in
inflans, inflating
inflatus, inflated
infossus, sunken
infra, lower, below
infundibuliformis, infundibuliform, funnel-shaped
infuscatus, darkened
initio, at first
initium, ii, n., the beginning
innatus, innate
innotesco, to become clear
innumerus, innumerable
inordinatus, without order
inquinans, blackening
inquinatus, dirty
inquirendus, to be investigated
insculptus, insculptate, hollowed
insectum, i, n., insect
insertio, onis, f., insertion
insertus, inserted
insidens, seated upon
insitus, ingrafted
inspersus, scattered
inspissatus, thickened
instar, like
instructus, built up
insuetus, unusual
insula, ae, f., an island
integer, whole
intense, intensely
intercalary, in the midst of, between
interdum, sometimes
interim, meanwhile
intermedius, intermediate
intermixtus, mixed with
internervius, between the nerves
internus, internal
interspersus, interspersed, scattered
interstitium, ii, n., a space
intertextus, intertwined
intus, within
intracellularis, within the cell
intrans, entering
intricatus, intertwined
intumescens, swelling
intus, within
invasus, invaded
inversus, inverted
investiens, covering
invicem, in turn, mutually
involutum, i, n., involucre
ipse, self
irregularis, irregular
irregulariter, irregularly
irrepens, creeping in
irroratus, bedewed
isabellinus, isabel-colored

isogamete, one of two similar sex-cells

isthmus, *i, m.*, a connection

itaque, therefore

iteratus, repeatedly

J

jacio, to throw

jamdudum, this long time

jodicus, of iodine

jodus, *i, m.*, iodine

junior, younger, young

jus, juris, *n.*, law, right

juvenilis, young

juxta, near

L

labiatus, lipped

labium, *ii, n.*, lip

labrum, *i, n.*, a lip

labyrinthus, labyrinthian, tortuose

laccatus, milky

lacerans, tearing

laceratus, lacerate, torn

lacerus, torn

lacinia, *ae, f.*, a tear

laciniatus, lacinate, torn, lobed

lacrimiformis, tear-like

lactens, milky

lactescens, milky

lactiginosus, filled with milk, milky

lacuna, *ae, f.*, a hole

lacunosus, lacunose, with hollows

lac, lactis, *n.*, milk

lacus, *us, m.*, a lake

laeticolor, bright-colored

laetus, bright

laevis, smooth

lageniformis, flask-shaped

lamella, *ae, f.*, gill

lamina, *ae, f.*, scale, layer, blade

laminaris, leaf-like

lanatus, woolly

lanceolatus, lance-shaped

languens, withering

lanosus, woolly

lanuginosus, woolly

laricinus, of larch

larva, *ae, f.*, larva

lateritius, brick red

latitudo, *inis, f.*, width

latiusculus, somewhat wide

latus, *eris, n.*, the side

latus, broad, wide

laxus, loose

lectus, collected

lego, to collect

leiosporus, with smooth spores

leniter, slightly, gently

lenticularis, lenticular, lens-shaped

lentiformis, lens-shaped

lentus, tough, flexible

leporinus, of a hare

leptodermus, thin-walled

leprosus, scab-like

leucosporus, with white spores

levis, light, smooth

levitas, *atis, f.*, smoothness

liber, free

liberatus, freed

lichenicola, lichenicole, growing on lichens

lichenoides, lichen-like

ligneus, woody

lignatilis, of wood

lignicola, lignicole, growing on wood

lignum, *i, n.*, wood

lilacinus, lilac-colored

limbatus, bordered

limbum, *i, n.*, limb, border

limes, *itis, m.*, limit

limitatus, limited

limoniformis, lemon-shaped

linea, *ae, f.*, line

linearis, linear

lineola, *ae, f.*, little line

linguiformis, tongue-shaped

liquifaciens, liquifying

liquo, to melt

lirella, *ae, f.*, furrow

lirelliform, furrow-like

lividus, livid, purple

lobulatus, somewhat lobed

locandus, to be located

locatus, located

locellatus, with chambers

locellus, *i, m.*, a little cell

loco, to place, locate

loculiferus, containing hollows

loculus, *i, m.*, locule, place, cell, hollow

locus, i, m., place
 longicollus, with long beaks
 longior, longer
 longitrorsum, longitudinally
 longitudinalis, lengthwise
 longus, long
 lophus, i, m., a crest
 lubricus, slippery
 lucidus, clear, lucid
 ludibundus, playful
 lumen, inis, n., opening
 lunatus, crescent-shaped
 lunulate, crescent-shaped
 luridus, lurid
 luteus, yellow
 lutescens, yellowish
 lux, lucis, f., light

M

maceratus, softened
 macro-, large
 macrostylospora, ae, f., large stylo-
 spore
 macula, ae, f., a spot
 macularis, spotted
 maculicola, dwelling on spots
 maculiformis, spot-shaped
 madidus, moist, wet
 magis, more
 magniguttatus, with one or two large
 globules
 magnitudo, inis, f., size
 magnus, great, large
 majusculus, somewhat large
 male, poorly
 mamillaris, protuberant
 mamilliformis, shaped like a papilla
 manifestus, evident
 mappa, ae, f., a map
 marcescens, withering
 marginatus, margined
 margo, inis, m., and f., margin
 marmoratus, marble-like
 massa, ae, f., mass
 massula, ae, f., a little mass
 matricalis, belonging to the matrix
 matrix, icis, f., matrix, layer or tis-
 sue
 maturus, mature

matrescens, ripening
 maxime, greatly
 mazaedium, i, n., a dough-like mass
 of spores and paraphyses
 medietas, atis, f., middle
 mediocris, average
 mediocriter, moderately
 medius, i, m., medium
 medulla, ae, f., the pith, medulla
 medullary, belonging to the pith or
 medulla
 medullatus, stuffed, pithy
 melanosporus, with black spores
 melioideus, meliola-like
 melius, better
 melleus, honey-colored
 mellinus, honey-colored
 membrana, ae, f., membrane
 membranaceus, membranaceous, mem-
 branous, thin or membrane-like
 memoria, ae, f., memory
 mens, mentis, f., mind
 merenchymaticus, with many cells
 merens, deserving
 meridionalis, southern
 mesogenus, mesogenous, borne in the
 middle
 mesopodes, with stem in the middle
 mesopus, with central stalk
 metageneticus, metagenetic
 metallicus, metallic
 metiens, measuring
 metulaeformis, pyramid-shaped
 metuliformis, pyramid-shaped
 micro-, small
 microconidiophorus, bearing small
 conidia
 microcystis, small-celled
 micronemeus, with short hyphae
 micropycnidium, ii, n., small pycnidi-
 um
 microscopium, ii, n., microscope
 microstylospora, ae, f., microstylo-
 spore
 migro, to move
 miniatus, bright red
 minimum, least
 minor, smaller
 minuties, ei, f., detail
 minutus, minute

mitis, pleasant, mild
mitratus, mitre-shaped
mobilis, mobile, moving
molecularis, molecule-like
molliusculus, somewhat smooth
mollis, smooth
moneo, to caution, warn
monile, is, n., a chain, necklace
moniliformis, chain-like
monoascus, with one ascus
monocephalus, monocephalic, one-headed
monocyclus, with one cycle
monoicus, monoecious
monoplastus, uniform, with one pro-toplast
monospermus, one-spored
monosporus, one-spored
monostichus, monostichous, in one row
mons, tis, m., a mountain
monstrosus, monstrous
montanus, mountainous
montosus, mountainous
morbosus, diseased
moriens, dying
mos, m^{oris}, m., manner
motilis, motile, able to move
movens, moving
mox, at length
mucedineus, white and cottony
mucilago, inis, f., mucilage
mucosus, mucose, slimy, mucous
mucus, i, m., mucus
mucro, onis, m., a point
mucronatus, pointed
mucronulatus, with a little point
mucronulus, i, m., a little point
multifidus, multifid, many-divided
multiguttatus, with many oil-drops
multilocularis, many-celled
multiloculatus, with many cells
multinucleate, with many nuclei
multisporus, many-spored
multizonatus, with many zones
multoties, many times, often
multus, much
munitus, furnished
muralis, muriform
muriculatus, muriculate, spiny

muriformis, muriform, with cross and longitudinal walls
murinus, mouse-colored
murus, i, m., wall
muscosus, mossy
mutans, changing
mutatus, changed
muticus, muticate, not pointed
muto, to change
mutue, mutually
mutuus, mutual
mycelialis, mycelial
mycelicus, mycelial
mycelium, ii, n., mycelium
mycogenus, dwelling on fungi
mycologus, i, m., a student of fungi
myochrous, mouse-colored
myriosporus, with many spores
mytiliform, shell-like

N

nascens, arising
nascor, to be born
natalis, native
naufragium, ii, n., shipwreck
navel, point of attachment
navicularis, boat-shaped
nebulosus, nebulous, cloudy, dark
nec, not
nectriaceus, Nectria-like
nemorosus, woody, shady
neque, and not
nervicola, growing on veins
nervi-sequus, **nervi-sequens**, following the veins
nidulans, nesting
nidulor, to nest
niduo, to nest
niger, black
nigredo, inis, f., blackness
nigresco, to grow black
nigricans, blackening
nigrifactus, blackened
nigrificatus, made black
nigrolimitatus, black-lined
nigropilus, black-hairy
nigropunctulatus, black-dotted
nigrostrigosus, black-hairy
nimum, too, too much
nisi, unless

nitens, shining
niteo, to shine
niveus, snow-white
nobilis, grand
nodosus, with joints
noduliferus, bearing knots
nodulosus, with joints
nodus, i, m., a joint, knot
nomen, inis, n., a name
non, not
nondum, not yet
nonne, not
nonnihil, somewhat
nonnisi, except
nonnullus, some
normalis, normal
notatus, marked
notus, known
novus, new
nubecula, ae, f., a little cloud
nubilosus, cloudy
nucleatus, nucleate
nucleiferus, nucleus-bearing
nucleolus, nucleole
nucleus, i, m., center, nucleus
nudiusculus, somewhat naked
nudus, naked
nullimodus, in no wise
nullus, none
numerosus, numerous
numerus, i, m., a number
numquam, never
nunc, now
nutiquam = **ne-utiquam**, by no means
nuto, to incline
nutrix, icis, f., host
nux, nucis, f., a nut

O

ob, for, toward, on account of
obclavatus, reversed club-shaped
obconicus, reversed-conical
obducens, covering
obduco, to cover
oblique, obliquely
obliterans, disappearing
obliteratus, lost, destroyed
oblongatus, oblong
oblongus, oblong
obpyriformis, obpiriform, reversed
 pear-shaped
obrutus, covered
obscurus, dark
observandum, to be observed
observatus, found
obsessus, surrounded
obsolesco, to become obsolete
obsoletus, obsolete, lacking
obtectus, covered
obtegens, covering
obturaculum, i, n., opening
obtusangulus, with obtuse angles
obtusatus, obtuse
obtusus, obtuse
obtutus, us, m., a looking at
obvallatus, surrounded
obvelo, to cover
obvius, clear, open
obvolvens, enveloping
occellatus, with openings
occulo nudo, with unaided eye
occupans, occupying
ochraceus, pale yellow, ochreous
ochrosporus, with yellow or yellow-
 brown spores
octavus, eighth
octo, eight
octonus, in eights
octoseptatus, with eight cross-walls
octosporus, eight-spored
oleosus, oily, with oil drops
oligosporus, few-spored
olim, formerly
olivascens, olivascent, becoming olive
olivaceus, olive
omissus, omitted
omnino, everywhere, entirely
oosporous, with resting spores formed
 by the union of unlike sex-cells, e.g.,
 of egg and sperm
opacus, opaque
opalinus, clear
operculatus, operculate, with a lid
operculiformis, lid-shaped
operculum, i, n., a cover, lid
oppidum, i, n., a town
oppletus, filled
oppositus, placed
orbicularis, orbicular, round
orbiculatim, circularly

orbis, is, m., a circle
ordo, inis, m., order
organicus, organic
organum, i, n., an organ
oriens, arising
orientalis, eastern
orificium, i, n., opening
originalis, original
origo, inis, f., origin
orior, to arise
ornatus, furnished
orthotropus, straight
ortus, arisen
os, oris, n., mouth
oscillans, oscillating
osculum, i, n., mouth
ostendo, to show
ostiolatus, ostiolate, with a mouth
ostiolum, i, n., ostiole, opening
ovalis, oval
ovaricola, growing in ovaries
ovatus, egg-shaped
ovinus, of or belonging to a sheep
ovoideus, nearly egg-shaped

P

pachydermaticus, thick-walled
pachypleurus, thick-walled
paene, nearly
paenultimus, next to the last
pagina, ae, f., page, side
paliformis, paliform, stake-shaped,
 palisade-like
pallescent, turning pale
pallidus, pale
palmatus, palmate, hand-like, palm-
 like
palmicola, growing on palms
palpebra, ae, f., eyelid
paludosus, marshy
palumbinus, dove-colored, grayish
palus, udis, f., a marsh, swamp
panícula, ae, f., a panicle
paniculatus, paniculate, branched
panis, is, m., bread
pannosus, pannose, ragged
pannum, i, n., a rag, cloth
papillaris, papillate
papillatus, with papilla, papillate
papilliformis, like a papilla
papillula, ae, f., a little papilla
papillulatus, with a very small nipple
 or papilla
papulosus, with many pustules
papyraceus, papery
paradoxus, strange, contrary
parallelus, parallel
parasiticus, parasitic
parcus, few, scanty
parenchymaticus, parenchyma-like
paries, etis, m., a wall
paritas, atis, f., equality
paroecchia, ae, f., parish
pars, partis, f., a part
partitus, divided
parum, too little
parvulus, small
parvus, small
pascuum, i, n., pasture
passim, everywhere
patellaris, dish-like
patelliformis, shaped like a dish
patens, spreading
patenter, openly
pator, to support, endure
patulus, spreading
paucilocularis, few-celled
paucus, few
paulatim, gradually
paulisper, for a little while
paulo, a little
pectinatus, comb-like
peculiaris, peculiar
pedatus, foot-like
pedicellatus, with a pedicel
pedicellus, i, m., pedicel
pediculatus, pedicelled
pedunculatus, stalked
pedunculicola, growing on peduncles
pellicle, skin, covering
pellicula, ae, f., a little skin
pelliculosus, with a covering
pelluciditas, atis, f., clearness
pellucidus, pellucid, clear
peltatus, shield-shaped
pendo, to hang
pendulus, hanging
penetrans, penetrating
penicillate, brush-like
penicilliformis, brush-like

- pentagonus**, pentagonal
per, through
peraffinis, closely related
perbrevis, very short
percursus, run through
perdurans, resting
perduro, to last
perennans, perennial
perennis, perennial
perexiguus, very thin
perexilis, very slender
perfectus, complete, perfect
perforans, perforating
perforatus, perforated
perfossus, hollowed out
pericarpium, ii, n., pericarp, covering
peridermicus, belonging to the periderm
peridermium, ii, n., periderm
peridium, ii, n., peridium
periphericus, peripheral around the edge
peristomium, ii, n., mouth
perithecialis, perithecial
perithecigerus, perithecium-bearing
perithecioid, perithecium-like
peritheciophorus, bearing perithecia
peronatus, rough, rough-booted
perparum, very little
perrumpens, breaking through
persicinus, peach-colored
persistans, persistent
perspicuus, transparent
perspicuus, clear
persuasus, convinced
pertenuis, very thin
pertineo, to belong
pertusus, protruded
pes, pedis, m., foot
petiolum, i, n., petiole
petrifactus, made like rock, hardened
pezizoideus, pezizoid, cup-fungus-like, cup-like
phacidiodeus, like *Phacidium*, black and disk-like
phaeophragmeus, with dark transeptate spores
phaeosporus, with dark, one-celled spores
phaseoliformis, bean-shaped
phomatoideus, *Phoma*-like
phyllogenus, phyllogenous, borne on leaves
phyllostictodeus, *Phyllosticta*-like
phytogenus, growing on plants
phytographus, i, m., a botanist
phytophilus, phytophilous, growing on plants
pictura, ae, f., a painting
pictus, colored
pileatus, cap-shaped
pileus, i, m., a cap
pilosellus, somewhat hairy
pilosus, pilose, with hairs
pilum, i, n., a hair
pineus, piny
pingo, to paint
pinna, ae, f., a leaflet
pinnatus, pinnate
piperatus, peppery, pungent
piscis, is, m., a fish
pisum, i, n., pea
placenta, ae, f., placenta
placentiformis, placenta-like
plaga, ae, f., a spot
plagula, ae, f., a little spot
plaguliformis, spot-like
planta, ae, f., a plant
plantula, ae, f., a little plant
planus, plane, flat
plasma, atis, n., plasm, mass
plasmodium, ii, n., protoplasm-like mass
pleiosporus, many-spored
plenus, full
plerumque, for the most part
pleuroacrogenus, borne at the tip and at the sides
pleurogenus, pleurogenous, borne on the walls or sides
plica, ae, f., a fold
plicatus, plicate, folded
pliciformis, fold-form
plumbeus, lead-colored
plures, many
pluriarticulatus, many-celled
pluriciliate, with many cilia
plurifurcatus, many forked
pluriguttulatus, many guttulate
plurilocellatus, with many hollows

pluriperforate, with several openings
pluristratosus, many-layered
poculiformis, cup-shaped
podetium, i, n., a stalk-like or cup-like erect thallus
polaris, polar
politus, polished
polleo, to be able, avail
pollex, icis, m., thumb
pollicaris, thumb-like, an inch long
polus, i, m., a pole
poly-, many
polyascus, with many asci
polyblastus, many-celled
polycephalus, polycephalous, with many heads
polyedricus, polyhedral
polygonus, with many angles
polyrrhizus, with many roots
polystichus, polystichous, in many rows
pondus, eris, n., weight
populus, i, f., poplar
porosus, with pores
porrigo, to stretch out
porus, i, m., a pore
positus, placed
possum, to be able
postea, hereafter
postice, at the back
postremus, last
potius, rather
praecedens, preceding
praecipue, especially
praeclarus, distinguished
praecox, early, abundant
praeditus, furnished
praeferendum, preferred
praelongus, very long
praeprimis, especially
praesens, present
praesertim, particularly
praestans, distinguishing, excelling
praesumptus, assumed, presumed
praetereaue, besides, moreover
praeteritus, past
pratium, i, n., a meadow
primitivus, primitive
primitus, at first
primus, first

prioritas, atis, f., priority
prismaticus, prismatic
privus, without, deprived
pro, for
probabilis, probable
procerus, tall
processus, projection
procumbens, procumbent, prostrate
prodeuns, projected
productus, carried out, produced
proficiscor, to begin, arise
profunditas, atis, f., depth
profundus, deep
projectus, thrown off
proles, is, f., a race, offspring
proliferus, proliferous, produced, proliferate
proliger, bearing offspring
prolongatio, onis, f., prolongation; lengthening
promycelium, i, n., promycelium
prope, near
proper exciple, an apothecial covering or wall without algae
propius, proper
propinquus, adjacent
propulsus, expelled
proratione, comparatively
prorsus, forwards, exactly
prorumpo, to break through
prosenchymaticus, prosenchymatic, consisting of long cells or filaments
proteus, changing, variable
protractus, extended
protrudens, projecting
provectus, prolonged, advanced
proveniens, coming
pruinulosus, somewhat powdery
pruinosis, powdery, pruinose
pseudo-, false
pseudoparaphyses, false paraphyses
pseudoparenchyma, false parenchyma, a tissue looking like parenchyma but formed of threads
pseudoperidium, a covering
pseudoplasmodium, ii, n., a false plasmodium
pseudopodium, ii, n., false foot, lobe
pseudostiolum, i, n., false ostiole

pseudostroma, *atis*, *n.*, a false stroma
pseudostromaticus, resembling a stroma
ma
pseudothallus, *i*, *m.*, false thallus
puberulus, somewhat hairy
pubescens, hairy
pubes, *is*, *f.*, hair
puccinoideus, puccinia-like
pulchellus, beautiful
pulcher, beautiful
pulchre, beautifully
pulpa, *ae*, *f.*, pulp, mass
pulveraceus, powdery
pulverulentus, powdery
pulvinatus, cushioned
pulvinulus, *i*, *m.*, a little cushion
pulvis, *eris*, *m.*, powder
punctiformis, punctiform, dot-like
punctulans, dotting
punctulatus, punctate, dotted
purpurascens, becoming purple
purus, pure
pusillus, tiny
pusio, *onis*, *m.*, a growth
pustula, *ae*, *f.*, a mass
pustulate, pertaining to a swollen
mass
putamen, *inis*, *n.*, a shell
putredo, to decay
putrescens, decaying
putris, decaying
pycnidicus, pycnidial
pyramidatus, pyramidal
pycnidium, *i*, *n.*, pycnidium
pyreniformis, pyreniform, shaped like
a nut
pyriformis, pear-shaped
pyxidatus, like a box

Q

quadricoccus, of four round cells
quadripartitus, four-divided
quadriflorus, four-spored
quadratum, *i*, *n.*, a square
qualis, like
quam, than
quandoque, whenever, at some time
quartus, fourth
quasi, almost
quater, four times

quaternus, by fours
quattuor, four
quercinus, oaken
quia, because
quinqueseptatus, five septate
quisque, each
quisquillae, *arum*, *f.*, dirt, trash
quoad, as long as, as much as
quod, that
quoque, also

R

racemulus, *i*, *m.*, a little raceme
racemus, *i*, *m.*, a bunch of grapes, raceme
rachis, *is*, *f.*, axis
radians, radiating
radiatim, radiately
radiatus, radiate
redicalis, basal
radicans, root-like, rooting
radicatus, radicate, more or less rooted
radiciformis, root-shaped
radicosus, having many roots
radix, *icis*, *f.*, a root
ramicola, ramicole, living on twigs
ramosus, much branched
ramulus, *i*, *m.*, a little branch
ramus, *i*, *m.*, a branch
rarius, more rarely
raro, rarely
rasus, leveled
reabsorptus, reabsorbed
recedo, to recede, differ
recensio, *onis*, *f.*, a reviewing
recludens, opening
recognoscens, recognizing
rectangularis, rectangular
rectangulus, rectangular
rectus, straight
reddo, to return, restore
refractus, turned back
refringens, refringent
regio, *onis*, *f.*, region
relatus, related
relinquens, leaving
relinquo, to leave
reliquus, left, remaining
remote, distantly

remotiusculus, somewhat distant
reniformis, reniform, kidney-shaped
repandus, turned back
repens, creeping
reperio, to find
repertorium, ii, n., an inventory, catalogue
repertus, found
repetite, repeatedly
repetitus, repeated
repletus, full
repo, to crawl
res, rei, f., a thing
resolvens, breaking up
resorptus, absorbed
resupinatus, resupinate, horizontal,
the hymenium turned up
reticulatus, reticulate, net-like
reticulum, i, n., a net
retiformis, net-like
retineo, to retain, keep
retis, is, f., a net
retrorsus, backward
retusus, with a little sinus
revelo, to reveal, uncover
revivescens, reviving
revoco, to recall
revolutus, folded back
rhabarbarinus, yellow
rhizoid, root
rhizoideus, root-like
rhizomorphaeus, root-like
rhizophilus, growing on roots
rhodosporus, with rose-colored spores
rhombius, rhombic
rhomboideus, rhomboid
rhytismoideus, Rhytisma-like
ricciiformis, like Riccia, a liverwort
rigens, stiff, rigid
rigidulus, somewhat stiff
rigidus, stiff
rima, ae, f., cleft
rimosus, rimose, cleft, cracked,
ripa, ae, f., bank
rite, rightly, fitly, well
rivulosus, with channels
rivus, i, m., brook
robustus, robust
roridus, like dew
ros, roris, m., dew

roseolus, somewhat rosy
roseus, rose-colored
rostellatus, somewhat beaked
rostratus, rostrate, beaked
rostriformis, beak-like
rostrum, i, n., beak
resulatus, rosette-like
rotundatus, rounded
rubeolus, somewhat reddish
ruber, red
rubellus, somewhat reddish
rubescens, growing red
rubiginosus, rust-colored
rubricosus, reddish
rufescens, becoming reddish
rufus, reddish
rugosiusculus, more or less wrinkled
rugulosus, furrowed, roughened
rumpens, breaking
ruptus, broken
rursus, backward
rutilus, red

S

saccatus, saccate, sac-like
saccharinus, sugary
saccharum, i, n., sugar
sacciformis, sac-shaped
sacculiformis, like a little sac
sacculus, i, m., a little sac
saepe, often
salicinus, of willow
salmonicolor, salmon colored
salmonius, salmon-colored
saltem, at least
samara, ae, f., key fruit
samariform, key-shaped
sanguineus, bloody, blood-colored
sapidus, filled with sap, savory
sapor, oris, m., flavor
saprogenus, saprogenous, growing on
decayed matter
saprophilus, growing on decaying
matter
saprophyticus, saprophytic
sarciniformis, sarciniform, packet-
like
sarmentum, i, n., twig
satis, sufficient
saturatus, saturated

- scaber*, rough
scabridus, rough
scabriusculus, somewhat rough
scalaris, of a ladder, or staircase
scaliformis, ladder-like
scariosus, thin, papery
scheda, æ, f., sheet of paper
scio, to know
scissilis, splitting
sclerotiformis, sclerotium-like
sclerotioideus, sclerotoid, sclerotium-like
sclerotium, i, n., sclerotium, a hard black mass
scolecosporus, with thread shaped spores
scopulate, like a brush
scrobiculatus, roughened, furrowed
scrotiformis, bladder-like
scruposus, rough
scrutator, oris, m., an investigator
scutatus, shield-shaped
scutellatus, like a small shield
scutiformis, shield-shaped
secedens, separating
secernibilis, separable
sectio, onis, f., a section
secundarius, secondary
secundum, according to
secus, otherwise
sed, but
sedulus, diligent, careful
segmentiformis, segment-like
sejunctus, separate
semel, once
semen, inis, n., a seed
semi, half
semiexertus, half extended
semiimmersus, half immersed
semiinfossus, (cf. *infossus*)
semiinsculptus, (cf. *insculptus*)
seminalis, seed-like
seminicola, growing on seeds
semipellucidus, half-pellucid
semiteres, half columnar
semiuncialis, a half inch
semper, always
senescens, growing old
sensim, gradually
sensus, us, m., opinion, sense
separabilis, separable, separating
separo, to separate
sepimentum, i, n., partition
sepono, to separate
septatus, septate, divided into cells
septentrionalis, northern
septulum, i, n., a little septum
sepulchrum, i, n., grave
sequens, following
sericellus, somewhat silky
sericeus, silky
series, ei, f., a series
serotinus, late
serpens, creeping
serpentinus, serpentine
serratus, serrate
serus, late
sesqui, by a half
sesquilinea, one inch and a-half
sesquipedalian, very long
sessilis, seated, without a stalk
seta, æ, f., a bristle
setaceus, bearing one or more bristles
setiformis, bristle-shaped
setiger, bristle-bearing
setosus, setose, with bristles
setula, æ, f., a little bristle
setulose, with bristles or spines
seu, or
sexilocularis, with six cells or locules
sexsporus, six-spored
sexsulcatus, six-furrowed
siccans, drying
siccus, dry
sigillatim, seal-like
sigmoideus, sigmoid, s-like
signatus, marked
sileo, to be silent
silva, æ, f., a forest
similaris, like
similis, similar
simple, not branched; one-celled (of spores)
simplex, icis, simple
simul, at the same time
simulate, apparently
simulo, to imitate, copy, represent
sine, without
singularis, peculiar, not in chains
singulus, each

sinuatus, sinuate
sinuosus, crooked
sistens, comprising
situs, placed
socia, *ae*, *f.*, society
sociatus, grouped together
soleo, to be accustomed
solidiusculus, somewhat solid
solitarius, solitary
solitus, usual
sollertus, distinguished
solubilis, dissolving
solutus, dissolved
sordes, *is*, *f.*, dirt
sordidus, dirty
sorus, *i*, *m.*, spore mass
spadiceus, brownish
spatha, *ae*, *f.*, a spathe
spargo, to scatter
sparsus, scattered, sparse
spathulatus, spathulate
spatium, *i*, *n.*, space
specialis, special
species, *ei*, *f.*, species
spectans, looking
specto, to look
spermagonium, *ii*, *n.*, a pycnidium-like body
spermatiferus, spermatia-bearing
spermatiformis, like a spermatium
spermatioideus, spermatium-like
spermatium, *ii*, *n.*, a conidium-like body
spero, to hope
sphaericus, spherical
sphaeroideus, nearly spherical
sphaerula, *ae*, *f.*, a sphere
spica, *ae*, *f.*, a point, ear
spicatus, spike-like
spiculosus, spiny
spiculum, *i*, *n.*, a little spine
spiniformis, spiny
spinuligerus, spine-bearing
spinulosus, with little spines
spira, *ae*, *f.*, a spiral
spiralis, spiral
spiraliter, spirally
spiritus, *us*, *m.*, a spirit
spissus, thick
splendens, splendid

spongilliformis, sponge-like
spongiosus, spongy
sponte, spontaneously
sporangiferus, bearing sporangia
sporangioliiferus, bearing small sporangia
sporangiolum, *i*, *n.*, a little sporangium
sporangiophore, the stalk of a sporangium
spore-print, the spore mass obtained by placing the cap of a mushroom flat on a piece of white paper
sporicus, sporal
sporidiolum, *i*, *n.*, a little spore
sporidium, *i*, *n.*, a spore
sporiferus, spore-bearing
sporodochium, a compact, conidial body, mass of sporophores
sporomorphus, spore-shaped
sporophora, *ae*, *f.*, sporophore
spurius, false
squama, *ae*, *f.*, a scale
squamosus, scaly
squarrose, with spreading scales or hairs
statura, *ae*, *f.*, stature
status, *us*, *m.*, stage
stellatus, stellate, star-like
stelliformis, star-shaped
stercoratus, manured
stercus, *oris*, *n.*, dung
sterigma, *atis*, *n.*, stalk
stilbeus, stilbum-like, mallet like
stilbiformis, stalk-like
stilboid, with a stalked-head, Stilbum-like
stipatus, crowded
stipes, *itis*, *m.*, a stalk
stipitatus, stipitate, stalked
stipitellus, *i*, *m.*, a little stalk
stipitiformis, stalk-like
stoloniferous, producing runners
stoloniformis, runner-like
stramineus, straw-colored
stratosus, in layers
stratum, *i*, *n.*, a layer
strenuus, prompt, vigorous
stria, *ae*, *f.*, a line

strigosus, strigose, long or coarsely hairy
striiformis, line-like
strobilus, i, m., a cone
stroma, atis, n., a covering, layer
stromaticus, stromatic
stromatiferus, bearing a stroma
structura, ae, f., a structure
stupposus, tow-like
stylospora, ae, f., a stylospore
suadens, persuading
suavis, pleasant
sub, affix meaning somewhat, slightly
subacutus, somewhat acute
subaequans, nearly equal
subalbus, nearly white
subalutaceus, somewhat yellow
subastomous, more or less mouthless
subbulbosus, somewhat bulbous
subcarbonaceus, slightly carbonaceous
subcarnulosus, slightly fleshy
subclavatus, subclavate
subclypeate, somewhat shield-shaped
subcolumelliformis, somewhat like a columella
subconoideus, slightly conical
subcrustose, somewhat crust-like
subcuboideus, somewhat cubical
subcutaneus, under the epidermis
subdeterminatus, limited
subdiscoideus, somewhat disc-shaped
subelevatus, somewhat raised
suberosus, suberose, corky
subfuscus, subfuscous, somewhat dark
subglobosus, subglobose
subiculum, i, n., subicle, a compact cottony mycelium
subimmersus, slightly immersed
subinde, presently, forthwith, now and then
subito, suddenly
subnullus, nearly lacking
substantia, ae, f., substance
subterraneus, subterranean
subtilis, thin, slender
subtilitas, atis, f., fineness, thinness
subulatus, subulate, awl-shaped
subuliformis, awl-shaped
subvitro, under the lens

succresco, to grow under
suffultus, supported
sulcatus, sulcate, furrowed
sulcula, ae, f., a little furrow
sulcus, i, m., a furrow
sulphurellus, sulphurish
sulphureus, sulphur-colored
summa, ae, f., highest point, sum
superans, exceeding
superficialis, superficial
superficies, ei, f., the surface
superimpositus, superimposed
superpositus, superposed
superus, upper
supremus, uppermost
surculus, i, m., a shoot
sursum, upward
suspensor, supporting cell or group of cells
sustinens, supporting
sylva, ae, f., a forest (see *silva*)
sympodice, sympodially
synnema, atis, n., an erect fascicle of hyphae, as in *Stilbaceae*

T

tabesco, to melt
tactus, touched
taeniola, ae, f., a little band
talis, such
tamen, however, yet
tandem, at length
tantillus, so little
tapetum, i, n., nourishing layer
tarde, slowly, late
tartareus, powdery
tectus, covered
tegens, covering
tegmen, inis, n., a cover
teleutospora, ae, f., a teleutospore
teleutospiriferus, bearing teleutospores
tenacellum, somewhat tenaceous
tenellus, delicate
tentacula, ae, f., a tentacle
tentaculiformis, tentacle-shaped
tenuatim, drawn out
tenuis, slender
ter, three times
terete, cylindrical

- teretiusculus*, round, cylindric
terminalis, terminal
terminatus, terminated
ternate, in threes
ternus, three-fold
terra, ae, f., soil, earth
terrestris, terrestrial
tertius, third
testa, ae, f., a shell, coat
testaceus, brick-colored
tetradidymus, four-fold
tetragonus, four-angled
tetrasporus, four-spored
thalamium, i, n., a room
thallicola, growing on a thallus
thalliformis, thallus-like
thalline exciple, applied to an exciple containing algae
thallus, a more or less definite mass of hyphae parasitic on algae
thelephoroideus, thelephora-like
tigrinus, like a tiger
tinctus, tinged
tingens, tinging
tomentellus, hairy
tomentosus, hairy
tornatus, rounded-off
toruloideus, chain-like
torulosus, torulose, necklace-like
tortuosus, flexuous
tortus, twisted
totaliter, totally
totus, all
trabs, is, f., a beam
tractus, us, m., a tract
trahendum, to be drawn
trama, ae, f., a pathway
transeptate, with all cross-walls transverse
translucidus, clear
trapezoideus, trapezium-like
transiens, temporary
transversalis, transversal
tremelloideus, tremelloid, gelatinous
tremellosus, jelly-like
triangularis, triangular
tribus, us, f., a tribe
tricornutus, with three horns
trifoveolatus, with three hollows
trigonus, trigonous, three-angled
trilobus, three-lobed
trinacriformis, three-pronged
tripartitus, three-divided
tripedalis, three feet long
tripollicaris, three inches
triquetrus, three-cornered
trisporus, three-spored
tristichus, in three rows
tropicus, tropical
truncatus, cut-off
truncicola, growing on trunks
trunculus, i, m., a little trunk, stem
truncus, i, m., a trunk
tuber, eris, n., tuber, a swelling
tubercularinus, Tubercularia-like
tubercularioid, Tubercularia-like, warted
tubercularoideus, Tubercularia-like
tuberculiformis, wart-like
tuberculosus, roughened
tuberiform, tuber-like
tuberiformis, tuberiform, tuber-shaped
tubulosus, tubular
tubulus, i, m., a tube
tumescens, swelling
tumidulus, somewhat swollen
tumifactus, swollen
tunc, then
tunica, ae, f., cloak, coating
tunicatus, tunicate, covered
turbinatus, turbinate, top-shaped
turgescens, swollen
turgidus, swollen
turriiformis, shaped like a tower
turritus, turreted, tower-like
typice, usually, characteristically
typus, i, m., a type

U

- uber*, rich
ubi, where
ubiquemque, everywhere
udus, wet
uliginosus, rich, muddy
ullus, any
ultimus, last
ultra, beyond or more
-ulus, a, um, suffix, meaning small
umbellatus, umbellate, umbelled
umbelliformis, like an umbel

umbilicatus, umbilicate, with a navel, sunken in the center, somewhat funnel-form.

umbilicus, i, m., navel

umbonatus, umbonate, with a boss

umbra, ae, f., shade

umbrinus, brown

umbrosus, shady

uncia, ae, f., an inch

uncialis, an inch long

uncinatus, hooked

unde, whence

undique, in all directions

undulatus, wavy

uniarticulatus, one-jointed

unicus, single

uniformis, of one form

unilateralis, one-sided

unilocular, with a single cavity or cell

uniserialis, one-rowed

uniseriatus, one-rowed

unitus, joined

unquam, ever

urceolatus, pitcher-shaped

uredinicola, growing on rusts

uredospora, uredospore

uredosporiferus, bearing uredospores

urniformis, urn-shaped

uromorphus, tail-like

usque, up to

usurpatus, usurped

ut, as

uterque, both

ut-plurimum, for the most part

utriculiformis, bladder-shaped

utrimque, on both sides, in both directions

uvidus, moist, wet

V

vaccinus, pertaining to a cow

vacuus, empty

vage, vaguely

vagina, ae, f., a sheath

vaginatus, sheathed

vagus, vague

valde, strongly

validiusculus, more or less stout

valseus, valsous, valsooid, Valsa-like, with the perithecia in a circle in the stroma

valva, ae, f., a valve

valvatim, valvate, with valves

variabilis, variable

varie, variously

variegatus, of different colors

varius, different

-ve, or

vegetus, fresh, vegetating

venementer, strongly

vel, or

velatus, veiled

vellus, eris, n., fleece, wool

velo, to cover

velocitas, atis, f., swiftness

velum, i, n., a veil

veluti, as

velutinus, velvety

vena, ae, f., a vein

venenatus, poisonous

veniformis, vein-like

ventricosus, swollen

vere, truly

vergo, to approach

verisimiliter, apparently

vermicularis, worm-like

vermiformis, vermiform, worm-shaped

vernalis, vernal, of or belonging to spring

vero, truly

verruciformis, wart-like

verruculosus, verrucose, warted

versatus, poured

versicolor, of different colors

versiformis, of different forms

versus, towards

vertens, turning

vertex, icis, m., the tip

verticalis, vertical

verticillatim, in whorls

verticillatus, verticillate, whorled

vescus, small, weak

vesicula, ae, f., vesicle, swollen cell

vesiculosus, vesiculose, swollen, bladder

vestiens, covering

vestigium, i, n., remnant, vestige

vestio, to cover

vestitus, furnished, covered
vetustus, old
vibrans, changing
videor, to seem
vigens, growing
villosulus, somewhat woolly
villus, i, m., a hairy covering
vinarius, of wine
vineus, of or belonging to wine
vinum, i, n., wine
violaceus, violet
violascens, turning violet
virens, becoming green
virgatus, rod-shaped
viridarium, i, n., greenhouse
virgultum, i, n., bush, copse
viridifuscus, greenish brown
viridulus, greenish
viscidulus, viscid, somewhat sticky
visibilis, visible
visus, seen
vitellinus, yellow
vitreus, glassy
vivens, living
vividus, vivid
vivus, alive
vix, hardly

volva, ae, f., a cup-like sheath at the base of a stem
volvaceus, with a volva
volvatus, with a volva
vulgatus, common
vulgo, commonly
vulpinus, of a fox

X

xylogenus, xylogenous, growing on wood
xylophilus, growing on wood

Z

zona, ae, f., a zone
zonula, ae, f., a little zone
zoogenus, on animals
zoogonid, a motile propagative cell
zoospora, ae, f., zoospore
zoosporangium, ii, n., zoosporangium
zoosporiferus, producing zoospores
zygosporiacus, pertaining to a zygo-spore
zygosporous, with resting spores formed by the conjugation of similar sex cells
zymogenus, ferment-producing

INDEX

A

- Abrothallus, 69
 Absidia, 13
 Acallomyces, 19
 Acanthostigma, 34
 Acanthothecis, 59
 Acanthothecium, 59, 134
 Acarospora, 80
 Acerbia, 38
 Acerbiella, 38
 Acetabula, 89
 Achlya, 15
 Achlyogeton, 16
 Acinula, 164
 Acladium, 141
 Acolium, 71
 Acompsomyces, 19
 Acontium, 140
 Acremoniella, 148
 Acremonium, 141
 Acrocylindrium, 143
 Acroscyphus, 72, 92
 Acrospermum, 58
 Acrospira, 148
 Acrostalagmus, 143
 Acrotheca, 147
 Acrothecium, 152
 Actidium, 56
 Actiniceps, 155
 Actiniopsis, 34, 52
 Actinomma, 163
 Actinonema, 125
 Actinoplaca, 75
 Actinoscypha, 69
 Actinothecium, 131
 Actinothyrium, 132
 Acurtis, 105
 Adermatis, 79
 Aecidiella, 99
 Aecidiolum, 99
 Aecidium, 99
 Aegerita, 159
 Aegeritopsis, 161
 Agaricaceae, 110
 Agaricales, 102
 Agaricus, 114
 Aglaospora, 35
 Agyriella, 135
 Agyriellopsis, 133
 Agyrina, 67
 Agyriopsis, 67
 Agyrium, 67
 Ahlesia, 67
 Alboffiella, 115
 Albuginae, 17
 Albugo, 17
 Aldona, 56
 Aldridgea, 106
 Alecatoria, 82
 Aleuria, 88
 Aleurina, 89
 Allantonectria, 43
 Allantospora, 145
 Allarthonia, 58
 Allarthothelium, 58
 Allescheria, 22
 Allescheriella, 142
 Alliospora, 140
 Alternaria, 154
 Alveolaria, 101
 Amallospora, 163
 Amanita, 111
 Amanitopsis, 111
 Amaurascus, 93
 Amblyosporium, 140
 Ameghiniella, 66
 Amerosporium, 133
 Amoebocytrium, 11
 Amorphomyces, 19
 Amphichaeta, 137
 Amphisphaeria, 32
 Ampullaria, 129
 Anaptychia, 84
 Anapyrenium, 42
 Ancylistaceae, 16
 Ancylistae, 16
 Ancylistes, 17
 Anellaria, 114
 Anema, 73
 Angelinia, 56, 66
 Angiopoma, 126
 Anixia, 23
 Anxiopsis, 22
 Annularia, 112
 Anthina, 164
 Anthomyces, 101
 Anthostoma, 29
 Anthostomella, 28
 Anthracoderma, 123
 Anthracoidea, 101
 Anthracophyllum, 115
 Anthracothecium, 40
 Anthurus, 116
 Antromycopsis, 157
 Anzia, 81
 Aphanascus, 92
 Aphanomyces, 15
 Apiospora, 30
 Apiosporium, 23
 Aplacodina, 31
 Aplanes, 15
 Apodachlya, 16
 Aponectria, 45
 Aporophallus, 115
 Aposphaeria, 122
 Arachniotus, 93
 Arachnium, 119
 Arachnomyces, 23
 Arachnopeziza, 87
 Araeospora, 16
 Arcangelia, 30
 Arcangeliella, 120
 Arctomia, 73
 Arenaea, 87
 Areolaria, 119
 Argopsis, 78
 Argynna, 24
 Armillaria, 111
 Arrhenia, 111
 Arrhytidia, 105
 Arthonia, 58
 Arthoniactis, 76
 Arthoniae, 58
 Arthoniopsis, 58

Arthotheliopsis, 75
 Arthothelium, 58
 Arthrimum, 147
 Arthrobotrys, 144
 Arthrobotryum, 157
 Arthropryrenia, 41
 Arthrosporium, 156
 Aschersonia, 129
 Aschersoniopsis, 130
 Ascobolaceae, 92
 Ascobolae, 92
 Ascobolus, 92
 Ascochyta, 125
 Ascocorticaceae, 93
 Ascocorticium, 93
 Ascodes, 94
 Ascodesmis, 92
 Ascoideaceae, 93
 Ascoidea, 93
 Ascomycetella, 95
 Ascomycetes, 18
 Ascophanae, 92
 Ascophanus, 92
 Ascopolyporus, 48
 Ascotricha, 23
 Aseroe, 116
 Aspergillae, 140
 Aspergillus, 140
 Aspidopyrenium, 40
 Aspidothelium, 40
 Asterella, 53
 Asteridiella, 53
 Asteridium, 24, 53
 Asterina, 23, 53
 Asterinae, 52
 Asteristium, 76
 Asteroconium, 138
 Asterodon, 108
 Asteroma, 122
 Asteromella, 122, 131
 Asteromidium, 126
 Asteronia, 153
 Asterophora, 142
 Asteroporum, 41
 Asterosporium, 137
 Asterostroma, 107
 Asterothyrium, 75, 132
 Asterula, 23, 53
 Astrocystis, 28

Astrodochium, 162
 Astrotheliae, 42
 Astrothelium, 42
 Atractiella, 155
 Atractina, 152
 Atractium, 156
 Auerswaldia, 49
 Aulaxina, 59
 Aulographum, 56
 Aureobasidium, 107
 Auricularia, 103
 Auriculariae, 103
 Auriculariella, 103

B

Bacidia, 77
 Bacillus, 8
 Bacteriaceae, 8
 Bacteriales, 7
 Bacterium, 8
 Bactridiopsis, 159
 Bactridium, 161
 Bactroboletus, 109
 Bactrosphaeria, 38
 Bactrospora, 70
 Baculospora, 44
 Baeomyces, 78
 Baggea, 57
 Bagnisiella, 48
 Balansia, 47
 Balansiella, 47
 Balladyna, 24
 Balsamia, 97
 Balzania, 43
 Barclayella, 100
 Bargellinia, 93
 Barlaea, 88
 Bartalinia, 126
 Basiascum, 136
 Basidiella, 156
 Basidiobolus, 14
 Basidiomycetes, 102
 Basidiophora, 17
 Basisporium, 148
 Battarea, 117
 Battareopsis, 117
 Battarinia, 43
 Baumanniella, 105
 Baumiella, 34
 Beccariella, 106
 Beggiatoa, 7
 Beggiatoaceae, 7
 Belonia, 40
 Belonidium, 85
 Beloniella, 86
 Belonium, 87
 Belonopsis, 85
 Belonoscypha, 87
 Belospora, 87
 Beltrania, 151
 Beniowskaia, 159
 Berkelella, 46
 Berlesia, 36
 Bertia, 30
 Bertiella, 33
 Biatora, 76
 Biatorella, 68, 77
 Biatorellia, 68
 Biatorina, 77
 Bispora, 150
 Bivonella, 46
 Bizzozera, 25
 Bizzozeriella, 160
 Blasdalea, 51
 Elastenia, 83
 Blastodesmia, 40
 Blastomyces, 141
 Elastotrichum, 145
 Blennoria, 135
 Blennothallia, 74
 Blitrydium, 70
 Bloxamia, 135
 Blumenavia, 116
 Boerlagella, 36
 Bolacotricha, 149
 Bolbitus, 113
 Boletinus, 109
 Boletium, 108
 Boletopsis, 109
 Boletus, 109
 Bolinia, 29
 Bombardia, 29
 Bombardistrum, 33
 Bombyliospora, 77
 Bommerella, 29
 Bonia, 107
 Bonordeniella, 163

Bonplandiella, 162
 Bostrichonema, 144
 Botrytidae, 141
 Botryodiplodia, 126
 Botryosphaeria, 28
 Botryosporium, 140
 Botryotrichum, 149
 Botrytis, 142
 Bottaria, 42
 Boudiera, 92
 Boudierella, 92
 Bovilla, 38
 Bovista, 118
 Brachysporium, 152
 Brefeldiella, 51
 Bremia, 17
 Bresadolella, 44
 Bresadolia, 110
 Briardia, 63
 Briarea, 141
 Brigantiella, 54
 Briosia, 157
 Broomeia, 118
 Broomella, 46
 Brunchorstia, 132
 Bryophagus, 80
 Bryopogon, 82
 Buellia, 84
 Bulgaria, 67
 Bulgariaceae, 66
 Bulgariella, 67
 Bulgariopsis, 67
 Bullaria, 136
 Bulliardella, 56
 Burrillia, 102
 Buseella, 140
 Byssocystis, 122
 Byssonectria, 43

C

Cacosphaeria, 31
 Caenomyces, 21
 Caecoma, 99
 Calathiscus, 116
 Calcarisporium, 143
 Caldesia, 69
 Caldesiella, 108
 Calenia, 79
 Caliciaceae, 70

Caliciopsis, 71
 Calicium, 71
 Calloria, 67
 Calocera, 105
 Caloderma, 118
 Calonectria, 16
 Caloplaca, 83
 Calosphaeria, 26
 Calospora, 34
 Calostilbe, 46
 Calothricopsis, 38
 Calvatia, 118
 Calycidium, 71
 Calyptospora, 100
 Camarops, 32
 Camarosporium, 127
 Camillea, 29
 Campanella, 111
 Camposporium, 152
 Campsotrichum, 148
 Campptomyces, 19
 Campptosphaeria, 27
 Campptoum, 147
 Campylothelium, 41
 Candelaria, 81
 Candelariella, 79
 Cantharellus, 111
 Cantharomyces, 18
 Capillaria, 164
 Capnodaria, 25
 Capnodiaceae, 25
 Capnodiastrum, 124
 Capnodiella, 29
 Capnodiopsis, 25
 Capnodium, 25
 Capronia, 36
 Carestiella, 64
 Caryospora, 35
 Castoreum, 118
 Catastoma, 118
 Catenaria, 11
 Catenularia, 149
 Catharinae, 36
 Catillaria, 77
 Catinula, 133
 Cauloglossum, 117
 Celidium, 58
 Cenangella, 66
 Cenangium, 66

Cenococcaceae, 96
 Cenococcum, 96
 Cephaliphora, 145
 Cephalodochium, 160
 Cephalosporiae, 139
 Cephalosporium, 140
 Cephalotheca, 23
 Cephalothecium, 144
 Cephalotrichum, 147
 Ceracea, 105
 Ceraiomyces, 20
 Ceratocarpia, 24
 Ceratocladium, 150, 157
 Ceratomyces, 21
 Ceratophorum, 152
 Ceratosphaeria, 33
 Ceratosporium, 154
 Ceratostoma, 28
 Ceratostomella, 27
 Cercospora, 152, 154
 Cercosporiella, 145
 Cercosporidium, 152
 Ceriospora, 34
 Ceriosporiella, 33
 Cerocorticium, 106
 Cesatiella, 46
 Cetraria, 81
 Ceuthospora, 123
 Chaconia, 98
 Chaenoderma, 117
 Chaenotheca, 71
 Chaetocladiae, 14
 Chaetocladium, 14
 Chaetoconidium, 142
 Chaetoconis, 125
 Chaetodiplodia, 125
 Chaetodochis, 163
 Chaetomella, 124
 Chaetomidum, 23
 Chaetomium, 29
 Chaetomyces, 21
 Chaetopeltis, 132
 Chaetophoma, 122
 Chaetopsis, 150
 Chaetospermum, 160
 Chaetosphaeria, 35
 Chaetostroma, 162
 Chaetostromella, 163
 Chaetotheca, 22

- Chaetothyrium, 52
 Chaetozythia, 129
 Chalará, 150
 Chamonixia, 120
 Charonectria, 44
 Charrinia, 33
 Chiasmospóra, 129
 Chilonectria, 43
 Chiodectae, 60
 Chiodectum, 60
 Chiromyces, 154
 Chitonía, 114
 Chitoniella, 114
 Chitonis, 114
 Chitonomyces, 19
 Chitonospora, 34
 Chlamydobacteriaceae, 7
 Chloridium, 150
 Chlorocaulum, 78
 Chlorodothis, 50
 Chloropeltis, 75
 Chlorophyceae, 8
 Chlorosplenium, 86
 Choanophora, 14
 Choanophorae, 13
 Choeromyces, 98
 Chondromyces, 8, 155
 Choriactis, 65
 Chromosporium, 138
 Chrysobasidium, 107
 Chrysomyxa, 100
 Chrysopsora, 100
 Chrysothrix, 72
 Chrysotrichaceae, 72
 Chytridiaceae, 9
 Chytridium, 11
 Ciboria, 86
 Cicinnobella, 124
 Cicinnobolus, 122
 Ciliciopodium, 155
 Ciliella, 85
 Ciliofusarium, 163
 Ciliopora, 128
 Cintractia, 101
 Circinastrum, 124
 Circinella, 13
 Circinotrichum, 150
 Cirromyces, 150
 Citromyces, 141
 Cladosterigma, 156
 Cladobotryum, 143
 Cladochytriae, 11
 Cladochytrium, 11
 Cladoderris, 106
 Cladonia, 78
 Cladoniaceae, 78
 Cladorrhinum, 150
 Cladosphaeria, 34
 Cladosporium, 151
 Cladothrix, 7
 Cladotrichum, 151
 Clarkeinda, 114
 Clasterosporium, 151
 Clastopsora, 98
 Clathrella, 116
 Clathrogaster, 119
 Clathroporina, 40
 Clathrospora, 37
 Clathrus, 116
 Claudopus, 113
 Clavaria, 105
 Clavariaceae, 105
 Clavariopsis, 104
 Claviceps, 47
 Clavogaster, 118
 Clavularia, 155
 Cleistosoma, 43
 Cleistotheca, 25
 Clematomyces, 20
 Clethrídium, 36
 Clinconidíum, 159
 Clintoniella, 45
 Clithris, 62
 Clitocybe, 111
 Clitopilus, 113
 Clonostachys, 143
 Clypeolum, 51
 Clypeosphaeria, 34
 Coccaceae, 8
 Coccobotrys, 164
 Coccocarpia, 83
 Coccodiscus, 51
 Coccoidea, 50
 Coccoideaceae, 50
 Cocomyces, 62
 Cocconia, 61
 Coccopeziza, 63
 Coccophacidium, 62
 Coccospóra, 138
 Coccosporella, 138
 Coccosporium, 153
 Coccotrema, 40
 Coelomyces, 118
 Coelosphaeria, 26
 Coemansia, 143
 Coemansiella, 140
 Coenogonium, 72
 Coleoma, 99
 Coleopuccinia, 99
 Coleosporium, 100
 Collema, 74
 Collemataceae, 72
 Collematae, 73
 Collemopsis, 73
 Collemosporium, 73
 Colletotrichum, 135
 Collocystis, 129
 Collodendrum, 104
 Collodochium, 159
 Collonema, 127
 Collopezis, 103
 Collybia, 111
 Collyria, 105
 Colpoma, 57
 Colus, 116
 Combea, 60
 Ccmesia, 86
 Comoclatris, 57
 Completoria, 14
 Compsomyces, 20
 Confervales, 18
 Conida, 58
 Conidiascus, 93
 Conidiobolus, 14
 Coniocybe, 71
 Coniophora, 107
 Coniophorella, 107
 Conioscypha, 149
 Coniosporium, 146
 Coniothecium, 153
 Coniothyriella, 133
 Coniothyris, 133
 Coniothyrium, 124
 Conotrema, 79
 Cookella, 25, 96
 Coprinus, 114
 Coprolepa, 28
 Cora, 107

- Corallodendrum, 155
 Corallomyces, 45
 Cordana, 151
 Cordella, 146, 148
 Cordieritaceae, 92
 Cordierites, 92
 Corditubera, 118
 Cordyceps, 47
 Coremium, 155
 Coreomyces, 21
 Corethromyces, 20
 Corethropsis, 140
 Cornuella, 102
 Cornularia, 127
 Coronella, 139
 Coronophora, 26
 Corticium, 106
 Cortinarius, 114
 Corymbomyces, 143
 Coryne, 68
 Corynelia, 54
 Coryneliaceae, 54
 Coryneliella, 54
 Coryneum, 137
 Cerynogaster, 118
 Coscinaria, 47
 Cosmariospora, 161
 Costantinella, 149
 Coutinia, 28
 Couturea, 126
 Crandallia, 132
 Craterellus, 106
 Craterocolla, 104
 Crenothrix, 7
 Crepidotus, 113
 Crinula, 66
 Crocicreas, 122
 Crocynia, 72
 Cronartium, 99
 Crotonocarpia, 37
 Crucibulum, 121
 Crumenula, 66
 Cryphonectria, 45
 Cryptica, 97
 Cryptocoryneum, 151
 Cryptoderis, 33
 Cryptodiscus, 64
 Cryptomela, 136
 Cryptomyces, 61
 Cryptophallus, 115
 Cryptoporus, 109
 Cryptosphaerella, 26
 Cryptosphaeria, 26
 Cryptosphaerina, 35
 Cryptospora, 38
 Cryptosporella, 28
 Cryptosporium, 137
 Cryptostictis, 126
 Cryptothecium, 22
 Cryptothele, 72
 Cryptovalsa, 26
 Ctenomyces, 93
 Cubonia, 92
 Cucurbitaria, 37
 Cudonia, 91
 Cudoniella, 91
 Curreya, 50
 Cyanobaeis, 78
 Cyanocephalum, 45
 Cyathicula, 86
 Cyathus, 120
 Cycloconium, 150
 Cycloderma, 117
 Cyclomyces, 110
 Cycloschizum, 56
 Cyclostomella, 55
 Cylindrina, 37
 Cylindrium, 139
 Cyliandrocephalum, 140
 Cyliandrocladium, 144
 Cyliandrocolla, 160
 Cyliandrodendrum, 142
 Cylindrophora, 142
 Cylindrosporium, 137
 Cylindrotrichum, 141
 Cymatella, 112
 Cyphelium, 71
 Cyphella, 107
 Cyphina, 130
 Cystolobis, 81
 Cystophora, 148
 Cystotheca, 23
 Cystothyrium, 132
 Cystotricha, 125
 Cytodiplospora, 125
 Cytoplea, 124
 Cytospora, 123
 Cytosporella, 123
 Cytosporina, 128
 Cytosporium, 127
 Cyttaria, 94
 Cyttariaceae, 94
- D
- Dacrymycella, 159
 Dacryobolus, 121
 Dacryodochium, 160
 Dacryomitra, 105
 Dacryomyces, 104
 Dacryomycetaceae, 104
 Dacryopsis, 105
 Dactylaria, 145
 Dactylella, 145
 Dactylina, 82
 Dactylium, 145
 Dactylosporium, 153
 Daedalea, 110
 Daldinia, 29
 Dangardiella, 49
 Darbishirella, 60
 Darluca, 125
 Darwiniella, 49
 Dasyscypha, 88
 Dasypezis, 88
 Dasypthotheca, 45
 Dasyscypha, 88
 Dasyscyphae, 87
 Dasyspora, 99
 Davincia, 87
 Debaryella, 46
 Deconica, 114
 Delacourea, 37
 Delastria, 97
 Delitschia, 32
 Delitschiella, 32
 Delortia, 104
 Delpinoella, 55
 Delpontia, 64
 Dematiaceae, 146
 Dematiaceae, 162
 Dematium, 149
 Dendrodochium, 159
 Dendrogaster, 120
 Dendrographa, 60
 Dendrographium, 158
 Dendrophoma, 122
 Dendrostilbella, 155

Dendryphium, 152
Dermatea, 66
Dermateaceae, 65
Dermatiscum, 77
Dermatocarpae, 42
Dermatocarpum, 42
Desmazierella, 90
Desmidiospora, 154
Detonia, 88
Diachora, 48
Diaphanium, 158
Diaporthe, 31
Diaporthopsis, 28
Diarthonis, 58
Diatrype, 26
Diatrypella, 26
Diatrypeopsis, 26
Dibaeis, 78
Dibelonis, 86
Diblepharis, 18
Dichaeina, 56
Dichaeenopsis, 134
Dichlaena, 129
Dichodium, 73
Dichomera, 127
Dichomyces, 19
Dichoporis, 40
Dichosporium, 34
Dicoccum, 150
Dicollema, 74
Dicranidium, 161
Dicranophora, 13
Dictyobole, 115
Dictyocephalus, 119
Dictyographia, 59
Dictyonia, 68
Dictyophora, 115
Dictyorinis, 84
Dictyosporium, 153
Dictyuchus, 15
Dicyma, 147
Didymaria, 144
Didymella, 30
Didymobotryopsis, 156
Didymobotryum, 157
Didymochaete, 125
Didymocladium, 144
Didymopsis, 144
Didymospora, 100

Didymosira, 99
Didymosphaeria, 32
Didymosporium, 136
Didymostilbe, 156
Dielsiella, 56
Dietelia, 98
Digraphis, 59
Dilophia, 38
Dilophospora, 128
Dimargaris, 140
Dimerisma, 39
Dimerium, 24
Dimeromyces, 18
Dimerosporiopsis, 32
Dimerosporis, 32
Dimerosporium, 23
Dimorphomyces, 18
Dinemasporium, 133
Dioecomyces, 20
Diorchidium, 99
Diphaeis, 77
Diphaeosticta, 81
Diphanis, 77
Diphanosticta, 81
Diphloeis, 76
Diplocarpa, 88
Diplocladium, 144
Diplococcium, 151
Diplocryptis, 63
Diplocystis, 118
Diploderma, 117
Diplodia, 126
Diplodiella, 126
Diplodina, 125
Diplodiopsis, 126
Diplogramma, 59
Diplomyces, 20
Diplonaevia, 63
Diplopeltis, 132
Diplophlyctis, 11
Diplophysa, 11
Diplorhinotrichum, 144
Diploschistes, 79
Diplosporis, 27
Diplosporium, 144
Diplothea, 36
Diplotomma, 84
Diplozythia, 129
Dipodascus, 93

Diporina, 40
Dipyrenis, 40
Dipyrgis, 71
Dirina, 59
Dirinae, 59
Dirinaria, 84
Dirinastrum, 60
Discella, 134
Discina, 89
Discocolla, 161
Discocyphella, 106
Discomycetopsis, 124
Discomycopsella, 131
Discosia, 132
Discostroma, 50
Discula, 133
Dispira, 140
Distichomyces, 21
Dithelopsis, 40
Titola, 105
Ditopella, 27
Ditremis, 41
Ditylis, 71
Doassansia, 102
Doratomyces, 140
Dothichiza, 133
Dothidea, 49
Dothideaceae, 48
Dothidella, 49
Dothiopsis, 123
Dothiora, 62
Dothiorea, 123
Drepanospora, 152
Dufourea, 82
Duplicaria, 62
Durella, 70
Dyslachnum, 87
Dyslecanis, 79
Dysrhynchis, 82
Dysticta, 81
Dystictina, 82

E

Eccilia, 113
Echinobotryum, 147
Echinodontium, 108
Echinodothis, 48
Echinophallus, 115
Echinothecium, 31

- Ectinomyces, 20
 Ectolechia, 75
 Ectostroma, 164
 Ectrogella, 10
 Eidamella, 94
 Elaphomyces, 96
 Elaphomycetaceae, 96
 Elasmomyces, 116
 Eleutheromyces, 43
 Ellisiella, 149
 Elsinoe, 93
 Empusa, 14
 Enarthromyces, 19
 Encephalographa, 59
 Enchnoa, 25
 Enchnosphaeria, 34
 Endobasidium, 107
 Endobotrya, 127
 Endocarpum, 42
 Endocena, 82
 Endoconidium, 158
 Endodesmia, 160
 Endogonaceae, 96
 Endomyces, 94
 Endomycetaceae, 93
 Endophyllum, 100
 Endothia, 31
 Endoxyla, 26
 Endoxylina, 33
 Engleromyces, 44
 Englerula, 24
 Enterodictyum, 60
 Enterostigma, 61
 Entoloma, 113
 Entomophthora, 14
 Entomophthoraceae, 14
 Entomosporium, 132
 Entonaema, 44
 Entophlyctis, 10
 Entorrhiza, 101
 Entyloma, 101
 Eolichen, 39
 Eomycenella, 111
 Eoterfezia, 97
 Eoterfeziaceae, 96
 Ephebae, 74
 Ephebe, 74
 Ephebeia, 74
 Ephelina, 66
 Ephelis, 134
 Epichloe, 47
 Epiclinium, 163
 Epicoccum, 162
 Epicymatia, 30
 Epidochiopsis, 160
 Epidochium, 162
 Epigloea, 39
 Epilichen, 69
 Epochnium, 151
 Eremascus, 93
 Eremothecium, 94
 Erikssonia, 55
 Erinella, 88
 Erioderma, 83
 Eriopeziza, 86
 Eriosphaera, 118
 Eriosphaeria, 31
 Eriospora, 128
 Eriosporina, 127
 Eriothyrium, 131
 Erysibaceae, 21
 Erysibe, 22
 Erysibella, 22
 Erythrocarpum, 44
 Eubelonis, 87
 Eucantharomyces, 19
 Eucorethromyces, 20
 Eucronartium, 103
 Eucyphelis, 71
 Euhaplomyces, 19
 Eumollisiae, 84
 Eumonoecomyces, 19
 Euporthe, 31
 Eupropolis, 64
 Eurotiopsis, 128
 Eurotium, 22
 Euryachora, 48
 Eurytheca, 95
 Eutypa, 26
 Eutypella, 26
 Eustictidae, 62
 Euzodiomyces, 21
 Everhartia, 162
 Evernia, 82
 Everniopsis, 82
 Exascaceae, 93
 Exascus, 93
 Excipula, 133
 Excipulaceae, 133
 Excipularia, 134, 163
 Excipulina, 134
 Exidia, 104
 Exidiopsis, 104
 Exobasidium, 107
 Exosporina, 163
 Exosporium, 163
- ## F
- Fabraea, 86
 Farlowiella, 55
 Farriolla, 71
 Favillea, 119
 Favolus, 110
 Femsjonina, 104
 Fenestella, 37
 Feracia, 47
 Filoboletus, 109
 Fioriella, 131
 Fistulina, 109
 Fistulinella, 109
 Flaminia, 63
 Flammula, 113
 Fleischera, 48
 Floccimutinus, 115
 Fomes, 109
 Forssellia, 73
 Fracchiacea, 26
 Friesula, 106
 Fuckelia, 123
 Fuckelina, 150
 Fumago, 154
 Fungi Imperfecti, 121
 Fusariella, 151
 Fusarium, 161
 Fusella, 146
 Fusicladium, 151
 Fusicoccum, 123
 Fusicolla, 159
 Fusidium, 139
 Fusoma, 145
- ## G
- Gaillardiiella, 32
 Galactinia, 89
 Galera, 113
 Gambleola, 100
 Gamospora, 127

Gamosporella, 123
 Gasteromycetes, 115
 Gautiera, 120
 Geaster, 117
 Geasterae, 117
 Geasteropsis, 117
 Geisleria, 40
 Gelatinosporium, 128
 Geminispora, 27
 Genabea, 97
 Genea, 96
 Geoglossum, 91
 Geopora, 96
 Geopyxis, 89
 Geotrichum, 139
 Gibbera, 31
 Gibberella, 46
 Gibberidea, 35
 Gibellia, 28
 Gibellina, 32
 Gibellula, 156
 Gillettiella, 52
 Giulia, 132
 Glenospora, 148
 Gliobotrys, 140
 Gliocephalus, 139
 Gliocladium, 141
 Gloeocalyx, 67
 Gloeopeziza, 67
 Gloeoporus, 109
 Gloeosphaera, 143
 Gloeosporiella, 136
 Gloeosporium, 135
 Glomerella, 28
 Glomerularia, 138
 Gloniella, 56
 Gloniopsis, 57
 Glonium, 56
 Glossodium, 78
 Glutinium, 122, 157
 Glycophila, 139
 Glyphis, 60
 Glypholecia, 81
 Gnomonia, 30
 Gnomoniella, 27
 Gnomoniopsis, 33
 Godronia, 66
 Godroniella, 133
 Gomphidius, 115

Gomphyllus, 78
 Gonapodya, 18
 Gonatobotrys, 143
 Gonatobotrytae, 143
 Gonatobotryum, 147
 Gonatorrhodiella, 143
 Gonatorrhodum, 143
 Gongromeriza, 147
 Gongylia, 40
 Goniosporium, 147
 Gonohymenia, 73
 Gonotheceis, 75
 Gonytrichum, 150
 Gorgoniceps, 87
 Grammothele, 108
 Grandinia, 108
 Grandiniella, 107
 Granularia, 159
 Graphidaceae, 58
 Graphidiae, 58
 Graphina, 59
 Graphiothecium, 157
 Graphis, 59
 Graphium, 157
 Graphyllum, 57
 Guelichia, 160
 Guepinia, 105
 Guignardiella, 27
 Guillermondia, 95
 Gyalecta, 80
 Gyalectae, 80
 Gymnascaceae, 93
 Gymnascales, 93
 Gymnascus, 93
 Gymnoconia, 100
 Gymnoderma, 78
 Gymnodoichium, 160
 Gymnoglossum, 118
 Gymnographa, 58
 Gymnomycetes, 120
 Gymnosporangium, 100
 Gyrocephalus, 104
 Gyrocera, 147
 Gyrocratera, 97
 Gyrodon, 109
 Gyromitra, 91
 Gyrophora, 77
 Gyrophorae, 77
 Gyrostomum, 80

H

Habrostictis, 63
 Hadrortrichum, 148
 Haematomma, 79
 Haematomyces, 67
 Haematomyxa, 68
 Hainesia, 135
 Halobysus, 131
 Hapalophragmium, 101
 Haplaria, 141
 Haplariopsis, 144
 Haplobasidium, 147
 Haplocybe, 91
 Haplographium, 149
 Haplomyces, 19
 Haplopyrenula, 41
 Haplosporella, 124
 Haplosporium, 28
 Haplotrichum, 140
 Hariotia, 30
 Harknessia, 124
 Harknessiella, 95
 Harpidium, 79
 Harpocephalum, 157
 Harpochytriae, 11
 Harpochytrium, 11
 Harpographium, 157
 Hartigiella, 141
 Harziella, 143
 Hassea, 39
 Hebeloma, 113
 Helicobasidium, 107
 Helicocephalum, 149
 Helicogloea, 103
 Helicoma, 154
 Helicomycetes, 146
 Helicopsis, 154
 Helicosporium, 154
 Helicostilbe, 156
 Helicotrichum, 149
 Helicoum, 146
 Heliomyces, 112
 Heliscus, 161
 Helminthascus, 47
 Helminthocarpum, 59
 Helminthophana, 20
 Helminthosphaeria, 28
 Helminthosporium, 152
 Helostroma, 156

- Helotiaceae, 86
 Helotiae, 86
 Helotium, 86
 Helvella, 91
 Helvellaceae, 90
 Helvellae, 91
 Hemiglossum, 91
 Hemihysteriaceae, 54
 Hemileia, 98
 Hemileiosis, 101
 Hendersonia, 126
 Hendersonula, 127
 Henningsomyces, 32
 Henningsinia, 29
 Henriquesia, 55
 Heppia, 75
 Heppiae, 75
 Heptameria, 34
 Hercospora, 31
 Hericium, 108
 Herpocradiella, 13
 Herpomyces, 20
 Herpothrix, 35
 Herpotrichia, 34
 Heterobasidium, 107
 Heterobotrys, 146
 Heterocarpum, 42
 Heterocephalum, 155
 Heterochaete, 104
 Heterodea, 81
 Heteromyces, 78
 Heteronectria, 43
 Heteropatella, 133
 Heterosphaeria, 65
 Heterosporium, 152
 Heufleria, 42
 Hexagonia, 110
 Heydenia, 157
 Hiatula, 111
 Himantia, 164
 Hippoperdon, 118
 Hirneola, 103
 Hirneolina, 104
 Hirsutella, 105
 Hirudinaria, 154
 Hobsonia, 162
 Hoehneliella, 157
 Holcomyces, 132
 Holocoenis, 72
 Holocyphis, 71
 Holothelis, 40
 Holstiella, 34
 Holwaya, 68
 Homopsella, 74
 Homostegia, 49
 Homothecium, 73
 Hormiactella, 149
 Hormiactis, 144
 Hormiscium, 147
 Hormodendrum, 149
 Hormodochis, 163
 Hormosperma, 33
 Hormyllum, 135
 Humaria, 88
 Hussey, 117
 Hyaloceras, 137
 Hyaloderma, 25
 Hyalodothis, 48
 Hyalopsora, 98
 Hyalopus, 140
 Hyaloria, 104
 Hyalostilbae, 154
 Hyalothyridium, 127
 Hyalothyris, 127
 Hydnaceae, 107
 Hydnangium, 120
 Hydnobolites, 97
 Hydnochaete, 108
 Hydnocystis, 96
 Hydnofomes, 108
 Hydnohypsa, 108
 Hydnotrya, 97
 Hydnum, 108
 Hydraeomyces, 19
 Hydrothyria, 83
 Hygrophorus, 111
 Hymenella, 159
 Hymenogaster, 120
 Hymenogastraceae, 119
 Hymenogramme, 112
 Hymenomycetes, 102
 Hymenopsis, 162
 Hymenoscypha, 87
 Hymenula, 159
 Hyperphyscia, 84
 Hypha, 164
 Hyphaster, 163
 Hyphoderma, 141
 Hypholoma, 114
 Hyphomyceteae, 138
 Hyphoscypha, 88
 Hyphostereum, 159
 Hypocenia, 124
 Hypochnus, 107
 Hypocopra, 28
 Hypocrea, 45
 Hypocreaceae, 42
 Hypocrella, 48
 Hypocreodendrum, 129
 Hypocreopsis, 45
 Hypoderma, 57
 Hypodermella, 55
 Hypodermium, 135
 Hypodermopsis, 57
 Hypolyssus, 106
 Hypomyces, 45
 Hyponectria, 43
 Hypospila, 33
 Hypospilina, 30
 Hypoxyloopsis, 49
 Hypoxylum, 30
 Hysterangium, 120
 Hysteriaceae, 55
 Hysteriales, 54
 Hysterium, 57
 Hysterographium, 57
 Hysteromyxa, 130
 Hysteropatella, 69
 Hysteropsis, 57
 Hysterostomella, 55

I

- Icmadophila, 79
 Idiomycetes, 20
 Illosporium, 160
 Ingaderia, 60
 Inocybe, 113
 Inzengaea, 27
 Iotidea, 89
 Iridionia, 63
 Irpex, 108
 Isaria, 156
 Isariopsis, 157
 Isothea, 37
 Itajahya, 115
 Ithyphallus, 115

J

Janseella, 64
 Jansia, 115
 Jenmania, 73
 Johansonia, 69
 Jola, 103
 Jonaspis, 80
 Julella, 36

K

Kabatia, 131
 Kalchbrennera, 115
 Kalmusia, 35
 Karschia, 69
 Karstenia, 64
 Karstenula, 36
 Keithia, 62
 Kellermannia, 125, 126
 Kmetia, 159
 Kneiffia, 106
 Koerberia, 73
 Konradia, 48
 Kretschmaria, 29
 Kullhemia, 48
 Kusanoa, 95
 Kusanobotrys, 24

L

Laaseomyces, 23
 Laboulbenia, 21
 Laboulbeniaceae, 18
 Laboulbeniales, 18
 Labrella, 131
 Labridium, 132
 Laccocephalum, 109
 Lachnea, 90
 Lachnella, 88
 Lachnellula, 87
 Lachnocladium, 105
 Lachnodoichium, 160
 Lachnum, 87
 Lactariopsis, 111
 Lactarius, 111
 Laestadia, 27
 Lagenidiæ, 16
 Lagenidium, 16
 Lagerheimia, 69
 Lahmia, 70
 Lambro, 45

Lamia, 14
 Lamyella, 123
 Langloisula, 142
 Lanopila, 118
 Lanzia, 87
 Laquearia, 65
 Laschia, 109
 Lasiobolus, 92
 Lasiobotrys, 23
 Lasioderma, 155
 Lasiodiplodia, 126
 Lasiosphaeria, 33
 Lasiosphaeris, 35
 Lasiostrictis, 64
 Lasmenia, 131
 Laurera, 42
 Lauterbachella, 64
 Lecanactidae, 76
 Lecanactis, 76
 Lecania, 79
 Lecanidion, 70
 Lecaniella, 75
 Lecanora, 79
 Lecanorae, 78
 Lecidea, 76
 Lecideaceae, 76
 Lecideae, 76
 Lecideopsis, 58
 Leciographa, 70
 Lecithium, 46
 Lecoglyphis, 70
 Leciophysma, 73
 Leiosepium, 141
 Lemalis, 129
 Lembosia, 56
 Lembosiella, 55
 Lemmopsis, 73
 Lemonniera, 146
 Lentinus, 112
 Lentodiopsis, 112
 Lentodium, 112
 Lentomita, 30
 Lenzites, 112
 Leotia, 91
 Leotiella, 91
 Lepidocollema, 83
 Lepiota, 111
 Lepolichen, 42
 Lepraria, 164

Leprocollema, 73
 Leptinia, 100
 Leptodendrisium, 74
 Leptogidium, 74
 Leptogiopsis, 74
 Leptogium, 74
 Leptolegnia, 15
 Leptomitæ, 15
 Leptomitæ, 15
 Leptomitæ, 15
 Leptonia, 113
 Leptophyma, 95
 Leptorhaphis, 41
 Leptosphaeria, 34
 Leptosphaeropsis, 34
 Leptosphaerulina, 36
 Leptospora, 38
 Leptostroma, 131
 Leptostromataceae, 130
 Leptostromella, 133
 Leptothyrella, 131
 Leptothyrium, 131
 Leptotrema, 80
 Leptotrichum, 161
 Letendrea, 46
 Letharia, 82
 Leucangium, 97
 Leucocrea, 47
 Leucogaster, 120
 Leucopezis, 90
 Leucophleps, 120
 Levieuxia, 124
 Libertella, 138
 Libertiella, 129
 Lichenopsis, 126
 Lichenosticta, 121
 Lichina, 74
 Lichina, 74
 Lichinella, 74
 Lichinodium, 74
 Licopolia, 32
 Lilliputia, 97
 Limacina, 24
 Limnaeomyces, 19
 Lindauella, 63
 Linhartia, 85
 Linospora, 37
 Lisea, 45
 Lisiella, 43
 Listeromyces, 163

- Lithoecis, 39
 Lithographa, 59
 Lithothelium, 42
 Lituaria, 161
 Lizonia, 31
 Lizoniella, 30
 Lloydella, 107
 Lobaria, 81
 Lobarina, 83
 Locellina, 113
 Lopadiopsis, 75
 Lopadium, 77
 Lopharia, 108
 Lophidiopsis, 54
 Lophidium, 54
 Lophiella, 53
 Lophionema, 54
 Lophiosphaera, 53
 Lophiostoma, 54
 Lophiostomataceae, 53
 Lophiotrema, 54
 Lophiotricha, 53
 Lophium, 57
 Lophodermium, 57
 Lycoperdaceae, 116
 Lycoperdæ, 118
 Lycoperdales, 115
 Lycoperdon, 118
 Lycoperdopsis, 118
 Lysurus, 116
- M
- Macowanites, 120
 Macrobatis, 126
 Macrodiplodia, 125
 Macrophoma, 122
 Macropodia, 89, 90
 Macrosporium, 153
 Macrostilbum, 155
 Magnusia, 23
 Malbranchea, 139
 Malmeomyces, 46
 Manginia, 123
 Marasmiopsis, 112
 Marasmius, 112
 Marchaliella, 24
 Maronea, 80
 Marsonia, 136
 Martellia, 120
 Martensella, 142
 Martindalia, 155
 Martinella, 129
 Massalongia, 83
 Massalongiella, 25
 Massaria, 34
 Massariella, 31
 Massarina, 33
 Massarinula, 30
 Massariovalsa, 32
 Maseea, 87
 Maseella, 99
 Massospora, 138
 Mastigosporium, 145
 Mastodia, 42
 Mastomyces, 126
 Mattirolia, 47
 Matruchotia, 107
 Maurodothis, 49
 Maurya, 38
 Mazosia, 61
 Mazzantia, 48
 Megalonectria, 46
 Megalospora, 76
 Melachroia, 89
 Melampsora, 98
 Melampsorella, 98
 Melampyidium, 76
 Melanconiaceae, 135
 Melanconiales, 135
 Melanconiella, 32
 Melanconiopsis, 124
 Melanconis, 31
 Melanconium, 136
 Melanogaster, 120
 Melanomma, 35
 Melahops, 34
 Melanopsamma, 31
 Melanopsichium, 102
 Melanospora, 44
 Melanostroma, 135
 Melanotaenium, 101
 Melanotheca, 42
 Melasmia, 131
 Melaspilea, 59, 69
 Melchiora, 31
 Meliola, 24
 Meliopsis, 23
 Melittiosporium, 64
 Melittosporiopsis, 85
 Melittosporis, 85
 Melogramma, 35
 Melomastia, 33
 Melophia, 132
 Mendogia, 57
 Menispora, 150
 Merarthonis, 58
 Merodontis, 87
 Merodothidis, 128
 Merophora, 77
 Meroplacis, 84
 Merorinis, 84
 Merostictina, 82
 Merostictis, 64
 Merulius, 110
 Mesneria, 28
 Mesobotrys, 150
 Mesophellia, 117
 Metadothella, 62
 Metanectria, 45
 Metasphaeria, 33
 Metraria, 112
 Michenera, 106
 Micrascus, 28
 Microcera, 161
 Micrococcus, 8
 Microcylus, 49
 Microdiplodia, 125
 Microglæna, 40
 Microglossum, 91
 Micrographa, 59
 Micronectria, 47
 Micropeltis, 52
 Micropera, 128
 Microphiale, 80
 Microphyma, 95
 Microspatha, 155
 Microsphaera, 22
 Microspira, 8
 Microstelium, 58
 Microstroma, 138
 Microthelia, 41
 Microtheliopsis, 41
 Microthyriaceae, 51
 Microthyriae, 51
 Microthyrium, 52
 Micula, 128
 Midotiopsis, 65

Midotis, 65
 Milesia, 100
 Milowia, 144
 Minksia, 60
 Misgomyces, 21
 Mitromyces, 117
 Mitrula, 91
 Mitruliopsis, 91
 Moelleriella, 43
 Moellerodiscus, 91
 Mollieriella, 95
 Mollisia, 84
 Mollisiaceae, 84
 Mollisiella, 84
 Monacrosporium, 145
 Monascus, 23
 Monilia, 139
 Moniliaceae, 138
 Moniliales, 138
 Monoblepharidaceae, 18
 Monoblepharis, 18
 Monochaetia, 137
 Monoecomyces, 19
 Monographus, 48
 Monopodium, 141
 Monosporidium, 98
 Monosporium, 142
 Monotospora, 148
 Montagnella, 49
 Montagnites, 115
 Montagnula, 37
 Morchella, 91
 Morenoella, 55
 Morinia, 137
 Moriola, 39
 Moriola, 39
 Mortierella, 13
 Mortierellae, 13
 Moschomyces, 20
 Moutoniella, 62
 Mucedinae, 158
 Mucedineae, 138
 Mucor, 13
 Mucoraceae, 12
 Mucorae, 12
 Mucronella, 108
 Mucronoporus, 109
 Mucrosporium, 145

Müllerella, 28
 Munkia, 130
 Munkiella, 49
 Muricularia, 129
 Mutinus, 115
 Myceliophthora, 138
 Mycena, 111
 Mycobacidia, 70
 Mycobilimbica, 69
 Mycoblastus, 76
 Mycocalicium, 71
 Mycocitrus, 45
 Mycodendrum, 109
 Mycogala, 122
 Mycogone, 144
 Mycolecis, 70
 Mycomalus, 48
 Mycoporaceae, 50
 Mycoporellum, 50
 Mycoporis, 78
 Mycoporum, 50
 Mycorhynchus, 130
 Mycosphaerella, 30
 Mycosyrinx, 102
 Myiocoprura, 51
 Mylittopsis, 103
 Myriadoporus, 110
 Myriangiaceae, 95
 Myriangella, 95
 Myriangium, 95
 Myridium, 67
 Myrioblepharis, 18
 Myriogenospora, 50
 Myriolecis, 79
 Myriophysis, 162
 Myrmaecium, 31
 Myrmecocystis, 97
 Myrothecium, 162
 Mystrosporium, 153
 Mytilidium, 57
 Myxobacter, 8
 Myxobacteriaceae, 8
 Myxobactrales, 8
 Myxococcus, 8
 Myxodictyum, 79
 Myxomycidium, 105
 Myxonema, 160
 Myxormia, 135

Myxosporella, 135
 Myxosporium, 135
 Myxotrichum, 149
 Myzocyttum, 16

N

Naegeliella, 16
 Naemacyclus, 64
 Naematelia, 104
 Naemosphaera, 124
 Naemospora, 137
 Naevia, 63
 Naeviella, 63
 Napicladium, 152
 Naucoria, 113
 Necator, 159
 Nectria, 45
 Nectriella, 43
 Nectrioidaceae, 128
 Negeriella, 158
 Nematogonium, 143
 Nematosporangium, 16
 Neoarcangelia, 26
 Neobarclaya, 136
 Neocosmospora, 44
 Neomichelina, 152
 Neopectia, 32
 Neoravenelia, 101
 Neorehmania, 23
 Neoskofitzia, 45
 Neottiopezis, 90
 Neottiospora, 122
 Neovossia, 101
 Nephroma, 76
 Nephromopsis, 81
 Nephromium, 75
 Nesolechia, 69
 Nidula, 121
 Nidularia, 120
 Nidulariaceae, 120
 Nigrospora, 148
 Niorma, 84
 Niptera, 85
 Nitschkea, 26
 Nocardia, 7
 Nolanea, 113
 Nomuraea, 143
 Normandina, 42

Nostocotheca, 94
 Nothopatella, 124
 Nothostroma, 50
 Nowakowskia, 10
 Nowakowskiella, 11
 Nummularia, 30
 Nyctalis, 111

O

Obelidium, 11
 Ocellaria, 63
 Ocellis, 80
 Ocellularia, 80
 Ochrolechia, 79
 Octaviana, 120
 Odontia, 108
 Odontotrema, 65
 Odontura, 65
 Oedemium, 148
 Oedocephalum, 139
 Oedomyces, 101
 Ohleria, 35
 Ohleriella, 35
 Oidiopsis, 139
 Oidium, 139
 Oleina, 93
 Ollula, 130
 Olpidiae, 9
 Olpidiopsis, 10
 Olpidium, 10
 Olpitrichum, 141
 Ombrophila, 67
 Omphalia, 111
 Oncopodium, 153
 Oncospora, 134
 Onygena, 96
 Onygenaceae, 96
 Oochytriae, 11
 Oomyces, 47
 Oospora, 139
 Opegrapha, 59
 Opegraphella, 59
 Ophiobolus, 37
 Ophioceras, 38
 Ophiochaete, 38
 Ophiocladium, 138
 Ophiodictyum, 36
 Ophiodothis, 50

Ophiogloea, 68
 Ophiognomonina, 37
 Ophiomassaria, 37
 Ophiomeliola, 25
 Ophionectria, 47
 Ophiopeltis, 52
 Ophiotrichum, 152
 Orbicula, 23
 Orbilia, 67
 Oropogon, 82
 Orphniospora, 76
 Oscarbrefeldia, 94
 Ostreium, 57
 Ostropa, 57, 65
 Ostropae, 65
 Otidea, 88
 Otidella, 88
 Otthia, 32
 Otthiella, 31
 Oudemansiella, 112
 Ovularia, 142
 Ovulariopsis, 142
 Oxydothis, 50
 Ozonium, 164

P

Pachybasium, 143
 Pachyphiale, 80
 Pachyphloeus, 97
 Pactilia, 158
 Paepalopsis, 139
 Pampolysporium, 23
 Panaeolus, 114
 Pannaria, 83
 Pannariae, 83
 Panus, 112
 Papulospora, 140
 Paranectria, 46
 Paraspora, 145
 Parathalle, 70
 Paratheliae, 41
 Parathelium, 41
 Parmelia, 81
 Parmeliaceae, 78
 Parmeliae, 81
 Parmeliella, 83
 Parmeliopsis, 81
 Parmenteria, 42

Parmularia, 55
 Parmulariella, 55
 Parodiella, 24, 32
 Paryphedria, 67
 Passalora, 151
 Passeriniella, 34
 Passerinula, 45
 Patellaria, 70
 Patellariaceae, 68
 Patellea, 69
 Patellina, 130, 158
 Patellinae, 130
 Patinella, 69
 Patouillardia, 159
 Patouillardiella, 161
 Paulia, 73
 Paurocotylis, 119
 Paxillus, 114
 Pazschkea, 85
 Pazschkella, 125
 Peccania, 73
 Peckia, 123
 Peckiella, 40
 Pedilospora, 146
 Pedisordaria, 29
 Pellicularia, 142
 Pellioniella, 125
 Pelodiscus, 90
 Peloronectria, 46
 Peltidium, 91
 Peltigera, 75
 Peltigeromyces, 88
 Peltophora, 75
 Peltophoraceae, 75
 Peltophorae, 75
 Peltophoromyces, 88
 Peltosphaeria, 36
 Peltostroma, 131
 Pemphidium, 51
 Penicilliopsis, 44
 Penicillium, 141
 Peniophora, 107
 Pentagonella, 60
 Penzigia, 29
 Perforaria, 79
 Peribotryum, 156
 Pericladium, 99
 Periconia, 147

- Periconiella, 147
 Peridermium, 99
 Periola, 160
 Perisporiaceae, 22
 Perisporina, 24
 Perisporiopsis, 24
 Perisporium, 24
 Peronospora, 17
 Peroneutypa, 26
 Peroneutypella, 26
 Peronosporaceae, 17
 Peronosporae, 17
 Perisporiopsis, 24
 Perrotia, 88
 Pertusaria, 79
 Pertusariae, 79
 Pestalozzia, 137
 Pestalozziella, 135
 Pestalozzina, 136
 Petasodes, 133
 Petractis, 74
 Peyritschella, 19
 Peziza, 89
 Pezizaceae, 88
 Pezizae, 88
 Pezizales, 61
 Pezizella, 86
 Pezoloma, 86
 Phacidiaceae, 61
 Phacidium, 61
 Phacopsis, 58
 Phacopsora, 98
 Phaeangella, 66
 Phaeangium, 66, 97
 Phaeoclavulina, 105
 Phaeoconis, 148
 Phaeocreopsis, 46
 Phaeodiscula, 133
 Phaeodothis, 49
 Phaeoglaena, 40
 Phaeographina, 59
 Phaeographis, 59
 Phaeohygrocybe, 114
 Phaeolimacium, 114
 Phaeomacropus, 90
 Phaeomeris, 39
 Phaeonectria, 46
 Phaeopeltis, 52
 Phaeopeltosphaeria, 36
 Phaeopezia, 89
 Phaeophacidium, 61
 Phaeopterula, 105
 Phaeoradulum, 108
 Phaeosaccardinula, 52
 Phaeoscutella, 52
 Phaeosphaerella, 31
 Phaeospora, 34
 Phaeosporis, 39
 Phaeostilbae, 156
 Phaeothrombis, 40
 Phaeotrema, 80
 Phallaceae, 115
 Phallogaster, 116
 Phalodictyum, 77
 Phalomeris, 39
 Phalostauris, 39
 Phalothrix, 88
 Phanosticta, 81
 Phanotylum, 80
 Pharcidia, 30
 Phellorina, 119
 Phialea, 86
 Phillipsiella, 95
 Philocopra, 28
 Phlebia, 108
 Phlebophora, 103
 Phleboscyphus, 89
 Phleospora, 128
 Phloeconis, 164
 Phloeopeccania, 73
 Phlyctella, 79
 Phlyctidia, 79
 Phlyctaena, 128
 Phlyctis, 79
 Phlyctochytrium, 11
 Pholidota, 113
 Pholiotella, 113
 Phoma, 122
 Phomataceae, 121
 Phomatales, 121
 Phomatospora, 27
 Phomopsis, 121
 Phragmidiothrix, 7
 Phragmidium, 100
 Phragmographium, 56
 Phragmonaevia, 64
 Phragmopeltis, 132
 Phragmopyxine, 84
 Phragmopyxis, 100
 Phragmostele, 100
 Phragmothele, 39
 Phragmotrichum, 137
 Phycascus, 89
 Phycodiscis, 83
 Phycomyces, 13
 Phyllachora, 48
 Phyllactinia, 22
 Phylliscidium, 72
 Phylliscum, 72
 Phyllobathelium, 41
 Phyllophthalaria, 80
 Phylloporina, 41
 Phylloporis, 41
 Phyllopsora, 77
 Phyllopsorae, 77
 Phyllosticta, 121
 Phymatotrichum, 142
 Phymatosphaeria, 95
 Phymatosphaeriaceae, 95
 Physalacria, 105
 Physalospora, 27
 Physcia, 84
 Physciaceae, 83
 Physcidia, 81
 Physma, 73
 Physoderma, 11
 Physmatomyces, 67
 Physospora, 141
 Phytrophthora, 17
 Picoa, 97
 Piersonia, 97
 Piggotia, 131
 Pilacre, 155
 Pilacrella, 103
 Pilaira, 13
 Pilgeriella, 27
 Pilidium, 134
 Pilobolae, 13
 Pilobolus, 13
 Pilocarpum, 76
 Pilocratera, 90
 Pilophorum, 78
 Pilosace, 114
 Pimina, 150

- Pionnotes, 161
 Piptocephalis, 14
 Piptostoma, 51
 Piptostomum, 122
 Pirella, 13
 Piricularia, 145
 Pirobasidium, 155
 Pirostoma, 131
 Pirottaea, 85
 Pisomyxa, 22
 Pistillaria, 106
 Pithomyces, 161
 Pitya, 88
 Placidiopsis, 42
 Placographa, 68
 Placolecanis, 79
 Placosphaerella, 125
 Placosphaeria, 123
 Placothelium, 39
 Placynthium, 83
 Plagiotrema, 41
 Planococcus, 8
 Planosarcina, 8
 Plasmopara, 17
 Platygloea, 103
 Platysticta, 64
 Platystomum, 54
 Plearthonis, 58
 Plectania, 90
 Plectothrix, 142
 Plenodomus, 122
 Pleochaete, 22
 Pleococcum, 133
 Pleoconis, 73
 Pleochroma, 80
 Pleodothis, 49
 Pleogibberella, 46
 Pleoglonis, 56
 Pleolecis, 76
 Pleolpidium, 10
 Pleomassaria, 36
 Pleomeliola, 25
 Pleonectria, 46
 Pleophragmia, 37
 Pleopyrenis, 72
 Pleoravenelia, 101
 Pleorinis, 84
 Pleosphaeria, 37
 Pleosphaerulina, 36
 Pleospilis, 69
 Pleospora, 36
 Pleosporopsis, 29
 Pleostictis, 64
 Pleotrachelus, 10
 Pleurascus, 23
 Pleurocybe, 72
 Pleurotus, 111
 Pleurostoma, 26
 Pleurothelium, 41
 Pleurotrema, 41
 Plicaria, 89
 Plicariella, 89
 Ploettnera, 63
 Plowrightia, 49
 Pluteolus, 113
 Pluteus, 112
 Pocillum, 87
 Pocosphaeria, 34
 Podaleuris, 89
 Podaxae, 116
 Podaxon, 117
 Podocapsa, 94
 Podocapsium, 94
 Podochytrium, 11
 Podocrea, 45
 Podoloma, 117
 Podosordaria, 29
 Podosphaera, 22
 Podosporella, 158
 Podosporium, 158
 Podostictina, 82
 Podostroma, 44
 Poecilosporium, 102
 Polyangium, 8
 Polyascomyces, 19
 Polyblastia, 39
 Polyblastiopsis, 41
 Polycephalum, 155
 Polychidium, 74
 Polydesmus, 152
 Polygaster, 119
 Polynema, 133
 Polyphagus, 12
 Polyplocium, 116
 Polyporaceae, 108
 Polyporus, 109
 Polyrrhina, 11
 Polysaccopsis, 102
 Polysaccum, 119
 Polyscytalum, 139
 Polystictis, 109
 Polystigma, 43
 Polystigmina, 130
 Polystomella, 51
 Polystroma, 80
 Polythelis, 41
 Polythrincium, 151
 Polytrichia, 27
 Poria, 109
 Porina, 40
 Porocyphus, 74
 Poronia, 29
 Poropeltis, 131
 Poroptyche, 110
 Porotheium, 110
 Pragmopara, 70
 Prillieuxia, 107
 Prismaria, 146
 Pritzeliiella, 155
 Prophytroma, 149
 Propolidium, 64
 Propolina, 63
 Propolis, 63
 Prosthemiella, 136
 Prosthemium, 127
 Protococcales, 9
 Protoglossum, 119
 Protomerulius, 104
 Protomyces, 94
 Protomycetaceae, 93
 Protostegia, 134
 Protoventuria, 32
 Protubera, 116
 Psammia, 136
 Psathyra, 114
 Psathyrella, 114
 Pseudacolium, 71
 Pseudobeltrania, 151
 Pseudocenangium, 134
 Pseudodiplodia, 129
 Pseudogenea, 96
 Pseudographis, 56, 62
 Pseudohydnotria, 96
 Pseudolizonia, 31

Pseudomelasmia, 131
Pseudomeliola, 38
Pseudomonas, 8
Pseudopatella, 134
Pseudopeziza, 85
Pseudophacidium, 61
Pseudoplectania, 90
Pseudopyrenula, 41
Pseudostictis, 130
Pseudotryblidium, 70
Pseudotrype, 43
Pseudothia, 32
Pseudovalsa, 35
Pseudozythia, 129
Psilocybe, 114
Psilopezia, 91
Psilospora, 133
Psilothecium, 69
Psora, 77
Psorella, 77
Psoroglaena, 42
Psorographis, 59
Psoroma, 79
Psoromaria, 77
Psorotheciella, 85
Psorotheciopsis, 85
Psorotichia, 73
Pteridiospora, 31
Pterophyllus, 112
Pterula, 105
Pterygiopsis, 74
Pterygium, 74
Ptychographa, 59
Puccinia, 100
Pucciniastrum, 100
Pucciniopsis, 163
Pucciniosira, 99
Pucciniospora, 125
Pucciniostele, 100
Puiggariella, 45
Pulparia, 67
Pulveraria, 164
Pustularia, 89
Puttemansia, 90
Pycnochytrium, 10
Pycnographa, 61
Pycnostroma, 130
Pyrenastrum, 42

Pyrenidiace, 38
Pyrenidium, 38
Pyrenochaeta, 123
Pyrenocollema, 39
Pyrenopeziza, 85
Pyrenopezizae, 85
Pyrenophora, 37
Pyrenopsidae, 72
Pyrenopsidium, 72
Pyrenopsis, 72
Pyrenothamnia, 42
Pyrenotheca, 50
Pyrenotrichum, 122
Pyrenula, 40
Pyrenulace, 40
Pyrenyllum, 41
Pyrigidium, 71
Pyrigillus, 71
Pyronema, 89
Pyronemella, 90
Pythiae, 16
Pythiopsis, 15
Pythium, 16
Pyxine, 84

Q

Quaternaria, 26
Queletia, 117

R

Rabenhorstia, 123
Racodium, 72
Radulum, 108
Ramalina, 82
Ramonia, 80
Ramularia, 145
Ramulaspora, 142
Ravenelia, 101
Ravenelula, 69
Rebentischia, 34
Reessia, 9
Rehmiella, 30
Reinkella, 60
Resticularia, 17
Rhabdospora, 127
Rhachomyces, 20
Rhacodium, 164
Rhacophyllus, 112

Rhadinomyces, 20
Rhagadolobium, 61
Rhamphoria, 36
Rhamphospora, 101
Rhinocladium, 148
Rhinotrichum, 141
Rhipidium, 16
Rhipidonema, 107
Rhizidiace, 10
Rhizidiomyces, 11
Rhizidium, 11
Rhizina, 91
Rhizinae, 91
Rhizocarpum, 77
Rhizoctonia, 164
Rhizomorpha, 164
Rhizomyces, 20
Rhizomyxa, 16
Rhizophidium, 10
Rhizophlyctis, 10
Rhizopogon, 120
Rhizopus, 13
Rhombostilbella, 155
Rhopalidium, 136
Rhopalomycetes, 139
Rhopoglyphus, 49
Rhymnocarpus, 68
Rhynchodiplodia, 125
Rhynchomelas, 44
Rhynchomeliola, 32
Rhynchomyces, 152
Rhynchonectria, 44
Rhynchophoma, 125
Rhynchospaeria, 34
Rhynchosporium, 144
Rhynchostoma, 32
Rhyarobius, 92
Rhytidhysterium, 56
Rhytidopeziza, 70
Rhytisma, 62
Riccoa, 158
Richonia, 24
Rickia, 21
Rickiella, 95
Riessia, 158
Rimbachia, 111
Rinia, 27
Rinodina, 84

Robergea, 58, 65
 Robillardia, 125
 Roccella, 60
 Roccellae, 60
 Roccellaria, 60
 Roccellina, 60
 Roccellographa, 60
 Rodwaya, 109
 Roesleria, 71
 Roestelia, 99
 Rosellinia, 29
 Rosenscheldia, 49
 Rostrella, 27
 Rostrupia, 100
 Rotaea, 145
 Roumegueriella, 128
 Roussoella, 49
 Rozella, 10
 Ruhlandiella, 97
 Russula, 111
 Rutstroemia, 87

S

Saccardaea, 156
 Saccardia, 22
 Saccardinula, 52
 Saccardoella, 33
 Saccardomyces, 25
 Saccharomyces, 94
 Saccharomycetaceae, 94
 Saccoblastia, 103
 Saccobolus, 92
 Saccopyrenia, 39
 Sacidium, 131
 Sagiolechia, 80
 Samarospora, 22
 Santiella, 126
 Saprolegnia, 15
 Saprolegniaceae, 15
 Saprolegniae, 15
 Sarcina, 8
 Sarcinella, 154
 Sarcinodochium, 161
 Sarcographa, 60
 Sarcographina, 60
 Sarcomyces, 68
 Sarcopodium, 149
 Sarcoscypha, 90

Sarcosoma, 67
 Scaphidium, 134
 Sceptromyces, 143
 Schenckia, 24
 Schinzinia, 113
 Schismatomma, 76
 Schizacrosporum, 58
 Schizomycetes, 7
 Schizonella, 102
 Schizopelte, 60
 Schizophyllum, 112
 Schizosaccharomyces, 94
 Schizospora, 100
 Schizostoma, 53
 Schizothyrella, 134
 Schizothyrium, 55, 61
 Schizotrichum, 163
 Schizoxylum, 64
 Schneepia, 55
 Schroeteria, 102
 Schroeteriaaster, 98
 Schulzeria, 111
 Schweinitziella, 48, 50
 Sciodothis, 50
 Scirrha, 49
 Scirrhia, 48
 Sclerococcum, 163
 Sclerodepsis, 110
 Scleroderma, 118
 Sclerodermatae, 118
 Scleroderris, 65
 Sclerodiscus, 162
 Sclerogaster, 120
 Sclerographium, 158
 Sclerophytum, 60
 Scleroplea, 37
 Sclerospora, 17
 Sclerotinia, 86
 Sclerotiopsis, 122
 Sclerotium, 164
 Scoleactis, 76
 Scoleciocarpus, 119
 Scolecopeltis, 52
 Scolecosporis, 77
 Scolecosporium, 137
 Scolecotrichum, 151
 Scoliciosporum, 77
 Scopinella, 44

Scopularia, 150
 Scorias, 25
 Scoriomyces, 160
 Scortechinia, 28
 Scutellinia, 90
 Scutellinae, 89
 Scutellum, 52
 Scutula, 69
 Scutularia, 70
 Scytopezis, 65
 Sebacina, 104
 Secotium, 116
 Seismosarca, 105
 Selenotila, 138
 Selinia, 43
 Sepedonium, 142
 Septobasidium, 106
 Septocylindrium, 144
 Septodhideopsis, 128
 Septogloeum, 136
 Septomyxa, 136
 Septonema, 152
 Septorella, 127
 Septoria, 127
 Septosporiella, 128
 Septosporium, 153
 Septotrullula, 137
 Sepultaria, 90
 Seuratia, 25
 Seynesia, 52
 Seynesiopsis, 132
 Shiraia, 47
 Sigmoidomyces, 139
 Sillia, 38
 Simblum, 116
 Simonyella, 60
 Siphula, 82
 Sirentyloma, 101
 Siridiella, 137
 Siridium, 137
 Sirobasidium, 104
 Sirococcus, 123
 Sirocypis, 130
 Sirodesmium, 153
 Sirodothis, 123
 Siropatella, 134
 Sirothecium, 124
 Sirozythia, 128

- Sistotrema, 108
 Skepperia, 106
 Skierka, 99
 Stigmella, 153
 Solenia, 107
 Solenoplea, 30
 Solorina, 75
 Solorinella, 76
 Solorinina, 75
 Sordaria, 28
 Sorokinia, 68
 Sorosporella, 138
 Sorosporium, 102
 Sorotheia, 32
 Sparassia, 105
 Spathularia, 91
 Spegazzinia, 163
 Spegazzinula, 45
 Spermodermia, 162
 Sphacelia, 159
 Sphacelotheca, 102
 Sphaerella, 30
 Sphaeriaceae, 25
 Sphaeriales, 21
 Sphaericeps, 117
 Sphaeridium, 160
 Sphaerita, 9
 Sphaeroidaceae, 121
 Sphaerobolus, 121
 Sphaerocolla, 159
 Sphaerocybis, 117
 Sphaeroderma, 44
 Sphaerodes, 44
 Sphaerographium, 127
 Sphaeromyces, 163
 Sphaeronaema, 122
 Sphaeronaemella, 129
 Sphaeropezia, 62
 Sphaerophoropsis, 77
 Sphaerophorus, 72
 Sphaerophragmium, 101
 Sphaeropsidae, 121
 Sphaeropsis, 124
 Sphaerosoma, 91
 Sphaerospora, 90
 Sphaerosporium, 158
 Sphaerostilbe, 45
 Sphaerostilbella, 44
 Sphaerotheca, 22
 Sphaerulina, 33
 Sphaleromyces, 20
 Spheconisca, 39
 Sphinctrina, 71
 Spicaria, 143
 Spicularia, 140
 Spilomium, 162
 Spilonema, 74
 Spilopezis, 85
 Spinellus, 13
 Spira, 153
 Spirillaceae, 7
 Spirillum, 8
 Spirochaete, 8
 Spirographa, 59
 Spirogyrales, 12
 Spirosoma, 8
 Spolverinia, 27
 Spondylocadium, 152
 Sporendonema, 149
 Sporochisma, 152
 Sporocybe, 157
 Sporocystis, 161
 Sporoderma, 159
 Sporodesmium, 153
 Sporodinia, 13
 Sporoglena, 148
 Sporomega, 57
 Sporonema, 133
 Sporopodium, 75
 Sporormia, 35
 Sporostatia, 77
 Sporotrichella, 142
 Sporotrichum, 141
 Spragueola, 91
 Spumatoria, 30
 Squamotubera, 29
 Stachybotryella, 147
 Stachybotrys, 147
 Stachylidium, 150
 Stagonopsis, 129
 Stagonospora, 126
 Stamnaria, 67
 Starbaeckia, 69
 Staurochaeta, 123
 Staurothele, 39
 Steganosporium, 137
 Stegia, 61, 63
 Steinera, 74
 Stella, 117
 Stemmaria, 157
 Stemphyliopsis, 145
 Stemphylium, 153
 Stenocybe, 71
 Stephanoma, 164
 Stephensia, 97
 Stereochlamys, 41
 Stereocaulum, 78
 Stereum, 106
 Sterigmatocystis, 140
 Sterile mycelia, 164
 Stichomyces, 19
 Stichopsora, 100
 Sticta, 81
 Stictidaceae, 62
 Stictina, 82
 Stictis, 64
 Stictoclypeolum, 85
 Stictophacidium, 63
 Stigmatea, 30
 Stigmatella, 160
 Stigmatomyces, 20
 Stigmatula, 27
 Stigmella, 153
 Stigmina, 151
 Stilbaceae, 154
 Stilbocrea, 45
 Stilbomyces, 156
 Stilbonectria, 46
 Stilbospora, 137
 Stilbothamnium, 157
 Stilbum, 155
 Stirochaete, 150
 Strasseria, 122
 Streptococcus, 8
 Streptotheca, 92
 Streptothrix, 148
 Strigula, 41
 Strigulae, 41
 Strobilomyces, 109
 Stromne, 44
 Stropharia, 114
 Strumella, 162
 Stuartella, 35
 Stylobates, 111

Stypella, 104
 Stypinella, 103
 Stysanus, 157
 Suillus, 109
 Sydowia, 34
 Symphyosira, 156
 Synalissa, 72
 Synarthonia, 60
 Syncephalastrum, 14
 Syncephalidae, 14
 Syncephalis, 14
 Synchytriae, 10
 Synchytrium, 10
 Synglonium, 56
 Synsporium, 147
 Synthetospora, 145

T

Taeniophora, 134
 Tapellaria, 75
 Tapesia, 85
 Taphridium, 93
 Taphrina, 93
 Tarichium, 14
 Tarzetta, 89
 Telimena, 49
 Teratomyces, 21
 Terfezia, 98
 Terfeziopsis, 97
 Testicularia, 119
 Testudina, 24
 Tetrachytrium, 11
 Tetracium, 146
 Tetraccladium, 146
 Tetracoccusporis, 153
 Tetracoccusporium, 153
 Tetraploa, 153
 Thamnidium, 13
 Thamnolia, 82
 Thaxteria, 31
 Thecophora, 102
 Thecopsora, 100
 Thecospira, 160
 Thecostroma, 135
 Thelebolus, 92
 Thelenidia, 39
 Thelephora, 106
 Thelephoraceae, 106
 Thelidium, 39
 Thelocarpum, 43
 Theloporus, 110
 Thelopsis, 41
 Theloschistes, 83
 Thelotrema, 80
 Thelotremae, 80
 Thermomyces, 141
 Thermitis, 74
 Therrya, 38
 Thielavia, 23
 Thielaviopsis, 149
 Thiothrix, 7
 Tholurna, 72
 Thoracella, 125
 Thozetia, 160
 Thraustotheca, 15
 Thrombium, 40
 Thuemenella, 44
 Thwaitesiella, 108
 Thyrea, 73
 Thyridaria, 35
 Thyridella, 36
 Thyridium, 37
 Thyronectria, 46
 Thyrsidium, 136
 Thysanothecium, 78
 Tiarospora, 125
 Tichospora, 37
 Tichosporella, 36
 Tichothecium, 32
 Tilachlidium, 155
 Tilletia, 101
 Tilotus, 112
 Tirmania, 97
 Titaea, 146
 Titania, 35
 Tjibodasia, 103
 Tolypomyria, 142
 Tolyposporella, 102
 Tolyposporium, 102
 Tomasiella, 42
 Toninia, 77
 Torrendia, 119
 Torrubia, 47
 Torsellia, 123
 Torula, 147
 Torulina, 147
 Toxosporium, 137
 Trabutia, 28
 Trachyxyllaria, 32
 Tracyella, 131
 Trametes, 110
 Treleaseia, 45
 Treleaseiella, 129
 Trematosphaeria, 35
 Trematosphaeriopsis, 35
 Trematosphaeris, 35
 Tremella, 104
 Tremellaceae, 103
 Tremellae, 103
 Tremellastrum, 105
 Tremellodendron, 104
 Tremellodon, 107
 Tremellopsis, 105
 Tremotylum, 80
 Trichaegum, 153
 Trichaleuris, 90
 Trichaster, 118
 Trichobelonium, 85
 Trichobotrys, 147
 Trichocladium, 151
 Trichocollonema, 127
 Trichoconis, 145
 Trichocrea, 130
 Trichoderma, 140
 Trichodytes, 137
 Tricholoma, 111
 Trichopeltis, 52
 Trichopeltium, 131
 Trichopeltulum, 131
 Trichophila, 131
 Trichophyma, 95
 Trichopsora, 100
 Trichoscypha, 90
 Trichoseptoria, 127
 Trichosperma, 130
 Trichosphaerella, 27
 Trichosphaeria, 27
 Trichosporium, 148
 Trichostroma, 162
 Trichotheca, 158
 Trichothecium, 144
 Trichothelium, 42
 Trichothyrium, 53

Trichotrema, 41
 Trichurus, 157
 Tridentaria, 146
 Triglyphium, 161
 Trigonosporium, 122
 Trimmatostroma, 163
 Trimmatothele, 39
 Trinacrium, 146
 Triphragmium, 101
 Triplicaria, 162
 Tripospora, 54
 Triposporium, 154
 Trochila, 63
 Trogia, 112
 Troposporella, 164
 Troposporium, 162
 Trullula, 136
 Tryblidaria, 70
 Tryblidiaceae, 65
 Tryblidiella, 57, 66
 Tryblidiopsis, 65
 Tryblidis, 65
 Tryblidium, 56, 65
 Trypetheliae, 42
 Trypethelium, 42
 Tubaria, 113
 Tuber, 97
 Tuberaceae, 96
 Tuberales, 94
 Tubercularia, 159
 Tuberculariaceae, 158
 Tuberculina, 159
 Tubeufia, 47
 Tuburcinia, 102
 Tulasnellaceae, 107
 Tylophorella, 71
 Tylophorum, 71
 Tylopilus, 109
 Tylostoma, 117
 Tympanis, 66
 Tympanopsis, 29
 Typhula, 106

U

Uleomyces, 47
 Ulocolla, 104
 Umbilicaria, 77
 Uncigera, 143

Uncinula, 22
 Underwoodia, 91
 Uredinaceae, 98
 Uredinales, 98
 Uredinopsis, 100
 Uredo, 99
 Urnula, 65, 89
 Urobasidium, 107
 Uroconis, 126
 Urocystis, 102
 Urohendersonia, 126
 Uromyces, 98
 Uromycodes, 98
 Urophlyctis, 12
 Uropyxis, 99
 Urospora, 27
 Urosporella, 27
 Urosporium, 152
 Usnea, 82
 Usneae, 81
 Ustilaginaceae, 101
 Ustilaginula, 102
 Ustilago, 101
 Ustilagopsis, 102
 Ustilina, 29

V

Valsa, 26
 Valsaria, 33
 Valsella, 26
 Varicellaria, 80
 Vaucheriales, 14
 Velutaria, 86
 Venturia, 31
 Vermicularia, 123
 Vermiculariella, 125
 Verpa, 91
 Verrucaria, 39
 Verrucariaceae, 38
 Verrucariae, 39
 Verticicladium, 150
 Verticilliae, 142
 Verticilliopsis, 143
 Verticillium, 143
 Vialaea, 38
 Vibrisea, 91
 Virgaria, 148
 Vizella, 51

Volutella, 160
 Volutellaria, 160
 Volutina, 160
 Volvaria, 112
 Volvariella, 113
 Volvoboletus, 108

W

Wallrothiella, 27
 Weinmannodora, 124
 Willia, 94
 Winteria, 36, 37
 Winterina, 33
 Wojnowicia, 126
 Woodiella, 69
 Woronina, 10

X

Xanthocarpia, 84
 Xanthoria, 83
 Xenodochnus, 100
 Xenopus, 141
 Xenosporium, 153
 Xerotus, 112
 Xylaria, 29
 Xylariodiscus, 29
 Xylobotryum, 32
 Xyloceras, 32
 Xylocadium, 157
 Xylocrea, 44
 Xyloglyphis, 64
 Xylogramma, 64
 Xylographa, 59, 63
 Xylopodium, 119
 Xyloschistes, 59
 Xylostroma, 164

Y

Yoshinagaia, 51
 Ypsilonia, 122

Z

Zaghouania, 98
 Zignoella, 33
 Zignoina, 27
 Zimmermanniella, 48
 Zodiomyces, 21

Zopfia, 24	Zygochytriae, 12	Zygosporium, 150
Zukalia, 24	Zygochytrium, 12	Zythia, 128
Zukalina, 92	Zygodesmus, 148	Zythiaceae, 128
Zukaliopsis, 24	Zygosaccharomyces, 94	Zythiae, 128