TEXT-BOOK
IN
GENERAL PHYSIOLOGY
AND
ANATOMY

WALTER HOLLS EDDY

CORNELL
UNIVERSITY
LIBRARY

FROM

The Estate of

F.R. Macmillan

Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

A TEXT-BOOK
IN
GENERAL PHYSIOLOGY
AND
ANATOMY

BY
WALTER HOLLIS EDDY
CHAIRMAN OF THE DEPARTMENT OF BIOLOGY IN THE
HIGH SCHOOL OF COMMERCE, NEW YORK

NEW YORK ./. CINCINNATI ./. CHICAGO
AMERICAN BOOK COMPANY

laboratory course in zoölogy these chapters should be omitted, except for general reading.

This text is not intended to be used as a substitute for laboratory study. In my *Experimental Physiology and Anatomy* are outlined laboratory exercises which should be first performed by the pupil himself, if possible, before he is assigned a lesson in this text. While written as a supplement to the laboratory guide mentioned, the text is suited for use with any other laboratory manual and, where lack of laboratory facilities renders such preliminary work impossible, its description of the laboratory exercises is sufficiently full to permit its use where a textbook alone must serve.

The text meets both the optional and required work of the New York State Syllabus and the requirements of the Harvard entrance examinations.

Its topical arrangement and the printing of less important details in finer type make it adaptable to longer or shorter courses of study; and in these structural details the author has endeavored to make the book meet the needs of all classes of schools.

I wish to acknowledge the helpful suggestions given me by my colleagues of the High School of Commerce, and by Professor Geis of Columbia University and Dr. Darling of Harvard University. I am especially indebted to my colleague Mr. C. W. Hahn for his very careful reading of the proofs and his many helpful suggestions, and to my wife for assistance in the revision of grammatical arrangement and in the mechanical details of the work.

W. H. EDDY.

THE HIGH SCHOOL OF COMMERCE,
New York.

CONTENTS.

	Page
I. WHAT THE STUDY OF PHYSIOLOGY MEANS.	7
II. CHEMICAL COMPOSITION OF LIVING MATTER	12
III. THE STRUCTURAL UNITS OF LIVING MATTER	28
IV. STRUCTURAL UNITS OF HUMAN TISSUES — HISTOLOGY	39
V. FOODS AND NUTRITION	51
VI. FOOD ACCESSORIES; ALCOHOL AND TOBACCO	73
VII. DIGESTION	90
VIII. DIGESTION (<i>continued</i>)	107
IX. ABSORPTION	133
X. DIGESTION IN THE LOWER ANIMALS	146
XI. BLOOD AND LYMPH	159
XII. CIRCULATION	180
XIII. CIRCULATION (<i>continued</i>)	191
XIV. CIRCULATION IN THE LOWER ANIMALS	212
XV. THE SKELETON	221
XVI. THE SKELETON (<i>continued</i>)	244
XVII. SKELETONS OF THE LOWER ANIMALS	252
XVIII. MUSCLES	265
XIX. MUSCLES (<i>continued</i>)	278
XX. MOVEMENT IN LOWER ANIMALS	294

	Page
XXI. RESPIRATION	312
XXII. RESPIRATION IN LOWER ANIMALS	340
XXIII. EXCRETION	352
XXIV. SKIN STRUCTURE AND EXCRETION IN THE LOWER ANIMALS	374
XXV. THE NERVOUS SYSTEM	383
XXVI. THE NERVOUS SYSTEM (<i>continued</i>)	411
XXVII. THE NERVOUS SYSTEM IN THE LOWER ANIMALS .	429
XXVIII. THE SPECIAL SENSES	436
XXIX. THE SPECIAL SENSES (<i>continued</i>)	452
XXX. THE SPECIAL SENSES (<i>concluded</i>)	470
XXXI. SENSATION IN THE LOWER ANIMALS	480
XXXII. THE VOICE	486
XXXIII. BACTERIA AND SANITATION	493
INDEX	505

PHYSIOLOGY AND ANATOMY

I. WHAT THE STUDY OF PHYSIOLOGY MEANS.

FROM our earliest days we are familiar with living forms. We know that each living form must perform certain actions every day in order to live; but few of us have a clear idea of why these actions maintain life, or of the part each separate action plays in keeping us alive. For example, if asked why we eat, we answer, "To keep us alive," but that answer does not explain how food gives life. Again, we see gardeners watering plants, repotting them in new earth, taking care to place them in sunlight, and we recognize that all these actions are necessary to the life of the plants. We know that dogs and cats, horses and cows, must be fed and cared for to keep them alive and healthy. In fine, we know that to maintain life all living bodies, whether plant or animal, must repeat certain actions continually, or else die ; but we do not know how these actions produce life. It is this question of "how" and "why" that the present study of Physiology is to answer.

Method of study. — How shall we begin the study? An example will indicate the method. If we see a man moving the handle of a certain machine up and down and see a stream of water issuing from another part of the same machine, we know at once that the movement of the handle in some way causes the flow of water. In

order to answer the question "how" we should first pull our pump or a similar machine to pieces and find what parts it has and how they are placed in relation to one another. We should call this the study of the *structure* of the pump. Next, we should set the machine in motion and observe the action of the parts and how the movement of one affects the other. We should call this the study of *function*, or use of parts. Combining our knowledge of structure and function we might go still further, and, by experimenting, find that certain actions, such as lubricating the parts with oil, improve the action of the machine as a whole. We call this the study of the *conditions best suited to the action of the machine*.

Applied to the study of living objects such a method of investigation resolves itself into the following divisions of research.

First. The study of the structure of the living bodies. Such a study is called *anatomy* (*ana*, up; *temnein*, to cut) and is dependent upon dissection.

Second. The study of the use or functions of the various structures. Such a study is called *physiology* (*physiologia*, an investigation into the nature of things) and is dependent upon experimenting with living structures.

Third. The study of methods that will cause the parts to act to the best advantage. This study is called *hygiene* (*hygeia*, health) and like physiology is dependent upon experimentation.

Formulating these three directions we have the following definitions:

ANATOMY. — The study of the structure of a plant or animal to determine the parts present and the relative position of these parts.

PHYSIOLOGY. — The study of the function or use of the parts of a plant or animal body and their method of action.

HYGIENE. — The study of the means best adapted to maintain the parts and actions of the plant or animal body in a working or healthful state.

Anatomy.—Like a pump, a living body is made up of parts arranged in definite positions. It is manifestly impossible to locate all these parts from the study of the living body. To get at the internal structure it is necessary to cut or dissect the dead body. In the case of the human body such dissection is of course impossible for the high school pupil. It is here that a fact of great importance comes to our rescue, namely, that the arrangement of parts in our bodies is so similar to that in certain other animal forms that by studying these forms we may learn of our own structure. In other words, the dissection of the rat, cat, or rabbit, enables us to know our own structures.

Even with this difficulty overcome, however, many parts are so small that the eye cannot make them out. To remedy this there has been devised an instrument called the microscope; and so important is this side of anatomical study that it has received a special name.

Histology (*histos*, a tissue; *logos*, speech) is the microscopic study of plant and animal parts.

Finally, chemists tell us that the parts themselves are composed of simpler substances called *chemical elements*. This branch of our study is called *chemical anatomy*.

When we have studied a body from these three points of view, its gross structure, its microscopic structure, and its chemical composition, we shall have exhausted all that anatomy has to tell us.

Physiology. — Anatomy tells us only of the structure of the lifeless body. It is physiology that tells us of the action of the parts in the living body. Here the problem of investigation becomes difficult, but again we call to aid the fact that vital processes in plants and animals are similar to those in our own bodies, and the experiments we employ with the former tell us the workings of our own body. These experiments are supplemented by the reports of surgeons of their operations on human bodies, while another branch of science, called *physiological chemistry*, has taught us how to reproduce artificially many of the actions which take place normally in our bodies.

Hygiene. — The best engineer is the man who knows not only how his engine is made and acts, but also how to keep it in order and to run it to the best advantage. It is this knowledge — how to run our bodies to the best advantage, that is, to keep them healthy — that the study of hygiene gives us. It can be acquired only by taking to ourselves the experience of those most familiar with the running of our bodies. There are, however, certain of these experiences which have been formulated under the *laws of sanitation* which will give us a better foundation for our apprenticeship. Such laws are those that have been formulated by doctors, physicists, and bacteriologists, and we shall study them in their proper place.

Relation to other sciences. — The preceding pages indicate in a general way what the scope of our study is to be. What relation does this study bear to other sciences?

Biology (*bios*, life; *logos*, speech) is the name given to the study of living as distinct from lifeless matter. This subject naturally groups itself into the study of plants, *botany*, and the study of animals, *zoölogy*. From what has

been said already it will be seen that the terms anatomy, physiology, and hygiene refer to methods of study applied to either plants or animals, for example, *plant physiology*, *animal anatomy*, etc. — or they may be restricted to individual members of each group, *human physiology*, *horse anatomy*, *yeast physiology*, etc. In short, physiology, anatomy, and hygiene are merely one phase of the study of living matter (biology) and while our present problem is the human body, we must bear in mind that many of the laws that hold true of our bodies are merely special cases of laws which apply to all living organisms.

The great problem which all biologists, zoölogists, and physiologists are seeking to answer is, "What is life?" and in whatever field they may be working, their results are valuable and throw light upon this main, great problem.

II. CHEMICAL COMPOSITION OF LIVING MATTER.

Chemical elements. — The dissection of a plant or animal shows it to be made up of many different kinds of materials, some soft, some hard, some elastic. Dissection, however, can tell us nothing of the composition of these separate masses of material, and to find out what bone, muscle, and fat are made of, we must go to the chemist. As the result of methods which chemists have learned to apply to the problem, they tell us that muscle, fat, bone, wood, and in fact all the other substances of the universe are built up of simple substances which are called elements. By examining all the different kinds of matter, both living and lifeless, to be found in the universe, it has been found possible to reduce the number of these simple substances or elements to some seventy-six, and just as all the words in the language may be built up of the twenty-six letters of the alphabet, so all matter is built up by varying combinations of these seventy-six elements. Some of these elements are very rare, while others are exceedingly abundant. Some are never found in the pure state but always combined with other elements, while others are found both in a free state and in combination. Of the seventy-six known elements, twelve¹ are always found in living matter and six of the remaining sixty-four are sometimes present.

¹ The twelve that are always present in living matter are phosphorus, sulphur, carbon, oxygen, hydrogen, nitrogen, chlorine, potassium, sodium, calcium, magnesium, and iron.

Characteristics of chemical elements. — The distinguishing feature of a chemical element is that it cannot be resolved by any known means into a simpler substance. Thus while it is possible to resolve a substance like common salt into two substances known respectively as sodium and chlorine, no one has ever been able to separate sodium or chlorine into anything simpler, and these two substances are, therefore, called elements. Iron, sulphur, gold, silver, are familiar examples of elements. Since the chemical elements are the substances out of which all matter is formed, it will be well for us to examine a few of them to learn their properties and the manner in which they combine to form compounds. We shall select for examination those which are particularly concerned in the composition of our body materials.

Phosphorus. — (See Ex. III.) This element forms a large part of our bones; these are, in fact, composed mainly of a compound of this element with calcium and oxygen. In its pure state phosphorus has a yellow color and a waxy consistency. When exposed to the air it gives off white fumes which have a peculiar odor. These fumes are a compound of phosphorus with the oxygen of the air, and to keep phosphorus from combining with oxygen it is necessary to cover it with water.

These fumes teach us of a peculiar property of phosphorus, namely, its power to combine with another element. Such power on the part of an element is called its *chemical affinity*. All elements possess this affinity for certain other elements, but the strength of the affinity varies with the element and with the external conditions. To this affinity of elements all chemical compounds are due.

Chemical compounds distinguished from physical mixtures. — Most of the substances found in nature are mixtures of several substances. If we examine a piece of wood we can pick out several different kinds of matter which are intimately united in the compound wood. Similarly a piece of flesh shows pieces of fiber, and bits of fat, blood, and muscle, which are all intimately mixed together, but each of which possesses properties peculiar to itself. If we examine a bit of common salt and compare it with the elements sodium and chlorine, we find that the salt has properties which are distinct from either the sodium or the chlorine. In short the sodium and chlorine which have combined to form salt have lost their identity and individual properties, and the compound resulting from their union has entirely new properties.

These examples illustrate the distinction between a mechanical mixture and a chemical compound. The flesh is a mechanical mixture of fat, blood, etc. The salt is a chemical compound.

Whenever two or more elements combine, losing their individual properties and producing a substance with entirely new properties, the combination is called a *chemical compound*. The white fumes which result from the union of phosphorus with the oxygen of the air are entirely distinct in properties from the phosphorus and oxygen. In their formation, therefore, we have an example of the formation of a chemical compound. If the phosphorus is exposed to the air in a dark room a dull glow of light may be seen. This glow is due to the heat generated in the act of union of the phosphorus and oxygen. A certain amount of heat is always generated when chemical elements combine.

Sulphur. — (See Ex. IV.) Another important element in our body compounds is sulphur. Most of us have seen this element in its pure state and recall its yellow color. The sulphur used on sulphur matches is the element in its pure state, and we may recall the fact that when it burns it gives off suffocating fumes and a blue flame. In its free state, this element is found only around volcanic regions. Combined with other elements (chemical compounds of sulphur) it is extremely abundant. Muscle is such a compound of sulphur with other elements. Pure sulphur is odorless and tasteless.

When flesh decays it gives off a disagreeable odor. This odor is a property of a gas called hydrogen sulphide, which the sulphur forms as flesh decays, by combining with the element hydrogen. The ordinary hen's egg contains sulphur, and in decay this gas is formed abundantly. The blackening of a silver spoon used in eating an egg is due to a black compound called sulphide of silver, which the sulphur of the egg forms with the element silver.

The fumes from burning sulphur (a compound of sulphur and oxygen), the bad smelling gas (sulphur and hydrogen) and the black compound (sulphur and silver) bring out the fact that a chemical element may have an affinity for more than one element. The fact that it occurs free in the air and yet combines with oxygen to form a compound when heated is an illustration of the way in which external conditions may affect the affinity of an element, while the blue flame which results when sulphur combines with oxygen and which is absent in the formation of the other two compounds, shows that the amount of heat generated by the union of two elements may vary greatly with the elements.

Carbon. — (See Ex. V.) When we partly burn a piece of wood, bone, or flesh, we find it transformed into a black substance which we call *charcoal*. In nature we find this same substance in a much harder form and we call it *coal*. When we use our lead pencils the mark we make is due to another form of this substance called graphite. Finally in the diamond we have a substance very different in appearance from charcoal, coal, or graphite, but essentially like them in some respects. All of these substances are forms of a very abundant chemical element called carbon. So important a part of all living matter is this element that the study of its compounds is made a special branch of chemistry under the name Organic Chemistry.

In most of its forms, carbon shows a black color and is without odor or taste. In its free state it has so little affinity for other elements that farmers often char the ends of fence posts before setting them in order to insure against decay. When heated to a certain temperature, however, all forms of carbon burn and form a colorless gas, which may be readily detected by its power to turn clear lime water milky. This gas is a chemical compound of carbon with the oxygen of the air and is called *carbon dioxide*. It is this gas that is used to charge water in the manufacture of ordinary soda water. Just as in the union of phosphorus and sulphur with oxygen, the formation of this gas is accompanied by the generation of much heat, and this fact is made use of in our bodies as well as in our stoves and furnaces. If we blow our breath through clear lime water the milky color that results gives unmistakable evidence that this gas is being constantly formed in our bodies and given off with the exhaled air.

A study of the element carbon demonstrates three important facts.

First. It is present in large amounts as a part of all living matter.

Second. Its combination with oxygen to form carbon dioxide is always accompanied by the production of heat, and the formation of this gas in our bodies results in maintaining a constant body temperature.

Third. A chemical element in its free state may occur in forms which appear very different to the eye, and must be tested by other means than sight to determine its identity.

Oxygen. — (See Exs. VII. and VIII.) This important element occurs free in the air in the form of a colorless, odorless, and tasteless gas. In its combined form it is abundant in both living and non-living matter. It is of great importance to all living processes because it has a great affinity for other elements, and because, in combining with other elements, it sets free a great amount of energy in the form of heat.

While oxygen is present in the air in a free state, it is so mixed with other gases that special processes must be resorted to to obtain it pure. The simplest method of obtaining oxygen is to heat a compound which contains it. Some of these compounds when so heated break up and set free the pure gas. Two compounds which are especially favorable to this method are oxide of mercury (a compound of mercury and oxygen), and potassium chlorate (potassium, chlorine, and oxygen).

If we heat some oxide of mercury in a tube, as shown in Figure 1, we note that it gradually disappears. If, while the heating is going on, we insert the glowing end

of a charred match into the mouth of the tube the presence of the oxygen gas will become evident by the increased brilliancy of the glow, due to the rapid union of the charcoal with the liberated oxygen. The presence of the mercury will be seen in the drops which condense on the cooled sides of the tube. Such an experiment not only illustrates the manner in which a compound may be separated into its elements, but also demonstrates an important property of oxygen, namely, that in its pure state its affinity for carbon is much greater than when mixed with other gases in the air.

FIG. 1 — Apparatus for heating mercuric oxide.

If we heat a mixture of manganese dioxide¹ and potassium chlorate and collect the liberated gas over water as shown in Figure 2, it is possible to compare the union of sulphur and phosphorus with this pure product, with the result when these elements combined with the mixed oxygen of the air. In every case the union is much more violent than when ordinary air is present, and results in more heat being liberated.

FIG. 2 — Apparatus arranged for obtaining oxygen.

Oxidation and combustion. — Such unions of elements with oxygen as have been described in the preceding pages, are spoken of as *oxidations* and the compounds

¹ The manganese dioxide undergoes no change, but merely facilitates the decomposition of the potassium chlorate.

formed are called *oxides*. For example, a chemist would describe the formation of carbon dioxide as the *oxidation* of carbon to form an *oxide*. In every case the formation of an oxide is accompanied by the production of heat, but sometimes the union of the elements takes place much more rapidly than at others, and when this occurs the amount of heat produced is large and often accompanied by flames or light. Such rapid unions are called *rapid oxidation* or *combustion*, to distinguish them from slower unions or slow oxidation. All burning then may be explained as the rapid oxidation of the substance burned, and it is evident, therefore, that burning cannot take place in the absence of oxygen. Further, since in burning the oxygen combines with the substance to form an oxide, it follows that for a substance to continue burning implies a continuous supply of oxygen until the substance has entered completely into combination. It is for these reasons that we supply our stoves with draughts which will permit the continuous entrance of oxygen. This explanation of the true nature of combustion also explains the methods of extinguishing fires, all such methods aiming to separate, through the aid of some substance which does not combine with oxygen, the burning substance from its air or oxygen supply. In short, pouring water on a fire puts an incombustible substance between the burning matter and the air, and thus prevents further oxidation. Water is more frequently used than sand or dirt for this purpose because it is easier to handle, but in theory one is as effective as the other.

The special interest of oxidation to the physiologist consists in the fact that it furnishes a means for the production of heat, and we shall learn later how the body

makes use of oxidation to maintain the body temperature, while the necessity for a continuous oxidation explains the presence of carbon in the body and the continuous entrance of oxygen and exit of carbon dioxide in breathing.

Nitrogen. — Experiments with pure oxygen show that combustion takes place more rapidly in it than in air. This implies that air must contain something besides oxygen, which dilutes it and makes oxidation slower. What is this substance? A simple experiment answers this question. (See Ex. IX.). Place a piece of phosphorus the size of a pea in an evaporating dish that is floating on water. Light the phosphorus and thus raise its temperature to the combining point. Then cover the dish quickly with a bell jar (Fig. 3). The phosphorus

FIG. 3 — Apparatus arranged for obtaining nitrogen from air.

will at once begin to combine with the oxygen in the enclosed air, and when it stops burning it will be because there is no more oxygen to combine with it to form the white fumes of phosphorus oxide. If we let the fumes settle they will dissolve in the water and there will be left in the jar only what was in the air before the oxygen was taken out.

Further, since the water rises in the jar to take the place of the consumed oxygen the volume of gas left in the jar subtracted from its original volume will give us the proportion of oxygen present in a given volume of air; roughly one fifth of its volume.

What is this gas that is left? If we examine it by

inserting heated elements as we did oxygen, we find that while it is colorless, odorless, and tasteless like oxygen, it has not the power of that element to combine with these other elements or produce combustion. This residual gas which is so lacking in chemical affinity is an element called *nitrogen*. Like sulphur, however, its lack of affinity is due rather to lack of the right conditions for combination than to absence of that power. In nature, nitrogen occurs very abundantly in combination with other elements, and some of its compounds form important parts of our body material, notably muscle or flesh. Owing to the weakness of its affinity for most elements its compounds

are apt to break up readily, and on that account are spoken of as unstable compounds. Most of our explosives, such as gunpowder and nitroglycerine, owe their explosive power to the fact that they are unstable compounds of nitrogen.

FIG. 4—A, apparatus for obtaining hydrogen and oxygen with the aid of an electric current. B, method of collecting the gases.

Hydrogen. — (See Exs. X. and XI.). If we allow an electric current to pass through water (Fig. 4, A) which has had a little acid added to it to make it a conductor

of the current, we find that the current has the power to separate the water into two gases. One of the gases is twice as great in volume as the other and shows very different properties. If we collect these gases (Fig. 4, *B*) and examine them we find that oxygen is the smaller volume. The other gas is colorless, odorless, and tasteless, like the first, but when we introduce a lighted splinter into it, it takes fire with an explosion and burns with a very hot colorless flame. If, instead of using water and the electric current, we pour sulphuric acid upon zinc (Fig. 5)

FIG. 5 — Apparatus arranged for obtaining hydrogen from zinc and sulphuric acid.

the zinc will cause the acid to split and set free large quantities of this same substance which we may collect and examine more carefully. Such an examination will show that not only does this substance burn, but that when it does burn it forms water.

This peculiar gas is a chemical element called *hydrogen* and since burning is

simply the result of the combining of an element with oxygen, we must conclude that water is not an element but a compound of hydrogen and oxygen. The explosive force with which this combination takes place shows that the affinity of hydrogen for oxygen is very strong. Hydrogen has a strong affinity for many other elements beside oxygen, especially for the element carbon; and its compounds with this element form a very large part of living matter. Aside from being one of the most active of

chemical elements, hydrogen is also noteworthy as being the lightest chemical element known.

Metallic elements. — All the elements we have considered so far are called by chemists, *nonmetals*, to distinguish them from another group of elements called *metals*. Some of these metallic elements are very familiar to us under the names of iron, zinc, copper, gold, etc. Beside these more familiar metals, there are others that form important parts of our bodies but which very rarely occur uncombined owing to their greater affinity. Such are the elements calcium, potassium, sodium, and magnesium. The most important characteristic of metals is that when they combine, it is with a nonmetallic element, and very rarely with another metal. Such compounds of metals and nonmetals fall very readily under two classes, bases, and salts. All chemical elements are divisible into metals and nonmetals.

Acids, bases, and salts; neutralization. — When iron is exposed to the air it takes on a reddish coating that we call rust. This action is hastened by the presence of moisture. This rust is merely a compound of the metal iron with the oxygen of the air or water. Similarly other metals such as potassium, sodium, and calcium, etc., are able to unite with the oxygen of air or water and form *oxides*. If we dissolve some of these latter oxides in water the mixtures have peculiar properties. They have a slippery feel and a soapy taste, and when we put into them a piece of paper dyed red with the vegetable coloring matter, litmus, they change the color of the paper from red to blue. Such mixtures are not solutions but true chemical compounds of the oxides and the water, and are called bases. If we put the paper which was colored blue by these bases

into vinegar or lemon juice it will turn red again. The reason for this is that vinegar and lemon juice contain certain combinations of nonmetallic elements called *acids*. The acid present in vinegar is a combination of hydrogen, oxygen, and carbon, called acetic acid. Some other powerful acids are sulphuric acid (hydrogen, sulphur, and oxygen), nitric acid (hydrogen, oxygen, and nitrogen), and hydrochloric acid (hydrogen and chlorine). All acids have a sour taste, and contain no metals in their composition.

If we add drop by drop some of a base to an acid, we shall evidently come to a point when the tendency of the acid to turn litmus red is just counterbalanced by the tendency of the base to turn it blue. When this point is reached the base is said to have *neutralized* the acid, and the process is called *neutralization*. Finally, if we drive off the water of such a neutralized mixture by evaporation, we have left a product which turns litmus neither blue nor red and is called a salt. Analysis of such a product shows it to be composed partly of the metal which was in the base and partly of the nonmetals which were in the acid. You can make common table salt yourself by mixing hydrochloric acid and the base caustic soda (oxide of sodium) in this way. (See Ex. XII.) Salts, or compounds of metals and nonmetals, form an important part of the composition of our bodies, bones being a salt of calcium, phosphorus, and oxygen. Since most of the minerals in nature, limestone, quartz, etc., are metallic salts, such salts in the living matter are spoken of as mineral salts to distinguish them from compounds of purely nonmetallic elements.

CHEMICAL COMPOUNDS OF THE BODY.

The elements previously enumerated very rarely occur in an uncombined state in plant or animal matter; the only important exception to this rule being the element oxygen, and even that is more abundant in combined form than in the free state. The various compounds (acids, oxides, salts, and bases) are so numerous and varied that it would take too long to enumerate them all. Fortunately, they admit of classification into a few characteristic groups of life properties. These groups are called, respectively, minerals, gases, and carbon compounds.

Minerals. — Under this head are grouped water, mineral salts, and all acids that do not contain carbon. The only important acid in our bodies under this group is hydrochloric, and this is found in small quantities in the gastric juice of the stomach. Water forms more than fifty per cent, by weight, of all living matter. In the human body approximately 59% is water; the bones contain 22%, the liver 69%, the muscles 75%, and the kidneys 82%, while the fluids of the body are still richer in water content. Mineral salts will not burn, and by burning different parts of the body until nothing but the ash or mineral salts remain we cannot only demonstrate their presence in every part of the body, but also determine the form and proportion present. Such determination shows them to be of great variety as to kind. The most common are the chlorides, carbonates, and phosphates of the metals sodium, calcium, and potassium. These names indicate the predominant elements in these compounds. Bone ash also represents nearly 33% by weight of the entire bone,

and is composed mainly of calcium carbonate and calcium phosphate.

Gases. — Oxygen and carbon dioxide occur in all living matter. They may be free, as in the lungs, or absorbed in the fluids of the body. These two are the most important gases present, but others are present in smaller quantities.

Carbon compounds. — The compounds of this element in living matter are so numerous and so important that they have been given the name of organic compounds, or life compounds. They may be classified under two groups, nitrogenous and non-nitrogenous.

Nitrogenous carbon compounds. — As the name indicates, these compounds contain the element nitrogen in addition to the elements carbon, hydrogen, and oxygen. They are of many forms. *Proteid*, which contains sulphur, in addition to the above mentioned elements, is the name given to the form which is perhaps most abundant, and which under the name of albumin and fibrin in the blood, myosin in the muscles, and casein in milk, is very important in its relation to vital processes. When proteid burns it loses part of its carbon and the result is another nitrogen compound called urea, which is given off as a waste by the kidneys. Still another form of nitrogen compound is found in the gelatine of the bones and the keratin of the hair and nails.

Non-nitrogenous carbon compounds. — In this group are placed all carbon compounds found in living matter, which do not contain nitrogen. This group is divisible into two sub-groups, carbohydrates and fats.

Carbohydrates. — Besides carbon these compounds contain only hydrogen and oxygen, which are always in the

proportion in which they occur in water, namely, two parts of hydrogen to one of oxygen. Carbohydrates fall into two main divisions, the sugars and the starches. The most important of the sugars is glucose or grape sugar, found in the blood and in many other parts of the body. Starch occurs in the body in a form called glycogen, which is stored in the liver. When a carbohydrate is oxidized it liberates much heat, and the oxidation in the body of these compounds furnishes us much of our warmth.

Fat. — This substance is present in the body in varying amounts. Like the carbohydrates it is a compound of carbon, hydrogen, and oxygen, and like them it produces heat when oxidized and thus acts as a body fuel. It occurs largely in a liquid state in the living body, in the muscles, the blood, the bones, and connective tissues.

We shall consider the properties of these carbon compounds further, in our study of foods.

III. THE STRUCTURAL UNITS OF LIVING MATTER.

THE analyses of the chemist tell merely what elements or combinations of elements enter into the building of living matter. The principal interest in such analyses lies in the proof they give that all living matter, plant or animal, consists of essentially the same chemical elements.

FIG. 6— *A*, diagram of a human trunk split to show the internal skeleton. *B*, diagram of a grasshopper split to show the external skeleton. Hard parts are in black.

When we speak of the structural composition of a given plant or animal we do not refer to its chemical make up, but to the form which these compounds take. Thus the most evident structural forms these compounds take in our bodies we call eyes, ears, bones, flesh, etc., while in plants they receive the name of leaves, roots, stems, etc.

Gross structure of the body.

—Roughly, the body consists of a head, a trunk, and appendages (arms and legs). All these parts agree in being made up of soft matter supported by a framework of hard matter.

Lower animals, like the crayfish and grasshopper, also consist of hard and soft matter, but with them the hard matter, when present, is on the

outside and forms what is called an external skeleton, while in man it is internal with the soft parts packed upon it. If we examine a diagram (Fig. 7) of the human body split lengthwise the arrangement of these soft and hard parts is made clearer. Such a diagram shows that the head is really a box of bone resting on top of the backbone with an external covering of soft parts, the eyes, ears, cheeks, and scalp, and containing within a mass of soft material called the brain. This latter is continued down the center of the backbone in a long cord called the spinal cord. The trunk is divided into two cavities, one above the other. The upper cavity, the chest, is protected on each side by arches of bone called the ribs, on the front by the breastbone, and is separated from the lower cavity by a thin membrane called the diaphragm. The lower cavity is supported at the base by a broad basket of bones called the pelvis and at the back by the backbone. The upper cavity is filled with the heart and lungs, while the lower contains a number of soft organs (stomach, intestines, kidneys, etc.). When

FIG. 7 — Diagram of a human trunk split lengthwise; *a*, brain and cord cavity; *b*, nasal cavity; *c*, mouth cavity; *d*, regions of alimentary canal; *e*, heart in chest cavity or thorax; *f*, partition (diaphragm) separating the chest cavity from the abdominal cavity below.

split in the same manner the appendages show a central cylinder of bone surrounded by solid masses of soft matter called flesh. In addition to these general regions we find that the whole body is covered with a soft layer called skin, with openings at certain points for special structures such as eyes, ears, and mouth.

Character of the body parts. — When we examine the parts of the body more in detail, we find that they are capable of separation into portions that perform some definite kind of work. For example, in and on the head are parts called eyes, tongue, brain, etc., each of which has some special work to do. Again, the body cavities contain portions called the heart, the lungs, the stomach, etc., all with special work to perform. These specialized portions of matter which may be entirely soft or hard, or contain both soft and hard parts, are called *organs*. They are usually of a definite shape or outline and position. We may define an organ as a portion of the matter of the body which performs a special kind of work. For example, the eye is an organ for seeing, the ear for hearing, the heart for pumping, etc.

In some cases several organs combine and work to a common end, and we call such a collection of organs a system. For example, the brain, the spinal cord, and the nerves, are organs combined for the control of our actions. Likewise the stomach, liver, intestines, etc., all share in preparing food for the body, and hence constitute a system. We may define a *system* as a collection of organs so arranged as to accomplish some definite end by their combined actions. Systems are named, usually, according to the end attained, for example, the digestive system, the circulatory system, etc.

Composition of organs. — If we turn now to an examination of the organs, we find them made up of masses of material which differ both in appearance and action. For example, the nose consists of a hard framework of bone, an outer covering of skin, and underlying portions of red flesh, blood vessels, and nerves. Other organs, when similarly examined, are found to contain also these flesh, blood-vessel, and nerve layers, but in different positions. These substances of which the organs are built are called *tissues*, and the simplest structural units to which we can reduce the body with the aid of the knife are these tissues. Tissues then are the layers of matter of similar appearance which combine to form the organs.

If now we call the microscope to our aid, tissue, in turn, is seen to be made up of tiny bodies of definite structure and shape called *cells*.

The cell. — If we scrape the inside of the cheek with a

FIG. 8 -- Plant and animal cells; *A*, plant cell (spirogyra); *B*, animal cells scraped from the inside of the cheek; *C.W.*, cell wall; *C*, colored bodies; *P*, protoplasm; *S.B.*, spiral band; *N*, nucleus; *C.S.*, cell sap.

clean knife and mount the scrapings in a drop of water, for examination under the microscope (see Ex. XXII.), these

scrapings resolve themselves into a collection of irregular bodies whose most prominent structure is a round particle imbedded in what seems to be a granular liquid. The bodies are *cells*, the particle is a *cell nucleus*, and the granular liquid is called *protoplasm*.

If we examine in the same way a little of the common green pond scum (*Spirogyra*) we shall see that it also is made up of oblong bodies which contain a nucleus and protoplasm (see Ex. XXIII.). These plant cells differ from the cheek cells, in having a distinct boundary wall, and appear to be filled with long spiral bands of a green color.

Both the bodies mentioned above are typical cells,

FIG. 9 — Diagram of a cell. [After Wilson.] Notice in this diagram of a typical cell the cell protoplasm, composed of cell plasma and cell lymph, and containing cell foods, *C f.* and cell sap *C.s.* The large spherical nucleus has a true nucleolus *t.n.*, a false nucleolus *f.n.*, a nuclear network *N.n.*, and nuclear lymph *N.l.* *c* is the centrosome.

and all plants and animal tissues are built of such structures. For this reason the cell is called the *structural unit*

of all living matter. Assuming the fact that the above are typical cell structures we may then define a cell as follows: A cell is a boxlike body having an external wall that incloses a liquid called protoplasm, and in which floats a particle called a nucleus. Cells may vary in size, shape, and in the thickness of their walls, and may contain other substances beside the liquid protoplasm. Our examination of the other tissues of the body will determine whether this definition is complete (see pp. 39-51). Meanwhile let us examine this liquid protoplasm a little more fully.

Protoplasm. — Some one has said that "The study of physiology is the study of protoplasm." The meaning of this statement lies in the fact that the action of all living tissues depends upon the action of this substance. If that is the case, it is certainly worth while to learn what we can of its properties. Here, however, a difficulty arises. It is easy to obtain tissues whose cells are filled with this substance, but when these tissues are removed from the plant or animal body in which they belong, the protoplasm which they contain is killed and shows only a few of the properties of the living substance. Fortunately, however, there exists in nature a tiny animal form, only about a hundredth of an inch in diameter, which is made up almost entirely of protoplasm, uninclosed by any wall. Furthermore, this little animal, called the *amœba*, can be picked up from the surface of leaves in the still pools where it lives and be transferred alive, in a medicine dropper, to the slide of a microscope, where we can watch all its actions (see Ex. XXIV.).

Living protoplasm; the amœba. — The amœba is composed of liquid substance (protoplasm) of about the con-

sistency of the white of an egg. This liquid itself is colorless, but while it appears clear at the edges, its inner part is filled with tiny granules that give it a slightly grayish tinge. The clear outer part is called ectoplasm to distinguish it from the granular portion or endoplasm. These terms also (*ecto*, meaning outer; *endo*, meaning inner) serve to indicate the position of the two forms.

FIG. 10 — Amœba (*Amœba proteus*); *a*, food vacuole; *c*, contractile vacuole; *ec*, ectoplasm; *en*, endoplasm; *v*, water vacuoles; *n*, nucleus.

The endoplasm is seen also to contain many round bubble-like bodies (*vacuoles*) which are filled, some with water, some with small particles of food. Besides these we note a single dark body (*nucleus*) in which is a smaller dark spot (*nucleolus*). This nucleus is actually only another form of protoplasm, and the ectoplasm and endoplasm together are spoken of as cytoplasm to distinguish the nucleus from the liquid parts. In structure this tiny

body is identical with the particle scraped from the cheek, except that it has no boundary wall. It is, in fact, a single cell without a cell wall, and is one of the lowest forms of animal life known.

But we have not yet noted the most important feature of this tiny fluid body. If we watch it a moment we shall be able to see that it moves. The cytoplasm will be seen to flow out first in little projections (*pseudopodia*) which then fill as the rest of the mass flows into them, and in this way the body progresses across the slide. That this is not mere physical flowing like the running down hill of water is shown by the fact that it can change its direction. This power of automatic movement is possessed by no other fluid known, and in itself is sufficient to make this fluid remarkable.

If now we press down the cover glass with a needle or pin just above the animal, we shall note that the movement stops and that the animal draws back to form a round ball. If not disturbed further, it will expand and flow as before. In this simple action two more properties of protoplasm are shown; the power of expansion and contraction (*contractility*) and sensitiveness to a stimulus such as touch (*irritability*). Now watch the amœba until it comes in contact with a particle in the water. It flows over it and engulfs it, but curiously enough, if the particle is of vegetable matter it is carried along with the amœba, while if sand, it is soon cast out. The reason for this action is that the amœba, in selecting what to retain and what to reject, actually makes a distinction between foods and substances which are not foods. Further, if the particle is retained it is soon broken up in one of the round bubbles (food vacuoles) and this is slowly revolved

in the cytoplasm and continually decreases in size until it finally disappears entirely. What has happened? Simply this: This curious liquid has selected a food substance, broken it up, dissolved it, and distributed the dissolved product all over the body; while the fact that the product is no longer visible shows that it must have been transformed into protoplasm itself. In other words, protoplasm is capable of distinguishing and taking in food, of dissolving or digesting it, and of making new protoplasm out of it.

Finally, we may note that one of the vacuoles is larger than the others and expands and contracts rhythmically. As it expands this vacuole fills with a liquid that seeps from the protoplasm, and which the *contracting vacuole* forces out into the surrounding water. This liquid is the waste matter produced by the active protoplasm, and is formed by the breaking up of the chemical composition of the old protoplasm. Protoplasm, therefore, not only can assimilate food, that is, build it up into new protoplasm, but also has the power to dispose of its waste products.

Other experiments have taught that the amoeba absorbs oxygen as well as food, and that in the breaking up of the protoplasm the carbon which it contains unites with this oxygen and produces carbon dioxide with a consequent production of heat and power. The carbon dioxide is given off as a waste from the surface of the cytoplasm.

We may summarize then, as follows, the properties of this peculiar fluid, protoplasm, as exhibited in the amoeba:

- (a) Power of automatic movement.
- (b) Sensitiveness to stimuli.
- (c) Power to select food and to take it in.

(d) Power to dissolve and distribute food.

(e) Power to transform absorbed food into new protoplasm.

(f) Power to remove waste matter.

(g) Power to take in air and give off carbon dioxide.

A review of these properties shows that protoplasm performs in a simple way practically all the actions that we perform as human beings. If we recall that all cells contain protoplasm, and that tissues are made of cells, organs of tissues, and systems of organs, it is evident that the activity of this protoplasm must be very closely related to the actions of the organs and systems themselves, and hence to the activities of the organism that contains these systems.

Biological position of man. — Botanists and zoölogists (who study plants and animals) tell us that there are thousands of different kinds of plants and animals, but that when we examine the structure of these many forms of living matter, all are found to be capable of reduction to a common structural unit (the cell) and a common structural substance (the protoplasm). Furthermore, they tell us that the only essential difference in structure between high and low plants and animals is to be found in the greater number, kind, and complexity of arrangement of the cell units in the higher forms, as compared with the simpler arrangement, kinds, and numbers in the lower forms. Man, therefore, is to be considered as an animal composed of many cells of different kinds and action. He is placed highest in the living kingdom because in him we have a cellular organism in its highest development.

Evolution. — The theory has been set forth in what is known as the doctrine of evolution that all plants and animals have arisen from a common origin, namely, the substance protoplasm, and that each form of plant or animal has developed from a less complex form. According to this theory living matter once consisted of simple one-celled forms, and all the complicated forms we now know have been developed from these, much as the many kinds of houses in a city have been made by different arrangements of the bricks.

Classification. — In the animal world it is customary to group the various forms in an ascending series with the most complex form (man) at the head of the series. When this is done it is found that the series is roughly divisible into two groups, vertebrates, or animals with a backbone; and invertebrates, or those without a backbone. In the former class are included all the fishes, frogs, reptiles, birds, and mammals. In the latter are the one-celled protozoans, the worms, mollusks, insects, etc. Man, in common with horses, cows, rabbits, dogs, cats, etc., possesses hair, and glands for suckling its young (mammary glands), and hence belongs to the group of mammals. It is for this reason, namely, that all mammals are very much alike in structure, that it is possible for us to use the dissections of other mammals, such as the cat, rabbit, and the rat, to get a clear idea of our own structure. Many times in the pages to come we shall make use of this similarity of structure for illustrations of structural arrangement.

IV. STRUCTURAL UNITS OF HUMAN TISSUES — HISTOLOGY.

THE study of the structural units (the cells) that make up the human tissues, is called microscopic anatomy or *histology*. The methods of microscopy make it possible for us to cut very thin sections of tissue, to stain them with certain dyes so as to bring out the different parts of the cells, and finally, to mount them on glass slides in such a way that they may be examined with the compound microscope. This instrument, by magnifying many diameters, makes it possible for us to study details of structure that were otherwise invisible. The various tissues that are found in the body are found to be capable of classification into four general classes of tissues: epithelial, nervous, muscular, and connective. This classification is based upon the characters of the cells of which the tissues are formed.

Epithelial tissue. — All surfaces of the body, external and internal, are protected by layers of membranes. The external layer we call the skin; the layer which lines the mouth, digestive organs, and lungs we call the mucous membrane. When we examine these membranes under the microscope they are found to be composed entirely of cells united in layers by a thin cementlike substance, much as bricks are held together by mortar. Owing to the fact that the membranes which these cells form are covering or protective layers, the name of *epithelial* tissue has been given to them, and the cells of which they are

composed are called *epithelial cells*. As we have already noted, one of the functions of epithelial tissue is protection. A second function, exhibited only by certain layers of this tissue, is secretion. Thus the skin that covers the flesh and the mucous membrane that lines the mouth, both prevent injury to the underlying parts, but in addition to this protective action the mucous membrane secretes a liquid (the mucus) which keeps that membrane moist and flexible. All epithelial cells consist of a large oval nucleus floating in protoplasm of a granular or netted appearance, while the cell walls are extremely thin. Epithelial tissues are classified according to the shape and arrangement of the cells which compose them. Thus we speak of pavement, cubical, columnar, spheroidal epithelium, according as the cells which compose them are

FIG. 11 — Forms of epithelial tissue; *a*, 1, cylindrical cells; *a*, 2, squamous cells showing cement connection; *b*, ciliated epithelial cells; *c*, stratified epithelium, showing, 1, a pavement cell, 3, 4, spheroidal cells; *d*, columnar epithelium, composed of ciliated cylindrical and spheroidal cells.

flat, cubical, cylindrical, or spherical in shape (Fig. 11). When the layers are more than one cell layer deep we call it stratified epithelium. The cells scraped from the inside of the cheek (see Ex. XXII.) illustrate the flattened or

squamous form, while a section of the intestine lining (see Ex. XXV.) illustrates the columnar or cylindrical forms. Sometimes the cells are modified in various ways. For example, in the skin, hair, and teeth the external cells become hard or horny, while the cells that line the breathing passages may develop hairlike projections of protoplasm called cilia. These cilia wave back and forth and thus free the surface of tiny particles of dust, etc. In secretory epithelium the secreting cells are often goblet shaped.

Connective tissue. — If we attempt to pull apart a piece of beef we find that it resists this action by means of an intricate interlacing of white fibers that hold the separate parts together. These fibers make up what is called intermuscular tissue, which, in turn, is a form of connective tissue. These fibers are not cells, but are formed in the matter thrown off by certain cells. They are of two kinds, white inelastic, and yellow elastic, fibers, and their function is to surround and connect other layers. Hence the name, connective. It is evident then that the connecting is done by the substance given off by cells and not by cells themselves. This characteristic holds true of all forms of connective tissue. The cells are small and of importance only as makers of the connective part which is called inter-

FIG. 12 — Intermuscular connective tissue. Corpuscles are the cells which form the fibers.

cellular substance since it occurs between cells. The other important forms of connective tissue are cartilage and bone.

Cartilage. — An examination under the microscope (see Ex. XXVI. and Fig. 13) shows that cartilage, or gristle, consists of a solid mass surrounding pits very much after the manner of Swiss cheese. The solid mass is the elastic intercellular part of the connective tissue, and corresponds

FIG. 13 — Hyaline cartilage; *cap*, capsule in a lacuna; *m*, matrix formed by cells; *c*, cartilage cell; *n*, nucleus.

to the fibers of the intermuscular form. The pits or *lacunæ* are inclosed in *capsules*, and within these capsules lie the cells which produce the solid matter. These cells are the true structural units of this form of tissue, and the solid part is made by them. As this is thrown out by the protoplasm of the cells, it collects around them, and thus forms an important structural element of our bodies. For

example, the external ear and the nose owe their shape to underlying layers of this substance, while the ends of the bones are capped by layers of the same material, and thus protected from friction and jar.

Bone. — If, in addition to the gristly matter given off by the cartilage cells, there be added salts of lime, the intercellular mass becomes hardened, and we have a structure called bone. An examination under the microscope (see Ex. XXVI. and Fig. 14) shows that bone differs from cartilage in that the intercellular mass is arranged in rings

or *lamellæ*, about central canals (the *Haversian canals*). Further, the pits or *lacunæ* which contain the bone-forming cells are connected by fine tubes (*canaliculi*). The explanation of this difference of arrangement is found in the fact that the cells can give off only substances which they manufacture from the food brought to them. The Haversian canals contain blood vessels which bring to the

FIG. 14 — Bone tissue; *A*, piece of bone under the low power of the microscope, showing rings or *lamellæ* about central cavities (*Haversian canals*); *B*, piece of bone highly magnified, showing the *lacunæ* with the fine *canaliculi* radiating from them.

bone cells their food and lime. Food and lime ooze out of the blood vessels and finally reach the cells by way of the canaliculi. The protoplasm of the cells, then, manufactures from the food the cartilage material which it throws out with the lime, so that it accumulates in layers about the cells. The result of the similar action of many cells means a great accumulation of this material, and hence the production of supports for the soft parts of our body.

Connective tissue, then, consists of cells that manufacture and give off a substance which, in turn, binds together and supports other parts of the body. Besides the kinds mentioned, we have still other forms, named according to the character of the *intercellular material*. Thus, adipose tissue is a form of connective tissue in which the

thrown off matter consists of drops of oil or fat, held together by fibers (Fig. 15).

FIG. 15 — Adipose tissue;
a, connective fibers; b,
oil drops.

Muscular tissue. — This tissue, to which all our motions are due, consists of cells whose protoplasm has the power of contraction and expansion especially well developed. It is of two kinds and these kinds differ both in action and structure.

Striated, or *voluntary* muscular tissue, is the name given to the kind whose expansion and contraction is under our control. It is made up of cells that are very long in comparison with their width and are marked with transverse bands or striations. Such tissue forms the muscles of the legs, arms, back, etc. Nonstriated or *involuntary* muscle tissue is the name given to the form whose expansions and contraction are not under our direct control. It is made up of cells which are spindle shaped and without transverse striations. Such is the tissue which forms a part of the walls of our digestive organs.

Striated muscular tissue. An examination under the microscope (see Ex. XXVII. and Fig. 16) shows such

muscular tissue to consist of long fibers covered with a transparent covering or sarcolemma, within which may be made out several nuclei and two kinds of bands or *striæ*. These *striæ* appear as alternate broad, dim stripes and narrow, light stripes. Both these bands are made of protoplasm called *sarco-plasm* or muscle protoplasm. The numerous nuclei indicate that a fiber is a collection of several cells fused together with a common wall or sarcolemma.

Nonstriated muscular tissue. The

FIG. 17 — Muscle layer of the intestine (unstriated muscle cells); *n*, nucleus of a single cell.

muscle layers of the intestine (see Ex. XXVII. and Fig. 17)

show us that this tissue consists of a mass of spindle-shaped cells each containing a single large nucleus surrounded by a mass of protoplasm which is striated longitudinally but not transversely. Around each cell is a thin wall or membrane similar to the sarcolemma.

Muscle cells then may be described as cells containing protoplasm, which is so arranged that when it expands and contracts the whole cell is shortened or extended. By massing cells of this nature together this shortening and extension is made to cause movement of the parts attached. Evidently the extent of motion is also increased by this massing, the total short-

FIG. 16 — Striated muscle fiber; *a*, termination of muscle substance; *n*, nuclei.

ening being the sum of the shortenings of the respective cells.

Nerve tissue. — In every human being there are two organs called the brain and the spinal cord. From these there extends all over the body a lacework of fibers which are called nerves. Brain, cord, and nerves are built of the

substance called nerve tissue, whose principal purpose seems to be to originate and conduct stimulations to various parts of the body. These stimuli act upon the other tissues of the body and cause them to either increase, or cease their special kind of activity. In this way is controlled the action of the different organs that are made up of tissues. Nervous tissue is made up of cells just as are all other tissues. These cells are, however, peculiarly arranged and extended so as to be continually in contact. For example, while the muscles of the foot are entirely distinct

FIG. 18 — Nerve cell, showing nucleus in the center of the cell, and having one axis cylinder or *nerve process* and several *dendrites*.

and separated from those of the hand, the nerve cells of the hand are, on the contrary, connected with those of the foot and other parts of the body, in the same way that telegraph stations are connected. In other words, if we could take away all the other tissues of the body we should have left the nervous tissue in the form of an intricate network,

all parts of which were connected. It is this connection that allows us to become conscious of an itching sensation in the foot, and permits us to direct the muscles of the hand to rub the affected part. The manner of the connection of these cells is peculiar to this tissue, and its method we shall investigate further in a later chapter.

Nerve cells.—As shown under the microscope, a nerve cell consists (see Ex. XXVIII. and Fig. 18) of a mass of protoplasm with a nucleus. The protoplasm is drawn out into two forms of projections, *nerve processes* and *dendrites*. The rodlike nerve processes (*axis cylinders*) form the center of a fiber called a *nerve* and this latter may be of great length. These nerves may branch extensively, and through its nerve process permit a nerve cell located in the brain or cord to be in contact with widely distant points, say in the finger or foot. By means of the dendrites, one cell and its nerve may be brought in contact with another cell and nerve, and thus the entire set of cells brought into communication. A nerve cell with its processes and dendrites is called a *neuron*. (See Fig. 19.) Just as the protoplasm of a muscle cell has the power of expansion and contraction developed to a great degree, so the protoplasm of a 'nerve cell has developed the power of conducting impulses of various kinds.

FIG. 19 — Neuron; *np*, nerve process; *n*, nerve cell; *t*, terminal brush; *c*, *d*, nerve fiber with sheaths surrounding the axis cylinder.

In our study of the nervous system we shall investigate further this conductive power and the action of nerve tissue.'

GENERAL PROPERTIES OF TISSUES.

There are two general facts to be learned from the study of the structure of tissue. First, that all parts of our bodies are made up of tiny structural units or cells. Second, that the functions of a given form of tissue are due to the predominance of at least one of the properties of the protoplasm of its cells over the other properties. Thus, the protoplasm of the bone cell, like that of the muscle cell, has the power of absorbing food, building up new protoplasm, giving off wastes, and forming new cells. In these respects bone cells are like all other cells. They differ in that the protoplasm of the bone cell has developed its power of giving off waste matter over all the other properties, while the muscle cell has made a specialty of the power of contraction and expansion. The result is that a combination of bone cells becomes important in the body for the character of material it throws off, namely, bone, while the combination of muscle cells is equally important from the power of motion which it makes possible. A comparison may make this point clearer. Take two men, one a musician, one a carpenter. Each eats, moves, thinks, and performs all the ordinary actions of the other. Both have all the properties of a man. One, however, has trained his hands to one kind of action, the other, to a different kind, and to this specialization are due their respective callings. So tissues are but collections of cells whose essential substance is protoplasm.

All these cells possess the common attributes of protoplasm. The specialization of one of these properties: contraction, excretion, or conduction, makes them different, and gives its special tissue name to the collection of which these cells are a part.

What happens when we combine bodies of men of different specialties? Take a factory, for example. If it is a cotton mill, some are carders, some spinners, some weavers, and the like. Each body performs its own special work, and all together produce the common result, the production of cotton cloth. So in our bodies. The various organs when dissected are found to be composed of different tissues; that is, collections of specialized cells, which work together for some common result, such as seeing, digestion, smelling, and feeling. We may carry the comparison further. Just as different factories supply the needs of a community, so in the body we find organs combined into systems, each organ contributing its action to a common purpose. Thus the stomach, intestines, liver, etc., are organs which by their combined action prepare the food for the body, and hence we group them under the name of the digestive system. Finally, all these systems are placed under the control of a governing board in the form of the nervous system, and so their work is coördinated, and all work for the maintenance of the individual which they make.

The individual man consists of a collection of systems, organs, tissues, cells, and the total actions of the individual are but the sum of the actions of his individual cells. This division of the work of the body among different specialized cells is called *physiological division of labor*. The significance of this term lies in the fact that

each set of cells performs special work just as in a cotton mill, spinners, weavers, and others have their special tasks.

Reproduction of cells. — Before we leave the tissues, one feature of cell action which they all share must be considered more fully. Like the parts of a machine, the parts of our body are continually being broken down by use. To remedy this there must be continual repair or building up. Otherwise the body, like the machine, soon becomes

FIG. 20—Reproduction of a cell; *A*, diagram of a cell, showing centrosome (*a*), and nucleus (*n*) consisting of nuclear membrane inclosing granular substances in which are a spherical nucleolus and irregular masses of chromatin; *B* to *F*, changes in the course of reproduction.

worn out and useless. How is this brought about? The answer to this question is found in the discovery of a very peculiar property of protoplasm called reproduction. This means that an old cell has the power to produce new cells out of its protoplasm. The manner in which it does this is illustrated in Figure 20; thus a single cell or egg is able by repeated divisions to produce all the tissues of a body, and a tissue is able to renew its structure to maintain its efficiency. When the protoplasm of an individual loses this power the individual soon becomes worn out and dies. Life depends upon the reproductive power of protoplasm.

V. FOODS AND NUTRITION.

ONE of the earliest sensations that a child experiences is hunger or the desire for food. A child comes early to recognize that this hunger is merely a way the body has of telling us that it must have food at certain intervals. Further experience teaches that this food must be of a certain kind to enable the body to grow and maintain itself in a state of health and activity. An examination of other animals and of plants teaches that they too require food and that without food they die.

The foods of the different plants and animals differ greatly, however, and what supports life in one does not necessarily answer the same purpose in another. For example, the hay on which a horse feeds would be useless to us, and the fertilizer and water on which a plant flourishes would prove valueless to both the horse and ourselves. How then shall we define food as a general term which shall include all classes? Broadly, we may define it as follows: *Anything which an animal or plant takes into its body as nourishment is food.* Applying this definition to ourselves, food is anything which we eat or drink that nourishes our body. The number of such substances is very great. Chemists have analyzed all the various forms of food, however, and have found that they are composed of one or more of some six classes of compounds. These compounds then are the structural units of which all the varied forms of foods are built up. These

fundamental food units are spoken of as food stuffs, alimentary principles, or nutrients. We shall adopt the last term in speaking of them.

Nutrients. — Nutrients are usually classified under the following names: water, inorganic salts, proteids, carbohydrates, fats, and albuminoids. Since the importance of a food and its value as nourishment depends upon the amounts and kinds of these nutrients present, it will be well to consider some of their properties and the means by which their presence may be detected.

Proteids or nitrogenous compounds. — Different varieties and forms of proteid are found in nearly all kinds of animal and vegetable foods. The exact composition of proteid is unknown, but the elements of which it is built are nitrogen, carbon, sulphur, hydrogen, and oxygen. It occurs in a great variety of forms which receive different names; for example, myosin and syntonin in meats; egg albumin; casein of milk and cheese; gluten of grain. White of egg or egg albumin illustrates well the characteristic properties of proteids.

When egg albumin is heated or treated with acids (see Ex. XIII.), it is changed from a soluble, sticky liquid to a white solid insoluble in water. This change to a solid form, due to the heat or acids, is called coagulation, and is a characteristic of many proteids. The solidifying of a piece of meat under heat is due to the coagulation of the proteid in it by the heat, while the curdling of milk when sour is another case of coagulating proteid due to the presence of lactic acid in the milk.

There are a number of ways in which the presence of proteid in a food may be detected. The principal chemical tests are as follows (see Ex. XIII, *a-d.*):

(a) *Xanthoproteic test.* Proteid boiled in nitric acid is dissolved and the solution colored yellow by the action of the acid. If to this mixture is added enough ammonia to neutralize the acid the color turns a deep orange.

(b) *Millon's test.* Proteid, brought slowly to boiling with a solution of mercury in nitric acid, becomes rose red. The nitric acid solution of the mercury is called *Millon's reagent*.

(c) *Biuret test.* If a cold solution of proteid is first made alkaline, by the addition of caustic soda, and then a few drops of a very dilute solution of copper sulphate are added, a violet color appears which deepens in color with boiling.

Any one of these tests applied to a food which contains proteid will reveal its presence. Of the three, the xanthoproteic is best for common use.

Albuminoids. — These substances, an example of which is the gelatine found in bones, are closely allied to proteids in composition. This resemblance to proteid is confined to their composition and the presence of nitrogen as they vary from proteid, not only in failing to respond to proteid tests, but also in their action. In this group is included keratin which forms horn, hair, feathers, nails, and hoofs; elastin, which forms the elastic fibers of connective tissue; and certain bodies called enzymes which we shall investigate later. The different forms vary so much in characteristics that it is impossible to name any one feature that is distinctive. Of the common foods, soups and soup stock made from bone marrow are richest in this nutrient.

Carbohydrates. — Under this name are included the starches and sugars. These are compounds of carbon,

hydrogen, and oxygen, so combined that the hydrogen and oxygen are always in the same proportion as they occur in water.

Starch is found in all green plants in the form of granules and is frequently stored in the cells of grains, seeds, and stems, as a reserve food supply for the plant. To two of these plant storehouses, the potato and the corn, commerce is indebted for its market supply of the substance. The principal characteristics of starch are its insolubility in water, and its transformation to a sticky swollen mass or paste by the addition of boiling water. It gives a blue color when treated with a solution of iodine, and this latter means is usually employed as a test to detect its presence in foods. (See Ex. XIV.)

There are several different kinds of sugars; cane sugar or *saccharose* is the sugar of the market, and as its name indicates is obtained from the juice of the sugar cane. It is also obtained from the juice of the sugar beet. Grape sugar or *dextrose* (also called *glucose*) is found in almost all plant and animal tissues, and it derives its name of grape sugar from its abundance in grapes. Still another form is milk sugar or *lactose*, found in milk. Both *saccharose* and *lactose* can be changed to *dextrose* by boiling with certain acids, notably, hydrochloric. This process is called inversion.

Grape sugar is characterised by two important properties, the power to ferment and the power to take up oxygen. As a result of the first, certain substances, such as yeast, are capable of breaking it up into alcohol and carbonic acid gas; while to the second is due the means of detecting its presence in foods. The most common test used is a mixture called Fehling's solution. (See Ex. XV.)

When a solution containing grape sugar is treated with a few drops of this solution, a substance is precipitated, which, upon heating, gives up part of its oxygen to the sugar, and turns a reddish brown. If no sugar is present this loss of oxygen does not occur and there is no consequent colored precipitate. This reaction furnishes a simple and delicate means of detecting the presence of grape sugar in a compound.

Fats. — Under this name are included not only what is ordinarily understood by the term fat, but also animal and vegetable oils of all kinds. Like the carbohydrates, fats are composed of carbon, hydrogen, and oxygen, but in different proportions. They are all lighter than water and insoluble in it. All are readily made liquid by a moderate amount of heat, while some are liquid at ordinary temperatures. Thus body fat melts at a temperature of 98° Fahrenheit and is liquid in the living body. (See Ex. XVI. *a.*) Fats are soluble in ethers and benzene and when heated with alkalis form soap. Their presence may be detected in one of two ways. (See Ex. XVI., *b, c.*)

(*a*) Rub on paper the substance to be tested or place it on the paper and heat gently. If fat is present a grease spot will result, which gives the paper a translucent appearance and is readily detected.

(*b*) Soak the substance in ether (benzene will do as well). The ether will dissolve the fat if it is present. Then filter off the solution and allow the ether to evaporate. The fat does not evaporate and will remain. This second test enables us to extract and measure the exact quantity of fat present as well as to determine its presence.

Mineral salts. — Many forms of salts occur dissolved in water and are present in all living matter, while other insol-

uble forms enter into the structure of important parts of the body. For example, in the fluids of the body we find common salt (chloride of sodium) as well as chlorides of potassium, ammonium, etc., while important parts of our bones are the insoluble carbonates and phosphates of calcium. Mineral salts all agree in being unburnable, and it is by burning foods and examining the character and amount of the unburned residue that we are enabled to determine the kinds and amounts of salts present in a given food. (See Ex. XVII.)

Water. — This liquid forms a large proportion of all fresh vegetable and plant foods. It constitutes nearly 59% of the weight of our bodies. This latter fact indicates the great importance of water as a nutrient. The fact that water becomes a vapor and passes off in that form into the air when heated, enables us by drying a substance in a warm place to determine its water content. (See Ex. XVIII.) The loss of weight represents the water evaporated.

FIG. 21 — Apparatus arranged for nutrient experiment with a seedling.

COMPOSITION OF FOODS.

The extent to which these nutrients occur in various forms of foods (see also Ex. XX.) is indicated in Chart I., on page 57.

Sources of nutrients. — An examination of our ordinary food supply reveals the fact that for most of our nutrients we are dependent ultimately upon the vegetable world. Our food, in general, is animal, vegetable, or mineral, in

CHART I.

Food	COMPOSITION % OF NUTRIENTS						Energy in Calories per Pound	Average Cost per Pound
	Proteid	Starch	Other Carbo- hydrate	Fat	Water	Mineral		
Bread (White)	8.	47.	3.	1.	37.	2.	1280.	\$.04
Flour	11.	66.	4.2	2.	15.	1.7	1645.	.025
Oatmeal.	12.6	58.	5.4	5.6	15.	3.	1850.	.05
Rice	6.	79.	0.4	0.7	13.	0.5	1630.	.07
Beans	23.1	55.	2.	2.	12.6	3.1	1615.	.05
Potatoes	2.	18.	3.	0.2	76.	0.7	375.	.0125
Milk	4.	...	5.0	4.0	86.	0.8	325.	.035
Cheese	28.3	...	1.8	35.5	30.2	4.2	2070.	.16
Beef (Round)	20.5	...	...	10.1	68.2	1.2	805.	.14
Beef (Corned Flank)	14.2	...	...	33.	49.8	3.	1655.	.10
Mutton (Leg)	18.3	...	...	19.	61.8	0.9	1140.	.18
Veal (Shoulder)	20.2	...	...	9.8	68.8	1.2	790.	.20
Pork (Shoulder — fresh)	16.	...	...	32.8	50.3	0.9	1680.	.16
Pork (Ham)	16.7	...	...	39.1	41.5	2.7	1960.	.16
Pork (Salt Fat)	0.9	...	...	82.8	12.1	4.2	3510.	.12
Chicken	24.4	...	...	2.0	72.2	1.4	540.	.20
Eggs	14.9	...	...	10.5	73.8	0.8	721.	.18
Butter	1.	...	0.5	85.	10.5	0.3	3615.	.30
Cod fish	15.8	...	...	0.4	82.6	1.2	310.	.08
Mackerel	18.2	...	...	7.1	73.4	1.3	640.	.12
Oysters	6.	...	3.7	1.2	87.1	12.	230.	.25

origin, but if we consider for a moment the kinds of animals upon which we depend for food it is noteworthy that these animals themselves feed exclusively upon vegetable and mineral matter. We may say, then, that with the exception of some mineral salts and water, the animal world in general ultimately derives all its supply of nutrients from the vegetable world. When we eat flesh we are simply using the proteid, carbohydrate, and fat that these animals have themselves obtained from vegetable matter. It is in the plant, then, that the chemical compounds

which we call nutrients, are produced from the elements or their inorganic compounds. A few simple experiments with a seedling (see Ex. XIX.) demonstrate that a plant grows well when its diet is restricted to water, certain mineral salts, and carbon dioxide. The roots take up the salts and water as solutions, while the leaves absorb the carbon dioxide from the air. When analyzed, the plant thus restricted in diet is found to have produced proteid, starch, sugars, and oils. Plant physiologists tell us that the simple raw materials are broken up and recombined chemically into what we call the nutrients, and that a plant

is actually a sort of manufacturing chemical laboratory where these reassemblings of elements are continually taking

A

B

FIG. 22 — *A*, diagram illustrating the manufacture of nutrients by the plant; *B*, cycle illustrating the relation of living forms and the mineral world in regard to nutrient supply.

place. Some of this manufactured nutrient the plant uses for its own food and some it stores. The stored nutrients are in turn taken in by the animals, and by them

converted into animal proteid, and carbohydrate. If now we note what happens when animal bodies and plant bodies die, we find that in the process called decay these nutrients are again broken down into mineral salts and gases and thus complete a cycle. The plant world therefore stands between the mineral and the animal world as a manufacturer of nutrients.

Uses of nutrients. — Very careful experiments have been performed in the past and 'are being performed at the present time to determine the exact use of each kind of nutrient to the body. What is already known in this direction may be summarized as follows:

Nutrients serve two purposes in the body. First, they furnish the material out of which our tissues are built and repaired. Second, they may be oxidized in the body, and when this action occurs the result is the production of either action (movement) or heat; both action and heat being forms of *energy*. These two uses are usually expressed by saying that nutrients furnish the body with either building material or fuel.

Building material. — All the organs and tissues of the body are continually undergoing two processes, a building up and a tearing down. These two processes are spoken of collectively as *metabolism*. The building up is called *anabolism*, and the tearing down *katabolism*. Anabolism and katabolism vary in amount at different periods of our life. In childhood the anabolism is greater than katabolism and the body increases in size and weight, that is, it grows. In middle age the two processes are nearly equal in amount so that the new material merely serves to repair or replace the worn out parts. Finally, in old age, katabolism becomes the greater, the new material

fails to replace completely the worn out parts, the body loses in power, and finally its action ceases altogether, and what we call death follows. To furnish new material for anabolism is what is meant by the use of nutrients as building materials.

If we recall that tissues are composed of cells, and cells of protoplasm, we shall see that this function of nutrients really resolves itself into the building up and repair of broken down protoplasm. The particular nutrients that are best suited to use as building material are proteids or the nitrogen compounds. The fact that proteid and protoplasm are nearly identical in composition has a new significance in the light of the above facts. How far the other forms of nutrients share in this use as building material, is at present uncertain. In fact, experiment has shown that food free from inorganic salts will not support life, thus demonstrating that these substances are an essential part of the tissue structure.

Fuel. — Whether carbohydrates and fats are ever used as building material or not, this is not their principal function in the body. Their main purpose is to serve as fuel, and thereby to produce what we call energy, or the power to do work.

Meaning of energy. — Whenever our muscles move parts of the body, as in running, leaping, walking, and eating, we have examples of this power to do work, or energy. Whence comes this power? An example may help us to understand better the nature and production of energy. An energy producing machine with which we are all familiar is the steam engine. Such an engine derives its power to do things from the coal which we put in the fire-box. Here this fuel burns (that is, oxidizes, see p. 18),

and the result is the production of a form of energy which we call heat. This heat in turn acts upon water inclosed in a boiler, and changes it to steam. Steam has the power of expansion; that is to say, the heat energy has been transformed into expansive power, another form of energy. If the steam is admitted to the cylinder of the engine and allowed to expand against the piston, its form of energy (expansion) is transformed into still another form, thrusting movement. Further, if we attach the piston to a wheel this form of energy is changed once more into rotary movement, and that in turn may be applied to a saw, and thus the engine may be made to saw wood. In all these steps the process has simply consisted in transforming the energy derived from the burning fuel into first one form and then another form of energy. In other words, the steam engine is simply an apparatus for changing the form of energy derived from combustion. We may summarize what this example tells us as follows:

First. In the coal we see that energy may exist without manifesting itself. Coal, in other words, has the power to burn and produce heat even when it is not burning. This power becomes manifest as soon as the coal actually begins to burn or oxidize. We express this fact by saying that energy is of two forms, potential energy and kinetic energy. Energy is potential when, owing to its position or composition, a body possesses the power to do work without manifesting that power. When the energy manifests itself in the actual doing of work we have an example of kinetic energy.

Second. Potential energy may be transformed into kinetic energy, and kinetic energy into potential energy. Thus the potential energy of the coal to produce heat is

transformed into kinetic energy or actual heating when the coal is burned. Further, heat is transformed into potential energy or the power to expand, that steam has when it is not actually expanding.

Third. Kinetic energy can be transformed from one kind of kinetic energy into another. Thus heat is transformed into expansion, expansion into thrusting, thrusting into rotation, rotation into sawing, etc. Careful experiment has demonstrated that when energy is transformed from one form to another no energy is lost or created. The law which expresses this fact is called the law of the *conservation of energy*, and applies to all cases where the production of energy is concerned. It is usually stated as follows: Energy can be transformed from one kind to another, but can never be increased, diminished, destroyed, or created from nothing.

Fourth. The source of energy in a steam engine is the oxidation of fuel. That is to say, the energy of an engine is derived fundamentally from the oxidation of certain chemical compounds. Such oxidation is simply a chemical change, and chemical changes are the sources of most forms of energy.

In the light of this example let us examine again our body as a machine for the transformation of energy. Just as in the steam engine, we find that its source of energy is to be found in the oxidation of fuel. Such fuel is found in the carbon compounds known as carbohydrates and fats which form part of our food. The energy liberated by the oxidation of these compounds in the body takes many forms, such as heat, muscular activity, secretion, etc., and one form may be transformed into another, since the law of conservation holds here just as every-

where else. Thus the energy of a sleeping child is potential and becomes changed to various forms of kinetic energy when he wakes. Finally, all energy liberated in the body ultimately manifests itself either in action or heat, and hence we may say that our actions and our body temperature are merely manifestations of energy whose source is the oxidation of the nutrients, carbohydrate and fat. Since some part of our body is always in action, and all parts must be kept at a constant temperature, it follows that there must always be on hand a sufficient quantity of fuel to supply this energy. It is this fact that makes it so important that our daily diet contain a certain proportion of carbohydrate and fat, and explains why people whose work is largely muscular require more of these nutrients than those whose labors are less active.

Other substances than fats and carbohydrates may be used as fuel, notably both proteids and albuminoids, as both contain oxidizable carbon. The carbohydrates and fats, however, are the cheapest and easiest sources of this energy. It is interesting to note that the carbohydrates are capable of transformation into fat, and that it is in the latter state that the reserve energy of the body is stored. In starvation it is this stored fat that maintains life.

Another striking feature that can be deduced from the above statements is that *proteid is the only universal food*, since it is the only nutrient that combines in itself both tissue building and fuel qualities.

Value of common foods. — A study of Chart II., shows us that an important factor in determining the value of a food is its composition. A food like meat, for example, owing to its high percentage of proteid, is at once recog-

nized to be a valuable tissue builder. Potatoes with their abundant starch are in the same way put down as fuel or energy producers. If we consider our own experience, however, we realize that composition tells us only a little of the true value of a food. A food, for example, may contain all the nutrients necessary to the body, and yet be absolutely valueless to us. Hay, for instance, contains all the nutrients necessary to support life, and yet man cannot use it as food. Again, certain foods may be so rare that their cost prohibits their use. In general, then, we may say that the value of a food is determined only when we have considered its composition, its fitness for our use, and its cost.

The first point that may be noted from Chart II. is that most food as bought in the market is made up of nutrient only in part. The remainder may consist of bone, connective tissue, skin, bran, etc., all of which is included under the heading of refuse in the chart. The amount of refuse present is evidently an important factor as regards the value of a food. It is because of the fact that eggs, milk, and bread contain almost no refuse that they are such valuable foods.

A second point to be noted is the water content of foods. In this respect, animal foods, in general, contain more water than the vegetable foods. Exceptions to this rule are found in potatoes, onions, lettuce, and some other vegetables, in which the proportion of water is very high. In meat foods, too, the fatty forms contain less water than the lean meats. In general, other things being equal, the higher the proportion of water the less the food value. Fish on this account are less valuable as food than ordinary meats.

CHART II. — (From Farmers' Bulletin Number 23).

A calorie equals the amount of heat necessary to raise the temperature of a kilogram of water (2.2 lbs.) from 0 to 1° Centigrade, i.e. 1½° Fahrenheit. It is possible to determine the amount of heat energy that a given amount of any nutrient can produce in terms of these calories, and hence they furnish convenient units to express the fuel value or energy producing power of foods.

Again, the fuel value of a food is largely a matter of its fat and carbohydrate content, though the fact that proteid has a fuel value makes it a factor. Foods rich in these substances take high rank as energy producers. It is interesting to note that fats have about two and a quarter times the energy producing power of either carbohydrate or proteid. Foods, therefore, which are rich in fats and poor in water have the greatest fuel value. Meats differ greatly in this respect, and while the different cuts and kinds necessarily vary, veal is distinctly the poorest and pork the richest in fat, while mutton and lamb rank with the fatter cuts of beef. Chickens and turkeys rank low in fuel value but high in proteid in spite of their large proportion of refuse. Vegetable foods, though rich in carbohydrates, have less fuel value than fat meats.

A comparison of vegetable and animal foods from the chart is interesting, and the importance of dry seeds and grains with their small proportion of water and high proportion of nutrient is worthy of note. In most of such vegetable foods the proportion of nutrient is not far from 87%. Beans and peas are richest in tissue building material, cereals next richest, and fresh vegetables least rich. Vegetable foods, therefore, are good tissue builders, but lack the fuel value of meats owing to the small proportion of fat. They are in general less rich in proteid than meats. In this comparison the position of bread as a nearly ideal food is interesting. The flour or vegetable part with its large proportion of proteid and little water has its lack of fat and mineral matter remedied by the addition of lard or butter, milk, and salt, and thus combines the advantages of the cereal with that of the fat meat.

Fitness for use. — So many factors enter into this determination that it is difficult to generalize them. The presence of the necessary organs for digestion, sex, age, size, occupation, likes and dislikes, amounts necessary for nourishment, and conditions of the nutrients all bear on this question. If we leave out of account all personal peculiarities the most important of these factors may be comprised under the head of digestibility. We shall discuss digestion more in detail in a later chapter. For the present it is sufficient to state that by digestibility we mean the relative amount of a food that in a given time our body is able to put into tissue building or energy producing form. Without attempting to detail the methods by means of which digestibility is determined, we may state the general conclusions of such experiments as follows:

First. Vegetable proteid is much less readily digested than animal proteid. That is to say, animal foods will furnish us with more energy and tissue in a given length of time than will vegetable foods.

Second. The amount of fat present in a given animal food does not indicate exactly its energy producing power, since, as ordinarily eaten, nearly five per cent of such fat escapes digestion, fat requiring a long time for its digestion.

Third. Carbohydrates are ordinarily very completely digestible and hence foods rich in these nutrients become important energy producers even though their caloric value is less than that of fat.

Fourth. The amount of food digested in a given time (its digestibility) is not affected to the extent commonly supposed, by the addition of flavors or condiments, nor does the time differ much with people of different habits. The statements made above are, of course, general, and

are all subject to modification when we consider them in relation to individual tastes and needs. Thus certain forms of food may be distasteful to certain persons. In that case a less nutritious food may be more valuable because of its readier digestibility.

Cost of foods. — The money value of foods is a question that should have much more attention than is commonly given to it, and much time has been given to this side of food values by the Department of Agriculture at Washington. The table on the following page gives some of these results in a form which explains itself.

The principles which should guide in the selection of cheap foods may be stated as follows: "The cheapest food is that which supplies the most nutriment for the least money. The most economical food is that which is cheapest and at the same time best adapted to the wants of the eater."¹

These two statements indicate very clearly two of the guiding principles for the economical purchase of food, namely, nutritive value and palatableness. It is evident that only by the study of such tabulated statements as are given in Charts I. to III. can we determine the relative nutritive values of foods, and it is in supplying of such tables that the government is giving valuable aid to the purchaser.

Adulteration. — Another principle which the purchaser must consider is the quality or relative purity of foods. Especial care is necessary in the selection of brands of canned foods since many of the market brands contain adulterants or preservatives which are positively dangerous to the health of the user. Recent legislative measures

¹ Farmers' Bulletin, 23, p. 20.

CHART III. — Pecuniary Economy of Food.

(From Farmer's Bulletin Number 23.)

in the passing of Pure Food Laws are steps in the direction of safe-guarding the public in this respect. The refusal of the public to purchase any forms of foods except those which inspection proves to be of good quality will go far toward remedying the present tendency to adulterate.

Daily diet. — Observation experiments conducted in Europe and in the United States have given us figures which serve as standards in determining the amount of foods which must be consumed daily by people of almost every class in order to preserve good health. In Europe the figures given by Professor Voit of Munich, Germany, are considered most reliable. Those of Professor Atwater are considered better suited to conditions in our own country. The following table indicates the amounts determined by Professor Atwater:

Conditions	Proteid, Lbs.	Carbo- hydrate, Lbs.	Fat, Lbs.	Calories
Man with light muscular exercise	.22	.88	.22	2989.
Man with moderate work. . . .	.28	.99	.28	3570.
Man with active muscular work	.33	1.10	.33	4060.

By comparison with this table and the one on page 65, daily dietaries may be made up to fit all classes and at the same time secure variety of diet. (See Ex. XXI.)

Cooking. — Experience teaches us all the importance of cooking. In general, its scientific value lies in the fact that it either renders food more palatable or more digestible.

Methods of cooking meats. Meats are usually roasted,

boiled, broiled, or fried. Of these methods frying is the poorest as it adds too much fat and tends to make the meat less digestible. The advantage of roasting or broiling, over boiling, is that both of these former methods tend to retain the juices and thus prevent loss of nutrient and flavoring substances. If meat must be boiled, it should be plunged into boiling water at first, so as to coagulate the surface proteid, and thus keep in the juices. This, too, is the reason for a hot fire in broiling and roasting.

On the contrary soups obtain their value from the juices extracted from the meat and bones and hence the reverse process is necessary here. In this case the meat should be put in cold water and the temperature raised gradually. Very little nutrient is found in soups, and hence stews, which make use of the meat as well as the broth, are much better foods.

The value of meat cooking may be summarized as follows: (a) Heat kills bacteria and parasites, such as tape worms, which may be present, and thus makes the food safer for use. (b) It renders the meat more attractive and improves its flavor. (c) It renders the fibers softer and makes the digestion simpler for the organs. (d) It makes the time required for digestion shorter and thus enables the organs to digest more nutrient in a given time.

Vegetable cooking. The principal reason for cooking vegetables is found in the fact that it renders them more readily digestible and improves the flavor. Starch as found in raw vegetables is inclosed in grains which have a hard indigestible outer shell. Heat causes the starch to swell and burst these walls and thus enables the digestive fluids to get at it.

In boiling vegetables, care must be taken to prevent the soaking up of water, as this increases the proportion of water. For this reason the cooking should be as rapid and with as little water as possible. On this account steaming is preferable to immersion in water.

VI. FOOD ACCESSORIES; ALCOHOL AND TOBACCO.

IN ordinary diet we eat many things which merely add to the enjoyment of eating. These substances are to be regarded not as foods but as food accessories. They include the condiments (pepper, mustard, etc.), the flavors, the stimulants (tea, coffee, etc.), and the narcotics (tobacco and opium). Some of these substances are harmless, some are beneficial, and others positively injurious. Some are harmless in small quantities, but become distinctly injurious in larger amounts. None of these food accessories is absolutely necessary to the body, though few people are willing to eat food entirely without flavor.

Condiments and flavors.—Most of these substances have a directly beneficial effect either in making the food more palatable or in stimulating the flow of the fluids which digest food. Some hasten the absorption of the food into the blood. They are harmless if not used to excess and not used to disguise foods whose natural flavors would indicate their unfitness for use. A few, such as the salts and the sugars, have a food value also. Used in excess, they may result in overstimulation of the digestive fluids or the cultivation of habits of use which destroy the normal appetite. The continual use of highly seasoned food destroys the appetite for natural flavors.

The members of this group may be classified as follows: aromatic condiments (vanilla, cinnamon, nutmeg, etc.), peppers, alliaceous condiments (garlic, onion juice, mus-

tard, etc.), acid condiments (vinegar, lemon juice, pickles, etc.), the salts, and sugars.

Stimulants. — This is the name applied to substances which excite cells to action without furnishing them material with which to develop heat and energy. They produce their effect through action upon the nervous system. Of all the substances included in this group it may be said that, while in small amounts they are harmless and may even be beneficial, in excess they are dangerous to the best health of the body. The main reason for their injurious action lies in the fact that while a little stimulation is sometimes necessary, an excess soon causes a weakening of the powers of the cells affected, and leaves the body in a worse state than before the use of the stimulants. In this respect, their action is like the application of the whip to a horse. A little use may produce good results, but excess soon exhausts the powers of the animal. It is generally agreed by all health authorities that the use of stimulants should be avoided by the growing child.

Tea, coffee, and cocoa or chocolate owe their stimulating effect to certain principles which they contain (cafein in tea and coffee, theobromin in cocoa and chocolate). These principles cause an increase in the blood flow, the tired cells are supplied with more food from the blood, and the sense of fatigue disappears for the time. While such stimulation may be necessary at times for the adult, it is always better to secure the stimulation through natural means such as exercise or by allowing the cells to rest. Beef extracts were formerly supposed to contain much nourishment. It is now known that their sole claim to value lies in their action as a stimulant. In cases

of sickness they furnish the best amount of stimulation with the least amount of harm, but must never be considered as a substitute for nutriment.

ALCOHOL.

Relation of the study of alcohol to physiology. — Owing to the fact that the use of alcohol is so widespread among men, and that this use has been so productive of evil, many societies have been formed throughout the country to fight this tendency. One of the objective points of the work of these societies has been to inform the public of the effects of alcohol upon the human system. To this end, through legislation, they have made the study of the effects of alcohol a required part of the course in physiology in the public schools of the country.

Since the science of physiology has as its *sole* object the explanation of life processes, so far as the most careful experiments of competent investigators have made such explanation possible, we shall consider the effects of alcohol upon the human system from the following view points only:

(a) Its proper position among the substances which affect the action of the body organs.

(b) The statement of the effects, direct or indirect, which its use has upon the action or structure of the organs of the body.

In the present chapter we shall consider alcohol mainly from the first view point. The statements concerning these effects represent the views of highest authorities.

Position of alcohol. — Much time has been spent in trying to answer the question as to whether alcohol is a food

or a poison. The final answer to this question must rest upon experiment, and experiments have been conducted in the past and are still being conducted to establish the true answer to this question. The results of the past experiments and the questions involved are outlined in the following paragraphs:

A poison is defined by the "International Dictionary" as "any agent which, when introduced into the animal organism, is capable of producing a morbid, noxious or deadly effect upon it."

A food we have defined (see p. 51) as "anything which we eat or drink which nourishes the body." That is to say, anything from which tissue may be built, or energy produced *without harm to the tissues* of a person in a healthy condition.

The experiments of reliable investigators have established the following facts concerning this question:

First. Small quantities of grain alcohol (the amount varies with the individual) may be completely oxidized in the body, and by such oxidation energy is produced exactly similar in kind and effect to that produced by the oxidation of fats and carbohydrates.

Second. Small amounts of alcohol may be substituted for equivalent amounts of fats and carbohydrates in the diet without direct harm to the tissues, and when so substituted they protect the other nutrients of the body in practically the same way as do the nutrients for which they are substituted.

Third. Small amounts of alcohol are absorbed directly into the blood, that is, require no preparation for absorption.

Fourth. In large amounts alcohol acts as a drug, is always a poison, and causes direct harm to the tissues.

Fifth. As ordinarily taken, alcohol is not intended as a food, but for its exhilarating effect upon the nervous system. Such use is always attended with the danger of forming habits of excess.

Sixth. Alcohol is not necessary to people in health, and leads readily to habits of excess. In excess its effects are far worse than excess of ordinary food.

Seventh. Compared with ordinary food it is very expensive. The amount of alcohol necessary to produce a given amount of energy costs several times as much as the equivalent amounts of fat and sugar in ordinary foods.

Eighth. In small amounts, it increases the flow of digestive fluids and may be beneficial in sickness or disease.

These results teach us that we may consider small amounts of alcohol as true fuel food, even though it never acts as a tissue builder. Further, its use in small amounts does not necessarily mean immediate damage to our systems, that is, does not injure the tissues directly. Under these conditions then it is not a poison, but a true fuel food. The results, however, point still more strongly to the fact, first, that its use is unnecessary, since other foods furnish the same amount of energy at a much less cost and at the same time are attended with none of the dangers of acquiring habits of excess; second, that this habit which alcohol encourages is a very real and great danger, and that on that account its general use is to be discouraged. When we indulge in too much candy we are made sick, and the sickness tends to destroy our craving for such excess. When we indulge in too much alcohol the effect is not only to poison the system but to increase the craving. It is the fearful effect of this habit and the excesses it encourages

that should make us avoid this unnecessary and costly food, and on this point all investigators agree.

In stating the position of alcohol, then, we must bear in mind that while it is definitely established that alcohol in small quantities must be considered as a fuel food, it is equally well established that in large amounts it is a poison; that while it induces a flow of digestive fluids, and thus hastens digestion, its use for such purposes in a healthy person is unnecessary; that large amounts stop the action of the juices and thus really delay digestion; and finally that its tendency to encourage habits of excess makes it an extremely dangerous substance to take into our systems except under competent medical advice. If both sides of the above statement are kept in mind it will be seen that in determining the true position of alcohol the investigators have taught us a very forcible argument for total abstinence.

In fact, most of the great evils resulting from the use of alcohol have their beginning in the fact that many people who begin by its moderate use feel no evil effects, whatever, as would be the case if alcohol were a true poison in small amounts. The use, however, soon becomes a habit, the habit demands ever increasing amounts or the stimulation is no longer felt, the increased amounts soon pass the danger point, and all the effects of alcohol poisoning, with its attendant moral evils, follow. We might also suppose that if the amount was always limited no evil results would follow. The fact that people in general use alcohol for its stimulating effects and not for food, interferes with the safety of this view. In other words, while it is probably true that a small amount taken daily would be oxidized and produce energy without harm, it is a peculiarity of

nerve affecting substances that unless the amount be regularly increased the nerves fail to respond after a time, and hence more must be taken, or the object sought fails of accomplishment, the taker must increase his amount or lose the pleasing effect on himself. In this way the moderate drinker is often unable to remain moderate. Further, the safe amount is not the same for all individuals. It varies with the individual, the time of taking, the physical health, the presence of other foods, and all these influences make it almost impossible to prescribe a safe amount for even single individuals except for limited times, as in sickness.

In fine, all that we learn of the value of alcohol as a food and its action on the nerves unites to make more clear the uselessness and danger of this substance to the ordinary healthy individual.

Comparison of alcohol with ordinary foods. — Like fats and carbohydrates, alcohol contains no nitrogen. In comparison with other foods, therefore, it is associated with the above named nutrients as a fuel food as distinct from proteid or tissue building food. A comparison with fats and carbohydrates shows many similarities and differences. Some of the most important follow.

Before ordinary food can be used by the tissues it must be changed by digestion. Alcohol is a fermented product, and the process by which it is made fits it to be directly absorbed by the blood and used by the tissues without change. A food of this form is said to be predigested. The speed with which this absorption takes place and the alcohol reaches the tissues is one of its most valuable properties as a medicine.

Fats and carbohydrates can be transformed into body

fat and stored in the body for future use. Alcohol is never stored for any length of time in the body, but is oxidized at once. On this account fats and carbohydrates can serve as sources of reserve energy, alcohol, never.

Compared with the other fuel foods, the amount of energy that a given amount of alcohol can produce when completely oxidized is shown in the following table:

	Alcohol	Proteid	Fat	Carbo- hydrate
Fuel value per pound in calories . . .	3130	1820	4040	1820

In other words, the complete oxidation of a given amount of alcohol produces 1.73 times as much energy as the same amount of proteid or carbohydrate and 0.73 as much as the same amount of fat.

In ordinary digestion not all of the fat, carbohydrate, and other nutrients which are taken in as food are actually used by the tissue. The proportion of any substance which actually reaches the tissues and is used by them is called the availability of that nutrient. The availability of the above four nutrients as actual energy producers for the tissues is as follows: alcohol, 98%; proteid, 70%; fat, 95%; carbohydrate, 92%. In other words, 98% of all the alcohol taken, up to the safe limit, is at once transformed into energy as against 70%, 92%, and 95% of a similar amount of proteid, carbohydrate, or fat.

One of the uses of ordinary nutrients is to preserve other nutrients from oxidation. For example, the carbohydrate which we eat keeps our body from using up the proteid and fat of the body for its energy supply, or the

fat eaten keeps the body from using up carbohydrate and proteid, etc. This protective action is important, as otherwise, in times of need, the tissues of the body might be consumed in the demand for energy unless so protected. It has been conclusively demonstrated¹ that alcohol shares this protecting power with fats and carbohydrates. It was not until this important fact was demonstrated that alcohol could be considered as a food, since, if this were not so, its use might permit the destruction of tissues, and then the result would be harm to the tissues in every case. Compared with the other fuel foods, alcohol protects fat better than carbohydrates, but carbohydrates and fats both protect proteid better than alcohol. The fact that carbohydrates and fats can both be stored and held in reserve for this very purpose gives them an advantage over alcohol in this respect.

All energy liberated by fuel foods is ultimately transformed in the body into either heat or muscular energy. The availability of alcohol as a heat producer is greater than either fats or carbohydrates, but its value as a producer of muscular energy is not yet fully determined. All experiments so far conducted indicate that its action as a drug tends to interfere with its power in this direction and make it far less valuable in this respect than either fat or carbohydrate. To put a concrete case, a laborer does not get from a given amount of alcohol anywhere near the power of continued muscular effort that he can from other fuel, owing to the fact that the drugging effects which first stimulate the cells, and so weaken their later

¹ See Sub-Report of Committee of Fifty, Vol. II., and Howell, *Text-Book of Physiology*, 1906, p. 808.

efforts, make the total muscular energy produced less than it would sometimes appear.

Alcohol as a drug. — The real value of ordinary foods depends not only upon their tissue building or energy producing power but also upon their effect upon the organs of the body. All nutrients affect the action of the organs more or less directly, and their effects may be beneficial or harmful according to the conditions or amounts taken. In small quantities, the effects of alcohol are not harmful but in larger quantities it acts as a drug. A drug is a substance which acts upon a special set of cells either stimulating or retarding their natural action. As a drug the effect of alcohol is ultimately to impede digestion, dilate the blood vessels, cause loss of heat, and destroy sensibility. In this respect it is like a powerful narcotic, and to this narcotic action is due the bad effects of large amounts. (See Narcotics, p. 86).

Alcoholic beverages. — Alcohol is produced by a process called fermentation (see p. 102) in fluids which contain starches and sugars. In this process these two carbohydrates (starch and sugar) are ultimately changed into carbon dioxide and alcohol. The form of sugar which is most readily changed into alcohol is what we have called glucose (see p. 54). This glucose occurs abundantly in the juices of many fruits, and especially in that of grapes. Such juices naturally furnish one of the chief sources of alcoholic drinks or beverages. Drinks made by the fermentation of such fruit juices are called wines, and while the term is frequently restricted to fermented grape juice, it actually includes other fermented juices, such as cider, blackberry wine, etc.

Grains, potatoes, etc., are rich in starch. If a little of a

grain, such as barley, is moistened and kept in a warm place it develops a substance called diastase, which changes the grain starch into glucose. Such moistened grain may be heated after the change in the starch has gone on to a certain point and the action stopped. When this is done, the dried product with its diastase is called malt. If a little of this malt is added to ground grain and the whole mixture moistened, the action begins again and the starch of the ground grain is in turn transformed into sugar. If yeast is added, its action causes the mixture to ferment, and the sugar becomes partly changed to alcohol. In this manner grains, potatoes, and other starch-containing substances may be used to make alcoholic beverages, and the liquids thus produced are called malt liquors. Beers, ales, and porter belong in this class of alcoholic beverages.

Finally, if we add yeast to the malted mixture of starch-containing cereals, potatoes, etc., until all the starch and sugar are turned to alcohol, and then distil this liquid to remove some of the water, we get liquors of much higher proportion of alcohol. Grain mixtures treated in this way and distilled until they contain nearly 40% of alcohol produce whiskey. Distillation of wine produces brandy. Gin is produced by flavoring the distilled liquid of malted grains with juniper berries, while rum is produced by distilling fermented molasses. All these liquors are rich in alcohol, and are grouped together as distilled liquors. The kind of grain or juice used and the flavors developed determine the brands.

We may distil these liquors still further, and remove all or nearly all the water and flavoring materials, and thus obtain the alcohol in the free state or with only a small

proportion of water. Commercial alcohol contains 95% of alcohol and 5% of water. Absolute alcohol is about 99.8% pure alcohol.

In speaking of alcohol in the preceding pages we have reference to what is known as grain alcohol, cologne spirits, or ethyl alcohol. There are many other forms of alcohol, notably among them, methyl alcohol or wood alcohol. This latter, however, has only a few of the properties of ethyl alcohol and is a poison even in small quantities.

In summary, then, we may state that alcoholic beverages may be classified as wines, malted liquors, or distilled liquors. The range of alcohol in these forms varies. Wines contain from 5 to 25%, malted liquors from 2 to 5%, distilled liquors from 40 to 50%. All contain various other ingredients which are produced from the fruits and grains in manufacture and are called on this account *extractives*.

In addition to these recognized alcoholic beverages there is a fourth class of compounds which are sold under the name of patent medicines. While it is true that many drugs are insoluble unless alcohol is used, these patent medicines often contain amounts of alcohol which are out of all proportion to their use as solvents. In fact, the exhilarating effects of many of these so-called cures are due solely to the large proportion of free alcohol present, and the taker is fooled into thinking that he is being cured, when his "cure" is actually alcoholic exhilaration. He would scorn to take a similar amount of alcohol in a wine. Investigation exposed so many of these so-called medicines that the government came to recognize their danger to the public health, and in the Pure Food Law is a provision requiring all medicines to bear a placard, on which must be stated the exact amount of alcohol contained. People

should make sure that their medicine list includes none of these liquors in disguise, since their use may fasten upon one a habit difficult to overcome and dangerous in the extreme.

Nutritive value of alcoholic beverages. — The action of alcoholic beverages is influenced by two factors, first, the alcohol contained; second, the presence of extractives. It is undoubtedly true that in certain cases the character of the extractives may add to the harmful effects of the beverage, but careful experiment has demonstrated that in ordinary liquors this action of the extractives may be neglected, and the harmful effect of the beverage may be stated to be directly proportional to the amount of alcohol which it contains.

Summary of conclusions concerning alcohol. — All investigators agree that while alcohol may be used in small amounts by certain individuals, without necessarily causing harm, the amount varies so much with the individual that it is impossible to fix a universally safe limit.

Because of its tendency to form habits of use, drinks containing alcohol are much more dangerous for young people to use than for adults.

Statistics from business corporations, life insurance companies, and insane asylums all show that the use of alcohol, except as prescribed in cases of sickness, is dangerous. In other words, while inefficiency, insanity, pauperism, and ill health may result from other causes, so marked is the tendency of the use of alcohol to result in one or more of these evils that its use is recognized in the business world as a risk. For example, all corporations employing laborers guard against it by stringent rules, and life insurance companies increase the premiums on life insurance policies

for moderate drinkers and absolutely refuse to insure habitual toppers.

It is a very expensive form of fuel food, and all the energy which its use supplies can be secured with less expense and no danger from recognized foods.

Finally, its transition from a food in small amounts to a poison in larger amounts is due to its action as a narcotic drug.

NARCOTICS.

In the paragraphs on alcohol we have mentioned the fact that alcohol in large amounts is a narcotic. Whether it is a stimulant or a narcotic in small amounts is at present a disputed point (see p. 426). Narcotics is the class name applied to substances which in small doses relieve pain and produce a state of rest or sleep, but in larger amounts may produce stupor, convulsions, and even death. The sense of fatigue and pain is the method employed by the cells of our bodies to inform us of their need for rest, or relief from disturbing influences. Sometimes, as in cases of surgical operations or painful diseases, it becomes necessary to use narcotics to lessen this sensation of pain. In such cases, however, the use is purely temporary and to avoid greater evils. In cases of fatigue the best remedy and the only one which will completely restore the tired cells is natural sleep, or such change of occupation as will give the overworked cells a chance to recover. While the ordinary person never thinks of using chloroform or chloral for overcoming the sense of fatigue, there are two narcotics whose use is frequently resorted to; these, like alcohol, soon result in the formation of habits which, if

continued, may become dangerous to the health. One of these narcotics is tobacco, the other opium.

Tobacco.—This very commonly used substance owes its narcotic effect to the presence of a substance called nicotine. In a pure state nicotine is a colorless liquid. When heated in burning tobacco, the nicotine turns to vapor and is given off in the smoke. Nicotine is a powerful narcotic poison, two or three drops being sufficient to kill a man. In ordinary usage very tiny amounts actually enter the system. Its action when taken into the body for the first time is to produce sickness of the stomach (nausea) and general discomfort. In this way (by the nausea) the body tries to tell us of the presence of a poison. If, however, the use is persisted in, the body gradually becomes accustomed to its presence, the nerves which tell of fatigue are gradually dulled, and a grateful sense of restfulness is produced, which continues until the effect of the narcotic wears off. It is this pleasing sensation that constitutes its danger, for when the effect wears off the sense of weakness returns, and with it a desire for more, which if yielded to often results in the introduction into the body of amounts of nicotine great enough to injure the tissues permanently. The fact that the body is able to overcome the results of moderate use without permanent injury, and that pleasant sensations accompany such use, is responsible for the wide-spread use of tobacco.

Of the various methods of use those will evidently be least dangerous which introduce the least amount of nicotine. In this respect, smoking, when the smoke is not inhaled, is the least dangerous, and chewing, where more or less of the juice is swallowed, most dangerous. Fortunately, the frequent spitting which the chewing habit

induces renders such a method of use disgusting, and tends to discourage the method. Cigarettes are made of shreds of tobacco, and on account of their size may be quickly smoked and with less immediate effect than a cigar or pipe. The number required to produce nausea and their cheapness make their use particularly attractive to the young. This fact and the danger of habit forming make their use by young people one to be guarded against by every means possible. The governments of most states recognize this fact by forbidding their sale to minors. In weakening the cells the use of tobacco often prevents their full development. In this way the building of tissues may be hindered and the body fail to reach its full size. We call this the stunting effect on growth. It is this stunting that makes the use of tobacco very dangerous to young people who are growing, and forms a sufficient reason for prohibiting the use of tobacco to minors, were no other evils resultant from its use. In adults this effect is not so noticeable, but in times of sickness, when the full strength of the cells is required to resist the ravages of the disease, their weakening by tobacco often lowers their power to such an extent as to seriously hinder recovery, and a tobacco user may on this account succumb to a disease which a non-user would have no difficulty in conquering.

The cure of the tobacco habit is simply a matter of will power. It naturally becomes more difficult the more extensive the use. It is undoubtedly better never to have formed the habit, and every experiment to determine the effect of tobacco tends to show the needlessness of such use and the advantage of abstinence.

Opium.—This is a much more powerful narcotic than tobacco, and the knowledge of its evil effects and habit

forming tendencies is sufficient to keep most people from its use. The main danger in this respect comes from the same practice as the forming of the alcohol habit through the use of patent medicines. People suffering from headaches, or other painful maladies, sometimes seek relief in medicines whose composition they do not know. These may contain opium or other dangerous narcotics, and thus the use becomes a habit without their realizing it. The quickness with which these powerful drugs act and the rapid degeneration which results, make such unknown medicines a positive danger to the community. The best way to avoid such evils is to never take any medicine unless it is prescribed by a physician.

VII. DIGESTION.

THE purpose of food as shown in the preceding chapter, is to furnish the cells of the body with material out of which they may either make new cells or develop energy. It is evident that a piece of beef, while it may contain all the nutrients necessary to these two processes, is not in the condition to be taken up by the tiny cells of the body and used by them for these purposes. The beef also contains matter which is of value neither as tissue building material nor as fuel.

In general, then, all foods require special preparation in order to fit them for use by the cells of the body, and the steps included in such preparation are called digestive processes or collectively, *digestion*. The processes of digestion are of three kinds: First, the separation of a food stuff into its nutrient and non-nutrient parts. Second, the transformation of the nutrients into a liquid form which the cell can absorb. Third, the elimination of the non-nutrient matter. These three processes are preliminary to the use of food by any living cell, and while the means employed must vary with the food, digestion is common to all living forms as the first step in the manufacture of food into living matter and activity.

In man the processes of digestion take place in a part of the body known as the alimentary canal.

The alimentary canal.—This canal is a continuous tube extending through the body and having two openings, the mouth at one end and the anus at the other. In some of

PLATE I

the lower animals, such as the common earthworm, this tube is almost straight and of the same diameter throughout the body. In man it is very much twisted and doubled upon itself, so that its entire length in our body is over thirty feet. Its internal diameter in man varies at different

FIG. 23 — The alimentary canal.

points, forming at such points large cavities (mouth, stomach, etc.), which are connected by narrower portions (gullet and intestines). At certain points it gives out growths of various sizes called glands (liver pancreas, salivary glands). The main bulk of this tube and its outgrowths is packed and coiled in the abdominal cavity (see plate I) of the body.

The tube is made up of various kinds of tissue, muscular, connective, epithelial, blood, and nerve, arranged in such a way as to form layers of varying thickness. It is lined throughout its entire length with a layer of epithelial tissue called the mucous membrane. The various parts of the tract have been given special

names, and each part performs a distinct action in the preparation of food for the body.

While there are many of these parts each having a special kind of activity, their actions admit of classification under general headings. For example, the parts known as glands

all agree in secreting fluids called digestive fluids. These fluids, owing to the presence of certain chemical compounds called ferments, dissolve the nutrients with which they come in contact. Other parts, by rolling about and crushing the foods, cause them to become mixed with these fluids, and thus be more readily dissolved. In general, then, the functions of the parts or organs of the alimentary tract are to secrete digestive fluids, or to dissolve nutrients, or to crush and mix the nutrients with the fluids. We shall consider each of these actions as they occur in the various parts of the tract.

REGIONS OF THE ALIMENTARY CANAL.

The buccal cavity.—(See Fig. 24.) This is the name given to that part of the tract usually called the mouth. It is bounded in front and at the sides by the lips, teeth, and cheeks, below by the tongue, and above by the palate. The bony part of the palate which forms the roof of the mouth and is called the hard palate, is extended backward into a soft part (the soft palate) with a pendant

FIG. 24 — Relation of the buccal and nasal cavities to the esophagus; *n*, brain cavity; *l*, hard palate; *f*, uvula; *g*, opening of eustachian tube to ear; *e*, epiglottis; *k*, tongue; *c*, windpipe; *b*, esophagus.

called the uvula. The hard palate separates the buccal cavity from the nasal cavity. The uvula, which can be seen easily by opening the mouth and examining it with a mirror, forms a curtain which guards the opening from the mouth cavity into the pharynx or throat cavity. This opening at the back of the mouth cavity is flanked on either side by two swollen masses of flesh called the tonsils. All

these cavities are lined with a thin membrane which is continuous with the outside skin and of the same kind of tissue (epithelial). This membrane consists of a network of cells carrying blood vessels and nerves, and secretes a liquid called the mucus which moistens and lubricates the surface of the cavities. It is called the *mucous membrane*.

FIG. 25 — Teeth in place, showing place of nerve supply.

teeth and the tongue, which have a purely mechanical action in digestion; while connected with the mouth cavity are several outgrowths (the salivary glands) which secrete and pour into it a digestive fluid called saliva.

The teeth. — (See Fig. 25 and Ex. XXXIV.) The

In the mouth are two structures, the

teeth are hard bodies set at intervals in two semicircular rows of sockets. These sockets are formed for the purpose in the bones called the jaws. The teeth are surrounded by a layer of flesh which we call the gums. Though hard like bone they are not of the same structure but are actually hardened masses of epithelial tissue. In the life of a man or woman two sets of teeth are developed. The milk teeth are the first to appear, and are shed early in life. The second teeth, or permanent set, begin to appear before the milk teeth are all gone, and gradually replace them. These two sets agree in their origin and general structure but differ in number. There are twenty milk teeth and thirty-two permanent teeth.

Kinds of teeth. — (See Fig. 26.) Teeth of the same set

FIG. 26 — Kinds of teeth ; *a*, incisors ; *b*, canines ; *c*, bicuspids or premolars ; *d*, molars.

differ in shape. For example, in the permanent set there are found in front, eight teeth, with broad chisel-shaped

edges. These teeth, four on each jaw, are called *incisors*, and are shaped to cut up the food taken into the mouth. On each side of the incisors is found a tooth with a sharp point. These teeth, four in number, are called *canines* from their resemblance to the teeth of a dog. The shape of the canines fits them for tearing apart the food which cannot be easily cut. Next behind the canines, two on each side of the jaw, are teeth with two pointed surfaces. These are called the *bicuspid*s or *premolars*. They serve the same purpose as the remaining six teeth on each jaw which are called the *molars*. The twelve molars have rough broad surfaces for grinding the food into small pieces. The third molar on each side of a jaw sometimes does not appear until after the twentieth year, and these teeth are often spoken of as the wisdom teeth.

The milk teeth set differs from the permanent set in having no bicuspid and only eight molars. The numerical relation of the teeth is often represented by a fraction whose numerator gives the number of each kind on the upper jaw, and the denominator expresses the same relation for the lower jaw. Such a fraction is called a dental formula. The permanent set formula for man is:

$$\text{Dental formula} = \frac{2 + 2, 1 + 1, 2 + 2, 3 + 3}{2 + 2, 1 + 1, 2 + 2, 3 + 3} = 32 \text{ teeth; or, for}$$

$$\text{half a jaw, dental formula} = \frac{2 + 1 + 2 + 3}{2 + 1 + 2 + 3} = 16 \text{ teeth.}$$

$$\text{Milk teeth formula} = \frac{2 + 2, 1 + 1, 0 + 0, 2 + 2}{2 + 2, 1 + 1, 0 + 0, 2 + 2}.$$

No animal known has more kinds of teeth than man. The number on each jaw differs greatly among different

animals. For example, in the rabbit the dental formula is as follows:

$$\text{Rabbit dental formula} = \frac{2 + 2, 0 + 0, 3 + 3, 3 + 3}{1 + 1, 0 + 0, 2 + 2, 3 + 3} = 28.$$

Structure of the teeth.—(See Fig. 27.) The part of a tooth above the gum is called its *crown*. When a tooth is pulled we find that the crown is only a small part of the entire tooth. At the point where it enters the gum it narrows into a part called the *neck* and extends below in one or more projections called the *fangs* or *roots* which anchor it firmly in its bony socket. The incisors and canines have only one fang, the bicuspid may have two, while the molars may have as many as five and as few as two.

The internal structure of all human teeth is practically the same. If we split a tooth in two, lengthwise, examination shows that the crown is covered with an external layer of hard matter called *enamel*. This thins down at the neck and is succeeded by a fang covering called *cement*. The main bulk of the tooth is formed of a bonelike substance called *dentine* which surrounds a small central cavity called the pulp cavity. During life this cavity is filled with nerve fiber and blood vessels interlaced in a network of connective tissue, called the *pulp*. It is lined with a layer of epithelial cells which secrete the mineral matter of which the outer layers are formed. This cavity has a small

FIG. 27—Section of a tooth; *k*, crown; *e*, enamel; *n*, neck; *d*, dentine; *f*, fangs; *c*, cement; *p*, pulp cavity inclosed by dentine.

opening at the base of each fang through which the nerves and blood vessels communicate with the rest of the body. The blood supplies the epithelial cells with the material out of which they build the tooth, and the nerves probably direct this process. In some animals the teeth grow throughout life, but in man their growth ceases upon the attainment of a certain size.

Digestive action of the teeth. — The teeth grind and

FIG. 28 — The tongue.

break the food into small particles, and thus prepare it for the solvent action of the digestive juices. If the food is not properly chewed its solution is necessarily delayed just

as a lump of sugar dissolves less quickly than the same amount of pulverized sugar. The grinding action of the teeth is called *mastication*.

The tongue. — (See Fig. 28.) The tongue is a soft, fleshy mass of matter mainly composed of muscle with a covering of mucous membrane. It is fastened at its base to a bone called the hyoid. Its tip projects forward, and is free and very mobile. It has two functions, one mechanical, the other sensitive in nature. As a sense organ it is the seat of taste. As a mechanical organ it mixes the food with saliva and pushes it into the proper position for chewing and swallowing. It also serves as an indicator of the health of the digestive system by becoming coated with a "fur" or coat of yellowish mucus whenever the digestion is impaired. Its healthy color is red.

CHEMISTRY OF DIGESTION.

Both tongue and teeth are merely mechanical agents which prepare the food for the solvent action of a liquid called saliva. This liquid is a typical digestive fluid in that it is produced by a structure called a gland, and exerts a solvent action through chemical change induced by means of a contained ferment or enzyme. We shall understand its action better if first we make clear what a gland and a ferment are.

Glands and secretions. — (See Ex. XXXII.) Every cell has the power of taking in food and giving off waste matter. Evidently, then, the character of the food which a cell takes from the blood determines, in a way, the kind of waste which it shall give off. Sometimes the matter so given off is a solid, and the result is the formation of hard

structures like the bones and teeth. When it is liquid or semiliquid the matter is called a secretion. There are many kinds of secretion given out by the various cells of the body, and some of the secretions have an important part to play in the action of the body. So important are some of these secretions, in fact, that certain cells are set aside whose sole work is their preparation. These specialized cells often become grouped together in certain regions, and are then spoken of collectively as glands. A gland, then, is merely a collection of cells whose special duty is the giving off of a certain kind of secretion. To this group of secretions belong all the digestive fluids, and the cells or group of cells which secrete them are called the digestive glands. In this group are included the salivary glands, the pancreas, the liver, and the mucous membrane.

The form of tissue which furnishes most of these specialized cells is the epithelial, and the simplest form of a gland would be a layer of these cells so placed as to receive food from the blood and give off their secretion. Such a glandular structure is called a secreting membrane, and an example is the mucous membrane which lines all parts of the alimentary tract. (See Fig. 29.)

In so complicated a structure as the body, however, it is necessary to economize space, and it is easy to see that one of the simplest methods of doing this is to arrange a layer in a simple fold or sac. In this form (see Fig. 29) the secretion is poured out into the central cavity or *lumen*, which, when filled, overflows through the mouth or opening of the gland. Of such a character are the gastric glands of the stomach, and the simple glands (*crypts*) of the intestine. Evidently, the next step in saving space would be to collect a series of these simple

sacs about a common duct, and such arrangements are called *tubular*, *racemose* or *tubular-racemose* according as the separate elements are tubelike or branched or a combination of the two. Such glands are called compound

Fig. 29—Forms of glands; *A* and *B*, simple layers of mucous membrane cells; 1, a simple tubular gland; 2, a forked tubular gland; 3, a simple saccular gland with duct, *d*; 4, a saccular gland still more divided; 5, a saccular gland still more divided; 6, a compound racemose gland with common duct, *d*.

glands, and in this class are the salivary glands, liver, and pancreas.

Enzymes. — (See Ex. XXXI.) The secretions of the digestive glands owe their power of dissolving nutrients to the presence of certain complicated chemical compounds called enzymes. The action of these enzymes is little understood at the present time, but their effect is well known. In general, they all agree in causing a chemical change in a nutrient without being themselves consumed in the process.

For example, if we add a little malt (see p. 83), which contains the enzyme diastase, to some starch paste, the starch is gradually changed to sugar, but the amount of the diastase which causes the change is neither increased nor diminished by the process.

Similarly, yeast added to a sugar solution causes the sugar to break up into two simpler compounds, alcohol and carbon dioxide, while the number of yeast plants is not decreased by the process.

The name ferment was long used to describe these compounds, and hence the action of yeast or bacteria was described as fermentation. We now know that yeast and bacteria produce their effect through definite chemical compounds, and to all such compounds we give the name *enzymes*.

The salivary glands. — (See Fig. 30.) There are three pairs of these glands in man. They are called the parotid, the submaxillary, and the sublingual. All these glands are of the compound, racemose type, and resemble bunches of grapes in which the grapes are the sacs of secreting cells, and the connecting stems are the hollow ducts. The parotid glands lie in front of and under the ear on each

side of the head, and their ducts enter the mouth on the inner side of the cheeks, just about opposite the second upper molar teeth. It is the inflammation and enlargement of these glands that causes the disease known as the mumps.

The submaxillary glands lie below the two halves of the lower jaw, and their ducts open into the mouth at points near the middle line of the under side of the tongue.

The sublingual glands lie under the mucous membrane below the tongue and floor of the mouth. They open into the floor of the mouth by many small ducts.

Action of the salivary glands. — Each cell of a gland obtains its food supply from the blood vessels, and out of this food it manufactures the secretion known as saliva. Since the amount of saliva needed to digest the food we eat varies with the amount of food present in the mouth, it is evident that the secretion of the cells must be under control. In this respect the glands are like a factory whose output is regulated to meet the conditions of supply and demand. In the case of the cells this control is exercised by nerves whose general arrangement is indi-

FIG. 30 — Relation of salivary glands to the mouth cavity.

cated in the diagram. (See Fig. 31.) The results of this arrangement are as follows: The nerves may be stimulated in several ways: by the contact of the food in the mouth with the nerve ends, by the odor of the food which stimulates the nerve ends in the nose, or by the stimula-

FIG. 31.—Diagram to illustrate mouth control.

tions of the nerves at the brain end, as by thought of food. When the nerves are stimulated in any of these three ways the impulse they carry is communicated to two points. One impulse goes to the blood vessels and causes them to dilate, and thus supply more blood and food. The other impulse is given to the cells, and stimulates them to greater secretive activity. The result of these two directing impulses is an increased flow of saliva.

In a similar way, this threefold nerve connection also accounts for the dryness of the mouth when certain conditions are present. For example, fright, a bad smell, or unpleasant taste, may start impulses along the nerve which, instead of increasing the flow of blood and secretion, may constrict the blood vessel and diminish the secretive activity of the cells.

The saliva.—The secretion produced by the salivary glands is called the saliva. The liquid found in the mouth and constituting the spit is not pure saliva, but a mixture of this liquid with a secretion of the mouth lining called the mucus, together with a number of other more or less solid elements. This mixture is a colorless, cloudy,

slightly alkaline liquid, with a ropy consistency due to the mucin in it, and is often frothy from the presence of air bubbles. Pure saliva is a clear, viscid, colorless liquid consisting mainly of water and a certain enzyme called *ptyalin*. It is this enzyme which is responsible for the chemical action of saliva in digestion.

Action of saliva. — Of all the nutrients taken into the mouth starch is the only one acted upon by the saliva. If clear saliva is allowed to act upon starch it changes it to sugar in exactly the same way that the diastase of malt changes grain starch to sugar. This sugar is, of course, readily soluble in the watery part of the saliva, and thus the nutrient starch is forced in solution. That this transformation is actually produced may be readily demonstrated by chewing a little dry biscuit for a few moments. The dry, tasteless mass soon becomes sweet to the taste as its starch is changed to sugar by the saliva. The chemical changes involved in this transformation are rather complicated, but for our purpose it is sufficient to state that they have their origin in the action of the enzyme *ptyalin*, and we can, therefore, say that it is the *ptyalin* which produces the change.

Experiments with starch paste and clear saliva (see Exs. XXXV. *a* and XXXVI.) show that this transformation takes place best at the normal mouth temperature (98° F.), while high temperatures (above 145° F.), or very low temperatures (below 32° F.), or the presence of strong acids or alkalis are sufficient to stop completely the *ptyalin* action. Further, cooked starch is much more easily changed to sugar than uncooked, and requires less time for the action of the enzyme.

Saliva also aids digestion in a mechanical way by fit-

ting the food for swallowing. It is impossible to swallow dry food, and the moistening with the saliva makes this action possible, and at the same time acts as a lubricant and allows it to slip down more easily. In fact, so short is the time that food spends in the mouth in ordinary eating that probably very little starch is changed into sugar there, and the mechanical action, therefore, is the most important. Saliva, as we shall see later, also aids in producing the sensation of taste.

SUMMARY OF DIGESTION IN THE MOUTH.

1. The teeth grind the food into small particles and fit it for rapid and thorough mixing with the saliva. (Mastication.)
2. The tongue rolls the food about into position for mastication and mixes it with the saliva.
3. The saliva through the action of the ptyalin enzyme converts part of the starch into soluble sugar.
4. The saliva moistens and lubricates the food mass and thus fits it for swallowing.
5. The food leaves the mouth in proper shape for the action of other juices and with part of its starch converted into sugar.

VIII. DIGESTION (*continued*).

AFTER the action of the mouth the food is ready to be passed on to the stomach, where the next steps in digestion take place. The first step in this process is the act known as swallowing or *deglutition*. In this action the

FIG. 32—Swallowing or deglutition ; *a*, tongue ; *b*, pharynx ; *c*, food bolus ; *d*, soft palate ; *e*, uvula ; *f*, epiglottis.

A shows food being pushed back by the tongue. *B* shows position of uvula and epiglottis as food falls into the pharynx.

uvula at the back of the mouth is raised automatically to close the nasal passage, and the tongue simply pushes the food backward until it falls or slips into the pharynx cavity.

Structure of the pharynx cavity. — (See Fig. 24.) This cavity is a conical bag just back of the mouth cavity and about five inches in depth. It has seven apertures ; two into the nasal cavity at the top, two into the ear (the Eustachian tube openings), one into the mouth, one into the windpipe (glottis), and one into the esophagus or

gullet. The glottis opening is protected by a sort of trapdoor called the epiglottis, which is closed during swallowing but open at other times to permit the entrance and exit of air in breathing. The whole cavity is lined with mucous membrane, whose secretion of mucus moistens and makes flexible its walls. (Excess of this mucus produces catarrh.) When the food is swallowed the epiglottis is closed and the food falls upon the top of the gullet. Sometimes the epiglottis does not close quickly enough, and some food enters the air passage.

FIG. 33 — Diagrammatic section of the wall of the esophagus; *S. M.*, serous membrane; *L. M.*, longitudinal muscles; *C. M.*, circular muscles; *M. M.*, mucous membrane. Between the circular muscles and the mucous membrane is a thin layer of connective tissue.

This produces choking, that is to say, the forcible expiration of air in the attempt to force out the misplaced food. If properly swallowed, the food is forced along the gullet to the stomach.

The esophagus. —

The esophagus, or gullet, is the name given

to that part of the alimentary tract which begins at the lower end of the pharynx cavity, and passing through chest and diaphragm, enters the stomach. This portion is in the form of a tube of about nine inches in length and is formed by three layers of tissue. (See Fig. 33 and Ex. XLI.) The inside layer is a continuation of the mucous membrane, and the secretion of mucus here together with the saliva acts as a lubricator, making the food slip down easily. The outer layer is formed of muscles, while between this and the inside layer is a layer

of connective tissue. The muscular layer is actually a double layer, the inner fibers arranged in a circular manner, while the outer ones are extended lengthwise of the tube. By alternate contractions and expansions of these muscles the food is forced along the tube to the stomach. In this action the effect produced is similar to what happens when a ring slightly smaller than the tube is forced along a rubber tube; it forces any bodies inclosed in the tube in the same direction that the ring is moving. The contractions of the circular and longitudinal bands of muscles necessary to this action in the gullet have received the name *peristalsis*. Through peristalsis of the esophagus walls the food is always forced toward the stomach, regardless of the position of the body. The food undergoes no chemical change in either the pharynx cavity or esophagus.

THE ABDOMINAL CAVITY AND CONTENTS.

This large body cavity (see p. 29, and plate I), incloses the greater part of the alimentary canal. (See Ex. XXXIII.) It is separated from the chest at the top by a muscular partition called the diaphragm, while its base is formed of the basketlike bones of the pelvis. At the back are the spinal column and the lower ribs, while the sides and front are formed of thick sheets of muscle without bone support. This cavity is lined throughout with a very smooth membrane called the peritoneum. This peritoneum is so arranged that it not only lines the cavity but also, by extensions, forms folds which support the various parts of the alimentary canal. These folds are so arranged that the parts of the canal are hung in it like a

stone in a sling. (See Fig. 34.) The entire membrane is lubricated by a watery liquid or serum, and

FIG. 34 — Intestine supported by a fold of the peritoneum (mesentery), showing method of entrance of blood vessels.

for this reason it is called a serous membrane. It is also richly supplied with blood vessels. The organs sup-

FIG. 35 — External view of the stomach, showing covering of peritoneum and muscles; 1, esophagus; 2, cardiac end; 3, pyloric end; 4, small intestine; 5, a portion of peritoneum turned back; 6, longitudinal muscle fibers; 7, circular muscle fibers; 8, oblique muscle fibers.

ported by this membrane are the stomach and the many folded intestines.

The stomach.— (See Ex. XXXIII.) This organ is a bag-like swelling (see Fig. 35) of the alimentary canal lying just under the diaphragm and across the upper part of the cavity. The larger end is at the left side of the body, and into its upper surface the esophagus opens by an orifice called the *cardiac orifice*. The right end of the stomach tapers into a narrow neck which is connected with the beginning of the small intestine by an opening called the pylorus or *pyloric orifice*. The whole bag is supported in a fold of the peritoneum, whose two parts come together at the lower edge of the stomach and hang down like an apron over the front of the abdominal cavity. This apron is known as the greater omentum, and is frequently the seat of a great accumulation of fat.

The walls (see Ex. XLI.) of the stomach are composed of four coats or layers. (See Fig. 36.) The outermost is a serous membrane (the peritoneum just mentioned). Next inside this is a muscular coat composed of three layers of muscle fibers. Then comes a layer of connective tissue (the submucous coat), and lining the entire stomach is the mucous membrane. The blood vessels of the stomach enter between the folds of the peritoneum, and

FIG. 36 — Section of stomach wall; *s*, serous membrane (peritoneum); *l. m.*, longitudinal muscles; *c. m.*, circular muscles; *s. m.*, submucous coat; *m. m.*, mucous membrane; *d*, ducts in mucous membrane.

branch into capillaries in the submucous layer. The muscular coat is made up of unstriped muscle fibers, and varies greatly in thickness at different parts. It is thickest at the pyloric end. The outermost layer of fibers is a continuation of the longitudinal fibers of the esophagus and is

FIG. 37.—Longitudinal section of the stomach; *a*, esophagus; *b*, greater curvature; *c*, lesser curvature; *d*, sphincter pylorus; *h-i*, small intestine. The ridges in the stomach are folds of mucous membrane.

thickest over the curvatures of the stomach. The circular muscle fibers of the esophagus are likewise continued, and form the inner layer of muscle over the stomach. This circular layer becomes thickest at the pyloric end. Between these two layers is a very incomplete oblique layer whose fibers radiate from the cardiac orifice. The thick ring of circular fibers at the pyloric end is called the *sphincter pylorus*, and controls the entrance of food into the small intestine.

The mucous coat is a pink membrane which is not elastic, and, though smooth when the stomach is fully distended, is thrown into folds when the organ is empty. This membrane derives its pink appearance from the blood vessels underlying it and becomes much redder during digestion, as these submucous capillaries become distended with blood. When examined with a lens the mucous membrane is seen to be covered with shallow pits. These pits are the mouths of many simple tubular glands (the gastric glands) which secrete the digestive fluid of the stomach. This fluid is called the gastric juice. Between these special glands are the ordinary mucous glands which secrete the lubricating mucus.

Gastric glands. — These glands (see Fig. 38) are simple tubes lined with epithelial cells, formed by the folding of the mucous membrane. Between successive tubes project layers of the submucous capillaries. These capillaries furnish the food out of which the cells manufacture the fluids and ferments used in stomach digestion. The cells of these glands are of two kinds. Those which secrete the enzymes, *pepsin* and *rennin*, are called the *chief cells* to distinguish them from another group called the *border cells* which are supposed to secrete hydrochloric acid. The exact function of these border cells is still unknown. (For arrangement of these cells see Fig. 38.)

FIG. 38 — A peptic gland, from cardiac end of stomach. Very much magnified. A, central or chief cells, which make pepsin; B, border or parietal cells, which make acid. [From Miller's *Histology*.]

DIGESTION IN THE STOMACH.

Gastric juice.—The digestive action of the stomach is due mainly to the action of the digestive fluid produced by the gastric glands. This gastric juice is a thin, nearly colorless fluid, consisting mainly of water and holding in solution a small proportion of hydrochloric acid and two enzymes, pepsin and rennin.

As a result of the acid contained it gives a red color to blue litmus, and, since ptyalin can act only in an alkaline medium, the saliva swallowed with the food can exert no digestive action upon starch after the food is acidified.

The flow of the gastric juice begins with the entrance of food into the stomach, and the flow increases with the amount of food taken in. This flow may also be stimulated by certain food accessories, by the alkaline saliva, and by the flavor of the food. Normally, about three quarts of gastric juice is produced by the glands in twenty-four hours.

Digestive action of the enzymes.—*Pepsin.* This enzyme has the power to transform proteids to a soluble form called peptone. It acts only in an acid medium, and what is known as peptic digestion is therefore the result of the combined action of pepsin and hydrochloric acid. Peptic digestion varies with temperature. Low temperatures retard, while very high temperatures may stop completely the action of the pepsin. Pepsin can be obtained in a fairly pure state (see Ex. XXXV., b) and with the product so obtained it is possible to prepare artificial gastric juice whose action upon various foods can be observed with great accuracy. (See Ex. XXXVII.) In such experiments the artificial juice is allowed to act upon food prep-

arations contained in glass vessels. The results of such experiments may be summarized as follows:

The proteid thus treated first swells and softens and then is slowly transformed into a soluble form of proteid known as peptone.

This peptone is the result of the combined action of the pepsin and hydrochloric acid, the food proteids being broken gradually into simpler and simpler forms, until finally the soluble peptone results.

Peptic digestion is found to take place best at a temperature of from 98° to 102° Fahrenheit.

The action of peptic digestion is confined wholly to the proteid parts of the food, and does not directly affect any other nutrient. In fat meats the albuminous pockets are dissolved away in this action, and the result is the liberation of the fat globules inclosed by the pockets, but the pepsin does not change the fat in any way. In similar manner starch grains held in albuminous pockets may be liberated, but the starch itself is not affected.

All these actions of the pepsin and acid which are observed in experiments with glass vessels take place normally under similar conditions in the stomach.

Rennin. (See Ex. XXXVII., b.) Gastric juice contains a second enzyme called rennin. This enzyme plays an important part in the digestive action of the gastric juice upon the particular form of proteid found in milk. When milk proteid (*caseinogen*) comes in contact with gastric juice, the rennin enzyme causes it to coagulate and separate from the rest of the milk as curd or *casein*. This action is exactly similar to the curdling of milk by acid when it sours. This separation of milk proteid by the rennin prepares it for the action of the pepsin and

acid, and once separated, peptic digestion takes place as with ordinary proteid. In children whose diet is almost entirely of milk this rennin action becomes very important.

Hydrochloric acid. (See Ex. XXXIX.) Aside from the aid which this acid gives in peptic digestion its presence is important in that it destroys the germs of fermentation and disease, and probably dissolves some mineral salts. Its action in destroying germs permits the food to be stored in the stomach for some time without undergoing decay. In this way the stomach may act as a storehouse of food. While proteids are practically the only nutrients digested in the stomach, it is also true that some of the preparatory actions upon other nutrients take place here. For example, the heat of the stomach is sufficient to liquefy the fats of the food, and the digestion of the proteid covers of starch and fat particles liberates these nutrients and prepares them for future digestive changes. Sugar and certain mineral salts may also be dissolved here by the water and acid of the gastric juice. The mixture of all these substances together with the digested and undigested particles of food and the gastric juice results in the formation of a milky fluid called *chyme*. This chyme is from time to time forced out of the stomach into the small intestine.

Mechanical action of the stomach. — Once in the stomach the food is entirely cut off from the rest of the alimentary canal by the contraction of the cardiac and pyloric sphincter muscles. It was long thought that during the period of stomach digestion the food was forced in continuous circulation round and round the stomach until sufficiently ground, and mixed with gastric juice to permit of its discharge into the intestine. Later investi-

gators showed that while this mixing does take place the method is somewhat different from what was formerly supposed. According to these investigators, the stomach cavity may be divided into two parts, in only one of which does the actual mixing take place. The cardiac end of the stomach or *fundus* shows almost no movement and acts simply as a storehouse. The contractions start in the middle of the stomach and run toward the pylorus end, and only the portion in this end is actually in process of mixing. These contraction waves are repeated at intervals of two minutes or less, and finally reduce the food in this end to a thin liquid. At intervals depending upon the character of the food, the sphincter pylorus relaxes and portions of the liquid mass are spirted into the intestine while new material flows forward from the storage end to take its place. When the food is liquid these relaxations of the sphincter are very frequent, while for solid food they occur at longer intervals. The time required for the stomach to empty itself in this way is from two to five hours after a meal. The exact time, of course, depends upon the amount of food taken into the stomach, and ease with which it may be broken up. After the stomach has been empty for a certain length of time we feel a sense of hunger, and this is the signal for a renewal of the food supply. Meanwhile, the chyme which has entered the intestine is undergoing further digestive changes.

SUMMARY OF DIGESTION IN THE STOMACH.

1. The food enters the stomach by the cardiac orifice and by the peristaltic action of the muscular walls is mixed with the gastric juice.
2. The hydrochloric acid stops the action of the ptyalin and the digestion of starch, and makes possible the action of the pepsin.

It also tends to kill bacteria and prevent fermentation and aids in the solution of certain mineral salts.

3. The pepsin converts proteid into soluble peptone, part of which is absorbed directly by the blood vessels of the stomach.

4. The rennin coagulates the proteid in milk and prepares it for the action of the pepsin.

5. The partly digested proteids, and starch together with the undigested matter leaves the stomach as an acid mixture called chyme.

FIG. 39 — The stomach and intestines; 1, stomach; 2, duodenum; 3, small intestine; 4, end of ileum; 5, caecum; 6, vermiform appendix; 7, ascending colon; 8, transverse colon; 9, descending colon; 10, sigmoid flexure; 11, rectum; 12, anus.

DIGESTION IN THE INTESTINE.

While some starch is changed to sugar in the mouth and some proteid to peptone in the stomach, the real seat of digestion in the body is the small intestine. Here all kinds of nutrients are digested, and the final preparation of the food for use in the body takes place. Here, too, the digestible matter is separated from the indigestible. In short, even if there were no preliminary mouth or stomach action, the small intestine would be capable of performing all the digestive actions necessary to prepare the food for the body.

The intestine. — All of the alimentary canal between the stomach and the anus is known as the intestine. This part of the canal consists of a tube of varying diameter,

and to its different parts have been given appropriate regional names. The part next to the stomach, and continuing for some twenty feet from the pyloric orifice, is all included under the name small intestine. It varies in diameter from two inches at the stomach end to less than an inch where it enters the large intestine. By large intestine we include the five odd feet of tube extending from the point where the small intestine is considered to end to the anus. This portion of the tract has a diameter of from two and a half to one and a half inches. The relative position, size, and general arrangement of the two intestines are shown in Fig. 39.

The small intestine. — (See Ex. XXXII.) The entire part of the alimentary canal included under the name of small intestine is held in place by a fold of the peritoneum called the mesentery. This mesentery is about four inches in length at its back edge where it is fastened to the spinal column, and widens to twenty feet at its outer edge. The intestine is hung in this fold like an arm in a sling, and the outer edge of the mesentery and the intestine are very much coiled and folded to confine this long tube to the space adapted for it. The different regions of the small intestine are indicated by three names. The twelve inches nearest the stomach receive the name of *duodenum*, the succeeding two fifths is called the *jejunum*, and the remainder is called the *ileum*. These names are mainly useful in permitting us to designate special parts of the tube.

Like the stomach the small intestine has four coats (see Ex. XLI.) or layers, an external serous layer (peritoneum), next inside a muscular layer, inside of this a submucous coat, and a lining of mucous membrane. (See Fig. 40.)

The serous coat is formed of the fold of the mesentery. The muscular coat is in two layers, an outer one of longitudinal fibers and an inner of circular fibers.

FIG. 40—Wall of small intestine; *a, b*, mucous membrane with projecting villi; *c, d*, connective tissue layers (submucosa); *e*, circular muscle fibers; *e'*, longitudinal muscle fibers; *f*, serous membrane; *g*, spaces between villi; *h*, opening of *crypts* or simple intestinal glands; *i*, blood vessels in villi.

The submucous coat of connective tissue binds together the muscle and mucous membrane layers just as in the stomach, and receives the blood vessels which enter in a fold of the mesentery. The mucous coat which lines the whole intestine is pink, soft, and filled with minute vessels of the blood and lymphatic system. In the upper part of the intestine it is permanently folded into ridges known as the *valvulae conniventes*, which increase the digestive surface and delay the passage of the food particles. In

the lower part these ridges disappear, and to the eye it appears smooth. Examined carefully with a lens the entire inner surface of the intestine is seen to be raised in minute projections called villi, like the pile of velvet.

(See Fig. 41.) We shall consider the structure and function of these villi more in detail in our study of the absorption of food. (See p. 134.) Between the bases of these villi are the openings of tiny glands called the crypts of Lieberkühn.

FIG. 41 — Villi on surface of intestine (much magnified).

Digestive glands of the small intestine. —

The glands which secrete the digestive fluids of the small intestine are three in kind. Two of these are large and lie outside of the intestine, being connected with it by means of a duct. The other kinds of glands (the crypts) are very numerous and embedded in the mucous membrane itself. The two large glands are known as the liver and the pancreas, and the ducts from these two glands unite outside the intestine and pour their combined secretions into the intestine through a common duct at a point on the duodenum a few inches from its connection with the stomach. The secretions of these two large glands are known as the bile and the pancreatic juice. The relation of these two glands and the ducts is shown in the diagram (see Fig. 42). The secretion of the crypts is called the intestinal juice.

The liver. — (See Fig. 46 and Ex. XXXII.) This is the largest digestive gland in the body. It is a dark red mass located in the upper part of the abdominal cavity just under the diaphragm, and partly overlapping the stomach. The entire gland is covered with peritoneum, and is divided by a fissure into two unequal lobes. Of these two lobes,

the right is much the larger. These two lobes are connected on the under side by ducts which emerge from each lobe and fuse into a common duct called the hepatic

FIG. 42—Relation of bile duct and pancreatic duct to stomach and intestine.

duct. On the under side is also located a small sac (the gall bladder), in which the bile secreted by the liver is stored until needed by the intestine. From this sac another duct (the cystic), leads, and fuses with the hepatic duct to form the common bile duct which opens into the duodenum. Two large blood vessels (the portal

vein and the hepatic artery) carry blood to the liver, entering it on the lower side and breaking up inside the

FIG. 43 — The liver.

lobes into capillaries. These capillaries unite again, and the blood passes out of the liver by the hepatic veins.

FIG. 44 — Two liver globules; *A*, section across central hepatic vein; *B*, section splitting the central hepatic vein; *p*, capillaries of the portal vein which break up among the liver cells and finally unite in the central hepatic vein *h*.

The internal structure shows it to be composed of a mass of many sided bodies called lobules. Each lobule (see Fig. 44) consists of a mass of cells (hepatic cells) surrounded by a network of capillaries. Branches of the bile ducts also enter these lobules and receive the bile which the hepatic cells secrete from the blood. Each hepatic cell, therefore, is a true gland cell, and the combined secretion of all these cells is collected in the branched ducts and flows downward until it enters the common hepatic duct. From this point it may take one of two courses.

FIG. 45 — The pancreas; 1, central duct; 2, pancreatic duct; 3, stomach.

It either enters the common bile duct and flows directly into the intestine or it backs up and accumulates in the gall bladder. In this way bile is stored in this bladder so that when necessary it may be called on to supply the intestine with increased amounts of bile. In such case, bile flows from the lobes of the liver and from the gall bladder directly into the intestine. The bile is a thick golden-brown liquid of bitter taste. About a quart is produced daily, and performs very important functions in digestion.

The pancreas.— (See Fig. 45 and Ex. XXXII.) This

organ, often called the sweetbread, is an elongated racemose gland of a pinkish-yellow color lying along the greater curvature of the stomach. The cells of this gland secrete a digestive fluid called the pancreatic juice, which they pour into a central duct. This duct joins the bile duct close to its intestinal opening, and bile and pancreatic juice mingle and enter the intestine by the common opening. (See Fig. 42.)

Pancreatic juice.—This is a watery secretion which contains three very important enzymes. One (trypsin) changes proteid to peptone, one (amyllopsin) changes starch to sugar, and the third (steapsin) acts upon fats.

DIGESTION IN THE SMALL INTESTINE.

The partly digested food or chyme which is forced from time to time through the pyloric orifice of the stomach is now ready for the action of the various digestive fluids of the small intestine. (See Ex. XXXVIII.) The digestive action which takes place in the intestine is, therefore, due to the combined action of three digestive fluids, pancreatic juice, bile, and intestinal juice. Rhythmic peristaltic movements of the muscle walls of the intestine keep the contents of the intestine in constant motion, and thus mix them with these combined juices.

Action of the pancreatic juice.—This juice owes its digestive action to its three enzymes, trypsin, amyllopsin, and steapsin.

Trypsin. (See Ex. XXXVIII., *a.*) Trypsin is a more powerful enzyme than pepsin and, unlike that, acts best in an alkaline medium. Like pepsin, its action is affected

by temperature; it is more active at ordinary body temperature (98–102° F.), while it is completely destroyed at a temperature of 172° to 176° F. Its action upon proteid is very similar to that of pepsin in that it converts proteid into soluble peptone. It differs from pepsin in that there is no swelling of the proteid under its action. The process is rather an eating away of the proteid part, and in this action the indigestible matter is left in its original shape to be broken into fragments later by the churning action of the intestine. It also reduces some albuminoids to peptone. Most of the proteid nutrient of our food receives its final preparation for absorption in the small intestine under the action of this enzyme.

Amylopsin. (See Ex. XXXVIII., *b.*) Practically all the starch of foods is digested by the action of this enzyme. It is possible that some of the ptyalin of the saliva whose action was stopped by the acidity of the stomach shares in this digestive action upon returning to the alkaline medium of the intestine. It would be impossible to determine this definitely, since amylopsin and ptyalin are apparently identical in structure and action, and in German texts on physiology the name ptyalin is used for both enzymes. The use of the two names is, however, convenient in indicating their origin. Both convert starch to sugar under similar conditions and thus transform it into a soluble substance.

Steapsin. (See Ex. XXXVIII., *c.*) This enzyme (also called lipase), which is extremely sensitive to temperature, performs two distinct operations upon fats, saponification and emulsification.

In saponification the fat is first split into glycerine and fatty acids. These latter unite with the alkalies and

alkaline salt present, and form soaps which are, of course, soluble.

In emulsification the fat is separated into tiny droplets and surrounded by the fatty acid or alkaline material in such a way as to prevent these drops from uniting again. In this form they may be absorbed directly without further action. Whether emulsification precedes saponification or *vice versa* is still a disputed point.

In general, then, the pancreatic juice first transforms the acid chyme of the stomach into an alkaline fluid through the neutralizing action of the alkaline salts in the juice, and then by the action of the three enzymes it converts proteids, albuminoids, starches, and fats into substances fit for absorption. Chyme is neutralized mainly by bile salts.

Action of the bile. — (See Ex. XXXVIII., *d.*) It has been found by experiment that when pancreatic juice is mixed with bile the emulsifying power of the steapsin is greatly increased thereby. The alkaline bile salts are of aid in the splitting of fats and consequently in the saponification. Its presence is also supposed to prevent, to a certain degree, the putrefaction of foods, though this antiseptic power is disputed by some authorities. Finally, it takes a direct part in destroying the peptic digestive action of the acid chyme. It not only neutralizes the acid but also causes a precipitation or separation of the proteids in such a form that they may be acted upon directly by the trypsin.

Action of intestinal juice (*succus entericus*). — This juice is produced by the simple glands known as the crypts of Lieberkühn. It is a yellowish liquid with a strong alkaline reaction. This alkaline character it derives from the presence of sodium carbonate, and the presence of this

alkali is important in aiding the emulsification of fats. It varies in quantity in different regions of the intestine, being scarce in the upper part and very plentiful in the lower or ileac region. Its specific action is due mainly to the presence of two kinds of enzymes. One, similar to amylopsin, aids in converting the starches to sugar. The other (invertase) converts cane sugar and other forms of sugar to grape sugar so that all sugars and starches are finally absorbed as grape sugar.

SUMMARY.

1. The food enters the intestine as an acid mixture called chyme.

2. It is mixed in the intestine with three juices, bile, pancreatic juice, and intestinal juice.

3. The three juices transform the acid chyme into an alkaline mixture called chyle.

4. In this alkaline medium the various enzymes of the three juices transform proteids and albuminoids to peptone; fats to soap and emulsions; starches and sugars, to grape sugar.

5. The action of the enzymes is aided by the mechanical action of the muscle walls.

6. The resulting mixture is then ready for absorption.

DIGESTION OF THE LARGE INTESTINE.

That part of the alimentary tract included under the name of the large intestine is subdivided, like the small intestine, into regions. These regions are termed the *cæcum*, the *colon*, the *sigmoid flexure*, and the *rectum*.

The *cæcum* is the name applied to the saclike pouch into which the small intestine opens at the top by an orifice, guarded by a valve called the *ileocolic valve*. At

the bottom of this cæcum is a wormlike extension a few inches in length called the *vermiform appendix*. Inflammation of this appendix produces the disease known as appendicitis. The appendix has no known function, and is supposed to represent a remnant left in man's development. From the point where the small intestine enters the cæcum the intestine extends upward, along the right side of the abdomen, across the cavity just below the stomach and down the other side. These portions of the intestine have the greatest diameter ($2\frac{1}{2}$ inches) of any part of the intestine and are called the ascending, transverse, and descending colons according to position. At the base of the descending colon is an S-shaped bend called the sigmoid flexure, and from this a straight portion (the rectum) proceeds to the external opening or anus.

The large intestine shows the same number of coats as the small intestine. The muscular layer, however, is not uniformly developed but arranged in bands separated by spaces where it is wanting. This arrangement gives the tube a puckered appearance. The mucous coat has no valvulæ conniventes but is filled with closely set simple glands similar to the crypts of Lieberkühn.

Action of the large intestine.—By contractions of the small intestine its contents are forced through the ileocolic valve into the large intestine. Before these contents reach the large intestine, however, nearly all the nutrient part has been digested and absorbed, and what passes through the ileocolic valve consists mainly of indigestible material called fæces. These fæces consist mainly of water, cellulose, elastic tissue, and mucin. With these substances may be present some starch and fat. If this

is the case the digestion and absorption is continued in the large intestine until finally the residue is collected in the rectum and then passed out of the body. There are no distinctive enzymes secreted by the glands of the large intestine, and what little digestion goes on there is due either to secretions which pass through from the small intestine or to bacteria. These bacteria induce fermentation, and cause the putrefaction of certain parts of the fæces. The principal function of the large intestine, therefore, is the removal of indigestible material from the body. In case of over-feeding the fæces will contain more or less nutrient matter which the small intestine was unable to digest and absorb.

ALCOHOL AND DIGESTION.

Alcohol, in *small amounts*, stimulates the flow of gastric juice and saliva, and this increased flow is accompanied by an increase in the enzyme and hydrochloric acid content of these juices. Furthermore, when taken in small quantities, it is itself very quickly absorbed by the blood vessels. Hence, the amount which remains in the stomach at any given time is entirely insufficient to interfere with the action of the digestive fluid upon the nutrients. From these statements, then, it would appear that digestion is aided by the use of small quantities of alcohol. Wherein lies the danger? Aside from the danger of habit forming, which is great and in itself sufficient reason for avoidance of the use of alcohol, the fact that this flow of juices is due to unnatural stimulation is the dangerous feature of its use.

Natural foods stimulate the flow only to the extent

of a normal production of fluid. By normal production we mean a flow sufficient to use up the materials out of which the juices are made at a rate not faster than the body can naturally supply them for the purpose. Since the supply of this material remains fairly constant in amount it follows that any unnatural stimulation will cause this material to be used up faster than it can be supplied, and periods of increased flow must, therefore, be followed by periods of decreased flow. While the occasional use of alcohol, therefore, may, as in cases of sickness, be of advantage, its continued use results ultimately in decrease in supply of fluids.

Furthermore, the action of a stimulant is such that the secretive cells soon fail to respond to a slight stimulation, and hence the amount taken must be continually increased or fail of effect, thus leading to such increase as seriously to affect other parts of the body, such as the nerve centers. Finally, *in large amounts*, the effect of alcohol is to retard seriously the action of the fluids upon the foods, and to injure the digestive organs. While, at first, small amounts of alcohol may aid digestion, they soon fail of effect, and tend to encourage habits of excess which retard digestive processes and injure the body.

Alcoholic beverages also often contain other substances which counteract the advantage of increased flow. Wines, for example, contain acids, and while the alcohol may stimulate the flow of saliva, this increased flow is of no value since ptyalin cannot act in an acid medium, and simply drains the body of valuable fluid. Since the alcohol is mainly absorbed directly from the stomach its effect upon the intestinal secretives is practically nil, and may be neglected as a factor in digestive processes.

TABULAR SUMMARY OF DIGESTION. (See Ex. XL.)

Nutrient	Region of Alimentary Tract, Where Digested	Digestive Fluid	Enzyme	Digested Product
Starch	Mouth (very little) Sm. intestine Sm. intestine	Saliva Pancreatic juice Succus entericus	Ptyalin Amylopsin Amyloptic enzyme	Grape sugar Grape sugar Grape sugar
Proteid	Stomach Sm. intestine	Gastric juice Pancreatic juice	Pepsin and HCl ¹ Trypsin	Peptone Peptone
Albuminoids	Sm. intestine	Pancreatic juice	Trypsin	Peptone
Sugars other than Grape	Sm. intestine	Succus entericus	Invertase	Grape sugar
Fats	Sm. intestine	Pancreatic juice	Steapsin	Soap or emulsion
Soluble salts	Mouth, stomach, and intestine	Water of digestive juices		Salts in solution
Insoluble salts	Stomach	HCl of gastric juice		Soluble salt
Water		Requires no digestive action		

¹ HCl = hydrochloric acid.

IX. ABSORPTION.

IN the processes of digestion, food is broken up and transformed in various ways into soluble substances, such as peptone, sugar, etc. This food, however, is still in the alimentary canal, and far removed from the cells of the body which it is to repair and build up. In this respect, then, the alimentary canal is like a big factory whose whole action is centered in the preparation of food products. A delivery system is necessary for factory products to be made available for use, and, likewise, in the body, a delivery system is necessary to distribute to all parts of the body the food prepared by the alimentary canal. In the body such a system is found in the blood system, and it is evident that the first step necessary to distribution of food is the transfer of the digested food matter to the blood. In the business world such a transfer of goods to a delivery system is called shipping. In the body the processes which are concerned in transfer of digested food to the blood are all included under the name of absorption. The object of shipping and absorption is identical, namely, to get prepared products into circulation.

Relation of the blood vessels to the alimentary tract. — The transfer of food to the blood may take place in one of two ways. It may pass directly into the blood vessels which line the walls of the intestine and stomach, or it may enter another set of tubes called the lymphatics. These lymphatics are closed tubes like the blood vessels,

and the food which they collect from the intestine they convey, by a common duct, to a large vein at the base of the neck, and place in the blood circulation at that point. In either case the ultimate disposal of the food is the same, namely, the putting it into the blood. The knowledge of the relation of these collecting blood vessels and lymphatics to the alimentary canal is important to the understanding of the method of transfer.

In the study of the structure of the stomach and intestine it was noted that beneath the lining of the mucous membrane and separating it from the muscle layers was a coat of tissue called the submucous layer.

Further, it was noted that blood vessels entered this layer through a fold of the peritoneum and within the layer spread out over the mucous lining in a many branched network of fine tubes or capillaries. These tiny, thin walled capillaries, which are thus in contact with the mucous membrane in all parts of the intestine and stomach, are the direct receivers of the digested food of these tracts. These capillaries are all connected and finally fuse into a single large vein called the portal vein.

Again, we may recall that the lining of the small intestine was not smooth, except in appearance, and that the surface of the mucous lining was raised in tiny projections called villi. (See p. 120.) A lengthwise section of one of these villi (see Fig. 46) shows the following structure: Each villus is seen to be a tiny conical projection of the mucous membrane (one fiftieth to one eighth of an inch in length). The outside, or surface exposed to the digestive fluids, is composed of a single layer of mucous membrane cells, while the core of the villus is a loose network of blood capillaries and connective tissue cells, together with

one or more larger tubes or lymph capillaries. Like the blood vessels, these latter are branches of larger vessels in the submucous layer; and fluid which once enters the lymph capillaries of the villus flows first into these larger lymph vessels, while these in turn are collected and fuse into a single tube about the diameter of a goose quill, called the thoracic duct. This duct passes up in front of the spinal column and finally opens into a large vein in the neck under the left collar bone. Owing to the milky color of the fluid which is in these vessels and the absence of blood they have received the name of the *lacteals*, and this name also serves to distinguish

them from other parts of the lymphatic system. The blood capillaries as already mentioned are collected together and finally pour their contents into the portal vein. From this description it must be clear that the digested food of the stomach and intestine in order to enter the blood and lymph must pass through two very thin layers, namely, the single layer of mucous cells, which form the lining of the tracts, and

FIG. 46—Section of a villus; *a*, layer of mucous membrane cells which absorb food and carry it to the tubes within; *b*, an artery; *c*, blood capillaries; *d*, a lacteal.

the walls of the blood and lymph vessels themselves. The manner in which this is accomplished involves a discussion of certain peculiar physical processes known as *osmosis* and *dialysis*.

Osmosis and dialysis.—Tie a thin membrane, such as parchment or the skin which a butcher uses to hold sausage meat, tightly about the base of a student lamp chimney. Then, into the top of the chimney fit a cork with an eighth of an inch glass tube passing through a hole in its center. Before fitting in the cork and tube, fill the base of the chimney with water in which has been dissolved some colored salt, such as potassium bichromate. Then insert

the cork and tube. If the parchment is tightly tied there will be no leakage. Finally, suspend this whole apparatus in a jar of clear water so that the liquid in the jar and the chimney is at the same level. If these directions are carefully followed, the result will be an apparatus such as is pictured in Fig. 47, and the liquid in the chimney will have a reddish-yellow color. (See Ex. XXIX.) If we let this apparatus stand for a few hours, at the end of that time we shall be able to observe two curious

FIG. 47—Dialyzer.

results. First, the level of the liquid in the chimney will be higher than that in the jar; second, the clear water in the jar will have become colored like that in the chimney. Now it is evident that what has happened is not due to

ordinary leakage from the chimney, for while such explanation might account for the similarity in color of the two liquids, it would not account for the rise in liquid in the chimney. What has happened is as follows:

In some way, the soluble coloring matter in the chimney has passed through the membrane into the jar of clear water, while some of the water in the jar has passed through the membrane and into the chimney as evidenced by the higher level in the chimney. This tendency of two liquids to mix by passage through a separating membrane is called *osmosis* and the experiment just described is an example of osmosis.

If we continue the experiment and substitute for the potassium bichromate solution other solutions, such as grape sugar, white of egg, starch paste, peptone, etc., we shall obtain varying results. When we test the water in the jar we find that while grape sugar, peptone, and salt solutions pass readily through the membrane into the jar, white of egg and starch paste do not pass through it at all. In other words, peptone, grape sugar, and salt solutions are substances which mix with water by osmosis, while starch paste and albumin do not. Soluble substances which mix by osmosis are called crystalloids; those which do not are called colloids. Grape sugar, salt solution, potassium bichromate solution, and peptone are therefore crystalloids; starch paste and albumin are colloids. Furthermore, observations of the height to which the water rises in the chimney in each of the above cases shows that the rate of exchange is unequal, the water in each case passing through more rapidly than the crystalloid solution, and experiment has shown that this rate of passage is determined by the character of the crystalloid employed. This

relation of the crystalloid solution to the rate of exchange is spoken of as the osmotic pressure of the crystalloid. To say that potassium bichromate has a greater osmotic pressure than peptone means simply that the rate of flow of water through a membrane toward the potassium bichromate solution is greater than it is toward peptone solution. This process of osmosis enables us to separate a crystalloid from a colloid by putting the mixture in a membrane bag and suspending the bag in water. Under those conditions the crystalloid solution will pass through the membrane while the colloid will not. Separation of a crystalloid from a colloid by osmosis is called *dialysis*.

Relation of osmosis to absorption in the body.—If, now, we refer to the conditions in the intestine we shall note that the relation of the digested food to the fluid in the lymph capillaries and blood capillaries fulfils exactly the conditions for osmosis and dialysis. In other words, the mucous membrane and capillary walls furnish the thin membrane separating two fluids, and by dialysis and osmosis the soluble crystalloid substances in the intestine are separated from the colloids and pass into the blood.

In this explanation of the relation of digestion to dialysis is seen a new reason for the action of the various enzymes upon foods. In other words, they not only render them soluble but at the same time transform them into crystalloids in order to make possible this action of dialysis.

While the ordinary process of osmosis accounts for the passing of crystalloid material from the alimentary tract to the lymph and blood, it must be admitted that it does not fully explain the disposal of proteid matter. For example, if peptone were a simple crystalloid sub-

stance like grape sugar we should expect to find it in blood. The fact remains, however, that while we know that peptone leaves the alimentary tract as peptone it is not only *never found in the blood as peptone*, but when injected into the blood *it acts as a poison*. Now, since the only thing that intervenes between the alimentary tract and the blood and lymph are the cells which compose the membranes (mucous membrane and capillary wall cells), it must follow that in passing through these cells the protoplasm in them acts upon the peptone and converts it into another and harmless form of proteid and delivers it in this form to the blood. What the exact nature of this protoplasmic action is, is still unknown. We may, however, sum up the facts known as to transfer of foods as follows:

Summary. Digestion by transforming foods to crystalloids makes possible their transfer into the blood and lymph. This passage may or may not be accompanied by changes in the character of the substances in solution, according as the protoplasm does or does not act upon them in transit. Before leaving this subject, however, it must be stated that since the membranes concerned are made up of separate cells, there may be some seeping of liquid through the spaces between the cells. Such passage of liquids is called filtration, and is not to be confused with osmosis. Since these spaces are exceedingly minute, it is safe to say that the osmotic exchange accounts for the transfer of the major part of the food.

Where nutrients are absorbed.—*Proteid.* While some of the proteid is changed to peptone in the stomach, the amount of peptone absorbed by the blood vessels of the stomach is relatively small, as most of the peptone is

carried into the small intestine with the chyme. In the small intestine, therefore, is found the main seat of proteid absorption. A little may pass into the large intestine and be absorbed there, but this represents a very small amount of the whole, less in fact than in the stomach under ordinary conditions. Wherever it is absorbed it passes into the blood and never into the lymphatics or lacteal system. The reason for this selection is another of the facts not yet explained. In passing through the mucous membrane cells, as already mentioned, the protoplasm of those cells breaks down the peptone into a simpler form of proteid and delivers it in that form to the blood. While this action of the protoplasm is not yet fully explained, indications are that the change takes place in the mucous membrane cells, and is due to an enzyme present in the protoplasm. At present we know nothing of what this enzyme is or its composition. In view of the poisonous action of peptone upon the blood and the system, its importance is evident.

Starch and sugar. By the time digestion is completed in the small intestine, starches and sugars have all been changed to grape sugar, and this substance passes into the blood without change. While some may be absorbed in the stomach and large intestine, the main seat of absorption is the small intestine, as in the case of proteid. Like proteid, grape sugar enters the blood, and not the lacteals.

Fats. Fats are absorbed either as emulsions or soaps, and instead of entering the blood they are the only form of nutrient that passes into the lacteals. In passing through the mucous cells of the villi the soaps are changed back into emulsions so that the fat which enters the lac-

teals is all in the form of an emulsion. It is the milky character of this emulsion which accounts for the appearance and name of this branch of the lymphatics. The seat of absorption of fats is the villi of the small intestine, and the lacteals, which receive it finally, put it into the blood circulation at the opening of the thoracic duct into the left jugular vein. (See Fig. 48.)

Water and salts. These nutrients are absorbed directly into the blood mainly in the small and large intestine. The salts and the water pass through by osmosis owing

to the osmotic pressure of salts contained in the blood. In the case of water, the long time required by the food to pass through the large intestine permits the great bulk of water to become absorbed in this tract. This is further evidenced by the fact that while the chyle which enters the large intestine is very thin and watery, the matter that is passed out of the rectum is mainly solid. Very little water or salt is absorbed in the stomach:

Alcohol. Alcohol is very quickly and completely absorbed by the blood vessels of the stomach, very little ever passing into the intestine. This rapid absorption in the stomach accounts for the quick effect of this substance upon the body. It also has its beneficial side in that, as a solvent of certain drugs, it enables them to be put into circulation quickly.

FIG. 48—Diagram of absorption, showing how food reaches the blood circulation.

Summary of nutrient absorption.—In general, the seat of greatest digestion is the seat of greatest absorption, namely, the small intestine; the large intestine blood vessels taking up only water and such food as may have escaped absorption in the small intestine. Water, salts, grape sugar, and proteid enter the blood capillaries directly and, except in the case of the proteid, without change. All these nutrients are thus conveyed to a large vein called the portal vein which carries them to the liver.

Fats enter the jugular vein by way of the lymph capillaries and thoracic duct (the lacteals) in the form of an emulsion. They do not, therefore, enter the liver like the other nutrients. The nutrients which the portal vein carries to the liver are in the form of water, dissolved salts, blood proteid, and grape sugar.

Action of the liver as an absorbent organ.—The nutrients brought to the liver by the portal vein undergo still further changes before entering that part of the blood circulation which supplies the tissues with food. The most important of these changes is the transformation of grape sugar to a starch or glycogen (sometimes called liver starch). Another change which takes place in the liver is the elimination of poisonous compounds which may be absorbed with the foods, and which, if permitted to circulate to the tissues, would result in disease.

Glycogen.—The cells of the liver take up the grape sugar brought to it by the portal vein, which itself divides in the liver into many tiny capillaries. This grape sugar the liver changes into a carbohydrate of the same composition as vegetable starch, hence the name liver starch. This substance, like vegetable starch, may be changed into grape sugar again by the action of ptyalin and other diastasic enzymes. Acids also transform it directly into grape sugar. It may be distinguished from starch by its action on iodine solutions to which it gives a wine-

red color instead of the blue color of ordinary starch. The manner in which this glycogen is formed from the grape sugar of the portal vein is explained as follows: The protoplasm of the liver cells causes the sugar to lose water, and the result is a chemical change resulting in glycogen.

It has been demonstrated that blood which contains over a certain proportion of grape sugar throws off this excess through the kidneys, and it is thus lost to the body as a nutrient. To avoid this loss the liver cells take up this sugar and convert it into glycogen, which they store in the body of the cell and feed out to the tissue-supply system as it is needed. Before they feed it out they reconvert it into grape sugar, and thus the amount of sugar contained in the blood which supplies the tissues never exceeds a certain proportion (0.1% to 0.2%). The liver, therefore, acts as a storehouse of the carbohydrate or energy supplying nutrient of the body. How it changes the glycogen back to sugar is not fully known, but it is supposed that the liver cells contain an enzyme similar to diastase, which performs this action, as needed. When we consider that the carbohydrates furnish the main part of the energy of the body, this regulative action of the liver becomes very important to the activity of the body.

OTHER SOURCES OF GLYCOGEN.

Experiments have demonstrated that, to some extent, glycogen may be formed also from proteids. For example, when a person suffering from diabetes is fed upon purely proteid foods, all carbohydrates being excluded from his diet, sugar is still formed. This can be explained only on the assumption that it is manufactured from proteid. This manufacture is believed to take

place when the peptone passes through the mucous cells; that there a certain per cent of the proteid is split off as a sugar. If that is the case, it is evident that this sugar may be transformed in the liver to glycogen, and hence proteid is to a certain extent a glycogen former. Some evidence seems to indicate that fat may also be changed to glycogen, but this has not been satisfactorily proved at the present time.

Glycogen is formed in the muscles and in other parts of the body than the liver. In all these cases, it is made from the grape sugar in the blood and acts as a local supply of reserve energy. The great storehouse, however, is the liver.

Poisons.—In the intestines are found many forms of bacteria. These in turn may bring about fermentation of the food, and in this way are produced certain injurious substances or poisons which may be absorbed with the digested nutrients. In that case they are brought in the blood of the portal vein to the liver, and one of the important functions of this organ is to destroy these poisons or return them to the intestine with the bile. If, however, any interruption of the bile making of the liver occurs, it loses its power to destroy and give off these poisons, and the result is that they pass into the tissue-supply blood, and poison the body. The results of the entry of these poisons into the system usually manifest themselves in a coated tongue, loss of appetite, headache, and dullness, and such a condition is called biliousness. Certain drugs have the power to stimulate the liver cells and cause a great outpouring of bile which carries with it the poisons, and thus restores the liver to its normal action and sometimes removes the source of trouble.

Defæcation.—The removal of the fæces from the large intestine results not only in the removal from the body of indigestible material, but also has a vital connection with the poison removing power of the liver. When the bile formed in the liver is poured into the intestine by the gall duct it carries with it the poisons which the liver cells have separated from the blood of the portal vein. If these poisons are allowed to remain in the small intestine they will

again be absorbed, taken to the liver and so on indefinitely, or until the power of removal of the liver is exhausted. The only way in which this can be avoided is for the bile which contains them to be carried into the large intestine and out with the fæces. It is this fact that makes regular operations of the bowels necessary and makes constipation dangerous. In severe cases of constipation stimulants which increase peristalsis, and thus induce defæcation, are often given, but it must be borne in mind that all such remedies are purely stimulants and should be resorted to only under the advice of a physician. Regularity and care in diet will make their use unnecessary in most cases.

HYGIENE OF DIGESTION AND ABSORPTION.

The body requires the digestion and absorption of a certain amount of each kind of nutrient daily. (See p. 70.) If the amount of food taken by a person in a day is only a little in excess of what is required, this excess is readily taken care of by the action of the bowels. If, however, too much is taken, the intestines become clogged and over-worked, the food is not properly digested, poisons are developed and not removed, and all the evils of biliousness and indigestion follow. For this reason a balanced, moderate diet, taken at regular intervals, becomes absolutely necessary to a satisfactory operation of the digestive and absorbent system. Americans are especially prone to over-eat, and many of the ills of our nation have their source in this intemperance in eating.

X. DIGESTION IN THE LOWER ANIMALS.¹

THE process of digestion (as a preliminary step in the changing of food into living matter or protoplasm) is to be met with wherever we find living cells. In man the system has reached its most complex development, but if we examine the various groups of animals, we find that in every group the process of changing food into soluble substances by the aid of enzymes is quite as important and necessary a step in metabolism as it is in man, though the organs which perform this action may be much simpler of structure.

It is interesting, then, to study the methods of digestion in the lower forms, in a comparative way, from the light they throw upon the development of the complex human system. In the pages that follow a few types have been selected to bring out the manner in which the human system has been developed from lower types.

Digestion in one-celled animals.—In animals like the *amæba* and *paramæcium* (one-celled animals) the food is taken into the cell in the form of solid particles. In this respect the protozoans differ from all higher animals. Digestion proper, therefore, in these animals occurs in the cytoplasm of the cells themselves, and to distinguish it from that of other forms is called intracellular digestion.

¹ Chapters X, XIV, XVII, XX, XXII, XXIV, XXVII, and XXXI, are intended for study in connection with a course in zoölogy. In connection with laboratory work in zoölogy they furnish necessary data for comparison. When not preceded by such laboratory work they should be omitted.

While the methods of food taking of one-celled animals differs greatly, some engulfing the food like the amœba, others having a distinct opening or mouth for the purpose, and rows of cilia to direct the water currents toward the mouth, the digestion proper is practically similar in method in all. The amœba will serve for a type.

Digestion of the amœba. — With the ingested food is always taken in some water which collects about the food and forms a food bubble (food vacuole) in the cytoplasm. Soon the cytoplasm around the food vacuole begins to pour into it a secretion which contains hydrochloric acid and probably an enzyme. This secretion is therefore a sort of gastric juice. At the same time the vacuole is kept in motion by the streaming of the cytoplasm. Under the combined actions of this secretion and movement, the solid particle is broken up and dissolved and is then ready to be assimilated directly by the cytoplasm. In this way new protoplasm is built up, and the animal increases in size and strength.

In certain one-celled forms the food vacuoles move in a perfectly definite path (see Fig. 49) through the cytoplasm, and this movement probably serves to distribute the digested food uniformly, as well as to break it up. In such cases the resemblance to a primitive circulatory system is evident.

FIG. 49.—Path of food particles in a paramecium; *N*, nucleus; *M*, mouth; *F. V.*, food vacuoles. Note the decrease in size of the food vacuoles as the food is digested.

Another way in which these one-celled forms differ is in the different nutrients which they are able to dissolve.

The amœba, for example, cannot digest either starch or fats, while other forms have no difficulty in doing so. This is accounted for on the supposition that the amœba can produce no starch or fat dissolving enzymes, while the other forms do produce them.

Digestion in multicellular animals. — As an animal increases in number of cells it becomes manifestly more difficult for the food to reach the cells. In such animals we should expect to find various adaptations to remedy this difficulty.

Hydra and sponges. In the little animal known as the hydra, and in sponges, the cells which form the animal are arranged in layers about a central cavity. (See Fig. 50)

FIG. 50 — Longitudinal section of a Hydra; *b*, bud which will form a young hydra; *ba*, base by which it is attached when not creeping; *m*, mouth; *ov*, ovary with an egg; *sp*, spermary.

The cells lining this cavity are provided with hairlike projections, or cilia, which are in constant motion, and by their motion create a current which draws the water

into the cavity and causes it to circle about, thus bringing the food which it may contain in contact with all the cells which line the cavity. Many of these cells give off digestive secretions which dissolve the food matter contained in the cavity, and thus prepare it for absorption. Such cells represent simple glands, and since the solution takes place outside of the cells it is called extra-cellular digestion, and the cavity becomes a sort of digestive cavity or tract. All animals except the protozoans have extra-cellular digestion. In the forms mentioned (hydra and sponge) the waste is thrown out of the lining cells into the same cavity from which they absorb the food and is removed by the outgoing currents of water.

Worms. In these forms we have for the first time a definite alimentary tract or canal. In some, this canal

FIG. 51—Diagram of a longitudinal section of an earthworm, showing the central digestive tube.

is open only at one end, while in others it forms a continuous tube running the length of the body with a true mouth and anus. Such a continuous tube is found in the earthworm. In this animal the slitlike mouth at the front of the body opens into an expanded, muscular walled cavity just back of it called the pharynx. By the action of these muscular walls the food is sucked into the cavity through the mouth, after it has been

broken up by the lip. From this pharynx, the food is forced backward by true peristaltic movements of layers of muscles in the walls of the alimentary canal until it reaches the anal opening. In traversing this canal it passes through several modified regions where it is acted

FIG. 52—Cross section of earthworm, showing central intestine (*I*) and the dorsal fold or typhlosole (*T*); *D*, dorsal blood vessel; *V*, ventral blood vessel; *B. C.*, body cavity.

upon by special apparatus or secretions, and in this way the digestible material dissolved out and absorbed through the walls of the canal into the body cavity of the animal, where a fluid similar to the blood in our bodies receives it and distributes it to all parts of the body. In the earthworm the regional division of this tract is of par-

ticular interest in showing how the needs of the animal have brought about adaptations of a straight tube to digestive purposes. For example, the widening at the pharynx is a modification to suit the earth burrowing habits of the animal and permit the taking in of food of a solid character. The saclike enlargement back of the pharynx, called the crop, permits the food to be stored for a time in the tract, and enables the animal to eat more at one time than he can digest, and thus give the parts a chance for rest. The thick walled portion back of the crop, called the gizzard, is merely a portion whose muscle layers are thickened to form a special grinding organ for macerating the food, since the animal possesses no teeth. Furthermore, all these modifications are made in a tube of essentially the same general structure as our own tracts, since it has an inner layer of secreting cells, a middle muscle layer, and just outside it is a blood fluid to receive the absorbed food, as in our own body. In fact we might almost say that the earthworm's digestive tract is a living model of what a simple human tract might have been before it was modified into stomach glands, etc.

In the earthworm, therefore, we have a distinct advance over the hydra and sponge in that certain cells have been set aside to make a tube in which to digest the food for the whole body. Further, the manner in which this tube is constructed and modified indicates the way in which the complex system of glands, cavities, and organs of the higher animals could have been developed from such a simple foundation to fit the needs of the animal. As we pass from the earthworm to the higher animals the increased amounts of food necessary to furnish a much

larger number of body cells, the character of the food required by these cells, the opportunities for getting food, and many other factors combine to produce these greater modifications of structure. The nature of these adapta-

FIG. 53—Section of a fish ; *st*, stomach ; *i*, intestine ; *g*, gills. Davison, *Zoölogy*.

tions to increased needs, however, is not different in kind from that indicated in the earthworm. A few types will illustrate the direction which these modifications have taken in different animals.

Modifications in the digestive tube of the fish.—In this animal, the skin that lines the mouth has produced outgrowths of a horny character called teeth, which prevent the food from slipping backward when seized. These are not like our teeth in that they are not set in sockets, and are found all over the surface of the mouth. In the mouth also, is found a fleshy motionless projection called the tongue, which has in it nerves of taste. It has no movement and does not aid in swallowing like ours. Back of the mouth is a widened space or pharynx, in the sides of which four slits are developed for letting in water to the breathing organs or gills. From the pharynx the

food passes backward into a U-shaped portion or stomach. From this stomach the remainder of the tube or intestine passes at first forward and then turns and extends straight backward to the anus. Opening into this forward tube are a number of blind tubes called pyloric cæca. There is also a liver with a large gall bladder connected to this part of the intestine by a duct. The cæca and liver secrete digestive fluids which they pour into the alimentary tube.

The principal advance of this form of tract over that of the worm is seen in the two devices employed to increase the digestive surface, namely, by the folding and by the extension of the walls at points to form external glands. Other adaptations to the new needs are seen in the development of a tongue and teeth in the mouth, and in the increase in size of the stomach portion.

Modifications in the frog.—The most striking adaptations in the digestive tract of the frog are the mouth structures, the increased coiling of the intestine, and the division of the tube into the stomach, small and large intestine. In addition to the liver the frog also possesses a pancreas.

The mouth of the frog contains no teeth with sockets, but there is a single series of projections around the upper jaw which serve, as those of the fish do, to prevent the escape of the food when swallowing. In the frog, the tongue, unlike that of the fish, is movable, and plays a distinct part in swallowing. Unlike ours it is fastened at the front with the tip pointing down the throat. The end is sticky and can be extended to catch insects, which it snaps back into the throat.

The large intestine ends in an enlarged space called a cloaca into which open also the egg or sperm ducts and the

urinary tubes. The anus by which this cloaca opens in this animal, therefore, is the place of discharge of other matter besides indigestible food. The advance of this animal over the fish is seen in the increase of digestive surface

FIG. 54—Section of frog; *t*, tongue; *l*, mouth; *st*, stomach; *p*, pancreas; *d*, duodenum; *si*, small intestine; *r*, rectum, and *b*, bladder, opening into a common cavity, the cloaca; *ur*, ureter, and *ord*, oviduct, also opening into the cloaca; *k*, kidney; *ov*, ovary; *liv*, liver. Davison, *Zoölogy*.

by coiling, the placing of the teeth in relation to the jaws, and the greater tendency to regional arrangement.

Adaptations in warm-blooded animals.—In these animals (with the exception of birds) are found for the first time, true teeth in sockets. All such teeth may be divided into four classes (incisors, canines, premolars, and molars) as in man, and are placed uniformly upon the jaws. The kind which predominates in any animal is determined by the kind of food which it eats. Thus, squirrels and rats have the incisors greatly developed for gnawing, while the flesh eaters are marked by the predominance of canines, and the grass eaters by the molars.

In birds the teeth are replaced by horny sheaths, the mastication of the food being provided for by a thick walled portion of the digestive tract called the gizzard.

FIG. 55—Skulls, showing teeth; *A*, monkey; *B*, horse, a grass eater (note number of molars); *C*, dog, a flesh eater (note the prominent canines); *D*, rodent, a gnawer (note the prominent incisors).

While the other parts of the digestive tract are, in general, similar to that in man, the stomachs of the various forms show interesting modifications. The stomach of the rabbit and other rodents, and of the carnivora, is almost identical with that of man. The birds show a curious adaptation to compensate for the lack of teeth and the uncertainty of food supply. In these animals the stomach is in three parts. First, at the base of the gullet is a sac called the crop in which the food may be stored

until needed by the body. By means of this organ the bird is enabled to eat enough at one meal to last for some

time. Succeeding this is a compartment called the gizzard. When food is needed it is passed from the crop into this gizzard which has an outer wall of thick muscles and an inner horny lining. Inside it, too, are stones and other hard objects which the bird swallows for the purpose of grinding the food thoroughly to prepare it for the action of the digestive fluids. When thoroughly ground up, the food passes into the third compartment, or true digestive stomach, where the gastric juice is secreted.

FIG. 56 — Digestive tract of a bird ; *a*, esophagus ; *b*, crop ; *c*, proventriculus ; *d*, gizzard ; *e*, liver ; *f*, gall bladder ; *g*, pancreas ; *h*, duodenum ; *i*, small intestine ; *k*, caeca ; *m*, ureter ; *n*, oviduct ; *o*, cloaca.

show still another curious modification. In these animals, such as the cow, the stomach is divided into four compart-

The ruminants or cud-chewing animals

ments. The first and largest is called the rumen, and into this the grass mixed with saliva passes without mastication as it is eaten. From this rumen the food is passed into a second compartment with a honeycombed wall and called the reticulum. After the grazing is over the food is returned to the mouth from this reticulum in the form of boluses or round balls called the cud. This cud is then thoroughly masticated by the molars and finally swallowed again. This time it passes directly

FIG. 57 — Stomach of a sheep (ruminant); *e*, esophagus; *Ru*, rumen; *R*, reticulum; *Ps*, psalterium; *A*, abomasum; *P*, pylorus; *D*, duodenum.

into a third compartment called the psalterium, which is ridged, and strains the food into the fourth compartment, the abomasum or rennet stomach. In this abomasum the actual digestion takes place, and here is secreted the gastric juice. Such an arrangement as this is very evidently an adaptation to compensate for the greater difficulty of digestion of such food as grass and hay; but it is to be noted that in none of these cases is the purpose and general method of digestion essentially different.

In all, the various structures are working to one common end, namely, the transforming of food into soluble matter. And the ultimate method employed is the action of ferments or enzymes secreted by cells located in the walls or connected with the digestive tract.

In all the warm-blooded animals the intestines are much longer and more coiled than in the cold-blooded forms, and the time required for the digestion of vegetable matter is responsible for the greater length of the intestines in the herbivorous animals as compared with the carnivora.

In conclusion, we may note that all modifications of digestive methods are merely adaptations to meet special conditions. These adaptations may take the direction of specialized structures for producing digestive fluids or preparation of food for the action of fluids, or they may be directed toward increase in surface and the insurance of more complete digestion. In every case, however, the transformation of food into soluble matter as a preliminary to absorption is the end sought, and the ultimate chemical means is the action of special compounds called enzymes secreted by specialized cells.

XI. BLOOD AND LYMPH.

AFTER digestion and absorption the food is ready for the use of the cells. Absorption has placed this food in the blood for delivery to the cells, and it is in its character as a carrier of digested nutrients and other substances to the cells that we are to investigate the functions and structure of the blood.

Blood vessels. — The blood in our bodies is contained in a set of tubes called blood vessels, and these vessels are so constructed and connected as to form a completely closed system. In other words, while the blood absorbs matter into itself and gives out matter from itself to the cells of the body, this exchange takes place through the walls of the blood vessels, and never, except in case of accident (bleeding), does the blood (as blood) escape from the system of closed tubes. In this respect it is like a train on a circular track. We may load goods into this train and take goods from it, but the train itself keeps on continually and, except in case of accident, never leaves the track. In our body the blood is kept in constant movement within the tubes by the beating of the heart and for this reason the blood vessels and their contents are spoken of collectively as the *circulatory system*. If it were not for this constant motion it would be manifestly impossible for food taken in from the alimentary tract to reach the other parts of the body. Before considering the way in which this liquid is kept in circulation we shall first consider its composition.

Blood structure. — We may readily obtain human blood for study at any time by pricking the end of the finger with a clean needle. (See Ex. XLII.) If we examine a little of the blood so collected on a flat surface we note that the drop is not of the same color throughout. At the edge of the drop it shows a yellowish color while the center is dark red. With the compound microscope we find that this difference in color is due to the fact that blood consists of two distinct parts, a yellowish liquid called the plasma, and a vast number of tiny round disks called corpuscles. The red color of the blood is due to the presence of these corpuscles (red corpuscles). If we use the higher powers of the microscope we find that these red corpuscles are actually biconcave disks (see Fig. 58)

FIG. 58.—Blood corpuscles; *A*, red corpuscles in rouleaux; *a, a*, colorless corpuscles ($\times 400$); *B*, red corpuscles, highly magnified; *C*, view of edge; *D*, three quarters view; *E*, red corpuscle swollen with water; *F*, *G*, *H*, distorted red corpuscles.

massed together in bands or rouleaux. At the same time, we may be able to make out one or more colorless bodies of irregular shape floating among these bands of disks. These latter are called white corpuscles to distinguish them from the other more numerous form. They differ from the other corpuscles in having a nucleus. Both kinds of corpuscles are cells, but the red corpuscles have

lost their nucleus. The presence of these cells allows us to consider the blood as a true tissue.

In general, then, blood is a tissue consisting of a yellowish liquid (plasma) in which float a large number of non-nucleated cells (red corpuscles) and a few nucleated cells (white corpuscles).

Properties of blood. — If we mount a drop of blood on a glass slide and allow it to stand for a few moments, at the end of that time we note two distinct changes in its appearance. First, it is darker in color, and second, its surface is covered with a film which, when pricked with a needle, shows an elastic consistency. This formation is called a clot. If we examine the needle puncture on the finger we shall find that it has ceased to bleed, and is covered with a similar clot. If we wash this clot off the finger, bleeding begins anew, and this gives us a hint of an important use for this clot formation. An examination with the microscope of the clotted blood on the slide shows that it is actually a complicated network of slender threads in which are entangled the corpuscles. Evidently these threads must have their origin in the plasma, and to understand their nature it is necessary to make a careful examination of the plasma. For this purpose a larger quantity of blood is necessary.

Study of blood. — Fortunately for our purpose, the blood of all warm-blooded animals is practically identical in structure with human blood. This fact permits us to use fresh-drawn blood of a pig or cow. (See Ex. XLIII.) First, place some blood in a bottle and let it stand for a time exposed to the air. Put the rest in a basin and whip it vigorously with twigs. Pour the whipped blood into another bottle. While these two bottles are standing we

may examine the twigs with which we did the beating. These will be covered with a stringy elastic substance which, when washed, is seen to consist of glistening white strands or fibers. These fibers are composed of a substance called fibrin, and are identical in composition with those which were noted in the clot of human blood. Furthermore, if we test this fibrin with our nutrient tests, we shall find that it responds strongly to one test only, namely, that for proteid. In other words, fibrin is a coagulated proteid which was present in the plasma of fresh-drawn blood in a liquid state. If we examine the bottled blood after it has stood for awhile we shall find the contents of the bottle filled with fresh-drawn blood in a different condition from the whipped blood. The former will be found to have solidified into a clot of the same shape as the bottle, and to be floating in a small quantity of a straw-colored liquid (*serum*). The whipped blood will show no clot, and is in the same condition as when we put it in the bottle.

If we pour off the serum from the first bottle, and then break the bottle so as to remove the clot entire, we may examine this clot and study its structure. Such examination shows it to be made up of a solid mass of fibrin fibers which inclose masses of corpuscles. The corpuscles give it its color. The whipped blood fails to clot because we have removed its fibrin by whipping. From these observations we must conclude that the serum is plasma whose fibrin has been removed, and that the presence of this fibrin is necessary to the formation of a clot.

Our experiments thus far have succeeded in separating only one of the constituents of blood plasma, namely, fibrin.

If our nutrient tests are applied to the serum which was poured from the bottle containing the clot they furnish the following interesting results:

First. They show it to be rich in proteid.

Second. They indicate small amounts of fats and grape sugar.

Third. They show the presence of salts.

Fourth. They show that starch is absent.

Fifth. The serum gives a weak alkaline reaction.

These results justify the following conclusions in regard to the composition of blood plasma:

First. The proteid tests show that the plasma contains other proteid matter beside the fibrin, since that was removed by the clotting, and that this proteid matter does not coagulate like fibrin when the blood is removed from the living body.

Second. The small amount of grape sugar in circulation at any time is what we should expect from our study of the formation of glycogen in the liver (see p. 142), while the small amount of fat present shows that most of the fat poured into the blood by the lacteals does not enter into the general circulation but is used up by the living tissues.

Third. The absence of starch is accounted for by the fact that all the starch was changed by digestion into grape sugar and enters the blood in that form.

Fourth. The presence of salts explains the osmotic pressure of the blood, accounts for its alkaline reaction, and serves to keep certain proteids in solution.

Finally, the presence of these particular nutrients shows in an indisputable way that it is the plasma of the blood which is the carrier of digested nutrients.

While our experiments show in a general way the composition of the blood, we must go to the chemist for a careful analysis of its structure. Such chemical analyses show the plasma to be composed of the following substances:

COMPOSITION OF PLASMA.

Plasma	{	Water .	{	Fibrinogen	}	of	{	Sodium
		Proteids .		Paraglobulin (serum globulin)				
				Serum albumin				
		Extractives	{	Fats				
				Sugars				
				Urea				
				Acids				
	{	Salts . . .	{	Chlorides	}		{	Potassium
				Carbonates				Calcium
				Sulphates				Magnesium
				Phosphates				Iron
	{	Enzymes and unknown substances						

Blood proteids. — *Fibrinogen* is fibrin in the liquid state. It is, of course, absent in the serum. It varies in amount from .22% to .40% in human blood, and to it is due the clotting power of the blood. (The causes of clotting will be discussed under a separate heading, see p. 169.)

Serum albumin is the form of proteid which is supposed to have been formed from the digested proteids of the food. It forms 4.5% of the blood of man, and is supposed to be the form of proteid used in supplying the tissue cells.

Paraglobulin forms 3.1% of the blood of man. Its origin and function are at present unknown.

The two proteids, then, whose functions are definitely known are serum, albumin, and fibrin.

Blood extractives. — The fats and sugars are evidently absorbed from the digestive fluids, and as nutrients are carried to the tissues for use as energy producers.

Urea and acids are merely forms of waste matter given off by

the cells to the blood, and their presence indicates another function of blood, *viz.*, the removal of wastes. (See p. 352, Excretion.)

Blood Salts. The mineral salts are probably absorbed in the same way as the other nutrients, and their presence accounts for the osmotic pressure of the blood as well as for the alkaline reaction. This last is mainly due to the presence of the salt, sodium carbonate. Certain proteids (notably the globulins) are kept in solution by the presence of salts.

Water of blood. — This liquid is taken in by osmosis from the digestive tract and other parts of the body, and the liquid character of blood is mainly due to it. It forms 78% of the entire blood.

The solid matters in the blood. — Having now a general idea as to the nature of the liquid plasma, let us turn to the solid parts. We have already noted that blood contains two kinds of corpuscles, the red and the white. In addition to these solids, blood contains a third set of bodies called blood plates. These three solids are only slightly heavier than the plasma, and the movement of the blood keeps them pretty thoroughly mixed with the plasma. A knowledge of their origin and function is necessary to a clear understanding of the blood functions.

Red corpuscles. These bodies to which the red color of the blood is due, are shown by the microscope to be biconcave, circular disks which are usually banded together in rolls like a roll of coins. (See Fig. 58.) They are very small, having an average diameter of about .0077 millimeter, and while their number varies greatly with the condition of health the average is five million to a cubic millimeter of blood in males, and four million and a half per cubic millimeters in females. When viewed singly they are seen to be of a yellowish-red color, but in masses they appear deep scarlet. This red color is due to a pigment which they contain called hæmoglobin. Hæmo-

globin gives them an important function as carriers of oxygen from the lungs to the tissues, and their physiology is largely concerned with the action of this pigment.

In structure they are found to consist of a colorless substance called the stroma, which gives shape to the corpuscles, and in this stroma is deposited the hæmoglobin together with some water and salts. They contain no nucleus, and are to be considered as cells which have been modified to meet the needs of the blood.

Hæmoglobin. This pigment consists of two chemical compounds, hæmatin and a proteid body. The hæmatin has the power when in contact with oxygen to combine with it and to form an unstable compound which breaks down when it reaches the tissues, and gives up its oxygen to the cells. This power of hæmatin to unite with oxygen is due to the iron which forms a large part of its chemical composition. An interesting demonstration of the power of the red corpuscles to take up oxygen is furnished whenever we mix blood with air. For example, shake violently a tube half filled with blood. The increased redness of the shaken mixture is due to the taking up of oxygen. In the body the red corpuscles are brought in contact with the air in the lungs, and the oxygen which they pick up is carried by them to the tissues and there given up to the cells as needed. It is evident, then, that one function of the red corpuscles is to carry oxygen to the tissues. This, however, is only one of their functions.

Hæmoglobin forms compounds not only with oxygen but also with carbon dioxide. This compound is called *carbohæmoglobin* to distinguish it from *oxyhæmoglobin*. This compound is formed when the carbon dioxide is absorbed from the tissue cells. In this way the red corpuscles are

not only enabled to carry oxygen to the tissues but also to remove part of the waste carbon dioxide from them.

In summary, then, we may say that the two uses to the body of the red corpuscles are the carriage of oxygen to the tissues and the removal of carbon dioxide from them.

The red corpuscles of man and of all mammals, except the camel family, are biconcave, circular disks without nuclei. In the camel family, they are elliptical in shape. They vary, however, in size, and this variation often enables experts to determine the character of blood stains in murder trials. The red corpuscles of fishes, amphibia, reptiles, and birds are oval in shape and usually nucleated.

FIG. 59— Various types of red corpuscles.

Origin of red corpuscles. — The fact that the human corpuscles are without nuclei shows that they are cells which are in a degenerating state and hence cannot last long in circulation. Since

they are so necessary to the life of man it is evident that there must be a constant renewal of the supply from some source. This source is found in the red marrow of bones. Here are produced nucleated, colorless cells, called erythroblasts, which multiply and form daughter cells. These daughter cells finally produce hæmoglobin in their cytoplasm, and thus form nucleated red corpuscles. These lose their nucleus and finally enter the blood stream in the form we have recognized as red corpuscles. The old corpuscles are continually breaking down, and after undergoing dissolution in the blood stream, the liberated hæmoglobin is discharged by the liver as bile pigment. It was once thought that their destruction took place in the spleen, but there is no evidence to prove this and it is now believed that this destruction may take place anywhere in the blood stream. In a normal human being the supply and loss is so balanced as to maintain a fairly constant number of corpuscles in the blood.

The number of red corpuscles in the blood varies not only with the sexes, as already noted (see p. 165), but also with other conditions. They increase greatly with the altitude, supposedly to compensate for the reduction in air supply. They often decrease in number in disease causing a condition known as anemia. At present the remedies for lack of blood corpuscles usually take the form of tonics, that is, the giving of medicines, usually rich in iron, which stimulate the formation of both erythroblasts and hæmoglobin. Exercise in the open air is a far better remedy than any drug for this condition.

White Corpuscles.—These bodies are not nearly so numerous as the red corpuscles being only five thousand to seven thousand per cubic millimeter. They are colorless amœboid cells having a nucleus and no definite cell outline, moving about in the plasma in much the same manner as an amœba, which they strongly resemble. They are of two kinds: those with granular cytoplasm are more numerous and are called leucocytes; those without granules in the cytoplasm are called lymphocytes. The leucocytes

develop from bone marrow; the origin of lymphocytes is uncertain. The leucocytes have various functions. Some, called phagocytes, absorb disease-causing bacteria as their food, and thus protect the body from disease. Others aid in the absorption of fats and proteids from the digestive tract. They also aid in coagulation or clot forming and finally help to maintain the normal supply of proteid in the blood plasma.

Blood plates.—These are circular or elliptical bodies smaller than red corpuscles. They are more numerous than the white corpuscles, and have been shown to be capable of amœboid movements. They disintegrate rapidly when removed from circulation, and hence little is known of their structure. Their principal function appears to be their action in clotting, which we will discuss under that head.

FIG. 60—A phagocyte (white corpuscle), *A*, engulfing a bacterium, *B*; 1, 2, 3, steps in the process.

Theories of clotting.—In the beginning of the chapter we noted that blood removed from circulation tended to solidify into a clot. We noted also that the foundation of this clot was the coagulated fibrinogen or fibrin, of the blood plasma, and that the failure of whipped blood to clot was due to the removal of this fibrinogen by whipping. These experiments, however, merely give us an explanation of what the clot is made of, namely, of fibrinogen. The fibrinogen is made to coagulate and forms fibers of solid proteid matter called fibrin, between whose strands are caught the corpuscles, and to whose shrinking is due the squeezing

out of the serum or defibrinated plasma. What the cause is that suddenly produces this coagulation of fibrinogen, and the consequent formation of fibrin fibers, the experiment failed to explain. While this question has not yet been fully answered the theory which is generally accepted at the present time may be stated as follows:

The experimental facts upon which the theory is based are three. First, the coagulation of fibrin is not due to exposure to air, as it takes place with equal readiness when air is not present. Second, it occurs only when there are present calcium salts, fibrinogen, and an enzyme called thrombin. Third, the enzyme thrombin is not a part of the normal composition of plasma, but is furnished under special conditions by the combined action of the blood plates, the leucocytes, and the calcium salts.

The formation of the clot involves the following actions: First, the breaking down of the blood plates and leucocytes in the presence of calcium salts results in the formation of a chemical compound or enzyme called thrombin. The enzyme reaction of this thrombin causes the coagulation of fibrinogen into fibrin. The reason that this action occurs only when the blood is removed from the blood vessels is probably due to the fact that the blood plates and leucocytes do not disintegrate in sufficient numbers while the blood is in the body to produce much thrombin. It is also possible that in blood which is inclosed in blood vessels there may be compounds among the unknown constituents of plasma which neutralize the thrombin as fast as it is formed.

Conditions affecting the blood.—Since the blood furnishes the oxygen and food supply of the tissues, it is evident that it must be maintained in such a condition as to perform these functions. Anything, therefore, which interferes with the number of corpuscles, the presence of hæmoglobin, the oxygen content of the corpuscles, or the food content of the plasma, is bound to interfere with its normal functions. The conditions which have been found to be essential to a healthy condition of blood

are an abundance of fresh air to secure sufficient oxygen supply, plenty of sleep to prevent too great a drain by the tissues upon the food and oxygen it carries, exercise to allow the proper distribution of food and oxygen to all parts of the body and the removal of waste, and nutritious food for diet to maintain the proper food content of the plasma. This latter condition also affects the red corpuscles as well as the plasma, as without iron it is impossible to form hæmoglobin.

Amount of blood in the body.—We might expect that the constant giving up of food and oxygen by the blood would cause its quantity to vary. So far as known, however, the loss of these substances and the acquiring of wastes is so balanced that the quantity is fairly constant, and in a healthy person amounts to about one thirteenth of the total weight of the body. The amounts in different parts of the body, however, vary. Roughly, it is distributed in the following proportions:¹

Heart and Lungs	25%
Liver	25%
The Skeletal Muscles	25%
The Other Organs	25%

Blood Temperature.—Heat is constantly being produced in all of the animal tissues of the body, and at the same time heat is being given off to the outside air from the surface of the body. The amounts produced and lost vary in different parts of the body. Now, the blood is passing constantly from one part of the body to the other, and hence necessarily carries warmth from the parts where greater amounts of heat are produced to places where there is less. In this way it tends to equalize the temperature in all parts of the body, and the temperature of the

¹ Foster, *Physiology*.

blood is therefore the average temperature of the whole body. In man the production of heat and the loss to the outside of the body is so nicely balanced that the average or blood temperature remains almost constant. This being the case, any rise in this temperature indicates a more or less serious disturbance in the body. In some animals, such as the fishes, frogs, and reptiles, the loss of heat is so great compared with the production that the temperature of the blood is only slightly above that of the surrounding air or water. Such animals feel cold to the touch, and are spoken of as cold-blooded animals. They are also called variable temperature animals, since their average temperature varies with that of their surroundings. In birds and mammals, including man, the amount of heat produced is greater and the loss so nicely balanced that their average temperature is not only kept constant, but at the same time is higher than that of their surroundings and does not vary with the outside temperature. Such animals feel warm to the touch and are therefore called warm-blooded.

In man the temperature of a healthy body remains at about 98° Fahrenheit, and though it may increase to 103° or 105° in violent exercise, this is only temporary and it sinks in a few hours to normal. A continued temperature of over 100° is usually an indication of fever. This fact enables the physician to discover at once the presence of fever by taking a person's temperature. In birds the temperature is much higher than in man, these animals having an average temperature of 105° Fahrenheit.

Summary of blood action.—The blood is usually spoken of as the nutritive fluid of the body, since its main function is to supply food or nourishment to the tissues. It has, however, other func-

tions as we have seen, and we may summarize its functions as follows:

(a) It carries to the tissues the food which has been prepared by digestion, and introduced into the plasma by absorption.

(b) It carries oxygen, which it takes from the air in the lungs, to the tissues and there exchanges it for carbon dioxide which it carries away. The red corpuscles are the carriers in this case.

(c) It carries from the tissues to the lungs, kidneys, and skin, the wastes of the body. Such wastes include carbon dioxide, urea, acids, etc., and are removed from the body by the organs mentioned.

(d) It is the medium for the transportation of the internal secretions of certain glands.

(e) It distributes the heat produced in various parts of the body and thus tends to equalize the body temperature.

Blood in other animals.— Since the function of the blood is the distribution of food and oxygen, and the collection of wastes, it is evident that animals composed of a single or few cells will have no need of such a fluid. In animals like the *amœba*, and even in animals of more complicated cellular structure, such as the hydra, sponges, and jellyfishes, the digested food and the air are brought in direct contact with the cells, and hence in these animals there is no blood and no need of such fluid. As animals increase in complexity, some of their cells are necessarily so placed that the food is no longer brought in direct contact with them. In such animals some system of distribution becomes necessary, and hence most animals higher than those mentioned have some kind of fluid that answers the purpose of the blood in our bodies. In many this blood fluid is not inclosed in tubes and is without corpuscles. It is to be compared rather with the plasma than with the red blood of man. Crayfishes, crabs, and insects have such a fluid, and it is colorless. In the earthworm the blood fluid is red, but has no red corpuscles, the hæmoglobin being dissolved in the fluid itself. It is only when we reach the vertebrate or backboned animals that we find blood containing corpuscles specialized for the carrying of oxygen and carbon dioxide. In the lower forms of vertebrates these corpuscles are nucleated. Only in birds and mammals do we find the non-nucleated corpuscle.

THE LYMPH.

In the pages preceding we have attempted to make it clear that the materials required by the tissues are brought to them by the blood, and that the wastes (carbon dioxide, urea, etc.) given off by the tissues are finally conveyed by the blood to organs which remove them from the body. In the lower animals where the blood fluid is not inclosed in tubes, and bathes all the tissues, this exchange between tissue cells and blood fluid is a comparatively simple matter. In man and the higher animals the fact that the blood is contained always in closed tubes makes it less easy to see how this exchange takes place. The explanation of this exchange involves a study of another important liquid of the body known as the lymph. All the tissue elements of the body are bathed in this liquid. Furthermore, the lymph spaces are connected by a set of tubes known as the lymphatics, which permit the flow of the lymph from one part of the body to another.

Origin and structure of the lymph. — Examination of the composition of the lymph contained in the lymph tubes and spaces shows that it is practically identical in structure with the plasma of the blood, and in fact it is supposed to be plasma which has in some way passed out of the blood vessels and become a tissue bathing fluid. It was once thought that the lymph in the tubes and the lymph in the spaces were in direct communication. Recent investigation has shown that the lymph tubes are closed as completely as the blood vessels, but the lymph contained in them is identical in structure and origin with that in the spaces. How has the plasma that is contained in the blood vessels been able to escape through the walls

into the lymph spaces, and how does the lymph of these spaces pass into the lymph tubes? The answer to this question involves the whole mechanism by means of which the tissue elements receive their food and get rid of their wastes. In Figure 61 the relation of a lymph tube, blood tube, lymph space, and tissue element (cell) to one another is indicated in a diagrammatic way. Let us ex-

FIG. 61 — Diagram of lymph relation.

amine this diagram for a moment, to see what explanation it suggests for this problem.

The first point to be noted is that between the lymph in the space and the plasma in the blood tube there is only the thin membrane of the tube wall. Secondly, between the lymph in the space and the liquid contents of the cell is a second membrane, the cell wall. Finally, between the lymph in the space and that in the lymph tube is a third membrane, the lymph tube wall. If we recall what we learned in regard to dialysis and osmosis (see p. 136) it is evident that the above relation satisfies the conditions necessary to those processes. By osmosis the dissolved gases and some of the nutrients of the plasma together with the water, pass through the walls of the blood vessel into the lymph in the space, while in similar manner the wastes in this fluid pass into both the blood vessel and lymph tube. Furthermore, through the cell wall, the

nutrients of the lymph in the space are exchanged for the wastes of the cell. In this manner then the plasma of the blood tube becomes lymph and is in turn passed on into the tube, which drains it away to other parts of the body.

Another feature of this exchange is seen in the action of the corpuscles. It will be noted that the red corpuscles do not leave the blood tube, but the white corpuscles or leucocytes are able to pass through tiny spaces in the walls. The reason for this is that the leucocytes are able to change their shape and to push themselves through holes too small to let the rigid red corpuscles pass. Certain proteids which are not dialyzable also probably pass through similar spaces in the blood tube walls by filtration.

Another feature to be considered in this diagram is the disposal of the wastes. If the liquids were station-

FIG. 62—The lymphatic system; *a*, union of left jugular and subclavian veins, and point of union with the thoracic duct, *b*; *c*, right jugular and subclavian veins, and point of union of right lymphatic system; *d*, receptacle for food absorbed from the intestine; the oval bodies are glands.

ary the exchange would go on until all reached the same

composition. The fact that the lymph tubes are continually draining off the liquid they receive, and that the plasma is being constantly renewed, together with the varying activity of the cell, makes the composition of the three liquids subject to constant change. It is this change that gives a definite direction to the flow. While it is true that some substances pass from the lymph tubes into the space, and from the space into the blood tube, the prevailing direction of flow is from blood tube to space to lymph tube, and in this way most of the cell wastes are carried into the lymph tubes. What then becomes of these wastes which enter the lymph tubes?

The lymphatics. — The lymph tubes just referred to are only the ultimate tiny endings of a branching network of tubes called the lymphatics. From all parts of the body these tiny tubes (lymph capillaries) are collected together, and finally empty their contents into two large trunks on each side of the body. These trunks are of different size, the one on the left side with its branches being much the larger. It is this left branch that includes the system of branches called the lacteals (see p. 135). Both the right and left branches open finally into two veins on the right and left sides of the neck. In short, then, the waste laden lymph is finally drained from all parts of the body and poured into the blood again at such points that the wastes must pass through the lungs before entering the general circulation and are finally eliminated by the kidneys.

FIG. 63 — Lymph tubes, showing valves, *d*.

Lymph flow. — Unlike the blood system, there is no

apparatus for pumping lymph. The lymph vessels are, however, provided with valves which allow it to flow only toward the veins. While the flow is irregular and slow the pressure in the capillaries is sufficient to keep it in motion, and the movements of muscles that press upon the tubes also aid in driving it onward to the veins.

Since the lymph is laden with waste, it will be recognized that anything which interferes with this flow is bound to

disturb the health of the body by the over-accumulation of waste in the spaces. Muscular exercise is necessary, since it maintains the lymph flow.

At irregular intervals the tubes open into spongy bodies, known as lymph glands or nodes. These nodes are spongy masses of fibers filled with corpuscles similar to the white corpuscles of the blood. They are supposed to have two functions, first, the removal of certain wastes from the lymph, such as bacteria, which the corpuscles devour, and second, the production of the corpuscles themselves. In other words, these glands filter the lymph as it

FIG. 64—A lymph node.

passes through them, and tend to remove harmful bacteria in this way. Occasionally these nodes become overcharged with poisonous material, and break down, forming abscesses or boils.

Summary of the lymph relations. — From the preceding it is evident that the most important action of lymph is to act as a middleman for the exchange of nutrients and wastes between the blood on the one hand and the cells on the other. Furthermore, the waste matter is in turn taken up by the lymphatics, which drain it off into the veins. In cases of dropsy the failure of the drainage system to remove these wastes causes an accumulation of lymph in the spaces, and often causes the body to swell to an enormous extent. This relation of the lymph spaces and lymphatics also explains why medicines injected under the skin are so quickly distributed to all parts of the body, the lymphatics taking it up and distributing it to the tissues much more quickly than if it had entered the system through the digestive tract.

Certain parts of the body pour into the lymph more of their wastes and secretions than others, and this accounts for the variation in composition of lymph in different parts of the body, but the foundation of the fluid is the blood plasma.

XII. CIRCULATION.

ANATOMY OF THE HEART AND BLOOD VESSELS.

ENOUGH has been said in our study of the blood to make clear that its function is the distribution of food and oxygen to the tissue cells, and the removal of waste from these cells. In other words, its purpose in the body is distribution. Furthermore, it is clear that in order that distribution be made to all parts of the body, the blood must penetrate to all these parts, and its flow to and from these points must be continuous. Such conditions are satisfied by inclosing the blood in tubes of varying size and structure, so arranged as to enter all parts of the body, and connected in such a way that a pump placed in one part of the system will force the blood in a constant stream from the pump to these places and back to the pump again. In our body this closed system of blood vessels is spoken of collectively as the circulatory system, while the pump which keeps the blood in motion is called the heart. The various kinds of tubes are called veins, arteries, and capillaries, according to the direction of blood flow. Let us first examine the structure and workings of the heart, or pump.

Anatomy of the heart. —(See Ex. XLVII.) This organ is a hollow, muscular sac about the size of a large pear, lying in the chest cavity just above the diaphragm, and immediately behind the lower two thirds of the breast-bone (see Fig. 65). It is shaped somewhat like a cone

with the smaller end below and slightly to the left of the breastbone. It is this position of the point that makes the

FIG. 65—External view of the heart; *L, L*, lungs; *RA*, right auricle; *LA*, left auricle; *RV*, right ventricle; *LV*, left ventricle; *IVC*, inferior vena cava; *SVC*, superior vena cava; *Ao*, aorta; *PA*, pulmonary artery; *Tr*, trachea.

beating more distinctly felt on the left side of the body, but the bulk of the organ is in the middle of the cavity.

The heart is surrounded by a sac called the *pericardium*.

This is conical like the heart, but the broad end is downward and attached to the diaphragm. Both the inside of the pericardium and the outside of the heart are covered with smooth slippery membranes, while between these two layers is a fluid which moistens the surfaces and lessens

the friction between them as the heart beats. The heart is also lined internally with fibrous membrane while the walls between these outer and inner membranes are composed of a peculiar kind of striped muscle tissue whose cells differ markedly from the striped cells of ordinary voluntary muscle. These muscular walls contain nerves, blood vessels, and connective tissue. Their thickness varies markedly in different parts of the organ.

FIG. 66 — Relation of cavities in one side of the heart.

Internal structure of the heart. — (See Fig. 66.) On opening the heart, it is found to be completely separated into two distinct halves by a longitudinal partition or septum. Each of these halves, which are right and left in position,

is divided by transverse partitions into an upper and lower cavity. The cavities which are at the broad end of the heart are thin walled and flaplike in appearance. From a fancied resemblance to the ears of a dog they are called *auricles*. The other two cavities and the thick walls of muscle which surround them, make up the bulk of the heart. These latter cavities are called

ventricles. In the partition between the right auricle and right ventricle is an opening guarded by a three-cornered valve known as the *tricuspid valve*. The left auricle and left ventricle communicate by a similar opening, and the valve which guards this opening is called the *mitral valve* from its resemblance to a miter. The auricles receive the blood from veins, and this blood passes from the auricles into the ventricles through the valves. The ventricles in turn, by contraction of their muscular walls, force this blood out of the heart into arteries and thus over the body. It is the pumping action of these ventricles which constitutes what is known as the heart beat. The two sides of the heart beat in unison.

FIG. 67 — External view of the heart; 1, right ventricle; 2, left ventricle; 3, right auricle; 4, left auricle; 5, aorta; 6, pulmonary artery; 7, innominate artery; 8, carotid arteries; 9, subclavian artery; 10, superior vena cava 11, pulmonary veins.

External anatomy of the heart. — (See Fig. 67.) When the heart is examined from the outside, the position of these four internal cavities is seen to be outlined by grooves

on the surface. The two flaplike auricles at the top of the heart are separated from the ventricles by a deep groove, while a shallow furrow marks the division of the right and left ventricles, and the position of the septum.

On the outside may also be seen a number of large blood vessels which extend from the top of the heart. In addition to these there are smaller blood vessels (the coronary system) which branch

over the surface of the heart and supply it with blood. The outside is also more or less coated with fat.

Relation of blood vessels and heart cavities. — If we push bristle seekers through the external blood vessels, and then carefully dissect away the two sides of the heart we can make out the course of the blood through these vessels and the heart. Let us first examine the right side. (See Fig. 68.)

Following the course of the seekers we find that two large veins, called the *superior vena cava* and *inferior vena cava*, open into the right auricle, while at the back of the auricle is the opening of a small vein (the *coronary vein*).

Blood which enters the heart from these three veins passes

FIG. 68.—Right side of the heart (dissected); 1, superior vena cava; 2, inferior vena cava; 3, right auricle; 4, tricuspid valves; 5, papillary muscle fastening the threadlike chordæ tendinæ, which are, in turn, fastened to the edges of the valve; 6, pulmonary artery; 8, aorta; 9, branches of the coronary artery; 10, left auricle; 11, ventricle; 12, semilunar valves.

through the opening guarded by the tricuspid valve into the right ventricle. This tricuspid valve is so constructed as to open to admit the blood from the auricle when the ventricle walls expand, and to prevent its flow backward into the auricle when the ventricle contracts. It is in the form of a curtainlike flap which is fastened by chords (*chordæ tendinæ*) to raised prominences on the ventricle (*papillary muscles*).

If we follow the seekers from the right ventricle they extend outward through a single artery known as the *pulmonary artery*, since it conveys blood from the ventricle to the lungs. This artery has at the base series of valves or pockets of a half-moon shape (*semilunar valves*). These valves are so arranged as to permit the flow of blood from the ventricle into the artery when the ventricle contracts, and prevent its backward flow when the ventricle relaxes.

The left side of the heart shows the following connections (see Fig. 69): Into the left auricle open four veins (the *right and left pulmonary veins*) so-called because they bring blood to the auricle from the right and left lungs.

FIG. 69.—Left side of the heart (dissected); 1, pulmonary veins; 2, papillary muscles; 3, cut surface of the left ventricle walls; 4, semilunar valves at the entrance to the aorta; 5, pulmonary artery; 6, mitral valve; 7, aorta; 8, vena cava; 9, branches of the aorta.

From this auricle the blood passes into the left ventricle through an aperture similar to that between the right auricle and ventricle. This opening is guarded by the mitral valve, which is fastened and operates in a manner similar to the tricuspid valve. From the left ventricle a single large blood vessel opens, called the *aorta* or great artery. This, like the pulmonary artery, is guarded from back flow at its base, by semilunar valves. Near its base, also, separate two smaller arteries (the *coronary arteries*) which convey blood to the substance of the heart itself.

Arteries, veins, and capillaries. — (See Ex. XLIV.) In the preceding paragraphs we have used the terms artery and veins to indicate blood vessels, but without distinguishing clearly between them. What is the meaning of the terms artery, vein, and capillary as applied to blood vessels?

An examination of the back of the hand will show us a number of dark, bluish colored lines, which give no sensation of beating when we place the finger upon them. These lines are actually blood tubes lying underneath the skin, and their color and lack of beat or pulse indicates that they are veins. If now, we place our finger on the inside of the wrist we shall find that there is a certain spot where a regular beating or pulse is felt. That spot marks the presence of an artery near the surface, and if we look closely we can make out a thin line less blue than the surrounding veins.

While almost all parts of the body surface show blue veins the pulse spots are very few, the most prominent being located on the inside of the wrists, the temples, and the under side of the knee. This indicates that the arteries are less near the surface than the veins and are thus

better protected. The necessity for this is found in the fact that the bleeding of a cut artery with its pulsing blood is much more difficult to stop than a bleeding vein.

If we prick any spot on the skin with a needle we get bleeding, even though no arteries or veins are visible. This is due to the fact that a network of tiny tubes, too small to see with the eye, and called hair tubes or capilla-

FIG. 70—Capillary circulation in the web of a frog's foot ($\times 100$); *a*, *b*, small veins, *d*, capillaries in which the corpuscles are seen to follow one another in single series; *c*, pigment cells in the skin.

ries, because of their tiny dimensions, connect veins and arteries. The relation of arteries, veins, and capillaries may be observed to good advantage by stretching the living web of a frog's foot or tadpole's tail under the objective of a compound microscope. (See Ex. XLV. and Fig. 70.) The blood vessels, which are transparent, show

clearly the streams of blood plasma and corpuscles in motion. With closer attention we can make out which tubes are arteries, veins, and capillaries. The blood in the arteries flows in spurts or pulses. The capillaries show a steady streaming motion and are so small that the corpuscles move in almost single file through them. The veins are as large as the arteries but the blood in them flows in a steady stream. If, now, we observe the direction of flow in the several vessels, it will be seen that the blood in the arteries divides and passes into the small capillaries, while from them in turn it passes on into the veins and through these toward the body again. In other words, the capillaries are the connecting links between the arteries and veins, and in passing through them the blood loses its spurting method of flow and is converted into a steady continuous flow.

The difference in color between the arteries and veins is explained by the color of the blood in the two tubes. The blood in the arteries is bright red, being full of oxygen, while that in the veins has a darker color owing to the larger proportion of carbon dioxide present. This latter blood is not blue, and that color is due to certain light effects.

Summary. 1. Arteries convey blood from the heart, and veins return the blood to it

2. Arteries contain much oxygen and little carbon dioxide, veins contain much carbon dioxide and little oxygen.

3. Blood in arteries moves in spurts (that is, it pulses), while that in capillaries and veins moves in a steady stream.

4. Arteries are better protected than veins by being in general placed farther below the surface. (This does not apply to the main vein trunks.)

Structure of blood vessels. — (See Ex. XLVI.) *Arteries.* A cross section of an artery (see Fig. 71) shows it to be made up of three layers of tissue. The outer coat is composed of fibrous connective tissue, which is tough, and a strong protective layer. This coat contains also some elastic fibers that enable it to expand under the pulsing of the blood. The middle layer consists of a mixture of unstripped muscle cells and elastic fibers, and is thicker than the other two layers. The inner layer is

FIG. 71 — Cross section of an artery; *c*, fibrous connective tissue, outersheath; *m*, muscle cells and elastic fibers; *e*, epithelial cells, with bands of tissue separating them from the middle layer.

FIG. 72 — Cross section of a vein; *c*, outer fibrous sheath; *m*, middle muscle and elastic tissue layer; *e*, inner epithelial cell layer.

made of flattened, epithelial cells, and between it and the middle layer are a varying number of bands of membranous and elastic tissue.

Veins. The structure of the veins is practically identical with that of the arteries (see Fig. 72) except that the outer coat is thickest and the two inner layers are much less developed. Owing to this difference they are much

less elastic, and readily collapse when empty. The

toughness and thickness of the outer coat gives them great strength.

Many veins also possess semilunar valves formed by pockets located at intervals on the inner surface of the tubes. These are so arranged (see Fig. 73) that the edge toward the heart is free and prevents a backward flow in the tube by pressing out and bridging the tube under backward pressure. These valves are found chiefly in the superficial and muscle supply veins. The arteries have no such valves except at the entrance into the heart. The location of these valves may be recognized by the knotted appearance of the veins at these points.

FIG. 73 —
Vein, showing
semilunar
valves.

Capillaries. As the arteries decrease in size the outer and middle coats gradually disappear until, in the capillaries, only the inner layer is left. This is composed of flattened cells cemented together at the edges. Such a structure is evidently well adapted to the process of osmosis, dialysis, and filtration.

XIII. CIRCULATION (*continued*).

THE FLOW OF BLOOD, AND BLOOD SYSTEMS.

IN the preceding chapter we have seen the way in which the various parts of the circulatory system are put together. In the present chapter we shall investigate the cause of the blood flow and its course in the blood vessels.

Heart action. — (See Fig. 66.) Enough of the structure of the heart has been outlined to show that the actual driving of the blood into the arteries is due to the contractions of the ventricles. Before these ventricles contract it is necessary that they be filled with blood, and this requires a preliminary contraction on the part of the auricles. The order of these contractions is as follows: The action begins at the mouth of the large veins (the *venæ cavæ*). The blood is flowing steadily from these veins into the auricles and dilating them. Next the tricuspid and the mitral valves are forced open by the blood pressure in the auricles and the ventricles fill. Now the auricles contract and send a wave into the ventricles dilating them still further and checking the flow from the veins by back pressure, while the mitral and tricuspid flaps float upward. As the ventricle becomes fully filled, it, in turn, begins to contract, and these flaps are suddenly pressed upward until they cover the opening tightly and prevent further flow from the auricle into the ventricle and also back flow from the ventricle into the auricle. The *chordæ tendinæ* prevent the flaps from being forced back through the opening. The blood in the ventricles is now under

pressure and forcing down the semilunar valves at the mouths of the arteries; the blood from the ventricles enters these arteries with a spurt. While this ventricle contraction is going on, the auricles begin to expand and the blood flows from the veins into them. Before they finish expanding the ventricles also begin to expand, while the semilunar valves of the arteries close to prevent back flow from them. During this period of expansion the mitral and tricuspid valves begin to open, and both auricles and ventricles to fill with blood. This goes on until the auricles begin to contract again, and this cycle is repeated indefinitely. The position of the valves may be summarized as follows:

(a) As the auricles contract, the mitral and tricuspid valves are open; the semilunar valves are closed.

(b) As the ventricles contract, the mitral and tricuspid valves are shut tightly. The semilunar valves are open and blood is entering the arteries.

(c) During the pause before the next contraction the mitral and tricuspid valves are open to permit the filling of ventricles, while the semilunars are closed to prevent back flow from the arteries.

The name of *diastole* has been given to the period when the auricles and ventricles are expanding, while the period of contraction of these organs is called the *systole*. Between these two periods there comes a time when neither is contracting or expanding. Using these terms the various stages of a "cardiac cycle" or heart beat may be expressed as follows:

First. An auricular systole.

Second. A ventricular systole.

Third. Heart pause (both organs in diastole).

Sounds of the heart.—If one listens carefully to the heart beat, by placing the ear against the chest, two distinct sounds may be distinguished. The first is lower in pitch than the second and of longer duration. It is supposed to be produced by two factors, the vibration of the tricuspid and mitral valves in closing, and the vibration of the contracting muscle mass. The second sound is sharper and is due to the closing of the semilunar valves. The first sound is essentially a systolic sound, while the second is diastolic. These two sounds are of great aid to the physician in determining cases of heart trouble. For detecting variations in this sound an instrument called the stethoscope is used, and this method of examination is called auscultation.

Pulse.—When the ventricles contract the blood is forced into the arteries with a spurt. If the arteries, capillaries, and veins were rigid tubes this sudden pressure would be transmitted instantaneously throughout the entire system, and a quantity of blood would enter the auricles exactly equal to that which left the ventricles. Owing to the elasticity of the walls of the blood vessels this is not the case. Part of the pressure is used up in expanding the walls. Since the arteries are subject to the most direct effect of this systole the reason for their thicker walls is apparent. This elastic character of the arteries also results in the pressure being transmitted in a wave instead of directly. The results of this may be pictured as follows: As the ventricle contracts the expressed blood stretches that portion of the artery nearest the heart. As the ventricle expands and this pressure is removed this enlarged part of the artery begins to contract slowly, and the pressure is transferred to another point further along in the artery. As the next systole occurs this process is repeated, and thus the effect of the pressure or wave is gradually transmitted throughout the whole

arterial length. In this way each portion of the artery wall expands to a maximum and then contracts, at intervals corresponding to the systolic and diastolic periods of the heart. The disappearance of this wave in the capillaries and veins is explained when we consider how the arteries branch into smaller and smaller tubes, which are farther and farther removed from the seat of pressure until in the capillaries the wave has become so split up as to be barely perceptible, and in the veins is converted into a steady flow. In this respect it is like a water wave, which becomes less apparent the farther we go from the center of the disturbance. It is this wavelike character of the flow of the blood in the arteries that enables us to feel a pulse at points where the arteries are near the surface. What we actually count in taking the pulse is the time between successive periods of pressure and relaxation, and since these correspond to expansion and contraction of the heart they enable us to detect the rate of beat of the heart. The physician uses the pulse point in the wrists and temples to furnish him with valuable information about the heart and circulation.

Pulse rate.—The rate of the pulse or heart beat is subject to great variation under different conditions of health and disease. It is this fact that makes it a valuable indicator of the body condition to the physician.

Age, sex, size, atmospheric pressure, all have a distinct effect upon the pulse rate. The average rate for the adult man is seventy beats per minute, and for the adult woman seventy-eight to eighty per minute, showing that sex affects the rate. Tall individuals have a slower rate than short persons, while large bodies have a slower rate than small bodies. The human rate is most rapid in infancy, sinks rapidly at first, and then more slowly

to a minimum in adult life, rising slowly as we enter extreme old age. The following table gives the average rates:

At birth	140 beats per minute.	
Infancy	120	"
Childhood	100	"
Youth	90	"
Adults	75-70	"
Old age	70	"
Extreme age	75-80	"

SYSTEMS OF CIRCULATION.

From what has been said it follows that the blood which passes through any given chamber of the heart must ultimately return to that same chamber. Owing, however, to the division of the heart into two parts, which beat in unison, and the branching of the blood vessels, the actual course of the blood requires some further explanation. For convenience, the blood vessels that are concerned in the transfer of the blood from the right to the left side of the heart, and that have their branches in the lungs, are spoken of collectively as the pulmonary system. Those concerned with the transfer of blood from the left side of the heart to the right side, and whose branches take it all over the body, are spoken of as the systemic system.

Pulmonary circulation.—When the blood enters the right auricle it is partly exhausted of its oxygen and is dark and purplish in color, owing to the abundance of carbon dioxide which it contains. In this condition it enters the right ventricle, and when that organ contracts it is forced into the *pulmonary artery* which in turn breaks up into two branches leading to the right and left lung, and these into smaller branches within the lungs. These pul-

monary arteries convey the blood from the right ventricle directly to the lungs. In the many capillaries, into which

FIG. 74 — Diagram of the circulatory system.

the branches of the pulmonary artery divide in the lungs, the blood takes on a new supply of oxygen and loses its carbon dioxide. These capillaries are collected into larger

and larger veins and finally bring back the oxygen laden blood to the left auricle by four veins (two from each lung). These veins are called the right and left pulmonary veins respectively. From the left auricle the blood passes into the left ventricle, and in this manner the transfer of blood from the right to the left side of the heart is brought about. In our study of the lungs we shall discuss more in detail the manner in which it loses its carbon dioxide and takes on oxygen.

In this circulation it will be noticed that it is the pulmonary artery which contains the carbon dioxide laden blood, while the pulmonary veins contain the bright red, oxygen laden blood. Owing to this fact the only universal definition that will apply to all arteries and veins is one which is based on the direction of blood flow. Arteries convey blood away from the heart, veins toward the heart.

The systemic circulation.—From the left ventricle this oxygen laden blood is pumped into a large artery called the *aorta*. The *aorta* subdivides into many branches which carry the blood to all parts of the body. In the tissues these arteries subdivide into capillaries, and these, in turn, collect into veins through which the blood finally enters two large veins (the *superior* and *inferior venæ cavæ*) and is returned to the right auricle by them. In this system the blood is transferred over the body and from the left to the right side of the heart.

Since it is this system that carries food and oxygen to the tissues and transfers their wastes and carbon dioxide to the excretory organs, the relations of the branches is important to a clear understanding of the blood flow of this system. As it leaves the heart, the great *aorta* bends first to the left, then downward through the rear part of

FIG. 75 — The arterial system.

the chest cavity, finally piercing the diaphragm and entering the abdominal cavity. The change in direction as it leaves the heart results in an arch called the arch of the aorta. From the top of this arch on the right side is given off a large branch which, in turn, soon divides into two parts. One of these two parts (the right *carotid* artery) supplies blood to the right side of the head. The other part passes under the collar bone, down the arm, and at the elbow divides into the *radial* and *ulnar* arteries thus supplying the right arm with blood. It is the radial artery whose pulsings are felt in the wrist.

From the top of the arch on the left side are given off two branches, one of which (the left *carotid*) supplies blood to the left side of the head, while the other branches as on the right side and supplies blood to the left arm.

From the aorta, near the heart, arise also the coronary arteries which, branching over the surface of the heart, supply the tissues of that organ with blood. As the aorta curves downward through the chest cavity it gives off branches which supply the lung tissues, the esophagus, and the muscles of the ribs and spinal column.

From the aorta in the abdomen are given off five main branches. The first (*cœliac axis*) divides into three parts which supply the stomach, spleen, and liver respectively. Two other branches (the *mesenteries*) carry blood to the walls of the intestines. The remaining two branches carry blood to the kidneys. The main trunk of the aorta continues to the rear of the abdominal cavity and finally divides into two branches which each enter a leg and supply these extremities with blood.

All these branches of the aorta subdivide first into smaller arteries and then into capillaries, which are in turn col-

lected into veins by which the blood is returned to the right auricle of the heart.

The blood supplied to the head by the carotids is returned to the heart by two large veins (the *jugulars*) passing down each side of the neck; while from the arms come two other veins. These four veins finally unite to form the superior vena cava which opens into the right auricle of the heart. In a similar manner various veins return the blood from the lower part of the body and unite in one common vein (the inferior vena cava) which also opens into the right auricle of the heart. It is this lower part of the venous system which collects the food from the digestive system.

The portal circulation.—The veins which are found in the walls of the stomach and intestine do not convey the

FIG. 76—Diagram of the portal circulation.

blood directly to the inferior vena cava. After collecting the food from these organs these veins are collected into a single large vein (the *portal vein*) which passes directly to the liver. Here this vein splits up into capillaries again, which

in turn collect again into a second vein (the *hepatic vein*). This latter pours its blood into the inferior vena cava. It is to be noted in this case that the capillaries of the liver arise from and unite again into veins, and that the food is thus forced to pass through two separate sets of capillaries before entering the heart. This branching of the portal vein into capillaries is the only

example in the body of a vein's connection with another vein through capillaries. In all other cases the capillaries connect veins and arteries.

It is this arrangement of portal-vein capillaries which permits the liver to act as a storehouse of carbohydrate material, and to regulate the amount of that material which shall enter the general circulation at any given time.

The renal circulation. — The blood which is returned to the left side of the heart by the pulmonary veins has been freed of its carbon dioxide only. Hence the blood which passes into the aorta is still rich in urea and other body wastes brought to the heart by the *venæ cavæ*. Much of this waste is diverted into the artery branches which carry blood to the kidneys (the *renal* arteries). In these organs, by means which we shall discuss more fully under the excretory system, this waste is removed, and the *renal veins*, therefore, return to the inferior vena cava blood which is nearly free of body wastes.

Changes in blood composition. — In passing through the various systems of circulation, portal, renal, systemic, and pulmonary, the blood undergoes distinct changes in composition. Having in mind the parts of the body which these systems supply, we may now summarize these changes as follows:

The loading of the blood with nourishment. — 1. Food enters the blood at two points in its circulation.

- (a) All kinds of food, except the fats, enter the capillaries of the stomach and intestines and are collected and carried to the liver by the portal vein. In the liver, the capillaries into which this vein divides permit the liver cells to separate from the rest of the

blood the carbohydrate, together with certain poisons (see p. 142). The carbohydrate thus separated is either stored in the liver as glycogen, or given out as needed in the form of grape sugar to the hepatic vein. This hepatic vein passes this grape sugar and the other unchanged nutrients into the general circulation by opening into the inferior vena cava.

The poisons are passed into the liver secretion or bile and returned to the intestine with that fluid for removal with the fæces.

- (b) Fats are collected by the lacteals, and enter the veins by way of the thoracic duct at a point just under the left collar bone where the left jugular and left arm vein join. Thence they pass into the heart by the superior vena cava.

2. Oxygen enters the blood in the lungs, and is conveyed to the heart by means of the pulmonary veins. We shall consider this phase of the circulation more in detail in our study of respiration.

The loading of blood with waste.—All veins carry wastes (CO_2 , urea, etc.), which they collect from the tissues. This waste either enters the blood directly by way of the capillaries, or else enters the lymphatics and thus the veins by the thoracic duct or right lymphatic duct. The right lymphatic duct enters at the joining of the right jugular vein and the right arm vein.

Unloading of nourishment by the blood.—All arteries¹ carry food and oxygen to the tissues which they supply, and deliver these substances to the cells of the tissues

¹ The pulmonary veins and arteries are exceptional in that the pulmonary veins carry oxygen laden blood, and the pulmonary arteries, carbon dioxide laden blood.

through the thin walls of the capillaries in the manner already described (see p. 175). The plasma composition indicates that it is the part of the blood which carries the food to the tissue cells.

Unloading of wastes by the blood.—The carbon dioxide given off by the tissues is removed from the blood almost entirely in the lungs. It is noteworthy that the lungs do not free the blood of wastes in general, and on that account should not be said to *purify* the blood. In other words, their great function is to exchange oxygen for carbon dioxide or aerate the blood.

Most of the urea and true wastes of the body are removed by the kidneys and the skin, this waste being collected from all parts of the body by the veins and conveyed to these organs by special arteries, such as the renal artery to the kidneys.

Arterial and venous blood.—From the description given above it is readily seen that the blood in the arteries and veins varies in composition in different parts of the body. In general, however, we may say that the veins contain less oxygen and more carbon dioxide than the arteries. The redder color, due to the larger proportion of oxygen present, has led to the custom of representing arteries by red, and veins by blue lines in a diagram of circulatory systems. The notable exceptions to this rule are the pulmonary veins and arteries. The pulmonary vein contains very little carbon dioxide, while the pulmonary artery contains very little oxygen.

In general, then, the terms arterial and venous are used to indicate relative amounts of oxygen and carbon dioxide in the blood. In that sense the pulmonary vein contains arterial blood, and the pulmonary artery, venous blood.

REGULATION OF BLOOD FLOW.

Heart control.—The amount of blood which passes through a given part of the circulatory system at any one time is, of course, regulated primarily by the heart beat. This beat, which is produced by rhythmic contractions of the heart muscles, is ordinarily very regular. Its rate, however, is subject to certain external influences. For example, a decrease in blood pressure in the arteries is associated with an increase in heart rate and *vice versa*. Muscular exercise also increases the heart rate. Even so little an effort as a change in position is sufficient to change the rate, that of a standing position being higher than that of a sitting or reclining one.

The purpose of these changes in rate is readily understood when we consider that tissues which are active are using up more food and creating more waste than still tissues, and consequently require a greater flow to supply this demand.

Mental states and emotions, blood content, and temperature, all affect the rate of heart beat. Some lower this rate, others increase it. Some, like temperature, cause an increase up to a certain limit, and then above that limit produce a decrease. Physicians take advantage of these factors and introduce into the blood certain drugs which may accelerate or lower the rate of beat, and thus can control it to a certain extent in cases of disease.

The manner in which all these special conditions are able to affect the heart is through what we call nerve action. The muscle walls of the heart are connected by nerves to other parts of the body, and when one set of nerves is stimulated by any of the above means, the nerve-

ends in the heart muscle receive this impulse and increase the rate of contraction and expansion of the cells. When another set of nerves is stimulated, they, in turn, cause the contraction to be reduced in rate. These are like telegraph wires connecting stations. One station is the heart where messages are received. The others are found in the brain and in the body tissues, and the messages sent from these latter points control the action of the heart. Of the two kinds of nerves those that decrease the rate are called heart inhibitors; and those that increase the rate, heart accelerators. By their aid the rate of flow is adjusted to the needs and conditions of the tissue cells.

Artery control.—The walls of the arteries, as well as of the heart, are supplied with the ends of what are called the vaso-motor nerves. Under the normal action of these nerves the walls of the arteries have a certain elasticity which responds to the blood pressure; stretching when the pressure is increased and contracting as the pressure is removed. When these vaso-motor nerves are stimulated further the effect produced may be a still greater contraction of the artery walls and a consequent decrease in the blood content of the contracted artery, or a paralysis of the walls which causes them to lose their elastic tension and stretch. In this way it is possible for conditions, external or internal, to affect these nerves and produce stimuli, which, in turn, control the blood supply of any given part of the body. The nerves which, when stimulated, cause the contraction of the walls are called the vaso-constrictors. Those which, when stimulated, produce paralysis of the walls and thus allow them to stretch are called the vaso-dilators. This stretching and contract-

ing of the artery walls with the consequent increase and decrease in blood must necessarily cause the capillaries, whose walls are also elastic, to become stretched. Hence variation in the tension of the artery walls is always accompanied by similar variations in the blood supply to those capillaries which the arteries feed.

The conditions which may thus affect these vaso-motor nerves are many. Heat to a certain limit, exercise, the presence of food in the digestive tract, etc., all affect the vaso-dilator nerves and cause the artery walls to relax. Cold, on the contrary, affects the vaso-constrictors and causes the contraction of the walls. Mental emotions in some cases cause relaxation of the walls, and in other cases, contraction. Thus, blushing is due to overcharging of the facial capillaries from relaxation of those artery walls which supply these capillaries; while paleness is due in the same manner to contraction of the supplying artery walls. In both cases the mental emotions are responsible for the nerve stimulus.

Injury to tissue cells invariably results in a relaxation of the walls of the arteries supplying the affected part, and thus permits an increase in the blood supply to the injured area. Such an increase is spoken of as congestion.

Congestion and inflammation.—A simple case of congestion is seen when we scratch the skin with a pin, the scratched surface becoming reddened from the action of the injured cells upon the artery supply. If the injury is severe and the cells are killed, swelling often follows the redness, and painful sensations may result.

Such congestion, with its attendant swelling, is merely an indication of the first steps in the repair of the injury. The flow of blood to the injured part is soon followed, in

this case, by the outpouring of plasma and white blood corpuscles (leucocytes) through the capillary walls into the lymph spaces. This extraordinary flow is due to changes in the cement which holds together the capillary cells composing the walls, and may in some instances be accompanied by the outpouring of red corpuscles.

In simple injuries the result of this outpouring of plasma is to stick together the old cells and cause them to grow and unite, while the leucocytes and injured cells are absorbed and the injury thus repaired.

If, however, the injury is extensive or bacteria are introduced into the injured area, a more extensive action is involved. The growth of the bacteria or the nature of the injury may cause the death of other cells in the neighbourhood, and the formation of a mass of dead tissue results. In this event the white corpuscles proceed to kill or absorb the bacteria and dead tissue cells with the formation of a mass of matter usually designated as pus. At the same time new cells are formed and a new tissue grows in place of the old one. This pus may be removed by the lymphatics or absorption into the blood vessels, or it may soften the tissue about it and discharge, as in an abscess. In the latter event, the white corpuscles either devour the bacteria and cells, or give off ferments which dissolve them. Meanwhile the living cells about the injured part are excited to growth and reproduction, and thus new cells produced which unite and replace the old tissues.

The processes here are evidently two, the dissolving and removal of old matter through the agency of the white corpuscles, and the formation of new tissue cells to replace the injured ones. The manner in which the old matter is removed varies with the location of the injury. These

two processes of repair are called inflammatory processes, and inflammation is usually defined as "the local attempt at repair of an injury."

Examples of simple inflammatory processes.—When the mucous membrane is injured by exposure to cold, there results a congestion in the membrane. This congestion may be complicated by the presence of bacteria. In such congestion the formation of pus is accompanied by an increased activity and secretion of the mucous cells. The combined secretion and pus collects in the passages inclosed by these membranes and forms the discharge common to colds. This congestion of the mucous membrane may be brought about by the action of the cold upon the vaso-motor nerves which control the artery supply to the skin. In this event their action is to cut off the blood supply to the skin, and the excess of blood thus cut off is forced to accumulate in the mucous membranes and produce congestion. If this congestion and inflammatory process is confined to the nasal passages the result is called a cold in the head, or catarrh. If in the throat, it may result in swelled tonsils and a sore throat. If in the bronchia passages, it is called bronchitis, while in the lungs it may produce pneumonia. When certain bacteria attack the throat membranes the result may be severe. Such an instance is seen in diphtheria.

Another example of inflammatory processes is found in the pimples and boils. In these cases, bacteria are introduced in some way into the underlying layers of the skin where they cause the death of the surrounding cells. Congestion results, and the leucocytes dissolve the dead cells and devour the bacteria forming a greater or less quantity of pus. The pus thus formed softens the surface layers of skin and forms a head or weak spot on the surface through which the pus may ultimately be discharged. "Taking cold in a boil" is actually a case of accidental introduction of bacteria into a partly discharged boil, and the resultant formation of new pus, and has no connection whatever with cold.

The introduction, by the blood, of bacteria into the internal

tissues, internal disturbances of function, and many other causes, may bring about so extensive a congestion and accumulation of poisonous matter that the body is unable to remove it. In that event the presence of this matter in the body may produce various forms of disease, such as fevers, growths, etc.

Treatment of cuts and bruises.—When the capillaries are cut the bleeding is very slow and is usually easily checked by the natural clotting of the expressed blood. The cutting of an artery or vein, however, is a much more serious matter, and owing to the spurting of the blood the cut artery is more difficult to treat than the vein. If the artery or vein is small the mere pressing together of the walls with a bandage may check the flow sufficiently, to permit the formation of a clot. When the artery or vein is large the blood pressure is too great to allow the clot to form and special measures are necessary. The object of all such measures is to secure sufficient pressure upon the blood vessel to force the cut edges together and diminish the blood flow sufficiently to permit the formation of a clot. Owing to the direction of flow in an artery, this pressure must be applied on the side of the cut nearer the heart. In case of a vein the pressure should be applied on the side farther from the heart. In some cases simple bandaging fails to stop the flow and the clot is washed away as fast as it forms. In such cases an application called a tourniquet is resorted to. This consists of a bandage applied on the proper side of the cut and twisted tight with a stick so that the walls of the blood vessels are pressed together at a point above the cut. The blood being shut off from the cut ends, a dangerous loss of blood is thus prevented. In emergencies persons may thus be saved from bleeding to death since all the tools required are a

strip of cloth and a stick. The tourniquet is of course only a temporary measure and is to be employed only as an emergency remedy until the surgeon can treat the wound. Surgeons usually close large wounds by tying the ends of the cut blood vessels with clean catgut. Such bindings are called ligatures.

When the ends of a vein are cut they may grow together again and thus circulation be restored through the original channel. Artery ends, on the contrary, do not grow together but the ends close and the branches which are in communication above and below the cut enlarge and replace the original channel.

All cuts should be kept clean and free from bacteria. Weak solutions of certain germicides can be obtained from druggists and used to wash and protect the cut from these dangerous visitors. Blood poisoning is one of the serious results of neglect of these precautions.

The healing of a cut or bruise will take place without pain or inflammation if kept clean.

Effect of alcohol upon the circulation.—The most marked effect of small amounts of alcohol is the dilation of the surface capillaries and their filling with blood. The result is a sensation of warmth or glow due to the stimulation of the surface heat nerves. Alcohol produces this effect by its action on the nerve centers, which, in turn, stimulate the vaso-dilator nerves of the arteries that supply these capillaries, and not by any direct action upon the blood vessels.

In this case the sensation of warmth is deccitful, since the bringing of the blood from deeper parts of the body to the surface permits the escape of heat and a slight lowering of the body temperature.

In larger doses alcohol produces effects in one of two ways:

(a) It may cause the death or degeneration of the cells with which it comes in contact. For example, it may cause the cells of the heart to become charged with fat and thus weakened, or it may make the cells of the artery walls brittle and unable to withstand pressure, or finally, it may actually poison and destroy the cells of certain tissues.

(b) When brought in contact with the cells of the tissues it may so injure them that congestion and inflammation results. This inflammation may cause a disturbance of the natural action of the congested part due to the swelled condition, or it may stimulate unnatural growth on the part of the surrounding tissue cells. Another bad result of this congestion may be the stretching of the arteries to such an extent that they are unable to recover their natural size, and thus produce continued congestion in the part affected.

The extent of the injury which alcohol may produce in the body by either one of these means, that is, directly or indirectly, is a subject for a medical text, and we need only outline here the method of the action. In brief, then, it is evident that alcohol in either large or small doses acts upon the nervous centers and through them upon the vaso-dilator nerves, producing abnormal distribution of the blood, and that in large doses it is actually poisonous to the cells with which it comes in contact. While the extent of the injury produced by small doses may be only temporary, continued use or excess causes such changes in the distribution of blood as to be extremely dangerous to the system.

XIV. CIRCULATION IN THE LOWER ANIMALS.¹

IN the one-celled animals, such as the amoeba, the food is digested in the cell protoplasm itself, and oxygen is absorbed directly from the surrounding air or water. In like manner the wastes and carbon dioxide are removed from the cell and cast into the surrounding medium. In such animals a circulatory system is evidently unnecessary. In fact, the only suggestion of it is found in the movement of the protoplasm in the cell by means of which the absorbed food is distributed throughout the cell. This, however, is entirely an internal cell circulation, and not a system for bringing food to the cells as in man.

In the sponges and coelenterates each cell is able to obtain its air supply and food supply directly, and no special circulatory system is necessary. When we come to animals possessing many layers of cells it is evident that some cells will be so placed as to be unable to secure food and air directly or to rid themselves of waste. Some special apparatus, therefore, becomes necessary to supply these cells with food and to remove their wastes. In this way there arises the circulatory system as we have considered it in our definition. Some of the systems in the lower animals are interesting from the light they throw upon the development of our own complex network of tubes.

Insect circulation. — In the grasshopper, we see one of the simplest forms of a circulatory system. In this animal

¹ See Footnote on p. 146, Chapter X.

the food passes from the digestive tract directly through the walls of the stomach and intestine into the body cavity. Here it would remain were it not for a tube with several swellings along its surface extended along the dorsal surface of the body cavity. This tube, with its swellings, is a sort of muscular pump or heart, closed at the rear end and open in front. The swellings expand and contract rhythmically. As they expand the liquid of the body cavity is sucked in through openings along the sides. These openings are protected by valves which open inward. As the swellings contract these valves close, and the inclosed fluid is thus forced forward and out the front end of the tube toward the head. In this manner the body fluid is kept in motion, and by means of the cavities between the tissues it flows around and bathes all the tissues, and finally returns to its starting point, thus supplying the body cells with food and receiving their wastes. This waste is absorbed by kidney tubes which extend into the body cavity from the intestine, and the absorbed fluid is excreted through the intestine with the fæces. The liquid circulated contains no red corpuscles, as the air supply is furnished by an entirely independent set of tubes. There are no arteries or veins, the blood flowing through the spaces of the body from the mouth of

FIG. 77—Portion of an insect's blood tube (dissected); *a*, *b*, muscular walls; *d*, valves between the compartments; *c*, valve covering the general opening into the body cavity; through this valve the liquid enters.

the tube to its starting point. The heart is simply a contractile structure whose office is to keep the liquid in motion.

The earthworm.—In this animal we have a slight advance over the insect. It possesses a true closed system of tubes conveying a red blood. It consists of two tubes running lengthwise of the body, one dorsal and one ventral. The dorsal blood vessel can be easily made out through the transparent skin of the living earthworm, and observation shows that the red liquid which it contains moves in pulses toward the head. At the anterior

FIG. 78—Diagram of a longitudinal section of an earthworm; *h*, one of the fine pulsating arches which pump blood from the dorsal blood vessel into the ventral blood vessel.

end of the animal are five pairs of arches or blood vessels, which branch from each side of the dorsal blood vessel and, encircling the esophagus, open below it into the ventral blood vessel. These pairs are arranged, one behind the other, like rings from the sixth to the eleventh segment.

These five pairs of tubes are called the aortic arches, and they pulsate in unison, forcing the blood to flow from the dorsal blood vessel into the ventral. The course of the blood, then, is from the rear of the body along the dorsal blood vessel to the head; downward, through the arches to the ventral vessel, and through this toward the rear of the body again. Both dorsal and ventral blood

vessels give off branches, and while the exact course of the blood flow from the ventral vessel is at present unknown, it is supposed that in some way it is ultimately brought back into the dorsal vessel and thus completes its circuit. The branches of the two main vessels subdivide into capillaries, and by means of these the blood is brought into contact with the various tissues of the body, and thus food is distributed and wastes collected.

The red color of the blood is due to the hæmoglobin dissolved in the fluid, the only corpuscles present being a few white ones.

Besides this inclosed fluid, we find the body cavity filled with a similar liquid to that found in the grasshopper, and it is known that part of the absorbed food enters the inclosed system and part is distributed by the cavity fluid, so that we actually have two methods of circulation in this one animal.

The fish. — It is only in the vertebrate animals that we find a true blood with red corpuscles. The simplest vertebrate system is found in the fish. In this animal the heart is a simple structure consisting of a single ventricle and auricle, and is located in a pericardial cavity separated from the rest of the body organs by a thin membrane. The blood is driven from the ventricle through an artery (the aorta) to the gills. Here the artery branches into capillaries and through the thin walls of these is effected the interchange of oxygen and carbon dioxide between the water

FIG. 79 — Plan of circulation in a fish; *a*, auricle; *b*, ventricle; *c*, branchial artery; *e*, branchial veins; *d*, gills; *f*, aorta; *g*, vena cava.

and the blood. These capillaries ultimately return the blood to a dorsal artery which branches and carries the blood to all parts of the body, where the division of the branches into capillaries permits the giving up of food and the removal of the waste of the tissues. These capillaries are now collected into veins, and the blood is returned by these tubes to the auricle of the heart to repeat this cycle indefinitely. As in man the food of the digestive tract is absorbed by capillaries which unite to form the portal vein, and this absorbed food is carried to the liver by this vein. Here the portal vein produces capillaries, and these, after branching throughout the liver, unite again into the hepatic vein and are carried directly to the auricle by this tube.

In this circulation the single auricle and ventricle correspond to the right side of the human heart, and the heart receives and expels only the venous blood. The friction in the gill capillaries retards the flow so greatly that the circulation in the arteries of the fish is very sluggish.

The red corpuscles of the fish are nucleated, and it also possesses a lymphatic system.

The frog.—In the frog the blood is aerated in the lungs, and the heart has two auricles and one ventricle. The right auricle, as in the fish, receives blood from the veins which bring it from all parts of the body to this division of the heart. The left auricle receives aerated blood from the lungs. It is to be noted that the frog has advanced over the fish in having the aerated blood returned to the heart before pumping it over the body, and as we would expect, this arrangement allows a little faster flow through the arteries. Since, however, there is only one ventricle, the two kinds of blood found in the auricles must become mixed in this chamber, and the arteries in consequence are

filled with mixed blood. By an ingenious arrangement of arteries and valves the blood is not evenly mixed, but the part which is pumped to the head is richest in oxygen, that to the body organs next richest, and that to the lungs and skin is poorest in oxygen and richest in carbon dioxide. As in the fish, the red corpuscles are nucleated. In its early life the frog breathes by gills.

The reptiles.—These animals have, in general, the same sort of circulatory system as the amphibians. There is, however, one modification in the structure of the heart that shows an advance. In most reptiles the ventricle is partly divided by a partition, and this partition tends to keep separate the aerated and non-aerated blood so that the aorta which arises from the left side of the ventricle receives blood freer of carbon dioxide than the pulmonary arteries which arise from the right side. This partition is interesting as a step toward complete separation of the two kinds of blood. The red corpuscles are nucleated.

The birds.—In these vertebrates the ventricle is completely divided by the partition, and the bird heart, like that of man, is four chambered. This fact alone would enable us to classify birds as higher forms than reptiles. The rate of flow in birds is much more rapid than in man, and the temperature of the blood is much higher. This is due

FIG. 80.—Plan of circulation in amphibians and reptiles; *a*, right auricle receiving venous blood from the system; *b*, left auricle receiving arterial blood from the lungs; *c*, common ventricle; *d*, *f*, *c*, systematic arteries, capillaries, and veins; *g*, pulmonary artery; *k*, *h*, pulmonary capillaries and vein.

to the tremendous muscular exertions that the bird is compelled to make in the process of flying. Another change is the turning of the aorta to the right instead of to the left. The red corpuscles of this animal are not nucleated, and in this respect the bird is nearer man in the scale of animal life.

FIG. 81.—Plan of circulation in birds and mammals; *a*, right auricle receiving blood from the system; *b*, left auricle receiving arterial blood from the lungs; *c*, right ventricle; *c'* left ventricle; *d*, *e*, *f*, systemic arteries, veins, and capillaries; *g*, pulmonary artery; *k*, *h*, pulmonary capillaries and vein.

The mammals.—In all the mammals the circulatory system is practically identical with that of man, and the red corpuscles are without nuclei, but differ in shape.

Significance of these comparisons.—The lessons that these comparisons teach are two. First, they show us that while the apparatus may be different in construction in the different animals, the purpose of the structures found are identical in all animals, namely, to distribute food and oxygen and remove wastes. Second, the fact that they differ only in minor details and in adaptation to special conditions of life show us that man bears a very close relation to other animals and differs from them only in the greater perfection of his organs. It is by comparisons such as these that men have been enabled to trace the history of the development of living forms on the

earth, just as by studying all the different musical instruments that have ever been produced it is possible to learn the history of instrument making in the arts. In short,

each form of animal that shows an advance over another in the structure of its organs aids in locating the time of its development and its relation to other forms. The grouping of animals according to their relative development of structure is the basis of the classification of animals in the zoölogy of to-day, and the view that each form of animal has developed from a lower form is called the Doctrine of Evolution.

PLATE II — Hand. From an X-ray photograph. Overton, *Applied Physiology*.

XV. THE SKELETON.

IN the preceding pages we have considered at some length the means by which nutriment and oxygen are distributed to the various tissues of the body. It will be well for us to consider now two of these most important tissues, the bones and the muscles, since to their action we owe all our voluntary and involuntary movements.

The body framework.—If we examine certain of the lower animals, such as the jellyfishes and worms, we find that, while they have a body form, that form is subject to distinct changes when subjected to pressure. In other words, they are compressible to a great degree. This is owing to the absence of any hard matter in their bodies, that is, they have no skeleton or framework. Other animals, such as the insects, lobsters, and starfishes have a framework, but it is on the outside of the body. While such a skeleton is preferable to none at all, it is an inconvenient arrangement, since in most of these forms it does not permit any growth on the part of the animal, without splitting of the hard layers, and thus growth in these animals is attended with periods when the soft parts of the body are exposed. Skeletons such as these are called exoskeletons. In the higher animals the problem of securing rigidity and protection of the soft parts, and at the same time allowing growth, has been solved by the development of an inner framework or endoskeleton of bone.

By the aid of an apparatus known as the X-ray machine (see Plate II) it is possible to take a picture of parts of

the living body which shows only faint outlines of the soft parts and brings out clearly the bony framework of these parts. With the aid of this apparatus, or by the

FIG. 82.—Sea urchin shell, an exoskeleton.

removal of the soft parts, a definite framework of bone, cartilage, and gristle is revealed, to which the shape of the human body is due, and which we call the human skeleton.

This skeleton is composed of over two hundred separate bones, so joined as to fulfil the following distinct purposes: (a) They give shape and rigidity to the body. (b) They protect the delicate organs and soft parts of the body. (c) They provide levers on which the attached muscles may act, and thus produce motion. The skeleton of man and the higher animals is peculiar in the development of a special chain of bones called the *vertebræ*, which are so interlocked as to form a long flexible column, the spinal column, and this column is the central portion of the framework. All animals that possess such a structure are called *vertebrates*. Man is the highest type of a *vertebrate* animal.

Bone regions.— On the top of this long, curved spinal column is mounted a box-shaped collection of bony plates which is called the skull. This box incloses the delicate brain and forms the framework of the face. The chest, or thorax, is formed by a series of long, curved, flat bones called the ribs, which curve from the vertebræ at the back around to the front of the body. Here, they are united to a flat structure, half bone, half gristle, called the breastbone. This arrangement secures the protection of the heart and lungs. The abdominal cavity of the body has no side walls of bone, but at the bottom the soft parts are supported by a basket-shaped collection of bones connected to the lower end of the spinal column and known as the pelvis. Finally, the arms and legs contain a central axis of bones, which, in the case of the arms, are connected to the spinal column by a girdle known as the shoulder or pectoral girdle, and, in the case of the legs, by the pelvic bones or pelvic girdle. The

FIG. 84—Spinal column; *A*, side view; *B*, rear view; *C*, 1–7, cervical vertebrae; *D*, 1–12, dorsal vertebrae; *L*, 1–5, lumbar vertebrae; *S*, sacrum; *Co*, coccyx.

bones that compose the column and those that frame the thoracic and abdominal cavities are spoken of collectively as the axial skeleton. The bones which form the appendages (arms and legs) are called the appendicular skeleton.

THE AXIAL SKELETON.

The spinal column.

— The bones that make up this long, curved column are called the vertebrae. These vertebrae are twenty-four in number and are divided into three regions. The seven nearest the skull are called *cervical* vertebrae and constitute the cervical or neck region. The next twelve are called the *dorsal* vertebrae and carry the

twelve pairs of ribs. This is called the dorsal region. The remaining five are very large and are called the *lumbar* vertebræ. Together they make up the lumbar or loin region. Below these lumbar vertebræ is a single large bone called the *sacrum*. This is actually composed of five vertebræ, separate in the young child but fused in the adult. On the end of this sacrum is a curved tip of bone (the *coccyx*) formed of four separate bones fused together. (For regions, see Fig. 84.)

Structure of the vertebræ. — (See Ex. XLIX., c.) The vertebræ are all irregular bones with a central portion called the body, and many projections or processes. The first two cervical vertebræ are formed for a special purpose and are called the atlas and axis. The remaining vertebræ have the same general form, differing only in size. We shall first consider a typical vertebra and then note the variations in the other forms.

A dorsal vertebra. — Viewed from above (see Fig. 85),

FIG. 85—Dorsal vertebra; A, top view; B, side view; 1, centrum; 2, neural cavity; 3, spinous process; 4, transverse processes.

it shows the following regions: First, a central mass of bone called the body, or centrum, which, in the human body, extends toward the ventral side. From this on each side an arch of bone curves backward inclosing a cen-

tral or neural cavity, through which the spinal cord passes. This arch is called the neural arch, and from it are given off seven processes or projections of bone.

One extends from the very center of the arch dorsally, and is called the spinous process. It is the end of this process that one can feel as he passes his finger down the middle of the back. Two lateral processes extend from each side of the neural arch, and to these the ribs are attached. They are shorter than the spinous process. The remaining four processes are simply raised projections lined with cartilage, and are called articular processes. Two face dorsally, two ventrally, and they connect

FIG. 86 —I, cervical vertebra; II, lumbar vertebra; 1, centrum; 2, neural cavity; 3, spinous process; 4, transverse processes; *A*, top view; *B*, side view.

with similar processes on other vertebræ, thus permitting the bones to articulate and move upon one another. The centrum is smooth on each side, and between succes-

sive centra are inserted pads of cartilage. The centra and pads bear the weight of the body. On the centra are also two cartilage-lined sockets which support a part of the rib.

The cervical vertebræ. — These are very similar to the dorsal vertebræ except for the absence of the rib articulations. The transverse processes are poorly developed and are perforated. The centrum is small, and the spinal process short and often forked. They are specially developed to carry the head and give freedom of action.

The lumbar vertebræ. — These are the largest of all the vertebræ and have no rib attachments. Their spinous

FIG. 87—*A*, atlas; *B*, atlas and axis; *a*, *b*, articulations with occipital bone; *c*, transverse ligament; *o*, odontoid process.

and transverse processes are short and stout, and well adapted for the attachment of strong muscles. The heavy centra allow them to support weight.

Atlas and axis. — The two top cervical vertebræ have a special function to perform, namely, to support the skull and permit its free movement from right to left and forward and back. To satisfy this requirement they show a special modification. The top one, known as the atlas, is fastened to the skull. It has a small centrum and a large neural arch. This arch, or ring, is divided by a

transverse ligament into a dorsal and ventral portion. The spinal cord lies in the dorsal portion, while an upright process (the odontoid) of the second vertebra or axis projects through the lower larger portion. On the top of the atlas are two broad facets lined with cartilage on which the skull rests. By this arrangement, as the skull is turned to right or left by the muscles the atlas rotates about the peglike odontoid process of the axis, which thus forms a sort of pivot. In nodding, the surfaces *a* and *b* (see Fig. 87) of the atlas, permit the skull to rock back and forth on them. This odontoid process is a development of the centrum of the second vertebra.

The spinal column as a whole. — The total length of the column is about twenty-eight inches in the average man. Its curvature may be readily observed from a glance at the figure (see Fig. 84). This curvature and the cartilage pads between the centra give springiness to the column, while the articulations permit a certain amount of play to the bone. The curvature also prevents a sudden jar to the base of the brain. The greatest amount of movement is between the cervical vertebræ, but the possibility of bending the whole back indicates the flexibility of the whole column.

The placing of the heavier vertebræ at the bottom of the column is an adaptation to compensate for the greater weight upon that part. The neural arch protects the spinal cord, and the apertures between the arches and openings in the vertebræ themselves protect the nerve roots which branch from the cord.

The ribs and sternum. — The ribs are flat, double-curved bones, twenty-four in number, and arranged in pairs on each side of the twelve dorsal vertebræ. They are jointed

to the vertebræ by two processes, one of which rests against the transverse process and the other against the centrum. At the front of the body the curved ends of the first seven pairs end in cartilages which unite them to the flat breastbone, or sternum. The eighth, ninth, and tenth pairs end in similar cartilages, but these are fused together into one common cartilage which is attached to the lower part of the breastbone. The eleventh and twelfth pairs are not attached, and are called floating ribs.

The sternum is a flat, irregular bone formed of three parts in the adult. The upper piece is broad, and the collar bones fit into notches in the top. On the sides are two other notches to which the first pair of ribs is attached. The middle piece is narrower, and to it

FIG. 88 — Ribs and sternum.

are attached the fourth, fifth, and sixth pairs of ribs. The last piece has no rib attachments. The ribs are so arranged as to permit the expansion and contraction of the organs of the chest cavity and at the same time to protect the inclosed organs. We shall consider their action further in our study of respiration.

THE APPENDICULAR SKELETON.

The pectoral girdle.—Attention has been called to the fact that the ends of two bones called the collar bones (*clavicles*) fit into the notches at the top of the sternum. These bones may be readily located by the fingers in front and on each side of the neck. These bones extend from the sternum to the shoulders, where their ends form part of the socket for the upper arm bone. The other parts of the socket are formed by two large, triangular bones which

FIG. 89.—The pectoral girdle; *a*, backbone; *b*, scapula; *c*, shoulder joint; *d*, arm bone; *e*, clavicle; *f*, rib; *g*, breastbone (sternum).

fit against the back and are held in place by muscles and ligaments. These triangular bones are called the shoulder blades, or *scapulas* (see Fig. 89). The scapulas, clavicles, and the upper arm bones on each side of the body meet at a common point (the shoulder joint), and the only attachment of these three bones to the axial skeleton is the place where the clavicles are jointed to the sternum. Such an arrangement permits of great play to the arms and shoulders. Since the arrangement of clavicles and scapulars is such as to encircle the upper part of the thoracic cavity, these four bones are called the pectoral girdle.

The pelvic girdle.—On the sacrum are two large facets, or surfaces, to which are attached, one on each side, two

large, irregular bones called the iliac or pelvic bones. These bones curve forward to form a basketlike support for the soft parts of the abdominal cavity. Two other bones are

FIG. 90—The pelvic girdle; *A*, diagram of bone relations; *B*, the bones of the girdle; *a*, sacrum; *b*, hip joint; *c*, leg bones; *d*, pubis; *e*, ischium; *f*, ilium.

attached to the front of each ilium. One of these, curving downward, forms a loop and is called the ischium. The other fuses with the ilium and is called the pubis. The meeting of the edges of the pubes completes the arch, or

girdle. In each ilium is a deep socket into which fits the round head of the upper leg bone.

Like the arms, the legs are attached to the axial skeleton by a girdle of bones, but, unlike the pectoral girdle, this pelvic girdle is very rigid, and the movements of the legs are confined to the bones of these members.

The arms and legs.—The bones of the arms and legs correspond very closely in number and arrangement. The

FIG. 91 — Bones of the leg and the arm.

upper part of the arm contains a single long bone (the *humerus*), the upper end of which is jointed at the shoulder to the surface of the clavicle and scapula. To the lower end of the humerus are jointed two bones (*radius* and *ulna*) which form the forearm, and this joint is called the elbow.

In the leg the upper bone, or *femur*, is jointed to the pelvic bones; it corresponds in position to the humerus of the upper arm, but the socket where it joins the ilium (the hip joint) is much deeper than the shoulder joint. The joining of the two bones of the lower leg to the lower end of the femur forms the knee. These bones are called the *tibia* and *fibula*. While the radius and ulna are of nearly equal size, the tibia, on the contrary, is much larger than the fibula, and forms the main support of the lower leg.

The lower ends of the radius and ulna are jointed to eight irregular bones called the carpals, and these eight carpals are so joined as to permit of one sliding a little over the surface of the other. This sliding gives some flexibility to the wrist, of which these bones form the support.

In the ankle are seven irregular bones which correspond to the carpals and are called the tarsals. One of these tarsals is much larger than all the others, and forms the heel. The tarsals are less movable than the carpals.

Extending from the wrist are five relatively long bones called the metacarpals which form the hand. In the foot the instep is formed by five similar bones called the metatarsals.

The fingers and thumb are formed of fourteen bones called the phalanges. The thumb has only two phalanges, the fingers three. In the foot, the toe bones are also called the phalanges; the great toe showing two, the others three.

Bones of the head.—The bones which form the head are divided into two groups, those which form the brain box being called the *cranial* bones, while those that make up the face and jaws are called *facial* bones. The two together constitute the *skull*.

FIG. 92.—The skull; 1, frontal bone; 2, parietal bone; 3, occipital bone; 4, temporal bone; 5, nasal bone; 6, malar bone; 7, superior maxillary bone; 8, lachrymal bone; 9, inferior maxillary bone.

Cranium.—This is a box almost spherical, formed of eight bones. The bone that forms the forehead is called the *frontal* bone. Back of this, forming the top and part of the sides of the box, are two large *parietal* bones; while the back of the box is a single large bone, the *occipital*. The occipital has a large opening (the *foramen magnum*)

through which the brain communicates with the spinal cord. Two smaller bones, the *temporals*, and part of the wedge-shaped *sphenoid*, complete the sides of the box. In the temporals are cavities which contain the delicate parts of the ears. The lower part of the sphenoid bone, together with another bone called the *ethmoid*, form the base of the

FIG. 93.—The skull; 1, frontal bone; 2, parietal bone; 3, temporal bone; 4, sphenoid bone; 5, ethmoid bone; 6, occipital bone; 7, superior maxillary (upper jaw) bone; 8, malar bone; 9, lacrymal bone; 10, nasal bone; 11, inferior maxillary (lower jaw) bone.

cranium and separate it from the face portion. On each side of the foramen magnum are large facets which articulate with those of the atlas.

The face bones.—The face is composed of fourteen bones, of which twelve are paired. One pair, the *malars*, or cheek bones, form the base of the eye socket and the

upper part of the cheek. Another pair, the *nasals*, form the front and sides of the nose, while between these is a partition formed by a single bone, the *vomer*. The other paired bones of the face are: (a) The *maxillæ* or upper jaw bones which carry the upper teeth and form most of the hard palate. This palate separates the nasal cavity from the mouth cavity. (b) Back of these *maxillæ* are a pair of bones which complete the hard palate and are called the *palate* bones. (c) Two small, thin bones, which form the part of the eye socket nearest the nose, are called the *lachrymal* or tear bones. (d) The *inferior turbinate* bones, or the spongy bones which lie on the inside of the nose, form the outer wall of each nasal chamber.

A single large bone, the *mandible*, forms the lower jaw, and completes the list of face bones.

Besides these bones of the face and cranium, there are in the ear of the human being, three small, separate bones known from their shape as the *hammer*, the *anvil*, and the *stirrup* bone. Another separate bone, the *hyoid*, forms the support for the base of the tongue. The mandible is the only movable bone of the skull, all the others being dovetailed together by toothlike projections (*sutures*) into one rigid structure.

Uses of the skull.—This rigid box with its rounded surface forms an ideal protection for the delicate brain. The major part of the nose and ears is made of flexible cartilage, thus protecting these projecting parts from injury by blows. The sockets that hold the eyes are of just the right depth to catch a blow upon the frontal and malar bone without allowing it to strike the eye, while the delicate parts of the inner ear are protected by being placed in cavities in the temporal bone. The opening at

the base of the skull which communicates with the neural cavity of the spinal column permits the connection of the brain and spinal cord without exposing these delicate structures to blows or pressure; while the form of the joint permits free play of the head in all directions.

BONE STRUCTURE.

Shapes of bones.—These two hundred odd bones of the skeleton may be classified into three groups according to

FIG. 94.—Kinds of bones; *A*, long bone (femur); *B*, flat bone (scapula); *C*, irregular bones (tarsals).

shape. These are the long bones, the flat bones, and the irregular bones. To the first group belong the humerus, femur, radius, tibia, ulna, fibula, metacarpals, etc. To the second group belong the ribs, sternum, patellas, scapulae, ilia, and the skull plates. In the third group are included the carpals, tarsals, facial bones, vertebrae, sacrum, and coccyx.

Structure of a long bone.— (See Ex. XLIX., *b*.) A leg of lamb or a soup bone furnishes a good example for the study of such a bone. Examination shows the following external features: At each end is an enlargement called the head, connected by a relatively narrow, cylindrical shaft. The enlarged heads are usually more or less roughened except at the very ends to allow the attachment of muscles, while the ends are coated with layers of cartilage to form a smooth articulating surface where the bone is joined to another bone. The central shaft is smooth and smaller in diameter than the ends, and usually functions mainly for support. Very few muscles are attached to this part. All parts, except those covered with cartilage, are encased in a thin membrane (the *periosteum*). This periosteum is a layer of thin, tough, connective tissue which distributes blood and through whose agency most of the new bone is formed.

If we cut this bone lengthwise, we find the following internal structure: The two heads are filled with a spongy mass of bone tissue. Over the whole outside is a thin layer of very hard, compact bone, thinnest on the heads, and thickening to form the main part of the shaft walls. In the center of the shaft is a cavity

FIG. 95 — A long bone split lengthwise; *a*, head; *b*, shaft; *c*, sponge bone and marrow.

lined with a thin layer of spongy tissue and filled with a soft, fatty, reddish yellow marrow. This marrow also fills the spaces in the spongy tissue of the heads. It is this marrow that forms the red corpuscles of the blood. It is rich in fat and gelatin (an albuminoid), and it is this last substance that forms the thickening matter of soup stock. The structure of the long bone is excellently adapted to combining lightness with strength. It is a law of mechanics that a given amount of material in the form of a hollow cylinder produces a stronger structure than the same amount of material molded into a solid rod. The bone, therefore, by its hollow structure, secures the maximum strength with the minimum of material. While it is true that they would be stronger if they were kept of the same size but made solid throughout, it would require great effort to move such heavy structures. The structure of the heads, also, gives strength and lightness at the same time, while their enlarged surfaces and roughened projections fit them for the attachment of the various muscles and tendons. These adaptations fit the long bones to form the framework of movable parts, such as the legs and arms, that must be strong and, at the same time, light enough to admit of rapid movement.

Structure of a flat bone. — (See Ex. XLIX., *a*.) A sheep's rib serves well as a type of this form of bone. Such a bone may show slight enlargements or heads at one or both ends, but these heads are much smaller in proportion to the rest of the bone than in the case of the long bone, and are entirely lacking in case of the skull plates. In the rib, there is only one such head, namely, the one attaching the bone to the vertebræ of the spinal column. The other end terminates in a mass of cartilage.

The shaft, or body, of these bones is not cylindrical, but flattened like a blade. It is this character of the shaft that gives these bones their name. The vertebral end

FIG. 96 — Ribs.

of a rib shows two cartilage-covered surfaces or articulations that mark the points where the rib is jointed to the dorsal vertebræ. The rest of the bone is covered with periosteum like the long bone.

The internal structure is best examined by sawing the bone crosswise at different points. The structure appears to be uniform throughout. The entire interior is filled with the same spongy bone tissue that was present in the heads of the long bone, while the outer walls are composed of layers of hard bone. All flat bones show these layers of periosteum, hard bone, and spongy bone. In the case of the ribs, the spongy layer is usually filled with marrow, but the quantity is necessarily small as compared with that found in the long bones.

The flat bones of the skull and pelvis form excellent supports and protection to the inclosed parts; while the flatness and thinness of the ribs, and their flexibility, permit the covering and movement of the organs of the thorax without making the chest heavy and bulky.

Irregular bones.—The general features of the irregular bones, such as the vertebræ (see Ex. XLIX., c, and p. 225), have already been discussed. All irregular bones show adaptations for the attachment of muscles or articulations. They have an outer covering of either cartilage or periosteum, with inner coats of hard and spongy bone. They are without cavities except for the spaces between the spongy tissue.

Chemical composition of bone.—If we place a clean rib in a bottle with a 20% solution of hydrochloric acid and let it stand for a few days, we shall find, upon removing it from the acid, that its character has changed. (See Ex. L.) It has the same shape as before, but is so soft as to be easily cut with a knife, and so flexible as to permit of its being tied in a knot. The substance of which it is now composed, is cartilage or gristle, and this substance is identical in structure with the cartilage that formed the pads for the articulations. Evidently, the acid has taken away the stiffening material.

If we burn a rib in a fire we note another change. The bone is again without change in form, but is very white and brittle. If we put this brittle structure into hydrochloric acid it entirely dissolves.

These experiments demonstrate that bones are made of two substances: one, cartilage which will burn but is insoluble in acid; the other, a mineral salt, of lime formation, which will not burn but dissolves readily in acid. Another feature that the experiments teach us is that these two substances are so mixed in the living bone that the removal of one or the other does not alter the shape of the bone. To the cartilage component of bone is due evidently its flexibility, while the lime salt is responsible for its stiff-

ness. All bones have this compound structure, those in which the lime is most abundant being stiffer than those in which the cartilage predominates. Bones of old persons contain less cartilage and more lime than those of young people, and hence break more easily.

If we saw a soup bone lengthwise and boil it in water there will appear on the surface a rich collection of oil drops. These come from the marrow of the bone, which is very rich in fats. If, now, the water in which the bone was boiled is allowed to cool, a solid, jellylike mass forms whose jellylike character is due to the presence of albuminoid material. This albuminoid material, usually in the form of gelatin, forms the nutrient portion of soups. Food tests also show that bones contain a small proportion of other nutrients and some flavoring material, and when such a mixture is combined with vegetables the combination makes a very nutritious food. Bones alone, however, are not to be considered as very rich in nutrient material.

Nutrition of bones.—The bones of a child are much richer in cartilage than those of an adult. As the child grows to an adult the bones increase in size and in mineral matter. The manner in which this growth takes place becomes clear only when we understand the relation of the bone structure to the circulatory system. A glance at the microscopic structure of bone shows it to be penetrated by a series of canals (the Haversian canals) by means of which the blood vessels of the periosteum are able to penetrate to all parts of the bone. From these canals the blood is enabled to radiate by a series of cross channels, or canaliculi. These canaliculi are at right angles to the canals, and connect them with a series of cavities called the lacunæ which are arranged in concentric circles about

the canals as centers. (For location of these parts, see Fig. 97.) In these lacunæ are the living bone cells which absorb the food and transform it into bone material. The lime part of this secretion forms in concentric rings about these lacunæ and the layers of this material

FIG. 97—A, transverse section of bone ($\times 150$) *a, a*, Haversian canals, *b, b*, lacunar : B, longitudinal section of bone ($\times 100$) ; *a, a*, Haversian canals ; *b, b*, lacunar : C, bone lacunar ($\times 600$) ; *c*, lacunar ; *d*, canaliculi.

are called lamellæ. When the bones are young, the secreted matter from the cells is largely cartilage; later, the mineral part is given off more freely and the structure becomes more rigid. In the marrow, as already mentioned, are formed the red corpuscles which are added to the circu-

lating blood in the bones. In general, bone is formed by the bone cells located in the lacunæ, and the cartilage and lime out of which its main structure is formed are to be understood as secretions produced by these cells out of the food supplied to the cells by the blood. This blood is brought into the bones through the periosteum and the Haversian canals, and the canaliculi are simply channels for permitting the distribution of the capillaries and their contents to the bone cells.

In the child also, many bones are separate, which in the adult fuse together. In the skull, for example, the parietals and frontals do not meet at a point in the top of the skull for some time. This spot is called the fontanelle and often is not completely closed until the child has attained an age of two years. Again, the bones of the sacrum are five in number and separate in the child; likewise the four bones of the coccyx, and the breast bones. All of these bones fuse by growth, and such fusion is called *ankylosis*. It is this tendency of bones to fuse so readily in youth that makes a broken bone knit so much more quickly in a child than in an adult; while in old age the fusing power is almost entirely lost.

XVI. THE SKELETON (*continued*).

JOINTS.

IN the living body the various bones are closely united to one another by various kinds of articulations or joints. Some of these joints permit movement of the combining bones, while others are rigid and form a fixed or immov-

FIG. 98—Forms of joints; A, hinge joint (elbow); B, ball and socket joint (hip); C, immovable joint (skull sutures).

able joint. For convenience it is customary to classify all joints as movable or immovable. To the first class belong such unions as the shoulder and hip joints, the knee, elbow, wrist and ankle joints, the finger, toe, and jaw articulations, etc. Examples of the other class are found in the *sutures* of the skull plates and the *symphyses* of the pelvic bones with the sacrum. The word *sutures* is restricted to dovetailed joints, and *symphyses* to those in which the united bones have their edges simply pressed together.

Movable joints.—There are four classes of movable joints. To these, according to the method of joining, are given the names, ball and socket, hinge, gliding and pivot joints. The best examples of the first type are the joints of the arms and legs with the shoulder girdle and pelvic girdle respectively. The knee and elbow illustrate the second type. The third type is found in the joinings of the bones which make up the wrists and ankles (carpals and tarsals). These bones slide over one another and thus give flexibility to these parts. A fourth type is illustrated by the union of the axis and atlas and the ulna and radius at the elbow.

Structure of a joint.—The joint in a leg of lamb furnishes a very good example of the structure of a joint. After all the flesh is removed the two bones will be seen to be held together by tough bands of connective tissue called ligaments. These bands are so arranged as to permit of motion in certain directions, and at the same time

FIG. 99—Section of a hinge joint.

prevent too great a separation of the bone ends. Cutting away these bands, the smooth ends of the bones with their caps of cartilage are seen. These caps prevent friction where the two surfaces rub together. As the ligaments are cut there escapes a thin, slimy liquid of the consistency of a white of egg. This liquid is called the

synovial fluid. It is secreted by a thin membrane covering the inside of the ligaments (the synovial membrane), and acts as lubricating fluid to the heads of the bones of the joint. The movement of the bones is produced by muscles which are attached to them by inelastic cords called tendons. We shall discuss these attachments further in our study of the muscles. The roughened and enlarged heads of the joining bones are special adaptations to the need of surface for the attachment of the ligaments and muscles.

All movable joints show the above structure, namely, smooth cartilage pads where the bones are in contact, synovial fluid as a lubricant, ligaments to hold the bones in place, and a synovial membrane to line these ligaments and

FIG. 100— Section of ball and socket joint.

produce the synovial fluid. The direction of movement of a joint is determined by the form of union and the method

of attachment of the ligaments. In a ball and socket joint the rounded head of one bone fits into a more or less concave cavity in the other. Such a joint allows free movement about this socket in every direction. The hip and shoulder joints illustrate this type. In the hinge joint the bones are so connected as to permit of motion in only two directions, just as a door swings upon a hinge. The elbow and knee joints illustrate this type. In gliding joints the bones slide over the surfaces of one another. The union of the carpals in the wrist and of the tarsals in the ankle are examples in point. The lower jaw shows a combination of a hinge and a gliding joint. The up and down movements being a true hinge action while the power to protrude the lower jaw beyond the upper is due to its gliding action.

In the pivot joint one bone remains stationary and the other rotates upon it. When we turn the wrist from right to left this motion is made possible by such a pivotal joint at the wrist at the point where the ulna and radius are in contact. In this case, the radius rolls over the end of the ulna carrying the hand with it while the ulna remains stationary. The movement of the atlas over the surface of the fixed axis is another illustration of this type. (See p. 227)

Immovable joints.—These are of two kinds, sutures and symphyses. The interlocking of the skull plates illustrates the former type. Here the plates are held in close union by interlocking projections much as a board is dovetailed to fit another board. The projections of one bone fitting into the spaces between projections of the other.

The symphysis type is illustrated in the union of the ilia to the sacrum. Here the two smooth surfaces of the bones are pressed tightly together and held in place by muscles and ligament.

Still another form of union is seen in the joining of the ribs to the sternum. Here a band of cartilage connects the two bones, and the flexibility of the cartilage permits of some motion.

CLASSIFICATION OF JOINTS.

Kind of Joint	Names of Uniting Bones	Region of Joint
Hinge joints .	Humerus with ulna and radius	Elbow
	Carpals with metacarpals	Wrist
	Phalanges	Fingers and Toes
	Femur with tibia	Knee
	Tibia and fibula with tarsals	Ankle
	Mandible with temporal	Jaw
Ball and Socket joints. .	Humerus with clavicle and scapula	Shoulder
	Femur with ilia .	Hip
Gliding joints .	Carpals with carpals	Wrist
	Knee cap and femur . .	Knee
	Tarsals with tarsals	Ankle
	Vertebræ with vertebræ	Spinal column
	Mandible with temporal . .	Jaw
Pivot joints . .	Radius with ulna	Wrist
	Axis with atlas	Neck
Sutures . . .	Skull plates	Skull
Symphyses .	Ilia with sacrum	Back
Other joints	Ribs and Sternum	

Hygiene of the skeleton.—The formation of bone is directly dependent upon the amount of lime salts in the blood supply. On this account it is essential that children, whose percentage of bone stiffening matter is naturally small, should have a diet rich in these mineral salts. Milk is especially rich in such material, and on this account, as well as for its generally nutritious composition, furnishes an ideal food. The disease known as rickets is a result of such lack of bone-making material in the blood.

Bones do not increase in length and size uniformly, the

heads and shafts being separated by a cartilagenous area which does not solidify until late in life. On this account bones of the young are especially subject to distortion from pressure which, if continued, may result in permanent change of shape or deformity. Another factor which makes young bones sensitive to pressure is the small proportion of lime in them. Bow legs, for example, are a result of allowing children to walk too soon, and thus subjecting the bones to the weight of the body before they are sufficiently stiffened to resist this pressure. Sitting for long periods at a bench in schoolroom when the bench is too high to allow the feet to touch the floor may permanently bend the thigh bones. A still more common deformity is what is known as round shoulders. The bending of the body over books at school, or crouching attitudes in writing, helps to increase the natural curvature of the spine and produce this unsightly deformity.

The ease with which bones may be molded is illustrated in the practice of dentists in applying pressure to the jaw-bones to straighten teeth. Braces, to remedy round shoulders, are examples of this same plasticity of bone to external pressure.

One of the worst cases of deformity from pressure is found in the tight lacing of corsets as practiced by many women. As a result, the ribs are bent and the decrease in the size of the waist is accompanied by crowding of the chest and abdominal regions to such an extent as to hinder the action of the inclosed organs.

Still another effect of pressure applied to joints is injury to the tissues and ligaments which result in the formation of painful swellings, called bunions. Tight shoes are often responsible for these painful deformities.

Fractures.—When too great pressure is applied to a bone it can bend no farther, and the result is a break. Such a break tears the periosteum and injures the surrounding tissue cells, nerves, and blood vessels. The result is a painful and serious injury. Such a break is called a fracture. The surgeon remedies such fractures by placing the two ends of the bones together and binding them firmly in place with splints or a plaster cast. When kept in this position for a sufficient length of time, the bone cells produce new bone material and the two ends are soon as firmly united as before the break. In young people this union takes place very quickly, as their bone cells are in a naturally active condition. In older persons this power of growth has become weakened, and a long time is required for the knitting together of the ends. Owing to the large proportion of lime in the bones of old people, their bones are much more brittle than those of children and much more easily fractured.

If a bone is broken in two without splintering, it is called a simple fracture. When the break is not a clean break but splinters are formed, it is called a splintered fracture. If the broken end of the bone or bones work their way out through the flesh and are exposed, the fracture is called a compound fracture.

Sprains.—When joints are bent more than is natural the ligaments and surrounding muscles are stretched and torn. The result is the production of inflammation with swelling and pain. Such an injury is called a sprain. The remedy requires the reduction of the swelling and inflammation, and hot or cold applications will usually produce the desired result.

Dislocations.—When the bending of a joint is so great as to force the ends of the bones apart, the result is called a dislocation. Accompanying such dislocations is the tearing and stretching of ligaments and muscles and all the results of a sprain. The first step in the repair of such an injury is to replace the dislocated bones in their original position. This may often be done by simply pulling on the bone until it settles back into its place. This result is usually indicated by a snap as the ends come together. After the dislocation is repaired the swelling and inflammation may be reduced by methods similar to those employed in treatment of sprains.

Joint diseases.—Inflammation of the synovial membrane often produces a great flow of liquid and a painful distention of the joint. This often occurs in rheumatism. Again, waste matter often accumulates in the synovial membranes and cartilage, and results in another painful disease known as gout.

Sometimes bacteria enter the joint and produce a discharge of yellow matter with painful swelling. Such a case is that known as tuberculosis of the joints.

If the deposit of lime matter is too great it often accumulates in the joint and fuses the ends of the bones together, making them rigid and the joint immovable. Such a disease is called ankylosis of the joints.

XVII. SKELETONS OF THE LOWER ANIMALS.¹

THE most striking point of difference between the higher and the lower animals is the difference in their skeleton or supporting framework. In fact, it is customary to subdivide all animals into two groups (vertebrates and invertebrates) on the basis of the presence or absence of a backbone.

Invertebrate animals are characterized either by the entire absence of skeletal structures, as in the slugs, worms, jellyfishes, etc., or by the presence of an external protective layer called an external skeleton. A few, such as

FIG. 101 — Protozoans (foraminifera) with external skeleton.

the cuttle fishes, have an internal framework, but it is not in any way similar to the internal skeleton of the vertebrate animals. Vertebrates all possess an internal

¹ See Footnote, p. 146, Chapter X.

jointed skeleton whose central structure is the flexible column of bones called vertebræ.

Skeletons of Invertebrates.—In certain of the protozoans the cytoplasm secretes a layer of substance which, collecting over the surface of the animal, forms a protective outer membrane or wall. In certain cases this membrane becomes the seat of deposit for lime or siliceous material, and this deposit gives it a firm character which increases its power of support and protection. (See Fig. 101.)

In the sponges certain cells in the body produce spicules of lime, or silica, or spongin, which are embedded in the

FIG. 102 — Skeleton of horny sponge.

walls of the animal and form an interlacing network. The ordinary sponge of commerce consists of the spongin skeleton of such animals with the animal parts removed.

In certain of the coelenterates, such as the coral polyps, the walls secrete a limestone matter, which increases to such amounts that the animals which produce it finally lie in crevices of this hardened secretion, which thus forms masses often larger than the animals themselves.

These secreted skeletons are all more or less rigid, and while they function as supports, or protective material, they

FIG. 103 — A coral colony; living polyps in limestone which they have secreted.

are entirely without joints. Such a skeleton is evidently entirely unadapted for animals that are to move about in search of food, and whose parts must be capable of movement. Such animals require a modification of this rigid, one-piece skeleton, and in the starfish we find a simple change in this direction.

FIG. 104 — Arrangement of plates in a starfish; M, soft membrane; P, hard plates.

These animals are covered with an external limestone secretion which is in the form of plates, but these plates are small and are united by con-

nective tissue of a flexible character, so that, like a chain-mail shirt, the parts of the body may bend and still be protected and supported.

The mollusks (oysters, clams, snails, etc.) have a certain portion of the body, called the mantle, whose function it is to secrete hard material, and this material collects to form a structure called a shell. These shells are external and completely inclose the soft body. Those mollusks that require an opening for the entrance of food and air, and that, under ordinary conditions, are stationary, secure this result by having the shell hinged and provided with muscles which can open and close it when

FIG. 105 — Left shell of clam; *h*, hinge ligament; *t*, *c*, *t'*, hinge teeth; *a*, *a'*, points of attachment of the muscles, by means of which the clam closes its shell.

necessary. Others, such as the snails, have the shell of one piece, but so mounted that the animal can carry it on his back when moving about, and when disturbed can withdraw the body into it until the disturbing factor has disappeared. Still other forms of mollusks have the shell inside the body, where it serves to give the body a certain stiffness, as in the cuttlefish and squid. These internal shells are of no use in protecting the soft parts of the body.

In the arthropods (insects and lobsters, etc.) we have

the most perfect type of external skeleton found. In these animals the skeleton is formed of a horny material called chitin, which completely incloses all parts of the body, except the apertures for the mouth and other open-

FIG. 106 — External jointed skeleton (lobster).

ings. By making this secretion less stiff at points, very perfect joints are formed which allow a great freedom of movement to the legs and body. The arthropods are, in fact, animals encased in a suit of excellently jointed armor. This form of external skeleton, while excellent as a protection, and well adapted to movement, necessarily restricts the growth of the animal. In other words, when the crustacean increases in size he is in the same situation as the small boy in a tight suit of clothes, and the result is the same, he bursts the clothes and must have a new and larger suit supplied. The manner in which this is done is by withdrawing the body from the old, broken covering, and after expansion, allowing the new one to form from

secretions of the soft skin. Meanwhile the animal is subject to all sorts of dangers from its enemies, since between the time of shedding of the old skin and the forming of the new it is wholly unprotected. This is one reason why it so difficult to raise enough lobsters to supply the demands of the people, although the number of eggs hatched each year is enormous. The young are eaten between molts by all sorts of enemies.

Vertebrate skeletons. — Fishes, frogs, reptiles, birds, and mammals all agree in the possession of a jointed, internal framework, and differ only in the number of the bones, the composition of the bones, and the arrangement with regard to one another. All possess a central column or backbone made up of separate pieces called vertebræ, and on this column are hung the other bones of the body.

The fish skeleton.— The skeleton of the fish differs from that of the other vertebrates in containing a higher pro-

FIG. 107 — Skeleton of a fish ; note the great number of bones.

portion of cartilage. A few forms show remnants of an external skeleton in the form of the scales which in the garpike are so hard as to offer some protection to the underlying parts.

The absence of lime in the bones of the fish makes them extremely flexible, and fits the body for the bending necessary to swimming motion. Furthermore, the support which the water furnishes to the body makes such hardness unnecessary as a supporting structure. Another feature of the fish skeleton is the great number and small size of the bones. (See Fig. 107.)

The frog skeleton.—In the amphibians, of which the frog is a type, we find a marked difference in the composi-

FIG. 108—Skeleton of a frog.

tion and number of the bones, while the arrangement of the separate bones much more nearly follows the order in our own bodies. The explanation of this change is found in the adaptation necessary to fit the animal for a land existence. As soon as the animal is removed from the support of

water, the flexible, cartilage bones are unable to support the body, and we should expect, therefore, to find the bones of the frog more rigid than those of the fish. The new method of motion requires the modification of fins into legs and arms, and the front legs of the frog are supposed to have developed as modifications of what we call the pectoral fins of the fish. In the same way, the hind legs of the frog are modifications of the pelvic fins of the fish. Another reason for the rigid character of the frog skeleton

is the necessity for a stiff framework against which the legs may push in propelling the body. If the backbone were too flexible it would simply bend when the legs pushed against it, and the animal would not move any more than does an object when pushed with a feather.

The frog skeleton will be seen to differ from that of the higher forms of animals, though not so much as from that of the fish. For example, the frog has a much shorter backbone than is found in higher forms, the ribs are wanting, and the skull is made up largely of face bones with a very tiny cranium or brain box. As a machine for support and motion, it is very primitive compared with our skeleton. The legs are so attached that the thigh bones point outward from the body instead of being under the part to be supported. The shoulder girdle is rigid and the elbows point outward, which makes it awkward to use the front legs as pushing agents. In fact, in the frog these front legs are of small use in locomotion on land; the greatly elongated hind legs being so arranged as to make leaping the natural mode of progress.

Reptile skeletons.—The reptiles differ from the amphibians in possessing ribs (see Fig. 109), which have been de-

FIG. 109 — Skeleton of a crocodile.

veloped to protect the sides of the body and to aid in the operation of the lungs as breathing organs. In the snakes the legs are lacking and there is no breastbone. In all the

other forms, there are a breastbone and well developed legs. These legs, however, are poor organs of locomotion and support, for, like the frog's, the thighs point outward, and instead of raising the body from the ground the legs are so short that the belly rests on the ground and must be dragged

over the surface. The elbows point backward more than in the frog, and the arm, or foreleg, is, in this respect, an improvement over the similar structure in the frog.

Bird skeleton. — The bird skeletons are interesting as studies in adaptation. In these animals everything has been sacrificed to lightness and a flying habit. The bones are hollow and contain no marrow. The arms are developed as wings, the shoulder girdle is sturdy and

FIG. 110 — A bird skeleton (vulture).

rigid, to form a firm support for the wing joints, while the greatly elongated breastbone, with its deep keel, is an excellent adaptation for the attachment of the powerful flying muscles. Both the birds and reptiles have a larger cranium than the frogs or fishes. Another feature is the arrangement of the legs. In the bird the toes are the only part of the foot that is in contact with the ground, for the heel is

raised and the metacarpals are limited to one bone. The S-shaped position of the thigh bone, tibia and metacarpal forms a splendid arrangement for preventing a jar when the bird alights.

Mammal skeletons. — The most striking differences between the mammal structures and the human skeleton have to do with the adaptation of the body to an erect position, and the consequent freeing of the arms from the necessity of support. The development of the hand as a grasping organ, and the relations of the bones in the different forms of the arm and leg structure are interesting also as showing adaptations to different modes of use. The following figures illustrate the relations of the bones of the

FIG. 111 — Typical fore limbs ; A, of man ; B, of cow ; C, of cat ; D, of horse ; E, of whale ; F, of bat ; G, of mole.

arms and legs in animals of different habits; the important point to be observed is the manner in which the same

bones are modified to serve different uses. Nature's economy in this respect is remarkable. For example, in Figure 111 note the arrangement of the hands in the mole and their adaptations for digging. Note, also, in the cat the development of the finger and toe nails into claws for tearing. The flipper of the whale shows a structure very similar to that of the fins of the fish; in the horse and the cow, the development of the nail into hoofs and the reduction of the metacarpals into a single bone support,

FIG. 112—Typical skulls ; A, of man ; B, of cow ; C, of bat ; D, of mole ; E, of cat ; F, of horse ; G, of whale.

is interesting. In the bat, the enormously elongated phalanges serve as a framework for the membranous wings.

Note, also, the differences in the legs of these animals from that of man. In man, the whole foot rests on the ground; while in the cat, the horse, and the cow the toes alone support the body, thus permitting the development of powerful thigh and calf muscles at points where they will

not interfere with the movement by their bulk and yet give tremendous power to the pushing action of the legs.

Finally, a comparison of the skulls (see Fig. 112) is interesting as an illustration of the progression in size of the brain box, and the increase in intelligence as we go higher in the scale; while the jaws with their teeth modifications show every variety of adaptation to different foods and modes of eating.

General conclusions. — We may note, in conclusion, that the bones of all vertebrate animals are remarkably similar in arrangement and position, but that every animal shows modifications in structure and position, that fit his particular framework to his needs.

PLATE III — Muscles.

XVIII. MUSCLES.

KINDS, STRUCTURE, AND COMPOSITION.

IN the skeleton we have a well-jointed framework that of itself has no power of movement. In order to move the different bones and produce the different contractions and expansions of the organs of the body, a special tissue, called the muscles, is devised. Muscles, then, are responsible for all movement in our bodies.

Kinds of muscle tissue.—Every boy knows that when he bends his forearm at the elbow a swelling appears on the upper side of the upper arm. This swelling, which the boy speaks of as his “muscle,” is, in fact, a separate mass of muscle tissue and known technically as the biceps muscle. Its action illustrates two characteristics of muscle tissue. First, the production of motion by contraction. Second, that contractions are under control, for in this case the boy determines when they shall contract. The muscles which form the walls of the intestines and produce the grinding movements that we call peristalsis, do so by contraction also, but their action differs from the action of the biceps and triceps in that it is not under the control of the will. We cannot stop the action as we wish.

It was formerly customary to make this matter of control the basis of classification of muscles. Muscles whose contractions required no act of will on our part were called involuntary muscles. Those which are subject to our direct control, like the muscles of the fingers, arms, and legs, were called voluntary muscles. The difficulty of this

classification arises from the fact that some muscles are both voluntary and involuntary, for example, the muscles that close the eyelids.

Furthermore dissection has shown that all muscle tissue in one body may be put into one of three classes,

FIG. 113—Biceps muscle.

each with a distinct structural formation and it is this structural classification we shall adopt. These three classes are known as *striated*, *non-striated*, and *heart muscle*.

These three kinds of muscle tissue combined constitute nearly half of the entire weight of the body, and in their action is consumed the greater part of the food and oxygen supply of the body.

STRIATED MUSCLES.

Distribution and form.— (See Ex. LIII.) If we remove carefully the skin of any vertebrate animal, we find masses of flesh or meat just below this skin. This flesh is composed mainly of striated muscle tissue. We may note further, that this mass of flesh is composed of separate bundles of varying size and shape, each bundle covered with a glistening sheath and attached to the bones only at certain points. Such a bundle constitutes a separate muscle, and there are five hundred of such separate bundles or muscles in our body. In their sum they determine the form of the animal body. Although there are so many of these muscles, they admit of classification under comparatively few heads. The type forms are as follows:

Bellied muscles. These muscles, of which the biceps muscle already referred to is a type, receive their name from their shape. They are spindle-shaped bundles with a swelling or belly in the center, and taper at each end into one or more white, inelastic cords called tendons (see Fig. 114).

FIG. 114 — Bellied muscle; *a*, belly; *b*, *b*, tendons.

These tendons serve to attach the body of the muscle to the bones which it is to move. As these muscles contract, the belly swells and the spindle shortens, drawing the two attached ends nearer together. If these ends are attached to movable bones, these bones must necessarily move as the spindle shortens. In some cases, the bones at each end move. In others, only one of the bones moves. The end of the muscle that is attached to the bone

that moves most is called the *insertion* of the muscle. (See Fig. 115, A.) That attached to the end which moves

FIG. 115—A, bones of the arm, showing origin and insertion of biceps (a) and triceps (b) muscles; B, a digastric muscle.

least, is called the *origin* of the muscle. By following with the fingers the curve of the biceps muscle, the tendon that attaches it to the radius bone of the forearm and its place of attachment to this bone can be readily felt. The other end is covered with other masses of muscle, but dissection shows it to end in two tendons which are attached to the shoulder blade. In bending the forearm, it is evident that the radius moves most; hence the radius end of the muscle is its insertion in this case, and the shoulder blade its origin.

It must be noted, however, that these terms insertion and origin are merely relative terms. For example, in the operation of "chinning" oneself on a horizontal bar the radius bone remains fixed and the shoulder blade is the bone moved. In this case the radius end is spoken of as the origin of the muscle, and the shoulder-blade end as the insertion. Most muscles, however, are so arranged that one end is always the insertion and the other the origin.

Many bellied muscles have a single tendon at each end.

Others divide at one end into two tendons, and are called biceps or two-headed muscles on that account. Still others divide into three parts and are called triceps muscles. In another form of muscle there are two bellies attached by tendons and with tendons at each end. Such a form is called a digastric muscle (see Fig. 115, B); while others show as many as four bellied portions, and are called polygastric muscles.

Not all muscles end in tendons. In some cases the muscle ends are attached directly to the bones. Nearly all the muscles of the arms, legs, hands, and feet are of the bellied variety.

Flat muscles. On the trunk are found many muscles which are flat and attached directly to the bones. Such muscles are called flat muscles. They are of nearly uniform thickness throughout, and are found in places where the belly would interfere with the action of other organs or make the part too bulky. The muscles on the chest are of this variety.

Arrangement of striated muscles. — The bellied muscles of the body are usually arranged in pairs (see Fig. 115, A). Those which bend bones about a joint as pivot when they contract are called flexors. The biceps muscle is a type of a flexor muscle. Those, on the contrary, which by contraction, extend a bone about a joint as pivot, are called extensors. The triceps muscle, whose insertion is the end of the ulna and whose origin is the humerus and scapula, is an example of this type. The flexing and extending of the forearm is due to the alternate contraction and relaxations of these two muscles and illustrates the typical paired arrangement.

Gross structure of striated muscle. — (See Ex. LIV.).

A bellied muscle, dissected from the leg of a rat or frog, illustrates clearly the gross structure of a muscle. Exam-

FIG. 116. — Flat muscles of the back.

ination of such a dissected muscle shows it to be entirely inclosed in a connective tissue sheath called the *perimysium*. This sheath is thin and transparent, and, at the

edges of the muscle, fuses into a glistening, white, inelastic cord called the tendon. These tendons are very useful in the body, because they permit the contracting or belly part of the muscle to be placed at some distance from the part to be moved, and the inelastic cords transmit these contractions without loss of power to the parts to which they are attached, and, at the same time, this arrangement avoids bulkiness. Thus, the fingers are flexed and extended by muscles whose bodies are located in the forearm, and whose contractions are transmitted to the finger bones by tendons attached to the bones at one end and to the muscles at the other. The extensor tendons on the back of the hand can be readily followed to their origin and insertion. It is evident that such an arrangement gives much greater delicacy and ease of manipulation to the hand than would be the case if each finger were attached directly to a muscle. This tendon method of attachment is especially noticeable

in the limbs where bulkiness is most undesirable.

FIG. 117 — Bundles of striated muscles cut across ; *f, f*, several bundles bound together to make up the main muscle.

If we boil a piece of the dissected muscle or a piece of lean beef (the bellied muscle of a beef creature), it readily separates into a series of smaller prism-shaped bundles, and these may be easily picked apart. Examination of one of these prism-shaped bundles shows

it to be covered with the same sheathlike perimysium which in fact is merely a partitionlike extension of the outer peri-

mysium. If we proceed further, and cut across the end of one of these bundles, or fasciculi, we find it to be divided into smaller bundles by perimysium partitions, and these, in turn, into still finer partitions and bundles. These ultimate bundles may be teased apart into a certain number of microscopic muscle fibers too small to be examined with the naked eye. These fibers are the structural elements of the muscle. It is the combined contraction and expansion of these fibers that produces the complete muscle action.

Microscopic structure of striated muscle.—Each muscle fiber, when examined under the microscope, is found to be from $\frac{1}{3}$ inch to $1\frac{1}{2}$ inches in length, but only $\frac{1}{750}$ inch to $\frac{1}{450}$ inch in diameter. They are spindle-shaped and taper to a point at each end. They evidently do not extend the entire length of the bundle, which may be a foot or more in length, but are packed together, side by side, the pointed ends of one fiber fitting in between the ends of others, so that each fiber extends lengthwise of the bundle.

The fiber of a striated muscle may always be recognized by its striped or striated appearance. These stripes extend transversely across the fiber and pass entirely through it. They are of two kinds, a broad, bright band, and a narrower, darker band. These dark and bright bands alternate. With the high powers of the microscope can also be made out dim longitudinal striations.

If we press upon one of these fibers with a needle point it is seen to be covered with a transparent membrane or sarcolemma which surrounds the striped, contractile, inner substance. This outer coat or *sarcolemma* is a tough transparent membrane. Visible through the sar-

colemma and scattered over the body of the fiber are several oval bodies which indicate the true nature of the fiber. These oval bodies are nuclei and the fiber is actu-

FIG. 118 — A, muscle fiber with sarcolemma removed and broken at one end into its fibrillæ ; B, separate fibrillæ ; C, muscle fiber breaking into disks ; D, muscular fiber, the contractile substance of which (*a*) is torn, while the sarcolemma (*b*) has not given way ($\times 350$).

ally a composite muscle cell. Such a cell is called a multinuclear cell and the sarcolemma is the common cell wall.

Viewed as a multinuclear cell the above structures bear the following relations to the cell as we know it. The sarcolemma is the common cell wall. The contractile substance is protoplasm in which the power of contraction and expansion has become greatly developed. Such specialized protoplasm is called *sarcoplasm*, or muscle-

protoplasm. The oval bodies are nuclei. The faint, lengthwise striations indicate a peculiar, structural division of the sarcoplasm. Examination shows that it is divided into long hardened portions or *fibrils*, which extend lengthwise of the cell, and the spaces between these fibrils are filled with a liquid portion more like ordinary protoplasm. It is the faint outlines of these fibrils that give the lengthwise striations. These columns are also called *sarcostyles* to distinguish them from the liquid sarcoplasm. Each sarcostyle is divided transversely into disks of alternate dark and bright appearance, and these combined disks are what give the cross striated appearance. Both the sarcostyles and the sarcoplasm are merely modified forms of protoplasm. The part played in muscle action by these various elements of the fiber cell will be discussed later under muscle action. (See p. 278.) At present, all we need note is that the substance of a muscle cell has a highly specialized power of contraction, and that the combined contractions of many cells united into bundles and these into other bundles, is responsible for the ultimate action of the muscle, just as the combined pull of many men in a tug of war results in one motion of the rope. Most striated muscles are under the control of the will and were formerly included under the heading of *voluntary muscle*.

UNSTRIATED MUSCLES.

Muscles of this class are found in the walls of the digestive tract and organs, the blood vessels, the genital and urinary organs, etc. Such muscles show no tendons, but form expanded membranes that surround cavities. Therefore they have no definite origin or insertion. The fact

that their contractions and relaxations are controlled by nerve centers which are not under direct will controlled to their being called involuntary muscles.

Structure.—The membranes which unstriated muscles form are not in bundles like the striated muscles, but are composed of masses of single, spindle-shaped cells stuck together by a cementlike substance and held in place by an interlacing network of connective tissue. These cells are only about $\frac{1}{600}$ inch in length, and while they show a longi-

FIG. 119—A single unstriated muscle cell; N, nucleus.

tudinal striation *there are no cross striations.* The cells are covered with a cell wall of transparent membrane and inclose a single nucleus. They are, therefore, to be considered as single and not composite cells. The protoplasm which they contain has the power of contraction in the direction of their long axis, and since they are usually arranged about a cavity their contraction and relaxation results in the reduction or enlargement of the cavity.

FIG. 120
— Mus-
cle cells
of the
heart.

HEART MUSCLE.

The muscular walls of the heart are composed of a tissue which is intermediate in structure between that of the striated and unstriated types. It consists of single, nucleated, branched cells united to make a meshwork. These cells show cross striations, but these striations are less distinct than those of the skeletal muscles. The rate of contraction of the heart-muscle cells is less than

that of the ordinary striated tissue and greater than that of the unstriated. It is also remarkable for the rhythmic character of its contractions.

COMPOSITION OF MUSCLE.

Applying the common food tests to muscles we find them to be composed mainly of proteid and water with a small quantity of the salts, carbohydrates, fats, nitrogenous wastes, and enzymes.

Muscle proteids.—There are two forms of muscle proteid: one known as myosin, the other as myogen. The myogen is most abundant in living muscle, for it is present to three or four times the bulk of the other. Both are coagulated by heat, but the myogen requires a higher temperature (131°–149° F.) than the other (113°–122° F.). Both are similar to the fibrinogen of the blood, for when they are allowed to stand at ordinary temperatures they are capable of forming a clot, and of assuming an insoluble form called myogen-fibrin or myosin-fibrin, just as fibrinogen is transformed into fibrin by the clotting of blood. Dead muscle then consists mainly of clotted myosin and myogen, or myosin-fibrin and myogen-fibrin. This clotting is supposed to explain what is known as “rigor mortis” or the stiffening of the muscles after death.

Muscle carbohydrates.—The carbohydrates present in muscle are sugar and glycogen. It is supposed that muscle has the power to convert sugar into glycogen and glycogen into sugar as needed. In action the glycogen is first transformed into sugar and then consumed in the production of the energy. We can assume then that the glycogen in the muscle represents sugar brought to it by the blood, converted by it and stored in the form of glycogen until the muscle needs it. It is a reserve supply of fuel in the muscle.

Muscle Enzymes.—Muscles contain a number of enzymes. The following have been identified: A proteolytic enzyme capable

of digesting proteid, and an amylotic enzyme which changes glycogen to sugar, a glycolytic enzyme which destroys sugars, a fat enzyme or lipase which breaks up fats, and finally an enzyme which causes the clotting of myogen and myosin after death.

Inorganic muscle constituents. — Muscle contains many salts which aid in securing an interchange of nutrition by osmosis and dialysis with the lymph and blood through the sarcolemma, and they also give to the muscle its irritability and sensitiveness to nerve stimuli.

Water forms 75% of the living muscle.

Pigments and other components. — The red color of muscle is due partly to the blood it contains, and partly to a pigment, similar to hæmoglobin, called myohæmatin, which probably aids the muscle cell to absorb oxygen.

Muscle also contains several organic acids and nitrogenous wastes.

Nutrition of Muscles. — Muscles are penetrated everywhere by blood vessels. From the larger blood vessels smaller branches are given off, and these end at the capillaries which surround the finest bundles of the muscle mass. From the lymph which comes from these blood vessels, the muscle fibers select their food and give off their wastes to it. (See p. 175.) From the blood they also receive oxygen and give up the carbon dioxide to it.

XIX. MUSCLES (*continued*).

THEIR ACTION.

OUR study of the structure of the muscles has taught us that movement is produced by the contraction of the muscles which are connected to movable bones or so arranged as to inclose cavities. Further, we have learned that this power of contraction is located in tiny, microscopic fibers or cells, the sum of whose combined contractions is the movement of the entire bundle. Finally, we have seen that all parts of these tiny cells are not equally endowed with this power, but that it is limited to certain contractile substances found in the cell fibers.

FIG. 121—Frog's leg, dissected to show the muscle and nerve connections; *g*, gastrocnemius muscle; *sc*, sciatic nerve; *ad*, *pe*, etc., other muscles.

Study of a frog's muscle.—(See Ex. IV.) If we dissect away the belled muscle known as the gastrocnemius or calf muscle of a frog's hind leg we shall find a white threadlike fiber lying close to its surface. This fiber branches into many separate threads, and these, in turn, enter the body of the muscle. By means of certain stains it is possible to follow these branches and find that

ultimately they end in the walls of the muscle fibers themselves. The large thread, with its branches, is a nerve, and in the living frog the large end is continued backward to the spinal cord through which it connects with the brain. If we remove all the muscles from the leg of a freshly killed frog, except the gastrocnemius, then arrange such an apparatus as shown in Fig. 123, and bring the two ends of an electric circuit in contact with the nerve, the muscle will suddenly shorten and then relax again as though the

FIG. 122 — Nerve endings in striated muscle fibers.

FIG. 123 — Nerve muscle preparation; *s*, set screw; *c*, clamp, *f*, femur; *m*, gastrocnemius muscle; *n*, *n*, sciatic nerve; *h*, hook; *l*, lever; *e*, electrodes; *b*, battery.

frog were alive. If now we remove the ends of the wires from the nerve a second contraction and relaxation takes place. Evidently the shock of making and breaking the electric current is transmitted by the nerve to the muscle fiber and acts as a stimulus to the contractile substance which it contains. Furthermore, the experiment shows us that a stimulus, such as was given by the electric current that we

sent over the nerves to the fiber, is sufficient to set in contraction at one time all the muscle fibers of such a mass.

Such a simple experiment also enables us to understand how a desire formulated in the brain may be transmitted as a nerve impulse to the muscle cells and thus produce movement on the part of the muscle which receives such impulse. Every muscle in our body is supplied with a nerve connection, the ultimate ends of the nerves being embedded in the muscle fibers while the main trunks are collected into the great trunk called the spinal cord, and through that are in connection with the brain. The brain or cord, therefore, is the starting point of the impulse that causes the contractions of the muscle fibers, and this arrangement puts all the complicated movements of the body under their control.

The power of cells to respond to stimulus is called irritability. By irritability in general we mean the power of a cell when stimulated to exhibit that particular form of activity for which it has been specialized. Muscle cells, for example, have lost many of the characteristics of a free moving *amœba* but have developed the power of contraction to a high degree. A secreting cell, on the other hand, has little power of contraction but a highly developed power of secretion. Hence, the irritability of the muscle cell shows itself in contraction when stimulated; the gland cell, on the other hand, manifests increased secretory power when stimulated. In other words, each possesses irritability, but the expression of that irritability depends upon the manner in which the cell is specialized. Any impulse which can produce an expression of the cell's specialized power is said to excite its irritability.

If we plunge the ends of the electric wires directly into the muscle substance the mass will contract and relax as before, but not to so great an extent. It is evident from

this result that the nerves are especially well adapted for the carriage of impulses to the fibers, though they are not the sole means of stimulating the fibers. The sudden contraction of a muscle when struck is thus explained as merely a response of the cells to another form of stimulus. The nerves, however, constitute the normal means of transmitting an impulse to the muscle from outside.

Method of muscle contraction under stimulus. — If, in place of a simple apparatus, such as that described in the preceding paragraph, we substitute an instrument which can apply a rapid succession of shocks to the nerve end, it is possible to make the stimuli so rapid that the second contraction will begin before the first relaxation has ended. Under these conditions the muscle rapidly shortens until it finally reaches a point where it can contract no more. Successive stimuli will, then, merely serve to maintain it in a contracted state until finally the contractile power becomes exhausted, and extension occurs in spite of stimulation. Such a progressive series of contractions are spoken of collectively as a tetanic contraction. It is supposed that in the living body all nerve impulses are of this tetanic character, as the time of a simple contraction is only about a tenth of a second, and very few movements are of so short duration.

COMPARISON OF MUSCLE AND STEAM ENGINE AS ENERGY PRODUCERS.

The substance in the muscle cells has the power of irritability, and the nerves are the natural conductors of the stimuli which excite this irritability. What happens in a muscle fiber or cell when this stimulus is received? We

can best understand this by comparing the results with what takes place in an ordinary steam engine. The comparison of the methods of energy production in the two cases are shown in the following table.

The first step in the production of energy is combustion.

TABLE OF COMPARISON OF STEAM ENGINE AND MUSCLE CELL.

Conditions Necessary to Combustion in —

The Steam Engine	The Muscle Cell
<ol style="list-style-type: none"> 1. The presence of a combustible chemical compound, <i>i.e.</i>, one which can combine chemically with oxygen (Fuel). Coal, wood, or other carbon containing matter. 2. Some initial impulse (kindling) necessary to start rapid oxidation or combustion. Application of match flame or other kindling material. 3. There must be present a sufficient supply of oxygen to permit of rapid union with the fuel. Supplied by draughts of various kinds. 	<ol style="list-style-type: none"> 1. The presence of a combustible chemical compound, <i>i.e.</i>, one which can combine chemically with oxygen (Fuel). The carbon compounds, glycogen, sugar, proteid. 2. Some initial impulse necessary to start oxidation (kindling). The nervous impulse received by the cell substance. 3. There must be present a sufficient amount of oxygen to permit of rapid union with the fuel. Supply secured from the red corpuscles of the blood and from reserve supply stored in cells.

Results of Combustion in —

The Steam Engine	The Muscle Cell
1. Energy in form of heat.	1. Energy in form of heat.
2. Carbon dioxide as result of the union of the carbon in the fuel with the oxygen.	2. Carbon dioxide as the result of the union of the carbon in the fuel with the oxygen.
3. Wastes in the form of ashes or other unoxidizable material.	3. Wastes in form of lactic acid, urea, etc.

Disposal of the Products of Combustion in —

The Steam Engine	The Muscle Cell
1. Part of the heat energy transforms the water in a boiler to steam and shows itself in the power of expansion that this steam has. If this expansion is allowed to act upon a piston rod it is transformed into motion.	1. Part of the heat energy acts upon the fibrils (sarco-styles) or other contractile matter in the cell and causes it to shorten. The result is a shortening of the whole cell, <i>i.e.</i> , contraction or motion. When the heat supply stops, this contractile matter relaxes.
2. Part of the heat energy is lost as heat, and merely warms the surrounding air and parts of the engine.	2. Part of the heat energy escapes as heat, and thus raises the temperature of the body.
3. The carbon dioxide passes off with the smoke through the chimney or smoke stack.	3. The carbon dioxide passes off in the blood and is removed from the body by the lungs and skin.
4. The wastes pass out partly as smoke through the chimney, and partly as ashes into the fire box.	4. The wastes are carried off by the blood and lymph and removed from the body by the excretory organs, skin and kidneys.

Let us examine this analogy more closely to see how this explanation of the production of contraction and expansion in a muscle cell fits in with known conditions for muscular activity.

In the engine, if the supply of fuel becomes lessened, the amount oxidized is necessarily less and the power to do work becomes diminished. Likewise, in the muscle, any interference with the food supply must decrease the amount of fuel and lessen the working power of the muscles. It is essential, therefore, that the muscles be provided with a good supply of blood, and that this blood contain the proper kind of food for fuel. It is for this reason that athletes and others who use their muscles extensively are limited to a diet rich in the foods required for muscle building and combustion. Meats, coarse breads, eggs, vegetables, fruits, supply all the important carbohydrates and proteids. Fats and pastries are usually excluded, not because they do not contain energy producing material but because they are difficult to digest.

Tobacco and alcohol also tend to diminish the irritability of the cells, and should be excluded from the diet. What is good for the athlete is evidently well suited for the growing boy and girl, and, in general, the food of all people should contain the materials suited to the upbuilding of muscle cells and, at the same time, fitted to provide the fuel necessary for their action.

Again, if the fire of the engine is not supplied with a good draught, that is, a constant supply of oxygen, the production of heat must necessarily stop. So, in the body, the muscles must be supplied constantly, not only with food-laden blood, but also with oxygen-laden blood. This means that the lungs, from which this supply is

obtained, must be in good condition, and that the air which we take into them must be rich in oxygen. Bad ventilation diminishes the oxygen content of the blood, and diminishes the working power of the muscles by just so much.

Every one knows that it is necessary at times to let the engine fires go out in order that the accumulated wastes may be removed, and in order that the heated parts may cool. So, with the muscles, the constant use of the cells soon causes an accumulation of waste too great for the circulation to remove, the body becomes overheated, and soon the muscle loses its power to respond to a stimulus. Such a condition is called muscle fatigue. We feel tired, and this tired feeling is an indication that it is time to rest and allow the cells to recover from their activity. If we heed this call and permit the used part to rest for a while, the blood is able to remove the accumulated wastes during this period, and stock the cells with new fuel, and, in general, restore the muscle cells to their working condition. On this account, children and manual laborers who use their muscles much more constantly than brain workers, require much longer rest periods. Often a change of work will accomplish a restoration of the muscles, provided the new work calls into play a different set of muscles and allows the others to rest. A certain amount of sleep, however, is necessary for every healthy man, woman, or child.

Exercise.—Muscles possess another property besides contractility, namely, elasticity. For example, if we suspend weights at one end of a freshly dissected frog's muscle, the muscle readily stretches to accommodate the increase in weight, and as readily returns to its original length when the weight is removed, provided we do not

allow the weights to exceed a certain limit. In our bodies a healthy muscle is always in this state of elastic tension, as is shown by its shortening when the tendon which attaches it is cut. This tense condition makes the muscle respond much more quickly to stimuli, and is of great value in mechanical movements of the body. Muscles which are not used tend to lose this elastic tension and to become soft and flabby. The only method of remedy is to give them plenty of use or exercise. Exercise, therefore, is of value in keeping our muscles in a proper state of tension, and, at the same time, aids in the proper circulation of food and oxygen material to the used parts. Care should be taken in the selection of the exercise that it shall affect all the muscles of the body and not center upon any one set of muscles. Rowing and running are particularly valuable as forms of exercise which call into play many muscles; and walking is one of the best forms, since it exercises the body and leg muscles and the lung and heart muscles, and is not violent enough to over-fatigue any given set. What are called "setting up" exercises in the army and in our schools are particularly good, since they have been devised to fit every part of the body and to give a general toning up of the system rather than to develop any one set of muscles.

Another feature of exercise that should be borne in mind is that it should be regular. A little regular exercise each day, calling into play as many muscles as possible, is far more valuable than prolonged exercise one day, followed by a long period of no exercise. Further, since active muscle makes great demands upon the blood supply, we should not exercise immediately after eating when the blood is needed elsewhere.

THE MECHANICS OF MUSCLE ACTION.

In a steam engine the energy liberated by combustion is in the form of heat. Only part of this energy can be converted into motion and into the lifting of weights or overcoming of resistance of various kinds. The rest of the energy is lost as heat. In the best steam engines the amount of heat energy that can be converted into work represents only ten to fifteen per cent of the entire energy liberated by the combustion. We express this relation of the useful energy to the total energy liberated in any given machine, as the efficiency of the machine. A steam engine, then, has an efficiency of only 10% to 15%. Experiments with the muscles have demonstrated that the ordinary muscle has an efficiency of something like 33%. In other words, nearly a third of the energy liberated in the combustion of fuel in the muscle cell can be converted into useful work, while the rest is lost in heat. While, perhaps, this amount seems small compared with the heat loss, if we compare the muscle with the best steam engines it will be seen that it is a remarkably efficient machine. This efficiency represents an average, and differs in individuals.

The simplest method of expressing this useful energy in any machine is in terms of the weights or resistance which is overcome and the distance through which the action takes place. Thus, the energy necessary to raise a pound one foot is expressed as one foot pound of energy. In other words, work is equal to the force exerted, times the distance through which it acts. In formula: $\text{Work} = \text{Force} \times \text{Distance}$. It is evident from this formula that a given amount of energy may move a large body a short distance

and a small body a long distance without varying the total value of the work done. Thus with ten foot pounds of energy we may raise a ten-pound weight one foot or a one-pound weight ten feet.

Men have contrived various pieces of apparatus called machines by means of which weights, and the distances they are moved, may be varied without changing the total amount of energy required. One of the simplest of these machines is the lever.

Laws of the lever. — A lever is a rigid bar turning about a fixed point or pivot. There are three classes of levers known respectively as first, second, and third class levers.

FIG. 124 — Levers ; I, lever of the first class ; II, lever of the second class ; III, lever of the third class ; W, weight ; F, fulcrum, or pivot ; P, power.

In a first-class lever the pivot is between the weight to be raised and the force applied to raise it. (See Fig. 127.) In the second-class lever the weight is between the pivot and the power, while in the third class the power is between the pivot and the weight. In each of these classes the points nearest the pivot evidently move a less distance than those farther away, and by arranging at different distances from the pivot the power and weight applied, it is possible to make a small power balance a large weight and *vice versa*.

Expressed in terms of distance from the pivot this relation is known as the *law of levers*.

Weight times the perpendicular distance of its point of application from the pivot equals the power required to balance it times the perpendicular distance of its point of application from the pivot.

Thus, in a lever of the first class a pull of one pound applied two feet from the pivot will exactly balance a weight of two pounds applied one foot from the pivot. Further, in moving the two-pound weight through a circular arc one foot in length, the one-pound power moves through a two-foot arc, and the actual amount of energy exerted by the power and weight is the same. (See Ex. LVI.)

On this account levers are very useful in mechanical operations where they enable a person to move a large weight a small distance by applying a small power through a longer distance. The actual amount of energy exerted at each end of the lever remains the same.

Levers in the body. — When we come to examine the method of application of the muscles to the jointed bones of our body we find that nature has made use of this law of levers in a very remarkable way. For example, in the forearm we have a lever of the third class. (See Fig. 115, A.) Here the bones form the rigid bar, the elbow is the pivot, and the power is applied by the biceps muscle attached between the hand and elbow. With this arrangement the muscle is able by a short contraction to raise a weight in the hand no little distance. In this case the amount of force exerted by the muscle to raise a given weight can easily be calculated by applying the law of levers.

Furthermore, if we put in the hand a weight which the muscle is just able to raise, the calculation enables

us to get the maximum strength of the muscle. (See Ex. LVI.)

All classes of levers are illustrated in the muscle attachments to the bones in our bodies. (See Fig. 124.) Levers of the first class are least numerous. The action when the toes are pressed downward, and the head lifted are examples of this type. Those of the second class are comparatively few also. The action of the foot when the body is raised on the toes is a case in point. Here the Achilles' tendon, which connects the calf muscle with the heel, exerts the pull; the ankle supports the weight of the body, and the toes are the pivot. Levers of the third class are most abundant. The lifting of the forearm, as already described, the movement of the jaws in eating, the bending of the toes and fingers, and the bending of the knee, are all examples of this class. It is to be noted that while this form of lever requires relatively greater muscle pull to the amount of weight raised, it is of advantage in the body not only in the range of motion it permits, but also because it allows bulky muscle bellies to be placed where they will least interfere with freedom of action.

Standing, walking, and running.—It is a law of physics that a rigid body will remain in a given position of rest as long as a line drawn from its center of gravity toward the center of the earth falls inside of the supporting surface. If such a body is tipped it will return to this position of rest, provided it is not tipped so far that the line falls outside of this basal support. It is this second fact which makes it possible to tip a broad-based body farther than a narrow-based one without danger of upsetting it. Now, in the body, the center of gravity lies in the pelvic region near where the sacrum connects with the last lumbar ver-

tebra, and if the body were rigid it would remain erect as long as the line from the center of gravity fell within the area of the feet. As the feet were spread apart and the base support correspondingly increased, the more difficult it would be to upset the body. If, then, we had a rigid structure the only precaution necessary to secure erect position would be to fulfil this law of physics. But in our body there are many joints, and the tendency of the whole system is to bend and pitch the body forward. To remedy this it is necessary for the muscles that control the movements of the bones to be so fixed as to keep these joints rigid. This is accomplished by opposing the various flexor and extensor muscles so as to hold the bones of the legs and body in an upright position and prevent movement about the joints. To do this requires complete nerve control, and this control is only acquired by much practice as is evidenced by the difficulty which children experience in their first attempts to stand upright. The reeling of the drunken man is another evidence that the erect position is the result of nervous control since, in such case, the alcohol has disturbed the nerve centers only.

FIG. 125—Arrangement of some of the muscles which keep the body erect.

Some one has described walking as the act of falling down and recovering oneself repeated at regular intervals. This is an apt description, for as the flexors bend the leg muscle, the body falls forward upon the other foot as a

pivot, and this forward fall is stopped only by swinging the flexed leg forward and allowing the foot to strike the ground some distance ahead. Then, the body weight is shifted to this foot and the operation is repeated with the other leg. In ordinary walking not only the leg flexors are used but also those of the toes and foot, and the body is usually more or less bent at the waist while the arms swing freely about the shoulder sockets. The feature of this combination of movements which makes it difficult to the child is that these various movements must be so correlated that they take place in regular sequence, and the power to do this comes only with complete nerve control.

Running differs from walking in that the advancing foot alights on the toes alone, and there are instants when neither foot is in contact with the ground. The body is bent more, the pressure of the toes thrusts the body forward with more force and the rate of action of all the muscles is more rapid. Accompanying this greater rapidity of action is a greater demand for oxygen on the part of the muscles, and a consequent increase in the rate of breathing. For this reason running forms a valuable type of exercise for the development of the lung power and as a stimulant to circulation.

Effect of alcohol upon muscular activity.—The exact effect of alcohol upon muscle action is not yet fully determined. Sufficient evidence has been collected, however, to show that the combined action of alcohol upon the muscle cells and upon the nerve centers which control these cells is such as to make it less valuable than other fuel foods for the production of energy and activity. And it has been shown further that while the immediate effect of small doses may result in temporary stimulation, this period

of increased activity is soon followed by a period of exhaustion.

In general, then, the total effect of alcohol upon muscle activity is such as to make the total amount of work performed less with its use than without it, while through its effect upon the nerve centers which control the muscles its continued use tends to destroy the power of precise movement and to lower the user's endurance. In other words, it is a very poor source of energy as compared with other fuel foods.

XX. MOVEMENT IN LOWER ANIMALS.¹

AN examination of a muscle cell demonstrates that the power of contraction and expansion which it possesses is actually a property of the protoplasm of which the cell is composed. We should expect, then, to find in all living matter some power of movement, since all are built up of cells, that is, protoplasm. In other words, motion is an essential characteristic of all living protoplasm.

FIG. 126—A plant cell; the arrows indicate the streaming of protoplasm within the cell.

It does not follow, however, that this motion of protoplasm must necessarily result in the movement of the animal whose cells contain protoplasm, any more than the movement of water in a bowl necessarily results in the motion of the bowl. In fact, in many of the cells of our bodies and in those of plant and animal bodies in general, the protoplasm of the individual cells shows movement, while the cells in which it is inclosed remain stationary. It is necessary, therefore, in speaking of movement to distinguish between motion of protoplasm by itself and locomotion, or the movement of the body or cell from one position to another. The motion

of protoplasm within a stationary cell is very important as a means of distributing material within the cell, but

¹ See Footnote, p. 146, Chapter X.

in our present discussion we are more directly concerned with locomotion.

Causes of locomotion in animals.—The locomotion of all animal forms can usually be explained as due to one of four causes:

(a) A change of position due to the action of some force outside themselves, such as currents of water or air. This sort of movement is, of course, passive movement as far as the animals are themselves concerned, since they take no part in its production.

(b) A change of position due to change in the cell composition, as a result of which there is a gain or loss in specific gravity. Such change allows the buoyant force of the water to move the body up or down. This is not a passive movement, since the changes in composition determine the action of the buoyant force.

(c) A change in position due to active secretion on the part of the cells composing part or all of the animal, and an accumulation of this secretion in such amounts and in such a manner as to force the body forward.

(d) A change in position due to expansions and contractions of part or all of the cell bodies composing the animal, these movements being so arranged as to produce a movement of the whole animal.

The first cause needs little discussion, as it is due to no physiological action of the animals affected. The other three need some explanation to make their action clear.

Locomotion due to changes in specific gravity. — Changes of position of this character are usually confined to water-dwelling forms. Protoplasm is slightly heavier than water, and an animal, therefore, which rests upon the bottom and has no apparatus for raising itself must re-

main there unless it can produce such changes in its composition as to make it lighter than the water it displaces. Certain one-celled animals have the power to accumulate carbon dioxide within the cell bodies to such an extent as to decrease their specific gravity. Thus, the body becomes lighter than the liquid it displaces, and the buoyant force of the water floats it upward. Further, they can discharge this gas, increase their specific gravity, and sink, as they have become heavier than the liquid displaced. In this way many one-celled, water-dwelling animals are able to change their position without the aid or use of special organs of locomotion. In the higher animals,

FIG. 127—Swim bladder of the fish.

the swim bladder of the fish and the connection of the lungs of the bird with cavities of the body, by means of which air can be introduced into the body, are apparently parallel cases, but it must be noted that in these cases the decrease in body specific gravity is due to the action of special muscles and not to changes in the composition of the protoplasm of the cells. The result, however, is the same.

Locomotion due to secretion. — There are certain forms among the one-celled animals and plants that are able to creep over surfaces without contractile movements, or any apparent organs of locomotion. Examination shows that

in certain cases this creeping movement is due to active secretion. In one case which has been especially well studied (the one-celled gregarines), it has been shown that, out of the food supplied to it, the protoplasm of the cell secretes a slimy substance, just as the cells of the salivary glands produce saliva. This slimy substance is forced out of the cell body and accumulates at the rear of the animal. Here it sticks to the surface and hardens, and thus by its accumulation and resistance forces the body along slowly before it.

In the higher forms of animals we find no cases of motion due entirely to secretion. In the snails, however, the body secretes a slimy mucus which forms a surface for the action of the muscular foot and which we see as a shiny trail when it hardens. The common house fly also secretes a sticky fluid on the bases of the foot, by means of which it is enabled to walk over smooth surfaces. The web of the spider over which it travels, and by means of which it raises and lowers itself, is a secretion of certain glands of the body. These cases are evidently not cases of motion due to secretion alone, but simply instances of where secretion aids other methods. Motion due to secretion alone is confined to a few one-celled forms.

Locomotion due to contractility.—Practically all of the free-moving animal forms owe their ability to change position to the power of contractility exercised by the protoplasm of part or all of the cells of the body. We can distinguish three kinds of locomotion due to the contractility of protoplasm:

- (a) Amœboid movements.
- (b) Ciliary movements.
- (c) Muscular movements.

Amœboid movements. — In the study of the amœba it was noted that this animal moves by means of pseudopodia. These pseudopodia are merely extensions of the body protoplasm into which the rest of the protoplasm flowed. In other words, the amœba extends its proto-

FIG. 128 — Various shapes of an amœba in motion.

plasm at a definite point on the surface, and then, by contraction, forces the rest of the protoplasm to flow in that direction. The point where the pseudopods are produced varies with conditions and, since the animal has no cell wall, theoretically these pseudopodia may be produced anywhere. In general, however, they are so formed as to carry the animal toward food or away from irritating causes, and this fact shows that the expansions and contractions are in some way a response to a stimulus.

The result of this peculiar kind of contractility of protoplasm is to produce a slow creeping motion, and all animals which move in this slow creeping fashion with the aid of pseudopodia are said to show amœboid movement. In some of the forms of protozoans which have walls about the body, amœboid movement is still possible by extending the pseudopodia through openings in the walls. The little shelled form known as *arcella* and common in all fresh water is an example of such a type. (See Fig 130.)

In the higher forms we find no cases of amœboid movement on the part of the animals as a whole; but often the cells of a part of the body may exhibit this method of

motion. In our own body, for example, the movement of the white corpuscles is a case in point. To this amœ-

FIG. 129 — White corpuscles escaping from blood tube by amœboid motions.

boid movement these corpuscles owe their power to pass through spaces in the walls of the blood vessels through which the smaller but rigid, red corpuscles cannot pass.

Ciliary movements.—Amœboid movement is at best a very slow method of changing position, and is interesting mainly in showing us that free protoplasm has the power of contractility. We should expect, then, to find in the higher protozoans a more rapid method of progression. Such a modification in motile organs we do find in many of the protozoans, and these organs are called cilia.

In the arcella, movement is produced by pseudopodia which are projected through an opening in the cell wall, and these can be extended and withdrawn as necessity requires.

FIG. 130—An arcella creeping by pseudopod.

In the slipper animalcule or paramœcium (see Fig. 131),

FIG. 131 — Paramœcium; *c*, cilia; *cv*, contractile vacuule; *n*, micro-nucleus; *m*, macro-nucleus; *g*, gullet; *f*, food vacuole; *z*, modified cilia (trichocysts).

which is a higher type of protozoans than the arcella, the body wall is covered with permanent projections of protoplasm called cilia from their resemblance to hairs. These hairlike structures are composed of protoplasm like the pseudopodia, but, unlike them, they cannot be withdrawn, and in contracting and expanding they move at a different rate and in only two directions. As it contracts the cilium is bent quickly and sharply backward; relaxing, it returns much more slowly to an upright position. Many cilia, acting in concert all over the body, push the body through the water just as oars do a boat. A peculiarity of ciliary action, however, is that they do not all contract at once, but the movement of one follows that of the one preceding in definite order, so that in action

they resemble the waving of a field of grain in a wind, each cilium corresponding to a single stalk of the grain. Very rapid movement is made possible by these structures, and their

FIG. 132 — Ciliary movement.

permanent form permits the body to be completely covered with a wall, and at the same time this rigid wall furnishes a resistant structure which the cilia can push through the water much more readily than would be possible if the body were plastic.

Sometimes the cilia are replaced by a single long projection which performs movements of a whip-lash character. Such modifications are called flagella, and are found

FIG. 133 — A flagellated protozoan; *F*, flagellum; *N*, nucleus; *B*, body of cell.

FIG. 134—Colony of flagellated protozoans with two flagella.

in many one-celled plant and animal forms. (See Figs. 133 and 134.)

In many of the higher animals, including man, are found cells provided with similar, ciliary projections, but here the action of the cilia is not to produce movement of the cells themselves. On the contrary, their function here is to produce currents in the liquid that bathes them or to remove obstacles from the surface of the cells. Their action may be compared to the motion of the oars when

the boat is tied. In such case, the result is that the water, and not the boat, is moved. In many of the lower animals the digestive cavities are lined with such ciliated cells, and so the currents of water which they create cause the food, floating in the water, to be brought within reach by the currents set up. In our bodies, the walls

FIG. 135 — Ciliated epithelial cells lining the bronchial tubes; *a*, basal membrane; *b*, spherical epithelial cells; *c*, ciliated cells; *d*, cilia.

that line the bronchial tubes are lined with such ciliated cells, and here they serve to free the surface of the membrane from particles of dust which might produce harm if they were allowed to remain. In fact, it is only in the case of the protozoans and a few of the young stages of the lower animals that cilia are used to move the body of the animal itself. In all other forms their function is the setting up of currents or the removal of particles.

Cilia and flagella are to be considered, then, as merely a modified portion of protoplasm in which the power of rhythmic contraction and expansion has been specially developed. They are chiefly interesting, therefore, as indicating the manner in which protoplasm can become developed into a motile organ.

Muscular movement.—Muscle cells differ from ordinary cells in that they contain certain modifications of protoplasm called fibrillæ. These fibrillæ are thin threadlike fibers embedded in the fluid protoplasm of the cell, and may be variously arranged as to direction, but are always parallel to one another. In certain instances these fibrillæ are divided into segments by cross lines or striæ, and these

give the name of striated muscle fiber to muscle masses made up of such cells. Again, the fibrillæ may be entirely free from such cross lines, and the muscle masses composed of such cells are called non-striated muscle fiber. We have already noted in man that the muscles are distinguished by these differences.

The simplest form of a muscle fibrilla is the unstriated form found in the cell bodies of certain of the protozoans.

FIG. 136 — *Stentor*, a one-celled protozoan in three stages of contraction; the black lines are muscle fibers.

In the *stentor* (see Fig. 136) these fibrillæ are arranged parallel to one another so that by contraction they shorten the whole cell. In other one-celled forms, such as the *vorticella*, the fibrillæ are united into a bundle which extends the length of the stalk. The contraction of this

bundle is sufficient to draw back the whole body of the animal.

These simple forms of fibrillæ give us the key to their character. They are, in fact, merely specially developed portions of protoplasm in which the power of rapid and powerful contraction has been developed to a high rate, leaving the rest of the protoplasm of the cell to perform the ordinary functions of the cell. In the higher forms of animals the cells which contain these fibrillæ are almost completely filled by them, and such cells are set aside in the body to perform the action of contraction and expansion and produce movement, while the work of digestion, etc., is given to other cells. In

general, then, muscle cells are simply cells which have developed to a remarkable extent the modified forms of protoplasm called fibrillæ. It must be evident, then, that whether we have amœboid, ciliary, or muscle movement, the source of the power is the same, namely the contractile power of protoplasm. In this connection it may be noted that the striated fibrillæ have the power of rapid contraction much more highly developed than it is in the non-striated. The enormous extent to which this rapidity of

FIG. 137 — Vorlicella. This protozoan has a single fiber in the stalk, which by contracting draws the animal down as in B.

contraction may be developed is seen in the action of the striated wing muscles of the insects (three hundred to four hundred contractions per second in the fly).

Arrangement of muscle fibers.—The muscles of all animals lower than the vertebrates are parallel fibers, and are never united into bundles like those of man. In no case are they found bound together by a connective tissue, or united to hard parts by tendons.

All the muscles of the starfish, worm, and oyster family are of the unstriated variety. All those of the lobster, spider, and insect tribes (Arthropoda) are striated.

Organs of locomotion.—In many of the worms and lower forms, and even in some of the higher forms, such as the snake, motion is secured by undulations of the body as a whole. Most animals, however, have special organs of locomotion. The number of these organs varies, but, in general, the higher we go in the animal scale the greater the perfection and the fewer the number. Thus, while a starfish has hundreds of tube feet, and a centipede many legs, the two arms and legs of man are much more effective as means of progress. While it is impossible to consider in detail all the organs of locomotion in the animal world, we may consider for a moment the characteristics common to those adapted for special surroundings. All the organs of locomotion about to be described depend upon the action of variously arranged muscle layers, that is their movements are muscular movements.

Organs of locomotion for water life.—Since the water supports nearly all the weight of the body, we should expect to find the organs of locomotion of aquatic animals ill adapted for supporting the body. For example, the cilia of the protozoan, or the fins of the fish, are not intended to support the body, but to push it through the water; while even the webbed feet of the duck are placed

so far back on the body as to make walking awkward, though admirably placed for swimming.

All organs made for movement in the water are well

FIG. 138 — The fins of a fish.

adapted to pushing the body through the water. The broad tail of the lobster and the flexible abdomen enable it to shoot the body backward by bending this portion sharply forward, and then slowly straightening it. The fins of the fish and the flukes of the whale show the same paddlelike character, while the webbed feet of the frogs and aquatic birds are adaptations for pushing against water. In all these paddlelike organs, too, the broad surface is so constructed as to be opposed to the water in pushing, while by bending, it is enabled to return to position without offering much resistance to the water.

An interesting modification of the water motion is seen in the squid where the broad tail serves simply to steer with, while the motion is produced by ejecting spurts of water by contraction

FIG. 139 — Movement of the tail fin of a fish in swimming.

of the body walls, and forcing this ejected water through a funnel at the front of the body.

Organs of locomotion for a flying life.—All animals which fly require special adaptations of structure to lighten the body, and also very powerful muscles to keep the body in motion, while the form of paddle or wing used to push the body through the air must be much broader in proportion to the size than were those used in pushing against the water.

The bodies of flying insects are lightened by air sacs and cavities distributed through the body, while in birds similar air sacs and hollow light bones serve them for the same purpose. Feathers in the bird are adaptations for a body covering which secure lightness and protection at the same time.

The wings of insects are expansions of the skin or body

FIG. 140—Cross section of an insect; *h*, heart; *a*, alimentary canal; *n*, nerve ganglia; *t*, air tubes (tracheæ) which carry air to all parts of the body and form the veins of the wings.

covering, stretched upon a frame of hollow tubes of the same material. They are moved by muscles lying in the thorax, and these muscles are remarkable for their rapid rate of contraction. The number of wings in the insects varies from two to four, and they also vary in size from the tiny wings of the gnats to the relatively enormous surfaces of the butterfly wings. In some forms, such as the beetles, the front pair is adapted solely for a protective covering, while the flying is done with the second pair. The dragon fly is the most perfect flying machine known, being able to fly forward, backward, or remain poised in the air at will, while its movements are marked with great speed and accuracy.

In birds the arms are modified into wings by a special covering of feathers. These feathers are so arranged as to allow the air to pass through them as the wing is raised and to prevent such passage on the down stroke. This

FIG. 141 — Wings of the bat, formed by expansions of the skin over the lengthened finger bones.

arrangement enables them to strike the air very strongly at each down stroke, and thus to force the body upward and forward. The muscles which operate these wings are

powerful, and the breastbone is keeled to permit their attachment to it. The speed of some birds in flying is extremely great, a hawk being able to fly at the rate of one hundred and fifty miles an hour.

The only mammal that flies is the bat, and here the wings are formed by expansions of skin stretched over the arms and legs as frames, while the fingers are enormously developed for this framework. (See Fig. 141.)

Organs of locomotion for land and water-bottom life.— We find many curious adaptations in the animals which move on land or over the bottom of the sea or ponds.

The earthworm progresses by first contracting the circular muscles and thus extending the body. Then, as the longitudinal muscles are contracted, the bristles on the under side of the body are inserted into the surface over which it is creeping; as a result, the shortening of the body produced by the muscle contraction causes the body to be dragged forward.

The leeches use the same muscles, but fasten the ends of the body by means of suckers, so that their motion is a progression of loops.

Starfishes use their tube feet in a similar manner, first by lengthening the foot and attaching it by the sucking disk at the end, and then dragging the whole body forward by contraction.

In the mollusks, motion is produced by a creeping disk or foot which in the clam ploughs through the sand and forces the sharp-edged shell forward, while in the snail the muscles contract and expand in such a way as to wrinkle the bottom of the foot and thus push the body forward.

Many land animals and sea-bottom animals have jointed legs of some kind, operated by powerful muscles, usually

arranged in pairs, as in our own bodies. In the crustaceans these legs may be as many as twenty, some being moved backward as the others move forward. The motion of the six legs of the insect is illustrated in Figure 142.

FIG. 142—Motion of the legs of a fly in crawling.

Most land vertebrates, except the snakes, have four legs, the two front ones corresponding to our arms, and the two hind ones to our legs. In some forms the legs are so placed that the animal stands upon the tips of the toes as in the horse, cat, dog, etc. In others, as in man and the bears, the whole foot is on the ground. Nearly all

FIG. 143—Feet of mammals adapted to different methods of locomotion; A, foot of bear; B, foot of seal; C, foot of lion.

vertebrates have more powerful leg muscles than man, and can move much more rapidly. The legless snakes

move by writhing and by pushing the edges of the ventral scales against rough surfaces.

In all running vertebrates the hind legs are much more muscular than the fore legs or arms, and are the principal organs of locomotion; the front legs being mainly for the support of the front part of the body.

XXI. RESPIRATION.

THE activity of a muscle cell was seen to be dependent upon a constant supply of food and oxygen. What is true of a muscle cell is true of every cell in the body. All require food and oxygen, and all give off carbon dioxide and wastes. In our study of the digestive and circulatory

FIG. 144—Position of the lungs in the thorax; *A*, lung; *B*, heart; *D*, pulmonary artery; *E*, trachea.

systems we saw the manner in which food and oxygen reached the cells of the body and how the wastes were un-

loaded from them. Finally, we traced the source of the oxygen supply to the lungs of the body, and the wastes to the kidneys. The organs which are concerned in the supply of oxygen to the cells and in the removal of the carbon dioxide from the body are spoken of collectively as the *respiratory* or *breathing system*.

Respiratory organs.—The term respiration (*re*, again, and *spiro*, I breathe) involves two distinct actions. The taking of air into the body is called inspiration, or inhaling. The throwing out of the body of carbon dioxide and the air which has been freed of oxygen, is called expiration, or exhaling. These two processes of respiration are performed by various kinds of organs in the lower animals, but in man and in all the other vertebrates, except the fishes, the principal organs of respiration are the lungs. Connecting with the lungs, which are located in the thoracic cavity, are several passages and tubes which bear a part in the process and are to be included in our term respiratory organs. The relation of these parts to the lungs can be best understood

FIG. 145.—Relation of the buccal and nasal cavities to the esophagus; *n*, brain cavity; *l*, hard palate; *f*, uvula; *g*, opening of Eustachian tube to ear; *e*, epiglottis; *k*, tongue; *c*, windpipe; *b*, esophagus.

by following the course of the air as it passes through them.

Air enters the body by way of the nostrils or mouth and passes backward into the pharynx or throat cavity. Here, two passages are open to it: one, the gullet, leads to the stomach; the other, the trachea, or windpipe, leads to the lungs. As we inhale, the walls of the esophagus contract, and the air is sucked down the windpipe and enters the lungs. Just how this suction is produced we shall investigate later. For the present let us examine these passages a little more in detail. The figure (see Fig. 145) shows the arrangement of these parts with relation to one another. If we examine it carefully we note that the two external nostrils are merely openings into a large cavity (the nasal cavity) located just above the roof of the mouth and connecting at the back with the pharynx. The nostril entrance to this cavity is guarded by many hairs which serve to catch and remove dust particles from the air. The cavity itself is lined with a layer of mucous membrane, the secretion from which aids in catching particles of dust and germs that have escaped the hairs. This membrane is also lined with many blood vessels, and the heat which escapes from them serves to warm the air before it comes in contact with the delicate membranes of the lungs. These facts show clearly that the nose is adapted for the entrance of air to the body, and that while mouth breathing permits the air to reach the lungs, it has none of the safeguards against impurities and cold that are found in the nasal passage. Mouth breathing is a habit to be avoided.

The location of the top of the windpipe may be easily found by applying the fingers to the outside of the throat

where a hard swelling may be felt, called the Adam's apple. This is the *larynx*, or voice box, and is on the top of the windpipe. The *trachea* itself may be felt below this projection and recognised by the rings of cartilage which encircle it and prevent its collapse as the air is sucked in. The entrance to the larynx and trachea is guarded by a lid called the *epiglottis* which is always open to admit

FIG. 146—Diagram of trachea and connections.

air, except in the act of swallowing, when it closes to protect the tube from entrance of food.

At its lower end the trachea splits into two branch tubes, or *bronchi*, which, in turn, split into smaller and

smaller branches called *bronchial tubes*, and finally end in thin-walled sacs called *pulmonary sacs*. The bronchial tubes and the sacs are all inclosed in two baglike bodies covered with a membrane, and located in the chest cavity one on each side of the heart. These two bodies, the framework of which consists of the branches of the trachea, are what we designate by the name of lungs, and the pulmonary sacs of these lungs are the ultimate destination of the inhaled air. The structure of the windpipe and lungs of man are practically identical with those of all other mammals, and the dissection of the chest cavity of such a mammal (rat, cat, or rabbit) gives one a very clear idea as to the structure of our own organs, and their position with relation to the other organs of the body.

Position and structure of the lungs.— (See Ex. LVIII.) If we remove the skin from the chest and throat of a mammal the windpipe and the ribs will be laid bare. Between the ribs and connecting them are layers of muscle (the intercostals), and on moving the point of the breastbone up and down the layers will be seen to give a peculiar movement to the ribs. We can produce this same movement by inserting a glass tube in the entrance of the trachea and inflating the lungs with the breath. As we do this we shall also note a downward movement of a membrane which separates the chest cavity from the abdominal cavity. This membrane is the diaphragm, and, together with the ribs, it performs an important action in breathing movements. The relation of these boundary walls of the chest cavity are shown in Figure 144.

Cutting the ribs on one side close to the breastbone, and pressing them back, the lungs and heart which fill the cavity are seen. The lungs appear as two pinkish bags,

one on each side of the heart, and are held together at the top by the two bronchi or main branches of the trachea.

If we inflate these two bags with the aid of the glass tube and then tie the trachea tightly, the lungs may be separated from the cavity and floated upon water. This

FIG. 147—Ribs and muscles; *a*, ribs; *b*, cartilages; *c*, junction of bone and cartilage; *d*, bony portions, and *e*, cartilaginous portions of breastbone; *A*, external intercostal muscle; *B*, internal intercostal muscle. In the middle the external muscle has been removed.

power to float has earned them the name of “lights” from the butcher. The dissection of these separated lungs shows the following structural features.

Structure of the trachea.—The trachea is a stiff, cylindrical tube covered on the outside with a layer of fibrous, connective tissue in which are embedded the stiffening rings of cartilage. The interior of the tube is lined with a single layer of mucous membrane, the innermost cells of

which are provided with cilia. (See Fig. 135.) These cilia, together with the mucus secreted by the cells, serve to catch any particles of dust that may have passed the nasal cavity. The cilia are in constant motion, bending quickly upward and then slowly returning to place, and in this manner forcing the collected particles steadily upward toward the pharynx.

The cartilage rings do not completely inclose the tube, but are shaped like the letter C, the open part being at the back and leaving a portion of the tube soft and compressible. The gullet lies against this soft groove, and thus room for expansion is given as the food passes down the gullet.

Structure of the lungs.—These are merely membranous sacs with an internal framework composed of the branching bronchial tubes and air sacs. These treelike tubes are bound together with a soft, pink connective tissue, rich in elastic fibers, and over the whole is stretched an elastic membrane known as the pleura. The bronchi and bronchial tubes are continuations of the trachea, and the larger ones resemble it in structure with the exception that the cartilage rings are not so regularly arranged. As the tubes become smaller their walls become thinner, and finally the cartilage rings disappear entirely in the smaller tubes, while the mucous membrane

FIG. 148—Two air sacs (*b*) into which a bronchial tube (*a*) opens.

is reduced to a single layer of ciliated cells. The air sacs, in which the smallest tubes end, are composed of

a thin, elastic outer coat of connective tissue and a lining of flat, mucous-membrane cells without cilia. Between these two layers is a network of tiny, thin-walled capillaries whose blood is thus separated from the air in the sac by only the thin capillary wall and the layer of flat cells.

The whole structure is very elastic, and readily expands and contracts within the limits of the chest cavity as air enters and leaves it.

The pleura.—The pleura covers both the right and left lung. This pleura is a serous membrane, and is a complete cover for the lungs except at the point where the bronchi enter. Here, instead of fusing with the walls of the bronchi, it is turned back and lines the interior of the chest cavity. Between the pleural cover of the lung and the pleural lining of the chest cavity is thus formed a cavity which is so small as to require only a few drops of liquid to fill it. This cavity is filled during life with a liquid of the nature of lymph, secreted by the pleura, and as the lungs are inflated in breathing, this liquid serves to lubricate the pleural surfaces where they rub against one another and to prevent friction.

Breathing.—Two processes are involved in breathing, inhaling and exhaling. Inhaling consists of drawing the air from the outside through the nasal passages, trachea, bronchi, and finally distributing it through the bronchial tubes until it reaches every part of the lungs and fills the air sacs. Exhaling reverses this process and expels the air in the air sacs through the above passages in the reverse order to the outside. As the walls of the chest cavity enlarge in inhaling, the lungs fill with air; while in exhaling, the walls contract, the lungs are compressed,

and the air is forced out of them by the passages above noted. What causes the lungs to expand and fill with air, and, since the lungs themselves contain no muscles, what parts of the body are concerned in this process? The answer to this question is to be found in the structure and movements of the walls of the chest cavity.

Structure of the chest cavity.—The chest cavity is conical in shape, with the point of the cone at the top, and is completely inclosed on all sides. The upper part and sides of the cone are formed of the ribs, intercostal muscles, sternum, and backbone, all covered over with muscles and skin, while the floor of the cavity is composed of the membrane called the diaphragm. These walls are not rigid, but permit of movement in two definite directions. As we inhale, the sides of the cavity can be felt to move outward, while the diaphragm is pulled downward, forcing the contents of the abdominal cavity in the same direction, and causing the belly to swell under the pressure. As we exhale, the ribs move inward and the diaphragm moves upward, causing the belly to be drawn in. The causes of these movements are to be found by studying the shape of the ribs and the muscular connections.

In the dissection of the chest cavity, the muscles between

FIG. 149 — Diagram of rib movement in breathing; *AB*, vertebral column; *CD* and *EF*, ribs attached to movable breastbone, *DF*; *a*, *b*, external and internal intercostal muscles; *c*, neck muscles.

the ribs were noted under the name of intercostals. From the study of the skeleton it was noted that the ribs were jointed to the backbone at one end and to the breastbone, by cartilages, at the other; while the position of the ribs was such that the sternal ends were lower than the dorsal ends. Finally, muscles attach the upper ribs to the neck vertebræ. If, now, these latter muscles and the intercostals contract simultaneously, the effect is to lift the breastbone and the sternal ends of the ribs upward. To do this is to widen the chest cavity from front to back and from side to side. The action of these muscles is rhythmic and automatic. They account for the movement of the ribs, but not of the diaphragm.

The diaphragm is a membrane which forms a domelike roof for the abdominal cavity, with the top of the dome extending normally into the chest cavity. It is composed of thin, muscular tissue, and a tendon is attached at the apex of the dome. From this tendon, muscle fibers radiate downward and outward, and are fastened to the lower ribs, to the breastbone, and to the backbone at their outer ends. When

these fibers contract the result is that the dome of the membrane is pulled downward out of the chest cavity and compresses the contents of the abdominal cavity, thus increasing the depth of the chest cavity from top to bottom. This contraction of the diaphragm muscles and those which pull the ribs upward

FIG. 150—Position of the diaphragm (A) in breathing.

takes place simultaneously, as does the expansion. The combination of these two movements is responsible for the filling and emptying of the lungs with air. Now, it is easy to see how the return to position of the ribs and diaphragm might squeeze the lungs, and cause them to expel the air; but, as the walls of the chest cavity are not fastened to the walls of the lungs directly, it requires a more detailed study to understand how this outward movement causes the lungs to expand and become filled with air. How does the enlargement of the chest cavity cause the lungs to fill with air?

FIG. 151 — Apparatus to illustrate breathing movements and their effect upon the lungs. The rubber diaphragm corresponds to the diaphragm in the body; the handle to the tendon; the balloon to the lungs; the tube to the trachea; the bell jar to the walls of the thorax. As the handle is lowered the air flows down the tube and inflates the balloon.

Causes of inhaling and exhaling.—A simple mechanical device will make clear the relation of the chest enlargement to the inflation of the lungs. (See Ex. LIX.) An ordinary bell jar, open at the top and bottom, will represent the chest cavity. A piece of sheet rubber stretched across the bottom will represent the diaphragm. Finally, a toy rubber balloon tied to a glass tube will be the lung, and the tube the trachea.

To make the comparison complete we shall put the balloon into the bell jar with the tube fitted into the stopper. Our apparatus will now look like Figure 151. If we have made our connections

well, the jar is now air-tight, and the only connection of the interior of the balloon with the outer air is through the glass tube. This apparatus represents the relation of the lungs to the chest cavity, since the sides of the balloon are not in any way connected with the rubber sheet (diaphragm) or with the sides of the bell jar (the rib walls). Now pull the rubber sheet downward. The balloon becomes inflated as air enters through the tube (trachea). Let the sheet return to place, and the balloon collapses, expelling the air through the glass tube. What is the explanation? The answer is found in what the physicists call the laws of air pressure. They tell us that every square inch of surface exposed to the air is under a pressure of fifteen pounds. At the start of our experiment, therefore, both the inside and outside of the balloon are under equal pressure. Since no air can enter the bell jar, when we lower the diaphragm the air inside it must occupy a greater space, and its pressure on each square inch of the balloon surface becomes less. The only way in which the pressure can be equalized is for air to rush in from the outside and distend the balloon until the volume of air in the jar is the same as before. Then the inflow stops, since both surfaces of the balloon are again at the same pressure. When the diaphragm returns to position the air in the jar is compressed, and the surface of the balloon exposed to this compressed air is now under a greater pressure than the outside air can exert upon the inside through the glass tube, and the walls are gradually forced together with the outpouring of the contents of the balloon, until again the pressure on the outside of the balloon is the same as on the inside. Meanwhile, the balloon has performed all the movements of the lungs in

inhaling and exhaling, and this process may be repeated as often as we lower and raise the rubber diaphragm.

In the chest cavity the conditions are practically identical with that of our apparatus, but the lungs are much more sensitive to changes in pressure than are the balloon walls. One reason for this is that the chest cavity volume is increased not only by the downward movement of the diaphragm but also by the upward movements of the ribs. Furthermore, the lungs fit closely against the walls of the cavity, and thus a slight increase in volume is immediately felt by all parts of the lungs, with a correspondingly greater expansion.

The movements of contraction and expansion, on the part of the chest walls, take place rhythmically about fifteen to eighteen times per minute, which is the normal rate of breathing. In violent exercise this rate may increase to sixty or seventy times per minute. The rate also varies with the age, that for a baby being about forty, while this rate gradually decreases, until, at eighteen years of age, it is practically the same as for an adult. Much more muscular effort is exerted in inhaling than in exhaling, there being practically none in normal exhaling.

PHYSIOLOGICAL EFFECTS OF BREATHING.

Composition of inspired and expired air.—The study of ordinary air (see p. 20) shows it to be composed mainly of two gases, oxygen and nitrogen. Careful chemical analysis shows the presence of small quantities of other chemical elements and compounds, such as argon, krypton, carbon dioxide, and water. Of these, only the water and carbon dioxide have any physiological significance. Ex-

pressed in volume per cent the following table represents the composition of ordinary air:

*Inspired Air.*¹

Oxygen	20.96%
Nitrogen ¹	79.00%
Carbon dioxide	00.04%

If we examine the air which is expelled from the lungs we find certain remarkable changes in it. If we breathe upon a thermometer bulb or a mirror surface, two properties peculiar to expired air are apparent. The first is the increase in water content as shown by the film of vapor that collects upon the bulb. The second is the rise of temperature as indicated by the rise of the mercury column of the thermometer. A comparison with the body temperature shows that the expired air has been raised to this body temperature.

Evidently then, breathing involves a loss of water and heat to the body, and thus plays an important part in the regulation of the temperature and water content of the body.

If we blow expired air through a tube into a tumbler of limewater, the latter soon becomes milky, indicating that a large proportion of carbon dioxide is present. (See p. 16.) If we collect some of the expired air over water and introduce a lighted match into it, the flame soon goes out, indicating the loss of a large amount of its oxygen, and hence its power to support combustion. Expressed chemically,

¹ The very small per cent of argon and krypton is included in the nitrogen estimate. The percentage of water is so small as to be neglected in the above table.

the composition by volume of expired air is shown in the following table:

Expired Air.

Oxygen	16.02%
Nitrogen	79.00%
Carbon dioxide	4.38%
Water	0.60%

A comparison of this table with that for inspired air shows that in the lungs the air loses some 4.94% of oxygen, and takes on some 4.34% of carbon dioxide and 0.6% of water. The proportion of nitrogen remains unchanged. Expressed in terms of a day's breathing, the body obtains for the tissues, in a day, about two pounds of oxygen, and loses in that time some two pounds of carbon dioxide and a half a pound of water.

Expired air also contains traces of ill-smelling, organic compounds. These are what give the bad smell to a poorly ventilated room that has been occupied for some time by people. They render the air dangerous and poisonous to breathe. A bad digestion also adds to the ill-smelling compounds which, emerging from the gullet, mix with the expired air and make a fetid breath, while decaying teeth add still other odoriferous compounds.

Blood changes in the lungs.—The tables above indicate only the changes which take place in the inspired and expired air. They do not explain their purpose. To do this we must recall the structure of the lungs. It will be remembered that the inspired air ultimately reaches the air sacs and that these sacs are lined with thin walled capillaries. The blood, which has been circulating through the body tissues, comes to these capilla-

ries from the heart (pulmonary arteries). The corpuscles of this blood are laden with the carbon dioxide which they have absorbed from the tissues. When these corpuscles reach the lung capillaries their carbon dioxide passes, by diffusion, through the walls of capillary and air sac into the cavities of the sacs. At the same time, the hæmoglobin in the corpuscles absorbs the oxygen of the inspired air in the sacs. Similarly, by dialysis, the organic compounds and water are given off to the sacs from the blood plasma. In this way the blood in the capillaries is enabled to renew its supply of oxygen, rid itself of its burden of carbon dioxide and excess of water, and finally to return to the heart (pulmonary veins). From there it is sent out to the tissues again. This interchange, therefore, accounts for the manner in which the oxygen gets into blood, and carbon dioxide out of it. It is to be noted in this action that, except for the tiny amounts of organic compounds given off, no true wastes pass from the blood to the air in the air sacs. On this account, breathing should not be spoken of as the *purifying* of the blood. What actually happens is the aerating of the blood, that is, an exchange of carbon

FIG. 152.—Diagram of two air cells, showing the capillary network which covers them, and at *a* the structures which intervene between the air and the blood are indicated; 1, mucous membrane of the air cell; 2, submucous meshwork; 3, wall of capillary; 4, plasma in capillary; 5, red blood corpuscle.

dioxide for oxygen. The purifying or removal of wastes from the blood is performed by an entirely different set of organs, which we shall consider more in detail in a following chapter.

The whole respiratory system, then, is simply a device for getting the blood into thin walled tubes and these tubes in contact with air, in order that the process of osmosis may bring about the exchange of the two gases, oxygen and carbon dioxide.

BREATHING HABITS.

Costal and abdominal breathing.—In man, in quiet, normal breathing, the operation is mainly confined to diaphragm movement, the ribs expanding very little. In woman, on the contrary, quiet breathing is carried on mainly by the rib movements. These two kinds of breathing are spoken of usually as abdominal (diaphragm) and costal (rib) breathing. It is not believed that the two kinds are to be associated primarily with the sex, but, on the contrary, it is believed that the costal method of the woman has resulted from the tightness of her clothing as compared with that of a man. The use of corsets and tight fitting dress waists tends to restrict the expansion of the abdomen, and forces the use of the upper ribs for the purpose. If such tight dressing is carried to excess, as in tight lacing, not only is abdominal breathing prevented, but even rib action is interfered with, and thus the entire breathing action or chest cavity enlargement may be interfered with. Such interference results in insufficient air supply to the lungs and body tissues, and the distorting of the skeleton, and subjects the abdominal organs to such

great pressure that disease of these parts may result. Such tight clothing is evidently a menace to the health of the body and is to be deplored.

It is especially important that the growing child should have free play of the chest cavity, and in exercising the increased demands of the muscles for oxygen can be met only by allowing free play to the lungs and walls of the chest cavity.

Lung capacity.— The amount of air respired in a breathing depends, of course, upon the extent of chest enlargement, and the size of the individual. In ordinary breathing, the amount of air respired by all individuals represents only a small proportion of the total capacity of the lungs. Even in deep expirations, the lungs are never entirely emptied of air. The figures for an average adult are as follows:

FIG. 153 — Lung capacity.

After the fullest possible inspiration the lungs contain about three hundred and thirty cubic inches of air. After the fullest possible expiration following such an inspiration they still contain one hundred cubic inches of air.

The difference, two hundred and thirty cubic inches of air, represents the greatest possible amount of air that can enter and leave the lungs in a single respiration. This is called the *vital capacity* of the lungs. In quiet breathing

only about thirty cubic inches is inhaled and exhaled in a single respiration. This is called *tidal air*. The one hundred cubic inches left in the lungs after the fullest possible expiration is called *residual air*.

The significance of the small amount of tidal air and the presence of a certain constant amount of residual air is found in the fact that such a condition allows the blood exchange to take place continually instead of at intervals as would be necessary if the lungs were entirely emptied at each expiration. Further, the possibility of varying the amount of air content permits the lungs to accommodate the air supply to the demand of the tissues, that is, when exercising the increased amount of oxygen needed is met by deeper inspiration and expiration.

HYGIENE OF BREATHING.

We have already noted that in order that the air be properly warmed and cleared of dust before entering the lungs it should pass through the nasal cavities rather than through the mouth. We have also seen that the action of the lungs is dependent upon the enlargement of the chest cavity, and hence tight clothing or other interference with such enlargement is necessarily injurious.

Like muscles, too, the capacity of the lungs and their size are increased by use. Exercise increases both the blood rate and the breathing rate. Now, in quick, short breathing, only the air in the upper part of the lungs is changed, while the increased rate of blood flow shoots the corpuscles through the capillaries so rapidly that they do not have time to effect a satisfactory exchange of oxygen and carbon dioxide unless there is a large and constant

supply of fresh air to draw upon. Deep breathing exercises soon accustom the lungs to fuller expansions and contractions under these conditions, and thus bring the air to greater areas of capillaries. Runners, whose lungs have been so trained, can run much longer distances and use their muscles to greater advantage than those who have neglected such training. "Second wind" is simply the automatic response of the lungs by increase of rhythmical expansions and contractions. Again, since the breathing depends upon the action of the chest muscles and those of the diaphragm, exercise tends to develop these muscles and thus directly develop lung capacity. People who have well developed lungs and muscles are said to be long-winded, and are usually more healthy than short-winded ones.

Modifications of breathing.—When any obstacle such as dust or food enters the trachea, the abdominal muscles contract forcibly and compress the lungs. The glottis opens suddenly and the air forcing out the matter gives an effect called coughing. Sometimes inflammation of the lining of the bronchial tubes may cause this action in bronchitis. In like manner irritation of the nasal membranes causes the air to be forcibly expelled through the nose. Such an operation is called sneezing. Blowing is a long, forcible expiration through a small opening in the lips or through the nose. If the object which causes coughing is of sufficient size to stop the trachea, as when food slips by the epiglottis, the expirations increase in frequency and force, and we call the result choking. In this last, the supply of air to the lungs may be completely cut off if the obstacle is not soon removed, and strangulation, or death from lack of oxygen, result. When the air supply is cut off by inclosure in a room without ventilation, or by immersion in a gas or water, the result is spoken of as suffocation. Both strangulation and suffocation are the results of oxygen starvation.

Laughing, crying, and sobbing are merely rapid successions of

short expirations followed by a long inspiration. Hiccoughs are caused by a sudden contraction of the diaphragm. Yawning is a long inspiration and expiration through the open mouth while the throat muscles are held rigid. Sighing is similar, except that the inspiration is through the nose, and the muscles are suddenly relaxed after the inspiration, thus making a sound. Snoring is produced by the inspired air passing over the soft palate and setting up a vibration. Spitting is a use of expired air to force matter from the mouth. It may be done by the action of the mouth muscles alone.

In general, then, all these modifications of respiratory movements are essentially reflex, that is, they take place without any act or will on our part, and often in spite of our desire to the contrary. Spitting is an exception to this rule, though that may become a habit. Most of them, with the exception of hiccoughs, are under our control to a certain extent.

Asphyxiation and artificial respiration. — Asphyxiation is death from suffocation or the lack of oxygen supply. It may be brought about in various ways. Blocking the tracheal entrance by pressure applied to the outside of the throat, as in strangulation, is one method. Exposure in a medium containing no oxygen is another. Exposure to gases which have a greater affinity for the hæmoglobin of the blood than oxygen, and hence turn it out of the corpuscles, produces the same result. Carbon monoxide or the gas produced in burning charcoal is an example of such a cause. In general, any cause that deprives the lungs of their normal supply of oxygen will eventually produce asphyxiation.

When the tissues find their oxygen supply reduced in any of the above ways, the first visible symptom of this lack is seen in the deeper and more rapid breathing. Then the breathing loses its regularity, and soon convulsions follow. Gradually these convulsions spread until other muscles beside those concerned in breathing are affected, and the whole body is thrown into violent spasms. Finally the muscles become exhausted, the movements decrease and finally cease altogether, consciousness goes and the animal is soon still in death unless remedies of a powerful character are soon applied. What these remedies shall be is largely determined by the cause of the asphyxiation, and resource should

be had at once to a physician. In one method of asphyxiation, however, namely drowning, it often becomes necessary to apply remedies before the attendance of the physician can be secured. Owing to this fact, and because the methods illustrate the typical process, the next paragraph is devoted to consideration of such remedies.

Artificial respiration.— When a person has become unconscious from immersion in water, or is in a partly drowned state, it is often possible to imitate respiratory movements, and thus restore normal breathing and life. All such methods depend upon what is called artificial respiration. One of the simplest methods of the artificial breathing remedies is as follows: Remove all the clothing from the upper part of the body and turn the patient upon his face, place a roll of cloth under the pit of the stomach and then press down hard upon the back in order to force the contents of the stomach out of the mouth. Now turn the body on its back and set up artificial breathing movements by first grasping the arms above the elbows and raising them above the head. Next lower them gently to the sides, at the same time pressing the chest to expel the air and water in the lungs. Repeat these movements regularly about fifteen times to the minute. See that the tongue is well drawn out while this is going on. The first symptoms of life will show in a faint, pink color under the finger nails and in the lips. Continue these artificial breathing movements until the body takes them up automatically and regularly.

The action involved in these operations depends upon the fact that in raising and lowering the arms the chest walls are made to move in the same manner as in normal breathing, and gradually the muscles which control these movements take up their functions again and the patient is restored. The pulling forward of the tongue opens the larynx and stimulates the nerves which control the respiratory movements.

The above method is only one of many for the restoration of partially asphyxiated victims. All methods depend upon the forcing of air in and out of the lungs until the exhausted muscles are restored to their normal activity.

Ventilation.— Since the lungs are constantly exchanging large quantities of carbon dioxide laden air for oxygen, it is evidently

necessary that the air in the rooms in which we live be renewed frequently. If this air is not constantly renewed the lungs are forced to inhale air which has lost part of its oxygen, and the preliminary stages of asphyxiation are soon reached. A feeling of dullness, labored breathing, and headache are the warnings which tell us of the need of change of air in a close room, and these warnings should be heeded. All schoolrooms and public halls are now provided with systems of ventilation, that is, apparatus forcing in new air and removing the old. Houses not provided with such systems should have the windows opened frequently, and the air should be allowed to circulate freely until the renewal is secured. During sleep the demand for oxygen is even greater than in the waking hours, since that is the time for the restoration of exhausted tissue. No one should sleep in a room whose windows are tightly closed and the means of ventilation thus cut off.

The basis of all ventilation methods is found in the fact that cool air is heavier than warm air. For example, the air in a room is usually much cooler at the floor than near the ceiling. If, now we provide an exit for the warmer air at the top of the room, and an entrance for the outside air, the outside air will rush in and the warm air will be forced out. Thus the supply for the room will be renewed. Precaution against draughts is important, since the sudden chilling of any part of the body may produce colds. The greater the difference in temperature between the outside and inside air the more rapid the flow, hence in winter much smaller openings are necessary for ventilation than in summer.

All furnace-heated houses should be provided with cold air boxes, so that the air in the pipes may receive constant renewal of oxygen as it circulates.

Air in a room cannot be changed oftener than three times in an hour without producing dangerous draughts. On that account, schoolrooms where fans are used to force air into the room should have this rate carefully regulated, and also should not be so crowded as to make a more frequent change necessary.

Dusting.—We have seen that the approaches to the lungs are well provided with apparatus for removing dust from the inhaled air. Such dust irritates the delicate membranes of the lungs if allowed to come in contact with them and may also contain disease

germs. On this account sweeping and dusting should be so performed as to remove dust from the room and not to fill the air with it. Sweeping raises into the air the dust from the carpets and furniture. Before the room is used again plenty of time should be allowed for this dust to settle and the air to clear. Then the dust should be removed from the furniture on which it has collected, with a damp cloth, and not with a brush or feather duster, as these latter do not remove it but simply throw it into the air again. Opening windows while sweeping permits currents of air to remove most of the dust particles, and at the same time renews the oxygen.

Stone cutters and others who work in dust-laden rooms often suffer from serious diseases as the result of inhaling these particles.

DISEASE OF THE RESPIRATORY ORGANS.

Adenoids.—This is the name given to certain grapelike swellings or growths which develop in the upper part of the pharynx just behind the nasal opening. They often form in children, and if allowed to grow soon close the opening to the nose, and thus make mouth breathing necessary. They may even develop to such an extent as to close the Eustachian tube leading from the pharynx to the ear, and thus produce deafness. While they are not in themselves dangerous, and often shrink and disappear as the child grows older, they may indirectly cause much harm as well as annoyance to the growing child. In children, the bones are in process of formation, and mouth-breathing often produces an unsightly deformation of the jaws. Adenoids also render the child liable to colds, since air which enters by the mouth is not properly warmed, and thus may chill the delicate membranes, and so produce sore throats, tonsilitis, etc. They are growths which may be easily removed, and children suffering from them should receive proper treatment for their removal at once.

Inflammatory diseases.—Sudden changes of temperature affect the vaso-motor nerves, and thus the blood supply to certain parts of the body. When the skin is suddenly chilled in any way the blood vessels of the mucous membrane, which lines the air passages, become charged with blood. If under such conditions these membranes become infected with bacteria, inflammation results. When this inflammation is limited to the nasal cavity we call the result a cold in the head. When the pharynx membrane is inflamed a sore throat is the result. A chest cold is due to inflammation of the windpipe and bronchial linings. If this inflammation spreads to the linings of the bronchial tubes, we call it bronchitis, while infection of the air sacs themselves produces pneumonia. The reactions involved in these various types of inflammatory diseases of the mucous membrane have already been discussed in the paragraphs on inflammation. (See p. 206). All forms are accompanied by great secretion of mucus, which tends to clog the air passages and render breathing difficult.

In certain cases the pleura which lines the thoracic cavity becomes inflamed, the two surfaces rub together, producing severe pains, while the secretion of pleural liquid increases to an abnormal amount and produces pressure upon the lungs. Such a disease is called pleurisy, and can often be relieved only by tapping the pleural cavity and extracting the excess of fluid collected there.

The treatment for inflammatory diseases depends upon the reducing of inflammation and the restoration of the blood supply and secretion to their natural amount. In severe cases a physician should always be consulted.

Other germ diseases.—Inhaled air often contains disease germs of various kinds. If these germs lodge in the mem-

branes of the air passages, they may grow and produce colonies of bacteria. These colonies in their growth may break up the surrounding tissues or may produce poisons or toxins which are absorbed by the blood and carried to parts where they produce serious harm and may even produce death.

Tuberculosis or consumption is one of these bacterial diseases. The bacteria which cause it are rod-shaped bodies, known as the bacillus tuberculosis. These bacteria infest the air passages and sacs and gradually destroy the surrounding tissues. Unless removed, they eventually produce death. The sputum which a consumptive coughs up consists of masses of these bacteria and broken down tissue, and soon sets free the bacilli if allowed to dry on the floor or street; these may then be blown about by the air and thus be inhaled. It is on this account that Boards of Health make strenuous rules against spitting in public places, and are also coming to insist more and more upon the isolation of consumptives in special sanatoriums where they will not be able to spread this dangerous disease.

Diphtheria, membranous croup, and la grippe are all germ diseases produced by special kinds of bacteria which settle in the throat and grow there. These forms produce their effect by poisons called toxins, which they give off, and which enter the blood, sometimes with fatal results. An effective cure for diphtheria has been found in a liquid called diphtheria anti-toxin; when this is introduced into the circulation it destroys the toxin produced by the bacteria. All diseases that are produced by bacteria are called communicable diseases, since they may be transmitted from one person to another by the air which carries the dread germs. Some of these bacterial diseases are produced

by bacteria which enter the body through the blood or food; but it is evident that the air furnishes a natural highway for them, and this is another reason why we should make sure that our air supply is as pure as possible, and that we keep our respiratory organs in a healthy state, in order that the germs which enter may be removed before they have time to form colonies.

Alcohol and respiration.—It is often claimed that alcohol increases the rate of respiration, and is thus a valuable element in cases where great demands for oxygen are made by the tissues, as in heavy muscular exertions. Careful experiment has shown that alcohol does increase the rate of respiration and the amount of oxygen taken into the body for a period of an hour or more after its use; while highly flavored wines and brandies have an even greater effect in this direction than pure alcohol. When, however, the manner in which alcohol causes this increase in oxygen supply is investigated, it is seen to be a harmful rather than a helpful agent. The reason for this may be stated as follows:

Alcohol causes a congestion of the surface blood vessels and a consequent loss of heat to the body. It also causes temporary increase in the activity of the digestive tissues and the muscles. This loss of heat and action of the tissues creates a demand for more oxygen on the part of the body, and this demand acts upon the respiratory center and makes the rate of breathing faster, in order to furnish the increase in oxygen. In other words, if it were not for the action of the alcohol on the tissues there would be no increase in the breathing rate, and the increase in oxygen thus secured is more than used up in offsetting the loss of heat and in supplying the needs of these unnaturally active

tissues. Hence the tissues which are being used in exercise are worse off than before the alcohol was taken.

In short, the taking of alcohol as a means for increasing oxygen supply is like giving a man an increase of salary and at the same time increasing his expenses by a larger amount. In the end he is worse off than in the beginning.

It is the above result that has made athletes and those who have severe muscular efforts to perform abandon alcohol as a means of aid to their action.

XXII. RESPIRATION IN LOWER ANIMALS.¹

THE activity of tissue cells demands that they receive a constant supply of oxygen; while the result of oxidation in them is the production of quantities of carbon dioxide which must be as constantly removed to permit of their continued activity. In other words, respiration is a property of protoplasm, and wherever there is active protoplasm there must be respiration. Since plants and animals are all built up of cells containing active protoplasm, respiration becomes a property of all living matter. The rate of action of the cells depends upon the completeness and speed with which this exchange of oxygen and carbon dioxide takes place.

In all the lower organisms the mere contact of the cells with a medium which contains oxygen — such as air or water — is sufficient for the exchange to take place. When, however, the organisms increase in size and the number of cells, many cells are necessarily forced into places where such contact is impossible. To these cells the oxygen must be brought in some way, and so in most plants and animals there are found special structures whose purpose is the distribution of oxygen to the tissue cells and the collection of carbon dioxide from them. Such apparatus wherever found, whether in plant or animal, is called a respiratory system. We shall consider

¹ See Footnote, p. 146, Chapter X.

a few of these systems in the animals lower than man and see in what they resemble his or differ from his.

Sources of oxygen.—All animal forms¹ obtain their oxygen from one of two sources, —from air or from water.

Water has the power to dissolve certain gases with which it is in contact, and thus water which is exposed to air for any length of time is bound to contain more or less oxygen in solution. Animals which have the power to extract this dissolved oxygen from water for use by their cells, are often called water-breathing, to distinguish them from those forms which take their oxygen from the air or air-breathing forms. In both air and water breathers the process is the same, namely, the extraction of the oxygen for the use of the tissue cells.

TYPES OF RESPIRATION IN INVERTEBRATES.

Protozoans, sponges, and cœlenterates.—All protozoans (one-celled animals), in their active state, live in water, or fluids containing water, and extract their oxygen from this water. With them, there is no need for blood or other respiratory apparatus, since each cell is in contact with the water, and can take in oxygen and give off carbon dioxide as needed.

The sponges and cœlenterates, such as the jellyfishes, anemones, etc., are built up of many cells, but through the use of cilia or other means the currents of water which bring food to the internal cells are so directed as to bathe

¹ Certain plant forms are known, called the *Anærobia* and limited mainly to certain groups of bacteria, that are able to live in media that contains no oxygen. This does not mean that these organisms are able to live without oxygen, but it is supposed that they have the power to extract oxygen from the food which they receive.

every cell, and thus each cell can effect its exchange of gases directly, in the same manner as the single cell of the protozoan.

The starfish and its allies.—In the starfish we have the simplest kind of a respiratory system. In these animals a tube is found which conveys water to various parts of

FIG. 154. — Vertical section of one arm of a starfish; *b*, ampulla; *d*, stone canal; *i*, radial ambulacral vessel; *ft*, feet. — DAVISON ZOOLOGY.

the body. The same tube is used in distributing water to the tube feet (see p. 305), and this tube serves not only to make locomotion possible, but also, by the water which it carries, to distribute oxygen to various parts of the starfish body and to receive carbon dioxide from them. These starfishes also have soft membrandlike extensions of the skin that line the body cavity and project as pouches, or pockets, between the plates on the upper surface of the body. Oxygen is absorbed from the sea water into the body cavity through these thin, membranous pockets, and carbon dioxide passes out through them, also. The starfish, therefore, has two entirely distinct methods of respiration. It is to be noted that while this animal also possesses a partly developed circulatory system the oxygen distribution is not dependent upon it. This water-tube system of distribution is also entirely distinct from the

food-distributing system. Such an apparatus is a true adaptation for respiratory purposes.

Worms.—The worms obtain their oxygen supply by absorbing it from the air or water through the skin, which is thin and moist, and in direct contact with the air or water in which they live. In some cases this skin is produced into pouches or tufts in order to increase the respiratory surface, and such tufts are called gills. After the oxygen has passed into the body cavity through the gills it must be distributed to the tissue cells. In some worms this distributing is performed by the digestive fluids. In others, such as the earthworm, this oxygen passes through the thin walls of a blood tube where it unites with hæmoglobin dissolved in the blood fluid, and, in this case, the blood acts as a true distributor of the gas to the cells. This blood also collects the carbon dioxide, which is passed out through the skin in the same manner that the oxygen enters it.

In the earthworms we find the first example of the blood acting in connection with the skin as a medium for the distribution of oxygen and carbon dioxide. It is to be noted that in this case the skin is the respiratory organ and the blood plasma the distributor, since the blood contains no corpuscles. In order to absorb oxygen the skin must be very thin and moist.

Clams and other mollusks.—In the clam and the oyster there is a heart and a definite system of branching blood vessels. Some of the smaller branches pass into four flaps

FIG. 155.—
The lob-
worm,
show-
ing tuft-
ed gills.

of soft skin (in the clam two on each side of the body) called gills. These gills are thin-walled membranes or

FIG. 156 — Diagrammatic section of a clam (*Anodonta*); *a*, lobes of mantle; *b*, gills, showing transverse partitions; *c*, ventricle of heart; *d*, auricles; *e*, pericardium; *f*, *g*, kidneys; *h*, venous sinus; *k*, foot; *A*, branchial, or pallial, chamber; *B*, epibranchial chamber.

extensions of the skin which cover the body of the clam, and thus the water in which they lie and which passes over them is separated from the blood by only this thin skin and the thin walls of the vessels. Oxygen is able to pass easily through these membranes and enter the blood just as it does through the air-sac linings and capillary walls of our lungs. The blood of the clam contains no corpuscles and no hæmoglobin and is colorless. It does, however, contain a substance similar to hæmoglobin called hæmo-

cyanin, which takes up the oxygen and distributes it. At the same time the blood passes its carbon dioxide out to the water and is thus aerated. This aerated blood passes from the gills back to the heart, and is then pumped over the body to the tissues, finally returning to the gills, and so on indefinitely.

Here in the clam, then, we have a definite structure developed solely for respiratory purposes, and showing practically the same relation between blood and the oxygen-supplying medium as that seen in our air sacs. The clam, also, has a pair of tubes which cause a constant stream of water to pass into and out of the shell and over the gills,

so that the supply of oxygen and the removal of carbon dioxide is constant. In the oyster the system is similar, except that there are no tubes, as the movement of the sea water is sufficient for this renewal and removal.

In the air breathing mollusks, of which the snails are a type, the blood vessels line a cavity in the body, and this cavity is connected with the outside by an opening. This cavity is an infold of the skin instead of an outfold, but its action is identical with that of the gill. Into this aperture the snail draws a bubble of air, and even the water snails creep to the surface for this purpose, and sink to the bottom again after filling the cavity. The blood vessels which line this cavity effect an interchange of gases with this air just as the gills do with the water, and after all the oxygen is removed the walls of the cavity contract and force out the bubble, now laden with carbon dioxide, and take in a new supply of air. This is the simplest form of a lung known.

Insects.—In the insects we find air breathers having a respiratory system unlike that of any other animal. These air breathers have openings called spiracles along the sides of the abdomen. By movements of the body these spiracles can be made to open and close rhythmically. They are actually the mouths of a system of branching tubes or tracheæ which penetrate to all parts of the body, and thus carry air directly to the various body tissues. These tubes are strengthened internally by a spiral thread which prevents their collapse as the air rushes in and out. The insects are the only animals that carry air directly to all parts of the body. The movements of the abdomen maintain a constant circulation of air through these tubes, in and out of the spiracles. This system is entirely inde-

pendent of the blood system, and in these animals the blood has practically no part in the exchange of gases.

Gills.—All the higher animals, including fishes, frogs, reptiles, and mammals, breathe by the aid of gills or lungs according as they are water breathers or air breathers.

FIG. 157 — The respiration system of an insect ; *a*, an air sac ; *b*, a tracheal branch ; *c*, a spiracle or breathing pore.

A gill in its simplest form is a fold of skin, usually lined with capillaries and so placed that the oxygen-supplying water shall pass over it, while the thinness of the skin and capillary walls permits the exchange of oxygen and carbon dioxide between the blood and the water. The pouchlike extensions of the starfish membrane differs from this definition only in the lack of blood capillaries. In this case the exchange of gases is between the body cavity and the water. The blood tubes appear first in the earthworm where the whole skin acts as a gill.

In the clam we find the simple fold relation, exactly as in our definition, and the clam gill may, therefore, be taken as typical.

The fringed gill appears first in the animal scale, in the crayfishes, lobsters, and crabs. Here, folds of skin

FIG. 158. — The lobster, showing fringed gills (*h*).

lined with blood tubes are attached to the bases of the legs and inclosed in a cavity formed by the shell. Instead of a flat fold, however, the skin is broken up into tiny, cylinderlike projections or fringes, thus increasing enormously the absorbing surface without increasing the space occupied by the gills. This fringing, then, is an adaptation for increase of absorbent surface without increase in space occupation, and is made necessary by the increased demand for oxygen on the part of the animal.

In these forms, the cavity is constantly supplied with water, and the movement of the legs causes the gills to wave about and thus bring all surfaces of the fringes in contact with it. Though this gill chamber, and the apparatus for circulating water through it, may be called accessory respiratory apparatus, it bears the same relation to breathing as do the rib walls and diaphragm in our body.

The fringed gill reaches its highest development in the fishes. Here, the gills are placed in the sides of the head, supported on arches of bone and in such a position that the fish can maintain a steady stream of water over these gill

fringes by constantly taking in and forcing out water through slits in the side of the head. The rich supply of

blood capillaries in these fish gills is indicated by their red color. The blood of the fish passes from the ventricle of the heart into these capillaries; there exchanges its carbon dioxide for oxygen and then passes directly over the body without first returning to the heart, as in our bodies.

FIG. 159 — Gills and heart of the perch, exposed by removal of gill cover on left side; *a*, first of the four bony arches which carry the gills; *b*, gills; *b'*, lower edges of gills on the right side; *h*, heart.

While the gill is well adapted to gas exchange with water the small proportion of oxygen found in water makes that element a poor source of oxygen, and all animals higher than the fishes obtain their supply of this gas from a medium much richer in it, namely, air. Such animals extract their oxygen supply from air by structures called lungs.

Lungs.— Just as we defined a gill as an external fold of the skin, so we may define the lung as an internal fold of the skin lined with capillaries and adapted for the exchange of oxygen and carbon dioxide between the blood and the air.

The lung of the snail is the simplest type. Here, the single, internal pocket, lined with blood vessels, may be conceived of as having been developed by a single infold of the external skin. So slow is the exchange of gases in

this animal, and so small the demand for oxygen by the tissues, that a single bubble of air furnishes oxygen enough for a considerable period of time.

In the frog, which in its earlier stages breathes by gills, we find a much more complex lung than this. Here, inclosed in the thoracic part of the body cavity, are two sacs connected at the throat by a common tube and lined with blood vessels. These sacs have their inner surface raised in many ridges that increase their internal surface without increasing the size. As we go higher in the animal scale, these ridges increase in depth, until finally, in mammals, the interior of the lung sac is subdivided into thousands of tiny air sacs connected by tubes to the common trachea. The frog has no diaphragm, and in order to force air into the lungs he first fills the mouth cavity through the nasal passages, then, closing these passages and elevating the throat, forces this mouthful of air down the tube into the lungs. The air is expelled from these sacs by simply contracting the body walls. In the frogs, too, we find the first development of the trachea as a voice-producing organ.

In the reptiles, instead of a ridged sac with a central cavity, the ridges are developed so as to divide the sac into pockets, and these pockets connect with the trachea by tubes. In this case, the two branches of the trachea (the bronchi) extend through the lung sac, and the branches or bronchial tubes are given off to the pockets from the bronchus as lateral branches.

In birds we find the first development of true air sacs, and a branching system of bronchial tubes similar to ours. The bird differs from man in having the lungs connected, by extensions, with air cavities in various parts of the body, and through these it is able to give the body a buoyancy

which helps it in flight. Its method of breathing differs from ours, there being no diaphragm, and the expansions

and contractions are produced by the thoracic muscles. The voice box of these birds is at the bottom of the trachea instead of at the top, and is called a syrinx instead of a larynx.

The mammals have lungs practically identical with ours. As we ascend the animal scale, then, we see a constant increase in the absorbent surface of the respiratory organs, and this increase keeps pace with the increased de-

FIG. 160—Lungs of a bird; *a*, trachea; *b*, bronchi; *c*, lungs; *d*, apertures communicating with air cells.

mand for oxygen. All lungs are alike in having the blood capillaries separated from the air by a thin membrane through which the exchange of gases may take place, and in having some method of introducing new air into these sacs and removing it at regular intervals, in order to maintain a constant supply of oxygen to the animal. In all vertebrates the oxygen extracted from the air is carried to the tissues by the blood corpuscles which contain hæmoglobin for this purpose. The increase in the number of membranes in the lungs is simply an adaptation for protec-

tion, or is made necessary by the increase in surface by foldings.

Cold-blooded and warm-blooded animals. — By cold-blooded animals we mean those in which the temperature of the blood is only slightly higher than that of the surrounding environment. These animals feel cold to the touch, as they are of lower temperature than our bodies. This cold-blooded condition is dependent upon the amount of oxidation which takes place in the body, and fishes, frogs, and reptiles have a slower rate than that of birds and man. The higher temperature of birds and mammals is evidently due to the fact that the greater perfection of the respiratory apparatus permits a more rapid exchange of oxygen and carbon dioxide in these animals than in the cold-blooded forms. As a rule, cold-blooded animals are more sluggish in their movements than warm-blooded ones.

XXIII. EXCRETION.

IN the preceding chapters we have traced in detail the processes by means of which food and oxygen are obtained and distributed to the cells of our tissues. We have also seen that the function of some of this food is to make new protoplasm, while other food substances free energy by uniting chemically with oxygen in the process called oxidation, and thus give to the protoplasm its varied power of action, secretion, contraction, or the like. The making of food into new protoplasm we call assimilation, while the production of energy we call oxidation. We give the name of constructive metabolism or anabolism to all the processes which lead up to the production of activity on the part of the cells. When oxidation takes place two results follow, namely, the activity of the cell through the energy produced, and the production of wastes such as carbon dioxide and urea. This breaking up of the cell through oxidation is called destructive metabolism or katabolism. Anabolism and katabolism, which together make up metabolism, are going on continually as long as the cells remain active. We have seen in our study of respiration how the cells receive their oxygen. We have also noted that the blood carries the carbon dioxide back to the lungs, and that these organs remove it from the body. How are the other wastes removed from the cells? To answer this question we must study another set of organs whose main function is the removal from the body of waste products of the cell. All organs concerned in

this removal of body wastes are grouped together under the name of the *excretory system*. The lungs, by removing the carbon dioxide, may be included in this system, but the removal of the principal body wastes is the work of the kidneys and the skin.

The skin.—If we examine the surface of the body carefully we find it everywhere covered with a layer of flexible, elastic tissue which we call the skin. The hair and nails are modifications of this tissue and are included in the study of skin structure.

Examination of a section of the skin taken from any surface of the body shows it to consist of two distinct layers. The upper layer (epidermis, or cuticle) is thinner

FIG. 161 — Diagram of a section of skin.

than the lower layer (dermis, or corium), and contains no blood vessels. Hence, it is possible to prick and even tear off portions of the epidermis without causing bleeding. This epidermal layer varies greatly in thickness in different

parts of the body, being thickest at points where there is greatest friction, as on the sole of the foot and the palm of the hand.

The inner layer, or dermis, contains the blood vessels, nerves, and various kinds of glands, and is easily recognised by its pink color.

Epidermis.—Examination with a microscope shows this layer to be composed of many cells held together by a sort of cement. The lowest layer is made up of a single row of active, columnar, epithelial cells. These cells absorb nourishment from the blood and lymph of the dermis, and, by growth, produce other cells which are gradually pushed upward. In this upward movement

FIG. 162 — Portion of the epidermis, showing the transition from the lower living cells to the upper horny scales.

they gradually lose their columnar shape, and become flattened, so that the upper layers of the epidermis consist mainly of flattened cells. Gradually, too, the protoplasm of these cells dries up and they are transformed into flat, horny scales which make up the uppermost layer of the epidermis. These outer horny scales are being constantly worn away and renewed from below, and the epidermis, as a whole, shows transition stages of cells from the active columnar layers near the dermis, to the dead horny outer layer above. Scattered among these cells may be found occasional cells which contain pigment or coloring matter. These are called pigment cells, and in the negro race they are very abundant and give the black color to the skin. The distinction between a blonde and a brunette is merely a difference in the number of these pigment spots or cells. Freckles are due to the collecting of these cells in spots.

Examination of the epidermis on the hand shows it to be raised in ridges parallel to one another, and so arranged as to form various patterns. Since these patterns do not change throughout life they serve to identify criminals. A hand lens also shows that the epidermis is perforated at frequent intervals. These perforations, or pores, are the openings of tiny ducts which extend downward to the sweat glands of the dermis and discharge the perspiration. In other places the epidermis is pierced by hairs which also have their source in the dermis.

As has been already noted, no blood vessels enter the epidermis; the active cells receiving their food from the lymph of the dermis. Nerve fibers are wanting in the upper layers of the epidermis, but fibers do extend into the lower cells, so that when this layer is separated from the dermis, as in a blister, the prick of the needle can still be felt by the epidermal covering of the blister though it produces no sensation of pain.

The dermis or corium.—The dermis or true skin is composed mainly of a network of connective tissue rich in elastic and white fibers. In tanning operations it is this layer that is transformed into leather. The dermis is loosely connected to underlying parts by a layer of fibers called the subcutaneous layer, and this loose connection permits the skin to move freely over the underlying flesh. Sometimes this subcutaneous layer is the seat of fat deposits, especially in the layers that cover the abdomen. These fat layers tend to protect the body from cold. The surface of the dermis that is in contact with the epidermis is raised in a great many tiny projections called papillæ. These papillæ are arranged in rows and are the cause of the ridged appearance of the epidermis. Some contain a

knot of capillaries which are supplied with blood from a small artery and empty into small veins. They enable

FIG. 163 — Dermal papillæ; the black lines represent capillaries, and the white lines, nerves. The nerves end in bundles called corpuscles.

the blood to be brought near the active, epidermal cells. The other papillæ contain delicate sense organs connected by nerves with the central nervous system. The function of this latter kind will be discussed more fully in our study of the nervous system. Some are sensitive to touch, and some to cold, and to them is due the power of skin to respond to external sensory stimuli. They are especially abundant at the ends of the fingers, thereby giving to these members their delicate perception of touch. We may distinguish these two kinds of papillæ by calling one kind blood papillæ and the other nerve papillæ. Evidently the dermis is well provided with blood vessels and nerves. Except for these papillæ, the dermis is usually smooth, but in old age the absorption of fat from the subcutaneous layer permits the skin to shrink and form wrinkles.

Besides these papillæ the dermis shows other structures, sweat and sebaceous glands, while hairs and nails are sim-

ply modifications of normal skin layers, and have their origin in the dermis.

Sweat glands. — If we trace a spiral duct which leads downward from a pore in the epidermis, we shall find that it passes through the dermis also and ends finally in the subcutaneous layer as a coil. This knotted coil is surrounded by a network of capillaries, and the combination is called a sweat gland. These coils are lined with a secreting membrane of epithelial cells, and the power of the skin to excrete wastes in the form of sweat or perspiration is due to their activity. Their action is as follows: From the blood brought to them by the capillaries the cells select the salts and nitrogenous wastes (urea). These wastes, together with water and some salts, are passed on by the cells as a secretion to the inside of the tubes where they collect, and finally leave the body through the epidermal ends of the ducts or pores. This action may take place so slowly as to be invisible to the eye (insensible perspiration), or in summer, and in violent exercise, drops may collect on the surface of the body (sensible perspiration). By placing the skin against a cool surface, such as

FIG. 164 — Sweat gland; *a*, epidermis; *b*, dermis; *c*, subcutaneous layer; *d*, sweat gland; *e*, mouth of sweat duct (spiral duct).

a mirror, we can condense the perspiration, and thus demonstrate that even when invisible to the eye the process is always going on.

Sebaceous glands.—These are small, saclike ducts embedded in the dermis, and opening usually by a small duct near the mouth of the hair follicles. They are lined with epithelial cells whose substance becomes gradually charged with oil drops. Finally the protoplasm disappears, the cells break down and their oil is discharged through the duct, new cells taking the place of the old ones and continuing the process indefinitely. While these glands are always found associated with hair follicles, they are also to be found in parts of the skin free of hair. In these cases the oil discharged spreads over the skin and moistens and softens it. When such ducts become stopped by dirt the contents become hardened and the ducts become distended and form black spots on the skin.

Hairs.—A vertical section of a hair shows the following structure: The external hair is a long, hollow shaft made of an extension of the epidermis and embedded at its base in a pocket of the dermis called the hair follicle. This pocket or follicle extends downward into the subcutaneous tissue, and at its base is a special papilla which forms the hair and which is abundantly supplied with nerves and blood vessels. The follicle is lined with a layer of epidermal cells, and the upper part of the hair shaft is composed almost entirely of the horny layers of the epidermis, and hence this part is not sensitive to cutting. The nerves at its base in the papilla render the hair sensitive to touch, and give the sharp sensation of pain when the hair is pulled. The relation of the various parts of the hair to the normal skin layers is shown in Figure 165. The color of the hair

is due to pigment cells lying in the hair shaft.

Attached to the sides of the hair follicle are oblique strips of unstriped muscle by means of which the hair may be moved. When these muscles are contracted by cold, the skin becomes roughened, and this condition is due to the erection of the hairs. Opening into the side of each hair follicle is the duct of a sebaceous gland. The glossy appearance of the hair is due to the oil poured out by this gland.

Hairs may be found on practically all parts of the skin except the sole of the foot and the palm of the hand. Their development varies greatly with the individual. On the head they form a good protection from blows.

Nails.—The ends of the fingers and toes are protected by thickenings of the horny layer of the epidermis called nails. These nails are very similar to hairs in their formation. Their relation to the epidermal and dermal layers is shown in Figure 166. The base is embedded in a fold of the skin, and at its root is a portion of the dermis called the matrix which, by pushing out

FIG. 165. — Section of a hair; *h*, hair shaft showing central medulla or core; *s*, sebaceous gland; *w*, sheath of skin; *f*, fatty tissue. At the base of the hair is seen the papilla which forms it.

new cells, constantly increases the length of the nail. At the same time the dermis just beneath the bed of the nail forms new cells which add to its thickness. The matrix

FIG. 166—Sections of nails. A, median longitudinal section ($\times 4$); *a*, the fold at the base of the nail; *b*, the nail; *c*, the bed of the nail. B, transverse section ($\times 4$); *a*, lateral folds on each side of the nail; *b*, the nail; *c*, bed of nail showing papillæ. C, portion of bed of nail ($\times 200$); *e*, nail, formed of horny dermal scales; *d*, deep layers of epidermis; *c*, papillæ.

may be compared to the hair papilla, and injury to hair or nail will be repaired so long as the hair papilla or the nail matrix is uninjured. The nail shows the following areas: The part embedded in the dermal pocket is called the

root. The visible part consists of a body or bed of the nail which is attached to the dermis, and a free edge. Near the root is a half moon, or *limula*, which is whiter than the rest. This difference in color is due not to the nail itself which is transparent, but to the fact that the blood papillæ are less numerous under this part of the nail.

General physiology of the skin.—The principal uses of the skin may be classified under the following heads:

(a) It affords protection to the underlying muscles and organs, and at the same time, by its loose connection, permits free movement of these underlying parts.

(b) It permits the evaporation of water from the surface of the body, and thus regulates the temperature of the body. In hot weather, or when one is exercising, the excess of internal or external heat is attended with profuse perspiration, and this heat is used to evaporate the water of the perspiration, and hence does not raise the body temperature. By this means the body temperature is maintained at an even rate of about 98° Fahrenheit.

(c) Its underlying layers of fat also prevent too great a loss of heat from the underlying parts.

(d) It contains sensory papillæ of various kinds which are connected with the central nervous system (spinal cord and brain), and these papillæ permit the response of various parts of the surface to external stimuli.

(e) It is an excretory organ, and by means of the sweat and sebaceous glands it is able to extract wastes from the blood and remove them to the outside of the body.

Action of the skin as an excretory organ.—The precise composition of sweat is difficult to determine, as it is usually mixed with the secretion of the sebaceous glands

It is a very thin liquid, slightly heavier than water (Sp. Gr. 1.004), and usually shows an alkaline reaction from the dissolved salts present. These salts consist mainly of alkaline phosphates and sulphates and sodium chloride. The organic wastes (urea, uric acid, etc.), which are produced by the breaking down of the proteid parts of the tissue cells, are present only as mere traces, though as the result of muscular work the amount of urea thus thrown off may become considerable. The principal waste given off by the sweat glands, however, is water, and it is as a remover of water from the body that the skin is especially important. The amount of water thus removed varies greatly with conditions of temperature, and with the mental and physical states. It has been calculated that the average amount of sweat given off in twenty-four hours by a normal person is between two and three quarts.

The sebaceous glands contribute an oily liquid that sets in a cheesy mass upon exposure to the air. It contains fats, soaps, and some proteid materials. It varies in quality in different parts of the body, that of the ear constituting the ear wax. Its principal function is to soften the skin and hairs, and protect the skin from external moisture, while it also tends to prevent loss of heat by evaporation.

In some of the lower animals the skin also excretes carbon dioxide and takes in oxygen, thus acting as a respiratory organ. In man the amount thus given off by the skin is small under ordinary conditions. In profuse perspiration, however, the amount of carbon dioxide removed in this way becomes considerable. Under ordinary conditions the removal of carbon dioxide takes place in the lungs.

HYGIENE OF THE SKIN.

The various excretions of the skin tend constantly to deposit increasing layers of waste on its surface. The accumulation of this waste clogs the pores and prevents the escape of further wastes. Since these wastes must leave the body somewhere, the result is the overworking of the kidneys and lungs, while the lack of water evaporation permits an excessive increase of body temperature. If, however, the pores are clogged with material which conducts heat there results a lowering of body temperature that produces death when any considerable portion of the skin has been painted or varnished.

To avoid any of these dangers, it is evidently imperative that the skin be bathed frequently, and thus the excess of waste removed and the pores kept open. Cleanliness of person, therefore, becomes one of the first laws of health.

Baths. — The oily secretions and the grimed-in dirt require more than the mere application of cold water for their removal. Soap contains an alkali which dissolves these oils and loosens the dirt. On this account exposed parts of the body should be washed with soap and warm water at least three times a day, and the whole body at least once a week. Care, however, must be taken in selection of the soap used, as many contain an excess of alkali which tends to injure the skin. Baths not only serve to cleanse the skin, but, if properly taken, tend to strengthen and invigorate the whole system. For this purpose a cold bath is better than a warm one, since warm baths tend to open the pores more widely, to dilate the skin arteries, and thus increase the perspiration and render the body liable to colds. On the other hand, cold baths tend to contract the arteries at first, leaving the skin pallid. This effect is soon followed by a reaction, the arteries fill with blood, and the skin begins to glow. This glow may be increased by a thorough rubbing with a rough towel which at

the same time removes the scales of dead epidermis. If the bath is continued too long a second reaction takes place, the skin becomes pale and chilled, and the person becomes depressed as much as from excessive warm bathing. A warm bath followed by a cold shower is often beneficial in producing a healthy glow. When we say cold baths we do not mean that the temperature must be excessively low. Some persons cannot stand the shock of too cold water, and for them water which just feels cold to the skin may produce better results than colder water. The salts in sea water tend to stimulate the skin nerves, and thus aid in producing the glow. Salt baths may often be taken with good effect when cold fresh water depresses. Cold baths should not be taken when the body is depressed, as they cause too much loss of heat. They should never be taken immediately after a meal, as they draw to the surface the blood required for digestion, and thus may produce serious indigestion. Shower baths take less heat from the body than immersions, and also give a greater stimulus to the skin nerves. Rightly indulged in, the cold bath tends to keep the skin clean and to invigorate the system, at the same time rendering the person less liable to colds. A judicious mixture of cold and warm baths is best for maintaining the body in the best possible state.

Care of the complexion. — The use of paints, powders, and external applications for the beautifying of the complexion is injurious, for the reason that, like dirt, they clog the pores of the skin. Many of them, too, contain poisonous substances which may be absorbed through the pores and produce disease. Good exercise and frequent baths will keep the skin glowing in health, and render it far more beautiful than any artificial applications.

Care of the hair. — The hair is naturally kept soft and glossy by the secretion of the oil glands. If it is not washed frequently, and the scalp rubbed thoroughly dry after the washing, this oil will collect, mix with the perspiration and dead-skin scales, and clog the pores of the scalp. Washing, therefore, is important in removing this accumulation, while the rubbing stimulates the blood supply and thus furnishes food to the hair papillæ. Leaving the hairs wet at the roots tends to rot them.

Care of the nails. — Children are very apt to bite the nails. This not only gives the hands a ragged appearance, but also is apt to tear the nails loose from the bed and thus permit germs, which collect in the dirt under the nails, to get into the blood and produce sores. For this reason also it is better to remove this dirt with a dull instrument, as a sharp blade is apt to cut the skin. The epidermis around the base of the nail also tends to grow to the nail, and as the nail lengthens the skin is torn and hangnails are produced. When this is the case, they should be cut back close to the skin. They may be prevented by loosening the fold of skin about the base of the nail with a dull instrument.

Burns and blisters. — Blisters are caused by friction or heat which kills the lower layers of the epidermis and permits the lymph and water to collect between the dermis and the epidermis. If this distended epidermis is pricked with a clean needle and the water allowed to escape, the two surfaces will eventually knit together again. Care must be taken not to tear the upper layer, as this permits the dermis to become uncovered, and often painful sores result. Ordinarily the epidermis will not absorb poisonous germs. When this layer is removed or the dermis is cut the blood vessels are exposed, and in this case they should be covered with some germicide material to prevent the absorption of germs into the blood. The germs which cause lockjaw and blood poisoning are very frequently absorbed from lack of precautions in this direction, and may even produce fatal results. One of the simplest germicides for this purpose is a 1% to 2% solution of carbolic acid. A cut should be cleansed first with water and if dirty, with carbolic acid solution. Then bandage it with cloths which are strictly clean. (Small cuts should receive as much care as large ones in this respect.)

Burns tend to produce blisters and kill the skin cells. They also bring about inflammation. All burned places should be kept from contact with the air, because this treatment lessens the pain and reduces the inflammation. For this purpose baking soda or vaseline is a useful application. Acid burns should be treated with weak alkalis which neutralise the acids, while burns caused by alkalis should receive neutralisation with weak acids.

Corns. — When the skin on the feet or the hands is rubbed, the parts protect themselves by increased development of the epidermis and thus form horny spots or callouses. Sometimes,

FIG. 167 — Section of a corn. Note the thickened epidermis over one papilla, and the manner in which it presses into the sensitive dermis below. OVERTON, *Applied Physiology*.

on the feet, these spots grow inward, and a point of hardened cells forms which presses inward into the sensitive dermis. The result is a painful formation called a corn.

OTHER EXCRETORY ORGANS.

The urinary organs. — On each side of the backbone, in the part of the abdominal cavity a little below the lower ribs, lie two bean-shaped bodies about four inches long, two inches wide, and one inch thick. These two organs are the kidneys. They are embedded in a thick layer of fat, and from the curved-in portion of each emerges a tube called the ureter. These ureters are continued downward and enter a baglike body covered externally with muscle and lined with mucous membrane. This body is the blad-

der. From the bladder a common duct called the urethra extends downward to the external urinary opening.

Each kidney receives from the abdominal aorta an artery called the renal artery. This artery breaks up in the kidney into fine capillaries which are eventually collected into a vein called the renal vein. The two renal veins, one from each kidney, ultimately return the blood to the inferior vena cava. The renal arteries and veins enter and emerge from the kidneys at the same point as the ureters.

FIG. 168—The kidneys and connections; *k, k*, kidneys; *rv*, renal vein; *ra*, renal artery; *u, u*, ureters; *b*, bladder.

General functions of the kidney. -- The kidneys receive the waste-laden blood of the body through the renal arteries. Out of this blood they take the waste material (mainly urea) and form of it a secretion called urine (urea dissolved in water). The waste-freed blood then returns to circulation by way of the renal veins, while the urine is poured from the kidneys into the bladder as fast as it is formed. When a sufficient quantity has collected in this bladder, the muscular walls contract and the liquid is expelled from the body by way of the urethra and urinary

opening. Unlike the skin, the action of these bodies is purely excretory, and to them is due the purification of the blood.

Structure of the kidney.—The ordinary lamb's kidney which may be obtained from the butcher serves well to demonstrate the structure of this important organ.

Removing the fat from the outside we find the reddish-brown kidney within, still surrounded with a transparent membrane or capsule. This membranous capsule is called the renal capsule. Remove this and we have the kidney.

FIG. 169.—Median longitudinal section of a kidney; *Ct*, cortex; *M*, medulla; *P*, pelvis; *U*, ureter; *RA*, renal artery; *Py*, pyramids of Malpighi.

Viewed from the outside it shows a bean-shaped outline, being convex on one side and concave on the other. In the depression on the concave side may be made out the connecting renal vein and artery and the ureter at the point where they enter the body of the kidney. If we cut the kidney lengthwise the cut surface shows the following areas (see

Fig. 169). As it enters the kidney the narrow ureter is seen to widen into a cavity called the pelvis. From this pelvis extend branches forming what are called calices. The solid portions show two layers. The outer is called

the cortex, and is the darker and more compact of the two. The inner layer is called the medulla. This medullary substance is less red than the cortex, and is divided into pyramids whose small ends are toward the pelvis and whose bases are toward the cortex. These pyramids are called the pyramids of Malpighi.

The renal artery divides in the hilus (the name given to the depression), and its branches extend between the pyramids to the cortex. While they give off a few branches to the pyramids they end in a much richer network of capillaries in the cortex, and to this fact is due the deeper color of the cortex. These cortex capillaries are in turn collected into veins which ultimately empty into the common renal vein, following a path similar to that of the arteries through the kidneys.

The pyramids of the medullary layer are composed of fine tubular glands. At the points of the pyramids are found opening into the calices the mouths (the papillæ) of many tiny tubes (the uriniferous tubules). These tubules extend backward into the medullary layer, branching frequently, and thus forming the pyramid.

Ultimately these branches end in the cortex, where they swell out and form a spherical capsule (Malpighian capsule.)

FIG. 170—A uriniferous tubule; *r*, cortical portion of kidney; *g*, medullary portion; *p*, a pyramid; *I*, a Malpighian capsule; *II* and *V*, twisted portions of tubule; *III*, *IV*, *VI-VIII*, straight portions of tubule; *IX*, opening of tubule into pelvis at top of pyramid.

The entire length of the tube, its branches and capsules, is lined with epithelial gland cells. The relation of these parts is shown in Figure 172. Into each capsule enters a tiny branch of the renal artery and there breaks up into a twisted knot of capillaries called a glomerulus which

FIG. 171—A Malpighian capsule; *v. a*, small branch of renal artery breaking up in the glomerulus (*g. l*) into capillaries, which in turn unite into the renal vein (*v. e*); *c*, tubule; *a*, epithelium over the glomerulus; *b*, epithelium lining the capsule.

FIG. 172—Relation of parts in kidney; *a*, artery; *v*, vein; *c*, capillaries about the tubule; *g*, glomerulus; *t*, tubule.

nearly fills the capsule. These capillaries pour their contents into a tiny vein, which, after leaving the capsule, breaks up again in a network about the walls of the uriniferous tubules. This network fuses again into a single tube which connects with the renal vein.

Excretory action of the kidney.—The cells which line the capsules and upper parts of the tubules are true gland or secreting cells. Those in the lower part of the tubule are purely for protection. The cells of the capsule secrete only water and inorganic salts which they receive from the

blood of the glomeruli. It was once supposed that the glomerulus was simply a filter, and that the waste and salts passed through by filtration and osmosis. Present theories tend to prove that the cells of the capsule actually secrete this water and salt material first received from the blood. The cells of the upper parts of the tubules secrete the organic wastes (urea, etc.) which they obtain from the capillaries surrounding these parts and which is dissolved in the water to make the urine. The water, salts, and organic wastes accumulate slowly, and drop by drop they pass through the length of the tubes into the calices and pelvis. Here they enter the ureter, and the combined liquid is stored in the bladder pending removal. The bladder is covered with a layer of muscle and peritoneum, and lined with mucous membrane, and the mouth of the urethra is closed by a sphincter muscle which opens from time to time to permit the escape of the urine.

The urine.—Analysis of the urine shows it to consist of a yellowish liquid slightly heavier than water and consisting of about 96% water and 4% solids. The principal solid constituent is a nitrogen waste called urea, and the most important excretory action of the kidneys is their power to remove the nitrogen wastes of the body.

In the metabolism of the tissue cells the nutrients are broken up into water, salts, carbon dioxide, and nitrogen wastes. We have already seen that the water as sweat, is largely excreted by the skin, and the carbon dioxide by the lungs. The kidneys, therefore, are the great nitrogenous waste disposers. The total amount of water, salts, and nitrogenous wastes given off by the kidneys in twenty-four hours varies from forty to sixty fluid ounces or one thousand five hundred grams. Of this, water forms 96%,

urea 2.2%, other nitrogenous wastes 0.17%, mineral salts about 1.6%.

The liver as an excretory organ. — We learned (see p. 144) that through the bile the liver has the power to remove various poisons and other wastes. In this respect it is also an excretory organ. It is now believed that the urea or nitrogenous waste is not formed in the kidneys, but is present in the blood which flows to the kidneys, these simply secreting it from that blood. It has been further proved that urea is formed, in part at least, in the liver and then sent in the blood to the kidneys for removal. Just how the wastes of the cells are made into urea, and where it takes place, is still uncertain, but it is already demonstrated that the liver performs an important part in this operation.

Hygiene of the kidneys. — The skin excretes very little urea and much water. The kidneys, on the other hand, excrete much urea. When much water is given off by the skin, as in summer, less is given off by the kidneys; while in winter, when perspiration is less, the amount of water disposed of by the kidneys is correspondingly increased. The fact that more blood passes to the skin in summer also has an effect in this direction. Further, the perspiration and blood supply to the skin is governed mainly by the external temperature, while the amount of water excreted by the kidneys is governed partly by the amount of water swallowed. Now a decrease below a certain limit in the amount of water excreted by the kidneys tends to prevent the proper washing away of the urea, while, if not washed away, the accumulation of this urea colors the urine and tends to clog the action of the kidneys.

For the above reasons, persons suffering from kidney troubles may often remove this trouble by increasing the amount of water drunk. Urea itself is not poisonous, but when the nitrogen of the tissues is not properly oxidized other substances are produced which pass in the blood to the kidneys. The kidneys try to excrete these products, but often become overworked in the attempt, and these poisonous substances collect and produce the disease known as Bright's disease. An excess of sugar in the diet often brings about an imperfect oxidation, and patients with this disease often have their sugar supply cut down for that reason. Exercise, which maintains a healthy oxidation of the tissues, indirectly helps to maintain the health of the kidneys.

Alcohol and its effect upon the skin and kidneys.—One of the first effects of alcohol drinking is to cause congestion of the surface blood vessels. When the use is occasional, this congestion may pass off and the skin return to its normal condition. If continued, however, the constant use of alcohol causes the blood vessels of the skin to become permanently dilated, the skin becomes red and often blotched with pimples, due to poor nutrition of the skin cells and the over-accumulation of wastes.

Alcohol also causes more or less congestion in the blood vessels of the kidneys, and the result may be that the kidneys are overworked, with results seriously menacing the health of the individual. All these effects, of course, vary in proportion to the amount of alcohol consumed.

XXIV. SKIN STRUCTURE AND EXCRETION IN THE LOWER ANIMALS.¹

IN comparing the skin of man with that of the lower animals we shall have to modify our conception of the word skin. In man we used the word to describe a two-layered system of tissues whose function was fourfold: protection of underlying parts, excretion of wastes, the end organ of various sense nerves, and the regulator of the body temperature. When we speak of the skin of the lower forms, we have in mind only one characteristic of the human skin, namely, a protective layer of some kind. In the consideration of the different groups of animals it will be of interest to note the manner in which the body coverings of the various forms approximate to the complex structure of man.

Protozoans.—Since the protozoan consists of only one cell, it can evidently have no skin in the sense of a tissue structure. Many protozoans, however, do develop a wall or outer covering for their protoplasm which in some respects fulfils the functions of the skin of higher forms. For example, in the thin membrane or cuticle, which is secreted by the ectoplasm of the paramœcium, we have a layer which protects the contents of the cell, and at the same time permits the exchange of oxygen and carbon dioxide to take place through it. In permitting the escape of carbon dioxide it is as truly a means of excretion as the thicker membranes of other forms.

¹ See Footnote, p. 146, Chapter X.

Sponges and coelenterates.—In these animals the body is composed of only two layers of cells, and for the most part the action of these cells is on the same plane as that of the protozoan. In the sponge, the variation between the inner and outer cells results in division of labor, and the outer cells are more protective and less digestive in their action.

In some of the coelenterates, however, we find a simple form of covering which approaches closely in structure the skin of man. In certain of the polyps, for example, we find the outer ectoderm layer of cells more or less completely divided into two layers. Where this occurs the outer layer is composed of cells very similar to those of the epidermis, while the inner layer may contain muscle fibers, nerves, touch organs, and certain simple glands. Such a covering is evidently much more than a protective form, and in fact performs several functions identical in nature with those of the human skin.

Mollusks.—In the oyster, clam, snail, and other members of this group we find the covering of the body composed of a true two-tissued layer. The outer layer is composed of epithelial cells, while the underlying part is made up of connective tissue in which are found muscle fibers, blood vessels, and nerve ends. This skin also acts very effectively in the discharge of carbon dioxide from the body. It differs from other forms of skin by the presence of special shell-secreting glands and thus is the source of the hard exoskeleton which we call the shell.

Worms and arthropods.—In the members of these two great classes we find also a true skin. The outer body layer, or hypodermis, as it is called in these groups, has the power of secreting a thin layer of material, which may

be so thin, in the earthworm, as to permit of the action of respiration, while, in the insects or crustaceans, it may be so thick, or become so loaded with salts of lime, as to form a hard external skeleton for the animals. When we speak of an insect's shedding its skin, we mean that it sheds this outer secretion or exoskeleton. The underlying layer or hypodermis remains and serves to secrete the new layer. This hypodermis is also found to contain glands of various kinds, and may even show layers of nerve and muscle fibers. The hard layer of the insect and lobster is evidently not a satisfactory substance for permitting the excretion of wastes, and very little excretion or respiration takes place through the skin of such forms.

Body coverings of vertebrates.—All vertebrates possess a dermis and epidermis similar in structure and function to that of man. The epidermis always consists of epithelial cells only, while the dermis is made up mainly of connective tissue cells and fibers, and may also inclose muscle fibers. While the various glands of the skin are composed of epithelial cells they are found most often embedded in or under the dermis, and have their opening by ducts on the surface of the epidermis. The epidermis is in two layers, as in man; the outer being composed of

FIG. 173—Skin of fish (diagram); *a*, dermis with vertical fibers, *b*; *c*, gristly layer; *d*, surface scales; *e*, layer containing limestone; *f*, scale pit.

dead cells or cuticle, with an underlying layer of living cells. In some vertebrates this outer layer, or cuticle,

becomes so thick and horny as to form a true exoskeleton, namely, the scales of the crocodile and the horny layers of the turtle. Again, in the amphibians it may be so thin that respiration takes place through it with readiness.

The skins of vertebrates differ mainly in the structures produced by modifications of the epidermis and the forms of glands. For example, the scales of the fishes and the bony horn-covered plates of the turtle are of like origin. Both consist of a bony plate developed by the dermis, and covered with a horny portion of the outer layer of epidermis. The feathers of the bird and the hairs of man are likewise identical in origin, being merely modifications of the epidermal layers.

In like manner, by infolding of the epithelial cells of the lower epidermis, glands of various forms are built; such are the sebaceous and sweat glands of man, the mammary glands or milk glands of all mammals, the various scent glands of different forms, and the slime glands of the amphibians and some of the fishes. Some of these glands are purely excretory in function, while others are protective (scent glands) or produce secretions of value in animal operations; for example, the slime glands of the amphibians

FIG. 174—A feather; *a*, quill; *b*, shaft; *c*, vane; *d*, down; *e*, *f*, upper and lower orifice.

make the body difficult to grasp, and at the same time moisten the outer layer so as to permit exchange of carbon dioxide and oxygen. Claws, spines, fur, hoofs, horns, etc., are all modifications of the horny outer part of the epidermis, and each arises from a dermal papilla, or papillæ, just as do the hair and nails of man. The dermal layer of animals is often much thicker than that of man, and when tanned it forms what we call leather.

EXCRETORY ORGANS OF THE LOWER ANIMALS.

Aside from the use of the skin as an excretory organ, there are no structures comparable to kidneys in any of the lower animals until we reach the worms and mollusks. The protozoans carry on the removal of wastes by simply throwing out the wastes of metabolism into the surrounding water or liquid in which they live. As the animals become larger, the cells multiply in numbers, and it becomes more difficult for each cell to get rid of its wastes. If this waste thrown out by the cells remains in the body it may produce serious harm to the tissues. The solution of this difficulty is met in the development of glands whose function is to take the collected wastes of the cells from the circulatory system, to transform these wastes into a secretion of uniform character, and then remove this secretion to the outside of the body.

All such glands, of whatever character or structure, are identical in plan with the air sac of the lung, the sweat gland of the skin, and the kidney of man. In all, the cells which do the secreting are placed in such relation to the blood system that the body wastes (carbon dioxide, water, or urea) may be readily absorbed, and when transformed

into a secretion shall pass into some kind of a duct through which it escapes to the outside. The function of the kidney of man is the removal of nitrogenous wastes, that is, the removal of the wastes resulting from metabolism of proteid in the protoplasm. We have already discussed the adaptations of the lower animals for the removal of carbon dioxide under the head of the respiration. It remains to examine the nitrogen waste removers of these lower forms, and in the following paragraph we shall consider only the organs which perform the same action as the human kidneys.

Nephridia.—One of the simplest forms of kidney is found in the group of mollusks and earthworms. It is called a nephridium, and as the earthworm is perhaps the most familiar type we shall describe the organs as seen in this animal. If we turn the earthworm over and examine its ventral surface we can make out a double row of pores, two to a segment, extending from the head end to the tail end. If we dissect the worm and trace the internal connections of these pores we find them to be the mouths of long coiled ducts which, after twisting about in the body cavity, end in funnel-shaped, ciliated mouths in the segment just in front of the segment in which the pore is located. (See Fig. 176). Further examination shows this duct to be formed in part by secreting cells and surrounded

FIG. 175.—A nephridium; *S*, septum; *F*, funnel; *O*, pore or external opening.

by branching blood vessels. Its action is as follows. The waste from the body cells of the segment collect in the

FIG. 176—Cross section of earth worm showing nephridia in position; *B. C.*, body cavity; *c*, cutis; *e*, epidermis; *c. m.* & *l. m.* muscle layers; *N*, nephridia; *ex. p.* excretory pores; *S*, setæ; *D*, dorsal blood vessel; *I*, intestine; *V*, ventral blood vessel; *N. V.*, ventral nerve cord.

cavity or are absorbed into the blood circulation. Those wastes of a nitrogenous character which remain in the cavity of the segment pass directly into the ciliated end of one or the other of the nephridia, and, after action by the cells, pass out of the pore as a secretion. Similarly, the wastes which entered the blood are eventually brought to the branches which surround this tube, and, being absorbed from the blood by the secreting cells, are made into a secretion, and in turn pass down the duct and out the pore. In

this way each of these coiled tubes absorbs wastes either from the body cavity or blood, transforms it into a secretion, and pours it into a duct and out the pore. The action is the same in every way as that of a single tubule and capsule of the kidney. (See Fig. 171.) The kidney of the clam is a coiled tube almost identical in its structural features with this nephridium of the earthworm, while the green glands of the crustacean are supposed to act in a similar manner.

Kidneys of insects.—These animals show a modification of the nephridium plan in that the tubules are arranged in a ring about the intestine. The ends of the tubes float in the body cavity of this animal, and absorb the waste from that, while their mouths open, not to the outside, but into the intestine, so that the nitrogen wastes pass out with the fæces through the anus.

Vertebrate Kidneys.—In all the vertebrates we find some form of kidney. These kidneys differ from a nephridium mainly in that they are a combination of many

FIG. 177.—Kidney tubes of an insect; *h*, kidney tubes; *a*, esophagus; *b*, crop; *c*, *d*, stomach; *e*, intestine; *f*, large intestine; *g*, anus.

secreting tubules under one covering, and they differ from that of man only in the manner in which the ducts terminate. For example, the ducts of the bird and reptile kidney open into an enlargement of the rectum (lower end of intestine) called the cloaca, and in these forms the anus is

the common external opening for the kidney and the intestine. In mammals the openings are for the most part distinct, the kidney opening being called the urinary opening. The development of the bladder is simply a device for preventing continuous secretion and permitting the animal to control this operation of removal from the body.

In general, then, whether we are dealing with the nephridium of the worm or the sweat gland of man, the means of disposal of waste are practically identical in action, and the kidneys are to be considered as simply a collection of separate excretory glands enormously increased in number to provide for the increase in waste, and having a common duct for removal.

XXV. THE NERVOUS SYSTEM.

STRUCTURE.

THE study of the preceding pages has taught us the part played by the various organs and structures in our body in the operation of tearing down and building up the human machine, and the manner in which these operations keep it in running shape. We have also seen that all these systems are interrelated, the work of one being directly dependent upon the activity of another, while in our life we are conscious of the fact that these many parts are all acting harmoniously and with comparatively little attention on our part. For example, in so simple an operation as walking, every system in our body is concerned. The muscles contract and expand, and in so doing produce motion, consume oxygen and nutriment, and give off wastes and carbon dioxide. The necessity for food must be met by a quickened flow of blood, while the increase in appetite warns us to supply the digestive system with more nourishment. The lungs must expand and contract to furnish the blood its oxygen, and the excretory system must increase its activity a little to dispose of the wastes. Even in our sleep these systems are all more or less active, and the needs of one part of the body can be met only by special effort on the part of all the others.

Manifestly such a harmonious action would be utterly impossible unless the organs had some means of communication by means of which the needs of one might be

PLATE IV.—Nervous system.

communicated to the others. Some central system of control must also exist, in order that the various activities of the body may be directed by us. The system of organs in our body by means of which this direction and control are exercised is spoken of collectively as the *nervous system*. The relation of this system to voluntary and involuntary actions and the structure of the parts will be considered in the following paragraphs.

General composition of the nervous system.—The nervous system may be said to consist of three parts: nerve centers, nerves, and nerve-end organs. The nerve centers in our bodies are the brain, the spinal cord, and little patches of nervous matter scattered in different parts of the body, called ganglia. The nerves arise from these centers, and with their branches penetrate all organs and tissues of the body. Finally, each nerve ends in some form of organ called an end organ. Some of these organs are very simple in structure; others, such as those at the ends of the sight and hearing nerves, are exceedingly complex.

For convenience, the cord and brain, together with the nerves which arise in them, are spoken of as the central nervous system, while the sets of ganglia on each side of the spinal column, with their nerve connections, are called the sympathetic system. All nervous centers are, however, in communication, and these two systems are separated in name for convenience of study, and not because of any true separation in the systems.

In general, then, the brain and cord may be likened to the "central" of a telephone system, the nerves to the wires, and the end organs to the individual telephones. Such an arrangement makes it necessary that all communi-

cations from one organ to another, or from one part of an organ to another, be made through the brain and cord.

If we enlarge our idea of this arrangement and conceive of the brain and cord as not only switching messages from one part of the body to another, but also of being able to modify the character of these messages, we shall have a true picture of the manner in which the brain and cord control and harmonize the actions of all parts of our bodies.

The nervous system of the frog. — A dissection of the nervous system of the frog is easily made, and since it is essentially similar in kind to ours, a study of this system forms a good introduction to the study of the human system.

As in our body, the brain is inclosed in the skull, and is continued downward as the spinal cord through the spinal cavities of the vertebræ. In this position the system is well protected from external pressure.

FIG. 178 — Nervous system of frog; *a*, cerebral hemispheres (fore-brain) *b*, olfactory lobes; *c*, eyes; *d*, midbrain; *e*, optic lobes; *f*, hindbrain (cerebellum); *g*, medulla; *h*, *i*, spinal cord; *I* to *X*, first to tenth pair of cranial nerves; *1* to *10*, spinal nerves.

If this bony covering be carefully removed so as not to injure the contents or its connections, a structure, such as shown in Figure 178 is seen.

The frog's brain.—The brain may be divided into three parts, called respectively, the forebrain, midbrain, and hind-brain. From these parts of the brain are given off ten pairs of nerves called cranial nerves. These nerves are made up of fibers called nerve fibers. Those cranial nerves whose fibers bring impulses or messages to the brain are called afferent or sensory nerves. Those whose fibers con-

FIG. 179—Frog brain; *I*, top view; *A*, olfactory nerves; *B*, olfactory lobes; *C*, *D*, forebrain; *E*, midbrain; *F*, optic lobes; *G*, hindbrain; *H*, medulla; *I*, cord; *J*, *K*, *L*, *M*, *N*, nerves; *II*, ventral view; *III*, lateral view.

vey impulses from the brain are called efferent or motor nerves. Several of these nerves contain both efferent and afferent fibers, and such are called mixed nerves.

The forebrain.—This consists of two elongated bodies lying side by side and called the cerebral hemispheres. Together, they constitute the cerebrum. At their extreme front are two swellings, from which a pair of cranial nerves extend to the nose. The fibers of these nerves are afferent, and carry sensory impulses from the nose to these lobes of the brain. They give to the animal the sense of smell. On this account the lobes of the brain from which these nerves arise are called the olfactory lobes, and the nerves are called the olfactory nerves. The olfactory are the only nerves connected directly with the cerebral hemispheres.

The midbrain.—Just back of the hemispheres is the midbrain. This is covered on the top and sides by two swellings called the optic lobes, so named because the sensory nerves from the eye enter them. They are the receiving points of sensations of light. The optic, or sight nerves (the second pair of cranial nerves) enter these lobes on the under side, and are crossed so that the nerve from the right eye enters the left lobe, and *vice versa*. These optic lobes also communicate with the forebrain by means of which we will speak later, but the optic nerves have no direct connection with any part of the brain except these lobes.

The third pair of cranial nerves also arises from the midbrain, and, unlike the first two pairs, it carries impulses from the brain instead of to it (that is, it is a motor or efferent nerve). The nerve fibers of this third pair control four of the six muscles which move the eyeballs and also some of the internal eye muscles. They are called the oculo-motor nerves.

The hindbrain.—The rest of the brain, which extends

backward to the beginning of the spinal cord, is called the hindbrain. It shows two regions. The transverse ridge just back of the optic lobes is called the cerebellum, while the wedge-shaped portion back of this, from which the spinal cord arises, is called the medulla oblongata. Seven pairs of cranial nerves arise from this medulla. The fourth pair, known as the trochlears, are very small, and carry motor impulses to one of the muscles which move the eyeballs. The fifth pair, or trigeminals, are very large. They belong to the class of mixed nerves. Some of their fibers bring sensory impulses from the teeth and face to the brain. Other fibers carry motor impulses to the muscles of the jaws and eyelids. The sixth pair, or abducents, carry motor impulses to a muscle of the eyeball. The seventh pair, like the fifth, are facial nerves of a mixed character, and carry impulses to and from certain parts of the face. In the higher animals and in man this nerve is purely a motor nerve, and controls the muscles which give expression to the face. The eighth pair, or auditory nerves, are sensory, and carry impulses of hearing from the ears. The ninth pair, or glossopharyngeal, supply the tongue and pharynx, and are mixed nerves. The tenth pair, or the vagus nerves, have the widest distribution of any. They are also mixed nerves, and their fibers supply the lungs and air passages, the heart and blood vessels, and the organs of digestion.

Finally, the centers of all these cranial nerves are in communication with parts of the forebrain, and the consciousness of sensations has its seat in that organ, though the nerves are not directly attached to it.

The spinal cord of the frog.—The cord of the frog, which is merely an extension of the medulla, is much shorter than that of man. From it are given off some ten pairs

of spinal nerves whose fibers are distributed to various parts of the body. . Figure 178 illustrates the method of branching of these nerves.

Comparison of the frog's spinal cord and brain with that of higher forms.—The spinal cord and brain are to be considered as one structure, the brain lobes and swellings being simply modifications of the more uniform spinal cord. These modifications become much more marked in the higher forms, and instead of lying directly behind one another the increase in size and the lack of space compels these brain lobes to become more or less twisted and doubled upon one another. If we keep the parts of the frog's brain in mind, however, we shall have no difficulty in recognizing the homologous parts of the human central system, as, aside from the folding and variation in size, they are almost identical, part for part.

Let us now examine the human brain with this comparison in mind, and see if we can identify the structures found there.¹

The human brain.—The brain of man is inclosed in that part of the skull called the cranium, and it fills this part completely. At the base of this cranium is an opening called the foramen magnum through which it is continued downward as the spinal cord. Various apertures in this bony box permit the entrance and escape of the cranial nerves.

Brain membranes.—When the bony plates of the cranium are removed, the brain is found to be covered with three membranes. The outermost (*the dura mater*) has

¹ The brain of the sheep is easily obtainable, and is so nearly identical with that of man in structure that it serves well for laboratory work. (See Ex. LXIII.)

a roughened outer surface which fits closely to the inner surface of the bones, and forms the inner periosteum of these bones. The brain side of this membrane is smooth,

FIG. 180 — Human brain in place; *C, C*, wrinkled surface of the cerebrum; *Cb*, cerebellum; *M.Ob.*, medulla; *B*, bodies of vertebræ, and *S*, their spinous processes inclosing the spinal cord.

and the whole membrane is thick and fibrous. It lines the whole cranial cavity, and ingrowths from this membrane support and invest all parts of the brain. At the foramen magnum it is continued as the outer coat of the cord. The blood vessels which supply this membrane are, for the most

part, distributed on the side next to the bones. In fact, their principal function is to supply nourishment to the bones, not to the brain.

A second membrane, called the *arachnoid*, fits against the smooth inner surface of the *dura mater*. This is a delicate, transparent, serous membrane. Beneath it, enveloping the brain itself, is a third, thin membrane, called the *pia mater*. This *pia mater* is composed of a fine network of small blood vessels held together by tissue, and these blood vessels supply the nourishment and remove the wastes of the brain itself. It is practically impossible to remove this membrane, as it covers all the folds of the brain.

These three membranes and the cranial bones furnish a very efficient protection for the delicate matter of the brain proper.

The brain proper.—Like the frog brain, it consists of three portions, forebrain, midbrain, and hindbrain, but these parts are so bent and doubled upon one another that they form a figure similar to an interrogation point (?). The two hemispheres of the forebrain occupy the bulk of the cranial cavity and correspond in position to the top, front, and part of the back of the curved portion of the interrogation point. The olfactory lobes are tucked away at the very front of the curve. In man, the forebrain constitutes nine tenths of the bulk of the entire brain.

The midbrain, with its optic lobes, is projected forward in a position corresponding to the curve at the top of the stem of the interrogation point. When viewed from the under side, this projection brings the optic lobes just back of the olfactory lobes.

The hindbrain has a much more developed cerebellum

than the frog's has, and this organ is here divided into two lobes by a longitudinal partition. Underneath the cerebellum, and extending downward, is the wedge-shaped medulla. In comparison with our figure the hindbrain occupies the position of the stem of the interrogation point, and if this were continued downward the extension would represent the spinal cord.

External features of the human brain.—Viewed from the top, the two hemispheres of the forebrain (the *cerebrum*) are seen to be much convoluted and folded, thus increasing their surface without increasing the space occupied. The two hemispheres are pressed closely together, the line of separation being marked by a fissure (the *longitudinal fissure*). Viewed from the bottom we can make out the continuation of this fissure back to the midbrain, and on each side of it appear the *olfactory lobes*. (See Fig. 182.) Just back of these lobes are the crossed optic nerves (the *optic chiasma*) which enter the optic lobes. Back of this chiasma is a band of white matter (the *pons varolii*) which connects the two lobes of the cerebellum and covers the top of the medulla. Underneath this pons, the medulla divides into two stalks (the *crura cerebri*) which spread out to form the cerebral hemispheres.

FIG. 181.—Cerebral hemispheres viewed from above.

The cerebellum shows a wrinkled surface different in appearance from the folded cerebrum. Between its lobes

FIG. 182 — Human brain viewed from below.

is seen the lower part of the medulla whose lower end narrows and is continued to form the spinal cord.

Along the under side can also be made out twelve pairs of cranial nerves, two more than in the frog's brain. The first ten pairs are connected to the same regions as those of the frog and have the same functions. The eleventh and twelfth arise from the medulla and are known respectively as the spinal accessory and hypoglossal. The spinal accessory is a motor nerve and supplies the muscles of the shoulders. The twelfth, or hypoglossal, is also motor and

supplies the muscles of the tongue. The first spinal nerve of the frog corresponds to this nerve in action.

Internal features of the human brain.—If we split the brain in two by a vertical cut, parallel to and a little to the left of the longitudinal fissure, we can make out the following parts on examining the right side. (See Fig. 184.) The fissure is seen to continue downward to a white fibrous body which binds the two hemispheres together. This is called the *corpus callosum*. This band curves in such a way as to inclose a space actually outside the brain and often called the “fifth” ventricle. This fifth ventricle is further inclosed by two thin membranes (the *septa lucida*). The forward curve of this connecting body below the fifth ventricle forms the roof of the midbrain, and this part is called the *fornix*. Under the fornix is the cavity of the midbrain or the third ventricle. This third ventricle communicates with cavities in the right and left hemispheres (the second or *lateral ventricles*) by two openings in the fore part (the *foramen Monroe*). In the figure, the opening into the right ventricle only appears. The midbrain cavity is also extended forward in a funnel-shaped tube called the *infundibulum*.

FIG. 183.—Origin of cranial nerves; *H*, hemispheres; *C. S.* corpora striatum; *P.* pineal body; *Pl.* pituitary body; *C. Q.* corpora quadrigemina; *Cb.* cerebellum; *Th.* optic thalamus; *M.* medulla; *I* to *XII*, the pairs of cranial nerves; *Sp. 1, 2*, spinal nerves.

The lateral ventricles in man are frequently named the *first* and *second* ventricles.

Extending from one side to the other of the third ventricle is a solid cord called the median commissure. The sides of the cavity are formed by the optic lobes, or *thalami*,

FIG. 184 — Vertical section of brain.

and its floor is formed by the *crura cerebri*, or stalks, of the cerebral hemispheres. We have already noted that these stalks are the forward extensions of the medulla. At the rear of the third ventricle is a passage called the *iter*, or aqueduct, leading to the fourth ventricle, a small cavity in the hindbrain. The top and sides of this iter are formed

by four little bodies called the *corpora quadrigemina*. Its floor is the medulla and the *pons Varolii*.

The fourth ventricle projects upward toward the cerebellum, and the internal structure of this organ shows a branching, treelike core of white matter (the *arbor vitæ*) surrounded on the outside by gray matter. The lobes of the cerebellum are only partly separated, the fissure dividing them being very shallow. The floor and sides of the fourth ventricle are formed by the medulla, and this cavity is continued backward as the *neural cavity* of the spinal cord.

From the above description it will be seen that the

FIG. 185 — Section of brain at right angles to longitudinal fissure.

brain is actually a hollow organ with a continuous cavity surrounded by various outgrowths or walls and

connecting all regions of the brain with the neural cavity of the spinal cord. In life this cavity (the ventricles) is filled with a liquid known as the *cerebro-spinal fluid* and the entire cavity is lined with a layer of ciliated epithelium. The third, fourth, and lateral ventricles are also supplied with blood from a network of blood vessels called the *choroid plexus*. Rupture and hemorrhage of the blood vessels of the lateral ventricles produces pressure upon the inside of the hemispheres and is one cause of apoplexy.

FIG. 186.—The human spinal cord and its branches.

If we cut slices at right angles to the longitudinal fissure in either hemisphere, we find that, like the cerebellum, the solid part of the cerebral hemispheres is composed of two kinds of matter, white and gray, surrounding a central cavity (the lateral ventricle). The outside of the hemispheres and folds is called the cortex and is composed of gray matter; the white matter forming the core of this part of the brain.

The human spinal cord. — This cord, which is an extension of the medulla, extends from the foramen magnum through the spinal cavities of the vertebrae to the articulation of the first and second lumbar vertebrae. From

here it narrows off into a slender filament which runs back to the end of the neural canal behind the sacrum. It is nearly cylindrical in form, being slightly flattened

dorso-ventrally, and has an average diameter of three quarters of an inch, with a length of about seventeen inches. It is largest in the neck region from the third to the first dorsal vertebræ, and there is a second expansion

FIG. 187.—Cross section of spinal cord; 1, anterior fissure; 2, posterior fissure, 3, central canal; 4, 5, commissures; 9, posterior root; 10, anterior root; a, e, gray matter. The outer layer is the pia mater, covering the white matter and sending into it blood vessels.

in the last dorsal vertebra. Externally, it is covered with three membranes like the brain: the dura mater, arachnoid, and pia mater.

The dura mater is not in close contact with the bones,

as in the brain, but is separated from their periosteum by a layer of fat and a network of blood vessels. It does not send partitions into the cord as it does into the brain.

The arachnoid is also not in close contact with the pia mater, but forms a loose bag around it. The pia mater closely invests the surface of the cord and, as in the brain, supplies it with blood.

The cord itself shows two fissures, a wide shallow anterior one and a narrow deep posterior one, thus dividing it into two right and left halves. The posterior fissure is not a true groove but rather a partition of connective tissue.

Each half of the cord gives off at intervals, from the top and bottom, nerve roots which unite at a short distance

FIG. 188 — Section of spinal cord showing nerve relations.

from the cord into common nerve trunks. (See Fig. 188.) These trunks or spinal nerves are arranged in pairs in the cervical and dorsal regions and emerge through openings between the vertebræ. At the lower end of the cord they are crowded together in parallel bundles. There are, in all, thirty one pairs of these nerves. Each spinal nerve trunk splits up into many smaller nerves and these into still smaller nerves which are distributed to various parts

of the limbs and trunk, thus bringing all parts of the body in communication with the cord. They are also connected on each side of the cord with the system of nerve centers called the *sympathetic ganglia*. (See Fig. 189.)

Internal structure of the cord. — If we section a portion of the cord in such a manner as to split a pair of spinal nerves (see Fig. 188), we can make out the following details: Like the brain, the cord is seen to consist of white and gray matter with this difference, that, in the cord, the white matter is on the outside. The inner gray matter is in the form of a letter H. Two horns of this H

FIG. 189 — Roots of a dorsal spinal nerve and its union with the sympathetic system; *c, c*, anterior fissure of cord; *a*, anterior root; *p*, posterior root with spinal ganglion; *s*, sympathetic, and *e*, its double connections with spinal nerve *a'* by a white and a gray filament.

project forward (the anterior cornua) and two backward (the posterior cornua). The central part, corresponding to the cross bar of the H, shows a small cavity in its center which is the continuation of the ventricle cavities of the brain. From the posterior and anterior cornua emerge nerve fibers which unite to form the roots of the

spinal nerves. The places where these roots emerge appear as four grooves on the surface of the cord. The posterior roots show a swelling just back of the point where they unite with the anterior fibers. This swelling is called the spinal ganglion. Experiment has shown that the anterior roots are composed mainly of efferent or motor fibers and convey motor impulses to the muscles, while the fibers of the posterior root are afferent or sensory and bring sensory impulses to the cord. The spinal nerves which contain both kinds of fibers are, therefore, mixed nerves.

Structure of a spinal nerve trunk and fiber. — If we cut a cross section of a spinal nerve we find it made up of many bundles of nerve fibers. The entire trunk is surrounded with a sheath of connective tissue. Inside this sheath, each bundle of fibers is wrapped in a sheath of connective tissue called the *perineurium*. We may remove this sheath and separate the bundles into the nerve fibers themselves.

In these ultimate nerve fibers, thus separated, we have the actual conductors of impulses, and our dissection shows that the spinal nerve may aptly be compared to the cables of a telephone station where each cable consists of bundles of separate wires, both bundles and wires being carefully insulated from one another.

Examination of a single nerve fiber shows it also to be composed of layers. In the middle is a core, or axis, composed of modified protoplasm. This is called the *axis cylinder*, and it is this cylinder which transmits the impulse. Surrounding this is a sheath called the *medullary* or *myelin sheath*. This sheath varies greatly in thickness in different fibers and is often segmented, forming the so-called "*nodes*

of *Ranvier*." Outside this myelin sheath is a thin membrane called the *neurilemma*, and comparable to the sarcolemma of the muscle fiber. Lying under the neurilemma are found nuclei, one for each segment of the medullary layer. These two outer layers of the fiber do not conduct impulses, and while their function has not been absolutely determined, they may be compared to the insulation which surrounds an electric wire. The axis, evidently, is the wire. In some nerve fibers the medullary layer is lacking, the axis being surrounded by the neurilemma only.

All nerve fibers originate either in the nervous system or in the sympathetic ganglia. Those from the former are all medullated; some of those of the latter are non-medullated. All fibers admit also of classification as afferent or efferent, since, in a given fiber, the direction in which the impulse travels is always the same. Mixed nerves, therefore, are composite bundles of afferent and efferent fibers.

The words afferent and efferent simply indicate the direction in which the impulse travels. Owing to the fact that efferent fibers all carry impulses which produce motion, the words efferent and motor are equivalent in describing a given fiber, while for the same reason the words afferent and sensory may be interchangeably applied to fibers which convey sensory impulses.

Finally, the two classes of fibers may be subdivided again on the basis of the effect of their impulses. For

FIG. 190—Section of a nerve.

example, a motor fiber transmits an impulse which may retard or increase a motion, while a sensory fiber transmits impulses which may increase or decrease sensation. Those fibers, either motor or sensory, which excite to greater motion or sensation, are called excitatory. Those which

decrease motion or sensation are called inhibitory. This inhibition or excitation is supposed to be determined, not by the nature of the impulse, but by the tissue in which the fiber ends.

The plexus.—The fact just noted, that the spinal nerves are composed of bundles of fibers, explains how it is possible for them to split up into smaller nerves and ultimately into separate fibers, just as we may unravel the strands of a rope. In many regions of the body some of these unravelled

FIG. 191—The solar and hypogastric sympathetic plexuses.

nerve fibers from several spinal or ganglionic nerves interlace to form a meshwork, and such networks are called plexuses. In this fusion, new combinations of fibers are often made, and nerves extending from a plexus may contain fibers from several different spinal or ganglionic nerves. Evidently the stimulation of such a nerve will result in messages to several different parts of the central system, and *vice versa*. For example, the nerves supplying the legs arise from large plexuses, and are thus supplied

with fibers from many different parts of the central system. The advantage of such a combination is seen in the manner in which a single nerve is able to coördinate very complex movements, the messages from the different centers of control being concentrated, as it were, by this arrangement.

One of the largest plexuses is located just above the pit of the stomach, and is called the solar plexus because smaller plexuses radiate from it like the rays of the sun. A blow on this area paralyzes many regions, and may even produce death.

Cell theory of nerve structure; neura.—

The presence of nuclei under the neurilemma at once suggests the idea that nerve fibers bear some relation to cells. Examination of the gray matter of the cord or brain in which these fibers originate gives us further information on this point, and to the relations thus established is due the present theory of nerve structure and action.

Preparations of gray matter show it to be rich in nucleated cells of irregular outline. (See Fig. 192.) From the angles of this outline project fine branches of protoplasm, which may subdivide again and again to form treelike growths, or which may continue some distance without branching, in the form of a cylinder. The branching outgrowths are called dendrites, while the cylinders are called axis cylinders, or *axones*.

FIG. 192—Nerve cells from the cortex of the cerebral hemispheres; *Ax* axis cylinder.

Nerve cells, wherever found, agree in this general structure. In some the production of dendrites is much less than in others. Some also have one axon only, while others may have from two to several. Those cells with one

FIG. 193.—Scheme of a neuron; *a*, free axis cylinder; *b*, axis cylinder surrounded by neurilemma alone; *c*, axis cylinder surrounded by medullary sheath alone; *d*, axis cylinder surrounded by the sheath and neurilemma and divided into segments (by constrictions called the nodes of Ranvier).

axon are called unipolar, with two, bipolar, and with more than two, multipolar. If we trace the course of one of these axones we find that it may extend a long distance from the parent cell with or without branching, and, what is most important, such axones are the axis cylinders of the nerve fibers.

We see then that a nerve fiber is to be considered as an elongated, axis cylinder, or axon, which has become covered with one or more protective sheaths. When the axis cylinder branches, these branches are called collaterals, while the terminal branches in which they all end are called the terminal brushes. A nerve cell with its dendrites, axones, sheaths, collaterals, and terminal brushes, is spoken of collectively as a

neuron. Such a unipolar neuron is shown in Figure. 193.

Theory of neuron action.—The action of an amoeba

when touched with a needle point demonstrates that protoplasm is capable of responding to stimulation. Since the result is the same no matter where the protoplasm is touched, it follows that it is also capable of conducting impulses thus set up. In other words, responses to certain forms of stimulation and conduction of impulses is a property of protoplasm. When we examine the cytoplasm of a nerve cell we find it to be made up of two kinds of protoplasm. One kind resembles closely that of the amoeba. The other kind seems to be in the form of threadlike fibrils. The axon protoplasm is almost purely of the fibrillar kind. This fibrillar protoplasm is found to be able to transmit impulses much more rapidly than ordinary protoplasm. In other words, it is protoplasm with a highly developed power of conduction. The discovery of this structural peculiarity of nerve cells enables us to think of them as cells whose protoplasm has been specially developed for conducting impulses.

Keeping in mind this fact, and also the structure relations in a neuron, the formulation of the theory of neuron action may be expressed in the following laws:

First law. An impulse may be started wherever there is protoplasm, but externally stimulated impulses have their origin usually in the terminal brushes of the axones or in the dendrites, while the internally stimulated impulses arise in the body of the nerve cells. In general, then, the nerve fiber or axon is not a source of impulses, though in cases of inflammation of the sheaths it may become so.

Second law. An impulse started in any part of the protoplasm of a neuron is conducted by the fibrillar protoplasm to every other part of the neuron protoplasm,

Thus, an impulse started at the end of an axon is transmitted directly to the protoplasm of the cell body and through that to the extreme ends of the dendrites and any other axones that may arise from that cell body. Similarly, an impulse arising in the dendrites or cell body is likewise transmitted over the axones to their terminations. Bearing in mind that impulses usually originate in one of three places, the cell body, the dendrites, or the terminations of an axon, it will follow that the principal function of the nerve fiber is conduction. This characteristic is still further evidenced by the fibrillar character of the axon protoplasm.

When two points are connected by two neuria the arrangement is as follows: The dendrites of one neuron are so placed as to be brought in contact with the terminations of the axon of the second. In this relation the course of the impulse takes one of two directions, and this direction may be formulated in the third law.

Third law. An impulse which originates in the terminations of the axon of one neuron, and has been conducted to the dendrites of the nerve cell of that neuron, may pass from those dendrites into the terminals of the axon of the second neuron, and thus on to the dendrites in the final cell of the series. Similarly, an impulse originating in the cell body or dendrites of one neuron may pass, by way of the axon terminations, into the dendrites of the connecting neuron, and thus on to the final axon terminations of the series.

These three laws enable us to explain the method of impulse transmission in all forms of nerve action; while neuria may differ in the number of axones, in the length of the axones, and in location, the laws of action given above

hold true for all classes. In connection with the third law of action it may be noted that whenever there is a passage of impulse from one neuron to another there is at this point a delay in the transmission, and sometimes several impulses must reach a cell before it can overcome this resistance between dendrites and axon contacts and discharge into the next cell. We may compare this delay to the effect of placing a small barrier in the path of a stream. The effect is not to stop the stream, but to check it until the volume of water is great enough to push aside the barrier. The normal rate of nerve discharge is about ten impulses per second.

It will be noted finally that the above laws give no explanation of the manner in which nerve impulses originate. The manner in which externally stimulated impulses originate will be considered in our study of the sensory end organs. The manner in which impulses originate in the cell body is still unknown. It is believed, however, that just as oxidation of fuel and metabolism account for the contraction of muscle protoplasm, so these nerve impulses set up in the nerve cells result from certain chemical changes in the composition of the cell protoplasm. The fact that nerve cells require food and become fatigued by overwork justifies this belief, but the exact nature of the metabolic changes involved is unknown.

Distribution of neura.—In the central nervous system, the cell bodies of the neura which originate there are found mainly in the gray matter of the cord and in the cortex of the brain. The axones of these cells may extend far beyond the limits of this system, or again they may be confined within the central system. The spinal and cranial nerves represent bundles of axones which

originate from these central system cells and connect them with more or less distant parts of the body. Many of the axones found in the white matter of the cord and brain, however, never leave the central system, and the function of these is to connect the brain cells with the cells of the cord or with those of other parts of the brain. In general, then, the gray matter of the central system consists mainly of nerve cells, while the white matter is largely composed of nerve fibers. Each cell gives out from one to several axones, and the entire nervous system may be thought of as very complicated network of interconnected neura.

XXVI. THE NERVOUS SYSTEM (*continued*).

FUNCTION.

FROM the nervous system we gain control of the organs of the body, and to its action we owe our power to adjust ourselves to our surroundings in general and to maintain relations with other human beings and objects. In securing to us this adjustment to our environment, the nervous system manifests many forms of activity. For example, the interconnection of the fibers permits the transmission of messages of control and supervision. The end organs of various forms give rise to special kinds of impulses, and when these are transmitted to the centers, the structures found there are able to translate them into sensations and thus furnish information in regard to our surroundings which is of vital importance in determining the character of the control impulses. Finally, the nerve centers are the origin of those mental operations which we call will, emotion, thought, etc. In short, the very individuality of a man is a function of his nervous system.

The explanation of the many actions which we group under the common name of mind activities is properly the province of a science called psychology. In the study of the physiology of nerve action we are concerned only with the changes which take place in the nervous elements of structure in giving rise to these operations and the manner in which these changes occur. Owing to the

delicacy of the nervous structure and the difficulties in the way of experiment our physiological knowledge of the nervous system is extremely limited, and, in fact, much has still to be learned of the physiology of nerves and nerve cells. In the following pages we shall simply try to indicate a few of the facts which have been determined up to the present time, the manner in which such knowledge has been acquired, and its bearing upon our common activities.

Action, voluntary and reflex.—The action of every organ of our body is dependent upon nerve control. The simplest division that can be made of the actions of the body separates them into two classes. Those which require the exercise of the will and are associated with consciousness are called voluntary actions. Those which take place without any effort of will and may or may not be accompanied by consciousness we call reflex. All such actions as the picking up of objects with the fingers and the avoiding of obstacles in walking, in short, all actions which involve a conscious choice on our part belong to the voluntary type. All such actions as the regulation of the beating of the heart, the withdrawal of the hand from a hot object, etc., in which the action takes place without any consciousness of choice on our part, belong to the class of reflexes.

The relation which consciousness may bear to such actions is brought out in the two examples of reflex action cited. In the beating of the heart consciousness is entirely lacking; while in the withdrawal of the burned hand we may be conscious of the pain and the movement, but ordinarily the movement has occurred before the pain is felt, and hence, in this case, while conscious-

ness may accompany the action, it does not determine the action. Many actions which are at first voluntary may become reflex by repetition; and, in fact, one of the fundamental features of education is concerned in making reflex or habitual, actions which in the child are voluntary and require effort of will on his part.

The reflex arc.—The spinal cord is the seat of most of the reflex actions of the organs of the trunk and limbs.

FIG. 194. — A spinal reflex.

The relation of the nerve elements in simple reflexes which have their origin in the cord, is shown in Figure 194, and the relation there shown is typical of all reflex actions. In this figure it will be seen that the posterior root of the spinal nerve has on its surface a swelling which was called the spinal ganglion. This ganglion contains many bipolar nerve cells whose axones extend in different directions. One axon passes out and forms one of the sensory fibers of the spinal nerve. It terminates in some form of an end organ. The other terminates in a brush in the gray matter of the cord. Likewise, in the anterior cornua of the cord is found a unipolar nerve cell whose dendrites are in

contact with the terminal brush of the sensory fiber. The axon of this nerve cell passes out through the anterior root and becomes one of the motor fibers of the spinal nerve. It terminates in a muscle.

If, now, the sensory end organ of such a system is stimulated, as when the finger is brought in contact with a hot stove, the impulse passes up over the axon or fiber to the ganglion cell and through that is transmitted to the terminal brush of the other axon in the cord. From this brush it passes into the dendrites of the motor cell, and this cell in consequence discharges an impulse over its axon to the muscle and causes that to contract.

In this simplest form of reflex action only two neura are involved; one neuron conveying the sensory impulse and the other the motor impulse. Most of our reflexes are

FIG. 195—Diagram showing how one sensory nerve may stimulate several motor nerves in a simple reflex.

more complex than this, but their complexity consists in the fact that they involve more than two neura and not in any essential difference in action. The manner in which a single sensory impulse may stimulate several motor fibers is seen in Figure 195.

Here the sensory axon

splits up in the cord into collaterals, and each collateral discharges through a different motor cell. Such an arrangement explains how several muscles can be reflexly thrown into action by a single sensory impulse. It must be noted,

however, that the production of the impulses is essentially the same as in the simpler case.

The reflex arcs of the cranial nerves are similar in action to those of the cord, and involve sensory and motor neurones. Inasmuch as in this action the impulse travels in a circuit, the name of reflex arc has been given to such arrangements as above.

The voluntary arc.—When we come to examine what takes place in a voluntary action we find that the process

FIG. 196—The voluntary arc.

is not very different from that of the reflex action, except in the time required for transmission. Figure 196 shows a typical voluntary arc. In such a circuit the path of the

impulse is as follows: End organ to cord, cord to brain centers by connecting neura, brain centers to other brain centers by other connecting neura, and finally back from the last brain center to the motor cell in the cord and over its axon as a motor impulse to the muscles. In such a circuit it often happens that the cord motor cell is so near the sensory fiber brush that a division of the discharge is made, part of the impulse passing over the long circuit noted above, while the other part passes directly over to the motor cell and thus establishes a short-circuited reflex. In traveling the long circuit to the brain no motor discharge can take place until the impulse is returned from that region. Such a circuit is characteristic of all deliberate actions. In the case of the division of the impulse, while part of the impulse may continue on to the brain and produce consciousness, the short circuited impulse is able to produce the action perhaps even before consciousness is aroused. In the case of the burned finger the importance of this latter arrangement is evident in regard to speed of action. This short circuiting also illustrates how a voluntary action may become reflex, for if gradually the bulk of the discharge follows the shorter circuit the less necessary becomes the long circuits, and it may ultimately be abandoned entirely.

In general, then, a voluntary circuit differs from a reflex circuit only in the length and in the number of neura involved. The tracing of all these connecting neura and their distribution in the central nervous system forms a part of the study of the nervous system in which much work still remains to be done. Almost nothing is known as yet about the character of the changes which take place in the brain cells and the manner in which past experiences can be stored and modify the nervous discharges. By

experimenting with lower animals, and by the study of human beings whose nerve tracts have become diseased in part, it has been found possible to locate the tracts where sensory impulses are translated into consciousness, and also the points of origin of control impulses for different organs.

Localization of functions in the central nervous system.—Such experimental studies as noted above have demonstrated that a sensory impulse produces no conscious sensation unless it reaches a certain area of the brain, and if this area is destroyed and the transmitting fiber with its end organ preserved intact, no sensation results. Similarly, it has been found that certain organs and parts of the body are under the control of definite areas of the brain and cord, and these areas must be in a healthy condition or the action of these organs is seriously interfered with. The determination of these areas of sensation and control is called the localization of functions, and the method by which results have been obtained is largely through experiment with lower animals.

Spinal cord.—If we cut the spinal cord of the frog just back of the medulla, the frog will continue to live and the heart will continue to beat, but he ceases to breathe or swallow. The animal lies flat on his belly and loses all power of voluntary movement or conscious sensation. He is still able, however, to respond to external stimuli with definite movements. Thus, if we irritate the skin, the animal will make definite and skilful movements to remove the irritating body; if we pinch the toe, the foot is withdrawn. From these and similar experiments it has been conclusively demonstrated that the cord is the seat of those reflex movements which concern responses to

externally applied stimuli, and is also the path of sensory impulses to the brain. In higher animals similarly treated this reflex response is less apparent, and indicates that in such animals the reflex action of the cord is more dependent upon control from higher parts of the system.

Cerebellum.—Experiments upon the cerebellum of the pigeon develop the following facts concerning this organ. When all the cerebellum is removed the animal can still execute voluntary movements, but these movements instead of being definite and coördinated are sprawling, uncertain, and throw the animal into all sorts of grotesque positions, while it is totally unable to fly. From these experiments and others upon higher forms it appears that while the cerebellum does not originate motor impulses it does have the power to regulate and coördinate these impulses, and is thus responsible for orderly movement. An animal still retains sensation when the cerebellum is sliced away.

Cerebrum.—In the cortex of the cerebral hemispheres is the seat of all the psychical reactions which we include under the name of intelligence, memory, will, and the emotions. It is also the seat of conscious sensation, and in it originate the impulses which produce voluntary movements. Involuntary movements or reflexes may, however, take place when these hemispheres are removed. A pigeon whose hemispheres are removed becomes a stupid, drowsy creature, which, when thrown into the air, recovers its equilibrium and flies, and it may even be made to feed by properly applied external stimuli. If starved, however, it becomes restless and pecks aimlessly at the ground, but does not seem to be able to locate or recognize food for itself. In other words, it has lost all those responses

which depend upon memory of past experiences, that is, intelligent responses. It shows no signs of fear or pleasure, and while it responds to sensory stimuli, these responses are purely reflex and not accompanied by true consciousness.

History of cerebral localization.—When it was established that the cerebral cortex was the seat of psychical activities, the question at once arose as to whether all parts of the cortex were equally capable of exercising these faculties, or whether different parts controlled specific faculties. The idea of separate areas for separate functions was first presented by Franz Joseph Gall, who divided the cortex into areas of special activity, and put forth the theory that the more developed any mental quality was the larger and more prominent became the cortical area which produced it. Further, since the cortex fits closely to the cranium, the relative prominence would be indicated externally by the shape of the skull. From this position arose the practice of phrenology or the determination of mental qualities by examination of cranial prominences. Opposed to the position of Gall was the view of Flourens, who held that all parts of the cortex were capable of producing all kinds of mental qualities, and that when one part was removed the remaining parts supplied the qualities originally centering in that part. Without going into this controversy we may present the modern views on the subject as follows:

First. The general view that functions are localized has been definitely established, and it is possible to map the cortex of the brain and thus indicate areas which, when stimulated, will produce definite actions.

Second. The idea of Gall that the more marked the

development of a function the larger and more prominent the area has been proved false.

Third. The interconnection of areas has been shown to be so intimate that, although normally specific areas control specific functions, an injury to one part may affect all the others.

In other words, the cerebrum is composed of many organs intimately associated with one another and interdependent. In general, the left hemisphere controls the

FIG. 197 — Diagrams illustrating localization of functions in the cerebrum; *I*, outer surface of left hemisphere; *II*, inner surface of same. Motor areas are shaded; *Cc*, corpus callosum; *Fr*, frontal lobe; *Oc*, occipital lobe; *Pa*, parietal lobe; *Ro*, fissure of Rolando; *Sy*, Sylvian fissure; *Te*, temporal lobe.

right side of the body and *vice versa*. The chief motor and sensory areas of the hemispheres are shown in Figure 197.

Functions of the medulla.—This is the center of those reflexes which control the respiratory and circulatory organs. When a cut is made anterior to the medulla the animal continues to breathe and the heart to beat for some time after the operation. If, however, the cut is made at the point where the medulla joins the cord or it is destroyed entirely, heart beat and breathing soon cease and rapid death follows. Other important reflexes

are also located in this organ which is the origin of many of the cranial nerves.

Functions of the sympathetic system.—This name was originally given to the chain of ganglia extending along each side of the spinal column under the mistaken idea that it was a pathway for all the so-called sympathetic (reflex) actions of distant organs. This system was supposed to arise from the brain by branches connected with the fifth and sixth cranial nerves. It is now known that this system consists merely of a collection of ganglia or nerve cells which are connected with one another and with the spinal nerves. While it is customary to limit the term to the ganglia on each side of the spinal column, it actually includes many other ganglia with or without names which lie in different regions of the body. According to Professor Langley, the efferent fibers from the sympathetic and related ganglia supply the unstriped muscle tissues, the cardiac muscles and the glands, in short, the muscles of the involuntary organs of the body, such as the walls of the intestines. He has given the name of *autonomic* to this system to indicate that these fibers are to a certain extent independent of the central nervous system. All the sympathetic ganglia are connected with the central nervous system by medullated fibers from the spinal or cranial nerves, but these fibers do not pass directly to the unstriped muscle tissue. On the contrary, they transfer their impulses to a ganglion cell and the impulse passes out through a fiber (autonomic fiber) of this ganglion cell which is usually non-medullated.

The connection of these fibers with the spinal cord is illustrated in Figure 189. Unlike the fibers which supply the striated muscles, practically the entire set of autonomic

fibers is removed from the control of the will. Under ordinary conditions they are probably excited reflexly, but the course of the afferent fibers is at present little understood. Most of these reflexes are also unconscious actions, and include such movements as the dilation and contraction of the arteries, the movements of the intestines, and the secretory activities of the digestive glands. The entire field of action and method is still very little understood.

HYGIENE OF THE NERVOUS SYSTEM.

As has been stated, the manner in which nerve cells generate impulses or give rise to psychical reactions is unknown. We do know, however, that generation of impulses and mental activity is in some way dependent upon the metabolism of the nerve cells. From this, and the fact that nerve cells, like all other cells, are made of protoplasm, it follows that lack of nourishment or of air is bound to affect their activity, overwork to fatigue them, and rest to restore them. Comparison of the nerve cells of a child with those of an adult show that the latter are much more branched and more intimately connected than those of the child, which is another indication of the relation between increased nervous action and metabolism.

Nerve food.—It used to be thought that nerve cells required special kind of foods for their development, notably, phosphorus; and fish, which are rich in phosphorus, were supposed to constitute an ideal brain food. It is now known that their protoplasm is built up of identically the same sort of materials as are used by other cells, and hence a food which is nourishing to the laborer is equally so to the brain worker. The important point to be kept in mind is that there must be a sufficient quantity of food, and that the

organs of digestion and circulation, etc., must be kept in good order for its preparation and distribution. In short, whatever tends to produce a healthy body also tends, indirectly, to maintain a good condition in the nervous system. Exercise, therefore, is as valuable to the brain worker as it is to the laborer, and perhaps even more so, in that the latter is apt to neglect muscular activity, while the very work of the laborer tends to keep his system in order.

Air supply.—The chemical changes which take place in the nerve cells are essentially similar to those in the muscle cells, in that the production of energy or action involves the oxidation of cell substance and food. It follows, therefore that not only must the nerve cells receive food but also a plentiful supply of oxygen. The brain responds very quickly to a lack of air by a feeling of dullness and headache, and it is extremely important that our school and living rooms be well ventilated. Exercise that tends to develop the lung power is likewise valuable.

Rest.—Like the muscle cells, again, the nerve cells accumulate wastes and consume food by continued activity. It is imperative, therefore, that they be allowed periods of rest when the supply of food may be renewed and the excess of waste removed. Sometimes, as with the muscles, a change of work will permit one set of cells to rest while another is working, and thus a change of occupation is a rest. But no person can work continuously and maintain a healthy nervous system, and for that reason sleep is nature's greatest remedy for tired nerves. Children whose nerve cells are growing should have plenty of sleep in order that their nerve cells may store up sufficient food in these rest periods to meet the daily needs, and at the same time permit of growth and branching. On this account adults whose cells have attained their normal size need less sleep than children.

Nerve cells are like muscle cells in another particular, in that they degenerate by disuse and lose their irritability. The person who is too careful of overtasking his nervous system may carry this care to excess and lose his mental power entirely. It is this need of constant nerve exercise which gives especial value to a general education, since the variety of subjects in such a course insures the normal development of all parts of the system.

Diseases of the nervous system. — *Nerves.* The sheaths which inclose the axis cylinder may become the seat of inflammation. The result is intense pain and various forms of nervous disease, according to the extent and nature of the inflammation. Among diseases of this nature are included all forms of neuritis, toothache, etc.

Spinal cord. — Injuries to the spinal cord or the introduction of certain forms of bacteria may result in the destruction of the whole or part of this organ. The result is paralysis of the parts supplied by the affected portion. Such diseases are usually slow in action and often incurable.

Brain. — Anything which affects the nerve cells of the brain, such as pressure, narcotics, worry, lack of work or sleep, etc., is bound to result in disturbed mental action. Such influences may result in sleeplessness, hysteria, lack of self-control, irritability, and even in total loss of control or insanity. Interference with the blood supply and pressure may produce fits or apoplexy. In short, so sensitive is the brain to external influences that its encasement in the bony cranium is none too strong a protection for this delicate structure and any interference with its structure is attended with serious results.

ALCOHOL AND THE NERVOUS SYSTEM.

Stages in intoxication. — It is a fact known to everyone that alcohol in any quantity affects the nervous system almost immediately. It is noticeable that moderate quantities have apparently a stimulating effect upon the various centers of speech, thought, and action. The winebibber becomes exhilarated, his speech flows more rapidly, his wit appears keener, and he is conscious of a sense of freedom from restraint. If the amount of alcohol is increased continuously, this stage of apparent stimulation is soon followed by excessive exhilaration. The subject becomes still more talkative and his freedom from restraint manifests itself

in excesses of various kinds. He laughs immoderately at remarks only slightly humorous and in many ways shows that his powers of judgment are becoming dulled. Gradually this dullness spreads. His speech becomes thick and uncertain and a stage of dullness and drowsiness of all his mental faculties follows. He may still be able to walk, but his inclination is to be quiet, and, if permitted to do so, he passes rapidly into a stupid, drunken slumber. Finally, the loss of control over his speech spreads to include the centers of voluntary movement. He loses control of his legs and is no longer able to walk, staggering blindly when he attempts to do so. Such a condition represents what are usually the final stages of acute intoxication or drunkenness. If, however, the doses of alcohol are still continued they reach a point when the involuntary centers which control the breathing and movements of the heart become affected, and in this case death of course must result.

The problem of the physiologist.—The problem of the physiologist is to account for these well-known effects and to tell exactly what is the action of the alcohol at each step of the process. All are agreed that alcohol in large or continued doses ultimately dulls or paralyses the nervous centers, and that this narcotic effect is felt first in the higher centers. After these centers are dulled the paralysis gradually spreads until it includes the motor centers and ultimately the vital processes, such as breathing and heart beat. Thus, in the description given, it is easy to see that the centers of judgment are the first to become dulled, and that the effect of this is to remove the speech and motor centers from control so that they seem to be stimulated. That this is not a true stimulation, but the first step in

paralysis, is seen in the speed with which these centers become dulled, while the tendency to attack the higher centers first is shown by the fact that speech is lost before control of movements.

In regard to the effect of alcohol in small doses there are at present two different opinions. One school of physiologists holds that alcohol in small doses is a true brain stimulant, and that the exhilaration felt as a result of such doses is accompanied by increased keenness of judgment, and that, while temporary in its effect, this effect is not due to loss of control by paralysis of higher centers.

The other school claims that alcohol is never a stimulant, but always a narcotic, whether in large or small doses. They explain the evident exhilaration which follows small doses as due to the same causes as the excessive exhilaration, namely, that it results from a partial paralysis of the centers of judgment and control. They say that, by removing the inhibiting influences from the speech and other centers, such paralysis inevitably causes in these centers a freedom of restraint which manifests itself in apparent stimulation. In short, this school claims that alcohol from its entry into the system is never a stimulant, but always a narcotic in its effect upon the nervous system as a whole.

The determination of the true position in this regard must come as a result of experiment, and the results of experiment have so far been extremely conflicting. Not enough has been done as yet to settle the question for either side, and all that can be said at present is that by far the larger number of physiologists take the view that alcohol is always a narcotic, even in small amounts.

Alcohol and time reactions.—Many experiments have been made to determine the effect of alcohol on the time

required for a nervous impulse to travel over the fibers. All such experiments require the subject to perform a certain muscular action upon receiving a certain sensory stimulus, and the measure of the time which elapses between the receipt of the sensory impulse and the performance of the action is taken to indicate the rate of transmission of the impulse — the “reaction time” as it is called. By comparing the reaction time of a person under normal conditions with the time when under the influence of definite amounts of alcohol the effect of alcohol can be inferred.

The results of such experiments are not absolutely conclusive in regard to the effects of small amounts of alcohol. All, however, tend to show that larger doses of alcohol lengthen the reaction time for all mental processes. The importance of this fact has led all corporations employing men in places where quickness of action, clearness of perception, close attention to duty, and the like are required, to forbid absolutely the indulgence in alcoholic drinks while performing such duties. Such actions on the part of corporations are significant warnings to us of the danger of this alcohol habit and its tendency to destroy efficiency.

Alcohol and brain workers.—It follows directly from the preceding that alcohol is not of advantage to the brain worker, and leads to confusion of judgment. One unfortunate effect of its use is that in dulling certain control centers it often produces in the indulger the delusion that he is actually keener in perception, when actually this apparent keenness may be only an indication of impaired powers.

Alcohol and diseases of the nervous system.—Continued use of alcohol lowers the resistance of the body to disease, and it is a well-known fact that hard drinkers often suc-

cumb to diseases from which a normal person would easily recover.

Many cases of insanity are reported as due to alcohol, but while it is undoubtedly true that the use of alcohol is in many cases indirectly the cause, the exact relation to this mental disease is not yet definitely known. The only form of mental disturbance which is a direct result of alcoholic indulgence is the violent disturbance known as delirium tremens.

It is also a well-known fact that nervous weakness can be transmitted by heredity, and thus children of drinking parents often develop mental disease which is due to this inherited weakness.

XXVII. THE NERVOUS SYSTEM IN THE LOWER ANIMALS.¹

NERVOUS tissue, wherever it exists, is in the form of nerve cells and fibers, as in man. Frequently the nerve cells are collected in masses which are called ganglia, and

FIG. 198—A nerve ganglion. The round bodies are nucleated nerve cells; the fibers are nerves.

in this chapter the use of the word ganglion may be taken to mean a collection of nerve cells. The fibers are always developed as processes of the nerve cells.

Protozoans and cœlenterates.—The one-celled animals can evidently have no true nervous system. Their protoplasm, however, is sensitive to stimuli and conducts impulses. Such a condition is interesting mainly as showing the relation of ordinary protoplasm to the specialized protoplasm of the nerve cell and as indicating the origin of the nervous system. The first appearance of nerve cells and

¹ See Footnote, p. 146, Chapter X.

fibers is found in the cœlenterates. In the hydra are certain scattered cells which are partly muscular and partly nervous in function. In the jellyfishes we find a network of fibers and cells scattered over the disk. The general purpose of such systems seems to be to give sensitiveness to the animals and thus aid them in securing food.

Echinoderms.—In the starfish and its allies we have a true system. In these animals we find a ring of five ganglia surrounding the mouth and connected by a nerve cord. Each ganglion gives rise to radial nerves which extend

FIG. 199.—Diagram of nervous system of starfish; *r*, nervous ring around mouth; *n*, radial nerves to each arm, ending in the eye.

along the arm opposite the ganglion. The peculiarity of this system is that it actually consists of three systems, each of which has its own combination of cells and fibers and supplies a special part of the body. These three systems are absolutely independent of one another in action. We may compare the nervous control of the starfish to a city with three absolutely independent telephone companies,

each company with its own “central” and wires. Such an arrangement is absolutely unknown in any other group of animals.

Mollusks.—The nervous system of such animals as the clam and its allies consists of several ganglia located in different parts of the body and interconnected by nerve trunks. Each ganglion controls special parts of the body, and this arrangement is spoken of as a scattered system of control. The arrangement in the clam consists of two

compound ganglia (cerebro-pleural) located on the right and left sides of the esophagus and connected by a nerve cord called a commissure. Each of these ganglia actually consists of two cell masses. The two cerebral masses give off nerves which control the head parts while the pleural masses control the neighboring parts of the body. Each compound ganglion gives off also two nerve trunks which pass backward along the same side of the body as the ganglion from which they originate. The right and left trunks of one pair unite with a ganglion (*visceral*) near the siphon end of the animal. The right and left trunks of the other pair unite with a ganglion in the foot, called the *pedal* ganglion. The pedal ganglion sends out fibers which direct the movements of the foot, while the visceral ganglion, through its fibers, controls the internal organs and the parts near it. In such a system it is evident that we have control stations located near their seat of control and, unlike those of the starfish, all are interconnected into one system by the nerve trunks. We may compare such a system with a city telephone system in which each district has its own "central" but all the centrals are connected.

In the snail we find an interesting modification of the clam system. Here the main ganglia are collected in a ring about the esophagus, while additional ganglia are found scattered through the body. In the cuttlefish this

FIG. 200 — Nervous system of clam; *c*, cerebro-pleural ganglia; *c'*, cerebral commissure; *p*, pedal ganglion; *ps*, visceral ganglion; *p'* and *ps'*, nerve trunks.

ring of ganglia is actually inclosed in a cartilage box, thus giving it a still closer resemblance to a brain.

Worms.—In the worms we find two main ganglia usually located in the head end and united by a commissure. In the simpler worm forms these ganglia are the only ones in the body, and fibers from them supply all parts of the body.

In the earthworm these two ganglia are represented by a single two-lobed mass located just above the esophagus and called on this account the supra-esophageal

FIG. 201 — Nervous system of earthworm; *g*, ganglion; *v.n.* ventral nerve cord.

ganglion. From this compound mass or “brain” a ring of cord encircling the esophagus unites it with a ganglion mass under the esophagus called the sub-esophageal ganglion. Finally, a double cord extends from this latter ganglion along the ventral side of the body. This cord shows a series of small ganglia at intervals corresponding to segments. Each of these small ganglia is the source of motor and sensory fibers which supply that particular segment, and these segment ganglia are the seat of the reflex actions. It is not difficult, then, to compare such a system with the one in man. The supra-esophageal ganglion may be compared to the brain, the ventral cord to the spinal cord, and the smaller ganglia to the centers of reflex actions and sources of spinal nerves in the cord.

This brain ganglion and cord constitute a central system in distinction to the fibers which arise from the small ganglia and which correspond to our peripheral system.

Arthropods.—In the lower arthropods, such as the lower forms of crustaceans, and the wormlike larvæ of insects, the nervous system is practically identical with that of the earthworm. In the higher arthropods such as the higher crustaceans and insects, the plan is the same, but many of the ganglia are fused together in the head and thorax making this end of the system much more brainlike in structure. This head mass also gives nerves which supply the eyes and antennæ much as our cranial nerves are given off from the brain. The ventral cord shows fewer ganglia, and each ganglion furnishes motor and sensory fibers to larger areas than in the worms.

In short, while this system is similar in composition to that of the worms the tendency is to centralize control in the head.

Vertebrate systems.—In the vertebrates the most remarkable changes are the development of the true brain as an organ of mental operations, and the transfer of the whole system to the dorsal side of the body where it is protected in the skeletal structures.

The essential features of the systems found in the different forms of vertebrates are identical, the variation being in the direction of complexity. Figures 203 and 204 illustrate the manner in which the complex brain of man has been developed, and the relative stages in

FIG. 202.—Nervous system of a caterpillar.

development seen in the vertebrates lower than man. The feature which strikes the eye as most vital in this

FIG. 203—A comparison of vertebrate brains; *O.L.* olfactory lobes; *C.B.*, cerebral hemispheres; *O.L.* optic lobes; *C.L.* cerebellum; *M.* medulla; *S.C.* spinal cord.

comparison is that the changes have arisen mainly in order to accommodate the increasing brain surface to the limits of the inclosing cavities. For example, in the

FIG. 204—Comparative wrinkling of cerebral surface of cat (*A*), and man (*B*).

lower vertebrates the increase in size has been met by doubling and folding one part upon the other, while in the higher forms the wrinkling or furrowing of the cortex gives increased surface.

Along with these physical changes has come increased mental power, until, in man, we have an animal preëminent over all others by virtue of the superior mental powers.

XXVIII. THE SPECIAL SENSES.

TOUCH, TASTE, AND SMELL.

IN the preceding chapters we have discussed, mainly, the mechanism by which sensory and motor impulses are conveyed and controlled; and the areas of the central system where these sensory impulses are received and translated into conscious sensation. The manner in which impulses originate and their character requires a study of the arrangements at the surface ends of the afferent fibers. Such arrangements are grouped together under the name of end or sense organs. They are all specialized structures adapted for response to certain kinds of stimuli only, and are named according to the effect of the impulses upon the consciousness. Thus, organs which produce sensations of sight are called sight organs, those which produce sensations of touch, tactile organs, etc. It must be distinctly borne in mind, however, that all of these organs merely originate impulses, and it is only when these impulses have traveled over the afferent fibers and stimulated certain tracts of the brain cortex that we become conscious of the sensations which they produce. In short, the sense organ is simply a structure adapted for response to a special kind of stimulus; and the impulse sent out by this stimulus, the brain interprets as sight, pain, heat, etc., according to the quality of the impulses.

Laws of sensory impulses.—There are three laws of sensory impulses which are applicable to all kinds of sensory fibers.

First law. A sensory fiber wherever stimulated, carries only one kind of impulse. Thus a touch nerve fiber, whether stimulated in its end organ or elsewhere always produces the sensation of touch, never that of sight or hearing.

Second law. In consciousness an impulse is located as coming from the natural source of such an impulse. Thus, in an amputated leg, pressure upon the cut end of the nerve is felt by the brain as though it came from the original end organs of that nerve. In this way may be explained many of the instances where pains are felt in the toes or fingers even after these parts have been amputated.

Third law. The increase in any given sensation from a definite kind of impulse does not take place uniformly with the increase in stimulation. On the contrary the amount of stimulation has to increase by a definite quantity before any increase in sensation is felt. A man named Weber investigated this relation between stimuli and sensation, and the law is often called Weber's Law on that account. An illustration will make clear its application. If we hang a thirty-gram weight to the finger we get a certain feeling of pressure. If we now add a half-gram weight no increase in pressure is felt, and it is only when we add a whole gram that the increase is actually felt. If, now, we start with sixty grams we have to add two grams before the increase is felt; that is, one thirtieth of the original weight. In this case, the stimulus increase necessary to sensation is one thirtieth of the original weight.

The amount of increase in a stimulus necessary to increased sensation varies greatly with different fibers. It is possible to reduce by practice the increase in stimulus

required by any given fiber. This relation of amount of stimulus to sensation is one of the factors which determine the sensitiveness of a sense center. The tips of the fingers, which require much less increase in stimulus to produce sensation than the skin of the back requires, are said to be more sensitive than the latter.

Classification of sense organs.—The end of every afferent nerve is a sense organ, but for convenience these end organs are grouped under various heads according to the region where they end and the quality of their impulses. The following scheme presents the commonly accepted classification:

CLASSES OF SENSE ORGANS.

Name of Group	Location of End Organs	Quality of Impulse
Cutaneous .	Skin and mucous membrane of the mouth.	Pressure Warmth Cold Pain
Internal . .	Internal membranes and muscles	Pain Muscle sense Hunger and thirst
Gustatory .	Tongue	Taste
Olfactory .	Nasal passages	Smell
Visual . . .	Eyes	Sight
Auditory . .	Ears	Hearing

Cutaneous sensations.—It was formerly customary to group all the sensations which were produced by the impulses from the sensory nerve endings in the skin as touch sensations. Modern experiments have shown that touch sensations show four distinct qualities, namely, pressure, warmth, cold, and pain, and that this difference in

quality depends not upon the nature of the stimulus but upon the particular fiber which is stimulated. In other words, *the fibers and end organs which transmit impulses of warmth are as distinct from those which transmit impulses of pressure as they are from those of sight, hearing, and smell.* For this reason the old term, touch sensation, has been replaced by the word cutaneous (skin) sensations, and the variety of end organs included under this head are to be considered as four distinct kinds of sense organs.

FIG 205. — A touch corpuscle in a skin papilla; C, corpuscle; N, nerve fibers.

Cutaneous end organs. — The sensory fibers which end in the skin show several varieties of end organs. Some form networks in the outer dermis where the axis cylinders lose their sheaths, and from this network extend out into the epidermis as tiny naked filaments. Others show specialized structures which may be classified under three heads, as tactile cells, tactile bulbs, or tactile corpuscles. At present it is impossible to tell by the look of a cutaneous end organ what kind of impulse it will transmit. In other words, unlike the fibers which end in the eyes and ears, we cannot be sure from their structure whether a given form of end organ produces pressure impulses or heat impulses, pain or cold.

Distribution of cutaneous sense areas. — The most interesting fact concerning cutaneous senses is that all parts of the skin are not equally sensitive to all kinds of stimuli. For example, in a given area of skin there will be spots which respond only to pressure, and not to cold or heat. Other spots in the same area will respond to heat and not to pressure, cold, or pain. In short it is possible to map any given

portion of the skin and to locate on such a map by different colored dots the position of the different sources of sensation. The relation of these spots to the fibers may be summarized as follows: (a) The surface of the skin is divided into spots or areas in the center of which is the ending of a sensory fiber. (b) Each of the four cutaneous senses has its own areas and fibers, and stimulation of one of these spots produces only one kind of sensation, heat or cold, warmth or pressure. (c) The pain spots are most numerous and the warmth spots least numerous. The ends of the pain fibers are nearest the surface and those of the warmth fibers farthest below the surface. (d) Discrimination between two stimuli of the same kind depends upon the stimulation of two distinct fibers. Thus, we feel the pressure of two points as two only when each point stimulates a different pressure fiber from the other. (e) Fibers of a given kind are much nearer together in certain parts of the skin than in others, and sensitiveness in any given direction depends partly upon the nearness together of the fibers.

Pressure. — The pressure spots are, next to the pain spots, the most numerous. In those portions of the body which are covered with hair the pressure fibers terminate in a ring surrounding the hair follicle. These rings, therefore, form the end organs. In other parts of the body the tactile corpuscles seem to function as end organs. These corpuscles are particularly abundant on the tips of the fingers where they underlie the dermal papillæ. It is estimated that on the entire body exclusive of the head region there are about five million pressure spots. These spots are very close together on the tips of the fingers and less near in other parts of the body.

The delicacy of pressure sense can be measured in two ways, by determining the least amount of pressure necessary to arouse a sensation, and by locating the least distance apart at which two points can be felt as two. The method of determining the first consists in varying the pressure upon various portions of the body until the least amount necessary to produce a sensation in any given part is determined. The results of such tests show that the skin of the face, forehead, and temples is most sensitive of all the skin areas to the feeling of pressure. The back of the hand is more sensitive in this respect than the tips of the fingers. Two milligrams of pressure is necessary to arouse sensation on the forehead, while from five to fifteen milligrams is necessary when applied to the finger tips. Such tests simply determine the relative sensitiveness to stimuli of the parts tested.

To test the power of discriminating between two pressure stimuli, the method consists in applying to the surface two points and determining how near together these points can be placed and still felt as two. The relative sensitiveness of different areas of the body to such a test is given in the following table: ¹

Area	Distance of Points Apart
Tip of tongue	1.1 millimeters
Tip of finger	2.3 "
Middle of palm	8 to 9 millimeters
Forehead	22.6 millimeters
Back of hand	31.6 "
Forearm	40.6 "
Chest	45. "
Middle of back	67.7 "

¹ N.B. In both methods of test the subject is blindfolded.

The above results show that while less stimulus is necessary to arouse a sense of pressure in the face than in the fingers, the power to discriminate is particularly acute in the finger tips and tongue. This second method is also of service in locating the distribution of nerve ends, since to feel two points as two means that two nerves are being stimulated.

Another interesting feature of this discrimination sense is that it can be increased with practice, notably in the case of the blind. This power is probably due to increased branching in the nerve ends thus increasing the number of spots on a given area.

Temperature sensations.—These are of two kinds, heat and cold, each with separate spots of stimulation and separate fibers. Their location may be determined by passing gently over the skin of the blindfolded subject, metal points which are colder or warmer than the skin surface. As this is done there will be aroused at definite points distinct sensations of heat or cold, which are at the same time entirely distinct from pressure sensations. If a certain area be explored in this manner and the spots located by ink spots, of different color, say red for cold spots and black for warm, the extent of the nerve ends and the relative number of heat and cold nerves for a given area can readily be determined. Figure 206 shows such a map. The results of such tests applied to all areas of the body may be summarized as follows:

FIG. 206—A portion of the skin mapped for hot and cold spots; white spots, cold; black spots, hot.

The cold and hot spots are always distinct, some portions of the skin being sensitive only to cold, some to heat. The cold spots are much more numerous and more sensitive to stimuli than the heat spots. The sensitiveness to stimuli varies for different areas of the body, the order of sensitiveness to temperature being as follows: tip of tongue, eyelids, forehead, cheek, lips, limbs, trunk. Temperature sense is always relative to the temperature of the skin, that is, objects colder than the skin stimulate cold spots and feel cold, and *vice versa*. Hot and cold spots may be stimulated by mechanical and chemical stimuli. Menthol, for example, gives a cold sensation. They may also be stimulated from within, the skin feeling hot in case of fevers being a case in point. No distinctive end organs for these sensations have as yet been discovered.

Pain sensations.—It used to be thought that pain was the result simply of excessive stimulation of sensory fibers. Thus, if a pressure nerve was stimulated by too great a weight the result was a sensation of pain added to that of pressure but conveyed by the same impulse. It is now believed that pain sensations of the skin at least are produced by special impulses which pass over special fibers and that these fibers are distributed in spots as are the other cutaneous nerve ends. The results of careful experimentation in locating and determining the peculiarities of the skin pain spots may be summarized as follows:

Summary.—(a) Pain spots are distinct from and more numerous than other cutaneous sense spots. (b) Pain spots of different regions vary greatly in the amount of stimulus necessary to produce a sensation. Those of the eye being much more sensitive than those of the fingers. All, however, require much greater stimulus than those of temperature or pressure. (c) The nerve fibers

which supply the pain spots have no distinctive end organs so far as known. (d) They may be stimulated by any of the usual forms of cutaneous stimuli, such as temperature or pressure, provided these stimuli are of sufficient intensity. (e) The location of a pain stimulus by the brain is aided by the pressure and temperature stimuli, parts which have lost these centers still retaining the sense of pain but the subject has difficulty in locating the seat of stimulation.

INTERNAL SENSATIONS.

The membranes of the internal organs, such as the mucous membranes that line the alimentary tract and the lungs and the membranes that invest the muscles or cavities of the body, have no pressure, warmth, or cold fibers. Increase in internal temperature or pressure can be detected, therefore, only by its effect upon the cutaneous fibers. These membranes are, however, supplied with pain fibers and other fibers whose function it is to supply the brain with information as to the health and needs of these parts. Very little is known of these latter fibers or their end organs, their method of distribution or stimulation. The sensations of hunger, thirst, general healthfulness or weakness are all produced by impulses passing over these fibers, and their function as indicators of the condition of our internal organs is extremely important. At present, we know only of the presence of these fibers and admit our ignorance of their end organs or properties.

GUSTATORY SENSATIONS.

Distribution of fibers and end organs.—The sensation of taste is produced by impulses sent over special nerve fibers whose ends are distributed over various parts of the mouth cavity and particularly in certain regions of

the tongue. If the tongue is protruded and its surface examined with the aid of a mirror it is possible to dis-

FIG. 207 — The tongue; *K*, filiform papillæ; *I*, fungiform papillæ; *L*, circumvallate papillæ.

tinguish many raised points on the surface of its mucous membrane. These raised points are called papillæ, and are of three forms — the *circumvallate*, *fungiform*, and *filiform*. Of these, the circumvallate are the largest and least numerous. They appear as circular extensions of the membrane and are surrounded by a little depression or ditch. They are only seven to twelve in number, and lie on the upper surface near the root of the tongue.

The fungiform papillæ are so called from their resemblance to a mushroom, and consist of small, rounded elevations supported on short, slender stalks. They are found all over the middle and fore part of the tongue on the upper surface.

The filiform papillæ are most numerous and the smallest, and appear as little pointed projections scattered all over the upper surface and edges of the tongue.

FIG. 208—Section of circumvallate papilla.

In the walls of the circumvallate papillæ, in some of the fungiform papillæ, and in portions of the pharynx, palate, and epiglottis, and even in the vocal cords, are peculiar structures known as taste buds. These taste buds are oval bodies composed of a number of overlapping epithelial cells (protective) surrounding a core of elongated or taste cells which project through a pore at the top some stiff hairlike processes and are connected at the base with the taste nerve fibers. The hairlike extensions constitute the part of these organs which is directly stimulated by the tasted substance. The impulse thus originated is passed through the body of the taste cells and finally stimulates the terminal branches of the taste nerve fibers themselves. These taste buds, therefore, may be con-

ected with the taste buds. These taste buds are oval bodies composed of a number of overlapping epithelial cells (protective) surrounding a core of elongated or taste cells which project through a pore at the top some stiff hairlike processes and are connected at the base with the taste nerve fibers. The hairlike extensions constitute the part of these organs which is directly stimulated by the tasted substance. The impulse thus originated is passed through the body of the taste cells and finally stimulates the terminal branches of the taste nerve fibers themselves. These taste buds, therefore, may be con-

FIG. 209—A, isolated taste bud, from whose upper free end project the ends of the taste cells; B, supporting or protecting cell; C, taste cell.

sidered as the specialized end organs of the sensory taste nerve fibers.

The perception of taste is produced by the connection of these fibers and the brain through the fifth and ninth pairs of cranial nerves.

Stimulation and sensitiveness.—These taste buds can be stimulated only by substances in solution, and in dissolving solids the saliva of the mouth plays an important part in the production of taste sensations. Dry salt or sugar placed upon the tongue remains tasteless until dissolved by the saliva. The sensitiveness of the taste buds to stimuli is influenced by many external factors as well as by the concentration and character of the dissolved substance. The result of experiments in this direction may be summarized as follows:

Summary.—(a) The taste buds are most sensitive at a temperature of from 50 to 86° Fahrenheit. Very high or very low temperatures tend to paralyse the taste buds and thus lessen or destroy the sensation of taste.

(b) Rubbing the substance to be tasted against the tongue increases the sensitiveness of the taste buds. This accounts for our habit of rubbing the tasted food against the walls of the mouth with the tongue.

(c) The sensitiveness varies with the character of the tasted substance. The order of greater sensitiveness in this respect being bitterness, sourness, sweetness, saltiness.

Kinds of taste sensations.—While our taste sensations appear to be very varied in kind, it has been demonstrated that they are actually only four in number, namely, bitter, acid, sweet, and salt. The bitter taste is most developed at the back of the tongue, the sweet taste at the tip of the tongue, the acid or sour taste at the sides, and the salt taste is nearly equally distributed. Many of our taste sensations are combinations of these four primary sensations with odor or sight sensations. This fact can be

easily demonstrated by closing the nasal cavities and bandaging the eyes when the sensation of taste produced by certain foods will often disappear entirely. This fact also accounts for loss of taste in cases of colds in the head, the nasal passages being clogged by mucus. Certain substances taste sweet when first taken into the mouth and bitter when swallowed. This is due to the fact that these substances are capable of stimulating both sweet and bitter taste buds and also demonstrates that these primary sensations are carried by different afferent nerve fibers.

OLFACTORY SENSATIONS.

Distribution of fibers and end organs.—The nasal cavity is located just above the hard palate. It communicates

FIG. 210—Section of nose showing outer wall of right nasal cavity; *a*, *b*, *c*, *d*, interior of nose; *k*, olfactory bulb giving off nerves to inside of nose.

with the external air through the two nasal openings and with the pharynx by a single posterior opening. The

walls of this cavity are formed by the inferior maxillary palate, the nasal and ethmoid bones, and is lined throughout with mucous membrane. In this membrane are distributed the fibers of the first pair of cranial nerves (the

olfactories), and through these fibers impulses of smell are carried to the olfactory lobes. These fibers end in the upper part of the membrane of this cavity in special end organs of smell. The lining of the nose is, therefore, divided into a lower or respiratory membrane composed mainly of ciliated epithelial cells, and an upper or olfactory membrane, rich in nerve fibers, and devoid of ciliated cells. The end organs themselves consist of the ends of the fibers developed into long, olfactory cells, and these cells are protected on all

FIG. 211.—Olfactory cells from lining of nose; *O, O*, olfactory cells terminating nerve fibers; *E, E*, ordinary epithelial cells of the mucous membrane; *H*, sensitive cilia affected by odor.

sides by surrounding nucleated, epithelial cells. The olfactory cells show a long, nucleated cell body somewhat like a taste cell, and, like that, ending in tiny hair-like projections at one end, while the other is continuous with the nerve fiber. The hairlike processes are first stimulated, and transmit this stimulus to the fiber which conveys the impulses to the brain.

Stimulation and sensitiveness.—The explanation of the stimulation of an olfactory nerve fiber is as follows:

It is supposed that all odoriferous bodies give off particles in the form of vapor, which latter when dissolved in the liquids of the mucous membrane stimulate the sensitive hairs of the end organs. The vapor may be given off by a near or distant body, but the question of whether it be smelled or not is determined by the currents of air which carry this vapor. If the air is inhaled it carries this vapor through the nasal passages, where the dissolved particles produce the sensation in the manner described. Similarly, swallowed bodies may send their vapors up into the nasal passages through the posterior openings, and stimulate in the same manner.

The sensitiveness of the olfactory nerves to stimuli varies greatly in different persons and between persons and animals. While in man the sensitiveness is not so great as in dogs and many of the lower vertebrates, it is still very acute. By taking odoriferous substances and diluting them to known amounts it has been shown that the sense is so keen in the average man that one part of camphor in four hundred thousand parts of air can still be readily detected, while odors like musk and vanilla can be diluted to eight and ten million parts of air and still be detected. These olfactory cells, however, soon become exhausted by continuous stimulation and cease to react. Their sensitiveness is also affected by lack of secretion on the part of the membranes.

Kinds of olfactory sensations.—Smell sensations are usually classified as agreeable and disagreeable, according as their effect is pleasant or unpleasant, and these effects undoubtedly serve to guard the person from evil atmospheres. This is hardly a true

classification, however, as many odors are agreeable to some persons and intensely disagreeable to others. It has not yet been determined whether there are, as in taste, fundamental or primary odors out of which others are compounded, but it seems reasonable to suppose that such is the case, and that special fibers convey special kinds of odor impulses.

XXIX. THE SPECIAL SENSES (*continued*).

VISUAL SENSATIONS.

THE eyes are the external organs by means of which visual impulses are produced. So complex are these organs that we must first study their anatomy and some of the laws of light before we can appreciate the manner in which they produce visual impulses.

Form and protection of the eyeball.—The eyes con-

FIG. 212—Section of eyeball in orbit; 1, frontal bone; 2, cheek bone; 3, eyebrow; 4, eyelids; 5, folds of the conjunctiva lining eyelids and covering front of eye; 6, 7, 8, muscles moving eyeball; c, cornea; a, aqueous humor; i, iris; l, lens; v, vitreous humor; ch, choroid layer; r, retina; s, sclerotic layer; o, optic nerve.

sist of two nearly spherical bodies called the eyeballs, which, together with certain protective structures, completely fill the two sockets in the facial bones which we call the orbits or eye sockets. Both eyeballs and their accessory parts are identical in structure, hence a description of one will serve for both.¹ In front the eyeball is protected by movable folds of the

skin called the eyelids. These lids are lined with a

¹ A sheep's skull with the bony parts and eyeballs uninjured serves well for a demonstration of the structure of the eyeballs. (See Ex. LXX.)

mucous membrane called the *conjunctiva*,¹ which is also folded back to cover the front of the eyeball. The lids are movable, and when lowered form a protective curtain in front of the eyeball. Their edges are lined with sebaceous glands, which, by secreting oil, prevent their sticking together when closed. A row of hairs also fringes these lids and serves to keep out dust and furnish a slight shade. These lids are closed by the contractions of a circular muscle (the orbicularis) which acts in part reflexly, as when an object is suddenly brought toward the eye or head.

As these lids rest against the front of the eyeball, friction would naturally result between the two surfaces when they rub together. The mucus secreted by the membrane partly prevents this by acting as a lubricant. There is in addition a small gland about the size of an almond (the tear gland) located on the outer and upper side of the orbit, and this gland secretes a salty liquid called the tear fluid or the lachrymal fluid. The gland is also known as the *lachrymal* gland. This fluid flows over the front of the eyeball assisted in its progress by the motion of the lids until it reaches the inner angle of the lids next to the nose. Here it is drained off into a small duct (the *lachrymal duct*) which

FIG. 213 — View of right eye and tear apparatus; *G*, tear gland; *C, C*, upper and lower tear ducts; *B*, common nasal tear duct.

¹ Inflammation of this lining causes conjunctivitis, or pink eye.

empties it into the nasal passages. Not only does this fluid aid in lubricating the surface of the ball and eyelid, but in passing continuously across the exposed part of the eyeball it removes from the surface dust or other foreign particles which may tend to collect there. Sometimes this fluid is poured forth very copiously, as when the membrane covering the eyeball is irritated, or the person is under strong emotion. Under such circumstances the ducts cannot drain it off fast enough, and it accumulates and rolls over the edges of the lids down the checks. Under ordinary conditions of flow, the oily secretion at the edge of the lids is sufficient to prevent its escape in this way.

Back of the lids the eyeball itself rests upon a fatty layer which lines the orbit. Connected to it is the optic nerve which enters the eyeball from the back through an opening in the socket. This nerve carries the visual fibers to their centers in the brain. The eyeball is also held in place and moved by six muscles. These muscles are controlled by motor impulses sent from the brain along the fibers of the third, fourth, and sixth cranial nerves. The attachment of

FIG. 214—Muscles of the eyeball.

these muscles shows an ingenious mechanical adaptation to the structure of the orbit. (See Fig. 214.) Four (*rectus* muscles) are straight muscles attached at one end to the eyeball and at the other to the bottom of the eye socket. Two (the *internal* and *external rectus*) are attached to the

sides of the eyeball, and, by contracting and relaxing alternately, turn the eye toward or away from the nose. Two (the *inferior* and *superior rectus*) are attached to the lower

and upper surface of the eyeball respectively, and by their action turn the eyeball upward or downward. The other two muscles (the *inferior* and *superior oblique*) serve to rotate the eyeball outward or inward. The inferior oblique has its origin near the front of the orbit on the inner side and passing under the eyeball is inserted on the outer side. Its contraction rotates the eyeball outward. The superior oblique has its origin at the back of the orbit like the rectus muscles, passes forward just above the superior rectus and its tendinous end, after first passing through a loop or pulley formed by a fibro-cartilage on the upper edge of the orbit, turns outward over the top of the eyeball, and is inserted in the eyeball at a point on the inner side. By this arrangement the contraction of a straight muscle rolls the eye inward.

The combined action of these six muscles is responsible for all the movements of the eyeball. The muscles of the two eyeballs contract and relax in pairs, the internal rectus of one eye contracting at the same time as the external rectus of the other, and similarly for all the others. In this way, both eyes are turned in the same direction at the same time.

As has been already mentioned, the eyelids are closed by the orbicularis muscle. The upper lids, however, are provided with elevators muscles attached to the back of the orbit, and these muscles pull the upper lids open farther than would be possible by the extension of the orbicularis alone.

Structure of the eyeball.—The eyeball itself, when detached from its supports, is a nearly spherical sac of about an inch in diameter. It is covered, except at the points where the optic nerve enters and in front, with an opaque

white coat known as the white of the eye or *sclerotic coat*. In this layer are the blood vessels which supply the coat with

FIG. 215—Section of the eyeball; *Sc*, sclerotic coat; *Con*, conjunctiva; *C*, cornea; *I*, iris; *O*, optic nerve; *Ch*, choroid layer; *C.P.*, ciliary processes; *Cm*, ciliary muscles and ligament; *R*, retina; *L*, crystalline lens; *A*, aqueous humor; *V*, vitreous humor.

nourishment. At the front this layer is replaced by the transparent layer of mucous membrane (the infolded *conjunctiva*) and a transparent modification of the white of the eye called the *cornea*. Back of this cornea is a colored, circular, muscular curtain called the *iris*.

If we split the eyeball vertically so as to split the optic nerve it shows the following internal structure. The conjunctiva in front folded back from the eyelids. Over the whole surface, and covering the outside of the optic nerve, is the sclerotic layer just mentioned, with its transparent window or cornea in the front. Inside the sclerotic

is a second layer, called the *choroid*, composed of loose connective tissue, and colored black by pigment particles similar to those in the skin. At the front where the cornea joins the sclerotic this middle layer is thrown into plaits (*ciliary processes*) beyond which it is continued as a muscular curtain (the *iris*). This iris is composed of circular and radial unstriped muscle fibers, is colored by pigment, and has a circular aperture in the center (the *pupil*). At its outer edge it is firmly united with the sclerotic layer by a ligament called the *ciliary ligament*, and to the *ciliary process* of the choroid. The contraction of its circular fibers decreases the size of the pupil, while the contraction of the radial fibers reverses this action. By means of this variation in the pupil, the amount of light which enters the eye is kept under control.

The third layer of the eye is an extension of the optic nerve called the *retina*. It consists of the endings of the optic nerve fibers and a supporting layer of connective tissue. The entire layer is very thin and lines the whole interior of the eyeball with the exception of the front, where it stops at the edge of the ciliary process. This retina is the sensitive layer of the eye and the one in which the visual impulses originate.

Just behind the iris is a biconvex, transparent body, called the *crystalline lens*. The front surface of this lens is slightly flatter than the back and is in contact with the inner surface of the iris. Between it and the cornea is a space which is filled with transparent watery liquid called the *aqueous humor*. The entire cavity of the eyeball back of this lens is filled with a glassy semi-solid mass called the *vitreous humor*. The lens itself is surrounded by a thin membranous capsule, and is kept in place by a circular

band (the *suspensory ligament*) which attaches the edges of the capsule to the ciliary process of the choroid layer. Between the ligament and the edge of the choroid layer are a series of muscles (the *ciliary muscles*). When these muscles contract they draw forward the edge of the choroid layer, thus causing the suspensory ligament to relax. This contraction permits the lens to shorten its diameter and become thicker and more bulging. In this way the shape of the lens may be changed, and this affects its image-forming power.

Functions of the various eye structures.—The human eye is constructed essentially like a camera, and we can

FIG. 216—An eye model.

obtain an idea of the function of its parts by study of the parts of a camera or eye model such as shown in Figure 216. This shows a box painted black on the inside and having at the front a small aperture into which may be fitted perforated cardboard disks of different sized apertures. Inside the box is a biconvex lens of glass so mounted as to bring its center opposite the center of the cardboard aperture, and to be capable of movement forward and backward. It also contains a ground glass screen similarly mounted, and adjustable to forward and

backward movement. If, now, we place this box with the aperture pointing toward an illuminated object and move the lens backward or forward, we find that for a certain position of the lens there appears on the ground glass screen an inverted picture of that object. In other words, this combination of lens, screen, and aperture, with its light-excluding walls enables us to project inverted images of external objects upon the screen. If we should replace the screen in this apparatus by a sensitive photographic plate we should obtain a permanent impression of the image or picture. Our box would then fulfil all the functions of a camera.

Now, the human eye is essentially such a sort of image producer as this box or camera. The only difference is that, instead of a sensitive photographic plate capable of one exposure and producing one picture, we have a nervous structure, the retina, which receives these impressions and transmits them to the brain as nerve impulses. This retina is always ready to receive impressions; the human camera is always "loaded." But the resemblance to the camera does not stop with the retina. The pupil of the eye with its iris serves to regulate the amount of light which shall enter, just as the cardboard disks or stops of a camera do. The sclerotic layer corresponds to the wooden or metal walls of the box, and like those gives shape and strength to the structure. The black, inner choroid coat, by means of its pigment, absorbs light, and prevents internal reflection, just as does the black paint on the inside of the camera box. Finally, the cornea and lens correspond to the lens of the camera, and, like that, produce inverted images upon the sensitive surface or retina.

The points of difference between the eye and a camera

may be noted as follows: First, the retina and lens of the eye are fixed in position. The external aperture is protected by the transparent cornea and thus makes the eye dust-proof. The air is replaced by the transparent aqueous and vitreous humors, but this does not affect the comparison, since these humors are transparent. They are made necessary by the fact that layers of the eye are not stiff like the wood of the box and would collapse were it not for the presence of these fluids. Finally, the lens and retina are fixed in position, and the eye, therefore, must resort to special means for producing pictures of near and distant objects and bringing them all to focus upon the retina. To understand this last process it is necessary that we first understand how a lens produces images.

FIG. 217—Scattered images produced by rays distributed all over the receiving surface.

How pictures are formed by lenses. — Every point of a luminous body transmits light in the form of rays. These rays pass from the luminous body in straight lines and in every direction. Wherever a ray of light strikes it produces an image of the point from which it came. In the ordinary course of light rays, therefore, images are formed all over the receiving surface, and the total effect is a

jumbling together of all these images so that it is impossible to detect anything but a mass of light. (See

Fig. 217.) Now, each ray is supposed to be made up of vibrations of a substance which fills all space and is called ether. The rate at which this ether vibrates determines the color of the light ray. These vibrating lines or rays when they pass from one transparent substance into another of different density are bent in definite directions, the shape and density of the transparent body determining the direction of bending. Glass structures called lenses are simply masses of transparent matter so shaped as to control the direction of the bending in rays which pass from air into them. By means of them it is possible

FIG. 218—Effect of lenses on rays of light; *A*, a concave lens spreading the rays; *B*, a convex lens bending the rays to a point.

to concentrate the rays which pass through them (convex lenses) or to spread the rays (concave lenses). (See Fig. 218.)

The point to which rays are converged by a convex lens, or from which they seem to be spread by a concave lens, is called the focus of those rays and the action of the lens is called focussing. By its shape, a convex lens is able to focus the rays which it receives from a luminous body, so that the rays, which come from a certain point on the object shall, after passing through it, all converge at a certain spot. In this way the lens is able so to collect the

scattered rays of a luminous body that a sharp but inverted image of the object is thrown upon any screen placed at the point of convergence. (See Fig. 219.) In

FIG. 219—Convex lens of eye forming an image by converging the rays on the retina.

a camera a glass lens serves this purpose. In the eye, the cornea and crystalline lens accomplish the same result.

Accommodation.—In a camera with a lens of a definite shape the images formed by the converging rays are focussed at different distances back of the lens according as the objects which produce the rays are near to or far from the lens. To secure a sharp image for any given object such instruments are provided with mechanical adjustments by means of which either the lens or the screen may be moved until the distance between them is such that the screen is exactly at the point of convergence of the rays. Such an adjustment is called accommodation. This accommodation to distant and near objects may also be secured by varying the thickness of the lens, and in the eye it is this latter method that is employed. As has been seen, the cornea and retina (screen) are fixed. By means of the ciliary muscles and suspensory ligament,

however, the shape of the crystalline lens may be so changed as to secure this accommodation, and the involuntary movements of these muscles enables the human eye to accommodate itself automatically to near and distant objects. (See Fig. 220.)

The eye a self-supporting camera. — In all physical details, then, the eye acts as a camera, and forms inverted images of objects upon the surface of the retina, exactly as the camera does upon the sensitive plate. Being living tissue, however, it requires nourishment for its cells, and hence the choroid layer not only serves by its color to prevent internal reflection but through its blood vessels nourishes the tissue cells of the inner eye. The sclerotic layer also contains blood vessels which supply it with food.

Effect of the images upon the retina. — The actual sensation of sight is located not in the eye but in the occipital lobes of the cerebral hemispheres where the visual impulses of the optic nerves are ultimately received. The optic nerve fibers in fact originate in these regions, passing from here to the anterior corpora quadrigemina (see Fig. 184, page 396), and are continued from here to a junction called the optic commissure. At this point many of the fibers cross, thus connecting the left eye with the right hemisphere, and *vice versa*. Some, however, do not cross, so that if the center of vision in one of the hemispheres is destroyed, there will still be sight in both eyes. This crossing of the main fibers is called the optic chiasma. After passing through it the fibers inclosed in two right and left trunks (the optic

FIG. 220 — Diagram to illustrate changes of the lens in accommodation.

nerves) pass to the back of the orbits, where they enter the eyeball, and, spreading out their fibers, have their endings in

the end organs of the retina. The projection of an image upon the retina stimulates these end organs and thus arouses the impulses of vision. How the brain cells interpret the impulses produced by an inverted image into a sensation of sight is unknown.

Structure of the retina.—The optic nerve fibers enter the retina on its anterior surface. Near this surface the fibers connect with the axis cylinders of large ganglion cells. These cells give off other cylinders which connect with other cells (see Fig. 221), and these

FIG. 221—*A*, Diagram of structure of the retina seen under the compound microscope; *B*, the essential nervous elements; *C*, the direction of light stimuli; 1, internal membrane; 2, nerve fibers; 3, nerve cells; 4, terminal brushes of neuria; 5, inner nerve cell layer; 6, 2d set terminal brushes; 7, nerve cells; 8, external membrane; 9, rods and cones; 10, pigment layer.

cylinders finally terminate in modified nerve cells at the very back of the retina next to the choroid layer. This last

layer of cells has received from their shape the name of the rods and cones. They are separated from the choroid layer by a thin layer of pigment. It is these rods and cones which the light vibrations of the projected image stimulate, and after passing through this complicated system of cells the impulses finally enter the optic nerve fibers and pass on to the brain. A peculiar feature of this arrangement is seen in the fact that though the rays pass over the connecting cells to reach the rods and cones they produce no impulses until they reach this area. The rods and cones are believed to convey different qualities of visual impulses. The cones are believed to produce the impulses which give color sensations, while the rods produce merely degrees of light and darkness.

Peculiarities of the retina.—Not all parts of the retina are equally sensitive to light. The region where the optic nerve enters the eye is a little to one side of the middle line. This spot, of course, has no rods and cones, and objects whose images fall upon this spot are invisible. It is called, therefore, the blind spot. The rest of the retina has a varying sensitiveness. That part of it directly back of the pupil in the mid line, naturally receives continuously more rays than the other parts. At this point (see Fig. 215) is a yellow spot about a quarter of an inch in diameter, where the retina is thinnest, and the rods and cones most numerous. This is the region of keenest vision.

Peculiarities of sight sensations.—Sight sensations are produced almost immediately upon turning the eyes toward an object, but the retina has the power to retain images upon it for a short interval (about one eighth of a second). Owing to this fact, impulses which succeed one another rapidly give the effect of continuous sensation. For example, falling raindrops appear as lines, flames swung in a circle as continuous rings of light, and moving pictures give the effect of a continuous movement. Owing to this same cause, two revolving colors give a mixed sensation, black and white producing gray, etc.

The sensitiveness of the retina is readily exhausted. Hence, if we look at a bright light for a short time that portion of the retina upon which the image falls soon becomes exhausted and loses its power to transmit impulses. If, now, we turn the eye from the light to a white surface, there appears to be a black spot on it, due to the rays which fall upon the blinded spot, in the retina.

Color sensations are due mainly to a difference in the quality of the impulses, and these differences are due to different rates of vibration, each color having its own time of vibration. White light is a mixture of the impulses of all colors, blackness the absence of all light impulses. There are also many different shades of color in sensation which are produced by combining impulses from two primary colors. Thus, green is a mixture of yellow and blue. Lack of ability to distinguish colors is called color blindness. In persons thus afflicted the light waves of several different colors may produce the same sensation in the retina. This affliction is due to a defect in the cones of the retina.

In binocular vision, images of the object are projected on both the retinas, but the impulses thus set up and transmitted by the two sets of nerve fibers give us only a single sensation in the brain. This shows that the sensation of sight is not a simple action but the result of association and fusing of impulses in the brain cells.

The intensity of the illumination increases the intensity of the stimulus and determines the brightness of the sensation. Very strong light or very dim light tends to strain the eye and should be avoided. Individuals differ among themselves and from the lower animals in the amount of light necessary to arouse sensation. Cats, for example, are able to distinguish objects with an illumination far less than that required by human eyes. Certain animals, such as the owls, whose eyes respond to the weak light of the night, become almost blind in the bright light of day. The retinal end organs may be stimulated by other means than by light waves. Blows on the eyes or head may produce sensations of light, and we "see stars," as the saying goes. The shadows cast by blood corpuscles of the eye blood vessels often modify the effects of vision.

HYGIENE OF THE EYES.

The eyes are extremely sensitive organs, and their satisfactory working depends upon the health of the general system. Poor circulation, which results in lack of nourishment, as in disease, often affects the eyes. Biliousness often renders the eyeballs yellow and sore. Nervous diseases also disturb their functions. Filth and dirt or rubbing the eyes with dirty hands may often cause diseases of the eyeball and lids. To remove the dirt which accumulates, the eyes should be bathed as carefully as other parts of the body. For this purpose warm or tepid water is better than cold.

Reading in a strong light or dim light or overuse tends to strain the muscles which operate the lenses and eyeballs and thus weaken them. Flickering light especially requires overexertion on the part of these muscles in accommodation and should be avoided. Often the eye is defective in structure, and unless this defect be corrected may react upon other parts of the system producing severe headaches and other nervous troubles.

All eye troubles may be classed as eye defects or eye diseases. The most common defects of the eye are nearsightedness, farsightedness, and astigmatism.

Nearsightedness and farsightedness. — In a normal eye images of both near and distant objects are, through the accommodation of the lens, projected exactly upon the surface of the retina. When, however, the eyeball is too short or too long, that is, abnormal in shape, the image instead of being formed on the retina is projected either in front or back of it, and the result upon the vision is a blurring often accompanied by headache. When the image is formed in front of the retina the defect is called nearsightedness (*myopia*). The defect may be corrected by the use of spectacles made with divergent lenses, which spread the

rays which enter the eye just enough to overcome this defect. In farsightedness (*hypermetropia*) the image is projected back of the retina, and convergent glasses are necessary to remedy this difficulty. In both of these cases the amount of convergence or

FIG. 222.—Position of retina in near and far sight; *B*, in natural sight; *G*, in far sight; *C*, in near sight.

divergence exerted by the spectacle lenses should be only such as to counterbalance the defect of the lens in this particular. The selection of glasses, therefore, should always be the result of prescription by an oculist who has determined by careful examination just how much correction is needed. Many eyes are ruined by careless selection of glasses.

Astigmatism. — A very common defect in the eyes of children and adults is an irregular curvature in the surfaces of the cornea or lens, or in both. The result of this is that rays which pass through the affected parts are brought to a focus at a different point from those which pass through the unaffected parts. This produces a blurred image on the retina, and indistinct vision. The eye muscles become strained in the constant attempts they make to change the focus of the lens and secure a clear image, and this strain often results in chronic headache. This defect receives the name of astigmatism. It can be corrected only by the use of glasses whose surfaces are so formed as to correct these irregularities in surface, and the selection of these glasses should be based upon careful examination by an oculist.

Cataract. — Degeneration of the lenses at points renders them opaque. This opacity, whether it occurs in spots or over the whole surface of the lens or its inclosing capsule, is known as cataract. The extent determines whether it produce partial or

total blindness. Removal of the eye lens completely, and the wearing of special forms of spectacles, permits the spectacle lens to act as a substitute for this eye lens and form images on the surface of the retina as before. Since these glass lenses have no power of accommodation, a different pair must be used for reading and the examination of near objects than is used for distant objects. They are of course a makeshift, but infinitely preferable to blindness.

Presbyopia. — In old age the lenses become stiffened and do not bulge enough. Hence convergent lenses are necessary, as in farsightedness. Such stiffening is called *presbyopia*.

Eye diseases. — Diseases of the eye include the formation of ulcers or scars on the cornea, the failure of the muscles of the iris to expand or contract, inflammation of the conjunctiva (pink eye and granulated lids), and the wasting of the optic nerve. Most are curable provided the optic nerve is not permanently injured.

XXX. THE SPECIAL SENSES (*concluded*).

AUDITORY SENSATIONS.

As with visual sensations the sensory fibers of the nerves of hearing have their endings in a pair of external organs called the ears. By means of these ears the waves of sound are made to produce impulses of hearing just as the eyes transform waves of light into visual impulses.

Structure and form of the ears.—The ears are to external appearance merely two twisted flaps of skin-covered

FIG. 223—Diagram of the ear.

cartilage attached one on each side of the head. These external flaps are, however, only the external parts of the organ whose sensitive parts are embedded in cavities of the temporal bones. For convenience in description, we divide the ear into three parts, which we call the external ear, the middle ear, and the inner ear. Since both ears are

exactly alike, a description of one applies in every particular to the other.

The external ear.—The oval, folded, and roughly funnel-shaped external ear consists, as has been mentioned, of a projecting structure of cartilage covered with a layer of skin and blood vessels, and bound to the side of the head by bands of muscle. These muscles in some individuals and in many of the lower animals are capable of voluntary movement. The irregular outer surface of this ear converges near the center to form a funnel-shaped canal about an inch long. This canal is lined with skin whose sebaceous glands secrete a peculiar form of fluid which on exposure loses its water and thickens into a bitter yellow wax (*ear wax*). This wax serves to keep moist a membrane stretched across the bottom of the canal (called the tympanic membrane). This membrane is covered externally by a thin layer of skin, and is extremely flexible. The external flap collects the sound waves, acting in this respect like the horn of an ear trumpet, and concentrates them upon this ear drum which is thus set in vibration.

The middle ear.—The tympanic membrane forms the external covering of a small cavity called the middle ear, or *tympanum*. This cavity is lined with mucous membrane and filled with air. A narrow tube (*Eustachian tube*) connects it with the pharynx. In swimming, water often passes from the pharynx through this tube into the middle ear and produces ringing in the ears, until it is allowed to run out. On the inner surface of the cavity are two small openings covered with membrane. One is called, from its shape, the round window, or *fenestra rotunda*. The other which is oval in shape is called the *fenestra ovalis*. Between the tympanic membrane and the fen-

estra ovalis there is strung across the cavity a chain of three bones called from their shape the hammer (*malleus*), the anvil (*incus*), and the stirrup (*stapes*) bones. One end of the malleus rests against the tympanic membrane, while its other end rests against the incus. The incus, in turn, articulates at its farther end with the stapes, whose interior end rests against the membrane of the fenestra ovalis. These bones, together with the air, permit vibrations of the tympanic membrane to be transmitted across the middle ear to the membrane of the fenestra ovalis. The Eustachian tubes play a part in this transmission in permitting the escape of air into the throat when the pressure becomes too great, and by admitting it return the membrane to position. If it were not for this outlet a sudden increase of pressure on the tympanum might burst the drum or one of the inner membranes. Its action also increases the sensitiveness of the ear drum to vibrations,

as is seen in the deafness that results when this tube is clogged, as by mucus in a cold. The middle ear is called the tympanum.

The inner ear. — The fenestra ovalis and fenestra rotunda are the openings by which the tympanum communicates with an irregular cavity of the temporal bone called the

FIG. 224 — Bony labyrinth.

osseous, or *bony, labyrinth*. This labyrinth and its contents constitute the inner ear.

The entire labyrinth is lined with a membrane which

secretes a fluid called the perilymph. It contains a closed membranous sac, or capsule, called the *membranous labyrinth*. This sac fills almost completely the spaces of the bony labyrinth. The bony labyrinth is usually divided into three areas. The central part, about an eighth of an inch in diameter, is called the vestibule. The part back of this consists of three tubes called the *semicircular canals*. The part in front of the vestibule is a single tube coiled like a snail shell and called the *cochlea*. Into the osseous labyrinth enter the branches of the eighth pair of cranial nerves (auditory).

The bony labyrinth.—(See Fig. 224.) The vestibule lies just back of the tympanum, and communicates with it by the membrane-covered fenestra ovalis. The three semicircular canals lie toward the back of the head and communicate with the vestibule by five openings, two of the canals uniting before entering the vestibule. The cochlea lies forward of the vestibule. It is a spiral tube consisting of two and a half turns about a central pillar. The large end opens into the vestibule. It communicates with the tympanum by the membrane-covered fenestra rotunda.

The membranous labyrinth.—This closed membranous sac lies in the bony labyrinth surrounded by the perilymph of that cavity. It conforms very closely to it in shape. Internally it secretes a fluid called the endolymph, which fills it and gives it shape. In this liquid are tiny particles of lime called *otoliths*. The membranous vestibule is divided into two sacs, the *utricle* and the *sacculus*, connected by a small opening. The utricle is the larger and from it arise three membranous semicircular canals. The sacculus is smaller and communicates with the membranous cochlea. The membranous

semicircular canals are about one third the diameter of the bony canals which they occupy. One of these is horizontal when the body is upright. The others are vertical, but at right angles to one another. All three have swellings at the front where they join the utriculus, called *ampullæ*. At the back these swellings are absent, and the vertical canals fuse before entering the utriculus. The three ampullæ and the walls of the utriculus and sacculus are supplied with branches of the auditory nerves which penetrate the membrane and attach it at these points to the bony walls.

Into the bony cochlea projects from the central pillar a thin shelf of bone dividing it partially into two sections. Attached to the edge of this shelf is the membranous cochlea, forming a triangular tube with the base of the triangle against the opposite wall. It thus divides the bony cochlea into three compartments. One of these compartments (the vestibular passage) communicates with the bony vestibule. Another, the tympanic passage, has its base against the fenestra rotunda. The middle passage, or membranous cochlea (sometimes called the cochlear canal), communicates with the sacculus. The vestibular and tympanic passages are filled with perilymph, the cochlear canal with endolymph. Along a groove in the bony shelf passes a branch of the auditory nerves whose fibers penetrate the membranous cochlea and end in special end organs known as the fibers of Corti. These fibers are the seats of origin of sound impulses.

Nature of sound stimuli and path of sound in the ear. — Just as light stimuli are produced by waves of vibrating ether, so sounds are produced by waves of vibrating air. Any vibrating body can produce these waves, and if they

reach the ear they will stimulate the auditory nerve and produce the sensation of sound. When these waves of air reach the ear they are first collected by the external ear and concentrated upon the tympanic membrane. This membrane is thus set in vibration in unison with the vibrating body, which caused the waves. Its vibrations are in turn communicated by the bones and air of the middle ear to the fenestra ovalis of the inner ear. The membrane covering this in turn sets in vibration the perilymph which fills the spaces of the bony labyrinth and thus distributes these vibrations to all parts of the inner ear. The rotunda by bulging out when the ovalis moves in, and *vice versa*, permits the vibration of the otherwise incompressible perilymph. Finally, the vibrations of the perilymph are communicated to the membranous labyrinth and its endolymph and thus reach the fibers of Corti where they start the auditory impulses.

The organs of Corti and sound impulses.—While the auditory nerves send branches to all parts of the membranous labyrinth, the impulses of hearing appear to originate almost exclusively in the cochlear division of this nerve. This division gives off many fibers which terminate in the sensory cells of the organ of Corti.

This organ of Corti is located inside the membranous cochlea. It consists of three parts, the *rods*, the *sensory cells*, and the *supporting cells*. The arrangement of these parts is seen in the accompanying figure. (See Fig. 225.) Notice that the rods are arranged in pairs, three thousand to five thousand in all, and are so constructed as to inclose a passage, their bases being expanded while the bodies lean toward one another. Against these rods on either side lean the sensory cells with hairlike extensions reach-

ing into the endolymph. These latter cells receive the terminal brushes of the fibers and are supported below by the layer of cells called the supporting cells.

FIG. 225 — Diagram of cross section of a single coil of the cochlea.

The action of this organ is not fully understood. All that can be stated definitely of the action and the manner in which sound impulses are interpreted may be summarized as follows:

Summary.—(a) The vibration of the perilymph being communicated to the endolymph through the membrane, sets up in it similar vibrations. These vibrations, in turn, move the hairs of the sensory cells and stimulate the fiber brushes, thus starting impulses.

(b) Not all the hair or sensory cells are affected by external vibrations. The region inclosed by the rods, and the membrane upon which they rest, does not respond as a whole to the vibrations of the perilymph, that is to say, when a particular vibration is set up in the perilymph only certain regions of the rod inclosed portion vibrate with it. As a result, only certain ones of the sensory cells are stimulated, and it is supposed that ability of the brain to distinguish different sounds depends upon the fibers which are thus stimulated.

(c) The character of the vibrations produced by a vibrating body, and hence set up in the perilymph may vary in three ways, in pitch or tone, in loudness, and in quality. These variations are distinguishable by the brain through variations in the impulses which the sensory cells transmit to it.

(d) The final interpretation of sound impulses has been located in certain convolutions of the temporal lobe of the cerebrum, which is connected by association fibers with the medullary origin of the auditory nerves. The exact area is much more difficult to localize than the sight area.

(e) The two organs of hearing produce in the brain not two sensations but one. Their position on each side of the head aids, however, in locating the source of vibrations.

(f) The sensory cells become fatigued by continuous stimulation in much the same manner as do the retinal cells of the eye.

Functions of the membranous vestibule and the canals. — Divisions of the auditory nerve supply the utricle, saccule, and canals with fibers. The endolymph of these parts is also continuous with that of the cochlear part, and they are surrounded by the perilymph of the bony labyrinth. Further, the fibers of these divisions of the auditory nerve end in the ampullæ, utricle, and saccule in sensory hair cells similar to those of the cochlea. In spite, however, of these similarities of equipment, these hair cells do not appear to transmit ordinary sound impulses. The present theory of the action of these parts may be

stated as follows: As the body or head changes position, the endolymph contained in the various canals and the sacs is subjected to pressure from different points. It thus tends for a given position to press upon definite sensory cells of the sacs and ampullæ. This pressure is transmitted to the brain as an impulse by the nerve fibers attached to these cells, and these impulses enable the brain to become conscious of the relative position of the body, and thus to exert the necessary control upon the muscles which coördinate movements. The sensitiveness of these sensory cells to such pressure is increased by the presence in the endolymph of the tiny particles of lime (otoliths) which act upon the hairs. This sense of position is often classified as a separate sense, though it is located in the ear.

HYGIENE OF THE EAR.

The ordinary injuries to the ear consist mainly of damage to the outer or middle ear. The inner ear is so well protected that it is almost impossible for external causes to affect it. Among the injuries to the outer ear which may be avoided by care is the over-accumulation of ear wax or dirt on the tympanic membrane. Such accumulation increases the thickness and destroys the sensitiveness of the membrane to vibrations. Ear wax allowed to accumulate in this way often hardens, and thus produces deafness. Washing the ear frequently with tepid water will remove the cause and thus prevent this trouble.

Colds or throat troubles may cause stoppage of the Eustachian tube by mucus or even inflammation of its lining and that of the middle ear. This results sometimes in the formation of abscesses, which may be simply

painful, or, by weakening the membranes, may cause them to break, and thus produce total deafness. Loud noises, likewise, may cause too great pressure upon the drum, while adenoids or throat growths often clog the Eustachian tube and produce deafness.

XXXI. SENSATION IN THE LOWER ANIMALS.¹

SENSATION may be defined as the perception of stimulation. For example, when a hard body presses against our finger it stimulates a nerve impulse which passes to the brain and is there interpreted. As a result we perceive the presence and action of the hard body. In the higher forms of animals the seat of consciousness is the brain, and the various organs where the sensory nerve fibers end are connected by these fibers with the brain. It is to be noted here that sensation does not take place necessarily when a sensory fiber is stimulated. It is only when those impulses reach the brain and are perceived that we have a true sensation. In short, the seat of sensation is the brain, and not the sense organ, and we must distinguish sharply between mere stimulation of sensory impulses and their perception by the brain as sensations.

Sensation in the protozoans.—It has been known for a long time that the one-celled animals have in their protoplasm a very sensitive substance which reacts to various external stimuli. Whether these reactions indicate true sensation, that is, whether they are accompanied by any consciousness on the part of the animals or not, has caused much speculation in the past. In attempting to answer this question two theories have been advanced. One school holds that these tiny organisms do actually possess the rudiments of consciousness, and that their reactions to stimuli indicate the true conscious choice on their part. The other school

¹ See Footnote, p. 146, Chapter X.

holds that these reactions are simply the results of physical or chemical changes produced in the protoplasm by the stimuli, and that the movement toward food, and away from particles which are not food, can be fully explained by the laws of physics or chemistry, and no more indicate consciousness than movements of iron particles to a magnet prove consciousness in the particles.

The latter school has conducted exhaustive experiments to demonstrate its point and has proved pretty conclusively that many of these responses to stimuli on the part of protozoans are the result of the chemical or physical action of the stimulus upon the protoplasm. Thus the movement of an amœba toward a food particle they explain as due to chemical attraction between the compounds in the protoplasm and the food, and not to conscious choice of the amœba. Recent investigations (notably those of Professor Jennings of the University of Pennsylvania) have shown responses that are difficult to explain by these natural laws and the question is still unsettled.

At any rate, these reactions to stimuli result in movements beneficial to the organism, and hence, whether they be indications of consciousness or not, they correspond to the function of sensation in higher forms.

DEVELOPMENT OF THE SENSORY ORGANS.

The sense of touch.—This sense is found in all animal forms. In the coelenterates it is confined to a few sensory cells which are located in special regions of the body. In forms higher than the coelenterates it is pretty generally distributed over the whole body covering, though some parts are more sensitive than others. In every case

stimulus applied upon a given area of covering affects a sensory cell, or cells which are themselves the termination of touch nerve fibers and control the area stimulated.

Taste and smell.—The sense organs of taste and smell are very difficult to distinguish in the lower forms. When found they consist usually of pits or depressions in the body covering, lined with sensory cells which are ciliated, and are the terminations of taste or smell fibers. Often the location of these pits is the sole indicator as to their function.

In the higher forms of life these sensory cells are still seen in the hair cells of the taste buds or olfactory buds. They are, however, distributed in connection with different organs with a view to putting them in the path of odors or tastes. Thus, the ciliated taste buds of our tongue, while very similar in structure to the olfactory cells, are in a position to come in contact with food rather than odors, while the cells of the nose are in the path of inspired air which brings them odors rather than taste stimuli.

Sensitiveness of these taste or smell cells varies greatly in different animals. Many animals, for example, can detect the presence of an enemy or of food by smell alone, when man would be utterly unaware of its presence.

Hearing.—In the lower forms, the organs of hearing consist of vesicles or cavities which may be open to the outside air or water or may be inclosed in the body covering. These vesicles are lined with sensory ciliated cells, and filled with a fluid in which float usually one or more particles of limestone formation called otoliths. Each of the sensory cells is the termination of an auditory nerve fiber, and when a sound wave strikes the body it sets in vibration the fluid and otoliths of these cavities. This

movement on the part of the fluid and particles stimulates the sensory cells by rubbing against the hairs, and thus produces sensation. Such organs are called otocysts, and are to be compared with the vestibule and semicircular canal structures of our ear rather than with the cochlea.

In the vertebrates all, except the lowest fishes, possess a structure similar to the inner ear of man located usually in the head. The external ear, however, is a structure peculiar to mammals. The cochlea and canals of the human ear are supposed to have developed from the simple vesicle of the lower forms. It is supposed, for example, that the vestibule, with its utriculus and sacculus, corresponds to the original vesicle from which the canals and cochlea have arisen as outgrowths. The arrangement of the sensory cells, and the presence of otoliths in these parts, is very suggestive of the otocyst of the lower form.

Vision.—The simplest form of a visual organ consists of a pigment spot overlying a sensory cell which is the termination of a nerve fiber. The pigment spot absorbs the light and converts it into a stimulus to the sensory cell. Such spots are found on the disks of the jelly-fishes and in the ray tips of a starfish.

The next step is to cover the pigment spot with a transparent thickening or lens. This lens by forming an image permits the animal to distinguish form instead of simply shapes of light and darkness as when the pigment only is present. In the snail we have such simple eyes or ocelli. Here the lens is in the form of a sphere covered with a transparent cornea and its image is projected upon a nervous network which we may consider as a simple retina. Over this network is the pigment layer. In the

snail these simple eyes are two in number and are located at the ends of the tentacles. In the scallop they are numerous and are located on the edges of the mantle.

The next step in advance is found in the insect. Here thousands of ocelli are massed together side by side to form what is called a compound eye. (See Fig. 226.)

FIG. 226—Compound eye of insect; A, section; *a*, optic ganglion; *b*, secondary nerves; *c*, retina; *d*, pigment layer; *e*, optic nerves; B, group of ocelli; *f*, bulb of optic nerve; *g*, pigment layer; *h*, vitreous humor; *i*, cornea.

Each ocellus points at a different angle and hence while the whole eye is fixed the range of vision is extensive. In such an eye each ocellus forms its own image independently of its neighbours. These images are all collected by the brain into a single perception much as are the images of our two eyes.

The nearest approach to the structure found in man and in all other vertebrates among the invertebrate forms is the

eye of the squid. Here we find a lens with a humor-filled cavity, retina, and other layers much as in man. In the squid eye, however, the rods are on the side of the retina nearest the lens and they are supplied with optic fibers from the back of the eye.

The eyes of the vertebrates are essentially alike in structure. The principal differences are in the means used to secure accommodation and range of vision.

XXXII. THE VOICE.

IN the study of the ear we noted that sounds are produced by setting in vibration various objects, which vibrations are propagated as waves of air. When the air waves reach the ears they produce impulses of sound. In our study of the respiratory system we noted that in the human body the expired air was capable of producing sounds and that the various modifications of these sounds were called the voice. Combining these two observations we should expect to find in our bodies some structure in which vibrations of different pitch, loudness, and

FIG. 227 — The larynx; *A*, front view; *B*, side view; *C*, back view.

quality could be set up by expired air. Such an instrument, in fact, is found in the larynx or cartilage box at the

top of the trachea or windpipe. This box is prominent externally as a projection which we call the Adam's apple.

Structure of the larynx.—The larynx or voice box is composed of cartilages, membranes, and muscles. The front and sides are formed of a single large cartilage (the *thyroid*). The curved front of this cartilage has a projection at the top like the lip of a pitcher (the Adam's apple). It curves around on each side in broad, flat sheets, which end above and below as projecting points. The two upper points are attached by ligaments to the hyoid bone and the two lower projections fit into sockets in a second cartilage called the *cricoid*.

This cricoid cartilage is a complete ring. The broad part of this ring forms the back of the larynx, while the narrow part encircles the front of the box just below the thyroid. On the top of the broad part of the cricoid are two triangular cartilages (*the arytenoids*) whose bases form true joints with the cricoid and whose apices extend upward. By means of the joints all these cartilages can move freely upon one another. Beside these four hyaline cartilages there are five yellow cartilage structures. One, the *epiglottis*, is attached to the top and front of the thyroid, and projects upward to form a lid for the box. Two others form small horn-shaped projections on the top of the arytenoids and the remaining two are inclosed in folds of the membranous lining of the box.

The box is lined with a smooth layer of mucous membrane. This layer is smooth, except at the point where the glottis or pharynx opening is located. Here it is thrown outward in two sets of ridges. The upper ridges are called the false vocal cords. They have nothing to do with voice production, but help to close the glottis. The two lower

ridges are made up mainly of elastic tissue running in the membrane from the front angle of the thyroid cartilage to their attachments at the base of the arytenoid cartilages. These ridges are the true vocal cords. They have a glistening white color and sharp smooth edges. In ordi-

FIG. 228—I, The vocal cords in the larynx as seen with larynxoscope; A, when voice is being produced; B, when no voice is produced; *e*, epiglottis; *cv*, vocal cords; *cvs*, false vocal cords; *u*, *s*, *w*, elevations due to cartilages; *l*, root of tongue. II, A, the cords alone when voice is being produced; B, vocal cords when no voice is produced.

nary breathing they are separated, forming a V-shaped opening to the glottis.

The arytenoids are united to one another and to the cricoid cartilages by bands of muscle. By means of these muscles they may be pulled forward, backward, or to one side. These muscle movements affect the cords attached to the arytenoids, pulling them apart, or by their backward movement varying the cord tension.

Vocalizing.—In ordinary breathing the arytenoids are

drawn forward and to one side and the vocal cords are thus separated. In this position they do not vibrate, and hence produce no sound. In whispering they are in the same position and the sounds produced are in this case due entirely to the position of the lips and tongue.

In talking, shouting, singing, however, these cords are drawn inward and backward by the cartilage muscles, and the expired air is thus forced through the narrower slit between their edges. This sets the edges in vibration producing sound. How this sound is varied in pitch, quality, and loudness requires some knowledge of the laws of vibrating bodies.

Laws of sound vibration. — The production of sound depends upon two factors. The first is a vibrating body. The second is a layer of air between this body and the sense organs which produce the impulses. Under these conditions the vibrations of the body are transmitted by the air in the form of waves. If the body which vibrates does so regularly and continuously, a musical tone is produced. If irregularly or at irregular intervals, the effect is called noise. Sound sensations differ in three characteristics, pitch or tone, loudness, and quality. This variation in sensation finds its explanation in corresponding variations in the manner of vibration of the sounding body.

(a) *Loudness.* Of two vibrating bodies vibrating at the same rate, the one whose vibrations are widest are most audible. Thus, a body struck hard produces wider vibrations and hence a louder sound than when struck lightly.

(b) *Pitch.* The number of vibrations a body makes in a given time determines the pitch or tone of the sound. Thus, a tuning fork whose prongs vibrate two hundred and fifty-six times a second produces the pitch called middle C of the piano, while the next note above it in the musical scale requires two hundred and eighty-eight vibrations per second. Vibrating bodies differ in size and elasticity, and this difference determines their rate of vibration. A large-sized string vibrates slower, and thus gives a lower

tone than a small one, while a tightly stretched string gives a higher note than one of the same size but less taut.

(c) *Quality*. A piano string and a violin string may be of the same size and tension and be struck with the same force. In that event they will give the same pitch and have the same loudness. We are still able, however, to detect that one is a violin string and the other a piano string. This difference between sound producers of the same pitch and loudness we call quality. It is due to the fact that few bodies vibrate simply and as a whole. Most bodies have not only a vibration as a whole which gives what is called the fundamental note, but separate parts of the bodies vibrate more or less independently producing distinct waves, which combine with the fundamental and are called overtones. No two bodies produce these overtones in the same way, owing to differences in size, shape, or material, and hence the complex tone produced is characteristic of each vibrating body and enables us to distinguish between various sources of sound.

Resonance.—The quality of a sound may be changed and its loudness increased by reënforcing the original vibration with others of a similar pitch. Thus, a violin string stretched over a sounding board sets the air of the sounding board in vibrations of the same pitch, reënforcing and altering the quality and loudness of the original string vibrations. Such reënforcement is called resonance and the reënforcing body is called a resonator.

Application of sound laws to vocalizing.—The vocal cords furnish the vibrating bodies which are struck or set in vibration by the expired air. The air cavities of the throat, pharynx, nasal cavity, and mouth are the resonators.

By movements of the larynx cartilages the tension of these cords can be changed, and thus the pitch is regulated. The force with which the air is expired determines the width of vibration and the loudness, while loudness and the quality of the tones may be still further modified by changing the shape of the resonating cavities.

Articulate speech is the result of voluntary changes in these resonating cavities, and the organs most concerned in producing the changes are the lips, tongue, and teeth.

Vowels are open-mouth sounds, with corresponding changes in the lip and cheek positions. Consonants are closed-mouth sounds, and tongue, lips, teeth, and cheeks aid in their production. Harshness, softness, etc., are due to differences in the quality of the vocal cord vibrations and the character of the resonating cavities

In a child the resonating cavities and larynx are smaller than in adults. At about thirteen to fifteen years of age the larynx widens rapidly and the cords lengthen. The result is a deepening of tone and a change of voice.

The vibrations of the cords are under voluntary control, and it is possible by practice to control these vibrations and the shape of the resonators so as to improve greatly the clearness and quality of the voice.

HYGIENE OF THE VOCAL ORGANS.

Since the voice is under voluntary control it is important that children be taught to control it properly. This teaching must be mainly by example, and the best training is secured by giving the child good examples to follow. Singing lessons are also valuable in teaching by exercise the proper control of the vocal muscles, even though the child never become a singer. Exercise which develops the lung capacity also increases the volume and resonance of the voice.

If the lining of the larynx becomes inflamed through colds or irritation the vocal sounds are at once affected. Similarly, inflammation of the walls of the resonators or air

passages reacts upon the voice. On that account colds of all sorts manifest their presence almost immediately in the quality of the vocal sounds. The ordinary precautions against disease of these passages, therefore, act as preventives of vocal troubles.

XXXIII. BACTERIA AND SANITATION.

WHENEVER moist foods, such as milk, eggs, meats, bread, etc., are exposed to warm air for any length of time changes take place in their composition which render them unfit for food. These changes we designate as souring, fermentation, decay, mold formation, etc., according to the character of the change. Such changes are produced not by the action of the air, but by the presence in the air of certain vegetable organisms commonly called germs. Analysis of these germs shows that there are three forms of plants which may cause changes in food, and that these three forms belong to the classes *yeasts*, *molds*, or *bacteria*. While these three forms of plants are very different in their appearance and action, they show striking similarity in the manner in which they bring about changes in foods. Of these three forms, the bacteria are perhaps the most varied in their effect; and since they illustrate the action of all, as well as being vitally connected with the health and disease of the human animal, we are justified in devoting part of our study of physiology to the action of these plant forms.

Experimental study of bacteria. — (See Ex. LXXII.) If we prepare three test tubes by placing in one chopped hay and water, in another some milk, and in a third some wheat and water, and allow all three to stand uncovered for some time in a warm room, we may observe the following changes in the contents of the tubes:

The hay solution becomes turbid, and a scum forms on its surface, while from the mixture comes a disagreeable odor.

The milk gradually curdles and becomes sour to the smell and taste.

The meat gives off fetid odors and changes in color and appearance.

If, now, we mount some of the liquid from any one of these three tubes on a slide under a compound microscope, we find it swarming with curious-shaped little bodies myriad in number and with sharply defined outlines. These bodies are extremely small beings, only $\frac{1}{50000}$ to $\frac{1}{500000}$ of an inch in diameter. Examined with the highest powers of the microscope they are seen to consist of a cell wall surrounding a bit of protoplasm, while scattered through the protoplasm are tiny particles which color more strongly than the other parts, when stained, and are supposed to correspond to the nuclear material of other cells. These tiny bodies are *bacteria*.

Definition and forms of bacteria.—Study of preparations such as the above, and of other sources, shows that there are many different species of bacteria, all differing in shape and action, but all showing the characteristic wall of cellulose and the scattered nuclear material. It is this cellulose wall that inclines us to place these forms among the plants. We can, therefore, define bacteria as follows: One-celled vegetable organisms having a definite unchanging cell wall and containing nuclear material in the form of scattered fragments which is never combined in the form of a single nucleus. Certain forms show cilia and are capable of movement. Others move only as they are carried about by the air or liquids in which they are present.

In identifying and naming bacteria, use is made of the unvarying character of the outline. Experience has also shown that definite actions are always associated with defi-

nite shaped cells. Thus, bacillus tuberculosis is the name given to the particular bacterium which causes tuberculosis. In this name the bacillus refers to the shape of the cell (*bacillus* = rod shaped), while the second name indicates the specific action of this form. Similarly, staphylococcus pyogenes means spherical cells producing pus formation. Spirillum volutans means corkscrew-shaped cells which whirl in their motion through liquids, etc. (See Fig. 230).

FIG. 229 — Bacteria, highly magnified; a, the germ of typhoid fever, stained to show the cilia; b, a spiral ciliated form; c, a rod-shaped form, in chains; d, a spherical form.—a, b, from ENGLER and PRANTL.

Bacteria as living cells.—Bacteria contain no *chlorophyll* (or green coloring matter) and, therefore, cannot make starch. On this account they must find their food ready made. Living organisms which obtain their food at the expense of other organisms are called *parasites*, and the living organisms upon which they live are called their hosts. If, however, the source of food be no longer living, as was the case of the chopped hay in the tube, the organism which feeds upon it is called a *saprophyte*. Bacteria are all parasites or saprophytes. Those which take their food from living cells, such as the diphtheria and consumption forms, belong to the parasitic forms, while those which live upon dead matter, such as dead animal or plant structures, belong to the saprophytic forms.

Like all living cells, bacteria take in food and oxygen, and give out wastes. They are rendered inactive by cold and the presence of certain chemicals, while most of them are

killed by exposure to high temperature, such as the boiling-point of water. They require water to develop in, and when there is no water present they become inactive, and in a dried condition may be easily borne about by the winds. This drying, however, does not kill them, and when transplanted to a moist place they become active again and grow.

The most remarkable feature of their action is the rapidity with which they reproduce. Like all single cells, they reproduce by division. In bacteria, however, this process of division is much simpler than in higher forms of cells and is accompanied by no nuclear changes. When a bacterium has reached a certain size the walls begin to fold in at the middle, a partition forms, and the two halves break apart to form two individuals. Each of these halves absorbs food until it becomes of normal size and then splits again. In active forms this whole process may be repeated as frequently as once in thirty minutes, and it has been calculated that one bacterium can produce at this rate a colony of over 200,000,000,000,000 in twenty-four hours. It is this marvellous rate of development that makes the introduction of a single disease causing bacterium into our system so dangerous. A single diphtheria bacillus can alone produce little damage, but if allowed to develop the increase may become sufficient to cause serious damage in a short time.

Occurrence of bacteria and infection.—Bacteria are found in earth, air, and water, and in fact may be present practically everywhere on the earth's surface. Certain conditions are, however, more favorable to their development than others, and fortunately the malignant forms arise only under conditions which are subject to control.

How bacteria produce diseases and other effects. — In the act of obtaining their food bacteria may bring about chemical changes in the material from which they take it, thus transforming that material into new and simpler compounds.

Again, they may give off as wastes, materials which are poisonous to the host upon which they live. Such poisons are called *toxins*.

Finally, they may by rapid reproduction become so numerous as to clog effectually the part of the host where they are present, and thus interfere with the normal action of that part.

All bacterial effects may be traced to one of these three causes. For example, the souring of milk is brought about as follows: The lactic bacteria found in milk break up the sugar in the milk in the attempt to obtain food and transform it into lactic acid. This acid causes the proteid part of the milk to coagulate or curdle, and gives the sour taste and odor. Again, the bacteria which collect in meat break up the complex proteid matter into simpler compounds and thus reduce it to simple mineral compounds. Both of these examples illustrate the first mentioned effect of bacteria.

The action of the diphtheria bacterium illustrates the second method. In this case, the bacterium gives off a toxin, or poison, which enters the blood and disturbs the action of the organs of the body. In the case of diphtheria, a substance has been found which if injected into the blood destroys the effect of the toxin, and is called on that account an *anti-toxin*.

The formation of tubercles by the massing of the tuberculosis forms in the lungs, thereby breaking down the

tissues and clogging the air sacs, is an example of the third method of action. Frequently a given bacterium combines these methods of effect. In every case, however, the organism produces its effect through some result of its metabolism, and these secondary effects, such as poisoning or clogging, decay, and the like, are the result of the attempt of one organism to take its food from another.

Beneficial bacterial action.—It must not be thought that all bacteria are harmful. If it were not for decay of waste materials, for example, the farmer would be unable to raise any crops. Decay in his case means the transformation of dead plant and animal matter into simpler compounds, such as are acceptable as food to growing plants, and without bacteria the supply of such food in the soil would be soon exhausted. Again, in the manufacture of many dairy products, such as butter and cheese, the bacteria perform a very important part in the manufacture. Certain forms of bacteria have also been found which, if introduced into the soil with certain plants, can transform the nitrogen of the air into soluble nitrates which the plants can use, and such forms are extremely valuable.

Harmful bacterial action.—The forms of bacteria which are most harmful to man are those which produce disease, and the following list gives us some idea of the diseases which are now known to be due to bacterial action:

Bubonic plague	Diphtheria	Lockjaw
Asiatic cholera	Erysipelas	Consumption
Fowl cholera	Glanders	Typhoid fever
Hog cholera	Boils and carbuncles	

The forms of bacteria which are beneficial in causing decay of waste material may be harmful to food supplies. The

methods employed in the preservation of foods are devices employed to prevent the action of bacteria upon food stuffs.

Relation of business methods to bacteria.—In the paragraphs on the action of bacteria as a living cell we noted certain conditions which were favorable to their development, and *vice versa*. The methods used in the preservation of food depend upon a proper use of the facts there noted. For example, in cold storage methods we have the application of the principle that bacteria are rendered inactive by cold. Foods, therefore, will not decay so long as they are kept sufficiently cold to secure this inactivity. We also noted that bacteria need water to develop in. Dried meats, therefore, and foods preserved in salt or sugar, either are free from this necessary water or have the water replaced by solutions of salt or sugar, which are not favorable to the action of the plants. The packing of foods in air-tight cans which have been submitted to temperatures of boiling water meets two conditions unfavorable to bacteria. The heating of the food and can kills the germs that might be present, while the sealing prevents the bacteria in the air from entering after the foods are cooled. Certain chemicals also kill bacteria, and it has been customary in the past to make use of some of these as preservatives of food. This latter method should always be carefully inspected, since often the chemicals used may be in such quantity and of such a nature as to be themselves dangerous to health. The pure food law in some of its clauses covers just this ground.

Medical methods.—The human body is extremely well protected from disease-causing bacteria, since the only avenue of approach to most of its organs is through the blood;

and to get into the blood is a difficult matter for bacteria, since the skin is of such structure as to prevent passage of the germs. In the case of wounds or breaks in the skin, these serve as excellent passages into the blood for bacteria. The entrance of bacteria into the body is spoken of as *infection*, and the part where they enter is said to be infected. It becomes, therefore, extremely important that a wound which is healing be kept free from these dangerous germs. Medical men have discovered certain chemicals which when applied to wounds do not prevent healing but at the same time do prevent the development of bacteria. Such substances are called antiseptics. All wounds should be first washed as clean as possible with warm water and then bandaged with one of these antiseptics. In this way antiseptics such as 0.1 per cent and 0.05 per cent solutions of carbolic acid and corrosive sublimate, may prevent such serious complications as tetanus or lockjaw, and blood-poisoning. It should be borne in mind that small wounds are quite as dangerous as large ones in this respect and should be treated as carefully.

In operations requiring the use of instruments everything used must be freed from bacteria. The process usually consists in treating the instruments and clothing worn as well as the hands to the action of various chemicals or antiseptics. Thus, the instruments are frequently soaked in solutions of carbolic acid or corrosive sublimate or are submitted to the action of hot steam. Materials made bacteria free in this manner are said to be *sterilized* or *disinfected*. Both terms indicate the same result, but the word disinfected is more frequently applied to the result of chemical treatment, while sterilizing more often refers to the action of heat. In removing a splinter with a needle

a simple method of precaution consists in sterilizing the needle by holding it in a flame for a moment. The bandages used in binding up wounds should all be sterilized in some manner.

Disinfection.—While this term means to free from bacteria, it is frequently used to refer to the methods in which objects are freed from disease germs. After a patient has been sick of diphtheria or other germ disease it becomes necessary as a matter of safety to the community that all clothing, furniture, and surroundings of the sick person be freed of germs, since their presence there might mean their introduction into some other person. There are various methods of performing this disinfection, but one of the simplest consists in sealing up the room and burning sulphur in it. The fumes of burning sulphur are sufficient to kill all forms of harmful bacteria. There are other kinds of disinfectants beside sulphur, and any form which kills bacteria is entitled to the name. In use it is imperative that the disinfectant be brought in direct contact with the germs. The mere presence of a disinfectant in a room is not a sufficient safeguard, and often disinfectants are sold which give out an odor but do not necessarily come in contact with the germs for a sufficient length of time to kill them. It is quite as important to know how to use a disinfectant as to know its name. One of the best disinfectants is plenty of bright sunlight.

SANITATION.

It is important that we know how to treat cases of bacteria-caused disease. It is still better to prevent such disease altogether. To a certain extent every one is responsible for the proper care of his person and property. In a

community, however, there are always some people who are careless in this regard, and unless the enforcement of such care be placed in the hands of specific authorities and backed by legal penalties for disobedience the action of one individual may result in great harm to the community. The department of a government, either city, town, or state, which has in charge the care of the public health is called the Sanitary department. In a city the work of sanitation is usually divided among several departments, for example, The Board of Health, The Street Cleaning Department, etc. The extent to which a city board of health looks after the health of its citizens is seen in the following summary of the duties of such a board:

Summary of duties of a board of health.—The enforcement of all laws, ordinances, and regulations relating to the safety of human life and the care and protection of health.

Exclusive control of the treatment of infectious and contagious diseases and of the public hospitals used for such cases.

May cause unsafe or unsanitary buildings to be vacated.

May condemn and destroy buildings that are themselves unfit for habitation or that make adjacent buildings unfit.

May destroy tainted or dangerous food.

May prohibit the sale of adulterated milk or other drinks or foods.

May prescribe and enforce rules concerning the sanitation of tenements, lodging houses, shops, and dwellings.

May be called upon to deal with sources of disease and infection and to keep the streets and houses free from matter dangerous to health.

It must insure the purity of the drinking water.

In short, it has full power in any case where the action of an individual tends to menace the health of the community.

Department of street cleaning.—This department is entrusted with the care of the streets and public highways of the city, and the removal of garbage, rubbish etc., from such streets.

Conclusion.—To attempt to indicate all the ways in which a community cares for the health of its individuals would be impossible in the limits of this book. It should be borne in mind by every citizen, however, that unless he himself takes an interest in maintaining his own health such carelessness may react upon the community, and that it is the duty of every individual to inform himself of the laws which govern the healthy action of his own body.

INDEX.

A.

Abdominal cavity, 109, 223.
Abducent nerve, 389.
Abomasum, 157.
Abscess, 207.
Absorption, 133.
 definition of, 133.
 hygiene of, 145.
 of nutrients, 142.
 summary of, 142.
Accommodation of eye, 462.
Acids, 23, 24.
Adam's apple.
Adaptation of digestive organs,
 154.
 of limbs, 261.
 of skulls, 262.
Adenoids, 335.
Adipose tissue, 44.
Adulteration, 68.
Afferent nerves, 403.
Affinity, chemical, 13, 15.
Air, composition of, 20.
 expired, 324, 325.
 inspired, 324, 325.
Air breathers, 341.
Air sacs, 319.
Albumin, 241.
Albuminoids, 53, 241.
Alcohol:
 as a beverage, 82.
 as a drug, 82.
 as a food, 75, 76, 77, 79, 85,
 86.
 as a fuel, 78.
 as a producer of heat and
 energy, 78, 80, 81, 293.

Alcohol —

 as a solvent, 84, 141.
 effect upon brain workers, 427.
 effect upon circulation, 210,
 211.
 effect upon digestion, 130, 131.
 effect upon gastric juice, 130,
 131.
 effect upon muscular activity,
 284, 292.
 effect upon nerve control, 211.
 effect upon nervous system,
 427.
 effect upon respiration, 338.
 effect upon saliva, 131.
 effect upon the heart, 211.
 effect upon the intestines, 131.
 effect upon the kidneys, 373.
 effect upon time reactions,
 426.
 effect upon veins and arteries,
 211.
 narcotic or stimulant, 426.
 percentage in beverages, 84.
 percentage in patent medi-
 cines, 84.
 relation of study to physi-
 ology, 75.
Alimentary canal, 52, 90.
 functions of, 93.
 in earthworm, 150.
 regions of, 92, 93.
Alkalies in saponification, 126.
Amœba, circulation in, 212.
 digestion of, 146, 147.
 movement of, 298.
 study of, 33.
Amœboid movement, 297, 298.

Ampulla, 474.
 Amylopsin, 125, 126.
 Anabolism, 59, 352.
 Anærobia (footnote), 341.
 Anatomy, 11.
 Anemia, 168.
 Ankle, bones of, 233.
 Ankylosis, 243.
 of joints, 251.
 Antiseptics, 500.
 Anti-toxin, 337.
 Anus, 129.
 Anvil bone, 235.
 Aorta, 186, 197, 199.
 Aortic arches, 214.
 Apoplexy, 398.
 Appendages of the body, 30.
 Appendix, vermiform, 129.
 Aqueduct or iter, 396.
 Aqueous humor, 457.
 Arachnoid membrane, 392.
 Arbor vitæ, 397.
 Arms, bones of, 232.
 Arterial blood, 187, 188.
 Arteries, 186-189.
 control, 205.
 pulmonary, 185, 195.
 pulse in, 186.
 renal, 201, 367, 368.
 structure, 189.
 Arthropods:
 body covering, 375.
 circulation in, 212, 213.
 excretion in, 381.
 nervous system of, 433.
 respiration in, 346.
 Articulations, 244.
 Arytenoid cartilages, 487.
 Asphyxia, 332.
 Assimilation, 175.
 Astigmatism, 123.
 Atlas vertebra, 225, 227.
 Auditory sensation, 470-479.
 Auricles, 182-184.
 action of, 191, 192.
 Auscultation, 193.

Autonomic system, 421.
 Axial skeleton, 224.
 Axis cylinders, 47, 402.
 Axis vertebra, 225-227.

B.

Bacteria, 493-503.
 action of, 498.
 as living cells, 495.
 experimental study of, 493.
 forms of, 494.
 metabolism of, 496.
 occurrence, 496.
 relation to blood poisoning, 210.
 relation to business methods, 499.
 relation to disease, 497.
 relation to fermentation in large intestine, 130.
 relation to inflammation, 207, 208.
 relation to medical methods, 499.
 relation to nervous diseases, 424.
 relation to respiratory diseases, 337.
 reproduction of, 496.
 size of, 494.
 structure of, 494.
 Ball and socket joints, 230, 231.
 Bases, 23, 24, 25.
 Baths and bathing, 363.
 Beef extract, nutritive value of, 74.
 Bellied muscles, 267.
 Biceps muscle, 206, 209.
 Bicuspid teeth, 95, 96.
 Bile, 122, 124, 127.
 Bile duct, 124.
 Biliousness, 144.
 Binocular vision, 466.
 Biology, 10.

- Birds, circulation in, 217.
 digestive system of, 156.
 heart of, 217, 218.
 respiration in, 349.
 skeleton of, 260.
- Biuret test for proteid, 53.
- Bladder, 367, 371.
- Bleeding, 209.
- Blind spot in eye, 465.
- Blood, 159-179.
 absorption into, 134, 135.
 aerating of, 203, 327.
 arterial, 187, 188, 203.
 changes in lungs, 326.
 clotting of, 161.
 cold, 351.
 composition of, 201, 202.
 conditions affecting, 170.
 experimental study of, 161.
 extractives, 164.
 flow, 191, 194-196, 204.
 in other animals, 173.
 in veins, arteries, and capillaries, 187, 188.
 loading of, with nutrients, 201, 202.
 loading of, with wastes, 202.
 plasma, 161.
 poisoning, 210, 365.
 properties of, 161.
 purifying, 203, 327.
 quantity and distribution, 171.
 relation to circulation, 159.
 relation to respiration, 326.
 structure, 160.
 systems, 191.
 temperature, 171.
 unloading of nutrients from, 202.
 unloading of wastes from, 203.
 venous, 203.
 warm, 351.
- Blood corpuscles.
 red, 160, 165, 167, 238.
 white, 167.
- Blood plates, 165, 169.
- Blood proteids, 164.
- Blood vessels, 159.
 of the heart, 184.
 of the skin, 356.
 relation to alimentary canal, 130, 134.
See Arteries, Veins, Capillaries.
- Blowing, 331.
- Boards of Health, duties of, 502.
- Body, parts of, 30.
- Body cavity in earthworms, 150.
- Body covering of lower animals, 374-376.
- Body framework, 221.
- Boils, 208.
- Bones, 221-243.
 composition of, 240.
 diseases of, 248-249.
 effects of age upon, 241.
 forms of, 236, 237.
 fractures of, 250.
 functions of, 222.
 regions of, 223.
 structure, 42, 43, 237-243.
- Bone cells, 243.
- Border cells, 113.
- Botany, 10.
- Bow legs, 249.
- Bowel movements, 145.
- Brain:
 development, 434.
 diseases, 424.
 external features of, 393.
 forebrain, 388.
 frog, 388.
 hindbrain, 388.
 human, 390.
 internal features of, 395.
 membranes, 390.
 midbrain, 388.
 structure, 392-397.
- Breastbone, 223.
- Breathing, 328.
 abdominal, 328.
 causes of, 322, 323.

Breathing—

- costal, 328.
- habits, 328.
- hygiene of, 331.
- modifications of, 341.
- mouth, 314.
- movements, 320.
- rate of, 324.
- relation of sex to, 324.
- Bright's disease, 373.
- Broken bones, 250.
- Bronchi, 315.
- Bronchial tubes, 316.
- Bronchitis, 208, 336.
- Buccal cavity, 93.
- Bunions, 249.
- Burning (combustion), 19.
- Burns and blisters, 365.

C.

- Cæcum, 128.
- Cæcapylonic in fish, 153.
- Calcium, 13, 26.
- Calices of kidney, 368.
- Canals, semicircular, 473, 477.
- Canalicule, 43, 241.
- Canines (teeth), 95, 96.
- Capacity of lungs, 329.
- vital, 329.
- Capillaries, 186, 187, 188.
- in the lungs, 319.
- seat of absorption, 134.
- structure of, 189.
- Capsules (bony), 42.
- Carbohæmoglobin, 166.
- Carbohydrates, 26, 53.
- as fuel, 60.
- digestion of, 132.
- Carbohydrates in muscle, 276.
- oxidation of, 27.
- Carbon, 16, 25, 26.
- Carbon dioxide, 16, 26.
- excretion by skin, 362.
- See Respiration.*

- Cardiac cycle, 192.
- Cardiac muscles, 266, 275.
- Cardiac nerves, 205.
- Cardiac orifice (of stomach), 111.
- Carnivora, teeth of, 155.
- Carotid artery, 199.
- Carpals (bones), 232.
- Cartilage, a form of connective tissue, 42.
- articulations, 227, 240.
- in bones, 222, 240.
- Casein and caseinogen, 26, 115.
- Cataract, 468.
- Catarrh, 208.
- Cavities, body, 29.
- Cells, air, 319.
- auditory, 476.
- defined, 33.
- Cells, forms of, 31.
- in tissues, 31.
- muscle, 282, 283, 284.
- nerve, 47, 405, 406.
- smell, 449.
- structural units, 32.
- structure of, 32.
- taste, 446.
- Central nervous system, 385.
- Centrum (of vertebræ), 225, 226.
- Cerebellum, 393.
- Cerebellum, function of, 418.
- Cerebral hemispheres, 392.
- distribution of gray matter in, 398.
- Cerebral localization, 419.
- Cerebro-spinal fluid, 398.
- Cerebrum, 392, 393.
- functions of, 418.
- Cervical vertebræ, 224, 226, 227.
- Charcoal, 16.
- Cheek bones, 234.
- Chemical compounds defined, 14.

- Chemical compounds, distinguished from mixtures, 14.
 Chemical compounds in the body, 25.
 Chemical compounds, relation to affinity, 13.
 Chemical elements, 12, 13, 28.
 Chest cavity, 320.
 Chiasma optic, 393.
 Chief cells, 113.
 Chocolate, 74.
 Choking, 331.
 Chordæ tendinæ, 185, 19.
 Choroid layer of eyes, 457.
 Choroid flexus, 398.
 Chyle, 141.
 Chyme, 116.
 Cigarettes, 88.
 Cilia in epithelial tissue, 41.
 in hydra, 148.
 use in digestion, 147.
 use in lungs, 318.
 Ciliary ligament, 457.
 Ciliary movement, 297-299.
 in protozoans, 300, 301.
 in higher forms, 302.
 Ciliary muscles, 458.
 Ciliary processes, 457.
 Circulation of blood, 180, 211.
 Circulation effects of alcohol upon, 210.
 in lower animals, 211, 219.
 portal, 200.
 pulmonary, 195.
 renal, 201.
 significance of, 218.
 systemic, 197.
 systems, defined, 180.
 Circumvallate papillæ, 445.
 Clavicles, 229, 230.
 Cloaca, 153, 381.
 Clotting of blood, 161, 169.
 Coal, 16.
 Coccyx, 225.
 Cochlea, 473.
 Cocoa, 74.
 Cœlenterates, 341, 375, 429.
 Cœliac axis, 199.
 Coffee, 74.
 Cold-blooded animals, 35.
 Colds, 208, 336.
 Collar bone, 229, 230.
 Colloids, 137.
 Colon, 128, 129.
 Color production, 466.
 blindness, 466.
 Combustion, 18, 19.
 in muscles, 282, 283.
 Complexion, care of, 364.
 Condiments, 73.
 Cones and rods, 465.
 Congestion, 206, 207.
 Conjunctiva, 453, 456.
 Connective tissue, 41, 42, 43.
 Consciousness, 418.
 Conservation of energy, 62.
 Constipation, 145.
 Contractility, 35.
 as a cause of locomotion, 297.
 of muscles, 28.
 Cooking, 70.
 Coral, skeleton of, 254.
 Corium, 355.
 Cornea, 456.
 Corns, 366.
 Coronary arteries, 186.
 system, 184, 199.
 veins, 184.
 Corpora quadrigemina, 397.
 Corpora striata, 395.
 Corpus callosum, 395.
 Corpuscles, bone, 42.
 red, blood, 160, 165, 167, 238.
 white, blood, 167.
 Cortex of brain, 405.
 Cortex of kidney, 368.
 Corti, fibers of, 474.
 organs of, 475.
 Costal cartilages, 229.
 Coughing, 331.

Cranial bones, 233.
 Cranial nerves, 388, 389, 394.
 Cranium, 234, 390.
 Cricoid cartilages, 487.
 Crop, 151, 156.
 Croup, 337.
 Crura cerebri, 393.
 Crustacea, skeleton of, 256.
 Crying, 331.
 Crypts, 120.
 Crystalloids, 137.
 Cud, 156.
 Cutaneous sensations, 438, 439.
 Cuticle, 354.
 Cuts, 209, 365.
 Cystic duct, 122.
 Cytoplasm, 34.

D.

Defæcation, 144.
 Deglutition, 107.
 Delirium tremens, 428.
 Dendrites, 47.
 Dental formulæ, 96.
 Dentine, 97.
 Dentition, 96.
 Dermis, 354-356.
 Dextrose, 54.
 Diabetes, 143.
 Dialysis, 136, 138.
 Dialyzer, 136.
 Diamond, 16.
 Diaphragm, 109, 180, 316, 320, 321.
 Diastase, 83.
 Diastole, 192.
 Diet, daily, 70.
 Digestibility of nutrients, 67.
 relation to food value, 67.
 Digestion, 90-132.
 chemistry of, 99.
 defined, 90.
 extra cellular, 149.
 hygiene of, 145.

Digestion —
 in lower animals, Chapter X, 146.
 of the mouth, 106.
 of the large intestine, 128, 129.
 of the small intestine, 118, 125-128.
 of the stomach, 114.
 peptic, 114-116.
 summary of, 132.
 Diphtheria, 208, 337.
 Disinfection, 500, 501.
 Dislocation of joints, 250.
 Distillation, 83.
 Distilled liquors, 83.
 Division of labor, 49.
 Dorsal vertebræ, 224-226.
 Ducts bile, 124.
 cystic, 124.
 thoracic, 135, 141.
 Duodenum, 119.
 Dura mater, 390.
 Dust and dusting, 334.

E.

Ears, bones of, 235.
 drum, 471.
 external, 471.
 hygiene of, 478.
 inner, 472.
 middle, 471.
 passages of, 474.
 structure and form, 470.
 vestibule, 473.
 wax, 471.
 Earthworm, blood in, 215.
 circulation of, 214.
 digestion of, 149.
 excretion in, 379.
 Earthworm respiration in, 343.
 Ectoderm, 375.
 Ectoplasm, 34.
 Elastic fiber, 41.

Elbow joint, 232.
 Elements, chemical, 12, 13, 28.
 Emulsion, 127.
 Enamel (tooth), 97.
 Endoplasm, 34.
 Endoskeleton, 221.
 Energy, conservation of, 62.
 definition of, 60.
 kinetic, 62.
 muscular, 282, 283.
 potential, 61.
 sources of, 59, 61.
 transformation of, 61.
 Enzymes, defined, 102.
 in muscle, 276.
 Epidermis, 353, 354.
 absence of blood in, 305.
 cells of, 354.
 nerves of, 355.
 Epiglottis, 108, 315, 487.
 Epithelial tissue, 39, 40.
 Erythroblasts, 168.
 Esophagus, action of, 109.
 structure of, 108.
 Ethmoid bone, 234.
 Eustachian tube, 107, 471, 472.
 Evolution, 38, 219.
 Excretion, 352-373.
 in lower animals, 374-382.
 Exercise, forms of, 286.
 relation to muscles, 285, 286.
 relation to breathing, 329.
 Exhaling, 319, 322, 323, 324.
 Exoskeleton, 221.
 Extractives, alcoholic, 84.
 Eye, 452-469.
 diseases, 469.
 functions of, 458.
 hygiene of, 467.
 muscles of, 454.
 position of, 452.
 structure of, 452-455.
 Eyeballs, 452-455.
 Eyesockets, bones of, 235.
 Eyelids, 453.

F.

Face bones, 233-235.
 Facial nerve, 389.
 Fæces, 129, 144.
 Fangs (tooth), 97.
 Far sight, 468.
 Fatigue muscle, 285.
 Fatigue, nerve, 423.
 retinal, 466.
 smell, 450.
 Fats, 26, 27, 60.
 absorption of, 140.
 digestion of, 126, 127.
 emulsification of, 127.
 forms of, 55.
 in bones, 241.
 in muscle, 276.
 saponification of, 126.
 tests for, 55.
 Fatty acids, 126.
 Feathers a form of skin, 377.
 Fehling's solution, 54, 55.
 Femur, 232.
 Fenestra ovalis, 474.
 rotunda, 471.
 Fermentation, 82, 83, 102.
 Fermented drinks, 83.
 Ferments, 102.
 Fibers, muscle, 302, 303.
 nerve, 402, 404.
 Fibrus and fibrinogen, 26, 162.
 Fibula, 232.
 Filiform papillæ, 445.
 Filtration, 139.
 Fingers, bones of, 233.
 Fissures of brain, 393.
 Fish, bones of, 257.
 circulation of, 215.
 digestion of, 152.
 gills of, 215, 348.
 heart of, 215.
 respiration of, 348.
 Flagella, 301.
 Flavors, 73.
 Flat bones, 239.

Floating ribs, 229.
 Fontanelle, 243.
 Foods, 51, 89.
 cost of, 68, 69.
 composition, 56-59.
 determination of fitness, 67.
 preservation, 499.
 values, 63.
 Food accessories, 73.
 Food stuffs, 53.
 Food vacuoles, 147.
 Foot, bones of, 233.
 Foramen magnum, 234, 390.
 Foramen Monroe, 395.
 Forebrain, 388.
 Fornix, 395.
 Fractures, 250.
 Freckles, 354.
 Frog, bones of, 257, 258.
 brain of, 387.
 circulation of, 216.
 digestion of, 153.
 heart of, 216.
 nerves of, 386, 387.
 respiration of, 348.
 Frontal bone, 234.
 Fuel, 60, 282.
 Fundus of stomach, 117.
 Function defined, 8.
 Fungiform papillæ, 445.

G.

Gall bladder, 122.
 Gases in the body, 25, 26.
 Gastric glands, 113.
 Gastric juice, 114.
 effect of alcohol upon, 130, 131.
 Gelatin, 26, 241.
 Germicides, 210.
 Germs, 493.
 Gills, 343, 144, 346-348.
 Gizzard, 151, 155.
 Glands, 99-101.
 in hydra, 149.

Glands —

 lachrymal, 453.
 lymphatic, 178.
 oil (*see Sebaceous*).
 salivary, 94, 102, 103, 104.
 sebaceous, 358.
 sweat, 357.
 wax, 471.
 Glomerulus of kidney, 370.
 Glossopharyngeal nerve, 389.
 Glottis, 107.
 Glucose, 54.
 Gluten, 52.
 Glycerine, 126.
 Glycogen, 27.
 in liver, 142, 143.
 in muscles, 276.
 Grape sugar, 54, 143.
 Graphite, 16.
 Gustatory sensations, 444.

H.

Habit, 413.
 Hæmocyannin, 344.
 Hæmoglobin, 165, 166.
 Hair, 358.
 care of, 364.
 follicles, 359.
 nerves of, 358.
 papillæ, 358.
 Hammer bone, 225.
 Hand, bones of, 233.
 Hard palate, 93, 235.
 Haversian canals, 43, 241, 242.
 Head, bones of, 233.
 Headache cures, 89.
 Health, Board of, 502.
 Hearing, 470.
 in lower animals, 482.
 Heart beat, 183, 191, 204, 205.
 muscle, 266, 275.
 nerves, 205.
 relation to lungs, 312, 316.
 sounds, 192, 193.
 structure, 180, 183.

Heat generation, 14, 19.
 relation to oxidation, 19, 27.
 sensation, 442.
 Heel bone, 233.
 Hepatic artery, 123.
 cells, 124.
 duct, 122.
 vein, 123, 200.
 Herbivora, 158.
 Hiccoughs, 332.
 Hindbrain, 388.
 Hinge joint, 245.
 Hip bones, 232.
 joint, 230, 231.
 Histology, 9, 39.
 Host, 495.
 Humerus bone, 232.
 Humor, aqueous, 457.
 vitreous, 457.
 Hunger, 117.
 Hydrochloric acid, 113, 116.
 Hydrogen, 15, 21, 22.
 Hygiene, 8, 9, 10.
 of bones and joints, 248.
 of circulation, 206.
 of digestion, 145.
 of the ears, 478.
 of the eyes, 467.
 of muscles, 285, 286.
 of nervous system, 422.
 of respiration, 330.
 public, 501, 502.
 Hyoid bone, 375.
 Hypodermis, 375.
 Hypoglossal nerve, 394.

I.

Ileocolic valve, 128.
 Ileum, 119.
 Iliac bones, 231.
 Images, 460, 463.
 Incisors (teeth), 95, 96.
 Incus bone, 472.
 Indigestion, 145.

Infection, 496.
 Inflammation, 206, 208.
 Infundibulum, 395.
 Inhaling, 319, 322-324.
 Inhibitory nerves, 205, 404.
 Insects, circulation of, 212, 213.
 excretion of, 381.
 respiration of, 346.
 Inspiration, 319, 322, 324.
 Instep, bones of, 233.
 Intercellular material, 44.
 Intercostal muscles, 317.
 Internal sensations, 444.
 Intestine, small, 118, 119-121.
 large, 128-130.
 Intestinal juice, 121, 127, 128.
 Intoxication, 424.
 Invertose, 128.
 Involuntary muscle, 266.
 Iris, 456, 457.
 Irritability, 35, 280.
 Ischium, 231.
 Iter, 396.

J.

Jaw bones, 235.
 Jejunum, 119.
 Joints, 244-251.
 diseases of, 249, 250, 251.
 dislocations of, 250.
 forms of, 244-248.
 structure of, 245.
 Jugular veins, 141, 200.

K.

Katabolism, 59, 352.
 Keratin, 26.
 Kidneys, 366-372.
 of insects, 381.
 of vertebrates, 38.
 sugar, disposal of, 143
 Knee joints, 232.

L.

- Labyrinth of ear, 473.
- Lachrymal bones, 235.
 - duct, 453.
 - glands, 453.
- Lacing, 249.
- Lacteals, 135, 141.
- Lactose, 54.
- Lacunæ (bone), 42, 43, 241.
- La grippe, 337.
- Lamellæ (bone), 43.
- Large intestine, 128, 129.
- Larynx, 315, 486, 487.
- Lateral ventricles, 395.
- Laughing, 331.
- Leather, 355, 378.
- Legs, bones of, 232.
- Lens, action of, 460.
 - crystalline, 457.
- Leucocytes, 168, 207.
- Levers, 288, 289.
- Ligaments, function of, 245.
 - transverse, 227.
- Light, effect of, 460, 461, 463.
 - nature of, 461.
- Lime, in bones, 240.
- Lime water and CO_2 , 16.
- Limula, 361.
- Lipase, 126.
- Litmus, 23.
- Liver, as an absorbent organ, 142.
 - as an excretory organ, 142, 372.
 - as a storehouse of energy, 143.
 - structure of, 121-123.
 - veins of, 201.
- Liver cells, 123.
- Liver starch, 142.
- Lobes, optic, 388, 392.
- Locomotion, causes of, 295.
 - due to contractility, 297.
 - due to secretion, 296.
 - due to specific gravity, 295.

- Locomotion, organs of, 305-309.
- Loudness, 489.
- Lumbar vertebræ, 225, 226, 227.
- Lungs, blood changes in, 326.
 - capacity of, 329.
 - capillaries of, 319.
 - development, 348-350.
 - excretory action of, 353.
 - structure of, 312-319.
- Lymph, 174-179.
 - capillaries, 177.
 - exchange, 175.
 - flow of, 177, 178.
 - formation of, 174.
 - glands, 178.
 - nodes, 178.
 - relation to absorption, 135, 175.
 - relation to assimilation, 175.
 - relation to osmosis, 175.
 - relation to plasma, 174.
 - tubes, 177.
 - valves, 177.
- Lymphatics, in villi, 133, 134.
 - system of, 176.
- Lymphocytes, 168.

M.

- Malars (bones), 234.
- Malleus (bone), 472.
- Malpighi, capsules of, 369.
 - pyramids of, 369.
 - tubules of, 369.
- Malt, 83.
- Manuals, 218, 261, 350.
- Man, biological position of, 37.
- Mandibles (bones), 235.
- Manganese dioxide, 18.
- Marrow (bone), 237.
- Maxillæ (bones), 235.
- Meats, cooking of, 71.
- Medulla oblongata, 394, 420.

Medullary layer of kidney, 369.
 sheath of nerves, 402.
 Membrane, mucous, 94, 314.
 Membranes of brain and cord,
 392, 400.
 of tympanum, 472.
 synovial, 246.
 Mercuric oxide, 17.
 Mesenteric arteries, 199.
 Mesentery, 119.
 Metabolism, 59, 352.
 Metacarpals (bones), 232.
 Metatarsals (bones), 232.
 Metallic elements, 23.
 Microbes (*see Bacteria*).
 Midbrain, 388.
 Milk, 57, 64, 248.
 Milk glands, 377.
 Milk teeth, 96.
 Millon's reagent, 53.
 Minerals, 24, 25.
 Mineral salts, 55, 56.
 Mitral valves, 183, 186, 191.
 Mixtures distinguished from
 chemical compounds, 14.
 Molars (teeth), 95, 96.
 Mollusks, 255, 375, 430.
 Motor nerves, 403.
 Mouth digestion, 106.
 Movement, amœboid, 297, 298.
 ciliary, 297, 299.
 in lower animals, Chapter XX,
 294.
 muscular, 297.
 Mucous membrane, 94, 314.
 Mucus, 94.
 Muscles, 265-293.
 action of, 278, 287, 288.
 attachment to bones, 240,
 268.
 bellied, 267.
 biceps, 206, 209.
 comparison with engine, 281-
 283.
 composition of, 276, 277.
 contraction of, 278, 281.

Muscles —
 digastric, 268.
 efficiency of, 287.
 fatigue of, 285.
 forms of, 267-269.
 heart, 266.
 involuntary, 265, 266.
 nerve endings in, 279.
 non-striated, 266, 274, 275.
 papillary, 185.
 rib, 317.
 striated, 266, 267, 269.
 triceps, 269.
 voluntary, 265, 266, 274.
 Muscle cells, 282, 283, 284.
 Muscle fibers, 272, 273, 305.
 fibrils, 274.
 Muscular movements, 297-302,
 303, 304.
 Muscular tissue, 44, 265, 266.
 Musical sounds, 489.
 Myogen, 277.
 Myopia, 467.
 Myosin, 26, 276.

N.

Nails, 359, 360, 365.
 Narcotics, 86.
 Nasal (bones), 235.
 Near sight, 467.
 Nephridia, 379.
 Nerve cells, 47, 405, 406.
 Nerve control, 291.
 Nerve diseases, 424.
 Nerve endings in muscle, 279.
 Nerve fibers, 402-404.
 excitatory, 404.
 inhibitory, 404.
 medullated, 403.
 Nerve muscle preparation, 279.
 Nerve processes, 47.
 Nerves, afferent, 403.
 cranial, 389.
 dermal, 356.

Nerves —

- efferent, 403.
- epidermal, 354.
- motor, 403.
- sensory, 403.
- vaso-motor, 205.
- Nervous system, 382-428.
 - central, 385.
 - composition, 385.
 - diseases of, 424.
 - effects of alcohol upon, 424.
 - functions of, 411.
 - hygiene of, 422.
 - in lower animals, Chapter XXVII, 429.
 - sympathetic, 385.
- Nervous tissue, 46.
- Neura, 47, 405, 406.
 - theory of action, 406-409.
- Neural arch, 226.
- Neural cavity, 225, 226, 397.
- Neuralgia, 424.
- Neurilemma, 403.
- Neuron (*see* *Neura*).
- Neutralization, 23, 24.
- Nicotine, 87.
- Nitrogen, 20.
- Nitrogenous compounds, 26, 52.
- Nodes of Ranvier, 402.
- Non-metallic elements, 23.
- Non-nitrogenous compounds, 26.
- Non-striated muscle, 44, 45, 266, 274, 275.
- Nose, bones, 235.
- Nose, sense organs of, 448.
- Nucleus, 32, 34.
- Nucleolus, 34.
- Nutrients, 52.
 - absorption of, 139.
 - forms, 52,
 - in blood, 163.
 - in bone, 241.
 - in foods, 57.
 - sources of, 56.
 - uses of, 59.
- Nutrition, 51-72.

O.

- Oblique muscles, 110.
- Occipital bone, 234.
- Ocelli, 483, 484.
- Oculo-motor nerve, 388.
- Odontoid, process, 227.
- Odors, 450.
- Olfactory lobes, 392, 393.
 - nerves, 388.
 - sensations, 448.
- Omentum, 111.
- Opium, 88, 89.
- Optic chiasma, 393.
 - lobes, 388, 392.
 - nerve, 388.
 - thalami, 396.
- Orbit of eye.
- Organic chemistry, 16.
 - compounds, 26.
- Organs, 30, 31.
 - of Corti, 475.
- Osmosis, 136, 138, 163.
- Otocysts, 483.
- Otoliths, 473.
- Oxidation, 18, 19, 282 (*see* *Respiration*).
- Oxides, 19, 25.
- Oxygen, 13, 17, 18, 19, 26 (*see* *Respiration*).
- Oxyhæmoglobin, 166.

P.

- Pain, sensations, 443.
- Palate, bones, 235.
 - hard, 93, 235.
 - soft, 93,
- Pancreas, 124, 125.
- Pancreatic juice, 125, 127.
- Papillæ, dermal, 355, 356.
 - hair, 358.
 - nail, 360.
 - taste, 445.
- Papillary muscles, 185.

Paramœcium, 146, 147.
 Parasite, 495.
 Parietal bones, 234.
 Parotid glands, 102.
 Patent medicines, 64.
 Pectoral girdle, 223, 230.
 Pelvic girdle, 230, 231.
 Pelvis, 223, 232.
 of kidney, 368.
 Pepsin, 113, 115.
 Peptic digestion, 117.
 Peptone, 115, 137, 140.
 Pericardium, 181.
 Perilymph, 473.
 Perimysium, 270, 271.
 Perineurium, 402.
 Periosteum, 237, 240, 241.
 Peristalsis, 109, 125.
 Peritoneum, 109.
 Perspiration, 357, 362, 363.
 Phagocytes, 169.
 Phalanges, 232.
 Pharynx, 94, 107.
 Phosphorus, 13.
 Phrenology, 419.
 Physiology, 7, 8, 9, 10.
 Physiological chemistry, 10.
 Pia mater 392.
 Pigment cells, 354.
 Pitch, 489.
 Pivot joint, 247.
 Plant physiology, 11.
 as a source of nutrients, 58.
 Plasma, 161.
 Pleura, 319.
 Pleurisy, 336.
 Plexus, 404, 405.
 Pneumonia, 208, 336.
 Poisons, 144.
 Pons varolii, 393.
 Portal circulation, 200, 201.
 Portal vein, 123, 134, 142.
 Potassium chlorate, 17.
 Premolars (teeth), 95, 96.
 Presbyopia, 469.
 Pressure sensations, 440, 441.

Proteid, 52.
 absorption of, 138, 139.
 action of trypsin upon, 126.
 action of gastric juice upon,
 114-116.
 blood, 163, 164.
 composition of, 26.
 digestion of, 126, 114-116.
 forms of, 26, 52.
 muscle, 276.
 sugar formers, 144.
 tests for, 53.
 Protoplasm, 32-36.
 movement of, 294.
 Protozoans, 252, 341, 374, 429,
 480.
 Psalterium, 157.
 Pseudopodia, 35, 298.
 Ptyalin, 105.
 Pubis (bones), 231.
 Pulmonary artery, 185, 195.
 capillaries, 196.
 sacs, 316-319.
 veins, 185, 197.
 Pulp cavity (tooth), 97.
 Pulse, 193, 194, 195.
 Pupil of eye, 457.
 Pure Food Law, 84.
 Pus, 207.
 Pyloric orifice, 111.
 Pylorus, 112.
 Pyramids of Malpighi, 369.

Q.

Quality of sound, 490.

R.

Radial artery, 199.
 Radius bone, 232.
 Rectum, 128.
 Reflex action, 412.
 arc, 413.

Renal artery, 201, 367, 368.
 circulation, 201.
 veins, 201, 367, 368.
 Rennin, 113.
 Reproduction of cells, 50.
 Reptiles, 217, 259, 349.
 Residual air, 329.
 Resonance, 490.
 Respiration (*see Breathing*),
 312-339.
 artificial, 332, 333.
 effects of alcohol upon, 338.
 in lower animals, Chapter
 XXII, 340.
 organs of, 313, 314, 320, 324.
 Respiratory organs, 313, 314.
 diseases of, 335.
 Rest, 423.
 Reticulum, 157.
 Retina, 457, 463-465.
 Ribs, action in respiration, 320.
 cartilages of, 229.
 floating, 229.
 muscles of, 317.
 processes, 229.
 relation to thorax, 223.
 types of bone, 238, 239.
 Rickets, 248.
 Rigor mortis, 276.
 Rodents, teeth of, 155.
 Rods and cones, 465.
 Rods of Corti, 475.
 Round shoulders, 249.
 Rumen, 157.
 Ruminants, 156, 157.
 Running, 290, 291.

S.

Saccharose, 54.
 Sacculus, 473.
 Sacrum, 225, 231.
 Saliva, action of, 105.
 composition, 94, 104.
 effect of alcohol upon, 131.

Salivary glands, 94, 102, 103,
 104.
 Salts, absorption of, 141.
 blood, 163, 164.
 characteristics, 23-25.
 Sanitation, 10, 493, 501, 502.
 Saponification, 126.
 Saprophyte, 495.
 Sarcolemma, 45, 272, 273, 274,
 277.
 Sarcoplasm, 45, 273.
 Sarcostyles, 274.
 Scapula, 230, 232.
 Sclerotic layer of eye, 456.
 Sebaceous glands, 356, 358,
 471.
 Secretion, 99.
 a property of epithelium, 40.
 as a cause of motion, 296.
 Semicircular canals, 473, 477.
 Semilunar valves, 185, 186, 191,
 192.
 Selection of food, 36.
 Sensation, 436-485.
 auditory, 470.
 classification of, 438.
 cutaneous, 438.
 gustatory, 444.
 in lower animals, 480-485.
 internal, 444.
 olfactory, 448.
 pain, 443.
 pressure, 440.
 temperature, 442.
 visual, 452.
 Senses, special (*see Sensation*).
 Sensory organs, development of,
 481.
 Septa, lucida, 395.
 Serous membrane, defined, 110.
 Serum, 162, 163.
 Sheath, medullary, 402.
 Shoulder blades, 230.
 Shoulder joints, 232.
 Sighing, 332.
 Sight sensation, 452, 465.

- Sigmoid flexure, 128.
 Silver sulphide, 15.
 Skeleton, 221-251.
 appendicular, 230.
 axial, 224.
 bones of, 222.
 external, 29, 221, 252.
 hygiene of, 248.
 in lower animals, 252-263.
 internal, 29, 221, 252.
 Skin, 353-363.
 as an excretory organ, 361.
 of lower animals, 374-382.
 functions of, 361.
 hygiene of, 363.
 modifications of, 353-356.
 structure of, 353.
 Skull, 223, 233.
 uses of, 235, 236.
 Smell, sensation of, 448-450.
 in lower animals, 482.
 Snails, 345.
 Sneezing, 331.
 Soaps, 126, 140.
 Sobbing, 331.
 Sore throat, 208, 335.
 Sound, 474, 489, 490.
 Special senses (*see Sensation*).
 Speech, 486.
 Sphenoids, 234.
 Sphincter pylorus, 112, 117.
 Spinal accessory nerve, 394.
 Spinal column, 222, 224, 228.
 Spinal cord, 398.
 center reflex action, 417.
 connection with sympathetic system, 401.
 diseases of, 424.
 functions of, 417.
 membranes of, 399.
 of the frog, 389, 390.
 structure, 401.
 Spinal nerves, 400, 401, 402.
 Spinous process, 226.
 Spiracles, insect, 346.
 Spitting, 332.
 Spleen, 168.
 Sponges, 148, 212, 237, 253, 341, 378.
 Sprains, 250.
 Spyrogyra, 32.
 Squameous epithelium, 40.
 Standing, 290, 291.
 Stapes (bone), 235, 472.
 Starch, absorption, 140.
 composition, 27, 53, 54.
 digestion of, 105, 126.
 in blood, 163.
 Starfish, 254, 342.
 Steapsin, 125, 126.
 Sterilizing, 500.
 Sternum, 228, 229.
 Stimulants, 74.
 Stirrup bones, 235.
 Stomach, as a storehouse, 116.
 curvatures of, 112.
 enzymes of, 113.
 glands of, 113.
 method of action, 117.
 structure, 111.
 Strangulation, 331.
 Striated muscle, 266, 267, 269, 270, 272-274.
 Stroma, 166.
 Structure, 8, 28.
 Subcutaneous layer, 355.
 Sublingual glands, 102.
 Submaxillary glands, 102.
 Submucous coat, 111.
 Succus entericus, 127.
 Suffocation, 331.
 Sugar, composition, 27.
 forms, 53-55.
 in blood, 163, 164.
 in muscle, 277.
 tests for, 54, 55.
 Sulphur, 15.
 Suspensory ligament, 458.
 Sutures, 235, 243, 247.
 Swallowing, 107.
 Sweat, 356.
 Sweat glands, 356, 357.

Sympathetic nervous system, 421.
 Symphyses, 243, 247.
 Synovial fluid, 245, 246.
 membrane, 246.
 Systematic circulation, 198-200.
 Systems, defined, 30.
 Systole, 192.

T.

Tarsals (bones), 232.
 Taste sensation, 444-447.
 Taste buds, 446.
 Taste in lower animals, 482.
 Tea, 74.
 Tears, 453.
 Tear glands, 453.
 Teeth, 94-96.
 in lower forms, 152-155.
 structure, 97.
 use in digestion, 98.
 Temperature, body, 19, 171,
 172, 363.
 sensations, 442.
 Temporals (bones), 234.
 Tendons, 238, 267, 271.
 Theobromin, 74.
 Thoracic duct, 135, 141.
 Thorax, 223.
 Thyroid cartilage, 487.
 Thumb, bones of, 233.
 Tibia (bone), 232.
 Tidal air, 330.
 Time reactions, 426.
 Tissue, 31, 39, 48.
 Tobacco, 87, 88, 284.
 Toes, 233.
 Tongue, 94, 99, 444, 445.
 Tonsils, 94.
 Tonsillitis, 335.
 Touch, 440, 481.
 Tourniquet, 209.
 Toxins, 497.
 Trachea, 315, 317, 318.

Tracheæ, insect, 346.
 Triceps muscle, 269.
 Tricuspid valves, 183, 185,
 191.
 Trigeminal nerve, 389.
 Trochlear nerves, 389.
 Trunk, 29.
 Trypsin, 125, 126.
 Tuberculosis, pulmonary, 337.
 of the joints, 251.
 Turbinate bones, 235.
 Tympanum membrane, 471.
 Tympanum, 471.
 Typhlosole, 150.

U.

Ulna, 232.
 Ulnar artery, 199.
 Urea, 26, 352, 357, 362, 371,
 372.
 Ureter, 366.
 Urethra, 367.
 Urine, 371.
 Urinary organs, 366.
 Utriculus, 473.
 Uvula, 94.

V.

Vacuoles, 34.
 Vagus nerves, 389.
 Valvulæ conniventes, 120.
 Vaso-motor nerves, 205.
 Vegetables, cooking of, 71.
 Veins, 186-190.
 Vena cava, 184, 191, 197, 200.
 Venous blood, 203.
 Ventilation, 333.
 Ventricles (heart), 183, 148,
 191, 192.
 Ventricles of the brain, 395,
 396, 397.
 Vermiform appendix, 129.

Vertebrae, 222-240.
 cervical, 224, 226, 227.
 dorsal, 224, 225, 226.
 lumbar, 225, 226, 227.
 Vertebrates, 222, 252, 376, 433:
 Vestibule of ear, 473.
 Vestibular passage, 474, 476.
 Villi, 120, 134, 135.
 Viscera, 29.
 Vision, 452.
 in lower animals, 483.
 Visual sensation, 452.
 Vocal cords, 488.
 organs, 486.
 Vocalizing, 488.
 Voice, 486-492.
 Voice, hygiene of, 491.
 Voluntary action, 412.
 arc, 415.
 muscle, 266.
 Vomer, 235.

W.

Walking, 290, 291.
 Warm blood, 351.

Water, absorption of, 141.
 as a nutrient, 56.
 breathers, 341.
 composition, 22, 25.
 Wax glands, 471.
 Wind pipe, 315.
 Wisdom teeth, 96.
 Worms, 343, 375, 432.
 Wrinkles, 356.
 Wrist, 233.

X.

Xanthoproteic test, 53.
 X-ray, 220-221.

Y.

Yawning, 332.
 Yellow spot of eye, 465.

Z.

Zoölogy, 10.

OUTLINES OF BOTANY

\$1.00

By ROBERT GREENLEAF LEAVITT, A.M., of
the Ames Botanical Laboratory. Prepared at the request
of the Botanical Department of Harvard University

Edition with Gray's Field, Forest, and Garden Flora \$1.80
Edition with Gray's Manual of Botany 2.25

THIS book covers the college entrance requirements in botany, providing a course in which a careful selection and a judicious arrangement of matter is combined with great simplicity and definiteness in presentation.

¶ The course offers a series of laboratory exercises in the morphology and physiology of phanerogams; directions for a practical study of typical cryptogams, representing the chief groups from the lowest to the highest; and a substantial body of information regarding the forms, activities, and relationships of plants and supplementing the laboratory studies.

¶ The work begins with the study of phanerogams, taking up in the order the seed, bud, root, stem, leaf, flower, and fruit, and closing with a brief but sufficient treatment of cryptogams. Each of the main topics is introduced by a chapter of laboratory work, followed by a descriptive chapter. Morphology is treated from the standpoint of physiology and ecology. A chapter on minute structure includes a discussion of the cell, while another chapter recapitulates and simplifies the physiological points previously brought out.

¶ The limitations of the pupil, and the restrictions of high school laboratories, have been kept constantly in mind. The treatment is elementary, yet accurate; and the indicated laboratory work is simple, but so designed as to bring out fundamental and typical truths. The hand lens is assumed to be the chief working instrument, yet provision is made for the use of the compound microscope where it is available.

AMERICAN BOOK COMPANY

A NEW ASTRONOMY

\$1.30

By DAVID TODD, M.A., Ph.D., Professor of Astronomy and Navigation, and Director of the Observatory, Amherst College.

ASTRONOMY is here presented as preëminently a science of observation. More of thinking than of memorizing is required in its study, and greater emphasis is laid on the physical than on the mathematical aspects of the science. As in physics and chemistry the fundamental principles are connected with tangible, familiar objects, and the student is shown how he can readily make apparatus to illustrate them.

¶ In order to secure the fullest educational value astronomy is regarded, not as a mere sequence of isolated and imperfectly connected facts, but as an inter-related series of philosophic principles. The geometrical concept of the celestial sphere is strongly emphasized; also its relation to astronomical instruments. But even more important than geometry is the philosophical correlation of geometric systems. Ocean voyages being no longer uncommon, the author has given rudimental principles of navigation in which astronomy is concerned.

¶ The treatment of the planets is not sub-divided according to the planets themselves, as is usual, but according to special elements and features. The law of universal gravitation is unusually full, clear, and illuminating. The marvelous discoveries in recent years and the advance in methods of teaching are properly recognized, while such interesting subjects as the astronomy of navigation, the observatory and its instruments, and the stars and the cosmogony receive particular attention.

¶ The illustrations demand special mention; many of them are so ingeniously devised that they explain at a glance what many pages of description could not make clear.

AMERICAN BOOK COMPANY

TEXT-BOOKS ON GEOLOGY

By JAMES D. DANA, LL.D., late Professor of Geology
and Mineralogy, Yale University

Geological Story Briefly Told	\$1.15
Revised Text-Book of Geology. Fifth Edition (Rice)	1.40
Manual of Geology. Fourth Edition	5.00

THE present edition of the GEOLOGICAL STORY was the last considerable work of the eminent author's long life. As geology is emphatically an outdoor science, the student is urged to study the quarries, bluffs, and ledges of rock in his vicinity, and all places that illustrate geological operations. The prefatory suggestions are full of practical help, besides enumerating the tools and specimens desirable for this study.

¶¶ The REVISED TEXT-BOOK OF GEOLOGY is here brought down to the present time as regards its facts, but it still expresses the views of its distinguished author. While the general plan and the distinctive features have been preserved as far as possible, a few important changes have been made relating mainly to zoölogical and botanical classifications, to the bearings of geology and paleontology upon the theory of evolution, and to metamorphism. The order of the grand divisions of the science, physiographic, structural, dynamical and historical, remains the same as in previous editions.

¶¶ The present edition of the MANUAL OF GEOLOGY was wholly rewritten under the author's direction. Owing to the extensive recent investigations, new principles, new theories, new facts relating to all departments of the science, and widely diverging opinions on various questions have all made their contributions to this book. The later tracing of formations and mountain-making, the increased number of fossils, the study of cañons and other results of erosion, and the development of petrology are all prominently treated here.

AMERICAN BOOK COMPANY

C H E M I S T R I E S

By F. W. CLARKE, Chief Chemist of the United States Geological Survey, and L. M. DENNIS, Professor of Inorganic and Analytical Chemistry, Cornell University

Elementary Chemistry . \$1.10

Laboratory Manual . . \$0.50

THESE two books are designed to form a course in chemistry which is sufficient for the needs of secondary schools. The TEXT-BOOK is divided into two parts, devoted respectively to inorganic and organic chemistry. Diagrams and figures are scattered at intervals throughout the text in illustration and explanation of some particular experiment or principle. The appendix contains tables of metric measures with English equivalents.

¶ Theory and practice, thought and application, are logically kept together, and each generalization is made to follow the evidence upon which it rests. The application of the science to human affairs, its utility in modern life, is also given its proper place. A reasonable number of experiments are included for the use of teachers by whom an organized laboratory is unobtainable. Nearly all of these experiments are of the simplest character, and can be performed with home-made apparatus.

¶ The LABORATORY MANUAL contains 127 experiments, among which are a few of a quantitative character. Full consideration has been given to the entrance requirements of the various colleges. The left hand pages contain the experiments, while the right hand pages are left blank, to include the notes taken by the student in his work. In order to aid and stimulate the development of the pupil's powers of observation, questions have been introduced under each experiment. The directions for making and handling the apparatus, and for performing the experiments, are simple and clear, and are illustrated by diagrams accurately drawn to scale.

AMERICAN BOOK COMPANY

A BRIEF COURSE IN GENERAL PHYSICS

\$1.20

By GEORGE A. HOADLEY, A.M., C.E.,
Professor of Physics, Swarthmore College

A COURSE, containing a reasonable amount of work for an academic year, and covering the entrance requirements of all of the colleges. It is made up of a reliable text, class demonstrations of stated laws, practical questions and problems on the application of these laws, and laboratory experiments to be performed by the students.

¶ The text, which is accurate and systematically arranged, presents the essential facts and phenomena of physics clearly and concisely. While no division receives undue prominence, stress is laid on the mechanical principles which underlie the whole, the curve, electrical measurements, induced currents, the dynamo, and commercial applications of electricity.

¶ The illustrative experiments and laboratory work, introduced at intervals throughout the text, are unusually numerous, and can be performed with comparatively simple apparatus. Additional laboratory work is included in the appendix, together with formulas and tables.

HOADLEY'S PRACTICAL MEASUREMENTS IN MAGNETISM AND ELECTRICITY. \$0.75

THIS book, which treats of the fundamental measurements in electricity as applied to the requirements of modern life, furnishes a satisfactory introduction to a course in electrical engineering for secondary and manual training schools, as well as for colleges. Nearly 100 experiments are provided, accompanied by suggestive directions. Each experiment is followed by a simple discussion of the principles involved, and, in some cases, by a statement of well-known results.

AMERICAN BOOK COMPANY

DESCRIPTIVE CATALOGUE OF HIGH SCHOOL AND COLLEGE TEXT-BOOKS

Published Complete and in Sections

WE issue a Catalogue of High School and College Text-Books, which we have tried to make as valuable and as useful to teachers as possible. In this catalogue are set forth briefly and clearly the scope and leading characteristics of each of our best text-books. In most cases there are also given testimonials from well-known teachers, which have been selected quite as much for their descriptive qualities as for their value as commendations.

¶ For the convenience of teachers this Catalogue is also published in separate sections treating of the various branches of study. These pamphlets are entitled: English, Mathematics, History and Political Science, Science, Modern Languages, Ancient Languages, and Philosophy and Education.

¶ In addition we have a single pamphlet devoted to Newest Books in every subject.

¶ Teachers seeking the newest and best books for their classes are invited to send for our Complete High School and College Catalogue, or for such sections as may be of greatest interest.

¶ Copies of our price lists, or of special circulars, in which these books are described at greater length than the space limitations of the catalogue permit, will be mailed to any address on request.

¶ All correspondence should be addressed to the nearest of the following offices of the company: New York, Cincinnati, Chicago, Boston, Atlanta, Dallas, San Francisco.

AMERICAN BOOK COMPANY

Cornell University Library

arV141

A text-book in general physiology and an

3 1924 031 168 705

olin,anx