New York
State College of Agriculture
At Cornell University
Ithaca, N. Y.

Library

Cornell University Library
SH 11.V5R

Report.

3 1924 003 251 778

mann

Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

REPORT OF COMMISSIONERS

RELATIVE TO THE

RESTORATION OF SEA-FISH

TO THE

CONNECTICUT RIVER AND ITS TRIBUTARIES.

By Order of the Legislature of Vermont.

ANNUAL SESSION, 1866.

MONTPELIER :
FREEMAN STEAM PRINTING ESTABLISHMENT.
1866.

REPORT OF COMMISSIONERS

RELATIVE TO THE

RESTORATION OF SEA-FISH

TO THE

CONNECTICUT RIVER AND ITS TRIBUTARIES.

By Order of the Legislature of Vermont.

ANNUAL SESSION, 1866.

MONTPELIER :
FREEMAN STEAM PRINTING ESTABLISHMENT.
1866.

IN SENATE, October 16, 1866.

On motion of MR. BARSTOW, the Report of Albert D. Hager and Hon. Charles Barrett, Commissioners on the restoration of sea-fish to the Connecticut river, was ordered to lie, and the Secretary directed to procure the printing of one thousand copies for the use of the General Assembly.

HENRY CLARK, *Secretary*.

REPORT.

To his Excellency, Paul Dillingham,

Governor of the State of Vermont:

SIR :—The commissioners, appointed in pursuance of the joint resolution of the Legislature of 1865, relative to the restoration of sea-fish and the introduction of new varieties of fresh-water fish into the waters of the Connecticut and its tributaries, would respectfully submit the following report :

In order to accomplish successfully the desired object of restocking the Connecticut river and its tributaries with migratory fish, it seemed necessary that there should be a hearty co-operation and concurrent legislation on the part of the states of Vermont, New Hampshire, Massachusetts and Connecticut.

To secure this it was deemed advisable to confer with the other commissioners that had been appointed, and therefore by previous arrangement we met Hon. H. A. Bellows and Capt. W. A. Sanborn, commissioners for New Hampshire, and Col. Theodore Lyman, one of the commissioners for Massachusetts, at Lawrence, Mass., on the 30th day of November last. At this meeting the commissioners, together with his Excellency, Governor Smyth, of N. H., and several other persons whom we met, manifested much interest in the object for which the meeting was called, and expressed an

earnest desire for the return of shad and salmon to the rivers of New England, as in the days of our ancestors.

Numerous instances were cited where experiments had been made in Great Britain and France to restock streams that had become nearly or quite destitute of edible fish. In every instance where the efforts had been judiciously made the results were successful. The barren streams in many instances had become sources of wealth to the nation, and furnished an immense amount of wholesome food for the inhabitants thereof. Is it placing too high an estimate upon Yankee enterprise and New England character to say that as good and even better results may follow similar efforts to restock the streams of New England?

During the session of the legislature of Massachusetts in April last, we visited Boston twice, and appeared before the legislative committee to whom was referred the subject of restocking the Merrimac and Connecticut rivers with sea-fish. That committee and the leading members of both branches of the legislature seemed favorably impressed with the importance and feasibility of the project, provided it had the concurrent legislation and assistance of the other states interested.

But it could not be expected that the legislature of Massachusetts would cause the erection and maintenance of fishways over the dams at Lowell, Lawrence, Holyoke and Turner's Falls, which at a moderate estimate would cost over twenty thousand dollars, unless the other states interested signified a willingness to co-operate in the movement. The legislature manifested a desire to cause the fishways to be built, if in return Vermont and New Hampshire would supply and suitably protect the young fish, and Connecticut would cause the erection of a fishway at Enfield Falls, and enact and enforce laws to prohibit the fearful and unrestricted slaughter made with gill-nets within her borders.

We, as commissioners, felt unauthorized to make any *pledges*, but expressed the opinion that Vermont would cheer-

fully do her duty in the matter. Judge Bellows expressed the same opinion concerning New Hampshire. At that time Connecticut had no commissioners upon the subject, therefore no expression of opinion for that state was given.

Before Massachusetts could act consistently in the matter of building fishways over dams on the Connecticut river, it seemed desirable that she should receive some assurance of co-operation from Connecticut. This was impossible until the meeting of the Connecticut legislature, which would be after the legislature of Massachusetts had adjourned. But the legislature of Massachusetts, with its accustomed willingness to advance a worthy object, enacted a law providing for the construction and maintenance of fishways on the Merrimac and Connecticut rivers, but inserted a provision as follows :

"SEC. 13. Said commissioners may in their discretion delay the definite construction of fishways on the Connecticut river, until they shall be satisfied that such legislation has been adopted by the state of Connecticut, as shall in the opinion of said commissioners be necessary to secure the free passage of the fish above named through that part of said river as runs through said state of Connecticut."

The legislature also adopted the following preamble and resolution :

"WHEREAS, The Connecticut river, in this state, formerly furnished an inexhaustible supply of shad and other kinds of migratory fish, which have now partially disappeared from that portion of said river within this state ; and

WHEREAS, There is nothing to prevent the return of said fish from the sea to our waters, except the want of suitable fishways over the dam at Enfield Falls, in the state of Connecticut, in such numbers as largely to contribute to the supply of food for the inhabitants of this state ; therefore,

Resolved, That the attention of the state of Connecticut be invited to this subject, and that it be earnestly requested to take early measures to cause fishways to be constructed over the dam at Enfield Falls ; either requiring this work to be done by the proprietors of said dam or by the exercise of the right of eminent domain, as due alike to the relations

of comity between that state and this Commonwealth, to the obligations of international law, and to the local interests of that state.

Resolved, That his Excellency, the Governor, is hereby requested to transmit a copy of these resolutions to the governor of the state of Connecticut, with a request that the same be laid before the legislature of the same."

Col. Theodore Lyman of Brookline and Hon. Alfred R. Field of Greenfield were appointed commissioners to superintend the construction and maintenance of fishways in Massachusetts, and are to hold their offices for the term of five years. A fishway at Lowell twelve feet wide is completed, and the one at Lawrence is being built, and will be completed before the stream is closed with ice. The cost of the latter is estimated at \$8,500.

The legislature of Connecticut, at its session in June last, adopted a resolution empowering the governor of the state to appoint two commissioners to confer with similar appointees of Vermont, New Hampshire and Massachusetts, concerning the introduction and protection of sea-fish in the Connecticut river. In pursuance of this his Excellency, Governor Hawley, appointed as said commissioners Messrs. J. Hammond Trumbull of Hartford, and William H. Goodspeed of East Haddam.

In a communication apprising us of the appointments that had been made, Governor Hawley expressed confidence in the enterprise, and said he should "take great pleasure in aiding in the attainment of the ends sought by the general movement."

The commissioners of New Hampshire have been authorized to co-operate with those of Massachusetts in restocking the Merrimac river with shad and salmon, and also the Connecticut river if the other states in interest do their duty in the matter. At this time they have a man in their employ in New Brunswick to procure salmon spawn, and they are also

adopting measures for the construction of fishways on the Merrimac and its tributaries in that state.

From the foregoing it appears that our sister states are in earnest upon this subject. The interest and unity of feeling manifested induces the belief that concurrent legislation and united efforts on the part of all the states in interest will accomplish the desired result.

The willingness and promptitude with which Vermont has ever responded to all reasonable calls that have been made, induce the belief that it is needless for us to suggest that the legislature should follow the noble examples set by the legislatures of Massachusetts and New Hampshire, and provide for the introduction of spawn into all streams suitable for either shad or salmon and also to enact stringent laws for the suitable protection of fish.

It may be proper in this place to present a few facts relative to shad and salmon, upon which our belief in the feasibility of the project proposed is predicated.

Shad and salmon are gregarious and migratory. During the winter months they live in salt water and grow very rapidly. In the spring they are very fat and invariably return to spend the summer in their native streams. The females that are filled with spawn are said to be the first to resort to the fresh water. In a few weeks the males follow them. It is mainly from the fact that these migratory fish always return to fresh water to deposit their spawn and *to the identical spawning beds where they were hatched*, that we predicate our belief that with suitable effort they may again become abundant in the Connecticut and its tributaries.

The Germans and French were the first to engage in artificial fish culture. The latter have carried it to a great degree of perfection. During the present year there has been holden, under the auspices of the *Scientific Society of Arcachon* with the co-operation of the French Government, a National Fair or exposition of fishery and water culture at Arcachon. A report of this exposition will be addressed to

the Imperial Commission of the Universal Exposition of Paris in 1867, where it is expected the contributions resulting from the fish culture in France will form an important feature. The council of administration or directors under whose supervision the exposition was holden, in their circular say: "If we have borrowed from England the improved methods of farming, it is to France that the English have come to gain the very idea of water culture. So we may hope that the imperial commission of the coming universal exposition of Paris will think proper to make use of the exposition of Arcachon, to arrive in 1867, in the section of fishery and water culture, at the same remarkable result attained by the Imperial Commission of the universal exposition of London in 1862, for the section of agriculture."

The experiment of fish culture has also been successfully made in Great Britain. In a few instances streams that never had these fish in them until the spawn were introduced and fishways built, have become sources of wealth.

The head waters of the Claregalway in Ireland was cut off from the sea by a natural fall in the stream so that no sea-fish were able to ascend to them. But a fishway has been built and salmon spawn introduced, and the river affords one of the best salmon fisheries in Great Britain. The Robe river that empties into the Lough Mask in Ireland had no sea-fish in it till a fishway about two miles in length was built over the natural obstruction in the stream, and salmon ova were placed there and hatched. It now abounds in salmon.

The Tay river in Scotland formerly produced salmon, but being unprotected by suitable laws and regulations they became nearly extinct in that stream. But the persevering efforts of Messrs. Buist and Brown, who propagated the young fish at their establishment at Stormonfield, and the stringent laws relating to fishing in that country, have had the effect to make salmon very abundant in that stream. In

1863 the streams of Scotland produced salmon to the value of £500,000, and the salmon caught that year in Ireland were estimated at 300,000 pounds sterling.*

We will cite one more instance touching this point, taken from "*The American Angler's Book*," by Norris (pp. 233-4) as follows :

Loch Shin, a piece of water about twenty-one miles by fourteen, situated in the heart of the Sutherland Mountains, is the immediate feeder of the River Shin, noted for its salmon fecundity. The loch itself has four feeders, middling sized rivers, viz : the Terry, Fiack, Garvie, and Curry, in which, previously to the year 1836, not a salmon was ever seen, though many were in the habit of entering the loch or lake. In the year mentioned, at the request of his Grace of Sutherland and Mr. Loch, M. P., salmon were caught in the river Shin, shortly before the spawning season, and conveyed to the four rivers above named, amongst which they were distributed in due proportions. Mr. Young was the managing director on the occasion. In the wonted season all the fish spawned, each in its respective river.

Now mark one of the consequences ;—salmon at present, and ever since, come regularly to spawn, traversing the lake to do so, in all these heretofore salmonless rivers. Nay, more, the fish hatched in the Terry—at least those that survive long enough—return to the Terry, and the young of the other three rivers return from the sea to them, each grilse or salmon entering never-faillingly the stream that gave it birth. What wonderful and unerring instinct !”

With these well established facts and figures before us, we cannot longer question the expediency of making the attempt to restock our streams with sea-fish.

The salmon (*salmo salar*) has always been a favorite fish, and brings the highest price of any fish in our markets. Its habits are tolerably well known, and are similar to those of

* Buckland's Fisheries of Great Britain.

the common brook trout (*salmo fontinalis*). Like the latter it always seeks clear, cold and shallow water, in which it deposits its spawn.

A female salmon that weighs ten pounds is said to deposit each year ten thousand ova. Larger and smaller ones produce in the same proportion, that is, for every pound that the fish weighs she is estimated to produce one thousand eggs. The common trout is even more prolific than this. Salmon and trout usually deposit their spawn in the months of October, November and December. At the time of spawning the female trout in their natural state are always attended by a male of the same species. A clean gravelly spot is selected, and lest some light substance of a muddy nature should be mixed with the gravel, the fish are guided by their instincts to *sweep* it clean with their tails before the spawn is deposited. This precaution is necessary, for spawn will not retain life in muddy water. This is doubtless one reason why so few trout are hatched in streams subject to freshets. After the spawning bed is suitably prepared the female deposits a portion of her ova, and is soon followed by the male fish that quietly settles upon the bed just vacated by his mate, and emits a spermatic fluid that descends upon and impregnates the eggs. By the combined motions of the fish and the effect of the running water the eggs are soon covered by the gravel or small pebbles upon which they are deposited.

The trout selects a shady spot in the stream, and usually deposits her spawn in night time. It is said—whether in truth or not—that spawn long exposed to the direct rays of the sun will not hatch. From the fact that the trout in her native stream selects a shady place for her spawning bed, we incline to the belief that the assertion may be true, and therefore would suggest that the safer way would be to deposit spawn of all fish in places not exposed directly to the sunlight. The young fry should also have a shady place in which to pass the first month of their existence.

Mr. Andrew Young, a Scotchman, who devoted much time to the study of salmon, gives the following description of the spawning bed, and the manner in which the ova are deposited and impregnated :*

"The spawning bed," says he, "which may be called a continuation of nests, is never fashioned transversely or across the water current, but straight against it. The way the bed is formed has never before been accurately described. Some have affirmed that the male fish is the sole architect; others that the female does all the work; others again, that the tail is the only delving implement employed, and others write that the bed trenches are dug across the stream. A salmon spawning bed is constructed thus: The fish having paired, chosen their spot for bed-making and being ready to lie in, they drop down stream a little and then rushing back with velocity toward the spot selected, dart their heads into the gravel, burrowing with their snouts into it. This burrowing action, assisted by the power of the fins, is performed with great force, and the water's current aiding; the upper part or roof of the excavation is removed. The burrowing process is continued until a first nest is dug sufficiently capacious for a first deposition of ova. Then the female enters this first hollow link of the bed, and deposits therein a portion of her ova. That done she retires down stream and the male instantly takes her place, and pouring, by emission, a certain quantity of milt over the deposited ova, impregnates them. After this the fish commence a second excavation immediately above the first, and in a straight line with it. In making the excavations they relieve one another. When one fish grows tired of its work, it drops down the stream until it is refreshed, and then with renovated powers resumes its labors, relieving at the same time its partner. The partner acts in the same spirit, and so their labor progresses by alternate exertion. The second bed completed,

* See G. P. Marsh's report on the Artificial Propagation of Fish, 1857.

the female enters it as she did the first, again depositing a portion of ova, and drops a little down stream. The male forthwith enters the excavation and impregnates the ova in it. The different nests are not made on the same day, but on different days progressively. The ova in the first are covered with gravel and sand, dug from the second, being carried into it chiefly by the action of the current. The excavating process just described is day by day continued until the female has no more ova to deposit. The last deposition of ova is covered in by the action of the fish and water, breaking down some of the gravel crust above and over the nest. Thus is formed a complete spawning bed, not at once, not by a single effort, but piecemeal, at several intervals of greater or less duration, according to the size of the fish and quantity of ova to be deposited."

When the female salmon or trout is engaged in depositing her ova the male is very attentive, and fiercely attacks any fish that attempts to eat the spawn or approach his mate. It is a curious fact and a wise provision in nature that during the spawning period fish will rarely ever eat anything. Were it otherwise they would devour their own spawn, for every fisherman knows how greedily trout and other fish will eat spawn, even of their own species, when given to them away from their spawning beds. It is said* that neither shad nor salmon ever have any food in their stomach after reaching fresh water, and the presumption is that they gain most of their sustenance in the salt water after they make their first trip to it. It is well known that they invariably are fat when they enter fresh water and lean when they leave it.

The period of incubation for trout and salmon varies according to the temperature of the water in which the ova are deposited. From the testimony of an English writer † salmon spawn have been known to hatch in thirty days, while

* See Norris' American Angler.

† Frank Buckland.

others in water of a lower temperature have remained one hundred and forty days before they hatched.

From the careful experiments made by Mr. John Shaw in Great Britain, about thirty years ago, it is found that the eggs of salmon will hatch in one hundred and fourteen days when the water has a temperature of thirty-six degrees; one hundred and one days with a temperature of forty-three, and ninety days with a temperature of forty-five degrees.

The spawn when first deposited are quite translucent. If properly impregnated, a milky speck will appear in the egg, which will gradually expand into a whitish line—the germ of the spine. At one end of this two black specks appear that mark the spots for the eyes. These continue to grow till the embryo fish breaks from the shell, when the eyes appear disproportionately large. As the young fry emerges from the shell it is nearly transparent, its fins but imperfectly formed, and it is able to swim but little in consequence of the umbilical bladder that is attached to it just back of the gills. This bladder is nearly as large as the egg which contained it, and from it the young fry derives its entire support during the first month of its existence. It is when the fish are in the embryo state—just after the black spots that mark the eyes appear—that they can safely be transported and introduced into streams far distant from the one in which their progenitors lived. All that is required to insure their safe removal seems to be an uniform temperature, and a sufficient amount of aerated water to keep them wet. If they are placed in water it should be often changed, or air should be driven through it with a bellows or syringe. Moss from swamps is like a sponge and not only retains the water but also seems to impart oxygen to it. Spawn carefully packed in wet sphagnum moss can be safely transported to a considerable distance.

Soon after the umbilical bladder is absorbed the young fry seek food, and are very sprightly in their motions. They are

then called *parrs*. We will now briefly quote the words of another,* showing the growth, and giving the *names* that the salmon bear in the different stage of their existence. After the umbilical sac is absorbed, he says: "It is then a salmon *fry*, and so continues to its twelfth month, when it becomes a *smolt* with silvery coat, its length five inches or so, and its weight two or three ounces. It then migrates to salt water, feeds therein, and in the course of three or four months becomes a *grilse*, or maiden salmon, weighing not unfrequently eight pounds (rapidity of growth most marvelous and almost incredible!) In this *grilse* state it returns to its native river, spawns in due season upon its fords, and not long afterward migrates for the second time to the sea. Having fed there for two or three months it becomes an adult salmon weighing twelve, fourteen or sixteen pounds. It again returns to its native river, deposits in the spawning beds some ten thousand to fifteen thousand ova, which are impregnated by the male, and give life in all probability to seven or eight thousand salmon. Salmon will migrate and immigrate every year until they are captured, increasing in size, not in the same ratio as in their first years, annually, and breeding annually." Some may ask how these facts are known. To such we will say that repeated experiments have been made by cutting the fins of the young fish or attaching marked labels to them which were easily recognized upon the return of the salmon.

Mr. Andrew Young, of Sutherlandshire, England, made experiments which he reports as follows:†

"I have experimented on salmon" says he, "for upwards of thirty years. * * * * In 1834 and for a number of years following we marked spawned fish for the purpose of settling the question denied by many, of the return of salmon to their native rivers. This we did satisfactorily.

* See Fry's Treatise on Fish Breeding, pp. 159-60.

† See Fry's Treatise on Fish Breeding, pp. 117-18.

We, in 1835, marked smolts to ascertain and set at rest the following point, denied by many, viz : that the *smolts* returned *grilse* the same year that they first went from the river to the sea in the smolt state. The experiment proved this also, and specimens of the grilse that we marked when smolts, and which returned grilse from the sea to fresh water the year they were marked, may be now seen in the Museum of the Royal Society of Edinburgh. We continued these markings many years, invariably with similar results, and at the same period, continuing continuously during three years, we carefully watched the spawning operations and spawning beds in all their stages, and were fully convinced at last that the fry remained in the rivers *one whole year and no longer after having been hatched.*"

From the foregoing it appears that the parr or young salmon fry remains in its native stream one full year, and weighs only a few ounces ; that it then becomes a smolt, clothed with silvery scales, and runs to the sea, where it remains a few months and returns a grilse from fifteen to eighteen months old, filled with spawn and weighing from four to eight pounds each ! After the spawning season is past it returns to salt water lean and worthless. A few months elapse and it again returns to fresh water weighing from ten to sixteen pounds, and then they are called salmon.

The natural history of the shad (*alosa præstabilis*) is in some respects similar to the salmon. They ascend fresh water streams in May and June and soon deposit their spawn that hatch in a few weeks. The young shad the next fall run in immense numbers to the salt water where they spend the winter. In the spring they return fat to their native waters weighing from two to six pounds each ! They select their spawning ground in bodies of water deeper and warmer than those occupied by the salmon. The deep eddies below dams and waterfalls are generally selected by them. The eddy below Bellows Falls was formerly a favorite spawning ground for the shad. The one below Holyoke dam in Mas-

sachusetts is now occupied for that purpose, and thousands of shad are annually caught at that place.

Before the erection of the high dam across the Merrimac at Lowell, shad and salmon ascended that stream in great numbers as far as Franklin, N. H. At this point the company parted. The shad invariably took the stream for Winnipisiogee lake, and the salmon took the other branch that flows from the cold springs near the base of the White Mountains.

Prior to the building of the dams in the Connecticut river to furnish slack-water navigation, in 1798, shad and salmon were abundant in that stream, and furnished much of the food for the people of the country at that time. The "Pleasant Bard," * in his poem on "Shad Fisheries," says that those who read his truthful rhymes,

"Can scarce conceive what schools of shad
Made our old fisher fathers glad."

and further adds :

"A single haul would bring ashore
Some forty, fifty, sixty score.

* * *

When for some two and seventy pence,
You might have drawn a cart load thence
Of just the finest shad that ever
Swam this or any other river."

Shad were formerly abundant as far up the Connecticut as Bellows Falls, but never above that barrier.

But salmon, which are a very strong and active fish, ascended the falls and were caught abundantly as far up as the head waters of the Connecticut, and occasionally in its tributaries. But the building of a dam sixteen feet high across the Connecticut just below the mouth of Miller's river in Massachusetts in 1798, effectually stopped the passage of both shad and salmon. In a few years, as might have been expected, salmon entirely disappeared from the stream. But not so with the

* Joshua D. Canning, of Gill, Ms.

shad. Instead of requiring the clear, cold water at the head of streams, they were so well suited with the eddies of the Connecticut as to remain there and fulfil their work of procreation in the deep waters of that stream.

Notwithstanding the reckless and unrestricted slaughter that is annually made upon the shad of the Connecticut by gill nets and other devices, they are still quite abundant as far up as the dam at Holyoke, beyond which they cannot ascend. But were the obstructions in the Connecticut removed by the building of suitable fishways over them, shad could again be caught at Bellows Falls as in the days of yore ; provided spawn or young shad were placed there. Perhaps the best way would be to capture some shad just before the spawning season and place them in the eddy below the falls. Some are of opinion that merely the construction of suitable fishways would be sufficient to ensure their return ; but the safer way would be to introduce the live shad or spawn as proposed above. We are of opinion that shad may be produced successfully at points in the Connecticut above Bellows Falls, and even in the streams and ponds of eastern Vermont that are suited to their natures, provided suitable fishways be constructed over the falls and dams between their spawning beds and the ocean.

With these conditions complied with, we are of opinion that shad may be produced abundantly in the numerous ponds through which Black River flows in Plymouth, Ludlow, Cavendish and Springfield, and the same may be true of other streams in eastern Vermont. As a general rule we think shad would be best suited with deep, still water, while salmon would run to the clear, cold water of our mountain streams. But before we can have an abundance of shad and salmon in our streams, money must be expended in the construction of fishways, and in procuring spawn to raise the first stock of fish. The Connecticut river is wholly out of Vermont, the boundary line of New Hampshire being the west bank of that stream ; therefore the Legislature is only

called upon to provide for the construction of fishways over the tributaries of the Connecticut, and make suitable provision for the introduction of spawn or young fry into the streams of the State.

Fishways are not very expensive except over high dams upon large streams. They are made of plank and usually start from one end of the dam. A small gap, say four to twelve feet wide and from one to two feet deep, according to the size of the stream, is made in the dam, so that the waste water that otherwise would flow over the dam shall pass through this gap and enter the fishway, which is a trough or *chute* descending about one foot in ten. In this trough are bars that extend about half way across and are slightly inclined up stream to afford *resting places* for the fish as they go up the fishway. To make it more plain we introduce the following diagram, which represents a fishway as seen from below the dam :

It is important that the foot of the fishway be in a deep basin. If the only basin below the dam is too near it to be reached by a fishway as shown in the diagram, one may be made to be run down part of the way and then be turned so as to reach the basin, similar to the turn often made with a flight of stairs.

The question may arise whether the fish now in our streams may not devour the young shad and salmon so as to exterminate them or prevent their rapid increase. We have fears that the bull pout (*pimelodus vulgaris*) may destroy many of the shad spawn. But the salmon spawn, deposited in the clear, cold water, will be likely to escape them, as the pout seldom frequents such spots.

The pickerel (*esox reticulatus*) is a very voracious fish, and will doubtless destroy some of the young shad and salmon. But they rarely ever run up swift running rivulets where the young salmon are bred, therefore the latter would be comparatively safe. The spawning season for the pickerel is in the spring, just about the time that the young salmon have absorbed the substance of their umbilical sac and want food. Being comparatively safe from the attacks of the pickerel, (for they like other fish will eat but little or nothing during the spawning period,) the young salmon fry will eat the pickerel spawn or the young pickerel fry as they are hatched.

The black sucker (*catastomas nigricans*) will probably eat thousands of shad spawn, but as a compensation for this, the young sucker fry will furnish excellent food for the shad and salmon. The salmon spawn will be comparatively safe from the attacks of suckers. It is a true saying that "big fish will eat little ones." We must expect this. We must not count upon more than one full grown fish for every ten spawn. But even one fish for every fifty will, in our opinion, yield a greater percentage of food for the capital invested than from any other source in the State.

By the conditions of the joint resolution under which we received our appointment no allusion was made to the introduction of fish into Lake Champlain and the rivers upon the west side of the Green Mountains. But we feel unwilling to dismiss this subject without alluding to the desirableness of restocking Lake Champlain and the streams entering it with sea-fish. At the last session of the Vermont Legislature a law was passed prohibiting the catching of shad or white-fish from Lake Champlain during the months of September, October and November, to take effect after the state of New York and the Canadian Government had passed a similar act. The Provincial legislature of Canada have passed a similar law, but the legislature of New York have failed to do so, and therefore the law is a nullity.

The shad or white-fish (*coregonus albus*) of Lake Champlain is unlike the shad of the Connecticut, and does not increase in size or numbers as rapidly as the latter. Were Canada and New York to enact suitable laws for the introduction and preservation of sea-fish in Lake Champlain and the St. Lawrence river, we are of opinion that Lake Champlain would become one of the best locations for shad fisheries in the United States, provided the spawn of the shad (*alosa præstabilis*) were introduced therein and suitably protected.

In relation to the introduction of "black bass, land-locked salmon and other varieties," into the waters of the State, we will say that we have not done so for the reason that it would be useless to make the attempt unless there were laws to suitably protect them.

Our streams are well suited to the habits of the common brook trout, and before we would recommend the introduction of other varieties we would ask that suitable protection be extended to them. If properly protected by legislative enactments, and a little effort is made to preserve and increase the number of young fry, our brooks and rivers would soon

be filled with trout. The spawn from half a dozen pair of trout per annum would furnish young fry enough to make good fishing in any brook in the State. Several people in the State are making experiments in artificial trout raising with a good degree of success. But they have generally commenced with little or no instructions or experience upon the subject, and have often met with reverses that a better knowledge of the business would have prevented.

ARTIFICIAL TROUT RAISING.

To aid such as may desire information upon this subject, we will here introduce very clear and valuable instructions from the pen of Hon. S. H. Ainsworth of West Bloomfield, New York, who has been very successful in artificial trout raising.

BEST METHOD OF CONSTRUCTING PONDS.

"To grow trout successfully, there must be three ponds, one above the other, in the same stream if possible, as it saves much time in transferring the small fry from one pond to the other, as they increase in size. The object of these ponds is to keep the young trout out of the reach of the larger ones until they are of sufficient size to take care of themselves, which requires two years at least. Pound trout would soon exterminate a pondful of one year olds, if not well fed, and undoubtedly take more or less of them if well fed, so that they would soon run out if left together. The first or upper pond is to grow the small fry in the first year; the second one for the second year's growth. In the spring of the second year they will geneally be large enough to care for themselves—say from seven to nine inches long, and from four to five ounces in weight, if well fed. They may now be turned into the lower or large pond. The first pond may contain from one to eighty square rods, and be from one to three feet deep, according to amount of water, and number of trout one wishes to grow. The second pond from 300 to 400 square rods, and from two to eight feet deep.

The third, or last pond from forty square rods to ten or more acres, and from eight to twenty feet deep. The water in the last pond must be deep if large trout would be grown. From 8,000 to 10,000 trout may be grown in the smallest pond, and from 300,000 upward to 2,000,000 in the largest pond.

The dams for these ponds should be built of stone or earth, to be lasting. Earth is much the cheapest. Clay will hold all the water and make a permanent dam. If built of stone, the cheaper way is to build a double wall, laid in quick lime, with two inch space between, and fill the space with water-lime grouting, which will hold the water through all time. This wall should be in thickness in proportion to its height, say, from four to ten feet. These dams can be built partly of stone and partly of earth, as may be desired. At the bottom of each of these dams there should be a flume from six inches to four feet square, according to volume of water, to draw off the water and fish when desired, from one pond to the other; also, to catch and assort the fish for use in the lower pond, and to clear the ponds of grass and filth when necessary—perhaps in from four to eight years. There must also be a flume for waste water on top, with a wire screen to prevent the trout from running out, or they would soon leave the pond. To hold the young fry, the meshes in the screen must be little larger than the head of a pin, as these young trout are very tiny when the egg is absorbed, not more than one to one and one-fourth inch long. The first dam must be tight to keep them. The dams will cost from \$50 to \$4000, depending upon size and quality.

SPAWNING BEDS.

There must be a race leading into each of the two lower ponds from two to ten feet wide, and two to six rods long, with descent enough to produce a gentle, rippling current. The bottom must be covered two or three inches deep with well washed gravel, with a nice screen at each end, to be raised or closed at pleasure. These races should be so con-

structed that the top of the gravel at the lower end will be on a level with the water in the pond at its lowest stage ; and the dams are constructed in such a manner that the water may be raised at will from twelve to eighteen inches in the pond and lower end of race. The races are the spawning ground for the trout, in each of the ponds, and every trout will run into them during the spawning season, to deposit their ova—at which time the eggs may be procured by artificial means, by dropping the screens and taking the fish in a scoop-net. During this season the race should be covered with boards, and all intruders kept away, as they are very shy at this time, although quite tame in the ponds. In like manner races may be built at the lower ends of the ponds by taking the water out along the side of the bank. The trout will find their way down as well as up.

WATER.

The most important thing to be looked to in trout-growing is the *amount*, *purity* and *temperature* of the water. The volume of water should be ample, entirely free from all sediment and all surface water, and at least forty degrees Fahrenheit—still better at forty-eight degrees—the temperature of the water in the far-famed Caledonia Springs, where the trout naturally increase and grow faster than in any other place known in the world. This stream, only one mile in length, is fished constantly from the first of April to the first of September, furnishing thousands of pounds, annually, without any apparent diminution, which can only be accounted for by taking into consideration the exceeding purity as well as the high temperature of the water,—the water in the coldest day of winter being ^{was} forty-eight degrees in the spring, and forty-seven degrees in the spawning beds, and forty-two degrees generally, in the creek. The small ponds can be supplied by a spring that will fill a two-inch pipe, and so on in proportion to volume, up to the largest named ponds, which should, at least, fill a forty inch pipe. This will keep

the water pure and clear in all the ponds, and the fish in a healthy and vigorous condition.

HATCHING BOXES.

The only sure method of success in hatching the eggs, is to place them in troughs from eight to eighteen inches wide, according to the volume of water, six inches deep, and from ten to fifty feet long, with a slight descent—enough to produce a gentle current, but not enough to move the spawn when placed in them. The bottom of these troughs are to be covered with two inches of fine gravel, and clean, coarse sand. These troughs are completed with a lid to cover tightly, and with screens at each end. They are now ready for the water. The water should be from one and a half to two inches deep over the gravel and eggs. The number of these boxes may be increased with the amount of water and number of eggs. The nearer the boxes are to the springs the better. The water from the spring should run directly into a box placed at right angles with the hatching boxes, to be used, say from three to fifteen feet long, twenty inches deep and wide, and drawn from this box through fine screens and spouts into the hatching boxes, graduating the amount of water required in each hatching box, to give the required depth to the water over the eggs. The boxes are now ready for the eggs.

SPAWNING SEASON.

In my pond, the trout commenced spawning on the 12th of November in 1860, on the 5th in 1861, and on the 3d in 1862, and finished each year from the 10th to the 15th of January. I am informed that in some other sections they commence the first half of October. The female trout seldom come into the race until the very day they begin to spawn. As soon as they commence to deposit the ova they should be taken out and the spawn procured artificially.

HOW TO EXTRUDE THE EGGS.

Take out the trout in the race with a net, and place them in baskets, standing in the water, in some convenient place to handle them. Take a pan or pail with three or four inches of water in it from the spring, and place it near the baskets containing the trout. The eggs must be quickly extruded and the trout replaced in the water. This operation must not consume more than one minute if possible. All things being ready a female trout is taken out of the basket with one hand, with the other gently rub the abdomen from the gills downward, and the spawn will flow in a continuous stream into the vessel. Continue the rubbing until the spawn is wholly extruded, then quickly replace the trout in the race, or separate basket. One side of the egg has a small, white speck; here is where the impregnation takes place. This side of the egg being the lightest it always falls this side up ready to receive the milt. We have now in the pail from 400 to 8,000 spawn, according to the age and size of the female. Now take a male trout from the basket, and, in like manner, or by the thumb and finger on each side of the abdomen, which requires rather more pressure, exude the milt. The milt falls upon the water and settles upon the eggs. I usually take from two to four males to impregnate from 2,000 to 8,000 spawn. In like manner I serve all the trout in the baskets. I then place the spawn and milt in shallow vessels and put the dishes in the spring water, where I allow them in the milt from one to twenty-four hours. Probably one hour is sufficient to insure impregnation. I took from 300 to 10,000 spawn daily from the 3d of November to the 10th of January, making in all, about 130,000 spawn, attended with perfect success.

HOW TO PLACE THE EGGS IN THE HATCHING BOXES.

Commence at the lower end of the box, and lay the eggs on top of the sand as thick as you can place them without

touching each other. Proceed in this manner daily until you have filled the entire length. Should you choose, you can put partitions between each day's deposit, with date and number of eggs. A box one foot wide and twenty feet long will contain 100,000 eggs. It has been recommended to place the spawn in gravel from one to two inches deep, in imitation of the trout; but I find by practice that the first method is the better one, as it will enable you to examine the spawn daily. The reason will be seen under the head of

AFTER CARE.

The eggs will need to be examined every week or two, and all the dead or white ones picked out with a pair of forceps, made of No. 8 wire flattened at the ends. If the water is perfectly pure, and above forty-two degrees through the winter, but few will die. As the eggs die, a vegetable fungus, called *Byssus*, attaches itself to them, and throws out its little hairy fingers and clasps all the live eggs in their reach and soon kills them. Hence the necessity of having the eggs in sight. These hatching troughs should be covered with a house, containing a stove with fire, as it will make it more comfortable for the operator, and aid in keeping up the temperature of the water.

THE TIME OF INCUBATION

Depends upon the condition and temperature of the water. The water in my hatching boxes stands this winter at thirty-eight degrees, and at forty degrees in the springs. The springs are from eight to thirty rods from the box, brought in two-inch pipe tile, laid from two and a half to three feet under ground in water-line. In this water the eggs commenced hatching the 21st of January, seventy-eight days after they were put in the box, and they have been hatching in great numbers, daily, since. So far, my success has been beyond my most sanguine expectations, and should nothing befall them I shall have enough to stock a number of small

ponds, at least. I took a few eggs from the race after they had been in forty days, and put them in a tumbler in my house where the temperature ranged from fifty to sixty degrees. They hatched in twenty-six days after, twelve days sooner than those in the race. The egg has two skins or membranes; the trout is formed between them. The first appearance of the egg, as viewed through a magnifying glass, is a red speck on one side near the white speck before referred to. This, I think, is the heart. This is about the fortieth day. In two or three days a fine artery is seen running each way from the speck around the egg. About the forty-ninth day the eyes are perceptible; on the fifty-first day could see the formation of the head and body; on the sixtieth, could see the heart beat; and the arteries running in all directions. It now shows life, and in a few days bursts the outer covering, but is still firmly attached to the egg, which, in fact, is a part of the young trout. The young trout, when first hatched, is about half an inch long, and looks and acts more like a wiggler you often see in rain-water, than a trout. It has no fins except the pectorals, just back of the gills. He lies upon his side almost constantly. On the fifteen day all the rest of the fins are plainly seen, and he now takes on the appearance and action of a trout, and is constantly in motion, darting through the water with great rapidity. It lives upon this sack until it is all absorbed, which is about the fortieth day. At this time the young fish begins to seek its own living. He is now one and a quarter inches long.

FOOD.

I found, last spring, that the young trout ate and thrived well on the yolk of hens' eggs, boiled hard and crumbled fine. After a few weeks I fed them on hashed liver. I find this to be the best food for trout of all ages, as they grew rapidly upon it. I feed my trout once a day through the summer, and from two to three times a week through the winter, as

they eat less in cold weather than in warm, and grow in proportion. I have 1100 parent trout caught with a hook in the streams about. My largest pond contains sixty-one square rods ; water fourteen feet deep. Four hundred of these trout were put in last summer, four hundred one year ago, and about four hundred two years ago. About fifty have been taken out for table use, and about the same number have died from various causes. They were from one to two years old when placed in the pond, so that they now range from two to five years old. A few are older. I fed them the livers, &c., from two beeves a week during the last summer. This furnished food so abundantly that they paid but little attention to flies or other insects. It cost me about seven dollars for food last year. The fish are very tame, coming regularly for their food ; they take it from a spoon, or hand, and throw themselves clear out of the water, by the hundred, in their eagerness for it.

GROWTH.

The trout that hatched one year ago now, are from four to seven inches long, and are heavy compared with those in the streams in a wild state. Those two years old are from seven to nine inches ; three years, from nine to eleven inches ; four years, from eleven to thirteen inches ; and five years, from thirteen to fifteen inches long. Trout four years old will average one pound in weight. The trout in my pond will weigh from one quarter of a pound to two pounds each. With pure, cold water, and plenty of good, fresh food daily, trout will grow rapidly, and may attain a pound's weight in three years. Still, those of the same age vary much in size. There are Tom Thumbs and Lavinia Warrens, as well as Daniel Lamberts and Hannah Crouses, in this trout family, as well as in the human family.

TRANSPORTATION.

One year old trout may be carried a long distance in wa-

ter with ice, if kept constantly in motion, so as to keep up the supply of oxygen. I have brought 300 trout, one and two years old, twenty miles in a barrel, without changing water, but keeping it cold with ice. Had the water been still, they would have lived but a short time. The eggs, after impregnation, may be carried any distance in water, in bottles or pails, or packed in clean, wet sand. In this way all the streams and ponds in the country might be stocked in a few years. The spawn from a few thousand parent trout, thus sent broadcast over the land, would soon fill the streams, and furnish an immense amount of sport to the angler, and most delicious food to the country.

PROFITS.

The first question asked by many is, Can trout-growing be made profitable? My answer is, Yes. At the present price of trout in market, which is fifty cents per pound, and ready sale for all that can be bought at that, I know of no other business that can be made more remunerative. Let us make the calculations in the scale of the largest ponds here discussed, where there is an abundance of pure spring water, viz :

Cost of buildings, dams and fixtures.....	\$6,000
Cost of 5,000 parents for spawn, at 50c.....	2,500
Three men's labor for four years, at \$300 per year.....	3,600
Cost of feed for 1,000,000 trout for 4 years.....	20,000
“ “ “ 3 “	10,000
“ “ “ 2 “	4,000
“ “ “ 1 “	1,000
Total.....	<u>\$47,100</u>

Now for their value. The million of four year olds will average a pound each, and are worth at least 25 cents per pound in the pond, which makes the four years olds

Worth.....	\$250,000
1,000,000 3 years, $\frac{1}{2}$ lb. each.....	175,000

1,000,000 2 years, $\frac{1}{2}$ lb. each.....	87,500
“ 1 “ 3 oz “	30,000
<hr/>	
The worth of all the trout at the end of 4 years.....	\$542,500
Deduct the price of growing.....	47,100
<hr/>	
Total profit.....	\$495,400

But if the profit is too large to please, we will say that the trout are worth but one shilling a pound; and where is there a disciple of Sir Isaak in America, who would not be glad of the chance for such high sport, through the means of his fly, and pay his shilling per pound for all he caught? Yes, sir, the anglers of the country would soon take out all the trout in the ponds at that lay. A pound trout at the end of a 100-foot line and a ten foot fly-rod, rushing through the water with an almost lightning speed, first this way, then that, bending the rod nearly double with his power, and spinning the reel as he runs out the line to its full length, showing his broad, beautiful sides—now here, now there, now diving deep down in his crystal element, leaping out of the water in this and that direction—requiring a steady hand and great skill to bring him to land; even now, about to pick him up, he makes another powerful struggle for life, and is away again, with the velocity of thought, but this time soon yielding to a superior power—the well-practiced hand of his captor—and now lies motionless on the water, with his mouth extended, his glistening sides turning about in the water as he is exultingly reeled to land, soon to be placed in the delighted angler's basket. One shilling per pound for such a trout! Every trout fisherman will tell you that his trout cost him nearer one dollar per pound, let him go where he will for them, besides hardships innumerable when going into a dense wilderness. At one shilling per pound, they would give the nice little sum of \$224,150. But suppose, again, there could be but one-half of the number raised in the largest sized ponds, and the expense of growing them would be equal to the first calculation; still this would give a profit

of \$88,525. Suppose, finally, that they are worth only six cents per pound; this will give a profit of \$29,712.50, besides all the ponds, parent trout, and 300,000,000 eggs, or young fry, in hatching-boxes, could have been taken from the number of trout now in the ponds,—worth a fortune of themselves! I believe the first sum might be realized, with everything favorable. But they should be located near a large city, in order to obtain sufficient food. Small ponds will yield a profit in proportion. Nearly every farmer has a spring, if properly arranged, that will furnish trout for his table, with but little expense.”

In a letter received from Mr. Ainsworth he says: “I have hatched as high as ninety-nine out of every hundred put into my spawning boxes, and in one instance two hundred and nine out of two hundred and ten spawn put into a small wire box to test the number that I could hatch on a small scale.” Mr. Ainsworth estimates the number of spawn in the female trout to average as follows:

A one year old has.....	300
two “ “ “	700
three “ “ “	1200
four “ “ “	1800
five “ “ “	2800
six “ “ “	4000

He says: “I have taken 8000 spawn from a six year old trout weighing two pounds, and hatched them all as far as I could discern, mixed with others.”

The most precarious time in the life of a trout is just after the umbilical sac is absorbed. They then require food and pure aerated water, or they will starve for want of water to breathe or food to eat. But if taken from the hatching boxes and placed in running streams of water suited to their nature, they will nearly all live and furnish themselves with food. In this way the brooks of Vermont may nearly all be made good “trout brooks.”

If persons owning artificial ponds would remove the trout therefrom in early winter, the young fry would be found in great numbers the following spring. Mr. Seth Green of Munford, Monroe County, New York, is extensively engaged in raising trout. He not only sells the mature trout, but also the small fry and spawn for stocking streams or ponds. The latter he sells at ten dollars per thousand. The young fry, an inch long, he sells at fifty dollars per thousand. He sells the yearlings at one hundred dollars per thousand. It is much less trouble, and quite as safe, to transport the spawn instead of either the young fry or the yearlings. He says in his circular, that contains full directions for handling the spawn: "I can send them a fifty day's journey, packed in a box with moss. I place the moss box in a tin pail, with sawdust so that the spawn will not feel the change of heat or cold."

From what has been said in the foregoing pages, it is evident that, with proper management, trout raising may be made as profitable business, with the capital required, as any other industrial pursuit in which a person can engage. A few in Vermont have already commenced upon a small scale, and more will probably do so when the matter is properly presented to them. To aid such in their beginning, has been our object in presenting so many facts, and giving the *modus operandi* so much in detail.

In order to have the enterprise of fish culture a successful one, whether in private ponds or in public streams, legislative aid will be required. Stringent laws should be enacted and enforced, embracing the following conditions:

First, An absolute prohibition to take salmon, shad or trout in any of the rivers or ponds of the State for the space of five years from the time such fish are placed in the streams, under the penalty of fifty dollars for every fish caught, and thirty days imprisonment in the county jail, the fine to go to the informer.

Second, The prohibition, under the same penalties *forever* of taking any of the fish named during the months from October to March inclusive; or within forty rods of any fishway over which the fish may pass at any season of the year.

Third, The prohibition under the same penalties, of using any gill-nets, stake-wears or other devices for taking fish that shall extend more than one third the distance across the stream.

Fourth, That fishways be constructed and maintained over all dams more than three feet high in streams suitable for shad, salmon or trout. With these conditions complied with, and the concurrent legislation and action on the part of New Hampshire, Massachusetts and Connecticut, we are of opinion that success will attend the enterprise, and a new source will be opened from whence these luxuries can be obtained in such abundance that they will form an important articles of subsistence for the people of the State.

ALBERT D. HAGER, }
CHARLES BARRETT, } Commissioners.

Montpelier, Vt., Oct. 11, 1866.