ms. 5X
LS / ϕ - web

A
MONOGRAPH
OF THE
BUCEROTIDÆ,
OR
FAMILY OF THE HORN BILLS.

BY
DANIEL GIRAUD ELLIOT,

COMMANDER OF THE ROYAL ORDERS OF THE CROWN OF ITALY, OF FREDERIC OF WURTEMBERG, AND OF CHARLES THE THIRD OF SPAIN;
KNIGHT OF THE IMPERIAL AND ROYAL ORDERS OF FRANCIS JOSEPH OF AUSTRIA, OF THE DANNEBROG OF DENMARK,
OF THE ALBERT ORDER OF SAXONY, OF ST. MAURICE AND ST. LAZARE OF ITALY, OF ISABELLA THE CATHOLIC OF SPAIN,
OF CHRIST OF PORTUGAL, OF PHILIP THE MAGNANIME OF HESSE, ETC. ETC.
FELLOW OF THE ROYAL SOCIETY OF EDINBURGH, OF THE ZOOLOGICAL SOCIETY OF LONDON.
MEMBER OF THE IMPERIAL LEOPOLDINO-CAROLINA ACADEMY OF GERMANY, OF THE ACADEMY OF NATURAL SCIENCES OF PHILADELPHIA,
OF THE ACADEMY OF SCIENCES OF NEW YORK, OF THE BRITISH ORNITHOLOGISTS' UNION, OF THE SOCIÉTÉ D'ACCLIMATATION OF PARIS,
OF THE SOCIÉTÉ ZOOLOGIQUE OF FRANCE, OF THE NEW-YORK HISTORICAL SOCIETY;
HONORARY MEMBER OF THE NUTTALL ORNITHOLOGICAL CLUB; CORRESPONDING MEMBER OF THE ROYAL ACADEMY
OF SCIENCES OF LISBON, OF THE NATURAL-HISTORY SOCIETY OF BOSTON, ETC. ETC.

AUTHOR OF THE
'NEW AND HERETOFORE UNFIGURED BIRDS OF NORTH AMERICA,'
OF THE ILLUSTRATED MONOGRAPHS OF THE
PITTIDÆ, TETRAONINÆ, PARADISEIDÆ, PHASIANIDÆ, FELIDÆ, &c. &c.

PUBLISHED FOR THE SUBSCRIBERS BY THE AUTHOR.

1882.

ALERE FLAMMAM.

PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET.

f QW
696
C7E46
RB
SI

TO

Osbert Salvin, Esq., F.R.S., &c.,

THIS

MONOGRAPH OF THE BUCEROTIDÆ

IS DEDICATED,

AS A MARK OF ESTEEM FOR HIS ABILITIES AS AN ORNITHOLOGIST,

AND IN TESTIMONY OF

A FRIENDSHIP OF MANY YEARS.

PREFACE

COULD an interest in any ornithological group only be created by the beauty of dress or gracefulness of form of its various members, it might possibly happen that the BUCEROTIDÆ would not be selected as the subject for an illustrated monograph; and while their full value is always accorded to these attributes, perhaps occasionally even in an exaggerated degree, yet as Nature never made an ugly object (even the most repulsive thing so called being admirably and wonderfully fitted for the place it is destined to fill in life), beauty of plumage and symmetry of form are by no means the only causes that lead a naturalist to choose any one group as an especial object for study. The very peculiar appearance of the majority of the birds contained in this volume, as well as the extraordinary habits and structure common to all, which make them to differ from other feathered creatures, together with the generally meagre accounts of many of the species, only to be met with by searching numerous publications, were the chief reasons that induced me to select this family as the subject of my fifth illustrated monograph.

Scattered as the species are over many countries, it has not fallen to the lot of any one ornithologist to observe all of them in their native haunts; but beside what could be gathered from published accounts, to be found in various journals written in many languages, I have been most kindly aided by those who have enjoyed opportunities of observing certain species in the localities where the birds dwell. Although in the various articles accompanying the species in the body of this work I have endeavoured to express my thanks to those who have in any way helped me, I would nevertheless take this opportunity to repeat them, apologizing at the same time to any one of my friends whose name I may inadvertently omit. To Mr. A. O. Hume, who has at all times in the most generous manner given me the benefit of his extensive knowledge of the Indian species, and incurred the expense of a special expedition to procure for this work specimens of *ANORRHINUS TICKELLI*, of which no example

existed in any European museum, I am under particular obligations. The late Marquis of Tweeddale, in whose sudden death this science sustained a grievous loss, was always ready to aid me by the loan of specimens from his magnificent collection, otherwise unobtainable, and by any other assistance it was in his power to bestow. To my friends Osbert Salvin, Esq., Dr. Sclater, Prof. Bocage, Captain Ramsay, R. B. Sharpe, Esq., Captain Shelley, Prof. Cabanis, Prof. Schlegel, Mons. A. von Bemmelin, Prof. A. Milne-Edwards, and Mons. Oustelet I desire to acknowledge my indebtedness for assistance on many different occasions.

The drawings, the happy results of Mr. Keuleman's talented pencil, most characteristically depict the strange forms and attitudes of these curious birds.

The colouring of the Plates has been successfully accomplished by Mr. Smith; while the letterpress by Messrs. Taylor and Francis leaves nothing to be desired, and the printing of the Plates by Messrs. M. and N. Hanhart has been most carefully executed.

The length of time during which the publication of the various Parts has extended was caused by circumstances entirely beyond my control; but I would especially express my thanks to those who have honoured my work with their support, and patiently awaited its completion.

D. G. E.

INTRODUCTION.

THE Bucerotidæ, or Hornbills, on account of the large size of many of the species, together with their apparently unwieldy and curiously formed casques, are very conspicuous among the families which constitute the class Aves. Their habits are no less uncommon than is their aspect; and the unique one possessed by the males of enclosing the female in the hollow of some tree, firmly fastening her in by a wall of mud, and keeping her a close prisoner until the eggs are hatched, is not paralleled by the customs of any other species of birds known to naturalists. Another curious habit is that of casting up the entire epithelial lining of the gizzard in the form of a sac or pocket, filled with undigested particles of food or hardened objects that may have been swallowed. Owls, as is well known, eject little round pellets, composed of the fur and bones of the small mammals upon which they have fed; but these are not accompanied, as a rule, by any portion of the bird's internal structure. The Hornbills, however, apparently suffer no inconvenience from the deprivation of their gizzard's walls; and an individual has been observed to quietly investigate with the point of his bill the contents of one of these sacs, which he had just cast forth, and which he would have probably swallowed again had not the bag been removed. In the present Monograph I have endeavoured—first, to review the literature of the Family from the time of Linnæus to the present day; secondly, to recapitulate the various genera which from time to time have been proposed for the different species; thirdly, to discuss the classification and geographical distribution of the various distinct forms; and, fourthly, to give the synonymy and life-history of each one, so far as these are known to me. In the arrangement that I have thus proposed, the first subject that presents itself is the Bibliography of the Family, of which I commence the review with the twelfth edition of Linnæus's 'Systema Naturæ,' published in 1766.

1766. LINNÆUS, Systema Naturæ.

In this edition the great systematist has recorded the species of this family known to him, retaining them all in the genus *Buceros*. Four only are mentioned, viz. *B. BICORNIS*, *B. HYDROCORAX* (= *PLANICORNIS*, Merrem, *HYDROCORAX* having been established as a genus by Brisson), *B. RHINOCEROS*, and *B. NASUTUS*. All these are recognized at the present day. Species 4.

1781. J. R. FORSTER, Indische Zoologie.

In this publication, of the zoological portion of which Thomas Pennant is ostensibly the author (but which is probably Pennant's list of English names latinized by Forster), six species of Hornbills are given. They are *B. BICORNIS*, Linn., *HYDROCORAX*, Linn., *RHINOCEROS*, Linn., *VIGIL* and *PLICATUS*, named for the first time, and *rostratus*. This last is referred to Willughby's 'Ornithology;' but as no plate or page is specified, it is impossible to ascertain what bird may have been intended. Although some of the species had been previously described by Linnæus, the fact appears to have been unknown to Forster, as Edwards, Willughby, and Dampier are the only authors cited. Species 6.

1783. BODDAERT, Tableaux des Planches Enluminées de d'Aubenton.

Founding his species upon the plates of Buffon's work, to which he gives Latin names, the present author records four species not before noticed. They are *B. CORONATUS*, *B. PANINI*, *B. MANILLÆ*, and *B. ABYSSINICUS*; all valid. Species 10.

1786. SCOPOLI, Deliciæ Floræ et Faunæ Insubricæ.

Three species of Hornbills are given in this work, one of which is recorded for the first time, viz. *B. BIROSTRIS*, which is the *ginginianus* of authors. The remaining two are *B. panayensis*, (= *PANINI*, Bodd.) and *B. pica* (= *B. CORONATUS*, Bodd.). Species 11.

1788. GMELIN, edition of Linnæus's 'Systema Naturæ.'

Twelve species are enumerated by Gmelin, only one, however, for the first time. They are *bicornis*, *abyssinicus*, *africanus* (= *ABYSSINICUS*, Bodd.), *MALABARICUS* (first distinguished as distinct from *CORONATUS*, Bodd.), *hydrocorax* (= *PLANICORNIS*, Merrem), *RHINOCEROS*, *galeatus* (= *VIGIL*, Forst.), *panayensis* (= *PANINI*, Bodd.), *manillensis* (= *MANILLÆ*, Bodd.), *NASUTUS*, *albus* (an albino of some bird not belonging to this family), and *obscurus* (= *PLICATUS*, Forst.). Species 12.

1790. LATHAM, Index Ornithologicus.

Sixteen species of Hornbills are here given, only one named for the first time. Nine are valid, the remainder either impossible to determine with certainty, or synonymous with other previously described species. They are *B. RHINOCEROS*, *B. galeatus* (= *VIGIL*, Forst.), *BICORNIS*, *ABYSSINICUS*, *africanus* (= *ABYSSINICUS*, Bodd.), *MALABARICUS*, *hydrocorax* (= *PLANICORNIS*, Merrem), *panayensis* (= *PANINI*, Bodd.), *manillensis* (= *MANILLÆ*, Bodd.), *NASUTUS*, *PLICATUS*, *ginginianus* (= *BIROSTRIS*, Scop.), *GRISEUS*, named for the first time, and *albus*, *orientalis*, and *viridis* not of this family. Species 13.

1793. LICHTENSTEIN, Catalogus rerum naturalium rarissimarum auctionis lege distrahendarum Hamburgi.

TOCKUS MELANOLEUCUS is here described. Species 14.

1801. LE VAILLANT, Histoire Naturelle d'Oiseaux Nouveaux et Rares de l'Amérique et des Indes.

In this work ten different Hornbills are given, illustrated by twenty-three plates of heads and figures of the entire bird. The nomenclature is French; and so some species represented here for the first time were characterized afterwards by other writers under Latin names. The species given are:—BICORNIS, Linn.; PLANICORNIS, Merrem; CORONATUS, Bodd.; SYLVESTRIS, Vieill.; MALABARICUS, Gmel.; BIROSTRIS, Scopoli; PANINI, Bodd.; PLICATUS, Forst.; and GINGALENSIS, Shaw.

1806. LE VAILLANT, Histoire Naturelle des Oiseaux d'Afrique.

As in the preceding work, the author employs French names. Nine species are mentioned, illustrated by plates; but the opportunity was lost for giving any of them a specific rank in the science.

1808. WILKES?, Encyclopædia Londinensis.

In this publication, of which in the octavo edition five volumes are devoted to ornithology, numerous species figured by Le Vaillant in his different works have Latin names given to them, some for the first time; and did we but know the author, they would take precedence of all those subsequently bestowed upon the species. No author's name, however, having been given, unfortunately none of the appellations can be recognized. It is unnecessary, therefore, to make further reference to the list.

1811. SHAW, General Zoology.

A list of the species as known to the author is given, compiled chiefly from Latham and Le Vaillant, and Latin names affixed to those previously only recognized by French or English appellations. Twenty-five in all are noticed, only fourteen of which are valid. B. MALABARICUS, Gmelin, is twice renamed, as *B. monoceros* and *B. albirostris*; B. CORONATUS, Bodd., is called *B. violaceus*; B. BICORNIS, Linn., is styled *B. cavatus*; B. UNDULATUS is described for the first time; *B. javanicus* (=UNDULATUS); B. MELANOLEUCUS, Licht., is named *B. coronatus*; B. GINGALENSIS and B. FASCIATUS are given for the first time. Species 17.

1811. BECHSTEIN, Kurze Uebersicht aller bekannten Vögel.

A catalogue of seventeen species of *Buceros* is given, with descriptions. Four are not valid, viz. *B. africanus* (=ABYSSINICUS, Bodd.), *B. albus*, *B. orientalis*, and *B. viridis* (ex Latham) not belonging to this family. No new species are described.

1816. VIEILLOT, Nouveau Dictionnaire d'Histoire Naturelle.

In the fourth volume of this work is a list of the then known species, two of which are described for the first time, viz. the Javan form of B. RHINOCEROS, founded upon the *Calao à casque en croissant* of Le Vaillant, and called B. SYLVESTRIS, and B. LEUCOCEPHALUS. B. PLICATUS, Forst., is renamed twice, as *B. ruficollis* and *B. niger*; B. BICORNIS, Linn., is called

B. cristatus; *B. jubatus* and *B. orientalis* from New Holland are not Hornbills; and *B. ruber* and *B. viridis* are copied from Latham and do not belong to this family; *B. coronatus*, Bodd., is styled *B. violaceus* after Shaw; *B. MELANOLEUCUS*, Licht., is called *B. coronatus*, and *B. FASCIATUS*, Shaw, is renamed *B. melanoleucus*; *B. VIGIL*, Forst., is called *B. galeatus* after Latham. The valid species recorded by this author are:—*ABYSSINICUS*, Bodd.; *VIGIL*, Forst.; *PLICATUS*, Forst.; *UNDULATUS*, Shaw (as *Le calao javan*); *SYLVESTRIS* and *LEUCOCEPHALUS*, described for the first time; *MELANOLEUCUS*, Licht.; *BIROSTRIS*, Scop. (as *ginginianus*, Lath.); *GRISEUS*, Lath. (referred to New Holland, a wrong habitat); *PANINI*, Bodd. (as *manillensis*, Lath.); *hydrocorax*, Linn. (= *PLANICORNIS*, Merr.); *BICORNIS*, Linn.; *RHINOCEROS*, Linn.; *CORONATUS*, Bodd. (as *violaceus*, Vieill.), and *GINGALENSIS*, Shaw (as *gingala*, Vieill.). All Linnæus's names are referred to Latham, he being apparently the most ancient author consulted. Species 19.

1820-39. TEMMINCK, Planches Coloriées.

A monograph of this family as then known is here given, with numerous species described for the first time, the material being supplied from the rich collection of the Leyden Museum. The new species are ten in number, viz. *CASSIDIX*, *ATRATUS*, *ELATUS*, *CYLINDRICUS*, *GALERITUS*, *CONVEXUS*, *CORRUGATUS*, *EXARHATUS*, *ERYTHORHYNCHUS*, and *BUCCINATOR*. *B. SYLVESTRIS*, Vieill., is renamed *lunatus*; *B. GRISEUS*, Lath., is called *B. cinerascens*; *B. MALAYANUS*, Raffl., is renamed *B. antracicus*. The female of *CORRUGATUS* is described as *B. gracilis*. The other species recorded are:—*RHINOCEROS*; *sulcatus* (= *LEUCOCEPHALUS*, Vieill.); *MALABARICUS*, Gmel. (as *monoceros*, Shaw); *BICORNIS*; *ABYSSINICUS*; *hydrocorax* (= *PLANICORNIS*, Merrem); *VIGIL*, Forst. (as *galeatus*); *UNDULATUS*, Shaw (as *plicatus*, Lath.); *CORONATUS*, Bodd. (as *violaceus*, Shaw); *PANINI*, Bodd. (as *panayensis*); *GINGALENSIS*, Shaw (as *ginginianus*); *FASCIATUS*, Shaw; *MELANOLEUCUS*, Licht. (as *coronatus*, Shaw); *PLICATUS*, Forst. (as *ruficollis*); *COMATUS*, Raffl.; and *NASUTUS*, Linn. The articles are illustrated with coloured plates of the new species. Species 29.

1822. RAFFLES, in the 'Transactions' of the Linnean Society.

Two new species are described from Sumatra, *B. MALAYANUS* and *B. COMATUS*. Species 31.

1823. BONNATERRE and VIEILLOT, Tableau Encyclopédique et Méthodique des trois Règnes de la Nature.

A catalogue of the species is here given, apparently a repetition of that in the 'Nouveau Dictionnaire,' with the same errors continued.

1824. MERREM, in Ersch und Gruber's Encyclopädie.

BUCEROS HYDROCORAX, Linn., is named *PLANICORNIS*, which will stand, as *HYDROCORAX* is Brisson's genus for the same bird.

1827. WAGLER, Systema Avium.

All the known species of *Buceros* are here given, and their synonymy as the author understood it. Proper discrimination regarding the specific value of those enumerated does not appear to have been made, as the errors of previous authors are continued, and the same species in several instances appears under different names. No new ones are added.

1828. HEMPRICH and EHRENBERG, Symbolæ Physicæ seu Icones et Descriptiones &c.

In this work, in footnotes, four Hornbills are mentioned, viz. *B. (LOPHOCEROS) HEMPRICHII*, named for the first time, *B. (Alophius) erythrorhynchus*, var. *leucopareus* (=ERYTHRORHYNCHUS, Temm.), *B. (LOPHOCEROS) hemileucus*, and *B. (LOPHOCEROS) forskali* ♀ (both =NASUTUS, Linn.). Species 32.

1829. CUVIER, Règne Animal.

Following the edition of 1817, the present one merely gives a short list of the existing species arranged in two groups—those with a prominent casque, and those without. No attempt is made to investigate the validity of the different species given; and the lists are *résumés* of those of the authors who preceded Cuvier.

1831. LESSON, Traité Ornithologique.

This author places this family of birds in the genus *Buceros* (*Calao*) of his Bucéridées, including with them the Motmots. He divides the genus into three subgenera—TOCKUS, BUCEROS, and BUCORVUS. The first comprises three species, ERYTHRORHYNCHUS, NASUTUS (here called *hastatus*, Cuv.), and GRISEUS. The second contains MELANOLEUCUS, Licht. (called *coronatus*, Shaw), BIROSTRIS, Scop. (called *ginginianus*, Lath.), CASSIDIX, BUCCINATOR, MALABARICUS, BICORNIS, PLANICORNIS (called *hydrocorax*, Linn.), LEUCOCEPHALUS (called *sulcatus*, Temm.), UNDULATUS, RHINOCEROS, SYLVESTRIS, and VIGIL (called *galeatus*, Gmel.). The third subgenus has only ABYSSINICUS. No species are described as new.

1833. HODGSON, Asiatic Researches.

BUCEROS (ACEROS) NEPALENSIS described and figured. Species 33.

1835-40. RÜPPELL, Neue Wirbelthiere zu der Fauna von Abyssinien gehörig.

Three species are here mentioned, and two described for the first time, viz. BUCEROS (BYCANISTES) CRISTATUS and BUCEROS (TOCKUS) FLAVIROSTRIS, *B. limbatus* (=B. HEMPRICHII, Hemp. & Ehr.). Species 35.

1843. BLYTH, Journal of the Asiatic Society of Bengal.

In the volume for this year Mr. Blyth separates the bird without any lateral ridges upon the bill as distinct from *B. UNDULATUS*, Shaw, and names it *B. SUBRUFICOLLIS*, both species being found in India, the latter, however, apparently restricted to the Tenasserim

provinces. In the second part of the same journal the writer reviews the species of Asiatic Hornbills known to him, and makes some valuable communications regarding doubtful and bad species, and corrects some errors in synonymy. Species 36.

1845. GRAY AND MITCHELL, Genera of Birds.

In this great work the family BUCEROTIDÆ is composed of one subfamily, *Bucerotinae*, consisting of three genera, EURYCEROS, BUCEROS, and BUCORVUS. The first of these contains *E. prevosti* from Madagascar, which is not a Hornbill. The genus BUCEROS is made to contain all the species excepting ABYSSINICUS. Thirty-seven are given in all, of which thirty-two are valid—the incorrect ones being *B. homrai*, Hodgson (=BICORNIS, Linn.), *violaceus*, Shaw (=CORONATUS, Bodd.), *sulcatus*, Temm. (=LEUCOCEPHALUS, Vieill.), *lugubris*, Begbie (=COMATUS, Raffles), *cinerascens*, Temm. (=GRISEUS, Lath.), *pæcilorhynchus*, Lafr. (=NASUTUS juv., Linn.), *pica*, Scop. (=CORONATUS, Bodd.), *plicatus*, Lath. (=PLICATUS, Forst.), *ginginianus*, Lath. (=BIROSTRIS, Scop.). The genus BUCORVUS consists of a single species, ABYSSINICUS.

1847. BLYTH, Journal of the Asiatic Society of Bengal.

The female of ANTHRACOCEROS MALABARICUS is described as *Buceros nigrirostris*.

1849. SUNDEVALL, Ofversigt af Kongl. Vetenskaps-Akademiens Förhandlingar.

The female of *B. ELATUS*, Temm., described as *B. cultratus*.

1850. J. CASSIN, Journal of the Academy of Natural Sciences of Philadelphia.

A. FISTULATOR from Fantee and A. ALBO-CRISTATUS from the Gaboon described. Species 38.

1850. BLYTH, Journal of the Asiatic Society of Bengal.

A. MALABARICUS, Gmel., from the Deyra Doon, is named *B. affinis*, and Hutton quoted as the authority for the species.

1850. BONAPARTE, Conspectus Generum Avium.

The family BUCEROTIDÆ is made to contain two subfamilies, *Bucerotinae* and *Euricerotinae*. The latter contains but a single species, which is not a Hornbill. Many genera are created for the reception of the groups as selected, some of which it will be necessary to accept. The first is BUCORVUS, containing six species—ABYSSINICUS, ELATUS, ATRATUS, CRISTATUS, BUCCINATOR, and CYLINDRICUS. Next is BUCEROTURUS with a single species, VIGIL, Forst., called *galeatus*. BUCEROS, the third genus, has six species—*hydrocorax* (=PLANICORNIS, Merrem), BICORNIS, RHINOCEROS, *rhinoceroïdes*, *sublunatus* (these last two being races of RHINOCEROS), and *lunatus* (which is SYLVESTRIS, Vieill.). HYDROCISSA, the fourth genus, has also six species—*monoceros*, Shaw (=CORONATUS, Bodd.), *pica*, Scop. (=CORONATUS, Bodd.), GALERITUS, *violaceus*, Shaw (=CORONATUS, Bodd.), MALAYANUS, and EXARHATUS. CALAO is divided into three subgenera: the first, CASSIDIX, contains CASSIDIX and CORRUGATUS; the second is CALAO, with *plicatus*, ex Java (=UNDULATUS, Shaw), and *ruficollis* (=PLICATUS, Forst.); and the third, ACEROS, with NEPALENSIS.

BERENICORNIS is the sixth genus, with two species, viz. COMATUS and *macrourus* (=ALBOCRISTATUS, Cassin): this genus being identical with ANORRHINUS, Reich., and of a later date, cannot stand. TOCKUS has twelve species:—*sulcatus* (=LEUCOCEPHALUS, Vieill.), *ginginianus* (=BIROSTRIS, Scop.), GRISEUS, GINGALENSIS, *sulcirostris*, Wagl. (=PANINI, Bodd.), FASCIATUS, MELANOLEUCUS, NASUTUS, *pæcilorhynchus* (=NASUTUS juv., Linn.), ERYTHORHYNCHUS, *limbatus* (=HEMPRICHII, Ehrenb.), and FLAVIROSTRIS. The arrangement of the various species into groups is effected with good judgment; but some of the genera, and especially the subgenera, are unnecessary.

1854. BONAPARTE, Conspectus Volucrum Anisodactylorum.

A list of the BUCEROTIDÆ is given, similar to the one reviewed above, but with some new genera added. There are two subfamilies, as before; but BUCEROTINÆ is divided into three sections—a. *Bucorveæ*, b. *Buceroteæ*, and c. *Tockeæ*. The first includes the genus BUCORVUS, with two species, ABYSSINICUS and LEADBEATERI?, Vigors. I have been unable to find a description of this last form. The second section, *Buceroteæ*, contains nine genera and five subgenera. The author commences with CERATOGYMNA for ELATA; *Tmetoceros* for ATRATUS, CRISTATUS, CYLINDRICUS, BUCCINATOR, FISTULATOR, and *cultratus*, Sundev., which is the female of ELATUS. *Berenicornis* and *Buceroturus* contain the same species as in the 'Consp. Gen. Av.' given above. BUCEROS has RHINOCEROS, *rhinoceroïdes*, *sublunatus* (=RHINOCEROS, Linn.), and *lunatus* (=SYLVESTRIS, Vieill.). *Homrains* (=DICHOCEROS, Glog.) has BICORNIS. HYDROCERAX contains PLANICORNIS. HYDROCISSA has *monoceros*, Shaw (=CORONATUS, Bodd.), *pica*, Scop. (=CORONATUS, Bodd.), and furthermore is divided into two subgenera:—ANTHRACOCEROS, Reich., for *bicolor*, Eyton (=MALAYANUS, Raffl.), and GALERITUS; and ANORRHINUS, Reich. (instituted for quite another form), for *violaceus*, Shaw (=CORONATUS, Bodd.), MALAYANUS, and EXARHATUS. RHYTICEROS is divided into three subgenera, viz.:—a. ACEROS, with NEPALENSIS; b. *Cassidix*, with CASSIDIX and CORRUGATUS; c. RHYTICEROS, with *plicatus* (=UNDULATUS, Shaw) and *ruficollis* (=PLICATUS, Forst.). The third section, *Tockeæ*, has seven genera. The 1st, *Calao*, has *sulcatus* (=LEUCOCEPHALUS, Vieill.); the 2nd, PENELOPIDES, contains *sulcirostris* (=MANILLÆ, Bodd.) and PANAYENSIS (=PANINI, Bodd.); the 3rd, *Meniceros*, Glog. (forestalled by LOPHOCEROS, Hemp. & Ehrenb.), has *ginginianus*, Lath. (=BIROSTRIS, Scop.); the 4th, RHINOPLAX, Glog. (instituted for VIGIL, Forst.), contains GRISEUS and GINGALENSIS; the 5th, *Grammicus*, contains FASCIATUS, NASUTUS, *hastatus*, Cuv. (=NASUTUS, Linn.), and *limbatus* (=HEMPRICHII, Ehrenb.); the 6th, RHYNCHOCEROS, has MELANOLEUCUS; TOCKUS, the last genus, contains FLAVIROSTRIS and ERYTHORHYNCHUS. The second subfamily, *Eurycerotinae*, contains *Euryceros prevosti*, which is not a Hornbill and should not be included in this family. It will be seen that this author has employed the various genera in a way never intended by those who established them, and in several instances has placed together species belonging to entirely different genera.

1855. HARTLAUB, Journal für Ornithologie.

TOCKUS SEMIFASCIATUS described. Species 39.

1856. BLYTH, Journal of the Asiatic Society of Bengal.

TOCKUS TICKELLI described. Species 40.

1856. J. CASSIN, Proceedings of the Academy of Natural Sciences of Philadelphia.

TOCKUS CAMURUS described. Species 41.

1859. CABANIS and HEINE, Museum Heineanum.

A catalogue of the species contained in the collection of M. Heine is given. Many new genera are adopted, and some proposed, while the names of others are changed. The new genera are *Platyceros* for *hydrocorax* (= *PLANICORNIS*, Merrem), *SPHAGLOBUS* for *ATRATUS*, Temm., *BYCANISTES* for *BUCCINATOR*, Temm., and *CRANORRHINUS* for *CASSIDIX*, Temm. No new species are described.

1860. GOULD, Proceedings of the Zoological Society of London.

TOCKUS HARTLAUBI described. Species 42.

1862. SCHLEGEL, Museum d'Histoire Naturelle des Pays-Bas.

A list of the species in the Museum of Leyden is here given. Nearly all the known species are enumerated, and one described as new—*B. CAFER*. All are included in one genus, *BUCEROS*. *B. RHINOCEROS* is divided into three races—*B. rhinoceros sumatranus* from Sumatra, *B. rhinoceros borneoensis* from Borneo, and *B. rhinoceros lunatus* (= *SYLVESTRIS*, Vieill.) from Java. To *B. ABYSSINICUS* are also assigned three races, viz. *B. carunculatus abyssinicus* from Abyssinia, *B. carunculatus guineensis* from West Africa, and *B. carunculatus cafer* from Caffraria. The first and last of these only are in this Monograph considered entitled to specific rank. Species 43.

1862. SCHLEGEL, Nederlandsch Tijdschrift voor de Dierkunde.

TOCKUS HARTLAUBI redescribed as *Buceros nagtglasii*.

1863. SCHLEGEL, Nederlandsch Tijdschrift voor de Dierkunde.

BUCEROS (TOCKUS) CAMURUS redescribed as *B. pulchrirostris*.

1864. TICKELL, Ibis.

An account of the Hornbills of India and Burmah, with an interesting history of the habits of some of the species, is here given, and T. TICKELLI, Blyth, more particularly described.

1865. HARTLAUB, Proceedings of the Zoological Society of London.

TOCKUS MONTEIRI described. Species 44.

TOCKUS FLAVIROSTRIS redescribed as *T. elegans*.

1869. CABANIS, von der Decken's Reisen in Ost-Africa, part iii.

BUCEROS (TOCKUS) DECKENI described. Species 45.

1870. FINSCH and HARTLAUB, von der Decken's Reisen in Ost-Africa.

The species of this family inhabiting East Africa are here given, with their synonymy. Seven in all are recorded; no new ones described.

1870. SCLATER, Proceedings of the Zoological Society of London.

BUCEROS (BYCANISTES) SUBCYLINDRICUS described. Species 46.

1870. HARTLAUB and FINSCH, von der Decken's Reisen in Ost-Africa.

TOCKUS MELANOLEUCUS redescribed as *Buceros pallidirostris*.

1871. G. R. GRAY, Annals and Magazine of Natural History.

BUCEROS (PHOLIDOPHALUS) CASUARINUS described from a head. Species 47.

1872. T. C. JERDON, Ibis.

BUCEROS AUSTENI described. Species 48.

1873. R. B. SHARPE, Zoological Record for 1871.

BUCEROS (ANORRHINUS) LEUCOLOPHUS separated from A. ALBO-CRISTATUS, Species 49.

1873. D. G. ELLIOT, Ibis.

PHOLIDOPHALUS SHARPII described as *Buceros sharpii*. Species 50.

1873. BARBOZA DU BOCAGE, Proceedings of the Zoological Society of London.

A paper chiefly upon the three forms of B. ABYSSINICUS, two of which are considered good species, viz. B. ABYSSINICUS and B. CAFER. The third, *B. guineensis* of the author, is probably the young of B. PYRRHOPS, Elliot.

1873. A. HUME, Stray Feathers.

A small form of R. PLICATUS from Narcondam Island, described as R. NARCONDAMI. Species 51.

1874. MURIE, Proceedings of the Zoological Society of London.

A paper describing the nature of the food-sac ejected by the Hornbills, and proving it to be the epithelial lining of the gizzard.

1876. R. B. SHARPE, Transactions of the Linnean Society.

CRANORRHINUS WALDENI described. Species 52.

1877. D. G. ELLIOT, Annals and Magazine of Natural History.

BUCEROS PYRRHOPS from the Congo described as distinct, and the three conspecies of the genus, as given by Schlegel, considered as probably representing distinct species. Species 53.

1877. LORD TWEEDDALE, Proceedings of the Zoological Society of London.

The following Hornbills are described :—HYDROCORAX MINDANENSIS as *Buceros MINDANENSIS*, HYDROCORAX SEMIGALEATUS, also in the genus *Buceros*, and PENELOPIDES AFFINIS. Species 56.

1877. D. G. ELLIOT, Annals and Magazine of Natural History.

ANTHRACOCEROS FRATERCULUS from Cochin China described. Species 57.

1877. CABANIS and REICHENOW, Journal für Ornithologie.

BYCANISTES ALBOTIBIALIS described as BUCEROS ALBOTIBIALIS. Species 58.

1880. OUSTALET, Bulletin de l'Association Scientifique de France.

LIMONOPHALUS MONTANI described as BUCEROS MONTANI. Species 59.

1880. CABANIS, Journal für Ornithologie.

BUCEROS (BYCANISTES) SUBQUADRATUS, from Angola, described. Species 60.

GENERA.

The genera proposed for the various species of Hornbills are very numerous; and many have been adopted by those authors who have written upon these birds. At first sight it would appear that so many divisions were superfluous; but it depends in a great measure on what kind of characters, and how many of them, an author deems necessary before he is willing to establish a genus. The casques of the Hornbills are very peculiar; and there are many species which have this formation very similar to each other in appearance and structure; and such individuals naturally constitute separate groups. I do not see any impropriety in considering these entitled to generic rank. Some are certainly generically distinct, viz. BUCORVUS ABYSSINICUS, HYDROCORAX PLANICORNIS, and RHINOPLAX VIGIL with its solid bony casque. No one would, I think, place the three birds above named in the same genus; and having acknowledged this, it is impossible to stop with them, for there are many species that cannot be placed with either of the three mentioned. With but few exceptions, however, all the birds that have been selected in this Monograph as types of the different genera have others in the family closely allied to them; and thus groups are readily formed and their generic position rendered clear. In the following list of genera I commence with Brisson, whose work in this respect is accepted by ornithologists.

Type.

1760. HYDROCORAX, Brisson, Ornithologie, vol. iv. p. 565	<i>Hydrocorax planicornis</i> , Merrem.
1766. BUCEROS, Linnæus, Syst. Nat. vol. i. p. 153	<i>Buceros rhinoceros</i> , Linn.
1828. LOPHOCEROS, Hemp. & Ehrenb. Symb. Phys.	<i>Buceros nasutus</i> , Linn.
1828. ALOPHIUS, iid. ibid. (nec Entomol.)	<i>Buceros erythrorhynchus</i> , Temm.
1831. TOCKUS, Less. Trait. Ornith. p. 252	<i>Buceros erythrorhynchus</i> , Temm.
1831. BUCORVUS, id. ibid.	<i>Buceros abyssinicus</i> , Linn.
1842. DICHOCEROS, Gloger, Hand- und Hilfsbuch, p. 335	<i>Buceros bicornis</i> , Linn.
1842. RHINOPLAX, id. ibid.	<i>Buceros vigil</i> , Forst.
1842. MENICEROS, id. ibid.	<i>Buceros rhinoceros</i> , Linn.
1842. RHYNCHOCEROS, id. ibid.	<i>Buceros melanoleucus</i> , Bodd.
1844. ACEROS, Hodg. Gray's Zool. Misc. p. 85	<i>Buceros nepalensis</i> , Hodg.
1847. TMETOCEROS, Cab. Wieg. Archiv, p. 345	<i>Buceros abyssinicus</i> , Linn.
1849. ANORRHINUS, Reich. Syst. Av. pl. 49	<i>Buceros galeritus</i> , Temm.
1849. ANTHRACOCEROS, id. ibid. pl. 50	<i>Buceros malabaricus</i> , Gmel.
1849. RHYTICEROS, id. ibid.	<i>Buceros undulatus</i> , Shaw.
1849. CRANOCEROS, id. ibid.	<i>Buceros vigil</i> , Forst.
1849. GRAMMICUS, id. ibid. pl. 49,	<i>Buceros nasutus</i> , Linn.
1849. PENELOPIDES, id. ibid.	<i>Buceros panini</i> , Bodd.
1849. BUCORAX, Sundev. Cefvers. Kongl. Vetensk. Akad.
Förh. p. 161	<i>Buceros abyssinicus</i> , Linn.
1850. CASSIDIX, Bp. Consp. Gen. Av. p. 90	<i>Buceros cassidix</i> , Temm.

	Type.
1850. CALAO, Bp. (nec Less.) Consp. Gen. Av. p. 90 . . .	<i>Buceros cassidix</i> , Temm.
1850. HYDROCISSA, Bp. Consp. Gen. Av. p. 90 . . .	<i>Buceros exarhatus</i> , Temm.
1850. BUCEROTURUS, id. ibid. p. 89 . . .	<i>Buceros vigil</i> , Forst.
1850. BERENECORNIS, id. ibid. p. 91 . . .	<i>Buceros comatus</i> , Raffles.
1854. CERATOGYMNA, Bp. Consp. Voluer. Anisod. p. 2. . .	<i>Buceros elatus</i> , Temm.
1854. TMETOCEROS, Bp. (nec Cab.) Consp. Voluer. Anisod. p. 2 . . .	<i>Buceros atratus</i> , Temm.
1854. HOMRAIUS, Bp. Consp. Voluer. Anisod. p. 2 . . .	<i>Buceros bicornis</i> , Linn.
1859. PLATYCEROS, Cab. & Hein. Mus. Hein. p. 174 . . .	<i>Buceros planicornis</i> , Merrem.
1859. BYCANISTES, iid. Mus. Hein. p. 171 . . .	<i>Buceros buccinator</i> , Temm.
1859. SPHAGOLOBUS, iid. ibid.	<i>Buceros atratus</i> , Temm.
1859. CRANORRHINUS, iid. Mus. Hein. p. 173 . . .	<i>Buceros cassidix</i> , Temm.
1873. OXYCEROS, Hume, Nests and Eggs of Indian Birds . . .	<i>Buceros birostris</i> , Scop.
1878. PHOLIDOPHALUS, Elliot, Mon. Bucerotidæ . . .	<i>Buceros fistulator</i> , Cass.
1882. LIMONOPHALUS, id. ibid.	<i>Buceros montani</i> , Oust.

KEY TO GENERA AND SPECIES.

1. Casque curving from base of culmen, horn-shaped, opened
or closed in front	I. BUCORVUS.
<i>a.</i> Casque with large circular opening anteriorly	1. <i>abyssinicus</i> .
<i>b.</i> Casque with small circular opening anteriorly	2. <i>pyrrhops</i> .
<i>c.</i> Casque completely closed anteriorly	3. <i>cafer</i> .
2. Casque very large, rounded on upper edge, two thirds
length of bill	II. BUCEROS.
<i>a.</i> Casque very much turned up anteriorly	1. <i>rhinoceros</i> .
<i>b.</i> Casque nearly straight on upper edge for entire length	2. <i>sylvestris</i> .
3. Casque flat anteriorly, concave, corners turned slightly
upwards	III. DICHOCEROS.
<i>a.</i> General plumage black and white; tail white, with
black band on terminal third	1. <i>bicornis</i> .
4. Casque flat, widest posteriorly, rather pointed anteriorly	IV. HYDROCORAX.
<i>a.</i> General plumage chestnut-red and black; tail white.	
<i>a'</i> . Casque with anterior margin compressed and elevated.	
<i>a''</i> . Bill red	1. <i>planicornis</i> .
<i>b''</i> . Bill, apical half white; tail buff	2. <i>mindanensis</i> .
<i>b'</i> . Casque anteriorly curving gradually to the culmen	3. <i>semigaleatus</i> .
5. Casque solid	V. RHINOPLAX.
<i>a.</i> Median rectrices greatly exceeding the rest in length	1. <i>vigil</i> .

6. Casque large, compressed laterally, highest anteriorly
 a. Casque white, with black patch; outer tail-feathers white.
 a'. Black on casque not extending to maxilla
 b'. Black on casque extending onto the maxilla
 b. Casque white, with black patch; tail-feathers black, terminal third white.
 a'. Black on casque not extending to maxilla
 b'. Black on casque extending onto the maxilla
 c. Casque entirely white
7. Casque high, keel-shaped, nearly half the length of bill, corrugated laterally
 a. Tail white for its entire length
 b. Tail black for half its basal length, remainder chestnut
 c. Tail cinnamon; base and apical portion greenish white
 d. Tail white, with an apical black band
8. Casque small, upright, compressed; bill grooved
 a. Tail dark brown; bar of deep buff two thirds from base; apical black band
 b. Tail deep chestnut; apical black band. Base of maxilla grooved perpendicularly
 c. Tail rufous, with a black band at base; tip blotched with black. Base of maxilla smooth
9. Casque very large, extending over half the maxilla, flat in front, upright
 a. Tail, two central rectrices black, remainder white
10. Casque extending nearly all over the maxilla, upper edge longest
 a. Tail dark green; terminal third of rectrices, except two median ones, white
11. Casque extending over two thirds of the maxilla, narrowed, compressed anteriorly, tip at right angle to culmen
 a. Tail pure white. Head crested
12. Casque longer than bill, pointed anteriorly, curved like maxilla
 a. Tail with median feathers black, rest tipped with white.
 a'. Base of mandible yellow; secondaries all black
 b'. Mandible all black; secondaries tipped with white
 b. Median rectrices black, lateral ones white, with broad transverse black band near base.
 a'. Casque black and white
 b'. Casque wholly black
- VI. ANTHRACOCEROS.
 1. *coronatus*.
 2. *convexus*.
 3. *malabaricus*.
 4. *fraterculus*.
 5. *malayanus*.
- VII. CRANORRHINUS.
 1. *cassidix*.
 2. *corrugatus*.
 3. *waldeni*.
 4. *leucocephalus*.
- VIII. PENELOPIDES.
 1. *manillæ*.
 2. *panini*.
 3. *affinis*.
- IX. CERATOGYMNA.
 1. *elata*.
- X. SPHAGLOBUS.
 1. *atratus*.
- XI. LIMONOPHALUS.
 1. *montani*.
- XII. BYCANISTES.
 1. *cristatus*.
 2. *buccinator*.
 3. *subquadratus*.
 4. *subcylindricus*.

- c.* Tail white, with a broad transverse black band.
- a'.* Thighs black 5. *cylindricus*.
- b'.* Thighs white 6. *albotibialis*.
13. Basal half of culmen covered by a low transversely grooved casque-like protuberance XIII. PHOLIDOPHALUS.
- a.* Tail with the lateral feathers black tipped with white 1. *fistulator*.
- b.* Tail with the lateral feathers pure white 2. *sharpii*.
- c.* Tail —? casque highest posteriorly, swollen, corrugated. (Head alone known.) 3. *casuarinus*.
14. Base of culmen covered with overlapping scales XIV. RHYTIDOCEROS.
- a.* Top of head and occiput dark chestnut.
- a'.* Base of bill grooved laterally 1. *undulatus*.
- b'.* Base of bill smooth, without grooves 2. *subruficollis*.
- b.* Head and neck reddish orange.
- a'.* Base of bill smooth 3. *plicatus*.
- b'.* Diminutive form of above 4. *narcondami*.
15. Culmen compressed laterally, elevated into a low crest XV. ANORRHINUS.
- a.* Tail white 1. *comatus*.
- b.* Tail black, tipped with white.
- a'.* Wings spotted with white 2. *albocristatus*.
- b'.* Wings black, without white spots 3. *leucolophus*.
- c.* Tail white, terminal third green 4. *galeritus*.
- d.* Tail greenish black, with pale tips 5. *tickelli*.
- e.* Tail slaty-grey with a greenish gloss, and tipped with white 6. *austeni*.
16. Bill without casque; base of maxilla ridged transversely XVI. ACEROS.
- Tail black at base, remainder pure white 1. *nepalensis*.
17. Casque low, grooved its entire length XVII. HYDROCISSA.
- General plumage black; superciliary stripe, cheeks, and throat white 1. *exarhata*.
18. Casque low, pointed anteriorly XVIII. LOPHOCEROS.
- a.* Bill black, yellow triangular space on base of maxilla 1. *nasutus*.
- b.* Bill dusky, tips and outlines of both maxilla and mandible whitish 2. *birostris*.
19. Culmen compressed laterally, elevated into a low crest, sometimes obsolete XIX. TOCKUS.
- a.* Bill bright red, base yellow. Head dark brown, streaked with white 1. *melanoleucus*.
- b.* Bill yellowish, tip black.
- a'.* Two rectrices next to outermost one white for their entire length 2. *fasciatus*.

b. Two rectrices next to outermost one white for half their apical length	3. <i>semifasciatus</i> .
c. Bill yellow, tip reddish brown. Head and neck leaden grey	4. <i>flavirostris</i> .
d. Bill deep red. Head and neck blackish brown	5. <i>hemprichii</i> .
e. Bill deep red. Head and neck light grey	6. <i>monteiri</i> .
f. Bill pale red at base, yellowish at tip. Outer rectrices white for two inches from tips	7. <i>griseus</i> .
g. Bill greenish white, brown at base. Outer rectrices greenish black, apical third white	8. <i>gingalensis</i> .
h. Bill deep red. Head and neck leaden grey; superciliary stripe white	9. <i>erythrorhynchus</i> .
i. Bill rufous. Head, neck, and underparts white . . .	10. <i>deckeni</i> .
j. Bill black, tip red. Tail greenish black, outermost rectrices tipped with white	11. <i>hartlaubi</i> .
k. Bill bright red. Tail dark olive-brown, edged with rufous .	12. <i>camurus</i> .

CLASSIFICATION.

The position of the BUCEROTIDÆ among birds is generally conceded by ornithologists to be between the Kingfishers and the Hoopoes. As long ago as 1838 the late Mr. Blyth * compared the species of this family with the Hoopoes, and regarded them as a distinct tribe, to which he gave the name *Appendirostres*. The Hornbills have many peculiar structural characters, the most conspicuous of which is, of course, the formidable bill. This, however, with perhaps one exception (*RHINOPLAX VIGIL*), is formidable in appearance only, as it is mainly hollow and permeated with a network of bony fibres. Another peculiarity is the possession of long and strong eyelashes. These are unusual among birds; and they may be given to protect the eyes from falling particles of wood or fruit dislodged by the creature when using its bill. Prof. Owen † states that the tongue is extremely short, of a triangular form, and smooth, and the œsophagus, as in the Toucans, is very wide and of nearly equal diameter as far as the gizzard. This last has its coats thick, and is of a more elongated form than those of the Toucans; its cuticular lining is very tough, and disposed in longitudinal ridges. There are no cæca; and the intestines are arranged in long and narrow loops, as in the Raven. The whole length of the intestines is five feet. At that period Prof. Owen considered the nearest ally of the Hornbills to be the Toucans; but the presence of a gall-bladder, which the Toucans do not possess, places the BUCEROTIDÆ nearer to the Crows. Although many of the Hornbills are of such large size, their skeletons are extremely light, and completely permeated by air, the very phalanges of the toes being hollow to their extremities. Like the Hoopoes, the Hornbills possess but ten tail-feathers; the Kingfishers, excepting those of the genus *Tanysiptera*, have twelve. Wallace ‡,

* Mag. Nat. Hist. vol. ii. p. 589.

† Proc. Zool. Soc. 1833, p. 102.

‡ Ann. Nat. Hist. 1856, xviii. p. 193.

on the other hand, places these birds nearest the Kingfishers, because, on account of the structure of the feet and toes, and also from their habits, the Hornbills are Fissirostral birds, though of an abnormal form. "Their very short legs and united toes, with a broad flat sole, are exactly similar to those of the Kingfishers. They have powerful wings; but their heavy bodies oblige them to use much exertion in flight, which is therefore not very rapid, though often extended to considerable distances." "We look upon Hornbills, therefore, as one of the abnormal developments of Fissirostral birds, of which they are the largest, the least elegant, and the least gifted with faculties for locomotion and for obtaining food; and their nearest affinities lie in the direction of the Kingfishers." Nitzsch, in his 'Pterylographie'*, places the three genera *Buceros*, *Upupa*, and *Alcedo* in his family LIPOGLOSSÆ, agreeing together in the absence of an aftershaft on the contour feathers, and in the feathered tip of the oil-gland, but differing in the characters of their plumage and in the distribution of their feather-tracts. *Buceros* and *Upupa* agree in having ten rectrices; *Alcedo* has twelve. Eyton† makes the BUCEROTIDÆ consist of three subfamilies, *Momotinæ*, *Upupinæ*, and *Bucerinæ*. Prof. Huxley‡, in his paper on the classification of birds, places the BUCEROTIDÆ in the Desmognathous division of the Carinatae, and arranges them between *Podargus*, *Alcedo*, and *Dacelo*; while Dr. Murie §, in his able paper on the UPUPIDÆ and their relationships, considers the members of that family to be the nearest allies of the Hornbills. He says, in concluding his remarks, "Lastly, what in exterior appearance can be more opposed to each other than such a great unwieldy horned bird as the Rhinoceros Hornbill (*Buceros rhinoceros*) and the graceful Hoopoe? Yet when *Tockus* is reached, size and outward peculiarities dwindle till we have a form in which can be recognized semblance to certain of the UPUPIDÆ. There is still a gap; but the very manifold structural agreements and adaptations thereof to habits &c. are strong evidences of congruity It suffices to say that economy in general, pterylosis, geographical distribution in part, and anatomy taken all in all, turn the scale in favour of the Hornbills as the group bearing the closest relation to the Hoopoes and Irrisors." It will be seen by the foregoing references that the chief writers on comparative anatomy have nearly agreed in the position which this family should occupy among birds, and that the Kingfishers on one side and the Hoopoes on the other are apparently its nearest allies. It must be borne in mind that, although the number of their rectrices are different, yet many of the habits of the Hornbills and the Kingfishers are very similar; and in the classification of birds these should always have considerable influence in forming an arrangement. The Hornbills as they exist at the present day, exhibit to us probably but a remnant of the great family which once dwelt amid the forests of that mighty eastern continent of which a large portion is now beneath the waters. So, many gaps exist, not one only, we may presume; and the diversified forms that would supply the necessary links to complete an unbroken chain of connected species throughout the family have long since disappeared. Our means, then, for working out this difficult problem are but scanty; but probably, amid this ever-shifting, changing world, they will never be more ample than at the present time.

* Ray Soc., edited by Selater, 1867.

† Proc. Zool. Soc. 1867, p. 446.

‡ Osteol. Av. (1867) p. 59.

§ Ibis, 1873, p. 181.

The BUCEROTIDÆ appear to me susceptible of arrangement into two divisions or subfamilies, containing the GROUND- and TREE-Hornbills respectively. Of the first of these we have at the present day three species, which possibly are the sole remaining members of what may have been an extensive group, and differ from the rest of the family in many particulars, the most conspicuous of which are the long legs formed for walking, whereas all the other species have short legs and can only hop. Myologically, also, these birds differ from the rest of the BUCEROTIDÆ in having only two muscles of the thigh represented, viz. the semitendinosus and the accessory semitendinosus; while the ambiens, the femoro-caudal, and the accessory femoro-caudal are absent. In the genera BUCEROS and TOCKUS the accessory femoro-caudal muscle is present. For this subfamily I would propose the term BUCORVINÆ. The other, BUCEROTINÆ, contains the remaining species, amounting to over fifty in number, comprising various groups allied to each other in the form of their casque and general colour of plumage; and in many of them their natural affinities are further upheld by their geographical distribution. The genera for these various forms have been many, all authors seeming to appreciate the difficulty of retaining these birds in one or two genera, as the older writers did. Of the thirty-four genera proposed, I have employed nineteen, which, I consider, fairly represent the different forms existing at the present day. The single genus *Tockus* is made to contain perhaps some species which might be deemed at least subgenerically different from the typical style; but I do not think it essential to make any such divisions, as in the main the species are all really members of one genus, the more or less slight elevation of the culmen being about the only character which could be urged in behalf of a separation of the species.

I commence the arrangement of the family with the genus BUCORVUS, containing three species. These birds have many characteristics to separate them from the other members of BUCEROTIDÆ. They are essentially ground-birds, yet ascend to the branches of trees; but their home is the ground, over which they walk with ease, while their relatives are tree-loving birds, descending but rarely to the earth, upon which they move by awkward hops. Here at once is perceived a great difference of habit; and to accommodate the Ground-Hornbills to their different life, their anatomical structure is modified. The second subfamily, BUCEROTINÆ, contains the greater part of the family; and these all live in and among the trees. I begin with the genus BUCEROS as containing a typical form of the true Hornbill, with two species, the chief (and perhaps only) difference between them being the upturned casque of the one, and the straight casque of the other. The third genus, DICHOCEROS, was formed for the well-known *bicornis*, the anterior corner of whose horns turns slightly upward. Next to this, as belonging to the flat-casque group, I place the genus HYDROCORAX, with three species; to be followed by RHINOPLAX, containing the extraordinary bird with solid casque, bare neck, and elongated median rectrices. Between all these genera gaps exist, which at the present time we are unable to fill; but we may feel assured that the extinct species which once supplied the desiderata were no less wonderful in structure than are their relatives which remain with us. ANTHRACOCEROS now introduces the species bearing an upright, more or less lengthened, laterally compressed casque. It contains five species, possessing, for the size of the birds, quite large casques, highest anteriorly. Four

of the species resemble each other closely in appearance, both as to casque and plumage; but the fifth has a perfectly white casque and a conspicuous superciliary stripe, not observed in the others. The seventh genus is *CRANORRHINUS*, with four species, distinguished by high, upright, keel-shaped casques extending half the length of the bill, with several undulations on the sides; and this genus is followed by *PENELOPIDES* with three species, of much smaller size, lower casques, and grooved bills. The birds of the last three genera, although they do not follow each other in regular order, unknown forms being required to supply the void between each two genera, still are closely related, and seem to constitute a natural group in the family. I now come to birds also possessing high and largely developed casques, but which are natives of a different zoogeographical region, and, as might be supposed, are rather widely separated from the members of the previous genera, yet at the same time possess characters in common with them. Several links are needed here to complete the line between the last-named genus and *CERATOGYMNA*, the one I now reach. This has but a single species, though one that is very remarkable for the form of its casque, which is large, bulky, extending over half the maxilla, rather flattened on its anterior face, which is upright; large irregular grooves also run along the sides of the casque near the maxilla. Allied to this is the single species of the genus *SPHAGOLOBUS*, but with a casque differently shaped and even larger than the other. Following this I place the genus *LIMONOPHALUS* with a single species, succeeded by *BYCANISTES* with six species, having, as a rule, a casque longer than the bill, pointed anteriorly, and curving to the culmen. Next is the genus *PHOLIDOPHALUS* with three species without a casque, but with a roughened casque-like protuberance covering the basal third of the culmen. These last five genera with their species form a well-marked group, with similarly coloured plumage; but, like the others, the links are wanting by which the existing species could be more readily arranged. Through the species of the last genus we are led to those birds which have bills devoid of casques, or else the latter represented by flattened protuberances having the appearance of bony scales. These are contained in the genus *RHYTIDOCEROS*, and consist of four species, one of which, like the members of the last genus, has transverse grooves at base of bill. The species of the fourteenth genus do not possess casques; but, in lieu of this appendage, the culmen is compressed laterally, and elevated into a low crest descending gradually forward to the culmen. There are five species included in this genus (*ANORRHINUS*), all having long loose crests on the head and occiput. *ACEROS*, the next genus, contains but one though a large and fine species, without casque, the culmen swollen at the base of maxilla. With this bird the list of the large Hornbills is finished, those that remain decreasing in size until we arrive at the last one, no larger than a Jay. *Exarhata*, which is the sole species of the genus *HYDROCISSA*, has really no ally in the family, and is remarkable for its crest-like casque, hardly to be distinguished from the maxilla, and is moreover peculiar for the lateral grooves running its entire length. It is a Celebes form, and probably is the sole survivor of a subgroup of this family. *LOPHOCEROS*, the next genus, contains but two species, allied closely to the members of the succeeding one, but differs in having the low casque extended to a point at its anterior end, returning to the culmen by an acute angle. The last genus into which I have divided the family is *TOCKUS*, containing twelve species. Some of these

have the culmen compressed and elevated into a low crest; others have merely a median ridge along the maxilla. It contains the smallest species of the family; and they are those which, according to the views of Dr. Murie, incline towards the *UPUPIDÆ*.

GEOGRAPHICAL DISTRIBUTION.

The members of the family Bucerotidæ are found in but three of the zoogeographical divisions of the globe, viz. the Ethiopian, Oriental, and Australian regions. Of these the last possesses but few representatives, confined to its Austro-Malayan division; for the Bucerotidæ are preeminently an Arctogean family, and have but few offshoots, and these are chiefly found among the Notogean islands lying nearest to those composing the Indo-Malayan division of the Oriental Region.

Commencing with that part of the earth's surface in which these birds are found that lies furthest to the westward, we enter Africa, in the Ethiopian Region, and observe that the members of the family are natives of all the countries lying south of the Great Desert of the Sahara, from the west to the eastern coasts and down to the Cape colony. Many species have great ranges, some reaching quite across the continent, as they are met with on both coasts; while others of less extended dispersion still overrun several districts in which other very distinct species have their habitat. This, indeed, is the case with members of the family in all the different zoogeographical divisions in which they are found; and it thus becomes extremely difficult, if not indeed well nigh impossible, to accurately designate the proper distribution of many of the species. In Africa, the countries lying on the west coast contain the largest number of the species found in the Ethiopian Region, viz. twenty out of twenty-seven, several of which, however, extend their range quite across the continent. Of the remaining seven, four only (so far as our knowledge of them enables us to decide) are confined to the countries on the east coast, two have their range unknown or, at least, undetermined, and one is generally distributed all over the continent. Beginning in the north-west, in Senegambia, at the southern margin of the Sahara, we find that this district contains four species—viz. *C. ELATA*, *A. ALBOCRISTATUS*, *L. NASUTUS*, and *T. SEMIFASCIATUS*. The first two and the last extend their range to the Gaboon; but how far into the interior, is unknown. The *L. NASUTUS* is a common African species with a very extensive distribution, being met with in various portions of the continent, such as Abyssinia on the east coast, and Sennaar, as well as the region of the Limpopo river to Damaraland, on the south. It would appear to be an inhabitant of nearly all Africa south of the desert. Proceeding southward along the west shore, we find in Fantee, Ashantee, and the Gold Coast, seven well-marked species not before mentioned—viz. *T. HARTLAUBI*, *A. LEUCOLOPHUS*, *B. CYLINDRICUS*, *S. ATRATUS*, *B. ALBOTIBIALIS*, *T. CAMURUS*, and *P. FISTULATOR*. Of these *T. HARTLAUBI* is only known from the Gold Coast. *A. LEUCOLOPHUS* seems to be

confined to Fantee. *B. CYLINDRICUS* is found in both Fantee and Ashantee. *S. ATRATUS* extends its range to Fernando Po. *B. ALBOTIBIALIS* was procured at the Gaboon, while *T. CAMURUS* is met with from the Gaboon to the Congo. The next district on the west coast that contains a species different from those already given is Angola, which has *B. SUBQUADRATUS*, *P. SHARPII*, *T. FASCIATUS*, and *T. MELANOLEUCUS*. Of these the second and third are apparently restricted to this district, or at all events they are very local in their range; but the last has a very wide distribution, being met with to the southward in Ovampo and Damaraland, and also across the east coast in the valley of the Shiré of the Zambesi region up to 5° south of the equator, and perhaps still further north. Damaraland possesses three species not yet given, viz. *T. ERYTHORHYNCHUS*, *T. FLAVIROSTRIS*, and *T. MONTEIRI*. Two of these have a very extended range, *T. ERYTHORHYNCHUS* having been procured, according to Schlegel, in Senegal and Sennaar. It has also been obtained at Ondonga on the Okovango river, and at Lake Ngami. Kirk met with it in the Zambesi region on the eastern coast; and Sundevall gives it as an inhabitant of Caffraria. *T. FLAVIROSTRIS* is found generally from Damaraland to the Transvaal, and as far north as Abyssinia. *T. MONTEIRI*, however, is restricted to Benguela and Damaraland. Passing now to the east coast in the enumeration of species still to be mentioned, we come to the district of the Zambesi and find two fine species, *B. BUCCINATOR* and *B. CRISTATUS*. The first named is not uncommon, according to Dr. Kirk, in the mountains and plains through which the Zambesi flows; but the extent of its range is not satisfactorily ascertained as yet, while *B. CRISTATUS* has been met with in the valley of the Shiré river, was procured by Capt. Speke and Grant in Uganda, and is also a native of Abyssinia, thus appearing to be distributed throughout the length of the east coast from the Zambesi northward; but, like its relative, its range towards the interior is unknown. Two species only of those restricted to the eastern part of the African continent remain to be noticed—*T. DECKENI* and *T. HEMPRICHII*. The latter is not found out of Abyssinia; and the exact locality of the former is not given by its describer. *B. SUBCYLINDRICUS*, whose habitat was unknown to its describer, has been procured in Central Africa by Dr. Emin Bey, at a place called Gor Aju*. *B. ABYSSINICUS* is found in Abyssinia and Sennaar. *B. PYRRHOPS* is as yet only known from the Gaboon; and *B. CAFER* is an inhabitant of Angola, Damaraland, and Caffraria. One species remains, regarding whose range we have no knowledge, viz. *P. CASUARINUS*, described by the late G. R. Gray from a head found in the British Museum without any locality attached.

Passing from the Ethiopian Region to the eastward, we come to our second zoogeographical division, viz. the Oriental Region. This is divided into four subregions (which are quite sufficient for the purpose of this review)—Indian, Ceylonese, Indo-Chinese, and Indo-Malayan. The Himalayas are separated into north-west and south-east portions, the dividing line being in the neighbourhood of Kitmandu. In the first two of these subregions, the Hornbills are scattered throughout the whole of their extent, from the Himalayas in the north to and including Ceylon in the south. In the Indo-Chinese subregion they are met with in the south-eastern

* Pelzeln, Verh. k.-k. zool.-bot. Gesell. Wien, 1881, p. 153.

Himalayas, Assam, Burmah, Arakan, Tenasserim, and Siam. How far they extend into China or the interior of Burmah is unknown; but it is most probable, as suggested by Wallace, that they will be found wherever the great forests may reach in that direction. Throughout the Indo-Malayan subregion they are pretty generally distributed, especially in the Malay peninsula and the great islands adjacent to it. To pass rapidly in review the ranges of the various species accepted as such in this Monograph, we find that in the Indian subregion we meet *A. MALABARICUS* at the base of the north-east Himalayas; and the species extends eastward along the chain into the Indo-Chinese subregion, throughout Nepal, Assam, and Burmah, and southward through British Burmah. It is not found in the Indian peninsula, that I am aware. Another species, *L. BIROSTRIS*, is distributed generally throughout India, not found in Assam, and never to the eastward of the Bay of Bengal. A third, *D. BICORNIS*, has an even wider distribution than either of the above, being met with in the hill-forests of all India, through British Burmah, and in the Malay peninsula and Sumatra of the Indo-Malayan subregion. Three other species are found in the Indian peninsula—*A. CORONATUS*, *T. GRISEUS*, and *T. GINGALENSIS*. The first and last of these extend southward through the Ceylonese subregion. Passing to the north-east into the Indo-Chinese subregion (besides *A. MALABARICUS*, already mentioned) we find the south-eastern Himalayas are inhabited by the fine *A. NEPALENSIS*, which ranges through Assam and the Tenasserim provinces, where also *A. TICKELLI* has been procured. In Cochin China *A. FRATERCULUS* has been obtained; but its exact dispersion is not known. Entering now the Indo-Malayan division of the Oriental Region, we find the Malayan peninsula tenanted by various species, the majority of which also inhabit the neighbouring islands. They are *B. RHINOCEROS*, *R. VIGIL*, *A. MALAYANUS*, *A. COMATUS*, *R. UNDULATUS*, and *A. GALERITUS*. All these are also natives of Sumatra and Borneo, except *A. COMATUS*, which is not found east of Sumatra, and *R. UNDULATUS*, which is not met with beyond Java. The last-named island also contains two species, *B. SYLVESTRIS* and *A. CONVEXUS*, the first peculiar to itself, the latter also found, according to Schlegel, in Sumatra and Borneo. The last two islands also possess one species not found elsewhere, *C. CORRUGATUS*. The remaining portions of the division that have any species of this family are the Philippines and Narcondam Island. The first possesses nine species, viz. *H. PLANICORNIS*, *H. MINDANENSIS*, *H. SEMIGALEATUS*, *C. LEUCOCEPHALUS*, *P. PANINI*, *P. MANILLÆ*, *P. AFFINIS*, *C. WALDENI*, and *L. MONTANI*. Narcondam Island contains *R. NARCONDAMI*. Should, however, this eventually be considered only a local race of *R. PLICATUS*, mentioned below, then it would extend the range of that species, as Narcondam Island lies further to the westward than any recorded habitat of *R. PLICATUS*.

In the Australian Region the Hornbills are confined entirely to Celebes and the Moluccan and Papuan groups of the Austro-Malayan division, where but four species of the family are met with, one, however, having a very extensive dispersion. Beginning with the most western island of this division, save the one given above, it is found that Celebes possesses two species, *H. EXARHATA* and *C. CASSIDIX*; and neither of them is met with in any other locality. The Moluccas contain *R. PLICATUS*, ranging from Mysol, New Guinea, to the Solomon Islands, these forming the extreme eastern limit that any species of this family reaches.

From the foregoing review of the dispersion of the various species of the Hornbills it will be perceived that the BUCEROTIDÆ are pretty equally divided at the present day between the Ethiopian and Oriental Regions, the first having twenty-seven and the latter twenty-nine species, while but a few, apparently the remaining representatives of a numerous race, are scattered about the islands of the Malay archipelago. Wallace has so thoroughly exhibited the probable way in which the great Australian continent has been broken up and left in the condition we now see it, and, at a similar period, how the Asiatic continent extended its limit until it almost reached its great southern neighbour, that it will be quite unnecessary to discuss the subject here; but unquestionably, during the convulsions of earth and sea, with the rising and falling land, the BUCEROTIDÆ, like many other great families of birds, have become widely scattered from the original continent which was once their home. Species have become separated from each other; some that may have been local in their habitat have disappeared altogether, or else are found today inhabiting one island only, such as those now living upon Celebes, Java, and certain of the Philippine Islands; while others that were widely dispersed over the old continent are now met with in many of the islands of the archipelago. In both the Oriental and Ethiopian Regions the range of many species appears arbitrarily drawn, and is inexplicable to us; but our difficulty may arise, in a measure at least, on the African continent, from our lack of knowledge of the fauna of the interior of that great land; for many of the species we now regard as restricted in their habitat may in reality have a wide dispersion. The same remark will apply to certain divisions of the Oriental Region; for we know nothing of the extent of range or number of species of this family beyond the north-eastern portion of the Indian subregion. How many may inhabit the unknown forests stretching away to the northward, or how far these birds may penetrate into the Palæarctic Region, or even to what degree, if at all, the family may be represented there, has not yet been ascertained. We can only rely upon future research for a solution of the problem. At present our knowledge simply points out to us that the "geographical distribution of the Hornbills seems quite in accordance with the almost universal belief of modern naturalists, that the present position of every living thing upon the earth is the result of that long and complicated series of geological changes and organic modifications which the globe has ever been and still is undergoing."

The following list gives, in a condensed form, the geographical distribution of the genera and species of the BUCEROTIDÆ :—

1. BUCORVUS.

Range of the Genus.

Ethiopian Region.

Range of the Species.

1. *Bucorvus abyssinicus*. East Africa: Abyssinia and Sennaar.
2. *Bucorvus pyrrhops*. Gaboon.
3. *Bucorvus cafer*. Angola; Damaraland; Caffraria.

2. BUCEROS.

Range of the Genus.

Indo-Malay division of the Oriental Region.

Range of the Species.

1. *Buceros rhinoceros*. Malay peninsula; Sumatra; Borneo.
2. *Buceros sylvestris*. Java.

3. DICHOCEROS.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Dichoceros bicornis*. Extensive hill-forests of all India; Assam; Arakan; Tenasserim; Malay peninsula; Sumatra.

4. HYDROCORAX.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Hydrocorax planicornis*. Luzon, Philippine Islands. This species was always stated by the older writers, and even, among modern ones, by as good an authority as the late Mr. Blyth, to be an inhabitant of the Moluccas. It has, however, comparatively lately, been ascertained to be a native of Luzon in the Philippine archipelago; and from that fact it is assumed that it is not found on any of the Moluccas. So far as known at present, the species seems to be restricted to the island named above.

2. *Hydrocorax mindanensis*. Mindanao, Philippine Islands.

3. *Hydrocorax semigaleatus*. Leyte, Philippine Islands.

5. RHINOPLAX.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Rhinoplax vigil*. Malay peninsula; Sumatra; Borneo.

6. ANTHRACOCEROS.

Range of the Genus.

Oriental Region.

INTRODUCTION.

xxix

Range of the Species.

1. *Anthracoceros coronatus*. Peninsula of India; Ceylon.
2. *Anthracoceros malabaricus*. Northern India; Nepaul; Bengal; Birmah; Assam and Tenasserim.
3. *Anthracoceros fraterculus*. Cochin China.
4. *Anthracoceros convexus*. Java; Borneo; Sumatra (*Ramsay*, P. Z. S. 1880, p. 14).
5. *Anthracoceros malayanus*. Malay peninsula; Sumatra; Borneo.

7. CRANORRHINUS.

Range of the Genus.

Oriental and Australian Regions.

Range of the Species.

1. *Cranorrhinus cassidia*. Celebes.
2. *Cranorrhinus corrugatus*. Sumatra; Borneo.
3. *Cranorrhinus waldeni*. Panay, Philippines.
4. *Cranorrhinus leucocephalus*. Mindanao, Philippines.

8. PENELOPIDES.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Penelopides panini*. Guimaras, Panay, Philippines.
2. *Penelopides manillæ*. Luzon, Philippines.
3. *Penelopides affinis*. Mindanao, Philippines.

9. CERATOGYMNA.

Range of the Genus.

Ethiopian Region.

Range of the Species.

1. *Ceratogymna elata*. Sierra Leone to the Gaboon.

10. SPHAGLOBUS.

Range of the Genus.

Ethiopian Region.

Range of the Species.

1. *Sphaglobus atratus*. Ashantee to Fernando Po.

INTRODUCTION.

11. LIMONOPHALUS.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Limonophalus montani*. Sooloo archipelago.

12. BYCANISTES.

Range of the Genus.

Ethiopian Region.

Range of the Species.

1. *Bycanistes cristatus*. Abyssinia; Zambesi district. Exact range unknown.
2. *Bycanistes buccinator*. Zambesi; Caffraria.
3. *Bycanistes subquadratus*. Angola.
4. *Bycanistes subcylindricus*. Gor; Aju; Central Africa.
5. *Bycanistes cylindricus*. Ashantee; Fantee.
6. *Bycanistes albotibialis*. Gaboon.

13. PHOLIDOPHALUS.

Range of the Genus.

Ethiopian Region.

Range of the Species.

1. *Pholidophalus fistulator*. Fantee; Senegal to the Gaboon.
2. *Pholidophalus sharpii*. Angola.
3. *Pholidophalus casuarinus*. —?

14. ACEROS.

Range of the Genus.

Oriental Region.

Range of the Species.

1. *Aceros nepalensis*. South-east Himalayas; hill region of Assam; Tenasserim provinces.

15. RHYTIDOCEROS.

Range of the Genus.

Oriental and Australian Regions.

Range of the Species.

1. *Rhytidoceros undulatus*. Malay peninsula; Sumatra; Java.
2. *Rhytidoceros subruficollis*. Arakan; Tenasserim provinces.
3. *Rhytidoceros plicatus**. Ceram; Mysol; New Guinea; Solomon Islands.
4. *Rhytidoceros narcondami*. Island of Narcondam.

16. ANORRHINUS.

Range of the Genus.

Oriental and Ethiopian Regions.

Range of the Species.

1. *Anorrrhinus comatus*. Malay peninsula; Sumatra.
2. *Anorrrhinus albocristatus*. Sierra Leone to the Gaboon.
3. *Anorrrhinus leucolophus*. Fantee.
4. *Anorrrhinus galeritus*. Malay peninsula.
5. *Anorrrhinus tickelli*. Tenasserim hills.
6. *Anorrrhinus austeni*. North Cachar hills.

17. HYDROCISSA.

Range of the Genus.

Australian Region.

Range of the Species.

1. *Hydrocissa exarhata*. Celebes.

18. LOPHOCEROS.

Range of the Genus.

Oriental and Ethiopian Regions.

Range of the Species.

1. *Lophoceros nasutus*. Senegambia; Damaraland; Abyssinia; Sennaar; Limpopo river.
2. *Lophoceros birostris*. India generally.

19. TOCKUS.

Range of the Genus.

Oriental and Ethiopian Regions.

* In the article on this species the first line should read, "This bird was separated from the well-known *R. undulatus*, Shaw"—not *R. plicatus*, Forst.

Range of the Species.

1. *Tockus melanoleucus*. Angola; Ovampoland; river Shiré to 5° south of the equator.
2. *Tockus fasciatus*. Angola.
3. *Tockus semifasciatus*. Senegambia; Fantee.
4. *Tockus flavirostris*. Damaraland; Abyssinia.
5. *Tockus hemprichii*. Abyssinia.
6. *Tockus monteiri*. Damaraland.
7. *Tockus gingalensis*. Ceylon.
8. *Tockus erythrorhynchus*. Damaraland; Okovango river; Lake Ngami.
9. *Tockus deckeni*. Voi-Fluss; Taita, East Africa.
10. *Tockus griseus*. Malabar.
11. *Tockus hartlaubi*. Gold Coast.
12. *Tockus camurus*. Gold Coast to Gaboon.

LIST OF PLATES.

1. Bucorvus abyssinicus	Plate I.	31. Bycanistes albotibialis	Plate XXXI.
2. Bucorvus pyrrhops	II.	32. Pholidophalus fistulator	XXXII.
3. Bucorvus cafer	III.	33. Pholidophalus sharpii	XXXIII.
4. Buceros rhinoceros	IV.	34. Pholidophalus casuarinus	XXXIV.
5. Buceros sylvestris	V.	35. Rhytidoceros undulatus	XXXV.
6. Dichoceros bicornis	VI.	36. Rhytidoceros subruficollis	XXXVI.
7. Hydrocorax planicornis	VII.	37. Rhytidoceros plicatus	XXXVII.
8. Hydrocorax mindanensis	VIII.	38. Rhytidoceros narcondami	XXXVIII.
9. Hydrocorax semigaleatus	IX.	39. Anorrrhinus comatus	XXXIX.
10. Rhinoplax vigil	X.	40. Anorrrhinus albocristatus	XL.
11. Anthracoceros coronatus	XI.	41. Anorrrhinus leucolophus	XLI.
12. Anthracoceros convexus	XII.	42. Anorrrhinus galeritus	XLII.
13. Anthracoceros malabaricus	XIII.	43. Anorrrhinus tickelli	XLIII.
14. Anthracoceros fraterculus	XIV.	44. Anorrrhinus austeni	XLIV.
15. Anthracoceros malayanus	XV.	45. Aceros nepalensis	XLV.
16. Cranorrrhinus cassidix	XVI.	46. Hydrocissa exarrhata	XLVI.
17. Cranorrrhinus corrugatus	XVII.	47. Lophoceros nasutus	XLVII.
18. Cranorrrhinus waldeni	XVIII.	48. Lophoceros birostris	XLVIII.
19. Cranorrrhinus leucocephalus	XIX.	49. Tockus melanoleucus	XLIX.
20. Penelopides manillæ	XX.	50. Tockus fasciatus	} L.
21. Penelopides panini	XXI.	51. Tockus semifasciatus	
22. Penelopides affinis	XXII.	52. Tockus flavirostris	LI.
23. Ceratogymna elata	XXIII.	53. Tockus hemprichii	LII.
24. Sphagolobus atratus	XXIV.	54. Tockus monteiri	LIII.
25. Limonophalus montani	XXV.	55. Tockus griseus	LIV.
26. Bycanistes cristatus	XXVI.	56. Tockus gingalensis	LV.
27. Bycanistes buccinator	XXVII.	57. Tockus erythrorhynchus	LVI.
28. Bycanistes subquadratus	XXVIII.	58. Tockus deckeni	LVII.
29. Bycanistes subcylindricus	XXIX.	59. Tockus hartlaubi	LVIII.
30. Bycanistes cylindricus	XXX.	60. Tockus camurus	LIX.

TO THE BINDER.

The page with the definition of subfamily BUCORVINÆ to go before the Plate of BUCORVUS ABYSSINICUS, and that with the definition of the subfamily BUCEROTINÆ to go before the Plate of BUCEROS RHINOCEROS.

The volume to be bound as follows :—

Titlepage.

Dedication.

Preface.

Introduction.

List of Plates.

Generic Plates, with description of Genera.

Description of Subfamilies as stated above.

Plates and text according to the List of Plates.

J. Smit lith.

Hanhart imp.

1 BUCORVUS ABYSSINICUS.	5 RHINOPLAX VIGIL.
2 " RHINOCEROS	6 ANTHRACOCEROS CORONATUS.
3 DICHOCEROS BICORNIS	7 CRANORRHINUS CASSIDIX.
4 HYDROCORAX PLANICORNIS.	8 PENELOPIDES MANILLÆ.
9 CERATOGYMNA ELATA.	

GENERIC CHARACTERS.

Family BUCEROTIDÆ.

Subfamily Bucorvinæ.

Genus I. BUCORVUS.

Head crested. Nude skin around the eyes, on sides of neck, and throat, with a pendent nude gular pouch. Bill long, powerful, curved, terminating in a sharp point. Nostrils basal, exposed. Casque large, extending from base of culmen for about one third its length, rising considerably above the maxilla, curved or straight, and open or closed anteriorly. Wings ample, reaching one third the length of the tail. Fifth primary longest, fourth and sixth next longest and equal. Tail long, ample, rounded. Legs long; tarsi covered with scales; toes rather short; claws short, strong.

Subfamily Bucerotinæ.

Genus I. BUCEROS.

Head slightly crested. Nude skin around eye. Bill long, powerful, terminating in a sharp point; culmen much curved. Casque very large, two thirds the length of the bill, rounded on the upper edge, straight, or much curved upward anteriorly. Nostrils small, basal, lateral, and hidden. Wings rather short; fourth and fifth primaries nearly equal and longest. Tail long, rounded. Tarsi short, robust, covered with strong broad scales. Toes of moderate length, the outer and inner ones united at base. Claws strong, curved, acute.

Genus II. DICHOCEROS.

Feathers on occiput moderately lengthened. Bill very long, stout, powerful, pointed at tip; culmen and gonys much decurved. Casque very broad, covering basal half of culmen, flat, concave anteriorly, with the corners turned slightly upwards into miniature horns projecting forwards. Nostrils small, basal, open. Bare skin around eyes. Wings ample, reaching one third the length of the rectrices. Tail long, broad, rounded. Tarsi short, stout, covered with large scales. Toes short, united at base by a membrane. Claws strong, acute.

Genus III. HYDROCORAX.

Head moderate, crested. Bill rather short, stout, pointed, much curved toward the tip. Casque extending two thirds the length of the maxilla, flat, wide at the posterior end, rather pointed anteriorly. Nostrils basal, small. Nude skin around eyes. Wings ample, rather long; fourth, fifth, and sixth primaries equal and longest. Tail long and rounded. Tarsi short, stout, covered with strong scales. Claws strong, curved, acute.

Genus IV. RHINOPLAX.

Occiput crested. Face, throat, neck, and spot on centre of back naked. Bill short, straight, stout, pointed. Casque solid, covering half the maxilla, broad on its anterior face, curved on

GENERIC CHARACTERS.

top, extends over the eyes, and returns to the base of the maxilla by a moderate curve. Nostrils basal, exposed. Wings long, reaching a short distance beyond the upper tail-coverts. Tail long, much rounded; two median rectrices twice the length of the rest. Tarsi short, stout, covered with strong scales. Toes stout. Claws strong, curved.

Genus V. ANTHRACOCEROS.

Head covered by a lengthened crest. Skin around eyes and space at base of mandible nude. Bill long, abruptly curved at tip. Casque covers two thirds the length of maxilla, is compressed laterally, extends over the head to above the eyes, and curving forwards returns to culmen by an acute angle. Nostril on top of maxilla, at base of casque, small, round, exposed. Wings moderate; secondaries as long as primaries; third primary longest. Tail long, much rounded. Tarsi short, stout, covered with strong scales. Toes short, stout. Claws curved, strong.

Genus VI. CRANORRHINUS.

Skin around eyes and on throat bare. Bill long, rather slender, graduating to a sharp point, crossed at base by several prominent transverse ridges. Casque high, upright, keel-shaped, swollen posteriorly, flattened anteriorly, covering nearly half the maxilla; curves gradually from the highest point of the anterior terminus, backwards to centre of the head. Wings ample; secondaries nearly as long as the primaries. Tail long, rounded. Tarsi and toes short, stout, covered with strong scales. Claws stout and acute.

Genus VII. PENELOPIDES.

Head covered by a lengthened crest. Bill rather short; base covered by a plate, on the maxilla with broad transverse grooves, on the mandible with narrow diagonal grooves. Casque low, compressed laterally; rises at base and extends for two thirds the length of maxilla, and terminates abruptly at a right angle to the culmen. Bare skin around eyes and at base of mandible. Wings long; fourth primary longest; first primary falcate. Tail long, slightly rounded. Tarsi and toes short, stout, covered with strong scales; outer and inner toes united at base. Claws strong, curved, acute.

Genus VIII. CERATOGYMNA.

Occiput crested. Side of face around the eye, throat and front of neck, with the exception of a narrow line in the centre covered with short feathers, and large gular pouch naked. Bill rather stout, very short. Culmen curved, especially at the apical third. Gonys nearly straight until within a short distance of the tip, where it curves downward. Casque very large, flat on its anterior face, keel-shaped on top, and swollen towards its posterior margin; highest in front, it curves rapidly downwards to above the eyes; several longitudinal grooves run along the sides. Wings ample, rather short. Tail long, much rounded. Tarsi and toes stout, short, strong.

Genus IX. SPHAGLOBUS.

Head covered by a long bushy crest. Skin around eyes, sides of face, throat, and gular pouch naked. Bill rather short, stout, pointed. Maxilla curved; gonys nearly straight. Casque very large, swollen laterally and rounded on top; rises from the maxilla about one third its length from the tip at a rather acute angle, curves backwards over the head, and returns to the maxilla at a sharp angle. Wings rather long. Tail long, slightly rounded. Tarsi short, stout. Toes rather long, stout. Claws strong, acute.

e. Nodul
Tail long
out, cover

ndible and
compress
o culmen
sed. Wing
ch round
ong.

r to a sh
pright, hi
illa; corn
f the be.
Tarsi 2

ate, on u
s. Cas
macula
at base
g. shiny
oes with

except
ked. I
y shiny
ge, the
t in be
the sh
ng.

and 4
Cas
is the
on m
nd. 3

J. Smith lith.

1. SPHACOLOBUS ATRATUS.
2. BYCANISTES BUCCINATOR.
3. LIMONOPHALUS MONTANI.
4. PHOLIDOPHALUS FISTULATOR.
5. RHYTIDOCEROS UNDULATUS.

6. ANORRHINUS CALERITUS.
7. ACEROS NIPALENSIS.
8. HYDROCISSA ENARHATUS.
9. LOPHOCEROS NASUTUS.
10. TOKUS ERYTHORHYNCHUS.

Harbort imp.

GENERIC CHARACTERS.

Genus X. LIMONOPHALUS.

Head and occiput crested. Nude skin around eyes. Bill long, curved, stout, broad at base. Casque high, keel-shaped, smooth, commencing above the eyes, curving gradually forwards for two thirds the length of bill, and terminating at a right angle to the culmen, swollen posteriorly, compressed laterally, sharp at its anterior edge. Wings ample, but rather short. Tail long, broad, rounded, the outermost rectrices on either side much shorter than the middle ones. Feet stout; claws strong.

Genus XI. BYCANISTES.

Head covered by a long loose crest. Bill rather short, strong. Culmen curved to the tip. Gonys straight for two thirds the length from base; it then turns abruptly upwards, and follows the curve of the gape to the point. Casque enormous, longer than bill, swollen and rounded on top, compressed anteriorly, having several irregular transverse grooves; it rises a short distance from tip of maxilla at an acute angle, then curves backwards to the centre of the head, and returns to base of maxilla by a steep angle. Wings ample, long, second and third primaries equal and longest. Tail broad, long, much rounded. Nude skin around eyes. Tarsi short, stout, as are the toes, and both covered with strong scales.

Genus XII. PHOLIDOPHALUS.

Head crested. Bill short, stout, pointed. Mandible with several irregular narrow transverse grooves. Casque very low, covering base of maxilla, ridged transversely, and falling away posteriorly into the culmen. Wings rather short; third and fourth primaries nearly equal and longest. Tail rather long, much rounded. Tarsi and toes short, covered with strong scales.

Genus XIII. RHYTIDOCEROS.

Bill long, nearly straight, graduated to a point. Base of maxilla and mandible grooved transversely. Culmen covered for nearly half its basal length by a casque-like protuberance, apparently formed of large scales overlapping each other. Skin around eyes and on throat nude. Wings ample. Tail long and rounded. Tarsi short, stout. Toes rather long. Claws strong, curved.

Genus XIV. ANORRHINUS.

Head covered by a long loose bushy crest. Bill moderately long, stout, curved on culmen. Casque extending over half the maxilla, highest at its anterior end, whence it inclines gradually backward to the base of bill. Skin around eyes, cheeks, and throat naked. Wings long; fourth primary longest; secondaries nearly as long as the primaries. Tail long, broad, and rounded. Tarsi stout, short. Toes moderately long. Claws curved, acute.

GENERIC CHARACTERS.

Genus XV. ACEROS.

Head crested. Bill long, rather narrow, tapering to a point. Culmen swollen and slightly elevated at base. Casque none. Maxilla at base ridged, with several transverse grooves. Nude skin around eyes, at base of mandible, and on throat. Wings long; third, fourth, and fifth primaries nearly equal and longest. Tail long, slightly rounded. Tarsi rather short and stout. Toes long, covered with strong scales.

Genus XVI. HYDROCISSA.

Head slightly crested. Bill moderately long, curved, sharply pointed. Casque low, covers the entire culmen to within an inch of the tip of the maxilla, compressed laterally, and indented by three deep grooves for its entire length. Nude skin around eyes. Wings short. Tail rather short, rounded. Tarsi and toes short, stout, covered with strong scales.

Genus XVII. LOPHOCEROS.

Head with a short occipital crest. Bill long, curved, slender, graduated to a sharp point. Nostril round, open, and exposed. A small casque commences at base of culmen, and extends for a little over half its length, terminating in a projecting point. Wings long; third primary longest; first primary very short and narrow. Tail long, slightly rounded. Tarsi and toes short, stout, covered with strong scales.

Genus XVIII. TOCKUS.

Bill long, slender, much curved; maxilla compressed laterally along the culmen. No casque. Wings and tail long, the latter rounded. Tarsi and toes short, stout, covered with rather strong scales. Claws curved, strong, acute.

Subfamily BUCORVINÆ.

Bill long, powerful, curved, terminating in a sharp point. Casque high, and, with the exception of one species, open in front. Skin around eyes, on sides of neck, throat, and gular pouch naked. Wings reaching one third the length of the tail. Legs long, fitted for walking on the ground, upon which the species pass nearly all their time. Toes short and stout. The semitendinous and accessory semitendinous muscles of the thigh are represented; but the ambiens, femoro-caudal, and the accessory femoro-caudal are absent. The species are essentially birds which live upon the ground.

BUCORVUS ABYSSINICUS.

GROUND-HORNBILL.

a. BUCORVUS ABYSSINICUS.

- Buceros abyssinicus*, Bodd. Tab. Plan. Enlum. d'Auben. (no. 779); Gmel. ed. Linn. Syst. Nat. vol. i. p. 358. sp. 5 (1788); Lath. Gen. Syn. (1781) vol. i. p. 347. sp. 4; id. Ind. Ornith. (1790) vol. i. p. 143. sp. 4; id. 2nd Supp. (1801) p. 99; Shaw, Gen. Zool. (1811) vol. viii. p. 21; Temm. Plan. Col. (1824) vol. ii. sp. 5 (text); Vieill. Gal. Ois. vol. i. p. 321, t. 191; Sclat. Proc. Zool. Soc. (1865) p. 676; Bald. Naum. (1857) p. 109; Nitzs. Pterylog. Ray Soc. (1867) p. 102, pl. vi. figs. 1 & 2; Giebel, Thesaur. Ornith. vol. i. p. 495 (1872); Vieill. Ency. Méthod. vol. i. p. 302 (1823); Heugl. Journ. für Ornith. (1864) p. 270; Vieill. Nouv. Dict. Hist. Nat. vol. iv. p. 589 (1816); Bocage, Journ. Sc. Math. Lisb. vol. ii. p. 347. sp. 96 (1868); Eyton, Ost. Av. p. 63 (1867).
- Grand Calao d'Abyssinie*, Buff. Plan. Enlum. (1783) no. 779, juv.?
- Le Calao carunculé*, Levaill. Hist. Nat. Ois. d'Afr. (1806) vol. v. p. 109, pls. 230, 231 (adults), 232 (juv.).
- Buceros africanus*, Lath. Ind. Ornith. (1790) vol. i. p. 143. sp. 5; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 591; id. Ency. Méthod. (1823) p. 303.
- Buceros brachyotus*, Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 201.
- Bucorvus abyssinicus*, Less. Trait. Ornith. (1831) p. 256; G. R. Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 1; Layard, B. of S. Afr. (1867) p. 228; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856) vol. ii. p. 581; Sclat. Proc. Zool. Soc. (1864) p. 111; Gurney, Ibis (1868) p. 162; Ayres, Ibis (1869) p. 296; Bon. Consp. Gen. Av. (1850) p. 89. sp. 1; id. Consp. Vol. Anisod. (1854) p. 2; G. R. Gray, Hand-l. Birds (1870), pt. ii. p. 131. sp. 7919; Blanf. Geol. & Zool. Abyss. (1870) p. 320; Salvad. Cat. Uccelli Mar. Rosso e dei Bogos (1873) p. 57 (420).
- Tragopan abyssinicus*, Gray, List Gen. B. (1841) p. 65; Rüpp. Syst. Übers. (1845) p. 79. sp. 320; Heugl. Syst. Übers. p. 45. sp. 456.
- Tmetoceros abyssinicus*, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 175. no. 481; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 480.
- Buceros carunculatus abyssinicus*, Schleg. Mus. Pays-Bas (1862) p. 19.
- Bucorax abyssinicus*, Kirk, Ibis (1864) p. 325; Bocage, Proc. Zool. Soc. (1873) pp. 698-700, figs. 1 & 4; Sundev. Öfver. Kongl. Vetens. Akad. Förh. (1849) p. 161.
- Bucorax cafer*, Lay. B. of S. Afr., Sharpe ed. p. 122. sp. 116 (reference to pls.).
- Abba Gumba* (Bruce).

Casque large, high, much curved on top, horn-shaped; large circular opening in front.

HAB. Abyssinia (RÜPPELL, BLANFORD); Zambesi region (KIRK ?); Natal (AYRES ?).

b. BUCORVUS GUINEENSIS.

Bucorax abyssinicus, Hartl. Ornith. West-Afr. (1857) p. 165; Monteiro, Ibis (1862) p. 338. sp. 37; Gurney, Ibis (1861) p. 132 ?

Buceros carunculatus guineensis, Schleg. Mus. Pays-Bas (1862) p. 20.

Bucorvus abyssinicus, Sharpe, Ibis (1869), p. 385; Reichenow, Journ. für Ornith. (1875) pp. 12, 49 ?

Buceros abyssinicus, Gurney, Anderss. B. of Damara-Land (1872) p. 205.

Bucorax cafer, Bar. du Bocage, Proc. Zool. Soc. (1873) p. 698, figs. 2, 6 (adults), fig. 5 (juv.).

Casque small, straight on top, tube-shaped; small oblong opening in front.

HAB. Gold Coast (SCHLEGEL); Sennaar (BALDAMUS ?); Damara-Land (ANDERSS. ?); Angola (MONTEIRO ?).

c. BUCORVUS CAFFER.

Buceros carunculatus cafer, Schleg. Mus. Pays-Bas (1862) p. 20.

Bucorvus leadbeateri, Gray, Hand-l. Birds, pt. ii. (1870) p. 131. sp. 7920.

Bucorax guineensis, Bocage, Proc. Zool. Soc. (1873) p. 699, 701, figs. 3 & 7 (juv.).

Casque small, excessively compressed, completely closed in front.

HAB. Caffraria (SCHLEGEL); Mossamedes (ANCHIETA).

There appear to be three forms or geographical races of this species, exhibiting their characters chiefly in the shape of the casque. Some writers have considered these differences sufficient to establish separate specific ranks, while others regard them as merely indicating age or possibly sex. While our material is not ample enough for us to determine positively that there is more than one species of the Ground-Hornbill, yet we certainly are able to decide that the differences met with in the form of the casque of individuals are not those belonging solely to young birds or characteristic of sex. I have therefore preferred in this paper to keep the three forms as distinct in their synonymy as I may have been able, but to consider all three as but geographical races of one variable species. Prof. Schlegel was the first to draw attention to the shapes of the various casques, and gave names to the possessors of them as follows: those individuals which have a high horn-like casque and a large round opening in front he called *Buceros carunculatus abyssinicus*, this being the typical *Buceros abyssinicus* of authors; those with a smaller, straighter casque, and a small opening in front, he named *Buceros carunculatus guineensis*, and for the third, with an exceedingly compressed casque, entirely closed in front, he proposed the appellation of *Buceros carunculatus cafer*. These three *conspecies* of the Professor inhabit respectively Eastern Africa north of Caffraria (rare in Sennaar), Western Africa, and Caffraria. Prof. Barboza du Bocage, in the 'Proceedings' of the Zoological Society of London for 1873, p. 698, has published a very interesting paper on these three varieties, and decides that two are certainly distinct, but that the third is still doubtful from want of sufficient material to enable him to form a judgment. He figures the heads of the three styles, and exhibits fairly their different characters. Unfortunately the names of two are transposed, and the *B. guineensis* (Schlegel) with the small casque is called *cafer*, and the *B. cafer* (Schlegel) with the closed casque is named *B. guineensis*. In attempting to define the geographical distribution of these three species (if they really are such), we are at once met with several awkward facts, which, with our present knowledge, can hardly reconcile ornithologists to the belief that these forms are really specifically distinct. To indicate what this difficulty is, I will give the distribution of the species as laid down by Professors Schlegel and Bocage. The *B. abyssinicus* (typical) inhabits Abyssinia, according to

Schlegel, rare in Sennaar. Layard (*vide* B. of S. Africa) gives it as found in the Cape colony. The *B. cafer* (Schlegel) inhabits Caffraria; and, as given by Prof. Bocage (if the name has not been transposed), the same bird, called by him *guineensis*, was procured on the coast of Guinea, thus making it range across the continent. If these birds are distinct, we have an extraordinary geographical distribution for continental forms, as *B. abyssinicus* of Abyssinia is thus cut off from its relatives in South Africa by *B. cafer*, which crosses the continent in the region of the Limpopo river and Damaraland. If the distribution of the forms as here given is correct, I shall not be able to reconcile myself to the belief that there are three distinct species of this bird until more information is obtained respecting them, and the boundaries of each one are definitely fixed; for it would seem at least strange that on a continent one species should inhabit the northern and southern portions, and the central part be occupied by a totally different species, and yet each keep their respective characters intact; for there must of necessity be many points where individuals of the two forms would meet and, in such closely allied birds as these under consideration might not unnaturally interbreed. A possible solution of the difficulty may be, that Layard did not describe his *B. abyssinicus* from a *South-African specimen*, as he states he never succeeded in getting one while there, supposing, like most authors, that there was only one species, and that Prof. Bocage's *B. cafer*, in his paper in the 'Proceedings' of the Zoological Society, was really intended for *B. guineensis*. In such a case, we should have the range of the forms as follows:—*B. abyssinicus* from East Africa, Abyssinia, and Sennaar; *B. guineensis* from Acra, West Africa; and *B. cafer* from all the region south of and including Damaraland and the Zambesi district. I have not been able to examine an authentic adult South-African specimen.

I have found some considerable difficulty in working out the synonymy of the three forms as presented here, because of course I have not had access to the specimens mentioned by the various writers; but I believe that, in the main, they will be found to be tolerably correct.

The Ground-Hornbill, although a frequenter of plains, is also a mountain-loving species, and has been met with at a considerable height above the sea; for, as stated by Blanford in his 'Zoology of Abyssinia,' it is chiefly found in that country at an elevation of from 4000–5000 feet, although also met with lower down. Monteiro, who procured the bird in Angola, states it to be everywhere present in that land, but more abundant in the interior, and very common on the mountain-range where Pungo Andongo is situated. For some unexplained reason, this species is an object of superstitious dread to the natives of the various countries it inhabits. Monteiro was unable to induce them to attempt its capture; and on Andersson's asking a chief in Ondonga to get him the eggs, he received the reply that it was not to be done, as they were soft to the touch and would break in pieces on the least handling. The Fingoes, says Layard, object to shoot them, lest they should lose their cattle by disease.

The food of this Hornbill appears to consist of almost any thing edible that falls in its way. In Abyssinia Blanford found it to be almost entirely insectivorous, the stomachs of those he examined containing chiefly large beetles and locusts, in one instance the remains of scorpions and large spiders. In a certain case, however, he found fragments of bones, apparently of a tortoise. Once he observed this bird near some mule-carasses, but not feeding on them, so far as he could see. In Angola, Monteiro found them to be omnivorous, feeding upon reptiles,

birds, eggs, beetles and all other insects, and also on the mandioca-roots and ginguba, or ground-nuts. In confinement, he says, it would readily eat fish and entrails of fowls; and once one of these birds, meeting some young chickens in the yard, immediately gulped down five or six of them. The Abba Gumba (by which name this species is known) feeds readily upon snakes; and their mode of attacking these, as related by Mr. Ayres, is peculiar. When a large serpent is discovered, three or four of the birds stretch out their wings and advance sideways towards it. The reptile, becoming irritated, seizes one by the wing-feathers, when all close around and strike the snake with their powerful bills, retreating as soon as it leaves its hold. This proceeding is repeated until their enemy is dead. Should the reptile attack them, they throw both wings forward, completely covering their heads and other vulnerable parts. Two or three kinds of land-tortoises frequenting the district of the Limpopo river are eaten by this Hornbill, which neatly pecks out every portion of flesh from the shell, leaving it quite clean, even eating the legs and head of the unfortunate creature. According to Layard, this species feeds on carrion; and their bodies emit such a stench in consequence that he never could induce any of his correspondents to send him one. They also devour great quantities of grubs and locusts. The flight is heavy, rather feeble, and somewhat noisy, and rarely prolonged for more than half a mile. The voice of this bird is an extraordinary one, very deep and powerful, composed, according to Blanford, of two notes—one, uttered chiefly on the ground, resembling the syllables *hūm-hūm*, the first higher and longer than the last. The second note was a peculiar booming sound, so much like the roar of the lion that it was with difficulty the two could be distinguished apart. They utter this noise both when on the ground and on the trees, and it can be heard for a considerable distance, particularly at night. Ayres compares the notes to the syllables *coo-coo*, and says that he has heard it at a distance of nearly two miles. The call of the female is similar to that of the male, but pitched a note or two higher. The male always calls first, the female immediately answering; and this is continued at intervals while they are feeding.

The Ground-Hornbill is gregarious, going in flocks of from five or six individuals up to one or two hundred, as stated by Monteiro. They are usually very shy and wild, and when alarmed fly to the nearest trees, choosing those with the densest foliage, where they squat on the branches or stand upright, quite motionless, with open bills. As soon as the object of their fears is discovered, the usual cry is uttered by one of them, and all fly away to another tree. When on the ground, and the grass is short, it is almost impossible to approach them. At a distance they greatly resemble turkeys; and the males have the habit of raising, then opening, and closing their tails exactly like those birds; the red wattle on the neck is also inflated, and becomes very conspicuous. Although this species walks well, its gait is awkward, and it presents a curious appearance as it goes slowly along, looking for food, turning over lumps of earth in search of insects, or poking its bill at any frog or other reptile that it may meet. While the ground is the chief resort of this species, over which Andersson says he has seen individuals running at a great rate, yet at night it is accustomed invariably to resort to the trees to roost. This species does not, I believe, incubate in the holes of trees, like other members of this family, but, according to Monteiro, the nest is built on the very highest *Adansonia*, in the hollow or cavity formed at the base or junction of the branches.

Bill long, curved, black, with a large red spot at base of maxilla. A curved casque rises at base of culmen, and has a round opening in front; the sides are grooved, these following the curve of the casque; as this grows from behind, it sheds off in front portions in the shape of rings, and in this way retains a regular length. The entire plumage is black, with the exception of the primaries, which are buff. A bare skin around the eye, continuing to the throat and part of the neck in front, is deep blue, bordered on its outer edges along the neck with scarlet, and terminating in a kind of scarlet bag at its lowest portion. Tarsi long and, with the feet, blackish. Bill $9\frac{1}{2}$ inches, casque $3\frac{1}{2}$, height 2; wing 21; tail 14; tarsus 6.

The *B. guineensis* of Schlegel differs in being smaller, the casque slightly closed in front; wing 18 inches; tail 11; tarsus $4\frac{3}{4}$ -5; bill along gape $6\frac{2}{3}$, casque 1 inch 10 lines, height $1\frac{2}{3}$ inch.

The *B. cafer* has the casque entirely closed in front. Length of bill $6\frac{2}{3}$ - $7\frac{2}{3}$ inches, casque 1 inch 11 lines, height 11-12 lines.

My thanks are due to the Publication Committee of the Zoological Society of London and to Prof. Bocage for permission to use the woodcuts that illustrate this article.

The figures are one third the natural size.

BUCORVUS PYRRHOPS.

RED-FACED GROUND-HORNBILL.

Bucorvus pyrrhops, Elliot, Ann. & Mag. Nat. Hist. (September 1877) vol. xx. p. 171.

HAB. Region of the Congo, Cachen?

In my article on *Bucorvus abyssinicus* I placed as races the two forms of Ground-Hornbills called respectively by Prof. Schlegel *Buceros carunculatus guineensis* and *B. carunculatus cafer*. Subsequent to the publication of the Part which contained *B. abyssinicus*, Mr. Keulemans sent me a sketch of the head and neck of a Ground-Hornbill which he had met with in the Zoological Gardens at Amsterdam, that differed from any thing I had ever seen as regarded the shape and coloration of the bare skin of the head and neck, as well as in the form of the casque. The bird was fully adult, and had been received, as I was informed by Mons. A. von Bemmelen, the Director of the Rotterdam Gardens, from the region of the river Congo. Satisfied that the individual represented an unknown species, I described it in the 'Annals and Magazine of Natural History,' as given above. The only form of Ground Hornbill that has received a name with which the present can be compared, is the *B. carunculatus guineensis*, Schlegel. The type of this last-named style is an immature bird; that is to say, the casque is not more than half developed; yet even now the greatly curved form of the upper outline is plainly discernible, showing that when the casque has attained its full growth it will not differ in shape from that of the well-known *B. abyssinicus*. Prof. Schlegel, when writing to me lately about these birds, gave in his letter the comparative differences which he observed in the separate forms named by him; and he thus distinguishes *guineensis*:—"Casque as in *abyssinicus*, but the whole bird smaller, plaque light brown." The difference of size is hardly sufficient to constitute a specific difference; and I therefore regard *B. guineensis*, Schlegel, as properly a synonym of the *B. abyssinicus*. This therefore reduces the number of Ground-Hornbills to three species instead of four, as given by me in the table accompanying the description of the present species in the 'Annals and Magazine of Natural History.' By the kindness of Prof. Barboza du Bocage I am in possession of a sketch of the head of a Ground-Hornbill, received at the Lisbon Museum from Cachen, West Africa, referred by the Professor to *B. guineensis*, Schlegel. The head is unfortunately that of a young bird; but the casque, although already open in front, has its upper outline nearly straight, as is seen in *B. pyrrhops*, and of quite a different form from that exhibited by the type of *B. guineensis*, Schlegel. I am inclined to consider this bird the young of *B. pyrrhops*. The present species differs from all others of this genus in the shape of its casque, which is nearly straight above, with a large opening in front, somewhat resembling a fleur-de-lys in outline. The

colouring and shape of the bare skin on the head and neck are quite different from those of any other species known to me. The type at Rotterdam is the only example yet brought to Europe, so far as I am aware, unless my supposition should prove correct that the *guineensis* of Bocage (nec Schlegel) is the same species. With the slight knowledge yet possessed of the present species, it is impossible to give more than an approximate idea of its habitat. It probably, however, ranges upon the west coast of Africa, between the habitat of *B. abyssinicus* on the north and that of *B. cafer* on the south; but how far it may extend into the interior of the continent I have not at present any means of ascertaining. The type, although it has not been very long in captivity, is very gentle, and is greatly pleased when taken into any person's arms and its head is rubbed. Within the gular pouch, at its lower extremity, a hard round substance can be felt, which is capable of being moved upwards as far as the red skin extends.

Adult.—Bill black, with an orange-coloured plate on the side of the maxilla near the base, as in *B. abyssinicus*. Casque, rising from base of maxilla, extends backwards over the eye, and then inclines forwards (nearly straight on top), terminating in a wide opening in front, similar to that in *B. abyssinicus*, but smaller. Space around the eye, and extending backwards nearly to the occiput, bare of feathers, which, together with that beneath the feathers on the forehead, is orange-red. Bare skin on the sides of the neck and throat dark blue. Gular sac orange-yellow in front, orange-red behind, this latter colour running up for a short distance on the posterior side of the blue skin of the neck. Entire plumage of body lustrous black; primaries pure white; irides pearly white.

Length of bill from the angle of the mouth $6\frac{5}{8}$ inches, height of casque $\frac{3}{4}$, height of bill without casque $1\frac{7}{8}$, tail $11\frac{3}{8}$, tarsus $5\frac{1}{2}$.

J.G. Keulemans lith.

Hanhart imp

BUCORVUS CAFER

BUCORVUS CAFER.

CAFER GROUND-HORNBILL.

Buceros abyssinicus cafer, Schleg. Mus. Pays-Bas, *Buceros*, p. 20.

Bucorvus abyssinicus, Gurn. (nec Gmel.), Anderss. B. Damara-l. p. 205 (1872).

Tmetoceros abyssinicus, Heugl. Orn. N.O.-Afr. p. 731 (1872); Finsch & Hartl. Vög. Ost-Afr. p. 480 (1867).

Bucorax abyssinicus, Boc. Journ. Acad. Sc. Lisb. no. viii. (1870) p. 347.

Bucorax cafer, Boc. Proc. Zool. Soc. (1873) p. 698; id. Journ. Acad. Sc. Lisboa, no. xvi. (1873) p. 284, no. xvii.

(1874) p. 57, no. xix. (1876) p. 149; id. Orn. Angola, p. 111 (1877); Sharpe in Lay. B. S. Afr. p. 122 (1875).

Mucungungo (adult) and *Inaquendi* (young), Natives of the Humbe (ANCHIETA).

HAB. Angola, Quillengues, Humbe (ANCHIETA); Damara Land (ANDERSSON); Caffraria (SCHLEGEL).

This species of Ground-Hornbill apparently represents the genus in South Africa, as its range seems to be from Angola, through Damara Land, into Caffraria. The exact dispersion of the species, or the extent of its range is not as yet thoroughly ascertained, as most writers have usually considered that there was but one form of Ground-Hornbill, and therefore called all the three species recognized in this work, *abyssinicus*, the name usually employed for the high opened-casque bird only.

M. Anchieta, as quoted by Prof. Bocage in his work on the Ornithology of Angola, states that when this species is walking along, the young go behind the adults at a respectful distance and content themselves with seizing whatever may have escaped the voracity of their elders, and therefore the natives believe that they are the slaves of the old ones, are obliged to follow them, and eat whatever they may choose to leave. M. Anchieta also says that they go in small bands and feed upon Coleopterous and other insects. They nest in holes of trees, and lay two eggs.

In Ondonga, according to Andersson, this species was common, but very wild; and he had also observed it in the desert near Okavango. They run very rapidly, but perch on trees when flushed. In fact in its habits and economy this bird does not appear to differ in any way from its relatives of the same genus in the northern part of the continent.

In appearance the Cafer Ground-Hornbill differs from the other forms of *Bucorvus* mainly in the absence of the plaque at the base of the maxilla, and in the shape of the casque, which at all ages is compressed and closed in front.

The general plumage is black with bronze reflections, the primaries white. Bare skin around the eyes, throat, and gular pouch yellowish orange inclining somewhat to red, and occasionally dotted with spots of blackish blue. Bill and casque black. Feet black. Iris pale greenish yellow.

Wing 19 to 21 inches, tail 11, tarsus $4\frac{1}{2}$, bill 7.

In the young the general plumage is a reddish brown, and the primaries have a few brown spots near their tips; bill short and whitish; casque very small and greatly compressed, black; bare skin around eyes and on the throat greenish grey. The gular pouch is wanting.

Subfamily BUCEROTINÆ.

Bill generally powerful, covered with a casque or casque-like protuberance. Nude skin nearly always present around the eyes, occasionally also at base of bill and on the throat, but usually limited in extent. One species only has the face and neck bare. Wings broad and ample. Tail generally long and rounded. Legs and feet short, only fitted for perching, the species being incapable of walking, but progressing over the ground by awkward hops. The members of this subfamily are tree-dwellers, and only occasionally leave their abode among the branches to descend to the earth.

M. & N. H. Hart 1897

BUCEROTUS RHINOCEROS

J.G. Keulemans del.

BUCEROS RHINOCEROS.

RHINOCEROS HORNBILL.

Buceros rhinoceros, Linn. Syst. Nat. (1766) vol. i. p. 153; Gmel. Syst. Nat. (1788) vol. i. p. 360; Lath. Ind. Ornith. (1790) vol. i. p. 141; Shaw, Gen. Zool. (1811) vol. viii. p. 3, pl. 1; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 598; Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 198; Temm. Planch. Col. (1824) vol. ii. sp. 1; Ersch u. Grub. Ency. (1824) p. 287; Wagl. Syst. Av. (1827); Griff. Anim. King. (1829) vol. ii. p. 432; Less. Trait. Ornith. (1831) p. 255; Begbie, Malay Penins. (1834) p. 512; Müll. & Schleg. Verhand. Natur. Geschied. Nederl. Ind. (1839-44) pp. 21, 26; G. R. Gray, Gen. Birds (1849) vol. ii. p. 349. sp. 1; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 42; Bon. Consp. Gen. Av. (1850) p. 89. sp. 3; id. Consp. Voluc. Anisod. (1854) p. 184; Horsf. and Moore, Cat. Birds Mus. E.-Ind. Co. (1856) vol. ii. p. 582. no. 867; Moore, Proc. Zool. Soc. (1859) p. 446; Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 174; Schleg. Mus. Pays-Bas (1862) p. 3; Blyth, Ibis (1866) p. 352; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 127. sp. 7866; Sclat. Wolf's Zool. Sketch. 2nd Ser. pl. xxx.; Giebel, Thes. Ornith. (1872) vol. i. p. 502; Garrod, Proc. Zool. Soc. (1873) p. 464, (1875) p. 342; Gulliv. Proc. Zool. Soc. (1875) p. 490.

Rhinoceros hornbill, Lath. Gen. Syn. (1781) p. 342. sp. 1, vol. i.

L'oiseau rhinocéros, Buff. Planch. Enlum. (1783) no. 934 (bill).

Buceros africanus, Gmel. Syst. Nat. (1788) vol. i. p. 357. sp. 6; Shaw, Gen. Zool. (1811) vol. viii. p. 6.

Le calao rhinocéros, Le Vaill. Ois. d'Amér. (1801) pls. i., ii. (bill).

Buceros rhinoceros, var. *a. sumatranus*, Müll. & Schleg. Verh. Geschied. Neder. Ind. (1839-44) p. 22.

Buceros rhinoceros, var. *b. bornensis*, Müll. & Schleg. Verh. Geschied. Neder. Ind. (1839-44) p. 22.

Buceros rhinoceros, var. *c. indica*, Müll. & Schleg. Verh. Geschied. Neder. Ind. (1839-44) p. 22.

Buceros rhinoceros, var. *d. javanica*, Müll. & Schleg. Verh. Geschied. Neder. Ind. (1839-44) p. 22.

Buceros rhinoceroïdes, Temm. Mus. Lugd.; Bon. Consp. Gen. Av. (1850) p. 89. sp. 4; id. Consp. Vol. Anisod. (1854) p. 2 (ex Borneo).

Buceros sublunata, Temm. Mus. Lugd.; Bon. Consp. Gen. Av. (1850) p. 90. sp. 5; id. Consp. Vol. Anisod. (1854) p. 2; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 127. sp. 7867.

Buceros rhinoceros sumatranus, Schleg. Mus. Pays-B. (1862) p. 4.

Buceros rhinoceros bornensis, Schleg. Mus. Pays-B. (1862) p. 4.

HAB. Malay peninsula (BLYTH); Sumatra, Borneo (WALLACE, SCHLEGEL).

The Rhinoceros Hornbill is pretty generally distributed throughout the Malayan peninsula, and is also found in Sumatra and Borneo. The upturned casque renders it very conspicuous, and causes the bird to differ from all the members of the family—so much so that I have retained it, with its near relative from Java, as the representatives of Linnæus's genus *Buceros*, and restricted that term in this work to this form. Schlegel has separated the birds from Sumatra and Borneo as races or *conspicies* of the Malayan species, one of the characters being the width of the black band on the tail. I find that this band varies very much in different specimens, even from the

same locality, and does not appear to be of any specific importance. Thus, of two fine male examples in the Leyden Museum from Batang, Sumatra, named by Schlegel *B. rhinoceros sumatranus*, one has a quite broad band, the other a comparatively narrow one, while the shape and size of bills are equal. I have therefore placed Prof. Schlegel's names as synonyms, as I do not consider there are more than two species of this form, viz. the present and *B. sylvestris*, Vieill.

According to Wallace, the exertion this Hornbill is compelled to make when flying is so great that it is obliged to rest at intervals of about a mile. It feeds on fruits, and requires so much to appease its appetite that a tree is soon stripped of all that is within reach; for, on account of its weight, the bird is only able to take those that are growing close to the larger branches. It moves along these sideways, by a kind of shuffling hop, and seizes the fruit with the point of its long bill. When it wishes to swallow any thing it may have obtained, the morsel is jerked upwards into the air, the head held back, and the bill opened wide; and the food is thus cast into the throat. Except in localities where it has not been molested, this species is very shy, and the appearance of a man, even at a considerable distance, will cause it to take flight at once. The same fruit-tree is rarely visited more than two or three days consecutively; and in some districts it is probably a matter of no slight difficulty for this large bird to provide itself with a meal every day.

The male *B. rhinoceros* has the bill straw-yellow, a triangular space at the lower base of maxilla red, and one on the mandible black. A large casque, broad at base, sharp at top, extends from the centre of the head to about halfway on the culmen, where it turns abruptly upwards, in old individuals curving slightly backwards at the tip; this is black posteriorly, and also on the keel of the curved portion anteriorly. A narrow black line also runs along the side from above the eye to the upper anterior part of the casque, which, with the exceptions just given, is of a general red colour, becoming yellowish along the anterior edges. The head, which is slightly crested, the neck, back, wings, and breast, as far as the thighs, are of a uniform bluish black. Thighs, lower part of abdomen, crissum, rump, upper and under tail-coverts pure white. Tail pure white, with a black band crossing all the feathers, distant about one third from the tips. Iris yellowish white, surrounded by black; eyelids red. Feet and tarsi brownish black.

Length of bill $9\frac{1}{4}$ inches, casque $6\frac{1}{4}$, height of casque $2\frac{1}{2}$; total length $39\frac{1}{2}$, wing $18\frac{1}{4}$, tail 17.

The plumage of the female is like that of the male; but the back part of the casque is reddish like the sides, and it is also without the narrow black line running from above the eye.

J.G. Keulemans lith.

Hanhart imp.

BUCEROS SYLVESTRIS.

BUCEROS SILVESTRIS.

JAVAN RHINOCEROS HORNBILL.

Le Calao à casque en croissant, Levaill. Ois. d'Amér. (1801) pl. 13.

Buceros semilunaris, Wilkes ? Ency. Lond. (1808) vol. iii. p. 479.

Crescent Hornbill, Shaw, Gen. Zool. (1812) vol. viii. p. 7.

Buceros silvestris, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 592 ; Less. Trait. Ornith. (1831) p. 256. sp. 14.

Buceros diadematus, Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 203.

Buceros niger, Ersch u. Grub. Ency. (1824) p. 286.

Buceros lunatus, Temm. Plan. Col. (1830) no. 546 ; Bon. Consp. Gen. Av. (1850) p. 90. sp. 6 ; id. Consp. Vol. Anisod. (1854) p. 2 ; Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 175, no. 480 ; Gray, Hand-l. B. (1870) pt. ii. p. 127. sp. 7869.

Buceros rhinoceros, var. *d. javanica* *, Müll. & Schleg. Verh. Geschied. Neder. Ind. (1839-44) p. 22.

Buceros rhinoceros lunatus, Schleg. Mus. Pays-B. (1862) p. 5.

HAB. Java.

This bird, possessing a very differently shaped casque from that of *B. rhinoceros*, and being restricted to one island, where it is the only representative of the genus, may perhaps, with some propriety, be regarded as a distinct species. It was named as long ago as 1808, in the 'Encyclopædia Londinensis,' from plate 13 of Levaillant's 'Oiseaux Rares de l'Amérique et de l'Inde,' and called *B. semilunaris*, but, unfortunately, without any author's name attached ; and therefore the appellation bestowed by Vieillot (*l. c.*) in 1816 is the one that has priority over all others. Temminck, as in many other instances, disregarded the names already given to the bird, or was ignorant of them, and redescribed it, in the 'Planches Coloriées,' as *B. lunatus* ; and many ornithologists have adopted this appellation. It will, however, be obliged to give way and become a synonym of the one given by Vieillot. We know very little of this bird in its native haunts. It has at various times been an inmate of different zoological gardens, but is much more seldom seen, even in collections, than the species from Sumatra and the Malayan peninsula.

The *male* has the bill white, with the base of both maxilla and mandible black, edged with deep red anteriorly. The casque, in the shape of which the main difference between this species and *B. rhinoceros* lies, is straight, without any upward turn at the point, rising from the posterior part of the culmen just above the eyes, and extending nearly two thirds the length of the maxilla :

* This synonym was inadvertently published with those of *B. rhinoceros*. It properly belongs to this species only.

general colour red, becoming deep yellow towards the point; posterior terminus black, as is also a narrow line running along the culmen. The plumage is the same as that of *B. rhinoceros*, being bluish black, with white abdomen, thighs, and upper and lower tail-coverts. The tail is also white, with a broad black band crossing it about two thirds from the base. Iris olive.

Length of bill along gape 8 inches, of casque $5\frac{1}{2}$, height of casque 2. Total length 3 feet, wing $17\frac{1}{2}$ inches, tail $15\frac{1}{4}$, tarsus 2.

The *female* resembles the male, except that there is no black upon the bill or casque, the posterior terminus of the latter being red.

My description and figure were taken from specimens in the British-Museum collection.

black
white
red
blue
green
yellow
orange
purple
brown
pink
gray
tan
olive
teal
lavender
mint
coral
turquoise
indigo
violet
cyan
magenta
fuchsia
amaranth
crimson
scarlet
ruby
emerald
sapphire
diamond
opal
pearl
shell
stone
metal
wood
glass
leather
fabric
paper
ink
oil
water
air
fire
earth
sky
sea
land
mountain
river
lake
forest
field
city
village
town
country
world

Hartwegt amp.

JG Keulemans lith.

DICHOCEROS BICORNIS.

THE HOMRAI.

Buceros bicornis, Linn. Syst. Nat. (1766) vol. i. p. 153; Gmel. Syst. Nat. (1788) vol. i. p. 358; Lath. Ind. Ornith. (1790) vol. i. p. 142; Shaw, Gen. Zool. (1811) vol. viii. p. 16; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 597; Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 198; Vieill. Ency. Méth. (1823) p. 301; Temm. Plan. Col. (1824) vol. ii. sp. 4 (text); Ersch u. Grub. Ency. (1824) p. 283; Wagl. Syst. Av. (1827) sp. 5; Less. Trait. Ornith. (1831) p. 254. sp. 9; Müll. & Schleg. Verh. Geschied. Nederl. Ind. (1839-44) p. 27. sp. 11; G. R. Gray, Gen. Birds (1849) vol. ii. p. 349; Bon. Consp. Gen. Av. (1850) p. 89. sp. 2; Hartl. Journ. für Ornith. (1859) p. 288. sp. 28; Baker, Journ. Asiat. Soc. Beng. (1859) p. 292; Schleg. Mus. Pays-B. (1862) p. 3; Sclat. Wolf's Zool. Sketch. 2nd Ser. pl. xxix.; G. R. Gray, Hand-l. B. pt. ii. (1870) p. 127. sp. 7870; Blanf. Ibis (1870) p. 466. sp. 33; Giebel, Thes. Ornith. (1872) vol. i. p. 496; Garrod, Proc. Zool. Soc. (1873) p. 464.

Le Calao à casque concave, Le Vaill. Ois. d'Amér. (1801) pls. 3, 4 (heads), and 5.

Le Calao bicolore, Le Vaill. Ois. d'Amér. pls. 7, 8 (bill), (1801).

Buceros cavatus, Shaw, Gen. Zool. (1811) vol. viii. p. 18; Raffles, Trans. Linn. Soc. vol. xiii. (1822) p. 291; Ersch u. Grub. Ency. (1824) p. 285; Vig. Appen. Mem. Raffl. (1830) p. 666; Jerd. Madr. Journ. (1840) p. 37; Blyth, Journ. Asiat. Soc. Beng. (1842) vol. xii. pt. ii. p. 986; Gould, Cent. Birds, pl.; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 42; Blanf. Ibis (1870) p. 466. sp. 32.

Buceros cristatus, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 591; id. Ency. Méth. (1823) p. 399.

Buceros homrai, Hodg. Journ. Asiat. Soc. Beng. (1832) p. 251; id. Asiat. Research. (1833) p. 102; J. E. Gray, Zool. Misc. p. 85; G. R. Gray, Gen. Birds (1849) vol. ii. p. 349. sp. 3; id. Hand-l. Birds (1870) pt. ii. p. 127. sp. 7871.

Dichoceros cavatus, Glog. Hand u. Hilfs. Naturgesch. (1842) p. 335; Hume, Str. Feath. vol. v. (1875) p. 20; Bourdillon, Str. Feath. (1876) p. 384.

Homraius bicornis, Bon. Consp. Vol. Anisod. p. 2, gen. 7. sp. 17 (1854); Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 583. no. 868; Moore, Proc. Zool. Soc. (1859) p. 447; Jerd. B. Ind. (1862) vol. i. p. 242; Blyth, Ibis (1866) p. 349; Jerd. Ibis (1872) p. 4.

Dichoceros homrai, Hume, Stray Feath. vol. ii. p. 470 (1874), and vol. iii. (1875) p. 55.

Dichoceros bicornis, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 173. no. 477; Hume, Stray Feath. vol. ii. (1874) p. 470; Blyth, Cat. Mamm. & Birds Burma, Journ. Asiat. Soc. Beng. (1875) p. 68. sp. 68.

Homrai in Nepaul, *Ban-rao* at Mussoree (JERDON). *Ingghen Papan* of the Sumatrans (RAFFLES). *Kugrong* of the Lepchas, *Gogrong* of the Bhotecas (JERDON). *Burong Oondan* of the Malays (RAFFLES).

HAB. Indo-Chinese countries, Malayan peninsula and Sumatra (BLYTH, SCHLEGEL).

This Hornbill was described by Linnæus (*l. c.*), who cited among the first of his synonyms Willughby's 'Ornithology,' t. 17. fig. 1. This author correctly figures the head and bill of the present species, with the two sides of the casque turned slightly upwards at their anterior termina-

tion. Linnæus also refers to Brisson's *Hydrocorax philippensis*, 'Ornith.' vol. iv. p. 568. Of this bird Brisson expressly states that he had only seen the *head and the bill*, which undoubtedly, from his description, belonged to the bird represented on the opposite plate. Where he got his idea of the plumage of the body is not stated; but the head was in the cabinet of M. de Réaumur. Linnæus, in his description of the plumage, followed Brisson in a great measure, and evidently confused two species together; for the tail of his bird belongs either to the *A. coronatus* or *A. convexus*; it is impossible to say which. As, however, both Willughby and Brisson, whom Linnæus quotes, accurately describe the head of the present species, the only portion they were acquainted with, and the first-named also figures it, there can be no doubt as to which bird Linnæus applied the term *bicornis*; and his diagnosis, "*B. fronte ossea plana antrorsum bicorni*," can only be intended for the species now under consideration, as there is no other Hornbill known for which this description would answer. There is a great difference observable in the form of the casques of individuals, the anterior margin in some assuming the shape of a semicircle, while in others it is almost flat. These peculiarities are found to exist among specimens from all localities, and do not indicate separate species. The Sikhim birds, as stated by Mr. Hume, are apparently somewhat larger than those from Northern Tenasserim; but some from South Tenasserim equal in size the Sikhim ones, and there is nothing to indicate two species.

I have been unable to distinguish any differences between the Indian and Malayan forms of this bird, sufficient to constitute them distinct, and have therefore placed Mr. Hodgson's *B. homrai* among the synonyms of *D. bicornis*.

This fine species, with one exception the largest of the family, has quite a wide range. Indeed, how extensive this is we do not exactly know, for it may be an inmate of the great forests that stretch away into China. Being met with generally throughout India, its habits have been observed by many of the excellent naturalists of that country, and consequently we are pretty familiar with its mode of life.

It is a forest-loving bird, and is usually found on the mountain-ranges at a height of from 3000 to 5000 feet. According to Hodgson, it is also met with upon the lower ranges of hills near the plains, preferring the open and cultivated spots in the wilds it inhabits, which places are usually restricted to the banks of the rivers. It is gregarious, twenty or thirty individuals, as stated by Hodgson, being commonly found in the same vicinity, six or seven on one tree, although Jerdon says he never saw more than five or six together, either in Southern India or the Sikhim Himalayas, and there only rarely; while Mr. Bourdillon states that three are the greatest number he has seen at one time in the Travancore hills; and Mr. Oates gives five or six as the number in a flock in the Evergreen Forests in Upper Pegu, where it is a common bird, but extremely wary and difficult to approach, keeping in the highest branches. They will remain perched for hours, almost motionless, uttering at intervals a low croak as though conversing with each other.

Usually it is rather a silent bird, but when wounded or taken captive, it utters the most extraordinary sounds imaginable. The voice is very harsh and grating, and the noise it makes is compared by Wallace to something between the bray of a jackass and the shriek of a locomotive, and is not to be surpassed probably, in power, by any sound that an animal is capable of making.

Tickell says that its roar reechoes through the hills to such a degree that it is difficult to assign the noise to a bird, and Wallace states that this is kept up so continuously as to be absolutely unbearable. He heard it plainly a mile away. The flight of this species is heavy, and performed by repeated flappings of its huge wings. It usually proceeds in a straight line, and sails only when about to alight upon some tree. The strokes are made with great force, and the noise of its wings can be heard for more than a mile. As a rule the food of this Hornbill is strictly fruits—certainly so, says Hodgson, at certain seasons, as in the months of January and February, when he found the stomachs contained nothing but the fruit of the Pipal tree. Tickell states that it eats lizards readily, not only from the hand, but will search for them and seize them. With this exception, authors generally agree in regarding fruit as the sole food of this bird.

It breeds in the holes of large trees, and, like the other members of this Family, the male plasters the female in, and never allows her to leave the nest until incubation is accomplished, keeping her well supplied with food in the meanwhile. The egg, as described by Tickell, is a dirty yellowish brownish white, spindle-shaped or pointed at both ends, and of a coarse surface, indented with numerous pores. Longitudinal and transverse axis $2\frac{1}{8}$ " and $1\frac{7}{8}$ " respectively.

"The newly hatched young," says Wallace, "is as large as a pigeon, destitute of plumage, exceedingly plump and soft, with a semitransparent skin, so that it looks more like a bag of jelly with head and feet stuck on than a real bird. Frequently the wing-spots, rump, nuchal crest, and parts of the bill are stained yellow. This colour comes from a bundle of stiff feathers protruding from the sac at the root of the tail, which exudes an oily secretion, with which the bird dresses its plumage." "When first shot," says Blyth, "this colour comes off the *bill* in considerable quantities, and the yellow substance continues to exude from the brush long after the specimen is prepared and dry." Mr. Oates states, in 'Stray Feathers' (1875) p. 55, in reference to this colouring-matter on the plumage, that he does not think the "yellow on the head and neck is entirely due to the secretion of the uropygial gland. It does not come off in any quantity when the bird is killed." Mr. Inglis, who met with this species in Cachar, states that in the dry weather they are continually migrating to the south, and, during the rains, to the north. They go mostly in flocks of five and seven, but sometimes thirty are seen together. A windy day is the best for shooting them, and they afford "splendid eating, far superior to any fowl or pheasant." Among the inhabitants of the forest, this bird is stated to be sacred to Vishnu, and the Nepaulese name for this species, "Homrai," is derived from the notes it is accustomed to utter.

Male.—Bill much curved. Maxilla yellow, reddish towards the tip; mandible ivory-white in adults, the base of both black. Casque very broad, flat, extending backwards over the head for about half the length of the bill, and descending to the maxilla at a right angle. Posterior termination black, its upper edge reddish; anterior edge black, this colour reaching to the bill and continuing along the culmen to the point. Rest of casque waxen-yellow. Cutting-edges of bill also black. Head and base of bill black, as are also the back, breast, and abdomen. Occiput, neck, thighs and tips of upper tail-coverts, crissum, and under tail-coverts white. Wings black, with the ends of the greater coverts, secondaries, primaries, and a spot on these last, about halfway

towards the shoulders, white. Tail pure white, with a broad black band about two thirds from the base. Sometimes the neck and the white on the wing are yellowish, caused by the secretion from the uropygial gland. Iris crimson; bare skin around eyes black.

Length of bill 10 inches, from point to end of casque 13, of casque $7\frac{1}{2}$, breadth 3, height of bill and casque at base 4. Total length of bird 4 feet, wing 20 inches, tail 17, tarsus $2\frac{1}{2}$.

The female resembles the male, except that there is no black on the casque, and its posterior end is red. Bare skin round the eyes red.

The specimens here described and figured were living in the Regent's-Park Gardens, London.

HYDROCORAX PLANICORNIS.

FLAT-CASQUE HORNBILL.

Buceros hydrocorax, Linn. Syst. Nat. (1766) vol. i. p. 153; Gmel. Syst. Nat. (1788) vol. i. p. 359; Lath. Ind. Ornith. (1790) vol. i. p. 144; id. Gen. Syn. (1790) vol. i. p. 351. sp. 7, 2nd Suppl. (1801) p. 100; Shaw, Gen. Zool. (1811) vol. viii. p. 31; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 597; id. Ency. Méth. (1823) p. 304; Temm. Planch. Col. (1824) vol. ii. sp. 6 (text); Wagl. Syst. Av. (1827) sp. 7; Less. Trait. Ornith. (1831) p. 254. sp. 10; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399. sp. 15; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 43. sp. 176; Bon. Consp. Gen. Av. (1850) p. 89. sp. 1; Schleg. Mus. Pays-B. (1862) p. 2; Von Mart. Journ. für Ornith. (1866) p. 18; Giebel, Thesaur. Ornith. (1872) p. 499; Wald. Trans. Zool. Soc. (1873) vol. ix. p. 164.

Calao des Moluques, Buff. Plan. Enlum. (1783) no. 283.

Buceros planicornis, Merrem, Ersch u. Grub. Ency. (1824) p. 287; Bon. Consp. Vol. Anisod. (1854) p. 184.

Buceros platyrhynchus, Pears. Jour. Asiat. Soc. Beng. (1841) p. 652.

Hydrocorax planicornis, Bonap. Consp. Vol. Anisod. (1854) p. 2. gen. 8. sp. 18.

Platyceros hydrocorax, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 174. no. 478.

Buceros (Hydrocorax) hydrocorax, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7884.

HAB. Moluccas (BLYTH); Philippines (SCHLEGEL).

This remarkable species was one of the few described by Linnæus in his 'Systema Naturæ,' and is a native of the Moluccas and also the Philippine Islands. In the shape of its casque it is nearest allied to *D. bicornis*, but differs so materially both from that species and all others of the family, that it does not seem inappropriate to retain it in a separate genus by itself. Brisson, in his 'Ornithologie' (1760) vol. iv. p. 565, instituted his genus *Hydrocorax* for this species; Cabanis and Heine, in 1860, also proposed the generic term *Platyceros*. Although Brisson is not deemed an authority for species, his genera have usually been employed in ornithology, and I have therefore retained his term to the exclusion of the later one by Cabanis. But little is known of the habits and economy of this bird, its habitat being one not often penetrated by Europeans, even Mr. Wallace, the most eminent of the naturalists who have visited portions of the country in which it is found, having had but little to relate concerning it. This is perhaps more to be wondered at as it must be a very conspicuous object in its native woods, and it could not move from one tree to another without betraying itself by the noise of its wings, which has been compared to the puffing of a locomotive just starting with a train.

Bill brick-red; a broad flat casque extends from about one third the length of the maxilla from its point, along the culmen, increasing in width on its upper surface, and projects over the head to behind the eyes; this, like the bill, is brick-red, darkest on its sides. Front, cheeks, and chin black. Head, neck, and upper part of breast chestnut-red. Lower part of the throat white.

Back and wings dark brown, edges of secondaries buff, primaries black. Flanks and abdomen black. Thighs and under tail-coverts rufous. Tail white. No difference between the sexes. Length of bill of male $6\frac{1}{2}$ inches, of casque $6\frac{1}{2}$, height of casque $3\frac{1}{2}$. Total length of bird $29\frac{3}{4}$ inches; wing 17; tail $14\frac{1}{2}$; tarsus $2\frac{1}{4}$.

Young have the casque and bill black. Head, neck, and underparts dirty white. Back and wings dark brown, scapulars tipped with white, also secondaries edged with buff. Tail white, the outer rectrices with a narrow bar of black on their terminal third.

The figures in the Plate represent the adult and young.

JG. Keulemans hth.

Hanbart imp.

HYDROCORAX MINDANENSIS.

HYDROCORAX MINDANENSIS.

MINDANAO FLAT-CASQUE HORNBILL.

Buceros mindanensis, Lord Tweeddale, Proc. Zool. Soc. (1877) pp. 543, 823, (1878) pp. 278, 946.

HAB. Island of Mindanao, Valley of Pasananca, Zamboanga (EVERETT).

This apparently distinct species was procured by the officers of the 'Challenger' in the valley of Pasananca, island of Mindanao, about seven miles from Zamboanga, and described by the late Marquis of Tweeddale (*l. c.*). Three examples were obtained—a male, a female, and a young female. Mr. Murray, the naturalist of the expedition, furnished the following notes regarding them:—"The stomachs of all contained seeds and fruits, and grubs, centipedes, grasshoppers, &c. These birds make a loud sound, like a crow somewhat, and frequent the highest trees. Several times in the early morning we came upon them on the ground under the trees; and I rather think they scrape at the roots of trees for food."

Though closely allied to the *H. planicornis*, the present species is distinguished by having a corrugated plate on the sides of the mandible at its base, by the casque being narrower, by the half of the maxilla and three fourths of the mandible anteriorly being white, and by the rufous of the plumage being darker. The sexes do not differ in the colour of their plumage.

Adult.—Casque, basal half of maxilla, and crura of mandible deep red. The corrugated plates on mandible are a darker red; rest of bill pale yellowish white. A black band surrounds the eye, extends down the face, and covers the chin. Upper part of throat tawny. Breast black. Rest of plumage of head and body, except tail-coverts, deep rufous. Interscapulars and wing-coverts brown. Upper tail-coverts ferruginous; under coverts rust-colour. Primaries black; secondaries margined on the outer web with ferruginous. Rectrices buff. The colour of the iris seems to vary from bright yellow to a pale green in examples from different parts of the island. Orbital skin blackish, with a yellow streak under the eye. Bare skin of throat yellow. Feet and tarsi coral-red.

Wing $15\frac{1}{2}$ inches; tail 14; bill along gape 6.62; length of casque 6, breadth 2; tarsus 3; middle toe 1.60.

Immature Bird.—Has the casque not so fully developed anteriorly, and not overhanging the true culmen. This latter is grey-brown, and the top of the casque red, as is also the base of mandible. The grooved lateral plates have not appeared; but there are indications of grooves on the mandible at base. Feet and tarsi red. Bill blackish, tinged with red. Irides blue.

Length, wing $14\frac{1}{4}$ inches, tail $13\frac{1}{2}$, bill along gape 5.62, casque $4\frac{1}{2}$, breadth of same $1\frac{1}{2}$, tarsus 3, middle toe $1\frac{3}{4}$.

Female has the iris white, feet light coral-red.

J. G. Keulemans. lith.

Hanhart imp.

HYDROCORAX SEMIGALEATUS.

HYDROCORAX SEMIGALEATUS.

SEMI-HELMETED HORNBILL.

Buceros semigaleatus, Tweeddale, Proc. Zool. Soc. (1878) p. 279.

HAB. Ampavo, island of Leyte, and Panaon, Philippines (EVERETT).

This local form of *H. planicornis* was described by the late Marquis of Tweeddale (*l. c.*) from specimens procured by Mr. Everett at Ampavo, in the extreme south of the island of Leyte, and on Panaon, both belonging to the Philippine archipelago. In the colouring of the bill and plumage this bird presents no difference from *H. mindanensis*; and its claim for specific distinction rests solely on the shape of the casque. This, instead of being flat upon its superior plane and terminating in a compressed and elevated anterior margin at a right angle to the culmen, like those of *H. planicornis* and *H. mindanensis*, has this portion arched, and the crown of this arch upon its anterior half forming an acute ridge, while the casque itself falls away anteriorly, and is lost in the culmen. In the two species mentioned above, the contour of the plane of the casque is oval, with the posterior end rounded, and the anterior pointed. In this bird the posterior end is rounded, and the sides are parallel for two thirds the length of the casque, where they form corners, and then recede rapidly to the culmen. Nothing was recorded regarding the economy or habits of this peculiar form, though several specimens were procured, representing the adult and immature birds. The plumage of the old individuals was the same as that of *H. mindanensis*, with which this form is most closely allied. The iris appears to vary in colour, that of a male from Leyte being light yellow, while one from Panaon was pale blue. Orbital skin almost black; gular skin indian-yellow; feet coral-red, nails dark brown-grey. At the base of the mandibles of adult examples there are slightly corrugated plates.

Male.—Length of wing 15·80 inches, tail 15, tarsus 2·20, bill at gape 5·85; length of casque 4·90, breadth 2, sides 3·35.

Female.—Length of wing 14·30 inches, tail 14·50, tarsus 2·20, bill at gape 5·30; length of casque 4·75, breadth 1·65, sides 3·30.

The young in first plumage have the head, neck, breast, abdomen, thighs, and vent greyish fulvous-white, the feathers being ferruginous at their insertions, and grey at their tips. Upper tail-coverts also same colour, with brown bases. Back and wings brown; quills dark brown with greyish fulvous edgings. Median pair of rectrices pale brown for two thirds their length, with the apical third creamy white; remaining rectrices pale rusty-brown. Bill black, with the tip yellow. Iris dark brown; orbital skin greenish yellow; gular skin yellow; legs and feet dark orange.

The figures in the Plate are taken from the type specimens in Captain Ramsay's possession, to whom I am indebted for the opportunity of introducing them into this work.

J. G. Keulemans lith.

RHINOPLAX VIGIL.

M & N. Hanhart imp

RHINOPLAX VIGIL.

HELMETED HORNBILL.

Buceros vigil, J. R. Forster, Zoolog. Indica (1781) p. 40.

Helmeted Hornbill, Lath. Gen. Syn. (1781) p. 343. sp. 2.

Buceros scutatus, Bodd. Tab. Pl. Enlum. d'Aubent. (1783) sp. 933, p. 55; G. R. Gray, Gen. Birds (1846) vol. ii. p. 399. sp. 17; Giebel, Thesaur. Ornith. (1872) p. 503.

Calao à casque rond, Buff. Pl. Enlum. (1783) no. 933 (head).

Buceros galeatus, Gmel. ed. Linn. Syst. Nat. (1788) vol. i. p. 360; Lath. Ind. Ornith. (1790) vol. i. p. 142; Shaw, Gen. Zool. (1811) vol. viii. p. 24; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 593; Dumont, Dict. Sc. Nat. (1817) p. 204; Vieill. Enc. Méth. (1823) p. 301; Temm. Pl. Col. (1824) vol. ii. sp. 7 (text); Ersch u. Gruber, Enc. (1824) p. 284; Wagl. Syst. Av. (1827) sp. 4; Less. Trait. Ornith. (1831) p. 256. sp. 18; Begbie, Malay Penin. (1834) p. 513; Blyth, Cat. B. Mus. Asiat. Soc. (1849) p. 45. sp. 192; Schleg. Mus. Pays-Bas (1862) p. 1. sp. 1.

Rhinoplax scutatus, Glog. Hand- und Hilfsb. (1842) p. 335; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) vol. ii. p. 581. no. 866; Moore, Proc. Zool. Soc. (1859) p. 446.

Buceroturus galeatus, Bon. Consp. Gen. Av. (1850) p. 89. sp. 1; id. Consp. Vol. Anisod. (1854) p. 2.

Buceros (Rhinoplax) scutatus, G. R. Gray, Hand-l. Birds (1870) part ii. p. 130. sp. 7908.

HAB. Malayan peninsula and archipelago (BLYTH); Borneo, Sumatra (SCHLEGEL).

Forster, in his 'Zoologia Indica' (1781), gave the name of *vigil* to this bird, founded upon the bill figured in Edwards's 'Birds,' plate 281c, and thus antedated Boddaert by two years.

This extraordinary bird is remarkable even among the unusual forms met with amid the members of the Bucerotidæ. Not only does it possess a large bill and casque, but, unlike all the other species, this latter structure, instead of being hollow, is perfectly solid. This fact causes it to be quite heavy; and a blow from the bill of a full-grown male upon a man's head might easily produce very serious, if not fatal, results. For what especial reason this bird is endowed with the solid casque has not yet been ascertained: but it is probably for detaching bark or fruit from the trees; for it would have nearly as great an effect when struck against any object as a blow from a hammer in the hands of a man. Another peculiar character possessed alone by this species among the Hornbills is the great length of the two median rectrices, which extend for a considerable distance beyond the rest of the feathers of the tail. With its bony casque, bare neck, and lengthened tail it seems fairly to be entitled to a distinct generic rank; and I have therefore retained for it the term *Rhinoplax*, first proposed by Gloger in 1842. Nothing is known of its economy or habits.

Male.—Bill straight, pointed, yellow, brick-red at base. About midway on the culmen rises a solid bony casque, orange-red on its anterior half, brick-red for the remainder, wide on its anterior face, curved on top, to which it narrows from the sides, but is nowhere compressed,

extends over the head to the eyes, and returns to the base of the bill by a moderate curve. Top of head and occiput black, a patch of feathers on the sides edged with rufous. Neck and spot on back bare, brick-red. Back, breast, and wings brownish black; primaries and one or two of the secondaries tipped with white; the median tail-feathers are double the length of the rest, and, like them, are white, with a black bar near the tip. Iris indian-red. Legs and feet dark red. Length of bill $5\frac{1}{2}$ inches, casque $3\frac{1}{2}$; height of bill and casque at base $4\frac{1}{2}$. Total length of bird 3 feet, to end of median tail-feathers 3 feet 6 inches; wing 19 inches; tail $15\frac{1}{2}$; tarsus $2\frac{1}{2}$.

ANTHRACOCEROS CORONATUS.

CROWNED HORNBILL.

Calao de la côte Malabar, Sonn. Voy. aux Indes (1782) p. 215, pl. 121.

Calao des Philippines, Buff. Plan. Enlum. (1783) no. 873.

Buceros coronatus, Bodd. Tab. Plan. Enlum. D'Auben. (1783) p. 53; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399; Hartl. Journ. für Ornith. (1859) p. 289. sp. 29; Schleg. Mus. Pays-B. (1862) p. 6; Blyth, Ibis (1866) p. 352; Nitz. Pteryl. Ray Soc. (1867) p. 102; Giebel, Thes. Ornith. (1872) p. 497; Garrod, Proc. Zool. Soc. (1872) p. 464; Sclat. Rev. List Vert. Anim. (1872) p. 171. sp. 274.

Buceros pica, Scop. Faun. et Flor. Insub. (1786) p. 87. sp. 32; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 43. sp. 177; Tickell, Ibis (1864) p. 179; Eyton, Osteol. Av. p. 62 (1867).

Buceros malabaricus, var. β & γ , Lath. Ind. Orn. (1790) vol. i. p. 144. sp. 6; Dumont, Dict. Sc. Nat. (1817) p. 205.

Buceros dschindschicus, Lath. & Davies, Faunula Indica (1795) p. 6. no. 10.

Le Calao unicorne, Le Vaill. Ois. d'Amér. (1801) pls. 9, 10 (bill), & 11.

Le Calao violet, Le Vaill. Ois. d'Amér. (1801) pl. 19.

Buceros violaceus, Wilkes? Ency. Lond. (1808) vol. iii. p. 479; Shaw, Gen. Zool. (1811) vol. viii. p. 19; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 599; Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 211; Wagl. Syst. Av. (1827) sp. 9; Temm. Plan. Col. vol. ii. sp. 9 (text); G. R. Gray, Gen. Birds (1849) p. 399; Blyth, Journ. Asiat. Soc. Beng. (1849) p. 803; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7878.

Buceros monoceros, Shaw, Gen. Zool. (1811) vol. viii. p. 8; Ersch u. Grub. Ency. (1824) p. 285; Wagl. Syst. Av. (1827) sp. 2; Begbie, Malay Penin. (1834) p. 513; Temm. Plan. Col. (1838) vol. ii. sp. 11 (text); Müll. & Schleg. Verh. Gesch. Ned. Ind. (1839-44) p. 22. sp. 111.

Buceros malabaricus, Tickell, Journ. Asiat. Soc. Beng. (1833) vol. ii. p. 579; Jerd. Madr. Journ. (1840) p. 38; Blyth, Journ. Asiat. Soc. Beng. (1843) p. 993; Layard, Ann. & Mag. Nat. Hist. vol. xiii. 2nd ser. p. 260.

Anthracoceros coronatus, Reich. Syst. Av. pl. 49 (1849).

Hydrocissa monoceros, Bon. Consp. Gen. Av. (1850) p. 90. sp. 1; id. Consp. Vol. Anisod. (1854) p. 2.

Hydrocissa pica, Bon. Consp. Gen. Av. (1850) p. 90. sp. 2; id. Consp. Vol. Anisod. (1854) p. 2.

Hydrocissa violaceus, Bon. Consp. Gen. Av. (1850) p. 90. sp. 4.

Annorrhinus violaceus, Bon. Consp. Vol. Anisod. (1854) p. 2. sp. 23.

Hydrocissa coronata, Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-58) vol. ii. p. 588. no. 869; Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 170. no. 470; Jerd. B. Ind. (1862) vol. i. p. 245. sp. 141; Blyth, Ibis (1866) p. 349; Holdsw. Proc. Zool. Soc. (1872) p. 425; Ball, Stray Feath. (1874) p. 387, & (1875) p. 290.

Buceros (Hydrocissa) coronatus, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 127. sp. 7872.

Bagnea-dunes in Bengal. *Wayera*, Mahratta. *Kamari*, Coucan, *Peshta-ganda* of the Gonds. *Kuchla-Kha*, Goomsoor.

HAB. Indian peninsula, Ceylon (BLYTH); Chota Nagpur (BALL).

The present species, the *A. malabaricus*, and the *A. convexus* have been confused one with another by different writers to an extraordinary degree. The *A. convexus* and the *A. coronatus* are, indeed, closely allied; but there is a considerable difference in their size, as well as in the marking and shape of their casques, while their geographical distribution is not at all the same. The bird now under consideration was first described by Boddaert (*l. c.*) in 1783, from Buffon's plate in the 'Planches Enluminées' of the *Calao des Philippines*. Scopoli followed shortly afterwards (1786) with his *Buceros pica*, founded upon Sonnerat's plate of the *Calao de la côte Malabar*. This figure, having the lateral rectrices entirely pure white, is evidently the same as Boddaert's *B. coronatus*. Latham noticed that this was a different species from the *B. malabaricus* of Gmelin, but contented himself by merely marking it var. β and γ . Shaw (1811) conferred two more names upon the luckless bird, viz. *B. monoceros* and *B. violaceus*, both of which have been about equally followed by succeeding writers. Many authors have also employed Gmelin's term of *malabaricus*, given to a very different, although allied species, one that never has the lateral rectrices entirely white.

The range of this species is confined to the peninsula of India and Ceylon. In the latter country, Layard observed it at Tangalla, and near St. Pedro, generally in large flocks; and Mr. Holdsworth also met with it a few miles from Aripo; and in the forest on the road between Kandy and Trincomalie, he frequently saw small parties of this Hornbill. Jerdon says that five or six usually compose the flocks seen in the Indian peninsula, and they generally keep in the thick forest, and near water.

The food of these birds is fruit—especially that of the Kuchla (*Strychnos nux vomica*), and berries, which they swallow whole. They are also fond of the fruit of the banian and other figs. In the jungles of Borabhum, Tickell says, they were very common, frequenting the Pipul trees, the fruit of which formed their principal food. The allied species, *A. malabaricus*, is stated by Mr. Inglis ('Stray Feathers,' vol. v. p. 20) to be very fond of live fish, which it catches in shallow pools. The way he discovered this predilection for an abnormal diet was as follows. He possessed a tame Otter and three tame Hornbills. At feeding-time the Otter was placed in a tub containing live fish; and these, when closely pressed, would jump out to escape from their pursuer, and were immediately swallowed by the Hornbills. Mr. Inglis has also found bones of fish in the stomachs of those birds which he had shot; and the Nagas affirm that, when these Hornbills are intent on fishing, they can be approached sufficiently close to be killed with a stick. I have not seen it anywhere recorded that the present species is a fish-eater; but it is not unlikely that it would accept this kind of food as readily as the *A. malabaricus*. The eggs are deposited in holes of trees, the female being, as usual, fastened in by her mate. The young keep with their parents many months after leaving the nest. The voice of this species is loud and harsh; and it is a very noisy bird, at the same time shy and wary. According to Tickell, it is never met with in high rocky lands, nor in barren tracks of saul-jungle, but abounds in the rich meadows composing the valley of the Soubourika. Occasionally, says Jerdon, in forest country, they come out in open spaces in the jungle to large fruit-bearing trees; and Layard states they appear to be much on the ground seeking food. The casque on the bill is not developed until after the first year.

In Chota Nagpur this species, according to Mr. Ball, affects certain localities, where it may generally be found in flocks of from six to ten individuals. He shot it in Manbhum, Singhbhum, and Sirguja, and also in the jungle on the road from the Ranchi plateau to Purula. He also shot at Lobloi, near Lanigarh, in the south of the Division, a specimen of the allied species, *A. malabaricus*, which formed one of a small party in a grove of mango trees; and in the same country he met with the present species, the limits of the two evidently overlapping. He states that the iris is orange-maroon, while Jerdon says it is crimson in the adult, brown in the young bird.

Some authors have used for this bird and its allies the generic term *Hydrocissa*, proposed by Bonaparte in his 'Conspectus Avium' for various species not nearly related beyond the fact that they belong to the same Family, among which the present one is included. I have not adopted this term, for the following reasons. Reichenbach proposed, in 1849 (of which date I believe there is no doubt), in his 'Syst. Nat. Av.,' the terms *Anthracosceros* (citing alone the *B. malabaricus*, Gmel., as its type) and *Anorrrhinus* (with the *B. galeritus*, Temm., as its type), and gave drawings of those characters in each considered by him generic. Although the portion of Bonaparte's 'Conspectus Avium' containing the Hornbills is dated 1849, I am unable to ascertain whether it appeared in that year separately, or with the rest of the first volume, which has its preface dated April 1850. In any event, on the score of priority, Bonaparte has no advantage over Reichenbach. The genus *Hydrocissa*, Bon., of which no definition whatever is given, is made to include such species as *coronatus*, Bodd. (called *monoceros*, Shaw), *galeritus*, Temm., *malayanus*, Raffl., and *exarhatus*, Temm. The other two, called *pica*, Scopoli, and *violaceus*, Shaw, are synonyms of *coronatus*, Bodd. The *Anthracosceros*, Reich., with *coronatus* as its type, would naturally include also the *B. malayanus*, Raffl.; and the *B. galeritus*, Temm., is appropriated by *Anorrrhinus*, Reich. There is thus left of Bonaparte's species only the *B. exarhatus*, Temm., to represent his genus *Hydrocissa*.

Male.—Bill curved, yellowish-white, base black, extending forward on the mandible. A large laterally compressed casque rises from base of maxilla, extends backwards over the head, and then curves forward for two thirds the length of maxilla, and returns by an acute angle to the culmen. A large patch of black covers the anterior three fourths of the casque, *but never reaches the maxilla*. In some specimens this patch is only a broad irregular black band. The posterior terminus is also black. Head covered by a loose, rather long crest, and, together with the neck, breast, and wings, black. Underparts and tips of primaries and secondaries white. Tail pure white, with the exception of the central pair, which are black. Naked skin on throat pale flesh-colour; iris crimson, brown in the young; feet dark grey.

Total length 3 feet, wing 13 inches, tail 14, bill along gape 7, casque 6, height of bill and casque 4.

Female same as male, but smaller. No black on posterior terminus of casque.

Specimens described and figured are in my own collection, and in the Paris Museum. The individual represented with the very large and black casque may be regarded as perhaps rather exceptional, as the majority of specimens have casques more like the others figured in the plate. This black-casque example is in the Paris Museum.

J.G.Keulemans del

Hanhart imp

ANTHRACOCERUS CONVEXUS.

ANTHRACOCEROS CONVEXUS.

TEMMINCK'S PIED HORNBILL

Buceros albirostris, Horsf. (nec Shaw), Trans. Linn. Soc. vol. xiii. p. 175.

Buceros malabaricus, Raffles (nec Gmel.), Trans. Linn. Soc. vol. xiii. p. 291.

Buceros convexus, Temm. Plan. Col. (1830) vol. ii. pl. 530; Gray, Gen. Birds (1849) vol. ii. p. 399; Motley & Dillw. Nat. Hist. Labuan (1855) p. 53; Schleg. Mus. Pays-B. (1862) p. 7; Gray, Hand-l. B. pt. ii. (1870) p. 127. sp. 7873; Sclat. Rev. List Vert. Anim. (1872) p. 172. sp. 276.

Buceros violaceus, Lord A. Hay (nec Shaw), Madr. Journ. (1844) vol. xiii. p. 148.

Buceros intermedius, Blyth, Journ. Asiat. Soc. Beng. vol. xvi. p. 994, and vol. xviii. p. 803; id. Cat. B. Mus. Asiat. Soc. (1849) p. 43. sp. 180; Wall. Ann. & Mag. Nat. Hist. 2nd ser. vol. xv. (1855) p. 98.

Hydrocissa convexa, Horsf. & Moore, Cat. Birds Mus. E.I. Co. (1856-8) vol. ii. p. 591. sp. 871; Moore, Proc. Zool. Soc. (1859) p. 448; Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 170. no. 469; Salvad. Ucc. di Borneo (1874) p. 80; Sharpe, Ibis (1879) p. 246.

Angka-Angka of the Sumatrans (RAFFLES).

HAB. Malacca, Java, Sumatra and Borneo (SCHLEGEL); Labuan (MOTLEY AND DILLWYN); Lâwas river, Borneo (USSHER, TREACHER).

The *A. convexus* was first separated from its relatives and described as distinct by Temminck (*l. c.*). It bears a very close resemblance in the colour of its plumage to the *A. coronatus*. In fact, in this respect the two birds are precisely alike; but there are differences, in the size and in the shape and markings of the casque of this species, which, together with its habitat, seem to point it out as distinct. It is a much smaller bird; the casque is swollen and not nearly so much compressed on the sides as that of *A. coronatus*; and the black mark on the casque reaches the maxilla, which is never the case in its ally. It has received two or three names besides those bestowed by Temminck; and some writers have called it by both Gmelin and Shaw's names, confounding it with the *A. malabaricus*. It has a fair distribution, being found throughout the Malayan peninsula, in Sumatra, and also at Labuan, where it was procured by Messrs. Motley and Dillwyn—who state that it is not uncommon, but shy and wild, going usually in pairs and keeping in the deep and tall jungle. Their food is almost entirely fruits, although in captivity they will eat any thing, but were especially fond of plantain. According to the Malays, they frequent the beach in search of small fish. Mr. Motley's captive birds were fed on boiled rice made into balls; but on one occasion they swallowed some ducklings that incautiously came within reach of their bills. The voice is a harsh and discordant croak, and is stated to sound something between the bray of an ass and the croak of a landrail! Besides Labuan, this species has also been procured

on the Láwas River, whence specimens were sent to London by both Governor Ussher and Mr. Treacher, the acting Governor of Labuan.

Male.—Bill and casque yellowish white, excepting the base of the bill and patches on the anterior portion and posterior terminus of the casque, which are black. Casque about as high as the bill, narrow for its whole length, and terminating abruptly anteriorly at about two thirds the length of the bill. The black on the anterior end of the casque extends onto the maxilla. Underparts, about three inches of the tips of the primaries and secondaries, together with the whole of the lateral tail-feathers, pure white. Rest of plumage black with a purplish gloss. Space around the eyes white, tinged with blue at the margins. Iris red-brown. Feet lead-colour.

Total length 29 inches, wing $11\frac{1}{2}$, tail $12\frac{1}{2}$, tarsus $1\frac{1}{2}$; length of bill $5\frac{1}{2}$ inches, length of casque 5, height of casque $1\frac{1}{2}$.

There is no difference in the plumage of the sexes; but the posterior end of the casque of the female is yellowish.

Specimens described and figured are in my own collection.

J. G. Keulemans lith.

ANTHRACOCEROS MALABARICUS

M. & N. Hanhart sculp.

ANTHRACOCEROS MALABARICUS.

PIED HORNBILL.

- Pied Hornbill*, Lath. Gen. Syn. (1781) vol. i. p. 349. sp. 6, pl. xi.
Buceros malabaricus, Gmel. Syst. Nat. (1788) vol. i. p. 359. sp. 7; Lath. Ind. Orn. (1790), vol. i. p. 143. sp. 6;
M'Clell. Proc. Zool. Soc. (1839) p. 164; Temm. Plan. Col. (1824) vol. ii. sp. 11 (text); Vieill. Nouv. Dict.
Hist. Nat. vol. iv. (1816) p. 595; Less. Trait. Ornith. (1831) p. 254. sp. 8; Ersch u. Grub. Ency. (1831) p.
285; Müll. & Schleg. Verh. Geschied. Neder. Ind. iv. (1839-44) pp. 22, 28.
Buceros albicornis, Wilkes? Ency. Lond. vol. iii. (1808) p. 479.
Buceros albirostris, Shaw, Gen. Zool. vol. viii. (1811) p. 13; Vieill. Nouv. Dict. Hist. Nat. vol. iv. (1816) p. 590;
Wagl. Syst. Av. (1827) sp. 11; Blyth, Cat. B. Mus. Asiat. Soc. Beng. (1849) pp. 43 & 803. sp. 179; Tickell,
Ibis (1864) p. 179; Blyth, Ibis (1864) p. 412; id. Journ. Asiat. Soc. Beng. (1843) vol. xii. pt. ii. p. 995;
Gray, Hand-l. B. pt. ii. (1870) p. 128. sp. 7874.
Buceros leucogaster, Blyth, Journ. Asiat. Soc. Beng. (1843) vol. x. p. 922 (juv.), vol. xii. p. 177 (1843).
Buceros nigrilbus, Hodg. Gray, Zool. Misc. (1844) p. 85.
Buceros affinis, Hutton, Journ. Asiat. Soc. Beng. vol. xviii. p. 802; Blyth, Cat. B. Mus. Asiat. Soc. Beng. (1849)
p. 43. sp. 178; Gray, Hand-l. B. pt. ii. (1870) p. 128. sp. 7875.
Hydrocissa albirostris, Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) vol. ii. p. 589. sp. 870; Cab. & Hein.
Mus. Hein. Th. ii. (1860) p. 171. no. 471; Jerd. B. Ind. vol. i. (1862) p. 247. sp. 142; Blyth, Ibis (1866)
p. 351; Blanford, Ibis (1870) p. 466; Jerd. Ibis (1872) p. 5. sp. 142; Schleg. Mus. Pays-Bas (1862) p. 6; Blyth,
B. of Burma (1875) p. 68. sp. 69; Sharpe, Proc. Zool. Soc. (1875) p. 102; Salvad. Cat. Ucelli Bor. p. 82.
Hydrocissa affinis, Jerd. B. Ind. vol. i. p. 247. sp. 143 (1862); Blyth, Ibis (1866) p. 349.
Hydrocissa coronata, Godwin-Austen, Journ. Asiat. Soc. Beng. (1870) p. 95.

HAB. Midnapore district, Rajmahal, Monghyr, Nepal, Assam, Sylhet, Arrakan, Tenasserim Provinces, Deyra
Doon, Burma, Tonghoo; Karen Hills (BLYTH); Borneo (SCHLEGEL); Labuan (Low).

Gmelin was the first to describe this bird; and although the colouring of the tail renders it
easy to be distinguished from its near relatives the *A. coronatus* and *A. convexus*, yet it has been
frequently confounded with them, and the synonymy has not been an easy task to unravel.
Shaw, as in the case of *A. coronatus*, conferred the name of *albirostris* upon the species,
evidently oblivious of Gmelin's prior name; and many writers have followed him, so that it is
perhaps better known by Shaw's appellation than by its rightful one of *malabaricus*. It differs
from both *coronatus* and *convexus* in having the lateral tail-feathers black to within a short distance
of the tips, which are white, whereas in the other species these feathers are wholly white; and
from *coronatus* it can in addition be distinguished by the black patch on the casque extending
onto the maxilla, which is never seen in examples of the allied species. It has a more northern
range than the *coronatus*. The two species meet in the jungles of Midnapore, whence the present
bird goes north to the base of the Himalayas, and then eastward as far as Burmah.

Blyth has separated the bird from the Deyra Doon as distinct, under the name of *B. affinis*,

simply on account of its larger size. Specimens vary greatly; and from the fact that as small ones are found among Deyra-Doon examples as are met with from other localities, I do not consider that there are two species. With respect to this so-called species of Blyth's, Dr. Jerdon remarks (Ibis, 1872, p. 5), "this is the Cachar bird noted by Major Godwin-Austen in his list of birds as *coronata*. His measurements quite agree with those of the next supposed species, *H. affinis*, Blyth, and confirm me in my doubts as to this bird being distinct. My suspicions were aroused by observations in the Dehra Doon, where I killed one or two individuals of the supposed species of much smaller size, nearly corresponding with the dimensions of *H. albirostris*." Tickell says that this bird is nearly omnivorous, but prefers fruit to other food. It breeds in Arakan and the Tenasserim provinces in July or August; and the female is stated to lay from two to four white eggs, in the hollow of a tree, without any nest.

Schlegel, in his catalogue of the birds of this family contained in the Leyden Museum, states that a skeleton of this species was obtained during the voyage of M. Diard, at Pontianak, in Borneo; and Mr. Low has sent it in a small collection of birds from Labuan.

Bill and casque yellowish white; base of casque, a broad patch at the anterior end of the same extending onto the maxilla, and the base of mandible jet-black. Casque about equal in height to the depth of the bill, narrow and sharp in front, swelling into a rounded form behind, and extending over the head at a rather sharp angle. The casque in the male extends forwards, following the line of the maxilla, and ends abruptly, being about a half inch in height [at its anterior termination. The nostrils are set in a groove on top of the maxilla, at the junction of the casque. Close to the base of the mandible is a bare flesh-coloured spot, and one behind the eye. Head and neck, back, wings, and tail, except the tips, black with a bluish gloss. Underparts, about two inches of the ends of primaries, and secondaries, except the uppermost ones, and ends of the lateral feathers of the tail pure white.

Total length 30 inches; wing 11; tail 12; tarsus $1\frac{3}{4}$; bill along gape 6, length of casque 5, height of bill and casque $3\frac{1}{4}$.

The female is like the male in plumage; but the casque does not extend so far forward on the maxilla, and slants towards the culmen at a different angle from that of the male. Its anterior portion, together with the tips of the bill, are black.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

J G Keulemans lith

Hanhart imp
33

ANTHRACOCEROS FRATERCULUS

ANTHRACOCEROS FRATERCULUS.

COCHIN-CHINA PIED HORNBILL.

Anthracoceros fraterculus, Elliot, Ann. & Mag. Nat. Hist. (1878) 5th ser. vol. i. p. 85.

HAB. Cochin China.

This species bears the same relationship to *A. malabaricus* as *A. convexus* does to *A. coronatus*, and apparently represents the first-named in Cochin China. It is a smaller bird than *A. malabaricus*, and has a differently shaped casque, much more compressed both anteriorly and posteriorly. The black mark on the anterior part does not reach the maxilla, but is confined to the casque, as is witnessed in *A. coronatus*; this mark in the other two species, however, always extends onto the maxilla. The lateral rectrices, by being white upon their apical third, show that the affinities of this species are with *A. malabaricus*, and not with the others mentioned. The type (which is fully adult), and a second specimen exactly resembling it (received at the Paris Museum since my description was published), both came from Cochin China; but the exact locality was not indicated. It is very interesting to notice the distribution of these four species, and observe that, while their habitats are contiguous, yet each is restricted to its own range of country, although the physical qualities of the different regions must resemble each other in very many particulars. Thus *A. malabaricus*, which has the most northern range of the four, is distributed from Siam and Burmah on the east, along the base of the Himalayas to the jungles of Midnapore, on the west of the Bay of Bengal, its southern limit. The Siamese examples of this species, of which I have lately seen a series that were received at the Paris Museum, are very much smaller than Indian specimens in all their dimensions, with small casques that appear like miniatures of those possessed by their larger brethren. The colour of their plumage is precisely like that of *A. malabaricus*; and the black mark on the casque extends onto the maxilla. They seem to be a diminutive race, but cannot be separated specifically from *A. malabaricus*. In Midnapore this species meets its ally *A. coronatus*, the two mingling together; and the last-named thence ranges throughout the Indian peninsula into Ceylon. *A. convexus* (the near ally of *A. coronatus*) is more particularly an island species, confined, so far as is known, to Malacca, Java, Sumatra, and Borneo, and differs from its continental relative in size as well as in the shape and marking of the casque. It is a singular fact that there should be four such closely allied species, two of which differ from their larger relatives in precisely the same way, illustrating in this instance that the influences peculiar to an insular existence, as exhibited in *A. convexus*, have produced the same effect in regard to its specific characteristics as have those of the continent upon *A. fraterculus*.

Adult Male.—Bill light yellow, with a black spot at base of mandible. A casque rises from the base of the culmen, extends backwards over the centre of the head, then curves forwards, and returns to the culmen by almost a right angle, at about one third its length from the tip of the maxilla. This is compressed laterally at both its anterior and posterior terminations, swelling outwards in the centre, but inclining to a keel-shape on top along its entire length. This casque is yellow like the bill, black on its anterior face, and with a broad black patch occupying nearly half the anterior portion, but which *does not reach to the maxilla*. Naked skin around the eye and on sides of the throat flesh-colour. Head, neck, throat, upper part of breast, back, wing, and central tail-feathers black, with dark green reflections. Entire underparts, thighs, and tips of the secondaries and primaries pure white. Lateral tail-feathers have their apical third pure white, rest black, with green reflections. Tarsi and feet black.

Total length from base of maxilla to end of central rectrices $23\frac{1}{2}$ inches, wing $10\frac{3}{4}$, tail $11\frac{1}{4}$, bill along gape $5\frac{1}{8}$, casque on top $4\frac{7}{8}$, height at base of maxilla $1\frac{1}{4}$, height of bill and casque at base $2\frac{3}{4}$, tarsus $1\frac{5}{8}$.

The figures in the Plate were drawn from the type in the Paris Museum.

1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900

ANTHRACOCEROS MALAYANUS

ANTHRACOCEROS MALAYANUS.

WHITE-BILLED HORNBILL.

Buceros malayanus, Raffl. Trans. Linn. Soc. vol. xiii. p. 292; Müll. and Schleg. Verh. Geschied. Nederl. Ind. (1839-44) pp. 23, 29; Hay, Madras Journ. (1844) vol. xiii. p. 151, ♀; Blyth, Journ. Asiat. Soc. Beng. (1847) p. 995; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 43. sp. 181; Wall. Ann. & Mag. Nat. Hist. (1855) vol. xv. p. 98; Schleg. Mus. Pays-B. (1862) p. 7; Blyth, Ibis (1866) p. 352; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7879; Giebel, Thes. Ornith. (1872) p. 500; Sclat. List Vert. Anim. (1877) p. 225. sp. 352.

Buceros antarcticus, Swain. Class. Birds, vol. ii. p. 296.

Buceros antracicus, Temm. Plan. Col. vol. ii. no. 529.

Buceros bicolor, Eyton, Proc. Zool. Soc. (1839) p. 104; Blyth, Journ. Asiat. Soc. Beng. (1843) vol. xii. pt. ii. p. 995, ♀.

Buceros elliotti, Lord A. Hay, Madras Journ. (1844) vol. xiii. pt. 2, p. 152; id. Journ. Asiat. Soc. Beng. vol. xvi. p. 995.

Buceros nigrirostris, Blyth, Journ. Asiat. Soc. Beng. (1847) p. 995, (1849) p. 803 (♀); id. Cat. Birds Mus. Asiat. Soc. (1849) p. 44. sp. 182; Wall. Ann. & Mag. Nat. Hist. (1855) vol. xv. p. 98; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7881.

Hydrocissa malayana, Bon. Consp. Gen. Av. (1850) p. 90. sp. 5; Horsf. and Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 592. no. 872; Moore, Proc. Zool. Soc. (1859) p. 448; Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 170. no. 468.

Hydrocissa nigrirostris, Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 593. sp. 873; Moore, Proc. Zool. Soc. (1859) p. 449.

HAB. Malayan peninsula, Sumatra, Borneo (BLYTH).

The female of this species was described by Blyth as *B. nigrirostris*; but Wallace states that he shot both the yellow- and black-billed birds on the same tree, and on dissection the black-billed ones proved to be females. There are four specimens now before me, collected by Mr. Wallace—two yellow-billed and two with black bills. One of each of these possess the broad white superciliary stripe; the other two have not this mark. Whether this is assumed when old, or is a character of the nuptial state only—whether there may be two species, one with this conspicuous mark and one without it, I am unable to state. All four specimens are apparently fully adult; and the two white-billed ones, having the posterior terminus of the casque black, are without doubt males; for it is seemingly a law in birds of this family with light-coloured bills that the female should have the hind part of the casque of a different colour from that observed in the male; and if the bird with the white casque and without the stripe were a female, the hind part of the casque would be white like the bill. Believing Mr. Wallace's statement to be correct regarding the sex, we must wait further information as to when or at what period of their life the white superciliary stripe is assumed by both sexes. Possibly it indicates the seasonal dress.

Adult Male.—Bill and casque white, the latter about equal in height to the bill. Posterior margin of casque hidden by feathers, black. The casque projects back over the head from the base of the maxilla, and curves forwards for about two thirds the length of the bill, terminating in an abrupt angle, the upper end inclining forwards nearly to a point; it is broadest at the posterior margin, and much compressed anteriorly. In some specimens a broad white superciliary stripe begins just forward of, and above, the eye, and extends to the occiput. Rest of plumage jet-black, with the exception of about three inches of the tips of the lateral tail-feathers, which are white. Bare space around the eye blue. Total length 30 inches, wing 13, tail 13, tarsus $1\frac{3}{4}$, bill along gape $5\frac{3}{4}$, height of bill and casque at base $5\frac{1}{2}$, length of casque $5\frac{1}{4}$.

Female.—The bill and casque black, the latter moderate in size, and terminating abruptly anteriorly. A broad white stripe in some examples commences at the base of the casque, goes above the eye, and extends to the occiput. Rest of plumage jet-black, with the exception of the tips of lateral tail-feathers, which are white. Bare skin around the eyes livid; iris greyish brown. Total length 24 inches, wing 12, tail 12, bill along gape 5, length of casque $2\frac{1}{2}$, height of bill and casque $2\frac{1}{2}$.

J.G. Keulemans lith.

CRANORRHINUS CASSIDIX.

M & N Hanhart sculp.

CRANORRHINUS CASSIDIX.

CELEBES HORNBILL.

Buceros cassidix, Temm. Planch. Col. vol. ii. pl. 210; Wagl. Syst. Av. (1827) sp. 3; Griff. Anim. King. (1829) vol. ii. p. 434, pl.; Less. Trait. Ornith. (1831) p. 253. sp. 6; id. Man. Ornith. vol. ii. p. 105; Mull. & Schleg. Verh. Geschied. Nederl. Ind. (1839-44) p. 24; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399. sp. 22; Schleg. Mus. Pays-B. (1862) p. 9; Gieb. Thesaur. Ornith. (1872) p. 497.

Calao cassidix, Bon. Consp. Gen. Av. (1850) p. 90. sp. 1.

Cassidix cassidix, Bon. Consp. Vol. Anisod. (1854) p. 3. gen. 10. sp. 27.

Cranorrhinus cassidix, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 173. no. 476; Wald. Trans. Zool. Soc. (1872) vol. viii. p. 47. sp. 58.

Buceros (Cranorrhinus) cassidix, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 129. sp. 7889.

HAB. Celebes: Tondano (REINWARDT); Menado (WALDEN); District of Maros, Macassar (WALLACE).

In the eighth volume of the 'Transactions of the Zoological Society,' in his paper on the birds of Celebes, Lord Walden gives woodcuts of the heads of the adult and young of both sexes of the species, illustrating the development of the casque. This in the young male is very much swollen posteriorly, but slopes rapidly forwards to the culmen, forming a graceful curve. The mandibles are quite smooth. In another, rather older individual, the casque has taken somewhat the form of the adult's, and is upright at its anterior end, and more separated from the culmen along its lower line. The base of the mandibles is covered by a plate; but there are no grooves. In the adult male the casque is nearly of the same thickness throughout, being less compressed anteriorly and less swollen posteriorly, while it inclines anteriorly to the culmen by an acute angle. The casque possesses several distinct undulations. The base of the mandibles is grooved deeply, diagonally, forming two folds on the maxilla, and three on the mandible. The cutting-edges of the bill are much broken on their anterior half. The adult female has a similar casque, but smaller; the anterior edge stands at a right angle to the culmen, and the grooves on the mandibles form three folds on each. The bill appears to increase in length, even after the wings and tail have become fully grown.

Of the economy and habits of this fine species, I have not been able to find much recorded. Mr. Wallace met with it during his sojourn in Celebes; and from his observations it would appear to eat insects; for he states that he has taken fragments of large long-horned *Batocera* from the gullet of individuals, and once he saw a *C. cassidix* capture one of these insects, beat it repeatedly against a branch, and then swallow it. In flying, like other members of this family, it makes a great noise with its wings.

Male.—Bill large, curved, orange-yellow, except the basal portion, which is dark red and

crossed by three or more prominent transverse ridges. A high, upright, keel-shaped casque, swollen in the posterior portion, descending rapidly from the anterior terminus, rises from the centre of the culmen about one third its length from the base, and extends backwards to the middle of the head; this casque is of a uniform deep red. Bare skin around the eye blue; that on throat bluish white, crossed on the lower part by a narrow black line. Occiput and nape chestnut. Sides of face and neck and upper part of breast light tan-brown colour. Entire rest of plumage of body and wings black, with greenish reflections. Tail pure white.

Total length 40 inches; wing 18; tail 14; bill along the gape 9, height at base with casque 5; length of casque $4\frac{1}{2}$, height in front from culmen $2\frac{3}{4}$. These measurements are taken from a very fine adult specimen, and possibly exceed the average.

Female.—Bill, casque, bare skin around eyes and throat, and the tail the same as the male, rest of plumage black.

Total length $29\frac{1}{2}$ inches, wing $15\frac{1}{2}$, tail 11, bill $6\frac{1}{2}$, casque 4.

1000000

J.G. Keulemans lith.

Hanhart del.

CRANORRHINUS CORRUGATUS

CRANORRHINUS CORRUGATUS.

WRINKLED-BILL HORNBILL.

- Buceros corrugatus*, Temm. Plan. Col. (1830) vol. ii. no. 531; Müll. & Schleg. Verh. Gesch. Ned. Ind. (1839-44) pp. 24, 31, sp. x.; Gray, Gen. Birds, (1849) vol. ii. p. 399. sp. 20; Schleg. Mus. Pays-B. (1862) p. 9; Sclat. Proc. Zool. Soc. (1868) p. 261; Bartl. Proc. Zool. Soc. (1869) p. 142; Flow. Proc. Zool. Soc. (1869) p. 150; Murie, Proc. Zool. Soc. (1874) p. 420.
- Buceros gracilis*, Temm. Plan. Col. (1832) no. 535, ♀.
- Buceros rugosus*, Begbie, Ann. & Mag. Nat. Hist. (1847) vol. xvii. p. 405.
- Calao corrugatus*, Bon. Consp. Gen. Av. (1850) p. 90. sp. 2.
- Cassidix corrugatus*, Bon. Consp. Vol. Anisod. (1854) p. 3.
- Hydrocissa migratoria*, Maing. Proc. Asiat. Soc. Beng. (1868) p. 196.
- Buceros (Cranorrhinus) corrugatus*, Gray, Hand-l. Birds, (1870) pt. ii. p. 129. sp. 7890.
- Cranorrhinus corrugatus*, Wald. Trans. Zool. Soc. vol. viii. p. 51 (1872); Sharpe, Ibis, (1879) p. 246; Salvad. Ucc. di Borneo, (1874) p. 86.

HAB. Malacca (DIARD, BEGBIE); Sumatra (MÜLLER); Borneo (SCHLEGEL, DIARD); Banjarmassing (SCHWANER); Sarawak (DORIA, BECCARI); Lâwas river (Ussher).

This Hornbill was described by Temminck in his Planches Coloriées, but not figured; and the female was also given in the same work as a distinct species under the name of *Buceros gracilis*. It is a native of Malacca, Sumatra, and Borneo, very rare in collections; and nothing is known of its economy or habits in the wild state. A specimen living in the London Zoological Gardens was stated by Mr. Sclater (*l. c.*) to have the end of the tail-feathers white instead of rufous; I have also noticed this as occurring in occasional specimens, but am inclined to regard it as only an individual peculiarity, as the majority of the examples that I have seen have had the end of the rectrices rufous or light chestnut. Salvadori, in his fine work on the Birds of Borneo (*l. c.*), states that the male of this species has the bare skin of the throat canary-yellow, and the iris dark red, while the iris of the female is yellowish brown.

In the 'Proceedings of the Zoological Society of London' for 1869, Mr. Bartlett gives an account of the sac, containing undigested food, cast up by a Hornbill of this species, at that time an inmate of the Gardens in Regent's Park. This curious object was submitted to Dr. Murie for examination, who ascertained it to be simply the "thickened semichondrified lining membrane of the gizzard. All the puckerings and indentations were more or less exactly represented, though less sharp in outline than is ordinarily the case. The mucous surface of the inner wall of the bag was slimy, otherwise perfectly identical with the same structure in a healthy bird. The surface outside, on which might be said to be the submucous tissue, was moist, comparatively uninjured,

and free from any effusion or disease. The rim of the mouth of the bag was irregular and shreddy, and thinned away at its free edge." This sac contained seven or eight discoloured grapes, in an undigested state, but, from their appearance, had been acted upon by the gastric juice. Somewhat alarmed at finding the individual thus deprived of the coating of its gizzard, Mr. Bartlett kept a close watch upon the bird, and in a day or two afterwards obtained another specimen of this bag of fruit, which in every way was like the first. As the Hornbill evidently suffered no inconvenience from the loss of this epithelial lining of the gizzard, it was not the result of disease; and Mr. Bartlett's conclusion was, that it was the means by which the male Hornbill supplies the female with food during incubation when she is walled up in the hollow of a tree. At such times the female generally becomes quite fat, while, according to Dr. Livingstone (Missionary Travels in South Africa, p. 613), "the poor slave of a husband gets so lean that on the sudden lowering of the temperature, which sometimes happens after a fall of rain, he is benumbed, falls down, and dies." This exhaustion, Mr. Bartlett naturally considers, is caused "not only by the constant and continued reproduction of the actual food taken by the male, but also by the supply of nutritive secretion in which the same is enveloped." Many kinds of birds are in the habit of casting up their food, at times, in a partially digested state for the nourishment of the young, or, as in the case of Raptorial birds, in the shape of pellets formed of feathers, bones, or hair of the creatures they have made their prey; but in no instance that I am aware of is this accompanied by the stomach's walls, save among the species of the Bucerotidæ; and this may be naturally regarded as the most wonderful habit of the members of this extraordinary family.

Male.—Bill curved, yellow. A bright-red casque rises from the culmen nearly halfway from the tip, and projects back onto the head above the eye, with two deep plaits on the anterior terminus. Skin around the eye blue; bare skin of throat red, with blackish mottlings. Top of head and back of neck, wings and body black. Sides and front of neck to breast white. Tail, basal half black, rest light chestnut.

Bill along culmen 7 inches, length of crest on maxilla 3, height of maxilla and crest $2\frac{1}{8}$, width of mandible $1\frac{1}{8}$, height of crest 1, from eye to end of bill $5\frac{3}{4}$, wing $14\frac{1}{2}$, tail $11\frac{1}{2}$, tarsus 2.

Female has the bill and crest yellow, the latter much smaller than that of the male; bare skin of eye and throat blue. Entire plumage black, except terminal half of tail, which is light chestnut.

Bill from eye to tip $4\frac{3}{4}$ inches, width of bill $2\frac{5}{8}$, length of crest $2\frac{3}{4}$, wing $13\frac{1}{2}$, tail 10.

My description and figures were taken from specimens in the collection of the British Museum.

J.G. Keulemans lith.

M & N Hanhart imp.

CRANORRHINUS WALDENI.

CRANORRHINUS WALDENI.

LORD WALDEN'S HORNBILL.

Cranorrhinus waldeni, Sharpe, Trans. Linn. Soc. 2nd ser. Zool. vol. i. (1877).

HAB. Island of Panay, Philippines.

This fine and interesting species was obtained in the island of Panay, one of the Philippines, by Professor Steere, who lately brought to England a collection of birds from the islands forming that group. Mr. Sharpe described this species in a memoir upon the birds of the Philippines, published in the 'Transactions of the Linnean Society,' and named it after the President of the Zoological Society, Viscount Walden.

Professor Steere has given a short account of his meeting with this species, which is all we know about it; and for the opportunity of inserting it here I am indebted to Mr. Sharpe, who kindly sent me a copy of his MS. The discoverer of this bird states:—"I shot this Hornbill on the highest ridge of the mountains west of Ilo-Ilo. This is the only place where any of the virgin forest is left; and here alone I saw these birds. They were not very scarce; but I could only get a single specimen, as they flew so high in the trees that my gun could scarcely reach them."

The *C. waldeni* belongs to a small group in this family, now composed of four species, characterized by a rather small upright casque, corrugated laterally. It appears to be nearest allied to the *C. cassidix*, and, like its immediate relatives, is rather a showy-looking bird. The type, which is unique, has been taken to America by Professor Steere. Bill red, the basal half of the mandible crossed by several prominent curved ridges. An upright keel-shaped casque, broadest on its posterior end, rises above the eye and extends for nearly two thirds the length of the maxilla, its anterior end being at a right angle to the culmen; this casque is deeply grooved along its sides for two thirds its length from the anterior terminus. General colour above and below greenish black, wings also the same. Head and neck dark chestnut, fulvous above the eye and on the ear-coverts. Tail cinnamon, apical portion and basal third greenish black. The figure is two fifths the natural size.

The female is not known; but she will probably be found to have a general black plumage, with the tail, and perhaps the casque, similar to those of the male. In size she will be slightly smaller.

J.G. Kwak lith.

Hanhart imp.

CRANORRHINUS LEUCOCEPHALUS.

CRANORRHINUS LEUCOCEPHALUS.

WHITE-HEADED HORNBILL.

Buceros leucocephalus, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 592.

Buceros sulcatus, Temm. Plan. Col. vol. ii. pl. 69; Wagl. Syst. Av. (1827) sp. 10; Less. Trait. Ornith. (1831) p. 254. sp. 11; Gray, Gen. Birds (1849), vol. ii. p. 400. sp. 25; Schlegel, Mus. Pays-B. (1862) p. 10; Gray, Hand-l. B. pt. ii. (1870) p. 129. sp. 7891.

Tockus sulcatus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 1.

Calao sulcatus, Bon. Consp. Vol. Anisod. (1854) p. 3.

Buceros (Penelopides) sulcatus, Von Mart. Journ. für Ornith. (1866) p. 18.

Cranorrhinus leucocephalus, Wald. Trans. Zool. Soc. (1875) p. 165, pl. 27. fig. 1 (♂), fig. 2 (♀); id. Proc. Zool. Soc. (1877) p. 824. sp. 28.

HAB. Mindanao (SCHLEGEL); Island of Camiguin (BRIT.-MUS. specimen).

The adult of this species is known to ornithologists chiefly from Temminck's plate, taken from the specimen in the Leyden Museum. It seems to be allied to *C. cassidia* from its elevated casque and deep transverse grooves at the base of the mandible. It was first described by Vieillot, as given above; but Temminck afterwards called it *sulcatus*, by which name it has usually been recognized, Vieillot's description having been overlooked. Although the latter states that the plumage is *black* generally, yet he says, at the same time, that his description was taken from Temminck's specimen. He probably had a female before him.

Male.—Bill curved, red. An elevated crest of a bright red colour, corrugated with deep furrows, runs along the culmen for about one half its length. At base of mandible are four deep diagonal ridges. Top of head and back of neck dark foxy red. Throat bare at base of mandible for about an inch; this is fiery orange-red; around the skin the feathers are whitish. Bare skin around the eyes bright orange-red. Side of neck and upper part of breast deep buff. Rest of body and wings black. Tail buffy white, probably darker in life, with an apical black band. Iris crimson; feet dark brown.

Bill from eye to tip 5 inches, length of casque $2\frac{3}{4}$, height of casque $1\frac{1}{4}$ inch, from edge of maxilla to top of casque 2, width of mandible 1 inch, wing 14, tail 10, tarsus 2. (*Ex* type, Leiden Museum.)

Female.—Jet black. Tail buffy white, with an apical black band.

Total length $20\frac{1}{2}$ inches, wing 11, tail 9, bill $4\frac{1}{2}$. (*Ex* specimen in British Museum.)

J.G. Keulemans hth

PENELOPIDES MANILLÆ.

Hanhart imp.
24

PENELOPIDES MANILLÆ.

MANILLA HORNBILL.

- Manilla Hornbill*, Lath. Gen. Syn. (1781) vol. i. p. 354. sp. 9.
Calao de Manille, Buff. Pl. Enlum. (1783) no. 891 (juv. ♂).
Buceros manillæ, Bodd. Tab. Pl. Enlum. D'Aubent. (1783) p. 54.
Buceros manillensis, Gmel. edit. Linn. Syst. Nat. (1788) vol. i. p. 361; Lath. Ind. Ornith. (1790) vol. i. p. 145;
Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 208.
Le Calao à bec ciselé, Levaill. Ois. d'Amér. (1801) pl. 18 (juv. ♂).
Buceros manillensis, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 596; id. Enc. Méthod. (1823) p. 305.
Buceros sulcirostris, Wagl. Syst. Av. (1827) sp. 13 (♂ ♀).
Tockus sulcirostris, Bon. Consp. Gen. Av. (1850) p. 91. sp. 5.
Penelopides sulcirostris, Bon. Consp. Vol. Anisod. (1854) p. 3.
Buceros panayensis, Schl. Mus. Pays-Bas, (1862) p. 11.
Buceros (Penelopides) panini, v. Martens, Journ. für Ornith. (1866) p. 18.
Penelopides manillæ, Wald. Trans. Zool. Soc. (1875) vol. ix. p. 168; id. Proc. Zool. Soc. (1877) p. 692. sp. 25.

HAB. Philippines, Luzon (EVERETT).

The present bird and the *P. panini* have been continually confounded together by all writers on ornithology up to a very late date. It was supposed that there was but a single species in the Philippine Islands; and the synonymy was all placed under one head. But the fact is, there are two forms which are very distinct, similar somewhat in the plumage of the body, but with the tails very differently coloured, so that they may be recognized at once. These well-marked tail-feathers are common to both sexes, so that the females can be readily assigned to their proper males. The present species was first described by Boddaert, from Buffon's plate of the *Calao de Manille*, representing a young bird. The term *sulcirostris* was afterwards applied to it by Wagler and Bonaparte. It is still a rare species, but few museums possessing examples. The figures in the accompanying Plate were drawn from specimens kindly loaned to me by Lord Tweeddale from his collection.

Male.—Bill. A low crest, much compressed on the sides anteriorly, rises at base of culmen, follows the curve of the maxilla for about two thirds its length, and terminates abruptly at a right angle to the culmen. The basal half of the maxilla is covered with a jet-black plate, with five shallow transverse orange-yellow grooves. The mandible has a similar plate; but the grooves are narrower and deeper, and cross the plate diagonally. The remaining portion of the bill and the casque are dark brown. Bare space around the eye, at base of mandible, and upper part of throat

white. Ear-coverts black, the feathers long, and mingled with those of the crest. A black band encircles the throat, following the line of the nude skin. Head crested, and, together with the neck and entire underparts, is silky yellowish white. Back and rump dark olive-brown, the latter with the edges of some feathers orange. Wings dark shining green, the edges of the webs of primaries and secondaries rufous brown. The tail has the outer rectrices the same colour as the wings, with a broad orange-red bar on the terminal third; the median feathers rufous brown; on the four central rectrices the orange bar is darkest in colour, becoming lighter as it goes to the outer feathers, and is confined to the inner web of the outside one, not passing the shaft. Irides crimson. Feet and nails dull black.

Total length without bill $18\frac{1}{2}$ inches, wing $9\frac{1}{2}$, tail 9, bill $3\frac{1}{2}$; casque on top $2\frac{1}{2}$, height at anterior end $\frac{1}{2}$; tarsus $1\frac{1}{2}$.

Female.—Bill like that of the male, but smaller. Head and crest leaden grey. Cheeks black; and a blackish band follows the line of the nude skin, which is distributed in the same way as that of the males. Neck and underparts mouse-colour, tinged with rufous brown. Wings and tail like those of the male.

The colouring of the soft parts is taken from Mr. Everett's notes, made at Luzon from the birds in the flesh, and kindly communicated to me by Lord Tweeddale.

J.G. Keulemans lith.

PENELOPIDES PANINI.

Hanhart comp.
21

PENELOPIDES PANINI.

PANAYAN HORNBILL.

- Le Calao mâle à bec cizelé de l'Isle Panay*, Sonn. Voy. Nouv. Guin. (1776) p. 123, pl. 82 ♀.
Le Calao femelle à bec cizelé de l'Isle Panay, Sonn. Voy. Nouv. Guin. (1776) p. 124, pl. 83 ♂.
Panayan Hornbill, Lath. Gen. Syn. (1781) vol. i. p. 353. sp. 8.
Calao de l'Isle Panay, Buff. Plan. Enlum. (1783) no. 780 ♀, 781 ♂.
Buceros panini, Bodd. Tab. Plan. Enlum. D'Aub. (1783); G. R. Gray, Gen. Birds (1849) vol. ii. p. 340. sp. 23;
Eyton, Osteol. Av. p. 63 (1867); Giebel, Thes. Ornith. (1872) p. 502.
Buceros panayensis, Scop. Faun. et Flor. Insub. (1786) p. 87. sp. 30; Gmel. Syst. Nat. (1788) vol. i. p. 360; Lath.
Ind. Ornith. (1790) vol. i. p. 144; Shaw, Gen. Zool. (1811) vol. viii. p. 33; Vieill. Nouv. Dict. Hist. Nat.
(1816) vol. iv. p. 594; id. Ency. Méth. (1823) p. 304; Temm. Plan. Col. (1824) vol. ii. sp. 12; Ersch u.
Grub. Ency. (1824) p. 287; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 45. sp. 188.
Le Calao à bec cizelé mâle, Le Vaill. Ois. Rares d'Amér. (1801) vol. i. pl. 16, ♀ ad.
La femelle du Calao à bec cizelé, Le Vaill. Ois. Rares d'Amér. (1801) vol. i. pl. 17, ♂ ad.
Penelopides panayensis, Bon. Consp. Vol. Anisod. (1854) p. 3.
Penelopides panini, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 168. sp. 465 (partim); Wald. Trans. Zool. Soc.
(1875) vol. ix. p. 166, pl. 28. fig. 1 ♂, fig. 2 ♀.
Buceros (Penelopides) panini, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 129. sp. 7892.

HAB. Moluccas, Philippines (BLYTH).

The synonymy of this bird has been even more confused than that of its ally, the *P. manillæ*, and from the same cause, viz. the supposition that there was only one species. Lord Walden in his paper "on the Birds of the Philippine Islands," gives a figure of this one, with correct synonymy and description.

The deep-chestnut tail, with apical black band glossed with green, will always serve to distinguish it from its relative, the *P. manillæ*. It was first described and figured by Sonnerat, who confounded the sexes, mistaking the male for female, as the *Calao à bec cizelé de l'Isle Panay*, and afterwards by Buffon in the 'Planches Enluminées,' under Sonnerat's name. Boddaert gave to this figure the Latin appellation of *Buceros panini*. It is a very rare species in collections, though lately some good series of specimens have been obtained.

Of its habits and economy nothing has as yet been recorded by any of those persons who have obtained it in the countries it inhabits.

Male.—Casque slightly elevated, deep red in colour, sharply defined on top and ending abruptly anteriorly. Two thirds of the basal portion of the maxilla is covered by a black plate, indented

by six perpendicular orange-yellow grooves. Base of mandible covered by a similar plate with six diagonal orange-yellow grooves, narrower than those on the maxilla. Apical part of bill apparently dark red, like the casque. Bare skin around the eye and upper part of throat flesh-colour. A broad black line starts from beneath the eye, extends backwards just beyond the ears, and follows the lower margin of the bare skin on the throat, becoming broadest in the centre. Rest of head, neck, and underparts buffy white. Abdomen ferruginous; thighs and upper tail-coverts, vent, and rump chestnut-red. Tail deep chestnut, until within two inches of the tip, when it becomes black with a greenish gloss.

Total length without bill 20 inches, wings 10, tail $9\frac{1}{2}$, bill from eye to point 4, length of casque $1\frac{3}{4}$, tarsus $1\frac{1}{2}$.

Female.—Entire plumage black, with the exception of the tail, which is like that of the male. Bill similar but smaller.

I am indebted to the Marquis of Tweeddale for the loan of the specimens here described and figured.

J.G. Keulemans lith.

PENELOPIDES AFFINIS.

Hanhart imp.

PENELOPIDES AFFINIS.

ALLIED HORNBILL.

Penelopides affinis, Tweeddale, Ann. & Mag. Nat. Hist. ser. 4, vol. xx. p. 534 (1877); id. Proc. Zool. Soc. (1877) p. 824. sp. 29, (1878) pp. 108, 341, 947.

HAB. Zamboanga, Butuan, Surigao, Island of Mindinao, Dinagat, Amparo, South Leyte, Philippines (EVERETT).

This bird, described by the late Lord Tweeddale (*l. c.*), is a representative form of *P. panini*, and differs in not having the perpendicular grooves on the lateral plates of the maxilla, these being smooth, and also by having the grooves on the mandible straight instead of curved as in *P. panini*. The bill is altogether less powerful. The abdominal region and under tail-coverts are coloured like the breast, and not rufous as in its ally; and the rectrices have a black band at their insertion. Lord Tweeddale states that "in one example some of the black upper tail-coverts are tipped and margined with dirty rusty. An old Surigao female has several of the secondary quills margined with ferruginous. A young Surigao male has a small part of the outer webs of the fourth, fifth, and sixth primaries edged albescent. An old Surigao male has the thigh-coverts, ventral region, and under tail-coverts pale rufous, and most of the upper tail-coverts as in *P. panini*, otherwise as in typical *affinis*. In a Butuan and a Surigao male, the middle pair of rectrices, with the exception of the basal black band, are rufous throughout, the normal black terminal band being only indicated by black blotches. The new rectrices are pale fulvous white (instead of ferruginous) when they first come in. The number of grooves on the mandible varies from three to four. The extreme altitude of the bill, ending at the nostril, is 1.06 inch, whereas in *P. panini* it is 1.37.

"♂. Wing 9.25 inches, tail 9.00, tarsus 1.75, culmen 3.40.

"♀. Wing 8.75 inches, tail 8.37, tarsus 1.60, culmen 3.25.

"*a*, ♂, Butuan, May: iris crimson; orbital skin, nude portion of skin, and unfeathered part of the rami white; casque, smooth lateral plates of maxilla, and the ungrooved triangular space at base of mandible dark brown; the whole apical part of the maxilla from the smooth lateral plates, and of mandible from the grooved base, pale brown; grooves on mandible dark brown; intermediate smooth spaces light ochreous; feet greenish lead-colour; nails greyish black. *b*, ♀, Butuan, May: iris crimson; orbital skin, nude part of chin, and unfeathered part of the rami dark blue; bill brown; feet dark greenish leaden. *c*, *d*, ♂ ♀, Surigao, May."

The specimens figured are the types of Lord Tweeddale's species, kindly loaned to me by Captain Ramsay, their present owner.

J. G. Leuclercq lith.

Hanhart imp
31

CERATOGYMNA ELATA.

CERATOGYMNA ELATA.

YELLOW-CASQUE BLACK HORNBILL.

- Buceros elatus*, Temm. Plan. Col. (1830) vol. ii. no. 521, fig. 1 (bill); Cass. Journ. Acad. Nat. Sc. Phil. (1849-50) vol. i. p. 135; G. R. Gray, Gen. Birds (1849), vol. ii. p. 390. sp. 13; Sundev. Öfvers. Kongl. Vetensk. Akad. Förh. (1849) p. 161; Hartl. Journ. für Ornith. (1854) p. 126. sp. 386, (1861) p. 261; Hartl. Ornith. West-Afr. (1857) p. 161; J. H. Gurney, Ibis (1859) p. 153; Schleg. Mus. Pays-Bas (1862) p. 18; Eyton, Osteol. Av. (1867) p. 63; Sharpe, Ibis (1872) p. 67; Sclater, Rev. List Vert. Anim. (1872) p. 72.
- Buceros cultratus*, Sundev. Öfvers. Kongl. Vetensk. Akad. Förh. (1849) p. 160 ♀; Jard. Contr. Ornith. (1852) p. 161; Hartl. Journ. für Ornith. (1854) p. 127. sp. 390; Hartl. Ornith. West-Afr. (1857) p. 161.
- Bucorvus elatus*, Bon. Consp. Gen. Av. (1850) p. 89. sp. 2.
- Ceratogymna elata*, Bon. Consp. Vol. Anisod. (1854) p. 2.
- Buceros (Ceratogymna) elatus*, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 130. sp. 7909.

HAB. Sierra Leone to the Gaboon (SCHLEGEL).

This truly fine species is a native of the west coast of Africa, where it is met with from Sierra Leone to the Gaboon. It is nearest allied to *Sphagolobus atratus*; and the colour of the plumage of the two species is not unlike; but the shape of the casques and the hues of the rectrices will always readily distinguish them. Although a large and conspicuous bird, nothing has been recorded about its economy or habits; and it is not very frequently seen in collections. Even among the extraordinary forms met with in this family, the present species is remarkable for the great development of its casque and its powerful bill. On account of this, as well as the peculiar shape of the casque, Bonaparte placed this species in a separate genus; and his term *Ceratogymna* I have adopted, as the bird seems to possess characters which fairly entitle it to a distinct generic rank. Occasionally individuals of this species have been brought alive to Europe and exhibited in zoological gardens. In 1871 a male was an inmate of the Regents'-Park Gardens.

Male.—Bill large, curved, black. From about its centre a high casque rises, rather flat on its anterior face, keel-shaped on top, and curves back over the head, becoming swollen towards its posterior margin; along the sides of this casque are several deep longitudinal grooves; the general colour is yellow, with the exception of the posterior terminus and lower edges of the sides next the maxilla, which are black. The bare skin around the eye and the greater part of the throat is dark blue; lowest portion of throat scarlet. Sides of the neck buffy white, central portion of feathers black. Head crested, and, together with the feathers of the body and wings, black. Two central rectrices black, remainder white. Feet black.

Bill from eye to tip 7 inches, length of casque along culmen 6, height 3, from lower edge of mandible to top of casque 5, wing $16\frac{1}{8}$, tail $14\frac{1}{8}$, tarsus $2\frac{1}{4}$.

Female.—Bill and casque yellow. The latter is a modified form of the male's, sloping forward gradually to the culmen. Head, neck, and crest ferruginous; rest of plumage like that of the male. The scarlet on the lower part of the throat is apparently wanting.

Bill 6 inches, height $2\frac{3}{4}$, wing $14\frac{1}{2}$, tail $11\frac{1}{2}$.

The specimens described and figured are in the British-Museum collection.

SPHAGOLOBUS ATRATUS.

BLACK HORNBILL

- Buceros atratus*, Temm. Pl. Col. (1830) vol. ii. pl. 558; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399. sp. 11; Hartl. Journ. für Ornith. (1854) p. 127. sp. 387; id. Ornith. West-Afr. (1857) p. 162; Schleg. Mus. Pays-B. (1862) p. 18; Bocage, Journ. Sc. Math. Lisb. (1868) vol. ii. p. 347. sp. 97; Sharpe, Ibis (1872) p. 67; Giebel, Thesaur. Ornith. (1872) p. 496; Garrod, Proc. Zool. Soc. (1873) p. 464.
- Bucorvus atratus*, Bon. Consp. Gen. Av. (1850) p. 89. sp. 3.
- Tmetoceros atratus*, Bon. Consp. Vol. Anisod. (1854) p. 12.
- Buceros poensis*, Fras. Ann. & Mag. Nat. Hist. (1855) p. 136 (♀).
- Hydrocissa atratus*, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 171. no. 472.
- Sphagolobus atratus*, Hein. Journ. für Ornith. (1864) p. 188. sp. 149; Cab. & Hein. Mus. Hein. Th. ii. p. 171.
- Buceros (Sphagolobus) atratus*, G. R. Gray, Hand-l. Birds (1870) ii. p. 131. sp. 7914.

HAB. Ashantee (PEL); Fernando Po (FRASER); Gold Coast (SCHLEGEL).

Although a close ally of *B. (Ceratogymna) elatus*, the present species is always to be distinguished from it by the shape of the casque and the colouring of the rectrices. The casques of the two birds, indeed, bear no resemblance to each other—that of the present species being swollen and rounded on its upper edge, with both the posterior and anterior ends presenting a steep angle with the culmen. The females of the two are similar in appearance; but the colouring of the tail will indicate the species. Both are natives of the west coast of Africa, their range into the interior being unknown.

Male.—Maxilla and tip of mandible black; mandible apparently reddish on its basal half; casque large, swollen and rounded on top, curving similarly to the culmen, rises from the maxilla about one third from the tip, inclines forward at a steep angle, and extends backward over the head, and returns to the base of the maxilla at a rather acute angle. Lower portion along the bill black; upper part and anterior end apparently light horn-colour. Head covered by a long crest, which with the entire plumage of the body is black glossed with dark purple. Wings and tail dark green; of the latter the terminal third of all the feathers except the median ones white. Tarsus black. Length of crest $7\frac{1}{4}$ inches, height $2\frac{1}{4}$; length of bill $6\frac{1}{8}$ inches; height of mandibles 2 inches; wings $15\frac{1}{2}$ inches; tail 14 inches. Total length without bill 26 inches.

Female.—Bill and casque much smaller than the male; the latter rises upright from the culmen, is rounded on top, increases in width towards its posterior end, and is lost in the crest of the head. Maxilla and point of mandible black; the base of the mandible is apparently red; a white spot near the base of maxilla extending onto the mandible. Casque black on sides, horn-colour on top. Head, neck, and upper part of breast brick-red. Plumage of body black with purple reflections. Wings and tail dark green; all the feathers of the latter, except the median ones, have their terminal third white. Length of bill $4\frac{1}{2}$ inches, height to top of crest $2\frac{1}{4}$; wing $13\frac{1}{8}$ inches; tail $10\frac{1}{2}$ inches. Total length 24 inches.

J. G. Keulemans lith.

Hanhart imp

LIMONOPHALUS MONTANI.

LIMONOPHALUS MONTANI.

SOOLOO HORNBILL.

Buceros montani, Oust. Bull. Hebd. Assoc. Scien. France, p. 193 (1880).

HAB. Sooloo Island, Sooloo archipelago.

This interesting species, described by my friend M. Oustalet, Aide-Naturaliste to the Museum of the Jardin des Plantes, Paris, was procured by MM. Montano and Rey, who were sent by the French Government on a scientific mission to the Philippine archipelago. M. Oustalet, in his short article on the species, compares it to *Sphagolobus atratus* from Africa. I have not seen this Sooloo specimen, which I believe is unique; but, judging from Mr. Keuleman's drawing of the type, I am led to think it represents a different genus from that containing the African species above mentioned, and, indeed, cannot be placed in any of those in this monograph. I have therefore proposed for it the term *Limonophalus**. Nothing is recorded of its habits. M. Oustalet's description, which is very short, states the total length to be about 80 centim. (32 inches), the bill black, and surmounted by a casque of 3 centim. ($1\frac{1}{4}$ inch) height, also black. Head and body black, with green reflections, tail white. The Plate will give a good idea of the appearance of this new species. A description of the generic characters is given in the text accompanying the Plates which illustrate the various genera.

* λειμὼν, smooth surface; φάλος, helmet.

J. G. Keulemans lith

M & N. Hanhart imp
15

BYCANISTES CRISTATUS.

BYCANISTES CRISTATUS.

CRESTED HORNBILL.

- Buceros cristatus*, Rüpp. Faun. Abyss. (1835) vol. i. p. 3. tab. 1; id. Syst. Ueb. Vög. Nord-Ost-Afr. (1845) p. 79. sp. 321; G. R. Gray, Gen. Birds (1849) vol. ii. p. 399; Von Heugl. Syst. Uebers. (1855) no. 457; Schleg. Mus. Pays-B. (1862) p. 16; Sclat. Proc. Zool. Soc. (1864) p. 111; Kirk, Ibis (1864) p. 326; Von Heugl. Journ. für Ornith. (1864) p. 270; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 482; Giebel, Thes. Ornith. (1872) p. 497; Elliot, Ibis (1873) p. 178; Layard, B. of S. Afr. Sharpe's ed. (1875) p. 126. sp. 18.
- Bucorvus cristatus*, Bon. Consp. Gen. Av. (1850) p. 89. sp. 4.
- Tmetoceros cristatus*, Bon. Consp. Vol. Anisod. (1854) p. 2.
- Bycanistes cristatus*, Cab. Mus. Hein. (1860) p. 172 (note); id. Von d. Deck. Reis. vol. iii. p. 38.
- Buceros (Bycanistes) cristatus*, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 331. sp. 7912.

HAB. Uganda (SPEKE); Abyssinia (RÜPPELL, HEUGLIN); Zambesi district (KIRK).

This large and handsome bird, one of the most striking of the genus to which it belongs, has as yet only been met with in eastern Africa,—probably, however, extending its range far into the interior; for Capt. Speke obtained it in Uganda. As it has not been procured anywhere on the western coast, we may feel pretty confident it does not cross the continent. This species has never received a second appellation, and its synonymy is without confusion. According to Kirk it is a local species, common on the river Shiré, goes in large flocks, and roosts regularly in the same place. The natives say that the female hatches her eggs in a hole underground, in which she is fastened by the male. On coming to roost they appear always to keep in pairs, and perch on the branches of the large trees, and generally fly away an hour before sunrise. A male that Dr. Kirk shot at, though uninjured, fell from fear, and was secured and taken on board ship. The female came for her mate every morning and evening, hovering over the vessel and calling for him to follow. Although at first he took food, after five days he became sulky, would eat nothing, and died.

Male.—Bill curved, black, yellow at base; gonys straight for two thirds its length from base, when it turns abruptly upwards, and then follows the curve of the gape to the point. An enormous casque rises a short distance from the tip of the bill, inclines forwards, and then curves backwards to near the centre of the head, returning to the base of the maxilla in a steep angle. The anterior end of this, together with the lower sides and posterior terminus, is black, the remaining portion bright yellow. Towards the base of the bill and reaching up the sides of the casque are several irregular roughened striæ, disappearing towards the ridge. Head covered with a rather long loose crest, and, together with the mantle, breast, and wings, black, with green reflections. In some adult specimens the centres of the feathers on the cheeks and sides of crest are greyish white.

Ear-coverts greyish. Back, rump, abdomen, thighs, upper and lower tail-coverts white. The tail has two central feathers greenish black for their entire length; rest greenish black for two thirds their length from base, remaining portion white.

Bill from eye to tip $5\frac{3}{4}$ inches; length of casque along curve 8, height $2\frac{3}{4}$; height of bill and casque $4\frac{3}{8}$; wing 15; tail 13.

Female.—Precisely like the male in plumage, with a less-developed crest. The bill is black, yellow at base; and the casque is more like a swelling of the culmen, rising gradually and extending over the centre of the head, and returning to the maxilla by a steep angle. Casque black like the bill, striated on the sides. Length of bill 5 inches, height to top of casque 3, wing $14\frac{1}{4}$, tail $11\frac{1}{2}$.

J.G. Keulemans lith.

BYCANISTES BUCCINATOR.

Hankart imp

BYCANISTES BUCCINATOR.

TRUMPETER HORNBILL.

Buceros buccinator, Temm. Plan. Col. vol. ii. pl. 284; Wagl. Syst. Av. (1827) sp. 12; Less. Trait. Ornith. (1831) p. 253. sp. 7; Gray, Gen. Birds, (1849) vol. ii. p. 399; Gurney, Ibis, (1861) p. 133; Schleg. Mus. Pays-B. (1862) p. 17; Kirk, Ibis, (1864) p. 326; Layard, B. of S. Afr. (1867) p. 226; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 484; Giebel, Thesaur. Ornith. (1872) p. 496; Elliot, Ibis, (1873) p. 179; Layard, B. of S. Afr. Sharpe's ed. (1875) p. 125. sp. 117.

Bucorvus buccinator, Bon. Consp. Gen. Av. (1850) p. 89. sp. 5.

Tmetoceros buccinator, Bon. Consp. Vol. Anisod. (1854) p. 2, gen. 3. sp. 7.

Bycanistes buccinator, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 171; id. von der Deck, Reis. vol. iii. p. 38.

Buceros (Bycanistes) buccinator, Gray, Hand-l. B. (1870) pt. ii. p. 130. sp. 7910.

Native name *Kakomira e none* (KIRK).

HAB. Zambesi region, valley of the Shiré (KIRK); Caffraria (SCHLEGEL).

Temminck was the first to describe this species, which is the type of Cabanis's genus *Bycanistes*, made to include all the great helmeted Hornbills of Africa with black-and-white plumage. Although remarkable for the extent of its casque, which reaches nearly to the end of the bill, it is exceeded in size by several others of the section to which it belongs. Its synonymy is very simple, only one specific name having been given to it, although it has been treated to several generic ones, Bonaparte having tried two, both of which had been employed before, viz. *Bucorvus*, which was given by Lesson to *B. abyssinicus*, and *Tmetoceros*, a term used by Cabanis for the Ground-Hornbill. In his 'Birds of the Zambesi Region,' Kirk says that this species is "rather common in the mountains and plains, frequenting the forests and living in small flocks. It returns every night to the same roosting-place, leaving in the day in search of fruit. It is very fond of a drupaceous species of *Strychnos* (resembling the *S. potatorum* of India), the fruit of which is said to be poisonous to man. Its cry is loud and harsh." Layard says, "it was procured by Mr. Ayres at Natal, and also included in Victorin's list. It has been forwarded to me by Mr. Henry Bowker, who procured it in the Transkei country. His sister (Mrs. Barber) informs me that it is a very wild and shy bird and very difficult to shoot."

Male.—Bill black, gonys straight for about one third from the base, then rising to the point of mandible in a sharp angle. A casque proceeds from the base of the maxilla, projects backwards over the head, then inclines forwards nearly to the point of the bill, and returns to the culmen at an acute angle. This is yellowish towards the upper part, remainder black, the yellow colour being widest at the posterior terminus. Head, neck, upper part of breast, flanks, and wings black

with greenish gloss. Abdomen, thighs, rump, upper and lower tail-coverts, and tips of secondaries pure white. Head covered by a rather full crest, with occasional feathers spotted with white. Tail black, glossed with green; all the rectrices except the central ones tipped with white.

Length of bill $4\frac{1}{2}$ inches, length of casque 5, height of bill and casque 3, wing 12, tail 10, tarsus 2.

Female the same as the male in colour of plumage. The bill and casque black. The latter rises from the culmen about one third its length from the tip, and inclines gradually backwards to the head. It is occasionally roughened on the sides by longitudinal creases or depressions running the entire length; this is probably the evidence of age.

Length of bill $3\frac{1}{4}$ inches, height $2\frac{1}{4}$, wing $10\frac{1}{2}$, tail $6\frac{1}{2}$, tarsus $1\frac{1}{2}$.

Specimens described and figured are in the collection of the British Museum.

J. G. Keulemans lith.

Hanhart imp

BYCANISTES SUBQUADRATUS.

BYCANISTES SUBQUADRATUS.

BLACK-AND-WHITE-CASQUE HORNBILL.

Bycanistes subquadratus, Cabanis, Journ. für Ornith. (1881) p. 350, pl. i.

HAB. Angola.

This Hornbill was described by Professor Cabanis, in the 'Journal für Ornithologie' as above quoted, from a specimen procured by Herr Otto Schütt. It resembles in its colour and markings the species described as *Buceros subcylindricus* by Dr. Sclater, from a specimen living in the Zoological Gardens, Regent's Park, and is apparently the adult male of that form. Sclater's bird was immature; and at the time his description was published the casque was small and sloped forwards to the culmen; but after some time had elapsed it had grown considerably, and nearly covered the entire length of the maxilla. This difference in the casque is shown by a comparison of the plate published in the 'Proceedings of the Zoological Society' for 1870, pl. 39, with the one published in this work. The sex of *B. subcylindricus* is not known; but it may be a female, which might account for the casque being smaller than that of Cabanis's bird, a figure of which is given in the opposite Plate. This casque presents the chief characters for separating the two birds as distinct species. In *B. subquadratus* it is high, commencing in the feathers of the crest above the head, and extends forwards in a gradual curve over two thirds the length of the maxilla, where it terminates with the anterior edge at a right angle to the culmen. The casque is black, with a white patch covering two thirds of the upper portion, with a projection in the form of a U extending downwards on the side a little over halfway to the culmen. This mark may be characteristic of the species, or it may be a peculiarity of certain individuals, as is seen in some specimens of *Anorrhinus galeritus*, and may possibly indicate age. I have noticed this yellowish-white marking upon the bills of *A. galeritus*; but as it occurred on specimens obtained in the same localities with others having all black bills, I never deemed it of any specific value. It certainly is a very peculiar mark in the bird described by Herr Cabanis (which bears every indication of being an old male), and may be in this instance a specific character; but it would be desirable to have more examples for examination before regarding *B. subquadratus* as distinct from *B. subcylindricus*. At present we can only represent the two birds under the names their describers gave them, trusting for additional information in the future to decide the question of whether they are distinct or not. Of course, should they prove to belong to the same species, the name *subquadratus* will become a synonym of that bestowed by Sclater on his species.

Casque as described above. Bill black, with a narrow band at the base yellowish. A long greenish-black crest covers the top and back of the head. Sides of the head greenish black, each feather edged with ash-grey. Ear-coverts light ash-grey. Mantle, wings, and breast metallic

greenish-black. Secondaries for half their apical length, rump, thighs, lower part of abdomen and under tail-coverts white. Median tail-feathers and a broad band near the basal end of the lateral ones metallic greenish-black; rest of tail white. Skin around eye dark brown. Iris red-brown.

Cabanis says it is larger than either *B. albotibialis* or *B. subcylindricus*.

Bill from point to base of mandible 5 inches, breadth at base $1\frac{3}{4}$; length of casque on top $4\frac{3}{4}$, height at base $2\frac{3}{4}$, height anteriorly 2; wing $10\frac{3}{4}$; tail 10.

J. H. Audubon del.

M & N Hanhart imp.

BYCANISTES SUBCYLINDRICUS.

BYCANISTES SUBCYLINDRICUS.

SCLATER'S HORNBILL.

Buceros subcylindricus, Sclat. Proc. Zool. Soc. (1870) p. 668, pl. 39; Giebel, Thesaur. Ornith. (1872) p. 503; Elliot, Ibis (1873) p. 179; Murie, Proc. Zool. Soc. (1874) p. 422.

HAB. ?

This form, described by Sclater (*l. c.*), is only known from an example living in the Zoological Gardens. It is allied to *B. cylindricus*, but differs from it in the great amount of white exhibited upon the secondaries, and the black median tail-feathers, these last being nearly all white in its relative. Its exact habitat is not known. Unfortunately the individual, after moulting its tail, has never since produced a perfect one, so that I have not seen the bird with that appendage as represented in my Plate, and also in that published in the 'Proceedings of the Zoological Society' in 1870. I consider that the principal claim this bird has to entitle it to be considered a distinct species is in the peculiar coloration of the tail-feathers; and it is to be regretted that the individual in question is unwilling or unable to present us with the proof of his claim to a distinctive rank.

Immature specimen.—Bill black; a small casque rises at base of the culmen and slopes anteriorly to the maxilla, also black. Head crested. Head, neck, back, and breast black with green reflections, the feathers of the crest being spotted with cinereous grey. Abdomen, under tail-coverts, thighs, and the greater portion of the secondaries pure white. Tail white; a broad band near the central part, and the whole of the two median feathers except their tips, black.

Total length 24 inches, wing 11·5, tail 10, bill along gape 4.

J.G. Keulemans lith

BYCANISTES CYLINDRICUS.

Hanhart imp

BYCANISTES CYLINDRICUS.

CYLINDRICAL HORNBILL.

Buceros cylindricus, Temm. Plan. Col. (1830) vol. ii. no. 521, fig. 2 (bill); Gray, Gen. Birds, (1849) vol. ii. p. 399. sp. 14; Hartl. Journ. für Ornith. (1854) p. 127. sp. 388; id. Ornith. West. Afr. (1857) p. 162; Schleg. Mus. Pays-B. (1862) p. 17; Sharpe, Ibis, (1872) p. 67; Giebel, Thes. Ornith. (1872) p. 498; Elliot, Ibis, (1873) p. 178.

Bucorvus cylindricus, Bon. Consp. Gen. Av. (1850) p. 89. sp. 6.

Tmetoceros cylindricus, Bon. Consp. Vol. Anisod. (1854) p. 2.

Bycanistes cylindricus, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 172. no. 1 (note.); Heine, Journ. für Ornith. (1860) p. 188. sp. 150.

Buceros (Bycanistes) cylindricus, Gray, Hand-l. Birds, (1870) pt. ii. p. 131. sp. 7911.

HAB. Ashantee (PEL); Fantee (SHARPE).

Bycanistes cylindricus was first described by Temminck in the 'Planches Coloriées,' and the bill figured. The species bears some resemblance to *B. cristatus*, but can easily be distinguished by the shape of its bill and colouring of its tail. Like all the members of this section, there is no difficulty with the synonymy, the bird having been always designated by the name first given. It is a native of the western coast of Africa, and apparently restricted to the Ashantee and Fantee countries. At least these are the points from which specimens have been received; but it is probable that, when the distribution is fully known, it will be found to be considerably more extensive than at present supposed. Nothing whatever, I believe, has been recorded regarding the economy or habits; and we can only conjecture that they would resemble those of the other members of the genus.

Bill and casque brownish yellow, tip of bill light yellow. The casque rises from the base of culmen, projects backwards over the head, and then inclines forward for two thirds the length of the bill, terminating at a right angle to it. The casque is divided into two portions by a deep furrow running from the posterior to the anterior face, above which it is rounded. The sides have several deep irregular grooves. The mandible has also four or five diagonal furrows at its base. Head and occiput covered by a long greenish black crest. Neck, back, breast, wings, and thighs black with bluish green reflections. Apical portion of secondaries and primaries for one third their length, abdomen, upper and under tail-coverts, and lower part of rump pure white. Tail white, with a very broad central black band with greenish reflections. Tarsi and feet black.

Total length 31 inches, bill at gape 5, casque on top $4\frac{1}{2}$, height over nostril $2\frac{1}{4}$, height of bill and casque at base $3\frac{1}{2}$, wing 13, tail $10\frac{1}{4}$, tarsus 2.

The *female* differs from the male only in the colour of bill and shape of casque. Both of these are black. The casque rises upright from the culmen about one third from the point of the maxilla, and extends to and a little over the head, and returns to the base of the maxilla by an abrupt angle. It is sharp at its anterior face, but swells outwards towards the posterior termination.

Length of bill $3\frac{3}{4}$ inches, height $2\frac{1}{4}$; length of casque $4\frac{1}{4}$, height $2\frac{1}{4}$; wing $10\frac{1}{2}$, tail $6\frac{1}{2}$, tarsus $1\frac{1}{2}$.

The specimens described are in the British Museum.

BYCANISTES ALBOTIBIALIS.

LOANGO WHITE-THIGHED HORNBILL.

Buceros albotibialis, Cab. & Reich. Journ. für Ornith. (1877) p. 103.

HAB. Loango (CAB. & REICH.).

This species was described by Herren Cabanis and Reichenow as above. It is singularly like *B. cylindricus*, but differs in the shape of the casque, and in having the thighs pure white instead of black. A drawing of the type, now in the Berlin Museum, was forwarded to me by Dr. Cabanis, and is reproduced in the smaller figure of the plate. Its casque differs in shape from that of an adult specimen of the same species in the British Museum, represented in the chief figure in the illustration: in this last the casque is smooth, and projects forward to a point over the culmen; the bill also has the furrows fewer and slighter. This last specimen is stated to have come from the Gaboon. The plumage of *B. albotibialis* is black, with green reflections only, and without any of the bluish tints seen on the feathers of its relative. Both species are very rare in collections, but few museums possessing examples; and but little is known of their economy or habits; in fact, as regards the present one we know absolutely nothing.

Male. Bill black, with the exception of apical third, which is yellow. A casque rises from base of the maxilla, inclines slightly backwards over the head, then extends by a gradual curve forwards over the point of the maxilla, and returns by a very acute angle to the culmen, which it reaches at about the commencement of the black colour. The posterior termination and sides of the casque along the maxilla are black. Head covered with a loose, rather full crest, and, together with the neck, back, and breast, it is black, with green reflections. Wings same colour as the back, with the exception of two thirds of the secondaries and one third of the primaries (from their tips), which are pure white. Rump, abdomen, thighs, upper and under tail-coverts pure white. Tail also pure white, crossed near the centre by a very broad black band glossed with green. Tarsi and feet black.

Length of bill $5\frac{1}{4}$ inches, height $3\frac{5}{8}$; length of casque along curve $5\frac{1}{2}$, height $2\frac{1}{8}$; wing $13\frac{1}{2}$, tail $11\frac{1}{2}$, tarsus 2.

J.G. Keulemans bth.

PHOLIDOPHALUS FISTULATOR.

Hanhart imp
29

PHOLIDOPHALUS FISTULATOR.

CASSIN'S HORNBILL.

Buceros fistulator, Cass. Proc. Acad. Nat. Scien. Phil. (1850) p. 68; Hartl. Journ. für Ornith. (1854) p. 127. sp. 389; J. & E. Verr. Rev. Mag. Zool. (1855) p. 175; Hartl. Ornith. West-Afr. (1857) p. 162; Hein. Journ. für Ornith. (1860) p. 188. sp. 151; Schleg. Mus. Pays-B. (1862) p. 16; Sharpe, Ibis (1872) p. 67; Giebel, Thesaur. Ornith. (1872) p. 498; Bocage, Proc. Zool. Soc. (1873) p. 702; Elliot, Ibis (1873) p. 179.

Buceros leucostigma, Temm. Mus. Lugd.

Tmetoceros fistulator, Bon. Consp. Vol. Anisod. (1854) p. 2.

Buceros (Bycanistes) fistulator, Gray, Hand-l. Birds (1870) ii. p. 131. sp. 7913.

HAB. Fantee (SHARPE); Senegal to the Gaboon (VERREAUX).

But little has been recorded of the habits and economy of this species. It was first described by Mr. Cassin, in the 'Proceedings of the Philadelphia Academy of Sciences,' in 1850, from an immature specimen. In the 'Revue et Magasin de Zoologie' MM. E. and J. Verreaux give the following short account of this bird as observed by them in Africa:—"We notice that in this species the male is always distinguished, not only by its greater size, but also by the bill, which is larger and heavier, and which varies considerably, before attaining its full development, as is the case in most of the species. The present bird, which is quite abundant at the Gaboon, is also found upon the greater part of the coast as far as Senegal, wherever the forests afford it sufficient shelter and abundant nourishment, which last consists of insects, larvæ, often even berries. It does not refuse young birds, which it seeks in their nests. Generally it is found in pairs. The cry is strong, heard throughout the morning and evening, and just before rain. The females deposit their eggs in the holes of trees to the number of two; we do not know their colour, our hunter having only found the young in October, which, with the exception of the size of the bill, resembled their parents in every particular. During the heat of the day these birds perch in the thickets, and remain quiet; but they pass the night in the holes which they have chosen, and to which they always return."

The *P. fistulator*, and the bird which I have named *P. sharpii*, by not possessing upright casques like the other African species to which they are related, apparently require a distinct generic term; and therefore I propose for them that of *Pholidophalus**, or scaly helmet, their low casques seeming to be composed of flattened scales overlapping each other.

Bill has the culmen covered on its basal half by a low casque, deeply ridged transversely. Head, neck, back, and wings black, with a greenish lustre. Underparts white. Secondaries black, except for about an inch and a half from their ends, which are white. Tail black, the lateral feathers with a narrow line at their bases, and their apical third white.

Total length $21\frac{1}{2}$ inches, wing 11, tail $9\frac{1}{4}$; bill 4, height at base 2 inches.

I am indebted to Captain Shelley for the male figured in the Plate, loaned from his collection, and to Mr. R. B. Sharpe for the female.

* *φολῖς*, scale; *φάλος*, helmet.

J G Keulemans lith

Hanhart imp

PHOLIDOPHILUS SHARPII.

PHOLIDOPHALUS SHARPII.

SHARPE'S HORNBILL.

Buceros fistulator, Cassin, Proc. Acad. Nat. Scien. (1859) p. 139. sp. 161; Sharpe, Proc. Zool. Soc. (1871) p. 139.

Buceros sharpii, Elliot, Ibis, (1873) p. 177; Barb. du Bocage, Proc. Zool. Soc. (1873) p. 702; id. Ornith. Ang. (1878) p. 114.

HAB. Angola (HAMILTON); rivers Camma, Muni, and St. Paul's (DuCHAILLU).

I described this bird from a fine adult male from Angola in the collection of Mr. Sharpe, which is now in the British Museum. It is nearest allied to the *B. fistulator* of Cassin, but may be easily distinguished by the white lateral rectrices—those of its relative being black, merely tipped with white. Mr. Cassin evidently had this species before him, in a collection sent to Philadelphia by DuChaillu, a list of which was published in the 'Proceedings of the Academy of Natural Sciences' for 1859; but he imagined it to be the adult of *B. fistulator* instead of a new species, and so described it. I have seen several specimens of this species in various museums; and in nearly every instance it was confounded with *B. fistulator*. Of the economy and habits nothing has been recorded; but this species will undoubtedly in both resemble its near ally *B. fistulator*.

Bill without casque; mandible swollen into a prominent ridge, broad, and corrugated at the base, becoming narrow anteriorly, and gradually falling into the outline of the culmen; the base, which ends abruptly about half an inch above the head, has four rather deep rounded plaits. The mandible is rugged, and furrowed with diagonal grooves nearly to the end, with the cutting portion serrated. The culmen and tips of bill yellowish horn-colour; a broad stripe for about two thirds the length of the maxilla, and nearly all the mandible (save the tip above mentioned), black. Base of maxilla beneath the nostril and spot on base of mandible yellowish white. Head covered with a long crest; feathers of cheeks and throat also long and loose; all these parts, together with the back, black with a bright greenish lustre. Primaries black. Secondaries, with the exception of the three innermost ones, white for three fourths their entire length, greenish black at base; the innermost ones same colour as the back. Under wing-coverts, lower part of breast, under tail-coverts, rump, and legs pure white. Tail, the two median feathers black for their entire length; the ones next to the median on either side white for half their length from the tip, remaining portion like the central ones; the lateral feathers pure white for their entire length, with the exception of the outermost ones, which have the basal half of the outer webs black.

Total length 21 inches, wing 11, tail $9\frac{1}{4}$.

My description and figure were taken from the type in the British Museum.

J Smit lith

Hanbart imp.

PHOLIDOPHALUS CASUARINUS.

PHOLIDOPHALUS CASUARINUS.

GRAY'S HORNBILL.

Buceros casuarinus, G. R. Gray, Ann. & Mag. Nat. Hist. (1871) vol. viii. pp. 437, 438, pl. xvii.; Barb. du Boc.
Proc. Zool. Soc. (1873) p. 703.

HAB. —?

This form, described by Gray from a head in the British Museum, although bearing evidence of belonging to a fully adult individual, appears to have a stunted abnormal casque, which, perhaps, may have been injured in some way when the bird was young. The casque, and mandible at its base, are deeply grooved diagonally; and the cutting-edges of both maxilla and mandible are broken, which is only seen in adult birds. It is impossible to say whether or not the specimen belongs to one of the known species or is a new form, as but a small bunch of feathers remains attached to the head. It is evidently closely allied to the subgroup which contains *P. fistulator* and *P. sharpii*, and may possibly be one of those species. Mr. Gray's description of this head is as follows:—"Bill broad at base, laterally compressed at the tip; casque elevated posteriorly and extending somewhat backwards over the eyes, rather compressed along the culmen, which is flat and grooved along the middle for two thirds of its length; the sides of the casque shelving to the nasal channel, and furnished with six deep oblique grooves; the sides below the former are comparatively smooth, and with three apparent scales near the eyes; the nostrils are large, and deeply imbedded in a broad channel, which runs along the sides of the maxilla for about two thirds of its length, in which they are situated at its base; the mandible has the gonys long and curved to the tip, the sides are furnished with four very obliquely placed grooves, advancing towards each other beneath the gonys; the margins of both mandibles are dentated in the middle. The length from the upper part of the base of the casque to the tip of the maxilla is five inches and three lines."

J. G. Keulemans lith

M. N. Hanhart imp

RHYTIDOCEROS UNDULATUS.

RHYTIDOCEROS UNDULATUS.

WREATHED HORNBILL.

Le Calao à casque festonné, Levaill. Ois. Rares (1801) pls. 20, 21.

Le calao javan, Levaill. Ois. Rares (1801) pl. 22; id. Hist. Nat. Ois. Afr. (1806) p. 124, pl. 239; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 595.

Buceros javanensis, Wilkes?, Ency. Lond. (1808) vol. iii. p. 479.

Buceros undulatus, Shaw, Gen. Zool. (1811) vol. viii. p. 26; Less. Trait. Ornith. (1831) p. 255. sp. 12; Horsf. Trans. Linn. Soc. vol. xiii. p. 175; Vig. App. Mem. Raffl. p. 666.

Buceros javanicus, Shaw, Gen. Zool. (1811) vol. viii. p. 28; Vieill. Nouv. Dict. Hist. Nat. vol. iv. (1816) p. 595; id. Ency. Méth. p. 401, t. 4, juv.

Buceros niger, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 592; id. Ency. Méth. p. 400. sp. 21.

Buceros javanus, Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 215.

Buceros annulatus, Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 210; Drapiez, Dict. Class. Hist. Nat. (1828) vol. iii. p. 32.

Buceros plicatus, Temm. Plan. Col. (1824) vol. ii. sp. 8 (text, partim); Ersch u. Grub. Ency. (1824) p. 287 (partim); Wagl. Syst. Av. (1827) sp. 8 (partim); Müll. & Schleg. Verh. Gesch. Ned. Ind. (1839-44) pp. 24, 30 (Aves); G. R. Gray, Gen. Birds (1847) vol. ii. p. 399; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 45. sp. 191; Hartl. Journ. für Ornith. (1859) p. 289. sp. 30; Bernstein, Journ. für Ornith. (1861) p. 113; Schleg. Mus. P.-Bas (1862) p. 2; Eyton, Ost. Av. (1867) p. 60?; Sclat. Rev. List Vert. Anim. (1872) p. 170, fig. 20; Garrod, Proc. Zool. Soc. (1873) p. 464 (nec Lath.).

Buceros ruficollis, Blyth (nec Vieill.), Journ. Asiat. Soc. Beng. (1843) vol. xii. p. 176.

Buceros pucoran, Blyth, Journ. Asiat. Soc. Beng. vol. xii. pt. ii. p. 990.

Buceros pasuran, Blyth, Cat. Birds Mus. Asiat. Soc. Beng. (1849) p. 45.

Rhyticeros plicatus, Reichenb. Syst. Av. (1849) pl. 50; Bon. Consp. Vol. Anisod. (1854) p. 3; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-58) vol. ii. p. 598. sp. 882; Moore, Proc. Zool. Soc. (1859) p. 451; Godwin-Austen, Journ. Asiat. Soc. Beng. (1870) pt. ii. p. 96, ♀; Jerd. Ibis (1872) p. 5 (nec Lath.).

Rhytidoceros obscurus, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 172; Salvad. Ucc. Born. (1874) p. 85 (nec Gmel.).

Buceros (Rhyticeros) obscurus, Gray (nec Gmel.) Hand-l. Birds (1870) pt. ii. p. 129.

Aceros, sp. indet.? Godwin-Austen, Journ. Asiat. Soc. Beng. (1870) pt. ii. p. 96. sp. 1466.

Aceros plicatus, Blyth (nec Lath.) Cat. Mamm. & Birds Burma (1875) p. 69. sp. 71.

HAB. Malayan peninsula, northward to Sylhet, Sumatra, Java (BLYTH); North Cachar (GODWIN-AUSTEN); Borneo.

Very great confusion exists in the synonymy of this species; and it has been the recipient of a large number of names. It has generally been known as the *Buceros plicatus* of Latham; but this title was bestowed upon the bird from Ceram and the neighbouring islands, called by ornithologists *Buceros ruficollis*. In my article on this latter species, this question of synonymy

will be found fully worked out. After Latham's, the next name that has priority is that of *undulatus*, Shaw, which is the one necessary to adopt for the species.

A very curious character of this bird is its casque formed of a succession of overlapping scale-like ridges: these grow from the base, and push forward those in front, the lengthening of the casque beyond a certain limit being prevented by the anterior ridges scaling off. In the young these scale-like protuberances are larger, and thinner in substance, growing narrower and closer by degrees.

According to Horsfield this bird is very abundant in the forests of Blambaugan, in the eastern end of Java, where it is called *barong chimburuan* (or "jealous bird") on account of the watchfulness displayed by the male during the period the female is immured in the hollow of a tree performing the duties of incubation; for, as stated by the natives, should another male approach the nest during the husband's absence, he leaves the unfortunate wife to perish with hunger, as she is unable to get out of her walled prison.

Dr. Cantor states that "the male has the bill (yellowish) white; iris pale crimson; gular pouch rich gamboge yellow, with two transverse black bars; feet blackish. And the female has the iris narrow, golden round the pupil, the rest golden vandyke; eyelids brick-colour; pouch dirty azure, with two transverse black bars; feet blackish grey. The young male has the iris mother-of-pearl colour; bill yellow at the point, and bluish green at the base; space around the eyes, and pouch, yellow, with the transverse black bars indistinct; feet bluish black."

"This Hornbill," says Jerdon, "has been killed in North Cachar by Major Godwin-Austen. It has not hitherto been recorded from further north than Arakan. He gives the dimensions as follows:—length 3 feet 2 inches; wing $18\frac{1}{4}$ inches; extent 5 feet 2 inches; tail 12 inches; bill $6\frac{1}{4}$ inches; depth of bill 3 inches."

This measurement gives the length of the species greater than I have been able to find in the dried skins; and it seems to me that the specimen described above must have been unusually large, although, with the exception of the length, it does not differ much from the average.

Male.—Bill with a roughened casque-like protuberance projecting from the base of the culmen for nearly half its length, in the form of large scales, indented by several transverse grooves; base of maxilla and mandible transversely grooved; bill white, with its base deep chestnut-red, this being more extensive on the mandible, where it reaches on the gonys nearly half its length. The scaly protuberance on the culmen also chestnut, becoming yellowish towards its anterior terminus. Bare skin around the eye pinkish; that of the throat rich yellow, with a conspicuous broken ring of dark blue on its lower part. Sides of the head yellowish; lower portion of neck to the breast pure white. Top of head and occiput dark chestnut. Back of neck and entire plumage of the body black, with green reflections. Tail pure white. Total length $29\frac{1}{2}$ inches; wing $18\frac{1}{2}$; tail 13; bill $6\frac{1}{2}$, height at base 3; tarsus $2\frac{1}{4}$.

Female.—The naked skin of throat and around the eyes bright blue. Entire plumage black. Tail white. Total length 28 inches, wing $16\frac{1}{2}$, tail 11, bill $6\frac{1}{2}$, tarsus 2.

J.G. Keulemans lith.

Hanhart imp.
34

RHYTIDOCEROS SUBRUFICOLLIS.

RHYTIDOCEROS SUBRUFICOLLIS.

BLYTH'S WREATHED HORNBILL.

- Buceros subruficollis*, Blyth, Journ. Asiat. Soc. Beng. (1843) vol. xii. p. 177, (1849) p. 320; Barbe, Journ. Asiat. Soc. Beng. vol. x. p. 922.
- Buceros plicatus*, Blyth, Journ. Asiat. Soc. Beng. vol. xii. p. 991, vol. xvi. p. 998; id. Cat. Birds Mus. Asiat. Soc. (1849) p. 45 (nec Forster).
- Rhyticeros subruficollis*, Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 600. no. 883; Hume, Nest & Eggs Ind. Birds, pt. i. (1873) p. 115; id. Str. Feath. vol. ii. (1874) p. 470; Ramsay, Ibis (1877) p. 455.
- Rhytidoceros subruficollis*, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 172. no. 474; Tweeddale, Ibis (1877) p. 295.
- Buceros (Rhyticeros) pusaran*, Tickell, Ibis (1864) p. 180.
- Buceros (Rhyticeros) subruficollis*, Gray, Hand-l. B. (1870) pt. ii. p. 129. sp. 7886.
- Aceros subruficollis*, Blyth, Cat. Mamm. & Birds Burma, (1875) p. 69. sp. 72.

HAB. Arakan, Tenasserim Provinces (BLYTH); Southern Shan States of Siam (TICKELL); Lawas river, N.W. Borneo (USSHER).

This bird was separated by Blyth from the well-known *R. plicatus*, Forst., on account of the base of the mandible being smooth and without the transverse ridges so conspicuous in Forster's species. In *R. subruficollis* the mandible never possesses at any age of the bird this roughened surface. As this difference appears to be a permanent one, it would seem to be sufficient to give the two birds distinct specific rank. Their geographical distribution also is not the same, the present species ranging rather more to the northward than its ally, and never going into the Malayan peninsula. *R. subruficollis* may also be distinguished by never having the black transverse bar on the gular skin of either sex, that part being yellow in the male, blue in the female. Lieut. Wardlaw Ramsay observed this difference between the species to have been constant in a large series of examples collected by him in the district of Tonghoo, Tenasserim.

In regard to the curious scaly casque possessed by this bird and its allies, Mr. Blyth found, after examining numerous specimens, that in no instance did the number of transverse ridges exceed seven; and he deemed it obviously apparent that they scaled off anteriorly as they were pushed forward by the growth of those at the base of the maxilla.

Tickell says, speaking of this bird under the name of *B. pusaran*, that it is very numerous in the Tenasserim Provinces and in the inland forests of Arakan. Its voice is a short gruff croak like "kukkuk," repeated at intervals. The flight is slow and regular, capable of being continued for a great distance; and the noise made by the air rushing through the feathers of the wings is so

great that it can be heard half a mile distant. Its food consists of various fruits; and it is particularly fond of the figs of the Banian and other trees of the *Ficus* family. Except when feeding, it is very shy and wary, but at such times will frequently return to the same tree even after having been shot at. The female lays two or three coarse white eggs in the hollow of a tall tree, and, according to the Karens, is not immured while sitting. The egg is 2" by 1½" in size. When feeding, this species is usually silent, and on the lofty fruit-trees mixes indiscriminately with numerous species of fruit-eating pigeons (*Carpophaga*, *Treron*, &c.), as well as monkeys (*Hylobates lar*, *Semnopithecus phayrei*, *Inuus nemestrinus*, &c.), which are similarly occupied, and form a picture so singular and strange as not easily to be forgotten when once seen. Tickell relates that on one occasion he was on his way down the Hounghthran river, "a clear pretty stream shaded by lofty timber, eddying in deep pools, under high gravelly banks, breaking into foam and tumbling over boulders of sandstone, or rippling along shallow beds of clean pebbles and silvery sand. To the last-named spots, just before or during the short twilights of a tropical evening, these Hornbills used to resort in great numbers, allowing my boat to approach pretty near. As it glided down the stream I could thus watch them on the little sand flats, hopping freely enough along the ground, and delving their beaks in, as if searching for worms or molluscs, while some stood up to their bellies in the water, apparently much enjoying their bath. As the dusk gathered over the river, I remarked them resorting to roost on the loftiest trees fringing its course. The Karens, who live in these virgin forests, say that between the 'Yowng-yowng' (*B. bicornis*) and the 'Owkhennet' (the present species) there is always open war; and, in truth, I do not remember to have remarked the two species anywhere together." He had observed parties of five or six of this species flying over the sea at a great height, and had watched them until they "melted from sight into the horizon, as if they had finally left the shore." As the species is not found on the western shores of the Bay of Bengal, he could not imagine where these excursions ended. I have lately seen, however, in Mr. Sharpe's possession, several specimens of this Hornbill, sent from Borneo by Governor Ussher, which proves that the species inhabits that island; and it is possible it may also dwell upon some of the intervening ones between Borneo and Siam, where it is common. The individuals seen by Tickell may have been on their way to some of the islands in the Bay of Bengal; and long flights are very possibly frequently undertaken by the species.

Lieut. Ramsay says that the Burmese have many legends about the Hornbills, and that the name is continually occurring in their poetry and plays, whilst the female Hornbill is regarded as a model of virtue. They also imagine that the plaster with which the bird closes the entrance to its nest is made of earth brought from the four quarters of the globe, and mixed with a gum extracted from trees; and this composition is much esteemed for its medicinal properties.

Male.—Bill curved, pointed, greenish white; base of bill and ribbed part of maxilla vinous brown; cutting-edges in adults much broken. About a third of the culmen from its base is covered by five or more deep plaits. This casque-like addition to the culmen is cream-colour, with the furrows earthy. Bare skin of face and rim around the eye bright vinous brown; eyelids pale greenish; skin of the throat bright lemon-yellow, without any black bars as in *R. undulatus*.

Top and sides of head, neck, and upper part of breast rich yellow. Centre of head, occiput, and middle of neck behind deep reddish chestnut. Rest of plumage black, with green reflections. Tail pure white. Legs black.

Total length 31 inches, wing 16, tail 11, bill along gape 7, height at base $3\frac{1}{2}$, tarsus $2\frac{1}{4}$.

Female.—Entire plumage of body black, with green reflections; tail white. The casque-like protuberance on the culmen is smaller than in the male, with the scales broader and fewer. Bare skin of throat turquoise-blue.

Total length 27 inches, wing 14, tail 10; bill along gape $5\frac{1}{4}$, height at base $2\frac{1}{4}$.

The specimens figured are in my own collection.

J. S. Keulemans hth

RHYTIDOCEROS PLICATUS.

Hanhart imp
28

RHYTIDOCEROS PLICATUS.

PAPUAN WREATHED HORNBILL.

Buceros plicatus, Forst. Ind. Zool. p. 40 (1781), ex Dampier; Lath. Ind. Orn. (1790) vol. i. p. 146; Shaw, Gen. Zool. (1811) vol. viii. p. 38; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 593; Dumont. Dict. Sc. Nat. (1817) vol. vi. p. 210; Vieill. Ency. Méth. (1823) vol. i. p. 306.

Wreathed Hornbill, Lath. Gen. Syn. (1787) Supp. p. 71.

Buceros obscurus, Gmel. Syst. Nat. (1788) vol. i. p. 362; Giebel, Thesaur. (1872) 501 (partim).

Buceros ruficollis, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 600; Temm. Plan. Col. (1830) pl. 557; Müll. & Schleg. Verh. Gesch. Ned. Ind. (1839-44) pp. 24, 30. sp. ix.; Gray, Gen. Birds (1849) vol. ii. p. 399. sp. 19; id. Proc. Zool. Soc. (1860) p. 356, (1861) p. 436; Bernst. Journ. für Ornith. (1861) p. 118; Schleg. Mus. Pays-B. (1862) p. 3; Wall. Intell. Observ. (1863) vol. iii. p. 314; Sclat. Proc. Zool. Soc. (1869) p. 122; Giebel, Thes. Ornith. (1872) p. 502.

Calao ruficollis, Bon. Consp. Gen. Av. (1850) p. 90. sp. 4.

Rhyticeros ruficollis, Bon. Consp. Vol. Anisod. (1854) p. 3.

Calao papuensis, Rosenb. Natuurk. Tijdsch. Nederl. Ind. (1863) p. 229. sp. 57; id. Journ. für Ornith. (1864) p. 117. sp. 57.

Buceros (Rhyticeros) ruficollis, Gray, Hand-l. B. (1870) pt. ii. p. 129.

Rhytidoceros ruficollis, Salvad. Ann. Mus. Civ. de Stor. Nat. Genova, (1876-77) vol. ix. p. 19.

Rhytidoceros plicatus, Salv. Ann. Mus. Civ. St. Nat. Genova. (1877) vol. x.

Rhytidoceros plicatus var. *ruficollis*, Salv. Ann. Mus. Civ. St. Nat. Genova. (1877) vol. x.

Massouahou-Boro in Guabé (QUOY & GAIM.); *Wama* in Lobo (S. MÜLLER); *Oedooi* in Andai (VON ROSENBERG.).

HAB. Solomon Islands (BENNETT); New Guinea (BECCARI); Waigiou, Mysol (WALLACE); Halmahera, Morty, Batchian, Ceram (SCHLEGEL); Amboina (TEMM.); Salwatti (BRUIJN); Ravak, Grebèh, Batanta (BECCARI).

The first Latin name given to this species is in the "Faunula Indica" of Forster's 'Indische Zoologie,' where the appellation of *plicatus* is conferred upon the Hornbill the head of which is represented in Dampier's plate, published in the third volume of his 'Voyages,' page 231. This bird was said to have come from Ceram; and, from the fact that the bill had no lateral ridges upon either maxilla or mandible, it is without doubt the present species, called afterwards *ruficollis* by Vieillot, and generally known to ornithologists by the latter name. The other prominent synonyms of the species are *obscurus*, Gmelin, and *Wreathed Hornbill* or *Buceros plicatus*, Latham.

The "Wreathed Hornbill" of Latham (*l. c.*) was founded upon a drawing of a bill given in Ray's translation of Willughby's 'Ornithology,' and which, having no lateral grooves, must have belonged to the present species. To this Gmelin gave the name of *obscurus* (*l. c.*); and two years afterwards Latham also gave to his Wreathed Hornbill a Latin title, and called it in his 'Index

Ornithologicus' (*l. c.*) *Buceros plicatus*. Both of these Latin names, however, must become synonyms of *plicatus* given in Forster's work above mentioned.

This bird is a native of several of the islands of the Eastern archipelago, Ceram being apparently its western limit; but it goes the furthest eastward of all the members of the family, having been obtained by Mr. Bennett in the Solomon Islands. Of its habits and economy we know very little, the few remarks given below, published by Mr. Wallace in the 'Intellectual Observer,' being all that I have met with upon this subject.

Wallace states that "on one occasion he shot a Moluccan Hornbill (*Buceros ruficollis*), to the roof of whose beak was sticking a large lump of bees' comb, showing that he had been making a meal off bees' brood and honey. They are also positively declared by the natives to eat eggs and young birds occasionally. Yet, notwithstanding this varied bill of fare, there seems little doubt but that periodical scarcity of food is the most efficient check to their increase, as shown by the fact that the larger kinds are always scarce, even where the smaller ones abound."

Dr. Beccari says that individuals of this species obtained in New Guinea have the neck of a much lighter colour than those from the islands of Ceram and Amboina; and D'Albertis records in his report on the ornithology of the Fly River that it was common all over the country. He saw some bills in the natives' huts that were much smaller than the usual style, which made him suspect that there was a second species; but he could not procure the bird itself.

Bill long and curved, yellowish white, red at the base. At the base of the culmen, instead of a casque is a moderate elevation divided into several deep prominent ridges of the same colour as the bill, and also red at the base. Bare skin of throat and round the eyes pale blue. Head, neck, and upper part of breast reddish orange. Entire rest of body and wings black with a greenish tinge. Tail pure white. Iris salmon-colour. In old birds the cutting edges of the bill are frequently much broken towards the point. Feet black.

Total length 29 inches, wing 17, tail 11; bill along gape 7, height at base 3; tarsus 2.

The female has the body jet-black, with the bare skin blue; iris chestnut; tail white; feet black.

The specimen from which my figure and description were taken is in my own collection.

G. Keulemans lith.

RHYTIDOCEROS NARCONDAMI

Hanhart imp.

RHYTIDOCEROS NARCONDAMI.

NARCONDAM ISLAND WREATHED HORNBILL.

Rhyticeros narcondami, Hume, Str. Feath. vol. i. (1873) p. 411, vol. ii. (1874) pp. 108, 110.

HAB. Island of Narcondam, Bay of Bengal.

This species was described by Mr. Hume (*l. c.*) from specimens procured in his expedition to the islands in the Bay of Bengal. The colour of its plumage is precisely like that of *R. plicatus*, Forst.; but in all its dimensions the present is a much smaller bird. Mr. Hume states (in litt.) that "I entertain no doubt of the distinctness of this species. I can tell you nothing further about the bird" (beyond what is recorded in 'Stray Feathers'); "scarcely any one else has ever landed on the island; all I can say is that they were in great numbers, keeping everywhere to the tops of the trees, from the water's edge to the very summit of the island, and that most certainly nothing like them occurs anywhere in any of the Nicobars, Andamans, Cocos, Preparis, Barren Island, Pegu, or Tenasserim, nor, as far as we yet know, in the Mergui archipelago, where *R. undulatus* is very common; but we have not yet explored this archipelago so as to be able to assert that *narcondami* occurs nowhere on any of the other odd thousand islands.

"I have no specimen of *ruficollis* (*plicatus*, Forst.); but with both *undulatus* and *subruficollis* before me certain differences immediately strike the eye. In the first place *narcondami* is not one third the bulk of *subruficollis*: an ordinary male of the latter weighs 3.75 lb.; our male *narcondami*, an adult, weighed 1 lb. 3 oz. There is much less bare gular skin; and this ends square across the throat, and not in a curve as in the two species above named. The head and neck in the male are a bright chestnut, the upper part of the throat only being buffy, changing to chestnut. The bill is straighter and not so much curved as in *undulatus* even, and *à fortiori* not so much as in *subruficollis*. The casque projects proportionally more on either side of the mandible than in either of those two species. The metallic reflections of the black of the mantle are pure green; in both the other species there is a decided blue tinge in some lights, though not in others, whereas *narcondami* in any light that does not show it pure green is a bronze-green."

Narcondam Island, the abode of this species, is almost inaccessible; for, according to Mr. Hume, not only no vessels ever visit it, but for at least 340 days out of the year it is impossible to land on account of the heavy surf that breaks around its shores. For this reason it is most probable that *R. narcondami* will always be a great rarity in collections. Two specimens, I believe, were all that were obtained; and these are now in Mr. Hume's magnificent museum.

In both sexes the irides are pale red, legs and feet brown; bill pale horny yellow, brownish red at base of both mandibles; posterior plate of casque reddish brown, margin horny yellow,

succeeding ridges horny yellow, furrows blackish brown; bare orbital skin pale smalt-blue; gular skin white, tinged with greenish blue.

"In the female the entire plumage (except the tail, which is pure white) is jet-black, glossed on the wings and back with dull metallic green; the male differs in having the front of the neck below the bare gular patch a somewhat pale rufous buff, while the whole of the rest of the head and neck are bright chestnut-coloured" (*Hume*, loc. cit.).

The following are the measurements as given by Mr. Hume:—

"*Male*.—Length 26 inches, expanse 41.5; tail from vent 8.25; wing 12; wings when closed reach to within 4.5 of end of tail; tarsus 2.05; middle toe to root of claw 2.5; claw only, straight from point to base, 0.7; bill from nostril to point 4.5, bill from posterior margin of casque to point, straight, 5.0; length of casque only, 2.67; greatest height of upper mandible including casque 1.4, of casque only 0.6. Weight 1 lb. 3 oz.

"*Female*.—Length 24 inches, expanse 37.75; tail from vent 8.0; wing 10.75; tarsus 1.8; middle toe to root of claw 1.4; claw only, straight from point to base, 0.55; bill from nostril to point 3.6; bill from posterior margin of casque only, 2.6; greatest height of upper mandible including casque 1.25, of casque only 0.6. Weight 1 lb."

J.E. Keulemans lith.

ANORRHINUS COMATUS.

ANORRHINUS COMATUS.

LONG-CRESTED HORNBILL.

Buceros comatus, Raffles, Trans. Linn. Soc. vol. xiii. p. 339; Temm. Plan. Col. (1830) vol. ii. (text); G. R. Gray, Gen. Birds (1849) vol. ii. p. 399. sp. 16; Blyth, Journ. Asiat. Soc. Beng. (1847) p. 996, pl. xlv. fig. 2, and Cat. B. Mus. Asiat. Soc. (1849) p. 44. sp. 186; Schleg. Mus. Pays-B. (1862) p. 8; Giebel, Thesaur. Ornith. (1872) p. 497; Mull. & Schleg. Verh. Geschied. Neder. Ind. (1839-1844) pp. 23, 29, sp. vii.; Hay, Madr. Journ. vol. xiii. p. 149.

Buceros lugubris, Begbie, Malay. Penin. (1834) p. 513; id. Ann. & Mag. Nat. Hist. (1846) p. 405.

Berenicornis comatus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 1; Horsf. & Moore, Cat. B. E.-Ind. Co. (1856-58) vol. ii. p. 594. sp. 875; Moore, Proc. Zool. Soc. (1859) p. 450; Bon. Consp. Vol. Anisod. (1854) p. 2.

Buceros (Berenicornis) comatus, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7882.

HAB. Malayan peninsula, Sumatra (BLYTH).

This very distinct species was first described by Raffles in the 'Linnean Transactions' for 1822, from a young male, which differs from the adult in the brown colour of its back, wings, and tail, and dusky hue of bill and casque. It is a very peculiar-looking bird, and is the type of Bonaparte's genus *Berenicornis*. As I do not consider this species to be generically distinct from *A. galeritus*, which is the type of Reichenbach's genus *Anorrhinus*, I have placed Bonaparte's term among the synonyms. This genus includes six species, according to my views, two of which are natives of Africa, and four of the Oriental region. But a single other name besides the one it bears has been given to this Hornbill, viz. *B. lugubris* of Begbie; and therefore its synonymy is happily freed from all complication. The last-named author says:—"This bird is melancholy in its disposition, but withal voracious in its habits. I had one in confinement for some time, feeding it principally on plantains, which it devoured greedily, and never attempted to escape from its perch. I destroyed it on account of the filth and stench it generated." This short account is all that I have met with regarding the habits of this species—not sufficiently attractive to make further investigation desirable. This species is confined to the Malayan peninsula and the island of Sumatra.

Male.—A low casque, being more like the elevation of the culmen, commences on the maxilla, about one third of its length from the point, extends backwards to the base, and disappears in the loose feathers of the crest. Head, neck, crest, breast, abdomen, tail, end of primaries and secondaries pure white. Wings, back, and thighs black. Length of bill 6 inches, height $2\frac{1}{2}$; length of crest on bill $4\frac{1}{4}$; wing $15\frac{3}{4}$; tail 20; tarsus $2\frac{1}{4}$.

Female.—Crest, end of secondaries and primaries, and the tail pure white; all the rest black. Length of bill $5\frac{5}{8}$ inches; wing $15\frac{1}{2}$; tail $19\frac{1}{2}$; tarsus 2.

J.C. Keulemans lith

ANORRHINUS ALBOCRISTATUS

M&N Hanhart imp.

ANORRHINUS ALBOCRISTATUS.

WHITE-CRESTED HORNBILL.

Buceros albocristatus, Cassin, Journ. Acad. Nat. Scien. Phil. (1850) vol. i. p. 135, pl. 15; Hartl. Ornith. West-Afr. (1857) p. 163; Cassin, Proc. Acad. Nat. Sci. Phil. (1859) p. 139. sp. 162; Schleg. Mus. Pays-B. (1862) p. 9; id. Handl. d. Dierk. Vog. pl. 3. fig. 39; Giebel, Thesaur. Ornith. (1872) p. 495; Reich. Jour. für Ornith. (1875) p. 12.

Buceros macrourus, Temm. Mus. Lugd.

Berenicornis macrourus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 2; id. Consp. Vol. Anisod. (1854) p. 2.

Berenicornis albocristata, Hartl. Journ. für Ornith. (1854) p. 127. sp. 391; Heine, Journ. für Ornith. (1860) p. 188. sp. 148; Sharpe, Ibis (1869) p. 385.

Buceros (Berenicornis) albo-cristatus, G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 128. sp. 7883.

HAB. Sierra Leone, St. Paul's River (CASSIN); Rio Boutry, Ashantee (PEL); Gaboon (DU CHAILLU).

Cassin, in the Journal of the Philadelphia Academy (*l. c.*), states:—"The specimen above described I received several years since from Robert Mac Dowell, M.D., Surgeon, attached to the Colonial Government of Sierra Leone, an enthusiastic naturalist, who obtained it on the banks of the St. Paul river. This species resembles no other that I have seen, or of which I can find a description, and may at once be recognized by its white erect crest and long tail." Bonaparte, in his 'Conspectus Generum Avium,' includes this species under the name of *macrourus*. This is attributed by Schlegel and others to Temminck. I cannot ascertain that such a name was ever published, but was probably merely a manuscript one attached to some specimen in the Leyden Museum. This of course cannot be noticed; and Mr. Cassin's name therefore has priority, and is the one that has been generally adopted. The present species is still rare in collections, although doubtless not uncommon in its native haunts. Its long white crest renders it very conspicuous among the Bucerotidæ, and makes it one of the most attractive species of the family. It ranges from Sierra Leone to the Gaboon, though perhaps it is not found in Fantee.

Bill black, a triangular yellow spot at base of maxilla. A small thin casque, with a prominent ridge on the side extends along the culmen for nearly its entire length, its anterior end being upright. Head covered by a long loose full crest, white; the feathers on top of head having black shafts and tips. Plumage of body black. Wings black with a green reflection; lesser coverts, secondaries, and primaries tipped with white. Tail dark green, tipped with white.

Total length 24 inches; wing $8\frac{1}{2}$; tail 16.

Female like the male, but with a smaller bill and crest.

J.G. Keulemans lith.

ANORRHINUS LEUCOLOPHUS.

M & N Hanhart 1891

ANORRHINUS LEUCOLOPHUS.

FANTEE WHITE-CRESTED HORNBILL.

Berenicornis albo-cristatus, Sharpe, Proc. Zool. Soc. (1871) p. 604. sp. 7.

Berenicornis leucolophus, Sharpe, Zool. Rec. (1873) p. 54.

HAB. Fantee (SHARPE).

In his list of the Birds of Cameroons, published in the 'Proceedings' of the Zoological Society, as above cited, Mr. Sharpe calls attention to the difference between specimens of this Hornbill from Fantee and the Gaboon. He states:—"I must remark that all the specimens received from Fantee differ conspicuously in being much smaller, and in having the wing scarcely tipped with white. The bill also is very different, being much smaller in the Fantee bird. Although I have a good series of these Hornbills, I must wait for further evidence before describing the Fantee bird as new, as Cassin's original type came from Sierra Leone, and agrees with the Gaboon bird, and not with that from Fantee, as one would expect." In the 'Zoological Record' for 1871, he remarks in a footnote, "I have since seen many specimens from Fantee, and find the character constant. The bird from this locality must be separated as *B. leucolophus*." The Fantee birds certainly do vary both in size and shape of bill, as well as in the colour of the plumage, as pointed out by Mr. Sharpe; but the great difficulty that exists against its being deemed distinct from the *albocristatus* of Cassin, is its geographical distribution. The species just named extends from Sierra Leone to the Gaboon. Of this dispersion there can hardly be any doubt, as the type specimen came from Sierra Leone, it was procured by Pel on the Rio Boutry in Ashantee, and by Du Chaillu at the Gaboon. Now it appears strange that nearly in the centre of this range a closely allied but distinct species should exist. I have never seen any of the *albocristatus* from Fantee; but that does not prove that it is not also found there with the *leucolophus* style. It is possible that Mr. Sharpe's bird is simply a local race of the widely distributed *albocristatus*.

Bill black, with a triangular yellow spot near the base of the maxilla. A keel-shaped casque rises above the maxilla at its base, runs for nearly its entire length, and terminates at a right angle to the culmen. Head and occiput covered by a very long loose crest; and these, together with the neck, are white, with the shafts of the feathers and a spot on their tips black. Entire rest of body black. Wings black with a green gloss. Tail cuneate (median feathers very long), black, glossed with green and tipped with white. Feet and tarsi black.

Total length 26 inches, wing 9, tail 17, bill $3\frac{1}{2}$, tarsus $1\frac{3}{4}$.

The specimens described and figured were in Mr. Sharpe's collection.

J. H. Koulmans lith.

Hanhart imp.

ANORRHINUS GALERITUS.

ANORRHINUS GALERITUS.

BUSHY-CRESTED HORNBILL.

- Buceros galeritus*, Temm. Plan. Col. (1830) vol. ii. pl. 520; Müll. & Schleg. Verh. Gesch. Ned. Ind. (1839-44) pp. 23, 28, v; Gray, Gen. Birds, vol. ii. (1847) p. 399. sp. 12; Low, Sarawak, p. 411 (1848); Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 44. sp. 185; Schleg. Mus. Pays-B. (1862) p. 8.
- Buceros carinatus*, Blyth, Journ. Asiat. Soc. Beng. vol. xiv. pt. i. (1845) p. 187 (juv.).
- Anorrhinus galeritus*, Reich. Syst. Av. (1849) pl. 49; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) vol. ii. p. 594. sp. 874; Moore, Proc. Zool. Soc. (1859) p. 450; Cab. & Heine, Mus. Hein. vol. ii. (1860) p. 594. sp. 874; Salv. Ucc. di Borneo, p. 79 (1874); Lord Tweedd. Ibis (1877) p. 292; Hume, Str. Feath. (1878) pp. 109, 500.
- Hydrocissa galerita*, Bon. Consp. Gen. Av. (1850) p. 90. sp. 3; Sclat. Proc. Zool. Soc. (1863) p. 214.
- Anthracoceros galeritus*, Bon. Consp. Vol. Anisod. (1854) p. 2. gen. 9. sp. 22.
- Buceros (Anorrhinus) galeritus*, Gray, Hand-l. B. pt. ii. (1870) p. 128. sp. 7877.
- Anorrhinus galeritus*, Sharpe, Ibis (1879) p. 246.

HAB. Malayan peninsula (BLYTH); Uwalabo, Malawoon, Bankasoon, Tenasserim (HUME); Sumatra (S. MÜLLER); Borneo (SCHLEGEL, DIARD); Lâwas river (USSHER); Monte Sakoembang, Lumbidan, Borneo (TREACHER, LAWUT, S. MÜLLER); Banjermassing (MOTLEY); Sarawak (DORIA, BECCARI).

This species is Reichenbach's type of his genus *Anorrhinus*, distinguished by the keel-shaped casque or crest covering the base of the culmen and sloping gradually forward to the base of the maxilla. It is a fine species, and cannot be separated generically from *A. comatus*, which Bonaparte placed in his genus *Berenicornis*. It was described by Temminck, and a fair figure given in his 'Planches Coloriées.' The synonymy presents no difficulties, the species never having received a second appellation.

I am indebted to Messrs. Hume and Davison for the following account of this bird, which is especially welcome from the fact that but very little has been recorded of its habits by any of those ornithologists who have met with it in its native haunts:—

These naturalists state:—"Though this species was not uncommon in the forests around Malawoon and Bankasoon, yet it was so very wary and difficult of approach, that only one specimen (a male) was shot by ourselves. We saw them almost daily, always in small parties of five or six, keeping to the densest portions of the forest and the tops of the highest trees. They never fly together, but always one after the other in a string or line. When about to start, they set up a sort of gabbling chorus; and after a few seconds, perhaps half a minute, of vociferous altercation, one flies away, followed immediately by another and another, till all have left.

"Their note is very similar to that of *H. albirostris (malabarica)*; and, like these, they continually utter it at short intervals so long as they remain perched.

"Their flight is almost noiseless, with none of the metallic clang so conspicuous in that of *R. undulatus* and *subruficollis*; nor has it the swish of *H. cavatus*, Shaw, or *B. rhinoceros*. In fact, it is just like that of *H. affinis*, *malabaricus*, or *R. scutatus* (Vigil)—a few rapid strokes of the wing, a short sail with outspread wings, again a few rapid strokes, again a sail, and so on. Although the flight of all these is comparatively noiseless, you still can, when sufficiently close, detect the sound of their wing, unlike *B. comatus*, whose flight is utterly noiseless, almost owl-like.

"They are strictly arboreal in their habits, never descending to the ground as does *B. comatus* and the small black-billed *H. nigristrostris*, Blyth, of the Malayan peninsula, which is generally considered the female of the much larger and white-billed *H. malayana*, but which we never met with in company with any white-billed specimens.

"This species is, we believe, strictly frugivorous; the only specimen whose stomach we dissected contained only fruit. Of course many Hornbills, like *B. comatus* for instance, are quite omnivorous; we have taken both lizards and birds out of the stomach of these; but we are inclined to think that it is chiefly the species that habitually descend to the ground that feed in this fashion, and *galeritus* most certainly does not so descend, as, although able to procure but few specimens, we saw the bird constantly, not only in Southern Tenasserim, but also at Kuroo, Nealys, Johore, and other places in the south of the Malayan peninsula.

"The following were the dimensions and colours of the soft parts recorded from a fine fresh specimen of an adult male:—

"Length 33·0, expanse 46·74, tail from vent 13·0, wing 14·25, tarsus 2·0; bill from gape 5·75, from posterior margin of casque along ridge and culmen to tip 6·75; of casque only, 4·0; height of upper mandible and casque at centre of casque 1·46, of upper mandible at the same place 0·79; weight 2·5 lb.

"The legs, feet, and claws were black; the irides lake-red; the bill entirely black. Gular and orbital skin pale blue, darkest just in front of the eye, at the angle of the gonys and junction with the feathers of the throat. Eyelids mottled black and white."

Male.—Bill black. At a point about the centre of the maxilla a casque slopes gradually upwards from the culmen and inclines backward to the head. This is highest at its anterior end, and is black throughout. Bare skin around the eyes, and also on the cheeks and throat, pale blue. Head covered with a long loose crest, which, together with the neck and mantle, is dark green, the feathers very light along their edges. Wings have the secondaries bronzy green, edges whitish. Rest of body light brown, the edges of many feathers buff. Tail is bluish white for two thirds of its length from the base, rest dark green. Iris lake-red. Eyelids mottled with black and white. Feet greenish grey.

Length 35 inches, wing 14, tail 13, tarsus 2, bill $5\frac{1}{8}$, casque $3\frac{1}{8}$; height of bill and casque $2\frac{1}{2}$.

Female.—Precisely like the male, with the exception of the casque and bill, which are shorter and smaller, and the bare skin on throat, which is yellow.

Wing $12\frac{1}{2}$ inches, tail $11\frac{3}{4}$, tarsus $1\frac{1}{2}$, bill $4\frac{1}{4}$.

Specimens described and figured are contained in the collection of the British Museum.

J.G. Keulemans del.

Hanhart imp.

ANORRHINUS TICKELLI.

ANORRHINUS TICKELLI.

TICKELL'S HORNBILL.

- Buceros tickelli*, Blyth, Journ. Asiat. Soc. Beng. vol. xxiv. pp. 266, 285, vol. xxviii. p. 412; Jerd. Ibis, (1872) p. 5.
Toccos tickelli, Tickell, Ibis, (1864) p. 173, pl. iii.
Meniceros tickelliae, Hume, Str. Feath. vol. ii. (1874) p. 470.
Anorrhinus tickelli, Blyth, Cat. Mamm. & Birds Burma, (1875) p. 69. sp. 70.
Cranorrhinus corrugatus, Blyth, Cat. Mamm. & Birds Burma, Journ. Asiat. Soc. Beng. vol. xliii. p. 69 (1875), juv.
Anorrhinus tickelli, Godw.-Aust. Ibis, (1878) p. 206.
Ocyeros tickelli, Davids. Str. Feath. vol. vi. (1878) p. 103; Hume, Str. Feath. vol. vii. (1879) p. 499.
Ocyeros tickelliae, Bingham, Str. Feath. vol. viii. (1879) p. 194.

HAB. Tenasserim Hills (TICKELL); Mountains of Amherst Province, Burma, up to 400 feet of elevation (BLYTH); Khasi Hills (GODWIN-AUSTEN); Thoungyah Hills, Kan Karyit (DARLING).

The *A. tickelli* was first described by the late Mr. Blyth from an immature specimen obtained by Tickell in the Tenasserim Hills. Blyth placed it in the genus *Buceros*, while Tickell and Jerdon deemed it a *Toccos*. Its proper generic position, however, is with *Anorrhinus*, and it is a closely allied form to *A. galeritus*.

Tickell found this species from about the base of the Tenasserim Hills up to a height of 4000 feet. The birds were most abundant on the Siamese side of the range, appearing in pairs or small flocks of five or six individuals. Their voice was loud and plaintive, resembling "*whey-whey*," and whilst feeding they continually kept up a low cackle like parrots. The flight was performed by regular flappings of the wings, like *R. subruficollis*, and not with alternate flappings and sailings, like some other species of this family, and they generally flew at high elevations. The specimen figured in 'The Ibis' (*l. c.*) was procured at Thengangyee sakán, a place on the wild path used by travellers going from the Shan States of Yahan, in Siam, to Moulmein. The food of this species is fruits. Some years afterwards Col. Tickell met this Hornbill a second time in the flat forest lying along the south side of the Hounghtrau river, to the south of Thengangyee sakán, and on a considerably lower level. They were very wild and could not be approached; but on his last day's march through the forest he met with three individuals near a Karén clearing, two of which one of his followers succeeded in securing. They both proved to be males, not differing, however, in plumage from the specimen he first obtained, which was a female.

No other examples of this species were obtained, since Tickell met with it, until last year, when Mr. Hume, knowing my great desire to present illustrations of the adult in this work, most generously deputed his assistant curator, Mr. G. Darling, to visit the only place known to him which they frequented; and after much trouble he succeeded in securing a series of both sexes.

An account of this expedition is given in 'Stray Feathers' (*l. c.*), and the following is a summary:— Six males and six females were procured, the birds having been first met with on the Thoungyah Hills, some fifteen miles from Kan Karyit on the way to the Yahine Territory, *viâ* Meawuddee. They were very shy and difficult of approach, always in heavy forest, and it was wonderful how quickly they discovered a man who endeavoured to get near to them. When feeding they keep up an incessant cackling; but when frightened, a shrill scream is given and then they fly away one after another in an extended line; they do not fly far, and recommence their cackling when they alight. When feeding they come to within thirty or forty feet of the ground, but if disturbed fly to the tops of the highest trees. They are constantly flying and sailing about, never remaining more than a minute in one spot; and their chief food seemed to be berries, although the remains of some kind of fly were discovered in their stomachs. At a latter period Captain Bingham records, in 'Stray Feathers' (*l. c.*), that he found this species not uncommon in the Kyoong Choung Reserve, high up on the Zainee River, again between Meawuddee and Kan Karyit; and in July 1879 he saw a flock between Yunbine on the Salaween River and Koosaik on the Thoungyeen. Everywhere they kept to the tops of the very highest trees, completely out of the range of an ordinary gun.

Male.—Darling states that the adult male has the bill yellowish white, slightly browner on the casque, with a patch of saffron or orange-yellow at the base, and usually a small dusky patch at the tip of the mandible, this last at times running as a narrow band backwards along the commissure. Head bistre-brown, with the shafts of feathers paler. Upper parts umber-brown tinged with olive, and in certain lights exhibiting a greenish gloss. Secondaries and primaries greenish black, the latter with their outer margins for a short space in the centre as well as their tips whitish. Some of the greater wing-coverts are also tipped with white, which is not the case with the female. Occasionally the white tippings of the quills extend to the tertiaries. Chin, throat, sides of neck, entire lower surface of body, thighs, and lower tail-coverts bright ferruginous rufous. Two median rectrices like the back, with pale tips; remainder greenish black, tips pale. Irides bright brown. Legs and feet brownish black, claws horny black. Orbital skin and bare spot at base of mandible bluish white or blue, not unfrequently pink.

Length 30 inches, wing 13, tail $12\frac{1}{2}$, tarsus $1\frac{3}{4}$, bill from gape $4\frac{2}{3}$.

Female.—Bill brownish black. Upper parts and tail resemble the male; but the chin, throat, sides of neck, entire lower surface of body, thighs, and lower tail-coverts are grey earthen brown, partially tinged with dull ferruginous rufous. Irides and soft parts coloured as in the male; but Darling records that one female had black irides. Legs and feet brownish black. Primaries black, with a white patch on outer webs; secondaries dark green.

Length 28 inches, wing 12, tail $11\frac{1}{4}$, tarsus 2, bill from gape $4\frac{1}{3}$.

A young male shot by Captain Bingham had the bill whitish tinged with greenish yellow; bare skin on chin and above and behind the eyes purplish pink, rest of bare facial skin fleshy white. Irides with an inner and an outer circle of grey. Legs and feet greenish plumbeous, claws horny. The plumage resembles that of the female, except the entire lower surface, which is a

ruddy buff, palest on the throat; the feathers of the throat, and many of those on the breast and abdomen, with white centres. Forehead, crown, and occipital crest dark ferruginous brown, the feathers, especially the crests, strongly tinged with grey, as are also those of the thighs. Greater wing-coverts, outer margins of primaries, and some of the secondaries margined with whitish buff. Tail-feathers and primaries tipped with white.

The specimens figured were obtained by Mr. Darling, and most kindly presented to me by Mr. Hume, to whom I owe my sincere thanks for the efforts made to procure them.

ANORRHINUS AUSTENI.

GODWIN-AUSTEN'S HORNBILL.

(NOT FIGURED, NO SPECIMEN OBTAINABLE.)

Anorrhinus galeritus, Godw.-Aust. (nec Temm.) Journ. Asiat. Soc. Beng. (1870) p. 96, vol. xxxix., juv.?

Anorrhinus austeni, Jerd. Ibis (1872) p. 6, juv.; Hume, Str. Feath. vol. v. p. 117 (1877) vol. vii. pp. 167, 499 (1878).

Anorrhinus tickelli, Godw.-Aust. Ibis (1878) p. 208, juv.

HAB. North Cachar hills (GODWIN-AUSTEN).

A presumably young bird of some species of Hornbill was procured by Colonel Godwin-Austen in the North Cachar hills, and was referred by him to *A. galeritus*, Temm. Jerdon later described the specimen as belonging to an undescribed species, and named it *austeni*, after its discoverer. Blyth, in his 'Mammals and Birds of Burmah,' considered the bird to be the same as *Cranorrhinus corrugatus*, having examined a head which he supposed belonged to Godwin-Austen's specimen. In this, however, he was mistaken, as Lord Tweeddale states (Ibis, 1878, p. 208) that this head had no reference to Godwin-Austen's bird. The latter gentleman, in the communication to 'The Ibis' to which Lord Tweeddale's remarks given above were attached, believes that his example is the young of *A. tickelli*, Blyth. Mr. Hume contends, however, that this conclusion is erroneous; and I am inclined to agree with him, and believe that Godwin-Austen's bird most probably represents a distinct species. Fortunately, Mr. Hume has been able to procure a very fine series of *A. tickelli* of both sexes; and none of them possess the peculiar coloration of plumage described by Godwin-Austen in his bird. In no specimen of *A. tickelli*, as stated by Mr. Hume, are *the throat and sides of the neck white*, nor does either sex have *the base of the primaries white*, or the primaries themselves *barred with white*, as is the case with *A. austeni*. The bill being yellowish white would seem to prove that the specimen of *A. austeni* was a male, as the females of this genus have a blackish-brown bill. In size *A. austeni* and *A. tickelli* are about the same. I have not been able to see this example of *A. austeni*, which was deposited by Godwin-Austen in the British Museum; and the ticket having been changed upon it, identification of the specimen seems to be impossible. Under these circumstances, and judging from the description, I consider that *A. austeni* cannot be referred to *A. tickelli*, and therefore leave it as a distinct species until the future acquisition of examples may define beyond question its proper position.

The following is Colonel Godwin-Austen's description of his specimen:—

"The whole of the upper part pale slaty grey, having in certain lights a greenish tinge; throat and sides of neck white, dull rufous on the breast and belly, thighs, and under tail-coverts. Primaries greenish black, tipped and barred with white; a white spot formed by the tips of the outer wing-coverts, the bases of the primaries being also of this colour; secondaries edged whitish; tail tipped white; centre feathers same colour as the back. Bill yellowish white. Length about 31 inches, wing 13, tail 13, bill to gape $4\frac{1}{2}$, depth 2."

JG Keulemans lth.

ACEROS NEPALENSIS.

Hanhart imp.
27

ACEROS NEPALENSIS.

RUFIOUS-NECKED HORNBILL.

Buceros nepalensis, Hodgs. Calc. Glean. Sc. (1829) vol. i. p. 249; id. Proc. Zool. Soc. (1832) p. 15; id. Asiat. Research. (1833) p. 178, pls. 1, 2; Blyth, Journ. Asiat. Soc. Beng. (1843) vol. xii. pt. ii. p. 989, (1847) p. 997. pl. 44. fig. 1; G. R. Gray, Gen. Birds, (1849) vol. ii. p. 400. sp. 24; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 45. sp. 189; Schleg. Mus. Pays-B. (1862) p. 10; Giebel, Thes. Ornith. (1872) p. 501.

Aceros nepalensis, Hodgs. Gray's Zool. Misc. (1844) p. 85; Bon. Consp. Vol. Anisod. (1854) p. 3; Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 601. sp. 884; Jerd. B. Ind. (1862) vol. i. p. 250; Blyth, Ibis (1866) p. 350; Jerd. Ibis (1872) p. 5; Blyth, Cat. Mamm. & B. Burma (1875), p. 69. sp. 73; Gammie, Str. Feath. (1875) p. 210.

Calao nepalensis, Bon. Consp. Gen. Av. (1850) p. 90. sp. 5.

Buceros (Aceros) nepalensis, Tickell, Ibis (1864) p. 182; Gray, Hand-l. B. (1870) pt. ii. p. 129. sp. 7888.

Khulut of the Lepchas, *Gog-biah* of the Bhotees of Darjeeling (JERDON).

HAB. S.E. Himalayas, hill-ranges of Assam and Munipur, Kachar, Tenasserim Provinces (BLYTH).

This handsome bird, one of the finest of the family, has been made the type of the genus *Aceros* by Mr. Hodgson. Although belonging to the group of large Hornbills, it differs conspicuously from them in being destitute of a casque, that remarkable addition to the bills of its relatives; instead of this the culmen is swollen at the base, and slightly elevated. There is not, fortunately, any confusion existing in the synonymy of this species, the bird being too striking in its appearance to leave much room for errors in its identification to occur.

The range of this fine bird appears to be along the South-eastern Himalayas to the mountains of Tenasserim. Tickell shot one on the spur leading to Mooleyit Peak, at a height of 3500 feet, which is the only specimen obtained in British Burma that I have seen recorded. It did not differ from the Nepaulese specimens. The flight of this species is performed by irregular beats of the wings, with retracted neck and lowered tail. The voice of the adult is a hoarse croak; but when the bird is excited, it resembles a dog's bark, and is very noisy. The young utter a sound like the clucking of a brood-hen, according to Hodgson, falling occasionally into the shriller notes of the guinea-fowl. The species feeds chiefly on fruits, but will not refuse reptiles if hungry. When tame it will eat meat, both raw and cooked, and also boiled rice. Every thing it takes it swallows whole. Its flesh is eaten by the mountaineers, and much esteemed by them. It nests in a hole in a tree, excavating it, says Hodgson, out of the solid wood, and closing the aperture by an *ingeniously contrived door!* so that it is with difficulty discovered. It is said to go in pairs, and not to be gregarious, although Jerdon speaks of observing it in small flocks. He says that this species is easily alarmed, and that the people of Cachar and Munnipore take advantage of this

trait by shouting, beating drums, and firing matchlocks whenever they see these birds flying over their villages. They immediately become bewildered, and descend to the nearest tree for shelter, sometimes even to the ground, and are killed by arrows, occasionally by sticks. Mr. Hodgson's statement that this species closes its abode in a tree by a curiously contrived door I have not seen corroborated by any other naturalist. In 'Stray Feathers' for 1875, p. 210, Mr. Gammie describes a nest and egg of this species at Poomong, in Sikhim. The tree was a species of *Dysoxylon*, about 80 feet high; and under the lowest branch there was a slit, which was the entrance to the bird's house. On each side of the slit was a quantity of plaster, which had evidently been placed there by the female, and which did not meet in any part. At the top of the slit was a round hole; and from this to the bottom the opening was about two inches broad. The entrance, after the plaster was removed, was seventeen inches in length, by four and a half in breadth, and the hollow of the tree seventeen inches in diameter. In the hole were merely a few of the bird's feathers. The egg was an oval, compressed at one end, and slightly pyriform. The shell was strong and thick, coarse, without gloss, and pitted with minute pores. In colour it was dirty white, with a yellowish tinge, and everywhere obscurely stippled with minute purer white specks. It measured 22.5 by 1.75. The plaster, as examined by Dr. D. Cunningham under the microscope, was entirely composed of vegetable tissue, cells, fibres, oil-globules, &c., and contained no clay or mineral matter of any kind. The vegetable tissue appeared semidigested, many cells being wholly or partially emptied of their contents, and free granules and globules of a bright yellow oily-looking matter abounding. The most abundant and characteristic forms of cells present were:—1st, small, totally empty, thick-walled cells, scattered or still holding together in small patches; 2nd, large rounded cells, full of the yellow oily matter so abundant in the free state, and of a deep brown colour. Their contents were rather of a gummy than oily nature, as boiling with *liquor potassæ* reduced the material to a glutinous mass, of a deep brown colour. There were also present in small numbers fragments of feathers, spores of fungi, &c. From this analysis it would seem evident, as stated by Mr. Hume, that the plaster was but the bird's own ordure, with which she closed the opening to her nest, leaving an aperture only sufficiently wide to receive food from the male. A heap at the foot of the tree of rejected droppings was of the same composition as the plaster, but with fewer of the gummy globules and a larger proportion of feathers, &c. Nothing is mentioned of a door; and in this case, at least, it is evident the species closed the entrance to the nest in the same manner as do the other members of this family. The male was seen to feed the female with the fruit of the *Dysoxylon* tree.

The usual position assumed by this Hornbill, according to Tickell, is a kind of squat, with ruffled neck-feathers, the neck drawn in between the high shoulders of the wings, and the tail either erect like a Magpie's, or dropped down. At such times the bird has a very stupid appearance. It is, however, when in the act of dressing its plumage that this handsome species appears to most advantage. Then it rises on its feet, exposing the strong legs, the shoulders of the wings are projected, and the neck extended and arched backwards, giving to the bird "some of the graces and even terrors of the noble birds of prey." In disposition this Hornbill is gentle, but not deficient in spirit; and when captive, although it dislikes them, it views with indifference

the near approach of either man or dog. It soon becomes tame, its quiet habits enabling it to easily endure confinement.

Male.—Bill yellow, without casque, culmen slightly elevated and swollen at base. Sides of basal portion of maxilla ridged transversely, these grooves chestnut. Naked skin around the eye and at base of bill light blue, that of the throat bright scarlet. Head, neck, and lower parts bright rufous; back and wings black, glossed with green; the third to eighth primaries broadly tipped with white. Tail white for more than half its terminal length; remainder black.

Length 4 feet, wing $18\frac{1}{2}$ inches, tail 18, bill 9; height 3.

The female has her entire plumage black, glossed with green, excepting the ends of the primaries and terminal portion of the tail, which are white.

Length 3 feet 6 inches, wing $17\frac{1}{2}$, tail 17, bill $7\frac{1}{2}$.

The specimens from which my description and figures were taken are contained in the British-Museum collection.

J.G. Keulemans lith.

HYDROCISSA EXARHATUS.

Hanhart imp.
32

HYDROCISSA EXARATA.

WHITE-CHEEKED HORNBILL.

- Buceros exarhatus*, Temm. Plan. Col. vol. ii. pl. 211 ♀; Wagl. Syst. Av. (1827) sp. 15; Müll. and Schleg. Verh. Gesch. Ned. Ind. (1839-44) p. 23; Gray, Gen. Birds (1846) vol. ii. p. 400. sp. 26; Schleg. Mus. Pays-B. (1862) p. 10; Gray, Hand-l. B. (1870) pt. ii. p. 128. sp. 7880.
- Buceros exaratus*, Blyth, Cat. B. Mus. Asiat. Soc. (1849) p. 44. sp. 187; Wald. Trans. Zool. Soc. (1871) vol. viii. p. 47. sp. 57; Giebel, Thesaur. Ornith. (1872) p. 498.
- Hydrocissa exarata*, Bon. Consp. Gen. Av. (1850) p. 90. sp. 6.
- Anorhinus exaratus*, Bon. Consp. Vol. Anisod. (1854) p. 2.

HAB. Malacca, Celebes (MEYER).

Temminck first described and figured this species in the 'Planches Coloriées.' It is a native of Malacca and Celebes. Some authors have placed this bird in the same genus with *Anthracoceros coronatus*, *A. malabaricus*, &c. From such species, however, the present one is separated by the shape of its casque. Properly speaking it has none, but the culmen is covered by a ridged crest; while the species above named all possess well-developed, peculiarly shaped and conspicuous casques, having nothing in common with that exhibited by *Hydrocissa exarata*. As I have shown in my article on *Anthracoceros coronatus*, the present species is the only one that can be selected to represent Bonaparte's genus *Hydrocissa*, the other species included by him in that term having previously been placed in distinct genera by Reichenbach. Of the habits and economy of this species nothing has been recorded.

Male.—The bill has the maxilla rufous, the point and base yellowish white. A crest runs from the base of maxilla to within an inch of the point, compressed laterally, its sides indented for their entire length by three deep grooves. The anterior end slopes rapidly to the culmen. The mandible is yellowish white, a black band starts from the base, runs upwards along the gape for about one third its length, then becomes apparently dark red for another third part. Superciliary stripe, cheeks, and throat white; rest of plumage jet black; wings and tail with a greenish gloss. Feet and tarsi black. Bare skin on throat near base of mandible flesh-colour.

Total length 22 inches; wing 10; tail 10; bill along gape 4; length of crest $2\frac{1}{2}$, height of bill and crest at base $1\frac{3}{4}$.

Female.—Bill like the male, but smaller, as is also the crest, which is deeply grooved, precisely in the same manner. The anterior end slopes gradually to the culmen. Bare skin of throat, apparently, black or bluish-black. The entire plumage is black, wings and tail glossed with green.

Total length 21 inches, wing 8, tail 8, tarsus $1\frac{1}{4}$.

Specimens described and figured are in my own collection, procured from Mr. Wallace.

3
4

J.G. Keulemans lith.

LOPHOCEROS NASUTUS.

M & N Hanhart
17

LOPHOCEROS NASUTUS.

BLACK-BILLED HORNBILL.

- Buceros nasutus*, Linn. Syst. Nat. vol. i. (1766) p. 154; Gmel. Syst. Nat. (1788) vol. i. p. 361; Lath. Ind. Ornith. (1790) vol. i. p. 145; Shaw, Gen. Zool. (1811) vol. viii. p. 30; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 600; Temm. Plan. Col. (1824) vol. ii. sp. 18 (text); Ersch u. Grub. Ency. (1824) p. 286; Wagl. Syst. Av. (1827) sp. 19; Less. Trait. Ornith. (1831) p. 252. sp. 2; G. R. Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 32; Hartl. Ornith. West-Afr. (1857) p. 164; Schleg. Mus. Pays-Bas (1862) p. 13; Layard, B. S. Afr. (1867) p. 227; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 486; Nitzs. Pteryl. Ray Soc. (1867) p. 102; Finsch, Trans. Zool. Soc. (1869) vol. vii. p. 277; Ayres, Ibis (1871) p. 260; Giebel, Thes. Ornith. (1872) p. 501.
- Black-billed Hornbill*, Lath. Gen. Syn. (1781) vol. i. p. 354. sp. 10.
- Calao à bec noir du Sénégal*, Buff. Planch. Enlum. (1783) no. 890.
- Le calao nasique*, Le Vaill. Hist. Nat. Ois. d'Afrique. (1806) vol. v. p. 120, pl. 236.
- Buceros nasutus*, var. *caffer*, Dumont, Dict. Sci. Nat. (1817) vol. vi. p. 213; Sundev. Æfvers. Vet. Akad. Förhdlg. (1850) p. 108.
- Buceros (Lophoceros) forskalii*, Hemp. & Ehrenb. Symb. Phys., Av. (1828) fol. z, ♂.
- Buceros hemileucus*, Hemp. & Ehrenb. Symb. Phys., Av. (1828) fol. aa, ♀; Dresser & Blanf. Ibis (1874) p. 338.
- Buceros hastatus*, Cuv. Règn. Anim. (1829) vol. i. p. 446 (note).
- Buceros pæcilorhynchus*, La Fres. Rev. Zool. (1839) p. 237, juv.; Hartl. Ornith. West-Afr. (1857) p. 164; Schleg. Mus. Pays-B. (1862) p. 14; Layard, B. S. Afr. (1867) p. 226.
- Tockus nasutus*, Rüpp. Syst. Uebers. Vög. N.O.-Afr. (1845) p. 79. sp. 321; Bon. Consp. Gen. Av. (1850) p. 91. sp. 8; Hartl. Journ. für Ornith. (1854) p. 128. sp. 39; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) p. 596. sp. 878; Von Heugl. Journ. für Ornith. (1864) p. 271; Gurney, Anderss. B. Damaraland (1872) p. 206; Dresser & Blanf. Ibis (1874) p. 337; Layard, B. S. Afr. Sharpe ed. (1875) p. 133. sp. 123.
- Tockus pæcilorhynchus*, Bon. Consp. Gen. Av. (1850) p. 91. sp. 9; Hartl. Journ. für Ornith. (1854) p. 193; Von Heugl. Journ. für Ornith. (1864) p. 271.
- Grammicus nasutus*, Bon. Consp. Vol. Anisod. (1854) p. 3.
- Grammicus hastatus*, Bon. Consp. Vol. Anisod. (1854) p. 3.
- Lophoceros forskali*, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 167.
- Toccus pæcilorhynchus*, Kirk, Ibis (1864) p. 327.
- Toccus nasutus*, Salvad. Cat. Uccelli Mar. Ross. e dei Bogos, p. 55 (418).
- Buceros forskalii*, Dress. & Blanf. Ibis (1874) p. 337.

HAB. Senegambia, Sennaar, Abyssinia, Limpopo River (AYRES); Damaraland (ANDERSSON).

The *L. nasutus* and the *L. birostris*, on account of having casques continued anteriorly to a sharp point, and projecting over the maxilla, cannot be retained in the genus *Tockus*, whose members have merely the culmen slightly elevated into a crest-like ridge. I have therefore adopted for them the term *Lophoceros*, proposed by Hemprich and Ehrenberg in 1828 as a sub-genus for the *L. nasutus*. The present bird has received many names; and its synonymy has been

greatly confused. The authors just quoted conferred two names upon it—*B. forskalii*, given to a male, and *B. hemileucus*, to a female. Cuvier called it *hastatus*, and La Fresnaye bestowed the title of *B. pœcilorhynchus*, which term was adopted by Bonaparte and some others. It is a bird of very wide distribution, as it is found in Abyssinia on the east, and is also an inhabitant of Damaraland on the west coast. It probably extends its range pretty much over the continent.

Andersson states that it does not occur in Great Namaqualand. It goes in small flocks of about six individuals, roosts on large trees, perching upon branches about halfway up, and returns nightly to the same place. It rarely alights upon the highest branches. During the hot weather it suffers apparently greatly from the heat, and seeks the shadiest part of the forest, where it gasps as if for breath, and may be easily approached. When flying, it occasionally utters short piercing cries.

Although almost omnivorous, the principal food of this species consists of berries, young shoots, and insects. The irides are reddish brown. Mr. Ayres found this bird common along the Limpopo; and he also received it from the Megaliesberg, a range of wooded mountains about seventy miles from Potchefstroom. Once he was greatly astonished to observe one of these birds perched on a tall tree singing very prettily, with a voice like that of a Thrush.

Messrs. Dresser and Blanford having examined Ehrenberg's type of *Buceros forskalii*, state (*l. c.*) that it is scarcely distinguishable from *B. nasutus*, but appears to be larger and has a longer wing. The type is said to have come from Arabia! There were three specimens in the collection; and one from Abyssinia had the longest wing, but in all its other measurements it was the smallest. These slight variations in size do not seem to be sufficient for a specific difference, as they are evidently not constant in examples even from any one locality; and I have not hesitated to place *B. forskalii* among the synonyms of *L. nasutus*.

In his notes attached to Dr. Finsch's article on the birds of North-eastern Abyssinia and the Bogos Country (Trans. Zool. Soc. vol. vii. p. 278), Mr. Jesse states that this species was found from Ain to the Anseba, and its note was so variable that it was a source of constant trouble and disappointment, as it led him always to expect to find some different species. He noticed it hawking for insects in a very awkward manner; it would also eat fruit.

The adult has the bill black, with the exception of a long triangular-shaped bright-yellow mark from the base, reaching halfway along the sides of the maxilla. On the mandible are three or four fine transverse lines also bright yellow. The culmen is swollen at the base, forming a diminutive casque that extends along the bill for a little over half its length, terminating in a projecting point. Head and throat dark grey, darkest on the throat, but graduating to a light brown upon the breast. A very broad whitish superciliary stripe from the base of the bill to the occiput. Back and wings dark brown, every feather broadly margined with white. Entire underparts pure white. Tail blackish brown, all the feathers except the two central ones tipped with white. Tarsus brownish black. Iris reddish brown. Total length, including bill, 18 inches; wing 8; tail 9; bill along gape $3\frac{3}{4}$; length of casque $2\frac{1}{4}$; tarsus 1. Sexes alike.

The young have the basal portion of the maxilla and transverse lines on mandible yellowish, and the tip of the bill for about an inch red, remainder black. No casque, but the basal portion of the culmen slightly elevated.

LOPHOCEROS BIROSTRIS.

GINGI HORNBILL.

Calao de Gingi, Sonn. Voy. aux Ind. (1782) p. 214. pl. 120; Le Vaill. Ois. Amér. (1801) pl. 15.

Buceros birostris, Scopoli, Delic. Flor. et Faun. Insub. (1786) p. 87. sp. 31.

Gingi Hornbill, Lath. Gen. Syn. (1787) p. 71.

Buceros ginginianus, Lath. Ind. Ornith. (1790) vol. i. p. 146. sp. 13; Shaw, Gen. Zool. (1811) vol. viii. p. 36; Vieill. Nouv. Diet. Hist. Nat. (1816) vol. iv. p. 594; Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 207; Vieill. Ency. Méth. (1823) vol. i. p. 306; Temm. Plan. Col. (1824) vol. ii. sp. 13 (text); Ersch u. Grüb. Ency. (1824) p. 285; Less. Trait. Ornith. (1831) p. 253. sp. 5; Blyth, Journ. Asiat. Soc. Beng. (1843) vol. xii. pt. ii. p. 996; Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 27; Schleg. Mus. Pays-B. (1862) p. 11.

Buceros oxyurus, Wagl. Syst. Av. (1827) p. 14.

Buceros cinerascens, Hodg. Gray's Zool. Misc. p. 85 (1844).

Buceros birostris, Blyth, Journ. Asiat. Soc. Beng. (1847) vol. xvi. p. 995; id. Cat. B. Mus. Asiat. Soc. (1849) p. 44. sp. 183.

Tockus ginginianus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 2.

Meniceros ginginianus, Bon. Consp. Vol. Anisod. (1854) p. 3. gen. 13. sp. 34; Ball, Str. Feath. p. 388 (1874).

Tockus bicornis, Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 597. sp. 881.

Penelopides ginginianus, Cab. & Heine, Mus. Hein. (1860) Th. ii. p. 169. no. 466.

Meniceros bicornis, Jerd. B. Ind. (1862) vol. i. p. 248. sp. 144; Blyth, Ibis (1866) p. 350; Horne, Proc. Zool. Soc. (1869) p. 241; Marshall, Str. Feath. vol. iii. (1875) p. 331; Butl. Str. Feath. (1877) vol. v. p. 218.

Ocyrceros ginginianus, Hume, Nests and Eggs of Indian Birds, (1873) pt. i. p. 113.

Chakotra, Hindostanee; *Puttial dhanes* in Bengal; *Dhamuar* or *Lamdar* in Upper Provinces; *Rundu-mukala guwa*, Tel. (*i. e.* Two-billed Pigeon); *Manumukala kaka* (*i. e.* Three-billed Crow), Tam.; *Selagilli* at Sangor (JERDON).

HAB. India generally, not Assam, never on the eastern side of the Bay of Bengal, tolerably common in the tank country north of Ahmedabad (BUTLER).

This is the species generally known as *Buceros ginginianus* of Latham; but it was described previous to that author by Scopoli, who conferred the term *birostris* upon it, which is the one it must bear. It is the second species of the genus *Lophoceros*, distinguished by the pointed casque, and is common throughout India, but never goes to the eastern side of the Bay of Bengal. The following short extract from Jerdon's 'Birds of India' gives a very fair idea of the distribution of the species, and its mode of life:—

"This small Hornbill is spread throughout all India in well-wooded districts, but does not appear to extend into Assam, nor into any of the countries to the eastward. It is not found in thick forests, and is almost unknown in the woods of Malabar, but frequents open and thick forest-

jungle, groves of trees, gardens, and especially the fine old avenues of banian and other trees that abound in the extreme south of India, and are said to have been mostly planted by Hyder Ali. It is almost always found in small parties, occasionally in pairs, living chiefly on fruit, and especially the figs of the banian, peepul, and other fig-trees, sometimes, however, feeding on large insects. On such occasions I found that it had eaten Mantides and Locustæ. It has a loud sharp cry. In some parts of India the flesh is used medicinally to alleviate the pangs of child-birth."

Mr. Ball says that in Chota Nagpur this species is occasionally met with on the borders of heavy jungle, and in open country, where mhowa trees abound. He shot it in Manbhum, Singhbhum, and Hazaribagh, but did not see it in Sirguja. It was apparently more abundant in the Rajmehals, and in the Satpura hills very common. A female, procured in the Rajmehal hills on the 14th April, had in her stomach a quantity of grasshoppers, in addition to some fruits of a species of *Ficus*.

Mr. Horne, in the 'Proceedings of the London Zoological Society' for 1869, gives an interesting account of the breeding-habits of this species, as observed by him at Mainpuri, in the North-west Provinces of India. He states that, on account of the neck, bill, and tail being long, and the wings comparatively short, the flight is rather undulating, and the flapping frequent, and that when in the air the bird often utters its harsh note. When feeding, this species is very fearless; and it has the habit of climbing by the bill as a parrot does. "It would also extract the oranges piece by piece, leaving the skins hanging apparently entire upon the twigs." Upon the lawn surrounding Mr. Horne's house was a fine sissou-tree (*Dahlbergia sissoo*), with a large hole in the trunk beneath the first fork, for the possession of which Rollers and Parrots were always contending. At last a pair of this Hornbill made up their minds to occupy it. The hole was about a foot deep; and on the 29th April the female went into it, and did not come out again. The next day he observed her working hard to close up the entrance with her own ordure, plastering it right and left with her bill, as with a trowel. The male was most assiduous in supplying her with food, bringing generally the small peepul-fig. He would first alight near by, then fly to the hole, and, holding on to the bark by his claws, knock with his bill, when that of the female would appear and receive the fruit. The hole in the tree was at first about 6 inches high, and 3 to 4 wide; but when closed the opening at its widest part was only sufficiently large to admit the finger. It took two or three days to reduce the opening to this size, after which the ordure was thrown out. On the 7th May he took the female out, and obtained three eggs. She was very fat, and scarcely able to fly at first, but succeeded in a short time. The natives say that as soon as the newly-hatched young need food the female digs herself out.

In 'Stray Feathers' (*l. c.*) the Messrs. Marshall call attention to the difference in the plumage of the sexes of this species, from examples collected and the sex ascertained by themselves. In the female and immature birds the white tips are wanting to the first six primaries, and sometimes to the seventh, while on all but the first two the white streaks are more developed than in the adult male. The casque is lower, and has not the sharp-pointed horn.

In the male the head, face, and tail become blacker with age.

General plumage grey, darkest on top of head, paler on breast, and becoming almost white on abdomen and lower tail-coverts. Ear-coverts plumbeous. Back pearly grey with a pinkish tinge. Primaries and secondaries brownish black, tipped broadly with white. Inner secondaries pearly grey with a pinkish tinge. The first three primaries with a white line on their outer web near the centre. Tail pale rufous, with a subterminal blackish-brown bar, and tipped with white. Bill has the anterior third part of the culmen and tips, together with the underpart of mandible, yellowish white, rest black. Casque low, greatly compressed; ridge extending for two thirds the length of the culmen, to which it returns at a very acute angle, forming a sharp point on the end of the casque. The posterior portion does not extend over the head, and is hidden in the feathers. Iris red-brown, feet plumbeous.

Total length 21 inches, wing 8, tail 10, bill $4\frac{1}{2}$, casque $2\frac{1}{2}$, tarsus $1\frac{3}{4}$.

I desire to acknowledge my indebtedness to Lord Tweeddale for the loan of the specimens described and figured.

J.G. Keulemans lith.

Hanhart imp

TOCKUS MELANOLEUCUS.

TOCKUS MELANOLEUCUS.

BLACK-AND-WHITE HORNBILL.

- Buceros melanoleucus*, Licht. (nec Vieill.), Cat. Rer. Nat. rar. Hamb. (1793) p. 8. no. 90; Bechst. Lath. Uebers. vol. ii. (1794) p. 362; id. Kurze Uebers. p. 170; Licht. Verz. Doublet. (1823) p. 21; Wagl. Syst. Av. (1827); Gray, Gen. B. (1847) vol. ii. p. 400. sp. 29; Hartl. Ornith. Ost-Afr. (1857) p. 164; Schleg. Mus. Pays-B. (1862) p. 12.
- Le Calao couronné*, Le Vaill. Hist. Nat. Ois. Afriq. (1806) vol. v. p. 117, pls. 234 & 235.
- Buceros coronatus*, Wilkes? Ency. Lond. (1808) vol. iii. p. 480; Shaw (nec Bodd.), Gen. Zool. (1811) p. 35; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 593; id. Ency. Méth. (1823) vol. i. p. 401, pl. 240. fig. 4; Temm. Plan. Col. (1824) vol. ii. sp. 16 (text); Ersch u. Grub. Ency. (1824) p. 285; Wagl. Syst. Av. (1827) sp. 17; Less. Trait. Ornith. (1831) p. 253. sp. 4; Swains. Zool. Illust. 1st ser. vol. iii. pl. 178; id. B. W. Afr. vol. ii. p. 257; Layard, B. of S. Afr. p. 225; id. Ibis (1869) p. 372.
- Tockus melanoleucus*, Bon. Consp. Gen. Av. (1850) p. 91. sp. 7; Hartl. Journ. für Ornith. (1854) p. 128. sp. 393; Horsf. & Moore, Cat. Birds Mus. E.-Ind. Co. (1856-8) vol. ii. p. 597. sp. 881; Gurney, Anderss. Birds Damara Land (1872) p. 208; Layard, B. of S. Afr., Sharpe's ed. (1875) p. 127. sp. 119; Bocage, Ornith. Angola (1878) p. 116; Nichols. Proc. Zool. Soc. (1878) p. 358.
- Rhynchoceros melanoleucus*, Bon. Consp. Vol. Anisod. (1854) p. 3, gen. 16. sp. 41.
- Lophoceros melanoleucus*, Cab. & Heine, Mus. Hein. (1860) p. 168.
- Toccus coronatus*, Gurney, Ibis (1861) p. 133.
- Toccus melanoleucus*, Gurney, Ibis (1862) p. 157; Sclat. Proc. Zool. Soc. (1864) p. 111; Kirk, Ibis (1864) p. 327; Bocage, Journ. Sc. Math. Lisb. (1868) vol. ii. p. 347. sp. 98; Gulliv. Proc. Zool. Soc. (1872) p. 16, (1875) p. 490; Sclat. Rev. List Vert. Anim. (1872) p. 172. sp. 281.
- Buceros pallidirostris*, Hartl. & Finsch, Von der Deck. Reis. Ost-Afr. (1870) p. 871; Bocage, Journ. Sc. Math. Lisb. (1870) vol. ii. p. 348. sp. 100; Gray, Hand-l. B. pt. ii. (1870) p. 130. sp. 7907.
- Buceros (Lophoceros) melanoleucus*, Gray, Hand-l. B. pt. ii. (1870) p. 130. sp. 7906.
- Tockus pallidirostris*, Bocage, Ornith. Angola (1878) p. 117.

Sunguandondo in Angola (BOCAGE).

HAB. River-Shiré valley, east coast, in open bush and low trees (KIRK); Darra-Salam (NICHOLSON); west coast, in Angola, Ovampo-land, and Damara-land (ANDERSSON).

This is one of the species of this family whose synonymy is very much confused. It was first described by Lichtenstein in 1793 (*l. c.*) as *Buceros melanoleucus*, and afterwards by Shaw (*l. c.*) as *B. coronatus*. A pale variety subsequently received the name of *pallidirostris* from Finsch and Hartlaub (*l. c.*). The majority of authors employed Shaw's name for the species, which was confounded with *coronatus* of Boddaert, a name bestowed upon an entirely different bird. It

seems to be more an inhabitant of Southern Africa, never having been obtained, I believe, north of Angola on the west coast, although it goes nearly to the equator on the east, as Layard states that he found it very abundant at Kisiludini, flying in small flocks or families, and feeding on berries and fruits. At the Knysna he states that it is also common. Andersson says that in Damara Land the food of this species consists of beetles and lizards, very different fare from that of the same species on the other side of the continent.

Male. Bill at base yellow; remainder bright red, with a narrow line of dark red along the apical portion of the cutting edge. The maxilla is broad and swollen at the base, becoming greatly compressed towards its anterior end, and rising into a minute casque, in old birds, that terminates at a right angle to the culmen. Head covered by a rather full, not long, crest, dark brown, with the feathers on sides and lower part of neck streaked with white. Cheeks, throat, and upper part of breast lead-colour. Wings and back dark brown; feathers of the former margined with light brown. Underparts white. Tail brownish black, all the feathers broadly tipped with white, except the median ones, which have but a small amount of white at their ends.

Total length without bill 16 inches, wing $9\frac{1}{2}$, tail 8; bill along gape $3\frac{1}{4}$, height at base 1 inch, tarsus 1.

The female is like the male, with a smaller bill.

My description and figures were taken from specimens in my own collection.

JG Keulemans lith.

TOCKUS SEMIFASCIATUS. TOCKUS FASCIATUS.

Hanhart imp.
30

TOCKUS FASCIATUS.

BLACK-AND-WHITE-TAIL HORNBILL.

Le Calao longibande, Le Vaill. Hist. Nat. Ois. Afr. (1806) vol. v. p. 115. pl. 233.

Buceros longibandus, Wilkes ? Ency. Lond. (1808) vol. iii. p. 480.

Buceros fasciatus, Shaw, Gen. Zool. (1811) vol. viii. p. 36 ; G. R. Gray, Gen. Birds, (1816) vol. iii. p. 400. sp. 28 ; Temm. Plan. Col. (1824) vol. ii. sp. 15 ; Ersch u. Grub. Ency. (1824) p. 284 ; Hartl. Ornith. West-Afr. (1857) p. 163 ; Gurn. Ibis, (1859) p. 153 ; Schleg. Mus. Pays-B. (1862) p. 12 ; Eyton, Osteol. Av. (1867) p. 62 ; G. R. Gray, Hand-l. Birds (1870), pt. ii. p. 130. sp. 7897 ; Giebel, Thes. Ornith. (1872) p. 498 ; Reichenb. Journ. f. Ornith. (1876) p. 445.

Buceros melanoleucus, Vieill. (nec Licht.) Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 595 ; Wagl. Syst. Av. (1827) sp. 16.

Tockus fasciatus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 6 ; Hartl. Journ. f. Ornith. (1854) p. 127. sp. 392 ; Cass. Proc. Acad. Nat. Scien. (1856) p. 319. sp. 27, (1859) p. 140. sp. 163 ; Bocage, Ornith. Angola, (1878) p. 123.

Lophoceros fasciatus, Cab. & Hein. Mus. Hein. (1850) Th. ii. p. 168. sp. 7 ; Heine, Journ. f. Ornith. (1860) p. 188.

Grammicus fasciatus, Bon. Consp. Vol. Anisod. (1854) p. 3.

HAB. Angola? (SCHLEG.) ; Congo, Old Calabar, and the borders of the Casamanze (HARTLAUB) ; Cape Lopez (DU CHAILLU).

The *T. fasciatus* was first made known to ornithologists by Le Vaillant (*l. c.*) as *le Calao longibande*, in his work entitled 'L'Histoire naturelle des Oiseaux d'Afrique,' and a figure published. Unfortunately he did not confer upon the bird a Latin name ; and therefore the species has always been known by that of *B. fasciatus*, bestowed upon it by Shaw in 1811. It is true, a name was given to the species previously in the 'Encyclopædia Londinensis ;' but, as the author is not known, it cannot be accepted. Mr. Cassin, in the 'Proceedings of the Academy of Natural Sciences of Philadelphia' for 1867, called the attention of naturalists to the list of birds in the work just named, and made them generally known ; but there is no evidence to show that Mr. Wilkes, who is claimed as the author of the ornithological portion, ever had any thing especially to do with it, although he may have been the editor of the work ; and there is no signature attached to the articles to indicate who the writers were. Under these circumstances, although the names bestowed upon several of the Hornbills in the publication are the earliest known, they cannot be received ; for to make a name valid that of its author should go with it.

Nothing has been recorded concerning the economy and habits of the present species. It is easily recognized from its near ally (*T. semifasciatus*) by the two lateral rectrices being pure white for their entire length. It does not seem to be uncommon in the localities it frequents, and is often sent to Europe in collections. Professor Schlegel gives the locality of a specimen in the

Leyden Museum as *Angola* ; but, according to Professor Bocage, if it occurs there it must be very rare, and probably is only found in the districts north of the Quanza. M. Anchieta during his long residence in Angola never met with the species.

Bill yellowish white, brownish at the tips of the mandibles ; a brown line also runs along the commissure for half its length from the tip. A low crest rises at the base of the culmen and extends two thirds its length, with grooves on either side. Head, neck, upper part of breast, back, wings, and flanks black with a slight greenish gloss. Tail black, with the exception of the two lateral rectrices next to the outermost ones, which are pure white. Tarsus black.

Total length 18 inches, wing $9\frac{1}{4}$, tail 10, bill $3\frac{1}{4}$.

Specimens described and figured are in my own collection.

TOCKUS SEMIFASCIATUS.

ALLIED HORNBILL.

Buceros semifasciatus, Hartl. Journ. für Ornith. (1855) p. 356; id. Ornith. West-Afr. (1857) p. 163; Schleg. Mus. Pays-B. (1862) p. 13; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 130. sp. 7902; Giebel, Thesaur. Ornith. (1872) p. 503; Reichen. Journ. für Ornith. (1876) p. 43.

Tockus semifasciatus, Sharpe, Ibis (1869) p. 192. sp. 41; id. Cat. Afr. Birds (1871) p. 9. sp. 70; Reich. Journ. für Ornith. (1875) p. 13.

HAB. Senegambia to the Gaboon (SCHLEGEL); Fantee (USSHER).

Although closely resembling the *T. fasciatus*, the present bird may always be distinguished from it by having the two lateral rectrices next the outermost ones black on their basal half, instead of being white for their entire length. The geographical distribution of the two species is very similar. Of the habits and economy of the present one I have not found any thing recorded.

Bill.—maxilla for half its length from the point, tip of mandible, and a line along the commissure, running downwards towards the base, black; rest of bill bright yellow; the maxilla is much swollen at base, and rises into a low crest with a deep lateral groove on each side, descending rapidly towards the tip. Head, neck, back, wings, and flanks black, with a greenish gloss on back and wings. Tail black, with a greenish gloss, the two lateral rectrices next to the outermost one white for nearly the apical half of their length.

Total length $18\frac{1}{2}$ inches, wing $9\frac{1}{2}$, tail 10; bill $3\frac{1}{4}$, height at base $1\frac{1}{2}$.

Specimen figured is in the collection of the Paris Museum.

J.G. Keulemans lith.

TOCKUS FLAVIROSTRIS.

M & N Hanhart imp
9

TOCKUS FLAVIROSTRIS.

YELLOW-BILLED HORNBILL.

- Buceros flavirostris*, Rüpp. Faun. Abyss. (1835) vol. i. p. 6, tab. 2. fig. 2; G. R. Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 36; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 490; Ayres, Ibis (1871) p. 260. sp. 128; Schleg. Mus. Pays-B. (1862) p. 14; Von Heugl. Journ. für Ornith. (1864) p. 271; Finsch, Trans. Zool. Soc. vol. vii. p. 279; Giebel, Thesaur. Ornith. (1872) p. 498.
- Tockus flavirostris*, Bon. Consp. Gen. Av. (1850) p. 92. sp. 12; id. Consp. Vol. Anisod. (1854) p. 3; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) vol. ii. p. 596. sp. 877; Gurney, Anderss. Birds of Damara Land (1872) p. 210; G. R. Gray, Hand-l. Birds (1870) pt. ii. p. 129. sp. 7894; Sharpe, Cat. Afr. Birds, no. 77; Rüpp. Syst. Uebers. Vög. N.O.-Afr. (1845) p. 79. sp. 324; Layard, B. of S. Afr., Sharpe's ed. (1875) p. 130. sp. 121.
- Toccos elegans*, Hartl. Proc. Zool. Soc. (1865) p. 86, pl. iv.; Bocage, Journ. Sc. Math. Lisb. (1868) vol. ii. p. 347. sp. 99.
- Toccos flavirostris*, Barb. du Bocage, Journ. Sc. Math. Lisb. tom. iii. (1871) p. 270. sp. 16.
- Tockus elegans*, Bocage, Journ. Sc. Math. Lisb. tom. i. (1866) p. 335.

HAB. Abyssinia, Namaqualand, Damaraland (ANDERSSON); Transvaal (AYRES); Kuruman, Kanye (EXTON).

This bird has a very wide distribution, having been obtained from Abyssinia on the east to Damaraland on the west; and probably it will be found throughout the interior as our knowledge of the African avifauna is increased by future explorers. It is a well-marked bird, and conspicuous for its broad yellow bill. Hartlaub redescribed it in the 'Proceedings' of the Zoological Society as *T. elegans*, from specimens obtained by Monteiro. Andersson states that it is the most common of the Hornbills in Middle and Southern Damaraland. It is not gregarious, but is seen singly or in pairs, and, being rather fearless, is easily approached, especially during the heat of the day. At such times it resorts to the top of some tree, and keeps up a subdued chattering note of "töc, töc, töcke töcke, töcke töc," in a tone not unlike the yelpings of young puppies, at the same time flapping its wings and bowing its head. He says that in adult birds there is a considerable difference of size; the irides are yellow, legs and toes very dark brown. Speke states that this species is common in the Somali country, where it is fond of being in the jungle-trees, and is very noisy about sunrise, making a sound not unlike that of a frog.

Adult.—Bill much curved, no casque; upper part of maxilla much compressed laterally; orange-yellow, except along the commissure and tips, which are reddish brown. Top of head and neck dark leaden grey. Over the eye a broad white stripe. Throat and side of neck bare, yellow. Back brownish black, a line of white down the centre. The feathers of upper part of breast and wing-coverts are black, with a conspicuous white blotch near the tip. Secondaries light rufous-brown on the three innermost ones; next two white, irregularly marked with blackish brown;

remainder brownish black, as are also the primaries, the last having a small white spot upon both webs midway of their length. Underparts pure white. Tail has four central rectrices brownish black, remainder white, with black base and black bar near the tip; the outermost feather nearly all white, with very little black at the base; the white is very much less on the second; and on the third the black of the base extends upwards for more than half the length of the feather. Iris yellow. Total length without bill 16 inches, wing 7, tail $8\frac{1}{2}$, bill along gape $3\frac{1}{2}$, tarsus 1.

There is no difference in the plumage of the sexes. Young birds often have very dark bills.

A. G. Cammermeyer del.

TOCKUS HEMPRICHII

M. & N. Hanhart imp.
19

TOCKUS HEMPRICHII.

HEMPRICH'S HORNBILL.

- Buceros hemprichii*, Ehrenb. Symb. Phys. Av. (1828) fol. *aa*; Finsch, Trans. Zool. Soc. (1869) vol. vii. p. 317; Dresser & Blanf. Ibis (1874) p. 338.
- Buceros limbatus*, Rüpp. Faun. Abyss. (1835) vol. i. p. 5, tab. 2. fig. 1; G. R. Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 35; Schleg. Mus. Pays-B. (1862) p. 13; Von Heugl. Journ. für Ornith. (1864) p. 270; Finsch, Trans. Zool. Soc. (1869) vol. vii. p. 279; Giebel, Thes. Ornith. (1872) p. 500.
- Tockus limbatus*, Rüpp. Syst. Uebers. Vög. N.O.-Afr. (1845) p. 79. sp. 325; Bon. Consp. Gen. Av. (1850) p. 92. sp. 11; Horsf. & Moore, Cat. B. Mus. E.-Ind. Co. (1856-8) vol. ii. p. 596. sp. 879.
- Grammicus limbatus*, Bon. Consp. Vol. Anisod. (1854) p. 3.
- Lophoceros limbatus*, Cab. Mus. Hein. (1859-60) p. 168. sp. 6 (note).
- Tockus hemprichii*, Blanf. Geol. & Zool. Abyss. (1870) p. 326; Salvad. Cat. Uccelli Mare Rosso e dei Bogos, p. 54 (417).

HAB. Abyssinia, Senafé (JESSE).

This species was first described by Ehrenberg (*l. c.*) as *Buceros hemprichii*. This has been referred to the *B. (Lophoceros) nasutus* by authors; but Messrs. Dresser and Blanford, having examined the type in the Berlin Museum, stated in 'The Ibis,' 1874, that it was the same as the *B. limbatus*, Rüppell, described some seven years afterwards. Of course this last will have to become a synonym of the older name. It may be as well to remark that Mr. Blanford, in his 'Geology and Zoology of Abyssinia' (1870), had already stated that this species was the *B. hemprichi* of Ehrenberg. The latter's type was obtained in Abyssinia. It is a fine bird, rather large for the genus compared with the other members, and has not been met with, so far as I am aware, beyond the borders of Abyssinia. Cabanis and Heine, in their list of the Bucerotidæ in the Museum Heineanum, include the present species in *Lophoceros*; but as it has not like the members of that genus an elevated casque, it should not be placed with them.

Bill rather long, curved, compressed laterally, deep red. Head, neck, upper part of breast and back blackish brown. Wings black, margined with white. Underparts pure white. Tail blackish brown, the two lateral rectrices next to the outermost ones pure white. Iris brown.

Total length 22 inches, wing 12, tail $11\frac{1}{4}$, bill $5\frac{1}{4}$.

The naked space on the sides of the throat, according to Roth's notes in Horsfield and Moore's Catalogue of Birds, is black in the male and yellow in the female. With this exception both sexes are alike.

JG Keulemans lith.

M & N Hanhart imp.

TOKUS MONTEIRI.

TOCKUS MONTEIRI.

MONTEIRO'S HORNBILL.

Toccus monteiri, Hartl. Proc. Zool. Soc. (1865) p. 87, pl. 5.

Tockus monteiri, Gurn. in Anderss. Birds of Damara Land (1872) p. 208; Sharpe, ed. Lay. B. of S. Afr. (1875) p. 129. sp. 120.

Buceros monteiri, G. R. Gray, Hand-l. Birds (1870) part ii. p. 130. sp. 7900; Giebel, Thesaur. Ornith. (1872) p. 501.

HAB. Damara Land (ANDERSSON); Benguela, Angola (MONTEIRO).

This species is but little known, as it appears to be a rare bird even in its own country. It was described by Hartlaub (*l. c.*) from specimens obtained by Mr. Monteiro in Benguela, West Africa. It has also been procured by Andersson in Damara Land. A great difference exists in the size of the bill between the male and female, that of the latter being quite small in comparison with that of her mate.

Mr. Monteiro, in his work on Angola and the river Congo, states:—"The *Toccus elegans* (*T. flavirostris*) and the *T. monteiri* are very odd birds in appearance and habits. I found that their food consisted of grubs, grasshoppers, and other insects, hornets' nests and hard seeds. They dig in the sand with their long curved bills when seeking their food, throwing the sand behind them with their legs. They look very comical when sitting on a tree, their soft feathers puffed out like those of an Owl; and they raise and depress their crest-feathers, uttering loud, long-drawn, unearthly cries, like the squall of a sick baby." He goes on to say that the natives state that the female imprisons the male in a hollow of some tree, and obliges him to hatch out the eggs, thus reversing the habits of the other species of this family, and of most families generally. He endeavoured to verify this statement, and offered a large reward for a nest; but the natives were unable to discover one for him. It would seem probable that they had mistaken the sex, although the great difference in the size of the bill between the sexes of *T. monteiri* would cause either to be readily identified even at a considerable distance.

According to Andersson this species is not common in Damara Land, where it is generally found in pairs, though occasionally half a dozen may be seen in proximity to each other. Being shy, it is a difficult bird to approach, except in hot weather. It perches on the tops of trees, but at the slightest alarm betakes itself to flight, seldom, however, going far at a time, and progresses by dipping and rising alternately like a Woodpecker. He had often found much gravel in the stomach, and flushed it frequently from the ground, whither it resorts to pick up sand as well as food.

Adult Male.—Bill very long, curved, yellowish red, turning to dark purple near the tip. A deep groove runs along the entire length of the maxilla, following the curve of the culmen, which

is compressed laterally. Head and cheeks light grey, darkest on the edges of the occiput. Throat and breast plumbeous, each feather with a central streak of black. Wing light brown, a round white spot on the tip of each feather; innermost secondaries light brown, remainder white; primaries black, a white spot midway on the webs, and the tips slightly margined with white. Back and rump dark brown. Underparts and thighs pure white. The tail has the four central rectrices blackish brown, next one white for half its length from the tip, remaining ones pure white. Iris nut-brown. Feet and tarsus horn-colour. Total length without bill 17 inches; wing $8\frac{1}{4}$; tail 10; bill along culmen $5\frac{1}{2}$; tarsus 2.

The female resembles the male; but her bill is much shorter and more feeble, averaging only $3\frac{1}{4}$ inches.

JG Keulemans lith.

Hanhart imp.
22

TOCKUS GRISEUS.

TOCKUS GRISEUS.

JUNGLE GREY HORNBILL.

Grey Hornbill, Lath. Gen. Syn. (1787) Supp. p. 72.

Buceros griseus, Lath. Ind. Ornith. (1790) vol. i. p. 147; Shaw, Gen. Zool. (1811) vol. viii. p. 40; Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 594; Dumont, Dict. Sc. Nat. (1817) p. 212; Vieill. Ency. Méth. (1823) vol. i. p. 307; Less. Trait. Ornith. (1831) p. 252. sp. 3; Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 37; Schleg. Mus. Pays-B. (1862) p. 11; Gray, Hand-l. B. (1870) pt. ii. p. 131. sp. 7917; Giebel, Thesaur. Ornith. (1872) p. 499.

Buceros cinerascens, Temm. Plan. Col. vol. ii. (text); Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 31.

Tockus griseus, Bon. Consp. Gen. Av. (1850) p. 91. sp. 3.

Rhinoplax griseus, Bon. Consp. Vol. Anisod. (1854) p. 3.

Toccus gingalensis, Jerd. B. Ind. (1862) vol. i. p. 250 (partim); id. Ibis (1866) p. 350, (1872) p. 5.

Toccus griseus, Bourdillon, Str. Feath. (1876) vol. iv. p. 387.

Kaldal-haki, Can., *Chalotra*, Hindoostanee (JERDON).

HAB. Malabar.

This is the species described by Latham (*l. c.*) and mentioned by Jerdon in his 'Birds of India' as *Toccus gingalensis*. It is apparently quite a common bird in the localities it frequents; and the following extract from Dr. Jerdon's work sufficiently indicates the habits of the species. He says:—The "Jungle Grey Hornbill is found in the forests of Malabar and Ceylon. I have seen it nowhere else than in Malabar; and it is most abundant perhaps in the extreme south, in Travancore and in Ceylon. Its food consists of fruit of various kinds. It is rather a shy bird, is found singly or in pairs, occasionally in small parties, has a harsh call like the others, and is said to breed in the same way."

Ceylon, mentioned above as a habitat of this bird, is incorrect, as it is not found there, but is represented by the allied species *T. gingalensis*, Jerdon having confounded the two together. Mr. Bourdillon, who procured this species in the Travancore hills, says it is very common in heavy jungle at an elevation of from 1000 to 3000 feet. Its flight is rapid and easy; and the bird is very shy, and has a very ludicrous call, reminding one of a Punch-and-Judy show. The male had bright red irides; and a supposed female, in the Trevandrum Gardens, had pale-grey irides. Mr. Hume remarks that females procured in the Wynaad and on the Cardamon hills only differ in being paler and duller above and below, have less of the green gloss, and only three pairs of the rectrices are tipped with white. "The young bird," he continues, "has the irides dirty yellow; the bill is entirely pale yellow and smaller, and not overspread with the reddish brown tint of the adult, and has a black patch at the base of the lower mandible, and another on the basal portion of the culmen."

Bill strong, curved, culmen slightly swollen. Basal half pale red, becoming yellowish at tip, and black along the commissure. Top of head, back of neck, and back dark ashy-grey, each feather with a central white streak. Superciliary stripe pale greyish-brown. Front of neck and underparts dark grey, each feather with a central streak of greyish white. Primaries black, with an inch and a half of their tips white, and a white line on the central part of the edge of the outer webs. Secondaries black, broadly edged with grey on their outer webs. Rectrices black, margined with dark grey, the three outermost ones white for about two inches of their tips. Iris bright red. Feet dark grey.

Length 22 inches, wing $8\frac{1}{2}$, tail $8\frac{1}{2}$, bill $4\frac{1}{8}$, height at base $1\frac{1}{2}$.

Female.—Has the plumage like the male; the bill, however, is yellow, with the exception of the basal two thirds of the culmen, spot on lower edge of maxilla at base, and base of mandible, all of which are black.

The specimens figured and described were kindly loaned to me by Lord Tweeddale, to whom I desire to express my thanks.

JGKeulemans hth.

Hanhart im
20

TOCKUS GINGALENSIS.

TOCKUS GINGALENSIS.

CEYLONESE GREY HORNBILL.

Le Calao gingala, Le Vaill. Ois. Rares d'Amér. (1801) pl. 23.

Buceros gingala, Wilkes? Ency. Lond. (1808) vol. iii. p. 480.

Buceros gingalensis, Shaw, Gen. Zool. (1811) vol. viii. p. 37; Ersch u. Grub. Ency. (1824) p. 282; Temm. Plan. Col. (1824) vol. ii. p. 17; G. R. Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 30; Blyth, Cat. Birds Mus. Asiat. Soc. (1849) p. 44. sp. 184; Layard, Ann. & Mag. Nat. Hist. (1854) vol. xiii. p. 260; Schleg. Mus. Pays-B. (1862) p. 12; Gray, Hand-l. Birds (1870) pt. ii. p. 131. sp. 7918; Giebel, Thes. Ornith. (1872) p. 499.

Buceros gingala, Vieill. Nouv. Dict. Hist. Nat. (1816) vol. iv. p. 600; Dumont, Dict. Sc. Nat. (1817) vol. vi. p. 214.

Buceros pyrrhopygus, Wagl. Syst. Av. (1827) sp. 18.

Tockus gingalensis, Bon. Consp. Gen. Av. (1850) p. 91. sp. 4; Blyth, Ibis (1867) p. 296; Holdsw. Proc. Zool. Soc. (1872) p. 425; Legge, Ibis (1874) p. 14.

Rhinoplax gingalensis, Bon. Consp. Vol. Anisod. (1854) p. 3.

Buceros (Penelopides) gingalensis, Von Mart. Journ. für Ornith. (1866) p. 18.

Toccus gingalensis, Jerd. Ibis (1872) p. 5. sp. 145.

HAB. Ceylon, never Bengal (BLYTH).

The name of *Buceros gingala* was given to this species in the 'Encyclopædia Londinensis,' founded upon Le Vaillant's plate in his 'Oiseaux Rares d'Amérique et des Indes.' As the author of the ornithological part of the publication is not certainly known, the name cannot stand; and therefore that of Shaw, who described the bird as *B. gingalensis*, is the one that has priority.

The Ceylonese Grey Hornbill has been more fortunate than many of its relatives; for as most writers have employed Shaw's name, there is not much confusion in its synonymy. It is restricted to Ceylon, never having been found in Bengal, according to Blyth. My friend the late Dr. Jerdon, in his 'Birds of India,' confounded the species with its Indian ally, and spoke of both birds as *T. gingalensis*. In 'The Ibis' for 1872, p. 5, however, he corrected this error. In size and appearance the two birds are not unlike, the chief differences between them being found, as is the case with several allied species of Hornbills, in the markings and coloration of the tail-feathers.

Layard states that "the Lesser 'Hornbill' is common in the Wanny about Anarajahpoora and in the Mookalane jungles of the southern provinces. It feeds, on the tops of the loftiest trees, upon fruits and berries, which it swallows whole. It is a wary and shy bird; and although its presence is often revealed by its loud harsh call, it rarely falls before the hunter's gun; and the best way to procure it is to lie concealed near a tree in fruit, if it be such as it feeds upon. The irides are reddish, and when partly hidden by the long stiff black eyelashes have a very peculiar appearance. In some specimens the bill is white, with a black patch extending from the naked space round

the eye about three fourths of an inch along the lower half of the upper mandible ; the bill is $3\frac{3}{4}$ inches long. The three outer tail-feathers are white, the fourth half black from the quill, the fifth black. The head has a rufous tinge.

"In other specimens the head wants the rufous tinge, the first tail-feather is white, with the outer shaft black up two thirds of its length, and slightly tinged up one third of the inner web ; second and third feathers black on both sides up two thirds of their length ; fourth black up to an inch from the top ; fifth black altogether. This species (?) is found in the southern provinces about the base of the hills ; the former in the Anarajahpoora Wanny."

Lieut. Legge states, "This species ranges from the highest down to the lowland forests, frequenting the tallest trees in them, and is more numerous in the latter than in the mountains or in any part of Ceylon that I have as yet explored."

Male.—Bill greenish white, becoming brownish at base. Head crested, feathers moderately long, dark brown, with a white central line. Back of neck and upper parts brownish grey. Wings light leaden grey, each feather margined with black. Primaries greenish black, the second, third, fourth, and fifth tipped with white ; secondaries greenish black, the outer webs margined with leaden grey. Throat, breast, and abdomen buffy white, becoming dark buff on the lower part of abdomen and under tail-coverts. Tail greenish black, the three outermost rectrices having their apical third pure white, the one next the central pair tipped with white. Irides reddish brown.

Total length without bill, average $19\frac{1}{2}$ inches ; wing 8, tail 10, bill along culmen 3, tarsus $1\frac{1}{2}$.

Female like the male in the plumage, but has the bill black, with a yellowish-white line occupying the lower half of the maxilla along the commissure and reaching to within an inch of the tip.

I am indebted to the Marquis of Tweeddale, who kindly loaned to me from his collection the specimens from which my figures and descriptions were taken.

J.G. Keulemans lith.

TOCKUS ERYTHRORHYNCHUS.

Hanhart imp.
36

TOCKUS ERYTHORHYNCHUS.

RED-BILLED HORNBILL.

- Calao à bec rouge du Sénégal*, Buff. Plan. Enlum. (1783) no. 260.
- Buceros nasutus*, var. B, Gmel. Syst. Nat. (1788) p. 361; Lath. Ind. Ornith. (1790) vol. i. p. 145.
- Le Calao Toc*, Le Vaill. Hist. Nat. Ois. d'Afrique. (1806) vol. v. p. 122, pl. 238.
- Buceros nasutus*, Vieill. (nec Linn.) Ency. Méth. (1823) vol. i. p. 305, pl. 10. fig. 3.
- Buceros erythrorhynchus*, Temm. Plan. Col. (1824) vol. ii. sp. 19 (text); Ersch u. Grub. Ency. (1824) p. 284; Wagl. Syst. Av. (1827) sp. 20; Less. Trait. Ornith. (1831) p. 252. sp. 1; Thom. Exp. Niger, vol. i. p. 250; Gray, Gen. Birds (1849) vol. ii. p. 400. sp. 34; Hartl. Ornith. West-Afr. (1857) p. 165; Schleg. Mus. Pays-B. (1862) p. 14; Layard, B. of S. Afr. (1867) p. 227; Finsch u. Hartl. Reis. Ost-Afr. (1867) p. 491; Ayres, Ibis (1869) p. 296; Finsch, Trans. Zool. Soc. (1869) vol. vii. p. 278; Giebel, Thesaur. Ornith. (1872) p. 498.
- Buceros (Alopius*, non Schönh. 1826, Coleopt.) *erythrorhynchus*, var. *leucopareus*, Hemp. & Ehrenb. Sym. Phys. Av. Dec. fol. aa, note 1.
- Buceros leucomelas*, Licht. Verz. Säugeth. und Vög. (1842) p. 17; Gray, Hand-l. B. (1870) pt. ii. p. 130. sp. 7895.
- Tockus erythrorhynchus*, Rüpp. Syst. Übers. Vög. N.O.-Afr. (1845) p. 79. sp. 322; Bon. Consp. Gen. Av. (1850) p. 92. sp. 10; id. Consp. Vol. Anisod. (1854) p. 3; Hartl. Journ. für Ornith. (1854) p. 193; Horsf. & Moore, Cat. Birds Mus. E.I. Co. (1856-8) vol. ii. p. 575. sp. 876; von Heugl. Journ. für Ornith. (1864) p. 271; Kirk, Ibis (1864) p. 327; Bocage, Journ. Sc. Math. (1866) p. 335. sp. 33, (1868) vol. ii. p. 348. sp. 101; Gray, Hand-l. Birds (1870) pt. ii. p. 129. sp. 7893; Gurney, Anderss. B. Damara L. (1872) p. 211; Sclat. Rev. List Vert. Anim. (1872) p. 172; Salvad. Uccelli, Mar. Rosso e di Bogos, p. 54 (1874); Layard, B. of S. Afr., Sharpe's ed. (1875) p. 131. sp. 122; Bocage, Ornith. Angola (1878) p. 120.
- Buceros rufirostris*, Sundev. Öfvers. Kongl. Vetensk. Akad. Förh. (1850) p. 108.
- Buceros erythrorhynchus*, var. *caffer*, Sundev. Öfvers. Kongl. Vet. Akad. Förh. (1850) p. 50; Von Heugl. Ibis (1859) p. 343. sp. 58.
- Tockus pæcilorhynchus*, Bon. Consp. Gen. Av. (1850) p. 91. sp. 9.
- Rhynchoceros erythrorhynchus*, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 166. sp. 1.
- Rhynchoceros leucomelas*, Cab. & Hein. Mus. Hein. (1860) Th. ii. p. 166. sp. 2.
- Tockus erythrorhynchus*, Buckley, Ibis (1874) p. 365.
- Kicumbiandondo* in Angola (ANCHIETA).

HAB. Senegambia to Sennaar (SCHLEGEL); Damaraland, Ondonga, Okovango river, Lake Ngami (ANDERSSON); Zambesi region (KIRK); Angola (ANCHIETA).

This is a well-known bird, and has received many appellations, as a glance at the above list of synonyms will testify. Its range is very extensive, quite across the African continent from Sennaar

on the east to Damaraland on the west. Sundevall gives it as a native of Caffraria; Layard says he had not heard of it there, but had received it in considerable numbers from the Zambesi; Kirk states that the native name for this species is "Kope," and that they are common in all woodland districts, feed on fruit and insects, and breed in hollow trees. Andersson, in his 'Notes on the Birds of Damara Land,' edited by Mr. Gurney, speaks of the differences observable in the plumage of this species from separate localities, for one of which Sundevall proposed a specific name.

He says "This species is common in Ondonga, at Okovango River, and for some distance to the south of that stream; and I have also obtained specimens from Lake Ngami. I have also met with it in Damara Land proper, at Objimbinque and Schmelen's Hope; but specimens from these two last-named localities differ considerably from those found in more northern parts. Thus in the former the whole of the underparts, the forehead, a broad band under the eyes continued down the sides of the neck, the ears, cheeks, chin, and throat are of a uniform silky white; whilst in the more northern bird the colour of these parts, as well as of the breast, is mingled with blackish grey, and there is also less white about the wings and tail. This Hornbill is frequently seen searching for food upon the ground; and the way in which it swallows some kinds of food is peculiar, raising its head and pitching the morsel into the air, receiving it again into its bill, and repeating the process several times, perhaps with the object of softening the food or reducing it to a pulp."

It is difficult to say what causes the variation in the plumage spoken of by the author in the specimens from Objimbinque and Schmelen's Hope. I have examined examples from both these localities, and do not consider that I should be justified in recognizing two species of them. Prof. Sundevall has regarded the bird with grey cheeks as the Caffrarian race, and proposed the term *rufirostris* for it; I have not adopted his name, as it does not appear to me to be necessary to make a specific distinction between the forms. The late M. Jules P. Verreaux, as quoted by Mr. Gurney, considered that there is only one species, and that the pure white cheeks are assumed only in the nuptial season. This is very possibly the fact.

In 1828 Ehrenberg proposed the term *Alophius*, as a subgenus, for this species. But in 1826 Schönherr gave the same name to a genus of Coleoptera; and as this has been generally employed in Entomology, it cannot be adopted for this bird and its allies.

Adult.—Bill curved, no casque, maxilla deep red, much compressed laterally along the culmen. The mandible for two thirds its length from the base is very dark red, almost black, remainder like the maxilla. Upper part of head and neck leaden-grey. A white superciliary stripe extends along the side of the head and down the neck. Cheeks and neck beneath the white band leaden-grey mingled with white; in some specimens these parts are pure silky white. Wings blackish brown, each feather broadly tipped with white. Innermost secondaries light brown, next two blackish, broadly tipped and marked with white; remainder black margined with white. Primaries black, first two spotted midway with white on inner web, next four on both webs, remainder uniform black. Entire underparts pure white. The tail has the two median feathers brownish black, next

on either side jet-black; remainder black, broadly tipped with white, the outermost being all white, with the exception of a short distance from the base (chiefly on the inner web) and a narrow bar on the apical third part. Iris light yellow.

Total length without the bill 16 inches, wing 8, tail 9, bill along gape 3.

There is no difference in the plumage of the sexes. The female is a little smaller, with a shorter bill.

The description and figures were taken from specimens in my own collection.

J.G. Keulemans lith.

Hanhart imp.

TOCKUS DECKENI.

TOCKUS DECKENI.

VON DER DECKEN'S HORNBILL.

Buceros (Rhinoceros) deckeni, Cab., v. d. Decken, Reis. vol. iii. p. 37, t. vi.; Finsch & Hartl. Reis. Ost-Afr. (1867) p. 489.

Buceros (Tockus) deckeni, G. R. Gray, Hand-l. Birds, (1870) pt. ii. p. 130. sp. 7896.

Rhinoceros deckeni, Cab. Journ. für Ornith. (1870) pl. 2.

Buceros deckeni, Giebel, Thesaur. Ornith. (1872) p. 498.

HAB. East Africa (FINSCH & HARTLAUB); Voi-Fluss (Taita), East Africa (HILDEBRANDT).

This very distinct species was procured by Baron von der Decken during his journey into Eastern Africa, and described by Finsch & Hartlaub (*l. c.*). Its white cheeks, neck, and underparts make it impossible to be confounded with any other species of this genus. Nothing is known of its economy or habits. The specimen described and figured is in the collection of my friend Captain Shelley, who most kindly loaned it to me to be figured in this work.

Top of head and back black. Superciliary stripe, sides of the head, neck, stripe down middle of the back, and underparts white, tinged with buff on throat and breast. Ear-coverts grey. Primaries black, with an oblong white spot upon the outer web near the centre. Secondaries black, edged with white, some of them pure white, with an irregular black mark in the centre, the innermost ones brownish. Four central rectrices greyish black, the next two black for two thirds their basal length, with the remaining portions white; the external rectrix black for only one third the basal length, rest pure white. Bill deep red for half its basal length, rest yellow, with the cutting-edges and tips brownish black. Maxilla has the basal half covered by a scale, with slight horizontal grooves.

Total length $17\frac{1}{2}$ inches, bill on culmen 4, wing $7\frac{3}{4}$, tail $9\frac{3}{4}$, tarsus $1\frac{3}{4}$.

J.G. Keulemans lith.

TOCKUS HARTLAUBI.

Hanhart imp.

TOCKUS HARTLAUBI.

HARTLAUB'S HORNBILL.

Tockus hartlaubi, Gould, Proc. Zool. Soc. (1860) p. 380.

Buceros nagtglasii, Schleg. Neder. Tijdsch. voor de Dierk. (1862) vol. i. p. 56, pl. 2; id. Mus. Pays-B. (1862) p. 16;

Sclater, Ibis, (1863) p. 359, (1864) p. 398; G. R. Gray, Hand-l. Birds, (1870) pt. ii. p. 130. sp. 7904.

Tockus hartlaubi, Sharpe, Ibis, (1870) p. 485. sp. 182.

HAB. Gold Coast (SCHLEGEL).

This is one of the most diminutive species of this family, and was described first by Mr. Gould (*l. c.*). Afterwards Schlegel (*l. c.*) redescribed it as *nagtglasii*. These are the only names that have been given to the bird. It is very distinct, and not to be confounded with any other species. Thus far it has only been obtained on the Gold Coast.

Bill slate-colour, red at tip; culmen much compressed at base, and rising into a low keel-like crest. Head black, a bare spot on cheek and throat, in dried skin yellow. Superciliary stripe and feathers on the occiput white. Underparts dark grey, each feather tipped with silvery white. Back greyish black. Wing dark green; primaries dark purplish brown. Tail greenish black, the outer rectrices tipped with white.

Total length 14 inches, wing 6, tail $7\frac{1}{2}$, bill $2\frac{1}{2}$, height at base 1.

The specimen described and figured is in the British-Museum collection.

J.G. Keulemans lith

TOCKUS CAMURUS.

Hanhart imp.

Tockus camurus
(1860)
Buceros pulchellus
pl. 4, O.
Tockus pulchellus
für Orn.
Buceros camurus
p. 497.

HAB. G.

Mr. Ca.
in 1856; an
'script' for
Africa, and
species that
appears to
but is much

Bill bri
Head, breast
white, form
both webs:
shafts and c
Total le
Specime

TOCKUS CAMURUS.

CASSIN'S PYGMY HORNBILL.

Tockus camurus, Cassin, Proc. Acad. Nat. Sci. (1856) p. 319, (1859) p. 140. sp. 164; Heine, Journ. für Ornith. (1860) p. 188. sp. 146; Reich. Journ. für Ornith. (1875) p. 13.

Buceros pulchrirostris, Schlegel, Soc. Nat. Art. Mag. Amster.; id. Ned. Tijds. voor d. Dierk. vol. i. (1862) p. 74, pl. 4, Ois.; id. Mus. Pays-Bas, (1862) p. 15; G. R. Gray, Hand-l. Birds, pt. ii. (1870) p. 130. sp. 7903.

Tockus pulchrirostris, Sharpe, Ibis, (1870) p. 485. sp. 181; id. Proc. Zool. Soc. (1871) p. 604. sp. 8; Reich. Journ. für Ornith. (1875) p. 13.

Buceros camurus, G. R. Gray, Hand-l. Birds, pt. ii. (1870) p. 130. sp. 7905; Giebel, Thesaur. Ornith. vol. i. (1872) p. 497.

HAB. Gold Coast (SCHLEGEL); Cape Lopez (DU CHAILLU).

Mr. Cassin first described this species in the 'Proceedings of the Academy of Natural Sciences' in 1856; and it was subsequently renamed *pulchrirostris* by Schlegel in the 'Nederlandsch Tijdschrift' for 1862. Mr. Cassin's type was obtained by Du Chaillu near Cape Lopez, in Western Africa, and sent with other rare and new birds to Philadelphia. Mr. Cassin remarks of this species that it "is the smallest bird of the genera *Buceros* and *Tockus* that I have ever seen, and appears to be the smallest known species. It resembles in some measure *T. melanoleucus*, but is much smaller. Three specimens are in the collection, essentially alike."

Bill bright red; maxilla greatly compressed laterally; culmen elevated into a low crest. Head, breast, and back rufous brown. Wings dark brown, lesser and greater coverts tipped with white, forming two rows. Primaries brownish black, with an oval buff spot about the centre on both webs; secondaries blackish brown, broadly margined with deep buff. Tail dark olive-brown, shafts and edges of webs and tips rufous brown. Underparts white. Feet dark brown.

Total length $12\frac{1}{2}$ inches, wing 6, tail $6\frac{3}{4}$, bill $2\frac{1}{4}$.

Specimen figured is in the British-Museum collection.

2/7/-

1/20/51

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00286200 1

crisrb f0L696.C7E46

A monograph of the Bucerotidae, or family