SF

487

.U6

1919 a

Class SF 487

Book .U6

1919a

Culling the Poultry Flock

Why, When, and How

The Hen That Lays is the
Hen That Pays

UNITED STATES DEPARTMENT OF AGRICULTURE
DEPARTMENT CIRCULAR 31

Contribution from the Bureau of Animal Industry
Animal Husbandry Division

Washington, D. C.

Issued May 2, 1919; reprint, October, 1919

WASHINGTON : GOVERNMENT PRINTING OFFICE : 1919

Monograph

LIBRARY OF CONGRESS
RECEIVED
OCT 30 1933
DIVISION OF DOCUMENTS

SF487

.76

1919a

CULLING THE POULTRY FLOCK.

WHY CULL?

Intelligent culling of the poultry flock is of the utmost importance for success in poultry keeping. Culling serves three purposes: First, it increases the profits by insuring that the feed will be consumed by the better-producing hens, the profit payers, and will not be consumed by the poor producers which are kept at a loss. Second, it makes it possible to save those best suited for breeders, both on account of their better production and on account of their superior strength and vitality, which have enabled them to stand up under the severe strain of heavy laying. Third, weeding out the poor hen gives those left more room and a better chance. Where trap nesting is practiced, culling is a comparatively simple process, as the recorded egg record enables one to determine whether a hen should be retained or discarded.

WHEN TO CULL.

Culling should be continuous throughout the year. This continuous culling should consist in weeding out, when discovered, any hen which is sick, which is very thin, or which shows evidences of non-production, weakness, or poor vitality.

The whole flock should also be given a careful and systematic culling at least once during the year. The hens should be handled individually and gone over carefully with the object of dividing them into two lots, one the better producers and the other the poorer producers. From the better producers it is also desirable to pick out and band or otherwise mark as many of the best as will be needed for breeders. Market those selected as the poor producers. Save for laying and breeding those selected as the better producers.

When a single systematic culling is made, the best time to do it is in August or September. At this time it is easier to form a close estimate of the relative value of a hen as an egg producer and to weed out the poor producers. Hens which show indications of laying at this time are those which on the average have been the better producers for the year. It must be remembered, too, that the better producers during the first laying year are usually those which will be the better producers in subsequent years. Hens showing indications of having been good producers throughout the year should be

retained for the next year regardless of their age, but relatively few hens will prove to be profitable producers beyond their second laying year if of the heavier breeds, such as the Plymouth Rock, Rhode Island Red, Wyandotte, or Orpington, or beyond their third laying year if of the lighter breeds, such as Leghorn, Ancona, or Minorca. Preliminary culling during July is also desirable in order to eliminate hens which have started to molt and have stopped laying at that time.

HOW TO CULL.

In going over the entire flock for the purpose of culling there are a number of points or characteristics which should receive special attention in selecting the layers from those which are not laying at the time mentioned. If the different characteristics or several of them in the case of any individual agree as indicating good production or poor production, selection is comparatively accurate. If they do not agree, judgment must be used in deciding which should receive the greatest weight. The following are the main points to consider:

Molt.—This is probably the most valuable and most easily applied test of production. Hens cease laying completely or almost completely during the molt. The better producers lay late in the fall, and therefore molt late. Late molters also molt rapidly as a rule, while early molters molt slowly. Therefore, save the hens which have not molted by August or are only just beginning to molt late in September or in October, and discard those which have finished molting or are well into the molt. The hens which molt latest, provided they are otherwise desirable, are the best to save for breeders. A hen which has not molted at that time is characterized by soiled and worn or broken plumage, which is especially evident in the tail plumage, while those molted or molting show clean, fresh plumage or pinfeathers or partly grown feathers. Culling the early molters just as soon as they begin to molt is one of the easiest ways to weed out the poor layers.

Shank color.—In those breeds having yellow legs, the color fades out slowly as the laying season advances. The rapidity and degree to which the yellow color is lost depends to a considerable extent upon the heaviness of laying. The yellow color goes out from the scales on the front of the shanks first and finally from the scales on the rear.¹ Hens on grass range do not lose the shank color so quickly or completely as those in bare yards. Some soils also tend to bleach the color of the legs. Hens showing strong or medium yellow-shank color in August and September are usually poor layers, but occasionally poor layers may show pale or white shanks. A sick hen or

¹ Taken from an outline on judging fowls for egg production, prepared at a judging school held at Cornell University.

one in poor condition may also show pale shanks. When hens stop laying they begin to regain the yellow-shank color. Hens which have been broody and are therefore not laying, may show more or less yellow in shanks, but if they are not molting should not be discarded on that account. In breeds such as the Rhode Island Red where the shanks may show horn color as well as yellow, the horn must not be confused with the yellow. The rear and sides of the shank show little of this horn color.

Beak color.—In breeds having yellow beaks, the same principle applies as in the case of the yellow shanks. However, the color is lost from the beak more quickly than from the shanks and is also regained more quickly. The lower beak bleaches faster than the upper, but may be used where the upper is obscured by horn or black.¹

Vent color.—In yellow-skinned breeds the yellow color of the skin immediately about the vent is quickly lost with laying, and is quickly regained after laying ceases. White or pink vent color generally indicates that the hen is laying, while a yellow vent means that the bird is not laying.

Appearance of vent.—The vent of a hen laying heavily is large, expanded, and moist, while that of a hen not laying is comparatively small, hard, puckered, and dry.

Comb.—When a hen is laying or about to lay her comb is waxy,² full of blood, and red in color. When not laying, the comb is smaller and shrunken, paler or duller red in color, comparatively hard, and covered with whitish scales. A dark or bluish color usually indicates sickness. The changes in the wattles and ear lobes are similar to those of the comb, but not so marked. It is necessary to keep in mind the difference in size of comb of the various breeds.

Pelvic bones.—As a hen stops laying there is a tendency for her to take on fat. This is noticeable in examining the pelvic bones, the two bones which can be felt as points on either side of the vent. When the hen is laying, these bones are comparatively thin and flexible. When she is not laying, they feel thicker and less flexible, due to the fat which has accumulated there. Often poor producers are somewhat beefy in type and, therefore, show stiffer, thicker pelvic bones even when in laying condition than do good producers. The spread or distance apart of these pelvic bones is also a valuable indication of whether the hen is laying. When laying they are wider apart than when the hen is not laying. The spread can be roughly measured for practical purposes by determining how many fingers can be laid between the bones. If the spread measures two fingers or less, the

¹ Taken from an outline on judging fowls for egg production, prepared at a judging school held at Cornell University.

² Cornell Extension Bulletin 21.

probabilities are that the hen is not laying, while if the spread is greater, she is probably laying.

Distance from pelvic bones to keel bone.—A hen laying well is a good eater. Her intestines are, therefore, fuller and more distended, and require more room than when she is not laying and not eating so much.¹ When laying, the ovary and oviduct are of greater size and require more room. To provide this extra room the distance from the rear end of the keel to the pelvic bones increases, with a consequent increase in the size of the abdomen. A spread of 3 or more fingers in the smaller breeds, such as the Leghorn, and 4 or more fingers in the larger breeds, such as the Plymouth Rock, indicates that the hen is in a laying condition. A spread of less than 3 fingers in the smaller breeds and less than 4 fingers in the larger breeds indicates that she is not in a laying condition. The comparative size or spread of the abdomen of hens in laying condition, together with the length of the keel, is a measure of value with respect to the capacity or ability to lay. Mere spread of abdomen may be misleading, as a hen with a short keel may show a large spread of abdomen and yet have poorer capacity than one with a longer keel but smaller spread of abdomen.

Flexibility of abdomen.—When the hen is laying the greater size of the abdomen, together with the lessened tendency to accumulate fat at this point, results in a soft, flexible abdomen, suggesting, when handled, the texture of a partly milked-out udder of a cow. When laying has ceased, the abdomen grows smaller, or contracts, and it feels, when handled, harder and less flexible.

In culling the flock remember that it is safer to depend upon the agreement of a combination of several characteristics rather than to select by any one alone. It is also necessary to give consideration to the general condition of the flock and the care and feeding which they receive and the conditions under which they are kept. For example, a flock which has free range and plenty of green feed may have produced well and still show considerable yellow-shank color. In culling such a flock not so much weight should be given the presence of yellow in shanks as in one which has been confined to a bare yard and where in consequence a general paler leg color has resulted. With this in mind—

CULL THESE HENS.

Sick, weak, lacking vigor, inactive, poor eaters, early molters, with small, puckered, hard, dry vents; with small, shriveled, hard, scaly, dull-colored combs; with thick or coarse, stiff pelvic bones, that are close together, small spread between pelvic bones and rear

¹ Cornell Extension Bulletin 21.

end of keel, and hard, small abdomen. In breeds with yellow skin and shanks, the discarded hens should also show yellow or medium-yellow shanks and yellow beaks and vents.

SAVE THESE HENS.

Healthy, strong, vigorous, alert, and active; good eaters; not molting or just beginning to molt in September or October; with large, moist vents; with full, red combs; thin, pliable pelvic bones well spread apart, widespread between pelvic bones and rear end of keel, and large, soft, pliable abdomen. In breeds with yellow skins and shanks, the hens saved should also show pale or white shanks, and pale or white beaks and vents.

KEEP STANDARDBRED POULTRY

Standardbred poultry is more uniform in size, type, and color.

Standardbred poultry is more attractive in appearance and appeals more strongly to purchasers of stock and eggs.

Standardbred poultry offers a greater combination of practical and utility qualities suitable to the needs of the farmer and poultry keeper.

Standardbred products are more uniform in quality, are in greater demand, and bring better prices.

Standardbred poultry means greater success and better profits.

LIBRARY OF CONGRESS

0 002 851 188 7 ●