1932

College
of New York.

Library.

GIVEN BY

Dr. N. L. Brittin.

Contents.

589.2
24

- No. 1. Fries, O.R. - Om svampbildningar på menniskans hud. 1867.
- No. 2. ✓ Peck, C.H. - ... New species of fungi.
- No. 3. " , " - Colorado fungi.
- No. 4. ~~✓~~ " , " - New species of fungi.
- No. 5. ✓ Johnson, A.E. M.D. - Fungi.
- No. 6. ✓ " , " " - Botany for 1877.
- No. 7. ✓ Cooke, M.C. - Maine fungi.
- No. 8. ✓ " , " - Hyphomycetous fungi of the U.S.
- No. 9. ✓ Barbey, W. - Champignons. 1881.
- No. 10. Harkness, H.W. M.D. & Moore, J.P. - Pacific coast fungi. 1880. QK605.5. P3 H37
- No. 11. ✓ Seymour, A.B. - ... Fungi... along the Northern Pacific railroad.
- No. 12. ✓ " , A.B. - ... Fungus diseases on small fruits.
- No. 13. ✓ Ward, H.M. - ... Sexuality of the fungi.
- No. 14. ✓ Scribner, F.L. - Fungicides. 1888.
- No. 15. ✓ Pound, R. - Synchrony.
- No. 16. Giard, A. - De insectorum morbis qui fungis parasitis efficiuntur.
- No. 17. ✓ Seymour, A.B. - Injuries produced by parasitic fungi. 1887.
- No. 18. ✓ Bessey, C.E. - Saccardo's Sylloge fungorum. 1890.

- No. 19 ✓ Staten Island fungi. 1890.
- No. 20 ✓ Fairman, C.E. - Fungi of western N.Y.
- No. 21 - Voglino, P. - Nota micologica. skip
- No. 22 ✓ Chester, F.D. - Three new... diseases of plants.
- No. 23. Pammel, L.H. - Bekämpfung der pilzkrankheiten mit Bordeauxmischung. 1891. skip.

ÖM

SVAMPBILDNINGAR PÅ MENNISKANS HUD

OCH

DERAF FÖRORSAKADE SJUKDOMAR.

AF

OSCAR ROBERT FRIES.

Upsala

Edquist & Berglund. 1867.

Om Svampbildningar på människans hud och deraf förorsakade sjukdomar.

Af

Oscar Robert Fries.

Under den mäktiga utveckling, som Medicinen på sednare tider snabbare än någonsin erfarit — sedan allvarlig, rastlös forskning allestädes intagit de lockande men tomma hypotesernas plats —, har den skyndat att från sina hjälpvetenskaper tillägna sig och vidare utveckla resultaterna af de äfven der allt djupare trängande undersökningarne. Här af förklaras, att nästan hvarje vigtigare upptäckt på naturvetenskapernas område äfven inom Medicinens historia betecknas af en mer eller mindre ingripande förändring, ett större eller mindre steg framåt.

Till de talrika beröringspunkter mellan Medicin och Botanik, hvilka redan från deras första barndom nära förenat de begge vetenskaperna med hvarandra, kom, sedan microscopet öppnat insigt i de lägre organismernas byggnad och lefnadsförhållanden, genom Bassi's upptäckt af muscardinet (1835) ett nytt uppslag i frågan om vissa sjukdomsformers ursprung och rätta natur. Väl hade redan förut någon gång på djur anträffats svamparter (t. ex. *Sporendonema muscæ*, *Isariæ*), men först genom de svåra härjningar, som den af *Botrytis Bassiana* bland silkesmasken förorsakade farsot åstadkom, fästes större uppmärksamhet vid detta ämne. Härigenom gafs impulsen till Schönlein's och Gruby's m. fl. upptäckter af hudparasiterna, för hvilka iakttagelsers vidare utveckling vi i denna uppsats skola redogöra, Berg's af torsksvampen o. s. v. — Snart gick man likväl för långt och lät hypoteserna ersätta facta — och sedan Henle, utan stöd af observationer, genom uppställande af en botanisk parasit-theori sökt förklara de miasmatiske och contagiösa sjukdomarne, inträdde en reaction, som nästan helt och hållet uppgaf undersökningarne på detta fält och tillerkände ætiologisk betydelse egentligen endast åt hudparasiterna, stundom ej ens åt dessa.

Emellertid började för något mer än ett tiotal år tillbaka, i synnerhet genom Tulasne's upptäckter, en ny uppfattning af de lägre svamparnes metamorphoser göra sig gällande, hvaraf väcktes en ny, särdeles liflig verksamhet i deras studium. Samtidigt härmed verkställde Pasteur sina berömda, epochgörande undersökningar, hvilkas resultat, sammanfattadt i orden: "utan svampar ingen jäsnings, ingen förruttelse", af talrika efterföljare blifvit bekräftadt (ehuru visserligen äfven motsagdt) och naturligtvis mäktigt måste ingripa ej mindre i Medicinen, än i mångfaldiga förhållanden inom det borgerliga lifvet.

Det dröjde icke länge, förrän äfven inom Medicinen verkningarna af detta nyväckta lif förspordes — och af de på sista tiden hopade iakttagelserna om växtparasiter¹⁾, såsom orsakande eller åtminstone åtföljande flera de mest olikartade sjukdomar, skulle man kunna frestas att tro Jahn-Stark's dunkla och sväfvande parasit-theori — om ock i förändrad gestalt — börja blifva verklighet. Vi vilja korteligen omnämna de viktigaste af dessa uppgifter, som ligga utom vårt egentliga ämne.

Fullkomligt afvikande från alla de öfriga och genom talrika observationer, såsom det synes, satt utom tvifvel är den märkvärdiga s. k. "*Madura-Foot*", "*Mycetoma*"²⁾, en sjukdom, som i vissa delar af Ostindien förekommer ej sällsynt. Här skall svampen (*Chionophye Carteri* Berk.) uppträda i den subcutana bindväfven på foten, under bildande af smärre, fasta, rörliga knölar, och derifrån i alla riktningar sprida sitt mycelium — likasom en *Rhizomorpha* —, hvarunder foten enormt förtjockas och antager en oformlig, rundad gestalt. Under tiden uppkomma ulcerationer (mellan hvilka huden är normal), och ur dermed communicerande långa, sinuösa canaler uttömmas, blandade med var, en mängd svartgråa svampconglomerater af ett knappålhufvuds till en bösskulas storlek. Dylika träffas allestädes i bindväfven mellan musklerna och i den spongiösa bensubstansen. Sjukdomen kan vara 10 år och deröfver, sprider sig till underbenet samt

¹⁾ Af olika författare hänföras de ifrågavarande microscopiska organismerna dels till de lägre djuren, under namn af "vibrioner", "monader", "bacterier" o. s. v., dels till växtriket; och der antingen till Algerna ("*Palmellæ*", "*Leptothrix*") eller Svamparne. Emellertid är det enligt Richter, Hallier m. fl. sannolikt, att de i flertalet fall äro af vegetabilisk natur och tillhöra Svamparne (såsom af Hallier redan om *Leptothrix* m. fl. är bevisadt). — För öfrigt hänvisa vi med afseende på denna fråga äfvensom rörande mögelsvamparnes utvecklingshistoria m. m. dem, som sakna tillgång till den botaniska litteraturen i ämnet, till Richters korta sammandrag i "Schmidt's Jahrb." 1867, *M* 7.

²⁾ Se Hirsch i "Virch. Arch." b. XXVII.

medför marasm och död, der ej den enda hjälpen — amputation — mellankommer.

I *Cholera*-uttömningarne, der redan 1849 af Swayne, Brittan m. fl. svampar anträffades men ånyo råkade i glömska, hafva på sista tiden samtidigt af Thomé¹⁾ och Klob²⁾ rikliga svampbildningar constant blifvit iakttagna och närmare studerade samt framställda såsom egande ett intimt (troligen caussal)-sammanhang med sjukdomen. Hallier³⁾ kommer på grund af en mängd odlingsförsök till den åsigt, att svampen är en utvecklingsform af den på sädesslagen parasitiska *Ustilago* (*Tilletia*), hvilken form för sin utveckling fordrar en högre grad af värme och fugtighet. Härmed förbinder han åtskilliga lösliga hypoteser, såsom att densamma utom tarmcanalen sannolikt endast i Indiens heta klimat kan utvecklas på risväxten (eget nog har äfven cholera kallats "morbus oryzeus") o. s. v.⁴⁾

Äfven vid *Cholera nostras*, *Diarrhé*, *Dysenteri*, *Typhus* m. fl. äro svampbildningar anträffade i uttömningarne (jfr Richter & Klob ll. cc.) —, hvilket dock föga bevisar, då, enligt Hallier, dylika, ehuru i ringa mängd, äfven pläga åtfölja normala faeces.

Öfver "*Palmellæ*" såsom orsak till *fross-miasma* har Salisbury⁵⁾ meddelat talrika märkvärdiga iakttagelser. "Sporerna", hvilka genom inandning eller med dricksvattnet inkomma i organismen, öfvergå i circulationen, fortfara att utveckla sig och aflägsnas hufvudsakligen med urinen. De skola isynnerhet inverka fiendtligt på epithelialcellerna och förgifta deras producter, företrädesvis mjeltens, lefverns och mesenterial-körtlarnes, samt derigenom medelbarligen hela organismen, hvars reaction mot det giftiga ämnet yttrar sig genom paroxysmerna. S. har äfven genom öfverflyttande af dylika *Palmellæ* hos friska personer i en frisk trakt framkallat sjukdomen. Liknande observationer, hvilka tala för frossans beroende af alg-intoxication, äro gjorda af Morren, v. d. Corput, Lemaire m. fl. (se Richter & "Schm. Jahrb." ll. cc.). — Härmed sammanhänger nära Binz's⁶⁾ iakttagelse

1) "*Cylindrotracium cholerae asiaticæ*, ein neuer, in den Cholera-ausleerungen gefundener Pilz", i "Virch. Arch." b. XXXVIII.

2) "Path.-Anat. Studien über das Wesen d. Cholera-Processes." Leipz. 1867.

3) "Das Cholera-contagium." Leipz. 1867.

4) Såsom ett olyckligt försök att förklara Cholerans ætiologi må anföras Debey's uppsats i "Deutsche Klin." 1867 *N* 1 & 2, der han såsom den sammans orsak framställer en "ny alg", "*Cholero-phyton*", hvilken uppgift han likväl i *N* 5 nödgas modifiera derhän, att "algen" utgör — ägg af *Ascaris lumbricoides*.

5) Se "Schmidt's Jahrb." b. CXXXI.

6) "Centralbl. f. d. Med. Wissensch." 1867 *N* 20.

om chininets exquisita, ej ens af sublimat öfverträffade, parasitdödade verkan.

En stor mängd fall af en *messlingliknande utslagssjukdom*, framkallad af en på ruttande halm växande svamp, uppgifver sig samme Salisbury ¹⁾ iakttagit och t. o. m. genom direct inoculation lyckats frambringa.

Vid *Mjeltbrand* är af flera forskare (Davaine, Brauell m. fl.) konstateradt, att blodet innehåller "bacterier", hvilka äfven temligen allmänt antagas vara sjukdomens rätta orsak (jfr Klob l. c.).

Den i foglarnes respirationsorganer ofta observerade mögelbildning förekommer äfven någon gång hos människan. Medan Virchow ²⁾ i sina såsom "Broncho"- och "Pneumono-mycosis" anförda fall anser mögelsvampen (*Aspergilli* sp.) uppträda secundärt vid lobulära pneumonier o. s. v., hafva Leyden & Jaffé, Rosenstein m. fl. meddelat fall, der svampbildning skulle vara orsak till putrid *Bronchitis* och *Lungbrand* (Richter). Äfven vid *laryngeal-* och *bronchial-catarrher* äro svampar observerade i sputa.

Af Hallier ³⁾ har i *diphtheritiska* och "*pseudo-diphtheritiska*" membraner mögelbildning blifvit iakttagen.

I *blåsan* uppväcka jästsvampar, genom orena cathetrar eller på annat sätt införda, envisa *catarrher* (Schönbein, Traube m. fl. ⁴⁾).

Samtliga ofvan anförda iakttagelser lida emellertid mer eller mindre af det betänkliga felet att ej vara tillräckligt pröfvade för att såsom erkända sanningar kunna ingå i vetenskapen. Annorlunda är förhållandet med den väl begränsade sjukdomsgrupp, med hvilken vi i denna afhandling närmare skola sysselsätta oss —, om hvars vegetabiliska ætiologi knappt någon meningsskiljaktighet vidare kan sägas råda. Härifrån bildar ännu den föröfrigt utmärkte dermatologen och anatomen Wilson ett enstaka stående, märkvärdigt undantag. Han ser hos "*Nosophyterodermata*" i hela sjukdomsprocessen endast en "*aberration of cellformation*" ⁵⁾, i det de primitiva anlagen till hår- och epidermis-celler kvarstå på sin embryonala utvecklingsgrad, till följe hvaraf de, ehuru varande af animalisk natur, få ett vegetabiliskt ut-

¹⁾ Enl. "Schm. Jahrb." b. CXXI.

²⁾ "Virch. Arch." b. IX.

³⁾ "Die pflanzl. Parasiten des menschl. Körpers." Leipz. 1866 och "Virch. Arch." b. XXXVI.

⁴⁾ Jfr Bardeleben, "Lehrb. d. Chir." b. IV. p. 229 och "Ups. Läkaref. Förh." b. II. p. 260.

⁵⁾ "Brit. and For. Med. Chir. Rev." b. XXXIII. p. 200.

seende (!). En annan mening hyllas af Devergie¹⁾, Cazenave²⁾ och några andra franska författare, att nämligen svampbildningen endast är ett tillfälligt secundärt fenomen vid de sjelfständigt uppträdande, oftast från dyscrasier härledda exanthemerna. — Af samtliga nutidens dermatologer föröfrigt erkännes dessa sjukdomars parasitiska natur, hvilken blifvit satt utom allt tvifvel genom talrika iakttagelser och försök, till hvilka vi i det följande flerstädes återkomma.

Äfven här hafva den nyare Mycologiens läror blifvit tillämpade — och äfven om forskningen med afseende på parasitsvamparnes rätta ursprung m. m. ännu ej kommit till några vissa resultater och stundom t. o. m. synes råkat på något äfventyrliga afvägar, torde man med skäl kunna hoppas, att inom kort vetenskapen i detta ämne skall lyckas sprida full klarhet.

För någon tid tillbaka ifrigt sysselsatt med Botanikens studium omfattade jag med förkärlek den del deraf, som behandlar växtriketets lägsta former, Svamparne. Såsom f. d. dilettant inom Mycologien var det naturligt, att under mina medicinska studier de på svampparasiter beroende sjukdomarne skulle för mig erbjuda ett egendomligt intresse — ty här borde väl Celsi ord: "*Ipsa quoque naturæ rerum contemplatio quamvis non faciat medicum, aptiorem tamen medicinæ reddit*" egentligast ega sin tillämpning. — Efterföljande uppsats grundar sig visserligen till någon del på eget iakttagande af de der skildrade svamp- och sjukdomsformer, men är dock egentligen att betrakta såsom ett försök att med tillgodogörande af den rikhaltiga litteraturen i ämnet lemna en kortfattad och åskådlig framställning af deras vigtigaste förhållanden. Om derföre i densamma intet nytt anträffas, hoppas jag, att den åtminstone skall vara en temligen trogen skildring af åsigternas successiva utveckling och frågans närvarande ställning.

¹⁾ "*Traité pratique des maladies de la peau.*" Paris 1857.

²⁾ "*Traité des Maladies du Cuir chevelu.*" Paris 1850.

I.

De på människans hud parasitiskt förekommande
svampbildningar.1. *Achorion Schoenleinii* Remak ¹⁾.

Synon. Porrigophyton, Mycodermis, Mycoderma Favi Gruby; Oidium Schoenleinii Lebert.

Historik. Remak skall varit den förste, som (1837) i Favus iakttog mögeltrådar, men utan att tilldela dem någon betydelse. Schönlein, genom Bassi's upptäckt af muscardinet föranledd att noggrannare undersöka Favus, visade först dess beroende af svampbildning ²⁾. Gruby ³⁾ lemnade den första riktiga och fullständiga beskrifning af möglet och dess "capsel", framställde dess förhållande till epidermis och säger sig lyckats genom ympning öfverflytta den på trä. Inoculationer på människor och djur äro med framgång gjorda af Remak, Bennett, Köbner m. fl. — Om dess närmare kännedom hafva i synnerhet Remak, Bennett ⁴⁾, Lebert ⁵⁾, Wedl ⁶⁾, som först anträffade densamma äfven *in* håret, Robin ⁷⁾ samt sednast Hallier gjort sig förtjenta.

Beskrifning. Mycelium består af sammanflätade, förgrenade, n. jemnbreda, böjda, färglösa trådar, 0,002—0,004 m.m. breda, i början långsträckta, utan synbart innehåll, småningom tilltagande i bredd, articularade, med allt kortare leder, som i början innehålla en enkel

¹⁾ Vi bibehålla här de gamla, allmänt kända benämningarne — churude, såsom nedan visas, utan tvifvel endast beteckna lägre utvecklingsstadier af andra mögelsvampar —, då undersökningarne ännu ej kunna anses kommit till *bestämd* visshet, till hvilka arter de rättast böra hänföras.

²⁾ I bref till J. Müller, tryckt i dennes Archiv 1839, jemte en ofullkomlig teckning af densamma — endast mycelium.

³⁾ I "Compt. Rendus des Séances de l'Academie d. Sciences de Paris." b. XIII. 1841. m. fl. st.

⁴⁾ "On the vegetable nature of Tinea favosa" (i "Transact. of the Roy. Soc. of Edinburgh", b. XV. 1842).

⁵⁾ "Memoire sur la Teigne" (i "Physiologie pathol." II. Paris 1845).

⁶⁾ "Grundzüge d. pathol. Histologie." Wien 1854.

⁷⁾ "Histoire naturelle des Végétaux parasites." Paris 1853.

rad små korn, hvilka ej vidröra hvarandra men mot grenarnes ändar ("receptacula") småningom tillväxa i storlek, närma sig och slutligen nå hvarandra samt ändtligen synas sammansmälta med den omgifvande väggens membran och afsnöras — så att öfvergången till conidier ("sporer") är omärklig; conidierna aflånga (vanl.), runda l. mera sällan oregelbundna, 0,003—0,010 m. m., perlbandslikt förenade (moniliformia) l. oregelbundet hopade; de äro starkt ljusbrytande, endast de större innehållande en liten kärna; deras antal betydligt öfvervägande myceliitrådarnes.

Ann. Beskrifningarne äro hos författarne vanligen orediga och differera ej obetydligt, hvilket hufvudsakligen beror på bemödandet att med stränga gränser skilja mycelium och "receptacula", mellan hvilka öfvergången är successiv.

Svampen börjar såsom en fin punktering på ett antal epidermisceller, n. alltid, såsom det synes, i nedre delen af hårets epidermiscanal; de härifrån utväxande mycelii-trådarne gifva upphof till de characteriska conglomerater, hvilkas form och utseende vi nedan (under Favus) skola beskrifva. Här må blott anföras, att i det inre af dessa "favi" finnas conidier i otalig mängd, mot periferien blandade med myceliet, och ytterst ett tunnt, amorph lager eller en capsel ("stroma", "gangue amorphe"), som (enl. de flesta författare) bestämdt tillhör svampen och ej den omgifvande epidermis eller deraf producerats. — För öfrigt intränger äfven svampen i sjelfva hårfollikeln samt mellan hårets längsfibriller (t. o. m. ända ner i bulben: Hoffmann) — eller mellan nagellamellerna.

Naturalhistoria ("moderväxt" etc.). Ehuru redan förut af Hebra och andra uttalats den åsigten, att den endast borde anses för en outvecklad form af "vanligt mögel", gissningsvis *Penicillium crustaceum* Fr., har först på sista tiden frågan om dess egentliga ursprung och förhållande till öfriga hudparasiter börjat vetenskapligt behandlas, så att den för närvarande utgör, så att säga, en af Mycologiens brännande frågor. — Redan 1845 skall Remak på ett äpple ur Favus-svampen uppdragit *Penicill. crust.* (under hvilka försigtighetsmått känna vi ej). Lowe¹⁾ och Köbner²⁾ gjorde nya, men ofullkomliga odlingsförsök, på grund af hvilka den förre identifierade *Achorion* med *Aspergillus glaucus* Lk., den sednare hänförde den till släktet *Penicillium*. — På klinisk väg

¹⁾ "Ann. & Mag. of Nat. Hist." b. XX.

²⁾ "Klin. und exper. Mittheilungen aus der Dermatologie und Syphil," Erl. 1864.

(genom inympning af *Penic.* på huden ¹⁾) har Pick ²⁾) ensam lyckats komma till liknande resultat: efter ympningen uppkom en sjukdomsform, identisk med "das herpetische Vorstadium des Favus" (se längre fram!), en gång t. o. m. med svampbildning, liknande den hos Favus. Äfven iakttog han, att vid yppig vegetation hos Favus-svampen någon gång utvecklas fructifications-organer, tillhörande *Penicillium* och (tillfälligtvis?) en *Aspergillus*-art.

Sednast har af Hallier ³⁾) frågan blifvit med stor flit och grundlighet behandlad. Genom talrika, under behöriga försigtighetsmått utförda odlingsförsök och i en mängd uppsatser har han sökt — och såsom oss synes lyckats — ådagalägga, att *Ach. Schoenl.* verkligen är en på lägre utvecklingsgrad stående form ("Oidium-bildning") af *Penicillium crustaceum* ⁴⁾), hvilken på människokroppen, i det kolsyrerika mediet, ej kan normalt utveckla sig. Hans bevis härför äro: 1) *Achorion*-conidier, öfverförda på citronskifvor l. dyl., producera inom några dagar *Penicillium* — ja, han säger sig under microscopet timma för timma följt dess vegetation från det groende conidiet ända till börjande penselbildning; 2) *Penicillii*-conidier, odlade på blod, ägghvita m. m., frambringa liknande *Achorion*-bildningar; 3) i sällsynta fall finnes hos *Ach. Schoenl.* spår till penselbildning . . .

Härmed kunde något hvar tycka saken böra vara afgjord, då helt nyligen Herm. Hoffmann ⁵⁾), en af nutidens mest framstående mycologer, stödd på egna försök, med bestämdhet förklarar den

¹⁾ Sättet, hvarpå denna, såsom Köbner benämner den, "epidermoidala inympning" tillgår, är följande: svamperustorna macereras några timmar i vatten; derpå befrias huden på ett eller annat sätt från det yttre epidermis-lagret och fuktas med några droppar af svampvattnet, som sedan får intorka, hvarefter anlägges ett fast sittande förband med små fuktiga compressor.

²⁾ "Untersuchungen über die pflanzl. Parasiten" (i "Verhandl. d. Zool. Bot. Gesellschaft in Wien" b. XV).

³⁾ "Die pflanzl. Parasiten", "Gährungserscheinungen." Leipz. 1867 m. m. Af det förra arbetet är meddeladt ett referat i "Ups. Läkare-Fören. Förh." b. I. p. 419.

⁴⁾ Jfr dock de Bary (i "Handb. d. physiol. Botanik v. Hofmeister" b. II. p. 244), som erinrar om *Pen. crust.* "allestädesnärvarelse" och möjligheten att förväxla under odlingen uppträdande jästceller och myceliitrådar af *Penicillium* — hvars fullständiga utestängande han anser för "ein Ding der Unmöglichkeit" — med parasitsvampen samt hänföra de från de förra utgående utvecklingsformerna till den sednare o. s. v.

⁵⁾ "Ueber den Favuspitz" ("Botan. Zeitung" ²/₈ 1867).

sannskyldige stamfadren vara *Mucor racemosus* Bull., hvilken han — enligt vår öfvertygelse med fullt skäl — anser omöjligens kunna betraktas såsom en vegetationsform af *Penicillium*.

På grund af denna skiljaktighet i resultater kunde möjligen någon falla på den tanken, att den leder sitt ursprung från flera olika slag af mögel. Häremot talar dock dess constanta, egendommiga utvecklingssätt, dess aldrig varierande microscopiska characterer o. s. v.

För att skaffa oss en egen öfvertygelse i saken, hafva vi äfven företagit några odlingsförsök. Då dessa emellertid ej under tillräckliga cauteler blifvit utförda — hvilket hardt när gränsar till det omöjliga — och till följe deraf i sjelfva verket ingenting bevisa, vilja vi ej dervid uppehålla oss, utan anföra endast vår derifrån hemtade subjectiva åsigt, enligt hvilken vi obetingadt ansluta oss till Hallier's mening. Denna är äfven så mycket sannolikare, som dels *Achorion* onekligen i sin byggnad företer större likhet med *Penicillium*, än med *Mucor* (& *Aspergillus*), dels *Pen. crusti* växesätt, med sina grönaktiga crustor, såsom oss synes, rätt nära erinrar om "favi."

Svampen ger upphof till den under namn af *Favus* kända sjukdom (hvilken, då den angriper naglarne, blifvit kallad *Onychomycosis*).

2. *Trichophyton tonsurans* Malmsten.

Synon. *Trichomyces tons.* Malmst.; *Achorion Leberti* Robin; *Mentagrophyta* Gruby, *Microsporon mentagrophytes* Robin; *Trichomaphyton*, *Mycoderma plicæ* Günsburg (enl. Robin).

Historik. Malmsten, som iakttagit svampbildning såsom orsak till Herp. tonsdens¹⁾, meddelade sig härom med Gruby, hvil-

¹⁾ Redan förut (1843) hade Günsburg (Compt. R. b. XVII) beskrifvit en i *Plica polonica* tillfälligtvis förekommande svamp, som sannolikt hit bör hänföras — och ännu året förut (C. R. b. XV) Gruby i *Mentagra* funnit den af Robin *Microsporon mentagrophytes* benämnda mögelbildning. Denna förklarades visserligen sedan af Robin ej vara annat än den genom epidermis-cellernas upprullning kring håret förorsakade nätlika bildning (se nedan, anm. under beskr.), hvaremot Bazin, Köbner m. fl. identifierade densamma med *Trichophyton*, en åsigt till hvilken Robin sjelf omsider öfvergått. — Såsom bevis på den här rådande villervalla må anföras, att Küchenmeister ("Die in und an dem Körper des leb. Menschen vork. Parasiten." Leipz. 1855), som ännu anser sig böra bibehålla *Microsp. Men-*

ken i sitt svar säger sig äfven observerat densamma. Den sednares iakttagelser blefvo först publicerade¹⁾; kort derefter lemnade den förre en noggrann beskrifning jemte afbildning af svampen och gaf den dess allmänt antagna namn²⁾. Begge ansågo den likväl uteslutande tillhöra håren; dess allmänna förekomst bland epidermis har först af Hebra, Bazin och Bäremsprung (se under Herp. tonsens!) framhållits. Slutligen har dess kännedom gjort stora framsteg genom Köbner, som företrädesvis behandlat dess förhållande till Syccosis, och Borch³⁾. Inoculations-försök hafva talrika och med framgång verkställt; bland andra hafva Köbner, Borch och Ziemssen på sig sjelfva öfverflyttat svampen med thy åtföljande sjukdom.

Beskrifning. Svampbildningens hufvudmassa utgöres af ett i en mängd perlbandslikt förenade, klotrunda, ovala eller om-sesidigt tryck kantiga och vanl. lika stora (omkr. 0,004 m. m.) leder [eller, om man så vill, ännu ej afskilda conidier] deladt, färglöst, sällan grenadt (schizo-)mycelium. Spridda deremellan förekomma isolerade conidier af samma form och storlek, de större med stor kärna. Derjemte finnas, ehuru sällsynt, något smalare, oledade trådar, med l. utan små kärnor. Dylika trådar utgöra (enl. Borch), då möglet förekommer på ytan af epidermis, större delen deraf och bilda ett tätt nät af sammanflätade, enkla l. grenade mycelitrådar, hvilka äfven sprida sig ner i hårfolliklarne; mellan epidermis-cellerna utväxa de, blifva ledade och afsnöra conidier. Enl. Köbner äro äfven i naglarne mycelitrådarne talrika.

Mögelbildningen har sitt säte såväl bland epidermis-cellerna, som mellan hårets fibrer, hvilka ofta ända från roten äro h. o. h. undanträngda af de i hårets längdriktning radbandslikt utväxande conidii-kedjorna, tills den några m. m. ofvan hufvudsvålen orsakar hårens afbrytande. Äfven bildar den omkring håret, både inom follikeln och straxt ofvan utträdet ur svålen, tjocka slidor af tätt lagrade conidier och mycelitrådar, och härifrån intränger den i håret, vanl. i roten men äfven från sidorna. (Jfr Borch's utförliga och sakrika framställning l. c.).

tagr. som egen art, sedan han ord för ord efter Gruby beskrifvit denna, såsom afbildning bifogar en af Gudden på pubishåren funnen *Aspergillus*, som med den förra ej har aflägsnaste likhet (!). Jfr Köbner ("Vchw Arch." b. XXII).

¹⁾ "Compt. Rend." b. XVIII (1 Apr. 1844).

²⁾ Föredrag vid Skand. naturforsk.mötet i Christiania 1844 — tryckt, jemte tillägg, i Hygiea 1845 *N* 6 & 8.

³⁾ "Bidr. til Læren om de planteparasitiske Hudsygdomme." Kjöbenh. 1865.

Anm. Den stundom såsom svampen tillhörig beskrifna "nätbildning" beror på partiellt lossnande och upprullning af hårets epidermis-celler, hvilka dervid synas bilda ett mer l. mindre reguliärt nät.

Naturalhistoria. Med afseende på dess ursprung återgifva vi här i största korthet Halliers åsigt, hvilken, så vidt vi känna, är den ende, som gjort några egentliga försök. Efter att förut¹⁾, förledd af den yttre likheten, ansett den såsom en vegetations-form ("Acrosporen-Hefe") af *Penicillium crustaceum*, redogör han i sitt nyaste arbete häröfver²⁾ för de försök, som föranledt honom att ändra åsigt. Vid dessa uppkom under odling af *Trichophyton* först en *Oidium*-form ("*Oidium albi-ans*" Auctt.), som i myceliitrådarnes ändar afsnörade bruna, runda sporer, identiska med [?] sporerne af *Ustilago Carbo* Tul. (sv. "Kolbrand"); dessa grodde sedan och alstrade *Aspergillus glaucus*, äfvensom ett *Eurotium*. Af *Aspergillus* säger han sig omvänt uppdragit *Ustilago* — och anser han på grund häraf *Trichophyton* vara en "*Oidium*-bildning i omogen form (*Torula*)" af serien *Aspergillus-Eurotium-Ustilago*³⁾.

Frågan om svampens identitet med Achorion Schoenleinii eller icke är ännu oafgjord. Hebra⁴⁾, som tror sig sett *Favus* uppkomma ur en efter långvarigt bruk af compressor spontant utvecklad *Herpes tonsdens*, W. Boeck m. fl. hafva uttalat sig för densamma, under det att isynnerhet franska dermatologer fasthållit skilnaden eller t. o. m. sökt urskilja flera arter. På sednare tider hafva Köbner, Strube⁵⁾ och Stark⁶⁾ sökt vederlägga Hebra's åsigt, stödjande sig på inoculationer, dervid de aldrig ur *Favus* lyckats frambringa *Herpes tonsdens*, men väl "ein herpetisches Vorstadium des Favus", som lätt förvexlas med *Herpes*; ej heller ur *Herpes tonsdens* någon *Favus*. — Å andra sidan anser sig Pick genom experimenter bevisat motsatsen (identiteten). Enligt denne

1) "Bot. Zeit." 1865 **Nr** 49 & "Pflanzl. Paras."

2) "Gährungserscheinungen etc."

3) Vi erinra härvid, att dessa undersökningar förefalla oss vara den moderna Mycologien — på hvars "utvecklingsserier" föröfrigt ännu framstående mycologer tvifla — drifven till sin spets, men måste föröfrigt, då vi ej haft tillfälle att anställa egna försök, afhålla oss från något bestämt omdöme.

4) "Hebra's klin. Foredrag over Hudsygdommene ved Marcus." Kjöbenh. 1862.

5) "Exanthemata phytoparasitica eodemne fungo efficiantur, quæritur." Diss. Berl. 1863.

6) "Jenaische Zeitschr. f. Med. & Naturw." b. II.

— som sätter enda skilnaden emellan de begge sjukdomarne i den större eller mindre massan af svampelementer och deraf beroende förstöring af väfnaderna — uppkommer efter ympling af *Favus* först en Herpes-eruption (Köbner's "herp. Vorstadium"); denna utvecklar sig allt efter mer eller mindre gynnande yttre omständigheter till *Favus* eller Herpes tondens. Efter inoculation af *Trichophyton* uppkommer vanligen endast Herp. tondens, men stundom utvecklar sig en sjukdomsform, identisk med Favi "herp. Vorstadium." Denna dess obenägenhet att bilda *Favus* skulle vara att söka antingen i hårets olika beskaffenhet hos olika individer ¹⁾ eller deri, att svampen, såsom stående på ett lägre stadium i generations-vexlingen, i början endast förmår frambringa likartade bildningar och dem motsvarande reactionsfenomenen på huden (hvilket dock synes stå i strid med hvad nyss är anfördt om uppkomsten af Favi "herp. Vorstadium"). — Deri öfverensstämma emellertid alla de nämnda försöken, att hittills aldrig ur *Trichophyton* någon verklig *Achorion-crusta* ("favus", "scutulum") uppkommit. Äfven då begge arterna samtidigt inympas på samma person, utveckla sig, enl. Köbner, de resp. sjukdomarne i sina karakteristiska former, så att man ej gerna kan tala om beroende af "disposition", "yttre omständigheter" o. s. v.

De sjukdomar, som af svampen förorsakas, kunna under det gemensamma namnet *Herpes tondens* sammanfattas; derunder höra *H. tondens sens. propr.*, *H. circinatus* (der den beror på svampbildning) jemte "*Eczema marginatum*" (enl. Köbner), den smittosamma formen af *Sycosis* (*S. parasitica*) samt *Onychomycosis* (delvis) — sannol. ock den af Ellinger observerade, på parasitbildning beroende *Blepharitis ciliaris* samt Bäremsprung's "*Erythrasma*."

3. (?) *Microsporon Audouini* Gruby.

Ann. Vi upptaga här för fullständighetens och bättre redas skull denna "art", hvars tillvaro från flera håll, åtminstone såsom orsakande en

¹⁾ Med anledning häraf yttrar Klob, sedan han erinrat om, huru behöfligt det varit att här få veta, hvori det herpet. Vorstadium hos *Favus* skiljer sig från det hos Herp. tondens: "Zieht man in Schlussfolgerungen so problematische Dinge wie das herpetische Vorstadium des Favus oder Dispositionen einzelner Individuen u. s. f. als Prämissen herein, dann darf auch der Schluss selbst nur als hypothetischer und nicht mit so apodiktischer Schärfe hingestellt werden, soll anders die Wissenschaft aus solchen Arbeiten Nutzen ziehen" (l. c. p. 35). (Hallier, hvilken polemiken äfven gällar, har, såsom vi redan sett, samtidigt öfvergifvit sin förra åsigt).

särskild sjukdomsform, drages i tvifvelsmål. Beskrifningarne äro ock hos olika författare mycket afvikande och häntyda delvis uppenbart på föregående, med hvilken den numera ofta — och troligen rättast — förenas.

Synon. Trichophyton & Trichomyces decalvans Malmst.; Microsporon decalvans Bazin.

Historik. *Microsporon Audouini* synes blifvit upptäckt af Gruby redan 1841¹⁾, men beskrefs först 1843²⁾; den har vidare studerats af Bazin³⁾, som skilde den i tvenne arter, *M. Audouini* och *M. decalvans*, hvilka han dock numera åter förenar, samt af Robin, hvilken likväl i sina sednare arbeten förenar den med *Trichophyton*.

Beskrifning. Mycelium består af talrika, korta, vågformigt krökta, 0,002—0,003 m. m. breda trådar, som sins emellan anastomosera till ett tätt nät, hvilket omgifver håret; utanpå detta finnas lagrade talrika små runda (0,002—0,005 m. m.) eller ovala "sporer", genomskinliga, utan synbart innehåll. Svampbildningen börjar på hårets yta, 1—2 m. m. från hufvudsvålen, omgifver detta såsom en (0,015 m. m. tjock) slida, hvilken så intimt är fästad vid hårets epidermisbeklädnad, att detta förr brister, än slidan aflossas, och så att säga stryper eller afbryter håret. Den sprider sig vidare till närstående hår, hvarvid äfven hufvudets epidermisceller blifva säte för sjukdomen...

Sådant är hufvudinnehållet af Gruby's framställning, om hvilken dock Bazin, som för närvarande nästan ensam fasthåller *Microsp. Aud.* såsom egen art, säger, att den "thyvär till större delen är en roman." — Enligt denne skiljer sig *Microsp. Aud.* från *Trichophyton* hufvudsakligen genom sina smärre och mindre talrika sporer samt rikligare mycelium — äfvensom genom sättet för dess utbredning på håret. "Sporerna" bilda näml. på dess yta små isolerade grupper och anordna sig gerna nätformigt. Sjelfva håren visa *sphäriska eller ovala uppdrifningar*, inom hvilka massor af sporer träffas. Den finnes äfven bland epidermis (se vidare under Porriigo decalvans!).

Om dess **Naturalhistoria** är intet vidare känt.

Den sjukdomsform, till hvilken svampen skall gifva upphof, kallas af Bazin *Tinea pelada* (Porriigo decalvans pr. p.?). Äfven skall (enl. Kleinhans) Bazin hafva observerat ett af densamma förorsakadt fall af *Onychomycosis*.

¹⁾ "Compt. Rend." b. XIII. p. 388.

²⁾ "Compt. R." b. XVII.

³⁾ "Die parasitäre Hautaffectionen nach Bazin's Leçons theor. et cliniques etc. von Kleinhans." Erlangen 1864.

4. *Microsporon Furfur* Robin.

Synon. Mycoderma Eichstädtii Richter; Epidermophyton Bazin.

Historik. Upptäcktes 1846 af Eichstädt, som jemte Sluyter och Richter m. fl. vidare studerat densamma. Sitt namn har den erhållit af Robin, som likväl erkänner sig aldrig hafva sett den (!). — Äfven med denna hafva inoculationer företagits och lyckats, t. o. m. på djur, der den eljest ej uppgifves förekomma (Köbner l. c.).

Beskrifning. Mycelium utgöres af en mängd i alla riktningar vridna och sins emellan oregelbundet sammanflätade, långa, h. o. d. förgrenade, 0,002—0,003 m. m. breda trådar, än tomma, än försedda med kärnor, och på sidorna l. ändarne afsnörande conidier; dessa äro klotrunda, 0,004—0,006 m. m. (uppgifvas ofta mycket mindre), starkt ljusbrytande, med *en* eller *två* kärnor, samlade i större, drufklaslika hopar, så att de stundom genom ömsesidig tryckning blifva polygonala och förefalla såsom en sammanhängande väfnad.

Den har sitt säte i och mellan de ytligare lagren af epidermisfjäll, hvilka den i alla riktningar genomdrager, men intränger ej i hårfolliklarna; undre lagret utgöres af myceliet, det öfre af conidier.

Naturalhistoria. Sådan den på människokroppen uppträder, borde den otvifvelaktigt hänföras till släktet *Collarium* Lk., hvars character: "Sporidia globosa, simplicia, in acervos conglutinata, floccis mucedineis (mycelio) cincta vel intertexta"¹⁾ tyckes vara just efter densamma uppsatt.

Äfven här är Hallier den ende, som genom odling sökt utforska den egentliga moderväxten — och återgifva vi här nedan de resultater, till hvilka han kommit, med samma reservation som under *Trichophyton*.

Efter att genom odling af *Aspergillus*-sporer på gummilösning erhållit bildningar, fullkomligt liknande *Microsporon*, och omvänt ur den sednare (ehuru endast med svårighet, då dess conidier gro ytterst långsamt) uppdragit *Aspergillus*, har han nyligen genom nya försök bekräftat och utvecklat sin lära. Vid odling af *Microsporon* i stärkelseklistor uppkom ett *Stemphylium* [?], hvilken bildning i samma medium uppdrogs äfven af *Aspergillus* och *Ustilago Carbo*. Mest bevisande för sambandet mellan *Asperg.* och *Microsp.* är dock ett odlingsförsök i vatten, der af den sednare uppkom *Aspergillus*—

¹⁾ Fries "Syst. Mycol." b. III. p. 440.

och *Eurotium*-knölar, fullkomligt öfverensstämmande med de på möglande växter förekommande (äfvensom med de af *Trichophyton* frambragta). På grund häraf räknar han *Microsporon Furfur* såsom äfven tillhörande serien *Aspergillus-Eurotium-Ustilago* och anför den såsom "Hefe der *Oidium*-sporen" ¹⁾. Sannolikt anser han vara, att Pityriasis versicolor uppkommer ur *Ustilago*-sporer, och finner högst troligt, att dessa öfverföras genom att vid stärkande af linne använda stärkelse, bemängd med sporer af "Kolbranden."

Af hvad ofvan blifvit anfördt synes, att Hallier anser *Trichophyton* och *Microsporon* för olika vegetationsformer af samma serie. (Han säger sig äfven af *Trichophyton* hafva uppdragit *Microsporon Furfur*). Härvid möter dock den frågan, hvarför ej alltid Pityriasis versic. ledsagar Herpes tondens och tvärtom. Förklaring häraf tror han kunna sökas i den långa tid (flera veckor eller månader), som åtgår för att öfverföra formerna i hvarandra, inom hvilken tid epidermis redan hunnit affjälla.

För fullständighetens skuld må nämnas, att Hebra åtminstone förr trott äfven denna art vara identisk med *Achorion* och *Trichophyton* och endast afvika genom olika local, hvilken åsigt Pick ännu delar. Af samma tanke är äfven Hutchinson, som bland annat stödjer sig på fall, der han sett Pityriasis versicolor uppkomma hos kvinnor, som gäfvat di åt barn med Favus! (Köbner).

Svampen är orsak till *Pityriasis versicolor*.

Tillfälligtvis funna svampar.

"*Puccinia Favi*", af Arndtsen ²⁾ funnen vid Favus, skall enligt W. Boeck ³⁾ sedermera upprepade gånger i Norge anträffats beledsagande densamma. v. Düben ⁴⁾ har vid Pityriasis versicolor funnit ett exemplar. Utomlands synes den knappt blifvit observerad. — Också kan den svårligen anses för annat än en tillfällig inblandning — liksom stundom svampsporer på näsans slemhinna —, ej ens växande bland "favi", såsom Robin antager.

¹⁾ Med "Hefe" förstår han enkla, rundade celler med en eller få kärnor, som ej fortplanta sig genom utsändande af trådar, utan genom afsnörning — utan att — såsom vanligt — dervid fästa begreppet af gasutveckling ("Botan. Zeit." 1867 *M* 10).

²⁾ "Om en ny Planteform i Favus" ("Norsk Mag. for Lægevidensk." ser. 2, b. V p. 316.

³⁾ "Norsk Mag. for Lægev." ser. 2. b. VII. p. 221.

⁴⁾ "Sv. Läk. Sällsk. nya Handl." b. IX. p. 14.

Aspergilli spec., iakttagna i yttre hörselgången af Mayer och Pacini, i naglarne af Meissner (?) o. s. v.

Penicillii sp. ("*Sterigmatocystis antacustica*"), äfven anträffad i yttre örat, af Cramer.

"*Trichophyton sporuloides*" Robin, af Walther funnen i *Plica polonica*, växande bland den håren sammanklibbade vätskan — och således ej hithörande, likasom

"*Trichophyton ulcerum*" Rob. et

Mucor Mucedo L. (?), den förra funnen bland crustorna på ett gammalt bensår, den sednare vid gangräna senilis.

"*Oidium albicans*" Robin — egentligen en epitheliophyt — förekommer någon gång omkring mun, anus, på mammæ och genitalia.

["*Microsporon Hardii*" ¹⁾, uppgifven såsom förekommande i *Acne punctata* et "*varioliiformis*", är enligt "Schm. Jahrb." b. CXXIV. p. 387 endast en "*Parasiten-schwindel*", härledande sig från bomulls- och linnetrådar, kring hvilka moleculer ur varet grupperat sig.]

Hvart den af Fuchs ²⁾ i "*Alphus*" iakttagna svampbildning rättast bör hänföras, måste lemnas of afgjort.

(Om "*Microsporon minutissimum*" Bäreuspr. se under *Herpes tonsdens*!)

II.

De af svampbildning på människans hud förorsakade sjukdomar.

1. *Favus* ³⁾.

Synon. *Tinea vera*, *favosa*, *ficosa*, *lupina*, *scutulata*; *Porri*go, *P. favosa*; *Mycosis achorina*. [*Fr.* *Teigne vraie*, *faveuse*; *Engl.* *Scall-Head*; *Tysk.* *Aechter Grind*, *Erbgrind*; *Sv.* *Ondskorf*].

Historik. Det kan synas underligt, säger Cazenave ⁴⁾, för den, som noggrannt studerat *Favus* och blifvit frapperad af den bestämda, egendomliga sjukdomsbild, som dess symptom framställa, att en sjukdom — utan tvifvel bland de på hufvudets hårbevuxna del

¹⁾ Jfr Chausit i "l'Union médicale" b. XIX. **M** 101.

²⁾ "Die krankhaften Veränd. d. Haut." b. II. p. 538.

³⁾ Namnet *Favus* är betecknande för rufvornas utseende, *Tinea* för sjukdomens härjande inverkan (eller, enligt andra, för dess envishet); benämningen *Porri*go (motsvarande grek. *πυργιασος*) är mindre exact, såsom egentligen afseende hudsjukdomar med torr kliartad affjällning.

⁴⁾ "Malad. du cuir chevelu", p. 210.

förekommande den viktigaste — så tydligt begränsad till sin form och utveckling, ej i alla tider tillräckligt fästad läkarnes uppmärksamhet för att beskrivas under eget namn och densamma uteslutande tillhörig synptomatologi. — Så är emellertid förhållandet. Hos de äldsta författarne finnes intet, som *med säkerhet* kan tolkas såsom hithörande.

Celsus begagnar visserligen uttrycket "favi", men endast för att liknelsevis characterisera en utslagsform, benämnd "Meliceria", hvilken med sin "hvitaktiga, honungstjocka, klibbiga vätska" ej kan vara Favus, utan sannolikt en Impetigo. Galenus talar om skallighet, orsakad af "vätskornas förskämning" och föregången af exsudation och ulceration. Archigenes och Asclepiades beskrifva ulcererande utslag, hvilka medföra skallighet och kräva energisk behandling. Apollonius omnämner elakartade "achorer." Här af vill synas, att de gamle likväl kände Favus och åtminstone antydde dess viktigaste symptom — bland hvilka för dem hårets förlust var det förnämsta — ehuru de förbisett contagiositeten (Cazenave).

Föga mera bestämda äro Arabernas beskrifningar. Avicenna — i hvars "sahafati" man oriktigt velat spåra sjukdomen — anser skallighet i vissa fall uppkomma af frätande sår, beroende på ett *egendomligt ämne*, som finnes bland håret. Hos följande författare beskrivas vanligen under namnet Tinea hudsjukdomar bland hufvudets hårbetäckning, tillhörande barnaåldern, och upptagas allt efter olika åsikter under vexlande namn ett olika antal arter. Dessa inskränka sig ofta till 2, en botlig och en obotlig, hvilken sednare företrädesvis torde betecknat Favus. Ambr. Paré skilde mellan 3 slags Tineæ, bland hvilka "T. fcosa" skall motsvara Favus; hos Plenck finnes den under "Scabies capitis lupina" och "Tinea vera" antydningsvis skildrad; Lorry antager blott *en* Tinea, "T. vera", och afskiljer de öfriga såsom "Pseudo-tineæ."

Först med Alibert bragtes större reda i ämnet. Han lemnade en naturtrogen skildring af sjukdomens väsendtliga characterer (bland hvilka han likväl endast med yttersta inskränkning erkände contagiositeten, hvilken först framhölls af Mahon) och uppförde den (i sina sednare arbeten) bland classen Tineæ såsom eget slagte, Favus, med arterna vulgaris och scutiformis. — Hittills hade såsom characteristiskt för Tineæ ansetts den gemensamma platsen samt bildningen af crustor. Men med Willan, hvilken till grund för sitt system lade affectionens primitiva form, förlorade crustorna, såsom secundära producter, sin betydelse. Han ansåg Porrigo (hvilket namn han i st. f. Tinea upptog) för en pustulös sjukdom och ställde den närmast Impetigo, från hvilken den endast genom contagiositeten skulle afvika. Ba-

teman's arbete utgör deremot ett steg tillbaka, såsom sammanförande — och delvis sammanblandande — alla hudsjukdomar bland håret under Porrigio. I Frankrike upptogs Willan's åsichter och utvecklades vidare af Bielt m. fl.¹⁾

Genom Schönlein's upptäckt och Gruby's noggranna undersökningar spriddes ljus öfver sjukdomens rätta natur — och derefter kan man säga, att parasitsvampens historia är sjukdomens. Efter att förut varit under afdelningen Pustulæ eller Squamæ uppförd i de dermatologiska systemerna, upptages den nu jemte de följande allmänt i en särskild grupp; (Dermatophytæ²⁾).

Beskrifning. ("Symptomer och Förlopp").

Favus är en, vanligen till hufvudets hårbeklädda del inskränkt, contagiös hudsjukdom, beroende på uppträdande inom epidermislagret af en mögelart, som bildar egendomliga, halngula conglomerater ("favi"), hvilka medföra irritation och — vid längre duration — atrofi af huden och obliteration af hårfollikeln eller degeneration af papillen, med åtföljande (beständig) förlust af håret.

Sjukdomen börjar med en mer eller mindre häftig klåda, åtföljd af en lindrig, circumscrip³⁾ eller (vanl.) diffus rodnad samt ökad affjällning af epidermis. Stundom uppkommer en pustulös eruption, som föregår uppträdandet af "favi" men någon gång äfven kvarstår, då dessa framträda, hvilket gifvit anledning till förvexling dem emellan, som dock ej bör vara svår att undvika⁴⁾. Svampbildningen visar sig

¹⁾ Jfr angående Favi historia Cazenave och Bazin ll. cc. samt Hebra, "Atlas der Hautkrankheiten." Wien 1856—66.

²⁾ Härifrån gör likväl fortfarande en eller annan författare undantag; bland andra Cazenave, som oaktadt sina vidlyftiga forskningar ännu alltjemnt — såsom förr Sauvages, Mahon m. fl. — anser Favus bero på en af hypersecretion åtföljd inflammation af glandulæ sebaceæ i hårfollikelns mynning.

³⁾ Egentligen då sjukdomen tar plats på bålen, är rodnaden circumscrip^t och antager gerna en reguliärt afrundad form. Dessa erythematösa ringar kunna lätt förvexlas med de af *Trichophyton* förorsakade Herpesringarne, men skilja sig constant genom mindre diameter och fullständig likformighet (Bazin). Jfr om detta "herpet. Vorstadium" ofvan under *Trichoph.* samt Köbner: "Klin. und exp. Mittheil."

⁴⁾ Favus-crustan kan lossas från sin plats utan læsion af underliggande väfnad, hvilket naturligtvis ej eger rum med pusteln. Emellertid uppstår stundom vid Favus en liflig inflammation, då svampbildningen kan maskeras af det intorkade varet ur pustlerna. Man har därför uppgifvit en stor mängd sätt att med säkerhet skilja "favi" från intorkadt var, epidermisconglomerater o. s. v. (Jfr Küchenmeister "Paras."). Detta låter sig bland annat lätt göra på kemisk väg, genom användande af medel (t. ex.

nu, i början som en punktformig upphöjning under epidermis. Aflossas epidermis, framträda små, i midten af ett (någon gång 2—3) hår genomborrade och nedtryckta crustor¹⁾. Dessa tillväxa temligen hastigt, upplyfta och afspränga delvis den betäckande epidermis-lamellen och visa sig nu under form af runda, torra, sköra, skålförmiga, "kräftstenslika" bildningar, som med den convexa, undre sidan — ofta försedd med en mucro — ligga nedtryckta i huden. På insidan äro de vanligen ojemna och visa antydning till concentriskt ordnade lager, af hvilka det yttersta upplyfter den omgifvande epidermis, hvari skålen är liksom infattad, och ofta höjer sig 1—2 m.m. öfver den angränsande hudytan. Färgen är smutsigt hvitgul, mot periferien något mörkare; stundom, i synnerhet då, såsom ofta händer, patienten sjelf afkliat epidermislaget, är den genom extravaseradt, intorkadt blod förändrad. Crustornas bredd är vanligen 2—6 m.m., men kan någon gång uppgå ända till 1—2 c.m. Under deras vidare tillväxt försvinner den ursprungliga formen: de blifva oregelbundna, sprickiga, här och der genomdragna af hår och sammanflyta stundom med närgränsande, då det, för att igenkänna sjukdomens ursprungliga form, är nödvändigt att genom cataplasmer aflägsna de gamla rufvorna, hvar efter nya framträda, företeende den egendomliga formen. Efter någon tids bestånd (ett par veckor — månader) bli de mörkare, med upphöjda, liksom inrullade kanter, lossna allt mer och affalla för att ersättas af nya; stundom upplösa de sig delvis i ett smutsgult pulver (conidier²⁾).

Aflägsnas en Favus-rufva försigtigt, visar sig i den underliggande huden en jemn och glatt, rodnad depression, som dock — der sjuk-

alkalier) som lösa de sednare, men ej inverka på de förra — eller genom andra reactioner. Vi förbigå likväl dessa, då hela utvecklingen — och i nödfall microscopet — ej gerna kan lemna något tvifvel öfrigt om deras rätta natur.

¹⁾ Svampen synes börja sin utveckling rundt omkring håret i nedre ändan af hårets epidermis-canal och derifrån utsända mycelii-trådar företrädesvis periferiskt mellan epidermis-lagren. Då epidermiscanalen fast sammanhänger med håret, hindras möglet att i denna riktning fritt utveckla sig — och detta är, enligt Bazin, orsaken till den nävelförmiga fördjupningen. (Se Küchenmeister l. c.)

²⁾ Såsom egendomligt för svampbildningen har man äfven anført dess vidriga lukt och genom allehanda liknelser, såsom katturin, härskt mjöl, träskluft o. s. v. sökt förtydliga densamma. — Då den åtföljes af pustulösa eruptioner, sprider dessas i förruttnelse stadda innehåll en ytterst vämjelig stank; eljest synes oss lukten ej afvika från den, vanligt mögel, samladt i massor, sprider.

domen ej varat för länge — inom kort (stundom inom en timma) höjer sig i niveau med den öfriga huden. Voro "favi" sammanflytande, synas en mängd linsstora fördjupningar, mellan hvilka skilnaden antydes genom smala upphöjda linier.

Vi öfvergå nu till de sjukliga förändringar, håren på ett af sjukdomen angripet ställe undergå. Dessa blifva i början torra och glanslösa och contrastera derigenom mot de omgifvande friska. Snart antaga de en grå (stundom rödlett) färg, blifva tunnare, ojemna och sköra, så att de stundom brista i jernhöjd med crustorna. Vid lindrig dragning lossna de utan smärta, eller ock affalla de sjelfmant (primitiv alopeci). Under microscopet visa de sig mörka, opaka, ej transparenta; vanligen medföljer vid utdragningen den sjukligt förändrade rotslidan såsom ett färglöst gelatinöst öfverdrag, omgifvande hårroten. På sjelfva hårets yta finnas stundom sporer samlade i större eller mindre grupper ända ned till hårroten; ofta anträffas äfven svampen mellan de sins emellan splittrade längsfibrerna¹⁾. — Slutligen förstöras hårpapillerna, folliklarne oblitereras delvis eller helt och hållet, hvaraf (definitiv) skallighet blir följden²⁾. Nu försvinna äfven svampcrustorna, troligen af brist på näring — kvarlemnande kala, bleka, nedtryckta ärr.

Sådant är sjukdomens vanliga förlopp, der ej verksam hjälp melankommer. I svårare fall beledsagas den af envisa pustulösa eruptioner och ulcerationer, från hvilka inflammationen någon gång sprider sig till underliggande väfnader, så att verkliga abscesser uppstå; ja, man skall t. o. m. sett cranii ben blottas, carieras och hjernans hinner angripas. — Oftare, i synnerhet när den har sitt säte i temporal-regionen eller bakhufvudet, medför den angioleucit med inflammation af hals- och nackkörtlarne. (Någon gång uppgifves den äfven orsaka otitis, ophthalmi o. s. v., äfvensom den ansetts inverka hämmande på kroppens och själens utveckling).

Vanligen förlöper den likväl h. o. h. localt och utan att medföra några allmänna symptom. Dock torde väl den svåra klåda, som medföljer sjukdomen, underhållen och förökad af de skaror af löss,

¹⁾ Om de microscopiska förändringarne handla utförligt Robin l. c. samt Hoffmann i "Bot. Zeit." 1867 №. 31.

²⁾ Hårens förändring beror i början af talgkörtlarnes upphörda secretion, sedan hufvudsakligen af förändringar i papillen, hvilka åter hafva sin grund, förutom i den compression, som "favi" utöfva, i de mera djupliggande mögelbildingar, som kring densamma utvecklas. Dessas närvaro förklarar äfven den envishet, med hvilken svampen recidiverar, äfven sedan alla crustor blifvit aflägsnade.

som ofta frodas bland crustorna, i synnerhet på förut nedsatta subjecter, kunna genom förhindrande af sömn och hvila menligt inverka¹⁾, likasom å andra sidan exempel anföras, att dels till följe häraf, dels genom det begrepp af skam och vanära, som medföljer denna sjukdom, sinnesbeskaffenheten och t. o. m. förnuftet taga skada.

Sedan äldre tider har man plägat skilja mellan tvenne, till utseendet skilda *former*, *Tinea favosa* och *scutulata*. Den förra, för hvilken vi helst skulle vilja antaga benämningen *Favus urceolaris* (= *disseminatus*, *sparsus*, *isolatus*, *alveolaris*, *vulgaris* etc.) är hufvudformen, och hafva vi i föregående beskrifning företrädesvis på denna fäst afseende. Den intager ofta större delen af hufvudets hårbevuxna del, hvilken den betäcker med en mängd skilda, skålformigt urholkade crustor, erinrande om cellerna i en honungskaka. Då crustorna utväxt, sammanflyta de visserligen stundom (= "*Tin. favosa cohærens*" Baz.), men i omgifningen finnas dock alltid characteristiskt *concava* crustor — och *dylika uppkomna äfven, om de gamla* genom emollierande omslag *aflägsnas*. — Denna form är den enda, hvarunder sjukdomen visar sig utom hufvudets hårbeklädnad.

Den andra formen, *Favus scutulatus* (= *nummularis*, *figuratus*, *annularis* etc.) är mera sällsynt; den intar vanligen mindre yta och är inskränkt till hufvudets hårbetäckning, så att den, om den derifrån sprider sig till ansigtet, alltid öfvergår till *F. urceolaris*. Det i början uppträdande erythemet bildar regelbundet rundade, känsliga fläckar och efterföljes af en stark och långvarig epidermis-affjällning. Efter någon tid iakttages bland fjällen en mängd små, gulaktiga punkter, hvilka hastigt sammansmälta och bilda en sammanhängande, afrundad, tunn, ojemnt skroflig, af hår genomdragen massa, *men utan likformiga fördjupningar*²⁾; storleken kan uppnå 1—4 c.m. Aflägsnas de gamla crustorna, *bilda sig nya, fullkomligt liknande*. Någon gång sammanflyta flera sålunda uppkomna "plaques" och kunna då

¹⁾ En målande, ehuru visserligen starkt färglagd teckning häraf lemnar Sauvages: "Propter sorditiem & purulentiam putredinem acerrime foetet caput tinea affectum & ob pectinationis defectum pediculi generantur, inde pruritus et scalptus dirissimus, inde pervigilium, macies, quandoque cum hectica, faciei pallore etc." ("*Nosologia method.*" II. p. 580).

²⁾ Lebert (l. c.) ansåg på grund af denna olikhet och det i öfrigt skiljaktiga förloppet denna form bero på en egen mögelart. Enligt Bazin skall dess uppkomst betingas deraf, att möglet utvecklar sig omkring tätt stående, t. ex. ur samma follikel uppväxande hår, så att "favi" genom det ömsesidiga trycket förlora sin ursprungliga form och sammanflyta till en *crusta*, *redan innan de trädt i dagern*. I sjelfva verket skall den äfven efter epilation försvinna och lemna rum för *F. urceolaris*.

betäcka hela hjessan med en slags calott, i hvars periferi man dock nästan alltid återfinner eruptioner i form af större eller mindre cirkel-segmenter (Bazin). — Sjukdomsförloppet är här långsammare, möglet synes ej tränga så djupt, hvarför äfven hårens sjuklighet är mindre markerad; utbredningen derjemte vanligen mindre. På grund häraf anses den i allmänhet såsom mera godartad och lättare curabel än hufvudformen.

Föröfrigt urskiljas af författarne mer eller mindre oväsentliga former, t. ex. *F. suberinus*, *achatinus*, *murinus* o. s. v. Bland dessa omnämna vi här endast Bazin's "*Tinea squarrosa*", egentligen en form af *F. scutulatus*, som företrädesvis utvecklar sig uppåt omkring håren; de härigenom kring dessa bildade slidor sammansmälta och bilda irreguliära upphöjningar, liknande bergen på en reliefkarta.

Vi hafva hittills egentligen talat om sjukdomens förlopp, då den har sin vanliga plats — bland hufvudhåren. Understundom — och företrädesvis i invetererade fall — sprider den sig dock till den öfriga kroppen, ansigte, bål och extremiteter¹⁾, ja, den har af flera observatörer ertappats på sjelfva glans! Den väljer dock med förkärlek extremiteternas sträcksidor, såsom mera hårbevuxna (Bazin). Dess förlopp är här mera godartadt: föregångna af en circumscrip rodad uppträda de här alltid skällika favi, sammanflyta stundom, men affalla vanligen inom en månad, hvarmed sjukdomen — på detta ställe — plägar upphöra (Hebra).

Blott på *en* local, hvilken för möglets fortkomst erbjuder liknande vilkor som håren, näml. *naglarne*, är den af mera ingripande natur — och äfven oftare anträffad. Vi skola här i största korthet framställa dessa på sednare tider af flera läkare studerade sjukliga förändringar, som förorsakas både af *Achorion* och *Trichophyton* (enligt Bazin äfven af *Microsp. Audouini* och enl. Virchow möjligen af *Aspergillus* eller andra). Att vi här sammanföra dem på ett ställe, har sin förklaring deri, att i de — för öfrigt föga viktiga — sjukdomarne, så vidt hittills känt är, de med microscopet diagnosticerade olika mögelarterna ej medföra några skiljaktiga symptom.

Redan länge hade man vid *Favus* iakttagit, att naglarne ofta företedde ett sjukligt förändradt utseende. Meissner var den förste, som (1853) påvisade mögelbildning såsom orsak till degeneration af

¹⁾ Prof. Malmsten har iakttagit fall, der hos barn sjukdomen haft sitt säte på kroppen, utan att hufvudet varit angripet ("Hygiea" b. XXV. *N* 10).

naglarne, hvarefter flera, särdeles Virchow¹⁾ och Kleinhans (l. c.) deråt riktat sin uppmärksamhet²⁾.

Naglarne förlora sin glans och antaga en smutsgrå färg, blifva skrofliga och spröda, stundom splittriga, samt tilltaga betydligt i tjocklek, i synnerhet de yttersta $\frac{2}{3}$. De fortfara likväl att tillväxa; lunula är vanligen oförändrad. I allmänhet inställa sig lindriga inflammations-symptomer i omgifningen. På undre sidan af nageln och ofta mellan lamellerna träffas en halfmjuk, smutsgul massa, som microscopiskt undersökt visar sig innehålla mögel-elementer. — Huru dessa, åtminstone på händerna, blifvit öfverförda, är ej svårt att inse.

Ætiologi och Förekomst. Sjukdomens sjelfva väsende beror just på närvaron af en svampbildning. Häraf följer med nödvändighet — derest man ej vill hylla läran om "generatio æquivoca" — att den aldrig spontant uppstår, utan måste utifrån tillföras. Också erkännes numera af alla författare dess smittosamhet, hvilken äfven genom direct inoculation blifvit ådagalagd. Deremot är möjligheten af dess uppkomst ur några af de öfverallt i luften sväfvande svampsporer ("Panspermismus"), som ej tillhört Favus — ehuru i hög grad sannolik — ännu ej med säkerhet bevisad. Fall af sjukdomens öfverförande från djur till människor äro observerade.

Den företrädesvis — och måhända enda — prædisponerande orsaken till dess uppkomst är osnygghet och vanskötsel af håret, under det att antagandet af någon särskild constitution eller diathes (t. ex., såsom ännu af Bazin m. fl. fasthållas, den scrophulösa), såsom därför mera gynnsam, synes oberättigadt. Att dess ärftlighet måste förkastas, faller af sig sjelf.

Sjukdomen, som i synnerhet bland de lägre klasserna ej sällan anträffas, förekommer visserligen i alla åldrar, men sällsyntare under de första lefnadsåren och efter puberteten; allmännast är den mellan 6—9:de året (Barthez & Rilliet). Någon väsentlig olikhet mellan de olika könens mottaglighet förefinnes ej — dock säga sig flera författare funnit den frequentare hos gossar, hvilka ock möjligen mera utsätta sig för smittan.

Hvad särskildt dess spridning inom Sverige angår, är den här sällsyntare än på continenten. Enligt de officiella läkare-rapporterna förekommer den i de nordliga provinserna och på Gotland mera sällan

¹⁾ "Arch." b. IX.

²⁾ Af Virchow benämndes sjukdomen Onychomycosis, af Köbner ("Virch. Arch." b. XXII) Herpes unguium och af Kleinhans — Phytoparasitosis tonsurans unguicularis (!).

— inom Jemtlands län är den ännu ej observerad. I Dalarne deremot, men synnerligast i Bohuslän och Norra Halland är den relativt allmän ¹⁾).

Prognos. Favus är en med rätta för sin långvarighet och en-
vishet fruktad sjukdom. Lemnad åt sig sjelf torde den ganska sällan
upphöra, förrän håret till större delen är förstördt. (Har den sitt säte
på kroppen, plägar den deremot, såsom vi ofvan sett, inom kort sjelf-
mant försvinna ²⁾). Någon fara för lifvet torde den likväl högst säl-
lan — och då genom sina complicationer — medföra. Prognosen be-
ror på dess säte och utsträckning, tiden för dess duration — samt
på möjligheten att underkasta den rationell behandling, hvarunder ut-
sigten för dess botande är säker, ehuru lång tid dertill erfordras ³⁾).

Behandlingen har till uppgift att aflägsna eller döda mögel-
svampen. Det härvid numera allmänt brukliga förfarandet — hvilket
redan i den uråldriga beklufvan har sin förebild — är epilationen.
Denna tillgår i korthet på följande sätt. Sedan håret blifvit kort-
klippt och rufvorna genom repeterad ingnidning af olja vederbörligen
uppmjukats, aflägsnas de fullständigt, och hufvudet tvättas med såp-
vatten. Härpå utdragas med pincett, strå efter strå, de på det sjuka
stället och i dess närmaste omgifning befintliga hår, hvarje dag på en
eller annan kvadrattums yta. Så snart på detta sätt hela den an-
gripna fläcken — under förnyade såptvättningar — blifvit renplockad,
börjar man ånyo med den plats, hvarifrån man utgick — o. s. v. ända
till dess all fjällbildning upphört, då möglet är att anse såsom utro-
tadt. Detta förfaringssätt är säkerligen det enklaste och användes i
allmänhet på de svenska sjukhusen. Utomlands förenas dermed van-
ligen application af "parasitdödande medel", såsom sublimat, oleum
cadinum, arsenik, veratrin, creosot, terpentin o. s. v. Bland dessa
ingår sannolikt sublimatet till större delen kemisk förening med de
ytligare hudlagren och tränger svårigen på djupet, likasom ej heller
oleum cadinum. Deremot torde de lättflytande terpentinoljan och cre-
osoten utan svårighet nedtränga i djupet af hårfollikeln och — då de

¹⁾ Jfr Troilius: "Om Favus, Herpes tonsens och Alopecia areata samt deras företeelser inom Sverige" ("Hygiea" b. XXV. *M* 10).

²⁾ I "Berl. Klin. Wochenschr." 1866 *M* 42 meddelas likväl ett fall, der sjukdomen, som inalles varat 44 år (från första barndomen), efter att i 26 år endast hållit sig till hufvudet, spridde sig till bålen och nedre extremiteterna, der den slutligen antog enorma dimensioner.

³⁾ För de under åren 1853—62 på Serafimer-Lazarettet i Stockholm vårdade Favus-sjuka uppgick medelunderhålls-tiden till 233 dagar, hvilket dock till någon del berott af brist på personer för att verkställa epilationen (Troilius).

vid direct beröring dödande inverka på svampelementerna — framför andra medel förtjena att användas. (Jfr Hoffmann l. c.)

Vid "Onychomycosis" har behandlingen, der en sådan vidtagits, vanligen bestått i afskrapande af nagelns yta samt indränkning med veratrin- eller sublimat-lösning — stundom med godt resultat ¹⁾.

2. *Herpes tonsdens.* (sens. lat.)

Synon. [vid förekomst bland hufvudhåren] Squarus tonsdens; Herpes, Porrigo, Mycosis tonsurans; Porrigo circinata, tonsoria; Tinea, Trichonosis furfuracea; Rhizo-phyto-alopecia; Trichosis, Trichophytia [Fr. Teigne tondante; Engl. Ringworm p. p.; Sv. Hårskärande mögel (Malmsten)].

Historik. Hvilkendera kommit sanningen närmare, Cazenave, hvilken i Celsi "Area" och Galeni *ὀφίασις* trott sig igenkänna Herpes tonsdens, eller Bateman (med hvilken Bazin förenat sig), som hänför dessa namn till Porrigo decalvans, kan på grund af beskrifningarnes ottydlighet ej utrönas. Lika litet låter sig hos författarne ända till nyare tider afgöras, om eller under hvilket slag af Alopecia, Tinea, Pityriasis o. s. v. den innefattats. Man kan således betrakta den såsom en ny sjukdom, och synes Plumbe varit den förste, som deraf haft en riktig uppfattning. Han beskriver hårens egendomliga afbrytning, omtalar dess smittosamhet samt uppträdande allestädes på kroppen och skiljer derifrån en icke smittosam Herp. circinatus. (Förut hade i England t. ex. af Willan och Bateman under benämningen "Ringworm" varit sammanförda denna sjukdom och Favus scutulatus, troligen äfven delvis Porr. decalvans). Emellertid tillskrifves äran af dess upptäckt vanligen Mahon, hvilken på ett sätt, som vunnit följande observatörens enstämmiga bifall, tecknade dess utseende och förlopp, påpekade dess contagiositet och svårbotlighet, men ansåg den för en constitutionell, ärftlig sjukdom. Han benämnde densamma "Teigne tondante." Cazenave, som iakttog, att den ofta förekom "i förbindelse med" Herp. circinatus, gaf den namnet H. tonsurans ²⁾.

Ett jättesteg framåt gjorde sjukdomens kännedom genom upptäckten af dess rätta orsak, på grund hvaraf Gruby benämnde den "Rhizo-phyto-alopecia", hvilket namn äfven Malmsten samtidigt föreslog i sin på upptäckten af parasitsvampens närvaro grundade, noggranna utredning af dess egendomliga natur ³⁾. Den har allt sedan, af Bazin, Bäremsprung m. fl. med förkärlek studerats.

Redan före denna upptäckt hade Gruby iakttagit sin "Mentagrophyta" (se under *Trichophyton*!) och på grund häraf uppställt en

¹⁾ Ripping i "Deutsche Klinik" 1865 *N* 38.

²⁾ Jfr ang. den äldre historiken Malmsten, Cazenave, Hebra, Borch II. cc.

³⁾ "Trichophyton tonsurans (hårskärande mögel)" i "Hygiea" 1845 *N* 6 & 8.

contagiös sjukdom, "Phytomentagra" ¹⁾). Emellertid gick han för långt, då han härmed förenade *all* Sycosis, hvori han efterföljdes af Küchenmeister, Neligan m. fl. och i början af Robin och Bazin. Den sistnämnde — hvars begär att framhålla sina egna förtjenster och fördölja sina misstag här på ett eclatant sätt framlyser — har ändtligen efter flera mer eller mindre sväfvande och orediga framställningar jemkat sin åsigt derhän, att Sycosis, der den orsakas af mögelbildning, är identisk med sista stadiet af Herpes tonsdens. Tillvaron af en Sycosis med parasitiskt ursprung antages föröfrigt af Anderson, Hardy, Koch, Kleinhans m. fl., hvaremot Hebra, Bäremsprung och många andra förneka den. Såsom afgörande frågan torde kunna anses Köbner's tvenne sakrika uppsatser ²⁾, der han — som spårar den parasitiska Mentagrans historia ända till Plinius — stödd på 230 observerade fall, lemnar en klar framställning af sjukdomen. Han skiljer mellan 2 slags Sycosis: en "Folliculitis barbæ" och en "Trichomycosis nodosa", beträffande hvilken sednare vi längre fram skola återgifva hans åsikter.

Prioritetsfrågan med afseende på upptäckten af identiteten mellan Herpes tonsdens och *H. circinatus* (ehuru den visserligen redan af Plümbe och Cazenave var genomskådad) har varit tvist underkastad. Utslaget har fallit till förmån för Bazin, som 1854 påpekade närvaro af *Trichophyton* i vissa fall af Herp. circ. ³⁾), hvilket han följande år utsträckte till alla, samtidigt med Bäremsprung's publication om förekomsten af en svampbildning i densamma. Sedermera har dock Hebra ⁴⁾ tillägnat sig äran, såsom varande den förste, som (1854) visade tillvaron af en maculös form af Herp. tonsdens och dess identitet med *H. circinatus*. Medan Bazin och Bäremsprung härleda *all* *H. circinatus* från *Trichophyton*, har Hebra iakttagit en egen, af densamma oberoende *H. circin.* — Bäremsprung — hvars arbete öfver Herp. tonsdens allmänt erkännes såsom det grundligaste — uppställde jemte flera andra former äfven en "*H. inguinum*" hvilken han likväl sedermera uppförde såsom en särskild sjukdom, förorsakad

¹⁾ Synonymer äro: *Mentagra contagiosum*, *Tinea Mentagra*, *Sycosis parasitica*, *Trichomycosis nodosa*. — Att den af G. ursprungligen beskrifna sjukdom var en verklig Sycosis, synes mindre troligt; sannolikare en ytlig Herpes tonsdens.

²⁾ "Ueber Sycosis und ihre Beziehungen zur Mycosis tonsurans" ("Virch. Arch." XXII) och "Klin. und exper. Mittheil. aus d. Dermatologie und Syphil." Erl. 1864.

³⁾ Här af uppkomna synonymer äro: *Dermatomyces tonsurans* och *Trichophytia circinata*.

⁴⁾ "Hautkrankheiten" p. 523.

af en egen svamp, förslagsvis kallad *Microsporon minutissimum*. Köbner¹⁾ är dock böjd att förena den med Herp. circ., likasom han äfven genom intressanta observationer och directa försök synes ådagalagt, att Hebra's "Eczema marginatum" bör föras hit. Hebra förnekar emellertid närvaron af någon svampbildning i sin sjukdom²⁾.

Beskrifning. Såsom vi ofvan sett, sammanfattas numera under benämningen Herpes tondens flera sjukdomsformer, hvilka förr under särskilda namn beskrifvits. För att vinna mera reda, skola vi därför, efter att lemnat en öfversigtlig bestämning af sjukdomen, särskildt beskrifva densamma, allt efter dess förekomst bland hufvudets hårbeklädnad, i skäggets hårfolliklar eller på den öfriga kroppen. (De sjukliga förändringar, den medför af naglarne — hvilka redan af Mahon observerades — hafva vi under Favus berört).

Herpes tondens är en på alla kroppens epidermis-bildningar förekommande, i hög grad smittosam sjukdom, orsakad af en (n. uteslutande af conidier bestående) mögelsvamp, hvilken har sitt säte i och omkring hårrötterna eller bland epidermis och bildar ett characteristiskt, hvitt, flockigt beslag; den medför hårets afbrytande (och någon gång dess totala undergång) samt åtföljes — allt efter säte och utvecklingsstadium — af symptomatiska, vanl. circulärt ordnade, erythematösa, vesiculösa, pustulösa eller tuberculösa eruptioner.

a) *Herpes tondens* (sens. strict.)

kallas sjukdomen företrädesvis, när den angriper hufvudets hårbeklädda del. Dess förlopp är här följande. På en liten, rodnad fläck uppkommer, under stundom ganska liffig klåda, en mängd särdeles små vesikler, hvilka hastigt resorberas (hvarföre de ofta förbises) och efterträdas af nya, periferiskt sig spridande dylika, hvarjemte en förökad epidermis-afstötning eger rum. Håren blifva glanslösa, missfärgade, torra och bräckliga samt afbrytas slutligen splittrigt, kvarlemnande endast en ojemn stubb af 1—3 m.m. höjd, hvarigenom de sjukliga ställena antaga det egendomliga utseende, som af Mahon träffande liknats vid skälhud. Vid microscopisk undersökning af de bräckliga hårresterna visar sig en djupgripande förstöring af både cortical- och medullar-substansen, hvilka ofta ej vidare kunna urskiljas;

¹⁾ "Klin. und Exp. Mittheil." — Öfver samma ämne har K. i "Berl. Med. Centralzeit." ²⁸/₃ 1867 skrifvit en "vorläufige Mittheilung", hvilken vi dock ej varit i tillfälle att jemföra.

²⁾ På Hebra's plancher synas äfven, såsom Borch påpekat, håren o-skadda kvarstå — åtminstone till större delen.

mellan fibrerna, af hvilka stundom endast rester kvarstå, finnas hopade otaliga conidier; bulben — der den vid hårets utryckande medföljer — är vanligen böjd, skrumpen och liksom förvissnad. — Mellan hårstumparne är huden torr, sträf (liknande "cutis anserina"), till färgen vanligen blå- eller gråhvit, mellan de afstöta epidermisresterna betäckt med ett egendomligt, fint flockigt eller lamellöst, hvitt, "sammetslikt" stoft (mögelsvampen). Detta förenar sig omkring de framträngande håren och omgifver dem med fina, hvita, mattglänsande ("asbestlika") sidor.

Fläckarne vidga sig småningom och nå storleken af en 5-öresslant och deröfver. Stundom börjar sjukdomen samtidigt på flera ställen, och de uppkomna fläckarne sammanflyta, bildande en någon gång öfver hela hufvudet utbredd "tonsur." När den ett ställe, som saknar hårbeklädnad, såsom pannan eller nacken, bildar den der röda cirkelsegmenter och öfvergår i den form, till hvilken vi under c) skola återkomma.

Stundom försvinner sjukdomen, om endast tillbörlig renlighet iakttages, själfmant, ehuru visserligen alltid först efter en längre tids förlopp, och de angripna ställena betäckas ånyo af fullkomligt normalt hår; stundom — der den ej underkastas behandling och i synnerhet osnygghet gynnar dess utveckling — tränger den djupare och medför inflammation af hårfolliklarne. Härefter uppkomma confluerande pustulösa eruptioner, efter hvilkas intorkande de sjuka fläckarne betäckas med mörkgula, glänsande crustor¹⁾. Då dessa omsider försvinna, kan till följe af hårfollikelnas destruction permanent skallighet inträda eller (oftare) endast ofullkomligt utvecklade hår alstras.

b) "*Sycosis parasitica*"²⁾.

Icke hvarje fall af Herpes tondens barbæ är tillika en Sycosis parasitica — lika litet som hvarje Sycosis bör räknas hit — utan derunder innefattas endast de fall, der parasitsvampen (som i de första stadierna äfven här medför förändringar, identiska med dem på öfriga hårbeklädda delar) tränger djupare ned och på grund af localens

¹⁾ Jfr. Bazin l. c. — I detta stadium, der man på grund af crustornas utseende lätt kan taga sjukdomen för en Impetigo (l. Favus), är diagnosen ej lätt. Utom den hjälp, anamnesen och microscopet lemna, bör med synnerlig uppmärksamhet närvaron af Herp. circinatus på öfriga kroppsdelar efterspanas.

²⁾ Beskrifningen är hufvudsakligen i korthet sammanfattad efter Köbner, Bazin (ll. cc.) samt Ziemssen: "Sycosis oder Mentagra" (i "Greifsw. med. Beiträge" b. II. Danzig 1864.)

egendomliga anatomiska förhållanden förorsakar mera ingripande förändringar, hvilka — äfven dessa — hittills blifvit under collectivbenämningen *Mentagra* eller *Sycosis* inbegripna. I denna period utvecklar sig parasiten hufvudsakligen mellan håret och inre rotslidan, afpräparerar denna småningom ända ner till bulben, som genom dess tryck atrofieras, åstadkommer inflammation och varbildning i follikeln och dess omgifning samt förorsakar slutligen uppkomsten af furunklar och "fungösa plaques" eller tuberkler, beroende af hypertrophi af papillarkroppen.

Sjukdomen plägar uppträda på haka, kinder, läppar och främre delen af halsen, på hvilka ställen den föregås af symptom, liknande dem, vi ofvan beskrifvit såsom vanliga vid utvecklingen af *Herp.* tondens bland hufvudhåren, men äfven närmande sig den character, de förete, då densamma har sitt säte på den öfriga kroppen. Tiden för det ondas öfvergång i detta stadium plägar variera mellan 6:te veckan och 5:te månaden från dess första uppkomst.

Vanligen visa sig redan under de första månaderna omkring några af de sjukliga håren röda, linsstora papler, omkring andra smärre pustler. Begge tilltaga i antal och storlek samt confluera, hvarigenom af de sednare uppstå brungula crustor, här och der genomdragna af degenererade hårstumpar, af de förra bön — hasselnötstora, kopparfärgade eller livida tuberkler, hvilka ofta i sin ordning på ytan ulcerera och i den omkringliggande huden åstadkomma djupa, vidsträckta indurationer. De visa ofta en kretsformig anordning och omgifvas åt alla håll på en vidsträckt yta af *Herpes*-eruptioner i olika stadier. — Genom den i hårfollikeln uppkomna inflammationen lossnar håret och bortfaller stundom; blir det deremot qvarsittande, bidrager det, likt en sequester, att öka inflammationens häftighet. Den ymniga varbildningen medför äfven parasitens död och upplösning, till följe hvaraf det ofta är svårt, ja omöjligt att i de kvarblifna hårresterna med microscopet upptäcka densamma (hvilket deremot alltid lyckas med närstående, mindre angripna hår).

Slutligen — ofta först efter flera års förlopp — upphör sjukdomen, vanligen först sedan den medfört obliteration af hårfolliklarne.

Orsaken till den mera intensiva form, sjukdomen (liksom föröfrigt andra, eljest ytliga eruptioner) på detta ställe antager, beror på hårfolliklarnes storlek och djupa läge, corii tjocklek samt den subcutana, laxa bindväfvens kärl- och nerf-rikedom. (Äfven i axillar- och pubis-hårens folliklar — "pudendagra" — närmar sig stun-

dom symptomernas intensitet denna form¹⁾. — Att dess uppkomst orsakas af svampens närvaro (hvilken ej är en tillfällig complication) visar bland annat Ziemssen's lyckade experiment att på sig sjelf inoculera densamma jemte åtföljande Sycosis; dess identitet med Herpes tondens ådagalägges till öfverflöd af deras samtidiga uppträdande och öfvergång i hvarandra samt inoculationer, der man af den ena frambragt den andra.

Den differentiella diagnosen mellan denna form och den rent inflammatoriska Sycosis bör företrädesvis stödja sig på anamnesen (den föregående och vanligen partiellt qvarstående Herpes-eruptionen), tuberkelnas djupare säte och större spridning, hårets macro- och microscopiska förändringar samt uteblifvandet af recidiver, sedan den en gång blifvit ordentligt botad.

c) "*Herpes (tondens) circinatus.*"

På icke hårbetäckta delar af kroppen, der svampen hufvudsakligen har sitt säte bland epidermis, visar sig det onda under en mindre characteristisk, teml. varierande form²⁾. Åtföljda af en oftast särdeles häftig klåda uppkomma antingen på en liten (ärt — skillings-stor), något upphöjd erythematös fläck en mängd små vesikler, hvilka hastigt resorberas eller brista, qvarlemnande en papperstunn, snart aflossande, gul skorpa, medan i den rodnade periferien nya små vesikler allt vidare uppstå — eller visar sig en liten grupp tätt stående, föga upphöjda, knapphålshufvudstora papler, som betäckas med små hvita fjäll och likaledes sprida sig periferiskt. Under sjukdomens vidare utveckling observeras vanligen endast en röd, mer l. mindre bred, upphöjd, kretsformig, med små vesikler, papler l. crustor besatt strimma (någon gång 2—3 utanför hvarandra), hvilken beständigt vidgar sig, medan huden på det innanför belägna partiet antingen är normal, starkare pigmenterad, rodnad eller betäckt med små epidermisfjäll; — korteligen sjukdomen kan framställa bilden af en Herpes circinatus, H. Iris, H. squamosus, Erythema marginatum, Lichen circumscriptus o. s. v. På detta sätt sammanflyta ofta flera på skilda håll uppkomna eruptioner och gifva upphof till hvarjehanda figurer i form af ett

¹⁾ Hit bör äfven sannolikt — troligare än till Favus, såsom förf. sjelf anser — föras en af Ellinger beskrifven envis Blepharitis ciliaris, förorsakad af en i ciliernas rötter vegeterande svamp ("Virch. Arch." b. XXIII).

²⁾ Jfr Bazin och Borch ll. cc., Hebra "Atlas d. Hautkrankh.", Köbner "Klin. und Exp. Mitth." m. fl. — Bäremsprung's berömda uppsats har ej varit oss tillgänglig.

klöfverblad, en 8 o. s. v. Äfven här angripas håren, men gifva genom sin ringa utveckling parasiten föga näring.

Denna form väljer gerna till sitt säte ansigtet samt händernas och underarmens ryggidor, men förekommer för öfrigt öfver hela kroppen och sprider sig ofta på en vidsträckt yta, öfver större delen af ansigtet, halsen och bröstet eller från lårens insida öfver buk, rygg o. s. v.¹⁾ Till följe af localens beskaffenhet har sjukdomen en ojemförligt mildare character och försvinner vanligen af sig sjelf, åtminstone inom 1—2 månader ("Eczema marginatum" — om det bör föras hit — är dock ganska envist), utan att lemna några spår efter sig och blott undantagsvis recidiverande.

Från den ej på parasitisk grund stående *H. circinatus* är denna på yttre utseendet omöjlig att skilja, äfven om vesiklerna här vanl. äro något smärre. Diagnosen får afgöras af microscopet, hvartill äfven anamnesen (bevislig smitta), längre duration (den ej parasitiska *H. circ.* varar sällan öfver 14 dagar: Hebra) samt större utbredning kan bidraga. Föröfrigt är *H. circinatus* i 9 fall af 10 beroende af parasitsvampen (Hebra).

Ætiologi och Förekomst. Äfven här beror sjukdomens uppkomst på mögelsvampens öfverförande. Hebra, som flera gånger sett densamma uppträda på kroppen efter användande af compressor, som ej varit rentvättade och till följe deraf troligen bemängda med mögelbildningar, drar deraf den sannolika slutsatsen, att de sednare, utvecklade på huden, härtill äro orsaken²⁾. Dess stora smittosamhet är genom talrika iakttagelser ådagalagd; ofta angripas flera eller alla medlemmarne af en familj. Med afseende på förekomsten visa de olika formerna en viss olikhet. Herpes circinatus — den vanligaste — angriper begge könen, likväl oftare mankönet och företrädesvis mellan 30—50:de året (Köbner); H. tondens visar sig nästan uteslutande hos barn under 10 år och oftare hos gossar; Sycosis tillhör mannaåldern. Häraf förklaras, att en med Sycosis behäftad familjfader meddelar sin hustru H.

¹⁾ I "Berl. Klin. Wochenschr." 1865 *N* 14 omtalas ett fall, der sjukdomen spred sig från hufvudet och intog hela kroppen samt medförde hårets nästan totala förlust.

²⁾ Här må ock anföras Lowe's märkliga uppgift, att hos bryggeriarbetare, sysselsatta med *jästberedning*, uppkomma hudutslag ("Psoriasis annulata" och Mentagra), i hvilka anträffas svampmycelium. Han säger visserligen detta likna Favus-svampen, men efter sjukdomens säte (haka, händer och armar) hör den sannolikt hit ("Ann. & Mag. of Nat. Hist." b. XX p. 307).

circinatus, medan barnen få H. tondens. — Fall af sjukdomens complication med Favus hafva förekommit.

Såsom ett särdeles vanligt medel till infection anklagas rakknifvar, efter hvilkas bruk inom 3—5 dagar erythemet och derefter vidare symptomter inställa sig. I de flesta fall öfverföres likväl smittan direct genom intimare beröring på ett eller annat sätt — och den såsom Eczema marginatum hittills upptagna formen skall enligt Köbner's observationer ofta genom coitus fortplantas¹⁾. — Hos djur, i synnerhet hornboskap, förekommer sjukdomen ganska ofta, och hos Gerlach²⁾ och Borch äro samlade en mängd fall af dess öfverförande från dessa på menniskor. Äfven i Sverige är den, i synnerhet under fuktiga år, flerstädes hos hornboskapen iakttagen, enl. meddelande af läraren i Veterinär-medicin vid Ultuna landbruks-institut, Hr C. Lindqvist. Densamme har äfven på egendomen Marielund nära Upsala, der större delen af koladugården var angripen, hos tvenne ladugårdspigor observerat de för sjukdomen egendomliga Herpes-eruptioner, hvilka hos begge hade sin plats på begge underarmarnes insida³⁾.

Någon särskild gynnande orsak till sjukdomens uppkomst känna vi knappt; åtminstone synes ej osnygghet här hafva samma stora inflytande som vid Favus, då den lika ofta, om ej oftare, träffas hos de förmögnares renliga och väl skötta barn, som bland de fattigas. Enligt Bazin skall Syphilis, enligt Hebra fugtig bostad⁴⁾ gynna dess uppkomst.

Med afseende på dess utbredning synes den, i synnerhet i Frankrike, vara mycket spridd, äfven i form af Sycosis, hvaraf i Tyskland endast några fall blifvit iakttagna. I Sverige är den ganska sällsynt⁵⁾. "Endast undantagsvis", säger Troilius, "spri-

¹⁾ Detta kan möjligen förklara, hvarför Hebra ("Hautkrankh." p. 363) länge misstänkte den stå på syphilitisk botten.

²⁾ "Die Flechte des Rindes" (i "Mag. f. die gesammte Thierheilkunde" b. XXIII h. 3).

³⁾ Härstädes tillvaratagna och förf. till närmare undersökning benäget meddelade hår och epidermis-rufvor hade på fuktigt ställe blifvit förvarade. Det papper, hvari de voro inneslutna, befanns i deras närhet uppmjukadt och sönderföll vid beröring i smulor, hvilka vid microscopisk undersökning visade characteristiska conidii-kedjor af *Trichophyton*, ofta förgrenade och sparsamt genomdragna af smalare, längledade trådar.

⁴⁾ Orsaken skulle vara öfverförande genom fuktigt, mögelbeväxt linne. — Om ock hypotesen är obevisad, är Bärensprung's skämt olämpligt, "att lika omöjligt är, att en tryffel skulle uppkomma af en champignon."

⁵⁾ Åren 1853—62 vårdades på Serafimer-Lazarettet endast 3 fall och

der den sig utom hufvudets hårbeklädda del"; något i Sverige diagnosticeradt fall af *Sycosis parasitica* är oss ej bekant.

Prognos. Såsom vi ofvan sett, är sjukdomen på olika localer och i olika stadier af betydligt olika betydenhet. Visserligen kan prognosen i allmänhet sägas vara god — vida bättre än vid *Favus* — der ej osnygghet och vanskötsel gynna det ondas utveckling. Likväl är sjukdomen, med undantag af *Herpes circinatus*, hvilken själfmant plägar försvinna, mycket långvarig: *Herpes tonsdens* räcker ofta ett år och deröfver, *Sycosis parasitica* 10 gånger så länge. Äfven med energisk behandling — som här i hårens bräcklighet möter en särskild svårighet — kräfver dess botande tid. Emellertid medför den sällan (*Sycosis paras.* undantagen) hårets förlust, och något menligt inflytande på det allmänna helsestillståndet torde näppeligen ens vid sjukdomens vidsträcktaste utbredning kunnat förmärkas.

Behandling. Vid denna sjukdom skulle, måhända mera än vid de öfriga här afhandlade, profylactiska försigtighetsmått till förekommande af beröring mellan friska och sjuka individer — kanske t. o. m. mellan friska och sjuka kroppsdelar — kunna ifrågakomma. — För en enkel *Herp. circinatus* äro vanligen några såp-tvättningar tillräckliga; vid intensivare former (t. ex. "*Ec. margin.*") användas kraftigare medel, alkaliska tvättningar i förening med *oleum cadinum*, bensen, sublimat-lösning o. s. v. *Herp. tonsdens* behandlas af Bazin på samma sätt som *Favus* med epilation och parasitocida, medan andra, t. ex. Cazenave, Malmsten m. fl. genom renlighet (flitig tvättning och kamning) samt application af medel, som medföra lifligare regeneration af huden, såsom alkohol, borax, alkalier m. m. säga sig vinna godt resultat. Vid *Sycosis parasitica* är repeterad epilation nödvändig och föregås lämpligen af *oleum cadinum*, som, utom sin parasitdödande, äfven genom hårens lossande får en anæstheserande verkan (Bazin). Efter epilationen appliceras en svag sublimat-lösning eller præcipitat-salva. Ofta kvarstå efter svampens utrotande röda tuberculösa indurationer, hvilka lättast resorberas under bruk af jodsvafvelsalva (Klein-hans), ytliga lapistoucheringar (Köbner) l. dyl.

1853—60 på samtliga rikets öfriga allmänna sjukhus intet (*Troilius*), hvartill dock otvifvelaktigt den ringa vigt, som fästes vid sjukdomen, är anledning. På Läns-Lazarettet i Upsala hafva, enligt meddelande af prof. Mesterton, på sednare åren förekommit 6 fall. (Under tjänstgöring på Seraf.-Laz. samt i Upsala har förf. haft tillfälle att observera 3 fall).

3. (?) *Porrigo decalvans.* (?)

Synon. Trichosis Area; Alopecia areata l. circumscripta; Vitiligo Capitis; — Phyto-alopecia; Tinea pelada, T. achromatosa & decalvans.

Historik. Redan under Herpes tondens nämndes, att man sökt leda äfven denna sjukdoms anor ända från de gamles Area och *ὀφίασις*. Huru härmed än må vara, är Bateman den förste, som tydligt urskilde sjukdomen och placerade den bredvid Favus under släktet Porrigo. Sedan Gruby i densamma upptäckt en mögelbildning, uppstod i frågan om dess rätta natur en villervalla, som ännu herrskar. Under det några fullständigt anslutit sig till Gruby's åsichter och andra, egentl. Bazin, antagit desamma, men med betydande reformation af beskrifningen på svampens utvecklingsätt, hafva andra sökt sjukdomen i former af Herp. tondens, medan slutligen andra — hvilkas antal är i tilltagande — h. o. h. förneka dess parasitiska natur. Bland dessa sednare äro flera, som stödja sig på en mängd fall, der de förgäfves efterspanat parasiten, såsom Bärensprung, Hutchinson, Ziemssen, till hvilka Hebra (ehuru först på sednare tider) och Malmsten slutit sig. — Bazin urskiljer två former, Pelada achromatosa & decalvans, men har återkallat sin förra åsigt om deras beroende på olika arter...

Denna fortfarande meningsskiljaktighet, som visserligen till någon del eger sin grund i sjukdomens sällsynthet, är ej lätt att förklara. Likväl synas oss — då man ej gerna kan antaga, att öfvade observatörer så ofta skulle förbisett parasiten — endast två möjligheter finnas: *antingen* har sjukdomen i vissa fall varit *complicerad* med närvaron af någon parasit (t. ex. *Trichophyton*¹⁾ *eller* äro under namnet Porrigo decalvans sammanblandade *tvänne olika sjukdomar*²⁾. För den sednare händelsen vore måhända lämpligare att upptaga Bazin's namn, Tinea pelada, för den på parasitbildning beroende — såvida den ej sammanfaller med Herp. tondens.

Beskrifning (i sammandrag efter Bazin). Sjukdomen börjar, likasom de föregående, med klåda och en mer eller mindre markerad förändring af håret, hvilket på en inskränkt plats får ett

¹⁾ Tilbury Fox (se "Canst. Jahresbericht 1864" p. 268) säger sig stundom vid P. decalv. tillfälligtvis anträffat "sporer och trådar af *Trichophyton*."

²⁾ Detta synes, med afseende fästadt vid Bazin's afvikande beskrifning, ej vara alldeles osannolikt. — Köbner, som ej vill fälla något bestämdt omdöme om Porrigo decalvans, säger sig på Bazin's egen klinik sett flera fall af Porrigo decalvans, der ingen parasit stod att upptäcka.

glanslöst, smutsigt utseende och synes liksom beströdt med stoft. Derefter visar sig parasiten mellan håren i form af obetydligt, gråhvitt stoft; samtidigt synes hufvudsvålen stundom betydligt för tjockad. Hårens utseende blir alltmera sjukligt, de affalla och regenereras omvexlande — under fortfarande klåda — men bli allt ofullkomligare utvecklade, tills de slutligen fullständigt försvinna och lemna rum för en permanent skallighet, medan huden på det sjuka partiet skrumprar. — Degenerationen af håren visar sig i form af runda eller ovala ansvällningar på vissa afstånd från hvarandra, uppkomna genom utbugtning af längs fibrerna och inuti fyllda af en mängd små "sporer." Håren brista i jemnhöjd med dessa uppdrifningar, mellan hvilka substansen synes frisk. Äfven hårroten är sjukligt förändrad, atrofierad.

De angripna fläckarne tillväxa i antal och utsträckning, förena sig och intaga stundom hela hufvudet eller sprida sig t. o. m. "ej sällan" öfver alla kroppsdelar.

Vid Tinea pelada achromatosa är hårets sjukliga utseende mer i ögonen fallande; der uppstå vanligen ovala fläckar, på hvilka huden är affärgad, mjölkhvit. Den upphör stundom af sig sjelf, utan hårets förlust. — T. pelada decalvans har mycket snabbare förlopp, så att ofta inom en månad huden öfver hela kroppen angripes; håren affalla utan någon tydligt utpreglad förändring på fläckar af regelbunden form, på hvilka huden bibehåller sin färg. Den medför vanligen definitiv skallighet. —

Ofvanstående beskrifning, *vilken vi ansett oss här böra föredragesvis följa, då denna uppsats gäller på svampbildning beroende sjukdomar*, afviker — förutom hvad som angår parasitsvampens närvaro — i åtskilligt från den, andra författare (Bateman, Hebra, Cazenave m. fl.) lemna af Porrigio decalvans, och närmar sig mera Herpes tonsdens. Någon klåda omnämnas ej af dessa — tvärtom skall sensibiliteten på fläcken vara minskad; ej heller omtalas någon successiv regeneration af hår, utan sedan dessa en gång affallit, förblir fläcken en längre tid kal och glatt. Likväl är kalheten i de flesta fall ej permanent, utan efter någon tid (stundom efter ett eller annat års förlopp) börjar håret vanl. med ens att återväxa och blir då vanl. åter normalt (stundom grått), ehuru i början ofta mera tunnt och mjukt. Likaså är dess utbredning oftast inskränkt (vanl. till en mindre fläck på bakhufvudet) — och utom hår och skägg observeras den ytterst sällan.

Ætiologi och Förekomst. Såsom vi sett, är sjukdomens orsak dunkel — dess parasitiska natur tvifvelaktig. Enligt Bären-

sprung¹⁾ skall den bero på stördt trophiskt nervinflytande. De flesta författare neka dess contagiositet, hvilken deremot påyrkas af Bazin och af Ziemssen's²⁾ och Hillier's³⁾ observationer synes vinna stöd. Fuktig bostad och Syphilis skall äfven här gynna dess uppkomst (Bazin). Enligt densamma skall den snarare tillhöra de förmögna än de fattigare klasserna, hvaremot Ziemssen tror otillräcklig och dålig föda vara ett viktigt ætiologiskt moment. Den förekommer i alla åldrar, oftast dock hos barn; hos äldre skall den oftare angripa skägget.

Sjukdomen hör, med undantag, som det synes, af Frankrike, till de sällsynta. I Sverige är den ytterst rar⁴⁾ — och någon parasitbildning har der ej blifvit iakttagen.

Prognos och Behandling. Äfven häruti skilja sig Bazin's åsikter betydligt från öfriga författares. Enligt honom är prognosen här sämre än vid de öfriga "Tineæ", då obotlig skallighet är den vanliga följden och curen fordrar lång tid och ändock är oviss. Han använder äfven här epilation och "antiparasitica."

Öfriga auctorer ställa prognosen god och öfverlemna vanligen åt naturen och tiden att bota sjukdomen. Då den vanligen curerar sig sjelf och den sjuke lätteligen råkar ut för någon charlatan, om han ej får någon behandling af läkaren, råder Hebra att ordinera något "oskyldigt medel", såsom Oleum Macidis l. dyl. Cazenave recommenderar tvättningar med spiritiosa, quininlösning o. s. v., Bärensprung föreslår electricitet.

4. *Pityriasis versicolor.*

Synon. Dermatomycosis microsporida.

Historik. Efter att förut jemte åtskilliga andra anomalier i hudens pigmentering sammanfattats under benämningarne "Chloasma", "Lentigo", "Pannus hepaticus", "Ephelis hepatica" ("lefverfläckar") m. fl.⁵⁾, blef sjukdomen först genom Eichstädt's upp-

¹⁾ "Hautkrankheiten" p. 113.

²⁾ "Area Celsi" i "Greifsw. Med. Beiträge" b. II.

³⁾ Se "Canst. Jahresb." 1864 p. 267.

⁴⁾ Prof. Malmsten med sin vidsträckt erfarenhet om hudsjukdomarne i vårt land har (1863) ej sett mer än 8 fall (Troilius). — Under förflutna tjänstgöring på Serafimer-Lazarettet förevisades der ett fall.

⁵⁾ Utan tvifvel betecknas ännu sjukdomen ofta med dessa namn; dock går Bazin för långt, som med densamma identifierar alla "Chloasmata", "maculae gravidarum", ja t. o. m. ephelides.

täckt af dess parasitiska natur rätt uppfattad och bestämdt begränsad. Af sednare författare har i synnerhet Gudden sysselsatt sig med dess studium.

Beskrifning. *Pityriasis versicolor* är en, nästan uteslutande till bålen inskränkt, smittosam hudsjukdom, orsakad af en i de ytliga epidermislagren förekommande mögelbildning, som föranleder uppkomsten af (vanl. gulaktiga) fläckar, på hvilka den degenererade epidermis är stadd i beständig affjällning, men som hvarken angriper håren eller beledsagas af några inflammatoriska eruptioner.

I sjukdomens början upptäcker man vanligen på bröstet eller ryggen flera eller färre, oftast ljust smutsgula, hampröstora, spridda fläckar, hvilka efter hand vidga sig och sammanflyta till större l. mindre, oregelbundna grupper eller figurer. Dessa nå ofta en betydlig utsträckning, så att de intaga nästan hela bröstet och ryggen samt derifrån sprida sig uppåt hals, axlar och öfverarmar, nedåt buken, klinkorna och mera sällan låren. Yttre delarne af extremiteterna angripas ytterst sällan, och troligen aldrig ansigte, händer och fötter.

Färgen är oftast ljusgul ("Pit. lutea"), men stundom äfven mörkare, brun eller n. svart ("P. nigra"), stundom mera grå eller rödaktig. Fläckarne, som ej äro upphöjda öfver hudens niveau, kännas torra, något ojemna och affjälla, i synnerhet vid skrapning, i mängd ett smutshvitt, kliliknande, småfjälligt stoft, bestående af parasiten, blandad med epidermisrester; då detta bortskrapas, synes den underliggande huden fullkomligt frisk. Det enda symptom, som föröfrigt åtföljer fläckarne, är en mer l. mindre häftig klåda, som under arbete eller efter förtärande af spirituosa (vid lifligare circulation) säges tilltaga.

Sjukdomen — som oftast öfverlemnas åt sig sjelf — plägar kvarstadna der den en gång tagit sin plats; likväl iakttager man stundom ett omvexlande till- och aftagande i fläckarnes utbredning. Ej sällan ser man parasitsvampen spontant utdö, då efter affjällning den gula färgen försvinner på fläckarne, hvilka derefter någon tid synas gråa, pigmentfattiga.

Ätiologi och Förekomst. Den mögelbildning, som ligger till grund för sjukdomen, öfverföres genom direct smitta från person till annan. (Hallier's hypothes om dess ursprung ur "Kolbrandens" sporer hafva vi förut (p. 17) omnämnt. Äfven här verkar bristande renlighet och vård af huden prädiposnerande, ehuru exempel finnas på dess förekomst äfven hos de renligaste personer.

Mera tvifvelaktig är dess förkärlek för personer med lymphatiskt temperament och som lida af bröstsjukdomar. Koch ¹⁾ fäster (eget nog) stor vikt vid ymnig svettafsöndring såsom ætiologiskt moment, under det Niemeyer säger sjukdomen sannolikt på grund af den rikligare svettsecretionen vanligen förbättras under sommarens. Egendomlig är dess afsky för alla blottade hudställen. — Män äro oftare angripna än qvinnor; hos barn är den knappt observerad.

Sjukdomen skall vara allmännare i heta climater, men är äfven i Sverige särdeles vanlig, ehuru — och med skäl — föga uppmärksammas.

Prognosen är alltid god. — Utan **Behandling** kvarstår visserligen sjukdomen, men gör just ej heller någon skada; underkastad behandling försvinner den lätt, recidiverar likväl ej sällan. Diverse medel äro härvid använda, såsom lösningar af borax, sublimat, svafvelkalium etc.; det enklaste och mest practiska är utan tvifvel några grundliga tvättninglar med såpvatten.

¹⁾ "Beitr. zur Lehre von den Epiphyten" ("Virch. Arch." b. X).

47
2.5
1.02
11.100

V. Descriptions of New Species of Fungi

BY CHAS. H. PECK.

[Read before this Society June 6th, 1873.]

HYMENOMYCETES.

Agaricus (Amanita) russuloides, Peck.

Pileus at first ovate, then expanded or convex, rough with a few superficial warts or entirely smooth, viscid when moist, widely striate-tuberculate on the margin, pale yellow or straw color; lamellae close, free, narrowed toward the stem, white; stem firm, smooth, stuffed, annulate, equal or slightly tapering upward, bulbous; annulus thin, soon vanishing; volva fragile, subappressed; spores broadly elliptical, .0004* long, .0003' broad.

Plant 2'-3' high, pileus 1.5'-2' broad, stem 3"†-5" thick.

Grassy ground in open woods. Greenbush, June. This species is remarkable for the thin striate-tuberculate margin of the pileus which causes it to resemble some species of *Russula*.

Agaricus (Lepiota) fuscousquameus, Peck.

Pileus hemispherical or convex, rough with numerous erect pointed blackish-brown scales; lamellae close, white, free; stem equal, thickened at the base, hollow or stuffed with a cottony pith, floccose, brown; spores .0003' × .00014'.

Plant 2'-3' high, pileus 1.5'-2' broad, stem 3" thick.

Ground in woods. Croghan. September.

Agaricus (Lepiota) oblitus, Peck.

Pileus convex or expanded, subumbonate, smooth or obscurely squamose from the breaking up of the veil, viscid, alutaceous inclining to tawny, the umbo generally darker; lamellae crowded, free, whitish or yellowish, some of

*One accent (') = inch or inches.

†Two accents (") = line or lines.

them forked; stem equal or slightly tapering upward, floccose, viscid, smooth at the top, hollow or containing a cottony pith; annulus obsolete; spores .00016' $\times$.00012'.

Plant 2'-3' high, pileus 2'-3' broad, stem 3" thick.

Ground in frondose woods. Lowville. September.

Agaricus (Armillaria) ponderosus, Peck.

Pileus thick, compact, convex or subcampanulate, smooth, white or yellowish, the naked margin strongly involute beneath the slightly viscid persistent veil; lamellae crowded, narrow, slightly emarginate, white inclining to cream color; stem stout, subequal, firm, solid, coated by the veil, colored like the pileus, white and furfuraceous above the annulus; flesh white; spores nearly globose, .00016' in diameter.

Plant 4'-6' high, pileus 4'-6' broad, stem about 1' thick.

Ground in woods. Copake. October.

The veil for a long time conceals the lamellae and finally becomes lacerated and adheres in shreds or fragments to the stem and margin of the pileus.

Agaricus (Tricholoma) rubicundus, Peck.

Pileus convex, then expanded or centrally depressed, viscid, slightly tomentose on the margin when young, smooth, or sometimes with a few scales either on the disk or on the margin, red; lamellae close, white, becoming spotted with red, some of them forked; stem firm, equal, solid, slightly pruinose, white, often stained with red; spores .00028' $\times$.00016'.

Plant 3'-5' high, pileus 3'-5' broad, stem 6"-8" thick.

Ground in woods. New Scotland. October.

Agaricus (Tricholoma) flavesceus, Peck.

Pileus firm, convex, often irregular, dry, smooth, sometimes cracking on the disk into minute scales, white or pale yellow, minutely tomentose on the margin when young; lamellae close, floccose on the edge, white or pale yellow; stem firm, solid, often unequal, central or eccentric, colored like the pileus; spores subglobose, .0002' in diameter.

Plant caespitose, 2'-3' high, pileus 2'-3' broad, stem 4"-6" thick.

Old pine stumps. Bethlehem and North Greenbush. October.

Agaricus (Tricholoma) decorosus, Peck.

Pileus firm, at first hemispherical, then convex or expanded, coated with numerous brownish subsquarrose tomentose scales, dull ochraceous or tawny; lamellae close, rounded and slightly emarginate at the inner extremity, the

edge suberenulate; stem solid, equal or slightly tapering upward, white and smooth at the top, elsewhere tomentose, scaly and colored like the pileus; spores broadly elliptical, $.0002' \times .00015'$.

Plant subcaespitose, 2'-4' high, pileus 1'-2' broad, stem 2"-4" thick.

Rotten logs in woods. Catskill Mountains and Rock City. September and October.

Agaricus (Tricholoma) multipunctus, Peck.

Pileus brittle, broadly convex, sometimes centrally depressed or subumbilicate, densely dotted with minute brown or blackish scales, yellowish-brown, the disk often darker; lamellae close, slightly emarginate, yellow, sometimes with a darker edge; stem subequal, squamulose-punctate, hollow, colored like the pileus; spores suborbicular, $.00016'$ in diameter.

Plant subcaespitose, 1'-2' high, pileus 1'-2' broad, stem 2"-4" thick.

Rotten logs in woods. Sandlake and Adirondack Mountains. July and August.

This species is related to *Ag. rutilans*.

Agaricus (Tricholoma) lacunosus, Peck.

Pileus convex or expanded, dry, lacunose, densely furfuraceous, bright golden yellow; lamellae subdistant, white, the interspaces sometimes veiny; stem firm, solid, equal or slightly tapering downwards, scaly or furfuraceous, colored like the pileus.

Plant 1'-2' high, pileus 1' broad, stem 1" thick.

Fallen branches and decaying wood. Savannah. August.

The colors are well retained in the dried specimens. The lacunae of the pileus give it a somewhat reticulated appearance.

Agaricus (Tricholoma) laterarius, Peck.

Pileus convex or expanded, sometimes slightly depressed in the center, pruinose, whitish, the disk often tinged with red or brown, the thin margin marked with slight subdistant short radiating ridges; lamellae narrow, crowded, white, prolonged in little decurrent lines on the stem; stem nearly equal, solid, white; spores globose, $.00018'$ in diameter.

Plant 3'-4' high, pileus 2'-4' broad, stem 3"-5" thick.

Ground in woods. Worcester. July.

Agaricus (Tricholoma) Limonium, Peck.

Pileus thin, smooth, yellowish; lamellae crowded, narrow, not forming decurrent lines on the stem, lemon-yellow; stem tapering downwards, smooth, striate, rooting.

Plant 3'-4' high, pileus 2'-3' broad, stem 3"-4" thick.

Ground in woods. Worcester and Croghan. July and September.

The lemon color of the lamellae and the root-like prolongation of the stem characterize this species.

Agaricus (Tricholoma) virescens, Peck.

Pileus convex or expanded, sometimes depressed centrally, moist, smooth, dingy-green, the margin sometimes wavy or lobed; lamellae close, gradually narrowed toward the outer extremity, rounded or slightly emarginate at the inner, white; stem subequal, stuffed or hollow, thick but brittle, whitish, sometimes tinged with green; spores broadly elliptical, $.0002' \times .00015'$.

Plant 3'-5' high, pileus 3'-5' broad, stem 6"-12" thick.

Mossy ground in open woods. North Elba. July.

Agaricus (Tricholoma) fumidellus, Peck.

Pileus subumbonate, smooth, moist, dingy-white or clay colored, clouded with brown; lamellae close, subventricose, whitish; stem equal, smooth, solid, whitish; spores, $.00018' \times .00015'$.

Plant 2'-3' high, pileus 1'-2' broad, stem 2"-3" thick.

Ground in woods. New Scotland. October.

The disk is generally darker than the margin. The pileus becomes paler in drying. The stem splits easily.

Agaricus (Tricholoma) fallax, Peck.

Pileus firm, convex or expanded, rarely depressed in the center, moist, smooth, dull saffron color; lamellae crowded, narrow, tapering toward the outer extremity, rounded at the inner, yellow; stem short, smooth, stuffed or hollow, usually tapering toward the base, colored like the pileus; spores minute, subelliptical, $.00012'$ long.

Plant gregarious, 1'-1.5' high, pileus 6"-15" broad, stem 1" thick.

Ground under spruce and balsam trees. North Elba. July.

Agaricus (Tricholoma) thujinus, Peck.

Pileus convex or centrally depressed, smooth, hygrophanous, pale alutaceous, the margin generally irregular, wavy or lobed; lamellae crowded, thin, abruptly emarginate, alutaceous; stem slightly thickened at the top, smooth, hollow, concolorous, whitish-villous at the base.

Plant 2' high, pileus 2' broad, stem 2"-3" thick.

Swampy ground under *Thuja occidentalis*. Memphis. August.

Agaricus (Tricholoma) Hebeloma, Peck.

Pileus broadly conical or subcampanulate, obtuse, thin, hygrophanous, striatulate brown and dark on the disk when moist, grayish when dry; lamellae broad, rounded behind and deeply emarginate, yellowish; stem equal, hollow, smooth, pallid; spores $.00028' \times .00016'$.

Plant 1' high, pileus 6" broad, stem scarcely 1" thick.

Ground in woods. Worcester. July.

This plant closely resembles some species of the subgenus *Hebeloma*, but the color of its spores forbid its reference to that subgenus.

Agaricus (Clitocybe) connexus, Peck.

Pileus thin, subumbonate, clothed with a minute appressed silkiness, white, the margin sometimes faintly tinged with blue; lamellae crowded, narrow, white inclining to yellowish; stem equal or tapering downwards, solid, whitish.

Plant 2'-3' high, pileus 2'-3' broad, stem 2" thick.

Ground in woods. Croghan. September.

The lamellae are not strongly decurrent and sometimes terminate abruptly, hence it might easily be mistaken for a *Tricholoma*. The margin of the pileus is sometimes marked with slight ridges as in *Ag. laterarius*. The odor is weak but aromatic and agreeable.

Agaricus (Clitocybe) albissimus, Peck.

Pileus convex or expanded, dry, smooth, soft, pure white; lamellae crowded, short-decurrent, white, some of them forked at the base; stem equal, smooth, solid, white.

Plant growing in rings, 2'-3' high, pileus 2'-3' broad, stem 2"-3" thick.

Ground in woods. Croghan. September.

The pure white color and soft texture is retained in the dried specimens.

Agaricus (Clitocybe) maculosus, Peck.

Pileus centrally depressed, smooth, marked with numerous watery spots when moist, having slight short radiating ridges on the margin; lamellae crowded, narrow, long-decurrent, pallid or yellowish, some of them forked; stem slightly thickened at the base, smooth, stuffed or hollow, colored like the pileus.

Plant 2'-3' high, pileus 1.5'-2' broad, stem 2"-3" thick.

Ground in woods. Croghan. September.

The spots of the pileus resemble those of *Ag. marmoreus*. They generally disappear as the plant becomes dry.

Agaricus (Clitocybe) Truncicola, Peck.

Pileus thin, firm, expanded or centrally depressed, smooth, dry, white; lamellae narrow, thin, crowded, adnate-decurrent; stem equal, stuffed, smooth, often eccentric and curved, whitish.

Plant 1' high, pileus 1' broad, stem 1" thick.

Trunks of frondose trees, especially maples. Croghan. September.

Agaricus (Clitocybe) subzonalis, Peck.

Pileus thin, centrally depressed or subinfundibuliform, marked with two or three obscure zones, with a slight appressed silkiness, pale yellow; lamellae close, narrow, equally decurrent, some of them forked, pallid or yellowish; stem equal, slightly fibrillose, stuffed, pale yellow.

Plant 2' high, pileus 2'-3' broad, stem 2"-4" thick.

Ground in woods. Croghan. September.

Agaricus (Clitocybe) Gerardianus, Peck.

Pileus thin, funnel-form, hygrophanous, striatulate when moist, brown, rough with scattered blackish points; lamellae decurrent, close, a little paler than the pileus, some of them forked; stem rather long, flexuous, smooth, stuffed, concolorous, white at the base.

Plant 2'-3' high, pileus 8"-12" broad, stem .5"-1" thick.

Sphagnous marshes. Sandlake and New Paltz. June.

Agaricus (Collybia) coloreus, Peck.

Pileus convex, subumblicate, slightly fibrillose, hygrophanous, yellow, sometimes tinged with red, the margin exceeding the lamellae; lamellae moderately close, emarginate, yellow; stem equal, smooth, hollow, sometimes eccentric, yellow.

Plant 1'-2' high, pileus 8"-12" broad, stem 1" thick.

Decaying wood. Croghan. September.

Agaricus (Collybia) succosus, Peck.

Pileus firm, convex or campanulate, minutely tomentose, cinereous or brownish-gray, the margin generally exceeding the lamellae; lamellae thin, close, emarginate and slightly decurrent-toothed, tapering toward the outer extremity, whitish; stem firm, equal or slightly tapering upward, minutely tomentose, containing a whitish pith; spores minute, subglobose, .00015' in diameter; flesh subcartilaginous, abounding in a thin watery or serum-like juice, changing to purplish or black when cut.

Plant 1'-3' high, pileus 6"-12" broad, stem 1" thick.

Decaying prostrate trunks of trees in woods. Portville and Croghan. September.

This is a very remarkable and somewhat aberrant species. In color it resembles dark forms of *Hydnum gelatinosum*. The stem is sometimes eccentric. The juice exudes from wounds as in species of *Lactarius*.

Agaricus (Collybia) myriadophyllus, Peck.

Pileus very thin, broadly convex, then expanded, sometimes umbilicate, hygrophanous, watery-brown when moist, pale ochraceous or alutaceous when dry; lamellae very numerous, crowded, narrow, rounded at the stem and slightly emarginate, brownish-lilac; stem equal, smooth, stuffed, reddish-brown; spores subelliptical, minute, .00012' long.

Plant subcaespitose, 1'-1.5' high, pileus 8''-12'' broad, stem .5'' thick.

Decaying wood and fallen branches in woods. Portville. September.

The color of the lamellae is remarkable.

Agaricus (Mycena) subcaeruleus, Peck.

Pileus very thin, convex or campanulate, obtuse, smooth, striate, pale bluish-green; lamellae narrow, close, tapering outwardly, white; stem slender, equal, pinkish-white, slightly pruinose; spores subglobose, .00025' in diameter.

Plant caespitose, 2' high, pileus 4''-8'' broad.

Trunks of beech trees in woods. Adirondack Mountains. July.

The disk is more highly colored than the margin and the pileus has a separable cuticle.

Agaricus (Mycena) minutulus, Peck.

Pileus convex or campanulate, smooth, striatulate, papillate; lamellae broad, subdistant, with a slight decurrent tooth; interspaces reticulated by transverse veinlets which descend on the lamellae; stem short, slender, firm, smooth or sprinkled with minute mealy particles.

Plant gregarious, white throughout, 8''-12'' high, pileus 2''-4'' broad.

Bark of prostrate trunks in woods. Portville. September.

Agaricus (Mycena) roseocandidus, Peck.

Pileus convex or broadly campanulate, subpapillate, striate nearly to the apex, white or rosy-red; lamellae close, uncinata, colored like the pileus; stem slender, smooth, white.

Plant 2' high, pileus 4''-6'' broad.

Among mosses in woods. Adirondack Mountains. July.

Usually the whole plant is pure white, but sometimes the pileus has a delicate rosy hue except on the apex and the margin. The striations of the pileus remain in the dried specimens. The papilla is sometimes very prominent, sometimes wanting.

***Agaricus (Mycena) miratus*, Peck.**

Pileus thin, campanulate, umbilicate, smooth, striate, cinereous; lamellae narrow, slightly uncinuate, whitish; stem long, filiform, smooth, whitish, radicating, villous at the base.

Plant 1.5'-2' high, pileus 3''-4'' broad.

Among fallen leaves in copses. Center. October.

This species may be known by the umbilicate pileus and the long striae which extend to the umbilicus.

***Agaricus (Omphalia) olivarius*, Peck.**

Pileus convex, umbilicate, smooth, yellowish-olive; lamellae arcuate, decurrent, subdistant, pale yellow; stem equal, short, smooth, hollow, colored like the pileus; spores subglobose or broadly elliptical, .00026' long.

Plant 1'-1.5' high, pileus 1' broad, stem 1'' thick.

Burnt ground under balsam trees. North Elba. July.

***Agaricus (Omphalia) rugosodiscus*, Peck.**

Pileus thin, convex, then expanded, smooth, hygrophanous, striatulate when moist, brown, rugose-wrinkled on the disk, the thin margin often wavy; lamellae narrow, close, arcuate, decurrent, white; stem equal, short, smooth, hollow, often curved, whitish.

Plant 1'-1.5' high, pileus 6"-12" broad, stem .5" thick.

Decaying prostrate trunks of trees in woods. Croghan and Worcester. July and September.

***Agaricus (Pluteus) sterilomarginatus*, Peck.**

Pileus broadly convex or expanded, with a slight appressed tomentum, white faintly tinged with pink, the thin margin exceeding the lamellae; lamellae close, subventricose, free, minutely eroded on the edge, pale flesh color; stem short, equal, solid, smooth, whitish; spores subglobose, angular, with a central nucleus, .00025' in diameter.

Plant 1' high, pileus 6"-12" broad, stem .5" thick.

Decaying wood and sticks in woods. Portville. September. The pileus is sometimes cracked and then has the appearance of being coated with a thin scaly paste.

Agaricus (Pluteus) granularis, Peck.

Pileus convex or expanded, subumbonate, rugose-wrinkled, sprinkled with minute blackish granules, varying in color from yellow to brown; lamellae rather broad, close, ventricose, free, whitish, then flesh-colored; stem equal, solid, pallid or brown, usually paler at the top, velvety, with a short close plush; spores subglobose, about .0002' in diameter.

Plant 2'-3' high, pileus 1'-2' broad, stem 1"-2" thick.

Old logs in woods. Pine Hill and Worcester. July.

The granules form a sort of plush which is more dense on the disk of the pileus and its wrinkles than on the margin.

Agaricus (Entoloma) cyaneus, Peck.

Pileus convex, dry, minutely scaly, brown or brownish-violaceous; lamellae whitish, then tinged with flesh color; stem subequal, hollow, scaly and violaceous toward the top; spores angular, .00033' $\times$.00025'.

Plant 2' high, pileus 1'-1.5' broad, stem 1" thick.

Decaying wood and old mossy trunks in woods. Pine Hill and Worcester. June and July.

Agaricus (Leptonia) foliomarginatus, Peck.

Pileus convex, umbilicate, scabrous on the disk, bluish-brown, the disk a little darker; lamellae broad, subdistant, plane, whitish, then flesh-colored, the edge entire and colored like the pileus; stem smooth, equal, solid below, with a small cavity above, concolorous.

Plant 1'-2' high, pileus 6"-10" broad, stem .5" thick.

Ground and decaying wood in groves. Maryland. July.

Agaricus (Nolanea) fuscifolius, Peck.

Pileus thin, conical or campanulate, papillate, smooth, hygrophanous, dark brown and striatulate when moist, grayish-brown and shining when dry; lamellae ascending, narrowed toward each end, brown; stem equal, stuffed, smooth, concolorous, with a white mycelium at the base; spores irregular, nucleate, .00033' $\times$.00025'.

Plant 1' high, pileus 3"-6" broad, stem 5" thick.

In woods on old logs. Maryland. July.

Agaricus (Pholiota) albocrenulatus, Peck.

Pileus fleshy, firm, convex or campanulate, subumbonate, viscid, rough with dark-brown or blackish floccose scales, yellowish-brown; lamellae broad, subdistant, emarginate, white crenulate on the edge, grayish, then ferruginous;

stem firm, equal or slightly tapering upward, stuffed or hollow, squamose and pallid below the evanescent ring, white and slightly furfuraceous above; spores subelliptical, $.00045' \times .00025'$.

Plant 3'-5' high, pileus 2'-3' broad, stem 3"-5" thick.

Mossy base of maple trees in woods. Adirondack Mountains. July and August.

Under a lens the lamellae appear to be beaded on the edge with minute milky globules.

Agaricus (Pholiota) Acericola, Peck.

Pileus broadly convex, glabrous, rugose-reticulated or corrugated, hygrophanous, yellow; lamellae close, emarginate, grayish, then ferruginous-brown; stem equal or thickened at the base, hollow, fibrillose-striate, white; annulus large; spores elliptical, $.00035' \times .00025'$.

Plant 3'-4' high, pileus 2'-3' broad, stem 3"-5" thick.

Mossy trunks of maple trees in woods. North Elba. August.

Agaricus (Pholiota) discolor, Peck.

Pileus thin, convex, then expanded or slightly depressed, smooth, viscid hygrophanous, watery-cinnamon and striatulate on the margin when moist, bright ochraceous yellow when dry; lamellae close, narrow, pallid, then pale ferruginous; stem equal, hollow, fibrillose-striate, pallid; annulus distinct, persistent; spores elliptical, $.00028' \times .0002'$.

Plant subcaespitose, 2'-3' high, pileus 8"-16" broad, stem 1" thick.

Old logs in woods. Greig. September.

The change in color when passing from the moist to the dry state is very marked.

Agaricus (Pholiota) cerasinus, Peck.

Pileus broadly convex, smooth, hygrophanous, watery-cinnamon when moist, yellow when dry; lamellae close, emarginate, yellow, then cinnamon color; stem solid, equal, often curved, furfuraceous at the top; annulus slight, fugacious; flesh yellow; spores elliptical, rough, $.0003' \times .0002'$.

Plant caespitose, 2'-4' high, pileus 2'-4' broad, stem 2'-4" thick.

Old prostrate trunks of trees in woods. Sterling. August.

When fresh it has a strong cherry-like or amygdaline odor.

Agaricus (Hebeloma) pallidomarginatus, Peck.

Pileus brittle, broadly convex, sometimes irregular, smooth, hygrophanous, brown with a pale margin when moist, ochraceous and subatomaceous when dry; lamellae close, thin, rounded and slightly emarginate at the stem, taper-

ing outwardly, ochraceous-brown; stem usually long and flexuous; equal or tapering upward, hollow, a little paler than the pileus, white-floccose at the base; spores subelliptical, $.0004' \times .0002'$.

Plant gregarious 1'-3' high, pileus 6"-12" broad, stem 1" thick.

Ground in swamps and wet places. Sandlake. September.

Agaricus (Hebeloma) stellatosporus, Peck.

Pileus convex, dry, rough with numerous squarrose or erect scales, brown; lamellae pallid becoming brown; stem equal, scaly, concolorous; spores subglobose, rough with little nodules, $.0003'$ in diameter.

Plant 2' high, pileus 1' broad, stem 1" thick.

Ground in woods. Croghan. September.

This plant bears a close resemblance to *Ag. mutatus*, but the persistent scales and rough spores distinguish it.

Agaricus (Hebeloma) griseoscabrosus, Peck.

Pileus hemispherical or convex, dry, rough with scales and appressed fibres, cinereous, the margin whitish when young; lamellae close, broad, whitish, then ochraceous-brown; stem firm, solid, fibrillose or slightly scaly, subconcolorous; spores smooth, $.00035' \times .0002'$.

Plant gregarious, 1.5'-2' high, pileus 6"-10" broad, stem 1"-1.5" thick.

Ground in open pine woods. Bethlehem. October.

Agaricus (Naucoria) bellulus, Peck.

Pileus thin, convex, moist, smooth, bright watery-cinnamon; lamellae crowded, narrow, emarginate, yellow, becoming darker with age; stem equal, hollow, often curved, smooth, reddish-brown; spores, $.0002' \times .00014'$.

Plant 1' high, pileus 6"-12" broad.

Decaying hemlock trunks in woods. Lowville and Sandlake. September.

It is sometimes caespitose. It is difficult to find a mature specimen of this plant in which the lamellae have not a stained or spotted appearance as if bitten by some small insect.

Agaricus (Naucoria) geminellus, Peck.

Pileus convex, even, firm, dry, yellowish-red, the margin paler; lamellae crowded, emarginate, pale yellow; stem equal, smooth, containing a white pith or a small cavity, colored like the pileus; flesh white; spores $.00033' \times .0002'$.

Rotten wood. Croghan. September.

The dimensions and habit are the same as in the last species, to which this is clearly related. Its lamellae also have the same peculiar appearance.

Agaricus (Naucoria) discomorbidus, Peck.

Pileus thin, convex or expanded, smooth, slightly viscid, reddish-brown or dull chestnut; lamellae narrow, crowded, minutely serrulate, white or pallid, then brownish; stem equal, stuffed, smooth, slightly mealy at the top, white; flesh white; spores nucleate, $.0004' \times .00025'$.

Plant 2'-3' high, pileus 1'-1.5' broad, stem 1''-2'' thick.

Ground in woods. Croghan and Copake. September and October.

In the dried specimens the disk has a dark discolored appearance as if beginning to decay.

Agaricus (Galera) expansus, Peck.

Pileus submembranaceous, expanded or depressed, viscid, plicate striate on the margin, brownish-ochre, sometimes tinged with yellow and pink hues; lamellae close, ferruginous; stem long, equal, hollow, slightly pruinose, faintly striate, yellow; spores $.00045' \times .00028'$.

Plant 3'-4' high, pileus 1' broad, stem 1'' thick.

Decaying wood. Sandlake and Memphis. August.

Agaricus (Galera) callistus, Peck.

Pileus thin, expanded, subumbonate, smooth, viscid, striatulate on the margin, olivaceous or ochraceous, the umbo or disk bright chestnut color; lamellae thin, close, ventricose, easily separating from the stem, yellowish, becoming bright ferruginous; stem equal, hollow, pruinose, yellow, spores $.00035' \times .0002'$.

Plant 1'-1.5' high, pileus 6''-10'' broad, stem .5'' thick.

Exsiccated water-holes in wooded swamps. Croghan. September.

In the dried specimens the lamellae are white on the edge and the pileus has assumed a dull metallic green color.

Agaricus (Galera) Coprinoides, Peck.

Pileus membranaceous, soon expanded, often split on the margin, plicate-sulcate to the small even disk, yellowish inclining to ochre; lamellae close, slightly rounded behind, concolorous; stem equal, short, hollow, minutely hairy-pruinose, white; spores $.00028' \times .0002'$.

Plant 1' high, pileus 6'' broad, stem .5'' thick.

Grassy ground. Sterling. August.

The appearance of the pileus is suggestive of some of the smaller Coprini.

Agaricus (Crepidotus) Herbarum, Peck.

Pileus thin, at first resupinate, with the margin incurved, clothed with white down, at length somewhat reflexed, less downy, the margin spreading; lamellae narrow, not crowded, diverging from a naked lateral or eccentric point, white, then tawny; spores slightly curved, $.00028' \times .00014'$.

Pileus 2"-4" broad.

Dead stems of herbs. North Greenbush. October.

Agaricus (Psalliota) diminutivus, Peck.

Pileus expanded or centrally depressed, sometimes with a slight umbo, dry, alutaceous, the disk rosy-brown and spotted with small appressed silky scales; lamellae close, thin, free, ventricose, brownish-pink, becoming black; stem equal or slightly tapering upward, hollow or stuffed with a whitish pith, smooth, pallid; annulus thin, persistent, white; spores $.0002' \times .00015'$.

Plant 1.5'-2' high, pileus 1'-1.5' broad, stem 1"-2" thick.

Ground in woods. Croghan. September.

Sometimes the whole pileus is reddish-brown. The flesh is quite brittle.

Agaricus (Stropharia) Howeanus, Peck.

Pileus convex, then expanded, fragile, smooth, subumbonate, yellowish; lamellae close, thin, rounded behind, eroded on the edge, whitish becoming ferruginous-brown; stem smooth, hollow, slightly thickened at the base; annulus thin, fugacious, sometimes adhering to the margin of the pileus; flesh white; spores $.00033' \times .0002'$.

Plant 3'-4' high, pileus 2'-3' broad, stem 2"-4" thick.

Center. June.

The surface of the pileus sometimes cracks into areas. The taste is bitter. The color of the spores is not a decided brown, and the plant might with almost equal propriety be referred to the subgenus *Pholiota*.

Agaricus (Hypholoma) hirtosquamulosus, Peck.

Pileus hairy-squamulose, hygrophanous, grayish-brown when moist, gray when dry; lamellae narrow, rounded at the stem, gray, then brown; stem short, firm, equal, hollow, slightly hairy-squamulose and colored like the pileus; spores subelliptical, nucleate, $.00025'$ long.

Plant 1' high, pileus 6"-10" broad, stem .5" thick.

Prostrate trunks of maple trees in woods. Portville. September.

Agaricus (Hypholoma) phyllogenus, Peck.

Pileus firm, convex, sometimes slightly umbonate, hygrophanus, reddish-brown when moist, alutaceous when dry; lamellae plane, broad, close, brown, white on the edge; stem equal, fibrillose, stuffed or hollow, expanded at the base into a thin flat disk; spores pale-brown, subglobose, .0002' in diameter.

Plant 8"-12" high, pileus 2"-4" broad, stem .5" thick.

Fallen leaves in woods. Worcester. July.

This is a very small but distinct species, remarkable for the disk-like base of the stem by which it is attached to the leaves on which it grows.

Coprinus variegatus, Peck.

Pileus fleshy, fragile, oblong-ovate, then campanulate, obtuse, hygrophanous, pale watery-brown when moist, whitish or cream colored when dry, variegated by scales and patches of a superficial ochraceous tomentum, the margin finely striate; lamellae lanceolate, crowded, ascending, free, white, then rosy-brown, finally black; stem equal, brittle, hollow, white, at first peronate-annulate, then floccose-pruinose, with white branching root-like threads at the base; spores .00033' long.

Plant densely caespitose, 3'-5' high, pileus 1'-1.5' broad, stem 2"-4" thick.

Thin soil and decaying leaves covering rocks. Slope of Crows' Nest near West Point. June.

Allied to *C. atramentarius*. When young the whole plant is coated by an abundant superficial tomentum. This soon breaks up into loose scales or patches which peel off in flakes, revealing the smooth pileus beneath. The slight abrupt annulus soon vanishes.

Coprinus insignis, Peck.

Pileus campanulate, thin, sulcate-striate to the disk, grayish fawn-color, the smooth disk sometimes cracking into small areas or scales; lamellae ascending, crowded; stem hollow, slightly fibrillose, striate, white; spores rough, .0004' x .00028'.

Plant 4'-5' high, pileus 2'-3' broad, stem 3" thick.

About the roots of trees in woods. Worcester. July.

Coprinus angulatus, Peck.

Pileus thin, hemispherical or convex, plicate-sulcate, the disk smooth; lamellae subdistant, whitish, then black; stem equal, smooth, whitish; spores compressed, angular, subovate, .0004' x .00033'.

Plant 1'-2' high, pileus 6"-12" broad, stem .5" thick.

Woods. Croghan. September.

The specific name has reference to the angular character of the spores.

Cortinarius (Myxacium) sphaerosporus, Peck.

Pileus convex, smooth, very viscid, pale ochraceous; lamellae close, nearly plane, slightly emarginate, whitish, then cinnamon; stem tapering upward, solid, floccose, viscid, subconcolorous, white at the top; flesh white; spores nearly globose, about .0003' in diameter.

Plant 2'-4' high, pileus 2'-3' broad, stem 3"-5" thick.

Ground in woods. Croghan. September.

Cortinarius (Phlegmacium) longipes, Peck.

Pileus convex or expanded, slightly fibrillose, viscid, yellowish or pale ochraceous; lamellae close, plane, brownish-olivaceous, then cinnamon; stem long, slightly fibrillose, tapering upwards, whitish.

Plant 6' high, pileus 2'-3' broad, stem 4" thick.

Ground in woods. Croghan. September.

Cortinarius (Inoloma) lilacinus, Peck.

Pileus firm, hemispherical, then convex, minutely silky, lilac; lamellae close, lilac, then cinnamon; stem stout, bulbous, silky fibrillose, solid, whitish tinged with lilac; spores nucleate, .0004' $\times$.00025'.

Plant 4'-5' high, pileus 3' broad, stem 4"-6" thick.

Low mossy ground in woods. Croghan. September.

Cortinarius (Inoloma) Clintonianus, Peck.

Pileus convex or expanded, with a few appressed silky fibrils, reddish-brown tinged with gray; lamellae close, dull violaceous, then cinnamon; stem solid, silky-fibrillose, tapering upwards, violaceous at the top; spores .0003' $\times$.00025'.

Plant 2'-3' high, pileus 1'-2' broad, stem 2"-3" thick.

Ground in woods. Croghan and New Scotland. September.

Cortinarius (Inoloma) modestus, Peck.

Pileus convex or expanded, subfibrillose, even or slightly rugose-wrinkled, alutaceous; lamellae close, nearly plane, pallid, then cinnamon; stem bulbous, subfibrillose, hollow or with a white pith, concolorous; flesh white; spores .00033' $\times$.00025'.

Plant 2' high, pileus 1'-1.5' broad, stem 2" thick.

Ground in woods. Croghan. September.

It is distinguished from the preceding species by its paler color, more bulbous stem, and the entire absence of violaceous hues in the lamellae.

Cortinarius (Telamonia) lignarius, Peck.

Pileus smooth, hygrophanous, dark watery cinnamon when moist, paler when dry; lamellae close, thin, concolorous, when young concealed by the copious white webby veil; stem equal, silky-fibrillose, hollow or with a whitish pith, subannulate, with a dense white mycelium at the base; spores .00028' $\times$.0002'.

Plant subcaespitose, 1'-2' high, pileus 8''-12'' broad, stem 1'' thick.

Rotten wood. Catskill mountains. June.

Cortinarius (Telamonia) nigrellus, Peck.

Pileus at first conical, then convex or expanded, obtuse or subumbonate, minutely silky, hygrophanous, blackish chestnut when moist, paler when dry; lamellae close, narrow, emarginate, brownish-ochre, then cinnamon; stem subequal, silky fibrillose, pallid, often flexuous; annulus slight, evanescent; spores .00028' $\times$.00016'.

Plant 2'-3' high, pileus 1'-2' broad, stem 2''-3'' thick.

Mossy ground in woods. New Scotland. October.

When moist the pileus has the color of boiled chestnuts, when dry, of fresh ones. The incurved margin of the young pileus is whitened by the veil. The lamellae are darkest when young.

Cortinarius (Hygrocybe) pulcher, Peck.

Pileus conical, then broadly convex, umbonate, often irregular, hygrophanous, ochraceous, shining and sometimes striatulate when moist, pale ochraceous when dry; lamellae subdistant, broad, emarginate, uneven on the edge, ochraceous, stem equal, solid, subflexuous, silky-fibrillose, whitish or pale ochraceous; spores .00033' $\times$.0002'.

Plant gregarious, 2' high, pileus 1'-1.5' broad, stem 1''-2'' thick.

Ground in woods. New Scotland. October.

Paxillus strigosus, Peck.

Pileus dry, convex or expanded, brittle, strigose with scattered stiff hairs, whitish; lamellae close, narrow, subdecurrent, whitish, then pale cinnamon color, some of them forked; stem equal, solid, pruinose, concolorous; spores brownish-ochre, subglobose, .00018' in diameter.

Plant 2' high, pileus 1'-1.5' broad, stem 1''-1.5'' thick.

Ground among fallen leaves in woods. Can^{ro}aghan. September.

The young plant might readily be mistaken for a species of Clitocybe. Owing to the very brittle character of the pileus, the lamellae are not easily separated from it. The hairs of the pileus are either erect or appressed.

Lactarius regalis, Peck.

Pileus convex, deeply depressed in the center, viscid when moist, often corrugated on the margin, white tinged with yellow; lamellae close, decurrent, whitish, some of them forked at the base; stem stout, short, equal, hollow, smooth; taste acrid; milk sparse, white quickly changing to sulphur-yellow; spores .0003'.

Plant 4'-6' high, pileus, 4'-6' broad, stem 1' thick.

Ground in woods. Croghan. September.

This interesting plant rivals *L. piperatus* in size and closely resembles it in general appearance, but the viscid pileus and sparse milk quickly changing to yellow, as in *L. chrysorrheus*, clearly distinguish it.

Lactarius Geradii, Peck.

Pileus expanded or centrally depressed, dry, rugose-wrinkled, often with a minute umbo or papilla, sooty-brown, the thin spreading margin sometimes wavy or irregular; lamellae broad, distant, decurrent, white, the interspaces uneven; stem equal, solid, colored like the pileus; flesh and spores white; taste mild; milk white and unchangeable.

Plant 3'-5' high, pileus 2'-4' broad, stem 4''-6'' thick.

Ground in woods and groves. Poughkeepsie, *W. R. Gerard*. Albany and Croghan. September.

In the color of the pileus and stem, this species is like the large variety of *L. fuliginosus*, but its real relationship is with *L. distans*, from which it is separated by its color and its longer equal stem, characters which may prove to be only varietal.

Russula sordida, Peck.

Pileus firm, convex, centrally depressed, dry, sordid white, sometimes clouded with brown; lamellae white, some of them forked; stem equal, solid, concolorous; spores globose, .0003'; taste acrid; flesh changing color when wounded, becoming black or bluish-black.

Plant 4'-5' high, pileus 3'-5' broad, stem 6''-12'' thick.

Ground under hemlock trees. Worcester. July.

The whole plant turns black in drying.

Marasmius semihirtipes, Peck.

Pileus thin, tough, nearly plane or depressed, smooth, sometimes striate on the margin, hygrophanous, reddish-brown when moist, alutaceous when dry, the disk sometimes darker; lamellae subdistant, reaching the stem, slightly

venose-connected, subcrenulate on the edge, white; stem equal, hollow, smooth above, velvety-tomentose toward the base, reddish-brown.

Plant gregarious, 1'-2' high, pileus 6"-9" broad, stem .5" thick.

On and among fallen twigs and leaves. West Point. June.

***Marasmius umbonatus*, Peck.**

Pileus thin, tough, expanded, umbonate, smooth, even or substrate, alutaceous, the margin at first incurved; lamellae narrow, subdistant, reaching the stem, venose-connected, sometimes branched toward the outer extremity, white; stem equal, solid, velvety-tomentose, tawny below, paler above.

Plant gregarious, 1'-1.5' high, pileus 6"-9" broad, stem .5" thick.

Ground under balsam trees. North Elba. July.

***Marasmius caespitosus*, Peck.**

Pileus fleshy, convex, even, brown, with a lilac tint, the thin margin exceeding the lamellae; lamellae close, free, somewhat united with each other at the stem, narrowed outwardly, white; stem sometimes compressed at the top, stuffed or hollow, pruinose.

Plant caespitose, 1'-2' high, pileus 6"-10" broad.

Birch stumps in woods. Richmondville, June.

***Marasmius longipes*, Peck.**

Pileus thin, convex, smooth, finely striate on the margin, tawny-red; lamellae white; stem tall, straight, equal, hollow, pruinose-tomentose, radicate, brown or fawn color, white at the top.

Plant 2'-5' high, pileus 4"-6" broad, stem .5" thick.

Among fallen leaves in woods. Savannah and Bethlehem. August and October.

The long straight slender stem is a characteristic feature of this plant.

***Marasmius glabellus*, Peck.**

Pileus membranaceous, convex, then expanded, distantly striate, often uneven on the disk, dingy-ochraceous; lamellae broad, distant, unequal free, ventricose, whitish, the upper margin and the interspaces venose; stem corneous, equal, smooth, shining, hollow, reddish-brown or chestnut, whitish at the top, with a thick mycelium at the base.

Plant 1'-2' high, pileus 6"-10" broad, stem .5" thick.

Fallen leaves in woods. Worcester and Croghan. July and September.

Marasmius straminipes, Peck.

Pileus membranaceous, hemispherical or convex, smooth, striate, whitish; lamellae distant, unequal, white; stem corneous, smooth, shining, filiform, inserted, pale straw color.

Plant 1'-2' high, pileus 1"-3" broad.

Fallen leaves of the pitch pine, *Pinus rigida*. Center. October.

Lenzites vialis, Peck.

Pileus coriaceous, sessile, dimidiate or elongated, sometimes confluent, obscurely zoned, submentose, brown or grayish-brown, the margin cinereous; lamellae thin, anastomosing abundantly, pallid, cinereous-pruinose on the edge when fresh.

Pileus 6"-12" long.

Old railroad ties. North Greenbush and Center. October.

Boletus separans, Peck.

Pileus thick, convex, smooth, shining, sometimes deeply lacunose, brownish-lilac; tubes plane or slightly depressed around the stem, at first quite closed and attached to the stem, then by the expansion of the pileus usually torn from it, small, subrotund, yellow or brownish-yellow; stem solid, nearly equal, distinctly reticulated, dull lilac; spores $.00055' \times .00022'$; flesh white, unchangeable.

Plant 3'-4' high, pileus 3' broad, stem 6"-10" thick.

Grassy ground in open woods. Greenbush. August.

In dry weather the separation of the tubes from the stem does not always take place.

Boletus affinis, Peck.

Pileus dry, minutely tomentulose, even or slightly rugose, chestnut colored, soon fading to tawny or ochraceous, the cuticle sometimes cracking into areas; tubes plane or convex, attached to the stem and sometimes depressed around it, at first white and closed, then yellow, small, unequal, angular or subrotund; stem solid, unequal, smooth, rarely reticulated at the top, pallid or tinged with dull red; spores $.00035' \times .00016'$; flesh white, unchangeable.

Plant 2'-3' high, pileus 2'-3' broad, stem 6"-10" thick.

Grassy ground in open woods. Greenbush. July.

Boletus modestus, Peck.

Pileus firm, often irregular, dry, yellowish-brown; tubes nearly plane, attached and subdecurrent, pale ochraceous, angular and compound; stem

equal, brown, reticulated with darker lines; spores $.0004' \times .0002'$; flesh gray or pinkish gray.

Plant 2' high, pileus 2' broad, stem 2"-4" thick.

Grassy ground in open woods. Greenbush. August.

***Boletus pallidus*, Frost.**

Pileus soft, viscid when moist, smooth, pale alutaceous; tubes plane, attached to or sometimes slightly depressed around the stem, small, subangular, pale yellow, slightly changing color when wounded; stem subequal, smooth, solid, pallid; spores $.00045' \times .00022'$.

Plant 2'-5' high, pileus 2'-4' broad, stem 4"-6" thick.

Ground in woods. North Greenbush. August.

***Boletus ampliporus*, Peck.**

Pileus broadly convex or expanded, sometimes slightly umbonate, dry, squamulose-tomentose, pinkish-brown; tubes convex, attached or slightly decurrent, very large, angular, compound, yellow; stem equal, solid, yellowish-brown, paler at the top, and marked by the decurrent walls of the tubes; flesh whitish tinged with yellow, unchangeable; spores pale ochraceous, with a greenish tinge, $.00035' \times .00016'$.

Plant 3'-5' high, pileus 3'-4' broad, stem 3"-6" thick.

Low mossy ground in woods. North Elba and Sandlake. August and September.

***Polyporus caeruleoporus*, Peck.**

Pileus fleshy, broadly convex, submentose, moist or hygrophane, brown; pores short, angular, decurrent, grayish-blue; stem central or eccentric, solid, colored like the pileus, sometimes tinged with the color of the pores; flesh white.

Plant gregarious or subcaespitose, 2' high, pileus 1'-2' broad, stem 2"-3" thick.

Shaded banks. Copake. October.

This and the three following species belong to the section *Mesopus*.

***Polyporus griseus*, Peck.**

Pileus fleshy, firm, convex, often irregular, smooth or with a minute appressed silkiness, dry, gray; pores small, short, unequal, subangular, pallid, the mouths white; stem central, thick, short, concolorous; flesh pinkish-gray.

Plant 2'-3' high, pileus 3'-5' broad, stem 6"-10" thick.

Shaded banks. Copake. October.

***Polyporus flavidus*, Peck.**

Pileus fleshy, tough, depressed or funnel-form, smooth, rarely a little villous on the disk, zonate, yellow with darker bands, the margin sometimes lobed or wavy; pores short, minute, angular, yellow; stem central, solid, slightly tapering downwards, smooth, subconcolorous.

Plant 3'-5' high, pileus 2'-4' broad, stem 3"-4" thick.

Ground in woods. Worcester. July.

***Polyporus splendens*, Peck.**

Pileus thin, coriaceous, expanded, subumbilicate, slightly zonate, silky, shining, dark ferruginous when moist, tawny ferruginous when dry, the margin deeply fimbriate; pores small, angular, short, subconcolorous; stem slender, equal, tomentose, concolorous.

Plant 1' high, pileus 6"-10" broad, stem .5"-1" thick.

Much decayed stumps. Center. August.

***Polyporus attenuatus*, Peck.**

Resupinate, effused, very thin, separable from the matrix, pinkish-ochre, the margin whitish; pores minute, subrotund, with thin acute dissepiments.

Prostrate trunks of deciduous trees. Croghan. September.

The pores are scarcely visible to the naked eye.

***Craterellus caespitosus*, Peck.**

Pileus fleshy, tough, irregular, expanded, centrally depressed or funnel-form, smooth, moist, variable in color, greenish-yellow, pinkish-brown, or blackish; the margin sometimes decurved and lobed; hymenium at first smooth, then rugose-wrinkled, the folds decurrent on the short, solid, tough stem which is either central or eccentric; spores oblong, obtuse, sometimes slightly curved, .00035'-.00045' long.

Plant caespitose, 6"-12" high, pileus 6"-10" broad.

Decaying wood in swamps. Portville. September.

The pilei sometimes grow together, forming an intricate irregular tuft.

***Grandinia coriaria*, Peck.**

Effused, membranaceous-tomentose, separable from the matrix, under side and margin tawny-yellow, upper side and minute crowded granules greenish or dingy olivaceous; spores globose, rough, .0003' in diameter.

Forming patches 1'-3' in diameter on old scraps of leather in damp places. Greenbush. August.

Thelephora Willeyi, *Clinton*.

Pileus funnel-formed, thin, smooth, obscurely zoned, white, the margin entire or laciniately toothed and lobed; hymenium smooth, concolorous; stem central, equal, solid, white.

Plant 1'-1.5' high, pileus 6"-12" broad, stem .5"-1" thick.

Ground in woods. Buffalo, *G. W. Clinton*. Lowville. September.

Sometimes the pileus is split on one side down to the stem.

Stereum radiatum, *Peck*.

Resupinate or slightly reflexed, suborbicular, blackish-brown; hymenium uneven, marked with thick corrugations or ridges radiating from the center, cinnamon color.

Old hemlock logs. Catskill Mountains. June.

Corticium bicolor, *Peck*.

Thin, membranaceous, resupinate, flaccid, smooth, separable from the matrix, under surface greenish-yellow, upper surface white.

Rotten wood. Center. October.

Clavaria pusilla, *Peck*.

Stem slender, solid, rather tough, much and irregularly branched; branches unequal, divergent, tips acute.

Plant scarcely 1' high, yellowish.

Ground under spruce and balsam trees. North Elba. September.

Clavaria clavata, *Peck*.

Simple, straight, clavate, obtuse, smooth, not hollow, yellow when fresh, rugose-wrinkled and orange colored when dry.

Plant 4"-6" high.

Damp shaded banks by roadsides. Sandlake. June.

The surface of the ground where it grows is covered by a stratum of green confervoid filaments. The species is related to *C. mucida*.

Tremella colorata, *Peck*.

Plant gregarious, swollen subglobose or irregular soft pulpy and raisin-colored when moist, externally black and internally brownish-pink when dry; filaments colored in the mass; spores globose, colored like the hymenium when mature, .0005'-.0007' in diameter.

Bark of dead ash trees. Tyre. September.

Exobasidium Azaleae, Peck.

Gall subglobose, often lobed or irregular, succulent, fleshy, solid, smooth pale green or glaucous becoming pruinose; spores oblong, straight or curved, obscurely uniseptate, white, .0006'-.0008' long.

Terminal on living twigs of the pinxter plant, *Azalea nudiflora*, transforming the flower buds.

North Greenbush and New Scotland. May and June.

Exobasidium Andromedae, Peck.

Gall flattened or somewhat cup shaped, more or less lobed, smooth, pale green or green varied with red, becoming paler and pruinose with age, hollow, the cavity containing shreds of loose soft cottony filaments; spores narrow, oblong, simple, often curved near one end, white, .0007'-.0009' long.

Lateral on living branches of *Andromeda ligustrina*, transforming the leaf buds.

Center. May and June.

GASTEROMYCETES.**Lycoperdon pedicellatum, Peck.**

Subpyriform, whitish, the outer peridium persistent, forming dense angular spinose processes which are smaller toward the base of the plant; capillitium and spores greenish ochre or dingy olivaceous; spores smooth, pedicellate, globose, .00016'-.00018' in diameter, the pedicel three to five times as long.

Ground and rotten wood. Croghan and Center. September and October.

The spores resemble those of species of *Bovista*.

Diderma crustaceum, Peck.

Effused or circumambient, crowded, sessile, subglobose, smooth, white, outer peridium crustaceous, resembling the shell of some small egg, the inner delicate, appearing cinereous to the naked eye, iridescent under the microscope; columella none; spores globose, black, .0005' in diameter.

Diderma farinaceum, Peck.

Effused or circumambient, crowded, sessile, subglobose, plumbeous when moist or young, white rugulose and farinaceous when dry; spores globose, brown, black in the mass, .0004' in diameter.

Incrusting mosses and fern stems in low woods. Croghan. September.

***Diderma Mariae-Wilsoni*, Clinton.**

Scattered or crowded, sessile, subglobose, smooth, white or pinkish-white, outer peridium crustaceous, within at the base brownish-pink, inner peridium delicate; columella subglobose, rugulose, slightly colored; spores globose, blackish-brown, .0004' in diameter.

Fallen leaves, sticks, moss, etc. Buffalo, *Clinton*. Memphis, Center and Sandlake. August and October.

***Didymium connatum*, Peck.**

Peridium depressed or subglobose, cinereous, furfuraceous, stipitate; stems mostly connate at the base, tapering upward, longitudinally wrinkled, whitish or cream color; spores subglobose, black, .0004' in diameter.

Decaying fungi. Portville. September.

The subfasciculate mode of growth is a marked feature in this species.

***Physarum pulcherripes*, Peck.**

Peridium globose, variable in color, ochraceous, gray, brown or black; stem slender, equal or slightly tapering upwards, vermilion; spores globose, brown, .00033' in diameter.

Rotten wood. Richmondville and Worcester. July.

The bright color of the stem is quite conspicuous, notwithstanding the small size of the plant.

***Physarum caespitosum*, Peck.**

Peridia aggregated in tufts or clusters, crowded, sessile, smooth, brown or blackish-brown; spores dingy ochre, smooth, globose, .00025' in diameter.

Decaying wood. Greenbush. August.

***Craterium obovatum*, Peck.**

Peridium obovate, rugose-wrinkled, glabrous, lilac-brown; flocci whitish; stem colored like the peridium; spores smooth, globose, black, .0005'-.0006' in diameter.

Decaying wood and fallen leaves. Center, Sandlake and Croghan. August and September.

The operculum is not always distinct, the peridium appearing frequently to be irregularly ruptured at the apex.

***Stemonitis herbatica*, Peck.**

Densely fasciculate; capillitium slender, cylindrical, brown when moist, ferruginous-brown when dry; stem black, arising from a membranaceous

hypothallus, penetrating to the apex of the capillitium; spores globose, .0003'-.00035' in diameter.

Plant 2"-3" high, growing on living leaves of grass and herbs. Albany. June.

The color of this plant is almost the same as that of *S. ferruginea*, but the spores are much larger, surpassing even those of *S. fusca*.

***Trichia reniformis*, Peck.**

Peridia gregarious or clustered, sessile, subglobose or reniform, small, brown; flocci few, short, sparingly branched; spores globose, minutely echinulate, yellow-ochre, sometimes tinged with green, .0005' in diameter.

Dead bark of striped maple, *Acer Pennsylvanicum*. Portville. September.

The branches and apices of the flocci are sometimes without spiral markings and slightly nodulose.

***Perichaena flavida*, Peck.**

Yellow throughout; peridia crowded, clustered, sessile, variable in size and shape, shining; flocci few, short, subnodulose, obtuse, sparingly branched; spores globose, echinulate, .00045' in diameter.

Mosses. Sandlake. August.

The bright golden yellow color renders the clusters conspicuous.

CONIOMYCETES.

***Dinemasporium Robiniae*, Gerard.**

Perithecia cup-shaped, bristly, black; spores hyaline, .0002' long, the terminal bristles about as long as the spore.

Dead wood of locust trees. Poughkeepsie, Gerard.

***Dinemasporium acerinum*, Peck.**

Perithecia small, pezizoid, black, hispid with short straight scattered black hairs; spores unequally elliptical, .0003' long, the terminal bristles scarcely one-third the length of the spore.

Dry maple wood. Buffalo, Clinton. April.

***Puccinia pulchella*, Peck.**

Spots yellow or greenish-yellow, orbicular, rarely confluent; sori small, circinating, sometimes confluent, blackish-brown; spores .001'-.0013' long, .0006' broad.

Upper surface of leaves of *Ribes prostratum*. North Elba. July.

***Puccinia Cryptotaeniae*, Peck.**

Spots small, pallid or yellowish, sometimes tinged with purple, dotted by the sori, occasionally confluent; sori minute, clustered, at first covered by the epidermis, then surrounded by its pale ruptured remains which continue in the form of a small pustule with a contracted subcircular opening at the apex, reddish-brown; spores subelliptical, scarcely constricted, crowned with a hyaline pustule, .0011'-.0016' long, .0006' broad.

Under surface of leaves of *Cryptotaenia Canadensis*. North Greenbush. June.

***Puccinia Mariae-Wilsoni*, Clinton.**

Amphigenous; spots none; sori scattered or clustered, unequal, reddish-brown; spores subelliptical, scarcely constricted, crowned with a pustule, .0013'-.0018' long, .0007'-.0008' broad.

Leaves and stems of *Claytonia Caroliniana*. Buffalo, Clinton. Knowersville. May.

***Puccinia Lobeliae*, Gerard.**

Sori minute, scattered or confluent, tawny-brown; spores oblong-elliptical, slightly constricted at the septum and easily separating into two parts, pale, .0013'-.0016' long; pedicel short or obsolete.

Under surface of leaves of *Lobelia syphilitica*. Poughkeepsie, Gerard. The fragile spores are peculiar.

***Puccinia obtecta*, Peck.**

Cauline; sori unequal, often very large, angular or orbicular, scattered or confluent, slightly elevated, long covered by the epidermis, black; spores oblong or oblong-clavate, sometimes curved, constricted, obtuse or obtusely pointed, .0018'-.0024' long, .0008' broad; pedicel colored, seldom half as long as the spore.

Stems of *Scirpus validus* and *S. pungens*. Watkins, Montezuma Marshes and Albany. September and October.

***Puccinia linearis*, Peck.**

Amphigenous; sori very narrow, deep seated, oblong or linear, parallel, crowded, long covered by the epidermis, black; spores oblong, slightly tapering toward the base, not constricted, very obtuse or truncate, .0018'-.0024' long, .0006' broad; pedicel colored, very short.

Leaves and sheaths of grasses. Watkins. September.

This is related to *Puccinia coronata*, but it is without the apical teeth of the spore.

***Puccinia angustata*, Peck.**

Hypogenous; spots pallid or none; sori oblong or linear, sometimes regularly arranged at equal intervals in long parallel lines, narrow, black; spores narrow, oblong-clavate or elongated, septate above the middle, strongly constricted, having the lower cell more narrow than the upper, and cylindrical or slightly tapering downwards, .00018'-.0024' long, .0006' broad; pedicel colored, thick, very short.

Leaves of *Scirpus sylvaticum* and *S. Eriophorum*. West Albany and Watkins. September.

***Protomyces Erythronii*, Peck.**

Spots stained with red or purple; spores growing in the tissues of the leaf, scattered or crowded, most often arranged in short series and erumpent through narrow chinks in the epidermis, large, globose, at length black, .002'-.0026' in diameter.

Leaves and petioles of *Erythronium Americanum*. Greenbush. May.

The leaf is most frequently affected at the base of the lamina or blade.

***Ustilago Erythronii*, Clinton.**

Produced on the leaves in oblong or irregular vesicular patches, half an inch or more in length; spores globose, rough, rather large, .0006'-.00075' in diameter, black in the mass.

Leaves of *Erythronium Americanum*. Goat Island, Clinton.

***Uredo Ledicola*, Peck.**

Spots small, definite, rarely confluent, suborbicular, reddish-brown, sometimes with a darker border; sori subrotund or irregular, surrounded by the ruptured epidermis; spores subglobose, rough, .0012' in diameter, orange, with a thick hyaline epispore.

Upper surface of leaves of *Ledum latifolium*. Mt. Marcy. July.

Apparently quite distinct from *U. Ledi* A. & S.

Peridermium Cerebrum, Peck.

Peridia large, convex, erumpent, irregularly confluent, forming brain-like convolutions, white, rupturing irregularly, the cells radiate-striate on the margin; spores ovate-elliptical or subglobose, rough, yellow, .0008'-.0011' long.

Trunks and branches of young pine trees, *Pinus rigida*, forming excrecences half an inch to two inches in diameter. Center, J. A. Lintner.

Roestelia aurantiaca, Peck.

Peridia cylindrical, fragile, soon lacerated, fugacious, white; spores subglobose, bright orange, about .001' in diameter, with a thick hyaline epispore.

Unripe fruit of *Amelanchier Canadensis*. New Baltimore, J. L. Zabriskie. Keene. July. Also on the unripe fruit of *Crataegus*. Buffalo, Clinton.

The color of the spores will enable this species to be easily recognized.

Aecidium Gerardiae, Peck.

Spots small, suborbicular, scattered, yellowish-green, peridia usually few, small, short, the mouth fringed with spreading or recurved teeth; spores orange, .0008' in diameter.

Leaves of *Gerardia quercifolia*. Highlands near Cold Spring. June.

Aecidium album, Clinton.

Spots none; peridia scattered, short, white, the margin subentire; spores subglobose, white, about .0008' in diameter.

Under surface of leaves of *Vicia Americana*. Buffalo, Clinton.

Aecidium Lycopi, Gerard.

Spots yellow; subiculum more or less thickened; peridia short, scattered or crowded, margin crenate; spores pale yellow.

Leaves, stems and petioles of *Lycopus Europaeus*. Poughkeepsie, Gerard. Buffalo, Clinton. June.

Aecidium Hydrophylli, Peck.

Spots small, few, yellow, with a pale greenish border; subiculum thickened whitish; peridia few, generally crowded, short, the margin subcrenate; spores bright yellow or orange; spermatogonia central, on the opposite side.

Under surface of leaves of *Hydrophyllum Canadense*. Catskill Mountains. June.

HYPHOMYCETES.

***Stilbum ramosum*, Peck.**

Head subglobose, whitish or pale yellow; stem thick, smooth, branched, white above, pallid or brownish below, sometimes creeping and sending up branches at intervals; spores minute, oblong.

Dead larvae of insects buried in rotten wood. Sterling. September.

***Periconia Azaleae*, Peck.**

Plant small, .03'-.04' high, black; stem slightly tapering upward; head globose; spores subglobose or elliptical, colored, .0002'-.0003' long.

Twigs, capsules and old galls of *Azalea nudiflora*. New Scotland. June.

***Macrosporium Chartarum*, Peck.**

Flocci long, jointed, flexuous, branched, colored; branches widely spreading, somewhat nodulose; spores variable, subglobose, elliptical, obovate or pyriform, black, shining, one to three septate, with one or two longitudinal septa, .0006'-.001' long.

Damp paste board. Albany. November. It forms indefinite black spots or patches.

***Clasterisporium pedunculatum*, Peck.**

Flocci erect, opaque, septate; spores terminal, nearly straight, multiseptate, colored, mostly subfusiform or lanceolate, about .003' long, the terminal cell hyaline.

Cut surface of wood. Savannah. October.

***Streptothrix abietina*, Peck.**

Tufts pulvinate, scattered or crowded, blackish-brown; flocci branched, pale, echinulate; spores globose, minutely rough, .00025'-.0003' in diameter.

Bark of prostrate trunks of spruce trees. Sandlake. September.

The larger rough spores and echinulate threads separate this species from *S. atra*.

***Aspergillus fuliginosus*, Peck.**

Creeping flocci white, septate; fertile flocci erect, not septate, crowned with a globose head which is rough with projecting processes; spores globose, sooty-black, smooth, .00016' in diameter.

Rice paste and apple. Albany.

ASCOMYCETES.

Microsphaera Russellii, Clinton.

Amphigenous; mycelium arachnoid, evanescent; appendages 8-18, very long, flexuous, colored, paler toward the tips which are simple or one to three times divided; sporangia ovate, 4-8; spores 4, elliptical, .0007'-.0008' long.

Leaves and petioles of *Oxalis stricta*. Buffalo, Clinton. Poughkeepsie, Gerard. North Greenbush. October.

This might with almost equal propriety be referred to the genus *Erysiphe*.

Erysiphe Euphorbiae, Peck.

Mycelium thin; conceptacles small, .0035' in diameter; appendages few, long, flexuous, colored; sporangia broadly ovate, 3-4; spores 3-4, large, .001' x .00065'.

Leaves of *Euphorbia hypericifolia*. Greenbush. October.

Geoglossum simile, Peck.

Plant 1'-2' high, black, minutely hairy; club obtuse, generally compressed, sometimes with a broad shallow groove on one side, tapering into the stem; asci broad; spores fasciculate, elongate, slightly curved, seven-septate, colored, .003'-~~.004~~.004' long; paraphyses slightly thickened at the tips, septate, sometimes branched.

Damp mossy ground in swamps and in peat bogs. Ft. Edward, Howe. Sandlake. September.

It is scarcely possible to separate this species from *G. hirsutum* without microscopical examination.

Vibrissea lutea, Peck.

Plant 6"-12" high, yellow, receptacle subglobose, smooth, the margin slightly lobed, inflexed, free; stem nearly equal, solid, a little more highly colored than the receptacle, longitudinally wrinkled when dry; asci clavate or cylindrical; spores long, filiform.

Prostrate mossy trunks of trees and among fallen leaves in woods. North Elba. August.

Peziza Solenia, Peck.

Cups minute, nearly cylindrical, hairy, brown, opening by a contracted, white-margined mouth; spores oblong, crowded or biseriate, uniseptate usually with four nuclei, subhyaline, .0005' long; paraphyses filiform.

Dead stems of *Eupatorium ageratoides* in damp shaded places. Watkins Glen. September.

The cups are a little longer than broad, and appear like some minute *Solenia*.

Rhytisma linearis, Peck.

Linear, here and there interrupted or constricted, black ; asci broad, clavate, eight-spored ; spores very long, obtuse, strongly narrowed in the middle, involved in mucus, .002'-.003' long.

Under surface of leaves of pine trees, *Pinus Strobus*. Guilderland, Greenbush and Sandlake. June.

It forms a thick black line on the under surface of the leaf, often extending the entire length. The spores appear to consist of two oblong parts connected by a narrow neck.

Hypomyces polyporinus, Peck.

Perithecia minute, ovate or subconical, seated on a pallid subiculum, smooth, yellowish, or pale amber ; asci narrow, linear ; spores fusiform, acuminate at each end, nucleate, .0006'-.0007' long.

On *Polyporus versicolor*. Worcester and Croghan. July and September.

Nectria Apocyni, Peck.

Conidia. Subhemispherical or irregular, small, pale red ; spores fusiform, straight, .0005'-.0006' long.

Ascophore. Caespitose or scattered, dull red, perithecia minute, pale ochraceous, and subglobose when moist, dull red collapsed or laterally compressed and rough with minute whitish scales when dry ; spores biserial, uniseptate, fusiform, nucleate, .00065'-.0008' long.

Base of dead stems of *Apocynum cannabinum*. North Greenbush. October.

Nectria mycetophila, Peck.

Perithecia crowded or scattered, minute, smooth, subglobose, pale yellow when young, then pinkish-ochre ; ostiole minute, papillate, distinct, darker colored ; asci subclavate ; spores oblong, simple, .0005' $\times$.00016'.

Decaying fungi. New Scotland. October.

Sphaeria Staphylina, Peck.

Perithecia minute, black, covered by the epidermis, which at length ruptures in a stellate manner or irregularly; spores biseriate, colorless, constricted in the middle, three to five septate, .0009'-.001' long, the two parts formed by the central septum unequal in diameter.

Dead twigs of *Staphylea trifolia*. Helderberg Mountains. May.

Sphaeria Desmodii, Peck.

Perithecia scattered or seriatly placed, minute, covered by the epidermis, which is pierced by the acute ostiole, black; asci clavate; spores biseriate, fusiform, colorless, four nucleate, .00035'-.0004' long.

Dead stems of *Desmodium*. Garrisons. June.

occupying the whole lower surface of the leaf, ferruginous-brown; spores ovate or oblong-ovate, umbonate at the apex, .001-.0013 inch long; pedicel hyaline, short.

Living leaves of *Sophora sericea*, Pursh.

Mr. Brandegee remarks concerning this fungus that it is very common at Canon City, about one-half the *Sophora* plants being affected by it. The plants attacked by it become more erect in habit and do not blossom. The sori sometimes occur sparingly on the upper surface of the leaf also. The specific name is suggested by the hyaline membrane that at first covers them.

PUCCINIA NIGRESCENS, *Pk*.—Spots none; sori rather prominent, cauline, oblong, blackish-brown or black; spores smooth, elliptical or oblong, obtuse or bluntly pointed, slightly constricted at the septum, .0015-.0018 inch long, .0008-.001 broad; pedicel colorless, two to four times as long as the spore.

Stems and branches of *Salvia lanceolata*, Willd.

No leaves of the host plant accompanied the specimens sent, so that I am not able to say whether the fungus ever attacks the leaves or not.

The two following species have been received from the sources given at the end of the descriptions:

DIATRYPELLA FROSTII, *Pk*.—Stroma verruciform, rather prominent, convex or hemispherical, scarcely one line broad, sometimes confluent, partly covered by the closely adhering fragments of the ruptured epidermis, black externally, greenish within, forming a white spot on the wood beneath; perithecia sub-globose, generally ten to fifteen; ostiola obscure; asci sub-cylindrical, scarcely pedicellate, polysporous; spores yellowish in the mass, cylindrical, curved, .00025-.00035 inch long.

Dead maple branches. Brattleborough, Vermont. *C. C. Frost*.

Externally this fungus resembles small forms of *Diatrypella verruciformis*, from which it is easily separated by the greenish color of the stroma. *D. betulina*, which also has a greenish stroma, differs from this species in the larger size, different form and naked disk.

SOROSPORIUM DESMODII, *Pk*.—Spore balls irregular or sub-globose, compact, rough, .0012-.0016 inch in diameter, color in the mass reddish-brown inclining to cinnamon; spores sub-globose, rough, .0006-.0007 inch in diameter, four to ten in a group, not easily separating from each other.

Seeds of *Desmodium acuminatum*, DC. Closter, New Jersey. *C. F. Austin*. September.

I am not aware that any representative of this genus has before been detected in the United States. Nor is it surprising that this species has so long escaped notice, for the fungus is wholly concealed, being produced in the seeds while they are yet inclosed in the pod. For a time there is no external evidence of the presence of the fungus, but at length a slight discoloration of the part of the pod immediately over the affected seeds indicates that all is not right within.

The whole seed, except perhaps the thin external coat, appears to be transformed into the fungus spores. The color of these is similar to that of the spores of *S. Saponaria*, but the spore masses are smaller and less easily separated into their component spores. Every seed in a pod and nearly every pod on an affected plant, so far as shown by the specimens sent me, is attacked. From this it would appear that the fungus must be quite effective in preventing the increase of this *Desmodium* by the seed in localities where it abounds.

MORE ABOUT LOBELIAS.—In the April number (1877) of the GAZETTE I mentioned having found a white plant of *Lobelia syphilitica*. I now have to report not only a red flowered plant of the same species, (differing in no particular, except in color, from the common form,) but a remarkable plant which appears to be a hybrid between *L. syphi-*

COLORADO FUNGI. (NEW SPECIES.)—By CHAS. H. PECK.—The species of fungi here described were collected in Colorado by Mr. T. S. Brandegee, and forwarded to me for examination by Mr. E. S. Rau, of Bethlehem, Pennsylvania.

SPHÆROPSIS RAUI, *Pk.*—Spots indefinite, brown; perithecia numerous, gregarious, minute, phomatoid, sub-membranaceous, black, .006–.007 inch in diameter; spores obovate or oblong, colorless, .00065–.00075 inch long.

Living leaves of *Artemisia scopulorum*, Gray.

LECYPHAEA SPECIOSA, *Pk.*—Spots suborbicular, one to two lines in diameter, yellow; sori few, crowded, prominent, hypophyllous orange-colored; abortive spores elongated, curved, obtuse, nearly cylindrical; spores subglobose or elliptical, .001–.0012 inch long, rough, the endochrome bright orange.

Living leaves of *Rubus deliciosus*, Torr.

This is probably the early or *Uredo* form of some species of *Phragmidium* which should be sought later in the season. It is distinguished from its allies by the deep orange color of the sori, and the narrow elongated abortive spores.

ÆCIDIUM BRANDEGEEI, *Pk.*—Peridia amphigenous, scattered or aggregated, sometimes seriatly placed, elongated; spores subglobose, orange colored, .0008–.001 inch in diameter.

Living stems and leaves of *Asclepias verticillata*, L.

The peridia resemble those of *Æ. Thalictri*, but the spores and habit of the plant are quite different. It occupies the stem and leaves in patches of considerable extent and changes the aspect of the host plant.

ÆCIDIUM HEMISPHERICUM, *Pk.*—Peridia prominent, scattered or crowded, hemispherical, opening by a small aperture; spores orange, subglobose or elliptical, .0008–.001 inch long, the epispore thick, hyaline.

Living stems and leaves of *Mulgedium pulchellum*, Nutt.

The peridia are generally more scattered on the stem than on the leaves, the whole lower surface of which is often occupied by them.

ÆCIDIUM POROSUM, *Pk.*—Spots none; cups crowded, deep-seated, broad, wide-mouthed, occupying the whole lower surface of the leaf to which they give a porous appearance; spores orange-colored, sub-angular, .0008–.001 inch in length.

Living leaves of *Vicia Americana*, Muhl.

This species is quite distinct, both in its spores and in the size and character of the peridia, from *Æ. album*, Clinton, which also inhabits *Vicia Americana*.

ÆCIDIUM ABUNDANS, *Pk.*—Spots yellow, numerous, small, sub-orbicular, sometimes confluent; peridia few, hypophyllous, sub-circumating, pustulate; spores sub-globose, orange, .001 inch in diameter; spermogonia on both sides of the leaf on the same or on different spots.

Living leaves of *Symphoricarpos oreophilus*.

The leaves of the host plant are beautifully mottled by the spots. The peridia in the specimens seen are nearly all closed, from which it is probable that they are scarcely mature.

ÆCIDIUM BIGELOVIAE, *Pk.*—Spots thickened, yellow; peridia pustulate, hemispherical, opening by a small aperture; spores sub-globose, bright yellow or orange, .001–.0013 inch in diameter.

Living stems and leaves of *Bigelovia Douglasii*, Gray.

The peridia resemble those of *Æ. hemisphericum*, but the species is at once distinguished by the yellow spots and larger spores.

UROMYCES HYALINUS, *Pk.*—Spots none; sori numerous, hypophyllous, at first covered by the hyaline membranous epidermis, at length exposed, effused, nearly or quite

[Reprinted from THE BULLETIN OF THE TORREY BOTANICAL CLUB, Vol. 22, No. 5, May, 1895.]

New Species of Fungi.

BY CHAS. H. PECK.

LEPIOTA FULVODISCA. Pileus thin, convex or nearly plane, obtuse or umbonate, viscid when moist, white, with the disk or umbo fulvous or tawny-brown; lamellae narrow, close, free, white; stem slender, flexuous, viscid, hollow, white or whitish, the base abruptly bulbous, the annulus thin, membranous, pure white; spores ovate-elliptical, .0003 to .0004 in. long, .00016 to .0002 in. broad, usu-

ally containing a shining nucleus and furnished with a slight apiculus at one end.

Pileus 1 to 1.5 in. broad; stem 2 to 3 in. long, 1 to 1.5 line thick.

Plant fragile, growing among fallen leaves in woods. Pasadena, California. January. Prof. A. J. McClatchie.

From *L. illinita* Fr. this species is separated by its tawny disk of the pileus, its membranous annulus, the bulbous base of the stem and the different shape of the spores.

CLITOCYBE PUSILLA. Pileus at first hemispherical or convex, then nearly plane, obtuse, sometimes with a very small umbo, dry, pruinose, grayish, flesh thin, whitish; lamellae narrow, close, adnate or decurrent, white; stem short, solid, pruinose, grayish; spores subglobose or very broadly elliptical, .00016 to .0002 in. long.

Pileus 3 to 8 lines broad; stem about 6 lines long, scarcely 1 line thick.

Densely gregarious or subcaespitose. On manure. Pasadena. February. McClatchie.

Apparently belonging to the tribe Disciformes and related to *C. Bresadolae* Schulz, but an exceedingly small plant for that tribe.

COLLYBIA ALBOGRISEA. Pileus fleshy, thin, convex or nearly plane, often somewhat irregular on the margin, glabrous, whitish or grayish, flesh white; lamellae broad, distant, adnate, white or whitish, the interspaces often venose; stem nearly equal, hollow, sometimes twisted, whitish or grayish, the lower part covered with a dense whitish, grayish or yellowish tomentum, the upper part naked or merely white-pruinose; spores broadly elliptical, .0002 to .00024 in. long, .00016 broad, generally containing a single shining nucleus.

Pileus about 1 in. broad; stem 1.5 to 2 in. long, 1 to 2 lines thick.

Plant often caespitose and then the pileus irregular from mutual pressure. Among fallen leaves. Pasadena. January. McClatchie.

The species belongs to the tribe Vestipedes.

MYCENA ELEGANTULA. Pileus membranous, conical or campanulate, sulcate-striate, brown or purplish-brown; lamellae distant, adnate with a decurrent tooth, whitish or pallid with the edge purplish-brown, the interspaces (in the dried specimens) venose or transversely wrinkled; stem slender, hollow, glabrous, sometimes

with a loose grayish fibrillose tomentum at the base ; spores oblong-elliptical, .00035 to .0005 in. long, .00016 to .0002 broad, usually containing one or two small nuclei.

Pileus, 4 to 10 lines broad ; stem 1 to 2 in. long, .5 a line thick.

Gregarious or caespitose. Among fallen leaves under trees. Pasadena. December. McClatchie.

This plant may be distinguished from the closely related *M. atromarginata* Fr. by its smaller size and the purplish tint to the edge of the lamellae, and from *M. purpureofusca* Pk. by its differently shaped, longer spores.

OMPHALIA SEMIVESTITES. Pileus very thin, deeply convex or subcampanulate, glabrous, grayish-brown in the dried state, paler when fresh ; lamellae rather broad, distant, arcuate, decurrent, white or whitish ; stem hollow, white or whitish, the upper half glabrous, the lower half clothed with a white mycelioid tomentum ; spores elliptical, .0002 to .00024 in. long, .00012 to .00016 broad.

Pileus 6 to 12 lines broad ; stem about 1 in. long, 1 to 2 lines thick.

Growing on much decayed wood. Newfoundland. May. Rev. A. C. Waghorne.

This plant is apparently related to such species as *O. setipes* Fr. and *O. grisea* Fr. and belongs to the tribe Mycenariae. It is easily recognized by the peculiar character of the stem. In the dried examples the upper glabrous part of the stem is shriveled and longitudinally striate, but the lower tomentose part is plump and even as in the fresh plant.

HYGROPHORUS ELEGANTULUS. Pileus convex or nearly plane, glabrous, viscid, grayish-yellow or slightly tawny, flesh white ; lamellae distant, slightly decurrent, white ; stem equal, solid, slightly floccose-squamulose at the top, elsewhere glabrous, glutinous, white or whitish, sometimes abruptly pointed at the base ; spores elliptical, .0004 in. long, .0002 to .00024 broad.

Pileus 1 to 2 in. broad ; stem 2 to 3 in. long, 3 to 4 lines thick.

Woods. Maryland. November. T. Taylor.

The species belongs to the tribe Limacium, and is related to *H. discoideus* Fr., from which it may be separated by its solid stem and larger spores.

ENTOLOMA FERRUGINANS. Pileus fleshy, convex, obtuse or umbonate, often irregular, hygrophanous, glabrous, shining, dark fuliginous or broccoli-brown, flesh whitish, fibrous and colored at

the surface; lamellae 4 to 7 lines broad, adnexed, easily splitting transversely, grayish-salmon, becoming clay-color; stem solid, glabrous, yellowish or cream-color, blunt at the base or sometimes attenuated and radicating; spores subglobose, irregular or angular, .0003 to .0004 in. long.

Pileus 2 to 6 in. broad; stems 3 to 4 in. long, 4 to 8 lines or more in thickness.

Under oak trees. Pasadena. February. McClatchie.

This plant is related to *E. rhodopolium*, but is separated from it by its darker color and solid stem. According to Prof. McClatchie, the fresh plants have a strong odor resembling that of ferric chloride, and chemical tests showed the presence of iron. It is pronounced edible by him.

LEPTONIA EDULIS. Pileus thin, convex or centrally depressed, with or without an umbo, velvety, dark-gray; lamellae rather broad, subventricose, adnexed; moderately close, at first whitish or light drab, becoming flesh-color; stem slender, hollow, colored like the pileus, often with an abundant white mycelioid tomentum at the base; spores subglobose, angular, apiculate at one end, .0003 to .0004 in. long, containing a single large nucleus.

Pileus 6 to 18 lines broad; stem 12 to 18 lines long, .5 to 1 line thick.

Among grass and weeds. Pasadena. January.

According to Prof. McClatchie, this plant when fresh has a nutty flavor and is edible. The velvety appearance of the pileus has disappeared from the dried specimens. In some, the margin of the pileus is striate, but in the fresh plant the margin is said to be even.

ECCILIA NIGRICANS. Pileus thin, convex, umbilicate or centrally depressed, subzonate, unpolished, grayish-black; lamellae broad, distant, decurrent, light-drab or brownish, becoming tinged with flesh-color; stem short, hollow, grayish-black, commonly with an abundant white mycelium; spores angular, .0004 in. long, nearly as broad, containing a single large nucleus.

Pileus 6 to 18 lines broad; stem about 1 in. long, .5 to 1 line thick.

Grassy ground. Pasadena. January.

Prof. McClatchie's notes say that this plant has the odor and flavor of butternuts and that it is delicious when cooked. Also that when fresh the pileus is tomentose and the margin even, but these characters are not clearly shown in the dried specimens.

PHOLIOTA ANOMALA. Pileus at first hemispherical or subconical, then convex, glabrous, hygrophalous, broccoli-brown when moist, pale-yellow or cream-color when dry; lamellae adnate or slightly decurrent, subarcuate, pale becoming brownish-ferruginous, often white on the edge; stem cavernous or hollow with irregular transverse partitions, sometimes containing a cottony tomentum, whitish, with a slight evanescent annulus; spores elliptical, .0003 to .0004 in. long, .00016 to .0002 broad.

Pileus 8 to 18 lines broad; stem 1.5 to 2.5 in. long, 1 to 3 lines thick.

Sticks and leaves lying on the ground. Pasadena. January. McClatchie.

The species belongs to the tribe Truncigenae, section Hygrophanae. It is well marked by its fugacious annulus, subdecurrent lamellae and peculiar cavernous stem.

HEBELOMA FOEDATUM. Pileus fleshy, convex becoming plane or centrally depressed, glabrous, very viscid or glutinous, reddish cinnamon, flesh yellowish-white; lamellae subventricose, emarginate with a decurrent tooth, cinnamon-color, becoming mummy-brown; stem solid, equal or slightly thickened at the base, fibrillose, paler than the pileus; spores broadly elliptical, .00024 to .0003 in. long, .00016 to .0002 broad.

Pileus 1.5 to 3 in. broad; stem 1.5 to 2.5 in. long, 2 to 4 lines thick.

Streets of Pasadena. December. McClatchie.

The species is apparently related to *H. firmum*, *H. testaceum* and *H. glutinosum*, from all of which it is separated by its small spores. Its viscid pileus causes dirt to adhere to it in such quantity as to give the plant a very defiled, unattractive appearance.

FLAMMULA ANOMALA. Pileus deeply umbilicate or infundibuliform, often irregular, glabrous, whitish; lamellae narrow, close, decurrent, pale-ferruginous; stem short, irregular, whitish; spores globose, brownish-ferruginous, .00024 in. broad.

Pileus about 1 in. broad; stem 6 to 12 lines long.

Plant commonly caespitose. Ground. Trexlertown, Pennsylvania. October. Dr. William Herbst.

A whitish umbilicate pileus is unusual among species of *Flammula*. This plant appears to belong to the tribe Gymnotae.

TUBARIA PALLESCENS. Pileus fleshy but thin, convex or nearly plane, sometimes slightly depressed in the center, glabrous, hygrophalous, brick-red when moist, yellowish or cream color when

dry; lamellae broad, adnate or slightly decurrent, tawny-buff, becoming brownish-ferruginous; stem slender, hollow, yellowish, with a white mycelium at the base; spores elliptical, .0003 in. long, .00016 broad.

Pileus 5 to 10 lines broad; stem 12 to 18 lines long, .5 to 1 line thick.

Sticks and leaves under trees. Pasadena. January. McClatchie.

When young, slight vestiges of a veil are visible, connecting the incurved margin of the pileus with the stem.

PLUTEOLUS LUTEUS. Pileus thin, at first subovate, then convex or subcampanulate, glabrous, viscid, slightly striate on the margin, yellow; lamellae numerous, close, free or but slightly adnexed, yellowish becoming ferruginous; stem slender, hollow, slightly thickened toward the base, striate at the top and there sprinkled with mealy particles, yellowish; spores elliptical, .0004 to .0005 in. long, .00024 to .0003 broad.

Pileus 6 to 12 lines broad; stem 1.5 to 2.5 in. long, 1 to 2 lines thick. Plant very fragile, gregarious. Under trees. Pasadena. December. McClatchie.

The yellow color and viscid pileus are prominent characters of this species. In some of the dried specimens the lamellae appear free, in others slightly adnexed, but because of the viscid pileus I have referred the plant to the genus *Pluteolus*.

CORTINARIUS VIRGATUS. Pileus thick, fleshy, hemispherical or convex, obtuse or subumbonate, slightly viscid, ochraceous tinged with olive-buff, conspicuously virgate with reddish fibrils, flesh dingy-white; lamellae subdistant, adnexed, at first subcinnamon, then ochraceous-russet; stem short, stout, solid, enlarged and fibrillose at the base, pale-ochraceous; spores subglobose or broadly elliptical, .00024 to .0003 in. long, .0002 to .00024 broad.

Pileus 3 to 4 in. broad; stems about 2 in. long, 8 to 12 lines thick.

Under oak trees. Pasadena. February. McClatchie.

This species is well marked by its stout habit and by the reddish fibrils of the pileus.

AGARICUS CALIFORNICUS. Pileus at first subconical, becoming convex, minutely silky or fibrillose, whitish, tinged with purple or brownish-purple on the disk, flesh whitish; lamellae close, free, pink becoming purplish, then blackish-brown; stem rather long, solid or stuffed, equal or tapering upward, distinctly and rather abruptly narrowed above the entire externally silky annulus, pallid

or brownish; spores broadly elliptical, .0002 to .00025 in. long, .00016 to .0002 broad.

Pileus 1 to 3 in. broad; stem 1.5 to 3 in. long, 2 to 4 lines thick.

Under oak trees. Pasadena. January. McClatchie.

This fungus is similar in size, shape and habitat to *A. hemorrhoidarius*, but it is unlike that species in color, in the adornment of the pileus and in its color not changing where bruised or broken.

STROPHARIA BILAMELLATA. Pileus fleshy, convex, even, whitish or yellowish, flesh pure white; lamellae close, adnate, purplish-brown when mature; stem short, solid, white, with a well-developed pure white annulus which is striately lamellate on the upper surface; spores elliptical, purplish brown, .0004 in. long, .0002 to .00024 broad.

Pileus 1 to 2 in. broad; stem about 1 in. long, 3 to 4 lines thick.

Streets of Pasadena. January. McClatchie.

This fungus is remarkable for the lamellated upper surface of the rather thick membranous annulus. These lamellae are uneven on the edge and in some cases they appear to extend upward on the stem till they meet the true lamellae. The plant is said by its discoverer to be edible. The color of the young lamellae is not shown by the examples.

HYPHOLOMA LONGIPES. Pileus thin, campanulate, even or obscurely striate on the margin, fibrillose becoming glabrous, hygrophanous, yellowish-brown when moist, brown or isabelline-brown when dry, the margin appendiculate with the very white floccose fugacious veil; lamellae narrow, close, adnate, white or whitish, becoming nearly black, often whitish on the edge; stem slender, long, hollow, striate at the top, white, with a white mycelioid tomentum at the base; spores elliptical, .0005 in. long, .0003 broad.

Pileus 1 to 1.5 in. broad; stem 2 to 5 in. long, 1 to 2.5 lines thick.

Plant fragile, growing among fallen leaves in very wet weather. Pasadena. September. McClatchie.

The disk of the pileus is so thin and the stem so completely hollow to the apex that in the dried specimens there is a depression or umbilicus in the center of the pileus.

PANAEOLUS INTERMEDIUS. Pileus campanulate or convex, even, glabrous, moist or hygrophanous, grayish-brown; lamellae ascending or subarcuate, subdistant, adnate, black when mature; stem slender, often elongated, hollow, grayish-brown, white-pruinose at the top; spores oblong-elliptical, .0005 to .0006 in. long, .00025 to .0003 broad.

Pileus 6 to 12 lines broad; stem 2 to 4 in. long, .5 to 1 line thick.

Rich soil along gutters or in cañons. Pasadena. January. McClatchie.

The margin of the pileus does not extend beyond the lamellae, and this character with the slender stem suggests the genus *Psathyrella*, but because of the absence of striae on the pileus it seems best to refer the plant to the genus *Panaeolus*.

PANAEOLUS DIGRESSUS. Pileus hemispherical or convex, glabrous, bay-red; lamellae very broad, plane, distant, adnate, purplish black with a white edge; stem short, floccose-fibrillose toward the base, striate at the apex, hollow, a little paler than the pileus; spores broadly elliptical, .0005 to .0006 in long, .00035 to .0004 broad.

Pileus 4 to 6 lines broad; stem about 1 in. long, 1 line thick. On dung. Pasadena. July. McClatchie.

This plant also diverges from the generic character in its lamellae extending quite to the margin of the pileus, and in its unpolished stem.

COPRINUS CALYPTRATUS. Pileus when mature adorned with a few grayish floccose scales and crowned with a persistent stellately split membranous dingy-yellow or subtawny calyptra, radiate striate to the disk, grayish-flocculent along the ridges of the striae, blackish; lamellae free, dark lead color becoming black; stem equal, hollow, white, becoming blackish in drying except at the base, neither annulate nor distinctly volvate; spores elliptical, black, .0006 to .0008 in. long, .00045 to .0005 broad.

Pileus about 2 in. broad; stem 3 to 4 in. long, 2 to 3 lines thick. Open cultivated ground. Rockport, Kansas. August. E. Bartholomew.

This species is well marked by the persistent membranous calyptra that adheres to the summit of the pileus. Its margin is split into four to six broad rays. The change of color in the stem is similar to that ascribed to the stem of *C. sterquilinus* Fr., but our plant differs from that in its calyptra and in the absence of an an-

nulus and volva at the base of the stem. Only mature specimens were seen, consequently the characters of the young plant remain unknown and the description to that extent is defective.

COPRINUS JONESII. Pileus submembranous, campanulate becoming broadly convex or expanded and split or revolute on the margin, very blunt or truncate at the apex when young, everywhere covered with tawny-gray or pale-cervine floccose scales which wholly or partly disappear with age revealing the striate surface beneath; lamellae crowded, linear, free, at first white or whitish, becoming black; stem equal or slightly tapering upward, minutely floccose, hollow, white; spores black, broadly elliptical, .0003 to .00035 in. long, .00025 broad, with an apiculus at one end.

Pileus 1 to 2 in. broad; stem 2 to 3 in. long, 2 to 3 lines thick.

Plant fragile, sometimes caespitose. In a cellar. Vermont. April. Prof. L. R. Jones.

The species is closely related to *C. finetarius*, of which it might easily be considered a variety, but it is easily distinguished by the truncate apex of the young pileus, the differently colored scales and the smaller spores. *C. soboliferus* Fr. has the pileus truncated at the apex, but it is a very different species.

COPRINUS APICULATUS. Pileus membranous, campanulate or deeply convex, acute or apiculate, furfuraceous, plicate-striate to the disk, grayish; lamellae few, subdistant, reaching the stem, black; stem filiform, glabrous, white; spores elliptical, black, .0003 in. long, .00016 broad.

Pileus about 3 lines broad; stem 1 to 1.5 in. long, scarcely half a line thick. Lewiston, Pennsylvania. Mrs. E. B. Noyes.

BOLETINUS BOREALIS. Pileus fleshy, convex, obtuse or subumbonate, brownish yellow, obscurely and somewhat reticulately streaked with reddish-brown lines; pores large, angular, unequal, slightly decurrent, brownish-yellow; stem short, equal or slightly tapering upward, brownish-yellow with a whitish mycelioid tomentum at the base; spores oblong, .0004 to .0005 in. long .00016 to .0002 broad.

Pileus 1 to 2 in. broad; stem about 1 in. long.

Sandy soil. Capstan Island, Labrador. October. Waghorne.

The markings of the pileus appear as if due to the drying of a glutinous substance. The radiating lamellae and the transverse partitions of the interspaces are very plainly shown. Described from two dried specimens.

BOLETUS INFLEXUS. Pileus convex, glabrous, viscid, yellow, often red or reddish on the disk, the margin thin, inflexed, concealing the marginal tubes, flesh whitish, not changing color where wounded; tubes rather long, adnate, yellowish, becoming dingy-yellow with age, the mouths small, dotted with reddish glandules; stem rather slender, exannulate, solid, viscid, dotted with livid-yellow glandules; spores yellowish, .0004 to .0005 in. long, .00016 to .0002 broad.

Pileus about 1 in. broad; stem about 2 in. long, 2 to 4 lines thick.

Open woods. Trexlertown. September. Herbst.

This *Boletus* belongs to the tribe Viscipelles. It is remarkable for and easily recognized by the inflexed margin of the pileus, which imitates to some extent the appendiculate veil of *Boletus versipellis*. It sometimes grows in tufts. The paper in which fresh specimens were wrapped was stained yellow. *Boletus Braunii* Bres. has an inflexed margin, but that is a much larger plant with a yellowish-brown pileus, a fibrillose stem and much smaller spores.

POLYPORUS ANCEPS. Effuso-reflexed or resupinate, inseparable from the matrix, firm, subcorky but flexible, white; pileus narrow, about 6 lines broad, laterally elongated or confluent, minutely downy, sometimes rugosely pitted; pores minute, subrotund, commonly 2 to 3 lines long, the dissepiments obtuse; mycelium white, permeating the bark and wood.

Dead trunk of hemlock, *Tsuga Canadensis*. Stony Brook, Massachusetts. October and November. Prof. E. A. Burt.

The plants are commonly resupinate, but sometimes reflexed, forming a narrow pileus about half an inch broad but extending laterally for several inches. They are suggestive of the first year's growth of *P. connatus* Fr., but they do not revive the next year, and they have a different habitat. Though differing somewhat in texture they are apparently related to such species as *P. semisupinus* and *P. semipileatus*, and with them they serve to connect the genus *Polyporus* with the genus *Poria*.

SPARASSIS HERBSTII.—Plants much branched, forming tufts 4 to 5 in. high and 5 to 6 in. broad, whitish, inclining to creamy-yellow, tough, moist, the branches numerous, thin, flattened, concrescent, dilated above and spatulate or fan-shaped, often somewhat longitudinally curved or wavy, mostly uniformly colored, rarely with a few indistinct, nearly concolorous, transverse zones

near the broad entire apices; spores subglobose or broadly elliptical, .0002 to .00025 in. long, .00016 to .0002 broad.

Trexlerstown. August. Herbst.

The species is evidently closely allied to *S. spathulata* Schw., but differs especially in its paler color, with no rufescent hues, in its much more branching habit and in the absence of any distinct zones.

BATTARREA ATTENUATA. Exoperidium unknown; endoperidium 2 in. or more in breadth, the basal part hard, thick, even and concave beneath, convex above and somewhat coarsely reticulate by the bounding walls of broad shallow pits; stem 8 to 10 in. long, gradually attenuated toward the base, hard, almost woody, solid, rough except at the top with rather coarse spreading or reflexed scales, brown externally, rusty-brown within; spores globose, ferruginous, .0003 in. broad; threads of the capillitium destitute of spiral thickenings.

Plants commonly growing in tufts of 3 to 5 individuals.

Dry sandy soil. Nevada. Collected by C. W. Irish; communicated by Dr. Thomas Taylor.

The single dried specimen from which, with notes kindly communicated by the collector, the above description was derived, was past maturity and destitute of any volva or exoperidium. The upper part of the endoperidium, which is apparently membranous, had nearly all disappeared, and but a mere remnant of the spores and capillitium remained. It is very unsatisfactory to attempt the description of a species from such imperfect data, yet the characters seen are so peculiar and distinct that I have been willing to strain a point in order to make this remarkable plant known. It does not agree rigidly with the characters ascribed to the genus *Battarrea*, differing apparently in the solid stem, the absence of spiral thickenings in the capillitium threads and in the coarsely pitted subreticulate hymenial substratum, so that it might easily be taken as the type of a distinct genus.

The dried specimen has a strong, unpleasant odor, indicating its relationship to the Phalloideae. From the notes of Mr. Irish we learn that the long stem, which is about half an inch thick at the base and one and a half at the top, is almost wholly buried in the soil, and that the plants appear above the surface only in seasons after heavy snow falls, whose gradual melting has moistened the earth deeply.

TYLOSTOMA SEMISULCATUM. Peridium subglobose, usually a little longer than broad, 6 to 8 lines broad, 9 lines long, glabrous above, ferruginous-tomentose on the lower half; ostium entire, stem equal, about 2 in. long, even and glabrous or but slightly furfuraceous on the upper part, the lower part longitudinally sulcate, whitish; spores ferruginous, globose, .00016 to .0002 in. broad; threads of the capillitium colorless, not septate.

Sandy soil. Nevada. Collected by C. W. Irish; communicated by T. Taylor.

This species is separated from *T. mammosum* Fr. by its peridium, which is tomentose on the lower half and not depressed, and by its stem, which is distinctly furrowed in its lower half.

LYCOPERDON BELLII. Peridium about 2 in. broad, subglobose, sessile, grayish or yellowish-brown, exterior peridium continuous, adorned with numerous persistent hard angular irregular or substellate warts which are smaller toward the base, separable at maturity from the inner peridium and falling away in flakes or patches; the inner peridium thinner, paler, glabrous; capillitium composed of long slender interwoven, slightly colored and occasionally branched filaments, a little broader than the diameter of the spores; spores globose, olive-brown, .00024 to .00028 in. broad.

Rocky ground. Digges Island, Hudson's Bay. August. Collected by R. Bell; communicated by Prof. J. Macoun.

This puff-ball is peculiar in its continuous but warty exterior peridium, which is less friable than that of *L. separans*. The peridium appears to rupture irregularly, as in *Calvatia*. It is also well marked by the dark olive-brown spore mass.

EXCIPULINA OBSCURA. Perithecia cupulate, $\frac{1}{4}$ to $\frac{1}{2}$ line broad, sessile, black, the margin at first incurved, then erect, thin; hymenium subgelatinous; spores subfiliform, curved, plurinucleate or obscurely pluriseptate, hyaline, .0012 to .0018 in. long, .00016 broad, supported on short sporophores.

Bark of hemlock trees, *Tsuga Canadensis*. Newfoundland. Waghorne.

MELASMA IMITANS. Perithecia hypophyllous, membranous, variable in size and shape, commonly subelliptical or oblong, somewhat confluent in nerve-following lines, rugosely uneven, black, opening irregularly; spores subcylindrical, straight or nearly so, colorless, sometimes obscurely plurinucleate, .0008 to .0012 in. long, .00016 to .0002 broad, oozing out and forming pallid globular or irregular masses or short thick tendrils.

Lower surface of living fronds of *Fteris aquilina*. California. Prof. M. A. Howe.

This fungus imitates *Phyllachora pteridis* in habitat and general appearance, and might easily be supposed to be the conidial state of it. The lines of the perithecia are more narrow and less prominent. The masses of discharged spores are rather large and very numerous and partly conceal the black perithecia beneath them.

CAEOMA ABERRANS. Pustules suborbicular, slightly elevated, .5 to 1 line broad, at first covered by a whitish membrane which finally disappears revealing the mass of orange-yellow spores; spores subglobose or elliptical, smooth or nearly so, .0007 to .0009 in. long.

Bark of living alder. Newfoundland. May. Waghorne.

This fungus is peculiar in its habitat. The spores do not easily separate from each other, but possibly this is due in some measure to immaturity.

ASPERGILLUS SUBGRISEUS. Grayish; sterile hyphae creeping; fertile erect, continuous, simple, .0003 in. thick, terminating above in an inflated subglobose vesicle .0012 to .0016 in. broad; sterigmata none or obsolete; spores globose, .00016 in. broad.

On *Corticium amorphum*. Newfoundland. Waghorne.

This species is separated from the related *A. griseus* by its larger spores, continuous fertile hyphae and by the absence of distinct sterigmata.

LEPTOGLOSSUM LATUM. Club 3 to 6 lines long, nearly or quite as broad, soft when moist, rather fragile when dry, compressed and somewhat irregular, black; stem about as long as the club, black; asci clavate-cylindrical, .0045 to .005 in. long, about .0005 broad, 8-spored; spores crowded in the ascus, oblong or cylindrical, straight or slightly curved, continuous, obtuse, colorless, .001 to .0016 in. long, .0002 to .00024 broad; paraphyses colored, thickened at the top and sometimes recurved.

Sandy soil. Labrador. September. Waghorne.

A species very peculiar in its broad compressed club. The colored paraphyses are conspicuous under the microscope. They project slightly above the surface of the receptacle and give it a soft, almost velvety appearance.

VALSA BREVIS. Pustules numerous, rather prominent, perithecia 10 to 20 or more in a pustule, nestling in the inner bark; ostiola even, black, barely emerging from and dotting or obliterated.

ting the orbicular or elliptical erumpent subpulverulent disk; asci .0008 to .001 in. long; spores allantoid .00024 to .0003 in. long.

Bark of balsam fir, *Abies balsamea*. Labrador. Waghorne.

This fungus is allied to but differs from *Valsa Friesii* in its larger pustules, more numerous perithecia and shorter asci and spores. The color of the disk is grayish or grayish-green.

HYGROPHORUS NIGRIDIVS. Pileus fleshy, convex, obtuse or subumbonate, glabrous, grayish-brown or black-brown, often a little darker in the center, flesh white; lamellae distant, decurrent, white; stem rather slender, solid, brownish, white at the top; spores elliptical, .0004 to .0005 in. long, .00024 to .0003 broad.

Pileus 1 to 2 in. broad; stem 1 to 2 in. long, 2 to 4 lines thick.

Gregarious or rarely two or three united at the base. Pine and fir tree woods. Prince Edwards Island. October and November. J. MacSwain.

This fungus differs from *H. caprinus* Fr. in its smaller size, glabrous pileus and larger spores. A description of it was published in the Country Gentleman of November 29, 1894, but one is here given that it may be more readily accessible to mycologists.

In many of the foregoing descriptions I have been obliged to give the colors indicated by the dried specimens. It is not likely therefore that they will in all cases agree rigidly with the colors of the fresh plant.

SURVEY OF MINNESOTA.

logical effects are very obscure, and when the use of simple remedies is followed to follow the use of complex mixtures dissolved in large quantities of water, but little satisfaction can be gained from theoretical speculations that attribute them to the presence of one or the other ingredient of the mixture. But little more can be said than that the water contains small quantities of substances that give, when found in large proportions, the characteristic of seltzer, chalybeate, and white-sulphur springs, and that its use in many instances has been attended with beneficial results.

In accordance with your request, that I should furnish you the results of the analyses that were made in the fall of 1875 for Messrs. Kindred and Culver, I submit the following:

I. Three specimens supposed to contain silver were sent to Dr. P. B. Rose to be assayed. The first two were from J. B. Culver, Esq., of Duluth, and were reported as containing silver. The third specimen was from C. F. Kindred, Esq., of Brainerd. It was reported as containing "52.32 oz. of silver to the ton of ore, which at \$1.30 per oz. = \$68.01 per ton."

II. A sample of iron ore, supposed to contain tin, from the neighborhood of Duluth, was sent to Prof. F. W. Morley, of Hudson, Ohio, a gentleman of some experience in iron assays. He reported two assays as yielding 10.9 per cent. and 16.17 per cent. "Tin not present in any appreciable quantity. The ore appeared to contain quite a notable quantity of Titanium. Titanium is now supposed to be injurious. * * * The amount of iron in the ore is so small that even produced from it could hardly compete with richer ores, even if flux and coal were both near at hand. One would not like to say the ore is valueless without knowing the relative amount of silica and phosphorus in the ore; but the probability that they are in such proportions as to render the addition of a flux needless, is very slight."

Another sample of iron ore from near Duluth, was sent Prof. Morley with instructions to determine the quantity of any thing he might find in it of commercial value. After giving the details of the process of analysis, he says: "You will see that everything of any commercial value is pretty thoroughly excluded. It is an iron—magnesium—sodium silicate; with not enough iron to make it of value as an iron ore."

The first mentioned iron ore resembles an iron ore found in the northern part of Rhode Island, where it occurs in a protruded mass, in some respects resembling porphyry.

Respectfully submitted,

S. F. PECKHAM,

MINNEAPOLIS, Jan. 9, 1877.

State Chemist.

V.

BOTANY. 1877

The survey has received a hearty response to the following circular, which was issued in conformity with the instructions of the Regents, in the spring of 1876 :

THE GEOLOGICAL AND NATURAL HISTORY SURVEY OF
MINNESOTA.

(BOTANY.)

To the Botanists of the State :

At the annual meeting of the Board of Regents, held in St. Paul in December, 1875, action was taken ordering the commencement of a thorough and systematic examination of the flora of the state. This was done in conformity to the law ordering a geological and natural history survey of the state, and placing it in charge of the Regents of the University. One clause of that law reads as follows :

“SEC 3. The natural history survey shall include, first, an examination of the vegetable productions of the state, embracing all trees, shrubs, herbs and grasses native or naturalized in the state; second, a complete and scientific account of the animal kingdom as properly represented in the state, including all mammalia, fishes, reptiles, birds and insects.”

In the prosecution of this examination it is expected that the Regents will have the assistance of the botanists of the state, and it is for their information that the following suggestions are made concerning the collection and preservation of information, and especially of botanical specimens.

1st. Make as full notes as possible on the flora of your own locality, not only naming species, but mentioning peculiarities and variations of structure, habitat, color, and relative abundance. Recollect that there is no published text book that professes to give an account of the flora of the country west of the Mississippi in this latitude, and that you are very largely

an independent observer. Therefore repeated verifications of an observed variation or peculiarity, or of a species not named in the familiar text-books, should be made *before reporting such observation as a fact.*

2d. Collect and preserve as many specimens as possible. They will exemplify the local flora of your region, and will serve as duplicates for exchange with other portions of the state and with foreign botanists. The most valuable portion of your contribution to the survey will be, after all, the collections which you may make and forward for careful identification.

3d. It is the design of the Regents to collect together at the University, a complete representation of the plants of the state as ordered by the law, and to have them so arranged and exhibited that they can be seen for comparison by any of the botanists of the state that may wish to examine them.

4th. For aid in the study of the flora of the state, the following works will be useful:

Gray's *Manual of Botany.*

Torrey & Gray's *Flora of North America*, 2 vols., to *Lobeliaceæ.*

De Candolle's *Prodromus*, 18 vols., down to *Endogenae.*

U. S. Geological Exploration of the Fortieth parallel. Vol. V, Botany. (Watson.)

Pacific R. R. Reports.

Transactions of the St. Louis Academy.

Proceedings of the American Academy, Philadelphia.

Catalogue of the Plants of Minnesota, by I. A. Lapham, published in the Annual Report of the State Horticultural Society for 1875. Furnished by the Secretary of the Society.

Hayden's *Nebraska Plants.*

Hooker's *Fl. Bot. Am.*, 2 vols.

Flora of Colorado, 1 vol., Porter & Coulter.

Nuttall's *Genera*, 2 vols.

Bentham's *Genera*, 4 parts.

5th. It is evident that there is no botanist, nor even any public library in the state that possesses the books necessary for the thorough study and satisfactory determination of the species of our flora. Within certain limits our flora can be studied, but the burden of our first efforts must be the collection of specimens. Their exhaustive study can only be done by experts, with the fullest facilities for comparison.

6th. The present design of the survey is to act as a means of communication between botanists of the state, to enable them to compare specimens, and as a depository for duplicates. To this end exchanges will be made with such as desire to compare species, and any aid or information will be rendered that it is possible to give. Lists of the local flora in different parts of the state should be made out as thoroughly as possible, to judge of the distribution of species. The areas that are covered with timber in the various counties, or townships, and the kinds of trees, should be stated. It is highly desirable that a local botanist be assigned to the working up of each county. For this purpose, the survey should be furnished with the

names of such botanists as are known to be interested in our state flora. Annual reports made by such local botanists would indicate the progress of the work.

N. H. WINCHELL.

THE UNIVERSITY OF MINNESOTA,

May 1st, 1876.

The survey has received several hundred specimens representing the flora of the state, and others have been gathered by Mr. Leonard and Mr. Herrick, assistants on the survey. Nothing systematic has been attempted in the way of identifications, but through the kindness of others, chiefly Mr. O. E. Garrison, of St. Cloud, and Dr. A. E. Johnson, of Minneapolis, valuable assistance has been rendered in determining our local flora.

F U N G I .

From Fifth annual report, 1877

BY DR. A. E. JOHNSON.

QK

2.5

.0B

V. 100

Prof. N. H. Winchell:—I believe no attempt prior to this has been made to collect and list the Mycologic Flora of Minnesota. As a small contribution to the Natural History of Minnesota, I present the following list of plants, in the Mycologic Flora of our state, for such use as you may be pleased to make of it.

The list has been collected mainly from Hennepin county; some from Anoka, Ramsey and Wright counties.

I have placed the Gen. or Sub-Gen. before each species; as for example, Sub-Gen. 1. *Amanita*. Hence, in reading *Agaricus vaginatus*, the word *Amanita* should be supplied thus: *Agaricus* (*Amanita*) *vaginatus*. *Bull.* And so with all the species under each Gen. or Sub-Gen. Following the specific name of the species are initials or letters used by botanists to designate the name of the author of the species, and immediately following is the common name of the plant if it has received one; then follow such words as, wood, woods, ground, sticks, stumps, epiphytal, terrestrial, etc., indicating the habitat, and lastly the month or months in which the plant is found in our climate. If there is but one month named, the plant has only been observed in that month, but if two months are named, as June, Nov. the plant has been observed in both months and most always during the intervening months:

DIVISION 1. Sporifera.

FAMILY 1. Hymenomycetes.

ORDER 1. Agaricini.

SERIES 1. Leucospori. Spores white.

SUB-GEN. Amanita. Pers.

GEN. Agaricus. L.

1. *Agaricus vaginatus*. Bull. Slick Amanita. Woods. August.
2. *A. Ceciliae*. B. & Br. Grey, ringless Amanita. Woods. Aug., Sept.
3. *A. adnatus*. Smith. Adnate-gilled Amanita. Woody places. Aug.
4. *A. vernus*. Bull. Spring Amanita. Woods, open places. June to Aug
5. *A. phalloides*. Fr. Stinking Amanita. Woods. August, Nov.
6. *A. mappa*. Batsch. Delicate Amanita. Under trees. Poisonous.
7. *A. muscarius*. L. Fly Amanita. Thin woods. Aug. & Sept. Poisonous.
8. *A. excelsus*. Fr. Tall Amanita. Woods. Aug. and Sept.
9. *A. pantherinus*. D. C. Spotted Amanita. Woods and open ground.
10. *A. asper*. Fr. Rough Amanita. Woods. June to Sept.

SUB-GEN. Lepiota. Fr.

11. *A. procerus*. Scop. Parasol mushroom. Woods. Aug. to Sept.
12. *A. excoriatus*. Schæff. Flaky Lepiota. Woods. May to Sept.
13. *A. gracilentus*. Krombh. Slender Lep. Ground. Sept. Edible.
14. *A. acutesquamosus*. Wm. Squarrose Lepiota. Ground. July.
15. *A. clypeolarius*. Bull. Fragrant Lepiota. Woods. Ju.
16. *A. Americanus*. Peck. Prairies. Aug. to Sept.
17. *A. cristatus*. Fr. Stinking Lepiota. Woods, fields. Aug.
18. *A. naucinus*. Fr. Large spored Lepiota. Woods. Aug. to Sept.
19. *A. granulatus*. Batsch. Granular Lepiota. Forests. July.
20. *A. polystictus*. Berk. Little Brown Lepiota. Ground. July.

SUB-GEN. Armillaria. Fr.

21. *A. melleus*. Vahl. Honey-coloured Armillaria. W. Aug to Sept.

SUB-GEN. Tricholoma. Fr.

22. *A. vaccinum*. Pers. Scaly Tricholoma. Ground, in woods. Sept.
23. *A. crassifolium*. Berk. Thick-gilled Tricholoma. Larch swamps.
24. *A. variegatum*. Scop. Variegated Tricholoma. Ground. June, July.
25. *A. sulfureum*. Bull. Sulphury Tricholoma. Woods. July, Aug.
26. *A. gambosum*. Fr. St. George's Mushroom. Grassy ground. June.
27. *A. melaleucum*. P. Changeable Tricholoma. Cleared ground. Sept.

SUB-GEN. *Clitocybe*. *Fr.*

28. *A. nebularis*. *Batsch*. Clouded Clitocybe. On ground in woods. Sept.
29. *A. Adirondackensis*. *Peck*. Smooth Clitocybe. Woods. Aug., Sept.
30. *A. candicans*. *Fr.* Whitish Clitocybe. Woods. Sept.
31. *A. phyllophilus*. *Fr.* Leaf-loving Clitocybe. Woods. Sept.
32. *A. dealbatus*. *P.* Ivory Clitocybe. Woods. July.
33. *A. giganteus*. *Fr.* Giant Clitocybe. Woods. Sept.
34. *A. cyathiformis*. *Fr.* Cup-shaped Clitocybe. Woods. Aug., Sept.
35. *A. laccatus*. *Scop.* Waxy Clitocybe. Woods. June-Oct.
36. *A. cerussatus*. *Fr.* White lead Clitocybe. Larch. May.
37. *A. radio-zonaria*. *n. sp.* Radiated Clitocybe. Ground, wood. June.

A full description in Bulletin of the Minnesota Academy of Natural Sciences for 1876.

SUB-GEN. *Pleurotus*. *Fr.*

38. *A. ostreatus*. *Jacq.* Oyster Pleurotus. Trunks in woods. July.
39. *A. seratinus*. *Schrad.* Yellowish Pleurotus. Larch. May.
40. *A. mastrucatus*. *Fr.* Imbricated Pleurotus. On wood. June.
41. *A. atrocæruleus*. *Fr.* Blue-black Pleurotus. On wood. July.
42. *A. sulfuroides*. *Peck*. Pine logs, &c. Sept.
43. *A. serotinoides*. *Peck*. Trunks in woods. Nov.

SUB-GEN. *Collybia*. *Fr.*

44. *A. radicans*. *Relh.* Rooting Collybia. Ground, stumps. July.
45. *A. velutipes*. *Curt.* Velvet-stemmed Collybia. Stumps. May.
46. *A. confluens*. *P.* Confluent Collybia. Woods. July.
47. *A. cirrhatus*. *Schum.* Cirrhate Collybia. June, September.
48. *A. tuberosus*. *Bull.* Tuberous Collybia. Ground, &c. Aug., Oct.
49. *A. acervatus*. *Fr.* Tufted Collybia. Woods. Aug., Oct.
50. *A. xanthopus*. *Fr.* Yellow-stemmed Collybia. Ground. July, Aug.
51. *A. dryophilus*. *Bull.* Wood Collybia. Epiphytal. June to Oct.
52. *A. exsculptus*. *Fr.* Sulphur-gill Collybia. Ground. Sept.
53. *A. plexipes*. *Fr.* Twisted Collybia. On trunks.
54. *A. stipitarius*. *Fr.* Fibrillose Collybia. Epiphytal. July to Aug.

SUB-GEN. *Mycena*. *Fr.*

55. *A. praelongus*. *Peck*. Epiphytal. Swamps. June to July.
56. *A. paluster*. *Peck*. Larch swamps. Sphagnum. June.
57. *A. radius*. *n. sp.* Pale-yellow Mycena. Ground. May.

See Bulletin of the Minn. Acad. of Natural Sciences. 1876.

58. *A. strobilinus*. *Pers.* Fir-cone. Mycena. Epiphytal.
59. *A. purus*. *P.* Amethyst Mycena. Larch swamps. June to July.

- 60. *A. pauperculus*. *Berk.* Little-stump *Mycena*. July.
- 61. *A. sanguineolentus*. *A. & S.* Stinking *Mycena*. June to Oct.
- 62. *A. crocatus*. *Schrad.* The Stainer. Woods. June to July.
- 63. *A. epipterygius*. *Scop.* Yellow-stem *Mycena*. July to Aug.
- 64. *A. stylobates*. *P.* Discoid *Mycena*. Epiphytal. Woods. Aug.
- 65. *A. corticola*. *Schum.* Bark *Mycena*. July, Aug.
- 66. *A. capillaris*. *Schum.* Capillary *mycena*. Leaves. Aug.

SUB-GEN. *Omphalia*. *Fr.*

- 67. *A. pyxidatus*. *Bull.* Variable *Omphalia*. Ground. July.
- 68. *A. affricatus*. *Fr.* Hairy Bog *Omphalia*. Ground. July.
- 69. *A. muralis*. *Sow.* Wall *Omphalia*. Ground. July.
- 70. *A. umbelliferus*. *L.* Common *Omphalia*. Ground. July. Sept.
- 71. *A. rufulus*. *B. & Br.* Reddish *Ompahlia*. Ground. July.
- 72. *A. campanella*. *Batsch.* Tawny *Omphalia*. Larch swamps.
- 73. *A. chryseus*. *Peck.* Logs in woods. August.
- 74. *A. fibula*. *Bull.* Button *Omphalia*. Ground. June, Oct.
- 75. *A. gracillimus*. *Weinm.* Delicate *Omphalia*. Epiphytal. Aug.
- 76. *A. integrellus*. *P.* Little-white *Omphalia*. Epi. Aug., Sept.

SERIES. 2. *Hyporhodii*. *Fr.*SUB-GEN. *Volvaria*. *Fr.*

- 77. *A. bombycinus*. *Schæff.* Silky *Volvaria*. Epiphytal. July, Aug.
- 78. *A. volvaceus*. *Bull.* Stove *Volvaria*. Ground. July, Aug.
- 79. *A. Taylori*. *Berk.* Taylor's *Volvaria*. Ground. Aug., Sept.
- 80. *A. gloiocephalus*. *Fr.* Umbonate *Volvaria*. Ground. June.

SUB-GEN. *Chamæota*. *Smith.*

- 81. *A. cretaceus*. *Fr.* Chalky *Chamæota*. Ground. Aug.

SUB-GEN. *Pluteus*. *Fr.*

- 82. *A. cervinus*. *Schæff.* Fawn *Pluteus*. Ground. May, Oct.
- 83. *A. nanus*. *P.* Mealy *Pluteus*. Logs in woods. Aug.
- 84. *A. petasatus*. *Fr.* Broad-capped *Pluteus*. Epiphytal. Aug.
- 85. *A. leoninus*. *Schæff.* Yellow *Pluteus*. On wood. Aug., Oct.
- 86. *A. chrysophæus*. *Schæff.* Dingy *Pluteus*. On wood. Aug.
- 87. *A. phlebophorus*. *Ditm.* Veined *Pluteus*. On wood. Aug.

SUB-GEN. *Entoloma*. *Fr.*

- 88. *A. strictior*. *Peck.* Ground. Sept., Oct.
- 89. *A. sinuatus*. *Fr.* Woods. Ground. Poisonous. July.
- 90. *A. prunuloides*. *Fr.* Plum-like *Entoloma*. Woods, ground. Sept.

91. *A. helodes*. *Fr.* Moor Entoloma. Ground, woods. Sept.
92. *A. repandus*. *Bull.* Repand Entoloma. Ground, woods. July.
93. *A. ardociacus*. *Bull.* Meadow Entoloma. Ground, woods. Sept.
94. *A. sericellus*. *Fr.* Silky Entoloma. Woods. Aug., Sept.
95. *A. rhodopolus*. *Fr.* Rosy Entoloma. Ground, woods. Aug., Sept.
96. *A. costatus*. *Fr.* Costate Entoloma. Ground. Sept.
97. *A. cuspidatus*. *Peck.* Swamps. Sphagnus marshes. Sept.

SUB-GEN. *Clitopilus*. *Fr.*

98. *A. prunulus*. *Scop.* Plum Clitopilus. Woods, ground. Season.
99. *A. cretatus*. *Berk.* Chalky Clitopilus. Woods, ground. Aug., Sept.
100. *A. noveboracensis*. *Peck.* Dingy white Clitopilus. Ground. Aug.
101. *A. Woodianus*. *Peck.* On ground, and wood. Sept.
102. *A. undatus*. *Fr.* Waved Clitop. Aug., Sept.

SUB-GEN. *Claudopus*. *Smith.* *Seem.* Journal.

103. *A. euosmus*. *Berk.* Tarragon Clau. Wood. June.
104. *A. depluens*. *Batsch.* Ground Clau. Ground. Sept.

SUB-GEN. *Leptonia*. *Fr.*

105. *A. chalybæus*. *P.* Steel-blue Leptonia. Prairies. July, Sept.
106. *A. incanus*. *Fr.* Hoary Leptonia. Prairies. Dayton. Aug.

SUB-GEN. *Nolanea*. *Fr.*

107. *A. pascuus*. *P.* Pasture Nola. Wet prairies. June.
108. *A. rufo-carneus*. *Berk.* Red-brown Nola. Marshes. Aug., Sept.
109. *A. Babingtonii*. *Blox.* Babington's Nola. Woods. Sept.
110. *A. conicus*. *Peck.* On rotten wood. Swamps. Aug.
111. *A. delicatulus*. *Peck.* Delicate Nola. Larch swamp. Aug.
112. *A. Clintonianus*. *Peck.* Clinton's Nola. Swamps. Aug.

SERIES 3. *Dermini*. *Fr.*

SUB-GEN. *Pholiota*. *Fr.*

113. *A. præcox*. *P.* Spring Phol. Prairies. May.
114. *A. comosus*. *Fr.* Hairy Phol. Decaying trunks. Aug.
115. *A. squarrosus*. *Mull.* Scaly Phol. Trunks. Aug.
116. *A. flammans*. *Fr.* Yellow scaly Phol. Ground. June.
117. *A. temnophyllus*. *Peck.* Brownish Phol. Ground. June, July.

SUB-GEN. *Hebeloma*. *Fr.*

118. *A. punctatus*. *Fr.* Ground in woods. Sept.
119. *A. crustuliniformis*. *Bull.* Ring Hebeloma. Woods. Sept.

120. *A. fastibilis*. *Fr.* Ochrey Hebe. Woods. July-Oct. Common.
121. *A. stellatosporus*. *Peck.* Stellate Hebeloma. Ground. Sept.
122. *A. griseo-cabrosus*. *Peck.* Rough Hebeloma. Popple groves. Sept.
123. *A. illicitus*. *Peck.* Smooth Hebeloma. Popple groves. Sept.
124. *A. ascophorus*. *Peck.* Viscid Hebeloma. Burned ground. Sept.
125. *A. mutatus*. *P.* Changeable Hebeloma. Ground. July, Aug.
126. *A. pyriodorus*. *P.* Pear-scented Hebeloma. Woods. Sept.
127. *A. obscurus*. *P.* Violet Hebeloma. Ground. July.
128. *A. flocculosus*. *Berk.* Flocculose Hebeloma. Ground. Sept.
129. *A. deglubens*. *Fr.* Peeling Hebeloma. Woods. Aug., Sept.
130. *A. fastigiatus*. *Fr.* Peaked Hebeloma. Woods. July.
131. *A. rimosus*. *Bull.* Cracked Hebeloma. Ground. June, Sept.
132. *A. trechisporus*. *Berk.* Rough-spored Hebeloma. Ground. Aug.
133. *A. auricomus*. *Batsch.* Golden-haired Hebeloma. Woods. July.
134. *A. flocculentus*. *Poll.* Woolly Hebeloma. Ground. July.

SUB-GEN. *Flammula*. *Fr.*

135. *A. polychrous*. *Berk.* Reddish *Flammula*. Ground, wood. Aug., Sept.
136. *A. gummosus*. *Lasch.* Viscid *Flammula*. Wood. July.
137. *A. sapineus*. *Fr.* Bright-Spored *Flam.* Ground and wood. Aug.

SUB-GEN. *Crepidotus*. *Fr.*

138. *A. mollis*. *Schæff.* Soft *Crepidotus*. On wood. July-Oct.

SUB-GEN. *Naucoria*. *Fr.*

139. *A. semiorbicularis*. *Bull.* Half-round *Naucoria*. Dung. June.
140. *A. vernalis*. *Peck.* Decaying wood. June.
141. *A. lignicola*. *Peck.* Decaying wood. June.
142. *A. erinaceus*. *Fr.* Hedgehog *Naucoria*. Wood. Nov.

SUB-GEN. *Galera*. *Fr.*

143. *A. ovalis*. *Fr.* Oval *Galera*. Cow-dung. Aug.
144. *A. tener*. *Schæff.* Slender *Galera*. Dung. July-Oct.
145. *A. sparteus*. *Fr.* Meadow *Galera*. Mossy ground. June.
146. *A. hypnorum*. *Batsch.* Moss-loving *Galera*. July. Sept.
147. *A. sphagnum*. *Pers.* Bog-moss *Galera*. July, Sept.

SUB-GEN. *Tubaria*. *Smith. Journ., 1870.*

148. *A. inquilinus*. *Fr.* Little *Tubaria*. On wood, swamps. June, Sept.
149. *A. furfuraceus*. *P.* Mealy *Tubaria*. Chips, wood. July, Sept.

SERIES 4. *Praetellæ*. *Fr.*SUB-GEN. *Psalliota*. *Fr.*

150. *A. campestris*. *L.* Common Mushroom. Ground. August.
Var. pratensis. *Vitt.* Rich muck. Aug.
Var. silvicola. *Vitt.* Ramsey and Wright counties. Aug.
 151. *A. silvaticus*. *Schæff.* Wood *Psalliota*. Woods. Aug.
 152. *A. Johnsonianus*. *Peck.* Woods. Sept.

SUB-GEN. *Pilosace*. *Fr.*

153. *A. eximius*. *Peck.* Decaying logs in woods. Sept.

SUB-GEN. *Stropharia*. *Fr.*

154. *A. stercorearius*. *Fr.* Dung *Stropharia*. May, Sept.
 155. *A. semiglobatus*. *Batsch.* Semiglobose *Stropharia*. June, Sept.

SUB-GEN. *Hypholoma*. *Fr. Hab.* Generally on stumps.

156. *A. sublateritius*. *Fr.* Brick-red *Hypholoma*. Sept.
 157. *A. fascicularis*. *Hud.* Tufted Yellow *Hypholoma*. Sept.
 158. *A. lacrymabundus*. *Fr.* Weeping Hyph. July, Nov.
 159. *A. velutinus*. *P.* Velvety Hyph. Trunks. July.
 160. *A. perplexus*. *Peck.* Ground about stumps. Sept.
 161. *A. phyllogenus*. *Peck.* Fallen leaves. July

SUB-GEN. *Psilocybe*. *Fr. Hab.* Mostly on the ground.

162. *A. limicola*. *Peck.* Aug., Sept.
 163. *A. spadiceus*. *Schæff.* Bay *Psilocybe*. Wood. Ground. Aug.
 164. *A. cernuus*. *Mull.* Nodding *Psilocybe*. Wood. Ground. Sept.
 165. *A. foenisecii*. *P.* Brown *Psilocybe*. Ground. Sept.

SUB-GEN. *Psathyra*. *Fr. Hab.* Ground. Wood.

166. *A. mastiger*. *Berk & Br.* Peaked *Psathyra*. Ground. Aug., Sept.
 167. *A. corrugis*. *P.* Wrinkled *Psathyra*. Ground. Aug.
 168. *A. obtusatus*. *Fr.* Obtuse *Psathyra*. Ground. June.

SERIES 5. *Coprinarii*. *Fr.* Spores black.SUB-GEN. *Panæolus*. *Fr.* Mostly on dung.

169. *A. separatus*. *L.* Ochrey *Panæolus*. June, Sept.
 170. *A. leucophanes*. *B. & Br.* Shiny-White *Panæolus*. May, Aug.
 171. *A. fimiputris*. *Bull.* Dark-gray *Panæolus*. June, Aug.

172. *A. campanulatus*. *L.* Campanulate *Panæolus*. June, Aug.
 173. *A. papilionaceus*. *Bull.* Butterfly *Panæolus*. June, July.
 174. *A. solidipes*. *Peck.* July.
 175. *A. fimicola*. *Fr.* Dung *Panæolus*. June.

SUB-GEN. *Psathyrella*. *Fr.*

176. *A. gracilis*. *Fr.* Slender *Psathyrella*. Sept.
 177. *A. pronus*. *Fr.* Stooping *Psathyrella*. Ground. Sept.
 178. *A. atomatus*. *Fr.* Sprinkled *Psathyrella*. Manure. June, July.
 179. *A. disseminatus*. *Fr.* Clustered *Psathyrella*. Woods. July, Sept.
 180. *A. odoratus*. *Peck.* Manure heaps. May, June.

GEN. 2. *Coprinus*. *Fr.*

181. *C. comatus*. *Fr.* Shaggy *Coprinus*. Rich ground. Sept.
 182. *C. atramentarius*. *Fr.* Inky *Coprinus*. Dung. June, July.
 183. *C. fuscescens*. *Fr.* Brownish *Coprinus*. Wood. June, July.
 184. *C. picaceus*. *Fr.* Magpie *Coprinus*. Ground. Sept.
 185. *C. similis*. *B. & Br.* Striate *Coprinus*. Wood. Sept.
 186. *C. fimetarius*. *Fr.* Shaggy-dung *Coprinus*. June, July.
 187. *C. tomentosus*. *Fr.* Downy *Coprinus*. Dung. May, June.
 188. *C. miveus*. *Fr.* Snowy *Coprinus*. May, June.
 189. *C. micaceus*. *Fr.* Glistening *Coprinus*. May, Sept.
 190. *C. deliquescens*. *Fr.* Deliquescent *Coprinus*. June.
 191. *C. lagopus*. *Fr.* Hare's foot *Coprinus*. Dung. July.
 192. *C. radiatus*. *Fr.* Delicate *Coprinus*. Dung. May, June.
 193. *C. ephemerus*. *Fr.* Ephemeral *Coprinus*. May, June.
 194. *C. insignis*. *Peck.* About roots of trees. July, Aug.
 195. *C. angulatus*. *Peck.* Ground in woods. Aug., Sept.
 196. *C. plicatilis*. *Fr.* Rich ground. June, July.
 197. *C. hemerobius*. *Fr.* Collared *Coprinus*. July.
 198. *C. silvaticus*. *Peck.* Ground. Sept.
 199. *C. semilanatus*. *Peck.* Dung. Aug.

GEN. 3. *Bolbitius*. *Fr.* Manuré, rich soil.

200. *B. Boltoni*. *Fr.* Bolton's *Bolbitius*. Dung. June, Sept.
 201. *B. fragilis*. *Fr.* Fragile *Bolbitius*. Epiphytal. May, Aug.
 202. *B. titubans*. *Fr.* Wavering *Bolbitius*. Ground. Forest. Open. Ju., Oct.
 203. *B. apicalis*. *Smith.* Two-coloured *Bolbitius*. Ground. June, July.
 204. *B. nobilis*. *Peck.* Noble *Bolbitius*. Ground. Woods. Sept.

GEN. 4. *Cortinarius*, *Fr.*SUB-GEN. 1. *Phlegmacium*. *Fr.*

205. *C. varius*. *Fr.* Variable Cort. Ground. Woods. Aug., Sept.

- 206. *C. cyanipes*. *Fr.* Blue-stemmed Cort. Woods. July, Aug.
- 207. *C. russus*. *Fr.* Ruddy Cort. Woods. Sept.
- 208. *C. coloratus*. *Peck.* Amongst moss. Larch swamps. Sept.
- 209. *C. communis*. *Peck.* Woods. Sept.
- 210. *C. multiformis*. *Fr.* Multiform Cort. Woods. Sept.
- 211. *C. glaucopus*. *Fr.* Brown-zoned Cort. Sept.
- 212. *C. callochrous*. *Fr.* Tawny-viscid Cort. Woods. Aug., Sept.
- 213. *C. cœrulescens*. *Fr.* Azure-blue Cort. Woods. Sept.
- 214. *C. turbinatus*. *Fr.* Top-shaped Cort. July, Sept.
- 215. *C. scaurus*. *Fr.* Club-footed Cort. Woods. Sept.
- 216. *C. corrugatus*. *Peck.* Woods. June.

SUB-GEN. 2. *Myxadium*. *Fr.*

- 217. *C. collinitus*. *Fr.* Smeared Cort. Woods. Sept.
- 218. *C. sphærosporus*. *Peck.* Woods. Sept.

SUB-GEN. 3. *Inoloma*. *Fr.*

- 219. *C. violaceus*. *Fr.* Violet Cort. Woods. Aug.
- 220. *C. camphoratus*. *Fr.* Strong-scented Cort. Ground. Sept.
- 221. *C. callisteus*. *Fr.* Stout Cort. Woods. August.
- 222. *C. pholideus*. *Fr.* Scaly Cort. Woods. Sept.
- 223. *C. sublanatus*. *Fr.* Woolly Cort. Woods. Sept.
- 224. *C. lilacinus*. *Peck.* Woods. Sept.
- 225. *C. squamulosus*. *Peck.* Scaly Cort. Woods. Aug., Sept.
- 226. *C. asper*. *Peck.* Newly cleared places. Sept.

SUB-GEN. *Dermocybe*. *Fr.*

- 227. *C. anomalus*. *Fr.* Thin-capped Cort. Woods. Sept. Oct.
- 228. *C. spilomeus*. *Fr.* Scaly-stemmed Cort. Woods. Sept.

SUB-GEN. *Telamonia*. *Fr.*

- 229. *C. bulbosus*. *Fr.* Bulbous Cort. Woods. Aug., Sept.
- 230. *C. lignarius*. *Peck.* Decayed wood. June.
- 231. *C. limonius*. *Fr.* Lemon Cort. Decaying pine. June.
- 232. *C. hinnuleus*. *Fr.* Fawn Cort. Woods. June.
- 233. *C. psammocephalus*. *Fr.* Little Tawny Cort. Woods. Aug., Sept.
- 234. *C. illeopodius*. *Fr.* Tan-colored Cort. Woods. June, Sept.

SUB-GEN. *Hygrocybe*. *Fr.*

- 235. *C. Armeniacus*. *Fr.* Peach Cort. Woods.
- 236. *C. vernalis*. *Peck.* Spring Cort. Ground. June.
- 237. *C. castaneus*. *Fr.* Chestnut Cort. Woods. Sept.
- 238. *C. Reedii*. *Berk.* Reed's Cort. Shores of lakes. June.

239. *C. leucopus*. *Fr.* White-stemmed Cort. Woods. May.
 240. *C. decipiens*. *Fr.* Deceptive Cort. Woods. Sept.

GEN. *Lepister*. *Smith*. *Sieem.* Jour. 1870.

241. *L. nudla*. *Bull.* Amethyst Lep. Woods. Prairies. Aug. Sept.
 242. *L. personata*. *Fr.* Purple-stemmed Lep. Ground. Aug. Sept.

GEN. *Paxillus*. *Fr.*

243. *P. involutus*. *Fr.* Involute Pax. Ground in woods. Aug. Sept.
 244. *P. strigosus*. *Peck*. Hairy Pax. Ground, woods. Sept.

GEN. *Hygrophorus*. *Fr.* Terrestrial. Woods, on ground.

245. *H. ebureus*. *Fr.* Ivory Hygroph. Woods. Sept., Oct.
 246. *H. cossus*. *Fr.* Goat-moth Hygroph. Woods. Sept., Oct.
 247. *H. cerascinus*. *B.* Waxy Hygroph. Woods. Sept., Oct.
 248. *H. aromaticus*. *B.* Aromatic Hygroph. Woods. Sept.
 249. *H. mesotephrus*. *B. & Br.* Brown-disk Hygroph. Woods. Aug., Sept.
 250. *H. hypothecus*. *Fr.* Pine-wood Hygroph. Sandy soil. Woods. Sept.
 251. *H. olivaceo-albus*. *Fr.* Olive Hygroph. Woods. Sept.
 252. *H. leporinus*. *Fr.* Hare-colored Hygroph. Terrestrial. Sept.
 253. *H. pratensis*. *Fr.* Pasture Hygroph. Open places, woods. Sept.
 254. *H. niveus*. *Fr.* Snow-white Hygroph. Mossy ground. Aug., Oct.
 255. *H. ceraceus*. *Fr.* Wax-like Hygroph. Woods. Sept., Oct.
 256. *H. miniatus*. *Fr.* Vermillion Hygroph. Woods. Aug., Sept.
 257. *H. coniscus*. *Fr.* Conical Hygroph. Terrestrial. Aug., Oct.
 258. *H. nitidus*. *B. & R.* Amongst moss in wet places. Aug., Sept.

GEN. *Gomphidius*. *Fr.* Terrestrial mainly.

259. *G. viscidus*. *Fr.* Viscid Gomphidius. On wood. Aug.

GEN. *Lactarius*. *Fr.* Terrestrial. Mainly in woods.

260. *L. torminosus*. *Fr.* Woolly Lactarius. Aug., Oct.
 261. *L. cilicioides*. *Fr.* Tomentose Lact. Sept.
 262. *L. turpis*. *Fr.* Dirty Lact. July.
 263. *L. pubescens*. *Schrad.* Pubescent Lact. Aug., Sept.
 264. *L. zonarius*. *Fr.* Zoned Lact. Aug., Oct.
 265. *L. blennius*. *Fr.* Slimy Lact. Aug., Sept.
 266. *L. pyrogalus*. *Fr.* Pear-scented Lact. Aug., Sept.
 267. *L. plumbeus*. *Fr.* Lead-coloured Lact. Aug., Sept.
 268. *L. chrysorrhæus*. *Fr.* Yellow juiced Lact. July, Aug.
 269. *L. piperatus*. *Fr.* Peppery Lact. July, Sept.
 270. *L. vellereus*. *Fr.* Woolly-white Lact. July.
 271. *L. deliciosus*. *Fr.* Delicious Lact. Aug., Oct.

- 272. *L. pallidus*. *Fr.* Pallid Lact. Aug., Sept.
- 273. *L. theiogalus*. *Fr.* Sulphur-juiced Lact. Aug.
- 274. *L. cyathala*. *Fr.* Cup-like Lact. Aug., Sept.
- 275. *L. glyciosmus*. *Fr.* Scented Lact. Aug., Oct.
- 276. *L. serifluus*. *Fr.* Thin-juiced Lact. Sept.
- 277. *L. fuliginosus*. *Fr.* Dingy Lact. July, Oct.
- 278. *L. affinis*. *Peck.* Viscid Lact. Sept., Oct.
- 279. *L. volemus*. *Fr.* Orange-brown Lact. July, Sept.
- 280. *L. platyphyllus*. *Peck.* Aug., Sept.
- 281. *L. sordidus*. *Peck.* Sandy soil. Sept.
- 282. *L. griseus*. *Peck.* Low ground. Aug.

GEN. *Russula*. *Fr.* Terrestrial. Usually late Summer and Autumn.

- 283. *R. nigricans*. *Fr.* Blackish Rus. July, Aug.
- 284. *R. aduster*. *Fr.* Scorched Rus. Sept., Oct.
- 285. *R. delica*. *Fr.* Whitish Rus. Woods. Aug.
- 286. *R. furcata*. *Fr.* Forked Rus. Woods. Sept.
- 287. *R. sanguinea*. *Fr.* Blood-red Rus. Woods. July.
- 288. *R. rosacea*. *Fr.* Rosy Rus. Woods. July.
- 289. *R. sardonina*. *Fr.* Changeable Rus. Woods. July.
- 290. *R. depallens*. *Fr.* Bleached Rus. Woods. July.
- 291. *R. virescens*. *Fr.* Greenish Rus. Woods. July, Sept.
- 292. *R. lepida*. *Fr.* Scaly Rus. Woods. July, Aug.
- 293. *R. rubra*. *Fr.* Red Rus. Woods. July, Aug.
- 294. *R. fœtens*. *Fr.* Fœtid Rus. Woods. July, Sept.
- 295. *R. emetica*. *Fr.* Emetic Rus. Woods. July, Oct.
- 296. *R. fragilis*. *Fr.* Fragile Rus. Woods. July, Aug.
- 297. *R. integra*. *Fr.* Entire Rus. Woods. July, Aug.
- 298. *R. decolorans*. *Fr.* Discolored Rus. Woods. Sept.
- 299. *R. veteriosa*. *Fr.* Sleepy Rus.
- 300. *R. nitida*. *Fr.* Shining Rus. Woods. Sept.
- 301. *R. alutacea*. *Fr.* Tan-colored Rus. Woods. July, Aug.
- 302. *R. lutea*. *Fr.* Yellow Rus. Woods. August.
- 303. *R. chamæleontina*. *Fr.* Chameleon Rus. Woods. Sept.
- 304. *R. marlæ*. *Peck.* Woods. July, Aug.

GEN. *Cantharellus*. *Adams' Fung.*

- 305. *C. cibarius*. *Fr.* Edible Chantarelle. Woods. July.
- 306. *C. aurantiacus*. *Fr.* False Chant. Ground and wood. Aug.
- 307. *C. umbonatus*. *P.* Umbonate Chant. Ground. July.
- 308. *C. tubæformis*. *Fr.* Tubæform Chant. Woods. July.
- 309. *C. infundibuliformis*. *Fr.* Funnel-shaped Chant. Ground. July, Aug.
- 310. *C. minor*. *Peck.* Ground in woods. July, Aug.
- 311. *C. dichotomus*. *Peck.* Damp ground. Woods. July, Aug.

GEN. *Nyctalis*. *Fr.* Gen. Hymen.

- 312. *C. asterophora*. *Fr.* Star-bearing *Nyctalis*. Dead Fungi. Sept.

GEN. MARASMIUS. *Fr.* Epiphytal. Terrestrial.

- 313. *M. oreades.* *Fr.* Fairy-ring. Champignon. May, Oct.
- 314. *M. fusco-purpureus.* *Fr.* Purple brown Maras. Woods. June, July.
- 315. *M. Wynnei.* *B. & Br.* Wynne's Maras. Leaves. June, July.
- 316. *M. erythrophus.* *Fr.* Pallid Maras. On ground and wood. July.
- 317. *M. terginus.* *Fr.* Clustered Maras. In woods on wood. June.
- 318. *M. alliaceus.* *Fr.* Onion-scented Maras. Wood. July, Aug.
- 319. *M. rotula.* *Fr.* Collared Maras. Ground. June, Oct.
- 320. *M. androsaceus.* *Fr.* Black stemmed Maras. June, Sept.
- 321. *M. insititius.* *Fr.* Horny stemmed Maras. Aug., Oct.
- 322. *M. epiphyllus.* *Fr.* Leaf Maras. Woods. June, Oct.
- 323. *M. saccharinus.* *Fr.* Granular Maras. Epiphytal. Woods. June, July.
- 324. *M. spodoleucus.* *B. & Br.* Stemless Maras. Epiphytal. Woods. June, Sept.
- 325. *M. velutipes.* *B. & C.* Woolly Maras. Woods. July.
- 326. *M. plancus.* *Fr.* Hairy Maras. Woods. June, Oct.
- 327. *M. subvenosus.* *Peck.* Leaves in woods. Aug., Oct.
- 328. *M. campanulatus.* *Peck.* Leaves in woods. Aug.
- 329. *M. cæspitosus.* *Peck.* Decaying branches, woods. June.
- 330. *M. longipes.* *Peck.* Among fallen leaves, woods. Aug., Oct.
- 331. *M. glabellus.* *Peck.* Amongst leaves, woods. July, Sept.
- 332. *M. anomalus.* *Peck.* Sticks, leaves in woods. July.

GEN. LENTINUS. *Fr.* Generally on wood.

- 333. *L. tigrinus.* *Fr.* Tiger-spot Len. June, Oct.
- 334. *L. Dunalii.* *Fr.* Dunal's Len. June, Sept.
- 335. *L. lepidius.* *Fr.* Scaly Len. June, July.
- 336. *L. cochleatus.* *Fr.* Shell Len. June, July.
- 337. *L. vulpinus.* *Fr.* Strong-scented Len. May, Aug.

GEN. PANUS. *Fr.* Epiphytal. Stumps.

- 338. *P. torulosus.* *Fr.* Twisted Pan. Sept.
- 339. *P. conchatus.* *Fr.* Shell Pan. May, Oct.
- 340. *P. salicinus.* *Peck.* On *Salix discolor.* Michx. Sept., Oct.
- 341. *P. strigosus.* *B. & C.* Aug.

GEN. TROGIA. *Fr.*

- 342. *T. crispa.* *Fr.* Crisped Trogia. On dead branches. Sept

GEN. SCHIZOPHYLLUM. *Fr.*

- 343. *S. commune.* *Fr.* Common Schiz. On dead wood. The season.

GEN. Lenzites. *Fr.*

344. *L. betulina*. *Fr.* Birch Lenzites. Stumps. Perennial.
 345. *L. floccida*. *Fr.* Flaccid Lenzites. Stumps. Sept., Oct.

ORDER II. Polyporei.

GEN. Boletus. *Fr.* Terrestrial.

346. *B. elegans*. *Schum.* Elegant Boletus. Woods. June, Oct.
 347. *B. flavus*. *With.* Bright-yellow Bolet. Woods. July.
 348. *B. badius*. *Fr.* Bay Boletus. Woods. Wright county. August.
 349. *B. striæpes*. *Sec.* Striate Bolet. Aug.
 350. *B. chrysenteron*. *Fr.* Red-cracked Bolet. Sept.
 351. *B. subtomentosus*. *L.* Yellow-cracked Bolet. Woods. Aug.
 352. *B. pachypus*. *Fr.* Thick-stemmed Bolet. Aug., Sept.
 353. *B. edulis*. *Bull.* Edible Bolet. Woods. Aug.
 354. *B. æstivalis*. *Fr.* Early Bolet. Woods. Esculent.
 355. *B. purpureus*. *Fr.* Purple Bolet. Woods. Aug.
 356. *B. scaber*. *Fr.* Shaggy Bolet. Woods. Aug. Esculent.
 357. *B. alutarius*. *Fr.* Tan-colored Bolet. Woods. Aug.
 358. *B. felleus*. *Bull.* Bitter Bolet. Woods. Sept.
 359. *B. cyanescens*. *Bull.* Sibthorp's Bolet. Woods. Aug.

GEN. Polyporus. *Fr.*

360. *P. leptocephalus*. *Fr.* White-pored Poly. On wood. June, July.
 361. *P. perennis*. *Fr.* Perennial Poly. Ground, stumps. Aug., Oct.
 362. *P. Rostkovii*. *Fr.* Rostkovius' Poly. Rotton wood. June, Sept.
 363. *P. elegans*. *Fr.* Elegant Poly. Trunks, branches, woods. July.
 364. *P. quercinus*. *Fr.* Oak Poly. Old oaks. June.
 365. *P. sulfureus*. *Fr.* Sulphury Poly. Logs, stumps. June, Sept.
 366. *P. alligatus*. *Fr.* Connected Poly. Base of stumps. June, Sept.
 367. *P. heteroclitus*. *Fr.* Grand Poly. Ground. Wright county. Aug.
 368. *P. salignus*. *Fr.* Willow Poly. On Willows. July, Nov.
 369. *P. chioneus*. *Fr.* Soft white Poly. Roots of stumps. Aug.
 370. *P. cassius*. *Fr.* Blue-gray Poly. On pine logs. July, Oct.
 371. *P. destructus*. *Fr.* Destructive Poly. Larch, ground. July, Sept.
 372. *P. rutilans*. *Fr.* Reddish Poly. Branches, woods. June.
 373. *P. adustus*. *Fr.* Scorched Poly. Stumps.
 374. *P. hispidus*. *Fr.* Hispid Poly. Living oaks. Perennial.
 375. *P. spumeus*. *Fr.* Oozing Poly. Trunks, branches. July.
 376. *P. nigricans*. *Fr.* Black-hoof Poly. Living birch. Perennial.
 377. *P. annosus*. *Fr.* Imbricated Poly. Larch stumps. Perennial.
 378. *P. connatus*. *Fr.* Connate Poly. Crab trunks. Miss R. A. Johnson.
 379. *P. hirsutus*. *Fr.* Bristly Poly. Dead trunks. Woods. July.
 380. *P. versicolor*. *Fr.* Common Zoned Poly. Stumps, &c. Persistent.

- 381. *P. abietinus*. *Fr.* Whitish Fir Poly. Larch. July.
- 382. *P. contiguus*. *Fr.* Contiguous Poly. Decayed wood, &c. June, Sep.
- 383. *P. ferruginosus*. *Fr.* Rusty Poly. Posts, &c. June, Sept.
- 384. *P. Armeniacus*. *Berk.* Buff Fir Poly. June.
- 385. *P. incarnatus*. *Fr.* Flesh-Colored Poly. Larch. July, Aug.
- 386. *P. violaceus*. *Fr.* Violet Poly. Poplar Larch. July, Aug.
- 387. *P. medulla-panis*. *Fr.* Crumb of Bread Poly. Wood. June, Sept.
- 388. *P. obducens*. *Fr.* Incrusting Poly. Rotten wood. Perennial.
- 389. *P. vulgaris*. *Fr.* Common-effused Poly. Dead wood.
- 390. *P. vaporarius*. *Fr.* Creeping Poly. Fallen branches.
- 391. *P. glomeratus*. *Peck.* On *Acer saccharinum*. *Wang.* Aug.
- 392. *P. Gordoniensis*. *B. & Br.* Gordon's Fir Poly. Decaying wood. Sep.

GEN. *Trametes*. *Fr.*

- 393. *T. pini*. *Fr.* Fir trunk *Trametes*. Pine trunks. Perennial.
- 394. *T. odora*. *Fr.* Small pored *Trametes*. Willows.

GEN. *Dædalea*. *Fr.*

- 395. *D. unicolor*. *Fr.* One-coloured *Dædalea*. Stumps. Trunks.
- 396. *D. latissima*. *Fr.* Effused *Dædalea*. On fallen dead branches.

GEN. *Merulius*. *Fr.*

- 397. *M. tremellosus*. *Schrad.* Tremellose *Merulius*. Sept.
- 398. *M. corium*. *Fr.* Leathery *Merulius*. Dead trunks.
- 399. *M. malluscus*. *Fr.* Thin *Merulius*. Dead wood.
- 400. *M. rufus*. *P.* Rufous *Merulius*. Dead oak branches.
- 401. *M. serpens*. *Fr.* Creeping *Merulius*. Dead wood. June, Sept.

GEN. *Porothelium*. *Fr.*

- 402. *P. Friesii*. *Mont.* Fries' *Porothelium*. Pine wood.

ORDER III. *Hydnei*.

GEN. *Hydnum*. *Linn.*

- 403. *H. repandum*. *L.* Spreading Hyd. Ground. Aug., Sept.
- 404. *H. zonatum*. *Batsch.* Zoned Hyd. Ground. Woods. Sept.
- 405. *H. tomentosum*. *L.* Tomentose Hyd. Ground. Woods. July.
- 406. *H. erinaceus*. *Hull.* Hedgehog Hyd. Living Oak. Sept., Oct.
- 407. *H. niveum*. *P.* Snowy Hyd. Dead wood. Leaves. Sept.
- 408. *H. farinaceum*. *P.* Mealy Hyd. Decaying wood. July, Sept.

ORDER IV. *Auricularini*. *Fr.*

GEN. *Cratorellus*. *Fr.*

- 409. *C. cornucopioides*. *Fr.* Horn-like *Cratellus*. Ground.

GEN. Thelephora. *Fr.*

410. *T. cæsia*. *P.* Ash-gray Thelephora. Incrusting grass.
 411. *T. arida*. *Fr.* Dry Thelephora. Decaying pine.

GEN. Stereum. *Fr.*

412. *S. purpureum*. *Fr.* Purple Stereum. Trunks. Perennial.
 413. *S. hirsutum*. *Fr.* Hairy Stereum. Stumps. Peren.
 414. *S. acerinum*. *Fr.* Maple Stereum. Living maple trunks.

GEN. Hymenochæte. *Lev.*

415. *H. rubiginosa*. *Lev.* Rubiginous Hymenochæte.

ORDER V. Clavariici.

GEN. Clavaria. *L.*

416. *C. amethystina*. *Bull.* Amethyst Clavaria. Sept.
 417. *C. fastigiata*. *D. C.* Fastigiate Clavaria. Woods. Aug., Oct.
 418. *C. coralloides*. *L.* White Coral Clavaria. Woods. Aug., Sept.
 419. *C. umbrina*. *Berk.* Umber Clavaria. Woods. Aug., Sept.
 420. *C. cinerea*. *Bull.* Cinereous Clav. Woods. Sept.
 421. *C. cristata*. *Holmsk.* Crested Clav. Woods. Sept.
 422. *C. rugosa*. *Bull.* Wrinkled Clav. Woods. Sept.
 423. *C. Kunzei*. *Fr.* Kunze's Clav. Woods. Sept.
 424. *C. aurea*. *Schæff.* Golden Clav. Open woods. Sept.
 425. *C. formosa*. *Pers.* Beautiful Clav. Aug., Oct.
 426. *C. crocea*. *P.* Saffron-yellow Clav. Decaying wood. May.
 427. *C. purpurea*. *Mull.* Purple Clav. Woods. Sept.
 428. *C. inæqualis*. *Mull.* Unequal Clav. Woods. Sept.
 429. *C. vermiculata*. *Scop.* White-tufted Clav. Aug., Sept.
 430. *C. fragilis*. *Holmsk.* Brittle Clav. Woods. Sept.
 431. *C. contorta*. *Fr.* Contorted Clav. Branches. Aug.

GEN. Calocera. *Fr.*

432. *C. glassoides*. *Fr.* Soft Calocera. Decayed stumps. Sept.

ORDER VI. Tremellini.

GEN. Tremella. *Fr.*

433. *T. fimbriata*. *Pers.* Fringed Tremella. Dead branches. June.
 434. *T. frondosa*. *Fr.* Large Pale Tremella. Ground. Aug., Oct.
 435. *T. foliacea*. *P.* Foliaceous Tremella. Stumps. Aug.
 436. *T. lutescens*. *Fr.* Yellowish Tremella. Old stumps. Aug., Sept.
 437. *T. mesenterica*. *Retz.* Orange Tremella. Sticks. Aug., Oct.

438. *T. vesicaria*. *Bull.* Bladdery Tremella. Ground. Aug., Sept.
 439. *T. albida*. *Hud.* Whitish Tremella. Logs. June, Aug.
 440. *T. intumescens*. *Sow.* Contorted Tremella. Wood. July, Sept.
 441. *T. indecorata*. *Somm.* Dingy Tremella. Dead willows. Aug.
 442. *T. tubercularia*. *Berk.* Horny Tremella. Branches. Sept., Oct.
 443. *T. torta*. *Willd.* Twisted Tremella. Oak. July, Sept.
 444. *T. epigæa*. *B. & Br.* Ground Tremella. Ground. Sept.

GEN. *Exidia*. *Fr.*

445. *E. glandulosa*. *Fr.* Witches'-Butter *Exidia*. Oak branches. Aug.

GEN. *Hirneola*. *Fr.*

446. *H. Auricula-Judae*. *Berk.* Jew's-ear *Hirneola*. Wood.

GEN. *Dacrymyces*. *Nees.*

447. *D. stillatus*. *Nees.* Orange *Dacrymyces*. Pine logs.

GEN. *Apyrenium*. *Fr.*

448. *A. lignatile*. *Fr.* Wood loving *Apyrenium*. Decayed wood.

HYPOGÆI.

FAMILY II *Gasteromycetts.*ORDER VII. *Hypogæi.*GEN. *Hymenogaster*. *Tul.*

449. *H. muticus*. *B. & Br.* Cracking *Hymenogaster*. Sept.
 450. *H. luteus*. *Vitt.* Yellow *Hymenogaster*. Woods. July, Sept.
 451. *H. decorus*. *Tul.* Comely *Hymenogaster*. Woods. Aug. Sept.

PHALLOIDEI.

ORDER VIII. *Phalloidei.*GEN. *Phallus*. *Linn.*

452. *P. impudicus*. *Linn.* Common Stink-horn. Sept., Oct.

TRICHOASTRES.

ORDER IX. *Trichogastres.*GEN. *Tulostoma*. *Pers.*

453. *T. mammosum*. *Fr.* Nippled *Tulostoma*. Ground.

GEN. Geaster. *Mich.*

454. *G. fornicatus*. *Fr.* Vaulting Geaster. Ground. Sept., Oct.
 455. *G. striatus*. *D. C.* Striate Geaster. On sand.
 456. *G. Bryantii*. *Berk.* Bryant's Geaster. Sandy soil.
 457. *G. hygrometricus*. *P.* Hard-coated Geaster. Ground. Woods. Oct.
 458. *G. lageniformis*. *Vitt.* Flask-like Geaster. Woods. Oct.

GEN. Bovista. *Dill.*

459. *B. nigrescens*. *P.* Blackish Bovista. Prairies. May.
 460. *B. plumbea*. *P.* Lead-colored Bovista. Prairies. Common.
 461. *B. ammophila*. *Lev.* Rooting Bovista. Woods. Sept.

GEN. Lycoperdon. *Tourn.*

462. *L. giganteum*. *Batsch.* Giant Puff-ball. Pastures. Oct.
 463. *L. cælatum*. *Fr.* Collapsing Puff-ball. Prairies. Aug., Sept.
 464. *L. atropurpureum*. *Vitt.* Purple-spored Puff-ball. Aug.
 465. *L. pusillum*. *Fr.* Little Puff-ball. Prairies. June, Sept.
 466. *L. saccatum*. *Vahl.* Elongated Puff-ball. Thickets. July.
 467. *L. gemmatum*. *Fr.* Warted Puff-ball. Meadows. Prairies. Aug.
 468. *L. pyriforme*. *Schæff.* Pear-shaped Puff Ball. Stumps. Sept.

GEN. Scleroderma. *P.*

469. *S. vulgare*. *Fr.* Common Scleroderma. Borders of woods. Aug.
 470. *S. bovista*. *Fr.* Thin-coated Scleroderma. Aug., Sept.
 471. *S. verrucosum*. *Pers.* Warty Scleroderma. Prairies. Aug.

GEN. Polysaccum. *D. C.*

472. *P. olivaceum*. *Fr.* Olive Polysaccum. Ground, woods. Aug.

ORDER X. Myxogastres.

GEN. Lycogala. *Mich.*

473. *L. epidendrum*. *Fr.* Stump Lycogala. July, Oct.

GEN. Reticularia. *Bull.*

474. *R. maxima*. *Fr.* Large Reticularia. Trunks, fallen trees.
 475. *R. atra*. *Fr.* Black Reticularia. Pine logs. Aug.
 476. *R. umbrina*. *Fr.* Umber Reticularia. Stumps. July.

GEN. Æthalium. *Link.*

477. *A. septicum*. *Fr.* Æthalium. Woods. Decaying wood.

GEN. Spumaria. *Fr.*

478. *S. alba*. *D. C.* White Spumaria. Living grass. June.

GEN. Ptychogaster. *Ca.*

479. *P. albus*. *Corda*. White Ptychogaster. On ground. July.

GEN. Diderma. *P.*

480. *D. farinaceum*. *Peck*. Invests fern stems in low woods.
 481. *D. Marlae-Wilsoni*. *Clinton*. Sticks, woods. Aug.
 482. *D. globosum*. *Fr.* Globose Diderma. Dead leaves. Sept.

GEN. Didymium. *Schrad.*

483. *D. melanopus*. *Fr.* Black-stemmed Didymium. Sticks. Aug.
 484. *D. connatum*. *Peck*. Decaying *Russula*. Sept.
 485. *D. furfuraceum*. *Fr.* Scurfy Did. Rotten wood. July, Aug.
 486. *D. squamulosum*. *A. & G.* Scaly Did. Dead leaves, &c. Aug.
 487. *D. farinaceum*. *Fr.* Mealy Did. Dead leaves. Aug.
 488. *D. pertusum*. *Berk.* Pierced Did. Stumps. Oct.

GEN. Physarum. *P.*

489. *P. pulcherripes*. *Peck*. Rotten wood. July.
 490. *P. cæspitosum*. *Peck*. Rotten wood. Aug.
 491. *P. atrum*. *Fr.* Black Physarum. Dead Branches. Aug.

GEN. Angioridium. *Grev.*

492. *A. sinuosum*. *Grev.* Twisted Angioridium. Sept.

GEN. Craterium. *Trent.*

493. *C. mutabile*. *Fr.* Changeable Craterium. Bark. July.

GEN. Diachæa. *Fr.*

494. *D. elegans*. *Fr.* Elegant Diachæa. Dead leaves. Aug.

GEN. Stemonitis. *Gled.*

495. *S. fusca*. *Rath.* Brown Stemonitis. Dead wood. June.
 496. *S. ferruginea*. *Ehnb.* Rusty Stemonitis. Dead wood. July.
 497. *S. ovata*. *P.* Ovate Stemonitis. Rotten wood. June.
 498. *S. obtusata*. *Fr.* Obtuse Stemonitis. Rotten wood. June.

GEN. *Arcyria*. *Hill*.

499. *A. nutans*. *Fr.* Nodding *Arcyria*. Rotten wood. June.

ORDER XI. *Nidulariacei*. *Tul.*GEN. *Polyangium*. *Link.*

500. *P. vitellinum* *Ditm.* Egg-yellow *Polyangium*. Stumps.

FAMILY III. *Coniomycetes*.ORDER XII. *Sphaeronemei*.GEN *Phoma*. *Fr.*

501. *P. ampelinum*. *B. & C.* Dead grape vines. Woods. July.
 502. *P. exiguum*. *Desm.* Little *Phoma*. Elder shoots. Aug.
 503. *P. glandicola*. *Lev.* Acorn *Phoma*. Acorns. Sept.

GEN. *Discella*. *B. & Br.*

504. *D. carbonacea*. *B. & Br.* Black *Discella*. Dead twigs.

ORDER XV. *Pucciniæi*.GEN. *Phragmidium*. *Link.*

505. *P. mucronatum*. *Link.* Rose Brand. Living Rose leaves. Aut.
 506. *P. gracile*. *Greve.* Raspberry Brand. Rasp. leaves. Aut.
 507. *P. obtusum*. *Link.* Strawberry Brand.
 508. *P. graminis*. *Pers.* Corn mildew. Leaves of corn. Aut.
 509. *P. striola*. *Link.* Sedge Mildew. Rushes. Autumn.
 510. *P. coronata*. *Corda.* Coronated Mildew. Grasses.
 511. *P. vaginalium*. *Link.* Knot-grass Brand. Aut.
 512. *P. primulae*. *Grev.* Primrose Brand. Primroses. June.
 513. *P. variabilis*. *Grev.* Variable Brand. Taraxacum. July.

GEN. *Gymnosporangium*. *D. C.*

514. *G. juniperi*. *Lk.* Living branches.

ORDER XVI. *Cœomacei*.GEN. *Tilletia*. *Tul.*

515. *T. caries*. *Tul.* Bunt. On wheat filling the grains. Aut.

GEN. *Ustilago*. *Link.*

516. *U. carbo*. *Tul.* Corn smut. Autumn.
 517. *U. antherarum*. *Fr.* Anther Smut. Lychens, &c.
 518. *U. violae*. *B. & Br.* Violet Smut. Violet leaves. August.
 519. *U. occulta*. *Preus.* Rye smut. On culms of rye.

GEN. *Uredo*. *Lev.*

520. *U. Quercus*. *Brand.* Oak-leaf Uredo. Sept.
 521. *U. bifrons*. *Grev.* Twin-faced Uredo. On *Rumex*. Aug., Sept.

ORDER XVII. *Æcidiacei*.GEN. *Æcidium*. *Pers.*

522. *A. euphorbiæ*. *Pers.* Spurge Cluster-Cups. May, June.
 523. *A. urticæ*. *D. C.* Nettle Cluster-Cups. June.

FAMILY IV. *Hyphomycetes*.ORDER XVIII. *Isariacei*.GEN. *Isaria*. *Fr.*

524. *I. arachnophila*. *Ditm.* Spider Isaria. Dead spiders.
 525. *I. citrina*. *P.* Lemon-colored Isaria. Decaying fungi. Aug.
 526. *I. intricata*. *Fr.* Intricate Isaria. Decaying fungi. Sept.

ORDER XIX. *Stilbacei*.GEN. *Tubercularia*. *Tode.*

527. *T. granulata*. *P.* Granulate Tubercularia. Dead branches.

DIVISION II. *Sporidiifera*.FAMILY VI. *Physomycetes*.ORDER XXIV. *Mucorini*.GEN. *Ascophora*. *Tode.*

528. *A. elegans*. *Corda.* Elegant Ascophora. Fowls' dung.

GEN. *Mucor*. *Mich.*

529. *M. ramosus*. *Bull.* Branched Mucor. Decaying fungi. Aug.
 530. *M. Mucedo*. *L.* Common Mucor. Decaying fruits.
 531. *M. caninus*. *P.* Dog's dung Mucor. Dung of dogs.

FAMILY VII. Ascomycetes.

ORDER XXVIII. Elvellacei.

GEN. Morchella. *Diel.*

532. *M. esculenta*. *Pers.* Common Morel. Woods. May, June.

GEN. Helvella. *Linn.*

533. *H. gigas*. *Kromb.* Large Helvella. Ground, woods. Spring.
 534. *H. crispa*. *Fr.* Pallid Helvella. Ground, woods. Early Summer.
 535. *H. sulcata*. *Afz.* Sulcate Helvella. Ground, woods. Oct.

GEN. Verpa. *Swartz.*

536. *V. digitaliformis*. *Pers.* Finger-shaped Verpa. Wood.

GEN. Spathularia. *P.*

537. *S. flavida*. *Prs.* Yellow Spathularia. Woods. July.

GEN. Leotia. *Hill.*

538. *L. lubrica*. *Pers.* Slimy Leotia. Woods. Sept.

GEN. Geoglossum. *P.*

539. *G. viride*. *P.* Green Geoglossum. Decayed wood.
 540. *G. glutinosum*. *P.* Glutinous Geoglossum. Woods.

GEN. Peziza. *Linn.*

541. *P. macropus*. *Pers.* Long-stemmed Peziza. Ground. July.
 542. *P. badia*. *P.* Large Brown Peziza. Pond margin. July.
 543. *P. aurantia*. *Fr.* Orange Ground Peziza. June.
 544. *P. lutea-nitens*. *B. & Br.* Bright Yellow Peziza. Ground.
 545. *P. fibrillosa*. *Curr.* Woolly Orange Peziza. Oct.
 546. *P. repanda*. *Wahl.* Spreading Peziza. Ground. June.
 547. *P. trachycarpa*. *Curr.* Rough-spored Peziza. Woods.
 548. *P. leiocarpa*. *Curr.* Smooth-spored Peziza. Ground.
 549. *P. cupularis*. *L.* Scolloped Peziza. June.
 550. *P. subhirsuta*. *Schum.* Hirsute Peziza. Ground.
 551. *P. humosa*. *Fr.* Ground Peziza. Woods.
 552. *P. scutellata*. *L.* Shield-like Peziza. Woods. May, Sept.
 553. *P. unicisa*. *Peck.* Ground in woods. Sept., Oct.
 554. *P. echinosperma*. *Peck.* Ground in woods. June.
 555. *P. rubra*. *Peck.* Burnt ground. June.
 556. *P. tiliae*. *Peck.* Dead branches. *Tilia Americana*. July.
 557. *P. coccinea*. *Jacq.* Carmine Peziza. Wood. Nov.

ORDER XXXI. Sphæriacei.

GEN. Valsa. *Fr.*

558. *V. pulchella. Fr.* Beautiful Valsella. Cherry and birch.

With three or four exceptions, the plants constituting the foregoing list of *Fungi*, have been found by the writer.

Respectfully contributed,

A. E. JOHNSON.

MINNEAPOLIS, January 1st, 1877.

VI.

ORNITHOLOGY.

REPORT OF DR. P. L. HATCH.

Prof. N. H. Winchell:

In accepting the position of Ornithologist on the Geological and Natural History Survey of the state to which I have been appointed by the Board of Regents, I desire to express to you and the Board my appreciation of the honor thus conferred upon me.

When first proposed to me to undertake this work, its objects were so accordant with my inclinations that I had only to harmonize its requirements with the exacting duties of my profession, to enter enthusiastically upon it. This our mutual arrangements have enabled me to do more satisfactorily than I had hoped. I therefore permit myself to expect to have something to report to you at the end of another year appropriate to an embodiment in the permanent records of your comprehensive work.

To do this I must be permitted to rely upon the co-operation of not only yourself and the other members of your staff, but of the Board, and all collectors temporarily or permanently associated with the survey.

I have had sufficient experience in the work before me to realize the necessity of aid from reliable sources, and this department of your survey has been so long delayed that it must be rigorously prosecuted if it shall keep its place alongside of the others in the years to come. I do not commence it, however, entirely *de novo*, having had some nineteen years observations in which I have accumulated some notes on birds in my vicinity, with occasional explorations into other sections as fully represented by them. The rapid settlement of the state has changed the relative representation of this class of its fauna very materially, and the increasing extent of the cultivation of the soil for varied productions is at present not only changing the aggregate numbers of birds that come here to breed, and those that permanently reside here, but

Bull. Minn. Acad. 1.325-144, 1878.

REPORT OF THE COMMITTEE ON BOTANY,
FOR 1877.

BY A. E. JOHNSON, M. D.

MR. PRESIDENT :—Since my last report, one year ago, I have collected and identified one hundred and two species of plants. Of the Fungi, I have collected ninety-five species new to our State, and fourteen species new to science, the descriptions of which I have endeavored to give fully. I have also collected quite a number of Algæ, seven of which I have been able to identify with the aid of the elaborate descriptions and fine colored plates of Prof. Wood on the "Fresh Water Algæ of the United States," published in the "Smithsonian Contributions to Knowledge, vol. xix."

The summer of 1877 was noted for extreme heat all over the world, and in our State, as one of extreme drought, there having been scarcely any rain during the whole of July and August. These conditions are very unfavorable to the production of fungi. Agarics of every genus were very scarce, not one was to be seen where thousands existed in previous seasons. Scarcely a *boletus* has been observed this season; only one or two *hydna*; a very few *polypori*; scarcely a *clavaria* or *tremella*; not a *phalloidei*, and but a very few *trichogastres* towards the close of the season; and so with nearly all other orders of fungi. Drought and extreme heat seem to effect the production of fungi more than almost any other plants.

558. AGARICUS (TRICHOLOMA) RUTILANS.

Schæff.

"Red-haired Tricholoma."

Our plant is not as large as the book description. Not necessarily on pine stumps. On decayed wood. Oct.

559. AGARICUS (LEPIOTA) HOLOSERICEUS. *Fr.*
 "Silky Lepiota."

On the ground. October. Scarce in 1877.

560. AGARICUS (TRICHOLOMA) ALBUS. *Fr.*
 "White Tricholoma."

Stipe, gills and pileus white; gills very broad, thin, and appear as if scooped out from the stem $\frac{1}{8}$ of an inch in width.

On the ground in woods. Ramsey county. October.

561. AGARICUS (CLITOCYBE) ELIXUS. *Sow.*
 "Sodden Clitocybe."

On the ground. Nicollet Island. September and October. Very scarce in 1877.

562. AGARICUS (CLITOCYBE) MAXIMUS. *Fr.*
 "Sowerb's Clitocybe."

Pileus fleshy, thick, firm, scaly, disk compact; not umbonate; stem compact, attenuated downwards; slightly fibrillose; gills largely decurrent, when old nearly blood red.

Pileus 7 inches broad, with an ocraceous tinge; stipe, 3 inches high, 2 inches thick at top, one inch at base; strong cadaveric odor when old. In thin oak woods.

October. Rare.

563. AGARICUS (PLEUROTUS) SALIGNUS. *Fr.*
 "Willow Pleurotus."

On trunks of trees and stumps. October.

564. AGARICUS (COLLYBIA) FAMILIA. *Peck.*

Height about 2 inches; breadth of pileus $\frac{3}{4}$ to $1\frac{1}{2}$ inches; stipe 1 to 2 lines thick. On decayed wood in woods May and September.

565. AGARICUS (COLLYBIA) COLOREUS. *Peck.*

Stem in our plant is pale, brownish yellow below, and lightish or white above.

On decaying basswood stumps. • June and September, and October.

566. AGARICUS (MYCENA) IRIS. *Berk.*

"Blue Mycena."

On bits of decayed wood, beside an old oak stump in oak thickets. October. Rare.

567. AGARICUS (MYCENA) PROLIFEROUS. *Sow.*

"Proliferous Mycena."

When matured, the cuticle of the pileus frequently contracts towards the center, producing a cup shape, and sometimes contracts to a globe shape, showing nothing but the gills turned up over the pileus.

As far as observed our plants are gregarious or solitary, not cæspitose. On rich soil amongst grass in open woods. Nicollet Island. June. Not common.

568. AGARICUS (MYCENA) VITREUS. *Fr.*

"Glassy Mycena."

Our plant is not very fragile. Stem nearly 4 inches long, nearly one line thick, and very brittle.

In woods amongst moss. Often attached by mycelium. October.

569. AGARICUS (MYCENA) TENUIS. *Bolt.*

"Slender Mycena."

Pileus and stem brownish; stem 1 to 2 inches long, not a line thick,

Amongst moss on the ground in damp woods. June. Scarce.

570. AGARICUS (MYCENA) VITILIS. *Fr.*
 "Flexile Mycena."
 Amongst grass in woods. June.
571. AGARICUS (MYCENA) SPEIREUS, *Fr.*
 "Opaque Mycena,"
 Amongst moss on sandy ground. Solitary. May.
572. AGARICUS (MYCENA) VULGARIS. *P.*
 "Common Mycena."
 On the ground and decaying wood. October.
573. AGARICUS (MYCENA) SETOSUS, *Sow.*
 "Setose Mycena."
 White or a whitish gray; stem fine as a hair; under a glass, somewhat pubescent, which distinguishes it from *Agarcus capillaris*. Often beneath leaves. Easily overlooked, because of its excessive minuteness. Pileus 1-20 to 1-16 of an inch broad; stem 1 to 1½ inches high.
 On and under dead leaves in woods. October.
574. AGARICUS (OMPHALIA) BELLÆ. *Fohnst.*
 "Cup-like Omphalia."
 In woods, on decaying wood. Sept. and Oct.
575. AGARICUS (PLUTEUS) UMBROSUS. *Pers.*
 "Brown Pluteus,"
 On the ground. Sept.
576. AGARICUS (ENTOLOMA) PLACENTA. *Batsch.*
 "Brown Entoloma."
 On the ground in woods. October.

577. AGARICUS (ENTOLOMA) FLAVUS. *n. sp.*

Gold-yellow Entoloma.

Pileus, fleshy, conical, expanded, depressed, smooth, bright orange when fresh, flesh white; stem short, solid, slightly attenuated upward, white; gills bright gold color when fresh, broad externally, regular, thin, sharp edged, narrow at the stipe, slightly attached.

Pileus, 2 inches broad; stipe $\frac{1}{2}$ inch high, 4 lines thick. Spores irregular, large, spherical, papilous and papilate; bright reddish yellow.

In woods on the ground. Last of October. Scarce. The spores lose their reddish cast after long exposure to the air and light.

578. AGARICUS (PHOLIOTA) AURIVELLUS. *Batsch.*

"Golden Pholiota."

On fallen basswood trunks in woods. Excelsior. July.

579. AGARICUS (PHOLIOTA) SPECTABILIS. *Fr.*

"Orange Pholioto."

Subcæspitose. Pileus firm, brittle, convex rather than plain or depressed, dry, cuticle torn into scales, fibrous or silky, sometimes patchy points turning up and darker, giving the surface a tigrine appearance; stem first solid then hollow, fibrous, striated, equal, rooting, curved; gills adnate, ventricose, entire, divides easily into two parts, and wavy on the free surfaces, crowded, rounded behind, ferruginous.

Pileus, 3 to 4 inches broad, fleshy and bright, rhubarb color; stem 3 inches long, $\frac{1}{2}$ inch thick, annulated. Taste pleasant, not bitter.

On a decayed spot on a living locust. June. Scarce.

580. AGARICUS (GALERA) OVALIS.

Fr.

"Brick-red Galera."

Stipe, 3 to 5 inches high; pileus 1 inch and more high.

On the ground amongst grass in open woods. September to October.

581. AGARICUS (STROPHARIA) SQUAMOSUS.

Fr.

"Scaly Stropharia."

Pileus, ocraceous or reddish, viscid, spotted with yellowish scales, nearly plain, not very thin, quite fleshy; stem slender, stuffed with a whitish pith, more or less squamose below the distant, not well defined annulus; gills purplish brown with whitish edge, very broad, adnate.

Pileus from $\frac{3}{4}$ to $1\frac{1}{2}$ inches broad; stipe 2 to 3 inches long, nearly 2 lines thick, hollow when old.

On the ground, attached to decaying wood, in woods. October.

582. AGARICUS (HYPHOLOMA) CAPNOIDES.

Fr.

"Fir Wood Hypholoma."

Pileus usually yellowish, nearly plain, smooth like kid leather, dry, rather fleshy; stem smooth, pale yellow, even, sub-festulose; gills purplish or grayish, dry, not crowded, broad, adnate, very various in size.

Pileus, from 1 to 3 inches or more broad; stem 2 to 3 inches long, 2 to 4 lines thick, equal, sometimes curved.

In Lake Harriet woods. Not necessarily in fir woods. May. Not common. 1877.

583. AGARICUS (HYPHOLOMA) APPENDICULATUS. *Bull.*

"Appendiculate Hypholoma."

Pileus $1\frac{1}{2}$ to 3 inches broad, veil attached to the margin in patches; stem 2 to 3 inches long, 2 to 3 lines thick.

Amongst moss on rotten stumps, in woods. June.

584. AGARICUS (HYPHOLOMA) EGENULUS. *Berk.*
 "White Hypholoma."
 By the road side, amongst grass. May.

585. AGARICUS (PSILOCYBE) AREOLATUS. *Klotsch.*
 "Patchy Psilocybe."
 Nicollet Island. September to October.

586. AGARICUS (PSATHYRA) BIFRONS. *Berk.*
 "Changing Psathyra."
 So called from its many changes of color.
 Nicollet Island. September to October.

587. AGARICUS (PSATHYRA) SEMIVESTITUS. *Berk & Br.*
 "Half-clothed Psathyra."
 Pileus $\frac{1}{4}$ to $\frac{1}{2}$ inch broad; stipe 1 to 1 $\frac{1}{2}$ inches long.
 Nicollet Island. October.

588. AGARICUS (PSATHYRA) GORDONI. *Berk.*
 "Gordon's Psathyra."
 Amongst grass in woods. September. October.

COPRINUS.

589. COPRINUS FLOCCULOSUS. *D. C.*
 "Flocculose Coprinus."
 Nicollet Island. September.

590. CAPRINUS EXTINGTORIUS. *Fr.*
 "Extinguisher Coprinus."
 Stem more or less rooting. Sometimes the root is very long.

On dung heaps and rich, soft loam. June to July.

591. *COPRINUS ARATUS.*

Berk & Br.

"Umber Coprinus."

Our plant is gregarious; pileus 1 to 4 inches broad; stipe 3 to 5 inches high, 2 to 6 lines thick, fibrous.

On the top of a decaying elm stump. Lake Harriet woods. May.

592. *COPRINUS RADIANS.*

Fr.

"Radiating Coprinus."

Pileus 1 to 2 inches broad; stem $1\frac{1}{2}$ to 2 inches high.

On sandy soil. May. Very scarce. 1877.

593. *COPRINUS MACROCEPHALUS.*

Berk.

"Stunted Coprinus."

Pileus 1 inch or less broad, cylindrico-campanulate; stipe $\frac{1}{2}$ to 2 inches high, 2 to 3 lines thick.

Solitary or gregarious. On stumps and horse manure. April.

594. *COPRINUS FILIFORMIS.*

Berk & Br.

"Filiform Coprinus."

Stipe $\frac{1}{2}$ to 1 inch high; pileus 1 to 2 lines high. Conical.

In woods on the ground. September to October.

CORTINARIUS.

595. *CORTINARIUS (PHLEGMACIUM) DIBAPHUS.*

Fr.

"Spotted Cortinarius."

Stem winding; spores kidney shaped with an apiculus at both ends.

On the ground in woods. October.

596. *CORTINARIUS (PHLEGMACIUM) FULGENS.* *Fr.*

"Shining Cortinarius."

Very fleshy; flesh white; stipe solid, bulb depressed, marginate. The whole stem wooly, which becomes ferruginous when the spores drop.

In woods. October. Not common.

597. *CORTINARIUS (MYXACIUM) ELATIOR.* *Fr.*

"Tall Cortinarius."

The gills in our plant are not only venose connected, but strongly costate, every alternate one being dimidate and excessively rugose. The characters seem almost sufficient to base a new species on.

In woods. October. Not common. 1877.

598. *CORTINARIUS (MYXACIUM) FACICULARIS.* *n. sp.*

Tufted Cortinarius.

Cæspitose, gregarious or solitary; pileus plain, fleshy, viscid or glutinous when moist, smooth when dry, reddish yellow, edge thin and dotted with a few fragments of the veil; gills cinnamon, crowded, broadly attached, sometimes sub-decurrent, easily separating from the hymenophora; stipe white, solid, fibrous, straight or flexurous, attenuated downward. When old the flesh becomes yellowish white, the stem brownish.

Pileus $\frac{1}{2}$ to 2 inches broad; stipe 1 to 2 inches long, 1 to 3 lines thick.

On the ground in open places. October.

599. CORTINARIUS (MYXACIUM) MACULATUS. *n. sp.*
 "Egg-yellow Cortinarius."

Pileus thin, membranaceous, soft, thin except the disk which is thick, firm, broadly umbonate, smooth, shining, excessively viscid, bright egg yellow, dotted here and there with iron rust colored spots, cuticle tough, with a kid leather feel when dry; gills attached with a tooth, at first whitish, then yellow, then dark cinnamon brown, sharp, smooth edges; stipe, equal, stout, not annulate or bulbous, whitish, fibrilose, more or less curved.

Breadth of pileus, 2 to 5 inches; stipe 3 to 5 inches high, 2 to 6 lines thick.

In woods on decaying *Acer* stumps. Nicollet Island. October.

Spores large, regular spheres, appearing frequently nucleated.

600. CORTINARIUS (MYXACIUM) RADICIBUS. *n. sp.*
 Rooting Cortinarius.

Pileus sub-viscid when moist, polished when dry, shining, fleshy, convex, then expanded; stem solid, fibrous, cylindrical, smooth, straight or crooked, annulated, annulus $\frac{1}{4}$ inch from top of stipe standing up, upper surface color of the gills, base sending out long, white, fibrous roots; gills broad, sub-decurrent, lightish or gray, then cinnamon.

Pileus $\frac{3}{4}$ to $1\frac{1}{2}$ inches broad; stem 2 to 3 inches high, 2 to 3 lines thick.

Spores reddish ochre, or Ferri Carb. color, very small, egg shape.

On the ground in woods. May.

601. CORTINARIUS (DERMOCYBE) OCHROLEUCUS. *Fr.*
 "Pallid Cortinarius."

Pileus not very fleshy; stipe sometimes festulose at the top.

In woods. June. Not common. 1877.

602. CORTINARIUS (DERMOCYBE) CINNAMOMEUS. *Fr.*

"Cinnamon Cortinarius."

In thin woods on mucky soil. May.

603. CORTINARIUS (DERMOCYBE) APPENDICULATUS. *n. Sp.*

Bank-loving Cortinarius.

Pileus fleshy, thin, smooth, expanded, border appendiculate with a fugaceous veil, yellowish brown, becoming nearly black when dry; gills adherent, sub-decurrent, crowded, brown, then black; stem equal, smooth, festulose, whitish, becoming brown when dry.

Pileus 1 inch broad; stipe 1 to 2 inches high, about 2 lines thick.

Spores small, smooth, nearly spherical.

On low banks in woods. May.

604. CORTINARIUS (DERMOCYBE) TIGRINUS. *n. sp.*

Tiger Spotted Cortinarius.

Pileus thin but fleshy, convex, spotted with small, brown scales when fresh, rugose and brown when dry; gills numerous, crowded, adnate, pale clay when young, becoming cinnamon, then dark brown; stem whitish, attenuated upwards, sub-bulbose.

Pileus $\frac{3}{4}$ to 1 inch broad; stipe one inch or more high, one line thick.

On decaying wood in woods. June. Not common. 1877.

605. CORTINARIUS (TELAMONIA) EVERNIUS.

Fr.

"Tufted Cortinarius."

About stumps in open places. September and October.

606. CORTINARIUS (HYGROCYBE) VENOSUS.

n. sp.

Venous-Gilled Cortinarius.

Pileus smooth, thin, yellowish brown, when mature so completely everted that the gill surface only appears; gills cinnamon, sub-ventricose, rather narrow, attached, profusely venosed and venose connected; stipe smooth, equal, fibrous, not bulbous, sometimes split or forked at the top.

Pileus 1 to 2 inches broad; stipe, 3 inches long, 3 lines thick.

On the ground amongst poplar leaves in woods. October. Scarce.

607. HYGROPHORUS UNGUINOSUS.

Fr.

"Dingy Hygrophorus."

Gills adnate, decurrent, connected by veins.

On the ground in woods. October.

608. HYGROPHORUS LACTUS.

n. sp.

Grayish-white Hygrophorus.

Pileus hemispherical, irregular, grayish white, cuticle covered more or less with a milk white gluten, flesh thick, white; gills white, arcuate, adnate, decurrent, edges smooth, entire, sharp; stipe smooth, fibrous, white, mealy above, sharply attenuated downward.

Pileus 1 to 1½ inches broad; stipe 2 to 3 inches high, 3 to 5 lines thick.

On wet clay soil amongst dead poplar leaves. October. Rare.

609. GOMPHIDIUS VISCIDUS.

Fr.

"Viscid Gomphidius."

On the top of a decaying hickory stump. Meeker woods
August.

LACTARIUS. Fr.

610. LACTARIUS SACCHARIUM.

n. sp.

Sweet-milk Lactarius.

Pileus at first convex, papilate, pruinose, brownish, or slate, at length expanded, depressed, infundibuliform, reddish orange; gills largely decurrent, crowded, narrow, sinuate, dimidate or entire, pale orange, becoming brownish when old; stem irregular, compressed, curved or straight, orange.

Pileus and upper part of stipe milky, gills not milky.

Pileus $\frac{1}{2}$ to $1\frac{1}{2}$ inches broad, stem 1 to 2 inches high, 1 to 3 lines thick.

Plant gregarious or cæspitose. Milk white. Slightly sweet.

Amongst moss and grass, beneath poplar trees. July and August.

611. MARASMIUS CAULICINALIS.

Fr.

"Mealy Stemmed Marasmius."

Scarcely striate. Pileus 1 to 2 inches broad; stipe $1\frac{1}{2}$ to 2 inches high.

On the ground in woods. October.

612. MARASMIUS SULPHUREUS.

n. sp.

Sulphur Colored Marasmius.

Pileus, when matured, reddish brown, expanded, border rugose, umbo darkest; gills numerous, entire, dimidate, adnate, sub-decurrent, venose connected, pale orange; stem sulphur color, tomentose, equal, festulose, curved, base bulbous.

613. *MARASMIUS SEMIHIRTIPES.**Peck.*

"Pale Brown Marasmius."

On the ground in woods. June. Not common.

*LENTINUS. Fr.*614. *LENTINUS MAXIMUS.**n. sp.*

Ringed Lentinus,

Pileus thick, compact, coreaceous, convex, plain, unequal, pale straw-color surface, dotted with dark scaly spots; stem stout, hard, rooting, scaly, annulate; gills $\frac{3}{4}$ to 1 inch broad, sinuate, decurrent, forming lines on the stem down to the ring, whitish, with torn or tattered borders.

Pileus 3 to 7 inches broad, stem 1 to 2 inches long, an inch and more thick.

On the square end of a tamarack log. Aug.

*LENZITES.**Fr.*615. *LENZITES FLACCIDA. Fr.*

"Flaccid Lenzites."

On stumps of nearly all kinds of wood.

*POLYPORUS.**Fr.*616. *POLYPORUS SQUAMOSUS. Fr.*

"Scaly Polyporus."

On oak trunks and fallen branches. Oct.

A difficult species to determine.

617. *POLYPORUS VARIUS. Fr.*

"Variable Polyporus."

On fallen trunks and branches. In swamps.

618. *POLYPORUS TOMENTOSO-QUERCINUS.**n. sp.*

Woolly Oak-Polyporus.

Pileus at first, soft, compact, spongy, tomentose, pulvinate, dimidate, sessile, very thick, divergently fibrous within, broad surface of attachment, dirty grayish white when

young, pale straw or sub-ferruginous when old, hard, coreaceous, woody at maturity; pores large, irregular, toothed or fringed, easily separated, from $\frac{3}{4}$ to 1 inch long, varying in color from straw to bright orange.

Nearly always on the north side of living oaks. Pileus 1 to 2 inches thick, 2 to 5 inches broad. Spores numerous, white, globose, very small. Drops its spores in May or early June. Plant is persistent, lasting the whole year.

Its characters are decided and unmistakable when once observed. Very scarce, only seen occasionally.

619. POLYPORUS INTYBACEUS.

Fr.

"Gray-branched polyporus."

Our plant has an anise-like odor, and on the upper surface of some of the pilei there are pores developed in small patches. In all other respects fully answers book descriptions.

On an old oak stump close to the ground. In woods September.

TREMELLA *Fr.*

620. TREMELLA EPIGAEA. *B. & Br.*

"Ground Tremella."

Effused, white, gelatinous within, externally dusted with the white spores.

On the ground, spreading over leaves, sticks, &c. Lake Harriet woods. June and July.

DIDERMA *P.*

621. DIDERMA FLORIFORME. *P.*

"Flower-like Diderma."

At first whitish, becoming milk-white, when full grown shining and fading through all shades to a yellowish brown; stem slender, peridium spherical, splitting in all shapes, and sometimes throwing off a complete cap.

Spores yellowish brown, and may be seen under a glass

thrown out with much force so as to fall some distance away.

On decaying trunks of trees. Oct.

ASPERGILLUS. *Mich.*

622. *ASPERGILLUS GLAUCUS.* *Lk.*

"Blue Mould."

On decaying leather and other decomposing substances.
Sept.—Oct.

623. *ASPERGILLUS VIREUS.* *Lk.*

"Green-branched aspergillus."

On decaying *Agarics* and *Boleti* Oct.

HAPLARIA. *Link.*

624. *HAPLARIA GRISEA.* *Lk.*

"Gray Haplaria."

On various decaying plants. Sept.—Oct.

PILOBOLUS. *Tode.*

- 625 *PILOBOLUS CRYSTALLINUS.* *Tate.*

"Crystalline Pilobolus."

At first appearing like a small, yellow *Sclerotium*, soon forming a stem, which becomes inflated at the apex, which is terminated by a black and somewhat spherical cap; at this period of development the yellow hue disappears.

Densely tufted; very short lived:

On horse dung. July.

UNCINULA. *Lev.*

626. *UNCINULA ADUNCA.* *Lev.*

"Willow Blight."

On leaves of willows. Sept. Very abundant.

MICROSPHÆRIA. *Lev.*

627. *MICROSPHÆRA EXTENSA.* *C. & Pt*

On upper surface of oak leaves. Sept. Common.

MORCHELLA. *Dill.*

628. **MORCHELLA CRASSIPES.**

Pers.

"Gigantic Morel."

Our plant is not so tall as the book descriptions, which make 9 inches or more. Ours is 6 or 7 inches high.

In woods. May. Esculent.

629. **MORCHELLA PATULA.**

Pers.

"Spreading Morel."

Obtusely conic, pits irregular, stem $1\frac{1}{2}$ to 2 inches high, even, sometimes with an undulated exterior, thick, hollow.

In woods. May. Not common.

GEOGLOSSUM.

P.

630. **GEOGLOSSUM ALBUS.**

n. sp.

White Geoglossum.

Fasciculate or solitary, smooth; clubs cylindrical, confluent with the stem; stem abruptly black, shading to greenish white; hymenium smooth, white; spores white, oblong, $\frac{3}{80}$ of an inch long.

Plant $\frac{1}{2}$ to 2 inches high. On ground and decayed wood, in woods. May.

PEZIZA. *Linn.*

631. **PEZIZA RADULA.**

B. & Br.

"Black warted Peziza."

Cup depressed, sessile, $\frac{1}{2}$ to 1 inch in breadth, dark brown or black, covered with small warts on its upper surface, amidst which the asci are seen. Very brittle. Beneath silver maple trees on my lawn. July.

632. **PEZIZA ARGILLACEA.**

Sow.

"Clay Peziza."

Our plant is dusky, or brownish, or slightly yellow within the cup, dirty white externally.

Varies in size from $\frac{3}{4}$ to 2 inches in breadth. On clay in Marr's woods. May.

633. PEZIZA PULLULUS.

n. sp.

Blackish-brown *Peziza*.

Thin, the cup infundibuliform, bright brown or black, not turning pale, smooth, border incurved, crenate, extremely smooth, black; stem costate, flat, festulose; cup 1 to $1\frac{1}{2}$ inches broad, 1 to $1\frac{1}{2}$ inches deep; stem 1 to $1\frac{1}{2}$ inches high; whole plant 2 to 3 inches high.

Solitary or gregarious. On the ground or partly buried sticks.

Spores oblong, about 1-1000 by 1-500 of an inch, smooth, nucleated. In woods. May.

634. PEZIZA SALMONICOLOR.

B. & Br.

"Salmon-Color *Peziza*."

One inch broad, gregarious; cups hemispherical, pale salmon-color; asci 4 to 6 times as long as thick; sporidia, nucleate, elliptic, biseriate.

On burnt ground, in woods. Oct.

635. PEZIZA CILIARIS.

Schrad.

"Fringed white *Peziza*."

On decayed wood, leaves and bark; in woods. May.

636. PEZIZA NIVEA.

Fr.

"Snowy *Peziza*."

On decayed and partly buried wood; in woods. May.

637. PEZIZA VIRGINEA.

Batsch.

"Virgin White *Peziza*."

Stipitate; under a glass of 50 diameters glistening white, hemispherical, externally densely covered with patulous hairs. Cup about 8 lines broad under a magnifying glass of 50. Margin ciliated, often covered with little drops of water. Spores fusiform, .0002 of an inch long and half as broad.

On the under side of half-buried rotten logs, in woods.
May.

638. PEZIZA BICOLOR.

Bull.

"Two-colored *Peziza*."

Subsessile, nearly spherical, heavily tomentose, externally white; disc little tinged with yellow, very small, with a glass of 50 appears about 6 lines broad; stipe about 8 lines high, attenuated downwards.

On decayed twigs, buried beneath rotten logs, in woods.
May.

639. PEZIZA DIPLOCARPA.

Curr.

"Double fruited *Peziza*."

Our plant is pale orange or brown, not greenish olive; gregarious, perfectly covering, sometimes, large patches of earth.

On recently burned ground, Highland Park. Oct.

CALYCELLA. *Fries.*

640. HELOTIUM TUBA.

Fr.

"Tube *Helotium*."

Plant $\frac{1}{2}$ inch high; on twigs on the ground. July to September.

FRESH-WATER ALGÆ.

GENUS DOLICHOSPERMUM. *Thwaites.*

1. D. SUBRIGIDUM.

Wood.

This species I have found growing in the scum floating on water in ditches and in the coffer dam on Nicollet Island.

GENUS GLOIOTRICHIA, *J. Ag. 1842.*

2. G. INCRUSTATA.

Wood.

Found attached to small water plants along the Hennepin Island shore of the east branch of the Mississippi River.

GENUS **ZONOTRICA**.

3. **Z. MOLLIS.** *Wood.*
Found attached to the rocks at Minnehaha Falls.

GENUS **DASYACTIS.** *Ktz.*

4. **D. MOLLIS.** *Wood.*
Found in a bog north of Sandy Lake.

GENUS **PALMELLA.**

5. **P. DURA.** *Wood.*
This is in form of little blackish balls varying in size from $\frac{1}{4}$ of a line to 1 inch in diameter, firm and hard, the largest appearing as if they were aggregations of multitudes of small ones. Found in slow moving water in the coffer-dam on Nicollet Island.

GENUS **PENIUM.** *Preb. 1848.*

6. **P. DIGITUS, (Ehrb).** *Preb.*
Found in small lakes and ponds.

GENUS **VAUCHERIA.**

7. **V. VELUTINA.** *Ag.*
Found growing on mud in shallow ditches, after the subsidence of water from a heavy shower. Sept.

REPORT ON ORNITHOLOGY FOR 1877.

In some respects the season has been rather unfavorable to the observation of the birds; more especially as regards their migrations. The year previous was exceptionably characterized by a great tidal wave of bird-life that reached us here in its vernal flow about the 10th to the 15th of May. It was emphatically an ethereal wave of song as it was constituted of the typical warblers. Although less so, yet to a considerable extent, a similar characteristic was manifest through the entire migration of the other groups, determining the measure of collections only by the possibility of taking adequate care of the material obtained. Widely different from this, the birds of the past year have arrived in such inconsiderable number, at or about any one period, that it has required the utmost vigilance to note their presence, and then only as stragglers, stealing their way to their secluded places for resting and rearing their young. It is presumable that the diversity between the two seasons has been principally determined by the contrast in their meteorological characteristics, as affecting the conditions of transit, supply of food, &c. It is to be acknowledged that with all that has been learned of the birds, very little is settled as to the causes and modifications of their migrations. More has been written, perhaps, during the year, upon this matter, than upon any previous one in the history of ornithology.

Of course, the additions to our list of newly recognized species have been very meagre, consisting of Leconte's Sparrow—*Coturnicuius lecontei* (Bor.), collected by Mr. Williams near the city, I think, and the Buff-breasted Sandpiper, *Tryngites rufescens*, (Cob.), collected by the Messrs. Roberts at Sand Lake, near the northern limits of the East Division.

The former, originally discovered and named by Audubon *Emberiza leconti* has long been held doubtful as a distinct species, having been supposed to be either *passcrinus* or *henslowii*.

Numerous collections, made in northern Dakota by Dr. Cones, when associated with the Northern Boundary Survey in 1873. and by others at other localities, have determined its habitation to be the central plains from northern Dakota to Texas. Its presence here shows our formal relationship to the province that lies to the west of us, and together with other species previously obtained, puts us on the alert for exceptional forms hitherto restricted to areas beyond our borders.

The Sandpiper obtained by the Messrs. Roberts has been sought for through many long years without success. Its known breeding places far to the north of us and the sparcity of its numbers in migration along the Atlantic coast led to the anticipation that it would long since have been observed in migration through the state.

We are at last rewarded through the vigilance of these gentlemen who have contributed so much to our already large list, embracing about as many as have been identified in New England, after a century's observation by a legion of experts.

Although offering so little in this Annual Report that is new, the year has not been an idle one in obtaining an extension of our knowledge of the habits and characteristics of species that breed in our district. In this, perhaps, no year preceding has given us nearly so rich results, the proof of which will become evident in the future reports involving life-histories of local species.

THE CRETACEOUS IN MINNESOTA.

BY N. H. WINCHELL.

There are frequent exposures of the Cretaceous along the Minnesota river from Big Stone Lake to Mankato; and at several places lignitic coal occurs in beds varying from six inches to three or four feet in thickness. It is found on the Cottonwood river, a large tributary of the Minnesota from the south, and on the Blue Earth. There are evidences of it under the drift at points between Mankato and Mendota. It occurs in Steele county, also in Freeborn, Fillmore and Mower. There are indirect evidences of it in Dakota county, particularly at Empire City. It is scattered largely through the drift in the western part of Hennepin county. In the central and eastern parts of Wright county it has excited expectations of coal, and some shafting has been done. Pieces of lignite coal have been found along the Mississippi river as far north as Fort Ripley; and in Stearns county a considerable interest was excited a few years ago by exposures of lignite in the valley of the Sauk river, at several places, particularly at Richmond. It is met with in Faribault county in sinking deep wells, also in Kandiyohi. It is distributed abundantly through the drift in the valley of the Red river of the North, both in the form of "slate" and lignitic fragments. Pieces of lignite are said to have been found on the St. Louis river, near the mouth of Swan river. According to H. R. Schoolcraft, lignitic coal was discovered in the country south of the Lake of the Woods in 1858 or 1859. J. A. Wheelock refers to this in the "First Annual Report of the Commissioner of Statistics," Jan. 1st, 1860. In the summer of 1875, Cretaceous lignite was discovered in the country between Rainy Lake and Vermillion Lake, and a hundred pounds of what

was taken for coal were brought to Duluth from that point. During the last Summer (1878) further information has been obtained of lignite along the international boundary line on Nemakon Lake, east of Rainy Lake.

It thus appears that fragments of the Cretaceous, which cannot bear long transportation, are found throughout nearly the whole State, and that the beds *in situ* have been discovered in a sufficient number of places to enable us to conclude that they extend, or did prior to the spreading of the drift, from the Iowa state line to that of the British Possessions. The only portions of the State where these indications do not exist, are the triangle north of Lake Superior, and the so-called driftless area in the southeastern corner of the State. These are both many hundred feet lower than the central and western portions where the Cretaceous is known to exist, and must have been buried under the Cretaceous ocean unless by some great warping of the crust they were relatively much higher then than now. There is no evidence of such warping; hence it is more likely that unfavorable circumstances have prevented the preservation of the beds or concealed them, so as to delay their discovery.

A line drawn from the west end of Hunter's Island, on the Canadian boundary line, southward to Minneapolis, and thence southeastwardly through Rochester, to the Iowa state line would, in general, separate that part of the State in which the Cretaceous is not known to exist from that in which it does. It is not here intended to convey the idea that the whole State west of this line, is spread over with the Cretaceous, because there are many places where the drift lies directly on the Silurian or earlier rocks; but throughout this part of the State, the Cretaceous exists at least in patches, and perhaps once existed continuously.

The beds of this formation that occur in Minnesota, belong to the lower portion. The Dakota Group of Dr. F. V. Hayden, supposed to be the equivalent of the Nishnabotany sandstone of Iowa, is probably represented by the white sandstones at Austin and on the Blue Earth, and by the stone

quarried at Austin and at points below in the banks of the Cedar river, as well as by the sandstone seen at Fritz' quarry on the Minnesota, a few miles below New Ulm. There is a pebbly deposit in Fillmore county, associated with a white sandstone, that also probably belongs to the Dakota Group. The Fort Benton Group has been identified in the Sauk river valley by a few fossils found in explorations for coal a few years ago. This rests on the authority of F. B. Meek. There are also extensive clay and shale beds exposed along the Minnesota, and on the Cottonwood, embracing beds of lignite that doubtless belong to the Fort Benton. At New Ulm is a calcareous member, burned for quicklime, that probably represents the Niobrara.

Wherever this formation has been explored for coal in Minnesota, the results have not been encouraging. The lignite varies from a cannel coal to charcoal, and is often injured by clayey impurities, and by the presence of what appears to be ashes. The difficulty and cost of mining are also increased by the non-indurated condition of the overlying layers, rendering it necessary to line by timbers, or otherwise support the upper part of every opening that is made. Some of the clays have been successfully employed in making pottery, and also in the manufacture of fire-brick.

To the drift the Cretaceous has contributed largely in the State of Minnesota. Its *debris* is distinguishable in nearly every county west of the above line. It is the source of much of the clayey material of the boulder clay, and its alkaline qualities are very perceptible in much of the surface water. Superficially the Cretaceous has produced a greater effect on Minnesota than any other geological formation; yet its beds *in situ* are less frequently seen than any others.

EXCHANGES RECEIVED DURING THE YEAR 1877.

Transactions of the Academy of Science, St. Louis, Vol. III, Nos. 1 and 2.

Transactions of the Academy of Science and Arts, New Haven, Vol III, Part 1.

Bulletins of the American Geographical Society, New York, Nos. 4 and 5.

Bulletins of the Buffalo Society of Natural Sciences, Vol. I, No. 3, Vol. II, No. 3.

Fourth Annual Report of the Zoological Society of Philadelphia.

Fifth Annual Report of the Zoological Society of Philadelphia.

Catalogue of Paintings from the Historical Society of Pennsylvania.

Constitution and By-Laws of the Antiquarian Society of Philadelphia.

Circular of the Vermont Historical Society, Montpelier.

Smithsonian Contributions to Knowledge, 50, 172, 192, 197.

Geology of the Uintah Mountains, with Atlas, by Powell.

Bulletins of the U. S. National Museum, Nos. 6, 7, 8, 9.

Bulletins of the U. S. Entomological Commission, Nos. 1 and 2.

Animal Resources of the United States, G. B. Gove, 1875-6.

North American Mustelidæ Coues.

North American Rodentia Coues.

North American Ethnology, Dall & Gibbs.

Annual Report of Geological Survey of Canada, 1874-5.

Descriptive Catalogue of Minerals, &c., Geological Survey of Canada.

Proceedings of the Liverpool Geological Society, 1875-6.

Report of the Zoological Society of London, 8168, 1873—6 numbers.

Constitution and By-Laws of the Zoological Society of London.

Philosophical Transactions of the Royal Geological Society of Ireland Vol. 165, Part 2.

Report of the Ohio State Board of Agriculture, 1874.

Proceedings of the Nova Scotia Institute of Natural Sciences, 1870-1.

Proceedings of the Nova Scotia Institute of Natural Sciences, 1871-2.

Proceedings of the Nova Scotia Institute of Natural Sciences, 1872-3.

Proceedings of the Nova Scotia Institute of Natural Sciences, 1873-4.

Proceedings of the Nova Scotia Institute of Natural Sciences, 1877-8.

Boletin de Sociedad Mexicana de Geografia Y. Estadistico.

Biografica Ann. Institutio Historico Geographico, E. Ethnographico, Rio de Janeiro, de Brazil S. A., Vols. II and III.

Informa de los Esploradoras del Territorios de San Martin, 1871.

Cataloga del Estado de Antiog—Each contributed by the Sociedad de Naturalistes Nee Granadiros, Bogota New Granada.

Circular of the Am. Association of Anthropology.

Bulletin of the Illinois Museum of Natural History, No. 1.

Transactions of the Indianapolis Academy of Natural Sciences, 1870.

Transactions of the Indianapolis Academy of Natural Sciences, 1872.

Celebration of Humbolt's Centennial, contributed by the Iowa Institute of Sciences and Arts of Dubuque.

Transactions of the Davenport Academy of Natural Sciences, Vol. I.

Third Annual Report of the Minnesota Fish Commissioners.

Report of the New York State Agricultural Society, Vol. XXIII, Nos. 9, 10.

Report of the New York State Agricultural Society Awards, 1874.

Journal of the New York State Agricultural Society, 1873.

Proceedings of the Cincinnati Society of Natural History, No. 1.

Transactions of the Wisconsin Academy Sciences, Arts and Letters, 1874-5.

Birthplace of Americus.

Early Outposts of Wisconsin.

Transactions of the Wisconsin Academy of Sciences, Arts and Letters, 1875-6.

Journal of Science of the Canadian Institute, Toronto, Can., Vol. XIV, No. 1.

Proceedings of the Royal Society of New South Wales, Vol. X.

Progress and Resources of New South Wales.

Report of the Mining Department, New South Wales, 1876.

Climate of New South Wales.

Report of Railways, New South Wales.

Inauguration of President Gilman, Johns Hopkins University

Eleventh Annual Report of the Minnesota Hospital for the Insane.

Fourteenth Annual Report of the State Normal School, Winona.

Fifteenth Annual Report of the Minnesota State Institute for Deaf, Dumb and Blind.

Eighteenth Annual Report of the Superintendent of Public Instruction.

Eleventh Annual Report of the Minnesota State Reform School.

Annual Report of the Minnesota State Adjutant General, 1877.

Topography, Climate, etc., of Michigan, Prof. A. Winchell.

The Fifth Annual Report on the Geological and Natural History Survey of Minnesota.

DECADES OF MAINE FUNGI.

By M. C. COOKE, *London, England,*

(Corresponding Member.)

(DEC. I, II.)

Since the time of Schweinitz, little has been added to our knowledge of North American Fungi except by the labors of H. W. Ravenel of South Carolina, and Dr. Curtis. As far as I am aware, no attempt has been made to secure a list of the Fungi of any state except South Carolina and Ohio, (Cincinnati, T. G. Lea;) and certainly no effort has hitherto been made in favor of the state of Maine. Through the kindness of the Rev. E. C. Bolles, I hope gradually to obtain specimens, at least of the more minute, and less perishable Fungi, from the neighborhood of Portland, so that in course of time a respectable list may be constructed of the Fungi of Maine. Without waiting till such list can be perfected, I have consented to notice and describe the several consignments which I receive from time to time, and to give under each species a brief character, sufficient for its identification.

The two Decades now forwarded are confined to Coniomycetal Fungi, those in which the spores are the main features, and which are developed principally upon the green parts of living plants. These include the *Æcidacei*, the *Pucciniae*, and the *Coomacei*, as described fully in my "Microscopic Fungi." Circumstances have transpired to strengthen the views I therein expressed that the majority of species which constitute the genera *Lecythea* and *Trichobasis* are but unicellular forms of species belonging to other genera, and with these latter forms I shall associate them whenever I can do so with confidence. Instances will occur in which I am unable to associate species of *Trichobasis* or *Lecythea* with their higher forms, especially in extra-European species:—in such cases I must content myself with recording the species under their provisional names, in the hope that the future may unite them to their lawful spouses.

Di-morphism is well illustrated in the rust and mildew of wheat. Early in the season the 'red rust' appears. The orange spores are nearly globose, and one-celled, under which form they have been regarded as a species of *Trichobasis*. Later in the year the brown lines of the wheat mildew put in an appearance. The spores are elongated, attached to a long stalk or peduncle, are divided by a septum into two cells, and then known as *Puccinia graminis*. It is not uncommon to find spores of both forms in the same pustule and springing from the same mycelium. It has been suspected, and even asserted, that species of *Uromyces* and *Æcidium* are further developments from *Trichobasis* and *Puccinia*, and form with them an alternation of generations. The assertion that *Æcidium*

Berberidis is an associate-form of *Puccinia graminis*, rests for the present upon the observation of one individual, and requires confirmation before it can be accepted.

ÆCIDIACEI.

The Family *Æcidiacei* consists, as far as our knowledge of North American species is concerned, of three genera, thus briefly characterized :

Spores contained within a peridium.

Peridium elongated,
separating in threads—*Ræstelia*.
rupturing irregularly—*Peridermium*.

Peridium abbreviated or semi-immersed—*Æcidium*.

RÆSTELIA, *Rebent.*

Peridium elongated, at length opening by lateral fissures, or a terminal lacerated mouth. *Spermogonia* on the opposite surface; on the same, or on different leaves.

Two European species have already been found in Maine. Schweinitz records four in his 'Synopsis,' but two of these I have not seen.

Ræstelia lacerata, Tul. Peridia clustered in tufts, brown, elongated, splitting to the base in segments: spores light brown.

On the under surface of the leaves, and on the petioles and fruit of Hawthorn, (*Crataegus*) &c. Westbrook: E. C. B.

(*Ræstelia cylindrites*—(in part)—Schweinitz Tyn. Car. No. 432. Fung. Am. Bor. No. 2899.—*Æcidium laceratum*. Ravenel, Fungi. Car. Ex. V, No. 96.)

Ræstelia cancellata, Reb. Spots yellow, then red, prominent; peridia split to the base into laciniae which remain united at the apex. On Pear leaves &c. (Westbrook: E. C. B.)

(*Æcidium cancellatum*, Schweinitz Syn. Car. No. 433. *Ræstelia rastelites*, Schw. Fung. Am. Bor. No. 2900.)

ÆCIDIUM, *Pers.*

Peridium seldom elongated, opening by a terminal mouth, surrounded by a fringe of recurved teeth, or when short bursting irregularly. Spores disposed in chains. *Spermogonia* on the same or the opposite surface, clustered or scattered, central or intermixed.

A large genus, well represented in the United States.

Æcidium Geranii, DC. Spots yellow and purple; peridia in circinating clusters. Spores yellow; at length brown.

On the under surface of leaves of *Geranium*, Westbrook: E. C. B.

(*Æcidium Geranii maculati*, Schweinitz Syn. Car. No. 443. *Æcidium Geraniatum*. Schw. Fung. Am. Bor. No. 2879.)

Æcidium crassum, Pers. Spots yellow-brown, subiculum thickened; peridia crowded into a roundish heap, at first globose, yellow, at length open; spores orange.

On leaves of *Rhamnus* &c. (Canton, N. Y.; Maine.)

(Ravenel, Fungi Car. exsic. I. No. 92.)

Æcidium Hypericatum, Schwz. Spots small, roundish, orange; Peridia cylindrical, elevated, collected in small tufts, whitish when dried. Spores oblong, whitish, pellucid.

On leaves of *Hypericum perforatum* (Westbrook) also on *Hypericum frondosum* (fide Schweinitz).

(Schweinitz Syn. Car. No. 451.—Fung. Am. Bor. No. 2883.)

Whether the spores of the species on leaves of *H. perforatum*, received from the Rev. E. C. Bolles, are whitish, or yellow when fresh, could not be determined from the dried specimens. It is referred here to Schweinitz's species, with which it accords in other respects. Although the *Hypericum* is common in England, neither the *Æcidium* nor *Uromyces* has yet been found on it.

Æcidium asteratum, Schwz. Spots pallid purple, sometimes confluent. Peridia clustered and semi-immersed. Spores white.

On leaves and stems of *Aster*, (Westbrook). On *Aster cordifolius* (New York, W. W. Denslow).

(Schweinitz Syn. Car. No. 444.—Syn. Fung. Am. Bor. No. 2870.)

The only appreciable difference between this and *Æcidium compositarum*, D. C. appears to be in the colour of the spores, and it is doubtful whether this is as reliable a distinction as hitherto supposed.

PUCCINLÆI.

a. Spores stipitate :

Spores multiseptate—*Aregma*.

“ biseptate—*Triphragmium*.

“ uniseptate—*Puccinia*.

b. Spores immersed in Gelatine :

Stroma tremelloid and expanded—*Gymnosporangium*.

“ clavate, or club-shaped—*Podisoma*.

AREGMA Fr.

Spores cylindrical, multiseptate, borne on a long peduncle.

Three American species are well known. Two of these are European. One only has yet been sent from Maine.

Aregma mucronatum, Fr.

Uredo-form. (*Lecythea Rosæ*, Lev.)

Spots yellow, small, scattered; spores sub-oval, sometimes minutely pedicellate, orange.

Septate-form—Hypogenous, scattered over the leaves in minute tufts; spores 5 to 7-septate, terminal joint mucronate; peduncles incrassated below, fusiform.

On leaves of Roses (Westbrook, E. C. B.)

Curtis in Silliman's Journ. 1848, vi. p. 353, No. 26.

PUCCINIA, Pers.

Spores of two kinds. Early spores, or Uredo spores, globose or nearly so, unicellular, at first with a short peduncle which ultimately falls away, yellow or brown. Later spores uniseptate, supported on a distinct peduncle.

This genus is largely represented in North America. Very few are yet known from Maine.

Puccinia graminis, Pers.

Uredo-form, (*Trichobasis rubigo-vera*, Lev.)

Spots yellow, heaps oval, scattered; generally on the upper surface; epidermis at length bursting longitudinally; spores sub-globose, reddish brown, easily dispersed. Ravenel, Fung. Car. Ex. i. No. 100.

Septate form—Spots pale, diffuse; sori linear, confluent, amphigenous; spores at length black, clavate, very slightly constricted, uniseptate. On the leaves and culms of corn and grass. (Westbrook, E. C. B.; New York, W. W. D.)

(*Uredo Rubigo*. Schweinitz Fung. Am. Bor. No. 2817.)

(*Puccinia graminis*. Schw. Syn. Car. No. 492. Fung. Am. Bor. No. 2905. Lea, Plants of Cincinnati, p. 70. Ravenel, Fungi Car. Ex. i. No. 99.)

Puccinia striola, Link.

Uredo-form—(*Trichobasis Caricina*, B.)

Spots reddish; sori oval, minute, scattered, surrounded by the ruptured epidermis; spores sub-globose, reddish, then brown.

Septate form—Spots pallid; sori linear, crowded, distinct, sub-convex; spores at length black, oblong and slightly constricted, or obovate and not constricted.

On sedges, rushes, &c. (Westbrook, E. C. B.)
(Schweinitz, Fung. Am. Bor. No. 2906.)

Puccinia Circææ, Pers.

Uredo-form—Uncertain.

Septate form—Spots obliterated; sori minute, semiglobose, crowded into subrotund patches, occasionally confluent, generally covered with the epidermis.

On leaves of *Circæa*. (Westbrook, E. C. B.)
(Schweinitz Syn. Car. No. 491. Fung. Am. Bor. No. 2938. Lea, Plants of Cincinnati, p. 70.)

Puccinia Violarum, Link.

Uredo-form (*Trichobasis Violarum*, Berk.) Spots yellowish; sori subrotund, scattered, generally on the under surface; epidermis ruptured, persistent; spores sub-globose, brown.

Septate form—Spots yellowish; sori hypogenous, small, crowded, covered with the epidermis, then surrounded by it; spores brown, elliptic, or broadly elliptic, slightly constricted, uniseptate.

On Violet leaves, (Scarboro', Westbrook, E. C. B.)

Both forms of spores are often present in the same pustule.

Puccinia Pyrolæ, Cooke.*Uredo-form*—Unknown.

Septate-form—Spots large, pallid, orbicular; sori round, often confluent and then elongated, concentric, bullate and compact, nearly black, surrounded by the epidermis; spores ovate or elliptic, scarcely constricted, peduncles elongated, clear umber brown, upper cell darkest.

On leaves and petioles of *Pyrola*, (Cape Elizabeth, E. C. B.)

This is a most distinct and unmistakable species, and apparently undescribed. The external appearance, habit and color, is very much that of *Puccinia Umbilici*, Guép.

Puccinia Xanthii, Schw.*Uredo-form*—Uncertain.

Septate form—Spots orbicular, pallid, on the opposite surface yellowish brown with a pallid margin. Sori compact, large, concentric, confluent, orbicular, forming a blackish brown, convex mass. Spores elliptic, slightly constricted, brownish; peduncles elongated.

On leaves of *Aster*. (Westbrook E. C. B.) On leaves of *Xanthium strumarium*. (Schweinitz.)

(Schweinitz Syn. Car. 500. Fung. Am. Bor. No. 2927.)

PODISOMA, Link.

Peduncles extremely long, agglutinated by gelatine into a common stem, spreading out above into a clavariæform mass. Spores mostly uniseptate.

Gymnosporangium is represented in South Carolina, and besides *Podisoma Juniperi* a distinct form which has been named *Podisoma macropus* is known from the same State. Only the true *Podisoma Juniperi* has yet been found in Maine.

Podisoma Juniperi, Fr.

Orange, clavariæform, somewhat branched; stroma simple; spores very long, lanceolate, filled with elliptic granules.

On living branches of *Juniper*, (Cape Elizabeth, E. C. B.)

(Schwz. Fung. Am. No. 3095.)

The spores are very much longer than in the form named *Podisoma macropus*, Schw., which has been referred to this species. They are longer than in British specimens, which again are longer than in *P. macropus*. The figure in this, as in all the species figured, are from American specimens.

CÆOMACEI.

Spores of one order;

Simple and free;

Produced in separate cells,

Superficial, yellow or brown—*Uredo*.

Not inclosed in separate cells—*Lecythæa*.

Stipitate;

With permanent peduncles,—*Uromyces*.

Spores of two orders;

1. Spherical,
2. Wedge-shaped, compact, } *Melampsora*.

UREDO, Lev.

Stroma composed of little irregular cells forming a lentiform disc, whose surface is covered with many layers of cells, each of which encloses a spore. Spores simple, always without any appendage.

Uredo Potentillarum, D. C. Spots yellowish; sori subrotund or oval, bullate, aggregate, open, confluent; spores sub-globose, sub-coherent, orange.

On various Rosaceæ (Westbrook E. C. B. New York, W. W. D.)

(Schweinitz, Syn. Car. No. 461. Fung. Am. Bor. No. 2834. Lea, Plants of Cincinnati, p. 71. Ravenel, Fung. Car. Exsic. ii. No. 97.)

Uredo Filicum, Desm. Spots yellowish, sori subrotund, bullate, scattered and aggregate, on the under surface; epidermis at length bursting; spores sub-globose, yellow.

On Ferns. (Scarboro', E. C. B.)

(Schweinitz, Fung. Am. Bor. No. 2836. Ravenel, Fung. Car. Exsic. iv. No. 97.)

I have received *Uredo solidaginis* Schw. from New York, and Newport, Rhode Island, but not at present from Maine.*

UROMYCES, Lev.

Spores unilocular, attached permanently to a decided peduncle of greater or less length.

Uromyces apiculosa, Lev. Spots yellow or brown. Sori sub-rotund, scattered, surrounded by the ruptured epidermis; spores ovoid, brown, shortly pedunculate.

On leaves of various plants, (Peak's Island, E. C. B.)

(*Uredo apiculosum*. Schweinitz, Syn. Car. No. 478. Fung. Am. Bor. No. 2844.)

Uromyces triquetra, Cooke.

Spots claret-colored; Sori small, scattered, hypogenous, surrounded by the remains of the ruptured epidermis. Spores angular, often three sided.

On leaves of *Hypericum*, (Scarboro, Westbrook, E. C. B.)

Whether this is the same as the species described by Schweinitz under the name of *Uredo Hyperici* is somewhat uncertain, since his specimens are recorded as occurring on the stems, forming elevated bullate sori, and the present is confined to the leaves, and the sori are not bullate. Neither of them are the yellow *Uredo Hypericorum*, D. C., so common in Britain.

MELAMPSORA, Cast.

Spores of two orders. Summer spores globose or oval, yellow. Winter spores crowded into a dense compact mass, with or without a covering, wedge-shaped.

*Found by Mr. C. B. Fuller at Brunswick, Maine.

Melampsora populina, Lev. Hypophyllous, epiphyllous, or amphigenous, at first yellow or orange.

Summer spores—obovate oblong, attenuated or truncate, echinulate, paraphyses obovate, capitate or claviform, abundant in fully ripened sori.

Winter spores—Sori at first tawny yellow, becoming black during the winter, swelling in the spring, and becoming of a cinnamon color, hypophyllous, roundish or oblong; spores prism-shaped, five or six together, yellowish, smooth.

On Poplar leaves. (Westbrook, E. C. B.)

(*Uredo cylindricum*. Schweinitz, Fung. Am. Bor. No. 2855.)

LECYTHEA, Lev.

Stroma surrounded or sprinkled with elongated abortive spores. Spores free, invested with their mother-cell or concatenate.

Probably not a good genus, most of the species having been found to be only conditions of species in other genera. It is only inserted here provisionally to include species not yet traced to their associate forms.

Lecythea gyrosa, Berk. Spots obliterated; sori minute, confluent, and forming a small distinct ring; epiphyllous, epidermis bursting; spores globose, and elongato-pyriform, yellow or pale.

On the upper surface of Bramble and Raspberry leaves. (Westbrook, E. C. B.; New York, W. W. D.)

(*Uredo gyrosum*. Schweinitz, Fung. Am. Bor. No. 2854.)

It is also doubtful whether this is an autonomous species.

Lecythea epitea, Lev. Sori roundish, scattered, at first tawny, at length growing pale, surrounded by the ruptured epidermis; spores subrotund and pyriform.

On Willow leaves. (Chicago, Illinois; Niagara Falls, E. C. B.)

(*Uredo epiteum*. Schweinitz, Fung. Am. Bor. No. 2856. Ravenel, Fung. Car. Ex. v. No. 98.)

N. B. This is probably only the *Uredo*-form of some species not yet associated with it, and as such is recorded here provisionally under the above name assigned to it by Leveille. At present I have not received it from any locality in Maine.

THE CLIO BOREALIS ON THE COAST OF MAINE.

By DR. W. WOOD.

This animal was brought to the notice of the Society by Mr. C. B. Fuller, at its regular meeting, May 7, 1868. He stated that whilst looking for *Acalephs*, as he has long been in the habit of doing, he obtained a single specimen of the *Clio*, for the first time April 6th, with *Bolina*, *Pleurobrachia*, *Idyia*, *Nanomia* and other jelly fishes, which at that time were very abundant. On April 28th the *Nanomia*, *Tiaropsis*, *Oceania*, *Bolina* with the *Sagitta* were taken

in abundance, and with them a few Clios near the end of the wharf at House Island in Portland Harbor. On May 2d, the Clios were abundant, coming in apparently with the high spring tides, and from that time they continued to become more and more numerous, until finally the water appeared to be alive with them. After May 7th, they began to decrease and to drop off with the low tides, and entirely disappeared with the approach of the rainy weather, so that in few days more not one could be obtained.

The novelty of this sudden appearance and disappearance of an animal regarded and described as a denizen of the Arctic seas, its inherent beauty, graceful, and seemingly joyous movements, at once excited a lively interest on the part of the members present, none of whom had before had an opportunity of seeing it.

Questions arose as to the causes of its appearance at this time, and the frequency of its visits to these waters, and the most southern limit of its present and past migrations? These are questions which do not admit of an entirely satisfactory conclusion, but it was generally conceded that this visitation was in some way connected with the Arctic character of the winter through which we had just passed, it having been one of unusual severity, not once yielding to a thaw.

Upon examination, no allusion is made to this animal in Gould's Invertebrata of Massachusetts, nor in any of the volumes of the Journal or Proceedings of the Boston Society of Natural History, these publications dating back to 1841; and no mention is made of it in the publications of the Philadelphia Academy of Nat. Sciences. The name occurs in Stimpson's Check list of the shells of North America, and is mentioned by Packard, in his list of animals, dredged off Southern Labrador, as being quite abundant in that region. The only approach to an answer to the above questions that could be obtained from any source within our reach, was derived from the New York Reports. De Kay, there gives a brief description of the animal and a figure which are generally correct. He says that "this species occurs in almost incredible numbers in the Northern Atlantic, where it forms the ordinary food of whales. It has been observed occasionally in great numbers, in our bays. In April, 1833, they were very abundant and of a blood red color. After a few days they all disappeared."

We cannot find that their presence in our waters has been noticed by any one since that date, down to the time of their being discovered by Mr. Fuller, nor can we learn that *this* visit of the Clio, was observed in any other place along the coast. This would leave room for the supposition, that their visits might be more frequent than the known facts would indicate, since it is probable that at that season of the year, there are but few abroad watching the waters with the carefulness necessary to detect the presence of the Clio. Certain it is, however, that for a series of years, no such event has occurred, or it would have been announced by Mr. Fuller, who at all times and seasons has scanned the waters with a keen and critical eye in search of every thing

pertaining to the study of Natural History, but especially of the minute and transparent acalephs or jelly fishes, which, clear as the waters in which they float, are very much more difficult to detect than these colored Clios.

That naturally the shores of Labrador are the most southerly bounds of its migrations, its visits to our waters being as rare as already indicated, is rendered still further probable by the fact that the genus *Clio*, was originally instituted by Brown, in his "Natural History of Jamaica", to include a *tropical* species, but the name was subsequently adopted and modified by Linnaeus and applied to the only representative of the class known by him, the *Clio borealis*, the genus *Cleodora* having been at a later date created to receive the original *Clio* of Brown. This animal belongs therefore, to a division of the Mollusca, called Pteropoda or "Wingfoot", from the swimming appendages which resemble the organs of flight in birds, and forms one of the six genera into which it is divided, each genus being represented by a single or a very limited number of species.

The figure accompanying this article, drawn by Mr. Fuller, about double the natural size, gives a very accurate idea of the general form of the animal.

It is of a soft jelly like consistency, having a pearly or opaline color with the extremities or shaded portions, as seen in the figure, of a bright red color. It is so nearly transparent, that the internal organs, the stomach, liver &c. can be plainly seen occupying but a limited portion of the sac of which the body is composed. Seen as delineated in the figure, one would be puzzled to determine by what process it contrives to nourish itself. But a little observation shows that the head-piece is capable of longitudinal division from before backwards, the two upright tentacles becoming horizontal, whilst from the inner surface of each hemisphere protrude three other tentacles which are at once understood to be the organs by which its food is directed to the mouth lying at the bottom of the fissure made by the division of the head-piece. The formidable

array of numerous long curved teeth with which this mouth is armed seems to indicate a method of existence quite at variance with the "angelic" cut of its vestments, and is rather suggestive of the wolf in sheep's clothing. But after close observation of them for many days, in glass jars and tanks, no food could be detected going into the mouth, even when examined with a lens, but the accumulation of a dark substance in the stomach in which Mr. Fuller detected the presence of diatomaceous shields, and of a considerable quantity of excrementitious matter at the bottom of the jar, proved that they were busy in separating from the waters the invisible pabulum of life.

Their movements were almost incessant, and were constantly those of ascending and descending, almost always maintaining a vertical position even when moving laterally; at rare intervals they would suddenly pitch over and start headlong for the bottom with hurried movement, as if bent upon a frolic, but they usually recovered their upright position and measured step before reaching it.

The motion of the wing like appendages was monotonous and always jerkily from before backwards, making the tips nearly meet in both directions, sweeping through the arc of a semicircle. No variation in the direction of this regular flip-flap motion could be detected by which the upward or downward movements of the animal could be predicated. Why they should go up or why they should go down could not be determined by any visible variation in this singular movement of the wings. These movements were almost unceasing, whether observed by day or by night, the only variation being, that at long intervals they would keep near the surface of the water, bobbing apparently for a bubble of air which they would contrive to swallow, or lying perfectly motionless, and apparently dead upon the bottom of the jar, for a longer or shorter period. In this manner they lived in the jar for several weeks and even for sometime after they had entirely disappeared from the waters of the harbor.

This animal has been described by numerous writers, but evidently by a few only, from their own observations. The "Anatomical Researches upon the *Clypeus borealis*," by D. F. Eschricht however, published at Copenhagen in 1838, illustrated by numerous minute beautiful and accurate figures, seem to preclude the necessity for any better description of its intricate structure.

By reference to a diagram, made by Mr. A. W. Longfellow, of the Coast Survey, for the purpose of determining at a glance the mean temperature of Portland Harbor for a series of years, it would appear that the winter and spring of 1832-3 were below the average range, and may perhaps have been similar in other respects to that of 1867-8. This, as has been already stated, was characterized by an early, steady, uninterrupted cold, though not extreme, but unbroken by a single thaw till spring was well advanced. Whether the same peculiarity attended that of 1832, we have not been able altogether to ascertain, nor can we of course positively decide that it is to this peculiarity of the season that we are indebted for this visit of the *Clio*.

XIV. The Hyphomycetous Fungi of the United States

BY M. C. COOKE, M. A.

[Read before this Society, Feb. 2, 1877.]

THIS division of the Fungi of the United States is in the most unsatisfactory condition, requiring complete examination and verification. The present list must therefore be regarded as involving no responsibility for accurate determination, but as a preliminary step towards securing verification:

Order I. ISARIACEI.

1. *Isaria farinosa* Fr. S. M. iii, 270. On dead pupae. Car. (Curt.); N. Eng. (Murray).
[*Stilbum ramosum* Peck, probably belongs to this species.]
2. *Isaria sphingum* Schw. (syn. Car. 1298), Am. Bor. 3055. On *Phalaena*. Car. (Curt.).
3. *Isaria araneorum* Schw., Am. Bor. 3057. On spiders. Car. (Curt.).
4. *Isaria nigripes* Schw., Am. Bor. 3068. On chrysalids. Car. (Curt.).
5. *Isaria phalangiphila* Schw. (syn. Car. 1299), Am. Bor. 3056. Car. (Schw.).
6. *Isaria crassa* Schw., Am. Bor. 3054. On chrysalids. Penns. (Schw.).
7. *Isaria furcata* Schw. Am. Bor. 3067. On chrysalids. Bethlehem (Schw.).
All the foregoing are conidia of species of *Torrubia*, although entered here.
8. *Isaria brachiata* Schum. Curt. Cat. p. 125. On *Agarics*. Car. (Curt.).
9. *Isaria agaricina* Pers. Schw., Am. Bor. 3058. On *Agarics*. Bethlehem (Schw.).
[Perhaps the same as the preceding].
10. *Isaria radiata* B. & C., Grevillea iv, 584. On pine wood. Ala. (Peters).
11. *Isaria brevis* B. & C., Grevillea iv, 585. On decayed wood. Penns. (Mich.).
12. *Isaria congesta* B. & C., Grevillea iv, 586. On *Sphaeria deusta*. Car. (Curt.).
13. *Isaria umbrina* Schw., Am. Bor. 3064. On rotten wood. Car. (Curt.).
[The conidia of *Hypoxyylon coccineum*.]
14. *Isaria citrina* Pers. Schw., Am. Bor. 3066. On rotten trunks. Car. (Schw.; Curt.).

15. *Isaria carnea* P. Schw., Am. Bor. 3069. On stems and leaves. Car. (Curt.).
16. *Isaria clavata* Ditm. Schw., Am. Bor. 3060. On rotten chips.
17. *Isaria bulbosa* Nees. Schw., Am. Bor. 3061.
18. *Isaria hydroides* Lk. Schw., Am. Bor. 3062. On trunks.
19. *Isaria glaucocephala* Lk. Schw., Am. Bor. 3063. On branches.
20. *Isaria furcellata* Mart. Schw., Am. Bor. 3065. On rotten chips.
21. *Isaria candida* Schw., Am. Bor. 3070. On rotten wood.
22. *Isaria globosa* Schw., Am. Bor. 3071. On the ground.
23. *Isaria bicolor* Schw., Am. Bor. 3072. On stems lying on the ground.

Some of these species are entered solely on the authority of Schweinitz, and require verification.

24. *Ceratium hydroides* A. & S. Schw., Am. Bor. 3074. On dead wood. Car. (Curt.); Ala. (Peters); N. Y. (Peck).
25. *Ceratium porioides* A. & S. Schw., Am. Bor. 3076. On pine wood. Car. (Schw.; Curt.); N. Y. (Peck).
26. *Ceratium pyxidatum* Schw., Am. Bor. 3075, is probably the same as the last.
27. *Ceratium crustaceum* B. & C., Grevillea iv, 587. On decayed wood. New Jersey (B.).
28. *Ceratium insociabile* Gerard. Bull. Torr. Club. On leaves.
29. *Dacrina cinnabarinum* Nees. Schw., Am. Bor. 3078. On bark of *Juglans*. Car. (Curt.).
30. *Dacrina hydnoideum* Schw., Am. Bor. 3079. On *Rhododendron* wood.
31. *Gliotrichum fuligo* Fr. (*Dematium fuligo* Schw., Am. Bor.).
32. *Peribotryon coccineum* Fr. (*Coremium coccineum* Schw., Am. Bor.).
33. *Peribotryon subiculatum* Fr. (*Coremium subiculatum* Schw., Am. Bor.).
34. *Peribotryon camptotrichum* Fr. (*Camptotrichum splendidum* Schw.).

Order II. STILBACEI.

35. *Stilbum tomentosum* Schr. Schw., Am. Bor. 3715. On *Arcyria* and *Trichia*. Car. (Schw.; Rav.); N. Y. (Peck).
36. *Stilbum fimetarium* B., Grevillea iv, p. 63. On dung. Car. (Curt.).
37. *Stilbum byssinum* P. Schw., Am. Bor. 2716. On putrid fungi. Car. (Schw.; Curt.).
38. *Stilbum bulbosum* Tode. Schw., Am. Bor. 2718. On stems. Car. (Schw.; Curt.).
39. *Stilbum piliforme* A. & S. Schw., Am. Bor. 2717. On wood. Car. (Schw.; Curt.).
40. *Stilbum lateritium* Berk, Lond. Jour., 1843, 642. On bark. Car. (Curt.).
41. *Stilbum cinereo-rubrum* B. & C. On *Cyrilla* and *Hibiscus*. Car. (Curt.).

42. *Stilbum carcinophthalmum* B. & C., Grevillea iv, 590. On dead stem of parsnip. Car. (Curt.).
43. *Stilbum leucocephalum* B. & C., Grevillea iv, 596. On bark of *Carya*. Car. (Curt.).
44. *Stilbum aurifilum* Gerard, Bull. Torr. Club. On old *Daedalea*.
45. *Stilbum giganteum* Peck., 24th Report, p. 93. On old logs. N. Y. (Peck).
46. *Stilbum pellucidum* Schrad. (fide Peck). On rotten wood. N. Y. (Peck).
47. *Stilbum rhoidis* B. & C., Grevillea iv, 595. (*Sporocybe Rhois* Curt.). On bark of *Rhus*. Car. (Curt.); N. Y. (Peck).
48. *Stilbum vulgare* Tode. Schw., Am. Bor. 3713. On dead wood. Maine (Sprague); Car. (Curt.); N. Eng. (Frost).
49. *Stilbum gelatinosum* P. Schw., Am. Bor. 2714. On trunks. Car. (Schw.; Curt.).
50. *Stilbum rigidum* P. Schw., Am. Bor. 2719. On dead twigs of *Carya*. Car. (Curt.).
51. *Stilbum turbinatum* Tode. Schw., Am. Bor. 2720. On wood. Car. (Schw.; Curt.).
52. *Stilbum rubicundum* Tode. Schw., Am. Bor. 2721. On beech stumps. Car. (Schw.; Curt.).
53. *Stilbum spraguei* B. & C., Grevillea iv, 589. On cabbage stalks. N. Eng. (Sprague).
54. *Stilbum annulatum* B. & C., Grevillea iv, p. 63. On cabbage stalks. Car. (Curt.).
55. *Stilbum lichenoideum* B. & C., Grevillea iv, 592. On dead wood of *Juniperus*. Penns. (Mich.).
56. *Stilbum vellereum* B. & C., Grevillea iv, 593. On decayed wood. Penns. (Mich.).
57. *Stilbum aleuriatum* B. & C., Grevillea iv, 594. On decayed wood of *Acer*. Texas (Wright).
58. *Stilbum roseum* Schw., Am. Bor. 2722. On rotten wood.
59. *Stilbum inquinans* Schw., Am. Bor. 2723. On branches.
- 59½. *Chordostylum capillare* Fr. Schw., Am. Bor. 2712. On leaves.
60. *Atractium fusisporum* B. & C., in Curt. Cat. 126. On bark of *Acer*. Car. (Curt.).

This and *Atractium flammeum* are doubtless forms of *Sphaerostilbe*.

61. *Hyalopus mucorinus* B. & C., Grevillea iv, 598. On hens' dung. Car. (Curt.).
62. *Hyalopus parasitans* B. & C., Grevillea iv, 597. On decayed *Merulius*. Car. (Curt.).
63. *Hyalopus griseus* B. & C., Grevillea iv, 599. On wood, under the bark of *Nyssa*. Car. (Curt.).

64. *Graphium explicatum* B. & C., Grev. iv, 620. On twigs of *Quercus Catesbei* and *nigra*. Car. (Curt.).
65. *Graphium clavisorum* B. & C., Grevillea iv, 619. On vine leaves. Car. (Curt.).
66. *Chondromyces crocatus* B. & C., Grev. iv, 600. On decayed melons. Car. (Curt.).
67. *Tubercularia vulgaris* Lk. Schw., Am. Bor. 2999. On branches. (Conidia of *Nectria*, as also are the majority of species in this genus.) Car. (Schw.; Curt.); N. Y. (Peck); Ohio (Lea.); Maine (E. C. B.).
68. *Tubercularia granulata* Schw., Am. Bor. 3003. On branches. Car. (Schw.; Curt.); N. Eng. (Frost).
69. *Tubercularia minor* Schw., Am. Bor. 3006. On *Castanea*. Car. (Schw.; Curt.).
70. *Tubercularia nigricans* Schw., Am. Bor. 3006. On wood and bark of *Fraxinus*. Car. (Curt.); N. Eng. (Frost); N. Y. (Peck).
71. *Tubercularia microsperma* B. & C. Curt. Cat., p. 126. On pine cones. Car. (Curt.).
72. *Tubercularia dubia* Schw., Am. Bor. 3004. On branches. Car. (Curt.).
73. *Tubercularia persicina* Ditm. Curt. Cat., p. 126. On *Uredo luminata*. Car. (Curt.).
74. *Tubercularia pezizoidea* Schw., Am. Bor. 3013. On poplar. N. Y. (Schw.); N. Eng. (Sprague).

The following require revision:

75. *Tubercularia subdiaphana* Schw., Am. Bor. 3020. On *Vitis*.
76. *Tubercularia calycanthi* Schw., Am. Bor. 3019. On branches of *Calycanthus*.
77. *Tubercularia fatescens* Schw., Am. Bor. 3018. On branches.
78. *Tubercularia radicalis* Schw., Am. Bor. 3017. On roots of trees.
79. *Tubercularia difformis* Schw., Am. Bor. 3016. On spines of *Crataegus*.
80. *Tubercularia celastri* Schw., Am. Bor. 3015. On branches of *Celastrus*.
81. *Tubercularia subpedicellata* Schw., Am. Bor. 3014. On branches of *Lycium*.
82. *Tubercularia pusilla* Fr. Schw., Am. Bor. 3012. On young branches.
83. *Tubercularia sarmentorum* Lk. Schw., Am. Bor. 3011. On *Zea*.
84. *Tubercularia menispermii* Schw., Am. Bor. 3010. On branches of *Menispermum*.
85. *Tubercularia herbarum* Schw., Am. Bor. 3009. On stems of *Artemisia*.
86. *Tubercularia liceoides* Schw., Am. Bor. 3008. On cabbage stalks.
87. *Tubercularia ciliata* Schw., Am. Bor. 3007. On rotten wood.
88. *Tubercularia floccosa* Schw., Am. Bor. 3005. On *Rhus typhina*.
89. *Tubercularia mirabilis* Schw., Am. Bor. 3002. On branches of *Morus*.
90. *Tubercularia lutescens* Schw., Am. Bor. 3000. On branches of *Aralia*.

91. *Cheiromyces stillatus* B. & C., Grevillea iv, 601. On *Scirpus Eriophorum*. Penns. (Mich.); Ala. (Beaum.).
92. *Stigmatella aurantiaca* B. & C., Grevillea iv, 602. On *Sphaeria Hibisci* and lichens. Car. (Curt.; Rav.).
93. *Volutella ciliata* Fr. Schw., Am. Bor. 3023. On stems. Car. (Curt.).
94. *Volutella pulchra* B. & C., Grevillea iv, 603. On willow branches. Conn. (B.).
95. *Fusarium roseum* Lk. Schw., Am. Bor. 3022. On various substances. Car. (Curt.); N. Y. (Peck).
96. *Fusarium lateritium* Schw., Am. Bor. 3026. On various plants. Car. (Curt.; Rav.); Ohio (Lea); N. Y. (Peck).
97. *Fusarium aurantiacum* Cda. On *Acacia julibrissin* and gourds. Car. (Curt.).
98. *Fusarium marginatum* B. & C., Grevillea iv, 604. On stems of *Smilax*. Ala. (Peters).
99. *Fusarium microspermum* B. & C., Grevillea iv, 605. On fig. Santee River (Rav.).
100. *Fusarium heliotioides* B. & C., Grevillea iv, 606. On *Ilex prinoides*. Ala. (Peters).
101. *Fusarium Mikaniae* B. & C., Grevillea iv, 607. On stems, etc., of *Mikania*. Santee River (Rav.; Curt.).
102. *Fusarium nitidum* B. & C., Grevillea iv, 608. On *Aralia spinosa*. Penns. (Mich.).
103. *Fusarium erubescens* B. & C., Grevillea iv, 609. On dead bark. Ala. (Beaumont); N. Y. (Peck).
104. *Fusarium semitectum* B. & Rav. Grevillea iv, 610. On petioles of banana. Penns. (Rav.).
105. *Fusarium cymbiferum* B. & C., Grevillea iv, 611. On herbaceous stems. Car. (Curt.).
106. *Fusarium anomalum* B. & C., Grevillea iv, 612. On *Gleditschia*. N. Eng. (Sprague).
107. *Fusarium Berkeleyi* Mont. Grev. iv, 612.* On fig. *Hibiscus, cornus*. Car. (Curt.; Rav.). Ala. (Peters).
108. *Fusarium sticticum* B. & C., Grev. iv, 613. On peach twigs. Car. (Curt.).
109. *Fusarium arcuatum* B. & C., Grev. iv, 614. On apple bark. Car. (Curt.).
110. *Fusarium pyrinum* Schw., Am. Bor. 3028. On apple. Penns. (Schw.).
111. *Fusarium pallens* Lk. Schw., Am. Bor. 3021. On bark. N. Y. (Schw.).
112. *Fusarium Curtisii* (*Fusarium pallens* B. & C., in Grevillea iv, 615. Not Link). On inflorescence of *Juncus*. Car. (Curt.).
113. *Fusarium heterosporium* Lk. Schw., Am. Bor. 3027. On *Paspalum*. Penns. (Schw.).

114. *Fusarium oxysporum* Schlecht. Schw., Am. Bor. 3025. On legumes of *Phaseolus*.
115. *Fusarium stilbaster* Lk. Schw., Am. Bor. 3024. On roots.
116. *Fusarium effusum* Schw., Am. Bor. 3029. On branches of *Hypericum*.
117. *Fusarium coccineum* Schw., Am. Bor. 3031. On bark of *Castanea*.
118. *Fusarium capitatum* Schw., Am. Bor. 3032. On wood of *Pinus Canadensis*.
119. *Fusarium insepitatum* Schw., Am. Bor. 3030. On bark of *Mezereum*.
120. *Myrothecium convexum* B. & C., Grevillea iv, 616. On decaying grapes. Penns. (Mich.).
121. *Myrothecium roridum* Tode. Grev. iv, 616* (*M. carmichaeli* Schw., Am. Bor. 2389). On leaves, etc., of *Syringa*. Car. (Curt.); N. Eng. (Murray).
122. *Myrothecium inundatum* Tode. Schw., Am. Bor. 2388 (*M. fungicola* Peck). On Agarics. N. Y. (Peck).
123. *Myrothecium verrucaria* Ditm., Schw., Am. Bor. 2390. On leaves, grass, etc. Car. (Curt.). Ala. (Beaumont); Penns. (Mich.).
124. *Dermosporium atrum* Schw., Am. Bor. 3035. On hickory wood.
125. *Dermosporium roseum* Schw., Am. Bor. 3034. On rotten wood.
126. *Dermosporium decolorans* Schw., Am. Bor. 3033. On stems of plants.
127. *Myxormia atroviridis* B. & Br. Grevillea iv. On bark of *Rubus*. Ala. (Beaumont).
128. *Epicoccum scabrum* Corda. Icon. i, fig. 81. On dry roots of *Nyssa*. Car. (Curt.).
129. *Epicoccum neglectum* Desm. Cooke, Hdbk., No. 1678. On male inflorescence of *Zea*. Car. (Curt.).
130. *Epicoccum micropus* Corda. Icon. iii, 82. On old pods of *Phaseolus*. Penns. (Mich.).
131. *Epicoccum simplex* B. & C., Grev. iv, 617. (*E. sphaerospermum* Curt.). On leaves of *arundinaria*. Car. (Curt.).
132. *Epicoccum nigrum* Schw., Am. Bor. 3037. On stems of *Ricinus*.
133. *Epicoccum purpurascens* Schw., Am. Bor. 3638. On stems of asparagus.
134. *Epicoccum caricicola* Schw., Am. Bor. 3039. On leaves of *Carices*.
135. *Sphaerosporium lignatile* Schw., Am. Bor. 3036. On bark of oak and willow.
136. *Illosporium coccineum* Fr. Schw., Am. Bor. 2406. On lichens. Car. (Curt.).
137. *Illosporium roseum* Fr. Cooke, Hdbk., No. 1680. On *Physcia*. N. Y. (Peck).
138. *Illosporium carneum* Fr. Cooke, Hdbk., No. 1681. On *Peltigera*. N. Y. (Peck).
139. *Illosporium persicinum* Fr. Curt. Cat., p. 126. On pine palings. Car. (Curt.).

140. *Illosporium caesium* Schw., Am. Bor. 2407. On wood of walnut.
 141. *Aegerita candida* E. Schw., Am. Bor. 3040. On decayed bulb. Car. (Curt.).
 142. *Aegerita ovula* Schw., Am. Bor. 3043. On very rotten wood. Car. (Curt.).
 143. *Aegerita lutescens* Schw., Am. Bor. 3042. On wood.
 144. *Aegerita caesia* Schw. Am. Bor. 3041. Syn. Car. 1275. On wood. Car. (Schw. Curt.).

Order III. DEMATIEI.

145. *Pachnocybe rosea* B. & C., Grevillea iv, 588. On bark of *Robinia*. Car. (Curt.).
 146. *Pachnocybe subulata* B. (*Periconia*, Schw., Am. Bor. 3045.) On wood of *Acacia*. Car. (Curt.); Penns. (Schw.); N. Eng. (Sprague).
 147. *Periconia gracilis* Schw., Am. Bor. 3047. On capsules of *Staphylea*.
 148. *Periconia Azaleae* Peck. 25th Report, p. 93. On capsules of *Azalea*. N. Y. (Peck).
 149. *Periconia epiphylla* Schw., Am. Bor. 3047. On leaves.
 150. *Periconia nana* Lk. Schw., Am. Bor. 3046. On pine leaves.

We have two or three other forms of *Periconia*, but the specimens are insufficient for accurate description.

151. *Sporocybe byssoides* Fr. (*Periconia*, Schw., Am. Bor. 3044.) On dead herbaceous stems, etc. Car. (Curt.). Penns. (Schw.); New York (Peck).
 152. *Sporocybe Persicae* Fr. Curt. Cat., p. 126. On bark of *Persica* and *Cerasus*. Car. (Curt.). N. Y. (Peck).
 153. *Sporocybe calicioides* Fr. Grev. iv, No. 618. On bark of poplar. Car. (Rav.). Maine (Rev. J. Blake); N. E. (Sprague).
 154. *Sporocybe fasciculata* (*Periconia*, Schw., Am. Bor. 3049). Curt. Cat., p. 126. On stems. Car. (Curt.).
 155. *Sporocybe bulbosa* (*Periconia*, Schw., Am. Bor. 3051). Curt. Cat., p. 126. On stems. Car. (Curt.).
 156. *Sporocybe macularis* (*Periconia*, Schw., Am. Bor. 3050). Curt. Cat., p. 127. On naked wood. Car. (Curt.).
 157. *Sporocybe alternata* B. in Grev. iv, No. 618. On damp paper. Car. (Curt.).
 158. *Stachybotrys atra* Corda. Icon. i, fig. 278. On damp paper. Car. (Curt.).
 159. *Cephalotrichum rigescens* Lk. Schw., Am. Bor. 3052. On wood. N. E. (Russell).
 160. *Cephalotrichum monilioides* Lk. Schw., Am. Bor. 3053. On trunks of *Pinus Canadensis*.

161. *Dendryphium fasciculare* B. & Rav. Grev. iv, 618. On bark of *Liquidambar*. Car. (Rav.).
162. *Monotospora setosa* B. & C., Grev. iv, 621. On decayed wood in moist places. Car. (Curt.).
163. *Monotospora triseptata* Peck. 24th Report, p. 94, is *Helminthosporium apicale* B. & Br.
164. *Oedemium atrum* Fr. Schw., Am. Bor. 2610. On leaves and twigs. Car. (Curt.).
165. *Oedemium sparsum* B. & Rav. Grev. iv, 622. On bark of *Platanus*. Car. (Rav.).
166. *Actinocladium penicillus* Fr. Curt. Cat., p. 127. On leaves of *sassafras*. Car. (Curt.).
167. *Glenospora Curtisii* B. & Desm. Journ. Hort. Soc. Lond. On branches of *Nyssa*, *Quercus*, etc. Car. (Curt.).
168. *Glenospora Melioloides* Curt. Cat., p. 127. On living leaves of *Galax*. Car. (Curt.).
169. *Helminthosporium tiara* B. & Rav. Grev. iv, 640. On fallen branches. Car. (Rav.).
170. *Helminthosporium princeps* B. & C., Curt. Cat. 127. On living bark of *Quercus*. Car. (Curt.).
171. *Helminthosporium macrocarpum* Grev. Sc. Cr. Fl. t. 148. On branches. Car. (Curt.).
var. *caudatum* B. in *Grevillea* (*Helminthosporium caudatum* Curt.). On *Castanea* and oak. Car. (Grav.; Curt.).
172. *Helminthosporium corticale* Schw. Am. Bor. 2627. On bark of *Platanus*. Car. (Curt.).
173. *Helminthosporium apicale* B. & Br. (*Monotospora triseptata* Peck). Cooke, Hdbk., No. 1723. On rotten wood. N. Y. (Peck).
174. *Helminthosporium obtusissimum* B. & C., Grev. iv, 632. On dead wood. Car. (Curt.).
175. *Helminthosporium attenuatum* C. & P. (*clasterisporium* Peck). On cut surface of wood. N. Y. (Peck).
176. *Helminthosporium arbuscula* B. & C., Grev. iv, 630. On bark of various trees. Car. (Curt. Rav.).
177. *Helminthosporium melanosporum* B. & C., Grev. iii, 636. On fallen branches. Car. (Curt.). N. Eng. (Sprague).
178. *Helminthosporium siliquosum* B. & C., Grev. iv, 624. On branches of *Vitis* and *Smilax*. Car. (Curt.).
179. *Helminthosporium apiculatum* Corda. (*Helm. lanceolatum* Curt.). Cooke, Hdbk., No. 1722. On *Cornus* and oak. Car. (Curt.).
180. *Helminthosporium fragilissimum* B. & C., Grev. iv, 627. On dead twigs of *Smilax*. Car. (Curt.).
181. *Helminthosporium rectum* B. & C., Grev. iv, 631. On oak wood. Car. (Curt.).

182. *Helminthosporium molle* B. & C., Grev. iv, 633. On *Passiflora* and *Dolichos*. Car. (Curt.).
183. *Helminthosporium Ravenelii* Curt. Grev. iv, 628. (*Helminthosporium Hoffmanni* Berk. MSS.) On inflorescence of *Sporobolus*. Car. (Curt.; Rav.).
184. *Helminthosporium nodosum* B. & C., Grev. iv, 629. On inflorescence of *Eleusine Indica*. Car. (Curt.).
185. *Helminthosporium Petersii* B. & C., Grev. iv, 626. On leaves of *Smilax*, etc. Car. (Curt.). Ala. (Peters).
186. *Helminthosporium interseminatum* B. & Rav. Grevillea iv, 624. On *Phytolacca* and *Cicuta*. Car. (Rav.). Penns. (Mich.).
187. *Helminthosporium Pruni* B. & C., Grev. iv, 635. On bark of *Prunus serotina*. Car. (Curt.).
188. *Helminthosporium dubium* Schw., Am. Bor. 2626. On branches.
189. *Helminthosporium Berkeleyi* (*Helminthosporium dubium* B. & C., not Schw. in Grev. iv, 637). On leaves of *Viburnum* (?). Ala. (Beaum.).
190. *Helminthosporium inflatum* B. & Rav. Grev. iv, 638. On branches of *Myrica*. Car. (Rav.).
191. *Helminthosporium subfuscum* B. & C., Grev. iv, 639. On dead branches. N. J. (M. J. B.).
192. *Helminthosporium olivaceum* B. & R., Grev. iv, 625. On leaves of *Gleditschia*. Car. (Rav.). Penns. (Schw.).
193. *Helminthosporium foliculatum* Corda. Grev. iv. Cooke, Hdbk., No. 1707. On bark, stems, etc. Car. (Curt.). Penns. (Mich.); N. Eng. (Murray).
194. *Helminthosporium velutinum* Lk. Schw., Am. Bor. 2618. On dead twigs. Car. (Rav.). Penns. (Schw.).
195. *Helminthosporium clavariarum* Desm. Cooke, Hdbk., No. 1720. On *Clavaria*. N. Eng. (Sprague).
196. *Helminthosporium simplex* N. Schw., Am. Bor. 2623. On willow wood.
197. *Helminthosporium nanum* Lk. Schw., Am. Bor. 2621. On rotten wood.
198. *Helminthosporium subulatum* Lk. Schw., Am. Bor. 2619. On branches.
199. *Helminthosporium corniculatum* Schw., Am. Bor. 2633. On oak bark.
200. *Helminthosporium brassicola* Schw., Am. Bor. 2632. On cabbage leaves.
201. *Helminthosporium Sorghi* Schw., Am. Bor. 2631. On leaves of *sorghum*.
202. *Helminthosporium herbarum* Schw., Am. Bor. 2630. On stems of *Euphorbia*.

203. *Helminthosporium crustosum* Schw., Am. Bor. 2629. On young branches of *Catalpa*.
204. *Helminthosporium virgultorum* Schw., Am. Bor. 2628. On branches of *Morus*.
205. *Helminthosporium fasciculatum* Schw., Am. Bor. 2625. On leaves.
206. *Helminthosporium tenuissimum* Lk. Schw., Am. Bor. 2624. On stems of plants.
207. *Helminthosporium pellucidum* Lk. Schw., Am. Bor. 2622. On branches.
208. *Helminthosporium minus* Lk. Schw., Am. Bor. 2620. On branches.
- 208 a. *Septosporium velutinum* C. & Ellis in Grevillea v, p. 33. On maple wood. New Jersey (Ellis).
209. *Fusicladium dendriticum* Vekl., Syn. Myc. (*Cladosporium dendriticum* Walls). On apples and apple leaves. Car. (Curt.). Canada (Poe).
210. *Fusicladium pyrorum* Vekl. (*Fusidium pyrinum* Mart.) (*Helminthosporium pyrorum* Desm.). On pear leaves. Car. (Rav. Curt.).
211. *Podosporium rigidum* Schw., Am. Bor. 2608. On bark of various trees. Car. (Curt. Rav.).
212. *Podosporium briareus* B. & C., Grev. iv, 623. On dead branches. Car. (Curt.).
213. *Podosporium praelongum* Curt. Cat., p. 127. On branches of *Vaccinium*. Car. (Curt.).
214. *Podosporium glandicola* Schw., Am. Bor. 2609. On acorns and cups.
215. *Clasterisporium caricinum* Schw., Am. Bor. 2998. On leaves of *Carex*. N. Y. (Peck).
[*Clasterisporium pedunculatum* Peck, 25th Report is *Helminthosporium attenuatum* C. & P.]
216. *Mystrosporium Curtisii* B., in Grevillea iv, 644. (*Mystrosporium Spraguei* B. & C.) On fallen branches. Car. (Curt.). N. Eng. (Sprague).
217. *Mystrosporium rubigo* B. & C., Curt. Cat., p. 128. On *Phytolacca* and grasses. Car. (Curt.).
218. *Stemphylium fuligo* B. & C., Curt. Cat., p. 128. On branches of *Rhus*. Car. (Curt.).
219. *Triposporium elegans* Corda. Curt. Cat., p. 128. On rotten stems. Car. (Curt.).
220. *Ceratosporium fuscescens* Schw., Am. Bor. 2996. On epidermis of *Rhus*.
221. *Drepanispora pannosa* B. & C., Grev. iv, 641. Forming a cloth-like stratum on cut surface of stump. Car. (Curt.).
222. *Spondylocladium fumosum* Lk. Schw., Am. Bor. 274. On wood.
223. *Helicosporium vegetum* Lk. Schw., Am. Bor. 2634. On wood.

- 223 a. *Helicosporium olivaceum* Peck (probably a form of the last). New Jersey (Ellis).
- 223 b. *Helicosporium brunneolum* B. & C., in Grevillea, No. 540. On *Helminthosporium*. New Eng. (Sprague).
224. *Helicosporium griseum* B. & C., in Grevillea, No. 541. On fallen limbs. Car. (Curt.).
225. *Helicosporium fuscum* B. & C., in Grevillea, No. 539. Under pine wood. Car. (Curt.).
226. *Helicosporium aureum* Corda. On oak. Car. (Curt.).
227. *Helicotrichum pulvinatum* Lk. Schw., Am. Bor. 2574. On stems.
228. *Helicoma Berkeleyi* Curt. Grevillea iv, 645. (*Helicoma binale* B. & C.) On *Liquidambar* and *Fraxinus*. Car. (Curt. Rav.).
229. *Helicoma Curtisii* Berk. Grevillea iv, 646. On *Salix Babylonica* and poplar. Car. (Curt.); N. Y. (Peck).
230. *Cladotrichum scyphophorum* Corda. Pracht. f. 23. On rotten wood. Car. (Curt.).
231. *Cladotrichum uniseptatum* Cooke, in Grevillea. On branches. N. Y. (Gerard).
232. *Polythrincium trifolii* Kze., Schw., Am. Bor. 2611. On clover leaves. Car. (Curt.). N. E. (Sprague); Canada (MacLagan).
233. *Cladosporium herbarum* Lk. Schw., Am. Bor. 2588. On decayed stems, etc. Car. (Curt.); Ohio (Lea).
234. *Cladosporium effusum* B. & C., Grev. iv, 648. On *Polygonum*, *Lobelia*, etc. Car. (Curt.). Penns. (Mich.).
235. *Cladosporium compactum* B. & C., Grev. iv, 649. On leaves of *Arun-dinaria*. Car. (Curt.).
236. *Cladosporium stenosporum* B. & C., Grev. iv, 650. On apple leaves, etc. Car. (Curt.).
237. *Cladosporium microspermum* B. & C., Grev. iv, 631. On leaves of *Quercus obtusiloba*. Car. (Curt.).
238. *Cladosporium cubisporum* B. & C., Grev. iv, 632. On *Ribes*. N. E. (Frost); Maine (Rev. J. E. Blake).
239. *Cladosporium personatum* B. & C., Grev. iv, 647. On leaves of *Arachis hypogaea*. Car. (Rav.).
240. *Cladosporium episphaeria* Schw., Am. Bor. 2607. On old *Sphaeriae*.
241. *Cladosporium simplex* Schw., Am. Bor. 2606. On ash leaves.
242. *Cladosporium virgultorum* Schw., Am. Bor. 2605. On branches and twigs.
243. *Cladosporium iridicola* Schw., Am. Bor. 2604. On leaves of *Iris virginica*.
244. *Cladosporium typhae* Schw., Am. Bor. 2603. On leaves of *Typha*.
245. *Cladosporium sparsum* Schw., Am. Bor. 2602. On scapes of onion.
246. *Cladosporium lignatile* Schw., Am. Bor. 2601. On putrid wood.

247. *Cladosporium bignoniae* Schw., Am. Bor. 2600. On legumes of *Bignonia*, etc.
248. *Cladosporium maculaus* Schw., Am. Bor. 2599. On *Jasminum*.
249. *Cladosporium graminum* Lk. Schw., Am. Bor. 2596. On grasses.
250. *Cladosporium epiphyllum* Lk. Schw., Am. Bor. 2597. On leaves. N. Y. (Peck).
251. *Cladosporium grumosum* Lk. Schw., Am. Bor. 2596. On *Pteris aquilina*.
252. *Cladosporium fuscatum* Lk. Schw., Am. Bor. 2595. On stems.
253. *Cladosporium tenerrimum* Lk. Schw., Am. Bor. 2594. On bark.
[*Cladosporium fumago* Lk. Schw., Am. Bor. 2593. Consists of species of *Antennaria*, *Capnodium*, etc.]
254. *Cladosporium fuscum* Lk. Schw., Am. Bor. 2592. On stems.
255. *Cladosporium polysporum* Lk. Schw., Am. Bor. 2591. On willow branches.
256. *Cladosporium atrum* Lk. Schw., Am. Bor. 2590. On stems and cabbage stalks.
257. *Cladosporium solutum* Lk. Schw., Am. Bor. 2589. On stems of *Euphorbia*.
258. *Scolicotrichum virescens* Lk. Schw., Am. Bor. 2578. On branches.
259. *Scolicotrichum candidum* Lk. Schw., Am. Bor. 2579. On fallen leaves.
260. *Myxotrichum chartarum* Lk. Schw., Am. Bor. 2581. On paper.
261. *Myxotrichum simile* B. & C., Grev. 691. On decaying *Arundinaria*. Car. (Curt.).
262. *Macrosporium punctiforme* Berk. Hook, Journ. On dead stems of *Rubus*. Ohio (Lea).
263. *Macrosporium pinguedinis* Berk. Curt. Cat., p. 128. On dead stems. Car. (Curt.); Ohio (Lea).
264. *Macrosporium chartarum* Peck. 25th Report, p. 93. On paste-board. N. Y. (Peck).
265. *Macrosporium brassicae* Berk. (Doubtless only *Macrosporium cheiranthi* Fr.). On cabbage leaves. N. Y. (Peck).
var. *CIRCINANS* B. & C. (*Macrosporium circinans* Curt.). On cabbage leaves. Car. (Curt.).
var. *ECHINELLUM* B. & C. (*Macrosporium echinellum* Curt.). On leaves of *Platanus*. Car. (Rav.).
267. *Macrosporium antennaeforme* B. & C., Grev. iv, 642. On leaves of *Celtis*. Ala. (Peters).
268. *Macrosporium stilbosporoideum* B. & C., Grev. iv, 643. On leaves of *Crataegus*. Ala. (Beaumont).

269. *Streptothrix atra* B. & C., Grev. iv, 654. On Juniper, etc. Ala. (Peters); Mass. (Sprague); Penns. (Mich.); Car. (Curt.); N. Y. (Gerard).
270. *Streptothrix abietina* Peck (is not a *Streptothrix*). On spruce. N. Y. (Peck).
271. *Colletosporium atrum* Lk. Schw., Am. Bor. 2558. On trunks of *Smilax*.
272. *Colletosporium umbrinum* Lk. Schw., Am. Bor. 2557. On rotten bark.
273. *Arthrimum carieicola* Lk. Schw., Am. Bor. 2612. On *Carices*.
274. *Sporophlaeum graminum* Lk. Schw., Am. Bor. 2615. On leaves of *Zizania*.
275. *Gonatosporium puccinioides* Lk. (Schw., Am. Bor. 2614.) On leaves of *Carex*.
276. *Conoplea hispidula* Lk. Schw., Am. Bor. 2616. On *Panicum*.
277. *Camptoum curvatum* Lk. Schw., Am. Bor. 2613. On *Scirpus lacustris*.
278. *Sporodum atropurpureum* B. & C., Grev. iv, 653. On roots of *Arun- dinaria*. Car. (Curt.).
279. *Chloridium candidissimum* Schw., Am. Bor. 2587. On putrid wood.
280. *Chloridium griseum* Lk. Schw., Am. Bor. 2586. On putrid bark.
281. *Chloridium viride* Lk. Schw., Am. Bor. 2585. On rotten wood.
282. *Acladium densissimum* Schw., Am. Bor. 2584. On trunks and chips.
283. *Acladium microspermum* Lk. Schw., Am. Bor. 2583. On rotten branches.
284. *Acladium conspersum* Lk. Schw., Am. Bor. 2582. On trunks.

It is uncertain what the following may be which are referred by Schweinitz to species of *Dematium* :

285. *Dematium cinnamomeum* Schw., Am. Bor. 2787.
286. *Dematium lanatum* Lk. Schw., Am. Bor. 2786. Interior of trunks.
287. *Dematium corium* P. Schw., Am. Bor. 2785. In fissures of trunks. Penns. (Schw.); Car. (Curt.)
288. *Dematium aeruginosum* P. Schw., Am. Bor. 2784. Penns. (Schw.); Car. (Curt.).
289. *Dematium papyraceum* Lk. Schw., Am. Bor. 2783. On rotten wood. Car. (Schw.).
290. *Dematium episphericum* Schw., Am. Bor. 2782. On old *Sphoeria*.
291. *Dematium smilacis* Schw., Am. Bor. 2781. On *Smilax*. Car. (Schw.).
292. *Dematium ramosum* Schw., Am. Bor. 2780. On branches of *Andro- meda*.
293. *Dematium lentaceum* Schw., Am. Bor. 2779. On trunks.
294. *Dematium aluta* Lk. Schw., Am. Bor. 2778. Interior of trunks.
295. *Dematium strigosum* Lk. Schw., Am. Bor. 2777. On rotten walnut wood.

- 296. *Dematium subcosticale* Lk. Schw., Am. Bor. 2776. On bark.
- 297. *Dematium cornutum* Lk. Schw., Am. Bor. 2775. On willow branches.
- 298. *Dematium muscorum* Lk. Schw., Am. Bor. 2774. On mosses. N. E.
(Murray).
- 299. *Dematium rupicola* Lk. Schw., Am. Bor. 2773. On lichens.
- 300. *Dematium rupestre* Lk. Schw., Am. Bor. 2772.
- 301. *Dematium jubatum* Lk. Schw., Am. Bor. 2771.
- 302. *Dematium badium* Lk. Schw., Am. Bor. 2770. On branches.
- 303. *Dematium mollissimum* Lk. Schw., Am. Bor. 2769. On roots.
- 304. *Dematium nigrum* Lk. Schw., Am. Bor. 2768. On bark.

It is probable that very few, if any, of the last twenty species are autonomous fungi.

= No. 9 =

CHAMPIGNONS

Rapportés en 1880 d'une excursion botanique en Egypte et en
Palestine

Par M. WILLIAM BARBEY

(Extrait de la *Revue Mycologique*, janv. 1881, t. III, p. 23-25).

CHAMPIGNONS

Rapportés en 1880

D'UNE

EXCURSION BOTANIQUE EN EGYPTE ET EN PALESTINE

PAR

M. WILLIAM BARBEY

PARIS

J.-B. BAILLIÈRE et FILS

LIBRAIRES DE L'ACADÉMIE NATIONALE DE MÉDECINE

Rue Hautefeuille, 19

—
1881

28
2.5
0.5
1.00

CHAMPIGNONS

Rapportés en 1880 d'une excursion botanique en Egypte et en Palestine

Par M. William BARBEY

M. William Barbey, de Valleyres (Suisse), a accompli au printemps de cette année un voyage botanique dans le Levant. Il a parcouru le désert d'Egypte en Syrie, la Palestine méridionale, les terres qui avoisinent la mer Méditerranée, la Judée et les rives du Jourdain, en remontant au Sud-Est vers le rivage stérile de la mer Morte. Le but principal de l'exploration de M. Barbey était la récolte de plantes phanérogames. La cryptogamie, et notamment les champignons, ont tenu une bien petite place dans les préoccupations du collecteur. Le bouquet qu'il a fait un peu à notre intention le témoignerait; mais les proportions modestes de ce bouquet nous causent un vif regret à cause de l'importance réelle qu'il conserve. « Tout cela est bien peu, nous disait M. Barbey le 4 novembre dernier, en nous envoyant ses *Fungi*, mais vous savez la difficulté qu'on rencontre pour récolter de front des phanérogames et des cryptogames, surtout lorsqu'on chemine parfois au galop du cheval..... » La mycologie de la Palestine et de l'Egypte est pour ainsi dire encore à traiter. Le sujet est neuf et très vaste, malgré les tentatives heureuses du Dr Haussknecht, remontant à une douzaine d'années, et celles du Dr Georg. Schweinfurth, déterminées par M. de Thumen (*Grevillea* VI p. 102; VIII, p. 49 et *Flora* 1880) (1).

Sur douze numéros de sa récolte, M. W. Barbey compte trois nouveautés. Les deux premières ont été étudiées de concert avec nous, par notre savant ami, M. Carol. Kalchbrenner, et nous avons la bonne fortune d'être le premier à les faire connaître. Elles rappelleront le souvenir de MM. Barbey et Boissier si sympathiques à tous les botanistes. Notre planche XV, esquissée par M. Kalchbrenner, et reproduite en

(1) Voir l'analyse de ces documents dans notre *Revue*, t. I, p. 9 et t. III, p. 42.

couleurs par l'habile dessinateur Ch. Cuisin, montre ces nouvelles espèces.

C. ROUMEGUÈRE.

Agaricus (Naucoria) verruacti Fr. Hym. Eur. Ed. 11. p. 260. — In arena deserti Ægyptiaco-Syriaci, leg. W. Barbey, martio 1880. n° 1065. — Ab. *Ag. pèdiade* Fr. Stipite brevior, crassior, sursum attenuato distinctus.

Coprinus Barbeyi Kalch. *sp. nov. Revue Myc. Tab. XV f. I.*

Subsolitarius, pileo carnoso-membranaceo, hemisphærico-ovato, demum expanso, 1-2 unc. lato, squamis magnis, crassiusculis persistentibus pulvescentibus imbricato: stipite firmo, fistuloso, cylindrico ad duas unc. alto, 3-4 lin. crasso aequali vel sursum deorsumve attenuato hinc subventricosus, albido fuscus, ipsa basi plerumque in discum dilatato et mycelio radiceformi arenam in conum obversum conglobante; lamellis uncinato adnatis, lanceolatis, confertis diu persistentibus, atris. Spor. 0,013 × 0,020 mm. amplis.

In arena deserti Ægypt.-Arabici ad Oasim « *Atsum-Monga* » in arena deserti Ægypt.-Syriaci ad « *Bir, Abid, Rûk* » et « *Bir et Abid*; — in arena Amalekitorum, ad fines Palestinæ meridionalæ Legit W. Barbey, sub. n°s 1057, 1058, 1059, 1060 Martio 1880.

Fungus hic in arena desertorum, per Camelos etc. Stercorata haud raro obvius videtur. Fors ac *Coprinus imbricatus* Rabent. (*Sitzungsbericht der Ges. Isis in Dresden* 1870. IV.) quem D^r Haussknecht in deserto Mesopotamiæ (1867) legit et cum quo nostram speciem jungerem, ni ab illo pileo haud conico et stipite haud striato differret.

N° 1063 (coll. Barbeyi), pertinet etiam ad *Cop. Barbeyi* Kbr. (vid. Tab. XV, f. 1 b.) Sed fungus immaturus est, lamellis adhuc albis et stipite quoque floccoso-squamoso.

Tulostoma Boissieri Kalch. *sp. nov. Revue Myc. Tab. XV f. 2.*

Proximum *Tulostomati fimbriato* Fries, a quo statura robustiore et stipitis: parte superiore longitudinaliter sulcato striato differt. — Peridium globoso-depressum, nucis Avellanæ majoris amplitudine, glabriusculum, flavescens, basi appendiculi prominente stipitem amplexans. Stipes 2 poll. longus, 2 lin. et ultra crassus, nudiuseculus albus, ad basim volva membranacea, lacera faciliq. disparente auctus. Sporæ ex argillaceo-sordide rubentes.

In arena deserti Ægyptiaco-Syriaci. *El arish* leg. W. Barbey (n° 1062), 25 mars 1880.

Uromyces concentricus Lev. Ann. sc. nat. — In fol. *Scillæ maritimæ* L. — « *Bir Mabrouhy* » In deserto Ægyptiaco-Syriaco, 23 martio 1880; coll. Barbey, n° 1053. Obs. Teleutosp. paulo majores (quam in *U. muscari*), hyal. 25-18.

Ustilago Vaillantii Tul. Ann. sc. nat., 1847. XII T. 3 f. 15-19 — In antheris *Scillæ Trifoliatæ* Desf. — In cultis derelictis calcareis « *Mariut* » prope Alexandriam Ægypto Infer. 4 Martio Coll. Barbey, n° 1051. — Spor. irregul. 11-5; 9-10; 11-10, oliv.

Ustilago Carboi Tul. Ann. sc. nat. (1847) T. 3 f. 1-12. Var. Barbeyi Nob. (Sporæ gossæ 5-6 micr., d., leves, fuliginæ, diu intra glumas

inclusæ). In ovariis *Triticæ vulgaris*? Oasis « *Aïoun Monça* » ni deserto Ægyptico-Arabico. 16 martio. — Coll. Barbey, n° 1055.

Puccinia paraphysaria Bagn. p. p. (*Monogr. Pucc.* p. 75 fig. 323-324. in *Kæleria*) excl. synonym. — *P. Striformis* f. *Laguri* Sacc. Mich. II. 151. *P. Lolii* Pass. Funghi Parm. Ustil. e Ured. p. 250, nec West. (que formam *coronalam* spectat, teste auctore Cf. Kickx Flandr. II. p. 56). *P. Anomala* Rost. in Thiim M. U. n° 831! paraphysibus prædita est et a *P. paraphysaria* non differt nisi teleutosporis simplicibus copiosioribus, quæ tamen etiam in *P. paraphysaria* non desunt. — Hab. in foliis *Kæleria Berythriæ* in Ægypt. Apr. 1880, n° 1108 (Barbey). — Teleutosporæ clavatæ, 50-60 $\approx$ 22, apice obluso incrassatæ et obscuriores, cinnamomeæ, loculo inferiore cuneato; spicite tereti 20 $\approx$ 6 rufescente suffultæ et paraphysibus densis paliformibus initio hyalini dein cinnamomei, sursum incrassatis cinctæ.

Melampsora Euphorbiæ. (Pers) Cast. T. mars. T. 5. — Ad fol. Euphorbiar. variar. Cultis arenosis, a « Heliopolis » (Ægypt. Inf.) prope « Hahiram » 12 martio — coll. Barbey, n° 1045.

Æcidium Euphorbiæ Pers Syn. p. 211 f. E. punctatæ (præcipue punctiperidiis candidis differt.) Ad fol. *Euphorbiæ punctatæ* Desf. — Oasis « *Hatich* » in deserto Ægyptiaco-Syriaco. 22 martio — coll. Barbey, n° 1050.

Æcidium cressæ de. fl. fr. ad fol. *Cressæ creticæ* L. — In arvis, circa Larnaca Cypri; 28 aprile 1880. — Coll. Barbey, n° 1052.

Æcidium (e sect. *Tubularium* Bon.) *Barbeyi* Roum. *Oomyces Barbeyi*, *Revue mycol.* 1880, p. 196 (1) Tab. XV f. 3. (extremis. d'une feuille grand. nat.; fragment de la même feuille et acidiospores grossis. — Coll. Barbey. n° 850.

Cellulæ pseudoperidii angulosæ 20-25 micr. d., punctulatæ, sub hyalinæ — *Æcidiosporæ* globoso-inaequales 18-20 micr. d. subhyalinæ, guttis 1-4 aureis fætæ.

In caul. et fol. viv. *Asphodeli fistulosæ* L. — Palestina-Syria « Jéricho » et Mer Morte » 9 aprile.

Cladosporium herbarum Nees Syst. 1 p. 66, 67; Tab. V. f. 64. B. f. *Phœnicis* (Stat. obsoletum) in petiol. *Phœnicis dactyl.* Vagans ad Arenam deserti Ægyptiaco-Lybii, ad occident « Gizzet » Pyramidum. 11 martio. — Coll. Barbey. n° 1066.

(1) C'est la plante que nous avons reçue au mois d'octobre dernier de M. W. Barbey, en un échantillon fort peu avancé en maturité. Les exemplaires plus développés, ceux que nous avons reçus depuis de notre généreux correspondant et que nous distribuons en ce moment dans notre *Enciclopedia*, n° 1172 ont été analysés de nouveau et nous avons pu ranger définitivement cette nouveauté mycologique dans la section *Tubularium* Bonn. du genre *Æcidium*. Nous saisissons cette occasion pour rectifier aussi l'indication donnée par nous à propos de la récolte de l'*Oomyces*? que M. E. Boissier de Genève, accompagnait M. W. Barbey. M. E. Boissier, notre savant correspondant, n'a point fait partie des récentes excursions entreprises par M. W. Barbey. Il n'est pas revenu en Orient depuis son voyage bien connu de l'année 1846.

Harkness & Moore

Catalogue of the Pacific Coast Fungi.

[Read before the California Academy of Sciences Feb. 2, 1880.]

INTRODUCTION.

Previous to the year 1876 nothing had been accomplished in the study of the fungi of this Coast, with the exception of a few species collected by the Wilkes expedition, which were at that time referred to Professor Torrey for identification. But four or five of these were determined and of these but one or two can now be traced. Since that period, although the flowering plants have been most thoroughly studied, the fungi of this Coast have been entirely neglected.

Fully realizing the importance of the undertaking, not only for its scientific value, but also from an economic stand-point, we, four years since, determined to devote our leisure to making a collection of the Pacific Coast Fungi. To those who have given their time to scientific pursuits we need not say that this has been a work of ever-increasing interest. While we have not been able to visit every portion of our Coast, it will be observed that our explorations have extended over quite a wide range of territory—from Mt. Shasta on the north to Fort Yuma on the south—from the seashore to the eastern limits of the Sierras.

In reference to the geographical distribution of the fungi, it will be seen that most of our species found upon the low lands are common to both Europe and America, whilst of those growing along the Sierras, many are the same as the Alpine species of Europe; others are peculiar to our own Coast. Again, it will be seen, that the hot and arid desert at the south yields species common to Africa. The distribution we have aimed to make clear by giving in every instance the locality where found. We have, in the case of the Hymenomycetes and other edible fungi, indicated the fact by appending an *E*. We have also added the habitat of our *parasitic fungi*, to aid the student, and have, as will be observed, indicated the *new species*.

We do not claim for this catalogue that it contains anything like a full list of the fungi of this Coast, or even of our own collection. It is merely a catalogue of such as we know to be found here, and such as we have fully determined. A large mass of material still remains in our hands for study and determination, as our leisure may allow. We give this list, the first of the kind ever published on this Coast, in the hope that it may be of service to others and incite them to a study of this most interesting department of botanical research. In the cave fungi, or fungi of our mines, comparatively little has been done. Want of a literature on this subject has been a great hindrance to us in this work. The single cave fungus placed in the addenda is so striking that we have deemed it worthy of an insertion. There are others which we are at work upon, and in due time hope to report.

We have followed in the first part of this catalogue the arrangement of the genera as given by Fries in his Hymenomycetes Europaei. For the rest, we have adopted the order, as far as practicable, as given in Cooke's

Hand-book. Special mention should be made here of the valuable assistance rendered us in the determination of species, by Messrs. Cooke, Plowright, Phillips and Vize of England. To these gentlemen our hearty thanks are tendered. To the San Francisco Microscopical Society, for the use of its valuable library and plates, we owe much.

H. W. HARKNESS, M. D.
JUSTIN P. MOORE, A. M.

Family I. HYMENOMYCETES.

Order I. AGARACINI.

I. AGARICUS.

Series I. LEUCOSPORI.

I. AMANITA.

I. (AMANITA.)

- adnatus. Smith. San Raf.
- excelsus. Fr. San Raf.
- nitidus. Fr. San Raf.
- pantherinus. D. C. San Fran.
- phalloides. Fr. San Raf.
- strangulatus. Fr. San Raf.
- strobiliformis. Fr. rare. San Raf.
- vagitanus. Bull. San Raf.
- vernus. Bull. San Raf.
- virosus. Fr. San Raf.

II. LEPIOTA.

A. (LEPIOTA.)

- cristatus. Fr. San Raf.
- excoriatus. Schaeff. E. San Raf.
- illinatus. Fr. E. San Raf.
- procerus. Scop. E. San Fran.
- rachodes. Vltt. E. San Raf.

III. ARMILLARIA.

A. (ARMILLARIA.)

- constrictus. Fr. San Raf.
- melleus. Vahl. E. San Raf.

IV. TRICHOLOMA.

A. (TRICHOLOMA.)

- albus. Schaeff. San Raf.
- carneus. Bull. San Raf.

colossus. Fr. Nevada.
 columbetta. Fr. San Raf.
 compactus. Fr. San Raf.
 equestris. Linn. San Raf.
 gambosus. Fr. San Raf.
 imbricatus. Fr. San Raf.
 leucocephalus. Kromb. San Raf.
 panæolus. Fr. San Raf.
 personatus. Fr. San Raf.
 pessundatus. Fr. San Raf.
 portentosus. Fr. San Raf.
 resplendens. Fr. San Raf.
 sejunctus. Sowerb. Howards.
 sudus. Fr. San Raf.
 sulfureus. Bull. San Raf.
 terreus. Schaeff. San Raf.
 tigrinus. Fr. San Raf.

V. CLITOCYBE.

A. (CLITOCYBE.)

cyathiformis. Fr. Howards.
 dealbatus. Sowerb. *E.* San Raf.
 fragrans. Sowerb. common. San Raf.
 fumosus. Pers. San Raf.
 geotropus. Bull. *E.* San Raf.
 infundibuliformis. Schaeff. San Raf.
 laceatus. Scop. San Fran.
 maximus. Fr. San Raf.
 nebularis. Batsch. *E.* San Fran.
 odorus. Bull. *E.* San Raf.
 phyllophilus. Fr. San Raf.
 trullæformis. Fr. San Raf.
 tumulosus. Kalchbr. San Raf.

VI. COLLYBIA.

A. (COLLYBIA.)

butryaceus. Bull. San Raf.
 cirrhatus. Fr. San Raf.
 esculentus. Wulf. *E.* San Raf.
 exculptus. Fr. San Raf.
 fusipes. Bull. *E.* San Raf.
 longipes. Bull. *E.* San Raf.
 muscigenus. Schum. San Raf.
 ocellatus. Fr. San Raf.
 racemosus. Pers. San Raf.
 radicans. Rabh. *E.* San Raf.
 tuberosus. Bull. San Raf.
 velutipes. Curt. San Raf.

VII. MYCENA

A. (MYCENA.)

aciula. Schaeff. San Raf.
amictus. Fr. San Raf.
capillaris. Schum. San Fran.
citrinellus. Pers. San Raf.
corticola. Fr. San Raf.
debilis. Fr. San Raf.
dilatatus. Fr. San Raf.
echinipes. Lasch. Blue Cañon.
filopes. Bull. San Raf.
hiemalis. Fr. San Raf.
Iris. Berk. Sausalito.
juncicola. Fr. San Raf.
lacteus. Fr. San Raf.
purus. Pers. Howards.
pterigenus. Fr. San Raf.
rugosus. Fr. San Raf.
sanguineolentus. A. & S. San Fran.
stipularis. Fr. San Fran.
strobilinus. Fr. Howard.

VIII. OMPHALIA.

A. (OMPHALIA.)

fibula. Bull. San Raf.
gracillimus. Weinm. Howards.
integrellus. Pers. San Raf.
oniscus. Fr. *E.* San Raf.
pyxidatus. Bull. San Raf.
scyphoides. Fr. San Raf.

IX. PLEUROTUS.

A. (PLEUROTUS.)

applicatus. Batsch. Sausalito.
circinatus. Fr. San Raf.
hypnophilus. Berk. San Raf.
limpidus. Fr. San Raf.
mutilus. Fr. San Raf.
ostreatus. Jacq. *E.* San Raf.
pometi. Fr. *E.* San Raf.
salignus. Abbild. San Raf.
serotinus. Schrad. Sausalito.
sestonoides. Cke. Blue Cañon.
striatulus. Sausalito.
ulmarius. Bull. *E.* San Raf.

Series II. HIPORHODII.

X. VOLVARIA.

A. (VOLVARIA.)

- bombycinus. Schaeff. *E.* San Raf.
 gloiocephalus. D. C. San Raf.
 speciosus. Fr. San Raf.

XI. ANNULARIA.

XII. PLUTEUS.

A. (PLUTEUS.)

- cervinus. Schaeff. San Raf.

XIII. ENTOLOMA.

A. (ENTOLOMA.)

- clypeatus. Huds. San Raf.
 madidus. Fr. San Raf.
 repandus. Bull. San Raf.
 sericellus. Fr. San Raf.

XIV. CLITOPILUS.

A. (CLITOPILUS.)

- orcella. Bull. *E.* San Raf.
 prunulus. Scop. *E.* San Raf.

XV. LEPTONIA.

A. (LEPTONIA.)

- chalybæus. Pers. San Raf.
 serrulatus. Pers. San Raf.

XVI. NOLANEA.

XVII. ECCILIA.

XVIII. CLAUDOPUS.

A. (CLAUDOPUS.)

- depluens. Batsch. San Fran.
 variabilis. Pers. San Raf.

Series III. DERMINI.

XIX. PHOLIOTA.

A. (PHOLIOTA.)

- marginatus. Batsch. Sausalito.
 mutabilis. Schaeff. *E.* rare. San Raf.
 radicosus. Bull. San Raf.
 squarrosus. Müll. *E.* San Fran.

XX. INOCYBE.**A. (INOCYBE.)**

- cæsariatus*. Fr. San Raf.
fastigiatus. Schæff. San Fran.
rimosus. Bull. Sausalito.
sambucinus. Fr. San Raf.

XXI. HEBELOMA.**A. (HEBELOMA.)**

- crustuliniformis*. Bull. San Raf.
glutinosus. Lindgr. San Raf.
mesophæus. Fr. San Raf.
sinapizans. Fr. San Raf.
versipelles. Fr. San Raf.

XXII. FLAMMULA.**A. (FLAMMULA.)**

- lupinus*. Fr. San Fran.
sapinus. Fr. San Raf.

XXIII. NAUCORIA.**A. (NAUCORIA.)**

- erinaceus*. Fr. San Raf.
semiorbicularis. Bull. San Raf.
triscopus. Fr. San Raf.

XXIV. PLUTEOLUS.**XXV. GALERA.****A. (GALERA.)**

- confertus*. Bolt. San Raf.
hypnorum. Batsch. San Raf.
lateritius. Fr. San Raf.

XXVI. TUBARIA.**A. (TUBARIA.)**

- muscorum*. Hoff. San Raf.

XXVII. CREPIDOTUS.**A. (CREPIDOTUS.)**

- mollis*. Schæff. San Raf.
rubi. Berk. San Raf.

Series IV. PRATELLI.

XXVIII. CHITONIA.

XXIX. PSALLIOTA.

A. (PSALLIOTA.)

- arvensis. Schæff. *E.* common. San Raf.
 campestris. Lin. *E.* San Raf.
 cretaceus. Fr. *E.* San Raf.
 elvensis. Berk. San Raf.
 pratensis. Schæff. *E.* common. San Raf.
 silvaticus. Schæff. *E.* San Raf.

XXX. STROPHARIA.

A. (STROPHARIA.)

- æruginosus. Curt. San Raf.
 caput medusæ. Fr. rare. San Raf.

XXXI. HYPHOLOMA.

A. (HYPHOLOMA.)

- appendiculatus. Bull. San Raf.
 capnoides. Fr. San Raf.
 coronatus. Fr. San Raf.
 dispersus. Fr. San Raf.
 epixanthus. Fr. San Raf.
 fascicularis. Fr. common. San Raf.
 lacrymabundus. Bull. San Raf.
 subalteritius. Schæff. San Raf.
 velutinus. Pers. San Raf.

XXXII. PSILOCYBE.

A. (PSILOCYBE.)

- corneipes. Fr. rare. San Raf.
 ericæus. Pers. rare. San Raf.
 foenisecii. Pers. rare. San Raf.
 spadiceus. Fr. San Raf.
 squalens. Fr. San Raf.

XXXIII. PSATHYRA.

A. (PSATHYRA.)

- corrugis. P. San Raf.
 noli-tangere. Fr. San Raf.

Series V. COPRINARIÆ.

XXXIV. PANÆOLUS.

A. (PANÆOLUS.)

- campanulatus. Linn. San Raf.
 fimiputris. Bull. San Raf.
 fimicola. Fr. San Raf.

XXXV. PSATHYRELLA.

A. (PSATHYRELLA.)

- subatratus. F. San Raf.

II. MONTAGNITES.

III. COPRINUS.

C.

- atramentarius. Fr. *E.* common. San Fran.
 comatus. Fr. *E.* common. San Fran.
 deliquescens. Fr. San Raf.
 fimitarius. Fr. San Fran.
 micaceus. Fr. San Fran.
 ovatus. Fr. San Raf.

IV. BOLBITUS.

B.

- conocephalus. Fr. San Raf.
 fragilis. Fr. San Raf.
 tener. Berk. San Raf.

V. CORTINARIUS.

Tribe I. PHLEGMACIUM.

C. (PHLEGMACIUM.)

- callochrous. Fr. San Raf.
 purpurascens. Fr. San Raf.
 scaurus. Fr. San Raf.

Tribe II. MYXACIUM.

C. (MYXACIUM.)

- collinatus. Fr. San Raf.

Tribe III. INOLOMA.

C. (INOLOMA.)

- violaceus. Fr. San Raf.

Tribe IV. DERMOCYBE.**C. (DERMOCYBE.)**

cinnamomeus. Fr. *E.* San Raf.

Tribe V. TELAMONIA.**Tribe VI. HYDROCYBE.****C. (HYDROCYBE.)**

sp. (?)

VI. GOMPHIDIUS.**G.**

gracilis. Berk. San Raf.

viscidus. Fr. San Raf.

VII. PAXILLUS.**P.**

giganteus. Fr. *E.* rare. San Raf.

involutus. Fr. San Raf.

nuda. Bull. San Raf.

VIII. HYGROPHORUS.**H.**

calyptræformis. B. & Br. rare. San Raf.

distans. Berk. rare. San Raf.

eburneus. Fr. *E.* San Raf.

pratensis. Fr. San Raf.

puniceus. Fr. common. San Raf.

IX. LACTARIUS.**L.**

chrysorrhæus. Fr. San Raf.

controversus. Pers. San Raf.

deliciosus. Linn. *E.* common. San Raf.

insulus. Fr. *E.* San Raf.

mitissimus. Fr. San Raf.

piperatus. Fr. *E.* San Raf.

volemus. Fr. *E.* San Raf.

zonarius. Fr. San Raf.

X. RUSSULA.**R.**

adusta. Fr. *E.* San Raf.

alutacea. Fr. *E.* San Raf.

decolorans. Fr. San Raf.

delica. Fr. San Raf.

emetica. Fr. San Raf.

heterophylla. Fr. *E.* San Raf.
 lactea. Fr. San Raf.
 lepida. Fr. *E.* San Raf.
 lutea. Fr. San Raf.
 ochracea. Fr. San Raf.
 rubra. Fr. San Raf.
 sanguinea. Fr. San Raf.
 vesca. Fr. San Raf.

XI. CANTHARELLUS.

C.

aurantiacus. Fr. San Raf.
 cibarius. Fr. *E.* San Raf.
 crispus. Fr. San Raf.

XII. ARRHENIA.

XIII. NYCTALIS.

XIV. MARASMIUS.

M.

epiphyllus. Fr. San Raf.
 oreades. Fr. *E.* San Raf.

XV. LENTINUS.

XVI. PANUS.

P.

stipticus. Fr. San Raf.

XVII. XEROTUS.

XVIII. TROGIA.

T.

crispa. Fr. San Raf.

XIX. SCHIZOPHYLLUM.

S.

commune. Fr. San Raf.

XX. LENZITES.

L.

abietina. Fr. Howards.
 confragosa. Fr. Howards.
 flaccida. Fr. Howards.
 sepiaria. Fr. Howards.

XXIII. POLYPORUS.

P.

- abietinus. Fr.
 On cedar. Big Trees.
 adustus. Fr. San Raf.
 annosus. Fr.
 On logs. Big Trees.
 argillaceus. n. sp. Cke.
 On rotting oak. Colfax.
 brumalis. Fr. San Raf.
 carneus. Nees.
 On fir trees. Streetens.
 cervinus. Pers. San Raf.
 cinnabarinus. Fr. San Raf.
 confluens. Fr. San Raf.
 ferruginosus. Fr.
 On charred redwood. Streetens.
 fomentarius. Fr.
 On laurel. San Raf.
 hirsutus. Fr.
 On laurel. San Raf.
 hispidus. Fr. San Raf.
 hybridus. B. & Br. San Raf.
 igniarius. Fr.
 On laurel. San Raf.
 Kalakaua. n. sp. Moore.
 On a boat, sea side Sandwich Islands.
 labyrinthicus. Schw.
 On pinus contorta. Summit.
 obvolutus. n. sp. Cke.
 On fir logs. Blue Cañon.
 perennis. Fr.
 On oak. San Raf.
 quercinus. Fr.
 On oak. San Raf.
 sulphureus. Fr. E.
 On redwood. San Raf.
 varius. Fr.
 On oak. San Raf.
 velutinus. Fr.
 Oak branches. San Raf.
 versicolor. Fr.
 On oak. Colfax.
 violaceus. Fr.
 On oak. San Raf.
 vulgaris. Fr.
 On oak. San Raf.

XXIV. TRAMETIS.**T.**

cinabarinum.

On oak. San Raf.

Bulliardii. Fr. San Raf.

serpens. Fr. San Raf.

XXV. DAEDALEA.**D.**

quercina. Pers.

On oak. San Raf.

unicolor. Fr.

On wood. common. San Raf.

vorax. Harkness.

On libocedrus. Blue Cañon.

XXVI. HEXAGONA.**XXVII. FAVOLUS.****XXVIII. MERULIUS.****M.**

confluens. Schw.

On decaying spruce. Summit.

corium. Fr.

On oak bark. San Raf.

lacrymans. Fr.

Common. San Raf.

molluscus. Fr.

On redwood. San Raf.

rufus. Pers.

tremellosus. Schw.

On logs. Summit.

XXIX. POROTHELIUM.**XXX. SOLENIA.****S.**

candida. Pers.

On liber of fir. Streetens.

fasciculata. Pers.

On redwood. San Raf.

ochracea. Hoff.

On oak bark. San Fran.

Order III. HYDNEI.

XXXI. HYDNUM.

H.

alutaceum. Fr. San Raf.

caput ursi. Fr. San Raf.

chlorinum. n. sp. Cke.

On maple. Colfax.

coralloides. Scop.

E. common. San Raf.

cyathiforme. Schaeff. San Raf.

niveum. P.

On deadwood. Big Trees.

ochraceum. P. San Raf.

ramaria. Fr.

On oak. San Raf.

repandum. L. *E.* San Raf.

XXXII. HERICIUM.

XXXIII. TREMELLODON.

T.

gelatinosum. Pers.

On redwood logs. San Raf.

XXXIV. SISTOTREMA.

XXXV. IRPEX.

I.

Johnstonii. Berk.

On various woods. San Raf.

paradoxus.

On alder. San Raf.

XXXVI. RADULUM.

R.

molare. Fr.

On dead laurel. San Raf.

quercinum. Fr.

On oak. San Raf.

XXXVII. PHLEBIA.

XXXVIII. GRANDINIA.

G.

ocellata. Fr.

On madroño. Streetens.

" n. s. (?)

On redwood. Streetens.

XXXIX. ODONTIA.**XL. KNEIFFIA.****XLI. MUCRONELLA.****Order IV. THELEPHOREI.****XLII. CRATERELLUS.****C.**

- cornucopioides. Pers.
On wood. Berkeley.

XLIII. THELEPHORA.**T.**

- arida. Fr.
On pine wood. common. Blue Cañon.
— Harknessii. n. sp. Pb.
On ground among redwood. San Raf.
pallida. Pers.
On ground. Sierras.
Sowerbii. Berk.
On ground. Sierras.
terrestris. Ehrh.
On ground. Sierras.

XLIV. STEREUM.**S.**

- evolvens. Fr.
On prunus demissa. Blue Cañon.
hirsutum. Fr.
On dead logs. San Raf.
purpurellum. Fr.
On willow bark. Sac.
purpureum. Fr.
On elder. Streetens.
tabacinum. Mont.
On Baccharis. San Raf.

XLV. HYMENOCHÆTE.

- rubiginosa. Lev.
On stumps. Streetens.
tabacina. Lev.
On redwood. Streetens.
sp. (?)
On charred redwood. San Raf.

XLVI. CORTICIUM.**C.**

- caeruleum. Fr.
On redwood. San Raf.
- calceum. Fr.
On madroño. San Raf.
- carneum. n. sp. B. & Cke.
On pinus contorta. Summit.
- comedens. Fr.
On laurel. Blue Cañon.
- evolvens. Fr.
On cherry. Sac.
- incarnatum. Fr.
On cornus. Blue Cañon.
- laeve. Pers.
On laurel. San Raf.
- lactescens. B.
On willow. Sac.
- lacteam. Fr.
On redwood. Streetens.
- quercinum. P.
On oak. San Raf.
- salicinum. Fr.
On willow. Summit.
- sambuci. Fr.
On elder. Streetens.
- sanguineum. Fr.
On redwood. San Raf.
- scutellare. B. & C.
On adenostoma. Colfax.
- sulfureum. Fr.
Var. ochroideum. on elder. Sac.
- umbrinum. A. & S.
On charred redwood. Howards.

XLVII. CYPHELLA.**C.**

- capula. Fr.
On dead weeds. Berkeley.
- galatea. Fr.
On mosses. Fr.
- ochroleuca. B. & Br.
On bramble twigs. San Raf.
- villosa. Pers.
On dead weeds. Berkeley.

Order V. CLAVARIEI.

XLVIII. SPARASSIS.

XLIX. CLAVARIA.

C.

- abietina. Schum.
 On fir. San Raf.
 coralloides. Linn. San Raf.
 cristata. P.
 E. on foliage Big Trees. Big Trees.
 fastigiata. Linn.
 E. On ground. San Raf.
 flava. Schaeff.
 E. on ground under oak. San Raf.
 inquanda. P. (?) San Raf.
 pistillaris. L.
 On ground. San Raf.
 ——— n. sp. (?)

L. CALOCERA.

LI. PTERULA.

LII. TYPHULA.

T.

- filiformis. Fr.
 On leaves. San Raf.

LIII. PISTILLARIA.

Order VI. TREMELLINI.

LIV. TREMELLA.

T.

- albida. Hud.
 On dead bark. Blue Cañon.
 lutescens. Fr. common. San Raf.
 mesenterica. Retz.
 E. on redwood. San Raf.

LV. EXIDIA.

E.

- auricula Judæa. Fr.
 On wood. Sierras.
 glandulosa. Fr.
 On cherry. Sac.
 recisa. Fr.
 On pine twigs. Blue Cañon.
 saccharina. Fr. Summit.

LVI. HIRNEOLA.**H.**

- auricula Judæa, Berk. Blue Cañon.
polytricha. Mont. Sandwich Is.

LVII. FEMSJONIA.**LVIII. NAEMATelia.****N.**

- encephala. Fr.
On redwoods. Streetens.

LIX. GUEPINIA.**G.**

- helvelloides. Fr.
On ground. common. San Raf.
(?) cyphella. Fr.
On ground. rare. San Raf.
spathularia. Fr.
On ground. Sac.

LX. DACRYMYCES.**D.**

- chrysocomus. Tul.
On dead spruce. Blue Cañon.
deliquescens. Fr.
On pinus contorta. Summit.
stillatus. Nees.
On dead sequoia gigantea. Yosemite.
sp. (?)
On redwood. Streetens.

LXI. DITIOLA.**D.**

- radicata.
On dead pine branches. Blue Cañon.

Family II. GASTEROMYCETES.**Order VII. HYPOGÆI.****MELANOGASTER.**

- variegatus. Ful. E.
On or under the ground under oaks. San Raf.

Order VIII. PHALLOIDEI.**PHALLUS.**

- impudicus. Fr.
On ground. Oakland.

Order IX. TRICHOCASTRES.

BATARREA.

phalloides. P.

On sand hills south of San Fran. and at Phoenix, Arizona.

PODAXON.

carcinomale. Fr. Colorado desert.

loandensis. W. & C. Colorado desert.

GEASTER.

fimbriatus. Fr.

On ground. San Raf.

hygrometricus. P.

On ground. San Raf.

mammosus. Chw.

On ground. San Raf.

striatus. D. C.

On ground. San Raf.

BOVISTA.

ammophila. Lev.

On ground. San Raf.

nigresceus. P.

On ground. San Raf.

plumbea. P.

On ground. San Raf.

LYCOPERDON.

asperrimum. W. & C.

On ground. San Raf.

bicolor. W. & C.

On ground. San Raf.

cæspitosum. W. & C.

On ground. San Raf.

giganteum. Balsch.

E. on ground. San Raf.

gemmatum. Fr.

On ground. San Raf.

pyriforme. Schw.

On ground. San Raf.

radicata. W. & C.

On ground. San Raf.

SCLERODERMA.

verrucosum. Pers.

vulgare. Fr.

Order X. MYXOMYCETES.

LYCOGALA.

- epidendrum. Fr.
On decaying wood. Sierras.

RETICULARIA.

- maxima. Fr.
On pine stumps. Sierras.
umbrina. Fr.
On oak stumps. Stockton.

ÆTHALIUM.

- septicum. Fr.
On boards of tan vats. San Fran.

DIDERMA.

- albescens. n. sp. Ph.
On pine and oak bark. Blue Cañon.
— brunneola. n. sp. Ph.
On oak bark. San Fran.
— geasteroides. n. sp. Ph.
On dead bark. Colfax.
granulatum. Fr.
On decaying leaves. Sausalito.
— lacineatum. n. sp. Ph.
On decaying wood. Cushings.
testaceum. Schw.
On decaying wood. Cushings.
vernicosum. P.
On decaying twigs. San Fran.

CHONDRIODERMA.

- geasteroides. Ph.
On rotten wood. Colfax.

DIDYMIUM.

- cinerium. Fr.
On dead ferns. San Fran.
clavus. A. & S.
On decaying bark. San Raf.
— glaucum. n. sp. Ph.
On decaying wood. San Fran.
— granuliferum. n. sp. Ph.
On decaying wood. Blue Cañon.
squamosum. A & S.
On decaying foliage of redwoods. San Raf.

PHYSARUM.

nutaus. P.

On decaying wood. Cushings.

BADHAMIA.

inaurata. Curr.

On dead pine bark. Blue Cañon.

DIACHÆA.

elegans. Fr.

On living strawberry leaves. Sac.

COMATRICHA.

Friesiana. DeBary.

On rotten wood. Big Trees.

typhina. Roth.

On dead oak. Colfax.

STEMONITIS.

arcyriodes. Som.

On oak bark. Blue Cañon.

ferruginea. Ehr.

On dead laurel. San Raf.

fusca. Roth.

On cedar bark. Yosemite.

globosa. Schum.

On dead oak. Blue Cañon.

nigra. Fl. Dan.

On spruce chips. Blue Cañon.

obtusata. Fr.

On pinus contorta. Summit.

typhoides. D. C.

On decaying wood. Blue Cañon.

DICTYDIUM.

cernuum. Pers.

On dead pine. Yosemite.

ARCYRIA.

nutaus. Fr.

On dead oak. Sac.

punicea. Pers.

On dead laurel. Angel Is.

On pine. Yosemite.

umbrina. Schum.

On dead twigs. San Fran.

— versicolor, *n. s. p.* Ph.

On bark sequoia gigantea. Big Trees.

— vitellina, *n. s. p.* Ph.

On pinus contorta. Summit.

TRICHIA.

- chrysosperma. D. C.
 On balm of gilead, Sac.
 fallax. Pers.
 On decaying wood. Cushings.
 nigripes. Pers.
 On decaying wood. Cushings.
 turbinata. With.
 On decaying wood. Cushings.
 varia. P.
 On Pinus Lambertiana. Colfax.

Order XI. NIDULARIACEI.

CYATHUS.

- striatus. Wild.
 On ground. Sac.

CRUCIBULUM.

- vulgare. Tul.
 On dead pine. Yosemite.

NIDULARIA.

- campanula. With.
 On wood. Sierras.

SPHÆROBOLUS.

- stellatus. Tode.
 On decaying oak. Sauc.

Family III. CONIOMYCETES.

Order XII. SPHERONMEI.

LEPTOSTROMA.

- flicinum. Fr.
 On pteris aquilina. San Raf.

MACROFLODIA.

- arctostaphyli. n. sp. Vize.
 On manzanita.
 — sambucina, n. sp. Cke.
 On sambucus niger. Sac.

PHOMA.

- sp. (?)
 On bramble stems. Sac.
 concentricum. Desm.
 On yucca. Sac.

- sequoiæ, n. sp. Pl.
On dead wood. Big Trees.
verbascola. Schw.
On dead mullein stalks. Sac.

ACTINOTHYRIUM.

- graminis. Kze.
On clover leaves. Sausalito.

CRYPTOSPORIUM.

- lupini. Cke.
On lupins. San Fran.

SHHÆROPSIS.

- mutica. B. & Br.
On elder. Sac.

DIPLODIA.

- aceris. Fekl.
On sycamore twigs. Sac.
salicis. Mst.
On willows. Sac.

HENDERSONIA.

- sp. (?)
On salicornia herbacea. common. San Raf.

VERMICULARIA.

- dematium. Fr.
polygoni. Schw.
On polygonum and pie plant. Sac.

MELASMIA.

- arbuticola. n. sp. Vize.
On leaves of madroño. San Raf.

SEPTORIA.

- xanthii. Des.
On xanthium strumarium. Sac.
sp. (?)
On leaves Darlingtonia Californica. Sac.

EXCIPULA.

- hispidula. Sch.
On dead grass. Sac.
strigosa. Fr.
On dead elder branches. Streetens.

DINEMASPORIUM.

graminum. Lev.

On wild oats. Berkeley.

ASTEROMA.

Rosæ. D. C.

On living rose leaves. San Fran.

CYTISPORA.

sp. (?)

On buckeye husks. San Raf.

fugax. Bull.

On willow bark. Santa Cruz.

leucosperma. Fr.

On twigs wild cherry. Streetens.

populina. Pers.

On poplar bark. San Raf.

salicis. Rabh.

On willow. Summit.

sp. (?)

On liber redwood. Streetens.

Order XIII. MELANCONEI.

PESTALOZZIA.

planimi. n. s. p. Vize.

On leaves of Enonymus. San Raf.

NEMASPORA.

crocea. Fr.

On dead oak bark. Streetens.

GLÆSPORIUM.

carpigenum.

On buckeye. San Raf.

Order XIV. TORULACEI.

SPORIDESMIUM.

induratum. n. s. p. Cke.

On dead manzanita leaves. Yosemite.

On dead roots sequoia. Big Trees.

lepraria. B. & Br.

On dead oak bark. San Fran.

velutinum. Cke.

On dead twigs. San Fran.

Order XV. PUCINIÆI.

PHRAGMIDIUM.

- bulbosum. Sch.
On bramble leaves. San Raf.
- gracile. Grev.
On Rubus Nutkanus. San. Raf.
- mucronatum. Link.
On rose leaves. San Fran.

PUCCINIA.

- amorphæ. B. & Curt.
On amorphæ Cala. Common. San Raf.
- anemones. Schw.
On anemone. San Raf.
- angustata. Peck.
On juncus. Sac.
- artemisiæ. Fekl.
On artemisia. Sac.
- asparagi. D. C.
On asparagus stems. Sac.
- Berberidis. Mont.
On Berberis aquifolium. Fresno River.
- coronata. Cord.
On oats. San Fran.
- gayophiti. *n. s. p.* Vize.
On gayophitum. Blue Cañon.
- graminis. Pers.
On wheat and sugarcane. San Fran. and S. Is.
- Harknessii. *n. s. p.* Vize.
On zigodesmus spinosa. Mt. Rosa, Nev., 7,000 ft.
- helianthii. Schu.
On helianthus. Sac.
- hieracii. Mart.
On crepis glaucus. Alta.
- malvacearum. Mont.
On malva. San Fran.
- menthæ. Pers.
On menthæ. Sac.
- obtusa. Sch.
On white sage. Tehachepi.
- œnothera. *n. s. p.* Vize.
On œnothera densiflora. Colfax.
- polygonorum. Link.
On polygonaceæ. Sac.
- prunorum. Link.
On plum leaves. Sac.

saniculæ. Grev.

On sanicula Menziesii. San Raf.

— splendens. *n. s. p.* Vize.

On tessaria. Colorado Desert.

— striata. *n. s. p.* Cke.

vaginalium. Lk.

On polygonum.

Xanthii. Schw.

On Xanthium strumarium. Sac.

GYMNOSPORANGIUM.

biseptatum. Ellis.

On living branches of zibrocedrus. Yosemite.

Order XVI. CÆOMACEI.

SYNCHITRIUM.

fulgens. Schoeter.

On œnothera biennis. San Raf.

papillatus. Far.

On alfilaria. San Raf.

USTILAGO.

carbo. Tul.

On oats. San Fran.

bromivora. F. de W.

On grass. St. Helena.

Maydis. Tul.

On corn. Los Angeles.

utriculosa. Ful.

On polygonum aviculare. Sac.

UROCYSTIS.

occulta. Preuss.

On grass. Sac.

UROMYCES.

apiculatus. Lev.

On clover. San Raf.

appendiculata. Lev.

On vicia. Lake Co.

Betæ. Kuhn.

On beet leaves. San Fran.

Ficariæ. Lev.

On ranunculus Cal. San Raf.

Howei. Pk.

On asclepias. Lake Co.

— intricata. *n. sp.* Cke.

On gayophytum. Blue Cañon.

junci. Tul.

On juncus. Sac.

“ var. Scirpi.

On scirpus. San Raf.

limonii. Lev.

On statice limonium. San Raf.

lupini. Sac.

On lupins. San Fran.

— oblonga. n. sp. Vize.

On burr clover. Sac.

prunorum. Lk.

On peach leaves. Sac.

“ var. Amygdali.

On peach leaves. Sac.

COLEOSPORIUM.

— Madiæ. n. sp. Ck.

On madia. San Raf.

miniatum. Lev.

On wild rose. Yosemite.

MELAMPSORA.

populini. Lev.

On poplar leaves. San Raf.

salicina. Lev.

On willow leaves. Sac.

CYSTOPUS.

candidus. Lev.

On cabbage leaves. Sac.

UREDO.

convolvuli. Str.

On convolvulus. Sac.

lupini. B. & C.

On lupins. Sac.

quercus. Brond.

On oak leaves. Sierras.

scirpina. West.

On scirpus. Sac,

TRICHOBASIS.

epilobii. D. C.

On epilobium. San Fran.

helianthi. Schw.

On helianthus. Sac.

polygonorum. Lev.

On polygonum. Sac.

rubigo vera. Lev.

On wheat. San Fran.

Order XVII. *ÆCIDIACEI*.**LYCETHEA.**

populina. Lev.

On populus moniloides. Sac.

RÆSTELIA.

cancellata. Ret.

On pear leaves. San Raf.

PERIDERMIMUM

— Harknessi. n. sp. Moore.

On pinus ponderosa. Colfax.

pini. Fr.

On pinus insignis. San Fran.

ÆCIDIUM.

Berberidis. Pers.

On grass. Sac.

— gayophyti. n. sp. Vize.

On gayophitum ramosissimum. Blue Cañon.

Family IV. *HYPHOMYCETES*.Order XVIII. *ISARIACEI*.**CERATIUM.**

hidneoides. A. & S.

On rotten oak. Streetens.

Order XIX. *STILBACEI*.**OZONIUM.**

aureconium. Lk.

On rotten pine. Colfax.

On dead fir branches. Blue Cañon.

STILBUM.

fimetarium. B. & Br.

On ground. Sac.

pellucidum. Schr.

On decaying polyporus. Sac.

VOLUTELLA.

setosa. B.

On twigs. San Fran.

TUBERCULARIA.

- nigricans. Lk.
On dead fir bark. Big Trees.
vulgaris. Tode.
On locust. Sac.

FUSARIUM.

- personatum. n. sp. Cke.
On dead laurel leaves. San Raf.

Order XX. DEMATIEI.

HELMINTHOSPORIUM.

- delicatulum. B. & Br.
On umbelliferae. Sac.
macrocarpum. Grev.
On dead twigs. Sac.
nanum. Nees.
On dead herbs. Sac.
rhabdiferum. B. & Br.
On ripe peaches. Sac.
turbinatum. B. & Br.
On dead wood. Blue Cañon.
velutinum. Lk.
On dead twigs. San Raf.

CERCOSPORA.

- inquinans. n. sp. Cke.
On gymnocarpus. San Raf.

RAMULARIA.

- obovata. Fekl.
On leaves rumex. Sac.

MACROSPORIUM.

- brassicæ. B.
On decaying cabbage leaves. Sac.

POLYTHRINCIUM.

- trifolii. Kze.
On dead clover leaves. Sac.

CLADOSPORIUM.

- delicatulum. Cke.
On dead leaves. San Fran.
dendriticum. Wall.
On pear leaves. Sac.

- epiphyllum. Cord.
 On poplar leaves. San Raf.
 fumago. Lk.
 On laurel leaves. San Raf,
 herbarum. Lk.
 On typha latifolia. Sac.

Order XXI. MUCEDINES.

ASPERGILLUS.

- candidus. Lk.
 On decaying substances. Common.
 dubius. Cord.
 On rabbit dung. Sac.
 glaucus. Lk.
 On decaying leaves pinus contorta. Summit.
 mollis. B.
 On dead leaves. Blue Cañon.
 virens. Lk.
 On decaying agarics. San Raf.

PERONOSPORA.

- effusa. Grev.
 On spinach. San Fran.
 ficariæ. Tul.
 On ranunculus repens. San Raf.
 gangliiformis. B.
 On lettuce leaves. Sac.
 grisea. Ung.
 On veronica. San Raf.
 infestans. Mont.
 On potato leaves. Coast.
 nivea. Ung.
 On umbelliferae. San Raf.
 obliqua. Ck.
 On dock leaves. Sac.
 parasitica. Pers.
 On cruciferae. Sac.
 potentillæ. Sch.
 On potentilla. Sac.
 pygmæ. Ung.
 On anemone leaves. San Raf.
 sordidæ. B.
 On scrophularia Californica. San Raf.
 sparsa. B.
 On rosa Californica. San Raf.
 trifoliorum. De B.
 On clover. San Fran.

urticæ. Casp.

On nettle. San Raf.

vicæ. B.

On pear leaves. San Raf.

PENICILLIUM.

bicolor. Fr.

On decaying substances. Sac.

chartarum. Cke.

On damp wall paper. Sac.

crustaceum. Fr.

On decaying substances. Sac.

OIDIUM.

aureum. Lk.

On rotten wood. Sierras.

fasciculatum. Berk.

On decaying oranges. Sac.

favorum. B. & Br.

On honeycomb. Sac.

fructigenum. Schrad.

On apples and pears. San Fran.

Tuckeri. Berk.

On wild and cultivated grape leaves. Common.

REISSIA.

semiophora. Fresen.

On pine. Yosemite.

MYXOTRICHUM.

ochraceum. B. & Br.

On liber of redwood. Streetens.

BOTRYOSPORIUM.

diffusum. Cord.

On dead twigs. San Raf.

Order XXII. SEPEDONIEI.

SEPEDONIUM.

chrysospermum. Lk.

On decaying fungi. San Raf.

roseum. Fr.

On decaying fungi. San Raf.

FUSISPORIUM.

atrovirens. B.

On onions. Sac.

fæni. B. & Br.

On decaying grass. Sac.

roseolum. Stph.

On rotting potatoes. Sac.

Division II. SPORIDIFERA.

Family V. PHYSOMYCETES.

Order XXIII. ANTENNARIEI.

ANTENNARIA.

Guavæ. n. sp. Cke.

On leaves of guava. Sandwich Islands.

Order XXIV. MUCORINI.

MUCOR.

amethysteus. Bon.

On rotting pears. Sac.

caninus. Pers.

On dog's dung. Sac.

clavatus. Lk.

On fruit. Sac.

fusiger. Lk.

On decaying fungi. Sac.

mucedo. Linn.

On preserves. Sac.

phycomyces. B.

On oil, etc. Sac.

ramosus. Bull.

On fungi. Sac.

stolonifer. Ehrb.

On melons. Sac.

subtillissimus. B.

On onions. Sac.

tenerrimus. B.

On twigs. Sac.

TRICHODERMA.

viride. Pers.

On dead bark. Sausalito.

PILOBOLUS.

cristallinus. Tode.

On cow dung. Sac.

roridus. Schum. On cow dung. Sac.

RHYPARORIBUS.

dubius. Boud.

Cookei.

On dog's dung.

Woolhopenses. Renny.

On bird's dung. San Raf.

ACROSTALAGMUS.

cinnabarinus. Cord.

On dead weeds. Sausalito.

ENDOgone.

pisiformis. Lk.

On oak bark, San Fran.

Order XXV. **SAPROLEGNIEI.****SAPROLEGNIA.**

ferax. Gr.

On living salmon and dead flies. common.

Family VI. **ASCOMYCETES.**Order XXVI. **ONYGENEI.****ONYGENA.**

equina. Pers.

On dead horse's hoof. Sac.

Order XXVII. **PERISPORIACEI.****SPHAEROTHECA.**

castagnei. Lev.

On hop leaves. Sac.

pannosa. Lev.

On rose leaves. Sac.

PHYLLACTINIA.

guttata. Lev.

On oak leaves. Alta.

UNCINULA.

adunca. Lev.

On willow. Sac.

bicornis. Lev.

On maple leaves. San Raf.

PODOSPHÆRIA.

Kunzei. Lev.

On plum leaves. Sac.

MICROSPHÆRIA.

comata. Lev.

On euonymus. San Raf.

— fulvo-fulera. n. sp. Cke.

On spiræa dumosa. San Raf.

Grossulariæ. Lev.

On wild gooseberry. Sierras.

pencillata. Lev.

On alder leaves. San Raf.

ERYSIPHE.

communis. Schl.

On leguminosae. Sac.

graminis. D. C.

On grass. Sac.

lamprocarpa. Lev.

On plantain. Sac.

Martii. Lke.

On peas, beans, etc. common.

montagnei. Lev.

On sesbania. Fort Yuma

CHÆTOMIUM.

elatum. Kze.

On dead ash wood. Sac.

glabrum. B.

On damp straw. Sac.

EUROTIIUM.

herbarorium. Lke.

On dead bark. San Fran.

CAPNODIUM.

citri. B. & Desm.

On orange leaves. Los Angeles.

elongatum. B. & Desm.

On pear twigs. Sac.

salicini. H.

On willow leaves. Sac.

Order XXVIII. ELVELLACEI.**MORCHELLA.**

esculenta. Pers.

E. under oaks. Streetens.

conica.

E. On charred redwood. Streetens.

GYROMITRA.

esculenta. Fr.

E. under oaks. San Raf.

HELVELLA.

Californica. n. sp. Ph.

E. on ground. Blue Cañon.

crispa. Fr.

E. under oaks. San Raf.

lacunosa. Afz.

On hillsides.

VERPA.

digitaliformis. Pers.

Among redwood leaves. rare. Streetens.

SPATHULARIA.

flavida. Var. Californica. n. var. Moore.

On redwood. San Raf.

VIBRISSEA.

truncorum. Fr.

On sticks in water. Sierras.

PEZIZA.

acutipila. Kast.

On *Rubus Nutkanus*. San Raf.

acuum. Fr.

On fir leaves. Streetens.

Agassizii. var. rufipes. n. var. Ph.

On *pinus contorta*. Blue Cañon.

— alutipes. n. sp. Ph.

On dead cedar leaves. Blue Cañon.

— arida. n. sp. Ph.

On pine bark. Blue Cañon.

aurantia. Fr.

On ground. Sausalito.

badia. P.

On spruce boards. Truckee.

bicolor. Bull.

On *Rubus Nutkanus*. Blue Cañon.

bufonia. Pers.

On cow dung in dense woods. Streetens.

bulgarioides. Rab.

On dead cones sequoia. Big Trees.

carbonaria. A. & S.

On dead leaves sequoia. Big trees.

caucus. Reben.

On decaying cedar leaves. Alta.

“ var. fusco-purpurea. n. var. Ph.

- cinerea.* Batsch.
 On decaying twigs. Colfax and Streetens.
citrina. Hedw.
 On rotten wood. Streetens.
clandestina. Bull.
 On oak. Summit.
coprinaria. Cke.
 On cow dung. Truckee.
corticalis. P.
 On twigs. San Fran.
crenulata. B. & Br.
 On ground. Streetens.
cupressina. Batsch.
 On dead leaves redwood. Streetens.
cyathoidea. Bull.
 On *aralia racemosa*. Blue Cañon.
elaphines. B. & Br.
 On decaying wood. Angel Island.
epixantha. n. sp. Cke.
 On oak. Summit.
escharoides. B. & Br.
 On sticks. Blue Cañon.
fulgens. P.
 On ground. Yosemite.
fusca. P.
 On dead redwood. San Raf.
gemmea. n. sp. Ph.
 On dead redwood leaves. Streetens.
granulata. Bull.
 On rotting straw. Streetens.
hinnulea. B. & Br.
 On decaying wood. Colfax.
hyalina. P.
 On boards. Sac.
leiocarpa. Curr.
 On charred wood and earth. Streetens.
leporina. Batsch.
 On ground. Sausalito.
leucostigma. Fr.
 On oak. Colfax.
macrocystis. Cke.
 On charred wood. Streetens.
maculosa. n. sp. Ph.
 On cow dung. Colfax.
molesta. n. sp. Ph.
 On redwood chips. Streetens.
nigrescens. Ck.
 On willow bark. Summit.

- nivea*. Fr.
 On oak chips. Blue Cañon.
 — *obscura*. n. sp. Ck.
 On oak. Summit.
omphalodes. Bull.
 On ground. Oakland.
 — *phymatodes*. n. sp. Ph.
 On dead weeds. Blue Cañon.
pithya. P.
 On redwood bark. Streetens.
relicina. Fr.
 On aralia stems. Colfax.
rosæ. P.
 On dead twigs. San Fran.
rutilans. Fr.
 On damp wall. San Fran.
sanguinea. P.
 On dead oak. Yosemite.
 — *scabro-villosa*. n. sp. Ph.
 On *Rubus Nutkanus*. Blue Cañon.
scubalonta. Oke.
 On cow dung. San Raf.
scutellata. L.
 On damp ground. Streetens.
 — *secreta*. n. sp. Ph.
 On ground—woods. Blue Cañon.
 — *sequoiæ*. n. sp. Ph.
 (Hymenosecyphæ.)
 On dead leaves sequoia. Big Trees.
 — *sequoiæ*. n. sp. Ph.
 (Sarco-secyphæ.)
 On dead leaves sequoia. Big Trees.
 — *setigera*. n. sp. Ph.
 On dead stem *Saxifraga peltata*. Blue Cañon.
stercorea. P.
 On cow dung. Colfax.
strobiliina. Fr.
 On pine cones. Blue Cañon.
subhirsuta. Schw.
 On cinder heaps. San Fran.
subtillissima. C.
 On decaying spruce. Summit.
sub-urceolata. n. sp. Ph.
 On ground under sequoia. Big Trees.
 — *succina*. n. sp. Ph.
 On dead oak. Blue Cañon.
sulphurea. P.
 On dead weeds. Berkeley.

- thelebeloides. A. & S.
 On cow dung. Sac.
 uncinata. n. sp. Ph.
 On oak twigs. Blue Cañon.
 ustorum.
 On redwood. Streetens.
 venosa. Pers.
 On ground. Blue Cañon.
 vesiculosa. Bull.
 On dunghill San Fran.
 villosa. Pers.
 On dead weeds. Berkeley.
 vinosa. A. & S.
 On ground. Sierras.
 virginea. Batsch.
 On dead leaves. San Fran.

HELOTIUM.

- æruginosum. Fr.
 On dead birch. Sierras.
 agaricinum. Berk.
 On decaying wood. Yosemite.
 aureum. Pers.
 On dead laurel. Colfax.
 citrinum. Fr.
 On dead oak. Yosemite.
 claro-flavum. B.
 On oak bark. San Fran.
 epiphyllum. Fr.
 On dead leaves. Yosemite.
 fagineum. Fr.
 On decaying twigs. Blue Cañon.
 fibuliforme. Fr.
 On elm branches. Sac.
 herbarum. Fr.
 On dead stems. Blue Cañon.
 lutescens. Fr.
 On dead twigs. Blue Cañon.
 pallescens. Fr.
 On dead laurel. San Raf.
 salicellum. Fr.
 On willow. Summit.
 serotinum. Fr.
 On sticks. Summit.
 subtile. Fr.
 On fir foliage, dead. Summit.
 virgultorum. Fr.
 On dead wood. San Fran.

PATELLARIA.

- abietina. n. sp. Cke.
On fir. San Raf.
- olivacea. Batsch.
On rotten willow. Summit.
- rhabarbarina. B.
On dead bramble. Blue Canon.

TYMPANIS.

- alnea. P.
On alder twigs. Summit.
- conspersa. Fr.
On raspberry twigs. Sac.

DERMATEA.

- flavo-cinerea. n. sp. Ph.
On Cornus. Blue Cañon.

CENANGIUM.

- leptospermum. B. & C.
On fir. San Raf.
- prunastri. Fr.
On wild cherry. Yosemite.
- Ribis. Fr.
On currant twigs. Sac.
- Rubi. Fr.
On raspberry twigs. Alta.
- sequoiæ. n. sp. Pl.
On bark sequoia. Big Trees.

ASCOBOLUS.

- serugineus. Fr.
On horse dung. Sac.
- atro-fuscus. n. sp. Ph. & Pl.
On dung. Sac.
- carneus. Pers.
On decaying pine wood. Colfax.
- ciliatus. Boud.
On horse dung. Yosemite.
- furfuraceus. Pers.
On cow dung. Warm Springs.
- Kerverni. Crouan.
On cow dung. Sac.
- immersus. Pers.
On cow dung. Sac.
- incanus. n. sp. Ph.
On cow dung. Colfax.

- papillatus. Boud.
 On cow dung. Colfax.
 patellatus. Wall.
 On cow dung. Sac.
 pilosus. Boud.
 On dog dung. Sac.
 — rari-pilus. n. sp. Ph.
 On cow dung. San Raf.

BULGARIA.

- inquinans. Fr.
 On oak. San Raf.
 sarcoides. Fr.
 On bark. Sauc.

STICTUS.

- Berkleyana. De B. & Lev.
 On rotten wood. Cushings.
 — calcera. n. sp. Ph.
 On dead willow. Tehachepi.
 radiata. Fr.
 On oak twigs. Streetens.
 versicolor. Ph.
 On liber redwood. San Raf.

ASCOMYCES.

- deformans. Berk.
 On peach leaves. Sac.

Order XIX. PHACIDIACEI.

PHACIDIUM.

- internum. n. sp. Ph.
 On libocedrus. Yosemite.
 pini. Schm.
 On pine bark. Blue Cañon.
 ranunculi. Desm.
 On ranunculus. San Raf.
 Rubi. Fr.
 On Rubus Nutkanus. San Raf.
 trifolii. Boud.
 Clover leaves. Sac.

RHYTISMA.

- acerinum. Fr.
 On maple leaves. San Raf.
 — arbuti. n. sp. Ph.
 On madroño. San Raf.

- punctatum. Fr.
 On sycamore. Sac.
 salicinum. Fr.
 On willow leaves. Sac.
 On balm of gilead. Yosemite.

HYSTERIUM.

- acerinum. Frost.
 On maple. San Raf.
 angustatum. A. & S.
 On redwood. Streetens.
 arundinaceum. Schrad.
 On grass. Blue Cañon.
 commune. Fr.
 On dead stems. Summit.
 foliicolum. Fr.
 On oak leaves. Summit.
 — formosum. n. sp. Ck.
 On juniper bark. Summit.
 pinastri. Schrad.
 On pine leaves. Summit.
 virgultorum. D. C.
 On bramble stems. San Raf.

LABRELLA.

- pomi. M.
 On apples in damp grass. Sac.

Order XX. SPHÆRIACEI.

TORRUBIA.

- militaris. Fr.
 On pupa of caterpillar. Sausalito.

CLAVICIPS.

- purpurea. Tul.
 On grass seeds. Sac.

HYPOCREA.

- lenta. Fr.
 On fir boards. Sac.
 rufa. Fr.
 On redwoods. San Raf.

HYPOMYCES.

- aurantius. Tul.
 On dead fungi. Sausalito.
 luteo-virens. Fr.

- On dead fungi. Sausalito.
 rosellus. Tul.
 On dead fungi. San Raf.
 ochraceus. Tul.
 On dead fungi. San Raf.

NECTRIA.

- coccinea. Fr.
 On bark lupins. San Fran.
 cinnabarina. Fr.
 On bark. San Fran.
 cucurbitula. Fr.
 On bark. Cushings.
 episphæria. Fr.
 On bark. San Fran.
 ochraceo-pallida. R. & Br.
 On elder twigs. Streetens.
 peziza. Fr.
 On lupins. San Fran.
 umbrina. Fr.
 On decaying bean stalks. Sac.

XYLARIA.

- hypoxyton. Grev.
 On roots. Sausalito.

USTULINA.

- vulgaris. Tul.
 On dead trees. Sausalito.

HYPOXYLON.

- coccineum. Bull.
 On bark. San Fran.
 cohærens. Fr.
 On twigs. Sausalito.
 concentricum. Grev.
 On dead ash. Clear Lake.
 fuscum. Fr.
 On oak. San Raf.
 malleolus. B. & Rav.
 On oak. Sausalito.
 rubiginosum. Fr.
 On laurel logs. San Raf.
 rufa. Fr.
 On wild cherry. Yosemite.
 stigmatum. n. sp. Cke.
 On dead bark. Tehachapi.
 serpens. Fr.
 On dead laurel. San Raf.

NUMMULARIA.

- Bulliardi. Tul.
On live oak. Yosemite.

DIATRYPELLA.

- tocciana. De Not.
On bark dead alder. Streetens.

DOTHIDEA.

- graminis. Fr.
On grass. Sac.
junci. Fr.
On juncus. Sac.
potentillæ. Fr.
On potentilla glandulosa. San Raf.
ribesia. Pers.
On currant twigs. Sac.
Rosæ. Fr.
On rose. Sac.

DIATRYPE.

- aspera. Fr.
On oak. Streetens.
bullata. Fr.
On willow. Streetens.
disciformis. Fr.
On buckeye. San Raf.
hystrix. Fr.
On sycamore. Sac.
incaerata. B. & Br.
On rose. Sac.
strumella. Fr.
On currant. Sac.

MELANCONIS.

- longipes. Tul.
On oak branches. San Fran.

VALSA.

- coronata. Fr.
On oak. Sausalito.
salicina. Fr.
On willow. Santa Cruz.
sordida. Fr.
On poplar bark. Sac.
vitis. Schw.
On grape cuttings. San Raf.

DIAPORTHE.

arctii. Nke.

On heracleum. Streetens.

CUCURBITARIA.

confluens. n. sp. Plow.

On oak bark. Colfax.

macilenta. Cke.

On cedar. Blue Cañon.

SORDARIA.

bombardiodes. Awd.

On colt's dung. Mt. Shasta.

Californica. n. sp. Plow.

On cow dung. San Raf.

coprophila. De Not.

On cow dung. Sac.

fimicola. Rab.

On horse dung. Sac.

BYSSOPHÆRIA.

aquila. Fr.

On rotten wood. San Francisco.

PSILOSPHÆRIA.

moriformis. Tode.

On hazel. Howards.

myriocarpa. F.

On elder. Tehachapi.

pulveracea. Ehr.

On oak. Yosemite.

LASIOSPHÆRIA.

canescens. Pers.

On redwood. Streetens.

SPHÆRIA.

Arctostaphylos, n. s. Pl.

On dead manzanita. Yosemite.

bombardioides. Awd.

On cow dung. Colfax.

commanipula. B. & Br.

On Aralia Cala. Colfax.

confertissima. n. sp. Plow.

On redwood cones. Streetens.

conficta. n. sp. Cke.

On oak leaves. Streetens.

Herbarum. Pers.

On soap plant. St. Helena.

- megalocarpa*. n. sp. Plow.
 On maple. Colfax.
morbosa. Schw.
 On living wild cherry. Yosemite.
palmacea. n. sp. Cke.
 On palms. Los Angeles.
permunda. n. sp. Cke.
 On Soap plant. Blue Cañon.
picea. Pers.
 On *Chenopodium alba*. Sacramento.
pulvis-pyrius. pers.
 On *pinus contorta*. Blue Cañon.
rostellata. Fr.
 On *Ribes ursinus*.
rubella. Pers.
 On *heracleum*. San Fran.
salicella. Fr.
 On willows. Summit.
sambuci. n. sp. Pl.
 On elder. Streetens.
sequoiæ. n. sp. Pl.
 On *sequoia*. Big Trees.
spiculosa. Pers.
 On ash bark. Streetens.
sub-moriformis. n. sp. Pl.
 On bark. San Raf.
sustenta. n. sp. Pl.
 On manzanita. Blue Cañon.
tumulata. n. sp. Cke.
 On *pinus contorta*. Summit.
Vizeana. Cke.
 On *lathyrus venosa*. Sac.
Yucca. Sch.
 On *yucca communis*. Sac.
Yuccaegena. n. sp. Cke.
 On *yucca angustifolia*. Mohave.

GNOMONIA.

- alni*. n. sp.
 On alder. San Raf.

SPHÆRELLA.

- brassicola*. De Not.
 On cabbage leaves. Sac.
conglomerata. Wall.
 On alder leaves. Howards.

- erysipheana. B.
 On hop leaves. Sac.
 latebrosa. Cke.
 On fallen sycamore leaves. Sac.
 salicicola. W.
 On willow leaves. Summit.
 simulans. Cke.
 On oak leaves. San Raf.
 sparsa. Awd.
 On chestnut leaves. Sac.

VENTURIA.

- sequoiæ. n. sp. Pl.
 On decaying leaves, sequoia. Big Trees.

STIGMATEA.

- chætonium. Fr.
 On raspberry leaves.
 Geranii. Fr.
 On leaves of geranium.
 polygonorum. Fr.
 On living leaves of Polygonum. San Raf.
 potentillæ. Fr.
 On living leaves of Potentilla. Sac.
 ranunculi. Fr.
 On Rannuculus Cala. San Raf.

OZHOMA. (Sp?)

- On pine wood. Blue Cañon.

ADDENDA.

STEPTOTHRIX.

atra. B. & C.

On grape cuttings. San Raf.

AGARICUS. (Sub. Gen?)

tridens. n. sp. Moore.

CAVE FUNGUS.

Pileus fleshy, conchiform; skin very thick, coriaceous; color light buff. Edge entire, involute. The stem is *three feet and four inches long*, and is attached to the timber on which it grows, and from which it hangs suspended by a disk several inches in diameter. This disk is clothed with a thick, soft brown felt, which extends down and covers the stem for several inches.

At nine inches from the point of attachment the stem divides into three branches, which again unite, singularly enough, at just nine inches from the point of division, and immediately send off two lateral branches.

The main stem continues downward, enlarging to the diameter of nearly three inches. It is no longer covered with the felty mass, but is smooth and hard to the point at which the shell-shaped pileus is thrown off. Here a most singular growth has taken place. From all sides short branches, resembling the young antlers of a stag, are pushed out. Two of these becoming more vigorous than the rest have extended downward, the larger of the two dividing into three branches, thus *terminating the whole in a perfect trident*.

The gills, which are distinct, notched, sinuate, and of a pale straw color, run for some distance up the stem. The spores are ovate, exceedingly minute and borne on true basidia.

The plant was found attached to a timber in one of the abandoned drifts of the Yellow Jacket Mine, Virginia City, Nevada, at the depth of 400 feet below the surface. J. P. M.

Order XXI. BOLETUS.**B.**luteus. Lin. *E.* rare. San Raf.

flavidus. Bull. San Raf.

piperatus. Bull. San Raf.

variegatus. Swartz. Howards.

subtomentosus. Linn. *E.* San Raf.edulis. Bull. *E.* San Raf.

impolitus. Fr. San Raf.

aestivalis. Fr. San Raf.

Satanas. Lenz. San Raf.

luridus. Schæff. San Raf.

erythropus. Pers. San Raf.

versipelles. Fr. *E.* San Raf.scaber Fr. *E.* San Raf.granulatus. L. *E.* San Raf.bovinus. L. *E.* San Raf.

castaneus. Bull. San Raf.

Order XXII. FISTULINA.

*Compliments of A. B. Seymour,
Cambridge, Mass.*

Issued November 8, 1889.

No. 115

QK

2.5

.03

v. 100

[From the PROCEEDINGS OF THE BOSTON SOCIETY OF NATURAL HISTORY,
VOL. XXIV, 1889.]

LIST OF FUNGI, COLLECTED IN 1884 ALONG THE
NORTHERN PACIFIC RAILROAD.

BY A. B. SEYMOUR.

THE fungi enumerated in this list were collected by the author
in August and September, 1884, along the line of the Northern

Pacific Railroad. Beginning at St. Paul, Aug. 21, short stops and hurried collections were made at various points until Sept. 23rd. The dates and localities of collecting are as follows :

- August 21, St. Paul, Minnesota,
- “ 22, Brainerd and Detroit, Minnesota,
- “ 23, Fargo, Dakota,
- “ 25, Valley City, Dakota,
- “ 26, Jamestown, “
- “ 28, 29 and 30, Bismarck, Dakota.

- September 2, Medora, Dakota,
- “ 3, Miles City, Montana,
- “ 4, Billings, “
- “ 6, Livingston, “
- “ 8, Bozeman, “
- “ 11 and 12, Helena, Montana,
- “ 15, Sand Point, Idaho,
- “ 17, Wallula, Washington,
- “ 19, Thompson Falls, Montana,
- “ 23, Minneapolis and Lake Minnetonka, Minnesota.

A few other localities are mentioned in the list, represented by specimens picked up at points where the train stopped. Species distributed in *exsiccati* from this collection are so noted.

For aid in the determinations, I am indebted to Drs. W. G. Farlow and P. Magnus and especially to the lamented Dr. Winter.

CHYTRIDIACEÆ.

Synchytrium decipiens, Farl., on *Amphicarpæa monoica*, plentiful ; Bismarck, Dakota.

PERONOSPOREÆ.

Peronospora Halstedii, Farl., on *Rudbeckia laciniata* ; Fargo and Valley City, Dak.

Peronospora ribicola, Schrt., Jahresbericht d. Schles. Gesellsch. 1883, p. 179 ; *Plasmopara ribicola*, Schrt. Krypt. Fl. Schles. Vol. III, p. 238, where it is described as follows : “Rasen fleckenweise, sehr locker, weiss. Conidienträger straff aufrecht, 2-4 mm. hoch, mit 3-5 aufrecht abstehenden graden Aesten, meist in eine Endspitze auslaufend ; Aeste grade, die untern mit 3-5 graden Seitenästen ; Endästchen grade, verschmälert, abgestutzt. Conidien kurz elliptisch, 15-20 μ lang, 11-13 μ breit, mit flacher Papille. Keimung

nicht beobachtet. Oosporen unbekannt. Auf Blätten von *Ribes rubrum*."

The Minnesota specimens show conidia $11.7-17\mu$ by $13-21\mu$; oospores smooth or obscurely roughened, nearly filling the oogonium and closely invested by its thin, fuliginous wall, 25.5 to 35.7μ . The papilla of the conidium is seen with difficulty and appears more like an inward thickening of the wall than as an external prominence. On leaves of *Ribes hirtellum*. Detroit, Minn.

Peronospora parasitica, (P.) Tul., on *Lepidium*; Jamestown, Dak.

Peronospora Arthuri, Farl., on *Oenothera biennis*; Sand Point, Idaho. Oospores 15.6 to 39μ .

Since this species was first published as occurring in Iowa and Minnesota, it has been reported from Illinois, Wisconsin, Missouri, Kansas and New York. To these localities may be added Nantasket Beach, Mass., where it was found by the author, June 23, 1885 and Grand Manan, Aug. 13, 1888, K. Miyabe.

Peronospora Trifoliorum, D By. On *Astragalus*; Bismarck, Dak., Livingston, Mont.

Exs.: Fungi Europæi 3576.

Cystopus Bliti, (Biv.) Lev. On *Amarantus retroflexus*; St. Paul and Brainerd, Minn., Jamestown, Dak.

ERYSIPHEÆ.

DETERMINED BY F. S. EARLE.

Sphærotheca Humili, (DC.) Burrill., on *Viola*; Valley City, Dak.

Sphærotheca pannosa, (Wallr.) Lev., on *Rosa*; Valley City, Dak.

Sphærotheca Mors-uvæ, (S.) B. & C., on *Ribes floridum*; Fargo, Dak.

Sphærotheca Castagnei, Lev., on *Erechtites hieracifolia* and *Nabalus* sp. Lake Minnetonka, Minn. *Hydrophyllum Virginicum*; Valley City, Dak.

Erysiphe communis, (Wallr.) Schl., on *Ranunculus Cymbalaria*; Livingston, Mont. *Lathyrus*?; Lake Minnetonka, Minn. *Oenothera biennis*; Kootenai, Idaho. *Oenothera* sp.; Bismarck, Dak.

Erysiphe Cichoracearum, DC., on *Aster*, Valley City, Dak. *Solidago* and *Cnicus*; Bozeman, Mont. *Helianthus*; Fargo, Dak.

Erysiphe Linkii, Lev., on *Artemisia dracunculoides*; Bozeman, Mont. *Artemisia* sp.; Valley City, Dak. Also on *A. Ludoviciana*; Helena, Mont., Sept., Rev. F. D. Kelsey.

Erysiphe Galeopsidis, DC., on *Stachys palustris*; Detroit, Minn.
Uncinula parvula, B. & C., on *Celtis*; Wallula, Washington.
Uncinula Salicis, (DC.) Wint., on *Salix*; Lake Minnetonka, Minn., Jamestown and Bismarck, Dak., Bozeman, Mont.

Phyllactinia suffulta, (Reh.) Sacc., on *Betula*?; Lake Minnetonka, Minn. Also on *Cornus stolonifera*; Helena, Mont., Sept. 1888, Rev. F. D. Kelsey. Remarkable for occurring on both sides of the leaf.

Podosphaera Oxyacanthæ, (DC.) D By., on *Prunus* (cult.?) ; Livingston, Mont. *Spiræa* sp., Sand Point, Idaho. Also on *Prunus Virginiana*; Helena, Mont., Sept., 1888, Rev. F. D. Kelsey.

Microsphaera Alni, (DC.) Wint., on *Ceanothus Americanus*; Brainerd, Minn. *Syringa vulgaris* and *Betula*; Lake Minnetonka, Minn.

Microsphaera diffusa, C. & P., on *Lespedeza capitata*; Brainerd, Minn.

Microsphaera Ravenelii, B. on *Lathyrus*?; Detroit, Minn.

Microsphaera Symphoricarpi, Howe, on *Symphoricarpus*; Valley City, Dak., Bozeman, Mont., Sand Point, Idaho, Cheney, Washington.

UREDINEÆ.

Uromyces Terebinthi, (DC.) Wint., on *Rhus Toxicodendron*; Valley City, Dak.

Uromyces Lupini, B. & C., on *Lupinus argenteus*; Bozeman, Mont.

Uromyces Genistæ-tinctoriæ, (P.), on *Trifolium*; Thompson Falls, Mont. *Lupinus argenteus*; Livingston, Mont.

This is the same as *U. Lupini*, Sacc., and is characterized by warty apiculate spores with fragile pedicels. *U. Lupini*, B. & C. has smooth spores and is very different. It is erroneously referred here by Dr. Winter in "Die Pilze," 1, 146, on the authority of Thuemen, Myc. Univ. 842. The form on *Trifolium* has somewhat smaller spores but is not specifically distinct.

Uromyces Psoraleæ, Pk., on *Psoralea lanceolata*; Medora, Dak.

Uromyces argophyllæ, Seymour, in Ell. and Ev. N. Am. Fung. 1862.

Spots none; sori amphigenous, small, scattered, inconspicuous, obscured by the hairs of the leaf; teleutospores subglobose to oblong, broadest at, above or below the middle, $17-25 \times 22-45 \mu$; epispore thin, of uniform thickness or slightly thickened and papillate at the apex; pedicel fragile. Differs from *U. Psoraleæ*, Pk. chiefly in longer and narrower spores and more deciduous pedicels.

On *Psoralea argophylla*; Valley City and Bismarck, Dak.

Uromyces sp. on *Glycyrrhiza lepidota*; Jamestown and Bismarck, Dak., Miles City and Livingston, Mont. This was called *U. Trifolii* by Dr. Winter (in litt.) and distributed under that name in Ell. & Ev. N. Am. Fung. 1876, but I have been unable to satisfy myself with the determination.

Uromyces Viciæ, (P.) D By., on *Lathyrus*; Brainerd, Minn.; Valley City, Jamestown and Bismarck, Dak.

Uromyces Rudbeckiæ, Arth. and Holw., on *Rudbeckia laciniata*; Fargo and Valley City, Dak., Livingston, Mont.

Exs., Fungi Europæi 3412.

Uromyces Euphorbiæ, C. & P., on *Euphorbia marginata*; Miles City, Mont. *Euphorbia* sp.; Medora, Dak.

Uromyces Polygoni, (P.) Fckl., on *Polygonum aviculare*; Medora, Dak.

Uromyces Eriogoni, Ell. Hark., on *Eriogonum* sp.: II, Helena, Mont.; III, Medora, Dak.; II, III, Livingston, Mont.

Exs.: Ell. Ev. N. Am. Fung. 1871.

Uromyces Junci, (S.) Tul., on *Juncus*; Bozeman, Mont.

Uromyces Scirpi, Burrill, on *Eleocharis*; Jamestown, Dak.

Uromyces acuminatus, Arth., on *Spartina cynosuroides*; Valley City and Bismarck, Dak. *Spartina gracilis*; Medora, Dak.

Uromyces Alopecuri, n. sp.

II. Sori epiphyllous, small, scattered, inconspicuous, mostly covered by the epidermis; spores subglobose to elliptical; epispore rather thick, warty; size, $12-20 \times 15-24\mu$.

III. Sori epiphyllous, small, blackish, covered by the epidermis; spores obovate to elliptical, broadly rounded or truncate at the apex, $17-23 \times 23-32\mu$; epispore of nearly equal thickness throughout; pedicel persistent, as long as the spore or shorter.

On *Alopecurus geniculatus*, var. *aristatus*; Brainerd, Minn.

Puccinia Anemones-Virginiæ, S., on *Anemone patens*, var. *Nuttalliana*; Livingston, Mont. *Anemone cylindrica*; Valley City, Dak.

Puccinia mirabilissima, Pk., on *Berberis repens*: II, Helena, Mont.; II, III, Thompson Falls, Mont.

Puccinia Violæ, DC., on *Viola*; Valley City and Jamestown, Dak.

Puccinia Arenariæ, (Schum.) Wint., on *Cerastium arvense*; Livingston, Mont.

Puccinia Malvastri, Pk., on *Malvastrum coccineum*: II, III, Al-

mont, Dak.; III, Jamestown and Bismarck, Dak.; Billings and Helena, Mont.

Exs.: Ell. and Ev. N. Am. Fung. 1850; Fungi Europæi 3515.

Puccinia Petalostemonis, Farl., on *Petalostemon violaceus*: II, III, Jamestown, Dak.; III, Valley City and Bismarck, Dak.

Exs.: Ell. and Ev. N. Am. Fung. 1844.

Puccinia Amorphae, M. A. Curtis, on *Amorpha canescens*; Brainerd and Lake Minnetonka, Minn., Bismarck, Dak. *Amorpha fruticosa*; Valley City, Dak.

Puccinia Pruni-spinosæ, P., on *Prunus (Americana?)*; Valley City, Dak.

Puccinia Epilobii, DC., on *Epilobium paniculatum*; Bozeman, Helena and Thompson Falls, Mont.

Exs.: Ell. and N. Am. Fung. 1846.

Puccinia Pimpinellæ, Lk., on *Osmorrhiza brevistylis*; Bozeman, Mont.

Puccinia Jonesii, Pk., on *Cymopterus bipinnatus*; Livingston, Mont.

Puccinia Galiorum, Lk., on *Galium boreale*; Valley City and Jamestown, Dak.; Helena, Mont.

This is apparently the form called *P. rubefaciens*, Johanson; see Lagerheim in Hedw. 1889, p. 106, fig. 5.

The sori are elliptical or oblong, large, dark, convex. Those commonly seen on *Galium concinnum* are punctiform; on *G. Aparine*, squarish and usually remaining covered by the epidermis. The uredospores have been rarely if ever found in the United States and the æcidia are rarely if ever seen except on *G. Aparine*. The host name *G. triflorum* in ILLINOIS UREDINEÆ, p. 181, is an error for *G. Aparine*.

Puccinia Kuhniae, S., on *Kuhnia eupatorioides*; Valley City and Bismarck, Dak.

Puccinia Asteris, Duby, on *Aster conspicuus*; Helena, Mont. *Aster* sp.; Fargo, Dak.

Puccinia Grindeliæ, Pk., on *Grindelia squarrosa*; Livingston and Helena, Mont. Exs.: Fungi Europæi 3513.

Puccinia Helianthi, S., on *Helianthus annuus*; Miles City, Mont. *Helianthus*, sp.; Lake Minnetonka, Minn. Fargo, Valley City, Jamestown, Bismarck, and Medora, Dak., Billings and Miles City, Mont.

Puccinia Tanacetii, DC., *Artemisia dracunculoides*: II, III, Me-

dora, Dak. ; III, Bismarek, Dak., Livingston and Bozeman, Mont. *Artemisia Ludoviciana*, Jamestown, Dak.

Exs. : Ell. and Ev. N. Am. Fung. 1848.

Puccinia Xanthii, S., on *Xanthium Canadense*; St. Paul, Minn., Bismarek and Medora, Dak., Miles City, Mont. *Ambrosia psilostachya*; Valley City and Medora, Dak., Billings, Mont.

Exs. : Ell. and Ev. N. Am. Fung. 1853.

Puccinia variolans, Hark., on *Actinella acaulis*; Livingston, Mont.

Puccinia Balsamorhizæ, Pk., on *Balsamorhiza sagittata*; Thompson Falls, Mont.

Puccinia cladophila, Pk., on *Stephanomeria minor*; Livingston, Mont. Differs slightly from the original specimen.

Also collected in Washington Terr., Sept., 1883, by Suksdorf; Spokane Falls, Sept. 24, 1880, Watson; on *Stephanomeria runcinata*; foot hills of Santa Rita Mountains, Ariz., June 1, 1884, Pringle. Exs. : Ell. and Ev. N. Am. Fung. 1839.

Puccinia flosculosorum, (A. & S.) Roehl., on *Oniscus*; Fargo, Dak., Billings, Mont. *Hieracium*; Fargo and Bismarek, Dak., Billings and Thompson Falls, Mont. *Taraxacum*; II, Fargo, Dak.

Puccinia Menthæ, P., on *Mentha* sp.; II, III, Brainerd and Lake Minnetonka, Minn. *Monarda fistulosa*, Jamestown, Dak., Livingston, Bozeman and Helena, Mont.

Puccinia plumbaria, Pk., on *Gilia* sp.; Bozeman, Mont.

Puccinia Convolvuli, Cast., on *Calystegia sepium*; II, III, Fargo; III, Valley City, Dak.

Puccinia Gentianæ, (Strauss) Lk., on *Gentiana Oregana*; Helena, Mont. *Gentiana* sp.; Jamestown, Dak.

Puccinia Polygoni-amphibii, P., on *Polygonum amphibium* or allied form; II, III, Brainerd and Lake Minnetonka, Minn., Bozeman, Mont.

Puccinia angustata, Pk., on *Scirpus Eriophorum*; Brainerd and Lake Minnetonka, Minn.

Puccinia Caricis, (Schum.) Reb., on *Carex*; Bismarek, Dak., Helena, Mont.

Puccinia Phragmitis, (Schum.) Körn., on *Spartina cynosuroides*; Valley City and Bismarek, Dak. In the specimens gathered at Valley City the spores are often acuminate.

Puccinia Stipæ, Arth., on *Stipa viridis*; Southeart, Dak. *Stipa viridis* (?); Jamestown, Dak., Helena, Mont.

Puccinia vexans, Farl., on *Bouteloua racemosa*; Valley City and Jamestown, Dak.

Puccinia graminis, P., on *Bouteloua oligostachya* (?); Jamestown, Dak. *Ammophila longifolia*; Jamestown, Bismarck and Sweet Briar, Dak. Wheat; Fargo, Dak., scarce. Oats: II, Billings, Mont.; III, Jamestown, Dak. *Hordeum jubatum*; Lake Minnetonka, Minn.

Puccinia Rubigo-vera, (DC.), on Wheat; Fargo, Valley City and Jamestown, Dak. *Agropyrum*; Bismarck, Dak., Thompson Falls, Mont. *Hordeum jubatum*; Fargo and Valley City, Dak. *Elymus Canadensis*; Bismarck, Dak. In some cases these specimens have the ordinary paraphyses with the teleutospores and different capitate paraphyses with the uredospores. *Elymus*; Fargo and Jamestown, Dak., Livingston, Mont. In the Livingston specimens, the teleutospores have apical points, suggesting *P. coronata*, Cda.; Specimens on an unknown grass from Thompson Falls, Mont., resemble *P. coronata*, but have paraphyses like *P. Rubigo-vera*.

P. coronata, Cda., on Oats; Fargo, Jamestown, Bismarck, Dak.

Puccinia tomipara, Trelease, on *Bromus ciliatus*; Detroit, Minn. Exs.: Ell. and Ev. N. Am. Fung. 1842.

Puccinia emaculata, S., on *Panicum capillare*; Lake Minnetonka, Minn.

Puccinia Andropogi, S., on *Andropogon furcatus*; Brainerd, Minn., Jamestown and Bismarck, Dak. *Andropogon scoparius*; Jamestown and Bismarck, Dak.

Phragmidium Fragariae, (DC.), on *Potentilla Norvegica*; III, Helena, Mont. *Potentilla gracilis*, var. *rigida*; III, Bozeman, Mont. *Potentilla* (near *P. gracilis*); III, Helena, Mont.

Phragmidium Potentillae, (P.), on *Potentilla Pennsylvanica*; III, Helena, Mont. This species is characterized by dark colored firm spores and firm pedicels.

Phragmidium mucronatum, (P.) Lk., on *Rosa* sp.: II, III, Brainerd, Minn., Fargo, Dak., Miles City, Livingston and Bozeman, Mont.; III, Detroit, Minn., Jamestown, Dak., Helena and Thompson Falls, Mont.

Phragmidium Rubi-idaei, (P.) Wint., on *Rubus strigosus*; Helena, Mont. *Rubus Nutkanus*; Helena and Thompson Falls, Mont.

Coleosporium Sonchi-arvensis, (P.) Lev., on *Aster macrophyllus*; II, III, Brainerd, Minn. *Aster conspicuus*; III, Helena, Mont. *Aster* sps.; II, Brainerd, Minn., Fargo and Valley City, Dak., Bozeman

Mont. ; II, III, Thompson Falls, Mont. *Solidago* : II, Medora, Dak., Miles City and Helena, Mont. ; II, III, Detroit, Minn.

Melampsora Lini, Tul., on *Linum* sp. ; II, III, Livingston, Mont.

Melampsora Epilobii, (P.) Fekl., on *Epilobium angustifolium* ; II, III, Brainerd, Minn., Helena and Thompson Falls, Mont., Kootenai and Sand Point, Idaho. *Epilobium coloratum* : II, Bozeman, Mont., II, III ; Livingston and Helena, Mont.

Melampsora populina, Lev., on *Populus monilifera* ; II, III, Jamestown and Bismarck, Dak. *Populus balsamifera* ; II, III, Helena, Mont. *Populus tremuloides* ; II, III, Livingston, Mont. Very injurious to young cottonwood trees planted on tree claims.

Melampsora salicina, Lev., on *Salix* sps. : II, Detroit, Minn., Fargo and Bismarck, Dak., Billings, Livingston and Helena, Mont. ; II, III, Bozeman, Mont.

Cronartium asclepiadeum, (Willd.) Fr., var. *Thesii*, B., on *Comandra umbellata* ; II, III, Livingston, Mont.

Cronartium asclepiadeum, (Willd.) Fr., var. *Quercuum*, B & C., on *Quercus coccinea* ; Brainerd, Minn. Exs. : Ell. & Ev. N. Am. Fung. 1881 ; Fungi Europæi 3418.

Æcidium Euphorbiæ, P., on *Euphorbia* sp. ; Miles City, Mont.

Roestelia lacerata, (Sow.) Fr., on *Cratægus* sp. ; Bozeman, Mont. *Amelanchier alnifolia* ; Livingston and Bozeman, Mont.

USTILAGINEÆ.

Ustilago subinclusa, Körn., on *Carex trichocarpa*, var. *Deweyi* ; Bismarck, Dak. Exs. : Ell. & Ev. N. Am. Fung. 1889.

Ustilago segetum, (Bull.) Ditm., on Oats ; Sand Point, Idaho ; Billings, Mont.

Ustilago Rabenhorstiana, Kühn, on *Panicum glabrum* ; Minneapolis.

Ustilago Syntherismae, S., on *Cenchrus tribuloides* ; St. Paul.

Sorosporium Ellisia, Wint., var. *occidentalis*, Seymour, on *Andropogon furcatus* ; Bismarck, Dak.

In the original specimens described by Dr. Winter the undeveloped flowers are infested before the expansion of the inflorescence. In these specimens, individual flowers of the fully expanded inflorescence are affected. The spores measure 9–10 μ . This has been referred to *Ustilago Andropogonis*, Kell. and Swingle (Journ. Myc. Vol. v, p. 12, pl. 1, figs. 12–26) but the measurements of the spores are less than those of the Kansas specimens as there stated, also

somewhat less than those given by Dr. Winter (Torr. Bull. x, 7). The spores certainly fall apart very easily, but the glomerules may be seen by careful examination. This was positively stated by Dr. Winter (in litt. July 9, 1885) to be his *U. Ellisii*. I alone am responsible for the varietal name.

Entyloma Menispermii, Farl. and Trelease, on *Menispermum Canadense*; Fargo and Valley City, Dak.

Entyloma Compositarum, Farl., *Erigeron Philadelphicum*; Valley City, Dak.

Entyloma Physalidis, (Kalch. & Cke.) Wint., on *Solanum triflorum*; Bismarck, Dak.

Exs.: Ell. & Ev. N. Am. Fung. 1491*b* after 1900.

The Secretary called attention to peculiar calcareous formations from the Bermudas known as "Fossil Palms."

SALEM PRESS, July, 1889.

EVENING SESSION.

WEDNESDAY, JANUARY 20, 1886.

The meeting was called to order at 7 o'clock by President Smith.

The following lecture was then delivered by Prof. A. B. Seymour, of the State University at Madison, Wisconsin:

SOME FUNGUS DISEASES OF SMALL FRUITS.

Mr. President, Ladies and Gentlemen:

Before speaking of any particular disease, I wish to show how and why fungi produce disease in plants. It is well known that the green coloring matter occurring in all our common plants has the power of converting inorganic matter taken from the soil and air into food materials for the plant. All plant food and indirectly animal food is produced in this way. The plants known as fungi, however, have no leaf-green and therefore are not able to provide themselves with food from soil and air. They must derive their nourishment from either living or dead organic matter.

Different kinds of plants grow on different soils. Fungi choose places of growth in much the same way that other plants do, but much more closely. Some grow on almost any decaying vegetable matter; others on the wood of some particular tree. A very large number grow on living plants and commonly each species only on plants of one particular family or species. Wheat rust grows on various cereals, the orange rust of berry plants, only on raspberries and blueberries, while corn rust is not known to occur on any plant but Indian corn.

Where do fungi come from? How do they get there? They never appear spontaneously but always from reproductive bodies called spores, serving the purpose of seeds. These are very minute and so light that they are borne in the air like dust, and some kinds are carried by wind. Under favorable conditions if they are brought to the right kind of plant, as corn smut spores to corn, they germinate and send out a tender tube which answers the purpose of a root, and makes its way into the plant, often through breathing pores. Once inside, it grows and branches, sending its threads in various directions among the plant cells. Frequently special branches are produced, which enter the cell cavities and act as suckers to take up nourishment. Fungi have different stages of development, reminding one

rather of insects in their transformations than of plants. Each stage has its own kind of spores. Frequently different stages occur on different host plants, as in wheat rust. This produces on the wheat itself yellow summer spores and blackish spores which live over winter. The summer spores retain their vitality only a short time; only the blackish spores can carry the fungus through the winter. In spring they may germinate on a different plant, the barberry, and produce a third stage there; the spores from this stage, in their turn, produce yellow rust on the wheat.

It has been claimed that where fungi grow on plants, the plants first become diseased and weakened and the fungus comes afterwards, following and not causing the disease. This may be true in some cases, but in all ordinary cases of this kind the fungus is the true cause of the disease. Certain conditions of weather are favorable to its development, just as certain conditions are favorable to the development of any other plant; but the conditions which favor the fungus may be unfavorable to the plant it grows upon and so give the former a double advantage. Moreover plants have a greater power and resist disease when abundantly supplied with food materials, when they are not exhausted by fruiting, when every condition is favorable to robust growth; but the fungus may grow and produce the disease in spite of all this. A man in robust health is less likely to succumb to a contagious disease, as small pox, but still he is likely to take it.

There are several ways in which a fungus may produce injury to the plant. One way, common to all, is by taking away the food of the plant for its own growth. On green parts, they destroy part of the leaf-green and so reduce the power of the plant to supply itself with food; in many cases the leaves fall, as from premature ripeness. Frequently the fungus causes an abnormal development of plant tissues as in the black knot of the plum tree and the curl of the peach leaves; or they arrest development, as in the orange rust of blackberry, so that rusted leaves are smaller than healthy ones. In many cases the flower or fruit alone is destroyed, as in the smut of wheat and oats, "double blossom" of blackberries and the swelling of young plums.

ORANGE RUST.

The most striking and most destructive fungus disease of ^{Small}swell fruits is the orange rust, (*Cæoma nitens*, Schw.) which occurs on raspberry and blackberry leaves, and is especially destructive on the latter. This has been most thoroughly studied by Professor T. J. Burrill, of the University of Illinois, and many of the following facts are from his

investigations. This rust appears as a thick orange coating on the under surface of the leaves and attains its greatest development in June.

Figure 1—Orange Rust Fungus, *Cecoma nitens*, Schw. Cross-section of a diseased leaf; *a* spores; *b*, immature spores or sterile bodies; *c* bed of fungus issue; *d* mycelium threads; *e* suckers; *g* spermatogones; *h* cells of leaf; *m* epidermis of upper leaf surface. (After Burrill.)

The leaves do not reach their full size, are unusually rigid, and have a sickly appearance; something of this may be seen even before the yellow coating appears, and the latter at first appears in definite and paler patches, somewhat like scales, standing out slightly on the under leaf surface, covered by the epidermis, through which they afterwards burst. In this early stage, on the upper surface may be seen also yellowish specks, which on closer examination prove to be protuberances. There seems to be an exudation from these, which attracts insects.

Microscopic examination shows that the orange portion consists of a great number of roundish spores. Their surface is roughened with sharp points, by means of which it is possible that they adhere to insects crawling over the leaf and are carried by them to other plants.

It is certain that they may be carried by the wind from place to place. The chains are produced in chains arising perpendicular to the leaf surface. The spores toward the inner end of the chain are either immature spores or sterile cells. The spore chains arise from a bed or cushion of fungus tissue, from which may be traced the threads of mycelium running through the leaf tissues.

Some have special branches entering into the interior of cells and there forming coils, acting as suckers to take up nourishment. Some also extend to the upper surface, where they are connected with the yellowish bodies, *spermagones*, previously mentioned as occurring there. One of the epidermal cells is greatly enlarged and protrudes. In its cavity a thread produces a number of branches which extend nearly parallel toward the top and bear at their ends minute spore-like bodies, *spermatia*. Their office is not well known.

It has been believed by cultivators that this fungus lives over winter in the roots, and spreads to the stem and leaves in the spring; but this is not the case. It extends scarcely at all beyond the areas covered by the orange spores. More than this, it is found that the spores will lose their power of germination if deprived of moisture for a few days and will not live over winter. Hence it is believed that some other kind of spores is produced corresponding to the black spores of wheat rust, which carry the fungus through the winter. These have been carefully sought for, but heretofore without success.

Prof. Burrill has however at last found some evidence as to what they are, and it is hoped that his culture experiments, now in progress, will settle the matter definitely and lead to important practical results.

Some varieties of blackberries, especially the Snyder, are seldom, if ever attacked by this rust.

Remedies: Cut out and burn all diseased parts as soon as the disease appears, and as a preventive measure destroy any useless vines, as wild ones in fence corners or elsewhere, which might nourish the disease.

OTHER RUSTS OF BLACKBERRIES AND RASPBERRIES.

There are two other forms of yellow rust, one on blackberries only, the other on raspberries only, both inconspicuous compared with the preceding and never likely to be mistaken for it. They are quite similar to each other and bear yellow spores in minute pustules scattered over the inferior leaf surface. They sometimes do considerable damage but never approach the *Cæomas* in that respect. The general mode of growth is much the same as in the *Cæomas* (Orange Rust) but the sperma-

gonies are absent and the spores are borne on stalks, from which they easily fall, instead of being in chains.

FIGURE 2.—Summer or Uredo spores of the raspberry rust, *Phragmidium rubi-idaei*. Spore cluster surrounded by club-shaped sterile bodies, paraphyses. (After Winter.)

winter spores. The yellow or summer spores of the two kinds differ but little from each other, but the form on blackberries is a little the more conspicuous. The winter spores are long, cylindrical, nearly black and tipped with a sharp point. The kind on blackberries is finely warty and divided by cross partitions into five or six cells; that on raspberries is coarsely warty and divided into seven to nine cells.

In this region the black raspberries are seldom attacked, but the red ones, especially in a wild state, quite commonly. The cultivated varieties of the red, Turner, Brandywine and Cuthbert, are reported to have suffered seriously from this disease at Jacksonville, Illinois.

The yellow spores come in August or September and the black ones soon follow. On the blackberry, the yellow spores come somewhat later, and the black ones have seldom been seen in this country.

FIGURE 3.—Winter or telento-spores of raspberry rust, *Phragmidium rubi-idaei*.

Figure 4—Leaf of currant showing diseased spots.

A CURRANT DISEASE.

In Western Massachusetts last summer (1885) the currant bushes were badly injured by a fungus (*Septoria ribis*, Desm.) growing on the leaves. The same occurs in this region on wild currants and gooseberries, on the cultivated currant in Ohio, and on gooseberries in Kentucky. The disease appears as brown spots on the leaves, angular spots limited by the veins and soon becoming dead and dry. The spots are good sized and when large or numerous, destroy nearly the whole leaf. It certainly has an exhausting effect on the leaf and many leaves become entirely exhausted and fall off.

The microscope shows the presence of a fungus growing among the leaf tissues in the usual way, but its mode of fruiting is quite different from those described before. Within the tissues and somewhat protruding on the surface are minute blackish postules or spore-cases (perithecia). At the base of the perithecia, within, arise threads which bear the spores. The spores are colorless, very slender and thread-like, curved, pointed at the free end.

Figure 5—Spore of currant disease, highly magnified.

The life history of this fungus is unknown, but it is known that many of this group are only the summer stages of species which live through the winter and mature in late winter or spring, either on the same spot with the summer form, as in the black knot of the plum trees, or on different parts where the summer spores have

been carried and have germinated. Many fungi of this class produce their winter spores on fallen leaves or twigs, for instance the black blotch fungus of maple leaves.

It is probable that this currant fungus has some winter form on the dead vegetable matter about the bushes. Hence the remedy should be sought in burning the leaves that fall and any other rubbish that might harbor the fungus.

A similar species *Septoria rubi*, West. is very common on blackberries all over the country. It produces roundish spots with a red or purple margin and a brown center.

DOUBLE BLOSSOM.

“Double blossom” is the name given to a blackberry disease that is quite prevalent and destructive in some regions. It is a disease of the blossoms, as the name implies, but they are not actually double. The several parts are abnormally swollen through the effect of the fungus, so that at a casual glance the flower appears double. It is killed of course and no fruit is formed. The mycelium runs through the tissue of the floral organs, swelling and distorting them, and at length sends out on the surface clusters of short, colorless threads, which bear the spores on their ends and produce a whitish, mealy appearance over the flower. The spores are slender and pointed, frequently curved, and divided by cross-partitions into two to four cells.

This disease was first observed and studied by Mr. F. S. Earle. He sent specimens to Dr. Winter of Germany and the latter has recently described and published it as a new species, which he refers with doubt to the genus *Fusisporium* and calls *Fusisporium rubi*.

Figure 6—Spores of double blossom fungus, *Fusisporium rubi*, on blackberry. (After Earle.)

RASPBERRY CANE RUST.

The cane rust of raspberries and blackberries is rather inconspicuous but very destructive. It was first studied by Professor T. J. Burrill, who published an excellent account of it in the *Agricultural Review* for November, 1882. He states that it was first observed in 1878; since that time it has spread rapidly and caused great damage. One instance is given of a blackberry field that yielded a profit of four hundred dollars a year with promise of increase. This disease reduced it so that it scarcely paid expenses the next year.

The spots on the canes are rounded, of a grayish color and surround-

ed by a definite and slightly raised border. The fungus fruits by sending out clusters of very short, unbranched threads, each of which produces at the tip a minute, colorless spore, about twice as long as broad. It has never been definitely classified, but Prof. Burrill is inclined to class it *Sphaceloma ampelinum*, which produces one kind of black rot in grapes.

Remedy: Cut out the canes as soon as the berries are picked and keep the field clean.

In conclusion, I will say a few words about the treatment of fungus diseases in general. The thing to be continually and most earnestly sought is a knowledge of the life histories of the different species. This is essential to intelligent action, and this is what all earnest students are seeking. Some progress has been made, with good practical results. But it requires an amount of time and patience, care and accuracy, that one can scarcely conceive of till he has tried it and one can scarcely hope for success unless he has an absorbing interest in the subject for its own sake, as a science. Most of the world's valuable discoveries have been made by men moved by the love of science. No mercenary motive will answer in its place.

Besides interest and ability, a man must have means to work with. He must know what others have done and what methods they have used to gain their results; hence, he must have books. He must have microscopes and other apparatus for his investigations. Considering the interests involved, the amount of damage done to crops by fungi, the desirability of increasing the knowledge of the subject and of educating the people with regard to it, it would seem that the State might well afford to furnish the means for subsistence and for investigation to someone who will give his best efforts and interest to the work, such interest as only the work itself will satisfy, and which is essential to success. Yet results can not be gauranteed and much patient waiting and disappointment may be unavoidable.

In the meantime the best general mode of treatment is to destroy all diseased plants or part of plants and any useless living plants or dead matters that might harbor disease.—*From the Minnesota Horticultural Report, vol. xiv, 1886.*

Also we find here, in a noted cañon of ancient cliff-dwellings near San Francisco Mountains, a large *Cystopteris*, uniformly bearing bulblets near the apex of the fronds. If this is the species *C. bulbifera* it has not before been reported so far west as Arizona.

Fort Moroni, near Flagstaff, Ariz., July 30, 1884.

On the Sexuality of the Fungi.¹

BY H. MARSHALL WARD.

I propose to show that it is probable that the sexuality of the higher Fungi has disappeared, because its purpose has been equally well or better attained otherwise than by means of sexual organs.

Preliminary to this it will be necessary to be quite clear as to what sexual organs and the sexual process essentially are.

The two points common to all the cases of sexual reproduction which have been directly observed are the following:

1. A larger or smaller quantity of protoplasmic material passes from one portion (the male organ) of the same or another individual, into the protoplasm contained in another portion (the female organ).

2. The protoplasm contained in the female organ therefore becomes capable of further development; either at once, or, more generally, after undergoing a period of rest.

It is not necessary to quote the numerous cases of observed analogies between the sexual reproduction of animals and plants; but will suffice to note that the essential in the sexual process is always the addition of a portion of protoplasm from the male, to the protoplasm of the female.

But this is not all. It is now well established in embryology that the normal ovum, or female mass of protoplasm, is incapable of further development until it has received the protoplasm of the male; that the latter, in fact, incites the former to further development.

The outcome of all we know of these matters leads to the conviction that we have in the germination or development of an

¹ The statement of the important hypothesis hereby presented is somewhat abbreviated from the concluding portion of a long and interesting article by Professor Ward, given under the same title. The review of the historic progress of our knowledge of sexuality in fungi, and the present state of such knowledge, with the numerous illustrative diagrams are necessarily omitted for want of space.—EDS.

oospore—and the same is true for an egg, etc., the terms being different—simply a *renewal* of the growth of the organism; and from this and other convictions follows the result that the formation of an oosphere, although it may take place after an accumulation of large quantities of food, implies a condition of weariness—if the term may be allowed—on the part of the protoplasm for the time being. No doubt the molecular energy of the protoplasm forming the oosphere, is less than that of the rest of the plant for the time being; the access of the antherozoid or male protoplasm, however, reinvigorates the sluggish mass, and renewed life ensues. This may require some time, however, and we may possibly not be far wrong if we imagine that interval to be occupied in molecular rearrangements in the mass.

But, although we can sum up the foregoing by saying that, after a time, protoplasm requires reinvigorating by the addition of fresh protoplasm from another source, it is extremely improbable that the protoplasm of the male and female organs is at all similar.

It now remains to be seen if we can throw any light on the curious disappearance of sexual organs and sexuality in the Fungi—curious, because the sexual process appears to be all but universal in all organisms excepting the very lowest.

A hypothesis which suggests itself, and which Eidam¹ favors, and which is certainly supported by some analogies, is to the effect that the apogamous Fungi, *i. e.*, those in which the sexual organs are totally suppressed, are not always apogamous. We know that many forms only produce their sexual organs at comparatively long and rare intervals. The *Mucors*, for instance, may be propagated through numerous generations by means of the asexual spores; the sexual organs only arising now and again under favorable conditions.

Moreover, the cases of polyembryony—where several embryos arise in an embryo sac, although only one oosphere is fertilized—favor the view that the effect of fertilization may be extensive; and we can not doubt that such is the case where adventitious covering branches arise after the conjugation of certain *Mucorini*, and in the *Orchideae*, where fertilization or even the mere growth of the pollen tube affects the whole flower.

The sexual act, however, consisting as it does simply or mainly in the reinvigoration of protoplasm by the addition of protoplasm of a different nature (though we do not know the kind or limit of difference), it may be that an explanation of what occurs

¹ Cohn's Beitr. zur Biologie, etc., B. iii, H. iii.

in the Fungi is afforded by their mode of life. The Fungi in which sexual organs seem to be most certainly absent are those which are most highly specialized as parasites. Now, we have every reason to believe, first, that parasitism is a matter of degree, and secondly, that the most highly specialized form of parasitism consists in directly obtaining those contents of the cells of the host which are chemically most complex, and therefore contain most energy.

I need not dwell on the degrees of parasitism exemplified by plants which merely rob their hosts of space or moisture, or which have obtained a hold so intimate that they break it up and feed on the rotting *debris*, but may at once pass on to consider a few consequences which follow from the mode of life of those highly specialised parasites which have become so closely adapted to their host, that they exist for a time as all but an organic part of its tissues and substance.

It can scarcely be doubted that the protoplasm of a higher plant, such as a phanerogam, differs from that of a lower cryptogam in being capable of doing more work, and that the great advantage derived by a parasitic Fungus which has its life so adapted that it can tax the cells of a phanerogamous host plant, is that it contains its food materials in a condition more nearly approaching that of its own substance, than would be the case if it had to work these materials up from inorganic matters.

Now it seems not improbable that the protoplasmic substance of a higher phanerogam may contain so much energy that it can not only supply the vegetative mycelium of a parasitic fungus with all that it requires for its immediate growth, but also suffices to enable that fungus to store up enough energy in its asexual or apogamous spores to last until the next generation of the fungus gains its holdfast on another (and it may be distant) source of life-giving substance.

Let us take the case of a uredinous fungus parasitic in the leaves of a phanerogam. We know that the substances necessary for the whole growth of the phanerogam are formed in the cells of the leaf; not only so, the matters which eventually find their place in the reproductive organs must be formed there also, potentially at least. The leaf of a phanerogam so attacked, moreover, is able to support the parasitic fungus for a long time uninjured, as I have convinced myself by experiment, and there can be no doubt that substances pass into the fungus which would normally have passed into other parts of the host plant itself.

But we may imagine even this to fail after a time—we may

suppose that at length the Fungus derives too little benefit to be able to go on; or the season during which the host plant flourishes is drawing to an end.

No doubt we have in heterœcism the salvation of such a Fungus. Not only is it carried through a dangerous period, by seeking relief at the hands of a second host, but—and which I believe to be far more important—it obtains reinvigoration by the new protoplasm with which it comes in contact. We may not inaptly compare the sojourn of the Fungus on its second host, to a trip to the seaside, where the weary and enfeebled organism enjoys fresh diet and associations for a time, which in their turn pall and prepare the recipient to renew old modes of life.

We have seen that the disappearance of the sexual organs, leading to apogamy, commences especially in the lower *Ascomycetes*, and it may be more than a coincidence that epiphytic forms, which show a tendency to produce one kind of spore while on the living leaf and develop their asci on the fallen leaf are common here; such forms suggest how the parasitism and heterœcism of higher forms may have begun, and it is remarkable that the apogamy becomes more and more complete as we ascend through the latter.

It is not pretended that the hypothesis embodied above at once explains all the cases possible, and it will be well to state a few of the difficulties at once. The *Basidiomycetes* I shall not dwell upon, since our knowledge of them is still very imperfect.

The difficulty may suggest itself to many that there are parasitic fungi—such as the *Peronospora*—which nevertheless develop the sexual organs in the condition typical and perfect for the group to which they belong. I have already referred to the fact that many of these forms are really saprophytes, and that others break down and destroy the tissues of their hosts—clumsily killing their prey, and then feeding on the rotten mass—and have pointed out that this is a much less specialized form of parasitism than that of the higher Fungi and *Ustilagineæ*.

Nevertheless, the sexuality shows signs of disappearance in extreme members. De Bary¹ shows that in *Phytophthora* and *Peronospora* there is a less evident passage over of protoplasm from the antheridium to the oosphere than in *Pythium*; and that in some cases, indeed, the quantity passing over is too small to be observed. I will not attempt to lay stress on the coincidence that in *Phytophthora infestans* (the fungus of the potato disease) no sexual act has yet been discovered.

¹ Beitr. zur Morph., etc., der Pilze, iv, p. 72.

Another obvious objection may be raised as follows:—The *Saprolegniae* are in the main saprophytes, and yet they are said to be advanced towards apogamy—parthenogenetic, at any rate. The answer may be that they are saprophytic chiefly on animal protoplasm, which contains more potential energy than does vegetable protoplasm. At the same time, some *Saprolegniae* are parasitic on plants, and *S. ferax* now appears to be parasitic on fish¹.

I may say, in conclusion, that it was during the study of the parasitic fungus of the coffee disease (*Hemileia vastatrix*)² in Ceylon that I was first led to speculate on the enormous amount of energy displayed by an organism which shows not the remotest satisfactory trace of sexuality, but which reproduces itself through many generations exclusively by means of asexual spores. That this energy of reproduction is derived from the coffee tree there can be no doubt, and that it is at the cost of the reproduction of the host is sadly evident; the clear inference from the fact that the coffee leaf supplies substance for the reproduction, etc., of a fungus at the expense of its own fruit, is that the fungus takes matters which are very rich in energy, so rich, indeed, that the fungus is not necessitated to sort these substances in special reproductive organs, and to secrete sexual elements, one of which would then reinvigorate the other, but may employ them forthwith for the purposes of its own relatively simpler existence and reproduction — *Quart. Jour. Mic. Sc.*, April, 1884.

GENERAL NOTES.

Polarity of Lettuce Leaves.—The orientation of the leaves of *Lactuca Scariola*, which has made it one of the two best known “compass” plants, is repeated in a less degree in the leaves of the common garden lettuce. The polarity is scarcely apparent until the lettuce begins to throw up the flowering stem. It is very weak in the curled and wrinkled varieties, but it is well marked in the Cos varieties, which have flat narrow leaves much like the wild *L. Scariola*. The observation was made on over one hundred varieties of lettuce grown the present season in the garden of the New York Agricultural Experiment Station.—J. C. A.

Hibiscus Moscheutos and H. roseus.—Dr. J. Guillaud, of Bordeaux, sends a pamphlet containing his investigations resulting in the identification of

¹ Prof. Huxley, ‘*Quart. Jour. Mic. Sc.*’, 1882. [It may be found upon otherwise healthy salmon, according to the investigations of Mr. George Murray. ‘*Science*’, IV, p. 27.—Eds.]

² [‘*Quart. Jour. Mic. Sc.*’, Jan. 1882; noticed and figured in ‘*Am. Nat.*’, July, 1882.—Eds.]

Hibiscus roseus of Thore—a species supposed to be indigenous to the southeastern coast of France, also found in Italy—with our Marsh Hibiscus. He is not aware that the same identification has been made by Mr. Daydon Jackson, and published a year or two ago in the nineteenth volume of the *Journal of the Linnean Society*, London. Dr. Guillaud has had the advantage of seeing the two plants growing spontaneously, ours in the neighborhood of New York, the other in the marshes of the Landes. *H. roseus* has also been found in North Italy, in the marshes of the Po and lagunes of the Adriatic, and, according to Dr. Guillaud, specimens have been received from Asia Minor, but no mention is made of it in Boissier's *Flora Orientalis*.

Is this species indigenous to Europe as well as to the Atlantic coasts of North America? Is it a survival from the time when the floras of Europe and Eastern America had more common elements than they now have? Or has it somehow been conveyed across the Atlantic, and if so, whether at some early period, or within historic times? Questions not easily answered. If the first, then this plant, like a few others that might be named, is in Europe what *Convallaria majalis*, *Littorella lacustris*, *Marsilia quadriflora*, *Scolopendrium* and perhaps *Calluna* are in North America. In favor of the second view, and even of a late and casual introduction, it is to be said, as Dr. Guillaud notes, that Thore found the plant on the coast of France only at the beginning of this century; that it was unknown to Tournefort, who botanized around Bayonne in the autumn of 1688; that the plant has disappeared from the particular stations where Thore found it and where it was said to abound, and that it is now more rare than formerly. Its spread from the Atlantic coast to that of the Adriatic may be owing to the carriage of seeds by marsh birds. Indeed, Dr. Guillaud thinks it may have been brought to Europe by sea birds. On the other hand, since it is now found in the district near Mantua, he quotes the lines in Virgil's *Eclogues*, in which the stems of *Hibiscus* are twice mentioned, in a way by no means *mal-a-propos*; but he thinks they might as well apply to Marsh-mallow. It appears that the specific name *Moscheutos* came to Linnaeus through Cornuti, from a "*Rosa Moscheutos*" of Pliny, some kind of Rose-mallow, we may suppose. Since the two Linnean species are clearly one, it is a pity that the name *H. palustris* was not chosen. Torrey and Gray are responsible for that. The reason of the choice was, that *H. Moscheutos* stands first in the book, and *H. palustris* is merely differentiated from that—reasons which need not have prevailed.

A. GRAY.

Vincetoxicum.—Following some authority, which it is now not worth while to look up, it appears that in the Synoptical Flora of N. America, I had derived this name from "*vincies*, that serves for binding" and *toxicum*. Dr. Hance, in Britten's *Journal of Botany* for May, 1883, notes, (1) that the only authority for this adjective is a line of Plautus in which *vincea* is now known to have been a mistake of some copyist for *junceæ*, and (2), that the old herbalists, Fuchs and Matthioli, clearly indicate that the Latin part of this hybrid name is from *vincere*, to conquer.

Stipules in Saxifragaceæ are of small account, as Prof. Coulter's pupil show me by sending *Mitella diphylla* with good stipules between the cauline leaves. It seems to be regularly so.

"Breweria minima," Gray, in Proc. Am. Acad. xvii, 228, is *Convolvulus pentapetaloides* of Linnaeus, and doubtless was introduced into California from the Mediterranean region, probably with grain. It turns up from various parts of California of late. The style and stigmas are truly as in *Convolvulus*.

A. GRAY.

(Reprinted from BULLETIN OF THE TORREY BOTANICAL CLUB, Vol. XIII., No. 11.)

A Letter from Commissioner Colman to the Botanists of the A. A. A. S.

DEPARTMENT OF AGRICULTURE, }
WASHINGTON, D. C., October 20th, 1886. }

To the Members of the Botanical Club of the American Association for the Advancement of Science :

LADIES AND GENTLEMEN : I have noted with extreme gratification the interest you have taken in the work established by me in this Department, relative to the investigation of the fungus diseases of plants ; and the resolutions you have passed commending my action and assuring me of your support and aid in securing the necessary means for the continued and successful prosecution of this most important undertaking are fully appreciated, and I wish to thank you on behalf of the farmers and fruit growers of the country, in whose interest and for whose direct benefit this work is designed.

As you are well aware, only a few of the more important plant diseases have been thoroughly worked out by scientists, and the little that has been done—little when compared with what there is to do, but a great deal when considered by itself—has been the result of private effort on the part of some of your own well-known members. Such obscure diseases as the peach yellows, the cotton rust, and the “foot rot” of the orange tree, demand immediate attention, and, for their proper elucidation, we need to command the services of our most skillful investigators, giving them opportunities to make special studies in the field until the knowledge desired is gained. As you have well suggested, a liberal supply of funds is required for this work.

In addition to the assistance in this particular, to which you have so generously pledged yourselves, I beg leave to call your attention to the fact that you, as botanists, knowing our cultivated and native plants and the fungus parasites infesting them, may do much valuable service as *observers* in your respective localities, by recording such facts as may come to your notice relating to this subject, and by collecting and transmitting to the Department material useful in the investigations, or that may serve to record the distribution of the injurious species of fungi.

Facilities for this work, and a free use of the mails, will be accorded those who may have such notes or materials to transmit, and the source of all matter that may be used for publication will be properly credited.

Again thanking you for your hearty commendation of my course in relation to the subject, and assuring you that I shall continue to do all in my power to further the work, I remain,

Yours respectfully,

NORMAN J. COLMAN,
Commissioner.

FUNGICIDES, OR REMEDIES FOR PLANT DISEASES.

U. S. DEPARTMENT OF AGRICULTURE, BOTANICAL DIVISION,

SECTION OF VEGETABLE PATHOLOGY—CIRCULAR NO. 5.

In order to bring to the attention of the Directors of the Agricultural Experiment Stations, as well as to lay before others directly interested in the treatment of the fungus diseases of plants, the following notes on the subject, and formulas of fungicides of real or supposed value, have been prepared by my direction for distribution.

NORMAN J. COLMAN,

Commissioner of Agriculture.

21004 FU—1

QK
2.5
108
1.100

SIR: I submit herewith a circular on fungicides, prepared in accordance with your instructions.

Respectfully,

F. LAMSON SCRIBNER.

Hon. NORMAN J. COLMAN,
Commissioner of Agriculture.

FUNGICIDES.

INTRODUCTORY.

In combating the various fungus diseases of plants many chemicals and chemical compounds have been tried from time to time, and not a few nostrums of the quack order have been widely advertised for this or that, or for every distemper that plants are heir to. A little knowledge of vegetable anatomy and of the character and habits of the fungi to be treated would enable us to avoid the impositions of quacks and assist us in determining the *rationale* of any proposed treatment. The diseases in plants caused by fungi are simply the effects produced by other plants of parasitic habits, and we must keep the two—the parasite and the plant attacked—distinct in our minds in our efforts to protect the one from the evils produced by the other.

For some of these so-called diseases there is no remedy but the knife or the complete destruction of the infested plant. It is important to understand the cases of this character, not only that we may avoid wasting time and money in vain efforts to treat them otherwise, but in order that prompt action may be taken and sources of infection be quickly destroyed, for all fungus diseases may be regarded as infectious. A large class of these diseases, many occasioning heavy annual losses, may be mitigated or entirely overcome by the application of certain remedial or preventive agents. Those remedies or preventives which have apparently yielded positive results are here enumerated, together

with directions for their preparation, mention also being made of the diseases for which they have been employed.

APPLYING THE REMEDIES.

In applying the remedies we must consider the period and manner of the attack made by the parasite and the manner of action of the substance applied. Fungi living within the tissues of the host must be prevented from gaining an entrance to these tissues; fungi which live upon the surface of plants or having their bodies soon exposed through the breaking up of the epidermis, like the apple-scab fungus or the fungus of bird's-eye rot of grapes, may be treated by curative methods. The compounds having sulphate of copper for a base are, so far as we know to-day, the best preventive remedies: sulphur, sulphur and lime, sulphide of potassium, hyposulphite of soda, are to be classed as curatives.

Destructive treatments are available between the periods of vegetation (winter season), and consist in destroying all infectious material and in washing the plants to be protected with strong caustic solutions, *e. g.*, solutions of sulphate of iron or copper and sulphuric acid.

During the growing season the strength of the solutions used is governed by the power of the green tissues to resist their action. In the early part of the season while the shoots and leaves are yet tender, weaker solutions than those which may safely be applied later in the season, must be employed. There is great diversity of susceptibility to the caustic action of the remedies between different species of plants and even between different varieties of the same species. The conditions of the weather at the time the applications are made may be of importance in this connection. Sulphur alone, applied when the weather is very hot and the sun bright, may cause a burning of the foliage. The same is true of sulphatine and also of eau celeste.

Explicit directions can not be given for these cases. Much has yet to be learned by experiment, and each one must exercise judgment and govern himself by surrounding circumstances.

Avoid making the applications excessive; do not drench the plants with the fluids nor plaster them with the powders. With a suitable spraying apparatus, which projects a fine, mist-like spray, merely wet the plant surfaces, and employ bellows which will discharge the powder evenly and in such a manner that the plants may be enveloped in a cloud of dust, which, settling upon all parts, becomes just perceptible.

The necessity for good and easily-worked spraying pumps and bellows is evident. For small plantations and general vineyard use, the knapsack form of sprayer, having the reservoir and pump combined, to be carried on the back of the operator like a knapsack, is the best. For spraying fruit-trees more powerful appliances are required.

Nixon's Climax nozzle is excellent for spraying clear liquids, but its use demands considerable power in the pumps.

FIG. 1.—Nixon's Climax Nozzle.—This is made in three sizes, differing in size of jet and "no." of the gauze used to cover the top.

The Vermorel modification of the eddy chamber or cyclone nozzle is a most excellent pattern for both clear and pasty or thick liquids. The degorger combined with it renders the spraying of the latter possible.

FIG. 2.—The Vermorel Nozzle with Lance.—The hose is attached to the lance at H. C., stop-cock. d, exposed end of the degorger; at the other end of the degorger is a pin which may be made to pass through the opening of the nozzle, clearing it when clogged. The degorger is pushed in by the rod R, sliding through the fixed bearings b, b, b. The force of liquids under pressure will bring it back into position.

LIQUIDS.

1. Simple solution of sulphate of copper.

(Comptes Rendus, Paris, October 5, 1885; Bull. 2, Bot. Div. U. S. Dept. of Agr., 1886, p. 89; Report of the Mycologist, 1886, p. 100; Circular No. 3, Sect. of Veg. Pathology, April, 1887.)

Dissolve 1 pound of pure sulphate of copper in 25 gallons of water. For treatment of downy mildew and oïdium of the vine. For treatment of downy mildew and black-rot of the grape.

2. Simple solution of sulphate of copper.

(Diseases of Field and Garden crops, Smith, 1884, p. 261; Report of the Assistant Botanist, U. S. Dept. of Agr., 1885, p. 79; Report of the Kentucky Expr. Sta., 1887, p. 14; Bull. No. 3, Central Ex. Farm, Ottawa, Canada, p. 14.)

Simple solution in water, 5 to 8 pounds to 10 gallons.

For soaking grains previous to sowing to destroy the spores of smuts.

3. Copper mixture of Gironde, Bordeaux mixture.

(La Vigne Americaine, June, 1886, p. 193; Bull. 2, Bot. Div., U. S. Dept. of Agr., 1886, p. 15; report of the Mycologist, U. S. Dept. of Agr., 1886, p. 100; Circular No. 3, U. S. Dept. of Agr., April, 1887; Circular No. 4, U. S. Dept. of Agr., July, 1887; Le Mildiou, G. Patrigeon, 1887, p. 98; Traitement du Mildiou, Millardet, 1887, p. 35; Manuel Pratique pour le Traitement des Maladies de la Vigne, Viala and Ferrouillat, 1888, p. 26.)

Original formula.—Dissolve 16 pounds of sulphate of copper in 22 gallons of water, in another vessel slake 30 pounds of lime in 6 gallons of water. When the latter mixture has cooled it is slowly poured into the copper solution, care being taken to mix the fluids thoroughly by constant stirring. It is well to have this compound prepared some days before it is required for use. It should be well stirred before applying.

Numerous modifications in the preparation of this compound have been suggested, chiefly for the purpose of reducing the amount of copper.

A solution containing the ingredients in the following proportions has been recommended for general use:

Sulphate of copper	pounds..	4
Lime	do....	4
Water.....	gallons..	22

The copper is dissolved in 16 gallons of water, while the lime is slaked in 6 gallons. When cool the solutions are mixed as described above.

For treatment of mildew. For downy mildew and black-rot of the grape. For blight and rot of the tomato and potato.

4. Eau Celeste, Audouinaud process.

(Progrès Agricole et Viticole, March 28, 1886; Report of the Mycologist, 1886, p. 102; Circular No. 3, Sect. of Veg. Pathol. U. S. Dept. of Agr., April, 1887; Circular No. 4, Sect. of Veg. Pathol. U. S. Dept. of Agr., July, 1887; Sixth Report of the New York Experiment Station, 1887, p. 106; Le Développement et le Traitement du Mildiou et de L'Anthracnose, Millardet, 1887.)

Dissolve 1 pound of sulphate of copper in 2 gallons of hot water; when completely dissolved and the water has cooled, add 1½ pints of commercial ammonia (strength 22° Baumé); when ready to use dilute to 22 gallons. The concentrated liquid should be kept in a keg or some wooden, earthen, or glass vessel.

For downy mildew. For treatment of downy mildew and black rot of the grape. For treatment of mildew and anthracnose. For blight and rot of the tomato and potato. For apple scab.

5. Eau Celeste, modified formula.

(Progrès Agricole et Viticole, July, 1887, p. 18; Etc.)

Sulphate of copper	pounds..	2
Carbonate of soda	do....	2½
Ammonia (22° Baumé)	pints..	1½
Water.....	gallons..	22

Dissolve the sulphate of copper in 2 gallons of hot water, in another vessel dissolve the carbonate of soda in a similar manner; mix the two solutions, and when all chemical reaction has ceased add the ammonia; then dilute to 22 gallons.

6. Solution of ammoniacal carbonate of copper.

(Progrès Agricole et Viticole, August, 1887, p. 114; Etc.)

Prepared as follows: Into a vessel having a capacity of two quarts or more pour one quart of ammonia (strength 22° Baumé), add 3 ounces carbonate of copper, stir rapidly for a moment and the carbonate of copper will dissolve in the ammonia, forming a very clear liquid. The concentrated liquid thus prepared may be kept indefinitely. For use, dilute to 22 gallons.

For peronospora of the vine.

7. Sulphate of iron.

(La Vigne Américaine, June, 1884, p. 181; Bull. 2, Bot. Div. U. S. Dept. of Agr., 1886, pp. 38 and 121; Report of Mycologist, U. S. Dept. of Agr., 1886, p. 114; Etc.)

Simple solution in water 4 to 8 pounds to the gallon, to be used only as a wash.

For anthracnose.

8. Chloride of iron.

According to Nature (vol. xxxvii) a very dilute solution of the above-named substance has been used with success in combating the coffee disease due to *Hemileia vastatrix*. This fungus is a member of the group *Uredineæ* (rusts), and it is said that the chloride of iron not only cures the disease, but also prevents its recurrence. The solution is applied to the under surface of the leaves by means of a pulverizator or spraying apparatus. Its sticky nature causes it to adhere for two months.

9. Skawinski's sulphate of iron and sulphuric acid solution.

(Traitement des Maladies de la Vigne, Viala et Ferrouillat, 1888, p. 84.)

Sulphate of iron.....	pounds..	110
Sulphuric acid (53°).....	pints..	1½
Warm water.....	gallons..	22

"In order to avoid sputtering it is necessary to be very careful in pouring the sulphuric acid upon the crystals of iron sulphate, and not to add the warm water until after this operation. Besides, it is best to use sulphuric acid of only 53° Beaumé, because it is less dangerous to handle. I am careful to use this mixture the same day and while warm, for when it becomes cold the sulphate of iron crystalizes and the solution has no longer the strength necessary in order to be efficacious."

10. Sulphide of potassium, Liver of sulphur.

(Garden 1885, p. 379; Gardeners' Chronicle, 1885, Vol. XXIII., pp. 276, 352, 378; Vol. XXIV., p. 379; Bull. 2, Bot. Div. U. S. Dept. of Agr., 1886, p. 67; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 121; sixth report of the New York Experiment Station 1887, pp. 348, 349, 350; Etc.)

Simple solution in water, ¼ to 1 ounce to the gallon.

For mildew in greenhouses. For mildew on roses. For oidium and erinose of the vine. For orange-leaf scab. For celery-leaf blight. For pear and apple scab.

11. Solution of hyposulphite of soda.

(Report of the New York Experiment Station, 1885, pp. 231, 232; 1886, pp. 177, 178; 1887, pp. 99, 100; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 120; Etc.)

Simple solution made by dissolving 1 pound of the soda in 10 gallons of water. Must be used as soon as prepared.

For apple scab. For celery-leaf blight. For orange-leaf scab.

12. Liquid Grison. Eau Grison.

(American Agriculturist, 1885, p. 295; Bull. No. 2, Bot. Div. U. S. Dept. of Agr., November, 1886, p. 26; Report of the Mycologist Sect. of Veg. Pathology, 1886, p. 21; Etc.)

Prepared by boiling 3 pounds each of flowers of sulphur and lime in 6 gallons of water until reduced to 2 gallons, when settled pour off the clear liquid and bottle it. When used, mix 1 part of the clear liquid in 100 parts water.

For mildew on grape vines. For powdery mildew of the vine.

13. Milk of lime.

(Bull. No. 2, Bot. Div. U. S. Dept. of Agr., 1886, p. 70; Report of the Mycologist Sect. of Veg. Path. U. S. Dept. of Agr., 1886, p. 100; Développement et le Traitement du Mildiou et de L'Anthracnose, Millardet, 1887; Etc.)

Simple solution in water, 2 to 6 parts lime to 100 parts water.

For peronospora of the vine. For anthracnose.

14. Phenic acid. Carbolic acid.

(La Vigne Americaine, July, 1884, p. 197; Journal de Agr. Pratique, p. 76, 1884; Etc.)

Simple solution in water one-half pint to 10 gallons.

For powdery mildew of the vine.

15. Phenic acid in solution diluted with soap-suds and glycerine.

(Report of Mycologist U. S. Dept. of Agr., 1886, p. 121.)

Soap-suds 10 gallons. glycerine 1 pound, carbolic acid one-half pint; mix thoroughly to form an emulsion.

For orange-leaf scab?

POWDERS.

16. Sulphur.

(Bull. No. 2, Bot. Div. U. S. Dept. of Agr., 1886, pp. 26, 27; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 108; Etc.)

For grape mildew. For powdery mildew of the vine.

17. Sulphur and lime.

A mixture of sulphur and lime in equal parts by weight.

For treatment of anthracnose during the growing season.

18. Blight powder, Sulphated sulphur.

(Progrès Agricole et Viticole, 1887, p. 39; Report of the Mycologist U. S. Dept. Agr., 1886, p. 103; Circular No. 4, Section of Veg. Pathology, U. S. Dept. of Agr., July, 1887.)

Prepared by thoroughly mixing from 3 to 8 pounds of anhydrous sulphate of copper with 90 to 100 pounds of flowers of sulphur.

For simultaneous treatment of oïdium and the downy mildew. For downy mildew of the vine. For tomato and potato blight and rot.

19. Sulphatine, the Estève process.

(Bull. de Cent. Soc. de France, 1887, p. 316; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 102; Circular No. —, U. S. Dept. of Agr., April, 1887; Circular No. 4, U. S. Dept. of Agr., July, 1887; Etc.)

Mix 2 pounds of anhydrous sulphate of copper with 20 pounds of flowers of sulphur and 2 pounds of air-slaked lime. The proportions may be varied.

For the treatment of mildew. For the treatment of the downy mildew and black rot of the grape. For the treatment of the tomato and potato for blight and rot.

20. Skawinski's powder.

(La Vigne Américaine, Nov., 1886; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 104; Cours Complet de Viticulture, G. Foex, 1888, p. 485; Progrès Agricole et Viticole, Mar., 1888, p. 222; Etc.)

Mix 22 pounds of finely powdered sulphate of copper with 33 pounds of soot or alluvial earth and 165 pounds of coal dust.

For simultaneous treatment of oïdium and downy mildew of the vine. For treatment of mildew.

21. Sulfosteatite or cuprique steatite.

(Le Mildiou, par Dr. G. Patrigeon, 1887, p. 134, Journal d'Agriculture Pratique, 1887, p. 768; Etc.)

An exceedingly fine bluish powder composed of steatite, or talc, and sulphate of copper, the proportion of the latter substance amounting to about 10 per cent. Very easily applied; considered the most adherent of all the powders.

For the treatment of mildew (*Peronospora*).

22. David's powder.

(Journal d'Agr. Pratique, Nov. 25, 1886; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 102; Nouvelles Recherches sur le Développement et le Traitement du Mildiou et de L'Anthracnose, Millardet, Feb., 1887, p. 22; Circular No. 3, Sect. of Veg. Pathol., U. S. Dept. Agr., April, 1887; Etc.)

Dissolve 4 pounds of sulphate of copper in the least possible amount of hot water, and slake 16 pounds of lime with the smallest quantity of water required. When the copper solution and slaked lime are completely cooled mix them together thoroughly; let the compound dry in the sun, crush and sift. Apply with a sulphuring bellows furnished with an outside receptacle for the powder. The copper coming in contact with the leather will soon destroy it.

For downy mildew. For downy mildew and black rot of the grape.
For mildew and anthracnose.

23. Podechard's powder.

(Bull. Soc. Agr. Herault, France, 1886, p. 59; *La Vigne Américaine*, June, 1886, p. 193; Bull. 2, Bot. Div. U. S. Dept. of Agr., Nov., 1886, pp. 80, 81, 82; Report of the Mycologist U. S. Dept. of Agr., 1886, p. 101; *Le Développement et le Traitement du Mildiou et de L'Anthracnose*, Millardet, Feb., 1887; Etc.)

	Pounds.
Air-slaked lime	225
Sulphate of copper	45
Flowers of sulphur	20
Ashes	30

Dissolve the sulphate of copper in the water; when thoroughly dissolved pour the solution upon the lime, which is surrounded by the ashes to keep the liquid from spreading; after twenty-four hours add the sulphur, thoroughly mix the compound, ashes and all, and when dry sift through a sieve with meshes of one-eighth of one inch. This preparation may be made several months before it is required for use.

For the downy mildew of the vine. For the treatment of mildew and anthracnose.

WASHINGTON, D. C., April 25, 1888.

Compliments of
Roscoe Pound

No. 15

The Sychytria, Mucoraceae, and Entomophthoraceae
OF NEBRASKA.

By ROSCOE POUND.

EXTRACTED FROM
FLORA OF NEBRASKA,
Part I.

18

25

9E

1.100

Family.—CHYTRIDIACEAE.*

"Mycelium lacking, or developed in the form of delicate protoplasm-threads, more seldom clearly hyphae-like, unicellular. Sporangioophores wanting or but poorly developed. Sporangia always forming swarm spores, thin walled and quickly ripening, or thick walled and resting for a period (resting sporangia). Sexual spore formation only developed in a few forms; difference between antherid and oogone scarcely marked." (Schroeter).

This group contains about 35 genera and 160 species of minute, microscopic fungi, which are almost all parasites. They are found parasitic chiefly on algae, but also on flowering plants, fungi, and some lower animals (worms). They seem to be closely allied to the algae of the *Protococcoideae*, to which the simpler forms have many striking resemblances.

The family is divided into 6 sub-families, of which we have as yet found representatives of but 1 in the state.

SUB-FAM.—**Synchytriaeae**.—Swarmsporangia formed by simultaneous division, united into a sorus or arranged in a row. Restingsporangia formed directly from the whole fruiting body, or by division, producing a sorus of restingsporangia.

1. SYNCHYTRIUM DeBary & Woronin Bericht. d. Naturf. Ges. Friburg. III. (Ex Fischer). 1863.

Parasites in the epidermal cells of flowering plants forming galls. Sori of swarmsporangia (summer sori) formed directly from the vegetative body, surrounded by the colorless membrane of the mother-cell, consisting of a number of close laid sporangia which by pressure are polygonal; swarmspores globular, one ciliate; resting spores round or elliptical, with a thick brown exospore.

Etymology: Greek *συν*, together, and *χυτρίς*, pot.

Synchytrium fulgens SCHROETER. Hedwigia XII, 141. 1873.

Spots minute, purple; galls small, orange red; sori bright yellow, spherical or elliptical, produced singly in the epidermal cells which they completely fill, 60–100 μ in diam.; resting spores spherical, with a smooth, thick, brown exospore, 66–82 μ .

On leaves of *Oenothera biennis*. Not uncommon.

Synchytrium peckii (THUEMEN).

Uredo acididioides PECK. 24th Rep. 88, 1870. Not *U. acididioides* DC. Fl. Fr. II., 236, (1815) = *Melampsora acididioides* SCHROETER.

Uredo peckii THUEMEN Mycotheca Universalis no. 538. 1876.

S. fulgens decipiens FARLOW Bull. Buss. Inst. II. 229. 1878.

S. decipiens FARLOW Bot. Gaz. 1885, p. 240.

Spots bright yellow, galls hemispherical, sori spherical, bright yellow, 180–200 μ in diam.; sporangia very numerous, 15 μ in diam.; resting spores unknown.

On leaves of *Falcata comosa*, Lincoln. Pl. XV., Fig. 1, a. b.

Order 4.—CONJUGATAE.—Unicellular or filamentous, ~~chlorophyll-green or colorless; propagation by cell division; reproduction by the union of similar cells (isogametes).~~

* By Roscoe Pound.

1. MUCOR LINNE Spec. Pl. II., 1655. 1753.

Saprophytic; mycelium spreading in and upon the substratum; sporangio-phores springing up here and there on the mycelium, simple or branched; sporangia round, many-spored; zygosporos borne on the mycelium naked, the copulating branches (suspensors) without outgrowths.

Etymology: Latin *mucor*, mould.

This was the name of one of the eleven genera under which Linne in his Genera Plantarum included all fungi.

Mucor mucedo LINNE l. c. (in part).

Sporangiophores erect, rigid, simple, 2-15 cm. high; sporangia large, round, 100-200 μ in diam., the membrane quickly disappearing, leaving a small collar-like fragment at the base; columella high-arched, cylindrical or truncate-conical, 70-140x50-80 μ ; spores rounded, cylindrical, or long ellipsoid, 6-12x3-6 μ or sometimes larger, colorless or light yellow.

In my specimens the spores are regularly 8-10 μ , about half as wide and rather strongly tinged with yellow.

On excrement of animals and various decaying substances the world over. Quite common on decaying insects in the water around Lincoln. Pl. XIV., Fig. 1, a, b, c.

Mucor racemosus FRESENIUS Beitrage 12. 1850.

Chlamydomucor racemosus BREFELD Untersuch. 1890.

Sporangiophores erect, of various sizes, 5-40 mm. high or small and frail, richly and irregularly branched, each branch terminating in a sporangium; sporangia small, round, of various sizes, depending on the nourishment, 20-70 μ in diam., the membrane not dissolving but splitting; columella broad clavate or obovate; spores round or short ellipsoid, smooth, colorless singly but in mass yellowish, 6-10x5-8 μ .

When grown in a solution it forms septa rapidly and grows by budding. In this condition it forms ellipsoid or rounded-oblong chlamydospores here and there in the hyphae and even in the sporangiophores. In its budding state it is a ferment.

On decaying organic substances the world over. On paste in the botanical laboratories at the University and very common in solutions, neglected culture-media, etc., in the laboratory. Pl. XIV., Fig. 2, a, b, c.

2. PHYCOMYCES KUNZE Mycol. Hefte II., 113. 1823.

Mycelium radiate; sporangiophores simple, arising singly, bronze-green, strongly metallic, terminated by a large sporangium; sporangia round many-spored, the membrane dissolving; columella pear-shaped; conjugating branches tong-shaped, the suspensors producing dichotomously branched, dark brown projections.

Etymology: Greek *φυκος*, a'ga, and *μυκης*, fungus.

Phycomyces nitens (AGARDH) KUNZE l. c.

Uva nitens AGARDH. 1817.

The characters of the genus. Sporangio-phores 7-30 cm. long; sporangia very large, about 1 mm.; spores ellipsoid, 16-30x8-15 μ .

A beautiful species, quickly recognized by its metallic appearance. The sporangiophores have the look of small, flattened wires.

On greasy, oily substances. Also found on a squash at Lincoln. Pl. XIV. Fig. 3, a, b, c.

[*Spinellus rhombosporus* (EHRB.), *S. fusiger* (LK.) VAN TIEGH., is found on decaying agarics. It may be distinguished by its aerial mycelium which is covered with single or 2-4 verticillate, pointed, thorn-like branchlets. This species is reported for North America, but has not been met with as yet in this state.]

[*Syzygites aspergillus* (SCOPOLI), *Sporodinia aspergillus* (SCOP.) SCHROETER, is a parasite or saprophyte on fleshy fungi. The sporangiophores are dichotomously branched above, and the zygospores are produced in large numbers on specialized hyphae. It is not yet certainly reported from North America, but is very liable to be found.]

TRIBE.—RHIZOPEAE.—Mycelium of two sorts, the vegetative growing on the substratum, and the fertile or aerial mycelium which grows by stolons and upon which the sporangiophores are borne.

3. ASCOPHORA TODE Fung. Mecklenb. I., 13. 1790.

Fertile mycelium at first white, then brown or brownish black, growing in all directions by stolons which fasten here and there by rhizoids and at these points produce one or more sporangiophores and other stolons, sporangiophores swelling just below the sporangia; sporangia hemispherical, the membrane entirely disappearing; columella hemispherical, forming with the terminal swelling of the sporangiophore, a club-shaped head which collapses and has the appearance of an umbrella; zygospores naked.

Etymology: Greek *ασκος*, sac, and *φορεω*, to bear.

Ascophora mucedo TODE l. c.

Mucor stolonifer EHRB. Sylv. Myc. Berol. 25. 1818.

Rhizopus nigricans EHRB. Nov. Act. Acad. Leopold. X., 1, 198. 1820.

Mucor clavatus Lk. Sp. Pl. VI., 1, 92. 1824.

Stolons creeping here and there over the substratum, quickly covering it, at first colorless, then brown; branches 1-3 cm. or longer; rhizoids more or less branched; sporangiophores rarely single, usually in clusters of 3-5 or more on each node, $\frac{1}{2}$ to 4 mm. high; sporangia hemispherical, 100-350 μ wide; columella broad hemispherical, high-arched, with the swelling of the sporangiophore forming a clavate, cylindrical head reaching almost to the tip of the sporangium, usually collapsing after the dissolution of the sporangium membrane, and remaining a long time covered with spores; spores of various sizes and shapes, irregularly globose, or oval, with one or two truncated corners, somewhat longer than broad, thick walled, finely striate, averaging 6-17 μ .

On all kinds of decaying organic matter—one of the commonest of fungi. Quickly recognizable by its mode of growth and the peculiar umbrella-like appearance of the collapsed columella.

Mucor clavatus Lk., Webber's Catalogue No. 129, belongs here. The specimen there referred to is principally *A. mucedo*, but it seems to have grown over another mould which from the spores is doubtless *Mucor mucedo*, and the spores of the *Ascophora* are somewhat larger than usual. Pl. XIV., Fig. 4, a, b, c, d.

TRIBE.—PILOBOLEAE.—Membrane of sporangium of two parts; the upper half cuticularized and permanent, the lower thin and quickly dissolving.

4. HYDROGERA WIGGERS Fl. Holsat. 110. 1780.

Sporangiophores simple, arising singly from swellings in the mycelium, colorless or orange, above expanding into a large ellipsoid swelling; sporangia hemispherical or lens-shaped, many-spored; the membrane above black and cuticularized, the lower half quickly disappearing and leaving the upper part resting on the conical columella; both at maturity thrown off by tension of the terminal swelling of the sporangiophore; zygospores, naked, borne on tong-shaped branches.

Etymology: Greek *ὕδωρ*, water, and Latin *gero*, to carry.

Hydrogera obliqua (SCOP.) OK. Rev. Gen. 855. 1891.

Mucor obliquus SCOPOLI Flor. Carniol. II., 494. 1772.

Hydrogera crystallina WIGGERS l. c.

Pilobolus crystallinus TODE Schrift. Naturf. Freund. Berl. V. 46. 1784. (Ex Fischer).

Sporangiophores arising singly from a bladder-like swelling of the mycelium, 5-10 mm. long, the terminal swelling ellipsoid or ovoid, .85-1.30x.60-.85 mm.; sporangia plano-convex, resting on the side of the terminal swelling 300-400x100-150 μ ; columella conical; spores elliptical 5-10x3-6 μ , colorless, but in mass greenish yellow.

On dung, on ground in greenhouse, not uncommon. Pl. XIV, Fig. 5, a.

[*Mortierella* has a distinct fertile mycelium from which the sporangiophores arise singly or in groups, the bases being enveloped in a mass of short branches. The sporangia are many-spored and have no columella. The zygospores are covered with a dense mass of hyphae, which branch off from the suspensor-cells and the branches from which the latter arise. *M. polycephala* Coemans, distinguished among other things by its branched sporangiophores, grows on dung and on decaying pore fungi. It has been reported from the United States, and should be found here.]

SUB-FAM.—**Chaetocladiaceae**.—Asexual reproduction by conidia which are borne singly (i. e., not in chains) in groups on the swollen middle portion of branches of the conidiophores, the ends of which are sterile.

Through the *Thamnidieae*, one of the tribes of the *Mucoreae*, not represented in our flora, this group is connected with the *Eumucoreae*. The gradations shown by other forms and produced by cultivation make it reasonably certain that the conidia are to be regarded as reduced one-celled sporangia.

5. CHAETOCLADIUM FRESSENIUS Beitrage 97. 1853.

Parasitic upon other *Mucoraceae*, mycelium thin, colorless, forming clusters of short, thick haustoria at the point of attachment with the hyphae of the host; sporangiophores creeping, verticillately branched, ending in a long, sterile, pointed tip, the branches short with sterile tips, bearing on the swollen portion large numbers of single conidia.

Etymology: Greek *χαίτη*, hair, and *κλαδίον*, branch.

Chaetocladium brefeldii VAN TIEGH. & LEMON. Ann. Sc. Nat. Bot. 5, XVII., 342. 1873.

Characters of the genus; conidia globose or globose-elliptical, smooth, colorless, 2-5 μ .

Parasitic on *Mucor mucedo* and *Ascophora mucedo*. I have found it but once—at Lincoln in 1888 on an onion with *Ascophora mucedo*. Pl. XIV., Fig. 6, a.

[*C. jonesii* (BERK. & BR.) FRES., distinguished by its larger conidia with finely echinate exospore, blue in mass, grows on dung with other *Mucoraceae*. It has been reported from North America, and ought to be found here.]

Family.—ENTOMOPHTHORACEAE.*

“Mycelium mostly parasitic on living animals (insects), more rarely on plants, or saprophytic, richly branched, often falling apart in bits, at first unicellular. Asexual reproduction by conidia, which are produced singly on the ends of unbranched threads growing up out of the substratum, and at maturity are abscised; without special, stalked conidiophores. Zygosporangia on the mycelium.” (Fischer, in Rabh. Krypt. Flor.)

A small group, chiefly parasitic on insects, containing 5 genera and about 40 species. The resting spores, which are either zygosporangia or azygosporangia, as in the *Mucoraceae*, point to some relationship with that group. The two groups are for that reason usually placed near together by systematic writers. However, they do not seem to have any immediate connection.

1. ENTOMOPHTHORA FRESenius Bot. Zeit. XIV., 883. 1856.

Empusa COHN Nov. Act. Acad. Caesar. Leopold. Carol. XXV., I., 317. 1855. (Ex Winter), not *Empusa* LINDLEY, 1824=*Liparis* RICH.

Parasitic on insects; the characters of the family.

Etymology: Greek *εντομος*, insect, and *φθορη*, death.

Empusa COHN, the name adopted by Berlese & DeToni in the *Sylloge Fungorum* and by Thaxter in his *Entomophthoraceae* of the United States, must be rejected on account of the older *Empusa* LINDLEY, one of the orchids, in accordance with the Rochester Rules.

The name *Entomophthora* was formerly restricted to the conidial stage of these fungi, the resting spore stage being placed in a genus *Tarichium*.

Entomophthora muscae (FRIES.) FRES. l. c.

(?) *Sporodonema muscae* FR. Syst. Mycol. III., 435. 1829.

Empusa muscae COHN l. c.

Conidia bell-shaped or nearly spherical, with a broad subtruncate base and sharply pointed apex; 18–25x20–30 μ ; containing usually a single large oil globule, and surrounded after discharge with a mass of protoplasm. Conidiophores simple, broad and stout, tapering gradually to a narrow base; emerging in white rings between the segments of the host, without coalescing over its body. Secondary conidia like the primary, or more commonly subovoid, small, rounded at the apex and formed by direct budding from the primary form. Resting spores, azygosporangia, produced laterally or terminally from hyphae within the host; spherical, colorless, 30–50 μ in diameter. (Winter.) Host attached to substratum by proboscis.—(Thaxter.)

On house flies.—*Musca domestica*. Very common in the winter, when the flies affected may be found attached to the walls and ceiling indoors. Pl. XV., Fig. 2, a.

The resting spores, described by Winter in Rabh. Krypt. Flor. v. Deutschl., etc., have not been observed in this country.

*By Roscoe Pound.

Entomophthora grylli FRES. l. c.

E. calopteni BESSEY Am. Nat. XVII, 1280 and 1286. 1883.

Resting spores spherical, colorless, 30-45 μ .

On grasshoppers.—*Melanoplus differentialis*, *M. bivittatus*, and *M. femurrubrum*; very common in autumn. Pl. XV., Fig. 3, a, b.

The affected grasshoppers climb to the tops of weeds and die there attached to the stem. They are readily known by their tight and rigid grasp, due to contraction of the limbs.

According to Thaxter, *E. calopteni* is not distinct from the European *E. grylli* which is found on crickets.

Order 5.—SIPHONEAE.—Typically unicellular, chlorophyll-green or colorless, filiform (sometimes branched), saccate, or foliaceous, one to plurinucleate; chloroplasts disk-shaped, parietal; propagation by cell-division; or by zoogonidia; reproduction either heterogametic or isogametic.

SYNOPSIS.

FAM.—**Vaucheriaceae.**—Terrestrial or aquatic; thallus filiform, elongated, often branched; reproduction by antherids and oogones borne laterally on the filament; propagation by zoospores, zoospores large, arising from an apical inflation of the filament.

FAM.—**Hydrogastraceae.**—Terrestrial; thallus a globose or pyriform cell attached to the earth by branching, hyaline rhizoids; reproduction by copulation of zoogonidia.

FAM.—**Saprolegniaceae.**—Aquatic fungi, mostly saprophytes, rarely parasites; asexual reproduction chiefly by biciliate zoospores arising in zoosporangia; sexual reproduction by oogones and antherids generally borne upon short lateral branches.

FAM.—**Peronosporaceae.**—Fungi growing parasitically in the subdermal tissues of flowering plants, piercing the cell walls by means of haustoria; asexual reproduction by conidia; sexual reproduction by oogones and antherids borne laterally upon mycelial filaments.

Family.—VAUCHERIACEAE.

Apparently unicellular, filiform, terrestrial or aquatic, chlorophyll-green; forming elongated, tubular filaments, simple or pseudo-dichotomously branched, often attached to substratum at base by hyaline rhizoids; propagation by motile or non-motile cells formed in apical inflations; reproduction by antherids and oogones, usually on the same filament; oogones lateral, sessile, or borne on a more or less elongated, simple or branched pedicel, cytoplasm converted into a large oospore; antherids lateral, sessile, or cut off by a septum from the upper portion of a lateral branch, producing antherozoids internally, which, being emitted, penetrate the apex of the oogone; antherozoids oblong, furnished with two unequal cilia.

1. VAUCHERIA DC. in Vauch. Hist. Confer. 25. 1803.

The characters of the family; chloroplasts minute, numerous, parietal; nuclei globose, numerous, small; protoplasm containing numerous oil globules.

Etyymology: dedicated to the phycologist Vaucher.

No. 16

EXTRAIT

DU

BULLETIN SCIENTIFIQUE

DE LA FRANCE

ET DE LA BELGIQUE,

PUBLIÉ PAR

ALFRED GIARD,

*Chargé de cours à la Sorbonne (Faculté des Sciences),
Maître de Conférences à l'École Normale Supérieure.*

DE INSECTORUM MORBIS
QUI FUNGIS PARASITIS EFFICIUNTUR,

PAR

J. KRASSILTSCHIK.

Analyse critique par A. GIARD.

PARIS,

OCTAVE DOIN, Éditeur,

8, Place de l'Odéon, 8.

1889

BULLETIN SCIENTIFIQUE DE LA FRANCE ET DE LA BELGIQUE, (REVUE MENSUELLE)

III^e SÉRIE, 2^e ANNÉE.

I-IV, JANVIER-AVRIL.

SOMMAIRE :

	Pages.
A. GIARD. — Leçon d'Ouverture du Cours d'Évolution des Êtres organisés.	1
P. PELSENER. — Sur la classification phylogénétique des Pélécy-podes [Communication préliminaire] (4 fig. dans le texte)	27
A. GIARD. — Sur la transformation de <i>Pulicaria dysenterica</i> en plante dioïque (3 fig. dans le texte et Planche I)	53
N. SOROKIN. — Un nouveau parasite de la Chenille de la Betterave, <i>Sorosporella agrotidis</i> (fig. dans le texte) [avec Note de A. GIARD]	76
A. GIARD. — Première liste de Galles du Nord de la France par M. Foc- KEU [Analyse critique]	84
F. BLOCHMANN. — Les globules polaires chez les œufs d'insectes se dé- veloppant sans fécondation.	93
A. GIARD. — Sur la signification des globules polaires.	95
E. SAUVAGE. — Contribution à la connaissance de la faune du Pas-de- Calais et des parties voisines de la Mer du Nord et de la Manche.	104
A. GIARD. — De insectorum morbis qui fungis parasitis efficiuntur, par J. KRASSILTSCHICK [Analyse critique]	120
P. GARNAULT. — La Castration parasitaire chez <i>Helix aspersa</i> (Planche II).	137
A. GIARD. — Paléontologie fantaisiste : un reptile en bois !	143
O. BUTSCHLI. — Devons-nous admettre un accroissement du plasma par intussusception ?	145
A. MAGNIN et A. GIARD. — Notes sur la castration parasitaire de <i>Melan- dryum vespertinum</i> (<i>Lychnis dioica</i>).	150
E. METSCHNIKOFF. — Sur le rôle phagocytaire des cellules géantes du tubercule [Analyse]	161
A. GIARD. — Fragments biologiques (suite) : XV. Sur l'orientation des Bo- pyres relativement à leurs hôtes. — XVI. Sur l'habitat de Phreoryctes Menkeanus. — XVII. Sur les espèces de <i>Sepiola</i> des côtes de France. — XVIII. Une station de <i>Mutilla Europaea</i> L. dans le nord de la France. — XIX. Sur le Phragmidiotrix incrustans n. sp. — XX. Sur la présence du <i>Thon</i> <i>Thynnus vulgaris</i> L. dans la mer du Nord. — XXI. Les Odonates du département du Nord	167
C. DE BOSSCHÈRE. — L'exposition internationale de géographie bota- nique, commerciale et industrielle à Anvers (1899).	185
A. GIARD. — BIBLIOGRAPHIE : VAYSSIÈRE : Atlas d'Anatomie comparée des Invertébrés. — CH. BRONGNIART. Sur le <i>Pleuracanthus Gaudryi</i>	192

PRIX DE L'ABONNEMENT :

Pour la France et l'Étranger, UN AN, 15 FRANCS.

Les abonnements partent du 1^{er} Janvier de chaque année.

PRIX DU NUMÉRO, 1 FRANC 25.

Adresser tout ce qui concerne la Rédaction au Professeur
A. GIARD, 14, rue Stanislas, PARIS.

DE INSECTORUM MORBIS QUI FUNGIS PARASITIS EFFICIUNTUR,

PAR

J. KRASSILSTSCHIK (1).

Analyse critique par A. GIARD.

Dans le précédent fascicule du *Bulletin* (2), nous avons reproduit une note très intéressante d'un naturaliste russe, J. KRASSILSTSCHIK, sur l'application des champignons parasites à la destruction des insectes nuisibles. Cette note, écrite par l'auteur lui-même à la demande de notre ami le Prof. MARION, est le résumé partiel d'un mémoire plus général publié en 1886, dans les Comptes-rendus de la Société des naturalistes d'Odessa. Elle donne surtout une idée du côté pratique de la question, et dans cette direction, KRASSILSTSCHIK paraît avoir obtenu des résultats très remarquables. Le terrain était d'ailleurs admirablement préparé par les travaux antérieurs de CIENKOWSKY et de METSCHNIKOFF. La culture artificielle de la forme conidienne (*Isaria*) de certains Pyrénomycètes du groupe des Nectriées paraît se faire avec autant de facilité que celle des *levures* ou des Schizomycètes. C'est là un succès bien encourageant et qui doit nous engager à poursuivre de nouvelles recherches sur le groupe des Entomophthorées jusqu'à présent rebelle à toute tentative de culture dans un milieu non vivant (3).

Mais alors même que cette difficulté de cultiver les Entomophthorées sur un substratum artificiel se montrerait insurmontable, ce ne serait pas une raison pour abandonner toute idée d'utilisation intensive de ces précieux auxiliaires.

(1) Publié en russe dans les *Zapiski* de la Société des Naturalistes de la Nouvelle-Russie, Odessa, t. XI, 1886, 1^{er} fascicule, pp. 75-172.

(2) Voir *Bulletin scientifique*, t. XIX, 1888, p. 461.

(3) Les expériences de EIDAM sur *Basidiolobus ranarum* et les miennes sur *Entom. calliphoræ* prouvent que l'on pourra peut-être vaincre cette résistance.

Les recherches de THAXTER ont montré en effet que certaines *Entomophthora* sont moins exclusives dans le choix de leur hôte que je ne l'avais supposé autrefois. Telles sont par exemple deux des espèces les plus utiles : *Entomophthora grylli* FRES. et *E. sphærosperma* FRES. Ces deux espèces infestent surtout en Europe, la première les Acridiens, la deuxième les chenilles de la Piéride du chou.

Mais elles peuvent, d'après THAXTER, s'inoculer soit naturellement soit artificiellement à un certain nombre d'insectes d'ordres différents. Ne pourrait-on, si les expériences de THAXTER se vérifient, cultiver par exemple l'*E. grylli* sur les chenilles d'*Arctia* si communes et si faciles à élever, puis se servir des spores ainsi produites pour infester les Acridiens et arrêter leurs ravages. Cette manière de procéder serait comparable à la pratique depuis longtemps en usage pour le champignon de la vaccine que nous cultivons sur les vaches pour le transporter ensuite sur l'homme. La culture des *Entomophthorées* sur des milieux vivants convenablement choisis, pour rendre facile un ensemencement ultérieur sur des êtres différents ouvrirait ce me semble un champ nouveau aux investigateurs et permettrait d'utiliser d'une façon régulière ces terribles ennemis des insectes nuisibles. Dès 1879, j'ai attiré sur cette question l'attention des agriculteurs, mais j'avais alors recommandé surtout l'ensemencement *direct* des spores durables (*Tarichium*) procédé auquel l'expérience m'a forcé de renoncer. (Voir *Bulletin scientifique*. t. XI, p. 361 et suiv.)

Comme dans les questions de ce genre, plus encore que partout ailleurs, la pratique doit être guidée par la théorie, nous avons lu avec le plus grand soin le mémoire russe de KRASSILTSCHIK et nous nous proposons d'analyser brièvement ce travail en indiquant les points que nous pensons inexacts ou tout au moins contestables.

D'après KRASSILTSCHIK, J. V. LIEBIG aurait le premier dès 1820 attribué à un *contagium vivum* certaines maladies, telle que la peste, la petite vérole, le syphilis, le typhus, etc. (1).

L'importance attachée à quelques passages des *Lettres sur la chimie* est, je crois, bien exagérée. Comme l'a très justement fait observer VIRCHOW (2), la théorie du *contagium animatum* se perd dans la nuit du moyen-âge.

(1) LIEBIG, *Chemische Briefe*. Brief 21.

(2) VIRCHOW, La liberté de la science dans l'état moderne, *Revue scientifique*, t. 20, 1877, p. 538.

Nous avons reçu ce mot de nos ancêtres : il surgit déjà nettement au XVI^e siècle. Mais c'est seulement au XIX^e siècle qu'on a commencé à trouver effectivement morceau par morceau des *contagia animata*, et c'est dans la seconde dizaine de ce siècle qu'on a peu à peu découvert des preuves de plus en plus positives à l'appui de cette explication des maladies infectieuses.

Le naturaliste qui a certainement contribué le plus à répandre la notion de l'étiologie parasitaire d'un grand nombre de maladies est F.-V. RASPAIL.

A une époque où les recherches microscopiques n'étaient pas en honneur comme elles le sont aujourd'hui, RASPAIL eut le mérite de découvrir la cause réelle d'un certain nombre de maladies dues à des contagés vivants et de recommander deux des meilleurs parasitocides qu'on connut alors, le camphre et le goudron.

Les belles recherches de C.-J. DAVAINÉ méritent aussi d'être citées d'une façon spéciale.

En ce qui concerne plus spécialement les maladies des insectes, les naturalistes commencèrent à s'en préoccuper lorsque la muscardine vint menacer l'industrie de la sériculture. ROBIN nous a fourni un excellent résumé des découvertes successives, auxquelles a conduit l'étude de ce fléau (1).

FOSCARINI montra le premier (1819) que la muscardine se communique aux vers à soie, par contact et inoculation.

CONFLIGLIACCHI et BRUGNATELLI annoncèrent ensuite (1820) que ce qu'on appelait l'efflorescence muscardinique était véritablement une production de la nature des moisissures. En 1829, BONAFOUS, enfin converti à la théorie de la contagion qu'il avait repoussée d'abord, reconnaît que la muscardine se transmet des vers à soie malades aux vers à soie sains, et que de plus elle peut être communiquée artificiellement à des chenilles de papillons différents (*Cucullia verbasci*) mises en contact avec des vers morts de cette affection. Bientôt (de 1836 à 1839) grâce aux recherches de MONTAGNE, TURPIN, AUDOUIN et BONAFOUS, les exemples de transmission du parasite à des insectes de genres et même d'ordres différents se multiplièrent, en même temps que les travaux de BALSAMO et de MONTAGNE permettaient de définir plus complètement la nature de la muscardine.

Ce champignon fut appelé d'abord par BALSAMO *Botrytis para-*

(1) ROBIN, Végétaux parasites, etc., 1853, pp. 592 et suiv.

doxa, puis *Botrytis Bassiana* en l'honneur de Bassi, qui avait engagé BALSAMO à entreprendre l'étude botanique de l'*efflorescence* des vers à soie.

Pendant toute cette première période, on ne s'inquiéta des champignons parasites des insectes que pour chercher le moyen de les détruire et de faire disparaître les épidémies de l'insecte utile (ver à soie). C'est dans le même esprit qu'on étudia plus tard la *pébrine* (due à une Psorospermie) et la *flacherie* (due à un Schizomycète).

Mais depuis 1878-1879 la préoccupation des naturalistes s'est tournée vers un autre but. On a songé que, pour quelques insectes utiles, il y en avait des milliers de nuisibles, et qu'il serait peut être possible d'employer contre ces derniers les redoutables cryptogames parasites qu'on avait appris à connaître.

E. METSCHNIKOFF est certainement le premier naturaliste qui soit entré dans cette voie féconde : il écrivit un mémoire très important, sur les *Maladies des hannetons du blé*, publié en langue russe à Odessa en janvier 1879.

Dans ce mémoire, le savant zoologiste fit connaître une nouvelle sorte de muscardine trouvée par lui sur la larve de l'*Anisoplia austriaca* HERBST., qui ravageait les champs de blé du Midi de la Russie. En octobre 1878 METSCHNIKOFF avait découvert dans les larves d'*Anisoplia* une affection bactérienne et une *muscardine verte* provoquée par un champignon du genre *Isaria* qu'il nomma *Isaria destructor*.

L'analogie de cette muscardine avec celle du ver à soie permettait de supposer une propagation facile.

Aussi METSCHNIKOFF n'hésita-t-il pas à proposer d'ensemencer les spores de l'*Isaria destructor* dans les champs attaqués par l'*Anisoplia*. En avril 1879, CIENKOWSKY se ralliait à cette proposition et affirmait de plus la possibilité d'une culture de l'*Isaria* en dehors de l'organisme vivant. Cette culture fut obtenue d'abord dans la trempe de bière de maïs en automne 1879 par METSCHNIKOFF aidé des conseils de M. WERIGO, chimiste à Odessa.

Les belles recherches de CIENKOWSKY et de METSCHNIKOFF ne furent connues en France, que dans le courant de 1880. Grâce à la présence à Marseille du Professeur KOWALEVSKY, notre ami MARION put donner un excellent résumé des travaux des savants Russes dans son *Rapport sur l'application du sulfure de carbone aux vignes phylloxérées* (4^e année, Paris, 1880, p. 24-28).

De notre côté et d'une façon tout à fait indépendante, nous nous

occupions depuis 1878 de recherches de même nature (1) mais, au lieu de recourir comme METSCHNIKOFF et CIENKOWSKY aux entomogènes (2) du groupe des *Ascomycètes*, nous avons porté toute notre attention sur la famille des Entomophthorées.

Les épidémies occasionnées chez les insectes par ces Basidiomycètes inférieurs sont tout aussi meurtrières que celles produites par les Isariées ; peut être même amènent-elles plus rapidement la mort de l'insecte infesté. Mais le maniement pratique de ces champignons présente des difficultés plus grandes. Je dois dire d'ailleurs que j'ai été peu aidé dans mes tentatives, soit par les botanistes, soit par les entomologistes, soit par les agriculteurs.

Toutefois, ainsi que je l'ai dit ci-dessus, les obstacles que j'ai rencontrés ne sont pas insurmontables, et je fais un nouvel appel au concours de tous les intéressés pour me faciliter la continuation de ces essais.

Dans le mémoire qu'il a publié en 1886, KRASSILSTSCHIK traite d'une manière générale de tous les champignons entomogènes : il est visible toutefois que, suivant l'exemple de METSCHNIKOFF, il a principalement étudié les *Isaria* et ne connaît que fort imparfaitement les Entomophthorées.

La partie historique de ce travail renferme de nombreuses lacunes et cette insuffisance bibliographique enlève une partie de leur valeur aux conclusions tirées de la statistique des champignons insecticides.

Du reste, ces considérations statistiques devront être entièrement remaniées par suite des documents nouveaux que nous a apportés la belle monographie des Entomophthorées d'Amérique par R. THAXTER, publiée postérieurement au mémoire de KRASSILSTSCHIK.

Mais il y a dans le travail du savant russe un défaut plus grave que les lacunes bibliographiques : c'est l'insuffisance des études préalables nécessaires pour mener à bien l'étude si complexe des cryptogames parasites des insectes.

Des recherches de cette nature, pour être irréprochables au point de vue scientifique, exigent du naturaliste qui les entreprend, non

(1) Voir *Bulletin scientifique*, t. XI, 1879, pp. 353 et suiv.

(2) Pour me conformer à l'usage créé par les botanistes, j'emploierai quelquefois le mot d'*entomogène*, mais il faut remarquer que ce mot est très mal fait et qu'il vaudrait mieux le remplacer par celui d'*entomophyte* ou d'*insecticide*.

seulement une grande habileté d'expérimentateur, mais encore une double compétence en mycologie et en entomologie.

Au point de vue de cette double compétence, le travail de KRASSILSTSCHIK est loin d'être à l'abri de toute critique. Les entomologistes s'étonneront en le lisant de voir classer parmi les Orthoptères les fourmis (*Atta cephalotus* FAB. et *Formica* sp.) dans le tableau des insectes infestés (pp. 114-115). Ils s'étonneront aussi de voir citer comme espèce spéciale la *Vespa crinita* FELTON (p. 130) (1). D'autre part, les botanistes admettront difficilement que les *Elaphomyces* qui sont des Tubéracés, c'est-à-dire des *Ascomycètes*, soient rangés au nombre des *Gastromycètes* qui représentent un type analogue par convergence dans le groupe des *Basidiomycètes*.

Quoiqu'il en soit et malgré ces lapsus regrettables, il est juste de reconnaître que le travail de KRASSILSTSCHIK, en dehors de son mérite pratique incontestable, nous fait connaître un certain nombre de faits scientifiques nouveaux et intéressants.

La découverte par VEDHALME de l'*Entomophthora grylli* FRES. sur le *Stenobothrus nigromaculatus* aux environs de Kazan ne fait qu'ajouter un nom nouveau à la liste des insectes qui hébergent ce parasite, liste considérablement augmentée, peut-être même trop augmentée, depuis par R. THAXTER.

D'un intérêt très grand est au contraire l'observation d'une Entomophthorée parasite de *Cecidomya destructor* SAY que KRASSILSTSCHIK et VEDHALME ont rencontrée assez largement répandue à Kourskaia et en Bessarabie.

Cecidomya destructor est un insecte des plus nuisibles et il serait très important de pouvoir le combattre d'une manière efficace. KRASSILSTSCHIK a rencontré l'Entomophthorée de la Cecidomye à l'état de conidies et à l'état de *Tarichium*, il en fait à tort deux formes distinctes. Cela tient, comme nous allons le voir, à la con-

(1) Je signale seulement en passant une méprise linguistique assez singulière. KRASSILSTSCHIK donne comme noms de genres *Amphiorrycis*, *Aucericis* (p. 130). Ces noms sont les génitifs de *Amphiorryx* et *Aucerix*. D'ailleurs, l'ignorance des langues anciennes est un trait caractéristique de presque tous les naturalistes russes. Cela tient sans doute à la réforme de l'enseignement qui, pendant un certain temps, a fait supprimer dans ce pays d'une façon complète l'étude des langues classiques. L'exemple est à citer à ceux de nos compatriotes qui voudraient nous voir entrer dans la même voie. Au point de vue scientifique international, une diagnose en mauvais latin vaut mieux qu'une description en français ou en russe d'une parfaite élégance.

naissance imparfaite qu'avait ce naturaliste de l'évolution des Entomophthorées. La description de cette *Entomophthora* est d'ailleurs tout à fait insuffisante et pour ainsi dire nulle.

KRASSILTSCHIK a découvert également une *Entomophthora* sur les chenilles de l'*Agrotis segetum* noctuelle qui, dans les environs d'Odessa, attaque surtout le seigle. Mais constamment les chenilles infestées lui ont présenté la forme conidiale ou *Empusa* de l'Entomophthorée et jamais la forme *Tarichium*. « Ce fait est très intéressant, observe-t-il, car COHN a trouvé sur les mêmes chenilles un *Tarichium* sans jamais rencontrer de conidies et, s'il s'agissait d'un même champignon, on ne voit pas pourquoi il se produirait ici la forme *Tarichium* et ailleurs la forme *Entomophthora* (*Empusa*), le substratum restant le même. »

L'observation est très intéressante à coup sûr, mais on ne peut, je pense, en tirer la conclusion que cherche à déduire KRASSILTSCHIK. Ce n'est pas le seul exemple que nous ayons d'un végétal qui, dans une zone géographique déterminée, perd plus ou moins complètement un de ses modes de reproduction. Une Hépatique très commune dans le Nord de la France, la *Lunularia vulgaris*, ne se reproduit jamais que par progagules dans cette région, tandis qu'en Italie et dans les pays méditerranéens, elle développe un appareil de reproduction sexuée déjà signalé par MICHELI. Sans sortir du groupe des champignons entomogènes, le *Cordyceps militaris* nous présente un fait du même genre. Dans le midi de la France, d'après ROUMEGUÈRE, l'état conidiophore de ce champignon (*Isaria farinosa*) est beaucoup plus commun que la forme thécigère. C'est l'inverse, au contraire, dans les contrées du Nord : KARSTEN (*Mycol. Finland.*) déclare n'avoir jamais rencontré l'*Isaria* dans la Finlande (1).

Dans la France septentrionale, aux environs de Valenciennes comme aux environs de Fécamp, je n'ai jamais rencontré que la forme *Tarichium* de l'*Entomophthora* de l'*Agrotis*. Cependant ayant placé un jour quelques chenilles momifiées dans une chambre humide maintenue à une température assez élevée, je les ai vues se couvrir d'un duvet blanchâtre analogue à l'état conidial ordinaire des Entomophthorées : malheureusement l'expérience fut interrom-

(1) ROUMEGUÈRE. Les sphériacées entomogènes (*Revue mycolog.*, t. VI, 1884, p. 149).

pue par un accident, sans que j'aie pu constater la formation des conidies.

En tout cas, l'observation de KRASSILTSCHIK me paraît insuffisante pour prouver que l'*Empusa* trouvée à Odessa sur les chenilles de l'*Agrotis* diffère spécifiquement du *Tarichium megaspermum* COHN.

Par suite, je ne vois pas non plus de raisons sérieuses pour considérer comme espèces distinctes l'*Entomophthora* et le *Tarichium* que VEDHALME et KRASSILTSCHIK ont rencontrés à Kourskaia et en Bessarabie sur des individus différents (larves ou nymphes) de *Cecidomya destructor* SAY.

La même observation s'applique évidemment à l'*Entomophthora* et au *Tarichium* que KRASSILTSCHIK et VEDHALME ont observé en Bessarabie sur le *Thrips solanacearum* WIDH. (1) (larves et imago).

Quant au champignon décrit par KRASSILTSCHIK, sous le nom de *muscardine rouge* ou de *Tarichium uvella* et trouvé sur la larve de *Cleonus punctiventris* GERM. dans les environs de Kiew, il m'est impossible d'y reconnaître un *Tarichium*. Déjà R. THAXTER a proposé de le rapprocher du genre *Massospora* PECK, mais il me paraît encore plus voisin du parasite de l'*Agrotis segetum* décrit récemment par SOROKIN, sous le nom de *Sorospora agrotidis* (2).

KRASSILTSCHIK ne fait que citer sans le décrire un champignon indéterminé trouvé par VEDHALME sur le *Schizonentra radicicola* aux environs de Sarato-Galbina (district de Bender). Ce champignon attaquait les pucerons à l'état aptère. C'est là une espèce qu'il serait bien désirable de voir étudier de plus près en raison des services qu'elle pourrait rendre peut-être contre le *Phylloxera*.

C'est surtout, comme nous l'avons dit, sur les Entomogènes Ascomycètes que KRASSILTSCHIK a porté son attention.

La découverte de *Botrytis Bassiana* BALS sur deux hôtes nouveaux (*Musca domestica* et *Athalia berberidis*?), découverte faite par KOWALEVSKY aux environs d'Odessa méritait d'être signalée. On sait que METSCHNIKOFF avait déjà observé la muscardine blanche sur l'*Anisoplia austriaca* dans le district de Tcherkassy (G^t de Kiew).

(1) Voir t. IV des Travaux de la Société entomologique d'Odessa, 1884, p. 97 (en russe).

(2) Voir ce Bulletin, tome XX, p. 76.

KRASSILSTSCHIK a retrouvé le même parasite dans le gouvernement de Kiew sur le *Cleonus punctiventris* GERM. Les insectes, si redoutés des cultivateurs de la betterave de la Russie méridionale, peuvent donc être combattus efficacement par trois champignons : la muscardine verte, *Isaria destructor* METSCH. ; la muscardine rouge, *Soros-porella uvella* KRASSIL. ; et la muscardine blanche *Botrytis Bassiana* BALS.

VEDHALME et KRASSILSTSCHIK ont découvert aussi une espèce nouvelle d'*Isaria* sur un *Lixus* indéterminé (larve et nymphe) qui vit sur l'*Onopordon acanthium* L., aux environs d'Odessa, *Isaria* qu'on pourrait utiliser sans doute pour se débarrasser des divers Curculionides qui attaquent les Carduacées.

Enfin l'*Isaria destructor*, si bien étudiée par METSCHNIKOFF sur le *Cleonus punctiventris*, a été retrouvée par VEDHALME, près d'Odessa, sur les larves d'un *Agriotes* et sur les chenilles d'*Heliothis dipsaceus* L.

Mais le champignon que KRASSILSTSCHIK a étudié le plus complètement est celui qu'il a rencontré dans la Bessarabie méridionale sur les œufs du Criquet voyageur (*Pachytylus migratorius* L.) et qui a été observé également par A. KOWALEVSKY et J. VEDHALME. C'est une forme conidiophore que KRASSILSTSCHIK croit pouvoir rapporter à l'*Isaria* de *Cordyceps* (*Torrubia*) *ophioglossoïdes* EHR. TULASNE (1) (Selecta fungorum Carpol. III, p. 20-22, Pl. II, fig. 9). C'est aussi, paraît-il, l'avis émis par les Professeurs CIENKOWSKY et REINHARDT qui ont vu des préparations de ce champignon. L'aspect extérieur rappelle assez bien le *Penicillium glaucum*, mais la forme des spores qui sont cylindriques ne permet pas la confusion.

On sait que la plupart des *Cordyceps* vivent en parasite sur les insectes. Un très petit nombre d'espèces seulement (*Cordyceps ophioglossoïdes* EHR. TUL. et le *Cordyceps capitata* TUL.) se rencontrent exclusivement sur les champignons du genre *Elaphomyces*.

(1) Ce Champignon fut décrit par EHRHART, qui, après LINNÉ et VAILLANT, signala également *Sphaeria* (*Torrubia*) *militaris*. L'abréviation EHR. trompa ROBIN qui la traduisit par EHRENBURG (Végétaux parasites, p. 647, *Sph. militaris* EHRENBURG). La même erreur a été commise par un cryptogamiste distingué, M. LEBRETON, dont le *Bulletin de la Soc. des Amis des Sciences de Rouen* a publié une Note sur le *Torrubia ophioglossoides* EHRENBURG (année 1876, 2^e semestre, p. 199).

Si l'*Isaria* des œufs de Criquets est bien l'*Isaria ophioglossoïdes*, nous nous trouvons en présence d'un fait très curieux dont KRASSILTSCHIK n'a pas saisi toute l'importance.

« Quoique, dit-il (p. 104), *Cordyceps ophioglossoïdes*, dont notre champignon se rapproche le plus, soit du nombre des espèces de ce genre qui vivent en parasite sur les *Elaphomyces* et jamais sur les insectes, il ne faudrait pas, sous prétexte qu'il n'y a rien des éléments de l'*Elaphomyces* dans les œufs de Criquets, admettre ici un double parasitisme, car, d'une part, il est bien invraisemblable que les *Elaphomyces*, ces *gastromycètes* (1), puissent végéter sur des œufs de criquets, et d'un autre côté, il serait encore plus invraisemblable qu'un *Elaphomyces* qui est ordinairement cent fois plus grand qu'un œuf de criquet, puisse jouer le rôle de ce dernier. Par conséquent, il nous paraît certain que notre champignon vit immédiatement et directement aux dépens des œufs de criquets. »

Cette conclusion n'explique rien. La question est plus complexe et d'un intérêt plus général. Les œufs de criquets fournissent évidemment au champignon une nourriture insuffisante pour lui permettre de développer son appareil reproducteur le plus parfait, l'appareil thécigère. Mais comment se fait-il que dans certaines localités le *Cordyceps ophioglossoïdes* abandonne les œufs d'Acridiens pour végéter d'une manière plus complète sur les *Elaphomyces* ?

On sait que les *Elaphomyces* sont très sujets à être envahis par de nombreux parasites et, en particulier, par les larves de diptères du genre *Helomyza*.

Or les diptères de ce groupe sont fréquemment infestés à leur tour par des Sphaeriacées entomogènes. QUELET a décrit un *Cordyceps*, le *C. Forquignonii* QUEL. qui vit sur l'*Helomyza rufa* FAB. (2) et GADEAU DE KERVILLE a signalé le *Stilbum Kervillei* QUELET comme parasite très fréquent de *Leria cæsia* MEIG. aux environs de Rouen et d'Elbœuf. (3)

(1) Nous avons relevé ci-dessus cette erreur : les *Elaphomyces* sont des *Tuberacés* et non des *Gastromycètes*.

(2) QUELET. Quelques espèces critiques ou nouvelles de la flore mycologique de France (*Association française*. Congrès de Toulouse, 1887, p. 591, Pl. xxi, fig. 18).

(3) H. GADEAU DE KERVILLE. Note sur une nouvelle espèce de Champignon entomogène (*Stilbum Kervillei* QUEL.) (*Bulletin de la Soc. des Amis des Sciences nat. de Rouen*, 1884).

Il me paraît bien probable que les *Cordyceps*, parasites des *Elaphomyces*, ont vécu d'abord à l'état d'*Isaria* sur les larves de diptères qui attaquent ces Tuberacées, puis ont étendu leur mycelium sur l'*Elaphomyces* lui-même, et ont pu ainsi, grâce à cette nourriture plus large, développer leur appareil thécigère.

Il est même possible que les choses se soient passées ainsi autrefois dans l'évolution phylogénique, et qu'aujourd'hui les *Torrubia ophioglossoides* et *capitata* envahissent directement les *Elaphomyces*. Cependant les observations de KRASSILSTSCHIK montrent que, pour la première de ces espèces, le parasitisme sur les insectes s'observe encore en certaines localités au moins, pour la forme *Isaria*.

La suppression d'un hôte intermédiaire introduisant un parasite sur un substratum tout différent de celui qu'exploitent ses congénères, n'est pas un fait absolument rare, et il me serait facile d'en citer bien des exemples.

J'ai signalé ailleurs, avec détail, le cas si curieux des Rhizocéphales considérés comme hôtes intermédiaires entre les Décapodes et les Bopyriens.

Un exemple non moins intéressant, et peut-être plus facile à saisir, est celui que nous offrent certains hyménoptères, de la famille des Chalcidites. Les *Eurytoma*, auxquels je fais allusion en ce moment, sont généralement parasites d'autres insectes, et certains d'entre-eux vivent notamment aux dépens de Diptères gallicoles, tels que les Cécidomyes. Mais il en est qui forment directement des galles sur les végétaux. L'*Eurytoma longipennis*, par exemple, produit sur le *Psamma arenaria* une galle assez voisine de celles que les Cécidomyes occasionnent sur d'autres Graminées. L'*Eurytoma calleyae* étudié par RILEY, est considéré par les horticulteurs d'Amérique, et même de Belgique (où il s'est introduit récemment), comme un insecte nuisible aux Orchidées.

Il est bien évident que ces hyménoptères ont été autrefois des insectes utiles, mais ils ont fini par prendre la place du parasite, aux dépens duquel ils vivaient primitivement. Le même fait s'est produit, sans doute, chez les hyménoptères capricificateurs, d'après les observations de FRITZ MUELLER (Voir ce *Bulletin*, p. 88).

De pareils exemples me paraissent jeter une certaine lumière sur les rapports des *Isaria* et des *Elaphomyces*, et nous permettent

d'expliquer rationnellement des particularités qui semblaient jusqu'ici fort embarrassantes.

KRASSILTSCHIK a résumé son travail dans une série de tableaux donnant la liste (par ordres) des insectes infestés, les noms des champignons parasites, ceux des naturalistes qui les ont découverts, l'étendue des épidémies, les tentatives de culture artificielle et les indications bibliographiques relatives à chaque cas.

C'est là une entreprise fort louable, et qui rendra de grands services à ceux qui voudront étudier cette question si intéressante des champignons parasites des insectes.

Toutefois, comme nous l'avons déjà dit, ces tableaux sont très incomplets, et l'auteur a négligé en particulier la littérature mycologique française. Les travaux admirables des frères TULASNE, les notes de CORNU et les miennes sont seuls cités. Tout en étant très flatté de me trouver au petit nombre des élus, je ne puis m'empêcher de regretter que KRASSILTSCHIK ait laissé dans l'ombre une série d'observations intéressantes (1).

J'en rappelle quelques-unes au hasard :

1° *Polistes pallipes* LEP., parasité par *Torrubia (Cordyceps) sphecocephala* KL., Antilles (PEREZ);

2° *Vespa crabro* L., parasité par *Cordyceps Dittmari* QUEL., Jura (QUELET). — La forme conidiophore de *Cordyceps Dittmari* est très probablement l'*Isaria sphecocephala* DITTMAR (*Ceratonema crabronis* PERSON) trouvée également sur *Vespa crabro* et rattachée par SOROKIN, sans raison valable ce me semble, à *Cordyceps sphecocephala* KL.;

3° et 4° *Myrmica rufa* L. et *Formica nigra* L., parasitées par *Cordyceps myrmecophila* CESATI, Italie, Finlande, Angleterre, Amérique septentrionale, Ceylan, Sarawak;

5° *Melolontha vulgaris* L., parasité par *Cordyceps militaris* L., France, Aude (ROUMEGUÈRE);

6° *Rhizotrogus solstitialis* FAB., parasité par *Cordyceps entomorrhiza* (DICKS.), France : Super Bagnères (ROUMEGUÈRE et FOURCADE);

(1) Voir notamment ROUMEGUÈRE, Les Sphériacées entomogènes (*Revue mycologique*, VI, 1884, p. 148 et suiv.).

7° *Lucanus cervus* L., parasité par *Isaria eleutheratorum* NÉES.
France : Sàone-et-Loire (GROGNOT).

Cet *Isaria* est peut-être la forme conidiophore de *Cordyceps entomorphiza* ;

8° *Zabrus gilbus* FAB., parasité par *Cordyceps cinerea* TUL
France : Valmigère (Aude) (ROUMEGUÈRE) ;

9° *Acronycta aceris* L., parasité par la variété *prolifera ramosa*
ROUMEGUÈRE (Fungi Gallici exsiccati, N° 1361) de l'*Isaria* de *Tor-
rubia militaris*. Environs de Lyon (J. THIERRY) ;

10° *Sphinx atropos* L., parasité par *Cordyceps sphingum*.

L'état conidifère de cette espèce est l'*Isaria sphingum* SCHW.

11° *Leria caesia* MEIG., parasité par *Stilbum Kervillei* QUELET.
Sur les parois de la cavée de Dieppedalle, près Rouen, et grottes
calcaires d'Orival, près Elbeuf (GADEAU DE KERVILLE).

Je n'ai cité dans cette liste que des cas observés antérieurement
à la publication de KRASSILSTSCHIK.

Depuis 1876 le nombre des observations s'est encore fortement
accru, et un travail d'ensemble devient de jour en jour plus néces-
saire.

KRASSILSTSCHIK termine son mémoire par des considérations
générales sur les champignons parasites des insectes.

D'après lui, il convient de ne considérer comme réellement ento-
mogènes que les six genres suivants : *Cordyceps*, *Isaria*, *Stilbum*,
Botrytis, *Entomophthora* et *Tarichium*.

Nous avons vu que KRASSILSTSCHIK désigne sous le nom de
Tarichium, un champignon bien différent de ceux auxquels COHN a
donné ce nom ; et d'autre part nous avons montré ailleurs que les
Tarichium de COHN ne constituent pas un genre autonome. Ils
répondent à une sorte de fructifications (spores durables) des *Ento-
mophthora*, de même que l'ancien genre *Empusa* correspond à
l'appareil conidien des mêmes champignons.

Peut-être aussi KRASSILSTSCHIK accepte-t-il avec un peu trop de
timidité le lien qui rattache les *Isaria* aux *Cordyceps*. Mais toutes
réserves faites, et en laissant de côté la question de savoir s'il ne
conviendrait pas de subdiviser les grands genres *Cordyceps* et

Entomophthora, nous acceptons volontiers les vues de l'auteur, dont voici la traduction :

« Bien que dans ces derniers temps DE BARY penche à admettre l'opinion de TULASNE que l'*Isaria* est seulement la forme conidiale de *Cordyceps* et considère comme *très probable* que *Isaria farinosa* appartient au développement de *Cordyceps militaris*, il faut faire attention que le genre *Isaria* comprend un nombre énorme de formes et que l'union de ce genre avec les *Cordyceps* étant seulement *probable* (1) pour *une* espèce, il est utile pour le moment de conserver le genre *Isaria* et de désigner chaque forme par un nom spécial.

« Quant à *Botrytis Bassiana*, DE BARY a admis autrefois (ainsi que BREFELD) que ce n'est autre chose que la forme conidiale d'un Pyrénomycète (*Melanospora parasitica*); toutefois après de nombreuses cultures et recherches, il paraît avoir renoncé à cette idée (2).

« Le genre *Stilbum* a seulement un représentant parmi les champignons muscardiniens et ne fait que peu de victimes. C'est le *Stilbum Buqueti*; quoique ce dernier ne soit pas complètement bien connu et que pour cette raison on puisse se demander si tous les stades de son développement se passent sur les insectes, quoique, d'après BUQUET, le champignon se développe seulement sur des insectes morts, d'après la description et les bons dessins de ROBIN, on peut admettre avec ce dernier que le *Stilbum* se développe déjà du vivant de l'insecte et qu'il achève son évolution après la mort de celui-ci, en un mot qu'il se comporte comme une vraie muscardine (3).

« Quant à ce qui concerne *Tarichium*, j'ai déjà dit que ce genre doit suivre les *Entomophthora*. La difficulté des cultures artificielles des *Tarichium* et l'impossibilité de l'infestation artificielle des insectes avec les spores de ces champignons gênent très fort l'étude de ce groupe d'organismes parasitaires intéressant au plus haut degré.

« Au nombre des champignons qui figurent sur notre liste comme

1) A. DE BARY. Vergleichende Morphologie und Biologie der Pilze, etc. Leipzig 1884, p. 402.

(2) A. DE BARY, l. c., p. 274.

(3) C. ROBIN. Hist. nat. des végét. parasites, etc., 1853, p. 644.

se développant accidentellement sur les insectes avec le rôle d'entomogène se trouve un représentant du genre *Cladosporium*. Quoique beaucoup d'espèces du genre *Cladosporium* soient connues comme parasites des végétaux et que dans le monde des animaux les *Cladosporium* ne se montrent pas en qualité de parasites, il faut noter la présence sur notre liste d'un cas de ce genre, constaté par un mycologue aussi expérimenté et aussi instruit que le Prof. SALENSKY de Kazan. Il serait un peu étrange à première vue que l'existence d'un parasite aussi utile que *Cladosporium parasiticum* SOROK., qui vit sur le coleoptère *Polyphylla fullo*, put être dans le cas donné interprété comme si nous n'avions affaire qu'à un champignon saprophyte accidentel.

« Il est tout à fait inutile de parler des *Penicillium* considérés comme entomogènes. Le *Penicillium glaucum* qui dans l'opinion de LOHDE serait parasite d'un papillon, *Bryophila raptricula*, ne peut en aucune manière être compris au nombre de champignons entomogènes, et si, sur les cadavres de chrysalides réellement muscardinisées, il peut en effet se développer du *Penicillium glaucum*, c'est tout simplement comme phénomène consécutif et comme saprophyte sur des chrysalides déjà mortes. Il n'y a aucun doute que les chrysalides dont parle LOHDE, étaient déjà frappées par un autre champignon parasite, à la suite duquel le *Penicillium glaucum* s'était développé postérieurement.

« Tout aussi inexact, quoique beaucoup moins intelligible, est le cas cité par HAAGEN, à savoir que BURNS aurait détruit le coléoptère ennemi de la pomme de terre (*Doryphora*) avec de la levure de bière (!!?)

« Les champignons des levures n'ont rien de commun avec les muscardines et si on pouvait croire autrefois, comme l'a fait BAIL, que la mouche d'appartement était tuée par une levure, nous savons aujourd'hui après les belles recherches de BREFELD, que la prétendue levure de la mouche d'appartement, n'est autre chose qu'une *Entomophthora* à un certain stade de développement. Mais entre un *Saccharomyces* et une *Entomophthora* il y a certainement très loin. Peut-être si HAAGEN et BURNS ont constaté seulement le fait de la présence de levures dans le corps de *Leptinotarsa decemlineata*, peut-on conjecturer que vraisemblablement ils ont eu affaire à une *Entomophthora* qu'ils n'ont pas reconnue.

« Mais s'ils affirment qu'une espèce déterminée *la levure de bière* peut détruire le *Doryphora*, cela est évidemment une erreur (1).

« Il faut signaler encore le genre *Metarhizium* dont certains représentants figurent aussi sur notre liste. Ce genre nouveau a été établi par le professeur N. SOROKIN pour la muscardine verte, découverte par METSCHNIKOFF sur les larves d'*Anisoplia austriaca* et nommée par lui *Entomophthora anisopliae*. Comme les caractères de ce champignon ne concordent pas absolument avec ceux des *Entomophthora*, le Prof. SOROKIN a proposé de l'appeler *Metarhizium*. Mais comme plus tard METSCHNIKOFF a donné à la muscardine verte le nom d'*Isaria destructor*, comme mes longues observations sur ce champignon et un grand nombre de cultures pures m'ont prouvé que le champignon de la muscardine verte se rapproche tout à fait du genre *Isaria*, le nom de *Metarhizium* devenait inutile, d'autant plus que les autres représentants de ce genre étaient insuffisamment établis. Le champignon de la muscardine verte prend aussi l'aspect typique d'une plante arborescente d'*Isaria* sur les larves de *Cleonus punctiventris* lorsque celles-ci sont placés sur du sable humide. Dans les cultures artificielles, l'*Isaria destructor* est connu en outre sous la forme de *Coremium*.

« Quant aux noms ingénieux que LEBERT a donné à diverses formes de champignons entomogènes, tels que *Verticillium*, *Polistophthora*, *Acanthomyces*, etc., ils ne représentent nullement des formes nouvelles, mais ne sont que des synonymes de *Cordyceps* et *Isaria*.

« Il reste à voir dans combien de cas les muscardines parasites des insectes se sont développées dans des conditions artificielles. On constate que les expériences ont été faites sur 24 espèces différentes d'insectes (les expériences de BURNS non comptées, bien entendu) et ont toujours donné des résultats favorables. En outre, dans quatre autres cas la contagion, quoique observée aussi dans des conditions artificielles (non à ciel découvert), ne provenait pas de spores intentionnellement semées.

« Dans ces derniers cas le parasite se développait sur des insectes renfermés dans des bocaux ou des boîtes. Il est intéressant de noter que

(1) Chez les Daphnies seulement on a trouvé (METSCHNIKOFF) une maladie causée par un champignon du groupe des levures.

presque tous les cas d'infestation artificielle sont dus à des champignons des genres *Isaria* et *Botrytis*, c'est-à-dire des champignons dont les cultures artificielles réussissent à présent à merveille.

« Au contraire les infestations avec les genres *Cordyceps* et *Entomophthora* sont en nombre plus restreint. Jusqu'à présent les cultures artificielles de ces derniers genres n'ont aucunement réussi. Pour le genre *Cordyceps* nous n'avons qu'une expérience de DE BARY et pour les *Entomophthora* trois expériences de BREFFELD. Avec *Stilbum* aucune expérience n'a été tentée, mais à en juger par la structure de ce champignon si semblable à celle des *Isaria*, il est probable que la culture artificielle de *Stilbum* et les infestations réussiraient aussi bien qu'avec les *Isaria*. Pour ce qui concerne le genre *Tarichium* toutes les expériences de contagion artificielle ont complètement échoué et par conséquent toute culture artificielle s'est montrée également impossible.

« Si des recherches ultérieures confirmaient le cycle de développement de *Tarichium uvella* mihi, et si un développement conforme s'obtenait dans les cultures artificielles aussi pour les autres représentants de ce genre, il y aurait alors possibilité de tenter des expériences d'infestations avec les spores obtenues par culture artificielle.

« Les spores ordinaires des *Tarichium*, ainsi qu'il a été indiqué, mises en contact de toute façon avec le corps des insectes ne produisent jamais de contagion. »

Il va sans dire que je me joins très volontiers et sans aucune restriction à l'appel chaleureux que KRASSILTSCHIK adresse en terminant aux botanistes, aux entomologistes, voire même aux chimistes, pour leur demander d'unir leurs efforts en vue d'un but commun très digne d'exciter leur émulation : la victoire des agriculteurs contre le fléau ruineux des insectes nuisibles qui ravagent tour à tour nos champs, nos vergers et nos bois.

Paris, le 5 Avril 1889.

BULLETIN SCIENTIFIQUE

COLLECTION DES PREMIÈRES SÉRIES

(En vente chez O. DOIN, Éditeur, 8, place de l'Odéon, PARIS).

PREMIÈRE SÉRIE,

Dirigée par MM. GOSSELET, DESPLANQUE et DEHAISNE.

			Prix :
Tome I.	— 1869	(Quelques volumes).	10 fr.
» II.	— 1870	(Épuisé).	
» III.	— 1871	Id.	
» IV.	— 1872	Id.	
» V.	— 1873	(Quelques volumes).	10 fr.
» VI.	— 1874	Id.	—
» VII.	— 1875	Id.	—
» VIII.	— 1876	(Épuisé).	
» IX.	— 1877	Id.	

DEUXIÈME SÉRIE,

Dirigée par ALFRED GIARD

Tome X.	— 1878	(Épuisé).	
» XI.	— 1879	Id.	
» XII.	— 1880		8 fr.
» XIII.	— 1881		—
» XIV.	— 1882		—
» XV.	— 1883		—
» XVI.	— 1884-85		—
» XVII.	— 1886	(Quelques volumes).	15 fr.
» XVIII.	— 1887	Id.	—

TROISIÈME SÉRIE,

Dirigée par ALFRED GIARD.

Tome XIX.	— 1888		20 fr.
-----------	--------	--	--------

No. 17, no cards.

QK
2.5
OB
v. 100

(EXTRACTED FROM THE AMERICAN NATURALIST, DECEMBER, 1887.)

BOTANY.⁵

Character of the Injuries produced by Parasitic Fungi upon their Host-Plants.⁶—The first and most general injury which is produced upon plants by parasitic fungi results in consequence of the essential character of the fungus. Green plants have the power of converting inorganic materials into food for themselves. Fungi have not this power, but depend for their food entirely upon materials that have been previously elaborated by some green plant. Parasitic fungi take their food from their host while it is yet alive. The consequence to the plant, therefore, is a loss of food, a withdrawal of formed nutritive material, resulting, to greater or less extent, in starvation. In individual cells the contents usually suffer the greater injury, but the wall is also injured, as when penetrated by the haustoria, or suckers,—*e.g.*, upon the mycelium of *Peronospora*. But the special office of the haustoria is to absorb nutriment from the cell-contents. In some cases, as the above, the cell may live to be the continual prey of the parasite, and in some cases the cell is killed.

(2) While the food-supply of the plant is reduced, its power to replenish it is at the same time impaired,—*i.e.*, in case the fungus grows upon the green parts, as it does most frequently. The passage of light to the cells may be obstructed. The white moulds, or mildews, must obstruct it to some extent, black moulds growing over the surface, much more. The latter may occur either as true parasites, or as saprophytes upon “honey-dew.”

Impaired nutrition weakens the physiological power of the chlorophyll. The life of the cells preyed upon is frequently

⁵ Edited by Prof. CHARLES E. BESSEY, Lincoln, Nebraska.

⁶ Read before Section F, A. A. A. S., August, 1887.

destroyed, and they become brown and dry, as in many species of *Septoria*, or crumble and fall out, leaving holes in the leaf, as frequently in *Puccinia asteris*. Where the injury is not so severe, spots of various colors, often characteristic of the species, mark more or less definitely the extent of the mycelium. Exposure to light and the activity of the chlorophyll may be lessened in another way, by a change in the position of the infested stems and leaves. A very common effect of a parasite is a more erect and rigid growth, as in blackberries diseased with *Cæoma nitens*, the orange-rust, and in the garden spurges, *Euphorbia maculata* and *E. hypericifolia*, infested with the cluster-cup, *Æcidium euphorbiæ*. These spurges normally grow prostrate, with leaves widely spreading, allowing the greatest possible exposure to sunlight. In the diseased condition the stems are erect and the leaves less favorably spread.

(3) Growth may be abnormally accelerated or retarded, and both these effects may be produced in different cases by the same fungus. *Peronospora parasitica* growing upon the common peppergrass, *Lepidium virginicum*, forms a thick, felted coating of hyphæ upon the under surfaces of the leaves, and causes the latter to be somewhat contracted. The same fungus growing upon the stems of *Sisymbrium canescens* causes swelling and distortion. The *Æcidium* on *Sambucus canadensis* distorts the stem by accelerating growth, the infested side becoming swollen and the stem bent.

Exobasidium vaccinii on species of *Vaccinium* causes the leaves to shrink; forms believed to belong to the same species, growing on Azalea and Andromeda, produce inflated sacs, often several inches in diameter.

A distortion usually consists of the fungus producing it and the abnormal tissues of the host, as illustrated in the smut of maize and the curl of peach-leaves. In the latter extra layers of cells are formed on one side.

Æcidium euphorbiæ accelerates the growth of the stems of its host and retards that of the leaves,—an effect resembling that of insufficient light upon plants, as upon vines on the shaded side of a trellis, or potato-sprouts in a cellar. The orange-rust on blackberries retards the growth of the leaves, as does also the cluster-cup upon *Ranunculus abortivus*.

Fusicladium, the scab-fungus of apples and pears, retards the growth often of one side, while that of the other continues, producing unilateral development and curvature of the axis of the fruit.

(4) Fungi are not confined to the chlorophyll-bearing parts of plants; woody stems, roots, flowers, seeds, and fleshy fruits all have their parasites. One of the *Myxomycetes*, *Plasmodiophora brassicæ*, causes the disease called club-root in the roots of cabbage. The appearance of the club-root of roses, which is

troublesome in American green-houses, suggests a similar cause. Among those on woody stems are the black knot on species of *Prunus* and the *Gymnosporangia* on cedars.

The flower is attacked in the case of the "double blossom" of blackberries, the smuts of many grasses (such as *Ustilago rabenhorstiana* on *Panicum sanguinale*), and other cases that will suggest themselves. Often the entire inflorescence is arrested in its development. Smut of Indian corn attacks both staminate and pistillate flowers, and the green parts as well. *Ustilago antherarum* upon Caryophyllaceæ produces its spores in the anthers. *Tilletia caries*, the bunt of wheat, destroys the inner structure of the seed, without much changing the external appearance. The majority of Ustilagineæ attack the inflorescence, flowers, and fruit.

The young ovary is affected in various ways. Ergot distorts it and produces an abnormal, horn-like mass of tissue; smuts convert it into powdery spores. *Exoascus* changes that of *Prunus* into a pouch. *Fusicladium* kills the newly-formed pear.

(5) The last case is also an example of the premature falling of diseased parts, apparently by the formation of the separative layer before its time. Such a falling of leaves is well illustrated by *Septoria ribis* upon the currant. On the other hand, diseased parts may remain green after the other parts have matured in the usual way. Ovaries of *Eragrostis poæoides* filled with *Ustilago spermophorus* are most readily detected by this sign. Probably this may be classed as retarded development.

(6) Many fungi cause decay of ripe fruit, both while attached to its plant and after removal, while still alive. From a practical stand-point, the injury to fruit may be of two kinds; it may affect simply the appearance, and hence the market-value, or impair the quality, when it does not entirely destroy. The scab of apples is injurious in either of these degrees.

Some fungi begin their growth as parasites under ordinary conditions, to continue it under conditions more like those which accompany saprophytic growth; in this connection may be mentioned the potato-rot, beginning its growth upon the green plant and sometimes completing it in the cellar, but upon the still living tuber.

(7) Some valuable plants are liable to infection from others of less value; a knowledge of this fact in individual cases may be of great practical utility, as a remedy may be more easily and effectively applied. It is easier to destroy wild grapes and *Ampelopsis* than to destroy *Peronospora* upon cultivated grapes which have suffered infection from the wild ones. This remedy especially commends itself when one stage of the fungus grows upon one kind of plant and another stage upon another kind, as do cedar-apples and orchard-rusts.

Perhaps no line of investigation in economic mycology prom-

ises better practical results than this. The relations of a few of the Uredineæ are known, as are also the relations of Ustilagineæ and the yeast-like forms which may be looked for in manure ; but of the life-histories of the great group of "Fungi imperfecti," including many injurious species, almost nothing is known.

It is often difficult to estimate the extent of the injury to be attributed to any one of these causes, as several commonly operate together. Canada thistle attacked by *Puccinia suaveolens* wilts in hot sunshine, while healthy plants remain fresh ; and this appears to be due chiefly to the first cause, the loss of nutriment, which weakens the physiological power. Is it possible that the breaking of the epidermis by the sori, in this or other cases, increases the rapidity of transpiration ?

Farmers say the red rust of wheat produces comparatively slight injury, while the black rust is destructive. The black rust appears on the stems, which at that stage of growth have little chlorophyll, even if healthy ; the formation of winter spores appears to make a greater drain upon the resources of the host than the formation of summer spores, which have much weaker vitality.

Certain groups of plants are the prey of certain groups of fungi. Gymnosporangia grow upon Cupressineæ ; Ræstelie upon Pomaceæ ; Phragmidia upon the sub-order Rosaceæ proper. Numerous species of Uromyces are parasitic upon Leguminosæ, and the grasses support more Pucciniæ than does any other family of phanerogams. In these two genera no species is known to produce its teleutospores upon host-plants belonging to more than one order. The large majority of Uredineæ grow upon herbaceous plants, as do Ustilagineæ and Peronosporæ. In these groups a given species commonly inhabits but few species of host-plants, while some Erysipheæ inhabit a large number.

Many Erysipheæ grow upon woody plants. Fungi imperfecti are perhaps least of all to be classified with reference to hosts, though many of their genera may be. Woody plants have fewer active parasites than herbaceous plants. Water-plants and those of wet places are freer from attack than those which grow upon common soil.

To recapitulate, parasitic fungi injure their host-plants by (1) depriving them of nourishment ; (2) impairing the power of assimilation ; (3) abnormally accelerating or retarding growth, causing distortion ; (4) not only green parts are affected, but roots, stems, inflorescence, flowers, and fruit ; (5) leaves and fruit fall prematurely ; (6) decay is produced in ripe fruits before and after removal from the plant ; (7) valuable plants receive injury from those of less value by ordinary infection or heterœcism.

The amount of injury due to one cause is uncertain, since several act together.

Certain groups of phanerogams are liable to be attacked by certain groups of fungi.—*A. B. Seymour, Cambridge, Mass.*

Jan. 30, 1888.

From the AMERICAN NATURALIST, July, 1890.

BOTANY.

Some Elementary Botanies.—That there is a dissatisfaction with the commoner methods of teaching botany as set forth in the text books, is indicated by the numerous attempts of various teachers to give us better books. In no other science is there to-day such diversity of opinion as to the best method of introducing the pupil to the subject to be studied. As a result we have a multiplication of elementary books, each designed to lead the pupil into his work by a different route. For twenty years the little books by Miss Youmans have stood as a sort of protest against mere "book botany." Though faulty in many particulars, they were valuable in showing that there are other ways of teaching botany than the stereotyped ones. A recent book, "Descriptive Botany," by the same author, has much in it to commend. At the very beginning the pupil is told to supply himself with his own material for examination. He is told to "gather a variety of leaves; and to begin their study by comparing them," etc., etc. Further on we find this: "Pull up any herb which has a distinct stem, and compare the stem with the root," and so on repeatedly. This is excellent, and the pupil cannot fail to be greatly benefited by such a course. There is too marked an emphasis given to technical terms, which are needlessly printed in italics, and too frequently there is a dictionary-like brevity, as when we read that "The leaf of a fern is called a *frond*," and "The stalk or petiole of a frond is called a *stipe*." Why this is so is not hinted. The "Popular Flora" is just what it pretends to be—popular—and will be useful to the beginner who has prepared himself aright to take it up. It contains brief but plain descriptions of the more common flowering plants including cultivated as well as wild species. It is pleasant to note that the Gymnosperms are assigned to their proper place between Angiosperms and the Pteridophytes. It is not so pleasant, however, to note that the explanation of the structure of the flowers of the Conifers (the sole representatives of the Gymnosperms) is wholly erroneous. There has been an attempt to carry the old and discarded ideas as to floral structure over into the new classification. The Conifers as described in this book should go back into their old position, sandwiched between the Monocotyledons and Dicotyledons!

In many points the "High School Botany," prepared by H. B. Spotton for the use of Canadian students, has a considerable resem-

blance to Miss Youmans's work noticed above. There is the same admiration and following of Henslow's "schedules" for analysis, the same examination of representative plants, while in each there are several chapters given to generalizations. Most of the work of preparing this volume appears to be well done, but there are evidences here and there of haste. Thus while the true nature of the lichens is recognized on page 202, a little further on (p. 206) we have the old statement that "the lichens, from their peculiar constitution, may be regarded as transitional between the Algæ and the Fungi." Here we have a little new wine (p. 202) in a very old bottle (p. 206). The "Flora for the Use of Beginners" in this book is much like Miss Youmans's "Popular Flora." It is really a very useful little manual.

A recent English book, Edmonds's "Elementary Botany," has been placed before the American public by Longmans of New York. It is a much more scientific book than either of the preceding, although like them it teaches botany by observation. The principal difference is that the observation in this book is more profound, and is directed to essential rather than to superficial characters. The student is brought to study the plant as a living thing, rather than an object to be classified and labeled. We find that while 153 pages are given to structure and physiology, but 35 are devoted to classification. A few orders are selected, and in each a typical plant is suggested for study, while a few others are cited as common examples. The book is a very good one.

Dr. Campbell's "Structural and Systematic Botany," which has recently been brought out by Ginn & Co., is an attempt to supply a small and handy introduction to all parts of the vegetable kingdom. It is based upon, and to a certain extent is an abridgement of, Gœbel's "Outlines of Classification and Special Morphology of Plants," a work of great usefulness to the student in spite of its considerable cost. This introduction will be welcome to many a teacher and student who cannot afford the larger work.

After a few introductory pages devoted to methods of work, and a brief examination of the cell, the Protophytes are taken up, nine pages being devoted to them. Then follow in order the Algæ, Fungi, Bryophytes, Pteridophytes, and Spermaphytes. Very good outline drawings, largely original, accompany the text. A commendable feature of the work is the adoption of Eichler's arrangement of the flowering plants.

Here and there slips, due doubtless to haste, are noticed. Thus the figures B, C, and D, on page 132, are certainly not of "year-old cones of Scotch Pine," and on page 131 the relation of scale and "ovule-bearing leaf" are badly confused. There is also a looseness in

the quotation of the titles of works of reference and the names of authors and publishers on pp. 235-6. We note further that Ustilago is persistently spelled Ustillago.—CHARLES E. BESSEY.

The Completion of Saccardo's Sylloge Fungorum.—Eight years ago the first volume of this great work appeared, and this has been followed by others in rapid succession until now we have the eighth and final volume of the series. In these thick volumes, which aggregate more than eight thousand pages, nearly thirty-two thousand species have been described (exactly, 31,927). The completion of so great a labor in so brief a space of time must excite at once our wonder and admiration. We have here a work of vast extent, whose first and last volumes are near enough together in time, so that they are not appreciably separated by any change in plan, due to a change of view on the part of the authors. Whatever we may say of the plan of the work, and however much we may wish that a different one had been adopted, it is comfortable to know that here at least is a book completed upon the lines laid down by its author less than a decade ago. It is cheerful, also, to think that a generation has not died during the publication of the work, but that nearly all who saw its beginning have seen its completion. Thus the depressing influence of De Candolle's "Prodromus," dragging its way through fifty years to incompleteness, is counteracted, and we may again hope to see great undertakings inaugurated.

If we take the great masses of families as worked in this book, and make a distribution in an approximately natural system, we get a better idea of the numbers and extent of the fungi. For convenience of reference the number of species in each family is given, and the total number in each order or class.

PROTOPHYTA.

MYXOMYCETÆ.—(Vol. VII.)—Monadinaceæ, 49 species; Soro-phoraceæ, 9; Myxomycetaceæ, 383. Total, 441 species.

SCHIZOPHYTA.—Schizomycetaceæ, 659 species. (Vol. VIII.)

ZYGOPHYTA.

CONJUGATÆ.—Protomycetaceæ, 19; Chytridiaceæ, 132. (Vol. VIII.); Entomophthoraceæ, 20; Mucoraceæ, 200. (Vol. VII.); Total, 371 species. (Vol. VII.)

OÖPHYTA.

CŒOELOBLASTEÆ.—Saprolegniaceæ, 80; Peronosporaceæ, 96. Total, 176 specie s. (Vol. VII.)

CARPOPHYTA.**ASCOMYCETÆ.**

PYRENOMYCETÆ.—(Vols. I., II., and Add.)—Perisporiaceæ, 481; Sphæriaceæ, 5448; Coryneliaceæ, 2; Hypocreaceæ, 640; Dothidiaceæ, 351; Microthyriaceæ, 65; Lophiostomaceæ, 213; Hysteriaceæ, 372; *Hermihysteriaceæ*, 3. Total, 7575 species.

HYPODERMEÆ.—(Vol. VII.)—Uredineæ, 1224; Ustilagineæ, 284. Total, 1508 species.

— ? —

Phymetosphæriaceæ, 16; Onygenaceæ, 6; Laboulbeniaceæ, 15. Total, 37 species. (Vol. VIII.)

SPHÆROPSIDEÆ.—(Vol. III.)—Sphærioidaceæ, 3690; Nectrioidaceæ, 44; Leptostromaceæ, 203; Excipulaceæ, 143. Total, 4080 species.

— ? —

Melanconiaceæ, 606 species. (Vol. III.)

HYPHOMYCETÆ.—(Vol. IV.)—Mucedinaceæ, 1147; Dematiaceæ, 1579; Stilbaceæ, 344; Tuberculariaceæ, 594. Total, 3664 species.

SACCHAROMYCETÆ.—Saccharomycetaceæ, 30 species. (Vol. VIII.)

DISCOMYCETÆ.—(Vol. VIII.)—Caliciaceæ, 78; Gymnoascaceæ, 51; Cordieritaceæ, 5; Patellariaceæ, 161; Phacididiaceæ, 268; Stictaceæ, 229; Bulgariaceæ, 152; Dermateaceæ, 255; Ascobolaceæ, 130; Pezizaceæ, 1948; Heloellaceæ, 169; Cyttariaceæ, 7. Total, 3453 species.

TUBEROIDEÆ.—(Vol. VIII.)—Elaphomycetaceæ, 21; Cenococcaceæ, 1; Tuberaceæ, 102; Endogonaceæ, 6. Total, 130 species.

BASIDIOMYCETÆ.

GASTEROMYCETÆ.—(Vol. VII.)—Hymenogastraceæ, 78; Lycoperdaceæ, 426; Nidulariaceæ, 61; Phallaceæ, 81. Total, 646 species.

HYMENOMYCETÆ.—(Vols. V. and VI.)—Tremellaceæ, 258; Clavariaceæ, 371; Thelephoraceæ, 884; Hydniaceæ, 427; Polyporaceæ, 1972; Agaricaceæ, 4639. Total, 8551 species.

There are thus 1100 species of Protophytes; 371 of Zygomycetes; 176 of Oöphytes, and 30,280 of Carpophytes. Of the latter again there are 12,703 Ascomycetæ; 9,197 Basidiomycetæ, with 8,380 probably, but not certainly, imperfect stages of the former.

CHARLES E. BESSEY.

PROCEEDINGS OF THE NATURAL SCIENCE ASSOCIATION OF STATEN ISLAND.

SPECIAL No. II.

August, 1890.

LIST OF STATEN ISLAND FUNGI IN THE COLLECTION OF THE ASSOCIATION.

The following list of fungi, in the cabinet of the Association, was prepared from specimens collected from time to time by our members and finally transmitted to Mr. J. B. Ellis, of Newfield, N. J., to whom we are indebted for their determination.

ARTHUR HOLLICK.
N. L. BRITTON.

BASIDIOMYCETES.

- Agaricus (Collybia) velutipes*, Curtis. On *Ailanthus* stump.
Agaricus (Pleurotus) ostreatus, Jacq. New Dorp.
Agaricus (Hypholoma) sublateritius, Schæff. (probably). New Dorp.
Lentinus Lecontei, Fr. On sweet birch, Clove Lake.
Panus stipticus (Bull.), Fr.
Schizophyllum commune, Fr. On *Ailanthus*, New Dorp; on wild cherry.
Lenzites betulina (L.), Fr. On white oak, Richmond; New Dorp.
Polyporus lucidus (Leyss), Fr. On a dead oak, New Dorp.
Polyporus sulphureus, Fr. On dead wood.
Polyporus obtusus, Berk. (probably). On *Hicoria ovata*.
Polyporus adustus (Willd.), Fr. On *Fagus*. Woods of Arden.
Polyporus applanatus (Pers.), Fr. On *Salix fragilis* and *Quercus alba*.
Polyporus cinnabarinus, Fr. On dead wood.
Polyporus pergamenus, Fr. On dead wood.
Polyporus fomentarius (L.), Fr.
Polyporus betulinus (Bull.), Fr. On white birch.
Polyporus hirsutus (Wulf.), Fr. On *Ailanthus*, New Dorp; on *Castanea*, Richmond.
Polyporus versicolor (L.), Fr. On an old pear stump and dead sticks.
Polyporus conchifer, Schw. On a chestnut log, New Dorp.
Polyporus rimosus, Berk. On locust.*
Mucronoporus gilvus (Schw.), E. & E. On scarlet oak, beech or hornbeam (Bull's Head), and on chestnut.
Dadelea quercina (L.), Pers. On white oak stump, New Dorp.
Dadelea confragosa (Bolt.), Pers. New Springville.
Merulius corium, Fr. On dead wood, Bard Ave.
Fistulina pallida, B. & Rav. On chestnut, Richmond.†
Trpex sinuosus, Fr. On *Ailanthus* and chestnut sapling.
Stereum frustulosum (Pers.), Fr.
Stereum versicolor, Fr. On trees.
Corticium scutellare, B. & C. (probably a young specimen). On alder.
Corticium salicinum, Fr. On alder, Bull's Head.‡
Tremella foliacea, Pers.
Hirneola Auricula-Judæ (L.), Fr. On a chestnut log, New Dorp.
Geaster hygrometricus, Pers. South Beach, on the ground.
Bovista pila, B. & C.
Lycoperdon gemmatum, Fr. (probably). On dead stumps.
Lycoperdon cyathiforme, Bosc. Four Corners.||
Scleroderma vulgare, Fr.

ASCOMYCETES

- Valsa (Euvalsa) glandulosa*, Cke. On dead *Ailanthus*, old fort at Richmond.
Xylaria polymorpha, (Pers.) On a cherry stump, New Dorp.
Daldinia concentrica (Bolt.), Ces. & De Not. On beech, after fires; on black birch.

HYPHOMYCETES.

- Verticillium candidum*, Sacc. On black walnut.

MYXOMYCETES.

- Trichia chrysosperma* (Bull.), D. C. (probably).

The following notes are extracts from the letters of Mr. Ellis, accompanying the specimens :

* “*Polyporus rimosus*, Berk. Yours is just the same as specimens collected in Ohio by Prof. Morgan and determined by Cooke. *Polyporus salicinus* is darker and smoother, margin and all, and does not crack open like this. It is closely allied to *P. igniarius*, Fr. Your specimen is the first I have seen from this region.”

† “*Fistulina Hepatica* is credited to this country, and it is not absolutely impossible that your No. 8 is that species. It appears, however, to be the same as I find here, and which Cooke says is the *F. pallida*, B. & Rav.”

‡ “No. 12 must be *C. salicinum*, I think. I see no difference between your specimen and mine on willow. Yours is marked ‘on alder.’ Is it not on willow?”

|| “*Lycoperdon cyathiforme*, Bosc. . . . When young and white like cheese curd, this species is good eating. Peel off the outer bark, slice and fry it.”

89.2
[No. 20.]

CONTRIBUTIONS
TO THE
MYCOLOGY OF WESTERN NEW YORK.

I. THE FUNGI OF WESTERN NEW YORK.

BY CHARLES E. FAIRMAN, A. M., M. D.

MEMBER OF THE AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF
SCIENCE. CORRES. MEMBER OF THE ROCHESTER
ACADEMY OF SCIENCE.

[Extracted from the Proceedings of the Rochester Academy of Science,
Vol. I, Aug., 1890.]

GA

2.5

.03

v. 100

THE FUNGI OF WESTERN NEW YORK,

BY DR. CHARLES E. FAIRMAN.

In the introductory portion of this paper a review of the progress of mycology in the United States was given at length, and a resumé of the work of the late Hon. G. W. Clinton, of Buffalo, the pioneer investigator of the fungi of this section. A synopsis of the contributions of Clinton, contained in the Reports of the New York State Museum from the 23rd to the 39th, ended the historic treatment of the subject.

For the past five years the author has been investigating the mycologic flora of Orleans county, N. Y., and has collected over 425 different species during that time, which may be classified as follows :

Pyrenomycetes,	-	-	-	-	87	species.
Sphærosideæ,	-	-	-	-	61	"
Hyphomycetes,	-	-	-	-	43	"
Hymenomycetes,	-	-	-	-	96	"
Gasteromycetes,	-	-	-	-	7	"
Phycomycetes,	-	-	-	-	10	"
Myxomycetes,	-	-	-	-	17	"
Uredineæ and Ustilagineæ,	-	-	-	-	51	"
Discomycetes,	-	-	-	-	24	"
Imperfect and unclassified,	-	-	-	-	29	"

In the fungus flora of Orleans county, so far as known, the Hymenomycetes, or mushrooms and their allies, head the list with the greatest number of species, viz: 96, or nearly one-fourth the number of recorded species. The Pyrenomycetes come next with 86 species. The Hymenomycetes, being mostly of large size, were collected rapidly at first, while members of the other orders, for the most part microscopic, were overlooked. Lately, however, the Pyrenomycetes are coming to the front and will, doubtless, greatly outnumber the other fungi when our mycologic flora has been thoroughly investigated. The list of fungi found in Orleans county includes forms new to the state or, at least, not enumerated in the reports of Prof. C. H. Peck, and several new species and varieties which are set forth at the conclusion of this article. (One of the rarer forms is *Pleospora subsulcata*, E and E. See plate 4, fig. 1 and 2.)

All of the families into which Prof. Saccardo divides the Pyrenomycetes are represented in the mycologic flora of Orleans county except one (the Microthyriaceæ). Nearly forty (40) genera are found in the list of black fungi, the common genera, *Valsa*, *Hypoxylon*, *Eutypa*, *Rosellinia*, *Diatrype* and *Diaporthe*, having the greatest number of species to their credit. The "black fungi" previously mentioned belong to the saprophytes and exist on dead and decaying substances.

We now turn to consider some species of Pyrenomycetes which are parasitic on living plants, the mildews or Perisporiaceæ, the first family of Pyrenomycetes in the *Sylloge Fungorum*. Since the publication of Vol. I of Saccardo's *Sylloge*, there has been published a paper on the "Mildews of Illinois," by Dr Burrill, which reduces many species of the former work to synonyms. Our mildews (Orleans county) by the arrangement adopted by Peck in his reports, or by Saccardo in the *Sylloge*, are 19 in number, and by the revision of Burrill became reduced to 14 species. Among the host plants in Western New York attacked by mildew, we find cherry, horsechestnut, grape, lilac, honeysuckle, phlox, violet, larkspur, woodbine, aster, viburnum, elder, elm, beech, maple and gooseberry. The mildew on *Agrimonia eupatoria* (which has heretofore been referred to *Spherotheca Castagnei*, Lev.) is called by Burrill *Spherotheca Humuli*, (D. C.). Lyndonville specimens of a *Spherotheca* on common agrimony show perithecia larger, appendages shorter and more delicate, ascus and spores larger than the common *S. Castagnei*. Therefore *Spherotheca Humuli* (D. C.), Burrill, seems an appropriate name for our species (Plate 3, fig. 6).

The Sphærospideæ have sixty (60) representatives which are distributed in many genera, the principal ones being *Septoria*, with 13

species, *Diplodia*, 8, *Sphæroopsis* 5, *Phyllosticta* 4. *Pestalozzia* is represented by one species, the rare *Pestalozzia insidens* of Zabriskie, found on elms, which from the beauty and perfection of its spores has been called the "Prince of Pestalozzias" (Plate 4, fig. 9). We find a *Septoria* on common chickweed—*Septoria Stellariæ*, R. and D., and *Phyllosticta Cirsii*, Desm., on Canada thistle, new to the United States, until discovered at Lyndonville. *Haplosporella Nerii* which grows on oleander stems, *Phoma cyprena* on bittersweet branches, and *Phyllosticta filipendulina* on cultivated spiræa are forms which are new to New York State.

The Hyphomycetes are represented in the collection by a number of species (also distributed among many genera) including *Cercospora*, *Fusicladium*, *Ovularia* and *Ramularia*, parasitic on living plants, and *Botrytis* and *Verticillium* on dead or decaying substances. While not exerting as destructive action upon vegetation as some other fungi, this group affords many delicate microscopic forms. Two members of this group, viz: *Fusicladium dendriticum* or apple-peel fungus, and *Ramularia Fragariæ* or the strawberry leaf blight, have been found with us, seriously injuring their respective hosts.

The Hymenomycetes comprize nearly one-fourth the collection, and our flora will be found quite rich in these forms. In the Agaricus family we find many genera present with us, headed by *Agaricus* proper with 22 species, and including members of such genera as *Russula*, *Lactarius*, *Marasmius*, *Coprinus*, etc. The striking feature is that no specimens of the large genus *Cortinarius*, have, as yet, been found, although looked for with assiduity. Elsewhere in the eastern portion of New York State *Cortinarii* have been plentifully gathered. The family Polyporei is represented by a number of genera and species. A few *Boleti* have been found, in the months of July and August, in our locality, but *Polyporus* has the larger number of species to its credit (about 20).

It may be noted in passing that it has long been known to students of fungi, that *Polyporus applanatus* often attains a large size and that its pores are ferruginous, with a white orifice, which causes the inferior surface of the plant to appear as if white-washed. It has also been known that these whitened surfaces when bruised or scratched show quickly the brown underlying color. I have seen this property made use of in a decorative sort of way, which I have nowhere seen mentioned. Pictures of trees, houses and various objects are drawn upon the white inferior surface of the fungus and are permanent. In many country houses these prepared objects are found plentifully adorning mantels and bric-a-brac receptacles.

Among the Hydneæ we have only the very common genera *Hydnum*, *Irpex*, and *Grandinia*.

The Thelephorei (4th family) are also well represented by species of *Corticium*, *Stereum* and other genera. We have the rare *Corticium lividocæruleum*, Karst, new to the state, and, probably to the country, which agrees with an authentic specimen from Karsten, and a new species, the *Corticium rhodellum*, Peck, of a beautiful rose color.

The Clavariæ are only sparingly represented by a few common forms.

The peculiar gelatinous family Tremellini is represented by species of *Tremella*, *Exidia* and *Dacrymyces*, which are commonly found on moist hemlock stumps.

The next order of fungi the Gasteromycetes or puff-balls is known only by a few common puff-balls, one member of the offensive Phalloids and *Nidularia pulvinata*, Schw, a species rarely occurring in the state.

The order Phycomycetes is represented by a number of "blights" and "moulds." In wet seasons the blights are a source of great annoyance from the wide spread damage done to field and garden crops. It is only necessary to mention the *Peronospora viticola* or grape blight, and the *Peronospora gangliiformis*, DeBy, the lettuce blight, to prove this.

The Myxomycetes or slime moulds, a group of remarkable organisms, which have occupied debatable positions in various systems of classification, are represented by 17 species. During the vegetative portion of their life history these organisms consist of naked masses of protoplasm, the so-called plasmodia, variously colored, and possessing the power of changing their forms and places, and hence are said to be motile. Under proper conditions, this protoplasmic vegetative state ceases, and a resting state is entered upon, with the formation of spores in spore cases, or sporangia, which assume resemblances to puff-balls, or other fungi. They have always been objects of great interest not only from their curious life history, but also from the delicacy of their fructification and the beauty of their spores and threads, when investigated by the microscope. We have some of the rare forms of this order, as well as the common genera *Fuligo*, *Trichia* and *Arcyria*. We have also one new species a *Didymium* (*Didymium Fairmani*, Sacc.) found growing on the lower surfaces of leaves of *Smilacina bifolia*, which will be noticed later on.

Taking up for our next consideration the rusts and smuts we find the list includes 51 species. The genera represented are *Ecidium*, *Puccinia*, *Uromyces*, *Phragmidium*, *Melampsora*, *Coleosporium*, *Roestelia*

and *Cæoma*. This group is of considerable interest to agriculturists on account of the damage done to cultivated plants. Some of the species in the list belong to common forms, such as wheat rust and corn smut, and others to rare members of the order. Only a few features can be here noted in a general way. We have found only one species of *Roestelia* (*Roestelia lacerata*). These roestelia forms, as proven by the cultures of Halsted, Thaxter and Farlow are the æcidia or cluster cup stage of species of *Gymnosporangium*, or the so-called "cedar apples," which are found upon species of cedar and juniper. The *Roestelia* of our flora was found upon *Cratægus* leaves, but a search for the "cedar apples" upon Juniper has proved fruitless, nor have I found evidence of the occurrence of species of *Gymnosporangium* in Western New York.

While this review was in course of preparation I noticed along the roadside some patches of *Malva rotundifolia* whose leaves were spotted. A removal of some of the spotted leaves and a microscopic examination enables me to announce the unfortunate presence with us of *Puccinia Malvacearum*, or as it is commonly termed, the hollyhock disease. And next I examined my hollyhocks and found them attacked. There are many features of interest about this parasitic rust. When first reported from Australia it was said to be very destructive. It has been recorded in this county in scattered localities in Vermont, Massachusetts and California, and unfortunately seems to be spreading. It has only recently been found in this state. Some specimens were lately sent to Prof. Peck, from Geneva, which he has pronounced to be this fungus, and in an article in the "Country Gentleman" recommended prompt measures to be taken for its suppression; but it probably has secured too firm a foothold to be stamped out. "As far as the attacks on *Malva rotundifolia* are concerned it may be a good thing, but not so with its attacks upon cultivated hollyhocks," says Prof. Peck in a recent letter. Prof. Farlow of Harvard College, has compared our specimens with others and pronounced them as the same fungus found in Massachusetts, Vermont and Central New York. "The spots are much lighter colored than the Western and California form which is not the true *Puccinia Malvacearum* but *Puccinia Malvastri*, Peck." The color of the latter is black brown, while that of the former is yellow brown. The depth of the apparent color in *P. Malvacearum* is dependant to some extent on the fact of the production of promycelia or not, as this species is a member of the sub-genus in which the spores germinate at once *in situ*, if the promycelia are just starting the color appears lighter, if they have not begun to appear it is somewhat darker. (Sec. Farlow in litt.)

Among the Discomycetes we have a wide range of forms from our largest morel—*Morchella esculenta* var *conica*—to the microscopic *Pezizas*. The genera represented are *Peziza*, *Morchella*, *Propolis*, *Stictis*, *Patellaria*, *Dermatea*, *Tapesia*, *Encolia*, *Helotium*, *Ascophanus* and *Ascobolus*. Our earliest species is *Peziza coccinea*, Jacq., which occurs at the same time as the spring beauty (*Claytonia*) and the *Hepatica*. Some of our species are quite rare, as for instance *Tapesia Rosæ*, (Pers.), found on dead stems of wild rose. Mr. Ellis, to whom specimens were sent, says that he had never before had *Tapesia Rosæ* from this country. We have also, recently, found a *Dermatea* growing on twigs of *Lindera benzoin*, which is apparently near *D. Viburnicola* Ell. in Torr. Bulletin (*D. purpurea*, Ell. olim).

The following list contains the new species and varieties which we have added to the mycologic flora of Western New York.

1. Fenestella amorphia, E. and E.

Jour. Mycol, vol. 4, page 58, vol. 5, page 79.

On fallen branches of hickory (*Carya*) in the spring. We have attempted to trace the development of the sporidia of this fungus in an article prepared for Jour. Mycol, vol. 6, on the "Development of Some Fenestrate Sporidia." See also plates I and II, Jour. Mycol. vol. 6, for figures of the *Fenestella*.

2. Didymosphæria accedens, Sacc., *n. sp.*, (Plate 4, fig. 12).

"Peritheciis gregariis, tectis, $\frac{1}{4}$ – $\frac{1}{3}$ m. m. lat., ostiolo laeve papillato erumpento, nucleo primo albo. Ascis tereti-clavatis $120 \times 10 \mu$., paraphysibus obsoletis. Sporidiis senis octonisve, ellipsoideis, utrinque rotundatis, 1 septatis, leviter constrictis, $20-22 \times 9-11 \mu$., fuligineis monostichis. Affin. *D. Rhamni* et *D. incarcerato* ostiolo ascisque diversa." Saccardo in litt. Hab. on dead branches of some tree (Ash?).

April and May, 1889, Lyndonville, N. Y.

3. Anthostomella eructans, E. and E., *n. sp.*, (Plate 4, figs. 7–8).

Perithecia gregarious, globose, $\frac{1}{2}$ – $\frac{3}{4}$ m. m. diam., with thick coriaceous walls, buried in the wood, abruptly contracted above into a short neck with an obtuse-conical erumpent ostiolum. Asci cylindrical, $75-80 \times 7 \mu$. (p. sp.) with abundant paraphyses. Sporidia uniseriate, brown, continuous, rather acutely elliptical, $10-15 \times 5-7 \mu$. (mostly $12 \times 5 \mu$.). The surface of the wood is uniformly blackened and the sporidia when mature are discharged as in *Massaria*.

On decorticated (maple?) limb.

Lyndonville, N. Y., May, 1889. Fairman, No. 42.

4. **Lophiostoma Pruni**, E. and E., (Plate 4, figs. 10-11).

Jour. Mycol, vol. 4, page 64. Also figured in Berlese, Icones Fungorum, Fasc. I Part I, Tab. VI, fig. 3.

On *Prunus serotina*, Lyndonville, April, 1888.

5. **Lophiostoma rhopaloides**, Sacc., var. *pluriseptata*, n. var.

Differs from the type in having sporidia three to five septate, instead of three septate.

On dead branches of *Maple*, May, 1889.

6. **Pseudovalsa Fairmani**, Ellis and Everhart, n. sp. (Plate 3, figs. 1, 2, 3, 10, 11).

*Stromata convex pulvinate 1-1½ m. m. diam., formed of the slightly altered substance of the inner bark, the surface only sub-carbonized and blackened, not surrounded by any distinct circumscribing line, covered by the epidermis which is pierced by the stout, short, cylindrical or conical ostiola, with smooth or quadrisulcate tips. Perithecia 4 to 8 in a stroma, closely packed, ovate or subangular from compression, about ⅓ m. m. diam. with whitish, waxy contents. Asci (p. sp.) 75-85x20 μ., mostly only six (6) spored. Sporidia oblong cylindrical, yellowish, 3 septate, 30-40x5-7 μ., slightly constricted at the septa. The young stromata contain an abundance of *pycnidial* spores, (about the size and shape of the *ascospores*) borne on stout or branching sporophores about as long as the spores themselves.

On dead hickory limbs (*Carya*), Lyndonville, N. Y.

(In one specimen 5-septate sporidia were found.)

7. **Vermicularia phlogina**, Fairman.

Botanical Gazette, March, 1887.

On leaves of *Phlox divaricata*, Ridgeway, N. Y.

8. **Vermicularia solanoica**, n. sp.

Perithecia superficial, numerous, black, 150-175 m. m. in diam. Bristles few, mostly uniseptate, at times continuous, of various lengths, the longer ones gradually attenuated to an acute tip, brownish, with tips subhyaline, 75-100 μ. long and 5 μ. wide at the base. Conidia oblong, fusoid, subarcuate 25-30x2½-3 μ. Endochrome light green, continuous or faintly divided near the center, granular, nucleate.

*In the pycnidial stage the fungus might be taken for a *Hendersonula*. In this stage the stroma is orbicular, depressed, black outside, whitish waxy, horny within, divided into ovate, globose or angular cells from the surface of which spring the sporophores (40-60x3 μ.) bearing at their extremities the oblong cylindrical, subhyaline mostly 3 septate sporules (35-50x6-9 μ.) generally slightly constricted at the septa. The surface of the stroma in this pycnidial or the *Hendersonula* stage is at times indistinctly papillose from the slightly prominent ostiola. This stroma is also seated on the inner bark and erupts through the epidermis. At first the pycnidial spores are only granular and nucleate, but soon become from three to four septate in the progress of development.

On dead stems of *Solanum dulcamara*, Ridgeway, Orleans Co., N. Y., May, 1889.

This might be referred to *V. Dematium*, but until the limits of that species are fixed we propose this as new.

9. *Phoma Weldiana*, n. sp.

Perithecia few, scattered, black, shining, small, oval or rotund. 25-30 μ . diam. Spores oblong, pointed at one end abruptly, or both at times rounded, 10-15x3 μ .

On decorticated wood of *Euonymus atropurpureus*. Lyndonville, N. Y., May, 1889.

Name (Weldiana) after Miss L. A. Weld, who has assisted me in the determination of host plants and in the preparation of the plates.

10. *Phoma albovestita*, n. sp.

Cortical spots mostly surrounded by a white zone, most apparent before rupture.

Perithecia small, erumpent, nestling in the inner bark, occasionally clustered, dull black. Spores oblong, ends obtusely rounded. 4-7x1 $\frac{1}{2}$ -2 μ .

On bark of *Juglans cinerea*, May, 1889.

11. *Phoma Lyndonvillensis*, n. sp., (Plate 3, fig. 4).

Occupying faded spots on the stem. Perithecia ostiolate filled with minute oval or oblong sporules, sometimes nucleated, 3-6 μ . diam.

On stems of *Malva rotundifolia*, April, 1888. Found on the stems of plants which had been attacked the previous fall by *Septoria malvicola* E. and M. Both the *Phoma* and *Septoria* may be connected as states of some higher or ascomycetous fungus. In the spring, as early as the snow has gone, one can find, on the green stems of the *Malva*, here and there, bleached or dead spots, generally one inch long on which the *Phoma* grows. There is a *Phoma Malvacearum*, but the above peculiar growth has induced me to separate our plant.

12. *Phoma Rudbeckiæ*, n. sp.

Perithecia numerous, erumpent, globose depressed, ostiolate, light black. Sporules oblong, rounded at the ends, 4-6 μ . long, 2-3 μ . broad, hyaline.

On dead stems of *Rudbeckia laciniata*, Lyndonville, N. Y., April, 1888.

13 *Septoria Fairmani*, E. and E.

Jour Mycol., vol. 5, page 151.

On living leaves of hollyhock (*Althæa rosea*), Lyndonville, N. Y., June, 1889.

14. Septoria malvicola, E. and M.

Jour. Mycol., vol. 3, page 65.

On leaves of *Malva rotundifolia*.

Prof. Peck has referred our specimens to *Septoria heterochroa*, Desm.

15. Diplodia spiræicola, E. and E., *n. sp.*, (*in litt.*) (Plate 4, fig. 13). Spores smaller than either of the described species on Spiræa, measuring $8-10 \times 3-4 \mu$.

D. Spiræae has spores $14-20 \times 8 \mu$, and *D. Spiræina*, Sacc., has spores $20-22 \times 10 \mu$. (Mr. Ellis now thinks this species may come under *D. Spiræae*.)

On dead stems of cultivated *Spiræa hypericifolia*, June, 1889.

16. Diplodia maura, C. and E., *n. var.*

Var. *Americana* Ell. *in litt.*

On dead limbs of mountain ash, *P. Americana*.

17. Morthiera Thuemenii, Cooke, var. *sphaerocysta*, Peck, (*in litt.*).

Jour. Mycol., vol. 5, page 79.

On *Crataegus* at North Ridgeway on the County line road between Orleans and Niagara counties.

18. Haplosporella Ailanthi, E. and E. (Plate 3, fig. 5).

Jour. Mycol., vol. 5, page 147.

On dead *Ailanthus glandulosus*.

19. Sphaeropsis Lappæ, E. and E., *n. sp.*

Perithecia scattered, subglobose, $\frac{1}{3}$ m. m. in diameter, at first covered by the cuticle, soonerumpent superficial. Sporules elliptical, brown, with a single large nucleus, $15-20 \times 8-10 \mu$.

On dead stems of burdock (*Lappa major*), May, 1889.

20. Sporonema pallidum, E. and E.

Jour. Mycol., vol. 5, page 153.

On decorticated maple, Ridgeway, N. Y., May, 1889.

21. Sporiodesmium toruloides, E. and E., *n. sp.*, (Plate 4, fig. 3).

Forming small (1 m. m.) gregarious, cushion-like, black tufts, sometimes subefused. Conidia various, mostly toruloid, forming simple or branched chains of cells, $12-25 \times 5-7 \mu$. Most of the component cells divided by a longitudinal septum, or also subglobose $5-7 \times 5-7 \mu$. This is closely allied to *Septonema toruloideum*, C. and E., and to "*Coniothecium*" *toruloideum*, B. and C., but differs from both of these in its longitudinally divided (muriform) cells, which are also slightly muricate-roughened.

On dead wood and bark of *Cornus*, Lyndonville, June, 1889.

22. Discella pilosula, E. and E.

Jour. Mycol., vol. 5, page 153.

On a decorticated maple, Lyndonville, April, 1889.

23. Septoria divaricata, E. and E.

Jour. Mycol., vol. 3, page 85, vol. 5, page 151.

On living leaves of *Phlox divaricata*, Lyndonville, N. Y., May, 1889.

24. Coniosporium Fairmani, Sacc. (Plate 4, figs. 14-15).

Jour. Mycol., vol. 5, page 78.

Ab affin *C. Apiosporiade* differt conidiis multis minoribus, 5-7 μ . d., globosis, levibus, fuliginosis, subinde 1-nucleatis. On cortex of Hubbard squash, exposed to the weather, Lyndonville, N. Y., February 10, 1886. The fungus covers the rind with black sooty patches.

25. Didymium Fairmani, Sacc., (Plate 3, figs. 7, 8, 9).

Jour. Mycol., vol. 5, page 78.

Dignostitus peridiis, sparsis, sessilibus, floccis hyalinis laxè reticulatis, sporis levibus, 8-10 μ . diam, crystallis eximie stellatis. Columella subglobosa, fuscella.

On the lower surfaces of leaves of *Smilacina bifolia*. Aug., 1886.

In the original publication of the species, named by Dr. Saccardo, there was an error in translation, and I have thought best to insert here the original diagnoses.

26. Mucor Tæniæ, n. sp., (Plate 4, figs. 4, 5, 6).

Sporangiferous hyphæ erect, rarely, if ever branched, septate, yellow, 7 μ . diam.

Sporangia globose, brownish or yellow brown, smooth, mostly 40 μ . in diam. Columella elliptical or sub-sphaeroidal, at times with contraction at the base, brownish.

Spores globose, or ellipsoid, light yellow, 3-5 μ . in diam., with smooth epispore. Zygosporos not observed.

On segments or joints of tape worm (*Tænia solium*). The fungus forms a felted mat on the affected joints, of a sordid yellow color. At times the hyphæ have contents contracted into bands.

The parasitism of the mucor upon the entozoon or intestinal parasite is curious and interesting. It seems to me distinct from *Mucor mucedo* in smooth sporangia and spores, and from *Mucor erectus*, Bainier, by the color of the spores.

27. Pseudohelotium Fairmani, (E. and E.), Sacc. page 302,

No. 1,262, Sylloge Fungorum, vol. 8.

Peziza (mollisia) Fairmani, E. and E. Jour. Mycol., vol. 4, page 56.

On inner surface of bark (Oak ?) lying on the ground. Ridgeway and Lyndonville, N. Y., April, 1888.

28. *Helotium fumosum*, E. and E.

On dead stems of *Leonurus cardiaca* and *Lappa major*, May, 1889. Specimens were furnished for distribution in N. A. F.

29. *Camarosporium acerinum*, E. and E., n. sp.

Differs from the allied *C. subfenestratum*, B and C., in its broader (15-26x8-10 μ .), triseptate spores, and more prominent perithecia.

On dead maple limbs, Lyndonville, N. Y.

C. subfenestratum was described by Berkeley from specimens on *Robinia*, and no other host was given. A form found at Lyndonville, on *Ailanthus glandulosus* would be *C. Berkeleyanum*, (Lev.) of some authors, but Mr. J. B. Ellis informs us that Prof. Farlow, who has the original specimens, says this is scarcely distinct from *C. subfenestratum*. Evidently the three species are very closely allied.

30. *Tubulina cylindrica*, (Bull.) var. *acuta*, Peck, (*in litt.*)

Differs very noticeably from the type in having the peridia *acute* instead of "*rounded*" at the apex. Otherwise no marked differences exist. Hab. on rotten stumps in woods.

The illustrations were drawn from camera lucida sketches, by the author and Miss L. A. Weld.

PLATE 3.

Pseudovalsa Fairmani, E. and E.

- Fig. 1. Portion of hickory bark with fungus (slightly enlarged).
Fig. 2. An ascus and sporidia.
Fig. 3. Sporidia.

Pycnidia state of the above named *Pseudovalsa*.

- Fig. 10. A branched sporophore, with two pycnidiospores.
Fig. 11. Another form of pycnidiospore.

Phoma Lyndonvillensis, Fair.

- Fig. 4. Spores of the fungus.

Haplosporella Ailanthi, E and E.

- Fig. 5. Group of spores.

Sphærotheca Humuli, Burrill.

- Fig. 6. *Ascus* and *Sporidia*.

Didymium Fairmani, Sacc.

- Fig. 7. *Flocci* and spores.
Fig. 8. Crystals.
Fig. 9. Leaf of *Smilacina Bifolia*, with groups of the *Didymium*
(Nat. size).

PLATE 4.

(Scale the same as Plate I.)

Pleospora subsulcata, E. and E.

- Fig. 1. Ascus and sporidia.
Fig. 2. A sporidium (*involved in mucus*).

Sporidesmium toruloides, E. and E.

- Fig. 3. Spores of the fungus.

Mucor Tæniæ, Fairman.

- Fig. 4. A sporangium.
Fig. 5. Spores.
Fig. 6. Hypha with contracted contents.

Anthostomella eructans, E. and E.

- Fig. 7. Ascus with sporidia.
Fig. 8. A group of sporidia.

Pestalozzia insidens, Zabriskie.

- Fig. 9. Two spores of the fungus.

Lophiostoma Pruni, E. and E.

- Fig. 10. A single sporidium.
Fig. 11. An ascus with sporidia.

Didymosphaeria accedens, Saccardo.

- Fig. 12. Sporidia of the fungus.

Diplodia spiræicola, E. and E.

- Fig. 13. Spores of the fungus.

Coniosporium Fairmani, Saccardo.

- Fig. 14. Rind of squash (nat. size) bearing the fungus.
Fig. 15. The spores of the *Coniosporium*.

No. 215 no cards.

Estratto dal *Bullettino della Società botanica italiana*
(Adunanza della Sede di Firenze dell' 11 Gennaio 1891)
nel *Nuovo Giorn. Bot. Ital.*, Vol. XXIII, N. 2, Aprile 1891.

NOTA MICOLOGICA DEL DOTT. P. VOGLINO.

Fin dai primi giorni del mio arrivo in Casale il distinto botanico cav. F. Negri mi parlava di una regione poco distante da Casale ricchissima per vegetazione.

Colta l'occasione di alcune belle giornate del passato ottobre potei infatti colla saggia compagnia del chiarissimo avv. Negri esplorare il bosco della Badia di Lucedio in territorio di Trino e notai, per quanto alla sfuggita, che l'asserzione del Negri era verissima.

Tralascio per ora di parlare delle *Fanerogame*, già studiate in parte dal Rosellini e dal Negri e che d'altronde erano già in istato molto avanzato. Di queste pubblicheremo fra non molto un esteso catalogo, essendovi delle specie molto interessanti.

Mi limito a ricordare alcuni *Imenomiceti* non avendo potuto visitare il bosco che due volte e solo in ottobre. Le diverse specie però che ho trovato sono per la maggior parte abbastanza importanti ed alcune nuove per l'Italia, per il che ho creduto opportuno di darne subito una breve comunicazione.

Il bosco in generale è formato da *quercie*, *acacie*, *pioppi* ed *ontani*.

Frequentissime potei notare l'*Amanita virosa* Fr., l'*A. phalloides* Fr., l'*A. pantherina* Fr., l'*A. rubescens* Fr., l'*A. spissa* Fr., e l'*A. venenosa* var. *fulva* = *A. Mappa* Fr., che rassomigliava per l'aspetto generale alle figure date dal Vittadini (*Funghi mangerecci e velenosi*) dell'*Agaricus citrino-albus*, ad eccezione di uno sviluppo di gran lunga superiore. Dappertutto trovai l'*Amanitopsis vaginata* Bull., tanto la forma cinerina che l'aranciata. Comunissima vi è la *Lepiota procera* Scop., così anche trovai diversi esemplari di *L. erminea* Fr. riscontrata da me

per la prima volta in Italia nel boschetto dell'Orto botanico di Torino, e di *L. naucina* Fr., nonché di *L. glioderma* Fr., trovata unicamente sin ora verso S. Germano Chisone (Pinerolo). Comune assai nei boschetti di acacia, l'*Armillaria mellea* Valh. forma grande e piccola. Frequenti il *Tricholoma spermaticum* Fr., ritenuto fin ora non troppo comune in Italia, il *T. terreum* Schaeffer, il *T. saponaceum* Fr., *T. sulphureum* Bull., *T. acerbum* Bull., *T. melaleucum* Pers. e la *Clitocybe odora* Bull. di un bel colore verde-azzurro.

Abbastanza comuni la *Clitocybe dealbata* Sow., la *C. flaccida* Sow. e la *C. nebularis* Batsch.

Intorno ai tronchi di quercia specialmente nella parte più umida del bosco trovai molti esemplari di *Collybia radicata* Rell., *C. fusipes* Bull., *C. dryophila* Bull. e *C. muscigena* Schum.

In due diversi punti nel fitto dei boschi di acacia che si trovano nella parte centrale e più umida della località potei raccogliere alcuni esemplari di *Collybia ventricosa* Bull., specie che non si era ancora riscontrata in Italia: tutti gli individui erano somigliantissimi alla figura data dal Cooke nelle *Illustrations of british fungi*, tav. 145 A.

Negli stessi punti ora ricordati riscontrai anche la *C. rancida* Fr. rappresentata da individui con un pileo non superiore ai 2-3 cm. e con uno stipite lungo dai 12 ai 20 cm. e per il colore e disposizione delle lamelle quasi simili alla figura data dal Cooke (loc. cit., tav. 153). È questa la prima volta ch'io trovo questa specie, anch'essa nuova per l'Italia.

Frequentissime poi notai le *Mycenae* più comuni fra le quali la *Mycena pura* Pers., *M. lineata* Bull., *M. flavo-alba* Fr., *M. lactea* Pers., *M. galericulata* Scop. colle diverse varietà, *M. polygramma* Bull., la *M. filipes* e la *M. corticola* Schum.

Fra le *Omphaliae* abbastanza comune l'*Omphalia umbilicata* Schaeffer, *O. stellata* Fr., *O. setipes* Fr., ed *O. Fibula* Bull.

Trovai pure alcuni esemplari di *Pleurotus corticatus* Fr.: comuni specialmente nella parte periferica del bosco, il *P. olearius* Dec., il *P. ostreatus* Jacq. ed il *P. perpusillus* Fr.

Degli *Hygrophorus*, l'*H. eburneus* Bull. e l'*H. fusco-albus* Lasch.: notai comunissimo in tutto il bosco, specialmente nelle parti umide, l'*H. virgineus* Wulf.

Poco rappresentati i *Laclarius* e le *Russule* delle quali

potei solo raccogliere le specie più comuni dell'alta e media Italia.

Sopra alcuni esemplari di *Clitocybe nebularis* Bat. in istato di putrefazione trovai la *Volvaria Loveiana* Berk., non ancora riscontrata in Italia. I cinque esemplari che potei studiare somigliavano perfettamente alla figura data dal Berkeley (*Outl. of Brit. Fung.*, tav. 7, fig. 2).

Comune riscontrai il *Pluteus cerrinus* Schaeffer ed in alcuni punti, specialmente della parte interna, alcuni esemplari di *P. nanus* Pers.

La specie che più attirò la mia attenzione per la sua grande diffusione è l'*Entoloma nidorosum* Fr., del quale si trova addirittura coperto tutto il bosco. Avevo già notato, specialmente presso Alba e Massa, che questa specie si presentava sempre con numerosi esemplari ma non avevo ancora potuto notare una località nella quale si potesse raccogliere per l'estensione di parecchi chilometri continui e numerosissimi esemplari.

Abbastanza frequente notai pure il *Clitopilus Prunulus* Scop.

Del genere *Pholiota* non trovai che alcuni esemplari di *Ph. radicata* Bull., simili alla figura data dal Cooke, loc. cit., tav. 361.

Comune assai l'*Inocybe dulcamara* Alb. et Schw.: meno l'*I. obscura* Pers. Potei raccogliere anche alcuni esemplari che corrispondevano all'*I. phaeocephala* Bull., la quale risulterebbe nuova per l'Italia. Comunissime specialmente lungo le rive delle strade l'*I. rimosa* Bull. e l'*I. fastigiata* Schaeffer.

Potei pure raccogliere l'*Hebeloma fastibile* Fr., l'*H. testaceum* Batsch, l'*H. longicaudum* Pers., l'*H. mesophaeum* Fr. e l'*H. crustuliniforme* Bull.: i due ultimi si trovano comuni quasi dappertutto.

Sopra un tronco putrido di quercia notai alcuni esemplari di una *Flammula* che, dopo averli attentamente studiati, potei riferire alla *F. fusa* Batsch e che rassomigliavano alla figura dell'*Agaricus hybridus* di Bull. (*Champignons de France*, tavola 398). Comunissima trovai la *Naucoria cerodes* Fr., la *N. sideroides* Bull. e la solita *Galera tenera* Schaeffer.

Fra i *Cortinarius* tutte le solite specie; ma credo ce ne debbano essere in buon numero avendo notato in molti luoghi degli esemplari già putrefatti.

Notai alcuni esemplari di *Stropharia aeruginosa* Lasch., *S.*

coronilla Bull, *S. merdaria* Fr. e *S. stercoraria* Fr. Comuni
l' *Hypholoma silaceum* Pers. e l' *H. sublateralitium* Schaeffer,
comunissimi l' *H. elaeodes* Fr. e l' *H. fasciculare* Huds.

Fra le diverse specie mangereccie si trovano comuni l' *Ama-
nita caesarea* Scop. ed il *Boletus edulis*, che vengono raccolte
quotidianamente in grande quantità e portate specialmente nel
mercato di Casale.

Notes on Three New or Noteworthy Diseases of Plants.

By F. D. CHESTER.

During the work of the past summer my attention has been called to three diseases of cultivated plants, caused by fungi, which are apparently new.

A brief description of them will be given here, while a fuller account, with figures, will be reserved for one of the regular publications of the Experiment Station.

ANTHRACNOSE OF THE TOMATO.

This disease appeared during the past summer upon the grounds of the Experiment Station, where it has caused a considerable destruction of fruit.

So far as observed it does not affect the green tomato, but rather at the point when it just begins to color, and from that on to complete ripening. When, however, an attack is once made, the malady spreads so rapidly as to occasion serious loss before the fruit can be gathered.

The disease shows itself upon the tomato as sunken, discolored spots, each with a dark center, becoming black. These spots increase in size, or by confluence cover a large portion of the decaying fruit. Over this area the fruit is black and shrunk, flattened or depressed, surrounded by a shrunk, corrugated, discolored skin; the dark centers due to the gregarious acervuli with their dark setæ.

The disease is easily and quickly produced by introducing the spores within a puncture made by a sterilized needle, but no results have yet come from repeated attempts to produce the disease by sowing the spores upon the uninjured surface of either ripe or green tomatoes. The fungus causing the trouble is a species of *Colletotrichum*. It is clearly distinct from *C. nigrum* E. & Hal., found by Dr. B. D. Halsted on cultivated peppers, which, however, it somewhat resembles. Attempts to grow the tomato fungus on peppers, even by introducing the spores of the latter beneath the skin, were unsuccessful.

The following botanical description of the fungus is appended:

Colletotrichum Lycopersici, n. sp.

Spots depressed, circular, slightly discolored, center black, 5–10 μ in diameter, afterwards becoming irregular and confluent. Acervuli abundant, densely gregarious, rusty brown to black, applanate, 95–150 mm. in diameter. Setæ abundant, fuliginous, generally curved, rarely undulate or straight, often geniculate in places, gradually tapering, septate, length 65–112 μ , about 5 μ at base. Spores oblong 16–22 μ ×4, average 18–20×4, ends subacute, hyaline, generally containing two to three oil drops which stain brown with osmic acid. Basidia short, slender, 30–40 μ , arising from a well-developed stroma. On fruit of cultivated tomato.

A LEAF SPOT OF CELERY.

Not long since some celery leaves were brought into my laboratory by a local gardener with the complaint that his celery plants were suffering badly. To general appearance the affection seemed to be the ordinary leaf blight, (*Cercospora Apii*, Fres.), but closer observation showed the spots to be covered with numerous black pustules, which upon further examination proved to be the pycnidia of a *Septoria*.

Examinations were at once made in other celery gardens, and in all the same disease was found. The effect upon the plant is very similar to that of the well known leaf blight, causing in extreme cases a complete wilting of the leaves, and in the case of younger growths a complete destruction of the entire plant.

The spores, which germinate readily in water, were sown upon healthy leaves with the result that in fifteen days the characteristic spots were produced at the points of infection, upon which were developed pycnidia containing the spores of this fungus. The fungus of the disease is apparently distinct from *Septoria Pastinacæ*, West, and from *S. pastinacina*, Sacc., both found on *Pastinaca sativa*, L.; the former species having larger and septate spores, while the *Septoria* on celery is to all appearance non-septate. The latter species differs in having much smaller spores (20–30 μ ×7–1 μ) and larger pycnidia (120–150 μ).

The fungus, according to the opinion of Mr. J. B. Ellis, is more closely related to *Septoria Petroselini*, Desm., but is probably not identical with it.

Through Dr. D. H. Halsted I am informed that Briosi has named a variety of the last species, i. e. *S. Petroselini*, var. *Apii*.

which may be the form in question, but in the absence of specimens, the writer has no way of proving the point.

Whether the form in question is a new species, in which case it might be named *Septoria Apii*, or simply the above variety, is unimportant. The fact remains that the presence of this disease is noteworthy and it is therefore brought to the attention of mycologists.

The fungus is described as follows :

Spots amphigenous, white to tawny, irregular, becoming confluent, the entire leaf finally wilting. Pycnidia black, innate, scarcely erumpent, amphigenous, 74-100 μ in diameter, mostly 90 μ , globose-subglobose, loosely gregarious. Sporules hyaline, curved-flexuose, 25-40 $\mu \times$ 2-2.5 μ , apparently non-septate or septulate, eguttulate, one end commonly attenuate.

On leaves of cultivated celery.

BLIGHT OF WATERMELON VINES.

For some years past growers of watermelons in the southern part of Delaware have suffered badly from a disease which seems heretofore to have escaped the notice of mycologists.

During the present season a noted grower in this section experienced almost a complete failure of his crop through the same trouble. The malady was not brought to my notice until rather late in the season, and hence it was difficult to make the extended field observations which would be advisable.

Present observations have demonstrated, however, that the trouble is due to a parasitic fungus, most destructive in its habit, and capable of widespread financial disaster wherever it gains foothold. Young plants, 10 feet or less in length, attacked by the fungus, generally go down completely, while older plants suffer through the more or less complete destruction of their foliage and other tender parts of the vine, preventing completely the further development of the melon. Nor do the young fruit, up to the size of marbles, and fruit buds escape, such parts turning black, and finally developing the mature pycnidia of the fungus.

In short, the disease attacks leaf, leaf stalk, stem, tendrils, fruit buds and blossoms. It shows itself upon the leaves as black, circular or irregular spots, marked by concentric ridges. These spots increase in size and coalesce so as to cover a large

part or all of the leaf. On leaf stalks and tendrils as elongated dark lines, which increase in breadth so as to involve the part and cause it to turn black and shrivel. Similar elongated spots, which afterwards become white in the center, mark its stems, but in the case of these woody growths, the damage I have not found sufficient in the specimens examined to be serious, the injury being mostly confined to the tenderer parts of the vine.

The spores of the fungus, which germinated readily, were sown upon a young watermelon plant, grown from seed in the greenhouse; in three days the plant thus affected began to show signs of disease; in eighteen days the plant was completely dead. An examination of the blackened and wilted leaves showed the pycnidia of the fungus containing the characteristic and well developed sporules.

The check plant uninfected remained healthy, with no signs of the disease.

The fungus causing the above trouble is a member of the genus *Phyllosticta*, although from the character of the sporules, which are sometimes uniseptate and hyaline, it is questionable whether it might not, following Saccardo, be classed as an *Ascochyta*. It seems to differ from either *P. orbicularis*, E. and E., or *P. curbitraccarum*, Sacc., found on *Cucurbita Pepo*, L., and is here described as new.

Phyllosticta Citrullina, n. sp.

Spots circular, irregular, black, concentrically ridged, becoming confluent. Pycnidia amphigenous, brown, immersed, scarcely crumpled, membranaceous, lenticular 75-131 μ , average of many measurements 107 $\mu \times$ 67 μ . Sporules 9-10.7 μ , average about 10 $\mu \times$ 3.5 μ generally continuous, sometimes uniseptate, straight, slightly curved, ends obtuse, often biguttulate, hyaline.

On leaves and other parts of watermelon.

Delaware College, Newark, Del., Oct. 27, 1891.

No. 23, no cards.

Sonderabdruck aus der „Zeitschrift für Pflanzenkrankheiten.“

(Verlag von Eugen Ulmer in Stuttgart.)

Versuche über die Bekämpfung der Pilzkrankheiten mit Bordeauxmischung und Ammoniak-Kupferlösung,

ausgeführt auf der Jowa Versuchsstation (Ames. N. A.) im Jahre 1891
von

L. H. Pammel.

(Botanist. Jowa Agricultural Experiment Station.)

Es ist schon früher von mehreren Forschern nachgewiesen worden, dass die Bräune der Birnenblätter (Pear leaf blight), verursacht durch *Entomosporium maculatum*, sowie der Mehltau der Kirschen (*Podosphaera Oxyacanthae*), die Fleckenkrankheit der Kirschen (*Cylindrosporium Padi*) und die Fleckenkrankheiten der Johannisbeeren (*Cercospora angulata* und *Septoria Ribis*) mit obengenannten Kupfermitteln erfolgreich zu bekämpfen sind.

a.

Das *Entomosporium maculatum*¹⁾ verursacht in einzelnen Teilen, von Nord-Amerika sehr bedeutende Verluste; namentlich stark heimgesucht werden die Baumschulen. Im Juli und August sind manche Bäume schon gänzlich entlaubt. Bereits im Jahre 1889 hatte Galloway²⁾ gezeigt, dass die Bordelaiser-Mischung ein gutes Bekämpfungsmittel der Krankheit ist. Ich experimentierte 1890 mit dieser Mischung und der Ammoniak-Kupferlösung. Der Unterschied zwischen den unbeprengten und besprengten Pflanzen zu Gunsten der letzteren war ein so auffallender, dass ich in diesem Jahre den Versuch mit den nämlichen Pflanzen wieder aufnahm, aber nur die Ammoniak-Kupferlösung verwendete. Die Bespritzungen erfolgten am 8., 20., 27. Juni, am 9., 11., 18., 21. Juli und am 1. August. Das sehr zufriedenstellende Resultat ergibt sich aus folgender Tabelle.

1) *Entomosporium maculatum* Lév. wird in den deutschen Handbüchern vorzugsweise als *Morthiera Mespili* (DC.) Fekl. *Stigmatea Mespili* Sor. aufgeführt. Red.

2) Circular Nr. 8. Section of Vegetable Patholog. Departm. of Agriculture.

3) Bull. 13. Jowa Agricultural College, Experiment Station p. 43.

Besprengte Pflanzen				Nicht besprengte Pflanzen.		
Nr.	Höhe in engl. Zoll.	Zweige.	Blattzahl.	Höhe in engl. Zoll.	Zweige.	Blattzahl.
1	49	11	786	60	15	197
2	43	12	356	24	3	0
3	30	7	275	43	4	71
4	25	10	180	27	9	276
5	52	17	480	23	3	6
6	25	6	151	15	6	55
7	27	6	76	25	11	33
8	38	12	203	55	9	5
9	83	13	533	53	12	200
10	35	7	110	24	2	8
11	18	6	10	58	10	121
12	33	13	90	28	2	11
13	42	10	170	40	3	23
14	33	4	199	40	9	22
15	73	15	320	23	2	5
16	28	4	135	35	6	0
17	55	17	386	65	6	143
18	25	15	170	64	8	57
19	20	8	115	27	3	0
20	52	15	344	55	12	26
21	33	10	270	30	9	32
22	36	5	225	30	0	0
23	52	18	492	33	6	0
24	36	5	110	52	6	34
25	40	16	30	34	4	0
Summe:	980	252	6446	963	160	1325
Mittel:	39.2	10.4	257.8	38.5	6.4	5.3

b.

In Amerika tritt *Cylindrosporium Padi* Karst. ebenfalls besonders heftig in Baumschulen auf und nur mitunter auch auf grösseren Bäumen. Es wurden 3 Versuche mit Bordeaux-Mischung und Ammoniak-Kupferlösung unternommen.

Auf *Prunus Mahaleb* okulierte Kirschen in zweijährigen Exemplaren wurden am 6., 8., 20. und 29. Juni, am 8., 11., 18., 21. Juli und am 1. und 4. August bespritzt; die Krankheit begann am 6. Juni sich zu zeigen. Die Wirkung der Kupfermittel ist aus der Zahl der durch die Besprengung den Pflanzen erhalten gebliebenen Blätter ersichtlich; es ergaben nämlich 35 besprengte Bäumchen 4498 Blätter, während die nicht bespritzten nur 1195 behalten hatten. — Hierbei wurde auch bemerkt, dass die Mittel sehr wirksam gegen die Ausbreitung von *Podosphaera Oxyacanthae* sind. Nur wenige *Crataegus* Blätter zeigten den Mehltau, ausgenommen diejenigen, welche nach dem 4. August erschienen waren oder an niedrigen Pflänzchen sassen, welche von der Besprengung nicht erreicht wurden. — Da manche Kirschen auf *Prunus Mahaleb* oder andere Unterlage nicht okuliert werden können, muss man Sämlinge nehmen; aber gerade Sämlinge werden bisweilen durch

das *Cylindrosporium Padi* derart geschädigt, dass sie sehr wenig wachsen und oftmals zu Grunde gehen. Es wurden deshalb auch die Mittel bei diesjährigen Kirschsämlingen in Anwendung gebracht und es zeigte sich, dass im Durchschnitt (von 35 Pflanzen) die besprengte Pflanze 12 Blätter behielt, die unbesprengte dagegen nur 3. 4 Blätter.

c.

Gegen die *Cercospora angulata* auf Johannisbeeren wurden 7 Besprengungen mit Ammoniak-Kupferlösung und zwar am 6. 8. 20. 29. Juni, am 9. und 21. Juli und am 13. August ausgeführt. Während die mit der Kupferlösung behandelten Sträucher nur wenig kranke Blätter aufwiesen und bis 1. Oktober belaubt blieben, waren die daneben stehenden unbespritzten Sträucher bereits zu Anfang des August entblättert. —

Bei *Ribes nigrum* wird die Fleckenkrankheit hauptsächlich durch *Septoria Ribis* hervorgerufen und die Folge ist ebenfalls eine vorzeitige Entblätterung: nur ist der Verlust an Laub nicht so stark, wie bei *Ribes rubrum* und *R. aureum*, die vielleicht weniger widerstandsfähig sind. Immerhin zeigte sich bei der schwarzen Johannisbeere ebenfalls der günstige Einfluss des Kupferpräparates, indem die mit demselben behandelten Pflanzen fleckenärmer und blattreicher gegenüber den unbespritzten Exemplaren waren. Von letzteren besaßen 50 Zweige nur 1400, dagegen von ersteren 2100 Blätter.

New York Botanical Garden Library

3 5185 00215 7145

Made in Italy

10-09-857

8 0329 19 890070