ON

awkes.

TRANSFERRED TO
YALE MEDICAL LIBRARY

Digitized by the Internet Archive
in 2012 with funding from
Open Knowledge Commons and Yale University, Cushing/Whitney Medical Library

ORGANIC

MATERIA MEDICA,

INCLUDING THE STANDARD REMEDIES OF THE LEADING PHARMACOPŒIAS
AS WELL AS THOSE ARTICLES OF THE

NEWER MATERIA MEDICA

MORE RECENTLY BROUGHT BEFORE THE MEDICAL PROFESSION, WITH SHORT
NOTICES OF THEIR THERAPEUTICS AND DOSAGE COLLATED FROM THE
MOST RELIABLE SOURCES; AND OF THE PREPARATIONS MADE
THEREFROM BY

PARKE, DAVIS & CO.,

WITH WHICH IS INCORPORATED NUMEROUS FORMULÆ FOR THE EX-
TEMPORANEOUS PREPARATION OF TINCTURES, SYRUPS, OINTMENTS,
ETC., FROM THESE PRODUCTS. WITH AN APPENDIX CONTAIN-
ING A COMPLETE LIST OF THEIR CAPSULE, ELIXIR AND PILL
FORMULÆ, ETC., PHARMACEUTICAL SPECIALTIES, FINE
CHEMICALS AND ALKALOIDS; WITH TABLES OF
WEIGHTS AND MEASURES, POSOLOGY, ETC.

DETROIT, MICH., U. S. A.
1888.

Copyrighted by
GEORGE S. DAVIS.
1888.

RS154
888P

Gift of Dr. W. W. Newhall

PREFACE.

In the following pages are enumerated the crude drugs and such other articles of *materia medica* as are of organic nature and which at the same time are marketed by us either in the form of the crude article, or preparations therefrom, or both. Attached to the name of each is a brief statement of its reputed medicinal properties and its dose, together with a descriptive list of the derivative pharmaceutical preparations and compounds into which it enters as a therapeutic factor. At a glance can be thus seen almost all the methods or combinations known to modern pharmacy by which the exhibition of any desired remedy is made available. So far as is practicable, however, decoctions and infusions have been discarded since they are universally acknowledged to be obsolete. Some preparations indicated are not listed by us, but in these cases the formulæ are given by which they may be prepared extemporaneously with others which are listed. A complete and classified list of all our preparations and pharmaceutical specialties will be found in the appendix.

As Parke, Davis & Company, Manufacturing Pharmacists, we claim that all products of our Laboratory, chemical and pharmaceutical, are prepared in the most conscientious manner by the latest and most approved processes, and from most carefully selected drugs.

Of the various classes of pharmaceutical preparations mentioned special notes on fluid extracts, normal liquids and concentrations are alone considered necessary—these will be found below.

FLUID EXTRACTS—There is an urgent and general demand that the relative strengths of the fluid preparations of pharmacy be simplified and that many of the dilute ones be dropped altogether. Tinctures, infusions, wines, syrups, etc., have no uniform and well-understood ratio of strength either dosimetric or pharmacometric, and the recollection of dose, etc., is burdensome to the memory. The use of fluid extracts does away with this inconvenience. Having a strength so adjusted that each minim represents a grain, all that remains for a prescriber to know is the dose of the standard crude drug. Tinctures, wines, etc., may be thereupon made by simply using the proper diluent and to the amount necessary to exhibit a desired therapeutic effect, or to extemporaneously prepare a weaker preparation. This method, if generally adopted, would much simplify pharmacy and also—a much more important matter—relieve the physician and pharmacist of the burden of carrying so many different preparations of the same drug; and, by retaining one only and that the most accurate and scientific one, the probability of dispensing stale and deteriorated fluids would be at a minimum. We cannot here, however, do more than merely point out the economic and scientific advantages to be gained from a more extended use of this class of preparations.

It is occasionally a matter of some importance with physician and druggist to know the behavior of certain fluid extracts on mixing them with water—the fluid which is generally used as a diluent. When an option is allowed between the use of

two or more, if other things be equal, that one will preferably be taken which will make the most elegant compound. For the convenience and reference of prescribers and dispensers there is attached to the name of each fluid extract mentioned in the following pages an index of its miscibility with water. There are four different grades of classification indicated, namely: 1. Miscible with water; 2. Miscible with water without material precipitation; 3. Not miscible with water without some precipitation; 4. Not miscible with water.

NORMAL LIQUIDS.—The U. S. Pharmacopœia has omitted to provide for a scientific standard in the determination of the strength of the official fluid extracts. It is not sufficient that, prepared by its directions, one cubic centimeter of any fluid extract should represent one gramme of the drug employed. A crude drug, supposed to be of prime quality judged by its physical appearance, may be found to be either much above the average or notoriously inert when tested for the presence of its active principle, and since no test for the quality of the drug in this respect is demanded, it follows that the therapeutic exhibition of the average fluid extract must be accompanied with a similar uncertainty and variation in effects.

In this connection we particularly emphasize the fact that by the introduction of our normal liquids the superior standard of our fluid extracts is in no wise affected. As heretofore, they are prepared from full quantities and of the best materials, and manufactured with the utmost care and accuracy. Indeed, as an extra precaution over and above the requirements of the U. S. P., we test all crude drugs by assay wherever practicable. Whatever of discrimination is asked in favor of normal liquids is based solely upon the omission of the Pharmacopœia to recognize the fact that the only true and scientific test of a fluid extract is the presence of a standard amount of active principle actually ascertained in the finished product,—strength should not alone be determined, as at present, by the quantity of drug employed. Normal liquids were introduced by us to satisfy the demand for fluid extracts that could be depended upon to represent a definite and uniform alkaloidal strength and to possess a definite and invariable basis of therapeutic value.

CONCENTRATIONS.—These had their origin in the demand on the part of physicians for vegetable remedies of concentrated strength, in the form of permanent powders, in order that a well assorted supply might be carried in the pocket or saddle case. This class of preparations then was naturally introduced to notice by manufacturers, and their use extended rapidly among the profession who were not slow to perceive their unquestionable merits.

The nomenclature of concentrations, so-called, has been formed by adding the affix "*in*" to the root of the botanical name of the plant from which it is derived, thus conveying the impression that the preparation is the active principle of that plant, since chemists had already adopted this method of designating principles peculiar to any plant or genus of plants. Chemists now also distinguish further between alkaloids and neutral principles (glucosides, etc.) by adding in the former case a final *e* to the termination.

The names which have been adopted for the concentrations are, therefore, either wholly inappropriate and misleading, or they have a dangerous similarity to those of the active principles themselves of the same drugs, of which they actually contain only 5 to 10 per cent. We do not hold ourselves responsible for the present

PREFACE.

nomenclature, and a radical change therein can only be made by concerted action between the manufacturers and the medical profession. We have ventured in some cases to adopt certain innovations, not for the sake of distinguishing preparations of our own manufacture from others in the market, but simply as a safeguard against serious mistakes in dispensing. Thus we label the concentrations from aconite and belladonna, *ACONAPELLIN*, *not aconitin*, and *BELLADONNIN*, instead of *atropin*, since these latter in the ordinary nomenclature might easily be taken for the alkaloids aconite and atropine, articles at least ten times as powerful. We do not claim for these names any exclusive proprietorship.

The advantages offered by concentrations over other preparations of vegetable drugs are sufficiently obvious. They are, briefly: 1st. Concentration of strength; 2d. The form of a permanent powder; 3d. Convenience for dispensing at the bedside of the patient.

Valid objections are urged against them on the ground of their unscientific nomenclature and of the lack of any authoritative standard to which even their physical properties must conform. Preparations from different manufacturers bearing the same label are totally unlike in properties and composition, each one being a law unto himself. We have aimed in our concentrations to present all the active constituents of each drug, excluding as far as possible, inert matter; and when our preparations differ from those of other manufacturers, we are willing that they should be judged by their therapeutic efficiency.

Connected with our Laboratory is a Scientific Department of experts in the various branches of the science of pharmacology the members of which are devoting a life-time to the careful study of the physical, chemical and therapeutic properties of drugs. The object of this department is also to interest men of science abroad in the investigation of new therapeutic agents, for which we have frequently donated the material for experimentation, in order that they might be introduced on a scientific basis with as full and correct physiological reports as possible, for the prescriber's guidance. We can point with satisfaction to the results as published in reports on investigations made with our assistance by such men as Professors Dragendorff, Podwissotzki, Gerhardt, Schatz, Liebreich, Langgaard, Senator, Kobert, etc., and to the numerous drugs of value which we have been instrumental in introducing, such as Black-haw, Cascara sagrada, Kava-kava, Grindelia robusta, Pichi, Yerba santa, etc. In the compilation of this book the dosage of our preparations, botanical nomenclature employed, and other information contained have received special and careful revision by our Scientific Department and we trust, in presenting this to our friends, to have rendered them a service by placing in their hands a book of reference to which they may frequently and confidently refer for information.

A large, elegant handwritten signature in cursive script that reads "Parke, Davis & Co." The signature is written in dark ink and occupies a significant portion of the lower half of the page.

DETROIT, MICH., U. S. A., MARCH 1, 1888.

ORGANIC MATERIA MEDICA.

NOTES.

Any article sought for under a synonym or alphabetical arrangement other than that in which it is hereinafter described may be at once located by referring to General Index.

Authorities are quoted with reference to many of the Newer Materia Medica in order that the value of the statements given may be determined by the professional eminence of the originators.

A complete and numbered list of Capsule, Elixir and Pill Formulæ, and to which frequent reference is made in the following pages, will be found in appendix.

ABSCESS ROOT. *Polemonium reptans*, *Linné*. (Root)

Synonyms.—American Greek Valerian, False Jacob's Ladder, Sweat-root.

Properties.—Alterative, astringent, diaphoretic and expectorant. The fluid extract taken in warm water will produce copious perspiration; it will be found serviceable in pleurisy, bronchial and laryngeal affections, also in febrile and inflammatory diseases.

Abscess Root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Abscess Root. Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

ACONITE. *Aconitum*, U.S.P. (rt.) *Aconitum Napellus*, *Linné*. (Lvs: rt.)

Synonyms.—Monkshood, Wolfsbane; Aconit, Coqueluchon, Fr.; Eisenhut, Sturmhut, Ger.

Properties.—POISONOUS. A powerful nervous and (indirectly) arterial sedative. Employed in fevers as a diaphoretic, to reduce temperature, and to decrease the heart's action. These effects are best secured by minute doses. It is also employed with advantage in cardiac hypertrophy, neuralgia, rheumatism, gout, etc.

ANTIDOTES—None reliable known. Use strong emetics or the stomach-

ACONITE—Continued.

pump, promptly. Sustain vitality by stimulants—whisky, ammonia, strong coffee, the battery, etc. Digitalis has been found useful as an antidote.

N. B.—Aconite leaves are no longer recognized by the U. S. P. Physicians should specify the root in prescribing, to prevent mistakes, the root being five times more active than the leaves.

Aconite Leaves, compressed in ounce packages for retailing purposes.

Aconite Root, U. S. P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLES.

Aconitine, pure alkaloid (amorphous).

Dose, 1-500 to 1-100 grain (.00013 to .00065 Gm.).

Aconitine, pure alkaloid (crystals, Duquesnel's). A specific in facial neuralgia. Solubility, 1 part dissolves in 150 of cold water; sufficient to admit of its use hypodermically. 1-100 grain is equal to about 1 grain Aconite root.

Dose, 1-500 to 1-100 (.00013 to .00065 Gm.).

Napelline.—An alkaloid obtained from Aconitum Napellus, more soluble in water than Aconitine, but its physiological and therapeutic action are not yet sufficiently differentiated from those of Aconitine; claimed however to possess hypnotic as well as analgesic properties.*

Dose, 1-100 grain (.0006 Gm.), cautiously increased.

PREPARATIONS.

Abstract Aconite, U. S. P. See below for formula.

Dose, 1-6 to 1 grain (0.01 to 0.065 Gm.), 1 part equals 2 of root.

Aconapellin, a so-called **Concentration†**—a powdered preparation from aconite root.

Synonym.—Aconitin. Dose, 1-25 to 1-12 grain (0.003 to 0.006 Gm.)

Colloidion, Aconite.—Each fl. oz. represents 15 grs. Aconite root.

Extracts.

Fluid Extract Aconite Leaves.

Dose, 2 to 5 minims (0.13 to 0.33 C. c.), diluted.

Fluid Extract Aconite, U. S. P. (root.)

Dose, 1-3 to 2 minims (0.02 to 0.13 C. c.)

{ Not miscible with water without some precipitation.

Powdered Extract Aconite Leaves.—Made of a uniform strength by assay. Prepared by evaporating at a low temperature the solid extract, and replacing the moisture with powdered aconite leaves.

Dose, 1-4 to 1 grain (0.016 to 0.065 Gm.)

Powdered Extract Aconite Root.—1 part equals 5 of root, made of a uniform strength by assay, and equal to solid extract aconite root. Prepared by evaporating at a low temperature the solid extract, and replacing the moisture with powdered aconite root.

Dose, 1-20 to 1-5 grain (0.003 to 0.013 Gm.)

Solid Extract Aconite Leaves.—1 part equals 5 of the leaves, made of a uniform strength by assay.

Dose, 1-4 to 1 grain (0.016 to 0.064 Gm.)

Solid Extract Aconite, U. S. P. (root).—1 part equals 5 of the root, made of a uniform strength by assay.

Dose, 1-20 to 1-5 grain (0.003 to 0.013).

Hypodermic Tablet. Aconitine (crystals, DUQUESNEL'S), 1-120 gr.

A safe and ready means of exhibiting this powerful alkaloid in subcutaneous injection.

Liniment, Aconite, B. P. See below for formula.

Normal Liquid* Aconite Root.

Standard; contains 0.75 % alkaloid estimated by Mayer's reagent and Squibb's physiological test; made of a uniform strength by assay.

Dose, 1-3 to 2 minims (0.02 to 0.13 C. c.)

Ointments.

Aconite, P., D. & Co.'s. (1 part solid extract aconite root in 8.)

Aconitine, B. P. See below for formula.

* Dujardin-Beaumetz, "New Medications," p. 274. (edition of Geo. S. Davis, 1886).

† See note in Preface on Concentrations.

ACONITE—Continued.

Oleate Aconitine (in solution) containing 2% of the alkaloid.
Advantageously employed where the local effect of aconite is desired;
more cleanly in exhibition than the ointment.

Pills.

Aconite Extract (leaves), 1-4, 1-2, 1 grain.

Aconite Root, 1-20 grain.

Aconitine (crystals, Duquesnel's), 1-500, 1-200 grain.

Neuralgie, Dr. Gross', 2 1-2 grains.

Quinine sulph., 2 gr.

Ext. Aconite leaves, 1-2 gr.

Morphine sulph., 1-20 gr.

Strychnine, 1-30 gr.

Arsenious Acid, 1-20 gr.

Neuralgie Idiopathic, Brown-Sequard's, 3 1-2 grains.

Ext. Henbane, 2-3 gr.

Ext. Conium seed, 2-3 gr.

Ext. Ignatia, 1-2 gr.

Ext. Opium, 1-2 gr.

Ext. Aconite leaves, 1-3 gr.

Ext. Cannabis Indica, 1-4 gr.

Ext. Stramonium seed, 1-5 gr.

Ext. Belladonna, 1-6 gr.

Consult numbers 280, 282, 301, 479, in appended Pill Formulæ list
for other pills containing Aconite.

Plaster Aconite. In half-pound rolls or cans.

This may be made extemporaneously. See below for formula.

Tablet Triturates. **Aconite** (fluid extract), 1-10 and 1 minim.

Aconitine (crystals, Duquesnel's), 1-500 and 1-200 grain.

Tinctures.

Aconite, B. P. (root). See below for formula.

Dose, 5 to 15 minims (0.3 to 0.92 C. c.)

Aconite, Fleming's (root). See below for formula.

Dose, 1-2 to 3 minims (0.03 to 0.18 C. c.)

Aconite "German Tincture,"—We import this tincture of the
green plant; it is not the official Ger. Pharmacopœial preparation,
however; for the latter see below, Aconite, P. G.

Dose, 10 to 35 minims (0.6 to 2.16 C. c.)

Aconite Leaves, U. S. P. 1870. See below for formula.

Dose, 10 to 40 minims (0.65 to 2.6 C. c.)

Aconite, P. G. (root). See below for formula.

Dose, 7 to 18 minims (0.42 to 1.12 C. c.)

Aconite, U. S. P. (root). See below for formula.

Dose, 1 to 6 minims (0.06 to 0.4 C. c.)

FORMULÆ.**Abstract Aconite, U. S. P.**

Normal liquid aconite root, P., D. & Co.'s.....1 fl. oz.

Sugar of milk, q. s. to make.....228 grains.

Evaporate the fluid extract nearly to dryness by a very gentle heat;
add 175 grains of the sugar of milk; incorporate thoroughly with the ex-
tract; reduce to complete dryness; weigh and add sufficient sugar of
milk to bring the weight to 228 grains; finally powder the product. Keep
in a well corked bottle.

Liniment Aconite, B. P.

Fluid extract aconite root, P., D. & Co.'s.....20 fl. ozs.

Camphor.....1 oz.

Ointment Aconitine, B. P.

Aconitine.....8 grains.

Alcohol.....1-2 fluidrachm.

Benzoated lard.....1 oz. av.

Dissolve the Aconitine in the alcohol and mix solution thoroughly with
the lard.

Plaster Aconite.

Solid extract aconite root, P., D. & Co.'s.....2 ozs.

Resin plaster.....14 ozs.

Thin the extract with sufficient alcohol, then add to plaster, (previ-
ously melted) mix intimately, and stir till cold.

* See note in Preface on Normal Liquids.

ACONITE—Continued.**Tinctures.****Aconite, B. P.**

Fluid extract aconite root, P., D. & Co.'s1 fl. oz.
 Alcohol.....7 fl. ozs.

Aconite, Fleming's.

Fluid extract aconite root, P., D. & Co.'s.....2 fl. ozs.
 Alcohol.....1 fl. oz.

Aconite Leaves, U. S. P. 1870.

Fluid extract aconite leaves, P., D. & Co.'s.....2 fl. ozs.
 Dilute alcohol.....14 fl. ozs.

Aconite, P. G.

Fluid extract aconite root, P., D. & Co.'s.....1 fl. oz.
 Dilute Alcohol.....9½ fl. ozs.

Aconite, U. S. P. (root.)

Fluid extract aconite root, P., D. & Co.'s.....6 fl. ozs.
 Alcohol.....11½ fl. ozs.

ADDER'S TONGUE. *Erythronium americanum, Smith, (Leaves).*

Synonyms.—Dog's Tooth Violet, Yellow Snake Leaf; Dent de chien jaune, Fr.; Amerikanischer Hundszahn, Ger.

Properties.—Alterative and in large doses emetic. Useful as a poultice to inflamed scrofulous tumors; and internally to remove impurities of the blood.

Adder's Tongue Leaves, compressed in ounce packages for retailing purposes.

ADONIS VERNALIS, Linné, (Herb.)

Synonym.—False Hellebore.

Properties.—Poisonous. It owes its medicinal activity to the glucoside, adonidin, the physiological action of which seems to be identical with that of digitalin. Like digitalis, it is employed in dropsy and heart disease. A recent study of its therapeutic properties, made in the wards of Prof. Botkin, at St. Petersburg, shows that under its use the heart's action becomes slower, more regular, and more forcible, the urinary secretion being at the same time greatly increased. Dr. Bubnow, who reports these experiments, believes that Adonis is positively preferable to digitalis in organic heart disease, and he finds that it is not cumulative in action. **ANTIDOTES.**—Emetics if sufficiently early. Alcoholic stimulants, opium, rest in recumbent position.

ACTIVE PRINCIPLE.

Adonidin. Resembles Digitalin and employed almost exclusively for its action upon the heart.

Dose, 1-10 to 1-3 grain (.006 to .021 Gm.)*

PREPARATIONS.**Fluid Extract Adonis Vernalis.**

Miscible with water without material precipitation.

Dose, 1 to 2 minims (0.065 to 0.13 C. c.), cautiously increased.

Pill Adonidin, 1-10 grain.

A convenient and exact means of exhibiting Adonis.

ADRUE. *Cyperus articulatus, Linné, (Rhizome.).*

Synonym.—Anti-emetic root.

Properties.—"Adrue is of surprising worth in checking emesis, be it the vom-

* Durand, Bulletin General de Therapeutique, January 30, 1885.

ADRUÉ—Continued.

iting of pregnancy, of yellow fever, or indigestion."* It has a fine aromatic flavor and produces a general feeling of warmth and comfort in the stomach. In atonic dyspepsia it will be found a pleasant and efficient remedy.

PREPARATION.**Fluid Extract Adué.**

Miscible with water without material precipitation.
Dose, 30 minims (2 C. c.)

AGRIMONY. *Agrimonia Eupatoria, Linné.* (Herb.)

Synonyms.—Cocklebur, Stickwort; Odermennig, Leberklette, Ger.

Properties.—Mild corroborant and astringent. Recommended in bronchial disorders, bowel complaints—especially chronic diarrhoeas, and in passive hæmorrhages. Prescribed in leucorrhœa and catarrhal diseases of the mucous membranes. Employed as a gargle in ulcerations of the mouth and throat.

Agrimony herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Agrimony.—Not miscible with water.
Dose, from 1-2 to 2 drachms (2 to 8 C. c.)

AILANTUS GLANDULOSA, Desfontain. (Bark.)

Synonyms.—Tree of Heaven, Chinese Sumach; Götterbaum, Ger.

Properties.—A powerful nervous depressant; purgative, anthelmintic and neurotic. Used in palpitation of the heart, obstinate hiccough, spasmodic asthma, twitching of the muscles, and epilepsy. Given in small doses it is "the Chinese remedy for diarrhoea" and dysentery.† In Europe it has been used to advantage as an anthelmintic, especial for tape worm.

PREPARATION.**Fluid Extract Ailantus Glandulosa.**

Not miscible with water without some precipitation.
Dose, from 10 to 30 minims (0.6 to 2 C. c.)

AJOWAN. *Ptychotis Ajwan, DeCandolle.* (Seeds.)

Synonyms.—Bishop's Weed, Ajava; Ajwain, Hind.

Properties.—Carminative, antiseptic. Contains much thymol, to which it probably owes its medical properties. Dr. Waring regards this, of all umbelliferous seeds, as the most efficient carminative, and bespeaks for it a trial in cholera, in which, it is alleged, even in East Indian empirical practice it has done wonders. Strongly recommended as a remedy for dipsomania or alcoholism. Applied externally, as a constituent of poultices, it relieves pain.

PREPARATION.

Fluid Extract Ajowan.—Not miscible with water.
Dose, 10 to 30 minims (0.6 to 2 C. c.)

ALKANET. *Alkanna tinctoria, Tausch.* (Root.)

Properties.—Astringent, but now used exclusively for coloring purposes.

ACTIVE OR COLORING PRINCIPLE.

Alkannin, a convenient preparation for coloring tinctures, ointments, oils etc. Insoluble in water.

* Morse, "New Therapeutical Agents," p. 187.

† Dolan, "New Remedies," p. 15.

ALLIGATOR PEAR. *Persea gratissima. Gartner. (Seeds.)*

Synonyms.—Aguacate, Avocado Pear.

Properties.—Recommended highly by the residents of Southern Mexico, in intercostal neuralgia and as an anthelmintic: In *Intercostal Neuralgia*.—Rub the chest and back with a dry towel until a good superficial circulation is obtained, then apply from 1 1-2 to 2 ounces of the fluid extract, using gentle friction, and cover the part with a dry flannel to protect the clothing from stain.

Great anthelmintic properties are claimed for the seeds by the natives of Tehuantepec. A case is reported by Dr. Henry Frøehling, of Baltimore, in which a tapeworm with head attached, was expelled from a man within five hours. Give three separate doses, of a fluid drachm each, in space of one hour, and follow with a dose of castor oil.

PREPARATION.**Fluid Extract Alligator Pear Seeds.**

Not miscible with water without some precipitation.
Dose, 30 to 60 minims (2 to 4 C. c.)

ALOES. Aloe, U. S. P. Inspissated juice of Aloe Socotrina. *Lamarck.*

Synonyms.—Socotrine Aloes, Aloés, Fr.

Properties.—Tonic, purgative, emmenagogue, anthelmintic, and cholagogue. Useful in constipation, combining tonic with laxative action; habitual use does not sensibly diminish its activity. Useful in jaundice from hepatic torpor, in amenorrhœa, in menorrhagia arising from debility, etc. It has a special tendency in its action to the lower portion of the intestinal canal.

NOTE.—A very much larger dose of Aloes can be given in solution than in the solid form.

ACTIVE PRINCIPLE.**Aloin.**

Twice the strength of a fair sample of Aloes.
Dose, 1-2 to 2 grains (0.03 to 0.13 Grm.)

PREPARATIONS AND FORMULÆ.**Extracts.**—**Fluid Aloes.**

16 fluidounces imperial represent 5 1-4 ozs. av. of Socotrine Aloes.
Not miscible with water; incompatible with acid solutions.
Dose, 3 to 20 minims (0.2 to 1.3 C. c.)

Fluid Aloes and Myrrh.—One pint represents 6 2-5 ozs. av. each of Aloes and Myrrh; 4 times stronger than Tincture of Aloes and Myrrh, U. S. P., for the extemporaneous preparation of which it is specially adapted.

Not miscible with water; incompatible with acid solutions.
Dose, 15 to 30 minims (1 to 2 C. c.)

Powdered Extract Aloes.—Of the same strength as the recent extract; produced by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Sugar of Milk.
Dose, from 1 to 5 grains (.06 to .3 Gm.)

Solid Extract Aloes, U. S. P.

Dose, from 1 to 5 grains (.06 to .3 Gm.)

Pills.**Aloes U. S. P., 4 gr.**

Purified Aloes, 2 gr.

Soap, 2 gr.

Aloes and Asafetida, U. S. P., 4 gr.

Purified Aloes, 1 1-3 gr.

Soap, 1 1-3 gr.

Asafetida, 1 1-3 gr.

Aloes and Iron,

Purified Aloes, 1 gr.

Iron sulph., exsic., 1 gr.

Jamaica Ginger, 1-2 gr.

Ext. Conium seed, 1-2 gr.

ALOES—Continued.

- Aloes and Mastich, U. S. P.**
 Purified Aloes, 2 gr. Powd. Red Rose, 1-2 gr.
 Mastich, 1-2 gr.
- Aloes and Myrrh, U. S. P.,** 3 1-2 gr.
 Purified Aloes, 2 gr. Myrrh, 1 gr.
 Aromatic Powder, 1 2 gr.
- Aloes and Nux Vomica, 2 gr**
 Purified Aloes, 1 1-2 gr. Ext. Nux Vomica, 1-2 gr.
- Aloes Nux Vomica and Belladonna, 2 1-4 gr.**
 Purified Aloes, 1 1-2 gr. Ext. Nux Vomica, 1-2 gr.
 Ext. Belladonna, 1-8 gr.
- Aloin, 1-10, 1-5, 1-4, 1-2, 1 gr.**
- Aloin Co., 1-2 gr.**
 Aloin, 1-8 gr. Ext. Belladonna, 1-4 gr.
 Podophyllin, 1-8 gr.
- Aloin Comp. and Strychnine 1-2 gr.**
 Aloin, 1-8 gr. Strychnine, 1-80 gr.
 Podophyllin, 1-8 gr. Oleoresin Capsicum, 1-10 gr.
 Ext. Belladonna, 1-8 gr.
- Aloin and Strychnine, 1-4 gr.**
 Aloin, 1-5 gr. Strychnine, 1-60 gr.
- Aloin Strychnine and Belladonna, No. 1.**
 Aloin, 1-5 gr. Strychnine, 1-60 gr.
 Ext. Belladonna, 1-8 gr.
- Aloin Strychnine and Belladonna Comp.**
 Aloin, 1-5 gr. Strychnine, 1-60 gr.
 Ext. Belladonna, 1-8 gr. Ext. Rhamnus Purshiana, 1-2 gr.
- Eccoprotic, 2 1-2 gr.**
 Ext. Soc Aloes, 2 gr. Podophyllin, 3-10 gr.
 Ext. Nux vomica, 1-5 gr. Oil Cloves, 1-10 gr.
- Lapactic, 12 gr.**
 Aloin, 1-4 gr. Ext. Belladonna, 1-8 gr.
 Strychnine, 1-60 gr. Ipecac, 1-16 gr.
- The above represent the principal aloetic pills; for other combinations see appended Pill Formulae list for numbers 9, 18b, 27, 34, 35, 44, 45, 101, 105, 113, 125, 145, 166-168, 178-181, 186, 187, 194, 223, 254, 260, 261, 233, 234, 302-304, 335, 392, 393, 423, 430.
- Tablet Triturates.—Aloin, 1-10, 1-4 and 1-2 grain.**
- Tinctures.**
- Aloes and Canella (Hiera Picra.)**
 Fluid Aloes, P., D. & Co.'s 2 fl. ozs.
 Fluid Extract Canella, P., D. & Co.'s 1 fl. oz.
 Dilute Alcohol 13 fl. ozs.
 Dose, 1 to 2 fluidrachms (4 to 8 C. c.)
- Aloes and Myrrh, U. S. P. (Elixir proprietatis.)**
 Fluid Aloes and Myrrh, P., D. & Co.'s 4 fl. oz.
 Alcohol 12 fl. ozs.
 Dose, 1 to 2 fluidrachms (4 to 8 C. c.)
- Aloes U. S. P.**
 Fluid Aloes, P., D. & Co.'s 6 fl. ozs.
 Fluid Extract Licorice, P., D. & Co.'s 10 fl. ozs.
 Dilute Alcohol 5 fl. ozs.
 Dose, 10 to 60 minims (0.6 to 4 C. c.)
- Wine Aloes U. S. P.**
 Fluid Aloes, P., D. & Co.'s 2 1-2 fl. ozs.
 Fluid Extract Cardamon, P., D. & Co.'s 1 fl. drachm.
 Fluid Extract Ginger, P., D. & Co.'s 1 fl. drachm.
 Stronger White Wine 10 fl. ozs.
 Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.)

ALSTONIA CONSTRICTA, *F. Mueller.* (Bark.)

Synonym.—Australian fever bark.

Properties.—Tonic, antiperiodic. Useful in those cases of remittent fever which will not yield to the salts of Cinchona bark. It strengthens the heart's action.

ALSTONIA CONSTRICTA—Continued.**PREPARATION.****Fluid Extract *Alstonia Constricta*.**

Not miscible with water without some precipitation
Dose, from 2 to 8 minims (0.13 to 0.50 C. c.)

ALUM ROOT. *Heuchera americana*, *Linné*. (Root.)

Synonym.—Amerikanische Sanickel-Wurzel, Ger.

Properties.—A most powerful astringent. Useful, in small doses, in all cases where astringents are indicated. Externally, in hæmorrhages, epistaxis, wounds, foul and indolent ulcers, etc. As a gargle for aphthous sore mouth, and for sore throat. Internally, in diabetes and in bleeding piles; in the last complaint in the form of an injection.

PREPARATION.

Fluid Extract *Alum-root*.—Not miscible with water.

Dose, from 10 to 20 drops (0.65 to 1.3 C. c.)

ALVELOZ MILK. The milky juice of the plant *Euphorbia heterodoxa*. *Müller*.

Synonyms.—Aveloz, Arveloz.

Properties.—"Exhibits the combined action of a caustic and of papaine. In canceroid ulceration it promptly destroys the affected tissues, layer by layer."* A profuse suppuration follows its application to a granular surface. The action is irritating, producing a spreading dermatitis without much pain, and said to result in the destruction of the morbid tissue, which is replaced by healthy granulations, doing the work, in fact, of the chloride of zinc paste. "Destined to play a considerable rôle in the therapeutics of canceroid and syphilitic ulcers."†

AMERICAN BLUE VERVAIN. *Verbena hastata*, *Linné*.
(Herb, root.)

Synonyms.—Blue Vervain, Vervain, Wild Hyssop, Simpler's Joy.

Properties.—Tonic, emetic, expectorant and sudorific. The hot infusion is employed as a sudorific in colds, obstructed menstruation, etc.

American Blue Vervain herb, compressed in ounce packages for retailing purposes.

American Blue Vervain root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract *American Blue Vervain*.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

AMERICAN CENTAURY. *Sabbatia angularis*, *Pursh*. (Herb.)

Synonyms.—Centaury, Rose Pink, Bitter Clover, Wild Succory, Red Centaury.

Properties.—Long and popularly employed as a prophylactic and remedy in our autumnal intermittent and remittent fevers. Certain physicians express themselves strongly in favor of its use as a substitute for quinine in malarial diseases, especially remittents. Particularly applicable to that condition existing between the paroxysms when the remission is such

* New York Medical Times (Medical Age, 1885, p. 240).

† Dr. Clemente Ferreira, of Rezende, Brazil, in the Bulletin Général de Thérapeutique, Dec. 15, 1885.

AMERICAN CENTAURY—Continued.

as to call for tonics. Useful as a tonic in slow convalescence, and in dyspepsia and other diseases of debility.

American Centaury herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract American Centaury,—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

AMERICAN COLUMBO. *Frasera carolinensis*, *Walter*. (Root.)

Properties.—Tonic and slightly aperient. In overdoses purgative and emetic.

Useful in stomachic and general debility, and in all cases where a mild tonic is indicated, particularly in irritated conditions of the digestive organs.

American Columbo root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract American Columbo.

Miscible with water without material precipitation.

Dose, 20 drops to a fluidrachm (1.3 to 4 C. c.)

Fraserin, a so-called **Concentration**,* a powdered preparation from American Columbo root.

Dose, 1 to 3 grains (0.06 to 0.19 Gm.)

AMERICAN HELLEBORE. *Veratrum viride* (U. S. P.), *Aiton*. (Rhizome.)

Synonyms.—Green Hellebore, Swamp Hellebore, Itch Weed, Indian Poke; Veratre Vert., Fr.; Grüner Germer, Ger.

Properties.—Poisonous. The chief use of this agent is to depress the action of the heart and to lower the vaso-motor tonus. Very favorable results are claimed in the treatment of simple cardiac hypertrophy, irritable heart, aneurism, and in acute parenchymatous congestions; indicated in cerebral hyperæmia. The hypertrophy of the heart and the abdominal arterial tension which accompany the chronic form of albuminuria, are said to be alleviated by its use. When there are valvular lesions, and when the heart is enfeebled from any cause, this agent is inadmissible.—Bartholow.

ADMINISTRATION.—As the effect of the drug quickly reaches its maximum, the dose must not be at longer intervals than two hours. *The recumbent posture must be rigidly enforced*, and the drug should only be given by competent hands.

ANTIDOTE.—Morphine and laudanum, in full doses, given in strong coffee, and alcoholic stimulants.

American Hellebore Root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Ext. Amer. Hellebore, U.S.P.—Not miscible with water.

Dose, 1 drop (0.03 C. c.), gradually increased until the desired effect on the circulation is produced.

Solid Extract American Hellebore.—1 part equals 5 of root.

Dose, 1-8 to 1-4 grain (.008 to .016 Gm.)

Normal Liquid† American Hellebore.

Standard, 10 C. c. of the Normal Liquid require for complete precipitation of the alkaloids 3 C. c. of Mayer's solution.

Dose, one drop (0.03 Gm.) gradually increased if necessary.

Pills Veratrum Viride Extract, 1-4, 1-2 grain.

American Hellebore is also an ingredient of pills, numbers 260 and 261, see list of Pill Formulæ appended.

AMERICAN HELLEBORE—Continued.**Tincture, American Hellebore, U. S. P.**

Fluid Extract American Hellebore, P., D. & Co.'s.....10 fl. ozs.

Alcohol12 fl. ozs.

Dose, to commence, 1 minim (0.06 C. c.)

This preparation is sometimes known as Norwood's Tincture.

Veratrin, a so-called **Concentration**,* a powdered preparation from American Hellebore root.**Synonym.**—Veratrin (not to be confounded with Veratrine, the active principle of Cevadilla seed.)

Dose, 1-16 to 1-2 grain (0.004 to 0.032.)

AMERICAN IVY. *Ampelopsis quinquefolia*, Michaux. (Bk. of rt.)**Synonyms.**—Woodbine, Virginia Creeper, Five-leaves, False Grape; Vigne Vierge, Fr.; Amerikanischer Epheu, Ger.**Properties.**—Alterative, tonic, astringent and expectorant. It has been found very useful in dropsy, by promoting absorption. Recommended also in bronchitis and other pulmonary complaints.**American Ivy**, compressed in pound packages for retailing purposes.**PREPARATIONS.****Ampelopsin**, a so-called **Concentration**,* a powdered preparation from American Ivy.

Dose, 2 to 4 grains (.12 to .25 Gm.)

Fluid Extract American Ivy.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

AMERICAN SARSAPARILLA. *Aralia nudicaulis*, Linné. (Rhizome.)**Synonyms.**—False Sarsaparilla, Wild Sarsaparilla, Small Spikenard, Shot-bush; Aralia nude, Fr.; Nackte Aralienwurzel, Ger.**Properties.**—Alterative. Frequently substituted in practice for Sarsaparilla (*smilax officinalis*). It also acts as a gentle stimulant and diaphoretic.**American Sarsaparilla Root**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract American Sarsaparilla.**

Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.)

AMERICAN WORMSEED. *Chenopodium*, U. S. P. *Chenopodium ambrosioides*, Linné; var. *anthelminticum*, Gray. (Seed.)**Synonyms.**—Jerusalem Oak; Chenopode anthelmintique, Fr.; Amerikanischer Wurmsamen, Ger.**Properties.**—Anthelmintic and antispasmodic; one of the most efficient indigenous remedies for the expulsion of the round worms in children. A dose of it is usually given before breakfast in the morning, and at bed time in the evening, for three or four days successively and then followed by some brisk cathartic. Repeat this procedure if necessary.**PREPARATIONS AND FORMULA.****Capsule, Wormseed Oil**, in soft Gelatin.

Oil American Wormseed2 minims.

Olive Oil8 minims.

Fluid Extract American Wormseed.

Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.)

* See note in Preface on Concentrations.

† See note in Preface on Normal Liquids.

AMMONIAC. Ammoniacum, U. S. P. Gum-resin from Dorema Ammoniacum. *Don.*

Properties.—Antispasmodic, stimulant of circulation and nutrition, expectorant. Useful in chronic catarrh, bronchitic affections and asthma. Externally as a rubefacient in plasters for the relief of chronic rheumatism and where indicated for a superficial and sustained irritation of the skin.

PREPARATIONS.

Pills.

Ipecac and Squill, 3 gr.

Powd. Ipecac, co. 2 gr.

Ammoniac, 2-3 gr.

Powd. Squill, 2-3 gr.

Squill Comp. 3 gr.

Powd. Squill, 3-8 gr.

Powd. Soap, 1 1-8 gr.

Powd. Jamaica Ginger, 3-4 gr. Ammoniac, 3-4 gr.

Plasters. In half-pound rolls or cans.

Ammoniac, U. S. P.

Ammoniac with Mercury, U. S. P.

Applied as a discutient in syphilitic cases.

ANGELICA. { (American) *Archangelica atropurpurea*, *Hoffmann*. (Lvs., Seed.) (European) *Archangelica officinalis*, *Hoffmann*. (Root.)

Synonyms.—(Root.) Masterwort, Garden Angelica, Engelwurzel, Ger.; *Angelica archangelica*, *Linné*.

Synonyms.—(Leaves, Seed.) Masterwort, High Angelica.

Properties.—Aromatic, stimulant, diaphoretic and emmenagogue. Used in flatulent colic and is a popular remedy, in infusion, to promote menstruation. "Useful in the chronic bronchial catarrh of aged and feeble persons, under circumstances where Senega is usually prescribed." *

Angelica leaves, compressed in ounce packages for retailing purposes.

Angelica root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Extracts.

Fluid Extract Angelica root.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

Fluid Extract Angelica seed.—Not miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C. c.)

ANGUSTURA. *Cusparia trifoliata*, *Willdenow*. (Bark.)

Synonyms.—*Galipea Cusparia*, *St. Hillaire*; *Galipea officinalis*, *Hancock*; *Cusparia febrifuga*, *Humboldt*.

Properties.—A stimulant tonic, with febrifuge properties, in large doses emetic and cathartic. Said to be peculiarly efficacious in bilious diarrhoeas and dysenteries, and recommended in dyspepsia and other diseases requiring tonic treatment.

PREPARATION.

Fluid Extract Angustura, not miscible with water.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

ANISE. Anisum, U.S.P. *Pimpinella Anisum*, *Linné*. (Fruit.)

Synonyms.—*Anisum Vulgare*, *Clusius*; Aniseed, European Anise; Anis, Fr. and Ger.

Properties.—A grateful aromatic carminative. Said to have the property, in

* Napheys, "Modern Medical Therapeutics," p. 179.

ANISE—Continued.

warm infusion, of increasing the flow of milk. Used in flatulency, flatulent colic of infants, and to remove nausea.

PREPARATION.

Fluid Extract Anise.—Not miscible with water.

Dose, 10 to 20 minims (0.6 to 1.3 C. c.)

NOTE.—It enters into several preparations as a flavoring agent merely.

APPLE-TREE. *Pyrus malus*, *Linné*. (Bark.)

Synonym.—Apfelbaumrinde, Ger.

Properties.—Tonic and febrifuge; used with advantage in intermittent, remittent and bilious fevers.

Apple-tree bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Apple-tree bark.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

ARBOR VITÆ. *Thuja* (U.S.P.). *Thuja occidentalis*, *Linné*. (Tops.)

Synonyms.—False white cedar, arbre de vie, Fr.; Lebensbaum, Ger.

Properties.—Expectorant, febrifuge, emmenagogue, anthelmintic, resembles savin in its general character. Recommended in intermittent and remittent fevers, cough, scurvy and rheumatism.

Arbor Vitæ leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Elixir Arbor Vitæ, each fl. oz. represents 120 grs.

Dose, 1 to 2 fluidrachms (4 to 8 G. c.)

Fluid Extract Arbor Vitæ.—Not miscible with water.

Dose, $\frac{1}{4}$ to 1 fluidrachm (1 to 4 C. c.)

Glycerole Arbor Vitæ.—Each fluidounce represents 120 grains.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

ARECA. *Areca Catechu*, *Linné*. (Seeds.)

Synonyms.—Betel Nut; Noix d'Arec, Fa.; Arekanisse, Ger.

Properties.—Astringent and vermifuge. Successfully used for the expulsion of tape worm in doses of from four to six drachms. *The Forest and Farm* recommends it highly as a vermifuge for young dogs, as both effective and safe—the dose being graded to two grains (or two minims of the fluid extract) to each pound of the animal's weight. Whether used as a vermifuge for man or animal, the bowels should be first cleansed by fasting and castor oil.

PREPARATION.

Fluid Extract Areca Nuts.—Not miscible with water.

Dose, as a vermifuge, 2 fluidrachms (8 C. c.) and upwards.

ARNICA. { *Arnicae flores*, U.S.P. } *Arnica montana*, *Linné*. (Flowers,
{ *Arnicae radix*, U.S.P. } Rhizome.)

Synonyms.—Leopard's Bane, Mountain Tobacco; Falkkraut, Ger.; Arnique, Fr.

Properties.—Stimulant; in large doses irritant. Has been employed as a remedy in typhoid and other adynamic fevers, and in dysentery which often presents a typhoid type.

Preparations of the flowers are employed externally for the relief of pain from bruises, sprains, etc., and to prevent or remove ecchymosis.

Arnica flowers, U.S.P., compressed in ounce packages for retailing purposes.

Arnica root, U.S.P., compressed in pound packages for retailing purposes.

ARNICA—Continued.

PREPARATIONS AND FORMULÆ.

Collodion Arnica.—Each fl. oz. represents 24 grains arnica root

Extracts.**Fluid Extract Arnica flowers.**

Miscible with water without material precipitation.

Dose, from 5 to 25 minims (0.3 to 1.5 C. c.)

Fluid Extract Arnica root, U.S.P.

Miscible with water without material precipitation.

Dose, 5 to 20 minims (.3 to 1.3 C. c.)

Solid Extract Arnica flowers.—One part equals 5 of the flowers.

Dose, 1 to 5 grains (0.065 to 0.33 Gm.)

Solid Extract Arnica root, U.S.P.—One part equals 5 of the root.

Recently made official in place of the extract of the flowers, which is more irritating.

Dose, 1 to 5 grains (.065 to .33 Gm.)

Infusion Arnica flowers.

Fluid Extract Arnica flowers, P. D. & Co.'s.....1 fl. oz.

Hot water4 fl. ozs.

This strength is considered the best for local use; preferably made from the flowers themselves.

Pill Arnica flowers, 1-5 grain.

Plaster Arnica, U.S.P.—In half pound rolls or cans. May be prepared extemporaneously as follows:

Solid Extract Arnica root, P., D. & Co.'s.....1 oz.

Resin plaster2 ozs.

Thin the extract with sufficient alcohol, then add to plaster (previously melted) mix intimately and stir till cold.

Soap, Arnica Medicated, contains 10 per cent. Extract Arnica. Useful in cleansing boils, carbuncles, and in many of the pustular skin affections. An elegant bath application for abrasions, wounds, and bruises of the integument.

Tinctures.**Arnica flowers, U.S.P.**

Fluid Extract Arnica flowers, P., D. & Co.'s.....3 fl. ozs.

Dilute Alcohol13 fl. ozs.

Dose, 10 to 30 minims (.6 to 2 C. c.)

NOTE.—Uniform in strength with that of the Fr. Cod., double that of the Pharm. Ger.

Arnica Root, "German Tincture."—An imported unofficial preparation of the green root.

Dose, 12 to 15 minims (.7 to 3 C. c.)

Arnica root, U.S.P.

Fluid Extract Arnica root, P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol19½ fl. ozs.

Dose, 20 to 60 minims (1.25 to 4 C. c.)

NOTE.—Double the strength of the B. P. tincture.

AROMATIC.

Fluid Extract Aromatic, U.S.P.—Not miscible with water.

Each pint represents 5¾ oz. each of cinnamon and ginger, and 2½ oz. each of cardamom and nutmeg; presents in fluid form the well-known aromatic powder. Carminative, and gently stimulant; employed for the relief of nausea, flatulence, colic or diarrhoea, and with other remedies as a corrigent and adjuvant.

Dose, 10 to 30 minims (0.65 to 2. C. c.)

Wine Aromatic, U.S.P.

One part in 100 of Lavender, Origanum, Peppermint, Rosemary, Sage, Wormwood

Only used externally—as a vulnerary to recent wounds, or to stimulate indolent ulcers. Almost similar to the Tincture Vulneraria (Alcoolature Vulnéraire), Fr. Cod., which has the same range of application. A preparation now seldom used.

ASAFETIDA, U.S.P. Gum-resin from *Ferula Narthex*, *Boissier*.

Properties.—A general and local tonic stimulant, carminative and antispasmodic. Useful in all hysterical affections, chorea, whooping cough; in chronic bronchitis, for the relief of tympanites and constipation, and as a soothing remedy for infants.

PREPARATIONS AND FORMULÆ.

Fluid Asafetida.—Not miscible with water.

Each pint represents 8 ounces of select drug.

Dose, 6 to 30 minims (0.4 to 2 C.c.).

Mixtures.**Asafetida, U.S.P.**

Fluid Asafetida, P., D. & Co.'s.....1 fl. oz.

Water.....11 fl. ozs.

Dose, $\frac{1}{2}$ fl. oz. (15 C.c.) and upwards.

Magnesia and Asafetida, U.S.P. DEWEE'S Carminative.

Fluid Asafetida, P., D. & Co.'s.....40 minims.

Tincture Opium.....14 minims.

Simple Syrup.....2½ fluidrachms.

Magnesium Carbonate.....70 grains.

Water sufficient to make.....5 fluidounces.

Dose, for an infant, 20 minims (1.3 C.c.).

Pills.**Aloes and Asafetida, U.S.P.**, 4 gr.

Purified Aloes, 1 1-3 gr.

Soap, 1 1-3 gr.

Asafetida, 1 1-3 gr.

Asafetida, 2, 3, 4, and 5 gr.**Asafetida Comp.**, 2 1-2 gr.

Opium, 4-5 gr.

Asafetida, 4-5 gr.

Ammonium carb., 4-5 gr.

Asafetida and Iron, 3 gr.

Asafetida, 2 gr.

Iron sulph. exsic., 1 gr.

Asafetida and Nux Vom., 3 1-4 gr.

Asafetida, 3 gr.

Ext. Nux vomica, 1-4 gr.

Asafetida and Rhubarb, 3 Gr.

Asafetida, 1 gr.

Iron by hydrogen, 1 gr.

Powd. Rhubarb, 1 gr.

Galbanum Co., U.S.P., 3 1-2 gr.

Galbanum, 1 1-2 gr.

Myrrh, 1 1-2 gr.

Asafetida, 1-2 gr.

Syrup Asafetida.

Fluid Asafetida, P., D. & Co.'s.....5 fluidrachms.

Simple Syrup, q. s. to make.....8 fluidrachms.

Dose, for an infant, $\frac{1}{2}$ to 1 teaspoonful (2 to 4 C.c.).

Tincture Asafetida, U.S.P.

Fluid Asafetida, P., D. & Co.'s.....6 fluidounces.

Alcohol.....11 fluidounces

Dose, $\frac{1}{2}$ to 1 fluidrachm (2 to 4 C.c.).

NOTE.—Uniform in strength with those of the Pharm. Ger. and Fr. Cod.

ASPARAGUS. *Asparagus officinalis*, *Linné*. (Rhizome.)

Synonyms.—Asperge, Fr.; Spargel, Ger.

Properties.—Diuretic, aperient, deobstruent and slightly diaphoretic.

"Asparagus is undeniably a cardiac sedative and is palliative in all heart diseases attended with severe action."* In overdoses it is liable to irritate the mucous membrane of the urinary passages, but useful, however, as an adjunct to other diuretics.

Asparagus root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Asparagus.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

* Morse, "New Therapeutical Agents," p. 60

BAEL. *Ægle marmelos*, *Correa de Serra*. (Fruit.)

Synonyms.—Bengal quince, *Crataeva marmelos*, *Linné*.

Properties.—Claimed to be almost a specific in chronic diarrhœa. It has remarkable astringent properties which render it useful in diarrhœa, dysentery with debility of the mucous membrane, and other diseases of the bowels with relaxation, all of which it relieves without constipation. It is much used in India.

PREPARATION.

Fluid Extract Bael Fruit.—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.)

BALMONY. *Chelone glabra*, *Linné*. (Leaves.)

Synonyms.—Turtle Head, Turtle Bloom, Snake Head, Shell-flower, Salt-Rheum Weed.

Properties.—Tonic, cathartic and anthelmintic. Useful in the treatment of jaundice, hepatic disorders, debility of the digestive organs, and during convalescence from febrile and inflammatory diseases. Popularly used as an application to painful and inflamed tumors, inflamed breasts, piles, etc.

Balmoney leaves. compressed in ounce packages for retailing purposes.

PREPARATIONS.

Chelonin, a so-called **Concentration**.*—A powdered preparation from Balmoney leaves.

Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Fluid Extract Balmoney.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

BALSAM FIR. *Abies balsamea*, *Marshall*. (Bark.)

Synonym.—Balsam Spruce.

Balsam of Fir or Canada balsam, oleoresin obtained from same.

Properties.—The bark is stimulant, diuretic and anthelmintic, in large doses laxative. From its direct stimulating action on the mucous tissues, it has been recommended in gonorrhœa, gleet, chronic inflammation of the bladder, etc. The oleoresin is used successfully in the treatment of chronic bronchitis.

PREPARATIONS.

Capsules, Balsam of Fir, 10 minims. In soft gelatin.

Fluid Extract Balsam Fir bark.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

BALSAM OF PERU. *Balsamum Peruvianum*, U. S. P. Obtained from *Myroxylon Pereiræ*, *Klotzsch*.

Properties.—Stimulant to the circulation with special tendency to mucous membranes. Useful in limiting the secretion in chronic bronchitis. Has been of service in asthma, chronic dysentery, gonorrhœa, leucorrhœa and other discharges from mucous surfaces. Externally is much used in Europe in the treatment of scabies as being equally effective and more agreeable than sulphur in its application.

Dose, 8 to 30 minims (.5 to 2 C.c.)

Capsules, Balsam of Peru, 10 minims. In soft gelatin.

* See note in Preface on Concentrations.

BALSAM POPLAR. *Populus balsamifera, Linné; var. canadensis, Gray.* (Buds.)

Synonyms.—Balm of Gilead; Balsampappel, Ger.

Properties.—Stimulant to mucous membranes, diuretic and anthelmintic.

An efficacious remedy in subacute gonorrhœa, gleet, chronic inflammation of the bladder, as also in chronic bronchitis. •

PREPARATION.

Fluid Extract Balsam Poplar (Balm of Gilead) buds.—Not miscible in water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

BAMBOO BRIER. *Smilax Pseudo-China, Linné.* (Rhizome.)

Synonym.—Virginia Sarsaparilla.

Properties.—Alterative and tonic. By many physicians it is preferred to the imported sarsaparillas in constitutional syphilis.

PREPARATIONS.**Extracts.**

Fluid Alterative Compound.—Not miscible with water.

Synonyms.—Bamboo Brier Comp., Succus Alterans, Mist. Smilacis Comp.

Seventeen fl. ozs. (imperial) represent 4 ozs. Av. each of Bamboo Brier, Stillingia, Burdock and Poke roots and 1 oz. Prickly Ash bark.—A combination of alteratives which was once extensively used in the treatment of syphilitic cases

Dose, 1 to 4 fluidrachms (4 to 15 C. c.) in water.

NOTE.—We have recently introduced a preparation known as Syrup Trifolium Compound, which is deemed superior to the above for the purposes indicated; for formula, etc., see Red Clover.

Fluid Extract Bamboo Brier Root.—Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.)

Solid Extract Alterative Compound.

For formula see Fluid Bamboo Brier Comp. above. One grain is equivalent to five minims of fluid Alterative Co.

Dose, 10 to 40 grains (0.65 to 2.6 Gm.).

Pill Alterative Compound, 3 grains. Dose, 3 pills.

Ext. Bamboo Brier rt, 2-3 gr.

Ext. Poke Root, 2-3 gr.

Ext. Stillingia sylvat, 2-3 gr.

Ext. Prickly Ash bark, 1-3 gr.

Ext. Burdock, 2-3 gr.

Syrup Alterative Compound.

Each fluidounce contains two drachms Fluid Alterative Comp.

Dose, 2 fluidrachms (8 C. c.) and upwards.

BANANA. *Musa sapientum, Linné.* (Root.)

Properties.—Said to have valuable alterative properties, and is used in the treatment of strumous affections.

PREPARATION.

Fluid Extract Banana root.—Miscible with water

Dose, 10 to 30 minims (0.65 to 2 C. c.).

BARBERRY. *Berberis vulgaris, Linné.* (Bark.)

Synonyms.—Epine-vinette, Fr.; Saurach, Ger.

Properties.—Stomachic, tonic, cholagogue, laxative, anti-periodic. It increases the secretion in the intestinal mucous membrane and its glandular appendages, hence it is useful in atonic dyspepsia. From its alterative action on the mucous membrane, it is especially indicated in all torpid conditions of the liver. Useful in jaundice and chronic malarial poisoning.

Barberry Bark, compressed in pound packages for retailing purposes.

BARBERRY—Continued.**PREPARATIONS.**

Berberin a so-called **Concentration**.*—A powdered preparation from Barberry bark.

One grain contains the alkaloidal principles, berberine and oxyacanthine, representing ten grains of the root bark of barberry.

Dose, 1 to 3 grains (.06 to 0.2 Gm.).

Fluid Extract Barberry.—Not miscible with water.

Dose, $\frac{1}{2}$ to 2 fluidrachms (2 to 8 C.c.)

BASSWOOD. *Tilia americana*, *Linné*. (Bark.)

Synonyms.—*Tilia glabra*, *Ventenat*; Linden, Lime-tree, Whitewood.

Properties.—Emollient, mucilaginous, vulnerary.

Basswood bark, compressed in pound packages for retailing purposes.

BAYBERRY. *Myrica cerifera*, *Linné*. (Bark.)

Synonyms.—Candle berry, Wax-myrtle; *Arbre à suif*, Fr.; Wachsbäum, Wachsgagel, Ger.

Properties.—Astringent, acrid, in large doses emetic, anti-spasmodic, errhine.

Useful in bowel complaints, jaundice and scrofula; employed, in decoction, as a gargle for sore mouth and throat, and as an injection in leucorrhœa.

Bayberry bark, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Bayberry.—Miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.)

Myricin a so-called **Concentration**.*—A powdered preparation from Bayberry bark.

Valuable as a catarrh snuff in addition to its other properties.

Dose, 1 to 5 grains (.06 to .3 Gm.)

BAY LAUREL (1). *Laurus nobilis*, *Linné*. (Leaves.)

Synonyms.—Bay tree, Sweet Bay; *Laurier commun*, Fr.; Lorbeer, Ger.

Properties.—Stimulant, narcotic, astringent; at one time a favorite remedy in scalp eruptions and as an injection in vaginal relaxation with leucorrhœa.

Bay Laurel leaves, compressed in pound packages for retailing purposes.

BAY LAUREL (2). *Myrcia acris*, *DeCandolle*, (Leaves)

Synonyms.—Bay, Wild Clove, Wild Cinnamon

Properties.—Only used for its fragrance.

PREPARATION AND FORMULA.**Bay Rum, Spirit Myrcia, U. S. P.**

Extract Bay Laurel, concentrated, P., D. & Co.'s 2 fl. ozs.

Alcohol..... $\frac{1}{2}$ gallon

Water..... $\frac{1}{2}$ gallon

Dissolve the extract in half of the alcohol and allow the solution to stand

24 hours before adding the remainder of the alcohol and water previously

mixed.

Used chiefly as a refreshing perfume and lotion in the sickroom and else-

where.

Extract Bay Laurel Concentrated.

For the extemporaneous preparation of the above, q. v.

* See note in Preface on Concentrations.

BAYCURU. *Statice brasiliensis, Boissier. (Root.)**Synonym.*—Biacuru.*Properties.*—One of the most powerful of astringents. It may be used wherever Kino and Catechu are employed, but its special indication is as an application to aphthous and ulcerative affections of the mouth and pharynx, to ulcers and serofulous sores, and as an injection in leucorrhœa.**PREPARATION.****Fluid Extract Baycuru.**—Not miscible with water.Incompatible with the mineral acids, salts of iron and lead, opium, etc.
Dose, 30 to 60 minims (2 to 4 C.c.)**BEARSFOOT.** *Polymnia Uvedalia, Linné. (Root.)**Synonyms.*—Leaf Cup, Yellow Leaf Cup.*Properties.*—Laxative, stimulant, tonic, and secernant; increases nutrition.

Externally it seems to combine valuable discutient and anodyne properties particularly in glandular enlargements. It is highly esteemed in enlargement of the spleen, the result of malaria, "ague cake," and caked breast. For these purposes it is employed in ointment form and used freely two or three times daily.

Bearsfoot root, compressed in pound packages for retailing purposes.**PREPARATIONS AND FORMULA.****Extracts.****Fluid Extract Bearsfoot.**—Not miscible with water.

Dose, 3 drops every three hours, gradually increased.

Solid Extract Bearsfoot.—One part equals 5 parts of root.

Dose, 1-2 to 2 grains (.03 to .13 Gm.).

Ointment Bearsfoot, Benzoated.

May be extemporized as follows:

Solid Extract Bearsfoot, P., D. & Co.'s 1 oz.

Benzoated Lard. 9 ozs.

Thin the solid extract with alcohol q. s. and mix thoroughly with the lard.

BEECH. *Fagus ferruginea, Aiton. (Bark, leaves.)**Synonyms.*—American Beech; Hetre ferrugineux, Fr.; Americanische Buche, Ger.*Properties.*—Astringent and slightly tonic. Useful in mild diarrhœa.**Beech bark,** compressed in pound packages for retailing purposes.**Beech leaves,** compressed in ounce packages for retailing purposes.**BEECH-DROP.** *Epiphegus virginiana, Barton. (Plant.)**Synonyms.*—Cancer root; Orobanché de Virginie, Fr; Krebswurz, Sommerwurz, Ger.*Properties.*—Astringent; used with benefit in hemorrhages of the bowels and uterus, and in diarrhœa. In erysipelas, it may be given freely and the parts bathed with a decoction of the plant. The decoction is also useful as a topical application in leucorrhœa, gleet, etc.**Beech-drop,** compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Beech-drop** (Cancer-root).—Not miscible with water.

Dose, from 10 to 30 minims (0.65 to 2 C.c.).

BELLADONNA. { *Belladonnæ folia, U.S.P.* } *Atropa Belladonna,*
{ *Belladonnæ radix, U.S.P.* } *Linné. (Lvs., rt.)**Synonyms.*—Deadly Nightshade, Dwale, Black Cherry; Tollkirsche, Ger.*Properties.*—Poisonous. Mydriatic, anodyne, excito-motor; suppresses the

BELLADONNA—Continued.

secretions. A physiological antidote to opium in opium poisoning, stimulating the respiratory centre. Used in incontinence of urine, in night sweats of hectic fever, in whooping cough, croup, chronic constipation, etc.; applied externally may be used to diminish the secretion of milk—to dilate the pupil of the eye, etc.

ANTIDOTES.—In cases of poisoning, the stomach should be promptly emptied by emetics (sulphate of zinc or copper), and the nervous disturbances combatted, as they arise, by pilocarpine or morphia.

NOTE.—Incompatible with caustic alkalies.

Belladonna lvs., compressed in ounce packages for retailing purposes.

Belladonna root, compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLES.

Atropine, U. S. P. (pure; crystals.) Its therapeutic uses are identical with those of the crude drug. Generally administered in the pill form, being only sparingly soluble in water (1 in 600).
Dose, 1-200 to 1-50 grain (.0003 to .0013 Gm.).

Atropine Sulphate, U. S. P. (neutral; crystals.) Freely soluble in water. While the pure alkaloid may be administered medicinally in the form of pills or may easily be brought into solution in mixtures containing free acid, some soluble salt like the sulphate must be employed for neutral solutions for application to the eye, etc. The medicinal properties are the same as those of the pure alkaloid.
Dose, 1-200 to 1-50 grain (.0003 to .0013 Gm.).

Homatropine Hydrobromate. This alkaloid dilates the pupil of the eye promptly and powerfully, but the effect is not so enduring as that of atropine; hence oculists employ it in making ophthalmoscopic examination of the eye.

Generally used in a 3 grains to the ounce solution.

PREPARATIONS.

Abstract Belladonna root, U. S. P. See below for formula.

Dose, $\frac{1}{2}$ a grain as a commencing dose (.03 Gm.). One part equals 2 of the root.

Belladonnin, a so-called **Concentration,*** a powdered preparation from Belladonna leaves.

Synonym.—Atropin.

Dose, 1-24 to 1-4 grain (0.003 to 0.016 Gm.).

Collodion Belladonna,—Each fl. oz. represents 20 grs. Belladonna lvs.

Extracts.

Fluid Extract Belladonna leaves,—Miscible with water; contains 0.4 % atropine by weight.

Dose, 1 to 4 minims (0.06 to 0.25 C.c.).

Fluid Extract Belladonna root, U. S. P.—Miscible with water without material precipitation; contains 0.5 % atropine by weight.

Dose, 1 to 3 minims (0.06 to 0.2 C.c.).

Powdered Extract Belladonna leaves,—One part equals 5 parts of leaves; made of a uniform strength by assay, and equal to Solid Extract Belladonna leaves.

Prepared by evaporating the solid extract at a low temperature and replacing the moisture with powdered Belladonna leaves.

Dose, from $\frac{1}{8}$ to $\frac{1}{2}$ grain (.008 to .03 Gm.).

Solid Extract Belladonna alcoholic, U. S. P. (leaves),—

One part equals 5 of leaves; made of uniform strength by assay.

Dose, $\frac{1}{8}$ to $\frac{1}{2}$ grain (0.008 to 0.03 Gm.).

Solid Extract Belladonna leaves, strictly alcoholic.

N. B.—This preparation is about half as strong again as the U. S. P.

Extract Belladonna, with which it must not be confounded.

One part equals about $7\frac{1}{2}$ parts of leaves.

Dose, 1-12 to 1-3 grain (.005 to .03 Gm.).

Hypodermic Tablets.

Atropine Sulphate, 1-100 and 1-50 grain.

Morphine and Atropine, No. 1.

Morphine sulph., $\frac{1}{8}$ gr.

Atropine sulph., 1-200 gr.

Morphine and Atropine, No. 2.

Morphine sulph., $\frac{1}{4}$ gr.

Atropine sulph., 1-150 gr.

BELLADONNA—Continued.**Liniment Belladonna, U. S. P.** See below for formula.

An efficient preparation for allaying the local pain of neuralgic rheumatism; advantageously combined with aconite.

Normal Liquids.*

Belladonna leaves.—Standard, 0.4 % atropine by weight.
Dose, 1 to 4 minims (.06 to .25 C.c.).

Belladonna root.—Standard, 0.5 % by weight atropine, or one-fourth higher than that adopted for normal liquid belladonna leaves.
Dose, 1 to 3 minims (0.06 to 0.18).

Ointment Belladonna, U. S. P.—One part solid extract Belladonna leaves in 10.

Oleate Atropine. A 2-per-cent. solution of the alkaloid in oleic acid. More active from its deeper penetration, and more cleanly in application, than the ointment.

Pills.

Aloin, Strych. and Belladonna Comp., 1 gr.

Aloin, 1-5 gr.	Strychnine, 1-60 gr.
Ext. Belladonna, 1-8 gr.	Ext. Cascara Sagrada, 1-2 gr.

Atropine, 1-120, 1-100 and 1-60 gr.

Belladonna Extract, 1-8, 1-4, 1-2 and 1 gr.

Cascara Sagrada, Nux vomica and Belladonna, 2 1-4 gr.
Ext. Cascara Sagrada, 2 gr. Ext. Belladonna, 1-16 gr.
Ext. Nux vomica, 1-8 gr.

Morphine and Atropine, 1-8 gr.

Morphine sulphate, 1-8 gr.	Atropine sulphate, 1-100 gr.
----------------------------	------------------------------

Morphine Sulphate and Belladonna, 1-2 gr.

Morphine sulph., 1-4 gr.	Ext. Belladonna, 1-4 gr.
--------------------------	--------------------------

Phosphorus and Belladonna, 1-8 gr.

Phosphorus, 1-100 gr.	Ext. Belladonna, 1-8 gr.
-----------------------	--------------------------

Quinine and Extract Belladonna, 1 1-2 gr.

Quinine sulph., 1 gr.	Ext. Belladonna, 1-2 gr.
-----------------------	--------------------------

Consult numbers 13, 15, 17, 18, 33, 37, 95, 130, 272, 283, 306, 352, 353, 448, and 453 in appended Pill Formula List for other pills containing belladonna.

Plaster Belladonna root, U. S. P. In half-pound rolls or cans. See below for formula for extemporaneous preparation.

Solution Atropine Sulphate, B. P. See below for formula.

Dose, 1 to 4 minims = 1-100 to 1-25 grain (.0006 to .0026 Gm.).

Excellent for ophthalmic use. The camphor water employed prevents decomposition better than distilled water.

Tablet Triturates.

Atropine Sulphate, 1-500, 1-200 and 1-100 grain.

Belladonna, Fluid Extract, 1-2 and 1 minim.

Morphine and Atropine, No. 1.

Morphine sulph., 1-8 gr.	Atropine sulph. 1-150 gr.
--------------------------	---------------------------

Morphine and Atropine, No. 2.

Morphine sulph., 1-4 gr.	Atropine sulph., 1-120 gr.
--------------------------	----------------------------

Morphine and Atropine, No. 3.

Morphine sulph., 1-6 gr.	Atropine sulph., 1-180 gr.
--------------------------	----------------------------

Tinctures.

Belladonna "German Tincture."

We import this tincture of the green root; it is not a German Pharmacopœial preparation.

Dose, 2 to 10 minims (.12 to .6 C.c.).

Belladonna, U. S. P. (Leaves.) See below for formula.

Dose 4 to 15 minims (.24 to 1 C.c.).

FORMULÆ.

Abstract Belladonna root, U. S. P.

Normal liquid or fl. extract Belladonna root, P., D., & Co.'s. 1 fl. oz.
Sugar of milk, q. s. to make. 228 grains.
Evaporate the normal liquid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract;

* See note in Preface on Normal Liquids.

BENZOIN—Continued.**Tinctures.****Benzoin, U. S. P.**

Fluid Benzoin, P. D. & Co.'s..... 4 fl. ozs.
 Alcohol..... 12 fl. ozs.
 Dose, 1-3 to 1-2 fluidrachm (1.25 to 2 C.c.)

Benzoin Compound, U. S. P.

Fluid Benzoin compound, P. D. & Co.'s..... 4 fl. ozs.
 Alcohol..... 12 fl. ozs.
 Known as Friar's balsam, Jesuits' drops, Turlington's balsam, etc.
 Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.) well diluted.

BERBERIS AQUIFOLIUM, Pursh. (Root.)

Synonyms.—Oregon Grape-root, Mountain Grape, Holly-leaved Barberry.

Properties.—Berberis Aquifolium is one of the most powerful alteratives and tonics. It stimulates digestion and assimilation, and excites absorption. "Advantageously combined with Cascara Sagrada in the treatment of constipation."† See Cascara (Cordial).

In constitutional syphilis and in scrofulous complaints its use is attended with remarkably good results, and pityriasis, psoriasis and other cutaneous diseases of like type often yield to its influence.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Berberis Aquifolium.—Miscible with water without material precipitation.
 Dose, 10 to 30 minims (0.6 to 2 C.c.), three or four times a day.

Powdered Extract Berberis Aquifolium.—Of same strength as the solid extract: produced by evaporating at a low temperature the solid extract, and replacing the moisture with powdered drug.
 Dose, from 2 to 6 grains (0.13 to .4 Gm.).

Solid Extract Berberis Aquifolium.—One part equals 5 of root.
 Dose, 2 to 6 grains (.13 to .4 Gm.).

Malt Extract Combinations.

Berberis Aquifolium, 1-2 drachm Berberis in each fl. oz.
 Dose, 2 to 4 fluidrachms (8 to 15 C.c.).

Cascara Sagrada and Berberis Aquifolium, 1-2 drachm each of Cascara and Berberis in each fluidounce.
 Dose, 2 to 4 fluidrachms (8 to 15 C.c.).

Pills.

Berberis Aquifolium Extract, 3 grains.

Berberis Comp., 3 grains.

Ext. Berberis aquifol., 2 gr. Ext. Cascara Sagrada, 1 gr.

BETH-ROOT. Trillium erectum, Linné. (Rhizome.)

Synonyms.—Bath Fower, Ground Lily, Birth Root, Lamb's Quarter, Wake-robin.

Properties.—Astringent and tonic. Good results are reported from its use in hæmoptysis and in bronchorrhœa. Said to facilitate parturition when given for some time prior to the event.

Beth-root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Beth-root.—Miscible with water.
 Dose, 30 to 60 minims (2 to 4 C.c.).

Pill, Trillium, 1-8 grain.

Trillin, a so-called **Concentration**,* a powdered preparation of Beth-root.
 Dose, 2 to 4 grains (0.12 to 0.25 Gm.).

* See note in Preface on Concentrations.

† Morse, "New Therapeutical Agents," p. 177.

BISTORT. *Polygonum bistorta*, *Linné*. (Rhizome.)

Synonyms.—Snake weed; couleuvrine, Fr.; Natterwurz, Ger.

Properties.—A simple, mild astringent, employed in the same complaints as other vegetable astringents. Thus it is prescribed in hemorrhages, passive diarrhoeas, leucorrhœa, and as a local application in relaxed conditions of the throat, anus, vagina, etc.

PREPARATION.

Fluid Extract Bistort.—Not miscible with water.
Dose, 20 to 40 minims (1 3 to 2 6 C.c.).

BITTER BUGLEWEED. *Lycopus europæus*, *Linné*. (Herb.)

Synonyms.—Bitter Bugle, European Bugleweed.

Properties.—Tonic and alterative, exercising its influence most markedly in the glandular enlargements and splenic engorgement resulting from chronic malarial poisoning. Valuable as a prophylactic in exposure to malaria.

Bitter Bugleweed, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Bitter Bugleweed.—Miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

BITTER ORANGE. *Aurantii Amari cortex*, U. S. P. Rind of the fruit of *Citrus vulgaris*, *Risso*.

Synonym.—Pomeranzenschale, Ger.

Properties.—A mild tonic, carminative and stomachic. A pleasant and useful addition to preparations of the bitter tonics, such as quassia, gentian, columbo, etc.

PREPARATIONS AND FORMULÆ.

Fluid Extract Bitter Orange Peel, U. S. P.—Miscible with water without material precipitation.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Syrup Bitter Orange Peel, B. P.

Fluid Extract Bitter Orange Peel, P. D. & Co.'s.....1 fluidrachm.

Syrup7 fl. ozs.

Dose, ad libitum; a flavoring agent only.

Tincture Bitter Orange Peel, U. S. P.

Fluid Extract Bitter Orange Peel, P. D. & Co.'s.....2 fl. ozs.

Dilute Alcohol8 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

NOTE.—Double the strength of the B. P. tincture.

BITTER-ROOT. *Apocynum androsæmifolium*, *Linné*. (Root.)

Synonyms.—Dog's Bane, Wandering Milkweed.

Properties.—Tonic, diaphoretic, laxative, and in large doses emetic. Has been found valuable in the treatment of chronic hepatic torpidity, and in conjunction with Yellow Parilla in dyspepsia and amenorrhœa. As a diaphoretic it should be combined with opium.

Bitter-root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Apocynin, a so-called **Concentration**,* a powdered preparation from Bitter-root.

Dose, 1-2 to 2 grains (.03 to .13 Gm.).

* See note in Preface on Concentrations.

BITTER-ROOT—Continued.**Extracts.**

Fluid Extract Bitter-root.—Not miscible with water without some precipitation.

Dose, as an emetic, 1-2 fluidrachm (2 C.c.); tonic and diaphoretic, 10 to 20 minims (0.6 to 1.3 C.c.).

Solid Extract Bitter-root.—One part equals 5 of root.

Dose, 1 to 4 grains (.06 to .26 Gm.).

BITTERSWEET. Dulcamara, U. S. P. *Solanum Dulcamara*, Linné.
(Leaves, twigs.)

Synonyms.—Woody Nightshade; Douce-amere, Fr.; Bittersüss-Stengel, Ger.

Properties.—In large doses narcotic, but used principally as an alterative diuretic, diaphoretic and discutient. Employed in the treatment of cutaneous eruptions, particularly those of a scaly nature. It is also prescribed in chronic rheumatism and chronic catarrh, and it is said to have useful anaphrodisiac properties.

Bittersweet leaves, compressed in ounce packages for retailing purposes.

Bittersweet twigs, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Bittersweet twigs, U. S. P.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Solid Extract Bittersweet.—One part equals 5 of the twigs.

Dose, 5 to 15 grains (.3 to 1 Gm.).

Ointment Discutient, prepared after King's formula; contains in 10 parts of ointment 1 each of Bittersweet, Stramonium, Conium leaves, Belladonna, Yellow Dock and 2 parts of Venice Turpentine.

BLACK ALDER. Prinos, U. S. P. *Prinos verticillatus*, Linné. (Bark.)

Synonyms.—Winterberry, Fever-bush.

Properties.—Tonic, astringent and alterative. Used with success in intermittent fevers and in diarrhoea, and is recommended as an alterative in cases of skin diseases. Properly diluted it forms a useful application for gangrenous and ill conditioned ulcers, and some cutaneous eruptions.

Black Alder bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Black Alder.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

BLACK ASH. Fraxinus ambucifolia, Lambert. (Bark)

Properties.—Tonic and astringent. May be used for all internal and external purposes, where a combination of astringency with tonic influence is indicated.

Black Ash bark, compressed in pound packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Black Ash.—Miscible with water without material precipitation.

Dose, 1 to 4 fluidrachms (4 to 15 C. c.).

Solid Extract Black Ash. One part equals 5 of bark.

Dose, 10 to 20 grains (.65 to 1.3 Gm.).

BLACKBERRY. *Rubus*, U. S. P. *Rubus villosus*, *Aiton*. (Root, bark of root.)

Properties.—Astringent and tonic. Used in diarrhoea, dysentery, and as an ingredient in compounds for cholera infantum and in a relaxed condition of the bowels of children; also as an astringent gargle and lotion.

Blackberry, bark of root, U. S. P., compressed in pound packages for retailing purposes.

Blackberry root, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Blackberry, U. S. P.—Not miscible with water. Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Blackberry Compound. (Blackberry Aromatic.)—Not miscible with water.

Each pint represents: 14 ozs. Blackberry root; 1 1-3 ozs. Cloves; 1 1-3 ozs. Cassia.

Particularly adapted to the treatment of the diarrhoea of infants and young persons.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Powdered Extract Blackberry.—One part equals 5 parts of the root bark; of the same strength as the solid extract.

Prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered blackberry root.

Dose, 3 to 10 grains (.19 to .65 Gm.).

Solid Extract Blackberry.—One part equals 5 of the root bark.

Dose, 3 to 10 grains (.19 to .65 Gm.).

Syrups.

Blackberry, U. S. P.—Each fl. oz. represents 120 grs. of root bark.

May also be made extemporaneously as follows:

Fluid Extract Blackberry, P., D. & Co.'s. 4 fl. ozs.

Syrup. 12 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

Blackberry Aromatic.—Each fl. oz. represents 120 grs. Blackberry root bark combined with aromatics.

May also be made extemporaneously as follows:

Fluid Extract Blackberry Compound, P., D. & Co.'s. 4 fl. ozs.

Syrup. 12 fl. ozs.

Dose, 1 to 3 fluidrachms (4 to 12 C. c.).

BLACK BIRCH. *Betula lenta*, *Linné*. (Bark.)

Synonyms.—Cherry birch, Sweet birch.

Properties.—Popularly used to prepare diaphoretic infusions.

Black Birch bark, compressed in pound packages for retailing purposes.

BLACK COHOSH. *Cimicifuga*, U. S. P. *Cimicifuga racemosa*, *Nuttall*. (Rhizome.)

Synonyms.—*Actæa racemosa*, *Linné*; Black Snake Root, Rattle-snake Root, Squaw Root; *Actée à grappes*, Fr.; *Schwarze schlangenwurz*, Ger.

Properties.—Tonic, nervine and antispasmodic. *Cimicifuga* has acquired no little reputation in the treatment of chorea, convulsions, nervous excitability, asthma, and other spasmodic diseases, in remittent and intermittent fevers, and in acute and chronic rheumatism. A valuable uterine tonic, and useful in the headaches attendant on uterine derangement.

Black Cohosh root, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.

Cimicifugin, a so-called **Concentration**,* a powdered preparation from Black Cohosh root.

Synonym.—Macrotin.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

* See note in Preface on Concentrations.

BLACK COHOSH - Continued.**Elixirs.****Cimicifuga** (120 grains Black Cohosh in each fl. oz).

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Cimicifuga Compound.—Each fluidounce represents: 30 grains Black Cohosh; 8 grains Seneka; 10 grains Licorice root; 4 grains Ipecac; 8 grains Cherry bark.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Extracts.**Fluid Extract Black Cohosh, U. S. P.**—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Black Cohosh Compound.—Not miscible with water.

Each pint represents: 8 ozs. Black Cohosh; 2 ozs. Seneka; 3 ozs. Licorice; 1 oz. Ipecac; 2 ozs. Cherry bark.

Nervous and arterial sedative; in larger doses acting as a diaphoretic and causing increased secretion from the bronchial mucous membrane. Useful in chronic rheumatism, both articular and when existing in the form of lumbago; in atonic amenorrhœa; chronic bronchitis.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Powdered Extract Black Cohosh.—One part equals five of the root; of the same strength as the solid extract. Prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Black Cohosh.

Dose, 3 to 10 grains (0.19 to 0.65 Gm.).

Solid Extract Black Cohosh.—One part equals 5 parts of root.

Dose, 3 to 10 grains (0.19 to 0.65 Gm.).

Pills.**Cimicifugin, 1 gr.****Dysmenorrhœa,** Alvord's, 1 1-2 gr.

Morphine sulph., 1-10 gr.

Quinine sulph., 2-3 gr.

Cimicifugin, 2-3 gr.

Female, Amenorrhœa, 4 grs.

Ext. Black Cohosh, 1 gr.

Ext. Cotton root, 1 gr.

Iron sulph. exsic., 1 gr.

Purified Aloes, 1 gr.

Podophyllin Co., Eclectic, 1-2 gr.

Podophyllin, 1-8 gr.

Leptandrin, 1-16 gr.

Juglandin, 1-16 gr.

Macrotin (Cimicifugin) 1-32 gr.

Oleo-resin Capsicum, 1-32 gr.

Tinctures.**Black Cohosh, U. S. P.**

Fluid Extract Black Cohosh, P., D. & Co.'s..... 4 fl. ozs.

Alcohol 14 fl. ozs.

Water 5 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Black Cohosh, B. P. (Tincture of Actæa.)

Fluid Extract Black Cohosh, P., D. & Co.'s..... 1 fl. oz.

Dilute Alcohol..... 7 fl. ozs.

Dose, 1 fluidrachm and upwards (4 C.c.).

BLACK HAW. Viburnum, U. S. P., Viburnum prunifolium, *Linné*.
(Bark of root)

NOTE.—The bark is the part directed for use by the U. S. P.; we employ the bark of the root since experience has demonstrated it to be more efficient.

Properties.—Held in high esteem as a uterine tonic and sedative, especially valuable in threatened abortion and as a corrective of the chronic disposition to miscarry. In the latter class of cases it should be given continuously in moderate doses for at least a month prior to the usual time at which the miscarriage takes place. In dysmenorrhœa due to almost every cause, it may be prescribed with benefit, commencing its administration some days before the menstrual crisis. In after-pains it is also valuable.

BLACK HAW—Continued.

Black Haw bark of root, compressed in pound packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Black Haw.—Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.), several times a day.

Powdered Extract Black Haw.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Black Haw. Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Solid Extract Black Haw.—One part equals 5 of bark of root.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Pill Black Haw Extract, 3 grains.

BLACK HELLEBORE. *Helleborus niger*, *Linné*. (Rhizome.)

Synonyms.—Christmas Rose; Ellébore noir, Fr.; Schwarze Nieswurzel, Weinnachtswurzel, Ger.

Properties.—Poisonous. Drastic cathartic, anthelmintic and emmenagogue. In overdoses a violent gastro-intestinal irritant, producing death by hyper-catharsis with convulsions and collapse.

Antidotes.—Evacuate the stomach, administer stimulants and external heat. Opium may be given to allay the irritation of the intestines.

Black Hellebore root, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Black Hellebore.—Not miscible with water without some precipitation.

Dose, as a purgative, 5 to 20 minims (0.3 to 1.3 C.c.).

Powdered Extract Black Hellebore.—Of the same strength as the solid extract. Prepared by evaporating the solid extract, at a low temperature, to dryness, and replacing the moisture with powdered black hellebore. Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Solid Extract Black Hellebore.—One part equals 5 of root.

Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Pills.

Black Hellebore Extract, 1 gr.

Emmenagogue, Improved, 4 1-2 gr.

Ergotin, 1 gr.

Purified Aloes, 1 gr.

Oil Savin, 1-2 gr.

Ext. Black Hellebore, 1 gr.

Iron sulph. exsic., 1 gr.

Female, Hooper's, 2 1-2 gr.

Purified Aloes, 1 gr.

Iron sulph. exsic., 1-2 gr.

Ext. Black Hellebore, 1-4 gr.

Powd. Jam. Ginger, 1-8 gr.

Gum Myrrh, 1-4 gr.

Castile Soap, 1-8 gr.

Powd. Canella, 1-8 gr.

BLACK OAK. *Quercus tinctoria*, *Bartram*. (Bark.)

Synonym.—Quercitron bark.

Properties.—Astringent and tonic. Generally used externally as a wash in prolapsus ani and hæmorrhoidal affections and as an injection in leucorrhœa.

Black Oak bark, compressed in pound packages for retailing purposes.

BLACK PEPPER. *Piper*, U. S. P. *Piper nigrum*, *Linné*. (Fruit.)

Properties.—Well-known as a condiment; a powerful stimulant to the stomach and intestinal canal. In intermittents it may be used as an adjuvant

BLACK PEPPER—Continued.

to more powerful febrifuges, when the stomach is not acted upon by Quinine, as in the case of drunkards. Useful in colic, cholera morbus, and flatulency.

PREPARATIONS.

Fluid Extract Black Pepper.—Not miscible with water.
Dose, 5 to 20 minims (.3 to 1.25 C.c.), diluted with water.

Pills.

Ague , 3 grs.	
Chinoidin, 2 grs.	
Ext. Coloc co., 1-3 gr.	Oleoresin Black Pepper, 1-6 gr.
	Iron sulph. exsic., 1-2 gr.
Anti-Chill , 4 grs.	
Chinoidin, 1 gr.	Arsenious Acid, 1-20 gr.
Iron ferrocyanide, 2 gr.	Oleoresin Black Pepper, 1 gr.
Also the above Anti-chill 1-16th strength ("Pink Granule")	
Anti-Malarial , McCaw's, 2 grs.	
Quinine sulph., 1 gr.	Gelsemerin, 1-4 gr.
Iron sulph. exsic., 1-4 gr.	Podophyllin, 1-8 gr.
Arsenious Acid, 1-80 gr.	Oleoresin Black Pepper, 1-16 gr.
Chinoidin Comp. , 3 1-2 grs.	
Chinoidin, 2 grs.	
Piperin, 1-2 gr.	Iron sulph. exsic., 1 gr.
Piperin , 1-20 gr.	

BLACK WALNUT. *Juglans nigra*, *Linné*. (Bark, leaves.)

Properties.—Alterative and deobstruent. It modifies favorably all chronic cachectic conditions, such as those of scrofula and syphilis, and promotes the healing of indolent ulcers, specific and non-specific. It is employed freely in the form of infusion as a local application, and is also taken internally, the effect in chronic cases becoming manifest sometimes only after its persistent use for some weeks. It may frequently be combined to advantage with mercurial alteratives or with iodide of potassium.

Black Walnut bark, compressed in pound packages for retailing purposes.

Black Walnut leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Black Walnut Leaves.—Not miscible with water without some precipitation.
Dose, 20 to 30 minims (1.25 to 2 C.c.).

BLACK WILLOW. *Salix nigra*, *Marshall*. (Bark, buds.)

Properties.—Willow bark is a tonic and antiperiodic, depending for its activity on the salicin it contains.

The buds are a bitter tonic and they are also said to possess pronounced anaphrodisiac action and to exert a peculiar sedative influence upon the whole sexual apparatus.

Black Willow bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Black Willow buds.—Not miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

BLADDER-WRACK. *Fucus vesiculosus*, *Linné*. (Plant.)

Synonyms.—Sea-wrack, Kelp-ware, Black-tang, Varech versiculeux, Fr.; Blasentang, Ger.

Properties.—This plant has long had the reputation of a deobstruent and resolvent, and been given in goitre and scrofulous swellings—its virtues possibly depending on the iodide which it contains. Has been employed

BLADDER-WRACK—Continued.

with benefit in some cases of morbid obesity. Its use must be persisted in for some weeks before its effects are apparent, and in some instances even after a faithful trial of the drug no benefit has been derived from it. The remedy has proved efficient in so many cases, however, that it is worthy of a trial where obesity appears to be associated with serous plethora.

"The cases in which *Fucus vesiculosus* shows its most decided beneficial effects are women, in whom there exists usually some menstrual derangement, as menorrhagia and leucorrhœa, owing to an atonic and flabby condition of the uterine tissue. In such cases an improvement in these local derangements usually precedes the general reduction of fat and the improved tonicity of the general system."*

PREPARATIONS.**Extracts.**

Fluid Extract Bladder-wrack.—Miscible with water without material precipitation

Dose, 1-2 to 4 fluidrachms (2 to 15 C. c.) three times a day.

Powdered Extract Bladder-wrack.—Of the same strength as the solid extract. Prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Bladder-wrack.

Dose, 5 to 30 grains (0.3 to 2 Gm.).

Solid Extract Bladder-wrack.—One part equals 5 of the plant.

Dose, 5 to 30 grains (.3 to 2 Gm.).

Pill, *Fucus vesiculosus* ext., 3 grs.

BLESSED-THISTLE. *Centaurea benedicta*, Linné. (Herb.)

Synonyms.—*Cnicus benedictus*, Linné; Chardon bénit, Fr.; Benedictendistel, Ger.

Properties.—Tonic, diaphoretic and emetic. In small doses as a mild tonic in debility of the stomach, principally in dyspepsia where there is hepatic congestion; in larger doses, administered in warm water, will produce copious perspiration to the patient in bed; in maximum dose, it is said to be emetic.

Blessed-Thistle herb, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Blessed-Thistle.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.)

Solid Extract Blessed-Thistle.—One part equals 5 of the herb.

Dose, 3 to 10 grains, (.2 to .6 Gm.).

BLOOD-FLOWER. *Asclepias curassavica*, Linné. (Herb.)

Synonym.—Bastard ipecacuanha.

Properties.—Astringent, styptic and vermifuge. Given on an empty stomach it acts as a very prompt tænicide. Also used with benefit as an injection in vaginal leucorrhœa.

PREPARATION.

Fluid Extract Blood-flower.—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

BLOOD-ROOT. *Sanguinaria*, U. S. P. *Sanguinaria canadensis*, Linné. (Rhizome)

Synonyms.—Red puccoon, Tetterwort, Indian paint; Blutwurz, Ger.

Properties.—In small doses tonic, increasing secretions; in larger doses

* Mulheron, J. J., "Obesity: a few thoughts on its Nature and Treatment," p. 2.

BLOOD-ROOT—Continued.

nauseates and reduces pulse, and acts as an expectorant; if medicinal dose is exceeded, it produces sometimes emesis, oftener burning in the stomach, vertigo, impaired vision, alarming prostration, etc. Principally used as a stimulating expectorant in bronchitis, croup and asthma, and as a remedy in atonic dyspepsia.

Blood-root, compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLE, Sanguinarine.

Sanguinarine Nitrate.—Salt of alkaloid. Expectorant, nauseant, deobstruent, etc., used especially as a stimulating expectorant in chronic bronchitis and asthma.

Dose, 1-10 to 1-4 grain (.006 to .016 Gm.).

Sanguinarine Sulphate.—Resembles closely in physical and therapeutic properties the nitrate, being, however, somewhat more soluble in water.

Dose, 1-10 to 1-4 grain (.006 to .016 Gm.).

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Blood Root, U. S. P.—Not miscible with water without some precipitation.

Dose, as an emetic, 10 to 20 minims (0.6 to 1.3 C. c.); as an expectorant, 2 to 5 minims (0.12 to 0.3 C. c.).

Solid Extract Blood Root.—One part equals 5 of the root.

Dose, 1 to 5 grains (.06 to .3 Gm.).

Pills.

Bilious, Junge's, 2 grs.

Manganese iod . 1-2 gr.

Leptandrin, 3-10 gr.

Juglandin, 3-10 gr.

Sanguinarin, 1-5 gr.

Ext. Henbane, 3-5 gr.

Eucalyptus Comp., 1 5-8 grs.

Ext. Eucalyptus, 1 gr.

Sanguinarin, 1-8 gr.

Ext. Canadian Hemp, 1-3 gr.

Sanguinaria Ext., 1-2 gr.

Sanguinarin (conc.), 1-2 and 1 gr.

Sanguinarin, a so-called **Concentration**.*—A powdered preparation from Blood root.

Dose, 1-8 to 1 grain (0.008 to 0.065).

Tincture Blood Root, U. S. P.

Fluid Extract Blood Root, P., D. & Co.'s 3 fl. ozs

Dilute Alcohol. 19 fl. ozs

Dose, as an expectorant, 10 to 40 minims (0.6 to 2.5 C. c.)

Vinegar Blood Root, U. S. P.

Fluid Extract Blood Root, P., D. & Co.'s ... 2 fl. ozs

Dilute Acetic Acid. 18 fl. ozs

Dose, as an alterative and expectorant, 10 to 40 minims (0.6 to 2.5 C. c.).

BLUE CARDINAL-FLOWER. *Lobelia syphilitica*, *Linnaeus* (Leaves.)

Synonyms.—Great Lobelia; Gemeine Lobelie, Ger.

Properties.—Cathartic and emetic. Useful in dropsy to assist the removal of fluids.

Dose, a wineglassful (60 C. c.) of an ounce to the pint infusion.

Blue Cardinal-flower leaves, compressed in ounce packages for retailing purposes.

BLUE COHOSH. *Caulophyllum*, U.S.P. *Caulophyllum thalictroides*, *Michaux.* (Rhizome.)

Synonyms.—Papoose Root, Squaw Root, Blueberry Root.

Properties.—Diuretic, diaphoretic and anthelmintic. Used principally as an emmenagogue, parturient and anti-spasmodic.

It is used in rheumatism, dropsy, colic, cramps, hiccup, epilepsy,

* See note in Preface on Concentrations.

BLUE COHOSH—Continued.

hysteria, uterine inflammation, uterine leucorrhœa, amenorrhœa, dysmenorrhœa, etc. Used in decoction for several weeks previous to delivery, it is said to facilitate parturition.

Blue Cohosh root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Caulophyllin, a so-called **Concentration**.*—A powdered preparation of Blue Cohosh root.

Dose, 1 to 5 grains (0.06 to 0.3 Gm.).

Extracts.

Fluid Extract Blue Cohosh.—Not miscible with water.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

Fluid Extract Blue Cohosh Comp.—Not miscible with water. Nine fl. ozs. (imperial) represents 4 ozs. av. Blue Cohosh, 2 ozs. each of Ergot and Water Pepper, and 1 oz. of Savin. A useful emmenagogue remedy.

Dose, 5 to 20 minims (0.35 to 1.3 C. c.).

Solid Extract Blue Cohosh.—One part equals 4 of the root.

Dose, 2 to 4 grains (.13 to .26 Gm.).

Pill, Caulophyllin, 1-8 gr.

BLUE FLAG. Iris, U.S.P. Iris versicolor, *Linné*. (Rhizome.)

Synonyms —Flag Lily, Liver Lily, Snake Lily; Verschiedenfarbige schwert-lilie, Ger.

Properties.—Cathartic, cholagogue and alterative. The experiments of Dr. Rutherford have demonstrated that Blue Flag is one of the most direct cholagogues of the materia medica. Indicated in all affections depending directly or indirectly upon hepatic torpor. Combined with podophyllin, in the form of a pill, it is recommended for dropsy and chronic visceral affections.

Blue Flag, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixirs.—Active ingredients stated per fluidounce.

Blue Flag, 2 drachms.

Wahoo and Blue Flag, 80 grs. Wahoo, 40 grs. Blue Flag.

Stillingia Compound: Stillingia, 30 grs.; Turkey corn, 30 grs.; Pipsissewa, 15 grs.; Elder flowers, 15 grs.; Blue flag, 15 grs.; Prickly ash berries, 7½ grs.; Coriander seed, 7½ grs.

Extracts.

Fluid Extract Blue Flag, U.S.P.—Not miscible with water.

Dose, 10 to 20 minims (0.65 to 1.3 C. c.).

Fluid Extract Stillingia Comp.—Not miscible with water.

Ingredients in same proportion as Elixir Stillingia above, but 4 times stronger.

Solid Extract Blue Flag, U.S.P.—One part equals 5 of root.

Dose, 2 to 4 grains (.13 to .26 Gm.).

Irisin, a so-called **Concentration**.*—A powdered preparation of Blue Flag root.

Dose, 1 to 3 grains (.06 to 0.2 Gm.).

Pills.

Hepatic Eclectic, 1-2 gr.

Podophyllin, 1-4 gr.

Leptandrin, 1-2 gr.

Irisin, 1-4 gr.

Irisin, 1-2 and 1 gr.

Irisin Comp., 1-2 gr.

Irisin, 1-4 gr.

Strychnine, 1-40 gr.

Ext. Nux vomica, 1-16 gr.

Powd. Capsicum, 1-3 gr.

Podophyllin, 1-10 gr.

* See note in Preface on Concentrations

BLUE FLAG—Continued.**Irisin Comp. and Henbane.**

Irisin, 1-4 gr.

Strychnine, 1-100 gr.

Popophyllin, 1-10 gr.

Ext. Henbane, 1-2 gr.

Leptandrin Comp., 13-8 gr.

Leptandrin, 1 gr.

Podophyllin, 1-8 gr.

Irisin, 1-4 gr.

Syrup Stillingia Comp.—Active ingredients same as Elixir Stillingia Comp. above.**BLUE GENTIAN.** *Gentiana puberula, Michaux.* (Root.)*Synonyms.*—American Gentian, *Gentiana Catesbæi, Elliott.**Properties.*—Tonic and stomachic in impaired digestion and anorexia. It may be profitably employed in convalescence from acute diseases and in chronic debility of the system.**Blue Gentian root,** compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Blue Gentian.**—Miscible with water.

Dose, from 10 to 40 minims (.6 to 2.5 C.c.).

BOLDO. *Boldoa fragrans, Gay.* (Leaves)*Synonym.*—Peumus Boldus, *Molina.**Properties.*—Employed chiefly as a tonic. By virtue of its essential oil it acts as a stimulant and antiseptic. It is also an anthelmintic. Especially valuable in atonic dyspepsia and in blenorhagia and chronic catarrh of the bladder it has been found useful.

Employed in cases of chronic hepatic torpor, and atony of various organs where quinine cannot be tolerated. Large doses produce emesis, and act as an irritant to the bowels.

PREPARATION.**Fluid Extract Boldo.**—Not miscible with water.

Dose, 1 to 4 minims .065 to .25 C.c.) gradually increased.

BONESET. *Eupatorium, U. S. P. Eupatorium perfoliatum, Linné.* (Leaves and Tops.)*Synonyms.*—Thoroughwort, Indian sage; Durchwachsener Wasserhanf, Ger.*Properties.*—Tonic, diaphoretic and, in large doses, emetic and laxative. A favorite remedy, especially when given in warm infusion, for colds accompanied with febrile symptoms.

As a tonic it is useful in the convalescent stages of remittent, intermittent and typhoid fevers, and in dyspepsia and general debility.

Boneset herb, U. S. P., compressed in ounce packages for retailing purposes.**PREPARATIONS.****Eupatorium,** a so-called **Concentration.***—A powdered preparation from Boneset.

Dose, 1 to 3 grains (.06 to .2 Gm.)

Extracts.**Fluid Extract Boneset, U. S. P.**—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.)

Solid Extract Boneset.—One part equals 5 of the herb.

Dose, 10 to 25 grains .6 to 1.6 Gm.).

BORAGE. *Borago officinalis, Linné.* (Leaves.)*Properties.*—Somewhat diuretic and refrigerant. Contains much mucilage and is valuable as a demulcent, emollient and protective.**Borage leaves,** compressed in ounce packages for retailing purposes.

BORAGE—Continued.**PREPARATION.**

Fluid Extract Borage.—Miscible with water without material precipitation.

Dose, 1 fluidrachm (4 C. c.)

BROOM. *Scoparius*, U. S. P. *Cytisus scoparius*, *Link.* (Tops and Twigs.)

Synonyms.—Scotch, or Irish Broom; *Sarothamnus scoparius*, *Koch*; *Genét à balais*, Fr.; *Besenginster Pfriemenkraut*, Ger.

Properties.—Diuretic, laxative, and in larger dose, cathartic. A valuable remedy in dropsical complaints, especially those of cardiac origin and also in anasarca of chronic kidney disease. Used successfully in scarlatinal albuminuria.

Broom tops, U. S. P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLE.

Sparteine Sulphate.—Sparteine is a volatile alkaloid obtained from broom tops. The sulphate is a crystallizable salt, readily soluble in water. Its properties are similar to those of digitalin or convallamarine; "Sparteine seems especially to be a regulator of the beatings of the heart."*

Dose, 1-10 to 1-3 grain (0.007 to 0.022 Gm.).

PREPARATION.

Fluid Extract Broom-tops.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.)

BUCHU, U.S.P. *Barosma betulina*, *Bartling.* (Short-leaved.) (Lvs.)

Properties.—Buchu stimulates the appetite, relieves nausea and flatulence, increases perspiration and augments the flow of the urine. Hence it is largely used in gastric debility, dyspepsia, colic, etc. Principally recommended in chronic irritation and catarrh of the bladder, hæmaturia, incontinence of urine, gravel, chronic rheumatism, cutaneous affections and dropsy.

PREPARATIONS AND FORMULA.

Barosmin, a so-called **Concentration**.†—A powdered preparation from Buchu leaves.

Dose, 1 to 3 grains (0.065 to 0.20 Gm.).

Capsule; Copaiba, Cubeb and Buchu, in soft gelatin.

Copaiba, Para, 6 minims; Oleoresin Cubeb, 2 minims; Ext. Buchu, 2 grs.

Elixirs.

Buchu, 1-4 strength of Fluid Extract Buchu.

Buchu Compound, 1-4 strength of Fluid Extract Buchu Comp.

Buchu, Juniper and Potassium Acetate.—Each fluidounce contains: Buchu, 45 grs.; Juniper, 12 grs.; Potassium acetate, 16 grs.

Buchu and Pareira Brava, 1-8 strength of Fluid Extract Buchu and Pareira Brava.

Extracts.

Fluid Extract Buchu, U. S. P.—Not miscible with water.

Dose, 15 to 30 minims (1 to 2 C. c.) diluted.

Fluid Extract Buchu Compound.—Not miscible with water.

Sixteen fluidounces, imperial, of this fluid extract represents: Buchu, 8 ozs.; Cubeb, 2 ozs.; Juniper berries, 2 ozs.; Uva Ursi, 2 ozs.; with 2 fl. ozs. Spirit of Nitrous Ether.

Principally used in chronic catarrhal diseases of the genito-urinary organs

Dose, 30 to 60 minims (2 to 4 C. c.).

*Dujardin-Beaumetz, "New Medications," p. 41. (edition of Geo. S. Davis, 1886.)

†See not in Preface on Concentrations.

BUCHU—Continued.**Fluid Extract Buchu and Juniper with Potassium****Acetate.**—Not miscible with water.

Fifteen fluidounces, imperial, represents: Buchu leaves, 12 ozs.;

Juniper berries, 3 ozs.; Potassium acetate, 432 grs.

A very efficient diuretic. Has proved valuable in leucorrhœa, dropsy, chronic inflammation of the bladder and urinary passages, etc.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Buchu and Pareira Brava.—Not miscible with water.

Each pint represents: Buchu, 8 czs., troy; Pareira Brava, 2 ozs.; Stone

root, 2 ozs.; Juniper Berries, 4 czs.

An efficacious combination in general chronic affections of the genito-urinary mucous membrane.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Solid Extract Buchu.—One part equals 5 of Buchu leaves.

Dose, 3 to 10 grains (.2 to .6 Gm.)

Tincture Buchu, B. P.

Fluid Extract Buchu, P., D. & Co.'s..... 1 fl. oz.

Alcohol..... 7 fl. ozs.

Dose, 1 to 4 fluidrachms (4 to 15 C. c.) diluted.

BUCKBEAN. *Menyanthes trifoliata*, *Linné*. (Leaves, rhizome.)*Synonyms.*—Bogbean, Marsh trefoil, Trefle d'eau (de marais), Fr.; Fieber-klee, Bitterklee, Ger.*Properties.*—Tonic, purgative, anti-scorbutic, emmenagogue and vermifuge; in larger dose emetic. Much used in chronic cutaneous eruptions and also with advantage in dyspepsia attending the digestive disturbances common to malarial districts.**Buckbean leaves**, compressed in ounce packages for retailing purposes.**Buckbean root**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Buckbean root.**—Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

BUCKEYE. *Æsculus glabra*, *Willdenow*. (Bark.)*Properties.*—Has a special action on the portal circulation and the liver.

"It is especially valuable in the constipation attending hemorrhoids. It relieves hepatic congestion admirably and seems to promote the biliary secretions."*

PREPARATION.**Fluid Extract Buckeye bark.**

Miscible with water without material precipitation.

Dose, 3 to 5 minims (.2 to .3 C. c.).

BUCKHORN BRAKE. *Osmunda regalis*, *Linné*. (Root stock.)*Synonyms.*—Buckhorn, Royal Flowering Fern.*Properties.*—Demulcent, tonic and styptic. Useful in coughs, diarrhœa and dysentery. Also as a tonic during convalescence from exhausting diseases and in leucorrhœa.**Buckhorn Brake**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Buckhorn Brake.**—Not miscible with water without some precipitation.

Dose, 1 to 3 fluidrachms (4 to 12 C. c.).

* Morse, "New Therapeutical Agents," p. 82.

BUCKTHORN BARK. *Frangula*, U. S. P. *Rhamnus Frangula*, *Linné*.

Synonyms.—Alder buckthorn; Bourdaine, Fr.; Faulbaumrinde, Ger.

Properties.—A mild, but not a very certain, laxative or cathartic. On account of its freedom from drastic and nauseating properties it is recommended as a substitute for senna as a cathartic for children. Especially valuable in the treatment of hemorrhoids, given in full laxative doses at bed time.

NOTE.—There are three species of *Rhamnus* in the market, viz: *Rhamnus Frangula*, *Rhamnus cathartica* and *Rhamnus Purshiana* (*Cascara Sagrada*), and it is important that they be not confounded.

Buckthorn bark, compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixir Buckthorn bark.—Each fl. oz. represents 120 grs. of bark.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Extracts.

Fluid Extract Buckthorn bark.—Not miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Buckthorn bark.—One part equals 5 of bark.
Dose, 3 to 10 grains (0.2 to 0.64 Gm.).

BUCKTHORN BERRIES. *Rhamnus cathartica*, *Linné*.

Synonyms.—*Cervispina cathartica*, *Moench*; Nerprun, Fr.; Kreuzdornbeeren, Ger.

Properties.—Powerfully cathartic, and useful as a hydragogue in dropsy, rheumatism and gout, but apt to cause griping and nausea. The best method of administration is in the form of syrup, or in combination with aromatics to modify its action.

PREPARATIONS AND FORMULA.

Fluid Extract Buckthorn berries.—Miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Syrup Buckthorn berries.—Each fluidounce represents 120 grains of the berries in combination with aromatics.

Dose, for a child, 1-2 to 1 fluidrachm (2 to 4 C.c.).

This syrup may be prepared extemporaneously as follows:

Fluid Extract Buckthorn berries, P., D. & Co.'s.	2 fl. ozs.
Fluid Extract Ginger, P., D. & Co.'s.	1 1/2 fl. drms.
Fluid Extract Pimento, P., D. & Co.'s.	1 1/2 fl. drms.
Simple Syrup	15 fl. ozs.

BUGLEWEED. *Lycopus virginicus*, *Linné*. (Herb.)

Synonyms.—Sweet Bugle, Paul's Betony, Water Horehound.

Properties.—Narcotic and astringent. Valuable in pulmonary hemorrhage. It diminishes the frequency of the pulse, quiets irritation, and allays cough.

Bugleweed herb, compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Bugleweed.—Miscible with water.
Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Solid Extract Bugleweed.—One part equals 5 of herb.
Dose, 5 to 15 grains (0.32 to 1 Gm.).

Lycopin, a so-called **Concentration**,* a powdered preparation from Bugleweed.
Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

* See note in Preface on Concentrations.

BURDOCK. Lappa, U. S. P. Lappa officinalis, *Allioni*. (Leaves, Root, U. S. P., Seeds.)

Properties.—Alternative, aperient, diuretic and sudorific. Used in gout, rheumatism, syphilis, leprosy, and other similar disorders, and is preferred by some to sarsaparilla.

Burdock seed, according to Dr. Reiter, of Pittsburg, has almost a specific effect in curing certain chronic skin diseases, especially psoriasis. A reprint of his article sent on application.

Burdock Leaves, compressed in ounce packages for retailing purposes.

Burdock root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Extracts.

Fluid Extract Alternative Comp.—Miscible with water.

Synonyms.—Bamboo Brier Comp., Succus Alterans, Mist. Smilacis Comp.

Seventeen fluidounces imperial represent 4 ozs. av. each of Bamboo Brier, Stillingia, Burdock and Poke root, and 1 oz. Prickly Ash bark.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.) in water.

NOTE.—Syrup Trifolium Comp. is rapidly supplanting this formula as an alternative, see below.

Fluid Extract Burdock root.—Miscible with water without material precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Fluid Extract Burdock seed.—Not miscible with water.

Dose, as a tonic, 10 to 30 minims (0.6 to 2 C.c.); as an alternative, 30 to 60 minims (2 to 4 C.c.), well diluted, half an hour before meals.

Solid Extract Alternative Comp.—One grain is equivalent to 5 minims Fluid Extract Alternative Comp., see above for formula.

Dose, 10 to 40 grains (0.65 to 2.6 Gm.).

Solid Extract Burdock root.—One part equals 5 of root.

Dose, 6 to 20 grains (0.4 to 1.3 Gm.).

Pill, Alternative Compound, 3 gr.

Ext. Smilax Sarsap., 2-3 gr.

Ext. Stillingia sylvat., 2-3 gr.

Ext. Burdock, 2-3 gr.

Ext. Poke root, 2-3 gr.

Ext. Prickly ash bark, 1-3 gr.

Syrups.—Each fluidounce representing ingredients mentioned.

Syrup Alternative Compound, 2 drachms Fluid Extract Alternative Comp.: see above for formula.

Dose, 2 fluidrachms (8 C.c.) and upwards.

Syrup Trifolium Comp.

Red clover, 32 grs.; Stillingia, 16 grs.; Burdock root, 16 grs.; Poke root, 16 grs.; Berberis aquifolium, 16 grs.; Cascara amara, 16 grs.; Prickly ash bark, 4 grs.; Potassium iodide, 8 grs.

Recommended by Dr. Rush as superior in efficacy to the combination of alternatives that has recently acquired considerable notoriety under the name of Bamboo Brier root compound, Succus Alterans or Alternative compound. Its restorative action in secondary syphilis is prompt and unequivocal.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.) three times a day.

BUTTERNUT. Juglans, U. S. P. Juglans cinerea, *Linné*. (Bark of root, U. S. P., leaves)

Synonyms.—White Walnut, Oil Nut; Noyer gris, Fr.; Graue Wallnusrinde, Ger.

Properties.—A gentle and agreeable cathartic, causing no griping, nor subsequent weakness of the intestines. Is valuable in cases of habitual constipation, colo-rectitis and several other forms of intestinal derangement attended by hepatic torpor.

Butternut bark of root, U. S. P., compressed in pound packages for retailing purposes.

Butternut leaves, compressed in ounce packages for retailing purposes.

BUTTERNUT—Continued.**PREPARATIONS.****Extracts.**

Fluid Extract Butternut.—Not miscible with water.
Dose, 1 to 2 fluidrachms (1 to 8 C. c.).

Powdered Extract Butternut.—Of the same strength as the official extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Butternut.
Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Solid Extract Butternut, U.S.P.—One part equals 5 of the bark.
Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Juglandin, a so-called **Concentration**,* a powdered preparation from Butternut.
Dose, 1 to 5 grains (0.06 to 0.32 Gm.).

Pill, Anti-Constipation, Goss', 2 1-4 gr.

Podophyllin, 1-4 gr.

Ext. Coloc., 1-4 gr.

Ext. Butternut, 1-2 gr.

Ext. Nux vomica, 1-8 gr.

Ext. Henbane, 1-8 gr.

Ext. Gentian, 1-4 gr.

Ext. Cascara sagrada, 1-4 gr.

Powd. Canadian Hemp, 1-2 gr.

Consult Nos. 63 and 356 in Pill Formulæ list appended for other pills containing Butternut.

BUTTON-BUSH. *Cephalanthus occidentalis*, *Linne*. (Bark.)

Synonyms.—Pond dogwood, Globe flower, Button-wood, Crane willow.

Properties.—Tonic, febrifuge, aperient and diuretic. Also claimed to be expectorant and diaphoretic. It has been used with asserted success in intermittent and remittent fevers, and in gravel. It seems to exert a special influence over the respiratory and digestive organs, relieving the former by promoting digestion and assimilation, effecting a healthy action of the skin, and relieving any local congestion of the lungs or bronchi—hence its use in consumption, bronchitis, catarrh, whooping-cough, croup, and coughs generally.

PREPARATION.

Fluid Extract Button-bush.—Not miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

BUTTON SNAKEROOT. *Liatis spicata*, *Willdenow*. (Root.)

Synonyms.—Gay Feather, Devil's Bit.

Properties.—Diuretic, with tonic, stimulant and emmenagogue properties.

It is very efficacious in gonorrhœa, gleet and nephritic diseases. Said to be beneficial in scrofula, dysmenorrhœa, amenorrhœa, after-pains, etc.; also in Bright's disease, in connection with bugleweed and unicorn. May be used as a gargle in sore throat, or as an injection in leucorrhœa.

Button Snakeroot, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Button Snakeroot.—Not miscible with water.
Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

CALABAR BEAN. *Physostigma*, U.S.P. *Physostigma venenosum*, *Balfour*. (Seeds)

Synonyms.—Ordeal Bean; Fève de Calabar, Fr.; Kalabarbohne, Ger.

Properties.—POISONOUS. Spinal sedative, acting especially on the motor centers. Employed with success in traumatic tetanus, and in strychnine poisoning, also in cerebro-spinal meningitis, and in constipation due to

* See note in Preface on Concentrations.

CALABAR BEAN—Continued.

torpor of the bowels, in which case it is advantageously combined with Belladonna and Nux vomica. Contracts the pupil.

ANTILOTES.—Give emetics (apomorphine, zinc sulphate, mustard); in severe cases, atropine hypodermically (1-60 grain, every 15 minutes, until pupils dilate), chloral, strychnine, stimulants, artificial respiration.

NOTE.—Incompatible with vegetable astringents and caustic alkalies.

ACTIVE PRINCIPLE AND DERIVATIVES.

Physostigmine.—Alkaloid (pure crystals). *Synonym.*—Eserine.

Insoluble in water, and on this account the sulphate or salicylate is preferred in most cases.

Dose, 1-300 to 1-100 grain (0.0002 to 0.0006 Gm.).

Physostigmine Salicylate, U. S. P.—This has the advantage over the sulphate in that it forms permanent crystals, and its solutions in water are more stable. Chiefly employed as an application to the eye, but may be used wherever Calabar bean is indicated.

Dose, 1-300 to 1-100 grain (0.0002 to 0.0006 Gm.).

Physostigmine Sulphate.—An extremely deliquescent salt, which is rapidly giving place to the more manageable salicylate.

Dose, 1-300 to 1-100 grain (0.0002 to 0.0006 Gm.).

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Calabar bean.—Miscible with water.

Dose, 1 to 4 minims (0.06 to 0.25 C.c.), cautiously increased.

Powdered Extract Calabar bean.—Of same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered drug.

Dose, 1-8 grain (0.008 Gm.), cautiously increased.

Solid Extract Calabar bean, U. S. P., Assayed.—One grain represents 25 grains of calabar bean, and contains 5 per cent. alkaloid.

Dose, 1-8 grain (0.008 Gm.), cautiously increased.

Hypodermic Tablet, Physostigmine Sulphate, 1-100 grain.

Pills.

Calabar Bean Extract, 1-12 gr.

Physostigmine Salicylate, 1-100 gr.

Podophyllin, Belladonna and Calabar, 3-4 gr.

Podophyllin, 1-4 gr.

Ext. Calabar bean, 1-4 gr.

Ext. Belladonna, 1-4 gr.

Tablet Triturates, Physostigmine Salicylate, 1-500, 1-400, 1-300, 1-200, 1-100 grain.

Tincture Physostigma, U. S. P.

Fluid Extract Calabar Bean, P., D. & Co.'s.....1 fl. oz.

Alcohol.....11 fl. ozs.

Dose, 10 to 40 minims (0.65 to 2.6 C.c.).

CALENDULA, U. S. P. *Calendula officinalis, Linné.* (Flowers.)

Synonyms.—Marigold; Souci, Fr.; Ringelblume, Ger.

Properties.—Slightly stimulant and diaphoretic. It was formerly employed in spasmodic affections, strumous maladies, etc.; but its chief use is as a local application to ulcers and wounds, as it is claimed to promote cicatrization.

Dr. Clary says: "As a local remedy after surgical operations it has no equal in the *Materia Medica*."

Calendula flowers, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Calendula flowers.—Not miscible with water.

Dose, internally, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract Calendula flowers.—One part equals 5 of flowers.

Dose, 5 to 10 grains (0.3 to 0.6 Gm.).

CALENDULA—Continued.

Glycerole Calendula.—Each fluidounce represents 120 grs. Calendula.

Tincture Calendula, U. S. P.

Fluid Extract Calendula, P., D. & Co.'s.....4 fl. ozs.

Alcohol10 fl. ozs.

Water7 fl. ozs.

After standing for a day or two decant the clear tincture.

CALIFORNIA FEVER-BUSH. *Garrya Fremontii, Torrey.* (Lvs.)

Properties.—Used as a cure for chills and fever since the occupation of California by Americans. Dr. Q. C. Smith, of Cloverdale, Cal., says, concerning it: "I consider *Garrya* leaves a valuable remedy, deserving of notice by the medical world, and simply invaluable as a remedy for remittent fevers, chills, etc., and other malarial ailments." He reports cases which had resisted the ordinary treatment, but which yielded promptly to this remedy. It contains a bitter alkaloid closely resembling quinine in its physiological and therapeutic action. As a tonic, *Garrya* is valuable in chronic diseases connected with debility.

PREPARATION.

Fluid Extract California Fever-bush (*Garrya*).—Not miscible with water without some precipitation.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

CALIFORNIA LAUREL. *Umbellularia californica, Nuttall.* (Lvs.)

Synonym.—Spice Tree.

Properties.—The use of this drug in neuralgic headache, and in cerebrospinal meningitis is followed by favorable results. In atonic diarrhoea and intestinal colic it is said to possess curative properties.* Is used externally in rheumatism.

PREPARATION.

Fluid Extract California Laurel.—Not miscible with water without some precipitation.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

CAMPHOR. A stearopten from the wood of *Cinnamomum camphora*, *Fr. Nees & Ebermaier.*

Properties.—An efficient remedy in summer diarrhoea. It will allay cough and promote expectoration, and is serviceable as a stimulant in chronic bronchitis, capillary bronchitis, typhoid pneumonia, etc. In larger dose it allays maniacal and sexual excitement. Externally it is a mild counter-irritant. Solubility 1 in 900 of water.

Dose, 5 to 20 grains (0.3 to 1.3 Gm.).

DERIVATIVE.

Camphor Monobromated.—Almost insoluble in water, freely in alcohol. Has been used to advantage in whooping cough, neuralgia, chorea and hysteria.

Dose, 1 to 10 grains, (0.065 to 0.56 Gm.).

PREPARATIONS.**Pills.****Blennorrhagic.**

Terebinth. alb., 1 1-2 gr.

Ext. Humuli, 3-4 gr.

Camphor monobr., 3-4 gr.

Podophyllin, 1-8 gr.

Camphor, 1-20 gr.**Camphor Comp., 3 grs**

Camphor, 1 gr.

Powdered opium, 1 gr.

Powd. Kino, 1 gr.

Ext. Capsicum, 1-16 gr.

* Morse, "New Therapeutical Agents," p. 152.

CAMPHOR—Continued.**Camphor and Henbane, 2 gr.**

Camphor, 1 gr.

Ext. Henbane, 1 gr.

Camphor monobromated, 1, 2 and 3 grs.**Camphor and Opium, 1-4 gr.**

Camphor, 1-6 gr.

Powd. Opium, 1-12 gr.

Camphor and Opium, 3 gr.

Camphor, 2 grs.

Powd. Opium, 1 gr.

Diaphoretic, 1 1-2 gr.

Morphine acetate, 1-25 gr.

Powd. Potass. nitrate, 1 gr.

Powd. Ipecac, 1-4 gr.

Powd. Camphor, 1-4 gr.

Consult Nos. 87-90, 457, in appended Pill Formulæ list for other pills containing camphor.

Soap, Medicated Camphor—Containing 10 per cent. or 168 grains of Camphor. This soap has primarily a stimulant, and at times an irritant action, which is succeeded by an anodyne effect. It is most frequently used for the relief of pruritus that attends eczema, chilblains, and other irritable affections of the skin.

CANADA THISTLE. *Cnicus arvensis, Hoffmann.* (Rhizome.)*Synonym.*—*Cirsium arvense, Scopol.*

Properties.—Popularly employed for its astringent action in diarrhœa, dysentery and leucorrhœa.

Canada Thistle root, compressed in pound packages for retailing purposes.

CANADIAN HEMP. *Apocynum, U. S. P.* *Apocynum cannabinum, Linné.* (Root.)

Synonyms.—Black Indian Hemp; Chanvre du Canada, Fr; Canadische Hanfwurzel, Gr.

Properties.—Powerfully emetic and cathartic, possessing marked diaphoretic, expectorant and diuretic powers. It produces much nausea, diminishes the frequency of the pulse, and appears to induce drowsiness. Acting as a powerful hydragogue cathartic, it is of value in dropsy, especially the anasarca of Bright's disease and in ascites, but is recommended also in intermittent and remittent fevers, and pulmonary affections. Dr. Cawthorn considers its anti-periodic powers scarcely inferior to those of quinine.

NOTE.—The name Indian Hemp ought not to be applied to this drug, since this name is officially recognized as the designation of *Cannabis Indica* (U. S. P., 1880). It is better, in prescribing either of these drugs, to write the Latin name in full.

Canadian Hemp, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Canadian Hemp.—Not miscible with water.

Dose, 5 to 30 minims (0.32 to 1.2 C. c.); and as an emetic, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Powdered Extract Canadian Hemp.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Canadian Hemp.

Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Solid Extract Canadian Hemp.—One part equals 5 of root.

Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Pills.

Apocynum Extract, 2 grs.

Eucalyptus Comp., 1 5-8 gr.

Ext. Eucalyptus, 1 gr.

Sanguinaria, 1-8 gr.

Ext. Canadian Hemp, 1-2 gr.

See also No. 36 in Pill Formulæ list appended.

CANELLA. *Canella alba, Murray.* (Bark.)

Properties.—Aromatic and stimulant. Used as an adjuvant to tonics and purgatives.

PREPARATION AND FORMULA.

Fluid Extract Canella.—Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Tincture Aloes and Canella (*Hiera Pica*).

Fluid Aloes, P. D. & Co.'s 2 fl. ozs.

Fluid Extract Canella, P., D. & Co.'s 1 fl. oz.

Dilute Alcohol 13 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

CANTHARIDES. *Cantharis, U.S.P. Cantharis vesicatoria, De Geer.*
(The insects.)

Properties.—Poisonous. Internally, *Cantharides* is a powerful stimulant, with a peculiar direction to the urinary and genital organs. In moderate doses it sometimes acts as a diuretic, and generally excites some irritation in the urinary passages, which, if its use be persevered in, often amounts to violent strangury. In larger doses it is a violent poison, 20 grains of the powder having been known to cause death. Medicinally it has been used with success in the treatment of dropsy, amenorrhœa, obstinate gleet, leucorrhœa, seminal weakness, diabetes and incontinence of urine. Externally it is only used for blistering purposes.

Antidotes.—Emetics or the stomach-pump, promptly; opiates by the stomach and rectum; animal charcoal in fine powder, mucilaginous or aluminous drinks. To relieve strangury, warm sitz baths, etc.

PREPARATIONS AND FORMULÆ.**Cerates.**

Cantharides, U. S. P. (*Emplastrum Cantharides, B. P.*)

Cantharides Extract, U. S. P.

Collodion Cantharidal, U. S. P. (*Blistering Collodion.*)

Each fluidounce represents 240 grs. cantharides.

Fluid Extract Cantharides.—Used externally as a blistering fluid; the Tincture only is suited for internal use, see below.

Liniment Cantharides, U. S. P.

Fluid Extract Cantharides, P., D. & Co.'s 1 fl. oz.

Oil Turpentine 7 fl. ozs.

Ointment Cantharides, U. S. P., 1870.

A dilution (1 in 3) of *Cerate Cantharides Extract, U.S.P., 1880*, is a much more elegant substitute.

Pills.

Cantharides, 1-50 gr.

Phosphorus and Cantharides Comp., 2 grs.

Phosphorus, 1-50 gr.

Powd. *Nux vomica*, 1 gr.

Cantharides, 1 gr.

Phosphorus and Cantharides Comp., Formula "B." 2 grs.

Phosphorus, 1-33 gr.

Powd. *Nux vomica*, 1 gr.

Cantharides, 1 gr.

Plasters.

Calafacens (Emp. Picis c. Cantharide, U. S. P.).—In half pound rolls or cans.

Issue Plasters.—A valuable counter-irritant which may be used in acute cases as a simple epispastic, but superior to the common fly blister in their remarkable uniformity of action and non-liability to produce strangury, nor to exhibit unwonted harshness.

In chronic cases the plasters may be applied daily to the same spot, thus maintaining with little if any pain, an issue or suppurating ulcer. This is the use to which they are particularly adapted.

Cantharides (active constituent) 6 parts.

Camphor (active constituent) 6 parts.

Powd. Opium (active constituent) 2 parts.

Ext. *Mezereum* (active constituent) 4 parts.

White Wax and Cocoa butter, q. s. to make 42 parts.

CANTHARIDES—Continued.**Tincture Cantharides, U. S. P.**

Fluid Extract Cantharides, P., D. & Co.'s.....1 fl. ozs.

Alcohol.....23 fl. ozs.

Dose, 3 to 20 minims (0.2 to 1.3 C. c.) employed cautiously.

NOTE.—This is 4 times the strength of the B. P. tincture, but only half the strength of the P. G. and Fr. Cod. tinctures.

CAPSICUM, U. S. P. Capsicum fastigiatum, *Blume*. (Fruit.)*Synonyms*.—Bird Pepper, Cayenne Pepper, Red Pepper; Spanischer Pfeffer, Ger.*Properties*.—Stimulates the mucous lining of the stomach, and promotes secretion of gastric juice. It is useful in cases of enfeebled and languid stomach, dyspepsia, atonic gout attended with flatulence and in flatulent colic. An excellent addition to quinine in intermittents; used with some success in Asiatic cholera; and is a sovereign remedy in delirium tremens. Externally it is a rubefacient.**PREPARATIONS AND FORMULÆ.****Collodion Capsicum**.—Each fluidounce represents 16 grs. Capsicum.**Extracts.****Fluid Extract Capsicum, U. S. P.**—Not miscible with water.

Dose, 5 to 15 minims (0.3 to 1 C. c.), diluted with sweetened water.

Solid Extract Capsicum.—One part equals 4 of the drug.

Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Pills.**Anti-Periodic**, 2 grs.

Cinchonidine sulph., 1 gr.

Iron sulph. exsic., 1-2 gr.

Podophyllin, 1-20 gr.

Strychnine sulph., 1-33 gr.

Gelsemerin, 1-20 gr.

Oleoresin Capsicum, 1-10 gr.

Capsicum, 1 gr.**Capsicum Extract**, 1-2 gr.

Consult Nos. 17, 28, 83, 95, 100, 260, 261, 356, 446, in Pill Formulæ list appended for other pills containing capsicum.

Tablet Triturate, Capsicum, 1 gr.**Tincture Capsicum, U. S. P.**

Fluid Extract Capsicum, P., D. & Co.'s.....1 fl. oz.

Alcohol.....22 fl. ozs.

Water.....1 fl. oz.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

NOTE.—Half the strength of the P. G. tincture.

CARAWAY. Carum, U. S. P. Carum Carvi, *Linné*. (Seed.)*Properties*.—Chiefly used as a flavoring agent, but also efficient in the flatulent colic of children.**PREPARATION.****Fluid Extract Caraway seed**.—Not miscible with water.

Dose, 20 to 60 minims (1.3 to 4 C. c.).

CARDAMOM. Cardamomum, U. S. P. Elettaria Cardamomum, *Maton*. (Seeds.)*Properties*.—An agreeable carminative and stomachic. Frequently added to tonic and stimulant preparations but chiefly employed as a flavoring agent.**PREPARATIONS AND FORMULÆ.****Extracts.****Fluid Extract Aromatic Powder**.—Not miscible with water.

Each pint represents: Cinnamon and Ginger, 5 3-4 ozs. each; Cardamom and Nutmeg, 2 1-2 ozs. each.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

CARDAMOM—Continued.

Fluid Extract Cardamom seed.—Not miscible with water.
Dose, 5 to 15 minims (0.3 to 1 C.c.).

Fluid Extract Cardamom Comp.—Not miscible with water.
Sixteen fluidounces, imperial, represent: Cardamom seeds and Cinnamon, 2 1-2 ozs. each; Caraway, 1-4 oz.; Cochineal, 5-8 oz.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Tinctures.**Cardamom, U. S. P.**

Fluid Extract Cardamom Seed P., D. & Co.'s..... 2 fl. ozs.
Alcohol..... 9 fl. ozs.
Water..... 3 fl. ozs.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Cardamom Compound, U. S. P.

Fluid Cardamom Comp., P., D. & Co.'s..... 2 fl. ozs.
Dilute Alcohol..... 13 fl. ozs.
Glycerin..... 1 fl. oz.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).
A favorite remedy for flatulent colic.

CARICA PAPAYA, Linné.

Synonyms —Melon tree, Papaw (true Papaw, and wholly different from the common Papaw, *Asimina triloba*, of the southern States).

Properties.—The inspissated juice of the unripe fruit possesses the remarkable power of digesting albuminous substances just as pepsin does, and it is employed as a substitute therefore, especially in the treatment of diphtheria. A committee of the Royal Society of New South Wales appointed to investigate the properties of this drug report recently,* and claim without reserve, the greatest efficiency for carica papaya as a galactagogue. Their experiments show that the mammary secretion could be stimulated, in cases where the flow had commenced to fail, either by internal administration of the drug or by topical application. They also report noticing its action as an abortifacient when administered to pregnant patients; its use therefore requires caution.

PREPARATIONS.

Papain.—(Papaw juice.) A true digestive ferment. A ten-per-cent. solution in water and glycerin, equal parts, applied to the diphtheritic membrane with a camel's-hair pencil frequently causes it to speedily soften and become detached. It may be employed to assist digestion, and also as a galactagogue.

Dose, 1 to 5 grains (0.065 to 0.35 Gm.).

Papayotin.—The active principle of Papaw juice. Its uses are the same as those of the crude juice.

For use in diphtheria, mix one part of papayotin with 4 parts of water and 4 parts of glycerin, and apply hourly with brush or feather.

Dose, 1-4 to 1 grain (0.016 to 0.065 Gm.).

GARNAUBA. Copernicia cerifera, Martius. (Root.)

Properties.—Alterative. Used in Brazil, to which country it is indigenous, for the same purposes for which Sarsaparilla, Stillingia, etc., are employed with us. It is, however, regarded as a more pronounced alterative than these.

PREPARATION.

Fluid Extract Carnauba.—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

CAROBA. Jacaranda procera, Sprengel. (Leaves.)

Properties.—A valuable and effective anti-syphilitic. It is said to possess alterative, diuretic, sudorific and tonic properties, to a marked degree.

Hence its value in the treatment of syphilis in all its forms.

* Australian Medical Gazette, Feb., 1887 (Therapeutic Gazette, May 16, 1887.).

CAROBA—Continued.

Caroba may be combined with iodide of potassium with good results, in cases distinguished by pains in the limbs. Its principal effect seems to be directed to the cure of old syphilitic ulcers, to which it may be applied topically, in addition to internal treatment.

PREPARATION.

Fluid Extract Caroba.—Not miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

CARPENTER'S SQUARE. *Scrophularia nodosa*, *Linné*. (Herb.)

Synonyms.—Figwort; Knotenwurz, Ger.

Properties.—Alterative, diuretic and anodyne. It is highly beneficial in hepatic diseases, scrofula, cutaneous diseases, dropsy and as a general deobstruent to the glandular system. The warm infusion given freely is said to restore lochial discharges when suppressed, and to relieve the pains attending difficult menstruation.—King.

Carpenter's Square (Figwort) herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Carpenter's Square (Figwort).—Miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

CARRION-FLOWER. *Smilax herbacea*, *Linné*. (Leaves.)

Synonym.—Jacob's Ladder.

Properties.—Alterative. Popularly employed in scrofulous diseases, and as a general purifier of the blood.

Dose, a wineglassful (60 C. c.), of an ounce to the pint infusion.

Carrion-flower (Jacob's Ladder) leaves, compressed in ounce packages for retailing purposes.

CASCARA AMARGA. *Picramnia* (Sp. indetermined). (Bark.)

Synonym —Honduras bark.

Properties.—Alterative; credited with achieving cures in chronic cases of syphilitic eruptions, syphilitic tubercles, chronic eczema and gummy tumors. The discovery, in our laboratory, of an alkaloid in this drug, confirms us in our belief that it is in reality a valuable remedy, and the clinical evidence in its favor is accumulating.

"Though it is a drug of slow action, it is almost a specific for syphilis. The remedy seems to eliminate the specific poison from the system. It is equally as good in chronic nephritis"*

PREPARATIONS.**Extracts.**

Fluid Extract Cascara Amarga.—Not miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Powdered Extract Cascara Amarga. One part equals 5 of bark; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Cascara Amarga.
Dose, 5 to 10 grams (0.3 to 1 Gm.).

CASCARA SAGRADA. *Rhamnus Purshiana*, *DeCandolle*. (Bk.)

Synonyms.—Sacred Bark, Sacred Tree Bark.

Properties.—Laxative, cathartic and intestinal tonic. This drug has been aptly called a "Tonic Laxative," because of its physiological action, which seems to be directed, as a tonic, to the vaso-motor nervous system,

* Morse: "New Therapeutical Agents," p. 179.

CASCARA SAGRADA—Continued.

stimulating the secretory apparatus of the alimentary canal, and increasing the peristaltic action of the intestines, thus restoring to the bowels their normal activity. Its use is not attended with the unpleasant after effects following the employment of cathartics; neither does it, as a rule, cause griping or other unpleasant symptoms. It is as nearly a specific for chronic constipation as any article of the materia medica is a specific for any disease. To be of service, however, it must not be given in cathartic doses. It relieves this condition indirectly by restoring tonicity to the bowels.

"The effects are quite satisfactory this new medicine deserves to retain a place in therapeutics as a remedy in habitual constipation." *

PREPARATIONS.

Capsules, in soft gelatin.

Cascara Sagrada, Solid Ext., 2 gr., Castor oil, q. s. 10 minims.

Cascara Sagrada, Solid Ext., 3 gr., Castor oil, q. s. 10 minims.

Cascara Cordial.—An elegant and palatable combination for the exhibition of the remedies composing it, and, unlike many preparations of this character, intended solely for use on the prescription of the physician. The great desideratum in the treatment of chronic constipation has heretofore been a remedy which possessed not only a temporary specific action on the bowels, but which would be followed by no reaction and no repression of secretion, and which could be administered for a sufficient length of time to permit of restoration of normal function. This preparation of *Cascara sagrada* fully meets these indications.

Each litre contains: *Cascara sagrada* bark (true *Rhamnus Purshiana*), 100 grms.; *Berberis aquifolium*, true, 37 grms.; Diluted Alcohol, 233 grms.; Sugar, 288 grms.; Water, q. s. ad. 1 litre. Elegantly flavored with carminative aromatics.

Dose, as a laxative, 1 fluidrachm (4 C. c.) twice a day, night and morning; as a cathartic, 1-2 fluidounce (15 C. c.) twice a day, night and morning; for constipation and dyspepsia, from 1-4 to 1 fluidrachm (1 to 4 C. c.) four times a day for a week. If no improvement ensues, increase the dose till a normal or laxative passage is obtained, when the dose should be gradually decreased in amount and frequency until the benefit is permanent.

Cascara Cordial has also become widely known to the profession as a vehicle for nauseous medicines. It not only effectually disguises the taste of such inherently bitter medicines as quinine, but, on account of its laxative properties, is a most valuable adjuvant or corrigent to any prescription containing drugs, such as iron or opium, the continued administration of which tends to constipate.

Elixirs.

Cascara Sagrada, 120 grs. (in each fluidounce).

Cascara Sagrada Compound.

Cascara sagrada, 60 grs.
Licorice, 60 grs.

Berberis aquifolium, 30 grs.
Aromatics, q. s.

Cathartic Compound.

Senna, 16 grs.
Cascara sagrada, 32 grs.
Culver's root, 16 grs.

Butternut, 16 grs.
Rochelle salts, 32 grs.

Extracts.

Fluid Extract Cascara Sagrada, B. P. (*Rhamnus Purshiana*.)

—Not miscible with water.

- Dose, as a laxative, 3 to 15 minims (0.2 to 1 C. c.), 3 times a day; as a cathartic, 20 to 45 minims (1 3 to 3 C. c.), morning and evening; as a stomachic, 4 to 10 minims (0.3 to 0.6 C. c.), 4 times a day.

Malt Extract with Cascara Sagrada. (60 grs. in each fl. oz.)

Malt Ext. with Cascara Sagrada and Berberis Aquifol.

Each fluidounce represents: *Cascara sagrada*, 30 grs.; *Berberis aquifolium*, 30 grs.

* Dujardin-Beaumetz: "New Medications," p. 106 (edition of G. S. Davis, 1886.).

CASCARA SAGRADA—Continued.

Powdered Extract Cascara Sagrada.—Of same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing moisture with powdered sugar of milk. Dose, as a laxative, 1 to 2 grains (0.06 to 0.13 Gm.); as a cathartic, 3 to 8 grains (0.2 to 0.5 Gm.).

Solid Extract Cascara Sagrada, B. P.—One part equals 4 of the bark.

Dose, as a laxative, 1 to 3 grains (0.06 to 0.2 Gm.); as a cathartic, 3 to 8 grains (0.2 to 0.5 Gm.).

Pills.

Aloin, Strych. and Belladonna Comp., 2-3 gr.

Aloin, 1-5 gr.

Strychnine, 1-60 gr.

Ext. Belladonna, 1-8 gr.

Ext. Cascara sagrada, 1-2 gr.

Anti-Constipation, Carson's, 2-3-4 grs.

Ext. Cascara sagrada, 1 gr.

Ext. Rhubarb, 1 gr.

Ext. Nux vomica, 1-4 gr.

Aloin, 1-3 gr.

Anti-Constipation, Goss', 2 1-4 gr.

Podophyllin, 1-4 gr.

Ext. Coloc., 1-4 gr.

Ext. Butternut, 1-2 gr.

Ext. Nux vomica, 1-8 gr.

Ext. Henbane, 1-8 gr.

Ext. Gentian, 1-4 gr.

Ext. Cascara sagrada, 1-4 gr.

Powd. Canadian Hemp, 1-2 gr.

Berberis Compound, 2 grs.

Ext. Berberis aquifolium, 2 grs.

Ext. Cascara sagrada, 1 gr.

Cascara Comp. Pellets, D'Ary's, 1 gr.

Cascara sagrada, 4-15 gr.

Ext. Nux vomica, 1-30 gr.

Ext. Belladonna, 1-60 gr.

Euonymin, 1-5 gr.

Xanthoxylin, 4-15 gr.

Oleo-resin Capsicum, 1-20 gr.

Cascara Sagrada Extract, 2 grs.

Cascara Sagrada and Nux vomica, 2 1-5 grs.

Ext. Cascara sagrada, 2 grs.

Ext. Nux vomica, 1-5 gr

Cascara Sagrada, Nux vomica and Belladonna, 2 1-4 grs.

Ext. Cascara sagrada, 2 grs.

Ext. Nux vomica, 1-8 gr.

Ext. Belladonna, 1-16 gr.

CASCARILLA, U. S. P. Croton Eluteria, *Swartz.* (Bark)

Properties.—Aromatic, stimulant, tonic and febrifuge. Used advantageously with Cinchona in intermittent fever, in epidemic dysentery, and as an ingredient in tonic bitters. "Much used in Germany in the advanced stages of diarrhoea" *

Cascarilla bark, U.S.P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Cascarilla.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 teaspoonful (2 to 4 C.c.).

NOTE —Incompatible with lime water and mineral acids.

Solid Extract Cascarilla.—One part equals 6 of bark.

Dose, 5 to 10 grains (0.3 to 0.64 Gm.).

Tincture Cascarilla, B. P.

Fluid Extract Cascarilla, P., D. & Co.'s.....1 fl. oz.

Dilute Alcohol.....7 fl. ozs.

CASSIA. Cinnamomum, U. S. P. Cinnamomum of other species than *C. zeylanicum*, *Breyne.* (Bark.)

Synonym.—Chinese Cinnamon.

Properties.—Highly aromatic and somewhat astringent. Much used to check diarrhoea in combination with opium, chalk, or some vegetable infusion. Efficacious in internal hemorrhage.

* Napheys: "Medical Therapeutics," p. 354.

CASSIA—Continued.

PREPARATIONS AND FORMULA.

Extracts.**Fluid Extract Aromatic, U. S. P.**—Not miscible with water.

Each pint represents: Cinnamon and Ginger, 5 3-4 ozs. each; Cardamom and Nutmeg, 2 1-2 ozs. each.

Dose, 10 to 30 minims (0.65 to 2 C. c.).

Fluid Extract Cassia.—Not miscible with water.

Dose, 10 to 20 minims (0.65 to 1.25 C. c.).

Tincture Cinnamon, U. S. P.

Fluid Extract Cassia, P., D. & Co.'s.....2 fl. ozs.

Alcohol.....13 fl. ozs.

Water.....7 fl. ozs.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

CASTOR OIL PLANT. *Ricinus communis, Linné.* (Seeds; oil, U. S. P., leaves)*Synonyms.*—Palma christi; Ricin, Fr.*Properties.*—The leaves are galactagogue; used to accelerate or increase the secretion of milk in cases where it is tardy in appearing or deficient in quantity. They may be applied to the breast in poultice and a decoction or the fluid extract given internally at the same time, although Dr. D. S. Clark, of Rockford, Ill., states that he finds the fluid extract equally efficacious when administered alone.

The beans or seeds are powerfully cathartic in large doses, and sometimes emetic. The alcoholic fluid extract of the seeds is said to be superior to Castor Oil, as it acts in smaller doses, is more agreeable to the taste, and has less disposition to irritate the bowels, or to cause vomiting. It may be substituted for the oil in cases where the action of the alcohol is not detrimental.

Castor leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Capsules.—An efficient means of administering this otherwise "best of simple purgatives" without offending the palate.**Castor Oil,** 10 minims. In both soft and hard gelatin.**Castor Oil and Podophyllin,** in both soft and hard gelatin.

Castor Oil, 10 minims. Podophyllin, 1-8 grain.

Castor Oil, 2 1-2, 5, 10, and 15 grms. In soft gelatin.**Castor Oil and Oil of Fleabane,** in soft gelatin.

Castor Oil, 5 minims. Oil Fleabane, 5 minims.

Male-fern and Castor Oil, 2 1-2 grms.

Oleoresin Male-fern, 1 grm. Castor Oil, 1 1-2 grm.

Santonin and Castor Oil, 2 1-2 grms.

Santonin, 1-4 gr. Castor Oil, 2 1-2 grm.

Extracts.**Fluid Extract Castor beans.**—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Castor leaves.—Miscible with water without material precipitation.

Dose, 1 fluidrachm (4 C. c.).

CATECHU, U. S. P. An extract from the wood of *Acacia Catechu, Willdenow.**Synonyms.*—Cutch, Terra japonica, Japonica; Cachou, Fr.; Katechu, Ger.*Properties.*—Gently tonic and powerfully astringent. Employed locally as an application to relaxed or congested mucous membranes accompanied by excessive discharge, and for checking hemorrhages. Combined with opium it answers a good purpose as an internal remedy in chronic diarrhœa, chronic catarrh, colliquative diarrhea and chronic dysentery.

CATECHU—Continued.**PREPARATIONS AND FORMULÆ.****Fluid Catechu.**—Not miscible with water.

Each pint represents 8 1-2 ozs., av., of the crude drug—one-half the strength of a fluid extract.

Dose, 10 to 45 minims (0.6 to 3 C.c.).

Lozenge, Catechu, 2 grs.**Tinctures.****Catechu, Phar. Ger. and Fr. Cod.**

Fluid Catechu, P., D. & Co.'s..... 2 fl. ozs.

Dilute Alcohol..... 3 1-2 fl. ozs.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Catechu Compound, U. S. P.

Fluid Catechu, P., D. & Co.'s..... 3 fl. ozs.

Fluid Extract Cinnamon, P., D. & Co.'s..... 1 fl. oz.

Dilute Alcohol..... 8 fl. ozs.

Dose, 30 minims to 3 fluidrachms (2 to 12 C.c.).

NOTE.—The B. P. tincture is almost identical.**CATNEP.** *Nepeta cataria, Linné.* (Leaves and Tops.)*Synonyms.*—Catmint; Cataire, Fr.; Katzenminze, Ger.*Properties.*—Antispasmodic and slightly tonic. Useful in nervous febrile derangements. Much used for the relief of flatulent colic in infants, and as a popular remedy in retarded or painful menstruation.**Catnep leaves,** compressed in ounce packages for retailing purposes.**PREPARATION.****Fluid Extract Catnep.**—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

CEDRON SEED. *Simaba Cedron, Planch.* (Seeds.)*Properties.*—Poisonous. Cerebral sedative, anti-spasmodic, anti-periodic; long valued in Central America and adjacent countries as a remedy for the bite of poisonous serpents and insects. Also used successfully in intermittent fevers, spasm of the stomach and bowels, and neuralgia of the face.* Dr. F. B. Thompson, of London, has found it useful in gout.**PREPARATION.****Fluid Extract Cedron seed.**—Not miscible with water.

Dose, 1 to 8 minims (0.06 to 0.5 C. c.).

For serpent bites it is recommended in six-drop doses, and also to dress the bite with fluid extract.

CELERY. *Apium graveolens, Linné.* (Seeds.)*Properties.*—Carminative, stimulant, and slightly aperient. Used successfully, combined with guarana, in the treatment of sick-headache (migraine). Also used as a flavoring agent.**PREPARATIONS.****Elixir Celery and Guarana;** 60 grains of each to the fluidounce.

Dose, 1 fluidrachm (4 C.c.).

Extracts.**Fluid Extract Celery.**—Not miscible with water.

Dose, 5 to 15 minims (0.3 to 1.25 C. c.).

Solid Extract Celery.—One part equals 5 of seed.

Dose, 1 to 3 grains (0.065 to 0.2 Gm.).

* Morse, "New Therapeutical Agents," p. 171.

CEVADILLA. *Veratrum Sabadilla*, *Schlechtendal*. (Seeds.)

Synonyms.—*Asagroea officinalis*, *Lindley*; *Schoenocaulon officinale*, *Gray*.

Properties.—Poisonous. A drastic and irritant cathartic. It has been used in articular rheumatism, dropsies, and as a cardiac sedative, but it is somewhat uncertain in action and requires to be exhibited with caution. Externally it finds favor in superficial neuralgias, myalgia, sciatica, and in some forms of pruritus.

Antidotes.—None reliable known. Use emetics (zinc sulphate, apomor; phine, mustard), stimulants, coffee; combat symptoms as they arise—recumbent position.

ACTIVE PRINCIPLE.

Veratrine.—This alkaloid is employed almost exclusively in the form of an oleate, as a topical application in neuralgias, especially in sciatica. Dose, 1-40 grain (0.0015 Gm.) cautiously increased.

PREPARATIONS.

Fluid Extract Cevadilla.—Not miscible with water.

Dose, 2 to 4 minims (0.13 to 0.26 C. c.).

Oleate Veratrine, containing 10 per cent. alkaloid.

Pills, Veratrine, 1-60, 1-32 and 1-12 gr.

Tablet Triturates, Veratrine, 1-32 and 1-12 gr.

CHAMOMILE. *Anthemis*, U. S. P. *Anthemis nobilis*, *Linné*. (Flwrs.)

Synonym.—Roman Chamomile.

Properties.—Anti-spasmodic, stomachic and tonic. It is useful in the spasmodic and pseudo-neuralgic affections of hysterical women. The warm infusion is largely employed as a stimulant diaphoretic in incipient inflammations of mucous tracts, and as an emmenagogue when the flow is absent or scanty owing to cold.

PREPARATIONS.**Extracts.**

Fluid Extract Chamomile.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Solid Extract Chamomile.—One part equals 4 of flowers.

Dose, 8 to 12 grains (0.5 to 0.8 Gm.).

Pill, Anthemis Extract, 2 grains.

Soaps, Medicated.

Chamomile Soap, containing 10 per cent. or 168 grs. Ext. Chamomile. A mild stimulant and astringent, and particularly valuable both for its medicinal virtues and aromatic odor, in all conditions of excessive secretion and ill-smelling sores. An excellent soap to use in intertrigo or chafing, and in dermatitis, seborrhœa, hyperidrosis and bromidrosis.

Chamomile and Sulphur Soap, containing 10 per cent. or 168 grs. Ext. Chamomile, and 5 per cent. or 84 grs. Sulphur. Uses same as above, but frequently more effective by the addition of the sulphur. The greasy, shining complexion attended with numerous small elevations and pimples, a form of acne common to young women, from either gastric or menstrual disorders, often yields to applications of it, and it assists in restoring the skin to its healthy, transparent state.

CHARCOAL. Wood charred by exposure to a red heat without access of air.

Properties.—Absorbent of intestinal gases, antiseptic and deodorant. Useful in flatulent dyspepsia, epidemic dysentery, gastralgia, etc.

Dose, 20 grains (1.4 Gm.) and upwards.

CHARCOAL—Continued.**PREPARATIONS.****Pill, Charcoal,** (Willow) 3 grs.**Lozenges.****Bismuth and Charcoal,** 2 grs.

Bismuth subnitrate, 1 gr

Charcoal, (willow) 1 gr.

Charcoal, (Willow) 4 grs.**CHAULMOOGRA OIL.** The fixed oil expressed from the seeds of *Gynocardia odorata*, *R. Brown.*

Properties.—This oil is said to be the most powerful alterative known to medical science, and the only reliable remedy for leprosy. In England it has been found of great service in many complaints, especially rheumatism, serofulous swellings, sprains, lupus and various skin diseases.

"Is a very successful remedy in eczema of the third stage."*

Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

PREPARATION.**Capsules, Chaulmoogra Oil,** 10 minims. In soft gelatin.

A method of internal administration which does not offend the palate.

CHEKAN. *Eugenia chequen*, *Hooker & Arnott.* (Leaves.)

Synonyms.—Cheken, Myrtus Chekan, *Molina.*

Properties.—Introduced from Chili as a remedy in chronic catarrhal inflammation of the respiratory mucous lining. Dr. Murrell, of London, especially commends it in winter cough, that annoying and intractable affection of elderly people. Dr. Dessaur, of the German hospital of Valparaiso, speaks highly of its effects in the purulent form of bronchitis, and others who have tested its virtues, recommend it as a valuable addition to the list of remedies in chronic bronchitis.

"A tonic expectorant with diuretic qualities."†

PREPARATION.**Fluid Extract Chekan.**—Not miscible with water.

Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

CHESTNUT. *Castanea*, U. S. P. *Castanea vulgaris*, *Lamarck*, var. *americana*, *A. DeCandolle.* (Leaves.)

Synonyms.—Châtaignier, Fr.; Kastanienblätter, Ger.

Properties.—An efficient remedy in relieving the symptoms of whooping cough. It not only alleviates the severity of attacks, but even breaks up the paroxysms, leaving merely a cough, attended with mucous expectoration, which gradually yields to ordinary expectorants.

Chestnut leaves, U. S. P., compressed in ounce packages for retaining purposes.

PREPARATION.**Fluid Extract Chestnut leaves, U. S. P.**—Miscible with water.

Dose, 5 to 60 minims (0.33 to 4 C.c.).

CHEWSTICK. *Gouania domingensis*, *Linné.* (Bark.)

Properties.—Expectorant, tonic, laxative. A grateful aromatic, bitter tonic, useful in cases of debility, as a stomachic in feeble digestion, and also in gonorrhœa, from its tonic effect on the mucous membranes.

Diluted with water it may be used as a mouth-wash in cases of salivation,

* Dymock, "Materia Medica of Western India, p. 727.

† Napheys, "Medical Therapeutics," p. 194.

CHEWSTICK—Continued.

etc., being at once astringent and soothing in its effect; or as a dentifrice, imparting tone to the gums.

PREPARATION.

Fluid Extract Chewstick.—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

CHIAN TURPENTINE. Oleoresin of *Pistacia terebinthus*, *Linné*.

Properties.—Claimed to cure "cancer of the generative organs,"* and also mammary cancers. "It appears to act upon the periphery of the growth with great vigor, causing the speedy disappearance of what is usually termed the cancerous infiltration."

Dose, 5 to 20 grains (0.3 to 1.3 Gm.).

PREPARATION.

Emulsion Chian Turpentine.—Prepared after the formula of Prof. Clay, its introducer, as a remedy in cancer. It is less liable to disturb the stomach in emulsion than when given in pill form, and is moreover more readily absorbed. Contains 5 grains Chian Turpentine in the fluidounce. Dose, 1 fluidounce (30 C.c.) three times a day.

CHICKWEED. *Stellaria media*, *Smith*. (Herb.)

Synonyms.—*Stellaire*, Fr.; *Sternkraut*, Ger.

Properties.—Demulcent and emollient. Used externally in the form of poultice in ophthalmia and in bruises, inflammations, etc.

Chickweed herb, compressed in ounce packages for retailing purposes.

CHIRATA, U. S. P. *Ophelia Chirata*, *Grisebach*. (Herb.)

Properties.—A simple bitter tonic. It is recommended in dyspepsia, in the debility of convalescence, and generally in cases where corroborant measures are indicated. In Hindoostan it is held to be stomachic, tonic, febrifuge, cholagogue and deobstruent, as well as efficient in chronic bronchitis.

NOTE.—One of the drugs which may be prescribed with iron.

PREPARATION AND FORMULA.

Fluid Extract Chirata, U. S. P.—Miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

Tincture Chirata, U. S. P.

Fluid Extract Chirata, P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

CHOKE CHERRY. *Prunus virginiana*, *Linné*. (Bark of tree.)

NOTE.—Not the *P. virginiana* of the U. S. P.; see Wild Cherry.

Properties.—Tonic, anti-spasmodic, mildly astringent, and anti-periodic. Employed in the treatment of intermittent fevers, and as a tonic in all conditions of debility, convalescence from acute diseases, etc.

PREPARATION.

Cerasin, a so-called **Concentration**,† a powdered preparation from Choke Cherry bark.

Synonym.—*Cerascin*.

Dose, 2 to 8 grains (0.13 to 0.52 Gm.).

* *Lancet*, March 27; *Therapeutic Gazette*, 1880, p. 210.

† See note in Preface on Concentrations.

CINCHONA. Pale. *C. officinalis*, *Hooker*; Red, *Cinchona rubra*, U. S. P., *C. succirubra*, *Pavon*; Yellow, *Cinchona fiava*, U. S. P., *C. Calisaya*, *Weddell*. (Bark)

Synonyms.—Pale—Crown bark, Loxa bark, Pale Peruvian bark; Braune (grau) Chinarinde, Ger.

Red—Red bark; Quinquina rouge, Fr.; Rothe Chinarinde, Ger.

Yellow—Yellow Cinchona; Quinquina jaune, Fr.; Königschina, Ger.

Properties.—Tonic, febrifuge, anti-periodic. A tonic having a wide range of use, especially indicated wherever there is any malarious influence at work, and in convalescence from acute diseases. All the symptoms which result from marsh miasm are amenable to treatment by cinchona and its derivatives.

Any species is official which contains 3 per cent. total alkaloids.

Pale bark contains most Cinchonine and least Quinine; Yellow bark contains most Quinine and least Cinchonine, while Red bark contains these in nearly equal proportions.

NOTE.—Cinchona salts are precipitated from solution by alkalis.

ACTIVE PRINCIPLE (derivative).

(For others see preparations below.)

Quinine and Urea Muriate.—(Quinia bimuriatica carbamidata.)

This double salt is remarkable for its ready solubility in water, hence it has been selected for hypodermic use. The concentrated aqueous solution contains a grain in each minim, and, being perfectly neutral, causes comparatively little irritation. The effects produced are precisely those of the sulphate of quinine, and it is employed in the same doses.

Dose, 1 to 10 grains (0.065 to .65 Gm.).

PREPARATIONS AND FORMULÆ.

Capsules, in soft gelatin.

Cinchonidine Sulphate, 1, 2, 3, 4 and 5 grs.

Quinine Sulphate, 1, 2, 3, 4 and 5 grs.

“**Warburg’s Tincture**,” concentrated, 10 minims representing 2 fluidrachms of tincture.

Elixirs, each fluidounce represents ingredients stated.

Calisaya bark, 40 grs.

Calisaya and Iron Pyrophosphate.

Calisaya bark (alkaloids), 40 grs. Iron pyrophosphate, 8 grs.

Calisaya, Iron and Strychnine.

Calisaya bark (alkaloids), 40 grs. Iron pyrophosphate, 8 grs.

Strychnine, 8-100 gr.

Iron, Quinine and Strychnine Phosphates.

Iron phosphate, 16 grs. Strychnine phosphate, 8-60 gr.

Quinine phosphate, 4 grs.

Owing to the large number in which cinchona or its derivatives enter it is impossible to name here more than these leading elixirs; for others see appended Elixir Formulæ list numbers 9, 19, 21-23, 25-31, 33-39, 44, 49-52, 79, 80, 82, 84-88, 90, 111, 114, 119, 125, and 129-132.

Extracts.

Fluid Extract Cinchona Aromatic.—Not miscible with water.

Sixteen fluidounces, imperial, represents: Cinchona bark, 13 ozs; Sweet Orange peel, 2 1-6 ozs; Cardamom seeds and Cassia, 1-3 oz. each; Caraway seeds, 1-6 oz.

Combined with pleasant aromatics to modify the intense bitterness; can be used wherever the fluid extract of the bark is indicated.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Cinchona, Pale.—Not miscible with water.

Said to be superior to either Calisaya or Red bark as a tonic, being less obnoxious to the stomach and intestines.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Cinchona, Pale, Comp.—Not miscible with water.

Each pint represents: Cinchona bark (pale), 8 ozs.; Bitter Orange peel, 6 ozs.; and *Serpentaria*, 1 1-2 oz.

Suited for the extemporaneous preparation of Huxham’s Tincture.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

CINCHONA—Continued.

Fluid Extract Cinchona, Red.—Not miscible with water.

Prepared from the selected, true Red Cinchona.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Cinchona, Red, Comp.—Not miscible with water.

Each pint represents: Red Cinchona bark (true), 7 1-2 ozs.; Bitter Orange peel, 6 ozs.; and Serpentaria, 1 1-2 oz.

Five times the strength of the Compound Tincture Cinchona. U. S. P. No preparation of Cinchona bark is more in demand than the compound tincture. According to the U. S. Pharmacopœia, 1880, this is now made from Red Cinchona bark.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Cinchona, Red, Detannated.—Not miscible with water

The ordinary fluid extracts of Cinchona bark, containing much cincho-tannic acid, are wholly incompatible with the salts of iron. This detannated extract, therefore, will be found convenient for the extemporaneous preparation of elixirs, etc., in which it is required to combine cinchona bark with iron. The resulting elixirs will be found much superior in flavor to those in which alkaloids are substituted for the bark.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Cinchona, Yellow (Calisaya), U. S. P.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Powdered Extract Calisaya bark.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Calisaya bark.

Dose, 5 to 25 grains (0.3 to 1.6 Gm.).

Solid Extracts Cinchona, Pale, Red, and Yellow, U.S.P.

These extracts are made of a uniform strength by assay, one grain representing five grains of Cinchona bark.

Dose, 5 to 25 grains (0.3 to 1.6 Gm.).

Granular Effervescent Preparations containing Cinchona alkaloids.—Each drachm contains the quantity of ingredient specified.

Iron and Quinine Citrate, U. S. P., 3 grs.

Iron and Quinine Citrate with Arsenic.

Iron and quin. cit., U.S.P., 3 grs. Arsenious acid, 1-60 gr.

Iron Phosphate, Quinine and Strychnine.

Iron phosphate, 2 grs. Strychnine, 1-60 gr.

Quinine, 1 gr.

Quinine Citrate, 1 gr.

Normal Liquids.*

Cinchona Calisaya (Yellow bark).

Assayed standard 2 per cent. Quinine (U.S.P., 1880.).

Dose, 15 to 60 minims (1 to 4 C. c.).

Cinchona, Red.

Assayed standard 3.5 per cent. total alkaloids. Ten C.c. yields 0.35 Gm. mixed alkaloids (anhydrous) of which not more than 0.15 Gm. is Cinchonine.

Dose, 15 to 60 minims (1 to 4 C. c.).

Malt Extracts.—Each fluidounce represents ingredients specified.

Malt Extract Ferrated with Quinine.

Iron and Quinine citrate, 8 grs

Malt Extract with Iron, Quinine and Strychnine.

Iron and Quinine citrate, 8 grs. Strychnine, 4-100 gr.

Malt Extract with Quinine.

Quinine sulphate, 2 grs.

Oleate Quinine, containing 25 per cent. of the alkaloid.

Readily absorbed in local application.

Pills containing Cinchona or derivatives.

Anti-Malarial, McCaw's, 2 grs.

Quinine sulph., 1 gr.

Iron sulph. exsic., 1-4 gr.

Arsenious acid, 1-80 gr.

Gelsemin, 1-4 gr.

Podophyllin, 1-8 gr.

Oleoresin Black Pepper, 1-16 gr.

*See note in Preface on Normal Liquids.

CINCHONA—Continued.**Calisaya Alkaloids**, 2 grs.

Quinine sulph., 1-2 gr.

Cinchonine sulph., 1-2 gr.

Quinidine sulph., 1-2 gr.

Cinchonidine sulph., 1-2 gr.

Chinoidin, 1-2, 1, 2, and 3 grs.**Cinchonidine Salicylate**, 2 1-2 grs.**Cinchonidine Sulph.**, 1-2, 1, 2, 3, 4, and 5 grs.**Cinchonine Sulphate**, 1 1-2 and 3 grs.**Quinidine Sulphate**, 1, 2, and 3 grs.**Quinine Bisulphate**, 1-4, 1-2, 1, 2, 3, 4, and 5 grs.**Quinine Bromide**, 2 grs.**Quinine Sulphate**, 1-10, 1-4, 1-2, 1, 2, 3, 4, and 5 grs.**Quinine Valerianate**, 1-2 gr.**Tonic, Hæmætic**, Andrews, 2 3-4 grs.

Quinine sulph. 1 gr.

Iron by hydrogen, 1 1-2 gr.

Arsenious acid, 1-40 gr.

Powd. Ipecac, 1-8 gr.

Strychnine sulph., 1-40 gr.

“Warburg’s Tincture,” each pill representing 1 fluidrachm.

For other combinations see appended Pill Formula list for numbers 4, 5, 32, 41, 47, 99, 112, 116—119, 122, 162, 172—174, 214, 232, 281, 320—322, 330, 331, 334—338, 367, 369, 370, 372—385, 424, 426, 427, 447, and 490.

Syrups.—Each fluidounce represents ingredients specified.**Hæmætic Hypophosphites**, Improved.

A scientific and non-secret substitute for Fellow’s Hypo-phosphites and containing the same ingredients.

Adapted to a great variety of diseased conditions, in debility or depraved nutrition, anæmia, nervous prostration, and in all scrofulous and tubercular affections.

Potassium hypophos., 1 1-2 gr.

Manganese hypophos., 1 gr.

Strychnine hypophos., 1-16 gr.

Iron hypophos., 1 1-4 gr.

Calcium hypophos., 1 gr.

Quinine hypophos., 7-16 gr.

Iron and Quinine Citrate, 10 grs.**Iron, Quinine and Strychnine Phosphates.**

Iron phosphates, 16 grs.

Quinine phosphate, 8 grs.

Strychnine phosphate, 8-60 gr.

Tablet Triturates.**Quinine Bisulphate**, 1-4, 1-2, and 1 gr.**Quinine Sulphate**, 1-4, 1-2, and 1 gr.**Tinctures.****Cinchona, U. S. P.**

Fluid Extract or Normal Liquid Calisaya, P., D. & Co.’s.....4 fl. ozs.

Alcohol.....13 fl. ozs.

Water.....4 1-2 fl. ozs.

Glycerin.....1 fl. oz.

Dose, 1 to 4 fluidrachms (4 to 15 C. c.).

NOTE.—This is uniform in strength with the tinctures of the European pharmacopœias; the B. P. is made with Red Cinchona.

Cinchona Comp., U. S. P.

Fluid Extract Cinchona, Red, Comp., P., D. & Co.’s.....3 fl. ozs.

Alcohol.....10 fl. ozs.

Water.....1 fl. oz.

Glycerin.....1 fl. oz.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

“Huxham’s Tincture,” after the original formula is made from the Pale bark. Prepared extemporaneously by employing the formula of the U. S. P. compound tincture as above, but substituting Fluid Extract Cinchona, Pale, Comp.

“Warburg’s Tincture.”—Each fluidounce represents 12 grains of Quinine sulphate, 3 grains Camphor, in combination with numerous aromatics. Said to frequently prove successful in malarial fever where quinine alone has failed.

The rules for the use of the remedy, as given by Dr. Maclean, are as follows: “One-half ounce is given alone without dilution, after the bowels have been evacuated by any convenient purgative, all drink being withheld; in three hours another half ounce is administered in the same way. Soon afterwards, particularly in hot climates, profuse, but seldom exhausting, perspiration is produced; with this there

CINCHONA—Continued.

is a rapid decline of temperature, immediate abatement of frontal headache—in a word, complete deferrescence, and it seldom happens that more is required.

Wines.—Each fluidounce represents ingredients specified.

Calisaya Bark, true, 40 grs.

Calisaya and Iron.

Calisaya bark, 40 grs.

Iron pyrophosphate, 8 grs.

Iron, Bitter, U. S. P.

Calisaya bark, alkaloids, 40 grs.

Iron and Ammonium cit., 16 grs.

CLEAVERS. Galium Aparine, *Linné.* (Herb.)

Synonyms.—Goose Grass, Bedstraw; Caille-lait, Fr.; Labkraut, Ger.

Properties.—Aperient, diuretic, alterative. Used chiefly as a diuretic in dropsy, in jaundice, and in treating skin diseases of a scrofulous nature.

Cleavers herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Cleavers.—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

CLOTBUR. Xanthium strumarium, *Linné.* (Leaves.)

Synonyms.—Cocklebur, Clotweed.

Properties.—Given internally as a hæmostatic where the bleeding surface can be reached only through the medium of the circulation. It is also a styptic when applied locally. Said to be an antidote to the bite of poisonous insects.

PREPARATION.

Fluid Extract Clotbur (Cocklebur).—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

GLOVES. Caryophyllus, U. S. P. Caryophyllus aromaticus, *Linné.* (Flower-buds.)

Properties.—Stimulant, anti-spasmodic, locally anæsthetic. Used as a corrective with griping purgatives, and as a flavoring article of diet and medicines.

PREPARATION.

Fluid Extract Cloves.—Not miscible with water.

Incompatible with salts of iron and alkaloids.

Dose, 8 to 30 minims (0.5 to 2 C. c.).

COCA. Erythroxylon, U. S. P. Erythroxylon Coca, *Lamarck.* (Leaves.)

Properties.—The physiological effects of coca are due to the presence in the drug of the alkaloid cocaine, whose effects as a local anæsthetic have come into such prominence. Probably the drug owes to this anæsthetic action its power of preventing the sensation of fatigue under extraordinary exertion, etc., The drug produces in health effects analogous to those of tea or coffee, but more powerful.

A writer in the *Centralblatt für Klinische Medicin* summarizes its therapeutic applications thus: 1. As a stimulant, if one wishes to do extra physical or mental work. 2. In gastric indigestion. 3. In the cachexiæ. 4. In combating the effects of morphine and alcohol. 5. In asthma. 6. As an aphrodisiac. 7. As a local anæsthetic.

Its utility has thus far been more fully demonstrated as a local anæsthetic than in any other rôle. It is certainly the best agent at the command of the physician in facilitating minor surgical procedures, examinations, and operations of all kinds.

"Cocaine remains thus far the only local anæsthetic of mucous membranes, and this fact renders the introduction of this alkaloid into medicine one of the most precious therapeutic acquisitions of this age."*

* Dujardin-Beaumetz, "New Medications," p. 310 (edition of Geo. S. Davis, 1886).

COCA—Continued.

Coca leaves, U. S. P., compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLE AND ITS SALTS.

Cocaine.—Crystals. The remarkable power of coca to sustain the system, under extraordinary exertion is due to this alkaloid. The drug itself, however, in the form of fluid extract, is better adapted in most cases than the alkaloid, for producing this effect.

Dose, 1-8 to 1 grain (0.008 to 0.065 Gm.), or more.

Cocaine Citrate, in 4 per cent. solution only; extremely hygroscopic.

Cocaine Hydrobromate, crystals, and also in 4 per cent. solution.

Cocaine Hydrochlorate, crystals, and also in 2 and 4 per cent. solutions.

The remarkable discovery has recently been made that this and other soluble salts of cocaine, have the property, when applied to the mucous membrane, of completely deadening sensibility. A few drops of a two-per-cent. solution when applied to the eye produce complete anaesthesia of the cornea and conjunctiva, so that operations can be performed without the need of chloroform or ether. The effect is transient, passing off within half an hour. The pupil of the eye is at the same time dilated, so that under the influence of this agent, ophthalmoscopic examinations can be made, and this effect too passes off in a few hours.

Dose, 1-8 to 1 grain (0.008 to 0.065 Gm.) or more.

Cocaine Salicylate, in 4 per cent. solution only; extremely hygroscopic.

PREPARATIONS.**Coca-leaf Cigars and Cigarettes.**

Coca smoked seems to produce the same effect on the system as coca taken internally, but the effects are not so marked. Used for the relief of hay-fever and many irritable throat affections.

Coca Cordial.—This cordial presents coca in palatable form, commending it especially to the large class of persons of delicate nervous organization, for whom it has most frequently to be prescribed. In its preparation the astringent and bitter constituents of the drug, which would exert only a disturbing influence, are rejected, while care is taken to retain the true active principle, cocaine. One fluidounce represents 60 grains of Coca leaves of good quality. Frequently advantageously combined with liquid acid phosphate (liquor acidis phosphoricis).

Dose, 2 to 4 fluidrachms (8 to 16 C. c.).

Elixir Coca.—Each fluidounce represents 120 grs.

Dose, 2 fluidrachms (8 C. c.).

Extracts.

Fluid Extract Coca, U. S. P.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

Powdered Extract Coca leaves.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Coca leaves.

Dose, 10 to 25 grains, (0.6 to 1.6 Gm.).

Solid Extract Coca.—One part equals 5 of leaves.

Dose, 10 to 25 grains (0.6 to 1.6 Gm.).

Hypodermic Tablet, Cocaine Hydrochlorate, 1-4 gr.

Normal Liquid Coca,*—So great is the variability in the coca leaves that reach this market, that little dependence can be placed on any preparation not known to be made from a standard drug, or to be adjusted to a fixed standard of strength. In this respect our Normal Liquid of Coca may claim superiority to the products of the majority of manufacturers; standard provisionally 0.5 per cent. cocaine.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

Oleate Cocaine, 5 per cent.

Pills.

Aphrodisiac Comp., 2 3-4 grs.

Ext. Coca, 1 gr.

Cinchonidine sulph., 1-2 gr.

Phosphorus, 1-200 gr.

Ext. Nux vomica, 1-8 gr.

Iron bromide, 1 gr.

Coca Extract, 3 grs.

Cocaine Hydrochlorate, 1-16 and 1-8 gr.

* See note in Preface on Normal Liquids.

COCA--Continued.**Sedative,** Mann's, 2 1-2 gr.

Ext. Coca, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Cannabis Indica, 1-4 gr.

Vallet's Mass, 1 gr.

Arsenious acid, 1-80 gr.

Strychnine 1-80 gr.

Codeine, 1-8 gr.

Tablet Triturates.**Cocaine Hydrobromate**, 1-16 and 1-8 gr.**Cocaine Hydrochlorate**, 1-16 and 1-8 gr.**Wines.**—Each fluidounce represents ingredients specified.**Coca**, 60 grs.**Coca and Beef.**

Coca, 60 grs.

Beef, 2 ozs.

Coca, Beef and Iron.

Coca, 60 grs.

Iron and Ammonium cit., 4 grs.

Beef, 2 ozs.

COCILLANA. *Sycocarpus Rusbyi*, Britton. (Bark.)*Properties.*—A new drug from Bolivia possessing powerful emetic and expectorant properties; somewhat cathartic.

It has not yet been fully tested by the profession, and reports are solicited from physicians having clinical experience in its use.

PREPARATION.**Tincture Cocillana, Concentrated.**—Miscible with water. Four parts of tincture equals one of the bark.

Dose, 1-4 to 2 fluidrachms (1 to 8 C.c.) to be modified if further investigation necessitates.

COD-LIVER OIL. A fixed oil from the fresh livers of *Gadus Morhua*, *Linné*.*Properties.*—A supplementary food—nutrient and alterative. It is the most easily digested and also the most rapidly diffusible through animal membrane of any known oil or fat. In its use there is an increase of red blood-corpuscles and of the body weight, also general stimulation of healthy cell-formation. Externally it is successfully used to allay the itching of eczema and to assist in its cure.**PREPARATIONS.****Capsules**, in soft gelatin.**Cod-Liver Oil, Norwegian:** 10 minims (this size also in hard gelatin); 2 1-2, 5, 10, and 15 grms.**Cod-Liver Oil and Iodine:** Oil, 10 minims; Iodine, 1-4 gr. (also in hard gelatin.)**Cod-Liver Oil and Iodoform:** Oil, 10 minims; Iodoform, 2 grs.**Cod-Liver Oil and Iron:** Oil, 10 minims; Iron iodide, 1-2 gr.**Cod-Liver Oil and Phosphorus (1-30):** Oil, 10 minims; Phosphorus, 1-30 gr.**Cod-Liver Oil and Phosphorus (1-60):** Oil, 10 minims; Phosphorus, 1-60 gr.**Emulsions.****Cod-Liver Oil with Dialyzed Iron, Glycerin and Catawba Wine**, Dr. Stewart's formula.

Each 1-2 fluidounce, or ordinary dose, contains:

Cod-Liver Oil, 120 minims. Dialyzed Iron (glyc'd), 7 1-2 mins.

Mucilage Gum Arabic, 22 1-2 mins. Catawba Wine, 45 minims.

C P. Glycerin, 45 minims. Flavoring, 2 minims.

Cod-Liver Oil with Hypophosphite of Lime and Soda.

Each 1-2 fluidounce, or ordinary dose, contains:

Cod-Liver Oil, 2 fl. drms. Sodium hypophosphite, 2 grs.

Calcium hypophosphite, 4 grs. Gum, Sugar, Water, Flavor, q. s.

COD-LIVER OIL—Continued.

Malt Extracts.—Each fluidounce represents ingredients specified.

Malt Extract with Cod-Liver Oil.

Malt ext., 320 minims.

Cod-liver oil, 160 minims.

Malt Extract with Cod-Liver Oil and Hypophosphites.

Malt ext., 4 fluidrachms.

Calcium hypophos., 4 grs.

Cod-liver oil, 2-3 fluidrachms.

Sodium hypophos., 2 grs.

Malt Extract with Cod-Liver Oil and Iron Iodide.

Malt ext., 320 minims.

Iron iodide, 4 grs.

Cod-liver oil, 160 minims.

Malt Extract with Cod-Liver Oil and Phosphorus.

Malt ext., 320 minims.

Phosphorus, 1-25 gr.

Cod-liver oil, 160 minims.

COFFEE. *Coffea arabica*, *Linné*. (Seeds.)

Properties.—The physiological effects of coffee as a sustaining and restorative agent are well known, and depend mainly upon the caffeine contained in the seed. Indicated in all conditions in which caffeine is prescribed, particularly those of nervous headache, opium narcosis, spasmodic asthma, etc. More stimulating than coca, but less sustaining.

ACTIVE PRINCIPLE.

Caffeine Citrate, crystals.

Dose, 1 to 3 grains (0.065 to 0.2 Gm.).

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Coffee (from freshly roasted Java coffee).—Miscible with water. Chiefly used to make coffee syrup for flavoring purposes, but may be used medicinally.

Fluid Extract Coffee (from the green berry).—Miscible with water without material precipitation. Originally introduced as a substitute for Fluid Extract Guarana, which is a much more expensive preparation, but also depending upon the caffeine contained therein for its medicinal quality.

Dose, 1-2 to 1 1-2 fluidrachms (2 to 6 C. c.).

Granular Effervescent Preparations.—Each drachm represents ingredients specified.

Caffeine and Potassium Bromide.

Caffeine, 1 gr.

Potassium bromide, 5 grs.

Caffeine Citrate, 1-2 gr.

Pill, Caffeine Citrate, 1 gr.

Syrup, Coffee.

Fluid Extract Coffee P., D. & Co.'s.....2 fl. ozs.

Hot Syrup.....14 fl. ozs.

Tablet Triturates.

Caffeine, 1-2 and 1 gr.

Caffeine Citrate, 1-2 and 1 gr.

COLCHICUM. { *Colchici radix* } U. S. P. *Colchicum autumnale*,
{ *Colchici semen* } *Linné*. (Corm, seed.)

Synonym.—Meadow Saffron.

Properties.—Poisonous. Almost specific in acute gout, combination with an alkali assisting its action. Small doses are nauseant, cathartic, diuretic, and diaphoretic, thus generally exciting the secretory organs. It increases tissue waste and promotes elimination of the products of such waste.

Antidotes.—Tannin, to prevent absorption. Emetics—zinc sulphate. Apomorphine, mustard. Demulcent drinks and opium, to allay irritation. Stimulants, to prevent collapse.

Colchicum root, U.S.P., compressed in pound packages for retailing purposes.

COLCHICUM—Continued.

ACTIVE PRINCIPLE.

Colchicine, crystals. Uses the same as those of the crude drug, chiefly the treatment of gout and acute rheumatism. Soluble in water and in alcohol.

Dose, 1-50 to 1-10 grain (0.0013 to 0.006 Gm.).

PREPARATIONS AND FORMULÆ.

Extracts.

Fluid Extract Colchicum root, U.S.P.—Miscible with water.
Dose, 2 to 8 minims (0.12 to 0.5 C.c.).

Fluid Extract Colchicum seed, U. S. P.—Not miscible with water without some precipitation.

Dose, 2 to 8 minims (0.13 to 0.5 C.c.).

Powdered Extract Colchicum root, Acetic.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered sugar of milk.

Dose, 1-3 to 1 1-2 grains (0.02 to 0.1 Gm.).

Solid Extract Colchicum root, U. S. P.—One part equals 5 of the corm.

Dose, 1-3 to 1 1-2 grains (0.02 to 0.1 Gm.).

Hypodermic Tablet, Colchicine, 1-30 gr.

Normal Liquids.*

Colchicum root, standard, 1 per cent. Colchicine.

Dose, 2 to 8 minims (0.12 to 0.5 C.c.).

Colchicum seed, standard, 1 per cent. Colchicine.

Dose, 2 to 8 minims (0.12 to 0.5 C.c.).

Pills.

Colchicine, 1-60 gr.

Colchicum Extracti, 1-2 gr.

Rheumatic, 3 grs.

Ext. Coloc. co., 1 1-2 gr.

Ext. Colch. acet., 1 gr.

Ext. Henbane, 1-3 g

Calomel, 1-3 gr.

Tablet Triturates, Colchicine, 1-100 and 1-50 gr,

Tinctures.

Colchicum seed, P. G.

Fluid Extract Colchicum seed, P., D. & Co.'s.....1 fl. oz.

Dilute Alcohol.....10 fl. ozs.

Dose, 25 to 60 minims (1.5 to 4 C.c.).

Colchicum seed, U. S. P.

Fluid Extract Colchicum seed, P., D. & Co.'s.....2 fl. ozs.

Alcohol.....8 fl. ozs.

Water.....4 fl. ozs.

Dose, 15 to 60 minims (1 to 4 C.c.).

Wines.—Each fluidounce represents ingredients specified.

Colchicum root, U. S. P., 180 grs.

A medicinal equivalent may be extemporaneously prepared as follows:

Fluid Extract Colchicum root, P., D. & Co.'s.....4 fl. ozs.

Sherry (or native) Wine.....6 fl. ozs.

Dose, 5 to 20 minims (0.33 to 1.33 C.c.).

Colchicum seed, U. S. P., 70 grs.

A medicinal equivalent may be extemporaneously prepared as follows:

Fluid Extract Colchicum seed, P., D. & Co.'s.....2 fl. ozs.

Stronger White Wine.....11 fl. ozs.

Dose, 15 to 60 minims (1 to 4 C.c.).

COLOCYNTH. Colocynthis, U. S. P. Cucumis Colocynthis, Linné. (Fruit, pulp.)

Synonyms—Citrullus Colocynthis, *Schrader*; Bitter Apple, Bitter Cucumber, Coloquinte, Fr.; Coloquintenaffel, Ger

Properties.—**POISONOUS.** A powerful drastric hydragogue cathartic, producing, when given in overdoses, violent griping, with dangerous inflam-

* See note in Preface on Normal Liquids.

COLOCYNTH—Continued.

mation. It is seldom given alone, but when given with other cathartics as an adjunct to ensure their action, it is considered a standard remedy. It has been employed alone in obstinate dropsy, amenorrhœa, and in cerebral derangements.

ANTIDOTES.—Emetics (zinc sulphate, apomorphine, etc.), if the case is seen early. Later, demulcents and opiates, with stimulants to combat the tendency to collapse.

ACTIVE PRINCIPLE.

Colocynthin.—Glucoside. Uses the same as those of the crude drug. Purgative in doses of 1-8 grain (0.008 Gm.) and upwards.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Colocynth.—Not miscible with water without some precipitation.

Dose, 2 to 5 minims (0.13 to 0.33 C.c.).

Powdered Extract Colocynth, U. S. P.

Dose, 1 to 2 grains (0.06 to 0.12 Gm.).

Powdered Extract Colocynth Comp., U. S. P.—Each 100 parts represents: Ext. Colocynth, 16; Aloes, 50; Cardamom, 6; Resin of Scammony, 14; and Soap, 14 parts.

Dose, 5 to 20 grains (0.3 to 1.3 Gm.).

Pills.

Anti-Bilious, 2 3-4 grs.

Ext. Coloc. comp., 2 1-2 grs.

Podophyllin, 1-4 gr.

Aperient, 3 grs.

Ext. Nux vomica, 1-3 gr.

Ext. Henbane, 1-2 gr.

Ext. Coloc. comp., 2 grs.

Calomel and Extract Coloc. Comp., 3 1-2 grs.

Calomel, 1 gr.

Ext. Coloc. comp., 2 1-2 grs.

Coloc. Comp., Extract, U. S. P., 3 grs.

Colocynthin, 1-8 gr.

Laxative, Cole's, 4 gr.

Podophyllin, 1-10 gr.

Ext. Coloc. comp., powd., 3 grs.

Calomel, 1 gr.

Ox Gall, Craig's, 3 grs.

Ox Gall, 2 grs.

Ext. Coloc. comp., 1 gr.

For other combinations containing Colocynth see Pill Formulæ list appended for numbers 4, 27, 36, 37, 44, 56, 64, 82, 102-106, 130-140, 205, 255, 388, 460, 461.

Tablet Triturates, Colocynthin, 1-20 gr.

Tincture Colocynth, P. G.

Fluid Extract Colocynth, P., D. & Co.'s..... 1 fl. oz.

Alcohol..... 11 fl. ozs.

Dose, 25 to 60 minims (1.5 to 4 C.c.).

COLTSFOOT. Tussilago Farfara, Linné. (Leaves, rhizome.)

Synonyms.—Hufelttig, Rosshuf, Ger.

Properties.—Demulcent and slightly tonic. Generally used for coughs and colds, as indicated by its name (*tussis*, L. a cough).

Coltsfoot leaves, compressed in ounce packages for retailing purposes.

Coltsfoot root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Coltsfoot leaves.—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

Lozenges, 2-3 grain in each.

COLUMBO (foreign). Calumba, U. S. P. Jateorrhiza Calumba, *Miers.* (Root.)*Synonym.*—Cocculus Palmatus, *DeCandolle.*

Properties.—A mild tonic without astringency, with very little stimulating power, and generally acceptable to the stomach; it answers admirably as a remedy in simple dyspepsia, and in the debility of convalescence, especially when the alimentary canal is left enfeebled, *e. g.*, after remittent fever, dysentery, diarrhoea, cholera morbus and cholera infantum. The absence of irritating properties renders it also an appropriate tonic in the hectic fever of phthisis and kindred affections. For vomiting unconnected with inflammation of the stomach, as in the sickness of pregnancy it is often of service.

NOTE.—Contains no tannin, it may therefore be prescribed with iron.

Columbo root, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Columbo root.—Miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C. c.).

Powdered Extract Columbo.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Columbo.

Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

Solid Extract Columbo.—One part equals 5 of root.

Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

Tincture Columbo, U. S. P.

Fluid Extract Columbo, P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol.....20 fl. ozs.

Dose, 1 fluidrachm (4 C. c.)

COMFREY. *Symphytum officinale, Linné.* (Root.)

Synonyms.—Consoude, Fr.; Schwartzwurz, Beinwurz, Ger.

Properties.—Demulcent, slightly astringent and tonic. Useful in catarrhal affections of the mucous membranes, as in bronchial irritation, leucorrhœa, endometritis, etc.

Comfrey root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Comfrey root.—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

CONDURANGO. *Gonolobus Cundurango, Triana.* (Bark.)

Properties.—Said to be tonic, emetic, diuretic and deobstruent. It is unquestionably an alterative of considerable value, and has been employed in the treatment of chronic rheumatism, syphilis, non-specific cutaneous affections, etc. It is now official in the German Pharmacopœia, and it promises to obtain a permanent place in the materia medica.

It was first introduced to the notice of the profession as a specific in the treatment of cancer.

Condurango bark, compressed in pound packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Condurango bark.—Not miscible with water without some precipitation.

Dose, 30 to 60 minims (2 to 4 C. c.).

Powdered Extract Condurango.—One part equals 5 of bark.

Dose, 6 to 10 grains (0.4 to 0.65 Gm.).

CONIUM, U. S. P. *Conium maculatum, Linné.* (Leaves, seed.)

Synonyms.—Poison Hemlock, Poison Parsley. Grande ciguë, Fr.; Schierlingsfrüchte, Ger.

Properties.—POISONOUS. Conium is anodyne, antispasmodic, anaphrodisiac and discutient. In overdoses it produces motor paralysis. It is of value as a palliative in the treatment of scirrhus cancerous ulcers, mammary tumors, chronic enlargements of the liver, painful scrofulous tumors and ulcers, also in various diseases of the skin, especially strumous. Its most important use is in controlling excitement in maniacs, and in the treatment of chorea, whooping-cough and other spasmodic affections.

ANTIDOTES.—Tannin (strong tea, decoction of oak bark or sumach). Emetics (apomorphine, zinc sulphate, mustard). Sustain the vital powers by stimulants, external heat, faradization; employ artificial respiration persistently as long as there is any sign of cardiac action.

N. B.—Conium seed is now official in place of the leaves, which are less active. Prescribers should always specify which is meant.

Conium leaves, compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLE.

Coniine.—Volatile liquid alkaloid. Preparations of conium are notoriously untrustworthy, the drug varying greatly in strength even when fresh, and rapidly deteriorating with age. The alkaloid commends itself as a substitute invariable in medicinal activity, but best employed in the form of a salt, hydrochlorate or hydrobromate, since those compounds are readily crystallizable and freely soluble in water; the alkaloid itself can have only a limited use owing to its highly volatile character, but an alcoholic solution of it—1 to 10—may be preserved indefinitely, is constant in strength and may easily be introduced into mixtures. The therapeutic properties are those of the crude drug. Medium dose, 1-16 gr (0.004 Gm.), but this must be rapidly, and often largely, increased to produce the desired effect.

Coniine Hydrobromate, crystals.

Dose, 1-20 to 1-5 grain (0.003 to 0.013 Gm.), cautiously increased.

Coniine Hydrochlorate, crystals.

Dose, 1-20 to 1-5 grain (0.003 to 0.013 Gm.), cautiously increased

PREPARATIONS AND FORMULÆ.**Abstract Conium seed, U. S. P.**

Normal Liquid or Fluid Extract Conium seed, P., D. & Co.'s.....1 fl. oz.
Sugar of Milk, q. s. to make228 grs.
Evaporate the fluid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains, finally powder the product. Keep in a well corked bottle.

Dose, 2 to 10 grains (0.13 to 0.65 Gm.). One part equals 2 of seed

Extracts.

Fluid Extract Conium leaves.—Miscible with water.

Dose, 5 to 20 minims (0.33 to 1.3 C.c.), cautiously increased.

Fluid Extract Conium seed, U. S. P.—Miscible with water.

Dose, 5 to 20 minims (0.33 to 1.3 C.c.)

Powdered Extract Conium leaves.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Conium lvs.

Dose, 2 to 5 grains (0.13 to 0.33 Gm.)

Solid Extract Conium leaves.—One part equals 5 of prime lvs.

Dose, 2 to 5 grains (0.13 to 0.33 Gm.)

Solid Extract Conium seed, U.S.P.—One part equals 6 of seed.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.)

Hypodermic Tablet, Coniine Hydrobromate, 1-8 gr.

The subcutaneous use of Coniine is indicated in tetanus, strychnine poisoning and acute mania.

Normal Liquid,* Conium seed.—Standard .6 per cent. Coniine.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

* See note in Preface on Normal Liquids.

CONIUM—Continued.**Pills.****Conium Extract** (*of the seed*), 1-2 and 1 gr.**Conium Extract and Ipecac**, 1 gr.

Ext. Conium seed, 1-2 gr. Powd. Ipecac, 1-2 gr.

Syphilitic, Ricord's, Modified, 2 1-2 grs.

Mercury protiodide, 1-2 gr. Ext. Opium, 1-10 gr.

Lactucarium, 1-2 gr. Ext. Conium seed, 1 1-2 gr.

See numbers 8 and 279, also containing Conium, in Pill Formulæ list appended.

Tablet Triturates.**Coniine Hydrobromate**, 1-20 and 1-10 gr.**Coniine Hydrochlorate**, 1-20 and 1-10 gr.**Tinctures.****Conium, U. S. P.**

Fluid Extract Conium Seed, P., D. & Co.'s.....2 1-2 fl. ozs.

Dilute Alcohol14 1-2 fl. ozs.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.), increased if necessary.

"German Tincture" Conium seed.

An imported preparation of the green fruit (not official).

Dose, 8 to 20 minims (0.5 to 1.3 C.c.).

COOLWORT. *Mitella nuda*, Linné. (Leaves.)*Synonyms.*—*Mitella cordifolia*, Lamarck; Mitrewort.*Properties.*—Diuretic. Useful in strangury, and as a remedy in inflammatory and catarrhal affections of the kidneys and bladder.**Coolwort leaves**, compressed in ounce packages for retailing purposes.**COPAIBA.** An oleoresin chiefly obtained from *Copaifera Langsdorffii*, Desfontaines.*Properties.*—In the treatment of chronic catarrh of the mucous membrane it is the most active and efficient of the balsams and oleoresins. Given in gonorrhœa and cystitis after the acute inflammatory symptoms have subsided; also in leucorrhœa, bronchorrhœa, and in chronic skin diseases such as leprosy and psoriasis.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.), best administered in gelatin capsules; see below.

PREPARATIONS.**Capsules**, in hard and in soft gelatin.**Copaiba, Para**, 10 minims.**Copaiba, Cubeb and Sandal.**

Copaiba, Para, 6 minims.

Oil Cubeb, 2 minims.

Oil Sandal, East India, 2 minims.

Copaiba and Oil of Cubeb.

Copaiba, Para, 7 minims.

Oil Cubeb, 3 minims.

Copaiba and Oleoresin of Cubeb.

Copaiba, Para, 7 minims.

Oleo-resin Cubeb, 3 minims.

These are the leading copaiba combinations in capsules, for others see appended Capsule Formulæ list for numbers 18, 20, 21, 22, 24, 25, 27, 37, 38, 39, 41, 69, 105-108, 114, 115.

Pills.**Copaiba**, 3 and 4 grs.**Copaiba Compound**, 3 grs.

Pil. Copaiba, 1 1-2 gr.

Resin Guaiac, 3-8 gr.

Iron citrate, 3-8 gr.

Ext. Cubeb, 5-8 gr.

Copaiba and Oleo-Resin Cubeb, 3 grs.

Pil. Copaiba, 2 grs.

Oleo-resin Cubeb, 1 gr.

Copaiba, Cubeb and Citrate of Iron, 3 gr.

Pil. Copaiba, 3-4 gr.

Ext. Cubeb, 1 1-2 gr.

Iron citrate, 3-4 gr.

Gonorrhœa, 3 grs.

Powd. Cubebs, 1 1-4 gr.

Solid. Copaiba, 1 1-4 gr.

Iron sulph. exsic., 1-4 gr.

Venice Turpentine, 1-4 gr.

CORIANDER. *Coriandrum*, U. S. P. *Coriandrum sativum*, *Linné.*
(Seed.)

Properties.—An agreeable stimulant aromatic and carminative. Principally used as a flavoring to other remedies.

PREPARATION.

Fluid Extract Coriander seed.—Not miscible with water.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

CORN (Maize). *Ustilago*, U. S. P. *Ustilago Maydis*, Corn smut or Corn ergot: *Stigmata*, Corn-silk (pistils), *Zea Mays*, *Linné.*

Properties.—Corn ergot closely resembles in its medicinal properties the ergot of rye; and has been considered, by many practitioners, to be quite as efficient and more uniform than the latter. The uterine contraction from ergot is tonic; that from *ustilago* seems to be regularly intermittent. If this proves to be a constant peculiarity in the action of the drug, it is likely to prove a more serviceable agent in labor than ergot itself. It will give less pain to the mother, less danger to her in its administration, and also less danger to the child.

It has been pronounced superior to ergot in passive hemorrhages, and also very effectual in spermatorrhœa, psoriasis, eczema, fibroid tumors and allied affections.

Corn-silk has been recently introduced on the recommendation of very eminent authorities as a remedy in inflammatory affections of the bladder, acute or chronic, idiopathic or traumatic. It is diuretic, demulcent and antiseptic in its action, and is reported to be especially valuable in cases in which decomposition of the urine takes place in the bladder, giving rise to vesical irritation.

PREPARATIONS.

Extracts.

Fluid Extract Corn Ergot (Ustilago).—Miscible with water.
Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

Fluid Extract Corn-silk.—Miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Syrup, Corn-silk.—Each fluidounce represents 120 grains Corn-silk.

Wine, Corn-silk.—Each fluidounce represents 120 grains Corn-silk.

COTO. *Nectandra* (Sp. indet.). (Bark.)

Synonym.—Para-Coto bark.

NOTE.—Two distinct kinds of coto bark are now brought from Bolivia, distinguished as "coto" and "paracoto," respectively. The therapeutic properties ascribed to the two drugs are similar. There is reason to believe that the true coto bark is at least as efficient a medicine as the paracoto, but since our patrons have so generally expressed a preference for the latter, after having made its acquaintance under the name of Coto, we invariably supply this article when coto bark is ordered.

Properties.—An efficient remedy in cholera morbus, diarrhœa, acute or chronic dysentery, colic, and is also recommended in gout, rheumatism and neuralgic toothache. Cases are reported of long-standing diarrhœa, intestinal catarrh and dysenteric disease which had been unsuccessfully treated with tannin, opium and lead acetate, but making complete recovery in from 12 hours to six days by hourly doses of 1-10 grain cotoin.*

ACTIVE PRINCIPLES.

Cotoin.—Neutral principle derived from Coto bark. A valuable remedy in chronic diarrhœas, such as those that complicate mental derangement, phthisis, etc. Less apt to produce nausea than the fluid extract.
Dose, 2 to 3 grains (0.13 to 0.20 Gm.).

* Napheys: "Medical Therapeutics," p. 352.

COTO—Continued.

Paracotoïn.—Neutral principle derived from Paracoto bark. Resembles cotoïn in its uses, but is less powerful.
Dose, 2 to 5 grains (0.13 to 0.3 Gm.).

PREPARATIONS.

Elixir Coto bark.—Each fluidounce represents 30 grains of bark.

Fluid Extract Paracoto (Coto) bark.—Not miscible with water.
Dose, 5 to 20 minims (0.3 to 1.3 C.c.), four to six times a day.

In doses of 10 drops and upwards, it is liable to cause gastric irritation, hence it is best to begin with the minimum dose, increasing, if it is well borne, until the desired effect is produced. In all cases it should be well diluted before it is administered.

COTTON. *Gossypium*, U.S.P. *Gossypium herbaceum*, *Linné*. (Bark of root, hairs of the seed, leaves.)

Synonym.—Baumwolle, Ger.

Properties.—The bark and also the leaves of the Cotton Root are emmenagogue, oxytotic and abortifacient.

It is said to promote uterine contraction with as much efficiency and more safety than ergot. It has been used with success for chlorosis, amenorrhœa, dysmenorrhœa etc.

The hairs of the seed are employed as surgical dressing, and indirectly in the preparation of collodion.

Cotton-root bark, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Collodions.—Each fluidounce represents ingredients specified.

Aconite, 15 grs.

Anodyne (Morphine), Morphine, 4 grs.

Arnica, root, 24 grs.

Belladonna, leaves, 20 grs.

Cantharidal, U. S. P., 240 grs.

Capsicum, 16 grs.

Carbolic Acid, 30 grs.

Caustic, Corrosive sublimate, 68 grs.

Ferruginous, Solution Iron chloride, U. S. P., 60 minims.

Flexible, U. S. P.

Iodized, Iodine, 15 grs.

Simple, U. S. P.

Styptic, U. S. P., Tannin, 30 grs.

Extracts.

Fluid Extract Cotton leaves.—Miscible with water.

Dose, 1 to 4 fluidrachms (4 to 16 C.c.).

Fluid Extract Cotton Root bark, U. S. P.—Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Fluid Extract Cotton Root bark (green bark).—Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Solid Extract Cotton root.—One part equals 10 of root bark.

Dose, 3 to 15 grains (0.2 to 1 Gm.).

Gossypin, a so-called **Concentration**,* a powdered preparation from the fresh inner bark of the root.

* See note in Preface on Concentrations.

COTTON—Continued.**Pills.****Emmenagogue, No. 4,** 4 1-2 grs.

Ergotin, 1 gr.

Ext. Cotton root, 1 gr.

Purified Aloes, 1 gr.

Iron sulph. exsic., 1 gr.

Oil Savin, 1-4 gr.

Female, Amenorrhœa, 4 grs.

Ext. Black Cohosh, 1 gr.

Iron sulph. exsic., 1 gr.

Ext. Cotton root, 1 gr.

Purified Aloes, 1 gr.

COUCH-GRASS. Triticum, U. S. P. Triticum repens, *Linné*.
(Rhizome.)*Synonyms.*—Dog-grass, Knot-grass, Quickens; Chiendent, Fr.; Queckenwurz, Ger.*Properties.*—A feeble diuretic, but chiefly used for its demulcent and emollient qualities in cystitis and irritable bladder.**Couch-grass, U. S. P.,** compressed in ounce packages for retailing purposes.**PREPARATIONS.****Extracts.****Fluid Extract Couch-grass, U. S. P.**—Miscible with water.

Dose, 1 to 6 fluidrachms (4 to 24 C. c.).

Solid Extract Couch-grass.—One part equals 2 of drug.

Dose, 20 grains (1.3 Gm.), and upwards.

CRAMP-BARK. Viburnum Opulus, *Linné*. (Bark.)*Synonyms.*—Cranberry tree, High Cranberry.*Properties.*—Tonic, anti-spasmodic and alterative, popularly known as "Cramp Bark" from its influence over painful spasmodic muscular contractions. It has been employed with benefit in asthma and in the painful muscular spasms so frequently attacking the extremities of pregnant females. Particularly beneficial in the treatment of dysmenorrhœa of the neuralgic variety, and in that irritable condition of the uterus which predisposes to abortion. Useful in the treatment of after-pains.**Cramp bark,** compressed in pound packages for retailing purposes.**PREPARATIONS.****Fluid Extract Cramp bark.**—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Viburnin, a so-called **Concentration,*** a powdered preparation from Cramp-bark.

Dose, 1 to 3 grains (0.065 to 0.2 Gm.).

CRANESBILL. Geranium, U. S. P. Geranium maculatum, *Linné*.
(Rhizome.)*Synonyms.*—Crowfoot, Spotted Geranium.*Properties.*—A powerful astringent. Used in dysentery, diarrhœa, cholera infantum, and in excessive mucous discharges; also as a gargle for sore mouth, sore throat, and relaxation of the uvula. Highly recommended as a substitute for tannic acid, as unlike that agent, it does not cause a dryness of the mucous surfaces with which it comes in contact, but produces its therapeutical influences upon them without entirely checking their secretion.**Cranesbill root, U. S. P.,** compressed in pound packages for retailing purposes.**PREPARATIONS.****Extracts.****Fluid Extract Cranesbill, U. S. P.**—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

* See note in Preface on Concentrations.

CRANESBILL—Continued.

Solid Extract Cranesbill.—One part equals 5 of root.

Dose, 5 to 10 grains (0.32 to 1.65 Gm.).

Geranin, a so-called **Concentration**,* a powdered preparation from

Cranesbill root.

Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

Pill, Geranin, 1 gr.

CRAWLEY. *Corallorhiza odontorhiza*, *Nuttall.* (Rhizome.)

Synonym.—Coral-root.

Properties.—A prompt, powerful and certain diaphoretic, with sedative properties. It promotes perspiration without causing excitement of any kind in the system. Hence its use in typhus and other fevers, certain inflammatory diseases, acute erysipelas, cramps, flatulency, pleurisy, and night-sweats. It relieves hectic fever without debilitating the patient. Combined with Blue Cohosh, it forms an excellent agent in amenorrhœa, dysmenorrhœa, after-pains, suppression of the lochia and febrile symptoms which occur at the parturient period.

Crawley root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Crawley root.—Not miscible with water without some precipitation.

Dose, 15 to 30 minims (1 to 2 C. c.) in hot water.

CROTON OIL. The oil expressed from the seeds of *Croton Tiglium*, *Linné.*

Properties.—A violent irritant purgative. On account of the smallness of the dose and rapidity of action, it is found very useful in coma, tetanus, and the like, where the patient is unable to swallow; and in phrenitis where a strong counter-action is required. Also useful in obstinate constipation where, after the lower bowel has been emptied by enemata, milder purgatives fail. Rubefacient externally.

Dose, 1-3 to 2 minims (0.02 to 0.12 C. c.).

PREPARATIONS.**Pills.**

Croton Oil, 1-50 gr.

Triplex, Dr. Francis', 4 grs.

Purified Aloes, 1 1-5 gr.

Blue Mass, 1 1-5 gr.

Oil Caraway, 1-5 gr.

Resin Scammony, 1 1-5 gr.

Croton Oil, 1-20 gr.

Tr. Aloes and Myrrh, 1-4 gr.

CUBEB. *Cubeba*, U.S.P. *Piper cubeba*, *Linné.* (Unripe fruit.)

Synonym.—*Cubeba officinalis*, *Miguel.*

Properties.—Local irritant, with a general stimulant action, directed especially to the mucous tissues. Stimulant diuretic and expectorant. Used successfully in gonorrhœa, gleet, leucorrhœa, and in chronic catarrhal inflammation of the mucous tracts.

PREPARATIONS.**Capsules.**

Copaiba and Oil of Cubeb.—In hard and in soft gelatin.

Copaiba, Para, 7 minims.

Oil Cubeb, 3 minims.

Copaiba and Oleoresin of Cubeb.—In soft gelatin.

Copaiba, Para, 7 minims.

Oleoresin Cubeb, 3 minims.

Copaiba and Oleoresin of Cubeb.—In hard gelatin.

Copaiba, Para, 8 minims.

Oleoresin Cubeb, 2 minims.

* See note in Preface on Concentrations.

CUBEB—Continued.

Copaiba, Cubeb, Matico and Sandal.—In soft gelatin.
 Copaiba, Para, 2 minims. Oleoresin Matico, 1 minim.
 Oleoresin Cubeb, 3 minims. Oil Sandal, East India, 3 minims.
Copaiba, Cubeb and Sandal.—In hard and in soft gelatin.
 Copaiba, Para, 6 minims. Oil Sandal, East India, 2 minims.
 Oil Cubeb, 2 minims.

Cubeb Oil, 10 minims.—In soft gelatin.

Cubeb Oleoresin, 10 minims.—In soft gelatin.

For other capsules containing Cubeb see appended Capsule Formulæ list for numbers 20—22, 24, 25, 38, 40, 41, 48, 108, 115.

Extracts.

Fluid Extract Cubeb.—Not miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C. c.).

Solid Extract Cubeb.—One part equals 5 of the fruit.

Dose, 2 to 8 grains (0.13 to 0.52 Gm.).

Lozenges.—Each containing ingredients specified.

Bronchial (with Cubeb).

Ext. Licorice, 4-5 gr.
 Balsam Tolu, 1-20 gr.

Powd. Cubeb, 5-8 gr.
 Oil Sassafras, 1-40 gr.

Cubeb.

Ext. Licorice, 3-4 gr.
 Oil Sassafras, 1-32 gr.

Oleoresin Cubeb, 1-16 gr.

Pills.

Cubeb Extract, 2 gr.

Cubeb and Alum, 3 grs.
 Oleoresin Cubeb, 2 grs.

Powd. Alum, 1 gr.

Cubeb Extract, Rhatany and Iron, 3 grs.

Ext. Cubeb, 1 1-2 gr.
 Iron sulph. exsic., 1 gr.

Ext. Rhatany, 1-2 gr.

For others containing Cubeb see appended Pill Formulæ list for numbers 147—149, 199.

Tablet Triturate, Ammonia Muriate Comp.

Ammonia muriate, 1-4 gr.
 Powd. Cubeb, 1-8 gr.

Powd. Ext. Licorice, 1-10 gr.

Tincture Cubeb, U. S. P.

Fluid Extract Cubeb, P., D. & Co.'s..... 2 fl. ozs.
 Alcohol..... 20 fl. ozs.
 Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

NOTE.—Stronger alcoholic menstruum than the Pharmacopœa directs is necessary in order to obtain solution of the oil and oleoresin.

CUCUMBER-TREE. *Magnolia acuminata, Linné.* (Flowers.)

Synonym.—*Magnolia* Flowers.

Properties.—Used as a substitute for Cinchona bark in the treatment of remittent and intermittent fever. Its curative agency is said to be favored by the diaphoretic action following its administration. The flowers have the general character of an aromatic bitter tonic, and are particularly recommended in dyspepsia and chronic rheumatism.

PREPARATION.

Fluid Extract Cucumber-tree (magnolia) flowers.—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

CULVERS. *Leptandra, U.S.P. Veronica virginica, Linné.* (Rhizome.)

Synonyms.—*Leptandra virginica, Nuttall*; Blackroot; *Leptandra*, Fr. and Ger.

Properties.—A mild cathartic somewhat resembling podophyllin in its operation. It is indicated in diseases of the intestinal canal, and in diseases in which there is constipation, with insufficiency of biliary and intestinal secretions. An invaluable remedy to prevent the return of intermittents after they have been broken up by quinine.

CULVERS—Continued.

Culver's root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixir Cathartic Comp.—Each fl. oz. represents ingredients specified.

Senna, 16 grs.	Butternut, 16 grs.
Cascara Sagrada, 32 grs.	Rochelle Salts, 32 grs.
Culver's root, 16 grs.	

Extracts.

Fluid Extract Culver's root.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Powdered Extract Culver's root.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Culver's root.
Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Solid Extract Culver's root.—One part equals 5 of root.

Dose, 3 to 10 grains, (0.2 to 0.6 Gm.).

Leptandrin, a so-called **Concentration**,* a powdered preparation from Culver's root.

Dose, in acute cases, 1-4 to 1 grain (0.016 to 0.065 Gm.). In chronic cases, 1 to 4 grains (0.065 to 0.26 Gm.).

Pills.

Dandelion and Leptandrin, 2 grs.

Ext. Dandelion, 1 1-3 gr. Leptandrin, 2-3 gr.

Leptandrin, 1-16, 1-8, 1-4, 1-2 and 1 gr.

Leptandrin Comp., 1 3-8 gr.

Leptandrin, 1 gr. Irisin, 1-4 gr.
Podophyllin, 1-8 gr.

Podophyllin and Leptandrin, 1 1-2 gr.

Podophyllin, 1-2 gr. Leptandrin, 1 gr.

For other pills containing Culver's root see Pill Formulæ list for numbers 63, 101, 102, 206, 260, 261, 263, 356, 460.

Tablet Triturate, Leptandrin, 1-16 gr.

CURARE. An extract employed as an arrow poison by the natives of South America, and obtained from several plants, one of which is a *Strychnos* and another a *Cocculus*.

Synonym.—Worara.

Properties.—Produces, when introduced into the circulation, powerful toxic effects, the most notable symptom of which is muscular paralysis.

Medicinally, curare acts as a powerful anti-spasmodic, and is employed with success in the treatment of tetanus. It is also sometimes prescribed in epilepsy, hydrophobia, etc.

Commencing dose 1-10 grain (0.006 Gm.), which should be administered hypodermically to secure its full action.

DAMIANA. *Turnera microphylla, DeCandolle.* (Leaves.)

Synonym.—*Turnera aphrodisiaca, L. F. Ward.*

Properties.—Stimulant, tonic, laxative, diuretic and distinctly aphrodisiac.

The drug has been extensively prescribed with reference to the last named property, and is less objectionable than the intestinal irritants to which the same property is ascribed. A more important property is probably the general tonic and roborant action of the drug, which appears to be exerted primarily on the nervous system.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Damiana, true, 120 grs.

Damiana, Iron and Phosphorus.

Damiana, true, 60 grs.	Phosphorus, 4-100 gr.
Iron pyrophosphate, 8 grs.	

* See note in Preface on Concentrations.

DAMIANA—Continued.

Damiana, Phosphorus, Nux Vomica and Iron.
 Damiana, true, 60 grs. Iron pyrophosphate, 8 grs.
 Nux vomica, 8 grs. Phosphorus, 4-100 gr.

Damiana, Phosphorus and Nux Vomica.
 Damiana, 60 grs. Phosphorus, 4-100 gr.
 Nux vomica, 8 grs.

Extracts.

Fluid Extract Damiana.—Not miscible with water.
 Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Powdered Extract Damiana.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Damiana.
 Dose, 5 to 15 grains (0.3 to 1 Gm.).

Solid Extract Damiana.—One part represents 5 of the leaves.
 Dose, 5 to 15 grains (0.3 to 1 Gm.).

Glycerole; Damiana, Phos. and Nux Vomica Comp.—Each fluidounce represents ingredients specified.
 Damiana, 60 grs. Nux vomica, 4 grs.
 Phosphorus, 1-25 gr.
 Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Pills.

Damiana Extract, 3 grs.
Phosphorus, Nux Vomica and Damiana, 2 1-4 grs.
 Phosphorus, 1-100 gr. Ext. Damiana, 2 grs.
 Ext. Nux vomica, 1-8 gr.

DANDELION. Taraxacum, U. S. P. Taraxacum officinale, *Weber.*
 (Herb; root, U.S.P.)

Synonyms.—Taraxacum Dens-leonis, *Desfontaines*; Dent-de-leon, Fr.; Löwenzahn, Ger.

Properties.—Tonic, alterative, diuretic and laxative. Dandelion is chiefly employed as a hepatic stimulant, especially in chronic affections in which the functions of the liver are not properly performed. It is especially of use in dyspepsia, and in the treatment of cutaneous eruptions associated with hepatic derangements, and is also employed in gout, hectic fever, etc.

Dandelion herb, compressed in ounce packages for retailing purposes
Dandelion root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Dandelion, 180 grs.

Dandelion Compound.

Dandelion, 10 grs.

Cherry, 10 grs.

Licorice, 20 grs.

Aromatics, q. s.

Extracts.

Fluid Extract Dandelion, U. S. P.—Miscible with water.
 Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

Fluid Extract Dandelion Comp.—Not miscible with water without some precipitation.
 Each pint represents: Dandelion, 13 1-2 ozs.; Mandrake, 2 1-4 ozs.;
 Conium leaves, 1 1-8 oz.
 Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Fluid Extract Dandelion and Senna.—Miscible with water.
 Sixteen fluidounces (imperial) represent: Dandelion, 8 ozs.; Senna, 8 ozs. Tonic and laxative, with slight action on the liver.
 Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Fluid Extract Sarsaparilla and Dandelion.—Miscible with water.
 Each pint represents: Sarsaparilla, 8 ozs.; Dandelion, 8 ozs. A useful alterative in eruptive diseases depending on derangement of the hepatic functions.
 Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

DANDELION—Continued.

Powdered Extract Dandelion.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered dandelion. Dose, 10 to 30 grains (0.6 to 2 Gm.).

Solid Extract Dandelion, U. S. P.—One part equals five of the root. Dose, 10 to 30 grains (0.6 to 2 Gm.).

Pills.

Dandelion and Leptandrin, 2 grs.
Ext. Dandelion, 1 1-3 gr. Leptandrin, 2-3 grs.

Quinine Compound and Extract Dandelion, 4 1-2 grs.
Quinine bisulph., 1 1-4 gr. Arsenious Acid, 1-24 gr.
Iron sulph. exsic., 2 gr. Ext. Dandelion, 1 1-4 gr

Taraxacum Extract, 3 grs.

DEER-TONGUE. *Trilisa odoratissima, Cassini.* (Leaves.)

Synonyms.—*Liatrix odoratissima, Willdenow*; Vanilla leaf.

Properties.—Aromatic, stimulant, tonic, diaphoretic. An agreeable addition to nauseating or griping mixtures. The infusion (one ounce to a pint of hot water), taken freely, is useful in colds with suppressed secretions.

Deer-Tongue, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Deer-Tongue.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

DITA. *Alstonia scholaris, R. Brown.* (Bark.)

Properties.—In India Dita bark is regarded as a valuable anti-periodic and tonic. It has even been pronounced a perfect substitute for cinchona bark, and it may prove useful in some cases of remittent and intermittent fever which do not yield to the ordinary remedies. Waring says that it has proved valuable in chronic diarrhoea and in the advanced stages of dysentery. It is also prescribed during convalescence from fevers, etc., and is no doubt a valuable general tonic.

Alstonia bark (Dita), coarse powder.

PREPARATION.

Fluid Extract Dita bark.—Miscible with water.

Dose, 2 to 8 minims (0.13 to 0.5 C.c.).

DITTANY. *Cunila mariana, Linné.* (Herb.)

Properties.—Carminative and diaphoretic. Used popularly in colds and in flatulency, and also as an emmenagogue.

Dose, a wineglassful (60 C.c.) of a one ounce to the pint infusion.

Dittany herb, compressed in ounce packages for retailing purposes.

DOGWOOD (Cornus, U.S.P.). *Cornus florida, Linné.* (Bark of root, U.S.P.; flowers.)

NOTE.—Not Jamaica Dogwood, q. v.

Properties.—Tonic, astringent, slightly stimulant, and anti-periodic. May be used where quinine is indicated and cannot be administered, owing to idiosyncrasy or other causes, and is valuable in all cases where tonics are required, in periodical and typhoid fevers, and in pyrosis, etc. It increases the strength and frequency of the pulse.

Dogwood bark of root, U. S. P., compressed in pound packages for retailing purposes.

Dogwood flowers, compressed in ounce packages for retailing purposes.

DOGWOOD—Continued.**PREPARATIONS.**

Cornin, a so-called **Concentration**,* a powdered preparation from Dogwood bark of root.
Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Extracts.

Fluid Extract Dogwood, U. S. P.—Miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Dogwood.—One part equals 6 of bark.
Dose, 2 to 8 grains (0.13 to 0.5 Gm.).

Pills.

Cornin, 2 grs.

Cornus Florida Extract, 2 grs.

Tonic, Waxham's, 2 grs.

Cinchonine sulph., 1 gr.

Ext. Prickly-ash bark, 1-4 gr.

Ext. Cinchona bark, red, 1-4 gr.

Ext. Dogwood bark, 1-4 gr.

Powd. Capsicum, 1-4 gr.

DUBOISIA MYOPOROIDES, R. Brown. (Leaves.)

Properties.—Poisonous. An efficient substitute for atropia. The solid extract and the alkaloid Duboisine are now largely employed as being less irritating to mucous membranes, more prompt in mydriatic action, and effects of shorter duration than those of atropia. The fluid extract meets the demand for a preparation adapted to internal administration. Constitutional effects are similar to those of belladonna, but it is less of a cerebral excitant and more of a calmative and hypnotic. It has been found useful in relieving night-sweats of phthisis, puerperal mania, vesical tenesmus, etc.

NOTE.—Antagonistic to morphine.

ANTIDOTES.—Emetics (sulphate of zinc, mustard, etc.). Morphine to combat nervous disturbances as they arise.

ACTIVE PRINCIPLE.

Duboisine Sulphate.—Alkaloid. Chiefly employed to dilate the pupil of the eye; its effects being more transient than those of atropine. Like daturine, this alkaloid is probably identical with hyoscyamine.
Dose, 1-100 grain (0.0006 Gm.) and upwards.

PREPARATIONS.

Fluid Extract Duboisia.—Miscible with water without material precipitation.

Dose, 5 to 10 minims (0.3 to 0.6 C.c.)

Solid Extract Duboisia.

Dose, 1-4 to 3-4 grain (0.016 to 0.05 Gm.).

DUMB CANE. Dieffenbachia seguine, Schott.

Synonym.—Caladium seguinum, *Willdenow.*

Properties.—Expectorant and diaphoretic; somewhat similar to Indian turnip in action.

PREPARATION.

"German Tincture" Dumb Cane.—An imported unofficial preparation from the green plant.

Dose, 2 to 5 minims (0.12 to 0.3 C.c.).

DWARF ELDER. Aralia hispida, Ventenat. (Rhizome.)

Synonyms.—Wild Elder, Bristly Sarsaparilla.

Properties.—A valuable diuretic. Employed in dropsy, gravel, suppression of urine and other urinary disorders.

Dwarf-Elder root, compressed in pound packages for retailing purposes.

* See note in Preface on Concentrations.

DWARF ELDER—Continued.**PREPARATION.****Fluid Extract Dwarf Elder.**—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

ELATERIUM. The sediment deposited by the expressed juice of *Momordica Elaterium*, *Linné*.*Properties.*—The most powerful of the hydragogue cathartics, in large doses produces nausea, vomiting and great depression. Most frequently employed in abdominal dropsy. Should be used with great caution.

Dose, 1-10 to 1-2 grain (0.006 to 0.03 Gm.).

ACTIVE PRINCIPLE.**Elaterin, U. S. P.** A crystalline neutral principle obtained from *Elaterium*, the substance deposited by the juice of the fruit on standing. About four times the strength of an average sample of *Elaterium*. Insoluble in water; more active in alcoholic solution than in powder.

Dose, 1-20 to 1-10 grain (0.003 to 0.006 Gm.).

PREPARATIONS.**Pills.****Elaterin** (crystals), 1-20 gr.**Elaterium**, Clutterbuck's, 1-10, 1-8 and 1-4 gr.**Tablet Triturates.****Elaterin** (crystals), 1-40, 1-20 and 1-10 gr.**ELDER.** *Sambucus*, U. S. P. *Sambucus canadensis*, *Linné*. (Bark; flowers, U.S.P.)

See also European Elder.

Properties.—Bark: hydragogue cathartic; has been long in use as a remedy in dropsy. Flowers: stimulant, diaphoretic, and somewhat diuretic.

Employed externally as fomentations in erysipelas, glandular engorgements etc.

Elder bark, compressed in pound packages for retailing purposes.**Elder flowers, U. S. P.**, compressed in ounce packages for retailing purposes.**PREPARATIONS.****Fluid Extract Elder leaves.**—Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

Solid Extract Elder flowers.—One part equals 4 of the flowers.

Dose, 10 to 20 grains (0.6 to 1.3 Gm.).

ELECAMPANE. *Inula*, U. S. P. *Inula Helenium*, *Linné*. (Root.)*Synonyms.*—*Radix Helenii*, P. G.; *Helenenwurz*, Ger.; *Aunée*, Fr.*Properties.*—Aromatic, stimulant, expectorant, emmenagogue, and diaphoretic. Used principally in pulmonary affections, attended with weakness of the digestive organs, or with general debility. It has also been highly recommended both as an external and an internal remedy in tetter, psora, and other diseases of the skin.**Elecampane root, U. S. P.**, compressed in pound packages for retailing purposes.**PREPARATIONS.****Fluid Extract Elecampane.**—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Inulain, a so-called **Concentration**,* a powdered preparation from *Elecampane* root.

Dose, 2 to 5 grains (0.13 to 0.3 Gm.).

NOTE.—This contains so large a proportion of helenin, or elecampane cam-

* See note in Preface on Concentrations.

ELECAMPANE—Continued.

phor, that the contents of the bottle will often show an efflorescence of crystals of this substance, which must not be mistaken for mold. The name Inulin, adopted by some, should not be applied to this preparation, since it is already employed by chemists to designate an inert substance, a variety of starch, which is one of the constituents of the drug.

ELEPHANT'S-FOOT. *Elephantopus tomentosus*, *Linné.* (Herb.)

Properties.—Diaphoretic, expectorant, and, in large doses, emetic. Introduced on the authority of Dr. Newlon, of Kansas, who reports excellent results from its use as an expectorant.

PREPARATION.

Fluid Extract Elephant's-foot.—Miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C.c.).

EPHEDRA ANTISYPHILITICA, *C. A. Meyer.* (Herb.)

Synonyms.—Whorehouse tea; Canutillo, Tepopote, Span.

Properties.—Long familiar under the names above given to the residents of Arizona and adjacent country as a remedy in venereal affections. It seems to be especially indicated as a remedy in gonorrhœa.

PREPARATION.

Fluid Extract Ephedra Antisyphilitica.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

ERGOT. *Ergota*, U. S. P. *Claviceps purpurea*, *Tulasne.*

Synonyms.—Spurred Rye; Secale Cornutum, Blé cornu, Fr.; Mutterkorn, Ger.

Properties.—Uterine motor stimulant; causes tonic contraction of involuntary muscular fibre, and is thus hemostatic, through action on capillaries. Used to aid parturition, to control internal hemorrhages, wherever located; to relieve local congestions, and to destroy morbid growths.

ACTIVE PRINCIPLES.

Ergotin.—Bonjean's formula; almost identical with the extract of ergot of the German pharmacopœia.

Dose, 1-3 to 3 grains (0.02 to 0.2 Gm.).

Ergotin, Purified, for hypodermic injection.—One minim represents 5 grains of selected Ergot. It may be diluted with water, if desired, for administration.

Dose, 2 to 6 minims (0.13 to 0.4 C.c.).

Sclerotic Acid.—Ergotinic acid of Kobert. According to Dr. Podwysotky, it is the chief active constituent of ergot. One grain represents about 25 grains of good ergot.

Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.). Solution 18 grains to 1 fluidounce corresponds in strength to the fluid extract of ergot.

PREPARATIONS.**Extracts.**

Fluid Extract Ergot, U. S. P.—Miscible with water.

Dose, 5 to 60 minims (0.3 to 4 C.c.). In inertia uteri give 20 to 60 minims (1 3 to 4 C.c.), repeated every 15 to 20 minutes until the desired effect is obtained.

Fluid Extract Ergot, Etherial.—Miscible with water.

Doses same as U. S. P. preparation above.

Solid Extract Ergot, U. S. P.—One part equals 5 of ergot.

Dose, 1 to 5 grains (0.06 to 0.3 Gm.).

Normal Liquid * Ergot.—An assayed preparation.

Dose, 5 to 60 minims (0.3 to 4 C.c.). In inertia uteri give 20 to 60 minims (1.3 to 4 C.c.), repeated every 15 to 20 minutes until the desired effect is obtained.

* See note in Preface on Normal Liquids.

ERGOT—Continued.**Pills.****Emmenagogue, Improved, 4 1-2 grs.**

Ergotin, 1 gr.

Purified Aloes, 1 gr.

Oil Savin, 1-2 gr.

Ext. Black Hellebore, 1 gr.

Iron sulph. exsic., 1 gr.

Emmenagogue, No. 4, 4 1-2 grs.

Ergotin, 1 gr.

Ext. Cotton root, 1 gr.

Purified Aloes, 1 gr.

Iron sulph. exsic., 1 gr.

Oil Savin, 1-4 gr.

Ergotin, 1-10, 1-2, 1, 2 and 3 grs.**Ergotin and Cannabis Indica, 1 1-2 gr.**

Ergotin, purified, 1 gr.

Ext. Cannabis Indica, 1-2 gr.

Solution, Sclerotic Acid.—Composed of Sclerotic Acid two parts and Scleromucir one part, both constituents of Ergot of Rye. One fluidrachm represents 1-2 grain of this combination, the proportion of each being the same as in Ergot. Successfully used by prominent specialists in the form of subcutaneous injections as parturients.

Tablet Triturates, Ergotin, 1-10 and 1-4 gr.**Wine, Ergot, U. S. P.**—Each fluidounce represents 68 grains Ergot.

Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

EUCALYPTUS GLOBULUS, U. S. P., Labillardière. (Lvs., U. S. P.; oil.)

Synonyms.—Australian Gum Tree, Blue Gum Tree, Fever Tree.

Properties.—Stimulant, aphrodisiac, anti-spasmodic and eminently antiseptic in its action. It is recommended in the treatment of septic fevers, diphtheria, chronic gastric and intestinal catarrh, chronic bronchitis, etc. As an anti-spasmodic it is useful in asthma, but its chief uses depend upon its antiseptic character. Thus it is employed in the treatment of foetid breath, ulcers (syphilitic and otherwise), purulent catarrhal affections of the bladder, urethra and vagina, spongy and bleeding gums, etc. Externally, suitably diluted, the fluid extract is employed as a disinfectant lotion in gangrenous or foetid suppuration, foul ulcers and offensive discharges of all kinds.

Eucalyptus Oil.

Dose, 5 to 30 minims (0.3 to 2 C.c.).

PREPARATIONS.**Capsules, Oil Eucalyptus, in soft gelatin.**

Oil Eucalyptus, true, 5 minims.

Oil Sweet Almond, 5 minims.

Also in hard gelatin, substituting Olive for the Almond oil.

A favorite method with Prof. H. C. Wood of exhibiting this remedy as a stimulating expectorant. He recommends that it be given 4 times daily in 10 minim doses.

Extracts.**Fluid Extract Eucalyptus, U. S. P.**—Not miscible with water.

Dose, 15 to 60 minims (1 to 4 C.c.).

Powdered Extract Eucalyptus.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Eucalyptus leaves.

Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Solid Extract Eucalyptus.—One part equals 5 of leaves.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Elixir Eucalyptus Compound.—Each fluidounce represents ingredients specified.

Eucalyptus Globulus, 15 grs.

Wild Cherry, 16 grs.

Gentian, 4 grs.

Licorice, 15 grs.

Dandelion, 20 grs.

Syr. Yerba Santa Arom., 30 mins.

Inhalant, Eucalyptus, No. 7 of P., D & Co.'s series of inhalants.

Oil Eucalyptus, 1 fluidrachm.

Used with advantage in fetid bronchitis, diphtheria, etc.

Lozenges, Eucalyptus, containing:

Ext. Eucalyptus, true, 3 grs.

Ext. Licorice, 1 gr.

EUCALYPTUS—Continued.**Pills.****Eucalyptus Compound**, 1 5-8 gr.

Ext. Eucalyptus, 1 gr.

Ext. Canadian Hemp, 1-2 gr.

Sanguinariin, 1-8 gr.

Eucalyptus Extract, 2 grs.**Fever and Ague**, 2 grs.

Ext. Eucalyptus, 1-4 gr.

Chinoidin, 1 gr.

Iron ferrocyanide, 1-2 gr.

Powd. Capsicum, 1-4 gr.

Arsenious acid, 1-200 gr.

Soap, Medicated Eucalyptol, containing 5 per cent. or 84 minims Oil of Eucalyptus. This soap has not only stimulating and astringent, but also pleasant balsamic qualities. It is, therefore, a useful disinfectant application to all foul-smelling wounds and ulcers. In bromidrosis, or fetid perspiration, it is very efficacious.

EULACHON OIL. The pure oil from the entire fish *Thaleichthys pacificus*, *Girard*.

Synonyms.—Oolachon, Houlachan, etc.—imitating the sound of the Indian name; Candlefish—from the large amount of oil which the fish contains.

Properties.—"As an efficient medicinal agent in the treatment of phthisis and other wasting diseases, it bids fair to become a rival of that well-established remedy, cod-liver oil.* No difficulty in digesting the oil is experienced. It is also less disagreeable in flavor than cod-liver oil, and said to be equally as efficacious in promoting nutrition in scrofulous and tuberculous subjects, and may be administered in the same doses and forms of combination.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

PREPARATION.**Emulsion, Eulachon Oil**, 50 per cent. oil.

Dose, for an adult, 1-2 fluidounce (15 C.c.) 3 or 4 times a day.

EUPHORBIA PILULIFERA, *Linné*. (Herb.)

Properties.—This herb is becoming a favorite domestic remedy in Australia, where it is a common roadside weed. It is regarded as an infallible remedy for colds, coughs, bronchial affections, in short, all diseases of the respiratory tract; but it is more especially esteemed for the prompt and complete relief it is said to give sufferers from asthma.

PREPARATION.**Fluid Extract Euphorbia Pilulifera**.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

EUROPEAN CENTAURY. *Erythræa Centaurium*, *Persoon*. (Herb.)

Synonyms.—Herba Centaurii, P. G.; Centaurée, Fr.; Tausendgueldenkraut, Ger.

Properties.—Diaphoretic and slightly tonic Useful in the debility following malaria, and as a hot drink in colds.

Dose, of an ounce to the pint infusion, a wineglassful (60 C.c.).

European Centaury, compressed in ounce packages for retailing purposes.

* New York Medical Journal, Nov. 29, 1884.

EUROPEAN ELDER. *Sambucus nigra*, *Linné.* (Bark)

Properties.—Alterative, and in sufficient doses cathartic. It is regarded as a valuable adjunct to tonics in the convalescence from malarial fever, and in chronic, rheumatic, gouty and syphilitic affections.

PREPARATIONS.

Fluid Extract European Elder.—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract European Elder.—One part equals 8 of bark.

Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

EVENING PRIMROSE. *Oenothera biennis*, *Linné.* (Leaves.)

Properties.—A mild, but efficient, sedative to nervous sensibility, acting more especially upon the pneumogastric nerve. Indicated in such cases of respiratory or gastric troubles as involve a morbid sensitiveness either in the laryngeal, pulmonary or gastric branches of that nerve, whether of an acute or chronic character. Worthy of trial in the treatment of such affections as whooping-cough, spasmodic asthma and certain morbidly sensitive conditions of the stomach, interfering with a healthy digestion. An ointment of Evening Primrose has been found very efficacious in curing porrigo (tetter) and other cutaneous affections of infants. It has also proved of value as an application to ulcers.

Evening Primrose, compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.

Fluid Extract Evening Primrose.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Glycerole Evening Primrose.—Each fluidounce represents 120 grs.

Dose, 2 fluidrachms (8 C.c.).

Ointment Evening Primrose.

Fluid Extract Evening Primrose, P., D. & Co.'s.....4 fl. ozs.

Lard.....1 pound.

Heat the lard till melted, add the fluid extract, and stir till cold.

EYEBRIGHT. *Euphrasia officinalis*, *Linné.* (Leaves.)

Synonym.—Augentrost, Ger.

Properties.—Of utility in catarrhal conjunctivitis through its mild astringency, and also in chronic catarrhs. It mitigates the catarrhal symptoms of hay-fever, and it will abort an attack of nasal catarrh with lachrymation.* For the latter purpose it should be given every two hours.

Dose, of an ounce to the pint infusion, 1-2 to 2 fluidounces (15 to 60 C.c.).

Eyebright leaves, compressed in ounce packages for retailing purposes.

FALSE BITTERSWEET. *Celastrus scandens*, *Linné.* (Bk. of rt.)

Synonyms.—Climbing Bittersweet, Fever Twig, Staffvine, Wax-work.

Properties.—Has considerable reputation in domestic practice as a diaphoretic, diuretic, anti-syphilitic and discutient.

False Bittersweet bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract False Bittersweet.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

FENNEL. *Foeniculum*, U. S. P. *Foeniculum officinale*, *Allioni.*

Synonyms.—Foeniculum vulgare, *Gaertner*; Fenouil, Fr.; Fenchelsamen, Ger.

Properties.—Carminative and aromatic. Used to advantage in flatulent colic, and as a corrigent of unpleasant medicines, particularly senna and

*Potter, "Materia Medica and Therapeutics," p. 189.

FENNEL—Continued.

rhubarb. It is recommended for these purposes because of the absence of any exciting property.

PREPARATION.

Fluid Extract Fennel seed.—Not miscible with water.
Dose, 10 to 30 minims (0.65 to 2 C.c.).

FEVERFEW. *Pyrethrum Parthenium, Smith.* (Herb.)

Synonyms.—Feather-few; Matricaire, Fr.; Mutterkraut, Ger.

Properties.—Carminative, emmenagogue, tonic. Much used in cases of flatulent or atonic dyspepsia, amenorrhœa, dysmenorrhœa, hysteria, and in some febrile diseases. Externally as a fomentation to local inflammations.

Feverfew herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Feverfew.—Not miscible with water without some precipitation.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

FIREWEED. *Erechthites hieracifolia, Rafinesque.* (Leaves.)

Properties.—Tonic and astringent. It has acquired a reputation in diseases of the mucous tissues, whether of the lungs, stomach or intestinal canal. Has been successfully used in summer complaint of children, when other means have failed. Its greatest reputation has been obtained as a remedy for piles, administered internally as a fluid extract, and externally in the form of an ointment.

Fireweed leaves, compressed in ounce packages for retailing purposes.

PREPARATION AND FORMULA.

Fluid Extract Fireweed.—Miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Ointment Fireweed.

Fluid Extract Fireweed, P., D. & Co.'s.....	4 fl. ozs.
Lard	.8 ozs.

Melt the lard, add the extract, and stir till cold.

FISH-BERRIES. *Anamirta cocculus, Wight & Arnott.* (Fruit.)

Synonyms.—Cocculus Indicus; Coque du Levant, Fr.; Fischkörner, Ger.

Properties.—Poisonous. Chiefly used in nervous diseases. Efficacious in partial palsies accompanied by giddiness, as also in hysterical, choreic and epileptic hemiplegia, and in nervous vomiting.*

Externally it is used as a parasiticide, particularly to destroy vermin in the hair, but great caution must be exercised in its use for this purpose.

ANTIDOTES.—None reliable known. Use emetics (apomorphine, zinc sulphate, mustard), with copious draughts of warm water, etc. Treat symptoms as they arise on general principles. Anæsthetics antagonize its spasm-producing action.

ACTIVE PRINCIPLE.

Picrotoxin, U. S. P.—An ointment containing 10 grains to the ounce has been used successfully in tinea capitis. Should not be applied when the skin is abraded. Dr. Murrell, of London, Eng., finds Picrotoxin in doses of 1-60 grain (0.001 Gm) an effective remedy for pathological sweating, e.g., in the night-sweats of phthisis, etc. It is prescribed also, in minute doses, in certain forms of dyspepsia, and as a remedy for leucorrhœa, dysmenorrhœa, and other uterine disorders.
Dose, 1-100 to 1-50 grain (0.0006 to 0.0013 Gm.).

* Napheys, "Medical Therapeutics," p. 119.

FISH-BERRIES—Continued.**PREPARATIONS.**

Fluid Extract Fish-berries (Cocculus Indicus).—Not miscible with water.

Dose, 1-4 to 1 minim (0.016 to 0.065 C.c.), cautiously increased.

Hypodermic Tablet, Picrotoxin, 1-50 gr.

Pill, Picrotoxin, 1-60 gr.

FIVE-FINGER. *Potentilla canadensis, Linné.* (Herb.)

Synonyms.—Cinquefoil; Fünffingerkraut, Ger.; Quintefeuille, Fr.

Properties.—Astringent and febrifuge. Useful in diarrhoea and other complaints for which astringents are usually indicated.

Dose, wineglassful (60 C.c.) of an ounce to the pint infusion.

Five-finger (Cinquefoil) herb, compressed in ounce packages for retailing purposes.

FIVE-FLOWERED GENTIAN. *Gentiana quinqueflora, Lamarck.* (Herb.)

Synonyms.—Gall of the Earth, Bilious Weed, Frostblow,

Properties.—This plant has been found of much value in liver affections, jaundice, etc. "It is well adapted to chronic ague where there is much debility and it has succeeded where quinine has failed. In intermittent fever no preparatory treatment need precede its use. In hepatic disorders there are few medicines of more worth a new and reliable chologogue."*

PREPARATION.

Fluid Extract Five-flowered Gentian.—Not miscible with water without some precipitation.

Dose, 10 to 30 minims (0.6 to 2 C.c.) in water four times a day.

FLEABANE. *Erigeron canadense, Linné.* (Leaves; oil.)

Synonyms.—Canada Fleabane, Colt's Tail, Pride Weed, Horseweed, Bitter Weed, Fire Weed; Vergerette, Fr.; Berufkraut, Ger.

Properties.—Slightly tonic, with more active diuretic and astringent properties. Useful in diarrhoea, and in dropsical affections. It seems to act more particularly on the uterus, and the urinary organs, and is prescribed for uterine hemorrhage, strangury, dysuria of children, and in many nephritic affections. It owes its virtue to the volatile oil it contains.

Fleabane leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Capsules.—In soft gelatin.

Castor Oil and Oil of Fleabane.

Castor Oil, 5 minims.

Oil Fleabane, 5 minims.

Erigeron Oil, 10 minims.

Fluid Extract Fleabane.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

FLORIDA ALLSPICE. *Calycanthus floridus, Linné.* (Bark.)

Properties.—Aromatic stimulant, making an agreeable and valuable addition to diarrhoea mixtures, and in cases of capulous colic.

PREPARATION.

Fluid Extract Florida Allspice.—Not miscible with water without some precipitation.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

* Morse, "New Therapeutical Agents," p. 123.

FLOWERING SPURGE. *Euphorbia corollata*, *Linné*.

Synonyms.—Wild ipecac, Wild Hippo, Euphorbe à Grand Fleurs, Fr.; Wolfsmilchwurzel, Ger.

Properties.—Emetic, diaphoretic, expectorant and epispastic. In large doses it produces irritation of the stomach and bowels, with hypercatharsis. It has been used with success in chronic bronchitis, laryngitis and vesical catarrh. In obstinate bilious dysentery it has proved effectual, and in dropsical diseases, especially hydrothorax and ascites, it will sometimes evacuate the water when other agents prove useless.

PREPARATION.

Fluid Extract Flowering Spurge.—Miscible with water.

Dose, as emetic, 20 minims (1.3 C. c. ; as cathartic, 10 minims (0.65 C. c.); as diaphoretic 5 minims (0.3 C. c.).

FLY AGARIC. *Agaricus muscarius*, *Linné*.

Synonym.—Intoxicating Mushroom of Siberia.

Properties.—Poisonous. Respiratory and cardiac depressant, intoxicant to the cerebrum. Its physiological action is similar to that of physostigmine, and is antagonistic to that of atropine. It stimulates powerfully the salivary and cutaneous secretions, contracts the pupil, etc. It has proved valuable in the night-sweats of phthisis, in diabetes insipidus, incipient pulmonary congestion, and in constipation due to intestinal torpor and deficient secretion. Generally exhibited in the form of the alkaloid salts.

Antidote.—Atropine is a perfect antagonist, so are also to some extent digitalin and pilocarpine.

ACTIVE PRINCIPLE.

Muscarine Sulphate.—Derivative salt from the alkaloid. Very hygroscopic.

Dose, 1-50 to 1-15 grain (0.0013 to 0.004 Gm.).

PREPARATION.

Pill, Muscarine Sulphate, 1-100 gr.

FOXGLOVE. *Digitalis*, U. S. P. *Digitalis purpurea*, *Linné*. (Leaves.)

Properties.—Poisonous. In small doses a cardiac tonic, diminishing the frequency and increasing the force of the heart's contractions. In overdoses poisonous, and peculiarly liable to produce toxic effects from so-called cumulative action. Used in cardiac affections generally where the heart's action is abnormally weak. Indirectly it is diuretic also, and so is especially useful in dropsies depending on cardiac embarrassment or renal obstruction.

Antidotes.—Tannin (strong tea, decoction of oak bark). Emetics, early, if needed. Alcoholic stimulants, opium, rest in recumbent position; aconite.

Foxglove leaves, U. S. P., compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLE.

Digitalin.—Amorphous. (Merck's.) Various preparations, representing more or less perfectly the active properties of foxglove, have been offered under the name of digitalin. These are all, in fact, mixtures of several compounds (digitalin, digitonin, digitalein, digitoxin, and derivatives from these). So great are the differences between the different articles sold under the name of digitalin, that the U. S. P., 1880, omits the preparation entirely. There is, however, no doubt that both the French and German digitalin are very active preparations, representing at least partially the therapeutic virtues of the drug, and physicians may prescribe them with advantage, if careful to specify the variety they have learned to use.

Dose, 1-50 grain (0.0013 Gm.), cautiously increased.

Digitalin.—Crystallized, Fr. Codex. This is the digitalin of M. Nativelle. Possibly not in the strict sense the active principle of foxglove, being per-

FOXGLOVE—Continued.

haps a derivative compound. It has the advantage, however, over any amorphous preparation of constancy of composition, and it is believed that it represents very perfectly the therapeutic activity of digitalis.

Dose, 1-50 grain (0.0013 Gm.), cautiously increased.

PREPARATIONS AND FORMULÆ.**Abstract Digitalis, U. S. P.**

Normal Liquid or Fluid Extract Foxglove, P., D. & Co.'s 1 fl. oz.
Sugar of milk, q. s. to make 228 grs.

Evaporate the fluid extract nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well corked bottle.

Dose, 1-2 to 1 grain (0.03 to 0.06 Gm.). One part equals 2 of drug.

Digitalisin, a so-called **Concentration**,* a powdered preparation from Foxglove leaves.

Dose, 1-16 to 1-4 grain (0.004 to 0.016 Gm.).

Extracts.

Fluid Extract Foxglove, U.S.P.—Not miscible with water without some precipitation

Dose, 1 to 2 minims (0.06 to 0.13 C.c.), increased if necessary.

Powdered Extract Foxglove.—Of the same strength as the solid extract, U. S. P., prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Digitalis.

Dose, 1-4 to 1-2 grain (0.016 to 0.032 Gm.), cautiously increased.

Solid Extract Foxglove, Strictly Alcoholic.—One part equals 7 of the leaves.

Dose, 1-6 to 1-3 grain (0.01 to 0.02 Gm.), cautiously increased.

Solid Extract Foxglove, U. S. P.—One part equals 5 of leaves.

Dose, 1-4 to 1-2 grain (0.016 to 0.032 Gm.), cautiously increased.

Hypodermic Tablet Digitalin (Amorphous), 1-100 gr.

Infusion Foxglove, U. S. P.

Fluid Extract Foxglove, P., D. & Co.'s..... 2 fluidrachms.

Tincture of Cinnamon, P., D. & Co.'s..... 2 fl. ozs.

Hot Water..... 14 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Normal Liquid† Foxglove.

Dose, 1 to 2 minims (0.06 to 0.13 C.c.), cautiously increased, if necessary.

Pills.

Digitalis Leaves, 1-20 gr.

Digitalin, pure, 1-60 gr.

Digitalis Extract, 1-2 gr.

Digitalis Compound, 4 grs.

Powd. Digitalis, 1 gr.

Powd. Potass. nitrate, 2 gr.

Powd. Squill, 1 gr.

Phosphorus, Quinine and Digitalis Comp., 1 1-2 gr.

Phosphorus, 1-50 gr.

Powd. Digitalis, 1-2 gr.

Quinine sulph., 1-2 gr.

Powd. Opium, 1-4 gr.

Powd. Ipecac, 1-4 gr.

For other pills containing Foxglove see Pill Formulæ list appended for numbers 314—317, 332, 333.

Tablet Triturates.

Digitalin (crystals), 1-100, 1-75 and 1-50 gr.

Digitalis (Fluid Extract), 1-10, 1-2 and 1 minim

Tinctures.**Digitalis, U. S. P.**

Fluid Extract Foxglove, P., D. & Co.'s..... 1 fl. oz.

Dilute Alcohol..... 7 fl. ozs.

Dose, 8 to 25 minims (0.5 to 1.5 C.c.).

Digitalis "German Tincture."

An imported preparation of the green herb (unofficial).

Dose, 8 to 25 minims (0.5 to 1.5 C.c.).

* See note in Preface on Concentrations.

† See note in Preface on Normal Liquids.

FOXGLOVE—Continued (Tinctures).**Digitalis, Pharm. Ger.**

Fluid Extract Foxglove, P., D. & Co.'s..... 1 fl. oz.
 Dilute Alcohol..... 9 1-2 fl. ozs.
 Dose, 8 to 25 minims (0.5 to 1.5 C.c.).

Digitalis, U. S. P.

Fluid Extract Foxglove, P., D. & Co.'s..... 2 fl. ozs.
 Dilute Alcohol..... 12 fl. ozs.
 Dose, 5 to 20 minims (0.3 to 1.3 C. c.).

FRINGE-TREE. *Chionanthus virginica, Linné.* (Bark of root.)

Synonym.—Old Man's Beard.

Properties.—Aperient, alterative and diuretic, with some narcotic properties.

It has been found efficacious in bilious remittent fevers, as well as in obstinate intermittents, and an excellent tonic after convalescence from exhausting diseases. It is a cholagogue of some value, and according to Dr. J. A. Henning, one of the most eligible remedies in jaundice, associated with podophyllin or leptandrin.* In the form of a poultice, it is an excellent local application in external inflammations, ulcers and wounds.

Fringe-tree bark of root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Fringe-tree.—Not miscible with water without some precipitation.
 Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Fringe-tree.—One part equals 3 of drug.
 Dose, 5 to 20 grains (0.32 to 1.3 Gm.).

FROSTWORT. *Helianthemum canadense, Michaux.*

Synonyms.—Frostweed; Heliantheme de Canada, Fr.; Canadisches sonnenroschen, Ger.

Properties.—Astringent, tonic, alterative. Has long been used, either alone or in combination, in scrofulous diseases. In large doses it acts as an emetic. Diluted with water, the fluid extract is used as a gargle in sore throat, particularly in scarlatina; also as a lotion in prurigo.

Frostwort herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Frostwort (Frostweed).—Miscible with water without material precipitation.
 Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

GALANGAL. *Alpinia Galanga, Linné.* (Rhizome.)

Properties.—A stimulant aromatic, somewhat resembling ginger. Used as a stomachic to promote digestion and relieve flatulence.

Dose, of an ounce to the pint infusion, 2 fluidounces (60 C.c.).

Galangal root, compressed in pound packages for retailing purposes.

GALBANUM. Gum-resin from *Ferula galbaniflua, Boissier & Buhse.*

Properties.—Stimulant, carminative, expectorant. Chiefly used in chronic affections of the bronchial mucous membranes, and in uterine and vaginal catarrh; externally, in plaster form, is mildly stimulant and irritant.

Dose, 5 to 20 grains (0.3 to 1.3 Gm.).

PREPARATIONS.

Pill Galbanum Comp., U. S. P., 3 1-2 grs.
 Galbanum, 1 1-2 gr. Myrrh, 1 1-2 gr.
 Asafetida, 1-2 gr.

Plaster Galbanum, U. S. P.—In half pound rolls or cans.

*Therapeutic Gazette, p. 846, 1886.

GAMBOGE. Cambogia, U. S. P. Gum-resin from *Garcinia Hanburii*, *Hooker filius*.

Properties.—Cathartic, apt to occasion much sickness and griping, but acts better in combination. Useful in obstinate costiveness, amenorrhœa, and dropsy attended with torpidity of the bowels.

Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

PREPARATIONS.**Pills.**

Cathartic Compound, U. S. P., 3 1-2 gr.

Ext. Coloc. co., 13-10 gr.

Ext. Jalap, 1 gr.

Calomel, 1 gr.

Powd. Gamboge, 1-4 gr.

Cochia, 3 1-2 grs.

Ext. Coloc. co., 1 gr.

Purified Aloes, 1 gr.

Powd. Potass. sulph., 1-2 gr.

Powd. Gamboge, 1-2 gr.

Powd. Scammony, 1-2 gr.

Gamboge, 1-32 gr.

Gamboge Compound, 3 grs.

Powd. Gamboge, 3-5 gr.

Purified Aloes, 3-5 gr.

Powd. Cinnamon comp., 3-5 gr.

Soap, 1 1-5 gr.

For other pills containing Gamboge see Pill Formulæ list appended for numbers 27, 101, 103, 104, 260, 261, 263, 461.

GARDEN CELANDINE. Chelidonium, U. S. P. Chelidonium majus, *Linné*. (Herb.)

Synonyms.—Great Celandine, Tetterwort, Herbe à l'hirondelle, Fr.; Schöllkraut, Ger.

Properties.—A drastic purgative, resembling Gamboge in its action. Applied externally to subdue traumatic inflammations.

Garden Celandine herb, U.S.P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Garden Celandine.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

"German Tincture" Garden Celandine.—An imported preparation of the green herb (unofficial).

Dose, 8 to 60 minims (0.5 to 4 C.c.).

GARDEN LETTUCE. Lactuca sativa, *Linné*. (Leaves.)

Synonyms.—Lettuce, Eng.; Salat des Jardins, laitue sative, Fr.; Gartensalat, Gartenlattich, Ger.

Properties.—Faintly medicinal; narcotic and antispasmodic. Used popularly to allay nervous irritability, and excitement from mental worry.

Dose, a wineglassful (60 C.c.) of a one ounce to the pint infusion.

Garden Lettuce, compressed in ounce packages for retailing purposes.

GARLIC. Allium, U. S. P. Allium sativum, *Linné*. (Bulb.)

Synonyms.—Ail, Fr.; Knoblauch, Ger.

Properties.—A general stimulant, it quickens the circulation, excites the nervous system, promotes expectoration in debility of the lungs, produces diaphoresis or diuresis according as the patient is kept warm or cool, and acts upon the stomach as a tonic or carminative. Also said to be emmenagogue and anthelmintic. It has been employed with advantage in dyspepsia, flatulence, chronic catarrh, and other pectoral affections, in the nervous and spasmodic coughs peculiar to very young children, old atonic dropsies, calculous disorders, ascariides and pertussis. Externally, it acts as an irritant and rubefacient, besides imparting its effects on the system by absorption. It is applied to the soles of the feet as a revulsive in disorders of the head, and is used for quieting restlessness in the febrile

GARLIC—Continued.

complaints of children. The fluid extract diluted may be applied as a liniment in infantile convulsions, etc. Dropped on cotton and placed in the ear, it is efficacious in atonic deafness.

PREPARATION AND FORMULA.

Fluid Extract Garlic.—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

Syrup Garlic, U. S. P.

Fluid Extract Garlic, P., D. & Co.'s.....6 fl. ozs.

Simple Syrup.....23 fl. ozs.

Acetic Acid3 fl. ozs.

Add the acid to the fluid extract, and finally to the syrup; filter.

Dose for a child a year old, one teaspoonful.

GELSEMIUM, U.S.P. *Gelsemium sempervirens, Aiton.* (Rhizome.)

Synonyms.—Yellow Jasmine, Yellow Jessamine.

Properties.—POISONOUS. Gelsemium and its alkaloids produce not only paralysis of sensibility, but also of motility.* Contra-indicated where cardiac action is weak; useful whenever exaltation of function exists in sensory or motor nerves. Recommended in neuralgia, especially of the trigeminus, in ovarian neuralgia, in dysmenorrhoea, after-pains, etc.

ANTIDOTES.—Emetics (apomorphine, zinc sulphate, etc.). Sustain vital powers by stimulants (ammonia, alcohol, etc.). Morphine is the most complete physiological antagonist; atropine is also recommended.

Gelsemium root, U.S.P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLE AND DERIVATIVES.

Gelsemine.—Alkaloid, insoluble in water (1 in 650); its salts, mentioned below, are therefore to be preferably prescribed. The most important use of these is in the treatment of obstinate neuralgias, which yield to hypodermic injections of the remedy.

Dose, 1-100 to 1-40 grain (0.0006 to 0.0016 Gm.).

Gelsemine Hydrochlorate.—This crystalline salt is permanent in the air; solubility in water, 1 in 39.

Dose, 1-100 to 1-40 grain (0.0006 to 0.0016 Gm.).

Gelsemine Sulphate.—This is the most soluble of the gelsemine salts; permanent in the air, but not crystallizable.

Dose, 1-100 to 1-40 grain (0.0006 to 0.0016 Gm.).

Gelsemine Tartrate.—A readily soluble and very permanent salt.

Dose, 1-100 to 1-40 grain (0.0006 to 0.0016 Gm.).

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Gelsemium, U. S. P.—Not miscible with water without some precipitation.

Dose, 1 to 10 minims (0.06 to 0.6 C. c.).

Fluid Extract Gelsemium.—From the *green* root.

This is not so satisfactory a preparation to use as that of the U. S. P. from the dried root, see above.

Dose, 1 to 15 minims (0.06 to 1. C. c.).

Solid Extract Gelsemium, assayed.—One part equals 5 of root.

Dose, 1-5 to 2 grains (0.013 to 0.13 Gm.).

Gelsemperin, a so-called *Concentration*,† a powdered preparation from Gelsemium (*Synonym.*—Gelsemin).

Dose, 1-8 to 1 grain (0.008 to 0.065 Gm.).

Hypodermic Tablet, Gelsemine Hydrochlorate, 1-50 gr.

Normal Liquid‡ Gelsemium.—Standard, 10 C. c. of the normal liquid require for complete precipitation of the alkaloid 3 C. c. of Mayer's solution.

Dose, 1 to 10 minims (0.06 to 0.6 C. c.).

* Dujardin-Beaumetz, "New Medications," p. 277, (edition of Geo. S. Davis, 1886).

† See note in Preface on Concentrations.

‡ See note in Preface on Normal Liquids.

GELSEMIUM—Continued.**Pills.****Dysentery,** 2 1-6 grs.

Blue Mass, 1 gr.

Powd. Ipecac, 1 gr.

Gelsemperin, 1-6 gr.

Gelsemperin, 1-8 and 1-4 gr.**Gelsemium root,** 1-50 gr.

For other pills containing Gelsemium see appended Pill Formulæ list for numbers 5, 40, and 41.

Tinctures.**Gelsemium, B. P.**

Fluid Extract Gelsemium, P., D. & Co.'s..... 1 fl. oz.

Dilute Alcohol..... 7 fl. ozs.

Dose, 5 minims (0.3 C.c.), gradually increased.

Gelsemium, U. S. P.

Fluid Extract Gelsemium, P., D. & Co.'s..... 2 fl. ozs.

Dilute Alcohol..... 12 fl. ozs.

Dose, 5 minims to 1-2 fluidrachm (0.3 to 2 C.c.).

GENTIAN. Gentiana, U.S.P. *Gentiana lutea, Linné.* (Root.)

Properties.—Gentian possesses in a high degree, the tonic powers which characterize the simple bitters. It excites the appetite, invigorates digestion, moderately increases the force of the circulation and operates in fact as a general stimulant tonic. In very large doses, however, it is apt to disturb the stomach and irritate the bowels. Without reference to the disease, gentian is indicated in all atonic conditions unattended with active fever.

Gentian root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.

Elixirs.—Each fluidounce represents ingredients specified.

Gentian, 20 grs.**Gentian and Iron Chloride.**

Gentian, 16 grs.

Tr. Iron chlor. (tasteless), 16 mins.

Gentian, Iron Chloride and Lactated Pepsin.

Gentian, 16 grs.

Tr. Iron chlor. (tasteless), 16 mins.

Lactated pepsin, 20 grs.

Gentian and Iron Pyrophosphate.

Gentian, 20 grs.

Iron pyrophosphate, 8 grs.

Gentian and Lactated Pepsin.

Gentian, 20 grs.

Lactated pepsin, 20 grs.

Extracts.

Fluid Extract Gentian, U. S. P.—Not miscible with water without some precipitation.

Dose, 10 to 40 minims (0.6 to 2.5 C.c.).

Fluid Extract Gentian Compound.—For the extemporaneous production of Tincture Gentian Comp., U.S.P. Fourteen fluidounces, imperial, represent: Gentian, 8 ozs.; Orange peel, 4 ozs.; Cardamom seed, 2 ozs.

Powdered Extract Gentian.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Gentian.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Solid Extract Gentian, U. S. P.—One part equals 3 of root.

Dose, 5 to 10 grains, (0.3 to 0.65 Gm.).

Pills.**Carminative,** 3 1-4 grs.

Strychnine, 1-20 gr.

Powd. Black Pepper, 1 1-2 gr.

Powd. Ipecac, 2-3 gr.

Ext. Gentian, 1 gr.

Gentian Compound, 3 grs.

Ext. Gentian, 2-3 gr.

Powd. Rhubarb, 1 1-3 gr.

Purified Aloes, 2-3 gr.

Oil Caraway, 1-10 gr.

Gentian Extract, 2 grs.

See also numbers 36 and 102 in Pill Formulæ list appended.

GENTIAN—Continued.**Tinctures.****Gentian Compound, U. S. P.**

Fluid Extract Gentian Comp., P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol.....13 fl. ozs.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Gentian, Pharm. Ger.

Fluid Extract Gentian, P., D. & Co.'s.....3 fl. ozs.

Dilute Alcohol.....12 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

GERMAN CHAMOMILE. *Matricaria*, U. S. P. *Matricaria Chamomilla*, *Linné.* (Flowers.)*Properties.*—Anthelmintic, antispasmodic. In popular use as a diaphoretic, emmenagogue, etc., in warm infusion.**PREPARATIONS.****Fluid Extract German Chamomile.**—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

“German Tincture” Chamomile.—An imported preparation from the fresh flowers (unofficial).

Dose, 12 to 30 minims (0.74 to 2 C. c.).

GINGER. I. JAMAICA. *Zingiber*, U. S. P. *Zingiber officinale*, *Roscoe*, (Rhizome.)*Synonyms.*—Gingembre, Fr.; Ingwer, Ger.*Properties.*—A grateful stimulant and carminative, often employed in dyspepsia, flatulent colic, and the feeble state of the alimentary canal attendant upon atonic gout. It is an excellent addition to bitter infusions and tonic mixtures, imparting to them an agreeable warmth and exercising a cordial influence on the stomach. It is much used in diarrhœa, dysentery, cholera morbus, cholera infantum, hysteria, and to relieve nausea, pain and cramps in the stomach and bowels, and to obviate tenesmus.**PREPARATIONS AND FORMULÆ.****“Ale, Ginger”**—A wholesome summer beverage.

Fluid Ginger, soluble, P., D. & Co.'s.....4 fluidounces or q. s.

Sugar.....2 lbs. av.

Citric Acid.....100 grs.

Essence Lemon.....30 drops.

Water.....2 gals.

Mix the essence of lemon with the sugar, add the fluid extract ginger, then the water. The citric acid may be dissolved separately in a fluid-ounce of water, and added when the sugar is dissolved.

Fluid Extract Ginger, U. S. P.—Not miscible with water.

Dose, 5 to 20 minims (0.35 to 1.35 C. c.) diluted.

Fluid Ginger, Soluble.—Miscible with water.

One-half the strength of the official fluid extract. May be substituted to advantage in extemporaneous prescriptions for the official fluid extract or tincture, and affords a ready means of preparing ginger ale, ginger wine, syrup ginger, U. S. P., or a stronger syrup for flavoring soda water.

Lozenge, Ginger, 1 1-4 gr. (official lozenge only 1-3 this strength.)**Syrup Ginger, U. S. P.**—(15 grs. to the fl. oz.)

Our preparation may be extemporized by the two formulæ below. No. 1 gives the most satisfactory product.

No. 1.

Fluid Ginger, Soluble, P., D. & Co.'s.....1 fl. oz.

Simple Syrup.....15 fl. ozs.

No. 2.

Fluid Extract Ginger, P., D. & Co.'s.....1 fl. oz.

Sugar, in coarse powder.....26 troy ozs.

Water.....15 fl. ozs.

Rub the fluid extract with 10 troy ounces of the sugar, allow the alcohol to evaporate at a temperature not above 140° F., mix with the water, agitate till dissolved, and filter; add the remainder of the sugar, dissolve by agitation without heat, and strain.

Properties.—A valuable tonic, possessing also pronounced antiperiodic properties. It also increases the secretion of the intestinal mucous membrane and its glandular appendages, and promotes the flow of bile. As a result the stools become softer and more frequent. It is one of the best remedies for the gastric catarrh of chronic alcoholism, and the best sub-

GOLDEN-SEAL—Continued.

stitute for alcoholic stimulants when their use is to be abandoned. Useful in stomatitis, follicular pharyngitis, chronic coryza and syphilitic affections of the mouth, throat and nares, and atonic dyspepsia. Its action on the mucous tissues seems to depend more particularly on the white alkaloid, hydrastine. Its general tonic influence is derived from its yellow alkaloid, berberine [frequently called hydrastine (yellow) alkaloid]. A concentrated powdered preparation of Golden-seal, representing all the properties of the drug, is prepared under the name of hydrastin, see below.

Golden-seal root, U. S. P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLES.

Berberine Hydrochlorate.—Berberine is an alkaloid which occurs in a number of different plants, notably in the various species of *Berberis*, in golden-seal, gold thread, yellow-root, and in columbo. The yellow color and the bitterness of all of these is dependent, in part at least, on the presence of this alkaloid. From its occurrence in golden-seal, it has sometimes been improperly called hydrastine. Properties: tonic, hepatic, stimulant, anti-periodic, etc. Solubility 1 in 500.
Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

Berberine Phosphate.—More soluble in water than the last named. Its properties, therapeutic uses, and dose are the same. Solubility 1 in 10.

Berberine Sulphate.—Neutral. Of equal solubility with the Phosphate. Properties and dose similar to the above.

Hydrastine.—The white alkaloid of *Hydrastis canadensis*. It exerts a peculiar stimulating and alterative action upon the mucous membrane, and is a valuable remedy in atonic conditions of the bowels, and in all catarrhal affections of the mucous tracts. Especially useful in chronic affections of the bladder and genito-urinary system.
Dose, 1-16 to 1-4 grain (0.004 to 0.016 Gm.).

Hydrastine Hydrochlorate.—Pure hydrastine is soluble in alcohol and chloroform, but not in water. The hydrochlorate, or muriate, is freely soluble, and is therefore to be preferred in prescribing. It is much used by oculists in solutions containing 1 to 5 grains to the ounce. Also a valuable local remedy in gonorrhœa (1-4 to 1 grain, or more to the oz).
Dose, 1-16 to 1-4 grain (0.004 to 0.016 Gm.).

Hydrastine Sulphate.—Properties and uses identical with those of hydrastine hydrochlorate, and used in the same doses.

PREPARATIONS AND FORMULA.

Elixirs.—Each fluidounce represents ingredients specified.

Berberine and Iron.

Berberine phosphate, 2 grs.

Iron pyrophosphate, 4 grs.

Rhubarb and Potassa.

Rhubarb, 60 grs.

Potassium bicarbonate, 60 grs.

Golden-seal, 30 grs.

Extracts.

Fluid Extract Golden-seal, U. S. P.—Not miscible with water.
Dose, 10 to 30 minims (0.65 to 2 C.c.).

Fluid Extract Golden-seal, without Alcohol.—Miscible with water.

Dose, 10 to 30 minims (0.65 to 2 C.c.).

Fluid Golden-seal, Colorless.—Miscible with water.

Each pint of this fluid contains, in solution, 20 grains of the white alkaloid hydrastine, upon which the peculiar alterative action almost solely depends. As its name implies, it is a colorless preparation, entirely free from berberine to which the characteristic yellow color of golden-seal is due. When employed as a topical remedy, it has a marked advantage over the fluid extract in that no stains are produced. It is also devoid of bitterness.

Dose, 10 to 60 minims (0.65 to 4 C.c.).

Powdered Extract Golden-seal.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Golden-seal.
Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

Solid Extract Golden-seal.—One part equals 5 of root

Dose, 2 to .0 grains (0.13 to 0.65 Gm.).

GOLDEN-SEAL—Continued.

Hydrastin, a so-called **Concentration**,* a powdered preparation from Golden-seal, containing the proximate principles of the root, including the alkaloids Hydrastine and Berberine.
Dose, 1 to 3 grains (0.065 to 0.2 Gm.).

Pills.

Berberine Hydrochlorate, 1 gr.

Berberine Sulphate, 2 grs.

Berberine and Podophyllin, 1-4 gr.

Berberine sulph., 1-4 gr. Podophyllin, 1-30 gr.

Female, Leucorrhœa, 3 grs.

Hamamelin, 2 grs.

Hydrastin, 1-2 gr.

Senecin, 1-2 gr.

Hydrastin ("Concentration"), 1-30 and 1 gr.

Ox Gall Compound, 2 1-3 grs.

Ox Gall, 2 grs.

Purified Aloes, 1-10 gr.

Ext. Stramonium seed, 1-6 gr.

Berberine hydrochlor., 1-12 gr.

Tablet Triturates Hydrastine (white alkaloid), 1-25, 1-16, 1-8, 1-4 gr.

Tincture Golden-seal, U. S. P.

Fluid Extract Golden-seal, P., D. & Co.'s..... 4 fl. ozs.

Dilute Alcohol..... 17 fl. ozs.

Dose, 1-4 to 2 fluidrachms (2 to 8 C.c.).

GOLDTHREAD. *Coptis trifolia*, *Salisbury*. (Plant.)

Synonyms.—Mouthroot; Coptide, Fr.; Gelbe Niesswurz, Ger.

Properties.—A pure, bitter tonic, without astringency, resembling in its effects Columbo and Gentian. The infusion is in general use as a wash or gargle for ulceration of the mouth and fauces.

Goldthread, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Goldthread.—Miscible with water without material precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

GRAVEL-PLANT. *Epigæa repens*, *Linné*. (Leaves.)

Synonyms.—Trailing Arbutus, Ground Laurel, Winter or Mountain Pink, Mayflower.

Properties.—This plant is a diuretic and astringent, and is said to be highly beneficial in lithic acid gravel, in strangury and vesical catarrh. Has been freely used to advantage in diseases of the urinary organs, and of the pelvic viscera generally, particularly in those cases of local irritation where uva ursi and buchu are indicated. It has given relief where both buchu and uva ursi have failed.

Gravel-plant leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Gravel-plant.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

GREAT LAUREL. *Rhododendron maximum*, *Linné*. (Leaves.)

Synonyms.—Deer-tongue Laurel, Wild-rose Bay.

Properties.—Expectorant, stimulant, astringent. It is highly recommended by Dr. Mulholland in cases of obstinate cough.

PREPARATION.

Fluid Extract Great Laurel.—Miscible with water without material precipitation.

Dose, 5 to 15 minims (0.3 to 1 C.c.).

* See note in Preface on Concentrations.

GREEN-DRAGON. *Arisæma Dracontium*, Schott. (Corm.)

Synonyms.—*Arum Dracontium*, Linne; Dragon Root.

Properties.—Expectorant and diaphoretic. Said to be a certain remedy in dry, hacking coughs where there is considerable irritation. Is best administered in small doses well diluted.

PREPARATION.

Fluid Extract Green-dragon.—Miscible with water.

Dose, 1 to 10 minims (0.06 to 0.6 C.c.).

GREEN OSIER. *Cornus circinata*, L'Heritier. (Bark.)

Synonyms.—Round-leaved Dogwood, Round-leaved Cornel; Cornouiller à feuilles arrondies, Fr.; Canadische cornelrinde, Ger.

Properties.—Astringent, tonic, febrifuge. It is chiefly employed as a stomachic tonic. Its medicinal properties are similar to those of *Cornus florida*. (See Dogwood.)

Green Osier bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Green Osier.—Miscible with water without material precipitation.

Dose, 20 to 60 minims (1.3 to 4 C.c.).

GRINDELIA, U. S. P. *Grindelia robusta*, Nuttall. (Herb.)

Properties.—*Grindelia robusta* has attained a good deal of reputation in late years in the treatment of asthma, giving prompt relief, it is said, when other remedies have failed. It has also been recommended in bronchial affections, in pertussis, and in some renal maladies.

PREPARATIONS.

Elixir Grindelia Robusta Aromatic.—Each fluidounce represents 120 grains of *Grindelia*.

Extracts.

Fluid Extract Grindelia Robusta.—Miscible with water.

The drug is exhausted with an alkaline aqueous menstruum, thereby making it practicable to combine our fluid extract with syrups and aqueous fluids without precipitation—an obvious advantage.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Grindelia Compound.—Miscible with water.

Eighteen fluidounces, imperial, represent: *Grindelia Robusta*, 12 ozs.; *Senna* and *Rhubarb*, 3 ozs. each.

Efficacious in the treatment of asthma where there is a torpid condition of the bowels, upon which the asthmatic attacks are frequently immediately dependent.

Dose, 1 fluidrachm (4 C.c.), repeated every 3 or 4 hours as required.

Solid Extract Grindelia Robusta.—One part equals 5 of herb.

Dose, 5 to 15 grains (0.3 to 1 Gm.).

Glycerole Grindelia Robusta.—Each fluidounce represents 120 grs.

Pill Grindelia Robusta Extract, 3 grs.

GRINDELIA SQUARROSA, Dunal. (Herb.)

Synonym.—Ague Weed.

Properties.—The Spanish American remedy for intermittent fevers. "In paludal diseases of any kind it is of worth. For cystitis, leucocythæmia, enlargement of spleen, torpidity of the liver, and, (in combination with yerba santa,) for anasarca, it is of growing value."* It has also been found useful in chronic rheumatic affections.

PREPARATIONS.

Fluid Extract Grindelia Squarrosa.—Not miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

Solid Extract Grindelia Squarrosa.—One part equals 4 of drug.

Dose, 4 to 8 grains (0.26 to 0.5 Gm.).

* Morse, "New Therapeutical Agents," p. 126.

GROUND IVY. *Nepeta Glechoma, Bentham.* (Herb.)

Synonyms.—Gill-over-the-ground, Gill Run, Catfoot; Lierre terrestre, Fr.; Gundelrebe, Gundermann, Ger.

Properties.—Said to be stimulant, tonic, and pectoral; and has been recommended in diseases of the kidneys, asthma, jaundice, and hypochondria. Painters use it as a remedy for, and to prevent, lead colic.

Dose, wineglassful (66 C.c.) of an ounce to the pint infusion.

Ground Ivy, compressed in ounce packages for retailing purposes.

GUACO. Leaves from *Mikania guaco, Humboldt & Bonpland*: Botanical source of the root not yet determined.

Properties.—Guaco leaves are somewhat tonic in their action, stimulating also the secretions. Used as a prophylactic in epidemic cholera and for their remedial action in chronic diarrhoea, also with success in chronic rheumatism and atonic deafness. Dr. E. W. Pritchard highly recommends this remedy in connection with local applications in gouty paroxysms.

Guaco root is regarded in Mexico as a specific in rheumatic affections.

PREPARATIONS.

Fluid Extract Guaco leaves.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Guaco root.—Miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

GUAIAIC. Guaiaci Lignum, U. S. P. *Guaiacum officinale, Linné.* (Resin, wood.)

Properties.—Alterative, stimulant, diaphoretic, anti-rheumatic and anti-syphilitic. Formerly held in high repute as a remedy in constitutional syphilis, but is now replaced by potassium iodide. It is found efficient in promoting menstruation in all atonic conditions of the uterine system, and hence in dysmenorrhoea, and amenorrhoea of that origin. For rheumatism it should be taken in conjunction with Rochelle salt. The patient should be covered warm in bed, and abundantly provided with warm drinks, to excite profuse perspiration.

PREPARATIONS.

Fluid Extract Guaiac Wood.—Not miscible with water

Dose, 10 to 20 minims (0.65 to 1.3 C.c.).

Pills.

Calomel Compound, 3 grs.

Calomel, 3-4 gr.

Resin Guaiac, 1 1-2 gr.

Sulphurated Antimony, 3-4 gr.

Dupuytren, 3 1-4 gr.

Powd. Guaiac, 3 gr.

Powd. Opium, 1-8 gr.

Corrosive Sublimate, 1-10 gr.

For other combinations see Pill Formulæ list appended for numbers 39, 147, and 254.

Guaiac also enters into Syrup Sarsaparilla Comp., U. S. P., q. v.

GUARANA, U. S. P. *Paullinia sorbilis, Martius.* (A dried paste from the seeds.)

Synonym.—Brazilian Cocoa.

Properties.—The special use of paullinia is in the treatment of sick headache or migraine. It is adapted to the so-called nervous form of sick headache, and is less efficient when the attacks are due to stomachal troubles. As it possesses, directly or indirectly, restorative powers, it may be employed to promote constructive metamorphosis. Administered with this view, it may be given with advantage in the convalescence from acute maladies, incipient phthisis, and in wasting diseases generally.*

* Bartholow, "Materia Medica and Therapeutics," p. 326.

GUARANA—Continued.**PREPARATIONS.****Elixirs.**—Each fluidounce represents ingredients specified.**Celery and Guarana.**

Celery seed, 60 grs.

Guarana, 60 grs.

Guarana, 120 grs.**Extracts.****Fluid Extract Guarana, U. S. P.**—Not miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C.c.).

Powdered Extract Guarana, assayed.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered sugar of milk.

Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Solid Extract Guarana, assayed.—One part equals 3 of the dried paste.

Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Pills.**Guarana Extract**, 3 grs.**Headache**, 3 grs

Lactated Pepsin, 1 gr.

Guarana, 1-2 gr.

Sodium bicarb., 1 gr.

Cypripedin, 1-2 gr.

GURJUN BALSAM. An oleoresin obtained from various trees of the Dipterocarpaceæ.*Synonyms.*—Wood-oil, Balsamum Dipterocarpi.*Properties.*—Similar to those of Copaiba, but less nauseous and communicates no unpleasant odor to the breath. One of the best dressings in blennorrhagic vaginitis (one part of the balsam to two of lime-water).* Has been successfully used in leprosy.**Capsules Gurjun Balsam**, 10 minims. In soft gelatin.**HAIRCAP MOSS.** *Polytrichum juniperinum, Hedwig.* (Plant.)*Synonyms.*—Bearsbed, Ground Moss, Robin's Rye.*Properties.*—A powerful diuretic. In full doses every half hour it is said to have removed from a dropsical patient twenty to forty pounds of water in twenty-four hours. It may be used in connection with hydragogue cathartics, or even alone, in dropsies, and is useful in uric acid and phosphatic lithiasis.**Haircap Moss**, compressed in ounce packages for retailing purposes.**PREPARATION.****Fluid Extract Haircap Moss.**—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

HARDHACK. *Spiræa tomentosa, Linné.* (Leaves.)*Synonyms.*—Meadow-sweet, White-leaf, Steeple-bush.*Properties.*—Astringent and tonic. An excellent remedy in summer complaint of children, being less offensive to the stomach than other astringent remedies. Well adapted to disordered digestion, and used in diarrhoea and dysentery. The fluid extract is useful as an astringent lotion.**Hardhack leaves**, compressed in ounce packages for retailing purposes.**PREPARATION.****Fluid Extract Hardhack.**—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

*Dujardin-Beaumetz, "Clinical Therapeutics," p. 338.

HELIANTHELLA TENUIFOLIA, *Torrey & Gray*. (Root.)

Properties.—An aromatic expectorant, anti-spasmodic, diuretic, diaphoretic and in large doses emetic. Indicated in chronic bronchitis, asthma, as an expectorant in consumption, and in dropsy. Advantageously added to syrupy cough mixtures.

PREPARATION.

Fluid Extract Helianthella.—Not miscible with water without some precipitation.
Dose, 5 to 30 minims (0.3 to 2 C.c.).

HELONIAS. *Chamælrirum carolinianum, Willdenow*. (Rhizome.)

NOTE.—The names Helonias and Stargrass are now used for False Unicorn and Unicorn respectively as being liable to less confusion than the two latter.

Synonyms.—False Unicorn, Starwort, Blazing Star; Helonias dioica, Pursh.

Properties.—Tonic and diuretic. In large doses, emetic. Has been found to be beneficial in dyspepsia, loss of appetite, and in atonicity of the generative organs. Used with success in spermatorrhœa caused by excesses, accompanied by nocturnal emissions, gastric derangement, impaired memory, mental apathy, etc. Used as a uterine tonic in leucorrhœa, amenorrhœa and dysmenorrhœa.

Helonias root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Helonias, 120 grs.

Helonias Compound.

Helonias, 15 grs.

Squaw Vine, 60 grs.

Cramp Bark, 15 grs.

Blue Cohosh, 15 grs.

Extracts.

Fluid Extract Helonias (False Unicorn).—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Helonias Compound (or Squaw Vine Comp.).—Not miscible with water.

Fourteen fluidounces, imperial, represent: Squaw Vine, 8 ozs.; False Unicorn, Cramp-bark, and Blue Cohosh, 2 ozs. each.

The several remedies composing it possess tonic, alterative and anti-spasmodic virtues, and act particularly upon the female organs of reproduction. Said to render parturition "safe and easy."

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Helonin, a so-called **Concentration**,* a powdered preparation from Helonias root.

Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Pill Helonin, 1-8 gr.

HEMLOCK SPRUCE. *Tsuga canadensis, Carrière*. (Bark.)

Synonyms.—Hemlock, *Abies canadensis, Michaux*.

Properties.—Astringent and tonic. Beneficial as an injection in leucorrhœa and prolapsus uteri. Administered internally and also in enema, it has been found useful in chronic diarrhœa due to simple relaxation. It is also of service as a local application in gangrene.

Hemlock bark, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Hemlock bark.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Glycerole Hemlock bark.—Each fluidounce represents 120 grs. of bk.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

*See note in Preface on Concentrations.

HENBANE. *Hyoscyamus*, U.S.P. *Hyoscyamus niger*, *Linné.* (Lvs.)

Synonyms.—Jusquiame noir, Fr.; Bilsenkraut, Ger.

Properties.—Poisonous. Deliriant narcotic, anodyne, anti-spasmodic and hypnotic. It is employed chiefly to relieve pain, and quiet irregular nervous action. Often prescribed for children who do not bear opiates well. Employed in asthma, whooping-cough, functional palpitation of the heart, chorea, hypochondriasis, mania, etc. Like belladonna, it dilates the pupil, and occasionally produces a cutaneous rash. Employed as a sedative when opium, from its constipating properties, is not advisable.

Antidotes.—Same as for belladonna. Emetics (sulphate of zinc or copper), warmth to extremities, cold to the head. Stimulants, particularly coffee and morphia hypodermically in small and repeated doses.

Note.—Caustic alkalies decompose the active principles of Henbane.

Henbane leaves, U. S. P., compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLES.

Hyoscyne Hydrobromate.—A recent study of hyoscyne by Prof. H. C. Wood, shows that it is to this alkaloid, rather than to hyoscyamine, that the characteristic effects of henbane as a hypnotic and calmative agent are due. It is a powerful sedative, particularly useful in quieting maniacal excitement. It is also useful, in minute doses, as an anti-spasmodic, in asthma, severe colic, pertussis, etc.

Dose, 1-400 to 1-100 grain (0.00016 to 0.00065 Gm.).

Hyoscyamine.—Crystals. Produces the peculiar effects of henbane. Is used especially in quieting excitement in insane patients.

Dose, 1-100 grain (0.00065 Gm.), cautiously increased.

PREPARATIONS AND FORMULÆ.**Abstract Henbane, U. S. P.**

Normal Liquid or Fluid Extract Henbane, P., D. & Co.'s..... 1 fl. oz.

Sugar of milk, q. s. to make..... 228 grs.

Evaporate the fluid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well-corked bottle.

Dose, 2 to 5 grains (0.13 to 0.3 Gm.). One part equals 2 of the drug.

Extracts.

Fluid Extract Henbane, U.S.P.—Not miscible with water without some precipitation.

Dose, 5 to 10 minims (0.3 to 0.6 C.c.), gradually increased.

Powdered Extract Henbane.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Henbane.

Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Solid Extract Henbane, U.S.P., assayed.—One part equals 5 of leaves.

Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Hyonigrin, a so-called **Concentration**,* a powdered preparation from the leaves. *Synonym.*—Hyoscyamin.

Dose, 1-16 to 1-2 grain (0.004 to 0.032 Gm.).

Hypodermic Tablets.

Hyoscyne Hydrobromate, 1-100 gr.

Hyoscyamine, 1-50 gr.

Normal Liquid† Henbane.

Dose, 5 to 10 minims (0.3 to 0.6 C.c.), gradually increased.

Pills.

Camphor and Henbane, 2 grs.

Camphor, 1 gr.

Ext. Henbane, 1 gr.

Camphor, Opium and Henbane, 2 1-2 grs.

Camphor, 1 gr.

Ext. Henbane, 1 gr.

Powd. Opium, 1-2 gr.

* See note in Preface on Concentrations.

† See note in Preface on Normal Liquids.

HENBANE—Continued (Pills).

- Camphor, Valerian and Henbane**, 2 grs.
 Camphor, 1 gr. Ext. Henbane, 1-2 gr.
 Powd. Valerian, 1-2 gr.
- Colocynth Comp. Ext. and Henbane**, 5 grs.
 Ext. Coloc. co., 3 1-3 grs. Ext. Henbane, 1 2-3 gr.
- Colocynth Co., Henbane and Blue Mass**, 5 grs.
 Ext. Coloc. co., 3 grs. Blue Mass, 1 gr.
 Ext. Henbane, 1 gr.
- Coloc. Co., Henbane and Podoph.**, Formula "A.", 4 1-4 gr.
 Ext. Coloc. co., 3 grs. Podophyllin, 1-4 gr.
 Ext. Henbane, 1 gr.
- Henbane Extract**, 1-8, 1-4, 1-2 and 1 gr.
- Hyoscine Hydrobromate**, 1-400 gr.
- Hyoscyamine**, pure, 1-200 gr.
- Irisin Compound and Henbane**, 1 gr.
 Irisin, 1-4 gr. Podophyllin, 1-10 gr.
 Strychnine, 1-100 gr. Ext. Henbane, 1-2 gr.
- Phosphorus, Digitalis and Henbane**, 3 grs.
 Phosphorus, 1-33 gr. Ext. Henbane, 2 grs.
 Powd. Digitalis, 1 gr.

The above represent the principal pills containing Henbane; for others see Pill Formulæ list appended for numbers 28, 33, 35, 36, 43, 44, 46, 63, 64, 82, 100-103, 113, 134, 138, 205, 255, 279, 314, 355, 388, 406, 446, 448, and 460.

Tablet Triturates, Hyoscyamine (crystals), 1-200 and 1-100 gr.

Tinctures.**Henbane, B. P.**

- Fluid Extract Henbane, P., D. & Co.'s.....1 fl. oz.
 Dilute Alcohol.....7 fl. ozs.
 Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Henbane, U. S. P.

- Fluid Extract Henbane, P., D. & Co.'s.....2 fl. ozs.
 Stronger Alcohol.....8 fl. ozs.
 Water.....4 fl. ozs.
 Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

"German Tincture" Henbane.—An imported unofficial preparation from the green leaves.
 Dose, 8 to 20 minims (0.5 to 1.25 C.c.).

HOANG NAN. *Strychnos Gauthieriana, Pierre.* (Bark.)

Properties.—Poisonous. An alterative and spicant. It has the reputation of possessing specific virtues in leprosy, while the eruptions of constitutional syphilis, eczema and various skin diseases have yielded to it with promptness. "The most important use to which it is applied is in the treatment of paralysis. In relaxation of the members in hemiplegia, and where paralysis is recent, this remedy is peculiarly efficacious. In rheumatismal, paludal or syphilitic paralysis it achieves important results."*
Antidotes.—Emetics or stomach-pump, followed by perfect quiet. Treat with strychnine antagonists, chloral, tobacco, chloroform inhalations or ether, etc.

Hoang Nan, powdered bark, for retailing purposes.

PREPARATIONS.

Pill, Hoang Nan, 3 grs.

Tincture Hoang Nan, concentrated.—Not miscible with water.
 Four parts of tincture equal one of bark.
 Dose, 3 to 10 minims (0.2 to 0.65 C.c.).

HOLLYHOCK. *Althæa rosea, Cavanilles.* (Flowers.)

Properties.—Exclusively those of a demulcent. Used in irritation and inflammation of the mucous membranes.

Dose, wineglassful (60 C.c.) of an ounce to the pint infusion.

Hollyhock flowers, compressed in ounce packages for retailing purposes.

*Morse, "New Therapeutical Agents," p. 139.

HOPS. { Humulus, U.S.P. Humulus Lupulus, *Linné.* (Strobiles,
 } Lupulinum, U.S.P. (glandular powder from the strobiles.)

Properties.—Tonic, hypnotic, and said to possess anthelmintic, anti-lithic, anti-periodic and diuretic properties. Hops are principally used for their sedative or hypnotic action, producing sleep and removing restlessness, and are prescribed in nervous tremors, wakefulness, and in the delirium of drunkards. Have been found useful as an antaphrodisiac in the treatment of gonorrhœa, spermatorrhœa and other irritated conditions of the genito-urinary apparatus.

NOTE.—Lupulin preparations are the more active.

Hops, U. S. P., compressed in 1 oz., 4 oz. and 8 oz. packages for retailing purposes.

ACTIVE PRINCIPLE.

Lupulin, U. S. P.—Being convinced that no preparation can be prepared which shall contain in the form of a permanent powder a larger proportion of the oleo-resinous constituents to which hops chiefly owe their activity, we have omitted from our list the "concentration" humulin, and supply in its place a select article of lupulin, of a recent crop.
 Dose, 5 to 10 grains (0.3 to 0.6 Gm.).

PREPARATIONS AND FORMULA.

Elixirs.—Each fluidounce represents ingredients specified

Hops, 60 grs.

Lupulin, 16 grs.

Extracts.

Fluid Extract Hops.—Not miscible with water.

Dose, 15 to 60 minims (1 to 4 C.c.).

Fluid Extract Lupulin, U. S. P.—Not miscible with water.

Dose, 10 to 15 minims (0.6 to 1 C.c.).

Fluid Extract Lupulin Compound.—Not miscible with water.

Each pint represents: Scullcap, Lettuce and Lupulin, 5 1-2 ozs. each. Largely used as a therapeutic substitute for opium, and as a corrective of the opium habit. It is tonic, anti-spasmodic, nerve and hypnotic, and though possessing the most important properties of opium, leaves none of the objectionable sequelæ, *i. e.*, nervous exhaustion, disordered stomach and constipation. May be used to advantage in chorea, delirium tremens, nervous irritability, wakefulness, neuralgia, teething of children, etc.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Powdered Extract Hops.—Of the same strength as the solid ext.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Solid Extract Hops.—One part equals 5 of hops.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Malt Extract with Hops.—Each fluidounce represents 30 grs. hops

Pills.

Blennorrhagic, 3 1-8 grs.

Terebinth. alb., 1 1-2 gr.

Ext. Humuli, 3-4 gr.

Camphor monom., 3-4 gr.

Podophyllin, 1-8 gr.

Lupulin, 3 grs.

Phosphorus and Zinc Compound, 2 grs.

Phosphorus, 1-50 gr.

Lupulin, 1 gr.

Zinc sulph., 1 gr.

Tincture Hops, U. S. P.

Fluid Extract Hops, P., D. & Co.'s.....4 fl. ozs.

Dilute Alcohol.....17 fl. ozs.

Dose, 1 to 4 fluidrachms (1 to 15 C.c.).

HOREHOUND. Marrubium, U. S. P. Marrubium vulgare, *Linné.*
 (Leaves.)

Properties.—A bitter tonic, laxative when given in large doses, and said to increase the secretions from the skin and kidneys. It is employed in catarrh and other chronic affections of the lungs attended with cough and copious expectoration.

HOREHOUND—Continued.

Horehound leaves, U.S.P., compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Horehound.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Horehound Comp.—Not miscible with water without some precipitation.

Each pint represents: Horehound, 8 ozs.; Comfrey, 4 ozs.; Seneka and Wild Cherry, 2 ozs. each. A valuable combination in coughs and colds. Its mild diaphoretic and sedative properties particularly fit it for administration to children.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Horehound.—One part equals 5 of the drug.

Dose, 5 to 15 grains (0.3 to 1 Gm.).

Syrup Horehound Compound.—Each fluidounce represents: Elecampane, Spikenard, Comfrey, Wild Cherry bark and Horehound, 20 grs. each; Blood Root, 10 grs. This may be extemporized by employing the fluid extracts of the drugs named, reading minims for grains, and adding simple syrup to make one fluidounce.

Tincture Horehound.

Fluid Extract Horehound, P., D. & Co.'s.....1 fl. oz.
Dilute Alcohol.....4 fl. ozs.

Dose, 1-4 fluidounce (7.5 C.c.) and upwards.

Too weak a therapeutic preparation to be of service.

HORSE-CHESTNUT. *Æsculus Hippocastanum*, *Linné*. (Bark, fruit.)

Properties.—Horse-chestnut bark is tonic, astringent, febrifuge, narcotic and antiseptic. Given in doses of a teaspoonful from four to six times a day, it has effected cures in intermittent fever. This bark contains a principle isomeric with quinovine—a principle existing in cinchona leaves—and has been long used in Europe as a substitute for Peruvian bark.

The properties of the nuts are similar to those of the bark. They are tonic, slightly astringent, anti-spasmodic, and have been employed with especial benefit in neuralgic affections.

PREPARATIONS.

Fluid Extract Horse-chestnut bark.—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Fluid Extract Horse-chestnut nuts.—Not miscible with water without some precipitation.

Dose, 5 to 15 minims (0.3 to 1 C.c.).

HORSEMINT. *Monarda punctata*, *Linné*. (Leaves.)

Properties.—Stimulant, diaphoretic, diuretic and carminative. In popular use, in hot infusion, in affections arising from colds, such as catarrh, rheumatism, diarrhoea, etc., and also as an emmenagogue.

Dose, a wineglassful (60 C.c.) of a one ounce to the pint infusion.

Horsemint leaves, compressed in ounce packages for retailing purposes.

HORSE-RADISH. *Cochlearia Armoracia*, *Linné*. (Leaves, root.)

Synonyms.—Raifort sauvage, Moutarde des moines, Fr.; Merrettig, Ger.

Properties.—Stimulant, diuretic, anti-scorbutic and rubefacient. It stimulates the urinary secretions, and excites the flow of gastric juice in gastric debility. It has been efficaciously employed in chronic atony of the digestive apparatus and as a stimulant in general systemic debility and as an anti-scorbutic in scurvy. Rubefacient externally.

Horse-radish leaves, compressed in ounce packages for retailing purposes.

Horse-radish root, compressed in pound packages for retailing purposes.

HORSE-RADISH—Continued.**PREPARATION AND FORMULA.**

Fluid Extract Horse-radish root.—Miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Spiritus Armoracæ Comp., B. P.

Fluid Extract Horse-radish, P., D. & Co.'s.....	4 fl. ozs.
Fluid Extract Bitter Orange Peel, P., D. & Co.'s.....	4 fl. ozs.
Fluid Extract Nutmeg, P., D. & Co.'s.....	48 minims.
Dilute Alcohol.....	36 fl. ozs.
Distil 32 fluidounces. Dose, to 2 fluidrachms (4 to 8 C.c.).	

HOUND'S TONGUE. *Cynoglossum officinale*, *Linné*. (Leaves.)

Properties.—Hound's tongue is said to be anodyne, demulcent and astringent.

It is recommended as valuable in the treatment of coughs, catarrhs, hemoptysis and dysentery. The fluid extract, diluted with its volume of water, applied on cloths to scrofulous tumors, burns, goitre, recent contusions or inflammations, excoriated feet, and irritated, chafed or bruised parts, will afford relief, sometimes permanently.

Hound's Tongue leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Hound's Tongue.—Miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

HYDRANGÆA ARBORESCENS, *Linné*. (Root.)

Synonyms.—Seven Barks, Wild Hydrangæa.

Properties.—Cathartic, diuretic, diaphoretic and a solvent of gravelly deposits in the bladder. Alleviates the pain attendant on the passage of calculi through the ureter. It is especially useful when there is an alkaline condition of the urine, occasioning a tendency to phosphatic deposits. It is useful "in biliary calculi causing jaundice and other hepatic disorders its free elimination constitutes it curative of cutaneous diseases, and as valuable in all strumous disorders."*

Hydrangæa root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Hydrangæa.—Not miscible with water without some precipitation.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

HYSSOP. *Hyssopus officinalis*, *Linné*. (Leaves.)

Synonyms.—Hysope, Fr.; Ysop, Ger.

Properties.—Stimulant, aromatic, carminative and tonic. Principally used in quinsy and other sore throats as a gargle, combined with sage and alum and sweetened with honey. Also recommended in atonic and flatulent dyspepsia, chronic bronchitis—especially in old persons, in amenorrhœa, etc. It may also be applied externally for the relief of muscular rheumatism, as well as of bruises, ecchymoses, etc.

Hyssop leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Hyssop.—Not miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

IGNATIA, U. S. P. *Strychnos Ignatii*, *Lindley*. (Seeds.)

Synonyms.—Ignatius bean; Fève ignasurique, Fr.; Ignazbohne, Ger.

Properties.—Poisonous. Similar to those of *Nux vomica*, and depend on the

* Morse, "New Therapeutical Agents," p. 165.

IGNATIA—Continued.

presence of the same alkaloids, strychnine and brucine (see *nux vomica*).
A powerful tonic, indicated especially where there is impairment of nerve function—hence useful in all forms of general or local paralysis, in neuralgia, impotence, neurasthenia, chronic constipation, etc.

ANTIDOTES.—The same as in poisoning by strychnine. Emetics or stomach pump, followed by perfect quiet. Chloral, tobacco, ether, and chloroform (inhaled) are its true physiological antagonists.

NOTE.—Bromides, chlorides and iodides precipitate strychnine from solutions.

PREPARATIONS AND FORMULÆ.**Abstract Ignatia, U. S. P.**

Fluid Extract Ignatia, P., D. & Co.'s.....1 fl. oz.
Sugar of milk, q. s. to make.....228 grs.
Evaporate the fluid extract nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well-corked bottle.

Dose, 1-2 to 5 grains (0.03 to 0.3 Gm.). One part equals 2 of drug.

Extracts.

Fluid Extract Ignatia.—Not miscible with water without some precipitation.

Dose, 1 to 10 minims (0.06 to 0.6 C.c.), cautiously increased.

Powdered Extract Ignatia, assayed.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered sugar of milk.

Dose, 1-6 to 1 grain (0.01 to 0.06 Gm.).

Solid Extract Ignatia, assayed.—One part representing 5 of the bean; contains 10 per cent. alkaloids, principally strychnine.

Dose, 1-6 to 1 grain (0.01 to 0.06 Gm.).

Pills.

Bismuth and Ignatia, 41-4 grs.

Bismuth subnitrate, 4 grs. Ext. Ignatia bean, 1-4 gr.

Dyspepsia, 2 grs.

Ext. Ignatia bean, 1-4 gr. Ext. Rhubarb, 1-4 gr.
Ext. Cinchona bark, 1 gr. Powd. Capsicum, 1-2 gr.

Ignatia Amara Extract, 1-4, 1-2 and 1 gr.

Neuralgic Iodipathic, Brown-Sequard's, 3 1-2 grs.

Ext. Henbane, 2-3 gr. Ext. Conium seed, 2-3 gr.
Ext. Ignatia, 1-2 gr. Ext. Opium, 1-2 gr.
Ext. Aconite leaves, 1-3 gr. Ext. Cannabis Indica, 1-4 gr.
Ext. Stramonium seed, 1-5 gr. Ext. Belladonna, 1-6 gr.

Tablet Triturates.

Ignatia, powdered, 1-20 and 1-10 gr.

Iron and Arsenic Compound.

Iron, reduced, 1 gr. Powd. Ignatia, 1-40 gr.
Arsenious acid, 1-100.

Tincture Ignatia, U. S. P.

Fluid Extract Ignatia, P., D. & Co.'s.....1 fl. oz.

Alcohol.....10 fl. ozs.

Water.....3-4 fl. oz.

Dose, 10 to 60 minims (0.6 to 4 C.c.).

INDIAN BLACK-ROOT. *Pterocaulon pycnostachyum, Elliott.*

Properties.—This remedy is used largely among the Indians as an alternative.

It is also said to be cathartic, but its value as a therapeutic agent has not yet been scientifically defined.

PREPARATION.

Fluid Extract Indian Black-root.—Miscible with water

Dose, 15 to 20 minims (1. to 2. C.c.).

INDIAN CANNABIS. *Cannabis Indica*, U. S. P. *Cannabis sativa*, (var. *indica*) *Linné*. (Flowering tops.)

Synonyms.—Foreign Indian hemp; Hashish, Chanvre indien, Fr.; Indischer Hanf, Ger.

Properties.—A powerful, but notoriously uncertain narcotic, producing in full doses exhilaration, intoxication and delirious hallucinations. Recommended in delirium tremens, and in certain forms of insanity, in delirium after fever, softening of the brain and other conditions involving anæmia of the brain; in some cases of migraine; to allay the itching of eczema; to prevent the griping tendency of certain cathartics, etc.

ANTIDOTE.—An overdose produces symptoms which are alarming rather than dangerous, and are to be combated by faradization of the respiratory muscles, and by the cautious use of strychnine.

Indian Cannabis, U. S. P., compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLES.

Cannabin Tannate.—Glucoside. An efficient hypnotic, producing a quiet and refreshing sleep. It does not disturb the secretions as most narcotics do, nor does it derange digestion.
Dose, 1 to 5 grains (0.065 to 0.33 Gm.).

Cannabinon.—A preparation of Cannabis with properties apparently identical with those of the ordinary extract, and apparently equally uncertain in its action.
Dose, 1-2 to 1 1-2 grains (0.03 to 0.1 Gm.).

PREPARATIONS AND FORMULÆ.

Chlor-anodyne.—Each gramme (ordinary adult dose) contains:

Tr. Capsicum.....	.0025	Morph. muriate.....	.0060
Acid Hydrocyanic, dil....	.0170	Tr. Indian Cannabis.....	.0800
Alcohol.....	.3000	Chloroform.....	.1350
Glycerin.....	.4570	Oil Peppermint.....	.0025

An elegant and efficient remedy for colic, cholera morbus, neuralgia, spasmodic pains, etc. Concentrated in form and rapid in action—of great value where anti-spasmodics or anodynes are indicated.

Extracts.

Fluid Extract Indian Cannabis (*Cannabis Indica*), U. S. P.—Not miscible with water.

Dose, 3 to 10 minims (0.2 to 0.6 C.c.), beginning cautiously.

Powdered Extract Indian Cannabis (*Cannabis Indica*).

NOTE.—Powd. Ext. Indian Cannabis as found in the market is not equivalent, grain for grain, to the ordinary solid extract. Physicians complain that they do not obtain from it the effects of the drug. This is no doubt due to the fact that chemical agents are employed to solidify the resin. We prefer to simply dilute the solid extract, and so avoid the injurious influence of excessive heating, and of the addition of chemicals. This saccharated extract we supply is one-fourth the strength of the ordinary extract.
Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Solid Extract Indian Cannabis (*Cannabis Ind.*), U.S.P.

One part represents 9 of crude drug.

Dose, 1-8 to 1 grain (0.008 to 0.065 Gm.).

Normal Liquid* Indian Cannabis.

Dose, 3 to 10 minims (0.2 to 0.6 C. c.), beginning cautiously.

Pills.

Indian Cannabis (*Cannabis Ind.*) Extract, 1-4, 1-2 and 1 gr.

Ergotin and Indian Cannabis, 1 1-2 gr.

Ergotin, purified, 1 gr. Ext. Indian Cannabis, 1-2 gr.

Neuralgic Idiopathic, Brown-Sequard's, 3 1-2 grs.

Ext. Henbane, 2-3 gr. Ext. Conium seed, 2-3 gr.

Ext. Ignatia, 1-2 gr. Ext. Opium, 1-2 gr.

Ext. Aconite leaves, 1-3 gr. Ext. Indian Cannabis, 1-4 gr

Ext. Stramonium seed, 1-5 gr. Ext. Belladonna, 1-6 gr.

Phosphorus and Indian Cannabis, 1-4 gr.

Phosphorus, 1-50 gr. Ext. Indian Cannabis, 1-4 gr.

Phosphorus and Indian Cannabis, Formula "B.", 1-4 gr.

Phosphorus, 1-25 gr. Ext. Indian Cannabis, 1-4 gr.

* See note in Preface on Normal Liquids.

INDIAN CANNABIS—Continued (Pills).**Sedative, 2 grs.**

Ext. Musk root, 1-2 gr.

Ext. Henbane, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Indian Cannabis, 1-10 gr.

Sedative, Mann's, 2 1-2 grs.

Ext. Coca, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Indian Cannabis, 1-4 gr.

Vallet's Mass, 1 gr.

Arsenious acid, 1-80 gr.

Strychnine, 1-80 gr.

Codeine, 1-8 gr.

Strychnine Compound, Formula "B.", 2 grs.

Strychnine, 1-100 gr.

Phosphorus, 1-100 gr.

Ext. Indian Cannabis, 1-16 gr.

Iron proto-carb. (Vallet's), 1 gr.

Powd. Ginseng, 1 gr.

Tablet Triturates, Indian Cannabis Ext., 1-20 and 1-10 gr.**Tincture Indian Cannabis (Cannabis Indica), U. S. P.**

Fluid Extract Indian Cannabis, P., D. & Co.'s..... 4 fl. ozs.

Alcohol.....

Dose, 10 to 60 minims (0.6 to 4 C.c.)..... 19 fl. ozs.

INDIAN PHYSIC. *Gillenia trifoliata, Manch.* (Bark of rhizome.)*Synonym.*—American Ipecac.*Properties.*—A mild and efficient emetic and cathartic, said to possess tonic properties. Used by some practitioners as a substitute for Ipecac, which it is said to resemble in action.**Indian Physic**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Indian Physic.**—Not miscible with water.

Dose, 10 to 40 minims (0.6 to 2.5 C.c.).

INDIAN TURNIP. *Arisæma triphyllum, Torrey.* (Corm.)*Synonyms.*—Arum, Dragon Root, Wake Robin.*Properties.*—Acrid, expectorant and diaphoretic. Recommended in croup, whooping-cough, asthma, chronic laryngitis, bronchitis; to relieve flatulence and colic, and as an application to aphthous sores of the mouth.**Indian Turnip root**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Indian Turnip.**—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

IPECAC. *Ipecacuanha, U. S. P. Cephælis Ipecacuanha, A. Richard.* (Root.)*Properties.*—Emetic in large doses. In small doses expectorant and diaphoretic; and, in minute doses, tonic and stimulant, increasing the appetite and promoting digestion. In dysentery it exercises peculiar powers. As a diaphoretic, combined with opium, it is applicable to all cases not attended with much fever, cerebral diseases or nausea in which there is an indication for diaphoresis. Narcotics hinder its emetic action.**ACTIVE PRINCIPLE.****Emetine.**—Alkaloid. Expectorant, nauseant, emetic. Experiments show that emetine possesses in a remarkable degree the power of exciting the mucous secretion in the air passages. Hence its great value as an expectorant. The alkaloid has precisely the same therapeutic uses as the drug itself, but is sometimes to be preferred from the smallness of the dose, and from its uniformity of action.

Dose, as an expectorant, 1 100 to 1-25 grain (0.006 to 0.003 Gm.).

IPECAC—Continued.

PREPARATIONS AND FORMULÆ.

Extracts.**Fluid Extract Black Cohosh Comp.**—Not miscible with water.

Each pint represents: Black Cohosh, 8 ozs.; Licorice, 3 ozs.; Cherry Bark, 2 ozs.; Seneka, 2 ozs.; Ipecac, 1 oz.

Nervous and arterial sedative; in larger doses acting as a diaphoretic and causing increased secretion from the bronchial mucous membrane. Useful in chronic rheumatism and atonic amenorrhœa.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Dover's Powder.—See Tinct. Ipecac and Opium below.**Fluid Extract Ipecac, U. S. P.**—Miscible with water.

Dose, as an emetic, 25 minims (1 6 C.c.) repeated if necessary; as a diaphoretic, 1 to 3 minims (0.06 to 0.18 C.c.); as an expectorant, 1-4 to 1 minim (0.015 to 0.06 C.c.).

Fluid Extract Ipecac and Senega.—Miscible with water.

Each pint represents: Ipecac, 8 ozs.; Seneka, 8 ozs. Emetic in large doses. In small doses, a stimulating expectorant and diaphoretic.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

Powdered Extract Ipecac, assayed.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Ipecac.

Dose, 1-20 to 1-2 grain (0.003 to 0.03 Gm.).

Solid Extract Ipecac, assayed.—One part equals 5 of root.

Dose, 1-20 to 1-2 grain (0.003 to 0.03 Gm.).

Lozenges.**Ipecac, U. S. P.,** 1-4 gr.**Ipecac and Morphine.**

Powd. Ipecac, 1-8 gr.

Morphine sulphate, 1-32 gr.

Pectoral, Dr. Jackson's.

Ipecac, 1-20 gr

Morphine muriate, 1-33 gr.

Antimony, sulphurated, 1-40 gr.

Normal Liquid* Ipecac, assayed.—Standard, 1.5 per cent. Emetine.

Dose, as an emetic, 25 minims (1.6 C.c.), repeated if necessary; as a diaphoretic, 1 to 3 minims (0.06 to 0.18 C.c.); as an expectorant, 1-4 to 1 minim (0.015 to 0.06 C.c.).

Pills.**Alternative,** 1 1-4 grs.

Blue Mass, 1 gr.

Powd. Opium, 1-8 gr.

Powd. Ipecac, 1-8 gr.

Diaphoretic, 1 1-2 gr.

Morphine acetate, 1-25 gr.

Powd. Potass. nitrate, 1 gr.

Powd. Ipecac, 1-4 gr.

Powd. Camphor, 1-4 gr.

Dover's Powder, 1-4 gr. See also Ipecac and Opium.**Dysentery,** 2 1-6 grs.

Blue Mass, 1 gr.

Powd. Ipecac, 1 gr.

Gelsemperin, 1-6 gr.

Heim's, 2 grs.

Quinine sulphate, 1 gr.

Powd. Ipecac, 1-4 gr.

Powd. Digitalis, 1-2 gr.

Powd. Opium, 1-4 gr.

Ipecac, 1-50 gr.**Ipecac Extract,** 1-4 gr.**Ipecac and Opium,** 1-2, 1 and 2 grs.

Containing of these ingredients 1-4, 1-2 and 1 gr. each respectively.

Lapactic, 1-2 gr.

Aloin, 1-4 gr.

Ext. Belladonna, 1-8 gr.

Strychnine, 1-60 gr.

Ipecac, 1-16 gr.

Liver, Christopher's, 3 1-2 grs.

Calomel, 2 grs.

Powd. Ipecac, 1-2 gr.

Rhubarb, 1 gr.

Quinine, Ipecac and Opium, 2 1-2 grs.

Quinine hydrobromate, 2 gr.

Powd. Opium, 1-4 gr.

Powd. Ipecac, 1-4 gr.

* See note in Preface on Normal Liquids.

IPECAC—Continued (Pills).

Tonic Hæmatic, Andrews', 2 3-4 grs.
 Quinine sulph., 1 gr. Powd. Ipecac, 1-8 gr.
 Iron by hydrogen, 1 1-2 gr. Strychnine sulph., 1-40 gr.
 Arsenious acid, 1-40 gr.

For others containing Ipecac see Pill Formulæ list appended for numbers 35, 37, 45, 56, 73, 94, 133, 139, 143, 161, 255, 262, 332, 333, 337 and 426.

Syrups.

Dover's Powder.—Each fluidounce representing 40 grs. Dover's Powder, U. S. P., or '56 gr. Morphine alkaloid equivalent to 3-4 gr. Morphine sulph.
 Dose, for adults, 1 to 2 fluidrachms (4 to 8 C. c.).

Ipecac, U. S. P.—Each fluidounce represents 30 minims fluid extract; made extemporaneously as follows:

Fluid Extract Ipecac, P., D. & Co.'s..... 2 fl. ozs.
 Syrup 29 fl. ozs.

Dose, as an emetic for a young child, 30 to 60 minims (2 to 4 C. c.) repeated; as an expectorant for adults, to 60 minims (2 to 4 C. c.); as an expectorant for a child, 1 to 10 minims (0.07 to 0.7 C. c.).

Potassium Citrate and Ipecac.—Formula suggested by Prof. H. C. Wood, in the *Therapeutic Gazette*, Jan. 15, 1885. Each fluidounce represents: Potassium citrate, 80 grs.; Lemon juice, 160 minims; Syrup Ipecac, 40 minims.

Dose, for an adult, a tablespoonful (15 C. c.) 4 to 6 times a day.

Prof. Wood, as the result of long experience, has settled on this combination of potassium citrate with ipecac as the best sedative cough mixture he has ever used. If there is much cough Dr. Wood is accustomed to add to the mixture an opiate, and such an addition can easily be made whenever the case seems to demand it. The mixture is intended to be used only during the first or dry stage of the "cold." After the first two or three days the more stimulating expectorants come into play, and here Prof. Wood finds the oil of eucalyptus in capsules the most serviceable remedy. See Eucalyptus: capsules.

Tablet Triturates, Ipecac Powdered, 1-10, 1-4 and 1-2 gr.

Tincture Ipecac and Opium, U. S. P., assayed.—A liquid form of Dover's Powder, representing 1 grain in each minim.

Dose, 2 to 12 minims (0.13 to 0.75 C. c.).

This may be made extemporaneously as follows:

Fluid Extract Ipecac, P., D. & Co.'s..... 1 fl. oz.

Fluid Opium, aqueous, P., D. & Co.'s..... 10 fl. ozs.

Reduce the Fluid Opium to 8 1-2 fluidounces by careful evaporation, add the fluid extract Ipecac. Filter and add dilute alcohol through filter to make 10 fluidounces.

Wine Ipecac, U. S. P.—Each fluidounce representing 33 grains Ipecac; made extemporaneously as follows.

Fluid Extract Ipecac, P., D. & Co.'s..... 1 fl. oz.

Sherry or native wine..... 12 1-2 fl. ozs.

Alcohol..... 1 fl. oz.

Dose, 15 to 60 minims (1 to 4 C. c.).

IRON-WOOD. *Ostrya virginica*, Willdenow. (The heart-wood.)

Synonyms.—Hop Hornbeam, Lever wood; Hopfenheimbuche, Ger.

Properties.—Tonic, anti-periodic, alterative. Prescribed in intermittent fevers, neuralgic affections, dyspepsia, scrofula and wherever an anti-periodic tonic is indicated.

PREPARATION.

Fluid Extract Iron-wood.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

JABORANDI. *Pilocarpus*, U.S.P. *Pilocarpus pinnatifolius*, Lemaire. (Leaves.)

Properties.—A powerful diaphoretic and sialagogue, having also the power of increasing the bronchial, lachrymal and nasal secretions. It has been used with advantage in diabetes insipidus, uræmia and renal dropsies, night-sweats of phthisis, as a galactagogue, etc. It is apt to produce

JABORANDI—Continued.

nausea on an empty stomach. Decreases the temperature. Resembles atropine in its action on the pulse, flushing the face, and exerting a more decided influence on adults than children; but is diametrically opposed to it in its action on the salivary, sudoral and mammary secretions, on the pupil, on the minute arteries, and in delirium. The sweating may be prevented or checked by the subcutaneous injection of 1-100 of a grain of atropine.

NOTE.—Atropine and pilocarpine are true physiological antagonists.

ACTIVE PRINCIPLE.

Pilocarpine Hydrochlorate or Muriate.—Salt of alkaloid. Has the same action as the drug itself, increasing greatly the salivary secretions and producing profuse perspiration. Applied to the eye it causes contraction of the pupil, but not to the same extent as physostigmine. Used especially as a powerful eliminant in uræmia, ascites, etc., also as a physiological antidote to atropine.

Dose, 1-16 to 1-4 grain (0.004 to 0.016 Gm.).

Pilocarpine Nitrate.—Salt of alkaloid. The muriate of pilocarpine has been more commonly prescribed. The nitrate having the same therapeutic properties is to be preferred because it is not, like the muriate, a hygroscopic salt.

Dose, 1-16 to 1-4 grain (0.004 to 0.016 Gm.).

PREPARATIONS.**Extracts.**

Fluid Extract Jaborandi, U. S. P.—Not miscible with water without some precipitation.

Dose, 20 to 60 minims (1.3 to 4 C.c.), to be taken in warm water.

Powdered Extract Jaborandi.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered jaborandi.

Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Solid Extract Jaborandi.—One part equals 5 of leaves.

Dose, 3 to 10 grains (0.2 to 0.6 Gm.).

Hypodermic Tablets.

Pilocarpine Hydrochlorate, 1-3 gr.

Pilocarpine Nitrate, 1-3 gr.

Pills.

Jaborandi Extract, 3 grs.

Pilocarpine Hydrochlorate, 1-16 gr.

Pilocarpine Nitrate, 1-16 gr.

Tablet Triturates Pilocarpine Nitrate, 1-100, 1-50, 1-25 and 1-10 gr.

JALAP. Jalapa, U.S.P. *Ipomœa purga, Wenderoth.* (Tuber.)

Synonym.—*Exogonium purga, Benth.*

Properties.—A drastic hydragogue cathartic, operating quickly, and sometimes painfully, producing copious watery discharges. Its action is modified by combination with other cathartics. United with bitartrate of potassium its hydragogue properties are much increased, and this combination is often prescribed in dropsy. About ten grains of bitartrate of potassium may be added to the minimum dose of the fluid extract.

ACTIVE PRINCIPLE.

Jalapin.—Resin Jalap, U. S. P. It is now generally believed that the resin is the sole purgative principle of Jalap. To obviate its occasional harshness, it has been advised to triturate it with other substances calculated to separate its particles.

Dose, 2 to 5 grains, (0.13 to 0.32 Gm.).

PREPARATIONS.**Abstract Jalap, U. S. P.**

Fluid Extract Jalap, P. D. & Co.'s..... 1 fl. oz.

Sugar of milk, q. s. to make..... 228 grains.

Evaporate the fluid extract nearly to dryness by a very gentle heat; add

JALAP—Continued.

175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the prod. Keep in a well-corked bottle.

Dose, 2 to 12 grains (0.13 to 0.8 Gm.). One part equals two of drug.

Extracts.

Fluid Extract Jalap.—Not miscible with water.

Dose: laxative, 2 to 10 minims (0.13 to 0.65 C.c.); cathartic, 10 to 25 minims (0.65 to 1.63 C.c.).

Fluid Extract Jalap and Senna.—Not miscible with water.

This combination of Senna with Jalap modifies the severity of the action of the latter.

* Each pint represents 8 ozs. each of Senna and Jalap.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Powdered Extract Jalap.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Jalap.

Dose, 2 to 5 grains (0.13 to 0.32 Gm.).

Solid Extract Jalap.—One part equals 5 of tuber.

Dose, 2 to 5 grains (0.13 to 0.32 Gm.).

Pills.

Ballou, 3 1-4 grs.

Ext. Coloc. comp., 1 gr.

Ext. Jalap, 1 gr.

Calomel, 1 gr.

Powd. Ipecac, 1-8 gr

Cathartic Compound, U.S.P., 3 1-2 grs.

Ext. Coloc. comp., 1 3-10 gr.

Ext. Jalap, 1 gr.

Calomel, 1 gr.

Powd. Gamboge, 1-4 gr.

Jalap Compound, 3 1-4 grs.

Powd. Jalap, 3-4 gr.

Powd. Rhubarb, 3-4 gr.

Oil Caraway, 1-8 gr.

Castile Soap, 3-4 gr.

Pil. Calomel comp., 1 1-8 gr.

Jalap Extract, 1 gr.

Jalapin, 1-16 and 1 gr.

For others containing jalap see appended Pill Formulæ list for numbers 64, 101—104, 260, 261, 263, 460 and 461.

JAMAICA DOGWOOD. *Piscidia Erythrina, Linné.* Bark of it)

Properties.—Anodyne, hypnotic. A valuable substitute for opium, producing natural refreshing sleep, without subsequent depressing effects, suppression of the secretions, constipation, headache, etc. As an analgesic it is inferior to opium, which, indeed, stands as yet unrivalled in this regard; but as a remedy for the sleeplessness caused by nervous exhaustion, or by minor constitutional disturbances, it is of great utility. Applied locally, the fluid extract relieves toothache, and taken internally, often allays pain, rendering unnecessary the administration of more powerful narcotics.

ACTIVE PRINCIPLE.

Piscidin.—Sedative, hypnotic, etc. Clinical experience has not yet shown how far this substance represents the activity of the crude drug or in what dose it should be given.

PREPARATIONS.**Extracts.**

Fluid Extract Jamaica Dogwood.—Not miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.)

Powdered Extract Jamaica Dogwood.—Of same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Jamaica Dogwood.

Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

Solid Extract Jamaica Dogwood.—One part equals 7 of bark.

Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

Pill Jamaica Dogwood Extract, 2 grs.

JAMBU ASSU. *Ottonia anisum, Sprengel.* (Root.)

Synonyms.—*Ottonia Jaborandi, Guillemín; Serronia Jaborandi, Gaudichaud-Beaupré.*

Properties.—Information regarding the therapeutic uses of this drug is as yet very meagre. Said to be employed in Brazil as a stimulant and febrifuge, particularly in low forms of fever. It is believed to have a specific action on the uterus, and is employed for the relief of painful affections of that organ, and is even believed to be capable of inducing abortions. The drug contains an alkaloidal principle, but its peculiar pungent properties reside in its oleoresin, and doubtless closely connected with its medicinal activity. We solicit from physicians reports of such clinical experience as they may possess, or be able to record.

PREPARATION.

Fluid Extract Jambu Assu.—Not miscible with water.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

JAMBUL. *Eugenia jambolana, Lamarck.* (Bark.)

Synonyms.—*Syzygium jambolanum, DeCandolle; Calyptanthus jambolana, Willdenow; Java plum.*

Properties.—This drug is a product of a tree well known in the East Indies for its edible fruit. All parts of the plant are astringent. The juice of the fresh bark and leaves is much used by the native physicians in diarrheas, especially of children, and the bark is employed in the preparation of astringent injections, gargles and lotions. The root and seeds also are used in a similar manner. Recently the discovery was made that both the bark and seeds possess the unique property of arresting the formation of sugar in diabetes, the seeds, however, being slightly more efficient. Clinical experiments conducted both in England and in America confirm this discovery, which is one of great importance. The drug may not prove a specific in all cases of diabetes, but in those at least depending on increased production of glucose, it promises to be of the greatest value. Experiment has demonstrated that it has a positive inhibitory influence upon the action of diastatic ferments. Thus, in two experiments conducted under the same conditions, a given amount of malt extract converted 22.4 grains of starch into sugar in absence of jambul, but only 6.3 grains in its presence.

Jambul bark, } in powdered form for convenience of prescribers and
Jambul seeds, } dispensers.

Dose, 5 to 10 grains (0.3 to 0.6 Gm.) two to three times a day.

JEQUIRITY. *Abrus precatorius, Linné.* (Seed)

Properties.—Jequirity, when applied to the eye in the form of an infusion, produces a peculiar form of inflammation, attended with much pain and a copious muco-purulent discharge. By using weak solutions a moderate action may be induced, which can be easily held under control, and which has been found to be curative in cases of pannus and trachoma of long standing.

Jequirity seeds, in ounce packages; also powdered, in ounce bottles.

PREPARATION

Fluid Jequirity.—Miscible with water 100 parts represent 12 of seed.
DIRECTIONS.—Dilute with three to five times its volume of pure water, and apply two drops once a day until the desired effect is produced. It is best to dilute only as it is required for use, as after dilution it quickly spoils. If the action is too severe employ hot water freely, and very dilute solutions of corrosive sublimate to control it.

JERUSALEM OAK. *Chenopodium Botrys*, *Linné*. (Leaves.)

Synonyms.—Wormseed, Feather Geranium, Jerusalem Oak; Anserine Anthelmintique, Fr.; Wurmwidriger, Gemeines Traubenkraut, Ger.

Properties.—Anthelmintic. Most useful in the case of round worms in the form of a decoction, one ounce to the pint of hot milk.

Dose, a wineglassful (60 C.c.).

Jerusalem Oak leaves, compressed in ounce packages for retailing purposes.

JEWEL-WEED. *Impatiens pallida*, *Nuttall*. (Leaves.)

Synonyms.—Touch-me-not, Wild Celandine; Wildes Schoellkraut, Ger.

Properties.—Alterative and diuretic. Useful in scrofula and chronic rheumatism in the form of an infusion, one ounce to the pint.

Dose, a wineglassful (60 C.c.).

Jewel-weed (Wild Celandine) leaves, compressed in ounce packages for retailing purposes.

JUDAS-TREE. *Cercis canadensis*, *Linné*. (Bark.)

Synonym.—Red Bud.

Properties.—Astringent. Said to be useful in the treatment of diarrhoea and dysentery, especially where it has become chronic, as it can be administered where there is an irritable condition of the stomach without increasing the trouble. It also makes an excellent injection for leucorrhœa and gleet where there is an atonic condition of the mucous membrane.

PREPARATION.

Fluid Extract Judas-tree.—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

JUNIPER. *Juniperus*, U.S.P. *Juniperus communis*, *Linné*. (Fruit.)

Properties.—Diuretic and gently stimulant, producing, when largely taken, disagreeable irritation of the urinary passages. Chiefly employed as an adjuvant to more powerful diuretics, in dropsical complaints. Is also used in scorbutic and cutaneous diseases, catarrh of the bladder, and atonic conditions of the alimentary canal and uterus, gonorrhœa, cystorrhœa, etc.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Buchu Compound.

Buchu, 60 grs.

Cubeb, 15 grs.

Juniper, 15 grs.

Spirits nit. ether, 15 minims.

Buchu, Juniper and Potassium Acetate.

Buchu, 45 grs.

Juniper, 12 grs.

Potassium acetate, 16 grs.

Buchu and Pareira Brava.

Buchu, 30 grains.

Pareira brava, 7 1-2 grs.

Juniper berries, 15 grs.

Stone root, 7 1-2 grs.

Extracts.

Fluid Extract Juniper berries.—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Solid Extract Juniper.—One part equals 3 of the berries.

Dose, 20 to 30 grains (1.3 to 2 Gm.).

Phl, Diuretic, 3 grs.

Powd. Soap, 1 1-2 gr.

Oil Juniper berries, 1-16 gr.

Sodium carb., 1 1-2 gr.

KAMALA. Kamala, U. S. P. *Mallotus philippinensis*, *Müller Arg.*
(Capsule glands.)

Synonym.—*Rottlera tinctoria*, *Roxburgh.*

Properties.—Actively purgative in full doses. It has long been used in India in the treatment of tape-worm. Dr. C. MacKinnon, of the British Army, reports that having used it in fifty cases, he had failed in expelling the worm but in two, and the testimony of other physicians is equally favorable. In case the first dose fails to operate on the bowels, it may be repeated in four hours, followed by a dose of castor oil. It requires no preparatory treatment. As an external remedy Kamala is used in various affections of the skin, particularly scabies and herpes.

PREPARATIONS.

Capsules Male-fern and Kamala.—In hard and soft gelatin
Oil Male-fern, 7 minims. Kamala, sifted, 4 grs.

Fluid Extract Kamala.—Not miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

KAVA KAVA. *Piper methysticum*, *Forster.* (Root.)

Synonyms.—Ava Ava, Kawa, Yakona.

Properties.—Used beneficially in gout, bronchitis, catarrhal affections, erysipelatous eruptions, and especially, according to Dr. Sanné, in spasmodic cystitis, orchitis, and acute gonorrhœa.* The action of Kava appears to vary with the amount taken. In small doses it acts as a stimulant and tonic, but when taken in large doses it produces an intoxication which differs from that caused by alcohol, in being of a silent and drowsy nature accompanied by incoherent dreams, the drinker not being quarrelsome or excited. As a remedy for gonorrhœa, two or three doses should be given during the day, each with a full goblet of water. It is said that twenty minutes after the first dose, a pressing desire to urinate is experienced. The quantity of urine is abundant, and it becomes as limpid and as clear almost as water. The pain that was present during the previous micturitions disappears, and a sensation of comfort is experienced in urinating. Kava, moreover, acts like a bitter tonic. It is pleasant to take, stimulates the appetite, does not derange the digestive functions, and produces neither diarrhœa, nor constipation.

ACTIVE PRINCIPLES.

Kawahin.—The crystalline principal of the drug. Its therapeutics is not yet established.

Alpha Resin Kava Kava.—Lewin believes is the active principle. It is a powerful local anæsthetic, but this use is restricted by the fact that it is insoluble in water. Dr. Sanné recommends it in doses of 1-2 grains, equal 15 grains of the root, eight to twelve times a day.

Beta Resin Kava Kava.—Regarded as somewhat inferior to the Alpha; its uses are essentially the same.

PREPARATIONS.

Extracts.

Fluid Extract Kava Kava.—Not miscible with water.

Owing to its local anæsthetic action Dr. Lewin recommends that it be employed to mask the taste of bitter substances, such as quinine.†
Dose, 20 to 60 minims (1.3 to 4 C.c.).

Solid Extract Kava Kava.—One part equals 10 of the root.
Dose, 2 to 6 grains (0.13 to 0.4 Gm.).

* Bulletin Générale de Thérapeutique, March 15; Therapeutic Gaz., June 15, 1886.

† Deutsche Medicinal Zeitung, February 1, 1886.

KINO, U. S. P. Inspissated juice from the trunk of *Pterocarpus Marsupium*, *Roxburgh*.

Properties.—Kino probably owes its virtues entirely to a peculiar kind of tannic acid. It is employed in various complaints on account of its energetic astringent properties. It has been recommended internally in menorrhagia, and as a topical application in leucorrhœa, relaxed sore throat, and aphthæ of the mouth or fauces. It is also sometimes used as a styptic to stop hemorrhage, especially that from the Schneiderian membrane. Used also to check diarrhœa, and in the treatment of chronic dysentery.

PREPARATIONS AND FORMULA.

Fluid Kino.—Miscible with water. Each pint represents 8 1-2 ozs. av. of the crude drug.
Dose, 10 to 50 minims (0.6 to 3.3 C.c.).

Lozenge, Alum and Kino.—One grain each of Alum and Kino.

Pill Camphor Compound.

Camphor, 1 gr.

Powd. Opium, 1 gr.

Powd. Kino, 1 gr.

Ext. Capsicum, 1-16 gr.

Tincture Kino, U. S. P.

Fluid Kino, P., D. & Co.'s.....5 fl. ozs.

Strong Alcohol.....20 fl. ozs.

Glycerin.....3 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

KOLA. *Cola acuminata*, *R. Brown*. (Seeds.)

Synonym.—*Sterculia acuminata*, *Palisot de Beauvais*.

Properties.—Highly prized by the natives of Africa for its stimulating and sustaining properties, which are analogous to those of the South American Coca. The active constituents of the nuts are caffeine and theobromine, together with a variety of tannic acid. Therapeutically, the Kola resembles in its action guarana and coca, and is said to have almost a specific effect in combatting the intoxicating influence of alcohol. It is employed also in diarrhœas, and in all depressed conditions of the nervous system where caffeine is indicated.

PREPARATION.

Fluid Extract Kola nuts.—Not miscible with water.

Dose, 10 to 30 minims (0.65 to 2 C.c.).

KOOSSO. *Brayera*, U. S. P. *Brayera anthelmintica*, *Kunth*. (Flowers and unripe fruit.)

Properties.—Tæniifuge; of all the remedies for tape-worm none is more efficient or certain. Its other effects when taken internally are not very striking. It appears to act as a poison to the worms, and has been found equally effectual against both kinds of tape-worm. The Koosso should be taken in the morning upon an empty stomach, and a previous evacuation of the bowels with castor oil or other cathartic is recommended. It is also recommended that lemon juice or tamarind water be freely taken before and after the Koosso. The alcohol may be in part removed from the extract by evaporating at a low temperature, when the dose is large.

PREPARATION.

Fluid Extract Koosso flowers.—Not miscible with water.

Dose, 2 to 8 fluidrachms (8 to 30 C.c.).

LABRADOR TEA. *Ledum latifolium*, *Aiton*. (Leaves.)

Properties.—Demulcent, expectorant and tonic. May be employed with benefit in the different varieties of bronchitis requiring combined tonic and expectorant treatment.

LABRADOR TEA—Continued.

Labrador Tea, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Labrador Tea.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

LACTUCARIUM, U. S. P. *Lactuca virosa, Linné.* (Concrete juice.)

Synonyms.—Lettuce Opium; Giftlattichsaft, Ger.

Properties.—Reputed hypnotic and anodyne. For infants and for susceptible patients it is employed as a substitute for opium, especially for nervous insomnia. It is almost exclusively prescribed in the form of the syrup.

PREPARATIONS AND FORMULA.

Fluid Extract Lactucarium, U.S.P.—Not miscible with water.

Dose, 4 to 16 minims (0.25 to 1 C.c.).

Pill, Syphilitic, Ricord's, Modified, 2 1-2 grs.

Mercury protiodide, 1-2 gr.

Ext. Opium, 1-10 gr.

Lactucarium, 1-2 gr.

Ext. Conium seed, 1 1-2 gr.

Syrup Lactucarium, U. S. P.

Fluid Extract Lactucarium, P., D. & Co.'s..... 1 fl. oz.

Syrup..... 15 fl. ozs.

Dose, 2 to 4 fluidrachms (8 to 15 C.c.); for an infant, 30 minims (2 C.c.).

LADIES' BED-STRAW. *Galium verum, Linné.* (Herb.)

Properties.—Refrigerant and diuretic. Used in many diseases of the urinary organs, as suppression of urine, calculous affections, inflammation of the kidneys and bladder, and in scalding of the urine in gonorrhoea. Contraindicated in diseases of a passive character, on account of its refrigerant and sedative effects, but may be used freely in fevers and all acute diseases. It has also been recommended in scorbutic and nervous diseases. It may be used internally and as a wash in cutaneous diseases—as freckles, lepra, psoriasis, lichen, etc., especially if those diseases are combined with a strumous diathesis.

PREPARATION.

Fluid Extract Ladies' Bed-straw.—Miscible with water without material precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

LADIES' SLIPPER. *Cypripedium, U.S.P.* *Cypripedium pubescens, Willdenow.* (Rhizome.)

Synonyms.—Moccasin Flower, American Valerian.

Properties.—Tonic, stimulant, diaphoretic and anti-spasmodic. It is said to be valuable in all cases of nervous excitability unconnected with organic lesions, allaying the irritability, lessening pain, producing a calm and cheerful condition of body and mind, and consequently favoring mental tranquility, or sleep. It has been used in hysteria, chorea, nervous headache, wakefulness and morbid sensitiveness of the nervous system generally, especially of the eye. Valuable as a substitute for opium in the treatment of children.

Ladies' Slipper root, U.S.P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Cypripedin, a so-called **Concentration**,* a powdered preparation of Ladies' Slipper.

Dose, 1-2 to 3 grains (0.03 to 0.2 Gm.).

* See note in Preface on Concentrations.

LADIES' SLIPPER—Continued.**Extracts.**

Fluid Extract Ladies' Slipper, U. S. P.—Not miscible with water without some precipitation.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Sculleap Compound.—Not miscible with water. Each pint represents: Sculleap, 5 1-2 ozs.; Ladies' Slipper, 5 1-2 ozs.; Hops, 2 1-2 ounces; Lettuce, 2 1-2 ozs. Combines the tonic and nervine properties of its several ingredients. Indicated in the nervous disturbances of hysterical females, and in the debility of children following an attack of summer complaint.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract Ladies' Slipper.—One part equals 4 of the drug.
Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Pills.

Cypripedium Extract, 2 grs.

Headache, 3 grs.

Lactated Pepsin, 1 gr.

Guarana, 1-2 gr.

Sodium bicarbonate, 1 gr.

Cypripedin, 1-2 gr.

LANOLIN. The oil or fat of sheep's wool.

Properties.—According to Prof. Oscar Liebreich, of Berlin, who has recently perfected methods of extracting and purifying this cholesterin fat, the following advantages are claimed for it as a basis of ointments: 1. It is capable of absorbing and intimately blending with large amounts of water, standing, thus in marked contrast to popular notions of the mutual relations of water and fats. The lanolin of commerce is a combination of neutral cholesterin-fat, with about thirty per cent. of water. 2. It is neutral to test paper, is not liable to rancidity, and is not easily saponified by alkalis. 3. It is miscible with glycerin, unlike other fats. 4. It rapidly, and in a remarkable degree, possesses the power of penetrating the epidermis. 5. Hence, Lanolin greatly facilitates the absorption, through the skin, of drugs mixed with it. So marked is this that, with poisonous drugs, such as the toxic alkaloids, less (about half) than the usual proportion, should be prescribed in ointments. The vaselines and paraffins hinder, rather than favor, the passage of drugs into the skin.

LARKSPUR. *Delphinium consolida, Linné.* (Seed.)

Properties.—Poisonous. Acrid, irritant, in overdoses producing dangerous vomiting and purging. In minute doses, reputed diuretic, emmenagogue and anti-spasmodic. Has been found useful in spasmodic asthma and dropsy. It is employed externally to destroy vermin.

ANTIDOTES—None reliable known; emetics, then treat symptoms as they arise on general principles. Tannin or compound tincture of iodine may be given to retard absorption of the poison.

PREPARATION.

Fluid Extract Larkspur seed.—Not miscible with water.
Dose, 1 minim (0.065 C.c.) or less, gradually increased

LAVENDER. *Lavandula, U. S. P. Lavandula spica, Linné.* (Flowers.)

Synonym.—*Lavandula vera, DeCandolle.*

Properties.—Stimulant, aromatic, and carminative. The official compound tincture Lavender is chiefly employed as a remedy for nausea, flatulent colic and gastric distress after taking food.

Lavender flowers, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATION AND FORMULA.

Fluid Lavender Compound.—Not miscible with water. Four times the strength of Tincture Lavender compound, U. S. P. Each pint repre-

LAVENDER—Continued.

sents: Oil Lavender, 1-2 fl. oz.; Oil Rosemary, 1 fl. drm.; and the soluble constituents of: Cinnamon, 1 1-8 oz.; Cloves, 1-4 oz.; Nutmeg, 5-8 oz.; Red Saunders, 1-2 oz.

Dose, 8 to 15 minims (1-2 to 1 C.c.).

Tincture Lavender Compound, U. S. P.

Fluid Lavender Comp., P., D. & Co.'s.....4 fl. ozs.

Alcohol9 fl. ozs.

Water3 fl. ozs.

Dose, 30 to 60 minims (2 to 4 C.c.).

LEMON. Limonis cortex, U. S. P. Citrus limonum, *Risso*. (Peel of frt.)

Properties.—Aromatic. Used only for flavoring purposes.

PREPARATION.

Fluid Extract Lemon Peel.—Miscible with water.

Dose, as desired for flavor.

LEMON BALM. Melissa, U. S. P. Melissa officinalis, *Linné*. (Herb.)

Synonyms.—Balm, Sweet balm, mélisse, céline, herbe au citron, Fr.; citronen-kraut, Ger.

Properties.—Diaphoretic, antispasmodic, and moderately stimulant. Has been found beneficial in slight febrile conditions, in painful menstruation, and to assist the action of other diaphoretics.

Lemon Balm, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Lemon Balm.—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

LEVANT WORMSEED. Santonica, U. S. P. Artemisia maritima, *Linné* (Seed.)

Synonyms.—Barbotine, Fr.; Zittwersamen, Ger.

Properties.—A most certain anthelmintic against lumbricoid worms, and in a less degree against ascariides. Generally employed in the form of the active principle, Santonin. Inactive against tape-worm.

PREPARATIONS.

Capsules Santonin and Castor Oil, 2 1-2 grms.: in soft gelatin.

Santonin, 1-4 gr.

Castor Oil, 2 1-2 grms.

Lozenges.—A favorite form of exhibiting Santonin to children.

Chocolate Worm.

Santonin, 1-2 gr.

Calomel, 1-2 gr.

Santonin, 1-2 and 1 gr.

Santonin and Calomel.—In two colors, pink and white.

Santonin, 1-2 gr.

Calomel, 1-2 gr.

Santonin Compound.

Santonin, 1-2 gr.

Podophyllin, 1-20 gr

Calomel, 1-2 gr.

Santonin and Podophyllin.

Santonin, 1-2 gr.

Podophyllin, 1-20 gr

Pills.

Anthelmintic, 2 grs.

Santonin, 1 gr.

Calomel, 1 gr.

Santonin, 1-10, 1-2 and 1 gr.

Santonin and Calomel, 1 gr.

Santonin, 1-2 gr.

Calomel, 1-2 gr.

Cacao Butter, q. s.

Santonin and Podophyllin, 1-2 gr.

Santonin, 1-2 gr.

Podophyllin, 1-20 gr.

Tablet Triturates.

Santonin, 1-2 and 1 gr.

Santonin and Calomel.

Santonin, 1-2 gr.

Calomel, 1-2 gr.

LICORICE. Glycyrrhiza, U. S. P. Glycyrrhiza glabra, *Linné*. (Root.)

Synonym.—Süßholzwurzel, Ger.

Properties.—Licorice root is emollient and demulcent. It acts upon the mucous surfaces, lessening irritation, and is consequently useful in cough, catarrhs, irritation of the urinary organs, and in pain of the intestines in diarrhoea. Its chief use is in bronchitis or bronchial irritation. It is pleasant to the taste and has the property of slightly obtunding the gustatory nerves, hence one of its uses is to cover the taste of other medicines such as quinine, quassia, aloes, cascara sagrada, etc.

ACTIVE PRINCIPLE.

Ammonium Glycyrrhizate.—Supposed to be the form in which Glycyrrhizin exists in the root. Employed to mask the bitterness of bitter or nauseous drugs. It is incompatible with acid solutions, soluble in water and alcohol.

Dose, 5 to 15 grains (0.32 to 1 Gm.).

PREPARATIONS.

Elixir Licorice Aromatic, for Quinine.—Each fluidounce represents 120 grains of Licorice.

An elegant vehicle for the administration of quinine and other remedies possessing a disagreeable taste. Place the quinine in powder in a wine-glass, add a dessertspoonful of the elixir, stir briskly till thoroughly mixed, and then let the mixture be taken, followed immediately by about a teaspoonful of the clear elixir.

Extracts.

Fluid Extract Licorice, U. S. P.—Miscible with water.

Dose, 1 to 5 fluidrachms (4 to 20 C.c.).

Fluid Extract Licorice for Quinine, etc.—Miscible with water.

Identical in strength with the official preparation, but free from the bitter extractive which the latter contains, and therefore especially adapted for disguising the taste of bitter and nauseous medicines. It forms clear mixtures with aqueous and alcoholic solutions, but the glycyrrhizin it contains is precipitated by acids. Hence, in quinine mixtures containing licorice, acid must not be prescribed to effect solution of the alkaloids.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Lozenges.—Each containing ingredients specified.

Bronchial with Cubeb.

Ext. Licorice, 4-5 gr.

Powd. Cubeb, 5-8 gr.

Balsam Tolu, 1-20 gr.

Oil Sassafras, 1-40 gr.

Cubeb.

Ext. Licorice, 3-4 gr.

Oleoresin Cubeb, 1-16 gr.

Oil Sassafras, 1-32 gr.

Licorice.—Both plain and sugar-coated.

Opium and Licorice (Wistar's).

Powd. Opium, 1-10 gr.

Ext. Licorice, 1 gr.

Oil Anise, 1-32 gr.

Tinct. Tolu, 1-16 gr.

Yerba Santa Compound.

Ext. Licorice, 1 gr.

Powd. Cherry, 1-4 gr.

Powd. Senega, 3-16 gr.

Yerba santa, 1 gr.

Tar, 1-32 gr.

LIFE-EVERLASTING. Gnaphalium polycephalum, *Michaux*. (Herb.)

Properties.—Tonic, mildly astringent, anodyne. Sometimes used as a diaphoretic in fevers, and as an astringent tonic in chronic diarrhoeas. Locally employed in the treatment of unhealthy ulcers.

Life-everlasting herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Life-everlasting.—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

LIFE-ROOT. *Senecio aureus*, *Linné*. (Herb.)

Synonyms.—*Senecio gracilis*, *Pursh*; Groundsel, Ragwort, Squaw-weed, Female Regulator, *Unkum*; *Senecion*, Fr.; *Kreuzkraut*,

Properties.—Life-root is diuretic, expectorant, diaphoretic and tonic, and exerts a peculiar influence on the female reproductive organs. Is very efficacious in promoting the menstrual flow; and may be given alone or in combination with equal parts of asarum and savin, in amenorrhœa of a functional nature, that is when not due to pregnancy or to any structural change. It is useful, combined with raspberry leaves and cinnamon, in menorrhagia. It has also proved of value in dysmenorrhœa, gravel and other urinary affections, strangury and in dysentery.

Life-root herb, compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Life-root.—Miscible with water without material precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract Life-root.—One part equals 5 of drug.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Pill Female Leucorrhœa, 3 grs.

Hamamelin, 2 grs.

Senecin, 1-2 gr.

Hydrastin, 1-2 gr.

Senecin, a so-called **Concentration**,* a powdered preparation from Life-root.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

LILY OF THE VALLEY. *Convallaria majalis*, *Linné*. (Flowers, herb, root.)

Properties.—"This medicine is proved to be one of the most powerful diuretics known applicable, then, especially to mitral diseases with dropsy ... *Convallaria* diminishes the number of pulsations while augmenting the amplitude of the contractions. It calms also the palpitations and disorders of the heart which are purely functional in character, and if I add that the preparations of this medicament have no toxic action in man, I shall have pointed out their principal advantages."† Albuminuria prevents or lessens its diuretic action. No cumulative effects have been observed.

ACTIVE PRINCIPLE.

Convallamarin.—Glucoside. Cardiac tonic, claimed to be superior to digitalin. Soluble in water.

Dose, 1-12 grain (0.005 Gm.) and upwards, cautiously increased; authorities do not yet agree as to the limit of maximum dose.

PREPARATIONS.**Extracts.**

Fluid Extract Lily of the Valley flowers.—Miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C.c.), which may be gradually increased until effects are secured. This is claimed to be stronger than the fluid extract of the herb or root.

Fluid Extract Lily of the Valley herb.—Miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C.c.), gradually increased.

Fluid Extract Lily of the Valley root.—Not miscible with water without some precipitation.

Dose, 10 to 30 minims (0.6 to 2 C.c.), gradually increased.

Pill Convallamarin, 1-12 gr.

Tablet Triturate, Convallamarin, 1-50 gr.

* See note in Preface on Concentrations.

† Dujardin-Beaumetz, "New Medications," p. 23, (edition of Geo. S. Davis, 1886).

LIME JUICE. Expressed from the fruit of the *Citrus limetta*, *Risso*.

Properties.—A grateful refrigerant tonic and a powerful anti-scorbutic. It is a popular beverage, especially with persons of a "bilious" habit. We only list it under the combination:

Lime Juice and Pepsin.—Each fluidounce represents: Lime juice, pure, 1 fluidrachm; saccharated pepsin, U. S. P., 16 grains. This is an admirable remedy in conditions of general debility attended with loss of appetite and dyspeptic symptoms. It should be taken at meal time, clear or in a little water. Commended for all the minor derangements of digestion, and as a substitute for the alcoholic bitters and liquors so frequently used to promote it. Being strictly a physiological remedy it is adapted to a wide range of cases that do not require any medication, and physicians will find occasion frequently to prescribe it for this very reason.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

LIPPIA MEXICANA. *Lippia dulcis*, *Treviranus*. (Leaves.)

Properties.—Demulcent and expectorant, exercising also an alterative influence over the bronchial mucous membrane. It is an excellent remedy in chronic bronchitis as well as in acute catarrhal affections of the respiratory tract.

PREPARATIONS.

Concentrated Tincture Lippia Mexicana.—Miscible with water without material precipitation. Four parts of tincture equal one of drug.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

LIVERWORT. *Anemone Hepatica*, *Linné*. (Leaves.)

Synonym.—*Hepatica triloba*, *Chaix*.

Properties.—A mild, mucilaginous astringent, useful in fevers and lung complaints.

Liverwort leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Liverwort.—Miscible with water.

Dose, 2 to 3 fluidrachms (8 to 12 C.c.).

LOBELIA, U. S. P. *Lobelia inflata*, *Linné*. (Herb, U. S. P.; seed.)

Synonyms.—Wild or Indian Tobacco.

Properties.—**POISONOUS.** Lobelia is an emetic, occasionally cathartic, and in small doses diaphoretic and expectorant. It is also possessed of some narcotic properties. In large and repeated doses it produces severe vomiting, attended with distressing nausea, copious sweating and general relaxation, followed by extreme prostration, great anxiety and distress, convulsions and death. It has been successfully used in spasmodic asthma, catarrh, croup, pertussis and other laryngeal and bronchial affections, but should be used with caution. Its poisonous effects resemble those of tobacco.

ANTIDOTES.—Tannin or Iodine (compound tincture largely diluted) may be administered to retard absorption of the poison. Emetics if needed. Stimulants; alcohol, opium, external heat, friction, etc.; recumbent posture.

Lobelia herb, U. S. P., } compressed in ounce packages for retailing
Lobelia leaves, U. S. P., } purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Lobelia herb, U. S. P.—Miscible with water.

Dose, as an expectorant, 3 to 30 minims (0.2 to 2 C.c.); as an emetic, 10 minims to 1 fluidrachm (0.65 to 4 C.c.).

Fluid Extract Lobelia seed.—Miscible with water without material precipitation.

Dose, as an expectorant, 1 to 10 minims (0.06 to 0.65 C.c.); as an emetic, 10 minims to 1 fluidrachm (0.65 to 4 C. c.).

LOBELIA—Continued (Extracts).

Fluid Extract Lobelia Compound.—Not miscible with water. Sixteen fluidounces, imperial, represents: Lobelia herb, 8 ozs. av.; Blood root and Skunk Cabbage, 4 ozs. each. Diaphoretic, expectorant, anti-spasmodic, and emetic. Dose, as an expectorant, 3 to 20 minims (0.2 to 1.3 C. c.); as an emetic, 20 to 60 minims (1.3 to 4 C. c.).

Powdered Extract Lobelia.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Lobelia. Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Solid Extract Lobelia.—One part equals 5 of drug. Dose, 1-2 to 2 grains (0.03 to 0.13 Gm.).

Lobelin, a so-called **Concentration**,* a powdered preparation from Lobelia herb. *Synonym.*—Lobelin.

Dose, as an emetic, 1 to 3 grains (0.06 to 0.2 Gm.), repeated if necessary; as an expectorant, 1-4 to 1-2 grain (0.016 to 0.03 Gm.).

Tincture Lobelia, U. S. P.

Fluid Extract Lobelia herb, P. D. & Co.'s..... 2 fl. ozs.
Dilute Alcohol..... 8 1-2 fl. ozs.
Dose, as an antispasmodic, 1-4 to 1 fluidrachm (1 to 4 C. c.); as an emetic, 2 to 4 fluidrachms (8 to 15 C. c.).

NOTE.—Double the strength of the Tincture Lobelia, Phar. Ger.

LOGWOOD. *Hæmatoxylon, U. S. P. Hæmatoxylon campechianum, Linné.* (The heartwood.)

Properties.—Tonic and astringent; may be used with advantage in diarrhœa, dysentery, and in the relaxed condition of the bowels following cholera infantum.

PREPARATION.

Fluid Extract Logwood.—Not miscible with water.

Dose, 10 to 60 minims (0.6 to 4 C. c.).

LOVAGE. *Levisticum officinale, Koch.* (Leaves, root.)

Synonym.—*Ligusticum levisticum, Linné.*

Properties.—A stimulant aromatic, and in hot infusion is employed as a diaphoretic and emmenagogue. It is somewhat of a diuretic and is also of benefit in flatulent dyspepsia.

Lovage leaves, compressed in ounce packages for retailing purposes.

Lovage root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Lovage root.—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

LOW MALLOW. *Malva rotundifolia, Linné.* (Flowers.)

Properties.—Demulcent. An infusion of an ounce to the pint, dose, wine-glassful (60 C. c.), forms an excellent demulcent in coughs, irritation of the air passages, catarrhal affections of the bladder, etc. The flowers bruised form a good emollient cataplasm to boils and local inflammations.

Low Mallow flowers, compressed in ounce packages for retailing purposes.

LUNGMOSS. *Sticta pulmonaria, Acharius.* (Plant.)

Properties.—A simple bitter tonic having also expectorant properties. Used in chronic pulmonary affections with debility in dose of a tablespoonful or two (15 to 30 C. c.) of an ounce to the pint of hot infusion.

Lungmoss, compressed in ounce packages for retailing purposes.

* See note in Preface on Concentrations.

LUNGWORT. *Pulmonaria officinalis*, *Linné.* (Leaves.)

Properties.—Expectorant, demulcent and mucilaginous. It has been used in bleeding from the lungs, bronchial and catarrhal affections, and other affections of the respiratory organs.

Lungwort leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Lungwort.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

MACE. *Macis*, U.S.P. *Myristica fragrans*, *Houttuyn.* (Arillus of fruit.)

Properties.—Carminative stimulant; in large doses narcotic, producing effects similar to those of camphor. It is chiefly used for flavoring purposes, but is sometimes used as an anodyne and astringent in diarrhoeas and dysentery.

PREPARATION.

Fluid Extract Mace.—Not miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C.c.).

MAGNOLIA, U. S. P. *Magnolia glauca*, etc., *Linné.* (Bark.)

Synonyms.—Sweet Bay, White Bay, Beaver Tree, Swamp Sassafras.

Properties.—Diaphoretic, tonic, antiperiodic. In remittent and intermittent fevers it has proved of considerable value, succeeding occasionally where cinchona preparations fail.

Magnolia bark, U. S. P., compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Magnolia.—Not miscible with water without some precipitation.

Dose, 30 to 60 minims (2 to 4 C.c.).

MAIDEN-HAIR. *Adiantum pedatum*, *Linné.* (Leaves.)

Synonyms.—Capillaire, Fr.; Frauenhaar, Venushaar, Ger.

Properties.—Expectorant, demulcent, and slightly tonic. Is much used in pulmonary catarrh, hoarseness, bronchitis, etc.

Dose, wineglassful (60 C.c.) of an ounce to the pint infusion.

Maiden-hair, compressed in ounce packages for retailing purposes.

MALE-FERN. *Aspidium*, U.S.P. *Aspidium Filix-mas*, *Swartz.* (Rhiz.)

Synonyms.—Male Shield-Fern; Fougère Male, Fr.; Wurmfarnwurz, Ger.

Properties.—Tonic, astringent and anthelmintic. Is celebrated as a specific for tape-worm. The extract should be administered in the morning on an empty stomach, followed by an ounce and a half of castor oil at the interval of an hour.

Male-fern, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.**Capsules.**

Male-fern and Castor Oil, 2 1-2 grammes, in soft gelatin.

Oleo-resin Male-fern, 1 grm. Castor Oil, 1 1-2 grm.

Male-fern and Kamala, in soft and hard gelatin.

Oleo-resin Male-fern, 7 minims. Kamala, sifted, 4 grs.

Extracts.

Fluid Extract Male-fern.—Not miscible with water without some precipitation. Prepared with a menstruum of mixed ether and alcohol, to preserve its fluidity, as the ethereal extract soon resolves itself into the oleoresin.

Dose, 1 to 4 fluidrachms (4 to 16 C.c.).

Solid Extract Male-fern.—One part equals 8 of drug.

Dose, 20 to 40 grains (1.3 to 2.6 Gm.).

MANACA. *Brunfelsia Hopeana, DeCandolle.* (Root.)

Synonyms.—*Franciscea uniflora, Pohl; Mercurio Vegetal.*

Properties.—Antisyphilitic, anti-arthritic, purgative and diuretic. From extensive experience with its action in scrofula, syphilis, and chronic and acute rheumatism it has come to be considered one of the most powerful alteratives in the materia medica. "In its use for rheumatic arthritis it is employed with remarkable success."*

PREPARATIONS.

Elixir Manaca and Salicylates.—Each fluidounce represents: Manaca, 80 grs.; Sodium salicylate, 64 grs.; Potassium salicylate, 32 grs.; Lithium salicylate, 8 grs. In its native country, Brazil, Manaca is regarded as a specific for the cure of rheumatic affections, and the results of its use in this country have been very gratifying. This elixir combines its virtues with those of the salicylates of sodium, potassium and lithium, thereby rendering it especially valuable in the treatment of gout and rheumatism.

Fluid Extract Manaca.—Miscible with water without material precipitation.

Dose, 10 to 60 minims (0.65 to 4 C.c.).

MANDRAKE. *Podophyllum, U.S.P. Podophyllum peltatum, Linné.* (Rhizome.)

Synonyms.—May Apple; Fussblattwurzel, Ger.

Properties.—Cathartic or laxative according to dose. Somewhat apt to gripe if taken alone; one of our best remedies in chronic constipation, especially where the liver is torpid.

Mandrake root, U.S.P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLES.

Podophyllin.—Resina Podophylli, U. S. P. This preparation represents the properties of Mandrake root in a highly concentrated form. In small doses it possesses aperient and alterative properties. In large doses it is a valuable cathartic.

Dose, as a laxative, 1-8 to 1-4 grain (0.008 to 0.016 Gm.); as a purgative, 1-4 to 1 grain (0.016 to 0.06 Gm.).

Podophyllin, Phar. Ger.—This differs from the above in the use of water alone for precipitation of the resin. Although the color is much paler, its medicinal character and activity are the same. It answers the German pharmacopœia tests in every particular except one, in which that authority is itself at fault; perchloride of iron does not produce in the aqueous solution a brown color, and will not do so with any sample of pure resin.

Podophyllotoxin.—The active constituent of the resin of Podophyllum. It is about twice as strong as podophyllin.

Dose, 1-10 to 1-2 grain (0.004 to 0.033 Gm.).

PREPARATIONS AND FORMULÆ.

Abstract Mandrake, U. S. P.—One part equals 2 of drug.

Normal Liquid or Fluid Extract Mandrake, P., D. & Co.'s..... 1 fl. oz.
Sugar of milk, q. s. to make 228 grs.
Evaporate the fluid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 223 grains; finally powder the product. Keep in a well-corked bottle.

Dose, 5 to 10 grains (0.3 to 0.6 Gm.).

Capsule Castor Oil and Podophyllin.—In hard and soft gelatin.
Castor Oil, 10 minims. Podophyllin, 1-8 gr.

Elixir Mandrake Comp.—Each fl. oz. represents ingredients specified.
Mandrake root, 45 grs. Senna, 30 grs.
Culver's root, 30 grs. Jalap, 15 grs.

Extracts.

Fluid Extract Mandrake, U. S. P.—Not miscible with water.

Palatably administered in fluid extract licorice.

Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

* Morse, "New Therapeutical Agents," p. 168.

MANDRAKE—Continued (Extract).

Fluid Extract Mandrake Comp.—Not miscible with water. Each pint represents 16 ounces of the following in the proportions named: Mandrake, 6 parts; Culver's root, 6 parts; Senna, 4 parts; Jalap, 2 parts.
Dose, as a laxative, 1 fluidrachm (4 C.c.); as a cathartic, 1 to 3 fluidrachms (4 to 12 C.c.).

Powdered Extract Mandrake.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Mandrake.
Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Solid Extract Mandrake.—One part equals 5 of the drug.
Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Lozenge Santonin and Podophyllin.

Santonin, 1-2 gr. Podophyllin, 1-20 gr.

Normal Liquid * Mandrake.—An assayed preparation. Mandrake root contains 4 to 5 per cent. of Podophyllin, of which about 45 per cent. is Podophyllotoxin. Standard, 4.1-4 per cent total resin.
Dose, 10 to 20 grains (0.6 to 1.3 Gm.).

Pills.

Anti-Bilious, 2 3-4 grs.
Ext. Coloc. co., 2 1-2 grs. Podophyllin, 1-4 gr.

Anti-Constipation, Goss', 2 1-4 grs.
Podophyllin, 1-4 gr. Ext. Henbane, 1-8 gr.
Ext. Coloc., 1-4 gr. Ext. Gentian, 1-4 gr.
Ext. Butternut, 1-2 gr. Ext. Cascara sagrada, 1-4 gr.
Ext. Nux vomica, 1-8 gr. Powd. Canadian Hemp, 1-2 gr.

Cathartic Comp. Granules, 3-4 gr.
Jalapin, 1-16 gr. Powd. Gamboge, 1-32 gr.
Leptandrin, 1-16 gr. Powd. Capsicum, 1-64 gr.
Aloin, 1-8 gr. Ext. Henbane, 1-8 gr.
Podophyllin, 1-4 gr. Oil Peppermint, 1-128 gr.

Cholagogue, 3 grs.
Podophyllin, 1-4 gr. Purified Aloes, 1 gr.
Ext. Henbane, 1-2 gr. Powd. Rhubarb, 1 gr.
Powd. Capsicum, 1-4 gr.

Eccoprotic, 2 1-2 gr.
Ext. Soc. Aloes, 2 grs. Podophyllin, 3-10 gr.
Ext. Nux vomica, 1-5 gr. Oil Cloves, 1-10 gtt.

Hepatic, Eclectic, 1 1-2 gr.
Podophyllin, 1-4 gr. Ext. Nux vomica, 1-16 gr.
Leptandrin, 1-2 gr. Powd. Capsicum, 1-8 gr.
Irisin, 1-4 gr.

Liver, Dr. Taylor's, 2 4-5 grs.
Podophyllin, 1 gr. Powd. Camphor, 4-5 gr.
Powd. Ipecac, 1 gr.

Podophyllin, 1-40, 1-8, 1-4, 1-2 and 1 gr.

Podophyllin, Belladonna and Calabar, 3-4 gr.
Podophyllin, 1-4 gr. Ext. Calabar bean, 1-4 gr.
Ext. Belladonna, 1-4 gr.

Podophyllin and Blue Mass, 2 1-4 grs.
Podophyllin, 1-4 gr. Blue Mass, 2 grs

Podophyllin Compound, 3-4 gr.
Podophyllin, 1-2 gr. Ext. Henbane, 1-8 gr.
Ext. Nux vomica, 1-16 gr.

Podophyllin and Leptandrin, 1 1-2 gr.
Podophyllin, 1-2 gr. Leptandrin, 1 gr.

Podophyllotoxin, 1-8 gr.

Podophyllum Extract, 1 gr.

Triplex, 3 1-4 grs.
Purified Aloes, 2 grs. Podophyllin, 1-4 gr.
Blue Mass, 1 gr.

The above are a few of the principal pills containing Mandrake, for others see appended Pill Formulæ list for numbers 15, 17, 33, 40, 41, 60, 64, 71, 100, 102, 106, 137, 138, 140, 221, 222, 253-255, 257, 261, 263, 352, 356, 402, 448, 460.

Tablet Triturates, Podophyllin, 1-16, 1-8, 1-4 and 1-2 gr.

* See note in Preface on Normal Liquids.

MANDRAKE—Continued.

Tincture Podophyllin, B. P.—Simply a solution of the resin in alcohol, one grain to the fluidrachm.
Dose, 15 to 60 minims (1 to 4 C.c.).

MANGOSTEEN. *Garcinia Mangostana, Linné.* (The pericarp.)

Synonym.—Mango fruit.

Properties.—Astringent. Employed in the treatment of nasal catarrh, diarrhoea, dysentery, leucorrhœa; as a gargle in tonsillitis and, properly diluted, as a lotion for foul ulcers, and for prolapse of the rectum or vagina. It diminishes too profuse menstruation.

PREPARATION.

Fluid Extract Mangosteen (Mango).—Not miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).
NOTE.—The article sold as mango bark is in reality *Garcinia*.

MAN-ROOT. *Ipomœa pandurata, Meyer.* (Root.)

Synonyms.—*Convolvulus panduratus, Linné*; Man-in-the-ground, Man-of-the-earth, Wild Jalap, Wild Potatoe.

Properties.—Feebly cathartic, and claims are made for it in infusion as an excellent diuretic.

Man-root, compressed in pound packages for retailing purposes.

MANZANITA. *Arctostaphylos glauca, Lindley.* (Leaves.)

Properties.—Astringent and tonic, with specific direction to the urinary organs. It is employed in diarrhoea, but chiefly used in treating diseases of the genito-urinary organs. It is prescribed in gonorrhœa and gleet, vesical catarrh, incontinence of urine, chronic nephritis, diabetes insipidus, as well as in leucorrhœa and menorrhagia. It resembles *uva ursi* in its range of action.

PREPARATION.

Fluid Extract Manzanita leaves.—Not miscible with water.
Dose, 20 to 60 minims (1.3 to 4 C.c.).

MARSHMALLOW. *Althæa, U. S. P.* *Althæa officinalis, Linné.* (Flowers, leaves, root.)

Synonyms.—Racine de guimauve, Fr.; Elbischwurz, Ger.

Properties.—Chiefly used as a demulcent in inflammatory affections of the mucous tissues, as bronchitis, vesical catarrh, renal irritation, etc.

Marshmallow flowers, } compressed in ounce packages for retailing
Marshmallow leaves, } purposes.

Marshmallow root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.

Fluid Extract Marshmallow root.—Miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Lozenge Marshmallow.

Powd. Marshmallow, 1-3 gr.

Powd. Orris root, 1-4 gr.

Syrup Marshmallow, U. S. P.

Fluid Extract Marshmallow, P., D. & Co.'s..... 1 fl. oz.

Syrup..... 17 fl. ozs.

Should be made when required; does not keep well.

Dose, ad libitum.

MARSH ROSEMARY. *Statice Limonium*, var. *caroliniana*, Gray.
(Root.)

Synonyms.—Sea lavender, Ink root; Romarin des marais, Fr.; Strandvelke, Ger.

Properties.—A strong astringent, very efficacious in the treatment of passive and chronic diarrhoeas. The decoction (one ounce to the pint) or the fluid extract diluted may be used as a gargle in ulcerations of the mouth and throat, scarlatina anginosa, etc., and as an injection in chronic gonorrhoea, gleet, leucorrhoea, prolapsus ani and uteri, and in some ophthalmic affections.

Marsh Rosemary root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Marsh Rosemary.—Not miscible with water
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

MARY THISTLE. *Silybum marianum*, Gartner.

Synonym.—*Carduus marianus*, Linné.

Properties.—Has the reputation in Europe of being a depurative through its gradual and insensible modification of nutrition somewhat similar in action to burdock seed.

PREPARATION.

“German Tincture” Mary Thistle.—An imported unofficial preparation of the green plant.
Dose, 15 to 60 minims (1 to 4 C.c.).

MASTERWORT. *Heracleum lanatum*, Michaux. (Root.)

Synonym.—Cow Parsnip.

Properties.—Stimulant, anti-spasmodic and carminative. Used in flatulency and dyspepsia. In full doses, taken daily, it has proved efficacious in epilepsy. It is also recommended in asthma, colic, dysmenorrhoea, palsy, apoplexy, intermittents, etc.

Masterwort root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Masterwort.—Not miscible with water.
Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

MASTIC, U. S. P. Resin from *Pistacia Lentiscus*, Linné.

Properties.—Stimulant to the bronchial and urinary passages. In association with aloes in pill form its slower solubility delays the action of the aloes so that it may act more upon the large than the small intestine.

PREPARATIONS.**Pills.**

Aloes and Mastic, U. S. P., 2 grs.	
Purified Aloes, 2 grs.	Powd. Red Rose, 1-2 gr.
Mastic, 1-2 gr	
Dinner, Chapman's, 3 grs.	
Purified Aloes, 1 gr.	Powd. Rhubarb, 1 gr
Gum Mastic, 1 gr.	
Dinner, Lady Webster's, 3 gr.	
Purified Aloes, 1 4-5 gr.	Rose leaves, 2-5 gr.
Gum Mastic, 3-5 gr.	

MATICO, U. S. P. *Artanthe elongata*, Miquel; with other species sometimes accompanying. (Leaves.)

Properties.—MatICO has some reputation as an aromatic tonic and stimulant, and somewhat resembles cubeb in its tendency to the urinary organs.

MATICO—Continued.

It has been used with advantage in leucorrhœa, gonorrhœa, piles, catarrh of the bladder, menorrhagia and chronic mucous discharges; also in dyspepsia due to chronic mucous affection of the stomach.

Matico leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.

Capsules.—In soft gelatin.

Copaiba, Cubeb and Matico.

Copaiba, Para, 6 minims.

Oleoresin Matico, 1 minim.

Oleoresin Cubeb, 3 minims.

Copaiba, Cubeb, Matico and Sandal.

Copaiba, Para, 3 minims

Oleoresin Matico, 1 minim.

Oleoresin Cubeb, 3 minims.

Oil Sandal, East India, 3 minims.

Copaiba and Matico.

Copaiba, Para, 7 minims.

Oleoresin Matico, 3 minims.

Cubeb and Matico.

Oleoresin Cubeb, 7 minims.

Oleoresin Matico, 3 minims.

Matico Oleoresin, 10 minims.

Elixirs.—Each fluidounce represents ingredients specified.

Matico, 120 grs.

Matico Compound.

Matico, 40 grs.

Hydrangœa, 40 grs.

Uva ursi, 40 grs.

Fluid Extract Matico, U. S. P.—Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Tincture Matico, U. S. P.

Fluid Extract Matico, P., D. & Co.'s..... 2 fl. ozs.

Dilute Alcohol..... 19 fl. ozs.

Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

NOTE.—Since alcohol is contra-indicated where matico is used, the fluid extract should be preferred.

MAYWEED. *Anthemis Cotula, Linné.* (Leaves.)

Synonyms.—Wild or Dog-Chamomile, Dillweed; Camomile puante, Maroute, Fr.; Hundskamillenkraut, Gr

Properties.—Popularly used as a sudorific and antispasmodic in the form of infusion, one ounce to the pint, dose, wineglassful (60 C. c.). Flatulent colic and dysmenorrhœa are said to be relieved by it. Externally it is rubefacient.

Mayweed leaves, compressed in ounce packages for retailing purposes.

MENTHOL. A camphoraceous substance derived from the Japanese plant *Mentha arvensis, Linné.*

Properties.—It is employed externally for the relief of neuralgic pains, headache, toothache, etc., and is said to be valuable as an antiseptic inhalation in acute or infusorial catarrh. It is also valuable as a germicide in scarlet fever, diphtheria, typhoid fever, etc.

Menthol Pencils, or cones of menthol inserted in neatly turned wood boxes. These afford a means of carrying this remedy in the pocket and at the same time greatly facilitating its local application.

MERCURY-WEED. *Mercurialis annua, Linné.* (Herb.)

Properties.—This herb has been employed from the most ancient times as a purgative and emmenagogue. It has also been considered by some a diuretic, and has been used in the treatment of syphilitic affections.

PREPARATION.

Fluid Extract Mercury-weed.—Not miscible with water without some precipitation.

Dose, no definite statement has yet been published regarding it. We would be pleased to receive clinical reports of its therapeutics.

MEZEREUM, U. S. P. *Daphne Mezereum, Linné.* (Bark.)

Synonym.—Mezereon.

Properties.—Externally, Mezereum is used for vesicating purposes. Internally administered, it is a stimulant capable of being directed to the skin or kidneys, and in large doses is apt to excite purging, nausea and vomiting. In over-doses it has the effect of the acrid poisons. Is used in the treatment of secondary syphilis, scrofulous affections, chronic rheumatism, palsy and obstinate diseases of the skin.

Mezereum bark, U.S.P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Mezereum bark, U. S. P.—Not miscible with water without some precipitation.

Dose, 5 to 20 minims (0.3 to 1.25 C.c.).

Solid Extract Mezereum, U. S. P.—One part equals 8 of drug.

Used in external applications to increase rubefacient effect

Dose, 1 to 2 grains (0.06 to 0.13 Gm.).

MISTLETOE. *Phoradendron flavescens, Nuttall.* (Leaves.)

Properties.—Laxative, oxytocic and antispasmodic. Its antispasmodic properties are taken advantage of in asthma, epilepsy, palsy and the neuroses. Dr. W. H. Long claims for Mistletoe extraordinary virtues as an oxytocic. As compared with ergot, he says: 1. It acts more promptly and surely; 2. it produces intermittent contractions, instead of tonic—hence, may be used in any stage of labor, or in primiparæ, where ergot is not admissible; 3. it can always be obtained fresh, and does not deteriorate by keeping. Dr. Long has also used this drug with success in many cases of menorrhagia, hemorrhage from the uterus and post partum hemorrhages.

Mistletoe leaves, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Mistletoe leaves.—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (2 to 4 C.c.), repeated every 20 minutes when given in labor.

MOTHERWORT. *Leonurus Cardiaca, Linné.* (Leaves.)

Properties.—Emmenagogue, nervine and antispasmodic; usually given in warm infusions (one ounce to the pint, dose, a wineglassful—60 C.c.) in amenorrhœa from colds, or in suppressed lochia.

Motherwort leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Motherwort.—Miscible with water

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

MOUNTAIN-ASH. *Pirus americana, DeCandolle.* (Bark.)

Synonyms.—*Sorbus americana.* Marshall; American Mountain-ash, Rowan tree.

Properties.—Very astringent, and in infusion is efficacious whenever such an effect is indicated in throat, anus or vagina. It is also given internally to check diarrhœa.

Mountain-ash bark, compressed in pound packages for retailing purposes

MOUNTAIN LAUREL. *Kalmia latifolia, Linné.* (Leaves.)

Synonyms.—Big-leaved Ivy, Spoonwood, Calico Bush, and occasionally as Sheep Laurel, although this is properly *Kalmia angustifolia, Linné.*

Properties.—Poisonous. In medicinal doses it is antisiphilitic, sedative to the heart, and somewhat astringent. Efficacious in primary and

MOUNTAIN LAUREL—Continued.

secondary syphilis, febrile and inflammatory diseases, allaying all febrile and inflammatory action, and lessening the action of the heart. It is also useful in obstinate diarrhoeas. Diluted with water, the fluid extract may be used in the treatment of tinea capitis, psora and other cutaneous affections, but must be used with caution.

PREPARATION

Fluid Extract Mountain Laurel.—Not miscible with water.
Dose, 10 to 30 minims (0.65 to 2 C.c.).

MOUNTAIN-MINT. *Pycnanthemum montanum, Michaux.* (Lvs.)

Properties.—Stimulant, carminative and tonic. Useful particularly as an adjuvant in the treatment of intestinal troubles attended by tormina.

Mountain-mint leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Mountain-mint.—Miscible with water without material precipitation.
Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

MOUNTAIN SAGE. *Artemisia frigida, Willdenow.* (Herb.)

Synonym.—Sierra Salvia.

Properties.—First introduced as a substitute for quinine. In the treatment of periodic fevers, a teaspoonful of the fluid extract is given in a glass of strong, hot lemonade, one hour before the expected chill, and repeated in thirty minutes if the stage of perspiration is not exhibited. In rheumatism, scarlet fever, diphtheria, etc., it is used hot, as above, and repeated every half hour until perspiration and urination are certainly established.

PREPARATION.

Fluid Extract Mountain Sage (Sierra Salvia).—Not miscible with water without some precipitation.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

MUGWORT. *Artemisia vulgaris, Linné* (Leaves.)

Synonyms.—Armoise Commune, Fr.; Beifuss, Ger.

Properties.—Alterative, tonic, anthelmintic, narcotic. In considerable repute in Germany as an emmenagogue; in the treatment of epilepsy; and in connection with asafetida, in chorea. Locally in bruises, sprains, etc.

Mugwort leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Mugwort leaves.—Miscible with water without material precipitation.
Dose, 20 to 60 minims (1.3 to 4 C.c.).

Fluid Extract Feverbush (Spicewood) Berries Comp.—Miscible with water. 15 1-2 ounces, imperial, represent: Mugwort, 13 ozs. av.; Orange peel, 1 oz.; Feverbush berries, 1-2 oz.; Licorice root, 1 oz. A desirable remedy for the treatment of skin diseases, dependent upon peculiar conditions of the system, other than those of syphilitic or scrofulous origin. It possesses valuable tonic and alterative properties, and has been employed with great benefit in the treatment of acne, boils, etc.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

MULLEIN. *Verbascum Thapsus, Linné.* (Leaves, root.)

Synonyms.—Wollkraut, Königskerze, Ger.

Properties.—Demulcent, diuretic, anodyne and antispasmodic. Employed in coughs, catarrh, cystitis, diarrhœa, dysentery and hæmorrhoids. Its

MULLEIN—Continued.

diuretic properties are weak, yet it is useful in correcting by its demulcent action the acidity of the urine, which is present in many diseases. As a popular remedy in phthisis pulmonalis, mullein is said not only to relieve cough and the troublesome diarrhoea of tuberculosis, but to improve the general nutrition so that patients actually gain in weight.

Mullein leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Mullein leaves.—Miscible with water.
Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Fluid Extract Mullein root.—Miscible with water.
Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

MUSK-ROOT. Sumbul, U. S. P. *Ferula Sumbul*, *Hooker filius*.
(Root.)

Synonym.—Jatamansi.

Properties.—A nervous stimulant in typhoid conditions, in asthenic cases of dysentery and diarrhoea, and in malignant cholera. Exercises a calming influence in delirium superior to that of opium. Recommended in gastric spasms, hysteria, chlorosis, amenorrhoea, dysmenorrhoea, epilepsy and other nervous disorders. It has been found useful in chronic bronchitis, pneumonia slow of resolution, in the moist asthma of old, anæmic and scorbutic patients, leucorrhoea and hypochondriasis. It resembles valerian in the general range of its therapeutic indications, but is less disagreeable and more powerful.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Musk-root.—Not miscible with water.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

Solid Extract Musk-root.—One part equals 5 of root.
Dose, 2 to 5 grains, (0.13 to 0.32 Gm.).

Pill, Sedative, 2 grs.

Ext. Musk-root, 1-2 gr.

Ext. Henbane, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Indian Cannabis, 1-10 gr.

Tincture Musk-root, U. S. P.

Fluid Extract Musk-root, P., D. & Co.'s.....2 fl. ozs.

Alcohol.....21 fl. ozs.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

MYRRH. Myrrha, U. S. P. Gum-resin from *Balsamodendron Kataf*, *Kunth*.

Properties.—A stimulating tonic, with some tendency to the lungs, and perhaps to the uterus. It is employed as an expectorant and emmenagogue in debilitated states of the system, in the absence of febrile excitement or acute inflammation. Used in chronic catarrh, phthisis pulmonalis and other pectoral affections in which the secretion of mucus is abundant but not easily expectorated; chlorosis, amenorrhoea, and in various affections connected with the state of the uterine functions. It is generally combined with chalybeates and other tonics, and in amenorrhoea very frequently with aloes. Used also as an application to spongy gums, in aphthous sore throat of children, and various kinds of unhealthy ulcers.

PREPARATIONS AND FORMULÆ.

Fluid Aloes and Myrrh.—Not miscible with water.

Each pint represents 6 2-5 ozs. av. each of Aloes and Myrrh; four times stronger than tincture of Aloes and Myrrh, U.S.P., for the extemporaneous preparation of which it is specially adapted. Incompatible with acid solutions.

Dose, 15 to 30 minims (1 to 2 C.c.).

Fluid Extract Myrrh.—Not miscible with water.

Dose, 10 to 30 minims (0.65 to 2 C.c.).

MYRRH—Continued.**Pills.**

Aloes and Myrrh, U. S. P., 3 1-2 grs.
 Purified Aloes, 2 grs. Aromatic Powder, 1-2 gr.
 Myrrh, 1 gr.

Galbanum Compound, U. S. P., 3 1-2 grs.
 Galbanum, 1 1-2 grs. Myrrh, 1 1-2 grs.
 Asafetida, 1-2 gr.

Iron Compound, U. S. P., 3 grs.
 Myrrh, 1 1-2 grs. Iron sulphate, 3-4 gr.
 Sodium carbonate, 3-4 gr.

Rhubarb Compound, U. S. P., 4 1-2 grs.
 Powd. Rhubarb, 2 grs. Powd. Myrrh, 1 gr.
 Purified Aloes, 1 1-2 grs. Oil Peppermint, 1-10 gr.

For other pills containing Myrrh see Pill Formulæ list appended for numbers 187 and 392.

Tinctures.

Aloes and Myrrh, U. S. P.—(Elixir Proprietatis.)
 Fluid Aloes and Myrrh, P., D. & Co.'s..... 4 fl. ozs.
 Alcohol.....12 fl. ozs.
 Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Myrrh, U. S. P.
 Fluid Extract Myrrh, P., D. & Co.'s..... 2 fl. ozs.
 Alcohol.....9 1-2 fl. ozs.
 Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.)

NETTLE. *Urtica dioica, Linné.* (Flowers, leaves, root.)

Synonym.—Stinging nettle.

Properties.—Astringent, tonic, and one of the purest and most powerful diuretics in our materia medica. Its effects are increased, if rheumatic pains or much dropsy exist, by combining it with potassium acetate. It is also used to advantage in gravel and other nephritic complaints, in summer complaints of children, and bowel affections of adults. It is an excellent remedy in both passive and symptomatic uterine hemorrhage.

Nettle flowers, } compressed in ounce packages for retailing purposes.
Nettle leaves, }
Nettle root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Nettle root.—Miscible with water.
 Dose, 20 to 40 minims (1.3 to 2.6 C. c.).

NEW JERSEY TEA. *Ceanothus americanus, Linné.* (Root.)

Synonyms.—Jersey tea, Red root; Céanothe, Fr.; Seckelblumen-wurzel, Ger.

Properties.—Astringent, expectorant, sedative, anti-spasmodic and anti-syphilitic. Used in gonorrhœa and dysentery; and in asthma, chronic bronchitis, and other pulmonary affections. Successfully used as a gargle and wash in the aphthæ of children, sore mouth subsequent to fevers, and ulceration of the fauces attending scarlatina.

New Jersey Tea root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract New Jersey Tea root.—Not miscible with water.
 Dose, 10 to 30 minims (0.6 to 2 C. c.).

NIGHT-BLOOMING CEREUS. *Cereus grandiflorus, Haworth.* (Stems and flowers.)

Synonym.—Cactus grandiflorus, Linné.

Properties.—Sedative and diuretic. It is especially useful in functional diseases of the heart attended with much irregularity of action, in which it exerts a decided action, palliating or removing the symptoms and

NIGHT-BLOOMING CEREUS—Continued.

frequently giving prompt relief. It has been found serviceable in palpitation, angina pectoris, cardiac neuralgia, rheumatism, valvular disease, also of hæmoptysis, dropsy, and in threatened apoplexy.

PREPARATION.

Fluid Extract Night-blooming Cereus.—Miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

NUT-GALL. Galla, U.S.P. Excrescences caused by puncture of an insect, and deposited ova in the bark of *Quercus infectoria*, *Olivier.*

Synonyms.—Galls; Noix de galle, Fr.; Galläpfel, Ger.

Properties.—Powerfully astringent, owing to the large amount of tannic acid present. Seldom used internally, though occasionally prescribed in chronic diarrhoea and chronic dysentery, and used in cases of poisoning by the vegetable alkaloids. In the form of an infusion (1 to 30) used as an astringent gargle, lotion and injection.

PREPARATIONS AND FORMULA.

Fluid Extract Nut-gall.—Miscible with water.

Dose, 10 to 30 minims (0.6 to 2. C.c.).

Glycerole Tannic Acid, 5 grains in each fluidounce.

Dose, 2 fluidrachms (8 C.c.).

Lozenge Tannic Acid, U. S. P., 1 gr.

Pills.

Camphor, Opium and Tannin, 1-4 gr.

Camphor, 1-16 gr.

Opium, 1-64 gr.

Tannin, 1-8 gr.

Camphor, Opium and Tannin, 3 1-4 grs.

Camphor, 1 gr.

Opium, 1-4 gr.

Tannin, 2 grs.

Tannin, 1-20, 1, 3 and 5 grs.

Tincture Nut-gall, U. S. P.

Fluid Extract Nut-gall, P., D. & Co.'s 4 fl. ozs.

Dilute Alcohol..... 15 fl. ozs.

Glycerin..... 1 2-3 fl. ozs.

Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

NUTMEGS. Myristica, U. S. P. Kernel of the seed of *Myristica fragrans*, *Houttuyn.*

Properties.—Aromatic, stimulant stomachic. In large doses it acts as a narcotic. Although stimulating to the digestive process it is almost solely used for flavoring purposes.

PREPARATION.

Fluid Extract Nutmeg.—Not miscible with water.

Dose, 15 to 60 minims (1 to 4 C.c.).

NUX VOMICA, U. S. P. *Strychnos Nux-vomica*, *Linné.* (Seeds.)

Properties.—Poisonous. In small doses tonic, especially to nervous system.

It is used in general and local paralysis, in neuralgia, impotence, neurasthenia, in chronic constipation depending on deficient peristaltic action, and as a general roborant.

ANTIDOTES.—Iodine (compound tincture, diluted), tannin. Use emetics (apomorphine, zinc sulphate, mustard). Chloral with potassium bromide or, better, inhalations of chloroform to control the spasms. Artificial respiration.

NOTE.—The strychnine of *Nux vomica* preparations is precipitated from solutions on addition of bromides, chlorides, or iodides.

NUX VOMICA—Continued.**ACTIVE PRINCIPLES.**

Brucine.—One of the alkaloids of *Nuxvomica*. Given internally it produces effects similar to those of strychnine, but very much less active. Applied locally it has anæsthetic properties similar to those of cocaine. A five-per-cent. solution relieves the itching of pruritus, and mitigates the pain produced by local applications of iodine, silver nitrate, etc.
Dose, 1-10 to 1-3 grain (0.006 to 0.02 Gm.).

Strychnine.—Only handled in preparations, see below.

PREPARATIONS AND FORMULÆ.

Abstract Nux Vomica, U. S. P.—One part equals 2 of drug.

Normal Liquid or Fluid Extract *Nux vomica*, P., D. & Co.'s..... 1 fl. oz.

Sugar of milk, q. s. to make..... 228 grs.

Evaporate the fluid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well-corked bottle.

Dose, 1-2 grain (0.03 Gm.).

Elixirs.—Each fluidounce represents ingredients specified.

Ammonium and Strychnine Valerianates.

Ammon. valerianate, 16 grs. Strych. valerianate, 8-60 gr.

Bismuth and Strychnine.

Bismuth and Ammon. cit., 8 grs. Strychnine, 8-100 gr.

Calisaya, Iron and Strychnine.

Calisaya bark (alkaloids), 40 grs. Strychnine, 8-100 gr.

Iron pyrophosphate, 8 grs.

Damiana, Phosphorus, Nux Vomica and Iron.

Damiana, true, 60 grs.

Iron pyrophosphate, 8 grs.

Nux vomica, 8 grs.

Phosphorus, 4-100 gr.

Iron, Quinine and Strychnine Phosphates.

Iron phosphate, 16 grs.

Strychnine phosphate, 8-60 gr.

Quinine phosphate, 4 grs.

Phosphorus and Strychnine.

Phosphorus, 8-100 gr.

Strychnine, 8-100 gr.

Strychnine, 8-100 gr.

Strychnine Valerianate, 8-50 gr.

For others containing *Nux vomica* see appended Elixir Formulæ list for numbers 27, 28, 33, 36-39, 52, 59, 73, 83, 87, 88, 90, 91, 95, 96, 112, 115, 117, 120-123.

Extracts.

Fluid Extract Nux Vomica, U. S. P.—Not miscible with water without some precipitation.

Dose, 1 to 10 minims (0.06 to 0.6 C.c.).

Powdered Extract Nux Vomica, assayed.—Of same strength as the solid extract; prepared by evaporating, at a low temperature, the standard solid extract, and replacing the moisture with powdered sugar of milk.

Dose, 1-10 to 1 grain (0.006 to 0.06 Gm.).

Solid Extract Nux Vomica, U.S.P., assayed.—One part equals 5 of drug. It contains 15 per cent. of alkaloids, about one-half of which is strychnine.

Dose, 1-10 to 1 grain (0.006 to 0.06 Gm.).

Glycerole Damiana, Phos. and Nux Vomica Comp.

Damiana, 60 grs.

Nux vomica, 4 grs.

Phosphorus, 1-25 gr.

Granular Effervescent Preparations.—Each drachm represents ingredients specified.

Iron Phosphate, Quinine and Strychnine.

Iron phosphate, 2 grs.

Strychnine, 1-60 gr.

Quinine, 1 gr.

Pepsin, Bismuth and Strychnine.

Pepsin, saccharated, 5 grs.

Strychnine, 1-60 gr.

Bismuth and Ammon. cit., 2 grs.

Hypodermic Tablets Strychnine Sulphate, 1-100 and 1-50 gr.

NUX VOMICA—Continued.

Normal Liquid * Nux Vomica.—An assayed preparation; standard 1.5 per cent. total alkaloid.

Dose, 1 to 10 minims (0.06 to 0.6 C. c.).

Malt Extract Combinations.—Each fluidounce represents ingredients specified.

Malt Extract Ferrated with Strychnine.

Iron pyrophosphate, 8 grs. Strychnine, 4-100 gr.

Malt Extract with Iron, Quinine and Strychnine.

Iron and Quinine citrate, 8 grs. Strychnine, 4-100 gr.

Malt Extract with Pepsin and Strychnine.

Pepsin, sacch. U. S. P., 4 grs. Strychnine, 4-100 gr.

Malt Extract with Strychnine.

Strychnine, 4-100 gr.

Oleate Strychnine.—In solution, containing 2 per cent. of the alkaloid; for dermatic application.

Pills.—Containing Nux vomica or its alkaloids.

Aloes and Nux Vomica, 2 grs.

Purified Aloes, 1 1-2 grs.

Ext. Nux vomica, 1-2 gr.

Aloin, Strychnine and Belladonna Comp., 2-3 gr.

Aloin, 1-5 gr.

Ext. Belladonna, 1-8 gr.

Strychnine, 1-60 gr.

Ext. Cascara sagrada, 1-2 gr.

Anti-Dyspeptic, 4 1-4 grs.

Strychnine, 1-40 gr.

Blue Mass, 2 grs.

Ext. Belladonna, 1-10 gr.

Ext. Coloc. comp., 2 grs.

Powd. Ipecac, 1-10 gr.

Aphrodisiac Compound, 2 3-4 grs.

Ext. Coca, 1 gr.

Ext. Nux vomica, 1-8 gr.

Cinchonidine sulph., 1-2 gr.

Iron bromide, 1 gr.

Phosphorus, 1-200 gr.

Bismuth and Nux Vomica, 5 1-2 grs.

Bismuth subnitrate, 5 grs.

Ext. Nux Vomica, 1-2 gr.

Carminative, 3 1-4 grs.

Strychnine, 1-20 gr.

Powd. Black Pepper, 1 1-2 grs.

Powd. Ipecac, 2-3 gr.

Ext. Gentian, 1 gr.

Cascara Sagrada and Nux Vomica, 2 1-5 grs.

Ext. Cascara sagrada, 2 grs.

Ext. Nux vomica, 1-5 gr.

Eccoprotic, 2 1-2 grs.

Ext. Soc. Aloes, 2 grs.

Podophyllin, 3-10 gr

Ext. Nux vomica, 1-5 gr.

Oil Cloves, 1-10 gtt.

Iron, Quinine and Strychnine Citrate, 2 grs.

Iron and Quinine citrate, 2 grs.

Strychnine citrate, 1-50 gr.

Iron and Strychnine, 2 grs.

Strychnine, 1-60 gr.

Iron by hydrogen, 2 grs.

Lapactic, 1-2 gr.

Aloin, 1-4 gr.

Ext. Belladonna, 1-8 gr.

Strychnine, 1-60 gr.

Ipecac, 1-16 gr.

Nux Vomica, 1-50 gr.

Nux Vomica Extract, 1-8, 1-4 and 1-2 gr.

Phosphorus, Aloes and Nux Vomica, 3-4 gr.

Phosphorus, 1-50 gr.

Ext. Nux vomica, 1-4 gr.

Ext. Aloes, 1-2 gr.

Phosphorus Compound, No. 1, 1-4 gr.

Phosphorus, 1-100 gr.

Ext. Nux vomica, 1-4 gr.

Phosphorus, Iron, Quinine and Strychnine, 3 1-2 grs.

Phosphorus, 1-50 gr.

Quinine sulphate, 1-2 gr.

Iron by hydrogen, 3 grs,

Strychnine, 1-40 gr.

Strychnine, 1-100, 1-60, 1-50, 1-48, 1-40, 1-32, 1-30, 1-20 and 1-16 gr.

Strychnine Sulphate, 1-100 gr.

Strychnine Compound, Formula "B.," 2 grs.

Strychnine, 1-100 gr.

Phosphorus, 1-100 gr.

Ext. Indian Cannabis, 1-16 gr.

Iron proto-carb. (Vallet's), 1 gr.

Powd. Ginseng, 1 gr.

* See note in Preface on Normal Liquids.

NUX VOMICA—Continued (Pills).**Tonic, Hæmatiole,** Andrews', 23-4 grs.

Quinine sulphate, 1 gr.	Powd. Ipecac, 1-8 gr.
Iron by hydrogen, 1 1/2 grs.	Strychnine sulphate, 1-40 gr.
Arsenious Acid, 1-40 gr.	

For others see appended Pill Formulæ list for numbers 13, 16—18B, 33—36, 41, 43, 45, 53, 95, 98, 100, 130, 206, 231, 232, 236, 238, 255, 281, 294, 296, 297, 302, 304, 309, 310, 312, 313, 325—331, 335, 336, 338—342, 344, 355, 375, 380—382, 385, 407, 414, 426, 448, and 490.

Syrups.—Each fluidounce represents ingredients specified.

Hæmatic Hypophosphites, Improved.—A scientific substitute for Fellow's Hypo-phosphites, which it faithfully reproduces in all essential particulars. It contains the same medicinal ingredients as Fellow's, but is a permanently clear syrup, while that preparation throws down an unsightly precipitate. No certificates of the value of this preparation are offered; physicians are allowed to judge from its published formula what are its merits.

Potassium hypophos., 1 1/2 grs.	Iron hypophos., 1 1/4 grs.
Manganese hypophos., 1 gr.	Calcium hypophos., 1 gr.
Strychnine hypophos., 1-16 gr.	Quinine hypophos., 7-16 gr.

Iron, Quinine and Strychnine Phosphates.

Iron phosphate, 16 grs.	Strychnine phosphate, 8-60 gr.
Quinine phosphate, 8 grs.	

Iron and Strychnine Citrate, 16 grs.**Tablet Triturates.****Aloin and Belladonna Compound.**

Aloin, 1-5 gr.	Strychnine sulphate, 1-60 gr.
Fl. ext. Belladonna, 5-8 minim.	

Iron, Quinine and Strychnine.

Iron, reduced, 1-2 gr.	Strychnine sulphate, 1-120 gr.
Quinine sulphate, 1-2 gr.	

Iron and Strychnine Citrate, 1-4, 1-2 and 1 gr.**Iron and Strychnine Compound.**

Iron, reduced, 1-2 gr.	Arsenious acid, 1-100 gr.
Quinine sulphate, 1-2 gr.	Strychnine sulphate, 1-120 gr.

Nux Vomica, Powdered, 1-100, 1-50, 1-10 and 1-4 gr.**Strychnine Sulphate,** 1-100, 1-60, 1-50, 1-40, 1-30, 1-20 and 1-16 gr.**Tinctures.****Nux Vomica, Phar. Ger.**

Fluid Extract Nux vomica, P., D. & Co.'s.....	1 fl. oz.
Dilute Alcohol.....	10 fl. ozs.
Dose, 8 to 80 minims (0.5 to 5 C.c.).	
NOTE.—Only half the strength of the U. S. P. tincture.	

Nux Vomica, U. S. P.—Containing 0.3 per cent. alkaloids.

Fluid Extract Nux vomica, P., D. & Co.'s.....	4 fl. ozs.
Alcohol.....	19 fl. ozs.
Dose, 4 to 40 minims (0.25 to 2.5 C.c.).	

OPIUM, U. S. P. Concrete milky exudation from the capsules of *Papaver somniferum*, *Linné*.

Properties.—Poisonous. A stimulant narcotic; it diminishes pain (insensibility), causes sleep, and arrests secretions excepting that of the skin which it promotes. Apt to produce nausea and constipation. In overdoses death results from paralysis of the respiratory muscles.

ANTIDOTES.—Atropine, caffeine. Evacuate the stomach by emetics (apomorphine, zinc sulphate, mustard, with copious draughts of luke-warm water) or stomach pump. Maintain respiration (faradization, flagellation, cold douche, etc.). Give strong tea or coffee (fluid extracts best, in drachm doses); belladonna or atropine, watching effect on pupil and on respiration. Alcoholic stimulants in stage of depression; artificial respiration, so long as there are signs of cardiac action.

ACTIVE PRINCIPLES.

Apomorphine Muriate.—Amorphous. Derivative alkaloid from morphine. A prompt and efficient emetic, especially used in cases of poisoning. Lately its value as an expectorant has been pointed out by Dr. Rossbach, who considers it the most efficient remedy we possess in

OPIUM—Continued.

chronic catarrhal affections of the air passages, and in croup pronounced it invaluable. Physiological experiment has shown that it produces a more copious secretion of mucus from the trachea than any other drug which can be substituted for it, except ipecac.

Dose, as an emetic, 1-15 to 1-10 grain (0.004 to 0.006 Gm.); as an expectorant, 1-60 to 1-30 grain (0.001 to 0.002 Gm.).

Codeine.—Alkaloid. Less powerful than morphine and less liable to be followed by unpleasant secondary effects. Used chiefly as a cough remedy. Frequently cures nervous headaches.

Dose, 1-10 to 1 grain (0.006 to 0.06 Gm.) and upwards.

For others see preparations below.

PREPARATIONS AND FORMULÆ.

Bronchial Sedative, Formula of B. W. Palmer, M. D.—Each fluidounce represents: Ammonium chloride, 30 grs.; Fluid Tolu, soluble, 8 minims; Fluid Opium Camphorated, 4 minims; Elixir Licorice, aromatic, q. s. ad. 1 fluidounce. In simple uncomplicated bronchitis of children or adults, in which the cough is the most prominent and distressing symptom its action is almost specific, and in the severer forms of inflammation of the respiratory tract it allays the most urgent symptoms and promotes speedy resolution.

Dose, for an adult, 1 fluidrachm (4 C. c.).

Chlor-anodyne.—Each gramme (ordinary adult dose) represents:

Tr. Capsicum	.0025	Morphine muriate,	.0060
Hydrocyanic Acid, dil...	.0170	Tr. Indian Cannabis.....	.0800
Alcohol	.3000	Chloroform.....	.1350
Glycerin	.4570	Oil Peppermint	.0025

An efficient and elegant remedy for colic, cholera morbus, neuralgia, spasmodic pains, etc. It is of great value where antispasmodics or anodynes are indicated—concentrated in form and rapid in action.

Collodion, Morphine.—Each fluidounce contains 4 grs. Morphine.

Compressed Tablets, Dover's Powder, 5 grs.

Confection, Opium, U. S. P., 1870.—Confection Theriac. Thirty-eight grains equal 1 gr. Opium.

Elixirs.—Each fluidounce represents ingredients specified.

Ammonium and Morphine Valerianates.

Ammonium valerianate, 16 grs. Morphine valerianate, 1-2 gr.

Morphine Bromide, 1 gr.

Morphine Valerianate, 1 gr.

Extracts.

Fluid Opium, Aqueous and Deodorized.—See Tr. Opium, deodorized, U. S. P.

Fluid Opium Camphorated.—Not miscible with water without some precipitation. Eight times the strength of Tincture Opium Camphorated, U. S. P. (Paregoric), for the preparation of which it may be employed. Anodyne, carminative, anti-spasmodic; much used to allay cough in chronic catarrh, asthma, consumption, pertussis, etc., to relieve nausea and slight pains in the stomach and bowels; to check diarrhoea; in teething, and in irritative febricula of infants. Dose, for an adult, 8 to 20 minims (0.5 to 1.3 C. c.); for an infant, 1 to 3 minims (0.06 to 0.2 C. c.).

Powdered Extract Opium.—Contains 20 per cent. morphine by assay. Prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered sugar of milk.

Dose, 1-4 to 1 grain (0.016 to 0.65 Gm.).

Solid Extract Opium, U. S. P.—Contains 20 per cent. morphine by assay.

Dose, 1-4 to 1 grain (0.016 to 0.65 Gm.).

Hypodermic Tablets.

Apomorphine Muriate, 1-10 gr.

Morphine Sulphate, 1-8 and 1-4 gr.

Morphine and Atropine, No. 1.

Morphine sulphate, 1-8 gr. Atropine sulphate, 1-200 gr.

Morphine and Atropine, No. 2.

Morphine sulphate, 1-4 gr. Atropine sulphate, 1-150 gr.

OPIUM—Continued.**Lozenges.****Ammonia, Dr. Jackson's.**

Ammonia muriate, 1-2 gr.

Morphine muriate, 1-60 gr.

Ipecac and Morphine.

Powd. Ipecac, 1-8 gr.

Morphine sulphate, 1-32 gr.

Opium and Licorice, Wistar's Cough.

Powd. Opium, 1-10 gr.

Ext. Licorice, 1 gr.

Oil Anise, 1-32 gr.

Tinct. Tolu, 1-16 gr.

Pectoral, Dr. Jackson's.

Ipecac, 1-20 gr.

Antimony sulphurated, 1-40 gr.

Morphine muriate, 1-33 gr.

Oleates.**Morphine.**—Containing 10 per cent. of the alkaloid.**Morphine and Mercury.**—Containing 5 per cent. of the alkaloid and 20 per cent. of mercuric oxide. Useful where these remedies cannot be administered in the usual way, or when local action is desired.**Pills.**—Containing opium or its derivatives.**Apomorphine,** 1-60 gr.**Calomel and Opium,** 3 grs.

Calomel, 2 grs.

Opium, 1 gr.

Camphor Compound, 3 grs.

Camphor, 1 gr.

Powd. Kino, 1 gr.

Powd. Opium, 1 gr.

Ext. Capsicum, 1-16 gr.

Codeine, 1-16, 1-8 and 1-4 gr.**Dover's Powder,** 1-4 gr. See also Ipecac and Opium below.**Heim's,** 2 grs.

Quinine sulphate, 1 gr.

Powd. Ipecac, 1-4 gr.

Powd. digitalis, 1-2 gr.

Powd. Opium, 1-4 gr.

Ipecac and Opium, 1-2, 1 and 2 grs.

Containing of these ingredients 1-4, 1-2 and 1 gr. each respectively.

Morphine Acetate, 1-8 gr.**Morphine and Atropine,** 1-8 gr.

Morphine sulphate, 1-8 gr.

Atropine sulphate, 1-100 gr.

Morphine Hydrochlorate, 1-4 and 1-8 gr.**Morphine Sulphate,** 1-50, 1-32, 1-20, 1-16, 1-10, 1-8, 1-6, 1-4 and 1-2 gr.**Morphine Valerianate,** 1-8 gr.**Opium, powdered,** 1-40 and 1-2 gr.**Opium Extract,** 1-4, 1-2 and 1 gr.**Opium and Lead Acetate, Formula "B.,"** 2 grs.

Opium, 1-2 gr.

Lead acetate, 1 1-2 grs.

Quinine, Ipecac and Opium, 2 1-2 grs.

Quinine hydrobromate, 2 grs.

Pulv. Opium, 1-4 gr.

Pulv. Ipecac, 1-4 gr.

Sedative, Mann's, 2 1-2 gr.

Ext. Coca, 1-2 gr.

Arsenious Acid, 1-80 gr.

Ext. Valerian, 1-2 gr.

Strychnine, 1-80 gr.

Ext. Indian Cannabis, 1-4 gr.

Codeine, 1-8 gr.

Vallet's Mass, 1 gr.

For others see Pill Formulæ list appended for numbers 28, 51, 73, 81, 83, 86—89, 160, 161, 170, 172, 270, 275, 276, 278—281, 287, 289, 323, 324, 332, 333, 337, 410, 430, 446, 456 and 457.

Plaster Opium, U. S. P.—One in 8 of Opium. Convenient for local application to painful parts of limited extent.**Syrup Dover's Powder.**—Each fluidounce representing 40 grs. Dover's Powder, U. S. P., or 0.56 grs. morphine alkaloid, equivalent to 3-4 gr. morphine sulphate.

Dose, for adults, 1 to 2 fluidrachms (4 to 8 C.c.).

Tablet Triturates.**Apomorphine Muriate,** 1-100 and 1-50 gr**Codeine,** 1-4 and 1-2 gr.**Dover's Powder,** 1-4 and 1 gr.**Morphine and Atropine, No. 1.**

Morphine sulphate, 1-8 gr.

Atropine sulphate, 1-150 gr.

OPIUM—Continued (Tablet Triturates).**Morphine and Atropine, No. 2.**

Morphine sulphate, 1-4 gr. Atropine sulphate, 1-120 gr.

Morphine Hydrochlorate, 1-8 and 1-4 gr.**Morphine Sulphate**, 1-20 1-16, 1-8, 1-6, 1-4 and 1-2 gr.**Opium**, 1-4 and 1-2 gr.**Tinctures.****Ipecac and Opium**, assayed.—Miscible with water.

A liquid form of Dover's Powder, representing 1 grain in each minim.

Dose, 2 to 12 minims (0.13 to 0.75 C.c.).

This may be made extemporaneously by the following formula:

Fluid Extract Ipecac, P., D. & Co.'s 1 fl. oz.

Fluid Opium, aqueous, P., D. & Co.'s 40 fl. ozs.

Reduce the Fluid Opium to 8 1-2 fluidounces by careful evaporation; add the fluid extract Ipecac. Filter and add dilute alcohol through filter to make 10 fluidounces.

Opium, U. S. P.—Laudanum. Each fluidounce contains by assay 6 grs. Morphine, equivalent to 7 1-2 grs. Morphine sulphate.

Dose, corresponding to 1 grain powdered opium, 10 minims (0.65 C.c.).

NOTE.—Of the same strength as Tincture Opium, Phar. Ger.

Opium Camphorated, U. S. P.—Each fluidounce contains by assay 1-4 gr. of morphine, alkaloid.

Dose, for an adult, 1 to 2 fluidrachms (4 to 8 C.c.); for an infant, 5 to 20 minims (0.3 to 1.3 C.c.).

This may be made extemporaneously as follows;

Fluid Opium Camphorated, P., D. & Co.'s 4 fl. ozs.

Glycerin 1 fl. oz.

Dilute Alcohol 27 fl. ozs.

Opium Compound.—Diarrhoea mixture, Squibb's formula. Each fluidrachm represents: Tinct. Opium, 12 minims; Spirits Camphor, 12 minims; Tinct. Capsicum, 12 minims; Purified Chloroform, 4 1-4 minims; Alcohol, 19 1-2 minims.

Dose, for adult, 1-2 to 1 fluidrachm (2 to 4 C.c.) in water.

Opium Deodorized, U. S. P. (Fluid Opium, Aqueous and Deodorized).—Miscible with water.

A preparation of opium which is not liable to nauseate delicate stomachs, nor to produce the unpleasant after-effects so often experienced from other opiates. It is a substitute for McMunn's Elixir and in appropriate doses for Battley's Sedative Drops. Each fluidounce contains by assay six grains of morphine.

Dose, 4 to 15 minims, 5 to 20 drops (0.25 to 1 C.c.). (10.5 minims represent 1 grain of powdered opium.)

Vinegar Opium, U. S. P.—Each fluidounce contains by assay 6 grains of morphine, equivalent to 7 1-2 grains of morphine sulphate.

Dose, equivalent to one grain of powdered opium, 10 minims (0.65 C.c.).

Wine Opium, U. S. P.—Each fluidounce contains by assay 6 grains of morphine, equivalent to 7 1-2 grains of morphine sulphate.

Dose, equivalent to one grain of powdered opium, 10 minims (0.65 C.c.).

ORRIS. *Iris florentina*, Linné. (Rhizome.)*Synonyms.*—Orrice Root, White Flag.*Properties.*—Mild digestive stimulant, slightly cathartic. It has also been employed as a diuretic. It is largely used for its odoriferous properties.**PREPARATION.****Fluid Extract Orris root.**—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

OX-GALL. *Fel Bovis*, U.S.P. The gall or bile of *Bos Taurus*, Linné.*Properties.*—Purgative, antiseptic, tonic. Has been found serviceable in constipation; also in jaundice depending upon catarrh of the bile ducts, partly through its purgative action and by affording aid to the duodenal digestion. Too nauseous to be given in the natural state. The following should be directed to be taken some time after eating and in pill or capsule form, so as to delay admixture of the bile until the food is ready to pass into the duodenum.

OX-GALL—Continued.**PREPARATIONS.****Ox-gall, Desiccated.**—Scales.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Ox-gall, Inspissated, U. S. P.—Prepared by evaporating the fresh gall after straining, to the consistency of a solid extract.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Ox-gall, Inspissated, powdered.—Prepared by drying the above more thoroughly and powdering; suitable for administration in capsules.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Ox-gall, Purified, U. S. P.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Pills.**Ox-gall, 3 grs.**

Ox-gall, 2 grs.

Powd. Jamaica Ginger, 1 gr.

Ox-gall, Craig's, 3 grs.

Ox-gall, 2 grs.

Ext. Coloc. comp., 1 gr.

Ox-gall Compound, 2 1-3 grs.

Ox-gall, 2 grs.

Ext. Stramonium seed, 1-6 gr.

Purified Aloes, 1-10 gr.

Berberine hydrochlorate, 1-12 gr.

Ox-gall and Pepsin, Armor's, 4 1-8 grs.

Purified Aloes, 1 gr.

Iron sulphate, excis., 1-2 gr.

Ext. Nux vomica, 1-12 gr.

Ox-gall, 1 1-2 grs.

Pepsin, 1 gr.

PANCREATIN. Extractive from the pancreas or "sweetbread."

Properties.—It transforms starch into sugar, emulsifies fats, and peptonizes albuminoids. Its beneficial effects are generally best secured by causing it to act on the food before it enters the stomach, since the acid normally present in that organ suspends its activity until it passes into the duodenum, and meanwhile it may be itself digested and lost. Indicated in intestinal dyspepsia, defective nutrition, and convalescence from acute diseases.

Pancreatin, pure.—There is of course no such thing made as a pure pancreatin; this is simply a highly concentrated preparation containing the various digestive ferments of the pancreatic secretion. Five grains with 20 grains of sodium bicarbonate will peptonize a pint of milk in about half an hour at 110° F.

Dose, 1 to 2 grains (0.06 to 0.13 Gm.).

Pancreatin, saccharated.—Diluted with sugar of milk, 1-4 the strength of pure pancreatin.

Dose, 4 to 8 grains (0.26 to 0.5 Gm.).

PREPARATIONS.**Compressed Tablets.****Digestive.**

Pepsin, pure, 1 gr.

Pancreatin, pure, 1 gr.

Calcium lacto-phosphate, 2 grs.

Peptonizing.—Put up in boxes of two vials each.

One of these vials contains 30 tablets of sodium bicarbonate, 10 grs.; and the other 30 tablets of pure pancreatin, 2 1-2 grs., with sugar of milk q. s. ad 10 grs. They are used for preparing peptonized milk, gruel, beef-tea, etc.

Elixirs.—Each fluidounce represents ingredients specified.

Pepsin, Bismuth and Pancreatin.

Saccharated pepsin, 8 grs.

Pancreatin, 8 grs.

Bismuth and Ammon. cit., 8 grs.

Pepsin, Bismuth, Strychnine and Pancreatin.

Saccharated Pepsin, U.S.P., 8 grs. Strychnine, 8-100 gr.

Bismuth and Ammon. cit., 8 grs. Pancreatin, 4 grs.

Pepsin and Pancreatin.

Saccharated Pepsin, U.S.P., 8 grs. Pancreatin, 4 grs.

Liquid Pancreatin, concentrated.—One teaspoonful will peptonize a pint of milk.

Dose, 1-4 to 2 fluidrachms (1 to 8 C.c.).

PANSY. *Viola tricolor*, *Linné*. (Leaves of the wild-grown plant.)

Synonyms.—Heartsease, Eng.; Pensée, Fr.; Freisamkraut, Ger.

Properties.—Mucilaginous, emollient and slightly laxative. Possessing considerable reputation in Europe as a purifier of the blood, and as a remedy in bronchial and intestinal catarrh, and in cutaneous eruptions.

It has been recently demonstrated to be almost a specific in eczema, whether in adults or the crusta lactea of infants.* Also used as an expectorant in pectoral affections.

Pansy leaves, U. S. P., compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Pansy.—Miscible with water.

Dose, for young children, in acute forms of eczema, 1 to 5 minims (0.06 to 0.3 C.c.); in subacute and chronic forms, 10 to 15 minims (0.6 to 1 C.c.); for adults, 1-2 to 2 fluidrachms (1 to 4 C.c.). Administered on an empty stomach, and half an hour before meals if possible. When used in cases of eczema, if aggravation occurs, the remedy should be discontinued for a few days and resumed in smaller doses.

PAPAW (American). *Asimina triloba*, *Dunal*. (Seeds.)

Synonym.—Custard Apple.

Properties.—Said to be useful in asthma, and also employed as an emetic.

PREPARATION.

Fluid Extract Papaw seeds.—Not miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

PARAGUAY TEA. *Ilex paraguayensis*, *St. Hilaire*. (Leaves.)

Synonym.—Yerba Maté.

Properties.—Extensively consumed as a beverage in South America. The leaves are diuretic, exhilarant and, in large doses, emetic and drastic. The peculiar characteristic properties of this drug depend on the caffeine it contains. It contains a larger proportion of mineral salts than either tea or coffee. It does not exalt the peripheric nerves like tea, nor the cerebriac like coffee, but rather induces indolence and drowsiness; it accelerates the cardiac contractions and peristaltic movements.

PREPARATION.

Fluid Extract Paraguay Tea.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

PAREIRA BRAVA, U.S.P. *Chondodendron tomentosum*, *Ruiz & Pavon*. (Root.)

Properties.—Tonic, slightly aperient, and diuretic. Very useful in chronic cystitis, in allaying the irritability of the bladder, and correcting the disposition to profuse mucous secretion. It has now an established reputation in the treatment of all chronic affections of the urinary tract, especially those attended with muco-purulent deposits in the urine.

Pareira Brava root, U.S.P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixir Buchu and Pareira Brava.—Each fluidounce represents:

Buchu, 30 grs.

Juniper berries, 15 grs.

Pareira brava, 7 1-2 grs.

Stoneroot, 7 1-2 grs.

Extracts.

Fluid Extract Pareira Brava, U. S. P.—Not miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

* Piffard, H. G., "The Modern Treatment of Eczema" (Edition of Geo. S. Davis).

PAIREIRA BRAVA—Continued—(Extracts).

Fluid Extract Buchu and Pareira Brava.—Not miscible with water.

Sixteen fluidounces, imperial, represents: Buchu, 8 ozs. av.; Pareira brava and Stone root, 2 ozs. each, and Juniper berries, 4 ozs.

An efficacious combination in various diseases of the genito-urinary system. It has been found useful in leucorrhœa, dropsy, chronic inflammation of the bladder and urinary passages, etc.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

PARSLEY. *Petroselinum sativum*, *Hoffmann*. (Leaves, root, seed.)

Synonyms.—*Apium petroselinum*, *Linne'*; Persil, Fr.; Petersilie, Ger.

Properties.—Aperient and diuretic; often used in nephritic and dropsical affections. The root is the most diuretic part of the plant, and it is also prescribed for the relief of strangury produced by cantharides, and for the dysuria attending attacks of gravel. Parsley seed is sometimes used as a substitute for quinine in marsh miasm, and apiol, its essential oil, has been found of service in intermittent neuralgia.*

The most important use of the seed (or of apiol) is as an emmenagogue in amenorrhœa and dysmenorrhœa. In these affections the fluid extract should be taken night and morning, in doses of half a fluidrachm, in anticipation of the menstrual period, and in the latter affection during its continuance.

Parsley leaves, compressed in ounce packages for retailing purposes.

Parsley root, compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLE.

Apiol.—In small doses, diuretic and stimulant to the circulation. In full doses of 10 to 15 minims (0.6 to 1 C. c.) it is a very efficient emmenagogue in amenorrhœa and dysmenorrhœa and also in cases of fetid menstruation.

PREPARATIONS.**Capsules.**

Apiol.—In hard gelatin.

Apiol, 4 minims.

Olive oil, 6 minims.

Apiol.—In soft gelatin.

Apiol, 5 minims.

Olive oil, 5 minims.

Extracts.

Fluid Extract Parsley root.—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Parsley seed.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

PEACH. *Amygdalus persica*, *Linne'*. (Bark, leaves.)

Synonym.—*Persica vulgaris*, *DeCandolle*.

Properties.—Sedative, from the presence of hydrocyanic acid, mildly laxative, diuretic and anthelmintic. Recommended in morbid irritability of the bladder and urethra, ischuria, hæmaturia, in pertussis, in inflammation of the stomach and bowels, and for the relief of nausea. Used to allay the pain of nephritic colic, to promote the discharge of calculi by causing diuresis, and as a remedy in intermittent fevers. Poisonous effects may be produced by excessive doses, owing to the prussic acid, or amygdalin contained.

Peach bark, compressed in pound packages for retailing purposes.

Peach leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Peach leaves.—Not miscible with water.

Dose, 10 to 25 minims (0.6 to 1.5 C. c.).

* Napheys, "Medical Therapeutics," p. 108.

PELLITORY. *Pyrethrum*, U.S.P. *Anacyclus Pyrethrum*, *DeCandolle*. (Root.)

Synonyms.—Spanish Chamomile; *Pyrèthre*, Fr.; *Bertramwurzel*, Ger.

Properties.—A powerful irritant, used almost exclusively as a sialagogue in certain forms of headache, rheumatic and neuralgic affections of the face, or as a local stimulant in palsy of the tongue and throat, and in relaxation of the uvula. An excellent remedy for toothache. Used by dentists as a local application to carious teeth, with a view to its benumbing effect before plugging.

Pellitory root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATION AND FORMULA.

Fluid Extract Pellitory.—Not miscible with water without some precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.), to be held in the mouth but not swallowed.

Tincture Pellitory, U. S. P.

Fluid Extract Pellitory, P., D. & Co.'s.....	2 fl. ozs.
Alcohol.....	9 1-2 fl. ozs.

PENCIL FLOWER. *Stylosanthes elatior*, *Swartz.* (Herb.)

Properties.—This plant has long been employed in domestic practice to relieve the irritability of the uterus, and consequent abnormal pains, liable to occur during the latter months of gestation. It is also said to give tone to the uterine fibre, thus increasing expulsive effort, and facilitating parturition. It may thus be classed as a uterine sedative and tonic.

PREPARATION.

Fluid Extract Pencil Flower (Stylosanthes).—Not miscible with water.

Dose, 10 to 20 minims (0.6 to 1.3 C.c.) three times a day prior to confinement.

PENNYROYAL (American). *Hedeoma*, U. S. P. *Hedeoma pulegioides*, *Persoon.* (Leaves.)

Synonyms.—Tick Weed, Squaw mint; *Herbe de pouliot*, Fr.; *Amerikanisches Polei*, Ger.

Properties.—Stimulant, diaphoretic and emmenagogue. Used chiefly in the form of a hot infusion (1 ounce to the pint of boiling water) in threatened inflammation of the throat and bronchia, in muscular rheumatism, and especially in connection with hot hip and foot baths as an efficient and safe emmenagogue.

Pennyroyal leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Capsules Oil Pennyroyal, 1 and 2 minims; in soft gelatin.

Fluid Extract Pennyroyal.—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

PEONY. *Pæonia officinalis*, *Linné.* (Root.)

Properties.—Much employed at one time in epilepsy, as an emmenagogue and, from its astringency, also in diarrhoea. Very little information of a modern nature is to be found regarding it, but it is occasionally used as an antispasmodic and nervine sedative in epilepsy, chorea and also in whooping-cough.

Dose, 1-2 to 2 fluidounces (15 to 60 C.c.) of an ounce to the pint infusion.

Peony root, compressed in pound packages for retailing purposes.

PEPPERMINT. *Mentha piperita*, Linné. (Leaves and tops)

Properties.—Diffusible stimulant and antispasmodic. Used in flatulent colic and to check nausea and vomiting.

Peppermint leaves, U.S.P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Peppermint leaves.—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Pill Peppermint Oil, 1-128 gr.

PEPSIN. An article representing the active principle of the gastric juice.

Properties.—Pepsin is that digestive ferment which dissolves albuminoids such as white of egg, meat fibrin, etc. It differs from pancreatin in not being able to act upon starchy or fatty substances, and in being active only in an acid solution instead of an alkaline one. Used as a substitute for the natural digestive fluid in atony of the stomach from functional derangement or organic disease.

Pepsin, lactated.—This contains all of the principal agents contributing to the process of digestion in its several stages. Each dose of 5 grains contains: Pepsin, pure (equal to 5 grs. Sacch. Pepsin, U. S. P.), 1-2 gr.; Pancreatin, saccharated, 1-2 gr.; Lactic Acid, Maltose, Diastase, Hydrochloric acid.

Pepsin, pure (pig).—One grain will dissolve from 450 to 500 grains coagulated albumen under conditions of U. S. P. test.
Dose, 1-2 grain (0.03 Gm.) and upwards.

Pepsin saccharated, commercial.—One grain will dissolve 12 1-2 grains coagulated albumen.

Pepsin Saccharated, U. S. P.—One grain will dissolve 50 grains coagulated albumen.
Dose, 5 to 10 grains (0.3 to 0.6 Gm.).

Pepsin, Sheep, pure.—One grain will dissolve from 350 to 400 grains coagulated albumen. For certain forms of dyspepsia and in the treatment of infants, Sheep Pepsin has been deemed superior to that of the hog. This preference has originated because the food of the sheep is wholly vegetable, which fact seems to render the action of the pepsin more specific.
Dose, 1-2 grain (0.03 Gm.) and upwards.

Pepsin, Sheep, saccharated.—Of the same strength as the U. S. P. Sacch. Pepsin.

PREPARATIONS.**Compressed Tablet, Digestive.**

Pepsin, pure, 1 gr.

Calcium lacto-phosphate, 2 grs.

Pancreatin, 1 gr.

Elixirs.—Names only; want of space prevents all formulæ being given here, and a selection would be difficult to make where all are equally valuable. For this information see Elixir Formulæ list appended.

Calisaya, Iron, Bismuth and Pepsin.

Calisaya, Iron, Bismuth, Pepsin and Strychnine.

Calisaya, Iron and Pepsin.

Calisaya, Iron, Strychnine and Pepsin.

Calisaya and Lactated Pepsin.

Calisaya, Pepsin and Bismuth.

Calisaya, Pepsin, Bismuth and Strychnine.

Calisaya, Pepsin and Strychnine.

Gentian, Iron chloride and Lactated Pepsin.

Gentian and Lactated Pepsin.

Iron, Quinine, Strychnine and Pepsin.

Iron, Quinine and Strychnine phos. with Lactated Pepsin.

Lactated Pepsin.

Lactated Pepsin and Bismuth.

PEPSIN—Continued—(Elixirs.)

Lactated Pepsin, Bismuth and Strychnine.
Lactated Pepsin, Bismuth, Strychnine and Iron.
Lime-juice and Pepsin.
Pepsin.
Pepsin and Bismuth.
Pepsin, Bismuth and Iron.
Pepsin, Bismuth, Iron and Quinine.
Pepsin, Bismuth and Nux vomica.
Pepsin, Bismuth and Pancreatin.
Pepsin, Bismuth and Quinine.
Pepsin, Bismuth, Strychnine and Pancreatin.
Pepsin, Bismuth and Wafer Ash.
Pepsin, Damiana, Phosphorus and Nux vomica.
Pepsin and Pancreatin.
Pepsin and Quinine.
Pepsin and Strychnine.
Pepsin, Strychnine and Bismuth.
Pepsin, Strychnine and Bismuth with Iron.
Pepsin, Strychnine and Iron.
Pepsin and Wafer Ash.

Glyceroles.—Each fluidounce represents ingredients specified.

Pepsin, conc., 480 grs. Pepsin sacch., U. S. P.

Pepsin and Wafer Ash.

Pepsin, sacch., U. S. P., 10 grs. Wafer Ash, 30 grs.

Granular Effervescent Preparations.—Ingredients stated per drachm.

Pepsin, Saccharated, U. S. P., 5 grs.

Pepsin and Bismuth.

Pepsin, saccharated, 5 grs. Bismuth and Ammon. cit., 1 gr.

Pepsin, Bismuth and Iron.

Pepsin, 5 grs. Iron citrate, 1 gr.

Bismuth citrate, 2 grs.

Pepsin, Bismuth and Strychnine.

Pepsin, saccharated, 5 grs. Strychnine, 1-60 grs.

Bismuth and Ammon. cit., 2 grs.

Liquid Pepsin, U. S. P., represents 18 grains Sacch. Pepsin, U. S. P., per fluidounce
 Dose, 1-2 to 1 fluidounce (15 to 30 C. c.).

Lozenges.

Pepsin, 3 grs.

Pepsin and Bismuth, 4 grs.

Sacch. Pepsin, U.S.P., 2 grs. Bismuth subnitrate, 2 gr.

Pepsin and Iron, 3 grs.

Sacch. Pepsin, U.S.P., 1 1-2 grs. Iron pyrophosphate, 1 1-2 grs.

Pepsin and Lime lactophosphate, 5 grs.

Sacch. Pepsin, U.S.P., 2 grs. Calcium lactophos., 3 grs.

Malt Extract Combinations.—Ingredients stated per fluidounce.

Malt Extract with Lactophosphates and Pepsin.

Calcium lactophos., 6 grs. Sodium lactophos., 2 grs.

Iron lactophos., 4 grs. Potassium lactophos., 2 grs.

Pepsin, sacch., U.S.P., 8 grs. Manganese lactophos., 2 grs.

Malt Extract with Pepsin.

Pepsin, sacch., U.S.P., 4 grs.

Malt Extract with Pepsin and Bismuth.

Pepsin, sacch., U.S.P., 4 grs. Bismuth and Ammon. cit., 4 grs.

Malt Extract with Pepsin and Pancreatin.

Pepsin, sacch., U.S.P., 4 grs. Pancreatin, sacch., 4 grs.

Malt Extract with Pepsin and Strychnine.

Pepsin, sacch., U.S.P., 4 grs. Strychnine, 4-100 gr.

PEPSIN—Continued.**Pills.****Digestive Pellets,** Hager's, 3 1-2 grs.

Cinchonidine sulph., 1-5 grs.

Pepsin, 1 1-5 grs.

Powd. Ginger, 3-25 gr.

Powd. Cardamom, 3-25 gr.

Powd. Pimento, 3-25 gr.

Powd. Gentian root, 6-25 gr.

Powd. Marshmallow root, 6-25 gr.

Powd. Tragacanth, 6-25 gr.

Glycerin, 2-5 gr.

Acid hydrochloric, 2-5 gr.

Headache, 3 grs.

Lactated Pepsin, 1 gr.

Guarana, 1-2 gr.

Sodium bicarb., 1 gr.

Cypripedin, 1-2 gr.

Night-sweat, 3 grs.

Zinc oxide, 1-2 gr.

Salicin, 1 gr.

Ext. Belladonna, 1-25 gr.

Hydrastin, 1 gr.

Lactated Pepsin, 1-2 gr.

Ox-gall and Pepsin, Armor's, 4 1-8 gr.

Purified Aloes, 1 gr.

Iron sulph. exsic., 1-2 gr.

Ext. Nux vomica, 1-12 gr.

Ox-gall, 1 1-2 grs.

Pepsin, 1 gr.

Pepsin and Iron by hydrogen, 2 1-4 grs.

Pepsin, 1 1-2 grs.

Iron by hydrogen, 3-4 gr.

For others containing pepsin see Pill Formulæ list appended for numbers 251, 295-297, and 299.

Syrups.—Ingredients stated per fluidounce.**Acid Phosphate with Pepsin.**

Sacch. Pepsin, U. S. P., 40 grs.

Liquid acid phosphate, 2 drachms.

Calcium lactophosphate and Pepsin.

Calcium lactophosphate, 16 grs.

Pepsin, sacch., U.S.P., 16 grs.

Lactated Pepsin. 40 grs.**Lactated Pepsin Compound.**

Lactated pepsin, 24 grs.

Calcium phosphate, 8 grs.

Iron phosphate, 8 grs.

Potassium phosphate, 8 grs.

Sodium phosphate, 8 grs.

Tablet Triturates, Pepsin, 1-4, 1-2 and 1 gr.**Wines.**—Each fluidounce represents ingredients specified.**Beef, Iron and Wine with Pepsin.**

Beef, 2 ozs.

Sacch. Pepsin, U.S.P., 8 grs.

Iron and Ammon. cit., 4 grs.

Pepsin, Sacch., U. S. P., 10 grs.**Pepsin and Bismuth.**

Sacch. Pepsin, U.S.P., 8 grs.

Bismuth and Ammon. cit., 8 grs.

PERSIMMON. *Diospyros virginiana, Linné.* (Bark, green fruit.)

Properties.—Astringent; successfully employed in diarrhœa, dysentery, and uterine hæmorrhage, as a gargle in ulcerated sore throat and as a wash in stomatitis. The bark is claimed to be used advantageously in intermittents.

Persimmon bark, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Persimmon bark.—Miscible with water without material precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Persimmon fruit (green).—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

PICHI. *Fabiana imbricata, Ruiz & Pavon.* (Stems and leafy branchlets.)

Properties.—Diuretic, tonic, terebinthinate. "In cases of vesical catarrh, acute or chronic, following a mechanical cause, such as gravel or a uric diathesis, this remedy will quickly modify the urinary secretions, calm the irritability, and favor the expulsion of the gravel and calculi that can be passed through the urethra. It modifies and cures also chronic p ur-

PICHI—Continued.

lent mucous secretions. Its action on the liver must be attributed to its diuretic properties, though it is recommended for icterus, hydropsy, and dyspepsia due to biliary secretions. In this case the essential oil, when absorbed and carried into circulation, acts like a stimulant of the secretory apparatus in general.* Contra-indicated where there is organic kidney disease.

PREPARATION.

Fluid Extract Pichi.—Not miscible with water unless solution is made alkaline.

Dose, 10 to 40 minims (0.6 to 2.6 C.c.).

PIMENTO. Pimenta, U.S.P. *Eugenia Pimenta, DeCandolle.* (Fruit, leaves.)

Synonyms.—Allspice, Jamaica pepper.

Properties.—Carminative and stimulant. An agreeable addition to diarrhoea mixtures, and useful as a stomachic in atonic dyspepsia and flatulency. The fruit is largely used as a condiment.

PREPARATIONS.

Fluid Extract Pimento fruit.—Not miscible with water.

Dose, 10 to 40 minims (0.6 to 2.5 C.c.).

Fluid Extract Pimento leaves.—Not miscible with water without some precipitation.

Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

PINK-ROOT. *Spigelia, U.S.P. Spigelia marilandica, Linné.*

Synonyms.—Worm-grass, Carolina Pink.

Properties.—Pink-root is a powerful, anthelmintic. In overdoses it excites the circulation, and has a decided narcotic action, which may be counteracted by combining it with some cathartic, such as senna, etc.

Pink-root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Pink-root, U. S. P.—Miscible with water without material precipitation.

Dose, adults, 1 to 2 fluidrachms (4 to 8 C.c.); children, 10 to 20 minims (0.6 to 1.3 C.c.).

Fluid Extract Pink-root Compound.—Not miscible with water without some precipitation.

Sixteen fluidounces, imperial, represents: Pink-root and Senna, 7 ozs. each; Savin and Manna, 1 oz. each. An efficacious combination as a specific for worms; combining both anthelmintic and laxative properties.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Pink-root and Senna, U. S. P., 1870.—Miscible with water.

Sixteen fluidounces, imperial, represents: Pink-root, 10 ozs. av.; Senna, 6 ozs.; Anise and Caraway oils, 20 minims each. An excellent vermifuge, combining the anthelmintic action of Pink-root with the cathartic action of Senna.

Dose, for a child two years old, 30 to 60 minims (2 to 4 C.c.); for an adult, 2 to 4 fluidrachms (8 to 15 C.c.).

PIPSISSEWA. *Chimaphila, U. S. P. Chimaphila umbellata, Nuttall.* (Leaves.)

Synonyms.—Prince's Pine, Wintergreen, Ground Holly; Pyrole ombellée, Fr.; Doldenblütiges Harnkraut, Ger.

Properties.—Astringent, alterative, tonic and diuretic. Useful in scrofula, and in chronic rheumatic and nephritic affections; can be used as a substitute for uva ursi in urinary disorders, as it is less obnoxious to the stomach.

* Les Nouveaux Remèdes, Feb. 15, 1886; Medical Age, March 10, 1886, p. 118.

PIPSISSEWA—Continued.

Pipsissewa leaves, U.S.P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Chimaphilin, a so-called **Concentration**,*—A powdered preparation from Pipsissewa.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

Elixir Stillingia Comp.—Each fl. oz. represents ingredients specified.

Stillingia, 30 grains.

Turkey corn, 30 grs.

Pipsissewa, 15 grs.

Elder flowers, 15 grs.

Blue Flag, 15 grs.

Prickly ash berries, 7 1-2 grs.

Coriander seed, 7 1-2 grs.

Extracts.

Fluid Extract Pipsissewa, U. S. P.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Stillingia Comp.—Not miscible with water.

Formula in same proportions as Elixir Stillingia compound above, but four times stronger.

Solid Extract Pipsissewa.—One part equals 4 of drug.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Pill Chimaphila Extract, 3 grs.

Syrup Stillingia Compound.—Formula same as Elixir above.

PITCHER-PLANT. *Sarracenia purpurea, Linné.* (Rhizome.)

Synonyms.—Side-saddle Flower, Fly-trap, Huntsman's Cup, Water Cup; *Sarracenie, Fr., Ger.*

Properties.—"Stimulating diuretic."† A stimulant tonic to the mucous surfaces and also to the system generally. It is effective in diarrhoea, dyspepsia and gastralgia. Used in sluggish or torpid conditions of the stomach, intestines, liver, kidneys, or the uterus, which produce costiveness, dyspepsia, sick-headache, amenorrhoea and dysmenorrhoea.

Pitcher-plant root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Pitcher-plant.—Not miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

PLANTAIN. *Plantago major, Linné.* (Leaves.)

Synonym.—Grosser wegetritt, Ger.

Properties.—Alterative, diuretic, refrigerant, and hemostatic. Prescribed in diarrhoea, dysentery, hæmorrhoids, hæmaturia, menorrhagia, etc. A reputed antidote to bites of venomous serpents, spiders, etc. It is valuable as a remedy in ascites.

Plantain leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Plantain leaves.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

PLEURISY-ROOT. *Asclepias, U.S.P. Asclepias tuberosa, Linné.* (Root.)

Synonym.—Butterfly Weed.

Properties.—Diaphoretic, antispasmodic, tonic, diuretic and carminative. It has obtained a reputation in diseases of the respiratory organs, more particularly in pleurisy, inflammation of the lungs and catarrhal affections. Beneficial in acute rheumatism, in febrile diseases, and in flatus of adults and children.

Pleurisy-root, U. S. P., compressed in pound packages for retailing purposes,

* See note in Preface on Concentrations.
 † Lescher, "Recent Materia Medica," p. 59.

PLEURISY-ROOT—Continued.

PREPARATIONS.

Asclepidin, a so-called **Concentration**,* a powdered preparation from Pleurisy-root.

Dose, 1 to 5 grains (0.065 to 0.33 Gm.).

Fluid Extract Pleurisy-root.—Not miscible in water without some precipitation.

Dose, 20 to 60 minims (1.3 to 4 C.c.).

POISON OAK. *Rhus Toxicodendron*, Linné. (Leaves.)

Synonyms.—Poison Ivy, Sumach vénéneux, Fr.; Giftsumach, Ger.

Properties.—POISONOUS. Stimulant narcotic, producing, when swallowed, more or less irritation of the stomach and bowels, and promoting the secretory functions of the skin and kidneys, and producing a stupefying effect upon the nervous system. Successfully used in the treatment of obstinate cutaneous disorders, dropsy, nocturnal incontinence, chronic paralysis, chronic rheumatism, and some diseases of the eye. It is said it produces twitchings of paralyzed muscles, and prickings of the affected limb, similar to the effects of strychnine or nux vomica.

Antidotes.—Use emetics followed by demulcents and alkalies, and treat symptoms as they arise on general principles.

Poison Oak leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Extracts.

Fluid Extract Poison Oak.—Not miscible with water without some precipitation.

Dose, 3 to 30 minims (0.2 to 2 C.c.).

Powdered Extract Poison Oak.—One part equals 5 of drug.

Dose, 1-2 to 5 grains (0.03 to 0.33 Gm.).

"German Tincture" Poison Oak.—An imported unofficial tincture of the green drug.

Dose, 2 to 10 minims (0.12 to 0.6 C.c.).

POKE-ROOT. { *Phytolaccæ Bacca*, U.S.P. } *Phytolacca decandra*,
{ *Phytolaccæ Radix*, U.S.P. } Linné. (Berries, root.)

Synonyms.—Garget, Pigeon Berry.

Properties.—Emetic, cathartic, alterative and somewhat narcotic. Has acquired reputation as a remedy for syphilitic and rheumatic affections; is also recommended in acute and chronic catarrhal affections of the mucous membranes, and in headache, dysmenorrhœa, passive congestion of the uterus, liver, spleen and other organs.

Poke-root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Extracts.

Fluid Extract Alterative Compound.—Miscible with water.

Synonyms.—Bamboo Brier Comp., Succus Alterans, Mist. Smilacis Comp.

Seventeen fluidounces, imperial, represents: Bamboo Brier, *Stillingia*, Burdock, and Poke roots, 4 ozs. av. each; Prickly Ash bark, 1 oz.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.), in water.

NOTE.—Syrup Trifolium Comp. is rapidly supplanting this formula as an alterative; see below.

Fluid Extract Poke berries.—Miscible with water.

Dose, 10 minims to 1 fluidrachm (0.6 to 4 C.c.), according to the effect desired.

Fluid Extract Poke root.—Miscible with water.

Dose, as an alterative, 2 to 10 minims (0.13 to 0.65 C.c.); as an emetic, 20 to 30 minims (1.3 to 2 C.c.).

* See note in Preface on Concentrations.

POKE-ROOT—Continued—(Extracts).

Powdered Extract Poke root.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Poke-root.
Dose, 1 to 5 grains (0.06 to 0.33 Gm.).

Solid Extract Alternative Comp.—One grain is equivalent to 5 minims Fluid Extract Alternative comp., see above for formula.
Dose, 10 to 40 grains (0.65 to 2.6 Gm.).

Solid Extract Poke-root.—One part equals 4 of root.
Dose, 1 to 5 grains, (0.06 to 0.3 Gm.).

Phytolaccin, a so-called **Concentration**,* a powdered preparation from Poke-root.

Pills.

Alternative Compound, 3 grs.

Ext. Smilax Sarsap., 2-3 gr.

Ext. Stillingia sylvat., 2-3 gr.

Ext. Burdock, 2-3 gr.

Ext. Poke-root, 2-3 gr.

Ext. Prickly ash bark, 1-3 gr.

Phytolaccin, 1-2 gr.

Syrups.—Each fluidounce represents ingredients specified.

Alternative Compound.

Two drachms Fluid Extract Alternative comp., see above for formula.
Dose, 2 fluidrachms (8 C. c.) and upwards.

Trifollum Compound.

Recommended by Dr. Rush as superior in efficacy to the combination of alternatives that has recently acquired considerable notoriety under the name of bamboo brier root compound, succus alterans or alternative compound. Its restorative action in secondary syphilis is prompt and unequivocal.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.) three times a day.

Red clover, 32 grs.

Stillingia, 16 grs.

Burdock root, 16 grs.

Poke-root, 16 grs.

Berberis aquifolium, 16 grs.

Cascara amarga, 16 grs.

Prickly-ash bark, 4 grs.

Potassium iodide, 8 grs.

POLYPODY. *Polypodium vulgare*, Linné. (Leaves.)

Synonyms.—Rock brake; Gemeiner Tüpfelfarn, Ger.

Properties.—Expectorant, demulcent, purgative and anthelmintic.

Dose, 2 fluidounces (60 C. c.) of an ounce to the pint infusion.

Polypody leaves, compressed in ounce packages for retailing purposes.

POMEGRANATE. *Granatum*, U. S. P. *Punica Granatum*, Linné.
(Bark of root.)

Properties.—The bark of the pomegranate root has long been known as an efficient remedy for tape-worm. Without the aid of castor oil or other cathartic it effects the destruction and expulsion of the parasite. Should it not have a purgative effect a brisker cathartic than castor oil is advisable, so that the tænia may not be allowed time to recover from the effects of the pomegranate. Its virtues are due to the alkaloid pelletierine.

Pomegranate bark of root, U. S. P., compressed in pound packages for retailing purposes.

ACTIVE PRINCIPLE.

Pelletierine Sulphate.—A remedy for tape-worm.

Dose, as a teninuge, 5 to 12 grains (0.33 to 0.76 Gm.).

Pelletierine Tannate.—An efficient remedy for tape-worm, and commonly used in preference to the sulphate.

Dose, 10 to 25 grains (0.65 to 1.66 Gm.).

PREPARATION.

Fluid Extract Pomegranate bark of root.—Miscible with water.

Dose, 1-4 to 2 fluidrachms (1 to 8 C. c.).

* See note in Preface on Concentrations.

PONGAMIA GLABRA, *Ventenat.* (Fixed oil from the seeds.)

Properties.—Has long been employed by the natives of India as a local application for scabies, herpes, and sundry other affections of the skin and for the relief of rheumatism. "The oil is an excellent remedy in pityriasis versicolor and porrigo capitis."*

POPPY. *Papaver somniferum, Linné.* (Capsules, leaves.)

Properties.—Anodyne, calming irritation and producing sleep. Used in pulmonary affections. It is a favorite ingredient in soothing syrups for children. For other products from the Poppy, see Opium.

Poppy leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Poppy-heads.—Not miscible in water without material precipitation.

Dose, 10 to 40 minims (0.65 to 2.6 C.c.).

Solid Extract Poppy-heads.—One part equals 7 of drug.

Dose, 2 to 5 grains (0.13 to 0.3 Gm.).

Pill Poppy Extract, 2 grs.

Syrup Poppy-heads, B. P.

Fluid Extract Poppy-heads, P., D. & Co.'s..... 4 fl. ozs.

Syrup 11 fl. ozs.

Evaporate the fluid extract by a gentle heat to two fluidounces; add the syrup and mix thoroughly.

Dose, 1-2 to 2 drachms (2 to 8 C.c.).

PRICKLY ASH. *Xanthoxylum, U. S. P. Xanthoxylum americanum, Miller.* (Bark, berries.)

Synonyms.—Suter berry, toothache tree, clavalier, Fr.; Zahnwehholz, Ger.

Properties.—Stimulant, tonic, alterative, diaphoretic, sialagogue, carminative, emmenagogue and lactagogue. It is used as a stimulant in languid states of the system, and is said to have proved highly beneficial in chronic rheumatism, flatulent colic, syphilitic and hepatic affections.

Prickly Ash bark, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Corydalis Compound.

Turkey corn, 30 grs.

Stillingia, 30 grs.

Blue flag, 40 grs.

Prickly ash, 15 grs.

Potassium iodide, 8 grs.

Stillingia Compound.

Stillingia, 30 grs.

Turkey corn, 30 grs.

Pipsissewa, 15 grs.

Elder flowers, 15 grs.

Blue flag, 15 grs.

Prickly ash berries, 7 1-2 grs.

Coriander seed, 7 1-2 grs.

Extracts.

Fluid Extract Alterative Comp.—Miscible with water.

Synonyms.—Bamboo Brier Comp., Succus Alterans, Mist. Smilacis Comp.

Seventeen fluidounces, imperial, represents: Bamboo Brier, Stillingia, Burdock and Poke roots, 4 ozs. av. each; Prickly ash bark 1 oz.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

NOTE.—Syrup Trifolium Comp. is rapidly supplanting this formula as an alterative, see below.

Fluid Extract Prickly Ash bark, U. S. P.—Not miscible with water without some precipitation.

Dose, 10 to 40 minims (0.6 to 2.6 C.c.).

Fluid Extract Prickly Ash berries.—Not miscible with water without some precipitation

Dose, 5 to 10 minims (0.33 to 0.65 C.c.).

* Dymock, "Materia Medica of Western India, p. 197.

PRICKLY ASH—Continued—(Extracts).**Fluid Extract Stillingia Comp.**—Not miscible with water.

Formula in same proportion as Elixir Stillingia Comp. above, but four times stronger.

Dose, 30 to 60 minims (2 to 4. C.c.).

Solid Extract Alternative Comp.—One grain is equivalent to 5 minims Fluid Extract Alternative Comp., see above.

Dose, 10 to 40 grains (0.65 to 2.6 Gm.).

Solid Extract Prickly Ash bark.—One part equals 5 of bark.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Pills.**Ague, Improved,** 2 1-4 grs.

Cinchonidine sulph., 1 gr.

Xanthoxylin, 1 gr.

Gelsemerin, 1-16 gr.

Powd. Capsicum, 1-16 gr.

Alternative Compound, 3 grs.

Ext. Smilax Sarsap., 2-3 gr.

Ext. Stillingia sylvat., 2-3 gr.

Ext. Burdock, 2-3 gr.

Ext. Poke root, 2-3 gr.

Ext. Prickly ash bark, 1-3 gr.

Cascara Compound Pellets, D'Ary's, 1 gr.

Ext. Cascara sagrada, 4-15 gr.

Ext. Nux vomica, 1-30 gr.

Ext. Belladonna, 1-60 gr.

Euonymin, 1-5 gr.

Xanthoxylin, 4-15 gr.

Oleoresin Capsicum, 1-20 gr.

Tonic, Waxham's, 2 grs.

Cinchonine sulph., 1 gr.

Ext. Prickly-ash bark, 1-4 gr.

Ext. Cinchona bark, red, 1-4 gr.

Ext. Dogwood bark, 1-4 gr.

Powd. Capsicum, 1-4 gr.

Syrups.—Each fluidounce represents ingredients specified.**Alternative Compound.**Two drachms Fluid Extract Alternative Comp., see above for formula.
Dose, 2 fluidrachms (8 C.c.) and upwards.**Trifolium Compound.**Recommended by Dr. Rush as superior in efficacy to the combination of alternatives known as alternative compound or succus alterans. Its restorative action in secondary syphilis is prompt and unequivocal.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.) three times a day.

Red clover, 32 grs.

Stillingia, 16 grs.

Burdock root, 16 grs.

Poke root, 16 grs.

Berberis aquifolium, 16 grs.

Cascara amarga, 16 grs.

Prickly-ash bark, 4 grs.

Potassium iodide, 8 grs.

Stillingia.

Stillingia, 88 grs.

Prickly-ash berries, 44 grs.

Stillingia Compound.—Formula same as Elixir Stillingia Comp., see above.**Xanthoxylin,** a so-called **Concentration,*** a powdered preparation from Prickly-ash bark.

Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

PRIDE OF CHINA. Azedarach, U. S. P. *Melia Azedarach, Linné.*
(Bark of root.)**Properties.**—Cathartic and emetic, and in large doses is said to produce narcotic effects similar to those of Pink-root. It is also considered equally efficient as an anthelmintic for lumbricoid ascarides.**PREPARATION.****Fluid Extract Pride of China.**—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

PRIVET. *Ligustrum vulgare, Linné.* (Leaves.)**Properties.**—Astringent, valuable in chronic bowel complaints; ulceration of stomach and bowels; as a gargle for ulcers of mouth and throat, and as an injection for ulcerated ears with offensive discharges.

Dose, 2 to 4 ounces of an ounce to the pint infusion.

Privet leaves, compressed in ounce packages for retailing purposes.

* See note in Preface on Concentrations.

PULSATILLA, U. S. P. *Anemone Pulsatilla, Linné.* (Herb.)

Synonyms.—Pasque flower; Coquelarde, Fr.; Küchenschelle, Ger.

Properties.—Alterative and antispasmodic. Administered with benefit in the secondary manifestations of syphilis, and in chronic cutaneous eruptions of a non-specific nature. It is also highly esteemed as a remedy in chronic catarrhal affections of the mucous membrane wherever located. Its antispasmodic properties make it an important drug for the relief of the "nervousness" of exhaustion due to any cause, and especially in the variety found in women as a result of menstrual derangements and sedentary habits. In functional amenorrhœa its stimulating action on the mucous lining of the uterus makes it valuable. It has been found useful also in the various nervous manifestations of hysteria.

Pulsatilla herb, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Pulsatilla.—Miscible with water without material precipitation.

Dose, 1 to 5 minims (0.065 to 0.33 C.c.).

Powdered Extract Pulsatilla.—One part equals 5 of drug.

Dose, 1-4 to 1 grain (0.016 to 0.064 Gm.).

"German Tincture" Pulsatilla.—An imported unofficial tincture of the green drug.

Dose, 3 to 10 minims (0.18 to 0.6 C.c.).

PUMPKIN. *Pepo, U.S.P. Cucurbita Pepo, Linné.* (Seed.)

Properties.—Pumpkin seeds are said to be a specific for tape-worm; they are also mucilaginous and diuretic. Are recommended in the treatment of strangury and urinary affections; also in gastritis, enteritis, febrile diseases and gonorrhœa. When given for tape-worm, follow after two hours with a brisk cathartic.

PREPARATIONS.

Fluid Extract Pumpkin seed.—Miscible with water.

Dose, 1-2 to 2 fluidounces (15 to 60 C.c.).

For tape-worm, evaporate, at a gentle heat, two fluidounces of the fluid extract to one-half fluidounce; let the patient take this at one dose, fasting, in the morning.

Solid Extract Pumpkin seed.—One part equals 20 of the seed.

Dose, 15 to 30 grains (1 to 2 Gm.).

An efficient tæniifuge. May be combined with oleoresin of male fern or with other anthelmintics.

PURLANE. *Portulaca oleracea, Linné.* (Herb.)

Properties.—Refrigerant, diuretic. Indicated in chronic catarrhal affections of the genito-urinary tract. It is also recommended in scurvy, and is thought by some to possess anthelmintic properties. "Purslane claims a place in the treatment of ascites, and as a mild and efficient diuretic, fills it well." *

PREPARATION.

Fluid Extract Purslane.—Miscible with water.

Dose, 1 to 3 fluidrachms (4 to 12 C.c.).

QUASSIA, U. S. P. *Picræna excelsa, Lindley.* (The wood.)

Synonyms.—Quassia excelsa, Swartz; Bitter Wood, Bitter Ash; Bois amer, Fr.; Quassienholz, Ger.

Properties.—A purely bitter tonic, which invigorates the system without increasing the action of the heart. It is particularly adapted to that debilitated state of the digestive organs which sometimes succeeds acute diseases.

* Morse, "New Therapeutical Agents," p. 57.

QUASSIA—Continued.**PREPARATIONS AND FORMULA.****Extracts.****Fluid Extract Quassia, U. S. P.**—Miscible with water.

Dose, 5 to 15 minims (0.3 to 1 C.c.).

Powdered Extract Quassia.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Quassia.

Dose, 1 grain (0.6 Gm.) and upwards.

Solid Extract Quassia, U. S. P.—One part equals 12 of drug.

Dose, 1 grain (0.06 Gm.) and upwards.

Pills.**Iron, Quassia and Nux Vomica, 3 1-4 grs.**

Iron by hydrogen, 1 1-2 grs.

Ext. Nux vomica, 1-4 gr

Ext. Quassia, 1 gr.

Powd. Soap, 1-2 gr.

Quassia Extract, 1 gr.**Tincture Quassia, U. S. P.**

Fluid Extract Quassia, P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol.....19 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

QUEBROCHO. *Aspidosperma Quebracho, Schlechtendal.* (Bark.)*Synonym.*—Quebracho Blanco.*Properties.*—Quebracho has lately been introduced as a remedy in dyspnœa.

Given in proper doses it relieves the cyanosis and sense of suffocation due to embarrassed respiration, as in emphysema, capillary bronchitis, phthisis and chronic pneumonic processes, and in asthma. It appears to assist the oxygenation of the blood as well as to stimulate the respiratory centers.

ACTIVE PRINCIPLE.

Aspidospermine.—Quebracho bark contains at least six different alkaloids. The aspidospermine of commerce consists of all the alkaloids of the bark, and hence represents fairly well the activity of the drug, which is perhaps the most valuable remedy we possess for dyspnœa in its various forms.

Dose, 1-4 to 1-2 grain (0.016 to 0.03 Gm.).

PREPARATIONS.**Fluid Extract Quebracho.**—Not miscible with water.

Dose, 15 minims to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Quebracho.—One part equals 10 of drug.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

QUEEN-OF-THE-MEADOW. *Eupatorium purpureum, Linné.*
(Leaves, root.)*Synonyms.*—Gravel Root, Trumpet Weed.

Properties.—A most excellent diuretic; tonic, somewhat stimulant and astringent. It has acquired a reputation for gravel, stranguy and all chronic diseases of the urinary organs. It may be used with excellent effect in dropsical affections, hæmaturia, gout and rheumatism.

Queen-of-the-Meadow leaves, compressed in ounce packages for retailing purposes.

Queen-of-the-Meadow root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Eupurpurin, a so-called **Concentration**,* a powdered preparation from Queen-of-the-meadow root.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

Extracts.

Fluid Extract Queen-of-the-Meadow root.—Miscible with water without material precipitation.

Dose, 30 to 60 minims (2 to 4 C.c.).

Solid Extract Queen-of-the-Meadow.—One part equals 7 of root.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

* See note in Preface on Concentrations.

QUININE-FLOWER. *Sabbatia Elliottii, Steudel.* (Root.)

Properties.—Tonic, febrifuge, antiperiodic. In its habitat quinine flower is much employed by physicians for the cure of the different types of malarial fever, the drug being used until the patient feels the *tininitus aurium* peculiar to the effects of quinine. In this respect it seems to have the same properties as the Cinchona barks to a similar degree, hence its name. Its reputation as an antiperiodic was fully established during the late war, when owing to the scarcity of quinine, every opportunity was afforded for testing the relative value of the various substitutes.

PREPARATION.

Fluid Extract Quinine-flower root.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.) every two hours. To be increased in quantity in obstinate cases.

RAGWEED. *Ambrosia artemisiæfolia, Linné.* (Leaves.)

Synonyms.—Roman Wormwood, Bitterweed, Hogwood; Ambrosia, Fr.; Traubenkraut, Ger.

Properties.—Astringent. Employed as an application to fresh wounds and inflammation of the soft parts arising from injuries, and in hemorrhoids. It is said to exert a stimulant action in the typhoid state of febrile affections.

Ragweed leaves, compressed in ounce packages for retailing purposes.

RASPBERRY. *Rubus Idæus, U. S. P. Rubus strigosus, Michaux.* (Leaves.)

Properties.—Similar in medicinal properties to the bark of the root of the other varieties of the same genus—the Dewberry and Blackberry. It will be found useful in the treatment of diarrhoea, dysentery, cholera infantum, relaxed conditions of the intestines in children, passive hemorrhages from the stomach, bowels or uterus, and in colliquative diarrhoea. As an injection, serviceable in gonorrhoea, gleet, leucorrhoea and prolapsus uteri and ani. Combined with aromatics, useful in cholera morbus, cholera infantum and diarrhoea. It is said that Raspberry will, during labor, increase the activity of the uterine contractions, when these are feeble, even in instances where ergot has failed, and it has been found serviceable in after-pains.

Raspberry leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Raspberry leaves.—Miscible with water.

Dose, 20 to 60 minims (1.25 to 4 C.c.).

Solid Extract Raspberry leaves.—One part equals 4 of drug.

Dose, 5 to 10 grains (0.32 to 0.65 Gm.).

RED CLOVER. *Trifolium pratense, Linné.* (Blossoms.)

Properties.—Deobstruent and sedative. The blossoms of this plant have long been used in the form of an infusion in whooping-cough, which they are said to relieve promptly. It has been used with success in the treatment of scrofula and externally as a wash for ill-conditioned ulcers.

Red Clover tops, compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Red Clover tops.—Miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Solid Extract Red Clover tops.—One part equals 5 of drug.

Dose, 5 to 25 grains (0.32 to 1.6 Gm.).

RED CLOVER—Continued—(Extracts.)

Syrup Trifolium Compound.—This syrup is recommended by Dr. Rush as superior in efficacy to the combination of alteratives that has recently acquired considerable notoriety under the name of bamboo brier root compound, succus alterans, or simply alterative compound. The doctor's experience with this latter combination led him to believe that the formula could be greatly improved, and, after much clinical experience, he has adopted the combination represented in this preparation as that best adapted for general use. He prescribes it in all cases of secondary syphilis, and finds that its restorative action in nearly every case is prompt and unequivocal.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.) three times a day.

Red clover, 32 grs.

Stillingia, 16 grs.

Burdock root, 16 grs.

Poke root, 16 grs.

Berberis aquifolium, 16 grs.

Cascara amara, 16 grs.

Prickly-ash bark, 4 grs.

Potassium iodide, 8 grs.

RED COHOSH. *Actæa spicata, Linné; var. rubra, Aiton.* (Root.)

Synonym.—Red Baneberry.

Properties.—A violent purgative, which must be used with caution. Said to possess emmenagogue properties. Dr. Davies states that this root possesses a certain narcotic property which may prove useful. Diluted with water and applied locally, the fluid extract acts as an insecticide.

Red Cohosh root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Red Cohosh (Red Baneberry).—Not miscible with water.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

RED MAPLE. *Acer rubrum, Linné.* (Bark.)

Synonym.—Swamp maple.

Properties.—A mild astringent. It has long been used by the Indians as their favorite remedy in all cases of sore eyes. Its virtues otherwise are not known.

Red Maple bark, compressed in pound packages for retailing purposes.

RED OAK. *Quercus rubra, Linné.* (Bark.)

Properties.—Contains much tannin and therefore powerfully astringent. It is useful wherever an astringent is indicated, such as in chronic diarrhœa, chronic mucous discharges, etc., and externally to indolent ulcers, hemorrhoids. Useful in leucorrhœa in form of injection and as a gargle in relaxed uvula, etc.

Red Oak bark, compressed in pound packages for retailing purposes.

RED OSIER. *Cornus sericea, Linné.* (Bark.)

Synonyms.—Swamp Dogwood, Rose Willow, Silky Cornel.

Properties.—Tonic and astringent; it may be used as a substitute for *Cornus florida* (see Dogwood) in the same doses. Is said to check the vomiting common in pregnancy. Is also recommended in dropsy, ulcers, and malignant fevers.

Red Osier bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Red Osier.—Not miscible with water.

Dose, 30 to 60 minims (2 to 4 C.c.).

RED POPPY. *Papaver Rhœas*, *Linné*. (Petals.)

Synonyms.—Red Poppy Petals, Corn Rose, Corn Poppy; Coquelicot, Pavot rouge. Fr.; Klatschrosen, Ger.

Properties.—Said to be anodyne and carminative. An uncertain but very weak opiate.

Red Poppy flowers, compressed in ounce packages for retailing purposes.

FORMULA.**Syrup Poppy flowers, B. P.**

Flowers 1 oz.
Hot water 1-2 pint.
Macerate 12 hours, strain, evaporate to 1 1-2 ozs., add 3 ozs. sugar, and when cold add alcohol, 3 fluidrachms. Useful in catarrhal affections of children.
Dose, 1 to 4 fluidrachms (4 to 16 C.c.).

RHATANY. *Krameria*, U. S. P. *Krameria triandra*, *Ruiz & Pavon*. (Root.)

Properties.—Rhatany is a powerful astringent, with some slight tonic virtues. It may be employed internally with advantage in menorrhagia, hæmatemesis, passive hemorrhages, chronic diarrhœa, leucorrhœa, chronic mucous discharges, colliquative perspiration, and incontinence of urine. As an energetic styptic in epistaxis, hemorrhage from the cavity of an extracted tooth, or the surface of a wound, and as a local application in prolapsus ani, fissure of the anus and leucorrhœa, as also for spongy and bleeding gums, it will be found unsurpassed.

Rhatany Root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.

Capsules.—In soft gelatin.

Copaiba, Cubeb and Rhatany.

Copaiba, Para, 6 minims. Ext. Rhatany, 2 grs.
Oleoresin cubeb, 2 minims.

Copaiba and Rhatany.

Copaiba, Para, 8 minims. Ext. Rhatany, 2 grs.

Extracts.

Fluid Extract Rhatany, U.S.P.—Not miscible with water without some precipitation.
Dose, 15 to 30 minims (1 to 2 C.c.).

Powdered Extract Rhatany.—of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Rhatany.
Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Solid Extract Rhatany, U. S. P.—One part equals 8 of drug.
Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Pills.**Cubeb Extract, Rhatany and Iron, 3 grs.**

Ext. Cubeb, 1 1-2 grs. Ext. Rhatany, 1-2 gr.
Iron sulph. exsic., 1 gr.

Krameria Extract, 2 grs.**Syrup Rhatany, U. S. P.**

Fluid Extract Rhatany, P., D. & Co.'s 7 fl. ozs.
Hot Syrup 10 fl. ozs.
Mix and strain.
Dose, for a child, 10 to 30 minims (0.6 to 2 C.c.).

Tincture Rhatany, U. S. P.

Fluid Extract Rhatany, P., D. & Co.'s 4 fl. ozs.
Dilute Alcohol 17 fl. ozs.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

NOTE.—This also represents the tinctures of the Phar. Ger. and Fr. Cod.

RHUBARB. Rheum, U. S. P. Rheum officinale, *Baillon*; and other species. (Root.)

Properties.—Cathartic, tonic and astringent. Its astringency is exerted after its cathartic effect is produced, thus making it a very valuable remedy in bowel complaints. Its tonic power also makes it a valuable remedy for habitual constipation and dyspepsia.

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Rhubarb and Columbo.

Rhubarb, 90 grs. Columbo, 30 grs.

Rhubarb and Magnesia.

Rhubarb, 60 grs. Magnesium acetate, 10 grs.

Rhubarb and Potassa.

Rhubarb, 60 grs. Potassium bicarbonate, 60 grs.
Golden-seal, 30 grs.

Extracts.

Fluid Extract Rhubarb, U. S. P.—Not miscible with water.
Dose, 5 to 30 minims (0.3 to 2 C.c.).

Fluid Extract Rhubarb Aromatic.—Not miscible with water. Twice the strength of the official aromatic tincture of Rhubarb, U. S. P., which it otherwise resembles in every particular. It is employed to prepare extemporaneously the aromatic syrup of Rhubarb, and for this purpose is preferable to the official tincture, as it does not introduce into the syrup so large a proportion of alcohol. Sixteen fluidounces, imperial, represents: Rhubarb, 6 1-2 ozs. av.; Cinnamon, 1 1-4 ozs.; Cloves, 1 1-4 ozs.; Nutmeg, 2-3 ozs.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Rhubarb Comp. with Potass.—Not miscible with water.

Tonic, antacid and laxative. Used principally in diarrhoea, dysentery, cholera morbus, cholera infantum, and in the same diseases as the compound powder of rhubarb. It is frequently of use in correcting acidity of the stomach, and may be employed as a gentle laxative, especially during pregnancy, and when piles are present. Sixteen fluidounces, imperial, represents: Rhubarb, 8 ozs. av.; Golden-seal and Cassia, 4 ozs. each, with addition of 30 minims Oil of Peppermint and 1 oz. Potassium Carbonate.
Dose, 15 to 60 minims (1 to 4 C.c.).

Fluid Extract Rhubarb and Senna.—Not miscible with water. A mild cathartic, well adapted to cases of costiveness with gastric uneasiness, in persons of a gouty habit.

Sixteen fluidounces, imperial, represents: Rhubarb and Senna, 8 ozs. av. each.
Dose, 15 to 60 minims (1 to 4 C.c.).

Fluid Extract Rhubarb, sweet.—Not miscible with water.

Four times the strength of the official sweet tincture of rhubarb. Sixteen fluidounces, imperial, represents: Rhubarb, select, 5 ozs. av.; Licorice root and Anise seed, 2 1-2 ozs. each; Cardamom seed, 5-8 oz.
Dose, 1-4 to 1 1-2 fluidrachms (1 to 6 C.c.).

Powdered Extract Rhubarb.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Rhubarb.

Dose, 2 to 10 grains (0.13 to 0.6 Gm.).

Solid Extract Rhubarb, U. S. P.—One part equals 3 of drug.

Dose, 2 to 10 grains (0.13 to 0.6 Gm.).

Solid Extract Rhubarb Compound, Phar. Ger.—Consists of: Ext. Rhubarb, 6 parts; Ext. Aloes, 2 parts; Resin Jalap, 1 part; Soap, 4 parts.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

Normal Liquid* Rhubarb, assayed.—Standard arbitrary, 30 per cent. extractive.

Dose, 5 to 25 minims (0.3 to 1.5 C.c.).

Pills.

Anti-constipation, Carsons', 2 3-4 grs.

Ext. Cascara sagrada, 1 gr.

Ext. Rhubarb, 1 gr.

Ext. Nux vomica, 1-4 gr.

Aloin, 1-3 gr.

* See note in Preface on Normal Liquids.

RHUBARB—Continued—(Pills.)

- Asafetida and Rhubarb**, 3 grs.
Asafetida, 1 gr. Iron by hydrogen, 1 gr.
Powd. Rhubarb, 1 gr.
- Calomel and Rhubarb**, 1 1-2 grs.
Calomel, 1-3 gr. Ext. Coloc, comp., 1-2 gr
Ext. Rhubarb, 1-2 gr. Ext. Henbane, 1-6 gr.
- Liver**, Christopher's, 3 1-2 grs.
Calomel, 2 grs. Powd. Ipecac, 1-2 gr.
Rhubarb, 1 gr.
- Magnesia and Rhubarb**, 2 grs.
Magnesia calcined, 1 gr. Powd. Rhubarb, 1 gr.
- Rhubarb Extract**, 1 gr.
- Rhubarb, U. S. P.**, 3 grs.
- Rhubarb and Blue Mass**, 3 1-4 grs.
Blue Mass, 1 3-4 grs. Powd. Rhubarb, 1 gr.
Sodium bicarbonate, 1-2 gr.
- Rhubarb Compound**, 3 1-2 grs.
Powd. Rhubarb, 1 1-2 grs. Purified Aloes, 1 1-8 grs.
Powd. Myrrh, 3-4 gr. Oil Peppermint, q. s.
- Rhubarb Compound, U. S. P.**, 4 1-2 grs.
Powd. Rhubarb, 2 grs. Powd. Myrrh, 1 gr.
Purified Aloes, 1 1-2 grs. Oil Peppermint, 1-10 gr.
- Rhubarb and Iron**, 3 grs.
Powd. Rhubarb, 2 grs. Iron sulph. exsicc., 1 gr.

For others see appended Pill Formulæ list for numbers 45, 46, 104, 113, 145, 166, 173 and 174.

Rhein, a so-called **Concentration**,* a powdered preparation from Rhubarb root.
Dose, 1 to 10 grains (0.06 to 0.6 Gm.).

Syrups.—Each fluidounce represents ingredients specified.

Rhubarb, U. S. P.—See below for formula for extemporaneous preparation.
Rhubarb, 55 grs. Carbonate of potassium, 4 grs.
Cinnamon, 11 grs.
Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Rhubarb, aromatic, U. S. P.—See below for formula for extemporaneous preparation.
Rhubarb, 11 grs. Cloves, 2 grs.
Cinnamon, 2 grs. Nutmeg, 1 gr.
Dose, for an infant, 1 fluidrachm (4 C.c.) repeated every two hours till the passages indicate by their color that the medicine has operated.

Rhubarb and Potassium Comp.—Neutralizing cordial.
Fluid Extract Rhubarb comp. with Potassium carb., 2 fluidrachms.
See above for formula
Dose, for an adult, 1-2 fluidounce (15 C.c.) in water.

Tinctures.—See below for formulæ for extemporaneous preparation.

Rhubarb, U. S. P.
Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Rhubarb, aromatic, U. S. P.
Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Rhubarb, sweet, U. S. P.
Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Wine Rhubarb, U. S. P.—Each fluidounce represents:
Rhubarb, 45 grs. Calamus, 4 grs.
See below for formula for extemporaneous preparation.
Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

FORMULÆ.**Syrups.**

- Rhubarb, U. S. P.**
Fluid Extract Rhubarb, P., D. & Co.'s.....2 1-2 fl. ozs.
Fluid Extract Cassia, P., D. & Co.'s .. 1-2 fl. oz.
Carbonate of Potassium. 68 grs.
Syrup.....14 1-2 fl. ozs.
- Rhubarb, aromatic, U. S. P.**
Fluid Extract Rhubarb, aromatic, P., D. & Co.'s.....1 fl. oz.
Syrup.....15 fl. ozs.

* See note in Preface on Concentrations.

RHUBARB—Continued.**Tinctures.****Rhubarb, U. S. P.**

Fluid Extract Rhubarb, P., D. & Co.'s.....	1 1-2 fl. ozs.
Fluid Extract Cardamom, P., D. & Co.'s.....	1-4 fl. oz.
Dilute Alcohol.....	11 1-2 fl. ozs.

Rhubarb, aromatic, U. S. P.

Fluid Extract Rhubarb, aromatic, P., D. & Co.'s.....	8 fl. ozs.
Dilute Alcohol.....	8 fl. ozs.

Rhubarb, sweet, U. S. P.

Fluid Extract Rhubarb, sweet, P., D. & Co.'s.....	4 fl. ozs.
Dilute Alcohol.....	12 fl. ozs.

Wine Rhubarb, U. S. P.

Fluid Extract Rhubarb, P., D. & Co.'s....	2 1-2 fl. ozs.
Fluid Extract Sweet Flag, P., D. & Co.'s.....	1-4 fl. oz.
Stronger white wine, q. s. to make	24 fl. ozs.

RHUS AROMATICA. *Aiton.* (Bark of Root.)

Synonym.—Sweet Sumach.

Properties.—"An efficient remedy against that very annoying condition of the bladder which gives rise to incontinence of urine."* Its astringency makes it serviceable in passive hemorrhages, atonic diarrhoea, dysentery and summer complaints of children. It acts as an excitant on the non-striped muscles of the bladder, the uterus, and of the inferior portion of the digestive canal.

PREPARATIONS.

Fluid Extract *Rhus Aromatica*.—Not miscible with water.

Dose, 5 to 30 minims (0.3 to 2 C c.).

Solid Extract *Rhus Aromatica*.—One part equals 9 of drug.

Dose, 1 to 5 grains (0.06 to 0.3 Gm.).

ROSE. { Pale. U. S. P., *Rosa centifolia*, } *Linné.* (Petals.)
 { Red. U. S. P., *Rosa gallica*, }

Properties.—Pale rose petals are seldom used medicinally; those of the red rose are used for their pleasant astringency in aphthæ and other ulcers of the mouth, ears, anus, etc., inflammation of the eyes, hæmorrhages and wherever a mild astringent is indicated.

Rose petals, pale, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Confection Red Rose petals, U. S. P.

Used exclusively as an excipient.

Lozenge, Rose-leaf and Alum.

Powd. Rose leaves, 1-4 gr. Powd. Alum, 1-4 gr.

Ointment Rose-water (cold cream), U. S. P.

ROSEMARY. *Rosemarinus*, U. S. P. *Rosmarinus officinalis*, *Linné.*
 (Flowers, leaves, U. S. P.)

Properties.—Formerly considered useful in functional nervous disorders, and as an emmenagogue, galactagogue and diuretic. Externally is employed in fomentations for relief of local pains.

Rosemary flowers, { Compressed in ounce packages for retailing purposes.
Rosemary leaves, U. S. P., }

* Napheys, "Medical Therapeutics," p. 467.

ROSIN-WEED. *Silphium laciniatum*, *Linné*. (Herb, root.)

Synonyms.—*Silphium gummiferum*, *Elliott*; Compass Plant, Indian Cup Plant, Polar Plant.

Properties.—Tonic, diaphoretic, alterative and emetic. Very beneficial in dry, obstinate coughs and intermittent fevers. Its action is perhaps analogous to that of *grindelia robusta*, and it is said to cure the heaves (asthma) in horses.

Rosin-weed herb, compressed in ounce packages for retailing purposes.

Rosin-weed root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Rosin-weed herb.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

RUE. *Ruta graveolens*, *Linné*. (Leaves.)

Synonyms.—*Rue odorante*, Fr.; *Garten raute*, Ger.

Properties.—A powerful emmenagogue in functional inactivity of the uterus and ovaries. It is of service in hysteria, convulsions and flatulence, the oil of rue being used preferably for the latter.

Rue leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Rue.—Not miscible in water without some precipitation.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

Solid Extract Rue.—One part equals 5 of leaves.

Dose, 2 to 5 grains (0.13 to 0.32 Gm.).

Pill, Emmenagogue, Rigaud's, 33-4 grs.

Purified Aloes, 1 1-2 grs.

Powd. Rue, 3-4 gr.

Powd. Saffron, 3-4 gr.

Powd. Savin, 3-4 gr.

SABBATIA CAMPESTRIS, *Nuttall*. (Plant.)

Properties.—Said to be a valuable remedy in sick-headache, when caused by deranged digestion. Useful also as an antiperiodic and preventative of miasmatic fevers; and in drachm doses, every four hours, it is a reliable tonic in convalescence from fevers and general debility of the system from any cause.

PREPARATION.

Fluid Extract Sabbatia Campestris.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

SAFFLOWER. *Carthamus tinctorius*, *Linné*. (Florets.)

Synonyms.—*African or Dyer's Saffron*, *American Saffron*, *Bastard Saffron*, *Saffron Thistle*.

Properties.—Used in domestic practice to favor the efflorescence of measles and other exanthemata and as a diaphoretic (in warm infusion). Used to dissipate commencing catarrh and muscular rheumatism. Laxative in large dose.

Safflower, compressed in ounce packages for retailing purposes.

PREPARATION AND FORMULA.

Fluid Extract Safflower (Saffron, American).—Miscible with water without material precipitation.

Dose, 5 to 20 minims (0.3 to 1.2 C.c.).

Syrup Safflower (Saffron, American).

Fluid Extract Safflower, P., D & Co's.....1 fl. oz.

Hot Syrup.....7 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Used as a popular remedy for children teething.

SAGE. *Salvia*, U.S.P. *Salvia officinalis*, *Linné.* (Leaves.)

Synonyms.—Garden Sage; *Sauge officinale*, Fr.; *Salbei*, Ger.

Properties.—Stimulant aromatic, tonic and astringent. It is a well known diaphoretic in the form of "sage tea." Its astringency is of service in the form of a gargle where such is indicated for the mouth or throat; its aromatic quality is useful in flatulence.

Sage leaves, U. S. P., domestic and Italian, compressed in ounce, 1-4 and 1-2 pound packages for retailing purposes.

PREPARATION.

Fluid Extract Sage leaves.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

SAINT JOHN'S-WORT. *Hypericum perforatum*, *Linné.* (Herb.)

Properties.—Reputed diuretic, astringent and sedative. Has been employed in a great variety of diseases, but especially esteemed in affections of the urinary passages. Used in hot infusion as an emmenagogue.

Saint John's-wort herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Saint John's-wort.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

SAMPSON SNAKE-ROOT. *Gentiana ochroleuca*, *Frölich.* (Root)

Properties.—Astringent, tonic, antiperiodic. Useful in all cases of enfeebled mucous tissue attended with discharge as in chronic catarrhal affections. Especially applicable to subacute irritations of the alimentary canal, where the mesenteric glands are somewhat involved and attended with diarrhoea.

Sampson Snake-root, compressed in pound packages for retailing purposes

SANDALWOOD. *Santalum album*, *Linné.* (Wood; oil, U.S.P.)

Synonyms.—Yellow Saunders, White Saunders.

Properties.—Employed as a substitute for copaiba in the treatment of gonorrhoea; used in chronic bronchitis and chronic diarrhoea with advantage. The fluid extract of the wood is more readily tolerated by some patients than the oil.

PREPARATIONS.**Capsules.**

Copaiba, Cubeb, Matico and Sandal.—In soft gelatin.

Copaiba, Para, 3 minims.

Oleoresin Matico, 1 minim.

Oleoresin Cubeb, 3 minims.

Oil Sandal, East India, 3 minims.

Copaiba, Cubeb and Sandal.—In soft and in hard gelatin.

Copaiba, Para, 6 minims.

Oil Sandal, East India, 2 minims.

Oil Cubeb, 2 minims.

Copaiba and Sandal.—In soft gelatin.

Copaiba, Para, 5 minims.

Oil Sandal, East India, 5 minims.

Copaiba and Sandal.—In soft gelatin.

Copaiba, Para, 7 minims.

Oil Sandal, East India, 3 minims.

Cubeb and Sandal.—In hard gelatin.

Oil Cubeb, 5 minims.

Oil Sandal, East India, 5 minims.

Sandal Oil, East India, 10 minims.—In soft and in hard gelatin.

Sandal and Cassia.—In soft and in hard gelatin.

Oil Sandal, East India, 9 minims. Oil Cassia, 1 minim.

Fluid Extract Sandalwood.—Not miscible with water.

Cinnamon adds to its palatability.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Pill Sandalwood Compound, 3 grs.

Oil Sandalwood, 1 gr.

Balsam Copaiba, 1 gr.

Ext. Cubeb, 1 gr.

SARACA INDICA, *Linné*. (Bark.)

Synonyms.—*Joutsia* Asoka, *Roxburgh*; Asok, Hind. and Beng.; Asoka, Jasundi, Bomb.

Properties.—Astringent, uterine sedative. "Much employed by the Hindoo physicians in uterine affections, especially in menorrhagia."* Useful also in the treatment of hemorrhoidal tumors.

PREPARATION.

Fluid Extract *Saraca indica*.—Not miscible with water without some precipitation.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

SARSAPARILLA, U. S. P. (Root.)

Honduras: *Smilax officinalis*, *Humboldt, Bonpland & Kunth*.

Mexican: *Smilax medica*, *Chamisso & Schlechtendal*.

Para: *Smilax* (species indetermined).

Properties.—Its most extensive and useful application is in the treatment of secondary syphilis and syphilitic diseases, and that shattered condition of the system following the imprudent use of mercury. It may also be employed in chronic rheumatism, scrofulous affections and certain cutaneous diseases.

Sarsaparilla root, U. S. P., Honduras, and Mexican, each, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.

Capsule Copaiba, Cubeb and Sarsaparilla.—In soft gelatin.

Copaiba, Para, 6 minims.

Ext. Sarsaparilla, 2 grs.

Oleoresin Cubeb, 2 minims.

Extracts.

Fluid Extract Sarsaparilla, U. S. P.—Miscible with water.

Dose, 30 to 60 minims (2 to 4 C.c.).

Fluid Extract Sarsaparilla (Para).—Miscible with water.

Dose, 30 to 60 minims (2 to 4 C.c.).

This valuable species of Sarsaparilla is comparatively unknown in the American markets, the supplies being carried through the usual channels of trade from Brazil to Europe.

Fluid Extract Sarsaparilla Comp., U. S. P.—Miscible with water.

Each pint represents: Sarsaparilla, 12 ozs.; Licorice root, 2 ozs.; Sassafras, 1-2-3 ozs.; Mezereon, 1-2 oz.

Dose, 1 fluidrachm (4 C.c.)

NOTE.—The ingredients of this fluid extract differ somewhat from those of the Syrup Sarsaparilla Compound, U. S. P. For the especial purpose of convenience in making the latter see below.

Fluid Extract Sarsaparilla Comp. (for Syrup).—Miscible with water.

Four times the strength of the Syrup Sarsaparilla Comp., U. S. P.

Dose, 30 to 60 minims (2 to 4 C.c.).

Fluid Extract Sarsaparilla and Dandelion.—Miscible with water.

Each pint represents: Sarsaparilla and Dandelion, 8 ozs. each. Useful in eruptive diseases depending chiefly on derangements of the hepatic functions.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Powdered Extract Sarsaparilla.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Sarsaparilla.

Dose, 4 to 10 grains (0.26 to 0.6 Gm.).

Solid Extract Sarsaparilla.—One part equals 5 of drug.

Dose, 4 to 10 grains (0.26 to 0.6 Gm.).

Solid Extract Sarsaparilla Comp.—One part equals 4 of the following mixture of crude drugs, one pound of which contains: Dandelion, 9.6 ozs.; Yellow Dock, 2.4 ozs.; Sarsaparilla, 4.0 ozs.; Oil Anise, 1-5 minim; Oil Wintergreen, 2-5 minim.

Dose, 4 to 10 grains (0.26 to 0.6 Gm.).

* Dymock, "Materia Medica of Western India," p. 209.

SARSAPARILLA—Continued.**Pill Sarsaparilla Extract**, 3 grs.**Syrups**.—Each fluidounce represents ingredients specified.**Sarsaparilla.**

Fluid extract sarsaparilla, 2 drachms.

Sarsaparilla Compound, U. S. P.

Sarsaparilla, 97 grs.

Guaiacum wood, 12 grains.

Pale rose, 8 grs.

Licorice root, 8 grs.

Senna, 8 grs.

Sassafras, 4 grs.

Anise, 4 grs.

Wintergreen, 4 grs.

For preparing this extemporaneously:

Fluid Extract Sarsaparilla Comp., for Syrup, P., D. & Co.'s. 4 fl. ozs.

Simple Syrup.....12 fl. ozs.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.) three or four times a day.

Sarsaparilla Comp. with Potassium Iodide.

Sixteen grains Potassium iodide in each fluidounce of syrup sarsaparilla comp.

SASSAFRAS, U. S. P. Sassafras officinale, *Nees*. (Bark of root.)*Properties*.—Aromatic, stimulant, alterative, diaphoretic and diuretic; used in combination with other alternatives but principally to impart flavor.**PREPARATION.****Fluid Extract Sassafras**.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

SASSY-BARK. Erythrophlœum guineense, *Don*. (Bark.)*Synonyms*.—Erythrophlœum judiciale, *Procter*; Casca-bark, Doom-bark, Mancona-bark.*Properties*.—Poisonous. The ordeal poison of Angola. Experiments conducted with a view to fix its value in medicine have shown it to be nauseant, emetic, astringent and narcotic. It is said to have been successfully employed in diarrhœa, dysentery and in colic. "In heart affections, it strengthens and moderates the action of the heart like digitalin."**Antidotes*.—Emetics or stomach-pump, if sufficiently early. Use stimulants and treat on general principles. Nux vomica has been recommended as a physiological antidote.**PREPARATION.****Fluid Extract Sassy-bark**.—Not miscible with water.

Dose, 5 to 15 minims (0.3 to 1 C.c.) used with caution.

SAVIN. Sabina, U.S.P. Juniperus sabina, *Linné*. (Tops.)*Properties*.—Poisonous. Emmenagogue, diuretic and diaphoretic. In large doses it will produce gastro-enteritis. Contra-indicated where general or local inflammation is present, and it should never be used during pregnancy, on account of its tendency to cause abortion. Notwithstanding this, it is reputed efficacious in checking the tendency to abort, and to be beneficial in menorrhagia when carefully exhibited in small doses.*Antidotes*.—Emetics, castor oil, opiates, linseed meal poultice to abdomen.**Savin tops, U. S. P.**, compressed in ounce packages for retailing purposes.**PREPARATIONS AND FORMULA.****Extracts.****Fluid Extract Savin, U. S. P.**—Not miscible with water.

Dose, 5 to 20 minims (0.3 to 1.25 C.c.).

Solid Extract Savin.—One part equals 4 of drug.

Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

* Lescher, "Recent Materia Medica," p. 23.

SAVIN—Continued.**Pills.****Emmenagogue Improved, 4 1-2 grs.**

Ergotin, 1 gr.	Ext. Black Hellebore, 1 gr.
Purified Aloes, 1 gr.	Iron sulphate, exsic., 1 gr.
Oil Savin, 1-2 gr.	

Emmenagogue No. 4, 4 1-2 grs.

Ergotin, 1 gr.	Iron sulphate, exsic., 1 gr.
Ext. Cotton root, 1 gr.	Oil Savin, 1-4 gr.
Purified Aloes, 1 gr.	

Emmenagogue, Rigaud's, 3 3-4 grs.

Purified Aloes, 1 1-2 grs.	Powd. Saffron, 3-4 gr.
Powd. Rue, 3-4 gr.	Powd. Savin, 3-4 gr.

Savin Extract, 1 gr.**Tincture Savin, B. P.**

Fluid Extract Savin, P., D. & Co.'s.....	1 fl. oz.
Dilute Alcohol.....	7 fl. ozs.

Dose, 20 to 60 minims (1.25 to 4 C.c.).

SAW PALMETTO. *Serenoa serrulata, Bentham & Hooker.* (Fruit.)*Synonyms.*—*Sabal serrulata, Roemer & Schultes; Chamærops serrulata, Pursh.**Properties.*—Sedative, pseudo-narcotic and diuretic, said to improve digestion, increasing the flesh and strength. It is remedial of neuralgic disorders and it will also allay irritation of the mucous membrane of the throat, nose and larynx. It has been used with decided success in phthisis pulmonaris, bronchitis, acute and chronic laryngitis, etc.**PREPARATION.****Fluid Extract Saw Palmetto berries.**—Not miscible with water.
Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).**SAXIFRAGE.** *Pimpinella Saxifraga, Linné.* (Root.)*Synonyms.*—*Pimpinella, Small burnet saxifrage; Grand Boucage, Fr.; Pimpinelle, Biebernell, Ger.**Properties.*—Diuretic, diaphoretic and stomachic; useful in chronic catarrh, asthma, dropsy, amenorrhœa, etc. It is also recommended as a masticatory in toothache, and as a gargle in palsy of the tongue, and in collection of viscid mucus in the throat.**Saxifrage root,** compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Saxifrage.**—Not miscible with water without some precipitation.
Dose, 15 to 30 minims (1. to 2. C.c.).**SCAMMONY.** *Scammonium, U. S. P.* Resinous exudation from the root of *Convolvulus Scammonia, Linné.**Properties.*—A powerful intestinal, but a feeble hepatic irritant, with tendency to occasion griping. Said to be somewhat uncertain as a purgative; best given in combination. Only mentioned for the preparations into which it enters.

Dose, 5 to 10 grains (0.32 to 0.65 Gm.).

PREPARATIONS.**Pills.****Cathartic Compound, Vegetable, 2 1-2 grs.**

Ext. Coloc., 1-3 gr.	Purified Aloes, 1 1-4 grs.
Podophyllin, 1-4 gr.	Powd. Cardamom, 1-8 gr.
Powd. resin Scammony, 1-3 gr.	Powd. Soap, 1-8 gr.

Cochia, 3 1-2 grs.

Ext. Coloc. comp., 1 gr.	Powd. Gamboge, 1-2 gr.
Purified Aloes, 1 gr.	Powd. Scammony, 1-2 gr.
Powd. Potass. sulph., 1-2 gr.	

SCAMMONY—Continued—(Pills).**Triplex**, Dr. Francis', 4 grs.

Purified Aloes, 1 1-5 grs.

Blue Mass, 1 1-5 grs.

Oil Caraway, 1-5 gr.

Resin Scammony, 1 1-5 grs.

Croton Oil, 1-20 gr.

Tr. Aloes and Myrrh, 1-4 gr.

Powdered Extract Colocynth Comp. U. S. P.—Each 100 parts represents: Ext. Colocynth, 16; Aloes, 50; Cardamom, 6; Resin of Scammony, 14; Soap, 14. See Colocynth.
Dose, 5 to 20 grains (0.3 to 1.3 Gm.).

SCARLET PIMPERNEL. *Anagallis arvensis*, Linné. (Herb.)

Synonyms.—Red Chickweed, Weather-glass; Mouron rouge, Fr.; Gauchheil, Rothe Miere, Ger.

Properties.—This plant has won considerable reputation in California as a remedy in rheumatism. It has been used with some success in epilepsy, mania, and nervous affections.

PREPARATION.**Fluid Extract Scarlet Pimpernel.**—Miscible with water.

Dose, 1 fluidrachm (4 C.c.) gradually increased.

SCOURING-RUSH. *Equisetum hyemale*, Linné. (Plant.)*Synonyms.*—Shave grass; Winter Schachtelhalm, Ger.

Properties.—Diuretic and astringent. Taken freely in the form of decoction, it has been found useful in dropsy, suppression of urine, hematuria, gravel and nephritic affections, as well as in gonorrhœa and gleet.

PREPARATION.**Fluid Extract Scouring-rush.**—Miscible with water.

Dose, 20 to 60 minims (1.3 to 4 C.c.).

SCULLCAP. *Scutellaria*, U. S. P. *Scutellaria lateriflora*, Linné. (Lvs.)

Synonyms.—Blue Scullcap, Mad-dog Weed, Hoodwort; Scutellaire, F.; Helm-kraut, Ger.

Properties.—Tonic, nervine and antispasmodic. It is recommended in chorea, convulsions, tremors, neuralgia and nerve affections generally. In all cases of nervous excitability, restlessness, or wakefulness, either idiopathic or accompanying acute or chronic diseases, it may be freely administered with favorable results.

Scullcap leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.****Fluid Extract Scullcap.**—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Scullcap Comp.—Not miscible with water.

Each pint represents: Scullcap and Ladies' Slipper, 5 1-2 ozs. each; Hops and Lettuce, 2 1-2 ozs. each. Combines the tonic and nervine properties of its several ingredients. Indicated in the nervous disturbances of hysterical females, and in the debility of children following an attack of summer complaint.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Powdered Extract Scullcap.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Scullcap.

Dose, 4 to 15 grains, (0.26 to 1 Gm.).

Solid Extract Scullcap.—One part equals 5 of drug.

Dose, 4 to 15 grains (0.26 to 1 Gm.).

Scutellarin, a so-called **Concentration**,* a powdered preparation from Scullcap leaves.

Dose, 1 to 3 grains (0.065 to 0.2 Gm.).

* See note in Preface on Concentrations.

SEDUM ACRE, Linné. (The whole plant.)

Synonyms.—English moss, Stonecrop, Wall-pepper, Mossy stonecrop; Jaubarbe, Fr.; Mauerpfeffer, Ger.

Properties.—Emetic, local irritant, and astringent. Great success has been recently claimed for it in the treatment of diphtheria.* Whether given in weak solution and swallowed, or in a more concentrated form and applied locally to the throat, it appears to act in the following manner: 1. By infiltrating the false membrane and so loosening or destroying its connection with the mucous membrane that it is readily ejected by vomiting. 2. By cleansing or disinfecting the cleared mucous surface and also the healthy parts, new diphtheritic foci are prevented from forming. 3. By its astringency reducing any tumefaction. 4. By its superior emetic properties. The emesis it produces is copious, causing little pain or fatigue. Dr. P. O. Wagener gives the following formula† as yielding the most satisfactory results in his experience with the remedy:

Fluid ext. sedum acre, fl. ʒj.

Spirits turpentine.

Lactic acid.

Fluid ext. aconite, ℥ss fl. ʒij. Mix.

One application should be made with a brush every three minutes for twenty minutes, when emesis should be induced by a glass of warm water if it has not already commenced, and the membrane will be entirely expelled. To prevent liability of ulceration from any bleeding points they should then be touched with Monsel's solution.

NOTE.—Sedum acre has no specific power to cure diphtheria, consequently it is of no value in the first stage of the disease; it should only be employed when the false membrane has been developed.

PREPARATION.

Fluid Extract Sedum acre.—Miscible in water without material precipitation.

Dose, 15 to 30 minims (1 to 2 C.c.) diluted.

SENEGA, U. S. P. *Polygala Senega, Linné.* (Root.)

Properties.—Senega is a stimulating expectorant and diuretic, and in large doses emetic and cathartic. It appears, indeed, to excite more or less all the secretions, proving occasionally diaphoretic, emmenagogue, and increasing the flow of saliva. It has been employed as an expectorant in cases not attended with acute inflammatory action, or in which the inflammation has been in a great measure subdued. It is used with reputed advantage in the treatment of chronic catarrhal affections, secondary stages of croup, diseases of the bronchial passages, pneumonia, asthma, incipient croup, rheumatism, dropsy and amenorrhœa.

Senega root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Abstract Senega, U. S. P.**

Fluid Extract Senega root, P., D. & Co.'s..... 1 fl. oz.

Sugar of milk, q. s. to make..... 228 grains.

Evaporate the fluid extract nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well-corked bottle.

Dose, 5 to 10 grains (0.32 to 0.65 Gm.).

Elixir Cimicifuga (Black Cohosh) Compound.—Each fluid-ounce represents ingredients specified.

Cimicifuga, 30 grs.

Seneka, 8 grs.

Licorice root, 10 grs.

Ipecac, 4 grs.

Cherry, wild, 8 grs.

* Dr. Louis Duval in *El Siglo Medico*, Madrid, 7, xij, 84.

† *Therapeutic Gazette*, July 15, 1885.

SENEGA—Continued.**Extracts.**

Fluid Extract Black Cohosh Comp.—Not miscible with water. Each pint represents: Black Cohosh, 8 ozs.; Licorice, 3 ozs.; Cherry bark and Seneka, 2 ozs. each; Ipecac, 1 oz. Nervous and arterial sedative; in larger doses acting as a diaphoretic and causing increased secretion from the bronchial mucous membrane. Useful in chronic rheumatism, both articular and lumbago formi; and in atonic amenorrhœa.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Horehound Comp.—Not miscible with water without some precipitation.

Each pint represents: Horehound, 8 ozs.; Comfrey, 4 ozs.; Seneka and Wild Cherry, 2 ozs. each. Demulcent, and expectorant. A valuable combination in coughs and colds with mild diaphoretic and sedative properties; particularly fitted for administration to children.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Ipecac and Senega.—Not miscible with water. Each pint represents: Ipecac and Seneka, 8 ozs. each. Emetic in large doses. In small doses, a stimulating expectorant and diaphoretic.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

Fluid Extract Senega, U. S. P.—Miscible with water.

Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

Fluid Extract Squill Comp.—Miscible with water.

Sixteen fluidounces, imperial, represents: Squill and Senega, 8 ozs. av. each. In large doses emetic. In smaller doses expectorant, diuretic and diaphoretic. Much used in inflammatory affections of the air passages, asthma and incipient croup.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

Glycerole Squill Comp.—Each fl. oz. represents ingredients specified.

Squill, 40 grs. Senega, 40 grs. Antimony and Potass. tart., 7-8 gr.

Lozenges.**Tar Compound.**

Pine Tar, 7-18 gr.

Powd. Wild Cherry, 2 1-3 grs.

Powd. Senega, 7-24 gr.

Yerba Santa Compound.

Ext. Licorice, 1 gr.

Powd. Cherry, 1-4 gr.

Powd. Senega, 3-16 gr.

Yerba santa, 1 gr.

Tar, 1-32 gr.

Syrups.—Each fluidounce represents ingredients specified.

Senega, U. S. P., 120 grs.

For preparing this extemporaneously:

Fluid Extract Senega, P., D. & Co.'s.....4 fl. ozs.

Hot Syrup.....12 fl. ozs.

Aqua Ammonia.....45 minims.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Squill Compound, U. S. P.—"Hive Syrup," "Croup Syrup."

Squill and Senega, 34 grs. each, and 7-8 gr. tartrate antimony and potassium For preparing this extemporaneously:

Fluid Extract Squill Comp, P., D. & Co.'s.....2 fl. ozs.

Tartar Emetic.....12 grs.

Boiling Water.....3 fluidrachms.

Simple Syrup.....11 fl. ozs.

Dissolve the tartar emetic in the boiling water, and mix with the other ingredients.

Dose, for children, 10 minims to 1 fluidrachm (0.6 to 4 C.c.); for adults, 20 to 30 minims (1 3 to 2 C.c.) as an expectorant.

SENNA, U.S.P. (Alexandrian.) *Cassia acutifolia, Delile.* (Lvs.)

Properties.—A prompt, efficient and safe cathartic. Very useful in febrile diseases. It acts principally upon the small intestines. It is apt to cause griping pains, which are modified by combining it with aromatics as correctives, and also by the mode of preparing it for use.

PREPARATIONS AND FORMULÆ.

Confection Senna.—A combination of Senna with coriander, cassia fistula, tamarinds, etc., making an efficient and mild laxative in doses of about 2 drachms (8 Gm.).

SENNA—Continued.

Elixirs.—Each fluidounce represents ingredients specified.

Cathartic Compound.

Senna, 16 grs.

Cascara sagrada, 32 grs.

Culver's root, 16 grs.

Butternut, 16 grs.

Rochelle salts, 32 grs.

Laxative.

Senna, 60 grs.

Butternut bark, 30 grs.

Rochelle salts, 60 grs.

Mandrake Compound.

Mandrake root, 45 grs.

Culver's root, 30 grs.

Senna, 30 grs.

Jalap, 15 grs.

Extracts.**Fluid Extract Dandelion and Senna.**—Miscible with water.

Sixteen fluidounces, imperial, represents: Dandelion and Senna, 8 ozs. each. Tonic and laxative, with slight action on liver.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Fluid Extract Grindelia Comp.—Miscible with water.

Eighteen fluidounces, imperial, represents: Grindelia Robusta, 12 ozs. av.; Senna and Rhubarb, 3 ozs. each. Efficacious in the treatment of asthma where there is a torpid condition of the bowels.

Dose, 1 fluidrachm (4 C.c.), repeated every three or four hours.

Fluid Extract Mandrake Comp.—Not miscible with water.

Each pint represents 16 ounces of the following drugs in the proportions named: Mandrake and Culver's Root, 6 parts each; Senna, 4 parts; Jalap, 2 parts.

Dose, as a laxative, 1 fluidrachm (4 C.c.); as a cathartic, 1 to 3 fluidrachms (4 to 11 C.c.).

Fluid Extract Pink-root Comp.—Not miscible with water without some precipitation.

Sixteen fluidounces, imperial, represents: Pink-root and Senna, 7 ozs. each; Savin and Manna, 1 oz. each. Efficacious as a specific for worms; combines both anthelmintic and laxative properties.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Pink-root and Senna.—Miscible with water.

Sixteen fluidounces, imperial, represents: Pink-root, 10 ozs. av.; Senna, 6 ozs.; Anise and Caraway oils, 20 minims each. An excellent vermifuge.

Dose, for a child two years old, 30 to 60 minims (2 to 4 C.c.); for an adult, 2 to 4 fluidrachms (8 to 15 C.c.).

Fluid Extract Rhubarb and Senna.—Not miscible with water.

Sixteen fluidounces, imperial, represents: Rhubarb and Senna, 8 ozs. each. Especially adapted to cases of costiveness with gastric uneasiness, in persons of gouty habit.

Dose, 15 to 60 minims (1 to 4 C.c.).

Fluid Extract Senna, U. S. P.—Miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Fluid Extract Senna, Aqueous.—Miscible with water.

Less apt than the hydro-alcoholic (official) extract to cause griping pains.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Fluid Extract Senna Comp.—Not miscible with water without some precipitation.

Each pint represents: Senna and Jalap, 7 ozs. each; Coriander, 2 ozs. An excellent mild purgative, useful in constipation attended with flatulence.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Senna and Jalap.—Not miscible with water.

Each pint represents: Senna and Jalap, 8 ozs. each. Senna modifies the severity of the Jalap action.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Powdered Extract Senna.—Of the same strength as the solid extract; prepared by evaporating the solid extract, at a low temperature, to sensible dryness, and replacing moisture with powd. senna leaves.

Dose, 10 to 20 grains (0.65 to 1.3 Gm.).

Solid Extract Senna.—One part equals 6 of drug.

Dose, 10 to 20 grains (0.65 to 1.3 Gm.).

Pill Senna Extract, 2 grs.

SENNA—Continued.**Syrups.****Sarsaparilla Compound, U. S. P.**—See under Sarsaparilla.**Senna, U. S. P.**

Fluid Extract Senna, P., D. & Co.'s.....8 fl. ozs.

Fluid Extract Coriander, P., D. & Co.'s.....1 fl. oz.

Syrup.....11 fl. ozs.

Dose, 1 to 4 fluidrachms (4 to 15 C. c.).

NOTE.—Syrup Senna, Phar. Ger. is about 1-4 of this strength and flavored with fennel instead of coriander.

SERPENTARIA, U.S.P. *Aristolochia Serpentaria, Linné.* (Rhiz.)*Synonyms.*—Virginia Snakeroot, Serpentry Root; Serpentaire de Virginie, Fr.; Virginische Schlangenzwurzel, Ger.*Properties.*—A stimulant tonic, acting also as a diaphoretic or diuretic, according to the mode of its application. It is adapted to the treatment of typhoid fevers, whether idiopathic or symptomatic, and has been found useful in intermittent and other fevers, in dyspepsia and amenorrhœa. In infusion it is an excellent gargle for putrid sore throat.**Serpentaria, U. S. P.**, compressed in pound packages for retailing purposes.**PREPARATIONS AND FORMULA.****Extracts.****Fluid Extract Cinchona Comp.**—Not miscible with water.

Each pint represents: Cinchona bark, pale, 8 ozs.; Bitter Orange Peel, 6 ozs.; Serpentaria, 1 1-2 ozs.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Cinchona, Red, Comp.—Not miscible with water.Each pint represents: Red Cinchona Bark, true, 7 1-2 ozs. av.; Bitter Orange Peel, 6 ozs.; Serpentaria, 1 1-2 ozs. Five times the strength of the Tincture Cinchona Compound, U. S. P., *q. v.*

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Extract Serpentaria, U. S. P.—Not miscible with water without some precipitation.

Dose, 15 to 30 minims (1 to 2 C. c.).

Tinctures.**Cinchona Comp, U. S. P.**—See Cinchona.**Serpentaria, U. S. P.**

Fluid Extract Serpentaria, P., D. & Co.'s.....2 fl. ozs.

Dilute Alcohol.....19 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

SHEEP SORREL. *Rumex Acetosella, Linné.* (Leaves.)*Synonym.*—Field sorrel.*Properties.*—Refrigerant and diuretic. Useful in febrile and inflammatory diseases, and in scorbutic affections.**Sheep Sorrel leaves**, compressed in ounce packages for retailing purposes.**PREPARATIONS.****Fluid Extract Sheep Sorrel.**—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

Solid Extract Sheep Sorrel.—One part equals 4 of the leaves.

Dose, 10 grains (0.65 Gm.).

SHELL-BARK HICKORY. *Carya alba, Nuttall.* (Bark.)*Properties.*—Tonic and anti-intermittent. Recommended for its tonic properties in the debility following an acute attack of malaria, or owing to exposure to a prolonged influence of the poison. As an abortive of intermittent fever it has a high reputation in some localities.**PREPARATION.****Fluid Extract Shell-bark Hickory.**—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

SHEPHERD'S PURSE. *Capsella bursa-pastoris*, *Manch.* (Herb.)*Synonym.*—Hirtentäschlein, Ger*Properties.*—An active diuretic, possessing also tonic and stimulant powers.

It has been found useful in many forms of kidney complaint, ascites, œdema glottidis, chronic diarrhœa, etc.

PREPARATIONS.**Fluid Extract Shepherd's Purse.**—Miscible with water without material precipitation

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

"German Tincture" Shepherd's Purse.—An imported unofficial preparation of the green herb.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

SILKWEED. *Asclepias Cornuti*, *Decaisne.* (Rhizome.)*Synonyms.*—*Asclepias syriaca*, *Linne*; Milkweed; *Asclépiade à la Soie*, Fr.; Seidenpflanze, Ger*Properties.*—Tonic, alterative, expectorant, diuretic. It has been tested in scrofula and found of considerable value. Its expectorant powers render it beneficial in asthma and bronchial disorders. It is indicated in dropsies, scarlatina and rheumatism. "This plant has a specific power in dispelling dropsical effusions."***Silkweed root**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Silkweed.**—Not miscible with water without some precipitation.

Dose, 10 to 60 minims (0.65 to 4 C. c.).

SIMARUBA. *Simaruba officinalis*, *DeCandolle.* (Bark of the root.)*Synonym.*—Mountain damson.*Properties.*—Used with advantage in dysentery and chronic diarrhœa. It operates simply as a tonic; and, though beneficial in relaxed and debilitated states of the alimentary canal, would do harm if used indiscriminately in all dysenteric cases. In large doses causes purging and diaphoresis and sometimes diuresis.**PREPARATION.****Fluid Extract Simaruba bark.**—Not miscible with water

Dose, 10 to 30 minims (0.6 to 2 C. c.).

SKUNK CABBAGE. *Symplocarpus foetidus*, *Salisbury.* (Rhizome.)*Synonyms.*—Skunk Weed, Polecat Weed.*Properties.*—Stimulant, antispasmodic and narcotic. In large doses it occasions nausea and vomiting, with headache, vertigo and dimness of vision. Used in asthma, whooping-cough, nervous irritability, hysteria, and convulsive affections; it is also used in chronic catarrh, and bronchial and pulmonary affections, to allay the irritation.**Skunk Cabbage root**, compressed in pound packages for retailing purposes.**PREPARATIONS.****Fluid Extract Lobelia Comp.**—Not miscible with water

Sixteen fluidounces, imperial, represents: Lobelia, 8 ozs av: Blood-root and Skunk Cabbage, 4 ozs, each. Diaphoretic, expectorant, antispasmodic, emetic.

Dose, as an expectorant, 3 to 20 minims (0.2 to 1.3 C. c.); as an emetic, 20 to 60 minims (1.3 to 4 C. c.)

Fluid Extract Skunk Cabbage.—Not miscible with water

Dose, 10 to 40 minims (0.6 to 2.5 C. c.).

* Napheys, "Medical Therapeutics," p. 461.

SLIPPERY ELM. *Ulmus*, U. S. P. *Ulmus fulva*, *Michaux.* (Bark.)

Properties.—Highly demulcent, slightly astringent and tonic, and somewhat nutritious. Used wherever a demulcent is indicated, such as in diarrhoea, dysentery, and affections of the urinary passages. Externally in poultice form to relieve local inflammations.

Slippery Elm bark-flour, compressed in 1-8 lb. packages for retailing purposes.

SOAP-TREE. *Quillaia*, U.S.P. *Quillaia saponaria*, *Molina.* (Bark.)

Synonyms.—Soap Bark; Seifenrinde, Ger.

Properties.—Internally it is recommended by Dr. Kobert* as a preferable substitute for senega. His clinical experience shows it to be borne better by patients than senega, that it very seldom produces vomiting or diarrhoea; that its sweeter taste renders it more acceptable to children; that it is a more powerful and uniform expectorant, since quillaia contains five times the proportion of the same two glucosides that are present in senega, with the additional advantage of being more constant in quantity. When bruised and macerated in water, Soap-tree bark, in common with other substances containing saponin, such as senega and saponaria, imparts to that liquid the property of frothing like soap when agitated. It is used for cleansing grease from cloth, in washing silk or woolen stuff when it is desirable not to change their color, and also for cleansing the hair, which it is supposed to beautify and preserve.

Soap-tree bark, U. S. P., compressed in pound packages for retailing purposes.

PREPARATION AND FORMULÆ.

Fluid Extract Soap-tree bark.—Miscible with water.

Dose, for a child, 2 minims (0.12 C.c.); for adults, 8 minims (0.5 C.c.) taken hourly.

Hair Wash.

Fluid Extract Soap-tree bark, P., D. & Co.'s.....	2 fl. ozs.
Cologne	1 fl. oz.
Glycerin.....	1 fl. oz.
Water	12 fl. ozs.

Mix first three ingredients thoroughly, then add the water. Said to be a very valuable wash for the hair, promoting its growth and eradicating dandruff.

Tooth Wash.—Resembling and said to be superior to "Sozodont."

Fluid Extract Soap-tree bark, P., D. & Co.'s.....	1 fl. oz.
Alcohol	3 fl. ozs.
Glycerin.....	2 fl. ozs.
Water	10 fl. ozs.
Oil Wintergreen.....	6 drops.
Oil Cloves	4 drops.

Color with Tincture Cudbear. Dissolve the oils in the alcohol, add the glycerin and fluid extract, and finally the water.

SOAPWORT. *Saponaria officinalis*, *Linné.* (Leaves, root.)

Synonyms.—Soaproot; Seifenwurz, Ger.

Properties.—Diaphoretic, diuretic, and said to be, as an alterative, superior to sarsaparilla. Used in Germany as a remedy in venereal and cutaneous eruptions and visceral obstructions.

Soapwort leaves, compressed in ounce packages for retailing purposes.

Soapwort root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Soapwort root.—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.)

* Centralblatt für Klin. Med., No. 30 1885: Therapeutic Gazette, Sept. 15, 1885, March 15, 1886, and May 16, 1887.

SOLOMON'S SEAL. *Polygonatum officinale, Allioni.* (Rhizome.)*Synonym.*—*Convallaria Polygonatum, Linné.**Properties.*—Tonic, mucilaginous and mildly astringent. Found of much value in leucorrhœa, menorrhagia, female debility and pectoral affections. Very efficacious in irritable conditions of the intestines, piles, erysipelas, and cutaneous affections of an erysipelatous nature. Affords direct relief in maladies of the skin produced by the poison vine and other plants.**Solomon's Seal**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Solomon's Seal.**—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

SOUR-WOOD. *Oxydendrum arboreum, DeCandolle.* (Leaves.)*Synonyms.*—Sorrel Tree, *Andromeda arborea, Linné.**Properties.*—Tonic, refrigerant and diuretic, and containing free acid in large quantity. It has been employed successfully in the treatment of dropsy, acting in this case as an efficient diuretic.**PREPARATIONS.****Fluid Extract Sour-weed leaves.**—Miscible with water without material precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Solid Extract Sour-weed leaves.—One part equals 5 of drug.

Dose, 5 to 15 grains (0.33 to 1 Gm.).

SOUTHERNWOOD. *Artemisia Abrotanum, Linné.* (Herb.)*Synonyms.*—Old man; aurne des jardins, Fr.; Eberraute, Ger.*Properties.*—The properties of the various species of *Artemisia* seem to be similar. Southernwood is anthelmintic, tonic and narcotic. Used in intermittent fever, jaundice, atonic dyspepsia, amenorrhœa, chronic leucorrhœa, obstinate diarrhœa, and as a vermifuge. Combined with a fixed alkali it is said to become powerfully diuretic.**Southernwood herb**, compressed in ounce packages for retailing purposes.**PREPARATION.****Fluid Extract Southernwood.**—Miscible with water without material precipitation.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

SPANISH NEEDLES. *Bidens bipinnata, Linné.* (Herb.)*Properties.*—Claimed to have a special tendency, in its action, to the uterus, and that it is an efficient emmenagogue. It is also reported to be a stimulating expectorant.**PREPARATION.****Fluid Extract Spanish Needles.**—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

SPEARMINT. *Mentha viridis, Linné.* (Leaves.)*Properties.*—An aromatic stimulant, antispasmodic and carminative, of milder properties than peppermint, and therefore often preferred in infantile cases.**Spearmint leaves, U. S. P.**, compressed in ounce packages for retailing purposes.**PREPARATION.****Fluid Extract Spearmint leaves.**—Miscible with water.

Dose, 1 to 3 fluidrachms (4 to 11 C.c.).

SPEEDWELL. *Veronica officinalis*, *Linné*. (Herb.)

Synonym.—Veronique mâle, Fr.; Ehrenpreis Wundkraut, Ger.

Properties.—Alterative, diuretic and expectorant. Useful in catarrh of the kidney, and as an aromatic expectorant in acute colds with cough. Used in domestic practice in the form of infusion (one ounce to the pint of hot water). Give freely.

Speedwell herb, compressed in ounce packages for retailing purposes.

SPICE-BUSH. *Lindera Benzoin*, *Meissner*. (Bark, berries, leaves.)

Synonyms.—Benzoin odoriferum, *Nees*; Wild Allspice, Feverbush, Spice-wood; Laurier benzoin, Fr.; Benzoeloorbeer, Ger.

Properties.—All parts of the shrub are diaphoretic (in hot infusion), aromatic, stimulant and tonic. The bark is also used in the treatment of intermittent and typhoid forms of fever as an exhilarant and refrigerant. The berries have been employed as a substitute for allspice, they are used advantageously in flatulence, flatulent colic, etc., and externally in rheumatism.

Spice-bush leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS.**Extracts.**

Fluid Extract Spice-bush bark.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Fluid Extract Spice-bush berries.—Not miscible with water without some precipitation.

Dose, 20 to 60 minims (1.3 to 4 C.c.).

Fluid Extract Spice-bush berries Comp.—Miscible with water. Fifteen and one-half fluidounces, imperial, represents: Mugwort, 13 ozs. av.; Orange Peel, 1 oz.; Spicebush berries, 1-2 oz.; Licorice root, 1 oz. A desirable remedy for the treatment of skin diseases, dependent upon peculiar conditions of the system, other than those of syphilitic or scrofulous origin. It possesses valuable tonic and alterative properties, and has been employed with great benefit in the treatment of acne, boils, etc.

Dose, 1-2 to 1 fluidrachm (1 to 4 C.c.).

SPIKENARD. *Aralia racemosa*, *Linné*. (Rhizome.)

Synonyms.—American Spikenard, Pettymorrell.

Properties.—Gently stimulant, diaphoretic and alterative. Used in chronic rheumatism, syphilitic and cutaneous affections.

Spikenard root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Spikenard.—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Syrup Horehound Comp.—Each fluidounce represents: Elecampane, Spikenard, Comfrey, Wild Cherry bark, Horehound, 20 grs. each; Blood-root, 10 grs.

SPINY CLOTBUR. *Xanthium spinosum*, *Linné*. (Herb.)

Synonyms.—Spiny Burreed, Spiny Cocklebur.

Properties.—Diaphoretic, sialagogue and diuretic. It causes a rise of temperature and accelerates the circulation. Dr. Grzymala, of Olschanka, Podolia, writing to the *British Medical Journal*, states, that of several persons bitten by rabid dogs and treated by him in the wards of his hospital, such always recovered as were treated with this medicine. In nearly 100 cases of men bitten and in which *Xanthium spinosum* was tested he had not failed in one case to ward off hydrophobia. The drug deserves further investigation.

PREPARATION.

Fluid Extract Spiny Clotbur.—Miscible with water.

Dose, 10 to 20 minims (0.6 to 1.3 C.c.).

SQUAW-VINE. *Mitchella repens, Linné.* (Herb.)

Synonyms.—Checkerberry, Partridgeberry, Winterclover.

Properties.—Diuretic and astringent. Used in dropsy, suppression of urine and diarrhoea. and has been found useful in many forms of uterine derangement such as amenorrhoea, some forms of dysmenorrhoea, menorrhagia, congestion of the uterus, etc.

Squaw-vine herb, compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.

Elixir Helonias Compound.—Each fluidounce representing:

Helonias, 15 grs.	Cramp bark, 15 grs.
Squaw-vine, 60 grs.	Blue Cohosh, 15 grs.

Extracts.

Fluid Extract Squaw-vine.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Squaw-vine Comp. (or Helonias Comp.)

—Not miscible with water.

Fourteen fluidounces, imperial, represents: Squaw-vine, 8 ozs. av.; Cramp bark, Blue Cohosh and Helonias, 2 ozs. each. Uterine tonic and antispasmodic. May be used in all cases where the functions of the internal reproductive organs are deranged, as in amenorrhoea, dysmenorrhoea, menorrhagia, leucorrhoea; and to overcome the tendency to habitual abortion. Said to render parturition "safe and easy."

Syrup Squaw-vine Comp.—Mother's Cordial (King).

Fluid Extract Squaw-vine Comp., P., D. & Co.'s.....	4 fl. ozs.
---	------------

Syrup	12 fl. ozs.
-------------	-------------

Dose, 2 to 4 fluidrachms (8 to 15 C.c.) three times a day.

SQUILL. *Scilla, U.S.P. Urginea Scilla, Steinheil.* (Bulb.)

Synonyms.—Squills; Scille, Squille, Fr.; Meerzwiebel, Ger.

Properties.—In over-doses, poisonous, occasioning hypercatharsis, strangury, bloody urine and fatal inflammation of the stomach and bowels. In medicinal doses, diuretic, expectorant; reduces frequency and increases tension of the pulse. Used chiefly in the form of syrup as an expectorant. As a diuretic, it is often prescribed in dropsy, especially in cases where the pulse is weak and the heart feeble or obstructed. It is not well adapted to cases of renal origin.

ACTIVE PRINCIPLE.

Scillitoxin.—Cardiac sedative, also acting on the kidneys.

Dose, 1-300 grain (0.0002 Gm.) cautiously increased.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Squill, U. S. P.—Miscible with water.

Dose, 1 to 10 minims (0.06 to 0.6 C.c.).

Fluid Extract Squill Comp.—Miscible with water.

Each pint represents: Squill and Senega, 8 ozs. each. Expectorant, diuretic, diaphoretic and, in large doses, emetic. Much used in inflammatory affections of the air passages, asthma and incipient croup.

Dose, 5 to 20 minims (0.3 to 1.3 C.c.).

Glycerole Squill Compound.—Each fluidounce containing: Squill, and Senega, 40 grs. each; Antimony and Potass. Tart., 7-8 gr.

Pills.

Digitalis Compound, 4 grs.

Powd. Digitalis, 1 gr.

Powd. Squill, 1 gr.

Powd. Potass. nitrate, 2 grs.

Ipecac and Squill, 3 grs.

Powd. Ipecac comp., 2 grs.

Powd. Squill, 2-3 gr.

Ammoniac, 2-3 gr.

Scillitoxin, 1-300 gr.

Squill Compound, 3 grs.

Powd. Squill, 3-8 gr.

Powd. Jamaica Ginger, 3-4 gr.

Powd. Soap, 11-8 grs.

Ammoniac, 3-4 gr.

SQUILL—Continued.**Syrups.**

Squill, U. S. P.—Each fluidounce contains 1-2 fluidounce Vinegar of Squill, representing 23 grains Squill. An extemporaneous preparation may be made as follows:

Fluid Extract Squill, P., D. & Co.'s	1 fl. oz.
Acetic Acid	13-4 fl. ozs.
Syrup	17 fl. ozs.

Squill Compound, U. S. P.—Each fluidounce containing Squill and Senega, 34 grs. each; and 7-8 gr. Tartrate Antimony and Potassium. An extemporaneous preparation may be made as follows:

Fluid Extract Squill Comp., P., D. & Co.'s	2 fl. ozs.
Tartar Emetic	11 grs.
Simple Syrup	11 fl. ozs.
Boiling Water	3 fluidrachms

Dissolve the tartar emetic in the boiling water, and mix solution with the other ingredients.

Dose, for children, 10 minims to 1 fluidrachm (0.6 to 4 C.c.); for adults, 20 to 30 minims (1.3 to 2 C.c.) as an expectorant.

Tincture Squill, U. S. P.

Fluid Extract Squill, P., D. & Co.'s	2 fl. ozs.
Dilute Alcohol	12 fl. ozs.

Dose, 6 to 30 minims (0.4 to 2 C.c.).

Vinegar Squill, U. S. P.

Fluid Extract Squill, P., D. & Co.'s	2 fl. ozs.
Diluted Acetic Acid	18 fl. ozs.

Dose, 10 to 60 minims (0.6 to 4 C.c.).

STAR-GRASS. *Aletris farinosa*, Linné. (Rhizome.)

Synonyms.—**Unicorn root**, **Ague Root**, **Blazing Star**, **Colic Root**, **Star-wort**.

Properties.—Tonic, diuretic and vermifuge. Used extensively in diseases of the uterine organs, and is supposed to exert a specific influence upon the uterus itself. It imparts tone and vigor to both male and female reproductive organs.

Star-grass root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Aletrin, a so-called **Concentration**,* a powdered preparation from Star-grass (unicorn) root.

Dose, 1-2 to 4 grains (0.03 to 0.26 Gm.).

Extracts.

Fluid Extract Star-grass (Unicorn) root.—Not miscible with water without some precipitation.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

Solid Extract Star-grass (Unicorn) root.—One part equals 6 of drug.
Dose, 2 to 5 grains (0.13 to 0.32 Gm.).

STAVESACRE. *Staphisagria*, U. S. P. *Delphinium Staphisagria*, Linné. (Seed.)

Properties.—Poisonous. Emetic, cathartic, narcotic, vermifuge; parasiticide, frequently used to destroy lice in the hair. It is employed in various disorders of the urino-genital apparatus, as in the prostaticorrhœa resulting from masturbation; in chronic irritation of the neck of the bladder, especially when the result of gonorrhœa or cold, and associated with temporary enlargement or irritation of the prostate; in chordee, and in gonorrhœal prostatitis; in leucorrhœa, amenorrhœa, and in prolapsus uteri where there is feeble circulation.

ANTIDOTES.—Use emetics (mustard, sulphate of zinc or apomorphine) and combat symptoms as they arise. Stimulants are generally indicated.

PREPARATION.

Fluid Extract Stavesacre seed.—Not miscible with water.

Dose, 1 to 2 minims (0.065 to 0.13 C.c.), gradually increased.

* See note in Preface on Concentrations.

STILLINGIA, U. S. P. *Stillingia sylvatica*, *Linné*. (Root.)

Synonyms.—Queen's Root, Queen's Delight, Yaw Root, Silver leaf.

Properties.—In large doses emetic and cathartic, in some instances leaving a disagreeable sensation in the stomach and alimentary canal, with more or less prostration of the system. In small doses, a powerful alterative, and used extensively as a remedy for scrofulous and syphilitic affections.

Stillingia root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.

Elixirs.—Each fluidounce represents ingredients specified

Corydalis Compound.

Turkey Corn, 30 grs.
Stillingia, 30 grs.
Blue Flag, 40 grs.

Prickly Ash, 15 grs.
Potassium iodide, 8 grs

Stillingia, 120 grs.**Stillingia Compound.**

Stillingia, 30 grs.
Turkey Corn, 30 grs.
Pipissewa, 15 grs.
Elder Flowers, 15 grs.

Blue Flag, 15 grs.
Prickly Ash berries, 7 1-2 grs
Coriander seed, 7 1-2 grs.

Extracts.**Fluid Extract Alternative Compound.**—Miscible with water.

Synonyms —Bamboo Brier Comp., Succus Alterans, Mist. Smilacis Comp. Seventeen fluidounces, imperial, represents Bamboo Brier, Stillingia, Burdock, and Poke roots, 4 ozs. av. each; Prickly Ash bark, 1 oz.

Dose, 1 to 4 fluidrachms (4 to 15 C c.), in water.

Note.—Syrup Trifolium Comp. (see below) is to be preferred, and is rapidly supplanting this formula as an alternative.

Fluid Extract Stillingia Comp.—Not miscible with water.

Formula in same proportion as Elixir Stillingia Comp. above, but four times stronger.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Stillingia, U. S. P.—Miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

Powdered Extract Stillingia.—of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Stillingia.

Dose, 2 to 5 grains (0.13 to 0.3 Gm.).

Solid Extract Alternative Comp.—One grain is equivalent to 5 minims Fluid Extract Alternative comp., see above.

Dose, 10 to 40 grains (0.65 to 2.6 Gm.).

Solid Extract Stillingia.—One part equals 6 of root.

Dose, 2 to 5 grains (0.13 to 0.3 Gm.).

Pills.—Containing Stillingia or its preparations.**Alternative Compound, 3 grs.**

Ext. Smilax Sarsap., 2-3 gr.
Ext. Stillingia sylvat., 2-3 gr.
Ext. Burdock, 2-3 gr.

Ext. Poke root, 2-3 gr.
Ext. Prickly Ash bark, 2-3 gr.

Stillingin, 1 gr.**Stillingin**, a so-called **Concentration**,* a powdered preparation from Stillingia root.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

Syrups—Each fluidounce represents ingredients specified.**Alternative Compound.**

Fl. Ext. Alternative comp., 2 drachms. See above for formula.
Dose, 2 fluidrachms (8 C. c.) and upwards.

Corydalis Compound.

Turkey Corn, 30 grs.
Stillingia, 30 grs.

Blue Flag, 40 grs.
Potassium iodide, 8 grs.

* See note in Preface on Concentrations

STILLINGIA—Continued—(Syrups).**Trifolium Compound.**

Recommended by Dr. Rush as superior in efficacy to the combination of alteratives known as Alternative Compound or Succus Alterans. Its restorative action in secondary syphilis is prompt and unequivocal.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.) three times a day.
 Red Clover, 32 grs. Berberis aquifolium, 16 grs.
 Stillingia, 16 grs. Cascara amarga, 16 grs.
 Burdock root, 16 grs. Prickly-ash bark, 4 grs.
 Poke root, 16 grs. Potassium iodide, 8 grs.

Stillingia, King's formula.

Stillingia, 88 grs. Prickly-ash berries, 44 grs.
 May be prepared extemporaneously as follows:
 Fluid Extract Stillingia, P. D. & Co.'s, 3 fl. ozs.
 Fluid Extract Prickly-ash berries, P. D. & Co.'s, 1 1-2 fl. ozs.
 Hot Syrup, 13 1-2 fl. ozs.
 Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Stillingia Compound, King's formula; active ingredients same as Elixir Stillingia Compound, see above. It may be extemporaneously prepared as follows:

Fluid Extract Stillingia Comp., P., D. & Co.'s, 4 fl. ozs.
 Syrup, 12 fl. ozs.
 Dose, 1 to 4 fluidrachms (4 to 15 C.c.), three or four times a day.

STONE-ROOT. *Collinsonia canadensis, Linné.* (Rhizome.)

Synonyms.—Horsebalm, Horse-weed, Heal-all, Rich-weed, Ox-balm, Knab-root; Guérit tout, Baume de cheval, Tr.

Properties.—Seems to exert an alterative influence on the mucous tissues, and has been found beneficial in chronic catarrh of the bladder, leucorrhœa, debility of the stomach, clergymen's sore throat, chronic laryngitis, etc. It possesses gentle tonic and diuretic properties, and has been used with efficacy in the lithic acid diathesis, calculous deposits and dropsical effusions.

Stone-root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Collinsonin, a so-called Concentration,*—A powdered preparation from Stone-root.
 Dose, 1 to 4 grains (0.06 to 0.25 Gm.).

Elixir Buchu and Pareira Brava.—Each fluidounce representing:
 Buchu, 20 grs. Juniper berries, 15 grs.
 Pareira brava, 7 1-2 grs. Stone-root, 7 1-2 grs.

Extracts.

Fluid Extract Buchu and Pareira Brava.—Not miscible with water.

Each pint represents: Buchu, 8 ozs., troy; Pareira Brava and Stone-root, 2 ozs. each; Juniper berries, 4 ozs. Found useful in leucorrhœa, dropsy, chronic inflammation of the bladder and urinary passages, etc.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Stone-root.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract Stone-root.—One part equals 7 of drug.

Dose, 2 to 5 grains (0.13 to 0.33 Gm.).

STRAMONIUM. { *Stramonii folia*, U.S.P., } *Datura Stramonium*,
{ *Stramonii semen*, U.S.P., } *Linné.* (Lvs., seed.)

Synonyms.—Thorn Apple, Apple Peru, Jamestown-weed, Jimpsonweed.

Properties.—**POISONOUS.** A powerful narcotic. In medicinal doses it acts as an antispasmodic and anodyne without causing constipation. Its uses are the same as those of belladonna, but it has been more particularly recommended as a remedy for attacks of asthma.

* See note in Preface on Concentrations.

STRAMONIUM—Continued.

ANTIDOTES.—Promptly empty the stomach by emetics (sulphate of zinc or copper), and combat the nervous disturbances as they arise by physostigma and opium.

NOTE.—All official preparations are made from the seed.

Stramonium leaves, U. S. P., compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLE.

Daturine (pure crystals).—Alkaloid from stramonium seed; believed to be identical with hyoscyamine.

Dose, 1-100 grain (0.00065 Gm.) and upwards.

Daturine Sulphate.—Differs from the pure alkaloid only in the circumstance that it is readily soluble in water. The solution, one-half to two grains to the ounce, may be employed in place of atropine to dilate the pupil of the eye, its effect being similar but more transient.

Dose, 1-100 grain (0.00065 Gm.) and upwards.

PREPARATIONS AND FORMULÆ.**Extracts.**

Fluid Extract Stramonium leaves.—Not miscible with water without some precipitation.

Dose, 1 to 5 minims (0.06 to 0.3 C.c.).

Fluid Extract Stramonium seed, U. S. P.—Not miscible with water.

Dose, 1 to 4 minims (0.06 to 0.25 C.c.).

Powdered Extract Stramonium leaves (assayed).—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Stramonium leaves.

Dose, 1-4 to 1 grain (0.016 to 0.065 Gm.).

Solid Extract Stramonium leaves (assayed).—One part equals 5 of drug.

Dose, 1-4 to 1 grain (0.016 to 0.06 Gm.).

Solid Extract Stramonium seed, U. S. P. (assayed).—One part equals 8 of drug.

Dose, 1-8 to 1 grain (0.008 to 0.06 Gm.).

Normal Liquids.*

Stramonium leaves.—Not miscible with water without some precipitation.

Standard, .375 per cent. alkaloid, estimated by Dragendorff's method.

Dose, 1 to 4 minims (0.06 to 0.24 C.c.).

Stramonium seeds.—Not miscible with water. Standard, provisionally the same as that of the leaves, viz: .375 per cent. alkaloid.

Dose, 1 to 4 minims (0.06 to 0.24 C.c.).

Ointment Stramonium, U. S. P.—This may be extemporaneously prepared as follows:

Solid Extract Stramonium seed, P., D. & Co.'s..... 52 grs.

Water.....1-2 fluidrachm.

Benzoated Lard.....1 oz. tr.

Rub the extract with the water till uniformly soft, then gradually add the lard and mix thoroughly.

Pills.—Containing Stramonium or its preparations.

Daturine Sulphate, 1-100 gr.

Neuralgic Idiopathic, Brown-Sequard's, 3 1-2 grs., also 1-2 strength 1 3-4 grs.

Ext. Henbane, 2-3 gr.

Ext. Conium seed, 2-3 gr.

Ext. Ignatia, 1-2 gr.

Ext. Opium, 1-2 gr.

Ext. Aconite leaves, 1-3 gr.

Ext. Indian Cannabis, 1-4 gr.

Ext. Stramonium seed, 1-5 gr.

Ext. Belladonna, 1-6 gr.

Ox-gall Compound, 2 1-3 grs.

Ox-gall, 2 grs.

Purified Aloes, 1-10 gr.

Ext. Stramonium seed, 1-6 gr.

Berberine hydrochlorate, 1-12 gr.

Stramonium Extract (seed), 1-4, 1-2 and 1 gr.

* See note in Preface on Normal Liquids.

STRAMONIUM—Continued.**Tinctures.**

"German Tincture" Stramonium Seed.—An imported unofficial preparation from the fresh seeds.
Dose, 5 to 15 minims (0.3 to 1 C.c.).

Stramonium, B. P.

Fluid Extract Stramonium seed, P., D. & Co.'s.....1 fl. oz.
Dilute Alcohol.....7 fl. ozs.
Dose, 8 to 30 minims (0.5 to 2 C.c.).

Stramonium, U. S. P.

Fluid Extract Stramonium seed, P., D. & Co.'s.....2 fl. ozs.
Alcohol.....12 fl. ozs.
Water.....7 fl. ozs.
Dose, 15 to 30 minims (1 to 2 C.c.).

STRAWBERRY. *Fragaria vesca, Linné.* (Leaves.)

Synonyms.—Feuilles du Fraisier Sauvage. Fr.; Erdbeerblätter, Ger.

Properties.—Mildly astringent and diuretic. Employed in mild diarrhoea and in deficient secretion of urine.

Dose, 2 fluidounces (60 C.c.) of an ounce to the pint infusion.

Strawberry leaves, compressed in ounce packages for retailing purposes.

STROPHANTHUS KOMBE, Oliver. (Seeds.)

Properties.—Poisonous. A powerful cardiac stimulant, acting more energetically on the heart than digitalis, while its action on the blood-vessels is less intense. Drs. C. L. Dana and Beverley Robinson report numerous cases of its successful use after other cardiac tonics have failed to produce any result.*

Antidotes.—Emetics or stomach-pump if sufficiently early. Aconite and American hellebore have been suggested as physiologically antagonistic.

PREPARATION.

Tincture Strophanthus.—Not miscible with water without some precipitation.

Prepared after Prof. Fraser's formula, strength being 1 in 20, or one imperial pint representing the virtues of one ounce, av., of the drug; anything stronger has been found too concentrated for convenient use.
Dose, 1 to 10 minims (0.065 to 0.65 C.c.).

SUMACH. *Rhus glabra, Linné.* (Bark, berries U.S.P., leaves.)

Synonyms.—Upland Sumach, Pennsylvania Sumach; Glatter Sumach, Ger.

Properties.—Tonic, astringent and antiseptic. Used as a gargle in inflammation and ulceration of the throat. A valuable astringent in gonorrhoea, leucorrhoea, diarrhoea, dysentery, and in profuse perspiration from debility. The infusion is employed as an injection in prolapsus uteri and ani, etc. The berries are also diuretic and refrigerant, and are employed in febrile diseases, diabetes, etc., as a gargle in quinsy and in ulcerations of the mouth and throat.

Sumach bark, compressed in pound packages for retailing purposes.

Sumach leaves, compressed in ounce packages for retailing purposes.

Extracts.**PREPARATIONS.**

Fluid Extract Sumach bark.—Not miscible with water without some precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Fluid Extract Sumach berries, U. S. P.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Rhusin, a so-called Concentration,† a powdered preparation from Sumach bark.

Dose, 1 to 2 grains (0.06 to 0.13 Gm.).

* Medical Record, Dec. 18, 1886; Therapeutic Gazette, Feb., 1887.

† See note in Preface on Concentrations.

SUMMER SAVORY. *Satureia hortensis*, *Linné*. (Leaves)

Properties.—Stimulant carminative and emmenagogue. Beneficial in warm infusion in colds, menstrual suppression and flatulent colic. It is a gentle stimulating tonic, during convalescence from fevers.

Summer-savory leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Summer-savory.—Not miscible with water without some precipitation.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

SUNDEW. *Drosera rotundifolia*, *Linné*. (Plant.)

Properties.—A valuable pectoral; found very useful in pertussis, asthma, incipient phthisis, chronic bronchitis with dry spasmodic cough, nervous or sympathetic cough, whether from pulmonary, cardiac or gastric disease. Sundew has been recommended in certain phases of dyspepsia, especially as a relief for flatulence. It has effected cures in cases of severe gastric disorder, approaching to chronic ulcer of the stomach, and resembling cancer.

PREPARATIONS.**Extracts.**

Fluid Extract Sundew.—Not miscible with water.

Dose, 5 to 20 minims (0.3 to 1.25 C.c.).

Solid Extract Sundew.—One part equals 5 of drug.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

"German Tincture" Sundew.—An imported unofficial preparation from the green plant.

Dose, 15 to 60 minims (1 to 4 C.c.).

Pill Sundew Extract, 3 grs.

SUNFLOWER. *Helianthus annuus*, *Linné*. (Seeds.)

Properties.—Mild diuretic and expectorant; used with considerable benefit in pulmonary and laryngeal affections.

PREPARATION.

Fluid Extract Sunflower.—Miscible with water.

Best exhibited in syrup.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

SWEET BASIL. *Ocimum Basilicum*, *Linné*. (Leaves.)

Synonyms.—Common or Citron Basil; Basilic, Fr.; Basilienkraut, Ger.; Albahaca, Span.

Properties.—An aromatic tonic and stimulant. Used in Europe principally as a condiment, aiding digestion by stimulating the flow of gastric juice. It is much used in Spanish South America as a carminative and nervine.

In India it is employed "as a remedy in gonorrhœa, catarrh, dysentery and chronic diarrhœa." *

Sweet Basil leaves, compressed in ounce packages for retailing purposes.

SWEET CICELY. *Osmorrhiza longistylis*, *DeCandolle*. (Root.)

Properties.—Aromatic, stomachic, carminative and expectorant. It has proved useful in coughs, in flatulence, and as a gentle stimulant tonic to debilitated stomachs. The sweetened infusion (1 to 10) may be taken freely in coughs.

Sweet Cicely root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Sweet Cicely.—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

* Drury, "Useful Plants of India," p. 337.

SWEET CLOVER. *Melilotus officinalis*, *Willdenow.* (Herb.)

Synonyms.—Yellow Melilot; Mélilot, Fr.; Steinklee, Melilotenkee, Ger.

Properties.—Stimulant and antispasmodic. Useful in strong infusion (one ounce to the pint) in whooping-cough, in 1 to 2 fluidounce doses (30 to 60 C.c.). The herb is sometimes used as an anodyne application in local pains in the form of hot poultice or infusion. Used for its perfume when scattered among clothes and furs to protect them from moths.

Sweet Clover herb, compressed in ounce packages for retailing purposes.

SWEET FERN. *Comptonia asplenifolia*, *Aiton.* (Leaves.)

Synonyms.—Fern gale, Meadow fern.

Properties.—Astringent, carminative and alterative. Used in diarrhoea, dysentery, hæmoptysis, leucorrhœa, rheumatism, debility succeeding fever and in rachitis.

Sweet Fern leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Sweet Fern.—Not miscible with water.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

SWEET FLAG. *Calamus*, U.S.P. *Acorus Calamus*, *Linné.* (Rhiz.)

Synonyms.—Acore odorant, Fr.; Kalmus wurzel, Ger.

Properties.—Carminative, slightly tonic and excitant, and a useful adjunct to other tonics and stimulants. It may be used in cases of flatulent colic, dyspepsia, feebleness of the digestive organs, and to aid the action of bark and quinine in intermittents. Externally it is a valuable application to indolent ulcers, and to keep up the discharge from blistered surfaces and issues. In the flatulent colic of infants, it is best combined with magnesia.

Sweet Flag root, U.S.P., compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Sweet Flag, U. S. P.—Not miscible with water.

Dose, 5 to 40 minims (0.3 to 2.6 C.c.).

Exhibited in syrup it is stated to be used as a substitute for Godfrey's Cordial.

Sweet Flag is also an ingredient (four grains to the fluidounce) of *Wine Rhubarb*, U. S. P., q. v.

SWEET GALE. *Myrica Gale*, *Linné.* (Bark.)

Synonym.—Dutch Myrtle.

Properties.—Astringent and stimulant, used in debilitated conditions of the mucous membranes, scrofula, jaundice, diarrhoea, etc.

PREPARATION.

Fluid Extract Sweet Gale.—Not miscible with water.

Dose, 15 to 30 minims (1 to 2 C.c.).

SWEET GUM. *Liquidambar styraciflua*, *Linné.* (Bark.)

Properties.—One of the most valuable and useful mucilaginous astringents that we possess. It can be employed with advantage in cases of diarrhoea and dysentery, and for these purposes was much used during the late war in camp hospitals.*

Sweet Gum bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Sweet Gum bark.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

* Medical and Surgical History of the War, Part II Medical, p. 47.

SWEET MARJORAM. *Origanum Majorana*, Linné. (Leaves.)

Properties.—Stimulant, tonic and emmenagogue. The hot infusion is used to stimulate the menstrual function.

Dose, a wineglassful (60 C.c.) of an ounce to the pint infusion.

Sweet Marjoram, compressed in ounce packages for retailing purposes.

SWEET ORANGE. *Aurantii Dulcis Cortex*, U. S. P. *Citrus Aurantium* Linné. (Rind of the fruit.)

Synonyms.—*Citrus Aurantium*, Risso; *Apfelsinedschalen*, Ger.

Properties.—Aromatic, tonic. Usually combined with tonic bitters to impart an agreeable flavor, to correct their nauseating influence, and to assist their stimulating properties. As an agreeable flavor, the rind of the sweet orange is preferable; as a tonic, that of the Seville or bitter orange.

PREPARATIONS AND FORMULÆ.

Confection Orange, U. S. P., 1870.—One part grated orange peel to 3 of sugar.

Extracts.

Fluid Extract Sweet Orange.—Miscible in water without material precipitation.

Fluid Extract Sweet Orange Comp.—Miscible with water without material precipitation.

Each pint represents: Orange peel, sweet, 12 ozs. av.; Cloves, Mace, Cinnamon, Caraway, and Anise, 3-8 oz. each; Orris root, 1 1-2 ozs.; Tonka bean, 3-4 oz.

Both of these fluid extracts are used exclusively as flavoring agents.

Syrup Orange Peel, U. S. P.

Fluid Extract Sweet Orange, P., D. & Co.'s. 1 fl. oz.

Hot Syrup. 13 fl. ozs.

Used merely as a vehicle and flavor.

Tincture Sweet Orange Peel, U. S. P.

Fluid Extract Sweet Orange, P., D. & Co.'s. 2 fl. ozs.

Dilute Alcohol 8 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

TAG ALDER. *Alnus serrulata*, Willdenow. (Bark.)

Synonyms.—Red Alder, Smooth Alder, Ecorse d'aune, Fr.; Erlendrinde, Ger.

Properties.—Emetic, astringent and alterative. Useful in the scrofulous and syphilitic diathesis, and in the treatment of chronic cutaneous diseases due to a constitutional dyscrasia.

Tag Alder bark, compressed in pound packages for retailing purposes.

PREPARATIONS.

Alnulin, a so-called **Concentration**,* a powdered preparation from Tag Alder bark.

Dose, 1 to 8 grains (0.06 to 0.5 Gm.).

Fluid Extract Tag Alder.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

TAMARAC. *Larix americana*, Michaux. (Bark.)

Synonyms.—Black Larch, American Larch, Hackmatac.

Properties.—Astringent, and gently stimulant, acting especially on mucous membranes. Has been found particularly useful in chronic bronchitis, forming the basis of the popular compound elixir of larch, see below. It is recommended also in obstruction of the liver, rheumatism, jaundice, diarrhoea, dysentery, and certain cutaneous diseases.

Tamarac bark, compressed in pound packages for retailing purposes.

* See note in Preface on Concentrations.

TAMARAC—Continued.**PREPARATION AND FORMULA.**

Fluid Extract Tamarac.—Not miscible with water.
Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

Elixir Larch Compound.

Fluid Extract Tamarac, P., D. & Co.'s.....	2 fl. ozs.
Fluid Extract White Pine, P., D. & Co.'s.....	1 fl. oz.
Fluid Extract Balsam Fir bark, P., D. & Co.'s.....	1 fl. oz.
Simple Elixir.....	20 fl. ozs.

Dose, 2 to 4 fluidrachms (8 to 15 C.c.).
A favorite and efficacious remedy in chronic bronchitis.

TANSY. Tanacetum, U. S. P. Tanacetum vulgare, *Linné.* (Leaves.)

Synonyms.—Tanaïsie, Fr.; Rainfarn, Ger.

Properties.—Tonic, emmenagogue, anthelmintic and diaphoretic. In small doses it has proved beneficial in convalescence from exhausting diseases, in dyspepsia, with troublesome flatulence, hysteria, jaundice, and worms. In larger doses, administered in hot water, it has been found useful for intermittent fever, suppressed menstruation, tardy labor pains, and as a preventive of the paroxysms of gout.

Tansy leaves, U.S.P., compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Tansy.—Miscible with water without material precipitation.
Dose, 20 to 60 minims (1.3 to 4. C.c.).

TAR. Pix Liquida, U. S. P. A complex empyreumatic oleo-resin from the destructive distillation of wood of the genus *Pinus*, principally *Pinus palustris*, *Miller.*

Properties.—A general stimulant in small doses, quickening the pulse and augmenting the secretions of the skin and kidneys. Said to be of decided benefit in bronchitis, pharyngitis, laryngitis and winter cough. Externally, irritant, rubefacient or mildly stimulant according to degree of dilution.

Dose, 1 to 2 drachms daily (4 to 8 Gm.) in ten-grain doses.

DERIVATIVES AND ALLIED PRODUCTS.

Naphthalin.—A coal-tar product, found also generally in the destructive distillation of organic substances. Crystalline; insoluble in water. Antiseptic. Used as an ointment in scabies, herpes tonsurans, favus, etc. It has been recommended by Rossbach internally in typhoid fever and various intestinal disorders.

Dose, 5 to 30 grains (0.35 to 2 Gm.). Ointment may be made 1:20.

Naphthol.—Crystalline, very sparingly soluble in water, readily miscible with fats and oils. Has a faint, aromatic odor, not at all disagreeable. A powerful antiseptic. Used in the form of an ointment, 1:15 up to 1:8, in the treatment of scabies, psoriasis, chromoplegiosis, and in some forms of chronic eczema. A single application of the ointment, if thoroughly made, will cure scabies.

Xylol.—Has been largely and chiefly employed to moderate the angina and eruption in the throat and to lessen general fetid exhalation in smallpox. Applied by spray and by wash and internally in doses of 10 to 15 minims (0.6 to 1 C.c.).

PREPARATIONS.**Capsules.**

Tar, 10 minims.—In hard and soft gelatin.

Xylol, 5 minims.—In soft gelatin.

Xylol, 10 minims.—In hard gelatin.

Glycerole Tar.—Containing 30 grains to the fluidounce.
Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Inhalant, Tar.—Number 6 of P., D. & Co.'s series of inhalants.

Oil of Tar, 3 ss.

Inhalant No. 1, or neutral oil, ʒj.

TAR—Continued.**Lozenges.****Tar**, 1-7 gr.**Tar Compound.**

Pine Tar, 7-48 gr.

Powd. Senega, 7-24 gr.

Powd. Wild Cherry, 2 1-3 gr.

Yerba Santa Compound.

Ext. Licorice, 1 gr.

Powd. Senega, 3-16 gr.

Tar, 1-32 gr.

Powd. Cherry, 1-4 gr.

Yerba Santa, 1 gr.

Ointments.**Tar, U. S. P.**, 1 in 2.**Tar Compound.**—Equal parts of Tar and Sulphur Ointments.**Soaps, Medicated.****Naphthol Soap.**—Containing 5 per cent., or 84 grains Naphthol.

One of the most effective and practical of all medicated soaps. It has decided antiseptic properties and is of great utility in all offensive discharges from the skin. Useful in very many cutaneous diseases on account of its stimulating, astringent and somewhat anæsthetic effect, and also in being free from odor. Animal parasites and pediculi can be easily, cleanly, and quickly destroyed by its use, superseding the filthy mercurial ointment for this purpose.

Naphthol-Sulphur Soap.—Containing 3 per cent. Naphthol and 10 per cent. Sulphur. Uses the same as above but more specifically adapted for scabies.

Tar Soap.—Containing 10 per cent. tar or 168 grains.

TARO FLOUR. From the bulbous root or corm of *Arum esculentum*, *Linné*.

Properties.—Physicians residing in the Hawaiian Islands recommend for their dyspeptic and consumptive patients an exclusive diet of the cooked taro flour, or "poi," the native food of the aborigines. L. S. Thompson, M. D., Resident Surgeon Kohala, Hawaii, says: "'Poi,' as an article of table use, is beyond praise. It is in the sick room the food 'par excellence,' being perfectly safe in all and any case, and its great nourishing and sustaining power must, when more universally known, give it the highest rank as a diet for invalids."

Taro flour may be employed in a great variety of ways as an article of staple food, but to enlarge on its merits in this respect would be here out of place. We are glad to aid in its general introduction believing that the claims made for it as a dietetic agent entitle it to the attention of the medical profession. Printed matter giving detailed information regarding its use, etc., will be sent on application.

TEA. *Camellia Thea*. *Link.* (Leaves.)

Properties.—Well known as a sustaining and restorative agent, producing its effects chiefly through the caffeine it contains. It is decidedly astringent, and in China is employed, mixed with vinegar, as a remedy for diarrhoea. A valuable remedy in nervous headaches, exerting a remarkable influence as a restorative in all conditions of fatigue and exhaustion. It is one of the best antidotes in cases of opium poisoning, and it is believed that it may aid victims of the opium habit to free themselves from its bondage. Diluted with water it may also be employed as an astringent gargle or wash.

PREPARATIONS.

Fluid Extract Tea.—Not miscible with water without some precipitation.

Dose, 20 to 60 minims (1.3 to 4 C.c.).

Solid Extract Tea.—One part equals 4 of leaves.

Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

THAPSIA GARGANICA, Linné. (Root.)

Synonyms.—Bou-néfa, Faux fenouil, Fr.

Properties.—Employed chiefly as a counter-irritant in the localized pains of rheumatism, gout, bruises, and in dyspnoea. Internally its properties are tonic. In over-doses it is an irritant cathartic.

PREPARATION.

Fluid Extract Thapsia Garganica.—Not miscible with water.
Dose, 10 to 30 minims (0.6 to 2 C.c.).

THIMBLE-WEED, Rudbeckia laciniata, Linné. (Herb.)

Synonym.—Cone-flower.

Properties.—Diuretic, tonic and balsamic. It is especially recommended in chronic catarrhal inflammation of the urinary tract.

Thimble-weed herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Thimble-weed.—Miscible with water without material precipitation.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

THYME, Thymus vulgaris, Linné. (Leaves.)

Synonyms.—Garden Thyme; Römischer Quendel, Ger. Origanum (oil).

Properties.—Tonic, emmenagogue and antispasmodic. The essential oil is also an antiseptic of some value, due to the presence of thymol.

Thyme leaves, compressed in ounce packages for retailing purposes.

ACTIVE PRINCIPLE.

Thymol.—Crystals. As an antiseptic more powerful, more permanent, and less poisonous than carbolic acid. Its fragrance also renders it more acceptable in the sick-room. Solubility: 1 in 1200 of water, sufficiently strong to use as a dressing, etc.; 1 in 1 of alcohol. Internally it checks flatulence by hindering fermentation.
Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

PREPARATIONS.

Capsules Origanum Oil.—In soft gelatin. Each containing:
Oil Origanum, 5 minims. Oil Sweet Almond, 5 minims.

Fluid Extract Thyme.—Miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

TOBACCO, Tabacum, U. S. P. Nicotiana Tabacum, Linné. (Lvs.)

Properties.—Poisonous. A potent acro-narcotic. It possesses sedative, emetic, diuretic, expectorant, discutient and antispasmodic properties. Has been used for renal dropsy, tetanus, strangulated hernia, obstinate constipation from spasm of the bowels, etc. Used as an ointment in tinea capitis, psora, and other cutaneous affections, indolent tumors, etc. Care must be observed in its use. It is an efficient physiological antagonist in strychnine poisoning.

Antidotes.—Use an emetic, and after the evacuation of the poison, support the system by internal and external stimulants, allaying the irritation of the stomach with opiates.

PREPARATION.

Fluid Extract Tobacco.—Miscible with water.
Dose, 5 minims (0.3 C.c.) gradually increased.

TOLU. Balsamum Tolutanum, U. S. P. From Myroxylon Toluiferum, *Humboldt, Bonpland & Kunth.*

Properties.—Stimulant in chronic and subacute inflammations of the bronchial mucous membrane.

PREPARATIONS AND FORMULA.

Fluid Tolu, Soluble.—Miscible with water.

Containing, in a concentrated form, those constituents of balsam of tolu which give flavor and medicinal virtue to the official syrup, which may be prepared from it instantaneously, according to the formula below. It commends itself especially to the prescriber as being much more concentrated than the syrup, and hence particularly useful in extemporaneous cough mixtures.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Lozenge Bronchial (with Cubeb).—Each containing:

Ext. Licorice, 4-5 gr.

Powd. Cubeb, 5-8 gr.

Balsam Tolu, 1-20 gr.

Oil Sassafras, 1-40 gr.

Syrup Tolu, U. S. P.—Representing 24 grains Balsam Tolu to the fl. oz.

Prepared extemporaneously as follows:

Fluid Tolu, soluble, P., D. & Co.'s..... 1 fl. oz.

Simple Syrup..... 15 fl. ozs.

Dose, ad libitum.

TOMATO. Lycopersicum esculentum, *Miller.* (Fruit.)

Synonym.—Love Apple.

Properties.—The fluid extract is claimed to exert a special curative action over ulcerative affections of the mucous lining of the mouth and other cavities, "nurse's sore mouth," "canker," etc. It is to be given internally, and also applied locally to the affected parts.

PREPARATION.

Fluid Extract Tomato.—Miscible with water without material precipitation.

Dose, 30 to 60 minims, (2 to 4 C.c.).

TONCA-BEAN. Dipterix odorata, *Willdenow.* (Seeds.)

Properties.—Chiefly used as a flavoring agent for tobacco; medicinally for external use it is claimed to possess an anodyne effect.

PREPARATION.

Fluid Extract Tonca (Tonka)-bean.—Miscible with water without material precipitation.

TONGA. { *Premna Taitensis, Schauer.* } (Bark and root.)
 { *Raphidophora vitiensis, Schott.* }

Properties.—Long employed by the natives of the Fiji Islands as a remedy for neuralgia. Placed on the market as a remedy in neuralgic affections on the recommendation of Drs. Ringer and Murrell, of London, who have made very thorough experiments to determine its therapeutic value.

"The result of the experiments of Drs. Ringer and Murrell, corroborated by the trials of many practitioners in this country [France], demonstrate conclusively the great value of this remedy in neuralgic affections, especially in those of the cranial nerves."*

PREPARATIONS.

Fluid Extract Tonga.—Not miscible with water.

Dose, 30 minims to 1 fluidrachm (2 to 4 C.c.).

Powdered Extract Tonga.—One part equals 7 of drug.

Prepared by evaporating, at a low temperature, the solid extract, and replacing moisture with powdered Tonga.

Dose, 2 to 6 grains (0.13 to 0.4 Gm.).

* Dujardin-Beaumez, "New Medications," p. 289, (edition of Geo. S. Davis, 1886).

TORMENTILLA. *Potentilla Tormentilla, Scopoli.* (Rhizome.)

Properties.—Tonic and powerfully astringent; used for constringing relaxed tissues, arresting passive hæmorrhages, chronic diarrhœa, and as a wash for indolent ulcers.

Tormentilla root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Tormentilla.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

TRUMPET PLANT. *Sarracenia flava, Linné.* (Rhizome.)

Synonyms.—Fly-trap, Side-saddle plant.

Properties.—This plant, a native of Florida, was brought to public notice by Dr. J. Dabney Palmer, of Monticello, Florida, who having experimented with the root, ascertained that it possesses extraordinary power in the cure of diarrhœa. Some of the cases in which it was given were of long standing, and very obstinate, others recent, but all yielded to the remedy. A few doses were generally sufficient. In cases of uncomplicated diarrhœa this remedy has seldom failed to effect a speedy and permanent cure, although it has no influence over the course of dysentery. It has been found useful in dyspepsia, sick-headache, gastralgia, abdominal distention, etc.

PREPARATION.

Fluid Extract Trumpet Plant.—Not miscible with water.

Dose, 5 to 15 minims (0.3 to 1 C.c.).

TULIP TREE. *Liriodendron tulipifera, Linné.* (Bark.)

Synonyms.—White-wood, Yellow Poplar.

Properties.—Aromatic, stimulant, tonic and antiperiodic. Used in intermittents, chronic rheumatism and hysteria.

Tulip-tree bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Tulip Tree.—Miscible with water without material precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

TURKEY-CORN. *Dicentra canadensis, DeCandolle.* (Tubers.)

Synonyms.—Wild Turkey Pea, Stagger-weed, Squirrel corn; *Corydalis canadensis, Goldie.*

Properties.—Tonic, diuretic and alterative. In syphilitic affections and in scrofula, in which a combined tonic and alterative influence is indicated, it is valuable. Its tonic properties resemble those of gentian, columbo, and other simple bitter tonics.

Turkey-corn root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Corydalin, a so-called **Concentration**,* a powdered preparation containing all the active constituents of the Turkey corn, including its resins and the alkaloid corydaline.

Dose, 1 to 5 grains (0.06 to 0.3 Gm.).

Elixirs.—Each fluidounce represents ingredients specified.

Corydalis Compound.

Turkey corn, 30 grs

Stillingia, 30 grs.

Blue Flag 40 grs.

Prickly Ash, 15 grs.

Potassium iodide, 8 grs.

Stillingia Compound.

Stillingia, 30 grs.

Turkey corn, 30 grs.

Pipsissewa, 15 grs

Elder flowers, 15 grs.

Blue Flag, 15 grs.

Prickly-ash berries, 1-2 grs

Coriander seed, 7 1-2 grs,

* See note in Preface on Concentrations.

TURKEY-CORN—Continued.**Extracts.**

Fluid Extract Stillingia Comp.—Not miscible with water.

Formula same as Elixir Stillingia Comp. above, but 4 times stronger.
Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

Fluid Extract Turkey Corn.—Miscible with water without material precipitation.

Dose, 10 to 40 minims (0.6 to 2.5 C. c.).

Syrups.—Each fluidounce represents ingredients specified.

Corydalis Compound.

Turkey corn, 30 grs.

Stillingia, 30 grs.

Blue Flag, 40 grs.

Potassium iodide, 8 grs.

Stillingia Compound.—Formula same as Elixir above.

TURMERIC. *Curcuma longa, Linné.* (Rhizome.)

Properties.—An aromatic stimulant. At present it is only employed to impart color to pharmaceutical preparations and as a condiment in curries.

PREPARATION.

Fluid Extract Turmeric.—Not miscible with water without some precipitation.

Dose, 15 to 60 minims (1 to 4 C. c.).

TURPENTINE. An oleoresin obtained principally from the *Pinus australis, Michaux.*

Properties.—Seldom used; of value medicinally on account of its volatile oil (extracted by distillation). The Oil of Turpentine employed externally is a valuable rubefacient and counter-irritant. Internally it is valuable as an anthelmintic for the expulsion of tæniæ, combined with an efficient dose of castor oil, as a hæmostatic, in paralysis of the bladder, catarrh of the urinary passages, tympanites, etc. Best exhibited in gelatin capsules.

Dose, as hæmostatic, diuretic, etc., 5 to 30 minims (0.3 to 2 C. c.); as an anthelmintic 1-2 to 2 fluidounces (15 to 60 C. c.).

DERIVATIVES.

Terebene.—Prepared by the action of sulphuric acid on oil of turpentine.

Its properties are similar to those of the ordinary oil, but its odor is entirely different and not disagreeable. Dr. Murrell, of London, regards it as one of the best remedies for winter cough. It is expectorant and antiseptic, and of value both internally and in form of a spray. It is a remedy also for flatulence.

Dose, 5 to 20 minims (0.33 to 1.33 C. c.).

Terpin Hydrate.—From oil of turpentine by the joint action upon it of alcohol and nitric acid. Crystalline.

Sparingly soluble (1:200) in water, more freely in a mixture of alcohol and water; of a terebinthinate taste. It is said to be a very valuable expectorant especially in chronic bronchitis, also to act as a diuretic, liable, however, to produce albuminuria.

Dose, 3 to 9 grains (0.2 to 0.6 Gm.).

PREPARATIONS.**Capsules.**

Terebene, 10 minims — In soft gelatin.

Turpentine Oil, 10 minims. — In both soft and hard gelatin.

Inhalant Turpentine.—No. 20 of P., D. & Co.'s series of inhalants.

Terebene, 1 fluidrachm.

Oil Sandalwood, 1 fluidrachm.

Oil Cubeb, 1 fluidrachm.

Neutral Petroleum oil, 5 fl. drms.

Lozenge Terebene.—Each containing 2 minims.

TWIN-LEAF. *Jeffersonia diphylla, Persoon.* (Rhizome.)

Synonyms.—Ground-squirrel Pea, Rheumatism Root.

Properties.—Somewhat similar to senega: diuretic, alterative, antispasmodic, and a stimulating diaphoretic. It has been successfully used in secondary syphilis, in dropsy, many nervous affections, spasms, cramps, nervous excitability. As a gargle it has been beneficial in diseases of the throat, ulcers about the fauces, scarlatina, ophthalmia, and indolent ulcers.

Twin-leaf root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Twin-leaf.—Miscible with water without material precipitation.

Dose, 20 to 60 minims (1.3 to 4 C[°])

URECHITES SUBERECTA, J. Mueller. (Leaves.)

Synonyms.—Savana Flower, Yellow-flowered Nightshade; *Echites suberecta, Jacquin.*

Properties.—**POISONOUS.** According to Dr. Isaac Ott (*Therapeutic Gazette*, Oct., 1880), the effects of the drug bear a general resemblance to those of aconite. The symptoms observed are depression of circulation, salivation, diaphoresis, vomiting, diarrhoea. It has been tried with remarkable success in Jamaica in the treatment of intermittent fever, and it is also claimed to be of great value in yellow fever.

ANTIDOTES.—Use emetics or stomach-pump promptly. Sustain vitality by stimulants—alcohol, ammonia, strong coffee, the battery, etc. Digitalis and Strophanthus have been suggested as in the case of Aconite.

PREPARATION.

Fluid Extract Urechites Suberecta.—Not miscible with water without some precipitation

Dose, 2 to 10 minims (0.13 to 0.65 C.c.) cautiously increased.

UVA-URSI, U. S. P. *Arctostaphylos Uva-ursi, Sprengel.* (Leaves.)

Synonyms.—Bearberry, Mountain Cranberry, Mountain Box; Busserole, Fr.; Bärentraube, Ger.; Gayuba, Span.

Properties.—Astringent and tonic with a specific direction to the urinary organs. It has some reputation as an antilithic, and has undoubtedly been serviceable in gravel. It is used in chronic nephritis, catarrh of the bladder, incontinence of urine, gleet, leucorrhœa and menorrhagia.

ACTIVE PRINCIPLE.

Arbutin.—Glucoside. Bitter; soluble in water and alcohol. Treated with acids it is split into glucose and hydrochiæone, and a similar change takes place in the body when ingested. It is diuretic, and relieves catarrhal conditions of the bladder, and even renal hemorrhages. It exercises a distinct antiseptic influence in lessening the formation of pus in cases of purulent inflammation of the bladder and kidney.*

PREPARATIONS.

Elixirs.—Each fluidounce represents ingredients specified.

Buchu Compound.

Buchu, 60 grs.
Cubeb, 15 grs.
Juniper, 15 grs.

Uva-ursi, 15 grs.
Spirits nit. eth., 15 minims.

Matico Compound.

Matico, 40 grs.
Uva-ursi, 40 grs.

Hydrangea, 40 grs.

* *Therapeutic Gazette*, p. 618, 1886.

UVA-URSI—Continued.**Extracts.**

Fluid Extract Buchu Compound.—Not miscible with water.
Dose, 15 to 30 minims (1 to 2 C.c.).

Four times stronger than Elixir Buchu Comp. above, same formula.

Fluid Extract Uva-ursi, U. S. P.—Not miscible with water.
Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Solid Extract Uva-ursi.—One part equals 4 of the leaves.
Dose, 5 to 15 grains (0.3 to 1 Gm.).

Pill Uva-ursi Extract, 2 grs.

VACCINIUM CRASSIFOLIUM, Andrzejowski. (Plant.)

Properties.—Astringent and diuretic, resembling uva-ursi to some degree.

Useful in catarrhal inflammation of the genito-urinary tract, gleet, chronic cystitis, etc. In dropsy, from whatever cause, it is an active diuretic, assisting the removal of the effused fluid. It has also been given with benefit in chronic diarrhoea and dysentery.

PREPARATION.

Fluid Extract Vaccinium Crassifolium.—Not miscible with water.
Dose, 30 minims to 1 fluidrachm (2 to 4 C.c.).

VALERIAN, U. S. P. Valeriana officinalis, Linné. (Rhizome.)

Synonym.—Baldrain, Ger.

Properties.—Gently stimulant, with an especial direction to the nervous system, but without narcotic effects. It is useful in cases of irregular nervous action, when not connected with inflammation or an excited condition of the system. Recommended in hysteria, hypochondria, hemi-crania, delirium tremens and low forms of fever attended with restlessness, morbid vigilance and other nervous disorders. It has also been used with advantage in the treatment of intermittents and acute rheumatism (in hot water baths).

Valerian root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULÆ.**Abstract Valerian, U. S. P.**

Fluid Extract Valerian, P., D. & Co.'s 1 fl. oz.
Sugar of milk, q. s. to make 228 grains.
Evaporate the fluid nearly to dryness by a very gentle heat; add 175 grains of the sugar of milk; incorporate thoroughly with the extract; reduce to complete dryness; weigh, and add sufficient sugar of milk to bring the weight to 228 grains; finally powder the product. Keep in a well-corked bottle.

Capsule Valerian Oil.—In soft gelatin, each containing:

Oil Valerian, 2 minims. Oil Sweet Almond, 8 minims.

One a dose in hysteria, chorea, etc.

Elixir Valerian, 120 grains to the fluidounce.

Extracts.

Fluid Extract Valerian, U. S. P.—Not miscible with water without some precipitation.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Powdered Extract Valerian.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Valerian.
Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

Solid Extract Valerian.—One part equals 5 of drug.
Dose, 5 to 10 grains (0.3 to 0.65 Gm.).

VALERIAN—Continued.**Pills.****Anti-Epileptic, 3 grs.**

Iron ferrocyanide, 1-2 gr.

Quinine valerianate, 1 gr.

Zinc valerianate, 1-2 gr.

Ext. Valerian, 1 gr.

Camphor, Valerian and Henbane, 2 grs.

Camphor, 1 gr.

Powd. Valerian, 1-2 gr.

Ext. Henbane, 1-2 gr.

Phosphorus, Zinc and Valerian, 3 grs.

Phosphorus, 1-40 gr.

Zinc sulphate, 1 gr.

Ext. Valerian, 2 grs.

Sedative, 2 grs.

Ext. Musk root, 1-2 gr.

Ext. Henbane, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Indian Cannabis, 1-10 gr.

Sedative, Mann's, 2 1-2 grs.

Ext. Coca, 1-2 gr.

Ext. Valerian, 1-2 gr.

Ext. Indian Cannabis, 1-4 gr.

Vallet's Mass, 1 gr.

Arsenious acid, 1-80 gr.

Strychnine, 1-80 gr.

Codeine, 1-8 gr.

Valerian Extract, 2 grs.**Tinctures.****Valerian, U. S. P.**

Fluid Extract Valerian, P., D. & Co.'s.....2 fl. ozs.

Alcohol.....9 fl. ozs.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

NOTE.—Uniform in strength with the tinctures of the Phar. Ger. and Fr. Cod.

Valerian Ammoniated, U. S. P.

Fluid Extract Valerian, P., D. & Co.'s.....3 fl. ozs.

Aromatic Spirit of Ammonia.....14 fl. ozs.

Dose, 30 to 60 minims (2 to 4 C.c.) diluted.

Antispasmodic and stimulant.

VIRGINIA STONE-CROP. *Penthorum sedoides, Linné.* (Herb.)

Properties.—Astringent, demulcent, laxative. That class of diseases characterized by affection of the mucous membrane of the throat, nares, etc. —pharyngitis, catarrh, laryngitis—are relieved by this remedy. Recommended in diarrhoea (in 1-2 fluidrachm doses), chronic bronchitis with increased secretion, and as an injection in chronic vaginitis and diseases of the cervix uteri.

PREPARATION.**Fluid Extract Virginia Stone-crop.**—Not miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C. c.).

WAFER ASH. *Ptelea trifoliata, Linné.* (Bark of root.)

Synonyms.—Wingseed, Shrubby Trefoil, Hop Tree; Hopfenbaum, Kleebaum, Ger.

Properties.—Tonic, antiperiodic, and reputed anthelmintic. It is recommended highly in the debility following low fevers, especially those connected with gastro-intestinal irritation, to which it seems to be particularly applicable from its mild and non-irritating qualities. It exercises a soothing influence on the stomach, which receives it when rejecting other tonics; while it promotes the appetite, invigorates digestion, and helps greatly to re-establish health and convalescence.

Wafer Ash bark of root, compressed in pound packages for retailing purposes.

PREPARATIONS.**Elixirs.**—Each fluidounce represents ingredients specified.**Pepsin, Bismuth and Wafer Ash.**

Sacch. Pepsin, U. S. P., 8 grs.

Bismuth and Ammon. cit., 8 grs.

Wafer Ash, 60 grs.

Pepsin and Wafer Ash.

Sacch. Pepsin, U. S. P., 10 grs.

Wafer Ash, 60 grs.

Wafer Ash, 120 grs.

WAFER ASH—Continued.**Fluid Extract Wafer Ash.**—Not miscible with water.

Dose, 10 to 30 minims (0.6 to 2 C.c.).

Glycerole Pepsin and Wafer Ash.

Sacch. Pepsin, U. S. P., 10 grs.

Wafer Ash, 30 grs.

Of service where the absence of all alcohol is desirable.

Ptelein, a so-called **Concentration** * a powdered preparation from

Wafer Ash bark of root.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

WAHOO. *Euonymus*, U. S. P. *Euonymus atropurpureus*, *Jacquin*.
(Bark of root, bark of tree.)*Synonyms*.—Spindle Tree, Burning Bush.*Properties*.—The experiments of Rutherford have fixed this drug as one of the most direct cholagogues in the materia medica. It is also tonic, alterative and laxative, and is especially indicated in hepatic derangements, whether accompanying or preceding intermittents, or occurring independently of malaria. In constipation due to hepatic torpor it is highly recommended. It is not nearly so active an irritant of the intestines as Podophyllin.**Wahoo, bark of root, U.S.P.,** } compressed in pound packages for
Wahoo, bark of tree, U.S.P., } retailing purposes.**PREPARATIONS.****Elixirs.**—Each fluidounce represents ingredients specified.**Wahoo**, bark of root, 160 grs.**Wahoo and Blue Flag.**

Wahoo root bark, 80 grs.

Blue Flag, 40 grs.

Euonymin, a so-called **Concentration**. *—A powdered preparation from Wahoo: *Euonymin brown* being from bark of root, *Euonymin green* from the bark of the twigs.

Dose, 1-2 to 4 grains (0.03 to 0.26 Gm.).

Extracts.**Fluid Extract Wahoo.**—Not miscible with water without some precipitation.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Powdered Extract Wahoo.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Wahoo.

Dose, 5 to 15 grains (0.3 to 1 Gm.).

Solid Extract Wahoo, U. S. P.—One part equals 4 of drug.

Though the U. S. P. does not so specify, this extract is prepared from the bark of the root, without any addition of the bark of the twigs, as the former is much richer in active principle and results in a more active and desirable preparation.

Dose, 5 to 15 grains (0.3 to 1 Gm.).

Pellets Cascara Compound, D'Ary's, 1 gr.

Ext. Cascaraagrada, 4-15 gr.

Euonymin, 1-5 gr.

Ext. Nuxvomica, 1-30 gr.

Xanthoxyllin, 4-15 gr.

Ext. Belladonna, -60 gr.

Oleoresin Capsicum, 1-20 gr.

WATER AVENS. *Geum rivale*, *Linné*. (Root.)*Synonyms*.—Avens, Purple Avens.*Properties*.—Very astringent, and more or less tonic. Apt to derange the digestion and cause vomiting if too freely given. Employed in all forms of disease of the mucous membranes depending upon relaxation and attended with excessive and altered secretions, including atonic dyspepsia, diarrhoea, bronchorrhoea, leucorrhoea, etc. It has also been used in menstuous derangements depending on debility. If administered in the form of infusion or decoction it may be taken ad libitum.**Water Avens root**, compressed in pound packages for retailing purposes.**PREPARATION.****Fluid Extract Water Avens root.**—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

* See note in Preface on Concentrations.

WATER ERYNGO. *Eryngium yuccæfolium, Michaux.* (Rhizome.)

Synonyms.—*Eryngium aquaticum, Linné*; Rattlesnake's master, Corn-snake root, and occasionally as Button-snake root, although this is properly *Liatris spicata, Willdenow*.

Properties.—Diaphoretic, expectorant, in large doses occasionally emetic; used by some physicians as a substitute for senega, and is said to be diuretic and stimulant. Employed in dropsy, nephritis and calculous affections, also in scrofula and syphilis; as a diaphoretic and expectorant in pulmonary diseases, chronic laryngitis and bronchitis. It is said to exert an influence on the reproductive organs, lessening erectile power, and proving effectual in cases of seminal losses with urethral irritation pain in the testes, and irritation of the bladder.

Water Eryngo, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Water Eryngo.—Miscible with water.

Dose, 20 to 40 minims (1 3 to 2.6 Gm.).

WATER FENNEL. *Oenanthe Phellandrium, Lamarck.* (Seed.)

Synonyms.—Water Dropwort, Five-leaved Water Hemlock; Fenouil d'eau, Fr.; Wasser fenchel, Ger.

Properties.—A mild narcotic stimulant, expectorant, alterative, and diuretic.

In large doses it produces dizziness, inebriation, and dull pains in the head. Medicinally used, this drug is of value in chronic affections of the air passages, as bronchitis, laryngitis, asthma, hemoptysis, catarrh, etc., and in periodical febrile diseases, dyspeptic affections, and indolent ulcerations. In consumption and bronchitis, it is said to silence troublesome cough, to render expectoration smaller and easier, and to produce sleep at night.

PREPARATION.

Fluid Extract Water Fennel.—Not miscible with water without some precipitation.

Dose, 4 to 12 minims (0.25 to 0.75 C c.), cautiously increased.

WATER GERMANDER. *Teucrium Scordium, Linné.* (Herb.)

Synonyms.—Germandrée Aquatique, Fr.; Lachenknoblauch, Ger.

Properties.—The ancient writers on therapeutics attribute to this plant extraordinary virtues as an alexipharmic, sudorific and antiseptic. It is used as a corroborant in low forms of disease, and in cases where there is want of tone in the stomach, in flatulence, anasarca, amenorrhœa and humid asthma.

PREPARATION.

Fluid Extract Water Germander.—Miscible with water without material precipitation.

Dose, 1-2 to 1 fluidrachm (2 to 4 C c.).

WATER HEMLOCK. *Cicuta maculata, Linné.* (Leaves.)

Synonyms.—American Water Hemlock, Beaver-poison, Musquash root, Spotted Cowbane, Spotted Parsley, Spotted Water Hemlock.

Properties.—POISONOUS. Often confounded with conium, and has been employed as a substitute for that sedative, but its action appears to be entirely different, and therapeutic uses have not yet been established. It is recommended as a specific in nervous- and sick-headache.

ANTIDOTES.—Empty the stomach thoroughly by use of prompt emetics, and then support the system with stimulants.

NOTE.—Physicians sometimes write *cicuta* when they want conium; before dispensing the *cicuta* the druggist should assure himself that it is the remedy intended by the prescriber.

WATER HEMLOCK—Continued.**PREPARATION.**

Fluid Extract Water Hemlock.—Miscible with water without material precipitation.

Dose, 3 to 15 minims (0.2 to 1 C.c.).

WATER-MELON. *Citrullus vulgaris*, *Schrader*. (Seed.)

Synonyms.—*Cucumis Citrullus*, *Séringe*; *Cucurbita Citrullus*, *Linne*.

Properties.—Diuretic and demulcent. Useful in all cases of irritation of the kidneys or bladder, and especially in those of retention of urine produced by cold.

PREPARATION.

Fluid Extract Water-melon seed.—Miscible with water without material precipitation.

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

WATER-PEPPER. *Polygonum acre*, *Humboldt*, *Bonpland* & *Kunth*. (Herb.)

Synonyms.—*Polygonum punctatum*, *Elliott*; *Polygonum hydropiperoides*, *Pursh*; Smartweed, Water Smartweed.

Properties.—Stimulant, diuretic emmenagogue, antiseptic, diaphoretic and vesicant. It has been found efficacious in amenorrhœa (20 drops 4 or 5 times a day), gravel, colic, and colds. Externally it is said to be of value in choleraic seizures (the patient being wrapped in a sheet previously soaked in a hot decoction of the drug), in gangrene, chronic ulcers, hemorrhoidal tumors, chronic erysipelatous inflammations, and (as a fomentation) in tympanitis, and flatulent colic. As a wash in sore mouth of nursing women, and mercurial pytalism.

Water-pepper herb, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract Blue Cohosh Comp.—Not miscible with water.

Nine fluidounces, imperial, represents: Blue Cohosh, 4 ozs. av.; Ergot and Water-pepper, 2 ozs. each; Savin, 1 oz. A useful remedy in amenorrhœa, dysmenorrhœa, and other uterine affections.

Dose, 5 to 20 minims (0.35 to 1.3 C.c.).

Fluid Extract Water-pepper.—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

WHITE AGARIC. *Polyporus officinalis*, *Fries*. (The fungi.)

Synonyms.—Larch agaric, Purging agaric, Touchwood, Spunk; *Boletus laricis*, *Linne*; Agaric blanc, *Fr.*; Lärchenschwamm, *Ger.*

Properties.—In doses of 8 grains, gradually increased to a drachm, it has been found useful in checking the night-sweats of phthisis. It appears to restrain all the secretions, and is thus especially useful in the later stages of consumption. It is said also to hasten the drying up of the milk in weaning. In large doses it acts as a cathartic, but is not now employed for this purpose.

ACTIVE PRINCIPLE.

Agaricin.—Recently strongly recommended for the relief of the night-sweats of phthisis. Best given in pills. Its effects are produced only after an interval of five or six hours.

Dose, 1-15 to 1-6 grain (0.004 to 0.011 Gm.).

PREPARATIONS.

Fluid Extract White Agaric.—Not miscible with water.

Dose, 3 to 20 minims (0.2 to 1.3 C.c.) and upwards.

Tablet Triturate Agaricin, 1-20 gr.

WHITE ASH. *Fraxinus americana*, Linné. (Bark.)*Synonym.*—American ash.

Properties.—Introduced by Dr. Chas. P. Turner, as a remedy for many forms of uterine disease. This author advocates the theory that many uterine difficulties have as their foundation pathological plethora of the uterus. The plethora diminishes the cavity of the uterus by inward pressure, and lessens the calibre of the cervical canal, hence amenorrhœa or dysmenorrhœa. White Ash, according to Dr. Turner, attacks the benumbed or torpid vaso-motor nerve system, arouses it to new vitality, acts mildly but persistently in its secondary effects upon the absorbents, thus gradually reducing the size of the enlarged uterus, and bringing it back to its normal condition.

White Ash bark, compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.

Fluid Extract White Ash.—Miscible with water.

Dose, 1-2 to 4 fluidrachms (2 to 15 C. c.).

Wine White Ash.

Each fluidounce represents 120 grains White Ash bark.

Dose, a dessertspoonful (8 C. c.) and upwards

A medicinal equivalent for our preparation may be prepared extemporaneously as follows:

Fluid Extract White Ash bark, P., D. & Co.'s.....	11 fl. ozs.
Sherry Wine.....	29 fl. ozs.

WHITE BRYONY. *Bryonia*, U.S.P. *Bryonia alba*, Linné. (Root.)*Synonyms.*—Bryonia: Couleuvrée, Fr.; Zaubrübe, Gichrübe, Ger.

Properties.—Hydragogue cathartic, in large doses an irritant poison. Chiefly employed as a hydragogue in dropsy, resembling jalap in action. Also recommended in chronic intermittent fever with enlargement of the spleen, in chronic bronchitis, and in the catarrhal stages of whooping-cough.

White Bryony root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.

Fluid Extract White Bryony.—Not miscible with water without some precipitation.

Dose, 10 to 60 minims (0.6 to 4 C. c.).

"German Tincture" White Bryony.—An imported unofficial preparation from the green root

Dose, 5 to 15 minims (0.3 to 1 C. c.).

Tablet Triturate Bryony Tincture, 5 minims.

Tincture White Bryony, U. S. P.

Fluid Extract White Bryony, P., D. & Co.'s.....	2 fl. ozs.
---	------------

Alcohol	21 fl. ozs.
---------------	-------------

Dose, 1 to 2 fluidrachms (4 to 8 C. c.).

WHITE CLOVER. *Trifolium repens*, Linné. (Tops.)*Synonyms.*—White Trefoil, Dutch Clover, Shamrock; Kriechender weisser Klee, Ger.; Trèfle blancourampant, Fr.

Properties.—Employed in whooping-cough and other varieties of spasmodic cough. Generally used in the form of an infusion, one ounce to the pint.

Dose, ad libitum.

White Clover tops, compressed in ounce packages for retailing purposes.

WHITE COHOSH. *Actæa alba*, Bigelow. (Rhizome.)*Synonym.*—White Blueberry.

Properties.—Violently purgative, sometimes producing gastro-intestinal disorders by its irritant and emetic action.

White Cohosh root, compressed in pound packages for retailing purposes.

WHITE HELLEBORE. *Veratrum album*, *Linné*. (Rhizome.)

Properties.—**POISONOUS**. A vaso-motor depressant. Used in cases of irritable heart, hypertrophy of the cardiac muscle, etc. A violent irritant, occasioning when snuffed up in the nostrils, severe coryza, and when swallowed, severe vomiting, with profuse diarrhoea. When it proves fatal, narcotic symptoms are superadded. Great care should be taken in its use even externally. An infusion of 1-2 ounce to the pint is used as a wash in parasitic diseases of the skin, etc.

ANTIDOTES.—Coffee, as a drink, stimulants to overcome the depressed condition, and opiates and demulcents to relieve internal inflammation. Recumbent position.

White Hellebore root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract White Hellebore.—Not miscible with water.

Dose, 2 to 4 minims (0.12 to 0.24 C.c.).

WHITE INDIAN-HEMP. *Asclepias incarnata*, *Linné*. (Rhizome.)

Synonyms.—Swamp Silkweed, Flesh-colored *Asclepias*, Rose-colored Silkweed, Swamp Milkweed.

Properties.—**Anthelmintic**. It has been recommended as efficacious in rheumatic, asthmatic, catarrhal and syphilitic affections.

White Indian-hemp root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract White Indian-hemp.—Not miscible with water without some precipitation.

Dose, 10 to 40 minims (0.6 to 2.5 C.c.).

Solid Extract White Indian-hemp.—One part equals 5 of drug.

Dose, 2 to 8 grains (0.13 to 0.5 Gm.).

WHITE OAK. *Quercus alba*, *Linné*. (Bark.)

Synonyms.—Écorce de Chêne, Fr.; Eichenrinde, Ge.

Properties.—**Astringent** and somewhat tonic. Used internally with advantage in obstinate chronic diarrhoea and certain forms of passive hemorrhage. Externally used as an astringent wash, gargle or injection.

White Oak bark, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract White Oak bark.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract White Oak.—One part equals 5 of bark.

Dose, 3 to 10 grains (0.2 to 0.65 Gm.).

WHITE PINE. *Pinus Strobus*, *Linné*. (Bark.)

Properties.—Those of balsamic preparations generally. Useful as an emollient and expectorant in chronic affections of the air passages, and has been proved of service in catarrh of the bladder.

White Pine bark, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract White Pine bark.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

WHITE POND-LILY. *Nymphaea odorata, Aiton.* (Rhizome.)

Synonym.—White Water-lily.

Properties.—Astringent, demulcent, anodyne and anti-scrofulous. Used in dysentery, diarrhoea, gonorrhoea, leucorrhoea and scrofula, and, combined with wild cherry, in bronchial affections. Externally it may be applied to boils, tumors, scrofulous ulcers and inflamed skin. The fluid extract diluted is used as a gargle in ulcers of the mouth and throat, and as an injection in leucorrhoea.

White Pond-lily root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract White Pond-lily.—Not miscible with water
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

WHITE POPLAR. *Populus tremuloides, Michaux.* (Bark.)

Synonyms.—Poplar, American Aspen.

Properties.—Tonic and febrifuge; used in intermittent fever with advantage. It contains an active principle, populin, analagous to the salicin of willow bark. Diuretic properties are also ascribed to the bark, which is accordingly employed with asserted advantage in urinary affections, gonorrhoea, gleet, etc.

White Poplar bark, compressed in ounce packages for retailing purposes.

PREPARATIONS.

Fluid Extract White Poplar bark.—Not miscible with water without some precipitation.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Populin, a so-called **Concentration**,* a powdered preparation from White Poplar; not to be confounded with the crystallizable glucoside of that name which may be extracted from it.
Dose, 2 to 5 grains (0.13 to 0.33 Gm.).

WHITE SNAKE-ROOT. *Eupatorium aromaticum, Linné.* (Rhiz.)

Properties.—Diaphoretic, antispasmodic, and expectorant. It is used with advantage in pleurisy, chronic bronchitis, nervous irritability and flatulence, hysteria, in the wakefulness of low fevers of a typhoid character, and in chronic irritation of the bladder.

White Snake-root, compressed in pound packages for retailing purposes.

WHITE VERVAIN. *Verbena urticæfolia, Linné.* (Root.)

Synonym.—Nettle-leaved Vervain.

Properties.—Has been successfully used in the treatment of intermittent and remittent fever. It is credited with cure of the opium habit.

PREPARATION.

Fluid Extract White Vervain.—Miscible with water.
Dose, 30 to 40 minims (2 to 2.5 C.c.).

WHITE WILLOW. *Salix, U. S. P. Salix alba, Linné.* (Bark)

Properties.—Tonic, febrifuge, and astringent; possessing also antimalarial properties which have led to its use in intermittents with no little success. Its active principle, salicin, along with the derivative, salicylic acid, have, however, to a large extent taken the place of the bark on account of more condensed dosage: they have been used advantageously in rheumatism.

White Willow bark, U. S. P., compressed in ounce packages for retailing purposes.

* See note in Preface on Concentrations.

WHITE WILLOW—Continued.**ACTIVE PRINCIPLE (derivative).****Salicylic Acid, pure.**—Kolbe's process.**Salicylic Acid, pure, from Oil of Wintergreen.**—Although identical in chemical composition, these two products differ materially in appearance and flavor, and many physicians prefer the latter, as an article nature provides ready-made, to the former, an artificial imitation. Dose, 5 to 20 grains (0.32 to 1.4 Gm.).**PREPARATIONS.****Fluid Extract White Willow bark.**—Miscible with water. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).**Lozenge Salicylic Acid,** 1 grain.**Pills.****Night Sweat,** 3 grs.

Zinc oxide, 1-2 gr.

Salicin, 1 gr.

Ext. Belladonna, 1-25 gr.

Hydrastin, 1 gr.

Lactated Pepsin, 1-2 gr.

Salicin, 1, 2, 3, and 5 grs.**Salicylic Acid,** 1-10, 1, 2 1-2, 3, and 5 grs.**Tablet Triturates, Salicylic Acid,** 1-4 and 1-2 gr.**WILD BERGAMOT.** *Monarda fistulosa, Linné.* (Leaves.)*Synonym.*—Horsemint.*Properties.*—First introduced as a substitute for quinine in intermittents, and its use has been reported in a number of cases without a failure. In large doses it is a diaphoretic, and said to be inferior only to jaborandi in its power of exciting the sweat glands. It is used in flatulent colic, cholera infantum, and lower forms of fever, as also in liniments for relief of local paralyses.**PREPARATION.****Fluid Extract Wild Bergamot.**—Miscible with water. Dose, 15 to 60 minims (1 to 4 C.c.).**WILD CHERRY.** *Prunus virginiana, U.S.P.* *Prunus serotina, Ehrhart.* (Bark.)*Synonyms.*—*Cerasus serotina, Loiseleur-Deslongchamps; Prunus virginiana, Miller; Wildkirschenrinde, Ger.**Properties.*—Tonic and stimulating to the digestive apparatus, and possessing simultaneously a sedative action on the nervous system and circulation. Valuable in all cases where it is desirable to give tone and strength to the system without causing too great heart-action and strain on the blood-vessels, as during convalescence from pleurisy, pneumonia, acute hepatitis, etc. It is also useful in hectic fever, colliquative diarrhoea, some forms of dyspepsia, whooping-cough, irritability of the nervous system, and as a palliative in phthisis.**Wild Cherry bark, U.S.P.,** compressed in pound packages for retailing purposes.**PREPARATIONS AND FORMULÆ.****Elixirs.**—Each fluidounce represents ingredients specified.**Dandelion Compound.**

Dandelion, 10 grs.

Cherry, 10 grs.

Licorice, 20 grs.

Aromatics, q. s.

Eucalyptus Compound.

Eucalyptus globulus, 15 grs.

Wild Cherry, 16 grs.

Gentian, 4 grs.

Licorice, 15 grs.

Dandelion, 20 grs.

Syrup yerba santa, arom., 30 min.

Wild Cherry, 120 grs.**Wild Cherry and Iron.**

Wild Cherry bark, 60 grs.

Iron pyrophosphate, 8 grs

WILD CHERRY—Continued.**Extracts.**

Fluid Extract Black Cohosh Comp.—Not miscible with water. Each pint represents: Black Cohosh, 8 ozs.; Licorice, 3 ozs.; Wild Cherry and Senega, 2 ozs. each; Ipecac, 1 oz. Nervous and arterial sedative; in larger dose diaphoretic, and causing increased secretion from the bronchial mucous membrane. Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Horehound Comp.—Not miscible with water without some precipitation. Each pint represents: Horehound, 8 ozs.; Comfrey, 4 ozs.; Seneka and Wild Cherry, 2 ozs. each. Demulcent and expectorant. Its mild diaphoretic and sedative properties particularly fit it for administration to children. Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Wild Cherry, alcoholic.—Not miscible with water. This fluid extract, styled "alcoholic," differs from that of the Pharmacopoeia of 1880, in not containing glycerin. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Wild Cherry Comp.—Not miscible with water. Each pint represents: Wild Cherry, 8 ozs.; Wild Lettuce, 3 ozs.; Horehound, 3 1-2 ozs.; Blood Root, 1 oz.; American Hellebore, 1-2 oz. Indicated in convalescence from acute inflammatory and pulmonary diseases. Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Fluid Extract Wild Cherry, detannated.—Miscible with water. Chiefly employed in making elixirs of wild cherry bark, into which iron enters as an ingredient. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Wild Cherry for Syrup.—Miscible with water. Specially adapted for the preparation of a bright clear syrup of wild cherry of full medicinal activity. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Wild Cherry, U.S.P.—Not miscible with water. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Glycerole Yerbline Comp.—Each fluidounce represents:
 Yerba Santa, 60 grs. Wild Cherry bark, 30 grs.
 Potassium bromide, 10 grs. Tar, 5 grs.
 Licorice, 60 grs. Salicylic acid, 5 grs.
 Grindelia Robusta, 5 grs.

Lozenges.**Tar Compound.**

Pine Tar, 7-48 gr.
 Powd. Senega, 7-24 gr.

Powd. Wild Cherry, 2 1-3 grs.

Wild Cherry, 1 gr.

Prunin, a so-called **Concentration**,* a powdered preparation from Wild Cherry bark.
 Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

Syrups—Each fluidounce represents ingredients specified.

Horehound Compound.

Elecampane, 20 grs. Wild Cherry bark, 20 grs.
 Spikenard, 20 grs. Horehound, 20 grs.
 Comfrey, 20 grs. Blood Root, 10 grs.

This may be extemporized by employing the fluid extracts of the drugs named, reading minims for grains, and adding simple syrup to make one fluidounce.

Wild Cherry bark, U. S. P., 75 grs.

May be prepared extemporaneously after the following formula:
 Fluid Wild Cherry bark, P., D. & Co.'s. 6 fl. ozs.
 Syrup. 10 fl. ozs.
 Glycerin. 1 fl. oz.
 Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

Wines.—Each fluidounce represents ingredients specified.

Wild Cherry, 80 grs**Wild Cherry and Iron.**

Wild Cherry bark, 80 grs. Iron pyrophosphate, 8 grs.

* See note in Preface on Concentrations.

WILD GINGER. *Asarum canadense, Linné.* (Rhizome.)

Synonyms.—Canada Snakeroot; Assaret du Canada, Fr.; Canadische Haselwurz, Ger.

Properties.—Aromatic stimulant, carminative, with diaphoretic properties; used in low forms of fever as a substitute for serpentaria, which it resembles in effects. It has some reputation as an anthelmintic in ascariides, and as a diuretic it has met with some success in dropsy, with albuminous urine.

Wild Ginger (Canada Snakeroot), compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Wild Ginger (Canada Snakeroot).—Not miscible with water without some precipitation.
Dose, 15 to 30 minims (1 to 2 C.c.).

WILD INDIGO. *Baptisia tinctoria, R. Brown.* (Root.)

Synonyms.—Rattle-bush, Horse-fly Weed; *Sophora tinctoria, Linné.*

Properties.—Purgative, emetic, astringent, and antiseptic. Used principally on account of its antiseptic virtues. Used efficiently in the atonic varieties of rheumatism, pneumonia and dysentery, in scarlatina, typhus and all cases where there is a tendency to septicæmia. Externally as a wash, or in ointment form, for ill conditioned ulcers, scrofulous and syphilitic sores, etc., as an injection in fetid leucorrhœa, fetid discharges from the ears, etc.

Wild Indigo root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Baptisin, a so-called **Concentration,*** a powdered preparation from Wild Indigo root.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

Fluid Extract Wild Indigo.—Not miscible with water.

Dose, 5 to 15 minims (0.3 to 1 C.c.).

Ointment Wild Indigo.—One part Wild Indigo in 8.

WILD LETTUCE. *Lactuca canadensis, Linné.* (Leaves.)

Synonyms.—*Lactuca virosa, Linné;* Lettuce; *Laittue vireuse, Fr.;* Giftlattich, Ger.

Properties.—Mildly soporific, producing in susceptible patients a soothing hypnotic effect, which is often preferable to the powerful narcotic action of opiates. Prescribed oftenest for children.

Wild Lettuce leaves, compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Wild Lettuce.—Not miscible with water without some precipitation.

Dose, 20 minims (1.33 C.c.).

Powdered Extract Wild Lettuce.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Wild Lettuce.

Dose, 4 to 12 grains (0.25 to 0.75 Gm.).

Solid Extract Wild Lettuce.—One part equals 5 of leaves.

Dose, 4 to 12 grains (0.25 to 0.75 Gm.).

“German Tincture” Lettuce (*Lactuca virosa, Linné*).—An imported unofficial preparation of the green herb.

Dose, 2 to 10 minims (0.12 to 0.6 C.c.).

Pill Lettuce Extract, 2 grs.

Syrup Lettuce.

Fluid Extract Lettuce, P., D. & Co.’s.....2 fl. ozs.

Simple Syrup.....14 fl. ozs.

Dose, 1-2 to 2 fluidrachms (2 to 8 C. c.).

* See note in Preface on Concentrations.

WILD YAM. *Dioscorea villosa*, *Linné*. (Rhizome.)

Synonyms.—Colic root, Rheumatism root.

Properties.—Expectorant, diaphoretic, antispasmodic, and stimulating to the intestinal canal. A valuable remedy in bilious colic, giving prompt and permanent relief in the most severe cases. Useful in combination in flatulence, borborygmi, uterine diseases, cramp of the stomach, etc.

Wild Yam root, compressed in pound packages for retailing purposes.

PREPARATIONS.

Dioscorein, a so-called **Concentration**,* a powdered preparation from Wild Yam root.

Dose, 1-2 to 3 grains (0.03 to 0.2 Gm.).

Fluid Extract Wild Yam.—Not miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

WILLOW-HERB. *Epilobium angustifolium*, *Linné*. (Herb.)

Synonym.—Rose Bay.

Properties.—Tonic, astringent, demulcent and emollient. Said to be useful in chronic diarrhoea, dysentery, leucorrhoea, menorrhagia and uterine hemorrhage.

Willow-herb, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Willow-herb.—Miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

WINTERGREEN. *Gaultheria*, U. S. P. *Gaultheria procumbens*, *Linné*. (Herb.)

Synonyms.—Mountain Tea, Teaberry, Deerberry, Boxberry, Partridgeberry, Checkerberry.

Properties.—Stimulant, aromatic and astringent; also somewhat emmenagogue and galactagogue. The essential oil has been used preferably in cases of putrefactive fermentation. It is also of service in articular or acute rheumatism, on account of the large quantity of methyl salicylate it contains. The fluid extract is used in chronic diarrhoea, as an astringent; in dysuria, as a diuretic; and in infantile colic as a carminative.

Wintergreen leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS.

Capsules.—In soft gelatin.

Wintergreen Oil, 10 minims.

Wintergreen Oil, 5 minims.

Oil Wintergreen, 5 minims.

Olive Oil, 5 minims.

Fluid Extract Wintergreen.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C. c.).

WINTER'S BARK. *Drimys Winteri*, *Forster*. (Bark.)

Properties.—Stimulant, aromatic and tonic, resembling *Canella* and *Cinnamon* in its properties, for which drugs it may be substituted.

PREPARATION.

Fluid Extract Winter's bark.—Not miscible with water.

Dose, 1-4 to 1-2 fluidrachm (1 to 2 C. c.).

* See note in Preface on Concentrations.

WITCH-HAZEL. *Hamamelis*, U.S.P. *Hamamelis virginica*, *Linné*. (Bark; leaves, U.S.P.).

Synonyms.—Winter Bloom, Snapping Hazel, Spotted Alder.

Properties.—Tonic and astringent. A remedy of great value for hæmorrhoids, for which it may be used externally in the form of an ointment or decoction, combined with its internal use. It is also highly recommended in the treatment of varicocele, venous congestions and hæmorrhages of all kinds, as well as in diarrhoea, dysentery, excessive mucous discharges, menorrhagia, etc. In discussing threatening local inflammations it is much superior to arnica, especially when in the form of the distilled extract.

Witch-hazel bark, compressed in pound packages for retailing purposes.

Witch-hazel leaves, U. S. P., compressed in ounce packages for retailing purposes.

PREPARATIONS AND FORMULÆ.

Extracts.

Distilled Extract Witch-hazel (bark).—Miscible with water.

The best preparation for external use in sprains, bruises, prevention of ecchymosis, leucorrhœa, etc. It may be used internally in the same doses as the fluid extract.

Fluid Extract Witch-hazel, U. S. P. (leaves).—Not miscible with water.

Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

Powdered Extract Witch-hazel.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Sugar of Milk.

Dose, 5 to 15 grains (0.33 to 1 Gm.).

Solid Extract Witch-hazel.—One part equals 5 of the leaves.

Dose, 5 to 15 grains (0.3 to 1 Gm.).

Hamamelin, a so-called Concentration,* a powdered preparation from Witch-hazel leaves.

Dose, 1 to 3 grains (0.06 to 0.2 Gm.).

Ointment Witch-hazel.

Fluid Extract Witch-hazel, P., D. & Co.'s.....2 fl. ozs.
Lard.....14 ozs.

Melt the lard, add the fluid extract, and stir until cool.

Pill, Female, Leucorrhœa, 3 grs.

Hamamelin, 2 grs.

Hydrastin, 1-2 gr.

Senecin, 1-2 gr.

WOOD BETONY. *Betonica officinalis*, *Linné*. (Leaves.)

Properties.—Useful in atonic dyspepsia, chronic rheumatism, congestion of the portal circulation, and in catarrhal and malarial jaundice. It is also claimed that it has alterative properties which make it valuable in constitutional syphilis and scrofulosis.

Wood Betony leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Wood Betony.—Miscible with water.

Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

WORMWOOD. *Absinthium*, U. S. P. *Artemisia Absinthium*, *Linné*. (Leaves.)

Synonyms.—Absinthe commune, Armoise amère, Fr.; Wermuth, Ger.

Properties.—Tonic, anthelmintic, and narcotic. It is used in dyspepsia to promote the appetite and to restore tone to the digestive organs.

Wormwood leaves, U. S. P., compressed in ounce packages for retailing purposes.

* See note in Preface on Concentrations.

WORMWOOD—Continued.**PREPARATIONS.**

Fluid Extract Wormwood leaves.—Not miscible with water without some precipitation.

Dose, 5 minims to 1 fluidrachm (0.3 to 4 C.c.).

Solid Extract Wormwood.—One part equals 5 of leaves.

Dose, 1 to 10 grains (0.06 to 0.65 Gm.).

YARROW. *Achillea Millefolium*, *Linné.* (Herb.)

Synonyms.—Milfoil, Millefeuille, Fr.; Schafgarbe, Ger.

Properties.—Aromatic, tonic, mildly stimulant and astringent, exerting a special local action on the pelvic organs. In amenorrhœa, dysmenorrhœa, and all menstrual disorders, it is found useful. Also used in dysentery and in low forms of exanthematous fevers, intestinal colic, and infantile convulsions.

Yarrow leaves, compressed in ounce packages for retailing purposes.

PREPARATION.

Fluid Extract Yarrow.—Not miscible with water without some precipitation

Dose, 30 to 60 minims (2 to 4 C.c.).

YELLOW DOCK. *Rumex*, U.S.P. *Rumex crispus*, *Linné.* (Root.)

Synonyms.—Narrow Dock, Curled Dock; Patience frisée, Fr.; Grindwurz, Ger.

Properties.—Alterative, tonic, mildly astringent, but also somewhat laxative and not unlike rhubarb in its action. It is largely used in acne, favus, ichthyosis, the syphilitic skin diseases, scurvy, psora, and other cutaneous disorders. It is of service in all cases presenting the strumous diathesis.

Yellow Dock root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS AND FORMULA.**Extracts.**

Fluid Extract Yellow Dock, U.S.P. Not miscible with water.
Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Fluid Extract Yellow Dock Comp.—Not miscible with water. Each pint represents: Yellow Dock, 8 ozs.; False Bittersweet, 4 ozs.; American Ivy, 2 ozs.; Figwort, 2 ozs. Has long held a high place in domestic and scientific practice as an alterative compound in the form of the syrup, for the making of which it is specially adapted.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Solid Extract Yellow Dock.—One part equals 3 of root.
Dose, 5 to 20 grains (0.32 to 1.3 Gm.).

Rumicin, a so-called **Concentration**,* a powdered preparation from

Yellow Dock root.

Dose, 2 to 4 grains (0.13 to 0.26 Gm.).

Syrup Yellow Dock Compound.—Each fluidounce representing:

Yellow Dock, 60 grs.	Figwort, 15 grs.
Bitter Sweet, 30 grs.	Licorice, 30 grs.
American Ivy, 15 grs.	Cinnamon, 60 grs.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.)

An excellent excipient for exhibition of potassic iodide.

A medicinal equivalent may be prepared extemporaneously as follows:

Fluid Extract Yellow Dock Comp., P., D. & Co.'s.....	4 fl. ozs.
Hot Syrup.....	12 fl. ozs.

* See note in Preface on Concentrations.

YELLOW PARILLA. *Menispermum*, U. S. P. *Menispermum canadense*, *Linné*. (Rhizome.)

Synonyms.—Texas Sarsaparilla, Moonseed, Vine Maple, etc.

Properties.—Tonic, laxative, alterative and diuretic. In large doses it increases the actions of the pulse and bowels, and improves the appetite. In excessive doses produces vomiting and purging. Is said to be superior to the best imported sarsaparilla. Used as a remedy in scrofulous, cutaneous, arthritic, syphilitic and mercurial diseases. Employed in dyspepsia, general debility, and chronic inflammation of the viscera.

Yellow Parilla root, U. S. P., compressed in pound packages for retailing purposes.

PREPARATIONS.

Fluid Extract Yellow Parilla.—Not miscible with water.

Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

Menisperm, a so-called **Concentration**,* a powdered preparation from Yellow Parilla root.

Dose, 1 to 4 grains (0.06 to 0.26 Gm.).

YELLOW POND-LILY. *Nuphar advena*, *Aiton*. (Rhizome.)

Synonyms.—Spatterdock, Frog-lily.

Properties.—Astringent, anodyne, demulcent and antiscrofulous. Is used in dysentery, diarrhoea, gonorrhoea, leucorrhoea, scrofula and, combined with wild cherry, in bronchial affections. Externally, by means of cloths saturated with the diluted fluid extract, it is of value applied to boils, tumors, scrofulous ulcers and inflamed skin. In form of decoction it is used as a gargle in ulcers of the mouth and throat, and as an injection in leucorrhoea.

Yellow Pond-lily root, compressed in pound packages for retailing purposes.

PREPARATION.

Fluid Extract Yellow Pond-lily.—Not miscible with water.

Dose, 1-2 fluidrachm (2 C.c.).

YERBA BUENA. *Micromeria Douglassii*, *Bentham*. (Leaves.)

Properties.—A grateful aromatic stimulant, tonic, and anthelmintic. "Owing to its alterative, laxative, and tonic action, it has been employed in atonic dyspepsia."[†] Beneficial in gastrodynia, in allaying nausea, in expelling flatus, and in reducing the force and frequency of the pulse in fevers. It excites the menstrual flow, and hence is not without value as an emmenagogue. Yerba Buena is also an excellent anthelmintic, and is found to be well adapted to the expulsion of round worms in children.

PREPARATION.

Fluid Extract Yerba Buena.—Not miscible with water

Dose, 1-2 to 2 fluidrachms (2 to 8 C.c.).

YERBA MANSA. *Houttuynia californica*, *Bentham & Hooker*. (Root.)

Synonym.—Anemiopsis (*Anemopsis*) *californica*, *Hooker*.

Properties.—Stimulant, astringent and tonic. In malarial fever there is good evidence that it is valuable, particularly in the chronic varieties of the malarial infection. In diarrhoea and dysentery it is a remedy of pronounced value.

PREPARATION.

Fluid Extract Yerba Mansa.—Not miscible with water.

Dose, 15 to 60 minims (1 to 4 C.c.).

* See note in Preface on Concentrations.

[†] Dolan, "New Remedies," p 52.

YERBA REUMA. *Frankenia grandifolia*, Chamisso & Schlechtendahl. (Herb.)

Properties.—As its Spanish name (*flux herb*) implies, this plant is chiefly used in catarrhal affections. It contains a peculiar astringent principle, and a large percentage of saline ingredients, to the combination of which its therapeutical properties, as a topical application, are largely attributable. Used as an injection, it is curative in catarrhal affections of the nares and genito-urinary tract. In vaginal leucorrhœa this injection is peculiarly beneficial.

PREPARATION.

Fluid Extract Yerba Reuma.—Not miscible with water.

Dose, 10 to 20 minims (0.6 to 1.2 C. c.).

When used as an injection or a gargle, it must be diluted with three times its bulk of water.

YERBA SANTA. *Eriodictyon glutinosum*, Bentham. (Leaves.)

Synonyms.—Bear's Weed, Consumptive's Weed, Mountain-balm, Saint Herb.

Properties.—"A tonic expectorant, agreeable to the palate, and without nauseating properties."* It is especially indicated in chronic subacute inflammation of the bronchial mucous membrane, and in cases of paralysis of the bronchial muscles. Its effects in such cases are similar to the combined action of ipecac and balsam of Peru, its stimulating properties preponderating. One of its most remarkable properties is its power of completely destroying the bitterness of quinine, and it is often prescribed with reference to this property alone.

PREPARATIONS AND FORMULA.

Extracts.

Fluid Extract Yerba Santa.—Not miscible with water.

May be made to form a clear mixture with syrups and aqueous fluids by adding a little alkali.

Dose, 1-4 to 1 fluidrachm (1 to 4 C. c.).

Fluid Yerba Santa, aromatic.—Miscible with water.

The strength of the fluid is just four times that of the syrup of Yerba Santa, aromatic, and is specially adapted for the extemporaneous preparation of the latter, by simply diluting one part of the fluid with three parts of syrup. It enjoys a high reputation as a vehicle for quinine, completely overcoming the bitterness of this alkaloid.

Dose, 15 to 60 minims (1 to 4 C. c.).

NOTE.—This preparation is incompatible with acids and acid salts; exhibit quinine in suspension, not in solution.

Solid Extract Yerba Santa.—One part equals 4 of leaves.

Dose, 3 to 12 grains (0.2 to 0.8 Gm.).

Glyceroles.—Each fluidounce represents ingredients specified.

Yerba Santa, 60 grs.

Yerbine Compound.

Yerba santa, 90 grs.
Grindelia Robusta, 30 grs.
Wild Cherry, 30 grs.
Licorice, 90 grs.

Tar, 2 3-4 grs.
Salicylic acid, 2 grs.
Potassium carbonate, 2 1-2 grs.
Potassium bromide, 7 1-2 grs.

Inhalant Yerba Santa.—No. 3 of P. D. & Co.'s series of inhalants.

Fl. Ext. Yerba Santa, 3 j.
Glycerin, 3 j.

Tincture Opiumi, 3 j.
Alcohol, q.s. to make 3 liij.

Lozenge Yerba Santa Compound.—Each containing:

Ext. Licorice, 1 gr
Powd. Senega, 3-16 gr.
Tar, 1-32 gr.

Powd. Wild Cherry, 1-4 gr.
Yerba santa, 1 gr.

Malt Extract with Yerba Santa.—Sixty grains to the fluidounce.

Pill Yerba Santa Extract, 3 grs.

Syrup Yerba Santa, aromatic.—Each fluidounce representing 16 grains Yerba Santa in combination with aromatics. A teaspoonful completely masks the bitterness of 5 grains of quinine sulphate. It may be prepared extemporaneously from the above Fluid Yerba Santa, aromatic, q.v.

* Napheys, "Medical Therapeutics," p. 180.

See page 274

for

ADDENDA AND ERRATA.

APPENDIX.

NUMBERED FORMULA LIST.

OF SOME ELEGANT, CONVENIENT AND SCIENTIFIC

Pharmaceutical Preparations.

We solicit the patronage of physicians and pharmacists for these in consideration of:

1st. *Pharmaceutical accuracy*.—We guarantee that they contain all the medicinal agents, and in such proportions, as stated on our labels.

2d. *Palatability*.—They are as palatable as may possibly be exhibited with due regard for medicinal activity and pharmaceutical accuracy.

3d. *No information is afforded* on the labels or by means of circulars, that will induce or stimulate their use by the laity.

Our formulæ for masking the taste of the active principles in preparations such as elixirs, syrups, wines, etc., are the result of careful study and patient experiment, and so confident are we that they are unsurpassed for the object contemplated, that, when a preparation, claiming to contain an equal percentage of active medicinal principles, is more palatable, we at once suspect a discrepancy between the claims on the label and the actual contents of the bottle, and invite a comparative analysis. While we regard palatability as important, we hold medicinal virtue to be paramount, and deprecate the dishonesty which sacrifices the latter for the former.

CAPSULES.

There are many drugs from which to secure the best effects, it is necessary that they be administered in a more or less crude form. Something of therapeutic value is lost by attempting to render palatable, by incorporating with excipients or with aromatic diluents, such drugs as copaiba, male-fern, turpentine, cod-liver oil, etc., and in capsules alone is it possible for the physician to obtain their full and uncomplicated effects and at the same time refrain from offending the palate of his patient. The following are offered as desirable aids in the exhibition of this class of remedies.

Solubility.—This may readily be demonstrated by allowing one to lie loosely in the mouth; the contents will be found to escape in from 2 to 5 minutes, according to whether the capsule with which test is made be of soft or hard gelatin. There is not the remotest possibility of the capsule passing intact with the feces.

In Soft Gelatin, or "Soluble Elastic" Capsules.

List
No.

- I Apiol.
 Apiol, 5 minims.
 Olive Oil, 5 minims.
- 2 Balsam Fir, 10 minims.
- 3 Balsam Peru, 10 minims.
- 4 Castor Oil, 10 minims.
- 5 Castor Oil and Oil of Fleabane.
 Castor Oil, 5 minims.
 Oil Fleabane, 5 minims.
- 6 Castor Oil and Podophyllin.
 Castor Oil, 10 minims.
 Podophyllin, 1-8 grain.
- 7 Chaulmoogra Oil, 10 minims.
- 8 Chloroform, 10 minims.
- 9 Cod-liver Oil, Nor., 10 mins.
- 12 Cod-liver Oil and Iron.
 Cod-liver Oil, Nor., 10 minims.
 Iron oxide, 1-2 grain.
- 13 Cod-liver Oil and Iodine.
 Cod-liver Oil, Nor., 10 minims.
 Iodine, 1-4 grain.
- 14 Cod-liver Oil and Iodoform.
 Cod-liver Oil, Nor., 10 minims.
 Iodoform, 2 grains.
- 15 Cod-liver Oil and Phosphorus (1-30).
 Cod-liver Oil, Nor., 10 minims.
 Phosphorus, 1-30 grain.
- 16 Cod-liver Oil and Phosphorus (1-60).
 Cod-liver Oil, Nor., 10 minims.
 Phosphorus, 1-60 grain.
- 17 Copaiba, Para, 10 minims.
- 18 Copaiba and Iron.
 Copaiba, Para, 10 minims.
 Iron ammon. cit., 2 grains.
- 19 Copaiba and Oleoresin of Cubeb.
 Copaiba, Para, 7 minims.
 Oleoresin Cubeb, 3 minims.

List
No.

- 20 Copaiba, Cubeb and Buchu.
 Copaiba, Para, 6 minims.
 Oleoresin Cubeb, 2 minims.
 Ext. Buchu, 2 minims.
- 21 Copaiba, Cubeb and Iron.
 Copaiba, Para, 6 minims.
 Oleoresin Cubeb, 2 minims.
 Tr. chlor. iron (equiv.), 2 minims.
- 22 Copaiba, Cubeb and Rhatany.
 Copaiba, Para, 6 minims.
 Oleoresin Cubeb, 2 minims.
 Ext. Rhatany, 2 grains.
- 23 Copaiba, Cubeb and Sandal.
 Copaiba, Para, 6 minims.
 Oil Cubeb, 2 minims.
 Oil Sandal, East India, 2 minims.
- 24 Copaiba, Cubeb and Sarsaparilla.
 Copaiba, Para, 6 minims.
 Oleoresin Cubeb, 2 minims.
 Ext. Sarsaparilla, 2 grains.
- 25 Copaiba, Cubeb and Turpentine.
 Copaiba, Para, 4 minims.
 Oil Cubeb, 2 minims.
 Oil Turpentine, 4 minims.
- 26 Copaiba and Oil of Cubeb.
 Copaiba, Para, 7 minims.
 Oil Cubeb, 3 minims.
- 27 Copaiba and Rhatany.
 Copaiba, Para, 8 minims.
 Ext. Rhatany, 2 grains.
- 28 Copaiba and Sandal.
 Copaiba, Para, 5 minims.
 Oil Sandal, East India, 5 minims.
- 29 Petroleum Mass, 10 minims.
- 30 Cubeb Oil, 10 minims.
- 31 Cubeb Oleoresin, 10 minims.
- 32 Gurjun Balsam, 10 minims.
- 33 Haarlem Oil, 10 minims.
- 35 Linseed Oil, 10 minims.

CAPSULES—Continued.

<p>List
No.</p> <p>36 Matico Oleoresin, 10 minims.</p> <p>37 Copaiba and Matico.
Copaiba, Para, 7 minims.
Oleoresin Matico, 3 minims.</p> <p>38 Copaiba, Cubeb and Matico.
Copaiba, Para, 6 minims.
Oleoresin Cubeb, 3 minims.
Oleoresin Matico, 1 minim.</p> <p>39 Copaiba, Cubeb, Matico and Sandal.
Copaiba, Para, 3 minims.
Oleoresin Cubeb, 3 minims.
Oleoresin Matico, 1 minim.
Oil Sandal, East India, 3 minims.</p> <p>40 Cubeb and Matico.
Oleoresin Cubeb, 7 minims.
Oleoresin Matico, 3 minims.</p> <p>41 Copaiba Oil and Oil of Cubeb.
Oil Copaiba, 6 minims.
Oil Cubeb, 4 minims.</p> <p>42 Erigeron Oil, 10 minims.</p> <p>43 Eucalyptus Oil.
Oil Eucalyptus, true, 5 minims.
Oil Sweet Almond, 5 minims.</p> <p>44 Male-fern and Kamala.
Oil Male-fern, 7 minims.
Kamala, sifted, 4 grains.</p> <p>45 Origanum Oil.
Oil Origanum, 5 minims.
Oil Sweet Almond, 5 minims.</p> <p>46 Sandal Oil, East India, 10 mins</p> <p>47 Sandal and Cassia.
Oil Sandal, East India, 9 minims.
Oil Cassia, 1 minim.</p> <p>48 Cubeb and Sandal.
Oil Cubeb, 5 minims.
Oil Sandal, East India, 5 minims.</p> <p>49 Turpentine Oil, 10 minims.</p> <p>50 Wormseed Oil.
Oil Wormseed, 2 minims.
Olive Oil, 8 minims.</p> <p>51 } Phosphorated Oil, containing
52 } respectively 1-10, 1-30, 1-50,
53 } 1-60 and 1-100 of Phos-
54 } phorus in 10 minims of oil.
55 }</p> <p>56 Phosphorated Oil, compound.
1-60 gr. Phosphorus in 10 mins. oil.
Ext. Nux vomica, 1-4 grain.</p> <p>57 Tar, purified, 10 minims.</p> <p>59 Xylol.
Xylol, 5 minims.
Olive Oil, 5 minims.</p> <p>60 Castor Oil, 2 1-2 grammes.</p> <p>61 Castor Oil, 5 grammes.</p> | <p>List
No.</p> <p>62 Castor Oil, 10 grammes.</p> <p>63 Castor Oil, 15 grammes.</p> <p>64 Cod-liver Oil, 2 1-2 grammes.</p> <p>65 Cod-liver Oil, 5 grammes.</p> <p>66 Cod-liver Oil, 10 grammes.</p> <p>67 Cod-liver Oil, 15 grammes.</p> <p>68 Nitroglycerin.
Nitroglycerin, 1-100 grain.
Olive Oil, 8 minims.
Castor Oil, 2 minims.</p> <p>69 Copaiba Resin.
Copaiba, deprived of volatile oil,
5 minims.
Olive Oil, 5 minims.</p> <p>70 Male-fern and Castor Oil, 2 1-2 grammes.
Oleoresin Male-fern, 1 gramme.
Castor Oil, 1 1-2 grammes.</p> <p>71 Santonin and Castor Oil, 2 1-2 grammes.
Santonin, 1-4 grain.
Castor Oil, 2 1-2 grammes.</p> <p>72 Wintergreen Oil, 10 minims.</p> <p>73 Warburg's Tincture, 10 mins.
Each capsule of 10 minims represents 2 drachms of Warburg's tincture.</p> <p>74 Valerian Oil, 2 minims.
Oil Valerian, 2 minims.
Oil Sweet Almond, 8 minims.</p> <p>75 } Oil of Pennyroyal, containing
76 } respectively 1 and 2 mins.
with Olive oil, q.s. 10 mins.</p> <p>77 Wintergreen Oil, 5 minims.
Oil Wintergreen, 5 minims.
Olive Oil, 5 minims.</p> <p>78 Terebene, 10 minims.</p> <p>79 } Cascara Sagrada, containing
80 } respectively 2 & 3 grs. of the
solid ext. with Castor oil,
q. s. 10 minims.</p> <p>201 Quinine sulphate, 1 grain.</p> <p>202 Quinine sulphate, 2 grains.</p> <p>203 Quinine sulphate, 3 grains.</p> <p>204 Quinine sulphate, 4 grains.</p> <p>205 Quinine sulphate, 5 grains.</p> <p>206 Cinchonidine sulphate, 1 gr.</p> <p>207 Cinchonidine sulphate, 2 grs.</p> <p>208 Cinchonidine sulphate, 3 grs.</p> <p>209 Cinchonidine sulphate, 4 grs.</p> <p>210 Cinchonidine sulphate, 5 grs.</p> |
|---|--|

In Hard Gelatin, or "Hard Filled" Capsules.

List No.		List No.	
100	Apiol. Apiol, 4 minims. Olive Oil, 6 minims.	109	Haarlem Oil, 10 minims.
101	Castor Oil, 10 minims.	110	Eucalyptus Oil. Oil Eucalyptus, true, 5 minims. Olive Oil, 5 minims.
102	Castor Oil and Podophyllin. Castor Oil, 10 minims. Podophyllin, 1-8 grain.	111	Male-fern and Kamala. Oil Male-fern, 7 minims. Kamala, sifted, 4 grains.
103	Cod-liver Oil, Norwegian, 10 minims.	112	Sandal Oil, East India, 10 mins.
104	Cod-liver Oil and Iron. Cod-liver Oil, Nor., 10 minims. Iron iodide, 1-2 grain.	113	Sandal and Cassia. Oil Sandal, East India, 9 minims. Oil Cassia, 1 minim.
105	Copaiba, Para, 10 minims.	114	Copaiba and Sandal. Copaiba, Para, 7 minims. Oil Sandal, East India, 3 minims.
106	Copaiba and Iron. Copaiba, Para, 9 minims. Iron ammon. cit., 2 grains.	115	Copaiba, Cubeb and Sandal. Copaiba, Para, 6 minims. Oil Cubeb, 2 minims. Oil Sandal, East India, 2 minims.
107	Copaiba and Oleoresin Cubeb. Copaiba, Para, 8 minims. Oleoresin Cubeb, 2 minims.	116	Turpentine Oil, 10 minims.
108	Copaiba and Oil Cubeb (black). Copaiba, Para, 7 minims. Oil Cubeb, 3 minims.	117	Tar, purified, 10 minims.
		118	Xylol, 10 minims.

MEDICINAL ELIXIRS.

These are preparations which, in harmony with the advance of modern scientific pharmacy, aim at combining medicinal activity with palatability and pharmaceutical elegance.

Each fluidounce of any Elixir represents the quantity of ingredients specified in its formula.

Dose is uniformly 1 to 2 fluidrachms (4 to 8 C. c.).

1	Ammonium bromide, 40 grs.	10	Bismuth and Strychnine. Bismuth and ammon. cit., 8 grs. Strychnine, 8-100 grain.
2	Ammonium valerianate, 16 grs.	11	Blue Flag, 120 grains.
3	Ammonium and Morphine valerianates. Ammonium valerianate, 16 grs. Morphine valerianate, 1-2 grain.	12	Buchu, 120 grains.
4	Ammonium and Quinine valerianates. Ammonium valerianate, 16 grains. Quinine valerianate, 4 grains.	13	Buchu compound. Buchu, 60 grains. Cubeb, 15 grains. Juniper, 15 grains. Uva-ursi, 15 grains. Spirits nit. eth., 15 minims.
5	Ammonium and Strychnine valerianates. Ammonium valerianate, 16 grs. Strychnine valerianate, 8-60 gr.	14	Buchu, Juniper and Potassium acetate. Buchu, 45 grains. Juniper, 12 grains. Potassium acetate, 16 grains.
6	Arbor vitæ, 120 grains.	15	Buchu and Pareira Brava. Buchu, 30 grains. Pareira brava, 7 1-2 grains. Juniber berries, 15 grains. Stone root, 7 1-2 grains.
7	Berberine and Iron. Berberine phosphate, 2 grains. Iron pyrophosphate, 4 grains.	16	Buckthorn bark, 120 grains.
8	Bismuth. Bismuth and ammon. cit., 16 grs.	17	Calcium bromide, 40 grains.
9	Bismuth and Quinine. Bismuth and ammon cit., 16 grs. Quinine sulphate, 2 grains.	18	Calcium iodide, 16 grains.

ELIXIRS—Continued.

- 19 Calisaya alkaloids.
Representing Calisaya bark, 40 grains.
- 20 Calisaya bark, 40 grains.
- 21 Calisaya and Bismuth.
Calisaya bark (alkaloids), 40 grs.
Bismuth and ammon. cit., 8 grs.
- 22 Calisaya and Iron bromide.
Calisaya bark (alkaloids), 40 grs.
Iron bromide, 16 grains.
- 23 Calisaya and Iron protoxide.
Calisaya bark 40 grains and citrate of iron protoxide equivalent to 8 grains ferrous sulphate cryst.
- 24 Calisaya and Iron pyrophosphate.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
- 25 Calisaya, Iron and Bismuth.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
- 26 Calisaya, Iron, Bismuth and Pepsin.
Calisaya bark (alkaloids), 40 grs.
Sacch. pepsin, U. S. P., 8 grains.
Iron pyrophosphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
- 27 Calisaya, Iron, Bismuth, Pepsin and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
Sacch. pepsin, U. S. P., 8 grains.
Strychnine, 8-100 grain.
- 28 Calisaya, Iron, Bismuth and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine citrate, 8-100 grain.
- 29 Calisaya, Iron and Pepsin.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Sacch. pepsin, U. S. P., 8 grains.
- 30 Calisaya, Iron and Phosphorus.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Phosphorus, 4-100 grain.
- 31 Calisaya, Iron and Quinine.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 2 grains.
- 32 Calisaya, Iron and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 33 Calisaya, Iron, Strychnine and Pepsin.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
Sacch. pepsin, U. S. P., 8 grains.
Strychnine, 8-100 grain.
- 34 Calisaya and Lactated Pepsin.
Calisaya bark (alkaloids), 40 grs.
Lactated pepsin, 40 grains.
- 35 Calisaya, Pepsin and Bismuth.
Calisaya bark (alkaloids), 40 grs.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
- 36 Calisaya, Pepsin, Bismuth and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine, 8-100 grain.
- 37 Calisaya, Pepsin and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Sacch. pepsin, U. S. P., 8 grains.
Strychnine, 8-100 grain.
- 38 Calisaya, Quinine and Strychnine.
Calisaya bark (alkaloids), 40 grs.
Quinine sulphate, 2 grains.
Strychnine, 8-100 grain.
- 39 Calisaya and Strychnine.
Calisaya bark, 40 grains.
Strychnine, 8-100 grain.
- 40 Cascara Sagrada, 120 grains.
- 41 Cascara Sagrada Compound.
Cascara sagrada, 60 grains.
Licorice, 60 grains.
Berberis aquifolium, 30 grains.
Aromatics, q. s.
- 42 Cathartic Compound.
Senna, 16 grains.
Cascara sagrada, 32 grains.
Culver's root, 16 grains.
Butternut, 16 grains.
Rochelle salt, 32 grains.
- 43 Celery and Guarana.
Celery seed, 60 grains.
Guarana, 60 grains.
- 44 Chinoidin, 40 grains.
- 45 Chloral hydrate, 80 grains.
- 46 Chloral hydrate and Potassium bromide.
Chloral hydrate, 40 grains.
Potassium bromide, 40 grains.
- 47 Cimicifuga root, 120 grains.
- 48 Cimicifuga Compound.
Cimicifuga, 30 grains.
Licorice root, 10 grains.
Cherry, wild, 8 grains.
Seneka, 8 grains.
Ipecac, 4 grains.
- 49 Cinchona, detannated.
40 grains Red Cinchona bark, true, with choice aromatics; especially adapted for combination with the salts of iron.
- 50 Cinchona, ferrated.
Red Cinchona bark, true, 40 grs.
Iron citrate, 16 grains.

ELIXIRS—Continued.

- 51 Cinchonidine and Iron.
Cinchonidine sulphate, 4 grains.
Iron pyrophosphate, 8 grains.
- 52 Cinchonidine, Iron and Strychnine.
Cinchonidine sulph., 4 grains.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 53 Coca leaves, 120 grains.
- 54 Corydalis compound.
Turkey corn, 30 grains.
Stillingia, 30 grains.
Blue flag, 40 grains.
Prickly-ash, 15 grains.
Potassium iodide, 8 grains.
- 55 Coto bark, 30 grains.
- 56 Damiana, true, 120 grains.
- 57 Damiana, Iron and Phosphorus.
Damiana, true, 60 grains.
Iron pyrophosphate, 8 grains.
Phosphorus, 4-100 grain.
- 58 Damiana, Phosphorus, Nux Vomica and Iron.
Damiana, true, 60 grains.
Nux vomica, 8 grains.
Iron pyrophosphate, 8 grains.
Phosphorus, 4-100 grain.
- 59 Damiana, Phosphorus and Nux Vomica.
Damiana, 60 grains.
Nux vomica, 8 grains.
Phosphorus, 4-100 grain.
- 60 Dandelion 180 grains.
- 61 Dandelion Compound.
Dandelion, 10 grains.
Cherry, 10 grains.
Licorice, 20 grains.
Aromatics, q. s.
- 62 Eucalyptus Compound.
Eucalyptus globulus, 15 grains.
Wild cherry, 16 grains.
Gentian, 4 grains.
Licorice, 15 grains.
Dandelion, 20 grains.
Syrup yerba santa, aromatic, 30 minims.
- 63 Gentian, 20 grains.
- 64 Gentian and Iron chloride.
Gentian, 16 grains.
Tincture iron chloride (tasteless), 16 minims.
- 65 Gentian, Iron chloride and Lactated Pepsin.
Gentian, 16 grains.
Tincture iron chloride (tasteless), 16 minims.
Lactated pepsin, 20 grains.
- 66 Gentian and Iron pyrophosphate.
Gentian, 20 grains.
Iron pyrophosphate, 8 grains.
- 67 Gentian and Lactated Pepsin.
Gentian, 20 grains.
Lactated pepsin, 20 grains.
- 68 Grindelia Robusta Aromatic.
Grindelia robusta, 120 grains.
- 69 Guarana, 120 grains.
- 70 Helonias (False Unicorn), 120 grains.
- 71 Helonias compound.
Helonias, 15 grains.
Squaw vine, 60 grains.
Cramp bark, 15 grains.
Blue cohosh, 15 grains.
- 72 Hops, 60 grains.
- 73 Iron, Bismuth and Strychnine.
Iron pyrophosphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine, 8-100 grain.
- 74 Iron bromide, 16 grains.
- 75 Iron citrate and Arsenic.
Iron citrate, 8 grains.
Arsenious acid, 8-32 grain.
- 76 Iron phosphate, 8 grains.
- 77 Iron protoxide.
Equivalent to ferrous sulphate, 8 grains.
- 78 Iron protoxide and Bismuth.
Iron protoxide, equivalent to ferrous sulphate, 8 grains.
Bismuth and ammon. cit., 8 grs.
- 79 Iron protoxide, Calcium iodide and Cinchonine.
Calcium iodide, 2 grains.
Iron protoxide, equivalent to ferrous sulphate, 8 grains.
Cinchonine sulph . 4 grains.
- 80 Iron protoxide, Calcium iodide and Cinchonine with Arsenic.
Calcium iodide, 2 grains.
Iron protoxide, equivalent to ferrous sulphate, 8 grains.
Cinchona bark, 4 grains.
Fowler's solution, 40 minims.
- 81 Iron pyrophosphate, 16 grains.
- 82 Iron pyrophosphate and Quinine.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 2 grains.
- 83 Iron pyrophosphate and Strychnine.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 84 Iron, Quinine and Arsenic.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 2 grains.
Arsenious acid, 8-32 grain.
- 85 Iron and Quinine citrate, 16 grains.
- 86 Iron and Quinine phosphates.
Iron phosphate, 16 grains.
Quinine phosphate, 4 grains.
- 87 Iron, Quinine and Strychnine.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 2 grains.
Strychnine, 4-120 grain.

ELIXIRS—Continued.

- 88 Iron, Quinine, Strychnine and Pepsin.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 2 grains.
Strychnine, 4-100 grain.
Sacch. pepsin, U. S. P., 8 grains.
- 89 Iron, Quinine and Strychnine phosphates.
Iron phosphate, 16 grains.
Quinine phosphate, 4 grains.
Strychnine phosphate, 8-60 grain.
- 90 Iron, Quinine and Strychnine phosphates with Lactated Pepsin.
Iron phosphate, 16 grains.
Quinine phosphate, 4 grains.
Strychnine phosphate, 8-60 grain.
Lactated pepsin, 20 grains.
- 91 Iron and Strychnine citrate, 16 grains.
- 92 Iron valerianate, 8 grains.
- 93 Lactated Pepsin, 40 grains.
See p. 138 for formula.
- 94 Lactated Pepsin and Bismuth.
Lactated pepsin, 20 grains.
Bismuth and ammon. cit., 8 grs.
- 95 Lactated Pepsin, Bismuth and Strychnine.
Lactated pepsin, 20 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine, 8-100 grain.
- 96 Lactated Pepsin, Bismuth, Strychnine and Iron.
Lactated pepsin, 20 grains.
Bismuth and ammon. cit., 8 grs.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 97 Laxative.
Senna, Alexandria, 60 grains.
Butternut bark, 30 grains.
Rochelle salt, 60 grains.
- 98 Licorice, aromatic, for Quinine.
Licorice, 120 grains.
An elegant vehicle for the administration of quinine and other remedies possessing a disagreeable taste. Place the quinine in powder in a wine-glass, add a dessertspoonful of the elixir, stir briskly till thoroughly mixed, and then let the mixture be taken, followed immediately by about a teaspoonful of the clear elixir.
- 99 Lime Juice and Pepsin.
Lime juice, pure, 1 fluidrachm.
Sacch. pepsin, U. S. P. 16 grains.
Lime juice is a grateful refrigerant tonic, and a powerful anti-scorbutic. It aids digestion, and exerts a general benign influence over the system even in health, by promoting and regulating the secretions. This combination of pepsin with lime juice is an admirable remedy in conditions of general debility attended with loss of appetite and dyspeptic symptoms. It should be taken at meal-time, clear or in a little water. For all the minor derangements of digestion it is an admirable remedy, and especially commends itself as a substitute for the alcoholic bitters and liquors so frequently used to promote digestion.
Being strictly a physiological remedy it is adapted to a wide range of cases that do not require any medication, and intelligent physicians will find occasion frequently to prescribe it for this very reason. For patients of bilious habit it is a remedy that rarely comes amiss.
- 100 Lithium bromide, 40 grains.
- 101 Lithium citrate, 16 grains.
- 102 Lupulin, 16 grains.
- 103 Mandrake compound.
Mandrake root, 45 grains.
Culver's root, 30 grains.
Senna, 30 grains.
Jalap, 15 grains.
- 104 Matico, 120 grains.
- 105 Matico compound.
Matico, 40 grains.
Uva-ursi, 40 grains.
Hydrangea, 40 grains.
- 106 Morphine bromide, 1 grain.
- 107 Morphine valerianate, 1 grain.
- 108 Pepsin.
Sacch. pepsin, U. S. P., 10 grains.
- 109 Pepsin and Bismuth.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit. 8 grs.
- 110 Pepsin, Bismuth and Iron.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Iron pyrophosphate, 8 grains.
- 111 Pepsin, Bismuth, Iron and Quinine.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit. 8 grs.
Iron pyrophosphate, 8 grains.
Quinine sulphate, 8 grains.
- 112 Pepsin, Bismuth and Nux Vomica.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Nux vomica, 15 grains.
- 113 Pepsin, Bismuth and Pancreatin.
Sacch. pepsin, U. S. P., 8 grains.
Pancreatin, 8 grains.
Bismuth and ammon. cit., 8 grs.
- 114 Pepsin, Bismuth and Quinine.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Quinine sulphate, 2 grains.
- 115 Pepsin, Bismuth, Strychnine and Pancreatin.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine, 8-100 grain.
Pancreatin, 4 grains.
- 116 Pepsin, Bismuth and Wafer Ash.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Wafer-ash, 60 grains.

ELIXIRS—Continued.

- 117 Pepsin, Damiana, Phosphorus and Nux Vomica.
Sacch. pepsin, U. S. P., 8 grains.
Damiana, 60 grains.
Nux vomica, 8 grains.
Phosphorus, 4-100 grain.
- 118 Pepsin and Pancreatin.
Sacch. pepsin, U. S. P., 8 grains.
Sacch. pancreatin, 4 grains.
- 119 Pepsin and Quinine.
Sacch. pepsin, U. S. P., 10 grains.
Quinine sulphate, 2 grains.
- 120 Pepsin and Strychnine.
Sacch. pepsin, U. S. P., 10 grains.
Strychnine, 8-100 grain.
- 121 Pepsin, Strychnine and Bismuth.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Strychnine, 8-100 grain.
- 122 Pepsin, Strychnine, Bismuth and Iron.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 123 Pepsin, Strychnine and Iron.
Sacch. pepsin, U. S. P., 10 grains.
Iron pyrophosphate, 8 grains.
Strychnine, 8-100 grain.
- 124 Pepsin and Wafer-ash.
Sacch. pepsin, U. S. P., 10 grains.
Wafer-ash, 60 grains.
- 125 Phosphates Compound with Calisaya.
Calisaya bark (alkaloids), 40 grs.
Iron phosphate, 2 grains.
Calcium phosphate, with small excess of phosphoric acid, 2 grs.
- 126 Phosphorus, 8-100 grain.
- 127 Phosphorus and Strychnine.
Phosphorus, 8-100 grain.
Strychnine, 8-100 grain.
- 128 Potassium bromide, 40 grains.
- 129 Quinine and Arsenic.
Quinine sulphate, 8 grains.
Arsenious acid, 8-32 grain.
- 130 Quinine bromide, 4 grains.
- 131 Quinine sulphate, 8 grains.
- 132 Quinine valerianate, 4 grains.
- 133 Rhubarb and Columbo.
Rhubarb, 90 grains.
Columbo, 30 grains.
- 134 Rhubarb and Magnesias.
Rhubarb, 60 grains.
Magnesium acetate, 10 grains.
- 135 Rhubarb and Potassa.
Rhubarb, 60 grains.
Golden-seal, 30 grains.
Potassium bicarbonate, 60 grains.
- 136 Simple Elixir.
This article possesses a pleasant aromatic flavor and is suitable as a basis for the preparation of other medicinal elixirs. It is also useful as an adjuvant and corrective vehicle for the administration of nauseous medicines which may be either dissolved or suspended therein.
- 137 Sodium bromide, 40 grains.
- 137B Sodium Salicylate, 8 grains.
- 137C Sodium Salicylate and Iron.
Sodium salicylate, 8 grains.
Iron and ammon. cit., 8 grains.
- 138 Stillingia, 120 grains.
- 139 Stillingia compound.
Stillingia, 30 grains.
Turkey corn, 30 grains.
Pipsissewa, 15 grains.
Elder flowers, 15 grains.
Blue flag, 15 grains.
Prickly-ash berries, 7 1-2 grains.
Coriander seed, 7 1-2 grains.
- 140 Strychnine, 8-100 grain.
- 141 Strychnine valerianate, 8-50 grain.
- 142 Valerian, 120 grains.
- 143 Wafer Ash, 120 grains.
- 144 Wahoo (bark of root), 160 grs.
- 145 Wahoo and Blue Flag.
Wahoo-root bark, 80 grains.
Blue flag root, 40 grains.
- 146 Wild Cherry, 120 grains.
- 147 Wild Cherry and Iron.
Wild cherry bark, 60 grains.
Iron pyrophosphate, 8 grains.
- 148 Zinc valerianate, 4 grains.

ADDENDUM.

- 102B Manaca and Salicylates.
Manaca, 80 grains.
Sodium salicylate, 64 grains.
Potassium salicylate, 32 grains.
Lithium salicylate, 8 grains.

GRANULAR EFFERVESCENT PREPARATIONS.

These are of great value where cooling drinks are considered to be necessary or refreshing and where at the same time it is desired to incorporate medicines acceptably. Carbonic acid gas is of itself one of the most efficient remedies for flatulence, and from its slightly anæsthetic effect assists the stomach where there is a tendency to nausea or vomiting. There are also many salts which, to produce their best effects, require to be administered with a considerable quantity of water, and their disagreeable taste is much more thoroughly and palatably disguised by the effervescence of carbonic acid gas than by the addition of any flavoring agent merely.

Each drachm of any preparation contains the quantity of ingredient specified in its formula.

- 1 Ammonium bromide, 2 grains.
- 2 Bismuth and Ammon. cit., 2 grs.
- 3 Caffeine and Potassium bromide.
Caffeine, 1 grain.
Potassium bromide, 5 grains.
- 4 Caffeine citrate, 1-2 grain.
- 5 Carlsbad.
One drachm of the salt is equivalent to a pint of the natural water.
- 6 Cerium nitrate, 2 grains.
- 7 Cerium oxalate, 2 grains.
- 8 Congress.
Two drachms of the salt is equivalent to a pint of the natural water.
- 9 Crab Orchard.
One drachm of the salt is equivalent to a pint of the natural water.
- 10 Friederichshall.
Three drachms of the salt is equivalent to a pint of the natural water.
- 11 Iron bromide, 1 grain.
- 12 Iron carbonate, 2 grains.
- 13 Iron and Ammon. cit., 1 grain.
- 14 Iron citrate and Arsenic.
Iron and ammon. citrate, 2 grs.
Arsenious acid, 1-60 grain.
- 15 Iron and Quinine citrate, U. S. P., 3 grains.
- 16 Iron and Quinine citrate with Arsenic.
Iron and quinine citrate, U.S.P., 3 grains.
Arsenious acid, 1-60 grain.
- 17 Iron iodide, 1 grain.
- 18 Iron phosphate, 1 grain.
- 19 Iron phosphate, Quinine and Strychnine.
Iron phosphate, 2 grains.
Quinine, 1 grain.
Strychnine, 1-60 grain.
- 20 Kissingen.
Two drachms of the salt is equivalent to a pint of the natural water.
- 21 Lithium benzoate, 2 grains.
- 22 Lithium citrate, 4 grains.
- 23 Magnesium Aperient.
Magnesium sulphate, 5 grains.
Sodium tartrate and cit., 55 grs.
- 24 Magnesium citrate, 12 grains.
- 25 Pepsin, Saccharated, U. S. P., 5 grains.
- 26 Pepsin and Bismuth.
Sacch. pepsin, U. S. P., 5 grains.
Bismuth and ammon. cit., 1 gr.
- 27 Pepsin, Bismuth and Iron.
Sacch. pepsin, U. S. P., 5 grains.
Bismuth and ammon. cit., 2 grs.
Iron and ammon. citrate, 1 grain.
- 28 Pepsin, Bismuth and Strychnine.
Sacch. pepsin, U.S.P., 5 grains.
Bismuth and ammon. cit., 2 grs.
Strychnine, 1-60 grain.
- 29 Phosphates Compound (Chemical Food).
Iron phosphate, 1 grain.
Calcium phosphate, 2 grains.
Sodium phosphate, 1-2 grain.
Potassium phosphate, 1-2 grain.
- 30 Potassium benzoate, 2 grains.
- 31 Potassium bicarbonate, 10 grs.
- 32 Potassium bromide, 5 grains.
- 33 Potassium citrate, 5 grains.
- 34 Potassium iodide, 2 grains.
- 35 Potassium nitrate, 5 grains.
- 36 Pullna.
Three drachms of the salt is equivalent to one pint of the natural water.
- 37 Quinine citrate, 1 grain.
- 38 Rockbridge Alum.
One drachm of the salt is equivalent to a pint of the natural water.
- 39 Seidlitz mixture.
After effervescence has taken place, this preparation represents an equal quantity of Rochelle salt.
- 40 Seltzer.
One drachm of the salt is equivalent to a pint of the natural water.
- 41 Vichy.
One drachm of the salt is equivalent to a pint of the natural water.

GLYCEROLES.

This class of therapeutic agents is designed to permit of medication with the minimum use of alcohol, the presence of which is sometimes either contra-indicated or incompatible.

Each fluidounce of any Glycerole represents the quantity of ingredients specified in its formula.

1 Arbor vitæ, 120 grains.	12 Pepsin and Wafer Ash.
Sacch. pepsin, U. S. P., 10 grains.
Wafer ash, 30 grains.
2 Bismuth and Ammon. citrate,
16 grains.	13 Phosphorus, 2-25 grain.
3 Calendula, 120 grains.	14 Potassium chlorate, 8 grains.
4 Carbohc acid, 100 grains.	15 Sodium sulphite, 80 grains.
5 Damiana, Phosphorus and
Nux Vomica.
Damiana, 60 grains.
Phosphorus, 1-25 grain.
Nux vomica, 4 grains.	16 Squill Compound.
Squill, 40 grains.
Senega, 40 grains.
Antimony and potassium tartrate,
7-8 grain.
6 Evening Primrose, 120 grains.	17 Tannic Acid, 5 grains.
7 Grindelia Robusta, 120 grains.	18 Tar, 30 grains.
8 Hemlock bark, 120 grains.	19 Yerba Santa, 60 grains.
9 Hypophosphites.
Calcium hypophosphite, 6 grains.
Sodium hypophosphite, 5 grains.
Potassium hypophosphite, 3 grs.	20 Yerbine Compound.
Yerba santa, 90 grains.
Grindelia robusta, 30 grains.
Wild cherry, 30 grains.
Licorice, 90 grains.
Tar, 2 3-4 grains.
Salicylic acid, 2 grains.
Potass. carbonate, 2 1-2 grains.
Potass. bromide, 7 1-2 grains.
10 Lead subacetate, 160 grains.	
11 Pepsin, concentrated.
Sacch. pepsin, U. S. P., 480 grs.	

INHALANTS.

For use in atomizing inhalers for the treatment of nose, throat and lung disorders such as nasal catarrh, diphtheritis, tonsilitis, laryngitis, bronchitis, and even in the graver lung affections, pneumonia and consumption.

Since the germ theory of disease has become so well established, the application of inhalants must be greatly extended and the inhalation treatment be regarded as the most scientific method of medication for a very large class of diseases.

Inhalant No. 1.

A petroleum product especially adapted for inhalation and for use as a base, or diluent, for inhalants. It is free from all irritating properties, on account of which it is much to be preferred to similar preparations from which the sulphuric acid or other objectionable agents, used in process of purification, have been imperfectly removed.

Inhalant No. 2.

Balsam copaiba, 3 ss.
Ether, 3 j.
Inhalant No. 1, q. s. to make 3 iv.

Inhalant No. 3.

Fl. ext. yerba santa, 3 j.
Glycerin, 3 j.
Tinct. opium, 3 j.
Alcohol, q. s. to make 3 iij.

Inhalant No. 4.

Fl. ext. stramonium seed,
Fl. ext. hyoscyamus, 3 j.
Fl. ext. belladonna, 3 ss.
Glycerin, 3 j.
Alcohol, q. s. to make 3 iij.

Inhalant No. 5.

Tinct. benzoin comp.,
Glycerin,
Alcohol, of each 3 j.

Inhalant No. 6.

Oil of tar, 3 ss.
Inhalant No. 1, 3 j.

Inhalant No. 7.

Oil eucalyptus, 3 j.
Inhalant No. 1, 3 j.

INHALANTS—Continued.

Inhalant No. 8.

Tinct. iodine, 3 iss.
Glycerin, 3 j.
Alcohol, q. s. to make 3 iij.

Inhalant No. 9.

Fl. ext. blood-root, 3 j.
Fluid tolu, soluble, 3 ss.
Glycerin, 3 j.
Alcohol, q. s. to make 3 iij.

Inhalant No. 10.

Tinct. iodine, 3 ij.
Carbolic acid, 3 ij.
Fluid tolu, soluble, 3 j.
Glycerin, 3 j.
Alcohol, q. s. to make 3 iij.

Inhalant No. 11.

Fl. ext. cubeb, 3 j.
Fluid tolu, soluble, 3 j.
Tinct. iodine, 3 j.
Tinct. camphor, 3 j.
Carbolic acid, 3 j.
Glycerin, 3 j 1-3.
Alcohol, q. s. to make 3 iv.

Inhalant No. 12.

Ether,
Inhalant No. 1, of each 3 j.

Inhalant No. 13.

Tinct. iodine, 3 ss.
Glycerin, 3 ss.

Inhalant No. 14.

Carbolic acid, gr. x.
Inhalant No. 1, 3 j.

Inhalant No. 17.

Chloroform,
Inhalant No. 1, of each 3 j.

Inhalant No. 18.

Beechwood creasote, 3 j.
Inhalant No. 1, 3 ij.

Inhalant No. 19.

Iodoform, gr. xx.
Creasote, ℥x.
Oil eucalyptus, ℥xx.
Ether, 3 ij.
Oil sweet almond, q. s. to make 3 j.

Inhalant No. 20.

Terebene, 3 j.
Oil cubeb, 3 j.
Oil sandalwood, 3 j.
Inhalant No. 1, 3 v.

MALT PREPARATIONS.

A very extended use of the extract of malt by the profession of this country has demonstrated both the necessity of proper care in its preparation, and the great value of such properly prepared article as a digestive and reconstructive agent.

Our improved extract of malt is made after a process which a large experience has shown to be the most successful in deriving from the grain the largest proportion of its therapeutic and nutritive principles, and at the same time securing the permanency and keeping properties of the preparation. The best grade of Canada barley is employed in the manufacture.

Each fluidounce of any preparation represents the ingredients specified in its formula with malt extract as base or excipient.

Dose is uniformly 1 to 2 fluidrachms (4 to 8 C. c.).

1 Malt Extract Improved.

2 Malt Extract with Alteratives.

Iron iodide, 1 grain.
Potassium bromide, 5 grains.
Manganese iodide, 1-2 grain.
Manganese bromide, 1-2 grain.
Calcium iodide, 1 grain.

3 Malt Extract with Beef, Iron and Wine.

Iron and ammon. cit., 4 grains.
Virtues of one ounce of beef.
Malt ext. and wine, equal parts.

4 Malt Extract with Berberis Aquifolium.

Berberis aquifolium, 30 grains.

5 Malt Extract with Cascara Sagrada.

Cascara sagrada, 60 grains.

6 Malt Extract with Cascara Sagrada and Berberis Aquifolium.

Cascara sagrada, 30 grains.
Berberis aquifolium, 30 grains.

7 Malt Extract with Cod-liver Oil.

Cod-liver oil, 1-3 fluidounce.

7B Malt Extract with Cod-liver Oil and Hypophosphites.

Cod-liver oil, 1-3 fluidounce.
Calcium hypophosphite, 4 grains.
Sodium hypophosphite, 2 grains.

8 Malt Extract with Cod-liver Oil and Iron Iodide.

Cod-liver oil, 1-3 fluidounce.
Iron iodide, 4 grains.

9 Malt Extract with Cod-liver Oil and Phosphorus.

Cod-liver oil, 1-3 fluidounce.
Phosphorus, 4-100 grain.

10 Malt Extract with Dialyzed Iron.

Dialyzed iron, 1-2 drachm.

11 Malt Extract Ferrated.

Iron pyrophosphate, 8 grains.

12 Malt Extract Ferrated with Quinine.

Iron and quinine citrate, 8 grains.

MALT PREPARATIONS—Continued.

- 13 Malt Extract Ferrated with Strychnine.
Iron pyrophosphate, 8 grains.
Strychnine, 4-100 grain.
- 14 Malt Extract with Hops.
Hops, 30 grains.
- 15 Malt Extract with Hypophosphites.
Calcium hypophosphite, 4 grains.
Sodium hypophosphite, 4 grains.
Iron hypophosphite, 2 grains.
Potassium hypophosphite, 6 grs.
- 16 Malt Extract with Iodides Comp.
Iron iodide, 4 grains.
Manganese iodide, 2 grains.
- 17 Malt Extract with Iron Iodide.
Iron iodide, 2 grains.
- 18 Malt Extract with Iron protochloride.
Iron protochloride, 4 grains.
- 19 Malt Extract with Iron, Quinine and Strychnine.
Iron and quinine citrate, 8 grains.
Strychnine, 4-100 grain.
- 20 Malt Extract with Lactophosphates.
Calcium lactophosphate, 6 grains.
Iron lactophosphate, 4 grains.
Sodium lactophosphate, 2 grains.
Potassium lactophosphate, 2 grs.
Manganese lactophosphate, 2 grs.
- 21 Malt Extract with Lactophosphates and Pepsin.
Calcium lactophosphate, 6 grains.
Iron lactophosphate, 4 grains.
Sacch. pepsin, U. S. P., 8 grains.
Sodium lactophosphate, 2 grains.
Potassium lactophosphate, 2 grs.
Manganese lactophosphate, 2 grs.
- 22 Malt Extract with Pancreatin.
Pancreatin, saccharated, 8 grains.
- 23 Malt Extract with Pepsin.
Sacch. pepsin, U. S. P., 4 grains.
- 24 Malt Extract with Pepsin and Bismuth.
Sacch. pepsin, U. S. P., 4 grains.
Bismuth and ammon. cit., 4 grs.
- 25 Malt Extract with Pepsin and Pancreatin.
Sacch. pepsin, U. S. P., 4 grains.
Pancreatin saccharated, 4 grains.
- 26 Malt Extract with Pepsin and Strychnine.
Sacch. pepsin, U. S. P., 4 grains.
Strychnine, 4-100 grain.
- 27 Malt Extract with Petroleum.
Petroleum, crude, 2 drachms.
- 28 Malt Extract with Quinine.
Quinine sulphate, 2 grains.
- 29 Malt Extract with Strychnine.
Strychnine, 4-100 grain.
- 30 Malt Extract with Yerba Santa.
Yerba santa, 60 grains.

PILLS.

Sugar Coated and Gelatin Coated.

The following list comprises most of the official and popular formulæ known to the profession, and to which we have added many new and valuable combinations.

Our pills are made entirely by hand, from the purest materials and are coated by a method avoiding the application of any degree of heat which could impair their efficacy. They are unexcelled for superiority of finish, ready solubility and uniformity of coating.

We direct especial attention to the perfect distribution of the ingredients and the permanent softness of the mass. Particular care is taken to ensure the preservation of the drug.

They are put up in bottles of 100 and 500 each, except pills of the rarer alkaloids and glucosides, specially indicated, which only contain 25 each.

oA Acetanilid, 4 grains. 1 Aconite Extract (<i>of the leaves</i>) 1-4, 1-2 and 1 grain. 2 Aconitine, cryst., 1-500 and 1-200 grain. 3 *†Adonidin, 1-10 grain. 4 Ague.
Chinoidin, 2 grains.
Ext. Coloc. co., 1-3 grain.
Oleoresin Black Pepper, 1-6 gr.
Iron Sulph. excis., 1-2 grain. | <ol style="list-style-type: none"> 5 Ague, Improved.
Cinchonidine sulph., 1 grain.
Xanthoxyllin, 1 grain.
Gelsemperin, 1-16 grain.
Powd. Capsicum, 1-16 grain. 6 Aloes, U. S. P.
Purified Aloes, 2 grains.
Soap, 2 grains. 7 Aloes and Asafetida, U. S. P.
Purified Aloes, 1 1-3 grains.
Asafetida, 1 1-3 grains.
Soap, 1 1-3 grains. |
|---|--|

*Put up in bottles of 25 each.

†Gelatin-coated only.

PILLS—Continued.

- 8 Aloes and Iron.
Purified Aloes, 1 grain.
Iron sulph. exsic., 1 grain.
Jam. Ginger, 1-2 grain.
Ext. Conium seed, 1-2 grain.
- 9 Aloes and Iron, U. S. P.
Purified Aloes, 1 grain.
Iron sulph. exsic., 1 grain.
Aromatic Powder, 1 grain.
Confection Rose, q. s.
- 10 Aloes and Mastic, U. S. P.
Purified Aloes, 2 grains.
Mastic, 1-2 grain.
Powd. Red Rose, 1-2 grain.
- 11 Aloes and Myrrh, U. S. P.
Purified Aloes, 2 grains.
Myrrh, 1 grain.
Aromatic Powder, 1-2 grain.
- 11B Aloes, Myrrh and Iron.
Pulv. Aloes Soc., 2 grains.
Pulv. Myrrh, 1 grain.
Iron sulph. exsic., 1 grain.
- 12 Aloes and Nux vomica.
Purified Aloes, 1 1-2 grains.
Ext. Nux vomica, 1-2 grain.
- 13 Aloes, Nux vomica and Belladonna.
Purified Aloes, 1 1-2 grains.
Ext. Nux vomica, 1-2 grain.
Ext. Belladonna, 1-8 grain.
- 14 Aloin, 1-10, 1-5, 1-4, 1-2 and 1 gr.
- 15 Aloin Comp.
Aloin, 1-8 grain.
Ext. Belladonna, 1-4 grain.
Podophyllin, 1-8 grain.
- 16 Aloin and Strychnine.
Aloin, 1-5 grain.
Strychnine, 1-60 grain.
- 17 §Aloin Comp., and Strychnine.
Aloin, 1-8 grain.
Podophyllin, 1-8 grain.
Strychnine, 1-80 grain.
Ext. Belladonna, 1-8 grain.
Oleoresin Capsicum, 1-10 grain.
- 18 Aloin, Strychnine and Belladonna, No. 1.
Aloin, 1-5 grain.
Strychnine, 1-60 grain.
Ext. Belladonna, 1-8 grain.
- 18B Aloin, Strychnine and Belladonna, No. 2.
Aloin, 1-10 grain.
Strychnine, 1-50 grain.
Ext. Belladonna, 1-6 grain.
- 19 Aloin, Strychnine and Belladonna Comp.
Aloin, 1-5 grain.
Ext. Belladonna, 1-8 grain.
Strychnine, 1-60 grain.
Ext. Cascara sagrada, 1-2 grain.
Aloin, Strychnine, Belladonna and Ipecac (See Lapactic).
- 20 Alternative.
Blue Mass, 1 grain.
Powd. Ipecac, 1-8 grain.
Powd. Opium, 1-8 grain.
- 21 Alternative Compound.
Ext. Smilax Sarsap., 2-3 grain.
Ext. Stillingia sylvat., 2-3 grain.
Ext. Burdock, 2-3 grain.
Ext. Poke Root, 2-3 grain.
Ext. Prickly ash bark, 1-3 grain.
- 22 †Aluminium Compound.
Metallic Aluminium, 1 grain.
Hydrate Aluminium, 5-8 grain.
Calcium carb., 5-8 grain.
- 23 §Ammonium bromide, 1 grain.
- 24 §Ammonium chloride, 3 grains.
- 25 Ammonium picrate, 1-8, 1-4, 1-2, 1, 2 and 3 grains.
- 26 §Ammonium valerianate, 1 grain.
- 27 Anderson's Scot's.
Purified Aloes, 1 3-5 grains.
Powd. Colocynth, 4-15 grain.
Powd. Gamboge, 1-15 grain.
Powd. Soap, 1-15 grain.
Oil Anise, 1-30 grain.
- 28 Anodyne.
Camphor, 1 grain.
Ext. Henbane, 1 grain.
Morphine acetate, 1-20 grain.
Oil Capsicum, 1-20 grain.
- 29 Anthelmintic.
Santonin, 1 grain.
Calomel, 1 grain.
- 30 Anthemis Extract, 2 grains.
- 31 Anti-Bilious.
Ext. Coloc. co., 2 1-2 grains.
Podophyllin, 1-4 grain.
- 32 Anti-Chill.
Chinoidin, 1 grain.
Iron ferrocyanide, 2 grains.
Arsenious Acid, 1-20 grain.
Oleoresin Black Pepper, 1 grain.
- 33 Anti-Constipation, Brundage's.
Podophyllin, 1-10 grain.
Ext. Nux vomica, 1-4 grain.
Powd. Capsicum, 1-4 grain.
Ext. Belladonna, 1-10 grain.
Ext. Henbane, 1-4 grain.
- 34 †Anti-Constipation, Carson's.
Ext. Cascara sagrada, 1 grain.
Ext. Rhubarb, 1 grain.
Ext. Nux vomica, 1-4 grain.
Aloin, 1-3 grain.
- 35 Anti-Constipation, Palmer's.
Purified Aloes, 1 grain.
Ext. Henbane, 1 grain.
Ext. Nux vomica, 1-3 grain.
Powd. Ipecac, 1-10 grain.
- 36 Anti-Constipation, Goss'.
Podophyllin, 1-4 grain.
Ext. Coloc., 1-4 grain.
Ext. Butternut, 1-2 grain.
Ext. Nux vomica, 1-8 grain.
Ext. Henbane, 1-8 grain.
Ext. Gentian, 1-4 grain.
Ext. Cascara sagrada, 1-4 grain.
Powd. Canadian Hemp, 1-2 grain.

§Sugar-coated only.

†Gelatin-coated only.

PILLS—Continued.

- 37 Anti-Dyspeptic.
Strychnine, 1-40 grain.
Ext. Belladonna, 1-10 grain.
Powd. Ipecac, 1-10 grain.
Blue Mass, 2 grains.
Ext. Coloc. co., 2 grains.
- 38 Anti-Epileptic.
Iron ferrocyanide, 1-2 grain.
Quinine valerianate, 1 grain.
Zinc valerianate, 1-2 grain.
Ext. Valerian, 1 grain.
- 36 Antimony Co., U. S. P. (Plummer's Pill.)
Calomel, 1-2 grain.
Sulphurated Antimony, 1-2 grain.
Powd. Guaiac, 1 grain.
Mucilage Tragacanth, q.s.
- 40 Anti-Malarial, McCaw's.
Quinine sulph., 1 grain.
Iron sulph. exsic., 1-4 grain.
Arsenious acid, 1-80 grain.
Gelsemerin, 1-4 grain.
Podophyllin, 1-8 grain.
Oleoresin Blk Pepper, 1-16 grain.
- 41 Anti-Periodic.
Cinchonidine sulph., 1 grain.
Iron sulph. exsic., 1-2 grain.
Podophyllin, 1-20 grain.
Strychnine sulph, 1-33 grain.
Gelsemerin, 1-20 grain.
Oleoresin Capsicum, 1-10 grain.
- 42 §Anti-Syphilitic.
Potass. iodide, 2 1-2 grains.
Corrosive sublimate, 1-40 grain.
- 43 Aperient.
Ext. Nux vomica, 1-3 grain.
Ext. Coloc. co., 2 grains.
Ext. Henbane, 1-2 grain.
- 44 Aperient, Bauer's.
Ext. Henbane, 1-2 grain.
Ext. Aloes, 1 grain.
Ext. Coloc. co., 1 grain.
Potass. and Sod. tart., 1 1-2 grain.
- 45 Aperient, Drysdale's.
Powd. Rhubarb, 1 1-4 grains.
Powd. Ipecac, 5-12 grain.
Purified Aloes, 1 1-4 grains.
Powd. Nux vomica, 1-2 grain.
- 46 Aperient, Mild.
Ext. Coloc. co., 1-2 grain.
Ext. Henbane, 5-6 grain.
Powd. Rhubarb, 2 grains.
Oil Caraway, 1-20 grain.
- 47 Aphrodisiac Comp.
Ext. Coca, 1 grain.
Cinchonidine sulph., 1-2 grain.
Phosphorus, 1-200 grain.
Ext. Nux vomica, 1-8 grain.
Iron bromide, 1 grain.
- 48 Apocynum Extract, 2 grains.
- 49 Arsenious Acid, 1-100, 1-50, 1-50, 1-40, 1-32, 1-30, 1-20, and 1-12 gr.
- 50 Asafetida, 1, 2, 3, 4, and 15 grains.
- 51 Asafetida Comp.
Opium, 4-5 grain.
Ammonium carb., 4-5 grain.
Asafetida, 4-5 grain.
- 52 Asafetida and Iron.
Asafetida, 2 grains.
Iron sulph. exsic., 1 grain.
- 53 Asafetida and Nux vomica.
Asafetida, 3 grains.
Ext. Nux vomica, 1-4 grain.
- 54 Asafetida and Rhubarb.
Asafetida, 1 grain.
Powd. Rhubarb, 1 grain.
Iron by hydrogen, 1 grain.
- 55 Atropine, 1-120, 1-100 and 1-60 gr.
- 55 Ballou.
Ext. Coloc. co., 1 grain.
Ext. Jalap, 1 grain.
Calomel, 1 grain.
Powd. Ipecac, 1-8 grain.
- 57 Belladonna Extract, 1-8, 1-4, 1-2 and 1 grain.
- 58 Berberine hydrochlorate, 1 grain.
- 59 Berberine sulphate, 2 grains.
- 60 Berberine and Podophyllin.
Berberine sulphate, 1-4 grain.
Podophyllin, 1-20 grain.
- 61 Berberis Aquifolium Ext., 3 grs.
- 62 Berberis Compound.
Ext. Berberis aquifol., 2 grains.
Ext. Cascara sagrada, 1 grain.
- 63 Biliary, Junge's.
Manganese iodide, 1-2 grain.
Leptandrin, 3-10 grain.
Juglandin, 3-10 grain.
Sanguinarin, 1-5 grain.
Ext. Henbane, 3-5 grain.
- 64 Biliary, Wann's.
Ext. Coloc. co., 1 grain.
Podophyllin, 1-4 grain.
Ext. Jalap, 1-4 grain.
Ext. Henbane, 1-8 grain.
Powd. Capsicum, 1-4 grain.
- 65 Bismuth and Ignatia.
Bismuth subnitrate, 4 grains.
Ext. Ignatia bean, 1-4 grain.
- 66 Bismuth and Nux vomica.
Bismuth subnitrate, 5 grains.
Ext. Nux vomica, 1-2 grain.
- 67 Bismuth subcarbonate, 3 grains.
- 68 Bismuth subnitrate, 2, and 3 grs.
- 69 Black Haw Extract, 3 grains.
- 70 Black Hellebore Extract, 1 grain.
- 71 §Blennorrhagic.
Terebinth. alb., 1 1-2 grains.
Ext. Humuli, 3-4 grain.
Camphor monobrom., 3-4 grain.
Podophyllin, 1-8 grain.
- 72 Blue Mass, 1-2, 1, 2, 3 and 5 grs.
- 73 Blue Mass Comp.
Blue Mass, 1 grain.
Powd. Ipecac, 1-4 grain.
Powd. Opium, 1-2 grain.
- 74 Caffeine citrate, 1 grain.
- 75 Calabar bean Extract, 1-12 grain.

§Sugar-coated only.

†Gelatin-coated only.

§Sugar-coated only.

PILLS—Continued.

- 76 Calcium sulphide, 1-20, 1-10, 1-8, 1-5, 1-4, 1-2, 1 and 2 grains.
- 77 Calisaya Alkaloids.
Quinine sulph., 1-2 grain.
Cinchonine sulph., 1-2 grain.
Quinidine sulph., 1-2 grain.
Cinchonidine sulph., 1-2 grain.
- 78 Calomel, 1-10, 1-4, 1-2, 1, 2, 3, and 5 grains.
- 79 Calomel Comp.
Calomel, 3-4 grain.
Resin Guaiac, 1-2 grains.
Sulphurated Antimony, 3-4 grain.
- 80 Calomel and Extract Coloc. co.
Calomel, 1 grain.
Ext. Coloc. co., 2 1-2 grains.
- 81 Calomel and Opium.
Calomel, 2 grains.
Opium, 1 grain.
- 82 Calomel and Rhubarb.
Calomel, 1-3 grain.
Ext. Rhubarb, 1-2 grain.
Ext. Coloc. co., 1-2 grain.
Ext. Henbane, 1-6 grain.
- 83 Camphor Comp.
Camphor, 1 grain.
Powd. Opium, 1 grain.
Powd. Kino, 1 grain.
Ext. Capsicum, 1-16 grain.
- 84 Camphor and Henbane.
Camphor, 1 grain.
Ext. Henbane, 1 grain.
- 85 Camphor monob., 1, 2, and 3 grs.
- 86 Camphor and Opium.
Camphor, 2 grains.
Powd. Opium, 1 grain.
- 87 Camphor, Opium and Henbane.
Camphor, 1 grain.
Powd. Opium, 1-2 grain.
Ext. Henbane, 1 grain.
- 88 Camphor, Opium and Lead acet.
Camphor, 1 grain.
Opium, 1 grain.
Lead acetate, 1 grain.
- 89 Camphor, Opium and Tannin.
Camphor, 1 grain.
Tannin, 2 grains.
Opium, 1-4 grain.
- 90 Camphor, Valerian and Henbane.
Camphor, 1 grain.
Powd. Valerian, 1-2 grain.
Ext. Henbane, 1-2 grain.
- 91 Indian Cannabis Ext. (Cannabis Indica), 1-4, 1-2 and 1 grain.
- 92 Capsicum, 1 grain.
- 93 Capsicum Extract, 1-2 grain.
- 94 Carminative.
Strychnine, 1-20 grain.
Powd. Ipecac, 2-3 grain.
Powd. Black Pepper, 1-1-2 grains.
Ext. Gentian, 1 grain.
- 95 Cascara Comp. Pellets, D'Ary's.
Ext. Cascara sag., 4-15 grain.
Ext. Nux vomica, 1-30 grain.
Ext. Belladonna, 1-60 grain.
Euonymin, 1-5 grain.
Xanthoxyllin, 4-15 grain.
Oleoresin Capsicum, 1-20 grain.
- 96 Cascara sagrada Ext., 2 and 3 grs.
- 97 Cascara sagrada and Nux vomica.
Ext. Cascara sagrada, 2 grains.
Ext. Nux vomica, 1-5 grain.
- 98 Cascara sagrada, Nux vomica and Belladonna.
Ext. Cascara sagrada, 2 grains.
Ext. Nux vomica, 1-8 grain.
Ext. Belladonna, 1-16 grain.
- 99 Catarrh, Hager's.
Quinidine sulph., 3-8 grain.
Cinchonidine sulph., 3-8 grain.
Powd. Marshmallow, 1-4 grain.
Powd. Gentian, 1-4 grain.
Powd. red Sandalwood, 1-12 gr.
Powd. Tragacanth, 1-4 grain.
Glycerin, 1-4 grain.
Hydrochloric Acid, 1-4 grain.
- 100 Cathartic Comp. Cholagogue.
Podophyllin, 1-2 grain.
Ext. Henbane, 1-8 grain.
Oleoresin Capsicum, 1-8 grain.
Blue Mass, 1-4 grain.
Ext. Nux vomica, 1-16 grain.
- 101 Cathartic Comp. Granules.
Jalapin, 1-16 grain.
Leptandrin, 1-16 grain.
Aloin, 1-8 grain.
Podophyllin, 1-4 grain.
Powd. Gamboge, 1-32 grain.
Powd. Capsicum, 1-64 grain.
Ext. Henbane, 1-8 grain.
Oil Peppermint, 1-128 grain.
- 102 Cathartic Comp. Improved.
Ext. Coloc. co., 1 grain.
Ext. Jalap, 1-2 grain.
Podophyllin, 1-4 grain.
Ext. Gentian, 1-2 grain.
Leptandrin, 1-4 grain.
Ext. Henbane, 1-4 grain.
Oil Peppermint, q. s.
- 103 Cathartic Comp., Mild.
Ext. Coloc. co., 1 grain.
Jalap resin, 1-2 grain.
Calomel, 1-2 grain.
Gamboge, 1-4 grain.
Ext. Henbane, 1-2 grain.
Oil Peppermint, q. s.
- 104 Cathartic Comp. Modified.
Ext. Coloc. co., 1 grain.
Ext. Jalap, 3-4 grain.
Calomel, 3-4 grain.
Powd. Gamboge, 1-6 grain.
Powd. Rhubarb, 1-2 grain.
Powd. Jam. Ginger, 1-4 grain.
- 105 Cathartic Comp. U.S.P.
Ext. Coloc. co., 1 3-10 grains.
Ext. Jalap, 1 grain.
Calomel, 1 grain.
Powd. Gamboge, 1-4 grain.

PILLS—Continued.

- 106 Cathartic Comp. Vegetable.
Ext. Coloc., 1-3 grain.
Podophyllin, 1-4 grain.
Powd. resin Scammony, 1-3 grain.
Purified Aloes, 1 1-4 grains.
Powd. Cardamom, 1-8 grain.
Powd. Soap, 1-8 grain.
- 107 Caulophyllin, 1-8 grain.
- 108 Cerium oxalate, 1-4, 1 and 2 grs.
- 109 Charcoal, Willow, 3 grains.
- 110 Chimaphila Extract, 3 grains.
- 111 Chinoidin, 1-2, 1, 2 and 3 grains.
- 112 Chinoidin Comp.
Chinoidin, 2 grains.
Piperin, 1-2 grain.
Iron sulph. exsic., 1 grain.
- 113 Cholagogue.
Podophyllin, 1-4 grain.
Ext. Henbane, 1-2 grain.
Powd. Capsicum, 1-4 grain.
Purified Aloes, 1 grain.
Powd. Rhubarb, 1 grain.
- 114 Cimicifugin, 1 grain.
- 115 Cinchonidine salicylate, 2 1-2 grs.
- 116 Cinchonidine sulphate, 1-2, 1, 2, 3, 4 and 5 grains.
- 117 Cinchonine sulphate, 1 1-2 and 3 grains.
- 118 Cinchonidine Comp.
Cinchonidine sulph., 1 grain.
Arsenious acid, 1-32 grain.
Iron by hydrogen, 1 grain.
- 119 Cinchonidine Compound and Strychnine.
Cinchonidine sulph., 1 grain.
Arsenious acid, 1-20 grain.
Iron by hydrogen, 1 grain.
Strychnine, 1-20 grain.
- 120 Cinchonidine and Iron.
Cinchonidine sulph., 1 grain.
Iron by hydrogen, 1 grain.
- 121 Cinchonidine, Iron and Strychnine.
Cinchonidine sulph., 1 grain.
Iron proto-carb. (Vallet's), 2 grs.
Strychnine sulph., 1-60 grain.
- 122 Cincho-Quinine, 1, 2 and 3 grains.
- 123 Coca Ext., 3 grains.
- 124 †Cocaine muriate, 1-16 and 1-8 gr.
- 125 Cochia.
Ext. Coloc. co., 1 grain.
Purified Aloes, 1 grain.
Powd. Potass. sulph., 1-2 grain.
Powd. Gamboge, 1-2 grain.
Powd. Scammony, 1-2 grain.
- 126 Codeine, 1-16, 1-8, 1-5, 1-4, †1-2 and †1 grain.
- 127 *†Colchicine, 1-60 grain.
- 128 Colchicum Extract, 1-2 grain.
- 129 Colocynth comp. Ext., U. S. P., 3 grains.
- 130 †Colocynth compound Extract, Belladonna and Nux vomica.
Ext. Coloc. comp., 3 grains.
Ext. Belladonna, 1-4 grain.
Ext. Nux vomica, 1-2 grain.
- 131 Colocynth comp. Ext., and Blue.
Ext. Coloc. co., 2 1-2 grains.
Blue Mass, 1-2 grain.
- 132 Colocynth comp. Ext., and Blue.
Ext. Coloc. co., 2 1-2 grains.
Blue Mass, 2 1-2 grains.
- 133 Colocynth comp. Ext., Blue Mass and Ipecac.
Ext. Coloc. co., 2 grains.
Powd. Ipecac, 1-6 grain.
Blue Mass, 2 grains.
- 134 Colocynth compound Extract and Henbane
Ext. Coloc. co., 2 grains.
Ext. Henbane, 1 grain.
- 135 Colocynth compound Extract and Henbane.
Ext. Coloc. co., 3 1-3 grains.
Ext. Henbane, 1 2-3 grains.
- 136 †Colocynth comp. Henbane and Blue Mass.
Ext. Coloc. co., 3 grains.
Ext. Henbane, 1 grain.
Blue Mass, 1 grain.
- 137 Colocynth comp., Henbane and Podophyllin, Formula "A."
Ext. Coloc. Co., 3 grains.
Ext. Henbane, 1 grain.
Podophyllin, 1-4 grain.
- 138 Colocynth comp., Henbane and Podophyllin, Formula "B."
Ext. Coloc. co., 3 grains.
Ext. Henbane, 1 grain.
Podophyllin, 1-8 grain.
- 139 Colocynth comp. Ext., and Ipecac.
Ext. Coloc. co., 2 1-2 grains.
Powd. Ipecac, 1-2 grain.
- 140 Colocynth compound Extract, and Podophyllin.
Ext. Coloc. co., 2 1-2 grains.
Podophyllin, 1-4 grain.
Powd. Licorice, 1-4 grain.
- 141 *†Colocynthin, 1-8 grain.
- 142 Conium Extract (*of the seed*), 1-4, 1-2 and 1 grain.
- 143 Conium Ext., and Ipecac.
Ext. Conium seed, 1-2 grain.
Powd. Ipecac, 1-2 grain.
- 144 *†Convallamarin, 1-12 grain.
- 145 Cook's Pill.
Purified Aloes, 1 grain.
Powd. Rhubarb, 1 grain.
Powd. Soap, 1-2 grain.
Calomel, 1-2 grain.
- 146 Copaiba, 3, and 4 grains.

*Put up in bottles of 25 each.

†Gelatin-coated only.

*Put up in bottles of 25 each.

†Gelatin-coated only.

PILLS—Continued.

- 147 Copaiba Comp.
Pil. Copaiba, 1 1-2 grains.
Resin Guaiaac, 3-8 grain.
Iron citrate, 3-8 grain.
Ext. Cubeb, 5-8 grain.
- 148 Copaiba and Oleo-resin Cubeb.
Pil. Copaiba, 2 grains.
Oleo-resin Cubeb, 1 grain.
- 149 Copaiba, Cubeb and Iron cit.
Pil. Copaiba, 3-4 grain.
Ext. Cubeb, 1 1-2 grains.
Iron citrate, 3-4 grain.
- 150 Cornin, 2 grains.
- 151 Cornus florida Ext., 2 grains.
- 152 Corrosive Sublimate, 1-100, 1-60, 1-50, 1-40, 1-30, 1-20, 1-16, 1-12 and 1-8 grain.
- 153 Cubeb Extract, 2 grains.
- 154 Cubeb and Alum.
Oleo-resin Cubeb, 2 grains.
Powd. Alum, 1 grain.
- 155 Cubeb Ext., Rhatany and Iron.
Ext. Cubeb, 1 1-2 grains.
Iron sulph. exsic., 1 grain.
Ext. Rhatany, 1-2 grain.
- 156 Cypripedium Extract, 2 grains
- 157 Damiana Extract, 3 grains.
- 158 Dandelion and Leptandrin.
Ext. Dandelion, 1 1-3 grains.
Leptandrin, 2-3 grain.
- 159 *†Daturine sulphate, 1-100 grain.
- 160 Diaphoretic.
Morphine acetate, 1-25 grain.
Powd. Ipecac, 1-4 grain.
Powd. Potass. nitrate, 1 grain.
Powd. Camphor, 1-4 grain.
- 161 Diarrhoea Pellets.
Calomel, 1-8 grain.
Morphine sulph., 1-16 grain.
Powd. Capsicum, 1-16 grain.
Powd. Ipecac, 1-32 grain.
Powd. Camphor, 1-16 grain.
- 162 Digestive Pellets, Hager's.
Cinchonidine sulph., 1-5 grain.
Pepsin, 1 1-5 grains.
Powd. Ginger, 3-25 grain.
Powd. Cardamom, 3-25 grain.
Powd. Pimento, 3-25 grain.
Powd. Gentian root, 6-25 grain.
Powd. Marshmallow root, 6-25 gr.
Powd. Tragacanth, 6-25 grain.
Glycerin, 2-5 grain.
Acid hydrochloric, 2-5 grain.
- 163 Digitalin, pure, 1-60 grain.
- 164 Digitalis Ext., 1-2 grain.
- 165 Digitalis Compound.
Powd. Digitalis, 1 grain.
Powd. Squill, 1 grain.
Powd. Potass. nitrate, 2 grains.
- 166 Dinner, Chapman's.
Purified Aloes, 1 grain.
Gum Mastic, 1 grain.
Powd. Rhubarb, 1 grain.
- 167 †Dinner, Cole's.
Blue Mass, 1 1-5 grains.
Purified Aloes, 1 1-5 grains.
Powd. Jalap, 1 1-5 grains.
Ant. and Pot. tart., 1-50 grain.
- 168 Dinner, Lady Webster's.
Purified Aloes, 1 4-5 grains.
Gum Mastic, 3-5 grain.
Rose leaves, 2-5 grain.
- 169 Diuretic.
Powd. Soap, 1 1-2 grains.
Oil Juniper berries, 1-16 grain.
Sodium carb., 1 1-2 grains.
- 170 Dupuytren.
Powd. Guaiaac, 3 grains.
Powd. Opium, 1-8 grain.
Corrosive sublimate, 1-10 grain.
- 171 Dysentery.
Blue Mass, 1 grain.
Gelsemerin, 1-6 grain.
Powd. Ipecac, 1 grain.
- 172 Dysmenorrhoea, Alvord's.
Morphine sulph., 1-10 grain.
Cimicifugin, 2-3 grain.
Quinine sulph., 2-3 grain.
- 173 Dyspepsia.
Ext. Ignatia bean, 1-4 grain.
Ext. Cinchona bark, 1 grain.
Ext. Rhubarb, 1-4 grain.
Powd. Capsicum, 1-2 grain.
- 174 Dyspepsia, Moore's.
Cinchonidine sulph., 1-2 grain.
Ext. Capsicum, 1-3 grain.
Powd. Rhubarb, 1 grain.
- 175 Eccoprotic.
Ext. Soc. Aloes, 2 grains.
Ext. Nux vomica, 1-5 grain.
Podophyllin, 3-10 grain.
Oil Cloves, 1-10 gtt.
- 176 *†Elaterin, 1-20 gr.
- 177 Elaterium (Clutterbuck's), 1-10, 1-8 and 1-4 grain.
- 178 Emmenagogue, Improved.
Ergotin, 1 grain.
Purified Aloes, 1 grain.
Oil Savin, 1-2 grain.
Ext. Black Hellebore, 1 grain.
Iron sulph. exsic., 1 grain.
- 179 Emmenagogue, Mutter's.
Iron sulph. exsic., 1 1-2 grains.
Gum Turpentine, 1 1-2 grains.
Purified Aloes, 1-2 grain.
- 180 Emmenagogue No. 4.
Ergotin, 1 grain.
Ext. Cotton root, 1 grain.
Purified Aloes, 1 grain.
Iron sulph. exsic., 1 grain.
Oil Savin, 1-4 grain.
- 181 Emmenagogue, Rigaud's.
Purified Aloes, 1 1-2 grains.
Powd. Rue, 3-4 grain.
Powd. Saffron, 3-4 grain.
Powd. Savin, 3-4 grain.
- 182 Ergotin, 1-2, 1, 2 and 3 grains.

*Put up in bottles of 25 each.

†Gelatin-coated only.

*Put up in bottles of 25 each.

†Gelatin-coated only.

PILLS—Continued.

- 183 Ergotin and Indian Cannabis.
Ergotin purified, 1 grain.
Ext. Indian Cannabis, 1-2 grain.
- 183B Ergotin and Indian Cannabis.
Ergotin, purified, 1-2 grain.
Ext. Indian Cannabis, 1-4 grain.
- 184 Eucalyptus Comp.
Ext. Eucalyptus, 1 grain.
Ext. Canadian Hemp, 1-2 grain.
Sanguinariin, 1-8 grain.
- 185 Eucalyptus Extract, 2 grains.
- 186 Female, Amenorrhœa.
Ext. Black Cohosh, 1 grain.
Iron sulph. exsic., 1 grain.
Ext. Cotton root, 1 grain.
Purified Aloes, 1 grain.
- 187 Female, Hooper's.
Purified Aloes, 1 grain.
Iron sulph. exsic., 1-2 grain.
Ext. Black Hellebore, 1-4 grain.
Powd. Jam. Ginger, 1-8 grain.
Gum Myrrh, 1-4 grain.
Castile Soap, 1-8 grain.
Powd. Canella, 1-8 grain.
- 188 Female, Leucorrhœa.
Hamamelin, 2 grains.
Senecin, 1-2 grain.
Hydrastin, 1-2 grain.
- 189 Ferruginous, Blaud's 3 grains.
Iron sulph. exsic., 1 1-2 grains.
Potass. carb., 1 1-2 grains.
- 190 Ferruginous, Blaud's, 5 grains.
Iron sulph. exsic., 2 1-2 grains.
Potass. carb., 2 1-2 grains.
- 191 Fever and Ague.
Ext. Eucalyptus, 1-4 grain.
Chinoidin, 1 grain.
Iron ferrocyanide, 1-2 grain.
Powd. Capsicum, 1-4 grain.
Arsenious acid, 1-200 grain.
- 192 Fucus vesiculosus Ext., 3 grains.
- 193 Galbanum Comp., U. S. P.
Galbanum, 1 1-2 grains.
Asafetida, 1-2 grain.
Myrrh, 1 1-2 grains.
- 194 Gamboge Comp.
Powd. gum Gamboge, 3-5 grain.
Purified Aloes, 3-5 grain.
Powd. Cinnamon comp, 3-5 grain.
Soap, 1 1-5 grains.
- 195 Gelsemerin, 1-8 and 1-4 grain.
- 196 Gentian Comp.
Ext. Gentian, 2-3 grain.
Purified Aloes, 2-3 grain.
Powd. Rhubarb, 1 1-3 grains.
Oil Caraway, 1-10 grain.
- 197 Gentian Extract, 2 grains.
- 198 Geranin, 1 grain.
Glonoïn (see Nitroglycerin).
- 199 Gonorrhœa.
Powd. Cubeb, 1 1-4 grains.
Solid. Copaiba, 1 1-4 grains.
Iron sulph. exsic., 1-4 grain.
Venice Turpentine, 1-4 grain.
- 200 Grindelia robusta Extract, 3 grs.
- 201 Guarana Extract, 3 grains.
- 202 Headache.
Lactated pepsin, 1 grain.
Guarana, 1-2 grain.
Sodium bicarb., 1 grain.
Cypripedin, 1-2 grain.
- 202B Heim's (Quinine, Digitalis, Ipecac and Opium).
Quinine sulphate, 1 grain.
Powd. Digitalis, 1-2 grain.
Powd. Ipecac, 1-4 grain.
Opium, 1-4 grain.
- 203 Helonin, 1-8 grain.
- 204 Henbane Extract, 1-4, 1-2 and 1 gr.
- 205 Hepatica.
Blue Mass, 2 grains.
Ext. Henbane, 2-3 grain.
Ext. Coloc. co., 2-3 grain.
- 206 Hepatic, Eclectic.
Podophyllin, 1-4 grain.
Leptandrin, 1-2 grain.
Irisin, 1-4 grain.
Ext. Nux vomica, 1-16 grain.
Powd. Capsicum, 1-3 grain.
- 207 Hoang-Nan, 3 grains.
Hooper's (see Female).
- 208 Hydrastin (neutral), 1 grain.
(see also "*Berberine*,")
- 209 *§Hyoscine hydrobromate, 1-400 gr.
- 210 *†Hyoscyamine, pure, 1-200 grain.
- 211 Ignatia amara Extract, 1-4, 1-2 and 1 grain.
- 211B Intestinal Antiseptic.
Mercury protiodide, 1-8 grain.
Podophyllin, 1-16 grain.
Aloin, 1-16 grain.
Ext. Nux vomica, 1-16 grain.
Ext. Henbane, 1-16 grain.
- 212 Iodoform, 1 grain.
- 213 Iodoform and Iron.
Iron by hydrogen, 1 grain.
Iodoform, 1 grain.
- 214 Iodoform, Iron and Quinine.
Iodoform, 1 grain.
Iron proto-carb. (Vallet's), 2 grs.
Quinine sulphate, 1-2 grain.
- 215 Ipecac Ext., 1-4 gr.
- 216 Ipecac and Opium. } Equal to 2 1-2
Opium, 1-4 grain. } grs. Dover's
Ipecac, 1-4 grain. } Powder.
- 217 Ipecac and Opium. } Equal to 5
Opium, 1-2 grain. } grs. Dover's
Ipecac, 1-2 grain. } Powder.
- 218 Ipecac and Opium. } Equal to 10
Opium, 1 grain. } grs. Dover's
Ipecac, 1 grain. } Powder.
- 219 Ipecac and Squill.
Powd. Ipecac and Opium, 2 grs.
Powd. Squill, 2-3 grain.
Ammoniac, 2-3 grain.
- 220 Irisin, 1-2 and 1 grain.

*Put up in bottles of 25 each.

§Sugar-coated only.

†Gelatin-coated only.

PILLS—Continued.

- 221 Irisin Comp.
Irisin, 1-4 grain.
Strychnine, 1-40 grain.
Podophyllin, 1-10 grain.
- 222 Irisin Comp., and Henbane.
Irisin, 1-4 grain.
Strychnine, 1-100 grain.
Podophyllin, 1-10 grain.
Ext. Henbane, 1-2 grain.
- 223 Iron and Aloes.
Purified Aloes, 2 grains.
Iron sulph. exsic., 1 grain.
- 224 Iron bromide, 1 grain.
- 225 Iron citrate, 2 gr.
- 226 Iron citrate and Strychnine citrate.
Iron citrate, 2 grains.
Strychnine citrate, 1-50 grain.
- 227 Iron Compound, U. S. P.
Myrrh, 1 1-2 grains.
Sodium carb., 3-4 grain.
Iron sulph., 1-4 grain.
- 228 Iron ferrocyanide, 3 grains.
- 229 Iron by hydrogen (Q'venne's), 1 gr.
- 230 Iron by hydrogen (Q'venne's), 2 grs.
- 231 Iron iodide, U. S. P., 1 grain.
- 232 Iron iodide and Quinine.
Iron iodide, 1 1-2 grains.
Quinine sulph., 1-4 grain.
- 233 Iron lactate, 1 grain.
- 234 Iron and Manganese.
Iron carb., 2 grains.
Manganese carb., 1 grain.
- 235 Iron phosphate (ferrous), 2 grains.
- 236 Iron proto-carb., (Vallet's), 2, 3
5 grains.
- 237 Iron pyrophosphate, 1 grain.
- 238 Iron, Quassia and Nux vomica.
Iron by hydrogen, 1 1-2 grains.
Ext. Quassia, 1 grain.
Ext. Nux vomica, 1-4 grain.
Powd. Soap, 1-2 grain.
- 239 Iron and Quinine citrate, 1, 2, 3
and 15 grains.
- 240 Iron, Quinine and Strych. citrate.
Iron and Quinine citrate, 2 grs.
Strychnine citrate, 1-50 grain.
- 241 Iron and Strychnine.
Strychnine, 1-60 grain.
Iron by hydrogen, 2 grains.
- 242 Iron sulphate exsic., 1, 2 and 4 grs.
- 243 Iron valerianate, 1 grain.
- 244 Jaborandi Extract, 3 grains.
- 245 Jalap Compound.
Powd. Jalap, 3-4 grain.
Powd. Rhubarb, 3-4 grain.
Oil Caraway, 1-8 grain.
Castile Soap, 3-4 grain.
Pil. Calomel comp., 1 1-8 grains.
- 246 Jalap Extract, 1 grain.
- 247 Jalapin, 1 grain.
- 248 Jamaica Dogwood Extract, 2 grs.
- 249 Kermes mineral, 1-4 grain.
- 250 Krameria Extract, 2 grains.
- 251 Lactated Pepsin and Bismuth.
Lactated pepsin, 2 grains.
Bismuth subnitrate, 2 grains.
- 252 Lapactic.
Aloin, 1-4 grain.
Strychnine, 1-60 grain.
Ext. Belladonna, 1-8 grain.
Ipecac, 1-16 grain.
- 253 Laxative, Cole's.
Podophyllin, 1-10 grain.
Calomel, 1 grain.
Ext. Coloc. co., powd., 3 grains.
- 254 Laxative.
Purified Aloes, 1 grain.
Sulphur, 1-5 grain.
Podophyllin, 1-5 grain.
Resin Guaiac, 1-2 grain.
Syrup Buckthorn, q. s.
- 255 Laxative "Special."
Ext. Coloc. co. powd., 1 2-3 grains.
Ext. Henbane, 1 1-4 grains.
Soc. Aloes, powd., 5-6 grain.
Ext. Nux vomica, 5-12 grain.
Podophyllin, 1-12 grain.
Powd. Ipecac, 1-12 grain.
- 256 Leptandrin, 1-8, 1-4, 1-2, and 1 gr.
- 257 Leptandrin Comp.
Leptandrin, 1 grain.
Podophyllin, 1-8 grain.
Irisin, 1-4 grain.
- 258 Lettuce Extract, 2 grains.
- 259 Liver, Christopher's.
Calomel, 2 grains.
Rhubarb, 1 grain.
Powd. Ipecac, 1-2 grain.
- 260 Liver, Improved.
Purified Aloes, 1 grain.
Powd. Jalap, 1 grain.
Powd. Gamboge, 1-8 grain.
Leptandrin, 1-8 grain.
Calomel, 1-8 grain.
Oil Capsicum, 1-48 grain.
Tinct. Verat. Viride, 1-4 grain.
- 261 Liver, Improved, Vegetable.
Purified Aloes, 1 grain.
Powd. Jalap, 1 grain.
Powd. Gamboge, 1-8 grain.
Leptandrin, 1-8 grain.
Podophyllin, 1-8 grain.
Oil Capsicum, 1-48 grain.
Tinct. Verat. Viride, 1-4 grain.
- 262 Liver, Dr. Taylor's.
Podophyllin, 1 grain.
Powd. Ipecac, 1 grain.
Powd. Camphor, 4-5 grain.
- 263 Liver, Waxham's.
Ext. Culver's root, 1 grain.
Powd. Capsicum, 1-2 grain.
Podophyllin, 1-4 grain.
Ext. Jalap, 3-8 grain.
Powd. Gamboge, 1-8 grain.
- 264 Lupulin, 3 grains.
- 265 Manganese binoxide, 1 and 2 grs.

†Gelatin-coated only.

PILLS—Continued.

- 266 Magnesia calcined, 2 grains.
 267 Magnesia and Rhubarb, 2 grains.
 Magnesia calcined, 1 grain.
 Powd. Rhubarb, 1 grain.
 268 Mercury biniodide, 1-25, 1-16, 1-8,
 and 1-4 grain.
 269 Mercury protiodide, 1-40, 1-16, 1-10,
 1-8, 1-6, 1-5, 1-4, 1-3, and 1-2 gr.
 270 Mercury protiodide and Opium.
 Mercury protiodide, 1 grain.
 Powd. Opium, 1-3 grain.
 271 Morphine acetate, 1-8 and 1-4 gr.
 272 Morphine and Atropine.
 Morphine sulphate, 1-8 grain.
 Atropine sulphate, 1-100 grain.
 273 Morphine hydrochlorate, 1-8 and
 1-4 grain.
 274 Morphine sulphate, 1-32, 1-20, 1-16,
 1-10, 1-8, 1-6, 1-4, and 1-2 grain.
 275 Morphine sulphate and Belladonna.
 Morphine sulphate, 1-4 grain.
 Ext. Belladonna, 1-4 grain.
 276 Morphine sulphate comp.
 Morphine sulph., 1-4 grain.
 Antimony and Potass. tart., $\frac{1}{4}$ gr.
 Calomel, 1-4 grain.
 277 Morphine valerianate, 1-8 and
 1-4 grain.
 278 *†Muscarine sulphate, 1-100 grain.
 279 Neuralgic Idiopathic, Brown-
 Sequard's.
 Ext. Henbane, 2-3 grain.
 Ext. Ignatia, 1-2 grain.
 Ext. Aconite leaves, 1-3 grain.
 Ext. Stramonium seed, 1-5 grain.
 Ext. Conium seed, 2-3 grain.
 Ext. Opium, 1-2 grain.
 Ext. Indian Cannabis, 1-4 grain.
 Ext. Belladonna, 1-6 grain.
 280 Neuralgic Idiopathic, Brown-
 Sequard's, half-strength.
 281 Neuralgic, Dr. Gross'.
 Quinine sulph., 2 grains.
 Morphine sulph., 1-20 grain.
 Arsenious acid, 1-20 grain.
 Ext. Aconite leaves, 1-2 grain.
 Strychnine, 1-30 grain.
 282 Neuralgic without Morphine, Dr.
 Gross'.
 283 Nightsweat.
 Zinc oxide, 1-2 grain.
 Salicin, 1 grain.
 Ext. Belladonna, 1-25 grain.
 Hydrastin, 1 grain.
 Lactated pepsin, 1-2 grain.
 284 Nitroglycerin, 1-200, 1-100, 1-50,
 1-33, 1-25, and 1-10 grain.
 285 Nux vomica Extract, 1-8, 1-4, and
 1-2 grain.
 286 Opium 1-2 grain.
 Powd. Opium, 1-2 grain.
 Soap, 1-8 grain.
 287 Opium, U. S. P., 1 grain
 Powd. Opium, 1 grain.
 Soap, 1-4 grain.
 288 Opium Extract, 1-4, 1-2, and 1 gr.
 289 Opium and Lead acetate.
 Opium, 1 grain.
 Lead acetate, 1 grain.
 290 §Opium and Lead acetate, Formula
 "B."
 Opium, 1-2 grain.
 Lead acetate, 1 1-2 grains.
 291 Ox Gall.
 Ox Gall, 2 grains.
 Powd. Jamaica Ginger, 1 grain.
 292 Ox Gall, Craig's,
 Ox Gall, 2 grains.
 Ext. Coloc. co., 1 grain.
 293 Ox Gall Comp.
 Ox Gall, 2 grains.
 Ext. Stramonium seed, 1-6 grain.
 Purified Aloes, 1-10 grain.
 Berberine hydrochlorate, 1-12 gr.
 294 Ox-Gall and Pepsin, Armor's.
 Purified Aloes, 1 grain.
 Iron sulph. exsic., 1-2 grain.
 Ext. Nux vomica, 1-12 grain.
 Ox Gall, 1 1-2 grains.
 Pepsin, 1 grain.
 295 Pepsin and Bismuth.
 Pepsin, 1 1-2 grains.
 Bismuth subcarb., 3-4 grain.
 296 †Pepsin, Bismuth and Strychnine,
 No. 1.
 Pepsin, sacchr. 2 1-2 grains.
 Bismuth subnitrate, 2 1-2 grains.
 Strychnine, 1-60 grain.
 297 †Pepsin, Bismuth and Strychnine,
 No. 2.
 Pepsin pure, 1 grain.
 Bismuth subnitrate, 5 grains.
 Strychnine, 1-60 grain.
 298 Pepsin and Iron by Hydrogen.
 Pepsin, 1 1-2 grains.
 Iron by hydrogen, 3-4 grain.
 299 Pepsin and Iron Iodide.
 Pepsin, 1 1-2 grains.
 Iron by hydrogen, 1 1-2 grains.
 Iron iodide, 3-4 grain.
 300 Phosphorus, 1-100, 1-50, 1-33, 1-25,
 and 1-20 grain.
 301 Phosphorus and Aconite.
 Phosphorus, 1-50 grain.
 Ext. Aconite leaves, 1-16 grain.
 302 Phosphorus, Aloes, Iron and
 Strychnine.
 Phosphorus, 1-50 grain.
 Ext. Aloes, 1 grain.
 Iron sulph. exsic., 1 1-2 grains.
 Strychnine, 1-30 grain.
 303 Phosphorus, Aloes and Nux
 vomica.
 Phosphorus, 1-50 grain
 Ext. Aloes, 1-2 grain.
 Ext. Nux vomica, 1-4 grain.

*Put up in bottles of 25 each.

†Gelatın-coated only.

§Sugar-coated only.

†Gelatın-coated only.

PILLS—Continued.

- 304 Phosphorus, Aloes and Nux vomica, Formula "B."
Phosphorus, 1-20 grain.
Ext. Aloes, 1-2 grain.
Ext. Nux vomica, 1-2 grain.
- 305 Phosphorus and Belladonna.
Phosphorus, 1-100 grain.
Ext. Belladonna, 1-8 grain.
- 306 Phosphorus and Belladonna, Formula "B."
Phosphorus, 1-33 grain.
Ext. Belladonna, 1-4 grain.
- 307 Phosphorus and Indian Cannabis.
Phosphorus, 1-50 grain.
Ext. Indian Cannabis, 1-4 grain.
- 308 Phosphorus and Indian Cannabis, Formula "B."
Phosphorus, 1-25 grain.
Ext. Indian Cannabis, 1-4 grain.
- 309 Phosphorus and Cantharides co.
Phosphorus, 1-50 grain.
Cantharides, 1 grain.
Powd. Nux vomica, 1 grain.
- 310 Phosphorus and Cantharides co., Formula "B."
Phosphorus, 1-33 grain.
Cantharides, 1 grain.
Powd. Nux vomica, 1 grain.
- 311 Phosphorus Comp., No. 1.
Phosphorus, 1-100 grain.
Ext. Nux vomica, 1-4 grain.
- 312 Phosphorus Comp., No. 2.
Phosphorus, 1-60 grain.
Ext. Nux vomica, 1-4 grain.
- 313 Phosphorus Comp., No. 3.
Phosphorus, 1-50 grain.
Ext. Nux vomica, 1-8 grain.
- 314 Phosphorus and Digitalis comp.
Phosphorus, 1-50 grain.
Ext. Henbane, 1 grain.
Powd. Digitalis, 1 grain.
- 315 Phosphorus, Digitalis and Henbane.
Phosphorus, 1-33 grain.
Powd. Digitalis, 1 grain.
Ext. Henbane, 2 grains.
- 316 Phosphorus, Digitalis and Iron.
Phosphorus, 1-50 grain.
Powd. Digitalis, 1 grain.
Iron by hydrogen, 1 grain.
- 317 Phosphorus, Digitalis and Iron, Formula "B."
Phosphorus, 1-33 grain.
Powd. Digitalis, 1 grain.
Iron by hydrogen, 3 grains.
- 318 Phosphorus and Iron.
Phosphorus, 1-100 grain.
Iron proto-carb. (Vallet's), 1 gr.
- 319 Phosphorus and Iron, Formula "B."
Phosphorus, 1-50 grain.
Iron by hydrogen, 3 grains.
- 320 Phosphorus, Iron and Quinine.
Phosphorus, 1-100 grain.
Iron proto-carb. (Vallet's), 1 gr.
Quinine sulph., 1 grain.
- 321 Phosphorus, Iron and Quinine, Formula "B."
Phosphorus, 1-50 grain.
Iron by hydrogen, 3 grains.
Quinine sulph., 1-2 grain.
- 322 Phosphorus, Iron, Quinine and Strychnine.
Phosphorus, 1-50 grain.
Iron by hydrogen, 3 grains.
Quinine sulph., 1-2 grain.
Strychnine, 1-40 grain.
- 323 Phosphorus, Morphine and Zinc valerianate.
Phosphorus, 1-50 grain.
Morphine sulph., 1-12 grain.
Zinc valerianate, 1 grain.
- 324 Phosphorus, Morphine and Zinc valerianate, Formula "B."
Phosphorus, 1-25 grain.
Morphine hydrochlor., 1-12 grain.
Zinc valerianate, 1 grain.
- 325 Phosphorus and Nux vomica.
Phosphorus, 1-50 grain.
Ext. Nux vomica, 1-4 grain.
- 326 Phosphorus and Nux vomica, Formula "B."
Phosphorus, 1-33 grain.
Ext. Nux vomica, 1-2 grain.
- 327 Phosphorus, Nux vom. and Damiana.
Phosphorus, 1-100 grain.
Ext. Nux vomica, 1-8 grain.
Ext. Damiana, 2 grains.
- 328 Phosphorus, Nux vomica and Iron.
Phosphorus, 1-100 grain.
Ext. Nux vomica, 1-4 grain.
Iron phosphate precip., 1-2 grain.
- 329 Phosphorus, Nux vomica and Iron, Formula "B."
Phosphorus, 1-33 grain.
Ext. Nux vomica, 1-3 grain.
Iron by hydrogen, 3 grains.
- 330 Phosphorus, Nux vomica, Iron and Quinine.
Phosphorus, 1-100 grain.
Ext. Nux vomica, 1-8 grain.
Iron proto-carb. (Vallet's), 1 gr.
Quinine sulph., 1 grain.
- 331 Phosphorus, Nux vomica, Iron and Quinine, Formula "B."
Phosphorus, 1-25 grain.
Ext. Nux vomica, 1-3 grain.
Iron by hydrogen, 2 grains.
Quinine sulph., 1 grain.
- 332 Phosphorus, Opium and Digitalis.
Phosphorus, 1-50 grain.
Powd. Ipecac, 1-4 grain.
Powd. Opium, 1-4 grain.
Powd. Digitalis, 1-2 grain.
- 333 Phosphorus, Opium and Digitalis, Formula "B."
Phosphorus, 1-33 grain.
Powd. Ipecac, 1-4 grain.
Powd. Opium, 1-4 grain.
Powd. Digitalis, 1-2 grain.

PILLS—Continued.

- 334 Phosphorus and Quinine.
Phosphorus, 1-50 grain.
Quinine sulph., 1 grain.
- 335 Phosphorus, Quinine, Aloes and Nux vomica.
Phosphorus, 1-33 grain.
Quinine sulph., 1-2 grain.
Ext. Aloes, 1-2 grain.
Ext. Nux vomica, 1-3 grain.
- 336 Phosphorus and Quinine Comp.
Phosphorus, 1-50 grain.
Iron by hydrogen, 1 grain.
Quinine sulph., 1-2 grain.
Strychnine, 1-60 grain.
- 337 Phosphorus, Quinine and Digitalis Comp.
Phosphorus, 1-50 grain.
Quinine sulph., 1-2 grain.
Powd. Ipecac, 1-4 grain.
Powd. Digitalis, 1-2 grain.
Powd. Opium, 1-4 grain.
- 338 Phosphorus, Quinine and Nux vomica.
Phosphorus, 1-50 grain.
Quinine sulph., 1 grain.
Ext. Nux vomica, 1-4 grain.
- 339 Phosphorus and Strychnine.
Phosphorus, 1-50 grain.
Strychnine, 1-60 grain.
- 340 Phosphorus and Strychnine, Formula "B."
Phosphorus, 1-25 grain.
Strychnine, 1-30 grain.
- 341 Phosphorus, Strychnine and Iron.
Phosphorus, 1-100 grain.
Strychnine, 1-60 grain.
Iron proto-carb. (Vallet's), 1 gr.
- 342 Phosphorus, Strych. & Iron sulph.
Phosphorus, 1-100 grain.
Strychnine, 1-100 grain.
Iron sulph., 1 grain.
- 343 Phosphorus and Zinc comp.
Phosphorus, 1-50 grain.
Lupulin, 1 grain.
Zinc sulph., 1 grain.
- 344 Phosphorus, Zinc and Strychnine.
Phosphorus, 1-70 grain.
Zinc valerianate, 3-4 grain.
Strychnine, 1-30 grain.
- 345 Phosphorus, Zinc and Valerian.
Phosphorus, 1-40 grain.
Zinc sulph., 1 grain.
Ext. valerian, 2 grains.
- 346 *†Physostigmine salicylate, 1-100 grain.
- 347 Phytolaccin, 1-2 grain.
- 348 Picrotoxin, 1-60 grain.
- 349 *†Pilocarpine hydrochlorate, 1-30 and 1-16 grain.
- 350 Pilocarpine nitrate, 1-16 grain.
- 351 Podophyllin, 1-20, 1-10, 1-8, 1-4, 1-2, and 1 grain.

- 352 Podophyllin and Belladonna.
Podophyllin, 1-4 grain.
Powd. Capsicum, 1-2 grain.
Ext. Belladonna, 1-8 grain.
Powd. Sugar Milk, 1 grain.
- 353 Podophyllin, Bellad. and Calabar.
Podophyllin, 1-4 grain.
Ext. Belladonna, 1-4 grain.
Ext. Calabar bean, 1-4 grain.
- 354 Podophyllin and Blue Mass.
Podophyllin, 1-4 grain.
Blue Mass, 2 grains.
- 355 Podophyllin Comp.
Podophyllin, 1-2 grain.
Ext. Nux vomica, 1-16 grain.
Ext. Henbane, 1-8 grain.
- 356 Podophyllin Comp., Eclectic.
Podophyllin, 1-8 grain.
Juglandin, 1-16 grain.
Oleoresin Capsicum, 1-32 grain.
Leptandrin, 1-16 grain.
Macrotin, 1-32 grain.
- 357 Podophyllin and Leptandrin.
Podophyllin, 1-2 grain.
Leptandrin, 1 grain.
- 358 §Podophyllotoxin, 1-8 grain.
- 359 Podophyllum Extract, 1 grain.
- 360 Poppy Extract, 2 grains.
- 360B Post-partum, Fordyce Barker's.
Ext. Coloc. comp., 1 1-2 grains.
Calomel, 1 1-2 grains.
Ext. Nux vomica, 1-6 grain.
Powd. Aloes, 1-6 grain.
Powd. Ipecac, 1-6 grain.
Ext. Henbane, 1-3 grain.
- 361 §Potassium bromide, 1 and 5 grs.
- 362 §Potassium iodide, 2 and 5 grains.
- 363 Potassium and Iron tartrate, 2 grs.
- 364 §Potassium permanganate, 1-2, 1 and 2 grains.
- 365 Quassia Extract, 1 grain.
- 366 Quinidine sulphate, 1, 2, and 3 grs.
- 367 Quinine and Aloes.
Quinine sulph., 3-4 grain.
Purified Aloes, 1-4 grain.
- 368 Quinine bisulphate, 1-4, 1-2, 1, 2, 3, 4, and 5 grains.
- 369 Quinine bisulphate and Capsicum.
Quinine bisulph., 2 grains.
Powd. Capsicum, 1-2 grain.
- 370 Quinine and Blue Mass.
Quinine sulph., 1 grain.
Oleoresin Black Pepper, 1-4 gr.
Blue Mass, 2 grains.
- 371 Quinine bromide, 2 grains.
- 372 Quinine and Capsicum.
Quinine sulph., 2 grains.
Powd. Capsicum, 1-2 grain.

*Put up in bottles of 25 each.
Gelatin-coated only.

§Sugar-coated only.
†Gelatin-coated only.

PILLS—Continued.

- 373 Quinine Compound.
Quinine sulph., 1 grain.
Arsenious acid, 1-32 grain.
Iron by hydrogen, 1 grain.
- 374 Quinine Co. and Ext. Dandelion.
Quinine bisulph., 1 1-4 grains.
Iron sulph. excis., 2 grains.
Arsenious acid, 1-24 grain.
Ext. Dandelion, 1 1-4 grains.
- 375 Quinine Co. and Strychnine.
Quinine sulph., 1 grain.
Arsenious acid., 1-20 grain.
Iron by hydrogen, 1 grain.
Strychnine, 1-20 grain.
- 375B Quinine, Digitalis, Ipecac and Opium (Heim's).
Quinine sulphate, 1 grain.
Powd. Digitalis, 1-2 grain.
Powd. Ipecac, 1-4 grain.
Powd. Opium, 1-4 grain.
- 376 Quinine and Ext. Belladonna.
Quinine sulph., 1 grain.
Ext. Belladonna, 1-2 grain.
- 377 Quinine, Ipecac and Opium.
Quinine hydrobromate, 2 grains.
Pulv. Ipecac, 1-4 grain.
Pulv. Opium, 1-4 grain.
- 378 Quinine and Iron.
Quinine sulph., 1 grain.
Iron by hydrogen, 1 grain.
- 379 Quinine and Iron carbonate.
Quinine sulph., 1 grain.
Iron proto-carb. (Vallet's), 1 gr.
- 380 Quinine, Iron and Nux vomica.
Quinine sulph., 1 grain.
Iron proto-carb. (Vallet's), 2 grs.
Ext. Nux vomica, 1-4 grain.
- 381 Quinine, Iron and Strychnine phosphates.
Quinine phosphate, 1 grain.
Iron phosphate (precip.) 1 grain.
Strychnine phosphate, 1-60 grain.
- 382 Quinine, Iron and Strychnine.
Quinine sulph., 1 grain.
Iron proto-carb. (Vallet's), 2 grs.
Strychnine sulph., 1-60 grain.
- 383 Quinine and Iron valerianate.
Quinine sulph., 1 1-2 grains.
Iron valerianate, 1-2 grain.
- 384 Quinine, Iron and Zinc valerianates.
Quinine valerianate, 1 grain.
Iron valerianate, 1 grain.
Zinc valerianate, 1 grain.
- 385 Quinine and Strychnine.
Quinine sulph., 1 grain.
Strychnine, 1-60 grain.
- 386 Quinine sulphate, 1-4, 1-2, 1, 2, 3, 4, and 15 grains.
- 387 Quinine valerianate, 1-2 grain.
- 388 Rheumatic.
Ext. Coloc co., 1 1-2 grains.
Ext. Colch. acet., 1 grain.
Ext. Henbane, 1-3 grain.
Calomel, 1-3 grain.
- 389 Rhubarb Extract, 1 grain.
- 390 Rhubarb, U. S. P., 3 grains.
- 391 Rhubarb and Blue Mass.
Blue Mass, 1 3-4 grains.
Sodium bicarb., 1-2 grain.
Powd. Rhubarb, 1 grain.
- 392 Rhubarb Compound.
Powd. Rhubarb, 1 1-2 grains.
Powd. Myrrh, 3-4 grain.
Purified Aloes, 1 1-8 grains.
Oil Peppermint, q. s.
- 393 Rhubarb Compound, U. S. P.
Powd. Rhubarb, 2 grains.
Purified Aloes, 1 1-2 grains.
Powd. Myrrh, 1 grain.
Oil Peppermint, 1-10 grain.
- 394 Rhubarb and Iron.
Powd. Rhubarb, 2 grains.
Iron sulph. excis., 1 grain.
- 395 Salicin, 1, 2, 3, and 15 grains.
- 396 Salicylic Acid, 1, 2 1-2, 3, and 15 grains.
- 396B Salol, 2 1-2 and 5 grains.
- 397 Sandal Wood comp.
Oil Sandal Wood, 1 grain.
Ext. Cubeb, 1 grain.
Bals. Copaiba, 1 grain.
- 398 Sanguinaria Extract, 1-2 grain.
- 399 Sanguinariin (conc.), 1-2 and 1 gr.
- 400 Santonin, 1-2 and 1 grain.
- 401 Santonin and Calomel.
Santonin, 1-2 grain.
Calomel, 1-2 grain.
Cacao Butter, q. s.
- 402 Santonin and Podophyllin.
Santonin, 1-2 grain.
Podophyllin, 1-20 grain.
- 403 Sarsaparilla Extract, 3 grains.
- 404 Savin Extract, 1 grain.
- 405 *†Scillitoxin, 1-300 grain.
- 406 Sedative.
Ext. Musk root, 1-2 grain.
Ext. Henbane, 1-2 grain.
Ext. Valerian, 1-2 grain.
Ext. Indian Cannabis, 1-10 grain.
- 407 §Sedative, Mann's.
Ext. Coca, 1-2 grain.
Ext. Valerian, 1-2 grain.
Ext. Indian Cannabis, 1-4 grain.
Vallet's Mass, 1 grain.
Arsenious acid, 1-80 grain.
Strychnine, 1-80 grain.
Codeine, 1-8 grain.
- 408 Senna, Alex., Extract, 2 grains.
- 409 Silver iodide, 1-4 grain.
- 410 Soap and Opium.
Powd. Opium, 3-5 grain.
Soap, 2 2-5 grains.

*Put up in bottles of 25 each.

†Gelatin-coated only.

§Sugar-coated only.

†Gelatin-coated only.

PILLS—Continued.

- 411 Sodium bicarbonate, 4 grains.
 411B Sodium salicylate, 1 grain.
 412 Squill Comp.
 Powd. Squill, 3-8 grain.
 Powd. Jam. Ginger, 3-4 grain.
 Powd. Soap, 1 1-8 grains.
 Ammoniac, 3-4 grain.
 413 Stillingin, 1 grain.
 414 Stramonium Ext., (seed), 1-4, 1-2, and 1 grain.
 415 Strychnine, 1-100, 1-60, 1-50, 1-48, 1-40, 1-32, 1-30, 1-20, and 1-16 gr.
 416 Strychnine sulphate, 1-100 grain.
Strychnine Comp., Formula "A." see Quinine Comp. and Strychnine.
 417 Strychnine Comp., Formula "B."
 Strychnine, 1-100 grain.
 Ext. Indian Cannabis, 1-16 gr.
 Powd. Ginseng, 1 grain.
 Phosphorus, 1-100 grain.
 Iron proto-carb. (Vallet's), 1 gr.
 418 Sulphur iodide, 1-25 grain.
 419 Sundew Ext., 3 grains.
 420 Syphilitic, Ricord's, Modified.
 Mercury protiodide, 1-2 grain.
 Lactucarium, 1-2 grain.
 Ext. Opium, 1-10 grain.
 Ext. Conium seed, 1 1-2 grains.
 421 Tannin, 1, 3, and 15 grains.
 422 Taraxacum Extract, 3 grains.
 423 Tartar emetic, 1-16, 1-10, 1-8, and 1-4 grain.
 424 Tonic, Aiken's.
 Quinine sulph., 1 grain.
 Arsenious acid, 1-50 grain.
 Iron by hydrogen, 2-3 grain.
 Strychnine, 1-50 grain.
 425 Tonic, Hæmatic, Andrews'.
 Quinine sulph., 1 grain.
 Iron by hydrogen, 1 1-2 grains.
 Arsenious acid, 1-40 grain.
 Powd. Ipecac, 1-8 grain.
 Strychnine sulph., 1-40 grain.
 426 Tonic Hæmatic with Cinchonidine.
 Cinchonidine sulph., 1 grain.
 Powd. Ipecac, 1-8 grain.
 Arsenious acid, 1-40 grain.
 Iron by hydrogen, 1 1-2 grains.
 Strychnine sulph., 1-40 grain.
 427 Tonic, Waxham's.
 Cinchonine sulph., 1 grain.
 Ext. Prickly-ash bark, 1-4 grain.
 Ext. Cinchona bark, red, 1-4 gr.
 Ext. Dogwood bark, 1-4 grain.
 Powd. Capsicum, 1-4 grain.
 428 Trilliin, 1-8 grain.
 429 Triplex.
 Purified Aloes, 2 grains.
 Blue Mass, 1 grain.
 Podophyllin, 1-4 grain.
- 430 Triplex, Dr. Francis'.
 Purified Aloes, 1 1-5 grains.
 Blue Mass, 1 1-5 grains.
 Oil Caraway, 1-5 grain.
 Res. Scammony, 1 1-5 grains.
 Croton Oil, 1-20 grain.
 Tr. Aloes and Myrrh, 1-4 grain.
 431 Uva-ursi Extract, 2 grains.
 432 Valerian Extract, 2 grains.
 433 Veratrine, 1-60, 1-32, and 1-12 gr.
 434 Veratrum viride Ext., 1-4, and 1-2 grain.
 435 †Warburg's Tincture.
 Each pill equals one fluidrachm of the tincture.
 436 Yerba santa Extract, 3 grains.
 437 Zinc oxide, 1 grain.
 438 Zinc phosphide, 1-10, 1-8, 1-6, 1-4, and 1-2 grain.
 439 Zinc phosphide Comp.
 Zinc phosphide, 1-8 grain.
 Ext. Nux vomica, 1-8 grain.
 Ext. Indian Cannabis, 1-8 grain.
 440 Zinc phosphide and Nux vomica.
 Zinc phosphide, 1-10 grain.
 Ext. Nux vomica, 1-4 grain.
 441 Zinc valerianate, 1-2, 1, and 2 grs.

Pink Granules.

(Sugar-coated.)

These granules are intended to supply the needs of physicians in such cases as require minute or frequently repeated doses. *A glance at the list will show that they are distinguished from our regular line of pills in that they represent doses suitable for children.* This diminishes the risk there might otherwise be of giving to a child an overdose of some powerful remedy. At the same time their distinctive color renders them attractive to the class of patients for which they are intended.

- 442 Aconite root, 1-20 grain.
 443 Aloin, 1-10 grain.
 444 Alum, 1-10 grain.
 445 Ammonium chloride, 1-10 grain.
 446 Anodyne.
 Camphor, 1-8 grain.
 Ext. Henbane, 1-8 grain.
 Morphine acetate, 1-160 grain.
 Oil Capsicum, 1-160 grain.
 447 Anti-Chill.
 Chinoidin, 1-16 grain.
 Iron ferrocyanide, 1-8 grain.
 Arsenious acid, 1-320 grain.
 Oleo. Blk. Pepper, 1-16 grain.
 448 Anti-Constipation.
 Podophyllin, 1-40 grain.
 Ext. Nux vomica, 1-16 grain.
 Ext. Belladonna, 1-40 grain.
 Ext. Henbane, 1-16 grain.
 Powd. Capsicum, 1-16 grain.

†Gelatin-coated only.

†Gelatin-coated only.

PILLS—Continued.

- 449 Apomorphine, 1-60 grain.
 450 Arnica flowers, 1-5 grain.
 451 Arsenic iodide, 1-100 grain.
 452 Arsenious acid, 1-100 grain.
 453 Belladonna leaves, 1-20 grain.
 454 Calomel, 1-20 and 1-10 grain.
 455 Camphor, 1-20 grain.
 456 Camphor and Opium.
 Camphor, 1-6 grain.
 Powd. Opium, 1-12 grain.
 457 Camph. Opium and Tannin.
 Camphor, 1-16 grain.
 Tannin, 1-8 grain.
 Opium, 1-64 grain.
 458 Cantharides, 1-50 grain.
 459 Capsicum, 1-20 grain.
 460 Cathartic Comp., Improved, 1-4 gr.
 Ext. Coloc. co., 1-12 grain.
 Ext. Jalap, 1-24 grain.
 Podophyllin, 1-48 grain.
 Ext. Gentian, 1-24 grain.
 Leptandrin, 1-48 grain.
 Ext. Henbane, 1-48 grain.
 Oil Peppermint, q. s.
 461 Cathartic Comp., U. S. P., 1-4 gr.
 Ext. Coloc. co., 12-140 grain.
 Ext. Jalap, 1-14 grain.
 Calomel, 1-14 grain.
 Gamboge, 1-36 grain.
 462 Corrosive Sublimate, 1-100 grain.
 463 Croton Oil, 1-50 grain.
 464 Digitalis leaves, 1-20 grain.
 465 Dover's Powder, 1-4 grain.
 466 Ergotin, 1-10 grain.
 467 Gamboge, 1-32 grain.
 468 Gelsemium root, 1-50 gr.
 469 Henbane Extract, 1-8 grain.
 470 Hydrastin, 1-20 grain.
 471 Iodoform, 1-10 grain.
 472 Ipecac, 1-50 grain.
 473 Iron by hydrogen, 1-10 grain.
 474 Jalapin, 1-16 grain.
 475 Leptandrin, 1-16 grain.
 476 Mercury with chalk, 1-10 grain.
 477 Mercury protiodide, 1-40 and 1-20 grain.
 478 Morphine sulphate, 1-50 grain.
 479 Neuralgic, Dr. Gross', 1-4 grain.
 480 Nux vomica, 1-50 grain.
 481 Opium, 1-40 grain.
 482 Peppermint Oil, 1-128 grain.
 483 Phosphorus, 1-200 grain.
 484 Piperin, 1-20 grain.
 485 Podophyllin, 1-40 grain.
 486 Potass. arsenite, 1-100 grain.
 487 Potass. bromide, 1-5 grain.
 488 Potass. nitrate, 1-10 grain.
 489 Quinine sulphate, 1-10 grain.
 490 Quinine, Iron and Strychnine.
 Quinine sulph., 1-16 grain.
 Strychnine sulph., 1-960 grain.
 Iron proto-carb., 1-8 grain.
 491 Salicylic Acid, 1-10 grain.
 492 Santonin, 1-10 grain.
 493 Strychnine, 1-100 grain.
 494 Tannic Acid, 1-20 grain.
 495 Tartaric Acid, 1-10 grain.

Pills of the British Pharmacopœia.

SUGAR-COATED—OVIFORM.

To meet the demands of our trade in Great Britain and her colonies, we have prepared the following line of pills official in the British Pharmacopœia, 1835.

Aloes and Asafetida.

- Soc. Aloes, 1 grain.
 Asafetida, 1 grain.
 Soap, 1 grain.
 Confection Rose, 1 grain.

Aloes, Barbadoes.

- Barb. Aloes, 2 grains.
 Soap, 1 grain.
 Oil Caraway, 1-8 minim.
 Confection Rose, 1 grain.

Aloes and Iron.

- Iron sulph., 1-2 grain.
 Barb. Aloes, 2-3 grain.
 Cinnamon powd. co., 1 grain.
 Confection Rose, 1 1-3 grains.

Aloes and Myrrh.

- Soc. Aloes, 2 grains.
 Myrrh, 1 grain.
 Saffron, dried, 1-2 grain.

PILLS—Continued.

Aloes, Socotrine.

- Soc. Aloes, 2 grains.
- Soap, 1 grain.
- Oil Nutmeg, 1-8 minim.
- Confection Rose, 1 grain.

Asafetida Comp.

- Asafetida, 1 grain.
- Galbanum, 1 grain.
- Myrrh, 1 grain.

Colocynth Comp.

- Colocynth pulp, 1-2 gr.
- Barb. Aloes, 1 grain.
- Scammony, 1 grain.
- Potass. sulph., 1-8 grain.
- Oil Clove, 1-8 minim.

Colocynth and Henbane.

- Pil. Coloc. co., 2 grains.
- Ext. Henbane, 1 grain.

Gamboge Comp.

- Gamboge, 3-5 grain.
- Barb. Aloes, 3-5 grain.
- Cinnamon powd. co., 3-5 grain.
- Soap, 1 1-5 grains.

Hemlock Comp. (Conium).

- Ext. Conium leaves, 3-4 grain.
- Ipecac, 3-20 grain.

Ipecac and Squill.

- Ipecac powd. co., 2 grains.
- Squill, 2-3 grain.
- Ammoniac, 2-3 grain.

Iron carbonate.

- Iron carb. sacch., 2 1-2 grains.
- Confection Rose, 5-8 grain.

Iron Iodide, 1 grain.

Lead and Opium.

- Lead acetate, 2 grains.
- Opium powd., 1-3 grain.
- Confection Rose, 1-3 grain.

Mercurial pill, 3 and 5 grains.

Mercury subchloride Comp.

- Mercury subchloride, 3-4 grain.
- Antimony sulphurated, 3-4 grain.
- Guaiaic Resin, 1 1-2 grains.
- Castor Oil, 3-4 grain.

Phosphorus.

- Phosphorus, 1-30 grain.
- Bals. Tolu, 1 1-3 grains.
- Yellow Wax, 2-3 grains.
- Curd Soap, 1 grain.

Rhubarb Comp.

- Powd. Rhubarb, 1 grain.
- Soc. Aloes, 3-4 grain.
- Myrrh, 1-2 grain.
- Soap, 1-2 grain.
- Oil Peppermint, 1-16 grain.

Scammony Comp.

- Resin Scammony, 1 grain.
- Resin Jalap, 1 grain.
- Soap, 1 grain.
- Tinct. Ginger, 1 minim.

Soap Comp.

- Powd. Opium, 1-2 grain.
- Soap, 2 grains.

Squill Comp.

- Powd. Squill, 5-8 grain.
- Powd. Ginger, 1-2 grain.
- Powd. Ammoniac, 1-2 grain.
- Powd. Soap, 1-2 grain.

MEDICATED COLLODIONS.

These liquids evaporate rapidly on exposure to air, leaving a thin transparent film insoluble in water or alcohol. This is used as a protective covering for superficial burns, ulcers and wounds, cuts, cracked nipples, anal fissures, erysipelas, etc. Some medicinal agents are very readily incorporated with collodion, and physicians will appreciate the convenience of application suggested by the following combinations.

Simple (or contractile) collodion shrinks in drying and this property is taken advantage of where it is necessary to keep the edges of incised wounds in firm contact, or where compression is desired—otherwise flexible collodion is to be preferred.

Where medicinal agents are incorporated the quantity represented in each fluid-ounce is specified.

Aconite.

- Aconite root, 15 grains.

Anodyne (Morphine).

Arnica.

- Arnica root, 24 grains.

Belladonna.

- Belladonna leaves, 20 grains.

Cantharidal, U. S. P. (blistering).

- Cantharides, 240 grains.

Capsicum, 15 grains.

Carbolic.

- Carbolic Acid, 30 grains.

Caustic.

- Corrosive sublimate, 68 grains.

Contractile (simple).

Ferruginous.

- Solution Iron chloride, U. S. P., 60 minims.

Flexible, U. S. P.

Iodized.

- Iodine, 15 grains.

Morphine, 4 grains.

Simple, U. S. P.

Styptic, U. S. P.

- Tannic acid, 30 grains.

MEDICINAL SYRUPS.

Syrups prepared from crude drugs and known on our list as "medicated" will be found under the names of the articles from which they are prepared.

Each fluidounce contains the medicinal virtues of the quantity of ingredients specified in each formula.

- 1 Acid Phosphate with Pepsin.
Sacch. pepsin, U. S. P., 40 grs.
Liquid acid phosphate, 2 drachms.
 - 2 Calcium hypophosphite, 32 grains.
 - 3 Calcium iodide, 16 grains.
 - 4 Calcium lactophosphate, 16 grains.
 - 5 Calcium and Iron lactophosphates.
Calcium lactophosphate, 8 grains.
Iron lactophosphate, 4 grains.
 - 6 Calcium lactophosphate and Pepsin.
Calcium lactophosphate, 16 grs.
Sacch. pepsin, U. S. P., 16 grs.
 - 7 Calcium and Sodium hypophosphites.
Calcium hypophosphite, 24 grains.
Sodium hypophosphite, 16 grains.
 - 8 Calcium and Sodium lactophosphates.
Calcium lactophosphate, 16 grs.
Sodium lactophosphate, 8 grains.
 - 9 Calcium, Sodium, Iron and Potassium lactophosphates.
Calcium lactophosphate, 8 grains.
Sodium lactophosphate, 2 grains.
Iron lactophosphate, 4 grains.
Potassium lactophosphate, 2 grs.
 - 10 Chloral hydrate, 80 grains.
 - 11 Dover's powder.
Dover's powder, U. S. P., 1880, 40 grains; adjusted to a uniform strength by assay. Each fluidounce contains 0.56 grains morphine alkaloid, equivalent to 3-4 grain morphine sulphate.
 - 12 Hematic Hypophosphites, Improved.
A scientific substitute for Fellow's Hypo-phos-phites, which faithfully reproduces that preparation in all essential particulars. Our preparation contains the same medicinal ingredients as Fellow's, but is a permanently clear syrup, while that preparation throws down an unsightly precipitate. We offer no certificates of the value of this preparation, but allow the physician to judge for himself and from its published formula, what are its merits.
Potassium hypophos., 1-2 grs.
Manganese hypophosphite, 1 gr.
Strychnine hypophosphite, 1-16 gr.
Iron hypophosphite, 1-4 grains.
Calcium hypophosphite, 1 grain.
Quinine hypophosphite, 7-16 gr.
 - 13 Hydriodic acid, U. S. P.
1 per cent. absolute hydriodic acid.
 - 14 Hypophosphites, Lime, Soda and Potass., Churchill's.
Calcium hypophosphite, 16 grs.
Sodium hypophosphite, 16 grains.
Potassium hypophosphite, 8 grs.
 - 15 Hypophosphites Compound.
Calcium hypophosphite, 16 grs.
Sodium hypophosphite, 16 grains.
Potassium hypophosphite, 8 grs.
Iron hypophosphite, 2 grains.
- Hypophosphites Comp. with Quinine, Strychnine and Manganese (see Hematic Hypophosphites).*
- 16 Iron bromide, 60 grains.
 - 17 Iron citrate, 40 grains.
 - 18 Iron hypophosphite, 16 grains.
 - 19 Iron iodide, U. S. P.
Contains 10 per cent. (by weight) ferrous iodide.
 - 20 Iron lactate, 8 grains.
 - 21 Iron lactophosphate, 16 grains.
 - 22 Iron and Manganese hypophosphites.
Iron hypophosphite, 16 grains.
Manganese hypophosphite, 16 grs.
 - 23 Iron and Manganese iodides.
Iron iodide, 16 grains.
Manganese iodide, 8 grains.
 - 24 Iron phosphate, 24 grains.
 - 25 Iron protoxide, 25 grains.
Contains iron protoxide, equivalent to 8 grains of ferrous sulphate.
 - 26 Iron pyrophosphate, 24 grains.
 - 27 Iron and Quinine citrate, 10 grains.
 - 28 Iron, Quinine and Strychnine phosphates.
Iron phosphate, 16 grains.
Quinine phosphate, 8 grains.
Strychnine phosphate, 8-60 grains.
 - 29 Iron and Sodium albuminate, 4 grains.
 - 30 Iron and Strychnine citrate, 16 grains.
 - 31 Iron superphosphate, 10 grs.
 - 32 Lactated Pepsin, 40 grains.

MEDICINAL SYRUPS—Continued.

<p>33 Lactated Pepsin Compound.
Lactated pepsin, 24 grains.
Calcium phosphate, 8 grains.
Iron phosphate, 8 grains.
Potassium phosphate, 8 grains.
Sodium phosphate, 8 grains.</p> <p>34 Lactophosphates Compound.
Calcium lactophosphate, 16 grains.
Iron lactophosphate, 8 grains.
Manganese lactophosphate, 4 grs.</p> <p>35 Manganese hypophosphite, 8 grains.</p> <p>36 Manganese iodide, 16 grains.</p> <p>37 Manganese lactophosphate, 24 grains.</p> | <p>38 Phosphates Compound (Chemical Food).
Iron phosphate, 8 grains.
Calcium phosphate, 20 grains, and smaller quantities of sodium and potassium phosphates.</p> <p>39 Potassium Citrate and Ipecac.
Formula suggested by H. C. Wood, M. D., in the <i>Therapeutic Gazette</i>, Jan. 15, 1885.
Potassium citrate, 80 grains.
Lemon juice, 160 minims.
Syrup ipecac, 40 minims.</p> <p>40 Potassium lactophosphate, 16 grains.</p> <p>41 Sodium hypophosphite, 32 grs.</p> <p>42 Sodium lactophosphate, 16 grs.</p> <p>43 Yerba Santa, aromatic.
Yerba santa, 16 grains.
Aromatics, q. s.</p> |
|---|---|

HYPODERMIC TABLETS.

Soluble. Accurate. Permanent.

These tablets are surprisingly soluble and can be made ready for administration at a moment's notice. If dropped into water they disintegrate almost instantly and become dissolved on shaking or stirring to a perfectly limpid solution. The division of material is effected with the greatest exactness, so that each tablet can be relied upon to contain precisely the quantity of medicament stated on the label. Finally the tablets will keep indefinitely in any climate, and a full line of them with syringe can be carried without any trouble in a vest-pocket case.

The tablets weigh about 1-2 grain each, and are put up in tubes convenient for carrying in the pocket medicine case, each containing 25 tablets. The materials of which they are composed are absolutely non-irritating in character. They are to be dissolved in a definite quantity of water, forming thus a solution of known strength of which a part or the whole may be administered.

1 Aconitine (crystals), 1-120 gr. 2 Apomorphine muriate, 1-10 gr. 3 Atropine sulphate, 1-50 grain. 4 Atropine sulphate, 1-100 grain. 5 Cocaine hydrochlorate, 1-4 gr. 6 Coniine hydrobromate, 1-8 gr. 7 Colchicine, 1-30 grain. 8 Corrosive sublimate and Urea, 1-25 grain. 9 Digitalin (soluble), 1-100 grain. 10 Gelsemine muriate, 1-50 grain. 11 Hyoscine hydrobrom., 1-100 gr. 12 Hyoscyamine, 1-50 grain. 13 Morphine sulphate, 1-4 grain. 14 Morphine sulphate, 1-8 grain.	<p>25 Morphine sulphate, 1-6 grain.</p> <p>15 Morphine and Atropine, No. 1.
Morphine sulphate, 1-8 grain.
Atropine sulphate, 1-200 grain.</p> <p>16 Morphine and Atropine, No. 2.
Morphine sulphate, 1-4 grain.
Atropine sulphate, 1-150 grain.</p> <p>24 Morphine and Atropine, No. 3.
Morphine sulphate, 1-6 grain.
Atropine sulphate, 1-180 grain.</p> <p>17 Picrotoxin, 1-50 grain.</p> <p>18 Pilocarpine hydrochlor., 1-3 gr.</p> <p>19 Pilocarpine nitrate, 1-3 grain.</p> <p>20 Physostigmine sulph., 1-100 gr.</p> <p>21 Sodium arseniate, 1-10 grain.</p> <p>22 Strychnine sulphate, 1-50 gr.</p> <p>23 Strychnine sulphate, 1-100 gr.</p> |
|---|---|

Rules for Guidance in Hypodermic Medication.

1. Give only about one-third the usual dose by mouth.
2. See that a perfect solution only is used.
3. Avoid puncturing a vein.
4. Avoid inflamed tissues and bony protuberances.
5. Use freshly prepared solutions.

COMPRESSED TABLETS.

Each tablet contains the quantity of ingredients specified.

Ammonium Chloride, 3 grains.—Promotes secretion, principally of the mucous membrane of the mouth and throat. In chronic bronchitis, where dryness of the throat is a prominent symptom, the bronchial irritation, hoarseness and huskiness of voice, are promptly relieved. Especially indicated where a tenacious mucous secretion provokes an unavailing, persistent cough.

Dose, 1 to 2, allowed to dissolve in the mouth, not oftener than once an hour.

Borax, 5 grains.—Constitutional effect much less pronounced than that of ammonium chloride, but local effect on the throat somewhat similar. Its characteristic effect is soothing, while the former is stimulating. It possesses antiseptic action rendering it useful in aphthous affections. Very serviceable to public speakers and singers in irritable conditions of the throat, resulting either from chronic inflammation or excessive use of the voice.

Digestive (Pepsin, pure, 1 grain; Pancreatin, 1 grain; Calcium lacto-phosphate, 2 grains).—Useful in all forms of indigestion dependent upon any deficiency in the activity of these secretions or upon indiscretion in eating.

Dover's Powder, 5 grains.

Mercuric Iodide (soluble), 1 grain.—The soluble iodide of mercury is a germicide of extraordinary power—much more active than corrosive sublimate.

For a strong solution for disinfecting the hands, sponges, etc., dissolve one tablet in one-half pint of water. For a solution to be used as a spray in treating diphtheria, use one tablet to a pint of water. For intra-uterine injections or as an antiseptic surgical dressing, dissolve one tablet in one and one-half to two pints of water.

Peptonizing.—Put up in boxes of two vials each. One of these vials contains tablets of sodium bicarbonate, 10 grains, and the other tablets of pure pancreatin, 2 1-2 grains, with sugar of milk, q. s. ad. 10 grains. Recent experiments, however, have shown* that pancreatin is not alone active in alkaline solutions. These tablets are used for preparing peptonized milk, gruel, beef-tea, etc.

Potassium Bi-carbonate, 5 grains.—To be preferred to sodium bicarbonate, especially where there exists a tendency to rheumatism.

Potassium Bromide, 5 grains.—Sedative, producing its effects probably by its action on the cerebral circulation. A valuable remedy for congestive headaches, for the sleeplessness produced by excessive brain work, in the nervousness following alcoholic excesses, and also as an anaphrodisiac.

Potassium Chlorate, 5 grains.—Employed for hoarseness and bronchial irritation; exerts an alterative influence over chronic inflammations of the throat, as well as over all diphtheritic affections; of value in aphthous and ulcerative stomatitis, and in those follicular inflammations commonly known as ulcerated sore throat; has the important property of deodorizing offensive discharges.

In chronic cases patients should be cautioned against using more than six to ten tablets daily, as the remedy may induce renal irritation.

Potassium Chlorate (3 1-2 grains) and **Ammonium Chloride** (1 1-2 grains).—Each remedy produces its peculiar effect, the potassium chlorate acting as an alterative to the diseased mucous membrane, or as an antiseptic, while the ammonium chloride stimulates the mucous secretion and promotes absorption.

Potassium Chlorate (2 1-2 grains) and **Borax** (2 1-2 grains).—In this the soothing effect of borax is added to the alterative influence of potassium chlorate. An efficient remedy in many cases of chronic follicular pharyngitis.

Sodium Bi-carbonate, 5 grains.—Many of the distressing symptoms of dyspepsia, such as flatulence, heartburn, gastralgia and nausea, are due to excessive acidity. While medical treatment should in all cases be directed to a removal of the cause of the acid condition, this should not prevent the use in the meantime of a remedy in itself harmless, which is capable of affording immediate relief.

Soda Mint, 5 grains (Sodium Bi-carbonate and Oil of Peppermint).—This combination of a stimulant and carminative like the oil of peppermint with an antacid has a wide range of use in dyspeptic disorders. These tablets will often give prompt relief in cases of heartburn, flatulence, nausea from indigestion, sick-headache, etc. Especially useful when indigestion is the result of indiscretions in diet.

Yellow Oxide of Mercury, 1-200 grain (1-3 milligramme). Yellow oxide of mercury is a powerful antiseptic, preventing the putrefactive changes which often take place in ingested food, and thus indirectly promoting nutrition, restoring appetite, and removing dyspeptic symptoms. Useful in all acute or chronic derangements of the alimentary tract, in typhoid fever, phthisis, convalescence from surgical diseases, and wherever an intestinal antiseptic is indicated.

* Pharmaceutical Journal and Transactions, Aug. 30, 1887.

TABLET TRITURATES.

"An easy, economical and accurate method of dispensing medicines;" unsurpassed for permanency and also for solubility or ease with which disintegration is effected. We would direct especial attention to the fact that each tablet is stamped with its individual number, thus identifying it without possible chance of mistake.

List No.

- 187 Acid arsenious, 1-200 grain.
- 1 Acid arsenious, 1-100 grain.
- 2 Acid arsenious, 1-50 grain.
- 3 Acid arsenious, 1-30 grain.
- 4 Acid arsenious, 1-20 grain.
- 5 Fl. Ext. Aconite root, 1-10 minim.
- 6 Fl. Ext. Aconite root, 1 minim.
- 7 Aconitine crystals, 1-500 grain.
- 8 Aconitine crystals, 1-200 grain.
- 9 Agaricin, 1-20 grain.
- 10 Aloin, 1-10 grain.
- 11 Aloin, 1-4 grain.
- 12 Aloin, 1-2 grain.
- 181 Aloin and Belladonna Comp.
Aloin, 1-5 grain.
Ext. Belladonna, 1-8 grain.
Strychnine sulph., 1-60 grain.
- 13 Ammonium muriate, 1 grain.
- 182 Ammonium muriate Comp.
Ammonium mur., 1-4 grain.
Ext. Licorice, 1-10 grain.
Powd. Cube, 1-8 grain.
- 193 Antimony Comp. (Plummer's) 1-10 gr.
Antimony sulphurated, 1-40 gr
Calomel, 1-40 grain.
Powd. Guaiac, 1-20 grain.
- 180 Antiseptic, (Bernay's).
CAUTION.—For external use only.
Corrosive sublimate, 1 3-4 grains.
Citric acid, 87-100 grain.
- 14 Apomorphine muriate, 1-100 grain.
- 15 Apomorphine muriate, 1-50 grain.
- 16 Arsenic iodide, 1-100 grain.
- 17 Atropine sulphate, 1-500 grain.
- 18 Atropine sulphate, 1-200 grain.
- 19 Atropine sulphate, 1-100 grain.
- 20 Fl. Ext. Belladonna, 1-2 minim
- 21 Fl. Ext. Belladonna, 1 minim.
- 22 Bismuth subcarbonate, 2 grains.
- 23 Bismuth subnitrate, 2 grains.
- 188 Bryonia tincture, 5 minims.
- 24 Caffeine, 1-2 grain.
- 25 Caffeine, 1 grain.
- 26 Caffeine citrate, 1-2 grain.
- 27 Caffeine citrate, 1 grain.
- 28 Calcium sulphide, 1-10 grain.
- 29 Calcium sulphide, 1-4 grain.
- 30 Calcium sulphide, 1-2 grain.
- 31 Calomel, 1-100 grain.
- 32 Calomel, 1-10 grain.
- 33 Calomel, 1-4 grain.
- 34 Calomel, 1-2 grain.
- 35 Calomel, 1 grain.
- 195 Calomel, Ipecac and Sodium bicarb.
Calomel, 1-5 grain.
Ipecac, 1-10 grain.
Sodium bi-carb., 1 grain.

List No.

- 196 Chamomile tincture, 2 minims.
- 36 Cannabis Indica extract (Indian Cannabis), 1-20 grain.
- 37 Cannabis Indica extract (Innian Cannabis), 1-10 grain.
- 38 Capsicum, 1 grain.
- 40 Cerium oxalate 1 grain.
- 41 Cocaine hydrobromate, 1-16 grain.
- 42 Cocaine hydrobromate, 1-8 grain.
- 43 Cocaine hydrochlorate, 1-16 grain.
- 44 Cocaine hydrochlorate, 1-8 grain.
- 45 Codeine, 1-4 grain.
- 46 Codeine, 1-2 grain.
- 47 Colchicine, 1-100 grain.
- 48 Colchicine, 1-50 grain.
- 49 Colocynthin, 1-20 grains.
- 50 Coniine hydrobromate, 1-20 grain.
- 51 Coniine hydrobromate, 1-10 grain.
- 52 Coniine hydrochlorate, 1-20 grain.
- 53 Coniine hydrochlorate, 1-10 grain.
- 54 Convallamarin, 1-50 grain.
- 55 Corrosive sublimate, 1-1000 grain.
- 56 Corrosive sublimate, 1-500 grain.
- 57 Corrosive sublimate, 1-100 grain.
- 58 Corrosive sublimate, 1-50 grain.
- 59 Corrosive sublimate, 1-20 grain.
- 60 Corrosive sublimate, 1-16 grain.
- 61 Digitalin, pure, 1-100 grain.
- 62 Digitalin, pure, 1-75 grain.
- 63 Digitalin, pure, 1-50 grain.
- 64 Fl. Ext. Digitalis, 1-10 minim.
- 65 Fl. Ext. Digitalis, 1-2 minim.
- 66 Fl. Ext. Digitalis, 1 minim.
- 67 Dover's Powder, 1-4 grain.
- 68 Dover's Powder, 1 grain.
- 69 Elaterin, 1-40 grain.
- 70 Elaterin, 1-20 grain.
- 71 Elaterin, 1-10 grain.
- 72 Ergotin, 1-10 grain.
- 73 Ergotin, 1-4 grain.
- 197 Gelsemium tincture, 1 minim.
- 74 Hydrastine (white alkaloid), 1-25 gr.
- 75 Hydrastine (white alkaloid), 1-16 gr.
- 76 Hydrastine (white alkaloid), 1-8 gr.
- 77 Hydrastine (white alkaloid), 1-4 gr.
- 78 Hyoscyamine crystals, 1-200 grain.
- 79 Hyoscyamine crystals, 1-100 grain.
- 80 Ignatia, 1-20 grain.
- 81 Ignatia, 1-10 grain.
- 82 Ipecac, 1-10 grain.
- 83 Ipecac, 1-4 grain.
- 84 Ipecac, 1-2 grain.
- 189 Iron arseniate, 1-16 grain.
- 87 Iron arseniate, 1-10 grain.
- 85 Iron by hydrogen (Q'venne's) 1-2 gr

TABLET TRITURATES—Continued.

List No.

- 86 Iron by hydrogen (Q'venne's) 1 gr.
 183 Iron and Arsenic Comp.
 Iron by Hydrogen, 1 grain.
 Powd. Ignatia, 1-40 grain.
 Arsenious acid, 1-100 grain.
 88 Iron and Quinine citrate, 1-4 grain.
 89 Iron and Quinine citrate, 1-2 grain.
 90 Iron and Quinine citrate, 1 grain.
 91 Iron and Strychnine citrate, 1-4 grain.
 92 Iron and Strychnine citrate, 1-2 gr.
 93 Iron and Strychnine citrate 1 grain.
 198 Iron lactate, 1 grain.
 184 Iron, Quinine and Strychnine.
 Iron by hydrogen, 1-2 grain.
 Strychnine sulph., 1-120 grain.
 Quinine sulph., 1-2 grain.
 185 Iron and Strychnine Comp.
 Iron by hydrogen, 1-2 grain.
 Arsenious acid, 1-100 grain.
 Quinine sulph., 1-2 grain.
 Strychnine sulph., 1-120 grain.
 94 Leptandrin, 1-16 grain.
 95 Lithium carbonate, 1 grain.
 96 Manganese binoxide, 1-2 grain.
 97 Manganese binoxide, 1 grain.
 98 Mercury with chalk, 1-10 grain.
 99 Mercury with chalk, 1-5 grain.
 100 Mercury with chalk, 1 grain.
 101 Mercury biniodide, 1-100 grain.
 190 Mercury biniodide, 1-50 grain.
 102 Mercury biniodide, 1-16 grain.
 103 Mercury biniodide, 1-8 grain.
 104 Mercury protiodide, 1-100 grain.
 105 Mercury protiodide, 1-50 grain.
 106 Mercury protiodide, 1-16 grain.
 107 Mercury protiodide, 1-8 grain.
 108 Mercury protiodide, 1-6 grain.
 109 Mercury protiodide, 1-4 grain.
 110 Mercury protiodide, 1-2 grain.
 111 Mercury protiodide, 1 grain.
 191 Mercury, yellow oxide, 1-200 grain.
 112 Morphine and Atropine.
 Morphine sulphate, 1-8 grain.
 Atropine sulphate, 1-150 grain.
 113 Morphine and Atropine.
 Morphine sulphate, 1-4 grain.
 Atropine sulphate, 1-120 grain.
 114 Morphine hydrochlorate, 1-8 grain.
 115 Morphine hydrochlorate, 1-4 grain.
 116 Morphine sulphate, 1-20 grain.
 117 Morphine sulphate, 1-16 grain.
 118 Morphine sulphate, 1-8 grain.
 119 Morphine sulphate, 1-6 grain.
 120 Morphine sulphate, 1-4 grain.
 121 Morphine sulphate, 1-2 grain.
 122 Nitroglycerin, 1-100 grain.
 123 Nitroglycerin, 1-50 grain.
 124 Nux Vomica, 1-100 grain.
 125 Nux Vomica, 1-50 grain.
 126 Nux Vomica, 1-10 grain.
 127 Nux Vomica, 1-4 grain.
 128 Opium, 1-4 grain.

List No.

- 129 Opium, 1-2 grain.
 130 *Pepsin (pure), 1-4 grain.
 131 *Pepsin (pure), 1-2 grain.
 132 *Pepsin (pure), 1 grain.
 194 Pepsin, lactated, 1 grain.
 133 Physostigmine salicylate, 1-500 grain.
 134 Physostigmine salicylate, 1-400 grain.
 135 Physostigmine salicylate, 1-300 grain.
 136 Physostigmine salicylate, 1-200 grain.
 137 Physostigmine salicylate, 1-100 grain.
 138 Pilocarpine nitrate, 1-100 grain.
 139 Pilocarpine nitrate, 1-50 grain.
 140 Pilocarpine nitrate, 1-25 grain.
 141 Pilocarpine nitrate, 1-10 grain.
 142 Podophyllin, 1-16 grain.
 143 Podophyllin, 1-8 grain.
 144 Podophyllin, 1-4 grain.
 145 Podophyllin, 1-2 grain.
 199 Poison oak tincture, 1 minim.
 146 Quinine bisulphate, 1-4 grain.
 147 Quinine bisulphate, 1-2 grain.
 148 Quinine bisulphate, 1 grain.
 149 Quinine sulphate, 1-4 grain.
 150 Quinine sulphate, 1-2 grain.
 151 Quinine sulphate, 1 grain.
 200 Saccharin, 1-2 gr.
 152 Salicylic acid, 1-4 grain.
 153 Salicylic acid, 1-2 grain.
 154 Santonin, 1-2 grain.
 155 Santonin, 1 grain.
 156 Santonin and Calomel.
 Santonin, 1-2 grain.
 Calomel, 1-2 grain.
 157 Sodium arseniate, 1-25 grain.
 158 Sodium arseniate, 1-15 grain.
 159 Sodium arseniate, 1-10 grain.
 186 Sodium salicylate, 1 grain.
 160 Strychnine sulphate, 1-100 grain.
 161 Strychnine sulphate, 1-60 grain.
 162 Strychnine sulphate, 1-50 grain.
 163 Strychnine sulphate, 1-40 grain.
 164 Strychnine sulphate, 1-30 grain.
 165 Strychnine sulphate, 1-20 grain.
 166 Strychnine sulphate, 1-16 grain.
 167 Tartar emetic, 1-100 grain.
 168 Tartar emetic, 1-50 grain.
 169 Tartar emetic, 1-25 grain.
 170 Tartar emetic, 1-16 grain.
 171 Tartar emetic, 1-8 grain.
 172 Tartar emetic, 1-4 grain.
 173 Veratrine, 1-32 grain.
 174 Veratrine, 1-12 grain.
 192 Zinc phosphide, 1-20 grain.
 175 Zinc phosphide, 1-10 grain.
 176 Zinc phosphide, 1-8 grain.
 177 Zinc phosphide, 1-6 grain.
 178 Zinc phosphide, 1-4 grain.
 179 Zinc phosphide, 1-2 grain.

* One gr. pure pepsin equals 10 gra.
 sacch. pepsin, U. S. P.

MEDICATED LOZENGES.

A neat, eligible and convenient means of exhibiting many remedies. Lozenges are especially suited for administering remedies to children, and their use as palatable vehicles for cough or throat remedies is commended since in addition to the local action secured by slow solution in the saliva there is added that of their portability.

The formulæ given show the approximate quantity of active ingredients in each lozenge.

Alum and Kino.

Powd. Alum, 1 grain.
Powd. Kino, 1 grain.

Ammonia, Dr. Jackson's.

Ammonium muriate, 1-2 grain.
Morphine muriate, 1-60 grain.

Ammonium chloride, 2 grains.

Bismuth and Charcoal.

Bismuth subnitrate, 1 grain.
Charcoal, willow, 1 grain.

Bronchial (with Cubeb).

Ext. Licorice, 4-5 grain.
Balsam Tolu, 1-20 grain.
Powd. Cubeb, 5-8 grain.
Oil Sassafras, 1-40 grain.

Carbolic Acid, 1 grain.

Catechu, 2 grains.

Charcoal, Willow, 4 grains.

Chocolate and Iron.

Carb. Iron, 2 grains.

Chocolate and Quinine.

Quinine sulphate, 1-2 grain.

Chocolate Worm.

Santonin, 1-2 grain.
Calomel, 1-2 grain.
Chocolate, q. s.

Coltsfoot, 2-3 grain.

Cubeb.

Ext. Licorice, 3-4 grain.
Oil Sassafras, 1-32 grain.
Oleoresin Cubeb, 1-16 gr.

Eucalyptus globulus.

Ext. Eucalyptus, true, 3 grains.
Ext. Licorice, 1 grain.

Ginger, U. S. P., 2 grains tincture.

Ipecac, U. S. P., 1-4 grain.

Ipecac and Morphine.

Powd. Ipecac, 1-12 grain.
Morphine sulphate, 1-40 grain.

Iron carbonate, 4 grains.

Licorice.

Marshmallow.

Powd. Marshmallow, 1-3 grain.
Orris root, 1-4 grain.

Opium and Licorice, see Wistar's.

Pectoral, Dr. Jackson's.

Ipecac, 1-20 grain.
Morphine muriate, 1-33 grain.
Antimony sulphurated, 1-40 grain.

Pepsin, Sacch. U. S. P., 3 grains.

Pepsin and Bismuth.

Sacch. pepsin, U. S. P., 2 grains.
Bismuth subnitrate, 2 grains.

Pepsin and Iron.

Sacch. pepsin, U. S. P., 1 1-2 grs.
Iron pyrophosphate, 1 1-2 grains.

Pepsin and Lime lactophosphate.

Sacch. Pepsin, U. S. P., 2 grains.
Calcium lactophosphate, 3 grains.

Potassium chlor., 2 and 5 grains.

Potassium chlor. (Chocolate), 2 and 5 grains.

Potassium chlor. (Lemon), 2 grs.

Potassium chlor. (Wintergreen), 2 grains.

Potassium chlor. and Ammonium chlor. (Chocolate), 1 gr. each.

Rose-leaf and Alum.

Powd. Rose leaves, 1-4 grain.
Powd. Alum, 1-4 grain.

Salicylic Acid, 1 grain.

Santonin, 1-2 and 1 grain.

Santonin and Calomel (Worm Candy).

Santonin, 1-2 grain.
Calomel, 1-2 grain.

Santonin compound.

Santonin, 1-2 grain.
Calomel, 1-2 grain.
Podophyllin, 1-20 grain.

Santonin and Podophyllin.

Santonin, 1-2 grain.
Podophyllin, 1-20 grain.

Sodium bicarbonate, U. S. P., 3 grs.

Tannic Acid, U. S. P., 1 grain.

Tar, 1-7 grain.

Tar compound.

Pine Tar, 7-48 grain.
Powd. Senega, 7-24 grain.
Powd. Wild Cherry, 2 1-3 grains.

Terebene, 2 grains.

Wild Cherry, 1 grain.

Wistar's Cough.

Powd. Opium, 1-10 grain.
Oil Anise, 1-32 grain.
Ext. Licorice, 1 grain.
Tinct. Tolu, 1-16 grain.

Worm Candy, see Santonin and Calomel.

Yerba Santa compound.

Ext. Licorice, 1 grain.
Powd. Senega, 3-16 grain.
Tar, 1-32 grain.
Powd. Cherry, 1-4 grain.
Yerba santa, 1 grain.

MEDICINAL WINES.

Each fluidounce represents the medicinal virtues of the quantity of drug specified in the respective formulae.

- 1 American White Ash, 120 grs.
- 2 Antimonial, U. S. P.
Ant. and Pot. tartrate, 1 4-5 grs.
- 3 Aromatic, U. S. P.
This wine possesses strong tonic and aromatic properties, having an advantage over the aromatic and tonic tinctures in being less stimulating; used principally as an external application.
- 4 Beef and Wine.
Beef, 2 ounces.
- 5 Beef, Iron and Wine.
* A valuable nutrient tonic and stimulant. Most of the beef extracts in the market are of no value as food, being almost wholly destitute of proteids or nutrient principles. In this preparation we have substituted for Liebig's beef extract a Peptonized Extract of Beef of our own manufacture, which is a true food containing the proteids or tissue-forming constituents in the form of peptones, in addition to those mineral salts to which the ordinary beef extracts owe any virtue they may possess. Each fluidounce of our Beef, Iron and Wine contains the equivalent of an ounce of fresh beef, peptonized, and 4 grains citrate of iron and ammonium.
- 6 Beef, Iron, Wine and Cinchona.
Beef, 2 ounces.
Iron and ammon. cit., 4 grains.
Calisaya bark (alkaloids), 40 grs.
- 7 Beef, Iron and Wine with Pepsin.
Beef, 2 ounces.
Iron and ammon. cit., 4 grains.
Sacch. pepsin, U.S.P., 8 grains.
- 8 Calisaya bark, true, 40 grains.
- 9 Calisaya and Iron.
Calisaya bark (alkaloids), 40 grs.
Iron pyrophosphate, 8 grains.
- 10 Coca.
Each fluidounce contains the equivalent of 60 grains of coca. We have adopted an improved formula for this preparation, which is made of the same strength as formerly, although materially different in appearance and flavor.
- 11 Coca and Beef.
Coca, 60 grains.
Beef, 2 ounces.
- 12 Coca, Beef and Iron.
Coca, 60 grains.
Beef, 2 ounces.
Iron and ammon. cit., 4 grains.
- 13 Colchicum root, U. S. P., 180 grains.
- 14 Colchicum seed, U. S. P., 70 grains.
- 15 Corn silk, 120 grains.
- 16 Ergot, U. S. P., 70 grains.
- 17 Ipecac, U. S. P., 32 grains.
- 18 Iron, Bitter, U. S. P.
Calisaya bark (alkaloids), 40 grs.
Iron and ammon. cit., 16 grains.
- 19 Iron citrate, U. S. P.
Iron and ammon. cit., 18 grains.
- 20 Opium, U. S. P., assayed.
Vinum Opii, U. S. P., 1880; each fluidounce contains by assay 6 grains of morphine, equivalent to 7 1-2 grains of morphine sulphate.
- 21 Pepsin.
Sacch. pepsin, U. S. P., 10 grains.
- 22 Pepsin and Bismuth.
Sacch. pepsin, U. S. P., 8 grains.
Bismuth and ammon. cit., 8 grs.
- 23 Rhubarb, U. S. P.
Rhubarb, 45 grains.
Sweet flag, 4 grains.
- 24 Wild Cherry Bark, 80 grains.
- 25 Wild Cherry and Iron.
Wild cherry bark, 80 grains.
Iron pyrophosphate, 8 grains.

FLAVORED LOZENGES.

These are made of pure materials and the finest flavoring, and are especially adapted for druggists' use. They are made in two grades, x and xx, which differ only in strength of flavor, the same high quality of ingredients being used for both.

Capsicum.	Lemon.	Rose.
Cinnamon.	Musk.	Sassafras.
Cloves.	Peppermint.	Wintergreen.

OLEATES

Of the Principal Alkaloids and Metals.

These compounds, which have recently aroused much attention and which bid fair to enlarge the practice of dermic medication considerably, supplanting the less desirable agents heretofore employed, have been added to our manufactures under the belief that the employment of so serviceable a class of remedies will soon become popular. They are of very superior penetrative power, and are therefore very economical in the amount of material required to produce a specific effect.

Oleates of the alkaloids are only offered in the diluted form which clinical experience has proved to be the most efficient; they are held in solution in oleic acid.

Aconitine, 2 per cent.

Aluminium.

Used in ointments for astringent effect. Valuable in eczema and as a dressing for burns, etc.

Arsenic.

An ointment of this containing 20 grains of oleate to the ounce, is used as a caustic in the treatment of lupus, epithelioma, etc.

Atropine, 2 per cent.

Bismuth.

Used undiluted, as a soothing application.

Cocaine, 5 per cent.

Copper.

Employed in the form of an ointment, 10 to 20 per cent. oleate, for the treatment of ringworm, etc.

Iron.

A waxy solid, readily soluble in fats. Used as a local astringent and general tonic.

Lead.

An efficient substitute for Hebra's Diachylon Ointment.

Manganese, pure.

Recommended by Drs. Martin and McArthur, of Chicago, as an efficient emmenagogue. A 20 per cent. solution is commonly employed: See below.

Manganese, 20 per cent.

Emmenagogue. Applied by inunction to the abdomen. Said to be preferable to potassium permanganate.

Mercury.

A stable preparation. Produces all the therapeutic effects, local or constitutional, which are peculiar to mercury. Contains 28.3 per cent. of mercuric oxide.

Morphine, 10 per cent.

Morphine and Mercury.

Five per cent. morphine and 20 per cent. mercuric oxide.

Nickel.

An amorphous, waxy solid; good results are reported from its application in some cases of chronic eczema of the extremities, where the skin is hard and leathery.

Quinine, 25 per cent.

Silver.

An ointment containing 10 to 60 grains of the oleate to the ounce is employed to relieve itching and as an application in erysipelas.

Strychnine, 2 per cent.

Tin.

A remedy of considerable value for restoring the lustre of diseased nails.

Veratrine, 10 per cent.

Zinc.

An impalpable powder. Dusted over the surface in skin diseases. Also employed in the form of an ointment.

Ointments of Oleates.

The following ointments are offered as representing the most desirable strengths of the above. The percentages annexed indicate the proportion of *oleate* in the ointment.

Aluminium, 50 per cent.

Iron, 25 per cent.

Silver, 5 per cent.

Arsenic, 5 per cent.

Lead, 50 per cent.

Zinc, 25 per cent.

Copper, 20 per cent.

Mercury, 10 per cent.

CONFECTIONS.

Aromatic.

Orange.

Senna.

Black Pepper.

Rose, U. S. F

Theriac.

Opium, U. S. P., 1870.

OINTMENTS.

Aconite, benzoinated.	Discutient.	Stramonium, U. S. P., zoinated.
<i>Basilicon</i> (see Resin Cerate).	Iodine compound, benzoin- ated.	Sulphur, U. S. P., benzoin- ated.
Bearsfoot.	Mayer's.	Tar, U. S. P.
Belladonna, U. S. P., ben- zoinated.	Mercurial (1-2 Mercury), U. S. P.	Tar compound.
Benzoin.	Mezereon.	Wild Indigo.
Bittersweet.	Olive.	Zinc oxide, U. S. P., ben- zoinated.
Cantharides.	Rose-water (cold-cream), U. S. P.	
Citrine, U. S. P.		
Creasote.		

CERATES.

Cantharides, U. S. P.	Lead subacetate, U. S. P.	Simple, U. S. P.
Cantharides, ext., U. S. P.	Resin, U. S. P.	Soap.
Cetaceum, U. S. P.	Resin compound.	Zinc carbonate.

ROLL PLASTERS.

Aconite.	Calefaciens (Warming).	Mercury.
<i>Adhesive</i> (Resin).	Conium.	Opium, U. S. P.
Ammoniac.	Galbanum, U. S. P.	Pitch Burgundy.
Ammoniac with Mercury.	Iron (Roborans).	Pitch compound.
<i>Anodyne</i> (Opium, U. S. P.).	Lead.	Resin.
Arnica.	Logan's.	Soap.
Belladonna.		

GERMAN TINCTURES.

In order to supply a certain demand for tinctures of the green plant, which the advertisements of other houses have created, we have made direct importations of the following. While guaranteeing the correctness of their preparation according to the standard formulæ, we maintain, for reasons which we have not space here to specify, that they are inferior as therapeutic agents to the preparations of the properly dried plant.

Aconite.	Mary Thistle.
Aconitum Napellus, <i>Linne</i> .	Silybum marianum, <i>Gartner</i> .
Arnica.	Poison Oak.
Arnica montana, <i>Linne</i> .	Rhus Toxicodendron, <i>Linne</i> .
Belladonna.	Pulsatilla.
Atropa Belladonna, <i>Linne</i> .	Anemone Pulsatilla, <i>Linne</i> .
Conium.	Shepherd's Purse.
Conium maculatum, <i>Linne</i> .	Capsella bursa-pastoris, <i>Mench</i> .
Dumb Cane.	Stramonium (Thornapple).
Dieffenbachia seguine, <i>Schott</i> .	Datura Stramonium, <i>Linne</i> .
Foxglove.	Sundew.
Digitalis purpurea, <i>Linne</i> .	Drosera rotundifolia, <i>Linne</i> .
Garden Celandine.	White Bryony.
Chelidonium majus, <i>Linne</i> .	Bryonia alba, <i>Linne</i> .
German Chamomile.	Wild Lettuce.
Matricaria Chamomilla, <i>Linne</i> .	Lactuca virosa, <i>Linne</i> .
Henbane.	
Hyoscyamus niger, <i>Linne</i> .	

COMPOUND CEREBRAL SEDATIVE.

The properties of this preparation are sufficiently indicated by its name. It is offered as a scientific substitute for certain copyrighted preparations in the market, which are especially extolled as valuable remedies in the treatment of nervous affections. A *Compound Cerebral Sedative, Formula B*, differing from the formula given below only in the substitution of hyoscyamus for the opium is also on our list. Each fluidounce contains:

Chloral hydrate, 2 drachms.	Fl. ext. gelsemium, 50 minims.
Potassium bromide, 2 drachms.	Tincture opium, 40 minims.
Dose, a half fluidrachm (2 C.c.).	

MINERAL WATER SALTS.

The composition of these salts is based upon accurate analyses. They are designed for use by those having soda fountains, or in a position to charge solutions of the salts with carbonic acid gas—in this respect differing from the series of Granular Effervescing Preparations (see page 210), these latter containing in themselves the necessary ingredients for the production of this gas.

The numbers affixed to the names of the various salts indicate the quantity in gallons which one pound will produce, to exactly duplicate the natural mineral water desired.

Apollinaris, 30 1-2 gallons.	Hathorn, 8 1-2 gallons.
Carlsbad, 17 1-2 gallons.	High rock, 13 1-4 gallons.
Congress, 11 gallons.	Kissengen, 14 gallons.
Crab Orchard, 13 gallons.	Püllna, 2 1-4 gallons.
Excelsior, 15 1-4 gallons.	Seltzer, 28 gallons.
Friedrichshall, 3 1-4 gallons.	Star, 13 1-4 gallons.
Geyser, 9 gallons.	Vichy, 13 1-4 gallons.

LIST OF SPECIALTIES

MANUFACTURED BY US BUT NOT PREVIOUSLY MENTIONED IN THE FOREGOING PAGES;
ALSO THOSE ARTICLES FOR WHICH WE ARE MANUFACTURER'S
AGENTS. THEY ARE, BRIEFLY:

Amyl Nitrite Pearls; Aromatic Troches, for the breath; Boroglyceride, and also Boroglyceride Solution, 50 per cent.; Bronchial Sedative; Chamberlin's Utero-vaginal Syringe; Chloroform; Cocaine Case, improved; Capsules, empty; Capsule Fillers; Davenport's Unique and Reymond's Patent; Dialyzed Iron, plain and glycerated; Disinfectant and Antiseptic Preparations; Hypodermic Case; Iodoform Bougies, gelatin-coated, 1, 2, 3, 5 and 10 grains in boxes of 12 each; Liquid Acid Phosphate; Peptonized Extract of Beef; Petroleum Mass (used in asthma and bronchial affections); Physician's Pocket Reagent Case; Powdered Bone; Powder of Beef; Sanguis Bovinus Exsiccatus; Semple's Atomizing Inhaler; Sugar Test Flasks; Syringes; The New Ureometer; Urinary Test Papers; Urinary Test Tablets; Wilson's Eureka Abdominal Supporter; Rare Chemicals, Alkaloids, Glucosides, etc.

Several of these require no further comment; the others are briefly described below.

AMYL NITRITE PEARLS.

These pearls are shells of thin glass, each containing 2, 3, 4, 5, 8 or 10 drops of nitrite of amyl, packed in cotton, in boxes of one dozen. They are a serviceable method for the administration of nitrite of amyl. The patient may carry the box about his person without inconvenience, and on the approach of a paroxysm of the malady may crush a pearl in the handkerchief, and by inhaling the vapor find prompt relief. Its action on the heart is almost instantaneous, and it affords great relief in angina pectoris, spasmodic asthma, epilepsy, syncope, etc.

COCAINE CASE, IMPROVED.

This case is made in the best style of workmanship, of the finest morocco, velvet lining, and contains:

1. Five capsules, each containing exactly one grain of cocaine muriate in crystals.

2. A vial to contain a solution of cocaine muriate.

3. A minim pipette.

4. A camel's-hair pencil.

5. A place for holding a hypodermic syringe.

6. A card containing formulæ and directions for making 2 per cent. and 4 per cent. solutions of muriate of cocaine.

It is advisable in all cases to prepare solutions of cocaine salts fresh when they are required for use. To prepare a 4-per-cent. solution, it is only necessary to empty the contents of one of the capsules into the vial, and fill to the mark with distilled water.

We furnish these cases with or without a hypodermic syringe. Should the physician already have a hypodermic syringe, the latter case (which is provided with a place for holding it) will undoubtedly meet his requirements.

CAPSULES, IMPROVED EMPTY GELATIN:

In seven sizes for oral exhibition; three sizes for rectal and vaginal administration; Veterinary (horse) capsules, three sizes.

IMPROVED EMPTY GELATIN CAPSULES.

RECTAL CAPSULES.

Our empty capsules, for oral, rectal, and vaginal administration, and veterinary purposes are simply perfect as regards those qualities which commend to the practitioner these aids to the exhibition of medicines. Their solubility is perfect, and the fear of their passing undigested through the alimentary tract, as has been charged against other brands of these articles, is absolutely groundless. They cannot be excelled in quality by any in use. As a matter of professional interest we append a table showing the capacity of our Improved Empty Gelatin Capsules, when filled by hand. Larger quantities can be enclosed by using capsule fillers. (See article below.)

For further information upon capsules and other gelatin products, please apply to us for printed matter and samples.

HORSE CAPSULES.

APPROXIMATE CAPACITY OF EMPTY CAPSULES.

The capacity of empty capsules varies according to the degree of compression exerted in filling and the known variation in weight of the same bulk of different samples of the same powdered drug or alkaloid.

SIZES.	00	0	1	2	3	4	5
	grs.	grs.	grs.	grs.	grs.	grs.	grs.
Quinine sulphate.....	8	6	4	3	2	1½	¾
Cinchonidine sulphate	8	6	4	3	2	1½	¾
Powdered Ipecac Comp. (U. S. P., 1880.)	12	8	6	4	3	2	1
Powdered Aloes,	10	7	5	4	3	2	1
Bismuth subnitrate.....	20	14	10	8	6	4	2
Salicin.....	10	7	5	4	3	2	1
Powdered Rhubarb.....	12	8	6	4	3	2	1
Salicylic Acid.....	8	6	4	3	2	1½	¾

CAPSULE FILLERS.

Davenport's Unique Capsule Filler.

With this device gelatin capsules can be filled with powders more rapidly, and a larger quantity of medicine can be packed in a given size than by the ordinary method. By its use the outside of the capsules cannot become soiled with the bitter or nauseous medicines with which they are usually filled, and capsules that have become damp or soft can be filled as readily as any. The capsules being filled with a dry powder present a much more elegant appearance than when the tedious and disagreeable practice of first forming a mass is resorted to, and many persons who object to taking pills or capsules filled with pill mass, do not hesitate to take capsules when filled with the dry powder. The powder being packed in the body admits of the cap being entirely on the capsule, so that the powders in expanding are not liable to press the cap off after the prescription is placed in the hands of the customer, as often happens when both body and cap are filled.

Reymond's Capsule Filler.

The attention of pharmacists and physicians is called to this simple contrivance for filling capsules. No well equipped pharmacy should be without one. By its use the inconvenience of dispensing medicines in capsules is entirely removed, and it may be conveniently carried by the physician and used at the bedside, a little experience in its use making the filling of capsules nearly as expeditious as the enfolding of powders. Its advantages over other inventions of a similar nature are:

1. Simplicity previous experience is not required to operate it successfully
2. Economy in time; with other fillers only one capsule can be filled at a time; with the Reymond a dozen.
3. Economy in capsules; the method of filling the capsules is so simple that there is no possibility of breaking them.
4. Economy in material; the liability of waste while filling the capsules is reduced to a minimum.
5. Price; it is the cheapest filler in the market, the amount of work it will do being considered.

DIALYZED IRON.

The therapeutic merits of dialyzed iron, and its superiority over other ferruginous preparations, are now universally admitted. Each fluidounce contains 24 grains ferric oxide, or 16.79 metallic iron. It is free from disagreeable taste, has no injurious action on the teeth, stomach, or bowels, and is, therefore, suited to cases where other forms of iron are not tolerated. It is also an antidote for arsenical poisoning.

Dialyzed iron, however, has the disadvantage of great liability to congeal under exposure to low degrees of temperature, thus rendering its transportation during the winter months very hazardous and unprofitable. To overcome this difficulty we furnish a preparation of this article, known as our Glycerated Dialyzed Iron, which is a solution of the peroxide of iron, rendered perfectly pure and neutral by careful dialysis with distilled water, and united with a small percentage of chemically pure glycerin, which prevents it from congealing, and permits its shipment in any weather with perfect safety. It is permanent and will not gelatinize nor mould. The dose of this preparation is the same as that of the ordinary dialyzed iron, viz: from 5 to 30 drops, containing five per cent. of peroxide of iron.

DISINFECTANT AND ANTISEPTIC PREPARATIONS.

In offering these disinfectant and antiseptic preparations, we desire to say that we have aimed to meet three chief indications, the fulfilment of which we believe to be *a sine qua non* to the practical utility and popular use of agents of this nature. These indications are:

a. Adaptation for universal use wherever antiseptics or disinfectants are indicated.

b. Convenience of manipulation, which renders them possible of application by any person of ordinary intelligence

c. Inexpensiveness, which makes their use by all classes of people practicable.

The list comprises the following.

1. **Antiseptic Liquid.**—A colorless solution, free from disagreeable odor. It does not stain the most delicate fabrics. It is an efficient antiseptic and deodorizer, especially adapted to use about the house. Its constituents are the chlorides of zinc, aluminium, magnesium, calcium, and sodium, with oil of eucalyptus and oil of wintergreen (methyl salicylate). Put up in quart bottles

2. **Labarraque's Solution** (Solution Chlorinated Soda, U. S. P.)—A powerful oxidizing disinfectant and bleaching agent, capable of evolving chlorine, which attacks the foul gases that result from putrefaction. Put up in quart bottles.

3. **Sulphur Bricks.**—To be burned in place of crude brimstone to produce sulphurous acid gas; each brick contains one pound of sulphur.

4. **Antiseptic Cologne.**—May be employed in the sick-room in the form of a spray with an ordinary perfume atomizer, to overcome disagreeable odors; also used as a disinfectant for the hands of attendants in cases of infectious disease, or to remove the odor of Labarraque's solution, which is often employed for the same purpose. Applied to the face after shaving, it serves as a sure preventive of contagious affections, sometimes traceable to the barber-shop. Surgeons and physicians will find it to meet several important indications when they are attending patients suffering from infectious diseases; especially to prevent the possibility of their carrying the contagion from one patient to another. Put up in small (1-4 ounce) and large-sized (half-pint) bottles, neatly labeled, and with full directions for use.

5. **Solution Aluminium Acetate, 3 per cent.**—A non-poisonous deodorizing antiseptic. Dr. A. Rose, of New York, recommends this as the best antiseptic for the use of the surgeon. Applied to foul and gangrenous ulcers, it promptly destroys the odor, and restores the tissues to a healthy condition. It is to be preferred to carbolic acid, because, while equally efficient as an antiseptic it is free from the irritating character of that agent, and because it is inodorous. This solution should be diluted with from two to five times its volume of pure water for use in irrigating ulcers or wounds, or for an injection in gynaecological practice. For the latter purpose it was highly recommended by the late Dr. J. Marion Sims, and its use in surgery is favorably spoken of by Billroth.

LIQUID ACID PHOSPHATE.

This preparation is identical with that heretofore known as *Liquor Acidi Phosphorici*, and is superior to certain copyrighted and proprietary preparations of a similar nature upon the market. Physicians have long employed phosphoric acid, and various combinations of the acid with mineral bases, and are familiar with their effects upon the system. It is certainly, in an important sense, a nerve and brain food, and relieves symptoms of mental exhaustion, such as sleeplessness, melancholia, etc., and even increases the capacity for mental labor. This action is so well recognized that the acid phosphate is in considerable demand as a stimulating beverage, dispensed at the soda fountain in place of an alcoholic stimulant.

POWDER OF BEEF.

The experience of Dr. Debove, Professor in the Faculty of Medicine of Paris, at the Bictre and LaPitit Hospitals, in the treatment of diseases of inanition with powdered beef, as prepared according to his own formula, has excited great interest of late in the medical world. Dr. Dujardin-Beaumetz, of the Academy of Medicine and of the Hospital St. Antoine, and other physicians in Paris, report equally favorable results. Dr. Henry B. Millard, of New York, has enjoyed numerous opportunities of seeing it employed by Dr. Debove, and obtaining this author's experience, he says: "The class of ailments in which superalimentation is important are those characterized by emaciation from non-assimilation of food, chronic diarrhoea, nausea and vomiting, including the vomiting of pregnancy, chlorosis, anæmia, exhaustion following severe illness or hemorrhage, and, above all, phthisis pulmonalis;" and, he adds: "I have seen in Paris, and in my own practice at home, all these conditions relieved and cured." Descriptive circular and reprints of Dr. Millard's article above referred to furnished on application.

IMPROVED HYPODERMIC CASE.

We have recently replaced the morocco-covered case previously listed with a very handsomely plated metal one of a flattened-elliptical shape suited for carrying in the vest or coat pocket. The of its size and general appearance.

While we would call attention and elegance of the case itself, we mention the following points of ex-syringe with which the case is supplied.

1. The arrangement which prevents the plunger becoming dry. Attached to the piston-rod on the upper is a small nut which, on the rod being cap, fits there into a socket. If the

to the convenience would also specially celled in the plied.

vides against leak-through disuse.—side of the plunger drawn up to the rod is now turned towards the left, the nut, prevented from turning with the rod by the recess into which it fits, is forced against the leather to the degree found necessary to spread the plunger so as to prevent leakage.

2. The ease with which the needle is cleansed.—On insertion of the wire into the screw-cap it is immediately guided by bevelled sides to the opening into the needle proper. Other syringes have the floor of the cap chamber cut so squarely that it is often after many probings and vexatious delay that the wire can be guided to the opening.

3. The fortified needle. A reinforcement of the needle is effected by a sheath embracing the upper half, thus strengthening it in the part previously most liable to give way under pressure.

The case contains in addition: extra wires for cleaning purposes, and the following selection of hypodermic tablets in tubes of 25 each:

Apomorphine muriate, 1-10 grain.

Atropine sulphate 1-50 grain.

Morphine sulphate, 1-4 grain.

Morphine and Atropine, No. 2 (Morphine sulph., 1-4 gr.; Atropine sulph., 1-150 gr.)

Physostigmine sulphate, 1-100 gr. Strychnine sulphate, 1-50.

We furnish these cases *without syringes*, if desired. The syringes we supply with these cases, however, are of the best approved construction, and will be good investments, even to those who already have hypodermic syringes.

PEPTONIZED EXTRACT OF BEEF.

Analysis has demonstrated that this extract of beef contains from 30 to 40 per cent. of peptones; its nutritive value, therefore, is higher than that of any similar preparation in the market. It is readily soluble in water, contains a very large proportion of albuminoids in the form of peptones, and has a rich, appetizing flavor.

POWDERED BONE.

Bone flour is a true food, not a medicine. It can be taken most readily by incorporating it with the ordinary food. By digesting it a little while in strong vinegar it is rendered wholly soluble, and can then be used as a condiment, with or without addition of cayenne, mustard, walnut catsup, or curry flavor. It offers to the digestive fluids a supply of phosphoric acid much more easily assimilated than that which is contained in the tough structure of the outer coating of cereal grains; it is more digestible, and, we might add, more palatable than bran. Our powdered bone is prepared from carefully selected fresh beef bones deprived of fatty matter and adherent fibrous and muscular tissue. Its approximate composition is organic matter, chiefly collagen, 30; calcium phosphate, 60; magnesium phosphate, 1.7; calcium carbonate, 7.8; calcium fluoride, 0.5. Further descriptive matter furnished on application.

SANQUIS BOVINUS EXSICCATUS.

Dried bullock's blood, defibrinated. This preparation, with the addition of water, is in all respects similar to fresh bullock's blood. As a substitute for beef-tea and the various essences and extracts, it replaces, with a true aliment, what are at best but stimulants. The profession have been latterly much interested in the favorable reports of the use of blood as a rectal injection in low and wasting diseases. The article here presented affords a ready means for the extemporaneous preparation of the desired enema, thus placing this valuable method in the hands of those whose time and opportunity forbid resort to abattoirs for the natural fluid. Printed information furnished on application.

SEMPLE'S ATOMIZING INHALER.

The value of medicated vapor in the topical treatment of affections of the nose, mouth, larynx, bronchi and lungs, has long been recognized by the medical profession; but the various inhalers that have been heretofore placed before the profession have very imperfectly fulfilled the purpose for which they were designed, and physicians and patients have been discouraged by failures of apparatus to supply a spray which will admit of continuous inhalation without irritation of the membrane it is intended to medicate.

It is, therefore, with especial pleasure that we offer the profession Dr. Semple's apparatus, which is immeasurably superior to any now in use, and will, we believe, be largely employed in future, not only by specialists in diseases of the respiratory tract, but also by the general practitioner. This new invention is unquestionably the most perfect apparatus ever presented for the application of medicated vapor to the respiratory tract. It combines simplicity of construction, durability, and a fineness of spray unapproached by any atomizing inhaler hitherto introduced. It provides a constant supply of vapor in a reservoir, and secures more effectually than any other the topical effects of medicated vapor on the whole respiratory tract. In consumption, asthma, catarrh, bronchitis, hay fever, and indeed in all diseases in which it is desired to topically medicate the mucous membrane of the respiratory tract, this atomizing inhaler forms an indispensable addition to the armamentarium of the physician. For formulæ of inhalants suitable for use with this inhaler see page 211

SUGAR TEST FLASKS.

These flasks furnish a convenient and ready means of quantitatively determining the amount of grape sugar in diabetic urine. They are accompanied with explicit directions, and the process is simple. Each flask contains a sufficient quantity of copper solution to exactly reduce one-fifth of a grain of anhydrous glucose.

Unless hermetically sealed as in the flasks, a copper solution speedily deteriorates, and the physician, rather than be at the trouble of preparing his reagent for the occasion, often leaves his diagnosis in doubt, when, with the reagent at hand, he could easily arrive at certainty. Physicians find these flasks, therefore, a great convenience.

SYRINGES.

Chamberlin's Utero-vaginal Syringe.

This instrument shown in the cut, for which we are the manufacturer's sole agents, is constructed as follows: The vaginal tube is six inches in length; the surface is divided into ten longitudinal grooves one-eighth of an inch in depth, with an orifice at the termination of each at the base of the dome for a reverse stream. In the dome are orifices for direct and diverging streams. Being nearly an inch in diameter, it acts as a "repositor" in pro-

lapsus uteri. It presses the organ up to its normal position, at the same time distending the folds, by gently rotating the tube while the liquid is being injected. The grooves detach the tenacious mucus that adheres to the membrane, and the reverse streams passing down between the instrument and the vagina effectually removes it. As a consequence, two import-

ant results are obtained, viz: The unhealthy secretions are removed, and all injury therefrom obviated. The mucous membranes of the vagina and cervix uteri are effectually cleansed. If disease exist, medicine may now be directly and thoroughly applied to the affected part.

The Indispensable Cup Syringe.

This instrument has been constructed to meet the demand created for an improved method of irrigation in the treatment of uterine disorders, and is especially recommended, because:

It is a complete syringe for the *vagina* or *rectum*, as it is accompanied by an extra tube for rectal injections, and therefore no other syringe is necessary in the family

As a vaginal douche it can be used without rising from the bed.

Any quantity of water can be used without fatigue.

No assistant is needed.

It does not soil the bed or clothing

The syringe may be detached and the cups with the drainage tube forms a perfect urinal

THE NEW UREOMETER.

A simple, convenient, accurate and inexpensive apparatus for the determination of urea in the urine. The amount of urea excreted by the patient in a given time, furnishes the physician information of the greatest importance as bearing upon diagnosis, prognosis, and treatment. It is in general an index of the manner in which the various physiological functions of the organism are performed, and in particular, in certain kidney diseases, it enables the physician to foresee and guard against danger. No new principle is involved in our ureometer. The urea is decomposed as usual by the action of an alkaline solution of chlorine or bromine, and determined by the volume of gas (nitrogen) generated in the reaction. The original features in the apparatus are (a) the form of the receiver in which the gas is collected and measured, and (b) the graduation of this receiver in such a way that the results are read off directly in percentage of urea. The apparatus consists of:

1. A bottle provided with perforated rubber cork and delivery tube, in this the decomposition of the urea is effected.

2. A small test tube to contain the urine, graduated to hold 4 C. c., the quantity employed in each experiment.

3. A graduated jar for measuring the gas evolved. This jar is provided with an overflow tube, and a vent tube closed by a rubber cap, to secure accurate adjustment of the level of fluid in the jar at the commencement of the experiment.

Each ufeometer is accompanied by a descriptive circular containing full and explicit instructions; the estimation of urea by it consumes very little time, and the results for all practical purposes are as accurate as could be wished. Illustrated circular furnished upon application.

URINARY TEST PAPERS.

A happy suggestion was made by Dr. G. Oliver, of Harrogate, England, in 1883, that slips of paper might be impregnated with the various reagents employed in urinary tests, and that the physician could thus compress into the compass of a vest-pocket case all the reagents likely to be of service in making an examination of the urine. Such was introduced by us, and has met with unusual success. We would, however, call attention to an improvement upon these test papers which we have recently made: see Urinary Test Tablets next page.

The case contains; first, the indispensable litmus paper—a neutral shade, serving either for acidity or alkalinity of the secretion; papers impregnated with citric acid and with sodium carbonate, to be used in connection with the reagents for albumen and sugar, for the detection of glucose; papers impregnated with—1, indigo carmine and 2, picric acid, and the indigo paper. As tests for albumen the series includes four of the recently introduced reagents, whose value experience has already sufficiently established. These are: 1. Picric acid. 2. Potassio-mercuric iodide. 3. Potassium ferrocyanide, and 4. Sodium tungstate. All of these are to be used in connection with citric acid.

Full directions accompany each package. Our bound pamphlet containing Dr. Geo. Oliver's original article on "Bedside Urinary Tests," an article by Chas. W. Purdy, M. D., of Chicago, Ill., on "The Comparative Value of the Newer Tests for Albumen in the Urine," and one by the same author upon "Urinary Test Papers," will be forwarded, postpaid, to any address upon application.

Physician's Pocket Reagent Case.

Soon after the introduction of our Urinary Test Papers, *q. v.*, it became apparent that a small pocket case which should contain all the apparatus necessary for urine

analysis, at the bedside of the patient, was in large demand, and our Physician's Pocket Reagent Case was therefore placed upon the market. Our greatest expectations have been exceeded.

This case contains a complete set of chemical apparatus and reagents for urine analysis, qualitative and quantitative, to-wit: In addition to the series of urinary test papers issued by us, a comprehensive book of instructions for the use of Dr. Oliver's bedside tests, and the following apparatus:

1. Two test tubes, one of which is graduated.

2. A minim pipette.

3. A set of six specific gravity beads, corresponding with specific gravities respectively of 1.005, 1.010, 1.015, 1.020, 1.025, 1.030. These will be found very convenient to use.

A more compact arrangement it would be difficult to devise, and the physician cannot well forego the convenience of such a vest-pocket chemical laboratory.

URINARY TEST TABLETS.

The urinary tests in paper form as devised by Dr. G. Oliver, and introduced to the medical profession of this country by us, have had one serious objection urged against them; this defect being especially apparent in those tests depending upon the production of a precipitate.

If one of these reagent papers be boiled with a quantity of water, the agitation incident to ebullition is sufficient to disintegrate the paper, and the fibres floating through the liquid are liable to be mistaken for a precipitate, and consequently lead to a grave diagnostic error.

In the substitution, for these papers, of tablets consisting of a soluble inert base combined with the reagent, we have not only obviated this defect, but have in several instances been enabled to simplify the tests, at the same time enhancing the permanency thereof.

The following urinary tests are furnished in tablet form:

Tests for Sugar.—Three tests are furnished in this series:

1. INDIGO-AND-SODIUM-CARBONATE.
2. PICRIC ACID.
3. BISMUTH SUB-NITRATE (Boettcher's Sugar Test).

These reagents are all used in connection with tablets of SODIUM CARBONATE.

The diagnosis of glycosuria may be made with certainty by means of these test tablets, and an approximate estimation made of the quantity of sugar present.

The presence of albumen and of uric acid in the urine does not interfere with any of these tests, which are capable of detecting the smallest quantities of sugar that can be considered pathological.

Tests for Albumen.—Four different reagents for albumen are furnished in this series of test tablets.

1. POTASSIO MERCURIC IODIDE.
2. SODIUM TUNGSTATE.
3. POTASSIUM FERROCYANIDE.
4. PICRIC ACID.

These reagents are all used in connection with tablets of CITRIC ACID.

WILSON'S EUREKA ABDOMINAL SUPPORTER.

The best surgeons unanimously assert that the most appropriate and satisfactory treatment for the milder forms of prolapsus uteri is the continuous use of a properly adjusted supporter. It is claimed that when the procidentia is excessive, it is advisable to employ a stem pessary in addition, with a perforated cup for the reception of cervix uteri. By this combined pressure the organ is enabled to resume its natural position.

We are assured that in Dr. Wilson's Supporter are combined all the features essential to a satisfactory instrument, and beg to enumerate below the reasons which form the basis of our opinion.

The supporter (without the pessary) is perfect and complete in itself, consisting of back and abdominal pads, connected by elastic bands and perineal or thigh straps. It is thus admirably adapted to the less severe grades of displacements.

The stem of the pessary is made of German silver, plated with silver, and can be easily bent to fit any person. The cup, composed of celluloid, is so shaped as to carry the uterus with the greatest possible comfort, the support being natural and without pressure upon the os uteri, or urethra. Irritation of the parts, which so frequently results from the use of other instruments, is thus avoided.

The position of the pessary on the wire may be readily changed with little effort. In this manner it may be adjusted to meet the requirements of the varying degrees of retroversion and anteversion, a desideratum which the general practitioner will fully appreciate. Illustrated circular furnished on application.

RARE ALKALOIDS, GLUCOSIDES, CHEMICALS, ETC.,

REPRESENTING THE ACTIVE PRINCIPLES OF MANY IMPORTANT DRUGS, TOGETHER WITH A FEW CHEMICALS AND MEDICINAL AGENTS FROM OTHER SOURCES, WHICH, EXCEPT IN SOME OF THE LARGE CITIES, ARE FOUND DIFFICULT TO PROCURE.

Many of the articles are referred to pages where they have already appeared.

Acetanilid.—A prompt and powerful antipyretic; known also as Antifebrin. Dose, 8 to 12 grains (0.52 to 0.78 Gm.).

Acid Hippuric.—Found as a potassium and sodium salt in human urine, more especially after the administration of benzoic acid. For therapeutic properties see calcium hippurate.

Acid Metaphosphoric.—Glacial phosphoric acid. A convenient and portable test for albumen in the urine. A fragment as large as a small pea is dropped into the test-tube containing a fluidrachm of the urine to be tested and allowed to dissolve. If albumen is present, the acid, as it dissolves, will coagulate it, forming a white zone, more or less opaque, as in

RARE ALKALOIDS, Etc.—Continued.

the ordinary test with nitric acid. The only advantage claimed for this reagent over nitric acid is that it may be easily carried in the medicine case, and has no corrosive properties.

Acid Osmic.—Perosmic acid. Sparingly soluble in water, and its solution easily reduced by contact with organic matter. It is also volatile, and its vapor is very poisonous. Solutions should be freshly prepared when required. Its most important application thus far has been in the treatment of obstinate sciaticas and facial neuralgias. Five minims of a solution 1:100 are injected subcutaneously, and the operation repeated at intervals until permanent relief is obtained. Used with some success in the treatment of epilepsy. Dose, 1-40 to 1-10 grain (0.0016 to 0.006 Gm.).

Acid Pipitzahoinic.—This acid, which is recognized by the Mexican pharmacopœia, is prepared from the root of *Trixis fruticosa*, and is recognized as an active drastic. Dose, 3 to 5 grains (0.2 to 0.32 Gm.).

Acid Salicylic, page 193.

Acid Sclerotic, solution, page 74.

Acid Trichloroacetic.—A powerful antiseptic, without odor, and preferred to carbolic acid. As an application to gangrenous wounds and ill conditioned ulcers, a solution of 5 to 10 grains to the pint of water may be used. Internally it has been used in septic diarrhœa, gastric catarrh, and as a prophylactic against Asiatic cholera. Also employed as a reagent for the detection of albumen in the urine. A crystal of the acid is dropped into a test tube containing a fluidrachm of the urine and allowed to dissolve. If albumen is present a white zone is formed above the acid, as in the ordinary nitric acid test. The reaction is extremely delicate. The reagent, moreover, unlike nitric acid, is non-corrosive, and may be carried in the medicine case. Dose, 2 to 5 grains (0.13 to 0.32 Gm.) in dilute solution.

Aconitine, page 2.

Adonidin, page 4.

Agaricin, page 189.

Alkanin, page 5.

Aloin, page 6.

Aluminium acetate, solution, page 241.

Ammonium glycyrrhizate, page 113.

Ammonium picrate, or Ammonium Carbazotate.—Once highly recommended as a substitute for quinine, both as an antiperiodic and bactericide. Clinical experience, however, shows that it has little effect in antagonizing malarial poison, although as a remedy for whooping-cough, and other diseases depending on the presence of micrococci or bacteria, it has realized the hopes of those who have employed it on theoretical grounds. Dose, 1-8 to 1 grain (0.008 to 0.06 Gm.) and upwards.

Amyl nitrite.—An ethereal liquid having a peculiar fruit-like odor and a pungent aromatic taste. Usually given

by inhalation, but can be administered subcutaneously. It has proved very successful in angina pectoris and in asthma. Has also been used with advantage in a certain class of epileptic fits. See page 238. Dose, 2 to 5 minims (0.12 to 0.3 C.c.).

Amyl valerianate.—Amyl valerianic ether. This ether, in a dilute solution, has an agreeable apple-like odor, and is indeed sold under the name of apple oil. It is stimulant, antispasmodic anodyne and hypnotic. It has been found especially useful in nephritic and hepatic colic, in gastralgia, dyspnoea, dysmenorrhœa, angina pectoris, and in short, wherever there is pain associated with spasm. As a rapid diffusible stimulant, it is almost without a rival. Dose, 3 to 8 minims (0.2 to 0.5 C.c.).

Antipyrin.—An antipyretic, but not antiperiodic. A powerful and certain means of lowering the temperature in almost all febrile diseases. Dose, 15 grains (1 Gm.).

Apiol, page 136.

Apomorphine muriate, p. 130.

Arbutin, page 184.

Aspidospermine, page 148.

Atropine, pure, page 19.

Atropine sulphate, page 19.

Berberine hydrochlor., p. 88.

Berberine phosphate, p. 88.

Berberine sulph., acid, p. 201.

Berberine sulph., neutral, page 88.

Bismuth subiodide.—An efficient substitute for iodoform.

Boroglyceride.—Nearly colorless, transparent, and soluble in alcohol and water. Owing to its hygroscopic nature, it should be excluded from the air. Recommended as a valuable antiseptic.

Brucine, page 128.

Cadmium sulphate.—Astringent, irritant; resembling zinc sulphate, but much more powerful. Used chiefly as a local application in conjunctivitis, ulcers and opacities of the cornea, otorrhœa and gleet, in solution containing 1-2 to 4 grains to the fluidounce (1 to 8 grammes to the litre).

Caffeine citrate, page 58.

Calcium hippurate.—Dr. Poulet claims for this salt extraordinary virtues in the treatment of the urinary passages, and of the stomach and intestines, also in various disorders of nutrition. He bases his use of the salt on the hypothesis that hippuric acid rather than hydrochloric is the principal acid of the stomach. Said to produce remarkable effects in all affections associated with excess of uric acid. Dose, 5 to 10 grains (0.32 to 0.65 Gm.).

Cannabin tannate, page 100.

Cannabinon, page 100.

Cerium nitrate.—Recommended by Dr. Simpson as a nerve tonic, in chronic intestinal eruption, and chronic vomiting generally, as well as in that of pregnancy. Dose, 1 grain (0.065 Gm.).

Chinoline.—(Quinoline.) Derivative from coal-tar. A liquid not miscible with aqueous but soluble in alcoholic fluids. An antipyretic and antiseptic.

RARE ALKALOIDS, Etc.—Continued.

Chinoline salicylate.—The antiperiodic and febrifuge properties of the base in this salt are believed to be intensified by combination with salicylic acid, a powerful antizymotic. Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

Chinoline tartrate.—Freely soluble in water. A powerful antiseptic. Acts as an antipyretic and antiperiodic. It is said to relieve whooping-cough. Given in the same dose as quinine.

Cholesterin.—A white crystalline fatty substance, a constituent of bile, yolk of eggs, cod-liver oil, brain and nerve tissue, etc. Believed to be purely an excrementitious product, having no application in medicine.

Cocaine and salts of, page 56.

Codeine, page 131.

Coniine and salts of, page 62.

Colchicine, page 59.

Colocythin, page 60.

Convallamarin, page 114.

Cotin, page 64.

Curare, page 69.

Daturine and salt of, page 173.

Digitalin, page 80.

Duboisine and salt of, page 72.

Elaterin, page 73.

Emetine, page 101.

Eserine (Physostigmine), page 38.

Ethyl bromide.—The most pleasant, safe and prompt anæsthetic. For short operations, it certainly stands without a rival. Although in longer operations it cannot be substituted for ether or chloroform, it is a valuable adjunct to these. It may be employed as a preliminary to the administration of ether, or else used as one of the constituents of an anæsthetic mixture. Thus a favorite mixture in labor cases as well as in surgical operations is: bromide of ethyl, 1 part (by weight); chloroform, 1 part; alcohol, two parts.

Gelsemine and salts of, page 84.

Homatropine hydrobromate, page 10.

Hydrargyrum tannicum oxydulatum (Mercurous tannate).—One of the most efficient mercurials for the treatment of syphilitic affections. It is insoluble in water and in acids, but under the action of the alkaline secretions of the intestines, the mercury is reduced to the metallic state, in a condition of extremely minute subdivision, and it is thus very gradually and continuously dissolved and absorbed, producing a sustained effect on the system without deranging the functions of the stomach and bowels. Dose, 1 to 2 grains (0.065 to 0.13 Gm.).

Hydrastine and salts of, page 88.

Hydrochinon.—Hydroquinone. A coal-tar product, moderately soluble in cold water, readily soluble in hot water, alcohol, ether and acid solutions. It is antiseptic and antipyretic, resembling in action antipyrin. Without disagreeable secondary effects. Dose, 10 to 20 grains (0.65 to 1.3 Gm.).

Hyoscyne hydrobrom., p. 94.

Hyoscyamine, page 94.

Hypnone.—Phenyl-methyl-acetone, or acetophenone. Dujardin-Beaumez and other experimenters have found this to be a powerful, pure hypnotic that will act promptly in insomnia depending upon cerebral hyperexcitation, but wholly useless in insomnia caused by pain. Well borne by digestive tract. Disagreeable taste; best administered in capsules. Dose, 1 to 4 minims (0.06 to 0.25 C.c.).

Ichthyol.—A tar-like substance. It forms an emulsion with water, and is miscible with fats and oils. It contains ten per cent. of sulphur, and its applications in the treatment of skin diseases doubtless depend in a large degree on this circumstance. Used in the form of ointment, to sprains, painful tumefactions of joints, etc., either traumatic or of rheumatic origin; in muscular rheumatism, ichthyosis, psoriasis, etc.

Iodol.—Tetra-iodolpyrol. Contains 80 per cent. of iodine. In chemical constitution resembles and proposed as a substitute for iodoform in surgical practice, having the great advantage of being free from odor. Nearly insoluble in water; soluble in alcohol, ether and oils. It may be employed in powder or in solution in a mixture of: alcohol, 1 part; glycerin, 2 parts; or in ointments.

Kairine.—Oxyethylquinoline-hydrate hydrochloride. Artificial alkaloid derivative of coal-tar. Its action closely resembles that of quinine and it is given in similar doses especially as an antiperiodic. Said to be an excellent febrifuge.

Kava Kava resins, page 108.

Kavalin, page 108.

Lanolin, page 111.

Menthol, page 122.

Muscarine sulphate, page 80.

Napelline, page 2.

Naphthalin, page 178.

Naphthol, page 178.

Nitroglycerin.—One per cent. solution. Used with benefit in many forms of neuralgia, cerebral anemia, weak heart of the aged, chronic intestinal nephritis, etc. Said to give much more lasting benefit in angina pectoris than amyl nitrite. Best given in pill form. Dose, of 1% solution, 1-2 to 10 minims (0.03 to 0.6 C.c.).

Papain, page 43.

Papayotin, page 43.

Paracotin, page 65.

Paraldehyd.—A derivative of common ethylic alcohol. It has the same composition as aldehyde, but is much less volatile, and its properties are quite different. It is a hypnotic, resembling in its effects chloral hydrate. Dose, 45 grains (3 grammes). It may be mixed with any aromatic tincture.

Pelletierine salts, page 144.

Physostigmine and salts of, p. 38.

Picrotoxin, page 78.

Pilocarpine salts, page 104.

Piscidin, page 105.

Podophyllotoxin, page 118.

Pyridine.—Volatile alkaloid produced in destructive distillation of organic matter. It is believed to be the chief

RARE ALKALOIDS Etc.—Continued.

active constituent of tobacco smoke. It has been lately recommended by Prof. Germain Séé as a remedy for asthma, relieving the spasm more promptly and effectually than any other drug. Given by inhalation:—A half drachm or more of the fluid is placed in a saucer and the vapor caused to fill the air of a chamber in which the patient remains 15 to 30 minutes.

Quinine and Urea muriate, *Quinia bimuriatica carbamidata*, page 52.

Resorcin.—A phenol, with properties analogous to those of ordinary carbolic acid, especially as an antiseptic adapted to the wants of the surgeon. It is odorless, solid at ordinary temperature, and dissolves readily in water. Effectual in the treatment of epithelioma and eczema. It reduces pulse and temperature in febrile conditions and is used internally with asserted benefit in septic poisoning in whooping cough. Dose, 5 to 10 grains (0.32 to 0.65 Gm.).

Sanguinarine salts, page 30.

Scillitoxin, page 169.

Sodium choleate.—Its properties and uses are those of ordinary inspissated ox-gall, but dose is somewhat smaller. A mild laxative, especially useful in flatulent dyspepsia. Dose, 1 to 5 grains (0.06 to 0.32 Gm.).

Sodium nitrite.—Recommended in epilepsy in doses of about two grains, or just short of producing physiological effects. Cases in which bromide does good are not apt to be benefited by the nitrite, and *vice versa*; when the bromides begin to loose effect or cause bromism, it is useful as a substitute.

Sodium sulphocarbolate.—Soluble in 5 parts of water and in 132 parts of alcohol. Used with more or less success in fermentative dyspepsia and in diphtheria of the anginose form. Dose, 10 to 30 grains 0.6 to 2 Gm.), applied topically in a saturated aqueous solution.

Sodium tungstate.—Employed as a reagent for the detection of albumen in the urine. One part of the tungstate and two of citric acid are dissolved in four parts of water, and a few drops of the solution are added to the urine to be tested. See also Urinary Test Papers, page 245; and Urinary Test Tablets, page 245.

Sparteine sulphate, page 33.

Terebene, page 183.

Terpin hydrate, page 183.

Thalline.—A synthetic alkaloid, in properties intermediate between kairine and antipyrin. Its effects are not so lasting as those of antipyrin, but its action is more powerful. Dose, 4 to 10 grains (0.25 to 0.65 Gm.).

Thalline sulphate and tartrate.—Salts of the above with similar medicinal properties but more soluble.

Thymol, page 180.

Uranium nitrate.—This salt has been employed with alleged success in the treatment of diabetes mellitus. Dose, 1-6 to 1-2 grain (0.01 to 0.03 Gm.).

Urethane.—Ethyl carbamate. Without odor and nearly tasteless, freely soluble in water. A valuable hypnotic. It produces sleep of a seemingly perfect natural character and does not interfere, like opiates, with the secretions, or produce any unpleasant secondary effects. Dose, 10 to 20 grains (0.65 to 1.30 Gm.).

Xylol, page 178.

TABLES OF WEIGHTS AND MEASURES.

EQUIVALENTS OF WEIGHT UNITS.

1 gr. =	Gm.....	0648	1 oz. = av.	Gm.....	28.349
	scruple.....	.05		drachm.....	7.292
	drachm.....	.0167		oz. tr.....	.911
	oz. av.....	.00229		lb. tr.....	.076
	oz. tr.....	.00208		lb. av.....	.0625
	lb. tr.....	.000174		m.....	460.905
	lb. av.....	.000143		C. C.....	28.375
	m.....	1.0535		cu. in.....	1.783
	C. C.....	.0649		fl. oz.....	.96
	fl. dr.....	.0176		pint.....	.06
	cu. in.....	.00396		Liter.....	.0284
fl. oz.....	.00219				
pint.....	.000137				
.					
1 Gm. =	gr.....	15.432	1 oz. = tr.	Gm.....	31.103
	scruple.....	.772		oz. av.....	1.0971
	drachm.....	.257		lb. tr.....	.0833
	oz. av.....	.0353		lb. av.....	.0686
	oz. tr.....	.0322		m.....	505.679
	lb. tr.....	.00268		C. C.....	31.132
	lb. av.....	.0022		cu. in.....	1.901
	m.....	16.258		fl. oz.....	1.0535
	C. C.....	1.001		pint.....	.0658
	fl. dr.....	.271		Liter.....	.0311
	cu. in.....	.061			
fl. oz.....	.0339				
pint.....	.00212				
Liter.....	.001				
.					
1 sc. =	Gm.....	1.296	1 lb. = tr.	Gm.....	373.242
	drachm.....	.333		oz. av.....	13.166
	oz. av.....	.0457		lb. av.....	.823
	oz. tr.....	.0417		C. C.....	373.578
	lb. tr.....	.00347		cu. in.....	22.815
	lb. av.....	.00286		fl. oz.....	12.642
	m.....	21.07		pint.....	.79
	C. C.....	1.297		Liter.....	.374
	fl. dr.....	.351		gal.....	.0988
	cu. in.....	.0792			
	fl. oz.....	.0439			
pint.....	.00274				
.					
1 dr. =	Gm.....	3.888	1 lb. = av.	Gm.....	453.593
	oz. av.....	.137		oz. tr.....	14.583
	oz. tr.....	.125		lb. tr.....	1.215
	lb. tr.....	.0104		C. C.....	454.001
	lb. av.....	.00857		cu. in.....	27.726
	m.....	63.21		fl. oz.....	15.363
	C. C.....	3.891		pint.....	.96
	fl. dr.....	1.0535		Liter.....	.454
	cu. in.....	.238		gal.....	.12
	fl. oz.....	.132			
	pint.....	.00823			

EQUIVALENTS OF MEASURE UNITS.

1 m. =	gr	.949
	Gm	.0615
	scruple	.0475
	drachm	.0158
	oz. av.	.00217
	oz. tr.	.00198
	lb. tr.	.000165
	lb. av.	.000136
	C. C.	.0616
	fl. dr.	.0167
	cu. in.	.00376
	fl. oz.	.00208
	pint	.00013

1 C. C. =	gr	15.418
	Gm	.999
	scruple	.771
	drachm	.257
	oz. av.	.0352
	oz. tr.	.0321
	lb. tr.	.00268
	lb. av.	.0022
	m	16.243
	fl. dr.	.271
	cu. in.	.061
	fl. oz.	.0338
	pint	.00212

1 fl. dr. =	gr	56.953
	Gm	3.691
	scruple	2.848
	drachm	.949
	oz. av.	.13
	oz. tr.	.119
	lb. tr.	.00989
	lb. av.	.00814
	C. C.	3.694
	cu. in.	.226
	fl. oz.	.125
	pint	.00781
	Liter	.00369

1 c. in. =	gr	252.468
	Gm	16.36
	scruple	12.623
	drachm	4.208
	oz. av.	.577
	oz. tr.	.526
	lb. tr.	.0438
	lb. av.	.0361
	m	265.974
	C. C.	16.374
	fl. dr.	4.433
	fl. oz.	.554
	pint	.0346
	Liter	.0164
	gal	.00433

1 fl. oz. =	gr	455.625
	Gm	29.524
	oz. av.	1.0414
	oz. tr.	.949
	lb. tr.	.0791
	lb. av.	.0651
	C. C.	29.551
	cu. in.	1.805
	pint	.0625
	Liter	.0295

1 pt. =	gr	7290.
	Gm	472.384
	oz. av.	16.663
	oz. tr.	15.188
	lb. tr.	1.266
	lb. av.	1.0414
	C. C.	472.81
	cu. in.	28.875
	Liter	.473

1 L. =	gr	15418.44
	Gm	999.1
	scruple	770.922
	drachm	256.974
	oz. av.	35.242
	oz. tr.	32.122
	lb. tr.	2.677
	lb. av.	2.303
	m	16243.295
	C. C.	1000.
	fl. dr.	270.722
	cu. in.	61.027
	fl. oz.	33.84
	pint	2.115
	gal	.264

1 gal. =	lb. tr.	10.125
	lb. av.	8.331
	cu. in.	231.
	Liter	3.782

EQUIVALENTS OF IMPERIAL MEASURE UNITS.

IMPERIAL MEASURE, B. P.	WINE MEASURE, U. S. P.	APOTHECARIES' WEIGHT.	AVOIRDUPOIS WEIGHT.	METRIC WEIGHT AND MEASURE	CUBIC INCH.
1 minim	m. .96	gr. .912	gr. .912	Gm. C. C. { .0591 .0592 }	.00361
1 fl. dr.	{ m. fl. dr. .96	gr. dr. 54.69 .912	gr. oz. 54.69 .125	Gm. C. C. { 3.544 3.547 }	.217
1 fl. oz.	{ fl. dr. fl. oz. 7.681 .96	dr. oz. 7.292 .912	oz. 1.	{ Gm. C. C. 28.351 28.376 }	1.733
1 pint	{ fl. oz. pint. 19.202 1.2	oz. lb. 18.64 1.519	oz. lb. 20. 1.25	{ Gm. C C 567.012 567.522 Liter .568 }	34.659
1 gal.	{ pint. gal. 9.601 1.2	lb. 12.154	lb. 10.	{ Gm. C C. 4536.095 4540.177 Liter. 4.54 }	277.274

SPECIAL TABLES OF APPROXIMATE EQUIVALENTS.

Grains.	Grammes.	oz. troy.	Grammes.	C. C.	m.	fl. dr.
1-100	.00065	1	31.1	1	16	.27
1-64	.00101	2	62.2	2	32	.54
1-50	.00130	3	93.3	3	49	.81
1-40	.00162	4	124.4	4	65	1.28
1-36	.00180	5	155.5	5	81	1.35
1-30	.00216	6	186.6	6	97	1.62
1-24	.00270	7	217.7	7	114	1.90
1-20	.00324	8	248.8	8	130	2.17
1-18	.00360	9	279.9	9	146	2.44
1-16	.00405					
1-15	.00432					
1-12	.00540					
1-10	.00648					
1-8	.00810					
1-6	.01080					
1-5	.01296					
1-4	.0162					
1-3	.0216					
1-2	.0324					
3-4	.0486					
1	.065					
2	.130					
3	.195					
4	.260					
5	.324					
6	.389					
7	.454					
8	.518					
9	.583					
10	.648					
11	.713					
12	.778					
13	.842					
14	.901					
15	.972					
16	1.037					
17	1.102					
18	1.166					
19	1.231					
20	1.296					
30	1.944					
40	2.592					
50	3.240					
Drachms.	Grammes.	fl. oz.	C. C.	oz. tr.	lb. av.	
1	3.89	1	29.6	1-8	.00781	
2	7.78	2	59.1	1-4	.0156	
3	11.66	3	88.7	1-2	.0313	
4	15.55	4	118.2	3-4	.0469	
5	19.44	5	147.8	1	.063	
6	23.33	6	177.3	2	.125	
7	27.22	7	206.9	3	.188	
		8	236.4	4	.25	
		9	266.	5	.313	
				6	.375	
				7	.438	
				8	.5	
				9	.563	
				10	.625	
				11	.688	
				12	.75	
				13	.813	
				14	.875	
				15	.938	
oz. troy.	Grammes.	Grammes.	Grains.	oz. av.	lb. av.	
.1	48	1	15	1-8	.00857	
.2	96	2	31	1-4	.0171	
.3	144	3	46	1-2	.0343	
.4	192	4	62	3-4	.0514	
.5	240	5	77	1	.0686	
.6	288	6	93	2	.137	
.7	336	7	108	3	.206	
.8	384	8	123	4	.274	
.9	432	9	139	5	.343	
				6	.411	
				7	.481	
				8	.549	
				9	.617	
				10	.686	
				11	.754	
				12	.823	
				13	.891	
				14	.96	
				15	1.0286	
				16	1.0971	

WEIGHT AND MEASURE EQUIVALENTS.

These Tables, for the use of physicians, chemists, pharmacists and others, are designed to economize time and labor, and ensure accuracy in calculations relating to Weight and Measure.

A UNIT of each denomination of apothecaries' and avoirdupois weights, wine and imperial measures, as also the metric gram, cubic centimeter and liter, and the cubic inch are represented by equivalent quantities being given in every important denomination of each system; beginning with the lowest denomination, and following by the next larger successively. This plan and arrangement not only render the "tables" convenient and susceptible of all the varied uses that the quantities are capable of, but simplifies and shortens nearly all calculations to mere multiplication. Multiples of several of the more useful denominations are also tabulated.

When making use of the equivalents, for most practical purposes, one or more of of the figures beyond the decimal point may be dropped, - according to the degree of accuracy required - or, if any figure of the second or third decimal place is five or more, the preceding figure can be increased by one. The same method can apply to the results of calculations.

To Convert Troy Ounces into Avoirdupois Ounces.

It is required for example to convert 23 tr. ozs. into av. ozs. This may be done by ascertaining the tr. oz. equivalent from the table, that is, 1.0971 and this $23 \times 1.0971 = 25.233$. This differs but very slightly from 23×1.1 (which is the same as adding 1-10) $= 25.3$ so that for all practical purposes the addition of 1-10 will transform tr. ozs. into av. ozs.

To Convert Avoirdupois Ounces into Troy Ounces.

Similarly to the foregoing, instead of consulting the table of equivalents, it is useful at times to remember that deducting 1-11 from a number representing av. ozs. will give the equivalent weight in tr. ozs. Thus, taking the result of the previous example; 25.3 av. ozs., how many tr. ozs.? $25.3 - 1-11$ or $2.3 = 23$ tr. ozs.

To Estimate Cost.

Find the commercial equivalent of a unit of the given denomination; multiply by the required quantity, then by the cost; thus, a formula directs 30 Gm. of a substance valued at 80 cents an av. oz., the equivalent of 1 Gm. in av. oz., is .0353 and this $30 \times .0353 \times 80 = .847$, practically 85 cents cost. When formulæ express quantities in the troyounce denomination, and values are quoted for the av. lb., use the "special table" of troyounce in av. lb. equivalents; thus, 7 troyounces = .48 av. lb. $\times$ 1 lb. value. If a formula directs fluid measure, and the price is quoted for the pint, pursue the same general method; using the "special table" of fl. oz. in pint equivalents, when convenient.

To enlarge formulæ and prescriptions, and change their quantities to other equivalents follow the foregoing general method. Figures beyond the decimal point can be reduced to grains by means of the appropriate special tables.

To Ascertain the Weight of a Given Measure of any Liquid or Solid.

Find the water-weight of the measure in any desired denomination, which multiply by the specific gravity of the substance; thus, 1 pint of water $= 1.0414$ av. lb. $\times .85$ the supposed s. g. $= .885$ av. lb.; or if the equivalent in av. oz. is multiplied, the weight of 1 pint is practically 14 av. oz.

To Ascertain the Measure of a Given Weight of any Liquid or Solid.

Divide the weight by the specific gravity and multiply by the measure of the water-weight "unit" in any desired denomination; thus, 1 av. lb. $\div .85$ the supposed s. g. $\times .96$ pint (the measure of 1 av. lb. of water) $= 1.13$ pints; or if the equivalent in fl. oz. is employed, the measure is practically 18 fl. oz. Or, instead of dividing the weight by the s. g., multiply it by the specific volume if this is known: See below.

APPROXIMATE MEASURES.

These are subject to considerable elasticity of interpretation owing to the varying sizes of the domestic utensils in use, but the following are understood to be the quantities always intended when the popular names are employed:

A teaspoonful	1 fluidrachm.
A dessertspoonful	2 fluidrachms.
A tablespoonful	$\frac{1}{2}$ fluidounce.
A wineglassful	2 fluidounces.
A teacupful	4 fluidounces.
A tumblerful	8 fluidounces.

SPECIFIC VOLUME.

Specific volume is the ratio of the bulk of any liquid, contrasted with the bulk of an equal weight of water, which is taken as the standard. It is found by dividing unity by the specific gravity of the liquid whose sp. vol. is sought. Thus the sp. vol. of alcohol at 25° C. (77° F.) is 1.231, found thus $\frac{1}{.812}$ (sp. gr.). Since the U. S. P. directs all quantities by weight the utility of sp. vol. in calculations of the equivalent fluid measure is at once apparent. Thus, for example, 100 parts by weight of alcohol are directed to be taken for a given formula; by the weight unit equivalent (see table) 1 av. oz. = .96 fl. oz., therefore 100 av. ozs. = 96 fl. ozs. This is the fluid measure equivalent if the liquid happened to be water, but alcohol being so much lighter than water in the proportion of sp. gr. .812 : 1 a greater bulk of it in the proportion of the sp. vols. 1.231 : 1 will be equivalent to the given weight. The quantity of alcohol necessary for the formula is therefore $96 \times 1.231 = 118.176$ fl. ozs.

TABLE OF THERMOMETRIC EQUIVALENTS.
Centigrade and Fahrenheit Scales.

Centigrade	Fahrenheit	Centigrade	Fahrenheit	Centigrade	Fahrenheit	Centigrade	Fahrenheit	Centigrade	Fahrenheit
100	212	74	165.2	48	118.4	34	93.2	8	46.4
99	210.2	73	163.4	47	116.6	33	91.4	7	44.6
98	208.4	72	161.6	46	114.8	32	89.6	6	42.8
97	206.6	71	159.8	45	113	31	87.8	5	41
96	204.8	70	158	44	111.2	30	86	4	39.2
95	203.	69	156.2	43	109.4	29	84.2	3	37.4
94	201.2	68	154.4	42	107.6	28	82.4	2	35.6
93	199.4	67	152.6	41.66	107	27	80.6	1	33.8
92	197.6	66	150.8	41.11	106	26	78.8	0	32
91	195.8	65	149	41	105.8	25	77	-1	30.2
90	194	64	147.2	40.55	105	24	75.2	-2	28.4
89	192.2	63	145.4	40	104	23	73.4	-3	26.6
88	190.4	62	143.6	39.46	103	22	71.6	-4	24.8
87	188.6	61	141.8	39	102.2	21	69.8	-5	23
86	186.8	60	140	38.88	102	20	68	-6	21.2
85	185.	59	138.2	38.33	101	19	66.2	-7	19.4
84	183.2	58	136.4	38	100.4	18	64.4	-8	17.6
83	181.4	57	134.6	37.77	100	17	62.6	-9	15.8
82	179.6	56	132.8	37.50	99.5	16	60.8	-10	14
81	177.8	55	131	37.22	99	15	59	-15	5
80	176.	54	129.2	37	98.6	14	57.2	-17.78	0
79	174.2	53	127.4	36.66	98	13	55.4	-20	-4
78	172.4	52	125.6	36.11	97	12	53.6	-25	-13
77	170.6	51	123.8	36	96.8	11	51.8	-30	-22
76	168.8	50	122	35.55	96	10	50	-35	-31
75	167	49	120.2	35	95	9	48.2	-40	-40

To reduce Centigrade degrees to those of Fahrenheit:

Multiply by 9, divide by 5, and add 32.

To reduce Fahrenheit's degrees to those of the Centigrade scale:

Subtract 32, multiply by 5, and divide by 9.

POSOLOGY.

FROM THE PHYSICIAN'S PERFECT CALL-BOOK AND RECORD—GEO. S. DAVIS, PUBLISHER—
BY PERMISSION.

AXIOMS FOR REMEMBRANCE.

1. **Affinity of botanical characters** is no evidence of like therapeutic action; the same also holds good of *chemical composition*. Neither are *sensible properties* a test of physiological value.

2. **Like anatomical characteristics** are far from being evidence of like physiological functions. The action of medicines on lower animals, instead of being *guideposts*, are merely *landmarks*.

3. **Polypharmacy** is not medical science. The fewer ingredients in a prescription, and the more simple it can be made, the better.

4. **The action of medicines** is modified by *sex, age, temperament and diathesis, idiosyncrasies, habits of life, race, and mental impressions*.

5. **Women**, as possessed of more delicate organization than men, rarely require the same doses. Avoid aloes during *pregnancy*; sulphuric acid during *lactation*; and mercury in *anæmic chlorosis*.

6. **Old age and childhood** do not bear doses proportionately to manhood and youth. *Excepting under urgent circumstances, an opiate should never be administered to an infant.* Blisters and leeches make more decided impressions on children than adults,—all things being equal.—and the former are especially apt to induce ulceration and gangrene in the young. *Per contra*, mercury is better borne proportionately in childhood (and old age) than in middle life; it is all but impossible to salivate a child under ten years of age. The following is the scale of doses employed at Guy's Hospital, London, *one* being the unit or adult dose.*

AGE.	DOSE.	AGE	DOSE.
1 month.....	1-20	7 and 8 years.....	1-2
3 months.....	1-15	10 to 12 years.....	2-3
6 months.....	1-10	13 to 15 years.....	3-4
9 months.....	1-9	18 to 20 years.....	5-6
1 year.....	1-7	21 to 45 years.....	1
2 years.....	1-6	50 years.....	5-6
3 years.....	1-5	60 to 70 years.....	3-4
4 years.....	1-4	80 to 90 years.....	2-3
5 and 6 years.....	1-3	100 years.....	1-2

7. **Sanguine and sanguino-nervous temperaments** bear *antiphlogistics*, and *loss of blood*, readily; the reverse is true as to *stimulants*. *Antispasmodics* are more strikingly beneficial to *nervous temperaments*.

8. **Always inquire as to idiosyncrasies**, especially toward mercury and opium.

9. **Denizens of cities**, especially of overcrowded quarters and tenement-houses, cannot bear doses to which those of rural and out-door life are accustomed. (*Mark this as to children especially.*) The same active treatment necessary to relieve acute disease in the latter will probably tend to *fatality among the former*.

10. **Habit inculcates tolerance** even of poisons. Mustapha Shatoor was accustomed to take three drachms of opium at a dose; Suleyman Yenen a third as much corrosive sublimate during every twenty-four hours; and certain of the inhabitants of Northern India are accustomed to consume an entire *nux vomica* (about twenty grains), every twelve hours.—(*Bengal Dispensatory.*)

11. **Blondes require a larger dose than brunettes**. Indians, Negroes, Malays, Hindoos, and the dark and yellow races are for the most part amenable to *half Anglo-Saxon doses*.

12. **Passions and affections of the mind** markedly influence the action of remedies, especially narcotics. Faith in the *prescription*, or the *prescriber*, may remove *mountains of difficulties*, since faith is the foundation of the miraculous.

13. **Judicious combinations of medicines** are often more effective than remedies employed singly; but combinations are to be avoided except when great gain is to be had. *Digitalis* frequently fails to act as a diuretic until combined with squills or carbonate of ammonia; the *resinoids* are rendered more efficiently purgative by the action of mercury; and diaphoresis is more certainly induced by a combination of opium and ipecac than by either singly.

*What is known as Dr. Cowling's rule approximates this very closely and has the advantage of being easily remembered:—Given the dose for an adult, the dose for a child is obtained by dividing the number of the following birthday by 24. For example: the dose for an adult is 16 grains; for a child of two years of age the dose will be $3 \div 24 = 1-8$; 1-8 of 16 is 2 (grains). [Ed.]

14. **Chemical incompatibles** in the main are to be avoided; yet, because of such incompatibility, the resulting compound is not necessarily rendered inert. Here a trifle of chemical knowledge is sometimes invaluable. Of all unchemical combinations perhaps the most signally useful is that of opium and acetate of lead; they react, procuring the acetate of morphia and meconate of lead. (See page 261).

15. **Regulation of Dose.**—Almost every article in the *materia medica* operates differently when given in a large or small dose. **Tartar emetic**, for example, in dose of one-twelfth to one-sixth grain acts as a *diaphoretic* and *expectorant*; but increased to two or three grains is powerfully *emetic*. **Opium** is a *stimulant* in small, and *narcotic* in large, doses; and **oil of turpentine** in doses of *one or two drachms* acts as an acrid irritant of the kidneys and genito-urinary organs, whilst in dose of *one to two ounces* it operates freely upon the bowels without renal or vesical irritation. There is a source of error in the regulation of doses to which little attention is paid,—viz., the difference between the *drop* and the *minim*. Reference to the following list will show how erroneous is the common estimate that both are one-sixtieth of a drachm, and is convincing evidence of the value of *metric prescribing*:

APPROXIMATE TABLE OF DROPS TO FLUIDRACHM.

	DROPS.		DROPS.
Acid, Acetic	120	Oil of Orange	120
“ Hydrocyanic, dilute.....	45	“ Olive	76
“ Muriatic	54	“ Peppermint.....	103
“ Nitric	84	“ Rosemary.....	104
“ “ dilute	51	“ Sassafas.....	102
“ Sulphuric	90	“ Savine,	
“ “ dilute	51	“ Wintergreen.....	
Alcohol (rectified).....	138	Syrup of Acacia.....	58
“ (proof).....	120	“ Squills	85
Ammonia Water, strong.....	50	Tincture of Aconite.....	118
“ “ weak	40	“ Asafoetida.....	120
Creasote	91	“ Belladonna.....	110
Chloroform	180	“ Digitalis.....	120
Ether, Sulphuric	150	“ Guaiac	120
“ Hoffman's.....	90	“ Iodine	114
“ Spirits of Nitrous.....	90	“ Iron, Muriate.....	106
Glycerine	55	“ Opium.....	147
Liquor, Iodine Comp	76	“ “ Camphor.....	110
“ Donovan's Solution.....	52	“ Tolu.....	130
“ Fowler's Solution.....	60	Vinegar, Distilled.....	78
Oil of Sweet Almonds.....	120	“ Colchicum.....	80
“ Anise	85	“ Opium	89
“ Caraway	105	“ Squills	80
“ Cloves	103	Wine, Sherry, Madeira.....	78
“ Chenopodium.....	97	“ Antimony	72
“ Cinnamon.....	100	“ Colchicum.....	75
“ Croton	80	“ Opium	78
“ Cubebs	86	“ Aromatic.....	64
“ Fennel	103	Water, Distilled.....	45
“ Parsley (Apiol).....	55	“ Spring.....	64

N B.—Drops tend to vary according to the size of the lip of the bottle poured from. Patients should provide themselves with *minim pipette-tube*, and the physician may carry the same in a thermometer-case.

16. **The dose of any given medicine, particularly a narcotic or purgative**, should be regulated rather in accordance with the effect produced in each individual case than from the published tables; the latter are given as *guides* merely, and are not infallible.

17. **The character, period and form of disease** influence the operations of medicines in no slight degree. He who treats a disease by name only is, to say the least, guilty of great indiscretion. Again, in epidemics it is worthy of note that a remedy which is highly beneficial at one period appears to possess no efficacy at another. Usually while the onset and fastigium of an epidemic is marked by the multitude of deaths, with its wane recoveries are the rule; hence the *period* of an epidemic influences the action of medicines,—the physiology of which cannot here be discussed.

18. **Certain morbid conditions of body, or intensity of disease**, also modify the action of remedies. Note the tolerance of opium in spasmodic affections, and mercury in yellow fever or suppurative hepatitis.

19. The same is true of **deranged conditions of functions** also, especially where the digestive organs are involved. Perhaps quinine may cease to act as an antiperiodic, digitalis as a diuretic; or tonics prove distressing irritants.

20. **Diet.**—The influence of diet is marked. **Antiphlogistics** refuse to act in the presence of an excess of animal foods along with stimulants; the latter notoriously interfere with the action of mercurials. **Salt** retards the action of

nitrate of silver; **amylaceous foods** neutralize iodic preparations, and **iron salts** are inoperative when confronted with vegetable acids during febrile condition.

21. **The period of the day** at which remedies should be administered:—**Narcotics, emetics, and diaphoretics** operate most favorably when administered an *hour or two before bedtime*. **Diuretics** are best given during the day, when the body may be kept cool; **resinous cathartics** just *at bedtime*; other cathartics, such as neutral salts, oil, and those whose operation is speedy, *early in the morning*. **Quinine** acts best on an *empty stomach*; **arsenic** on a *full one*.

22. **Light, air, and exercise** in the main possess more therapeutical power than drugs,—at all events they are very essential adjuvants. **Confinement in close, dark, and ill-ventilated apartments effectually counteracts any benefit that otherwise might be had from tonics**. Exposure to the sun, however, hastens the physiological effect of nitrate of silver; and **violent physical action retards the effect of diuretics**.

23. **Season** also is a modifying agent. Some diseases, independent of all remedial measures, improve at certain seasons and retrograde at others: scrofulous patients almost invariably improve in *summer*, remain stationary in *winter* and decline in the spring. Again, **turpentine** should *never be given alone in large doses during cold or damp weather*, because it then tends to supply fuel for the evolution of animal heat rather than exert any therapeutic properties.

TABLE OF QUANTITIES

not to be exceeded in prescriptions without caution mark (!) to indicate that the dose is an unusual one.

ARTICLES.	SINGLE DOSES.	AGGREGATE IN ONE DAY.
Acid, Arsenious	1-6 grain.	$\frac{1}{2}$ grain.
Acid, Hydrocyanic, Diluted	10 minims.	30 minims.
Acid, Nitro-muriatic	5 minims.	30 minims.
Aconite Leaf, Extract of	1½ grains.	10 grains.
Aconite Root, Fluid Extract of	5 minims.	15 minims.
Aconite Root, Tincture of	6 minims.	15 minims.
Aconite Root, Fleming's Tincture of	3 minims.	10 minims.
Aconitine and its Salts	1-50 grain.	1-10 grain.
American Hellebore, Fluid Extract of	3 minims.	10 minims.
American Hellebore, Tincture of	5 minims.	12 minims.
Antimony and Potassium, Tartrate of	2 grains.	
Arsenate of Sodium	$\frac{1}{4}$ grain.	1 grain.
Arsenic, Solution of Chloride of	6 minims.	15 minims.
Arsenic, Iodide of	1-6 grain.	$\frac{1}{2}$ grain.
Arsenic and Mercury, Solution of Iodides of	20 minims.	60 minims.
Arsenite of Potassium, Solution of	15 minims.	45 minims.
Atropine and its Salts	1-60 grain.	1-20 grain.
Belladonna, Extract of	1 grain.	5 grains.
Calabar Bean, Extract of	$\frac{1}{4}$ grain.	1 grain.
Codeia	2 grains.	5 grains.
Coniine	1-16 grain.	$\frac{1}{2}$ grain.
Conium, Extract of	2 grains.	8 grains.
Croton Oil	2 minims.	
Cyanide of Potassium	$\frac{1}{4}$ grain.	1 grain.
Digitalin	1-50 grain.	1-20 grain.
Digitalis, Extract of	$\frac{1}{2}$ grain.	1 grain.
Hyoscyamus, Extract of	3 grains.	10 grains.
Ignatia, Extract of	3 grains.	10 grains.
Indian Cannabis, Extract of	1 grain.	5 grains.
Mercury, Corrosive Chloride of	$\frac{1}{4}$ grain.	$\frac{3}{4}$ grain.
Mercury, Iodide of, Red	$\frac{1}{4}$ grain.	1 grain.
Mercury, Iodide of, Yellow	1 grain.	3 grains.
Morphine and its Salts	$\frac{1}{2}$ grain.	2 grains.
Nux Vomica, Extract of	2 grains.	8 grains.
Opium	2 grains.	10 grains.
Opium, Extract of	1 grain.	4 grains.
Opium, Tincture of	30 minims.	120 minims.
Phosphorus	1-5 grain.	$\frac{1}{2}$ grain.
Stramonium Leaves, Extract of	2 grains.	5 grains.
Stramonium Seed, Extract of	1 grain.	3 grains.
Strychnine and its Salts	1-12 grain.	$\frac{1}{4}$ grain.

24. **The form** in which the remedy is administered may influence its action. Where a speedy effect is desired liquid form is usually desirable; especially is this the case with quinine and morphine. **Tannin**, when designed to act upon the stomach and bowels, is most suitably exhibited in form of pill; but solution is to be preferred if the general circulation is the goal. **Digitalis** in *tincture* is a direct heart sedative; in *infusion* a diuretic. Insoluble remedies, such as **calomel** and **iron peroxide**, are best given in form of powder or pill, and the pill form usually is improved by the addition of soap, as it hastens solution in the intestines and renders the action of purgatives milder and at the same time more certain. Powders may be given in honey, syrup, wafers, tissue-paper, or gelatin capsules,—the latter, however, are insoluble in the presence of an excess of acids.

25. **Disguising the taste of medicines** is often a matter of consideration. **Castor and cod-liver oils** are rendered palatable by chewing orange-peel or aromatics for a few moments before taking; **senna** may be drank as *tea*, with milk and sugar; **tannin** covers the taste of **quinine**;^{*} liquorice, of **aloes**; infusion of roses, **Epsom salts**.

N.B.—Drugs per rectum or vagina may be administered in *three times* the dose by the mouth. The same holds good in the preparation of oleates and unguents employed **endermically**.

For hypodermic use, drugs should be decreased until but one-sixth to one-fourth of the dose ordinarily administered by the mouth. Hypodermic fluids may be prepared as follows:

Morphia.—0.80 gramme to 32. grammes of camphor water; the acetate or muriate is to be preferred to the sulphate. 5 minims represents $\frac{1}{4}$ grain of morphia.

Atropia.—0.13 gramme in 4 grammes of water; 5 minims represents 1-45 grain. *Be cautious in administering to flaccid-haired, light-complexioned, nervous females.*

Hydrocyanic Acid.—Give 2 to 4 minims in water.

Physostigma.—Make fresh solution of solid extract, 0.13 gramme to 4 grammes of water, and filter; 5 minims represents $\frac{1}{4}$ grain.

Strychnia.—0.13 gramme in 8. grammes of water; 1 minim = 1-60 grain.

THERMOMETRY.

IN CHILDREN the temperature is normally *one to two degrees higher* than in adults.

THE TEMPERATURE IS NORMALLY ONE DEGREE HIGHER under the tongue than in the axilla. It is *highest* upon awakening in the morning; *lowest* at midnight.

A **rise of one degree** in temperature usually marks an increase of the pulse from six to ten beats a minute.

Continued temperature above 98.5° indicates *prostration and illness*; 101° to 105°, *severe fever*; 105° to 108°, *danger*; 108° to 109°, *impending death*.

A **temperature of 105° or 106°** on first day of illness is *prima facie* evidence of ephemeral fever: it is *not typhoid or typhus*, but probably *malarious*.

Though the typical evidences of *pneumonia* are present, if the thermometer fails to reach 101.7°, it may be concluded no soft infiltration of lungs is present.

A high temperature after the eruption of **measles** has faded indicates complications. An evening typhoid temperature of 103.5° indicates a mild course of fever. 105° in the evening or 104° in the morning, in the third week, indicates danger. A temperature of 104° and upwards, in pneumonia, indicates a severe attack. A temperature of 104° is always alarming in acute rheumatism; look for cardiac complications. In jaundice a rise of temperature is unfavorable. A rise of temperature in a puerperal female indicates the approach of pelvic inflammation. An increase of temperature in tuberculosis shows an advance of the disease or rise of complications.

Daily fluctuations of temperature are associated with malarial fever, typhus, typhoid, exanthemata, rheumatism, pyæmia, pneumonia, and acute tuberculosis. An even temperature from morning until evening is *favorable*. A high temperature from evening until morning is *unfavorable*. A falling temperature from evening until morning is *favorable*. A rising temperature from evening until morning is dangerous. The temperature of the body must be normal before convalescence begins.

N.B.—Mercurial thermometers increase in readings with age. During the first year of their manufacture this amounts to as much as one degree, and sometimes more. This is why manufacturers advertise their wares as "*Seasoned*," which, however, if true, does not altogether do away with the error.

Thermometers whose readings vary more than 0.25° at the same temperature are in every way unreliable.

^{*}In this instance lack of bitterness is lack of therapeutic value since the resulting quinine tannate is an almost insoluble salt (see *National Dispensatory*, iv. ed., p. 1291.) It would be better to recommend some article such as Yerba santa, *q. v.*, which effects no chemical reaction, but simply obtunds the nerves of taste. [Ed.].

RESPIRATION.

Two months to two years.....	35 per minute.
Two to six years	23 " "
Six to twelve years.....	20 " "
Twelve to fifteen years.....	18 " "
Fifteen to twenty-one years.....	16 to 18 " "

Respiration in the adult female is usually a trifle faster than in the male, especially during pregnancy.

RECTAL ALIMENTATION.

Remember that any excess of *four fluidounces* will promote disturbance. Half a teacupful is the usual amount, which may be repeated *not oftener* than every *eighth* hour.

FACTS REGARDING CHILDREN.

1. A healthy infant should **sleep** eighteen to twenty hours of the twenty-four.
2. As a rule a child cannot **raise its head** from the pillow before the *second or third* month, and cannot **sit erect** before the *fifth* month.
3. The *tenth* month is the usual time for walking to begin, but longer delay is no evidence of disease.
4. An infant should be **nursed** at *regular* intervals, and also supplied with **water**. Should be **weaned** from tenth to fourteenth month, *but not in early summer*.
5. A child does not **cry** from viciousness before the sixth month.
6. If **pulse** is 140 and child ill, cause *overfeeding*; if 40 and ill, cause *meningitis*.
7. **Tongue** furred with whitish curd, *intestinal irritation*; red, dry, and hot, *intestinal inflammation*; pale and flabby, *great debility*, and if tremulous, *nervous debility*; white fur, *fever*; yellow, *disturbance of liver and stomach*; brown and dry, *typhoid condition or blood-poisoning*; bright strawberry, look out for *scarlatina*; glazed, *indigestion*; glazed, with blue appearance, *congenital syphilis*.
8. **Congestion** of cheeks, without *cachexia* or chronic condition, means *inflammatory or fibrile condition*; of face, ears and forehead, of brief duration, along with strabismus, oscillation of iris, irregularity of pupil, and falling of upper lip, *cerebral affection*.
9. Progressive and marked **emaciation**, *grave subacute or chronic affection*.
10. **Bulbar hypertrophy of fingers and curving nails** are signs of *interference in functions of circulation*; hypertrophy of spongy portion of bones, or lividity of feet and hands with hypertrophy of connective tissue, *rachitis*.
11. **Pertinacious lividity**, or lividity procured by emotion and excitement, if respiration is normal, denotes *heart-fault or fault in trunk vessels*; temporary lividity, *grave acute disease*, probably of respiratory organs.
12. Absence of **tears** in children of six month and under is to suspect *disease that will prove fatal*; piercing and acute cries are suggestive of *cerebro-spinal trouble*.
13. **Thick, purulent secretion between the eyelids**, as derived from Meibomian glands, affords evidence of *grave prostration*; and passive congestion of conjunctival vessels, of *approaching death*.
14. **Irregular muscular movements**, partially controlled by the will, *chorea*; contraction of eyebrows and avoidance of light, *cephalgia*.
15. **When hand is constantly pressed to the head**, or head against parent or nurse, suspect *ear-disease*; if fingers are carried to mouth with great agitation, look for *abnormal condition of larynx*; and if head is constantly turned from side to side, suspect *laryngeal obstruction*.
16. **Hoarse and indistinct voice** is usually indicative of *laryngitis*; feeble and plaintive one, of *abdominal trouble*.
17. **Strong and sonorous cough**, *spasmodic croup*; hoarse and rough, *membranous croup*; suppressed and painful, *pleurisy or pneumonia*; convulsive, *pertussis*. Dry and painless cough may accompany *indigestion, difficult dentition, or febrile condition*.
18. **Respiration** slow and intermittent, accompanied by sighs, suggests *cerebral disease*; if intermittent and accelerated, *capillary bronchitis*; superficial and accelerated, *inflammatory condition of larynx and trachea*. The normal rate is, two months to two years, 35 per minute; two to six years, 23 per minute; six to twelve years, 20 per minute; twelve to fifteen years, 18 per minute.
19. **Normal pulse**, *in utero*, 140 to 150 per minute; of newborn 130 to 140; first year, 115 to 130; second year, 106 to 115; third year, 95 to 105; fifth to twelfth year, 80 to 90; thirteenth to twenty-first year, 75 to 85; twenty-first to sixtieth year, 70 to 75; in old age, 75 to 85.
20. **Acute fevers** always bring *remissions* in children; chronic fevers, *intermissions*.

21. **In early childhood** there is no relation between the intensity of the symptoms and the material lesion. The most intense fever, with restlessness, cries, and spasmodic movements, may disappear, without leaving a trace, in less than twenty-four hours.

22. **A child grows** most rapidly during its first year—six to seven inches; from fourth to sixteenth about *two inches* annually; thence to twentieth, *one inch*. Commonly, a child at two and a half years has attained half its ultimate adult stature. The diseases of youth always accelerate growth.

23. **First Dentition.**—The *central incisors* appear from the fifth to eighth month; *lateral incisors*, from seventh to ninth month; *first molars*, from tenth to twelfth month; *canines*, sixteenth to twentieth month; *second molars*, twentieth to thirtieth month; and, as a rule, teeth in the lower jaw precede corresponding teeth in the upper.

24. **Second Dentition.**—*First molars*, fifth to seventh year; *central incisors*, seventh to eighth year; *lateral incisors*, eighth to ninth year; *first bicusps*, ninth to tenth year; *second bicusps*, tenth to eleventh year; *canines*, eleventh to twelfth year; *second molars*, twelfth to thirteenth year; *third molars*, seventeenth to twenty-fifth year.

25. N. B.—While children bear *cathartics* and *mercurials* better proportionately to age than adults, *narcotics*, on the contrary, they do not bear so well; the latter are especially apt to act *cumulatively*.

26. Graduated Table for Administering Laudanum.

	MINIMS		MINIMS
At birth up to 1 month.....	1-2 to 1	1 to 10 years.....	5 to 15
Under 1 year.....	1-2 to 3	10 to 15 years.....	10 to 20
1 to 2 years	1 to 5	At 15 years.....	15 to 20
2 to 5 years	2 to 8	Adult	25 to 30

It is important that *laudanum* be of *known* strength. [Tinct. Opii should always be endorsed, U.S.P., when prescribing. Ed.]

INCOMPATIBILITY.

THIS MAY BE 1, CHEMICAL; 2, PHARMACEUTICAL; 3, THERAPEUTIC AND PHYSIOLOGICAL.

Chemical incompatibility, as has already been pointed out (Posology) need not and does not always signify therapeutic inertness. Substances which decompose others on admixture or are chemically incompatible are sometimes intentionally combined in order to obtain and make use of the resulting compound; such an instance occurs in the old familiar "Black Wash" or Lotio nigra, where Calomel and Lime water react on each other.

The infinite number of combinations possible makes it difficult to give but a few of the most general rules to be observed in exhibiting chemicals in order to avoid incompatibility of admixture.—

Alkalies and alkaline salts precipitate alkaloids and their salts.

Caustic alkalies decompose salts of the metals proper.

Tannic and gallic acids and substances containing them precipitate albumen and alkaloids and form 'inks' with tincture iron chloride.

Glucosides such as digitalin, cannabin, &c., are decomposed or incompatible with free acids and with emulsin.

Pharmaceutical incompatibility. This applies to articles physically incapable of mixing; such as spirits of nitrous ether with tincture of guaiacum—a gelatinous mass is the result; fixed oils and copaiba with aqueous liquids—unless excipients accompany; an acid with a quinine and licorice mixture—the acid precipitates the glycyrrhizin of the licorice and as the latter is present merely to mask the taste of the quinine, throwing it out of solution frustrates this object; fluid extracts marked in previous pages as "not miscible with water" contain resinous or other matter which will be precipitated on addition of aqueous liquids—they should be dispensed with acacia or some other emulsifier, or with simple elixir if the presence of alcohol is not contra-indicated.

Therapeutic or physiological incompatibility: Applied to substances which if administered together would neutralize each other's action more or less perfectly. Thus chloral hydrate is physiologically incompatible and should not be prescribed with nuxvomica. Antagonists like these are not necessarily incompatible in other respects. They are also sometimes intentionally combined as in the hypodermic administration of morphine, with atropine as a supposed safeguard—entirely a supposition by the way, according to recent authorities.* Space prevents here giving an extended list but very many incompatibles of this class will be found under Poisons and Antidotes, q. v.

* See *Therapeutic Gazette*, Sept. 15, 1887.

LATIN WORDS, PHRASES, ETC., USED IN WRITING
PRESCRIPTIONS; WITH THEIR CONTRACTIONS, AND
CORRESPONDING ENGLISH EQUIVALENTS.

WORD OR PHRASE.	AS GENERALLY WRITTEN.	ENGLISH EQUIVALENT.
Ad (with accus. case)	Ad	To, or up to.
Ad duas vices	Ad 2 vic.	At twice taking.
Ad libitum	Ad lib.	At pleasure.
Ad tertiam vicem	Ad 3tiam vic.	For three times.
Adde	Add.	Add.
Ad gratam aciditatem	Ad grat. acid.	To an agreeable sourness.
Adhibendus	Adhib.	To be administered.
Admove	Admov.	Apply.
Alternis horis	Alt. hor.	Every other hour.
Alvo adstricta	Alv. adst.	The bowels being confined.
Ana	ā	Of each.
Ante cibum	Ante cib.	Before eating.
Aqua	Aq.	Water.
Aqua astricta	Aq. astr.	Frozen water.
Aqua bulliens	Aq. bull.	Boiling water.
Aqua fervens	Aq. ferv.	Hot water.
Aqua fontana or fontis	Aq. font.	Spring water.
Bene	Bene	Well.
Bis in die, or dies	Bis die or Bisind.	Twice a day.
Bulliat or Bulliant	Bull.	Let boil.
Capiat	Cap.	Let him take.
Caute	Caute	Cautiously.
Cibus	Cib.	Food.
Cochleare magnum	Coch. mag.	A tablespoonful (3 ss).
Cochleare medium, or modicum.	Coch. med. or mod.	A dessertspoonful (3 ij).
Cochleare parvum	Coch. parv.	A teaspoonful (3 j).
Cola or Coletur	Col. or Colet.	Strain.
Collyrium	Collyr.	An eye-wash.
Coloretur	Coloret.	Let it be colored.
Congius	Cong.	A gallon.
Continuantur remedia	Cont. rem.	Let the medicines be continued.
Coque, Coquantur	Coq.	Boil, let them be boiled.
Cras, Crastinus	Crast.	To-morrow.
Cras mane sumendus	Cras mane sumend.	To be taken to-morrow morning.
Cujus, Cujus-libet	Cuj.	Of which, of any.
Cum (with abl. case)	C.	With.
Debita spissitudo	Deb. spiss.	A proper consistence
Decubitus	Decub.	Lying down.
De die in diem	De d. in d.	From day to day.
Diebus alternis	Dieb. alt.	Every other day.
Diebus tertiis	Dieb. tert.	Every third day.
Dilue, Dilutus	Dil.	Dilute (thou), Diluted.
Dimidius	Dim.	One-half.
Dividatur in partes æquales	D. in p. æq.	Let it be divided into equal parts.
Donec alvus soluta fuerit	Donec alv. sol. ft.	Until the bowels shall be opened.
Dosis	D.	A dose
Drachma	Dr. or 3	A drachm (60 grains).
Eadem (fem.)	Ead.	The same.
Ejusdem	Ejusd.	Of the same.
Fac or Fiat	F or Ft.	Make, let be made.
Fac pilulas duodecim	F. pil. xij.	Make 12 pills.
Fiant chartulæ xij	Ft. chart. xij.	Let 12 powders be made.
Fiat emplastrum ves- icatorium	Ft. emp. vesic.	Let a blister be made.
Fiat massa, et divide in pilulas xij	Ft. mas. div. in pil. xij.	Let a mass be made, and divide it into 12 pills.
Fiant pilulæ xij	Ft. pil. xij.	Let 12 pills be made.
Fiat pulvis in chartu- las xij dividenda	Ft. pulv. in chart. xij div.	Let a powder be made to be divid- ed into xij papers.
Gradatim	Grad.	By degrees, gradually.
Gutta, Guttae	Gtt.	A drop, Drops.

WORD OR PHRASE.	AS GENERALLY WRITTEN.	ENGLISH EQUIVALENT.
Guttatim	Guttat.	By drops.
Harum pilularum sumantur tres	Har. pil. sum. iij	Let three of these pil's be taken.
Haustus	Haust.	A draught.
Horâ decubitûs	Hor. decub.	At bedtime.
Horâ somni	Hor. som.	At the hour of sleep.
Horæ unius spatîo	Hor. j spat.	After one hour.
Idem	Id.	The same.
In dies	Ind.	Daily or From day to day.
Mane	Mane	In the morning.
Mica panis	Mic. pan.	Crumb of bread.
Misce	M.	Mix.
Mitte	Mitt.	Send.
Modo præscripto	Mod. præsc.	In the manner prescribed.
More dictu	Mor dict.	In the manner directed.
More solito	Mor. sol.	In the usual manner.
Ne tradas sine nummo	Ne tr. s. num.	Do not deliver without the money.
Nocte maneque	Noct. maneq.	At night, and in the morning.
Non repetatur	Non repetat.	Let it not be repeated.
Nox, Noctis	Nox, Noct.	The night, of the night.
Octarius	O., Oct.	A pint (½ xvj).
Omni horâ	Omn. hor.	Every hour.
Omni bihorio	Omn. bih.	Every two hours.
Omni quadrante horæ	Omn. quadr. hor.	Every ¼ hour.
Omni mane	Omn. mane	Every morning.
Omni nocte	Omn. noct.	Every night.
Partes æquales	Pt. æq.	Equal parts.
Penicillum camelinum	Penicil. cam.	A camel's-hair pencil, or brush.
Per. (with accus. case)	Per	Through, By.
Phiala prius agitata	P. P. A.	The bottle having been first shaken.
Post cibum	Post cib.	After eating.
Pro (with abl. case)	Pro	For.
Pro ratione ætatis	Pro rat. æt.	According to the age of the patient.
Pro re nata	P. r. n.	According to circumstances.
Quantum libet	Q. lib.	As much as you please.
Quantum sufficiat	Q. s.	As much as is sufficient.
Quâquâ horâ	Qq hor.	Every hour.
Quoque	Q q.	Also.
Quorum	Quor.	Of which.
Quotidie	Quotid.	Daily.
Recipe	R̄	Take.
Redactus in pulverem	Red. in. pulv.	Let it be reduced to powder.
Repetatur	Rept.	Let it be repeated.
Scrupulum	Scrup. or ʒ	A scruple (ʒ0 grains).
Secundum artem	S. A.	According to art.
Semis, or Semissis	Ss.	A half.
Semi-hora	Semih.	Half an hour.
Sesqui	Sesqui	As much again by half.
Sesquihora	Sesqh.	An hour and a half.
Signa	Sig.	Write, or Mark (thou).
Simul	Simul	Together.
Sine (with abl. case)	Sin.	Without.
Singulorum	Sing.	Of each.
Si opus sit	Si op. sit	If necessary.
Solve	Solv.	Dissolve.
Spiritus vini rectificatus	Spt. vin. rect.	Rectified spirit of wine (i.e. Alcohol).
Statim	Stat.	Immediately.
Stet, or Stent	St.	Let it (or them) stand.
Subinde	Subind.	Frequently.
Sumat talem	Sum. tal.	Take one such.
Sumatur	Sum.	Let it be taken.
Summo mane	Summo mane	Early in the morning.
Talis	Tal.	Such a one.
Ter in die, or Ter die	T. i. d., or T. d.	Thrice daily.
Tussis	Tus	A cough.
Ultimus præscriptus	Ult. præsc.	The last ordered.
Uncia	Unc. or ʒ	An ounce.
Ut dictum	Ut dict.	As directed.
Vel	Vel	Or.
Vesper, -eris	Vesp.	The evening.

DIFFERENTIAL DIAGNOSIS OF ERUPTIVE FEVERS.

NAME	PERIOD OF INCUBATION.	OCCURRENCE OF ERUPTION.	CHARACTER OF ERUPTION.	FADING OR DISAPPEARANCE OF ERUPTION.	DURATION OF ILLNESS.
Scarlatina *	4 to 10 days.	Usually second or third day of fever (within thirty-six hours of actual commencement of illness).	Bright scarlet and diffused.	Fifth day of fever, commonly.	8 to 15 days.
Measles	10 to 14 days.	Fourth day of fever, or after seventy-two hours of illness.	Small red dots like flea-bites; crescentic.	On seventh day of fever.	6 to 10 days.
Rötheln	7 to 14 days.	Fourth to sixth day.	Like measles, but less distinct; patches more irregular and brightest near center.	Fourth day after appearance. Often desquamates in bran-like scales, but is uncertain.	8 to 10 days.
Variola	12 days.	Third day of fever.	Small red pimples, becoming vesicular and subsequently pustular; umbilicated.	Pustules mature on eighth or ninth day; scabs follow on tenth or twelfth falling off two days later.	14 to 21 days.
Varicella	4 days.	Second day of fever.	Small rose pimples, becoming vesicles.	Slight scabs form on fourth day of fever.	6 or 7 days.
Erysipelas	3 to 7 days.	Second or third day.	Diffuse redness and swelling.	Uncertain.	Uncertain.
Typhus Fever ..	1 to 12 days.	Fourth to eighth day.	Mulberry color, usually over abdomen.	Uncertain.	14 to 21 days.
Typhoid Fever ..	10 to 14 days or (sometimes) suddenly.	Seventh to fourteenth day.	Rose colored spots, few in number and far apart.	Uncertain (may be wholly wanting).	20 to 30 days.

* Scarlatina presents a great variety of irregular forms; hence, the practitioner should ever be on his guard.

Therapeutic Suggestions; or Diseases and their Remedies.

This is necessarily a condensed list as well of the diseases as of their remedies. Pharmaceutical preparations are given only in a very few instances, since not only does space forbid but the physician himself can much better determine the suitability of the several preparations of any desired remedy to the case in hand.

The meaning is given of the majority of the terms used to designate diseases—rendering the list, in this respect, a GLOSSARY of the foregoing pages.

- ABORTION, THREATENED.**—Black Haw, Black Cohosh, Iron, Opium.
- ABSCCESS.**—Belladonna, Calcium sulphide, Calcium lactophosphate.
- ACIDITY OF STOMACH.**—Bismuth, Charcoal, Hydrochloric Acid, Liquid Acid Phosphate, Nux vomica, Sodium bicarbonate (temporary only).
- ACNE.**—Alkalies, Arsenic, Calcium sulphide, Hypophosphites, Phosphorus.
- ACUTE MANIA.**—Bittersweet, Bromides, Conium, Indian Cannabis, Jamaica Dogwood, Valerian.
- ADYNAMIA** (Debility; Deficiency of vital power).—Arsenic, Blood-root, Cinchona, Gentian, Golden-seal, Hypophosphites, Iron, Liquid Acid Phosphate, Strychnine.
- AFTER-PAINS.**—Black Cohosh, Button-snake root, Chloroform (liniment), Ergot, Gelsemium.
- AGUE.**—See Intermittent Fever.
- ALCOHOLISM.**—Ajowan, Black Cohosh, Capsicum, Cinchona, Coca, Lupulin.
- AMAUSIOSIS** (Impaired vision due to cerebral causes).—Arnica, Santonin, Strychnine.
- AMENORRHOEA** (Suppression of the menses).—Aconite, Aloes, Apioi, Black Cohosh, Blue Cohosh, Button-snake root, Cantharides, Iron, Potassium permanganate, Pulsatilla, Squaw-vine, Water-pepper.
- ANÆMIA** (Deficiency or insufficient formation of the red corpuscles of the blood).—Calcium lactophosphate, Hypophosphites, Iron, Quinine, Strychnine.
- ANEURISM** (An abnormal dilatation or rupture of the coats of an artery; also dilatation of the walls of the heart).—American Hellebore, Ergot, Foxglove, Potassium iodide.
- ANGINA PECTORIS** (An affection often connected with ossification of the heart or great vessels, characterized by most excruciating pain in chest and arms).—Aconite, Amyl nitrite, Arsenic, Nitroglycerin, Sodium nitrite.
- APHONIA** (Loss of voice).—Atropine, Borax, Ignatia, Ipecac, Inhalants—see page 211.
- APHTHÆ** (Small white ulcers confined to lips, mouth and intestinal canal).—Alum root, Agrimony, Baycuru, Borax, Potassium chlorate, Khatany, Tonics.
- APOPLEXY** (Effusion of blood into the substance of any organ, but generally applied to the brain—producing unconsciousness, loss of sensation and voluntary motion).—Aconite, Colocynth, Croton oil, Elaterium.
- ARTHRITIS** (Inflammation of the joints).—Arsenic, Cod-liver oil, Iodine, Oil of Wintergreen, Salicylates.
- ASCARIDES** (*Ascaris lumbricoides*—Round worm; *Ascaris vermicularis*—Thread worm).—American Wormseed, Pink-root, Santonin, Yerba buena.
- ASCITES** (Accumulation of serous fluid in the cavity of the peritoneum).—American Ivy, Canadian Hemp, Elaterium, Flowering spurge, Foxglove, Gravel plant, Hair-cap moss, Nettle, Plantain, Purslane.
- ASTHMA** (Spasmodic stricture of the bronchial muscle surrounding the excretory ducts or smaller bronchial tubes).—Ailanthus glandulosa, Amyl nitrite, Ammoniac, Black Cohosh, Blood-root, Euphorbia pilulifera, Grindelia robusta, Musk-root, Quebracho, Sundew, Water Fennel, Yerba santa.
- BILIQUENESS.**—Aloes, Blue Flag, Boldo, Jurubeba, Lime-juice and Pepsin, Mandrake, Mercury, Mineral acids, Wahoo.
- BLADDER, CATARRH OF.**—Balsam Fir, Balsam Poplar, Boldo, Buchu, Cantharides, Cubeb, Eucalyptus, Juniper berries, Kava Kava, Manzanita, Marshmallow, Pareira Brava, Pichi, Pipsissewa, Queen of the Meadow, Rhus aromatica, Stone root, Uva ursi.
- BLADDER, IRRITABLE.**—Belladonna, Buchu, Couch-grass, Cubeb, Indian Cannabis, Pichi, Stavesacre.
- BOILS.**—Arsenic, Calcium sulphide, Hypophosphites, Silver nitrate (to abort), Syrup Trifolium comp.
- BREATH, FETID.**—Charcoal, Eucalyptus, Potassium permanganate (as a wash for the mouth).

BRIGHT'S DISEASE.—Aconite; Ammonium and Potassium bromides, for restlessness (no opiates!); Button-snake root, Foxglove, Indian Cannabis, Milk diet, Tincture Iron chloride.

BRONCHIECTASIS (Dilatation of the bronchi).—Hypophosphites, Liquid Acid Phosphate, Quinine.

BRONCHITIS (Inflammation of the bronchi).—Aconite, Abscess root, Agrimony, Ammoniac, Ammonium carbonate and chloride, Angelica, Apomorphine (the best of all expectorants), Asafetida, Balsam Fir, Balsam of Peru, Balsam Poplar, Benzoin, Black Cohosh, Blood-root, Button-bush, Chekan, Eucalyptus, Grindelia robusta, Inhalations (see page 211), Indian Turnip, Ipecac, Kava Kava, Lippia Mexicana, Lobelia, Squill, Tolu, Yerba santa.

BRONCHORRHOEA (Catarrh of the bronchi, chronic bronchitis).—Asafetida, Ammonium iodide with Arsenic, Beth root, Eucalyptus, Grindelia robusta, Myrtol.

BRUISES.—Arnica, Distilled Ext. Witch Hazel.

BUNION.—Iodine tincture, Mercury plaster, Silver nitrate.

CACHEXIA (A vitiated or bad condition or habit of the body).—Arnica, Arsenic, Black Walnut, Cod-liver oil, Hypophosphites, Iron Quinine and Strychnine, Liquid Acid Phosphate, Syrup Trifolium Comp. (see Red Clover, page 149.)

CALCULI, BILIARY (Gall stones).—Alkaline mineral water, Sodium choleate, Sodium phosphate.

CALCULI, RENAL AND VESICAL (Gravel).—Ammonium benzoate, Buchu, Button-bush, Corn-silk, Dwarf Elder, Gravel-plant, Hair-cap Moss, Hydrangea, Iron succinate, Pichi, Potassium citrate.

CANCER.—Alveloz milk, Chian turpentine, Iodoform, Syrup Trifolium Comp. for systemic treatment (see Red Clover, page 149).

CATARRH, ACUTE NASAL (Coryza).—Aconite, Black cohosh, Eyebright, Jaborandi.

CATARRH, CHRONIC NASAL (Ozaena).—Ammoniac, Blood-root, Golden-seal, Inhalations (see page 211), Senega, Virginia Stone-crop.

CEREBRAL ANÆMIA.—Amyl nitrite, Hypophosphites, Iron, Liquid Acid Phosphate, Phosphorus, Sanguis bovinus exsiccatus.

CEREBRAL CONGESTION.—Aconite, American Hellebore, Belledonna, Bromides in large doses, Blood-letting occasionally serviceable, Cathartics, Gelsemium.

CHILBLAINS.—Tincture of Benzoin Comp., Tincture Iodine diluted, Turpentine.

CHLOROSIS (The green sickness; owing to deficiency of red blood-corpuscles. See also anæmia).—Arsenic, Hypophosphites, Iron Quinine and Strychnine, Phosphorus, Syrup Trifolium Comp.

CHOLERA and CHOLERA MORBUS.—Ajowan, Camphor, Chlor-anodyne, Coto-bark, Ginger, Jamaica Dogwood, Musk-root, Tincture Opium Comp.

CHOREA. (Involuntary motions of the muscles).—Ailantus glandulosa, Asafetida, Black Cohosh, Calabar bean, Cod-liver oil, Conium, Henbane, Hypophosphites with iron, Indian Cannabis, Iron, Ladies' Slipper, Mistletoe, Mugwort, Musk-root, Sculicap, Skunk Cabbage, Strychnine, Valerian.

COLIC INTESTINAL.—Ajowan, Asafetida, Blue Cohosh, Capsicum, Chamomile oil, Chloranodyne, Chloroform, Crawley root, Dewee's Carminative (for infants), Fish-berries, Jamaica Dogwood, Nux Vomica, Spirits of ammonia.

CONSTIPATION.—Aloes, Belladonna, Berberis aquifolium, Buckeye bark, Butternut, Calabar bean, Cascara sagrada, Culver's root, Five-flowered Gentian, Golden-seal, Jurubeba, Mandrake, Nux vomica, Wahoo.

CONVALESCENCE.—American Centaury, American Columbo, Balmoney, Blue Gentian, Boneset, Buckhorn Brake, Coca elixir, Eucalyptus, Fringe-tree, Golden-seal, Guarana, Hypophosphites, Liquid Acid Phosphate, Quinine.

COUGH.—Arbor vite, Blood-root, "Bronchial Sedative," "Bronchial" Lozenges, Buckhorn Brake, Button-bush, Gelsemium, Grindelia robusta, Helianthella tenuifolia, Ipecac, Lobelia, Pulsatilla, Senega, Soap-tree, Squill, Wild Cherry, Yerba santa.

CROUP, MEMBRANOUS.—Alum (as an emetic dislodges false membrane and prevents it reforming), Belladonna, Blood-root, Hot fomentations, Ipecac, Mercury subsulphate, Potassium chlorate with Iron chloride.

CYANOSIS (The general venous congestion with bluish appearance consequent upon affections of the lungs or heart).—Amyl nitrite, Quebracho.

CYSTITIS (Inflammation of the bladder).—Buchu, Cantharides, Corn-silk, Conchgrass, Cubeb, Eucalyptus, Kava Kava, Paraira Brava, Pichi, Pipsissewa, Uva ursi.

DELIRIUM TREMENS.—Belladonna, Black Cohosh, Capsicum, Henbane, Hops, Indian Cannabis, Jamaica Dogwood, Musk-root, Stramonium, Strychnine.

DIABETES INSIPIDUS (Excessive production of urine).—Alum root, Cantharides, Ergot, Jaborandi, Opium with gallic acid.

DIABETES MELLITUS (Saccharine urine).—Jambul seed, Rhus aromatica, Milk diet, Starches and sugars forbidden.

DIARRHŒA.—Agrimony, Ailantus glandulosa, Bael, Baycuru, Beech-drop (Cancer root), Bistort, Black Alder, Blackberry, Buckhorn Brake, Canada Thistle, Cascarilla, Catechu, Chlor-anodyne, Coto bark, Cranesbill, Hardhack, Ipecac, Judas-tree, Kino, Logwood, Mountain Laurel, Musk-root, Opium, Plantain, Rhatany, Rhubarb, Rhus aromatica, Squaw vine, Sumach, Trumpet plant, White Oak, White Pond-lily, Willow herb.

DIPHThERIA.—Belladonna, Brandy (as a stimulant freely), Carica Papaya, Corrosive Sublimate, Inhalants (see page 211), Pepsin, Potassium chlorate and iron, Sedum acre.

DROPSY.—American Ivy, Arsenic, Blue Cohosh, Blue Flag, Bryonia, Broom, Buchu, Buckthorn berries, Canadian Hemp, Cantharides, Carpenter's Square, Colocynthis, Digitalis, Dwarf Elder, Elaterium, Flowering spurge, Hair-cap moss, Hellebore, Jalap, Juniper berries, Pipsissewa, Shepherd's Purse, Silkweed, Squill, Strophanthus, White Bryony.

DYSENTERY (An affection of the mucous membrane of the large intestine, characterized by frequent bloody or mucous stools).—Ailantus glandulosa, Angustura, Bael, Benne, Benzoin, Blackberry, Buckhorn Brake, Canada Thistle, Cascarilla, Castor oil, Coto bark, Cranesbill, Ergot, Hardhack, Ipecac, Logwood, New Jersey tea, Nux vomica, Opium, Plantain, Rhus aromatica, Willow herb, Witch-Hazel, Yarrow.

DYSMENORRHŒA (Painful menstruation).—Aconite, Apiol, Belladonna, Black Cohosh, Black-haw, Black Willow, Blue Cohosh, Button-snake root, Cramp bark, Gelsemium, Helonias, Life-root, Squaw-vine.

DYSPEPSIA (Difficult or depraved digestion).—Ajowan, Angustura, Bismuth and charcoal lozenges, Blood-root, Boneset, Buckbean, Carica Papaya (Papain and Papayotin), Cascara sagrada, Chamomile oil, Chirata, Gentian, Golden-seal, Helonias, Nux vomica, Mercury yellow oxide (tablets), Pancreatin, Pepsin, Pulsatilla, Sweet-flag.

DYSPNŒA (Difficulty in breathing).—Amyl nitrite, Asafetida, Black Cohosh, Chloroform (a few inhalations), Grindelia robusta.

DYSURIA (Strangury; Difficulty or pain in passing urine).—Cantharides, Gelsemium, Indian Cannabis, Pipsissewa, Potassium citrate, Rhus aromatica, Spirits Nitrous Ether, Squill, Wintergreen.

ECCHYMOsis (Livid or black marking on the skin from extravasation of blood).—Ammonium chloride, Arnica, Witch-Hazel.

ECZEMA (An eruption of very minute aggregated vesicles on an inflamed base).—Arsenic, Burdock seed, Cascara amara, Cod-liver oil (internally and externally), Oil of Cade (externally—diluted with castor oil), Pansy (viola tricolor).

ENTERITIS (Inflammation of the intestines).—Aconite, Jamaica Dogwood, Opium.

ENURESIS (Incontinency of urine).—Belladonna, Buchu, Cantharides, Iron iodide, Rhus aromatica, Sculleap.

EPILEPSY (Convulsions accompanied with unconsciousness).—Amyl nitrite, Ailantus glandulosa, Blue Cohosh, Chloroform (inhalations), Fish-berries, Ignata, Indian Cannabis, Mistletoe, Mugwort, Nitroglycerin, Strychnine, Valerian.

EPISTAXIS (Hemorrhage from the nose).—Aconite, Alum root, Belladonna, Cranesbill, Ipecac, Witch-Hazel distilled extract (internally and externally).

ERYSIPELAS (An unhealthy form of contagious inflammation).—Aconite, Ammonium carbonate, Beech-drop (Cancer root), Belladonna, Crawley root, Iodized collodion, Poison oak, Silver nitrate lotion (20 grains to the fluidounce), Tincture iron chloride.

FEVER.—Acetanilid, Acid drinks, Aconite, American Hellebore, Antipyrin, Arsenic, Belladonna, Gelsemium, Quinine.

FLATULENCE.—Ajowan, Anise, Buchu, Columbo, Crawley root, Eucalyptus oil, Ginger, Nux vomica, Pepsin (to promote digestion), Wintergreen oil.

GASTRALGIA (Cardialgia or Heartburn).—Ajowan, Arsenic, Bismuth, Nux vomica, Pepsin, Soda mint tablets, Trumpet plant.

GASTRODYNIA (Spasmodic or neuralgic pain in the stomach).—Belladonna, Chloral, Jamaica Dogwood, Nitroglycerin, Nux vomica, Opium, Sulphuric Ether.

GASTRIC DILATATION.—Bismuth with Magnesia, Eucalyptus oil, Ignatia, Nux vomica, Wintergreen oil, Concentrated food free of starches and sugars.

GLANDS, ENLARGED (Parotitis).—Bearsfoot, Bitter Bugleweed, Calcium sulphide, Carpenter's Square, Corrosive sublimate, Golden-seal, Iodine and the iodides, Syrup Trifolium Comp.

GLEET (A serous, mucous, or muco-purulent discharge from the urethra, not accompanied with scalding).—Balsam Fir, Balsam Poplar, Beech-drop (Cancer root), Buchu, Button-snake root, Cantharides, Copaiba, Judas-tree, Kava Kava, Sandalwood oil, Scouring-rush, Yerba Leuma.

GONORRHOEA (A contagious discharge of muco-pus from the mucous membranes of the generative organs and accompanied with scalding).—Aconite (acute stage) Balsam Fir, Balsam of Peru, Balsam Poplar, Boldo, Buchu, Button-snake root, Chewstick, Copaiba, Golden-seal (Fluid Golden-seal Colorless best for injections), Gurjun balsam, Indian Cannabis, Iron chloride tincture, Manzanita, Matico, Pareira Brava, Pipsissewa, Pulsatilla, Sandalwood, Scouring-rush, Uva ursi, White Poplar, Yerba Reuma.

GOUT (An inflammation of the fibrous and ligamentous parts of the joints).—Aconite, Belladonna, Bittersweet, Burdock, Colchicum, Guaiac, Lithium citrate, Potassium iodide, Strychnine, Sulphides.

HAY FEVER.—Aconite, Arsenic, Belladonna, Cocaine (solution 4 to 20 per cent. painted or sprayed), Eyebright, Grindelia robusta, Ignatia, Ipecac, Lobelia, Quinine.

HEART DISEASE.—Aconite, Adonis vernalis, Amyl nitrite, American Hellebore, Arsenic, Black Cohosh (safer than Digitalis), Digitalis, Lily of the valley, Night-blooming Cereus, Strophanthus.

HEMATEMESIS (Vomiting of blood).—Alum, Ergot, Gallic acid, Ipecac, Iron subsulphate (Monsel's salt), Lead acetate, Tannin, Turpentine, Witch-Hazel.

HEMATURIA (Bloody urine).—Buchu, Ergot (Ergotin hypodermically), Gallic acid, Ipecac, Indian Cannabis, Iron chloride tincture, Pipsissewa, Queen of the Meadow, Scouring-rush, Sulphuric acid with gallic, Witch-Hazel.

HEMOPTYSIS (Spitting of blood).—Aconite, Arnica, Beth root, Bugleweed, Iron (sprayed), Water Fennel, also in addition remedies above under Hematemesis.

HEMORRHOIDS ("Piles")—Alum root, Black Oak, Buckthorn bark, Ergot, Golden-seal (internally and externally), Iron subsulphate (Monsel's salt), Nutgall, Nux vomica, Opium (externally), Saraca Indica, Stillingia, Sulphur and Sulphides, Witch-Hazel (distilled extract).

HEPATIC DISORDERS.—Aconite (in acute inflammation), Ammonium chloride (in chronic torpidity—20 grains every 4 hours), Balmony, Barberrry, Bitter root, Blue Flag, Boldo, Buckeye, Butternut, Colchicum, Colocynthus, Culver's root, Five flowered Gentian, Garden Celandine, Grindelia squarrosa, Ipecac, Stillingia, Wahoo, White Bryony, Wood Betony.

HYPOCHONDRIASIS (A species of mental alienation).—Arsenic, Asafetida, Black Cohosh, Bromides, Coca, Conium, Henbane, Ignatia, Jamaica Dogwood, Musk-root, Opium (stimulant doses), Valerian.

HYSTERIA.—Arsenic, Asafetida, Black Cohosh, Blood-root (in hysteria from chronic hepatitis), Blue Cohosh, Chamomile oil, Ether, Eucalyptus, Ignatia, Indian Cannabis, Iron, Ladies' Slipper, Musk-root, Phosphorus, Potassium bromide, Scullcap, Skunk Cabbage, Valerian, Zinc valerianate.

IMPOTENCE.—Arsenic, Blood-root, Cantharides, Coca, Damiana, Indian Cannabis, Iron, Liquid Acid Phosphate, Phosphorus, Strychnine, Water-pepper.

INSOMNIA (Sleeplessness).—Belladonna, Bromides, Chamomile, Chloral, Henbane, Hops, Hypnone, Ignatia, Indian Cannabis, Jamaica Dogwood, Ladies' Slipper, Musk-root, Paraldehyde, Phosphorus, Scullcap.

INTERMITTENT FEVER.—Alstonia constricta, American Centaury, Apioi, Apple-tree bark, Arbor vitae, Arsenic, Boneset, Black Alder, Black Cohosh, Button-bush, California Fever-bush, Canadian hemp, Cascarella, Cedron seed, Cinchona, Dita, Five-flowered Gentian, Golden-seal, Grindelia squarrosa, Nux vomica, Quinine flower, Salicin, Spice bush, Warburg's Tincture, White Poplar.

IRRITABILITY.—Asafetida, Black Cohosh, Bugleweed, Catnep, Chamomile, Ignatia, Ladies' Slipper, Pulsatilla, Scullcap, Skunk Cabbage, Strychnine, Twin-leaf.

JAUNDICE (Icterus; diffusion of bile into the blood).—Arsenic, Ammonium chloride, Balmony, Barberrry, Bayberry, Bitter root, Blue Flag, Boldo, Carpenter's Square, Colocynthus, Culvers' root, Garden Celandine, Golden-seal, Ipecac, Jurubeba, Rhubarb (chiefly for children), Sodium Phosphate, Stillingia, Wahoo.

LACTATION (Secretion of milk).—*Diminishing flow*: Belladonna, Camphor (saturated solution in oil), Potassium iodide, Quinine, Sulphuric acid (diluted), Tannin. *Increasing flow*: Anise, Carica Papaya, Castor-oil plant, Jaborandi, Tea (black), Prickly ash, Wintergreen.

LARYNGITIS (Inflammation of the larynx).—Abscess root, Aconite, Belladonna, Catechu, Indian Turnip, Inhalants (page 211), Ipecac, Sumach, Virginia Stonecrop, Water Eryngo, Water Fennel.

- LEUCORRHOEA** (A whitish or colorless discharge from the vagina).—Agrimony, Balsam of Peru, Bayberry, Beech-drop (Cancer root), Bistort, Button-snake root, Canada Thistle, Cantharides, Ergot, Golden-seal, Helonias, Hemlock, Judas-tree, Manzanita, Matico, Musk-root, Pulsatilla, Rhus aromatica, Sumach, Virginia Stone-crop, Water Avens, White Pond Lily, White-oak Bark, Willow herb, Witch-Hazel, Yerba Buena.
- LOCOMOTOR ATAXIA** (Loss of power to coördinate voluntary motion).—Belladonna, Calabar bean, Electricity, Phosphorus, Potassium iodide, Silver nitrate, strychnine, Systemic tonics—cod-liver oil, iron, etc.
- LUMBAGO** (Rheumatism of the loins).—Aconite (internally and externally), American Hellebore, Belladonna plaster, Black Cohosh, Capsicum (externally), Chloroform liniment, Electricity, Jamaica Dogwood, Morphine with atropine, Poison Oak, Potassium iodide (if of syphilitic origin), Potassium nitrate (10 grain doses if the urine is dark colored). Good diet and rest.
- MANIA** (Insanity).—American Hellebore, Belladonna, Black Cohosh, Chloral, Digitalis, Duboisia, Ergot, Gelsemium, Henbane, Hops, Indian Cannabis, Jamaica Dogwood, Potassium bromide, Stramonium.
- MASTITIS** (Inflammation of the breasts).—Belladonna (internally and externally), Conium, Iodine (to remove consequent induration), Poke-root (internally and externally).
- MELANCHOLIA**.—Black Cohosh, Caffeine citrate, Camphor monobromated, Chloral, Coca, Ignatia, Indian Cannabis, Iron chloride tincture, Liquid Acid Phosphate, Opium (in stimulant doses), Potassium bromide, Valerian.
- MENINGITIS** (Inflammation of brain membrane).—Aconite, American Hellebore, Belladonna, Calabar bean, California Laurel, Ergot, Gelsemium, Henbane, Opium, Potassium bromide, Potassium iodide, Pulsatilla. Depletion; purgatives; cold.
- MENORRHAGIA** (Profuse menstruation).—Aloes with iron, Black Cohosh, Black Haw, Cotton-root bark, Digitalis, Ergot, Fleabane oil, Indian Cannabis, Ipecac, Iron, Kino, Potassium bromide, Quinine, Rhatany, Saraca Indica, Witch-hazel.
- MYALGIA** (Muscular pain, cramp).—Ammonium chloride, Arnica (internally and externally), Belladonna with opium (externally), Black Cohosh, Coca, Chloroform liniment, Jamaica Dogwood, Prickly Ash.
- NERVOUS EXHAUSTION**.—Aromatic spirits of ammonia, Caffeine citrate, Coca, Hypophosphites, Liquid Acid Phosphate, Strychnine, Valerian, Zinc oxide.
- NEURALGIA**.—Aconite (internally and externally), Alligator pear (intercostal neuralgia), American Hellebore, Arsenic, Belladonna, Black Cohosh, Calabar bean (neuralgia of the eye-ball), California Laurel, Chamomile, Gelsemium, Horse-chestnut, Ignatia, Indian Cannabis, Jamaica Dogwood, Menthol, Osmic Acid, Phosphorus, Pink-root, Quinine, Saw Palmetto, Scullcap, Strychnine, Tonga, Zinc valerianate.
- NIGHT-SWEATS**.—Belladonna, Crawley root, Duboisia, Ergot, White Agaric.
- NYMPHOMANIA**.—Bittersweet, Black Willow, Bromides, Camphor monobromated, Henbane, Hops, Stramonium.
- OBESITY**.—Alkaline mineral waters, Bladder-wrack, Potassium permanganate, Avoidance of starches and sugars.
- ODONTALGIA** (Toothache).—Aconite, California Laurel, Cocaine, Coto, Etherial solution of tannin, Jamaica Dogwood.
- PARALYSIS** (Loss of the power of voluntary motion or sensation).—Belladonna, Calabar bean, Cantharides, Colocynth, Conium, Ergot, Fish-berries, Gelsemium, Henbane, Hoang-nan, Ignatia, Indian Cannabis, Mistletoe, Poison Oak, Phosphorus with cod-liver oil, Strychnine.
- PERTUSSIS** (Whooping-cough).—Ammonium bromide, Asafetida, Belladonna, Button-bush, Cochineal, Conium, Duboisia, Gelsemium, Horse-chestnut, Indian Turnip, Jamaica Dogwood, Lobelia, Pulsatilla, Saw Palmetto, Skunk Cabbage, Stramonium.
- PHARYNGITIS** (Inflammation of the pharynx).—Ammonium chloride, Black Cohosh, Glycerin pure or with Tannin (externally), Golden-seal (internally and externally), Inhalation (see Inhalants page 211), Pomegranate bark (gargle), Potassium chlorate, Prickly Ash, Virginia Stone-crop, Zinc sulphate (gargle).
- PHTHISIS** (Consumption).—Arsenic, Belladonna, Black Cohosh, Cod-liver oil, Eulachon oil, Inhalation (see Inhalants page 211), Iodine (internally and externally, Iron iodide, Mullein, Quebracho, Sundew, Water Fennel).
- PLEURITIS** (Pleurisy).—Abscess root, Aconite, American Hellebore, Crawley root, Digitalis, Iodine, Jaborandi, Opium, Pleurisy root, Potassium iodide, Squill, Sinapisms, White Bryony.
- PNEUMONIA** (Inflammation of lung substance).—Aconite, American Hellebore, Ammonium carbonate, Antipyrin (for pyrexia), Blood-root, Digitalis, Iodine and iodides, Ipecac, Opium, Phosphorus, Senega, Serpentina.

POLYURIA.—See Diabetes insipidus.

PSORIASIS (A cutaneous scaly disease of rough amorphous scales, generally chapped or fissured).—Arsenic, Berberis aquifolium, Burdock seed, Goa powder, Phosphorus, Tar.

PTYALISM (Salivation).—Belladonna, Iodine tincture (gargle) Sulphuric acid dilute (gargle), Tannin (gargle), Potassium chlorate.

PUERPERAL CONVULSIONS.—Aconite, American Hellebore, Amyl nitrite, Belladonna, Chloral, Chloroform inhalations.

PUERPERAL FEVER.—Aconite, American Hellebore, Opium, Stramonium, Warburg's Tincture.

PUERPERAL MANIA.—Aconite, Black Cohosh, Iron chloride tincture, Jamaica Dogwood, Potassium bromide, Stramonium.

PYÆMIA (Septicæmia, or blood-poisoning).—Potassium permanganate, Quinine, Salicin, Salicylic acid, Stimulants.

PYROSIS (Waterbrash; a hot sensation in the stomach with eruptions of an acid, burning liquid causing distressing sensations in the parts over which it passes).—Ajowan, Bismuth with Magnesia, Capsicum, Pulsatilla, Strychnine.

RACHITIS ("Rickets").—Cod-liver oil, Iron iodide, Lime-water, Sweet Fern, Phosphatic foods.

REMITTENT FEVER.—Aconite, Alstonia constricta, American Centaury, Black Cohosh, Button-bush, Canadian Hemp, Gelsemium, Morphine (hypodermically), Quinine, Serpentina, Warburg's Tincture.

RHEUMATIC ARTHRITIS (An affection characterized by pain, stiffness, and deformity of the joints, associated with deposition of new bone around them).—Aconite (internally and externally), Arsenic, Black Cohosh, Cod-liver oil, Colchicum, Guaiac, Iodine and iodides, Lithium salts, Manaca.

RHEUMATISM (Inflammation of the fibrous tissue).—Aconite, American Hellebore, Arnica, Belladonna, Bitter-sweet, Black Cohosh, Blue Cohosh, Buchu, Burdock, California Laurel, Chaulmoogra oil, Cod-liver oil (internally and externally), Colchicum, Conium, Coto, Digitalis, Guaiac, Hydriodic acid (syrup), Iron chloride tincture, Jaborandi, Kava Kava, Lithium salts, Manaca, Mezereum, Nettle, Poke-root, Poison Oak, Pongamia glabra oil, Potassium bicarbonate, Potassium iodide, Prickly Ash, Pulsatilla, Salicin, Salicylic acid and salicylates, Soapwort, Yellow Parilla, White Bryony.

SCABIES (The Itch).—Balsam of Peru, Ichthyol, Stavesacre, Sulphur, Sulphurous acid (gaseous), Tar ointment.

SCARLATINA (Scarlet fever).—Aconite, Ammonium carbonate, Chlorinated soda solution in "sloughing," Digitalis, Iron chloride tincture, Potassium chlorate, Potassium iodide, Potassium permanganate (internally and also locally to throat), Quinine, Sodium benzoate, Sulphurous acid inhalation.

SCIATICA (Pain which follows the course of the sciatic nerve).—Aconite, Belladonna (or preferably Atropine with Morphine hypodermically), Black Cohosh, Cocaine hypodermically, Guaiacum, Iodides, Osmic acid, Poison Oak, Tonga. Aquapuncture, Acupuncture.

SCROFULOSIS (Scrofula; a condition of constitutional debility with a tendency to ulcerative diseases).—American Sarsaparilla, Bayberry, Berberis aquifolium, Bittersweet, Black Walnut, Burdock, Button-snake root, Carpenter's Square, Carrion flower, Cod-liver oil, Eulachon oil, False Bittersweet, Hoang-nan, Iodides, Mezereum, Mountain Laurel, Poke-root, Prickly Ash bark, Red Clover tops, Sarsaparilla, Soapwort, Stillingia, Tag Alder, Turkey-corn, Twin-leaf, Water Eryngo, White Pond-lily, Yellow Dock, Yellow Parilla.

SPERMATORRHŒA (Involuntary emissions of spermatic fluid).—Belladonna, Black Cohosh, Black Willow, Cantharides, Conium, Damiana, Digitalis, Gelsemium, Helonias, Hops, Hygienic measures, Hypophosphites, Kava Kava, Potassium bromide, Star grass, Strychnine.

S P L E E N , ENLARGED.—Bearsfoot, Belladonna, Bitter Bugleweed, Ergot, Grindelia squarrosa, Iodine, Quinine.

SPRAINS.—Aconite, Ammonium chloride, Arnica, Mugwort, Witch-hazel (distilled extract).

STERILITY.—Damiana, Gold chloride, Hypophosphites. Os dilatation.

STOMATITIS (Inflammation of the mouth) - Bismuth subnitrate, Blackberry, Cranesbill, Eucalyptus, Golden-seal, Potassium chlorate in solution, Khatany, Compressed Tablets; Borax, Potassium chlorate, Potassium chlorate and borax.

SYCOSIS (Barber's Itch).—Boracic acid locally, Donovan's solution internally, Mercurial ointment or oleate externally, Sodium sulphite lotion.

SYNCOPE (Diminution or suspension of the heart's action, accompanied with cessation of respiration, and loss of sensation).—Ammonia (internally or inhaled), Amyl nitrite, Atropine hypodermically, Stimulants.

- SYPHILIS** (A virulent and specific venereal affection, dependent upon a distinct morbid poison; contagious).—Bamboo Brier, Berberis aquifolium, Burdock, Carnauba, Caroba, Cascara amarga, Iodides, Manaca, Mercurials, Mountain Laurel, Poke-root, Pulsatilla, Stillingia, Syrup Trifolium Comp. (see page 150), White Indian-hemp, Yellow Dock, Yellow Parilla, Yerba buena.
- TETANUS** (Characterized by the permanent contraction of some or all of the muscles without alteration or relaxation).—Aconite, Belladonna, Calabar bean, Chloral, Curare, Gelsemium, Henbane, Hoang-nan, Indian Cannabis, Potassium bromide, Strychnine.
- TIC DOULOUREUX** (Neuralgia of the branches of the fifth pair of nerves).—Aconite, Black Cohosh, Gelsemium, Phosphorus, Pink-root, Tonga (also see neuralgia).
- TONSILITIS** (Inflammation of the tonsils).—Aconite, Gargles of alum, tannin or capsicum, Guaiac, Silver nitrate (locally), Compressed tablets of Animonium chloride or Potassium chlorate.
- TYPANITES** (Inflammation of the lining membrane of the tympanum).—Asafetida, Capsicum, Fish-berries, Ginger, Opium, Turpentine.
- TYPHOID FEVER**.—Acetanilid or antipyrin for reducing hyperpyrexia, American Hellebore, Arnica, Belladonna, Boneset, Cinchona, Digitalis, Ergot, Golden-seal (for excessive sweating), Ipecac, Nitro-hydrochloric acid, Potassium bromide, Serpentina, Turpentine, Wild Indigo. Scalded-milk diet.
- ULCERS**.—Systemic remedies; Arsenic, Belladonna, Golden-seal, Syrup Trifolium Comp. (see page 150.) Local remedies; Chlorinated solution, Poke-root, Potassium chlorate, Witch-Hazel (distilled extract), Zinc sulphate, Zinc chloride (a very strong caustic).
- URÆMIA** (A condition of the blood in which it contains urea).—Colchicum, Digitalis, Hydragogue and saline cathartics, Manaca, Diaphoresis, Transfusion.
- URTICARIA** (Hives; distinguished by those elevations of the cuticle called wheals).—Acetic or Nitric acid largely diluted as a lotion, Benzoin Compound Tincture externally, Sodium salicylate, Strychnine.
- UVULA RELAXED**.—Bistort, Cranesbill, etc. diluted as gargles. Lozenges; Alum and Kino, Rose-leaf and Tannin. Amputation of part.
- VARICOSIS** (Varicose or enlarged and tortuous condition of a vein or veins).—Bandaing, Ergot, Witch Hazel (distilled extract).
- VARIOLA** (Small-pox).—Systemic remedies; Belladonna, Black Cohosh, Quinine. Ectrotic measures; Tincture of Iodine, Mercurial ointment, Collodion, Silver nitrate solution (20 grains to the fluidounce).
- VERTIGO** (An affection in which all objects appear to be turning round, and the individual affected finds difficulty in maintaining an erect posture).—Bromides, Cod-liver oil, Digitalis, Fish-berries, Skunk Cabbage, Strychnine with iron.
- VOMITING**.—Aconite, Adrue, Arsenic, Capsicum, Cerium oxalate, Golden-seal, Hydrocyanic acid, Quinine, Stavesacre.

OBSTETRIC CALENDAR.

NINE CALENDAR MONTHS.			TEN LUNAR MONTHS.	
FROM.	TO	DAYS.	TO	DAYS.
January..... 1	September..... 30	273	October..... 7	280
February..... 1	October..... 31	273	November..... 7	280
March..... 1	November..... 30	275	December..... 5	280
April..... 1	December..... 31	275	January..... 5	280
May..... 1	January..... 31	276	February..... 4	280
June..... 1	February..... 28	273	March..... 7	280
July..... 1	March..... 31	274	April..... 6	280
August..... 1	April..... 30	273	May..... 7	280
September..... 1	May..... 31	273	June..... 7	280
October..... 1	June..... 30	273	July..... 7	280
November..... 1	July..... 31	273	August..... 7	280
December..... 1	August..... 31	274	September..... 6	280

The above calendar may be read as follows: a patient has ceased to menstruate on the 1st of July; her confinement may be expected at soonest, about the 31st of March, [the end of nine calendar months] or, at latest, on the 6th of April [the end of ten lunar months]. Another has ceased to menstruate on the 20th of January, her confinement may be expected on the 30th of September, plus 20 days, [the end of nine calendar months], at soonest; or on the 7th of October, plus 20 days, [the end of ten lunar months], at latest.

POISONS AND ANTIDOTES.

Stomach-pump to be employed when available.

POISON.	ANTIDOTE.
ACIDS, MINERAL.....	{ Chalk; magnesia (plaster off wall in emergency); solution carbonate of soda; emollient drinks; fixed oils.
ACONITE.....	{ Emetics; stimulants externally and internally. atropine, hypodermically, or fluid extract belladonna by mouth.
ANTIMONY TARTRATE.....	{ Vegetable acids; tannic acid; catechu; strong coffee.
ARSENIC.....	{ Moist peroxide of iron (perchloride of iron and calcined magnesia); dialyzed iron; charcoal; ammonia; artificial respiration; cold affusion; stimulants.
ATROPINE.....	See BELLADONNA.
ARGENTI NIT. (silver nitrate).....	Solution of common salt in demulcent fluid.
BELLADONNA.....	{ Emetics; cold affusion, ammonia and other stimulants externally and internally; pilocarpine; opium.
CANNABIS INDICA.....	See INDIAN CANNABIS.
CANTHARIDES.....	{ Emetics; emollient drinks; opiates by mouth and rectum.
CARBOLIC ACID.....	{ Fixed oils; white of egg; stimulants; amyl nitrite inhalations; atropine hypodermically.
CHLOROFORM INHALATION.....	{ Artificial respiration; galvanism (interrupted current from stomach to larynx); amyl nitrite inhalation. Keep head lower than the rest of the body.
COLCHICUM.....	{ Emetics, followed by demulcent drinks and tannin frequently, or very strong tea. If coma supervene, brandy; ammonia; coffee.
CONIUM.....	{ Emetics; tannin; stimulants externally and internally; persistent artificial respiration.
CORROSIVE SUBLIMATE.....	{ Emetics; albumen (white of 1 egg to 4 grs. of sublimate); flour; milk.
CROTON OIL.....	{ Emetics, followed by mucilaginous fluids, containing opium.
COPPER SULPH.....	Albumen or white of egg; milk.
DIGITALIS.....	{ Recumbent posture after emetics; tannin; stimulants internally and externally.
ELATERIUM.....	{ Demulcent drinks and enemata; small doses of opium, and the warm bath.
FOXGLOVE.....	See DIGITALIS.
HYDROCYANIC ACID.....	{ Artificial respiration, with cold affusion; atropine hypodermically; interrupted galvanic current to heart region.
HYOSCYAMUS.....	See BELLADONNA.
INDIAN CANNABIS.....	{ Faradization of respiratory muscles and strychnine cautiously. Not often fatal.
IODINE.....	{ Emetics and demulcent drinks,—starch or flour diffused in water; amyl nitrite inhalations.

POISON.	ANTIDOTE.
LEAD SALTS.....	{ Sulphate or phosphate of soda; dilute sulphuric acid; epsom salts; emetics; afterwards opium and milk.
LOBELIA....	{ Tannin; stimulants externally and internally. Recumbent position.
MORPHINE.....	Same as OPIUM.
NUX VOMICA.....	See STRYCHNINE.
OPIUM.....	{ Emetic; stimulants externally and internally; strong coffee injection; cold affusion; galvanism; amyl nitrite inhalations; artificial respiration; forced exercise; atropine hypodermically.
PHOSPHORUS	{ Magnesia; emetics; epsom salts as a purgative.
POTASH AND SODA SALTS.....	{ Dilute acetic acid; vegetable acids; fixed oils; demulcents.
STRAMONIUM.....	See BELLADONNA.
STRYCHNINE.....	{ Iodine, followed by emetics; chloroform inhalations; belladonna; amyl nitrite inhalations; tobacco; chloral and potassium bromide.
TOBACCO.....	{ Emetic; stimulants externally and internally; strychnine.
ZINC SALTS	{ Carbonate of soda; emetics, afterwards tannin and warm demulcent drinks. Morphine.

ADDENDA AND ERRATA.

ADDENDA.

ALKANET ROOT.—(See page 5.) We are now prepared to supply this root compressed in pound packages for the convenience of retailing purposes.

BELLADONNA.—(See page 18.)

Powdered Extract Belladonna root.—One part equals 5 of the root. Of the same strength as the solid extract of the root; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Belladonna root. Dose 1-8 to 1-2 grain (0.008 to 0.03 Gm.).

Solid Extract Belladonna root.—One part equals 5 of the root. Dose, 1-8 to 1-2 grain (0.008 to 0.03 Gm.).

BLACK WILLOW.—(See page 28.) European experimenters* have lately reported most favorably on the use of Black Willow in cases of ovarian hyperæsthesia, uterine neuralgia, spermatorrhœa and nocturnal pollution. Claims for similar efficacy in the use of Black Willow buds have long been known to the medical profession on this side of the Atlantic, but since these clinical reports above referred to have been based on results obtained from use of the bark, we have in addition to the fluid extract of the buds, placed upon our list:

Fluid Extract Black Willow bark.—Not miscible with water. Claimed to be equal, if not superior, to potassium bromide as a powerful sexual sedative, but without the depressing effects which accompany the exhibition of the latter drug. Dose, 1-2 to 1 fluidrachm (2 to 4 C.c.).

BLOOD-ROOT.—(See page 29.)

Powdered Extract Blood-root.—Of the same strength as the solid extract, prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Blood-root. Dose, 1 to 5 grains (0.06 to 0.13 Gm.).

CASCARA SAGRADA.—(See page 44.) Having been instrumental in the introduction of Cascara Sagrada and identified with it since its advent in 1877, we long since recognized the necessity of a more thorough acquaintance with the true constitution of this valuable remedy. It is to be regretted that the many insidious attacks by interested parties upon its therapeutic merits, and upon our preparations of it, have raised a question of doubt as to the actual identity and properties of the drug in the minds of many physicians unfamiliar with its use. The statement has been made that preparations alleged to be from Cascara Sagrada are really compounds of Rhamnus Frangula, aloes, nux vomica and other improper additions. Cases of miscarriage, attributable to the use of certain preparations in the market purporting to be from Cascara Sagrada, have been clearly traced to the action of aloes. That certain preparations of Cascara in the market do contain aloes, we are in a position to prove. We are confident that our reputation is a sufficient guarantee that preparations which we offer are free from any foreign admixture. With this by way of apology, we have no hesitancy in asking the attention of physicians to what we further state upon the subject.

During the past year we have devoted our best energies towards obtaining a satisfactory solution of the peculiarities and sometimes contradictory behavior of this drug, both in the hands of the pharmacist and physician. This has induced a careful reconsideration and revision of all previous work, and we are pleased to say that we have succeeded in our object after an expenditure of considerable time and arduous labor. Our work has been an interesting one, and the results obtained are a matter of surprise, and in offering preparations manufactured after our new formula to the medical profession, we have to say that the principal object which has been secured is the removal of the extreme bitterness, which has been a very objectionable feature as urged against our regular fluid extract Cascara Sagrada. Our former products were bitter, changeable—owing to the peculiar characteristics of the bark itself—and but partially soluble in water; while occasionally, in individual cases, they would vary somewhat in therapeutic action.

Inasmuch as preparations in which the bitterness has been developed may at times be preferable or desirable, we shall continue to manufacture preparations of Cascara Sagrada by our old formula, which preparations can always be obtained from us or from the trade generally by proper specification. But that there may be no misunderstanding or confusion we shall designate our improved preparations, "manufactured after the formula of 1887."

The announcement of the detail of our work, and the results thereof, appeared

* Drs. J. Hutchinson (British Medical Journal, July 30,) and E. Hurry Fenwick (Lancet, September 21, 1887).

CASCARA SAGRADA—Continued.

in the *American Journal of Pharmacy*, of Philadelphia, February number of the current year (1888), and to those who are interested in the progress of pharmacy, or in the study of Cascara Sagrada, we will on request gladly mail, without charge, a reprint of this article. We now offer these improved preparations, formulæ of 1887, as follows:

Fluid Extract Cascara Sagrada (Formula of 1887).—Permanent and will not precipitate on standing, it is free from inert matter, not bitter, entirely soluble in water, and is a true representative of all the valuable constituents of the bark. Dose, same as fl. ext. old formula; see page 45.

Solid Extract Cascara Sagrada (Formula of 1887).—Permanent; will make a pill-mass which does not soften or decompose on keeping, and has the advantage of being readily soluble in water and in the liquids of the gastric juice. When made into pills alone, the only excipient needed is a small quantity of powdered marsh-mallow. Tragacanth, recommended by some, should be avoided, as pills made with it harden gradually, with impairment of their solubility. Dose, same as solid extract old formula, see page 46.

Cascarin (Concentration; in *Scale Form*; Formula of 1887).—This is not hygroscopic, is readily reduced to powder, almost tasteless, soluble in water and contains only the really active constituents of the plant. Dose, as a laxative, 1 to 3 grs. (0.06 to 0.2 Gm.); as a cathartic, 3 to 8 grs. (0.2 to 0.5 Gm.).

In addition to the above we have recently placed upon our list the following prepared from solid extract of our regular stock:

Capsule, Cascara Sagrada Solid Extract, 1 gr., Castor oil, q.s. 10 minims. In soft gelatin.

Pill, Cascara Sagrada Solid Extract, 1 gr.

DEWBERRY.—*Rubus canadensis*, Linné. (Root.) *Synonym*.—Low Blackberry. *Properties*.—Tonic and strongly astringent. May be employed with advantage wherever a vegetable astringent is indicated, but chiefly in summer complaint and infantile diarrhœa. It very closely resembles Blackberry (*q. v.*) in its medicinal properties. •

PREPARATION AND FORMULA.

Fluid Extract Dewberry root.—Not miscible with water. Dose, 1-4 to 1 fluidrachm (1 to 4 C.c.).

Syrup Dewberry root.—The popular form of its exhibition.

Fluid Extract Dewberry root, P., D. & Co.'s.....4 fluidounces.

Simple syrup.....12 fluidounces.

Dose, 1 to 4 fluidrachms (4 to 15 C.c.).

GELSEMIUM.—(See page 84.)

Powdered Extract Gelsemium.—Of the same strength as the solid extract; prepared by evaporating, at a low temperature, the solid extract, and replacing the moisture with powdered Gelsemium. Dose, 1-5 to 2 grains (0.013 to 0.13 Gm.).

GOLDEN-SEAL.—(See page 87.)

Berberine acid sulphate (crystals).—This has been inadvertently omitted from the enumeration of the active principles of Golden-seal which are upon our list. It is not so soluble a salt as the neutral sulphate, but is otherwise of the same medicinal value. Solubility, 1:75. Dose, 1 to 4 grains (0.065 to 0.26 Gm.).

JAMBUL.—(See page 106.) Since recent clinical reports have conclusively proved that the bark is much inferior to the seed as a curative agent in diabetes mellitus, we have withdrawn it from the market, and now list JAMBUL SEED only with the following preparation therefrom:

Fluid Extract Jambul seed.—Miscible with water. Dose, 5 to 10 minims (0.3 to 0.65 C.c.).

JURUBEBA.—*Solanum paniculatum*, Linné. (Root.) *Synonyms*.—Juuna, Jupeba pill. *Properties*.—Much valued in Brazil, its native country, as a remedy in all cases of obstruction of the abdominal viscera, particularly of the liver, and in vesical catarrh. It is used largely, and indeed its chief use is, in constipation. In larger doses it is employed as an active laxative, and as a diuretic and hydragogue cathartic in jaundice and dropsical affections. In small doses it is claimed to be tonic, alterative, antibleorrhagic and antisyphilitic. Combined with iron, Brazilians call it the "Tónico por excellencia."

PREPARATION.

Fluid Extract Jurubeba.—Not miscible with water without some precipitation. Dose, 5 to 30 minims (0.3 to 2 C.c.).

LIQUOR SEDANS.—Each fluidounce represents: Black Haw and Golden-seal, 60 grs. each; Jamaica dogwood, 30 grs., combined with aromatics q. s. A utero-ovarian sedative and anodyne. A scientific substitute for certain copyrighted preparations in the market. Dose, 1 to 2 fluidrachms (4 to 8 C.c.)

PEPSIN.—(See page 138.) We have been engaged for some time in elaborating an improved process of manufacturing pepsin. The result of our experiments now enables us to obtain this digestive ferment of a very high degree of concentration. We have recently placed upon our list:

Glycerole Pepsin, concentrated.—Each fluidounce represents 960 grains Pepsin sacch., U. S. P. Each minim will dissolve 150 grains of coagulated egg-albumen in accordance with the modification of the official test as fully described in our circular on Pepsin. It is thus twice the strength of Pepsin sacch., U. S. P., and about 50 times the strength of Liquid Pepsin, U. S. P. This is a very superior preparation, presenting the ferment of the gastric juice in a permanent and highly concentrated form. It is especially adapted for making extemporaneously weaker solutions, such as elixirs, wines, etc.

PICHI.—(See page 140.)

Capsule, Pichi Solid Extract, 5 grs., in soft gelatin.

Solid Extract Pichi.—One part equals 5 of drug. Dose, 2 to 10 grains (0.13 to 0.65 Gm.).

SALICYLIC ACID COMP. ELIXIR.—Each fluidounce represents: Salicylic acid, 40 grs.; Black Cohosh, 15 grs.; Gelsemium, 15 grs.; Potassium iodide, 4 grs. An efficient and convenient compound for many cases of rheumatism, gout, myalgia, etc. Each remedy well-known to be effective for these indications when taken singly has its powers intensified by the combination. Dose, 1 to 2 fluidrachms (4 to 8 C.c.).

TERPIN HYDRATE.—(See page 183.)

Pill, Terpin Hydrate, 2 grs., Sugar and gelatin coated.

"WARBURG'S TINCTURE" without aloes.

Pill, Warburg's Tincture, each pill representing 1 fluidrachm of the tincture without aloes. Sugar and gelatin-coated.

ERRATA.

- p. 5. **AJOWAN:** read *Carum copticum*, *Bentham*, instead of *Ptychotis Ajwan*, *DeCandolle*.
- p. 55. **CLOVES:** read *Eugenia caryophyllata*, *Thunberg*, instead of *Caryophyllus aromaticus*, *Linne*.
- p. 87. **GINSENG:** read *Aralia quinquefolia*, *Decaisne & Planchon*, instead of *Panax quinquefolium*, *Linne*.
- p. 95. **HOANG NAN:** read *Strychnos malaccensis*, *Bentham*, instead of *S. Gaultheriana*, *Pierre*.
- p. 207. Elixir formula No. 80; read Cinchonine sulphate 4 grains, instead of Cinchona bark 4 grains.
- p. 208. Elixir formula No. 88; read Strychnine 4-120 grain, instead of Strychnine 4-100 grain.
- " " Elixir formula No. 111; read Quinine sulphate 2 grains, instead of Quinine sulphate 8 grains.
- " " Elixir formula Nos. 113 and 115; read saccharated pancreatin, instead of pancreatin.

INDEX.

For the sake of brevity, pharmaceutical preparations of articles of organic materia medica, except in a few special instances, are not here indicated—these will easily be found either by referring to the group class of preparations, or under the name of the drug from which it is prepared. References, however, are given in the case of articles of an inorganic nature, since they are not treated of individually in the foregoing pages.

Cross-references are withheld wherever it is considered no inconvenience will be occasioned thereby. ACIDS, OILS, etc., are indexed under the adjective or qualifying part of their names—thus, Oil of Peppermint is placed under Peppermint only, Carbolic acid under Carbolic only; SALTS are only placed once, under the basic name—thus, Subiodide of Bismuth is indexed under Bismuth. The U. S. Pharmacopoeial names are not generally given, especially when they only differ in a Latinized termination from the name by which they are already indexed.

This index aims to be a complete Botanical Reference to all the plants mentioned in this work, or, in other words, of all those articles of organic materia medica handled by Messrs. Parke, Davis & Co. The Latin or botanical name of a plant may be found by looking up the vernacular or English name, or, *vice versa*, the popular name may be found by referring to the botanical one. If an article is sought for under an English synonym it is referred immediately to that name in the index to which the botanical name is attached. Botanical synonyms are in *italics*.

	PAGE.		PAGE.
Abbreviations of Latin used in pre-		Ail.....	Garlic 83
scriptions.....	262	Ailantus glandulosa.....	5
Abdominal supporter.....	238, 246	Ajava.....	Ajowan 5
Abies balsamea.....	Balsam fir 15	Ajowan.....	Carum copticum, 5, 276
<i>canadensis</i>	93	Ajwain.....	Ajowan 5
Abrus precatorius.....	Jequirity 106	Albumen, Tests for.....	246
Abscess root.....	Polemonium reptans 1	Alder, black.....	24
Absinthe commune.....	Wormwood 197	buckthorn... Buckthorn bark 35	
Absinthium.....	Wormwood 197	tag.....	177
Abstracts, see respective drugs.		red.....	Tag alder 177
Acacia Catechu.....	Catechu 47	smooth.....	Tag alder 177
Acer rubrum.....	Red maple 150	spotted.....	Witch-hazel 197
Acetanilid.....	246	Aletrin.....	Star-grass 170
Pills.....	213	Aletris farinosa.....	Star-grass 170
Acetophenone.....	248	Alimentation, rectal.....	260
Achillea Millefolium.....	Yarrow 198	Alkaloids, oleates of in solution.....	225
Acid Phosphate, Liquid.....	241	rare.....	246
Aconapellin.....	Aconite 2	Alkanet.....	Alkanna tinctoria, 5, 274
Aconite.....	Aconitum Napellus 1	Alkanna tinctoria.....	Alkanet 5
Aconitine.....	Aconite 2	Alkannin.....	Alkanet 5
Aconitum Napellus.....	Aconite 1	Alligator pear.....	Persea gratissima 6
Acore odorant.....	Sweet flag 176	Allium sativum.....	Garlic 83
Acorus Calamus.....	Sweet flag 176	Allspice.....	Pimento 141
Actæa alba.....	White cohosh 190	Florida.....	79
<i>racemosa</i>	25	wild.....	Spice-bush 168
<i>rubra</i>	Red cohosh 150	Alnuin.....	Tag alder 177
<i>spicata</i>	150	Alnus serrulata.....	Tag alder 177
Actæa à grappes.....	Black cohosh 25	Aloe Socotrina.....	Aloes 6
Adder's Tongue.....	Erythronium americanum 4	Aloes.....	Aloe Socotrina 6
Adhesive plaster.....	236	Aloin.....	Aloes 6
Adiantum pedatum.....	Maiden-hair 117	Alpinia Galanga.....	Galangal 82
Adonidin.....	Adonis vernalis 4	Alstonia constricta.....	7
Adonis vernalis.....	4	scholaris.....	Dita 71
Adrue.....	Cyperus articulatus 4	Alterative Compound, Bamboo brier, 16	
Ægle marmelos.....	Bael 15	Althæa officinalis.....	Marsh-mallow 120
Æsculus glabra.....	Buckeye 24	rosea.....	Hollyhock 95
Hippocastanum.....		Alum, Lozenges.....	233
Horse-chestnut.....	97	Pills.....	218, 225
African saffron.....	Safflower 155	and kino.....	Kino 109
Agaric blanc.....	White agaric 189	and rose-leaf.....	Rose 154
fly.....	80	Rockbridge. Effervescents.....	210
larch.....	White agaric 189	root.....	Heuchera americana 8
white.....	189	Aluminium acetate solution, 3%.....	241
Agaricin.....	White agaric 189	hydrate, Pills.....	214
Agaricus muscarius.....	Fly agaric 80	metallic, Pills.....	214
Agrimonia Eupatoria.....	Agrimony 5	oleate.....	235
Agrimony.....	Agrimonia Eupatoria 5	Alveloz milk, Euphorbia heterodoxa.....	8
Aguacate.....	Alligator pear 6	Ambrosia artemisiæfolia.....	Ragweed 149
Ague root.....	Star-grass 170	American ash.....	White ash 190
weed.....	Grindelia squarrosa 90	aspens.....	White poplar 192
		beech.....	Beech 18

PAGE.		PAGE.	
American blue vervain	8	Anise	Pimpinella 11
Verbena hastata	8	Aniseed	Mayweed 122
centaury,		Anisum	Anise 11
Sabbatia angularis	8	Anisum vulgare	11
columbo,		Anodyne collodion	237
Fraseria carolinensis	9	plaster	236
gentian	Blue gentian 32	Anserine anthelmintique,	
greek valerian,		Jerusalem oak	107
Abscess root	1	Anthemis Cotula	Mayweed 122
hellebore,		nobilis	Chamomile 49
Veratrum viride	9	Anti-emetic root	Adrue 4
ipecac	Indian physic 101	Antifebrin	246
ivy	Am-	Antimony and Potassium tartrate,	
pelopsis quinquefolia	10	Glyceroles	211
larch	Tamarac 177	Pills	218, 221
mountain-ash,		Wines	234
Mountain-ash	123	sulph'd, Lozenges	233
saffron	Safflower 155	Pills	215, 216
sarsaparilla,		Tablet triturates	231
Aralia nudicaulis	10	with ipecac and	
spikenard	Spikenard 168	morphine	102, 132
valerian	Ladies' slipper 110	Antipyrin	247
water hemlock,		Antiseptic Cologne	241
Water hemlock	188	Liquid	241
wormseed	Cheno-	Preparations	238, 241
podium anthelminticum	10	Tablets (Bernay's)	231
Amerikanische Buche	Beech 18	Apfelbaumrinde	Apple-tree 12
Sanickel-wurzel,		Apfelsinedschalen	Sweet orange 177
Alum root	8	Apiol	Parsley 136
Amerikanischer Epheu,		Capsules	203, 205
American ivy	10	Apium graveolens	Celery 48
Hundszahn,		petroselinum	136
Adder's Tongue	4	Apocynin	Bitter-root 25
Wurmsamen, Ameri-		Apocynum androsæmifolium,	
can wormseed	10	Bitter-root	23
Amerikanisches Polei	Pennyroyal 137	cannabinum,	
Ammoniac	Dorema Ammoniacum 11	Canadian hemp	40
Ammonium bromide Elixirs	205	Apollinaris	Mineral-water Salts 237
Effervescent	210	Apomorphine muriate	Opium 130
Pills	214	Apple Peru	Stramonium 172
carbazonate, see picrate		tree	Pyrus malus 12
carbonate, Pills	215	Aralia hispida	Dwarf elder 72
chloride, Bronchial		nude, American sarsaparilla,	10
Sedative	131	nudicaulis,	
Compressed		American sarsaparilla	10
tablets	230	quinquefolia	Ginseng, 87, 276
Lozenges	233	racemosa	Spikenard 168
Pills	214, 225	Araroba	Goa-powder 87
Tablet tritur-		Arbor vitæ	Thuya occidentalis 12
ates	231	Arbre à suif	Bayberry 17
glycyrrhizate	Licorice 113	de vie	Arbor vitæ 12
muriate, see chloride		Arbutin	Uva-ursi 184
picrate	247	Arbutus, Trailing	Gravel-plant 89
Pills	214	Archangelica atropurpurea, Angel-	
valerianate, Elixirs	205	ica,	11
Pills	214	officialis	Angelica 11
Amomum Zingiber, Ginger Mexican,	87	Arctostaphylos glauca	Manzanita 120
Ampelopsin	American ivy 10	Uva-ursi	Uva-ursi 184
Ampelopsis quinquefolia, Amer. ivy,	10	Areca	Areca Catechu 12
Amygdalus persica	Peach 136	Catechu	Areca 12
Amyl nitrite	247	Arekanisso	Areca 12
pearls	238	Arisæma Dracontium, Green dragon	90
valerianate	247	triphyllum	Indian turnip 101
Amylo-valerianic ether	247	Aristolochia Serpentaria, Serpentaria	164
Anacyclus Pyrethrum	Pellitory 137	Armoise amère	Wormwood 297
Anagallis arvensis, Scarlet pimpernel	160	Commune	Mugwort 124
Anamirta cocculus	Fish-berries 78	Arnica	Arnica montana 12
Andira Araroba	87	montana	Arnica 12
Andromeda arborea	167	Arnique	Arnica 11
Anemopsis californica	199	Aromatic, Fluid Extract, U. S. P.	13
Anemone Hepatica	Liverwort 115	Wine, U. S. P.	13
Pulsatilla	Pulsatilla 147	Troches	238
Angelica,		Arsenious acid, Elixirs	207, 209
Archangelica atropurpurea	11	Effervescent	210
Angustura	Cusparia trifoliata 11	Pills	214, 215,
		217, 219, 221, 224, 225,	236

	PAGE.		PAGE.
Arsenious acid, Tablet triturates, 231, 232		Balsam gurun.....Gurjun balsam	92
iron, quinine and		of fir.....Balsam fir	15
strychnine.....	130	of Peru, Myroxylon Pereira	15
Arsenic iodide, Pills.....	236	of Tolu.....Tolu	181
Tablet triturates.....	231	poplar.....Populus candicans	15
oleate.....	235	spruce.....Balsam fir	15
Artanthe elongata.....Matico	121	Balsamodendron Katak.....Myrrh	125
Artemisia abrotanum, Southernwood	167	Balsampappel.....Balsam poplar	16
absinthium.....Wormwood	197	Balsamum Dipterocarpi,	
frigida, Mountain sage	124	Gurjun balsam	92
maritima,		Bamboo brier..Smilax Pseudo-China	16
Levant wormseed	112	Banana.....Musa sapientum	16
vulgaris.....Mugwort	124	Baneberry, red.....Red cohosh	150
Arum.....Indian turnip	101	white.....White cohosh	190
Dracontium.....90		Baptisia tinctoria.....Wild indigo	193
esculentum.....Taro flour	179	Baptisin.....Wild indigo	193
triphyllum,		Barberry.....Berberis vulgaris	16
Arisema triphyllum	101	Barbotine.....Levant wormseed	112
Arveloz.....Alveloz milk	8	Bärentraube.....Uva-ursi	184
Asafetida.....Ferula Narthex	14	Barosma betulina.....Buchu	33
Asagrea officinalis.....49		Barosmin.....Buchu	33
Asaret du Canada.....Wild ginger	195	Basil, sweet.....175	
Asarum canadense.....Wild ginger	195	Basilic.....Sweet basil	175
Asclépiade à la Soie.....Silkweed	165	Basilienkraut.....Sweet basil	175
Asclepias Cornuti.....Silkweed	165	Basswood.....Tilia americana	17
curassavica..Blood-flower	29	Bastard saffron.....Safflower	155
incarnata,		Bath-flower.....Beth-root	22
White Indian-hemp	191	Baume de cheval.....Stone-root	172
syriaca.....165		Baumwolle.....Cotton	65
tuberosa.....Pleurisy root	142	Bay.....Bay laurel (2)	17
Asclepidin.....Pleurisy root	143	laurel (1).....Laurus nobilis	17
Ash, American.....White ash	190	(2).....Myrcia acris	17
black.....24		rose.....Willow-herb	196
bitter.....Quassia	147	rum.....Bay laurel (2)	17
mountain.....123		sweet.....Magnolia	117
prickly.....145		tree.....Bay laurel (1)	17
wafer.....186		white.....Magnolia	117
white.....190		wild-rose.....Great laurel	89
Asimina triloba, Papaw (American),	135	Bayberry.....Myrica cerifera	17
Asok.....Saraca indica	157	Baycuro.....Statice brasiliensis	18
Asparagus officinalis.....Asparagus	14	Bearberry.....Uva-ursi	184
Aspen, American.....White poplar	192	Bearsbed.....Haircap moss	92
Asperge.....Asparagus	14	Bearsfoot.....Polymnia Uvedalia	18
Aspidium Filix-mas.....Male-fern	117	Bear'sweed.....Yerba santa	200
Aspidosperma Quebracho,Quebracho	148	Beaver-poison.....Water hemlock	188
Aspidospermine.....Quebracho	148	tree.....Magnolia	117
Atomizing inhaler, Semp's.....243		Bedstraw.....Cleaver's	55
Atropa Belladonna.....Belladonna	18	Beech.....Fagus ferruginea	18
Atropine.....Belladonna	19	-drop.....Epiphegus virginiana	18
sulphate.....Belladonna	19	Beef and wine.....234	
Augentrost.....Eyebright	77	coca and.....234	
Aunée.....Elecampane	73	extract, Malt ext. preparations	212
Aurantii amari cortex, Bitter orange	23	iron and wine.....234	
dulcis cortex, Sweet orange	177	with pepsin.....234	
Aurone des jardins.....Southernwood	167	wine and cinchona.....234	
Austrahan gum-tree,		peptonized extract of.....238, 242	
Eucalyptus globulus	75	powder of.....238, 241	
fever bark,		Beifuss.....Mugwort	124
Alstonia constricta	7	Beinwurz.....Comfrey	61
Ava ava.....Kava kava	108	Belladonna.....Atropa Belladonna, 18,	274
Aveloz.....Alveloz milk	8	Belladonnin.....Belladonna	19
Avens.....Water avens	187	Benedictendistel.....Blessed thistle	59
Avocado pear.....Alligator pear	6	Bengal quince.....Bael	15
Avoirdu pois ounces into Troy ounces,		Benjoin.....Benzoin	21
to convert.....254		Benne.....Sesamum orientale	21
Azedarach.....Pride of China, 146		Benzoe.....Benzoin	21
		Benzoeorbeer.....Spice-bush, 168	
		Benzoin.....Styrax Benzoin	21
		odoriferum.....168	
Bael.....Egle marmelos	15	Berberin.....Barberry	17
Baldrain.....Valerian	185	Berberine.....Barberry	17
Balm.....Lemon balm	112	acid sulphate, Golden-seal	275
of Gilead.....Balsam poplar	16	hydrochlorate.....88	
Balmomy.....Chelone glabra	15	phosphate.....88	
Balsam copaiba.....Copaiba	63	Elixirs.....205	
fir.....Abies balsamea	15		

	PAGE.		PAGE.
Berberine sulphate.....	88	Bladder-wrack....	<i>Fucus vesiculosus</i> 28
Berberis aquifolium.....	22	Blasentang.....	Bladder-wrack 28
<i>vulgaris</i>	Barberry 16	Blazing star.....	<i>Helonias</i> 93
Bergamot, wild.....	193	<i>Stargrass</i>	170
Bernay's antiseptic tablets.....	231	Bl'd cornu.....	<i>Ergot</i> 74
Bertramwurzel.....	Pellitory 137	Blessed-thistle.....	<i>Centaurea benedicta</i> 29
Berufkraut.....	Fleabane 79	Blistering collodion.....	<i>Cantharides</i> 41
Besenginster pfriemenkraut.....	Broom 33	Blood, dried defibrinated.....	243
Betel nut.....	<i>Areca</i> 12	flower, <i>Asclepias curassavica</i>	28
Beth-root.....	<i>Trillium erectum</i> 22	root, <i>Sanguinaria canadensis</i> , 29, 274	
Betonia officinalis.....	Wood betony 197	Blue cardinal-flower, <i>Lobelia syphilitica</i>	30
Betula lenta.....	Black birch 25	cohosh.....	<i>Caulophyllum thalictroides</i> 30
Biacuru.....	<i>Bayacuru</i> 18	flag.....	<i>Iris versicolor</i> 31
Bidens bipinnata.....	Spanish needles 167	gentian.....	<i>Gentiana puberula</i> 32
Biebernell.....	<i>Saxifrage</i> 123	gum-tree.....	<i>Eucalyptus globulus</i> 75
Big-leaved ivy.....	Mountain laurel 123	mass, Pills.....	214
Bilious weed.....	Five-flowered gentian 79	215, 216, 217, 218, 219, 223, 224, 225	
Birch, black.....	Black birch 25	scullcap.....	Scullcap 160
cherry.....	Black birch 25	vervain, American blue vervain.....	8
sweet.....	Black birch 25	Blueberry root.....	Blue cohosh 30
Bird pepper.....	Capsicum 42	Blutwurzel.....	Blood-root 29
Birth-root.....	Beth-root 22	Bogbean.....	Buckbean 34
Bishop's weed.....	Ajowan 5	Bois amer.....	<i>Quassia</i> 147
Bismuth and ammonium citrate, Elixirs.....	205, 206, 207, 208, 209	Boldo.....	<i>Boldoa fragrans</i> 32
Effervescents.....	210	Boldoa fragrans.....	<i>Boldo</i> 32
Glycerols.....	211	<i>Boletus laricis</i>	189
Malt extract preparations.....	213	Bone, Powdered.....	238, 242
Wines.....	234	Boneset.....	<i>Eupatorium perfoliatum</i> 32
oleate.....	235	Borage.....	<i>Borago officinalis</i> 32
subcarbonate, Pills.....	215, 21	<i>Borago officinalis</i>	Borage 32
Tablet triturates.....	131	Borax, Compressed tablets.....	230
subiodide.....	247	Boroglyceride.....	238, 247
subnitrate, Lozenges.....	223	Bou-néfa.....	<i>Thapsia garganica</i> 190
Pills.....	215, 220, 221	Bourdaine.....	Buckthorn bark 35
Tablet triturates.....	231	Boxberry.....	Wintergreen 196
Bistort.....	<i>Polygonum bistorta</i> 23	Boxwood.....	Dogwood 71
Bitter apple.....	<i>Colocynth</i> 59	Brayera anthelmintica.....	Koosso 109
ash.....	<i>Quassia</i> 147	Brazilian cocoa.....	Guarana 91
bugle.....	Bitter bugleweed 23	Brier, bamboo.....	16
bugleweed, <i>Lycopus europæus</i>	23	Bristly sarsaparilla.....	Dwarf elder 72
clover.....	American centaury 8	British Pharmacopœia, Pills of.....	226
cucumber.....	<i>Colocynth</i> 59	Bronchial lozenge, cubeb.....	63
orange.....	<i>Citrus vulgaris</i> 23	sedative.....	131, 238
root, <i>Apocynum androsæmifolium</i>	23	Broom.....	<i>Cytisus scoparius</i> 33
weed.....	Fleabane 79	Brucine.....	<i>Nux vomica</i> 128
wood.....	Ragweed 149	Brunfelsia Hopeana.....	Manaca 118
Quassia.....	147	Bryonia, alba.....	White bryony 190
Bitterklee.....	Buckbean 34	Bryony, white.....	190
Bittersüss-Stengel.....	Bittersweet 24	Buchu.....	<i>Barosma betulina</i> 33
Bittersweet.....	<i>Solanum dulcamara</i> 29	Buckbean.....	<i>Menyanthes trifoliata</i> 34
climbing.....	77	Buckeye.....	<i>Esculus glabra</i> 34
false.....	77	Buckhorn.....	Buckhorn brake 34
Black alder.....	<i>Prinos verticillatus</i> 24	brake.....	<i>Osmunda regalis</i> 34
ash.....	<i>Fraxinus sambucifolia</i> 24	Buckthorn bark, <i>Rhamnus Frangula</i>	35
birch.....	<i>Betula lenta</i> 25	berries, <i>Rhamnus cathartica</i>	35
cherry.....	<i>Belladonna</i> 18	Bugleweed.....	<i>Lycopus virginicus</i> 25
cohosh.....	<i>Cimicifuga racemosa</i> 25	bitter.....	23
haw.....	<i>Viburnum prunifolium</i> , 26, 276	Burdock.....	<i>Lappa officinalis</i> 36
hellebore.....	<i>Helleborus niger</i> 27	Burning bush.....	Wahoo 187
Indian hemp, Canadian hemp.....	40	Busserole.....	<i>Uvi-ursi</i> 184
larch.....	<i>Tamarac</i> 177	Butterfly weed.....	<i>Pleurisy root</i> 142
oak.....	<i>Quercus tinctoria</i> 24	Butternut.....	<i>Juglans cinerea</i> 35
pepper.....	<i>Piper nigrum</i> 27	Button-bush.....	<i>Cephalanthus occidentalis</i> 37
root.....	Culver's 68	snake root.....	<i>Liatis spicata</i> 75
snake root.....	Black cohosh 25	wood.....	Water cryngo 188
tang.....	Bladder-wrack 28	Button-bush.....	37
walnut.....	<i>Juglans nigra</i> 28		
willow.....	<i>Salix nigra</i> , 28, 274		
Blackberry.....	<i>Rubus villosus</i> 25		
low.....	<i>Rubus canadensis</i> 275		
		Cachou.....	Catechu 47
		<i>Cactus grandiflorus</i>	136

	PAGE.
Cadmium sulphate.....	247
Caffeine.....Coffee	58
citrate.....Coffee	58
Effervescents.....	210
Caille-lait.....Cleaver's	55
Calabar bean,	
Physostigma venenosum	37
<i>Caladium seguinum</i>	72
Calamus, U. S. P.Sweet flag	176
Calcium carb., Pills.....	214
bromide, Elixirs.....	205
hippurate.....	247
hypophosphite, Glyceroles	211
Syrups.....	228
Malt ext.	
prep's.....	213
iodide, Syrups.....	228
Elixirs.....	205, 207
Malt ext. prepar's	212
lactophosphate, Syrups, 228,	229
Digestive	
tablets.....	230
Lozenges.....	233
Malt ext.	
prep's.....	213
phosphate, Elixirs.....	209
Effervescents.....	210
Syrups.....	229
sulphide, Pills.....	216
Tablet triturates.....	231
Calendula officinalis.....Calendula	38
Calico bush.....Mountain laurel	123
California fever-bush,	
Garrya Fremontii	39
laurel.....Um-	
bellularia californica	39
Calisaya.....Yellow cinchona	52
Calomel, Pills.....214, 215,	
216, 218, 220, 221, 223, 224,	226
Lozenges.....	233
Tablet triturates.....	231
Calumba, U. S. P.Columbo	61
Calycanthus floridus, Florida allspice	79
<i>Calyptanthus jambolana</i>	106
Cambogia, U. S. P.Gamboge	83
Camellia Thea.....Tea	179
Camomile puante.....Mayweed	122
Camphor.....Cinnamomum camphora	39
monobromated.....	39
Canada balsam.....Balsam fir	15
fleabane.....	79
snake-root.....Wild ginger	195
thistle.....Cnicus arvensis	40
Canadian hemp,	
Apocynum cannabinum	40
Canadische cornelrinde..Green osier	90
Hanfwurzel,	
Canadian hemp	40
Haselwurzel, Wild ginger	195
Canadisches sonnenroschen,	
Frostwort	82
Cancer-root.....Beech-drop	18
Candleberry.....Bayberry	17
Candlefish.....Eulachon	76
Canella alba.....Canella	41
Cannabin tannate.....	100
Cannabis Indica, U. S. P.,	
Indian cannabis	100
sativa.....Indian cannabis	100
Cannabinon.....	100
Cantharides.....Cantharis vesicatoria	41
Cantharis vesicatoria.....Cantharides	41
Canutillo.....Ephedra antispyphilica	74
Capillaire.....Maiden-hair	117

	PAGE.
Capsella bursa-pastoris,	
Shepherd's purse	165
Capsicum fastigiatum.....Capsicum	42
Capsules, Capacity of.....	239
Empty.....	238, 239
Capsule-fillers,	
Davenport's Unique	238, 240
Reymond's Patent,	238, 240
Caraway.....Carum Carvi	42
Carbolic acid, Collodions.....	227
Glyceroles.....	211
Inhalants.....	212
Lozenges.....	233
Cardamom.....Elettaria Cardamomum	42
Cardinal-flower, blue.....	20
<i>Carduus marianus</i>	121
Carica Papaya.....	43
Carlsbad, Effervescents.....	210
Mineral-water salts.....	237
Carnauba.....Copernicia cerifera	43
Caroba.....Jacaranda procera	43
Carolina Pink.....Pink-root	141
Carpenter's square,	
Scrophularia nodosa	44
Carrión-flower.....Smilax herbacea	44
Carthamus tinctorius.....Safflower	155
Carum Carvi.....Caraway	42
copticum.....Ajowan, 5,	276
Carya alba.....Shell-bark hickory	164
<i>Caryophyllus aromaticus</i>Cloves	55
Casca-bark.....Sassy-bark	158
Cascara amarga,	
Picramnia (Sp. indet.)	44
cordial.....Cascara sagrada	45
sagrada,	
Rhamnus Purshiana, 44,	274
Cascarilla.....Croton Eluteria	46
Cassia.....Cinnamomum (Sp. indet.)	46
acutifolia.....Senna	162
Castanea americana.....Chestnut	50
vesca.....Castanea vulgaris	50
Castor oil.....Castor-oil plant	47
plant.....Ricinus communis	47
Cataire.....Catnep	48
Catechu.....Acacia Catechu	47
Catfoot.....Ground ivy	91
Cathartic compound, Elixirs.....	206
Catmint.....Catnep	48
Catnep.....Nepeta cataria	48
Caulophyllin.....Blue cohosh	31
Caulophyllum thalictroides,	
Blue cohosh	30
Cayenne pepper.....Capsicum	42
Ceanothe.....New Jersey tea	126
Ceanothus americanus,	
New Jersey tea	126
Cedron seed.....Simaba Cedron	48
Celandine, garden.....	83
great.....Garden celandine	83
wild.....Jewel-weed	107
Celastrus scandens, False bitter-sweet	77
Celery.....Apium graveolens	48
Céline.....Lemon balm	112
Centaurea benedicta.....Blessed thistle	29
Centauree.....European centaury	76
Centaury, red.....American centaury	8
Centigrade and Fahrenheit Scales.....	255
Cephaelis Ipecacuanha.....Ipecac	101
Cephalanthus occidentalis,	
Button-bush	37
Cerascin.....Choke cherry	51
Cerasin.....Choke cherry	51
<i>Cerasus scrotina</i>	193
Cerates.....	236
Cercis canadensis.....Judas-tree	107

	PAGE.		PAGE.
Cerebral Sedative.....	237	Choke cherry.....	<i>Prunus virginiana</i> 51
<i>Cereus grandiflorus</i> ,.....		Cholesterin.....	248
Night-blooming cereus	126	Chondodendron tomentosum.....	
<i>Cerium nitrate</i> ,.....	247	<i>Pareira Brava</i>	135
Effervescent.....	210	Christmas rose.....	Black hellebore 27
oxalate, Effervescent.....	210	Chrysarobin.....	Goa powder 87
Pills.....	217	Cicely, sweet.....	175
Tablet triturates.....	231	<i>Cicuta maculata</i> ..	Water hemlock 188
<i>Cervipina cathartica</i> ,.....	35	<i>Cimicifuga</i>	Black cohosh 25
<i>Cevadilla</i>	<i>Veratrum Sabadilla</i> 49	<i>Cimicifuga racemosa</i> ..	Black cohosh 25
<i>Chamælorium carolinianum</i> ,.....		<i>Cinchona barks</i>	50
<i>Helonias</i>	93	<i>Cinchonidine sulphate</i>	52
<i>luteum</i> , <i>Chamælorium</i>		<i>Cinchonine sulphate</i>	52
<i>carolinianum</i>	93	<i>Cinnamomum camphora</i>	39
<i>Chamærops serrulata</i>	159	<i>U. S. P.</i>	Cassia 46
Chamberlin's Utero-vaginal Syringe		<i>zeylanicum</i>	46
.....	238, 244	Cinquefoil.....	Five-finger 79
Chamomile.....	<i>Anthemis nobilis</i> 49	<i>Cirsium arvense</i>	40
Tablet triturates.....	231	Citrine ointment, <i>U. S. P.</i>	236
dog.....	Mayweed 122	Citron basil.....	Sweet basil 175
German.....	86	Citronenkrout.....	Lemon balm 112
Spanish.....	Pellitory 137	<i>Citrullus Colocynthis</i>	59
Roman.....	Chamomile 49	<i>vulgaris</i>	Water-melon 188
wild.....	Mayweed 122	Citrus Aurantium.....	Sweet orange 177
Chanvre du Canada, Canadian hemp	40	<i>limetta</i>	Lime-juice 115
indien.....	Indian cannabis 100	<i>limonum</i>	Lemon 112
Charcoal.....	49	<i>vulgaris</i>	Bitter orange 23
Chardon bēnet.....	Blessed-thistle 29	Clavaliér.....	Prickly ash 145
Châtaigner.....	Chestnut 50	Claviceps purpurea.....	74
Chaulmoogra oil, <i>Gynocardia odorata</i>	50	Cleavers.....	<i>Galium Aparine</i> 55
Checkerberry.....	Squaw-vine 169	Climbing bittersweet,	
.....	Wintergreen 196	False bittersweet	77
Cheken.....	Chekan 50	Clotbur.....	<i>Xanthium strumarium</i> 55
Chekan.....	<i>Eugenia chequen</i> 50	spiny.....	168
Chelidonium majus,		Clotweed.....	Clotbur 55
Garden celandine	83	Clover, Dutch.....	White clover 190
<i>Chelone glabra</i>	Balmomy 15	red.....	149
Chelonin.....	Balmomy 15	sweet.....	176
Chemicals, rare.....	246	white.....	190
Chenopode anthelmintique,		Cloves.....	<i>Eugenia caryophyllata</i> , 55, 276
American wormseed	10	<i>Cnicus arvense</i>	Canada thistle 40
Chenopodium anthelminticum,		<i>benedictus</i>	29
Amer. wormseed	10	Coca.....	<i>Erythroxylon Coca</i> 55
<i>Botrys</i> , Jerusalem oak	107	cordial.....	55
Cherry.....	Wild cherry 193	Cocaine.....	Coca 55
birch.....	Black birch 25	Case.....	238
choke.....	51	citrate.....	Coca 56
wild.....	193	hydrobromate.....	Coca 56
Chestnut.....	<i>Castanea americana</i> 50	hydrochlorate.....	Coca 56
Chewstick ..	<i>Gouania domingensis</i> 50	salicylate.....	Coca 56
Chian turpentine,		Cocculus Indicus.....	Fish-berries 78
<i>Pistacia terebinthus</i>	51	<i>Palmatus</i>	61
Chickweed.....	<i>Stellaria media</i> 51	Cochlearia Armoracia,	
red ..	Scarlet pimpernel 160	Horse-radish	97
Chiendent.....	Couch-grass 66	Cocillana.....	<i>Sycocarpus Rusbyi</i> 57
Children, facts regarding.....	260	Cocklebur.....	Agrimony 5
Chimaphilin.....	Pipsissewa 142	spiny.....	Spiny clotbur 168
Chimaphila umbellata.....	Pipsissewa 141	Codeine.....	Opium 131
Chinese cinnamon.....	Cassia 46	Cod-liver oil.....	57
sumach, <i>Ailantus glandulosa</i>	5	Coffea arabica.....	Coffee 58
Chinoidin.....	53	Coffee.....	<i>Coffea arabica</i> 58
Elixirs.....	206	Cohosh, black.....	25
Chinoline.....	247	blue.....	30
salicylate.....	248	red.....	150
tartarate.....	248	white.....	190
Chionanthus virginica.....	Fringe-tree 82	<i>Cola acuminata</i>	Kola 109
Chirata.....	<i>Ophelia Chirata</i> 51	Colchicine.....	Colchicum 59
Chloral hydrate, Elixirs.....	206	Colchicum autumnale ..	Colchicum 58
Cerebral sedative ..	237	Cold-cream.....	Rose 154
Syrups.....	238	(Rose-water) ointment ..	236
Chlor-anodyne.....	100, 131	Colic root.....	Star-grass 170
Chlorinated Soda, <i>U. S. P.</i> , Solution.	241		Wild yam 196
Chloroform.....	238	Collinsonin.....	Stone-root 172
Capsules.....	203	Collinsonia canadensis.....	Stone-root 172
Inhalants.....	212		

	PAGE.		PAGE.
Colloids, Medicated.....	227	Cotton.....	Gossypium herbaceum 65
Colocynth.....	Cucumis Colocynthis 59	Couch-grass.....	Triticum repens 66
Colocynthin.....	Colocynth 60	Couleurvine.....	Bistort 23
Cologne, Antiseptic.....	241	Couleurée.....	White bryony 190
Coloquinte.....	Colocynth 59	Cowbane, spotted....	Water hemlock 188
Coloquintenapfel.....	Colocynth 59	Cow-parsnip.....	Masterwort 121
Coltsfoot.....	Tussilago Farfara 60	Crab Orchard, Effervescents.....	210
Coltsail.....	Fleabane 79	Mineral-water Salts.....	237
Columbo.....	Jateorrhiza Calumba 61	Cramp bark.....	Viburnum Opulus 66
American.....	9	Elixirs.....	207
Comfrey.....	Symphytum officinale 61	Cranberry, mountain.....	Uva-ursi 184
Common basil.....	Sweet basil 175	tree.....	Cramp bark 66
Compass plant.....	Rosin-weed 155	Crane willow.....	Button-bush 37
Compressed tablets.....	230	Cranesbill.....	Geranium maculatum 66
Comptonia asplenifolia.....	Sweet fern 176	Crateva marmelos.....	15
Concentrations, see respective crude drugs.		Crawley.....	Corallorhiza odontorhiza 67
Condurango, Gonolobus Cundurango 61		Creasote, Inhalants.....	212
Cone-flower.....	Thimble-weed 180	ointment.....	236
Confections.....	235	Croton Eluteria.....	Cascarilla 46
Congress, Effervescents.....	210	oil.....	Croton Tiglium 67
Mineral-water Salts.....	237	Tiglium.....	67
Conine.....	Conium 62	Croup syrup.....	Senega 162
hydrobromate.....	Conium 62	Crowfoot.....	Cranesbill 66
hydrochlorate.....	Conium 62	Crown bark.....	Pale cinchona 52
Conium maculatum.....	Conium 62	Cubeb.....	Piper cubeba 67
Consoude.....	Comfrey 61	Cubeba officinalis.....	67
Consumptive's weed.....	Yerba santa 200	Cucumber-tree, Magnolia acuminata 68	
Convallamarin.....	Lily of the Valley 114	Cucumis Citrullus.....	189
Convallaria majalis.....		Colocynthis.....	Colocynth 59
Lily of the Valley 114		Cucurbita Citrullus.....	189
Polygonatum.....	167	Pepo.....	Pumpkin 147
Convulvulus panduratus.....	120	Cunila mariana.....	Dittany 71
Scammonia.....	Scammony 159	Cup Syringe, The Indispensible.....	244
Coolwort.....	Mitella nuda 63	Curare.....	69
Copaiba.....	Copaifera Langsdorffii 63	Curcuma longa.....	Turmeric 183
Copaifera Langsdorffii.....	63	Curled dock.....	Yellow dock 198
Copernicia cerifera.....	Carnauba 43	Culvers.....	Veronica virginica 68
Copper oleate.....	235	Cusparia febrifuga.....	11
Coptide.....	Goldthread 89	trifoliata.....	Angustura 11
Coptis trifolia.....	Goldthread 89	Custard apple.....	Papaw (American) 135
Coque du Levant.....	Fish-berries 78	Cutch.....	Catechu 47
Coquelarde.....	Pulsatilla 147	Cynoglossum officinale, Hound's tongue 98	
Coquelicot.....	Red poppy 151	Cyperus articulatus.....	Adrie 4
Coqueluchon.....	Aconite 1	Cypripedin.....	Ladies' slipper 110
Coral-root.....	Crawley 67	Cypripedium pubescens, Ladies' slipper 110	
Corallorhiza odontorhiza.....	Crawley 67	Cytisus scoparius.....	Broom 33
Coriander.....	Coriandrum sativum 64		
Coriandrum sativum.....	Coriander 64		
Cornin.....	Dogwood 72		
Corn (maize).....	Zea Mays 64		
-ergot.....	Corn 64		
poppy.....	Red poppy 151	Damiana.....	Turnera microphylla 69
rose.....	Red poppy 151	Damson, mountain.....	Simaruba 165
-silk.....	Corn 64	Dandelion.....	Taraxacum officinale 70
-smut.....	Corn 64	Daphne Mezereum.....	Mezereum 123
-snake root.....	Water eryngo 188	Daturine.....	Stramonium 173
Cornel, silky.....	Red osier 150	Datura Stramonium.....	Stramonium 172
Cornouiller à feuilles arrondies, Green osier 90		Daturine sulphate.....	173
Cornus circinata.....	Green osier 90	Davenport's Unique Capsule Filler 240	
florida.....	Dogwood 71	Deadly nightshade.....	Belladonna 18
sericea.....	Red osier 150	Deerberry.....	Wintergreen 196
U. S. P.....	Dogwood 71	-tongue.....	Trilisa odoratissima 71
Corrosive sublimate, Colloids.....	227	laurel.....	Great laurel 69
Pills.....	215, 218, 226	Delphinium consolida.....	Larkspur 111
Tablet triturates.....	231	Staphisagria, Stavesacre 170	
and urea, Hypodermic tablets.....	229	Dent de chien jaune, Adder's tongue 4	
Corydalin.....	Turkey-corn 182	-de-leon.....	Dandelion 70
Corydalis canadensis.....	182	Dentition.....	261
Corypha cerifera, Copernicia cerifera 43		Devil's bit.....	Button-snake root 37
Costs, to estimate.....	254	Dewberry.....	Rubus canadensis 275
Coto.....	Nectandra (Sp. Indet.) 64	Dewee's Carminative.....	Asafetida 14
Cotoin.....	Coto 64	Dialyzed iron.....	238, 240
		glycerated.....	238, 240
		Malt Ext. Preparations.....	212
		Dicentra canadensis.....	Turkey-corn 182

	PAGE.		PAGE.
Dieffenbachia seguine....	Dumb cane 72	Ellébore noir.....	Black hellebore 27
Differential diagnosis of eruptive fevers.....	264	Elm bark.....	Ulmus fulva 166
Digestive Tablets.....	230	Emetine.....	Ipecac 101
Digitalin.....	Foxglove 80	Empty Capsules.....	239
Digitalis purpurea.....	Foxglove 80	Engelwurz.....	Angelica 11
Digitalisin.....	81	English moss.....	Sedum acre 161
Dillweed.....	Mayweed 122	Ephedra antisiphilitica.....	74
Dioscorea villosa.....	Wild yam 196	Epigaea repens.....	Gravel-plant 89
Dioscorein.....	Wild yam 196	Epilobium angustifolium.....	Willow-herb 196
Diospyros virginiana.....	Persimmon 140	Epine-vinette.....	Barberry 16
Dipterix odorata.....	Tonca-bean 181	Epiphegus virginiana.....	Beech-drop 18
Diseases and Their Remedies.....	265	Equisetum hyemale.....	Scouring rush 160
Disinfectant preparations.....	238, 241	Erdbeerblätter.....	Strawberry 174
Dita.....	Alstonia scholaris 71	Erechthites hieracifolia.....	Fireweed 78
Dittany.....	Cunila mariana 71	Ergot.....	Claviceps purpurea 74
Dock, yellow.....	193	Ergotin.....	Ergot 74
Dog chamomile.....	Mayweed 122	Ergotinic acid.....	Ergot 74
-grass.....	Couch-grass 63	Erigeron canadense.....	Fleabane 79
Dog's-bane.....	Bitter-root 23	oil, Capsules.....	203, 204
tooth violet.....	Adder's tongue 4	Eriodictyon glutinosum, Yerba santa	200
Dogwood.....	Cornus florida 71	Erlendrinde.....	Tag alder 177
Jamaica.....	105	Eruptive fevers, Differential diagnosis of.....	264
round-leaved.....	Green osier 90	Eryngium aquaticum.....	188
swamp.....	Red osier 150	yuccæfolium.....	Water eryngo 188
Doldenblütiges Harnkraut.....	Pipsissewa 141	Eryngo, water.....	188
Doom-bark.....	Sassy-bark 158	Erythraea Centaurium.....	European centaury 76
Dorema Ammoniacum.....	Ammoniac 11	Erythronium americanum.....	Adder's tongue 4
Douce-amère.....	Bittersweet 24	Erythrophloeum guineense.....	Sassy-bark 158
Dover's Powder.....	Ipecac, 102, 103	judiciale.....	158
Opium 131		Erythroxyton Coca.....	Coca 55
Compressed tablets.....	230	Eserine.....	Physostigmine 38
Syrups.....	228	Ether, Inhalants.....	211
Tablet triturates.....	231	spirits of nitrous, Elixirs.....	205
Dragon root.....	Green-dragon 90	Ethyl bromide.....	248
Indian turnip 101		carbamidate.....	249
Drimys Winteri.....	Winter's bark 196	Eucalyptus globulus.....	75
Drosera rotundifolia.....	Sundew 175	Eugenia caryophyllata.....	Cloves, 55, 276
Duboisia Myoporoides.....	72	chequen.....	Chekan 50
Duboisine sulphate.....	72	jambolana.....	Jambul 106
Dulcamara, U. S. P.....	Bittersweet 24	Pimenta.....	Pimento 141
Dumb cane.....	Dieffenbachia seguine 72	Eulachon oil.....	76
Durchwachsener Wasserhanf.....	Boneset 32	Euonymin.....	Wahoo 187
Dutch clover.....	White clover 190	Euonymus atropurpureus.....	Wahoo 187
myrtle.....	Sweet gale 176	Eupatorin.....	Boneset 32
Dwale.....	Belladonna 18	Eupatorium aromaticum.....	White snakeroot 192
Dwarf elder.....	Aralia hispida 72	perfoliatum.....	Boneset 32
Dyer's saffron.....	Safflower 155	purpureum.....	Queen of the meadow 148
Easton's syrup.....	Syrup Iron, quinine and strychnine phosph. 130	U. S. P.....	32
Eberraute.....	Southernwood 167	Euphorbe à Grand Fleurs.....	Flowering spurge 80
Echites suberecta.....	184	Euphorbia corollata.....	Flowering spurge 80
Ecorce d'anne.....	Tag alder 177	heterodoxa, Alveloz milk 8	
de Chene.....	White oak 181	pilulifera.....	76
Ehrenpreis Wundkraut.....	Speedwell 168	Enphrasia officinalis.....	Eyebright 77
Eichenrinde.....	White oak 191	Eupurpurin.....	Queen of the meadow 148
Eisenhut.....	Aconite 1	European anise.....	Anise 11
Elaterin, U. S. P.....	Elaterium 73	bugleweed.....	Bitter bugleweed 23
Elaterium.....	Momordica Elaterium 73	centaury.....	Erythraea Centaurium 76
Elbischwurz.....	Marshmallow 120	elder.....	Sambucus nigra 77
Elder.....	Sambucus canadensis 73	Evening primrose, Enothera biennis 77	
European.....	77	Excelsior Mineral-water Salts.....	237
Elecampane.....	Inula Helenium 73	Ecozonium purga.....	104
Elephant's-foot.....	77	Eyebright.....	Euphrasia officinalis 77
Elephantopus tomentosus.....	74		
Elephantopus tomentosus.....	74		
Elephant's-foot.....	74		
Elettaria Cardamomum, Cardamom 42			
Elixir formula.....	205		
proprietary.....	7, 126		
salicylic acid comp. Addenda 276			

	PAGE.		PAGE.
Fabiana imbricata	Pichi 140	Fœniculum officinale	Fennel 77
Facts regarding children	260	<i>vulgare</i>	77
Fagus ferruginea	Beech 18	Formulae of Pharmaceutical prepa-	
Fahrenheit and Centigrade Scales ..	255	arations.....	202
Fallkraut	Arnica 12	Fougère Male	Male-fern 117
False bittersweet , <i>Celastrus scandens</i>	77	Fowler's solution, Elixirs	207
grape.....	American ivy 10	Foxglove	<i>Digitalis purpurea</i> 80
hellebore.....	<i>Adonis vernalis</i> 4	Fragaria vesca	Strawberry 174
Jacob's ladder.....	Abscess root 1	<i>Franciscea uniflora</i>	118
sarsaparilla, American sarsaparilla.....	10	Frangula, U. S. P.	Buckthorn bark 35
unicorn.....	Helonias 93	Frankenia grandifolia, Yerba reuma	200
white cedar.....	Arbor vitæ 12	Fraseria carolinensis , American columbo.....	9
Faulbaumrinde	Buckthorn bark 35	Fraserin	American columbo 9
Faux fenouil	<i>Thapsia garganica</i> 180	Frauenhaar	Maiden-hair 117
Featherfew	Feverfew 78	Fraxinus americana	White ash 190
Feather geranium	Jerusalem oak 107	<i>sambucifolia</i>	Black ash 23
Fel bovis	Ox-gall 133	Freisamkraut	Pansy 135
Female Regulator	Life-root 114	Friar's balsam	Benzoin 22
Fenchelsamen	Fennel 77	Friederichshall, Effervescent	210
Fennel	<i>Fœniculum officinale</i> 77	Mineral-water Salts.....	237
water.....	188	Fringe-tree	<i>Chionanthus virginica</i> 82
Fenouil	Fennel 77	Frog lily	Yellow pond-lily 199
d'eau.....	Water fennel 188	Frostblow	Five-flowered gentian 79
Fern gale	Sweet fern 176	Frostweed	Frostwort 82
Fern meadow	Sweet fern 176	Frostwort, Helianthemum canadense	82
sweet.....	176	Fucus vesiculosus	Bladder-wrack 28
Ferula galbaniflua	82	Fünffingerkraut	Five-finger 79
Narthex.....	Asafetida 14	Fussblattwurzel	Mandrake 118
Sumbul.....	Musk-root 125		
Feuilles du Fraisier Sauvage , Strawberry.....	174	Gadus Morrhuæ	57
Fève de Calabar	Calabar bean 37	Galangal	Alpinia Galanga 82
Fève igasurique	Ignatia 98	Galbanum	<i>Ferula galbaniflua</i> 82
Fever-bush	Spice-bush 168	Gale, sweet	176
California.....	Black alder 24	Galipea Cusparia	11
tree.....	Eucalyptus globulus 75	<i>officinalis</i>	11
twig.....	False bittersweet 77	Galium Aparine	Cleavers 55
Feverfew	<i>Pyrethrum Parthenium</i> 78	Galium verum	Ladies' bed-straw 110
Fevers, Differential diagnosis of eruptive.....	264	Galla, U. S. P.	Nut-gall 127
Fiebertklee	Buckbean 34	Galläpfel	Nut-gall 127
Field sorrel	Sheep-sorrel 164	Gall-of-the-earth , Five-flowered gentian.....	79
Figwort	Carpenter's square 44	Galls	Nut-gall 127
Fine Chemicals	246	Gamboge	<i>Garcinia Hanburii</i> 83
Fir, balsam	15	Garcinia Hanburii	83
Fireweed	<i>Erechtites hieracifolia</i> 78	<i>Mangostana</i>	Mangosteen 120
Fleabane.....	78	Garden angelica	Angelica 11
Fischkörner	Fish-berries 78	celandine, <i>Chelidonium majus</i>	83
Fish-berries	<i>Anamirta cocculus</i> 78	lettuce.....	<i>Lactuca sativa</i> 83
Five-finger	<i>Potentilla canadensis</i> 79	sage.....	Sage 156
-flowered gentian, <i>Gentiana quinqueflora</i>	79	thyme.....	Thyme 180
leaves.....	American ivy 10	Garget	Poke-root 143
-leaved water hemlock, Water fennel.....	188	Garlic	<i>Allium sativum</i> 83
Flag, blue	31	Garrya Fremontii , California fever-bush.....	39
Lily	Blue flag 31	Gartenlattich	Garden lettuce 83
sweet.....	176	Gartenraute	Rue 155
Flasks, Sugar Test	243	Gartensalat	Garden lettuce 83
Flavored Lozenges	234	Gauchheil	Scarlet pimpernel 160
Fleabane	<i>Erigeron canadense</i> 79	Gaultheria procumbens, Wintergreen	196
Fleming's Tincture Aconite	3	Gay Feather	Button-snake root 37
Flesh-colored Asclepias , White Indian-hemp.....	191	Gayuba	Uva-ursi 184
Florida allspice, Calycanthus floridus	79	Gelatin Capsules, Improved empty ..	239
Flour, Taro	179	Gelbe Neisswurz	Goldthread 89
Flowering spurge , <i>Euphorbia corollata</i>	80	Gelsemine	84
Fluid Extracts, see respective drugs.		sulphate.....	84
Fly agaric	<i>Agaricus muscarius</i> 80	tartrate.....	84
-trap.....	Pitcher-plant 187	hydrochlorate.....	84
Trumpet-plant.....	182	Gelsemium	<i>Gelse-</i> <i>mium sempervirens</i> , 84, 275
		<i>sempervirens, Gelsemium</i>	84
		Tablet triturates.....	231
		Gelsemperin	84

	PAGE.		PAGE.
Gemeine Lobelie, Blue cardinal-flower	30	Grindelia robusta	90
Gemeiner Tüpfelfarn, Polypody	144	squarrosa	90
Gemeines Traubenkraut, Jerusalem oak	107	U. S. P., Grindelia robusta	90
Genét à balais, Broom	33	Grindwurz	198
Gentian, Gentiana lutea	85	Grosser wegetritt, Plantain	142
blue	32	Ground holly, Pipsissewa	141
five-flowered	79	ivy	91
<i>Gentiana Catesbæi</i>	32	laurel	89
lutea	85	lily	22
ochroleuca, Gentian	85	-moss	92
Sampson snake-root	156	-squirrel pea	184
puberula, Blue gentian	32	Groundsel, Life-root	114
quinqueflora, Five-flowered gentian	79	Grüner Germer, American hellebore	9
Geranium, Cranesbill	67	Guaco, Mikania guaco	91
Geranium maculatum, Cranesbill	66	Guaiac, Guaiacum officinale	91
German chamomile, Matricaria Chamomilla	86	Guaiacum officinale, Guaiac	91
tinctures	236	Guarana, Paulinia sorbilis	91
Germandrée aquatique, Water germander	188	Guérit tout, Stone-root	172
Geum rivale, Water avens	187	Gun Benjamin, Benzoin	21
Geyser Mineral-water Salts	237	sweet	176
Gichtrübe, White bryony	190	Gundelrebe, Ground ivy	91
Giflatlach, Wild lettuce	195	Gundermann, Ground ivy	91
Giflatichsaft, Lactucarium, U. S. P.	110	Gurjun balsam	92
Giftsumach, Poison oak	143	Gynocardia odorata	50
Gillenia trifoliata, Indian physic	101		
Gill-over-the-ground, Ground ivy	91	Hackmatac, Tamarack	177
Gill-run, Ground ivy	91	Hæmatic Hypophosphites, Syrup of	130
Gingembre, Ginger	86	Hæmatoxyton campechianum, Logwood	116
Ginger ale	86	Hair-wash, Soap-tree bark	166
Jamaica, Zingiber officinale	86	Polytrichum juniperinum	92
Mexican, Amomum Zingiber	87	Hamamelin, Witch-hazel	197
wild	195	Hamamelis virginica, Witch-hazel	197
Ginseng, Aralia quinquefolia	87, 276	Hardhack, Spiræa tomentosa	92
Glacial phosphoric acid	246	Hashish, Indian cannabis	100
Glatter sumach, Sumach	174	Hathorn, Mineral-water Salts	237
Globe flower, Button-bush	37	Haw, black	26
Glucosides, rare	246	Heal-all, Stone-root	172
Glycerole Pepsin concentrated	276	Haarlem oil, Capsules, 203, 205	
Yerbine comp.	211	Heartsease, Pansy	135
Glyceroles	211	Hedeoma pulegioides, Pennyroyal	137
Glycyrrhiza glabra, Licorice	113	Helenenwurz, Elecampane	73
Glycyrrhizin, Licorice	113	Helianthella tenuifolia	93
Gnaphalium polycephalum, Life-everlasting	113	Heliantheme de Canada, Frostwort	82
Goa powder	87	Helianthemum canadense, Frostwort	82
Golden-rod, Solidago odora	87	Helianthus annuus, Sunflower	175
-seal		Hellebore, American	9
Hydrastis canadensis	87, 275	black	27
Goldthread, Coptis trifolia	89	false, Adonis vernalis	4
Gonolobus Cundurango, Condurango	61	white	191
Goose-grass, Cleavers	55	Helleborus niger, Black hellebore	27
Gossypin, Cotton	65	Helmkraut, Scullcap	160
Gossypium herbaceum, Cotton	65	Helonias, Chamælorium carolinianum	93
Götterbaum, Ailantus glandulosa	5	dioica	93
Gouania domingensis, Chewstick	50	Helonin, Helonias	93
Granatum, U. S. P., Pomegranate	144	Hematic Hypophosphites, syrup, 54, 228	
Grand Boucage, Saxifrage	159	Hemlock, Hemlock spruce	93
Grande ciguë, Conium	62	five-leaved water,	
Granular Effervescent Preparations	210	Water fennel	188
Granules, pink	225	spruce, Tsuga canadensis	93
Graue Wallnusrinde, Butternut	86	water	188
Gravel-plant, Epigæa repens	86	Hemp, Canadian	40
root, Queen of the meadow	148	white Indian	191
Great celandine, Garden celandine	83	Herbane, Hyoscyamus niger	94
laurel, Rhododendron maximum	89	Hepatica triloba	115
lobelia, Blue cardinal-flower	30	Heracleum lanatum, Masterwort	121
Green-dragon, Arisæma Dracontium	90	Herba Centaurii, Pharm. Ger., European centaury	76
hellebore, American hellebore	9	Herbe à l'hirondelle, Garden celandine	83
osler, Cornus circinata	90	au citron, Lemon balm	112
		de poultot, Pennyroyal	137
		Hetre ferrugineux, Beech	18

	PAGE.		PAGE.
Heuchera americana	Alum root 8	Impatiens pallida	Jewel-weed 107
Hickory, shell-bark	164	Imperial measure, unit equivalents ..	252
Hiera hiera	Canella 41	Incompatibility	261
..... Tincture of	7	Indian black-root,	
High angelica	Angelica 11	Pterocaulon pycnostachyum ..	99
cranberry	Cramp bark 66	cannabis	Cannabis indica 100
-rock, Mineral-water Salts	237	cup plant	Rosin-weed 155
Hippuric acid	246	hemp, foreign,	
Hirtentaschlein	Shepherd's purse 165	Indian cannabis	100
Hive syrup	Senega 162	white	191
Hoang-nan,		paint	Blood-root 29
Strychnos malaccensis, 95, 276		physic	Gillenia trifoliata 101
Hogwood	Ragweed 149	poke	American hellebore 9
Hollyhock	Althæa rosea 95	sage	Boneset 32
Holly-leaved barberry,		tobacco	Lobelia 115
Berberis aquifolium	22	turmeric	Golden-seal 87
Homatropine hydrobromate,		turnip	Arisæma triphyllum 101
Belladonna	19	Indigo, wild	195
Honduras bark	Cascara amarga 44	Indischer Hanf	Indian cannabis 100
sarsaparilla	Sarsaparilla 157	Indispensable Cup Syringe	244
Hoodwort	Sculicap 160	Ingwer	Ginger 56
Hopfenbaum	Wafer ash 186	Inhalers, Semple's Atomizing ..	238, 243
Hopfenheimbuche	Iron-wood 103	Ink root	Marsh rosemary 121
Hop hornbeam	Iron-wood 103	Intoxicating mushroom of Siberia ..	60
tree	Wafer ash 186	Inula Helenium	Elecampane 73
Hops	Humulus Lupulus 96	Inulain	Elecampane 73
Horehound	Marrubium vulgare 96	Iodine, Capsules	203
Horse-balm	Stone-root 172	Collodions	227
Capsules	239	ointment, compound	236
-chestnut,		tincture, Inhalants	212
Æsculus Hippocastanum ..	97	Iodoform, Capsules	203
-fly weed	Wild indigo 195	Pills	219, 226
Horsemint	Monarda punctata 97	Bougies, gelatin-coated ..	238
-radish	Cochlearia Armoracia 97	Iodol	248
-weed	Stone-root 172	Ipecac	Cephaelis Ipecacuanha 101
.....	Fleabane 79	American	Indian physic 101
Houlachan	Eulachon 76	wild	Flowering spurge 80
Hound's tongue,		Ipomœa pandurata	Man-root 120
Gynoglossum officinale ..	98	purga	Jalap 104
Houttuynia californica, Yerba mansa ..	199	Iris	Blue flag 31
Hufattig	Coltsfoot 60	florentina	Orris 133
Humulus Lupulus	Hops 96	versicolor	Blue flag 31
Hundskamillenkraut	Mayweed 122	Irish broom	Broom 33
Huntsman's cup	Pitcher-plant 142	Irisin	Blue flag 31
Huxham's tincture	52, 54	Iron arseniate, Tablet triturates ..	231
Hydrangea arborescens	98	and ammonium citrate,	
Hydrargyrum tannicum oxydulatum ..	248	Capsules	203, 205
Hydrastin	Golden-seal 89	Effervescent	210
Hydrastine	88	Elixirs	209
hydrochlorate	88	Malt ext. preparations ..	212
sulphate	88	Wines	234
Hydrastis canadensis	Golden-seal 87	and quinine citrate	53, 54
Hydriodic acid, Syrups	228	Syrups	228
Hydrochinon	248	Effervescent	210
Hydrochloric acid, Pills	216, 218	Malt ext. preparations ..	212, 213
Hyonigrin	94	Pills	220
Hyoscine hydrobromate	94	Tablet triturates	232
Hyoscyamine	94	strychnine and malt ..	129
Hyoscyamus niger	Henbane 74	ext.,	
Hypericum perforatum,		with arsenic	53
Saint John's wort	156	and sodium albuminate, Syrups ..	228
Hypodermic Case, Improved	238, 242	and strychnine citrate	130
tablets	229	Syrups	228
Hyponæ	248	Elixirs	208
Hypophosphites, Syrup of Hæmatic ..	130	Tablet triturates	232
Hyssop	Hyssopus officinalis 98	bromide, Elixirs	206, 207
Hyssopus officinalis	Hyssop 98	Effervescent	210
.....		Syrups	228
.....		Pills	215, 220
Ichthyol	248	by hydrogen, Pills	215,
Ignatia	Strychnos Ignatii 98	217, 219, 220, 221,	
Ignatius bean	Ignatia 98	222, 223, 224, 225, 226	
Ignazbohne	Ignatia 98	Tablet triturates	231, 232
Ilex paraguaiensis	Paraguay tea 135		

	PAGE.		PAGE.
Iron carbonate, Effervescent	210	Jamaica dogwood, <i>Piscidia Erythrina</i>	105
Lozenges	233	ginger	86
Pills	230	pepper	141
chloride, Elixirs	207	Pimento	141
solution, U. S. P., Col-		Jambu assu	Ottonia anisum 106
iodious	227	Jambul	Eugenia jambolana, 106, 275
tincture, Capsules	203	Jamestown weed	Stramonium 172
citrate, Elixirs	206, 207	Japonica	Catechu 47
Pills	218, 220	Jasmin, yellow	Gelsemium 84
Syrups	228	Jasundi	Saraca indica 157
dialyzed	240	Jatamansi	Musk-root 125
glycerated	240	Jateorrhiza Calumba	Columbo 61
Malt ext. preparations	212	Jeffersonia diphylla	Twin-leaf 184
ferrocyanide, Pills	214, 225, 219, 220, 225	Jequirity	Abrus precatorius 106
hypophosphite, Syrups	228	Jersey tea	New Jersey tea 126
Malt ext. prep's	213	Jerusalem oak, <i>Chenopodium Botrys</i>	107
iodide, Syrups	228	American wormseed	10
Capsules	203, 205	Jessamine, yellow	Gelsemium 84
Effervescent	210	Jesuits' drops	Benzoïn 22
Malt ext. preparations		Jewel-weed	Impatiens pallida 107
Pills	212, 213	Jimsonweed	Stramonium 172
lactate, Syrups	228	John's-wort	St. John's-wort 156
Pills	220	<i>Joutsia Asoka</i>	157
Tablet triturates	232	Judas-tree	Cercis canadensis 107
lactophosphate, Syrups	228, 229	Juglandin	Butternut 37
Malt ext. prep's	213	Juglans cinerea	Butternut 36
oleate	235	nigra	Black walnut 28
phosphate, Syrups	228, 229	U. S. P.	Butternut 36
Effervescent	210	Juniper	Juniperus communis 107
Elixirs	207, 208, 209	Juniperus communis	Juniper 107
(ferrous), Pills	220	sabina	Savin 158
(precip.), Pills	222, 224	Jupeba nall.	Jurubeba 275
quinine and strychnine	53, 128	Jurubeba	Solanum paniculatum 275
plaster	236	Juuna	Jurubeba 275
proto-carb. (Vallet's) Pills	217, 219, 220, 222, 223, 224, 225, 226	Kairine	248
-chloride, Malt ext. prep's	213	Kalabarbohn.	Calabar bean 37
protoxide, Elixirs	206, 207	Kalmia angustifolia	123
Syrups	228	latifolia	Mountain laurel 123
pyrophosphate, Elixirs	205, 206, 207, 208, 209	Kalmus-wurzel	Sweet flag 176
Lozenges	233	Kamala	Mallotus philippinensis 108
Malt ext. prep's		Kastanienblätter	Chestnut 50
Pills	212, 213	Katechu	Catechu 47
Syrups	228	Katzenminze	Catnep 48
Wines	234	Kava kava	Piper Methysticum 108
and pepsin	139	Kawahin	Kava kava 108
strychnine and malt ext.	129	Kawa	Kava kava 108
quinine and strychnine	130	Kelp-ware	Bladder-wrack 28
phosphates, 54	130	Kerries mineral, Pills	220
(roborans) plaster	236	Kino	Pterocarpus Marsupium 109
sulphate, exsic. Pills	213, 214, 215, 217, 218, 219, 220, 221, 223, 224	Kissengen, Mineral-water Salts	237
superphosphate, Syrups	208	Effervescent	210
valerianate, Elixirs	208	Klatschrosen	Red poppy 151
Pills	220, 224	Kleebaum	Wafer ash 186
-wood	Ostrya virginica 103	Knoblauch	Garlic 83
Issue plasters	41	Knob-root	Stone-root 172
Itch weed	American hellebore	Knotenwurz	Carpenter's square 44
Ivy, American	10	Knot-grass	Couch grass 66
big-leaved	Mountain laurel 123	Kola	Cola acuminata 109
poison	Poison oak 143	Konigschina	Yellow cinchona 52
Jaborandi	Pilocarpus pinnatifolius 103	Königskerze	Mullein 124
Jacaranda procera	43	Kosso	Brayera anthelmintica 109
Jacob's ladder	Carion-flower 44	Krameria triandra	Rhatany 151
Jalap	Iponoea purga 104	Krebswurz	Beech-drop 18
wild	Man-root 120	Kreuzdornbeeren	Buckthorn berries 35
Jalapin	104	Kreuzkraut	Life-root 114
		Kriechender weisser Klee	White clover 190
		Küchenschelle	Pulsatilla 147
		Labarraque's Solution	241
		Labkraut	Cleavers 55

	PAGE.
Labrador tea <i>Ledum latifolium</i>	109
Lachenknoblauch, Water germander	188
Lactuca canadensis. Wild lettuce	195
Lactuca sativa Garden lettuce	83
<i>virosa</i>	195
<i>Lactucarium</i> , U. S. P.	110
Lactucarium, U. S. P., <i>Lactuca virosa</i>	110
Ladies' bed-straw. <i>Galium verum</i>	110
slipper, <i>Cypripedium pubescens</i>	110
Laitue sative. Garden lettuce	83
Laitue vireuse. Wild lettuce	195
Lamb's-quarter. Beth-root	23
Lanolin	111
Lappa officinalis. Burdock	36
Larch agaric White agaric	189
American. Tamarac	177
black. Tamarac	177
Lärchenschwamm White agaric	189
Large flowering spurge, <i>Flowering spurge</i>	80
Larix americana Tamarac	177
Larkspur Delphinium consolida	111
Latin used in prescriptions.	262
Laudanum Opium	133
Laurel, bay	17
California.	39
deer-tongue. Great laurel	89
great.	89
ground. Gravel-plant	89
mountain	123
Laurier benzoin Spice-bush	168
commun. Bay laurel (1)	17
Laurus nobilis Bay laurel (1)	17
Lavender Lavandula spica	111
sea Marsh rosemary	121
Lavandula, spica Lavender	111
<i>vera</i>	111
Laxative Elixir.	208
Lead acetate, Pills.	216, 221
oleate.	235
plaster.	236
subacetate cerate, U. S. P.	236
Glycerole.	211
Leaf-cup Bearsfoot	18
Lebensbaum Arbor vitæ	12
Leberklette Agrimony	5
Ledum latifolium Labrador tea	109
Lemon Citrus limonum	112
balm Melissa officinalis	112
Leonurus Cardiaca Motherwort	123
Leopards bane Arnica	12
Leptandra, U. S. P. Culvers	68
<i>virginica</i>	68
Leptandrin Culvers	69
Lettuce, garden.	83
opium, <i>Lactucarium</i> , U. S. P.	110
wild.	195
Levant wormseed, <i>Artemisia maritima</i>	112
Lever-wood Ironwood	103
Levisticum officinale Lovage	116
<i>Liatris odoratissima</i>	71
spicata	188
Button-snake root	37
<i>Lichen pulmonarius</i> , <i>Sticta pulmonaris</i>	116
Licorice Glycyrrhiza glabra	113
Lierre terrestre Ground ivy	91
Life-everlasting <i>Gnaphalium polycephalum</i>	113
-root Senecio aureus	114
<i>Ligusticum levisticum</i>	116
<i>Ligustrum vulgare</i> Privet	146
Lily of the valley, <i>Convallaria majalis</i>	114

	PAGE.
Lily white pond	192
yellow pond	199
Lime-juice Citrus limetta	115
tree. Basswood	17
Limonis cortex, U. S. P. Lemon	112
Linden Basswood	17
Lindera Benzoin Spice-bush	168
Linseed oil, Capsules	203
Lippia dulcis. Lippia Mexicana	115
Mexicana. Lippia dulcis	115
Liquor Acidi Phosphorici	241
sedans	276
Liquid Acid Phosphate	238, 241
Syrups	238
and pepsin.	140
Liquidambar styraciflua, Sweet gum	176
Liriodendron tulipifera Tulip tree	182
Lithium benzoate, Effervescent.	210
bromide, Elixirs	208
carbonate, Tablet triturates	232
citrate, Effervescent.	210
Elixirs	208
salicylate, Elixirs.	209
and manaca.	118
Liver lily Blue flag	31
wort. Anemone Hepatica	115
Lobelia inflata Lobelia	115
syphilitica, Blue Cardinal-flower	30
U. S. P. Lobelia inflata	115
Lobellin Lobelia	116
Logan's plaster	236
Logwood, <i>Hæmatoxylon campechianum</i>	116
Lorbeer Bay laurel (1)	17
Löwenzann Dandelion	70
Low blackberry Dewberry	275
Mallow. Malva rotundifolia	116
Lovage Levisticum officinale	116
Love-apple. Tomato	181
Loxa bark Pale cinchona	52
Lozenges, flavored.	234
medicated	233
Lung-moss. Sticta pulmonaria	116
wort. Pulmonaria officinalis	117
Lupulin Hops	96
Lycopersicum esculentum. Tomato	181
Lycopin. Bugleweed	35
Lycopus europæus, Bitter Bugleweed	23
virginicus. Bugleweed	35
Mace. Myristica fragrans	117
Macroton Black cohosh	150
Mad-dog weed. Scullcap	260
Magnesia, Pills	221
and asafetida. Asafetida	14
Magnesium acetate, Elixirs.	209
aperient, Effervescent.	210
citrate, Effervescent.	210
sulphate, Effervescent.	210
Magnolia acuminata, Cucumber-tree	68
flowers. Cucumber-tree	68
glauca. Magnolia	117
U. S. P. Magnolia glauca	117
Maiden-hair. Adiantum pedatum	117
Male-fern Aspidium Filix-mas	117
shield-fern. Male-fern	117
Mallotus philippinensis. Kamala	108
Mallow, low. Malva rotundifolia	116
Malt extract and its preparations.	212
Malva rotundifolia. Low mallow	116
Manaca Brunfelsia Hopeana	118
Mancona-bark Sassy-bark	158
Mandrake Podophyllum peltatum	118

	PAGE.		PAGE.
Manganese bromide, Malt ext. prep's	212	Mercury oleate	235
binocide, Pills	220	plaster	236
Tablet triturates	232	protiodide, Pills, 219, 221, 225,	236
carb., Pills	220	Tablet triturates	232
hypophosphite, Syrups	238, 239	weed.....Mercurialis annua	122
iodide	221	with chalk, Pills	226
Malt ext. prep's	212, 213	Tablet triturates	232
Syrups	239	yellow oxide of, Compressed	230
Pills	215	tablets	230
lactophosphate, Syrups	239	Tablet triturates	232
Malt ext. prep's	213	Merrettig	97
oleate	235	Metaphosphoric acid	246
Mango-fruit	120	Mexican ginger	87
Mangosteen	120	sarsaparilla	157
Garcinia Mangostana	120	Mezereon	123
Man-in-the-ground	120	Mezereum, U. S. P., Daphne mezereum	123
-of-the-earth	120	Micromeria Douglassii, Yerba buena	199
root	120	Mikania guaco	91
Manzanita	120	Milfoil	198
Arctostaphylos glauca	120	Milkweed	165
Maple, red	150	swamp, White Indian-hemp	191
swamp	150	Millefeuille	138
Marigold	38	Mineral-water salts	237
Calendula	38	Mint, mountain	124
Marjoram, sweet	177	Mistletoe	123
Maroute	122	Phoradendron flavescens	123
Marrubium vulgare	96	Mistura Smilacis Comp.	16
Marsh-mallow	120	Bamboo brier	169
Althæa officinalis	120	Mitchella repens	63
rosemary	121	Mitella cordifolia	63
trefoil	84	nuda	63
Mary thistle	121	Mitrewort	63
Silybum marianum	121	Moccasin flower	110
Masterwort	11	Momordica Elaterium	73
Angelica	11	Monarda fistulosa	193
Heracleum lanatum	121	Wild bergamot	97
Mastic	121	punctata	97
Pistacia Lentiscus	121	Monkshood	1
Matico	121	Aconite	1
Artanthe elongata	121	Mooneed	199
Matricaire	78	Morphine and its salts	130
Matricaria Chamomilla	86	Mossy stonecrop	161
German chamomile	86	Motherwort	123
May-apple	118	Leonurus Cardiac	123
Mandrake	118	Mountain-ash	123
-flower	89	Pirus americana	200
Gravel-plant	89	balm	184
weed	122	Yerba santa	184
Anthemis Cotula	122	box	184
Mayer's ointment	236	Uva-ursi	184
Meadow fern	176	cranberry	165
Sweet fern	176	Simaruba	165
Colchicum	58	grape, Berberis aquifolium	22
safron	92	Kalmia latifolia	123
Hardhack	92	laurel	123
Measures, approximate	235	mint	124
Medicated Collodions	237	Pycnanthemum	124
Lozenges	233	montanum	124
Medicinal Elixirs	205	pink	69
Syrups	238	Gravel-plant	124
Wines	234	Artemisia frigida	196
Meerzwiebel	169	sage	196
Melia Azedarach	146	Wintergreen	12
Méillot	176	Arnica	12
Sweet clover	176	Mouron rouge	160
Melilotenkle	176	Scarlet pimpernel	97
Melissa officinalis	112	Moutarde des moines	89
Mélisse	112	Horse-radish	89
Melon-tree	43	Mouthroot	124
Carica papaya	43	Goldthread	124
Menispermum	199	Mugwort	124
Yellow parilla	199	Artemisia vulgaris	124
Menispermum canadense	199	Mullein	74
Yellow parilla	199	Verbascum Thapsus	74
Mentha arvensis	122	Mutterkorn	74
Menthol	122	Ergot	74
piperita	138	Mutterkraut	74
viridis	167	Feverfew	16
Menthol	122	Musa sapientum	80
Mentha arvensis	122	Banana	80
Menyanthes trifoliata	34	Muscarine sulphate	60
Mercurial ointment, U. S. P.	236	Mushroom of Siberia, intoxicating	234
Mercurialis annua	122	Fly agaric	125
Mercuric iodide, Compressed tablets	230	intoxicating	234
Mercurio vegetal	118	(lozenges)	125
Mercurous tannate	248	Ferula Simbul	188
Mercury and morphine oleates	132	Musquash root	17
biniodide, Pills	221	Water hemlock	17
Tablet triturates	232	Bay laurel (2)	17
		cerifera	176
		Gale	17
		Sweet gale	17
		Myricin	17
		Bayberry	117
		Myristica fragrans	127
		Mace	127
		Nutmegs	127
		U. S. P.	127
		Nutmegs	127

	PAGE.
Myroxylon Pereiræ..Balsam of Peru	15
Toluiferum.....Tolu	181
Myrrh.....Balsamodendron Katak	125
Myrtle, Dutch.....Sweet gale	176
Myrtus chekan.....	50
Nackte Aralienwurzel,	
American sarsaparilla	10
Napelline.....Aconite	2
Naphthalin.....Tar	178
Naphthol.....Tar	178
Narrow dock.....Yellow dock	198
Natterwurz.....Bistort	23
Nectandra (sp. indet.).....Coto	64
Nepeta cataria.....Catnep	48
Glechoma.....Ground ivy	91
Nerprun.....Buckhorn berries	35
Nettle.....Urtica dioica	126
leaved vervain,White vervain	192
New Jersey tea,	
Ceanothus americanus	126
Nickel oleate.....	235
Nicotiana Tabacum.....Tobacco	180
Night-blooming cereus,	
Cereus grandiflorus	126
Nightshade, deadly.....Belladonna	18
yellow-flowered,	
Urechites suberecta	184
Nitroglycerin.....	248
Capsules.....	204
Pills.....	221
Tablet triturates.....	232
Noix d' Arec.....Arecia	12
de galle.....Nut-gall	127
Normal liquids, see respective drugs.	
Norwood's tincture,	
American hellebore	10
Noyer gris.....Butternut	36
Nuphar advena.....Yellow pond-lily	199
Nut-gall.....Quercus infectoria	117
Nutmegs.....Myristica fragrans	127
Nux vomica.....Strychnos Nux vomica	127
Nymphæa odorata...White pond-lily	192
Oak, black.....	27
Jerusalem,	
American wormseed	10
Chenopodium Botrys	107
poison.....	143
red.....	150
white.....	191
Ocimum Basilicum.....Sweet basil	175
Odermennig.....Agrimony	5
CEnanthe Phellandrium,	
Water fennel	188
CEnothera biennis, Evening primrose	77
Oil, Haarlem, Capsules.....	203, 205
linseed, Capsules.....	203
nut.....Butternut	36
Ointments.....	236
of oleates.....	235
Old man.....Southernwood	167
man's beard.....Fringe-tree	82
Oleates.....	235
ointments of.....	235
Oolachon.....Eulachon	76
Ophelia Chirata.....Chirata	51
Opium.....Papaver somniferum	130
Cerebral sedative.....	237
Orange, bitter.....	23
root.....Golden-seal	87
sweet.....	177
Ordeal bean.....Calabar bean	37

	PAGE.
Oregon-grape...Berberis aquifolium	22
Origanum Majorana,	
Sweet marjoram	177
oil.....Thyme	180
(Thyme) oil, Capsules....	204
Orobanche de Virginie...Beech-drop	18
Orrice root.....Orris	133
Orris.....Iris florentina	133
Osier, green.....	90
red.....	150
Osmic acid.....	247
Osmorrhiza longistylis..Sweet cicely	175
Osmunda regalis...Buckhorn brake	34
Ostrya virginica.....Iron-wood	103
Ottonia anisum.....Jambu assu	106
Jaborandi.....	106
Ox-balm.....Stone-root	172
-gall.....	133
Oxyacanthine.....Barberry	17
Oxydendrum arboreum...Sourwood	107
Oxyethylquinoline-hydride hydrochloride.....	248
Pale cinchona...Cinchona officinalis	52
Palma christi.....Castor-oil plant	47
Panax quinquefolium.....	87, 276
Pancreas.....Pancreatin	134
Pancreatin.....	134
Digestive tablets.....	230
Pansy.....Viola tricolor	135
Papain.....Carica Papaya	43
Papaver Rhœas.....Red poppy	151
somniferum.....Opium	130
.....Poppy	145
Papaw (American)..Asimina triloba	135
(true).....Carica Papaya	43
Papayotin.....Carica Papaya	43
Papoose root.....Blue cohosh	30
Para sarsaparilla.....Sarsaparilla	157
-coto bark.....Coto	64
Paracotoin.....Coto	64
Paraguay tea.....Ilex paraguaiensis	135
Paraldehyd.....	248
Pareira Brava,	
Chondodendron tomentosum	135
Parilla, yellow.....	199
Parsley.....Petroselinum sativum	136
spotted.....Water hemlock	188
Parsnip, cow.....Masterwort	121
Partridgeberry.....Squaw-vine	169
.....Wintergreen	196
Pasque flower.....Pulsatilla	147
Patience frisée.....Yellow dock	198
Paulinia sorbilis.....Guarana	91
Paul's betony.....Bugleweed	35
Pavot rouge.....Red poppy	151
Peach.....Amygdalus persica	136
Pearls, amyli nitrite.....	238
Pectoral lozenge.....	102
Pelletierine sulphate...Pomegranate	144
tannate...Pomegranate	144
Pellitory.....Anacyclus Pyrethrum	137
Pencil-flower.....Stylosanthes elatior	137
Pennsylvania sumach.....Sumach	174
Pennyroyal.....Hedeoma pulegioides	137
Pensée.....Pansy	135
Penthorum sedoides,	
Virginia stone-crop	186
Peony.....Pæonia officinalis	137
Pepo, U. S. P.....Pumpkin	147
Pepper, black.....	27
Peppermint.....Mentha piperita	138
oil of, Soda-mint tablets..	230
Pepsin.....	138, 276

	PAGE.		PAGE.
Pepsin, Digestive tablets	230	Podophyllotoxin	Mandrake 118
Sacch., Syrups	228	Podophyllum peltatum	Mandrake 118
Peptonized Extract of Beef	238, 242	Poh di Bahia	Goa powder 87
Peptonizing tablets	Pancreatin, 134, 230	Poison hemlock	Conium 62
Perosmic acid	247	ivy	Poison oak 143
Persea gratissima	Alligator pear 6	oak	Rhus Toxicodendron 143
Persica vulgaris	136	Tablet triturates	232
Persil	136	parsley	Conium 62
Persimmon	Diospyros virginiana 140	Poke-root	Phytolacca decandra 143
Peruvian bark	Cinchona 52	Polar-plant	Rosin-weed 155
Petersilie	Parsley 136	Polecat weed	Skunk cabbage 165
Petroleum, Capsules	203	Polemonium reptans	Abscess root 1
crude, Malt ext. prep's	213	Polygala Senega	Senega 161
mass	238	Polygonatum officinale	
Petroselinum sativum	Parsley 136	Solomon's seal	167
Pettymorrell	Spikenard 168	Polygonum acre	Water-pepper 189
Peumus Boldus	32	bistorta	Bistort 23
Phenyl-methyl-acetone	248	hydropiperoides	189
Phoradendron flavescens	Mistletoe 123	punctatum	189
Phosphate, Liquid Acid	241	Polymnia Uvedalia	Bearsfoot 18
Phosphoric acid, glacial	246	Polypodium vulgare	Polypody 144
Phosphorus, Capsules	203, 204	Polypody	Polypodium vulgare 144
Elixirs	206, 207, 209	Polyporus officinalis	White agaric 189
Glyceroles	211	Polytrichum juniperinum	
Malt ext. preparations	212	Haircap moss	92
Pills, 215, 221, 222, 223, 225, 226		Pomegranate	Punica Granatum 144
Physicians' Pocket Reagent Case	238, 245	Pomeranzenschale	Bitter orange 23
Physostigma venenosum		Pond dogwood	Button-bush 37
Calabar bean	37	-lily, white	192
Physostigmine	Calabar bean 38	yellow	199
salicylate	38	Pongamia glabra	145
sulphate	38	Poplar	White poplar 192
Phytolacca decandra	Poke-root 143	balsam	16
Phytolaccin	Poke-root 144	yellow	182
Pichi	Fabiana imbricata, 140, 276	white	192
Picraena excelsa	Quassia 147	Poppy	Papaver somniferum 145
Picramnia (sp. indet.)	Cascara amara 44	red	151
Picrotoxin	Fish-berries 78	Populin	White poplar 192
Pigeon berry	Poke-root 143	Populus candicans	Balsam poplar 16
Pill Formulæ List	213	tremuloides	White poplar 192
Pills of the British Pharmacopœia	226	Portulaca oleracea	Purslane 147
Pilocarpine hydrochlorate, Jaborandi	104	Posology	255
nitrate	Jaborandi 104	Potassium acetate, Elixirs	205
Pilocarpus pinnatifolius	Jaborandi 103	and iron tartrate, Pills	223
Pimento	Eugenia Pimenta 141	and sodium tartrate, Pills	215
Pimpinella	Saxifrage 159	arsenite solution, Elixirs	206
Anisum	Anise 11	Pills	227
Pine, white	161	benzoate, Effervescent	210
Pink Granules	225	bicarbonate, Elixirs	209
-root	Spigelia marilandica 141	Compressed	
Pinus australis	Turpentine 183	tablets	230
palustris	178	bromide, Cerebral seda-	
Strobis	White pine 191	tive	237
Piper cubeba	Cubeb 67	Compressed	
methysticum	Kava-kava 108	tablets	230
nigrum	Black pepper 27	Effervescent	210
U. S. P.	Black pepper 27	Elixirs	206, 209
Pipitzaohmic acid	247	Glyceroles	211
Pipsissewa	Chimaphila umbellata 141	Malt ext. prep's	212
Pirus americana	Mountain-ash 123	Pills	223, 226
Piscidia Erythrina		carbonate, Pills	219
Jamaica dogwood	105	chlorate, Compressed	
Piscidin	Jamaica dogwood 105	tablets	230
Pistacia Lentiscus	Mastic 121	Glyceroles	211
terebinthus	51	Lozenges	223
Pitcher-plant	Sarracenia purpurea 142	citrate, Syrups	229
Plantago major	Plantain 142	Effervescent	210
Plantain	Plantago major 142	and ipecac, Syrup	103
Plasters, Issue	41	hypophosphite, Malt ext.	
Roll	236	prep's	213
Pleurisy-root	Asclepias tuberosa 142	Syrups	228
Plummer's pill, see Antimony comp.		iodide, Elixirs	207
Pills and Tablet triturates		Effervescent	210
Pocket Reagent Case	245	Pills	215, 223
Podophyllin	Mandrake 118		

PAGE.	PAGE.
Potassium lactophosphate, Syrups. 228, 229	Quickens..... Couch-grass 66
Malt ext. 213	Quillaia saponaria..... Soap-tree 166
nitrate, Effervescents..... 210	Quinia bimuriatica carbamidata 52
Pills..... 218, 226	Quinine and urea muriate..... 52
permanganate, Pills..... 223	-flower..... Sabbatia Elliottii 149
phosphate, Effervescents 210	hypophosphite, Syrups..... 223
salicylate, Elixirs..... 209	phosphate, Syrups..... 223
and manaca.. 118	sulphate..... 52
Potatoe, wild..... Man-root 120	Quinquina jaune.... Yellow cinchona 52
Potentilla canadensis Five-finger 79	rouge..... Red cinchona 52
Tormentilla... Tormentilla 182	Quinoline..... 247
Powder of Beef..... 238, 241	Quintefeuille..... Five-finger 79
Powdered Bone .. 238, 242	
Powdered Extracts, see respective drugs.	Racine de guimauve.. Marshmallow 120
Premna Taitensis..... Tonga 181	Ragweed... Ambrosia artemisiifolia 149
Prescription abbreviations..... 262	Ragwort..... Life-root 114
Prickly ash.	Raifort sauvage..... Horse-radish 97
Xanthoxylum americanum 145	Rainfarn..... Tansy 178
Pride of China..... Melia Azedarach 146	Raphidophora vitiensis..... 181
weed..... Fleabane 79	Rare alkaloids, etc..... 246
Prince's pine..... Pipsissewa 141	Raspberry..... Rubus strigosus 149
Primrose, evening..... 77	Rattlebush..... Wild indigo 155
Prinos, verticillatus Black alder 24	Rattlesnake master... Water eryngo 188
Privet..... Ligustrum vulgare 146	root Black cohosh 25
Prunin..... Wild cherry 194	Reagent Case, Physicians' Pocket 238, 245
Prunus serotina..... Wild cherry 193	Rectal alimentation..... 260
virginiana..... 193	Capsules..... 239
virginiana..... Choke cherry 51	Red alder..... Tag alder 177
U.S.P., Wild cherry 193	baneberry..... Red cohosh 150
Ptelea trifoliata..... Wafer ash 186	bark..... Red cinchona 52
Ptelein..... Wafer ash 187	bud..... Judas-tree 107
Pterocarpus Marsupium..... Kino 109	centaury... American centaury 8
Pterocaulon pycnostachyum.	chickweed... Scarlet pimpernel 160
Indian Black-root 99	cinchona... Cinchona succirubra 52
<i>Ptychotis Ajwan</i> 5, 276	clover..... Trifolium pratense 149
Puccoon, yellow..... Golden-seal 87	cohosh..... Actæa rubra 150
Püllna, Mineral-water Salts..... 237	maple..... Acer rubrum 150
Effervescents..... 210	oak..... Quercus rubra 150
Pulmonaria officinalis..... Lungwort 117	osier..... Cornus sericea 150
Pulsatilla, U. S. P.	pepper..... Capsicum 42
Anemone Pulsatilla 147	poppy..... Papaver Rhæas 151
Pulse, the..... 260	puccoon..... Blood-root 29
Pumpkin..... Cucurbita Pepo 147	root..... New Jersey tea 126
Punica Granatum..... Pomegranate 144	Resorcin..... 249
Purging agaric..... White agaric 189	Respiration..... 260
Purple avens..... Water avens 187	Reymond's Capsule Filler..... 240
Purslane..... Portulaca oleracea 147	Rhamnus cathartica,
Pycnanthemum montanum,	Buckthorn berries 35
Mountain-mint 124	Frangula, Buckthorn bark 35
Pyrèthre..... Pellitory 137	Purshiana,
Pyrethrum Parthenium... Feverfew 78	Cascara sagrada 44
U. S. P..... Pellitory 137	Rhatany..... Kraineria triandra 151
Pyridine..... 248	Rhein..... Rhubarb 153
Pyrole ombellée..... Pipsissewa 141	Rheum officinale..... Rhubarb 152
Pyrus malus..... Apple-tree 12	Rheumatism root..... Twin-leaf 184
	 Wild yam 196
<i>Quassia excelsa</i> 147	Rhododendron maximum,
U. S. P..... Picræna excelsa 147	Great laurel 89
Quassienholz..... Quassia 147	Rhubarb..... Rheum officinale 152
Quebracho,	Rhus aromatica..... 154
Aspidosperma Quebracho 148	glabra..... Sumach 174
Queckenwurzeln..... Couch-grass 66	Toxicodendron... Poison oak 143
Queen-of-the-meadow,	Rhusin..... Sumach 174
Eupatorium purpureum 148	Rich-weed..... Stone-root 172
Queen's delight..... Stillingia 171	Ricin..... Castor-oil plant 47
root..... Stillingia 171	Ricinus communis... Castor-oil plant 47
Quercitron bark..... Black oak 27	Ringelblume..... Calendula 38
Quercus alba..... White oak 191	Robin's rye..... Haircap moss 92
infectoria..... Nut-gall 127	Roborans plaster..... 236
rubra..... Red oak 150	Rochelle salt, Elixirs..... 206, 208
tinctoria..... Black oak 27	Rock-brake..... Polypody 144
	Rockbridge alum, Effervescents..... 210
	Roll plasters..... 236

	PAGE.		PAGE.
Roman chamomile.....	Chamomile 49	Sanguinarin.....	Blood-root 30
wormwood.....	Ragweed 149	Sanguinarine nitrate.....	Blood-root 30
Romarin des marais, Marsh rosemary	121	sulphate.....	Blood-root 30
Römischer Quendel.....	Thyme 180	Sanguis Bovinus Exsiccatus.....	238, 243
Rose bay.....	Willow herb 196	Santalum album.....	Sandalwood 156
colored silkweed.....		Santonica, U. S. P., Levant wormseed	112
White Indian-hemp	191	Santonin.....	Levant wormseed 112
pale.....	Rosa centifolia 154	and podophyllin.....	119
pink.....	American centaury 8	Capsules.....	204
red.....	Rosa gallica 154	Saponaria officinalis.....	Soapwort 166
willow.....	Red osier 150	Saraca indica.....	157
Rosemary.....	Rosmarinus officinalis 154	Sarothamnus Scoparius.....	33
Rosin-weed.....	Silphium laciniatum 155	Sarracenia flava.....	Trumpet plant 132
Rosmarinus officinalis.....	Rosemary 154	purpurea.....	Pitcher-plant 142
Rosshuf.....	Coltsfoot 60	Sarracenie.....	Pitcher-plant 142
Rothe chinarinde.....	Red cinchona 52	Sarsaparilla, American.....	10
miere.....	Scarlet pimpernel 160	Ponduras.....	
Rottlera tinctoria.....	108	Smilax officinalis 157	
Round-leaved cornel.....	Green osier 90	Mexican.....	
dogwood.....	Green osier 90	Smilax medica 157	
Rowan tree.....	Mountain-ash 123	Para, Smilax (sp. indet.) 157	
Royal flowering fern, Buckhorn brake	34	Texas.....	Yellow parilla 199
Rubus canadensis.....	Dewberry 275	U. S. P.....	157
Idæus, U. S. P.....	Raspberry 149	Sassafras officinale.....	Sassafras 158
strigosus.....	Raspberry 149	swamp.....	Magnolia 117
U. S. P.....	Blackberry 25	U. S. P., Sassafras officinale 158	
villosus.....	Blackberry 25	Sassy-bark.....	
Rudbeckia laciniata.....	Thimble-weed 180	Erythrophloeum guineense 158	
Rue.....	Ruta graveolens 155	Satureia hortensis.....	Summer savory 175
odorante.....	Rue 155	Sauge officinalis.....	Sage 156
Rumex Acetosella.....	Sheep sorrel 164	Saunders, white.....	Sandalwood 156
crispus.....	Yellow dock 198	yellow.....	Sandalwood 156
Rumicin.....	Yellow dock 198	Saurach.....	Barberry 16
Ruta graveolens.....	Rue 155	Savanna flower.....	Urechites suberecta 184
Sabal serrulata	159	Savin.....	Juniperus sabina 158
Sabbatia angularis.....		Savory, summer.....	175
American centaury 8		Saw Palmetto.....	Serenoa serrulata 159
campestris.....	155	Saxifrage.....	Pimpinella Saxifraga 159
Elliottii.....	Quinine-flower 149	Scale of Doses by age.....	256
Sabina, U. S. P.....	Juniperus sabina 158	Scammony, Convolvulus Scammonia 159	
Saccharin, Tablet triturates.....	232	Scarlet pimpernel, Anagallis arvensis 160	
Sacred bark.....	Cascara sagrada 44	Schafgarbe.....	Yarrow 198
-tree bark.....	Cascara sagrada 44	Schierlingsfrüchte.....	Conium 62
Safflower.....	Carthamus tinctorius 155	Schænocaulon officinale.....	49
Saffron, African.....	Safflower 155	Schöllkraut.....	Garden celandine 83
American.....	Safflower 155	Schwartzwurz.....	Comfrey 61
bastard.....	Safflower 155	Schwarze Nieswurz.....	
dyer's.....	Safflower 155	Black hellebore 27	
thistle.....	Safflower 155	schlangenwurz.....	
Sage.....	Salvia officinalis 156	Black cohosh 25	
garden.....	Sage 156	Scilla, U. S. P.....	Squill 169
mountain.....	124	Scillitoxin.....	Squill 169
Saint herb.....	Yerba santa 200	Sclerotic acid.....	Ergot 74
John's wort.....		Scoparius, U. S. P.....	Broom 33
Hypericum perforatum 156		Scotch broom.....	Broom 33
Salat des Jardins.....	Garden lettuce 83	Scouring rush.....	Equisetum hyemale 160
Salbei.....	Sage 156	Scrophularia nodosa.....	
Salicin.....	White willow 193	Carpenter's square 44	
Salicylic acid.....	White willow 193	Scullcap.....	Scutellaria lateriflora 160
comp. Elixir, Addenda 276		Scutellaria lateriflora.....	Scullcap 160
Salix alba.....	White willow 192	Scutellarin.....	Scullcap 160
nigra.....	Black willow 298	Sea lavender.....	Marsh rosemary 121
Salol, Pills.....	224	-wrack.....	Bladder-wrack 28
Salt-rheum weed.....	Balmoney 15	Secale cornutum.....	Ergot 74
Salts, Mineral-water.....	237	Seckelblumenwurz, New Jersey tea 126	
Salvia officinalis.....	Sage 156	Sedans Liquor.....	276
Sambucus canadensis.....	Elder 73	Sedative cerebral compound.....	237
nigra.....	European elder 77	Sedum acre.....	161
U. S. P.....	Elder 73	Seidenpflanze.....	Silkweed 165
Sampson snake-root.....		Seidlitz, Effervescent.....	210
Gentiana ochroleuca 156		Seifenrinde.....	Soap-tree 166
Sandalwood.....	Santalum album 156	Seifenwurz.....	Soapwort 166
Sanguinaria canadensis.....	Blood-root 29	Mineral-water Salts.....	237
		Simple's Atomizing Inhaler.....	243

	PAGE		PAGE
Senecio	Life-root 114	Soap-wort	Saponaria officinalis 166
Senecio aureus	Life-root 114	Socotrine aloes	Aloes 6
<i>gracilis</i>	114	Soda Chlorinated solution, U. S. P.	241
Senecion	Life-root 114	-mint tablets	230
Senega, U. S. P.	Polygala Senega 161	and potassium tartrate, Elixirs	208
Senna (Alexandrian),		Sodium and potassium tartrate,	
<i>Cassia acutifolia</i> 162		Elixirs, 206, 208	
Serenoa serrulata	Saw palmetto 159	arseniate,	
Serpentaire de Virginie	Serpentaria 164	Hypodermic tablets 229	
Serpentaria, Aristolochia Serpentaria 164		Tablet triturates	232
<i>Serronia Jaborandi</i>	106	bicarbonate,	
Sesame	Benne 21	Compressed tablets 230	
Sesamum orientale	Benne 21	Pills	219, 224, 225
Seven barks, Hydrangea arborescens 98		Tablet triturates	231
Shamrock	White clover 190	Lozenges	233
Shave-grass	Scouring-rush 160	Elixirs	209
Sheep laurel	Mountain laurel 123	carbonate, Pills	218, 220
sorrel	Rumex Acetosella 164	choleate	249
Shell-bark hickory	Carya alba 164	hypophosphite, Glyceroles	211
-flower	Balmomy 15	Malt ext.	
Shepherd's purse,		prep's	213
<i>Capsella bursa-pastoris</i> 165		Syrups	228, 229
Shotbush	American sarsaparilla 10	lactophosphate, Malt ext.	
Slarubby trefoil	Wafer ash 186	prep's	213
Side-saddle flower	Pitcher-plant 142	Syrups	228, 229
plant	Trumpet plant 182	nitrite	249
Sierra salvia	Mountain sage 224	phosphate, Effervescent	210
Silkweed	Asclepias Cornuti 165	Syrups	229
rose-colored		salicylate, Elixirs	209
White Indian-hemp 191		Pills	225
swamp, White Indian-hemp 191		Tablet triturates	232
Silky cornel	Red osier 150	and manaca	118
<i>Silphium gummiferum</i>	155	sulphite, Glyceroles	211
<i>laciniatum</i>	Rosinweed 155	sulpho-carbolate	249
Silver iodide, Pills	224	tungstate	249
leaf	Stillingia 171	Solanum Dulcamara	Bitter-sweet 24
oleate	235	paniculatum	Jurubeba 275
Silybum marianum	Mary thistle 121	Solidago odora	Golden-rod 87
Simaba cedron	Cedron seed 48	Solid Extracts, see respective drugs.	
Simaruba	Simaruba officinalis 165	Solomon's seal,	
Simaruba officinalis	Simaruba 165	<i>Polygonatum officinale</i> 167	
Simple Elixir	209	Soluble Elastic Filled Capsules	203
Simpler's joy, American blue vervain 8		Solution aluminium acetate	241
Skunk cabbage,		sclerotic acid	74
<i>Symplocarpus foetidus</i> 165		Sommerwurzel	Beech-drop 18
weed	Skunk cabbage 165	<i>Sophora tinctoria</i>	195
Slippery elm	Ulmus fulva 166	<i>Sorbus americana</i>	123
Small burnet saxifrage	Saxifrage 159	Sorrel	Sheep sorrel 164
spikenard,		tree	Sourwood 167
American sarsaparilla 10		Souci	Calendula 38
Smartweed	Water-pepper 189	Sourwood	Oxydendrum arboreum 167
Smilax herbacea	Carrian-flower 44	Southernwood, Artemisia Abrotanum 167	
medica,		Spanischer Pfeffer	Capsicum 42
<i>Sarsaparilla, Mexican</i> 157		Spanish chamomile	Pellitory 137
<i>officinalis</i> ,		needles	Bidens bipinnata 167
<i>Sarsaparilla, Honduras</i> 157		Spargel	Asparagus 14
<i>peduncularis</i> ,		Sparteine sulphate	Broom 33
Smilax herbacea 44		Spatterdock	Yellow pond-lily 199
Pseudo-China, Bamboo brier 16		Spearmint	Mentha viridis 167
<i>sarsaparilla</i> ,		Specialties	238
Smilax Pseudo-China 16		Specific volume	255
Smooth alder	Tag alder 177	Speedwell	Veronica officinalis 168
Snake head	Balmomy 15	Spice-bush	Lindera benzoin 168
lily	Blue flag 31	-tree	California laurel 39
root, black	Black cohosh 25	-wood	Spice-bush 168
button	37	Spigelia marilandica	Pink-root 141
Canada	Wild ginger 195	Spikenard	Aralia racemosa 168
Sampson	156	Spindle tree	Wahoo 187
Virginia	Serpentaria 164	Spiny burreed	Spiny clotbur 168
white	192	clotbur	Xanthium spinosum 168
weed	Bistort 23	cocklebur	Spiny clotbur 168
Snapping hazel	Witch hazel 197	Spiraea tomentosa	Hardhack 92
Soap-bark	Soap-tree 166	Spirit myrcia, U. S. P.	Bay laurel (2) 17
-root	Soapwort 166	nitrous ether, Elixirs	305
-tree	Quillaia saponaria 166		

	PAGE.		PAGE.
Spiritus Armoracæ comp., B. P.		Supporter, Wilson's Eureka Abdom-	
Horse-radish	98	inal	238, 246
Spoonwood	123	Süssholz wurzel	113
Spotted alder	197	Licorice	113
Witch-hazel	197	Suter berry	145
cowbane	188	Swamp dogwood	150
Water hemlock	188	Red osier	150
geranium	66	Swamp hellebore	9
Cranesbill	66	American hellebore	9
parsley	188	maple	150
Water hemlock	188	Red maple	150
Water hemlock	188	milkweed	
White agaric	189	White Indian-hemp	191
Spunk	80	Magnolia	117
Spurge, flowering	80	sassafras	117
Spurred rye	74	silkweed, White Indian-hemp	191
Ergot	74	Sweet root	1
Squaw-root	30	Abscess root	1
Black cohosh	25, 30	Sweet balm	112
-mint	137	Lemon balm	112
Pennyroyal	137	basil	175
-vine	169	Ocimum	175
Mitchella repens	169	Basicum	175
Life-root	114	bay	17
Squill	169	Bay laurel (1)	17
Urginea Scilla	169	basil	117
Squirrel-corn	182	Magnolia	117
Turkey-corn	182	birch	35
Staffvine	77	Black birch	35
False bittersweet	77	bugle	25
Staggerweed	182	Bugleweed	25
Turkey-corn	182	cicely	175
Staphisagria, U. S. P.	170	Osmorrhiza longistylis	175
Stavesacre	170	clover	176
Star, Mineral-water Salts	237	Melilotus officinalis	176
grass	170	fern	176
Alettris farinosa	170	Comptonia asplenifolia	176
wort	93	flag	176
Heloniar	93	Acorus Calamus	176
Star-grass	170	gale	176
Statice brasiliensis	18	Myrica Gale	176
Baycuru	18	gum, Liquidambar styraciflua	176
caroliniana, Marsh rosemary	121	marjoram, Origanum Majorana	177
Stavesacre, Delphinium Staphisagria	170	orange	177
Steeple-bush	92	Citrus Aurantium	177
Hardhack	92	sumach	154
Steinklee	176	Sweetbread	134
Sweet clover	176	Cocillana	57
Stellaire	51	Sycocarpus Rusbyi	57
Chickweed	51	Symplocarpus foetidus	
Stellaria media	51	Skunk cabbage	165
Sterculia acuminata	109	Symphytum officinale	61
Chickweed	51	Comfrey	61
Sternkraut	51	Syringes	238, 244
Stickwort agrimony	5	Syrups, medicinal	228
Sticta pulmonaria	116	Syrup Trifolium Comp.	150
Lungmoss	116	Red clover	150
Stigmata Maidis	64	Syzgium jambolanum	106
Corn	64		
Stillingia	171	Tabacum, U. S. P.	180
Stillingia sylvatica	171	Table, graduated, for administering	
Stillingia	171	laudanum	261
Stillingin	171	of drops to fluidrachm.	257
Stinging nettle	125	of quantities not to be pre-	
Stonecrop	161	scribed without caution	
Sedum acre	161	mark	238
mossy	161	of prescription abbreviations	262
Virginia	186	of thermometric equivalents	255
Stone-root	172	of weights and measures	250
Collinsonia canadensis	172	Tablet triturates	221
Stramonium	172	Tablets, compressed	230
Strandvelke	121	hypodermic	229
Strawberry	174	Tag alder	177
Strophanthus Kombe	174	Alnus serrulata	177
Strychnine	128	Tamarac	177
Nux vomica	128	Larix americana	177
Malt ext. preparations	213	Tanacetum vulgare	178
hypophosphite, Syrups	238	Tanaisie	178
phosphate, Syrups	228	Tansy	178
Strychnos	95, 276	Tannic acid	127
Gaultheriana	95, 276	Nut-gall	127
Ignatii	98	Collodions	227
malaccensis		Glyceroles	211
Hoang-nan	95, 276	Lozenges	233
Nux vomica	127	Pills	225, 226
Sturmhut	1	Tansy	178
Aconite	1	Tanacetum vulgare	178
Stylosanthes elatior	137	Tar	70
Succus Alterans	16	Dandelion	70
Sugar Test Flasks	238, 243	Taro flour	179
Tests for	246	Tartar emetic, see also Antimony and	
Sulphur, Pills	220	Potassium tart.	
Bricks, Disinfectants	241	Pills	226
iodide, Pills	225	Tablet triturates	232
ointment, U. S. P.	236	Tartaric acid, Pills	226
Sumach	174	Tausendgueldenkraut,	
Rhus glabra	174	European centaury	76
Chinese, Ailantus glandulosa	5	Camellia Thea	179
sweet	154	New Jersey	126
Rhus aromatica	154	Teaberry	196
vénéneux	143	Wintergreen	196
Poison oak	143		
Sumbul, U. S. P.	125		
Musk-root	125		
Summer savory	175		
Satureia hortensis	175		
Sundew	175		
Drosera rotundifolia	175		
Sunflower	175		
Helianthus annuus	175		

	PAGE.		PAGE.
Tepopote.... Ephedra antisiphilitica	74	Trumpet plant..... <i>Sarracenia flava</i>	182
Terebene..... Turpentine	183	weed,	
Capsules.....	204	Queen-of-the-meadow	148
Inhalants.....	212	<i>Tsuga canadensis</i> Hemlock spruce	93
Terpin hydrate..... Turpentine,	183,	Tulip tree.... <i>Liriodendron tulipifera</i>	182
Terra japonica..... Catechu	47	Turkey-corn.... <i>Dicentra canadensis</i>	182
Test flasks, Sugar.....	243	pea, wild..... Turkey corn	182
papers, Urinary.....	238,	Turlington's balsam..... Benzoin	22
tablets, Sugar.....	238	Turmeric..... <i>Curcuma longa</i>	183
Urinary.....	238,	Indian..... Golden-seal	87
Tests for albumen.....	246	<i>Turnera aphrodisiaca</i>	69
for sugar.....	246	microphylla..... Damiana	69
Tetra-iodolpyrol.....	248	Turpentine..... <i>Pinus australis</i>	183
Tetterwort..... Blood-root	29	Chian.....	51
Teucrium Scordium..... Garden celandine	83	Turtle bloom..... Balmoney	15
Water germander	188	head..... Balmoney	15
Texas sarsaparilla.... Yellow parilla	199	Tussilago Farfara..... Coltsfoot	60
Thaleictchys pacificus.....	76	Twin-leaf..... <i>Jeffersonia diphylla</i>	184
Thalline.....	249		
sulphate.....	249	<i>Ulmus fulva</i> Slippery elm	166
tartrate.....	249	Umbellularia californica,	
Thapsia garganica.....	180	California laurel	39
Therapeutic Suggestions.....	265	Unicorn root..... Star-grass	170
Theriac confection.....	235	false..... Helonias	93
Thermometric Equivalents, Table of	255	Unkum..... Life-root	114
Thermometry.....	259	Upland sumach..... Sumach	174
Thimble-weed... <i>Rudbeckia laciniata</i>	180	Uranium nitrate.....	249
Thistle, blessed.....	29	Urea and quinine muriate.....	249
Canada.....	40	Urechites suberecta.....	184
Mary.....	121		238, 244
saffron..... Safflower	155	Urethane.....	249
Thorn apple..... Stramonium	172	Urginea Scilla..... Squill	169
Thoroughwort..... Boneset	32	Urinary Test Papers.....	238, 245
Thuja, U. S. P..... Arbor vitæ	12	Tablets.....	238, 246
Thuya occidentalis..... Arbor vitæ	12	Urtica dioica..... Nettle	126
Thyme..... <i>Thymus vulgaris</i>	180	Ustilago, U. S. P..... Corn	64
Thymol..... Thyme	180	Utero-vaginal Syringe, Chamberlin's	244
Thymus vulgaris..... Thyme	180	Uva-ursi.... <i>Arctostaphylos</i>	184
Tick-weed..... Pennyroyal	137		
Tilia americana..... Basswood	17	<i>Vaccinium crassifolium</i>	185
glabra.....	17	Vaginal capsules.....	203
Tin oleate.....	235	Valerian..... <i>Valeriana officinalis</i>	185
Tobacco..... Nicotiana Tabacum	180	American... Ladies' slipper	110
Indian..... Lobelia	115	<i>Valeriana officinalis</i> Valerian	185
wild..... Lobelia	115	Vanilla leaf..... Deer-tongue	71
Tollkirsche..... Belladonna	18	Varech versiculeux... Bladder-wrack	28
Tolu..... <i>Myroxylon Toluiferum</i>	181	Venushaar..... Maiden-hair	117
Tomato... <i>Lycopersicum esculentum</i>	181	Veratre vert.... American hellebore	9
Tonca-bean..... <i>Dipterix odorata</i>	181	Veratrin..... American hellebore	10
Tonga.....	181	Veratrin..... American hellebore	10
Tonka-bean..... Tonca-bean	181	Veratrine..... Cevadilla	49
Tornestilla... <i>Potentilla Tormentilla</i>	182	Veratrum album... White hellebore	191
Toothache tree..... Prickly ash	145	Sabadilla..... Cevadilla	49
Tooth wash..... Soap-tree	166	viride, American hellebore	9
Touch-me-not..... Jewel-weed	107	Verbascum Thapsus..... Mullein	124
Touchwood..... White agaric	189	Verbena hastata,	
Trailing arbutus..... Gravel-plant	89	American blue vervain	8
Traubenkraut..... Ragweed	149	urticefolia... White vervain	192
Tree of heaven... <i>Ailantus glandulosa</i>	5	Vergerette..... Fleabane	79
Trèfle blancourampant, White clover	190	Veronica officinalis..... Speedwell	168
d'eau (de marais).... Buckbean	34	virginica..... Culver's	68
Trichloracetic acid.....	247	Veronique mâle..... Speedwell	168
Trifolium compound, Syrup,		Verschiedenfarbige schwertlilie,	
Red clover	150	Blue flag	31
pratense..... Red clover	149	Vervain..... American blue vervain	8
repens..... White clover	190	nettle-leaved, White vervain	192
Trilisa odoratissima... Deer-tongue	71	white.....	192
Triblin..... Beth-root	22	Veterinary capsules.....	203
Trillium erectum..... Beth-root	22	Viburnin..... Cramp-bark	66
Trifolium repens..... Couch-grass	66	Viburnum Opulus..... Cramp-bark	66
Trixis fruticosa.....	247	prunifolium... Black haw	26
Troches, aromatic.....	238	U. S. P.....	26
compressed.....	230	Vichy, Mineral-water Salts.....	237
Troy Ounces into Avoirdupois Ounces	254		

	PAGE.		PAGE.
Vichy, Effervescent.....	210	White water-lily.....	White pond-lily 192
Vigne vierge.....	American ivy 10	willow.....	Salix alba 192
Vine, maple.....	Yellow parilla 199	-wood.....	Basswood 17
Viola tricolor.....	Pansy 135		Tulip tree 182
Virginia creeper.....	American ivy 10	Whorehouse tea,	
sarsaparilla.....	Bamboo brier 16	Ephedra antisiphilitica	74
snake-root.....	Serpentaria 164	Wild allspice.....	Spice bush 168
stone-crop,		bergamot.....	Monarda fistulosa 193
Penthorum sedoides	186	celandine.....	Jewel-weed 107
Virginische Schlangenzwurz,		chamomile.....	Mayweed 122
Serpentaria	164	cherry.....	Prunus serotina 193
Viscum flavescens,		cinnamon.....	Bay laurel (2) 17
Phoradendron flavescens	123	clove.....	Bay laurel (2) 17
Volume, Specific.....	235	elder.....	Dwarf elder 72
		ginger.....	Asarum canadense 195
		hippo.....	Flowering spurge 80
Wachsbau.....	Bayberry 17	hydrangea,	
Wachsgagel.....	Bayberry 17	Hydrangea arborescens	98
Wafer ash.....	Ptelea trifoliata 186	hyssop, American blue vervain	8
Wahoo.....	Euonymus atropurpureus 187	indigo.....	Baptisia tinctoria 195
Wake-robin.....	Beth-root 22	ipecac.....	Flowering spurge 80
	Indian turnip 101	jalap.....	Man-root 120
Wall-pepper.....	Sedum acre 161	lettuce.....	Lactuca canadensis 195
Walnut, black.....	28	potatoe.....	Man-root 120
Wandering milkweed.....	Bitter-root 23	rose bay.....	Great laurel 89
Warburg's tincture.....	52, 54	sarsaparilla,	
	Pills,.....225, 276	American sarsaparilla	10
Wasser fenchel.....	Water fennel 188	succory.....	American centaury 8
Water avens.....	Geum rivale 187	tobacco.....	Lobelia 115
-cup.....	Pitcher-plant 142	turkey pea.....	Turkey-corn 182
dropwort.....	Water fennel 188	yam.....	Dioscorea villosa 196
eryngo,		Wildes Schoellkraut.....	Jewel-weed 107
Eryngium yuccifolium	188	Wildkirschenrinde.....	Wild cherry 193
fennel,		Willow, black,	28
Enanthie Phellandrium	188	-herb,	
germander,		Epilobium angustifolium	196
Teucrium Scordium	188	rose.....	Red osier 150
hemlock.....	Cicuta maculata 188	white.....	192
five-leaved,		Wilson's Eureka Abdominal Sup-	
Water fennel	188	porter.....	238, 246
horehound.....	Bugleweed 25	wines, medicinal.....	234
-lily, white.....	White pond-lily 192	Wingseed.....	Wafer ash 186
-melon.....	Citrullus vulgaris 189	Winterberry.....	Black alder 24
pepper.....	Polygonum acre 189	Winter-bloom.....	Witch-hazel 197
smartweed.....	Water-pepper 189	-clover.....	Squaw-vine 169
Wax-myrtle.....	Bayberry 17	pink.....	Gravel-plant 89
work.....	False bittersweet 77	Schachtelhalm, Scouring-rush	160
Weather-glass.....	Scarlet pimpernel 160	Wintergreen, Gaultheria procumbens	196
Weight of any given measure, to		Pipsissewa	141
ascertain.....	254	Winter's bark.....	Drims Winter 196
Weinachtswurzel.....	Black hellebore 27	Witch-hazel.....	Hamamelis virginica 197
Wernuth.....	Wormwood 197	Wolfsbane.....	Aconite 1
White agaric.....	Polyporus officinalis 189	Wolfsmilchwurzel, Flowering spurge	80
baneberry.....	White cohosh 190	Wollkraut.....	Mullein 124
bay.....	Magnolia 117	Wood betony.....	Betonica officinalis 197
bryony.....	Bryonia alba 190	-bine.....	American ivy 10
cedar, false.....	Arbor vitæ 12	-oil.....	Gurjun balsam 92
clover.....	Trifolium repens 190	Woody nightshade.....	Bittersweet 24
cohosh.....	Actæa alba 190	Worara.....	Curare 69
flag.....	Orris 133	Worm-grass.....	Pink-root 141
hellebore.....	Veratrum album 191	Wormseed, American.....	10
Indian-hemp,			Jerusalem oak 107
Asclepias incarnata	191	Wormwood.....	Artemisia Absinthium 197
-leaf.....	Hardhack 92	Roman.....	Ragweed 149
oak.....	Quercus alba 191	Wurmfarnwurzel.....	Male-fern 117
pine.....	Pinus Strobus 191	Wurmwidriger.....	Jerusalem oak 107
pond-lily.....	Nymphaea odorata 192		
poplar.....	Populus tremuloides 192	Xanthium spinosum.....	Spikenard 168
saunders.....	Sandalwood 156	strumarium.....	Clothbur 55
snakeroot,		Xanthoxylum.....	Prickly ash 146
Eupatorium aromaticum	192		Prickly ash 145
trefoil.....	White clover 190	Xanthoxylum americanum,	
vervain.....	Verbena urtica-folia 192		Tar 178
walnut.....	Butternut 37		Capsules.....204, 205

	PAGE.		PAGE.
Yakona.....	Kava Kava 108	Yerba maté.....	Paraguay tea 135
Yam, wild.....	196	reuma, <i>Frankenia grandifolia</i>	200
Yarrow.....	<i>Achillea Millefolium</i> 198	santa, <i>Eriodictyon glutinosum</i>	200
Yaw root.....	<i>Stillingia</i> 171	Yerbine compound, <i>Glyceroles</i>	211
Yellow cinchona..	<i>Cinchona Calisaya</i> 52	Ysop.....	Hyssop 98
dock.....	<i>Rumex crispus</i> 198		
-flowered nightshade,			
<i>Urechites suberecta</i>	184		
jasmine.....	<i>Gelsemium</i> 84	Zahnwehholz.....	Prickly ash 144
jessamine.....	<i>Gelsemium</i> 84	Zaunrübe.....	White byrony 195
leaf-cup.....	<i>Bearsfoot</i> 18	Zea Mays.....	Corn 60
melilot.....	Sweet clover 176	Zinc carbonate, Cerate.....	236
oxide of mercury, Com-		oleate.....	235
pressed tablets.....	230	oxide, Pills.....	221, 225
parilla,		Ointment, U. S. P.....	236
<i>Menispermum canadense</i>	199	phosphide, Pills.....	225
pond-lily.....	<i>Nuphar advena</i> 199	Tablet triturates....	232
poplar.....	Tulip tree 182	sulphate, Pills.....	223
puccoon.....	Golden-seal 87	valerianate, Elixirs.....	209
root.....	Golden-seal 87	Pills, 215, 222, 223, 224,	225
saunders.....	Sandalwood 156	Zingiber officinale...Ginger Jamaica	86
snake-leaf..	Adder's tongue 4	Zittwersamen.....	Levant wormseed 112
Yerba buena..	<i>Micromeria Douglasii</i> 199		
mansa,			
<i>Houttuynia californica</i>	199		

Date Due

Demco 293-5

RS 154
838P