ROYAL ONTARIO MUSEUM LIBRARIES

3 1761 04305 7165

ANNOTATED LIST OF ONTARIO LEPIDOPTERA

J. C. E. RIOTTE

Digitized by the Internet Archive
in 2012 with funding from
Royal Ontario Museum

<http://archive.org/details/annotatedlistof00riot>

J. C. E. RIOTTE

**ANNOTATED LIST
OF ONTARIO
LEPIDOPTERA**

ROYAL ONTARIO MUSEUM
LIFE SCIENCES
MISCELLANEOUS PUBLICATIONS

ROYAL ONTARIO MUSEUM
PUBLICATIONS IN LIFE SCIENCES

The Royal Ontario Museum publishes three series in the Life Sciences:

CONTRIBUTIONS: a numbered series of original scientific publications.

OCCASIONAL PAPERS: a numbered series of original scientific publications, primarily short and of taxonomic significance.

MISCELLANEOUS PUBLICATIONS: an unnumbered series on a variety of subjects.

All manuscripts considered for publication are subject to the scrutiny and editorial policies of the Life Sciences Editorial Board, and to independent refereeing by two or more persons, other than Museum staff, who are authorities in the particular field involved.

LIFE SCIENCES EDITORIAL BOARD

Senior editor: M. D. Engstrom

Editor: D. Collins

Editor: K. Coates

External editor: C. S. Churcher

Manuscript editor: Judith Eger

Production editor: K. K. Mototsune

J. C. E. Riette is Research Associate, Department of Entomology, Bishop Museum, Honolulu, Hawaii, and research associate, Department of Entomology, Royal Ontario Museum.

The Royal Ontario Museum is an agency of the Ontario Ministry of Culture and Communications.

Canadian Cataloguing in Publication Data

Riette, J. C. E.

Annotated list of Ontario lepidoptera

(Life sciences miscellaneous publication)

Includes bibliographical references and index.

ISBN 0-88854-397-2

I. Lepidoptera - Ontario. I. Royal Ontario Museum.

II. Title. III. Series.

QL552.R56 1992 595.78'09713 C91-095769-X

Publication date: June 1992

ISBN 0-88854-397-2

ISSN 0082-5093

© 1992 Royal Ontario Museum

100 Queen's Park, Toronto, Canada M5S 2C6

Typesetting by True to Type Inc.

Printed and bound in Canada

Cover: Luna moth, *Actias luna*

PS
To
160
R56

TABLE OF CONTENTS

Figure 1	viii
Introduction	1
Materials and Methods	6
List of Species	9
Order Lepidoptera	9
Suborder Zeugloptera	9
Superfamily Micropterigoidea	9
Family Micropterigidae	9
Suborder Exoporia	9
Superfamily Hepialoidea	9
Family Hepialidae	9
Suborder Monotrysia	10
Superfamily Nepticuloidea	10
Family Nepticulidae	10
Superfamily Incurvarioidea	10
Family Incurvariidae	10
Subfamily Incurvariinae	10
Subfamily Adelinae	10
Suborder Ditrysia	11
Superfamily Tineoidea	11
Family Psychidae	11
Subfamily Taleporiinae	11
Subfamily Psychinae	11
Superfamily Gelechioidea	11
Superfamily Copromorphoidea	11
Family Alucitidae	11
Superfamily Yponomeutoidea	11
Family Yponomeutidae	11
Superfamily Sesiioidea	12
Family Sesiidae	12
Subfamily Tinthiinae	12
Subfamily Paranthreninae	12
Subfamily Sesiinae	12
Family Choreutidae	14
Subfamily Choreutinae	14
Superfamily Cossoidea	14
Family Cossidae	14
Subfamily Cossinae	14

Superfamily Tortricoidea	15
Superfamily Hesperioidea	15
Family Hesperiidae	15
Subfamily Pyrginae	15
Subfamily Heteropterinae	17
Subfamily Hesperinae	18
Superfamily Papilionoidea	22
Family Papilionidae	23
Subfamily Papilioninae	23
Family Pieridae	24
Subfamily Pierinae	24
Subfamily Anthocharinae	26
Subfamily Coliadinae	26
Family Lycaenidae	28
Subfamily Liphyrinae	28
Subfamily Lycaeninae	28
Subfamily Eumaeinae	29
Subfamily Polyommattinae	32
Family Riodinidae	34
Family Libytheidae	34
Subfamily Libytheinae	34
Family Nymphalidae	34
Subfamily Nymphalinae	34
Subfamily Argynninae	37
Subfamily Melitaeinae	39
Subfamily Limenitidinae	41
Subfamily Charaxinae	42
Family Satyridae	42
Subfamily Elymniinae	42
Subfamily Satyrinae	43
Family Danaidae	45
Subfamily Danainae	45
Superfamily Zygaenoidea	45
Family Zygaenidae	45
Family Limacodidae	45
Superfamily Pyraloidea	47
Family Pyralidae	47
(Series Crambiformes)	47
Subfamily Scopariinae	47
Subfamily Nymphulinae	48
Subfamily Odontiinae	49

Subfamily Glaphyriinae	49
Subfamily Evergestinae	50
Subfamily Pyraustinae	50
Subfamily Schoenobiinae	54
Subfamily Crambinae	54
(Series Pyraliformes)	57
Subfamily Pyralinae	57
Subfamily Chrysauginae	58
Subfamily Epipaschiinae	58
Subfamily Galleriinae	59
Subfamily Phycitinae	59
Subfamily Peoriinae	63
Superfamily Thyridioidea	64
Family Thyrididae	64
Subfamily Thyridinae	64
Superfamily Pterophoroidea	64
Family Pterophoridae	64
Subfamily Platyptilinae	64
Subfamily Pterophorinae	65
Superfamily Drepanoidea	66
Family Thyatiridae	66
Subfamily Thyatirinae	66
Family Drepanidae	67
Subfamily Drepaninae	67
Subfamily Oretinae	68
Superfamily Geometroidea	68
Family Geometridae	68
Subfamily Archiearinae	68
Subfamily Oenochrominae	68
Subfamily Geometrinae	68
Subfamily Sterrhinae	70
Subfamily Larentiinae	71
Subfamily Ennominae	79
Family Epiplemidae	95
Superfamily Mimallonoidea	96
Family Mimallonidae	96
Superfamily Bombycoidea	96
Family Apatelodidae	96
Subfamily Apatelodinae	96
Family Lasiocampidae	97
Subfamily Macromphaliinae	97

Subfamily Gastropachinae	97
Subfamily Lasiocampinae	98
Family Saturniidae	99
Subfamily Citheroniinae	99
Subfamily Hemileucinae	101
Subfamily Saturniinae	101
Superfamily Sphingoidea	104
Family Sphingidae	104
Subfamily Sphinginae	104
Subfamily Macroglossinae	109
Superfamily Noctuoidea	112
Family Notodontidae	112
Subfamily Pygaerinae	112
Subfamily Phalerinae	113
Subfamily Notodontinae	114
Subfamily Cerurinae	120
Family Arctiidae	121
Subfamily Lithosiinae	121
Subfamily Arctiinae	123
Subfamily Ctenuchinae	131
Family Lymantriidae	131
Subfamily Orgyiinae	131
Subfamily Lymantriinae	133
Family Noctuidae	133
Subfamily Herminiinae	133
Subfamily Hypenodinae	135
Subfamily Hypeninae	135
Subfamily Ophiderinae	136
Subfamily Catocalinae	138
Subfamily Plusiinae	141
Subfamily Euteliinae	143
Subfamily Sarrothripinae	143
Subfamily Nolinae	143
Subfamily Acontiinae	144
Subfamily Pantheinae	145
Subfamily Acronictinae	146
Subfamily Agaristinae	147
Subfamily Amphipyridae	148
Subfamily Cuculliinae	154
Subfamily Hadeninae	158
Subfamily Noctuinae	162

Subfamily Heliothinae	167
Acknowledgements	168
Literature Cited	169
Index of Scientific Names	176

FIGURE 1. The four main faunal regions of Ontario. The 0°C annual mean temperature isotherm is indicated, and has been adopted here as the southern limit of northern Ontario. See Introduction for discussion.

INTRODUCTION

This publication summarizes the current knowledge concerning the distribution of Ontario Lepidoptera, exclusive of the Copromorphoidea (in part), Gelechioidea, Tineoidea (in part), Tortricoidea, and Yponomeutoidea (in part). These superfamilies have been excluded because they have not yet been sufficiently studied and are not represented by reliably identified specimens in most collections.

In 1959 I published a "Revision of C. J. S. Bethune's list of the butterflies of the eastern provinces of Canada as far as northern Ontario is concerned" (Riotte, 1959). That list, which has since been revised (Riotte, 1962b, 1971a), was to be the first instalment of a distributional list of all Ontario Lepidoptera. My subsequent studies of the Lepidoptera of southern Ontario resulted in a checklist of Ontario skippers and butterflies (Riotte, 1970a). This present publication compiles the collection records of various institutions, the seasonal summaries published by the Toronto Entomologists' Association for 1969–1988, and other regional lists.

Ontario, the second largest Canadian province, extends through 15 degrees of latitude and encompasses the widest range of environments of any Canadian province, ranging from tundra on the shores of Hudson Bay to temperate deciduous forests on the northern shores of Lake Erie and Lake Ontario (see Munroe, 1956). Ontario's four main faunal regions (Snyder, 1939), shown in Figure 1, are based on both climatic and geologic factors. Region 4 encompasses the whole of southern Ontario, and the line A–B (Figure 3) represents the approximate northern limit of the Carolinian Zone in Ontario. Region 3, the boreal forest region, extends south of the 0° C annual mean temperature isotherm to the climatic boundary separating the Laurentian Plateau from the Lower Great Lakes Region (Figure 2, line C–D). Region 2, the taiga, is the next most northern region and is bounded in the south by the 0° C annual mean temperature isotherm. Region 1, the tundra, is restricted to a narrow band along the shores of Hudson Bay and James Bay.

It should be noted that the boundary between the deciduous and boreal forest regions is discontinuous. For example, the boundary between white pine (*Pinus strobus*)/hemlock (*Tsuga canadensis*) forests and the mixed forests of the northern hardwoods region traces the following lines: extends from southern Ontario over southcentral into northcentral Ontario; meets the eastern shore of Lake Superior at about Michipicoten; is found on Michipicoten Island in Lake Superior; is seen again along the western shore southwest of Thunder Bay (Neebing Township) where it apparently penetrates from Minnesota; and then continues from about the Nipigon River westward into Manitoba (Lake of the Woods). The western part of the region, although lying between 48° and 49° 36' north latitude, resembles more closely the southern section in the east below 46° 36' north latitude than the corresponding northern latitudes of the eastern part of the region.

FIGURE 2. Districts and selected localities in central, western, and northern Ontario. The line C-D approximates the 4.5° C annual mean daily temperature isotherm, and has been adopted here as the northern limit of southern Ontario.

FIGURE 3. Counties, districts, regional municipalities, and selected localities in southern Ontario. The dashed line indicates the approximate southern and eastern limits of the Canadian Shield in Ontario. The line A-B represents the approximate northern limit of the Carolinian Zone in Ontario.

Temperature seems to be the primary environmental factor that determines the faunal regions of Ontario. Table 1 shows representative temperatures for localities listed in Figures 2 and 3 arranged from north to south. For detailed information see the publications of the Canada Department of Transport, Meteorological Division (e.g., 1947, 1954, 1964).

Needless to say, these climatic differences strongly influence lepidopteran populations. It has been suggested that the climate of Ontario has become generally milder (Baker, 1955; Miller, 1956; Urquhart, 1957). This may explain the occasional newcomers to the fauna of southwestern Ontario and the extension of well-established southern species' ranges into the north (e.g., Munroe, 1957). On the other hand, northern influences and relicts are to be found far to the south in the acid bogs which, although numerous in the boreal forest region, are also found singly in the deciduous forest region, e.g., near Hamilton and London (Byron Bog).

There are fewer species encountered the farther north one travels. The

TABLE 1. Averages of daily mean temperatures in degrees Celsius.

Locality	January	July	Year
Trout Lake (Fort Severn)	-23.5	16.0	-3.0
Fort Frances	-16.0	20.0	3.3
Sioux Lookout	-19.4	18.3	1.3
Lansdowne House	-22.4	17.2	-1.4
Moose Factory (Moosonee)	-21.2	15.5	-1.7
Nakina	-18.8	16.5	0.2
Hornepayne	-20.0	15.0	0.0
Longlac (Geraldton)	-19.4	16.5	0.2
Nipigon	-13.8	17.2	1.1
Biscotasing	-16.6	16.5	0.0
Sault Ste. Marie	-11.1	17.2	3.3
Sudbury	-12.2	18.9	3.9
Manitoulin Island	- 8.8	17.7	4.4
Algonquin Park	-11.6	17.7	3.3
Ottawa	-11.1	21.1	5.6
Kingston (Chaffey's Locks)	- 7.2	20.0	6.6
Trenton	- 7.2	21.1	7.2
Muskoka (meteorological observation station)	-10.0	18.9	5.2
Orillia	- 8.3	20.0	5.6
Toronto	- 5.0	20.6	7.2
Owen Sound	- 6.6	20.6	7.2
Guelph	- 6.6	20.0	6.6
St. Catharines	- 3.3	21.6	8.9
Chatham	- 3.8	22.8	8.9
Pelee Island	- 3.8	23.9	9.4

greatest number of species is found in the deciduous forests of southwestern Ontario where several southern species reach their northern limit. But the mixed forests of southern Ontario are hardly "poor in species"; most of the species living in Ontario are represented there, too. Some species apparently do not cross the southern frontier of this region.

The boreal forests of central and western Ontario and of the southern parts of northern Ontario are still comparatively rich in moth species, but species richness, especially in the notodontids, decreases rapidly to the north. In fact, a reduction in species is already noticeable in the northeastern part of the mixed forest region. Most species are restricted to the boreal forests, but some species are collected on occasion in more southerly localities.

The presence of mixed forests west of Lake Superior—characterized by sumac (*Rhus* sp.), oak shrubs (*Quercus* sp.), and maple trees (*Acer* spp.)—as far north as Kenora, and of maple shrubs as far north as Macdiarmid, lends a special character to the fauna and flora of Ontario. These forests continue westward into Manitoba and support populations of some western species of Lepidoptera in northwestern Ontario. Here one might be surprised to find species that, east of Lake Superior, occur only much farther south.

The large forested areas of southern Ontario (e.g., Algonquin Park and Muskoka) are ideal breeding grounds for notodontids, noctuids, and geometrids. The area south of the Muskoka District, which 50 to 70 years ago yielded many species, is now largely under cultivation and supports fewer and fewer species.

MATERIALS AND METHODS

Compared to the butterflies, Ontario moths suffer a more serious lack of material, observations, and thorough collecting. Moreover, there is no previous list to build on, except the one found in the chapter dealing with insects and their allies by E. M. Walker (1913) in "The natural history of the Toronto region." Most of the earlier material in collections was acquired casually, although planned collection in Ontario can be traced for more than 100 years. The majority of available material has been collected over the last three or four decades. Only through the continued use of superactinic fluorescent ("blacklight") and mercury vapour lamps will we be able to obtain a more complete picture of the moths of the province.

The material used for compilation of the present list is accessible in the following collections: American Museum of Natural History, New York, New York 10024 (AMNH); Canadian National Collection, Biosystematics Research Centre, Ottawa, Ontario K1A 0C6 (CNC); Carnegie Museum of Natural History, Pittsburgh, Pennsylvania 15213 (CM); Field Museum of Natural History, Chicago, Illinois 60605 (FM); Lyman Entomological Museum, Macdonald College, Ste-Anne-de-Bellevue, Quebec H9X 3M1 (LEM); McMaster University, Hamilton, Ontario L8S 4L8 (McM); Queen's University, Kingston, Ontario K7L 3N6 (QU); Royal Ontario Museum, Toronto, Ontario M5S 2C6 (ROM); University of Guelph, Guelph, Ontario N1G 2W1 (UG); The University of Western Ontario, London, Ontario N6A 3K7 (UWO). Also used were Bethune's notebook (Beth) in the archives of the ROM, the seasonal summaries published by the Toronto Entomologists' Association for 1969–1988, and the following regional accounts: Freeman (1969), Judd (1963, 1970), Lafontaine (1972), Lamb (1967), Layberry, Lafontaine, and Hall (1980), and Worthington (1983). Distribution records taken from these last publications are indicated by an asterisk (*) in the checklist.

This study is based on material that can be traced to the following main sources:

- a) the historical collections of C. J. S. Bethune, W. Brodie, J. D. Evans, J. Fletcher, J. Macoun, J. A. Morden, H. S. Saunders, C. J. Tyers, and F. W. Warwick. Many specimens from these early collectors are preserved in the CNC and ROM. For further information about the historic development of entomology in Ontario, see Glen (1956).
- b) material collected in the first decades of this century and later by enthusiastic private collectors, especially H. V. Andrews (Toronto), H. Dawson (Hymers), P. Hahn (Toronto), H. S. Parish (Toronto), G. E. Scott (Severn Bridge), R. S. Scott (Toronto), I. M. Smith (Port Credit), and W. Plath (Hamilton, whose valuable geometrid collection is now in the ROM). Their material has found its way into the ROM and other institu-

tional collections either directly or subsequently by donation or purchase (e.g., the Buchholz collection in AMNH or B. P. Clark's sphingid collection in CM).

- c) material collected for institutions, especially by the following collectors: E. M. Walker (especially De Grassi Point on Lake Simcoe); F. P. Ide (especially Ottawa and vicinity); N. K. Bigelow in the 1920s (Macdiarmid and Low Bush); and R. V. Whelan in the 1930s (important collections in Smoky Falls). Members of ROM summer expeditions to selected northern Ontario localities provided some material: G. M. Neal, 1937 (Temagami Forest Reserve), and 1938 (Favourable Lake); S. C. Downing, 1939 (Fraserdale, Moosonee, and Onakawana); C. E. Hope, 1939 (Lansdowne House on Lake Attawapiskat), and 1940 (Fort Severn); F. A. Urquhart, 1942 (Big Piskwanish, Fort Albany, and Nettichi River); W. Y. Watson, 1945 and 1948 (Cape Henrietta Maria), and 1947 (Malachi, Nakina, and Sioux Lookout). I have included my own collecting: since 1952 in Geraldton, Nakina, and Nipigon (all in Thunder Bay District); since 1956 in Toronto, Sudbury, and One Sided Lake (in Rainy River District), and with the support of W. Hartley in (South) Neebing Township near Thunder Bay (material in AMNH, CNC, ROM); in 1963, 1964, and 1969–1975 in Chaffey's Locks (near Kingston); in 1965 in Rondeau Provincial Park and Algonquin Park; in 1966 in Geraldton and Chaffey's Locks; in 1967 in Kendal; in 1968 in Cawaja Beach. Finally, collectors include workers at the Algonquin Park Nature Museum for the Ontario Ministry of Natural Resources; workers at the Park Museum at Rondeau Provincial Park for the Ontario Ministry of Natural Resources; members of the Canadian Northern Insect Survey, organized by the former Division of Entomology of the Canada Department of Agriculture, D. F. Hardwick, 1949 (Moosonee); J. B. Wallis, 1952 (Ogoki Post); and A. B. Klots and F. H. Rindge of the AMNH, 1955 (Geraldton, Nakina, Nipigon, and western Ontario).
- d) material collected by members of the Biosystematics Research Centre, Canada Department of Agriculture, CNC. Here, particular mention should be made of the specimens collected by J. H. McDunnough, T. N. Freeman (especially southwestern Ontario), J. E. H. Martin (Marmora), E. G. Munroe, and employees of the entomological and other biological laboratories throughout the province.

Also deserving of special mention is the Forest Insect and Disease Survey (FIDS) established under the Forest Entomology and Pathology Branch of the former Canada Department of Forestry. Centred around the Forest Insect Laboratory in Sault Ste. Marie, Ontario, the FIDS assembles information about forest insects through the services of the Forest Biology Rangers, and publishes annual reports and distributional records (see McGugan, 1958;

Prentice, 1962, 1963, 1965; Raizenne, 1952). The present study would have been less comprehensive than it is were it not for the FIDS research, the availability of its reference collections in Ottawa and Sault Ste. Marie, and the original reports and reference cards prepared for its published distributional maps. For this study FIDS records were used primarily for data on rare and especially interesting species. Detailed information about food plants may be found in the publications mentioned above and is not repeated here.

In general, nomenclature and classification follows the "Checklist of the Lepidoptera of America North of Mexico (MONA)" (Hodges et al., 1983). Species are arranged alphabetically within genera. Subspecies are omitted except for consideration in the notes. Vernacular English and French names are added (English name/ French name) if listed in "Noms français d'insectes au Canada avec noms latins et anglais correspondants" (Benoit, 1975). Distributional information is given following the species name, by region (see Fig. 1). In some cases, an "s" or "n" is added to indicate occurrence in only the southernmost or northernmost parts of a region. Where only a single locality is known, this is given in the text. Ontario type localities are noted if known. For most families, detailed occurrences by county and/or district (with reference to the Ontario Gazetteer of 1974) are listed for each region (see Figs. 2 and 3). Remarks on specimens examined, flight period, number of generations, and food plant are appended following the distributional summary when available and relevant. In a few instances, information on crossbreeding and rearing experiments is included together with new evidence for the taxonomic status of Ontario populations. In the superfamily Pyraloidea and in the family Noctuidae, only the faunal regions are given; I feel that considerably more research and collecting would be necessary in these groups to provide a more detailed breakdown that does not create erroneous impressions of distribution. Further information on the Noctuidae is available in Rockburne and Lafontaine (1976). Additional information is also provided in the moth checklist for southeastern Ontario (Ward, Harmsen, and Hebert, 1974).

Throughout the list, the following abbreviations and terms are used: Dist. for District, Co. for County, Twp. for Township; the Latin *auctorum* to mean of various authors, *partim* to mean in part, and *teste* to indicate that one author's word is accepted for the record; the Toronto Entomologists' Association is identified as TEA.

The records from the Ontario tundra region are published here for the first time. This material is in the collection of the Royal Ontario Museum, which sent field parties to Fort Severn in 1940 and in Cape Henrietta Maria in 1948. Also remarkable are the several southern records (noted in the list) from a light trap operated by the Entomology Laboratory, Canada Department of Agriculture, in Chatham in southwestern Ontario.

LIST OF SPECIES

ORDER LEPIDOPTERA SUBORDER ZEUGLOPTERA

Superfamily Micropterigoidea Family Micropterigidae

Genus *Epimartyria* Walsingham

E. auricrinella Walsingham. 2: Cochrane. 4: Carleton; Simcoe. 4 May–18 July. CNC. Known only from a few localities.

SUBORDER EXOPORIA

Superfamily Hepialoidea Family Hepialidae

Genus *Sthenopsis* Packard

S. argenteomaculatus (Harris). 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Halton; Kent; Leeds; Nipissing; Simcoe; Waterloo. 21 June–29 July. AMNH, Beth, CNC, McM, ROM, UG. Common.

S. auratus Grote. 4: Carleton; Leeds; Muskoka; Peterborough. Mid July. CNC, ROM. Known from very few specimens.

S. purpurascens (Packard). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Essex; Leeds; Muskoka. 19 June–16 August. AMNH, CM, CNC, LEM, QU, ROM. Common.

S. quadriguttatus (Grote). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Hastings; Muskoka; Nipissing; York. 20 June–19 August. AMNH, Beth, CM, CNC, McM, ROM, UG. Common.

S. thule Strecker. 4: Carleton; Grenville; Hastings; Lanark; Leeds. 20 June–23 July. CNC, ROM, UG. Known only from a few specimens.

Genus *Korscheltellus* Börner [= *Hepialus* Fabricius, part.]

K. gracilis (Grote). 2: Kenora, Patricia Portion. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Carleton. 3 July–4 August. AMNH, CNC, ROM. Uncommon. The synonym *K. mustelinus* (Packard) from 4: Muskoka. 26 July–13 September. CM, ROM. Known only from a few specimens. See Wagner (1988).

Genus *Gazoryctra* Hübner [= *Hepialus* Fabricius, part.]

***G. noviganna* (Barnes and Benjamin).** 3: Thunder Bay. 11 August–9 September. Larvae as late as early November. Beth, CNC, LEM, ROM. Significant numbers of specimens from only two localities (Black Sturgeon Lake and Hymers) in the three collections; ROM specimens identified by N. B. Tindale, an Australian hepialid specialist; species referred to by Forbes (1923) as “*hyperboreus* Moeschler.” See Wagner and Tindale (1988).

SUBORDER MONOTRYSIA

Superfamily Nepticuloidea

Family Nepticulidae

Genus *Ectoedemia* Busck

***E. populella* Busck.** 3: Rainy River. 4: Carleton; Norfolk. CNC.

Superfamily Incurvarioidea

Family Incurvariidae

Subfamily Incurvariinae

Genus *Lampronia* Stephens [= *Incurvaria* Haworth]

***L. taylorella* (Kearfott).** 4: Muskoka (Gravenhurst, 23 June 1904). ROM.

Genus *Paraclemensia* Busck

***P. acerifoliella* (Fitch).** 4: mainly southeastern Ontario as far west as Sault Ste. Marie.

Subfamily Adelinae

Genus *Adela* Latreille

***A. purpurea* Walker.** 3: Thunder Bay. 4: Carleton; Muskoka (Severn Bridge, 27 April); Peel (Innis Lake, 5 May 1940). CNC, ROM.

***A. ridingsella* Clemens.** 4: Carleton; Simcoe. CNC.

SUBORDER DITRYSIA

Superfamily Tineoidea (in part)

Family Psychidae

Subfamily Taleporiinae

Genus *Solenobia* Dupont

S. walshella Clemens. 4: Carleton; Leeds; Ontario; Simcoe. 15 April–23 May. CNC, ROM. Known only from a few localities.

Subfamily Psychinae

Genus *Psyche* Schrank

P. casta (Pallas). 4: Peterborough (Owenbrook, 9–17 June 1955). CNC. An introduced European species, more common in Massachusetts and Pennsylvania; not recorded elsewhere in Canada.

Genus *Astala* Davis [= *Eurycyttarus* Hampson]

A. confederata (Grote and Robinson). 4: Essex; Halton; Victoria. July. CNC, ROM. Known only from a few localities.

Superfamily Gelechioidea (not included in this list)

Superfamily Copromorphoidea (in part)

Family Alucitidae

Genus *Alucita* Linné

A. hexadactyla Linné [= *huebneri* (Wallengren)]. 2. 4. CNC, ROM.

Superfamily Yponomeutoidea (in part)

Family Yponomeutidae

Genus *Atteva* Walker

A. punctella (Cramer) [= *aurea* Fitch]. 4: Essex*; Haldimand; Kent; Leeds; Simcoe. CNC, ROM.

Genus *Yponomeuta* Latreille

Y. atomocellus Dyar. 4s: Essex (Point Pelee). CNC.

Y. multipunctellus Clemens. 1: Essex. 4: Haldimand; Hastings; Lincoln; Middlesex; Simcoe; York. CNC, ROM.

Y. padellus (Linné). 4n: Carleton (Ottawa). CNC. Introduced.

Superfamily Sesiioidea

Family Sesiidae

See Engelhardt (1946).

Subfamily Tinthiinae

Tribe Pennisetiini

Genus *Pennisetia* Dehne

P. marginata (Harris). 3: Thunder Bay. 4: Carleton; Lincoln; Nipissing; Simcoe; Wellington. ROM.

Subfamily Paranthreninae

Tribe Paranthrenini

Genus *Paranthrene* Hübner

P. asilipennis (Boisduval). 4s: Wentworth (Hamilton, 30 May 1916). ROM.

Genus *Vitacea* Engelhardt

V. polistiformis (Harris). 3: Cochrane. 4: York. ROM.

Genus *Albuna* Hy. Edwards

A. fraxini (Hy. Edwards). 4s: Lincoln (Niagara Falls). CNC.

A. pyramidalis (Walker). 2: Cochrane; Kenora; Thunder Bay. 3: Cochrane; Sudbury; Thunder Bay. 4: Middlesex; Parry Sound; York. CNC, ROM.

Subfamily Sesiinae

Tribe Melittiini

Genus *Melittia* Hübner

M. cucurbitae (Harris). 4: York*.

Tribe Sesiini

Genus *Sesia* Fabricius

S. tibialis (Harris). 3: Thunder Bay*.

Tribe Synanthedonini

Genus *Synanthedon* Hübner

- S. albicornis* (Hy. Edwards). 4: Carleton; Essex; Leeds; Middlesex; Norfolk; Oxford; Prince Edward. CNC, ROM.
- S. acerni* (Clemens). 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Haldimand; Hastings; Leeds; Middlesex; Muskoka; Norfolk; Peterborough*; Simcoe; Welland; York. CNC, ROM.
- S. acerrubri* Engelhardt. 4: Simcoe (Orillia). CNC.
- S. bolteri* (Hy. Edwards). 3: Thunder Bay*.
- S. exitiosa* (Say). 3: Thunder Bay*. 4: Carleton; Lincoln; Middlesex; Wellington; Wentworth; York. CNC, ROM.
- S. fulvipes* (Harris). 3: Rainy River; Thunder Bay*. 4: Carleton; Lincoln. CNC.
- S. novaroensis* (Hy. Edwards). 2: Cochrane (Remi Lake). 3: Thunder Bay*. CNC.
- S. pictipes* (Grote and Robinson). 3: Thunder Bay*. 4: Carleton; Lincoln; Norfolk; Wentworth; York. CNC, ROM.
- S. pini* (Kellicott). 4: Carleton; Hastings; Renfrew. CNC.
- S. proxima* (Hy. Edwards). 3: Thunder Bay*. 4s: Middlesex; Oxford. CNC.
- S. pyri* (Harris). 4s: York. ROM.
- S. saxifragae* (Hy. Edwards). 3: Rainy River (One Sided Lake). CNC.
- S. scitula* (Harris). 4: Grenville; Lincoln; Norfolk; Peel; York. CNC, ROM.
- S. tipuliformis* (Clerck). 4: Hastings; Kent; Middlesex; Norfolk; Peel; York. CNC, ROM.
- S. viburni* Engelhardt. 4s: Lincoln; Wellington. CNC.

Genus *Podosesia* Möschler

P. syringae (Harris). 3: Thunder Bay*. 4n: Carleton. CNC.

Genus *Carmenta* Hy. Edwards

- C. bassiformis* (Walker) [= *aureopurpura* (Hy. Edwards)]. 4: Victoria. CNC.
- C. corni* (Hy. Edwards). 3: Thunder Bay*. 4n: Carleton. CNC.

C. pyralidiformis (Walker). 4: Hastings; Lincoln; Middlesex; Norfolk. CNC, ROM.

Genus *Penstemonia* Engelhardt

P. dammersi Engelhardt. 4: Lincoln; Norfolk. CNC.

P. edwardsii (Beutenmüller). 4: Lincoln; Simcoe. CNC.

Genus *Alcathoe* Hy. Edwards

A. caudata (Harris). 4: Hastings; Peel; York. CNC, ROM.

**Family Choreutidae
Subfamily Choreutinae**

Genus *Anthophila* Haworth

A. alpinella (Busck). 4: Carleton. CNC.

Genus *Prochoreutis* Diakonoff and Heppner

P. inflatella (Clemens). 4: Carleton. CNC.

Genus *Caloreas* Heppner

C. leucobasis (Dyar). 4: Carleton. CNC.

Genus *Tebenna* Billberg

T. onustana (Walker). 4: Carleton. CNC.

Genus *Choreutis* Hübner

C. diana (Hübner). 3: Thunder Bay. 4: Carleton. CNC.

C. pariana (Clemens). 4: Carleton; Frontenac; Simcoe; Wentworth. CNC.

**Superfamily Cossoidea
Family Cossidae
Goat-moths, carpenterworms.**

Subfamily Cossinae

Genus *Acossus* Dyar

- A. centerensis* (Lintner). Poplar carpenterworm/ver charpentier du peuplier. 3: Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Leeds; Middlesex; Muskoka; Nipissing; Simcoe; York. 10 June–26 July. AMNH, CNC, LEM, ROM.
- A. populi* (Walker). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 27 June–12 August. AMNH, CNC, LEM, ROM. Evidently more northern than the previous species.

Genus *Prionoxystus* Grote

- P. macmurtrei* (Guérin-Ménéville). Little carpenterworm/petit ver charpentier. 2: Thunder Bay. 3: Thunder Bay. 4: Durham; Frontenac; Huron; Kent; Leeds; York. 29 May–30 June. CNC, ROM. Known only from a few localities in very different faunal regions; the males are extremely rare in collections.
- P. robiniae* (Peck). 3: Algoma; Rainy River; Thunder Bay. 4: Carleton; Durham; Essex; Hastings; Kent; Leeds; York. 2 June–18 July. AMNH, CNC, ROM.

Superfamily Tortricoidea (not included in this list)

Superfamily Hesperioidea

Family Hesperiidae

Subfamily Pyrginae

Genus *Epargyreus* Hübner

- E. clarus* (Cramer). 3: Rainy River. 4: Brant*; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Halton; Huron; Kent; Lambton; Leeds; Lincoln; Manitoulin*; Middlesex; Muskoka; Norfolk; Northumberland; Peel; Peterborough; Prescott*; Renfrew; Russell; Simcoe; Waterloo*; Wentworth; York. End May–end September. AMNH, Beth, CNC, ROM, UG, UWO.

Genus *Achalarus* Scudder

- A. lyciades* (Geyer). 4: Essex (Sandwich West Twp.*, 9 June and 9 July 1988, Jeff Larson). Only two individuals known. Evidence suggests it may be a resident. See TEA Occasional Publications 1989, 20:11.

Genus *Thorybes* Scudder

- T. bathyllus* (J. E. Smith). 4: Elgin; Essex; Haldimand; Lambton; Middlesex; Norfolk; Wentworth; York. Early June–mid July. CNC, ROM, UG.

***T. pylades* (Scudder).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming*. 4: Carleton; Bruce; Dundas*; Durham; Elgin; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Kent; Lambton*; Lanark; Leeds; Lincoln; Manitoulin*; Muskoka; Nipissing; Norfolk; Ontario; Parry Sound; Peel; Peterborough; Renfrew*; Russell*; Simcoe; Stormont; Victoria; Wellington*; Wentworth; York. End May–end July. AMNH, Beth, CNC, ROM, UG, UWO.

Genus *Staphylus* Godman and Salvin

***S. hayhurstii* (Edwards).** 4: Essex (Pelee Island. Point Pelee). June–mid September. ROM. Forbes (1960:69) said, “Evans considered our species a race of the Mexican *S. mazans* (Reakirt) but the genitalia are strikingly different.” Rare straggler from south.

Genus *Erynnis* Schrank

***E. baptisiae* (Forbes).** 4: Essex (Leamington. Windsor, Ojibway Park); Lambton (Walpole Island*); Norfolk (St. Williams). June–end September. CNC. Known only from southwestern Ontario.

***E. brizo* (Boisduval and Leconte).** 4: Essex*; Grey*; Halton; Huron*; Lambton; Manitoulin*; Middlesex; Nipissing; Norfolk; Northumberland; Parry Sound; Peterborough; Victoria; Wentworth. End April–end June. ROM. Specimens cited by Scudder (see Burns, 1964) from Ottawa could not be located.

***E. horatius* (Scudder and Burgess).** 4: Essex (Point Pelee*); Lambton (Pinery Provincial Park); Norfolk. Early May–mid June and August. Known only from a few specimens in private collections until recently.

***E. icelus* (Scudder and Burgess).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Dufferin; Dundas; Durham; Elgin*; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton*; Hastings; Huron; Lambton; Lanark*; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Prescott; Renfrew; Russell; Simcoe; Victoria; Welland; Wellington; Wentworth; York. Early May–early July. AMNH, Beth, CNC, ROM, UG, UWO.

***E. juvenalis* (Fabricius).** 3: Algoma; Kenora; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peterborough; Renfrew; Russell; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Early May–early July. CNC, ROM, UG.

- E. lucilius* (Scudder and Burgess). 3: Algoma; 4: Bruce; Carleton; Durham; Elgin*; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Norfolk; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Russell; Victoria; Wellington; Wentworth; York. Early May–end August. CNC, ROM, UG, UWO.
- E. martialis* (Scudder). 4: Carleton; Essex*; Haldimand; Hastings; Kent; Lambton; Lincoln; Middlesex*; Norfolk; Victoria; Wentworth; York. Mid May–early August. CNC, ROM, UG.
- E. persius* (Scudder). 4: Lambton (Walpole Island*. Pinery Provincial Park*); Muskoka (Bracebridge*); Norfolk (Backus Woods*. St. Williams*). Early May–early June. See TEA Occasional Publications 1981, 12:39.
- E. zarucco* (Lucas). 4: Ontario (Rouge River, 20 June 1935, T. Irwin). CNC. Known only from one specimen that was out of range even as a straggler.

Genus *Pyrgus* Hübner

- P. centaureae* (Rambur). 1: Kenora, Patricia Portion (Winisk*). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Kenora; Sudbury. Early June–mid July. AMNH, Beth, CNC, ROM. The record “Favourable Lake” (Riotte, 1959) should be corrected to “Lansdowne House.” Not reported often.
- P. communis* (Grote). 4: Essex; Lincoln (St. Catharines); York (Toronto, Cedarvale Ravine, Bathurst Street at St. Clair, 1982, TEA Summary 1982). End August–end September. ROM. Rarely reported and not known to overwinter in Ontario.

Genus *Pholisora* Scudder

- P. catullus* (Fabricius). 4: Carleton; Elgin; Essex; Haldimand; Halton; Huron; Kent; Lambton; Lincoln; Middlesex; Nipissing; Muskoka; Norfolk; Peel; Peterborough; Wentworth; York. End May–end August. CNC, ROM, UG, UWO. Widely distributed but not common.

Subfamily Heteropterinae

Genus *Carterocephalus* Lederer

- C. palaemon* (Pallas). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Carleton; Dufferin; Dundas; Grenville; Haliburton; Halton; Hastings; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Ontario; Parry Sound; Peterborough; Prescott; Renfrew*;

Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth*; York. End May–early July. AMNH, Beth, CNC, ROM, UG, UWO. Subspecies *C. p. mandan* (Edwards).

Subfamily Hesperiiinae

Genus *Ancyloxypha* Felder

A. numitor (Fabricius). 2: Cochrane*; Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grenville; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Oxford; Peel; Peterborough; Perth; Prescott*; Prince Edward; Renfrew*; Russell; Stormont; Victoria; Wellington; Wentworth; York. End May–end July; in the south to early October. AMNH, CNC, ROM, UG, UWO.

Genus *Oarisma* Scudder

O. garita (Reakirt). 4n: Manitoulin (Great la Cloche Island only). 26 June–8 July. First Ontario record: P. M. Catling, 1976. ROM. See TEA Annual Summaries.

Genus *Thymelicus* Hübner

T. lineola (Ochsenheimer). 2: Cochrane (Hearst). 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grenville; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Prescott; Renfrew; Russell; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Mid June–early September. AMNH, CNC, ROM, UG, UWO. Introduced from Europe; first recorded in Ontario in Middlesex (Hyde Park Corner) in 1910. Subsequently spread south (Michigan, Ohio, Kentucky, Pennsylvania, New Jersey) and to northcentral Ontario (Sudbury, Sault Ste. Marie, and north), eastern Ontario (Ottawa, Brockville), and western Ontario to Thunder Bay (Fort William), and very recently to northern Ontario proper to Cochrane (Hearst) as well as to the western provinces of Canada. See Pengelly (1960, 1965), Arthur (1962, 1966), Riotte (1962a), and Burns (1966) for additional information. The pale form has been reported in Ontario but is rare.

Genus *Hylephila* Billberg

H. phyleus (Drury). 4: Elgin*; Essex; Haldimand; Halton; Kent (Rondeau Provincial Park); Lambton; Peel; Wentworth; York. End July–mid October. ROM, UWO. May overwinter in Ontario but only second brood reported so far. See Wormington (1983).

Genus *Hesperia* Fabricius

H. comma (Linné). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Sudbury; Thunder Bay; Timiskaming. 4: Grey; Manitoulin; Muskoka; Nipissing; Parry Sound; Renfrew. Early July–early September. AMNH, Beth, CNC, ROM, UG.

H. leonardus Harris. 3: Algoma; Sudbury. 4: Bruce; Carleton; Durham; Elgin*; Essex; Frontenac; Haldimand; Haliburton; Halton; Huron; Kent*; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prince Edward; Renfrew*; Russell; Wellington*; Wentworth; York. End July–end September. CNC, ROM, UG, UWO.

H. sassacus Harris. 2: Kenora. 3: Rainy River; Thunder Bay. 4: Carleton; Frontenac; Haldimand; Haliburton; Halton; Hastings; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Wentworth; York. End May–end July; in the north, end June–July. CNC, LEM, ROM. Subspecies *H. s. manitoboides* (Fletcher) in northern populations; Thunder Bay (Nipigon, type locality).

Genus *Polites* Scudder

P. coras (Cramer) [= *peckius* (Kirby)]. 2: Cochrane; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. Early June–early September. AMNH, Beth, CNC, ROM, UG, UWO.

P. mystic (Edwards). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury*; Thunder Bay; Timiskaming. 4: Brant; Carleton; Dufferin; Dundas; Durham; Elgin; Essex; Grenville; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria;

Waterloo; Wellington; Wentworth; York. Early June–early August.
AMNH, Beth, CNC, LEM, ROM, UG, UWO.

P. origenes (Fabricius) [= *manataaqua* (Scudder)]. 4: Brant; Carleton; Dufferin; Dundas; Durham; Elgin*; Essex; Frontenac; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Renfrew; Simcoe; Victoria; Waterloo; Welland*; Wellington; Wentworth; York. End June–early August. CNC, ROM, UG. Local.

P. themistocles (Latreille). 2: Kenora; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dufferin; Dundas; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Oxford; Parry Sound; Peel; Peterborough; Prince Edward; Renfrew; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. End May–end August. AMNH, Beth, CNC, ROM, UG, UWO.

P. vibex (Geyer). 4: York (Toronto, no date, Arthur Gibson). CNC. Known only from one specimen.

Genus *Wallengrenia* Berg

W. egeremet (Scudder). 3: Cochrane. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Haldimand; Haliburton; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Victoria; Wellington; Wentworth; York. Mid June–mid August. CNC, ROM, UG, UWO.

Genus *Pompeius* Evans

P. verna (Edwards). 4: Brant*; Elgin*; Essex; Haldimand; Halton; Huron; Kent; Lambton; Lincoln; Middlesex; Norfolk; Northumberland; Peel; Wellington; Wentworth; York. End June–end July. CNC, ROM, UG. Rare and local.

Genus *Atalopedes* Scudder

A. campestris (Boisduval). 4: Essex*; Middlesex (London, 21 September 1968, Ken Thorne). End June–end September. Known only from the one London female until 1988 when over 50 were recorded in Essex Co., many of which were from a subsequent local brood produced by the original migrants.

Genus *Atrytone* Scudder

***A. delaware* (Edwards).** 4: Brant; Essex; Halton; Huron; Kent; Lambton; Norfolk; Peel; Simcoe*; Waterloo; Wellington; Wentworth; York. End June–end July. CNC, ROM. Known only from a few localities; uncommon.

Genus *Poanes* Scudder

***P. hobomok* (Harris).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grenville; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. Mid May–mid July; in the north, early July. AMNH, Beth, CNC, ROM, UG, UWO. The female form *pocahontas* (Scudder) has been recorded in 3: Sudbury; Timiskaming*. 4: Carleton; Durham; Elgin; Haldimand; Leeds; Middlesex; Muskoka; Norfolk*; Parry Sound; Peel; Peterborough*; Simcoe; Victoria; Wentworth; York. CNC, Peabody Museum, ROM, UG.

***P. massasoit* (Scudder).** 4: Brant; Dundas; Elgin; Essex; Grey; Glengarry; Halton; Hastings; Huron*; Kent; Lambton; Lennox and Addington; Middlesex; Nipissing*; Russell*; Stormont; Waterloo; Wentworth. Early July–early August. ROM. Very local.

***P. viator* (Edwards).** 4: Brant; Carleton; Dundas; Elgin; Essex; Frontenac; Grenville; Haldimand; Halton; Huron; Kent; Lambton; Lanark; Leeds; Middlesex; Norfolk; Northumberland*; Renfrew; Russell; Stormont; Waterloo; Wellington; Wentworth; York. End June–early August. CNC, ROM, UG.

Genus *Euphyes* Scudder

***E. bimacula* (Grote and Robinson).** 3: Sudbury. 4: Brant; Carleton; Essex; Kent; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk*; Peterborough; Russell; Simcoe; Waterloo*; Wellington; Wentworth; York. Mid June–end July. AMNH, ROM, UWO. Known from isolated colonies.

***E. conspicua* (Edwards).** 4: Brant; Elgin*; Essex; Haldimand; Kent; Lambton; Lincoln; Norfolk; Middlesex; Waterloo; Wellington*; Wentworth (Dundas Valley). Early July–early August. ROM. Known only from the region immediately north of Lake Erie.

- E. dion* (Edwards). 4: Brant; Carleton; Dundas; Elgin; Essex; Frontenac; Glengarry; Haldimand; Huron; Kent; Lambton; Leeds; Middlesex; Norfolk; Northumberland; Peterborough; Russell; Waterloo; Welland; Wellington; Wentworth; York. End June–mid August. CNC, ROM, UG, UWO
- E. dukesi* (Lindsey). 4: Essex (McGregor, first Ontario specimen 11 July 1968; 9 July 1974, and 14 July 1975, A. M. Holmes, some specimens ROM. Point Pelee National Park. Other Essex localities); Kent (a few western localities); Leeds (Chaffey's Locks, 7 July 1975, summer students of Queen's University Biological Station on Lake Opinicon, swampy area north of old railway station, specimen lost). See also Riotte (1972b). End June–end August. Local.
- E. ruricola* (Boisduval). 2: Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grenville; Grey; Haldimand; Haliburton; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. End June–mid August. AMNH, Beth, CNC, ROM, UG, UWO. Subspecies *E. r. metacomet* (Harris) and *E. r. vestris* Boisduval.

Genus *Atrytonopsis* Godman

- A. hianna* (Scudder). 4: Lambton (Ipperwash. Pinery Provincial Park). End May–end June. ROM, UWO. L. Packer has proposed rare status for this species in Ontario (see TEA Occasional Publications 1988, 19:6).

Genus *Amblyscirtes* Scudder

- A. hegon* (Scudder) [= *samoset* (Scudder)]. 3: Sudbury; Timiskaming. 4: Carleton; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough*; Timiskaming*. End May–mid June. CNC, ROM, UG. Local and rare.
- A. vialis* (Edwards). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Dundas; Elgin; Frontenac; Haliburton; Hastings; Huron; Lambton; Lanark; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Parry Sound; Peterborough; Simcoe; Victoria; Wentworth; York. End May–early July. AMNH, Beth, CNC, ROM, UG.

Superfamily Papilionoidea

See Klots (1951), Ehrlich and Ehrlich (1961), and Opler and Krizek (1984).

Family Papilionidae
Subfamily Papilioninae
Tribe Troidini

Genus *Battus* Scopoli

B. philenor (Linné). Pipe vine swallowtail. 3: Thunder Bay (Caribou Island, Lake Superior, 12 September 1979, A. Wormington, a straggler from the south). 4: Elgin; Essex; Haldimand; Halton*; Huron; Kent; Northumberland; Ontario; Waterloo*; Wentworth; York. End May–early October. CNC, ROM. As its vernacular name suggests, breeding restricted to a small area on the north shore of Lake Erie and near Hamilton and Toronto where the foodplant is grown around houses, once and if migrants arrive. Not known to overwinter in Ontario, except on Pelee Island where the foodplant grows wild. Usually fairly rare.

Tribe Papilionini

Genus *Papilio* Linné

P. cresphontes Cramer. 4: Elgin; Essex; Haldimand*; Kent; Lambton; Lincoln; Middlesex; Norfolk*; Waterloo*; Welland; Wentworth*; York. End May–early September. CNC, ROM.

P. glaucus Linné. Tiger swallowtail/papillon glauque. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton*; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Mid April–mid October. CNC, ROM. Northern populations are smaller, light yellow, and one-brooded while southern populations are larger, deeper yellow, and have at least a partial second generation. Black form females occur in extreme southern Ontario—e.g., Pelee Island and Point Pelee—and are rare. Subspecies *P. g. canadensis* Rothschild and Jordan in the north; considered a separate species by some biologists.

P. "machaon Linné." 2: Cochrane (Nettichi River); Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Thunder Bay (Hymers [see Forbes, 1960]. City of Thunder Bay*, 1988. South Neebing Twp., 21 June 1963, W. Hartley). Early June–early July. CNC, ROM. Subspecies *P. m. hudsonianus* A. H. Clark. The question of whether nearctic "*machaon*"

is conspecific with the palaearctic nominate *machaon* is yet to be answered.

***P. polyxenes* Fabricius.** Black or parsnip swallowtail/papillon du céleri. 3: Algoma; Rainy River (Lake of the Woods shore); Sudbury; Thunder Bay (Nipigon). 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Glengarry; Grenville; Grey; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Oxford; Peel; Perth; Peterborough; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Mid April–end September. CNC, ROM. Subspecies *P. p. asterius* Stoll.

***P. troilus* Linné.** Spicebush swallowtail. 4: Elgin; Essex; Haldimand*; Huron; Kent; Lambton; Middlesex; Norfolk; Waterloo*; Welland; Wentworth*; York. Mid April–end September. CNC, ROM.

Genus *Eurytides* Hübner [= *Graphium* auct.]

***E. marcellus* (Cramer).** 4: Brant (reported flying in the Brantford area, 1978); Elgin*; Essex; Norfolk*; Northumberland*; Waterloo*; York. Mid June–early September. CNC, ROM. Recorded in Toronto as late as 1943 in the Humber Valley. May be seen at Point Pelee and other southerly locations. Not known to overwinter in Ontario. May breed here but the foodplant, pawpaw, is restricted in the province. Usually very rare. See Wormington (1983).

Family Pieridae Subfamily Pierinae Tribe Pierini

Genus *Pontia* Fabricius

***P. occidentalis* (Reakirt).** Checkered white cabbageworm/piéride occidentale du chou. 2: Kenora, Patricia Portion (Lansdowne House). 3: Rainy River (Atikokan); Thunder Bay. End June–September. ROM. Identity of specimens confirmed by dissection.

***P. protodice* (Boisduval and Leconte).** 2: Kenora (Minaki. Waldhof). 3: Rainy River; Thunder Bay. 4: Carleton; Elgin*; Essex*; Haldimand*; Halton*; Manitoulin; Norfolk; Northumberland; Peel; Simcoe; Waterloo*; York. End June–early October. CNC, ROM. Identity of specimens confirmed by dissection. Not common in northwestern Ontario; usually rare in the south.

Genus *Pieris* Schrank

This genus is kept intact following Ehrlich (1958), Forster and Wohlfahrt (1976), and Geiger (1980). Geiger (1980) used electrophoresis to show decisively that there is no support for the proposal to split the old genus *Pieris*.

***P. oleracea* Harris** [= *napi oleracea* Harris]. 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Frontenac; Glengarry; Grenville; Grey; Haliburton; Halton; Hastings; Huron*; Lambton; Lanark; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington*; Wentworth*; York. Mid April–mid September. CNC, ROM. Following Warren (1961, 1963) and Geiger (1986), *oleracea* was treated as a species distinct from the palaearctic *napi* (Linné). See also Eitschberger and Steiniger (1983).

***P. rapae* (Linné)**. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grey; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Early April–early November. CNC, ROM.

***P. virginiensis* Edwards**. 3: Algoma. 4: Brant*(?); Elgin; Frontenac; Halton; Leeds; Manitoulin; Middlesex (north Mosa Twp.); Peel; Waterloo*; Wellington*; [Wentworth (Hamilton); York (Etobicoke)]. Mid April–end May. CNC, ROM, UG, UWO. The counties in square brackets contain localities where the species is by now extinct. This was Ontario's only officially endangered butterfly; in 1990 it was removed from the list of endangered species. For detailed studies, see TEA Occasional Publications; see also Riotte (1967b).

Genus *Ascia* Scopoli

***A. monuste* (Linné)**. 4: Essex (Point Pelee, 16 June 1981, Park Naturalist A. Wormington, definitive sight record). See TEA Occasional Publications 1982, 13:35; Wormington, 1983. Only Ontario record.

Subfamily Anthocharinae

Genus *Euchloe* Hübner

- E. ausonides* (Lucas). 2: Kenora; Thunder Bay. 3: Rainy River; Thunder Bay. 4: Manitoulin. Early May–end June. CNC, ROM. See Riotte (1967a, 1968).
- E. olympia* (Edwards). 4: Carleton; Frontenac; Haliburton; Hastings; Lambton; Lanark; Leeds; Manitoulin; Muskoka; Northumberland; Peterborough; Parry Sound; Renfrew; Victoria. Mid April–mid June. CNC, ROM.

Subfamily Coliadinae

Genus *Colias* Fabricius

- C. eurytheme* Boisduval. Alfalfa caterpillar/coliede de la luzerne. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grenville*; Haldimand; Halton*; Hastings; Huron; Kent; Lambton; Lanark; Leeds*; Lennox and Addington; Lincoln; Manitoulin; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Stormont; Waterloo; Welland; Wellington; Wentworth; York. Early May–early November. CNC, ROM. The most common *Colias* in Ontario.
- C. gigantea* Strecker. 1: Kenora, Patricia Portion (Cape Henrietta Maria. Fort Severn. Winisk). 2: Cochrane (Fort Albany). End June–early August. ROM. Only found in the extreme north of Ontario.
- C. interior* Scudder. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Haliburton; Hastings*; Lanark; Manitoulin; Muskoka; Nipissing; Parry Sound; Peterborough*; Russell; Victoria; Waterloo* [?]; York. Mid June–mid August. CNC, ROM. Recently has been spreading to the south.
- C. palaeno* (Linné). 1: Kenora, Patricia Portion (Fort Severn. Winisk). 2: Kenora, Patricia Portion (Lansdowne House). July. CNC, ROM. Subspecies *C. p. chippewa* Edwards.
- C. pelidne* Boisduval and Leconte. 1: Kenora, Patricia Portion (Cape Henrietta Maria. Winisk). 2: Cochrane (Fort Albany. Moose Factory). End June–end July. CNC, ROM.
- C. philodice* Godart. Clouded sulphur/piéride du trèfle. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin;

Essex; Frontenac; Glengarry; Grenville; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. End April–early November. CNC, ROM.

Genus *Zerene* Hübner

Z. cesonia (Stoll). Dog face. 4: Essex (Point Pelee*, 24 June 1931); York (Toronto). ROM. Recorded at the Humber River, 11–15 June 1896 (see *Canadian Entomologist*, 28, July 1896). Two specimens recorded in Toronto, 6 June 1896, and a single specimen recorded in Toronto, 13 July 1906; another reportedly seen in a Toronto city park in 1973. Thought to occur in extreme southwestern Ontario near Leamington (in Essex Co.), but no specimens in collections are known. Rare straggler from the south. See also Wormington (1983).

Genus *Phoebis* Hübner

P. philea (Johannsson). 4: Essex (Windsor, 1896); Huron (1930); Lincoln (Vineland, 1930, 1944). 13 July–13 October. ROM. Very sporadic straggler from the south.

P. sennae (Linné). 4: Essex (Point Pelee. Windsor*, 1988); Kent (Rondeau Provincial Park); Peel (Port Credit*, 2 September 1985, from Hurricane Elena?); York (Toronto*). End May–early October. ROM. Subspecies *P. s. eubule* (Linné).

Genus *Eurema* Hübner

E. lisa Boisduval and Leconte. 3: Sudbury. 4: Carleton; Durham; Essex; Elgin*; Frontenac; Haldimand; Kent; Leeds*; Lincoln*; Manitoulin; Ontario*; Simcoe; Waterloo; Welland; York. Early June–end September. CNC, ROM. See TEA Occasional Publications 1934, 19:3; Wormington (1983). Almost yearly migrant from the south, which then produces as many as three broods. Varies in number from year to year. Usually not common. Not known to overwinter in Ontario.

E. mexicanum (Boisduval). 4: Essex (Point Pelee). CNC. Known in Ontario from only one specimen, 30 June 1882, W. E. Saunders. See Wormington (1983).

E. nicippe (Cramer). 3: Rainy River (Quetico Provincial Park*, 21 June 1978, photographed by Marie Nelson, a worn female on flower of *Corydalis sempervirens*). See TEA Occasional Publications 1979, 10:16, and 1980, 11. A suspect specimen with a label from 4: Carleton (Ottawa,

about 1900, W. E. Saunders), is probably just a Saunders collection label and unrelated to where the specimen was caught. CNC. See Lafontaine (1972).

Genus *Nathalis* Boisduval

N. iole Boisduval. 3: Algoma. 4: Peterborough (Bridgenorth, 1947, T. H. Johnson). CNC, ROM. Only two specimens known.

Family Lycaenidae

Subfamily Liphyrinae

Tribe Gerydini [= Spalgini]

Genus *Feniseca* Grote

F. tarquinius (Fabricius). 3: Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming*. 4: Carleton; Elgin*; Frontenac; Haliburton; Hastings*; Leeds; Muskoka; Nipissing; Norfolk; Northumberland; Ontario*; Peterborough*; Russell; Simcoe; Victoria; Waterloo*; Wentworth; York. Mid May–end August. CNC, ROM. Local.

Subfamily Lycaeninae

Genus *Lycaena* Fabricius

L. dorcas (Kirby). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Sudbury; Thunder Bay. 4: Bruce; Grey; Lambton; Manitoulin; Waterloo. End June–mid August. CNC, ROM.

L. epixanthe (Boisduval and Leconte). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Kenora; Rainy River. 4: Brant; Bruce; Carleton; Frontenac; Haliburton; Leeds; Middlesex; Muskoka; Nipissing; Oxford*; Prescott; Renfrew; Russell; Simcoe; Wentworth; York. Mid June–mid August. CNC, ROM. In southern Ontario restricted to acid bogs.

L. helloides (Boisduval). 2: Cochrane (west of Hearst). 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Haldimand*; Kent (Rondeau Provincial Park); Manitoulin; Parry Sound; Waterloo; York. End July–end August. CNC, ROM. Rare and local.

L. hyllus (Cramer) [= *thoe* (Guérin-Méneville)]. 3: Cochrane*; Kenora; Rainy River; Thunder Bay. 4: Brant; Bruce; Carleton; Dufferin; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln*; Manitoulin; Middlesex;

Norfolk; Northumberland; Ontario; Oxford; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Waterloo; Wellington*; Wentworth; York. Early June–early October. CNC, ROM.

L. phlaeas (Linné). 2: Cochrane. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand*; Halton; Hastings; Haliburton; Kent; Lambton*; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Prescott*; Russell; Simcoe; Victoria; Waterloo*; Welland; Wentworth; York. Mid May–early October. CNC, ROM. Subspecies *L. p. americana* Harris.

L. xanthoides (Boisduval). 3: Kenora (just east of Hwy 17 and 71, 29 July 1979, John Prideaux, Jr.), see TEA Occasional Publications 1983, 14. Ontario is at the extreme eastern edge of its range.

Subfamily Eumaeinae Tribe Strymonini

Genus *Harkenclenus* dos Passos

H. titus (Fabricius). 2: Kenora. 3: Algoma; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand*; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington*; Lincoln; Manitoulin; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel*; Peterborough; Simcoe; Waterloo; Wellington; Wentworth; York. End June–end August. CNC, ROM.

Genus *Satyrium* Scudder

S. acadicum (Edwards). 3: Algoma*; Rainy River; Sudbury. 4: Brant*; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grey; Haldimand*; Haliburton; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington*; Lincoln; Manitoulin*; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Russell; Simcoe; Victoria; Waterloo; Welland; Wellington*; Wentworth; York. Mid June–mid August. CNC, ROM.

S. calanus (Hübner). 3: Algoma*; Sudbury. 4: Brant; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry*; Grey; Haldimand*; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Russell; Simcoe; Victoria; Waterloo*; Welland; Wellington*; Wentworth; York. End June–mid August. CNC, ROM. Subspecies *S. c. falacer* (Godart) may be a species in its own right.

- S. caryaevorum* (McDunnough). 4: Carleton; Durham; Elgin; Essex; Frontenac; Glengarry; Halton; Hastings*; Kent; Lincoln; Leeds; Middlesex; Norfolk; Northumberland*; Ontario; Simcoe; Waterloo; Wentworth; York. End June–end July. CNC, ROM.
- S. edwardsii* (Grote and Robinson). 4: Brant; Durham; Elgin; Essex; Frontenac; Grey; Haldimand*; Halton*; Lambton; Leeds; Lennox and Addington*; Lincoln; Middlesex; Norfolk; Northumberland; Parry Sound; Renfrew; Simcoe; Waterloo*; Wellington*; Wentworth*; York. End June–mid August. CNC, ROM.
- S. liparops* (Leconte). 2: Kenora; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grey; Glengarry*; Haldimand*; Haliburton; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. End June–mid August. CNC, ROM. Subspecies *S. l. fletcheri* (Michener and dos Passos) in northern Ontario.

Genus *Mitoura* Scudder

- M. grynea* (Hübner). 4: Essex (Point Pelee, Pelee Island); Frontenac; Hastings; Northumberland; Prince Edward. End May–early August. CNC, ROM. Known for many years from Point Pelee; recorded near Marysville and near Shannonville (both in Hastings Co.) in 1965, and near Perth Road (in Frontenac Co.) in 1973.

Genus *Incisalia* Scudder

- I. augustus* (Kirby). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Grey; Haliburton; Lambton; Lanark; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Parry Sound; Peterborough; Prescott; Russell; Simcoe; York. Mid April–mid June. CNC, ROM. In southern Ontario restricted to acid bogs.
- I. eryphon* (Boisduval). 2: Kenora; Thunder Bay. 3: Rainy River; Thunder Bay; Timiskaming. 4: Durham (Port Hope); Frontenac; Haliburton; Manitoulin; Muskoka; Nipissing; Parry Sound; Renfrew. Early May–mid June. CNC, ROM. For many years not recognized and may be misidentified as *niphon* in some collections.
- I. henrici* (Grote and Robinson). 4: Carleton; Frontenac; Leeds; Muskoka*; Norfolk; Northumberland; Parry Sound; Peterborough; Prescott. Mid April–mid June. CNC, ROM. Local.

- I. irus* (Godart). 4: Norfolk (St. Williams). End April–early June. CNC, ROM. L. Packer proposed endangered status for this species in Ontario and it was placed on the endangered species list in 1990. See TEA Occasional Publications 1988, 19:6.
- I. lanoraieensis* Sheppard. 4: Prescott (Alfred Bog in Caledonia Twp., 22 May 1982; 29 May 1983). See TEA Occasional Publications 1984, 15. Ontario is at the extreme western edge of its range.
- I. niphon* (Hübner). Jack-pine hairstreak/lycaenide du pin gris. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Timiskaming. 4: Carleton; Durham; Frontenac; Haliburton; Halton*; Hastings*; Lambton; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Peterborough; Renfrew; Russell; Victoria; Wellington*; Wentworth; York. Mid April–mid June. CNC, ROM. Subspecies *I. n. clarki* (Freeman) in northern Ontario.
- I. polia* Cook and Watson. 2: Cochrane; Thunder Bay (Geraldton. Lukinto Lake. Nakina). 3: Rainy River; Thunder Bay (Marathon). 4: Bruce; Carleton; Frontenac (Perth Road); Lambton (Pinery Provincial Park); Lanark; Leeds (Chaffey's Locks); Manitoulin; Muskoka; Nipissing; Northumberland; Parry Sound; Peterborough; Renfrew; Victoria (Dalrymple). Early April–mid June. AMNH, CNC, ROM. Probably occurs in other localities where the foodplant *Arctostaphylos uvaursi* (bearberry) grows.

Genus *Euristrymon* Clench

- E. ontario* (Edwards). 4: Elgin (Port Stanley, 1868, type locality). CNC. This is the only specimen known from Ontario in any scientific collection. For other possible specimens, see TEA Occasional Publications 1986, 17:20, and 1987, 18:25.

Genus *Parrhasius* Hübner

- P. m-album* (Boisduval and Leconte). 4s: Essex (Point Pelee, 31 August 1960, two specimens. City of Essex, 11 September 1960). Known from only three specimens. CNC. See TEA Occasional Publications 1983, 14; Wormington (1983).

Genus *Strymon* Hübner

- S. melinus* Hübner. 4: Elgin; Essex; Frontenac; Huron*; Kent; Leeds; Lennox and Addington*; Nipissing; Norfolk*; Northumberland; Peterborough*; Renfrew*; Simcoe; Victoria. End April–early October. CNC, ROM. Not common.

Genus *Erora* Scudder

E. laeta (Edwards). 4: Leeds (Chaffeys Locks); Middlesex (London, type locality); Muskoka; Nipissing (Algonquin Provincial Park); Parry Sound; Simcoe; York. Mid May–mid June. CNC, ROM; holotype CM. Very rarely seen in known localities; never to be expected. See TEA Occasional Publications 1983, 14.

Subfamily Polyommatainae Tribe Everini

Genus *Everes* Hübner

E. amyntula (Boisduval). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. Early June–mid August. CNC, ROM. A partial second generation occurs in warm years.

E. comyntas (Godart). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Brant; Carleton; Durham; Elgin; Essex; Frontenac; Grenville; Haldimand*; Halton; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Nipissing; Norfolk; Northumberland; Ontario; Peel; Prince Edward; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. Early May–16 October. CNC, ROM. Wormington (1983) states there are up to four generations at Point Pelee. *Everes amyntula* and *comyntas* are not so much western and eastern as northern and southern in Ontario. Sudbury may be taken to be approximately the southernmost locality for *amyntula*. H. K. Clench (personal comment) now holds that the two are quite different species. See Klots (1951:163–164).

Tribe Lycaenopsini

Genus *Celastrina* Tutt

C. ladon (Cramer) [= *pseudargiolus* (Boisduval and Leconte)]. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth*; York. Early April–mid September. CNC, ROM.

Forms *lucia* (Kirby) and *marginata* (Edwards) are equally abundant in northern localities; a few specimens of form *violacea* (Edwards) are regularly intermingled. Form *neglecta* (Edwards) is local in early and hot summers: Thunder Bay (Nakina, 18 July 1955).

Tribe Scolitantidini

Genus *Glaucopsyche* Scudder

G. lygdamus (Doubleday). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Essex* (Point Pelee, 1981); Frontenac; Grey; Haliburton; Hastings; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Northumberland; Parry Sound; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Victoria. Early May–mid July. CNC, ROM. A partial second generation may occur in extremely hot summers in late August, e.g., Thunder Bay (Geraldton, 22 August 1955). Subspecies *G. l. couperi* Grote is questionably recorded in northern Ontario. For information on *G. l. lygdamus* (Doubleday) at Point Pelee, see Wormington (1983).

Tribe Polyommattini

Genus *Lycaeides* Hübner

L. idas (Linné) [= *argyrognomon* Bergsträsser]. 2: Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Thunder Bay. Early June–early August. ROM. Subspecies *L. i. nabokovi* Masters. See Higgins (1985).

L. melissa (Edwards). 4: Lambton; [Middlesex]; Norfolk; [York]. Early May–mid August. CNC, ROM. The counties in square brackets contain localities where the species is by now extinct. L. Packer proposed endangered status for this species in Ontario and it was placed on the endangered species list in 1990. See TEA Occasional Publications 1988, 19:6. Subspecies *L. m. samuelis* Nabokov.

Genus *Plebejus* Kluk

P. saepiolus (Boisduval). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Hastings; Muskoka; Nipissing; Parry Sound. Early June–early August. CNC, ROM. Subspecies *P. s. amica* (Edwards).

Genus *Agriades* Hübner

A. franklinii (Curtis) [= *glandon* auct. = *aquilo* auct.]. 1: Kenora, Patricia Portion (Cape Henrietta Maria. Brant River, Polar Bear Provincial Park). ROM. Recorded for the first time on the Ontario mainland near Hudson Bay on 30 July 1969, by T. O'Riley.

Family Riordinidae

None recorded in Ontario.

Family Libytheidae

Subfamily Libytheinae

Genus *Libytheana* Michener

L. bachmanii (Kirtland). 4: Bruce; Carleton; Elgin; Essex; Haldimand; Halton; Kent; Lambton; Lincoln; Manitoulin; Middlesex; Norfolk; Northumberland*; Prince Edward; Ontario; Waterloo; York. Mid May–early November. ROM. Rare outside of extreme southwestern Ontario.

Family Nymphalidae

Subfamily Nymphalinae

Tribe Nymphalini

Genus *Polygonia* Hübner

P. comma (Harris). 3: Algoma; Cochrane; Sudbury; Thunder Bay. 4: Bruce; Carleton; Dundas; Elgin; Essex; Frontenac; Haldimand*; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington*; Lincoln; Manitoulin; Middlesex; Muskoka*; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. Early March–early December. CNC, ROM.

P. faunus (Edwards). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Grey; Haliburton; Hastings; Lanark; Manitoulin; Muskoka; Nipissing; Parry Sound; Peel; Peterborough*; Renfrew; Russell; Wentworth. End April–early September. CNC, ROM.

P. gracilis (Grote and Robinson). 2: Thunder Bay. 3: Algoma; Cochrane; Sudbury; Thunder Bay. End May–end August. CNC, ROM. Not often reported.

P. interrogationis (Fabricius). 3: Rainy River; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Peel*; Peterborough; Renfrew; Russell; Simcoe; Stormont; Waterloo; Welland; Wellington; Wentworth; York. Early April–early November. CNC, ROM.

P. progne (Cramer). 2: Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant*; Bruce; Carleton; Dundas; Essex; Frontenac; Grenville; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton*; Lanark; Leeds; Lennox and Addington; Lincoln; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Victoria; Waterloo*; Wellington; Wentworth; York. End March–early October. CNC, ROM.

P. satyrus (Edwards). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay (Geraldton). 3: Algoma; Kenora; Rainy River; Thunder Bay; Timiskaming. 4: Manitoulin; Nipissing*. Mid May–end August. CNC, ROM. Subspecies *P. s. neomarsyas* dos Passos.

Genus *Nymphalis* Kluk

This genus is kept intact following Ehrlich (1961).

N. antiopa (Linné). 1: Kenora, Patricia Portion (Cape Henrietta Maria). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant*; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Haldimand*; Haliburton; Hastings; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Early March–early November. CNC, ROM.

N. milberti (Godart). 1: Kenora, Patricia Portion (Winisk). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Haldimand*; Halton; Huron; Kent; Lambton; Lanark; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prince Edward; Renfrew; Russell; Simcoe; Waterloo; Welland; Wentworth; York. Early March–early October. CNC, ROM.

N. vau-album (Denis and Schiffermüller). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Elgin*; Essex; Frontenac; Haldimand*; Haliburton; Halton; Huron*; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Muskoka; Nipissing; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Simcoe; Victoria; Waterloo; York. Early March–early October. CNC, ROM. The population can vary greatly. Subspecies *N. v. j-album* (Boisduval and Leconte).

Genus *Vanessa* Fabricius

- V. atalanta* (Linné). Red admiral/vanesse amirale. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grey; Haldimand*; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth*; Peterborough; Prescott; Prince Edward; Simcoe; Stormont; Russell; Victoria; Waterloo; Welland; Wellington*; Wentworth; York. End March–end October. CNC, ROM.
- V. cardui* (Linné). Painted lady/vanesse de l'artichaut. 1: Kenora, Patricia Portion (Winisk). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand*; Haliburton; Halton; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peterborough; Prince Edward; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. End April–mid November. CNC, ROM.
- V. virginiensis* (Drury). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grenville; Grey; Haldimand*; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. End March–mid October. CNC, ROM.

Tribe Hypolimnini

Genus *Junonia* Hübner

J. coenia (Hübner). 2: Thunder Bay. 3: Thunder Bay. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand*; Huron; Lambton; Leeds; Manitoulin; Nipissing; Norfolk; Northumberland; Ontario; Perth; Russell; Simcoe; Stormont*; Waterloo; Welland; Wentworth; York. Mid June–early November. AMNH, CNC, ROM. The only record from northern Ontario was in an abnormally hot year from Thunder Bay (Geraldton, 16 July 1955, A. B. Klots). Varies in number from year to year.

Subfamily Argynninae

Genus *Euptoieta* Doubleday

E. claudia (Cramer). Variegated fritillary/fritillaire panachée. 3: Algoma; Kenora; Rainy River; Thunder Bay. 4: Carleton; Elgin*; Essex; Haldimand*; Halton; Kent; Lambton; Norfolk; Perth; Waterloo; York. Mid June–end September. CNC, ROM. Uncommon. Reaches Ontario from the prairies in the west and from south of Lake Ontario and Lake Erie in the south.

Genus *Speyeria* Scudder

S. aphrodite (Fabricius). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Grey; Elgin; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Lambton; Lanark; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peel; Peterborough; Renfrew; Russell; Simcoe; Waterloo*; Welland; Wellington; York. End June–early September. CNC, ROM. Subspecies *S. a. winni* (Gunder). Subspecies *S. a. alcestitis* (Edwards) is known from southwestern Ontario (ROM).

S. atlantis (Edwards). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Dundas; Frontenac; Grey; Haliburton; Halton; Hastings; Huron; Lanark; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk*; Ontario*; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Waterloo*; Victoria; York. Mid June–early September. CNC, ROM.

- S. cybele* (Fabricius). 2: Kenora (Sioux Lookout). 3: Algoma*; Cochrane; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Early June–end September. CNC, ROM.
- S. idalia* (Drury). 4: Huron (Grand Bend); Kent (Rondeau Provincial Park); Middlesex (Hyde Park Corner); Norfolk*; Waterloo (Kitchener*). End July–end August. ROM. Very rare and not reported recently. See Lamb (1967).

Genus *Boloria* Moore

This genus is kept intact following Ehrlich (1961).

- B. bellona* (Fabricius). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk*; Northumberland; Ontario; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. Mid April–end September. CNC, ROM. Subspecies *B. b. toddi* (Holland) in northern Ontario.
- B. eunomia* (Esper). 1: Kenora, Patricia Portion (Cape Henrietta Maria. Fort Severn). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton (Mer Bleue*); Parry Sound; Nipissing. Early June–mid July. CNC, ROM. Very short flight period in any given area.
- Subspecies *B. e. tricularis* (Hübner) on the tundra; subspecies *B. e. dawsoni* (Barnes and McDunnough) in remaining localities; Thunder Bay (Hymers, type locality). See Freeman (1969) and Lafontaine (1972).
- B. freija* (Thunberg). 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Nipissing; Rainy River; Thunder Bay; Timiskaming (Kenogami). 4: Carleton (Mer Bleue*). Mid May–end June. CNC, ROM. See Freeman (1969) and Lafontaine (1972).
- B. frigga* (Thunberg). 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River. Mid May–end July. AMNH, CNC, ROM. The tundra population closely resembles

the population of Churchill, Manitoba, subspecies *B. f. saga* (Staudinger); the southern populations are distinctive with brighter basic wing colour and stronger black markings.

***B. selene* (Denis and Schiffermüller).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce*; Carleton; Dundas; Elgin; Frontenac; Grey; Haliburton; Halton*; Hastings; Huron; Lanark; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland*; Ontario; Parry Sound; Peel; Peterborough*; Prescott; Renfrew; Russell; Simcoe; Waterloo; Wellington*; Wentworth; York. End May–mid September. CNC, ROM. Subspecies *B. s. atrocistalis* (Huard) in the north; subspecies *B. s. myrina* (Cramer) in the south.

***B. titania* (Esper).** 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. Early July–end August. CNC, ROM. Subspecies *B. t. boisduvalii* (Duponchel) on the tundra. Subspecies *B. t. grandis* (Barnes and McDunnough) in the remaining localities; Thunder Bay (Hymers, type locality).

Subfamily Melitaeinae

Genus *Phyciodes* Hübner

The entry *P. campestris* (Behr) included in previous lists must be deleted. The male genitalia of some ROM specimens differ from *P. batesii* and *pascoensis*; however, it is not known to what extent this may be part of the normal range of variation in *batesii*.

***P. batesii* (Reakirt).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Elgin*; Frontenac; Halton; Lambton; Lanark; Kent; Manitoulin; Muskoka; Nipissing; Peterborough*; Simcoe; Waterloo; Welland; Wentworth; York. End May–early July. CNC, ROM.

***P. pascoensis* Wright** [= *P. morpheus* (Fabricius) = *P. selenis* W. Kirby]. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dufferin; Dundas; Durham; Elgin; Frontenac; Grey; Halton*; Hastings; Huron; Kent; Lambton; Lanark; Leeds (Chaffey's Locks); Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Parry Sound; Peel; Perth; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Stormont; Victoria; Waterloo; Welland; Wellington*;

Wentworth; York (Toronto). One brood, late June–early August. CNC, ROM. See also Opler and Krizek (1984); Scott (1986); and TEA Occasional Publications 1988, 19:14.

***P. tharos* (Drury).** 3: Kenora; Rainy River; Thunder Bay. 4: Essex (Point Pelee); Grenville; Haldimand; Haliburton; Kent; Leeds (Chaffey's Locks); Lincoln; Norfolk; Ontario; Peterborough; Simcoe; York (Toronto). ROM. Two broods in southern Ontario, May–early June and late July–September (P. A. Opler, personal comment; see also Opler and Krizek (1984); Scott (1986); and TEA Occasional Publications 1988, 19:14).

Genus *Charidryas* Scudder [= *Chlosyne* auct.]

***C. gorgone* (Hübner).** 4: York (Toronto). ROM. Three specimens from Toronto, Humber Valley, 6 June 1891; one specimen ex coll. Bethune without data, Toronto. Species now is considered to be extirpated in the province, but may eventually be found in other locations, especially near the western border.

***C. harrisii* (Scudder).** 3: Algoma; Sudbury; Thunder Bay. 4: Bruce; Carleton; Frontenac; Glengarry*; Grenville; Haliburton; Hastings; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peterborough; Prescott; Russell; Simcoe*; Waterloo*; Welland. Early June–end July. CNC, ROM.

***C. nycteis* (Doubleday).** 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Cochrane; Kenora; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Elgin; Essex; Haldimand*; Haliburton; Huron; Kent; Lambton; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peterborough*; Renfrew*; Simcoe; Victoria; Waterloo*; Welland; Wellington*; Wentworth*; York. End May–end July; a second brood has been reported in Windsor in August. CNC, ROM. Northern Ontario specimens resemble the subspecies *C. n. drusius* (Edwards) described from Jefferson County, Colorado. Other subspecies *C. n. reversa* (F. & R. Chermock).

Genus *Euphydryas* Scudder

***E. phaeton* (Drury).** 2: Thunder Bay. 3: Algoma*. 4: Brant; Carleton; Dundas; Elgin; Essex; Glengarry; Grenville; Grey; Haldimand*; Halton; Huron; Lambton; Lanark; Leeds; Lincoln; Middlesex; Norfolk; Northumberland; Ontario*; Peterborough*; Prescott; Russell; Simcoe*; Victoria; Waterloo; Welland; Wellington*; Wentworth; York. Mid June–end July. CNC, ROM. Local. The only record from Thunder Bay Dist. is a single large female (Macdiarmid, 22 July 1924, ROM).

Subfamily Limenitidinae
Tribe Limenitidini

Genus *Limenitis* Fabricius

L. archippus (Cramer). Viceroy/sylvain royal. 2: Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dufferin; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grenville*; Grey; Haldimand*; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario*; Oxford; Peel; Peterborough; Prince Edward; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria*; Waterloo; Wellington; Wentworth*; York. End May–mid October. CNC, ROM.

L. arthemis (Drury). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Nipissing; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Dundas; Durham; Frontenac; Glengarry; Grey; Haldimand*; Haliburton; Halton; Hastings; Huron; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Ontario; Parry Sound; Peel*; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; York. Early June–early September. CNC, ROM. The material from northwestern Ontario (Nakina–Geraldton–Nipigon and west) shows very strong traces of gradation to subspecies *L. a. rubrofasciata* Barnes and McDunnough in some specimens. This species is an excellent example of a very gradual and indistinct east-west cline.

The southern phenotype of this polymorphic species, *L. a. astyanax* (Fabricius), was found in the following counties: 4: Elgin; Essex; Haldimand; Kent; Lambton; Norfolk. End April–early September. CNC, ROM. A broad zone of intergradation between the two forms can be traced in the following counties: 4: Durham; Elgin; Essex; Haliburton; Halton; Huron; Kent; Lambton; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. CNC, ROM. For the solution of the *arthemis/astyanax* question see Platt and Brower (1968).

Subfamily Charaxinae
Tribe Apaturini

Genus *Asterocampa* Röber

- A. celtis* (Boisduval and Leconte). 4: Carleton (Nepean); Elgin*; Essex (Point Pelee, Pelee Island); Halton*; Middlesex. Mid June–mid September. CNC, ROM. See TEA Occasional Publications 1983, 14:6, and 1986, 17:20.
- A. clyton* (Boisduval and Leconte). 4: Brant (Brantford); Elgin; Essex; Huron; Kent (Bothwell); Lambton*; Middlesex (Byron); Victoria*; Waterloo. Early July–end August. ROM.

Family Satyridae
Subfamily Elymniinae
Tribe Parargini

Genus *Enodia* Hübner [= *Lethe* Hübner]

- E. anthedon* A. H. Clark. 2: Kenora. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry*; Grey; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. Mid June–mid August. CNC, ROM.

Genus *Satyrodes* Scudder [= *Lethe* Hübner]

- S. appalachia* (R. L. Chermock). 4: Brant; Carleton; Dundas; Elgin; Essex; Haldimand; Halton*; Huron; Kent; Lambton; Lanark; Manitoulin; Middlesex; Norfolk; Northumberland; Oxford; Peterborough; Prescott*; Waterloo; Wellington; Wentworth; York. Early June–early August. ROM. Probably more widely distributed in southern Ontario but not always recognized.
- S. eurydice* (Johansson). 2: Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Timiskaming. 4: Brant; Bruce; Carleton; Dufferin; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grey; Haliburton; Haldimand; Halton; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. Early June–mid September. CNC, ROM. Not common north of Sudbury.

Subfamily Satyrinae
Tribe Euptychiini

Genus *Megisto* Hübner [= *Euptychia* auct.]

***M. cymela* (Cramer).** 2: Kenora; 3: Algoma; Rainy River; Sudbury; Timiskaming; 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Glengarry; Grenville*; Grey; Halton; Haldimand*; Haliburton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth; York. End May–mid August. CNC, ROM. Essentially southern. There may be two cryptic species under this name, both of which may reach Ontario. See Opler and Krizek (1984).

Tribe Coenonymphini

Genus *Coenonympha* Hübner

***C. inornata* Edwards.** 1: Kenora, Patricia Portion (Winisk); 2: Cochrane; Thunder Bay; 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming; 4: Brant; Bruce; Carleton; Dufferin; Dundas; Durham; Frontenac; Glengarry; Grenville; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Muskoka; Nipissing; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Prescott; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Wellington; Wentworth; York. Mid May–early October. CNC, ROM. Recently has been spreading in southwestern Ontario. For information on supposed subspecies see TEA Occasional Publications.

Tribe Maniolini

Genus *Cercyonis* Scudder

***C. pegala* (Fabricius).** 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming; 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Grenville; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk;

Northumberland; Ontario; Parry Sound; Peel; Peterborough; Prescott*; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland*; Wellington; Wentworth; York. End June–early September. CNC, ROM.

Tribe Erebiini

Genus *Erebia* Dalman

- E. disa* (Thunberg). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Thunder Bay. End May–end June. CNC, ROM. Subspecies *E. d. mancinus* Doubleday and Hewitson.
- E. discoidalis* (Kirby). 1: Kenora, Patricia Portion (Fort Severn). 2: Kenora, Patricia Portion; Thunder Bay. 3: Cochrane; Kenora; Rainy River; Sudbury; Timiskaming. Early May–mid July. CNC, ROM.

Tribe Satyrini

Genus *Oeneis* Hübner

- O. chryxus* (Doubleday and Hewitson). 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haliburton; Hastings; Lanark; Leeds; Lennox and Addington; Manitoulin; Muskoka; Nipissing; Parry Sound; Peterborough; Renfrew; Simcoe; Victoria. Early May–mid June. CNC, ROM. Subspecies *O. c. strigulosa* McDunnough. The specimen from 2: Cochrane (Fraserdale, 13 June 1971, S. M. Daniels) may be subspecies *O. c. calais* (Scudder).
- O. jutta* (Hübner). 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Manitoulin; Nipissing; Parry Sound; Prescott (Alfred Bog); Russell. End May–end July. CNC, ROM. Subspecies *O. j. ascerta* Masters and Sorensen. The population on the tundra belongs to subspecies *O. j. alaskensis* Holland, known from Churchill, Manitoba.
- O. macounii* (Edwards). 2: Kenora; Thunder Bay. 3: Algoma; Rainy River; Thunder Bay (Nipigon, type locality). 4: Nipissing. End May–early August. CNC, ROM. Adults are said to fly only in even-numbered years.
- O. melissa* (Fabricius). 1: Kenora, Patricia Portion (Cape Henrietta Maria. Fort Severn). July. ROM. Subspecies *O. m. semplei* Holland.

Family Danaidae
Subfamily Danainae

Genus *Danaus* Kluk

***D. plexippus* (Linné).** Monarch butterfly/monarque. 2: Thunder Bay (Geraldton). 3: Algoma; Cochrane; Kenora (Minaki*); Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dundas; Durham; Elgin; Essex; Frontenac; Haldimand*; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Oxford; Parry Sound; Peel; Peterborough; Prince Edward; Renfrew; Russell; Simcoe; Stormont; Victoria; Waterloo; Welland; Wellington; Wentworth; York. Early May–early December. AMNH, CNC, ROM. Recorded only in abnormally hot years in Thunder Bay (Geraldton, 14 July 1955, A. B. Klots). In Nipigon, recorded normally every year in mid June. Not breeding in the north. Congregates along shores of Lake Ontario and Lake Erie and migrates in mass south to Mexico every winter. Descendants return to breed in early summer.

Superfamily Zygaenoidea
Family Zygaenidae

Genus *Harrisina* Packard

***H. americana* (Guérin-Ménéville).** 4: Essex; Haldimand; Kent; Norfolk. 19 June–23 July. CNC, McM, ROM. Occurs only in the southernmost part of the region along the north shore of Lake Erie.

Family Limacodidae

The tribes of Forbes (1923) are retained.

Tribe Cochliidiini

Genus *Tortricidia* Packard

***T. flexuosa* (Grote).** 4: Carleton; Durham; Elgin; Haldimand; Haliburton; Hastings; Kent; Leeds; Middlesex; Nipissing; Peterborough; Simcoe; Stormont. 8 June–29 July. CNC, LEM, ROM. A partial second generation may occur, e.g., Haldimand (Dunnville, 20 August). The form *caesonina* (Grote) was recorded at Chaffey's Locks.

T. pallida (Herrich-Schäffer). 4: Durham (Kendal); Leeds (Chaffeys Locks); Norfolk (St. Williams). 1–24 July. CNC, ROM. Known only from the three localities. Rare.

T. testacea Packard. 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Dufferin; Durham; Kent; Leeds; Peterborough; Simcoe; Wentworth. 5 June–20 July. AMNH, CNC, ROM.

Genus *Heterogenea* Knoch

H. shurtleffi Packard. 4: Leeds (Chaffeys Locks). 11–28 July. ROM.

Genus *Packardia* Grote and Robinson

P. elegans (Packard). 4: Norfolk; Peterborough. July. CNC, ROM.

P. geminata (Packard). 4: Carleton; Frontenac; Haldimand; Kent; Leeds; Norfolk; Peterborough. 14 June–6 July. CNC, ROM. Rare.

Genus *Lithacodes* Packard

L. fasciola (Herrich-Schäffer). 2: Kenora. 3: Rainy River; Sudbury. 4: Carleton; Durham; Haldimand; Halton; Hastings; Leeds; Muskoka; Nipissing; Peterborough; Simcoe; Wentworth. 23 June–27 July. AMNH, CNC, ROM.

Genus *Apoda* Haworth [= *Limacodes* Berthold]

A. biguttata (Packard). 4: Durham; Hastings; Huron; Leeds; Kent; Lambton; Peel; Wentworth. 23 June–28 July. CNC, ROM.

A. y-inversum Packard. 4: Bruce; Durham; Haldimand; Hastings; Leeds; Kent; Middlesex; Ontario; Peterborough; Wentworth; York. 16 June–24 July. CNC, ROM. Common.

Genus *Prolimacodes* Schaus

P. badia (Hübner) [= *scapha* (Harris)]. 4: Essex; Haldimand; Kent; Norfolk; Peterborough; Welland. July. CNC, ROM.

Tribe Phobetrini

Genus *Phobetron* Hübner

P. pitheciium (J. E. Smith). Hag moth/sorcière. 4: Carleton; Kent; Lambton; Leeds; Lincoln; Norfolk. June–July. CNC, ROM. Known only from a

few specimens from a few localities, mostly along the north shore of Lake Erie but also Chaffeys Locks and Ottawa.

Tribe Eucleini

Genus *Isa* Packard [= *Sisyrosea* Grote]

I. textula (Herrich-Schäffer). 4: Haldimand; Kent; Leeds; Norfolk; Wentworth. Mid July. CNC, ROM. Known from very few specimens from the southernmost part of the region, the north shore of Lake Erie, and Chaffeys Locks.

Genus *Adoneta* Clemens

A. spinuloides (Herrich-Schäffer). 4: Huron; Kent; York. 29 June–10 July. CNC. Known only from a few localities.

Genus *Euclea* Hübner

E. delphinii (Boisduval). 3: Thunder Bay. 4: Bruce; Carleton; Durham; Haldimand; Hastings; Kent; Leeds; Lincoln; Muskoka; Norfolk; Peel; Peterborough; Simcoe; Welland; Wentworth; York. 19 June–19 August. CNC, McM, ROM, UWO. Last date of capture usually at the end of July: Peel (Port Credit, 25 July); Bruce (Wiarton, 26 July); Chaffeys Locks, Hamilton, and Toronto, 17 July. A partial second generation may occur, e.g., Thunder Bay (Fort William, 19 August).

Superfamily Pyraloidea

Family Pyralidae

Pyralid moths/pyrales.

(Series Crambiformes)

Subfamily Scopariinae

Genus *Gesneria* Hübner

G. centuriella (Denis and Schiffermüller). 2. 3. 4. Subspecies *G. c. caecalis* (Walker).

Genus *Scoparia* Haworth

S. basalis Walker. 4.

S. biplagiata Walker. 3. 4.

- S. cinereomedia* Dyar. 4.
S. penumbralis Dyar. 4.

Genus *Eudonia* Billberg

- E. heterosalis* (McDunnough). 4.
E. lugubralis (Walker). 3. 4.
E. strigalis (Dyar). 4.

Subfamily Nymphulinae
Tribe Nymphulini

Genus *Neocataclysta* Lange

- N. magnificalis* (Hübner). 4.

Genus *Nymphula* Schrank

- N. ekthlipsis* (Grote). 3. 4.

Genus *Munroessa* Lange

- M. faulalis* (Walker). 4.
M. gyralis (Hulst). 3. 4. Subspecies *M. g. serralinealis* (Barnes and Benjamin).
M. icciusalis (Walker). 4.

Genus *Paraponyx* Hübner

- P. allionealis* Walker. 3. 4. Subspecies *P. a. itealis* (Walker).
P. badiusalis (Walker). 3. 4.
P. curviferalis (Walker). 4.
P. maculalis (Clemens). 2. 3. 4.
P. obscuralis (Grote). 4.

Tribe Argyractini

Genus *Petrophila* Guilding

- P. bifascialis* (Robinson). 4.
P. canadensis (Munroe). 4.
P. fulicalis (Clemens). 4.

Genus *Eoparargyractis* Lange

E. plevie (Dyar). 4.

Subfamily Odontiinae

Tribe Dichogamini

Genus *Metrea* Grote

M. ostreonalis Grote. 4.

Genus *Eustixia* Hübner

E. pupula Hübner. 4.

Tribe Odontiini

Genus *Microtheoris* Meyrick

M. ophionalis (Walker). 4. Subspecies *M. o. lacustris* Munroe.

Subfamily Glaphyriinae

Genus *Hellula* Guenée

H. rogatalis (Hulst). Cabbage webworm/tisseuse du chou. 4.

Genus *Nephrogramma* Munroe

N. reniculalis (Zeller). 4.

Genus *Stegea* Munroe

S. eripalis (Grote). 4.

Genus *Abegesta* Munroe

A. reluctalis (Hulst). 4.

Genus *Glaphyria* Hübner

G. glaphyralis (Guenée). 4.

G. sesquialis Hübner. 4.

Genus *Aethiophysa* Munroe

A. lentiflualis (Zeller). 4.

Genus *Xanthophysa* Munroe

X. psychialis (Hulst). 4.

Genus *Lipocosma* Lederer

L. sicalis (Walker). 4.

Genus *Lipocosmodes* Munroe

L. fuliginosalis (Fernald). 4.

Genus *Dicymolomia* Zeller

D. julianalis (Walker). 4.

Genus *Chalcoela* Zeller

C. iphitalis (Walker). 4.

C. pegasalis (Walker). 4.

Subfamily Evergestinae

Genus *Evergestis* Hübner

E. pallidata (Hufnagel). 4.

E. unimacula (Grote and Robinson). 4.

Subfamily Pyraustinae

Tribe Pyraustini

Genus *Saucrobotys* Munroe

S. fumoferalis (Hulst). 4.

Genus *Nascia* Curtis

N. acutella (Walker). 3. 4.

Genus *Crocidophora* Lederer

C. serratissimalis Zeller. 4.

C. tubercularis Lederer. 4.

Genus *Ostrinia* Hübner

O. marginalis (Walker). 4.

O. nubilalis (Hübner). European corn borer/pyrale du maïs. 4.

O. obumbratalis (Lederer). 4.

O. penitalis (Grote). 4.

Genus *Fumibotys* Munroe

F. fumalis (Guenée). 4.

Genus *Perispasta* Zeller

P. caeculalis Zeller. 3. 4.

Genus *Eurrhynx* Hübner

E. hortulata (Linné). 4.

Genus *Nealgedonia* Munroe

N. extricalis (Guenée). 3. 4.

Genus *Mutuuraia* Munroe

M. mysippusalis (Walker). 4.

Genus *Anania* Hübner

A. funebris (Ström). 4s: York (Toronto, 1986).

Genus *Hahncappsia* Munroe

H. mancalis (Lederer). 4.

H. marculenta (Grote and Robinson). 4.

H. neoblitalis Capps. 4.

H. pergilvalis (Hulst). 4.

Genus *Achyra* Guenée

A. rantalis (Guenée). 4.

Genus *Helvibotys* Munroe

H. helvialis (Walker). 4s.

Genus *Sitochroa* Hübner

S. chortalis (Grote). 4.

Genus *Uresiphita* Hübner

U. reversalis (Guenée). 4.

Genus *Loxostege* Hübner

L. anartalis (Grote). 2. 3. Subspecies *L. a. lulualis* (Hulst).

L. commixtalis (Walker). Alfalfa webworm/tisseuse de la luzerne. 4.

L. sticticalis (Linné). Beet webworm/tisseuse de la betterave. 4.

Genus *Pyrausta* Schrank

P. bicoloralis (Guenée). 4.

P. fodinalis (Lederer). 2. 3.

P. generosus (Grote and Robinson). 4s.

P. nicalis (Grote). 4.

P. orphisalis Walker. 4.

P. signatalis (Walker). 4.

P. socialis (Grote). 4.

P. subsequalis (Guenée) [= *borealis* Packard]. 2. 3. 4.

P. unifascialis (Packard). 4. Subspecies *P. u. subolivalis* (Packard).

Tribe Spilomelini

Genus *Udea* Guenée

U. inquinatalis (Zeller). 3.

U. itysalis (Walker). 3.

U. rubigalis (Guenée). Greenhouse leaf tier, celery leaf tier/lieuse des serres,
lieuse du céleri. 2. 3. 4.

U. sheppardi (McDunnough). 4.

Genus *Lineodes* Guenée

L. integra Zeller. 4s.

Genus *Loxostegopsis* Dyar

L. merrickalis (Barnes and McDunnough). 3. 4.

Genus *Mecyna* Guenée

M. luscitalis Barnes and McDunnough. 4.

M. submedialis (Grote). 4.

Genus *Diacme* Warren

D. adipaloides Grote and Robinson. 4s: Kent (Chatham).

Genus *Nomophila* Hübner

N. nearctica Munroe. Celery stalkworm/pyrale du céleri. 2. 3. 4.

Genus *Desmia* Westwood

D. funeralis (Hübner). Grape leaf roller/plieuse de la vigne. 3. 4.

D. maculalis Westwood. 3. 4.

Genus *Hymenia* Hübner

H. perspectalis (Hübner). 4.

Genus *Spoladea* Guenée

S. recurvalis (Fabricius). 4.

Genus *Diathrausta* Lederer

D. harlequinialis Dyar. 4.

D. reconditalis (Walker). 4.

Genus *Anageshna* Munroe

A. primordialis (Dyar). 4. Subspecies *A. p. vividior* Munroe.

Genus *Blepharomastix* Lederer

B. ranalis (Guenée). 4.

Genus *Diaphania* Hübner

D. hyalinata (Linné). Melonworm/pyrale du melon. 4. Occasional straggler.

Genus *Palpita* Hübner

P. aenescentalis Munroe. 4.

P. arsaltealis (Walker). 3. 4.

P. magniferalis (Walker). 3. 4.

P. quadristigmalis (Guenée). 4s; York (Toronto). Occasional straggler.

Genus *Polygrammodes* Guenée

P. flavidalis (Guenée). 4.

Genus *Pantographa* Lederer

P. limata (Grote and Robinson). Basswood leaf roller/enrouleuse du tilleul. 4.

Genus *Diastictis* Hübner

D. ventralis (Grote and Robinson). 4.

Genus *Framinghamia* Strand

F. helvalis (Walker). 3. 4.

Genus *Herpetogramma* Lederer

H. aeglealis (Walker). 4.

H. pertextalis (Lederer). 3. 4.

H. theseusalis (Walker). 4.

H. thestealis (Walker). 4.

Subfamily Schoenobiinae

Genus *Donacaula* Meyrick [= *Schoenobius* Duponchel]

D. longirostrella (Clemens). 4.

D. melinella (Clemens). 4. Subspecies *D. m. albicostella* Fernald.

Subfamily Crambinae

Tribe Ancyloleptini

Genus *Prionapteryx* Stephens

P. nebulifera Stephens. 4.

Tribe Crambini

Genus *Crambus* Fabricius

- C. agitatellus* Clemens. 2: Thunder Bay (Geraldton). 4.
C. albellus Clemens. 2: Thunder Bay (Geraldton). 4.
C. alienellus (Zincken). 2. Subspecies *C. a. labradoriensis* Christoph, restricted to acid bogs. 4. Subspecies *C. a. dissectus* Grote, uncertain status.
C. awemellus McDunnough. 2. 4: Lanark (Poland).
C. bidens Zeller. 2: Thunder Bay (Geraldton). 3: Rainy River (One Sided Lake).
C. caliginosellus Clemens. 4.
C. girardellus Clemens. 2. 3. 4.
C. hamellus (Thunberg). 2. 3.
C. laqueatellus Clemens. 4.
C. leachellus (Zincken). 2: Thunder Bay (Geraldton). 4.
C. luteolellus Clemens. 2: Thunder Bay (Geraldton). 4.
C. lyonsellus Haimbach. 2. Restricted to acid bogs.
C. pascuellus (Linné). 2: Thunder Bay (Geraldton). 3. 4. Subspecies *C. p. floridus* Zeller.
C. perlellus (Scopoli). 3. 4. Subspecies *C. p. innotatellus* Walker.
C. praefectellus (Zincken). 2: Thunder Bay (Geraldton). 3: Thunder Bay (Nipigon). 4.
C. unistriatellus Packard. 2. 3. 4.
C. whitmerellus Klots. 3. 4.
C. youngellus Kearfott. 4: Carleton (Mer Bleue Bog, type locality). Restricted to acid bogs.
C. zeellus Fernald. 4.

Genus *Chrysoteuchia* Hübner

- C. toparia* (Zeller). 2. 3. 4.

Genus *Arequipa* Walker

- A. turbatella* Walker. 4.

Genus *Rhaphiptera* Hampson

- R. argillaceella* (Packard). 4.

Genus *Platytes* Guenée

- P. vobisne* Dyar. 3: Kenora (Minaki).

Genus *Agriphila* Hübner

A. ruricolella (Zeller). 2. 4.

A. vulgivagella (Clemens). Vagabond crambus/pyrale vagabonde. 4.

Genus *Catoptria* Hübner

C. latiradiella (Walker). 2: Thunder Bay (Geraldton). 3. 4.

Genus *Pediasia* Hübner

P. abnaki (Klots). 3. 4.

P. dorsipunctella (Kearfott). 3.

P. trisecta (Walker). 4.

P. truncatella (Zetterstedt). 2.

Genus *Microcrambus* Bleszynski

M. biguttellus (Forbes). 4.

M. elegans (Clemens). 4.

M. minor (Forbes). 4.

Genus *Fissicrambus* Bleszynski

F. mutabilis Clemens. 4.

Genus *Thaumatopsis* Morrison

T. pexella (Zeller). 4: Muskoka (Rostrevor). Subspecies *T. p. gibsonella* Kearfott (Rostrevor, type locality).

Genus *Parapediasia* Bleszynski

P. teterrella (Zincken). 4.

Tribe *Argyriini*

Genus *Argyria* Hübner

A. sp. 4s. CNC.

Genus *Urola* Walker

U. nivalis (Drury). 4.

Genus *Vaxi* Bleszynski

V. auratella (Clemens). 3. 4.

V. critica (Forbes). 3. 4.

Tribe Chilini

Genus *Chilo* Zincken

C. plejadellus Zincken. 3. 4.

Genus *Thopeutis* Hübner

T. forbesellus (Fernald). 4.

Genus *Occidentalia* Dyar and Heinrich

O. comptulatalis (Hulst). 4.

Genus *Xubida* Schaus

X. panalope (Dyar). 4.

(Series *Pyaliformes*)

Subfamily *Pyalinae*

Genus *Pyalis* Linné

P. costiferalis Walker. 4.

P. disciferalis Dyar. 4.

P. farinalis Linné. 4.

Genus *Aglossa* Latreille

A. caprealis (Hübner). 4.

A. cuprina Zeller. 4.

Genus *Hypsopygia* Hübner

H. costalis (Fabricius). 3. 4.

Genus *Herculia* Walker

H. infimbrialis Dyar. 4: Simcoe.

H. intermedialis (Walker). 4.

H. olinalis (Guenée). 2. 4.

H. thymetusalis (Walker). 3. 4.

Subfamily Chrysauginae

Genus *Acallis* Ragonot

A. alticolalis (Dyar). 4: Carleton (Merivale).

Genus *Galasa* Walker

G. nigrinodis (Zeller). 3.

Genus *Tosale* Walker

T. oviplagalis (Walker). 4.

Genus *Arta* Grote

A. statalis Grote. 4.

Genus *Condylolomia* Grote

C. participalis Grote. 4.

Subfamily Epipaschiinae

Genus *Epipaschia* Clemens

E. superatalis Clemens. 4s: Kent (Chatham).

Genus *Oneida* Hulst

O. lunulalis (Hulst). 4n. 4s.

Genus *Tetralopha* Zeller

T. aplastella (Hulst). 3. 4.

T. asperatella (Clemens). 4.

T. baptisiella Fernald. 4s: Kent (Chatham)

- T. expandens* (Walker). 4.
T. maritimalis McDunnough. 4.
T. militella Zeller. 4.
T. robustella Zeller. 4.
T. vacciniivora Munroe. 3. 4.

Subfamily Galleriinae
Tribe Galleriini

Genus *Galleria* Fabricius

- G. mellonella* (Linné). Greater wax moth/teigne des ruches. 4.

Tribe Tirathabini

Genus *Aphomia* Hübner

- A. terrenella* Zeller. 4.

Genus *Corcyra* Ragonot

- C. cephalonica* (Stainton). 4. European dried food pest.

Subfamily Phycitinae

Genus *Acrobasis* Zeller

- A. amplexella* Ragonot. 3. 4.
A. angusella Grote. 4.
A. aurorella Ely. 4.
A. betulella Hulst. Birch tubemaker/pyrale tubicole du bouleau. 2. 3. 4.
A. caryae Grote. 4.
A. caryivorella Ragonot. 4.
A. comptoniella Hulst. 4.
A. demotella Grote. 4.
A. feltella Dyar. 4.
A. indigenella (Zeller). 4.
A. irrubriella Ely. 4.
A. juglandis (LeBaron). Pecan leaf casebearer/pyrale du noyer. 4.
A. kearfottella Dyar. 4.
A. ostryella Ely. 4.
A. palliolella Ragonot. 4: Carleton (Ottawa).

- A. rubrifasciella* Packard. Alder tubemaker/pyrale tubicole de l'aulne. 4.
A. secundella Ely. 4.
A. stigmella Dyar. 4.
A. sylviella Ely. 4.
A. tricolorella Grote. 4.
A. vaccinii Riley. Cranberry fruitworm/pyrale des atocas. 4n. 4s.

Genus *Myelopsis* Heinrich

- M. coniella* (Ragonot). 3. 4.
M. subtetricella (Ragonot). 2. 3. 4.

Genus *Apomyelois* Heinrich

- A. bistriatella* (Hulst). 2. 3. 4.

Genus *Etiella* Zeller

- E. zinckenella* (Treitschke). 4s: Welland (Port Colborne).

Genus *Glyptocera* Ragonot

- G. consobrinella* (Zeller). 4.

Genus *Immyrta* Dyar

- I. nigrovitella* Dyar. 4.

Genus *Oreana* Hulst

- O. unicolorella* (Hulst). 4.

Genus *Salebriaria* Heinrich

- S. engeli* (Dyar). 4.
S. tenebrosella (Hulst). 4.

Genus *Ortholepis* Ragonot

- O. myricella* McDunnough. 3. 4.
O. pasadamia (Dyar). 3. 4.

Genus *Meroptera* Grote

M. abditiva Heinrich. 4.

M. cviatella Dyar. 4.

M. pravella (Grote). Aspen leafroller/enrouleuse du tremble. 2. 3. 4.

Genus *Nephopterix* Hübner

N. basilaris Zeller. 3. 4.

N. carneella Hulst. 2. 3. 4.

N. celtidella (Hulst). 4.

N. fernaldi (Ragonot). 4.

N. subcaesiella (Clemens). 4.

N. subfuscella (Ragonot). 4.

N. vetustella (Dyar). 4.

N. virgatella (Clemens). 4.

Genus *Tlascala* Hulst

T. reductella (Walker). 4.

Genus *Tulsa* Heinrich

T. finitella (Walker). 4.

T. umbripennis (Hulst). 2: Cochrane (Smoky Falls).

Genus *Telethusia* Heinrich

T. ovalis (Packard). 4.

Genus *Pyla* Grote

P. aenigmatica Heinrich. 4.

P. fusca (Haworth). 2. 3. 4.

Genus *Dioryctria* Zeller

D. abietivorella (Grote). Fir coneworm/pyrale des cônes du sapin. 2. 3. 4.

D. banksiella Mutuura, Munroe, and Ross. 2. 3. 4.

D. disclusa Heinrich. 4.

D. reniculelloides Mutuura and Munroe. 4.

D. zimmermani (Grote). Zimmerman pine moth/pyrale des pousses du pin. 4.

Genus *Sarata* Ragonot [= *Megasis* Guenée]

S. caudellella (Dyar). 4n.

Genus *Canarsia* Hulst

C. ulmiarrosorella (Clemens). 4.

Genus *Psorosina* Dyar

P. hammondi (Riley). Appleleaf skeletonizer/squeletteuse du pommier. 4.

Genus *Homoeosoma* Curtis

H. stypticellum Grote. 3. 4.

Genus *Phycitodes* Hampson

P. albatella (Ragonot). 4. Subspecies *P. a. mucidella* (Ragonot).

Genus *Zophodia* Hübner

Z. convolutella (Hübner). Gooseberry fruitworm/pyrale des groseilles. 3. 4.

Genus *Euzophera* Zeller

E. semifuneralis (Walker). 4.

Genus *Eulogia* Heinrich

E. ochrifrontella (Zeller). 3. 4.

Genus *Ephesiodes* Ragonot

E. erythrella Ragonot. 4.

E. infimella Ragonot. 4.

Genus *Moodna* Hulst

M. ostrinella (Clemens). 4.

Genus *Vitula* Ragonot

V. edmandsii (Packard). 4.

Genus *Manhatta* Hulst

M. broweri Heinrich. 4.

Genus *Plodia* Guenée

P. interpunctella (Hübner). Indian meal moth/pyrale indienne de la farine. 4.
Common dried food pest.

Genus *Anagasta* Heinrich

A. kuehniella (Zeller). Mediterranean flour moth/pyrale méditerranéenne de la farine. 4. Common dried food pest.

Genus *Ephestia* Guenée

E. elutella (Hübner). Tobacco moth/pyrale du tabac. 4.

Genus *Cadra* Walker

C. cautella (Walker). Almond moth/pyrale des amandes. 4.
C. figulilella (Gregson). 4. Common dried food pest.

Genus *Eurythmia* Ragonot

E. angulella Ely. 3. 4.

Genus *Erelieva* Heinrich

E. parvulella (Ely). 4.

Genus *Coenochroa* Ragonot

C. illibella (Hulst). 4.

Subfamily Peoriinae

Genus *Peoria* Ragonot

P. approximella (Walker). 4.
P. gemmatella (Hulst). 4s: Kent (Chatham).
P. tetradella (Zeller). 4s: Welland (Port Colborne).

Superfamily Thyridioidea
Family Thyrididae
Subfamily Thyridinae

Genus *Thyris* Laspeyres

T. maculata Harris. 4.

T. sepulchralis Guérin-Ménéville [= *lugubris* Boisduval]. 4.

Superfamily Pterophoroidea
Family Pterophoridae
Subfamily Platyptiliinae

Genus *Sphenarches* Meyrick

S. ontario (McDunnough). 4: Norfolk (Normandale, type locality).

Genus *Geina* Tutt

G. buscki (McDunnough). 4: Muskoka (Go Home Bay, type locality).

G. periscelidactyla (Fitch). 4: Welland.

G. tenuidactyla (Fitch). 2: Cochrane (Moosonee). 3. 4.

G. n.sp. 4: Carleton (Ottawa); Essex; Haldimand; Hastings; Kent; Middlesex; Norfolk; Oxford; Peterborough; Russell; Victoria; Wellington; Wentworth; York (Toronto). CNC.

Genus *Capperia* Tutt

C. evansi (McDunnough). 3: Cochrane. 4: Bruce; Carleton (Ottawa); Frontenac; Haldimand; Kent; Leeds; Norfolk; Welland; Wellington. CNC.

Genus *Oxyptilus* Zeller

O. delawaricus Zeller. 4: Carleton (Ottawa).

Genus *Trichoptilus* Walsingham

T. lobidactylus (Fitch). 4.

Genus *Platyptilia* Hübner

P. carduidactyla (Riley). 4s: Essex (Point Pelee).

P. pallidactyla (Haworth). 2. 3. 4.

P. tesseradactyla (Linné). 4.

Genus *Amblyptilia* Hübner

A. pica (Walsingham). 4.

Genus *Stenoptilodes* Zimmerman

S. carolina (Kearfott). 4.

S. edwardsii (Fish). 4.

Genus *Stenoptilia* Hübner

S. coloradensis Fernald. 4: Carleton (Ottawa); Hastings. CNC.

Subfamily Pterophorinae

Genus *Adaina* Tutt

A. montana (Walsingham). 4.

Genus *Oidaematophorus* Wallengren

O. cretidactylus (Fitch). 4.

O. elliotii (Fernald). 4.

O. eupatorii (Fernald). 4.

O. glenni Cashatt. 4. CNC.

O. homodactylus (Walker). 2. 3. 4.

O. inquinatus Zeller. 4.

O. kellicottii (Fish). 4. CNC.

O. lacteodactylus (Chambers). 4.

O. mathewianus (Zeller). 4.

O. phacelliae McDunnough. 4.

O. stramineus (Walsingham). 4.

Genus *Emmelina* Tutt

E. monodactyla (Linné). 4.

Superfamily Drepanoidea
Family Thyatiridae
Subfamily Thyatirinae
Tribe Habrosynini

Genus *Habrosyne* Hübner

- H. gloriosa* (Guenée) [= *rectangulata* Ottolengui]. 2: Cochrane. 4: Bruce; Carleton; Haldimand; Manitoulin; Middlesex; Simcoe. 7 June–24 July. AMNH, CNC, ROM, UG. Known from only a few specimens
- H. scripta* (Gosse). 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Durham; Frontenac; Haldimand; Hastings; Kent; Manitoulin; Middlesex; Muskoka; Nipissing; Parry Sound; Peel; Peterborough; Simcoe; Wentworth; York. 22 May–September. AMNH, Beth, CNC, McM, ROM, UG. A partial second generation occurs in late August in western Ontario, and in September in southern Ontario. The species is sympatric with *H. gloriosa*, which is probably often overlooked.

Genus *Pseudothyatira* Grote

- P. cymatophoroides* (Guenée). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Northumberland; Peterborough; Renfrew; York. 5 May–30 August. AMNH, Beth, CNC, McM, ROM, UG. A partial second generation may occur.
- P. c.* form *expultrix* (Grote). 2: Cochrane; Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Lambton; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Wentworth; York. Same flight period as *P. cymatophoroides*. AMNH, Beth, CNC, McM, ROM, UG. In his revision of the thyatirids, Werny (1966) showed *cymatophoroides* and *expultrix* to be different species. Ferguson (1983), however, proved by rearing larvae obtained from eggs laid by a female of the nominate form (*cymatophoroides*) that the progeny of 37 adult moths split 16:21 between *cymatophoroides* and *expultrix*.

Tribe Macrothyatirini

Genus *Euthyatira* Smith

- E. pudens* (Guenée). 2: Cochrane; Thunder Bay. 3: Thunder Bay. 4: Carleton; Frontenac; Haldimand; Hastings; Leeds; Manitoulin; Middlesex; Muskoka; York. 2 May–6 June. AMNH, CNC, McM, ROM, UG.

Local and rare, with few specimens in collections. Werny (1966) considered *E. pudens* and *E. p.* form *pennsylvanica* Smith to be distinct species. Ferguson (1983) raises reasonable doubts and for the time being should be followed. The only known locality for form *pennsylvanica* in Ontario is Frontenac (Perth Road, near Kingston).

Family Drepanidae
Subfamily Drepaninae

Genus Drepana Schrank

- D. arcuata* Walker.** 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Manitoulin; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Welland; Wellington; Wentworth; York. AMNH, CNC, LEM, McM, ROM, UG, UWO. First generation 2 May (Sudbury Dist.), 12 May (Leeds Co.)–5 July (Nipissing Dist.), 29 July (Algoma Dist.); second generation 1 July (Wentworth Co.), 10 July (Sudbury Dist.)–16 August (Cochrane Dist.), 27 August (York Co.).
- D. bilineata* (Packard).** 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haliburton; Hastings; Kent; Lanark; Leeds; Muskoka; Nipissing; Peterborough; Simcoe; York. AMNH, CNC, QU, ROM, UG, UWO. First generation 1 May (Carleton Co.), 29 May (Sudbury Dist.)–28 June (York Co.), 14 July (Thunder Bay Dist.); second generation 26 June (Muskoka Dist.), 2 July (Sudbury Dist.)–14 August (York Co.), 21 August (Sudbury Dist.). Live specimens of a partial third generation were caught in Toronto (ROM) as late as 21 August, and even 18 November, 1954. They are indistinguishable from specimens of the spring generation.

Genus Eudeilinea Packard

- E. herminiata* (Guenée).** 2: Cochrane; Thunder Bay. 3: Rainy River; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Hastings; Kent; Middlesex; Norfolk; Simcoe; Victoria; Waterloo; Wentworth. 5 June–22 July. AMNH, Beth, CNC, ROM, UG. The form *biseriata* (Packard) occurs along with the nominate form.

Subfamily Oretinae

Genus *Oreta* Walker

O. rosea (Walker). 2: Cochrane; Kenora, Patricia Portion. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Frontenac; Haldimand; Hastings; Kent; Leeds; Norfolk; Renfrew; Simcoe; Victoria; Wentworth. AMNH, CNC, McM, ROM, UG. First generation 15 June (Haldimand Co.)–17 July (Carleton Co.); second generation 23 July (Carleton Co.)–18 August (Haldimand Co.). In the boreal forest region, only one generation occurs from 2 July, Smoky Falls (in Cochrane Dist.)–25 July, Vermilion Bay (in Kenora Dist.). The form *irrorata* (Packard) occurs along with the nominate form.

Superfamily Geometroidea

Family Geometridae

Subfamily Archiearinae

See McGuffin (1988).

Genus *Archiearis* Hübner

A. infans (Möschler). 3: Rainy River; Sudbury. 4: Carleton; Halton; Leeds; Nipissing; Ontario; Parry Sound; Peterborough; Simcoe; Wentworth; York. CNC, ROM.

Genus *Leucobrephos* Grote

L. brephoides (Walker). 3: Thunder Bay (Hymers). Known only from this locality. ROM.

Subfamily Oenochrominae

See McGuffin (1988).

Genus *Alsophila* Hübner

A. pometaria (Harris). Fall cankerworm/arpenteuse d'automne. 3: Kenora. 4: Brant; Carleton; Durham; Elgin; Haldimand; Hastings; Lambton; Leeds; Muskoka; Norfolk; Peel; Prince Edward; Simcoe; Victoria; Waterloo; Welland; York. CNC, ROM.

Subfamily Geometrinae

See Ferguson (1969); and McGuffin (1988).

Tribe Nemoriini

Genus *Nemoria* Hübner

- N. bistriaria* **Hübner**. 3: Sudbury. 4: Carleton; Hastings; Leeds; Renfrew. CNC, ROM. Subspecies *N. b. rubromarginaria* (Packard).
N. mimosaria (**Guenée**). 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Haliburton; Hastings; Leeds; Nipissing; Parry Sound; Russell; Simcoe; Victoria; Welland; York. CNC, ROM.
N. rubrifrontaria (**Packard**). 2: Cochrane. 3: Sudbury. 4: Carleton; Leeds; Muskoka; Simcoe; Victoria; York. CNC, ROM.

Genus *Dichorda* Warren

- D. iridaria* (**Guenée**). 4: Essex; Hastings; Leeds; Renfrew; Welland. CNC, ROM.

Tribe Synchlorini

Genus *Synchlora* Guenée

- S. aerata* (**Fabricius**) [= *liquoraria* Guenée]. 2: Thunder Bay. 3: Kenora; Rainy River; Thunder Bay. 4: Bruce; Carleton; Essex; Hastings; Leeds; Lincoln; Muskoka; Norfolk; Simcoe; Welland; York. AMNH, CNC, ROM. Subspecific status uncertain.

Tribe Hemitheini

Genus *Chlorochlamys* Hulst

- C. chloroleucaria* (**Guenée**). 3: Rainy River; Sudbury. 4: Bruce; Carleton; Essex; Haldimand; Hastings; Leeds; Norfolk; Renfrew; Simcoe; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.

Genus *Hethemia* Ferguson

- H. pistasciaria* (**Guenée**). 3: Algoma. 4: Carleton; Essex; Haldimand; Leeds. AMNH, CNC, ROM.

Genus *Mesothea* Warren

- M. incertata* (**Walker**). 3: Cochrane; Thunder Bay. 4: Carleton; Haliburton; Leeds; Middlesex; Parry Sound. AMNH, CNC, ROM.

Subfamily Sterrhinae

See McGuffin (1967).

Tribe Sterrhini

Genus *Idaea* Treitschke [= *Sterrha* Hübner]

- I. demissaria* (Hübner). 3: Sudbury. 4: Frontenac; Haldimand; Hastings; Huron; Lincoln; Peterborough. AMNH, CNC, ROM.
I. dimidiata (Hufnagel). 4: Carleton. CNC.
I. rotundopennata (Packard). 2: Kenora. 4: Frontenac; Hastings; Muskoka. CNC, ROM.
I. violacearia (Walker). 4: Lambton (*teste* Schweitzer).

Tribe Cosymbiini

Genus *Pleuroprucha* Möschler

- P. insulsaria* (Guenée). 3: Sudbury; Thunder Bay. 4: Carleton; Essex; Frontenac; Haldimand; Halton; Hastings; Kent; Leeds; Norfolk; Peterborough; York. AMNH, CNC, ROM.

Genus *Cyclophora* Hübner

- C. pendulinaria* (Guenée). 2: Cochrane; Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Grenville; Haldimand; Hastings; Leeds; Muskoka; Nipissing; Parry Sound; Peterborough; Renfrew; Simcoe; York. AMNH, CNC, ROM.

Tribe Timandrini

Genus *Haematopis* Hübner

- H. grataria* (Fabricius). 4: Bruce; Carleton; Essex; Frontenac; Grenville; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Norfolk; Peel; Simcoe; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.

Tribe Scopulini

Genus *Scopula* Schrank

- S. ancillata* (Hulst). 2: Thunder Bay (Geraldton). 3: Cochrane (Low Bush); Thunder Bay (Black Sturgeon Lake). AMNH, CNC, ROM.
- S. cacuminaria* (Morrison). 3: Rainy River; Sudbury. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Leeds; Victoria; Welland; York. AMNH, CNC, ROM.
- S. frigidaria* (Möschler). 2: Cochrane (Ogoki Post). CNC. *S. inductata* (Guenée). 2: Cochrane. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Essex; Frontenac; Haldimand; Hastings; Leeds; Muskoka; Norfolk; Ontario; Peel; Waterloo; Welland; Wentworth; York. AMNH, CNC, ROM.
- S. junctaria* (Walker). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Renfrew; Victoria; York. AMNH, CNC, ROM.
- S. limboundata* (Haworth). 2: Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Kent; Leeds; Muskoka; Norfolk; Peterborough; Renfrew; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.
- S. purata* (Guenée). 4s.
- S. quadrilineata* (Packard). 2: Thunder Bay. 4: Carleton; Frontenac; Hastings; Parry Sound; Victoria; Wentworth. AMNH, CNC.

Genus *Leptostales* Möschler [= *Xystrota* Hulst]

- L. ferruminaria* (Zeller). 4: Carleton; Frontenac; Hastings; Leeds; Victoria. AMNH, CNC, ROM.

Subfamily Larentiinae Tribe Hydriomenini

Genus *Dysstroma* Hübner

- D. citrata* (Linné). 2: Cochrane (Ogoki Post). 3: Thunder Bay (South Neebing Twp.). 4: Haldimand; Nipissing; Ontario; Peterborough; York. CNC, ROM.
- D. hersiliata* (Guenée). 2: Thunder Bay. 3: Rainy River; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Leeds; Norfolk; Peterborough; Welland; Wentworth; York.

AMNH, CNC, ROM. Subspecies *D. h. cervinifascia* (Walker); *D. h. manitoba* McDunnough.

- D. truncata* (Hufnagel). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Hastings; Middlesex; Nipissing; Peterborough. AMNH, CNC, ROM.
D. sp. 2: Thunder Bay (Geraldton). AMNH.

Genus *Eulithis* Hübner [= *Lygris* Hübner]

- E. destinata* (Möschler). 2: Cochrane; Thunder Bay. AMNH, CNC.
E. diversilineata (Hübner). Grapevine looper/arpenteuse de la vigne. 3: Thunder Bay. 4: Essex; Frontenac; Haldimand; Hastings; Huron; Kent; Lambton; Leeds; Muskoka; Peterborough; Wentworth; York. AMNH, CNC, ROM.
E. explanata (Walker). 2: Cochrane; Kenora. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Leeds; Muskoka; Nipissing; Peterborough. AMNH, CNC, ROM.
E. gracilineata (Guenée). 4: Carleton; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Nipissing; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.
E. molliculata (Walker). 4: Bruce; Haldimand; Lincoln; Middlesex; Simcoe. CNC, ROM.
E. propulsata (Walker). 2: Cochrane; Kenora. 3: Rainy River. 4: Carleton; Durham. CNC, ROM.
E. serrataria (Barnes and McDunnough). 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Peterborough. AMNH, CNC, ROM.
E. testata (Linné). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Frontenac; Haldimand; Hastings; Leeds; Wentworth; York. AMNH, CNC, ROM.

Genus *Eustroma* Hübner

- E. semiatrata* (Hulst). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Timiskaming; Thunder Bay. 4: Frontenac; Norfolk. AMNH, CNC.

Genus *Ecliptoptera* Warren [= *Diactinia* Warren]

- E. silaceata* (Denis and Schiffermüller). 2: Cochrane. 3: Thunder Bay; Timiskaming. 4: Halton; Hastings; Leeds; Peel; Peterborough; York. CNC, ROM. Subspecies *E. s. albolineata* (Packard).

Genus *Plemyria* Hübner

- P. georgii* Hulst. 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Peterborough. CNC, ROM.

Genus *Thera* Stephens

- T. contractata* (Packard). 3: Sudbury. 4: Frontenac; Leeds; Manitoulin; Muskoka. CNC.
T. juniperata (Linné). 4: Frontenac; Haldimand; Leeds; York. AMNH, CNC, ROM.

Genus *Hydriomena* Hübner

- H. divisaria* (Walker). Transverse-banded looper/arpenteuse à bandes transversales. 2: Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Lanark; Leeds; Nipissing; Renfrew; Victoria. AMNH, CNC.
H. furcata (Thunberg). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. AMNH, CNC.
H. perfracta Swett. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. AMNH, CNC.
H. renunciata (Walker). 2: Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Essex; Frontenac; Hastings; Leeds; Muskoka; Nipissing; Parry Sound; Peterborough; Simcoe; Wentworth; York. AMNH, CNC, ROM.
H. ruberata (Freyer). 3: Sudbury. 4: Carleton; Hastings; Middlesex; Parry Sound. AMNH, CNC, ROM.
H. transfigurata Swett. 3: Thunder Bay (Black Sturgeon Lake). CNC.

Genus *Triphosa* Stephens

- T. haesitata* (Guenée). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Simcoe; York. AMNH, CNC, ROM. Subspecies *T. h. affirmaria* (Walker).

Genus *Coryphista* Hulst

- C. meadii* (Packard). 4: Carleton; Essex; Frontenac; Hastings; Leeds; Peterborough; Wentworth; York. CNC. Subspecies *C. m. atlantica* Munroe.

Genus *Hydria* Hübner

- H. undulata* (Linné). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Algoma; Carleton; Essex; Frontenac; Haliburton; Hastings; Huron; Leeds; Muskoka; Nipissing; Norfolk; Ontario; Parry Sound; Peterborough; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Genus *Rheumaptera* Hübner

- R. hastata* (Linné). Birch looper/arpenteuse noire du bouleau. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Haldimand; Haliburton; Hastings; Leeds; Nipissing; Norfolk; Ontario; Wentworth; York. AMNH, CNC, ROM.
- R. subhastata* (Nolcken). 2: Cochrane. 4: Parry Sound. CNC. Subspecies *R. s. stygiata* McDunnough.

Genus *Entephria* Hübner [= *Dasyuris* auct.]

- E. aurata* (Packard). 2: Thunder Bay. 4: Carleton; Peterborough. AMNH, CNC, ROM.
- E. polata* (Duponchel). 1: Kenora, Patricia Portion (Cape Henrietta Maria). ROM. Subspecies not ascertained; subspecies *E. p. bradorata* (Munroe).

Genus *Mesoleuca* Hübner

- M. ruficillata* (Guenée). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Hastings; Leeds; Middlesex; Nipissing; Norfolk; Simcoe; York. AMNH, CNC, ROM.

Genus *Spargania* Guenée

- S. luctuata* (Denis and Schiffermüller). 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton. AMNH, CNC, ROM. Subspecies *S. l. obductata* (Möschler).
- S. magnoliata* Guenée. 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora; Thunder Bay. 3: Sudbury. 4: Carleton; Dufferin; Durham; Frontenac; Hastings; Middlesex; Muskoka; Parry Sound; Wentworth. AMNH, CNC, ROM.

Genus *Perizoma* Hübner

- P. basaliatum* (Walker). 2: Cochrane; Thunder Bay. 3: Rainy River; Thunder Bay. 4: Carleton; Dufferin; Hastings; Leeds. AMNH, CNC, ROM.

Genus *Anticlea* Stephens [= *Earophila* Gumpenberg]

- A. multifurcata* (Walker). 2: Cochrane; Thunder Bay. 3: Algoma. 4: Carleton; Durham; Haldimand; Hastings; Kent; Leeds; Middlesex; Norfolk; York. AMNH, CNC, ROM.

A. vasiliata Guenée. 2: Cochrane; Thunder Bay. 3: Algoma; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds. AMNH, CNC, ROM.

Tribe Stamnodini

Genus *Stamnodes* Guenée

S. gibbicostata (Walker). 4: Carleton; Middlesex; Welland. CNC, ROM.

Tribe Xanthorhoini

Genus *Xanthoroe* Hübner

X. abrasaria (Herrich-Schäffer). 2: Cochrane (Moosonee. Ogoki Post. Smoky Falls). 3: Thunder Bay (Black Sturgeon Lake). CNC.

X. algidata (Möschler). 2: Cochrane (Ogoki Post). CNC.

X. ferrugata (Clerk). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Frontenac; Hastings; Leeds; Norfolk; Victoria; Wentworth; York. AMNH, CNC, ROM.

X. iduata (Guenée). 2: Cochrane; Kenora. 3: Sudbury; Thunder Bay. 4: Frontenac; Leeds. CNC.

X. labradorensis (Packard) [= *designata* auct.]. 1: Kenora, Patricia Portion (Fort Severn). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Sudbury. 4: Carleton; Frontenac; Hastings; Leeds; Middlesex; Ontario; Renfrew; Simcoe; Wentworth; York. AMNH, CNC, ROM.

X. lacustrata (Guenée). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Lincoln; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.

X. munitata (Hübner). 2: Cochrane; Thunder Bay. 3: Rainy River; Thunder Bay. AMNH, CNC, ROM.

X. packardata McDunnough. 2: Cochrane. 3: Rainy River; Thunder Bay. 4: Carleton; Hastings; Norfolk; Parry Sound; Victoria; York. CNC.

Genus *Epirrhoe* Hübner

E. alternata (Müller). 1: Kenora, Patricia Portion (Cape Henrietta Maria). 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Leeds; Norfolk; York. AMNH, CNC, ROM.

E. sperryi Herbulot. 2: Cochrane (Fort Albany); Thunder Bay (Geraldton). AMNH, ROM.

Genus *Euphyia* Hübner

E. unangulata (Haworth). 2: Cochrane. 3: Rainy River; Sudbury. 4: Carleton; Essex; Frontenac; Haldimand; Hastings; Leeds; Muskoka; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM. Subspecies *E. u. intermediata* (Guenée).

Genus *Orthonama* Hübner [= *Nycterosea* Hulst]

O. centrostrigaria (Wollaston). 3: Algoma; Sudbury. 4: Bruce; Carleton; Durham; Essex; Frontenac; Hastings; Kent; Leeds; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.

O. evansi (McDunnough). 4: Hastings; Victoria. CNC.

O. obstipata (Fabricius). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Hastings; Kent; Leeds; Norfolk; Renfrew; Wellington; Wentworth; York. AMNH, CNC, ROM.

Tribe Asthenini

Genus *Hydrelia* Hübner

H. albifera (Walker). 2: Cochrane. 3: Sudbury; Thunder Bay. 4: Carleton; Haldimand; Hastings; Kent; Middlesex; York. AMNH, CNC, ROM.

H. condensata (Walker). 3: Algoma; Thunder Bay. 4: Carleton; Dufferin; Hastings; Middlesex; Parry Sound; Simcoe; Victoria. AMNH, CNC, ROM.

H. inornata (Hulst). 2: Cochrane. 3: Algoma; Thunder Bay. 4: Carleton; Nipissing; Renfrew. CNC.

Genus *Venusia* Curtis

V. cambrica Curtis. 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. AMNH, CNC.

V. comptaria (Walker). 3: Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Middlesex; York. AMNH, CNC, ROM.

Genus *Trichodezia* Warren

T. albovittata (Guenée). 2: Cochrane; Thunder Bay. 3: Rainy River; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Middlesex; Nipissing; Norfolk; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.

Tribe Operophterini

Genus *Epirrita* Hübner

- E. autumnata* (Borkhausen). 3: Sudbury. 4: Carleton; Frontenac; Hastings; Leeds; Middlesex. AMNH, CNC, ROM. Subspecies *E. a. henshawi* (Swett).

Genus *Operophtera* Hübner

- O. bruceata* (Hulst). 2: Thunder Bay. 3: Algoma; Timiskaming. 4: Carleton; Dufferin; Frontenac; Haldimand; Hastings; Leeds; Nipissing; Wellington; York. AMNH, CNC, ROM.

Tribe Eudulini

Genus *Eubaphe* Hübner

- E. mendica* (Walker). 3: Algoma; Rainy River. 4: Carleton; Dufferin; Durham; Essex; Frontenac; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Simcoe; York. AMNH, CNC, ROM.

Tribe Eupitheciini

Genus *Horisme* Hübner

- H. intestinata* (Guenée). 2: Cochrane. 3: Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Nipissing; Norfolk; Simcoe; Welland; York. AMNH, CNC, ROM.

Genus *Eupithecia* Curtis

- E. absinthiata* (Clark) [= *coagulata* Guenée]. 2: Thunder Bay. 4: Frontenac; Haldimand; York. CNC.
- E. affinata* Pearsall. 4: Carleton; Hastings; Lanark; Leeds. CNC.
- E. albicapitata* Packard. 3: Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Parry Sound. CNC.
- E. annulata* (Hulst). 2: Thunder Bay. 3: Algoma; Cochrane; Sudbury; Thunder Bay. 4: Carleton (Lanark Sound). CNC.
- E. anticaria* Walker. 2: Thunder Bay. 3: Sudbury. AMNH.
- E. bowmani* Cassino and Swett. 4: Manitoulin. CNC.
- E. bradorata* McDunnough. 3: Sudbury; Thunder Bay. CNC.
- E. cemicifugata* Pearsall. 4: Carleton; Hastings. CNC.

- E. coloradensis* (Hulst). 3: Thunder Bay. 4: Lanark. CNC.
- E. columbiata* (Dyar). 3: Thunder Bay (Nipigon). 4: Frontenac; Leeds. AMNH, CNC.
- E. cretacea* (Packard). 4: Manitoulin. CNC.
- E. fletcherata* Taylor. 2: Thunder Bay (Geraldton). 4: Frontenac. AMNH.
- E. fumosa* (Hulst). 4: Carleton; Hastings; Lanark; Simcoe. CNC.
- E. gelidata* (Möschler). 2: Cochrane. 3: Sudbury. 4: Carleton. CNC.
- E. interruptofasciata* Packard. 2: Cochrane. 4: Carleton; Bruce; Frontenac; Hastings; Muskoka. CNC.
- E. intricata* (Zetterstedt). 4: Leeds (Chaffey's Locks). AMNH. Subspecies *E. i. taylorata* Swett.
- E. johnstoni* McDunnough. 4: Hastings; Lanark. CNC.
- E. lariciata* Freyer. Fir-needle inchworm/petite arpenteuse du sapin. 3: Thunder Bay (Nipigon). 4: Frontenac. AMNH, CNC.
- E. miserulata* Grote. 3: Sudbury. 4: Frontenac; Leeds; York. AMNH, CNC.
- E. misturata* (Hulst). 2: Cochrane. 3: Sudbury. 4: Carleton; Hastings. CNC.
- E. mutata* Pearsall. 3: Cochrane; Thunder Bay. 4: Carleton; Grey; Lanark; Leeds; Nipissing; Norfolk; Simcoe. CNC.
- E. nimbicolor* (Hulst). 2: Thunder Bay (Geraldton). 3: Sudbury. AMNH.
- E. palpata* Packard. Small pine looper/petite arpenteuse du pin. 2: Thunder Bay (Nakina). 3: Rainy River. 4: Carleton; Frontenac; Hastings; Leeds. AMNH, CNC.
- E. perfusca* (Hulst). 3: Rainy River; Sudbury; Thunder Bay. 4: Frontenac; York. AMNH, CNC. Subspecies *E. p. youngata* Taylor.
- E. ravocostaliata* Packard. 2: Thunder Bay. 3: Sudbury; Thunder Bay. 4: Frontenac; Leeds. AMNH, CNC.
- E. russeliata* Swett. 3: Sudbury. 4: Frontenac. AMNH, CNC.
- E. satyrata* (Hübner). 2: Thunder Bay. 3: Sudbury. 4: Leeds. AMNH, CNC. Subspecies *E. s. fumata* Taylor.
- E. sheppardata* McDunnough. 3: Sudbury. 4: Carleton. CNC.
- E. strattonata* Packard. 3: Sudbury (Sudbury). AMNH.
- E. subfuscata* (Haworth) [= *castigata* (Hübner)]. 2: Thunder Bay. 3: Sudbury; Thunder Bay. 4: Leeds; York. AMNH, CNC.
- E. swetti* Grossbeck. 4: Carleton; Lanark; Leeds; Peterborough. CNC.
- E. tripunctaria* Herrich-Schäffer. 3: Thunder Bay. CNC.

Tribe Lobophorini

Genus *Carsia* Hübner

- C. sororiata* (Hübner). 2: Cochrane. 3: Rainy River. CNC. Subspecies not ascertained.

Genus *Acasis* Duponchel

A. viridata (Packard). 3: Thunder Bay. 4: Carleton. CNC.

Genus *Cladara* Hulst

C. atroliturata (Walker). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Frontenac; Leeds; Middlesex; Muskoka; Peterborough. AMNH, CNC, ROM.

C. limitaria (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury. 4: Carleton; Frontenac; Hastings; Leeds; Muskoka; Peel; York. AMNH, CNC, ROM.

Genus *Lobophora* Curtis

L. nivigerata Walker. 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Essex; Frontenac; Haldimand; Haliburton; Hastings; Leeds; Muskoka; Norfolk; Peterborough; Wentworth; York. AMNH, CNC, ROM.

Genus *Heterophleps* Herrich-Schäffer

H. refusaria (Walker). 4: Carleton; Durham; Essex; Huron; Middlesex; Norfolk. CNC, ROM.

H. triguttaria Herrich-Schäffer. 4: Carleton; Essex; Frontenac; Hastings; Huron; Leeds; Lincoln; Middlesex; Norfolk; Simcoe; Victoria; York. CNC, ROM.

Genus *Dyspteris* Hübner

D. abortivaria (Herrich-Schäffer). 4: Carleton; Durham; Essex; Frontenac; Hastings; Kent; Leeds; Norfolk; Ontario; Peel. CNC, ROM.

Subfamily Ennominae

See McGuffin (1972, 1977, 1981, 1987).

Tribe Abraxini

Genus *Heliomata* Grote and Robinson

H. cycladata Grote and Robinson. 4: Carleton; Haldimand; Leeds; Norfolk; Simcoe; Welland; York. AMNH, CNC, ROM.

Genus *Protitame* McDunnough

- P. virginalis* (Hulst). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haliburton; Hastings; Leeds; Peterborough; Prescott; Simcoe; Wentworth. AMNH, CNC, ROM.

Tribe Semiothisini

See McGuffin (1972).

Genus *Eumacaria* Packard

- E. latiferrugata* (Walker). 4: Carleton; Leeds; Muskoka; York. CNC, ROM.

Genus *Itame* Hübner

- I. abruptata* (Walker). 4: Bruce; Muskoka. CNC.
- I. anataria* (Swett). 2: Thunder Bay. 3: Kenora; Rainy River; Sudbury; Thunder Bay. 4: Haldimand; Leeds; Muskoka; Wentworth. AMNH, CNC, ROM.
- I. andersoni* (Swett). 2: Cochrane; Thunder Bay. 3: Kenora; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peterborough. AMNH, CNC, ROM. Subspecies *I. a. orientis* Ferguson.
- I. argillacearia* (Packard). 4: Hastings; Huron; Norfolk. CNC, ROM.
- I. bitactata* (Walker). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Hastings; Nipissing. AMNH, CNC.
- I. brunneata* (Thunberg). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Leeds. AMNH, CNC.
- I. coortaria* (Hulst). 4: Durham; Elgin; Essex; Halton; Hastings; Kent; Middlesex; Peterborough; Welland; Wentworth; York. AMNH, CNC, ROM. Subspecies *I. c. enigmata* (Burns and McDunnough).
- I. evagaria* (Hulst). 4: Frontenac; Lambton; Leeds. CNC, ROM.
- I. exaspicata* (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Kenora; Rainy River; Thunder Bay. 4: Carleton; Frontenac; Grey; Leeds; Nipissing; Norfolk; Simcoe. AMNH, CNC, ROM.
- I. flavicaria* (Packard). 4: Huron (Grand Bend). ROM. Only known locality.
- I. loricaria* (Eversmann). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Hastings; Middlesex; Victoria. AMNH, CNC. Subspecies *I. l. julia* (Hulst).
- I. occiduaria* (Packard). 4: Huron (Grand Bend); Lambton. CNC. Only known localities.
- I. pustularia* (Guenée). 3: Rainy River; Sudbury; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Kent; Leeds;

Lennox and Addington; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.

I. ribearia (Fitch). Currant spanworm/arpenteuse du groseillier. 3: Rainy River; Sudbury. 4: Carleton; Durham; Essex; Frontenac; Hastings; Lanark; Leeds; Lennox and Addington; Middlesex; Parry Sound; Peterborough; York. AMNH, CNC, ROM.

I. subcessaria (Walker). 4: Carleton; Durham; Essex; Frontenac; Hastings; Leeds; Lennox and Addington; Middlesex; Muskoka; Peterborough. AMNH, CNC, ROM.

I. sulphurea (Packard). 2: Cochrane; Thunder Bay. 3: Rainy River. 4: Frontenac; Huron; Leeds. AMNH, CNC, ROM.

Genus *Epelis* Hulst [= *Isturgia* auct.]

E. truncataria (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Timiskaming. 4: Carleton; Frontenac; Middlesex; Waterloo; Wellington. AMNH, CNC, ROM.

Genus *Mellilla* Grote

M. xanthometata (Walker). 4: Essex (Peelee Island). CNC. Only known locality.

Genus *Semiothisa* Hübner

S. aemulataria (Walker). 4: Frontenac; Hastings; Leeds; Waterloo; York. CNC, ROM.

S. banksianae Ferguson. 4: Leeds. ROM.

S. bicolorata (Fabricius). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Frontenac; Grey; Leeds; Muskoka; Nipissing; Oxford; Renfrew. AMNH, CNC, ROM.

S. bisignata (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Frontenac; Haliburton; Hastings; Leeds; Norfolk; Peterborough; Renfrew; Simcoe; Waterloo; Welland; York. AMNH, CNC, ROM.

S. denticulata Grote. 2: Cochrane; Thunder Bay. 3: Thunder Bay. 4: Huron. AMNH, CNC, ROM.

S. fissinotata (Walker). 3: Algoma; Sudbury. 4: Bruce; Carleton; Dufferin; Grenville; Grey; Hastings; Leeds; Middlesex; Muskoka; Norfolk; Oxford; Peel; Simcoe; Welland; Wellington; York. AMNH, CNC, ROM.

S. gnophosaria (Guenée). 4: Leeds; Middlesex; Welland; York. AMNH, CNC.

S. hebetata (Hulst). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. AMNH, CNC, ROM.

- S. mellistrigata* (Grote). 3: Algoma; Rainy River; Sudbury. 4: Carleton; Frontenac; Hastings; Kent; Leeds; Middlesex; Muskoka; Welland; Wentworth; York. AMNH, CNC, ROM
- S. minorata* (Packard). 3: Algoma; Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Hastings; Leeds; Nipissing; Norfolk; Parry Sound; Renfrew; Simcoe; Welland; Wentworth; York. AMNH, CNC, ROM.
- S. neptaria* (Guenée). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Leeds. AMNH, CNC.
- S. ocellinata* (Guenée). 4: Carleton; Durham; Essex; Frontenac; Hastings; Leeds; Norfolk; Northumberland; Prince Edward; Welland; Wentworth; York. AMNH, CNC.
- S. orillata* (Walker). 2: Cochrane. 3: Algoma; Sudbury. 4: Carleton; Frontenac; Hastings; Leeds; Nipissing; Norfolk; York. AMNH, CNC, ROM.
- S. oweni* (Swett). Owen's green looper/arpenteuse sombre. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Ontario. AMNH, CNC, ROM.
- S. pinistrobata* Ferguson. 3: Algoma; Rainy River; Sudbury. 4: Carleton; Hastings; Renfrew. CNC.
- S. sexmaculata* (Packard). Green larch looper/arpenteuse verte du mélèze. 2: Cochrane; Thunder Bay. 3: Algoma; Cochrane; Sudbury; Thunder Bay. 4: Brant; Carleton; Frontenac; Haldimand; Hastings; Leeds; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Stormont; York. AMNH, CNC, ROM.
- S. signaria* (Hübner). 2: Cochrane. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Haldimand; Hastings; Lanark; Leeds; Niagara; Nipissing; Parry Sound; Renfrew; Simcoe. CNC. Subspecies *S. s. dispunctata* (Walker).
- S. submarmorata* (Walker). 2: Thunder Bay. 3: Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Frontenac; Haliburton; Hastings; Lanark; Leeds; Nipissing; Simcoe; Welland; York. CNC, ROM.
- S. transitoria* (Walker). 3: Sudbury. 4: Durham; Leeds; Nipissing. AMNH, ROM.
- S. ulsterata* (Pearsall). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Middlesex; Norfolk; Welland; York. AMNH, CNC, ROM.

Genus *Enconista* Lederer

- E. dislocaria* (Packard). 4: Essex (Peelee Island. Point Pelee). CNC, ROM. Only known localities. Subspecies *E. d. malefactoria* (Barnes and McDunnough).

Tribe Boarmiini

Genus *Orthofidonia* Packard

- O. flavivenata* (Hulst). 2: Cochrane. 4: Frontenac; Leeds; Middlesex; Muskoka. CNC, ROM.
- O. tinctaria* (Walker). 3: Sudbury; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Muskoka; Parry Sound. CNC, ROM.

Genus *Hesperumia* Packard

- H. sulphuraria* Packard. 3: Sudbury. 4: Carleton; Grenville; Hastings; Leeds; Parry Sound; Peterborough. AMNH, CNC, ROM.

Genus *Ematurga* Lederer

- E. amitaria* (Guenée). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haliburton; Muskoka; Nipissing; Norfolk; Parry Sound. AMNH, CNC, ROM.

Genus *Stenoporpia* McDunnough

- S. polygrammaria* (Packard). 3: Algoma; Thunder Bay. 4: Frontenac; Kent; Leeds; Muskoka. CNC, ROM.

Genus *Aethalura* McDunnough

- A. intertexta* (Walker). 2: Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Ontario; Russell; Simcoe; Victoria; York. AMNH, CNC, ROM.

Genus *Anacamptodes* McDunnough

- A. ephyraria* (Walker). 2: Cochrane; Thunder Bay. 3: Sudbury; Timiskaming. 4: Carleton; Frontenac; Haliburton; Hastings; Huron; Leeds; Norfolk; Northumberland. AMNH, CNC, ROM.
- A. humaria* (Guenée). 2: Cochrane. 3: Thunder Bay. 4: Carleton; Huron. AMNH, CNC.
- A. vellivolata* (Hulst). 3: Rainy River; Sudbury. 4: Carleton; Frontenac; Grey; Huron; Leeds; Nipissing; Norfolk; Ontario; Victoria. AMNH, CNC, ROM.

Genus *Iridopsis* Warren

- I. larvaria* (Guenée). 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Haliburton; Hastings; Muskoka; Leeds; Parry Sound; Peterborough; Victoria; York. AMNH, CNC, ROM.

Genus *Anavitrinelia* McDunnough

- A. pampinaria* (Guenée). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Hastings; Huron; Lanark; Leeds; Nipissing; Norfolk; Renfrew; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.

Genus *Cleora* Curtis

- C. projecta* (Walker). 3: Sudbury; Timiskaming. 4: Carleton; Frontenac. AMNH, CNC.
C. sublunaria (Guenée). 4: Frontenac (Perth Road). CNC. Only known locality.

Genus *Ectropis* Hübner

Saddleback looper/arpenteuse bossue de la pruche.

- E. crepuscularia* (Denis and Schiffermüller). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Grey; Haldimand; Haliburton; Halton; Hastings; Kent; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Ontario; Peterborough; Renfrew; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Genus *Hypagyrtis* Hübner

See Rindge (1985).

- H. piniata* (Packard). Pine looper/arpenteuse du pin. 3: Algoma; Rainy River; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Halton; Hastings; Lanark; Leeds; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Renfrew; Wentworth; York. AMNH, CNC, ROM.
H. unipunctata Haworth. 2: Kenora. 3: Sudbury. 4: Frontenac; Grey; Hastings; Leeds; Muskoka; Nipissing; Norfolk; Victoria; York. AMNH, CNC, ROM.

Genus *Protoarmoia* McDunnough

P. porcelaria (Guenée). Dotted line looper/arpenteuse dodue. 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Thunder Bay. 4: Bruce; Carleton; Frontenac; Hastings; Leeds; Nipissing; Ontario; Simcoe. AMNH, CNC, ROM.

Genus *Epimecis* Hübner

E. hortaria (Fabricius). 4: Kent; Lincoln; Norfolk; Welland. CNC, ROM.

Tribe Melanolophiini

See McGuffin (1977).

Genus *Melanolophia* Hulst

M. canadaria (Guenée). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury. 4: Carleton; Dufferin; Durham; Frontenac; Haliburton; Hastings; Leeds; Middlesex; Norfolk; Ontario; Renfrew; Simcoe; Victoria; York. AMNH, CNC, ROM.

M. signataria (Walker). 3: Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haliburton; Hastings; Leeds; Renfrew. AMNH, CNC, ROM.

Genus *Eufidonia* Packard

E. convergaria (Walker). 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haldimand; Renfrew; Victoria. CNC.

E. discospilata (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Haliburton; Leeds; Parry Sound. AMNH, CNC, ROM.

E. notataria (Walker). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Dufferin; Frontenac; Haliburton; Hastings; Huron; Leeds; Nipissing; Norfolk; Simcoe; Victoria; Wellington; York. AMNH, CNC, ROM.

Tribe Bistonini

Genus *Biston* Leach

The generic name is of masculine gender.

B. betularius (Linné). Pepper-and-salt moth/arpenteuse du gadelier. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings;

Kent; Leeds; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Peterborough; Simcoe; Welland; Wellington; Wentworth; York. AMNH, CNC, ROM. Subspecies *B. b. cognatarius* (Guenée).

Genus *Lycia* Hübner

L. ursaria (Walker). 2: Cochrane. 3: Algoma; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Hastings; Lanark; Leeds; Lincoln; Middlesex; Muskoka; Victoria; Wentworth; York. AMNH, CNC, ROM.

Genus *Phigalia* Duponchel

P. strigataria (Minot). 4: Frontenac; Hastings; Leeds. CNC, ROM.

P. titea (Cramer). 4: Carleton; Frontenac; Haldimand; Hastings; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Ontario; Wentworth; York. AMNH, CNC, ROM.

Genus *Paleacrita* Riley

P. merricata Dyar. 4: Muskoka (Severn Bridge); York (Toronto). CNC, ROM.

P. vernata (Peck). Spring cankerworm/ arpentuse du printemps. 4: Carleton; Durham; Frontenac; Haldimand; Halton; Hastings; Leeds; Middlesex; Muskoka; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Genus *Erannis* Hübner

E. tiliaria (Harris). Linden looper/ arpentuse du tilleul. 2: Cochrane; Thunder Bay. 4: Brant; Carleton; Durham; Essex; Frontenac; Grenville; Grey; Haldimand; Hastings; Leeds; Lincoln; Middlesex; Muskoka; Northumberland; Peterborough; Prince Edward; Waterloo; Wellington; Wentworth; York. AMNH, CNC, ROM.

Tribe Baptini

Genus *Lomographa* Hübner

L. glomeraria (Grote). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Halton; Muskoka. AMNH, CNC, ROM.

L. semiclarata (Walker). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Haliburton; Halton; Hastings; Leeds; Muskoka; Parry Sound; Simcoe. AMNH, CNC, ROM.

L. vestaliata (Guenée). 2: Kenora, Patricia Portion. 3: Rainy River; Thunder Bay. 4: Brant; Carleton; Durham; Frontenac; Haldimand; Haliburton; Halton; Hastings; Kent; Leeds; Middlesex; Norfolk; Victoria; York. AMNH, CNC, ROM.

Tribe Caberini

Genus *Cabera* Treitschke

- C. borealis* (Hulst). 2: Thunder Bay (Longlac). AMNH, ROM.
- C. erythemaria* Guenée. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Peel; Victoria; Wentworth; York. AMNH, CNC, ROM.
- C. variolaria* Guenée. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Essex; Frontenac; Haldimand; Hastings; Kent; Lanark; Leeds; Norfolk; Victoria; Waterloo; Wentworth. AMNH, CNC, ROM.

Genus *Apodrepanulatrix* Rindge

- A. liberaria* (Walker). 4: Carleton; Leeds; Norfolk; Peterborough; Victoria; York. CNC, ROM.

Genus *Erastria* Hübner [= *Syrphodia* Hübner]

- E. coloraria* (Fabricius). 4: Hastings; Huron; Northumberland; York. CNC, ROM.
- E. cruentaria* (Hübner). 4: Huron; Lambton. CNC.

Tribe Angeronini

Genus *Aspitates* Treitschke

- A. orciferarius* (Walker). 1: Kenora, Patricia Portion (Fort Severn, 20 July 1940). ROM.

Genus *Lytrosis* Hulst

- L. unitaria* (Herrich-Schäffer). 4: Carleton; Frontenac; Haldimand; Hastings; Leeds; Lincoln; Muskoka; Norfolk. AMNH, CNC, ROM.

Genus *Euchlaena* Hübner

- E. effecta* (Walker). 3: Sudbury. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Welland; Wentworth; York. AMNH, CNC, ROM.
- E. irraria* (Barnes and McDunnough). 2: Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Hastings; Kent; Leeds; Lincoln; Muskoka; Norfolk; Peterborough; Victoria. AMNH, CNC, ROM.
- E. johnsonaria* (Fitch). 2: Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Frontenac; Grenville; Haldimand; Hastings; Huron; Leeds; Muskoka; Ontario; Parry Sound; Peterborough; Renfrew; Simcoe; Welland; York. AMNH, CNC, ROM.
- E. madusaria* (Walker). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton. AMNH, CNC.
- E. marginaria* (Minot). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Essex; Frontenac; Haldimand; Leeds; Middlesex; Muskoka; Nipissing; Ontario; Parry Sound; York. AMNH, CNC, ROM.
- E. muzaria* (Walker). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Haldimand; Muskoka; Nipissing; Peel; Simcoe; Wentworth. AMNH, ROM.
- E. obtusaria* (Hübner). 2: Cochrane; Thunder Bay. 3: Thunder Bay. 4: Carleton; Essex; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Muskoka; Peel; Renfrew; Welland; Wentworth; York. AMNH, CNC, ROM.
- E. serrata* (Drury). 3: Sudbury. 4: Carleton; Essex; Frontenac; Haldimand; Hastings; Huron; Leeds; Lennox and Addington; Middlesex; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.
- E. tigrinaria* (Guenée). 3: Thunder Bay. 4: Carleton; Halton; Nipissing; Renfrew; York. AMNH, CNC, ROM.

Genus *Xanthotype* Warren

- X. sospeta* (Drury). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Leeds; Middlesex; Nipissing; Norfolk; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Welland; Wentworth; York. AMNH, CNC, ROM.
- X. urticaria* Swett. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Halton; Hastings; Kent; Leeds; Lennox and Addington; Muskoka; Nipissing; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Tribe Azelinini

Genus *Pero* Herrich-Schäffer

- P. honestaria* (Walker). 2: Cochrane. 3: Thunder Bay. 4: Bruce; Carleton; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Norfolk; Peel; Simcoe; Welland; Wentworth; York. AMNH, CNC, ROM.
- P. hubneraria* (Guenée) [= *marmoratus* (Grossbeck)]. 4: Frontenac (Perth Road, P. S. Ward, only two specimens). ROM.
- P. morrisonaria* (Hy. Edwards). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Frontenac; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Simcoe; Wellington; Wentworth; York. AMNH, CNC, ROM.

Tribe Nacophorini

Genus *Nacophora* Hulst

- N. mexicanaria* (Grote). 4: Nipissing (Algonquin Provincial Park, 12 June 1963, D. Barr, Department of Entomology ROM, caught in a light trap). ROM.
- N. quernaria* (J. E. Smith). 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Halton; Huron; Kent; Lanark; Leeds; Lincoln; Muskoka; Wentworth; York. AMNH, CNC, ROM.

Tribe Campaeini

Genus *Campaea* Lamarck

- C. perlata* (Guenée). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Haliburton; Hastings; Huron; Kent; Leeds; Middlesex; Muskoka; Norfolk; Peterborough; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Tribe Ennomini

Genus *Ennomos* Treitschke [= *Deuteronomos* Prout]

The generic name can be of both masculine and feminine genders; however, it was used as feminine in the type species.

E. magnaria Guenée. Maple spanworm/arpenteuse de l'érable. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Lanark; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Peterborough; Simcoe; Victoria; Welland; Wellington; Wentworth; York. AMNH, CNC, ROM.

E. subsignaria (Hübner). Elm spanworm/arpenteuse de l'orme. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Halton; Hastings; Huron; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Peterborough; Simcoe; Waterloo; Welland; Wellington; Wentworth; York. AMNH, CNC, ROM.

Tribe Epirranthini

Genus *Epirranthis* Hübner

E. substriataria (Hulst). 2: Cochrane. 3: Sudbury; Thunder Bay. 4: Carleton; Muskoka; Northumberland; York. AMNH, CNC, ROM.

Tribe Lithinini

Genus *Petrophora* Hübner

P. subaequaria Walker. 2: Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Victoria; Wentworth; York. AMNH, CNC, ROM.

Genus *Tacparia* Walker

T. atropunctata (Packard). 3: Sudbury. 4: Frontenac; Hastings; Leeds. AMNH, CNC, ROM.

T. detersata (Guenée). 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Muskoka; Nipissing; Peterborough; Victoria; York. AMNH, CNC, ROM.

Genus *Homochlodes* Hulst

H. fritillaria (Guenée). 2: Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haldimand; Hastings; Kent; Leeds; Lincoln; Muskoka; Norfolk; Parry Sound; Simcoe; York. AMNH, CNC, ROM.

Genus *Gueneria* Packard

G. similaria (Walker). 4: Carleton; Frontenac; Leeds; Middlesex; Norfolk. AMNH, CNC.

Tribe Anagogini

Genus *Selenia* Hübner

S. alciphearia Walker. 2: Cochrane. 3: Sudbury; Thunder Bay. 4: Carleton; Frontenac. AMNH, CNC.

S. kentaria (Grote and Robinson). 3: Rainy River; Thunder Bay. 4: Carleton; Hastings. CNC.

Genus *Metanema* Guenée

M. determinata Walker. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Haldimand; Hastings; Lambton; Leeds; Middlesex; Muskoka; Nipissing; Welland; York. AMNH, CNC, ROM.

M. inatomaria Guenée. 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Lincoln; Muskoka; Nipissing; Norfolk; Peterborough; Simcoe; Victoria; Welland; Wellington; York. AMNH, CNC, ROM.

Genus *Metarranthis* Warren

See Rupert (1943).

M. amyrisaria (Walker). 4: Carleton (Constance Bay. South March). CNC.

M. apiciaria (Packard). 4: York (Toronto only, 1906 and 1911). ROM.

M. duaria (Guenée). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Frontenac; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Parry Sound; Victoria; Welland; York. AMNH, CNC, ROM.

M. hypochraria (Herrich-Schäffer). 3: Sudbury. 4: Carleton; Durham; Frontenac; Hastings; Leeds; Middlesex; Norfolk; Northumberland; Ontario; Peel; Simcoe; Victoria; Welland; York. AMNH, CNC, ROM.

M. indeclinata (Walker). 4: Durham; Frontenac; Leeds. AMNH, CNC, ROM.

M. obfirmaria (Hübner). 3: Sudbury. 4: Carleton; Frontenac; Leeds. AMNH, CNC, ROM.

M. refractaria (Guenée). AMNH, not yet segregated in collection.

M. warnerae (Harvey). 2: Cochrane. 3: Sudbury. 4: Carleton; Frontenac; Leeds; Nipissing; Wentworth. AMNH, CNC, ROM.

Genus *Cepphis* Hübner [= *Priocycla* Guenée]

- C. armataria* (Herrich-Schäffer).** 3: Rainy River. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Ontario; Peterborough; York. AMNH, CNC, ROM.
- C. decoloraria* (Hulst).** 4: Hastings; Leeds; Muskoka; Nipissing. CNC, ROM.

Genus *Anagoga* Hübner

- A. occiduaria* (Walker).** 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Halton; Leeds; Muskoka; Renfrew. AMNH, CNC, ROM.

Genus *Probole* Herrich-Schäffer

- P. alienaria* Herrich-Schäffer.** 2: Cochrane; Kenora. 3: Thunder Bay. 4: Dufferin; Durham; Frontenac; Leeds; Nipissing; Norfolk; Peel; York. CNC, ROM.
- P. amicara* (Herrich-Schäffer).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Essex; Frontenac; Haldimand; Haliburton; Hastings; Leeds; Lincoln; Muskoka; Nipissing; Norfolk; Parry Sound. AMNH, CNC, ROM.
- P. nyssaria* (Guenée).** 3: Thunder Bay.

Genus *Plagodis* Hübner

- P. alcoolaria* (Guenée).** 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Haldimand; Kent; Lambton; Leeds; Muskoka; Nipissing; Peterborough; Simcoe; York. AMNH, CNC, ROM.
- P. fervidaria* (Herrich-Schäffer).** 4: Peterborough; York (Toronto). ROM.
- P. kuetzingi* (Grote).** 3: Thunder Bay. 4: Carleton; Frontenac; Haliburton; Hastings; Leeds; Middlesex; Peterborough; Simcoe. AMNH, CNC, ROM.
- P. phlogosaria* (Guenée).** 2: Cochrane; Thunder Bay. 3: Algoma; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Halton; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Victoria; Wentworth; York. AMNH, CNC, ROM.
- P. serinaria* Herrich-Schäffer.** 3: Algoma. 4: Carleton; Dufferin; Frontenac; Haldimand; Leeds; Muskoka; Nipissing; Peterborough; Simcoe; York. AMNH, CNC, ROM.

Tribe Ourapterygini

Genus *Caripeta* Walker

- C. augustiorata* Walker. Brown pine looper/arpenteuse brune du pin. 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Leeds; Muskoka; Nipissing; Ontario; Peel; Peterborough; Simcoe; Waterloo; Wentworth; York. AMNH, CNC, ROM.
- C. divisata* Walker. Gray spruce looper/arpenteuse grise de l'épinette. 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Huron; Leeds; Muskoka; Nipissing; Ontario; Peterborough; Simcoe; Waterloo; York. AMNH, CNC, ROM.
- C. piniata* (Packard). 3: Algoma; Thunder Bay; Sudbury. 4: Durham; Frontenac; Haldimand; Hastings; Muskoka; Nipissing; Norfolk; Leeds; Peel; Peterborough; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Genus *Besma* Capps

- B. endropiaria* (Grote and Robinson). 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Frontenac; Grenville; Huron; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peterborough; Wentworth. AMNH, ROM.
- B. quercivoraria* (Guenée). 2: Cochrane; Thunder Bay. 3: Thunder Bay. 4: Brant; Frontenac; Haliburton; Hastings; Leeds; Norfolk; Renfrew; Wentworth. AMNH, CNC, ROM.

Genus *Lambdina* Capps

- L. fervidaria* (Hübner). 3: Algoma. 4: Carleton; Parry Sound. CNC.
Subspecies *L. f. athasaria* (Walker).
- L. fiscellaria* (Guenée). 2: Cochrane; Thunder Bay. 3: Algoma. 4: Bruce; Carleton; Dufferin; Durham; Frontenac; Glengarry; Halton; Hastings; Lanark; Leeds; Muskoka; Nipissing; Norfolk; Ontario; Peel; Peterborough; Prescott; Simcoe; Timiskaming; York. AMNH, CNC, ROM.

Tribe Cingiliini

Genus *Cingilia* Walker

- C. catenaria* (Drury). Chain-spotted geometer/arpenteuse catenaire. 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton;

Frontenac; Haldimand; Leeds; Middlesex; Muskoka; Nipissing; Parry Sound; Peterborough; Wentworth; York. AMNH, CNC, ROM.

Genus *Nepytia* Hulst

- N. canosaria* (Walker). False hemlock looper/ fausse arpenteuse de la pruche. 2: Cochrane. 3: Algoma; Sudbury. 4: Bruce; Carleton; Durham; Frontenac; Grey; Haliburton; Hastings; Leeds; Middlesex; Nipissing; Norfolk; Oxford; Parry Sound; Peel; Peterborough; Wentworth; York. AMNH, CNC, ROM.
- N. pellucidaria* (Packard). 4: Peterborough (Mt. Irwin, Crystal Lake); Renfrew (LaPasse. Ompah); York (Toronto). CNC, ROM.

Genus *Sicya* (Guenée)

- S. macularia* (Harris). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Halton; Hastings; Lambton; Leeds; Muskoka; Nipissing; Norfolk; Peterborough; Renfrew; Simcoe; Welland; York. AMNH, CNC, ROM.

Genus *Eusarca* Hübner [= *Apicia* Guenée]

- E. confusaria* Hübner. 3: Rainy River. 4: Carleton; Dufferin; Durham; Frontenac; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Peterborough; Simcoe; Wentworth; York. AMNH, CNC, ROM.

Genus *Tetracis* Guenée

- T. cachexiata* Guenée. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lanark; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Simcoe; Welland; Wellington; Wentworth; York. AMNH, CNC, ROM.
- T. crocallata* Guenée. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Norfolk; Peel; Victoria; York. AMNH, CNC, ROM.

Genus *Eugonobapta* Warren

- E. nivosaria* (Guenée). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Norfolk; Welland; Wentworth; York. AMNH, CNC, ROM.

Genus *Eutrapela* Hübner [= *Abbottana* Hulst]

***E. clemataria* (J. E. Smith).** 3: Sudbury. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Halton; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Ontario; Wentworth; York. AMNH, CNC, ROM.

Genus *Prochoerodes* Grote

***P. transversata* (Drury).** Large maple spanworm/arpenteuse transverse de l'érable. 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Kenora; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haliburton; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Simcoe; Timiskaming; Victoria; Waterloo; Welland; Wentworth; York. AMNH, CNC, ROM.

Genus *Antepione* Packard

***A. thisoaria* (Guenée).** 3: Rainy River; Sudbury. 4: Frontenac (Perth Road). CNC, ROM.

Genus *Nematocampa* Guenée

***N. limbata* (Haworth) [= *filamentaria* Guenée].** Horned spanworm/nématocampe. 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Hastings; Leeds; Middlesex; Norfolk; Peel; Wellington; Wentworth; York. AMNH, CNC, ROM.

Family Epiplemidæ

Genus *Callizzia* Packard

***C. amorata* Packard.** 2: Cochrane. 3: Timiskaming. 4: Carleton; Durham; Frontenac; Hastings; Leeds; Peterborough; Simcoe. 2 May–1 August. AMNH, Beth, CNC, ROM. Uncommon and very local.

Genus *Calledapteryx* Grote

***C. dryopterata* Grote.** 4: Carleton (Ottawa); Huron (Grand Bend); Wentworth (Hamilton). CNC, UG. Collected only in a few isolated localities.

Superfamily Mimallonoidea
Family Mimallonidae [= Perophoridae = Lacosomidae]

Genus *Lacosoma* Grote

The generic name is of neuter gender.

***L. chiridotum* Grote.** 4: Kent (Rondeau Provincial Park); Norfolk (St. Williams). 28 June–10 July. CNC, ROM. Known only from the two localities.

Genus *Cicinnus* Blanchard

***C. melsheimeri* (Harris).** 4: Haldimand; Kent; Middlesex; Norfolk; Welland; Wentworth. 6 June–2 July. CNC, ROM, UG. Known only from the north shore of Lake Erie and a few localities more inland.

Superfamily Bombycoidea
Family Apatelodidae
Subfamily Apatelodinae

Genus *Apatelodes* Packard

***A. torrefacta* (J. E. Smith).** 4: Essex (Harrow); Haldimand (Dunnville); Huron (Grand Bend); Kent (Rondeau Provincial Park, 25 July); Lincoln (Queenston); Middlesex (London); Norfolk (Normandale); Welland (Port Colborne, 7 May). CNC, ROM, UWO. Known only from the southernmost part of the region.

Genus *Olceclostera* Butler

***O. angelica* (Grote).** 4: Frontenac (Glenburnie. Perth Road); Haldimand (Dunnville); Hastings (Belleville. Trenton); Huron (Grand Bend); Kent (Rondeau Provincial Park); Leeds (Chaffey's Locks, 8 June–16 July); Lincoln (Queenston); Norfolk (Normandale); Simcoe. Peak in early July. CNC, QU, ROM. Reared easily on ash. Known from the area immediately north of Lake Erie and Lake Ontario.

Family Lasiocampidae
Subfamily Macromphaliinae

Genus *Tolype* Hübner

- T. laricis* (Fitch).** Larch lappet moth/lasiocampe du mélèze. 2: Cochrane. 3: Algoma; Sudbury; Thunder Bay. 4: Bruce; Carleton; Dufferin; Durham; Frontenac; Haldimand; Haliburton; Hastings; Huron; Leeds; Lennox and Addington; Middlesex; Muskoka; Nipissing; Ontario; Parry Sound; Peterborough; Renfrew; Simcoe; Stormont; Welland; Wellington; Wentworth; York. 20 July–17 September. Peak in August. AMNH, CNC, LEM, QU, ROM, UG, UWO. Reared on white pine in Chaffey's Locks. The larvae did not feed on fir but did well on spruce and larch. One larva was collected near Chaffey's Locks on hemlock.
- T. notialis* Franclemont.** 4: Kent (Rondeau Provincial Park, 17 August 1959). ROM. Known only from the southwesternmost part of the region. Two male specimens were found in the Park Museum at Rondeau Provincial Park in 1965. One of these specimens is now in the ROM and was dissected; the genitalia correspond to Franclemont's (1973) drawing.
- T. velleda* (Stoll).** Velleda lappet moth/lasiocampe à épaulettes. 4: Brant; Carleton; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Parry Sound; Peel; Peterborough; Welland; Wellington; Wentworth; York. 26 August–22 October. AMNH, Beth, CNC, LEM, McM, ROM, UWO. Larvae reared without difficulty on crabapple in Chaffey's Locks.

Subfamily Gastropachinae

Genus *Heteropacha* Harvey

- H. rileyana* Harvey.** 4: Kent (Chatham, 25 May 1931). CNC. Known only from a single specimen caught in a light trap at the Entomology Laboratory in Chatham.

Genus *Phyllodesma* Hübner

The generic name is of neuter gender.

- P. americanum* (Harris).** Lappet moth/papillon à épaulettes. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Ontario; Peterborough; Simcoe; Victoria;

Welland; Wellington; York. 19 April–17 August. AMNH, CNC, LEM, QU, ROM, UG, UWO. Specimens from late July and later in southern Ontario belong to a partial second generation and are redder in colour than the spring specimens. In more northern regions (i.e., Algoma, Cochrane, Rainy River, Sudbury, Thunder Bay) there is only one generation.

Larvae in the south prefer birch, oak, ash, apple, etc.; larvae in the north prefer aspen, poplar, and willow. Larvae from eggs produced by northern females refused foodplants other than those in the poplar complex. Larvae from southern females, however, may take *Populus* sp., but do not grow as fast as when on their usual foodplants.

Subfamily Lasiocampinae **Tribe Malacosomatini**

Genus *Malacosoma* Hübner

The generic name is of neuter gender.

- M. americanum* (Fabricius).** Eastern tent caterpillar/livrée d'Amérique. 3: Algoma; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Dufferin; Durham; Elgin; Essex*; Frontenac; Grenville; Haldimand; Haliburton; Hastings; Kent; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Perth; Peterborough; Renfrew; Simcoe; Victoria; Welland; Wellington; Wentworth; York. 7 June–23 August. AMNH, Beth, CNC, McM, ROM, UG, UWO.
- M. californicum* (Packard).** Western tent caterpillar/livrée de l'Ouest. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Muskoka; Nipissing. 8 July–3 August. CNC, LEM, ROM. Known from the region west of Lake Superior and east throughout the boreal forest, is now also in the northern part of the region east of Lake Superior and extending its range.
- M. disstria* Hübner.** Forest tent caterpillar/livrée des forêts. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Durham; Frontenac; Grenville; Haldimand; Haliburton; Kent; Lambton; Leeds; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peel; Peterborough; Simcoe; Wellington; York. 16 June–28 August. AMNH, Beth, CNC, McM, QU, ROM, UWO.

Family Saturniidae
See Ferguson (1971, 1972).

Subfamily Citheroniinae

Genus *Eacles* Hübner

***E. imperialis* (Drury).** Imperial moth/papillon impérial. This species is represented in Ontario by two populations considered by Ferguson (1971) to be two subspecies: *E. i. imperialis* proper and *E. i. pini* Michener. As there is a possibility that these two populations might prove to be two different species, we refer to them here under the subspecific names, especially as the pertinent locality records are readily available. In Ontario we have also a sizeable area in the Parry Sound District where a marked melanism regularly occurs in the population. Ferguson (1971, plate 1, no. 18; plate 2, no. 1) illustrated both melanic forms. He erred, however, when he said that there exist only three melanic specimens; there is an entire series of both forms in the ROM collection and new examples are found every year.

"*E. imperialis imperialis*." 4: Elgin; Essex; Haldimand; Lincoln; Middlesex; Norfolk; Peel; Waterloo; Wellington; Wentworth; York. One specimen cannot be easily assigned to either population—4: Durham (Port Hope).

"*E. imperialis pini*." 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Dufferin; Durham; Frontenac; Haliburton; Hastings; Leeds; Lennox and Addington; Muskoka; Northumberland; Ontario; Parry Sound; Peterborough; Renfrew; Simcoe; Victoria; York. 7 June–23 August. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. The exclusive larval foodplant is pine; larvae refused to feed on other trees offered them (W. Plath, Sr., personal comment) in Haldimand (Dunnville). For the foodplant problem see Ferguson (1971:26).

Genus *Sphingicampa* Walsh

***S. bicolor* (Harris).** 4: Brant (Brantford); Essex; Haldimand (Dunnville); Lincoln (St. Catharines); Peel (Bramalea); York (Woodbridge). May–July. ROM. Known only from the few localities above—see Riotte (1964)—and from Huron (Zurich. Grand Bend). See TEA Occasional Publications 1984, 15:56.

Genus *Dryocampa* Harris

***D. rubicunda* (Fabricius).** Green-striped mapleworm/anisote de l'érable. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury. 4: Brant; Carleton; Durham; Essex; Frontenac; Haldimand; Halton; Hastings; Huron;

Kent; Leeds; Lincoln; Middlesex; Nipissing; Norfolk; Northumberland; Perth; Renfrew; Simcomont; Welland; Wellington; Wentworth; York. 19 May–27 July. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. A partial second generation may occur in the field, e.g., Carleton (Ottawa, 5 September 1918, single specimen). Reared larvae regularly yield a partial second generation.

Specimens taken north of Muskoka usually show no yellow-cream colour and only a little pink on the wings; often they are pure white. Specimens from the north shores of Lakes Erie and Ontario may show intensive colouring, appearing along with more or less lightly coloured ones from the same area. In rearing experiments at Chaffeys Locks, we found that the offspring show the same colour pattern as the mother. The exclusive foodplant is maple. Oak leaves were offered to larvae in the laboratory, but they only nibbled on the leaves and refused them completely after a few days.

Genus *Anisota* Hübner

See Riotte and Peigler (1981).

- A. finlaysoni* **Riotte**. 4: Durham; Frontenac; Haldimand; Halton; Hastings; Lennox and Addington; Norfolk; Prince Edward; Victoria; Welland; Wentworth. Hastings (Shannonville, type locality). AMNH, CM, CNC, QU, ROM, UWO.
- A. senatoria* (**J. E. Smith**). Orange-striped oakworm/anisote sénatoriale. 4: Brant; Essex; Lambton; Middlesex; Oxford; Waterloo; Wellington; Wentworth. 16 May–2 July. CNC, ROM, UWO.
- A. stigma* (**Fabricius**). 4: Brant; Lincoln; Middlesex; Norfolk; Welland; Wentworth. 27 June–5 July. Beth, CNC, ROM, UG. Known only from a few localities on the north shore of Lake Erie. A FIDS record, “near North Bay in northern Ontario,” was based on misidentified larvae, which are regarded as belonging to the next species. The identification of the original report of the Forest Insect Survey, Canada, was changed to “*Anisota* sp.” (P. D. Syme, Canadian Forestry Service, Insect Pathology Research Institute, Sault Ste. Marie, in litt.). See also Riotte and Peigler (1981:171).
- A. virginiensis* (**Drury**). Pink-striped oakworm/anisote rose du chêne. 3: Algoma; Sudbury; Timiskaming. 4: Carleton; Elgin; Essex; Frontenac; Huron; Kent; Leeds; Lincoln; Manitoulin; Muskoka; Nipissing; Norfolk; Northumberland; Peel; Renfrew; York. 5 May–2 August; peak mid May–mid July. Beth, CNC, McM, QU, ROM, UWO.

Subfamily Hemileucinae
Tribe Hemileucini

Genus *Hemileuca* Walker

H. sp. 4: Carleton (Marlborough Twp.); Renfrew (McNab Twp., September). CNC. The species was discovered in autumn of 1977; larval foodplant is *Menyanthes trifoliata* Linné (buckbean). For details see TEA Occasional Publications 1980, 11:9; 1981, 12:36; 1982, 13:18.

Genus *Automeris* Hübner

***A. io* (Fabricius).** Io moth/automeris io. 3: Algoma; Rainy River; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grenville; Haldimand; Haliburton; Halton; Hastings; Kent; Lambton; Lanark; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Peel; Peterborough; Renfrew; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. 22 May–18 August; peak June–mid July. AMNH, Beth, CNC, McM, ROM, UG, UWO. A partial second generation occurs as late as September and October in the field (e.g., Rondeau Provincial Park, Toronto, and Ottawa), and from reared larvae. The most frequented foodplants are basswood and birch, although other trees and shrubs are recorded.

Subfamily Saturniinae
Tribe Saturniini

Genus *Antheraea* Hübner

***A. polyphemus* (Cramer).** Polyphemus moth/polyphème. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. 13 May–8 August; peak June–mid July. AMNH, Beth, CNC, McM, ROM, UG, UWO. Known throughout the regions extending to 53° north latitude. A partial second generation occurs in the field from early September, e.g., Leeds (Chaffey's Locks), and from reared larvae. The preferred foodplant is birch, especially in more northern latitudes. There are records of larvae feeding on other trees such as maple and basswood.

Genus *Actias* Leach

A. luna (Linné). Luna moth/papillon lune. 2: Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Haliburton; Hastings; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound*; Peel; Renfrew; Simcoe; Waterloo; Welland; Wellington; Wentworth; York. 12 May–31 July. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. Known throughout the regions extending to 53° north latitude. Reared larvae regularly yield a partial second generation, but there is no record of a partial second generation in the field. The main foodplant is birch, and basswood is occasionally utilized.

Tribe Attacini

Genus *Samia* Hübner

S. cynthia (Drury). 4: Middlesex (London, June 1903, one male); York (Toronto, 16 June 1936, R. B. Johnston, one female emerged from a pupa collected from among several on an ash tree at 112 Old Foresthill Road; 26 June 1958, G. B. Wiggins, newly emerged female collected at a midtown service station). ROM.

Genus *Callosamia* Packard

C. angulifera (Walker). 4: Kent (Rondeau Provincial Park, 23 June–24 July); Norfolk (Turkey Point, 9 August 1969). ROM. Known from only a few localities on the north shore of Lake Erie where the foodplant still occurs. First reported in 1965 in Rondeau Provincial Park where *Liriodendron* is a common tree.

C. promethea (Drury). Promethea moth/Prométhée. 4: Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton*; Hastings; Huron; Kent; Lambton; Leeds; Middlesex; Muskoka; Norfolk; Peel; Peterborough*; Renfrew; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. 2 May–10 July. Beth, CNC, McM, ROM, UG, UWO. A partial second generation may occur; e.g., Grey (Meaford, 1 August) and Norfolk (Normandale, 11 September). The foodplant is wild cherry (Chaffey's Locks). Larvae may also feed on other trees and shrubs, e.g., at Rondeau Provincial Park. The species does not always have high population densities and it may not be recorded from a certain locality for years until, during one summer, numerous specimens are observed.

In Chaffeys Locks one pair was found in copula on a wild cherry shrub, 16 June 1972, 1600 hrs. Two females were taken at superactinic lights at night, 18–19 June 1972, and one female at incandescent light at night, 21 June 1972, 0430 hrs. The offspring of these females emerged in the laboratory at Chaffeys Locks beginning 2 June 1973, and the newly emerged females attracted numerous wild males through the open window to their cage. The fertilized females were released. No females were caught during the summer of 1973 at lights.

Genus *Hyalophora* Duncan

H. cecropia (Linné). *Cecropia* moth/samia *cécropia*. 2: Cochrane; Kenora. 3: Rainy River; Sudbury; Timiskaming. 4: Brant; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Northumberland; Parry Sound*; Peel; Renfrew; Simcoe; Wellington; Wentworth; York. 19 May–18 August; peak mid June–mid July. AMNH, Beth, CNC, McM, QU, ROM, UG. Successfully reared in Toronto on lilac, and in Chaffeys Locks on birch and cherry. Larvae collected in Toronto on lilac.

H. columbia (S. I. Smith). 2: Kenora (Vermilion Bay, 22 June 1958, UG). 3: Kenora; Rainy River (Quetico Provincial Park, 1976, Q. Hess. Atikokan. Caliper Lake Provincial Park). 4: Carleton; Frontenac; Grey; Haliburton; Hastings; Middlesex; Muskoka (Fraserburg, 8 June 1968, L. Kohalmi); Nipissing; Parry Sound; Renfrew; Victoria. CM, CNC, ROM, UG. Most specimens represented in CNC emerged from pupae collected by T. N. Freeman in the vicinity of Ottawa. Kohalmi and Moens (1975) published an extensive paper concerning *Hyalophora*, and specimens of this research collection are now deposited in the ROM. The population of *H. columbia* in southern Ontario near London may be extinct. At other places the species seems to have made a comeback after the disaster of the larch sawfly infestations around midcentury.

H. columbia is known to feed only on larch in the east, while the northwestern population feeds on willow, the foodplant of *H. nokomis* (Brodie). A male specimen in the ROM collected at light in Caliper Lake Provincial Park, 11 June 1971, shows exterior characteristics of *H. nokomis*. As Kohalmi and Moens showed that *nokomis* and *columbia* intergrade in northwestern Ontario, specimens from Kenora (Rushing River Provincial Park) should not simply be ascribed to *H. nokomis* (as done in TEA Occasional Publications 1979, 10:70) but should be kept as *H. columbia* (see Kohalmi and Moens, 1975).

Superfamily Sphingoidea
Family Sphingidae
Subfamily Sphinginae
Tribe Sphingini

Genus *Agrius* Hübner

- A. cingulatus* (Fabricius). 4: Carleton; Essex; Middlesex; Wentworth; York. Mid-end September. Beth, CNC, ROM, UG. A straggler from the south.

Genus *Manduca* Hübner

- M. quinquemaculata* (Haworth). Tomato hornworm/sphinx de la tomate. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Norfolk; Renfrew; Simcoe; Stormont; Waterloo; Welland; Wellington; Wentworth; York. 13 June–8 October. Beth, CNC, McM, ROM, UG, UWO.
- M. sexta* (Linné). Tobacco hornworm/sphinx du tabac. 3: Timiskaming. 4: Essex; Kent; Middlesex. CNC, UG. A straggler from the south. Known to breed in Ontario only in the Chatham area since about 1933 (see Miller, 1956).

Genus *Dolba* Walker

- D. hyloeus* (Drury). 4: Carleton; Frontenac; Grenville; Lennox and Addington; Lincoln; Muskoka; Renfrew. 10–15 July. Beth, CNC, McM, ROM. Known from only a few specimens from very few localities.

Genus *Ceratomia* Harris

- C. amyntor* (Geyer). 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Hastings; Kent; Lambton; Lanark; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Renfrew; Simcoe; Victoria; Welland; Wellington; Wentworth; York. 30 May–29 July. AMNH, Beth, CNC, McM, QU, ROM, UG. A partial second generation may occur, e.g., Leeds (Chaffeys Locks, late August 1967).
- C. catalpae* (Boisduval). 4: Essex (Windsor). First known from a FIDS report from Windsor. The original report records as follows: "IX-1-1944, Windsor, 139 Elm Avenue, 500 larvae of *Ceratomia catalpae* on one tree, on *Catalpa* 12' to 15' size, W. R. Lapp, Plant Inspection Officer, 507 Canada Bldg., Windsor. Defoliation not severe as yet. Two trees

examined." However, only a few larvae were taken and none completed development. Recently collected as adult: Essex (Ruthven, 11 July 1981, P. D. Praff, CNC). Also, the Point Pelee Museum has one specimen collected at Point Pelee and another from Essex (Southeast Shoal, 7 September 1985, A. Wormington).

***C. undulosa* (Walker).** Waved sphinx/sphinx ondulé. 2: Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Haliburton; Hasjings; Huron; Kent; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Peel; Renfrew; Simcoe; Stormont; Victoria; Welland; Wellington; Wentworth; York. 27 May–7 August. AMNH, Beth, CNC, McM, QU, ROM, UG. A partial second generation may occur: e.g., Haliburton (Minden, 30 August); (Ottawa, 28 August); (Rondeau Provincial Park, 28 August); (Toronto, 5 September). The form *engeli* Chermock and Chermock is known from Haldimand (Dunnville), Welland (Port Colborne), and Rondeau Provincial Park. AMNH, CNC, ROM.

Genus *Peratrea* Grote

***P. plebeja* (Fabricius).** 4s: Wentworth (Caledonia, 25 June 1971, J. T. Troubridge). CNC.

Genus *Sphinx* Linné

***S. canadensis* Boisduval.** 3: Rainy River; Thunder Bay. 4: Brant; Carleton; Frontenac; Haliburton; Leeds; Lennox and Addington; Muskoka; Nipissing; Peterborough; Renfrew; Waterloo; Wentworth. 12 June–12 August. AMNH, CNC, ROM, UG. Common in the Rainy River District, otherwise local and never in large numbers. The larvae are said to feed on ash, not on blueberry (see Clayton, 1938), but refused both in our lab in 1970 and died.

***S. chersis* (Hübner).** Great ash sphinx/sphinx du frêne. 3: Algoma; Rainy River. 4: Brant; Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Haliburton; Hastings; Kent; Lanark; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 26 May–14 August. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO.

***S. drupiferarum* J. E. Smith.** 2: Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Lambton; Lanark; Leeds; Muskoka; Nipissing; Northumberland; Ontario; Renfrew; Simcoe; Waterloo; Welland; Wentworth; York. 12 May–31 July. AMNH, Beth, CNC, McM, ROM, UG.

- S. eremitus* (Hübner). Hermit sphinx/sphinx ermite. 4: Carleton; Durham; Essex; Halton; Hastings; Huron; Kent; Lambton; Lincoln; Leeds; Wentworth. 5 June–18 July. CNC, ROM. Known from isolated records but not a straggler.
- S. gordius* Cramer. Apple sphinx/sphinx du pommier. 4: Haldimand; Kent. June–July. Known only in the southernmost part of southern Ontario (see Riotte, 1980). The species name is a nomen proprium.
- S. kalmiae* J. E. Smith. Laurel sphinx/sphinx du laurier. 3: Algoma; Rainy River; Thunder Bay; Timiskaming. 4: Carleton; Dufferin; Durham; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Lambton; Lanark; Leeds; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Renfrew; Simcoe; Welland; York. 30 May–30 August. AMNH, Beth, CNC, McM, ROM, UG, UWO. Late August records may be of a partial second generation.
- S. luscitiosa* Clemens. 2: Cochrane; Thunder Bay. 3: Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Haliburton; Hastings; Lambton; Leeds; Lennox and Addington; Middlesex; Renfrew; Wellington; York. 3 June–7 July. AMNH, Beth, CM, CNC, LEM, ROM, UG. The only known species of the genus *Sphinx* extending its range into the Northwest Territories and Yukon. Uncommon. Larvae reared in our laboratory on willow and *Spiraea* sp.
- S. poecila* Stephens. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Lanark; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Parry Sound; Peel; Peterborough; Renfrew; Victoria; Waterloo; York. 24 May–31 July. AMNH, Beth, CM, CNC, FM, McM, QU, ROM, UG. A partial second generation may occur: e.g., (Geraldton, 29 August 1952); (Ottawa, 28 August 1906, two specimens).

The main foodplant is *Myrica gale* Linné and *Spiraea* sp., but species may also occur on birch; in the northern forests they often have been collected by FIDS from *Larix* sp. This species is the most common sphingid in Canada, from the east coast to the eastern slopes of the Rocky Mountains in British Columbia. It also extends south into the United States and flies every year just west of Ontario on the Manitoba and Saskatchewan prairies (very probably also on remnant Wisconsin prairies), at the same time and localities as *Sphinx gordius*, without producing any intermediate hybrids.

Genus *Lapara* Walker

Pine tree sphinx/sphinx du pin. For more information about this genus, see Riotte (1972a); D'Abrera (1986:43).

L. bombycoides Walker. 2: Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Glengarry; Haldimand; Haliburton; Hastings; Huron; Lambton; Leeds; Lennox and Addington; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Wentworth; York. 23 May–10 August. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG, UWO.

Tribe Smerinthini

Genus *Smerinthus* Latreille

S. cerisyi Kirby. 2: Cochrane; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Durham; Frontenac; Haliburton; Halton; Hastings; Huron; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Parry Sound*; Renfrew; Simcoe; York. 18 May–29 July. AMNH, Beth, CM, CNC, McM, ROM, UG, UWO. Widespread and comparatively common in northern Ontario, but only a few records from southern Ontario.

S. jamaicensis (Drury). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Elgin; Essex; Frontenac; Grenville; Haldimand; Hastings; Huron*; Kent; Lambton; Leeds; Lennox and Addington; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Welland; Wellington; Wentworth; York. 6 May–7 August. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG, UWO. A partial second generation may occur, e.g., Parry Sound (Lake Joseph, 21 August 1947). Most of the specimens are of the form *geminatus* Say. The nominate and *tripartitus* Grote forms are rare.

Genus *Paonias* Hübner

The generic name is of masculine gender.

P. excaecatus (J. E. Smith). 2: Kenora; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Brant; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Peel; Peterborough; Renfrew; Simcoe; Victoria; Waterloo; Welland; Wellington; Wentworth; York. 11 May–17 August. AMNH, Beth, CM, CNC, FM, McM, QU, ROM, UG, UWO.

P. myops (J. E. Smith). 3: Algoma; Cochrane; Rainy River; Sudbury. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand;

Haliburton*; Hastings; Huron; Kent; Lambton; Leeds; Lennox and Addington; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound*; Peel; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 20 May–29 July. AMNH, CM, CNC, LEM, McM, QU, ROM, UG.

Genus *Amorpha* Hübner [= *Cressonia* Grote and Robinson]

The previously used generic name *Cressonia* Grote and Robinson, 1865, has to be changed to *Amorpha* Hübner, 1809 (see Fletcher and Nye, 1982).

***A. juglandis* (J. E. Smith).** Walnut sphinx/sphinx du noyer. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 26 May–17 July. AMNH, Beth, CM, CNC, McM, QU, ROM, UG. A partial second generation occurs, e.g., Leeds (Chaffey's Locks, 18 August). Larvae feed freely on *Ostryia virginiana* Miller and relatives (*Corylus*, *Carpinus*, *Fagus*) besides the other well known foodplants of the walnut/hickory family, but they refused *Betula*.

The population west of the Nipigon River into Manitoba belongs to what was described as subspecies *A. j. manitobae* B. P. Clark. The foodplant of this population, however, is *Betula*; W. E. Sieker (personal comment) reported larvae in the field feeding on *Betula* in northern Wisconsin. The male genitalia of *manitobae* differ from *Ostryia*-feeding *juglandis*. As Hodges (1971) suggested, the entire group of *juglandis* populations should be intensively studied to determine the specific status of its components. Type species for this *Amorpha* is *Sphinx juglandis* J. E. Smith, designated by Hübner. See also D'Abrera (1986:94).

Genus *Pachysphinx* Rothschild and Jordan

***P. modesta* (Harris).** 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Essex; Durham; Frontenac; Grenville; Haliburton; Haldimand; Hastings; Huron; Kent; Lanark; Lambton; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound*; Peel; Renfrew; Simcoe; Waterloo; Welland; Wellington; Wentworth; York. 23 May–10 August. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO.

Subfamily Macroglossinae
Tribe Dilophonotini

Genus *Erinnyis* Hübner

- E. ello* (Linné). 4: Essex (Point Pelee, 20 September 1987, A. Wormington, in coll. A.W.); Middlesex (London, 28 September 1886); Wentworth (Hamilton); York (Toronto). ROM, UG. Casual straggler from the south.
- E. obscura* (Fabricius). 4: Essex (Point Pelee, early October 1988, A. Wormington, in coll. A.W.); Middlesex (London, 1897, UG); Welland (Ridgeway, 6 September 1887, collected on wild *Phlox* sp., UG); York (Toronto, 1917). ROM. Casual straggler from the south.

Genus *Aellopos* Hübner

- A. fadus* (Cramer). 4: York (Toronto, two specimens with neither month nor year). CNC. Casual straggler from the south.
- A. titan* (Cramer). 4: Kent (Rondeau Provincial Park, 12 July 1959, Rondeau Park Museum); Lincoln (Grimsby, two worn specimens with neither month nor year). ROM (from the Bethune collection).

Genus *Hemaris* Dalman

- H. diffinis* (Boisduval). Honeysuckle clear-wing sphinx/sphinx du chèvrefeuille. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Essex; Frontenac; Grenville; Haldimand; Haliburton; Hastings; Kent; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Parry Sound; Peel; Peterborough*; Renfrew; Victoria; Wellington; Wentworth; York. 7 May, Hastings (Marmora)–20 August, York (Rouge Hill); with two overlapping generations. AMNH, CM, CNC, McM, QU, ROM, UG, UWO.
- H. gracilis* (Grote and Robinson). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Muskoka; Norfolk. 31 May–21 June. AMNH, CNC, ROM. Known mainly from northern Ontario. Very few specimens in collections.
- H. thysbe* (Fabricius). Hummingbird clear-wing sphinx/sphinx colibri. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Parry Sound; Peterborough*; Prince Edward; Renfrew; Simcoe; Victoria; Welland; Wellington; Wentworth; York. 14 May–12 August. AMNH, CM, CNC, McM, ROM, UG, UWO.

Tribe Philampelini

Genus *Eumorpha* Hübner

- E. achemon* (Drury). Achemon sphinx/sphinx d'Achemon. 4: Carleton; Essex; Frontenac; Haldimand; Halton; Hastings; Huron; Lambton; Lennox and Addington; Middlesex; Ontario; Perth; Welland; Wentworth; York. 1 July–18 August. CNC, LEM, McM, ROM, UG.
- E. fasciata* (Sulzer). 4: Essex (Point Pelee, 4 September 1987, A. Wormington, worn specimen, in coll. A.W.).
- E. pandora* (Hübner). 4: Essex; Frontenac; Grenville; Haldimand; Hastings; Lambton; Leeds; Middlesex; Wentworth; York. 17 June–6 October. CNC, McM, QU, ROM, UG, UWO. This species is uncommon, except near Kingston. A partial second generation occurs, with peak occurrence after mid July.

Tribe Macroglossini

Genus *Sphecodina* Blanchard

- S. abbottii* (Swainson). 4: Essex; Frontenac; Haldimand; Halton; Huron; Lambton; Lanark; Leeds; Middlesex; Peel; Peterborough; Simcoe; Waterloo; Wentworth; York. 5–26 May. AMNH, CNC, LEM, ROM, UG. Local and rare.

Genus *Deidamia* Clemens

- D. inscripta* (Harris). 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Kent; Leeds; Middlesex; Norfolk; Ontario; Peel; Welland; Waterloo; Wentworth; York. 1 May–17 June. CNC, McM, ROM, UG, UWO.

Genus *Amphion* Hübner

- A. floridensis* B. P. Clark [= *Sphinx nesus* Cramer, preocc. by *Sphinx nesus* (= *Theretra nesus*) Drury]. 3: Rainy River. 4: Carleton; Essex; Frontenac; Haldimand; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Lennox and Addington; Lincoln; Middlesex; Muskoka; Norfolk; Parry Sound*; Peel; Victoria; Welland; Wentworth; York. 24 May–13 July. CM, CNC, McM, ROM, UG, UWO. A partial second generation may occur: e.g., Wentworth (Hamilton, 28 August 1950); York (Rouge Hill, 10 August 1937).

Genus *Proserpinus* Hübner

The generic name *Proserpinus* is derived from the Latin verb *proserpere*, to creep forwards, and is a masculine derivation. The specific name of the type species *proserpina*, however, is the name of a Greek goddess and therefore ends in "a."

***P. flavofasciatus* (Walker).** 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Thunder Bay; Timiskaming. 7 June–5 July. AMNH, CM, CNC, Nova Scotia Museum of Science, ROM, United States National Museum of Natural History. Occurs only in the boreal forest region. Rare.

Genus *Darapsa* Walker

***D. myron* (Cramer).** Virginia-creeper sphinx/sphinx de Myron. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Simcoe; Welland; Wellington; Wentworth; York. 5 May–26 July. AMNH, Beth, CM, CNC, LEM, McM, QU, ROM, UG, UWO. A partial second generation may occur, possibly overlapping. The species appears in two distinct colour forms, the normal dark olive green one, and an ochreous one without any green at all. The latter is known from Haldimand (Dunnville), London, Ottawa, Rondeau Provincial Park, and Toronto.

***D. pholus* (Cramer).** 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Essex; Frontenac; Grey; Haldimand; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Ontario; Parry Sound; Renfrew; Welland; Wellington; Wentworth; York. 18 May–15 August. AMNH, Beth, CM, CNC, McM, ROM, UG.

***D. versicolor* (Harris).** 4: Frontenac; Haldimand; Kent; Lanark; Leeds; Lennox and Addington; Peterborough; Welland; Wentworth; York. 19 June–26 July. CNC, ROM, UG. A partial second generation may occur, e.g., Leeds (Chaffey's Locks, late August 1967). Local and rare; more numerous only in Rondeau Provincial Park and Chaffey's Locks.

Tribe Choerocampini

Genus *Xylophanes* Hübner

***X. tersa* (Linné).** 4: Essex (Arner Stop near Kingsville, 24 September 1987, J. Larson. Point Pelee, 25 August–9 September 1987, A. Wormington, four specimens, in coll. A.W.); Middlesex (London, 12 October 1912, two specimens). ROM. Occasional straggler from the south.

Genus *Hyles* Hübner

H. euphorbiae (Linné). 4: Renfrew (Braeside). Imported from Central Europe and released by the Department of Agriculture for weed control. Apparently now well established.

H. gallii (Rottenburg). Bedstraw hawk-moth/sphinx du gaillet. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Thunder Bay; Timiskaming. 4: Carleton; Essex; Durham; Haliburton; Hastings; Huron; Middlesex; Muskoka; Nipissing; Northumberland; Welland; Wellington; Wentworth; York. 30 May–15 August. AMNH, Beth, CM, CNC, McM, ROM, UG. A partial second generation is known (e.g., Geraldton) and has also been experimentally demonstrated. The records from southern Ontario are not recent, and the species may now be considered to be northern.

H. lineata (Fabricius). White-lined sphinx/sphinx à lignes blanches. 2: Cochrane; Kenora, Patricia Portion. 3: Algoma; Rainy River; Thunder Bay; Timiskaming. 4: Brant; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Northumberland; Wentworth; York. 12 June–27 September. AMNH, Beth, CNC, McM, ROM, UG. A straggler all over Ontario; it is not known whether this species breeds in the southern part of the province.

Superfamily Noctuoidea

Family Notodontidae

Systematics of Forbes (1948).

The subfamily Cerurinae is retained at the end of the family.

Subfamily Pygaerinae

Genus *Clostera* Samouelle

C. albosigma Fitch. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Dufferin; Durham; Frontenac; Haldimand; Hastings; Huron; Kent; Lambton; Lanark; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Ontario; Peterborough; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 20 May–24 August. AMNH, Beth, CNC, McM, QU, ROM, UG. An overlapping, partial second generation occurs. The two broods show seasonal dimorphism, the first generation being lighter in colour, the second dark to very dark. In the north (Geraldton and northward), however, the first generation is nearly as dark as the second.

- C. apicalis* (Walker). 2: Algoma; Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Peel; Peterborough; Simcoe; Welland; Wellington; Wentworth; York. 6 May–18 July. AMNH, Beth, CNC, LEM, ROM, UG. A partial second generation occurs in the south (12 July–20 August).
- C. brucei* (Hy. Edwards). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Timiskaming. 4: Muskoka. 30 May–10 July. AMNH, CNC, ROM. Rare, restricted to the boreal, and more northern part of the mixed, forest regions except for the surprising finds in Muskoka District. Three AMNH specimens from the Buchholz collection labelled “Toronto” and “Charlton” with no year, but the same date, are suspect. All three show the same colour, which differs somewhat from that of *C. brucei* from Sudbury and Geraldton. *C. brucei* is otherwise known only from northern Ontario. See the remark under *Parasemia plantaginis*.
- C. inclusa* (Hübner). 4: Brant; Essex; Huron; Kent; Lambton; Norfolk; Welland. 22 May–11 June. CNC, ROM, UG.
- C. strigosa* (Grote). 2: Thunder Bay. 3: Algoma; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Huron; Leeds; Nipissing; Simcoe. 20 May–10 July. CNC, QU, ROM. A partial second generation occurs in the south: Haldimand (Dunnville, 3 August); Huron (Grand Bend, 9 August).

Subfamily Phalerinae

Genus *Datana* Walker

- D. angusii* Grote and Robinson. 4: Carleton; Durham; Frontenac; Hastings; Kent; Leeds; Lincoln; Middlesex; Norfolk; Peterborough; Welland; York. 3 May–27 July. CNC, ROM. A partial second generation occurs: Leeds (Chaffey's Locks, 15 August); Middlesex (Coldstream, 24 August). The remark in Prentice (1962), “distribution limited to the north shore of Lake Erie in southern Ontario,” is no longer valid.
- D. contracta* Walker. 4: Frontenac; Haldimand; Hastings; Huron; Kent; Leeds; Middlesex; Muskoka; Wellington; York. 22 June–21 July. AMNH, CNC, ROM, UG. *D. contracta* cannot be easily separated from *D. integerrima* without dissection. It is therefore possible that the records of *D. integerrima* include some of *D. contracta*, especially on the north shore of Lake Erie.
- D. drexelii* Hy. Edwards. 4: Bruce; Norfolk; Welland; Wentworth; York. 28 May–31 July. CNC, McM, ROM, UG.
- D. integerrima* Grote and Robinson. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Kent; Lambton; Leeds; Lincoln; Middlesex; Norfolk; Ontario; Peterborough; Simcoe; Welland;

Wentworth. 21 May–26 July. AMNH, CNC, McM, QU, ROM, UG, UWO. A partial second generation may occur in August.

D. major Grote and Robinson. 4: Norfolk (St. Williams, 1973, Walter Plath, Jr., only larvae found).

D. ministra (Drury). 2: Kenora. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Brant; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Peel; Peterborough; Renfrew; Simcoe; Victoria; Welland; Wentworth; York. 7 May–9 August. AMNH, Beth, CNC, McM, ROM, UG, UWO. A partial second generation occurs in the south at the end of July and in early August. The species is comparatively common in Ontario.

D. perspicua Grote and Robinson. 4: Brant; Bruce; Frontenac; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Muskoka; Simcoe; Wentworth. 11 July–12 August. AMNH, CNC, McM, ROM, UG. Larvae reared on *Rhus* species.

Subfamily Notodontinae Tribe Notodontini

Genus *Nadata* Walker

N. gibbosa (J. E. Smith). Green oak caterpillar/chenille verte du chêne. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Welland; Wentworth; York. 3 June–30 July. AMNH, CNC, McM, QU, ROM, UG. A partial second generation occurs in August (experimentally proven in Geraldton and Chaffey's Locks). A full second generation was observed in Rondeau Provincial Park.

Genus *Hyperaeschra* Butler

H. georgica (Herrich-Schäffer). 3: Rainy River. 4: Durham; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Muskoka; Nipissing; Simcoe; Welland; York. 26 May–22 July. CNC, McM, ROM, UG.

Genus *Peridea* Stephens

P. angulosa (J. E. Smith). 3: Rainy River; Sudbury. 4: Brant; Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Kent; Lambton; Leeds; Muskoka; Norfolk; Peel; Peterborough; Simcoe; Waterloo; York. AMNH, CNC, McM, QU, ROM, UG. The earliest known date

of appearance is 7 June (Toronto). The first generation appears to intermix with a partial second generation: Leeds (Chaffey's Locks, 18 August).

P. basitriens (Walker). 3: Algoma; 4: Carleton; Durham; Frontenac; Hastings; Kent; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Peel; Peterborough; Simcoe; Waterloo; Wentworth; York. 4 June–5 August. CNC, McM, ROM. The larvae, reared from eggs, were positively associated with this species for the first time in summer 1966 (Riotte, 1969).

P. ferruginea (Packard). 2: Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Peterborough; Simcoe; Welland; Wentworth; York. AMNH, CNC, McM, ROM, UG. The earliest known date of appearance is 16 May (Toronto, 1896), and the next earliest date recorded is 3 June (Ottawa). The latest date is the same as in *P. angulosa*: Leeds (Chaffey's Locks, 16 August); Thunder Bay (Neebing Twp., 20 August).

Genus *Pheosia* Hübner

P. rimosa Packard. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Parry Sound; Peterborough; Simcoe; Welland; Wellington; Wentworth; York. 8 May–28 July. AMNH, Beth, CNC, McM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 16 September (Ottawa).

Genus *Odontosia* Hübner

O. elegans (Strecker). 2: Cochrane; Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Kent; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Peterborough; Simcoe; Victoria; Wentworth. 4 June–27 July. AMNH, CNC, McM, ROM. A partial second generation occurs: Leeds (Chaffey's Locks, 13 August).

Genus *Notodonta* Ochseneimer

N. scitipennis Walker [= *stragula* Grote]. 2: Cochrane; Thunder Bay. 3: Cochrane; Rainy River; Sudbury; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Huron; Lambton; Leeds; Lennox and Addington; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Simcoe;

Wellington; Wentworth; York. 3 May–29 July. AMNH, Beth, CNC, McM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 17 August (Ottawa).

N. simplaria Graef. 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Huron; Leeds; Muskoka; Simcoe; York. 28 May–31 July. AMNH, CNC, ROM. A rare partial second generation occurs: Durham (Kendal, 4 September).

Genus *Nerice* Walker

N. bidentata Walker. 3: Timiskaming. 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Peel; Peterborough; Renfrew; Welland; Wentworth; York. 2 May–4 September. A partial overlapping second generation occurs.

Genus *Ellida* Grote

E. caniplaga (Walker). 3: Rainy River. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Peterborough; Simcoe; Waterloo; Welland; Wellington; Wentworth; York. 18 May–27 July. AMNH, CNC, ROM, UG. A partial second generation occurs: Leeds (Chaffey's Locks, 16 August).

Tribe Gluphisiini

Genus *Gluphisia* Boisduval

G. avimacula Hudson. 2: Cochrane. 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haliburton; Leeds; Muskoka; Nipissing; Renfrew; Simcoe. 13 May–20 June. AMNH, CNC, McM, ROM. The form *slossoniae* Packard may occasionally be found (CNC).

G. lintneri (Grote). 2: Cochrane. 3: Algoma; Thunder Bay. 4: Carleton; Frontenac; Halton; Hastings; Muskoka; Simcoe. 20 April–12 May. CNC, ROM.

G. septentrionis Walker. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Peterborough; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 29 May–31 July. AMNH, Beth, CNC, LEM, McM, ROM, UG, UWO. A partial second generation occurs in late July and August in southern Ontario, and in mid August in northern Ontario.

Tribe Nystaleini

Genus *Symmerista* Hübner

- S. albifrons* (J. E. Smith). Red-humped oakworm/chenille à bosse rouge de chêne. 4: Hastings; Leeds; Middlesex; York. 7 June–30 July. CNC, ROM, UG, UWO.
- S. canicosta* Franclemont. 3: Algoma; Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Haliburton; Hastings; Kent; Leeds; Muskoka; Nipissing; Norfolk; Renfrew; Russell; Simcoe; Welland. 4 June–14 July. CNC, ROM, UWO. Forbes (1948:223) interchanged the numbers of the aedoeagi of *leucitys* and *canicosta*.
- S. leucitys* Franclemont. Orange-humped mapleworm/chenille à bosse orangée. 3: Algoma. 4: Carleton; Durham; Frontenac; Haliburton; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Perth; Peterborough; Simcoe; Waterloo; Welland; York. 6 June–6 July. CNC, McM, QU, ROM. A partial second generation occurs (Rondeau Provincial Park, 24 August).

Genus *Dasylophia* Packard

- D. anguina* (J. E. Smith). 3: Rainy River; Thunder Bay. 25 June–27 July. CNC, ROM. This species only enters into westernmost Ontario.
- D. thyatiroides* (Walker). 4: Carleton; Durham; Frontenac; Haldimand; Huron; Kent; Leeds; Middlesex; Muskoka; Nipissing; Welland; Wentworth; York. 6 June–9 August. CNC, McM, ROM, UG. Pale specimens have been collected from Algonquin Provincial Park, Middlesex (Coldstream), London, Ottawa, Niagara (Ridgeway), and Toronto; dark (melanic) specimens have been collected in Chaffey's Locks, Haldimand (Dunnville), Huron (Grand Bend), Hamilton, Welland (Port Colborne), and Rondeau Provincial Park.

Tribe Heterocampini

Genus *Misogada* Harvey

- M. unicolor* (Packard). 4: Northumberland (Colborne, 6 July 1985, W. J. D. Eberlie). Only known record from Ontario.

Genus *Macrurocampa* Dyar

- M. marthesia* (Cramer). 3: Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Peel; Peterborough; Simcoe; Welland; Wentworth; York. 24 May–31

July. AMNH, CNC, McM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 16 August (Chaffey's Locks). McDunnough (1938) called the Rainy River District population subspecies *M. m. manitobensis*. In fresh specimens the conspicuous splashes are orange-brown, and not green as reported by Forbes (1948:235).

Genus *Heterocampa* Doubleday

- H. biundata* Walker. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Lambton; Leeds; Muskoka; Nipissing; Norfolk; Peterborough; Simcoe; Welland; York. 9 May–7 August. AMNH, CNC, ROM. Uncommon.
- H. guttivitta* (Walker). Saddled prominent/hétérocampe de l'érable. 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Peel; Peterborough; Simcoe; Welland; Waterloo; Wentworth; York. 6 May–11 July. AMNH, Beth, CNC, LEM, McM, ROM, UG, UWO. The melanic form *hugoi* Chermock and Chermock has been collected in Haldimand (Dunnville), Norfolk (Normandale), Durham (Orono), Ottawa, Rondeau Provincial Park, and Lincoln (Vineland).
- H. obliqua* Packard. 3: Sudbury. 4: Durham; Frontenac; Hastings; Huron; Lambton; Leeds; Muskoka; Peel; Peterborough. 23 June–27 July. CNC, ROM.
- H. subrotata* Harvey. 4: Essex (Peele Island South, 8 July 1965). ROM. Known only from Pelee Island.
- H. umbrata* Walker. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Durham; Frontenac; Hastings; Huron; Lambton; Leeds; Lennox and Addington; Muskoka; Nipissing; Parry Sound; Peterborough; Simcoe; Wentworth; York. 4 June–26 July. AMNH, Beth, CNC, QU, ROM, UG. The Ontario population is often referred to the subspecies *H. u. pulverea* Grote and Robinson, but it is doubtful that this subspecies has any validity.

Genus *Lochmaeus* Doubleday

- L. bilineatus* (Packard). Elm prominent/hétérocampe birayée. 4: Brant; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Leeds; Middlesex; Muskoka; Norfolk; Ontario; Peel; Simcoe; Welland; Wellington; York. 6 June–7 August. AMNH, CNC, ROM, UG, UWO. Uncommon. In laboratory cultures, approximately 90% of larvae went into diapause at the end of their development without forming a pupa. Only 10% pupated and released a partial second generation at the end of July. The remainder pupated the following spring.

L. manteo **Doubleday**. Variable oak leaf caterpillar/hétérocampe du chêne. 3: Algoma; Rainy River. 4: Carleton; Durham; Frontenac; Leeds; Middlesex; Muskoka; Simcoe; Wentworth; York. 2–31 July. AMNH, CNC, ROM, UG. Uncommon.

Genus *Schizura* Doubleday

S. apicalis (**Grote and Robinson**). 3: Rainy River; Sudbury. 4: Frontenac; Leeds; Muskoka; Peterborough. 3–29 July. AMNH, CNC, ROM. Rare.

S. badia (**Packard**). 3: Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Haldimand; Lambton; Leeds; Middlesex; Muskoka. 26 June–27 July. CNC, ROM, UG. Rare

S. concinna (**J. E. Smith**). Red-humped caterpillar/chenille à bosse rouge. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Grey; Kent; Leeds; Middlesex; Peterborough; Prince Edward; Simcoe; York. 27 June–6 August. AMNH, CNC, LEM, ROM, UG, UWO. Uncommon.

S. ipomoeae **Doubleday**. Oak and maple humped caterpillar/chenille bossue du chêne et de l'érable. 2: Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Kent; Lambton; Leeds; Lennox and Addington; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Welland; Wentworth; York. 8 June–9 August. A partial second generation may occur in early September, in Sault Ste. Marie and Welland (Port Colborne). The form *cinereofrons* (Packard) occurs together with the typical form, but is uncommon.

S. leptinoides (**Grote**). 2: Thunder Bay. 3: Rainy River; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Grey; Haldimand; Halton; Kent; Leeds; Lennox and Addington; Middlesex; Muskoka; Nipissing; Peterborough; Wentworth; York. 3 June–27 July. AMNH, CNC, LEM, McM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is in September (Toronto).

S. unicornis (**J. E. Smith**). Unicorn caterpillar/chenille licorne. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Grey; Haldimand; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Nipissing; Peel; Peterborough; Renfrew; Simcoe; Victoria; Wellington; Wentworth; York. 25 May–20 August. A partial overlapping second generation occurs in southern Ontario with many specimens collected in August.

Genus *Oligocentria* Herrich-Schäffer

O. lignicolor (**Walker**). 3: Algoma; Kenora; Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Kent; Lambton; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peterborough; Simcoe; York. 4 June–14

July. AMNH, Beth, CNC, LEM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 18 August in Algoma (Echo Lake).

O. semirufescens (Walker). 2: Kenora. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Frontenac; Hastings; Kent; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Peterborough; Simcoe; Wentworth; York. 14 June–6 August. AMNH, CNC, McM, ROM, UG, UWO. A partial overlapping second generation occurs. The latest known date of occurrence is 4 September in Thunder Bay (Neebing Twp.). This species has a tendency to produce melanic specimens in the deciduous forest region, light-coloured specimens with strong, dull red shading in the northern part of the eastern mixed forest region, and a more uniformly gray form in the mixed forest region west of Lake Superior.

Genus *Hyparpax* Hübner

H. aurora (J. E. Smith). 4: Huron (Grand Bend, 17 June 1936); Kent (Rondeau Provincial Park, 11 July 1965). CNC, ROM. Only two specimens are known from Ontario.

Subfamily Cerurinae

Genus *Furcula* Lamarck

F. borealis (Guérin-Ménéville). 3: Thunder Bay (Pukaskwa National Park). 4: Durham; Frontenac; Haldimand; Kent; Leeds; Middlesex; Muskoka; Northumberland; Peterborough; Renfrew; York. 15 June–3 August. Beth, CNC, ROM, UG. Rare in Ontario with few specimens in collections.

F. cinerea (Walker). 3: Rainy River; Sudbury. 4: Carleton; Durham; Frontenac; Grey; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Peterborough; Simcoe; Victoria; Welland; Wentworth; York. 27 May–29 July. AMNH, Beth, CNC, LEM, McM, ROM. A partial overlapping second generation occurs. The latest known date of occurrence is 13 August in Haldimand (Dunnville).

F. modesta (Hudson). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Victoria; Wentworth; York. 21 May–31 July. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 5 September (Toronto).

F. occidentalis (Lintner). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Nipissing; Norfolk; Simcoe; Welland; Wellington; Wentworth; York. 19 May–29 July. AMNH, Beth, CNC, LEM, McM, ROM, UG. A partial overlapping second generation occurs. The latest known date of occurrence is 22 August (Toronto).

F. scolopendrina (Boisduval). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Hastings; Muskoka. 2 June–27 July. AMNH, CNC, ROM. The Ontario population is often referred to subspecies *F. s. aquilonaris* (Lintner), which may be specifically distinct from *F. scolopendrina*. It is decidedly distinct and easily separable (even when greasy) from *F. modesta*, with a later flight period than *F. modesta* and apparently only one generation. It is almost exclusively northern and most numerous in the mixed forest region west of Lake Superior, e.g., in Thunder Bay (Neebing Twp.).

Genus *Cerura* Schrank

C. scitisscripta Walker. 2: Kenora. 3: Rainy River; Sudbury. 4: Carleton; Durham; Halton; Lambton; Leeds; Muskoka; Nipissing; Peterborough; Simcoe; Waterloo; Wellington*; Wentworth. 3–28 June. CNC, ROM. Known only from isolated localities. Subspecies *C. s. canadensis* McDunnough; *C. s. multisscripta* Riley.

Family Arctiidae Subfamily Lithosiinae Tribe Lithosiini

Genus *Eilema* Hübner [= *Lexis* auct.]

E. bicolor (Grote). Smoky moth/lexis bicolore. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Kenora; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Frontenac; Hastings; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Parry Sound; Simcoe. 3 July–23 August. AMNH, CNC, ROM, UG. The statement by McGugan (1958), "The absence of records north of Lake Superior . . . should be noted," is no longer valid.

Genus *Crambidia* Packard

C. casta (Packard). 3: Algoma; Sudbury; Thunder Bay. 4: Carleton; Leeds; Muskoka; Nipissing. 15 August–4 September. CNC, ROM.

- C. pallida* Packard. 3: Rainy River; Thunder Bay. 4: Carleton; Frontenac; Hastings; Huron; Lambton; Leeds; Middlesex. 15 July–3 September. AMNH, CNC, ROM, UG.
- C. pura* Barnes and McDunnough. 4: Frontenac. Glenburnie and Perth Road are the only known localities in Ontario. CNC, ROM. See Riotte (1970b).

Genus *Cisthene* Walker

- C. plumbea* Stretch. 4: Leeds (Chaffey's Locks, 18 July 1967; 13 July 1963); Middlesex (London, 12 August 1886). 13 July–12 August. CNC, ROM, UG. Only these three specimens have adequate data.

Genus *Lycomorpha* Harris

- L. pholus* (Drury). 2: Kenora; Thunder Bay. 3: Kenora; Rainy River; Thunder Bay. 4: Carleton; Frontenac; Hastings; Leeds; Lincoln; Muskoka; Nipissing; Parry Sound; Peel; Peterborough; Simcoe; Victoria; Wentworth; York. 1 July–7 September. CNC, ROM, UG.

Genus *Hypoprepia* Hübner

- H. fucosa* Hübner. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Lambton; Leeds; Lincoln; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Ontario; Peel; Simcoe; Stormont; Victoria; Welland; Wellington; Wentworth; York. 4 July–22 August. AMNH, Beth, CNC, QU, ROM, UG. The form *plumbea* Hy. Edwards has been collected only in the north.
- H. miniata* (Kirby). 3: Algoma; Rainy River; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Renfrew; Simcoe; Victoria; Wellington; Wentworth. 24 June–5 September. AMNH, Beth, CNC, McM, QU, ROM, UG.

Genus *Clemensia* Packard

- C. albata* Packard. 2: Cochrane; Kenora; Thunder Bay. 3: Kenora; Sudbury; Thunder Bay. 4: Frontenac; Hastings; Leeds; Muskoka; Ontario; Peterborough; Simcoe; Victoria; Wentworth. 8 July–4 August. AMNH, CNC, ROM, UG.

Subfamily Arctiinae
Tribe Callimorphini

Genus *Utetheisa* Hübner

U. bella (Linné). 4: Essex; Middlesex; Muskoka; Simcoe; York. 3 July–26 August. Beth, CNC, McM, ROM.

Genus *Haploa* Hübner

- H. clymene* (Brown). 4: Haldimand; Kent; Middlesex; Wentworth; York. 24 July–1 August. AMNH, CNC, ROM, UG. Rare.
- H. colona* (Hübner). 4: Kent (Rondeau Provincial Park, 19 July 1965). ROM. The specimen represents the form *fulvicosta* (Clemens).
- H. confusa* (Lyman). 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Elgin; Frontenac; Grenville; Haliburton; Halton; Hastings; Huron; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Ontario; Peel; Simcoe; Victoria; Wellington; Wentworth; York. 10 June–15 August. AMNH, Beth, CNC, McM, ROM, UG, UWO.
- H. contigua* (Walker). 4: Carleton; Durham; Frontenac; Halton; Hastings; Kent; Leeds; Muskoka; Peterborough; Simcoe; Victoria; Wentworth; York. 1 July–5 August. CNC, McM, QU, ROM.
- H. lecontei* (Guérin-Méneville). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Huron; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Peel; Peterborough; Renfrew; Wellington; Wentworth; York. 9 June–17 July. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG.
- H. militaris* (Harris). 4s: Essex (Peelee Island); Kent (Rondeau Provincial Park). Early July. ROM. Retained as a species in its own right following Forbes (1960). Only recorded in southernmost Ontario.
- H. reversa* (Stretch). 4: Lambton. CNC, ROM. Pinery Provincial Park and Grand Bend are the only known localities.

Tribe Arctiini

Genus *Holomelina* Herrich-Schäffer

- H. aurantiaca* (Hübner). 2: Kenora. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Hastings; Huron; Lambton; Leeds; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Wentworth; York. 7 June–21 August. AMNH, Beth, CNC, McM,

ROM, UG. A partial second generation may occur: Wentworth (Ancaster, 2 September); Essex (Point Pelee, 14 September).

H. ferruginosa (Walker). 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Huron; Lambton; Leeds; Manitoulin; Muskoka; Nipissing; Norfolk; Ontario; Parry Sound; Peel; Victoria; Welland; Wentworth; York. 1 June–18 August. AMNH, Beth, LEM, McM, ROM, UG.

H. laeta (Guérin-Méneville). 2: Kenora. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Hastings; Middlesex; Muskoka; Nipissing; Parry Sound; Simcoe. 2 June–2 August. AMNH, CNC, ROM, UG. Subspecies *H. l. treatii* (Grote).

H. lamae (T. N. Freeman). 2: Thunder Bay (Geraldton). 3: Rainy River (One Sided Lake). 6–23 July. AMNH, ROM. Known only from bog areas around Lake Superior and Lake Nipigon, with only two specimens in collections.

H. opella (Grote). 4: Kent; Leeds; Peterborough. 19 May–18 July. CNC, ROM. Recorded only in southernmost Ontario from Rondeau Provincial Park, Chaffey's Locks, Frontenac (Perth Road), and Peterborough (Kawartha Park).

Genus *Parasemia* Hübner

P. plantaginis (Linné) [= *scudderi* (Packard)]. 2: Cochrane; Kenora, Patricia Portion; Thunder Bay. 3: Thunder Bay; Timiskaming. 10 June–16 July. AMNH, CNC, LEM, ROM, UG. The record of two specimens in AMNH (from the Buchholz collection), labelled “Toronto VI-28” and “VII-20” without year, are questionable. This species has been recorded only from northern Ontario. Two other specimens, labelled “Charlton VI-10-1932” and “VI-18” without year, are also questionable; the labels are in the same handwriting as the Toronto specimens. See the remarks under *Clostera brucei* and *Spilosoma vestale*.

Genus *Pyrharctia* Packard [= *Isia* auct.]

P. isabella (J. E. Smith). 3: Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Frontenac; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Peel; Peterborough; Renfrew; Simcoe; Victoria; Wellington; Wentworth; York. 26 May–20 September. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO.

Genus *Estigmene* Hübner

E. acrea (Drury). Salt-marsh caterpillar/chenille des marais salants. 3: Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Middlesex; Muskoka; Norfolk; Northumberland; Ontario; Peel; Simcoe; Welland; Wellington; Wentworth; York. 12 May–end summer. A partial second generation may occur in the south (London, 3 October). One female specimen in the ROM is completely yellow (Toronto, 25 May 1898).

Genus *Spilosoma* Curtis

The generic name is of neuter gender and the species names must correspond to this. See International Code of Zoological Nomenclature (1985, article 30) where *-soma* is specially mentioned as neuter. See also Paclt (1979).

S. congruum Walker. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Kent; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Parry Sound; Peel; Simcoe; Wellington; York. 8 May–18 July. AMNH, CNC, McM, QU, ROM, UG. A partial second generation may occur.

S. dubium (Walker) [= *primum* Slosson]. 3: Algoma; Sudbury; Timiskaming. 4: Carleton; Durham; Frontenac; Hastings; Haliburton; Leeds; Manitoulin; Muskoka; Parry Sound; Simcoe; York. 15 May–23 June. AMNH, CNC, McM, ROM, UWO. A partial second generation may occur: Timiskaming (Larder Lake, 30 July).

S. latipennis Stretch. 4: Dundas; Durham; Elgin; Haldimand; Huron; Kent; Middlesex; Norfolk; Wentworth; York. 31 May–10 July. AMNH, CNC, LEM, ROM, UG.

S. vestale Packard. The AMNH specimen labelled “Charlton, 7 July” (from the Buchholz collection) is highly suspect.

S. virginicum (Fabricius). Yellow woollybear/diacrisie de Virginie. 2: Cochrane; Kenora; Kenora, Patricia Portion; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Brant; Bruce; Carleton; Durham; Elgin; Frontenac; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Oxford; Peel; Peterborough; Renfrew; Simcoe; Waterloo; Welland; Wellington; Wentworth; York. 6 May–28 August. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. A partial second generation may occur. Melanic specimens (form *fumosum* Strecker) from Geraldton and Longlac are in the ROM. Partially melanic specimens from Toronto and Renfrew (Petawawa) in the ROM, similar

to the picture in Holland (1915, plate XVI, fig. 7), were erroneously assigned to form *fumosum* Strecker.

Genus *Hyphantria* Harris

H. cunea (Drury). Fall webworm. 2: Cochrane; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Lambton*; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Peel; Renfrew; Simcoe; Waterloo; Wellington; Wentworth; York. 28 April–12 September. AMNH, Beth, CNC, McM, ROM, UG, UWO. A partial second generation may occur in the north in late July and early August, Rainy River (One Sided Lake) and Sudbury; at least two full generations occur locally in the south. All Ontario material is here referred to *H. cunea* (see McGugan, 1958; Heinicke, 1962; Morris, 1963). The pure white form dominates. A few checkered specimens exist in collections but are from many years ago, with the exception of one specimen with a few black spots from Leeds (Chaffey's Locks, 23 June 1963). An unnamed webworm with a light-headed caterpillar (see Morris, 1963) occurs sympatrically with *cunea* from New Brunswick to Georgia, but is not yet recorded in Ontario.

Genus *Ecpantheria* Hübner

E. scribonia (Stoll). 4: Leeds; Middlesex; Norfolk; Welland; Wentworth. CNC, ROM. One CNC specimen was referred to by Bethune (1894) in his manual: "larva found near London, May 6, 1903, under a log—sent to Dr. Fletcher, pupated May 9, imaged June 4, at Ottawa." The Leeds specimen was collected 25 June 1975 by J. J. Belwood, on an incandescent light run by the ROM at Chaffey's Locks, Queen's University Biological Station. More specimens have since been collected in Leeds. See TEA Occasional Publications 1985, 16:20.

Genus *Phragmatobia* Stephens

P. assimilans Walker. 2: Cochrane; Thunder Bay. 3: Algoma; Kenora; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Leeds; Muskoka; Parry Sound; Renfrew; Simcoe. 18 April–19 June. AMNH, CNC, LEM, ROM. A boreal species, only rarely found in the south.

P. fuliginosa (Linné). 2: Cochrane; Kenora; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Nipissing; Norfolk; Northumberland; Ontario; Peel; Renfrew; Simcoe; Welland; Wellington; Wentworth; York. 21 April–31 July. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG. A partial second generation

may occur in August–September. Subspecies *P. f. rubricosa* (Harris).
***P. lineata* Newman and Donahue.** 4: Haldimand; Kent. 29 June–20 July.
ROM, Rondeau Provincial Park Museum. Only three specimens
recorded from Ontario. Recorded in Canada from Calgary, Alberta,
and from Manitoba.

Genus *Platarctia* Packard

***P. parthenos* (Harris).** 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Kenora;
Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton;
Muskoka; Nipissing; Norfolk*; Wellington. 10 June–19 August.
AMNH, CNC, LEM, McM, ROM, UG.

Genus *Arctia* Schrank

***A. caja* (Linné).** 2: Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay.
4: Bruce; Carleton; Frontenac; Hastings; Leeds; Lincoln; Muskoka;
Nipissing; Northumberland; Parry Sound; Peel; Peterborough;
Renfrew; Simcoe; Wellington; York. 12 June–20 August. AMNH, Beth,
CNC, McM, ROM, UG, UWO. Subspecies *A. c. americana* Harris.

Genus *Apantesis* Harris

See Ferguson (1985).

***A. nais* (Drury).** 4: Essex; Frontenac; Haliburton; Hastings; Kent; Leeds;
Lincoln; Muskoka; Norfolk; Peel; Welland; Wentworth; York. 25
May–5 July. CNC, QU, ROM, UG. A partial second generation may
occur: Lincoln (Vineland, 9 September). Specimens of *A. nais* and *A.*
phalerata from over the entire range have been dissected before
assigning them to either of the two species. Specimens of *A. vittata*
(Fabricius), which have distinctive male genitalia, were not found
among the Ontario specimens surveyed.

***A. phalerata* (Harris).** 4: Brant; Bruce; Durham; Elgin; Frontenac;
Haldimand; Hastings; Kent; Leeds; Lincoln; Middlesex; Nipissing;
Norfolk; Welland; Wellington; Wentworth; York. 31 May–autumn.
CNC, ROM, QU. The only moth species in Ontario that may have
three overlapping generations; a third partial generation may occur.
First generation, end of May–mid July; second generation, mid
July–end August in Wentworth (Ancaster, 29 August), Durham (Port
Hope, 27 August. Trenton, 24 August); partial third generation, mid
September–mid October (Algonquin Provincial Park, 12 September),
(London, 8 October).

Genus *Grammia* Rambur

See Ferguson (1985).

***G. anna* (Grote).** 4: Brant; Frontenac; Haldimand; Hastings; Kent; Leeds; Lincoln; Middlesex; Norfolk; Peel; Welland; Wentworth; York. Earliest date: 4 June. Beth, CNC, McM, ROM, UG. A partial second generation may occur.

The normal form *persephone* (Grote) was collected at Brant (Brantford), Chaffeys Locks, Frontenac (Cole Lake), Haldimand (Dunnville), Halton (Erindale), Hamilton, Welland (Port Colborne), Rondeau Provincial Park, and Toronto. It occurs together with the nomotypical form.

***G. arge* (Drury).** 4: Bruce; Carleton; Durham; Frontenac; Haldimand; Hastings; Lambton; Leeds; Middlesex; Norfolk; Peel; Renfrew; Welland; Wellington; Wentworth; York. 5 May–23 September. AMNH, CNC, ROM, UG. A partial second generation may occur.

***G. celia* (Saunders).** 3: Sudbury. 4: Carleton; Durham; Frontenac; Huron; Leeds; Middlesex; Victoria; Wentworth; York. 2 June–7 August. Beth, CNC, McM, ROM. A partial second generation may occur. One ROM specimen has Saunders' labels and is marked *type*, from London. However, Bethune (1894) notes, "no. 110, *Arctia celia*, Toronto (Gore Vale by A.) 1862, TYPE, given to Dr. Fletcher October 1902." This specimen was found in 1966 in CNC.

***G. doris* (Boisduval).** 2: Cochrane; Thunder Bay. 3: Sudbury. 4: Muskoka; Simcoe. 3 June–20 July. AMNH, CNC, ROM.

***G. figurata* (Drury).** 3: Thunder Bay (Pukaskwa National Park). 4: York (Toronto, no other data). CNC. The questionable Toronto specimen belongs to form *preciosa* Nixon.

***G. gibsoni* McDunnough.** Single specimen recorded from Wellington (Guelph, June 1924, D. C. B. Duff), identified by E. G. Munroe (then McGill University, now CNC [ret.]). ROM. The record is questionable.

***G. oithona* (Strecker).** 4: Durham; Halton; Simcoe; Wellington. 23 June–9 September. CNC, ROM. A partial second generation may occur. The record from Simcoe Co. is only a note in Bethune (1894); there is no specimen representing this record. All preserved specimens belong to the normal form *rectilinea* (French).

***G. parthenice* (W. Kirby).** 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Frontenac; Hastings; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Ontario; Simcoe; Victoria; Wentworth; York. 8 June–4 September. AMNH, Beth, CNC, McM, ROM, UG, UWO. A partial second generation may occur. Subspecies *G. p. intermedia* (Stretch) has been collected at Geraldton and Hamilton.

- G. phyllira* (Drury). 3: Sudbury. 4: Bruce; Carleton; Durham; Frontenac; Grey; Hastings; Huron; Kent; Lambton; Middlesex; Muskoka; Norfolk; Prince Edward; Wellington; Wentworth; York. 14 June–5 October. AMNH, Beth, CNC, LEM, ROM, UG, UWO. A partial second generation may occur in the south.
- G. quenseli* (Paykull). 1: Kenora, Patricia Portion (Cape Henrietta Maria, 7 July 1948). ROM; represented by a single specimen in the collection. Subspecies *G. q. gelida* (Möschler).
- G. virgo* (Linné). 2: Cochrane; Kenora. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haliburton; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Northumberland; Ontario; Parry Sound*; Peterborough; Peel; Welland; Wellington; Wentworth; York. 9 June–5 September. A partial second generation may occur in the south.
- G. virguncula* (W. Kirby). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Lambton; Leeds; Lincoln; Manitoulin; Muskoka; Ontario; Welland; Wellington; Wentworth; York. 3 June–15 September. AMNH, Beth, CNC, McM, ROM, UG. A partial second generation may occur in the south. The form *otiosa* (Neumoegen and Dyar) is recorded from Carleton (Mer Bleue) and Toronto.
- G. williamsii* (Dodge). William's tiger moth/apantèse du blé. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Manitoulin; Muskoka; Nipissing; Parry Sound; Simcoe. 20 June–15 August. AMNH, CNC, LEM, McM, ROM, UG. Distinctly northern.

Tribe Phaegopterini

Genus *Halysidota* Hübner

- H. harrisii* Walsh. Sycamore tussock moth/halisidote du sycomore. 4: Carleton; Durham; Kent; Welland. FIDS reared the species from larvae on white oak, basswood, birch, and sycamore. The larvae are easily separated from those of *H. tessellaris*.
- H. tessellaris* (J. E. Smith). Pale tussock moth/halisidote du pommier. 3: Sudbury; Thunder Bay. 4: Bruce; Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Peel; Simcoe; Wellington; Wentworth; York. 6 June–14 August. AMNH, CNC, McM, ROM, UG, UWO. This species is mixed with *H. harrisii* in collections; the attempt has not been made to

separate the species on the basis of genitalia preparations, which would require the dissection of every available specimen.

Genus *Lophocampa* Harris

- L. caryae* Harris.** Hickory tussock moth/halisidote du caryer. 4: Carleton; Durham; Elgin; Frontenac; Haldimand; Hastings; Leeds; Middlesex; Norfolk; Victoria; Welland; Wentworth; York. 21 May–10 July. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. Bethune (1894) noted one specimen caught in Toronto alive on 13 March 1863.
- L. maculata* Harris.** Spotted tussock moth/halisidote maculée. 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Carleton; Durham; Haldimand; Hastings; Kent; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Northumberland; Ontario; Peel; Waterloo; Wellington; Wentworth; York. 29 May–19 August. AMNH, Beth, CNC, McM, ROM, UG, UWO. Normally the latest date of occurrence is in mid July; the Thunder Bay (Fort William) record of 19 August might represent a partial second generation.

Genus *Cynia* Hübner

- C. oregonensis* (Stretch).** 3: Sudbury. 4: Carleton; Durham; Essex; Frontenac; Hastings; Kent; Lambton; Leeds; Lincoln; Parry Sound; Simcoe; Wentworth; York. 20 May–29 July. AMNH, CNC, LEM, ROM.
- C. tenera* Hübner.** 2: Cochrane. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Elgin; Essex; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Muskoka; Norfolk; Peel; Simcoe; Welland; Wentworth; York. 3 June–22 July. AMNH, Beth, CNC, QU, ROM, UG. A partial second generation occurs in the south.

Genus *Euchaetes* Harris

- E. egle* (Drury).** 4: Bruce; Carleton; Durham; Essex; Frontenac; Haldimand; Hastings; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Peel; Renfrew; Simcoe; Welland; Wentworth; York. 11 May–June. AMNH, Beth, CNC, McM, QU, ROM, UG. A partial second generation overlaps with the first generation. The latest recorded date of occurrence is 23 August, in Frontenac (Westbrook).

Subfamily Ctenuchinae

Genus *Ctenucha* W. Kirby

- C. virginica* (Esper). 2: Cochrane; Kenora; Thunder Bay. 3: Algoma; Cochrane; Rainy River; Sudbury; Thunder Bay; Timiskaming. 4: Brant; Bruce; Carleton; Dufferin; Durham; Elgin; Essex; Frontenac; Grey; Haldimand; Hastings; Kent; Lambton; Leeds; Manitoulin; Middlesex; Muskoka; Nipissing; Norfolk; Northumberland; Ontario; Parry Sound; Peel; Peterborough; Renfrew; Simcoe; Victoria; Waterloo; Wellington; Wentworth; York. 26 May–19 August. AMNH, Beth, CNC, LEM, McM, QU, ROM, UG, UWO. A partial second generation may occur in the south: Brant (Brantford, September 1948).

Genus *Cisseps* Franclemont

- C. fulvicollis* (Hübner). 2: Cochrane; Thunder Bay. 3: Cochrane; Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Essex; Frontenac; Grenville; Haldimand; Halton; Hastings; Huron; Kent; Lambton; Leeds; Lincoln; Middlesex; Muskoka; Norfolk; Ontario; Parry Sound; Peel; Peterborough; Simcoe; Welland; Wellington; Wentworth; York. 6 June–5 October. AMNH, CNC, McM, QU, ROM, UG, UWO. A partial second generation may occur.

Family Lymantriidae

Subfamily Orgyiinae

Tribe Orgyiini

Genus *Dasychira* Hübner

- D. basiflava* (Packard). 4: Leeds; Wentworth. CNC, ROM.
- D. dorsipennata* (Barnes and McDunnough). 4: Elgin; Leeds; Peterborough. CNC, ROM. Larvae found and reared on aspen. The freshly emerged moth has a peculiar greenish shine whereas the preceding species is brownish.
- D. obliquata* (Grote and Robinson). 2: Cochrane. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Leeds; Nipissing; Norfolk; Peterborough; Prescott. AMNH, CNC, ROM.
- D. pinicola* (Dyar). 3: Rainy River. 4: Leeds; Muskoka; Nipissing; Simcoe. CNC, ROM.
- D. plagiata* (Walker). 2: Thunder Bay. 3: Algoma; Kenora; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Leeds; Muskoka; Nipissing; Parry Sound; Peel; Victoria; Wellington. AMNH, CNC, ROM.

D. vagans (Barnes and McDunnough). 2: Cochrane; Thunder Bay. 3: Rainy River; Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Grenville; Leeds; Muskoka; Nipissing; Timiskaming; Victoria. AMNH, CNC, ROM. Only since Ferguson's (1978) painstaking and definitive revision of the genus are we able to identify the North American species. As the basic work on this checklist was concluded in late 1975, only records from CNC (identified by Ferguson), ROM (compared to those), and AMNH (as far as "ex ROM") have been included. There certainly were *Dasychira* specimens also in other consulted collections; however, no certain identification was possible at the time.

Genus *Orgyia* Ochsenheimer

- O. antiqua* (Linné). 2: Cochrane; Thunder Bay. 3: Sudbury; Thunder Bay. 4: Carleton; Durham; Frontenac; Hastings; Lanark; Muskoka; Nipissing; Northumberland; Renfrew; Wellington; York. 29 July–23 September. AMNH, Beth, CNC, ROM, UG, UWO. Slight differences in the male genitalia exist between European and Canadian specimens. Subspecies *O. a. nova* Fitch.
- O. definita* Packard. 4: Carleton; Dufferin; Elgin; Grey; Haldimand; Halton; Hastings; Kent; Lambton; Leeds; Middlesex; Norfolk; Ontario; Oxford; Parry Sound; Wentworth. 7 August–4 October. CNC, ROM, UG. Common at Queen's University Biological Station at Chaffey's Locks, where the larvae frequent basswood. Numerous cultures were reared in the laboratory from egg masses produced by females set out in little cages near basswood and fertilized regularly by wild males in the late morning hours. Wild female larvae, recognizable by their larger size, were often observed leaving their native basswood trees for neighbouring ones for pupation. One of our first rearings was from one of these larvae. See also Riotte (1977).
- O. plagiata* (Walker). 3: Sudbury; Thunder Bay. 4: Algoma; Bruce; Carleton; Durham; Essex; Frontenac; Grey; Haldimand; Hastings; Huron; Kent; Leeds; Lincoln; Middlesex; Muskoka; Northumberland; Parry Sound; Peel; Renfrew; Victoria; Welland; Wellington; York. 5 July–4 October. AMNH, Beth, CNC, McM, QU, ROM, UG, UWO. Here regarded as full species (Ferguson, 1978), not as a subspecies of *O. leucostigma* (J. E. Smith). This species differs from *O. leucostigma* [= *O. meridionalis* Riotte] and *O. wardi* Riotte (from Nova Scotia) in characters of the female genitalia and early developmental stages. See Riotte (1971b, 1974, 1977, 1989); Chua, Riotte, and Gilmour (1976).

Subfamily Lymantriinae

Tribe Lymantriini

Genus *Lymantria* Hübner

L. dispar (Linné). 4: Essex (Point Pelee*); Frontenac (Wolfe Island); Glengarry; Northumberland; Peterborough; York. CNC, ROM. Collected for the first time in 1969, this species has spread to the mainland around Kingston.

Genus *Leucoma* Stephens

L. salicis (Linné). 4: Carleton (Ottawa and vicinity). CNC, ROM. Collected for the first time in 1973 by E. G. Munroe.

Family Noctuidae

This family is arranged according to Poole (1989).
The genera in the subfamilies and the species in the genera are arranged alphabetically.

Subfamily Herminiinae

Genus *Bleptina* Guenée

B. caradrinalis Guenée. 3. 4.

Genus *Chytolita* Grote

C. morbidalis (Guenée). 4.

C. petrealis Grote. 3. 4.

Genus *Dyspyralis* Warren

D. illocata Warren. 4.

D. nigella (Strecker). 4.

D. puncticosta (Smith). 4.

Genus *Idia* Hübner

I. aemula Hübner. Spruce epizeuxis/ noctuelle de l'épinette. 2. 3. 4.

I. americalis (Guenée). 2. 3. 4.

I. denticulalis (Harvey). 4: Frontenac (Sharbot Lake).

I. diminuendis (Barnes and McDunnough). 4.

I. forbesi (French). 4s: Frontenac (Perth Road); Leeds (Chaffey's Locks).

- I. julia* (Barnes and McDunnough). 4.
I. lubricalis (Geyer). 2. 3. 4.
I. majoralis (Smith). 4.
I. rotundalis (Walker). 4s: Frontenac (Perth Road); Leeds (Chaffeys Locks).
I. scobialis (Grote). 4.

Genus *Lascoria* Walker

- L. ambigualis* Walker. 4.

Genus *Lomanaltes* Grote

- L. eductalis* (Walker). 2. 3. 4.

Genus *Macrochilo* Hübner [= *Hormisa* Walker]

- M. absorptalis* Walker. 3. 4.
M. bivittata (Grote). 4.
M. litophora (Grote). 4.
M. louisiana (Forbes). 4s: Hastings (Trenton).
M. orciferalis Walker. 3. 4.

Genus *Palthis* Hübner

- P. angulalis* (Hübner). Spruce harlequin/arlequin de l'épinette. 2. 3. 4.
P. asopialis (Guenée). 3: Thunder Bay (Nipigon).

Genus *Phalaenophana* Grote [= *Philometra* Grote]

- P. pyramusalis* (Walker). 2. 3. 4.

Genus *Phalaenostola* Grote

- P. eumelusalis* (Walker). 4.
P. hanhami (Smith). 3. 4.
P. larentioides Grote. 4.
P. metonalis (Walker). 2. 3. 4.

Genus *Polypogon* Schrank [= *Zanclognatha* Lederer]

- P. jacchusalis* (Walker). 4.
P. laevigata (Grote). 4.
P. lituralis (Hübner). 4.
P. litalba (Smith). 3.
P. obscuripennis (Grote). 2. 3.

- P. ochreipennis* (Grote). 2. 3. 4.
P. pedipilalis (Guenée). 3. 4.
P. protumnusalis (Walker). 4.
P. theralis (Walker). 4.

Genus *Renia* Guenée

- R. factiosalis* (Walker). 2. 4.
R. flavipunctalis (Geyer). 3. 4.
R. sobrialis (Walker). 3. 4.

Genus *Tetanolita* Grote

- T. mynesalis* (Walker). 4s: Kent (Chatham).

Subfamily Hypenodinae

Genus *Hypenodes* Doubleday

- H. caducus* (Dyar). 4.
H. fractilinea (Smith). 4.
H. palustris Ferguson. 2: Thunder Bay (Geraldton).

Subfamily Hypeninae

Genus *Hypena* Schrank [= *Bomolocha* Hübner,
Ophiuche Hübner, *Parahypenodes* Barnes and McDunnough,
Plathypena Grote]

- H. abalienalis* (Walker). 4.
H. atomaria Smith. 4s: Essex (Point Pelee)
H. baltimoralis (Guenée). 4.
H. bijugalis (Walker). 4n: Carleton (Ottawa).
H. deceptalis (Walker). 4.
H. edictalis (Walker). 4.
H. humuli Harris. 2. 3. 4.
H. madefactalis (Guenée). 4.
H. manalis (Walker). 4.
H. minualis (Guenée). 4.
H. palparia (Walker). 3. 4.
H. quadralis Barnes and McDunnough. 4s: Hastings (Trenton).
H. scabra (Fabricius). Green clover worm/phalène des légumineuses. 2. 3. 4.
H. sordidula (Grote). 4.

Subfamily Ophiderinae [= Rivulinae]

Genus *Alabama* Grote

A. argillacea (Hübner). 2. 3. 4. Straggler.

Genus *Anomis* Hübner

A. erosa Hübner. 4s: Leeds (Chaffeys Locks). Occasional straggler.

A. flava (Fabricius). 4s: Norfolk (Delhi); York (Toronto). Occasional straggler.

Genus *Anticarsia* Hübner

A. gemmatalis Hübner. 4s: York (Toronto); Essex (Southeast Shoal*).
Occasional straggler.

Genus *Ascalapha* Hübner [= *Erebus* Latreille]

A. odorata (Linné). 3. 4. Occasional straggler as far as Thunder Bay.

Genus *Calyptra* Ochsenheimer

C. canadensis Bethune. 4.

Genus *Capis* Grote

C. curvata Grote. 2. 3. 4.

Genus *Cissusa* Walker

C. spadix (Cramer). 4s: York (Toronto). Rare straggler.

Genus *Colobochyla* Hübner [= *Salia* Hübner]

C. interpuncta (Grote). 4.

Genus *Drasteria* Hübner [= *Synedoida* Hy. Edwards]

D. adumbrata (Behr). 3. 4.

D. alleni (Grote). 3. 4.

D. grandirena (Haworth). 4.

Genus *Hypocala* Guenée

H. andremona (Cramer). 4s: Essex (Point Pelee); Kent (Chatham). Occasional straggler.

Genus *Ledaea* Druce

L. perditalis (Walker). 4.

Genus *Melanomma* Grote

M. auricinctaria Grote. 4.

Genus *Melipotis* Hübner

M. indomita (Walker). 4s: Lambton; York (Toronto).

M. jucunda Hübner. 4s: Essex (Point Pelee).

Genus *Metalectra* Hübner

M. discalis (Grote). 4.

M. quadrisignata (Walker). 4.

Genus *Mycterophora* Hulst

M. inexplicata (Walker). 2. 3.

Genus *Pangrapta* Hübner

P. decoralis Hübner. 2. 3. 4.

Genus *Panopoda* Guenée

P. carneicosta Guenée. 4.

P. repanda (Walker). 4s: York (Toronto).

P. rufimargo (Hübner). 4.

Genus *Phoberia* Hübner

P. atomaris Hübner. 4n: Hastings (Marmora).

Genus *Phytometra* Haworth

P. rhodarialis (Walker) [= *semipurpurea* (Walker)]. 4s.

Genus *Plusiodonta* Guenée

P. compressipalpis Guenée. 4.

Genus *Rivula* Guenée

R. propinqualis Guenée. 3. 4.

Genus *Scolecocampa* Guenée

S. liburna (Geyer). 4s.

Genus *Scoliopteryx* Germar

S. libatrix (Linné). 2. 3. 4.

Genus *Spargaloma* Grote

S. sexpunctata Grote. 2. 3. 4.

Genus *Spiloloma* Grote

S. lunilinea (Grote). 4s: Lincoln (Vineland).

Genus *Thysania* Dalman

T. zenobia (Cramer). 4s: Essex (Point Pelee); Kent (Chatham). Occasional straggler.

Subfamily Catocalinae

Genus *Allotria* Hübner

A. elonympha (Hübner). 4s: Essex (Windsor).

Genus *Argyrostromis* Hübner

A. anilis (Drury). 4.

Genus *Caenurgina* McDunnough

C. crassiuscula Haworth. 1. 2. 3. 4.

C. erechtea (Cramer). 2. 4.

Genus *Catocala* Schrank

- C. amatrix* (Hübner). 4s.
- C. amica* (Hübner) [= *lineella* Grote]. 4s.
- C. antinympha* (Hübner). 3. 4.
- C. blandula* Hulst. 2. 3. 4.
- C. briseis* Edwards. 2. 3. 4.
- C. cara* Guenée. 4s.
- C. cerogama* Guenée. 3. 4.
- C. clintoni* Grote. 4s.
- C. coccinata* Grote. 4s.
- C. coelebs* Grote. 3. 4.
- C. concumbens* Walker. 3. 4.
- C. connubialis* Guenée. 4s: Lambton (Pinery Provincial Park).
- C. crataegi* Saunders. 4s.
- C. dejecta* Strecker. 4s: Essex (Windsor).
- C. epione* (Drury). 4.
- C. grynea* (Cramer). 4s.
- C. habilis* Grote. 4.
- C. ilia* (Cramer). 4s.
- C. illecta* Walker. 4s: Essex (Harrow, near Leamington).
- C. innubens* Guenée. 4.
- C. insolabilis* Guenée. 4s.
- C. judith* Strecker. 4s: Essex (Windsor); Haldimand (Caledonia); Halton (Brockville).
- C. lacrymosa* Guenée. 4s: Norfolk (Delhi).
- C. meskei* Grote. 4s.
- C. micronympha* Guenée. 4s.
- C. mira* Grote. 3. 4s.
- C. nebulosa* Edwards. 4s: Essex (Windsor); Haldimand (Caledonia).
- C. neogama* (J. E. Smith). 4s.
- C. obscura* Strecker. 4.
- C. palaeogama* Guenée. 4s.
- C. parta* Guenée. 4s.
- C. piatrix* Grote. 4.
- C. praeclara* Grote and Robinson. 4s.
- C. relictata* Walker. 2. 3. 4.
- C. residua* Grote. 4s: York (Toronto).
- C. relecta* Grote. 4s.
- C. semirelictata* Grote. 2. 3. 4.
- C. serena* Edwards. 4.
- C. similis* Edwards. 4.
- C. sordida* Grote. 3. 4.
- C. subnata* Grote. 4s.
- C. ultronia* (Hübner). 3. 4

C. unijuga Walker. 2. 3. 4.
C. vidua (J. E. Smith). 4s.

Genus *Celiptera* Guenée

C. frustulum Guenée. 4s: Middlesex (Strathroy).

Genus *Euclidia* Ochsenheimer

E. cuspidata (Hübner). 2. 3. 4.

Genus *Euparthenos* Grote

E. nubilis Hübner. 4.

Genus *Mocis* Hübner

M. latipes (Guenée). 4s.

Genus *Parallelia* Hübner

P. bistriaris Hübner. 4.

Genus *Zale* Hübner

Z. aeruginosa (Guenée). 2. 3. 4.
Z. calycanthata (J. E. Smith). 4: Manitoulin.
Z. duplicata (Bethune). Pine zale/phalène du pin. 2. 3. 4.
Z. galbanata (Morrison). 4.
Z. helata (Smith). 4.
Z. horrida Hübner. 4.
Z. lunata (Drury). 2. 3. 4.
Z. lunifera (Hübner) [= *cingulifera* (Walker)]. 4.
Z. metatoides McDunnough. 4s.
Z. minerea (Guenée). 2. 3. 4.
Z. phaeocapna Franclemont. 4s.
Z. submediana Strand. 3. 4.
Z. undularis (Drury). 4.
Z. unilineata (Grote). 3. 4.

Subfamily Plusiinae

Genus *Abrostola* Ochsenheimer

A. urentis Guenée. 2. 4.

Genus *Agrapha* Hübner

A. oxygramma (Geyer). 4s: Norfolk (Delhi); York (Toronto). Southern straggler.

Genus *Allagrapha* Franclemont

A. aerea (Hübner). 4.

Genus *Anagrapha* McDunnough

A. falcifera (Kirby). Celery looper/ autographe du céleri. 2. 3. 4.

Genus *Argyrogramma* Hübner

A. verruca (Fabricius). 4s: Welland (Niagara Falls).

Genus *Autographa* Hübner

A. ampla (Walker). 2. 3. 4.

A. biloba (Stephens). 2. 4.

A. bimaculata (Stephens). 2. 3. 4.

A. flagellum (Walker). 2. 3. 4.

A. mappa (Grote and Robinson). 2. 3.

A. precationis (Guenée). 2. 3. 4.

A. pseudogamma (Grote). 3. 4.

A. rubida Ottolengui. 2. 4.

Genus *Chrysanympha* Grote

C. formosa (Grote). 3. 4.

Genus *Diachrysia* Hübner

D. aeroides (Grote). 2. 3. 4.

D. balluca Geyer. 2. 3. 4.

Genus *Eosphoropteryx* Dyar

E. thyatyroides (Guenée). 4.

Genus *Plusia* Ochsenheimer

P. contexta Grote. 4.

P. putnami Grote. 2. 3. 4.

P. venusta Walker. 2. 3. 4.

Genus *Pseudeva* Hampson

P. purpurigera (Walker). 3. 4.

Genus *Pseudoplusia* McDunnough

P. includens (Walker). 4.

Genus *Rachiplusia* Hampson

R. ou (Guenée). 2. 3. 4.

Genus *Syngrapha* Hübner

S. abstrusa Eichlin and Cunningham. 3. 4.

S. alias (Ottolengui). Spruce climbing cutworm/autographe de l'épinette. 2. 3. 4.

S. altera (Ottolengui). 2. 3: Thunder Bay (Nipigon, type locality).

S. epigaea (Grote). 2. 3. 4.

S. ignea (Grote). 2: Cochrane (Low Bush).

S. microgamma (Hübner). 2. 3. 4. Subspecies *nearctica* Ferguson.

S. montana (Packard). 3. 4.

S. octoscripta Grote. 3. 4.

S. rectangula (W. Kirby). Angulated cutworm/autographe à rectangle. 2. 3.

S. selecta (Walker). 2: Cochrane (Ogoki Post).

S. surena (Grote). 2: Cochrane (Low Bush. Smoky Falls).

S. u-aureum (Guenée). 3: Algonquin Provincial Park.

S. viridisigma Grote. Verdigris autograph/autographe verdâtre. 2. 3. 4.

Genus *Trichoplusia* McDunnough

T. ni (Hübner). Cabbage looper/arpenreuse du chou. 2. 3. 4.

Subfamily Euteliinae

Genus *Eutelia* Hübner

E. pulcherrima (Grote). 4s: Kent (Rondeau Provincial Park).

Genus *Marathyssa* Walker

M. basalis Walker. 4.

M. inficita (Walker). 4.

Genus *Paectes* Hübner

P. abrostoloides (Guenée). 4s: Waterloo (Kitchener).

P. oculatrix (Guenée). 4s: Welland (Port Colborne).

Subfamily Sarrothripinae

Genus *Baileya* Grote

B. australis (Grote). 4s: Haldimand (Dunnville).

B. dormitans (Guenée). 4.

B. doubledayi (Guenée). 3. 4.

B. ophthalmica (Guenée). 4.

Genus *Nycteola* Hübner

N. cinereana Neumoegen and Dyar. 2.

N. frigidana (Walker). 2. 3. 4.

N. scriptana (Walker). 3: Sudbury.

Subfamily Nolinae

Genus *Meganola* Dyar

M. minuscula (Zeller). 4: Carleton; Frontenac; Lanark; Leeds; Middlesex; Muskoka; Peterborough; Russell; Wentworth; York. Early July. CNC, ROM, UG.

M. sp. 4: Hastings (Marmora only). CNC. See Forbes (1960:53).

Genus *Nola* Leach [= *Celama* Walker]

- N. cilicoides* (Grote). 4: Carleton; Frontenac; Hastings; Huron; Leeds; Norfolk; Wentworth. July. CNC, ROM.
N. clethrae Dyar. 4: Carleton; Wentworth. CNC.
N. ovilla Grote. 4: Carleton; Simcoe. CNC.
N. triquetra (Fitch). 4: Carleton; Kent; Nipissing; Norfolk; Wentworth; York. 21 May–16 June. CNC, ROM, UG. Specimen from Norfolk (Simcoe) is labelled, “bred from witch hazel.”

Subfamily Acontiinae

Genus *Acontia* Ochsenheimer

- A. aprica* (Hübner). 4s: Essex (Pelee Island); Kent (Chatham).
A. terminimaculata (Grote). 4.

Genus *Amyna* Guenée

- A. octo* (Guenée). 4n: Carleton (Ottawa).

Genus *Exyra* Grote

- E. rolandiana* Grote. 3. 4.

Genus *Homophoberia* Morrison

- H. apicosa* (Haworth). 4.
H. cristata Morrison. 4.

Genus *Hyperstrotia* Hampson

- H. pervertens* Barnes and McDunnough. 3. 4.
H. secta (Grote). 4.
H. villificans Barnes and McDunnough. 4n: Carleton (Ottawa).

Genus *Leuconycta* Hampson

- L. diptheroides* (Guenée). 3. 4.
L. lepidula (Grote). 4.

Genus *Lithacodia* Hübner

- L. albidula* (Guenée). 2. 3. 4.
L. bellicula Hübner. 2. 4.

L. carneola (Guenée). 2. 3. 4.

L. muscosula (Guenée). 4.

Genus *Maliatha* Walker

M. concinnimacula (Guenée). 4.

M. synochitis (Grote and Robinson). 3. 4.

Genus *Nigetia* Walker

N. formosalis Walker. 4.

Genus *Spragueia* Grote

S. leo (Grote). 4s: Middlesex (London).

Genus *Tarachidia* Hampson

T. binocula (Grote). 4s: Wentworth (Ancaster).

T. candefacta (Hübner). 4.

T. erastrioides (Guenée). 4.

T. tortricina (Zeller). 4.

Subfamily Pantheinae

Genus *Charadra* Walker

C. deridens (Guenée). 2. 3. 4.

Genus *Colocasia* Hampson

C. flavicornis (Smith). 3. 4.

C. infanta (Smith). 2.

C. propinquilinea (Grote). 2. 3. 4.

Genus *Panthea* Hübner

P. acronyctoides (Walker). Tufted spruce caterpillar/phalène de l'épinette. 2. 3.
4.

P. furcilla (Packard). 4.

Genus *Raphia* Hübner

R. frater Grote. 2. 3. 4.

Subfamily Acronictinae

Dagger moths.

Genus *Acronicta* Ochsenheimer

- A. afflicta* Grote. 3. 4.
A. albarufa Grote. 4s: Huron (Grand Bend).
A. americana (Harris). American dagger moth/acronycte d'Amérique. 2. 3. 4.
A. clarescens Guenée. 4.
A. connecta Grote. 4s: Wentworth (Ancaster)
A. dactylina Grote. 2. 3. 4.
A. distans (Grote). 2. 3. 4.
A. fragilis (Guenée). 2. 3. 4.
A. funeralis Grote and Robinson. 2. 3. 4.
A. grisea Walker. 2. 3. 4.
A. haesitata (Grote). 3. 4.
A. hamamelis Guenée. 3. 4.
A. hasta Guenée. 4.
A. impleta Walker. 2. 3. 4.
A. inclara Smith. 3. 4.
A. increta Morrison. 4s: Wentworth (Ancaster).
A. innotata Guenée. 2. 3. 4.
A. interrupta Guenée. 3. 4.
A. laetifica Smith. 4.
A. lanceolaria (Grote). 3. 4.
A. leporina (Linné). 2. 3. 4.
A. lepusculina Guenée. 2. 3. 4.
A. lithospila Grote. 4.
A. lobeliae Guenée. 2. 4s: Lambton.
A. longa Guenée. 2. 3. 4.
A. modica Walker. 4s.
A. morula Grote and Robinson. 3. 4.
A. noctivaga Grote. 2. 3. 4.
A. oblinita (J. E. Smith). Smeared dagger moth/acronycte souillée. 2. 3. 4.
A. ovata Grote. 4s.
A. quadrata Grote. 3.
A. radcliffei (Harvey). 3. 4.
A. retardata (Walker). 3. 4.
A. rubricoma Guenée. 4s: Essex (Point Pelee).
A. sperata Grote. 2. 3. 4.
A. spinigera Guenée. 4s: Wentworth (Hamilton); York (Toronto). Syntypes:
Lincoln (St. Catharines). AMNH. See Poole (1989:29).
A. superans Guenée. 3. 4.
A. tristis Smith. 4s: Huron (Grand Bend).
A. tritona (Hübner). 2. 3. 4.
A. vinnula (Grote). 4.

Genus *Agriopodes* Hampson

A. fallax (Herrich-Schäffer). 4.

Genus *Anterastris* Sugi

A. teratophora (Herrich-Schäffer). 4.

Genus *Harrisimemna* Grote

H. trisignata (Walker). 2. 3. 4.

Genus *Polygrammate* Hübner

P. hebraeicum Hübner. 4.

Genus *Simyra* Ochsenheimer

S. henrici (Grote). 2. 3. 4.

Subfamily Agaristinae

Genus *Alypia* Hübner

A. langtoni Couper. 2. 3. 4.

A. octomaculata (Fabricius). Eight-spotted forester/ alypie à huit points. 4.

Genus *Androloma* Grote

A. maccullochii (Kirby). 2. 3.

Genus *Eudryas* Boisduval

E. grata (Fabricius). 4.

E. unio (Hübner). 4.

Genus *Psychomorpha* Harris

P. epimenis (Drury). 4s: Essex (Ojibway Park, 1987. Point Pelee*. Seen on Pelee Island); Halton (Campbellville); Waterloo (Kitchener).

Subfamily Amphipyrinae

Genus *Achatodes* Guenée

A. zae (Harris). 4.

Genus *Amolita* Grote

A. fessa Grote. 4.

Genus *Amhipoea* Billberg

A. americana (Speyer). 2. 3. 4.

A. interoceanica (Smith). 4s: Lambton; Wentworth (Ancaster).

A. velata (Walker). 3. 4.

Genus *Amhipyra* Ochseneimer

A. glabella (Morrison). 3. 4.

A. pyramidoides Guenée. 4.

A. tragopoginis (Clerck). 4.

Genus *Andropolia* Grote

A. aedon (Grote). 2. 3.

A. contacta (Walker). 3.

Genus *Anorthodes* Smith

A. tarda (Guenée). 2: Thunder Bay (Geraldton).

Genus *Apamea* Ochseneimer [= *Trichoplexia* Hampson]

A. alia (Guenée). 2. 3. 4.

A. amputatrix (Fitch). 2. 3. 4.

A. apamiformis (Guenée). 3. 4.

A. cariosa (Guenée). 4.

A. cogitata (Smith). 2.

A. commoda (Walker). 3.

A. devastator (Brace). Glassy cutworm/ver-gris vitreux. 2. 3. 4.

A. dubitans (Walker). 2. 3. 4.

A. extensa (Smith). 4n: Carleton (Ottawa).

A. finitima Guenée. 2. 3. 4.

A. helva (Grote). 4.

A. impulsiva (Guenée). 2. 3. 4.

- A. indocilis* (Walker). 2. 3. 4.
A. inficita (Walker). 3.
A. inordinata (Morrison). 4s: Lincoln (Queenston); Peterborough*.
A. lateritia (Hufnagel). 2. 3. 4.
A. lignicolora (Guenée). 4.
A. lutosa (Andrews). 4s.
A. mixta (Grote). 3: Sudbury (Westree).
A. nigrior (Smith). 4.
A. niveivenosa (Grote). 2. 3. 4.
A. plutonia (Grote). 3. 4.
A. verbascoides (Guenée). 3. 4.
A. vulgaris (Grote and Robinson). 4s: Lambton.
A. vultuosa (Grote). 3. 4.

Genus *Archanara* Walker

- A. laeta* (Morrison). 4n: Carleton (Mer Bleue).
A. oblonga (Grote). 3. 4.
A. subflava (Grote). 2. 3. 4.

Genus *Athetis* Hübner [= *Proxenus* Herrich-Schäffer]

- A. mendosa* (McDunnough). 2: Cochrane (Smoky Falls).
A. miranda (Grote). 2. 3. 4.

Genus *Balsa* Walker

- B. labecula* (Grote). 4.
B. malana (Fitch). 4.
B. tristrigella (Walker). 4.

Genus *Bellura* Walker

- B. densa* (Walker). 4s: Hastings (Trenton).
B. gortynoides Walker. 2. 3. 4.
B. obliqua (Walker). 2. 3. 4.

Genus *Callopistria* Hübner

- C. cordata* (Ljungh) [= *montifera* (Guenée)]. 3. 4.
C. mollissima (Guenée). 4.

Genus *Caradrina* Ochsenheimer

C. extima (Walker). 3: Sudbury.

C. meralis (Morrison). 4s: Frontenac (Perth Road); Hastings (Trenton).

C. morpheus (Hufnagel). 4n: Carleton (Ottawa).

C. multifera (Walker). 3. 4.

Genus *Catabena* Walker

C. lineolata Walker. 4.

Genus *Celaena* Stephens [= *Helotropha* Lederer]

C. reniformis (Grote). 2. 3. 4.

Genus *Cerma* Hübner

C. cerintha (Treitschke). 3. 4.

C. cora Hübner. 3. 4.

Genus *Chytonix* Grote

C. palliatricula (Guenée). 3. 4.

C. ruperti Franclemont. 4s: Huron (Grand Bend).

C. sensilis Grote. 4.

Genus *Cirrophanus* Grote

C. triangulifer Grote. 4s: Kent (Rondeau Provincial Park); Middlesex (London).

Genus *Condica* Walker [= *Platysenta* Grote]

C. sutor (Guenée). 4s: Wentworth (Ancaster).

C. vecors (Guenée). 4.

C. videns (Guenée). 4.

Genus *Conservula* Grote

C. anodonta (Guenée). 4.

Genus *Cosmia* Ochsenheimer

C. calami (Harvey). 3. 4.

Genus *Crambodes* Guenée

C. talidiformis Guenée. 4.

Genus *Dypterygia* Stephens

D. rozmani Berio [= *scabriuscula* (Linné)]. 3. 4.

Genus *Elaphria* Hübner

E. festivoides (Guenée). 2. 3. 4.

E. grata Hübner. 4s: Hastings (Trenton).

E. versicolor (Grote). 4.

Genus *Enargia* Hübner

E. decolor (Walker). 2. 3. 4.

E. infumata (Grote). 2. 3. 4.

E. mephisto Franclemont. 4s.

Genus *Euplexia* Stephens

E. benesimilis McDunnough. 2. 3. 4.

Genus *Fagitana* Walker

F. littera (Guenée). 4n: Carleton (Ottawa).

Genus *Galgula* Guenée

G. partita Guenée. 2. 3. 4.

Genus *Hydraecia* Guenée

H. immanis Guenée. Hop vine borer/perceur du houblon. 4.

H. micacea (Esper). Potato stem borer/perce-tige de la pomme de terre. 4.

H. stramentosa Guenée. 4.

Genus *Hypocoena* Hampson

H. basistriga McDunnough. 3: Thunder Bay.

H. defecta (Grote). 3. 4.

Genus *Hyppa* Duponchel

H. xylinoides (Guenée) [= *ancociscònensis* (Morrison)]. 3. 4.

Genus *Ipimorpha* Hübner

I. pleonectusa Grote. 2. 3. 4.

Genus *Lemmeria* Barnes and Benjamin

L. digitalis (Grote). 4.

Genus *Luperina* Boisduval

L. passer (Guenée). 2. 3. 4.

Genus *Macronoctua* Grote

M. onusta Grote. Iris borer/perceur de l'iris. 4.

Genus *Magusa* Walker

M. orbifera (Walker). 4.

Genus *Meropleon* Dyar

M. diversicolor (Morrison). 3. 4.

Genus *Nedra* Clarke

N. ramosula (Guenée). 4.

Genus *Ogdoconta* Butler

O. cinereola (Guenée). 4.

Genus *Oligia* Hübner

O. bridghami (Grote and Robinson). 3. 4.

O. chlorostigma (Harvey). 4s: Frontenac (Perth Road); Lambton*.

O. exhausta (Smith). 4n: Carleton (Dunrobin).

O. fractilinea (Grote). 3. 4.

O. illocata (Walker). 2. 3. 4.

O. mactata (Guenée). 3. 4.

O. marina (Grote). 2.

O. minuscula (Morrison). 4.

O. modica (Guenée). 3. 4.

O. obtusa (Smith). 4s: Welland (Port Colborne).

O. semicana (Walker). 4.

O. tonsa (Grote). 3. 4.

Genus *Papaipema* Smith

- P. apassionata* (Harvey). 4.
P. arctivorens Hampson. 4s.
P. astuta Bird. 2: Thunder Bay (Geraldton)
P. aweme (Lyman). 4s: Huron (Grand Bend).
P. cataphracta (Grote). Burdock borer/perce-tige de la bardane. 4.
P. cerina (Grote). 4s: Norfolk (Delhi).
P. cerrusata (Grote). 4s: Essex (Windsor area, J. Larson).
P. duovata (Bird). 4.
P. eupatorii (Lyman). 4.
P. furcata (Smith). 4.
P. harrisii (Grote). 3. 4.
P. impecuniosa (Grote). 4.
P. inquaesita (Grote and Robinson). 4.
P. leucostigma (Harris). Columbine borer/perce-tige de l'ancolie. 4.
P. lysimachiae Bird. 4s: Welland (Port Colborne).
P. marginidens (Guenée). 4.
P. nebris (Guenée). 4.
P. necopina (Grote). 4s: Kent (Chatham).
P. nelita (Strecker). 4.
P. nepheleptena (Dyar). 4n: Carleton (Ottawa). 4s: Norfolk (Delhi).
P. pterisii Bird. 3. 4.
P. speciosissima (Grote and Robinson). 4n: Carleton (Ottawa). 4s: Leeds (Chaffey's Locks).
P. unimoda (Smith) [= *frigida* (Smith)]. 3. 4.

Genus *Perigea* Guenée

- P. xanthioides* Guenée. 4s: Essex (Windsor); Kent (Chatham).

Genus *Phlogophora* Treitschke

- P. iris* Guenée. 2. 3. 4.
P. periculosa Guenée. 2. 3. 4.

Genus *Phosphila* Hübner

- P. miselioides* (Guenée). 4s: Essex (Windsor); Norfolk (Delhi).

Genus *Plagiomimicus* Grote

- P. pityochromus* Grote. 4s: Essex (Windsor).

Genus *Spartiniphaga* McDunnough

S. includens (Walker). 4.

S. panatala (Smith). 4.

Genus *Spodoptera* Guenée

S. exigua (Hübner). 4

S. ornithogalli (Guenée). 4.

Genus *Trachea* Ochsenheimer

T. delicata (Grote). 4.

Genus *Xylomoia* Staudinger

X. chagnoni Barnes and McDunnough. 4.

Subfamily Cuculliinae

Genus *Adita* Grote

A. chionanthi (J. E. Smith). 4.

Genus *Anathix* Franclemont

A. puta (Grote and Robinson). 2: Cochrane (Ogoki Post).

A. ralla (Grote and Robinson). 4.

Genus *Apharetra* Grote

A. dentata (Grote). 2: Thunder Bay (Geraldton). 3. 4: Carleton (Mer Bleue, type locality of the synonym *purpurea* McDunnough, see Poole, 1989:111).

Genus *Blepharita* Hampson [= *Mniotype* Franclemont]

B. miniota (Smith). 2. 3.

Genus *Brachionycha* Hübner

B. borealis (Smith). 3: Thunder Bay (Black Sturgeon Lake).

Genus *Brachylomia* Hampson

B. disciniga (Walker). 2: Cochrane (Ogoki Post).

Genus *Chaetagnaea* Franclemont

C. cerata Franclemont. 4s: Leeds (Chaffey's Locks).

C. sericea (Morrison). 4n: Carleton (Ottawa).

Genus *Copivaleria* Grote

C. grotei (Morrison). 4s.

Genus *Cucullia* Schrank

C. asteroides Guenée. 4.

C. convexpennis Grote and Robinson. Aster cutworm/cucullie de l'aster. 4.

C. florea Guenée. 2. 3. 4.

C. lucifuga (Denis and Schiffermüller) [= *intermedia* Speyer]. Goldenrod cutworm/cucullie de la verge d'or. 2. 3. 4.

C. omissa Dod. 2. 3. 4.

C. postera Guenée. 4n: Carleton (Ottawa).

C. speyeri Lintner. 4s.

Genus *Epignaea* Grote

E. apiata (Grote). 3. 4.

E. decliva (Grote). 4s.

Genus *Eucirroedia* Grote

E. pampina (Guenée). 3. 4.

Genus *Eupsilia* Hübner

E. devia (Grote). 4.

E. morrisoni (Grote). 4.

E. sidus (Guenée). 4.

E. tristigmata (Grote). 3. 4.

E. vinulenta (Grote). 4.

Genus *Feralia* Grote

F. comstocki (Grote). 2. 3. 4.

F. jocosus (Guenée). Green caterpillar/chenille à raies vertes. 4.

F. major Smith. 4.

Genus *Fishia* Grote

F. enthea Grote. 2. 3. 4.

Genus *Hillia* Grote

H. iris (Zetterstedt). 2. 4.

Genus *Homoglaea* Morrison

H. hircina Morrison. 2. 4.

Genus *Homohadena* Grote

H. badistriga (Grote). Honeysuckle budworm/phalène du chèvrefeuille. 3. 4.

H. infixa (Walker). 2. 3. 4.

Genus *Litholomia* Grote

L. napaea (Morrison). 2. 3. 4.

Genus *Lithomoia* Hübner

L. solidaginis (Hübner). 2: Cochrane (Ogoki Post).

Genus *Lithophane* Hübner

L. amanda (Smith). 2. 3. 4.

L. antennata (Walker). Green fruitworm/phalène du fruit vert. 4.

L. baileyi Grote. 3. 4.

L. bethunei (Grote and Robinson). 4.

L. disposita Morrison. 4.

L. fagina Morrison. 4.

L. georgii Grote. 2. 3. 4.

L. grotei Riley. 4.

L. hemina Grote. 4.

L. innominata (Smith). 3. 4.

L. laticinerea Grote. 4.

L. lemmeri (Barnes and Benjamin). 4: Frontenac (Arden).

L. lepida Grote. 3. 4.

L. oriunda Grote. 4.

L. patefacta (Walker). 4.

L. petulca Grote [= *ferrealis* Grote]. 3. 4.

L. pexata Grote. 2. 3. 4.

L. querquera Grote. 4s: Frontenac (Perth Road).

L. semiusta Grote. 4.

- L. tepida* Grote. 2: Timiskaming (Larder Lake). 4s: Frontenac (Perth Road).
L. thaxteri Grote. 2. 3. 4.
L. unimoda (Lintner). 3. 4.

Genus *Metaxaglaea* Franclemont

- M. inulta* (Grote). 4.

Genus *Oncocnemis* Lederer

- O. piffardi* (Walker). 2. 3.
O. riparia Morrison. 4s.
O. saundersiana Grote. 4.
O. viriditincta Smith. 4s: Leeds (Chaffey's Locks); Wentworth (Hamilton).

Genus *Parastichtis* Hübner

- P. discivaria* (Walker). 2. 4.

Genus *Platypolia* Grote

- P. anceps* Stephens. 3. 4.

Genus *Properigea* Barnes and Benjamin

- P. costa* Barnes and Benjamin. 4s: Frontenac (Perth Road); Leeds (Chaffey's Locks).

Genus *Psaphida* Walker

- P. electilis* (Morrison). 4s.
P. rolandi Grote. 4s: Kent (Chatham).
P. styracis (Guenée). 4s.
P. thaxteriana (Grote). 4s: Norfolk (Delhi).

Genus *Psectraglaea* Hampson

- P. carnosa* (Grote). 4s.

Genus *Pyreferra* Franclemont

- P. ceromatica* (Grote). 4s.
P. citromba Franclemont. 3.
P. hesperidago (Guenée). 4.
P. pettiti (Grote). 4.

Genus *Sunira* Franclemont

S. bicolorago (Guenée). 4.

Genus *Sutyna* Todd

S. privata (Walker). 2. 4.

S. profunda (Smith). 2.

Genus *Sympistis* Hübner

S. zetterstedtii (Staudinger) [= *kolthoffi* (Aurivillius)]. 1: Kenora (Cape Henrietta Maria).

Genus *Xanthia* Ochsenheimer

X. togata (Esper) [= *lutea* (Ström)]. 2. 3. 4.

Genus *Xylena* Ochsenheimer

X. cineritia (Grote). 2. 3. 4.

X. curvimacula (Morrison). 3. 4.

X. nupera (Lintner). 3. 4.

X. thoracica (Putnam-Cramer). 2. 3.

Genus *Xylotype* Hampson

X. acadia Barnes and Benjamin. 2. 3.

Subfamily Hadeninae

Genus *Achatia* Hübner

A. distincta Hübner. 4.

Genus *Aletia* Hübner

A. oxygala (Grote). 2. 3. 4.

Genus *Anarta* Ochsenheimer

A. cordigera (Thunberg) [= *luteola* (Grote and Robinson)]. 4n.

A. melanopa (Thunberg). 1: Kenora (Fort Severn).

Genus *Anartomima* Boursin

A. secedens (Walker). 3: Thunder Bay (Black Sturgeon Lake).

Genus *Anhimella* McDunnough

A. contrahens (Walker). 2. 3. 4.

Genus *Crocigrapha* Grote

C. normani (Grote). Climbing cherry cutworm/phalène du cerisier. 3. 4.

Genus *Discestra* Hampson

D. trifolii (Hufnagel). Clover cutworm/ver-gris du trèfle. 2. 3. 4.

Genus *Egira* Duponchel

E. alternans (Walker). 4s: Leeds (Chaffey's Locks).

E. dolosa (Grote). 3. 4.

E. purpurea (Barnes and McDunnough). 3.

Genus *Faronta* Smith

F. diffusa (Walker). Wheat head armyworm/légionnaire des épis de blé. 2. 3. 4.

Genus *Hadena* Schrank [= *Anepia* Hampson]

H. capsularis (Guenée). 4s: Lambton.

Genus *Homorthodes* McDunnough

H. furfurata (Grote). 3. 4.

Genus *Lacanobia* Billberg

L. atlantica (Grote). 2. 3. 4.

L. nevadae (Grote). 2. 3. 4.

L. radix (Walker). 2. 3. 4.

L. subjuncta (Grote and Robinson). 4.

Genus *Lacinipolia* McDunnough

L. anguina (Grote). 2. 3. 4.

L. implicata McDunnough. 4s: Leeds (Chaffey's Locks).

- L. lorea* (Guenée). 2. 3. 4.
L. lustralis (Grote). 3. 4.
L. meditata (Grote). 4.
L. olivacea (Morrison). 2. 3. 4.
L. renigera (Stephens). 2. 3. 4.
L. vicina (Grote). 2. 3. 4.

Genus *Lasionycta* Aurivillius

- L. albinuda* (Smith). 3: Thunder Bay (Black Sturgeon Lake).

Genus *Leucania* Ochsenheimer

- L. commoides* Guenée. 2. 3. 4.
L. inermis (Forbes). 4.
L. insueta Guenée. 2. 4.
L. multilinea Walker. 2. 3. 4.
L. phragmatidicola Guenée. 3. 4.
L. pseudargyria Guenée. 4.
L. ursula (Forbes). 4s.

Genus *Mamestra* Ochsenheimer

- M. curialis* (Smith). 3. 4.

Genus *Melanchra* Hübner

- M. adjuncta* (Boisduval). 2. 3. 4.
M. assimilis (Morrison). 3. 4.
M. picta (Harris). Zebra caterpillar/chenille zébrée. 4.
M. pulverulenta (Smith). 2. 3.

Genus *Morrisonia* Grote

- M. confusa* (Hübner). 4.
M. evicta (Grote). 4.
M. latex (Guenée). 2. 4.

Genus *Nephelodes* Hübner

- N. minians* Guenée. 2. 3. 4.

Genus *Orthodes* Guenée

- O. crenulata* (Butler). 3. 4.

- O. cynica* Guenée. 2. 3. 4.
O. goodelli (Grote). 4.
O. obscura (Smith). 2. 3. 4.

Genus *Orthosia* Ochsenheimer

- O. alurina* (Smith). 4n: Carleton (Dunrobin). 4s: Leeds (Chaffeys Locks).
O. hibisci (Guenée). 2. 3. 4.
O. revicta (Morrison). 2. 3. 4.

Genus *Papestra* Sukhareva

- P. cristifera* (Walker). 2. 3.
P. quadrata (Smith) [= *ingravis* Smith]. 3.

Genus *Polia* Ochsenheimer

- P. detracta* (Walker). 2. 3.
P. imbrifera (Guenée). 2. 3. 4.
P. nimbose (Guenée). 3. 4.
P. purpurissata (Grote). 2. 3. 4.
P. rogenhoferi (Möschler) [= *carbonifera* Hampson; = *leomegra* (Smith)]. 2:
 Cochrane (Ogoki Post). 3: Thunder Bay (Black Sturgeon Lake).

Genus *Protorthodes* McDunnough

- P. incincta* (Morrison). 4s: Essex (Windsor).
P. oviduca (Guenée). 2. 3. 4.

Genus *Pseudaletia* Franclemont

- P. unipuncta* (Haworth). Armyworm/légionnaire uniponctuée. 2. 3. 4.

Genus *Pseudorthodes* McDunnough

- P. vecors* (Guenée). 2. 3. 4.

Genus *Sideridis* Hübner

- S. congermana* (Morrison). 4.
S. maryx (Guenée). 2. 3.
S. rosea (Harvey). 2. 3. 4.

Genus *Spiramater* McCabe

S. grandis (Guenée). 2. 3. 4.

S. lutra (Guenée). 2. 3. 4.

Genus *Stretchia* Hy. Edwards

S. plusiiformis Hy. Edwards. 2. 3.

Genus *Trichoclea* Grote

T. artesta (Smith). 4s: Kent (Chatham); Welland (Port Colborne).

Genus *Trichodestra* McCabe

T. legitima (Grote). 2. 3. 4.

T. lilacina (Harvey). Flax cutworm/ver-gris du lin. 3. 4.

T. rugosa (Morrison). 4.

T. tacoma (Strecker). 2. 3.

Genus *Tricholita* Grote

T. signata (Walker). 3. 4.

Genus *Ulolonche* Smith

U. culea (Guenée). 3. 4.

U. modesta (Morrison). 4s.

Subfamily Noctuidae

Tribe Agrotini

Genus *Abagrotis* Smith

A. alternata (Grote). 3. 4.

A. barnesi (Benjamin). 4s: Northumberland (Presqu'île Provincial Park).

A. placida (Grote). 2: Thunder Bay (Geraldton).

Genus *Agrotis* Ochsenheimer

A. gladiaria Morrison. Clay-backed cutworm/ver-gris argile. 4.

A. ipsilon (Hufnagel). Black cutworm/ver-gris noir. 2. 3. 4.

A. mollis Walker. 3.

A. obliqua (Smith). 2. 3. 4.

A. stigmosa Morrison. 4.

- A. venerabilis* Walker. 4.
A. vetusta Walker. 3. 4.
A. volubilis Harvey. 2. 3. 4.

Genus *Anaplectoides* McDunnough

- A. prasina* (Denis and Schiffermüller). 2. 3. 4.
A. pressus (Grote). 2. 3. 4.

Genus *Aplectoides* Butler

- A. condita* (Guenée). 2. 3.

Genus *Cerastis* Ochsenheimer

- C. tenebrifera* (Walker). 4.

Genus *Chersotis* Boisduval

- C. juncta* (Grote). 2: Thunder Bay (Geraldton).

Genus *Choephora* Grote and Robinson

- C. fungorum* Grote and Robinson. 4s: Kent (Chatham); Welland (Port Colborne).

Genus *Cryptocala* Benjamin

- C. acadensis* (Bethune). 2. 3. 4.

Tribe Aniclini

Genus *Diarsia* Hübner

- D. dislocata* (Smith). 3.
D. freemani Hardwick. 2.
D. jucunda (Walker). 2. 3. 4.
D. rubifera (Grote). 2. 3. 4.

Genus *Euagrotis* McDunnough

- E. forbesi* Franclemont. 4.
E. illapsa (Walker). 3. 4.

Genus *Eueretagrotis* Smith

E. attenta (Grote). 2. 3. 4.

E. perattenta (Grote). 2. 3. 4.

E. sigmoides (Guenée). 4.

Genus *Eugraphe* Hübner

E. subrosea (Stephens) [= *opacifrons* (Grote)]. 2. 3. 4.

Genus *Eurois* Hübner

Generic name is of masculine gender.

E. astrictus Morrison. 2. 3. 4.

Genus *Euxoa* Hübner

Subgenus *Longivesica* Hardwick

E. divergens (Walker). 2. 3. 4.

E. messoria (Harris). Dark-sided cutworm/ver-gris moissonneur. 4.

E. sinelinea Hardwick. 2. 3.

Subgenus *Pleonectopoda* Grote

E. aurulenta (Smith). 4s: Welland (Port Colborne).

E. churchillensis (McDunnough). 1: Kenora (Cape Henrietta Maria).

E. pleuritica (Grote). 3. 4.

E. quebecensis (Smith). 2. 3.

E. scandens (Riley). White cutworm/ver-gris blanc. 2. 3. 4.

Subgenus *Chorizagrotis* Smith

E. lidia (Cramer). 4n: Carleton (Ottawa).

Subgenus *Euxoa* Hübner

E. albipennis (Grote). 4.

E. bostoniensis (Grote). 4.

E. campestris (Grote). 3. 4.

E. comosa (Morrison) [= *ontario* (Smith)]. 2. 3.

E. declarata (Walker). 3. 4.

E. detersa (Walker). Sand cutworm/ver-gris arénicole. 4.

E. fumalis (Grote). 4.

E. mimallonis (Grote). 3. 4.

- E. obeliscoides* (Guenée). 4.
E. ochrogaster (Guenée). Red-backed cutworm/ver-gris à dos rouge. 2. 3. 4.
E. redimicula (Morrison). 4.
E. scholastica McDunnough. 4.
E. servita (Smith). 3. 4.
E. tessellata (Harris). Striped cutworm/ver-gris rayé. 3. 4.
E. velleripennis (Grote). 4.

Subgenus *Orosagrotis* Hampson

- E. manitobana* McDunnough. 4s: Huron (Grand Bend).
E. perpolita (Morrison). 2. 3. 4.
E. ridingsiana (Grote). 2: Thunder Bay (Geraldton).

Genus *Feltia* Walker

- F. geniculata* Grote and Robinson. 2. 3. 4.
F. herilis (Grote). 3. 4.
F. jaculifera (Guenée) [= *ducens* (Walker)]. 3. 4.
F. subgothica (Haworth). Dingy cutworm/ver-gris terne. 4.
F. tricola (Lintner). 4s.

Genus *Graphiphora* Ochseneheimer

- G. haruspica* (Grote). 2. 3. 4.

Genus *Hemipachnobia* McDunnough

- H. monochromatea* (Morrison). 2. 4.

Genus *Heptagrotis* McDunnough

- H. phyllophora* (Grote). 2. 3. 4.

Genus *Loxagrotis* McDunnough

- L. sp.* 4: Manitouslin.

Genus *Metalepsis* Grote

- M. fishii* (Grote). 3: Thunder Bay (Black Sturgeon Lake). 4s: Frontenac (Perth Road).
M. salicarum (Walker). 2. 3. 4.

Genus *Ochropleura* Hübner

- O. fennica* (Tauscher). Black armyworm/légionnaire noire. 2. 3. 4.
O. plecta (Linné). 2. 3. 4.

Genus *Paradiarsia* McDunnough

- P. littoralis* (Packard). 2. 3. 4.

Genus *Peridroma* Hübner

- P. saucia* (Hübner). Variegated cutworm/ver-gris panaché. 2. 3. 4.

Genus *Protolampra* McDunnough

- P. brunneicollis* (Grote). 4.
P. rufipectus (Morrison). 2. 3. 4.

Genus *Rhynchagrotis* Smith

- R. anchocelioides* (Guenée). 4n: Carleton (Ottawa). 4s: Niagara Peninsula.
R. cupida (Grote). 2: Thunder Bay (Geraldton).

Genus *Spaelotis* Boisduval

- S. clandestina* (Harris). W-marked cutworm/ver-gris clandestin. 2. 3. 4.

Genus *Ufeus* Grote

- U. satyricus* Grote. 2. 3. 4.

Genus *Xestia* Hübner [= *Anomogyna* Staudinger]

- X. adela* Franclemont. Spotted cutworm/ver-gris tacheté. 2. 3. 4.
X. atrata (Morrison). 2. 3.
X. badicollis (Grote). 3. 4.
X. badinodis (Grote). 4.
X. bicarnea (Guenée). 3. 4.
X. bugrai Kocak. 2. 3. 4.
X. dilucida (Morrison). 4s: Frontenac (Perth Road).
X. dolosa Franclemont. 4s.
X. fabulosa Ferguson. 3: Thunder Bay (Pukaskwa National Park).
X. normaniana (Grote). 3. 4.
X. oblata (Morrison). 2. 3.
X. perquiritata (Morrison). Gray spruce cutworm/phalène grise de l'épinette. 3.
X. smithii (Snellen). 2. 3. 4.

- X. speciosa* (Hübner). 2.
X. tenuicula (Morrison). 2. 3. 4.
X. youngii (Smith). 4.

Subfamily Heliothinae

Genus *Eutricopis* Morrison

- E. nexilis* Morrison. 3. 4.

Genus *Helicoverpa* Hardwick

- H. zea* (Boddie). Corn earworm, tomato fruitworm/ver de l'épi du maïs, noctuelle de la tomate. 2. 4.

Genus *Heliothis* Ochsenheimer

- H. borealis* (Hampson). 2.
H. paradoxus Grote. 4s: Kent (Chatham).
H. phloxiphagus Grote and Robinson. 4.
H. virescens (Fabricius). Tobacco budworm/phalène verdoyante. 4s: Kent (Chatham).

Genus *Pyrria* Hübner

- P. exprimens* (Walker). 2. 3. 4.
P. umbra (Hufnagel). Rose budworm/pique-bouton du rosier. 3*. 4.

Genus *Rhodoccia* Hampson

- R. aurantiago* (Guenée). 4s: Huron (Grand Bend); Norfolk (Delhi).

Genus *Schinia* Hübner

- S. arcigera* (Guenée). 4.
S. florida (Guenée). 3. 4.
S. gloriosa (Strecker). 4s: Lambton (Pinery Provincial Park).
S. lynx (Guenée). 4.
S. meadi (Grote). 3: Rainy River (Emo).
S. nundina (Drury). 4s: Kent (Chatham).
S. obscurata Strecker. 4.
S. rivulosa (Guenée). 3. 4.
S. thoreauui (Grote and Robinson). 4s: Kent (Chatham).
S. trifascia Hübner. 4.

ACKNOWLEDGEMENTS

I would like to express sincere thanks to the institutions that made their collections available for the compilation of this list, and to the curators who provided all possible help for the study of their material: the late H. K. Clench (CM), D. M. Davies (McM), T. N. Freeman (then CNC), D. F. Hardwick (then CNC), W. W. Judd (UWO), A. B. Klots (AMNH), J. D. Lafontaine (CNC), W. C. McGuffin (then CNC), G. A. Moore (Lyman College), E. G. Munroe (then CNC), D. H. Pengelly (then UG), W. L. Sippell (FIDS), V. R. Vickery (LEM), A. S. West (QU), and D. M. Wood (then McM). Special thanks are extended to the following people: F. H. Rindge (AMNH), who identified thousands of specimens of Geometridae and Noctuidae collected by the author, and who checked and rechecked the Geometridae section; E. G. Munroe, who provided valuable advice over many years of work connected with this list (the section on the Pyralidae is virtually his); and P. R. Ehrlich, D. D. Murphy, and P. A. Opler for advice and assistance on the Papilionoidea section. Special thanks are extended also to the Lepidoptera section of Biosystematics Research Centre, in particular, J. D. Lafontaine, P. T. Dang, K. Bolte, B. Landry, and J.-F. Landry. They provided recent information on the distribution of Ontario Lepidoptera during the final stages of the preparation of the manuscript (in 1990).

Special acknowledgement must be given to the Department of Entomology, Royal Ontario Museum, especially to curators G. B. Wiggins, and D. Barr for their interested participation in the development of this publication, and for making possible the necessary fieldwork of the author, supported by the following summer student volunteers: I. Smith, P. Hébert, L. Kohalmi, L. Luciak, and P. Ward. Mrs. E. A. Perry-Easley, my field assistant for summer 1975, provided substantial help in the reorganization of the contents of this list from research and field notes. Nick Tzovolos (ROM) provided new county records and flight periods by researching various publications and by examining recently donated specimens in the ROM collection, and as well provided annotations and editorial work. The maps were produced by Zile Zichmanis and Patricia Stephens-Bourgeault of the ROM Department of Entomology; Figures 2 and 3 are based on maps provided by R. D. James (Department of Ornithology, ROM). Mr. E. R. Fuller (then ROM) deserves special thanks for transforming the entire list from the format of the McDunnough (1938, 1939) checklist to the format of the Moths of North America series. Very substantial editorial and secretarial contributions from Roslyn Darling and Susan Pasch are acknowledged with gratitude. And last but not at all least, very special thanks go to D. C. Darling, Department of Entomology, Royal Ontario Museum, who can very properly be called the *spiritus rector* of the final version of the checklist.

LITERATURE CITED

ARTHUR, A. P.

- 1962 A skipper, *Thymelicus lineola* (Ochs.) (Lepidoptera: Hesperiiidae) and its parasites in Ontario. *Canadian Entomologist* 94:1082-1089.
- 1966 The present status of the introduced skipper, *Thymelicus lineola* (Ochs.) (Lepidoptera: Hesperiiidae), in North America and possible methods of control. *Canadian Entomologist* 98:622-626.

BAKER, A. W.

- 1955 Changing faunal ranges. Annual Report of the Entomological Society of Ontario 86:23-29.

BENOIT, P., ed.

- 1975 Noms français d'insectes au Canada avec noms latins et anglais correspondants. 4ième éd. Québec, Agriculture Québec. 214 pp.

BETHUNE, C. J. S.

- 1894 The butterflies of the eastern provinces of Canada. Annual Report of the Entomological Society of Ontario 1894:29-44.

BURNS, J. M.

- 1964 Evolution in skipper butterflies of the genus *Erynnis*. University of California Publications in Entomology 37:1-214.
- 1966 Expanding distribution and evolutionary potential of *Thymelicus lineola* (Lepidoptera: Hesperiiidae), an introduced skipper, with special reference to its appearance in British Columbia. *Canadian Entomologist* 98:859-866.

CANADA DEPARTMENT OF TRANSPORT, METEOROLOGICAL DIVISION

- 1947 Climatic summaries for selected meteorological stations in the Dominion of Canada. 1. Ottawa. 63 pp.
- 1954 Addendum to volume I of climatic summaries for selected meteorological stations in Canada. Ottawa. 22 pp.
- 1964 Temperature normals for Ontario. Ottawa. 10 pp.

CANADIAN PERMANENT COMMITTEE ON GEOGRAPHICAL NAMES

- 1974 Gazetteer of Canada: Ontario. 2nd ed. Ottawa, Energy, Mines and Resources Canada. 823 pp.

CHUA, K. E., J. C. E. RIOTTE, and C. GILMORE

- 1976 Investigation of selected species of the genus *Orgyia* (Lymantriidae) using isoelectrofocusing in thin layer polyacrylamide gel. *Journal of Research on the Lepidoptera* 15:215-224.

CLAYTON, W. J.

- 1938 Description of and notes on the early stages of *Hyloicus canadensis* Bdv. (Lepidoptera: Sphingidae). *Entomological News* 49:69-72.

D'ABRERA, B.

- 1986 Sphingidae Mundi, Hawk Moths of the world, based on a checklist by Alan Hayes and the collection he curated in the British Museum (Nat. Hist.). London, E. W. Classey. 226 pp.

EHRlich, P. R.

- 1958 The comparative morphology, phylogeny and higher classification of the butterflies (Lep. Papilionoidea). University of Kansas Science Bulletin 39:305-370.

EHRlich, P. R. and A. H. EHRlich

- 1961 How to know the butterflies. Dubuque, W. C. Brown. 262 pp.

EITSCHBERGER, U. and H. STEINIGER

- 1983 Systematische Untersuchungen am *Pieris napi* - *bryoniae* Komplex (Lepidoptera: Pieridae). Herbiopoliana 1(1), 504 pp., 1(2), 601 pp. Published by the authors.

ENGELHARDT, G. P.

- 1946 The North American clear-wing moths of the family Aegeriidae. United States National Museum, Bulletin 190:1-207.

FERGUSON, D. C.

- 1969 A revision of the moths of the subfamily Geometrinae of America north of Mexico (Insecta, Lepidoptera). Yale University, Peabody Museum of Natural History, Bulletin 19:1-251.
- 1971 Bombycoidea (Saturniidae) comprising subfamilies Citheroniinae, Hemileucinae (Part). In Dominick, R. B., et al., eds., The moths of America north of Mexico, including Greenland. Fasc. 20.2A. London, E. W. Classey & R. B. D. Publications. 153 pp.
- 1972 Bombycoidea (Saturniidae) comprising subfamilies Hemileucinae (Conclusion), Saturniinae. In Dominick, R. B., et al., eds., The moths of America north of Mexico, including Greenland. Fasc. 20.2B. London, E. W. Classey & R. B. D. Publications. 275 pp.
- 1978 Noctuoidea (Part): Lymantriidae. In Dominick, R. B., et al., eds., The moths of America north of Mexico, including Greenland. Fasc. 22.2. London, E. W. Classey & The Wedge Entomological Research Foundation. 110 pp.
- 1983 On the status of *Pseudothyatira expultrix* (Grt.) and *Euthyatira pennsylvanica* J. B. Smith (Thyatiridae). Journal of the Lepidopterists' Society 37:179-180.
- 1985 Contributions toward reclassification of the world genera of the tribe Arctiini. Part 1—Introduction and revision of the *Neoarctia*-*Grammia* group (Lepidoptera: Arctiidae, Arctiinae). Entomography 3:181-275.

FLETCHER, D. S.

- 1979 Geometroidea. In Nye, I. W. B., ed., The generic names of moths of the world. Vol 3. London, Trustees of the British Museum (Natural History). 243 pp.

FLETCHER, D. S. and I. W. B. NYE

- 1982 Sphingoidea. In Nye, I. W. B., ed., The generic names of moths of the world. Vol 4. London, Trustees of the British Museum (Natural History). 192 pp.

FORBES, W. T. M.

- 1923 The Lepidoptera of New York and neighboring states. Primitive Forms, Microlepidoptera, Pyraloids, Bombyces. Cornell University Agricultural Experiment Station, Memoir 68:1-729.
- 1948 The Lepidoptera of New York and neighboring states. Part II.

- Geometridae, Spingidae, Notodontidae, Lymantriidae. Cornell University Agricultural Experiment Station, Memoir 274:1-263.
- 1960 The Lepidoptera of New York and neighboring states. Part IV. Agaristidae through Nymphalidae, including butterflies. Cornell University Agricultural Experiment Station, Memoir 371:1-188.
- FORSTER, W. and T. WOHLFAHRT
 1976 Die Schmetterlinge Mitteleuropas, 2. Tagfalter. Franckh, Stuttgart.
- FRANCLEMONT, J. G.
 1973 Mimallonoidea; Bombycoidea (in part). In Dominick, R. B., et al., eds., The moths of America north of Mexico, including Greenland. Fasc. 20.1. London, E. W. Classey & R. B. D. Publications. 86 pp.
- FREEMAN, T. N.
 1969 Some historical notes on collecting Lepidoptera in the Mer Bleue. Canadian Field-Naturalist 83:16-18.
- GEIGER, H. J.
 1980 Enzyme electrophoretic studies on the genetic relationships of pierid butterflies (Lepidoptera: Pieridae) I. European Taxa. Journal of Research on the Lepidoptera 19:181-195.
 1986 Eskalation im Streit um die systematische Stellung der Taxa in der *Pieris napi* (L.)-Gruppe. (Lep. Pieridae). Atalanta 17:247-253.
- GLEN, R.
 1956 Entomology in Canada up to 1956: a review of developments and accomplishments. Canadian Entomologist 88:290-371.
- HEINICKE, W.
 1962 *Hyphantria cunea* (Drury), 1770 (Lep.). Bibliographische Hinweise anlässlich eines Fundes dieser Art auf dem Gebiete der D. D. R. Mitteilungsblatt für Insektenkunde 6:7-14.
- HIGGINS, L. G.
 1985 The correct name for what has been called *Lycaeides argyrognomon* in North America. Journal of the Lepidopterists' Society 39:145-146.
- HODGES, R. W.
 1971 Sphingoidea. In Dominick, R. B., et al., eds., The moths of America north of Mexico, including Greenland. Fasc. 21. London, E. W. Classey & R. B. D. Publications. 158 pp.
- HODGES, R. W., et al., eds.
 1983 Check list of the Lepidoptera of America north of Mexico. London, E. W. Classey & The Wedge Entomological Research Foundation. 284 pp.
- HOLLAND, W. J.
 1915 The moth book, a popular guide to a knowledge of the moths of North America. Garden City, N.Y., Doubleday, Page. 479 pp. [Reprinted. New York, Dover Publications, 1968].
- HÜBNER, J.
 1809 Sammlung exotischer Schmettlinge. Augsburg, 1806-1824.
- INTERNATIONAL TRUST FOR ZOOLOGICAL NOMENCLATURE
 1985 International Code of Zoological Nomenclature. 3rd ed. London, International Trust for Zoological Nomenclature. 338 pp.

JUDD, W. W.

- 1963 Butterflies of Dunn Township, Ontario. Ontario Field Biologist 17:1-14.
1970 Additional observations on butterflies of Dunn Township, Ontario. Ontario Field Biologist 24:13-16.

KLOTS, A. B.

- 1951 A field guide to the butterflies of North America, east of the Great Plains. Boston, Houghton Mifflin. 349 pp.

KOHALMI, L. and P. MOENS

- 1975 Evidence for the existence of an intergrade population between *Hyalophora gloveri nokomis* and *H. columbia* in northwestern Ontario (Lepidoptera: Saturniidae). Canadian Entomologist 107:793-799.

LAFONTAINE, J. D.

- 1972 More butterflies. Trail and Landscape 6:94-95.

LAMB, L.

- 1967 A checklist of Waterloo County Lepidoptera, Papilionoidea. 9 pp.
Published by the author, Kitchener.

LAYBERRY, R. A., J. D. LAFONTAINE, and P. W. HALL

- 1980 Butterflies of the Ottawa District. Trail and Landscape 16:3-59.

MAYR, E.

- 1966 Animal Species and Evolution. The Belknap Press of Harvard University Press, Cambridge, Massachusetts. 783 + pp.

McDUNNOUGH, J.

- 1938 Check list of the Lepidoptera of Canada and the United States of America. Part I Macrolepidoptera. Memoirs of the Southern California Academy of Sciences 1:1-275.
1939 Check list of the Lepidoptera of Canada and the United States of America. Part II Microlepidoptera. Memoirs of the Southern California Academy of Sciences 2:1-171.

McGUFFIN, W. C.

- 1967 Guide to the Geometridae of Canada (Lepidoptera). I. Subfamily Sterrhinae. Memoirs of the Entomological Society of Canada 50:1-67.
1972 Guide to the Geometridae of Canada (Lepidoptera). II. Subfamily Ennominae 1. Memoirs of the Entomological Society of Canada 86:1-159.
1977 Guide to the Geometridae of Canada (Lepidoptera). II. Subfamily Ennominae 2. Memoirs of the Entomological Society of Canada 101:1-126.
1981 Guide to the Geometridae of Canada (Lepidoptera). II. Subfamily Ennominae 3. Memoirs of the Entomological Society of Canada 117:1-153.
1987 Guide to the Geometridae of Canada (Lepidoptera). II. Subfamily Ennominae 4. Memoirs of the Entomological Society of Canada 138:1-182.
1988 Guide to the Geometridae of Canada (Lepidoptera). III, IV and V. Subfamilies Archiarinae, Oenochrominae, and Geometrinae. Memoirs of the Entomological Society of Canada 145:1-56.

McGUGAN, B. M.

- 1958 Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Volume I - Papilionidae to Arctiidae. Canada Department of Agriculture, Forest Biology Division, Publication 1034:1-76.

- MILLER, L. A.
 1956 Changes in insect populations in southwestern Ontario. Annual Report of the Entomological Society of Ontario 87:15-19.
- MORRIS, R. F.
 1963 Synonymy and color variation in the fall webworm, *Hyphantria cunea* Drury (Lepidoptera: Arctiidae). Canadian Entomologist 95:1217-1223.
- MUNROE, E.
 1956 Canada as an environment for insect life. Canadian Entomologist 88:372-476.
 1957 Northward extension in recent years of the lettered sphinx moth. Canadian Field-Naturalist 71:37.
- OPLER, P. A. and G. O. KRIZEK
 1984 Butterflies east of the Great Plains. Baltimore, Johns Hopkins University Press. 294 pp.
- PACLT, J.
 1979 The nomenclature of Lepidoptera in the second edition of "Kloet and Hincks", Part 2: Genus-group names. Bollettino della Società Entomologica Italiana 111:121-125.
- PENGELLY, D. H.
 1960 *Thymelicus lineola* (Ochs.) (Lepidoptera: Hesperiiidae) a pest of hay and pasture grasses in southern Ontario. Proceedings of the Entomological Society of Ontario 91:189-196.
 1965 Oviposition sites and the viability of eggs of *Thymelicus lineola* (Ochs.) (Lepidoptera: Hesperiiidae). Proceedings of the Entomological Society of Ontario 95:102-105.
- PLATT, A. P. and L. P. BROWER
 1968 Mimetic versus disruptive coloration in intergrading populations of *Limentis arthemis* and *astyanax* butterflies. Evolution 22:699-718.
- POOLE, R. W.
 1989 Noctuidae. In Heppner, J. B., ed., Lepidopterorum Catalogus (New Series). Fasc. 118, part 1-3. Leiden, E. J. Brill/Flora and Fauna Publications. 1314 pp.
- PRENTICE, R. M.
 1962 Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Volume 2 - Nycteolidae, Noctuidae, Notodontidae, Liparidae. Canada Department of Forestry, Forest Entomology and Pathology Branch, Bulletin 128:77-281.
 1963 Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Volume 3 - Lasiocampidae, Thyatiridae, Drepanidae, Geometridae. Canada Department of Forestry, Forest Entomology and Pathology Branch, Publication 1013:282-543.
 1965 Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Volume 4 - Microlepidoptera. Canada Department of Forestry, Forest Entomology and Pathology Branch, Publication 1142:544-840.
- RAIZENNE, H.
 1952 Forest Lepidoptera of southern Ontario and their parasites received and

reared at the Ottawa Forest Insect Survey Laboratory from 1937 to 1948. Ottawa, Canada Department of Agriculture, Forest Biology Division. 277 pp.

RINDGE, F. H.

- 1985 A revision of the moth genus *Acronyctodes*, with a review of the New World Bistonini (Lepidoptera, Geometridae). American Museum Novitates 2807:1-24.

RIOTTE, J. C. E.

- 1959 Revision of C. J. S. Bethune's list of the butterflies of the eastern provinces of Canada as far as northern Ontario is concerned. Ontario Field Biologist 13:1-18.
- 1962a *Adopoea lineola* (Hesperiidae) now also in northern Ontario. Journal of the Lepidopterists' Society 16:62.
- 1962b First additions to the northern Ontario list of butterflies. Journal of the Lepidopterists' Society 16:243-245.
- 1964 *Syssphinx bicolor* (Saturniidae) in Ontario. Journal of the Lepidopterists' Society 18:89-90.
- 1967a New and corrected butterfly records for Ontario and for Canada. Journal of the Lepidopterists' Society 21:135-137.
- 1967b *Pieris virginiensis* Edwards in Ontario (Lepidoptera: Pieridae). Proceedings of the Entomological Society of Ontario 98:27-29.
- 1968 Notes on *Euchloe ausonides mayi* (Pieridae) in Ontario. Journal of the Lepidopterists' Society 22:40-41.
- 1969 Rearing and description of the early stages of the Nearctic species of *Peridea*, with special reference to *P. basitriens* (Lepidoptera: Notodontidae). Michigan Entomologist 1:351-356.
- 1970a Checklist of Ontario skippers and butterflies. Toronto Entomologists' Association, Occasional Publication 1:1-4.
- 1970b *Crembidia prura* (Arctiidae: Lithosiinae) new for Canada. Journal of the Lepidopterist's Society 24:227.
- 1971a Butterflies and skippers of northern Ontario. Mid-Continent Lepidoptera Series 2(21):1-20.
- 1971b Eine neue Art der Gattung *Orgyia* (Untergattung *Hemerocampa*) in Nordamerika (Lep.:Lymantriidae). Entomologische Zeitschrift 81:105-115.
- 1972a A review of the North American hawk moth genus *Lapara* (Lepidoptera: Sphingidae). Royal Ontario Museum, Life Sciences Contributions 79:1-40.
- 1972b On the distribution of some skippers in Ontario. Journal of Research on the Lepidoptera 11:81-82.
- 1974 Über *Orgyia meridionalis* (Lep., Lymantriidae). Entomologische Zeitschrift 84:149-157.
- 1977 Abschliessende Bemerkungen zu den Studien über nordamerikanische Arten der Gattung *Orgyia* (Lepidoptera: Lymantriidae). Entomologische Zeitschrift 87:9-12.
- 1980 *Sphinx poecila*, a valid North American hawkmoth species (Lepidoptera: Sphingidae). Great Lakes Entomologist 13:115-130.
- 1987 Designation of a neotype for *Sphinx poecila* Stephens (Lepidoptera: Sphingidae). Great Lakes Entomologist 20:87-91.
- 1989 Ammerkungen zur Nomenklatur einiger nordamerikanischer *Orgyia* (Lepidoptera: Lymantriidae). Entomologische Zeitschrift 99:177-179.

RIOTTE, J. C. E. and R. S. PEIGLER

- 1981 A revision of the American genus *Anisota* (Saturniidae). Journal of Research on the Lepidoptera 19:101-180.

ROCKBURNE, E. W. and J. D. LAFONTAINE

- 1976 The cutworm moths of Ontario and Quebec. Canada Department of Agriculture, Research Branch, Publication 1593:1-164.

RUPERT, L. R.

- 1943 A specific revision of the genus *Metarrhanthia* (Lepidoptera, Geometridae, Ennominae). Journal of the New York Entomological Society 51:133-159.

SCOTT, J. A.

- 1986 The butterflies of North America. Stanford, Stanford University Press. 583 pp.

SNYDER, L. L.

- 1939 A plan on Ontario subdivisions and their names for naturalists. Canadian Field-Naturalist 53:22-24.

URQUHART, F. A.

- 1957 Factors influencing the changing fauna of Ontario. In Urquhart, F. A., ed., Changes in the fauna of Ontario. Toronto, University of Toronto Press for the Division of Zoology and Palaeontology, Royal Ontario Museum, pp. 1-3.

WAGNER, D. L.

- 1988 Taxonomic status of *Korcheltellus* Börner in North America (Lep.: Hepialidae). Journal of the New York Entomological Society 96:345-354.

WAGNER, D. L. and N. B. TINDALE

- 1988 An appraisal of *Gazorycta* Hübner (Hepialidae) and description of a new species from Arizona and New Mexico. Journal of the Lepidopterist's Society 42:204-212.

WALKER, E. M.

- 1913 Insects and their allies. In Faull, J. H., ed., The natural history of the Toronto region, Ontario, Canada. Toronto, Canadian Institute, pp. 295-403.

WARD, P. S., R. HARMSSEN, and P. D. N. HEBERT

- 1974 An annotated checklist of the macroheterocera of southeastern Ontario. Journal of Research on the Lepidoptera 13:23-42.

WARREN, B. C. S.

- 1961 The androconial scales and their bearing on the question of speciation in the genus *Pieris* (Lepidoptera). Entomologisk Tidskrift 82:121-148.
1963 The androconial scales in the genus *Pieris*. 2. The nearctic species of the *napi* - group. Entomologisk Tidskrift 84:1-4.

WERNY, K.

- 1966 Untersuchungen über die Systematik der Tribus Thyatirini, Macrothyatirini, Habrosynini und Tetheini (Lepidoptera: Thyatiridae). Inaugural-Dissertation, Mathematisch-Naturwissenschaftliche Fakultät, Universität des Saarlandes, Saarbrücken, Germany. 463 pp.

WORMINGTON, A.

- 1983 The butterflies of Point Pelee National Park, Ontario. Ontario Field Biologist 37:1-27.

INDEX OF SCIENTIFIC NAMES

Generic and specific synonyms, preoccupied names, and names for species, subspecies, and forms mentioned in the text are enclosed in square brackets.

- A
- Abagrotis* Smith 165
abalienalis (Walker), *Hypena* 135
 [*Abbottana* Hulst = *Eutrapela* Hübner] 95
abbottii (Swainson),
 Sphecodina 110
abditiva Heinrich, *Meroptera* 61
Abegesta Munroe 49
abietivorella (Grote), *Dioryctria* 61
abnaki (Klots), *Pediasia* 56
abortivaria (Herrich-Schäffer),
 Dyspteris 79
abrasaria (Herrich-Schäffer),
 Xanthorhoe 75
Abraxini 79
Abrostola Ochseneheimer 141
abrostoloides (Guenée), *Paectes* 143
abruptata (Walker), *Itame* 80
absinthiata (Clark), *Eupithecia* 77
absorptialis Walker, *Macrochilo* 134
abstrusa Eichlin and Cunningham,
 Syngrapha 142
acadia Barnes and Benjamin,
 Xylotype 158
acadicum (Edwards), *Satyrium* 29
acadiensis (Bethune),
 Cryptocala 163
Acallis Ragonot 58
Acasia Duponchel 79
acerifoliella (Fitch),
 Paraclemensia 10
acerni (Clemens), *Synanthedon* 13
acerrubri (Engelhardt),
 Synanthedon 13
Achalarus Scudder 15
Achatia Hübner 158
Achatodes Guenée 148
achemon (Drury), *Eumorpha* 110
Achyra Guenée 51
Acontia Ochseneheimer 144
Acontiinae 144
Acrossus Dyar 15
- acrea* (Drury), *Estigmene* 125
Acrobasis Zeller 59
Acronicta Ochseneheimer 146
Acronictinae 146
acronyctoides (Walker), *Panthea* 145
Actias Leach 102
acutella (Walker), *Nascia* 50
Adaina Tutt 65
adela Franclemont, *Xestia* 166
Adela Latreille 10
Adelinae 10
adipaloides Grote and Robinson,
 Diacme 53
Adita Grote 154
adjuncta (Boisduval),
 Melanchnra 160
Adoneta Clemens 47
adumbrata (Behr), *Drasteria* 136
aedon (Grote), *Andropolia* 148
aeglealis (Walker),
 Herpetogramma 54
Aellopos Hübner 109
aemula Hübner, *Idia* 133
aemulataria (Walker),
 Semiothisa 81
aenescentalis Munroe, *Palpita* 54
aenigmatica Heinrich, *Pyla* 61
aerata (Fabricius), *Synchlora* 69
aerea (Hübner), *Allagrapha* 141
aereoides Grote, *Diachrysia* 141
aeruginosa (Guenée), *Zale* 140
Aethalura McDunnough 83
Aethiophysa Munroe 50
affinata Pearsall, *Eupithecia* 77
 [*affirmaria* (Walker), *Triphosa* *haesitata*] 73
afflicta Grote, *Acronicta* 146
Agaristinae 147
agitatellus Clemens, *Crambus* 55
Aglossa Latreille 57
Agrapha Hübner 141
Agriades Hübner 34
agrilaceella (Packard),
 Rhaphiptera 55

<i>Agriopodes</i> Hampson	147	Alucitidae	11
<i>Agriphila</i> Hübner	56	<i>alurina</i> (Smith), <i>Orthosia</i>	161
<i>Agrius</i> Hübner	104	<i>Alypia</i> Hübner	147
Agrotini	162	<i>amanda</i> (Smith), <i>Lithophane</i>	156
<i>Agrotis</i> Ochseneimer	162	<i>amatrix</i> (Hübner), <i>Catocala</i>	139
<i>Alabama</i> Grote	136	<i>ambigualis</i> Walker, <i>Lascoria</i>	134
[<i>alaskensis</i> Holland, <i>Oeneis</i>		<i>Amblyptilia</i> Hübner	65
<i>jutta</i>]	44	<i>Amblyscirtes</i> Scudder	22
<i>albarufa</i> Grote, <i>Acronicta</i>	146	<i>amercalis</i> (Guenée), <i>Idia</i>	133
<i>albata</i> Packard, <i>Clemensia</i>	122	<i>americana</i> (Harris), <i>Acronicta</i>	146
<i>albatella</i> Ragonot, <i>Phycitodes</i>	62	[<i>americana</i> Harris, <i>Arctia caja</i>]	127
<i>albellus</i> Clemens, <i>Crambus</i>	55	[<i>americana</i> Harris, <i>Lycaena</i>	
<i>albicornis</i> (Hy. Edwards),		<i>phlaeus</i>]	29
<i>Synanthedon</i>	13	<i>americana</i> (Guérin-Méneville),	
[<i>albicostella</i> Fernald, <i>Donacula</i>		<i>Harrisina</i>	45
<i>melinella</i>]	54	<i>americana</i> (Speyer),	
<i>albidula</i> (Guenée), <i>Lithacodia</i>	144	<i>Amphipoea</i>	148
<i>albifera</i> (Walker), <i>Hydrelia</i>	76	<i>americanum</i> (Fabricius),	
<i>albifrons</i> (J. E. Smith),		<i>Malacosoma</i>	98
<i>Symmerista</i>	117	<i>americanum</i> (Harris),	
<i>albinuda</i> (Smith), <i>Lasionycta</i>	160	<i>Phyllodesma</i>	97-8
<i>albipennis</i> (Grote), <i>Euxoa</i>	164	<i>amica</i> (Hübner), <i>Catocala</i>	139
<i>albicapitata</i> Packard, <i>Eupithecia</i>	77	[<i>amica</i> (Edwards), <i>Plebejus</i>	
[<i>albolineata</i> (Packard), <i>Ecliptopera</i>		<i>saepiolus</i>]	33
<i>silaceata</i>]	72	<i>amicaria</i> (Herrich-Schäffer),	
<i>albosigma</i> Fitch, <i>Clostera</i>	112	<i>Probole</i>	92
<i>albovittata</i> (Guenée), <i>Trichodezia</i>	76	<i>amitaria</i> (Guenée), <i>Ematurga</i>	83
<i>Albuna</i> Hy. Edwards	12	<i>Amolita</i> Grote	148
<i>Alcathoe</i> Hy. Edwards	14	<i>amorata</i> Packard, <i>Callizzia</i>	95
[<i>alcestis</i> (Edwards), <i>Speyeria</i>		<i>Amorpha</i> Hübner	108
<i>aphrodite</i>]	37	<i>Amphion</i> Hübner	110
<i>alciphearia</i> Walker, <i>Selenia</i>	91	<i>Amphipoea</i> Billberg	148
<i>alcoolaria</i> (Guenée), <i>Plagodis</i>	92	<i>Amphipyra</i> Ochseneimer	148
<i>Aletia</i> Hübner	158	Amphipyridae	148
<i>algidata</i> (Möschler), <i>Xanthorhoe</i>	75	<i>ampla</i> (Walker), <i>Autographa</i>	141
<i>alia</i> (Guenée), <i>Apamea</i>	148	<i>amplexella</i> Ragonot, <i>Acrobasis</i>	59
<i>alias</i> (Ottolengui), <i>Syngrapha</i>	142	<i>amputatrix</i> (Fitch), <i>Apamea</i>	148
<i>alienaria</i> (Herrich-Schäffer),		<i>Amyna</i> Guenée	144
<i>Probole</i>	92	<i>amyntor</i> (Geyer), <i>Ceratomia</i>	104
<i>alienellus</i> (Zincken), <i>Crambus</i>	55	<i>amyntula</i> (Boisduval), <i>Everes</i>	32
<i>Allagrapha</i> Franclemont	141	<i>amyrisaria</i> (Walker),	
<i>Allotria</i> Hübner	138	<i>Metarranthis</i>	91
<i>alleni</i> (Grote), <i>Drasteria</i>	136	<i>Anacamptodes</i> McDunnough	83
<i>allionealis</i> (Walker), <i>Paraponyx</i>	48	<i>Anagasta</i> Heinrich	63
<i>alpinella</i> (Busck), <i>Antophila</i>	14	<i>Anageshna</i> Munroe	53
<i>Alsophila</i> Hübner	68	<i>Anagoga</i> Hübner	92
[<i>altantica</i> Munroe, <i>Coryphista</i>		Anagagini	91
<i>meadii</i>]	73	<i>Anagrapha</i> McDunnough	141
<i>altera</i> (Ottolengui), <i>Syngrapha</i>	142	<i>Anania</i> Hübner	51
<i>alternans</i> (Walker), <i>Egira</i>	159	<i>Anaplectoides</i> McDunnough	163
<i>alternata</i> (Grote), <i>Abagrotis</i>	162	<i>Anarta</i> Ochseneimer	158
<i>alternata</i> (Müller), <i>Epirrhoe</i>	75	<i>anartalis</i> (Grote), <i>Loxostege</i>	52
<i>alticolalis</i> (Dyar), <i>Acallia</i>	58	<i>Anartomima</i> Boursin	159
<i>Alucita</i> Linné	11	<i>anataria</i> (Swett), <i>Itame</i>	80

<i>Anathix</i> Franclemont	154	<i>Apamea</i> Ochsenheimer	148
<i>Anavitrinella</i> McDunnough	84	<i>apamiformis</i> (Guenée), <i>Apamea</i>	148
<i>ancellata</i> (Hulst), <i>Scopula</i>	71	<i>Apantes</i> Harris	127
<i>anceps</i> (Stephens), <i>Platypolia</i>	157	<i>Apatelodes</i> Packard	96
<i>anchocelioides</i> (Guenée),		<i>Apatelodidae</i>	96
<i>Rhynchagrotis</i>	166	<i>Apatelodinae</i>	96
[<i>ancosisconensis</i> (Morrison),		<i>Apaturini</i>	42
<i>Hyppa</i>]	151	<i>Apharetra</i> Grote	154
<i>Ancylolomiinae</i>	54	<i>Aphomia</i> Hübner	59
<i>Ancyloxypha</i> Felder	18	<i>aphrodite</i> (Fabricius), <i>Speyeria</i>	37
<i>andersoni</i> (Swett), <i>Itame</i>	80	<i>apiata</i> (Grote), <i>Epiglaea</i>	155
<i>andremona</i> (Cramer), <i>Hypocala</i>	137	<i>apicalis</i> (Grote and Robinson),	
<i>Androloma</i> Grote	147	<i>Schizura</i>	119
<i>Andropolia</i> Grote	148	<i>apicalis</i> (Walker), <i>Clostera</i>	113
[<i>Anepia</i> Hampson = <i>Hadena</i>		[<i>Apicia</i> Guenée = <i>Eusarca</i>	
Schrank]	159	Hübner]	94
<i>angelica</i> (Grote), <i>Olceclostera</i>	96	<i>apiciaria</i> (Packard),	
<i>Angerlini</i>	87	<i>Metarrhantis</i>	91
<i>anguina</i> (Grote), <i>Lacinipolia</i>	159	<i>apicosa</i> (Haworth),	
<i>anguina</i> (J. E. Smith),		<i>Homophoberia</i>	144
<i>Dasylophia</i>	117	<i>aplastella</i> (Hulst), <i>Tetralopha</i>	58
<i>angulalis</i> (Hübner), <i>Palthis</i>	134	<i>Aplectoides</i> Butler	163
<i>angulella</i> Ely, <i>Eurythmia</i>	63	<i>Apoda</i> Haworth	46
<i>angulifera</i> (Walker), <i>Callosamia</i>	102	<i>Apodrepanulatrix</i> Rindge	87
<i>angulosa</i> (J. E. Smith),		<i>Apomyelois</i> Heinrich	60
<i>Peridea</i>	114-5	<i>appalachia</i> (R. L. Chermock),	
<i>angusella</i> Grote, <i>Acrobasis</i>	59	<i>Satyrodes</i>	42
<i>angusii</i> Grote and Robinson,		<i>appassionata</i> (Harvey),	
<i>Datana</i>	113	<i>Papaipema</i>	153
<i>angustiorata</i> Walker, <i>Caripeta</i>	93	<i>approximella</i> (Walker), <i>Peoria</i>	63
<i>Anhimella</i> McDunnough	159	<i>aprica</i> (Hübner), <i>Acontia</i>	144
<i>Aniclini</i>	163	[<i>aquilo</i> auct., <i>Agriades</i>]	34
<i>anilis</i> (Drury), <i>Argyrostromis</i>	138	[<i>aquilonaris</i> (Lintner), <i>Furcula</i>	
<i>Anisota</i> Hübner	100	<i>scolopendrina</i>]	121
<i>anna</i> (Grote), <i>Grammia</i>	128	<i>Archanara</i> Walker	149
<i>annulata</i> (Hulst), <i>Eupithecia</i>	77	<i>Archiearinae</i>	68
<i>anodonta</i> (Guenée), <i>Conservula</i>	150	<i>Archiearis</i> Hübner	68
<i>Anomis</i> Hübner	136	<i>archippus</i> (Cramer), <i>Limenitis</i>	41
[<i>Anomogyna</i> Staudinger = <i>Xestia</i>		<i>arcigera</i> (Guenée), <i>Schinia</i>	167
Hübner]	166	<i>Arctia</i> Schrank	127
<i>Anorthodes</i> Smith	148	<i>Arctiidae</i>	121
<i>antennata</i> (Walker), <i>Lithophane</i>	156	<i>Arctiinae</i>	123
<i>Antepione</i> Packard	95	<i>Arctiini</i>	123
<i>Anterastria</i> Sugi	147	<i>arctivorens</i> Hampson,	
<i>anthedon</i> (A. H. Clark), <i>Enodia</i>	42	<i>Papaipema</i>	153
<i>Antheraea</i> Hübner	101	<i>arcuata</i> Walker, <i>Drepana</i>	67
<i>Anthocharinae</i>	26	<i>Arequipa</i> Walker	55
<i>anticaria</i> Walker, <i>Eupithecia</i>	77	<i>arge</i> (Drury), <i>Grammia</i>	128
<i>Anticarsia</i> Hübner	136	<i>argenteomaculatus</i> (Harris),	
<i>Anticlea</i> Stephens	74	<i>Sthenopsis</i>	9
<i>antinympa</i> (Hübner), <i>Catocala</i>	139	<i>argillacea</i> (Hübner), <i>Alabama</i>	136
<i>antiopa</i> (Linné), <i>Nymphalis</i>	35	<i>argillacearia</i> (Packard), <i>Itame</i>	80
<i>antiqua</i> (Linné), <i>Orgyia</i>	132	<i>argillaceala</i> (Packard),	
<i>Antiphila</i> Haworth	14	<i>Raphiptera</i>	55

Argynniae	37	<i>Atrytonopsis</i> Godman	22
Argyractini	48	Attacini	102
<i>Argyria</i> Hübner	56	<i>attenta</i> (Grote), <i>Eueretagrotis</i>	164
Argyriini	56	<i>Atteva</i> Walker	11
[<i>argyrognomon</i> (Bergsträsser), <i>Lycaeides</i>]	33	<i>augustiorata</i> (Walker), <i>Caripeta</i>	93
<i>Argyrogramma</i> Hübner	141	<i>augustus</i> (Kirby), <i>Incisalia</i>	30
<i>Argyrostrotis</i> Hübner	138	<i>aurantiaca</i> (Hübner), <i>Holomelina</i>	123-4
<i>armataria</i> (Herrich-Schäffer), <i>Cepphis</i>	92	<i>aurantiagio</i> Guenée, <i>Rhodoecia</i>	167
<i>arsaltealis</i> (Walker), <i>Palpita</i>	54	<i>aurata</i> (Packard), <i>Entephria</i>	74
<i>Arta</i> Grote	58	<i>auratella</i> (Clemens), <i>Vaxi</i>	57
<i>artesta</i> (Smith), <i>Trichoclea</i>	162	<i>auratus</i> Grote, <i>Sthenopsis</i>	9
<i>arthemis</i> (Drury), <i>Limenitis</i>	41	[<i>aurea</i> Fitch, <i>Atteva</i>]	11
<i>Ascalapha</i> Hübner	136	[<i>aureopurpura</i> (Hy. Edwards), <i>Carmentis</i>]	13
[<i>ascerta</i> Masters and Sorensen, <i>Oeneis</i> <i>jutta</i>]	44	<i>auricinctaria</i> Grote, <i>Melanomma</i>	137
<i>Ascia</i> Scopoli	25	<i>auricrinella</i> (Walshingham), <i>Epimartyria</i>	9
<i>asilipennis</i> (Boisduval), <i>Paranthrene</i>	12	<i>aurora</i> (J. E. Smith), <i>Hyparpax</i>	120
<i>asopialis</i> (Guenée), <i>Palthis</i>	134	<i>aurorella</i> Ely, <i>Acrobasis</i>	59
<i>asperatella</i> (Clemens), <i>Tetralopha</i>	58	<i>aurulenta</i> (Smith), <i>Euxoa</i>	164
<i>Aspilates</i> Treitschke	87	<i>ausonides</i> (Lucas), <i>Euchloe</i>	26
<i>assimilans</i> Walker, <i>Phragmatobia</i>	126	<i>australis</i> (Grote), <i>Baileya</i>	143
<i>assimilis</i> (Morrison), <i>Melanchra</i>	160	<i>Autographa</i> Hübner	141
<i>Astala</i> Davis	11	<i>Automeris</i> Hübner	101
[<i>asterius</i> Stoll, <i>Papilio</i> <i>polyxenes</i>]	24	<i>autumnata</i> (Borkhausen), <i>Epirrita</i>	77
<i>Asterocampa</i> Röber	42	<i>avimacula</i> Hudson, <i>Gluphisia</i>	116
<i>asteroides</i> Guenée, <i>Cucullia</i>	155	<i>aweme</i> (Lyman), <i>Papaipema</i>	153
Asthenini	76	<i>awemellus</i> McDunnough, <i>Crambus</i>	55
<i>astrictus</i> Morrison, <i>Eurois</i>	164	Azelinini	89
<i>astuta</i> Bird, <i>Papaipema</i>	153	B	
[<i>astyanax</i> (Fabricius), <i>Limenitis</i> <i>arthemis</i>]	41	<i>bachmannii</i> (Kirtland), <i>Libytheana</i>	34
<i>atalanta</i> (Linné), <i>Vanessa</i>	36	<i>badia</i> (Packard), <i>Schizura</i>	119
<i>Atalopedes</i> Scudder	20	<i>badia</i> (Hübner), <i>Prolimacodes</i>	46
[<i>athasaria</i> (Walker), <i>Lambdina</i> <i>fervidaria</i>]	93	<i>badicollis</i> (Grote), <i>Xestia</i>	166
<i>Athetis</i> Hübner	149	<i>badinodis</i> (Grote), <i>Xestia</i>	166
<i>atlantica</i> (Grote), <i>Lacanobia</i>	159	<i>badistriga</i> (Grote), <i>Homohadena</i>	156
[<i>atlantica</i> Munroe, <i>Caryplusia</i> <i>meadii</i>]	73	<i>badiusalis</i> (Walker), <i>Paraponyx</i>	48
<i>atlantis</i> (Edwards), <i>Speyeria</i>	37	<i>Baileya</i> Grote	143
<i>atomaria</i> Smith, <i>Hypena</i>	135	<i>baileyi</i> Grote, <i>Lithophane</i>	156
<i>atomaris</i> Hübner, <i>Phoberia</i>	137	<i>balluca</i> (Geyer), <i>Diachrysia</i>	141
<i>atomocellus</i> Dyar, <i>Yponomeuta</i>	12	<i>Balsa</i> Walker	149
<i>atrata</i> (Morrison), <i>Xestia</i>	166	<i>balimoralis</i> (Guenée), <i>Hypena</i>	135
[<i>atrocostalis</i> (Huard), <i>Boloria</i> <i>selene</i>]	39	<i>banksianae</i> Ferguson, <i>Semiothisa</i>	81
<i>atroliturata</i> (Walker), <i>Cladara</i>	79	<i>banksia</i> Mutuura, Monroe and Ross, <i>Dioryctria</i>	61
<i>atropunctata</i> (Packard), <i>Tacparia</i>	90		
<i>Atrytona</i> Scudder	21		

Baptini	86	Biston Leach	85
<i>baptisiae</i> (Forbes), <i>Erynnis</i>	16	Bistonini	85
<i>baptisiella</i> Fernald, <i>Tetralopha</i>	58	<i>bistriaria</i> Hübner, <i>Nemoria</i>	69
<i>barnesi</i> (Benjamin), <i>Abagrotis</i>	162	<i>bistriaris</i> Hübner, <i>Parallelia</i>	140
<i>basaliatum</i> (Walker), <i>Perizoma</i>	74	<i>bistriatella</i> (Hulst), <i>Apomyelois</i>	60
<i>basalis</i> Walker, <i>Scoparia</i>	47	<i>bitactata</i> (Walker), <i>Itame</i>	80
<i>basalis</i> Walker, <i>Marathyssa</i>	143	<i>biundata</i> Walker, <i>Heterocampa</i>	118
<i>basiflava</i> (Packard), <i>Dasychira</i>	131	<i>bivittata</i> (Grote), <i>Macrochilo</i>	134
<i>basilaris</i> Zeller, <i>Nymphopteryx</i>	61	<i>blandula</i> Hulst, <i>Catocala</i>	139
<i>basistriga</i> McDunnough,		<i>Blepharita</i> Hampson	154
<i>Hypocoena</i>	151	<i>Blepharomastix</i> Lederer	53
<i>basitriens</i> (Walker), <i>Peridea</i>	115	<i>Bleptina</i> Guenée	133
<i>bassiformis</i> (Walker), <i>Carmenta</i>	13	Boarmiini	83
<i>batesii</i> (Reakirt), <i>Phyciodes</i>	39	[<i>boisduvalii</i> (Duponchel), <i>Boloria</i>	
<i>bathyllus</i> (J. E. Smith), <i>Thorybes</i>	15	<i>titania</i>]	39
<i>Battus</i> Scopoli	23	<i>Boloria</i> Moore	38
<i>bella</i> (Linné), <i>Utetheisa</i>	123	<i>bolteri</i> (Hy. Edwards),	
<i>bellicula</i> Hübner, <i>Lithacodia</i>	144	<i>Synanthedon</i>	13
<i>bellona</i> (Fabricius), <i>Boloria</i>	38	Bombycoidea	96
<i>Bellura</i> Walker	149	<i>bombycoides</i> Walker, <i>Lapara</i>	107
<i>benesimilis</i> McDunnough,		[<i>Bomolocha</i> Hübner = <i>Hypena</i>	
<i>Euplexia</i>	151	<i>Schrank</i>]	135
<i>Besma</i> Capps	93	<i>borealis</i> (Hampson), <i>Heliothis</i>	167
<i>bethunei</i> (Grote and Robinson),		<i>borealis</i> (Hulst), <i>Cabera</i>	87
<i>Lithophane</i>	156	[<i>borealis</i> Packard, <i>Pyrausta</i>]	52
<i>betularius</i> (Linné), <i>Biston</i>	85-6	<i>borealis</i> (Smith), <i>Brachionycha</i>	154
<i>betulella</i> Hulst, <i>Acrobasis</i>	59	<i>borealis</i> (Guérin-Ménéville),	
<i>bicarnea</i> (Guenée), <i>Xestia</i>	166	<i>Furcula</i>	120
<i>bicolor</i> (Grote), <i>Eilema</i>	121	<i>bostoniensis</i> (Grote), <i>Euxoa</i>	164
<i>bicolor</i> (Harris), <i>Sphingicampa</i>	99	<i>bowmani</i> Cassino and Swett,	
<i>bicolorago</i> (Guenée), <i>Sunira</i>	158	<i>Eupithecia</i>	77
<i>bicoloralis</i> (Guenée), <i>Pyrausta</i>	52	<i>Brachionycha</i> Hübner	154
<i>bicolorata</i> (Fabricius),		<i>Brachylomia</i> Hübner	155
<i>Semiothisa</i>	81	<i>bradorata</i> McDunnough,	
<i>bidens</i> Zeller, <i>Crambus</i>	55	<i>Eupithecia</i>	77
<i>bidentata</i> Walker, <i>Nerice</i>	116	[<i>bradorata</i> (Munroe), <i>Entephria</i>	
<i>bifascialis</i> (Robinson),		<i>polata</i>]	74
<i>Petrophila</i>	48	<i>brephoides</i> (Walker),	
<i>biguttata</i> Packard, <i>Apoda</i>	46	<i>Leucobrephos</i>	68
<i>biguttellus</i> (Forbes),		<i>bridghami</i> (Grote and Robinson),	
<i>Microcrambus</i>	56	<i>Oligia</i>	152
<i>bijugalis</i> (Walker), <i>Hypena</i>	135	<i>briseis</i> Edwards, <i>Catocala</i>	139
<i>bilineatus</i> (Packard),		<i>brizo</i> (Boisduval and LeConte),	
<i>Lochmaeus</i>	118	<i>Erynnis</i>	16
<i>bilineata</i> (Packard), <i>Drepana</i>	67	<i>broweri</i> Heinrich, <i>Manhatta</i>	63
<i>biloba</i> (Stephens), <i>Autographa</i>	141	<i>bruceata</i> (Hulst), <i>Operophtera</i>	77
<i>bimacula</i> (Grote and Robinson),		<i>brucei</i> (Hy. Edwards), <i>Clostera</i>	113,
<i>Euphyes</i>	21	124	
<i>bimaculata</i> (Stephens),		<i>brunneata</i> (Thunberg), <i>Itame</i>	80
<i>Autographa</i>	141	<i>brunneicollis</i> (Grote),	
<i>binocula</i> (Grote), <i>Tarachidia</i>	145	<i>Protolampra</i>	166
<i>biplagiata</i> Walker, <i>Scoparia</i>	47	<i>bugrai</i> Kocak, <i>Xestia</i>	166
[<i>biseriata</i> (Packard), <i>Eudeilinia</i>]	67	<i>buscki</i> McDunnough, <i>Geina</i>	64
<i>bisignata</i> (Walker), <i>Semiothisa</i>	81		

C

<i>Cabera</i> Treitschke	87	<i>Capis</i> Grote	136
Caberini	87	<i>Capperia</i> Tutt	64
<i>cachexiata</i> Guenée, <i>Tetracis</i>	94	<i>caprealis</i> (Hübner), <i>Aglossa</i>	57
<i>cacuminaria</i> (Morrison), <i>Scopula</i>	71	<i>capsularis</i> (Guenée), <i>Hadena</i>	159
<i>Cadra</i> Walker	63	<i>cara</i> Guenée, <i>Catocala</i>	139
<i>caducus</i> Dyar, <i>Hyphenodes</i>	135	<i>Caradrina</i> Ochseneheimer	150
[<i>caecalis</i> (Walker), <i>Gesneria</i>		<i>caradrinalis</i> Guenée, <i>Bleptina</i>	133
<i>centuriella</i>]	47	[<i>carbonifera</i> Hampson, <i>Polia</i>]	161
<i>caeculalis</i> Zeller, <i>Perispasta</i>	51	<i>cardui</i> (Linné), <i>Vanessa</i>	36
<i>Caenurgina</i> McDunnough	138	<i>carduidactyla</i> (Riley), <i>Platyptilia</i>	65
[<i>caesonia</i> (Grote), <i>Tortricidia</i>]	45	<i>cariosa</i> (Guenée), <i>Apamea</i>	148
<i>caja</i> (Linné), <i>Arctia</i>	127	<i>Caripeta</i> Walker	93
[<i>calais</i> (Scudder), <i>Oeneis</i>		<i>Carmenta</i> Hy. Edwards	13
<i>chryxus</i>]	44	<i>carneella</i> Hulst, <i>Nephopterix</i>	61
<i>calami</i> (Harvey), <i>Cosmia</i>	150	<i>carneicosta</i> Guenée, <i>Panopoda</i>	137
<i>calanus</i> (Hübner), <i>Satyrrium</i>	29	<i>carneola</i> (Guenée), <i>Lithacodia</i>	145
<i>californicum</i> (Packard),		<i>carnosa</i> (Grote), <i>Psectraglaea</i>	157
<i>Malacosoma</i>	98	<i>carolina</i> Kearfott, <i>Stenoptilodes</i>	65
<i>caliginosellus</i> Clemens, <i>Crambus</i>	55	<i>Carsia</i> Hübner	78
[<i>Callarctia</i> Packard = <i>Arctia</i>		<i>Carterocephalus</i> Lederer	17
Schränk]	127	<i>caryae</i> Grote, <i>Acrobasis</i>	59
<i>Calledapteryx</i> Grote	95	<i>caryae</i> Harris, <i>Lophocampa</i>	130
Callimorphini	123	<i>caryaeorum</i> (McDunnough),	
<i>Callizzia</i> Packard	95	<i>Satyrrium</i>	30
<i>Callopietria</i> Hübner	149	<i>caryivorella</i> Ragonot, <i>Acrobasis</i>	59
<i>Callosamia</i> Packard	102	<i>casta</i> (Packard), <i>Crambidia</i>	121
<i>Caloreas</i> Heppner	14	<i>casta</i> (Pallas), <i>Psyche</i>	11
<i>calycanthata</i> (J. E. Smith), <i>Zale</i>	140	[<i>castigata</i> (Hübner), <i>Eupithecia</i>	
<i>Calyptra</i> Ochseneheimer	136	<i>subfuscata</i>]	78
<i>cambrica</i> Curtis, <i>Venusia</i>	76	<i>Catabena</i> Walker	150
<i>Campaea</i> Lamarck	89	<i>catalpae</i> (Boisduval),	
Campaeini	89	<i>Ceratonia</i>	104-5
<i>campestris</i> (Boisduval),		<i>cataphracta</i> (Grote), <i>Papaipema</i>	153
<i>Atalopedes</i>	20	<i>catenaria</i> (Cramer), <i>Cingilia</i>	93-4
<i>campestris</i> (Grote), <i>Euxoa</i>	164	<i>Catocala</i> Schränk	139
[<i>campestris</i> (Behr), <i>Phyciodes</i>]	39	Catocalinae	138
<i>canadaria</i> (Guenée),		<i>Catoptria</i> Hübner	56
<i>Melanolophia</i>	85	<i>catullus</i> (Fabricius), <i>Pholisora</i>	17
<i>canadensis</i> Bethune, <i>Calyptra</i>	136	<i>caudata</i> (Harris), <i>Alcathoe</i>	14
<i>canadensis</i> Boisduval, <i>Sphinx</i>	105	<i>caudellella</i> Dyar, <i>Sarata</i>	62
[<i>canadensis</i> McDunnough,		<i>cautella</i> (Walker), <i>Cadra</i>	63
<i>Cerura</i>]	121	<i>cecropia</i> (Linné), <i>Hyalophora</i>	103
<i>canadensis</i> (Munroe), <i>Petrophila</i>	48	<i>Celaena</i> Stephens	150
[<i>canadensis</i> Rothschild and Jordan,		[<i>Celama</i> Walker = <i>Nola</i> Leach]	144
<i>Papilio glaucus</i>]	23	<i>Celastrina</i> Tutt	32-3
<i>Canarsia</i> Hulst	62	<i>celia</i> (Saunders), <i>Grammia</i>	128
<i>candefacta</i> (Hübner),		<i>Celipetia</i> Guenée	140
<i>Tarachidia</i>	145	<i>celtidella</i> (Hulst), <i>Nephopterix</i>	61
<i>canicosta</i> Franclemont,		<i>celtis</i> (Boisduval and LeConte),	
<i>Symmerista</i>	117	<i>Asterocampa</i>	42
<i>caniplaga</i> (Walker), <i>Ellida</i>	116	<i>centaureae</i> (Rambur), <i>Pyrgus</i>	17
<i>canosaria</i> (Walker), <i>Nepytia</i>	94	<i>centerensis</i> (Lintner), <i>Acossus</i>	15
		<i>centrostrigaria</i> (Wollaston),	
		<i>Orthonama</i>	76

<i>centuriella</i> (Denis and Schiffermüller), <i>Gesneria</i> 47		<i>Chysolita</i> Grote 133	
<i>cephalonica</i> (Stainton), <i>Corcyra</i> 59		<i>Chytonix</i> Grote 150	
<i>Cepphis</i> Hübner 92		<i>churchillensis</i> (McDunnough), <i>Euxoa</i> 164	
<i>Cerastis</i> Ochsenheimer 163		<i>Cicinnus</i> Blanchard 96	
<i>cerata</i> Franclemont, <i>Chaetaglaea</i> 155		<i>cilicoides</i> (Grote), <i>Nola</i> 144	
<i>Ceratomia</i> Harris 104		<i>cimicifugata</i> Pearsall, <i>Eupithecia</i> 77	
<i>Cercyonis</i> Scudder 43		<i>cinerea</i> (Walker), <i>Furcula</i> 120	
<i>cerina</i> (Grote), <i>Papaipema</i> 153		<i>cinereana</i> (Neumoegen and Dyar), <i>Nycteola</i> 143	
<i>cerintha</i> (Treitschke), <i>Cerma</i> 150		[<i>cinereofrons</i> (Packard), <i>Schizura</i>] 119	
<i>cerisyi</i> Kirby, <i>Smerinthus</i> 107		<i>cinereola</i> (Guenée), <i>Ogdoconta</i> 152	
<i>Cerma</i> Hübner 150		<i>cinereomedia</i> Dyar, <i>Scoparia</i> 48	
<i>cerogama</i> Guenée, <i>Catocala</i> 139		<i>cineritia</i> (Grote), <i>Xylena</i> 158	
<i>ceromatica</i> (Grote), <i>Pyreferra</i> 157		[<i>cinerosa</i> Grote, <i>Lithophane</i>] 156	
<i>cerrusata</i> (Grote), <i>Papaipema</i> 153		<i>Cingilia</i> Walker 93	
<i>Cerura</i> Schrank 121		<i>Cingiliini</i> 93	
<i>Cerurinae</i> 120		<i>cingulatus</i> (Fabricius), <i>Agrius</i> 104	
[<i>cervinifascia</i> (Walker), <i>Dysstroma</i> <i>hersiliata</i>] 72		<i>cingulifera</i> (Walker), <i>Zale</i> 140	
<i>cesonia</i> (Stoll), <i>Zerene</i> 27		<i>Cirrophanus</i> Grote 150	
<i>Chaetaglaea</i> Franclemont 155		<i>Cisseps</i> Franclemont 131	
<i>chagnoni</i> Barnes and McDunnough, <i>Xylomoia</i> 154		<i>Cissusa</i> Walker 136	
<i>Chalcoela</i> Zeller 50		<i>Cisthene</i> Walker 122	
<i>Charadra</i> Walker 145		<i>Citheroniinae</i> 99	
<i>Charaxinae</i> 42		<i>citrata</i> (Linné), <i>Dysstroma</i> 71	
<i>Charidryas</i> Scudder 40		<i>citromba</i> Franclemont, <i>Pyreferra</i> 157	
<i>chersis</i> (Hübner), <i>Sphinx</i> 105		<i>Cladara</i> Hulst 79	
<i>Chersotis</i> Boisduval 163		<i>clandestina</i> (Harris), <i>Spaelotis</i> 166	
<i>Chilini</i> 57		<i>clarescens</i> Guenée, <i>Acronicta</i> 146	
<i>Chilo</i> Zincken 57		[<i>clarki</i> (Freeman), <i>Incisalia</i> <i>niphon</i>] 31	
<i>chionanthi</i> (J. E. Smith), <i>Adita</i> 154		<i>clarus</i> (Cramer), <i>Eupargyreus</i> 15	
[<i>chippewa</i> Edwards, <i>Colias</i> <i>palaeno</i>] 26		<i>claudia</i> (Cramer), <i>Euptoieta</i> 37	
<i>chiridotum</i> Grote, <i>Lacosoma</i> 96		<i>clemataria</i> (J. E. Smith), <i>Eutrapela</i> 95	
<i>Chlorochlamys</i> Hulst 69		<i>Clemensia</i> Packard 122	
<i>chloroleucaria</i> (Guenée), <i>Chlorochlamys</i> 69		<i>Cleora</i> Curtis 84	
<i>chlorostigma</i> (Harvey), <i>Oligia</i> 152		<i>clethrae</i> Dyar, <i>Nola</i> 144	
[<i>Chlosyne</i> auct. = <i>Charidryas</i> Scudder] 40		<i>clintonii</i> Grote, <i>Catocala</i> 139	
<i>Choephora</i> Grote and Robinson 163		<i>Clostera</i> Samouelle 112	
<i>Choerocampini</i> 111		<i>clymene</i> (Brown), <i>Haploa</i> 123	
<i>Choreutidae</i> 14		<i>clyton</i> (Boisduval and LeConte), <i>Asterocampa</i> 42	
<i>Choreutinae</i> 14		[<i>coagulata</i> Guenée, <i>Eupithecia</i>] 77	
<i>Choreutis</i> Hübner 14		<i>coccinata</i> Grote, <i>Catocala</i> 139	
[<i>Chorizagrotis</i> Smith] 164		<i>Cochliidiini</i> 45	
<i>chortalis</i> (Grote), <i>Sitochroa</i> 52		<i>coelebs</i> Grote, <i>Catocala</i> 139	
<i>Chrysanympa</i> Grote 141		<i>coenia</i> (Hübner), <i>Junonia</i> 37	
<i>Chrysauginae</i> 58		<i>Coenochroa</i> Ragonot 63	
<i>Chrysoteuchia</i> Hübner 55		<i>Coenonympha</i> Hübner 43	
<i>chryxus</i> (Doubleday and Hewitson), <i>Oeneis</i> 44		<i>Coenymphini</i> 43	
		<i>cogitata</i> (Smith), <i>Apamea</i> 148	

[<i>cognatarius</i> (Guenée), <i>Biston</i> <i>betularius</i>]	86	<i>contigua</i> (Walker), <i>Haploa</i>	123
<i>Coliadinae</i>	26	<i>contracta</i> Walker, <i>Datana</i>	113
<i>Colias</i> Fabricius	26	<i>contractata</i> (Packard), <i>Thera</i>	73
<i>Colobochyla</i> Hübner	136	<i>contrahens</i> Walker, <i>Anhimella</i>	159
<i>Colocasia</i> Hampson	145	<i>convergaria</i> (Walker), <i>Eufidonia</i>	85
<i>colona</i> Hübner, <i>Haploa</i>	123	<i>convexipennis</i> Grote and Robinson,	
<i>coloradensis</i> Fernald, <i>Stenoptilia</i>	65	<i>Cucullia</i>	155
<i>coloradensis</i> (Hulst), <i>Eupithecia</i>	78	<i>convolutella</i> (Hübner), <i>Zophodia</i>	62
<i>coloraria</i> (Fabricius), <i>Erastria</i>	87	<i>coortaria</i> (Hulst), <i>Itame</i>	80
<i>columbia</i> (S. I. Smith),		<i>Copivaleria</i> Grote	155
<i>Hyalophora</i>	103	<i>Copromorphoidea</i>	11
<i>columbiata</i> (Dyar), <i>Eupithecia</i>	78	<i>cora</i> Hübner, <i>Cerma</i>	150
<i>comma</i> (Harris), <i>Polygonia</i>	34	<i>coras</i> (Cramer), <i>Polites</i>	19
<i>comma</i> (Linné), <i>Hesperia</i>	19	<i>Corcyra</i> Ragonot	59
<i>commixtalis</i> (Walker), <i>Loxostege</i>	52	<i>cordata</i> (Ljungh), <i>Callopietria</i>	149
<i>commoda</i> (Walker), <i>Apamea</i>	148	<i>cordigera</i> (Thunberg), <i>Anarta</i>	158
<i>commoides</i> Guenée, <i>Leucania</i>	160	<i>corni</i> (Hy. Edwards), <i>Carmenta</i>	13
<i>communis</i> (Grote), <i>Pyrgus</i>	17	<i>Coryphista</i> Hulst	73
<i>comosa</i> Morrison, <i>Euxoa</i>	164	<i>Cosmia</i> Ochsenheimer	150
<i>compressipalpis</i> Guenée,		<i>Cossidae</i>	14
<i>Plusiodonta</i>	138	<i>Cossinae</i>	14
<i>comptaria</i> (Walker), <i>Venusia</i>	76	<i>Cossoidea</i>	14
<i>comptoniella</i> Hulst, <i>Acrobasis</i>	59	<i>costalis</i> (Fabricius), <i>Hypsopygia</i>	57
<i>comptulatalis</i> (Hulst),		<i>costa</i> Barnes and Benjamin,	
<i>Occidentalia</i>	57	<i>Properigea</i>	157
<i>comstocki</i> Grote, <i>Feralia</i>	155	<i>costiferalis</i> Walker, <i>Pyralis</i>	57
<i>comyntas</i> (Godart), <i>Everes</i>	32	<i>Cosymbiini</i>	70
<i>concinna</i> (J. E. Smith),		[<i>couperi</i> Grote, <i>Glaucompsyche</i>	
<i>Schizura</i>	119	<i>lygdamus</i>]	33
<i>concinnimacula</i> (Guenée),		<i>Crambidia</i> Packard	121
<i>Malintha</i>	145	<i>Crambiformes</i>	47
<i>concumbens</i> Walker, <i>Catocala</i>	139	<i>Crambinae</i>	54
<i>condensata</i> (Walker), <i>Hydrelia</i>	76	<i>Crambini</i>	55
<i>Condica</i> Walker	150	<i>Crambodes</i> Guenée	151
<i>condita</i> (Guenée), <i>Aplectoides</i>	163	<i>Crambus</i> Fabricius	55
<i>Condylolomia</i> Grote	58	<i>crassiuscula</i> (Haworth),	
<i>confederata</i> (Grote and Robinson),		<i>Caenurgina</i>	138
<i>Astala</i>	11	<i>crataegi</i> Saunders, <i>Catocala</i>	139
<i>confusa</i> (Hübner), <i>Morrisonia</i>	160	<i>crenulata</i> (Butler), <i>Orthodes</i>	160
<i>confusa</i> (Lyman), <i>Haploa</i>	123	<i>crepuscularia</i> (Denis and	
<i>confusaria</i> (Hübner), <i>Eusarca</i>	94	Schiffermüller), <i>Ectropis</i>	84
<i>congermana</i> (Morrison),		<i>crephontes</i> Cramer, <i>Papilio</i>	23
<i>Sideridis</i>	161	[<i>Cressonia</i> Grote and Robinson =	
<i>congruum</i> Walker, <i>Spilosoma</i>	125	<i>Amorpha</i> Hübner]	108
<i>coniella</i> (Ragonot), <i>Myelopsis</i>	60	<i>cretaceata</i> (Packard), <i>Eupithecia</i>	78
<i>connecta</i> Grote, <i>Acronicta</i>	146	<i>cretidactylus</i> (Fitch),	
<i>connubialis</i> Guenée, <i>Catocala</i>	139	<i>Oidaematophorus</i>	65
<i>Conservula</i> Grote	150	<i>cristata</i> Morrison,	
<i>consobrinella</i> (Zeller),		<i>Homophoberia</i>	144
<i>Glyptocera</i>	60	<i>cristifera</i> (Walker), <i>Papestra</i>	161
<i>conspicua</i> (Edwards), <i>Euphyes</i>	21	<i>critica</i> Forbes, <i>Urola</i>	57
<i>contacta</i> (Walker), <i>Andropolia</i>	148	<i>crocallata</i> Guenée, <i>Tetraxis</i>	94
<i>contexta</i> (Grote), <i>Plusia</i>	142	<i>Crocidophora</i> Lederer	51
		<i>Crocigrapha</i> Grote	159

<i>cruentaria</i> (Hübner), <i>Erastria</i>	87	<i>dejecta</i> Strecker, <i>Catocala</i>	139
<i>Cryptocala</i> Benjamin	163	<i>delaware</i> (Edwards), <i>Atrytone</i>	21
<i>Ctenucha</i> W. Kirby	131	<i>delawaricus</i> Zeller, <i>Oxyptilus</i>	64
<i>Ctenuchinae</i>	131	<i>delicata</i> (Grote), <i>Trachea</i>	154
<i>Cucullia</i> Schrank	155	<i>delphinii</i> (Boisduval), <i>Euclea</i>	47
<i>Cuculliinae</i>	154	<i>demissaria</i> (Hübner), <i>Idaea</i>	70
<i>cucurbitae</i> (Harris), <i>Melittia</i>	12	<i>demotella</i> Grote, <i>Acrobasis</i>	59
<i>culea</i> (Guenée), <i>Ulolonche</i>	162	<i>densa</i> (Walker), <i>Bellura</i>	149
<i>cunea</i> (Drury), <i>Hyphantria</i>	126	<i>dentata</i> (Grote), <i>Apharetra</i>	154
<i>cupida</i> (Grote), <i>Rhynchagrotis</i>	166	<i>denticulalis</i> (Harvey), <i>Idia</i>	133
<i>cuprina</i> Zeller, <i>Aglossa</i>	57	<i>denticulata</i> Grote, <i>Semiothisa</i>	81
<i>curialis</i> (Smith), <i>Mamestra</i>	160	<i>deridens</i> (Guenée), <i>Charadra</i>	145
<i>curvata</i> Grote, <i>Capis</i>	136	[<i>designata</i> auct., <i>Xanthorhoe</i>	
<i>curviferalis</i> (Walker), <i>Paraponyx</i>	48	<i>labradorensis</i>]	75
<i>curvimacula</i> (Morrison),		<i>Desmia</i> Westwood	53
<i>Xylena</i>	158	<i>destinata</i> (Möschler), <i>Eulithis</i>	72
<i>cuspidata</i> (Hübner), <i>Euclidia</i>	140	<i>determinata</i> Walker, <i>Metanema</i>	91
<i>cviatella</i> Dyar, <i>Meroptera</i>	61	<i>detersa</i> (Walker), <i>Euxoa</i>	164
<i>cybele</i> (Fabricius), <i>Speyeria</i>	38	<i>detersata</i> (Guenée), <i>Tacparia</i>	90
<i>cycladata</i> Grote and Robinson,		<i>detracta</i> (Walker), <i>Polia</i>	161
<i>Heliomata</i>	79	[<i>Deuteronomos</i> Prout = <i>Ennomos</i>	
<i>Cyclophora</i> Hübner	70	Treitschke]	89
<i>Cynia</i> Hübner	130	<i>devastator</i> (Brace), <i>Apamea</i>	148
<i>cymatophoroides</i> (Guenée),		<i>devia</i> (Grote), <i>Eupsilia</i>	155
<i>Pseudothyatira</i>	66	<i>Diachrysia</i> Hübner	141
<i>cymela</i> (Cramer), <i>Megisto</i>	43	<i>Diacme</i> Warren	53
<i>cynica</i> Guenée, <i>Orthodes</i>	161	[<i>Diactinia</i> Warren = <i>Ecliptoptera</i>	
<i>cynthia</i> (Drury), <i>Samia</i>	102	Warren]	72
D		<i>diana</i> (Hübner), <i>Choreutis</i>	14
<i>dactylina</i> Grote, <i>Acronicta</i>	146	<i>Diaphania</i> Hübner	53
<i>dammersi</i> Engelhardt,		<i>Diarsia</i> Hübner	163
<i>Penstemonia</i>	14	<i>Diastictis</i> Hübner	54
<i>Danaidae</i>	45	<i>Diathrausta</i> Lederer	53
<i>Danainae</i>	45	<i>Dicestra</i> Hampson	159
<i>Danaus</i> Kluk	45	<i>Dichogamini</i>	49
<i>Darapsa</i> Walker	111	<i>Dichorda</i> Warren	69
<i>Dasychira</i> Hübner	131	<i>Dicymolomia</i> Zeller	50
<i>Dasylophia</i> Packard	117	<i>diffinis</i> (Boisduval), <i>Hemaris</i>	109
[<i>Dasyuris</i> auct. = <i>Entephria</i>		<i>diffusa</i> (Walker), <i>Faronta</i>	159
Hübner]	74	<i>digitalis</i> (Grote), <i>Lemmeria</i>	152
<i>Datana</i> Walker	113	<i>Dilophonotini</i>	109
[<i>dawsoni</i> (Barnes and McDunnough),		<i>dilucida</i> (Morrison), <i>Xestia</i>	166
<i>Boloria eunomia</i>]	38	<i>dimidiata</i> (Hufnagel), <i>Idaea</i>	70
<i>deceptalis</i> (Walker), <i>Hypena</i>	135	<i>diminuendis</i> (Barnes and	
<i>declarata</i> (Walker), <i>Euxoa</i>	164	McDunnough), <i>Idia</i>	133
<i>decliva</i> (Grote), <i>Epiglaea</i>	155	<i>dion</i> (Edwards), <i>Euphyes</i>	22
<i>decolor</i> (Walker), <i>Enargia</i>	151	<i>Dioryctria</i> Zeller	61
<i>decoloraria</i> Hulst, <i>Cepphis</i>	92	<i>diptheroides</i> (Guenée),	
<i>decoloris</i> Hübner, <i>Pangrapta</i>	137	<i>Leuconycta</i>	144
<i>defecta</i> (Grote), <i>Hypocoena</i>	151	<i>Dipterygia</i> Stephens	151
<i>definita</i> Packard, <i>Orgyia</i>	132	<i>disa</i> (Thunberg), <i>Erebia</i>	44
<i>Deidamia</i> Clemens	110	<i>discalis</i> (Grote), <i>Metalectra</i>	137
		<i>disciferalis</i> Dyar, <i>Pyrallis</i>	57
		<i>discinigra</i> (Walker), <i>Brachylomia</i>	155

<i>discivaria</i> (Walker), <i>Parastichtis</i>	157	<i>Dyspyralis</i> Warren	133
<i>disclusa</i> Heinrich, <i>Dioryctria</i>	61	<i>Dysstroma</i> Hübner	71
<i>discoidalis</i> (Kirby), <i>Erebia</i>	44	E	
<i>discospilata</i> (Walker), <i>Eufidonia</i>	85	<i>Eacles</i> Hübner	99
<i>dislocaria</i> (Packard), <i>Enconista</i>	82	[<i>Earophila</i> Gumpfenberg = <i>Anticlea</i>	
<i>dislocata</i> (Smith), <i>Diarsia</i>	163	Stephens]	74
<i>dispar</i> (Linné), <i>Lymantria</i>	133	<i>Ecliptoptera</i> Warren	72
<i>disposita</i> Morrison, <i>Lithophane</i>	156	<i>Ecpantheria</i> Hübner	126
[<i>dispunctata</i> (Walker), <i>Semiothisa</i>		<i>Ectoedemia</i> Busck	10
<i>signaria</i>]	82	<i>Ectropis</i> Hübner	84
[<i>dissectus</i> Grote, <i>Crambus</i>		<i>edictalis</i> (Walker), <i>Hypena</i>	135
<i>alienellus</i>]	55	<i>edmandsii</i> (Packard), <i>Vitula</i>	62
<i>distria</i> (Hübner), <i>Malacosoma</i>	98	<i>eductalis</i> (Walker), <i>Lomanaltes</i>	134
<i>distans</i> (Grote), <i>Acrionicta</i>	146	<i>edwardsii</i> (Beutenmüller),	
<i>distincta</i> Hübner, <i>Achatia</i>	158	<i>Penstemonia</i>	14
<i>Ditrysia</i>	11	<i>edwardsii</i> (Fish), <i>Stenoptilodes</i>	65
<i>divergens</i> (Walker), <i>Euxoa</i>	164	<i>edwardsii</i> (Grote and Robinson),	
<i>diversicolor</i> (Morrison),		<i>Satyrion</i>	30
<i>Meropleon</i>	152	<i>effecta</i> (Walker), <i>Euchlaena</i>	88
<i>diversilineata</i> (Hübner), <i>Eulithis</i>	72	<i>egeremet</i> (Scudder),	
<i>divisaria</i> (Walker), <i>Hydriomena</i>	73	<i>Wallengrenia</i>	20
<i>divisata</i> Walker, <i>Caripeta</i>	93	<i>Egira</i> Duponchel	159
<i>Dolba</i> Walker	104	<i>egle</i> (Drury), <i>Euchaetes</i>	130
<i>dolosa</i> Franclemont, <i>Xestia</i>	166	<i>Eilema</i> Hübner	121
<i>dolosa</i> Grote, <i>Egira</i>	159	<i>ekthlipsis</i> (Grote), <i>Nymphula</i>	48
<i>Donacaula</i> Meyrick	54	<i>Elaphria</i> Hübner	151
<i>dorcas</i> (Kirby), <i>Lycaena</i>	28	<i>electilis</i> (Morrison), <i>Psaphida</i>	157
<i>doris</i> (Boisduval), <i>Grammia</i>	128	<i>elegans</i> (Clemens),	
<i>dormitans</i> (Guenée), <i>Baileya</i>	143	<i>Microcrambus</i>	56
<i>dorsipennata</i> (Barnes and		<i>elegans</i> (Packard), <i>Packardia</i>	46
McDunnough), <i>Dasychira</i>	131	<i>elegans</i> (Strecker), <i>Odontotia</i>	115
<i>dorsipunctella</i> (Kearfott),		<i>Ellida</i> Grote	116
<i>Pediasia</i>	56	<i>elliottii</i> (Fernald),	
<i>doubledayi</i> (Guenée), <i>Baileya</i>	143	<i>Oidaematophorus</i>	65
<i>Drasteria</i> Hübner	136	<i>ello</i> (Linné), <i>Erinnyis</i>	109
<i>Drepana</i> Schrank	67	<i>elonympha</i> (Hübner), <i>Allotria</i>	138
<i>Drepanidae</i>	67	<i>elutella</i> (Hübner), <i>Ephestia</i>	63
<i>Drepaninae</i>	67	<i>Elymniinae</i>	42
<i>Drepanoidea</i>	66	<i>Ematurga</i> Lederer	83
<i>drexelii</i> Hy. Edwards, <i>Datana</i>	113	<i>Emmelina</i> Tutt	65
<i>drupiferarum</i> J. E. Smith,		<i>Enargia</i> Hübner	151
<i>Sphinx</i>	105	<i>Enconista</i> Lederer	82
[<i>drusus</i> (Edwards), <i>Charidryas</i>		<i>endopiaria</i> (Grote and Robinson),	
<i>nycteis</i>]	40	<i>Besma</i>	93
<i>Dryocampa</i> Harris	99	<i>engeli</i> (Dyar), <i>Salebriaria</i>	60
<i>dryopterata</i> Grote, <i>Calledapteryx</i>	95	[<i>engeli</i> Chermock and Chermock,	
<i>duaria</i> (Guenée), <i>Metarrhantis</i>	91	<i>Ceratomia</i>]	105
<i>dubium</i> Walker, <i>Spilosoma</i>	125	[<i>enigmata</i> (Burns and McDunnough),	
<i>dubitans</i> (Walker), <i>Apamea</i>	148	<i>Itame coartaria</i>]	80
[<i>ducens</i> (Walker), <i>Feltia</i>]	165	<i>Ennominae</i>	79
<i>dukes</i> (Lindsey), <i>Euphyes</i>	22	<i>Ennomini</i>	89
<i>duovata</i> (Bird), <i>Papaipema</i>	153	<i>Ennomos</i> Treitschke	89
<i>duplicata</i> (Bethune), <i>Zale</i>	140		
<i>Dyspteris</i> Hübner	79		

<i>Enodia</i> Hübner	42	<i>Eucirroedia</i> Grote	155
<i>Entephria</i> Hübner	74	<i>Euclea</i> Hübner	47
<i>enthea</i> Grote, <i>Fishia</i>	156	Eucleini	47
<i>Eoparargyractis</i> Lange	49	<i>Euclidia</i> Ochsenheimer	140
<i>Eosporopteryx</i> Dyar	142	<i>Eudeilinea</i> Packard	67
<i>Epargyreus</i> Hübner	15	<i>Eudryas</i> Boisduval	147
<i>Epelis</i> Hulst	81	Eudulini	77
<i>Ephestia</i> Guenée	63	<i>Eueretagrotis</i> Smith	164
<i>Ephestiodes</i> Ragonot	62	<i>Eufidonia</i> Packard	85
<i>ephyraria</i> (Walker), <i>Anacamptodes</i>	83	<i>Eugonobapta</i> Warren	94
<i>epigaea</i> (Grote), <i>Syngrapha</i>	142	<i>Eugraphe</i> Hübner	164
<i>Epiglaea</i> Grote	155	<i>Eulithis</i> Hübner	72
<i>Epimartyria</i> Walshingham	9	<i>Eulogia</i> Heinrich	62
<i>Epimecis</i> Hübner	85	<i>Eumacaria</i> Packard	80
<i>epimenis</i> (Drury), <i>Psychomorpha</i>	147	Eumaeinae	29
<i>epione</i> (Drury), <i>Catocala</i>	139	<i>eumelusalis</i> (Walker), <i>Phalaenostola</i>	134
<i>Epipaschia</i> Clemens	58	<i>Eumorpha</i> Hübner	110
Epipaschiinae	58	<i>eunomia</i> (Esper), <i>Boloria</i>	38
Epiplemlidae	95	<i>Euparthenos</i> Grote	140
Epirrhantini	90	<i>eupatorii</i> (Fernald), <i>Oidaematophorus</i>	65
<i>Epirrhantis</i> Hübner	90	<i>eupatorii</i> (Lyman), <i>Papaipema</i>	153
<i>Epirrhoe</i> Hübner	75	<i>euphorbiae</i> (Linné), <i>Hyles</i>	112
<i>Epirrita</i> Hübner	77	<i>Euphydryas</i> Scudder	40
<i>epixanthe</i> (Boisduval and LeConte), <i>Lycaena</i>	28	<i>Euphyes</i> Scudder	21
<i>Erannis</i> Hübner	86	<i>Euphyia</i> Hübner	76
<i>Erastria</i> Hübner	87	<i>Eupithecia</i> Curtis	77
<i>erastrioides</i> (Guenée), <i>Tarachidia</i>	145	Eupitheciini	77
<i>Erebia</i> Dalman	44	<i>Euplexia</i> Stephens	151
Erebiini	44	<i>Eupsilia</i> Hübner	155
[<i>Erebus</i> Latreille = <i>Ascalapha</i> Hübner]	136	<i>Euptoieta</i> Doubleday	37
<i>erectea</i> (Cramer), <i>Caenurgina</i>	138	[<i>Euptychia</i> auct. = <i>Megisto</i> Hübner]	43
<i>Erelieva</i> Heinrich	63	Euptychiini	43
<i>eremitus</i> (Hübner), <i>Sphinx</i>	106	<i>Eurema</i> Hübner	27
<i>Erinnyis</i> Hübner	109	<i>Eurostrymon</i> Clench	31
<i>eripalis</i> (Grote), <i>Stegia</i>	49	<i>Eurois</i> Hübner	164
<i>Erora</i> Scudder	32	<i>Eurrhynx</i> Hübner	51
<i>erosa</i> Hübner, <i>Anomis</i>	136	[<i>Eurycyrtarus</i> Hampson = <i>Astala</i> Davis]	11
<i>Erynnis</i> Schrank	16	<i>eurydice</i> (Johansson), <i>Satyrodes</i>	42
<i>eryphon</i> (Boisduval), <i>Incisalia</i>	30	<i>eurytheme</i> Boisduval, <i>Colias</i>	26
<i>erythemaria</i> Guenée, <i>Cabera</i>	87	<i>Eurythmia</i> Ragonot	63
<i>erythrella</i> Ragonot, <i>Ephestiodes</i>	62	<i>Eurytides</i> Hübner	24
<i>Estigmene</i> Hübner	125	<i>Eusarca</i> Hübner	94
<i>Etiella</i> Zeller	60	<i>Eustixia</i> Hübner	49
<i>Euagrotis</i> McDunnough	163	<i>Eustroma</i> Hübner	72
<i>Eubaphe</i> Hübner	77	<i>Eutelina</i> Hübner	143
[<i>eubule</i> (Linné), <i>Phoebis sennae</i>]	27	Euteliinae	143
<i>Euchaetes</i> Harris	130	<i>Euthyatira</i> Smith	66
<i>Euchlaena</i> Hübner	88	<i>Eutrapela</i> Hübner	95
<i>Euchloe</i> Hübner	26	<i>Eutricopis</i> Morrison	167
		<i>Euxoa</i> Hübner	164

[<i>Euxoa</i> Hübner]	164
<i>Euzophera</i> Zeller	62
<i>evagaria</i> (Hulst), <i>Itame</i>	80
<i>evansi</i> (McDunnough), <i>Capperia</i>	64
<i>evansi</i> (McDunnough), <i>Orthonama</i>	76
<i>Everes</i> Hübner	32
Evergestinae	50
<i>Evergestis</i> Hübner	50
Everini	32
<i>evicta</i> (Grote), <i>Morrisonia</i>	160
<i>exauspicata</i> (Walker), <i>Itame</i>	80
<i>excaecatus</i> (J. E. Smith), <i>Paonias</i>	107
<i>exitiosa</i> (Say), <i>Synanthedon</i>	13
<i>exhausta</i> (Smith), <i>Oligia</i>	152
<i>exigua</i> (Hübner), <i>Spodoptera</i>	154
Exoporia	9
<i>expandens</i> (Walker), <i>Tetralopha</i>	59
<i>explanata</i> (Walker), <i>Eulithis</i>	72
<i>exprimens</i> (Walker), <i>Pyrrhia</i>	167
[<i>expultrix</i> (Grote), <i>Pseudothyatira</i>]	66
<i>extensa</i> (Smith), <i>Apamea</i>	148
<i>extimia</i> (Walker), <i>Caradrina</i>	150
<i>extricalis</i> Guenée, <i>Nealgedonia</i>	51
<i>Exyra</i> Grote	144

F

<i>fabulosa</i> Ferguson, <i>Xestia</i>	166
<i>factiosalis</i> (Walker), <i>Renia</i>	135
<i>fadus</i> (Cramer), <i>Aellopos</i>	109
<i>fagina</i> Morrison, <i>Lithophane</i>	156
<i>Fagitana</i> Walker	151
[<i>falacer</i> (Godart), <i>Satyrium</i> <i>calanus</i>]	29
<i>falcifera</i> (Kirby), <i>Anagrapha</i>	141
<i>fallax</i> (Herrich-Schäffer), <i>Agriopodes</i>	147
<i>farinalis</i> (Linné), <i>Pyrallis</i>	57
<i>Faronta</i> Smith	159
<i>fasciata</i> (Sulzer), <i>Eumorphia</i>	110
<i>fasciola</i> (Herrich-Schäffer), <i>Lithacodes</i>	46
<i>faulalis</i> (Walker), <i>Munroessa</i>	48
<i>faunus</i> (Edwards), <i>Polygonia</i>	34
<i>feltella</i> Dyar, <i>Acrobasis</i>	59
<i>Feltia</i> Walker	165
<i>Feniseca</i> Grote	28
<i>fennica</i> (Tauscher), <i>Ochropleura</i>	166
<i>Feralia</i> Grote	155
<i>fernaldi</i> (Ragonot), <i>Nephopterix</i>	61
[<i>ferrealis</i> Grote, <i>Lithophane</i>]	156

<i>ferrugata</i> (Clerk), <i>Xanthorhoe</i>	75
<i>ferruginea</i> (Packard), <i>Peridea</i>	115
<i>ferruginosa</i> (Walker), <i>Holomelina</i>	124
<i>ferruminaria</i> (Zeller), <i>Leptostales</i>	71
<i>fervidaria</i> Herrich-Schäffer, <i>Plagodis</i>	92
<i>fervidaria</i> (Hübner), <i>Lambdina</i>	93
<i>fessa</i> Grote, <i>Amolita</i>	148
<i>festivoides</i> (Guenée), <i>Elaphria</i>	151
<i>figulilella</i> (Gregson), <i>Cadra</i>	63
<i>figurata</i> (Drury), <i>Grammia</i>	128
[<i>filamentaria</i> Guenée, <i>Nematocampa</i>]	95
<i>finitella</i> (Walker), <i>Tulsa</i>	61
<i>finitima</i> Guenée, <i>Apamea</i>	148
<i>finlaysoni</i> Riotte, <i>Anisota</i>	100
<i>fiscellaria</i> (Guenée), <i>Lambdina</i>	93
<i>Fishia</i> Grote	156
<i>fishii</i> (Grote), <i>Metalepsis</i>	165
<i>Fissicrambus</i> Bleszynski	56
<i>fissinotata</i> (Walker), <i>Semiothisa</i>	81
<i>flagellum</i> (Walker), <i>Autographa</i>	141
<i>flava</i> (Fabricius), <i>Anomis</i>	136
<i>flavicaria</i> (Packard), <i>Itame</i>	80
<i>flavicornis</i> (Smith), <i>Calocasia</i>	145
<i>flavidalis</i> (Guenée), <i>Polygrammodes</i>	54
<i>flavipunctalis</i> (Geyer), <i>Renia</i>	135
<i>flavivenata</i> (Hulst), <i>Orthofidonia</i>	83
<i>flavofasciatus</i> (Walker), <i>Proserpinus</i>	111
<i>fletcherata</i> Taylor, <i>Eupithecia</i>	78
[<i>fletcheri</i> (Michener and dos Passos), <i>Satyrium liparops</i>]	30
<i>flexuosa</i> (Grote), <i>Tortricidia</i>	45
<i>florea</i> Guenée, <i>Cucullia</i>	155
<i>florida</i> (Guenée), <i>Schinia</i>	167
<i>floridensis</i> B. P. Clark, <i>Amphion</i>	110
[<i>floridus</i> Zeller, <i>Crambus</i> <i>pascuellus</i>]	55
<i>fodinalis</i> (Lederer), <i>Pyrausta</i>	52
<i>forbesellus</i> (Fernald), <i>Thopeutis</i>	57
<i>forbesi</i> Franclemont, <i>Euagrotis</i>	163
<i>forbesi</i> (French), <i>Idia</i>	133
<i>formosa</i> (Grote), <i>Chrysanympha</i>	141
<i>formosalis</i> Walker, <i>Nigetia</i>	145
<i>fractilinea</i> (Grote), <i>Oligia</i>	135
<i>fractilinea</i> (Smith), <i>Hypenodes</i>	152
<i>fragilis</i> (Guenée), <i>Acrionicta</i>	146
<i>Framinghamia</i> Strand	54
<i>franklinii</i> (Curtis), <i>Agriades</i>	34

<i>frater</i> Grote, <i>Raphia</i>	145	<i>Geina</i> Tutt	64
<i>fraxini</i> (Hy. Edwards), <i>Albuna</i>	12	Gelechioidea	11
<i>freemani</i> Hardwick, <i>Diarsia</i>	163	[<i>gelida</i> (Möschler), <i>Grammia</i>	
<i>frei</i> ja (Thunberg), <i>Boloria</i>	38	<i>quenseli</i>]	129
<i>frigga</i> (Thunberg), <i>Boloria</i>	38-9	<i>gelidata</i> (Möschler), <i>Eupithecia</i>	78
[<i>frigida</i> (Smith), <i>Papaipema</i>]	153	<i>geminata</i> (Packard), <i>Packardia</i>	46
<i>frigidana</i> (Walker), <i>Nycteola</i>	143	[<i>geminatus</i> Say, <i>Smerinthus</i>]	107
<i>frigidaria</i> (Möschler), <i>Scopula</i>	71	<i>gemmatalis</i> Hübner, <i>Anticarsia</i>	136
<i>fritillaria</i> (Guenée),		<i>gemmatella</i> (Hulst), <i>Peoria</i>	63
<i>Homochlodes</i>	90	<i>generosus</i> (Grote and Robinson),	
<i>frustulum</i> Guenée, <i>Celiptera</i>	140	<i>Pyrausta</i>	52
<i>fuscata</i> Hübner, <i>Hypoprepia</i>	122	<i>geniculata</i> Grote and Robinson,	
<i>fulcalis</i> (Clemens), <i>Petrophila</i>	48	<i>Feltia</i>	165
<i>fuliginosa</i> (Linné),		Geometridae	68
<i>Phragmatobia</i>	126-7	Geometrinae	68
<i>fuliginosalis</i> (Fernald),		Geometroidea	68
<i>Lipocosmodes</i>	50	<i>georgica</i> (Herrich-Schäffer),	
<i>fulvicollis</i> (Hübner), <i>Ciseps</i>	131	<i>Hyperaeschra</i>	114
[<i>fulvicosta</i> (Clemens), <i>Haploa</i>]	123	<i>georgii</i> Grote, <i>Lithophane</i>	156
<i>fulvipes</i> (Harris), <i>Synanthedon</i>	13	<i>georgii</i> Hulst, <i>Plemyria</i>	72
<i>fumalis</i> (Grote), <i>Euxoa</i>	164	Gerydini	28
<i>fumalis</i> (Guenée), <i>Fumibotys</i>	51	<i>Gesneria</i> Hübner	47
[<i>fumata</i> Taylor, <i>Eupithecia</i>		<i>gibbicostata</i> (Walker), <i>Stamnodes</i>	75
<i>satyrata</i>]	78	<i>gibbosa</i> (J. E. Smith), <i>Nadata</i>	114
<i>Fumibotys</i> Munroe	51	[<i>gibsonella</i> Kearfott, <i>Thaumatoopsis</i>	
<i>fumoferalis</i> (Hulst), <i>Saucrobotys</i>	50	<i>pexella</i>]	56
<i>fumosa</i> (Hulst), <i>Eupithecia</i>	78	<i>gibsoni</i> McDunnough,	
[<i>fumosum</i> Strecker,		<i>Grammia</i>	128
<i>Spilosoma</i>]	125-6	<i>gigantea</i> Strecker, <i>Colias</i>	26
<i>funebri</i> s (Ström), <i>Anania</i>	51	<i>girardellus</i> Clemens, <i>Crambus</i>	55
<i>funeralis</i> Grote and Robinson,		<i>galbanata</i> (Morrison), <i>Zale</i>	140
<i>Acrionicta</i>	146	<i>glabella</i> (Morrison), <i>Amphipyra</i>	148
<i>funeralis</i> (Hübner), <i>Desmia</i>	53	<i>gladiaria</i> Morrison, <i>Agrotis</i>	162
<i>fungorum</i> Grote and Robinson,		[<i>glandon</i> auct., <i>Agriades</i>]	34
<i>Choephora</i>	163	<i>glaphyralis</i> (Guenée), <i>Glaphyria</i>	49
<i>furcata</i> (Smith), <i>Papaipema</i>	153	<i>Glaphyria</i> Hübner	49
<i>furcata</i> (Thunberg), <i>Hydriomena</i>	73	Glaphyriinae	49
<i>furcilla</i> (Packard), <i>Panthea</i>	145	<i>Glaucoopsyche</i> Scudder	33
<i>Furcula</i> Lamarck	120	<i>glaucus</i> Linné, <i>Papilio</i>	23
<i>furfurata</i> (Grote), <i>Homorthodes</i>	159	<i>glenni</i> Cashatt,	
<i>fusca</i> (Haworth), <i>Pyla</i>	61	<i>Oidaematophorus</i>	65
G		<i>glomeraria</i> (Grote), <i>Lomographa</i>	86
<i>Galasa</i> Walker	58	<i>gloriosa</i> (Guenée), <i>Habrosyne</i>	66
<i>galbanata</i> (Morrison), <i>Zale</i>	140	<i>gloriosa</i> (Strecker), <i>Schinia</i>	167
<i>Galgula</i> Guenée	151	<i>Gluphisia</i> Boisduval	116
<i>Galleria</i> Fabricius	59	Gluphisiini	116
Galleriinae	59	<i>Glyptocera</i> Ragonot	60
Galleriini	59	<i>gnophosaria</i> (Guenée),	
<i>gallii</i> (Rottenburg), <i>Hyles</i>	112	<i>Semiothisa</i>	81
<i>garita</i> (Reakirt), <i>Oarisma</i>	18	<i>goodelli</i> (Grote), <i>Orthodes</i>	161
Gastropachinae	97	<i>gordius</i> Cramer, <i>Sphinx</i>	106
<i>Gazorycta</i> Hübner	10	<i>gorgone</i> (Hübner), <i>Charidryas</i>	40
		<i>gortynoides</i> Walker, <i>Bellura</i>	149
		<i>gracilineata</i> (Guenée), <i>Eulithis</i>	72

<i>gracilis</i> (Grote), <i>Korscheltellus</i>	9	<i>hastata</i> (Linné), <i>Rheumaptera</i>	74
<i>gracilis</i> (Grote and Robinson), <i>Hemaris</i>	109	<i>hayhurstii</i> (Edwards), <i>Staphylus</i>	16
<i>gracilis</i> (Grote and Robinson), <i>Polygonia</i>	34	<i>hebetata</i> (Hulst), <i>Semiothisa</i>	81
<i>Grammia</i> Rambur	128	<i>hebraeicum</i> Hübner, <i>Polygrammate</i>	147
<i>grandirena</i> (Haworth), <i>Drasteria</i>	136	<i>hegon</i> (Scudder), <i>Amblyscirtes</i>	22
[<i>grandis</i> (Barnes and McDunnough), <i>Boloria titania</i>]	39	<i>helata</i> (Smith), <i>Zale</i>	140
<i>grandis</i> (Guenée), <i>Spiramater</i>	162	<i>Helicoverpa</i> Hardwick	167
<i>Graphiphora</i> Ochsenheimer	165	<i>Heliomata</i> Grote	79
[<i>Graphium</i> auct. = <i>Eurytides</i> Hübner]	24	<i>Heliothinae</i>	167
<i>grata</i> (Fabricius), <i>Eudryas</i>	147	<i>Heliothis</i> Ochsenheimer	167
<i>grata</i> Hübner, <i>Elaphria</i>	151	<i>helooides</i> (Boisduval), <i>Lycaena</i>	28
<i>grataria</i> (Fabricius), <i>Haematopis</i>	70	<i>Hellula</i> Guenée	49
<i>grisea</i> Walker, <i>Acronicta</i>	146	[<i>Helotropha</i> Lederer = <i>Celaena</i> Stephens]	150
<i>grotei</i> (Morrison), <i>Copivaleria</i>	155	<i>helva</i> (Grote), <i>Apamea</i>	148
<i>grotei</i> Riley, <i>Lithophane</i>	156	<i>helvalis</i> (Walker), <i>Framinghamia</i>	54
<i>grynea</i> (Cramer), <i>Catocala</i>	139	<i>helvialis</i> (Walker), <i>Helvibotys</i>	52
<i>grynea</i> (Hübner), <i>Mitoura</i>	30	<i>Helvibotys</i> Munroe	52
<i>Gueneria</i> Packard	91	<i>Hemaris</i> Dalman	109
<i>guttivitta</i> (Walker), <i>Heterocampa</i>	118	<i>Hemileuca</i> Walker	101
<i>gyralis</i> (Hulst), <i>Munroessa</i>	48	<i>Hemileucinae</i>	101
H		<i>Hemileucini</i>	101
<i>habilis</i> Grote, <i>Catocala</i>	139	<i>hemina</i> Grote, <i>Lithophane</i>	156
<i>Habrosyne</i> Hübner	66	<i>Hemipachnobia</i> McDunnough	165
<i>Habrosynini</i>	66	<i>Hemitheini</i>	69
<i>Hadena</i> Schrank	159	<i>henrici</i> (Grote), <i>Simyra</i>	147
<i>Hadeninae</i>	158	<i>henrici</i> (Grote and Robinson), <i>Incisalia</i>	30
<i>Haematopis</i> Hübner	70	[<i>henshawi</i> (Swett), <i>Epirrita</i> <i>autumnata</i>]	77
<i>haesitata</i> (Grote), <i>Acronicta</i>	146	<i>Hepialidae</i>	9
<i>haesitata</i> (Guenée), <i>Triphosa</i>	73	<i>Hepialoidea</i>	9
<i>Hahncappsia</i> Munroe	51	[<i>Hepialus</i> Fabricius = <i>Korscheltellus</i> Börner]	9
<i>Halysidota</i> Hübner	129	[= <i>Gazorycta</i> Hübner]	10
<i>hamamelis</i> Guenée, <i>Acronicta</i>	146	<i>Heptagrotis</i> McDunnough	165
<i>hamellus</i> (Thunberg), <i>Crambus</i>	55	<i>Herculia</i> Walker	58
<i>hammondi</i> (Riley), <i>Psorosina</i>	62	<i>herilis</i> (Grote), <i>Feltia</i>	165
<i>hanhami</i> (Smith), <i>Phalaenostola</i>	134	<i>herminiata</i> (Guenée), <i>Eudeilinea</i>	67
<i>Haploa</i> Hübner	123	<i>Hermiiniinae</i>	133
<i>Harkenclenus</i> dos Passos	29	<i>Herpetogramma</i> Lederer	54
<i>harlequinialis</i> Dyar, <i>Diathrausta</i>	53	<i>hersiliata</i> (Guenée), <i>Dysstroma</i>	71-2
<i>harrisii</i> (Grote), <i>Papaipema</i>	153	<i>Hesperia</i> Fabricius	19
<i>harrisii</i> (Scudder), <i>Charidryas</i>	40	<i>hesperidago</i> (Guenée), <i>Pyreferra</i>	157
<i>harrisii</i> Walsh, <i>Halysidota</i>	129	<i>Hesperiidae</i>	15
<i>Harrisimemna</i> Grote	147	<i>Hesperiinae</i>	18
<i>Harrisina</i> Packard	45	<i>Hesperioidea</i>	15
<i>haruspica</i> (Grote), <i>Graphiphora</i>	165	<i>Hesperumia</i> Packard	83
<i>hasta</i> Guenée, <i>Acronicta</i>	146	<i>Heterocampa</i> Doubleday	118
		<i>Heterocampini</i>	117
		<i>Heterogenea</i> Knoch	46
		<i>Heteropacha</i> Harvey	97
		<i>Heterophleps</i> Herrich-Schäffer	79

Heteropterinae	17	<i>Hyphantria</i> Harris	126
<i>heterosalis</i> (McDunnough),		<i>Hypocala</i> Guenée	137
<i>Eudonia</i>	48	<i>hypocharia</i> (Herrich-Schäffer),	
<i>Hethemia</i> Ferguson	69	<i>Metarrhantis</i>	91
<i>hexadactyla</i> Linné, <i>Alucita</i>	11	<i>Hypocoena</i> Hampson	151
<i>hianna</i> (Scudder), <i>Atrytonopsis</i>	22	Hypolimnini	37
<i>hibisci</i> (Guenée), <i>Orthosia</i>	161	<i>Hypoprepia</i> Hübner	122
<i>Hillia</i> Grote	156	<i>Hyppa</i> Duponchel	151
<i>hircina</i> Morrison, <i>Homoglaea</i>	156	<i>Hypsopygia</i> Hübner	57
<i>hobomok</i> (Harris), <i>Poanes</i>	21	I	
<i>Holomelina</i> Herrich-Schäffer	123		
<i>Homochlodes</i> Hulst	90	<i>icciusalis</i> (Walker), <i>Munroessa</i>	48
<i>homodactylus</i> (Walker),		<i>icelus</i> (Scudder and Burgess),	
<i>Oidaematophorus</i>	65	<i>Erynnis</i>	16
<i>Homoeosoma</i> Curtis	62	<i>Idaea</i> Treitschke	70
<i>Homoglaea</i> Morrison	156	<i>idalia</i> (Drury), <i>Speyeria</i>	38
<i>Homohadena</i> Grote	156	<i>idas</i> (Linné), <i>Lycæides</i>	33
<i>Homophoberia</i> Morrison	144	<i>Idia</i> Hübner	133
<i>Homorthodes</i> McDunnough	159	<i>iduata</i> (Guenée), <i>Xanthorhoe</i>	75
<i>honestaria</i> (Walker), <i>Pero</i>	89	<i>igneæ</i> (Grote), <i>Syngrapha</i>	142
<i>horatius</i> (Scudder and Burgess),		<i>ilia</i> (Cramer), <i>Catocala</i>	139
<i>Erynnis</i>	16	<i>illapsa</i> (Walker), <i>Euagrotis</i>	163
<i>Horisme</i> Hübner	77	<i>illecta</i> Walker, <i>Catocala</i>	139
[<i>Hormisa</i> Walker = <i>Macrochilo</i>		<i>illibella</i> (Hulst), <i>Coenochroa</i>	63
Hübner]	134	<i>illocata</i> (Walker), <i>Oligia</i>	152
<i>horrida</i> Hübner, <i>Zale</i>	140	<i>illocata</i> Warren, <i>Dyspyralis</i>	133
<i>hortaria</i> (Fabricius), <i>Epimecis</i>	85	<i>imbrifera</i> (Guenée), <i>Polia</i>	161
<i>hortulata</i> (Linné), <i>Eurrhyncha</i>	51	<i>immanis</i> Guenée, <i>Hydraecia</i>	151
<i>hubneraria</i> (Guenée), <i>Pero</i>	89	<i>Immyrta</i> Dyar	60
[<i>hudsonianus</i> Clark, <i>Papilio</i>]	23	<i>impecuniosa</i> (Grote),	
[<i>huebneri</i> (Wallengren), <i>Alucita</i>]	11	<i>Papaipema</i>	153
<i>humaria</i> (Guenée),		<i>imperialis</i> (Drury), <i>Eacles</i>	99
<i>Anacamptodes</i>	83	<i>implete</i> Walker, <i>Acronicta</i>	146
<i>humuli</i> Harris, <i>Hypena</i>	135	<i>implicata</i> McDunnough,	
<i>hyalinata</i> (Linné), <i>Diaphania</i>	53	<i>Lacinipolia</i>	159
<i>Hyalophora</i> Duncan	103	<i>impulsa</i> (Guenée), <i>Apamea</i>	148
<i>Hydraecia</i> Guenée	151	<i>inatomaria</i> Guenée, <i>Metanema</i>	91
<i>Hydrelia</i> Hübner	76	<i>incertata</i> Walker, <i>Mesothea</i>	69
<i>Hydria</i> Hübner	73	<i>incincta</i> (Morrison),	
<i>Hydriomena</i> Hübner	73	<i>Protorthodes</i>	161
Hydriomenini	71	<i>Incisalia</i> Scudder	30
<i>Hylephila</i> Billberg	19	<i>inclara</i> Smith, <i>Acronicta</i>	146
<i>Hyles</i> Hübner	112	<i>includens</i> (Walker),	
<i>hyllus</i> (Cramer), <i>Lycæna</i>	28-9	<i>Pseudoplusia</i>	142
<i>hyloeus</i> (Drury), <i>Dolba</i>	104	<i>includens</i> (Walker),	
<i>Hymenia</i> Hübner	53	<i>Spartiniphaga</i>	154
<i>Hypagyrtis</i> Hübner	84	<i>inclusa</i> (Hübner), <i>Clostera</i>	113
<i>Hyparpax</i> Hübner	120	<i>increta</i> Morrison, <i>Acronicta</i>	146
<i>Hypena</i> Schrank	135	[<i>Incurvaria</i> Haworth = <i>Lampronia</i>	
Hypeninae	135	Stephens]	10
<i>Hypenodes</i> Doubleday	135	Incurvariidae	10
Hypenodinae	135	Incurvariinae	10
<i>Hyperaeschra</i> Butler	114	Incurvarioidea	10
<i>Hyperstrotia</i> Hampson	144		

<i>indeclinata</i> (Walker), <i>Metarrhantis</i>	91	<i>interrogationis</i> (Fabricius), <i>Polygonia</i>	35
<i>indigenella</i> (Zeller), <i>Acrobasis</i>	59	<i>interrupta</i> Guenée, <i>Acronicta</i>	146
<i>indocilis</i> (Walker), <i>Apamea</i>	149	<i>interruptofasciata</i> Packard, <i>Eupithecia</i>	78
<i>indomita</i> (Walker), <i>Melipotis</i>	137	<i>intertexta</i> (Walker), <i>Aethalura</i>	83
<i>inermis</i> Forbes, <i>Leucania</i>	160	<i>intestinata</i> (Guenée), <i>Horisme</i>	77
<i>inexplicata</i> (Walker), <i>Mycterophora</i>	137	<i>intricata</i> (Zetterstedt), <i>Eupithecia</i>	78
<i>infans</i> (Möschler), <i>Archiearis</i>	68	<i>inulta</i> (Grote), <i>Metaxaglaea</i>	157
<i>infanta</i> (Smith), <i>Colocasias</i>	145	<i>io</i> (Fabricius), <i>Automeris</i>	101
<i>inficita</i> (Walker), <i>Apamea</i>	149	<i>irole</i> Boisduval, <i>Nathalis</i>	28
<i>inficita</i> (Walker), <i>Marathysa</i>	143	<i>iphitalis</i> (Walker), <i>Chalcoela</i>	50
<i>infimbrialis</i> Dyar, <i>Herculia</i>	58	<i>Ipimorpha</i> Hübner	152
<i>infimella</i> Ragonot, <i>Ephesiodes</i>	62	<i>ipomoeae</i> Doubleday, <i>Schizura</i>	119
<i>infixa</i> (Walker), <i>Homohadena</i>	156	<i>ipsilon</i> (Hufnagel), <i>Agrotis</i>	162
<i>inflatella</i> (Clemens), <i>Prochoreutis</i>	14	<i>iridaria</i> (Guenée), <i>Dichorda</i>	69
<i>infumata</i> (Grote), <i>Enargia</i>	151	<i>Iridopsis</i> Warren	84
[<i>ingravis</i> (Smith), <i>Papestra</i>]	161	<i>iris</i> Guenée, <i>Phlogophora</i>	153
<i>innominata</i> (Smith), <i>Lithophane</i>	156	<i>iris</i> (Zetterstedt), <i>Hillia</i>	156
<i>innotata</i> Guenée, <i>Acronicta</i>	146	<i>irraria</i> Barnes and McDunnough, <i>Euchlaena</i>	88
[<i>innotatellus</i> Walker, <i>Crambus</i> <i>perlellus</i>]	55	[<i>irrorata</i> (Packard), <i>Oreta</i>]	68
<i>innubens</i> Guenée, <i>Catocala</i>	139	<i>irrubriella</i> Ely, <i>Acrobasis</i>	59
<i>inordinata</i> (Morrison), <i>Apamea</i>	149	<i>irus</i> (Godart), <i>Incisalia</i>	31
<i>inornata</i> Edwards, <i>Coenonympha</i>	43	<i>Isa</i> Packard	47
<i>inornata</i> (Hulst), <i>Hydrelia</i>	76	<i>isabella</i> (J. E. Smith), <i>Pyrharctia</i>	124
<i>inquaesita</i> (Grote and Robinson), <i>Papaipema</i>	153	[<i>Isia</i> auct. = <i>Pyrharctia</i> Packard]	124
<i>inquinalialis</i> (Zeller), <i>Udea</i>	52	[<i>Isturgia</i> auct. = <i>Epelis</i> Hulst]	81
<i>inquinatus</i> Zeller, <i>Oidaematophorus</i>	65	<i>Itame</i> Hübner	80
<i>inscripta</i> (Harris), <i>Deidamia</i>	110	[<i>itealis</i> (Walker), <i>Paraponyx</i> <i>allionealis</i>]	48
<i>insolabilis</i> Guenée, <i>Catocala</i>	139	<i>itysalis</i> (Walker), <i>Udea</i>	52
<i>insueta</i> Guenée, <i>Leucania</i>	160	J	
<i>insulsaria</i> (Guenée), <i>Pleuroprucha</i>	70	[<i>j-album</i> (Boisduval and LeConte), <i>Nymphalis vau-album</i>]	36
<i>integerrima</i> Grote and Robinson, <i>Datana</i>	113-4	<i>jacchusalis</i> (Walker), <i>Polypogon</i>	134
<i>integra</i> Zeller, <i>Lineodes</i>	52	<i>jaculifera</i> (Guenée), <i>Feltia</i>	165
<i>interior</i> Scudder, <i>Colias</i>	26	<i>jamaicensis</i> (Drury), <i>Smerinthus</i>	107
[<i>intermedia</i> Speyer, <i>Cucullia</i>]	155	<i>jocosa</i> (Guenée), <i>Feralia</i>	155
[<i>intermedia</i> (Stretch), <i>Grammia</i> <i>parthenice</i>]	128	<i>johnsonaria</i> (Fitch), <i>Euchlaena</i>	88
<i>intermedialis</i> (Walker), <i>Herculia</i>	58	<i>johnstoni</i> McDunnough, <i>Eupithecia</i>	78
[<i>intermediata</i> (Guenée), <i>Euphyia</i> <i>unangulata</i>]	76	<i>jucunda</i> Hübner, <i>Melipotis</i>	137
<i>interoceanica</i> (Smith), <i>Amphipoea</i>	148	<i>jucunda</i> (Walker), <i>Diarsia</i>	163
<i>interpuncta</i> (Grote), <i>Colobochyla</i>	136	<i>judith</i> Strecker, <i>Catocala</i>	139
<i>interpunctella</i> (Hübner), <i>Plodia</i>	63	<i>juglandis</i> (J. E. Smith), <i>Amorpha</i>	108
		<i>juglandis</i> (LeBaron), <i>Acrobasis</i>	59

<i>julia</i> (Barnes and McDunnough), <i>Idia</i> 134		<i>langtoni</i> Couper, <i>Alypia</i> 147	
[<i>julia</i> (Hulst), <i>Itame loricaria</i>] 80		<i>lanoraieensis</i> Sheppard, <i>Incisalia</i> 31	
<i>julianalis</i> (Walker), <i>Dicymolomia</i> 50		<i>Lapara</i> Walker 106	
<i>juncta</i> (Grote), <i>Chersotis</i> 163		<i>Laphygma</i> (Guenée) 154	
<i>junctaria</i> (Walker), <i>Scopula</i> 71		<i>laqueatellus</i> Clemens, <i>Crambus</i> 55	
<i>juniperata</i> (Linné), <i>Thera</i> 73		<i>Larentiinae</i> 71	
<i>Junonia</i> Hübner 37		<i>larentioides</i> Grote,	
<i>jutta</i> (Hübner), <i>Oeneis</i> 44		<i>Phalaenostola</i> 134	
<i>juvenalis</i> (Fabricius), <i>Erynnis</i> 16		<i>lariciata</i> (Freyer), <i>Eupithecia</i> 78	
K		<i>laricis</i> (Fitch), <i>Tolyte</i> 97	
<i>kalmiae</i> J. E. Smith, <i>Sphinx</i> 106		<i>larvaria</i> (Guenée), <i>Iridopsis</i> 84	
<i>kearfottella</i> Dyar, <i>Acrobasis</i> 59		<i>Lascoria</i> Walker 134	
<i>kellicottii</i> (Fish),		<i>Lasiocampidae</i> 97	
<i>Oidaematophorus</i> 65		<i>Lasiocampinae</i> 98	
<i>kentaria</i> Grote and Robinson,		<i>Lasionycta</i> Aurivillius 160	
<i>Selenia</i> 91		<i>lateritia</i> (Hufnagel), <i>Apamea</i> 149	
[<i>kolthoffi</i> Aurivillius,		<i>latex</i> (Guenée), <i>Morrisonia</i> 160	
<i>Sympistis</i>] 158		<i>laticinerea</i> Grote, <i>Lithophane</i> 156	
<i>Korscheltellus</i> Börner 9		<i>latiferrugata</i> (Walker),	
<i>kuehniella</i> (Zeller), <i>Anagasta</i> 63		<i>Eumacaria</i> 80	
<i>kuetzingi</i> (Grote), <i>Plagodis</i> 92		<i>latipennis</i> Stretch, <i>Spilosoma</i> 125	
L		<i>latipes</i> (Guenée), <i>Mocis</i> 140	
<i>labecula</i> (Grote), <i>Balsa</i> 149		<i>latiradiella</i> (Walker), <i>Catoptria</i> 56	
<i>labradorensis</i> (Packard),		<i>leachellus</i> (Zincken), <i>Crambus</i> 55	
<i>Xanthorhoe</i> 75		<i>lecontei</i> (Guérin-Ménéville),	
[<i>labradoriensis</i> Christoph, <i>Crambus</i>		<i>Haploa</i> 123	
<i>alienellus</i>] 55		<i>Ledaea</i> Druce 137	
<i>Lacanobia</i> Billberg 159		<i>legitima</i> (Grote), <i>Trichodestra</i> 162	
<i>Lacinipolia</i> McDunnough 159		<i>lemmeri</i> (Barnes and Benjamin),	
<i>Lacosoma</i> Grote 96		<i>Lithophane</i> 156	
[<i>Lacosomidae</i> = <i>Mimallonidae</i>] 96		<i>Lemmeria</i> Barnes and	
<i>lacrymose</i> Guenée, <i>Catocala</i> 139		<i>Benjamin</i> 152	
<i>laetodactylus</i> (Chambers),		<i>lentiflualis</i> (Zeller), <i>Aethiophysa</i> 50	
<i>Oidaematophorus</i> 65		<i>leo</i> (Grote), <i>Spragueia</i> 145	
<i>lacustrata</i> (Guenée), <i>Xanthorhoe</i> 75		[<i>leomegra</i> (Smith), <i>Polia</i>] 161	
[<i>lacustris</i> Munroe, <i>Microtheoris</i>		<i>leonardus</i> Harris, <i>Hesperia</i> 19	
<i>ophionalis</i>] 49		<i>lepida</i> Grote, <i>Lithophane</i> 156	
<i>ladon</i> (Cramer), <i>Celastrina</i> 32		<i>lepidula</i> (Grote), <i>Leuconycta</i> 144	
<i>laeta</i> (Edwards), <i>Erora</i> 32		<i>leporina</i> (Linné), <i>Acronicta</i> 146	
<i>laeta</i> Guérin-Ménéville,		<i>leptinoides</i> (Grote), <i>Schizura</i> 119	
<i>Holomelina</i> 124		<i>Leptostales</i> Möschler 71	
<i>laeta</i> (Morrison), <i>Archanara</i> 149		<i>lepusculina</i> Guenée, <i>Acronicta</i> 146	
<i>laetifica</i> Smith, <i>Acronicta</i> 146		[<i>Lethe</i> Hübner = <i>Enodia</i> Hübner,	
<i>laevigata</i> (Grote), <i>Polypogon</i> 134		<i>Satyrodes</i> Scudder] 42	
<i>lamae</i> (T. N. Freeman),		<i>Leucania</i> Ochsenheimer 160	
<i>Holomelina</i> 124		<i>leucitys</i> Franclemont,	
<i>Lambdina</i> Capps 93		<i>Symmerista</i> 117	
<i>Lampronia</i> Stephens 10		<i>leucobasis</i> (Dyar), <i>Caloreas</i> 14	
<i>lanceolaria</i> (Grote), <i>Acronicta</i> 146		<i>Leucobrephe</i> Grote 68	
		<i>Leucoma</i> Stephens 133	
		<i>Leuconycta</i> Hampson 144	
		<i>leucostigma</i> (Harris),	
		<i>Papaipema</i> 153	

[<i>leucostigma</i> (J. E. Smith), <i>Orgyia</i>]	132		<i>littera</i> Guenée, <i>Fagitana</i>	151
[<i>Lexis</i> auct. = <i>Eilema</i> Hübner]	121		<i>littoralis</i> (Packard), <i>Pardiarsia</i>	166
<i>libatrix</i> (Linné), <i>Scoliopteryx</i>	138		<i>lituralis</i> (Hübner), <i>Polypogon</i>	134
<i>liberaria</i> (Walker), <i>Apodrepanulatrix</i>	87		<i>lobeliae</i> Guenée, <i>Acronicta</i>	146
<i>liburna</i> (Geyer), <i>Scolecocampa</i>	138		<i>lobidactylus</i> (Fitch), <i>Trichoptilus</i>	64
<i>Libytheana</i> Michener	34		<i>Lobophora</i> Curtis	79
Libytheidae	34		Lobophorini	78
Libytheinae	34		<i>Lochmaeus</i> Doubleday	118
<i>lidia</i> (Cramer), <i>Euxoa</i>	164		<i>Lomanaltes</i> Grote	134
<i>lignicolor</i> (Walker), <i>Oligocentria</i>	119-20		<i>Lomographa</i> Hübner	86
<i>lignicolora</i> (Guenée), <i>Apamea</i>	149		<i>longa</i> Guenée, <i>Acronicta</i>	146
<i>lilacina</i> (Harvey), <i>Trichodestra</i>	162		<i>longirostrella</i> Clemens, <i>Donacula</i>	54
[<i>Limacodes</i> Berthold = <i>Apoda</i> Haworth]	46		<i>Longivesica</i> Hardwick	164
Limacodidae	45		<i>Lophocampa</i> Harris	130
<i>limata</i> (Grote and Robinson), <i>Pantographa</i>	54		<i>lorea</i> (Guenée), <i>Lacinipolia</i>	160
<i>limbata</i> (Haworth), <i>Nematocampa</i>	95		<i>loricaria</i> (Eversmann), <i>Itame</i>	80
<i>limboundata</i> (Haworth), <i>Scopula</i>	71		<i>louisiana</i> Forbes, <i>Macrochilo</i>	134
Limenitidinae	41		<i>Loxagrotis</i> McDunnough	165
Limenitidini	41		<i>Loxostege</i> Hübner	52
<i>Limenitis</i> Fabricius	41		<i>Loxostegopsis</i> Dyar	53
<i>limitaria</i> (Walker), <i>Caldara</i>	79		<i>lubricalis</i> (Geyer), <i>Idia</i>	134
<i>lineata</i> (Fabricius), <i>Hyles</i>	112		[<i>lucia</i> (Kirby), <i>Celastrina</i>]	33
<i>lineata</i> Newman and Donahue, <i>Phragmatobia</i>	127		<i>lucilius</i> (Scudder and Burgess), <i>Erynnis</i>	17
[<i>lineella</i> Grote, <i>Catocala</i>]	139		<i>lucifuga</i> (Denis and Schiffermüller), <i>Cucullia</i>	155
<i>Lineodes</i> Guenée	52		<i>luctuata</i> (Denis and Schiffermüller), <i>Spargania</i>	74
<i>lineola</i> (Ochsenheimer), <i>Thymelicus</i>	18		[<i>lugubris</i> Boisduval, <i>Thyris</i>]	64
<i>lineolata</i> Walker, <i>Catabena</i>	150		[<i>lulualis</i> (Hulst), <i>Loxostege</i> <i>anartalis</i>]	52
<i>lintneri</i> (Grote), <i>Gluphisia</i>	116		<i>luna</i> (Linné), <i>Actias</i>	102
<i>liparops</i> (Boisduval and LeConte), <i>Satyrium</i>	30		<i>lunata</i> (Drury), <i>Zale</i>	140
Liphyrinae	28		<i>lunifera</i> (Hübner), <i>Zale</i>	140
<i>Lipocosma</i> Lederer	50		<i>lunilinea</i> (Grote), <i>Spiloloma</i>	138
<i>Lipocosmodes</i> Munroe	50		<i>lunulalis</i> (Hulst), <i>Oneida</i>	58
[<i>liquoraria</i> Guenée, <i>Synchlora</i>]	69		<i>Luperina</i> Boisduval	152
<i>lisa</i> Boisduval and LeConte, <i>Eurema</i>	27		<i>luscitialis</i> Barnes and McDunnough, <i>Mecyna</i>	53
<i>Lithacodes</i> Packard	46		<i>luscitiosa</i> Clemens, <i>Sphinx</i>	106
<i>Lithacodia</i> Hübner	144		<i>lustralis</i> (Grote), <i>Lacinipolia</i>	160
Lithinini	90		<i>lutaiba</i> (Smith), <i>Polypogon</i>	134
<i>Litholomia</i> Grote	156		[<i>lutea</i> (Ström), <i>Xanthia</i>]	158
<i>Lithomoia</i> Hübner	156		[<i>luteola</i> (Grote and Robinson), <i>Anarta</i>]	158
<i>Lithophane</i> Hübner	156		<i>luteolellus</i> (Clemens), <i>Crambus</i>	55
<i>litophora</i> (Grote), <i>Macrochilo</i>	134		<i>lutosa</i> (Andrews), <i>Apamea</i>	149
Lithosiinae	121		<i>lutra</i> (Guenée), <i>Spiramater</i>	162
Lithosiini	121		<i>Lycaeides</i> Hübner	33
<i>lithospila</i> Grote, <i>Acronicta</i>	146		<i>Lycaena</i> Fabricius	28
			Lycaenidae	28
			Lycaeninae	28
			Lycaenopsini	32

Lycia Hübner 86
lyciades (Geyer), *Achalarus* 15
Lycomorpha Harris 122
[lygdamus (Doubleday), *Glaucopsyche*
lygdamus] 33
[Lygris Hübner = *Eulithis*
Hübner] 72
Lymantria Hübner 133
Lxymantriidae 131
Lymantriinae 133
Lymantriini 133
lynx (Guenée), *Schinia* 167
lyonsellus Haimbach, *Crambus* 55
lysimachiae Bird, *Papaipema* 153
Lytrosis Hulst 87

M

m-album (Boisduval and LeConte),
Parrhasius 31
maccullochii (Kirby),
Androloma 147
machaon Linné, *Papilio* 23-4
macmurtrei (Guérin-Ménéville),
Prionoxystus 15
macounii (Edwards), *Oeneis* 44
Macrochilo Hübner 134
Macroglossinae 109
Macroglossini 110
Macromphaliinae 97
Macronoctua Grote 152
Macrothyatirini 66
Macrurocampa Dyar 117
mactata (Guenée), *Oligia* 152
maculalis (Clemens), *Paraponyx* 48
maculalis Westwood, *Desmia* 53
macularia (Harris), *Sicya* 94
maculata Harris, *Lophocampa* 130
maculata Harris, *Thyris* 64
madefactalis (Guenée), *Hypena* 135
madusaria (Walker), *Euchlaena* 88
magnaria (Guenée), *Ennomos* 90
magniferalis (Walker), *Palpita* 54
magnificalis (Hübner),
Neocataclysta 48
magnoliata Guenée, *Spargania* 74
Magusa Walker 152
major Grote and Robinson,
Datana 114
major Smith, *Feralia* 155
majoralis (Smith), *Idia* 134
Malacosoma Hübner 98
Malacosomatini 98
malana (Fitch), *Balsa* 149

[malefactaria (Barnes and
McDunnough), *Enconista*
dislocaria] 82
Maliatha Walker 145
Mamestra Ochsenheimer 160
manalis (Walker), *Hypena* 135
[manataqua (Scudder), *Polites*
origenes] 20
mancalis (Lederer), *Hahncappsia* 51
[mancinus Doubleday and Hewitson,
Erebia disa] 44
[mandan (Edwards), *Carterocephalus*
palaemon] 18
Manduca Hübner 104
Manhatta Hulst 63
Maniolini 43
[manitoba McDunnough, *Dysstroma*
hersiliata] 72
[manitobae B. P. Clark, *Amorpha*
juglandis] 108
manitobana McDunnough,
Euxoa 165
[manitobensis McDunnough,
Macrurocampa marthesia] 118
[manitoboides (Fletcher),
Hesperia] 19
manteo Doubleday, *Lochmaeus* 118
mappa (Grote and Robinson),
Autographa 141
Marathyssa Walker 143
marcellus (Cramer), *Eurytides* 24
marculenta (Grote and Robinson),
Hahncappsia 51
marginalis (Walker), *Ostrinia* 51
marginaria (Minot), *Euchlaena* 88
[marginata (Edwards),
Celastrina] 33
marginaria (Harris), *Pennisetia* 12
marginidens (Guenée),
Papaipema 153
marina (Grote), *Oligia* 152
maritimalis McDunnough,
Tetralopha 59
[marmoratus (Grossbeck), *Pero*] 89
marthesia (Cramer),
Macrurocampa 117-8
martialis (Scudder), *Erynnis* 17
maryx (Guenée), *Sideridis* 161
massasoit (Scudder), *Poanes* 21
mathewianus (Zeller),
Oidaematophorus 65
meadi (Grote), *Schinia* 167
meadii (Packard), *Coryphista* 73
Mecyna Guenée 53

<i>meditata</i> (Grote), <i>Lacinipolia</i>	160	<i>miconympha</i> Guenée, <i>Catocala</i>	139
<i>Meganola</i> Dyar	143	Micropterigidae	9
[<i>Megasis</i> Guenée = <i>Sarata</i>		Micropterigoidea	9
Ragonot]	62	<i>Microtheoris</i> Meyrick	49
<i>Megisto</i> Hübner	43	<i>milberti</i> (Godart), <i>Nymphalis</i>	35
<i>Melanchra</i> Hübner	160	<i>militaris</i> (Harris), <i>Haploa</i>	123
<i>Melanolophia</i> Hulst	85	<i>militella</i> Zeller, <i>Tetralopha</i>	59
Melanolophiini	85	Mimallonidae	96
<i>Melanomma</i> Grote	137	<i>mimallonis</i> (Grote), <i>Euxoa</i>	164
<i>melanopa</i> (Thunberg), <i>Anarta</i>	158	Mimallonoidea	96
<i>melinella</i> Clemens, <i>Donacaula</i>	54	<i>minosaria</i> (Guenée), <i>Nemoria</i>	69
<i>melinus</i> Hübner, <i>Strymon</i>	31	<i>minerea</i> (Guenée), <i>Zale</i>	140
<i>Melipotis</i> Hübner	137	<i>minians</i> Guenée, <i>Nephelodes</i>	160
<i>melissa</i> (Edwards), <i>Lycaeides</i>	33	<i>miniata</i> (Kirby), <i>Hypoprepia</i>	122
<i>melissa</i> (Fabricius), <i>Oeneis</i>	44	<i>miniota</i> (Smith), <i>Blepharita</i>	154
Melitaeinae	39	<i>ministra</i> (Drury), <i>Datana</i>	114
<i>Melittia</i> Hübner	12	<i>minor</i> (Forbes), <i>Microcrambus</i>	56
Melittiini	12	<i>minorata</i> (Packard), <i>Semiothisa</i>	82
<i>Mellilla</i> Grote	81	<i>minualis</i> (Guenée), <i>Hypena</i>	135
<i>mellistrigata</i> (Grote), <i>Semiothisa</i>	82	<i>minuscula</i> (Morrison), <i>Oligia</i>	152
<i>mellonella</i> (Linné), <i>Galleria</i>	59	<i>minuscula</i> (Zeller), <i>Meganola</i>	143
<i>melsheimeri</i> (Harris), <i>Cicinnus</i>	96	<i>mira</i> Grote, <i>Catocala</i>	139
<i>mendica</i> (Walker), <i>Eubaphe</i>	77	<i>miranda</i> (Grote), <i>Athetis</i>	149
<i>mendosa</i> (McDunnough),		<i>miselioides</i> (Guenée), <i>Phosphila</i>	153
<i>Athetis</i>	149	<i>miserulata</i> Grote, <i>Eupithecia</i>	78
<i>mephisto</i> Franclemont, <i>Enargia</i>	151	<i>Misogada</i> Harvey	117
<i>meralis</i> (Morrison), <i>Caradrina</i>	150	<i>misturata</i> (Hulst), <i>Eupithecia</i>	78
[<i>meridionalis</i> Riotte, <i>Orgyia</i>]	132	<i>Mitoura</i> Scudder	30
<i>Meropleon</i> Dyar	152	<i>mixta</i> (Grote), <i>Apamea</i>	149
<i>Meroptera</i> Grote	61	[<i>Mniotype</i> Franclemont = <i>Blepharita</i>	
<i>merriccata</i> Dyar, <i>Paleacrita</i>	86	Hampson]	154
<i>merrickalis</i> (Barnes and McDunnough),		<i>Mocis</i> Hübner	140
<i>Loxostegopsis</i>	53	<i>modesta</i> (Harris), <i>Pachysphinx</i>	108
<i>meskei</i> Grote, <i>Catocala</i>	139	<i>modesta</i> (Hudson), <i>Furcula</i>	120, 121
<i>Mesoleuca</i> Hübner	74	<i>modesta</i> (Morrison), <i>Ulolonche</i>	162
<i>Mesothea</i> Warren	69	<i>modica</i> (Guenée), <i>Oligia</i>	152
<i>messoria</i> (Harris), <i>Euxoa</i>	164	<i>modica</i> Walker, <i>Acronicta</i>	146
[<i>metacomet</i> (Harris), <i>Euphyes</i>		<i>molliculata</i> (Walker), <i>Eulithis</i>	72
<i>ruricola</i>]	22	<i>mollis</i> Walker, <i>Agrotis</i>	162
<i>Metalectra</i> Hübner	137	<i>mollissima</i> (Guenée),	
<i>Metalepsis</i> Grote	165	<i>Callopietria</i>	149
<i>Metanema</i> Guenée	91	[<i>monetifera</i> (Guenée),	
<i>Metarrhantis</i> Warren	91	<i>Callopietria</i>]	149
<i>metatoides</i> McDunnough, <i>Zale</i>	140	<i>monochromatea</i> (Morrison),	
<i>Metaxaglaea</i> Franclemont	157	<i>Hemipachnobia</i>	165
<i>metonalis</i> (Walker),		<i>monodactyla</i> (Linné), <i>Emmelina</i>	65
<i>Phalaenostola</i>	134	Monotrysia	10
<i>Metrea</i> Grote	49	<i>montana</i> (Packard), <i>Syngrapha</i>	142
<i>mexicanum</i> (Boisduval), <i>Eurema</i>	27	<i>montana</i> (Walshingham), <i>Adaina</i>	65
<i>mexicanaria</i> (Grote), <i>Nacophora</i>	89	<i>monuste</i> (Linné), <i>Ascia</i>	25
<i>micacea</i> (Esper), <i>Hydraecia</i>	151	<i>Moodna</i> Hulst	62
<i>Microcrambus</i> Bleszynski	56	<i>morbidalis</i> (Guenée), <i>Chytolita</i>	133
<i>microgamma</i> (Hübner),		[<i>morpheus</i> (Fabricius),	
<i>Syngrapha</i>	142	<i>Phyciodes</i>]	39

<i>morpheus</i> (Hufnagel), <i>Caradrina</i> 150		<i>nearctica</i> Munroe, <i>Nomophila</i> 53	
<i>morrisonaria</i> (Hy. Edwards), <i>Pero</i> 89		[<i>nearctica</i> Ferguson, <i>Syngrapha</i> <i>microgramma</i>] 142	
<i>morrisoni</i> (Grote), <i>Eupsilia</i> 155		<i>nebris</i> (Guenée), <i>Papaipema</i> 153	
<i>Morrisonia</i> Grote 160		<i>nebulifera</i> Stephens, <i>Prionapteryx</i> 54	
<i>morula</i> Grote and Robinson, <i>Acronicta</i> 146		<i>nebulosa</i> Edwards, <i>Catocala</i> 139	
[<i>mucidella</i> (Ragonot), <i>Phycitodes</i>] 62		<i>necopina</i> (Grote), <i>Papaipema</i> 153	
<i>multifera</i> (Walker), <i>Caradrina</i> 150		<i>Nedra</i> Clarke 152	
<i>multiferata</i> (Walker), <i>Anticlea</i> 74		[<i>neglecta</i> (Edwards), <i>Celastrina</i>] 33	
<i>multilinea</i> Walker, <i>Leucania</i> 160		<i>nelita</i> (Strecker), <i>Papaipema</i> 153	
<i>multipunctellus</i> Clemens, <i>Yponomeuta</i> 12		<i>Nematocampa</i> Guenée 95	
[<i>multiscripta</i> Riley, <i>Cerura</i> <i>scitisscripta</i>] 121		<i>Nemoria</i> Hübner 69	
<i>munitata</i> (Hübner), <i>Xanthorhoe</i> 75		<i>Nemoriini</i> 69	
<i>Munroessa</i> Lange 48		<i>neobliteralis</i> Capps, <i>Hahncappsia</i> 51	
<i>muscosa</i> (Guenée), <i>Lithacodia</i> 145		<i>Neocataclysta</i> Lange 48	
[<i>mustelinus</i> (Packard), <i>Korscheltellus</i>] 9		<i>neogama</i> (J. E. Smith), <i>Catocala</i> 139	
<i>mutabilis</i> (Clemens), <i>Fissicrambus</i> 56		[<i>neomarsyas</i> dos Passos, <i>Polygonia</i> <i>satyrus</i>] 35	
<i>mutata</i> Pearsall, <i>Eupithecia</i> 78		<i>nepheleptena</i> (Dyar), <i>Papaipema</i> 153	
<i>Mutuuraia</i> Munroe 51		<i>Nephelodes</i> Guenée 160	
<i>muzaria</i> (Walker), <i>Euchlaena</i> 88		<i>Nephoterix</i> Hübner 61	
<i>Mycterophora</i> Hulst 137		<i>Nephrogramma</i> Munroe 49	
<i>Myelopsis</i> Heinrich 60		<i>neptaria</i> (Guenée), <i>Semiothisa</i> 82	
<i>mynesalis</i> (Walker), <i>Tetanolita</i> 135		<i>Nepticulidae</i> 10	
<i>myops</i> (J. E. Smith), <i>Paonias</i> 107-8		<i>Nepticuloidea</i> 10	
<i>myricella</i> McDunnough, <i>Ortholepis</i> 60		<i>Nepytia</i> Hulst 94	
[<i>myrina</i> (Cramer), <i>Boloria</i> <i>selene</i>] 39		<i>Nerice</i> Walker 116	
<i>myron</i> (Cramer), <i>Darapsa</i> 111		[<i>nessus</i> Cramer, <i>Sphinx</i>] 110	
<i>mysippusalis</i> (Walker), <i>Mutuuraia</i> 51		[<i>nessus</i> Drury, <i>Theretra</i>] 110	
<i>mystic</i> (Edwards), <i>Polites</i> 19-20		<i>nevadae</i> (Grote), <i>Lacanobia</i> 159	
N		<i>nexilis</i> Morrison, <i>Eutricopis</i> 167	
[<i>nabokovi</i> Masters, <i>Lycaeides</i> <i>idas</i>] 33		<i>ni</i> (Hübner), <i>Trichoplusia</i> 142	
<i>Nacophora</i> Hulst 89		<i>nicalis</i> Grote, <i>Pyrausta</i> 52	
<i>Nacophorini</i> 89		<i>nicippe</i> (Cramer), <i>Eurema</i> 27-8	
<i>Nadata</i> Walker 114		<i>nigella</i> (Strecker), <i>Dyspyralis</i> 133	
<i>nais</i> (Drury), <i>Apantesis</i> 127		<i>Nigetia</i> Walker 145	
<i>napaea</i> (Morrison), <i>Litholomia</i> 156		<i>nigrinodis</i> (Zeller), <i>Galasa</i> 58	
[<i>napi</i> (Linné), <i>Pieris</i>] 25		<i>nigrior</i> (Smith), <i>Apamea</i> 149	
[<i>napi oleracea</i> Harris, <i>Pieris</i>] 25		<i>nigrovittella</i> Dyar, <i>Immyrta</i> 60	
<i>Nascia</i> Curtis 50		<i>nimbicolor</i> (Hulst), <i>Eupithecia</i> 78	
<i>Nathalis</i> Boisduval 28		<i>nimboza</i> (Guenée), <i>Polia</i> 161	
<i>Nealgedonia</i> Munroe 51		<i>niphon</i> (Hübner), <i>Incisalia</i> 30, 31	
		[<i>nipisiquit</i> McDunnough, <i>Coenonympha</i>] 43	
		<i>nivalis</i> (Drury), <i>Urola</i> 56	
		<i>niveivenosa</i> (Grote), <i>Apamea</i> 149	
		<i>nivigerata</i> Walker, <i>Lobophora</i> 79	
		<i>nivosaria</i> (Guenée), <i>Eugonobapta</i> 94	
		<i>noctivaga</i> Grote, <i>Acronicta</i> 146	
		<i>Noctuidae</i> 133	

Noctuinae	162	<i>oblonga</i> (Grote), <i>Archanara</i>	149
Noctuoidea	112	<i>obscura</i> (Fabricius), <i>Erinnyis</i>	109
[<i>nokomis</i> (Brodie),		<i>obscura</i> (Smith), <i>Orthodes</i>	161
<i>Hyalophora</i>]	103	<i>obscura</i> Strecker, <i>Catocala</i>	139
<i>Nola</i> Leach	144	<i>obscuralis</i> (Grote), <i>Paraponyx</i>	48
Nolinae	143	<i>obscurata</i> Strecker, <i>Schinia</i>	167
<i>Nomophila</i> Hübner	53	<i>obscuripennis</i> (Grote),	
<i>normani</i> (Grote), <i>Crocigrapha</i>	159	<i>Polypogon</i>	134
<i>normaniana</i> (Grote), <i>Xestia</i>	166	<i>obstipata</i> (Fabricius),	
<i>notataria</i> (Walker), <i>Eufidonia</i>	85	<i>Orthonama</i>	76
<i>notialis</i> Franclemont, <i>Tolype</i>	97	<i>obtusa</i> Smith, <i>Oligia</i>	152
<i>Notodonta</i> Ochseneheimer	115	<i>obtusaria</i> (Hübner), <i>Euchlaena</i>	88
Notodontidae	112	<i>obumbratalis</i> (Lederer), <i>Ostrinia</i>	51
Notodontinae	114	<i>Occidentalia</i> Dyar and Heinrich	57
Notodontini	114	<i>occidentalis</i> Lintner, <i>Furcula</i>	121
[<i>nova</i> Fitch, <i>Orgyia antiqua</i>]	132	<i>occidentalis</i> (Reakirt), <i>Pontia</i>	24
<i>novaroensis</i> (Hy. Edwards),		<i>occiduarua</i> (Packard), <i>Itame</i>	80
<i>Synanthedon</i>	13	<i>occiduarua</i> (Walker), <i>Anagoga</i>	92
<i>noviganna</i> (Barnes and Benjamin),		<i>ocellinata</i> (Guenée), <i>Semiothisa</i>	82
<i>Gazorycta</i>	10	<i>ochreipennis</i> (Grote),	
<i>nubilalis</i> (Hübner), <i>Ostrinia</i>	51	<i>Polypogon</i>	135
<i>nubilis</i> (Hübner), <i>Euparthenos</i>	140	<i>ochrifrontella</i> (Zeller), <i>Eulogia</i>	62
<i>numitor</i> (Fabricius),		<i>ochrogaster</i> (Guenée), <i>Euxoa</i>	165
<i>Ancyloxypha</i>	18	<i>Ochroleura</i> Hübner	166
<i>nundina</i> (Drury), <i>Schinia</i>	167	<i>octo</i> (Guenée), <i>Amyna</i>	144
<i>nupera</i> (Lintner), <i>Xylena</i>	158	<i>octomaculata</i> (Fabricius),	
<i>nycteis</i> (Doubleday), <i>Charidryas</i>	40	<i>Alypia</i>	147
<i>Nycteola</i> Hübner	143	<i>octoscripta</i> (Grote), <i>Syngrapha</i>	142
[<i>Nycterosea</i> Hulst = <i>Orthonama</i>		<i>oculatrix</i> (Guenée), <i>Paectes</i>	143
Hübner]	76	Odontiinae	49
Nymphalidae	34	Odontiini	49
Nymphalinae	34	<i>Odontesia</i> Hübner	115
Nymphalini	34	<i>odorata</i> (Linné), <i>Ascalapha</i>	136
<i>Nymphalis</i> Kluk	35	<i>Oeneis</i> Hübner	44
<i>Nymphula</i> Schrank	48	Oenochrominae	68
Nymphulinae	48	<i>Ogdoconta</i> Butler	152
Nymphulini	48	<i>Oidaematophorus</i> Wallengren	65
<i>nyssaria</i> (Guenée), <i>Probole</i>	92	<i>oithona</i> (Strecker), <i>Grammia</i>	128
O		<i>Olceclostera</i> Butler	96
<i>Oarisma</i> Scudder	18	<i>oleracea</i> (Harris), <i>Pieris</i>	25
[<i>obductata</i> (Möschler), <i>Spargania</i>		<i>Oligia</i> Hübner	152
<i>luctuata</i>]	74	<i>Oligocentria</i> Herrich-Schäffer	119
<i>obeliscoides</i> (Guenée), <i>Euxoa</i>	165	<i>olinalis</i> (Guenée), <i>Herculia</i>	58
<i>obfirmaria</i> (Hübner),		<i>olivacea</i> (Morrison), <i>Lacinipolia</i>	160
<i>Metarrhantis</i>	91	<i>olympia</i> (Edwards), <i>Euchloe</i>	26
<i>oblata</i> (Morrison), <i>Xestia</i>	166	<i>omissa</i> Dod, <i>Cucullia</i>	155
<i>oblinita</i> (J. E. Smith), <i>Acronicta</i>	146	<i>Oncocnemis</i> Lederer	157
<i>obliqua</i> Packard, <i>Heterocampa</i>	118	<i>Oneida</i> Hulst	58
<i>obliqua</i> (Smith), <i>Agrotis</i>	162	<i>ontario</i> (Edwards), <i>Euristrymon</i>	31
<i>obliqua</i> (Walker), <i>Bellura</i>	149	<i>ontario</i> (McDunnough),	
<i>obliquata</i> (Grote and Robinson),		<i>Sphenarches</i>	64
<i>Dasychira</i>	131	[<i>ontario</i> (Smith), <i>Euxoa</i>]	164
		<i>onusta</i> Grote, <i>Macronoctua</i>	152
		<i>onustana</i> (Walker), <i>Tebenna</i>	14

<i>[opacifrons</i> (Grote), <i>Eugraphe</i>]	164
<i>opella</i> (Grote), <i>Holomelina</i>	124
<i>Operophtera</i> Hübner	77
Operophterini	77
Ophiderinae	136
<i>ophionalis</i> (Walker), <i>Microtheoris</i>	49
<i>[Ophiuche</i> Hübner = <i>Hypena</i> Schrank]	135
<i>ophthalmica</i> (Guenée), <i>Baileya</i>	143
<i>orbifera</i> (Walker), <i>Magusa</i>	152
<i>orciferalis</i> Walker, <i>Macrochilo</i>	134
<i>orciferarius</i> (Walker), <i>Aspilates</i>	87
<i>Oreana</i> Hulst	60
<i>oregonensis</i> (Stretch), <i>Cycnia</i>	130
<i>Oreta</i> Walker	68
Oretinae	68
<i>Orgyia</i> Ochsenheimer	132
Orgyiinae	131
Orgyiini	131
<i>[orientis</i> Ferguson, <i>Itame</i> <i>andersoni</i>]	80
<i>origenes</i> (Fabricius), <i>Polites</i>	20
<i>orillata</i> (Walker), <i>Semiothisa</i>	82
<i>oriunda</i> Grote, <i>Lithophane</i>	156
<i>ornithogalli</i> Guenée, <i>Spodoptera</i>	154
<i>[Orosagrotis</i> Hampson]	165
<i>orphisalis</i> Walker, <i>Pyrausta</i>	52
<i>Orthodes</i> Guenée	160
<i>Orthofidonia</i> Packard	83
<i>Ortholepis</i> Ragonot	60
<i>Orthonama</i> Hübner	76
<i>Orthosia</i> Ochsenheimer	161
<i>ostreonalis</i> Grote, <i>Metrea</i>	49
<i>ostrinella</i> (Clemens), <i>Moodna</i>	62
<i>Ostrinia</i> Hübner	51
<i>ostryella</i> Ely, <i>Acrobasis</i>	59
<i>ou</i> (Guenée), <i>Rachiplusia</i>	142
Ourapterygini	93
<i>ovalis</i> (Packard), <i>Telethusia</i>	61
<i>ovata</i> Grote, <i>Acrionicta</i>	146
<i>oviduca</i> (Guenée), <i>Protorthodes</i>	161
<i>ovilla</i> Grote, <i>Nola</i>	144
<i>oviplagalis</i> (Walker), <i>Tosale</i>	58
<i>oweni</i> (Swett), <i>Semiothisa</i>	82
<i>oxygala</i> (Grote), <i>Aletia</i>	158
<i>oxygramma</i> (Geyer), <i>Agrapha</i>	141
<i>Oxyptilus</i> Zeller	64

P

Pachysphinx Rothschild and
 Jordan 108

<i>packardata</i> McDunnough, <i>Xanthorhoe</i>	75
<i>Packardia</i> Grote and Robinson	46
<i>padellus</i> (Linné), <i>Yponomeuta</i>	12
<i>Paectes</i> Hübner	143
<i>palaemon</i> (Pallas), <i>Carterocephalus</i>	17-8
<i>palaeno</i> (Linné), <i>Colias</i>	26
<i>palaegama</i> Guenée, <i>Catocala</i>	139
<i>Paleacrita</i> Riley	86
<i>palliatricula</i> (Guenée), <i>Chytonix</i>	150
<i>pallida</i> (Herrich-Schäffer), <i>Tortricidia</i>	46
<i>pallida</i> Packard, <i>Crambida</i>	122
<i>pallidactyla</i> (Haworth), <i>Platyptilia</i>	65
<i>pallidata</i> (Hufnagel), <i>Evergestis</i>	50
<i>palliolella</i> Ragonot, <i>Acrobasis</i>	59
<i>palparia</i> (Walker), <i>Hypena</i>	135
<i>palpata</i> Packard, <i>Eupithecia</i>	78
<i>Palpia</i> Hübner	54
<i>Palthis</i> Hübner	134
<i>palustris</i> Ferguson, <i>Hyperodes</i>	135
<i>pampina</i> Guenée, <i>Eucirroedia</i>	155
<i>pampinaria</i> (Guenée), <i>Anavitrinella</i>	84
<i>panalope</i> (Dyar), <i>Xubida</i>	57
<i>panatala</i> (Smith), <i>Spartiniphaga</i>	154
<i>pandora</i> (Hübner), <i>Eumorpha</i>	110
<i>Pangrapta</i> Hübner	137
<i>Panopoda</i> Guenée	137
<i>Panthea</i> Hübner	145
Pantheinae	145
<i>Pantographa</i> Lederer	54
<i>Paonias</i> Hübner	107
<i>Papaipema</i> Smith	153
<i>Papestra</i> Sukhareva	161
<i>Papilio</i> Linné	23
Papilionidae	23
Papilioninae	23
Papilionini	23
Papilionoidea	22
<i>Paraclemensia</i> Busck	10
<i>Paradiarsia</i> McDunnough	166
<i>paradoxus</i> Grote, <i>Heliothis</i>	167
<i>[Parahypenodes</i> Barnes and McDunnough = <i>Hypena</i> Schrank]	135
<i>pariana</i> (Clemens), <i>Choreutis</i>	14
<i>Parallelia</i> Hübner	140
<i>Paranthrene</i> Hübner	12
Paranthreninae	12
Paranthrenini	12

<i>Parapediasia</i> Bleszynski	56	<i>Pero</i> Herrich-Schäffer	89
<i>Paraponyx</i> Hübner	48	[<i>Perophoridae</i> = <i>Mimallonidae</i>]	96
<i>Parargini</i>	42	<i>perpolita</i> (Morrison), <i>Euxoa</i>	165
<i>Parasemia</i> Hübner	124	<i>perquiritata</i> (Morrison), <i>Xestia</i>	166
<i>Parastichtis</i> Hübner	157	[<i>persephone</i> (Grote), <i>Grammia</i>]	128
<i>pariana</i> (Clemens), <i>Choreutis</i>	14	<i>persius</i> (Scudder), <i>Erynnis</i>	17
<i>Parrhasius</i> Hübner	31	<i>perspectalis</i> (Hübner), <i>Hymenia</i>	53
<i>parta</i> Guenée, <i>Catocala</i>	139	<i>perspicua</i> Grote and Robinson,	
<i>parthenice</i> (Kirby), <i>Grammia</i>	128	<i>Datana</i>	114
<i>parthenos</i> (Harris), <i>Platarctia</i>	127	<i>pertextalis</i> (Lederer),	
<i>participalis</i> Grote, <i>Condylolomia</i>	58	<i>Herpetogramma</i>	54
<i>paritita</i> Guenée, <i>Galgula</i>	151	<i>pervertens</i> Barnes and McDunnough,	
<i>parvulella</i> (Ely), <i>Erelieva</i>	63	<i>Hyperstrotia</i>	144
<i>pasadamia</i> (Dyar), <i>Oritholepis</i>	60	<i>petrealis</i> Grote, <i>Chytolita</i>	133
<i>pascoensis</i> Wright, <i>Phyciodes</i>	39-40	<i>Petrophila</i> Guilding	48
<i>pascuellus</i> (Linné), <i>Crambus</i>	55	<i>Petrophora</i> Hübner	90
<i>passer</i> (Guenée), <i>Luperina</i>	152	<i>pettiti</i> (Grote), <i>Pyreferra</i>	157
<i>patefacta</i> (Walker), <i>Lithophane</i>	156	<i>petulca</i> Grote, <i>Lithophane</i>	156
[<i>peckius</i> (Kirby), <i>Polites</i>]	19	<i>pexata</i> Grote, <i>Lithophane</i>	156
<i>Pediasia</i> Hübner	56	<i>pexella</i> (Zeller), <i>Thaumatopsis</i>	56
<i>pedipilalis</i> (Guenée), <i>Polypogon</i>	135	<i>phaceliae</i> McDunnough,	
<i>pegala</i> (Fabricius), <i>Cercyonis</i>	43-4	<i>Oidaematophorus</i>	65
<i>pegasalis</i> (Walker), <i>Chalcoela</i>	50	<i>Phaegopterini</i>	129
<i>pelidne</i> Boisduval and LeConte,		<i>phaeocapna</i> Franclemont, <i>Zale</i>	140
<i>Colias</i>	26	<i>phaeton</i> (Drury), <i>Euphydryas</i>	40
<i>pellucidaria</i> (Packard), <i>Nepytia</i>	94	<i>Phalaenophana</i> Grote	134
<i>pendulinaria</i> (Guenée),		<i>Phalaenostola</i> Grote	134
<i>Cyclophora</i>	70	<i>phalerata</i> (Harris), <i>Apantesis</i>	127
<i>penitalis</i> (Grote), <i>Ostrinia</i>	51	<i>Phalerinae</i>	113
<i>Pennisetia</i> Dehne	12	<i>Pheosia</i> Hübner	115
<i>Pennisetiini</i>	12	<i>Phigalia</i> Duponchel	86
[<i>pennsylvanica</i> Smith,		<i>Philampelini</i>	110
<i>Euthyatira</i>]	67	<i>philea</i> (Johansson), <i>Phoebis</i>	27
<i>Penstemonia</i> Engelhardt	14	<i>philenor</i> (Linné), <i>Battus</i>	23
<i>penumbralis</i> Dyar, <i>Scoparia</i>	48	<i>philodice</i> Godart, <i>Colias</i>	26-7
<i>Peoria</i> Ragonot	63	[<i>Philometra</i> Grote = <i>Phalaenostola</i>	
<i>Peoriinae</i>	63	Grote]	134
<i>Peratrea</i> Grote	105	<i>phlaeas</i> (Linné), <i>Lycaena</i>	29
<i>perattenta</i> (Grote),		<i>Phlogophora</i> Treitschke	153
<i>Eueretagrotis</i>	164	<i>phlogosaria</i> (Guenée), <i>Plagodis</i>	92
<i>perditalis</i> (Walker), <i>Ledaea</i>	137	<i>phloxiphagus</i> Grote and Robinson,	
<i>perfracta</i> Swett, <i>Hydriomena</i>	73	<i>Heliothis</i>	167
<i>perfusca</i> (Hulst), <i>Eupithecia</i>	78	<i>Phoberia</i> Hübner	137
<i>pergilvalis</i> (Hulst), <i>Hahncappsia</i>	51	<i>Phobetritini</i>	46
<i>periculosa</i> Guenée,		<i>Phobetron</i> Hübner	46
<i>Phlogophora</i>	153	<i>Phoebis</i> Hübner	27
<i>Peridea</i> Stephens	114	<i>Pholisora</i> Scudder	17
<i>Peridroma</i> Hübner	166	<i>pholus</i> (Cramer), <i>Darapsa</i>	111
<i>Perigea</i> Guenée	153	<i>pholus</i> (Drury), <i>Lycomorpha</i>	122
<i>periscelidactyla</i> Fitch, <i>Geina</i>	64	<i>Phosphila</i> Hübner	153
<i>Perispasta</i> Zeller	51	<i>phragmatidicola</i> Guenée,	
<i>Perizoma</i> Hübner	74	<i>Leucania</i>	160
<i>perlata</i> (Guenée), <i>Campaea</i>	89	<i>Phragmatobia</i> Stephens	126
<i>perlellus</i> (Scopoli), <i>Crambus</i>	55	<i>Phyciodes</i> Hübner	39

Phycitinae	59	<i>pleuritica</i> (Grote), <i>Euxoa</i>	164
<i>Phycitodes</i> Hampson	62	<i>Pleuroprucha</i> Möschler	70
<i>phyleus</i> (Drury), <i>Hylephila</i>	19	<i>plevie</i> (Dyar), <i>Eoparargyractis</i>	49
<i>phyllira</i> (Drury), <i>Grammia</i>	129	<i>plexippus</i> (Linné), <i>Danaus</i>	45
<i>Phyllodesma</i> Hübner	97	<i>Plodia</i> Guenée	63
<i>phyllophora</i> (Grote),		[<i>plumbea</i> Hy. Edwards,	
<i>Heptagrotis</i>	165	<i>Hypoprepia</i>]	122
<i>Phytometra</i> Haworth	137	<i>plumbea</i> Stretch, <i>Cisthene</i>	122
<i>piatrix</i> Grote, <i>Catocala</i>	139	<i>Plusia</i> Ochsenheimer	142
<i>pica</i> (Walshingham), <i>Amblyptilia</i>	65	<i>plusiiformis</i> Hy. Edwards,	
<i>picta</i> (Harris), <i>Melanchra</i>	160	<i>Stretchia</i>	162
<i>pictipes</i> (Grote and Robinson),		<i>Plusiinae</i>	141
<i>Synanthedon</i>	13	<i>Plusiodonta</i> Guenée	138
Pieridae	24	<i>plutonia</i> (Grote), <i>Apamea</i>	149
Pierinae	24	<i>Poanes</i> Scudder	21
Pierini	24	[<i>pocahontas</i> (Scudder), <i>Poanes</i>]	21
<i>Pieris</i> Schrank	25	<i>Podosesia</i> Möschler	13
<i>piffardi</i> (Walker), <i>Oncocnemis</i>	157	<i>poecila</i> Stephens, <i>Sphinx</i>	106
<i>pini</i> (Kellicott), <i>Synanthedon</i>	13	<i>polata</i> (Duponchel), <i>Entephria</i>	74
[<i>pini</i> Michener, <i>Eacles</i>		<i>polia</i> (Cook and Watson),	
<i>imperialis</i>]	99	<i>Incisalia</i>	31
<i>piniata</i> (Packard), <i>Caripeta</i>	93	<i>Polia</i> Ochsenheimer	161
<i>piniata</i> (Packard), <i>Hypagyrtis</i>	84	<i>polistiformis</i> (Harris), <i>Vitacea</i>	12
<i>pinicola</i> (Dyar), <i>Dasychira</i>	131	<i>Polites</i> Scudder	19
<i>pinistrobata</i> Ferguson,		<i>Polygonia</i> Hübner	34
<i>Semiothisa</i>	82	<i>polygrammaria</i> (Packard),	
<i>pistasciaria</i> (Guenée), <i>Hethemia</i>	69	<i>Stenoporpia</i>	83
<i>pithecium</i> (J. E. Smith),		<i>Polygrammate</i> Hübner	147
<i>Phobetron</i>	46-7	<i>Polygrammodes</i> Guenée	54
<i>pityochromus</i> Grote,		<i>Polyommata</i> inae	32
<i>Plagiomimicus</i>	153	<i>Polyommata</i> ni	33
<i>placida</i> (Grote), <i>Abagrotis</i>	162	<i>polyphemus</i> (Cramer),	
<i>plagiata</i> (Walker), <i>Dasychira</i>	131	<i>Antheraea</i>	101
<i>plagiata</i> (Walker), <i>Orgyia</i>	132	<i>Polypogon</i> Schrank	134
<i>Plagiomimicus</i> Grote	153	<i>polyxenes</i> Fabricius, <i>Papilio</i>	24
<i>Plagodis</i> Hübner	92	<i>pometaria</i> (Harris), <i>Alsophila</i>	68
<i>plantaginis</i> (Linné), <i>Parasemia</i>	113,	<i>Pompeius</i> Evans	20
124		<i>Pontia</i> Fabricius	24
<i>Platartectia</i> Packard	127	<i>populella</i> Busck, <i>Ectoedemia</i>	10
[<i>Plathypena</i> Grote = <i>Hypena</i>		<i>populi</i> (Walker), <i>Acossus</i>	15
Schrank]	135	<i>porcelaria</i> (Guenée),	
<i>Platypolia</i> Grote	157	<i>Protoboarmia</i>	85
<i>Platyptilia</i> Hübner	65	<i>postera</i> Guenée, <i>Cucullia</i>	155
Platyptilinae	64	<i>praeclara</i> Grote and Robinson,	
[<i>Platysenta</i> Grote = <i>Condica</i>		<i>Catocala</i>	139
Walker]	150	<i>praeffectellus</i> (Zincken), <i>Crambus</i>	55
<i>Platytes</i> Guenée	55	<i>prasina</i> (Denis and Schiffermüller),	
<i>plebeja</i> (Fabricius), <i>Peratrea</i>	105	<i>Anaplectoides</i>	163
<i>Plebejus</i> Kluk	33	<i>pravella</i> (Grote), <i>Meroptera</i>	61
<i>plecta</i> (Linné), <i>Ochrolepura</i>	166	<i>precationis</i> (Guenée),	
<i>plejadellus</i> Zincken, <i>Chilo</i>	57	<i>Autographa</i>	141
<i>Plemyria</i> Hübner	72	[<i>preciosa</i> Nixon, <i>Grammia</i>]	128
<i>Pleonectopoda</i> Grote	164	<i>pressus</i> (Grote), <i>Anaplectoides</i>	163
<i>pleonectusa</i> Grote, <i>Ipimorpha</i>	152	<i>primordialis</i> (Dyar), <i>Anageshna</i>	53

[<i>primum</i> Slosson, <i>Spilosoma</i>]	125	<i>Psychomorpha</i> Harris	147
[<i>Priocyclus</i> Guenée = <i>Cepphis</i> Hübner]	92	<i>pterisii</i> Bird, <i>Papaipema</i>	153
<i>Prionapteryx</i> Stephens	54	Pterophoridae	64
<i>Prionoxystus</i> Grote	15	Pterophorinae	65
<i>privata</i> (Walker), <i>Sutyna</i>	158	Pterophoroidea	64
<i>Probole</i> Herrich-Schäffer	92	<i>pudens</i> (Guenée), <i>Euthyatira</i>	66-7
<i>Prochoerodes</i> Grote	95	<i>pulcherrima</i> (Grote), <i>Eutelia</i>	143
<i>Prochoreutis</i> Diakonoff and Heppner	14	[<i>pulverea</i> Grote and Robinson, <i>Heterocampa umbrata</i>]	118
<i>Proclissiana</i> Reuss, see <i>Boloria</i> Moore	38	<i>pulverulenta</i> (Smith), <i>Melanchra</i>	160
<i>profunda</i> (Smith), <i>Sutyna</i>	158	<i>punctella</i> (Cramer), <i>Atteva</i>	11
<i>progne</i> (Cramer), <i>Polygonia</i>	35	<i>puncticosta</i> (Smith), <i>Dyspyralis</i>	133
<i>projecta</i> (Walker), <i>Cleora</i>	84	<i>pupula</i> Hübner, <i>Eustixia</i>	49
<i>Prolimacodes</i> Schaus	46	<i>pura</i> Barnes and McDunnough, <i>Crambidia</i>	122
<i>promethea</i> (Drury), <i>Callosamia</i>	102-3	<i>purata</i> (Guenée), <i>Scopula</i>	71
<i>Properigea</i> Barnes and Benjamin	157	<i>purpurascens</i> (Packard), <i>Sthenopsis</i>	9
<i>propinqualis</i> Guenée, <i>Rivula</i>	138	<i>purpurea</i> Barnes and McDunnough, <i>Egira</i>	159
<i>propinquilinea</i> (Grote), <i>Colocasias</i>	145	[<i>purpurea</i> McDunnough, <i>Apharetra</i>]	154
<i>propulsata</i> (Walker), <i>Eulithis</i>	72	<i>purpurea</i> Walker, <i>Adela</i>	10
<i>Proserpinus</i> Hübner	111	<i>purpurigera</i> (Walker), <i>Pseudeva</i>	142
<i>Protiame</i> McDunnough	80	<i>purpurissata</i> (Grote), <i>Polia</i>	161
<i>Protoboarmia</i> McDunnough	85	<i>pustularia</i> (Guenée), <i>Itame</i>	80-1
<i>protodice</i> (Boisduval and LeConte), <i>Pontia</i>	24	<i>puta</i> (Grote and Robinson), <i>Anathix</i>	154
<i>Protolampra</i> McDunnough	166	<i>putnami</i> (Grote), <i>Plusia</i>	142
<i>Proorthodes</i> McDunnough	161	Pygaerinae	112
<i>protumnusalis</i> (Walker), <i>Polypogon</i>	135	<i>Pyla</i> Grote	61
[<i>Proxenus</i> Herrich-Schäffer = <i>Athetis</i> Hübner]	149	<i>pylades</i> (Scudder), <i>Thorybes</i>	16
<i>proxima</i> (Hy. Edwards), <i>Synanthedon</i>	13	Pyalidae	47
<i>Psaphida</i> Walker	157	<i>pyralidiformis</i> (Walker), <i>Carmenta</i>	14
<i>Psectraglaea</i> Hampson	157	Pyraliformes	57
<i>Pseudaletia</i> Franclemont	161	Pyralinae	57
[<i>pseudargiolus</i> (Boisduval and LeConte), <i>Celastrina</i>]	32	<i>Pyralis</i> Linné	57
<i>pseudargyria</i> Guenée, <i>Leucania</i>	160	Pyraloidea	47
<i>Pseudeva</i> Hampson	142	<i>pyramidalis</i> (Walker), <i>Albuna</i>	12
<i>pseudogamma</i> (Grote), <i>Autographa</i>	141	<i>pyramidoides</i> Guenée, <i>Amphipyra</i>	148
<i>Pseudoplusia</i> McDunnough	142	<i>pyramusalis</i> (Walker), <i>Phalaenophana</i>	134
<i>Pseudorthodes</i> McDunnough	161	<i>Pyrausta</i> Schrank	52
<i>Pseudothyatira</i> Grote	66	Pyraustinae	50
<i>Psorosina</i> Dyar	62	Pyraustini	50
<i>Psyche</i> Schrank	11	<i>Pyreferra</i> Franclemont	157
<i>psychialis</i> (Hulst), <i>Xanthophysa</i>	50	Pyrginae	15
Psychidae	11	<i>Pyrgus</i> Hübner	17
Psychinae	11	<i>pyri</i> (Harris), <i>Synanthedon</i>	13
		<i>Pyrrharctia</i> Packard	124
		<i>Pyrrhia</i> Hübner	167

Q

<i>quadrals</i> Barnes and McDunnough,	
<i>Hypena</i>	135
<i>quadrata</i> (Grote), <i>Acronicta</i>	146
<i>quadrata</i> (Grote), <i>Papestra</i>	161
<i>quadriguttatus</i> (Grote), <i>Sthenopis</i>	9
<i>quadrilineata</i> (Packard), <i>Scopula</i>	71
<i>quadrisingnata</i> Walker,	
<i>Metalectra</i>	137
<i>quadristigmalis</i> (Guenée),	
<i>Palpita</i>	54
<i>quebecensis</i> (Smith), <i>Euxoa</i>	164
<i>quenseli</i> (Paykull), <i>Grammia</i>	129
<i>quercivoraria</i> (Guenée), <i>Besma</i>	93
<i>quernaria</i> (J. E. Smith),	
<i>Nacophora</i>	89
<i>querquera</i> Grote, <i>Lithophane</i>	156
<i>quinquemaculata</i> (Haworth),	
<i>Manduca</i>	104

R

<i>Rachiplusia</i> Hampson	142
<i>radcliffei</i> (Harvey), <i>Acronicta</i>	146
<i>radix</i> (Walker), <i>Lacanobia</i>	159
<i>ralla</i> (Grote and Robinson),	
<i>Anathix</i>	154
<i>ramosula</i> (Guenée), <i>Nedra</i>	152
<i>ranalis</i> (Guenée),	
<i>Blepharomastix</i>	53
<i>rantalis</i> (Guenée), <i>Achyra</i>	51
<i>rapae</i> (Linné), <i>Pieris</i>	25
<i>Raphia</i> Hübner	145
<i>Raphiptera</i> Hampson	55
<i>ravocostaliata</i> Packard,	
<i>Eupithecia</i>	78
<i>reconditalis</i> (Walker),	
<i>Diathrausta</i>	53
<i>rectangula</i> (W. Kirby),	
<i>Syngrapha</i>	142
<i>[rectangulata]</i> Ottolengui,	
<i>Habrosyne</i>	66
<i>[rectilinea]</i> (French), <i>Grammia</i>	128
<i>recurvalis</i> (Fabricius), <i>Spoladea</i>	53
<i>redimicula</i> (Morrison), <i>Euxoa</i>	165
<i>reductella</i> (Walker), <i>Tlascala</i>	61
<i>refractaria</i> (Guenée),	
<i>Metarrhantis</i>	91
<i>refusaria</i> (Walker), <i>Heterophleps</i>	79
<i>relicta</i> Walker, <i>Catocala</i>	139
<i>reluctalis</i> (Hulst), <i>Abegesta</i>	49
<i>Renia</i> Guenée	135

<i>reniculalis</i> (Zeller),	
<i>Nephrogramma</i>	49
<i>reniculelloides</i> Mutuura and Munroe,	
<i>Dioryctria</i>	61
<i>reniformis</i> (Grote), <i>Celaena</i>	150
<i>renigera</i> (Stephens), <i>Lacinipolia</i>	160
<i>renunciata</i> (Walker),	
<i>Hydriomena</i>	73
<i>repanda</i> Walker, <i>Panopoda</i>	137
<i>residua</i> Grote, <i>Catocala</i>	139
<i>retardata</i> (Walker), <i>Acronicta</i>	146
<i>retecta</i> Grote, <i>Catocala</i>	139
<i>[reversa]</i> (F. & R. Chermock),	
<i>Charidryas nycteis</i>	40
<i>reversa</i> (Stretch), <i>Haploa</i>	123
<i>reversalis</i> (Guenée), <i>Uresiphita</i>	52
<i>revicta</i> (Morrison), <i>Orthosia</i>	161
<i>rhodarialis</i> (Walker),	
<i>Phytometra</i>	137
<i>Rheumaptera</i> Hübner	74
<i>Rhodoecia</i> Hampson	167
<i>Rhynchagrotis</i> Smith	166
<i>ribearia</i> (Fitch), <i>Itame</i>	81
<i>ridingsella</i> Clemens, <i>Adela</i>	10
<i>ridingsiana</i> (Grote), <i>Euxoa</i>	165
<i>rileyana</i> Harvey, <i>Heteropacha</i>	97
<i>rimosa</i> Packard, <i>Pheosia</i>	115
Riodinidae	34
<i>riparia</i> Morrison, <i>Oncocnemis</i>	157
<i>Rivula</i> Guenée	138
Rivulinae	136
<i>rivulosa</i> (Guenée), <i>Schinia</i>	167
<i>robiniae</i> (Peck), <i>Prionoxystus</i>	15
<i>robustella</i> Zeller, <i>Tetralopha</i>	59
<i>rogatalis</i> (Hulst), <i>Hellula</i>	49
<i>rogenhoferi</i> (Möschler), <i>Polia</i>	161
<i>rolandi</i> Grote, <i>Psaphida</i>	157
<i>rolandiana</i> Grote, <i>Exyra</i>	144
<i>rosea</i> (Harvey), <i>Sideridis</i>	161
<i>rosea</i> (Walker), <i>Oreta</i>	68
<i>rotundalis</i> (Walker), <i>Idia</i>	134
<i>rotundopennata</i> (Packard), <i>Idaea</i>	70
<i>rozmani</i> Berio, <i>Dipterygia</i>	151
<i>rubera</i> (Freyer), <i>Hydriomena</i>	73
<i>rubricoma</i> Guenée, <i>Acronicta</i>	146
<i>rubicunda</i> (Fabricius),	
<i>Dryocampa</i>	99-100
<i>rubida</i> Ottolengui, <i>Autographa</i>	141
<i>rubifera</i> (Grote), <i>Diarsia</i>	163
<i>rubigalis</i> (Guenée), <i>Udea</i>	52
<i>[rubricosa]</i> (Harris), <i>Phragmatobia</i>	
<i>fuliginosa</i>	127
<i>rubrifasciella</i> Packard, <i>Acrobasis</i>	60
<i>rubrifrontaria</i> (Packard), <i>Nemoria</i>	69

[<i>rubrofasciata</i> Barnes and McDunnough, <i>Limenitis arthemis</i>]	41
[<i>rubromarginaria</i> (Packard), <i>Nemoria bistriaria</i>]	69
<i>ruficollata</i> Guenée, <i>Mesoleuca</i>	74
<i>rufimargo</i> (Hübner), <i>Panopoda</i>	137
<i>rufipectus</i> (Morrison), <i>Protolampra</i>	166
<i>rugosa</i> (Morrison), <i>Trichodestra</i>	162
<i>ruperti</i> Franclemont, <i>Chytonix</i>	150
<i>uricola</i> (Boisduval), <i>Euphyes</i>	22
<i>uricolella</i> (Zeller), <i>Agriphila</i>	56
<i>russeolata</i> Swett, <i>Eupithecia</i>	78

S

<i>saepiolus</i> (Boisduval), <i>Plebejus</i>	33
[<i>saga</i> (Staudinger), <i>Boloria frigga</i>]	39
<i>Salebriaria</i> Heinrich	60
[<i>Salia</i> Hübner = <i>Colobochyla</i> Hübner]	136
<i>salicarium</i> (Walker), <i>Metalepsis salicis</i> (Linné), <i>Leucoma</i>	133
<i>Samia</i> Hübner	102
[<i>samoset</i> (Scudder), <i>Amblyscirtes</i>]	22
[<i>samuelis</i> Nabokov, <i>Lycæides melissa</i>]	33
<i>Sarata</i> Ragonot	62
Sarothripinae	143
<i>sassacus</i> Harris, <i>Hesperia</i>	19
Saturniidae	99
Saturniinae	101
Saturniini	101
<i>satyrata</i> Hübner, <i>Eupithecia</i>	78
<i>satyricus</i> Grote, <i>Ufeus</i>	166
Satyridae	42
Satyrinae	43
Satyrini	44
<i>Satyrium</i> Scudder	29
<i>Satyrodes</i> Scudder	42
<i>satyrus</i> (Edwards), <i>Polygonia</i>	35
<i>saucia</i> (Hübner), <i>Peridroma</i>	166
<i>Saucrobotys</i> Munroe	50
<i>saundersiana</i> Grote, <i>Oncocnemis</i>	157
<i>saxifragae</i> (Hy. Edwards), <i>Synanthedon</i>	13
<i>scabra</i> (Fabricius), <i>Hypena</i>	135
[<i>scabriuscula</i> (Linné), <i>Dipterygia</i>]	151
<i>scandens</i> (Riley), <i>Euxoa</i>	164

[<i>scapha</i> (Harris), <i>Prolimacodes</i>]	46
<i>Schinia</i> Hübner	167
<i>Schizura</i> Doubleday	119
Schoenobiinae	54
[<i>Schoenobius</i> Duponchel = <i>Donacaula</i> Meyrick]	54
<i>scholastica</i> McDunnough, <i>Euxoa</i>	165
<i>scitipennis</i> Walker, <i>Notodonta</i>	115-6
<i>scitisscripta</i> Walker, <i>Cerura</i>	121
<i>scitula</i> (Harris), <i>Synanthedon</i>	13
<i>scobialis</i> (Grote), <i>Idia</i>	134
<i>Scolecocampa</i> Guenée	138
<i>Scoliopteryx</i> Germar	138
Scolitantidini	33
<i>scolopendrina</i> (Boisduval), <i>Furcula</i>	121
<i>Scoparia</i> Haworth	47
Scopariinae	47
<i>Scopula</i> Schrank	71
Scopulini	71
<i>scribonia</i> (Stoll), <i>Ecpantheria</i>	126
<i>scripta</i> (Gosse), <i>Habrosyne</i>	66
<i>scriptana</i> (Walker), <i>Nycteola</i>	143
[<i>scudder</i> (Packard), <i>Parasemia</i>]	124
<i>secedens</i> (Walker), <i>Anartomima</i>	159
<i>secta</i> (Grote), <i>Hyperstrotia</i>	144
<i>secundella</i> Ely, <i>Acrobasis</i>	60
<i>selecta</i> (Walker), <i>Syngrapha</i>	142
<i>selenæ</i> (Denis and Schiffermüller), <i>Boloria</i>	39
<i>Selenia</i> Hübner	91
[<i>selenis</i> W. Kirby, <i>Phyciodes</i>]	39
<i>semitrata</i> (Hulst), <i>Eustroma</i>	72
<i>semicana</i> (Walker), <i>Oligia</i>	152
<i>semiclarata</i> (Walker), <i>Lomographa</i>	86
<i>semifuneralis</i> (Walker), <i>Euzophera</i>	62
<i>Semiothisa</i> Hübner	81
Semiothisini	80
[<i>semipurpurea</i> (Walker), <i>Phytometra</i>]	137
<i>semirelictæ</i> Grote, <i>Catocala</i>	139
<i>semirufescens</i> (Walker), <i>Oligocentria</i>	120
<i>semiusta</i> Grote, <i>Lithophane</i>	156
[<i>semplei</i> Holland, <i>Oeneis melissa</i>]	44
<i>senatoria</i> (J. E. Smith), <i>Anisota</i>	100
<i>sennæ</i> (Linné), <i>Phoebis</i>	27
<i>sensilis</i> Grote, <i>Chytonix</i>	150
<i>septentrionis</i> Walker, <i>Gluphisia</i>	116

<i>sepulchralis</i> Guérin-Ménéville,		<i>solidaginis</i> (Hübner),	
<i>Thyris</i>	64	<i>Lithomoia</i>	156
<i>serena</i> Edwards, <i>Catocala</i>	139	<i>sordida</i> Grote, <i>Catocala</i>	139
<i>serinaria</i> Herrich-Schäffer,		<i>sordidula</i> (Grote), <i>Hypena</i>	135
<i>Plagodis</i>	92	<i>sororiata</i> Hübner, <i>Carsia</i>	78
[<i>serralinealis</i> (Barnes and Benjamin),		<i>sospeta</i> (Drury), <i>Xanthotype</i>	88
<i>Munroessa gyralis</i>]	48	<i>spadix</i> (Cramer), <i>Cissusa</i>	136
<i>serrata</i> (Drury), <i>Euchlaena</i>	88	<i>Spaelotis</i> Boisduval	166
<i>serrataria</i> Barnes and McDunnough,		[<i>Spalgini</i> = <i>Gerydini</i>]	28
<i>Eulithis</i>	72	<i>Spargania</i> Guenée	74
<i>serratissimalis</i> Zeller,		<i>Spargaloma</i> Grote	138
<i>Crocidophora</i>	51	<i>Spartiniphaga</i> McDunnough	154
<i>servita</i> (Smith), <i>Euxoa</i>	165	<i>speciosa</i> (Hübner), <i>Xestia</i>	167
<i>Sesia</i> Fabricius	13	<i>speciosissima</i> (Grote and Robinson),	
<i>Sesiidae</i>	12	<i>Papaipema</i>	153
<i>Sesiinae</i>	12	<i>sperata</i> Grote, <i>Acronicta</i>	146
<i>Sesiini</i>	13	<i>sperryi</i> Herbulot, <i>Epirrhoe</i>	75
<i>Sesioidea</i>	12	<i>speyeri</i> Lintner, <i>Cucullia</i>	155
<i>sesquistrialis</i> Hübner, <i>Glaphyria</i>	49	<i>Speyeria</i> Scudder	37
<i>sexmaculata</i> (Packard),		<i>Sphecodina</i> Blanchard	110
<i>Semiothisa</i>	82	<i>Sphenarches</i> Meyrick	64
<i>sexpunctata</i> Grote, <i>Spargaloma</i>	138	<i>Sphingicampa</i> Walsh	99
<i>sexta</i> (Linné), <i>Manduca</i>	104	<i>Sphingidae</i>	104
<i>sheppardata</i> McDunnough,		<i>Sphinginae</i>	104
<i>Eupithecia</i>	78	<i>Sphingini</i>	104
<i>sheppardi</i> (McDunnough), <i>Udea</i>	52	<i>Sphingoidea</i>	104
<i>shurtleffi</i> Packard, <i>Heterogenea</i>	46	<i>Sphinx</i> Linné	105
<i>sicalis</i> (Walker), <i>Lipocosma</i>	50	<i>Spiloloma</i> Grote	138
<i>Sicya</i> (Guenée)	94	<i>Spilomelini</i>	52
<i>Sideridis</i> Hübner	161	<i>Spilosoma</i> Curtis	125
<i>sidus</i> (Guenée), <i>Eupsilia</i>	155	<i>spinigera</i> Guenée, <i>Acronicta</i>	146
<i>sigmoides</i> (Guenée),		<i>spinuloides</i> (Herrich-Schäffer),	
<i>Eueretagrotis</i>	164	<i>Adoneta</i>	47
<i>signaria</i> (Hübner), <i>Semiothisa</i>	82	<i>Spiramater</i> McCabe	162
<i>signata</i> (Walker), <i>Tricholita</i>	162	<i>Spodoptera</i> Guenée	154
<i>signatalis</i> (Walker), <i>Pyrausta</i>	52	<i>Spoladea</i> Guenée	53
<i>signataria</i> (Walker),		<i>Spragueia</i> Grote	145
<i>Melanolophia</i>	85	<i>Stamnodes</i> Guenée	75
<i>silaceata</i> (Denis and Schiffermüller),		<i>Stamnodini</i>	75
<i>Ecliptopera</i>	72	<i>Staphylus</i> Godman and Salvin	16
<i>similaria</i> (Walker), <i>Gueneria</i>	91	<i>statalis</i> Grote, <i>Arta</i>	58
<i>similis</i> Edwards, <i>Catocala</i>	139	<i>Stegae</i> Munroe	49
<i>simplaria</i> Graef, <i>Notodonta</i>	116	<i>Stenoporpia</i> McDunnough	83
<i>Simyra</i> Ochsenheimer	147	<i>Stenoptilia</i> Hübner	65
<i>sinelinea</i> Hardwick, <i>Euxoa</i>	164	<i>Stenoptilodes</i> Zimmerman	65
[<i>Sisyrosea</i> Grote = <i>Isa</i> Packard]	47	[<i>Stierha</i> Hübner = <i>Idaea</i>	
<i>Sitochroa</i> Hübner	52	Treitschke]	70
[<i>slossoniae</i> Packard, <i>Gluphisia</i>]	116	<i>Sterrhinae</i>	70
<i>Smerinthini</i>	107	<i>Sterrhini</i>	70
<i>Smerinthus</i> Latreille	107	<i>Sthenopis</i> Packard	9
<i>smithii</i> (Snellen), <i>Xestia</i>	166	<i>sticticalis</i> (Linné), <i>Loxostege</i>	52
<i>sobrialis</i> (Walker), <i>Renia</i>	135	<i>stigma</i> (Fabricius), <i>Anisota</i>	100
<i>socialis</i> (Grote), <i>Pyrausta</i>	52	<i>stigmella</i> Dyar, <i>Acrobasis</i>	60
<i>Solenobia</i> Dupont	11	<i>stigmosa</i> Morrison, <i>Agrotis</i>	162

[stragula Grote, Notodonta] 115-6
stramentosa Guenée, *Hydraecia* 151
stramineus (Walshingham),
Oidaematophorus 65
strattonata Packard, *Eupithecia* 78
Stretchia Hy. Edwards 162
strigataria (Minot), *Phigalia* 86
strigosa (Grote), *Clostera* 113
[strigulosa McDunnough, Oeneis
chryxus] 44
Strymon Hübner 31
Strymonini 29
[stygiata McDunnough, Rheumaptera
subhastata] 74
stypticellum Grote,
Homoeosoma 62
styracis (Guenée), *Psaphida* 157
subaequaria Walker, *Petrophora* 90
subcaesiella (Clemens),
Nephtopterix 61
subcessaria (Walker), *Itame* 81
subflava (Grote), *Archanara* 149
subfuscata (Haworth),
Eupithecia 78
subfuscella (Ragonot),
Nephtopterix 61
subgothica (Haworth), *Feltia* 165
subhastata (Nolcken),
Rheumaptera 74
subjuncta (Grote and Robinson),
Lacanobia 159
sublunaria (Guenée), *Cleora* 84
submarmorata (Walker),
Semiothisa 82
submedialis (Grote), *Mecyna* 53
submediana Strand, *Zale* 140
subnata Grote, *Catocala* 139
subrosea (Stephens), *Eugraphe* 164
subrotata Harvey, *Heterocampa* 118
subsequalis (Guenée), *Pyrausta* 52
subsignaria (Hübner), *Ennomos* 90
substriataria (Hulst), *Epirrhantis* 90
subtetricella (Ragonot),
Myelopsis 60
sulphuraria Packard,
Hesperumia 83
sulphurea (Packard), *Itame* 81
Sunira Franclemont 158
superans Guenée, *Acronicta* 146
superatalis Clemens, *Epipaschia* 58
surena (Grote), *Syngrapha* 142
sutor Guenée, *Condica* 150
Sutyna Todd 158
swetti Grossbeck, *Eupithecia* 78

sylviella Ely, *Acrobasis* 60
Symmerista Hübner 117
Sympistis Hübner 158
Synanthedon Hübner 13
Synanthedonini 13
Synchlora Guenée 69
Synchlorini 69
[Synedoida Hy. Edwards = Drasteria
Hübner] 136
Syngrapha Hübner 142
synchitis (Grote and Robinson),
Maliatha 145
syringae (Harris), *Podosesia* 13
[Syrrhodia Hübner = Erastria
Hübner] 87

T

tacoma (Strecker), *Trichodestra* 162
Tacparia Walker 90
Taleporiinae 11
talidiformis Guenée 151
Tarachidia Hampson 145
tarda (Guenée), *Anorthodes* 148
tarquinius (Fabricius), *Feniseca* 28
[taylorata Swett, Eupithecia
intricata] 78
taylorella (Kearfott), *Lampronia* 10
Tebenna Billberg 14
Telethusia Heinrich 61
tenebrifera (Walker), *Cerastis* 163
tenebrosella (Hulst), *Salebriaria* 60
tenera Hübner, *Cycnia* 130
tenuicula (Morrison), *Xestia* 167
tenuidactyla Fitch, *Geina* 64
tepida Grote, *Lithophane* 157
teratophora (Herrich-Schäffer),
Anterastris 147
terminimaculata (Grote),
Acontia 144
terrenella Zeller, *Aphomia* 59
tersa (Linné), *Xylophanes* 111
tesselaris (J. E. Smith),
Halysidota 129-30
tesselata (Harris), *Euxoa* 165
tesseradactyla (Linné), *Platyptilia* 65
testacea Packard, *Tortricidea* 46
testata (Linné), *Eulithis* 72
Tetanolita Grote 135
teterella (Zincken), *Parapediasia* 56
Tetracis Guenée 94
tetradella (Zeller), *Peoria* 63
Tetralopha Zeller 58
textula (Herrich-Schäffer), *Isa* 47

<i>tharos</i> (Drury), <i>Phyciodes</i>	40	<i>tonsa</i> (Grote), <i>Oligia</i>	152
<i>Thaumatopsis</i> Morrison	56	<i>toparia</i> (Zeller), <i>Chrysoteuchia</i>	55
<i>thaxteri</i> Grote, <i>Lithophane</i>	157	<i>torrefacta</i> (J. E. Smith),	
<i>thaxteriana</i> (Grote), <i>Psaphida</i>	157	<i>Apatelodes</i>	96
<i>themistocles</i> (Latreille), <i>Polites</i>	20	<i>Tortricidia</i> Packard	45
<i>Thera</i> Stephens	73	<i>tortricina</i> (Zeller), <i>Tarachidia</i>	145
<i>theralis</i> (Walker), <i>Polypogon</i>	135	<i>Tortricoidea</i>	15
<i>theseusalis</i> (Walker),		<i>Tosale</i> Walker	58
<i>Herpetogramma</i>	54	<i>Trachea</i> Ochsenheimer	154
<i>thestialis</i> (Walker),		<i>tragopoginis</i> (Clerk),	
<i>Herpetogramma</i>	54	<i>Amphipyra</i>	148
<i>thisoaria</i> (Guenée), <i>Antepione</i>	95	<i>transfigurata</i> Swett, <i>Hydriomena</i>	73
[<i>thoe</i> (Guérin-Ménéville),		<i>transitaria</i> (Walker), <i>Semiothisa</i>	82
<i>Lycaena</i>]	28	<i>transversata</i> (Drury),	
<i>Thopeutis</i> Hübner	57	<i>Prochoerodes</i>	95
<i>thoracica</i> (Putnam-Cramer),		[<i>treatii</i> (Grote), <i>Holomelina</i>	
<i>Xylena</i>	158	<i>lacta</i>]	124
<i>thoreaui</i> (Grote and Robinson),		<i>triangulifer</i> Grote, <i>Cirrophanus</i>	150
<i>Schinia</i>	167	<i>Trichoclea</i> Grote	162
<i>Thorybes</i> Scudder	15	<i>Trichodestra</i> McCabe	162
<i>thule</i> Strecker, <i>Sthenopsis</i>	9	<i>Trichodezia</i> Warren	76
Thyatridae	66	<i>Tricholita</i> Grote	162
Thyatrinae	66	[<i>Trichoplexia</i> Hampson = <i>Apamea</i>	
<i>thyatiroides</i> (Walker),		Ochsenheimer]	148
<i>Dasylophia</i>	117	<i>Trichoplusia</i> McDunnough	142
<i>thyatyroides</i> (Guenée),		<i>Trichoptilus</i> Walshingham	64
<i>Eosporopteryx</i>	142	[<i>tricularis</i> (Hübner), <i>Bellona</i>	
<i>Thymelicus</i> Hübner	18	<i>eunomia</i>]	38
<i>thymetusalis</i> (Walker), <i>Herculia</i>	58	<i>tricolorella</i> Grote, <i>Acrobasis</i>	60
Thyrididae	64	<i>tricosa</i> (Lintner), <i>Feltia</i>	165
Thyridinae	64	<i>trifascia</i> Hübner, <i>Schinia</i>	167
Thyridioidea	64	<i>trifolii</i> (Hufnagel), <i>Discestra</i>	159
<i>Thyris</i> Laspeyres	64	<i>triguttaria</i> Herrich-Schäffer	79
<i>Thysania</i> Dalman	138	[<i>tripartitus</i> (Grote),	
<i>thysbe</i> (Fabricius), <i>Hemaris</i>	109	<i>Smerinthus</i>]	107
<i>tibialis</i> (Harris), <i>Sesia</i>	13	<i>Triphosa</i> Stephens	73
<i>tigrinaria</i> (Guenée), <i>Euchlaena</i>	88	<i>tripunctata</i> Herrich-Schäffer,	
<i>tiliaria</i> (Harris), <i>Erannis</i>	86	<i>Eupithecia</i>	78
Timandrini	70	<i>triquetrana</i> (Fitch), <i>Nola</i>	144
<i>tinctaria</i> (Walker), <i>Orthofidonia</i>	83	<i>trisecta</i> (Walker), <i>Pediasia</i>	56
Tineoidea	11	<i>tresignata</i> (Walker),	
Tinithiinae	12	<i>Harrisimemna</i>	147
<i>tipuliformis</i> (Clerck),		[<i>tristata</i> auct., <i>Epirrhoe sperryi</i>]	75
<i>Synanthedon</i>	13	<i>tristigmata</i> (Grote), <i>Eupsilia</i>	155
Tirathabini	59	<i>tristis</i> Smith, <i>Acronicta</i>	146
<i>titian</i> (Cramer), <i>Aellopos</i>	109	<i>tristrigella</i> (Walker), <i>Balsa</i>	149
<i>titania</i> (Esper), <i>Boloria</i>	39	<i>tritona</i> (Hübner), <i>Acronicta</i>	146
<i>titea</i> (Cramer), <i>Phigalia</i>	86	Troidini	23
<i>titus</i> (Fabricius), <i>Harkenclenus</i>	29	<i>troilus</i> Linné, <i>Papilio</i>	24
<i>Tlascala</i> Hulst	61	<i>truncata</i> (Hufnagel), <i>Dysstroma</i>	72
[<i>toddi</i> (Holland), <i>Boloria</i>		<i>truncataria</i> (Walker), <i>Epelis</i>	81
<i>bellona</i>]	38	<i>truncatella</i> (Zetterstedt), <i>Pediasia</i>	56
<i>togata</i> Esper, <i>Xanthia</i>	158	<i>tubercularis</i> Lederer,	
<i>Tolyte</i> Hübner	97	<i>Crocidophora</i>	51

Tulsa Heinrich 61
turbatella (Walker), *Arequipa* 55
 U
u-aureum (Guenée), *Syngrapha* 142
Udea Guenée 52
Ufeus Grote 166
ulmiarrosorella (Clemens),
Canarsia 62
Ulolonche Smith 162
ulsterata (Pearsall), *Semiothisa* 82
ultronia (Hübner), *Catocala* 139
umbra (Hufnagel), *Pyrrhia* 167
umbrata (Walker),
Heterocampa 118
umbripennis (Hulst), *Tulsa* 61
unangulata (Haworth), *Euphyia* 76
undularis (Drury), *Zale* 140
undulata (Linné), *Hydria* 73
undulosa (Walker), *Ceratomia* 105
unicolor (Packard), *Misogada* 117
unicolorella Hulst, *Oreana* 60
unicornis (J. E. Smith),
Schizura 119
unifascialis (Packard), *Pyrausta* 52
unijuga Walker, *Catocala* 139
unilineata (Grote), *Zale* 140
unimacula (Grote and Robinson),
Evergestis 50
unimoda (Lintner), *Lithophane* 157
unimoda (Smith), *Papaipema* 153
unio (Hübner), *Eudryas* 147
unipuncta (Haworth),
Pseudaletia 161
unipunctata Haworth,
Hypagyrtis 84
unistriatellus Packard, *Crambus* 55
unitaria (Herrich-Schäffer),
Lytrosis 87
urentis Guenée, *Abrostola* 141
Uresiphita Hübner 52
Urola Walker 56
ursaria (Walker), *Lycia* 86
ursula Forbes, *Leucania* 160
urticaria Swett, *Xanthotype* 88
Utetheisa Hübner 123

V
vaccinii Riley, *Acrobasis* 60
vacciniivora Munroe, *Tetralopha* 59
vagans (Barnes and McDunnough),
Dasychira 132

Vanessa Fabricius 36
variolaria Guenée, *Cabera* 87
vasiliata (Guenée), *Anticlea* 75
vau-album (Denis and Schiffermüller),
Nymphalis 36
Vaxi Bleszynski 57
vecors (Guenée), *Condica* 150
vecors (Guenée), *Pseudorthodes* 161
velata (Walker), *Amphipoea* 148
velleda (Stoll), *Tolyte* 97
velleripennis (Grote), *Euxoa* 165
vellivolata (Hulst),
Anacamptodes 83
venerabilis Walker, *Agrotis* 163
ventralis (Grote and Robinson),
Diastictis 54
Venusia Curtis 76
venusta (Walker), *Plusia* 142
verbascoides (Guenée), *Apamea* 149
verna (Edwards), *Pompeius* 20
vernata (Peck), *Paleacrita* 86
verruca (Fabricius),
Argyrogramma 141
versicolor (Grote), *Elaphria* 151
versicolor (Harris), *Plarapsa* 111
vestaliata (Guenée), *Lomographa* 87
vestale (Packard), *Spilosoma* 122,
 125
 [vestris Boisduval, *Euphyes*
uricola] 22
vetusta Walker, *Agrotis* 163
vetustella (Dyar), *Nephoterix* 61
vialis (Edwards), *Amblyscirtes* 22
viator (Edwards), *Poanes* 21
vibex (Geyer), *Polites* 20
viburni Engelhardt,
Synanthedon 13
vicina (Grote), *Lacinipolia* 160
videns (Guenée), *Condica* 150
vidua (J. E. Smith), *Catocala* 139
villificans Barnes and McDunnough,
Hyperstrotia 144
vinnula (Grote), *Acronicta* 146
vinulenta (Grote), *Eupsilia* 155
 [violacea (Edwards), *Celastrina*] 33
violacearia (Walker), *Idaea* 70
virescens (Fabricius), *Heliothis* 167
virgatella (Clemens),
Nephoterix 61
virginalis (Hulst), *Protitame* 80
virginica (Esper), *Ctenucha* 131
virginicum (Fabricius),
Spilosoma 125-6
virginiensis (Drury), *Anisota* 100

<i>virginiensis</i> (Drury), <i>Vanessa</i>	36
<i>virginiensis</i> (Edwards), <i>Pieris</i>	25
<i>virgo</i> (Linné), <i>Grammia</i>	129
<i>virguncula</i> (Kirby), <i>Grammia</i>	129
<i>viridata</i> (Packard), <i>Acasis</i>	79
<i>viridisigma</i> Grote, <i>Syngrapha</i>	142
<i>viriditincta</i> Smith, <i>Oncocnemis</i>	157
<i>Vitacea</i> Engelhardt	12
[<i>vittata</i> (Fabricius), <i>Apantesis</i>]	127
<i>Vitula</i> Ragonot	62
[<i>vividior</i> Munroe, <i>Anageshna</i>]	53
<i>vobisne</i> Dyar, <i>Platytes</i>	55
<i>volubilis</i> Harvey, <i>Agrotis</i>	163
<i>vulgaris</i> (Grote and Robinson), <i>Apamea</i>	149
<i>vulgivagella</i> (Clemens), <i>Agriphila</i>	56
<i>vultuosa</i> (Grote), <i>Apamea</i>	149

W

<i>Wallengrenia</i> Berg	20
<i>walshella</i> Clemens, <i>Solenobia</i>	11
[<i>wardi</i> Riotte, <i>Orgyia</i>]	132
<i>warnerae</i> (Harvey), <i>Metarrhantis</i>	91
<i>whitmerellus</i> Klots, <i>Crambus</i>	55
<i>williamsi</i> (Dodge), <i>Grammia</i>	129
[<i>winni</i> (Gunder), <i>Speyeria</i> <i>aphrodite</i>]	37

X

<i>Xanthia</i> Ochsenheimer	158
<i>xanthioides</i> Guenée, <i>Perigea</i>	153
<i>xanthoides</i> (Boisduval), <i>Lycaena</i>	29
<i>xanthometata</i> (Walker), <i>Mellilla</i>	81
<i>Xanthophysa</i> Munroe	50
<i>Xanthorhoe</i> Hübner	75
<i>Xanthorhoini</i>	75
<i>Xanthotype</i> Warren	88
<i>Xestia</i> Hübner	166

<i>Xubida</i> Schaus	57
<i>Xylena</i> Hübner	158
<i>xylinoides</i> (Guenée), <i>Hyppa</i>	151
<i>Xylomoia</i> Staudinger	154
<i>Xylophanes</i> Hübner	111
<i>Xylotype</i> Hampson	158
[<i>Xystrota</i> Hulst = <i>Leptostales</i> Möschler]	71

Y

<i>y-inversum</i> Packard, <i>Apoda</i>	46
[<i>youngata</i> Taylor, <i>Eupithecia</i> <i>perfusca</i>]	78
<i>youngellus</i> Kearfott, <i>Crambus</i>	55
<i>youngii</i> (Smith), <i>Xestia</i>	167
<i>Yponomeuta</i> Latreille	12
Yponomeutidae	11
Yponomeutoidea	11

Z

<i>Zale</i> Hübner	140
[<i>Zanclognatha</i> Lederer = <i>Polypogon</i> Schränk]	134
<i>zarucco</i> (Lucas), <i>Erynnis</i>	17
<i>zea</i> (Boddie), <i>Helicoverpa</i>	167
<i>zeae</i> (Harris), <i>Achatodes</i>	148
<i>zeellus</i> (Fernald), <i>Crambus</i>	55
<i>zenobia</i> (Cramer), <i>Thysania</i>	138
<i>Zerene</i> Hübner	27
<i>zetterstedtii</i> (Staudinger), <i>Sympistis</i>	158
Zeugloptera	9
<i>zimmermani</i> (Grote), <i>Dioryctria</i>	61
<i>zinckenella</i> (Treitschke), <i>Etiella</i>	60
<i>Zophodia</i> Hübner	62
Zygaenidae	45
Zygaenoidea	45

Annotated List of Ontario Lepidoptera

J. C. E. Riotte

This checklist summarizes current knowledge about the distribution and taxonomy of the major groups of moths and butterflies in Ontario. The author, J. C. E. Riotte, is a research associate in the Department of Entomology, Bishop Museum, Honolulu, and also in the Department of Entomology, Royal Ontario Museum.

Books of related interest from the Royal Ontario Museum:

Revision of the Caddisfly Genus *Psilotreta* (Trichoptera: Odontoceridae)

Life Sciences Contributions 144

C. R. Parker and G. B. Wiggins

ISBN 0-88854-332-8, 1987, ill., 55 pp.

Revision of the World Species of *Spalangiopecta* (Hymenoptera: Chalcidoidea: Pteromalidae: Ceinae)

Life Sciences Contributions 155

D. Christopher Darling

ISBN 0-88854-395-6, 1991, ill., 48 pp.

Annotated Checklist of the Birds of Ontario Second Edition, Revised and Expanded

Ross D. James

Life Sciences Miscellaneous Publications

ISBN 0-88854-394-8, 1991, 128 pp.

Shrubs of Ontario

James H. Soper and Margaret L. Heimbürger

Life Sciences Miscellaneous Publications

ISBN 0-88854-283-6, 1982, ill., 528 pp.

ROM

**Royal Ontario Museum
Life Sciences
Miscellaneous Publications**

ISBN 0-88854-397-2

9 780888 543974