**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1998

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☒ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☒ Additional comments / Commentaires supplémentaires:

Pagination is as follows: p. [80]-86.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x		14x		18x		22x		26x		30x	
								☒			
	12x		16x		20x		24x		28x		32x

The copy filmed here has been reproduced thanks to the generosity of:

National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\longrightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\longrightarrow$ signifie "À SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrant la méthode.

MICROCOPY RESOLUTION TEST CHART

ANSI and ISO TEST CHART No. 2

APPLIED IMAGE Inc.

360 First Avenue
New York, New York 10014
Tel: (212) 512-1000
Telex: 208 512 1000

(Reprinted from THE COASTER Vol. XXI pp. 8-18, May, 1949)

THE SUMMER BIRDS OF HAZELTON, BRITISH COLUMBIA*

By P. A. TAVERNER

WITH ONE PHOTO

THE SUMMER BIRDS OF HAZELTON, BRITISH COLUMBIA*

By P. A. TAVERNER

WITH ONE PHOTO

HAZELTON, BRITISH COLUMBIA, is at the head of the Skeena River, at the forks where the stream is formed by the junction of the Bulkley and Babine rivers. It is the most northern point reached by the Grand Trunk Pacific Railroad, hence, with the exception of points on the new Hudson Bay Railroad, it is the most northern station reached by any of the main railway systems in America. In latitude $55^{\circ} 20'$, it is on line with the mouth of James Bay and slightly north of Hamilton Inlet on the Labrador coast. Situated as it is, at the head of navigation on the Skeena River, the supply route

*Published by permission of the Geological Survey, Ottawa, Canada

of all the northern parts of British Columbia, and on the line of the old Yukon telegraph line, Hazelton formerly enjoyed somewhat greater commercial importance than it does now, as but one of many stations on a transcontinental railroad system. It still remains, however, an important distribution center for the country to the north. Though geographically far north and in a latitude that in the east would exhibit arctic characters, the region about Hazelton is purely Canadian zone in its zoological and botanical aspects, though close to the Hudsonian boundary. It is situated at the inner edge of the coast range,

FIG. 19. THE BULKLEY RIVER, FROM THE SUSPENSION BRIDGE AT NEW HAZELTON, BRITISH COLUMBIA.

the lower altitude at about 800 feet. The surroundings are mountainous, Roche de Brulé, the highest mountain in the vicinity, rising 9000 feet above the sea level, while lesser elevations above the valley are numerous. The valleys are clothed with a dense growth of poplar, cottonwood and hemlock, with numerous Douglas firs and occasional particularly fine examples of birch. The mountain sides are mostly covered with Douglas fir, balsam and spruce. The upper limit of timber comes at about 5000 feet elevation. The whole country has suffered severely from fire, and, as a result, much of it is almost impassable with tangled second growth and fallen timber, making progress except along

trails or roadways arduous and slow. Fortunately the roads are exceptionally fine, or at least were during our season there; pack trails lead everywhere, giving access to all parts adjoining. There is little farming activity and most of the few clearings are devoted to the grazing of a few cattle and bunches of pack horses. The topography is rough and broken. At New Hazelton, four miles above the original settlement at the forks, the Bulkley River breaks through a ridge by way of a narrow canyon over three hundred feet deep. Across this gorge the wagon road is carried on a suspension bridge, one of the highest structures of its kind in the world. A few hundred yards up stream and at a little lower elevation are the remains of a unique bridge that fell this summer, built years ago by the original natives from wire from the abandoned transcontinental telegraph line.

This locality was collected in by two parties from the Canadian Geological Survey during the summer of 1917. The first, composed of J. M. Macoun and Wm. Spreadborough, was in the field from June 20 to July 21. They camp near (Old) Hazelton, on the shores of the Skeena River, and worked most of the mountains and valleys adjoining. The present writer joined a geological party for a week (August 23 to 29) at a point some two miles west of New Hazelton near the bridge across the Bulkley. Most of his work was done in the vicinity of the camp, and higher altitudes were scarcely visited. Birds were very scarce and retiring during this midsummer week and the work is worth reporting upon only as supplementary to that performed by Macoun and Spreadborough earlier in the season.

1. *Gavia immer*. Common Loon. A Loon was seen on the Skeena, August 22, by the writer, as the train he was on approached Hazelton. The species was noted almost daily by Spreadborough in June and July.
2. *Mergus americanus*. American Merganser. Not common at Hazelton. Spreadborough reports an adult and four young.
3. *Charitonetta albeola*. Bufflehead. An adult with four young seen by Spreadborough on a small lake near Hazelton, July 18.
4. *Actitis macularia*. Spotted Sandpiper. Reported by Spreadborough as a common breeder along the river; not seen about New Hazelton.
5. *Bonasa umbellus*. Ruffed Grouse. Several coveys and single birds seen at various times by both Spreadborough and myself. No specimens were taken, but *umbellus* is the form to be expected here.
6. *Accipiter velox*. Sharp-shinned Hawk. Three seen by Spreadborough, and single birds observed by the writer nearly every day.
7. *Buteo swainsoni*. Swainson Hawk. One doubtfully recorded by Spreadborough, July 18.
8. *Haliaeetus leucocephalus*. Bald Eagle. One taken June 29; several noted in July.
9. *Falco sparverius*. Sparrow Hawk. Spreadborough found the Sparrow Hawk fairly common about Hazelton, and the writer saw several birds near New Hazelton. Two females taken July 2 and 9. I refer these to the type form, *sparverius*.
11. *Otus asio*. Screech Owl. I thought I heard the distant quavers and trills of this species a couple of evenings, but too faintly to be absolutely certain of the identification.
12. *Bubo virginianus*. Great Horned Owl. One seen by Spreadborough July 20 or 21.
13. *Ceryle alcyon*. Belted Kingfisher. Several seen along the river at Hazelton and one seen by the writer on the Bulkley.
14. *Cryobates villosus*. Hairy Woodpecker. Reported as common by Spreadborough, and one taken by the writer August 26. Three specimens taken by Spreadborough. Two of these birds are of slightly questionable identity. They are quite smoky below, but whether this is due to contact with burnt timber or is a natural coloration is

a matter of some doubt. If the latter they must be regarded as showing a tendency toward *harrisi*. Otherwise all are plainly referable to *monticola*.

15. *Dryobates pubescens*. Downy Woodpecker. Spreadborough reports seeing what he took to be Gairdner Woodpecker (*Dryobates pubescens gairdneri*), July 18.

16. *Picoides arcticus*. Arctic Three-toed Woodpecker. Fairly common, and similarly to the next species, found at the edges of burnt timber. Four specimens taken by Spreadborough, July 11.

17. *Picoides americanus*. American Three-toed Woodpecker. Spreadborough reports it as fairly common at Hazelton along the borders of burnt timber, where it appeared to be feeding on larvae of the spruce-tree borer. Three specimens taken July 5 and 6. Being without comparable eastern material I include them under *fasciatus* upon geographical probability, though I can see no sign of longitudinal striping on the back.

18. *Sphyrapicus ruber*. Red-breasted Sapsucker. Three birds seen by the writer, and two juveniles taken. Spreadborough reports them as common at Hazelton, he taking five specimens there. The adult males of this series are considerably brighter and with the back less spotted than our only California specimen, hence I refer all to *notkensis*.

19. *Pileatus pileatus*. Pileated Woodpecker. One seen by Spreadborough at Hazelton.

20. *Colaptes auratus*. Flicker. Fairly common. Four specimens taken by Spreadborough and two by the writer. The former are hybrids but nearer *auratus* than to *cafer*. The mixture of blood shows mostly in the intrusion of grayish in the throat, but the yellow of the shafts on some is deepened to dull orange. The two August birds from New Hazelton are by plumage pure *auratus*.

21. *Chordeiles virginianus*. Nighthawk. Several seen at Hazelton by Spreadborough. Again noted by the writer the evenings of August 25 and 26, but not thereafter.

22. *Cypseloides niger*. Black Swift. Nearly every bright afternoon a flock of Black Swifts was seen over our camp. Keeping in a loose company they drifted up and down the valley, coming and going until near evening. They usually kept high up, but occasionally one stooped low enough to be collected. Two were so obtained August 29.

23. *Chaetura vauxi*. Vaux Swift. Reported by Spreadborough as fairly common and breeding at Hazelton. Not recognized by the writer.

24. *Selasphorus rufus*. Rufous Hummingbird. Several seen by Spreadborough at Hazelton.

25. *Nuttallornis borealis*. Olive-sided Flycatcher. Three or four seen by Spreadborough at Hazelton, and one by the writer at New Hazelton. Two specimens taken, July 17 and in August.

26. *Myiochanes richardsoni*. Western Wood Pewee. Several heard by the writer August 24-27. Specimens taken June 28 and 30, July 16 and August 24.

27. *Empidonax hammondi*. Hammond Flycatcher. Spreadborough reports the Hammond Flycatcher as common. We have four adult specimens taken July 27 and 30, and two juveniles August 26 and 30. Both these latter are in the yellow-bellied phase and are so very similar to *flaviventris* as to be readily mistaken for that species. However, the throats are distinctly whitish, the green above grayish, and the bills and feet quite distinctive.

28. *Empidonax wrighti*. Wright Flycatcher. Though not mentioned in Spreadborough's notes he took specimens June 28 and July 10. Another was taken August 25.

29. *Cyanocitta stelleri*. Steller Jay. But one seen by the writer, a specimen taken August 25. Spreadborough does not mention it in his notes, but took two specimens June 28. The only constant character I can see in *annectens* is the presence of the gray spot over the eye. The other particulars usually given as diagnostic are unstable and unsatisfactory for individual identification. One of these specimens, a male taken June 28, has the spot, therefore I call it *annectens*. The other two are without it and I can only regard them as *stelleri*. These birds are slightly bluer on the back than the majority of specimens, but can be matched by individuals in either race. These two forms have broad fields of overlapping in their distribution, and in such localities their separation is purely arbitrary.

30. *Perisoreus canadensis*. Canada Jay. Six, seemingly a pair with young, noted by Spreadborough, July 18, and three collected. These birds are similar to those taken in Jasper Park on the east slope of the Rockies. They have extensive white fore-crowns, and blue-black hind crowns. *Fumifrons* is a slightly defined and unstable race, otherwise

they might be referred to that subspecies. As it is, I prefer to regard them provisionally as intergrades between *capitalis* and *canadensis*.

31. *Corvus corax*. Raven. A few seen by Spreadborough near Hazelton in July.
32. *Corvus brachyrhynchos* (sp?). Crow. Not common. Only a few seen by Spreadborough at Hazelton. None taken, so identity cannot be established. I assume that *C. caurinus* is a coast form and *C. b. hesperis* is a scarcely recognizable race.

33. *Euphagus cyanocephalus*. Brewer Blackbird. One pair with young noted by Spreadborough July 15. Two specimens taken.

34. *Loxia curvirostra*. American Crossbill. Several flocks observed by Spreadborough at Hazelton.

35. *Leucosticte tephrocotis*. Rosy Finch. Five seen by Spreadborough July 16 at an elevation of 6000 feet. Two taken are both *littoralis*.

36. *Spinus pinus*. Pine Siskin. Only one seen at a high altitude (5000 feet), by Spreadborough, July 16. Observed in flocks in August at New Hazelton in the lowlands several times. Specimens taken August 25.

37. *Zonotrichia leucophrys*. White-crowned Sparrow. Not common. Spreadborough reports but two seen. One specimen, taken July 15, is typical *gambeli*.

38. *Zonotrichia coronata*. Golden-crowned Sparrow. Not common. Only one seen, by Spreadborough.

39. *Spizella passerina*. Chipping Sparrow. Reported as common at Hazelton by Spreadborough, but not noted by the writer. One specimen, taken June 30, referred to *arizonae*.

40. *Junco hyemalis*. Slate-colored Junco. One specimen falls under this species, no. 10963, adult male (skull granulated), August 26. It is straight *hyemalis* except for a slight but obvious amount of red edgings on the back against which the gray of the hind neck cuts sharply. There is no rufus on the sides at all and I regard it as *hyemalis* with a slight infusion of *oregonus* blood.

41. *Junco oregonus*. Oregon Junco. The commonest species present during the August visit. Spreadborough reports it as a common breeder. Six specimens were taken in recognizable plumage, June 26 and July 5, 13, and 18, and two juveniles August 24 and 25. Besides these, numerous striped juveniles were secured, whose exact identity can only be conclusively established from accompanying parents. The June and July birds are typical *connectens*, while the August specimens in first winter plumage show a strong tendency towards *oregonus* in the richer redness of the back. The fact that the heads and throats of these birds are grayish instead of black, somewhat similar to *hyemalis*, I attribute to age rather than racial affinity. It thus seems that we have three forms of *Junco* at Hazelton, nearly pure *hyemalis*, *connectens* and *connectens* tending toward *oregonus*.

42. *Melospiza melodia*. Song Sparrow. Reported by Spreadborough to be not common, and only half a dozen specimens noted during his stay. In August a few were seen daily, but most were juveniles or in worn plumage. Seven specimens taken. These all appear to me to be *rutina*.

43. *Melospiza lincolni*. Lincoln Sparrow. Two seen at Hazelton by Spreadborough, June 26 and after. Three taken June 26 to July 17. Though I can perceive a slightly more olivaceous cast in fall birds from the west compared with eastern specimens, the difference in my opinion is too slight for subspecific recognition. I include them therefore under the typical form *lincolni*.

44. *Piranga ludoviciana*. Western Tanager. Common breeder at Hazelton according to Spreadborough, though the species was not observed by the writer in August. Four specimens taken June 27 to July 9.

45. *Iridoprocne bicolor*. Tree Swallow. Reported fairly common by Spreadborough at Hazelton, though not seen by the writer later. Specimen taken July 15.

46. *Tachycineta thalassina*. Violet-green Swallow. According to Spreadborough, a common breeder at Hazelton in June and July, though not seen by the writer in August at New Hazelton.

47. *Stelgidopteryx serripennis*. Rough-winged Swallow. Reported by Spreadborough to have been fairly common on the river in June and July, but not observed by the writer in August. One specimen taken July 4.

48. *Vireosylva olivacea*. Red-eyed Vireo. Several seen by Spreadborough at Hazelton in June and July. Not noted at New Hazelton in August.

49. *Vireosylva gilva*. Warbling Vireo. Fairly common at both Hazelton and New Hazelton. Nine specimens taken. I cannot see that these birds are any darker than eastern ones. If anything, they are slightly lighter and brighter in color. They certainly do not show any differences in measurements. The bills are slightly smaller and lighter, but in so illusive a degree as to make this feature of little value in separating them from *gilva*.

50. *Vermivora celata*. Orange-crowned Warbler. One Orange-crown taken August 28. It is too juvenile a bird to make subspecific pronouncement upon.

51. *Dendroica aestiva*. Yellow Warbler. Several seen by Spreadborough in June and July. But one noted in August. Three specimens taken, June 24 and 30, and August 28. These specimens seem to exhibit the racial characters of *rubiginosa* better than do most of our British Columbia birds, and therefore may be referred to that race.

52. *Dendroica coronata*. Myrtle Warbler. One specimen, a juvenile female, taken August 28, may be of this species. By tail spotting it might be either *coronata* or *auduboni*, and I cannot decide whether a faint tint on the throat is stain or a suggestion of the yellow throat patch.

53. *Dendroica auduboni*. Audubon Warbler. Common in June and July, and two seen in August. Five specimens taken, June 24 to July 18, and August 28.

54. *Dendroica magnolia*. Magnolia Warbler. Not uncommon. Spreadborough reports seeing five pairs that, from their actions, were probably breeding. Two were taken in August. Five specimens in all were taken.

55. *Dendroica striata*. Black-poll Warbler. One juvenile female taken August 25.

56. *Dendroica townsendi*. Townsend Warbler. Two seen August 25.

57. *Seiurus noveboracensis*. Northern Water-Thrush. On June 28 Spreadborough took two, and thereafter noted a number, always in low wet woods. These specimens exhibit the rather faint characters of *notabilis* in their highest degree.

58. *Oporornis tolmiei*. Macgillivray Warbler. Common in low bushes on burnt ground in June and July, and seen again August 20 and 24. Four specimens taken.

59. *Geothlypis trichas*. Maryland Yellowthroat. One seen and collected August 25. This is a juvenile bird that cannot be confidently referred to a subspecies.

60. *Wilsonia pusilla*. Wilson Warbler. One seen August 25. As it is only a slight record no subspecific designation can be given. *Pileolata* is the geographic probability.

61. *Setophaga ruticilla*. Redstart. Common. Six specimens taken, June 24 to July 10, and August 26 to 29.

62. *Certhia familiaris*. Brown Creeper. Two collected by Spreadborough, July 7 and 17. As these are juveniles they are referred to *zelotes* by geographic probability.

63. *Sitta canadensis*. Red-breasted Nuthatch. Several seen by both Spreadborough and the writer. Specimen taken June 24.

64. *Penthestes atricapillus*. Black-capped Chickadee. One of the commonest birds. Five specimens taken, June 15 to 25, and August 26 to 28. All are *septentrionalis*.

65. *Regulus calendula*. Ruby-crowned Kinglet. A small flock seen August 25. The brush was very dense and birds shot could not be found, hence no specimens.

66. *Hylocichla ustulata*. Olive-backed Thrush. Reported by Spreadborough as common and breeding. In August thrushes were so shy and retiring that their specific identity was difficult to determine unless the birds were collected. Two specimens taken July 6, and three August 25 and 26. Though these birds are very slightly more olive than eastern specimens, I do not care to separate them from *swainsoni*.

66. *Hylocichla guttata*. Hermit Thrush. Thrushes were often heard and glimpsed in August, but seldom seen plainly enough for identification. Three specimens were taken August 26. All are juveniles, and but one sufficiently advanced to base subspecific identity upon. I cannot separate it from *pallasi*.

67. *Planesticus migratorius*. American Robin. Abundant in June and July, and a few seen daily in August. Four specimens taken. I do not see in them sufficient grounds for separation from eastern *migratorius*.

68. *Ixoreus naevius*. Varied Thrush. One seen by Spreadborough July 20.

69. *Sialia currucoides*. Mountain Bluebird. Reported by Spreadborough as being common in burnt timber in July. It was not seen by the writer personally in Aug-

ust, though reports were received, leaving little doubt as to its presence in the vicinity. Five specimens taken. July 5 and 6.

Geological Survey, Ottawa, Canada, April 5, 1918.