**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1994

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☐ Coloured covers/
Couverture de couleur

☐ Coloured pages/
Pages de couleur

☐ Covers damaged/
Couverture endommagée

☐ Pages damaged/
Pages endommagées

☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée

☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées

☐ Cover title missing/
La titre de couverture manque

☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées

☐ Coloured maps/
Cartes géographiques en couleur

☐ Pages detached/
Pages détachées

☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)

☒ Showthrough/
Transparence

☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur

☒ Quality of print varies/
Qualité inégale de l'impression

☒ Bound with other material/
Relié avec d'autres documents

☐ Continuous pagination/
Pagination continue

☒ Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure

☐ Includes index(es)/
Comprend un (des) index

☐ Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

Title on header taken from:/
Le titre de l'en-tête provient:

☐ Title page of issue/
Page de titre de la livraison

☐ Caption of issue/
Titre de départ de la livraison

☐ Masthead/
Générique (périodiques) de la livraison

☐ Additional comments:/
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
							☒				

The copy filmed here has been reproduced thanks to the generosity of:

Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

1	2	3
---	---	---

1	2
4	5

L'exemplaire filmé fut reproduit grâce à la
générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le
plus grand soin, compte tenu de la condition et
de la netteté de l'exemplaire filmé, et en
conformité avec les conditions du contrat de
filmage.

Les exemplaires originaux dont la couverture en
papier est imprimée sont filmés en commençant
par le premier plat et en terminant soit par la
dernière page qui comporte une empreinte
d'impression ou d'illustration, soit par le second
plat, selon le cas. Tous les autres exemplaires
originaux sont filmés en commençant par la
première page qui comporte une empreinte
d'impression ou d'illustration et en terminant par
la dernière page qui comporte une telle
empreinte.

Un des symboles suivants apparaîtra sur la
dernière image de chaque microfiche, selon le
cas: le symbole $\rightarrow$ signifie "A SUIVRE", le
symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être
filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être
reproduit en un seul cliché, il est filmé à partir
de l'angle supérieur gauche, de gauche à droite,
et de haut en bas, en prenant le nombre
d'images nécessaire. Les diagrammes suivants
illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1553 East Main Street
Rochester, New York 14609 USA
(7 6) 482 - 0300 - Phone
(7 6) 288 - 5989 - Fax

MAR 22 1920

Government of the Province of Saskatchewan
DEPARTMENT OF AGRICULTURE

ALFALFA IN SASKATCHEWAN

BY JOHN BRACKEN

Professor of Field Husbandry, University of Saskatchewan.

A hardy drought resistant leguminous hay crop has long been desired by the farmers in the older parts of Saskatchewan. Such a crop is a necessity from the point of view of soil conservation and very desirable as a supplement to the forage crops now grown. Some varieties of alfalfa when properly managed, combine all the qualities mentioned above. This crop promises to be a very useful one in many portions of Saskatchewan, both as a soil improver and as a forage crop.

HISTORY OF ALFALFA IN SASKATCHEWAN.

The record of alfalfa in this province is very brief. Previous to 1904 it was almost unknown here. At that time arrangements were made to investigate its possibilities at the Experimental Farm at Indian Head and in the same year the Fairs and Institutes Branch of the Provincial Department of Agriculture distributed seed of Turkestan alfalfa to farmers in all parts of the province through the co-operation of local agricultural societies. Since that time the experiment stations and many private farmers have done a considerable amount of experimental work with it and as a result we know today that there are varieties of alfalfa suited to the climatic and soil conditions found here.

SUITABILITY TO SOIL AND CLIMATE.

Alfalfa can be grown on practically all normal soils of the province. Alkali and low lying soils, which are subject to flooding are unfavourable to alfalfa, but on the normal loams and clays it finds a satisfactory environment. On the heavy soils it produces a larger yield of forage than on the lighter types; but on the other hand the production of seed is generally greatest on the warmer and earlier soils.

Eastern Saskatchewan is better adapted to the production of alfalfa hay than Western Saskatchewan. It is doubtful whether at the present price of seed, alfalfa can be grown profitably for hay in the drier portions of Western Saskatchewan. In these parts, however, alfalfa seed can be grown very satisfactorily if the crop is sown in rows two and one-half or three feet apart.

VARIETIES.

During the last five years approximately sixty strains of alfalfa have been tested by the Field Husbandry Department of the University of Saskatchewan. As a result we are able to say with some confidence

that Grimm, some strains of Sand Lucerne and some of Turkestan are, in the order named, among the most suitable for this province. Baltic, Ontario variegated and several strains of Siberian alfalfa are now under test. These promise satisfactory results. There is now no doubt but that with fair treatment the true Grimm alfalfa will withstand the low temperatures of all normal winters here. This variety has given us more satisfactory results than any other during the five years we have been testing the crop at Saskatoon.

SEEDING.

Our experience with alfalfa goes to show that it should be seeded in the early part of the rainy season, or between the middle of May and the middle of June. It gives best results when sown on fallow or on ground which has been in hoed crop the previous year. Soil which has been well manured before a hoed crop furnishes probably the best conditions for starting alfalfa. Grass is the crop's worst enemy and we are coming to believe that alfalfa should not be sown on land which has not been freed from the creeping rooted species of native or introduced plants. Thin seeding has given best results on soils free from weeds. Thick stands suffer most in periods of drought and on the average yield less than thinner ones. On weedy land thicker seeding is necessary because of the fact that many plants are smothered out before they are large enough to hold their own against the weeds. Thorough preparation of the soil before seeding to alfalfa is essential to the profitable production of the crop.

Broadcasting the seed without a nurse crop is quite often followed, and, if sown in the rainy season on well prepared soil that is not inclined to blow, it generally gives good results. The most successful farmers are now, however, drilling in the seed. Some use the grass seed attachment on the ordinary grain drill, while others use this drill without the attachment and increase the bulk of the seed with cracked wheat or barley or other material which has been sifted to a uniform size a little larger than alfalfa seed. By drilling the seed the conditions necessary for germination are more easily controlled and less seed is necessary. From eight to twelve pounds of good seed is quite sufficient for an acre. A nurse crop should not be used.

ALFALFA—A SOIL IMPROVER.

Alfalfa is a "legume." It has the power when "inoculated" of drawing upon the atmosphere for a part of the nourishment that would otherwise be taken from the soil. "Inoculation" can be effected by taking soil from an old productive alfalfa field and applying it to the field to be sown, or the soil may be mixed with the seed before sowing. The soil so used should not be allowed to dry out. After applying, the field should be harrowed immediately.

Artificial cultures of alfalfa bacteria may be purchased and applied to the seed before sowing. These often give better results than the soil method of inoculation, but sometimes the bacteria in the cultures are killed by unfavourable conditions during transportation. The soil method is the surest but the pure culture method is sometimes the most

effective. Increased yields of from three hundred to one thousand pounds per acre have been secured from the use of both inoculated soil and alfalfa culture.

"CLIP BACK" TO CONTROL WEEDS.

When seeded on land which has borne a crop of cereals the previous year many annual weeds come among the alfalfa plants. These can be controlled only by clipping back the crop. Clipping is not desirable except where it is necessary to control weeds. No crop is expected the year the seed is sown, but on the heavier soils in the more humid parts of the province and particularly in moist seasons one crop may be taken. It has been demonstrated conclusively time and time again that a growth of six to ten inches should be left on the field in the fall in order to protect the plant roots by holding snow in the winter. Invariably this practice results in increased yields the following year and in longer life to the field.

HARVESTING.

In harvesting alfalfa, three things should be remembered: First—That its feeding value is greatest if cut just after blooming has commenced. Second—That the leaves, which fall off with the slightest provocation when dry, contain by far the most valuable part of the crop, being equal in feeding value to wheat bran; and third—That rains not only discolour the hay and make it less palatable and less digestible, but also remove in solution much of the nutritious portion of the fodder.

The crop should not be cut while wet nor when dew lies upon it. When 10 per cent. or thereabouts of the plants are in bloom cutting should commence. It has been observed that if cut later than this the yield is less than it would otherwise be, due to the clipping back of the new sprouts which ordinarily appear at this time. Since quick drying in the sun makes the leaves brittle and causes them to break off easily, it has been found advisable to rake shortly after cutting and cure it in the cock. If left in the windrow, the outer layers become dry and brittle, causing great loss of leaves and deterioration in quality. This is better than leaving it in the swath, but not so good as curing in the cock. The latter takes more time and labour than when cured in the swath or windrow, but always gives a much superior quality of hay. It is often advisable to turn the cocks over a few hours before stacking or drawing to the barn in order to dry out the lower layer next the ground. The practice of curing in cock, may have some disadvantages in exposed locations in windy weather, but when the alfalfa is raked before it becomes too dry it will be found to settle down well and, except in the heaviest winds give very little trouble.

SURFACE CULTIVATE THE CROP IN SPRING.

In semi-arid regions the yield of a perennial forage crop is measured by the amount of precipitation conserved in the soil. The practice of disking, "renovating" or harrowing alfalfa in early spring when for three or four weeks after the snow goes the field is left practically bare and lifeless is to be recommended. It has given us increased yields at the University and on those plots where cultivation was given the

crop is in better condition now than where no cultivation was given. The harrows can do no damage after the first year, and the other implements named if used with care, will do but little harm to the plants and much good to the soil.

PRESENT STATUS OF ALFALFA PRODUCTION.

The present status of alfalfa production in Saskatchewan, while very promising, is not quite satisfactory. It has been amply demonstrated that certain varieties are hardy and are quite well suited to the conditions found here. In view of the knowledge of this fact, and the relatively small supply of seed of these varieties, the price is becoming almost prohibitive. In the southern and western part of the province where the precipitation is lighter the yield is often small particularly in a dry year.

To make the situation more difficult, practically no seed has been produced in a commercial way in the province. It is clearly the business of all agricultural interests to promote in every legitimate way the production of alfalfa seed. The south-western corner of the province seems particularly well adapted for this purpose. The farmers need the seed, and south-western Saskatchewan can produce it.

A considerable number of farmers are now sowing alfalfa in drills two, two and a half or three feet apart for the production of seed. The yield of seed is greater from fields sown in wide rows than from those sown in the ordinary way. When sown in rows two and a half feet apart, three or four pounds is ample for an acre. Yields of seed ranging from fifty to one hundred pounds have been secured in this province from crops sown in the ordinary way, and as much as three hundred pounds per acre has been harvested in south-eastern Alberta, where the crop was sown in rows two and a half feet apart.

It may be remarked incidentally that at the University larger yields of forage have been secured from seeding alfalfa in rows eighteen inches apart at six pounds per acre than by seeding it at any closer distance with more seed. In the dry year of 1914, rows 24 inches apart sown at $4\frac{1}{2}$ pounds per acre yielded more than any closer sowing. We do not at this time recommend farmers in the north and north-eastern parts of the province to undertake the production of alfalfa seed, but those in the south-western part can grow seed profitably if they will.

A circular on alfalfa seed production will be forwarded on request to anyone interested, by the College of Agriculture at Saskatoon, or the Department of Agriculture at Regina.