**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1994

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☒ Bound with other material/
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distorsion le long de la marge intérieure
- ☐ Blank leaves added during restoration may appear
within the text. Whenever possible, these have
been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Continuous pagination/
Pagination continue
- ☐ Includes index(es)/
Comprend un (des) index

Title on header taken from: /
Le titre de l'en-tête provient:

- ☐ Title page of issue/
Page de titre de la livraison
- ☐ Caption of issue/
Titre de départ de la livraison
- ☐ Masthead/
Générique (périodiques) de la livraison

- ☐ Additional comments: /
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☐	☒	☐	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaît sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., pouvant être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1853 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

259

62

BULLETIN No. 11.

DEPARTMENT OF AGRICULTURE

—OF—

BRITISH COLUMBIA.

WILD MUSTARD.

259

62

In view of the prevalence of Charlock, or Wild Mustard, throughout the Province, the Hon. J. D. Prentice, Minister of Agriculture, directs that the following Bulletin be published for distribution amongst the members of Farmers' Institutes.

J. R. ANDERSON,

Deputy Minister of Agriculture.

*Department of Agriculture of British Columbia,
Victoria, B. C., 30th January, 1903.*

WILD MUSTARD.

— 0 —

Wild Mustard (*Brassica sinapistrum*—Boiss), also known under the several vernacular names of Charlock, Yellow Weed, Shellock, Kerlock, or Kellock, is a vile weed introduced originally from Europe, and which has now spread in Canada from the Atlantic to the Pacific. It belongs to the natural order of Crucifers (*Crucifereæ*), to which also many other weeds, such as False Flax, Shepherd's Purse, Penny Cress, Pepper Grass, and others belong. Although not included specifically in any Noxious Weeds Act in this Province, it is probably the worst weed the farmer in British Columbia has to contend with to-day, and which causes more loss, and therefore no means should be left untried to eradicate it. The following cuts, kindly lent by the University of Wisconsin, give a very accurate idea of the appearance of this weed.

It is described in Bulletin No. 76 of the Agricultural Experiment Station of Wisconsin as follows:—

"The Wild Mustard is a coarse, rough, annual plant, resembling in general appearance the garden radish, with the exception that it has a more irregular and branching root. The stem and branches, which are sparsely clothed with leaves, terminate in clusters of yellow flowers, of which the lower ones are first to open, the stem in the meanwhile continuing to lengthen, forming a long, leafless raceme, with knotted pods towards the base, open flowers towards the summit, and a cluster of unopened flower-buds at the apex. The seeds resemble those of the cabbage, and have a harsh, biting taste. A portion of a plant of the Wild Mustard is shown in Fig. 1, and another portion of a plant, drawn to a larger scale, is shown in Fig. 2."

Being an annual, the usual methods recommended for the eradication of such weeds hold good in the case of Wild Mustard. Unfortunately, however, the seed has such vitality, owing to an essential oil it contains, that it may remain in the soil, when buried deeply, for an indefinite period and germinate when conditions are favourable, so that, although it will eventually succumb to the treatment of constant cultivation which is so successfully practised for annual weeds, the process of extermination is much slower and more troublesome, and, therefore, other methods are and have been recommended. Spraying with a solution of sulphate of copper has been found to be most effective, if properly applied, and the following account of the results obtained by experiment, and the method followed, by Mr. G. F. Strawson, of 71A, Queen Victoria Street, London, E. C., under date of 2nd April, 1902, and January, 1903, is therefore published for general information:—

FIG. 1.—An individual flower and a seed pod, about natural size, appear at the left, and at the lower left hand corner is shown a flower slightly enlarged.

L R Stowell del.

FIG. 2.—Flower and seed pods, slightly enlarged.

"DESTRUCTION OF CHARLOCK IN *CORN CROPS, 1901.

"In the place of issuing a third annual report on the Charlock spraying for 1901, it may prove more serviceable if I give a short account of the work done last year, with practical deductions drawn from it.

"During the last spring I took every opportunity of observing and enquiring into the operations being then carried on, and in addition to this I gave several public demonstrations in England and Scotland. The results fully confirm the main conclusions arrived at in my second annual report, which are repeated below.

"The most important difference I noted was in the age of the Charlock when sprayed, and I am fully convinced that quite 75 % of the operations were late; 50 % were much too late, and probably 25 % were so late that not more than half the advantages to be derived from spraying were realised. The importance of spraying early cannot be too strongly urged. Arrangements should be made to spray the Charlock soon after the rough leaf appears and whilst the plant is in a soft and tender state.

"Under these conditions I do not think a single plant of Charlock can escape destruction, providing it receives its proper proportion of fine spray resulting from 15 lbs. of pure copper sulphate dissolved in 50 gallons of water per acre.

"In the south the season was cold and late last year and not favourable to the growth of Charlock, but in Scotland the plant was luxuriant.

"The suggested use of nitrate of soda in place of copper sulphate has not proved successful on extended trials. Copper sulphate has proved to be altogether best, and is at present the only material worth using. I have added a fifth paragraph to the conclusions previously arrived at, which now stand as follows:—

"1. That Charlock can be destroyed in growing corn crops, without injury to the latter, by spraying with 50 gallons of three per cent. solution of pure sulphate of copper per acre.

"2. That, generally, the corn crops are much improved and give a better yield where Charlock is destroyed.

"3. That spraying early, when the weed is young and in soft fibre, is most successful.

"4. That the large sprayers are more successful than the smaller.

"5. The success of the operation is diminished in proportion to the delay which occurs after the Charlock is in the most suitable condition for being destroyed.

"The value of this new and successful agricultural process cannot be made too widely known in Charlock-infested districts.

* Corn crops or grain crops.

"G. F. STRAWSON."

"REPORT ON THE DESTRUCTION OF CHARLOCK IN CORN CROPS IN 1902.

"In publishing a fourth annual report on this subject, it is a matter for congratulation that most of the preliminary work necessary in developing the details of this valuable process into a practical form is now accomplished. By wide and extensive operations, the strength and quantity of the solution to be used, and the most effectual time to destroy the weed, has been thoroughly established.

"The conclusions arrived at in my annual report for 1901 have been thoroughly corroborated in 1902, and are now adopted by hundreds of farmers, and thousands of acres have been profitably sprayed.

"It is also interesting to observe a keener interest shown in the work by farmers, who have taken more care to be in good time with the spraying, with the result that the weed has been more effectually destroyed, and the increased yield of corn, from the absence of the weed, is larger.

"The Charlock season of 1902, though cold and wet at times, did not appear to be unfavourable to the growth of the plant, for, later in the year, unsprayed fields of its golden flowers were visible from almost all points between Inverness and the South of England.

"The details of the process, as established in 1901, and further confirmed by the practice of 1902, show no new developments of any importance.

"Several farmers have suggested to me that a good time to spray the Charlock is when it is just coming into flower, or actually in flower, because they have succeeded admirably in destroying it at that stage by using a large quantity of copper per acre. This suggestion is not on sound principles.

"It is now thoroughly proved that the Charlock can be destroyed at any period of its growth by copper sulphate, and that whilst the plant is young and in soft fibre a smaller quantity will effect the purpose than when the plant is older. This has an important bearing on the economical side of the question.

"Fifty gallons of 3 per cent. solution favourably applied to young plants will destroy 95 per cent. of the weed in an average-infested crop; in fact, it will destroy all except those few plants that are shaded by other leaves from the spray, and this quantity and strength will affect the corn crops so slightly that an increased yield of corn, more than sufficient to pay all the expenses of spraying, may with confidence be looked for.

"On the other hand, if the weed is sprayed when much older, more copper is required, and this gives a shock to the corn plant which prevents the likelihood of any material increase in the yield, besides which it must be remembered that the older the weed the more it has robbed the soil and shaded the sun from the corn.

"The spraying process is now firmly adopted by so large a number of intelligent farmers that there is little need of further special public demon-

strations. Indeed, these public demonstrations are beset with difficulties that often render them disappointing and valueless. One of the principal disadvantages is that a date must be fixed for them in advance, and the weed and the weather seem often to combine in order to defeat the best laid plans.

"The farmer has no such difficulty to contend with. He sees the crop daily from beginning to end, and, having provided himself with the means, may start any suitable day and make the destruction of the pest certain. No one is in so good a position as the farmer to give a demonstration for the benefit of his fellow farmers, and no one has so direct an interest in improving the annual value of the land. The operation is so simple and easy that an average farm labourer can be trusted to carry it out with half an hour's instruction from the master.

"It may be permissible to suggest that those farmers (and there are hundreds of such) that now know from experience the benefit of this work, could do much good to their neighbours by showing them the results on crops that have been sprayed under favourable conditions. In order to exhibit the striking effect of spraying, a breadth of the weed should be left unsprayed in each field, or, perhaps better, keep some plots unsprayed by covering them at the time with a sheet or old bags. Neighbouring farmers visiting these fields a week afterwards would have then an opportunity of seeing, in well-defined contrast, the difference between the sprayed and unsprayed crops.

"For successful working, the chief points to remember are:—

"To have everything in readiness well beforehand, and the labourer trained, so that it is only necessary to give the order to start when the weather is fine. The material must be pure and correctly measured and the spray fine so as to cover every plant.

"In the early development of this process, a good deal was heard of a smooth-leaved Charlock (Wild Radish) that is not so susceptible to the spray. The quantity of this kind of weed has evidently been much overestimated. During 1902 practically nothing has been heard of it, although the amount of spraying is now tenfold.

"Successful spraying should result in average increase in yield of corn from four to six bushels per acre, and the destruction of 95 per cent. of Charlock, but when the weed is in dense mats, one plant overlapping another, a smaller proportion will be killed, but the increased yield of corn will be greater.

"It is remarkable that spraying Charlock, though largely practised in some districts, is quite neglected in others which have quite as much need for it. In such districts farmers should not lose time in obtaining information about this valuable process with a view to its adoption.

"At the present time I see no reason to alter the main conclusions arrived at in 1901. They have been thoroughly corroborated by extensive experience in 1902, and, with slight additions, are as follows:—

- "1. That Charlock can be destroyed in growing corn crops, without injury to the latter, by spraying with 50 gallons of 3 per cent. solution of pure sulphate of copper per acre.
- "2. That, generally, the corn crops are much improved and give a better yield where Charlock is destroyed, and that young grass seeds and clover in the corn remain uninjured.
- "3. That spraying early, when the weed is young and in soft fibre, is most successful.
- "4. That the larger sprayers with powerful pumps are more successful than the smaller machines.
- "5. The profit of the operation is diminished in proportion to the delay which occurs after the Charlock is in the most suitable condition for being destroyed.
- "6. That the increased yield of corn by the destruction of Charlock leaves a profit after all the expenses of spraying have been defrayed, and that the annual value of all Charlock-infested land will increase until the extermination of the weed is complete.

"My sincere thanks are due to many gentlemen in England and Scotland, by whose kindness I have been offered facilities to pursue these investigations and by wide practice to justify the opinions I have expressed.

"G. F. STRAWSON."

In addition to the leaflet by Mr. Strawson, the following is taken from the Board of Agriculture of England Leaflet No. 63:—

"The injury induced by Charlock is partly direct and partly indirect. It competes with crops for light and air, that is to say, it overgrows more or less completely and smothers other plants with which it is associated. It also robs crops of a part of their nutriment, and prevents their deriving full benefit from the moisture of the soil. But, in other ways, though more indirectly, this weed may be the cause of much loss. The turnip 'fly,' for instance, would be unable to exist in early summer, when the cultivated crops on which it preys are, for the most part, absent from our fields, did it not find weeds like Charlock to supply it with food. Then, again, the microscopic fungus that causes finger-and-toe finds a congenial habitat in the roots of this plant, which may thus do much to carry the disease over the years that separate two turnip crops.

"In 1897 attention was called to the possibility of getting rid of Charlock in corn crops by means of the application of certain solutions which, it was contended, could destroy the weed without injuring the cereal. During the past two years this method of dealing with the pest has been extensively

tested in great Britain, and, as a whole, the results have been successful. The substances chiefly tried have been copper sulphate and iron sulphate, and good results have been got with both. Although the former at present costs about 30s. per cwt. as against 4s. to 5s. in the case of the latter, the solution of iron sulphate must be used so much stronger than the other that the difference in the cost of material is less per acre than would at first sight appear. Moreover, copper sulphate deteriorates less by keeping, is more easily manipulated, and does less injury to the clothes of the workmen.

"Experience indicates that good results will, as a rule, be got by dressing an acre with, at most, 40 gallons of a 4 per cent. solution of copper sulphate, or with a similar quantity of a 15 per cent. solution of iron sulphate. To make the former, dissolve 16 lbs. of copper sulphate, costing about 4s. 6d., in 40 gallons of soft water; while, in the latter case, 60 lbs. of iron sulphate, costing about 2s. 6d., are required in a similar quantity of water. Somewhat better results will be got by dressing an acre with 16 lbs. of copper sulphate dissolved in 60 gallons of water—thus making a $2\frac{3}{4}$ per cent. solution—but, although this entails no extra outlay for material, it implies an increased expenditure on account of labour. Then, again, in place of applying 40 gallons per acre of a 4 per cent. solution at a single operation, superior results may sometimes be got by applying 30 to 40 gallons per acre of a 2 per cent. solution at a somewhat early stage of the growth of the weed, and a similar quantity ten days or a fortnight later. As compared with a single dressing, this involves no extra expenditure on material, but it entails the application of about twice as much water, and the crop suffers more from mechanical injury. Good results have occasionally been got with weaker solutions and with smaller quantities than those indicated, but, on the whole, those recommended above have proved most effective.

"A convenient method of procedure is to have two 40-gallon barrels in use, so that while the contents of one are being distributed the other may be used for the preparation of a fresh solution.

"It may be noted that certain weeds closely allied to the common Charlock—especially Wild Radish, or White Charlock, and Smooth-leaved Charlock—are not infrequently met with, and these do not readily yield to treatment. Other weeds, such as docks and thistles, are more or less crippled, without being destroyed by the solutions.

"To obtain the best results, it is best to attend to the following points:—

- "1. The weed should not exceed three inches in height at the time of spraying, though fair success has sometimes attended treatment almost up to the time of flowering.
- "2. The solution must be made with clean and, if possible, soft water, and the vessels used in the process should be made of wood.

- "3. The material, especially copper sulphate, should be bought under a guarantee of purity of 98 per cent. It should be obtained powdered, not in crystals. This facilitates solution.
- "4. The solution must be applied by means of a machine that generates a fine spray under air pressure. A hand machine of good construction will dress three or four acres in a day, while a horse machine will cover nearly ten times as much. The latter distributes the solution more thoroughly and equally.
- "5. Rain immediately after spraying will interfere with success, and the calmer the weather the more evenly and effectively is the solution distributed. Moderate dampness of the crop at the time of spraying is no disadvantage, and better results will be got in dull weather than in bright sun."

The following method of eradication by cultivation and hand-pulling is taken from "The Weeds of Ontario," by Prof. F. C. Harrison, B.S.A., Ontario Agricultural College —

"ERADICATION.—Owing to the great vitality of the seed, Mustard is a very hard weed to eradicate. The seeds, once in the ground, live for years and continue to germinate as they are brought near the surface. Hence, it takes patience, a great deal of labour, and a long time to get rid of the weed when it once gets possession of the land. When present only in small amounts, hand-pulling is the best method, provided the pulling is done before seeds have formed; and as persons pulling in a hurry cannot wait to examine for seed, it is best to put the weeds, as they are pulled, in bundles where they can be burned when dry.

"When fields are overrun with the weed, it is best to proceed as follows: Harrow stubble ground early after harvest, or gang-plough and harrow. As soon as the weeds have had time to sprout, cultivate thoroughly; repeat cultivation at intervals, and rib up with a double mould board plough the last thing in the fall. Put in a hoed crop, either roots or corn, the following spring, and cultivate it thoroughly throughout the growing season. Cultivate and harrow well two or three times after roots or corn, having first run the plough along each row of corn roots to cut the roots and turn them up, and rib up before the frost. (If the plough is used after corn or roots, it is likely to bring more seed to the surface.) Sow a crop of grain the following spring and seed with clover. Pull weeds by hand out of the grain crop; take a crop or two of hay or pasture; plough the clover sod shallow, not more than four inches, early after harvest, say the 1st to the 15th of August, and harrow at once. Let it stand a couple of weeks, then cultivate, the same way as it was ploughed, two or three inches deep, with a spring-tooth cultivator. After a while, cross-cultivate a little deeper. If possible, cultivate a third or even a fourth time, going a little deeper each time; then, if you can manage to do so, rib it up with a double mould board plough, as

you would for a crop of turnips. When this is done, the available plant food (clover roots, etc.) is preserved in the centre of the drills, the water runs off early in the spring, and the drills can be levelled with the cultivator and harrow either for spring grain or hoed crops.

"This method will not only clean land but will greatly enrich it."

NOTE.—As both Copper Sulphate and Iron Sulphate are often called by the common names of blue stone and blue vitriol, farmers and others are particularly warned that only *pure copper sulphate* must be used or failure and consequent disappointment will result. This chemical should always be obtained from reliable chemists and not from ordinary store-keepers, who may possibly be ignorant of the difference between copper and iron sulphate. Be particular as to time of spraying, as on that depends the success of the work; a few days' delay may result in failure.

J. R. ANDERSON,

Deputy Minister of Agriculture.

Department of Agriculture,

Victoria, B. C., January 30th, 1903.

VICTORIA, B. C.:

Printed by RICHARD WOLFENDEN, Printer to the King's Most Excellent Majesty.
1903.