**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1994

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé la meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☒ Coloured covers/
Couverture de couleur

☐ Covers damaged/
Couverture endommagée

☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée

☐ Cover title missing/
La titre de couverture manque

☐ Coloured maps/
Cartes géographiques en couleur

☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)

☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur

☐ with other material/
avec d'autres documents

☒ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distorsion le long de la marge intérieure

☐ Blank leaves added during restoration may appear
within the text. Whenever possible, these have
been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans la texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.

☐ Additional comments: / Pagination is as follows: p. [413]-421.
Commentaires supplémentaires:

☐ Coloured pages/
Pages de couleur

☐ Pages damaged/
Pages endommagées

☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées

☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées

☐ Pages detached/
Pages détachées

☒ Showthrough/
Transparence

☐ Quality of print varies/
Qualité inégale de l'impression

☐ Continuous pagination/
Pagination continue

☐ Includes index(es)/
Comprend un (des) index

Title on header taken from: /
Le titre de l'en-tête provient:

☐ Title page of issue/
Page de titre de la livraison

☐ Caption of issue/
Titre de départ de la livraison

☐ Masthead/
Générique (périodiques) de la livraison

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☐	☐	☒	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Douglas Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche sheet contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Douglas Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaît sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482-0300 - Phone
(716) 288-5989 - Fax

UNIVERSITY OF TORONTO
STUDIES

PHYSIOLOGICAL SERIES

NO. 10: ON THE PROBABLE NATURE OF THE SUBSTANCE
PROMOTING GROWTH IN YOUNG ANIMALS, BY CASIMIR
FUNK AND ARCHIBALD BRUCE MACALLUM

(REPRINTED FROM THE JOURNAL OF BIOLOGICAL CHEMISTRY, VOL. XXIII)

THE UNIVERSITY LIBRARY: PUBLISHED BY
THE LIBRARIAN, 1916

NCE
SIMIR

UNIVERSITY OF TORONTO LIBRARY
COMMUNITAS DE MONTREAL

CHURCH, RICHARD L. (1911-1988) M.A., LL.D., D.D.
(President of the University)

CHURCH, W. L. (1911-1988) M.A.

CHURCH, W. H. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

CHURCH, J. J. (1911-1988) M.A.

STUDIES ON GROWTH.

II. ON THE PROBABLE NATURE OF THE SUBSTANCE PROMOTING GROWTH IN YOUNG ANIMALS.

BY CASIMIR FUNK¹ AND ARCHIBALD BRUCE MACALLUM²

(From the Cancer Hospital Research Institute, London, and the Biochemical Laboratory, University of Toronto.)

(Received for publication, October 8, 1915.)

Since the remarkable work of Osborne and Mendel,³ Hopkins,⁴ and McCollum and Davis,⁵ who have shown that a young animal requires something special besides the usual food constituents for its process of growth, there has been much discussion as to the exact nature of this product. In our first paper⁶ we discussed this subject and pointed out that most workers regard the growth factor as being closely associated with fats. The experimental evidence which led to this opinion was brought forward first by McCollum, but has become especially significant since Osborne and Mendel³ and later Osborne and Wakeman⁷ found that purified butter, which in their opinion was free from nitrogen, was still able to promote growth in young rats. In our earlier paper we found that even the purified butter contains traces of nitrogenous substances, and therefore might possibly contain traces

¹ The work was begun at the Cancer Hospital Research Institute, London, England; the experiments illustrated by the curves and tables were carried out in the Biochemical Laboratory, University of Toronto, during the tenure of a Senior Research Fellowship in the Department of Medical Research, and the expenses were defrayed by a grant from this department.

² Osborne, T. B., and Mendel, L. B., *Carnegie Institution of Washington, Publication No. 156*, pts. i and ii, 1911.

³ Hopkins, F. G., *Jour. Physiol.*, 1912, xlv, 425.

⁴ McCollum, E. V., and Davis, M., *Jour. Biol. Chem.*, 1913, xv, 167.

⁵ Funk, C., and Macallum, A. B., *Ztschr. f. physiol. Chem.*, 1914, xcii, 13

⁶ Osborne and Mendel, *Jour. Biol. Chem.*, 1913-14, xvi, 423.

⁷ Osborne, T. B., and Wakeman, A. J., *Jour. Biol. Chem.*, 1915, xxi, 91.

of vitamins, an opinion which is shared by McCollum and Davis. Recently MacArthur and Luckett⁹ have found that the growth-promoting substance is not contained in the ether-soluble fraction, and they also suggest the possibility of vitamins as a factor.

To complete our first paper we carried out a series of experiments on young rats with ordinary butter and purified butter fat as the fat fraction of the diet. Both with butter and purified butter fat all the animals died after five to seven weeks, although on the former diet a slight initial advantage was noticed. Repeating these experiments on pigeons we convinced ourselves of the inability of both diets to prevent the onset of beri-beri symptoms, indicating the absence of or an insufficient quantity of beri-beri vitamin.

In our subsequent experiments we increased the percentage of butter and pure butter fat in the diets from 12 to 30 per cent with the same negative result in all cases. On both diets the rats showed, twenty-four to forty-eight hours before death, a condition of spastic contraction resembling somewhat avian beri-beri or infantile tetany.

We also carried out experiments in which increasing amounts of starch were replaced by equivalent quantities of unpolished rice, with lard as the fat fraction of the diet. The results, which were tending to support the vitamin theory of growth of one of us,¹⁰ were still unsatisfactory, although the diet containing the largest percentage of unpolished rice proved to be much better both for maintenance and growth than those diets in which butter was used; yet this was finally inadequate as the animals declined after nine weeks.

Finally we made up diets, with butter used as fat, to which from 2 to 6 per cent of dried brewer's yeast was added. On this diet we have obtained successful growth and maintenance. As the growth-promoting factor is beyond question contained in yeast, we intend to fractionate the yeast and show which fraction contains the hypothetical growth substance.

The butter was purified as indicated in our earlier paper.⁸ The casein was purified by extraction with hot alcohol. The fuel

⁸ McCollum and Davis, *Jour. Biol. Chem.*, 1914, xix, 245.

⁹ MacArthur, C. G., and Luckett, C. L., *Jour. Biol. Chem.*, 1915, xx, 161.

¹⁰ Funk, C., *Ztschr. f. physiol. Chem.*, 1913, lxxxviii, 352.

value of the food and also the amount of the food absorbed from the intestine were controlled by means of an adiabatic calorimeter. The food mixture used, with the exception of the butter, was found to produce beri-beri in earlier experiments by one of us¹¹ on pigeons. Special experiments were carried out in order to ascertain the value of a food mixture consisting of casein, starch, cane-sugar, salt mixture, and butter for pigeons. All the pigeons fed on this food developed a typical beri-beri and no difference was noticed between the ordinary butter and the purified butter.

The results presented in this paper are a selection of a large number of experiments of uniform character. Two rats of the same sex were used in each experiment. The four diets used had the following composition.

Diet	I	II	III	IV
	per cent	per cent	per cent	per cent
Casein...	22	22	22	22
Sugar.....	10	10	10	10
Starch....	33	33	31	27
Butter (ordinary)	30		30	30
(purified).		30		
Agar.....	2	2	2	2
Salt mixture*..	3	3	3	3
Yeast (dry)....			2	6

* The composition of the salt mixture was the same as in the experiments of Osborne and Mendel.¹²

Experiment I.—Chart I. The curves represent the average weight of two rats each of which was fed on Diet I (ordinary butter). As represented by the upper curve the rats showed a slight initial gain in weight and maintenance for about 20 days; then a rapid decline set in with fatal termination after 36 days. The lower curve represents the average weight of two rats which were changed from Diet I to Diet IV on the twentieth day. The effect of the addition of dried brewer's yeast was striking; the rats suddenly recovered and grew normally up to the end of the experiment. The intake of food and the absorption from the intestine are recorded below.

¹¹ Funk, *Ztschr. f. physiol. Chem.*, 1914, lxxxix, 373.

¹² Osborne and Mendel, *Jour. Biol. Chem.*, 1913, xv, 311.

Days	Upper curve, Rats 23 and 26.				Lower curve, Rats 27 and 28.			
	Average weight.	Food intake.	Food.	Feces.	Average weight.	Food intake.	Food.	Feces.
	gm.	gm.	cal.	cal.	gm.	gm.	cal.	cal.
0	26.5				21.0			
4	32.25	35.1	120.5	3.67	27.5	18.15	80.2	3.0
8	36.5	19.6	95.4		26.5	15.0	74.7	
12	37.5	21.1	103.7	3.37	27.5	18.65	91.7	2.2
16	37.0	21.3	103.6		26.5	17.5	85.5	
20	33.0	21.0	125.1	2.66	27.0	25.65	134.2	5.4
24	30.5	21.5	111.2		38.0	43.2	220.5	
28	28.5	14.5	74.5	4.36	40.5	42.1	223.4	9.7
32	26.5	10.4	53.2		62.5	53.66	285.8	
36	23.5	10.2	52.2	2.21	74.5	57.95	308.8	16.00
40	Died				81.5		280.5	
44					85.0	54.15	288.5	
48					92.0	58.0	300.5	

This experiment was repeated on six male and two female rats with identical results.

Experiment II—Chart II. Here we have used purified butter fat. Each curve represents the average weight of two male rats. The results are similar to those of the first experiment, only the maintenance period was slightly shortened. The same marked recovery was observed on changing from Diet II to Diet IV. The details of the experiment are recorded below.

Days	Upper curve, Rats 31 and 32.				Lower curve, Rats 29 and 30.			
	Average weight.	Food intake.	Food.	Feces.	Average weight.	Food intake.	Food.	Feces.
	gm.	gm.	cal.	cal.	gm.	gm.	cal.	cal.
0	43.5				29.5			
4	49.0	32.2	176.8	8.34	36.5	25.85	141.2	4.25
8	52.0	26.35	143.0		39.5	18.3	99.0	
12	48.5	23.4	127.8	7.35	42.0	17.8	97.2	3.58
16	46.0	23.8	129.8		39.0	18.65	101.8	
20	44.7	23.75	129.6	8.00	36.0	17.75	96.9	3.15
24	42.0	24.1	131.5		31.5	17.65	96.6	
28	42.5	18.5	109.0	3.95	48.0	41.35	229.5	8.04
32	39.5	14.15	77.2		58.5	54.45	290.2	
36	36.5	11.85	68.2		74.5	61.85	324.3	13.09
40	32.5	6.70	36.79		80.0	53.5	285.0	
44	Died				94.5	57.1	304.2	
48					103.0	61.70	328.8	

This experiment was repeated on seven male rats and one female with the same result.

FIG. 1. Upper curve, Rats 25 and 26. Lower curve, Rats 27 and 28. On the curve at the point (*) Diet I was changed to Diet IV; Rats 25 and 26 died after 36 days on Diet I.

FIG. 2. Upper curve, Rats 31 and 32. Lower curve, Rats 29 and 30. At the point (*) on lower curve Diet II was changed to Diet IV. Rats 31 and 32 died after 40 days on Diet II.

Experiment III.—Chart III. In this experiment the upper curve illustrates the weight of two male rats, and the lower curve the weight of two female rats. Up to 20 days Diet III was used (with 2 per cent yeast), being then replaced by Diet IV. In one set this was continued to the end of the experiment (lower curve). In the other experiment Diet IV was replaced after 32 days, by Diet II (without yeast); the growth ceased abruptly and the animals rapidly declined. No marked difference was noticed between the diets containing 2 and 6 per cent of yeast. The details of the experiment are recorded below.

Days.	Upper curve, Rats 33 and 34.				Lower curve, Rats 35 and 36.			
	Average weight.	Food intake.	Food.	Feces.	Average weight.	Food intake.	Food.	Feces.
	gm.	gm.	cal.	cal.	gm.	gm.	cal.	cal.
0	20.0				18.5			
4	29.0	29.1	151.2	5.31	27.5	26.2	136.2	3.41
8	35.0	23.05	117.3		32.0	19.4	97.9	
12	37.0	25.95	132.4	7.30	37.0	26.65	137.1	4.39
16	41.5	26.60	137.2		39.5	29.85	148.1	
20	43.5	30.85	161.5	6.45	42.5	34.15	178.7	7.44
24	47.0	41.15	217.5		49.5	44.6	236.8	
28	59.0	44.0	233.5	8.06	58.0	43.5	230.0	11.13
32	69.0	49.5	262.6		69.5	48.75	259.5	
36	72.5	41.05	225.5	9.53	80.0	60.70	323.5	13.68
40	74.5	34.15	187.7		87.5	56.70	302.2	
44	71.0	30.85	169.6		90.0	51.90	276.5	
48	68.5	32.90	180.9		94.0	58.25	310.4	

This experiment was repeated with two additional female rats with the same result.

Our new series of experiments clearly show the absolute inability of either butter or purified butter fat to stimulate the growth of young rats. This result could only be expected from our experiments with the same diet on pigeons, which have shown the absence of vitamins in butter. From the tables we see a strict relationship between the growth observed and the food taken in. We also notice that the addition of yeast stimulates, directly or indirectly, the appetite. The calorific determination of the food and feces shows in all cases a practically complete absorption of the food by the digestive tract, as the calorific value of the feces represents but from 2.5 to 5 per cent of the calories taken in by

FIG. 3. Upper curve, Rats 33 and 34. Lower curve, Rats 35 and 36. To the left of the dotted line Diet III; to the right, Diet IV. At the point (*) Rats 33 and 34 were changed from Diet IV to Diet II.

the animals. Whether yeast alone without butter (replaced by lard) will produce normal growth in rats, and whether there are one or more components in yeast which stimulate growth, is under investigation.

UNIVERSITY OF TORONTO STUDIES

PHYSIOLOGICAL SERIES

- No. 1: The structure, micro-chemistry and development of nerve-cells, with special reference to their nuclein compounds, by F. H. SCOTT \$0.50
- No. 2: On the cytology of non-nucleated organisms, by Professor A. B. MACALLUM 0.75
- No. 3: Observations on blood pressure, by R. D. RUDOLF... 0.75
- No. 4: The chemistry of wheat gluten, by G. G. NASMITH... 0.50
- No. 5: The palaeochemistry of the ocean, by Professor A. B. MACALLUM 0.25
- No. 6: The absorptio of fat in the intestine, by G. E. WILSON 0.50
- No. 7: The distribution of fat, chlorides, phosphates, potassium and iron in striated muscle, by MAUD L. MINTEN..... 0.25
- No. 8: Surface tension and vital phenomena, by Professor A. B. MACALLUM 1.00
- No. 9: On the distribution of potassium in renal cells, by C. P. BROWN 0.25