2004 Annual Report

SUBLETTE MULE DEER STUDY (PHASE II): *Long-term monitoring plan to assess potential impacts of energy development on mule deer in the Pinedale Anticline Project Area.*

Prepared for:

Questar Exploration and Production Company
Independence Plaza
1050 17th St., Suite 500
Denver, CO 80265

TRC Mariah Associates Inc.
605 Skyline Road
Laramie, WY 82070

Bureau of Land Management
Pinedale Field Office
432 E. Mill Street
PO Box 768
Pinedale, WY 82941

Wyoming Game and Fish Department
Pinedale Regional Office
PO Box 850
Pinedale, WY 82941

Prepared by:

Hall Sawyer
Ryan Nielson
Lyman McDonald
Dale Strickland

Western EcoSystems Technology, Inc.
2003 Central Avenue
Cheyenne, WY 82001

November 2004

QL
737
.U55
S399
2004

ACKNOWLEDGEMENTS

Phase II of the Sublette Mule Deer Study has been a cooperative effort among agencies, industry, and private organizations. This project was largely funded by Questar Exploration and Production Company (QEP) and subcontracted through TRC Mariah Associates Inc. (TRC). The Pinedale Field Office of the Bureau of Land Management (BLM) provided additional funding. The Jackson/Pinedale Region of the Wyoming Game and Fish Department (WGFD) provided logistical support, project assistance, and was responsible for portions of the data collection. Many thanks to Ron Hogan (QEP), Jane Seiler (QEP), Steve Belinda (BLM), Keith Andrews (BLM), Karen Rogers (BLM), Pete Guernsey (TRC), Scott Smith (WGFD), Herb Haley (WGFD), Bernie Holz (WGFD), Scott Edberg (WGFD), Scott Werbelow (WGFD), Dan Stroud (WGFD), Dean Clause (WGFD), Dennis Almquist (WGFD), Brad Hovinga (WGFD), Fred Lindzey (University of Wyoming), Hawkins & Powers Aviation, and Gary Lust (Mountain Air). Thanks to John Amos (SkyTruth) for image processing.

LIST OF ACRONYMS

BACI	Before-After Control-Impact
BLM	Bureau of Land Management
CR	County Road
DAU	Data Analysis Unit
EIS	Environmental Impact Statement
GIS	Geographic Information System
GPS	Global Positioning System
HA	Hunt Area
JCR	Job Completion Report
MWRC	Mesa Winter Range Complex
NEPA	National Environmental Policy Act
NGO	Non-Government Organization
PAPA	Pinedale Anticline Project Area
PFWRC	Pinedale Front Winter Range Complex
QEP	Questar Exploration and Production Company
ROD	Record of Decision
RSPF	Resource Selection Probability Function
TPB	Trapper's Point Bottleneck
TRC	TRC Mariah Associates, Inc.
UD	Utilization Distribution
USGS	United States Geological Survey
UW	University of Wyoming
VHF	Very High Frequency
WEST	Western EcoSystem Technology, Inc.
WGFD	Wyoming Game and Fish Department

BLM Library
Don't to
Denver Federal Center
P.O. Box 25047
Denver, Colorado 80225

TABLE OF CONTENTS

	Page
1.0 OVERVIEW.....	1
2.0 SUBLETTE MULE DEER STUDY (PHASE II).....	3
2.1 INTRODUCTION.....	3
2.2 STUDY AREA.....	6
2.3 METHODS.....	6
2.3.1 Deer Capture.....	6
2.3.2 Winter Movement and Distribution Patterns.....	7
2.3.3 Population Characteristics.....	7
2.3.3.1 Abundance and Density Estimates.....	7
2.3.3.2 Reproduction.....	9
2.3.3.3 Adult female winter survival.....	9
2.3.3.4 Over-winter fawn survival.....	9
2.3.4 Direct Habitat Loss.....	9
2.3.5 Resource Selection.....	10
2.3.5.1 Study Area Delineation.....	10
2.3.5.2 Predictor Variables.....	10
2.3.5.3 Modeling Procedures.....	10
2.4 RESULTS.....	11
2.4.1 Deer Capture.....	11
2.4.2 GPS Data Collection.....	13
2.4.3 Winter Movement and Distribution Patterns.....	13
2.4.3.1 Treatment Area (PAPA).....	13
2.4.3.2 Control Area (PFWRC).....	14
2.4.4 Population Characteristics.....	28
2.4.4.1 Abundance and Density Estimates.....	28
2.4.4.2 Reproduction.....	30
2.4.4.3 Over-winter fawn survival.....	31
2.4.4.4 Adult winter survival.....	32
2.4.5 Direct Habitat Loss.....	36
2.4.5.1 Pre-Development.....	36
2.4.5.2 Year 1 of Development.....	36
2.4.5.3 Year 2 of Development.....	37
2.4.5.4 Year 3 of Development.....	38
2.4.5.5 Year 4 of Development.....	39
2.4.6 Resource Selection.....	40
2.4.6.1 Pre-Development: Winters 1998-99 and 1999-00.....	42
2.4.6.2 Year 1 of Development: Winter 2000-01.....	43
2.4.6.3 Year 2 of Development: Winter 2001-02.....	44
2.4.6.4 Year 3 of Development: Winter 2002-03.....	45
2.5 DISCUSSION AND FUTURE DIRECTION.....	46
2.6 PROJECT TIMELINE.....	50
3.0 LITERATURE CITED.....	51
APPENDIX A: Equations Used to Calculate Abundance and Density Estimates.....	55
APPENDIX B: Resource Selection Modeling Procedures.....	56
APPENDIX C: GPS locations and 60-mile seasonal migration of deer #863.....	59

BLM
Bldg. 510
Denver, CO 80225
P.O. Box 250
Denver, Colorado 80225

LIST OF TABLES

	<u>Page</u>
Table 2.1 Number and type of radio-collars functioning in treatment and control areas during the 2003-04 winter.....	12
Table 2.2 Summary statistics for abundance and density estimates in the treatment area during February 2002, 2003, and 2004.....	28
Table 2.3 Summary statistics for abundance and density estimates in the control area during February 2002, 2003, and 2004.....	29
Table 2.4 Mule deer fawn:doe ratios measured for treatment (Mesa) and control (Pinedale Front) areas by Wyoming Game and Fish Department, 1992-2004.....	30
Table 2.5 Mule deer count data and calculations for over-winter fawn survival in the control area (Pinedale Front), 1999-2004.....	31
Table 2.6 Mule deer count data and calculations for over-winter fawn survival in the treatment area (Mesa), 1999-2004.....	31
Table 2.7 Winter (2003-04) survival rates and summary statistics for adult female deer in treatment and control areas.....	32
Table 2.8 Summary of annual and cumulative direct habitat loss (i.e., surface disturbance) associated with road networks and well pads on the Mesa, 2000-2003.....	38
Table 2.9 Habitat variables and estimated coefficients for winter mule deer resource selection probability functions (RSPF), 1998-2003.....	40
Table 2.10 Optimal values for predictor variables with quadratic terms.....	41

LIST OF FIGURES

Figure 1.1 Location of Pinedale Anticline Project Area in western Wyoming (Fig.1 from BLM 2000).....	2
Figure 2.1 Location of the Sublette Mule Deer Data Analysis Unit, Wyoming Game and Fish Department Hunt Areas, and the Mesa and Pinedale Front Winter Range Complexes.....	4
Figure 2.2 Location of 38 quadrats used in control area during 2003 helicopter surveys.....	8
Figure 2.3 Location of 70 quadrats used in control area during 2004 helicopter surveys.....	8
Figure 2.4 GPS locations ($r = 923$) of deer #860 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.....	15

LIST OF FIGURES (continued)

	<u>Page</u>
Figure 2.5 GPS locations ($r = 1,367$) of deer #872 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.....	16
Figure 2.6 GPS locations ($r = 1,367$) of deer #878 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.....	17
Figure 2.7 GPS locations of all deer ($n = 3$) and land ownership of the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.....	18
Figure 2.8 GPS locations ($r = 924$) of deer #861 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.....	19
Figure 2.9 GPS locations ($r = 919$) of deer #869 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.....	20
Figure 2.10 GPS locations ($r = 1,978$) of deer #871 in the Pinedale Front Winter Range Complex, November 1, 2003 – April 15, 2004.....	21
Figure 2.11 GPS locations ($r = 1,969$) of deer #873 in the Pinedale Front Winter Range Complex, November 1, 2003 – April 15, 2004.....	22
Figure 2.12 GPS locations ($r = 1,942$) of deer #874 in the Pinedale Front Winter Range Complex, November 1, 2003 – April 15, 2004.....	23
Figure 2.13 GPS locations ($r = 1,378$) of deer #876 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.....	24
Figure 2.14 GPS locations ($r = 3,354$) of deer #871 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).....	25
Figure 2.15 GPS locations ($r = 3,344$) of deer #873 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).....	26
Figure 2.16 GPS locations ($r = 3,328$) of deer #874 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).....	27
Figure 2.17 December fawn:doe ratios in treatment and control areas, 1999-2004.....	30
Figure 2.18 Estimated over-winter fawn survival in treatment and control areas, 1999-2004.....	31
Figure 2.19 Winter survival rates of adult female radio-collared deer in treatment and control areas, 1998-2004.....	32
Figure 2.20 Satellite image of the Mesa on October 1999, prior to development of the Pinedale Anticline Project Area (PAPA).....	36

LIST OF FIGURES (continued)

	<u>Page</u>
Figure 2.21 Satellite image of the Mesa taken in August 2001, following 1 full year of gas development in the Pinedale Anticline Project Area (PAPA).....	37
Figure 2.22 Satellite image of the Mesa taken in October 2002, following 2.3 years of gas development in the Pinedale Anticline Project Area (PAPA).....	38
Figure 2.23 Satellite image of the Mesa taken in September 2003, following 3.25 years of gas development in the Pinedale Anticline Project Area (PAPA)..	39
Figure 2.24 Relationship between probability of habitat use and distance to well pad, for development years 1-3.....	41
Figure 2.25 Surface map depicting probability of habitat use for mule deer prior to development (1998-99 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).....	42
Figure 2.26 Surface map depicting probability of mule deer habitat use during Year 1 of development (2000-01 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).....	43
Figure 2.27 Surface map depicting probability of mule deer habitat use during Year 2 of development (2001-02 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).....	44
Figure 2.28 Surface map depicting probability of mule deer habitat use during Year 3 of development (2002-03 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).....	45

LIST OF PHOTOS

Photo 2.1 Winter deer mortality on south end of Mesa.....	33
Photo 2.2 Cross-section of mule deer femur. Bone marrow color (reddish) and consistency (gelatinous) suggests cause of death was chronic starvation...	33
Photo 2.3 Snow conditions in PFWRC along the Big Sandy River (view north), February 2004.....	34
Photo 2.4 Snow conditions in PFWRC along the Big Sandy River (view south), February 2004.....	34
Photo 2.5 Snow conditions in MWRC near Mount Airy (view north), February 2004..	35
Photo 2.6 Snow conditions in MWRC near Two Buttes (view south), February 2004....	35

1.0 OVERVIEW

In 1998 the Wyoming Cooperative Fish and Wildlife Research Unit began the Sublette Mule Deer Study, a collaborative effort with industry, agencies, and private organizations intended to examine movement patterns and population characteristics of the Sublette mule deer (*Odocoileus hemionus*) herd in western Wyoming. Although a variety of agencies and non-government organizations (NGOs) contributed to the study, it was funded largely by industry (Ultra Petroleum). Concurrently, the Bureau of Land Management (BLM), in compliance with the National Environmental Policy Act (NEPA), initiated an Environmental Impact Statement (EIS) to assess natural gas development in the 300-mi² Pinedale Anticline Project Area (PAPA) (BLM 2000) (Figure 1.1). Because the PAPA provides important winter range to a large segment of the Sublette mule deer herd, there were concerns about the potential effects gas field development may have on the deer population.

The Sublette Mule Deer Study was originally designed to have two phases. The first phase of the study was intended to gather information needed by agencies to improve management of the Sublette deer herd, including the identification of seasonal ranges, determination of migration routes, and estimation of survival rates (Sawyer and Lindzey 2001). Additionally, these data were collected so that pre-development information on the mule deer population would be available if Phase II of the study materialized. Phase II was envisioned as a long-term study that would examine the potential impacts of energy development on mule deer, using treatment and control areas, with energy development as the treatment. The BLM completed the PAPA EIS and released their record of decision (ROD) in July of 2000 (BLM 2000). Phase I of the Sublette Mule Deer Study was completed in March of 2001 (Sawyer and Lindzey 2001). Following a 1-year pilot study (Sawyer et al. 2002) funded by QEP, Phase II was initiated in December of 2002, as a Before-After/Control-Impact (BACI) study design (Green 1979, Morrison et al. 2001) that uses the PAPA as a treatment area and a portion of the Pinedale Front as the control area. Mule deer population characteristics (survival, reproduction, density) and habitat use in relation to development features will be measured in both areas, and over time, performance of mule deer in the PAPA will be compared to those in the control area, both before and after the treatment. This report summarizes the results from the 2004 study period.

Figure 1.1 Location of Pinedale Anticline Project Area in western Wyoming (from BLM 2000).

2.0 SUBLETTE MULE DEER STUDY

2.1 INTRODUCTION

Western Wyoming is home to the largest, most diverse ungulate populations in the Rocky Mountain region. Maintenance of these populations and protection of their habitats are primary concerns among the public and state and federal agencies. Because of their large numbers and economic importance, mule deer continue to be a top priority for the Wyoming Game and Fish Department (WGFD). The Sublette mule deer herd unit includes 15 hunt areas (HA) (130, 138-142, 146, 150-156, and 162) (Figure 2.1) and has a post-season population objective of 32,000 (WGFD 2002). Results from the Sublette Mule Deer Study (Sawyer and Lindzey 2001) indicate that these mule deer seasonally migrate 60-100 mi from winter range near Pinedale, Wyoming to summer in portions of the Salt River Range, Wyoming Range, Wind River Range, Gros Ventre Range, and Snake River Range. During the lengthy spring and fall migrations, mule deer spend a substantial amount of time, often 4-5 months out of the year, on mid-elevation transition ranges that connect summer and wintering areas. By late-fall, most mule deer annually converge in the Green River Basin to winter in one of two major complexes; the Mesa Winter Range Complex (the Mesa) and the Pinedale Front Winter Range Complex (the Pinedale Front) (Figure 2.1). Generally, the Mesa includes the PAPA and those wintering areas west of US 191, while the Pinedale Front includes those areas east of US 191 to the base of the Wind River Mountains.

Population parameters measured during the 3-year (1998-2000) Phase I study (WGFD 2002, Sawyer and Lindzey 2001) suggested the Sublette deer herd was a healthy and productive population prior to development of energy resources on the PAPA. Annual survival rates of radio-collared adult females ($n=149$) averaged 85% and were consistent with populations studied in other western states (Unsworth et al. 1999). Fawn:doe ratios, an indicator of reproductive success, were among the highest in the state, averaging >75 fawns per 100 does for the study period and approximately 70 fawns per 100 does over the last decade (WGFD 2002). Although the Sublette deer herd has been very productive in the past and recent studies have improved management, this deer herd is similar to others in the region in that habitat loss due to urban expansion and energy development continue to create major management concerns.

Natural gas production in Wyoming has steadily increased since the mid-1980s, particularly in the five counties that form the southwest quarter of the state: Sublette, Fremont, Lincoln, Uinta, and Sweetwater. This area of the state contains some of the largest and most productive gas fields in the nation, including the Jonah, Continental Divide/Wamsutter, Fontenelle, Big Piney-LaBarge, Moxa Arch, Riley Ridge, Desolation Flats, and the Pinedale Anticline. While coalbed methane development in the Powder River Basin has received much attention as of late, Sublette County continues to produce approximately 2.5 times more gas than Campbell County. Sublette County is the state's largest producer and accounts for >30% of all Wyoming gas production (BLM 2002). Because of renewed political and economic support for developing domestic energy reserves, natural gas exploration, development, and production are at an all time high in Wyoming and expected to increase.

Because the PAPA encompasses the Mesa, which is used by thousands of mule deer, pronghorn (*Antilocapra americana*), and sage grouse (*Centrocercus urophasianus*), development of this area may have adverse impacts on wildlife. Impacts to wildlife may be direct or indirect. Direct impacts include the loss of habitat to well pads, access roads, and pipelines. Indirect impacts may include changes in distribution, stress, or activity caused by increased human activity associated with energy development (i.e., traffic, noise, human presence). The best way to evaluate the impact(s) of energy development on wildlife populations is through long-term studies where pre-development data, such as, estimates

Figure 2.1 Location of the Sublette Mule Deer Data Analysis Unit, Wyoming Game and Fish Department Hunt Areas, and the Mesa and Pinedale Front Winter Range Complexes.

of survival and reproduction are available. Because these studies are by necessity observational, determining cause and effect relationships is very difficult. Simply documenting a behavioral response (e.g., avoidance, acclimation, dispersal) to a disturbance adds very little to our knowledge of the impact, if it cannot be linked to the survival or reproductive success of the species involved. And conversely, documenting a change in reproduction or survival does not add significantly to our understanding of the impact if the cause (e.g., weather, habitat loss, disease) of the change cannot be determined. Because

of the difficulty with designing and funding long-term studies, impacts of energy development on free-ranging ungulate populations are poorly understood and often debated. However, both direct and indirect impacts associated with energy development have the potential to affect ungulate population dynamics, particularly when disturbances are concentrated on winter ranges, where energetic costs are great and animals occur at high densities.

The major shortcoming of efforts to evaluate the impact(s) of disturbances on wildlife populations is that they seldom are addressed in an experimental framework, but rather tend to be short-term and are almost always observational. Brief, post-development monitoring plans associated with regulatory work generally result in little or no information that allow agencies and industry to assess impacts on wildlife or identify new, and potentially more effective, mitigation measures. On the other hand, long-term studies are difficult to implement because they are expensive and require interagency and industry cooperation and commitment. Additionally, the acquisition of pre-development data on movement patterns and population characteristics, and identification of suitable control and treatment areas is extremely uncommon. The situation in the PAPA and upper Green River Basin is unique because all the necessary information is available to conduct a BACI study to suggest if, and if so, how natural gas development affects the PAPA mule deer population.

The basic idea with a BACI study design is that the potentially impacted (treatment) site is sampled both before and after the time of the disturbance (e.g., energy development), and one or more control sites that do not receive any disturbance are sampled at the same time (Manly 2001). The assumption is that any naturally occurring changes will be similar at the control and treatment sites, and in the absence of the treatment the parameters of interest will be similar for both areas, or at least the magnitude of the differences will be relatively constant from year to year. Thus, potential changes at the treatment site may be attributed to the disturbance. It is not critical that the control and treatment populations be identical, only that the subpopulations are independent and that both respond to the same environmental factors.

For this study, energy development on the Mesa is considered the treatment and a portion of the Pinedale Front serves as the control area. The Pinedale Front consists mostly of federal lands located along the southwest portion of the Wind River Range, where no energy development is anticipated. The Pinedale Front is a suitable control site because: 1) there is little or no exchange of deer between the Mesa and Pinedale Front, 2) the two deer subpopulations use separate winter ranges, but share common transition and summer ranges, so they have comparable foods available during parturition and arrive on winter ranges in similar condition, 3) although the two deer subpopulations occupy distinct winter ranges, they are in close proximity to one another (15-30 mi), so both are exposed to similar weather patterns and environmental conditions, 4) habitat characteristics on both winter ranges are similar and dominated by sagebrush communities, and 5) population characteristics of the two subpopulations have consistently tracked one another prior to development of the PAPA.

We believe four population parameters should be monitored to detect the potential impacts of energy development on mule deer, including: 1) adult doe survival, 2) over-winter fawn survival, 3) reproduction, and 4) density. As these parameters are measured in treatment and control areas, comparisons can be made, and over time, the potential impacts of energy development on mule deer may be assessed. If mule deer in the PAPA continue to function as well as before development and as well as those in the control area it would suggest that energy development has no adverse impacts on mule deer in the region. If however, mule deer survival or reproduction in the PAPA decreases, while the same parameters in the control area remain unchanged or increase, then energy development may be the cause of those declines. Again, this does not demonstrate a cause-effect relationship; rather it is simply one piece in a weight of evidence approach, where the BACI study design examines several direct (e.g., survival, reproduction) and indirect (e.g., habitat use, displacement) parameters that are statistically analyzed and carefully interpreted.

Results from Phase I identified seasonal migration routes and distribution of deer in the Mesa and Pinedale Front (Sawyer and Lindzey 2001). Although mule deer migrations of >60 mi have been reported in parts of Idaho (Thomas and Irby 1990) and Montana (Mackie et al. 1998), mule deer on and adjacent to the PAPA are likely the most migratory deer in the western states, annually migrating 60-100 mi between winter and summer ranges. Because these deer are highly mobile and demonstrate strong fidelity to seasonal ranges, the potential for energy development, or other human disturbances, to disrupt migratory routes and/or winter distribution patterns exists. While changes in distribution or migratory patterns may not necessarily result in decreased deer survival or reproduction, it is useful to include within the monitoring plan to: 1) document if migration routes remain intact, 2) document if deer continue using pre-development winter ranges, 3) provide industry and agencies with accurate, precise movement data for site-specific analyses (e.g., seasonal range designation or comparison of effects of multiple well pads versus single well pad), 4) identify mitigation opportunities on and off-site treatment and control areas (e.g., migration corridors, habitat improvements), and 5) allow for analyses that estimate and describe indirect habitat loss (e.g., avoidance of roads or well pads) or changes in habitat use.

Properly designed long-term monitoring and examination of adult survival, over-winter fawn survival, reproduction, density, and seasonal distribution/movement patterns will allow for population-level inferences concerning the potential impacts of energy development on mule deer.

2.2 STUDY AREA

The PAPA is located in west-central Wyoming in Sublette County, near the town of Pinedale (Figure 1.1). The PAPA is characterized by sagebrush communities and riparian habitats associated with the Green and New Fork Rivers. Elevations range from 6,800 to 7,800 ft. The PAPA consists primarily of federal lands (80%) and minerals (83%) administered by the BLM. The state of Wyoming owns 5% (15.2 mi²) of the surface and another 15% (46.7 mi²) is private. Aside from the abundant energy resources, the PAPA is an important area for agriculture and provides winter range for 4,000-6,000 mule deer, 2,000-3,000 pronghorn, and 3,000-4,000 sage grouse. While the project area is fairly large, most deer occur in the northern portion of the PAPA, an area locally known as "The Mesa", which includes approximately 100-mi². In July of 2000, the BLM approved the development of 700 producing well pads in the PAPA and recognized that this may require as many as 900 well pads to be constructed and drilled (BLM 2000). Additionally, 401 mi of pipeline and 276 mi of access roads were approved for development of energy resources on the PAPA.

2.3 METHODS

2.3.1 Deer Capture

Helicopter net-gunning was used to capture deer across winter ranges in treatment (Mesa) and control (Pinedale Front) areas. Captured deer were fitted with collars supporting either a GPS or VHF radio transmitter. Both types of collars were equipped with mortality sensors that change pulse rate if the collar remains stationary for more than 8 hours. The VHF collars (Advanced Telemetry Systems, Isanti, MN) were duty-cycled to transmit signals October 1 through May 31. The GPS collars (Telonics, Mesa, AZ) were store-on-board units capable of storing approximately 3,000 locations and programmed to obtain fixes every 2 hours during winter months (November-April). Additionally, each GPS collar was equipped with a remote release mechanism programmed to activate at a specified time, so that collars could be retrieved and data downloaded.

2.3.1 Winter Movement and Distribution Patterns

Data collected from GPS-collared deer accurately identified winter distribution, movement patterns, and migration routes of the marked deer on and adjacent to winter ranges. Over time, these data will be used in conjunction with pre-development data to identify changes in mule deer movement patterns, habitat use and distribution relative to energy development activities. Because a portion ($n=11$) of GPS collars are to remain on the same deer for consecutive winters (2003-04 and 2004-05), some data for the 2003-04 winter will not be available until 2005.

2.3.3 Population Characteristics

2.3.3.1 Abundance and Density Estimates

Deer abundance and density were estimated in treatment (Mesa) and control (Pinedale Front) areas using aerial counts of deer in systematically sampled 1-mi² quadrat units. Winter distribution data collected from radio-collared deer in the study area between 1998 and 2003 was used to delineate 68-mi² and 70-mi² sampling frames for the treatment and control areas, respectively. Sampling frames were expected to contain high-densities of deer so stratification was unnecessary. We sampled 34 quadrats from each sampling frame, covering approximately 50% of the geographic area. Equations used to calculate abundance and density estimates were taken from Thompson et al. (1998) and are reproduced in Appendix A. Standard 90% confidence intervals were calculated using a Z statistic.

The size of the sampling frame in the control area has changed over the course of the study (See Section 2.4.4.1). During the first year of surveys (2002) we identified a 35-mi² sampling frame that we believed represented the core winter range in the Pinedale Front. During 2003 we made some slight modifications to improve our sampling and used a similar 38-mi² sampling frame (Figure 2.2). However, during the 2003 surveys many of our marked deer moved out of the sampled area. At this time it became apparent that these deer utilize a much larger area than we originally thought. To accurately adjust the size and extent of our sampling frame we conducted a telemetry flight prior to the 2004 survey to adjust the size of our sampling frame based on locations of marked deer. The new sampling frame was 70-mi² (Figure 2.3), nearly double the size of the 2002 and 2003 frames and approximately the same size as the sampling frame for the treatment area.

Group size and vegetative cover may significantly influence visibility bias in ungulate helicopter surveys (Samuel et al. 1987). However, the treatment and control areas for this study consist of homogenous sagebrush stands with no tree cover. Additionally, telemetry data from Phase I indicated male and female deer did not winter in areas with different habitat characteristics, so potential group size variation resulting from sexual segregation should not influence counts. Further, when survey areas contain large concentrations of animals that are widely distributed, recognition of individual groups may be near impossible. Attempting to determine visibility correction factors for groups is likely not feasible in these situations (Samuel et al. 1987). Counts of animals within the sampled quadrats are assumed to provide valid indices on density and abundance. That is, if not all animals present were detected, we assume the same visibility bias in both treatment and control areas over time.

Counts were conducted from a piston-powered Bell helicopter flown approximately 100-150 ft above ground and at speeds of 20-40 knots. The northeast UTM coordinates for each quadrat were programmed into a GPS unit on the helicopter. Quadrat perimeters were then flown clockwise, such that the observer was positioned on the inside, while the pilot navigated. A real-time flight path was traced into the on-board GPS and once the perimeter was established the quadrat interiors were systematically searched. Observer and navigator collectively detected deer groups and determined whether groups were inside or outside quadrat boundaries. Deer detected inside and moving out were considered in the quadrat, while deer detected outside and moving in the quadrat were considered out. Half of the deer

detected on perimeter boundaries were considered in the quadrat. For each quadrat, the observer recorded total number of deer, number of deer groups, and total search time.

Figure 2.2 Location of 38 quadrats used in control area during 2003 helicopter surveys.

Figure 2.3 Location of 70 quadrats used in control area during 2004 helicopter surveys.

2.3.3.2 Reproduction

Doe:fawn ratios are commonly used as an index to herd productivity or reproduction. Doe:fawn ratios were calculated from composition data collected during the WGFD's annual helicopter surveys in December, consistent with the previous 10 years of WGFD data collection (WGFD 2002). Sample sizes were adequate to obtain desired levels of precision in ratio estimates (Czaplewski et al. 1983).

2.3.3.3 Adult Female Winter Survival

Adult doe survival was estimated from telemetry records using the Kaplan-Meier procedure (Kaplan and Meier 1985). We attempted to maintain a sample of 30 marked deer in both control and treatment areas. Marked deer were located at least once per month, December through May.

2.3.3.4 Over-winter Fawn Survival

Deer from both the Mesa and Pinedale Front congregate on the northern ends of their respective winter ranges every spring which allows large numbers (>1,000) of animals to be counted and classified. Ground-based composition surveys conducted in April were used to calculate post-winter adult:fawn ratios. These data were used in conjunction with adult survival rates and December adult:fawn ratios to estimate over-winter fawn survival, using the change-in-ratio estimator from White et al. (1996):

$$\hat{S}_f = \hat{S}_a \times \frac{B}{A}, \text{ where } A = \text{count of December fawns/count of December adults}$$

$$B = \text{count of April fawns/count of April adults}$$

$$\hat{S}_a = \text{estimate of adult survival}$$

Adult survival rates were estimated from telemetry records, rather than carcass counts. The delta method (Seber 1982) was used to estimate variance.

2.3.4 Direct Habitat Loss

Satellite imagery and geographic information system (GIS) software were used to digitize road networks and well pads associated with natural gas development in the northern portion of the PAPA (i.e., The Mesa), from 2000 through 2003. Areas within the PAPA, but outside the Mesa were not considered. Landsat images were purchased from the USGS and processed by SkyTruth (Shepherdstown, WV). Images were generally obtained in early fall (i.e., September-October), after most annual construction activities (e.g., well pad and road building) were complete, but prior to snow accumulation. Pipelines and seismic tracks were not included in this analysis. Roads and well pads were digitized in ArcView[®] (ESRI, Redlands, California, USA). Length of road segments and size of well pads were calculated in ArcView[®] (ESRI, Redlands, California, USA). Acreage estimates associated with road networks were based on an average road width of 30 ft. We recognize there is some error associated with the digitizing process, however it is expected to be minimal and the resulting digital GIS coverages are considered the best available data. During the digitizing process we assumed full reclamation of well pads had not occurred, since the gas field is only 3 years old and successful reclamation (i.e., re-establishment of native plant species) of native shrub communities in arid environments is extremely difficult and unlikely to occur during a short time period.

2.3.5 Resource Selection

2.3.5.1 Study Area Delineation

The treatment study area was defined by mapping 30,626 GPS locations from 34 mule deer over a five-year period (1998 to 2002) and creating a minimum convex polygon (MCP), consistent with McClean et al. (1998)'s recommendation that study-area level of habitat availability should be based on the distribution of radio-marked animals. Additionally, the MCP generated from GPS data was consistent with winter distribution patterns documented for this deer population using >60 VHF radio-collars, between 1998 and 2000 (Sawyer and Lindzey 2001).

2.3.5.2 Predictor Variables

We identified five habitat variables as important landscape predictors of winter mule deer distribution: elevation (V_1), slope (V_2), aspect (V_3), road density (V_4), and distance to well pad (V_5). We did not include vegetation as a variable because the sagebrush-grassland vegetation is generally homogeneous and difficult to divide into finer vegetation classes across the study area. We believed differences in sagebrush characteristics could be largely explained by elevation, slope, and aspect. A GIS was used to measure variables V_1 to V_5 for sampled habitat units. The Spatial Analyst Extension for ArcView® (ESRI, Redlands, California, USA) was used to calculate slope and aspect from a 26 x 26 m digital elevation model (DEM) (USGS 1999). Grid cells with slopes >2 degrees were assigned an aspect value (i.e., NE = $2^\circ < \text{aspect} = 90^\circ$, SE = $90^\circ < \text{aspect} = 180^\circ$, SW = $180^\circ < \text{aspect} = 270^\circ$, NW = $270^\circ < \text{aspect} = 360^\circ$), while grid cells with slopes of $= 2$ degrees were considered flat and had no aspect value. Road and well pads were annually digitized from Landsat satellite images acquired from the U.S. Geological Survey and processed by SkyTruth (Shepherdstown, WV). Images were obtained from late-spring or early fall, prior to snow accumulation, but after most annual natural gas development activities were complete. Road density was calculated using three buffer sizes; 0.5 km radius, 1.0 km radius, and 1.5 km radius, and the modeling process was used to determine which had the most predictive power. A pairwise correlation analysis (PROC CORR; SAS Institute 2000) was conducted on the linear forms of the variables to identify possible multicollinearity (Neter et al. 1996) and determine if any variables should be dropped from the analysis prior to modeling.

No distinction was made between producing and developing well pads. The rationale for treating all well pads equally was based on the assumption that there are two potential factors that may affect deer habitat selection near well pads: 1) direct loss of habitat, usually in the form of a 3-4 acre surface disturbance and 2) human activity (e.g., traffic, noise) associated with the well pad. Because the PAPA occurs in a dry desert shrub community, re-establishment of native shrub communities in disturbed areas is difficult and unlikely to occur in < 10 years, following current reclamation standards and practices in the project area. Based on our experience in the PAPA and given development is in its early stages (i.e., < 5 years), we felt it was appropriate to assume similar habitat loss among all well pads. Additionally, there is no evidence from the PAPA that suggests the type of well pad (e.g., developing, producing) is an accurate indicator of the amount of human activity that occurs at the site. Without an accurate measure of human activity (e.g., traffic), we believed it was inappropriate to distinguish between producing and developing well pads in this analysis.

2.3.5.3 Modeling Procedures

Our modeling approach consisted of four basic steps: 1) estimate the utilization distribution's (UD) for GPS-collared deer during each winter, 2) use the height of the UD as a response variable in a multiple regression setting to model the probability of habitat selection for each deer, 3) develop a population-level model from the individual deer models for each winter, and 4) map predictions of population-

level models from each winter. Refer to Appendix B for modeling procedure details.

2.4 RESULTS

2.4.1 Deer Capture

We captured and radio-collared 21 adult female deer on December 15, 2003. Deer capture in the PAPA was restricted to those areas where deer congregate across the northern end of the PAPA in early winter, as they move onto the Mesa from Trapper's Point and/or the Ryegrass/Grindstone area. We assumed this represented a random sample of deer in the subpopulation because the deer were congregated on the north end, before they moved south to their respective winter ranges. For the same reason, deer capture in the Pinedale Front was restricted to the Big Sandy area; bounded to the north and west by the Big Sandy River, east to the Prospects, and south to Elk Mountain. Of the 21 deer captured, 18 were equipped with GPS radio-collars and 3 equipped with traditional VHF radio-collars. All GPS collars were store-on-board units equipped with VHF transmitters on 24-hour duty cycles, 8-hour mortality sensors, and remote-release mechanisms programmed to drop collars at 0800 hours on April 15, 2004 or April 15, 2005. The programming schedule for collars was as follows:

Generation II GPS collars:

- obtain 1 location every 3 hours December 20, 2002-April 15, 2003
- obtain 1 location every 3 hours December 20, 2003-April 15, 2004
- for a possible ~1,800 locations

Generation III GPS collars:

- obtain 1 location every 2 hours December 20, 2002-April 15, 2003
- obtain 1 location every 2 hours November 01, 2003-April 15, 2004
- for a possible ~3,300 locations

Consistent with previous years, our goal was to maintain a sample size of 30 deer in each area, including 10 GPS and 20 VHF radio-collars (Table 2.1).

Additionally, we captured deer #863 and recovered the GPS collar that had failed to drop off in 2003. This collar collected locations every 3 hours for 22 months (January 2002 - October 2003), including the spring and summer of 2003. Appendix C includes a map that illustrates the 60-mi seasonal migration of deer #863.

Table 2.1 Number and type of radio-collars functioning in treatment and control areas during the 2003-04 winter.

Treatment Area (The Mesa)		Control Area (Pinedale Front)	
Deer ID	Collar Type	Deer ID	Collar Type
810*	VHF	804	VHF
801	VHF	807	VHF
803	VHF	811	VHF
805*	VHF	812	VHF
806	VHF	814	VHF
809	VHF	816	VHF
813	VHF	820	VHF
814*	VHF	821	VHF
815	VHF	823	VHF
817	VHF	824	VHF
818	VHF	825	VHF
822	VHF	826	VHF
827	VHF	829	VHF
830	VHF	833	VHF
842	VHF	835*	VHF
844	GPS	836	VHF
845	VHF	848	VHF
849	VHF	850	VHF
852	VHF	851	VHF
853	VHF	861	GPS
854	VHF	864	GPS
855	GPS	867	GPS
8553	VHF	869	GPS
860	GPS	870	GPS
862	GPS	871	GPS
865	GPS	873	GPS
866	GPS	874	GPS
868	GPS	876	GPS
870*	VHF	958*	VHF
872	GPS	VHF = 20 GPS = 9 Total = 29	
878	GPS		
884	GPS		
886	VHF		
887	GPS		
889	GPS		
898*	VHF		
905*	VHF		
989	VHF		
946*	VHF		
976*	VHF		
VHF = 28 GPS = 12 Total = 40			

* Radio-collars left over from Phase I (Sawyer and Lindzey 2001).

2.4.2 GPS Data Collection

We intended to collect data from 9 GPS collars following the 2003-04 winter. All nine release mechanisms successfully dropped collars on April 15, 2004. However, because of several deaths due to the severity of the winter, we recovered 11 GPS collars, including 5 from the Mesa and 6 from the Pinedale Front. Four of the collars (#865, #871, #873, and #874) contained data for consecutive winters (2002-03 and 2003-04).

Of the 11 collars that were retrieved, six were Generation III and five were Generation II. All collars functioned properly and collected the expected number of locations. Generation II and Generation III collars averaged 923 and 1,521 successful fix attempts, respectively. Consistent with GPS performance in previous years (Sawyer et al. 2002), success rates for GPS fix attempts were very high (99%) and locations precise (~93% 3-D locations for GEN II and 73% 3-D for GEN III). A minimum of four satellites are needed to generate 3-D locations, which typically have less than 20-meter error (Di Orto et al. 2003).

2.4.3 Winter Movement and Distribution Patterns

2.4.3.1 Treatment Area (Mesa):

Individual maps were compiled for three GPS-collared deer (#860, #872, and #878) captured in the treatment area (Figures 2.4-2.6). Data from the eight remaining collars were not mapped because they will not be recovered until April 15, 2005. Data from deer #865 was not mapped because it died early in the winter (~January 15).

Deer locations were color-coded by month so that timing of winter movements and distribution patterns could be easily deciphered (Figures 2.4-2.6). Distribution and movement patterns were variable among deer. Despite unusually harsh winter conditions deer shifted areas of use through the winter and utilized a large portion of the Mesa. Locations from GPS-collared deer were similar with observations and locations of VHF-collared deer.

Most deer did not begin their northerly migration off the Mesa until late-March and early-April, presumably because of the above average snow conditions. Consistent with previous years, all deer traveled to the Cora Butte area via the Trapper's Point Bottleneck (TPB) (Sawyer and Lindzey 2001, Sawyer et al. 2003). With the exception of deer #872, rate of movement in and near the TPB increased, as evidenced by distance between locations (Figures 2.2-2.6). Deer #872 spent ten days (April 5-14, 2004) in the TPB.

Figure 2.7 includes locations ($r = 3,657$) from all three deer and illustrates the importance of BLM lands to this mule deer population. Boundaries between private and BLM lands generally correspond with habitat type and topography; with private lands consisting of flat river bottoms and agricultural areas, whereas BLM lands contain sagebrush hills in drier, more rugged terrain. Mule deer demonstrated a strong affinity to the sagebrush-dominated BLM lands.

Figure 2.7 clearly defines the TPB, located 7 mi west of Pinedale, near the junction of US 191, WYO 352, and CR 110. Sawyer and Lindzey (2001) defined bottlenecks as "those areas along migration routes where topography, vegetation, development and/or other landscape features restrict animal movements to narrow or limited regions." Bottlenecks create management concerns because the potential to disrupt or threaten established migratory routes are much greater in these areas. This naturally-occurring bottleneck is approximately 1 mi in width and length, restricted to the southwest by the Green River riparian complex and to the northeast by the New Fork/Duck Creek riparian complex. Sagebrush habitats

north and south of US 191 are used extensively by mule deer during certain times of the year (Sawyer and Lindzey 2001). Mule deer use the narrow strip of sagebrush connecting the 2 areas to cross US 191. Development of small, fenced house lots adjacent to BLM lands has narrowed the effective width of the TPB to < 0.5 mi.

2.4.3.2 Control Area (Pinedale Front):

Individual maps were compiled for six GPS-collared deer (#861, #869, #871, #873, #874, and #876) captured in the control area (Figures 2.8-2.13). Data from three collars (#867, #864, and #870) were not mapped because they will not be recovered until April 15, 2005. Again, deer locations were color-coded by month so that timing of winter movements and distribution patterns could be easily deciphered (Figures 2.8 – 2.13). Consistent among deer was their mobility and tendency to shift areas of use through the winter, utilizing areas that exceeded 100-mi². Similar to last year (Sawyer et al. 2003), deer moved outside the core winter range area around Buckskin Crossing to peripheral areas, such as Elk Mountain, Squaw Teat, and a large section of Little Sandy Creek east of the Prospects. While distribution patterns of deer were variable across the winter range, the migratory routes to and from the winter range were nearly identical among deer. The strong affinity deer demonstrate to this migration route is clearly illustrated by comparing locations from deer #871, #873, and #874. These three deer collected data for two consecutive years (2002-03 and 2003-04) and used the same migration route both years, during spring and fall movements (Figures 2.14-2.16).

Most deer began migrating north along the Pinedale Front in mid-March. Deer that winter along the Pinedale Front were known to migrate northerly along the Wind River Range to the New Fork Lake area before shifting their migration in a westerly route towards the Hoback Basin and adjacent mountain ranges (Sawyer and Lindzey 2001). Details of this migration route, in terms of size, width, specific location, and deer fidelity were unknown prior to GPS data collected over the last two years. Consistent with last year, all six GPS-collared deer captured in the control area migrated along what appeared to be a distinct movement corridor located at the base of the Wind River Range. While deer sometimes remained in one area for a number of days, they appeared to follow a well-defined route that rarely exceeded 1-mile in width and covered a distance of 50 mi.

The migration route deer followed from the Buckskin Crossing area took them north across the Big Sandy River, then northerly across the sagebrush flats below Sheep Creek and Muddy Creek. Deer then moved into slightly rougher terrain among the boulders and sagebrush draws east of CR 353, south of the East Fork, and west of Irish Canyon. Deer then moved northerly, crossing the East Fork and Pocket Creek approximately 2-3 mi east of CR 353. Once across Pocket Creek, deer contoured through the sagebrush slopes and aspen pockets, northerly through Cottonwood Creek and Silver Creek. From Silver Creek, deer continued northwesterly across Lovett and Scab Creek. Deer continued to contour across the sagebrush slopes below Soda Lake, towards the outlet of Boulder Lake. Deer crossed Boulder Creek near the outlet of Boulder Lake, and then moved north to Fall Creek, apparently to avoid an agricultural area between Fall Creek and Pole Creek. Deer crossed Fall Creek just below the confluence of Meadow Creek, and then moved northwesterly toward the outlet of Fremont Lake. Deer crossed Pine Creek at the Fremont Lake Bottleneck, as described by Sawyer and Lindzey (2001). GPS collars released on April 15, 2004 and no movements were recorded beyond the outlet of Fremont Lake. Last year (2003), however, spring conditions allowed deer to continue movements through the Fremont Lake Bottleneck, and north along the Willow Creek Road and Fremont Ridge before collars were released (Sawyer et al. 2003: Figure 2.11). Deer moved within ½-mi either side of the Willow Lake Road from Soda Lake to the outlet of Willow Lake. No movements were recorded beyond the outlet of Willow Lake for GPS collars released on April 15, 2003.

Figure 2.4 GPS locations ($n = 923$) of deer #860 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.

Figure 2.5 GPS locations ($n = 1,367$) of deer #872 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.

Figure 2.6 GPS locations ($n = 1,367$) of deer #878 in the Pinedale Anticline Project Area (PAPA), December 22, 2003 – April 15, 2004.

Figure 2.7. GPS locations of all deer ($n = 3$) and land ownership of the Pinedale Anticline Project Area (PAPA), December 22, 2003–April 15, 2004.

Figure 2.8. GPS locations ($n = 924$) of deer #861 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.

Figure 2.9. GPS locations ($n = 919$) of deer #869 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.

Figure 2.10 GPS locations ($n = 1,978$) of deer #871 in the Pinedale Front Winter Range Complex, November 1, 2003–April 15, 2004.

Figure 2.11 GPS locations ($n = 1,969$) of deer #873 in the Pinedale Front Winter Range Complex, November 1, 2003 – April 15, 2004.

Figure 2.12 GPS locations ($n = 1,942$) of deer #874 in the Pinedale Front Winter Range Complex, November 1, 2003–April 15, 2004.

Figure 2.13 GPS locations ($r = 1,378$) of deer #876 in the Pinedale Front Winter Range Complex, December 22, 2003 – April 15, 2004.

Figure 2.14 GPS locations ($n = 3,354$) of deer #871 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).

Figure 2.15 GPS locations ($n = 3,344$) of deer #873 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).

Figure 2.16 GPS locations ($n = 3,328$) of deer #874 in the Pinedale Front Winter Range Complex, December 20, 2002 – April 15, 2003 (green) and November 1, 2003 – April 15, 2004 (blue).

2.4.4 Population Characteristics

2.4.4.1 Abundance and Density Estimates

Helicopter flights were conducted on February 16-17, 2004 to count deer in selected 1-mi² quadrats of both treatment and control areas. Average flight time per quadrat was 10 minutes. Estimated deer abundance ($\hat{N}$) in the treatment area was $3,564 \pm 650$ and deer density ($\hat{D}$) was 52 ± 10 deer/mi² (Table 2.2). Deer abundance and density were lower than previous years. Since 2002, the standard errors and coefficients of variation have steadily decreased as the number of sampled quadrats (u) has increased.

Table 2.2 Summary statistics for abundance and density estimates in the treatment area during February 2002, 2003, and 2004.

February 2002, 2003, and 2004.				
Summary Statistics		Treatment Area (The Mesa)		
Year		2002	2003	2004
Total Quadrats (U)		68	66	68
Quadrats Sampled (u)		18	32	34
Deer Counted (N)		1,384	2,267	1,782
Density Estimate ($\hat{D}$)		77	71	52
Variance ($V\hat{a}r(\hat{D})$)		146	87	34
Standard Error ($SE(\hat{D})$)		12.07	9.30	5.82
90% Confidence Interval		(57, 97)	(56, 86)	(42, 62)
Abundance Estimate($\hat{N}$)		5,228	4,676	3,564
Variance ($V\hat{a}r(\hat{N})$)		673863	377132	156318
Standard Error ($SE(\hat{N})$)		821	614	395
90% Confidence Interval		(3878, 6578)	(3666, 5686)	(2914, 4214)
Coefficient of Variation ($CV(\hat{N})$)		16%	13%	11%

Estimated deer abundance ($\hat{N}$) in the control area was $2,411 \pm 541$ and deer density ($\hat{D}$) was 34 ± 8 deer/mi² (Table 2.3). Abundance and density estimates for the control area have been variable since 2002. Because the sampling frame in the control area did not reflect the area utilized by our marked population, abundance and density estimates are expected to be biased high during 2002 and biased low during 2003.

Table 2.3 Summary statistics for abundance and density estimates in the control area during February 2002, 2003, and 2004.

Summary Statistics	Control Area (Pinedale Front)		
	2002 ^a	2003 ^b	2004
Total Quadrats (U)	35	38	70
Quadrats Sampled (u)	7	18	34
Deer Counted (N)	810	849	1,171
Density Estimate ($\hat{D}$)	116	47	34
Variance ($\hat{V}\hat{a}r(\hat{D})$)	406	64	22
Standard Error ($SE(\hat{D})$)	20.14	8.01	4.70
90% Confidence Interval	(83, 149)	(31, 63)	(26, 42)
Abundance Estimate ($\hat{N}$)	4,050	1,792	2,411
Variance ($\hat{V}\hat{a}r(\hat{N})$)	496752	92661	108347
Standard Error ($SE(\hat{N})$)	705	304	329
90% Confidence Interval	(2891, 5209)	(1291, 2293)	(1870, 2952)
Coefficient of Variation ($CV(\hat{N})$)	17%	17%	14%

^a Abundance and density estimates expected to be high.

^b Abundance and density estimates expected to be low.

2.4.4.2 Reproduction

Year	Treatment	Control
	fawn:doe December	fawn:doe December
Pre-Development		
1992-93	62	61
1993-94	47	51
1994-95	61	72
1995-96	56	63
1996-97	73	75
1997-98	92	81
1998-99	67	76
1999-00	85	76
Average	68	69
Development		
2000-01	85	81
2001-02	69	71
2002-03	64	65
2003-04	78	78
Average	74	74

The WGFD conducted helicopter composition (buck:doe:fawn) surveys to collect pre-winter (December) information on the sex (i.e., buck or doe) and age (i.e., fawn or adult) structure of the population. A total of 9,704 deer were classified in December, 2003, including 3,264 on the Mesa and 6,440 on the Pinedale Front (S. Smith, WGFD, unpublished data). Estimated fawn:doe ratios were 78:100 in both areas (Table 2.4), well above the 12-year averages. Treatment and control areas have displayed similar trends in reproduction (fawn:doe ratios) prior to and since the PAPA ROD in 2000 (Figure 2.17).

Table 2.4 (Left) Mule deer fawn:doe ratios measured for treatment (Mesa) and control (Pinedale Front) areas by Wyoming Game and Fish Department, 1992-2004.

Figure 2.17 December fawn:doe ratios in treatment and control areas, 1999-2004.

2.4.4.3 Over-winter Fawn Survival

The WGFD conducted ground-based composition (adult:fawn) surveys to estimate post-winter fawn:adult ratios during April of 2004 (S. Smith, WGFD, unpublished data). A total of 1,534 and 2,212 deer were counted in the Mesa and Pinedale Front, respectively (Tables 2.5-2.6). Estimates of over-winter fawn survival were 0.36 and 0.33 in the Mesa and Pinedale Front, respectively (Tables 2.5-2.6). Until this year, over-winter fawn survival has generally been lower in the treatment area compared to the control, since the PAPA ROD in 2000 (Figure 2.18). Low fawn survival rates were indicative of the severe winter conditions during 2003-04 that appeared to affect both winter ranges.

Table 2.5 Mule deer count data and calculations for over-winter fawn survival in the control (Pinedale Front), 1999-2004.

Year	December Adults	December Fawns	April Adults	April Fawns	A*	B**	$\hat{S}_a$	$\hat{S}_f$
1999-00	2698	1517	959	494	0.56	0.52	0.83	0.76
2000-01	2853	1769	955	478	0.62	0.50	0.85	0.69
2001-02	4593	2455	790	300	0.53	0.38	0.85	0.60
2002-03	3565	1813	704	254	0.51	0.36	0.96	0.68
2003-04	3977	2463	1771	441	0.62	0.25	0.82	0.33

* A = count of December fawns/count of December adults

** B = count of April fawns/count of April adults

Table 2.6 Mule deer count data and calculations for over-winter fawn survival in the treatment (Mesa), 1999-2004.

Year	December Adults	December Fawns	April Adults	April Fawns	A	B	$\hat{S}_a$	$\hat{S}_f$
1999-00	2550	1547	1390	764	0.61	0.55	0.87	0.74
2000-01	2420	1458	1685	707	0.60	0.42	0.82	0.59
2001-02	2546	1275	1366	460	0.50	0.34	0.85	0.57
2002-03	1864	914	1489	470	0.49	0.32	0.88	0.57
2003-04	2063	1201	1215	319	0.58	0.26	0.79	0.36

Figure 2.18 Estimated over-winter fawn survival in treatment and control areas, 1999-2004.

2.4.4.4 Adult Winter Survival

Winter (December 15 – May 1) survival rates were estimated using the Kaplan-Meier technique (Kaplan and Meier 1985) and telemetry records of 68 radio-collared adult female deer, including 40 in the treatment and 28 in the control area (Table 2.1). Winter survival rates were 0.79 and 0.82 for the treatment and control areas, respectively (Table 2.7). Field necropsy of 10 animals indicated that starvation was the cause of death (Photos 1-2). Survival rates were lower in both areas than previous years (Figure 2.19) and likely a result of the severe winter conditions during the 2003-04 winter. Snow conditions during the February flights were the heaviest and most extensive since the inception of this study in 1998 (Photos 3-6). Most adult mortality occurred in March and April.

Table 2.7 Winter (2003-04) survival rates and summary statistics for adult female radio-collared deer in treatment and control areas.

Study Area	Time Period	N_1	N_2	$\hat{S}$	90% CI	SE
Pinedale Anticline (Treatment)	December 15, 2003 - April 15, 2004	40	8	0.79	0.71 to 0.86	0.06
Pinedale Front (Control)	December 15, 2003 - April 15, 2004	29	5	0.82	0.73 to 0.91	0.07

N_1 = number of available collars, N_2 = number of deaths, $\hat{S}$ = survival estimate, CI = confidence interval, SE = standard error

Figure 2.19 Winter survival rates of adult female radio-collared deer in treatment and control areas, 1998-2004.

Photo 2.1 Winter mule deer mortality on south end of Mesa.

Photo 2.2 Cross-section of femur from radio-collared mule deer on the Mesa, March 2004. Bone marrow color (reddish) and consistency (gelatinous) suggests chronic starvation.

Photo 2.3 Snow conditions in the Pinedale Front along the Big Sandy River (view north), February 2004.

Photo 2.4 Snow conditions in Pinedale Front along the Big Sandy River (view south), February 2004.

Photo 2.5 Snow conditions on the Mesa near Mount Airy (view north), February 2004.

Photo 2.6 Snow conditions on the Mesa near Two Buttes (view south), February 2004.

2.4.5 Direct Habitat Loss

2.4.5.1 Pre-Development

Prior to development, The Mesa portion of the PAPA was relatively undisturbed, with very few improved roads and approximately a dozen existing well pads (Figure 2.20).

Figure 2.20 Satellite image of the Mesa on October 1999, prior to development of the Pinedale Anticline Project Area (PAPA).

2.4.5.2 Year 1 of Development

The BLM's ROD for the PAPA was released in July, 2000. Accordingly, natural gas development was minimal during this year. Approximately 11 mi of new roads and 39 acres of well pads were constructed on the Mesa during 2000 (Table 2.8). Approximately 51% of total surface disturbance was associated with road building, while the other 49% was attributed to well pad construction (Table 2.8).

2.4.5.3 Year 2 of Development

2001 marked the first full calendar year of gas field development in the PAPA. Most development occurred along the central portion of the Mesa, adjacent to Lovatt Draw (Figure 2.21). Based on satellite imagery, approximately 13 mi of new roads and 113 acres of well pads were constructed on the Mesa during the first nine months 2001 (Table 2.8). Approximately 30% of total surface disturbance was associated with road building, while the other 70% was attributed to well pad construction (Table 2.8).

Figure 2.21 Satellite image of the Mesa taken in August 2001, following 1 full year of gas development in the Pinedale Anticline Project Area (PAPA).

2.4.5.4 Year 3 of Development

Similar to 2001, most development in 2002 occurred along the central portion of the Mesa, adjacent to Lovatt Draw, from the Paradise Road northwest to Stewart Point (Figure 2.22). Drilling activity was also evident on the northern Mesa, east of Stewart Point. Based on satellite imagery, approximately 18 mi of new roads and 201 acres of well pads were constructed on the Mesa between August 2001 and October 2002 (Table 2.8). Approximately 25% of total surface disturbance was associated with road building, while the other 75% was attributed to well pad construction (Table 2.8).

Figure 2.22 Satellite image of the Mesa taken in October 2002, following 2.3 years of gas development in the Pinedale Anticline Project Area (PAPA).

2.4.5.5 Year 4 of Development

Similar to 2001 and 2002, most gas development in 2003 occurred along the central portion of the Mesa, adjacent to Lovatt Draw, from the Paradise Road northwest to Stewart Point (Figure 2.23). Drilling activity was also evident on the northern Mesa, east of Stewart Point. Based on satellite imagery, approximately 14 mi of new roads and 237 acres of well pads were constructed on the Mesa between October 2002 and September 2003 (Table 2.8). Approximately 18% of total surface disturbance was associated with road building, while the other 82% was attributed to well pad construction (Table 2.8).

Figure 2.23 Satellite image of the Mesa taken in September 2003, following 3.25 years of gas development in the Pinedale Anticline Project Area (PAPA).

Table 2.8 Summary of annual and cumulative direct habitat loss (i.e., surface disturbance) associated with road networks and well pads on the Mesa, 2000-2003.

Year	Roads (mi)	Roads (acres) ^a	Well Pads (acres)	Total (acres)	% Roads	% Well Pads
2000	11.4	41	39	80	51%	49%
2001	13.1	48	113	161	30%	70%
2002	18.1	66	201	267	25%	75%
2003	13.9	51	237	288	18%	82%
Total	56.5	206	590	796	26%	74%

^a Based on an average road width of 30 feet.

2.4.6 Resource Selection

Note: Because some GPS-collars operate on the same deer for consecutive years, there is a lag time between when the data are collected and when they are analyzed. Resource selection analyses have been completed for data collected through the 2002-03 winter.

Population-level models (Table 2.9) and predictive maps (Figures 2.25-2.28) were estimated for four winter periods: Pre-Development (Winters 1998-99 and 1999-00), Year 1 of Development (Winter 2000-01), Year 2 of Development (Winter 2001-02), and Year 3 of Development (Winter 2002-03). Stepwise model building resulted in slightly different models for the three years of development. However, consistent predictor variables across years included slope, elevation, and distance to well pad. Road density and aspect were variable across years, in terms of their ability to predict deer use. Probability of mule deer habitat use increased as elevation increased. Slope, distance to well pad, and road density had non-linear (quadratic) effects in the models. When plotted against probability of use, the optimal values of the variables: slope, distance to well pad, and road density, occurs at the highest point of the curve, as illustrated in Figure 2.24. The optimal values can be approximated from the figures, but specific estimates were calculated using calculus (i.e. '1st derivative test') to find the value of the variable corresponding to the maximum probability of use. Table 2.10 lists the estimated optimal values for all predictor variables with quadratic terms in each model. These values were helpful in quantifying and interpreting how mule deer responded to predictor variables through time. Using the distance to well pad variable as an example, the optimal values increased as development progressed, indicating that mule deer were selecting for habitats farther away from well pads as development progressed from Year 1 to Year 3. This non-linear (quadratic) relationship suggests mule deer selected areas away from well pads, up to a certain distance. We believe this reflects the presence of less suitable habitats near the study area boundary, compared to the more suitable habitats in the central portion of the Mesa where most gas development has occurred. Additionally, habitat use near the study area boundary is likely influenced by factors outside the boundary, such as less desirable agricultural and riparian habitats (e.g., Green and New Fork Rivers), and other gas development infrastructure and activity (e.g., Sand Springs Draw).

Table 2.9 Habitat variables and estimated coefficients for winter mule deer resource selection probability functions (RSPF), 1998-2003.

Variable	Pre-Development ^a			Year 1			Year 2			Year 3		
	β	SE	P	β	SE	P ^b	β	SE	P ^b	β	SE	P ^b
Intercept	-29.736			-46.219	9.836	0.001	-44.398	6.655	<0.001	-76.067	8.681	<0.001
Elevation	0.009			0.014	0.004	0.009	0.013	0.003	<0.001	0.023	0.003	0.001
Slope	0.088			0.304	0.074	0.003	0.260	0.075	<0.001	0.250	0.095	0.069
Slope ²	-0.003			-0.018	0.005	0.004	-0.015	0.004	<0.001	-0.012	0.006	0.132
Well distance	0.116			4.327	1.903	0.967	2.834	1.283	0.017	7.593	2.430	0.04
Well distance ²	-0.007			-0.939	0.372	0.039	-0.408	0.174	0.012	-0.883	0.292	0.045
Road density	-0.185			-1.650	0.606	0.023	0.838	0.430	0.032	ns ^a		
Road density ²	-0.063			0.513	0.212	0.032	-0.442	0.161	0.005	ns		
Aspect = NE				ns			-0.481	0.151	0.007	ns		
Aspect = NW				ns			0.305	0.167	0.09	ns		
Aspect = SE				ns			0.051	0.154	0.743	ns		
Aspect = SW				ns			-0.321	0.165	0.072	ns		

^a Standard errors and P-values were not presented for the Pre-Development RSPF coefficients because stepwise model building was not performed

- ^b P-values for Year 1 based on student's t distribution with 9 degrees of freedom.
^c P-values for Year 2 based on student's t distribution with 14 degrees of freedom.
^d P-values for Year 3 based on student's t distribution with 6 degrees of freedom.
^e Not significant.

Figure 2.24 Relationship between probability of habitat use and distance to well pad, for development years 1-3.

Table 2.10 Optimal values for predictor variables with quadratic terms.

Year	Variable	Optimum Value
Year 1 of Development 2000-01	Distance to Well Pad ²	2.3 km
	Road Density ²	0 km/km ²
	Slope ²	8°
Year 2 of Development 2001-02	Distance to Well Pad ²	3.5 km
	Road Density ²	0.95 km/km ²
	Slope ²	9°
Year 3 of Development 2002-03	Distance to Well Pad ²	4.3 km
	Road Density ²	Ns
	Slope ²	10°

2.4.6.1 Pre-Development: Winters 1998-99 and 1999-00

The population-level model was estimated from a random sample of 4,469 habitat units and 953 VHF deer locations collected from 76 adult female mule deer during the winters (December 1- April 15) of 1998-99 and 1999-00. Because pre-development data consisted of many deer with relatively few locations, we could not treat the deer as the experimental unit and conduct the same model building process we did with the GPS-collared deer in subsequent years. Rather, we fit a model based on habitat variables that were consistently included in models from subsequent years (i.e., elevation, slope, distance to well, and road density). Consequently, we did not estimate standard errors or p-values for the coefficients (Table 2.9). Predicted probabilities indicate deer selected for areas with high elevation and moderate slopes.

Figure 2.25 Surface map depicting probability of mule deer habitat use prior to development (1998-99, 1999-00 winters). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).

2.4.6.2 Year 1 of Development: Winter 2000-01

The population-level model was estimated from a random sample of 4,489 habitat units and 18,706 GPS locations collected from 10 adult female mule deer during the winter (January 1- April 15) of 2000-01. The model included elevation, slope, distance to well pad, and road density (Table 2.9). Deer selected for areas with higher elevations, moderate slopes, low road densities, and away from well pads. Areas with the highest probability of use had elevations of 2,325 – 2,400 m, slopes of 15 degrees, road densities of 0.0 km/km², and were 2.3 km away from the nearest well pad. Predicted probabilities indicate deer use was lowest in those areas of the PAPA where well pads and associated road networks were developed (Figure 2.26). Portions of the study area that were classified as "high deer use" prior to development were classified as "low deer use" during year 1 of development. The model suggests these changes in deer distribution were due to well pad and road development.

Figure 2.26 Surface map depicting probability of mule deer habitat use during Year 1 of development (2000-01 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).

2.4.6.3 Year 2 of Development: Winter 2001-02

The population-level model was estimated from a random sample of 4,464 habitat units and 14,851 GPS locations collected from 15 adult female mule deer during the winter (January 4 – April 15) of 2001-02. The model included elevation, slope, aspect, distance to well pad, and road density (Table 2.9). Deer selected for areas with northwest aspect (positive coefficient, smallest p-value), high elevations, moderate slopes, low road densities, and away from well pads. Areas with the highest probability of use had northwest aspects, elevations of 2,325 – 2,400 m, slopes of 8 degrees, road densities <0.95 km/km², and were 3.5 km away from the nearest well pad. Predicted probabilities indicate deer use was lowest in those areas of the PAPA where well pads and associated road networks were developed (Figure 2.27). Portions of the study area that were classified as "high deer use" prior to development were classified as "low deer use" during year 2 of development. The model suggests these changes in deer distribution were due to well pad and road development.

Figure 2.27 Surface map depicting probability of mule deer habitat use during Year 2 of development (2001-02 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).

2.4.6.4 Year 3 of Development: Winter 2002-03

The model was estimated from a random sample of 4,462 habitat units and 5,131 GPS locations collected from 7 adult female mule deer during the winter (December 20 – April 15) of 2002-03. Our target sample of 10 marked animals was not met because several deer died early in the season and/or spent most of their time outside the study area. Deer were distributed farther north than most years (Sawyer et al. 2003). The RSPF included elevation, slope, and distance to well pad (Table 2.9). Deer selected areas with higher elevations, gentle slopes, and away from well pads. Areas with the highest probability of use had elevations of 2,325 – 2,400 m, slopes of 9 degrees, and 4.3 km away from the nearest well pad. Predicted probabilities indicate deer use was lowest in those areas of the PAPA where well pads were developed (Figure 2.28). Portions of the study area that were classified as "high deer use" prior to development were classified as "low deer use" during year 3 of development. The model suggests these changes in deer distribution were due to well pad development.

Figure 2.28 Surface map depicting probability of mule deer habitat use during Year 3 of development (2002-03 winter). Color-coded based on percentile of predictions (i.e., 0-25%, 25-50%, 50-75%, and 75-100%).

2.5 DISCUSSION AND FUTURE DIRECTION

Approximately 15,000 GPS locations were collected from 10 deer during the 2003-04 winter. Another 12 GPS collars should be recovered in 2005 and should provide an additional 24,000 locations from both the 2003-04 and 2004-05 winters. Although data collection is delayed a full year when GPS collars operate on the same deer for two consecutive winters, acquiring movement and distribution information from the same animal over a period of years provides useful year-to-year comparisons. Last year, for example, we were able to demonstrate that deer on the Mesa wintered farther north than in previous years (Sawyer et al. 2003). If we had not had GPS collars running for consecutive winters on the same deer, it could have been argued that we captured deer that only utilized the northern Mesa. And, this year we were able to demonstrate the strong affinity mule deer have for a 50-mile migration route along the Pinedale Front.

Basic distribution maps generated from GPS data illustrated winter use of the control and treatment areas, demonstrated the importance of BLM lands, and refined information on migration routes and seasonal ranges. Deer in the treatment area (Mesa) continued to utilize the Trapper's Point Bottleneck (TPB) as a migratory route between winter and spring/fall transition ranges. Deer movements through the TPB were quick, as evident by the distance between locations, but the TPB continued to function effectively during 2004. Agencies, industry, NGO's, and the public recognize the value of maintaining this movement corridor for the Sublette deer herd. Land-use decisions in and adjacent to the TPB should consider the migration routes and seasonal ranges of the Sublette deer herd.

Consistent with the 2002-03 winter, deer distribution and movement patterns in the control area (Pinedale Front) were variable during the 2003-04 winter. While deer spent some time in the original sampling frame (Figure 2.2), they also moved (10 – 15 mi) in all directions; south to Elk Mountain, southeast along the Big Sandy, easterly to the Little Sandy and Prospects, and northerly to Muddy Mountain. The ability to alter their rates of movements, to change their pathways, and occupy a variety of winter habitats as needed are behaviors that likely allow these deer to best exploit winter ranges. However, the unpredictable movement patterns have made calculation of abundance and density estimates difficult, as the size of the sampling frame has progressively increased over the study to encompass the range of our marked deer. Because the sampling frame did not reflect the area utilized by our marked population, abundance and density estimates are expected to be biased high during 2002, and biased low during 2003. Assuming the 2004 sampling frame accurately reflects the winter range extent used by marked deer, the abundance and density estimates should be more reliable than those calculated during 2002 and 2003.

Although winter distribution patterns varied among deer in the Pinedale Front, the migration route to northerly transition and summer ranges was surprisingly consistent. All GPS-collared deer captured in the control area migrated along a distinct movement corridor located at the base of the Wind River Range. Deer followed a well-defined route that rarely exceeded 1-mi in width and covered a distance of 50 mi. GPS data collected from individual deer for consecutive winters showed a strong affinity for this migration corridor during both spring and fall migrations. Deer that winter in the Pinedale Front were known to migrate northerly along the Wind River Range to the New Fork Lake area, before shifting their migration in a westerly direction towards the Hoback Basin and adjacent mountain ranges (Sawyer and Lindzey 2001). However, details of this migration route, in terms of size, width, specific location, and deer fidelity were unknown prior to GPS data collected over the last two years. Although these deer may migrate 100 mi between winter and summer ranges (Sawyer and Lindzey 2001), our GPS collars do not collect locations May – October, and therefore do not record the entire migration route(s). Deer management in the Sublette DAU is complicated by the long-distance migrations that occur through a variety of habitats and across a mix of land ownership. Knowledge of this migration route should provide agencies with the necessary information to maintain deer movements through the Pinedale Front, adjust harvest strategies accordingly, and prioritize habitat enhancement projects.

Because several thousand mule deer rely on this migration corridor to access their seasonal ranges, maintenance of the corridor should be a priority for agencies and conservation groups alike.

In addition to basic distribution and movement maps, GPS data can be used to conduct more rigorous scientific analyses, such as estimation of resource selection models (Manly et al. 2002) and site-specific analyses for industry (TRC Mariah 2003, Sawyer et al. 2004), agencies, or NGOs regarding permitting, stipulations, or other wildlife-related issues. Resource selection, as described by Manly et al. (2002), is a rapidly advancing methodology for analyzing, modeling, and interpreting wildlife field studies. Resource selection analyses have broad applications, and in the case of this study, were used to determine how mule deer use their habitats in relation to various habitat features, including well pads and road networks associated with energy development. Our basic approach to resource selection treats the GPS-collared deer as the experimental unit and estimates a population-level model (i.e., RSPF) so inference can be made to the entire population of mule deer on the Mesa. Sample size, in this case the number of collared mule deer, is an important consideration for statistical procedures that rely on simple random sampling to obtain population-level inference.

Numerous authors have calculated minimum sample sizes for home range analysis, survival estimates, and resource selection, using both simulations (Allredge and Ratti 1986) and empirical data (Powell et al. 2000). We recognize the number of marked animals in our analysis was smaller than the sample size of 25 recommended by Powell et al. (2000). However, we believe our sample adequately represents the Mesa deer population because of our *a priori* knowledge of deer movement and distribution patterns. Sawyer and Lindsey (2001) studied the movement and distribution patterns of this mule deer population for three years (1998-2000) prior to gas field development. Their results indicated that most deer congregate in the northern portion of the study area during early winter, before moving on to their respective winter ranges. Our sampling design incorporated this knowledge, in that we assumed obtaining a random sample was most likely to occur when deer were congregated on the north end. And, in fact when we compared the distribution (i.e., spatial extent) of 2,563 locations from 82 radio-collared deer for a 3-year period (1998-2000), to the distribution of 18,711 locations from 10 GPS-collared deer for a 1-year time period, the spatial extents were clearly similar (H. Sawyer, WEST, Inc., unpublished data). Thus, while our sample sizes may be less than those suggested by Powell et al. (2000), we believe our sampling strategy and model-building process adequately represents the PAPA deer population. Larger sample sizes should reveal the same relationships between the probability of mule deer habitat use and environmental conditions, but with higher precision.

Recent advances in GPS technology allow for the collection of thousands of locations per animal and thereby make it necessary to reassess the performance of resource selection methods in relation to the number of animals versus the number of observations per animal (Leban et al. 2001). When a large number (e.g., >1,000) of animal locations evenly spread over a study period are available, as is often the case with GPS-collar studies, the utilization distribution (UD) will provide an accurate estimate of the probability of use of habitat units for a particular animal. Accurate estimation of the UD is most likely to occur when precise, frequent, and unbiased animal locations are collected from a representative sample of study animals. Given these requirements, our GPS data appears well-suited for UD estimation, as they lack the inherent bias often associated with traditional telemetry methods because they are systematically collected, irrespective of human error, poor weather conditions, or time of day. Forest density and canopy cover may reduce the fix rates and positional accuracy of GPS collars (Rempel et al. 1995, Moen et al. 1996, D'Eon 2002, Di Orio 2003). However, the open shrublands and gentle topography of the PAPA allow for optimal satellite coverage and GPS performance. Fix-rate GPS bias associated with different habitat types has not been an issue in this study, as evidenced by high (99%) fix rate success and precision (80% 3-D locations). And, since the discontinuation of selective availability in May 2000, 3-D GPS locations typically have less than 20 m error (Di Orio 2003).

APPENDIX B: Resource Selection Modeling Procedures

We fit the same RSPF to each of the n individuals from each winter and estimated population-level RSPF coefficients by

$$\hat{\beta}_i = \frac{1}{n} \sum_{j=1}^n \hat{\beta}_{ij}, \quad (3)$$

where $\hat{\beta}_{ij}$ is the estimate of coefficient i for individual j . The variance of the estimated population-level RSPF coefficients was estimated by

$$\text{var}(\hat{\beta}_i) = \frac{1}{n-1} \sum_{j=1}^n (\hat{\beta}_{ij} - \hat{\beta}_i)^2. \quad (4)$$

A stepwise model building procedure (Neter et al. 1996) was used to estimate population-level models for winters 2000-01, 2001-02, and 2002-03. Significance levels for determining variable entry ($\alpha = 0.15$) and exit ($\alpha = 0.2$) from the RSPF were based on Hosmer and Lemshow (2000). The stepwise model building involved seven steps:

- 1) Each variable (X_i) was used to fit a simple UD model containing only that variable for each deer (n_i) within a winter.
- 2) The average and standard error of the n_i estimated coefficients were calculated for each variable.
- 3) A student's t-test for the statistical significance of each variable determined which variable entered the model first. The variable with the lowest p-value, provided it was ≤ 0.15 , was the first to enter the model.
- 4) The new probability model was fit to data from each deer; each model contained the most significant variable identified in step (3) and each of the other variables previously selected. Again, all remaining variables were considered.
- 5) The average and standard error of the n_i estimated coefficients were computed for those variables not identified in step (3) as most significant. A student's t-test for the statistical significance of those variables determined the next variable to enter the model. Again, the variable with the lowest p-value less than or equal to $\alpha = 0.15$ was chosen.
- 6) This new model containing two predictor variables was fit for each deer and the statistical significance of the variable chosen in step (3) was re-assessed. If the p-value from the t-test for this variable still met the criteria to be in the model, it was kept. If the p-value changed and exceeded the $\alpha = 0.2$ cutoff, it was dropped from the model.
- 7) This process was repeated, and variables were added or dropped from the model until no further variables entered or exited the model.

Our analysis correctly identified the deer as the experimental unit, and because it was necessary to have the same UD model (i.e., same predictor variables) for each deer within a winter, we used a stepwise (Neter et al. 1996) model building process to produce both individual and population level models for each winter, as recommended by Erickson et al. (2001). It is not possible to use Akaike's Information Criterion (AIC) and similar alternatives such as Akaike's Information Criterion second order variant (AICc) and Bayesian Information Criterion (BIC) (Burnham and Anderson 2002) for model selection in this context, since this could result in a different model for each deer within a winter and prevent estimation of population level models.

Stepwise model building was conducted for all winters except pre-development. Prior to development of the PAPA, the deer monitoring consisted of many VHF-collared deer with relatively few locations (Sawyer and Lindzey 2001), compared to the GPS collars used from 2000 to present. Because of the limited number of locations recorded for the VHF-collared deer we could not fit individual RSPF's to these deer. Therefore, we pooled data across deer ($n=76$) and winters (1998-99, 1999-00) to

APPENDIX B: Resource Selection Modeling Procedures

estimate a population-level model (RSPF) for the pre-development time period. Simple random samples of 30 locations were taken from deer with > 30 locations to ensure approximately equal weight was given to each deer in the analysis. All variables except *aspect* were included in the population level model for pre-development.

Predictions (i.e., probability of mule deer habitat use) of population-level models were mapped for each winter on 104 x 104 m grids that covered the study area. Predictions were first scaled so that the total probability corresponding to all grid cells in the study area totaled 1.0. To aid in mapping the estimated probability of use, each grid cell was assigned a value of 1 to 4. Grid cells with the largest 25% of values for predicted probability of use were assigned 1 (= high use), cells in the 51 to 75 percentiles were assigned 2 (= medium high use), cells in the 26 to 50 percentiles were assigned 3 (= medium low use), and cells in the 0 to 25 percentiles were assigned 4 (= low use).

APPENDIX C: GPS locations and 60-mile migration route of deer #863.

BLM Library
Bldg. 50
Denver Federal Center
P.O. Box 25047
Denver, Colorado 80225