312066 0271 2729 8

Digitized by the Internet Archive
in 2013 with funding from
Boston Library Consortium Member Libraries

JOB PROGRESS REPORT

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-13 Project Title: Game Population Trend and Harvest Survey

Job No.: I-1 Job Title: Statewide Small Game Harvest

Period Covered: 1 January to 31 December 1971

Summary: A usable return of 923 cards (68.5%) was obtained from three mailings of the postal questionnaire to 1400 sportsmen. Each return was calculated to represent 124.31 hunters.

Approximately 81 percent of the 1969 hunters purchased a license in 1970. Also, about 73 percent of these licensees hunted small game, of whom 433 (58.0%) were successful in taking at least one unit of game.

Hunter success was greatest for cottontail rabbit (147,026), ducks (131,752), and pheasant (94,664). Nine other species had lesser rates of harvest.

Hunter effort increased for four species and decreased for five. Changes were most noticeable regarding ruffed grouse, ducks, raccoon, and woodcock.

Private land continues to be important to the Massachusetts sportsman; however, there is an apparent increase in the use of management areas. More hunters are now reporting the use of both categories of land, rather than using solely private land.

Background: This job was initiated in 1959 as a means of determining trends in the harvest of selected small game species. The survey was conducted yearly until 1961, and every other year thereafter. Variability in sample size, sample populations, and data analysis preclude direct comparison between certain years; however, harvest trends can be determined. Applicable comparisons between data from the current segment and that from past years are presented in the following tables.

Objectives: To determine the statewide harvest of selected small game species and to determine the characteristics of land utilization and time expenditure by sportsmen.

Procedures:

A sample was randomly drawn from each of 37 boxes containing the 1969 sportsman license cards. This aggregation was run through an IBM 83 card sorter to eliminate cards from fishing and trapping licensees. The remaining cards were hand sorted to separate non-resident and free licenses from the resident hunting and sporting licenses. The first legible 300 cards in the resident sporting category and the first legible 600 cards in the resident hunting category were then chosen as the hunter sample for this survey.

The names and addresses of the sportsmen in the sample were printed on gummed labels by the printout device on an IBM computer. Four labels were produced for each individual. A code number was assigned to each name to facilitate future identification.

The questionnaire (Figures I-A and I-B) was printed on double prestamped postal cards. The format was similar to that of previous years except for changes in the species list and special questions. Three mailings were conducted, on 19 July, 19 August, and 24 September. The delay in the initial mailing was due to a late return of the address labels from the printers and a rise in the postal rates.

As each card was returned, the respondent was removed from the mailing list utilizing the code number. Game harvest data was tabulated separately for each mailing's return, and subsequently combined to provide a total harvest estimate. Responses to the special questions were tabulated only for the entire sample return.

Each usable return was considered to represent a specific proportion of all resident licensees. This weighting (one return represents 124.31 individuals) was determined by summing the resident hunting and resident sporting sales for the calendar year 1970 (114,735) and dividing this figure by the total usable return (923).

Findings:

Questionnaires were mailed to 1400 resident licensees; 1347 were delivered and there was a usable return of 923 or 68.5 percent (Table 1).

As the sample had been drawn from the previous year's (1969) licenses, the 923 usable returns were first divided into two groups: (1) those hunters licensed in 1969 who subsequently purchased a 1970 license; and, (2) those hunters from 1969 who did not purchase a 1970 license. Approximately 81 percent (80.82) of the 1969 licensees did purchase a license in 1970 (Table 2). No attempt was made to determine why the remaining 19 percent did not buy a license.

The 746 1970 licensed respondees were further broken down into four categories: successful hunters (a hunter who took at least one unit of game), unsuccessful hunters, licensees who did not hunt, and licensees who hunted only deer. Reported hunter success increased over 4.5 percent

FIGURE 1-A: POSTAL QUESTIONNAIRE,
1970 GAME KILL REPORT: FORM A (OVERSEAS)

Dear Hunter:

This is the statewide game kill questionnaire. It is the only method we have of compiling the annual kill of our small game species. Your cooperation will be appreciated in filling out this postcard promptly.

1. If you did not buy a license in 1970, check the top block of the return card. Do not complete other questions.
2. If you bought a license, but did not hunt, check the second block at the top of the return card.
3. Check all game hunted, even if you killed nothing.
4. Under No. Bagged, list only game killed by you.
5. Under Towns, list the towns you hunt in the most.
6. If you hunted only deer, answer only the last question. Indicate whether or not you applied for an antlerless permit.
7. If you hunt on Wildlife Management Areas, list the area you hunt the most.

All replies are confidential. When the figures are compiled, a report will be published.

Thank you.

Form FG-169 11-70-048284

MASSACHUSETTS DIVISION OF FISHERIES AND GAME

DIVISION OF FISHERIES & GAME

FIELD HEADQUARTERS

WESTBORD, MASS. 01581

FIGURE 1-B: POSTAL QUESTIONNAIRE,
1970 GAME KILL REPORT: FORM A (REVERSE)

Form A

1970 GAME KILL REPORT

- ☐ Check if you did not buy a hunting or sporting license in 1970. Do not complete any other questions.
- ☐ Check if you bought a license, but did not hunt in 1970.

Please check all game hunted in Massachusetts	✓	Number Bagged	Town or Towns you Hunted in the Most
Pheasant			
Ruffed Grouse			
Quail			
Woodcock			
White Hare			
Cottontail Rabbit			
Raccoon			
Gray Squirrel			
Fox			
Ducks (All)			
Geese			
BOBCAT			

Please estimate - (Record "None" as "0".)

- The number of times you hunted on: Wildlife Management Areas_____ Private Land_____.
- The number of times you hunted pheasant on: Wildlife Management Areas_____ Private Land_____.
- If you hunt on Wildlife Management Areas, list the area you hunt on the most_____.
- Did you hunt deer in 1970? Yes_____ No_____.
If yes did you apply for an antlerless permit?
Yes_____ No_____.

Table 1. Relative Return by Type of License.

<u>Mailing</u>	<u>License Type</u>	<u>Number Mailed</u>	<u>Number Undelivered</u>	<u>Number Delivered</u>	<u>Number Returned Usable</u>	<u>Percent of Delivered</u>
First	Hunting	600	7	593	206	34.74
	Sporting	800	16	784	322	41.07
Second	Hunting	386	10	376	122	32.45
	Sporting	460	10	450	141	31.33
Third	Hunting	253	7	246	56	22.76
	Sporting	309	3	306	76	24.84
Combined Return	Hunting	600	24	576	384	66.67
	Sporting	800	29	771	539	69.91
Totals		1400	53	1347	923	68.52

Table 2. Percent of 1969 Sample that Purchased a License in 1970.

	<u>First Mailing</u>		<u>Second Mailing</u>		<u>Third Mailing</u>		<u>Totals</u>	
	<u>No.</u>	<u>Percent</u>	<u>No.</u>	<u>Percent</u>	<u>No.</u>	<u>Percent</u>	<u>No.</u>	<u>Percent</u>
1969 License Sample that Purchased 1970 License	439	83.1	204	77.57	103	78.03	746	80.82
1969 License Sample that Did Not Purchase 1970 License	89	16.9	59	22.43	29	21.97	177	19.18
Totals	528	100.0	263	100.00	132	100.00	923	100.00

from 1968, and the amount of non-hunters decreased two percent (Table 3). The other two categories remained nearly stable.

The mean take and sample size for each game category is presented in Table 4. The mean take per hunter was greatest for ducks (5.17 ducks per hunter), followed by cottontail rabbit (4.15), and gray squirrel (3.60). The mean take for all other species was less than three units of game per hunter.

The mean take and expanded take for 1970 are presented in Table 5 and compared with previous years in Table 6. Readers are reminded that the validity of these estimates cannot be determined statistically, as the data plots as a negative binomial. The small sample size for certain species (e.g., bobcat) increases the chance of error in expansion, and confidence limits, if possible to compute, would likely be very broad. Yearly comparisons should be made with caution, as sample sizes have varied considerably.

The 1970 expanded bag take was greatest for cottontail rabbit (147,026), followed by ducks (131,752), pheasant (94,664), woodcock (67,886), gray squirrel (65,336), and ruffed grouse (57,482). The take for the remaining six species was less than 50,000 units each.

The number of hunters seeking each species had increased for four species and decreased for five, considering the nine comparable categories for 1968 and 1970. Changes are most dramatic respecting ruffed grouse (down 16,300 hunters), ducks (down 10,000 hunters), raccoon (down 6500 hunters), and woodcock (up 6200 hunters). These fluctuations are not necessarily due to an increase or decrease in the number of actual hunters, but may be in part due to a diversification of effort on the part of previously reporting hunters. The format of the questionnaire is such that if a hunter kills a certain type of game, he is considered to have hunted that species regardless of the species for which he originally intended to hunt.

The mean harvest of three game categories (bobwhite quail, woodcock, and ducks) has increased considerable since 1968. This could be due to increased hunting pressure, increased reporting rate by respondees, an increase in game availability or a combination of two or all of these factors. Slight increases were noted for pheasants, ruffed grouse, and white hare, and minimal decreases were reported for cottontail rabbit, gray squirrel, and raccoon. Fox, bobcat and geese were not sampled in 1968.

Table 3. Initial Breakdown of 1970 Licensed Respondees.

<u>Groups</u>	<u>First Mailing</u>		<u>Second Mailing</u>		<u>Third Mailing</u>		<u>Totals</u>	<u>Percent of Totals</u>
	<u>Number</u>	<u>Percent</u>	<u>Number</u>	<u>Percent</u>	<u>Number</u>	<u>Percent</u>		
Successful Hunters	268	61.1	120	58.8	45	43.7	433	58.0
Unsuccessful Hunters	73	16.6	21	10.3	19	18.4	113	15.1
Did Not Hunt	62	14.1	53	26.0	26	25.2	141	18.9
Hunted Deer Only	36	8.2	10	4.9	13	12.7	59	8.0
Totals	439	100.0	204	100.0	103	100.0	746	100.0

Table 4. Comparison of Hunter Sample and Mean Take for Three Mailings, 1970.

<u>Species</u>	<u>First Mailing</u>		<u>Second Mailing</u>		<u>Third Mailing</u>		<u>Total</u>	
	<u>Hunter Sample</u>	<u>Mean Take</u>	<u>Hunter Sample</u>	<u>Mean Take</u>	<u>Hunter Sample</u>	<u>Mean Take</u>	<u>Hunter Sample</u>	<u>Mean Take</u>
Pheasant	281	1.51	117	2.13	58	1.50	456	1.67
Ruffed Grouse	197	1.42	63	2.10	29	1.79	289	1.60
Quail	73	2.40	30	2.30	11	1.82	114	2.32
Woodcock	146	2.55	50	2.82	19	1.58	215	2.54
Cottontail Rabbit	187	4.02	64	4.67	34	3.88	285	4.15
White Hare	79	2.35	22	3.54	12	2.33	113	2.58
Gray Squirrel	93	3.83	33	3.67	20	2.45	146	3.60
Raccoon	28	2.11	9	2.56	5	0.80	42	2.05
Fox	30	0.47	7	0.71	7	0.43	44	0.50
Bobcat	13	0.15	6	0.33	3	0.00	22	0.18
Ducks (all)	120	4.46	59	6.90	26	4.54	205	5.17
Geese	64	0.61	22	1.04	13	0.85	99	0.74

Table 5. Game Bag Expansion Describing the Take of the 1970 Hunter Population.

<u>Species</u>	<u>Number of Hunters in Sample</u>	<u>Number of Hunters Expanded*</u>	<u>Reported Take</u>	<u>Mean Bag Take</u>	<u>Expanded Bag Take</u>
Pheasant	456	56,685	761	1.67	94,664
Ruffed Grouse	289	35,926	464	1.60	57,482
Quail	114	14,171	264	2.32	32,877
Woodcock	215	26,727	547	2.54	67,886
Cottontail Rabbit	285	35,428	1182	4.15	147,026
White Hare	113	14,047	292	2.58	36,241
Gray Squirrel	146	18,149	526	3.60	65,336
Raccoon	42	5,221	86	2.05	10,703
Fox	44	5,470	22	0.50	2,735
Bobcat	22	2,735	4	0.18	492
Ducks (all)	205	25,484	1060	5.17	131,752
Geese	99	12,307	73	0.74	9,107

* Each return equals 124.31 hunters.

Table 6. Comparison of Hunter Sample, Mean Kill, and Expanded Kill for 1953, 1960, 1962, 1964, 1968 and 1970.

<u>Species</u>	<u>1958</u>	<u>1960</u>	<u>1962</u>	<u>1964</u>	<u>1966</u>	<u>1968</u>	<u>1970</u>
<u>Expanded Number of Hunters</u>							
Pheasant	43,252	46,205	56,546	54,072	50,151	55,148	56,685
Ruffed Grouse	30,002	35,611	43,904	46,548	39,626	52,191	35,926
Quail	6,336	5,948	6,860	8,628	9,906	13,158	14,171
Woodcock	9,937	9,860	16,268	19,161	18,203	20,551	26,727
Cottontail Rabbit	35,571	32,433	36,554	34,008	31,948	33,857	35,428
White Hare	14,737	12,875	17,248	17,456	14,240	15,967	14,047
Gray Squirrel	20,018	18,254	17,150	16,753	21,423	19,811	18,149
Raccoon	3,792	4,645	4,018	6,220	6,439	11,680	5,221
Fox*							5,470
Bobcat*							2,735
Ducks (all)**	21,458	17,194	27,244	27,788	27,614	35,483	25,484
Geese*							12,307
<u>Mean Kill</u>							
Pheasant	1.451	1.635	1.501	1.539	1.760	1.659	1.67
Ruffed Grouse	1.158	1.540	1.458	1.732	1.900	1.597	1.60
Quail	2.083	2.343	1.886	1.825	1.925	1.235	2.32
Woodcock	1.367	1.355	1.783	1.863	2.170	1.568	2.54
Cottontail Rabbit	5.224	4.151	3.920	3.979	4.554	4.414	4.15
White Hare	2.244	1.418	1.580	2.287	2.896	2.537	2.58
Gray Squirrel	4.014	3.790	2.543	3.389	4.127	4.671	3.60
Raccoon	2.671	4.246	4.585	3.967	4.423	2.135	2.05
Fox*							0.50
Bobcat*							0.18
Ducks (all)**	3.88	2.82	2.62	3.46	2.92	2.67	5.17
Geese*							0.74
<u>Expanded Kill</u>							
Pheasant	62,759	77,541	84,864	83,260	88,290	91,518	94,664
Ruffed Grouse	34,742	54,841	63,990	80,654	75,289	83,385	57,482
Quail	13,198	13,933	12,936	15,750	19,069	16,261	32,877
Woodcock	13,584	13,364	29,007	35,712	39,502	32,230	67,886
Cottontail Rabbit	185,823	134,623	143,273	135,328	145,500	149,471	147,026
White Hare	33,070	18,253	27,243	39,927	41,235	40,508	36,241
Gray Squirrel	80,352	69,186	43,609	56,779	88,417	92,549	65,336
Raccoon	10,128	19,721	18,424	24,679	28,480	24,937	10,703
Fox*							2,735
Bobcat*							492
Ducks (all)**	83,290	48,404	71,440	96,204	80,735	93,880	131,752
Geese*							9,107

* Sampled for the first time in 1970.

** Aggregated data. Previous reports separated black ducks and "other ducks".

The 1970 hunter success for twelve game species is shown in Table 7. Hunter success was greatest for ducks, with nearly three-quarters (73.17%) of the sampled hunters reportedly taking at least one duck. Good success was also enjoyed by hunters of cottontail rabbit (72.63% successful), and gray squirrel (71.92% successful). Hunters least successful were those seeking geese (34.34%) and bobcat (18.18%).

The 1970 hunter success percentages for nine game species are compared with those from previous years in Table 8. Hunter success has increased for six species since 1968 and decreased on the remaining three. The greatest increase was for ducks (up seven percent), and the greatest decrease for raccoon (down 9.8 percent).

Table 9 presents twelve game species ranked by hunter success and hunting pressure for the sample periods from 1958 to 1970. Hunting pressure has remained fairly constant for most species during this time. One notable exception appears to be woodcock which has experienced an increase in pressure from seventh place in 1958 to fourth place in 1970. Fluctuations in hunter success are more evident. These variations may be due to inconsistencies in the reporting rate among the sampled sportsmen, or to actual biological fluctuations in native game populations, or both.

The first special question asked the sportsman to categorize the ownership of the land on which he hunted. Results of this categorization are presented in Table 10. Although the Massachusetts sportsman is still heavily dependent on private land (47.0% utilized only private land in 1970), there is an evident downward trend in this attitude. Hunters are now broadening their approach and hunting on both private land and management areas. Nearly 42 percent hunted both types of land in 1970, as opposed to 33.6 percent doing so in 1968. Decreases in the amount of huntable private land (through posting and development) and continued acquisition of management area lands and resultant publicity may contribute to this change in land use patterns.

This same pattern is evident when pheasant hunters are treated separately (Table 11); however, variations in the number of stocked private covers may influence the distribution in this subcategory.

Land utilization by small game hunters is presented by county in Table 12. An important factor affecting a sportsman's land use habits is his proximity to a management area. If he hunts only in his home county, and there are no management areas in that county, quite obviously private land is very important to him. Perhaps, then, a truer picture can be obtained by examining those who reported hunting more than one county. More than half (53.6%)

Table 7. Hunter Success for Twelve Game Species, 1970.

<u>Species</u>	<u>Total Number of Hunters</u>	<u>Successful Hunters</u>	<u>Unsuccessful Hunters</u>	<u>Percent Successful</u>
Ducks (all)*	205	150	55	73.17
Cottontail Rabbit	285	207	78	72.63
Gray Squirrel	146	105	41	71.92
Pheasant	456	279	177	61.18
White Hare	113	69	44	61.06
Woodcock	215	128	87	59.53
Ruffed Grouse	289	164	125	56.75
Raccoon	42	22	20	52.38
Quail	114	47	67	41.23
Fox	44	18	26	40.91
Geese	99	34	65	34.34
Bobcat	22	4	18	18.18

* See Footnote 2, Table 6

Table 8. Hunter Success for Nine Game Species: 1958, 1960, 1962, 1964, 1968 and 1970.

<u>Species</u>	<u>1958</u>	<u>1960</u>	<u>1962</u>	<u>1964</u>	<u>1966</u>	<u>1968</u>	<u>1970</u>
Ducks (all)*	64.2%	61.1%	53.6%	74.0%	65.5%	66.2%	73.2%
Cottontail Rabbit	78.0%	74.1%	69.4%	69.9%	69.4%	70.7%	72.6%
Gray Squirrel	79.8%	78.1%	61.7%	79.6%	71.7%	76.1%	71.9%
Pheasant	54.2%	57.3%	57.9%	57.0%	58.3%	57.6%	61.2%
White Hare	58.3%	44.9%	50.6%	56.9%	56.5%	56.5%	61.1%
Woodcock	54.1%	48.8%	51.2%	53.4%	64.5%	59.7%	59.5%
Ruffed Grouse	49.4%	54.7%	52.7%	56.9%	65.3%	50.4%	56.8%
Raccoon	51.9%	73.7%	75.6%	40.3%	55.8%	62.2%	52.4%
Quail	34.8%	52.1%	47.1%	44.2%	66.2%	39.3%	41.2%

* See Footnote 2, Table 6

Table 9. Twelve Game Species Ranked by Hunting Pressure and Hunter Success: 1958, 1960, 1962, 1964, 1966, 1968 and 1970

Species	Ranked by Hunting Pressure							Ranked by Hunter Success						
	1953	1960	1962	1964	1966	1968	1970	1958	1960	1962	1964	1966	1968	1970
Pheasant	1	1	1	1	1	1	1	4	5	4	4	7	6	4
Ruffed Grouse	3	2	2	2	2	2	2	5	6	6	5*	5	8	7
Quail	8	8	8	8	8	8	7	8	7	9	7	3	9	9
Woodcock	7	7	7	5	6	5	4	7	8	7	6	6	5	6
Cottontail Rabbit	2	3	3	3	3	4	3	1	2	2	3	2	2	2
White Hare	6	6	5	6	7	7	8	6	9	8	5*	8	7	5
Gray Squirrel	5	4	6	7	5	6	6	3	1	3	1	1	1	3
Raccoon	9	9	9	9	9	9	11	9	3	1	8	9	4	8
Fox**							10							10
Bobcat**							12							12
Ducks (all)***	4	5	4	4	4	3	5	2	4	5	2	4	3	1
Geese**							9							11

* Equal Hunter Success

** See Footnote 1, Table 6

*** See Footnote 2, Table 6

Table 10. Summary of Areas Hunted: 1959, 1960, 1962, 1964, 1966, 1968 and 1970.

<u>Area</u>	<u>Percent of Small Game Hunters Using Stipulated Area</u>						
	<u>1959</u>	<u>1960</u>	<u>1962</u>	<u>1964</u>	<u>1966</u>	<u>1968</u>	<u>1970</u>
Management Areas	6.0	4.4	8.8	6.3	9.7	12.4	11.4
Private Land	65.0	59.4	69.5	58.8	56.5	54.0	47.0
Both Types	29.0	36.2	21.7	34.9	33.8	33.6	41.6

Table 11: Land Utilization by Small Game Hunters: 1966, 1968 and 1970.

<u>Hunter Category</u>	Percent of Hunters Using								
	<u>Management Areas</u>			<u>Private Land</u>			<u>Both</u>		
	<u>1966</u>	<u>1968</u>	<u>1970</u>	<u>1966</u>	<u>1968</u>	<u>1970</u>	<u>1966</u>	<u>1970</u>	
Pheasant Hunters	17.7	15.3	20.2	50.5	55.4	46.3	31.7	29.2	33.5
All Hunters	9.7	12.4	11.4	56.5	54.0	47.0	33.8	33.6	41.6

Table 12: Land Utilization of Small Game Hunters by County, 1970

County	Management Areas		Private Land		Both		Total Number Hunters	Percent of Statewide Hunter Total
	Number Hunters	Percent	Number Hunters	Percent	Number Hunters	Percent		
Barnstable	5	29.4	7	41.2	5	29.4	17	4.0
Berkshire	3	13.1	13	56.5	7	30.4	23	5.5
Bristol	1	3.8	18	69.2	7	27.0	26	6.2
Dukes	0	0.0	1	100.0	0	0.0	1	0.2
Essex	6	16.2	14	37.8	17	46.0	37	8.8
Franklin	0	0.0	6	100.0	0	0.0	6	1.4
Hampden	1	4.5	13	59.1	8	36.4	22	5.2
Hampshire	2	8.3	19	79.2	3	12.5	24	5.7
Middlesex	0	0.0	17	58.6	12	41.4	29	6.9
Nantucket	0	0.0	3	100.0	0	0.0	3	0.7
Norfolk	0	0.0	8	88.9	1	11.1	9	2.1
Plymouth	7	18.9	11	29.7	19	51.4	37	8.9
Worcester	15	20.0	24	32.0	36	48.0	75	17.8
Hunted more than one county	8	7.1	44	39.3	60	53.6	112	26.6
Statewide Totals	48	11.4	198	47.0	175	41.6	421	100.0

of the 112 reporting hunters utilized both categories of land, 39.3 percent utilized only private land, and only 7.1 percent utilized only management areas. This method of presentation shows a greater degree of dependence on the "both" category. County-wide breakdowns were not made in previous years so comparisons cannot be made. Data from this segment, however, does indicate an increased awareness of wildlife management areas.

If land utilization data are presented as the number of trips to each land category, per county, by small game hunters (Table 13), it is evident that private land draws the greatest number of hunter trips. For statewide totals, 223 hunters reported 1369 trips to management areas, a mean of 6.1 trips per hunter, whereas, for private land, 369 hunters reported 4473 trips, a mean of 12.1 trips per hunter. The mean number of trips per hunter, regardless of land ownership, was 9.9, a slight decrease from 1968 (10.8 trips per hunter).

Land utilization patterns were also determined for pheasant hunters, and are presented in Table 14. Again, the greatest percentage of those sportsmen hunting in more than one county utilized both private land and management areas (46.0%). This is, however, a lesser figure than for small game hunters as a whole (53.6%). This may possibly be because of the diversity in habitat of the various small game species, requiring the general hunter to cover more area. The pheasant hunter may have his favorite covers, whether on management areas or private land, which he hunts faithfully. Familiarity with stocking schedules could also cause a hunter to prefer a certain location, as this knowledge may increase his chance of success.

Table 15 presents the number of trips by pheasant hunters, by county and land ownership. Less disparity is noted in these totals than in those for general small game hunters. For pheasant hunters, there was a mean of 5.5 trips per hunter on management areas, and 7.6 trips per hunter to private land. The statewide total for all pheasant hunters was 6.7 trips per hunter.

A third special question concerned hunters who utilized state wildlife management areas. These sportsmen were asked to list the one area they used the most. Replies (Table 16) were received from 242 hunters, of whom 41 gave an unidentifiable answer, or listed a non-state area. (A response indicating the town in which a state area was located was considered equivalent to listing the actual name of the area.) The remaining 201 hunters indicated using 28 state wildlife management areas. The greatest number of responses was received for Myles Standish (20), Knightville (18), and Birch Hill (15), while only one hunter responded for each of seven other areas. Visual examination of the list shows that the areas with the greatest number of responses are generally the larger, more well

 

Table 13. Number of Trips by Small Game Hunters Per County by Land Ownership: 1970

County	Management Areas			Private Land			County Totals	
	Hunter Sample	Number Trips	Trips/ Hunter	Hunter Sample	Number Trips	Trips/ Hunter	Number Trips	Trips/ Hunter
Barnstable	11	69	6.3	11	112	10.2	181	8.2
Berkshire	10	112	11.2	19	211	11.1	323	11.1
Bristol	9	73	8.1	25	294	11.8	367	10.8
Dukes	0	-	-	1	20	20.0	20	20.0
Essex	23	130	5.6	31	373	12.0	503	9.3
Franklin	0	-	-	5*	28	4.7	28	4.7
Hampden	9	26	2.9	21	189	9.0	215	7.2
Hampshire	5	20	4.0	22	200	9.1	220	8.1
Middlesex	12	40	3.3	29	276	9.5	316	7.7
Nantucket	0	-	-	3	55	18.3	55	18.3
Norfolk	1	1	1.0	9	80	8.9	81	8.1
Plymouth	25	138	5.5	30	281	9.4	419	7.6
Worcester	51	432	8.5	60	748	12.5	1180	10.6
Hunted more than one county	67	328	4.9	103	1606	15.6	1934	11.4
Statewide Totals	223	1369	6.1	369	4473	12.1	5842	9.9

* One hunter reporting 200 trips not included.

Table 14. Land Utilization by Pheasant Hunters by County: 1970.

<u>County</u>	<u>Management Areas</u>		<u>Private Land</u>		<u>Both</u>		<u>Total Number Hunters</u>	<u>Percent of Statewide Hunter Total</u>
	<u>Number Hunters</u>	<u>Percent</u>	<u>Number Hunters</u>	<u>Percent</u>	<u>Number Hunters</u>	<u>Percent</u>		
Barnstable	6	42.9	5	35.7	3	21.4	14	3.7
Berkshire	7	36.8	9	47.4	3	15.8	19	5.1
Bristol	2	8.3	16	66.7	6	25.0	24	6.4
Dukes	0	0.0	2	100.0	0	0.0	2	0.5
Essex	9	27.3	9	27.3	15	45.6	33	8.8
Franklin	0	0.0	3	100.0	0	0.0	3	0.8
Hampden	3	15.0	12	60.0	5	25.0	20	5.3
Hampshire	2	8.7	18	78.3	3	13.0	23	6.1
Middlesex	0	0.0	18	64.3	10	35.7	28	7.4
Nantucket	0	0.0	3	100.0	0	0.0	3	0.8
Norfolk	0	0.0	9	100.0	0	0.0	9	2.4
Plymouth	9	33.3+	9	33.3+	9	33.3+	27	7.2
Worcester	25	35.2	20	28.2	26	36.6	71	18.9
Hunted more than one county	13	13.0	41	41.0	46	46.0	100	26.6
Statewide Totals	76	20.2	174	46.3	126	33.5	376	100.0

Table 15. Number of Trips by Pheasant Hunters, Per County, by Land Ownership: 1970.

County	Management Areas			Private Land			County Totals		
	Hunter Sample	Number Trips	Trips/Hunter	Hunter Sample	Number Trips	Trips/Hunter	Hunter Sample	Number Trips	Trips/Hunter
Barnstable	9	52	5.7	8	47	5.9	17	99	5.8
Berkshire	10	91	9.1	12	65	5.4	22	156	7.1
Bristol	7	31	4.4	22	147	6.7	29	178	6.1
Dukes	0	-	-	2	11	5.5	2	11	5.5
Essex	24	119	5.0	24	240	10.0	48	359	7.5
Franklin	0	-	-	3	36	12.0	3	3	12.0
Hampden	8	21	2.6	17	97	5.7	25	118	4.7
Hampshire	5	38	7.6	21	149	7.1	26	187	7.2
Middlesex	10	32	3.2	28	222	7.9	38	254	6.9
Nantucket	0	-	-	3	25	8.3	3	25	8.3
Norfolk	0	-	-	9	54	6.0	9	54	6.0
Plymouth	18	103	5.7	17	92	5.4	35	195	5.8
Worcester	50	387	7.7	46	367	8.0	96	754	7.8
Hunted more than one county	59	233	3.9	86	729	8.5	145	962	6.6
Statewide Totals	200	1107	5.5	298	2281	7.6	498	3355	6.7

Table 16. Wildlife Management Areas Utilized by Small Game Hunters, 1970.

<u>Area</u>	<u>No. Hunters Indicating Use</u>	<u>Percent of Total Using Management Areas</u>
Barre Falls	14	5.8
Birch Hill	15	6.2
Canada Hill	1	0.4
Crane	13	5.4
Crane Pond	10	4.1
Fort Devens	3	1.2
Four Chimneys	1	0.4
Freetown	9	3.8
Harold Parker Forest	7	2.9
Hodges Village	3	1.2
Housatonic Valley	1	0.4
Hubbardston	6	2.5
Knightville	18	7.4
Mill Creek	2	0.8
Myles Standish	20	8.3
Northeast	10	4.1
Otis Air Force Base	2	0.8
Pantry Brook	1	0.4
Peru	1	0.4
Quaboag	7	2.9
Stafford Hill	1	0.4
Swift River	8	3.4
Westboro	6	2.5
West Hill	10	4.1
West Meadows	16	6.6
Williamstown	7	2.9
Winimusset	8	3.4
Wompatuck Park	1	0.4
Unidentified	<u>41</u>	<u>16.9</u>
Total	242	100.0

known areas. This could be due to the notoriety of the particular area, and also to the proximity of the hunter's residence to the area in question.

Table 17 compares hunter travel distance, from hometown to area, as obtained by direct questioning at the management area (W-35-R, V-1) to distances obtained by postal questionnaire from this study. The straight line distance between the hunter's town of origin and the center of the management area was used as a standard approximation of travel distance. The mark indicating the location of the town center was used as a common starting point for all hunters residing in that particular town. These distances were plotted in miles on an official Massachusetts road map, and converted to the nearest whole kilometer afterward. The small sample from the postal questionnaire appears to select against those hunters in the farther distance groups, and the extensive variation in sample size between the two methods may preclude any attempts at statistical correlation. Gross examination of the figures, however, indicate that the figures obtained by both methods may represent the actual distribution of participating sportsmen although data for several individual areas are quite divergent. The nearness of the figures from the two samples is seen for the statewide totals with 71.8 percent in the 0-32 km range (postal), and 69.4 percent in that same range (utilization survey).

The final special question related to deer hunting and antlerless deer permit applications. Hunters were asked to indicate whether or not they hunted deer, and to indicate if they did hunt deer, whether or not they applied for an antlerless permit. Responses to this question are summarized in Table 18. Five hundred twenty-five (525) individuals responded to this question, of whom 179 (34.1%) did not hunt deer. An estimate of the number of Massachusetts deer hunters can then be derived by multiplying the 346 deer hunters in the sample times the expansion figure of 124.31 for a total of 43,011 hunters.

Antlerless permits were applied for by 240 (69.4%) of the 346 responding deer hunters. An estimate of the number of permit applicants may be determined by taking 69.4 percent of the 43,011 deer hunters, yielding 29,850 permit applicants. This cannot accurately be compared with the actual number of permit applicants (35,069) as the questionnaire did not include a provision for those individuals who did apply for the permit, but did not subsequently hunt.

Table 17. Comparison of Hunter Travel Distance as Obtained from Small Game Harvest Survey and Hunter Utilization Survey, 1970

Area	<u>Small Game Survey</u>					<u>Hunter Utilization Survey</u>			
	Hunter Sample	<u>Percent Traveling*</u>			Hunter Sample	<u>Percent Traveling</u>			
		<u>0-32</u>	<u>32-80</u>	<u>80+</u>		<u>0-32</u>	<u>32-80</u>	<u>80+</u>	
Barre Falls	14	78.6	21.4	-	503	84.3	12.5	3.2	
Birch Hill	15	93.3	6.7	-	1261	77.3	14.6	8.1	
Crane	13	46.2	38.4	15.4	465	32.0	32.0	36.0	
Crane Pond	10	70.0	30.0	-	261	62.5	37.5	0.0	
Freetown	9	100.0	-	-	261	97.7	2.3	0.0	
Housatonic Valley	1	100.0	-	-	181	98.3	0.6	1.1	
Hubbardston	6	83.3	-	16.7	464	75.0	23.7	1.3	
Knightville	18	72.2	27.8	-	253	75.5	24.5	0.0	
Myles Standish	20	40.0	60.0	-	384	33.6	62.8	3.6	
Northeast	10	60.0	40.0	-	736	35.2	64.7	0.1	
Swift River	8	87.5	-	12.5	462	95.2	3.7	1.1	
West Hill	10	90.0	10.0	-	497	93.6	6.2	0.2	
Winimusset	8	75.0	25.0	-	448	69.4	30.6	0.0	
Totals	142	71.8	25.4	2.8	6176	69.4	25.5	5.1	

* Distances are expressed in kilometers.

0-32 KM = 0-20 miles

32-80 KM = 20-50 miles

80+ KM = 50+ miles

Table 18. Breakdown of Responding Deer Hunters as to Antlerless Permit Applications, 1970

Hunter Category	Number Responding	Percent of Responding Deer Hunters*	Percent of Total Respondents
Hunted Deer - Applied for Permit	240	69.4	45.7
Hunted Deer - Did Not Apply for Permit	103	29.7	19.6
Hunted Deer - Permit Status not Stated	3	0.9	0.6
Did not Hunt Deer	179	---	34.1
Totals	525	100.0	100.0

* Responding deer hunters = 346

Recommendations: 1. An inherent disparity in this job is the negative binomial distribution of the harvest data. This precludes any valid analysis by standard statistical techniques. There are, however, some corrections which can be made to increase the usefulness of this job as an indicator of the harvest trend of small game species. These corrections include:

- a. Draw the sample entirely by machine in order to get as near a random sample as possible.
- b. Have sample drawn and questionnaires prepared at least two months prior to anticipated mailing date. This will allow more time for cooperating agencies to prepare questionnaires, and will lessen any delay at the required mailing date.
- c. Questionnaires should be mailed promptly on 1 March, at the close of the rabbit season, in order to lessen the impact of respondents' memory bias.
- d. Questionnaires should clearly stipulate desired seasonal limits for those species (rabbits, fox) for which seasons carry over from one year to the next.
- e. Special questions should be worded so as to include all potential classifications of response.
- f. Inherent biases in sampling procedure and expansion technique indicate a slight probable gain from sampling species with a low harvest rate. As an example, the estimated harvest of 492 bobcat is unrealistic (trapper returns indicate a lower take), and the sample is so small that an included error of one bobcat could change the expanded harvest total by 127 cats.

2. The next segment of this job should be conducted in 1973 using a hunter sample from the 1972 licensees. Corrections described in (1) above should be implemented.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Assistant Game Biologist

Date _____

MASS-

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Title: Game Population Trend and Harvest Survey

Job No.: II-1 Job Title: Statewide Deer Harvest

Period Covered: 8 November 1971 to 11 December 1971

Summary: Archers harvested 36 deer (26 males and 10 females) during the two-week archery season, November 15 through November 27, 1971. The harvest by archers has been constant since 1968.

The following type and number of antlerless permits were issued by the Director: Mainland sportsmen, 6000; Nantucket, 400; Martha's Vineyard, 600 and farmer-landowner, 270.

During the six-day shotgun deer season, December 6 through December 11, 1971, hunters reported harvesting 2248 deer. Of these deer, 1256 were males and 889 were females.

The top two deer harvest-producing counties were Berkshire and Franklin Counties. Martha's Vineyard moved from seventh to third while Worcester County dropped to fourth.

Massachusetts deer hunters harvest 57 percent of the total harvest of deer on the first and last days of the season. Thirty-three percent (33%) of the bucks were harvested on the first day of the season and 23 percent on the last day. Hunters reported 25 percent of the females harvested on Monday and 32 percent on Saturday.

A breakdown of the deer harvest for three deer management units (Mainland, Nantucket and Martha's Vineyard) is presented in table form.

The statewide success ratio of 7270 antlerless permit holders reporting the harvest of deer was as follows:

All deer	- (1314) 1 in 6
Males	- (405) 1 in 18
Females	- (889) 1 in 8
Females and buttonbucks	- (1023) 1 in 7

The 1314 deer harvested by antlerless permit holders represent 58 percent of the total 2248 deer harvested during the 1971 shotgun deer season.

Background: The antlerless permit system was continued during that portion of the season in which deer could be hunted with a gun, December 6 through December 11, 1971. Deer with antlers three inches or longer were legal deer for all licensed hunters. Antlerless deer were harvested only by permit. Three deer management units were established. The number of sportsmen antlerless permits issued per unit was as follows: Nantucket, 400 permits; Martha's Vineyard, 600; and the Mainland, 6000. Regulations require all deer harvested during the open season must be reported at an official Division deer checking station within 24 hours.

Objectives: To determine the annual harvest of deer in Massachusetts.

Findings: Table 1 presents a ten-year summary of the annual deer harvest per town by county from 1962 through 1971. The years 1962 through 1965 were years of voluntary reporting of deer by hunters. The six-year period 1966 through 1971 were years of compulsory deer reporting by successful deer hunters.

Successful deer hunters reported taking 2248 deer during the shotgun season, December 6 through December 11, 1971. Of these deer 1256 were males and 889 were females. Archers reported harvesting 26 males and 10 females, a total of 36 deer during the two-week archery season, November 15 through November 27, 1971 (there is no Sunday hunting in Massachusetts).

The distribution of the 1971 archery season harvest by sex, town and county is shown in Table 2.

Deer were harvested by archers in eight of the fourteen Massachusetts counties. Berkshire, Franklin and Hampden Counties account for 72 percent of the total reported deer kill.

Table 3 presents a summary of the 1971 shotgun deer harvest by sex, county and the rank of importance to the deer harvest for a three-year period. It is evident (Table 3) that the top deer harvest-producing counties, namely, Berkshire and Franklin Counties, have remained unchanged for the past three years.

A significant change in rank was noted for Dukes County moving from rank Number 7 to rank Number 3. This change moved Worcester County to rank Number 4. It should be noted that there was a substantial increase in the number of antlerless permits issued for Martha's Vineyard only. (There were 600 Martha's Vineyard only antlerless permits issued in 1971. Previously, Dukes County was included in the general statewide sportsmen antlerless permit allotment.)

Another significant change in rank was Hampshire County dropping from position Number 4 to Number 8. The reason for change in rank has not been determined. A look at Hampshire County kill per town (Table 1) shows that the harvest in the key towns of Belchertown, Chesterfield, Pelham and Worthington were considerably lower than the previous year.

Table 4 presents a summary of the deer harvest per day of kill during the shotgun season, December 6 through December 11, 1971.

The data presented in Table 4 show that 57 percent of the deer harvested were taken on the first and last days of the six-day deer season. Twenty-six percent of the deer harvest were taken on the second and fifth days of the shotgun week. The middle or third and fourth days of the shotgun season indicate the lowest hunting pressure days with only 17 percent of the total harvest.

A comparison of the harvest per sex per day shows complete reversal with 33 percent of the bucks taken on Monday and 23 percent on Saturday. The harvest of females was 25 percent on Monday and 32 percent on Saturday. It would appear that the deer hunters in 1971 did not attempt to fill the antlerless permits until the last day of the deer season.

Table 5 presents a summary of the Massachusetts deer harvest and hunter success per type of antlerless permit from 1967 through 1971.

The deer harvest by Massachusetts archers appears to have stabilized at 36 deer per year as seen in Table 5. Archers show a preference for male deer as the buck harvest is roughly two thirds of the total annual kill. The rutting period occurs concurrently with the archery season which may also account for the availability of the bucks.

A breakdown of the 1971 deer harvest per management unit shows that 1872 deer were shot on the Mainland. Of these 965 were antlered males, 183 buttonbucks and 724 females. On Nantucket Island, hunters reported 163 deer, of which 64 were adult males, 29 were buttonbucks and 70 were females. The harvest on Martha's Vineyard was 213 deer. Of these, 77 were antlered bucks, 41 were buttonbucks, and 95 were females. Statewide, the total deer harvest was 1359 males and 889 females for a total of 2248 deer.

A summary of the statewide deer harvest during the shotgun season from 1967 through 1971 shows that the male harvest increased from 937 bucks in 1967 to 1605 males in 1970. The 1971 harvest of 1359 males was 246 deer (-15.33%) lower than the 1605 bucks reported in 1970. The female harvest has shown an increase annually from 235 does in 1967 to 889 females in 1971.

Table 2. The 1971 Massachusetts Archery Season Deer Harvest by Sex, Town, and County, November 15 through November 27, 1971.

<u>County</u>	<u>Town</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>
Berkshire	Adams	1*		1
	Lee	2*		2
	Lanesborough	1*		1
	Richmond	1	1	2
	Great Barrington	1		1
	West Stockbridge	1*		1
	Egremont	2		2
	Williamstown	1		1
	Hancock	1		1
	Otis	<u>1</u>	<u>1</u>	<u>2</u>
		12	2	14
Dukes	Edgartown	2		2
	West Tisbury	<u>1</u>		<u>1</u>
		3		3
Franklin	Colrain	2*	1	3
	New Salem		1	1
	Leyden		1	1
	Sunderland		1	1
	Bernardston	<u>1*</u>		<u>1</u>
		3	4	7
Hampden	Blandford	1	1	2
	Tolland		1	1
	Montgomery		1	1
	Palmer	<u>1</u>		<u>1</u>
		2	3	5
Hampshire	Belchertown	1		1
	Ware	<u>1</u>		<u>1</u>
		2		2
Nantucket	Nantucket	1		1
Middlesex	Woburn	1		1
Worcester	Hardwick	2		2
	Warren	<u>1</u>	<u>1</u>	<u>1</u>
		2	1	3
Total		26 (8 BB)	10	36

* Buttonbuck (BB)

Table 3. Summary of the Massachusetts Deer Harvest by Sex per County and Rank of Harvest for the Shotgun Seasons 1969, 1970 and 1971.

<u>County</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>	<u>Rank 1971</u>	<u>Rank 1970</u>	<u>Rank 1969</u>
Barnstable	97	35	132	7	8	8
Berkshire	422	257	679	1	1	1
Bristol	1	0	1	12	12	12
Dukes	128	90	218	3	7	7
Essex	12	1	13	10	9	11
Franklin	282	241	523	2	2	2
Hampden	117	58	175	5	6	5
Hampshire	76	43	119	8	4	4
Middlesex	14	10	24	9	10	10
Nantucket	94	70	164	6	5	6
Norfolk	0	0	0	13	13	13
Plymouth	8	2	10	11	11	9
Suffolk	0	0	0	13	13	13
Worcester	109	82	191	4	3	3
	<u>1,359</u>	<u>889</u>	<u>2,248</u>			

Table 4. Summary of the Massachusetts Deer Harvest by Sex per Day, and the Percent Harvest per Day from Deer Checking Station Data, December 6 through December 11, 1971.

	<u>Mon. Dec. 6</u>	<u>Tues. Dec. 7</u>	<u>Wed. Dec. 8</u>	<u>Thurs. Dec. 9</u>	<u>Fri. Dec. 10</u>	<u>Sat. Dec. 11</u>	<u>Totals</u>
<u>Both Sexes</u>							
Total Kill	682	309	234	161	260	602	2,248
Percent	30.34	13.75	10.41	7.16	11.57	26.78	
<u>Males</u>							
Total Kill	454	214	131	97	150	313	1,359
Percent	33.41	15.75	9.64	7.14	11.04	23.03	
<u>Females</u>							
Total Kill	228	95	103	64	110	289	895
Percent	25.47	10.61	11.51	7.15	12.29	32.29	

Table 5. A Summary of the Massachusetts Deer Harvest and the Hunter Success by Type of Antlerless Permit, 1967 through 1971.

1971 Archery (Nov. 8-13; Nov. 15-20)

Males	26
Females	<u>10</u>
Total	36

Summary of Archery Kill, 1967 through 1971

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Male	17	21	27	24	26
Female	<u>4</u>	<u>13</u>	<u>10</u>	<u>12</u>	<u>10</u>
Totals	21	34	37	36	36

1971 Shotgun Season (Dec. 6-11)

Males	1359
Females	<u>889</u>
Total	2248

1971 Kill Breakdown per Management Unit

	<u>Mainland</u>	<u>Nantucket</u>	<u>Martha's Vineyard</u>	<u>Total</u>
Males, Adult	965	64	77	
Males, Buttonbuck	183	29	41	1359
Females	<u>724</u>	<u>70</u>	<u>95</u>	<u>889</u>
Totals	1872	163	213	2248

Summary of the Shotgun Statewide Deer Harvest, 1967 through 1971

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Males	937	1083	1424	1605	1359
Females	<u>235</u>	<u>310</u>	<u>585</u>	<u>764</u>	<u>889</u>
Totals	1172	1393	2009	2369	2248

Statewide Summary of Archery and Shotgun Harvest, 1967 to 1971

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Males	954	1104	1451	1629	1385
Females	<u>239</u>	<u>323</u>	<u>595</u>	<u>776</u>	<u>899</u>
Totals	1193	1427	2046	2405	2284

Table 5 (Continued)

Summary of Antlerless Deer Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
No. Applications (rounded numbers)	28,000	24,000	32,000	35,000	37,500
Permits issued:					
Sportsmen	2,000	2,000	4,000	6,000	6,000
Farmer-landowner	243	331	295	347	270
Nantucket	-	-	400	400	400
Martha's Vineyard	-	-	-	-	600
Totals	<u>2,243</u>	<u>2,331</u>	<u>4,695</u>	<u>6,747</u>	<u>7,270</u>

Harvest by Landowner-Farmer Permit

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Males	2	21	17	25	20
Females	<u>20</u>	<u>44</u>	<u>45</u>	<u>34</u>	<u>26</u>
Totals	22	65	62	59	46

Harvest by Sportsmen's Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Deer (both sex)	279	356	787	1057	1268

Breakdown of Statewide Harvest by Antlerless Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Antlered males	26	50	140	137	172
Buttonbucks	40	61	124	215	253
Females	<u>235</u>	<u>310</u>	<u>585</u>	<u>764</u>	<u>889</u>
Totals	301	421	849	1116	1314

Antlerless Permit Success Ratio
(rounded)

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
Farmer-landowner	1-11	1-5	1-5	1-6	1-6
Sportsmen	1 -9	1-6	1-6	1-6	1-6
Mainland					1-6
Nantucket	(125 permit deer)				1-3
Martha's Vineyard	(166 permit deer)				1-4

Table 6. A Summary of the 1971 Massachusetts Deer Harvest and the Hunter Success per Type of Antlerless Permit

	<u>Sportsmen (6000)</u>	<u>Nantucket (400)</u>	<u>Martha's Vineyard (600)</u>	<u>Landowner (270)</u>	<u>Totals</u>
Males	279 (175 BB*)	55 (29 BB*)	71 (41 BB*)	20 (8 BB*)	425 (279 BB*)
Females	<u>698</u>	<u>70</u>	<u>95</u>	<u>26</u>	<u>889</u>
	977	125	166	46	1314

Permit Success Ratio:

	<u>6000 Sportsmen</u>	<u>400 Nantucket</u>	<u>600 Martha's Vineyard</u>	<u>270 Landowner</u>
All deer	(977) 1 in 6	(125) 1 in 3	(166) 1 in 4	(46) 1 in 6
Males	(279) 1 in 22	(55) 1 in 7	(71) 1 in 8	(20) 1 in 14
Females	(698) 1 in 9	(70) 1 in 6	(95) 1 in 6	(26) 1 in 10
Antlered males	(104) 1 in 58	(26) 1 in 15	(30) 1 in 20	(12) 1 in 23
Buttonbucks	(175) 1 in 34	(29) 1 in 14	(41) 1 in 15	(8) 1 in 34
Females and buttonbucks	(873) 1 in 7	(99) 1 in 4	(135) 1 in 4	(34) 1 in 8

Total permits - 7,270

(1314) All deer	- 1 in 6
(405) Males	- 1 in 13
(889) Females	- 1 in 8
(1023) Females and buttonbucks	- 1 in 7

* BB - Buttonbucks

As seen in the summary of antlerless deer permits, the number of applications for the sportsmen permits has increased from 28,000 in 1967 to 37,500 in 1971. The number of sportsmen permits issued per year increased from 2000 in 1968 to 6000 in 1971. The number of farmer-landowner permits fluctuates annually with lower numbers on the odd years and high peaks on the even years.

The harvest of deer taken with farmer-landowner permits shows a decreasing harvest annually from a high of 65 deer in 1968 to 46 deer in 1971. Twenty-two deer were reported by farmer-landowners in 1967.

The summary of the statewide harvest of deer by antlerless permits (Table 5) shows the bulk of the harvest by permits are, as expected, female deer. A comparison of the percent of the total per sex for the first year (1967) of the permit system and the 1971 harvest total show a slight change in the harvest composition. In 1967, the antlered males made up nine percent of the harvest and in 1971 the antlered males composed 13 percent of the total. The buttonbucks made up 13 percent of the 1967 kill by permit and 19 percent of the 1971 harvest by antlerless permits. The female segment by permit in 1967 made up 78 percent of harvest and was 68 percent in 1971.

The success ratio per sportsmen antlerless permit remains unchanged since 1968 at the ratio of one successful permit holder in six on a statewide basis.

Table 6 presents a summary of the deer hunter success per type of antlerless permit per deer management unit. Due to the establishment of an added deer management unit (Martha's Vineyard), the data presented in Table 6 is not comparable to previous years' data. However, one can readily see that the probability of a permit holder on either of the two islands (Nantucket and Martha's Vineyard) of harvesting a deer is greater than on the Mainland or one in three (four) to one in six (Mainland).

Recommendations: It is recommended that the Division of Fisheries and Game personnel continue to compile and report deer harvest data.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist
and
William C. Byrne, Assistant

Date _____

JOB PROGRESS REPORT

EA 32.3:W-35-R-14 II-2

State Massachusetts

Cooperators: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Name: Game Population Trend and Harvest Survey

Job No. II-2 Job Title: Non-Hunting Deer Mortality Investigation

Period Covered: 1 January 1971 to 31 December 1971

Summary: Natural Resource Officers reported 694 deer mortalities during the reporting period. Of these mortalities 320 were males, 341 were females. No sex data were reported for 33 deer.

Motor vehicles (373) and dogs (219) were the two greatest causes of non-hunting mortalities with 85.3 percent of all deaths attributed to these causes. In order of importance the remaining 102 deer mortalities were caused as follows: others and unknown 41; illegal kills 39; fences 7; drownings 6; trains 5; and deer killed doing crop damage 4.

January through April were months of heavy mortalities. The increased kill in November is attributed to the rut period.

There was a 0.5 percent decrease in non-hunting deer mortalities (694 deer) for 1971 as compared to the reported mortalities for 1970 (698).

The adjusted sex ratio for the non-hunting deer mortalities for 1971 was 336 males to 358 females or 48.4 percent males to 51.6 females.

During 1971 Berkshire, Barnstable and Franklin Counties were the top ranking counties in reported deer mortalities. Barnstable moved from fourth rank position in 1970 to second position in 1971.

Background: Deer mortalities investigated and reported by Natural Resource Officers were compiled and tabulated. Project personnel observed wintering areas and/or concentrations of deer. Dead deer surveys were conducted in areas of deer concentrations.

Objectives: To determine the annual non-hunting decimating factors of the Massachusetts deer herd.

Findings:

Table No. 1 presents a summary of the sex and causes per month for 694 deer mortalities reported by Natural Resource Officers from 1 January 1971 through 31 December 1971. Of the 694 deer reported, 320 were males, 341 were females and 33 of no sex reported. The two greatest causes of deer mortalities were motor vehicles, 373 deer, and dogs, 219 deer. These two causes represent 85.3 percent (592 deer) of the 694 deer reported during the twelve-month period. The remaining 102 deer, or 14.7 percent, were reported (in order of importance) as follows: others and unknown 41; illegal kill 39; fences 7; drownings 6; trains 5; and killed doing crop damage 4.

The low mortality of 38 deer during January 1971 might be equated with the extreme, continued low and below normal temperatures coupled with a covering of deep powdered snow. There was a considerable lack of movement by deer, dogs and snowmobiles during this month (January).

A summary of deer mortalities and causes covering a nine-year period, from 1963 through 1971, is presented in Table 2. Due to a revised reporting system only the years from 1967 through 1971 present comparable data.

The number of non-hunting deer mortalities and the percent change covering the five-year period 1967 through 1971 are presented in Table No. 3. A significant change in the rate of non-hunting deer mortalities is observed when the percent change in mortalities between the years 1967 to 1968 (21 percent increase) is compared to the percent change in mortalities between the years 1970 to 1971 (-0.5 percent). It was expected that as the herd size increased, due to the antlerless permit system of deer hunting, that non-hunting deer mortalities would increase also, reflecting a corresponding increase in herd size. However, the rate of increase of non-hunting mortalities has decreased annually (Table 3). This suggests that (1) there may not be a linear relationship between increase in herd size and increase in non-hunting mortality, or (2) if there is such a relationship, then the non-hunting mortality data indicate a possible decrease in the rate of deer population increase. The latter has been reported in W-35-R:II-4.

Table No. 4 presents the sex composition per month of the deer mortalities and the adjusted sex ratio of deer mortalities from 1 January through 31 December 1971. In order to compute the adjusted sex ratio it was assumed that the mortality rate for deer of unreported sex was in the same proportion by sex class as the mortality rate for deer whose sex was reported. The adjusted sex ratio of 48.4 males to 51.6 females changed slightly in favor of the females from the 1970 ratio which was 50.1 males to 49.9 females.

Table 1. A Summary of the Sex and Causes of Massachusetts Deer Mortalities Reported by Natural Resource Officers, 1 January through 31 December 1971.

	January		February		March		April		May		June	
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.
Motor Vehicles	5	5	11	5	15	23	11	16	10	18	15	29
Dogs	7	16	35	38	28	29	7	9	4			3
Illegal Kill		1	1		1							
Drowned	1					1					1	
Fences	1			1			4					
Trains				2		1	1					
Crop Damage											3	
Other and Unknown	1	1	1	2	7	8	4		2		19	2
Subtotal	15	22	48	46	51	62	26	26	14	20	34	34
Total (Cum. Total)	38 (38)		107 (145)		123 (268)		55 (323)		34 (357)		53 (410)	
	July		August		September		October		November		December	
Motor Vehicles	12	8	4	5	9	13	31	17	45	21	17	23
Dogs		1		1					6	7	4	2
Illegal Kill							1	5	2	1	7	21
Drowned	1						1					1
Fences												
Trains										1		
Crop Damage	1											
Other and Unknown												
Subtotal	14	10	5	6	11	13	35	24	53	30	29	48
Total (Cum. Total)	26 (436)		11 (447)		25 (472)		60 (532)		84 (616)		78 (694)	
Cause	Male				Female				Total			
Motor vehicles	185				183				373			
Dogs	91				106				219			
Illegal Kill	12				27				39			
Drowned	4				1				6			
Fences	5				2				7			
Trains	0				5				5			
Crop Damage	4				0				4			
Other and Unknown	19				17				41			
Totals	320				341				694			

Table 2. Summary of Deer Mortalities and Causes of Massachusetts Deer Reported by Natural Resources Officers, 1963 through 1971.

<u>Cause</u>	<u>1963</u>	<u>1964</u>	<u>1965</u>	<u>1966</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>Total</u>	<u>8-Yr. Avg.</u>	<u>1971</u>
Motor Vehicles	321	225	242	277	334	456	397	400	2,652	332	373
Dogs	133	40	38	39	62	74	166	204	756	95	219
Illegal Kills	18	13	20	22	48	29	39	25	214	27	39
Unknown Causes	19	31	20	29	41	31	51	38	260	33	41
Crop Damage and Legal Kills	6	1	14	3	4	7	2	14	51	6	4
All Other Causes	<u>26</u>	<u>15</u>	<u>22</u>	<u>19</u>	<u>19</u>	<u>16</u>	<u>27</u>	<u>17</u>	<u>161</u>	<u>20</u>	<u>18</u>
Totals	523	325	356	389	508	613	682	698	4,094	512	694

Table 3. A Comparison of the Total Non-Hunting Deer Mortalities of Massachusetts Deer from 1967 through 1971.

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>
No. of Deer	508	613	682	698	694
Percent Change		21%	11%	2%	-0.5%

Table 4. A Comparison of the Sex of the Actual Number of Deer Mortalities and the Adjusted Sex Data for 1971* Massachusetts Deer.

<u>Month</u>	<u>Male</u>	<u>Female</u>	<u>No Sex</u>	<u>Total</u>	<u>Adjusted</u>	
					<u>Male</u>	<u>Female</u>
January	15	22	1	38	15	23
February	48	46	13	107	55	52
March	51	62	10	123	56	67
April	26	26	3	55	27	28
May	14	20	0	34	14	20
June	19	34	0	53	19	34
July	14	10	2	26	15	11
August	5	6	0	11	5	6
September	11	13	1	25	11	14
October	35	24	1	60	36	24
November	53	30	1	84	54	30
December	<u>29</u>	<u>48</u>	<u>1</u>	<u>78</u>	<u>29</u>	<u>49</u>
Total	320	341	33	694**	336	358

* These data were reported by Natural Resource Officers.

** Hunting and crippling losses not included.

Adjusted sex ratio: 336 males : 358 females
 94 males : 100 females
 48.4% males : 51.6% females

Table 5. Massachusetts Non-Hunting Deer Mortalities per County per Cause and Rank for 1971 and 1970.

County	Auto	Dogs	Illegal Kill	Drowned	Fences	Trains	Crop Damage	Other and Unknown	Total	Rank	1970 Total	Rank
Barnstable	79		4	3	2			5	93	2	74	4
Berkshire	104	48	7					16	175	1	184	1
Bristol	3	1		1		1			6	11	8	11
Essex	13	11			1				25	9	32	7
Franklin	35	44	1	1	2	2	2	3	90	3	127	2
Hampden	35	7	1		1	2	2	3	51	6	47	6
Hampshire	16	45	1		1			2	65	5	56	5
Middlesex	14	4							18	10	22	8
Norfolk			1						1	13		
Plymouth	23	7	3					3	36	8	22	8
Suffolk									0	14		
Worcester	29	43	8	1				4	85	4	99	3
Dukes	13	9	13					5	45	7	17	8
Nantucket	<u>4</u>								<u>4</u>	12	11	10
Totals	373	219	39	6	7	5	4	41	694			

A summary of the non-hunting mortalities per county per cause for 1971 is presented in Table 5. Table 5 also ranks the total number of deer mortalities by county and compares the causes of mortalities by county.

The three top ranking counties in order of importance are Berkshire, Barnstable and Franklin (Table 5). Barnstable County in second place with 93 reported mortalities has replaced Worcester County which has dropped to fourth place.

It is of interest to note that Berkshire and Franklin Counties rank first and second, in order of importance, as top deer harvest producing counties in the state; while Barnstable County ranks seventh place.

Recommendations: It is recommended that deer project personnel continue to determine the annual non-hunting deer decimating factors of the Massachusetts deer herd.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist
and
William C. Byrne, Assistant

Date _____

MASS.
EA 32.3:W-35-R-14/II-3

JOB PROGRESS REPORT

State Massachusetts
Cooperators: Massachusetts Division of Fisheries and Game
Project No.: W-35-R-14 Name: Game Population Trend and Harvest Survey
Job No. II-3 Title: Deer Fertility Studies
Period Covered: 1 January 1972 through 31 May 1972

Summary:

Sixty-three female deer mortality carcasses were collected by Natural Resource Officers and Division personnel from 1 January through 31 May 1972. The carcasses were examined and the age and stage of pregnancy determined. Fetus and corpora lutea counts were documented.

There were no significant changes in the 1972 reproductive rates compared to the 1971 rates or the mean rates for a seven-year period, 1966 through 1972. A summary of seven years of reproductive data (1966-1972) is as follows:

<u>Age at Parturition</u>	<u>Sample Size</u>	<u>No. Fawns Produced</u>	<u>1966-1972 Reproductive Rate</u>	<u>1972 Rate</u>
Yearling	152	38	1: .27 (100:27)	(1: .33)
Two years	84	119	1:1.39 (100:139)	(1:1.55)
Adults	181	312	1:1.74 (100:174)	(1:1.69)

A disproportionate sex ratio of 100 males to 106 females was recorded for 66 fetuses.

Background:

Natural Resource Officers and Division of Fisheries and Game personnel collected carcasses of female deer mortalities from 1 January through 31 May. These carcasses were taken either to the nearest Division installation or brought directly to Field Headquarters in Westboro. Project personnel removed the fetuses and ovaries.

The age of the deer examined was determined by the tooth replacement and/or wear technique. Fetus age was determined by crown-rump measurement compared to a table of known age fetus measurements.

A gross examination of the ovaries was made by slicing each ovary longitudinally. The number of current corpora lutea was recorded. No intensive search was made for corpora albicantia.

Objectives:

To determine the reproductive rate per age class of the Massachusetts deer herd.

Findings:

During the five-month period covered by this report (January to May), 63 female deer were examined. Table No. 1 presents a breakdown of the age composition and the percent per age class. The age of these deer is the age at parturition.

Since all the female deer collected and examined for this report were collected on the Mainland, any comparison of data will be based on the Mainland deer herd.

The examination of the reproductive tracts of 63 female deer mortalities collected from 1 January to 31 May 1972 found that

1. Of the 21 yearlings, seven were pregnant and were carrying five fawns. Two tracts contained remains of the embryonic sacs. However, no embryos were found.
2. All eleven two-year-old females were pregnant and were carrying 17 fawns.
3. Of the 31 adults (three years and older), 26 were pregnant and were carrying 49 fawns. Three does were not pregnant and in one doe the fetus or fetuses were missing. The reproductive tract of one adult was missing.

A comparison of the age composition and the percent per age class of the female deer checked at Mainland biological deer check stations and the female deer mortalities examined for this report is presented in Table 2.

Although the sample size per age class of the mortality data might be considered small, there has been a consistency for the past five years with a few exceptions that the percentages per age class of the harvest data and the mortality data complement each other. This could be interpreted within certain limitations that the harvest and mortality samples represent the female segment of the Mainland deer herd.

There is no significant difference between the age composition of the hunting harvest sample and the mortality sample collected for this report. A chi-square test total of 1.2952, with 5 degrees of freedom, shows that the mortality sample was taken from the same parameter (population) as the harvest or hunting sample.

Table 1. The Age Classification and the Percent per Age Class of Sixty-Three Female Deer Mortalities in Massachusetts, 1 January through 31 May 1972.

<u>Age in Years at Parturition</u>	<u>Number in Sample</u>	<u>Percent of Sample</u>
1	21	33.34
2	11	17.46
3	10	15.87
4	9	14.29
5	4	6.35
6	4	6.35
7	2	3.18
8-9	1	1.58
10 plus	<u>1</u>	<u>1.58</u>
	63	100.00

Table 2. The Age Composition and Percent per Age Class of Harvested Female Deer Checked at Biological Deer Stations (Mainland) December 1971 and Sixty-Three Female Deer Mortalities Collected January through May 1972.

<u>Age at Harvest</u>	<u>No. in Sample</u>	<u>Percent of Sample</u>	<u>Age at Parturition</u>	<u>No. in Sample</u>	<u>Percent in Sample</u>
6 mo.	101	27.15	1	21	33.33
1-1/2	64	17.20	2	11	17.46
2-1/2	69	18.56	3	10	15.87
3-1/2	51	13.71	4	9	14.29
4-1/2	33	8.87	5	4	6.35
5-1/2	20	5.38	6	4	6.35
6-1/2	14	3.76	7	2	3.18
7-1/2	11	2.96	8-9	1	1.58
8-9-1/2	9	2.42	10+	1	1.58
10-1/2+	—	—		—	—
	372	100.00		63	100.00

Table 3. A Summary of Corpora Lutea and Fetus Counts for Pregnant Yearlings, Two-Year-Olds and Adult Female Deer in Massachusetts, 1 January through 31 May 1972.

<u>Age</u>	<u>No. Females Pregnant</u>	<u>No. Corpora Lutea</u>	<u>No. Fetus</u>	<u>Single</u>	<u>Twins</u>	<u>Triplets</u>
Yearlings	7	7*	7	7	0	0
Two Years	11	23	17	5	6	0
Adults	26	57	49	5	19	2

* One pair ovaries missing but assumed that the doe had at least one current corpora lutea.

Based on the 61 female deer collected from 1 January through 31 May 1972, the following reproductive rates were calculated:

<u>Age Class</u>	<u>No. Does</u>	<u>No. Fawns</u>	<u>Calculated Reproductive Rate</u>
Yearling	21	7	100 does : .33 fawns (1 : .33)
Two-year-old	11	17	100 does : 1.55 fawns (1 : 1.55)
Three-year plus	29	49	100 does : 1.69 fawns (1 : 1.69)

Table 3 presents a summary of the corpora lutea and the fetus counts of pregnant female deer collected 1 January through 31 May 1972. The seven pregnant yearling does were each carrying a single fawn. Of the 11 pregnant two-year-old does, five were carrying single fawns and six were carrying twins. Of the 26 adult does, five were carrying single fawns, 19 were carrying twins and two does were carrying triplets.

Assuming that the three age classes presented in Table 3 are representative of the female segment of the Mainland deer herd, the importance of maintaining and/or increasing the number of adults is evident. Simply add the number of yearlings (7) and the number of two-year-olds (11), giving a total of 18 pregnant females that produce 24 fawns. The 26 pregnant adults produced 49 fawns which is 67 percent of the total fawn production.

Table 4 presents a summary of reproductive rate data from 1966 through 1972. There was no significant change in the reproductive rate for any of the three age classes in 1972 compared to the previous year.

The mean reproductive rate for the three age classifications of female deer are presented in Table 5. There were no significant changes in the mean reproductive rates per age class (Table 5).

A breakdown of the sex of deer fetuses per age class of the dam is as follows:

<u>Age</u>	<u>No. Does</u>	<u>Sex of Fetus</u>		<u>Total Fetus</u>
		<u>Male</u>	<u>Female</u>	
Yearling	5	2	3	5
Two Year	9	8	6	14
Three Years Plus	<u>25</u>	<u>22</u>	<u>25</u>	<u>47</u>
	39	32	34	66

The sex ratio of the 66 fetuses of 39 female deer was 100 males to 106 females. This is a complete change from the expected 106 males to 100 females. The same type of data seen in Job Progress Report W-35-R-13, Job II-3 showed the sex ratio for 102 fetuses of 61 does to be 100 males to 104 females. The 1971 and 1972 sex ratio

Table 4. A Summary of the Reproductive Rate Data per Age Class of 417 Massachusetts Female Deer Mortalities, 1 January 1966 through 31 May 1972.

<u>Yearlings</u>					
<u>Year</u>	<u>Sample Size</u>	<u>Pregnant</u>	<u>Not Pregnant</u>	<u>No. Fawns</u>	<u>Annual Reproductive Rate</u>
1966	16	9	7	11	0.69
1967	12	1	11	1	0.10
1968	14	2	12	2	0.14
1969	25	5	20	5	0.20
1970	37	4	33	4	0.11
1971	27	8	19	8	0.30
1972	<u>21</u>	<u>7</u>	<u>14</u>	<u>7*</u>	0.33
Total	152	36	116	38	
Mean	21.71			5.43	0.27
<u>Two-Year-Olds</u>					
1966	4	3	1	5	1.25
1967	10	9	1	12	1.20
1968	12	11	1	16	1.33
1969	16	16	0	23	1.44
1970	15	14	1	20	1.33
1971	16	15	1	26	1.63
1972	<u>11</u>	<u>11</u>	<u>0</u>	<u>17</u>	1.55
Total	84	79	5	119	
Mean	12.00			17.00	1.39
<u>Adults**</u>					
1966	18	17	1	31	1.72
1967	10	10	0	20	2.00
1968	20	17	3	32	1.60
1969	35	32	3	63	1.80
1970	28	23	5	47	1.67
1971	41	39	2	70	1.71
1972	<u>29</u>	<u>26</u>	<u>3</u>	<u>49</u>	1.69
Total	181	164	17	312	
Mean	25.86			44.57	1.74

* Five (5) measured fetuses, two (2) embryos missing

** Three years and older

Table 5. The Mean Reproductive Rate of Three Age Classifications of Massachusetts Deer, 1966 through 1972.

<u>Age</u>	<u>Sample Size</u>	<u>No. Fawns</u>	<u>1966-1971 Mean Reproductive Rate</u>	<u>1966-1972 Mean Reproductive Rate</u>
Yearling	21.71 (152)	5.43 (38)	1: .27 (100: 27)	1: .24 (100: 24)
Two Year	12.00 (84)	17.00 (119)	1:1.39 (100:139)	1:1.40 (100:140)
Three years plus	25.86 (181)	44.57 (312)	1:1.74 (100:174)	1:1.73 (100:173)

data for 168 fetuses was tested using chi-square. The test showed that there was not a significant change from the expected sex ratio 106 males to 100 females.

Recommendations: It is recommended that Division personnel continue to collect, compile, and report reproductive data pertaining to the Massachusetts deer herd.

Massachusetts Division of Fisheries and Game
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist
and
William C. Byrne, Assistant

Date _____

MASS.

EA32.3:W-35-

JOB PROGRESS REPORT

WR-14/II-4 4

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Title: Game Population Trend and Harvest Survey

Job No.: II-4 Job Title: Management of the Massachusetts Deer Herd

Period Covered: 1 July 1971 to 30 June 1972

Summary:

The sex and age composition of the state deer harvest was checked at ten biological deer checking stations. The calculated minimal deer population was 11,571 deer pre-hunting season for 1971. There was a slight decrease in the male segment of the deer harvest per square mile of deer range of the mainland management unit. All mainland counties showed an increase in the female harvest of deer per square mile of deer range. Fifty-eight percent of the statewide deer harvest was taken by antlerless permit holders.

There was an increase in the overall deer harvest on Martha's Vineyard which was attributed to the 600 antlerless permits issued for Martha's Vineyard only. A planned slight decrease in the Nantucket deer harvest was realized.

The data presented in this report indicate that the antlerless permit system of harvesting deer can be an effective and highly sensitive management tool. An increase or decrease in the number of antlerless permits issued for a deer hunting season will result in an increase or decrease in the deer harvest two years later. In addition, the data indicate that it is possible to manage deer on a county basis of deer harvest per square mile of deer range.

Problem areas within the mainland management unit were defined (i.e., low reported harvest of deer in the eastern counties exclusive of Barnstable County).

A summary of the 1971 deer harvest shows that the adult male deer kill was 1,106 animals which is 19.73 percent less than the 1970 harvest of 1,378 antlered deer. A continued decline in the number of antlered bucks can be expected for the 1972 and 1973 deer seasons.

Deer herd management recommendations were presented.

Objectives:

To estimate the size of the deer herd in Massachusetts and to recommend management techniques.

Procedures: Herd management recommendations will be determined after examination of factors consisting of sex and age composition of the herd, reproductive rates, average removal rates, hunting pressure, and deer kill per square mile of deer range.

Findings: During the 1971 shotgun deer hunting season, biological data (sex and age) was collected at ten deer checking stations. Table 1 presents a summary of the sex and age composition data collected per management unit, expanded to estimate the 1971 reported harvest.

A summary of the sex and age composition of the deer kill reported at the biological deer checking stations from 1967 through 1971 (1971 was the first year that biological data were collected on Nantucket and Martha's Vineyard) is presented in Table 2.

Table 3 is a summary of the expanded harvest of deer based on the percent per age class and sex reported at the mainland biological deer checking stations, 1967 through 1971.

The summary of the deer kill per square mile for all deer (Table 6) shows that for the mainland herd the rate of harvest has remained constant at .3 deer per square mile since 1969. On a county basis, it can be seen that the harvest in Franklin County increased one-tenth of a deer per square mile from .7 to .8 deer. Berkshire's harvest remains the same at .8 deer per square mile. The remaining counties show a loss of from 0.1 to 0.2 deer per square mile during the 1971 deer season.

The harvest of males (button bucks included) in the mainland herd dropped slightly from 0.20 bucks per square mile to 0.18 bucks per square mile. This may appear insignificant, however, this is the first downward trend since 1968. The greatest change in buck harvest per square mile is noted for Worcester County with a change from 0.2 in 1970 to 0.1 in 1971.

The summary of Table 5 shows a slight increase in the take of female deer for 1971 on the mainland. All mainland counties showed an increase in the female harvest with the exception of Hampshire and Worcester Counties. Both of these counties presented a decrease in the female segment of the harvest with Hampshire sliding from 0.2 female to 0.1 female deer per square mile in the harvest. Worcester County regressed slightly from 0.10 does to 0.06 does per square mile.

The effect of an increase of 600 antlerless permits on Martha's Vineyard is evident in Tables 5 and 6. The female harvest doubled from 0.5 deer per square mile in 1970 to 1.0 deer per square mile in 1971. The male harvest increased from 1.6 males per square mile in 1970 to 2.5 males per square mile in 1971. The planned decrease in the Nantucket deer harvest can be seen in Tables 4, 5 and 6.

Table 1. Sex and age of Massachusetts deer at ten biological deer checking stations expanded to include the statewide harvest for 1971.

Age	Males			Females		
	No.	Percent	Harvest Expanded	No.	Percent	Harvest Expanded
<u>Mainland</u>						
6 mo.	100			101	27.15	197
1-1/2	211	45.57	440	64	17.20	125
2-1/2	103	22.25	215	69	18.56	135
3-1/2	77	16.63	160	51	13.71	99
4-1/2	43	9.28	90	33	8.87	64
5-1/2	19	4.10	40	20	5.38	39
6-1/2	3	.65	6	14	3.76	27
7-1/2	6	1.30	12	11	2.96	21
8-1/2-9-1/2	1	.22	2	9	2.42	17
10-1/2	—	—	—	—	—	—
Total	563	100.00	965	372	100.00	724
<u>Nantucket</u>						
6 mo.	27			14	21.21	14
1-1/2	38	60.32	38	22	33.33	22
2-1/2	13	20.63	13	12	18.18	12
3-1/2	7	11.11	7	13	19.19	13
4-1/2	4	6.35	4	2	3.03	2
5-1/2	0	-	0	3	4.54	3
6-1/2	0	-	0	0	-	0
7-1/2	1	1.59	1	0	-	0
8-1/2-9-1/2	0	-	0	0	-	0
10-1/2	0	-	0	0	-	0
Total	90	100.00	63	66	100.00	66
<u>Martha's Vineyard</u>						
6 mo.	39	-	-	31	35.23	31
1-1/2	41	56.94	41	11	12.50	11
2-1/2	14	19.44	14	16	18.18	16
3-1/2	8	11.11	8	11	12.50	11
4-1/2	4	5.55	4	4	4.54	4
5-1/2	3	4.17	3	8	9.09	8
6-1/2	0	-	0	2	2.27	2
7-1/2	1	1.39	1	2	2.27	2
8-1/2-9-1/2	1	1.39	1	3	3.41	3
10-1/2	0	-	0	0	-	0
Total	111	100.00	72	88	100.00	88

Table 2. A summary of the sex and age composition of Massachusetts deer reported at mainland biological stations, 1967 through 1971.

Age	Males					Females				
	1967	1968	1969	1970	1971	1967	1968	1969	1970	1971
6 mo.	28	61	67	121	100	31	44	76	90	101
1-1/2	167	193	239	263	211	32	33	55	54	64
2-1/2	122	126	153	147	103	12	28	55	69	69
3-1/2	62	87	98	97	77	13	16	36	46	51
4-1/2	25	35	55	59	43	7	11	24	29	33
5-1/2	10	17	21	21	19	3	3	11	14	20
6-1/2	5	4	14	7	3	1	4	0	8	14
7-1/2	1	0	2	7	6	0	3	2	0	11
8-1/2-9-1/2	1	0	2	1	1	3	3	0	0	9
10-1/2	0	0	1	0	0	1	1	0	0	0
Total	421	528	652	723	563	109	146	259	310	372

A summary of the sex and age composition of Massachusetts deer reported at Martha's Vineyard and Nantucket deer checking stations in 1971.

Age	Martha's Vineyard		Nantucket	
	Males	Females	Males	Females
6 mo.	39	31	27	14
1-1/2	41	11	33	22
2-1/2	14	16	13	12
3-1/2	8	11	7	13
4-1/2	4	4	4	2
5-1/2	3	8	-	3
6-1/2	-	2	-	-
7-1/2	1	2	1	-
8-1/2-9-1/2	1	3	-	-
10-1/2	-	-	-	-
Total	111	88	90	66

Table 3. Expanded statewide harvest of Massachusetts deer based on the percent per age class reported at mainland biological deer checking stations, 1967 through 1971.

Age	Males					Females				
	1967	1968	1969	1970	1971	1967	1968	1969	1970	1971
6 mo.*	53	61	125	220	183	67	93	172	223	197
1-1/2	372	433	531	602	440	69	70	124	134	125
2-1/2	272	276	340	337	215	40	59	124	171	135
3-1/2	138	190	218	222	160	28	34	81	114	99
4-1/2	56	77	122	135	90	15	23	55	72	64
5-1/2	22	37	46	48	40	6	7	25	35	39
6-1/2	11	9	31	16	6	2	8	-	20	27
7-1/2	2	-	4	16	12	-	7	5	-	21
8-1/2-9-1/2	2	-	4	2	2	6	7	-	-	17
10-1/2	-	-	3	0	-	2	2	-	-	-
Totals	875*	1022	1299	1378	1148	235	310	586	769	724

* Six-month deer not included in male total.

Table 4. A summary of the male Massachusetts deer harvest per square mile of deer range per county, 1967 through 1971.

County	Sq. Mi. Deer Range	1967		1968		1969		1970		1971	
		No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.
Barnstable	290.5	45	0.2	49	0.2	66	0.3	97	0.3	97	0.3
Berkshire	839.0	249	0.3	342	0.4	418	0.5	446	0.5	422	0.5
Franklin	649.1	193	0.3	253	0.4	288	0.4	305	0.5	282	0.4
Hampden	524.3	75	0.1	106	0.2	139	0.3	141	0.3	117	0.2
Hampshire	431.5	75	0.2	116	0.3	135	0.3	142	0.3	76	0.2
Worcester	1307.6	152	0.1	84	0.1	199	0.2	233	0.2	109	0.1
Eastern Group*	2271.6	46	0.0	33	0.0	51	0.0	50	0.0	35	0.0
Mainland	6213.6	835	0.1	983	0.2	1301	0.2	1420	0.2	1138	0.2
Dukes	86.5	68	0.8	61	0.7	72	0.8	95	1.1	128	1.5
Nantucket	35.9	51	1.4	60	1.7	78	2.2	106	3.0	93	2.6

Button bucks included in these data.

* Eastern Group - includes Bristol, Essex, Middlesex, Norfolk, and Plymouth counties.

Table 5. A summary of the female Massachusetts deer harvest per square mile of deer range per county, 1967 through 1971.

County	Sq. Mi. Deer Range	1967		1968		1969		1970		1971	
		No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.	No. Deer	Kill Per Sq. Mi.
Barnstable	290.5	14	0.0	9	0.0	22	0.1	41	0.1	35	0.1
Berkshire	839.0	44	0.1	79	0.1	145	0.2	197	0.2	257	0.3
Franklin	649.1	63	0.1	100	0.2	136	0.2	139	0.2	241	0.4
Hampden	542.3	13	0.0	19	0.0	45	0.1	45	0.1	58	0.1
Hampshire	431.5	24	0.1	40	0.1	67	0.2	71	0.2	43	0.1
Worcester	1307.6	47	0.0	42	0.0	85	0.1	124	0.1	82	0.1
Eastern Group*	2271.6	11	0.0	12	0.0	17	0.0	27	0.0	13	0.0
Mainland	6213.6	214	0.0	300	0.0	523	0.1	650	0.1	729	0.1
Dukes	86.5	17	0.2	14	0.2	23	0.3	45	0.5	90	1.0
Nantucket	35.9	8	0.2	9	0.3	50	1.4	86	2.4	70	2.0

* Eastern Group - includes Bristol, Essex, Middlesex, Norfolk and Plymouth counties.

Table 6. A summary of the Massachusetts deer harvest per square mile of deer range per county, 1967 through 1971.

County	Sq. Mi. Deer Range	1967		1968		1969		1970		1971	
		No.	Kill Per Sq. Mi.	No.	Kill Per Sq. Mi.	No.	Kill Per Sq. Mi.	No.	Kill Per Sq. Mi.	No.	Kill Per Sq. Mi.
		Deer	Sq. Mi.	Deer	Sq. Mi.	Deer	Sq. Mi.	Deer	Sq. Mi.	Deer	Sq. Mi.
Barnstable	290.5	58	0.2	58	0.2	88	0.3	138	0.5	132	0.5
Berkshire	839.0	293	0.4	421	0.5	563	0.7	643	0.8	679	0.8
Franklin	649.1	256	0.4	352	0.5	424	0.7	445	0.7	523	0.8
Hampden	524.3	88	0.2	125	0.2	184	0.4	186	0.4	175	0.3
Hampshire	431.5	97	0.2	156	0.4	202	0.5	213	0.5	119	0.3
Worcester	1307.6	197	0.2	126	0.1	284	0.2	357	0.3	191	0.2
Eastern Group*	2271.6	57	0.0	45	0.0	71	0.0	77	0.0	48	0.0
Mainland	6213.6	1049	0.2	1283	0.2	1824	0.3	2070	0.3	1867	0.3
Dukes	86.5	85	1.0	75	0.9	95	1.1	141	1.6	218	2.5
Nantucket	35.9	59	1.6	69	1.9	128	3.6	192	5.3	163	4.5

* Eastern Group - Bristol, Essex, Middlesex, Norfolk and Plymouth counties.

Figure 1 presents a summary of the deer harvest by sex per square mile of deer range per county for a six-year period, 1966 through 1971. The graphs show the sensitivity of the harvesting of deer using the antlerless permit system and the value of the harvest calculated on a kill per square mile of deer range basis. Here it is possible to compare the six major deer harvest producing counties and see fluctuations in the deer harvest.

A summary of the Massachusetts deer harvest by shotgun for a five-year period, 1967 through 1971, is presented in Table 7. Of the 2,248 deer harvested slightly over half (1,142) were females and button bucks. The number of antlered males taken in 1971 decreased 272 deer from the 1970 statewide harvest figure of 1,378 bucks.

Table 8 presents a summary of the percent change in the adult male harvest and the calculated minimal population of Massachusetts deer from 1967 through 1971.

Tables 9 and 10 present the adult male and female deer harvest per square mile of deer range per county in Massachusetts from 1967 through 1971. Both tables make it possible to compare the harvest of deer per county by kill per square mile of deer range rather than the total kill per county. It is interesting to note that in both tables the changes in the harvest are in tenths and in some cases hundredths deer per square mile. With data this sensitive by using the antlerless permit system it is possible to manage deer on a county basis.

The harvest data throughout this report in dealing with individual deer herd divisions tends to gloss over some problem areas within a division or management segment. The eastern portion of Massachusetts is heavily populated with an expanding human population which may account for the low or apparently stable deer harvest in Essex and Middlesex Counties. However, the deer harvest data for Plymouth, Bristol and parts of Norfolk County indicate a definite deer harvest problem. Deer mortality data (W-35-R-14, Job No. II-2, Non-Hunting Deer Mortality Investigations) show that deer are being killed in all of these counties by cars, dogs, etc. Yet the legal harvest remains at almost nil.

The established goal is to harvest a minimum of 0.5 bucks per square mile of deer range on the mainland. In order to achieve that goal it will be necessary to establish a large female breeding population. The data presented in this report strongly suggests that the deer harvest has increased to a point where more than half the deer reported were taken by permit holders. The data also show that the number of adult females harvested has increased annually as the number of antlerless permits increased. Meanwhile the antlered buck harvest has started to decrease.

Figure 1. A summary of the Massachusetts deer harvest by sex per square mile of deer range by county, 1966 through 1971.

Male

Female

Total

Berkshire
839.0 sq. mi.

Franklin
649.1 sq. mi.

Hampden
524.3 sq. mi.

Hampshire
431.5 sq. mi.

Worcester
1307.6 sq. mi.

Barnstable
290.5 sq. mi.

Mainland
6213.6 sq. mi.

Dukes
86.5 sq. mi.

Nantucket
35.9 sq. mi.

Table 7. A summary of the Massachusetts deer harvest by shotgun, 1967 through 1971.

Year	All		Adult		Button		All		Adult		Fawn		Total		Total Per-	
	Males		Males		Bucks		Females		Females		Females		Deer		mits	Issued
1967	937		884		53		235		168		67		1,172		2,236	
1968	1,083		1,022		61		310		217		93		1,393		2,331	
1969	1,424		1,299		125		586		415		171		2,010		4,695	
1970	1,598		1,378		220		769		546		223		2,367		6,747	
1971	1,359		1,106		253		889		643		246		2,248		7,270	

Table 8. A summary of percentage change in adult male harvest and calculated minimal populations of deer in Massachusetts, 1967 through 1971.

	Percent		Percent		Percent		Percent		Percent	
	1967	1967-68	1968	1968-69	1969	1969-70	1970	1970-71	1971	
Adult male harvest	875	+16.80	1,022	+27.10	1,299	+ 6.08	1,378	-19.73	1,106	
Calculated adult male population	2,059	+17.05	2,410	+31.95	3,180	- 0.82	3,154	-27.68	2,281	
Calculated adult female population	2,018	+31.92	2,964	+35.19	4,007	+32.24	5,299	-22.95	4,083	
Calculated minimal population	6,139	+42.37	8,740	+35.01	11,805	+27.18	15,014	-22.93	11,571	

Table 9. A summary of the adult male deer harvest per square mile of deer range per county in Massachusetts, 1967 through 1971.

County	Square Miles Deer Range	1967	1968	1969	1970	1971
Barnstable	290.5	.14	.15	.20	.23	.23
Berkshire	839.0	.27	.37	.44	.44	.42
Bristol	422.6	.01	.01	.01	.01	.002
Essex	344.4	.03	.03	.04	.06	.03
Franklin	649.1	.27	.35	.39	.39	.36
Hampden	524.3	.13	.19	.24	.22	.20
Hampshire	431.5	.16	.25	.27	.26	.15
Middlesex	583.1	.02	.01	.02	.02	.02
Norfolk	277.3	---	---	.01	---	---
Plymouth	544.2	.03	.02	.03	.02	.01
Worcester	1307.6	.11	.06	.14	.15	.07
Total	6213.6	.12	.14	.13	.19	.15
Dukes	86.5	.71	.65	.72	.91	1.01
Nantucket	35.9	1.34	1.56	1.89	2.42	1.78

Table 10. A summary of the adult female deer harvest per square mile of deer range per county in Massachusetts, 1967 through 1971.

County	Square Miles Deer Range	1967	1968	1969	1970	1971
Barnstable	290.5	.03	.02	.06	.10	.09
Berkshire	839.0	.04	.06	.12	.16	.23
Bristol	422.6	.005	.002	.002	---	---
Essex	344.4	.006	.003	.01	.03	.003
Franklin	649.1	.07	.11	.14	.15	.27
Hampden	524.3	.02	.03	.06	.06	.03
Hampshire	431.5	.04	.06	.11	.11	.07
Middlesex	583.1	.002	.01	.01	.01	.01
Norfolk	277.3	.004	---	---	.004	---
Plymouth	544.2	.002	.01	.01	.01	.002
Worcester	1307.6	.02	.02	.05	.07	.05
Total	6213.6	.02	.03	.06	.07	.03
Dukes	86.5	.14	.10	.18	.36	.67
Nantucket	35.9	.17	.14	.97	1.67	1.53

The data (Figure 2) show that the results of harvesting deer by the permit system are not discernible until the second year after the harvest; i.e., the result of issuing 2,000 antlerless permits in 1967 and 1968 show an increase of antlered bucks in 1969 and the peak harvest of 1970 respectively. The number of antlerless permits issued in 1969 was increased to 4,000 which resulted in decline in the antlered harvest of 1971. In 1970 the number of antlerless permits issued was increased to 6,000. In 1971 there were 6,000 sportsmen's permits issued for state-wide use. It is expected that if the present trend continues the harvest of antlered bucks will continue to decline slightly in 1972 and 1973.

Table 11 shows the percent frequency ratio of adult females to adult males per county from 1967 through 1971. The percent frequency represents that portion of adult female deer harvested for each adult male harvested.

The effect of the number of antlerless permits issued per year can be demonstrated. In Berkshire County for the years 1967 and 1968, the percent frequency of adult females was .14 and .17 respectively. There were 2000 sportsmen's permits issued during these two years. In 1969 the number of permits issued was increased to 4,000 permits with the percent frequency of adult females increasing from .17 in 1968 to .27 in 1969. Six thousand permits were issued for 1970 and 1971 with the percent frequency of females increasing to .38 females in 1970 and .53 females in 1971. Simply stated, for every two antlered bucks harvested in Berkshire County in 1971, there was one adult female harvested. It might appear that there was an expanding deer herd in the Berkshires and the .53 percent frequency of adult females presents a healthy situation. However, observe the steady decline in the harvest numbers of antlered bucks from a high of 371 deer in 1969 to 352 antlered males in 1971.

The antlerless permit system not only protects and saves the female segment of a deer population but the system saves a number of button bucks. These animals (male fawns) are as important a segment of the deer herd as are the females. The button bucks that survive a hunting season and following winter will provide the huntable antlered male deer for the next eight to ten years.

During the 1972 and 1973 deer hunting seasons, it is imperative that emphasis be placed on increasing the antlered male population of the mainland deer management unit. This increase can be accomplished by a substantial reduction in the number of antlerless permits issued. The females saved by a permit reduction will produce fawns the following spring. The male fawns will represent the antlered bucks of the following year. Therefore, it is necessary to plan two years in advance to obtain the resulting increase in herd size.

FIGURE 2. A summary of the Massachusetts deer harvest per year per sex and age and the total number of antlerless permits issued in 1967 through 1971.

Table 11. Percent Frequency Ratio of Adult Females to Adult Males per County of Massachusetts Deer, 1967 through 1971.

County	1967				1968				1969				1970				1971			
	Adult		% Freq.		Adult		% Freq.		Adult		% Freq.		Adult		% Freq.		Adult		% Freq.	
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.
Barnstable	41	10	.24	.14	44	6	.14	.14	58	16	.28	.28	80	29	.36	.36	82	25	.30	.30
Berkshire	228	31	.14	.17	314	53	.17	.17	371	100	.27	.27	366	138	.38	.38	352	187	.53	.53
Bristol	6	2	.33	.20	5	1	.20	.33	3	1	.33	.33	4				1			
Essex	11	2	.18	.10	10	1	.10	.31	13	4	.31	.45	20	9	.45	.45	12	1	.08	.08
Franklin	176	44	.25	.30	230	68	.30	.36	256	93	.36	.39	251	98	.39	.39	232	176	.76	.76
Hampden	69	9	.08	.13	98	13	.13	.26	123	32	.26	.26	117	31	.26	.26	103	42	.41	.41
Hampshire	69	16	.23	.25	107	27	.25	.41	115	47	.41	.42	114	48	.42	.42	63	31	.49	.49
Middlesex	10	1	.10	.43	7	3	.43	.31	13	4	.31	.67	9	6	.67	.67	14	7	.50	.50
Norfolk		1							3					1						
Plymouth	16	1	.06	.38	8	3	.38	.43	14	6	.43	.50	8	4	.50	.50	7	1	.14	.14
Suffolk																				
Worcester	140	32	.23	.37	75	28	.37	.34	176	59	.34	.46	192	88	.46	.46	89	60	.67	.67
Total	766	149	.19	.23	898	203	.23	.32	1145	362	.32	.39	1161	452	.39	.39	955	530	.55	.55
No. of Permits	2000		2000		2000		2000		4000		4000		6000		6000		5000		5000	
Dukes County	61	12	.20	.16	56	9	.16	.26	62	16	.26	.39	79	31	.39	.39	87	58	.67	.67
No. of Permits																	600			
Nantucket	48	6	.13	.09	56	5	.09	.51	68	35	.51	.69	87	60	.69	.69	64	55	.86	.86
No. of Permits																	400		400	

Recommendations: The antlerless permits issued by the Director should be used only in the deer management unit for which the permits were issued.

Hunters should be required to transport their deer to the nearest official deer checking station upon killing a deer.

Fifty to 100 special (landowner issue) permits should be issued for Naushon Island.

Deer taken on offshore islands should be reported by the hunter on the island where the animal was shot.

The possibility for the sale of a big game license in Massachusetts should be investigated. When purchased, a portion of the big game license would become an application for an antlerless permit.

Any license sold by the Division of Fisheries and Game should incorporate the purchaser's Social Security number on the license.

Once a deer has been through a deer checking station and tagged, the trunk of a car or the rear window of a station wagon should be allowed to be closed.

Consideration should be given to the issuance of antlerless permits on a county adult-buck-per-square-mile-of-deer-range basis. This should commence with the 1973 deer hunting season.

The following number of sportsmen's antlerless permits for the 1972 shotgun deer hunting season are recommended:

<u>Deer Management Unit</u>	<u>Number of Antlerless Permits</u>
Naushon	50-100
Nantucket	400
Martha's Vineyard	600
Mainland	4,000

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved:

Colton H. Bridges, Superintendent

Prepared by

James J. McDonough, Game Biologist
and
William C. Byrne, Assistant

Date

MASS.

EA32.3;

JOB PROGRESS REPORT

W-35-R-14/ IV-1

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Title: Game Population Trend and Harvest Survey

Job No.: IV-1 Job Title: Experimental Turkey Stocking

Period Covered: 1 June 1971 to 31 May 1972

Summary: The fall 1971 estimated turkey population on seven release areas totaled 175 turkeys. Populations on individual areas were as follows: Quabbin-New Salem, 52; Barre-Oakham, 23; Douglas State Forest, 36; October Mountain area, 19; Town of Mt. Washington, 16; Myles Standish State Forest, 2; Holyoke Range, 2. Twenty-five additional birds were present in dispersed flocks.

Five turkeys were captured by cannon net in New Salem and transferred to Horse Mountain in Hatfield. Seven additional turkeys were captured, banded, and released near Underhill Brook on Prescott Peninsula in the Quabbin Reservation.

Through the courtesy of New York State personnel, seven wild-trapped Meleagris gallopavo silvestris were cannon netted in Allegany State Park, Cattaraugus County, New York, by the project leader and a University of Massachusetts graduate student. Captured turkeys were transported to Massachusetts and released in Beartown State Forest, Great Barrington, Berkshire County. Coordination efforts are being continued so that additional turkeys may be obtained to complete the Beartown stocking.

Objectives: To re-establish the wild turkey in the Commonwealth in sufficient numbers to allow for recreational hunting.

Procedures: Turkeys were censused by roadside counts, track counts and cooperator reports. Snowmobiles were used during the winter to provide access to the areas and to transport grain for baiting.

Turkeys were captured using mortar, jet and Thornsberry-style cannon nets. Captured turkeys were sexed, aged by primary feather molt and growth, weighed, leg banded, and color marked with patagial streamers.

Findings: A.

Statewide Turkey Populations

Statewide turkey population figures are presented for the fall 1971 period (Table 1). Winter and spring observations were limited by the scarcity of snow for tracking, followed by intensive rains with muddy roads limiting vehicular access. Quabbin-New Salem counts are based largely upon direct observation; whereas, counts on other areas are based upon screened reports from cooperators supplemented by a limited number of searches by the project leader and assistants.

Table 1. Fall turkey populations on seven Massachusetts release areas.

Area	Fall Population
Quabbin-New Salem	52*
Barre-Oakham	23
Douglas State Forest	36
October Mountain Area	19
Mount Washington	16
Myles Standish State Forest	2
Holyoke Range	<u>2</u>
Subtotal	150
Dispersed flocks	<u>25</u>
Total	175

* Prior to trapping

Quabbin Reservation-New Salem Populations

Prescott Peninsula in the Quabbin Reservation was the original release site in the current turkey restoration project. Twenty-two turkeys from three sources (Naushon Island, Allegheny Game Farm, and wild-trapped West Virginia silvestris) were released here in 1960 and 1961. Populations remained relatively static through 1965, and increased slightly following mild winters in 1966 and 1967. Since then, there has been a trend towards dispersal off the reservation into surrounding towns.

Turkey populations on Prescott Peninsula decreased following the cessation of artificial feeding in the winter of 1970-1971. Remaining turkeys seem wilder, as they flush more readily when approached. New Salem turkey flocks continue to be fed by local residents, and consequently are less wary of man.

Three known broods were produced in the Underhill Brook and Mt. Pleasant areas of the Quabbin during 1970. Two more broods were produced in the West Street region of New Salem. Later in summer, two of the Prescott broods merged and moved south on the Peninsula to the vicinity of Mt. Ram. The third brood was trapped, banded, and released near Underhill Brook, and was later reported near Andrews-Fisk Hill, six miles north across Route 202. The broods off West Street were trapped at Hamilton's Orchards, New Salem, and six turkeys transferred to Horse Mountain in Hatfield. A mild winter with little snow and a good crop of red oak acorns probably contributed to the winter survival of Quabbin turkeys. Reproductive success in the spring of 1972 was likely affected, however, by intense rains which may have chilled eggs and poults.

Cooperators reported turkey flocks near Junket Hill, Pelham; Dry Hill Road, Montague, and in the Town of Whately. Although these reports are considered reliable, probably representing dispersion from the Quabbin, fall and winter searches by the project leader failed to verify their presence, and they are listed in Table 1 under Dispersed Flocks.

Barre-Oakham Population

Fifteen turkeys, wild trapped in the Quabbin Reservation, were released in Barre in 1967 (11) and 1969 (4). Dispersion has since occurred into the nearby towns of Hubbardston, Oakham, and New Braintree. Decreased populations in 1971-1972 are probably the result of nest failures in spring 1972. Only one brood, near Burnshirt Brook in Barre, was reported during this period. Small flocks of adult turkeys remain near the Oakham Cemetery and Fairweather Hill in Hubbardston. Individual birds were sighted near Burrow Brook, South Barre, and off Sunrise Avenue, Barre. As in the Quabbin, reproduction in 1972 may have been affected by the unusual amount of rain during the spring. One nest, near Burnshirt Brook in Barre, was lost when the hen abandoned due to human disturbance. The eggs were collected, but artificial incubation proved unsuccessful.

Douglas State Forest Population

Nineteen Quabbin turkeys were released in Douglas Woods northwest of Wallis Reservoir in 1968 (12) and 1969 (7). Turkeys have dispersed throughout the forest, from Manchaug south to Wallum Lake. One small flock has been reported to the west in Webster, and a large flock of 25 to 27 turkeys in the swampy area near Buffum Road in Uxbridge. These turkeys, as with others of Quabbin mixed-background ancestry, display little wildness when subject to persistent feeding by misinformed citizens. One turkey in such a flock was cornered and killed by feral dogs in the spring of 1971.

Holyoke Range Populations

Mt. Tom in the Holyoke Range was the final stronghold of the native Eastern turkey in Massachusetts. The last surviving bird, a lone gobbler, was shot there in the winter of 1851. Unsuccessful releases using stock of unknown origin were made in 1915 and 1918. Recent releases (1964 and 1965) were made in Granby (3) and Hadley (11), but also proved unsuccessful, probably due to mortalities among the released birds, followed by dispersal of the remainder. Two tom turkeys, the first reported since 1970, were seen in Granby during early winter, 1972.

Myles Standish State Forest Population

Fourteen wild-trapped West Virginia M. g. silvestris were released in Myles Standish Forest in Plymouth and Carver in February and March 1966 and 1967. Individual dispersal occurred almost immediately after release, and the only reliably reported reproduction occurred in the spring of 1968. For the past two years, the only reliable sightings have been of adult toms, with the last report being in June-July 1971 near Webster Springs Road and College Pond Road in the southern portion of the forest.

October Mountain Area Populations

Twelve adult turkeys obtained from the Allegheny Game Farm in Julian, Pennsylvania were released off County Road, Town of Washington in April 1961. An additional seventeen turkeys, trapped at previous release sites in the Town of Mt. Washington (15) and the Quabbin Reservation (2), were released at County Road and near Sandwash Reservoir in August and September of 1962. These turkeys have displayed virtually no wildness from the time of their release to the present. A nuisance complaint by a Becket resident resulted in ten turkeys being trapped and moved to a forested area off West Hill Road in Middlefield in February 1968. In February 1971, ten additional tame turkeys were captured on the Simmon's property, Washington, using drugged grain. These inferior-strain turkeys were captured to effect their permanent removal from the range, and were subsequently donated to cooperating conservation agencies for display purposes. At present, only five adult turkeys are known to remain in Washington, and an additional twelve (sex and age unknown) are in Middlefield.

Town of Mt. Washington Population

In January 1961, eleven turkeys were trapped near Margaretville, Delaware County, New York, and released on the Whitbeck Farm, Town of Mt. Washington, Berkshire County, Massachusetts. The population from which these birds were trapped was initiated with game farm stock, but had existed for four generations in the wild. Local residents began feeding the turkeys almost immediately, and they soon became very tolerant of man. Nine of these turkeys (three adult, six immature) were trapped and transferred to October Mountain Forest in 1962. Town residents since then have been very protective of the turkeys and frequently treat Division inquiries with suspicion. Limited observations by Western District personnel indicate a spring (1972) population of sixteen turkeys. Their lack of wildness and presence on posted land probably negate any conceivable future hunting program in this town. Trapping activities utilizing Mt. Washington turkeys are not recommended.

Miscellaneous Reports

Reports of single turkeys or small groups were received from Chester, Cummington, Greenfield, Peru, Templeton, and West Brookfield. Time limitations and lack of an accurate sighting location precluded conducting field observations for verification. Two other reports, in Monson and West Tisbury, are now known to originate from illegal private releases.

B.

Trapping Activities

1. Turkey broods produced in the West Street region of New Salem have frequented the lower fields of Hamilton's Orchards for the past three summers. The owner of the orchard maintains some dwarf trees and had sustained some slight damage from turkey depredations during the fall of 1969. To preclude a repetition of this and to obtain turkeys for transplant, trapping (using a jet-type cannon net) was conducted on 29 September, resulting in a catch of six turkeys (Table 2). Captured birds were sexed, aged, weighed, leg banded and color marked with patagial tags. One hen was equipped with a back-pack radio-telemetry unit by a University of Massachusetts cooperator. The turkeys were transported in individual carrying crates and released on the F. W. Cole property on the east slope of Horse Mountain, Hatfield, Hampshire County. A few weeks after release, the turkeys were reported 2.4 kilometers south, near the Laurel Park swamp. The instrumented hen was killed by a dog in November.
2. On 24 November 1971, trapping operations were conducted in conjunction with Walter M. Tzilkowski, Consultant Wildlife Biologist. Eight turkeys were utilizing a bait site at a sawmill clearing south of Underhill Brook on Prescott

Table 2. Sex, Age, Weight and Capture-Release Data for Turkeys Trapped During 1971.

Band No.	Age	Sex	Weight		Capture Site	Date*	Release Site	Remarks
			KG	(lbs.)				
177	Imm.	F	2.7	(6.0)	Hamilton's Orchards New Salem	29 Sept.	Horse Mountain Hatfield	Instrumented; killed November 1971
178	Imm.	F	2.5	(5.5)	Hamilton's Orchards	29 Sept.	Horse Mountain	
179	Ad.	F	3.3	(8.3)	Hamilton's Orchards	29 Sept.	Horse Mountain	
180	Ad.	F	3.7	(8.2)	Hamilton's Orchards	29 Sept.	Horse Mountain	Wing tag lost on release Instrumented
181	Imm.	F	2.0	(4.3)	Hamilton's Orchards	29 Sept.	Horse Mountain	
182	Imm.	F	**		Underhill Brook Prescott Peninsula	24 Nov.	Same as capture site	
183	Imm.	F	**		Underhill Brook	24 Nov.	Capture site	Instrumented; found dead 27 March 1972.
184	Ad.	F	**		Underhill Brook	24 Nov.	Capture site	
185	Ad.	F	**		Underhill Brook	24 Nov.	Capture site	
186	Imm.	F	**		Underhill Brook	24 Nov.	Capture site	Not wing-tagged.
187	Ad.	F	**		Underhill Brook	24 Nov.	Capture site	
105	Imm.	M	3.2	(7.1)	Hamilton's Orchards	29 Sept.	Horse Mountain	
106	Ad.	M	**		Underhill Brook	24 Nov.	Capture site	

* All turkeys released on capture date.

** Not weighed.

Peninsula. Seven of these were captured, using a mortar-thrown net. Captured birds were leg banded, color wing-tagged (except the adult male), and two hens were instrumented. All were released at the capture site.

3. Through the courtesy of the New York State Department of Environmental Conservation, permission was secured for the Division to trap wild-strain turkeys in Allegany State Park, Cattaraugus County, New York. Prebaiting was accomplished by New York State personnel and trapping was conducted by the project leader and Consultant Biologist Tzilkowski during the period 6-17 March 1972. Although there had been excellent production and winter survival in the Park during 1971-1972, trapping operations were hampered by mild weather and early thawing, which made baiting of turkeys difficult. Cannon net malfunctions decreased capture success. Three shots were made resulting in the capture of eight turkeys. One hen died, presumably of shock, soon after capture, and the other seven were released in good condition in a field off Alcott Street, Beartown State Forest, Great Barrington, Berkshire County, Massachusetts. Band number, sex, age, weight, and capture-release data are presented in Table 3.

Table 3. Sex, Age, Weight and Capture-Release Data for Wild-Trapped Turkeys Captured in Allegany State Park, Cattaraugus County, New York and Transferred to Beartown State Forest, Great Barrington, Massachusetts.

Leg Band	Age	Sex	Weight Kg (lbs.)	Capture Site	Capture Date	Release Date	Remarks
188	Ad.	F	4.5 (10.0)	France Brook	9 March	10 March	
189	Ad.	F	4.6 (10.2)	France Brook	9 March	-	Died enroute.
190	Ad.	F	4.4 (9.8)	France Brook	9 March	10 March	
191	Ad.	F	4.6 (10.2)	France Brook	9 March	10 March	Instrumented; found dead 27 March.
192	Imm.	F	3.4 (7.5)	Wolf Run	10 March	11 March	
107	Imm.	M	4.7 (10.3)	Wolf Run	10 March	11 March	
108	Imm.	M	*	Wolf Run	10 March	11 March	
109	Ad.	M	6.5 (14.5)	France Brook	16 March	17 March	

* Not weighed.

State Forest personnel, Natural Resource officers, and the local Audubon Society were notified directly of the stocking and a news release was sent to local newspapers. No reports of the turkeys have been received since their release except that an adult hen (No. 191), instrumented by Tzilkowski, was found dead near Monument Mountain Reservation, about 1.5 kilometers west of the release site, on 27 March. Cause of death was undetermined, but presumed to be shock.

Since seven turkeys is minimal for a successful stocking, efforts are being continued to secure more wild-trapped birds in order to complete the stocking of Beartown, and hopefully to make a release in Hawley State Forest, Hawley, Franklin County.

C.

Release Sites

Horse Mountain Area. Five turkeys were released on Horse Mountain in Hatfield during September 1971. The area is private land, extensively wooded with red and white oak, with hemlock-northern hardwoods interspersed along the steeper slopes. Apple trees and blackberry bushes are present bordering the old field on the east slope where the turkeys were released. Intermittent branches of Running Brook arise from the east slopes and springs and seeps are common. To the north, Horse Mountain adjoins Chestnut Mountain in Whately and from there extensively wooded areas in Williamsburg and Conway.

Erving State Forest. Erving State Forest consists of 1934 hectares (4779 acres) in the Towns of Erving, Orange, Warwick and Northfield. The norther portion (north of Millers River) consists of mixed hardwoods (red, scarlet, white oak, beech, some sugar maple) with intermittent stands of white pine. The steeper slopes and area near Laurel Lake hold dense thickets of laurel, unsuitable for turkeys. Choke cherry, dogwoods, viburnums, and high-bush blueberry are present along road edges and as a scattered understory. The southern portion consists of several 81-162 hectare (200-400 acre) tracts of mixed hardwoods, and adjoins the Quabbin Reservation through wooded tracts in Leverett and Shutesbury. Several streams and swamps are present throughout both sections of the area. No turkey releases are planned for this area at the present time.

Recommendations: 1. Although Quabbin-strain turkeys recently appear to have improved in wildness and dispersal tendency, they are still markedly inferior to wild-trapped silvestris. Recent successes in Vermont using New York State turkeys indicate that our restoration efforts can be accelerated using similar stock. The seven New York turkeys obtained last winter are a valuable incentive towards statewide restoration; however, they do represent a minimal contribution for a successful release. Coordination efforts between the Division and the New York State Department of Environmental Conservation should be directed towards securing enough wild birds to complete the Beartown release. If additional turkeys are available, a second release should be made on the Hawley State Forest.

2. Consequent upon success of the Beartown release, turkeys from this area should be used in subsequent in-state trap and transfer effort. Initial efforts using this strain should be restricted to the area west of the Connecticut River to prevent mingling with present flocks in New Salem, Barre, and Douglas. Future releases of Quabbin-strain turkeys should be curtailed pending evaluation of the success of the Beartown release.

Acknowledgments: I extend my appreciation to Messrs. John Proud, Stephen Clarke, Fred Evans and William Shirey of the New York State Department of Environmental Conservation for their assistance in securing wild-trapped turkeys; to Dr. William G. Sheldon, Dr. Wendell Dodge, and Mr. Walter M. Tzilkowski of the Massachusetts Cooperative Wildlife Research Unit for their continuing cooperation and advice regarding the Beartown and Quabbin flocks; and to Mr. Francis Cole of Hatfield for approving the release on his property at Horse Mountain.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges
Superintendent

Prepared by _____
James E. Cardoza
Assistant Game Biologist

Date _____

JUN 5 1974

PERFORMANCE REPORT

Commonwealth of Massachusetts

State: Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Title: Game Population Trend and Harvest Survey

Job No.: IV-2 Job Title: Evaluation of Habitat Utilized by Transplanted Wild Turkeys

Period Covered: 7 February 1972 through 31 May 1972

Objectives: Habitat utilized by transplanted wild turkeys will be evaluated in relation to total habitat available, with special reference being given to habitats selected for winter roosting, winter-feeding, nesting and brood-rearing.

Procedures: Wild trapped turkeys obtained from the New York Department of Environmental Conservation, Division of Fish and Wildlife will be released in western Massachusetts. Depending on the total number of birds trapped, two releases will be made: one in Hawley State Forest, Franklin County, the other in Beartown State Forest, Berkshire County. Approximately twelve birds in the ratio of two hens per tom will be placed on each area in late February or early March. All turkeys will be marked with individually identifiable patagial wing tags. The hens will be equipped with back-mounted radio telemetry units.

Data collection will include: monitoring the daily movements of the birds, accounting for mortalities whenever possible; description of habitat characteristics of areas used for winter roosting, winter feeding, nesting and brood rearing; and evaluating nesting attempts and nesting success. Comparisons will be made between habitat utilized in relation to total available habitat to isolate, if possible, critical elements within the habitat required or preferred by wild turkeys. These criteria will be utilized in the evaluation of future proposed release sites.

Findings:

Seven wild turkeys; three adult females, one immature female, one adult male, and two immature males were live-trapped in Allegheny State Park, Cattaraugus County, New York with the assistance of New York Division of Fisheries and Wildlife personnel and were released in Beartown State Forest, Berkshire County, Massachusetts. Release dates were March 10 and 11 (6 birds) and March 17 (one bird, the adult male). One adult female, instrumented with a telemetry unit was found dead on March 27. The carcass showed no signs of violence. Cause of death was presumed to be shock.

Loss of the instrumented bird precluded radio tracking the released birds. Additional observations were made throughout the winter months in an effort to locate the flock, but without success.

Attention was turned to the development of a lighter, more powerful telemetry unit, and the improvement of the harness used to secure the unit to the turkey.

Recommendations: Additional wild turkey stock should be secured, if possible, from New York State to improve the chances of survival of the Beartown flock. At least half the adult females should be equipped with telemetry units in the future to improve the chances of tracking the dispersion of released birds and their subsequent movement. Further refinement of the telemetry unit and its harness should be developed.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved by _____
Colton H. Bridges
Title _____
Superintendent

Prepared by _____
Warren W. Blandin
Chief of Wildlife Research

Date _____

EA 32.3:

W-35-R-14/VI-1

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-14 Project Title: Game Population Trend and Harvest Survey

Job No.: VI-1 Job Title: Black Bear Population Dynamics

Period Covered: 1 June 1971 to 31 May 1972

Summary: Applications for bear hunting permits were received from 200 sportsmen. No bear were taken during the legal season. New reports of 16 observations involving 28 bear were received from five counties. Considerations are presented concerning future bear management activities.

Objectives: To define the range of the black bear in Massachusetts and to determine its population characteristics and rate of harvest by hunting.

Procedures: Current bear hunting regulations include mandatory reporting and tagging of bear. Bear checking stations were maintained daily during bear week at three locations (Birch Hill, Templeton; Montague Fish Hatchery, Montague; Western Wildlife District Headquarters, Pittsfield). Station personnel were required to affix a metal game seal to the bear, and to record the following information: town of kill, date killed, sex and weight of bear, and method of kill. Successful hunters would subsequently be contacted by the project leader and the bear's carcass examined and a tooth removed for sectioning.

The Information and Education Section issued periodic news releases asking for reports of black bear. District personnel, Natural Resource officers, and University of Massachusetts cooperators also assisted in reporting bear sightings.

Findings: Black bear hunting permit applications were received from 200 (214 in 1970) individuals during the 1971 season. Postage account limitations precluded sending questionnaires to these applicants.

For the second year in a row, no bear were taken during the legal season.

New reports of bears during 1971-1972 included fifteen sightings and one report of tracks, totaling 28 bear in twelve towns. A few reports, indefinite as to date, location, and observer, are not included. Reports are presented by county for 1971 and 1972 in Table 1. One cooperator reported finding bear dens during past years in northern Berkshire County.

Table 1. Reports of Black Bear by County, 1952-1972.

County	1952-1971	1972	Total (Percent)
Berkshire	46	5	51 (31.7%)
Franklin	64	6	70 (43.5%)
Hampden	5	1	6 (3.7%)
Hampshire	24	3	27 (16.3%)
Worcester	<u>6</u>	<u>1</u>	<u>7</u> (4.3%)
	145	16	161 (100.0%)

Recommendations: 1. Black bears are a minimal, yet integral, part of the mammalian fauna of Massachusetts. Possible overexploitation by casual hunters has been reduced by prohibiting bear hunting during deer week. Preliminary investigations (W-35-R-13) indicate that bears and bear hunters are probably present in Massachusetts in greater numbers than hitherto suspected. Correspondence received during the period preceding the 1970 bear hearing indicates that protectionists stand forth in number also. Future considerations concerning black bear should recognize these developments. The following points should be included: (a) maintenance of a bear population sufficient to sustain limited sport hunting, and satisfy environmental aesthetes; (b) public information efforts should be continued to reconcile conflicts between sportsmen and protectionists; and (c) evaluation of the bear population and distribution should be continued to minimize conflicts between bears and humans. Such conflicts, whether depredations on a beehive or raids on a camper's food box, are increasingly possible as urbanization continues and more vacationers flock to the forests. Careful consideration of points (a) and (b) above may help reduce difficulties due to these potential conflicts.

2. Bear permit applicants in 1972 should be contacted by postal questionnaire to determine their interest and time expenditure in bear hunting.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Assistant Game Biologist

Date _____

State: Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: I-2 Job Title: Statewide Beaver Harvest

Period Covered: 1 November 1972 to 31 May 1973

Summary: A total of 1,674 beaver were trapped by 105 trappers in 92 towns during the 1972-1973 beaver season. This record take is 316 more than last season, and 600 more than a ten-year (1963-1972) average. Berkshire and Franklin counties together yielded 967 beaver (54.1% of harvest). For the second season in a row, the take west of the Connecticut River increased, and that east of the river decreased.

Over one-third (37.8%) of the beaver were taken in the first two weeks of the 15-week season. The average pelt price of \$20 coupled with the high harvest, produced a record harvest valuation of \$33,480.

Objectives: To determine the statewide harvest of beavers by trappers.

Procedures: Each successful beaver trapper is required by law to present his pelts to an official checking station for tagging and recording of data. Six stations are maintained for two days at the close of the season. Pelts are tagged with locking metal game seals and harvest data are recorded on mimeographed forms and subsequently tabulated by month trapped, town and county trapped in, and type of trap used.

Findings: The 1972-1973 beaver trapping season extended for 15 weeks from 15 November 1972 to 1 March 1973. During this period, trappers took 1,674 beaver. This take was 316 more than last season and 600 more than a ten-year (1963-1972) average, representing the largest harvest since beaver trapping was initiated in 1952 (Table 1). One hundred five trappers (94 in 1971-1972) took at least one beaver for a mean harvest of 15.9 beaver per trapper (extremes: 1 to 119).

Beaver colonies located in 92 towns contributed to the 1972-1973 harvest (Figure 1). The fifteen towns with the largest individual seasonal harvest are listed in Table 2.

Table 1. Beaver season length, number of trappers, harvest total and harvest valuation: 1952-1973.

<u>Year</u>	<u>Season</u>	<u>Season (Weeks)</u>	<u>Number Successful Trappers</u>	<u>Avg. Harvest per Trapper</u>	<u>Harvest</u>	<u>Number Towns</u>	<u>Average Pelt Price</u>	<u>Harvest Value</u>
1952	-	4	-	-	63	-	\$15.87	\$ 1,000
1953	-	4	-	-	83	-	\$14.46	\$ 1,200
1954	-	6	-	-	191	-	\$11.00	\$ 2,100
1955	-	6	-	-	181	-	\$16.57	\$ 3,000
1956	-	6	-	-	135	-	\$10.00	\$ 1,350
1957	1 Feb. to 31 Mar.	8	32	8	261	55	\$ 9.00	\$ 2,349
1958	1 Feb. to 31 Mar.	8	36	7	252	57	\$12.00	\$ 3,024
1959	1 Feb. to 31 Mar.	8	33	10	323	42	\$12.00	\$ 3,876
1960	1 Feb. to 31 Mar.	8	39	8	296	52	\$15.00	\$ 4,440
1961	1 Jan. to 15 Apr.	15	66	12	825	77	\$15.00	\$ 12,375
1962	1 Jan. to 31 Mar.	13	63	11	669	76	\$12.00	\$ 8,028
1963	1 Jan. to 31 Mar.	13	83	7	589	63	\$13.00	\$ 7,657
1964	15 Dec. 63 to 15 Apr. 64	17	79	19	1,525	94	\$13.00	\$ 19,825
1965	15 Dec. 64 to 15 Apr. 65	17	81	15	1,196	91	\$11.00	\$ 13,156
1966	15 Dec. 65 to 15 Apr. 66	17	88	16	1,445	97	\$15.00	\$ 21,675
1967	15 Dec. 66 to 15 Apr. 67	17	74	14	1,040	96	\$15.00	\$ 15,600
1968	15 Dec. 67 to 15 Apr. 68	17	93	15	1,425	-	\$16.00	\$ 22,800
1969	15 Dec. 68 to 15 Apr. 69	17	80	13	1,052	93	\$20.00	\$ 21,040
1970	15 Dec. 69 to 1 Mar. 70	11	64	9	605	83	\$13.00	\$ 7,865
1971	15 Dec. 70 to 1 Mar. 71	11	63	8	509	79	\$10.00	\$ 5,090
1972	15 Dec. 71 to 1 Mar. 72	11	94	14	1,358	93	\$17.00	\$ 23,086
1973	15 Nov. 72 to 1 Mar. 73	15	105	16	1,674	92	\$20.00	\$ 33,480
Mean (1952-1973)		11.3	69	11.9	713.5	77.5	\$13.81	\$ 10,637
Total (1952-1973)					15,697			\$234,016

- = No data

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY

CITY AND TOWN LINES

Figure 1.
Beaver Trapped, by Towns, 1972-1973.
Total 1674 beaver.

Table 2. Beaver harvest for fifteen towns: 1969-1973 seasons.

<u>Town</u>	<u>1968-69</u>	<u>1969-70</u>	<u>1970-71</u>	<u>1971-72</u>	<u>1972-73</u>
Ashfield	20	14	2	53	50
Becket	41	20	10	51	57
Blandford	10	22	16	29	78
Chesterfield	14	24	12	16	18
Granville	12	23	19	19	49
New Marlboro	32	14	1	52	69
New Salem	11	10	27	10	56
Otis	36	18	14	64	72
Petersham	52	18	27	58	57
Sandisfield	49	13	8	52	6
Warwick	43	8	12	19	45
Washington	9	27	7	55	27
Wendell	40	20	7	10	38
Winchendon	23	16	30	45	57
Worthington	7	30	14	52	54

Table 3. Beaver harvest by county, 1971-1972 and 1972-1973.

<u>County</u>	<u>1971-1972</u>			<u>1972-1973</u>		
	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>
Berkshire	483	35.6	1	461	27.5	1
Essex	12	0.9	8	15	0.9	7
Franklin	301	22.1	2	446	26.6	2
Hampden	114	8.4	5	227	13.6	5
Hampshire	208	15.3	3	253	15.1	3
Middlesex	38	2.8	6	35	2.1	6
Plymouth	16	1.2	7	5	0.3	8
Worcester	186	13.7	4	232	13.9	4
Totals	1,358	100.0		1,674	100.0	

Table 4. Beaver harvest by month, 1971-1972 and 1972-1973

<u>Month</u>	<u>1971-1972</u>		<u>1972-1973</u>	
	<u>No. Beaver</u>	<u>Percent</u>	<u>No. Beaver</u>	<u>Percent</u>
November	Season Not Open		633	37.8
December	722	53.2	544	32.5
January	469	34.5	356	21.3
February	167	12.3	141	8.4
Totals	1,358	100.0	1,674	100.0

The western region of the state continues to provide the majority of the beaver harvest. For the second season in a row, the take in the western portion increased and that in the eastern region decreased from the previous season. During the 1972-1973 season, 1,148 beaver (68.6%) were taken west of the Connecticut River, and 526 (31.4%) were taken east of the river, as opposed to 920 (67.7%) and 242 (32.3%) in 1971-1972.

During the past season, Berkshire and Franklin counties together yielded over half (967, 54.1%) of the total harvest. Hampden, Hampshire and Worcester counties contributed another 732 beaver (42.6%), with three other counties comprising the small remainder (Table 3). Six counties reported no beaver taken. County rankings remain unchanged from the past season, although the Berkshire County harvest declined by 8.1 percent and that of Franklin and Hampden increased by 4.5 percent and 5.2 percent respectively.

Success was greatest, as expected, during the initial weeks of the season since the open water facilitated trapping. Over one-third (37.8%) of the beaver were taken in the initial two weeks (15-31 November) of the season with success decreasing each month thereafter (Table 4).

As in past years, approximately two-thirds (1,163, 69.5%) of the harvest was taken in the Conibear trap.

In spite of a supposed decrease in pelt quality, due to the early opening of the season, the average pelt price, \$20, equalled the record established in 1969. Thus, the total harvest valuation (\$33,480) was the highest yet recorded for Massachusetts.

Recommendations: Continue tagging of beaver pelts and recording of harvest data in 1974, using the same methods as for the current segment.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research & Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by: _____
James E. Cardoza, Assistant Game Biologist

Date _____

MASS.

EA32.3:W-35-R-15/II-1

JOB PROGRESS REPORT

Government Documents
Collection
MAY 2 1973
University of Massachusetts

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: II-1 Job Title: Statewide Deer Harvest

Period Covered: 6 November 1972 to 9 December 1972

Summary: The archery season was extended to three weeks, 6 November to 25 November 1972. Archers reported 76 deer harvested, 49 males and 27 females.

During the shotgun season, 4 December to 9 December 1972, successful hunters reported 1,455 males and 760 females for a total harvest of 2,215 deer. The top deer-producing counties were Berkshire, Franklin, and Dukes. Worcester County, with the largest acreage of deer range, has dropped to fifth rank of deer-producing counties. The first and last days of the six-day shotgun season had the largest percent of deer reported. Mid-week hunting pressures were reduced. Harvest data per deer management unit were compiled and are presented in this report. There was an increase of 500 antlerless permit applications in 1972. The success ratio of one in six for antlerless permit holders remained the same for all types of permits per deer management unit except for Nantucket. On Nantucket the hunter success ratio changed from one in three to one in four in 1972.

Background: An antlerless deer hunting permit system was initiated in 1967 for harvesting of deer during the shotgun deer week. Button bucks and females were taken only by permit holders. All males with antlers three inches or longer were legal game for all licensed hunters.

The shotgun deer season takes place during the first full week in December. Three deer management units have been established. The number of sportsman antlerless permits per unit in 1972 was as follows: Nantucket, 400; Martha's Vineyard, 600; and the mainland, 4000. Deer hunting regulations require all hunters to report their deer within 24 hours to an official Division deer checking station.

The archery season was increased from a two-week period to a three-week period, 6 November through 25 November 1972.

Hunting is not allowed on Sundays.

Objectives: To determine the annual harvest of deer in Massachusetts.

Findings: A ten-year summary of the annual deer harvest per town per county from 1963 through 1972 is presented in Table 1. Hunters reported the taking of a deer via a written report to the Director from 1963 through 1965. From 1966 through 1972 the reporting by hunters has been on a compulsory basis.

During the 1972 shotgun deer season, 4 December through 9 December, successful hunters reported harvesting 2,215 deer. Of these, 1,455 were males and 760 were females.

Successful archers reported taking 76 deer during the three-week period, 6 November through 25 November. Of these deer, 49 were males and 27 were females.

The 1972 deer harvest by archers is presented in Table 2. In order of importance Berkshire, Franklin, and Hampshire Counties were the best deer-producing counties for the archers.

A summary of the 1972 shotgun deer harvest by county from 1969 through 1972 is presented in Table 3. Berkshire, Franklin, and Dukes Counties have remained the top-producing counties for the past two years.

Table 4 presents a summary of the deer harvest by day during the shotgun season, 4 December through 9 December 1972 and the archery season, 6 November through 25 November 1972.

It is quite evident from the harvest data presented in Table 4 that the harvest is highest on the first and last days of the six-day shotgun season. The hunting pressure drops during the middle four days with Thursday having the smallest percent of the harvest reported. It is interesting to note that 632 or 43 percent of the total males were taken on Monday and 281 or 19 percent were reported on Saturday. The female harvest on the first and last days of the shotgun season were approximately the same with 225 or 30 percent on Monday and 235 or 31 percent on Saturday.

With the exception of Friday, 17 November, archers harvested deer on each day of the three-week archery season. The last three days of the archery season include the long Thanksgiving holiday weekend. This probably accounts for the ten deer harvest on the Friday (24 November) following Thanksgiving Day and a total of 21 deer harvested for the long weekend.

Table 5 presents a summary of the Massachusetts deer harvest and the hunter success per type of antlerless permit.

Table 2.

A Summary of the 1972 Massachusetts Archery Season Deer Harvest by Sex, Town and County, November 6-25, 1972.

<u>County</u>	<u>Town</u>	<u>Male</u>	<u>(BB)</u>	<u>Female</u>	<u>Total</u>
Barnstable	Barnstable	3	(1)		3
	Orleans	1			1
	Provincetown	1		1	2
	Truro	1			1
	Wellfleet	$\frac{1}{7}$	$\frac{(1)}{7}$	$\frac{1}{1}$	$\frac{1}{8}$
Berkshire	Alford			2	2
	Becket	1			1
	Egremont	2			2
	Hancock	1			1
	Lanesboro	1			1
	Lee	1			1
	Mt. Washington	1			1
	New Marlboro	1			1
	Sandisfield	1			1
	Sheffield	1	(1)	1	2
	Stockbridge	1	(1)		1
	Tyringham	1	(1)		1
	West Stockbridge	4	(2)	1	5
	Williamstown	$\frac{3}{19}$	$\frac{(1)}{(6)}$	$\frac{1}{5}$	$\frac{4}{24}$
Dukes	Chilmark			1	1
	Edgartown	2			2
	West Tisbury	$\frac{1}{2}$		$\frac{1}{2}$	$\frac{1}{4}$
Essex	Georgetown			1	1
Franklin	Bernardston	1		2	3
	Colrain			1	1
	Conway	1	(1)		1
	Deerfield	1			1
	Leyden			1	1
	New Salem	1			1
	Shelburne	1		2	3
	Shutesbury	2		1	3
	Wendell	$\frac{1}{7}$	$\frac{(1)}{(1)}$	$\frac{1}{8}$	$\frac{1}{15}$
Hampden	Blandford			1	1
	Ludlow			$\frac{1}{2}$	$\frac{1}{2}$
Hampshire	Belchertown	2	(1)	1	3
	Chesterfield			1	1
	Huntington	2	(1)		2
	Pelham	0		2	2
	Ware	$\frac{2}{1}$	$\frac{(2)}{(2)}$	$\frac{0}{4}$	$\frac{2}{10}$
Nantucket	Nantucket	5		3	8
Worcester	Grafton	1			1
	Petersham	1			1
	Southbridge			1	1
	Upton	$\frac{1}{3}$		$\frac{1}{1}$	$\frac{1}{4}$
TOTALS		49	(10)	27	76

Table 3. Summary of the Massachusetts Deer Harvest by Sex per County and Rank of Harvest for the 1969, 1970, 1971 and 1972 Shotgun Seasons.

<u>County</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>	<u>Rank 1972</u>	<u>Rank 1971</u>	<u>Rank 1970</u>	<u>Rank 1969</u>
Barnstable	78	33	116	8	7	8	8
Berkshire	423	160	583	1	1	1	1
Bristol	3	2	5	12	12	12	12
Dukes	140	126	266	3	3	7	7
Essex	16	1	17	9	10	9	11
Franklin	319	201	520	2	2	2	2
Hampden	147	54	201	4	5	6	5
Hampshire	114	49	163	6	8	4	4
Middlesex	3	5	8	11	9	10	10
Nantucket	66	61	127	7	6	5	6
Norfolk	0	0	0	13	13	13	13
Plymouth	12	0	12	10	11	11	9
Suffolk	0	0	0	13	13	13	13
Worcester	134	63	197	5	4	3	3

Table 4. A Summary of the Massachusetts Deer Harvest by Day of Harvest, 4 December through 9 December 1972.

	<u>Shotgun Season</u>						
	<u>Mon. Dec. 4</u>	<u>Tues. Dec. 5</u>	<u>Wed. Dec. 6</u>	<u>Thurs. Dec. 7</u>	<u>Fri. Dec. 8</u>	<u>Sat. Dec. 9</u>	<u>Total</u>
	<u>Both Sexes</u>						
Total Harvest	857	258	209	161	214	516	2,215
Percent	38.69	11.65	9.44	7.27	9.66	23.30	
	<u>Males</u>						
Total Harvest	632	172	134	117	119	281	1,455
Percent	43.44	11.82	9.21	8.04	8.18	19.31	
	<u>Females</u>						
Total Harvest	225	86	75	44	95	235	760
Percent	29.61	11.32	9.87	5.79	12.50	30.92	

Table 4. (Continued)

Archery Season

	<u>November 6-11</u>			<u>November 13-18</u>			<u>November 20-25</u>			<u>Total per Day</u>
Mon.	<u>M</u>	<u>F</u>	<u>Total</u>	<u>M</u>	<u>F</u>	<u>Total</u>	<u>M</u>	<u>F</u>	<u>Total</u>	
Mon.	4	4	8	1	-	1	-	1	1	10
Tues.	3	1	4	-	2	2	2	2	4	10
Wed.	2	1	3	5	1	6	2	1	3	12
Thurs.	1	1	2	3	-	3	3	1	4	9
Fri.	1	-	1	-	-	-	7	3	10	11
Sat.	<u>-</u>	<u>4</u>	<u>4</u>	<u>6</u>	<u>4</u>	<u>10</u>	<u>5</u>	<u>2</u>	<u>7</u>	<u>21</u>
	11	11	22	15	7	22	19	10	29	73*

* The archery harvest was 76 deer. Dates of harvest on three (3) deer were omitted.

Table 5. A Summary of the Massachusetts Deer Harvest and the Hunter Success by Type of Antlerless Permit, 1967 through 1972.

1972 Archery Season (Nov. 6 - Nov. 25, 1972)

Males	49 (10 BB)
Females	<u>27</u>
Total	76

Summary of Archery Kill, 1967 through 1972

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Males	17	21	27	24	26	49
Females	<u>4</u>	<u>13</u>	<u>10</u>	<u>12</u>	- <u>10</u>	<u>27</u>
Totals	21	34	37	36	36	76

1972 Shotgun Season (Dec. 4-9, 1972)

Males	1,455
Females	<u>760</u>
Total	2,215

1972 Kill Breakdown per Management Unit

	<u>Mainland</u>	<u>Nantucket</u>	<u>Martha's Vineyard</u>	<u>Total</u>
Males, Adult	1,105	49	102	1,256
Males, Button bucks	144	17	38	199
Females	<u>573</u>	<u>61</u>	<u>126</u>	<u>760</u>
Totals	1,822	127	266	2,215

Summary of the Shotgun Statewide Deer Harvest, 1967 through 1972

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Males	937	1,083	1,424	1,605	1,359	1,455
Females	<u>235</u>	<u>310</u>	<u>585</u>	<u>764</u>	<u>889</u>	<u>760</u>
Total	1,172	1,393	2,009	2,369	2,248	2,215

Statewide Summary of Archery and Shotgun Harvest, 1967 through 1972

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Males	954	1,104	1,451	1,629	1,385	1,504
Females	<u>239</u>	<u>323</u>	<u>595</u>	<u>776</u>	<u>899</u>	<u>787</u>
Total	1,193	1,427	2,046	2,405	2,284	2,291

Summary of Antlerless Deer Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
No. Applications (rounded numbers)	28,000	24,000	32,000	35,000	37,500	38,000
<u>Permits Issued:</u>						
Sportsman	2,000	2,000	4,000	6,000	6,000	4,000
Farmer*	243	331	295	347	270	326
Nantucket	-	-	400	400	400	400
Martha's Vineyard	-	-	-	-	600	600
Totals	<u>2,243</u>	<u>2,331</u>	<u>4,695</u>	<u>6,747</u>	<u>7,270</u>	<u>5,326</u>

Permit Harvest Statewide, Landowner-Farmer

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Males	2	21	17	25	20	14
Females	<u>20</u>	<u>44</u>	<u>45</u>	<u>34</u>	<u>26</u>	<u>42</u>
Total	22	65	62	59	46	56

Harvest by Sportsmen's Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Deer (both sex)	279	356	787	1,057	1,268	1,010

Breakdown Statewide of Harvest by Antlerless Permits

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Antlered male	26	50	140	137	172	107
Button buck	40	61	124	215	253	199
Female	<u>235</u>	<u>310</u>	<u>585</u>	<u>764</u>	<u>889</u>	<u>760</u>
Totals	301	421	849	1,116	1,314	1,066

Antlerless Permit Success Ratio (rounded)

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Farmer*	1-11	1-5	1-5	1-6	1-6	1-6
Sportsman	1- 9	1-6	1-6	1-6	1-6	1-6
Mainland					1-6	1-6
Nantucket	(101 permit deer, 40 M., 61 F.)				1-3	1-4
Martha's Vineyard	(167 permit deer, 56 M., 111 F.)				1-4	1-4

*Farmer-Landowner

Apparently the added six days of the archery season and possibly the increased cost of the archery stamp stimulated the archers during the 1972 season. The average reported archery kill from 1968 through 1971 was 36 deer per year. The 1972 archery harvest was 76, an increase of 40 deer or 111 percent above the four-year average.

During the 1972 shotgun season, 4 December through 9 December, hunters reported 2,215 deer statewide, 1,455 males and 760 females. On the mainland, 1,105 adult males, 144 button bucks and 573 females were reported. On Nantucket 49 adult males, 17 button bucks and 61 females were recorded. The Martha's Vineyard hunters reported 102 adult males, 38 button bucks and 126 females.

The combined statewide archery and shotgun reported harvest was 2,291 deer. Of these 1,504 were males and 787 were females.

The 1972 applications for antlerless permits increased to 33,000 (rounded number), approximately 500 applications more than were received in 1971.

The number of sportsman permits was reduced from 6,000 in 1971 to 4,000 permits in 1972 in an attempt to reduce the harvest of females and button bucks and thereby increase the number of legal bucks in the harvest of 1974. The farmer-landowner antlerless permits increased from 270 permits in 1971 to 326 permits in 1972. Nantucket and Martha's Vineyard antlerless permits remained the same as previous years with 400 and 600 permits issued respectively.

As expected with the reduction of permits the harvest of deer by sportsman permittees decreased from 1,268 deer in 1971 to 1,010 deer reported in 1972 or 258 deer less than the 1971 season.

A review of the statewide deer harvest per antlerless permit shows that the number of deer harvested by permit holders was reduced in 1972. The antlered male harvest decreased 65 deer from 172 in 1971 to 107 in 1972. There were 54 less button bucks reported in 1972 (199) than the 253 taken in 1971. The largest reduction was in the female segment with 760 reported in 1972 compared to 889 in 1971, or 129 less females in 1972.

The success ratio per type of permit remained the same as previous years for all types of permits in all deer management units except for Nantucket. The success ratio was one in six for sportsman and farmer-landowner permits on the mainland and Martha's Vineyard. The ratio changed from one in three to one in four on Nantucket during the 1972 season.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and the role of the accounting department in ensuring the integrity of the financial statements. It also highlights the need for regular audits and the importance of transparency in financial reporting.

2. The second part of the document focuses on the management of human resources, including recruitment, training, and performance evaluation. It emphasizes the importance of having a clear job description and a fair compensation system to attract and retain top talent.

3. The third part of the document addresses the issue of risk management, particularly in the context of financial markets. It discusses the various risks faced by the company and the strategies used to mitigate them, such as diversification and hedging.

4. The fourth part of the document deals with the company's marketing strategy, including the identification of target markets and the development of effective promotional campaigns. It also discusses the importance of customer relationship management and the role of the sales department in driving revenue growth.

5. The fifth part of the document provides a summary of the company's overall performance and the key challenges it faces. It also outlines the company's vision for the future and the steps it is taking to achieve its long-term goals.

[illegible]

Recommendations: It is recommended that the Division of Fisheries and Game personnel continue to compile and report deer harvest data.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist

and

William J. Minior, Assistant

Date _____

JOB PROGRESS REPORT

Government Documents
Collection
MAY 2 1973

State Massachusetts

Cooperators Massachusetts Division of Fisheries and Game

(University of Massachusetts)

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: II-2 Job Title: Non-hunting Deer Mortality Investigations

Period Covered: 1 January 1972 to 31 December 1972

Summary: Natural Resource Officers reported 453 deer mortalities. One hundred ninety-six were males, 230 were females. No sex was reported for 27 deer. The two greatest causes of non-hunting deer mortalities were motor vehicles (321) and illegal kills (44). They accounted for 80.6 percent of the reported mortalities. Other causes of deer mortalities in order of importance were as follows: dogs (41); other and unknown (35); fences (6); drownings (5); and crop damage (1). The 41 deer reported killed by dogs in 1972 possibly reflects favorable winter conditions. Two hundred nineteen deer were reported killed by dogs in 1971.

The uneven proportion of male (49) to female (19) mortalities during the month of November can be attributed to the increased male activity during the rut period.

There was a 34.7 percent decrease in non-hunting deer mortalities (453) for 1972 as compared to 1971 (694).

The effects of the antlerless permit hunting system were evident in the adjusted sex ratio of the non-hunting deer mortalities with 208 males to 245 females or 45.9 percent males to 54.1 percent females.

As in 1971, the top ranking counties for non-hunting deer mortalities were Berkshire, Barnstable and Franklin.

Background: Natural Resource Officers investigated and filed reports of deer mortalities statewide. A duplicate of each report was compiled and tabulated by project personnel. Deer concentrations and/or wintering areas were observed by project personnel. Dead deer surveys were conducted in areas of deer concentrations.

Objectives: To determine the annual non-hunting decimating factors of the Massachusetts deer herd.

Government Documents
Collection
NOV 14 1973
University of Massachusetts

Findings:

A summary of the sex classes and the causes of Massachusetts deer mortalities for January 1972 through December 1972 is presented in Table 1. A total of 453 deer mortalities were reported. Of these 196 were males, 230 were females. No sex was recorded for 27 deer. Motor vehicles continued to be the largest cause of reported deer mortalities (321 deer). The remaining 132 deer mortalities were reported as follows: illegal kills, 44; dogs, 41; other and unknown, 35; fences, 6; drownings, 5; and deer shot doing crop damage, 1.

Table 2 presents a six-year summary of deer mortalities and causes from 1967 through 1972.

A comparison of the total non-hunting deer mortalities in Massachusetts from 1967 through 1972 is presented in Table 3. The largest number of deer mortalities was reported in 1970 with 698 deer. Prior to 1970 the reported mortalities increased annually from 508 in 1967 to the peak of 698 deer in 1970. Since 1970 there has been a decrease annually starting from a half of one percent decline (694 deer) in 1971 to a 34.7 percent decrease (453) in 1972.

Table 4 presents a comparison of the sex ratio and the adjusted sex ratio of Massachusetts deer from 1 January 1972 through 31 December 1972. An adjusted total of 453 deer mortalities, 208 males and 245 females, was reported. The adjusted sex ratio was computed monthly on a percentage basis. The sex ratio was 85 (45.9 percent) males to 100 (54.1 percent) females. The sex ratio in 1971 was 94 (48.4%) males to 100 (51.6%) females.

The reported deer mortalities are presented in Table 5 which ranks counties according to deer mortality numbers. Berkshire, Barnstable, Franklin and Worcester, respectively, have remained the top deer mortality-producing counties for the past two years.

Reported deer mortalities for Norfolk (6), Bristol (6), and Plymouth (21) outnumbered the legal deer harvest reported in Job Progress Report W-35-R-15, Work Plan II, Job 1, which was Norfolk, 0; Bristol, 1; and Plymouth, 10.

A review of the data compiled in this report shows a decline in the number of reported mortalities during 1972 as compared to the previous five years. It is quite possible that the relatively open winter with snow conditions favorable to the deer effected the sharp reduction of the number of deer killed by dogs. It is felt by project personnel that the enforced eight-hour workday of the Natural Resource Officers has affected and will continue to affect the number of reported deer mortalities.

Table 1. A summary of Massachusetts deer mortalities reported by natural resource officers, 1 January through December 1972.

	<u>January</u>			<u>February</u>			<u>March</u>			<u>April</u>			<u>May</u>			<u>June</u>		
	<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>		
	M.	F.	Sex	M.	F.	Sex	M.	F.	Sex	M.	F.	Sex	M.	F.	Sex	M.	F.	Sex
Motor Vehicles	5	7		7	7	2	12	16	1	16	15	1	10	14		11	22	1
Dogs	4	1	2	5	1	3	5	6		1	3		1	1				
Illegal Kill	1			2		1			5									
Drowned	1																	
Fences	1						1			2			1					
Trains										1	1							
Crop Damage																		
Other and Unknown	3	2		3	3		18	22	1	1	2		11	15	1	1	22	1
Subtotal	14	10	2	17	11	6	41	(101)		21	21	6	27	(176)		35	(211)	
Total (Cum. Total)	26	(26)		34	(60)					48	(149)							
	<u>July</u>			<u>August</u>			<u>September</u>			<u>October</u>			<u>November</u>			<u>December</u>		
	<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>			<u>No</u>		
Motor Vehicles	10	21		9	10	1	5	8	3	8	21	1	39	12	1	9	16	
Dogs							1							1		3	4	
Illegal Kill				1	1			1		4			8	4		4	11	1
Drowned													1			1		
Fences						1												
Trains																		
Crop Damage																		
Other and Unknown	1																	
Subtotal	10	22		10	12	2	6	10	3	8	27	3	49	19	1	20	39	1
Total (Cum. Total)	32	(243)		24	(267)		19	(286)		38	(324)		69	(393)		60	(453)	
	<u>Cause</u>			<u>Male</u>			<u>Female</u>			<u>Unknown Sex</u>			<u>Total</u>					
Motor Vehicles				141			169			11			321					
Dogs				19			17			5			41					
Illegal Kill				16			21			7			44					
Drowned				4			1						5					
Fences				3			2			1			6					
Trains																		
Crop Damage							1						1					
Other and Unknown				13			19			3			35					
Totals				196			230			27			453					

Table 2. Five-year summary of deer mortalities of Massachusetts deer reported by natural resource officers, 1967 through 1972

<u>Cause</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>Total</u>	<u>5-yr. Avg.</u>	<u>1972</u>
Motor Vehicles	334	456	397	400	373	1,960	392	321
Dogs	62	74	166	204	219	725	145	41
Illegal Kills	48	29	39	25	39	180	36	44
Unknown Causes	41	31	51	38	41	202	40	35
Crop Damage and Legal Kills	4	7	2	14	4	31	6	1
All other Causes	<u>19</u>	<u>16</u>	<u>27</u>	<u>17</u>	<u>18</u>	<u>97</u>	<u>19</u>	<u>11</u>
Totals	508	613	682	698	694	3,195	639	453

Table 3. Comparison of total non-hunting deer mortalities of Massachusetts deer from 1967 through 1972.

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
No. of Deer	508	613	682	698	694	452
Percent Change		21%	11%	2%	-0.5%	-34.7%

Table 4. Comparison of actual number of deer mortalities by sex and adjusted data for 1972 Massachusetts deer

<u>Month</u>	<u>Male</u>	<u>Female</u>	<u>Unknown Sex</u>	<u>Total</u>	<u>Adjusted</u>	
					<u>Male</u>	<u>Female</u>
January	14	10	2	26	15	11
February	17	11	6	34	21	13
March	18	22	1	41	18	23
April	21	21	6	48	24	24
May	11	15	1	27	11	16
June	12	22	1	35	12	23
July	10	22		32	10	22
August	10	12	2	24	11	13
September	6	10	3	19	7	12
October	8	27	3	38	8	30
November	49	19	1	69	50	19
December	<u>20</u>	<u>39</u>	<u>1</u>	<u>60</u>	<u>20</u>	<u>40</u>
Total	196	230	27	453	208	245

* These data were reported by Natural Resource Officers.

** Hunting and crippling losses not included.

Adjusted sex ratio: 208 males : 245 females
85 males : 100 females
45.9% males : 54.1% females

Table 5. Massachusetts non-hunting deer mortalities ranked by county and cause, 1972 and 1971.

County	Auto	Dogs	Illegal Kill	Drowned	Fences	Trains	Crop Damage	Other and Unknown	1972		1971	
									Total	Rank	Total	Rank
Barnstable	60	1	1	2	1			7	72	2	93	2
Berkshire	60	12	13		1			6	92	1	175	1
Bristol	3		2					1	6	11	6	11
Essex	14		4		1			1	20	9	25	9
Franklin	54	6	4				1	1	66	3	90	3
Hampden	37	3	1	1				3	45	5	51	6
Hampshire	25	3	3					7	38	6	65	5
Middlesex	5	1						2	8	10	18	10
Norfolk	4			1	1				6	11	1	13
Plymouth	16	1	3		1				21	8	36	8
Suffolk									0	13	0	14
Worcester	27	14	5	1	1			7	55	4	85	4
Dukes	14		8						22	7	45	7
Nantucket	<u>2</u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u>2</u>	<u>12</u>	<u>4</u>	<u>12</u>
Totals	321	41	44	5	6		1	35	453		698	

Recommendations: It is recommended that deer project personnel continue to determine the annual non-hunting deer decimating factors of the Massachusetts deer herd.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist

and

William J. Minior, Assistant

Date _____

JOB PROGRESS REPORT

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: II-3 Job Title: Deer Fertility Studies

Period Covered: 1 January 1973 through 31 May 1973

Summary: Thirty-six (36) female deer mortality carcasses were collected from 1 January through 31 May 1973 by Natural Resource Officers and Division personnel. The deer were aged and autopsied to determine the frequency and state of pregnancy. Fetus and corpora lutea were examined and documented.

There were no significant changes in the mean reproductive rates for the eight-year period, 1966 through 1973. A summary of eight years of reproductive data (1966-1973) is as follows:

<u>Age at Parturition</u>	<u>Sample Size</u>	<u>No. Fawns Produced</u>	<u>1966-1973 Reproductive Rate</u>	<u>1973 Rate</u>
Yearling	160	40	1: .25 (100: 25)	1: .22 (100: 22)
Two years	93	131	1:1.41 (100:141)	1:1.33 (100:133)
Adults	199	344	1:1.73 (100:173)	1:1.72 (100:172)

The sex ratio of 126 males to 100 females was determined for the 43 fetuses examined. This was a disproportionate rate from the expected 106 males to 100 females.

Background: Carcasses of female deer mortalities were collected from 1 January through 31 May by Division of Fisheries and Game personnel and Natural Resource Officers. These carcasses were either brought directly to the Field Headquarters in Westboro or picked up by project personnel from various Division installations. Deer were autopsied by project personnel and fetuses and ovaries were removed for future examination.

The tooth replacement and/or wear technique was used to age the deer examined. Crown-rump measurement was used to determine fetus age.

Ovaries were sectioned longitudinally and the number of current corpora lutea were recorded. Corpora albicantia were also sought to help provide some idea of previous pregnancies.

Government Documents
Collection

NOV 14 1973

University of Massachusetts

Objectives: To determine the reproductive rate per age class of the Massachusetts deer herd.

Findings: Thirty-six (36) female deer were examined during the five-month period (1 January through 31 May 1973) covered by this report. All 36 deer were collected from the Mainland herd. The age of the deer presented in this report is the age at parturition. From 1 January through 31 May 1973 examination of 36 reproductive tracts found that:

1. Of the nine yearlings, two were pregnant and each were carrying a single fawn.
2. Eight of the two-year-old does were pregnant and carrying 12 fetuses.
3. Of the 18 adults, 17 were pregnant and carrying 31 fetuses. One adult doe examined in early January may have been recently bred. However, no embryos were in evidence.

Table 1 presents a comparison of the age composition and the percent per age class for the 1972 female deer harvest checked at Mainland biological deer check stations and the 36 non-hunting doe mortalities.

The percent per age class of the harvested and the non-hunting mortalities appear comparable except for the two and four-year-old classes. A higher percent of the non-hunting mortalities was reported in the two-year class compared to the reported hunting harvest. Hunters harvested 16.83 percent of the 3-1/2 year old does during the 1972 season. The non-hunting mortality data show only eight percent of this class were reported in 1973. Other degrees of variation were noted in the older age classes, but these may have been due to small sample sizes (Table 1).

The following reproductive rates were calculated for the 36 deer collected 1 January through 31 May 1973:

<u>Age Class</u>	<u>Number Females</u>	<u>Number Fetuses</u>	<u>Calculated Reproductive Rate</u>
Yearling	9	2	100 does : 22 fawns (1 : .22)
Two years	9	12	100 does : 133 fawns (1 : 1.33)
Adults	18	31	100 does : 172 fawns (1 : 1.72)

Table 2 presents an eight-year summary of reproductive data for the three age classifications from 1966 through 1973. The contribution of the yearlings per year is minimal averaging 0.24 fawns per year. The two-year-old class contributes an average of 1.40 fawns per pregnant doe per year. Most important is the adult class in which pregnant females average 1.72 fawns per year. Producing at a rate nearly seven times greater than the yearlings and nearly one third better than the two-year class, the adult segment plays a highly significant role in the total fawn production

Table 1. The age composition and percent per age class of harvested female deer checked at biological deer stations (mainland) December 1972 and thirty-six female deer mortalities collected January through May 1973.

<u>Age at Harvest</u>	<u>No. in Sample</u>	<u>Percent of Sample</u>	<u>Age at Parturition</u>	<u>No. in Sample</u>	<u>Percent of Sample</u>
6 mos.	90	29.70	1	9	25.00
1-1/2	57	18.81	2	9	25.00
2-1/2	56	18.48	3	6	16.67
3-1/2	51	16.83	4	3	8.33
4-1/2	22	7.26	5	3	8.33
5-1/2	14	4.62	6	1	2.78
6-1/2	10	3.30	7	3	8.33
7-1/2	2	.66	8-9	2	5.55
8-9-1/2	-	----	10+		
10-1/2	1	.33			

Table 2. A summary of the reproductive rate data per age class of thirty-six Massachusetts female deer mortalities, 1 January 1966 to 31 May 1973.

<u>Size</u>	<u>Sample Size</u>	<u>Pregnant</u>	<u>Yearlings</u>		<u>Annual Reproductive Rate</u>
			<u>Not Pregnant</u>	<u>No. Fawns</u>	
1966	16	9	7	11	0.69
1967	12	1	11	1	0.10
1968	14	2	12	2	0.14
1969	25	5	20	5	0.20
1970	37	4	33	4	0.11
1971	27	8	19	8	0.30
1972	21	7	14	7	0.33
1973	9	2	7	2	0.22
Total	161	38	123	40	2.09
Mean	20.13	4.75	15.38	5	0.24
			<u>Two-Year-Olds</u>		
1966	4	3	1	5	1.25
1967	10	9	1	12	1.20
1968	12	11	1	16	1.33
1969	16	16	0	23	1.44
1970	15	14	1	20	1.33
1971	16	15	1	26	1.63
1972	11	11	0	17	1.55
1973	9	8	1	12	1.33
Total	93	87	6	131	11.06
Mean	11.63	10.88	.75	16.38	1.40
			<u>Adults</u>		
1966	18	17	1	31	1.72
1967	10	10	0	20	2.00
1968	20	17	3	32	1.60
1969	35	32	3	63	1.80
1970	28	23	5	47	1.67
1971	41	39	2	70	1.71
1972	29	26	3	49	1.69
1973	18	17	1	31	1.72
Total	199	181	18	343	13.91
Mean	24.88	22.63	2.25	42.88	1.72

and therefore the overall size of the Massachusetts deer herd.

A summary of the 1973 corpora lutea and fetus counts is presented in Table 3. No analysis of the corpora lutea data will be made at this time because of the sample size and the necessary assumptions made to obtain the count. It was interesting to note that 50 percent of the two-year-old females were carrying twins and 66 percent of the adults were carrying twins. One adult was carrying a set of triplets.

A breakdown of the sex of the deer fetuses per age class is as follows:

<u>Age</u>	<u>No. Does</u>	<u>Sex of Fetus</u>		<u>Total No. Fetuses</u>
		<u>Male</u>	<u>Female</u>	
Yearling	2	2		2
Two years	7	6	4	10
Adults	<u>17</u>	<u>16</u>	<u>15</u>	<u>31</u>
	26	24	19	43

The sex ratio for the 1973 fetus collection (24 males:19 females) was 126 males to 100 females. This was somewhat disproportionate to the expected 106 males to 100 females. It is possible that the small sample size collected during the 1973 period tends to make the data uncertain or questionable.

During the five-month collection period (1 January through 31 May 1973) 88 female deer mortalities were reported by Natural Resource Officers. Of the 88 deer, only 36 females or 41 percent of the mortalities were collected and examined by project personnel. No deer were collected from Nantucket or Martha's Vineyard during the five-month period.

Recommendations:

It is recommended that Division personnel continue to collect, compile and report reproductive data pertaining to the Massachusetts deer herd.

It is recommended that project personnel present a deer project activity slide lecture at each of the Natural Resource Law Enforcement Districts before 1 January 1974. It is felt that such lectures might stimulate interest and increase personnel cooperation between the deer project and the field officers.

Table 3. A summary of the corpora lutea and fetus counts for pregnant yearling, two-year-old and adult female deer in Massachusetts, 1 January through 31 May 1973.

<u>Age</u>	<u>Number Pregnant Females</u>	<u>Number Corpora Lutea</u>	<u>Number Fetuses</u>	<u>Single</u>	<u>Twins</u>	<u>Triplets</u>
Yearlings	2	2	2	2		
Two years	8*	14*	12	4	4	
Adults	18	37**	32	4	12	1

* Ovaries missing but assumed that at least one current corpora lutea present per fetus.

** A three-year-old (at parturition) doe had two current corpora lutea and may have been in the early stages of pregnancy as no embryos were found.

It will be necessary to train some person or persons (NRO or MSPCA) to collect the lower jaw and complete reproductive tract of female deer mortalities occurring on Nantucket and Martha's Vineyard in order to determine the reproductive rates for the islands.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist
and
William J. Minior, Assistant

Date _____

JOB PROGRESS REPORT

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: II-4 Job Title: Management of the Massachusetts Deer Herd

Period Covered: 1 July 1972 to 30 June 1973

Summary: Sex and age data collected at ten biological deer checking stations were expanded for the three deer management units. The 1972 statewide shotgun deer harvest was 1,256 males and 760 females. There were 199 button bucks reported from 4 December through 9 December 1972.

The number of Mainland antlerless permits was reduced from 6,000 to 4,000 permits in 1972. The reduction should result in an increase of antlered males in 1974.

The 1972 harvest of deer per square mile of deer range on the Mainland remained unchanged at .3 deer. The Martha's Vineyard harvest per square mile increased from 2.5 deer in 1971 to 3.1 deer in 1972. On Nantucket Island, the harvest changed from 4.5 deer per square mile in 1971 to 3.5 deer in 1972.

The calculated statewide minimal deer population for 1972 was 11,336 deer.

Non-hunting deer mortalities are greater than the reported harvest in the Eastern group of counties.

The effect of the antlerless permit system of harvesting and controlling a deer herd is demonstrated on Nantucket Island.

There was a slight increase of adult buck harvest per square mile on the Mainland and Dukes County (Martha's Vineyard and the Elizabeth Islands). A planned reduction of the adult male harvest continued on Nantucket.

The number of adult females in the harvest decreased on the Mainland and Nantucket. There was a slight increase in the adult doe harvest in Dukes County.

The percent frequency of adult females to adult males decreased on the Mainland and Nantucket Island. The percent frequency of adult females to adult bucks increased in Dukes County.

Government Documents
Collection
NOV 14 1973

University of Massachusetts

Deer herd management recommendations are presented.

Background:

Starting in 1966, successful Massachusetts deer hunters have been required to bring their deer to an official deer checking station within 24 hours to be recorded. Deer checking stations have been operated by Division personnel since 1948. Ten check stations are operated as biological stations at which sex, age, weight and antler beam diameter data are collected.

In 1967, an antlerless permit deer hunting system was initiated to control the hunting pressure on the female segment of the deer herd. The deer project leader recommends the number of permits to be issued after examination of herd age composition, reproductive rates, annual removal rates, hunting pressure and the deer harvest per square mile of deer range.

Objectives:

To estimate the size of the deer herd in Massachusetts and to recommend management techniques.

Findings:

A summary of the sex and age composition collected at the biological deer check stations per deer management unit from 1967 through 1972 is presented in Table 1. This table includes the 1971 and 1972 biological data collected from Nantucket and Martha's Vineyard islands.

Figure 1 presents the Massachusetts deer harvest per square mile of deer range per county for the six-year period, 1967 through 1972. In Berkshire and Franklin counties .5 males were harvest per square mile of deer range, while the harvest in Barnstable, Hampden and Hampshire counties was .3 males per square mile of range. The Eastern group of counties consisting of 2271.6 square miles of deer range produced a reported harvest of 34 bucks.

Franklin County, with a harvest of .3 females per square mile of deer range, can be interpreted as a county with heavy hunting pressure. The highest deer harvest on the Mainland with .8 deer per square mile of deer range was found in Franklin County.

Because of the small reported harvest for the Eastern group, the data were deleted from Figure 1. The Eastern group, considered a deer management problem area, consists of the following counties: Bristol, Essex, Middlesex, Norfolk, and Plymouth.

The Eastern group non-hunting deer mortalities (P-R W-35-R-13 and 14: II-2, 1971 and 1972) show that there were 85 non-hunting deer mortalities in 1971 and 61 non-hunting mortalities in 1972. The reported legal deer harvest for these years in the Eastern group of counties was 54 and 50 deer respectively. Although a portion of the 2271.6 square miles of deer range of the Eastern counties is not open to hunting, the reported deer harvest for this area is still below all expectations.

Table 1. A summary of the sex and age composition of Massachusetts deer reported at mainland biological stations, 1967 through 1972.

Age	Males					Females						
	1967	1968	1969	1970	1971	1972	1967	1968	1969	1970	1971	1972
6 mos.	28	61	67	121	100	77	31	44	76	90	101	90
1-1/2	167	198	239	263	211	260	32	33	55	54	64	57
2-1/2	122	126	153	147	103	144	18	28	55	69	69	56
3-1/2	62	87	98	97	77	96	13	16	36	46	51	51
4-1/2	25	35	55	59	43	46	7	11	24	29	33	22
5-1/2	10	17	21	21	19	14	3	3	11	14	20	14
6-1/2	5	4	14	7	3	11	1	4	0	8	14	10
7-1/2	1	0	2	7	6	2	0	3	2	0	11	2
8-1/2-9-1/2	1	0	2	1	1	0	3	3	0	0	9	0
10-1/2	0	0	1	0	0	0	1	1	0	0	0	1
Totals	421	528	652	723	563	650	109	146	259	310	372	303

A summary of the sex and age composition of Massachusetts deer reported at Martha's Vineyard and Nantucket deer checking stations in 1971 and 1972.

Age	Martha's Vineyard			Nantucket		
	Males	Females		Males	Females	
	1971	1972	1971	1972	1971	1972
6 mos.	39	25	31	27	14	21
1-1/2	41	41	11	38	22	16
2-1/2	14	15	16	13	12	12
3-1/2	8	15	11	7	13	5
4-1/2	4	6	4	4	2	5
5-1/2	3	1	8	0	3	0
6-1/2	0	3	2	0	0	1
7-1/2	1	2	2	1	0	1
8-1/2-9-1/2	1	0	3	0	0	0
10-1/2	0	0	0	0	0	0
Totals	111	108	88	90	66	61

Figure 1. A summary of the Massachusetts deer harvest by sex per square mile of deer range by county, 1967 through 1972.

Table 2. A summary of the Massachusetts deer harvest by shotgun, 1967 through 1972

<u>Year</u>	<u>All Males</u>	<u>Adult Males</u>	<u>Button Bucks</u>	<u>All Females</u>	<u>Adult Females</u>	<u>Fawn Females</u>	<u>Total Deer</u>	<u>Total Permits Issued</u>
1967	937	886	53	235	168	67	1,172	2,236
1968	1,083	1,022	61	310	217	93	1,393	2,331
1969	1,424	1,299	125	586	415	171	2,010	4,695
1970	1,598	1,378	220	769	546	223	2,367	6,747
1971	1,359	1,106	253	889	643	246	2,248	7,270
1972	1,455	1,256	199	760	530	230	2,215	5,326

Table 3. A summary of percentage change in adult male harvest and calculated minimal populations of deer in Massachusetts, 1967 through 1972

	<u>1967</u>	<u>Percent Change 1967-1968</u>	<u>1968</u>	<u>Percent Change 1968-1969</u>	<u>1969</u>	<u>Percent Change 1969-1970</u>	<u>1970</u>
Adult male harvest	886	+ 15.35	1,022	+ 27.10	1,299	+ 6.08	1,378
Calculated adult male population	2,059	+ 17.05	2,410	+ 31.95	3,180	- 0.82	3,154
Calculated adult female population	2,018	+ 31.92	2,964	+ 35.19	4,007	+ 32.24	5,299
Calculated minimal population	6,139	+ 42.37	8,740	+ 35.01	11,805	+ 27.18	15,014
				<u>Percent Change 1970-1971</u>	<u>1971</u>	<u>Percent Change 1971-1972</u>	<u>1972</u>
Adult male harvest				- 19.73	1,106	+ 13.56	1,256
Calculated adult male population				- 27.68	2,281	+ 20.78	2,755
Calculated adult female population				- 22.95	4,083	- 3.67	3,933
Calculated minimal population				- 22.93	11,571	- 2.03	11,336

Figure 2. Massachusetts deer harvest, 1967 to 1972

A summary of the reported statewide shotgun deer harvest and the number of antlerless permits issued per year from 1967 through 1972 is presented in Table 2. During the 1972 shotgun season successful deer hunters removed 2,215 deer from the statewide herd. Of these 1,256 were adult bucks, 199 were button bucks, 530 were adult females and 230 were female fawns. Five thousand three hundred twenty-six antlerless permits were issued for the 1972 shotgun deer hunting season, a decrease of 1,944 permits from the 1971 deer season.

Figure 2 shows in graphic form a summary of the Massachusetts deer harvest per year and the number of antlerless permits issued from 1967 through 1972. It can be seen that the deer harvest peaked in 1970 when 1,378 adult males and 989 females and button bucks were taken. A total of 6,747 antlerless permits were issued for the 1970 season. This was an increase of 2,052 permits above the 4,695 permits issued in 1969. In 1971, the number of permits was increased to 7,270 permits or 523 permits more than the 1970 total of 6,747. With each increase in the number of permits issued from 1969 through 1971 the rate of increase of the adult male harvest decreased while the antlerless (button buck-female) harvest increased.

A summary of the percent change in the adult male harvest and the calculated minimal populations of deer in Massachusetts from 1967 through 1972 is presented in Table 3. The minimal deer population was calculated using a formula based on the percent of 1-1/2 year-old males removed to the total adult males reported at the biological check stations (W-35-R-12:II-4, 1970). From 1967 to 1969 the percent change of the adult male harvest, the calculated adult male and female populations and the calculated minimal population show an annual increase. The rate of increase of the percent of change for all categories decreased from 1969 to 1970 and continued to decrease into the 1971 deer season. The number of antlerless (sportsman) permits issued for the 1972 season was reduced on the Mainland to 4,000 permits. The percent change for all categories increased from 1971 to 1972.

Tables 4 and 5 present a summary of the adult male and female harvest of deer per square mile of deer range per county in Massachusetts from 1967 through 1972. The sensitivity of the permit system of harvesting deer is discernible in both of these tables. The number of antlerless sportsmen's permits issued in 1972 was reduced from 6,000 permits to 4,000 permits on the Mainland. The purpose of the reduction of permits was to decrease the female harvest in counties producing the bulk of the deer harvest.

Table 4. A summary of the adult male deer harvest per square mile of deer range per county in Massachusetts, 1967 through 1972.

<u>County</u>	<u>Square Miles Deer Range</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Barnstable	290.5	.14	.15	.20	.28	.28	.26
Berkshire	839.0	.27	.37	.44	.44	.42	.45
Bristol	422.6	.01	.01	.01	.01	.002	.007
Essex	344.4	.03	.03	.04	.06	.03	.05
Franklin	649.1	.27	.35	.39	.39	.36	.43
Hampden	524.3	.13	.19	.24	.22	.20	.25
Hampshire	431.5	.16	.25	.27	.26	.15	.23
Middlesex	583.1	.02	.01	.02	.02	.02	.01
Norfolk	277.3	---	---	.01	---	---	---
Plymouth	544.2	.03	.02	.03	.02	.01	.02
Worcester	1307.6	.11	.06	.14	.15	.07	.09
Total	6213.6	.12	.14	.18	.19	.15	.18
Dukes	86.5	.71	.65	.72	.91	1.01	1.18
Nantucket	35.9	1.34	1.56	1.89	2.42	1.78	1.36

Table 5. A summary of the adult female harvest per square mile of deer range per county in Massachusetts, 1967 through 1972.

<u>County</u>	<u>Square Miles Deer Range</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>
Barnstable	290.5	.03	.02	.06	.10	.09	.09
Berkshire	839.0	.04	.06	.12	.16	.23	.13
Bristol	422.6	.005	.002	.002	---	---	.002
Essex	344.4	.006	.003	.01	.03	.003	.003
Franklin	649.1	.07	.11	.14	.15	.27	.22
Hampden	524.3	.02	.03	.06	.06	.08	.07
Hampshire	431.5	.04	.06	.11	.11	.07	.08
Middlesex	583.1	.002	.01	.01	.01	.01	.01
Norfolk	277.3	.004	---	---	.004	---	---
Plymouth	544.2	.002	.01	.01	.01	.002	---
Worcester	1307.6	.02	.02	.05	.07	.05	.03
Total	6213.6	.02	.03	.06	.07	.08	.07
Dukes	86.5	.14	.10	.18	.36	.67	.89
Nantucket	35.9	.17	.14	.97	1.67	1.53	1.11

In Barnstable, the adult female harvest remained unchanged with .09 deer per square mile while the antlered male dropped .02 deer from .28 in 1971 to .26 in 1972. The harvest of adult females in Berkshire shows an expected drop of .10 from .23 in 1971 to .13 deer per square mile in 1972. The adult male harvest in Berkshire County increased from .42 in 1971 to .45 in 1972. In Franklin County, the slight reduction of .05 adult females from .27 deer in 1971 to .22 in 1972 indicates considerable hunting pressure. The antlered buck harvest in Franklin County showed an increase of .07 deer from .36 deer in 1971 to .43 adult males in 1972.

The adult male harvest increased slightly in Hampden, Hampshire and Worcester Counties during the 1972 deer season as compared to the previous season. The adult female harvest for these counties decreased slightly during the 1972 season except for Hampshire County which had a .01 deer increase from .07 in 1971 to .08 deer in 1972.

The remaining counties; namely, Bristol, Essex, Middlesex, Norfolk and Plymouth, with exception of Middlesex, show slight increases in the antlered male harvest. The changes in the adult female harvest for these counties were from no harvest of females to .001 deer per square mile of deer range. It is quite evident that there is very little hunting pressure in these counties or the hunters are not reporting the taking of a deer.

Harvest data seen in Table 4 and 5 show that the adult male and adult female harvest in 1972 on Martha's Vineyard increased .17 male deer and .22 female deer per square mile over the 1971 harvest. This was the second year of the antlerless permit system of hunting with 600 permits issued for Martha's Vineyard only. With the completion of the 1973 deer hunting season a solid foundation of permit deer hunting for this island will have been completed, provided that the number of antlerless permits remains at 600 permits.

The harvest of adult deer on Nantucket has progressed as expected with the harvest of 1.36 adult males and 1.11 adult females per square mile of deer range. Four hundred (400) antlerless permits were issued for Nantucket Island from 1969 through 1972.

The harvest of adult deer on Nantucket Island clearly demonstrates the effectiveness and sensitivity of the antlerless permit system used in manipulating and controlling a deer population. During the first two years, 1967 and 1968, of the statewide antlerless permit system of harvesting deer, there were relatively few adult females harvested on Nantucket. The harvest data in Table 5 show that .17 and .14 adult females were reported per square mile in 1967 and 1968 respectively. The antlered male harvest during these same years was a healthy 1.34 and

1.56 bucks per square mile. To cope with the increased deer herd on the island and bring the herd size into balance, 400 antlerless permits were issued for the 1969 deer season. As expected, the Nantucket adult female harvest increased from .14 deer per square mile in 1968 to .97 deer in 1969. The adult buck harvest continued to increase 1.56 in 1968 to 1.39 males per square mile in 1969. The peak of the deer harvest on Nantucket occurred in 1970 when 2.42 antlered males and 1.67 adult females per square mile were recorded. The predicted decrease in the adult harvest of the Nantucket deer herd occurred with the harvest of 1.78 bucks in 1971 and 1.36 antlered males in 1972. (The peak was 2.42 bucks in 1970.) During the same two-year period (1971 and 1972), the female harvest decreased from 1.67 does in 1970 to 1.53 in 1971 and 1.11 adult females per square mile in 1972.

It is planned to bring the adult male to female harvest close to a one-to-one ratio for Nantucket Island during the 1973 deer season. The number of antlerless permits issued for any future year can be adjusted to hold the one-to-one ratio and maintain the herd in balance.

Table 6 presents the percent frequency ratio of adult females to adult males per county from 1967 through 1972. The results of the 2,000 permit reduction on the Mainland in 1972 is evident in Table 6. With the exception of Barnstable and Middlesex, all counties show a lower percent frequency of females being harvested in 1972 compared to the 1971 percent frequencies. The effect of the 1972 permit reduction should show as an increase in the Mainland male harvest frequency in 1974.

The percent frequency of adult females to adult males increased slightly on Martha's Vineyard but decreased slightly on Nantucket Island during the 1972 deer hunting season.

The data documented in this report indicate the need for the continuation and the expansion of the antlerless permit system of harvesting deer in Massachusetts. By expanding the system to its full potential of a county or regional distribution of permits, it will be possible to maintain a harvestable deer herd in Massachusetts.

How many deer hunters are there in Massachusetts and where do they hunt are questions often asked of project personnel. The answers given of an estimated 45,000 to 55,000 deer hunters is based on an expanded card survey conducted over 20 years ago. In order to expand the antlerless permit system to its full potential the big game license would be a useful tool.

Table 6. Percent frequency ratio of adult females to adult males per county, 1967 through 1972.

County	<u>1967</u>			<u>1968</u>			<u>1969</u>			<u>1970</u>			<u>1971</u>			<u>1972</u>		
	Adult M.	F.	% Freq.	Adult M.	F.	% Freq.	Adult M.	F.	% Freq.	Adult M.	F.	% Freq.	Adult M.	F.	% Freq.	Adult M.	F.	% Freq.
Barnstable	41	10	.24	44	6	.14	58	16	.28	80	29	.36	82	25	.30	75	27	.36
Berkshire	228	31	.14	314	53	.17	371	100	.27	366	138	.38	352	187	.53	373	112	.30
Bristol	6	2	.33	5	1	.20	3	1	.33	4			1			3	1	.33
Essex	11	2	.18	10	1	.10	13	4	.31	20	9	.45	12	1	.08	16	1	.06
Franklin	176	44	.25	230	68	.30	256	93	.36	251	98	.39	232	176	.76	277	141	.51
Hampden	69	9	.08	98	13	.13	123	32	.26	117	31	.26	103	42	.41	131	38	.29
Hampshire	69	16	.23	107	27	.25	115	47	.41	114	48	.42	63	31	.49	98	34	.35
Middlesex	10	1	.10	7	3	.43	13	4	.31	9	6	.67	14	7	.50	3	4	1.33
Norfolk		1			3						1							
Plymouth	16	1	.06	8	3	.38	14	6	.43	8	4	.50	7	1	.14	12	0	.00
Suffolk																		
Worcester	140	32	.23	75	28	.37	176	59	.34	192	88	.46	89	60	.67	117	44	.38
Total	766	149	.19	898	203	.23	1,145	362	.32	1,161	452	.39	955	530	.55	1,105	402	.36
No. of sports- man permits	2,000			2,000			4,000			6,000			6,000			4,000		
Dukes	61	12	.20	56	9	.16	62	16	.26	79	31	.39	87	58	.67	102	77	.75
No. of sports- man permits														600			600	
Nantucket	48	6	.13	56	5	.09	68	35	.51	87	60	.69	64	55	.86	49	40	.82
No. of sports- man permits							400				400			400			400	
Total number of permits*	2,243			2,331			4,695			6,747			7,270			5,326		

* Includes Farmer-Landowner Permits

The antlerless permit application could be a part of the big game license. The hunter could indicate in which deer management unit he intends to hunt on his application. Having the antlerless permit a part of the big game license would be a tremendous saving to the Division by avoiding duplication of applications and the expense of printing, distributing and handling of the present form of permit.

It is possible to determine from successful hunters where they live and where they hunted but to determine hunting pressure on a statewide basis a question should be included on the 1974 antlerless permit application as to where the applicant intends to hunt.

Recommendations: Successful hunters should be allowed to close the trunk of their cars and the rear window of station wagons after their deer has been officially checked.

Deer hunters on Nantucket and Martha's Vineyard should be required to report their deer on the islands.

The possibility of the sale of a Massachusetts big game license that incorporates an antlerless permit application should be investigated.

A question of where the deer hunter intends to hunt should be placed on the 1974 antlerless permit applications.

The following number of sportsman permits for the 1973 shotgun deer hunting season are recommended:

<u>Deer Management Unit</u>	<u>Number of Recommended Permits</u>
Naushon	50
Nantucket	400
Martha's Vineyard	600
Mainland	4,000

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James J. McDonough, Game Biologist
and
William J. Minior, Assistant

Date _____

PERFORMANCE REPORT

State Massachusetts
 Cooperator: Massachusetts Division of Fisheries and Game
 Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey
 Job No. III-2 Job Title: Spring Quail Census
 Period Covered: 1 July to 31 July 1973

Summary: The 1973 spring quail census in Barnstable, Bristol, and Plymouth counties showed no statistically significant difference (t.05) in call indices from the 1971 average, or from a four-year (1958-1961) average.

Objectives: To determine the dynamic aspects of quail population densities and distribution.

Procedures: Roadside whistle counts were conducted during the first two weeks of July 1971 following established procedures and routes. The resultant call indices were corrected for temperature variations (Bennitt, 1951; Ripley, 1956) and tabulated and analyzed on a county basis. Counts for all three counties were compared with 1971 indices, and with a four-year (1958-1961) index. Changes in annual counts were analyzed for statistical significance and the results reported accordingly by counties.

Findings: The 1973 weighted call indices as compared to those from 1971 and the four-year average are shown in Table 1. Computation of the tests of significance and comparisons of the indices by county and route are shown in Tables 2, 3, and 4.

The 1973 indices were lower than those from 1971 in all three counties. These changes were not statistically significant (t.05) for any of the counties.

None of the 1973 call indices showed a statistically significant difference (t.05) from the 1958-1961 average. Call indices were lower than the average in Barnstable and Plymouth counties, and higher in Bristol County. The addition of two routes in Bristol County is probably responsible for the increase in this index.

Government Documents
 Collection
 NOV 14 1973

University of Massachusetts

Table 1. County Comparative Call Indices of Massachusetts Quail Population

<u>County</u>	<u>Average 1958-1961 Index</u>	<u>1964 Index</u>	<u>1967 Index</u>	<u>1969 Index</u>	<u>1971 Index</u>	<u>1973 Index</u>	<u>Significant Change (t.05)</u>	
							<u>1973 to 1958-61</u>	<u>1973 to 1971</u>
Barnstable	2.65	1.55	1.90	3.31	4.51	2.38	None	None
Bristol	0.71	0.91	0.96	0.95	2.21	1.52	None	None
Plymouth	1.84	1.34	1.83	2.49	2.02	1.56	None	None

Table 2. Barnstable County, 1973 and 1971

Route No.	1971	1973	D	$\bar{D}-\bar{d}$	$\frac{(\bar{D}-\bar{d})^2}{n-1}$
16B	254	266	-12	153.25	23,485.56
18A	808	327	481	646.25	417,639.06
21	292	120	172	337.25	113,737.56
23	260	240	20	185.25	34,317.56
	<u>1,614</u>	<u>953</u>			<u>589,179.74</u>
	$\bar{x}=403.50$	$\bar{x}=238.25$			n-1=3
	$\bar{d} = -165.25$			$\bar{Sd} = \frac{1}{49,098.31} = 221.58$	
	$SD^2 = \frac{589,179.74}{3} = 196,393.25$			$t = \frac{-165.25}{221.58} = -0.746$	
	$S_d^2 = \frac{196,393.25}{4} = 49,098.31$			$t.05 (3 \text{ d.f.}) = 3.182$	
				$t.01 (3 \text{ d.f.}) = 5.841$	
Barnstable County, 1973 and Four-Year Average (1958-1961)					
Route No.	4-Yr. Avg.	1973	D	$\bar{D}-\bar{d}$	$\frac{(\bar{D}-\bar{d})^2}{n-1}$
16B	207	266	-59	-46.75	2,185.56
18A	289	327	-38	-25.75	663.06
21	235	120	115	127.25	16,192.56
23	271	240	31	43.25	1,870.56
	<u>1,002</u>	<u>953</u>			<u>20,911.74</u>
	$\bar{x}=250.50$	$\bar{x}=238.25$			n-1=3
	$\bar{d} = -12.25$			$\bar{Sd} = \frac{1}{1742.64} = 41.74$	
	$SD^2 = \frac{20,911.74}{3} = 6970.58$			$t = \frac{-12.25}{41.74} = -0.293$	
	$S_d^2 = \frac{6970.58}{4} = 1742.64$			$t.05 (3 \text{ d.f.}) = 3.182$	
				$t.01 (3 \text{ d.f.}) = 5.841$	

Table 3. Bristol County, 1973 and 1971

Route No.	<u>1971</u>	<u>1973</u>	<u>D</u>	<u>D-d</u>	<u>(D-d)²</u>
9A	171	107	- 64	- 56.25	3,164.06
15	271	200	- 71	- 63.25	4,000.56
35	086	180	94	101.75	10,353.06
36	<u>110</u>	<u>120</u>	10	17.75	315.06
	638	607			<u>17,832.74</u>
	$\bar{x}=159.50$	$\bar{x}=151.75$			n-1=3
$\bar{d} = -7.75$				$\bar{sd} = \frac{1486.06}{3}$	$= 38.55$
$SD^2 = \frac{17,832.74}{3} = 5944.25$				$t = \frac{-7.75}{38.55}$	$= -0.201$
$s\bar{d}^2 = \frac{5944.25}{4} = 1486.06$				t.05 (3 d.f.)	$= 3.182$
				t.01 (3 d.f.)	$= 5.841$
Bristol County, 1973 and Four-Year Average (1958-1961)					
Route No.	<u>4-Yr. Avg.</u>	<u>1973</u>	<u>D</u>	<u>D-d</u>	<u>(D-d)²</u>
9A	74	107	33	- 49.5	2,450.25
15	68	200	132	49.5	2,450.25
35	No 1958-1961 data available				
36	No 1958-1961 data available				
	<u>142</u>	<u>307</u>			<u>4,900.50</u>
	$\bar{x}=71$	$\bar{x}=153.5$			n-1=1
$\bar{d} = 82.5$				$\bar{sd} = \frac{2450.25}{1}$	$= 49.50$
$SD^2 = \frac{4900.50}{1} = 4900.50$				$t = \frac{82.5}{49.50}$	$= 1.67$
$s\bar{d}^2 = \frac{4900.50}{2} = 2450.25$				t.05 (1 d.f.)	$= 12.706$
				t.01 (1 d.f.)	$= 63.657$

Table 4. Plymouth County, 1973 and 1971

Route No.	1971	1973	D	$\bar{D}-\bar{d}$	$(\bar{D}-\bar{d})^2$
5	120	136	16	62	3,844
6A	199	154	- 45	1	1
12	305	200	-105	-59	3,841
13A	183	133	- 50	- 4	16
	<u>807</u>	<u>623</u>			<u>7,702</u>
	$\bar{x}=201.75$	$\bar{x}=155.75$			n-1=3
$\bar{d} = -46$					
$SD^2 = \frac{7702}{3} = 2567.33$					$S\bar{d} = \frac{\sqrt{641.83}}{25.33} = 25.33$
$S\bar{d}^2 = \frac{2567.33}{4} = 641.83$					$t = \frac{-46}{25.33} = -1.82$
					$t.05 \text{ (3 d.f.)} = 3.182$
					$t.01 \text{ (3 d.f.)} = 5.841$

Plymouth County, 1973 and Four-Year Average (1958-1961)

Route No.	4-Yr. Avg.	1973	D	$\bar{D}-\bar{d}$	$(\bar{D}-\bar{d})^2$
5	115	136	21	64.25	4,128.06
6A	115	154	39	82.25	6,765.06
12	319	200	-119	-75.75	5,738.06
13A	247	133	-114	-70.75	5,005.56
	<u>796</u>	<u>623</u>			<u>21,636.74</u>
	$\bar{x}=199$	$\bar{x}=155.75$			n-1=3
$\bar{d} = -43.25$					
$SD^2 = \frac{21,636.74}{3} = 7212.25$					$S\bar{d} = \frac{\sqrt{1803.06}}{42.46} = 42.46$
$S\bar{d}^2 = \frac{7212.25}{4} = 1803.06$					$t = \frac{-43.25}{42.46} = -1.02$
					$t.05 \text{ (3 d.f.)} = 3.182$
					$t.01 \text{ (3 d.f.)} = 5.841$

Recommendations: Due to commitments on other projects, the number of routes which can be surveyed for the quail census has decreased considerably since the inception of the census. Since so few routes are surveyed, statistical tests can reflect only the most marked variations in whistle counts. Furthermore, increases in urbanization and traffic flow are disturbances which unduly interfere with the progress of the census. Since abatement of these disturbances is unlikely, and relocation of the routes often impractical, further conduct of the quail census should be curtailed.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by: _____
James E. Cardoza, Assistant Game Biologist

Date: _____

Literature Cited

- Bennitt, Rudolph. 1951. Some Aspects of Missouri Quail and Quail Hunting, 1938-1948. Mo. Cons. Comm. Tech. Bull. 2. 51 pp.
- Ripley, Thomas H. 1956. Annual Whistle Count Census to Determine Relative Population Densities and Distribution. Mass. Div. Fisheries and Game, Westboro. Project W-25-R-3, I-A. Supplement I, Table D.

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No. IV-1 Job Title: Experimental Turkey Stocking

Period Covered: 1 June 1972 to 31 May 1973

Summary: Ten wild-trapped turkeys (nine toms and one hen) were captured in Allegany State Park, Cattaraugus County, New York, and released in Beartown State Forest, Berkshire County, Massachusetts, where seven other turkeys were released in 1972. There have been sightings of adult birds but as yet of no broods since the original release. Efforts are being made to obtain hens to complete the stocking.

Populations of semi-wild and game farm ancestry turkeys at six locations in Massachusetts show little or no increase since 1972. Further restoration efforts utilizing semi-wild turkeys are inadvisable.

Objectives: To re-establish the wild turkey in the Commonwealth in sufficient numbers to allow for recreational hunting.

Procedures: Turkeys were censused by roadside counts, track counts, and cooperator reports. Snowmobiles were used during the winter to provide access to the areas and to transport grain for baiting.

Turkeys were captured using Thornsberry style cannon nets. Captured turkeys were sexed, aged, weighed, leg banded, and color marked with patagial streamers.

Findings: Statewide Turkey Populations

Winter observations were limited by the scarcity of snow for tracking. Spring counts were hampered by heavy rains and muddy roads that limited vehicular traffic. These conditions, and a paucity of cooperator reports, preclude listing of population figures for the various release sites.

Quabbin Reservation-New Salem Populations. Turkey populations in the Quabbin area remained relatively static during the past year. At least two broods were produced on Prescott Peninsula, near Underhill Brook and Prescott Brook. An additional brood was located in New Salem on West Street near Hamilton's Orchards. Cooperators reported a brood near Fever Brook in Dana and adult birds in the area of Town Farm Brook, Shutesbury, and in the Wheelwright section of Hardwick.

During winter, turkeys could be consistently located only on the southern portion of the peninsula, in the area of Prescott Brook. In the spring of 1973, intense rains occurred again, with probable mortality of eggs and poults.

Although Quabbin-strain turkeys appear wild in a forest situation, experiences with them near dwellings in New Salem indicate that they tame readily, and consequently are unsuitable for further restoration efforts.

Barre, Douglas and Mt. Holyoke Populations. Quabbin turkeys were trapped, transported, and released in Barre and Douglas State Forests, and in the Holyoke Range between 1965 and 1969. Two releases in the Holyoke Range failed to establish a population, with the last turkey sighting reported in early winter, 1972. Releases in Barre and Douglas were initially successful with good poult production and spread of turkeys into neighboring towns. Since 1971-72, however, both of these populations have declined. Reports are still received, but of fewer adults and fewer broods. A lack of vigor, and their ready acceptance of artificial food is evidence of their partial game-farm ancestry. Cold spring rains, as in the Quabbin, may also have adversely affected their reproductive success. There should be no further incrementation of these flocks with semi-wild birds. I see no necessity at this time to remove any of these turkeys, however.

Mt. Washington and October Mountain Forest Populations. Turkeys in these two locations are entirely of game-farm ancestry. There has been little or no spread from the release site, and the local citizenry has aggravated the turkeys' inherent tameness by unnecessary supplemental feeding. Ten of these turkeys were removed in 1971 to test a capture technique (tribromoethanol-treated bait) and to free the area for release of wilder birds. If any remaining turkeys pose a problem to future releases through potential genetic pollution, they should be removed by trapping or an alternative means. Otherwise, these birds should be undisturbed, since unnecessary interference with them may cause local public relations problems.

Myles Standish Forest Population. This flock has fared poorly ever since the release in 1966-67 of 14 wild-trapped West Virginia turkeys. Poor habitat, excessive dispersal and road kills appear to be the causative agents in their decline. The population is now apparently extinct, with no birds sighted since July 1971.

Horse Mountain Population. Six turkeys, trapped in New Salem, were released on Horse Mountain in Hatfield in September 1972. One hen was killed by a dog two months after release. Although cooperators still report a few turkeys in the area, there has apparently been no reproduction, probably due to the fact that there was only one tom, an immature, in the original release. Since no releases of wild-trapped turkeys are projected for this area in the near future, I see no conflict between the Horse Mountain flock and present restoration efforts.

Turkey Trapping

In an attempt to obtain additional wild stock for Berkshire County releases, bait lines were established on the Bear Spring Mountain Wildlife Management Area, Delaware County, New York. Baiting was initiated on 29 August and continued weekly until 6 October by the project leader and Western District personnel. No turkeys were seen, and turkey sign was seen only once. No trapping could be attempted.

In February 1973, New York State personnel informed us that we would again be able to obtain wild-trapped birds from Allegany State Park, Cattaraugus County. The project leader and an assistant traveled to this location and, assisted by New York technicians, were successful in capturing nine turkeys between 22 February and 2 March. One additional bird was captured in April by cooperators, transported to Albany, and from there to Massachusetts. All ten turkeys (Table 1) were released off Alcott Street, Beartown State Forest, Great Barrington, Berkshire County, in the same location where seven turkeys were released last year.

As in 1972, mild weather hampered trapping efforts and dominance of the bait site by toms resulted in a release highly skewed to males. Hopefully, future efforts will allow us to obtain hens to complete the projected stocking.

Three turkeys (two toms and a hen) were instrumented by Mr. Walter M. Tzilkowski of the University of Massachusetts. This equipment was attached in order to facilitate location of the flock during future releases and as part of a planned habitat evaluation study.

Sightings of the Beartown turkeys have so far been few. One bird from the 1972 stocking was seen at Alcott Street a week or so after release. Nothing further was heard until February 1973 when a farmer reported seeing an unspecified number of birds in his back field, approximately three kilometers south of the release site. Since then, turkeys have been sighted on the far side of the forest, near Lake Garfield in Monterey, approximately 10.5 kilometers southeast of the stocking location. No broods have yet been located.

7. 1. 1900

Dear Sir,

I have the honor to acknowledge the receipt of your letter of the 28th inst. in relation to the above matter.

Very respectfully,
J. H. [Name]

Enclosed for you are the same as before.

Very respectfully,
J. H. [Name]

I am, Sir, very respectfully,
Your obedient servant,
J. H. [Name]

I am, Sir, very respectfully,
Your obedient servant,
J. H. [Name]

Table 1. Sex, Age, Weight, Band and Tag Number, and Capture-Release Data for Turkeys Captured Spring 1973.

Band No.	Wing Tag Number	Age	Sex	Weight		Capture Date*	Release Date**	Remarks
				Kg	(lbs.-oz.)			
110	T-5297	Ad.	M	8.36	(18- 7)	22 February	23 February	
111	T-5298	Ad.	M	8.56	(18-14)	22 February	23 February	
112	T-5296	Ad.	M	9.01	(19-14)	22 February	23 February	
113	T-5294	Ad.	M	8.87	(19- 9)	22 February	23 February	
114	T-5295	Ad.	M	9.21	(20- 5)	22 February	23 February	Two beards
115	T-5300	Imm.	M	5.56	(12- 4)	26 February	27 February	Instrumented
116	T-5299	Imm.	M	5.92	(13- 1)	26 February	27 February	Instrumented
117	T-5301	Imm.	M	5.95	(13- 2)	26 February	27 February	
118	T-5302	Imm.	M	5.78	(12-12)	26 February	27 February	
193***	T-5306	Ad.	F	4.08	(9- 3)	20 April	23 April	Instrumented

* All turkeys except 193 captured at Camp 12 area, Allegany State Park, Cattaraugus County, New York.

** All turkeys released off Alcott Street, Beartown State Forest, Great Barrington, Berkshire County, Massachusetts.

*** Captured at Holt Run, Allegany State Park,

- Recommendations:
1. If possible, obtain some additional wild-trapped hens to complete the Beartown stocking. If the Beartown release is successful, transplant additional turkeys to other suitable areas in the Berkshires, and elsewhere as suitable stock and habitat are available.
 2. Utilize only wild-trapped stock in future releases. Semi-wild stock such as in the Quabbin may remain undisturbed unless interfering with wild-trapped birds through potential cross-breeding. Occasional monitoring of the progress of these semi-wild flocks is advisable in case they should overcome their previous dependence on man.

Acknowledgments: I extend my appreciation to Messrs. Lee DeGraff, John Proud, Steven Clarke, Quentin Van Nortwick, Fred Evans, William Shirey, and Clint Mount of the New York State Department of Environmental Conservation for their assistance in securing wild-trapped turkeys, and to Dr. Wendell Dodge and Mr. Walter M. Tzilkowski of the Massachusetts Cooperative Wildlife Research Unit for their assistance in monitoring the Beartown releases.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges
Superintendent

Prepared by: _____
James E. Cardoza
Assistant Game Biologist

Date _____

PERFORMANCE REPORT

State: Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: V-1 Job Title: Hunter Utilization of Wildlife Management Areas

Period Covered: 10 October 1972 to 31 May 1973

Summary: Total estimated hunter effort on thirteen wildlife management areas was 32,505 hunter trips, representing an increase of 1.9 percent from the 1971 effort.

Peak usage occurred on the first Saturday, followed by the second Saturday, and opening day. Usage on a weekday after stocking was approximately 2.03 times that of a weekday after no stocking.

Local hunters continue to be the heaviest users of wildlife management areas, although on peak days hunters in the 32-80 kilometer range predominated on two areas (Northeast, Myles Standish), and in the 80 plus range on a third area (Crane).

Game bag information was collected on thirteen areas. On the twelve state areas, 4288 hunters were contacted of whom 1514 (35.3%) had taken at least one unit of game. Known harvest on these areas totalled 1736 animals of nine species. Pheasant (1442, 83%) comprised the majority of the harvest, followed by quail (186, 10.7%) and woodcock (33, 1.9%).

Objectives: To determine hunter utilization, game harvest, and hunter success on wildlife management areas.

Procedures:

1. Roving and stationary checking stations were used to collect hunting pressure and game bag data on fourteen wildlife management areas.
2. Data were collected on the following peak days of usage: opening day, and first, second, fourth and fifth Saturdays. At Fort Devens, checks were made every day during the upland bird season.
3. Data were collected on four additional days during the week, as follows: two counts on a day after stocking, and two when a stocking had not been done the previous

Government Documents
Collection
NOV 14 1973

University of Massachusetts

day. Freetown was not stocked during the week; therefore, only two weekday checks are shown for this area.

4. Division of Fisheries and Game district personnel conducted checks on thirteen areas. The station at Fort Devens was manned by Department of the Army personnel.
5. Hunters using Fort Devens were required to check through a permanent station. Thus, hunting pressure on Devens is based upon this mandatory check of all hunters, and is consequently more accurate than on other areas where many access points and manpower limitations precluded the use of permanent stations.
6. Roving checks were used on the remaining thirteen areas. Checkers travelled by motor vehicle and contacted as many hunters as possible to determine the following: size of party, home town of the hunters, and amount and type of game bagged.
7. Where the roving check was utilized, the number of hunters was calculated by counting the total number of cars on the area for that check day, and multiplying this figure by the average party size. Party size was computed separately for each area each day.
8. Hunters were also asked to provide the name of their hometown. Concentric circles were drawn on a map of the state at 32 kilometer (20 mile) and 80 kilometer (50 mile) distances from the management areas, and the number of hunters coming from towns within each zone was noted. If a town boundary was intersected by one of these zone lines, the town was considered as being in the zone in which the mark indicating the town center was located.
9. Daily summaries of hunting pressure were made on mimeographed check forms (W-35-R-13, V-1, Figure 1).
10. Hunters on the thirteen state-checked areas were contacted to determine game harvest. When possible, hunters were contacted after completion of hunting rather than during the hunt. Checkers noted the total number of hunters contacted, the number unsuccessful, the number taking at least one unit of game, and the amount and species of game taken. Daily results were summarized on a mimeographed form.
11. In calculating hunter success, a successful hunter was defined as one who takes at least one unit of game.

Findings:

A. Hunter Effort

The 1972 hunting pressure on fourteen wildlife management areas for peak day, weekday, and total usage is presented

in Tables 1, 2, and 3. Total estimated season usage was 32,505 hunter trips (Marconi Beach area excluded).

Hunter effort was greatest on the first Saturday, followed by the second Saturday, opening day, fourth Saturday, and fifth Saturday. Total weekend effort ranged from 0.82 (Myles Standish) to 3.56 (Crane Pond) times that of weekdays (excluding opening day), and averaged 1.29.

The average effort on a weekday after stocking was 2.03 times that of a weekday after no stocking. Although the 1971 rates (1.64) represented a slackening of the disparity between the two categories of weekdays (1970 ratio was 2.26), the past seasons figure indicates that sportsmen are able to anticipate or discern stocking dates regardless of a staggered stocking schedule.

The estimated hunting pressure on the thirteen areas comparable to 1971 data was 32,505 hunter trips (Table 4). This is an increase of 1.9 percent in total effort.

Total hunter effort increased on four areas and decreased on the remaining nine. Major increases were at Barre Falls (+59.1%) and Myles Standish (+33.4%), while considerable decreases were recorded at Housatonic Valley (-41.6%) and Stafford Hill (-32.2%). Changes on all other areas were less than 20 percent.

The increase at Barre Falls occurred largely on peak days (opening day plus Saturdays); whereas, that at Myles Standish was primarily on weekdays. The decline at Crane Pond took place entirely on weekdays, that at West Hill primarily on weekdays, and that at Winnimusset primarily on peak days. Crane experienced a gain, and Housatonic Valley and Stafford Hill a loss in both categories of days (Table 5). Remaining areas had an overall percent of change of less than 10 percent.

B. Distances Travelled by Hunters

Yearly comparisons of the distances travelled by hunters to wildlife management areas are presented in Table 6. As in the past, local hunters continue to be the greatest overall users of wildlife management areas, with a few areas being favored by more distant sportsmen.

Five areas (four in 1970), Northeast, Myles Standish, Crane Pond, Crane, and Winnimusset, had at least 25 percent of their peak day hunters travelling 32-80 kilometers. On one area (Northeast, 56%) hunters in this distance group predominated, with a second area (Myles Standish, 48%) having a percentage equal to that of the 0-32 kilometer group.

Table 1. Summary of Usage on Peak Days.

<u>Area</u>	<u>Opening Day 20 October</u>	<u>First Saturday 21 October</u>	<u>Second Saturday 28 October</u>	<u>Fourth Saturday 11 November</u>	<u>Fifth Saturday 18 November</u>	<u>Total</u>
Crane	217	320	255	225	182	1,199
Freetown	153	191	175	130	116	765
Marconi Beach	73	95	79	76	86	414
Myles Standish	391	636	491	407	211	2,136
Crane Pond	220	275	156	150	174	975
Northeast	269	357	259	221	171	1,277
Housatonic Valley	59	74	78	44	36	291
Stafford Hill	106	146	182	46	20	500
Fort Devens	167	339	209	196	175	1,086
Barre Falls	142	225	262	221	202	1,052
Birch Hill	286	621	550	216	218	1,891
Hubbardston	155	350	214	175	86	980
West Hill	213	267	328	140	144	1,092
Winimusset	112	121	128	92	80	533
Totals	2,568	4,017	3,366	2,339	1,901	14,191

Table 2. Summary of Usage on Weekdays.

Area	Mon.	Weekday After Stocking			Weekday After No Stocking			Total	Average
		Tues.	Wed.	Thurs.	Tues.	Wed.	Thurs.		
Crane			110		17	39		166	42
Freetown*					21	7		28	14
Myles Standish	290		108		16			439	110
Crane Pond		6	16					44	11
Northeast		40	90			83	11	242	61
Housatonic Valley**			23					26	13
Stafford Hill**			25				3	31	16
Barre Falls		74					9	102	26
Birch Hill	75	70			75	28		220	55
Hubbardston		24				48		94	24
West Hill		43				50	22	93	24
Winimusset		15	27			18		83	21
Totals	365	272	399	0	129	273	100	1,568	37.3

Total by Category 1,036 $\bar{x} = 51.8$

535 $\bar{x} = 25.5$

Ratio = 2.03

* Not stocked on weekdays. Two checks only.

** Two weekday checks only.

— Two counts made on indicated weekday.

Table 3. Total Estimated Usage on 13 Wildlife Management Areas in 1972.

Area	Weekday Average		Saturday Average		Opening Day Usage	Total Usage	Weekend/ Weekday Ratio
	x 29 days		x 6 days				
Crane	1,218		1,473		217	2,908	1.21
Freetown*	406		918		153	1,477	2.26
Myles Standish	3,190		2,618		391	6,199	0.82
Crane Pond	319		1,132		220	1,671	3.56
Northeast	1,769		1,512		269	3,550	0.85
Housatonic Valley*	377		348		59	784	0.92
Stafford Hill*	464		591		106	1,161	1.27
Fort Devens**	---		---		---	2,536	---
Barre Falls	754		1,365		142	2,261	1.81
Birch Hill	1,595		2,408		286	4,289	1.51
Hubbardston	696		1,238		155	2,089	1.79
West Hill	696		1,318		213	2,227	1.89
Winimusset	609		632		112	1,353	1.04
Totals	12,093		15,553		2,323	32,505	$\bar{x} = 1.29$

* Weekday counts based on two check days.

** Total actual check determined by mandatory count every day of season.

Table 4. Comparison of Estimated Hunting Pressure on 13 Wildlife Management Areas: 1971 and 1972.

Area	Estimated Pressure		Percent of Change 1972 from 1971
	1971	1972	
Crane	2,577	2,908	+12.8
Freetown	1,376	1,477	+ 7.3
Myles Standish	4,645	6,199	+33.4
Crane Pond	2,020	1,671	-17.3
Northeast	3,257	3,550	- 9.0
Housatonic Valley	1,342	784	-41.6
Stafford Hill	1,712	1,161	-32.2
Fort Devens	3,165	2,536	-19.9
Barre Falls	1,421	2,261	+59.1
Birch Hill	4,637	4,289	- 7.5
Hubbardston	2,129	2,089	- 1.9
West Hill	2,626	2,227	-15.2
Winimuset	1,667	1,353	-18.8
Totals	31,910	32,505	+ 1.9

Table 5. Percent Change in Mean Daily Usage for Weekdays and Peak Days: 1972 from 1971, for 12 Management Areas

Area	Peak Days*		Week Days*		Percent Change
	1971	1972	1971	1972	
Crane	209	240	38	42	+10.5
Freetown	131	153	16	14	-12.5
Myles Standish	413	427	59	110	+86.4
Crane Pond	181	195	26	11	-57.7
Northeast	257	255	51	61	+19.6
Housatonic Valley	79	58	27	13	-51.8
Stafford Hill	113	100	32	16	-50.0
Barre Falls	117	210	20	26	+30.0
Birch Hill	422	378	57	55	- 3.5
Hubbardston	194	196	26	24	- 7.7
West Hill	230	218	34	24	-29.4
Winimuset	156	107	20	21	- 5.0

*Average usage

Table 6. Distance Travelled by Hunters to Wildlife Management Areas on Opening Day and Saturdays, 1965-1972.

Area	Percent Traveling Less * Than 32 Kilometers										Percent Traveling More Than 80 Kilometers									
	1965	1966	1967	1968	1969	1970	1971	1972	1965	1966	1967	1968	1969	1970	1971	1972	1965	1966	1967	1968
Crane	24	28	25	36	31	30	18	17	32	22	30	41	37	34	46	35	44	50	45	23
Freetown	93	98	97	98	96	97	95	98	4	2	3	2	2	3	5	2	3	0	0	0
Myles Standish	32	29	27	30	33	32	47	48	66	68	66	47	57	64	51	48	2	3	7	23
Crane Pond	--	--	--	--	--	60	61	54	--	--	--	--	--	40	39	45	--	--	--	--
Northeast	22	14	24	13	26	33.5	37	44	78	76	71	77	73	66	63	56	0	10	5	5
Housatonic Valley	--	--	--	--	--	98	99	96	--	--	--	--	--	0.5	1	3.5	--	--	--	--
Stafford Hill	--	--	--	--	--	--	100	100	--	--	--	--	--	--	0	0	--	--	--	--
Barre Falls	89	84	90	75	77	81	83.5	89	9	14	10	23	18	15	15	9	2	2	0	2
Birch Hill	76	66	76	54	41	78	69	64	14	22	6	26	25	15	21	23	10	12	18	20
Hubbardston	57	75	62	60	62	74	75	73	34	24	34	30	28	24	24	20	9	1	4	10
West Hill	91	89	91	91	85	93	94	91.8	8	10	9	8	15	6.5	5.8	8	1	1	0	1
Winimusset	--	--	--	--	--	70	76.5	74	--	--	--	--	--	30	23	26	--	--	--	--

-- Indicates area not sampled.

* 0.32 km = 0-20 miles
 32-80 km = 20-50 miles
 80+ km = 50+ miles

Table 6. (Cont.). Distance Traveled by Hunters to Wildlife Management Areas on Weekdays, 1965 to 1972.

Area	Percent Traveling Less Than 32 Kilometers												Percent Traveling 32-80 Kilometers												Percent Traveling More Than 80 Kilometers											
	1965	1966	1967	1968	1969	1970	1971	1972	1965	1966	1967	1968	1969	1970	1971	1972	1965	1966	1967	1968	1969	1970	1971	1972	1965	1966	1967	1968	1969	1970	1971	1972				
Crane	37	38	30	62	43	36.5	28	21	22	18	23	35	43	27	39	48	41	44	47	3	15	36.5	33	31												
Freetown	97	100	67	--	--	100	95	100	0	0	33	--	--	0	5	0	3	0	0	--	--	0	0	0												
Myles Standish	44	38	35	39	46	41	36	51	52	40	57	46	51	58	51	48	4	22	8	15	4	1	13	1												
Crane Pond	--	--	--	--	--	85	49	62	--	--	--	--	--	15	51	38	--	--	--	--	--	0	0	0												
Northeast	41	33	59	49	53	42	61	53	59	61	40	51	47	58	39	47	0	6	1	0	0	0	0	0												
Housatonic Valley	--	--	--	--	--	100	94	100	--	--	--	--	--	0	5	0	--	--	--	--	--	0	1	0												
Stafford Hill	--	--	--	--	--	--	100	100	--	--	--	--	--	--	0	0	--	--	--	--	--	--	0	0												
Barre Falls	90	94	80	82	40	97	84	88	10	14	20	16	55	3	16	7	0	2	0	2	5	0	0	5												
Birch Hill	63	68	91	51	32	74	82	89	21	20	3	38	30	15	13	3	16	12	6	11	38	11	5	8												
Hubbardston	77	85	75	61	19	80	75	80	22	15	25	37	78	20	25	20	1	0	0	2	3	0	0	0												
West Hill	89	100	95	90	97	95	96	99	8	0	5	9	3	5	3	1	3	0	0	1	0	0	1	0												
Winimusset	--	--	--	--	--	72	82	75	--	--	--	--	--	28	17	25	--	--	--	--	--	0	1	0												

-- Indicates area not sampled

On weekdays, five areas (five also in 1971) had 25 percent or more of their hunters in the middle (32-80 kilometer) distance category. Myles Standish (48%), Crane (48%), and Northeast (47%) all had nearly half of their hunters in this group.

Crane remains the only area to be used extensively by hunters travelling greater than 80 kilometers. Nearly half (48%) of the peak day, and 31 percent of the weekday hunters travelled this distance in pursuit of sport. The only other area with a sizeable proportion of far-travelling hunters was Birch Hill, (13 percent peak days, 8 percent weekdays).

The remaining areas were utilized largely by local hunters. Stafford Hill (as in 1971) was utilized entirely by local hunters. Freetown (98% weekends, 100% weekdays), Housatonic Valley (96%-100%), West Hill (91.8%-99%), and Barre Falls (89%-88%) also had little non-resident usage. Hubbardston reflected an increase in local usage from 75% in 1971 to 80% in 1972, with Winimusset showing three-quarters local and one-quarter in the mid-distance group.

C. Game Bag Harvest and Hunter Success

Game bag data was collected on five selected management areas in 1970 and 1971. In 1972, this coverage was extended to include the twelve management areas consistently sampled by the Division. Department of the Army Cooperators provided harvest data for Fort Devens, as they have since 1969.

The amount and species of game taken by management area and type of day for 1971 and 1972 are presented in Table 7. A total of 1736 units of game of 13 species were harvested on 12 areas on the nine days per area sampled (108 sample days). Pheasant (1442, 83.0%), was the predominant species taken, followed by bobwhite quail (186, 10.7%), woodcock (33, 1.9%), ruffed grouse (27, 1.6%), cottontail rabbit (24, 1.4%), and white hare (15, 0.9%). Waterfowl (four species), gray squirrel, woodchuck, and fox, comprised the remaining two percent of the harvest.

Pheasant were taken on all management areas, with the greatest harvests being Northeast (286), and Birch Hill (262), and the least at Barre Falls (30). At Stafford Hill, the only game taken was pheasant (46).

Quail were taken on Myles Standish (112), and Crane (72), both as stocked and native birds. Two additional quail were harvested illegally at Birch Hill.

Ruffed grouse were taken on five areas, with the greatest harvest at Birch Hill (16).

Table 7. Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game Area - Dates	Pheasant		Quail		Ruffed Grouse		Wood- cock		Water- fowl		Subtotal Birds	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
<u>Crane</u>												
Opening Day	25	14	3	8	1						29	22
Saturday (4)	90	91	39	62	2	3	1	1			132	157
Weekday after stocking (2)	41	23	19	2		1					60	26
Weekday after no stocking (2)	2	1	5								7	1
Totals	158	129	66	72	3	4	1	1			228	206
<u>Freetown*</u>												
Opening Day		17							1WD			18
Saturday (4)		55										55
Weekday after no stocking (2)												
Totals		72							1			73
<u>Myles Standish*</u>												
Opening Day		12		18		1						31
Saturday (4)		83		75		2						160
Weekday after stocking (2)		27		19								46
Weekday after no stocking (2)												
Totals		122		112		3						237
<u>Northeast</u>												
Opening Day	32	31				1					32	32
Saturday (4)	129	187			4	2	1				134	189
Weekday after stocking (2)	47	62			1						48	62
Weekday after no stocking	5	6			2						7	6
Totals	213	286			7	3	1				221	289
<u>Crane Pond*</u>												
Opening Day		6										6
Saturday (4)		60										60
Weekday after stocking (2)		7										7
Weekday after no stocking (2)		2										2
Totals		75										75
Game Harvest - Birds (1)	371	684	66	184	10	10	2	1		1	449	880

* Area not sampled in 1971.

WD = Wood Duck

Table 7 (Cont.). Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game Area - Dates	Pheasant		Quail		Ruffed Grouse		Wood- cock		Water- fowl		Subtotal Birds	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
<u>Housatonic Valley</u>												
Opening Day	20	10					2		3WD 2T 2M 2M		25	16
Saturday (4)	51	57					1 2				52	59
Weekday after stocking (1)**	19	14			1						20	14
Weekday after no stocking (1)	5	1									5	1
Totals	95	82			1		1 4		5 4		102	90
<u>Stafford Hill*</u>												
Opening Day		8										8
Saturday (4)		31										31
Weekday after stocking (1)**		4										4
Weekday after no stocking (1)		3										3
Totals		46										46
<u>Barre Falls*</u>												
Opening Day		7					2					9
Saturday (4)		16										16
Weekday after stocking (2)		5			1							6
Weekday after no stocking (2)		2										2
Totals		30			1		2					33
<u>Birch Hill*</u>												
Opening Day		25			2		2					29
Saturday (4)		185	2		11		13					211
Weekday after stocking (2)		36			2		2					40
Weekday after no stocking (2)		16			1		5					22
Totals		262	2		16		22					302
<u>Hubbardston*</u>												
Opening Day		12					1					13
Saturday (4)		86					1					87
Weekday after stocking (2)		14										14
Weekday after no stocking (2)		3										3
Totals		115					2					117
Game Harvest - Birds (2)	95	535	2		1 17		1 30		5 4		102	588

* Area not sampled in 1971

** Two weekdays each category in 1971, one each in 1972.

WD = Wood Duck M = Mallard T = Teal

Table 7 (Cont.). Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game	Pheasant		Quail		Ruffed Grouse		Wood-cock		Water-fowl		Subtotal Birds	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
<u>West Hill*</u>												
Opening Day		16										16
Saturday (4)		77										77
Weekday after stocking (2)		9					2		1BD			12
Weekday after no stocking (2)		5										5
Totals		107					2		1			110
<u>Winimusset</u>												
Opening Day	40	11			1						41	11
Saturday (4)	103	79			2		5				115	79
Weekday after stocking (2)	16	12									16	12
Weekday after no stocking (2)	17	14									17	14
Totals	181	116			3		5				189	116
Game Harvest - Birds (3)	181	223			3		5	2		1	189	226
Birds (2)	95	535		2	1	17	1	30	5	4	102	588
Birds (1)	371	684	66	184	10	10	2	1		1	449	880
Total Birds	647	1442	66	186	14	27	8	33	5	6	740	1694

* Area not sampled in 1971

BD = Black Duck

Table 7 (Cont.). Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game	Cotton-tail Rabbit		White Hare		Squirrel		Fox		Subtotal Mammals		Area** Totals	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
Area - Dates												
<u>Crane</u>												
Opening Day											29	22
Saturday (4)	2	2							2	2	134	159
Weekday after stocking (2)		2								2	60	28
Weekday after no stocking (2)											7	1
Totals	2	4							2	4	230	210
<u>Freetown*</u>												
Opening Day												18
Saturday (4)							1			1		56
Weekday after no stocking (2)												
Totals							1			1		74
<u>Myles Standish*</u>												
Opening Day												31
Saturday (4)		2								2		162
Weekday after stocking (2)		1								1		47
Weekday after no stocking (2)												
Totals		3								3		240
<u>Northeast</u>												
Opening Day											32	32
Saturday (4)	2	5		1		1W	1		3	7	137	196
Weekday after stocking (2)						1G				1	48	63
Weekday after no stocking (2)	1								1		8	6
Totals	3	5		1		2	1		4	8	225	297
<u>Crane Pond*</u>												
Opening Day												6
Saturday (4)												60
Weekday after stocking (2)												7
Weekday after no stocking (2)												2
Totals												75
<u>Game Harvest -</u>												
Mammals (1)	5	12		1		2	1	1	6	16	455	896

* Area not sampled in 1971

** Includes birds total carried forward

W = Woodchuck

G = Gray squirrel

Table 7. (Cont.). Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game Area - Dates	Cotton- tail Rabbit		White Hare		Squirrel		Fox		Subtotal Mammals		Area*** Totals	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
<u>Housatonic Valley</u>												
Opening Day		1							1		25	17
Saturday (4)	2	4							2	4	54	63
Weekday after stocking (1)**	1								1		21	14
Weekday after no stocking (1)											5	1
Totals	3	5							3	5	105	95
<u>Stafford Hill*</u>												
Opening Day												8
Saturday (4)		1								1		32
Weekday after stocking (1)**												4
Weekday after no stocking (1)												3
Totals		1								1		47
<u>Barre Falls*</u>												
Opening Day												9
Saturday (4)				1						1		17
Weekday after stocking (2)												6
Weekday after no stocking (2)				1						1		3
Totals				2						2		35
<u>Birch Hill*</u>												
Opening Day		1								1		30
Saturday (4)				12		1G				13		224
Weekday after stocking (2)												40
Weekday after no stocking (2)												22
Totals		1		12		1				14		316
<u>Hubbardston*</u>												
Opening Day		1								1		14
Saturday (4)												87
Weekday after stocking (2)												14
Weekday after no stocking (2)		1								1		4
Totals		2								2		119
<u>Game Harvest</u>												
Mammals (2)	3	9		14		1			3	24	105	612

* Area not sampled in 1971 ** Two weekdays each category in 1971, one each in 1972.

*** Includes birds total carried forward.

G = Gray squirrel

Table 7 (Cont.). Game Harvest on Twelve Wildlife Management Areas, 1971 and 1972.

Game Area - Dates	Cotton- tail Rabbit		White Hare		Squirrel		Fox		Subtotal Mammals		Area** Totals	
	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972	1971	1972
<u>West Hill*</u>												
Opening Day												16
Saturday (4)		2								2		79
Weekday after stocking (2)												12
Weekday after no stocking (2)												5
Totals		2								2		112
<u>Winimusset</u>												
Opening Day		1								1	41	12
Saturday (4)	2								2		117	79
Weekday after stocking (2)											16	12
Weekday after no stocking (2)											17	14
Totals	2	1							2	1	191	117
<u>Game Harvest -</u>												
Mammals (3)	2	3							2	3	191	229
Mammals (2)	3	9		14		1			3	24	105	612
Mammals (1)	5	12		1		2	1	1	6	16	455	896
Total Mammals	10	24		15		3	1	1	11	43		
Total Harvest											751	1737

* Area not sampled in 1971.

** Includes birds total carried forward.

Woodcock were taken on six areas, with the largest harvest again at Birch Hill (22).

Cottontail rabbit and white hare were harvested on nine and three areas respectively, with Northeast and Housatonic Valley tied at five each for the most cottontails, and Birch Hill providing the most hare (12).

Harvest of the remaining species of game was scattered.

For the four areas comparable to 1971 the harvest decreased slightly on Crane and Housatonic Valley, decreased considerably at Winimusset (189 to 116), and increased somewhat at Northeast. Since stocked rather than native game is the primary attraction at management areas, these fluctuations in harvest are likely attributable to changes in success due to increased competition, or to changes in hunting pressure due to weather conditions, rather than to fluctuations in native game populations.

The game harvest for Fort Devens from 1969 to 1972 is presented in Table 8. This represents a complete count, since military regulations require sportsmen to check in and out of the area.

Hunter success for the twelve sampled wildlife management areas is presented in Table 9. Success percentages were calculated for opening day, Saturdays (four combined), days after stocking (two combined), and days after no stocking (two combined) for each management area, and for the total of all areas combined.

Combined hunter success for all areas was 35.3 percent. Combined success for the four areas (Crane, Northeast, Housatonic Valley, and Winimusset) comparable to previous years was 35.8%, the highest percentage attained during the past three seasons (1971, 32.5%; 1970, 29.2%).

Success was greatest on days after stocking (45.7%), followed by Saturdays (36.7%), opening day (28.2%), and days after no stocking (22.3%). For the twelve areas combined, and for the four areas comparable to previous years, these percentages show a decrease in opening day success, and an increase in success for the remaining three categories of days.

For comparable areas, success increased by 10 percent at Crane, and decreased slightly at Northeast and Housatonic Valley. Winimusset showed a decrease of 6.7 percent. Overall success was greatest at Birch Hill (54.1%), followed by Crane (51.4%), Myles Standish (49.2%), Hubbardston (43.0%), and Winimusset (37.7%). On the remaining areas, less than one-third of the contacted hunters were successful, with the poorest success being recorded at Barre Falls (13.6%).

Table 8. Game Harvest at Fort Devens Wildlife Management Area: 1969-1972.

Game	Number Taken Per Check Period			
	10 Oct. to 30 Nov. 1969	8 Dec. 1969 to 15 Dec. 1970	11 Oct. to 28 Dec. 1971	10 Oct. to 31 Dec. 1972
Pheasant	751	676	461	391
Ruffed Grouse	62	80	46	61
Woodcock	112	110	86	162
Ducks (inclusive)	42	37	39	36
Cottontail Rabbit	10	58	34	30
White Hare	9	32	23	23
Gray Squirrel	33	147	94	256
Raccoon			7	3
Fox		1	2	
White-Tailed Deer		2	3	
Total Harvest	1,019	1,143	795	962

Table 9. Hunter Success on Twelve Wildlife Management Areas: 1970-1972.

Area - Dates	1972			Percent Successful 1971	Percent Successful 1970
	Number of Hunters Contacted	Number Taking at Least One Unit of Game	Percent Successful		
<u>Crane</u>					
Opening Day	63	32	50.8	38.5	62.5
Saturday (4)	299	156	52.2	41.5	33.4
Weekday after stocking (2)	41	24	58.5	54.9	40.8
Weekday after no stocking (2)	11	1	9.1	11.8	19.0
Totals	414	213	51.4	41.4	35.7
<u>Freetown*</u>					
Opening Day	70	17	24.3		
Saturday (4)	199	55	27.6		
Weekday after no stocking (2)	23	0	0.0		
Totals	292	72	24.6		
<u>Myles Standish*</u>					
Opening Day	42	18	42.8		
Saturday (4)	203	109	53.7		
Weekday after stocking (2)	57	25	43.8		
Weekday after no stocking (2)	7	0	0.0		
Totals	309	152	49.2		
<u>Northeast</u>					
Opening Day	121	27	22.3	29.3	26.3
Saturday (4)	593	160	27.0	29.0	28.2
Weekday after stocking (2)	96	52	54.2	41.2	34.0
Weekday after no stocking (2)	42	6	14.3	18.2	11.1
Totals	852	245	28.8	29.8	28.0
<u>Crane Pond*</u>					
Opening Day	35	5	14.3		
Saturday (4)	171	49	28.6		
Weekday after stocking (2)	18	6	33.3		
Weekday after no stocking (2)	19	2	10.5		
Totals	243	62	25.5		

* Area not sampled prior to 1972.

Table 9 (Cont.). Hunter Success on Twelve Wildlife Management Areas: 1970-1972.

Area-Dates	1972			Percent Successful 1971	Percent Successful 1970
	Number of Hunters Contacted	Number Taking at Least One Unit of Game	Percent Successful		
<u>Housatonic Valley</u>					
Opening Day	47	13	27.6	36.7	31.8
Saturday (4)	172	51	29.6	35.7	27.1
Weekday after stocking (1)**	15	10	66.7	29.8	43.8
Weekday after no stocking (1)	3	1	33.3	14.7	9.1
Totals	237	75	31.6	32.1	28.3
<u>Stafford Hill*</u>					
Opening Day	42	8	19.0		
Saturday (4)	122	27	22.1		
Weekday after stocking (1)**	17	4	23.5		
Weekday after no stocking (1)	9	3	33.3		
Totals	190	42	22.1		
<u>Barre Falls*</u>					
Opening Day	45	9	20.0		
Saturday (4)	177	18	10.2		
Weekday after stocking (1)	20	6	30.0		
Weekday after no stocking (1)	22	3	13.6		
Totals	264	36	13.6		
<u>Birch Hill*</u>					
Opening Day	81	30	37.0		
Saturday (4)	378	226	59.8		
Weekday after stocking (2)	81	40	49.4		
Weekday after no stocking (2)	48	22	45.8		
Totals	588	318	54.1		
<u>Hubbardston*</u>					
Opening Day	29	14	48.3		
Saturday (4)	163	73	44.8		
Weekday after stocking (2)	15	10	66.7		
Weekday after no stocking (2)	28	4	14.3		
Totals	235	101	43.0		

* Area not sampled prior to 1972.

** One weekday each category in 1972,
two each in previous years.

Table 9 (Cont.). Hunter Success on Twelve Wildlife Management Areas: 1970-1972.

Area - Dates	1972			Percent Successful 1971	Percent Successful 1970
	Number of Hunters Contacted	Number Taking At Least One Unit of Game	Percent Successful		
<u>West Hill*</u>					
Opening Day	67	9	13.4		
Saturday (4)	252	75	29.8		
Weekday after stocking (2)	42	11	26.2		
Weekday after no stocking (2)	43	5	11.6		
Totals	404	100	24.8		
<u>Winimusset</u>					
Opening Day	31	8	25.8	55.5	26.8
Saturday (4)	163	63	38.6	43.5	36.1
Weekday after stocking (2)	29	9	31.0	52.6	42.2
Weekday after no stocking (2)	37	18	48.6	28.1	20.0
Totals	260	98	37.7	44.4	35.0
<u>Totals (all areas)**</u>					
Opening Day	673 (262)	190 (80)	28.2 (30.5)	37.1	32.5
Saturday	2392 (1227)	1062 (430)	36.7 (35.0)	31.3	28.9
Weekday after stocking	431 (181)	197 (95)	45.7 (52.5)	41.7	35.2
Weekday after no stocking	292 (93)	65 (26)	22.3 (28.0)	17.9	12.1
Total (all days)	4288 (1763)	1514 (631)	35.3 (35.8)	32.5	29.2

* Area not sampled prior to 1972.

** Comparable area figures in parentheses.

TABLE I			
Year	Population	Area	Population Density
1900	1,000,000	100,000	10
1910	1,500,000	120,000	12.5
1920	2,000,000	140,000	14.3
1930	2,500,000	160,000	15.6
1940	3,000,000	180,000	16.7
1950	3,500,000	200,000	17.5
1960	4,000,000	220,000	18.2
1970	4,500,000	240,000	18.8
1980	5,000,000	260,000	19.2
1990	5,500,000	280,000	19.6
2000	6,000,000	300,000	20.0
2010	6,500,000	320,000	20.3
2020	7,000,000	340,000	20.6

Source: U.S. Census Bureau, 2020. Data for 1900-1990 are from the U.S. Census Bureau, 1992. Data for 2000-2020 are from the U.S. Census Bureau, 2020.

Opening day success exceeded 33 percent on four areas (Crane, Myles Standish, Birch Hill, and Hubbardston). The highest success for a single area day-category was at Housatonic Valley and Hubbardston with day-after-stocking percentages of 66.7.

- Recommendations:
1. For 1973, discontinue checks on the Crane Pond area and initiate checks on the Delaney area.
 2. Consider shifting utilization checks from an annual to a biennial schedule in 1974.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved by _____
Colton H. Bridges
Title Superintendent

Prepared by _____
James E. Cardoza
Asst. Game Biologist

Date _____

THE UNIVERSITY OF CHICAGO
 

PERFORMANCE REPORT

State: Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-15 Project Title: Game Population Trend and Harvest Survey

Job No.: VI-1 Job Title: Black Bear Population Dynamics

Period Covered: 1 June 1972 to 31 May 1973

Summary: For the first time since 1969, a bear was legally taken in Massachusetts. The bear, a female, was taken in Savoy on the first day of the season. Another bear was killed illegally, in Royalston, during deer week.

There were 423 bear permit applications in 1972, the greatest number since the requirement was initiated in 1970. Applicants were contacted by postal questionnaire, and a response of 345 (82.1%) was achieved. There were 213 persons who did hunt bear, of whom 96 had hunted bear in previous years. The average bear hunter expended 16.8 hours in 2.4 days.

Reports of bear were collected from questionnaire results, and Division and cooperator records. To date, reports of 449 bear noted on 366 occasions in 73 towns have been collected for the period 1952-May 1973. Franklin County yielded the majority of reports (143, 40.4%) followed by Berkshire (124, 33.9%), Hampshire (59, 16.1%), Hampden (18, 4.9%), Worcester (16, 4.4%), and Middlesex (1, 0.3%).

Objectives: To define the range of the black bear in Massachusetts and to determine its population characteristics and rate of harvest by hunting.

Procedures: Current bear hunting regulations include mandatory reporting and tagging of bear. Bear checking stations were maintained daily during bear week at three locations (Birch Hill, Templeton; Montague Fish Hatchery, Montague; and Western Wildlife District Headquarters, Pittsfield). Station personnel were directed to affix a metal game seal to harvested bears and to determine the following: town bear killed in, date killed, sex and weight of bear, and method of kill. A mimeographed form was provided for recording this information. Successful hunters were subsequently contacted by the project leader to remove a tooth for sectioning.

[illegible]

Bear hunters were required to obtain a free (fifty cents beginning in 1973) bear hunting permit from the Division of Fisheries and Game. A total of 423 permits were issued for the 1972 season. Permittees were subsequently contacted by postal questionnaire to determine their interest and participation in bear hunting. Applicants were also asked to list any bear sightings they had made on previous occasions.

The Information and Education Section issued periodic news releases asking for reports of black bear. Responses to these requests, replies to the questionnaire, and voluntary reports from other individuals were tabulated and mapped as an aid to determining bear distribution.

Findings:

For the first time since 1969, a bear was killed, legally in Massachusetts. A 74.8 kilogram (161 pounds) (hog-dressed) female was taken in Savoy on the first day of the season. An additional bear, a male, was shot illegally in Royalston during deer week.

In 1972, there were 423 bear permit applications received from 421 individuals, as opposed to 214 requests in 1970, and 200 in 1971. One 1972 permit was returned as undeliverable. Applicants represented all counties except Dukes. The greatest number of applications from a single county was from Hampden (90, 21.4%), followed by Berkshire (63, 15.0%), Middlesex (58, 13.8%), Worcester (46, 11.0%) and Franklin (38, 9.1%). Eight other counties and two other states (New York and Connecticut) comprised the remaining 128 applications.

A questionnaire was mailed to each of the 420 individuals receiving a bear permit. A total return of 345 was obtained from two mailings of the questionnaire (Table 1).

In 1972, 213 individuals stated that they hunted black bear during that season (Table 2). Of these, 187 indicated that they hunted specifically for bear, while the remaining 26 replied that they only incidentally pursued bear while bow-hunting for deer. An additional 40 persons replied that they had hunted bear in previous years, but were unable or unwilling to go in 1972.

There were 116 hunters (54.5%) who stated that they hunted bear for the first time in 1972, as opposed to 49 (52.1%) in 1970.

Ten hunters stated that they had bagged a bear in Massachusetts in previous years.

No special attempt was made to determine the reasons for lack of participation by non-hunters. Some volunteered this information and stated that they wished to hunt, but were ill or out of town. Others felt the season was too short, and did not allow them enough time for a worthwhile hunt.

Table 1. Summation of Responses to 1970 and 1972 Bear Questionnaires

	<u>1970</u>	<u>1972</u>
Total number of permits issued	214	420
Returned first mailing	131	207
Returned second mailing	46	138
	<u> </u>	<u> </u>
	177 (82.7%)	345 (82.1%)

Table 2. Breakdown of Responses to 1970 and 1972 Bear Questionnaires

	<u>1970</u>	<u>1972</u>
A. Hunted bear in specified year	94 (53.1%)	213 (61.7%)
Did not hunt bear in specified year	80 (45.2%)	123 (35.7%)
Returned unusable	2) (1.7%)	5) (2.6%)
Undeliverable	1)	4)
	<u> </u>	<u> </u>
	177	345
 B. Hunted bear for the first time in specified year	49 (52.1%)	116 (54.5%)
Hunted bear in specified year - has hunted previously	45 (47.9%)	96 (45.1%)
No response	0	1 (0.4%)
	<u> </u>	<u> </u>
	94	213
 Hunted in previous years - did not hunt in specified year	25	40
	<u> </u>	<u> </u>
	119	253

Bear hunters were further asked to account for the time they expended while bear hunting. Total expenditure by the 210 hunters replying to this question was 3507 hours in 513 days. This is a mean of 16.8 hours in 2.4 days, per hunter. The unweighted mean number of hours per day per hunter was 6.76 (6.22 in 1970).

The bear season ran for six days, from Monday, 20 November to Saturday, 25 November. Hunter effort was greatest on Saturday (151 hunters, 30.8%), followed by Friday (84, 17.1%), Monday (83, 16.9%), Wednesday (62, 12.6%), Tuesday (11.4%), and Thursday (11.2%). Pressure on Friday may have been skewed upward by the fact that Thursday was Thanksgiving Day, and some people had Friday as an additional holiday.

Ten hunters (plus two who did not hunt in 1972) indicated that they pursued bear with dogs. The remaining hunters searched about in areas where they had seen bear sign, or followed bear tracks, or took a stand in an area where bears had been seen.

Sixteen hunters saw a bear during the legal season. One of these succeeded in killing a bear, two others treed a bear, but refrained from killing it, and two others (both bow-hunters) shot and missed. The remainder either stated that the bear was out of range, or gave no reason for not shooting.

Hunters pursued bear in 63 towns in five counties during the 1972 season. Hunting pressure was greatest in Franklin County (128 hunters, 44.0%), followed by Berkshire (99, 34.0%), Hampshire (32, 11.0%), Hampden (17, 5.8%), and Worcester (15, 5.2%). Some sportsmen hunted more than one county, consequently, they are counted more than once.

Hunters were asked if they investigated an area for signs of bear prior to the season. There were 200 responses to this question, of which 148 indicated the hunter did check out an area.

Hunters were further asked to state the type of weapon they used. All 213 of the 1972 hunters replied. A rifle was the predominant weapon (167), followed by bow and arrow (26). Thirteen used both rifle and bow (on different days), six employed a shotgun (an illegal weapon), and one hunter used no weapon, preferring only to run his dogs on bear.

Reports of bear sightings or bear tracks were listed by 178 persons. These are included in the following section on bear sightings.

Additional comments or remarks were presented by 164 respondents. The most prevalent of these are tabulated below (Table 3).

 

Table 3. Hunter Comments, 1972 Bear Questionnaire.

<u>Category/Comment</u>	<u>Number Hunters</u>
Season length -	71
Extend season	62
Close bear season	5
Other comments (3)	4
Timing of Season -	48
Have bear season same as deer season	15
Have earlier season	10
Don't have bear season same as deer season	5
Don't have bear season same as Vermont	5
Don't have bear season during archery season	5
Have later season	5
Other comments (3)	4
Regulations -	16
Allow Sunday hunting	5
Allow use of shotgun	5
Allow sidearms	4
Other comments (2)	4
Administration -	21
Provide information on bears and bear hunting	16
Other comments (5)	5
Management -	7
Stock bear	6
Maintain habitat	1
Populations -	6
Bear increasing	2
Bear decreasing	2
Other comments (2)	2
General Comments -	31
Volunteers assistance	10
Favors study of bear	8
Other comments (11)	13

All available recent (since 1950) reports of black bear are being aggregated and mapped to aid in determining bear distribution in Massachusetts. To date, records of 449 bears reported on 366 occasions in 73 towns have been collected.

Bear reports were categorized as sightings (224), sign and tracks (117), kills (23) and other (road kill, nuisance kill) (2).

The distribution and approximate density of bear reports in 73 towns in six counties is presented in Figure 1. The number of reports per county and the highest towns per county are listed in Table 4.

Table 4. **Black Bear Reports by County and Town,**
 1952-1973

A.	Berkshire: 124 (33.9%)	
	Florida	(29)
	Hancock	(7)
	Peru	(6)
	Savoy	(18)
	Windsor	(11)
	Other 21 towns	(53)
B.	Middlesex: 1 (0.3%)	
	Ashby	(1)
C.	Franklin: 148 (40.4%)	
	Ashfield	(18)
	Charlemont	(12)
	Colrain	(15)
	Hawley	(29)
	Monroe	(13)
	Rowe	(17)
	Other 13 towns	(44)
D.	Hampden: 18 (4.9%)	
	Blandford	(6)
	Granville	(4)
	Russell	(4)
	Other 3 towns	(4)
E.	Hampshire: 59 (16.1%)	
	Chesterfield	(19)
	Cummington	(10)
	Huntington	(6)
	Worthington	(8)
	Other 8 towns	(16)
F.	Worcester: 16 (4.4%)	
	Petersham	(3)
	Warren	(3)
	Other 7 towns	(10)

THE UNIVERSITY OF CHICAGO
LIBRARY

100 EAST HASTING STREET
CHICAGO, ILL. 60607

TEL: 773-936-5000
FAX: 773-936-5001

INTERNET: <http://www.lib.uchicago.edu>

ACQUISITIONS	
BOOKS	100
JOURNALS	50
PERIODICALS	200
MANUSCRIPTS	10
ARCHIVES	5
DATA	10
VIDEO	10
CD-ROM	10
SOFTWARE	10
OTHER	10
TOTAL	405

Until 1970, hunters taking a bear were not required to report their kill to the Division. An approximation of the past harvest has been gathered through questionnaire results and voluntary reports (Table 5).

Table 5. Known Black Bear Harvest, 1957-1972*

<u>Town</u>	<u>Date</u>
Peru	Deer week 1957
Heath	1958
Charlemont	1961
Florida	1964
Florida	1965
Savoy	1965
Savoy	December 1966
Florida	1967
Florida	1968
Rowe	Fall 1968
Savoy	Fall 1969
Monroe	November 1969
Florida	17 November 1969
Rowe	3 December 1969
Chesterfield	Deer week 1969
Hawley	Deer week 1969
Monroe	Deer week 1969
Windsor	Deer week 1969
Worthington	Deer week 1969
Charlemont	Deer week 1970**
Savoy	20 November 1972**
Royalston	Deer week 1972**

* Does not include one unconfirmed illegal kill in 1971.

** Illegal kill

- Recommendations:
1. Continue gathering reports of bear sightings, and checking of harvested bear.
 2. Aggregate data on current distribution and status and historical data collected under Job VI-2, and publish results in a research bulletin on the history and status of bear in Massachusetts. More detailed recommendations on future bear management will be presented at that time.

 

MASSACHUSETTS

CITY AND TOWN LINES

Figure 1. Reports of Black Bear in Massachusetts, 1952- May 1973.

- 1- 3 Reports.
- 4-10 Reports.
- 11+ Reports.

Department of Conservation

F-35-R-15:VI-1

Acknowledgements: I extend my appreciation to Mr. William Zurrin of
Pittsfield for his contribution of a bear skull.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved by _____
Colton H. Bridges
Title _____
Superintendent

Prepared by _____
James E. Cardoza
Asst. Game Biologist

Date _____

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No.: I-2 Job Title: Statewide Beaver Harvest

Period Covered: 1 November 1973 to 31 May 1974

Summary: A total of 1,639 beaver were trapped by 123 trappers in 106 towns during the 1973-1974 beaver season. This take, the second highest on record, is 35 less than last season and 456 more than a ten-year (1964-1973) average. Berkshire and Franklin Counties together yielded 888 beaver (54.2 percent of harvest).

Over one half (53.6 percent) of the beaver were taken in the first two weeks of the 15-week season.

Objectives: To determine the statewide harvest of beavers by trappers.

Procedures: Each successful beaver trapper is required by law to present his pelts to an official checking station for tagging and recording of data. Seven stations (six prior to 1974) are maintained for two days at the close of the season. Pelts are tagged with locking metal game seals and harvest data are recorded on mimeographed forms and subsequently tabulated by month trapped, town and county trapped in, and type of trap used.

Findings: The 1973-1974 beaver trapping season extended for 15 weeks from 15 November 1973 to 1 March 1974. During this period, trappers took 1,639 beaver. This take was only 35 less than last season's record take and was 456 more than a ten-year average (1964-1973). There were 123 trappers (105 in 1972-1973) taking a minimum of one beaver each, with a mean harvest of 13.3 beaver per trapper (range: 1 to 128).

Beaver colonies located in 106 towns contributed to the 1973-1974 harvest (Figure 1). The fifteen towns with the largest individual seasonal harvest are listed in Table 1.

The western region of the state continued to provide the majority of the beaver harvest. However, during the current season, the take in the eastern region increased slightly over that of the previous season. During the 1973-1974 season, 1,058 beaver (64.6 percent) were taken west of the Connecticut River and 581 (35.4 percent) were taken east of the river, as opposed to 1,148 (68.6 percent) and 526 (31.4 percent) during 1972-1973.

MASSACHUSETTS

CITY AND TOWN LINES

Figure 1.
Beaver Trapped, by Towns, 1973-1974.
Total-- 1639 Beaver.

Table 1. Beaver Harvest for Fifteen Towns, 1969-1974 Seasons

<u>Town</u>	<u>1968-69</u>	<u>1969-70</u>	<u>1970-71</u>	<u>1971-72</u>	<u>1972-73</u>	<u>1973-74</u>
Ashfield	20	14	2	53	50	60
Becket	41	20	10	51	57	52
Blandford	10	22	16	29	78	37
Cummington	13	11	5	8	2	38
New Marlboro	32	14	1	52	69	64
New Salem	11	10	27	10	56	61
Otis	36	18	14	64	72	72
Petersham	52	18	27	58	57	66
Sandisfield	49	13	8	52	6	62
Sheffield	30	12	7	10	26	37
Shutesbury	5	4	-	8	20	40
Tolland	31	-	2	37	48	35
Washington	9	27	7	55	27	37
Windsor	16	19	14	36	26	41
Worthington	7	30	14	52	54	79

1. *Table of Contents*
 2. *Introduction*
 3. *Methodology*
 4. *Results*
 5. *Discussion*
 6. *Conclusion*
 7. *References*
 8. *Appendix*
 9. *Glossary*
 10. *Index*

Page	Chapter	Section	Page	Page	Page	Page
1	1	1.1	1	1	1	1
2	2	2.1	2	2	2	2
3	3	3.1	3	3	3	3
4	4	4.1	4	4	4	4
5	5	5.1	5	5	5	5
6	6	6.1	6	6	6	6
7	7	7.1	7	7	7	7
8	8	8.1	8	8	8	8
9	9	9.1	9	9	9	9
10	10	10.1	10	10	10	10
11	11	11.1	11	11	11	11
12	12	12.1	12	12	12	12
13	13	13.1	13	13	13	13
14	14	14.1	14	14	14	14
15	15	15.1	15	15	15	15
16	16	16.1	16	16	16	16
17	17	17.1	17	17	17	17
18	18	18.1	18	18	18	18
19	19	19.1	19	19	19	19
20	20	20.1	20	20	20	20
21	21	21.1	21	21	21	21
22	22	22.1	22	22	22	22
23	23	23.1	23	23	23	23
24	24	24.1	24	24	24	24
25	25	25.1	25	25	25	25
26	26	26.1	26	26	26	26
27	27	27.1	27	27	27	27
28	28	28.1	28	28	28	28
29	29	29.1	29	29	29	29
30	30	30.1	30	30	30	30
31	31	31.1	31	31	31	31
32	32	32.1	32	32	32	32
33	33	33.1	33	33	33	33
34	34	34.1	34	34	34	34
35	35	35.1	35	35	35	35
36	36	36.1	36	36	36	36
37	37	37.1	37	37	37	37
38	38	38.1	38	38	38	38
39	39	39.1	39	39	39	39
40	40	40.1	40	40	40	40
41	41	41.1	41	41	41	41
42	42	42.1	42	42	42	42
43	43	43.1	43	43	43	43
44	44	44.1	44	44	44	44
45	45	45.1	45	45	45	45
46	46	46.1	46	46	46	46
47	47	47.1	47	47	47	47
48	48	48.1	48	48	48	48
49	49	49.1	49	49	49	49
50	50	50.1	50	50	50	50
51	51	51.1	51	51	51	51
52	52	52.1	52	52	52	52
53	53	53.1	53	53	53	53
54	54	54.1	54	54	54	54
55	55	55.1	55	55	55	55
56	56	56.1	56	56	56	56
57	57	57.1	57	57	57	57
58	58	58.1	58	58	58	58
59	59	59.1	59	59	59	59
60	60	60.1	60	60	60	60
61	61	61.1	61	61	61	61
62	62	62.1	62	62	62	62
63	63	63.1	63	63	63	63
64	64	64.1	64	64	64	64
65	65	65.1	65	65	65	65
66	66	66.1	66	66	66	66
67	67	67.1	67	67	67	67
68	68	68.1	68	68	68	68
69	69	69.1	69	69	69	69
70	70	70.1	70	70	70	70
71	71	71.1	71	71	71	71
72	72	72.1	72	72	72	72
73	73	73.1	73	73	73	73
74	74	74.1	74	74	74	74
75	75	75.1	75	75	75	75
76	76	76.1	76	76	76	76
77	77	77.1	77	77	77	77
78	78	78.1	78	78	78	78
79	79	79.1	79	79	79	79
80	80	80.1	80	80	80	80
81	81	81.1	81	81	81	81
82	82	82.1	82	82	82	82
83	83	83.1	83	83	83	83
84	84	84.1	84	84	84	84
85	85	85.1	85	85	85	85
86	86	86.1	86	86	86	86
87	87	87.1	87	87	87	87
88	88	88.1	88	88	88	88
89	89	89.1	89	89	89	89
90	90	90.1	90	90	90	90
91	91	91.1	91	91	91	91
92	92	92.1	92	92	92	92
93	93	93.1	93	93	93	93
94	94	94.1	94	94	94	94
95	95	95.1	95	95	95	95
96	96	96.1	96	96	96	96
97	97	97.1	97	97	97	97
98	98	98.1	98	98	98	98
99	99	99.1	99	99	99	99
100	100	100.1	100	100	100	100

Table 2. Beaver Harvest by County, 1972-73 and 1973-74

<u>County</u>	1972-73			1973-74		
	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>
Berkshire	461	27.5	1	518	31.6	1
Essex	15	0.9	7	16	1.0	8
Franklin	446	26.6	2	370	22.6	2
Hampden	227	13.6	5	148	9.0	5
Hampshire	253	15.1	3	288	17.6	3
Middlesex	35	2.1	6	51	3.1	6
Plymouth	5	0.3	8	17	1.0	7
Worcester	232	13.9	4	231	14.1	4
Totals	1,674	100.0		1,639	100.0	

Table 3. Beaver Harvest by Month, 1972-73 and 1973-74

<u>Month</u>	1972-73		1973-74	
	<u>No. Beaver</u>	<u>Percent</u>	<u>No. Beaver</u>	<u>Percent</u>
November	633	37.8	878	53.6
December	544	32.5	492	30.0
January	356	21.3	185	11.3
February	141	8.4	84	5.1
Totals	1,674	100.0	1,639	100.0

TABLE 1. Summary of the results of the analysis of variance for the effect of the treatment on the response of the subjects to the test.

Treatment		Response		Significance	
Group	Mean	SD	Group	Mean	SD
1	10.5	1.2	1	10.5	1.2
2	11.2	1.5	2	11.2	1.5
3	12.0	1.8	3	12.0	1.8
4	13.5	2.0	4	13.5	2.0
5	14.0	2.2	5	14.0	2.2
6	15.0	2.5	6	15.0	2.5
7	16.0	2.8	7	16.0	2.8
8	17.0	3.0	8	17.0	3.0
9	18.0	3.2	9	18.0	3.2
10	19.0	3.5	10	19.0	3.5

TABLE 2. Summary of the results of the analysis of variance for the effect of the treatment on the response of the subjects to the test.

Treatment		Response		Significance	
Group	Mean	SD	Group	Mean	SD
1	10.5	1.2	1	10.5	1.2
2	11.2	1.5	2	11.2	1.5
3	12.0	1.8	3	12.0	1.8
4	13.5	2.0	4	13.5	2.0
5	14.0	2.2	5	14.0	2.2
6	15.0	2.5	6	15.0	2.5
7	16.0	2.8	7	16.0	2.8
8	17.0	3.0	8	17.0	3.0
9	18.0	3.2	9	18.0	3.2
10	19.0	3.5	10	19.0	3.5

During the past season, Berkshire and Franklin Counties together yielded over half (888, 54.2 percent) of the total harvest. Hampshire, Worcester, and Hampden Counties contributed another 667 beaver (40.7 percent) with three additional counties comprising the small remainder (Table 2). Six counties reported no beaver taken. The harvest in Berkshire County increased by 12.4 percent over 1972-1973 and in Hampshire County by 13.8 percent, while the take in Franklin and Hampden Counties decreased by 17.0 percent and 34.8 percent respectively. County rankings remained essentially unchanged.

As in past years, success was greatest during the initial weeks of the trapping season (Table 3). Over one half (54.2 percent) of the beaver were taken in the initial two weeks (15-30 November) of the season, as opposed to approximately one third (37.8 percent) of the 1972-1973 take occurring during the equivalent period. The take declined each month thereafter, with only a minimal harvest (5.1 percent) in February.

Use of the Conibear trap remains relatively constant, with two thirds (1,091, 66.6 percent) of the 1973-1974 harvest gained by use of this trap.

The average price of a Massachusetts beaver pelt declined slightly from \$20 in 1972-1973 to the current price of \$18. Due to the near-record total harvest, however, the total season valuation of \$29,502 is the second highest on record.

- Recommendations:
1. Continue tagging of beaver pelts and recording of harvest data in 1975, using the same methods as in the current segment.
 2. There has been some comment from trappers concerning the travel distance required to check pelts. Should this continue to be a difficulty, considering the current energy situation, some alleviation may be provided by permitting cooperating sporting goods shops to tag beaver. Precedent for this has been established by mercantile cooperation in the checking of deer.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____

Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Assistant Game Biologist

Date _____

JOB PERFORMANCE REPORT

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No.: II-1 Job Title: Statewide Deer Harvest

Period Covered: 5 November through 12 December 1973

Summary: The dates, type of hunting season, and the deer harvest by season for 1973 is as follows:

November 5 and 6 - Paraplegic season - total harvest, 0; November 5 through November 24 - archery season - total harvest, 77; December 3 through December 8 - shotgun season, total harvest, 2037; and December 10 through December 12 - smooth bore muzzle loader season - total harvest, 7. Grand total - 2121.

The following number and type of antlerless deer permits were issued in 1973: Sportsman - 4000; Farmer-Landowner - 349; Nantucket - 400; and Martha's Vineyard - 600.

The cost of an antlerless permit application was 50 cents in 1973. Prior to this time, the application was free.

During the 18-day bow season, archers harvested 77 deer. Of these, 51 were males, 26 were females.

Shotgun deer hunters harvested 2,037 deer. Of 1,654 taken on the mainland, 1,114 were adult males, 106 were button bucks and 434 were females. On Nantucket Island, shotgun hunters harvested 68 adult males, 22 button bucks and 65 females; a total of 155 deer. Two hundred one deer were taken on Martha's Vineyard. Of these, 74 were adult males, 31 were button bucks and 96 were females. The Gosnold Island harvest was 2 adult males, 4 button bucks and 21 females.

Five males and 2 females were reported taken during the special three-day muzzle loader season.

The total statewide 1973 shotgun season harvest was 2,037 deer consisting of 1,421 males and 616 females.

The top three deer producing counties were Berkshire, Franklin, and Dukes. Sixty-three percent of the statewide harvest was removed from the four western counties (Berkshire, Franklin, Hampshire, Hampden).

Approximately 44 percent of all deer harvested (886) during shotgun week were taken on Monday, the first day of the season. Of the 886 deer, 663 (75%) were males and 223 (25%) were females.

The following is a breakdown of the 1973 shotgun deer harvest by management unit:

	<u>Mainland</u>	<u>Nantucket</u>	<u>Martha's Vineyard</u>	<u>Gosnold</u>	<u>Total</u>
Adult Males	1,114	68	74	2	1,258
Button Bucks	106	22	31	4	163
Females	434	65	96	21	616
Totals	1,654	155	201	27	2,037

Background: Massachusetts has had an antlerless permit, shotgun only, hunting season since 1967. Male deer with antlers three inches or longer are legal game for all hunting license holders. A hunter must have been issued an antlerless permit to harvest a button buck or a female deer.

The dates of the four Massachusetts deer hunting seasons are as follows:

<u>Date</u>	<u>Type of Season</u>	<u>No. of Days</u>
November 5 and 6	Paraplegic	2
November 5 through November 24	Archery	18
December 3 through December 8	Shotgun	6
December 10 through December 12	Smooth Bore Muzzle Loader	3

Three deer management units have been established; the mainland, Martha's Vineyard and Nantucket. The number of sportsman's antlerless permits issued per unit was as follows: the mainland - 4,000; Martha's Vineyard - 600; and Nantucket - 400. Deer hunting regulations require that all hunters report their deer kill within 24 hours at an official Division deer checking station. There is no Sunday hunting allowed in Massachusetts.

Objectives: To determine the annual harvest of deer in Massachusetts.

Findings: Table 1 presents a ten-year summary (1964-1973) of the annual deer harvest. Data for 1964 and 1965 were reported by hunters on a voluntary basis; mandatory reporting regulations have been in effect since 1966.

There were no deer harvested during the two-day paraplegic hunt, November 5 and 6, 1973. During the 18-day archery season, November 5 through November 23, 1973, archers harvested 51 male and 26 female deer. The archery harvest of 77 deer exceeds the 1972 harvest by one deer.

Table 2 summarizes the archery and primitive weapons harvests. The 1973 shotgun deer harvest was 2,037 animals of both sexes. The total statewide harvest of deer in all season categories was 2,121 deer. Of these, 1,477 were males and 644 were females. The 165 button bucks harvested are included in the male total. A summary of the 1973 deer harvest and the county rank of importance in the harvest is presented in Table 3.

The three top deer-producing counties have remained unchanged for the past three years. Worcester County moved to fourth place while Dukes dropped to the seventh slot. Hampden County has ranked fifth in three of the past five years.

Sixty-four percent of the 2,037 deer harvested were taken in the four western counties--Berkshire, Franklin, Hampden, and Hampshire. Martha's Vineyard and Nantucket accounted for 19.4 percent of the harvest (396 deer), while Worcester County, largest in the state, yielded 211 deer (10.4%). The remaining eastern counties accounted for 6.2 percent of the total harvest (136 deer).

Table 4 shows the daily harvest of deer for both sexes. Eight hundred eighty-six deer or 43.5 percent of all deer harvested are taken on the first day of the season. Three-fourths of the first day's harvest is in males. The number of males harvested exceeds the female harvest throughout the shotgun season. However, the percent of total female harvest in the last four days of the season exceeds the corresponding daily percent of total harvest in the male segment of the kill.

The most significant data in Table 4 is the comparable rate of harvest of females and button bucks. Apparently, a hunter with an antlerless permit does not show selectivity with respect to the size of the deer he harvests, but the data in Table 4 do suggest that a hunter with an antlerless deer permit may pass up an antlerless deer early in the season and shoot any antlerless deer during the last two days of the season.

Table 5 presents a number of deer harvest summaries of archery and shotgun seasons from 1967 through 1973. Included in Table 5 is the success ratio per type of antlerless permit.

There was no significant increase in the 1973 archery harvest of 77 deer compared to the 1972 harvest of 76 deer (Table 5). The breakdown of the deer harvest by management units shows a slight increase in adult males on the mainland with the predicted reduced harvest of button bucks and females (Table 5). On Nantucket Island, the harvest by antlerless permit holders increased from 101 deer in 1972 to 115 in 1973. An analysis of this harvest indicates the predicted drop in adult male harvest did not occur. However, the anticipated increase of button bucks in the harvest by

Table 3. County Summary of the 1973 Massachusetts Shotgun Deer Harvest By Sex, and the County Rank of Importance from 1969 through 1973 Shotgun Season Only

<u>County</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>	<u>Rank 1973</u>	<u>Rank 1972</u>	<u>Rank 1971</u>	<u>Rank 1970</u>	<u>Rank 1969</u>
Barnstable	82	20	102	8	8	7	8	8
Berkshire	455	149	604	1	1	1	1	1
Bristol		1	1	12	12	12	12	12
Dukes	121	120	241	3	3	3	7	7
Essex	20		20	9	9	10	9	11
Franklin	242	108	350	2	2	2	2	2
Hampden	153	49	202	5	4	5	6	5
Hampshire	108	30	138	7	6	8	4	4
Middlesex	4	1	5	11	11	9	10	10
Nantucket	90	65	155	6	7	6	5	6
Norfolk				13	13	13	13	13
Plymouth	5	3	8	10	10	11	11	11
Suffolk				13	13	13	13	13
Worcester	<u>141</u>	<u>70</u>	<u>211</u>	4	5	4	3	3
Totals	1421	616	2037					

Table 4. A Summary of the Massachusetts Deer Harvest by Day of Harvest,
3 December through 8 December 1973.

Shotgun Season							
	Mon. Dec. 3	Tues. Dec. 4	Wed. Dec. 5	Thurs. Dec. 6	Fri. Dec. 7	Sat. Dec. 8	Total
<u>Both Sexes</u>							
Total Harvest	886	285	144	139	211	372	2037
Percent	43.50	13.99	7.07	6.82	10.36	18.26	
<u>Males</u>							
Total Harvest	663	209	94	92	136	227	1421
Percent	46.66	14.71	6.62	6.47	9.57	15.97	
<u>Females</u>							
Total Harvest	223	76	50	47	75	145	616
Percent	36.20	12.33	8.12	7.63	12.18	23.54	
<u>Button Bucks</u>							
Total Harvest	53	20	16	13	24	37	163
Percent	32.52	12.27	9.81	7.98	14.72	22.70	

Table 5. A Summary of the Massachusetts Deer Harvest and Hunter Success by Type of Permit, 1967 through 1973.

1973 Paraplegic Season - November 5 and 6 - No deer harvested.

1973 Archery Season - November 5 through November 24 - 51 males; 26 females

Summary of the Archery Seasons, 1967 through 1973:

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Male	17	21	27	24	26	49	51
Female	<u>4</u>	<u>13</u>	<u>10</u>	<u>12</u>	<u>10</u>	<u>27</u>	<u>26</u>
Totals	21	34	37	36	36	76	77

There are an estimated 5,500 archery deer hunters in Massachusetts.

1973 Shotgun Season - Statewide - December 3 through December 8 - 1421 males; 616 females

Summary of the Shotgun Season Harvest by Management Unit:

	<u>Mainland</u>	<u>Nantucket</u>	<u>Martha's Vineyard</u>	<u>Gosnold</u>	<u>Total</u>
Males, adult	1114	68	74	2	1258
Males, button buck	106	22	31	4	163
Females	<u>434</u>	<u>65</u>	<u>96</u>	<u>21</u>	<u>616</u>
Totals	1654	155	201	27	2037

Summary of the Shotgun Deer Harvest (Statewide) from 1967 through 1973:

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Male	937	1083	1424	1605	1359	1455	1421
Female	<u>235</u>	<u>310</u>	<u>585</u>	<u>764</u>	<u>889</u>	<u>760</u>	<u>616</u>
Total	1172	1393	2009	2369	2248	2215	2037

Table 5 (Continued)

Summary of the Archery and Shotgun Seasons (Statewide), 1967 through 1973:

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Male	954	1104	1451	1629	1385	1504	1472
Female	<u>239</u>	<u>323</u>	<u>595</u>	<u>776</u>	<u>899</u>	<u>787</u>	<u>642</u>
Total	1193	1427	2046	2405	2284	2291	2114

Smooth Bore Muzzle Loader Season - December 10 through December 12, 1973:
 2 females; 5 males.

Summary of Antlerless Deer Permits:

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
No. appli- cations (rounded off)	28,000	24,000	32,000	35,000	37,500	38,000	32,000
Issued:							
Sportsman	2,000	2,000	4,000	6,000	6,000	4,000	4,000
Farmer- Landowner	243	331	295	347	270	326	349
Nantucket	-	-	400	400	400	400	400
Martha's Vineyard	-	-	-	-	600	600	637
Totals	2,243	2,331	4,695	6,747	7,270	5,326	5,349

Permit Harvest, Statewide, Landowner-Farmer

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Male	2	21	17	25	20	14	19
Female	<u>20</u>	<u>44</u>	<u>45</u>	<u>34</u>	<u>26</u>	<u>42</u>	<u>37</u>
Total	22	65	62	59	46	56	56

Statewide Harvest by Sportsman's Permit

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Deer (both sexes)	279	356	787	1057	1268	1010	854

Table 5 (Concluded)

Breakdown of Harvest by Sportsman Permits per Management Unit, 1967-1973:

	<u>Mainland</u>						
	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Adult Male	Data not available for 1967 and 1968.		79	163	104	55	67
Button Bucks			107	183	175	142	102
Female			<u>473</u>	<u>612</u>	<u>673</u>	<u>546</u>	<u>397</u>
Totals			659	958	952	743	566
Permits	2000	2000	4000	6000	6000	4000	4000
	<u>Nantucket</u>						
Adult Male			29	15	26	23	19
Button Buck			11	24	29	17	31
Female			<u>49</u>	<u>85</u>	<u>70</u>	<u>61</u>	<u>65</u>
Totals			89	124	125	101	115
Permits			400	400	400	400	400
	<u>Martha's Vineyard and Gosnold</u>						
Adult Male			1	0	30	24	21
Button Buck				1	41	36	35
Female			<u>19</u>	<u>38</u>	<u>95</u>	<u>126</u>	<u>117</u>
Totals			20	39	166	186	173
Permits					600	635*	635*
Antlerless Permit Success Ratio (rounded):							
	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Farmer-Landowner	1-11	1-5	1-5	1-6	1-6	1-6	1-6
Mainland			1-6	1-6	1-5	1-5	1-7
Nantucket			1-5	1-3	1-3	1-4	1-3
Martha's Vineyard					1-4	1-3	1-4

* Includes Naushon Island

TABLE 1

Summary of the results of the experiments conducted during the period from January 1, 1968, to December 31, 1968.

RESULTS

Run	Time	Temp.	Pressure	Flow	Rate	Efficiency
1	10.0	100.0	10.0	1.0	1.0	1.0
2	10.0	100.0	10.0	1.0	1.0	1.0
3	10.0	100.0	10.0	1.0	1.0	1.0
4	10.0	100.0	10.0	1.0	1.0	1.0
5	10.0	100.0	10.0	1.0	1.0	1.0

DISCUSSION

The results of the experiments conducted during the period from January 1, 1968, to December 31, 1968, are summarized in Table 1. The data show that the efficiency of the process is high, and that the rate of flow is constant. The temperature and pressure are also constant, and the time of the experiment is 10.0 minutes.

CONCLUSIONS

The results of the experiments conducted during the period from January 1, 1968, to December 31, 1968, are summarized in Table 1. The data show that the efficiency of the process is high, and that the rate of flow is constant. The temperature and pressure are also constant, and the time of the experiment is 10.0 minutes.

REFERENCES

1. Smith, J. D. (1968). The effect of temperature and pressure on the rate of flow. *Journal of Chemical Engineering*, 10, 1-10.
2. Jones, A. B. (1968). The effect of time on the efficiency of the process. *Journal of Chemical Engineering*, 10, 11-20.
3. Brown, C. D. (1968). The effect of flow rate on the efficiency of the process. *Journal of Chemical Engineering*, 10, 21-30.

antlerless permit holders did occur. The increase in the female harvest was not significant. In the event the harvest remains constant, not varying more than 30 deer, it can be said that the desired balance of the herd has been achieved. From that point in time, the control of the herd size can be obtained by raising or lowering the number of antlerless permits for Nantucket Island.

This is the third year the same number of permits has been issued for Martha's Vineyard. The 173 deer taken by permittees represent 86 percent of the 201 deer harvested on the Vineyard and Naushon Island. An unexplainable decline in the Martha's Vineyard harvest is noted. In 1972, there were 266 deer reported. In 1973, only 201 were reported.

The success ratio for farmer-landowner permit holders was 1:6 as in 1972. The mainland permit holder success ratio dropped from 1:5 in 1972 to 1:7 in 1973. Permit success ratios in 1973 were highest on Nantucket (1:3) and Martha's Vineyard (1:4).

Recommendations: It is recommended that the Division of Fisheries and Game personnel continue to compile and evaluate annual harvest data, and initiate plans, based on the preceding seven years' permit system data for a regional permit distribution program on the mainland.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____

Colton H. Bridges, Superintendent

Prepared by: _____

James J. McDonough, Game Biologist
and
William J. Minior, Assistant

Date _____

JOB PERFORMANCE REPORT

2000-2001
Collection
1 111

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No. II-2 Job Title: Non-Hunting Deer Mortality Investigations

Period Covered: 1 January 1973 to 31 December 1973

Summary: During the period covered by this report, Natural Resource Officers reported 420 deer mortalities, of which 188 were males, 207 were females and 25 with no sex reported. Motor vehicles accounted for 322 deer mortalities followed by 36 dog kills, 23 illegal kills. The remaining 39 mortalities were due to other and unknown causes (21): drowned-9, fences-5, trains-2, and shot doing crop damage-2.

The slightly decreased deer mortality of 420 deer for 1973 compared to the 453 deer mortalities for 1972 may again reflect an open winter. During an open winter, deer do not tend to concentrate in great numbers for any length of time. Therefore, there tends to be less movement thus reducing the motor vehicle accident rate. Dog kills are reduced also because snow conditions favor the deer.

There was a 7 percent decrease in the 1973 deer mortalities compared to a 35 percent decrease in 1972.

The adjusted sex ratio of the 1973 non-hunting deer mortalities was 48 percent male:52 percent females. The 1972 adjusted sex ratio was 46 percent males: 54 percent females.

Berkshire, Barnstable and Franklin (in order of importance) were the top ranking counties where Natural Resource Officers reported the largest number of deer mortalities.

Background: Deer mortalities investigated by Natural Resource Officers were reported in duplicate to the Law Enforcement office in Boston, Massachusetts. A copy of each report was compiled and tabulated at the Field Headquarters of the Division of Fisheries and Game by project personnel. A herd of 15 deer in a 400-acre enclosure at Westover Field, Chicopee, Massachusetts is being observed for natural mortalities.

Objectives: To determine the annual non-hunting decimating factors of the Massachusetts deer herd.

Findings:

Table 1 presents a summary of the sex classes and the causes of Massachusetts deer mortalities for 1 January through 31 December 1973. During the twelve-month period, there were 420 reported deer mortalities. Of these, 188 were males, 207 were females and 25 of unreported sex. With the exception of legally harvested deer, motor vehicle accidents remain the number one cause of reported deaths with 322 deer killed. In order of importance, the remaining 99 deer mortalities were as follows: dogs-36, illegal-23, other unknown causes-21, drowned-9, fences-5, trains-2, and crop damage-2.

A five-year summary of deer mortalities per cause, 1968 through 1973, is presented in Table 2. The 1973 reported deer mortalities were less than the average per cause for the previous five years.

Table 3 presents a comparison of the total non-hunting deer mortalities of Massachusetts deer from 1967 through 1973. From 1967 through 1970 there was a steady but declining percent increase in the reported deer mortalities. From 1970 through 1973 there was a decline in the number of deer reported and a fluctuating percent of decrease.

A comparison of the actual number of deer mortalities by sex per month and the adjusted sex ratios are presented in Table 4. The sex data is adjusted on a monthly basis, with sex of the largest percentage assigned any fractional part of a deer so that there are only whole deer in the adjusted sex ratio. The adjusted sex ratio of the 1973 non-hunting deer mortalities are:

<u>Males</u>		<u>Females</u>
200	:	220
91	:	100
47.6	:	52.4

(1972 sex ratio = 45.9% males:54.1% females)

Table 5 presents the Massachusetts non-hunting deer mortalities ranked by counties, 1967 through 1973. The top deer mortality-producing counties for the past seven years in order of importance are as follows:

1. Berkshire
2. Franklin
3. Barnstable
4. Worcester.

During the period covered by this report, Berkshire remains number one while Barnstable moved into the second slot. Franklin is third and Worcester remains number four. Non-hunting deer mortalities change slightly in the remaining nine counties from year to year.

The first of these is the fact that the
 
the thirteenth is the fact that the

Table 1. A Summary of Massachusetts Deer Mortalities Reported by Natural Resource Officers, 1 January through 31 December 1973.

	January			February			March			April			May			June		
	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No
Motor Vehicles	10	17		7	6	1	12	11	1	3	10	2	8	17	1	20	25	1
Dogs	8	5	3	3	3	1	1	1	1		1		1	1		1		
Illegal Kill	1												1	1				
Drowned				2				1		2				1				
Fences			1	1														
Trains		1																
Crop Damage					2			1										
Other and Unknown	1	4		3	1								2					2
Subtotal	19	28	4	13	14	3	13	14	2	10	11	2	9	22	1	21	25	3
Total (Cum. Total)		51		30	(81)		29	(110)		23	(133)		32	(165)		49	(214)	
	July			August			September			October			November			December		
	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No	M.	F.	No
Motor Vehicles	8	16		3	10		12	9	1	22	12	1	31	20	6	4	9	1
Dogs											2		1	1		2		
Illegal											4		1	1		9		6
Drowned										2			1					
Fences				1						1			1					
Trains														1				
Crop Damage																		
Other and Unknown																		
Subtotal	8	16		3	11		12	9	1	2	1	1	2	23	6	16	15	1
Total (Cum. Total)		24	(233)	14	(252)		22	(274)		48	(322)		66	(388)		32	(420)	
Cause	Male	Female	Total		Cause		Male	Female	Total		Cause		Male	Female	Total		Cause	
Motor Vehicles	145	162	322		Fences		3		3		Fences			1	1	2	5	
Dogs	17	14	36		Trains						Trains			2	2	2	2	
Illegal	10	13	23		Crop Damage						Crop Damage			2	2	2	2	
Drowned	7	2	9		Other and Unknown		6		6		Other and Unknown			11	4	15	21	
Totals			188		Totals						Totals			207	25	232	420	

Table 2. Five-Year Summary of Deer Mortalities of Massachusetts Deer Reported by Natural Resource Officers, 1968 through 1973.

<u>Cause</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>Total</u>	<u>5-yr. Avg.</u>	<u>1973</u>
Motor Vehicles	456	397	400	373	321	1,947	389	321
Dogs	74	166	204	219	41	704	141	36
Illegal Kills	29	39	25	39	44	176	35	23
Crop Damage	7	2	14	4	1	28	6	2
Unknown Causes	31	51	38	41	35	196	39	21
All Other Causes	16	27	17	18	11	89	18	16
Totals	613	682	698	694	453	3,140	628	420

Table 3. A Comparison of Total Non-Hunting Deer Mortalities of Massachusetts Deer from 1967 through 1973.

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
No. of Deer	508	613	682	698	694	453	420
Percent Change		21%	11%	2%	-.6%	-34.7%	-7.3%

Table 4. Comparison of Actual Numbers of Deer Mortalities by Sex* and Adjusted Data for Massachusetts Deer per Month, 1973.

<u>Month</u>	<u>Male</u>	<u>Female</u>	<u>Unknown Sex</u>	<u>Total</u>	<u>Adjusted</u>	
					<u>Male</u>	<u>Female</u>
January	19	28	4	51	20	31
February	13	14	3	30	14	16
March	13	14	2	29	14	15
April	10	11	2	23	11	12
May	9	22	1	32	9	23
June	21	25	3	49	22	27
July	8	16		24	8	16
August	3	11		14	3	11
September	12	9	1	22	13	9
October	27	19	2	48	28	20
November	37	23	6	66	41	25
December	16	15	1	32	17	15
Total	188	207	25	420	200	220

Adjusted Sex Ratio

200 males:220 females
 91 males:100 females
 47.6 males:52.4 females

* These data were reported by Natural Resource Officers.

Table 5. The Massachusetts Non-Hunting Deer Mortalities Ranked by County, 1967 through 1973.

County	1967		1968		1969		1970		1971		1972		1973	
	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank
Barnstable	52	4	55	5	63	4	74	4	93	2	72	2	76	2
Berkshire	103	1	134	1	168	1	184	1	175	1	92	1	109	1
Bristol	16	10	14	11	15	11	8	11	6	11	6	11	6	9
Essex	14	11	42	7	34	8	32	7	25	9	20	9	20	6
Franklin	35	8	81	3	90	3	127	2	90	3	66	3	73	3
Hampden	57	3	58	4	52	5	47	6	51	6	45	5	28	5
Hampshire	48	5	46	6	51	6	56	5	65	5	36	6	23	5
Middlesex	39	6	39	8	37	7	22	8	13	10	8	10	14	8
Norfolk	1	12	7	13	5	12			1	13	6	10	3	10
Plymouth	22	9	25	9	27	9	22	8	36	8	21	8	17	7
Suffolk	0	14	0	14	0	14	0	14	0	14	0	14	0	14
Worcester	35	2	86	2	104	2	99	3	99	3	55	4	37	4
Dukes			11	12	15	11	17	9	17	9	22	7	6	9
Nantucket			15	10	21	10	11	10	11	10	2	12	3	10

Three counties; namely, Norfolk, Bristol and Plymouth, produce more non-hunting deer mortalities than the number of deer reported taken by deer hunters:

<u>County</u>	1973	1973
	<u>Hunting Season</u>	<u>Non-Hunting Mortalities</u>
Bristol	1	6
Norfolk	0	3
Plymouth	3	17

I feel that the reported deer mortalities in this report are valid even though it is impossible to obtain data on every deer mortality statewide. These data indicate trends which must be evaluated with respect to weather conditions, the Natural Resource Officer's ability and interest in filing a deer mortality report, and the NRO's mandatory eight-hour working day.

In an effort to obtain greater cooperation and interest from the Natural Resource Officers, the project leader prepared and delivered a slide lecture to all Law Enforcement personnel in each of the seven Law Enforcement districts in Massachusetts.

Recommendations: It is recommended that deer project personnel continue to determine the annual non-hunting deer decimating factors of the Massachusetts deer herd.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by: _____
James J. McDonough, Game Biologist
and
William J. Minor, Assistant

Date _____

Received of the Treasurer of the
County of ... the sum of ...

for the purchase of ...
... ..

... ..
... ..
... ..

... ..
... ..
... ..

... ..
... ..

... ..
... ..

... ..
... ..

Witness my hand and seal this ... day of ...
1875
... ..

JOB PERFORMANCE REPORT

Section
11.1

State Massachusetts

Cooperator Massachusetts Division of Fisheries and Game

Project No. W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No. II-4 Job Title: Management of the Massachusetts Deer Herd

Period Covered: 1 July 1973 to 30 June 1974

Summary: Sex and age data collected at ten biological deer checking stations from 1967 through 1973 were analyzed.

Four thousand sportsman antlerless permits were issued for the mainland shotgun deer hunting season. Six hundred sportsman permits were issued on Martha's Vineyard; four hundred were issued on Nantucket.

Successful hunters removed 2,037 deer from the herd. Of these, 1,421 were males of which 1,258 were adults and 163 were button bucks. Six hundred sixteen female deer were harvested of which 444 were adult does and 172 were fawns. Antlerless permits issued totaled 5,349 including landowner permits. Thirty-eight percent of the shotgun harvest (2,037 deer) were taken by permit holders.

The calculated minimal herd size for the 1973 shotgun season was 11,431 deer. This is a .83 percent increase above the 1972 herd of 11,336 deer.

During the shotgun season, there were .18 adult males and .04 adult females harvested per square mile of deer range on the mainland. On Martha's Vineyard, adult males and females were harvested each at the rate of 1 deer per square mile.

There was an unexpected increase in the adult male (1.89) and adult female (1.22) deer harvest per square mile of deer range on Nantucket in the 1973 shotgun season.

The total harvest of deer per square mile of deer range in 1973 for the archery, muzzle loader and shotgun seasons was as follows:

<u>Location</u>	<u>Sq. Miles of Deer Range</u>	<u>Males per Sq. Mile of Deer Range</u>	<u>Females per Sq. Mile of Deer Range</u>	<u>Total Deer Per Sq. Mile of Deer Range</u>
Mainland	6213.6	.20	.07	.27
Dukes County	86.5	1.45	1.42	2.87
Nantucket	35.9	2.81	2.01	4.82

The percent frequency ratio of adult males to adult females was 0.29 on the mainland, 1.01 on Martha's Vineyard, and 0.69 on Nantucket. Berkshire, Franklin, Hampden, Barnstable and Worcester Counties, in descending order, produced the greatest deer harvest. The remaining six counties produced 0.01 percent deer or less per square mile of deer range.

Management recommendations for the distribution of antlerless permits on a county and/or regional (group of counties) basis is presented in this report.

Background:

A 24-hour compulsory reporting system for deer hunters was initiated in 1967. Deer checking stations have been operated by Division personnel since 1948. Ten checking stations are operated as biological stations at which sex, age, weight and antler beam diameter data are collected.

An antlerless permit deer hunting system, to control the hunting pressure on the female segment of the deer herd during the shotgun season, was initiated in 1967. The deer project leader recommends the number of permits to be issued after examination of the herd sex and age composition, the reproductive rates, annual removal rates, hunting pressure and the harvest per square mile of deer range.

Objectives:

To estimate the size of the deer herd in Massachusetts and recommend management techniques.

Findings:

Table 1 presents a summary of the sex and age composition of Massachusetts deer examined at the mainland, Nantucket and Martha's Vineyard biological deer check stations from 1967 through 1973.

Table 2 presents a summary of the Massachusetts shotgun deer harvest and the total number of antlerless permits issued from 1967 through 1973. Shotgun hunters removed 2,037 deer from the herd in 1973. Of these, 1,421 were male deer; 1,258 adults and 163 button bucks, being taken. There were 616 female deer harvested, of which 444 were adults and 172 were fawns. Although the data in Table 2 tend to mask the location of where the hunting pressure occurs, it is possible to point out the overall effect of the antlerless permit system in the state. As the

Table 1. A Summary of the Sex and Age Composition of Massachusetts Deer at Mainland Biological Deer Check Stations, 1967 through 1973.

Age	Male							Female						
	1967	1968	1969	1970	1971	1972	1973	1967	1968	1969	1970	1971	1972	1973
6 mos.	28	61	67	121	100	77	69	31	44	76	90	101	90	62
1-1/2	167	198	229	263	211	260	237	32	33	55	54	64	57	48
2-1/2	122	126	133	147	103	144	173	18	28	55	69	69	56	42
3-1/2	62	87	98	97	77	96	90	13	16	36	46	51	51	35
4-1/2	25	35	55	59	43	46	47	7	11	24	29	33	22	25
5-1/2	10	17	21	21	19	14	13	3	3	11	14	20	14	6
6-1/2	5	4	14	7	3	11	7	1	4	0	8	14	10	5
7-1/2	1	0	2	7	6	2	6	0	3	2	0	11	2	6
8-9-1/2	1	0	2	1	1	0	3	3	3	0	0	9	0	2
10-1/2	0	0	1	0	0	00	0	1	1	0	0	0	1	0
	421	528	652	723	563	563	650	109	146	259	310	372	303	231

Age	Martha's Vineyard					Nantucket				
	Male		Female		1973	Male		Female		1973
	1971	1972	1971	1972		1971	1972	1971	1972	
6 mos.	39	25	30	30	24	27	17	14	21	21
1-1/2	41	41	25	24	21	38	19	22	16	17
2-1/2	14	15	17	11	17	13	12	12	12	8
3-1/2	8	15	12	15	13	7	12	13	5	7
4-1/2	4	6	7	9	8	4	0	2	5	9
5-1/2	3	1	3	6	2	0	3	3	0	2
6-1/2	0	3	0	1	1	0	0	0	1	0
7-1/2	1	2	0	2	1	1	0	0	1	1
8-9-1/2	1	0	0	1	1	0	0	0	0	0
10-1/2	0	0	0	0	0	0	0	0	0	0
	111	108	94	99	88	90	63	66	61	65

number of antlerless permits issued per year increased the harvest of antlerless deer increased. In 1967 and 1968, there were 2,236 and 2,331 antlerless permits issued. During these years, the antlerless deer harvested by permit amounted to 25 percent (238) and 27 percent (371) of the total deer harvest. The adult male harvest increased 136 deer from 1967 (836) to 1968 (1,022).

In 1969, the number of permits issued was increased to 4,695 permits. The percent of antlerless deer harvested increased to 35 percent of the total harvest of the shotgun deer week. The adult male harvest increased 277 deer for a total of 1,299 adult males reported in 1969.

In 1970, the number of antlerless permits issued was increased to 6,747 permits. Forty-two percent of the deer harvested were button bucks and females. The adult male harvest increased only 79 animals with a total of 1,378 deer.

Seven thousand two hundred seventy permits were issued in 1971. The result was that 51 percent (1,142) of the shotgun harvest were antlerless deer. The adult male harvest dropped 193 animals for a total of 1,106 adult bucks reported.

In 1972, the number of antlerless permits issued was lowered to 5,326 permits. The combined button buck and female harvest dropped to 43 percent (959). The adult male harvest increased 150 animals for a total of 1,256 adult bucks.

In 1973, 5,349 antlerless permits were issued. The antlerless deer harvest dropped to 38 percent (799) of the total shotgun harvest. There was an insignificant increase of 2 bucks in the adult male harvest of 1,258 deer in 1973.

In summary, the data in Table 2 can be interpreted to demonstrate that on an overall basis the number of antlerless permits issued per year should be increased by relatively smaller increments than in past years, and only when the harvest of adult males warrants an increase of antlerless permits.

A summary of the percent change in the adult harvest and the calculated minimal deer population in Massachusetts, 1967 through 1973, is presented in Table 3. It is interesting to note that the percent change for the calculated adult female population and the calculated minimal population has been decreasing since the 1970 and 1971 deer seasons. It was during these seasons that the antlerless permits had been increased to 6,747 and 7,270 respectively.

The first of these is the fact that the
the second is the fact that the
the third is the fact that the

the fourth is the fact that the
the fifth is the fact that the

the sixth is the fact that the
the seventh is the fact that the

the eighth is the fact that the
the ninth is the fact that the

the tenth is the fact that the
the eleventh is the fact that the

the twelfth is the fact that the
the thirteenth is the fact that the

the fourteenth is the fact that the
the fifteenth is the fact that the

the sixteenth is the fact that the
the seventeenth is the fact that the
the eighteenth is the fact that the
the nineteenth is the fact that the
the twentieth is the fact that the

Table 2. A Summary of the Massachusetts Shotgun Deer Harvest, 1967 through 1973.

Year	All Males		Button Bucks	All Females		Adult Females	Fawn Females	Total Deer	Total Permits Issued	Button Bucks and Females	% of Total Deer
	Male	Female		Male	Female						
1967	937	886	53	235	168	67	1,172	2,236	288	25	
1968	1,083	1,022	61	310	217	93	1,393	2,331	371	27	
1969	1,424	1,299	125	586	415	171	2,010	4,695	711	35	
1970	1,598	1,378	220	769	546	223	2,367	6,747	939	42	
1971	1,359	1,106	253	889	643	246	2,248	7,270	1,142	51	
1972	1,455	1,256	199	760	530	230	2,215	5,326	959	43	
1973	1,421	1,258	163	616	444	172	2,037	5,349	779	38	

Table 3. A Summary of Percentage Change in Adult Harvest and Calculated Minimal Populations of Deer in Massachusetts, 1967 through 1973.

	<u>1967</u>	<u>Percent Change 1967-1968</u>	<u>1968</u>	<u>Percent Change 1968-1969</u>	<u>1969</u>	<u>Percent Change 1969-1970</u>	<u>1970</u>	<u>Percent Change 1970-1971</u>	<u>1971</u>
Adult male harvest	886	+ 15.35	1,022	+ 27.10	1,299	+ 6.03	1,378	- 19.73	1,106
Calculated adult male population	2,056	+ 17.05	2,410	+ 31.95	3,180	- 0.82	3,154	- 27.68	2,281
Calculated adult female population	2,018	+ 31.92	2,964	+ 35.19	4,007	+ 32.24	5,299	- 22.95	4,083
Calculated minimal population	6,139	+ 42.37	8,740	+ 35.01	11,805	+ 27.13	15,014	- 22.93	11,571
Adult male harvest					<u>1971</u>	<u>Percent Change 1971-1972</u>	<u>1972</u>	<u>Percent Change 1972-1973</u>	<u>1973</u>
Calculated adult male population					1,106	+ 13.56	1,256	+ 00.02	1,258
Calculated adult female population					2,231	+ 20.78	2,755	+ 11.80	3,080
Calculated minimal population					4,033	- 3.67	3,933	- 02.11	3,850
					11,571	- 2.03	11,336	+ 00.83	11,431

Tables 4 and 5 present a summary of the adult male and female deer harvest per square mile of deer range in Massachusetts from 1967 through 1973. Whereas it was felt that there was an overharvest in the female segment of the deer population during 1970 and 1971, the number of sportsman antlerless permits issued on the mainland was reduced to 4,000 permits in 1972. The same number (4,000) of permits was issued for the 1973 shotgun deer season.

The adult male harvest per square mile of deer range in Berkshire County, except for 1971, has shown a continuous rise from .27 antlered bucks in 1967 to .51 animals in 1973. The adult female harvest in 1967 was .04 deer and rose to a peak of .23 adult does in 1971. The harvest for 1972 and 1973 has remained at .13 adult does per square mile of deer range.

In Barnstable County, the adult male harvest increased slightly from .26 bucks in 1972 to .27 bucks in 1973. The adult female segment of the 1973 harvest dropped from .09 in 1972 to .05 for 1973.

There was a .10 drop in both adult age classes in Franklin County during the 1973 shotgun season. It is quite possible that the decrease was due to an overharvest of females from 1969 through 1972.

The harvest of adult deer in Hampden and Hampshire Counties appears as expected with the buck harvest increasing and the female segment remaining the same or slightly less than the 1972 harvest.

In Worcester County, the adult male harvest remained at .09 bucks per square mile of deer range while the adult female segment increased .01 deer to .04 females per square mile of range in 1973.

The remaining counties, namely Bristol, Essex, Middlesex, Norfolk and Plymouth, contributed an insignificant number of deer to the mainland harvest. These counties are called collectively the Eastern Group.

The adult male harvest in Dukes County decreased to 1 deer per square mile of deer range; the adult female harvest increased to 1 adult doe per square mile of deer range in 1973. Sixty-three percent of the deer harvested on Martha's Vineyard were button bucks and females.

The Nantucket Island adult male harvest increased from 1.36 in 1972 to 1.89 bucks in 1973. The increase in the male harvest was unexpected. There was a slight increase in the adult female harvest of .11 does during the 1973 shotgun season.

Table 4. Summary of the Adult Male Deer Harvest per Square Mile of Deer Range per County in Massachusetts, 1967 through 1973.

<u>County</u>	<u>Sq. Mi. Deer Range</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Barnstable	290.5	.14	.15	.20	.28	.28	.26	.27
Berkshire	839.0	.27	.37	.44	.44	.42	.45	.51
Bristol	422.6	.01	.01	.01	.01	.002	.007	.002
Essex	344.4	.03	.03	.04	.06	.03	.05	.05
Franklin	649.1	.27	.35	.39	.39	.36	.43	.33
Hampden	524.3	.13	.19	.24	.22	.20	.25	.28
Hampshire	431.5	.16	.25	.27	.26	.15	.23	.23
Middlesex	583.1	.02	.01	.02	.02	.02	.01	.01
Norfolk	277.3	---	---	.01	---	---	---	---
Plymouth	544.2	.03	.02	.03	.02	.01	.02	.01
Worcester	<u>1,307.6</u>	<u>.11</u>	<u>.06</u>	<u>.14</u>	<u>.15</u>	<u>.07</u>	<u>.09</u>	<u>.09</u>
	6,213.6	.12	.14	.13	.19	.15	.18	.18
Dukes	86.5	.71	.65	.72	.91	1.01	1.18	1.00
Nantucket	35.9	1.34	1.56	1.89	2.42	1.78	1.36	1.89

Table 5. Summary of the Adult Female Deer Harvest per Square Mile of Deer Range per County in Massachusetts, 1967 through 1973.

<u>County</u>	<u>Deer Range</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>
Barnstable	290.5	.03	.02	.06	.10	.09	.09	.05
Berkshire	839.0	.04	.06	.12	.16	.23	.13	.13
Bristol	422.6	.005	.002	.002	---	---	.002	.002
Essex	344.4	.006	.003	.01	.03	.003	.003	.00
Franklin	649.1	.07	.11	.14	.15	.27	.22	.12
Hampden	524.1	.02	.03	.06	.06	.08	.07	.07
Hampshire	431.5	.04	.06	.11	.11	.07	.08	.05
Middlesex	583.1	.002	.01	.01	.01	.01	.01	.002
Norfolk	277.3	.004	---	---	.004	---	---	---
Plymouth	544.2	.002	.01	.01	.01	.002	---	.004
Worcester	<u>1,307.6</u>	<u>.02</u>	<u>.02</u>	<u>.05</u>	<u>.07</u>	<u>.05</u>	<u>.03</u>	<u>.04</u>
	6,213.6	.02	.03	.06	.07	.08	.07	.05
Dukes	86.5	.14	.10	.18	.36	.67	.89	1.00
Nantucket	35.9	.17	.14	.97	1.67	1.53	1.11	1.22

Table 6 presents a summary of the total harvest of deer in Massachusetts per sex per county and the harvest of deer per square mile of deer range for the 1973 deer seasons (archery, muzzle loader and shotgun seasons). The six top ranking deer producing counties on the mainland in order of importance are: Berkshire, Franklin, Hampden, Barnstable, Hampshire and Worcester. The remaining counties produce .01 deer or less per square mile of deer range during the hunting seasons. Nantucket Island shows the largest reported harvest per square mile of deer range with 4.82 deer. Dukes County, with 2.87 deer per square mile of deer range, yields the second largest harvest on a unit area basis.

The percent frequency ratio of adult females to adult males from 1967 through 1973 is presented in Table 7. The data in Table 7 can be used, within certain limits, as the pulse of the antlerless permit system. In Dukes County, the harvest of adult females was slightly over 1 female for each adult male in 1973. The percent frequency ratio of adult females in the Nantucket harvest decreased from 0.82 to 0.69 percent in 1973. Please note the number of animals involved when interpreting the percent of adult females per adult male data for the mainland. For example, the .40 percent in Plymouth County involves 7 animals while in Worcester County the .43 percent was derived from a total of 170 animals (119 males:51 females).

The results of a reduction in the number of antlerless permits is evident in the data of Table 7. In 1970 and 1971, 6,000 sportsman permits were issued for the mainland. The percent of adult females to adult males harvested increased in all counties. The number of antlerless permits issued was reduced to 4,000 permits for the 1972 and 1973 deer seasons. The results seen in Table 7 show that the percent harvest of adult females to adult males declined for these years in all counties with the exception of Worcester County. The reduction in the number of permits issued removed pressure from the female segment of the deer herd. Based on the data in Table 7, the issuance of permits on a county basis is recommended.

Figure No. 1 graphs a summary of the total harvest of deer in Massachusetts by sex per county, and the harvest of deer per square mile of deer range from 1967 through 1973. Only the six major deer harvest producing counties are represented in the graph.

The sensitivity of the antlerless permit system can be observed in Figure No. 1. In 1967 and 1968, 2,000 sportsman antlerless permits were issued. The harvest of male deer increased during these years while the female harvest increased at a slower rate. The number of permits issued was increased to 4,000 in 1969. Both the male and female

Table 6. A Summary of the Total Harvest of Deer in Massachusetts (including Shotgun, Archery, and Muzzle Load Harvests) per County per Sex and the Harvest of Deer per Square Mile of Deer Range in Massachusetts for 1973

County	Males	Females	Total	Sq. Mi. of Deer Range	Harvest		
					Total Deer per Sq. Mi.	Males per Sq. Mi.	Females per Sq. Mi.
Barnstable	83	24	107	290.5	.36	.29	.08
Berkshire	472	150	622	839.0	.74	.56	.18
Bristol	0	1	1	422.6	.002	.00	.002
Essex	20	0	20	344.4	.06	.06	.00
Franklin	247	114	361	649.1	.56	.38	.13
Hampden	161	49	210	524.3	.40	.31	.09
Hampshire	114	33	147	431.5	.34	.26	.08
Middlesex	4	1	5	583.1	.009	.007	.001
Norfolk	0	0	0	277.3	.00	.00	.00
Plymouth	5	3	8	544.2	.01	.009	.005
Worcester	<u>145</u>	<u>74</u>	<u>219</u>	<u>1307.6</u>	<u>.17</u>	<u>.11</u>	<u>.06</u>
Total	1,251	449	1,700	6213.6	.27	.20	.07
Dukes	125	123	248	86.5	2.87	1.45	1.42
Nantucket	<u>101</u>	<u>72</u>	<u>173</u>	<u>35.9</u>	<u>4.82</u>	<u>2.81</u>	<u>2.01</u>
Total	1,477	644	2,121	6336.0	.33	.23	.10

Table 7. Percent Frequency Ratio of Adult Females to Adult Males, 1967 through 1973.

County	1967			1968			1969			1970		
	M.	F.	% Freq.	M.	F.	% Freq.	M.	F.	% Freq.	M.	F.	% Freq.
Barnstable	41	10	.24	44	6	.14	58	16	.28	80	29	.36
Berkshire	223	31	.14	314	53	.17	371	100	.27	366	138	.33
Bristol	6	2	.33	5	1	.20	3	1	.33	4		
Essex	11	2	.18	10	1	.10	13	4	.31	20	9	.45
Franklin	176	44	.25	230	68	.30	256	93	.36	251	98	.39
Hampden	69	9	.08	98	13	.13	123	32	.26	117	31	.26
Hampshire	69	16	.23	107	27	.25	115	47	.41	114	48	.42
Middlesex	10	1	.10	7	3	.43	13	4	.31	9	6	.67
Norfolk		1					3				1	
Plymouth	16	1	.06	8	3	.38	14	6	.43	8	4	.50
Suffolk												
Worcester	140	32	.23	75	23	.37	176	59	.34	192	88	.46
Total	766	149	.19	898	203	.23	1,145	362	.32	1,161	452	.39
No. of Sportsman Permits		2,000			2,000			4,000			6,000	
Dukes	61	12	.20	56	9	.16	62	16	.26	79	31	.39
No. of Sportsman Permits												
Nantucket	48	6	.13	56	5	.09	68	35	.51	87	60	.69
No. of Sportsman Permits								400			400	
Total Number of Permits*		2,243			2,331						6,747	

* Includes Farmer-Landowner Permits

Table 7. (Continued)

County	1971			1972			1973		
	M.	F.	% Freq.	M.	F.	% Freq.	M.	F.	% Freq.
Barnstable	82	25	.30	75	27	.36	77	15	.19
Berkshire	352	187	.53	373	112	.30	424	109	.26
Bristol	1			3	1	.33			
Essex	12	1	.08	16	1	.06	16	4	.25
Franklin	232	176	.76	277	141	.51	213	79	.37
Hampden	103	42	.41	131	38	.29	145	36	.25
Hampshire	63	31	.49	98	34	.35	101	22	.22
Middlesex	14	7	.50	3	4	1.33	4	1	.25
Norfolk									
Plymouth	7	1	.14	12	0	.00	5	2	.40
Worcester	89	60	.67	117	44	.38	119	51	.43
Suffolk									
Total	955	530	.55	1,105	402	.36	1,104	316	.29
No. of Sportsman Permits		6,000			4,000			4,000	
Dukes	87	58	.67	102	77	.75	86	87	1.01
No. of Sportsman Permits		600			600			600	
Nantucket	64	55	.86	49	40	.82	68	44	.69
No. of Sportsman Permits		400			400			400	
Total Number of Permits*		7,270			5,326			5,349	

* Includes Farmer-Landowner Permits

A Summary of the Total Massachusetts Deer Harvest by Sex per Square Mile of Deer Range by County. (1967 through 1973)

harvest increased with the female rate increasing at a slightly higher rate. Six thousand permits were issued for 1970 and 1971. The harvest of male deer in most cases peaked in 1970 and started to decline in 1971. The female harvest during these years continued to climb. The 1971 harvest reflects the increase of permits in 1969 in that it takes two years to put an antlered deer in the herd. The harvest of males in 1971 started to decline and the permit holders were taking more antlerless deer. The 1972 and 1973 harvests reflect the permit increase of 1970 and 1971. The 1974 harvest, therefore, should be higher than the 1973 harvest on the mainland because of the 1972 reduction in antlerless permits issued.

A summary of the total Massachusetts deer harvest by sex per square mile of deer range on the mainland, Dukes and Nantucket Counties from 1967 through 1973 is presented in Figure No. 2. The mainland data represents the bulk of the deer range in the state, but with the six poor deer producing counties included, the harvest of deer per square mile of range is quite low.

The total harvest of deer in Dukes County decreased in 1973. Whereas this is the third year of issuing 600 antlerless permits, further decline in the yield of this herd may necessitate a reduction in permits for the county.

There was an unexpected increase in both the male and female harvest of deer on Nantucket Island during the 1973 deer season. It is quite possible that the increased harvest was due to a build up of adult female deer surviving the 1967, 1968 and 1969 deer seasons. It had been anticipated that the male harvest would equal the female harvest in 1973 followed by a reduction in the number of permits to maintain an equal harvest of bucks and does.

Recommendations: Antlerless permit allocation should be modified to a county and regional basis of distribution. The following numbers of sportsman permits issued per county and/or region are suggested:

<u>County</u>	<u>Number of Sportsman Permits</u>
Barnstable	200
Berkshire	1,300
Franklin	700
Hampden	400
Hampshire	300
Worcester	700
Region I*	200
Region II**	200
Total	4,000

* Essex, Middlesex and Norfolk Counties

** Bristol and Plymouth Counties

 

—Total

Martha's Vineyard	600
Nantucket Island	400
Naushon	50

The sale of a Massachusetts big game license that incorporates an antlerless permit application should be investigated. When the antlerless application is submitted, the hunter will be required to indicate in which county or region he would like the application be drawn.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____

Colton H. Bridges, Deputy Director

Prepared by: _____

James J. McDonough
Game Biologist

William J. Minior
Assistant

Date _____

The first part of the history of the
the second part of the history of the
the third part of the history of the

The fourth part of the history of the
the fifth part of the history of the

The sixth part of the history of the
the seventh part of the history of the

The eighth part of the history of the
the ninth part of the history of the
the tenth part of the history of the

THE HISTORY OF THE

PERFORMANCE REPORT

State: Massachusetts

Cooperators: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Name: Game Population Trend and Harvest Survey

Job No. III-1 Job Title: Mourning Dove Census

Period Covered: 23 May to 13 July 1973

Summary: Calling doves were counted on three randomized routes in cooperation with the U. S. Fish and Wildlife Service's annual mourning dove breeding population census. The total number of calling doves for all three routes remained unchanged from 1972.

Objectives: To obtain an index of the spring breeding population of mourning doves.

Procedures: In accordance with instructions from the U. S. Fish and Wildlife Service, doves were censused on the three randomized routes used in 1967. Doves were censused by roadside coo-counts on these routes between 23 May and 1 June. Division personnel conducted two routes and a Fish and Wildlife Service cooperator conducted the third route.

Findings: Results of the 1973 call count of mourning doves are compared with previous years' data in Table 1.

The total number of calling doves heard on the three routes surveyed remained unchanged from 1972 totals. Taken individually, Route 8 decreased by 17 percent, Route 8A was unchanged, and Route 10 increased 100 percent from two to four doves.

The weighted mean number of doves heard per comparable Massachusetts route was 6.1 in both 1972 and 1973 (Ruos, 1973). Ruos further reports that the breeding population index for the Eastern Management Unit decreased 6.8 percent from 17.4 doves per route in 1972 to 16.2 doves per route in 1973. Long-term population trends also show a decline; regression analyses of the 1963-73 data indicate a highly significant downward population trend in the Eastern Management Unit (Ruos 1973:6).

Government Documents
Collection

NOV 14 1973

University of Massachusetts

Table 1. Doves heard on comparable routes: 1966-1973.

Route	<u>Doves Heard</u>							Percent 1973 Counts $\pm$ From							
	1966	1967	1968	1969	1970	1971	1972	1973	1966	1967	1968	1969	1970	1971	1972
3	22	26	11	2	11	10	12	10	- 54	- 62	- 9	-400	- 9	0	- 17
8A	*	1	1	1	2	1	2	2	*	-100	-100	-100	0	-100	0
10	3	7	0	1	3	1	2	4	+ 33	- 43	-400	-300	+ 33	+300	+100
Totals	25	34	12	4	16	12	16	16	- 36	- 53	+ 33	-300	0	+ 33	0
Mean	8.33	11.33	4.00	1.33	5.33	4.00	5.33	5.33							

*Route not conducted.

Recommendations: I recommend that the spring mourning dove census be continued in cooperation with the Bureau of Sport Fisheries and Wildlife.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Assistant Game Biologist

Date _____

Literature Cited

Ruos, James L. 1973. 1973 Mourning Dove Breeding Population Status.
USDI Bureau of Sport Fisheries and Wildlife, Laurel, Md. Administrative
Report. Xerox 1-10 pp.

JOB PERFORMANCE REPORT

State Massachusetts

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No. IV-1 Job Title: Experimental Turkey Stocking

Period Covered: 1 June 1973 to 31 May 1974

Summary: Twenty wild-trapped turkeys (five adult hens and fifteen poults) were captured in Allegany State Park, New York, and released in Beartown State Forest in western Massachusetts. To date, 37 turkeys have been released in this area.

Winter observations indicate the turkeys have adapted well to this area, with a good potential for broods this year.

Observations on semi-wild flocks were limited, but indicate slight improvement over previous years.

Objectives: To re-establish the wild turkey in the Commonwealth in sufficient numbers to allow for recreational hunting.

Procedures: Turkeys were censused by roadside counts, track counts and cooperator reports. Snowmobiles were used during the winter to provide access to the areas and to transport grain for baiting.

Turkeys were captured using Thornsberry style cannon nets. Captured turkeys were aged, sexed, weighed, wing-tagged, and color-marked with patagial streamers.

Findings: Turkey Trapping

Through the courtesy of the New York State Department of Environmental Conservation, Division of Fish and Wildlife, wild-trapped turkeys were again made available for transplanting to Massachusetts. Since adverse weather conditions had restricted trapping success during the winters of 1972 and 1973, New York biologists suggested that we try a fall release consisting of adult hens and their broods. Accordingly, bait lines were established in Allegany State Park, Cattaraugus County, and two captures were made by local technicians on 17 and 18 September. Twenty turkeys were captured (Table 1). The project leader and an assistant traveled to the park to pick up the turkeys, returning to Massachusetts the following day to release the birds. All turkeys were released off Alcott Street near Beartown State Forest, Great Barrington, Berkshire County, at the same location where turkeys had been set free the previous two springs.

Table 1. Sex, Age, Weight, Wing Tag Number, and Capture-Release Data for Turkeys Captured Fall 1973.

Wing Tag Number *	Age	Sex	Weight Kg (lbs-oz)	Capture Date	Capture Location**	Release Date***
T5226	Ad.	F.	3.88 (8-9)	17 September	A	19 September
T5876	Ad.	F.	4.24 (9-6)	18 September	B	19 September
T5877	Ad.	F.	3.94 (8-11)	18 September	B	19 September
T5878	Imm.	Unk.	-	18 September	B	19 September
T5879	Imm.	Unk.	-	18 September	B	19 September
T5880	Imm.	Unk.	-	18 September	B	19 September
T5881	Imm.	Unk.	-	18 September	B	19 September
T5882	Imm.	Unk.	-	18 September	B	19 September
T5883	Imm.	F	-	18 September	B	19 September
T5884	Imm.	Unk.	-	18 September	B	19 September
T5885	Ad.	F.	3.37 (7-7)	17 September	A	19 September
T5886	Ad.	F.	3.71 (8-3)	17 September	A	19 September
T5887	Imm.	M.	3.37 (7-7)	17 September	A	19 September
T5888	Imm.	F.	-	17 September	A	19 September
T5889	Imm.	F.	-	17 September	A	19 September
T5890	Imm.	F.	-	17 September	A	19 September
T5891	Imm.	F.	-	17 September	A	19 September
T5892	Imm.	F.	-	17 September	A	19 September
T5893	Imm.	M.	-	17 September	A	19 September
T5894	Imm.	M.	-	17 September	A	19 September

* All wing streamers white except T5226-orange.

** A, Wolf Run, Allegany State Park; B, Camp No. 10 area, Allegany State Park.

*** All turkeys released off Alcott Street, near Beartown State Forest, Great Barrington.

UNIVERSITY OF CHICAGO

LIBRARY

1911

UNIVERSITY OF CHICAGO

LIBRARY

1911

1911

UNIVERSITY OF CHICAGO

LIBRARY

UNIVERSITY OF CHICAGO

LIBRARY

Statewide Turkey Populations

A total of 37 wild-trapped turkeys (Table 2) have been released in Beartown State Forest between March 1972 and September 1973. Sixteen hens, fifteen toms, and six unsexed poults have been released. This should be quite sufficient to establish a sustaining population in this area.

Table 2. Sex, Age and Date Released of Turkeys Transferred to Beartown State Forest.

<u>Date</u>	<u>Ad. M.</u>	<u>Ad. F.</u>	<u>Imm. M.</u>	<u>Imm. F.</u>	<u>Imm. U.</u>	<u>Total</u>
10 March 1972		3*				3
11 March 1972			2	1		3
17 March 1972	1					1
23 February 1973	5					5
27 February 1973			4			4
23 April 1973		1				1
19 September 1973	—	<u>5</u>	<u>3</u>	<u>6</u>	<u>6</u>	<u>20</u>
	6	9	9	7	6	37

* One found dead 27 March 1972.

Since wild-trapped turkeys offer the best chance of establishing a viable and expanding turkey population in the Commonwealth, turkey census activities during this segment were concentrated on the Beartown flock. Observations in the remainder of the state were further hampered by a reduction in gasoline availability due to the energy crisis.

There have been few sightings of the Beartown birds during the past year. Western Wildlife District personnel reported tracking one adult tom near the Becket Wildlife Management Area (10.5 km/6.5 miles northeast of the release site) during the spring of 1973, and workers in the state forest were told by visitors that "a number" of turkeys were seen in the center of the forest near an abandoned Civilian Conservation Corps camp (2.8 km/1.75 miles east of the release site). The Project Leader and an assistant searched much of the forest and adjacent areas on snowmobiles and foot during the past winter. Thirteen of the twenty birds released in fall of 1973 were located, plus a few (two or three) toms. All birds were on the west slope above Ice Glen Road, 2.0 km (1.25 miles) north of the Alcott

Street release point. Snow cover was slight, seeps were open, and the turkeys were readily locating acorns, hickory nuts, and other foods. Hopefully, broods will be produced this year.

Turkeys in the Quabbin Reservation showed slight improvement over previous years. Broods continue to be produced in the southern portion of Prescott Peninsula, near Prescott Brook, and these birds flush readily when approached. On the other hand, birds of the same background, a few miles north in New Salem are very docile due to persistent feeding by landowners. Natural Resource officers reported a flock of turkeys in Montague, also heavily dependent on artificial feeding. Small flocks remain in Barre, Douglas and Hatfield, but little reproduction and virtually no dispersal has occurred during the past few years.

A small flock of game-farm ancestry turkeys remain near the Adams property in the Town of Mt. Washington, and a limited number of semi-wild birds are scattered about in the Washington-Becket area. These will probably be overwhelmed by the wild-strain Beartown birds should they disperse in this direction.

The Southeast District received an unverified report of turkeys near Camp Squanto in Plymouth during the spring of 1973. Should this be valid, it would be the first indication of these birds since the summer of 1971.

Recommendations: Continue evaluation of the Beartown release. Use these birds as a source of stock for future transplant operations once populations have grown enough to permit this.

Acknowledgments: Appreciation is extended to Messrs. Lee DeGraff, Fred Evans, and William Shirey of the New York State Department of Environmental Conservation and to Dr. W. Rasey of Westons Mills, New York for their assistance in securing and coordinating pick-up of wild-trapped turkeys, and to personnel of Beartown State Forest, the Quabbin Region of the Metropolitan District Commission and the Division of Law Enforcement for their help in reporting turkey sightings.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: Colton H. Bridges, Superintendent

Prepared by: James E. Cardoza, Game Biologist

Date _____

JOB PERFORMANCE REPORT

State MASSACHUSETTS

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No.: VI-1 Job Title: Black Bear Population Dynamics

Period Covered: 1 June 1973 to 31 May 1974

Summary: Applications for bear hunting permits were received from 309 sportsmen. No bears were reported taken during the open season. One bear cub was killed by an automobile. New reports of 18 observations involving 28 bear were received from five counties. Two instances of problem bears were investigated.

Objectives: To define the range of the black bear in Massachusetts and to determine its population characteristics and rate of harvest by hunting.

Procedures: Current bear hunting regulations include mandatory reporting and tagging of bear. Bear checking stations were maintained daily during bear week at three locations--Birch Hill Wildlife Management Area, Templeton; Montague Fish Hatchery, Montague; Western Wildlife District Headquarters, Pittsfield. Should a bear be presented for examination, station personnel were directed to affix a metal game seal to the bear, and to record the following information: town of kill, date killed, sex and weight of bear, and method of kill. Successful hunters would subsequently be contacted by the project leader and the bear's carcass examined and a tooth removed for sectioning.

The Information and Education Section issued periodic news releases asking for reports of black bear and the project leader contributed an article on bear to MASSACHUSETTS WILDLIFE. District personnel, Natural Resource officers and University of Massachusetts cooperators assisted in reporting bear sightings.

Findings: Black bear hunting permit applications were received from 309 individuals during the 1973 season (Table I). No hunter succeeded in bagging a bear.

Table 1. Number of Bear Permit Applications and Number of Bear Taken - 1970-1973

<u>Year</u>	<u>No. Permits</u>	<u>No. Bear Taken</u>	<u>Other Mortalities</u>
1970	214	-	1 illegal kill; 1 road kill
1971	200	-	
1972	423	1	1 illegal kill
1973	309	-	1 road kill; 1 captured bear

New reports of bears received during this segment included 15 sightings, one road kill, and one report of tracks, totaling 28 bear in 16 towns. Reports by county for the period 1952 to 1973 are presented in Table 2. Records of 477 bear noted on 384 occasions in 75 towns have been collected for the period 1952 to May 1973.

Table 2. Reports of Black Bear by County, 1952-1973

<u>County</u>	<u>1952-1972</u>	<u>1973</u>	<u>Total</u>	<u>Percent</u>
Berkshire	124	5	129	33.6
Franklin	148	6	154	40.1
Hampden	18	1	19	4.9
Hampshire	59	5	64	16.7
Middlesex	1	-	1	0.3
Worcester	<u>16</u>	<u>1</u>	<u>17</u>	<u>4.4</u>
	366	18	384	100.0

Four sow-cub groups were observed, two with two cubs each and two with three each.

Two problem situations were investigated. The first, in the summer of 1973, involved three cubs treed in a roadside park in Huntington, Hampshire County. One cub was captured by an onlooker who desired the cub for a pet. Investigation by the Division of Law Enforcement resulted in the retrieval of the cub and the prosecution of its captor. However, the cub was too young to survive on its own and had to be given to a zoo.

The second situation also involved a sow and cubs. On 20 September 1973, barking dogs treed a sow and three cubs in the Town of Clarksburg, Berkshire County. Broadcasts on a local radio station drew hundreds of curious onlookers who threatened the safety of the bears and themselves by their incautious behavior. Police and Natural Resource officers

dispersed the crowd and the bears later departed only to appear again later that night in the neighboring town of North Adams. The bears were again harassed by townspeople and the local fire department attempted to disperse the bears with fire hoses. State police and Resource officers again responded and succeeded in quieting the uproar and the bears were again able to depart. The cub hit by an automobile on Route 2 a month later may have been one of this group.

Recommendations:

1. Continue evaluation of bear harvest through checking stations and periodic questionnaires.
2. Continue determination of bear range and populations through recording of sightings.
3. Investigate nuisance complaints as necessary. Promote public education programs to lessen man-bear conflicts.
4. Incorporate data gathered to date in the planned bulletin on the history and status of the black bear.

Acknowledgments:

Efforts of the Division of Law Enforcement are appreciated, especially Officers Lawler, Kenary, Kulish and Ricardi in investigating complaints and sightings.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Game Biologist

Date _____

JOB PERFORMANCE REPORT

State MASSACHUSETTS

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-16 Project Title: Game Population Trend and Harvest Survey

Job No.: VI-2 Job Title: Historical Records of the Black Bear in Massachusetts

Period Covered: 1 June 1973 to 31 May 1974

Summary: A comprehensive literature search was conducted to locate historical references concerning black bear. A detailed report on the history and status of the bear is now in preparation.

Objectives: To determine trends in black bear populations and distribution in Massachusetts from pre-Colonial times to the present.

Procedures: The facilities of seven state and municipal libraries were utilized in conducting the literature search for this study. Indexing and abstracting journals were used as an initial source of reference concerning bear. Local and regional faunal lists, travel accounts, reports of sporting expeditions, and Division annual reports and correspondence files were examined for citations pertaining to bear. Town and county histories were a valuable source of data, and all histories available at the libraries visited were screened in addition to a few more histories obtained on Interlibrary Loan.

Abstracts were made of all pertinent references, a file-card bibliography was prepared, and maps were drawn diagramming fluctuations in bear distribution since Colonial times.

Findings: In cooperation with the University of Massachusetts, this study has been expanded to include adjacent New England states. A consolidated final report on the history and status of the black bear in this region is being prepared.

To date, 252 citations for the period 1634 to 1948 have been located for New England and component states. Of these, 125 pertain to Massachusetts.

Data in the published report will include findings from W-35-R, Job VI-1, Black Bear Population Dynamics, and will be broken down into four main sections: topography of the region and Indian influences on bear; trends in bear distribution; bears and man; status, management and recommendations.

A popular account of bears in the Commonwealth was published in MASSACHUSETTS WILDLIFE (Cardoza, 1973).

Recommendations: Consolidated findings of this job should be published as a Division research bulletin.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Game Biologist

Date _____

Literature Cited:

Cardoza, James E. 1973. Bay State Bruins--Past and Present. Mass. Wildl. 24(6):10-14.

 

U OF MASSACHUSETTS LIBRARY

State: MASSACHUSETTSProject No.: W-35-R-17Project Title: Game Population Trend and Harvest Survey

Project Type: Research and Surveys

Period Covered: 1 June 1974 to 31 May 1975

Work Plan I Statewide Game Harvest

Objectives: To determine the statewide harvest of selected small game and furbearer species and to present recommendations, based upon management practices and regulations, to increase the utilization of certain species:

Job I-1 Statewide Small Game Harvest

Job Objectives: To determine the statewide harvest of selected small game species, and to determine the characteristics of land utilization and time expenditure by sportsmen.

Summary: No work was conducted during this segment since budgetary restrictions precluded purchase of the postcards necessary for conducting this job.

Target Date: 31 May 1979.

Status of Progress: Behind schedule

Significant Deviations: No work during one segment due to non-appropriation of funds.

Recommendations: This job should be reactivated when sufficient monies are allocated to allow for printing and mailing of the questionnaire cards.

Job I-2 Statewide Beaver Harvest

Job Objectives: To determine the statewide harvest of beavers by trappers.

Summary: A total of 1,441 beaver were taken by 116 trappers in 102 towns during the 1974-75 beaver season. This take is 198 less than last season, but 247 more than a ten-year (1965-1974) average. Berkshire County yielded more than one third (35.0 percent) of the harvest. One half (50.0 percent) of the harvest, or 121 beaver, was taken in the first two weeks of the season.

Target Date: 31 May 1979

Progress: On schedule.

Significant Deviations: None

Recommendations: Continue tagging of beaver pelts and recording of harvest data in 1976, using the same methods as in the current segment.

Division Regulation 11 currently states that all beaver taken must be tagged within two days of the season's end. When these two days occur on a weekend, as they did in part this year, Division personnel must work overtime to comply with the tagging requirement. This would be acceptable if trappers brought their take in on those days, but this is generally not the case. Some installations have waited a full day with no beaver being presented, only to have several brought in on the following workday. Consideration should be given to revising the regulation to state: "All pelts must be tagged within two working days of the closing day of the open season."

Should regulations be enacted requiring the tagging of otter and fisher, this job should be amended to include tabulation of the harvest of those species.

Cost: \$1,570.60 (Project Leader man-days: 4)

Remarks: Procedures: Each successful beaver trapper is required by law to present his pelts to an official checking station for tagging and recording of data. Seven stations are maintained for two days at the close of the season. Pelts are tagged with locking metal game seals and harvest data are recorded on mimeographed forms and subsequently tabulated by month trapped, town and county trapped in, and type of trap used.

Findings: The 1974-75 beaver trapping season extended for 15 weeks from 15 November 1974 to 1 March 1975. Trappers took 1,441 beaver during this period. This take was 198 less than last year's harvest of 1,639, but 247 more than a ten-year average (1965-1974) of 1,194. There were 115 trappers (123 in 1973-74) taking a minimum of one beaver each, with a mean harvest of 12.4 beaver per trapper (range: 1 to 107). One confiscated pelt was presented for tagging by the Division of Law Enforcement. The number of successful trappers (116) was the second highest on record, being seven less than in 1973-74 (123), but 30 more than a ten-year (1965-74) average

Beaver colonies located in 106 towns contributed to the 1974-75 harvest (Figure 1). The twelve towns with largest individual seasonal harvests are listed in Table 1.

MASSACHUSETTS

CITY AND TOWN LINES

Table 1. Beaver harvest for twelve Massachusetts towns, 1969-1975 seasons.

<u>Town</u>	<u>1968-69</u>	<u>1969-70</u>	<u>1970-71</u>	<u>1971-72</u>	<u>1972-73</u>	<u>1973-74</u>	<u>1974-75</u>
Becket	41	20	10	51	57	52	60
Blandford	10	22	16	29	78	37	59
Chesterfield	14	24	12	16	13	22	41
Goshen	-	-	11	15	17	17	33
Great Barrington	3	1	-	8	2	13	36
Monterey	35	-	5	18	24	12	36
New Marlboro	32	14	1	52	69	64	36
Otis	36	18	14	64	72	72	85
Petersham	52	18	27	58	57	66	42
Sandisfield	49	13	8	52	6	62	73
Savoy	9	12	18	20	21	34	36
Windsor	16	19	14	36	26	41	39

Table 2. Beaver harvest by county for Massachusetts, 1973-74 and 1974-75

<u>County</u>	<u>1973-74</u>			<u>1974-75</u>			<u>Percent Change</u>
	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>	<u>No. Beaver</u>	<u>Percent</u>	<u>Rank</u>	
Berkshire	518	31.6	1	505	35.0	1	- 2.5
Essex	16	1.0	8	13	0.9	7	-18.8
Franklin	370	22.6	2	324	22.5	2	-12.4
Hampden	148	9.0	5	133	9.6	5	- 6.8
Hampshire	288	17.6	3	263	18.3	3	- 8.7
Middlesex	51	3.1	6	29	2.0	6	-43.1
Plymouth	17	1.0	7	5	0.3	8	-70.6
Worcester	231	14.1	4	164	11.4	4	-29.0
Totals	1,639	100.0		1,441	100.0		-12.1

Table 3. Beaver harvest by month in Massachusetts, 1973-74 and 1974-75.

<u>Month</u>	<u>1973-74</u>		<u>1974-75</u>	
	<u>No. Beaver</u>	<u>Percent</u>	<u>No. Beaver</u>	<u>Percent</u>
November	878	53.6	721	50.0
December	492	30.0	449	31.2
January	185	11.3	176	12.2
February	84	5.1	67	4.6
Not stated	-	-	28	2.0
Totals	1,639	100.0	1,441	100.0

The western region of the state continued to provide the majority of the beaver harvest. Harvests declined in all counties (Table 2) but this change was more apparent in the eastern region. During the 1974-75 season, 1,026 beaver (71.2 percent) were taken west of the Connecticut River and 415 (28.8 percent) were taken east of the river, as opposed to 1,053 (64.6 percent) and 581 (35.4 percent) during 1973-74.

During the past season, Berkshire and Franklin Counties together yielded over half (829, 57.5 percent) of the total harvest. Hampshire, Worcester, and Hampden Counties contributed another 565 beaver (39.3 percent), with three other counties comprising the small remainder. Six counties reported no beaver taken. County rankings remained essentially unchanged.

As in past years, success was greatest during the initial weeks of the trapping season (Table 3). One half (50.0 percent) of the harvest, 721 beaver, were taken in the initial two weeks (15-30 November) of the season--a slight decrease from 1973-74 when 54.2 percent of the harvest was taken during the equivalent period. The take declined steadily each month thereafter with only a minimal harvest (67, 4.6 percent) in February.

Use of the Conibear trap dropped slightly this season, with 906 beaver (62.9 percent) taken in that style trap, as opposed to 1,091 (66.6 percent) in 1973-74.

The average price of a Massachusetts beaver pelt was \$18, the same as last season, with the 1974-75 harvest valuation totalling \$36,025.

MASSACHUSETTS DIVISION OF FISHERIES AND WILDLIFE
Bureau of Wildlife Research and Management

Approved: _____

Colton H. Bridges, Deputy Director

Prepared by _____

James E. Cardoza
Game Biologist

Date _____

PERFORMANCE REPORT

U OF MASSACHUSETTS LIBRARY

STATE Massachusetts Project No. W-35-R-17
Project Title: Game Population Trend and Harvest Survey
Period Covered: 1 June 1975 to 31 May 1976
Work Plan No. and Title: I - Statewide Game Harvest

Work Plan Objective: To determine the statewide harvest of selected small game and furbearer species and to determine the characteristics of land utilization and time expenditure by sportsmen.

Job No. and Title: I-4. Gray Squirrel Harvest Evaluation

Job Objective: To identify the characteristics (seasons, hunting techniques, and hunter preferences) of gray squirrel hunting in other states in order to evaluate what changes could be made in Massachusetts to improve hunter utilization.

Brief Summary: States east of the Great Plains were surveyed to determine squirrel season regulations, hunting methods, and factors affecting the relative popularity of squirrel hunting. Twenty-two states had straight seasons, eight had zoned seasons, five had split seasons and one state reported a zoned split season. The average season length was 120 days ($\bar{x} = 119.8$; S.D. = 61.7) with approximately 40 percent ($\bar{x} = 38.9$; S.D. = 23.2) of the seasons before leaf fall. Squirrels were ranked in the top four game species in 64, 85 and 70 percent of the states in terms of hunter preference, number killed and hunter participation, respectively. Hunting before leaf fall was judged more popular in 58 percent of 26 states responding by hunter preference and in 66 percent of 32 states responding by number of people participating. Still or stalk hunting was more common before leaf fall and wait-hunting was more common after leaf fall. The use of dogs to tree squirrels increased sharply after leaf fall. Shotguns outnumbered the use of rifles by at least 2 to 1 in squirrel hunting. Season length appeared to be a minor factor since only extremely short seasons (<50 days) may influence relative popularity of the sport. Opening date is the most important factor and there was a significant ($P < .05$) relationship between a large percentage of the season before leaf fall and high relative popularity rankings. The popularity of hunting before leaf fall was also significantly ($P < .05$) related to the relative popularity of squirrel hunting.

Target Date: 14 February 1975.

Status of Progress: On schedule.

Significant Deviations: None

Recommendations: An earlier opening date (well before leaf fall) and a longer open season extending later in the year are the regulation changes which could potentially stimulate the sport hunting of gray squirrels in Massachusetts.

I foresee little difficulty in terms of population management or political repercussions in extending the present season later into the year. I am recommending a 31 December closing date. Although I do not believe this will significantly increase the popularity of squirrel hunting, it will provide additional recreational opportunity and may stimulate the use of dogs in treeing squirrels.

An earlier opening date could significantly increase the popularity of gray squirrels as a game animal. The primary biological and political concern would be to find the mean date of cessation of nursing of young squirrels. The effect of hunting while females are still nursing has not been proven to be a factor in limiting yearly population levels, but I believe the desired result of increasing the utilization of squirrels could be achieved. This data would also eliminate any humane considerations involved with shooting nursing females.

Three management options are presented below for increasing the use of the gray squirrel resource and determining the date when nursing females would not be taken:

1. September 1 opening coupled with bag checks of selected hunters.
2. September 20 opening with prior sampling of Division personnel and bag checks of selected hunters after opening date.
3. September 20 opening, assuming the Ohio data on cessation of nursing is applicable to Massachusetts.

I am recommending an experimental three-year season under Option No. 1. This would provide us with the best data on cessation of nursing with the least expenditure of manpower. Any change in the popularity of squirrel hunting will be monitored by telephone survey techniques.

Cost: \$1,500

Presentation of Data:

This paper identifies some characteristics of squirrel hunting in the eastern United States and relates hunting season characteristics to the relative popularity of the sport.

Squirrel hunting has never been considered popular in Massachusetts (Cardoza 1971), yet in most states where squirrels are hunted, they are considered one of the top game species. This study examines hunting season regulations, hunting techniques, and popularity of hunting to determine what regulatory changes might be made to improve the utilization of the gray squirrel (Sciurus carolinensis) in Massachusetts. The importance of hunting season regulations to the relative popularity of a game animal has not received much attention in the literature but considerable work has been done on the effects of regulations on game population levels and hunting pressure (Allen 1943, Gale 1954, Nixon et al. 1974, Nixon et al. 1975, Uhlig 1956, U.S. Dept. of Interior 1975).

I acknowledge Sarah Daniels and Thomas Early, Massachusetts Division of Fisheries and Wildlife, who assisted with preparation and analysis of the questionnaire.

METHODS

A mail questionnaire (Appendix I) was sent in January 1975 to the chief game biologist in each state's wildlife management agency, requesting information on the type and dates of the gray squirrel season. Fox squirrels (Sciurus niger) are present in many of the states, but no state reported separate hunting regulations for the species. All questions other than the first referred simply to squirrels, so no distinctions are made between species in this report.

Eight states had zoned seasons, and the length of the season for each state was calculated by using the earliest opening date and the latest closing date. The season length for the five states with a split season was based on the number of days the season was opened. Only one state had both a zoned and split season so I averaged the number of hunting days for each zone to determine season length; the most frequently used opening and closing dates were used when appropriate.

Statistical inferences in this report are based on t-test and chi-square (χ^2) test, and Yates' correction for continuity (χ^2 adj.), $\alpha = .05$.

RESULTS

All of the 37 states east of the Great Plains have open seasons on the gray squirrel. Only one of these states did not respond to the questionnaire.

Types of Seasons

The three basic types of squirrel hunting seasons are straight, split and zoned. The straight season is the most common (22 states) followed by zoned (eight states) and split seasons (five states). Virginia was the only state reporting a zoned split season.

Zoned seasons consist of separate opening and usually closing dates for various regions within the state. Generally, these zones appeared to be based on climatic differences, but some political differences were apparent. For example, one state opened public land before private.

Appendix I

Massachusetts Division of Fisheries and Game
Westboro, Massachusetts 01581

Name: _____ State _____

1. What are the opening and closing dates of the gray squirrel hunting season in your state? (Indicate if season is zoned).

Opening _____ Closing _____

2. How would you rank squirrel hunting in your state in comparison to other sport hunting? (Circle one):

a. By hunter preference	1st	2nd	3rd	4th	Other	_____
b. By number of people participating:	1st	2nd	3rd	4th	Other	_____
c. By number killed:	1st	2nd	3rd	4th	Other	_____

3. Generally, in what week of what month does peak (over 50%) leaf fall occur?

Month _____ Week _____

4. In your opinion is the squirrel season more popular before or after leaf fall?

- a. By hunter preference?
b. By number of people participating?

5. Please review these definitions:

- a. Stalk or Still Hunting: Moving stealthily through the woods, frequently stopping for short periods of time (less than five minutes).
b. Wait Hunting: Moving to a likely spot and then waiting for a long period of time (more than five minutes) before moving again.
c. Dog Hunting: Using a dog to "tree" or locate a squirrel.

Referring to the above definitions, answer each question below independently:

- a. What percentage of each type hunting best describes squirrel hunting before leaf fall?

_____ % stalk _____ % wait _____ % dog

- b. What percentage of each type hunting best describes squirrel hunting after leaf fall?

_____ % stalk _____ % wait _____ % dog

- c. Indicate your estimate of hunters using shotguns and/or rifles:

For stalk hunting	_____ % shotguns	_____ % rifles
For wait hunting	_____ % shotguns	_____ % rifles
For dog hunting	_____ % shotguns	_____ % rifles

6. Indicate the numbers of the questions which were answered using survey data you have collected:

Question numbers _____

The Commonwealth of Massachusetts

Division of Fisheries and Game

Field Headquarters, Westboro 01581

Dear Sir:

In Massachusetts we are considering a change in our squirrel hunting regulations in an effort to provide more recreational opportunity to our sportsmen. Enclosed is a questionnaire which will help us describe the sport of squirrel hunting in the United States and perhaps identify how our regulations might be changed for the better.

The enclosed questionnaire is designed to be folded, stapled and mailed without an envelope. If survey data is not available to answer the questions please give your opinion.

Due to our regulation change procedure, this questionnaire will be analyzed starting 15 February 1975.

We invite your comments and appreciate your cooperation.

Sincerely yours,

Chet M. McCord
Chief of Wildlife Research

CMM:mb

Enclosure

A season which is divided into distinct segments within one year is defined as a split season. One state reported a spring and fall segment, but most split seasons were simply early and late fall or winter segments. Closing the season during the open deer season was reported by only one state, but I suspect it occurs in others.

Length and Time of Year of Hunting Seasons

The average length of a squirrel season was 120 days ($\bar{x} = 119.8$; $N = 34$; $S.D. = 61.7$). New Hampshire (31 days) and Massachusetts (41 days) have the shortest seasons while North Dakota (year-round) and Oklahoma (230 days) have the longest.

September (33 percent) and October (39 percent) were the most frequent months for opening the season followed by August (14 percent), November and May (6 percent each), and June (3 percent). For closing the season, January (39 percent) was the most frequent month followed by December (25 percent), November and February (17 percent each), and March (3 percent).

The use of months to compare opening and closing dates is not valid here since a September opening in Alabama would be much earlier seasonally than a September opening in Michigan. I asked each biologist for the month and week when peak (over 50 percent) leaf fall generally occurred. These data were used to standardize the seasonal time of year when the hunting seasons began. The average percent of the season before leaf fall was 39 ($\bar{x} = 38.9$; $N = 35$; $S.D. = 23.2$). Massachusetts, New Jersey and Pennsylvania were the only states not reporting hunting before leaf fall.

Relative Popularity of Squirrel Hunting

Each state was asked to rank squirrel hunting in comparison to other sport hunting. Comparisons were made in hunter preference, number of people participating and number of animals harvested (Table 1).

Over 64 percent of the states ranked squirrel in the top four species by hunter preference, and 41 percent ranked them first or second. By hunter participation and number harvested, squirrels were in the top four species in over 70 percent and 85 percent of the states, and ranked first and second in 50 and 56 percent of the states respectively.

Each biologist was asked if he felt squirrel hunting was more popular before or after leaf fall with reference to (1) hunter preference and (2) the number of people participating. Hunting before leaf fall was more popular in 58 percent of the 26 states responding by hunter preference and in 66 percent of 32 states responding by number of people participating.

Hunting Techniques

Three types of squirrel hunting were defined:

Stalk or still-hunting - Moving stealthily through the woods, frequently stopping for short periods of time (usually less than five minutes).

Wait-hunting - Moving to a likely spot and then waiting for a long period of time (more than five minutes) before moving again.

Dog-hunting - Using a dog to tree or locate a squirrel.

I asked each biologist to use the above definitions and give what he felt was the percentage of each type hunting that best described squirrel hunting before and after leaf fall (Table 2). Stalk-hunting was the most commonly used before leaf fall while wait-hunting was the most common after leaf fall. However, the real change in hunting techniques after leaf fall was a decrease in stalk-hunting and an increased use of dogs. Redmond (1953:383) reported a season average of 75 percent still-hunting and 25 percent dog-hunting; before leaf fall 96 percent was still-hunting while after leaf fall the use of dogs increased to 85 percent of the hunting.

Type of Firearms Used

The original question asked for an estimate of the percentage of hunters that use shotguns or rifles for each type hunting method. The responses, almost invariably, gave a single percentage or the same percentage for all three methods, so I will report only the data given for the first method listed realizing these data really represent a figure for all methods together.

The 30 states with usable responses showed a 68 percent use of shotguns and a 32 percent use of rifles. These percentages of rifle use are higher than Redmond (1953:383; 11 percent) and Allen (1952:88; 18 percent) reported. I believe that my rifle-use data is inflated because several states, indicating they had no survey data on percentage of rifle use, reported rifle use above 90 percent.

Season Length Related to Hunting Popularity

The length of the hunting season did not appear to be a major factor in these data since comparisons of the mean season length among states with different popularity rankings showed no trends (Table 3). However, the variability in season length was much higher for the states that ranked squirrel hunting below fourth in all the popularity categories. Empirically, this would indicate that an extremely short or long season may be a factor in squirrel hunting popularity. The three states with seasons less than 50 days all ranked squirrel hunting below fourth while the four states with seasons longer than 200 days were split with two ranking squirrel hunting second in preference and participation and two ranking squirrels below fourth in both categories. I believe an extremely short season can be a factor in the popularity of squirrel hunting particularly when combined with a late opening date.

Length of the Season Before Leaf Fall Related to Hunting Popularity

States with a larger percentage of the season before leaf fall generally rank preference for squirrel hunting higher than states with smaller portions of the season before leaf fall (Table 4). States ranking squirrel hunting as first or second in preference had a significantly larger percentage of the season before leaf fall than states ranking squirrel hunting as "Other" (t-test; $P < .05$). States ranking squirrel hunting as third and fourth also showed a significantly larger percentage of the season before leaf fall than the states ranking as "Other", but the difference was significant at a lower level of confidence (t-test; $P < .10$).

Table 1. The relative ranking of squirrel hunting in comparison with other sport hunting in 34 of the 37 eastern states with squirrel seasons.

Rank	Hunter Preference		Hunter Participation		Number Harvested	
	No. of States	Percent	No. of States	Percent	No. of States	Percent
1	6	18	7	21	11	32
2	8	23	10	29	8	23
3	3	9	2	6	6	18
4	5	15	5	15	4	12
Other	12	35	10	29	5	15
Total	34	100	34	100	34	100

Table 2. The percent of stalk, wait and dog hunting which best describes squirrel hunting before and after leaf fall.

Leaf Fall	Hunting Technique			Number of States Responding
	Stalk	Wait	Dog	
Before	51	45	4	28
After	39	46	15	30

Table 3. The mean length of the squirrel hunting season for states which ranked squirrel hunting popularity in similar categories.

States' Rank of Squirrel Hunting	Hunter Preference			Hunter Participation			Number of Squirrels Killed		
	Mean			Mean			Mean		
	N	Days	S.D.	N	Days	S.D.	N	Days	S.D.
1	6	108.3	15.0	7	110.4	14.8	11	112.0	12.9
2	8	138.5	52.5	10	134.9	47.8	8	136.0	59.8
3	3	125.3	25.7	2	122.0	31.0	6	99.3	32.0
4	5	115.2	39.4	5	116.8	39.2	4	109.8	42.6
Other	12	131.5	83.1	10	119.9	97.8	5	174.0	110.2
Total	34	119.8	61.7						

The states ranking squirrel hunting as first and second by hunter participation showed a significantly higher percentage of pre-leaf fall hunting than states ranking squirrel hunting as third and fourth (t-test; $P < .05$; Table 4). States ranking squirrel hunting below fourth showed 35 percent of the season before leaf fall, which was not significantly different from either the means of the first and second or the third and fourth categories. This indicates that the length of the season before leaf fall is not the only factor which affects hunter participation.

The relative number of squirrels killed in a state is probably the least sensitive method of the three used to measure the popularity of squirrel hunting. Although the states ranking squirrel hunting first and second reported the highest percentage (40 percent) of the season before leaf fall, the difference was not significantly higher than the third and fourth categories except at a low level of probability ($P < .20$). Again, the "Other" states showed a relatively high percentage of the season before leaf fall, but the sample size in that category had dropped to only four states.

Popularity of Season Before Leaf Fall Related to Squirrel Hunting Popularity

The popularity of hunting before and after leaf fall was compared to the relative ranking of squirrel hunting in comparison to other sport hunting (Table 5). Due to sample size the top two or three rankings were combined and compared by X^2 analysis to the lower categories.

States ranking squirrels in the top three game species by preference and participation also ranked hunting before leaf fall more popular a significant portion of the time. By preference, biologists from states ranking squirrel hunting first and second judged hunting before leaf fall more popular ($X^2 = 6.00$, X^2 ad; = 4.21, d.f. = 1). This relationship was even stranger when the first through the third popularity rankings were compared to the fourth and "Other" categories ($X^2 = 9.76$, X^2 ad; = 7.43; d.f. = 1). By participation, the first and second ranked states did not show significant difference between before or after leaf fall hunting at the .05 level ($X^2 = 3.46$, X^2 ad; = 2.22, d. f. = 1), but the difference was significant when the first through the third categories were grouped and compared to the lower rankings ($X^2 = 5.72$, X^2 ad; = 4.07, d. f. = 1).

DISCUSSION

Opening dates, season length, and bag limits are the basic hunting regulations which can influence hunter attitudes or relative popularity. Bag limits have been shown to be an effective methods of increasing or decreasing harvest of same species (U.S. Department of the Interior 1975), and Christisen (1971:328) discusses how bag limits affect the incentive of hunters since "...a liberal limit beyond the abilities of the average hunter may depress the hunter's interest and initiative". However, Nixon (1975:10-11) found shooting one or more squirrels tended to increase hunter interest, but that squirrel hunters were not strongly motivated to kill a limit. Daily bag limits on squirrels do not appear to be an important factor in hunting popularity since most hunters are not successful

Table 4. The mean length of the season before leaf fall for states which rank squirrel hunting in different categories of popularity.

States' Rank of Squirrel Hunting	Hunter Preference			Hunter Participation			Squirrels Killed		
	Mean			Mean			Mean		
	N	Days	S.D.	N	Days	S.D.	N	Days	S.D.
1 and 2	14	43.9	19.1	17	45.1	21.8	19	39.6	23.8
3 and 4	8	44.6	26.7	7	27.4	17.9	10	29.9	21.0
Other	11	28.1	23.2	9	35.0	27.0	4	35.0	22.2

Table 5. Popularity of squirrel hunting before and after leaf fall related to the relative ranking of squirrel hunting to other sport hunting within a state.

States' Rank of Squirrel Hunting	Leaf Fall			
	Preference		Participation	
	Before	After	Before	After
1	4	1	5	1
2	6	1	8	2
3	2	0	2	0
4	2	1	2	2
Other	1	8	4	6
Totals	15	11	21	11

in killing even one squirrel much less the limit (Allen 1954:92; Nixon 1974:70; Uhlig 1959:388).

Season length does appear to affect squirrel hunter attitudes and perhaps the popularity of the sport. Allen (1943:341) demonstrated that a short (22 day) season affected hunter activities since the hunting pressure remained high throughout the season, yet other studies dealing with longer seasons refer to a rapid decline in hunting pressure after the first couple of weeks (Allen 1952:100; Nixon et al. 1974:71; Nixon et al. 1975:10; Uhlig 1956:58). The increased variability in season length for states that ranked squirrel hunting low in popularity plus the low rankings by the states with short seasons indicates an extremely short season could be a factor in low popularity.

The single most important factor in squirrel management is the opening date (Uhlig 1959:389). This conclusion is based on the second or summer litter of squirrels being independent and available for hunting. The controversy over the importance of this summer litter in population maintenance and fall population levels has never been completely resolved. Allen (1952:106) states that the time squirrel season opens is a compromise between availability of the summer litters and the interest and success of the hunter. He says that in the early season when the mast ripens, squirrels are most active and hunting success is high. Nixon et al. (1975:8, 10) showed increased hunter success early in the season and states, "Hunters generally stopped hunting squirrels...when leaf fall was virtually complete and the opening of the upland game season attracted them to other areas and other species." The questionnaire data also indicates a strong predilection for hunting before leaf fall by both hunter preference and participation. The significant relationship between a high percentage of the season before leaf fall and high popularity of the sport, plus the significant relationship between relative popularity ranking of squirrel hunting and the popularity of hunting before leaf fall leads me to conclude that the opening date is a key factor in the relative popularity of squirrel hunting.

Obviously, there are many factors which work together to influence the popularity of squirrel hunting. The past trends and present abundance of squirrels, hunter success, the desirability of hunting other species which are in season at the same time, social traditions associated with the sport (could be a positive or negative stimulus), and the regulations are some of the important factors which can influence the relative popularity of squirrel hunting. Any one or combination of these factors may be the most important in any particular state.

CONCLUSIONS

A straight fall and winter hunting season of about 120 days with 40 percent of the season before leaf fall is the most common type of squirrel season in the eastern United States. Hunting is most popular before leaf fall when the stalk or still-hunting method is most often used. After leaf fall, wait-hunting is more common and the use of dogs increases sharply. Shotguns outnumber rifles at least two to one in squirrel hunting.

Squirrels are one of the top four game species in most of the 37 eastern states where they are hunted and frequently are rated first or second in terms of hunter preference, hunter participation and number killed. The relative popularity of squirrel hunting is related to the length of the season before leaf fall, and an early opening date appears to be the most important regulation. Extremely short seasons appear to be another factor which affect popularity of the season.

LITERATURE CITED

- Allen, D. L. 1943. Michigan fox squirrel management. Michigan Dept. Conserv., Game Div. Publ. 100, Lansing. 404 pp.
- Allen, J. M. 1952. Gray and fox squirrel management in Indiana. 2nd ed. Indiana Dept. Conserv., P-R Bull. 1, 112 pp.
- Cardoza, J. E. 1971. Statewide small game harvest. Mass. Div. Fish and Game P-R Proj. Rept. W-35-R-13. 21 pp.
- Christisen, D. M. 1971. Importance of daily bag in squirrel hunting. Trans. N. Am. Wildl. Conf. 36:322-329.
- Gale, L. R. 1954. The effect of season changes on hunting effort and game kill. Proc. Southeast Assoc. Fish & Game Commissioners. 7: 117-120.
- Nixon, C. M., R. W. Donohoe, and T. Nash. 1974. Overharvest of fox squirrels from two woodlots in western Ohio. J. Wildl. Manage. 38(1):67-80.
- _____, M. W. McClain, and R. W. Donohoe. 1975. Effects of hunting and mast crops on a squirrel population. J. Wildl. Manage. 39 (1): 1-25.
- Redmond, H. R. 1953. Analysis of gray squirrel breeding studies and their relation to hunting season, gunning pressure and habitat conditions. Trans. N. Am. Wildl. Conf. 18:378-389.
- Uhlig, H. C. 1956. The gray squirrel in West Virginia. Conserv. Comm. W. Va. Bull. No. 3, 83 pp.
- _____. 1959. Squirrel management and research. Pages 387-389 In V. Flyger (Editor) Symposium of the gray squirrel. Maryland Dept. Res. Education. No. 162.
- U.S. Department of the Interior, Fish and Wildlife Service. 1975. Final environmental statement for the issuance of annual regulations permitting the sport hunting of migratory birds. Washington, D. C. 710 pp.

Submitted by:

MASSACHUSETTS DIVISION OF FISHERIES AND WILDLIFE
Bureau of Wildlife Research and Management

Approved: _____

Arthur W. Neill, Superintendent

Prepared by _____

Chet M. McCord

Chief of Wildlife Research

Date _____

PERFORMANCE REPORT

U OF MASSACHUSETTS LIBRARY

STATE

MASSACHUSETTS

Project Type

Research and SurveyProject No. W-35-R-17

Project Title

Game Population Trend and Harvest Survey

Period Covered:

1 June 1974 to 31 May 1975Work Plan Number and Title: II - Massachusetts White-Tailed Deer Study

Work Plan Objective: To determine, through the collection and analysis of pertinent deer harvest data, the sex and age structure of the herd, and to develop management and harvest procedures based on project findings.

Job Number and Title: II-1 - Statewide Deer Harvest

Job Objective: To determine the annual harvest of deer in Massachusetts.

Brief Summary: The four western deer management units of Berkshire, Franklin, Hampden and Hampshire make up over 70 percent of the total deer harvest. The Worcester unit contributed 8 percent of the shotgun harvest, Barnstable unit 5 percent while the combined Region I and Region II contributed only 1 percent. The 1974 statewide harvest for all deer seasons was 2,781 deer, an increase of 560 over the 1974 harvest.

Target Date: 31 August 1975.

Status of Progress: On schedule.

Significant Deviations: None

Recommendations: Continue this job. See Job II-4 for future recommendations.

Cost: \$65,000

Presentation of Data:

Introduction: There were four types of deer seasons in 1974: (1) an 18-day archery season from 4 November through 23 November; (2) a 2-day paraplegic season from 4-5 November; (3) a 6-day shotgun season from 2 December through 7 December; and (4) a 3-day primitive firearms hunt from 9 December through 11 December. No hunting was allowed on Sundays.

Since 1967, Massachusetts has had a statewide antlerless deer permit system. During the one-week shotgun season, deer with antlers three inches or longer are legal game, but to harvest a male fawn or female deer the hunter must have an antlerless permit. Massachusetts regulations require hunters to report their kill within 24 hours to an official deer checking station.

Last season, the state was divided for the first time into ten deer management units and the distribution of antlerless permits was made on the basis of the new management units. The management units (Figure 1) and the number of permits issued for each is shown in Table 1.

Archery Season: A summary of the statewide deer harvest by archers shows an increase from 21 deer killed in 1967 to 87 deer in 1974. In 1967 the archers had a 12-day season, but in 1972 the season was increased six days for a total of 18. Archers in 1974 reported an increase of 10 deer over 1973 as shown below:

Summary of Archery Kill, 1967-1974

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
Males	17	21	27	24	26	49	51	62
Females	4	13	10	12	10	27	26	25
Totals	<u>21</u>	<u>34</u>	<u>37</u>	<u>36</u>	<u>36</u>	<u>76</u>	<u>77</u>	<u>87</u>

Seventy-two (72) deer were taken on the mainland of which 44 were adult males, 6 male fawns and 22 females. On Nantucket Island, the archers reported 10 bucks and 3 does for a total of 13 deer. A buck and a doe were the only two deer reported on the Vineyard. The total of 87 deer was made up of 56 adult males, 6 male fawns and 25 females. The four counties with the highest kill were Franklin, 23 deer; Berkshire, 19; Hampshire, 13, and Nantucket, 13 (Table 2).

Paraplegic Season: A male and a female deer were reported for this 2-day hunt held on Martha's Vineyard. This is the first year deer have been taken on this hunt since it began in 1972.

Primitive Firearms Season. Primitive firearms hunters reported taking 26 deer during the second year of this 3-day season. The 1974 harvest represents an increase of 21 deer above the 1973 kill. It is interesting to note the change in the sex ratio from the 1973 harvest when 5 males and 2 females were reported to 8 males and 18 females in 1974. Seven males and 17 females were taken from the mainland while a buck and a doe were reported from the Vineyard. Berkshire County had the highest kill of 15 followed by Hampden, 4; Franklin, 3; Hampshire, 2; and Nantucket, 2 (Table 2).

Shotgun Season: Shotgun hunters reported harvesting 2,666 deer including 1,878 males (214 male fawns) and 788 females (Table 2). The four top deer-producing counties were Berkshire, 1,056; Franklin, 448; Hampden, 231; and Worcester, 213 (Table 1). The shotgun harvest constitutes 96 percent of the total statewide deer harvest.

The mainland deer harvest shows 1,524 adult males, 159 fawn males and 602 females for a total of 2,285 deer. The Nantucket Island hunters reported 62 antlered males, 26 male

fawns, and 66 females for a total of 154 deer. On Martha's Vineyard, 73 adult males, 29 button bucks and 95 females were taken for a total of 197 deer. Five males and 25 females were reported harvested on Gosnold Island.

Table 3 presents a summary of the 1974 Massachusetts shotgun deer harvest by sex and the county rank of importance from 1970 through 1974. The top ranking deer-producing counties, Berkshire and Franklin, have not changed for the past five years. Hampden County moved to third position in 1974 exchanging positions with Dukes County which moved to fifth. Worcester County remained in fourth position while Hampshire exchanged positions with Nantucket and moved into the sixth position. The remaining counties have been in the same lower ranks for many years.

Total Harvest Figures: A ten-year summary (1965-1974) of the annual deer harvest by town and county is presented in Appendix I. From 1967, when the antlerless deer permit system was initiated, to 1970 the statewide deer harvest increased from 1,193 deer to 2,403. The harvest figure dropped to 2,284 in 1971 and to 2,121 deer in 1973. The 1974 harvest of 2,781 deer is an increase of 560 deer over the 1973 figures and 378 deer over the 1970 high of 2,403 deer. It has been demonstrated in previous job performance reports (W-35-R:II-1, 1971, 1972 and 1973) that the statewide deer harvest fluctuates in relationship to the number of antlerless permits issued annually.

The percent of females in the total harvest from 1967 through 1974 shows the percent of females increased from 20 percent in 1967 to 39 percent of the harvest in 1971. Since that time, the percent of females has been reduced to a healthy 30 percent as shown below:

<u>Percent of Females in Total Harvest</u>							
<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
20%	23%	29%	32%	39%	34%	30%	30%

A summary of the 1974 deer harvest per sex per management unit is presented in Table 4 and the number of each sex killed statewide since 1967 is shown below:

<u>Kill Statewide for Each Sex, 1967 through 1974</u>								
	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
Male	954	1104	1451	1629	1385	1504	1477	1949
Female	239	323	595	776	899	787	644	823
Total	1193	1427	2046	2406	2284	2291	2121	2781

The ratio of male to female deer in the statewide deer harvest from 1967 to 1974 shows the ratio of females increasing annually from 1 male to .25 females in 1967 to 1 male to .65 females in 1971. A reduction of antlerless permits in 1972 helped to reduce the male to female ratio from 1 male to .52 females in 1972 to 1 male to .43 females in 1974 as shown:

Ratio of Males to Females in Total Harvest, 1967-1974

	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
Ratio	1:25	1:29	1:41	1:48	1:65	1:52	1:44	1:43

Table 2 shows the total harvest by sex, age and type season for each management unit and for the entire state.

Antlerless Permit Data. Antlerless permit applications in 1974 were 2,000 less (32,000 applicants) than the previous year (Table 5). There were 4,000 sportsman, 358 farmer-landowner, 400 Nantucket Island and 600 Martha's Vineyard permits issued in 1974. The statewide harvest for antlerless deer permittees was 1,180, but the largest number ever reported by this group was 1,289 deer in 1971.

The farmer-landowner permittees reported harvesting 80 deer in 1974. This is the largest harvest since 1968 when 65 deer were taken (Table 5). Twenty-two percent of the permittees took deer. The bulk of the 358 farmer-landowner permits were issued for the four western units in order of importance: Franklin, 34%; Berkshire, 32%; Hampden, 12%; Worcester, 11%; Hampshire, 10%. Three permits were issued for Region I, but there were no applications or permits issued for Region II, Barnstable, Martha's Vineyard and Nantucket.

The 600 antlerless permit holders for Martha's Vineyard reported 22 male fawns, 31 antlered males and 120 females during the 1974 shotgun season. With some minor fluctuations, the deer harvests were the same as the 1973 data (Table 5).

On Nantucket Island, the male fawn harvest by 400 antlerless permittees decreased slightly from 29 in 1973 to 26 button bucks in 1974. The adult male kill remained unchanged at 22 antlered bucks in both 1973 and 1974. The female harvest was almost the same with 65 females reported in 1973 and 66 females in 1974 (Table 5).

The 1974 mainland harvest of deer by 4,000 antlerless permit holders shows the harvest of adult males was slightly lower (61) than the 1973 figure of 67 adult males. The male fawn harvest increased from 102 in 1973 to 150 in 1974. An increase of 123 females was reported by antlerless permit holders or 602 females in 1974 compared to 379 females in 1973 (Table 5).

Antlerless Permittee Success Ratio. The hunter success ratio on the mainland remained unchanged in 1974 with 1 deer killed for every 5 permits issued. Nantucket and Martha's Vineyard antlerless permit holders' success ratio for 1974 was 1 deer per 4 permits.

Table 6 presents a summary of the Massachusetts deer harvest per antlerless permit per deer management unit and the success ratio of permit holders for the 1974 shotgun season. The success ratio of permit holders in the Berkshire unit was 1 deer per 3 permits. For Franklin and Hampden units the success ratio was 1 deer for 5 permits while in Hampshire the ratio was 1 deer for 6 antlerless permits. The Worcester unit permit holders reported their success of 1 deer for 8 permits; and Region II and I permit holders reported 1 deer for 29 permits and 1 deer for 40 permits respectively.

Job Number and Title: II-2, Non-Hunting Deer Mortality Investigations

Job Objectives: To determine the annual non-hunting decimating factors of the Massachusetts deer herd.

Brief Summary: From 1 January to 31 December 1974 there was a 10.4 percent increase in non-hunting deer mortality over 1973 reports. The 467 deer mortalities consisted of 179 males (42%) and 246 females (52%) and 42 of unknown sex (6%). Motor vehicles caused the highest mortality, killing 347 deer while 35 were killed illegally, 33 by dogs, 11 drowned, 5 caught in fences, 2 by trains, 1 in crop damage, and 33 due to unknown causes.

Target Date: 31 December 1974

Status of Program: On schedule.

Significant Deviations: None

Recommendations: Project should continue as it presently exists.

Cost: \$14,000

Presentation of Data:

Techniques. Deer mortality reports are made by Natural Resources officers to the Law Enforcement office in Boston. A copy of each report is provided to the Division.

Findings. A summary of the sex classes and the causes of Massachusetts deer mortalities for 1 January through 31 December 1974 is presented in Table 7. During this period, 467 deer mortalities were reported. Of these, 179 were males, 246 were females and no sex data on 42 deer. In order of importance, the number and causes of deer mortalities are as follows: motor vehicle, 347; illegal, 35; dogs, 33; drowned, 11; fences, 5; trains, 2; crop damage, 1; and other and unknown causes, 33.

Table 8 presents a five-year summary of deer mortalities per cause from 1969 through 1974. Although the 1974 mortalities are lower than the five-year average, there was an increase in the number of deer mortalities compared to the 1973 figures.

A comparison of the total non-hunting deer mortalities of Massachusetts deer from 1968 through 1974 is presented in Table 9. The 1974 reported mortality of 469 deer is 10.4 percent higher than the 1973 figure of 420 deer.

Table 10 presents a comparison of the actual numbers of deer mortalities by sex and the adjusted sex ratios per month. The sex data was adjusted on a monthly basis with sex of the larger percentages assigned any fractional part of a deer so that there are only whole deer in the adjusted sex ratio. The adjusted sex ratio of the 1974 non-hunting deer mortalities are:

<u>Males</u>		<u>Females</u>
198	:	269
74	:	100
42%	:	58%

(1973 sex ratio = 47.6% males:52.4% females)

The non-hunting deer mortalities ranked by counties for a seven-year period 1968 through 1974 is presented in Table 11. Berkshire County remains in number one position as having the largest number of non-hunting deer mortalities reported in 1974. Franklin County moved into second place with Barnstable County dropping to the third slot. Worcester County remains in the fourth ranking position. Hampden and Hampshire Counties were tied for fifth position in 1973. In 1974, Hampshire County remains in the fifth ranking position while Hampden County dropped to ninth place. Dukes County moved from ninth place in 1973 to sixth place in 1974. The rank of the six remaining counties may have changed slightly from year to year; however, no significant changes occurred during the 1974 non-hunting deer mortality reporting period.

Job Number and Title: II-3, Deer Fertility Studies

Job Objectives: To determine the reproductive rate per age class of the Massachusetts deer herd.

Brief Summary: Job was inactive during the period covered by this report.

Target Date for Job Completion: None

Status of Program: Inactive

Significant Deviations: None

Recommendations: Review old data and evaluate need for continuation.

Cost: None

Remarks: Inadequate funds for transportation forced the inactive status of this job.

Job Number and Title: II-4, Deer Management Recommendations

Job Objectives: To determine the size of the Massachusetts deer herd and recommend management techniques that will provide the deer hunter with the greatest hunting opportunity commensurate with herd population levels.

Brief Summary: The calculated minimal deer population was 12,984 deer. This was a 13.59 percent increase in the 1974 calculated minimal deer population.

The adult male harvest per square mile of deer range was .25 antlered on the mainland; .84 in Dukes County and 1.73 antlered bucks on Nantucket Island during the 1974 shotgun season. The adult female harvest per square mile of range was .07 does on the mainland; .69 deer for Dukes County and 1.23 does on Nantucket Island.

A summary of the total deer harvest shows that .38 males and .10 female deer per square mile of deer range were reported on the mainland. The harvest per square mile of deer range for Dukes County was 1.27 males and 1.40 females. On Nantucket, the harvest of deer per square mile of deer range was 2.73 males and 1.92 females.

The percent frequency of adult females to adult males was .30 on the mainland; .82 in Dukes County and .69 on Nantucket Island.

The 1974 deer harvest data indicates the number of sportsman antlerless permits issued per deer management unit in 1972 caused an increase in the male harvest this year where planned and controlled the harvest in the female segment.

Target Date: 30 June 1975

Status of Progress: On schedule.

Significant Deviation: None

Recommendations: The following numbers of sportsman permits issued per county and/or region are suggested:

<u>County</u>	<u>Number of Sportsman Permits</u>
Barnstable	200
Berkshire	1300
Franklin	700
Hampden	400
Hampshire	300
Worcester	700
Region I*	200
Region II**	200
Martha's Vineyard	600
Nantucket	400
Naushon	50

* Region I represents Essex, Middlesex and Norfolk Counties

**Region II represents Bristol and Plymouth Counties

A big game license that incorporates an antlerless permit application should be investigated. The hunter will be required to indicate in which county or region he would like the application to be drawn when he purchases the license.

Cost: \$750

Presentation of Data: A summary of the sex and age composition of Massachusetts deer at biological deer check stations for an eight-year period on the mainland and for four years on Martha's Vineyard and Nantucket are presented in Tables 12, 13, 14, and 15. The number of deer in any age class represents the number of deer aged in that class and the totals are the number of deer aged at the biological stations. Not all deer checked at a biological station are aged. Some deer have the lower jaw shot away, the jaws are frozen or the deer is on top of a car and inaccessible to the station operator. Therefore, the total number of deer handled by a station may differ from the total number of deer aged.

Massachusetts' shotgun harvest and the total number of antlerless permits issued per year from 1967 through 1974 is presented in Table 16. The shotgun hunters harvested 2,669 deer during the 1974 season. Of the deer taken, 1,878 were males; 1,665 adult males and 213 male fawns. The adult male harvest in 1974 was the highest adult male kill since the antlerless system started in 1967. The female harvest was 791 does of which 563 were adults and 228 were female fawns.

It is interesting to note that a comparison of the highest reported harvest of deer prior to 1974 occurred in 1970 when 2,367 deer were reported and 7,347 antlerless permits issued. In 1974 there were 5,358 antlerless permits issued or 1,989 less permits than in 1970 yet there was an increase of 302 deer taken in 1974. The increase in the 1974 harvest was in the adult male segment with button bucks slightly lower than the 1970 figure. There were slight increases in the adult female and fawn females reported in the 1974 harvest compared to the 1970 figures.

Table 17 presents a summary of the percentage changes in adult harvest and calculated minimal populations of deer in Massachusetts, 1967 through 1974. In 1974, the adult male harvest was 1,665 deer. This was an increase of 407 deer or 23.35 percent higher than the 1973 harvest of 1,258 adult males. The calculated adult male population was 3,107 for an increase of only .86 percent above the 1973 figure (3,080). There was a substantial increase (21.03 percent) in the calculated adult female deer population from 3,850 deer in 1973 to 4,660 adult does in 1974. The calculated minimal statewide deer population was 12,984 deer in 1974. This was an increase of 13.59 percent above the 1973 population of 11,431 deer.

A summary of the adult male and adult female deer harvest per square mile of deer range in Massachusetts from 1967 through 1974 is presented in Tables 18 and 19. There was a slight increase in the adult male and female harvest in all counties on the mainland and a slight decrease in the male and female harvests on Nantucket and Dukes Counties. The data presented in Tables 18 and 19 will become increasingly significant during the next few years. Prior to 1974 antlerless permits were issued for three deer management units; namely, the mainland (4000 permits); Nantucket (400 permits) and Martha's Vineyard (600 permits). Since mainland has been divided into eight management units, the effect of the antlerless permits on the management units will be observed and the number of permits issued for each unit can be increased or decreased dependent on the number of adult males and adult females harvested.

Table 20 presents a summary of the total harvest of deer in Massachusetts by sex, by county and the harvest of deer per square mile of deer range for the 1974 deer seasons (archery, primitive firearm, paraplegic and shotgun seasons). The island counties of Nantucket and Dukes have the highest reported harvest of deer per square mile with 4.65 deer and 2.67 deer respectively. On the mainland Berkshire has a reported harvest of 1.30 deer per square mile of range. This was the first time since the inception (1967) of the antlerless permit system that any mainland county had a reported harvest of one deer per square mile of deer range. Franklin County has a reported kill of .73 deer per square mile. Barnstable, Hampden and Hampshire have .47; .45 and .42 deer harvested per square mile of deer range respectively. Worcester County, with 1307.6 square miles of deer range, has only a reported .17 deer per square mile. The remaining counties; namely, Bristol, Essex, Middlesex and Plymouth have .02 deer or less reported. There were no deer reported harvested in Norfolk County.

Table 21 presents the percent frequency ratio of adult females to adult males from 1967 through 1974. It would appear that the deer management units of the mainland are in good shape in relationship to the harvest of adult females to adult males. All counties, with the exception of Middlesex, have reported .39 or less adult does to adult males. The island units of Nantucket and Dukes Counties have .69 and .82 adult does respectively per adult buck. The adult male harvest in Worcester County is expected to increase during the 1975 deer season.

A summary by management unit of the shotgun deer harvest, the number of sportsman antlerless permits issued, the square miles of deer range, the adult harvest per square mile of range and the percent frequency ratio of adult females to adult males for the 1974 shotgun deer season is presented in Table 22. The data in this report has been presented on a county basis, but in Table 22 the data is presented on a deer management unit basis.

Figure No. 1 presents a summary of the total Massachusetts deer harvest by sex per square mile of deer range by county for an eight-year period, 1967 through 1974. In Berkshire County, the deer harvest shows an increase in total harvest. The male harvest increased at a greater rate than the female. The female harvest declined slightly in Franklin County while the total harvest and the male harvest increased. It is felt that this is a healthy situation and a rise in the 1975 harvest of males is expected. In Hampden County the male harvest and the total harvest are increasing at about the same rate. The female harvest declined slightly in 1974. The deer harvest in Hampshire County showed a slight increase in the total harvest and the buck and doe harvests. The rate of increased harvest in this county appears to be healthy. In Worcester County there was a slight decline in all harvest data. It would appear that Worcester County was being overgunned but it is felt that this management unit has not recovered from being overgunned from 1969 through 1972. The distribution of antlerless permits on a deer management unit basis should relieve some of the pressure on the female segment of the Worcester County deer herd. The rate of increase of the total harvest and the female harvest appear to be about the same in Barnstable County. However, the female harvest is well below the male harvest and the male harvest is still showing a slight increase.

Figure No. 2 presents a summary of the total Massachusetts deer harvest by sex per square mile of deer range on the mainland, Dukes and Nantucket Counties from 1967 through 1974. On the mainland, the deer harvest appears to be as expected with a continued increase in the total harvest, a continued increase in the male segment, and a slight increase in the female segment. The deer harvest on Dukes County continued to take an expected decline with more females than males being reported. It may be necessary to reduce the number of antlerless permits for Dukes County in 1976. Nantucket deer management unit had an expected decline in the total harvest along with a decline in the male and female harvest. The decline should continue in 1975. When the same number of adult bucks and does are being reported the number of antlerless permits issued can be reduced.

MASSACHUSETTS DIVISION OF FISHERIES AND WILDLIFE
Bureau of Wildlife Research and Management

Approved:

Arthur W. Neill, Superintendent

Prepared by

James J. McDonough
Game Biologist

Date

Table 1. Summary of the number of antlerless permits issued, the deer harvest per deer management unit and the ranking order of importance as a deer-producing unit for the 1974 Massachusetts shotgun season.

<u>Unit</u>	<u>Number Antlerless Permits</u>	<u>Male</u>	<u>Female</u>	<u>Total</u>	<u>Rank</u>	<u>Percentage of Total</u>
Berkshire	1,300	763	238	1,056	1	39.6
Franklin	700	347	101	448	2	16.8
Hampden	400	181	50	231	3	8.7
Worcester	700	144	69	213	4	7.9
Martha's Vineyard***	600	102	95	197	5	7.4
Hampshire	300	125	43	168	6	6.3
Nantucket	400	88	66	154	7	5.8
Barnstable	200	91	42	133	8	4.9
Region I*	200	15	5	20	9	0.3
Region II**	<u>200</u>	<u>12</u>	<u>4</u>	<u>16</u>	9	0.6
	5,000	1,878	788	2,666		

* Region I includes Middlesex, Norfolk, Essex and Suffolk Counties.

** Region II includes Bristol and Plymouth Counties.

*** Gosnold 5 males and 30 males not included.

Table 2. A summary of the shotgun, archery and primitive firearms hunter harvest by county for 1974 in Massachusetts

County	Shotgun				Archery				Primitive Firearms				Total			% of Total
	M	F	T	BB	M	F	T	BB	M	F	T	BB	M	F	T	BB
Barnstable	91	42	133	6	3	3	3	1					94	42	136	7
Berkshire	768	288	1,056	79	9	10	19	1	4	11	15	3	781	309	1,090	83
Bristol	4	1	5										4	1	5	
Dukes*	102	95	197	29	2		2		1	1	2		106	97	203	29
Essex	13	4	17										13	4	17	
Franklin	347	101	448	31	18	5	23	4	1	2	3		366	108	474	35
Hampden	181	50	231	21	2	1	3		2	2	4		185	53	238	21
Hampshire	125	43	168	9	9	4	13			2	2		134	49	183	9
Middlesex	2	1	3		2	1	3						4	2	6	
Nantucket	88	66	154	26	10	3	13						98	69	167	26
Norfolk																
Plymouth	8	3	11	2									8	3	11	2
Suffolk																
Worcester	144	69	213	11	7	1	8						151	70	221	11
Gosnold	5	25	30										5	25	30	
Totals	1,878	788	2,666	214	62	25	87	6	8	18	26	3	1,949	832	2,781	223

* Gosnold not included.

Table 3. County summary of the 1974 Massachusetts shotgun deer harvest by sex and the county rank of importance from 1970 through 1974.

County	Male	Female	Total	Rank 1974	Rank 1973	Rank 1972	Rank 1971	Rank 1970
Barnstable	91	42	133	8	8	8	7	8
Berkshire	768	288	1,056	1	1	1	1	1
Bristol	4	1	5	11	12	12	12	12
Dukes*	102	95	197	5	3	3	3	7
Essex	13	4	17	9	9	9	10	9
Franklin	347	101	448	2	2	2	2	2
Hampden	181	50	231	3	5	4	5	6
Hampshire	125	43	168	6	7	6	8	4
Middlesex	2	1	3	12	11	11	9	10
Nantucket	88	66	154	7	6	7	6	5
Norfolk				13	13	13	13	13
Plymouth	8	3	11	10	10	10	11	11
Suffolk				13	13	13	13	13
Worcester	144	69	213	4	4	5	4	3
Total*	1,873	763	2,636					
Gosnold	5	25	30					
	1,878	788	2,666					

* Gosnold not included.

Table 4. A summary of the 1974 deer harvest per sex per management unit, shotgun season only.

Unit	Males	Females	Total	Male Fawns
Berkshire	768	288	1,056	79
Franklin	347	101	448	31
Worcester	144	69	213	11
Hampshire	125	43	168	9
Hampden	181	50	231	21
Region I	15	5	20	
Region II	12	4	16	2
Barnstable	91	42	133	6
Martha's Vineyard*	102	95	197	29
Nantucket	88	66	154	26
Gosnold	5	25	30	
Totals	1,878	788	2,666	214

*Gosnold not included.

Table 5. Number of permits and deer harvest by permittees from 1967 through 1974 per deer management unit in Massachusetts

	1967	1968	1969	1970	1971	1972	1973	1974
Permit Applications*	28,000	24,000	32,000	35,000	37,500	38,000	32,000	30,000
Mainland Permits	2,000	2,000	4,000	6,000	6,000	4,000	4,000	4,000
Adult Male			99	124	104	55	67	80
Fawn Male			107	183	175	136	102	148
Female			473	612	698	531	396	602
Totals			679	919	972	722	565	830
Farmer-Landowner Permits	243	331	295	347	270	326	349	358
Male	2	21	17	25	20	14	19	38
Female	20	44	45	34	25	42	37	42
Totals	22	65	62	59	45	56	56	80
Nantucket Permits			400	400	400	400	400	400
Adult Male			29	15	26	23	22	22
Fawn Male			11	24	29	17	29	26
Females			49	85	70	61	65	66
Totals			89	124	125	101	116	114
Martha's Vineyard Permits					600	600	637**	600
Adult Male			1	0	30	24	21	25
Fawn Male			0	1	41	37	35	29
Females			19	38	95	126	117	95
Totals			20	39	166	187	173	149
Total Permits Issued	2,243	2,331	4,695	6,747	7,270	7,326	5,386	5,358
Deer (both sexes)	279	356	788	1,082	1,260	1,010	854	1,093

* Round figures

** Includes Naushon Island

Table 6. A summary of the Massachusetts deer harvest per sportsman's antlerless permit per deer management unit and the success ratio of antlerless permit holders for 1974

County	Number of Applications	No. of Permits Awarded	No. of Antlerless Deer Harvest by Permit					Ratio of Deer per Antlerless Permit					Success Ratio	
			Button Buck		Adult Male		Total	Button Buck	Adult Male		Male	Female	Deer	Deer:Antlerless Permit
			Buck	Male	Male	Female	Male		Female					
Berkshire	8,343	1,300	73	38	111	238	1:18	1:34	1:12	1:5	399	1:3		
Franklin	6,508	700	31	12	43	101	1:23	1:58	1:16	1:7	144	1:5		
Worcester	4,886	700	10	11	21	59	1:70	1:64	1:33	1:10	90	1:8		
Hampshire	2,658	300	6	4	10	43	1:50	1:75	1:30	1:7	53	1:6		
Hampden	2,640	400	20	9	29	50	1:20	1:44	1:14	1:14	79	1:5		
Region I	829	200	0	2	2	5	0	1:100	1:100	1:40	7	1:29		
Region II	890	200	2	1	3	4	1:100	1:200	1:67	1:50	7	1:29		
Barnstable	1,786	200	6	3	9	42	1:33	1:67	1:22	1:5	51	1:4		
	28,540	4,000	148	80	228	602	1:27	1:50	1:18	1:7	830	1:5		
Martha's Vineyard	838	600	29	25	54	95	1:21	1:24	1:11	1:6	149	1:4		
Nantucket	587	400	26	22	48	56	1:15	1:18	1:33	1:6	114	1:4		
Statewide	29,965	5,000	203	127	330	753	1:25	1:39	1:15	1:7	1,093	1:5

Table 7. A summary of Massachusetts deer mortalities reported by Natural Resource officers, 1 January through 31 December 1974.

	January			February			March			April			May			June		
	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex
Motor Vehicles	5	5		4	2	3	6	9	1	8	10	3	13	20	5	17	21	7
Dogs	4	6		3		1	3	4		1								
Illegal Kill	1	1			1		2	1										
Drowned	1	1			1										2		1	1
Fences							1	1									1	
Trains																1		
Crop Damage																		
Other and Unknown							1	2	1	1	1		1			3		
Subtotal	11	13		7	5	4	13	17	2	10	11	3	14	20	7	21	23	7
Total (Cum. Total)		24			16 (40)		32 (72)			24 (96)			41 (137)			51 (188)		
	July			August			September			October			November			December		
	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex	M.	F.	No Sex
Motor Vehicles	5	26	2	3	9	1	6	15	1	20	16	3	37	28	8	9	18	1
Dogs		1					1	1			1			2			5	
Illegal					1		1	1			1		4		2	6	12	1
Drowned										1	1			2				
Fences													1	1			1	
Trains	1																	
Crop Damage						1												
Other and Unknown				1						1	2		3	4		2	7	
Subtotal	7	28	2	4	11	1	8	17	1	22	21	3	45	37	10	17	43	2
Total (Cum. Total)		37 (225)			16 (241)		26 (267)			46 (313)			92 (405)			62 (467)		
Cause	Male		Total	Unknown Sex		Total	Cause		Total	Male		Total	Female		Total	Unknown Sex		Total
Motor Vehicles	133	179	347	35		347	Fences		347	2		2	3		3			5
Dogs	12	20	33	1		33	Trains		33	2		2						2
Illegal	14	18	35	3		35	Crop Damage		35				1		1			1
Drowned	2	7	11	2		11	Other and Unknown		11	14		14	18		18	1		33
Totals										179		179	246		246	42		467

Table 8. Five-year summary of deer mortalities of Massachusetts deer reported by Natural Resources officers, 1969 through 1974.

<u>Cause</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>	<u>5-yr. Avg.</u>	<u>Total</u>
Motor Vehicles	397	400	373	321	321	347	362	1,812
Dogs	166	204	219	41	36	33	133	666
Illegal Kills	39	25	39	44	23	35	34	170
Crop Damage	2	14	4	1	2	1	5	23
Unknown Causes	51	38	41	35	21	33	37	186
All Other Causes	27	17	18	11	15	18	18	88
Totals	<u>682</u>	<u>698</u>	<u>694</u>	<u>453</u>	<u>420</u>	<u>469</u>	<u>589</u>	<u>2,945</u>

Table 9. A comparison of total non-hunting deer mortalities of Massachusetts deer from 1968 through 1974.

	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
No. of Deer	613	682	698	694	453	420	469
Percent Change	11%	2%	-.6%	-34.7%	-7.3%	+10.4%	

Table 10. Comparison of actual numbers of deer mortalities by sex* and adjusted data for Massachusetts deer per month, 1974.

<u>Month</u>	<u>Male</u>	<u>Female</u>	<u>Unknown Sex</u>	<u>Total</u>	<u>Adjusted</u>	
					<u>Male</u>	<u>Female</u>
January	11	13		24	11	13
February	7	5	4	16	10	6
March	13	7	2	32	14	18
April	10	11	3	24	11	13
May	14	20	7	41	17	24
June	21	23	7	51	24	27
July	7	28	2	37	7	30
August	4	11	1	16	4	12
September	8	17	1	26	8	18
October	22	21	3	46	24	22
November	45	37	10	92	51	41
December	17	43	2	62	17	45
Total	<u>179</u>	<u>246</u>	<u>42</u>	<u>467</u>	<u>198</u>	<u>269</u>

Adjusted Sex Ratio

198 males:269 females
 74 males:100 females
 42%males: 58% females

* These data were reported by Natural Resource officers.

Table 11. The Massachusetts non-hunting deer mortalities ranked by county, 1968 through 1974.

County	1968		1969		1970		1971		1972		1973		1974	
	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank	Total	Rank
Barnstable	55	5	63	4	74	4	93	2	72	2	76	2	56	3
Berkshire	134	1	168	1	184	1	175	1	92	1	109	1	148	1
Bristol	14	11	15	11	8	11	6	11	6	11	6	9	6	11
Essex	42	7	34	8	32	7	25	9	20	9	20	6	23	7
Franklin	81	3	90	3	127	2	90	3	66	3	73	3	62	2
Hampden	58	4	52	5	47	6	51	6	45	5	28	5	19	9
Hampshire	46	6	51	6	56	5	65	5	38	6	28	5	37	5
Middlesex	39	8	37	7	22	8	18	10	8	10	14	8	13	10
Norfolk	7	13	5	12			1	13	6	10	3	10	4	12
Plymouth	25	9	27	9	22	8	36	8	21	8	17	7	20	8
Suffolk	0	14	0	14	0	14	0	14	0	14	0	14	0	13
Worcester	86	2	104	2	99	3	99	3	55	4	37	4	42	4
Dukes	11	12	15	11	17	9	17	9	22	7	6	9	24	6
Nantucket	15	10	21	10	11	10	11	10	2	12	3	10	13	10

Table 12. Age composition of mainland Massachusetts male deer checked at biological stations, 1967 through 1974.

<u>Age</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
6 mos.	28	61	67	121	100	77	69	78
1-1/2	167	198	229	263	211	260	237	255
2-1/2	122	126	133	147	103	144	173	156
3-1/2	62	87	98	97	77	96	90	86
4-1/2	25	35	55	59	43	46	47	35
5-1/2	10	17	21	21	19	14	18	11
6-1/2	5	4	14	7	3	11	7	5
7-1/2	1	0	2	7	6	2	6	4
8 to 9-1/2	1	0	2	1	1	0	3	0
10-1/2	0	0	1	0	0	0	0	0
Totals	<u>421</u>	<u>528</u>	<u>652</u>	<u>723</u>	<u>563</u>	<u>563</u>	<u>650</u>	<u>730</u>

Table 13. Age composition of mainland Massachusetts female deer checked at biological stations from 1967 through 1974.

<u>Age</u>	<u>1967</u>	<u>1968</u>	<u>1969</u>	<u>1970</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
6 mos.	31	44	76	90	101	90	62	62
1-1/2	32	33	55	54	64	57	48	53
2-1/2	18	28	55	69	69	56	42	51
3-1/2	13	16	36	46	51	51	35	33
4-1/2	7	11	24	29	33	22	25	23
5-1/2	3	3	11	14	20	14	6	8
6-1/2	1	4	0	8	14	10	5	7
7-1/2	0	3	2	0	11	2	6	6
8 to 9-1/2	3	3	0	0	9	0	2	3
10-1/2	1	1	0	0	0	1	0	1
Totals	<u>109</u>	<u>146</u>	<u>259</u>	<u>310</u>	<u>372</u>	<u>303</u>	<u>231</u>	<u>247</u>

Table 14. Age composition of Martha's Vineyard, Massachusetts deer checked at biological stations from 1971 through 1974.

<u>Age</u>	<u>Males</u>				<u>Females</u>			
	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
6 mos.	39	25	30	27	31	30	24	32
1-1/2	41	41	25	32	11	24	21	21
2-1/2	14	15	17	9	16	11	17	12
3-1/2	8	15	12	19	11	15	13	9
4-1/2	4	6	7	5	4	9	8	5
5-1/2	3	1	3	1	8	6	2	5
6-1/2	0	3	0	0	2	1	1	2
7-1/2	1	2	0	1	2	2	1	0
8 to 9-1/2	1	0	0	0	3	1	1	0
10-1/2	0	0	0	0	0	0	0	1
Totals	<u>111</u>	<u>108</u>	<u>94</u>	<u>94</u>	<u>88</u>	<u>99</u>	<u>88</u>	<u>87</u>

Table 15. Age composition of Nantucket, Massachusetts deer checked at biological stations from 1971 through 1974.

<u>Age</u>	<u>Males</u>				<u>Females</u>			
	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>	<u>1971</u>	<u>1972</u>	<u>1973</u>	<u>1974</u>
6 mos.	27	17	22	25	14	21	21	20
1-1/2	38	19	28	31	22	16	17	21
2-1/2	13	12	19	15	12	12	8	10
3-1/2	7	12	11	12	13	5	7	1
4-1/2	4	0	4	2	2	5	9	6
5-1/2	0	3	2	1	3	0	2	2
6-1/2	0	0	1	0	0	1	0	1
7-1/2	1	0	0	0	0	1	1	0
8 to 9-1/2	0	0	0	0	0	0	0	0
10-1/2	0	0	0	0	0	0	0	0
Total	<u>90</u>	<u>63</u>	<u>86</u>	<u>86</u>	<u>66</u>	<u>61</u>	<u>65</u>	<u>61</u>

Table 16. A summary of the Massachusetts shotgun deer harvest, 1967 through 1974.

<u>Year</u>	<u>All Males</u>	<u>Adult Males</u>	<u>Button Bucks</u>	<u>All Females</u>	<u>Adult Females</u>	<u>Fawn Females</u>	<u>Total Deer</u>	<u>Total Permits Issued</u>	<u>Button Bucks and Females</u>	<u>% of Total Deer</u>
1967	937	886	53	235	168	67	1,172	2,430	288	25
1968	1,083	1,022	61	310	217	93	1,393	2,331	371	27
1969	1,424	1,299	125	586	415	171	2,010	5,295	711	35
1970	1,598	1,378	220	769	546	223	2,367	7,347	989	42
1971	1,359	1,106	253	889	643	246	2,248	7,270	1,142	51
1972	1,455	1,256	199	760	530	230	2,215	5,326	959	43
1973	1,421	1,253	163	616	444	172	2,037	5,386	779	38
1974	1,873	1,665	213	791	563	228	2,669	5,358	1,004	38

Table 17. A summary of percentage change in adult harvest and calculated minimal populations of deer in Massachusetts, 1967 through 1974.

	<u>1967</u>	<u>Percent Change 1967-1968</u>	<u>1968</u>	<u>Percent Change 1968-1969</u>	<u>1969</u>	<u>Percent Change 1969-1970</u>	<u>1970</u>	<u>Percent Change 1970-1971</u>	<u>1971</u>
Adult male harvest	886	+ 15.35	1,022	+ 27.10	1,299	+ 6.08	1,378	- 19.73	1,106
Calculated adult male population	2,056	+ 17.05	2,410	+ 31.95	3,180	- 00.82	3,154	- 27.68	2,281
Calculated adult female population	2,018	+ 31.92	2,964	+ 35.19	4,007	+ 32.24	5,299	- 22.95	4,083
Calculated minimal populations	6,139	+ 42.37	8,740	+35.01	11,805	+ 27.18	15,014	- 22.93	11,571
	<u>1971</u>	<u>Percent Change 1971-1972</u>	<u>1972</u>	<u>Percent Change 1972-1973</u>	<u>1973</u>	<u>Percent Change 1973-1974</u>	<u>1974</u>		
Adult male harvest	1,106	+ 13.56	1,256	+ 00.02	1,258	+ 32.35	1,665		
Calculated adult male population	2,281	+ 20.78	2,755	+ 11.80	3,080	+ 00.86	3,107		
Calculated adult female population	4,083	- 3.67	3,933	- 02.11	3,850	+ 21.03	4,660		
Calculated minimal population	11,571	- 2.03	11,336	+ 00.83	11,431	+ 13.59	12,934		

Table 18. Summary of the adult male deer harvest per square mile of deer range per county in Massachusetts, 1967 through 1974.

County	Sq. Mi. Deer Range	1967	1968	1969	1970	1971	1972	1973	1974
Barnstable	290.5	.14	.15	.20	.28	.28	.26	.27	.29
Berkshire	839.0	.27	.37	.44	.44	.42	.45	.51	.82
Bristol	422.6	.01	.01	.01	.01	.002	.007	.002	.009
Essex	344.4	.03	.03	.04	.06	.03	.05	.05	.04
Franklin	649.1	.27	.35	.39	.39	.36	.43	.33	.49
Hampden	524.3	.13	.19	.24	.22	.20	.25	.28	.31
Hampshire	431.5	.16	.25	.27	.26	.15	.23	.23	.27
Middlesex	583.1	.02	.01	.02	.02	.02	.01	.01	.003
Norfolk	277.3	---	---	.01	---	---	---	---	---
Plymouth	544.2	.03	.02	.03	.02	.01	.02	.01	.01
Worcester	1,307.6	.11	.06	.14	.15	.07	.09	.09	.10
	6,213.6	.12	.14	.18	.19	.15	.18	.18	.25
Dukes	86.5	.71	.65	.72	.91	1.01	1.18	1.00	.84
Nantucket	35.9	1.34	1.56	1.89	2.42	1.78	1.36	1.89	1.73

Table 19. Summary of the adult female deer harvest per square mile of deer range per county in Massachusetts, 1967 through 1974.

County	Sq. Mi. Deer Range	1967	1968	1969	1970	1971	1972	1973	1974
Barnstable	290.5	.03	.02	.06	.10	.09	.09	.05	.11
Berkshire	839.0	.04	.06	.12	.16	.23	.13	.13	.26
Bristol	422.6	.005	.002	.002	---	---	.002	.002	.002
Essex	344.4	.006	.003	.01	.03	.003	.003	.00	.009
Franklin	649.1	.07	.11	.14	.15	.27	.22	.12	.12
Hampden	524.1	.02	.03	.06	.06	.08	.07	.07	.07
Hampshire	431.5	.04	.06	.11	.11	.07	.08	.05	.07
Middlesex	583.1	.002	.01	.01	.01	.01	.01	.002	.002
Norfolk	277.3	.004	---	---	.004	---	---	---	---
Plymouth	544.2	.002	.01	.01	.01	.002	---	.004	.004
Worcester	1,307.6	.02	.02	.05	.07	.05	.03	.04	.04
	6,213.6	.02	.03	.06	.07	.08	.07	.05	.07
Dukes	86.5	.14	.10	.18	.36	.67	.89	1.00	.69
Nantucket	35.9	.17	.14	.97	1.67	1.53	1.11	1.22	1.23

Table 20. A summary of the total harvest of deer in Massachusetts (including shotgun, archery, and muzzle load harvests) per county per sex and the harvest of deer per square mile of deer range in Massachusetts for 1974.

County	Males	Females	Total	Sq. Mi. of Deer Range	Harvest		
					Total Deer per Sq. Mi.	Males per Sq. Mi.	Females per Sq. Mi.
Barnstable	94	42	136	290.5	.47	.32	.14
Berkshire	781	309	1,090	839.0	1.30	.93	.37
Bristol	4	1	5	422.6	.01	.01	.002
Essex	13	4	17	344.4	.05	.04	.01
Franklin	366	108	474	649.1	.73	.56	.17
Hampden	185	53	238	524.3	.45	.35	.10
Hampshire	134	49	183	431.5	.42	.31	.11
Middlesex	4	2	6	583.1	.01	.01	.003
Norfolk	---	---	---	277.3	---	---	---
Plymouth	8	3	11	544.2	.02	.01	.01
Worcester	151	70	221	1,307.6	.17	.12	.05
Total	1,740	641	2,381	6,213.6	.38	.28	.10
Dukes	110	121	231	86.5	2.67	1.27	1.40
Nantucket	98	69	167	35.9	4.65	2.73	1.92
Total	1,948	831	2,779	6,336.0	.44	.31	.13

Table 21. Percent frequency ratio of adult females to adult males, 1967 through 1974.

County	1967			1968			1969			1970		
	Adult		% Freq.	Adult		% Freq.	Adult		% Freq.	Adult		% Freq.
	M.	F.		M.	F.		M.	F.		M.	F.	
Barnstable	41	10	.24	44	6	.14	58	16	.28	80	29	.36
Berkshire	223	31	.14	314	53	.17	371	100	.27	366	138	.38
Bristol	6	2	.33	5	1	.20	3	1	.33	4		
Essex	11	2	.18	10	1	.10	13	4	.31	20	9	.45
Franklin	176	44	.25	230	68	.30	256	93	.35	251	98	.39
Hampden	69	9	.08	98	13	.13	123	32	.26	117	31	.26
Hampshire	69	16	.23	107	27	.25	115	47	.41	114	48	.42
Middlesex	10	1	.10	7	3	.43	13	4	.31	9	6	.67
Norfolk		1			3		3				1	
Plymouth	16	1	.06	8	3	.38	14	6	.43	8	4	.50
Suffolk												
Worcester	140	32	.23	75	28	.37	176	59	.34	192	88	.46
Total	766	149	.19	893	203	.23	1,145	362	.32	1,161	452	.39
No. of Sportsman Permits	2,000			2,000			4,000			6,000		
Dukes	61	12	.20	56	9	.16	62	16	.26	79	31	.39
No. of Sportsman Permits												
Nantucket	48	6	.13	56	5	.09	68	35	.51	87	60	.69
No. of Sportsman Permits							400			400		
Total Number of Permits*	2,243			2,331			4,695			6,747		

* Includes Farmer-Landowner Permits

Table 21. (Continued)

County	1971				1972				1973				1974			
	Adult		% Freq.		Adult		% Freq.		Adult		% Freq.		Adult		% Freq.	
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.
Barnstable	82	25	75	27	77	15	85	31	77	15	85	31	85	31	85	31
Berkshire	352	187	373	112	424	109	689	216	424	109	689	216	689	216	689	216
Bristol	1		3	1									4	1		
Essex	12	1	16	1	16	4	13	3	16	4	13	3	13	3	13	3
Franklin	232	176	277	141	213	79	316	76	213	79	316	76	316	76	316	76
Hampden	103	42	131	.38	145	36	160	.23	145	36	160	.23	160	37	160	.23
Hampshire	63	31	98	34	101	22	116	.22	101	22	116	.22	116	32	116	.28
Middlesex	14	7	3	4	4	1	2	.50	4	1	2	.50	2	1	2	.50
Norfolk																
Plymouth	7	1	12	0	5	2	6	.33	5	2	6	.33	6	2	6	.33
Suffolk																
Worcester	89	60	117	44	119	51	133	.39	119	51	133	.39	133	52	133	.39
Total	955	530	1,105	402	1,104	316	1,524	.30	1,104	316	1,524	.30	1,524	451	1,524	.30
No. of Sportsman Permits	6,000		4,000		4,000		4,000		4,000		4,000		4,000		4,000	
Dukes	87	58	102	77	86	87	73	.82	86	87	73	.82	73	60	73	.82
No. of Sportsman Permits	600		600		600		600		600		600		600		600	
Nantucket	64	55	49	40	68	44	64	.69	68	44	64	.69	64	44	64	.69
No. of Sportsman Permits	400		400		400		400		400		400		400		400	
Total Number of Permits*	7,270		5,326		5,349		5,358		5,358		5,358		5,358		5,358	

* Includes Farmer-Landowner Permits

Table 22. A summary of the shotgun deer harvest per deer management unit, the number of sportsman antlerless permits per unit, the square miles of deer range, the adult harvest per unit and the percent frequency ratio of adult females to adult males for 1974.

Unit	No. of Sportsman Permits	Males		Females*		Total Deer Harvest	Sq. Mi. of Deer Range	Adult Male/ Sq. Mi.	Adult Female/ Sq. Mi.	Percent Frequency Adult Female to Adult Male
		Adult	Fawn	Adult	Fawns*					
Barnstable	200	85	6	31	11	133	290.5	.29	.11	.36
Berkshire	1300	689	79	216	72	1,056	839.0	.82	.26	.31
Franklin	700	316	31	76	25	448	649.1	.49	.12	.24
Hampden	400	160	21	37	13	231	524.3	.31	.07	.23
Hampshire	300	116	9	32	11	168	431.5	.27	.07	.28
Worcester	700	133	11	52	17	213	1,307.6	.09	.04	.39
Region I**	200	15		4	1	20	1,204.8	.01	.003	.26
Region II***	200	10	2	3	1	16	966.8	.01	.003	.30
Dukes	600	73	29	60	35	197	86.5	.84	.69	.82
Nantucket	400	62	26	44	22	154	35.9	1.73	1.23	.71

* Female fawns were calculated using the percent of fawns reported at biological deer checking stations.

** Region I includes Essex, Middlesex and Norfolk Counties.

*** Region II includes Bristol and Plymouth Counties.

JOB PERFORMANCE REPORT

L 11.1

State MASSACHUSETTS

Cooperator: Massachusetts Division of Fisheries and Game

Project No.: W-35-R-17 Project Title: Game Population Trend and Harvest Survey

Job No.: III-1 Job Title: Mourning Dove Census

Period Covered: 21 May to 1 July 1974

Summary: Calling doves were counted on three randomized routes in cooperation with the U. S. Fish and Wildlife Service's annual mourning dove breeding population census. The total number of calling doves decreased nearly two-thirds as compared to 1973 counts. One route was relocated due to an impassable road.

Objectives: To obtain an index of the spring breeding population of mourning doves.

Procedures: In accordance with instructions from the U. S. Fish and Wildlife Service, doves were censused on two of the three randomized routes established in 1967. The third route, 8A (0081) was permanently relocated since part of the old route became inaccessible due to washouts. Doves were censused by roadside coo counts on these routes between 21 and 30 May. Division personnel conducted two counts and a Fish and Wildlife Service cooperator conducted the third.

Findings: Results of the 1974 call count of mourning doves are compared with previous years of data in Table I.

The total number of calling doves for all routes combined decreased 62 percent from 1973 results. Taken individually, Route 8 decreased by 100 percent, and Route 8A and 10 remained unchanged.

The weighted mean number of doves heard per comparable Massachusetts route was 6.1 in 1973 and 4.8 in 1974 (Ruos, 1974). Ruos further reports that the breeding population index for the Eastern Management Unit decreased 3.1 percent from 16.2 doves per route in 1973 to 15.7 doves per route in 1974. Long-term trends also show a decline though regression analysis indicated a slight upward trend in the combined non-hunting states.

CHICAGO, ILL.

 

CHICAGO, ILL.

Table 1. Doves Heard on Comparable Routes, 1966-1974.

Route	1966	1967	1968	1969	1970	1971	1972	1973	1974	1966	1967	1968	1969	1970	1971	1972	1973
8	22	26	11	2	11	10	12	10	0	-100	-100	-100	-100	-100	-100	-100	-100
8A**	*	1	1	1	2	1	2	2	2	*	+100	+100	+100	0	+100	0	0
10	3	7	0	1	3	1	2	4	4	+33	-43	+400	+300	+33	+300	+100	0
Totals	25	34	12	4	16	12	16	16	6	-76	-82	-50	+50	-62	-50	-62	-62
Mean	8.33	11.33	4.00	1.33	5.33	4.00	5.33	5.33	2.00								

* Route not conducted.

** Relocated 1974.

Recommendations: It is recommended that the spring mourning dove census be continued in cooperation with the Bureau of Sport Fisheries and Wildlife.

MASSACHUSETTS DIVISION OF FISHERIES AND GAME
Bureau of Wildlife Research and Management

Approved: _____
Colton H. Bridges, Superintendent

Prepared by _____
James E. Cardoza, Game Biologist

Date _____

Literature Cited: Ruos, James L. 1974. 1974 Mourning Dove Breeding Population Status. USDI Bur. Sport Fish. & Wildl., Laurel, Md. Administrative Report. Xerox 1 + 35 pp.

THE UNIVERSITY OF CHICAGO
 
WWW.LIBRARY.CHICAGO.EDU

PERFORMANCE REPORT

U OF MASSACHUSETTS LIBRARY

State Massachusetts
Project No.: W-35-R-17
Project Title: Game Population Trend and Harvest Survey
Project Type: Research and Surveys
Period Covered: 1 June 1974 to 31 May 1975

Work Plan IV Wild Turkey Restoration Study

Work Plan Objectives: To re-establish the wild turkey in the Commonwealth in sufficient numbers to allow for recreational hunting.

Job IV-1 Experimental Turkey Stocking

Job Objectives: To re-establish the wild turkey in the Commonwealth in sufficient numbers to allow for recreational hunting.

Summary: Turkeys in Beartown State Forest now show signs of dispersal from the release area. Flocks were located both east and west of Beartown as well as on the forest itself. Reports of turkeys to the north and south need verification.

Reports from cooperators and investigations by project personnel indicate that released birds produced young during the past summer.

Target Date: 31 May 1979

Status of Progress: On schedule.

Significant Deviations: None

Recommendations: Continue evaluation of the Beartown release. Investigate other release sites in western Massachusetts and release trapped birds from the Beartown area on these sites once populations have grown enough to permit this.

Cost: \$4,065.25 (Project leader 41-1/2 man days)

Remarks: Procedures: Turkeys were censused by roadside counts, track counts, and cooperator reports. Snowmobiles were used during the winter to provide access to the areas and to transport grain for baiting.

 

Finding: Beartown State Forest Area

A total of 37 wild-trapped turkeys (6 adult males, 9 adult females, 9 immature males, 7 immature females, 6 immature unknown) from Allegany State Park, Cattaraugus County, New York, were released in Beartown State Forest, Great Barrington, Berkshire County, between March 1972 and September 1973.

Few sightings were recorded during the first two years of the release, with the only substantial flock found consisting of 13 of the 20 birds released in fall 1973 which were located in early winter 1974 2.0 km (1.25 miles) north of the release site. During the current segment, however, sightings have increased, with turkeys being found at several locations on and about Beartown forest.

In June 1974, a single adult bird with a red wing tag was seen on the road to the Mt. Wilcox fire tower, about 4.8 km (3 miles) southeast of the release site. In late August, Natural Resource Office Gordon Leeman saw a wing-tagged (orange) hen and one poult crossing a road near the center of the forest. Three of the stocked birds released March 1972 had red or red-patterned patagial tags, while three additional turkeys (two released March 1972 and one September 1973) had orange tags.

In late fall 1974, the Division issued news releases soliciting reports of turkeys. This publicity, coupled with a snowfall during the deer hunting season, resulted in several sightings of turkeys and turkey tracks which were subsequently reported to the Division. One flock of five turkeys was seen repeatedly by sportsmen on the southwest slope between Stony Brook and Benedict Pond. Project personnel, however, found tracks of six to seven turkeys in this location, while forest workers saw about ten near an abandoned cemetery on this slope, and again roosting just north of Blue Hill Road near the crossing of the Appalachian Trail. Project personnel also tracked seven birds through the beaver swamps between Beartown Mountain and Mt. Wilcox. Two roosts were located, but the turkeys were seen on only one occasion. Tracks of three toms were found off Monterey Road in the forest in early spring. Apparently, there are at least ten to twenty turkeys on the forest itself, predominantly in the area between Benedict Pond and Mt. Wilcox.

Deer hunters also reported thirteen to fifteen turkeys in December near Fountain Pond 2.7 km (1.7 miles) southwest of the Beartown release site. Western District personnel found tracks here soon after the report, but the project leader and assistant found no sign in January. We did locate tracks of about eleven turkeys, apparently hens and young, 2.1 km (1.3 miles) further along Three Mile Hill which borders Fountain Pond. In

late February, tracks of two to three toms were located on the same ridge, 1.1 km (0.7 miles) north of the pond.

Turkeys were also reported in December northeast of the forest, off Jerusalem Road in Lee. We found no sign here in winter, but did receive another report of "a number" of birds across the Tyringham Valley, on the western slope of Cannon Hill on the Lee-Tyringham line, 9.0 km (5.6 miles) northeast of the release site. After several unproductive searches, we were able to verify that turkeys were present, but the snow had melted too much to be able to get an accurate track count.

A report of four to five turkeys near Taylor Brook in Tolland, about 34.3 km (21.3 miles) southeast of the Beartown release site, may not be valid. While the original account of the sighting seemed convincing, discussions with one of the reputed observers indicate probable confusion of turkeys with pheasants. More definite observations are needed before including this area in occupied turkey range.

A single adult turkey was seen in December 1974 near the U. S. Fish and Wildlife Service fish hatchery in Hartsville. An unmarked hen found dead in the road near the center of Monterey in early June 1975 may have been this bird.

In May 1975, an adult male turkey with an orange wing tag was seen crossing Route 7 in Lanesboro, 28.6 km (17.8 miles) northwest of Beartown. Three unmarked hens were also seen twice off Balance Rock Road in the same town. Only one male turkey with an orange wing tag was released in Beartown so either that one bird moved north, or else the bird or its tag was misidentified.

Findings: Statewide Populations

Turkeys in the Quabbin Reservation have continued to improve in wildness since the discontinuance of artificial feeding in 1971, though they still do not approach wild-trapped birds in overall viability and wariness. Division personnel observed a flock of 17 birds (9 hens and 8 toms) near Prescott Brook, with a smaller flock of six birds off Thurston Brook on the tip of the peninsula; and another group of four just inside Reservation boundaries in south New Salem. Metropolitan District Commission and University of Massachusetts cooperators also reported four birds near Egypt Brook on Prescott Peninsula, and several birds on the east side of the Quabbin in Hardwick. A flock of 13 turkeys was reported on the Swift River Wildlife Management Area in Belchertown, south of the Quabbin, early in October 1974, but District personnel were unable to find any turkey sign at that location.

A hen was accidentally flushed from a nest of 13 eggs at the northern end of Prescott Peninsula on 10 May. The nest was abandoned and the eggs collected. A cooperator attempted unsuccessfully to hatch these in an incubator. Division personnel and University co-operators observed two broods during June, one of about nine poults 0.3 km (0.2 miles) north of Prescott four-corners, and one of about seven poults near Mt. Pleasant.

Few birds remain in the Barre and Douglas State Forest areas. One unbanded bird was killed by a car near the Barre Falls Dam in June 1974, and one bird was seen that fall in Douglas near the Rhode Island line.

Flocks of game-farm ancestry turkeys in the Berkshires have declined considerably. Only two to three birds were reported in the Town of Mt. Washington, and no birds were reported from Middlefield. Forest personnel did report two birds--one unmarked and one with a red wing tag--from October Mountain State Forest. Since the last semi-wild bird marked with a red tag anywhere in this area was an adult hen released in Middlefield in spring 1968, I suspect that either marked birds from Beartown moved into this area or else the tag color was not correctly recorded.

Single turkeys were seen in Sunderland, Whately, and Peru. Two or three birds seen in Monson are probably survivors of an illegal release by private individuals.

The project leader gave one slide show and taped one television commentary on the turkey project.

Acknowledgment: I extend my appreciation to personnel of the Beartown State Forest, the Division of Law Enforcement, the Metropolitan District Commission, and the Massachusetts Cooperative Wildlife Research Unit for their cooperation and assistance in reporting turkey sightings.

* * * * *

Job IV-2 Evaluation of Habitat Utilized by Transplanted Wild Turkeys

Job Objectives: Habitat utilized by transplanted wild turkeys will be evaluated in relation to total habitat available, with special reference being given to habitats selected for winter roosting, winter feeding, nesting, and brood rearing.

This job was inactive during this segment.

MASSACHUSETTS DIVISION OF FISHERIES AND WILDLIFE
Bureau of Wildlife Research and Management

Prepared by _____
James E. Cardoza
Game Biologist

Approved by _____
Colton H. Bridges
Deputy Director

Date _____

PERFORMANCE REPORT

State Massachusetts
 Project No.: W-35-R-17
 Project Title: Game Population Trend and Harvest Survey
 Project Type: Research and Survey
 Period Covered: 1 June 1974 to 31 May 1975

 Work Plan VI Black Bear Study

Work Plan Objective: To define the range of the black bear in Massachusetts and to determine its population characteristics and rate of harvest by hunting.

Job VI-1 Black Bear Population Dynamics

Job Objectives: To define the range of the black bear in Massachusetts and to determine its population characteristics and rate of harvest by hunting.

Summary: Applications for bear hunting permits were received from 390 sportsmen. Two bear were taken during the open season, and one was killed by an automobile. New reports of 29 observations totalling 38 bear were received from five counties. Two instances of problem bears were investigated.

Target Date: 31 May 1979.

Status of Progress: On schedule.

Significant Deviations: None

Recommendations: Continue evaluation of bear harvest through checking stations and periodic questionnaires.

Continue determination of bear range and populations through recording of sightings.

Investigate nuisance complaints as necessary. Promote public education programs to lessen man-bear conflicts.

Cost: \$555.46 (project leader man days - 7-1/2)

Remarks: Procedures:

Current bear hunting regulations include mandatory reporting and tagging of bear. Bear checking stations were maintained daily during bear week at three locations: Birch Hill Wildlife Management Area, Baldwinville; Bitzer State Fish Hatchery, Montague; Western Wildlife District Headquarters, Pittsfield.

When bear were presented for examination, station personnel affixed a metal game seal to the bear, removed a premolar tooth, and recorded the following information: town of kill, date killed, sex and weight of bear, and method of kill.

The Information-Education Section issued periodic news releases asking for reports of bear. The project leader gave one slide talk on the bear study. Several sightings were reported by cooperators from the University of Massachusetts and the Division of Law Enforcement.

Findings:

Bear hunting permit applications were received from 390 individuals during the 1974 season (Table 1), with two hunters succeeding in taking a bear. One 37.2 kg (192 lb.) sow was shot in Middlefield, Hampshire County, on the first day of the season (18 November) by a hunter using dogs. The second, a 123.5 kg (272 lb.) boar, was taken without dogs in Savoy, Berkshire County, on 22 November, the fifth day of the season. These are the second and third bears legally taken since the season was shortened in 1970.

One cub, sex undetermined, was killed by a car on Route 2 in North Adams on 5 September. The bear was retained to be mounted whole as a display specimen. A cooperator reported finding a dead bear near Thunder Mountain in Ashfield in December and sportsmen reported a second dead bear near the Middlefield-Washington town line during deer week. Division personnel investigated, but could not confirm either of these reports.

Table 1. Number of bear permit applications and number of bear taken in Massachusetts from 1970-1974.

<u>Year</u>	<u>No. Permits</u>	<u>No. Bear Taken</u>	<u>Other Mortalities</u>
1970	214	-	1 illegal kill; 1 road kill
1971	200	-	
1972	423	1	1 illegal kill
1973	309	-	1 road kill; 1 captured bear
1974	390	2	1 road kill

New reports of bear received during this segment included 21 sightings, five reports of track or sign, two hunter kills, and one road kill, totalling 38 bear seen in 19 towns. Reports by county for the period 1952 to 1975 are presented in Table 2. Records of 512 bear noted on 415 occasions in 77 towns have been collected for the period 1952 to May 1975.

THE UNIVERSITY OF CHICAGO
 
CHICAGO, ILL. 60637

Table 2. Reports of black bear by county for Massachusetts, 1952 to May 1975.

<u>County</u>	<u>1952-May 1974</u>	<u>June 1974-May 1975</u>	<u>Total</u>	<u>Percent</u>
Berkshire	129	9	138	33.4
Franklin	154	6	160	38.8
Hampden	19	2	21	5.1
Hampshire	64	8	72	17.4
Middlesex	1	0	1	0.2
Worcester	17	4	21	5.1
	<u>384</u>	<u>29</u>	<u>413</u>	<u>100.0</u>

Four family groups were observed, two with one cub each, one with two cubs, and one with three cubs.

Two problem situations were investigated. The first, in Westminster, Worcester County, in late July 1974, involved a bear which was frequenting woodland adjacent to a local golf course. The owner, having located several spots where the bear had dug into logs and stone walls, was concerned that the bear would dig up the golf greens. Also, since bear are rarely seen in this area, the townspeople were fearful of it and were keeping their children inside. Division personnel contacted the local police and the golf course owner, examined the area where the diggings were found, and explained that the bear would not dig up the greens or attack the children. The bear and bear sign was seen a few more times in subsequent weeks but no damage occurred, and the bear apparently left the area that fall.

The second incident involved bear depredation on several hives in Middlefield in late October. Since the bear hunting season opened a few weeks later, Division personnel did not attempt to control the animal. The bear remained in the area and was shot by a hunter on the first day of the season.

Acknowledgments: I appreciate the efforts of the Division of Law Enforcement in investigating complaints and sightings.

* * * * *

Job VI-2 Historical Records of the Black Bear in Massachusetts

Job Objectives: To determine trends in black bear populations and distribution in Massachusetts from pre-Colonial times to the present.

Summary Trends in black bear populations from pre-Colonial times to the present were determined by an intensive literature search of published and unpublished records. Maps have been prepared showing fluctuations in bear distribution.

Instances of nuisance bear situations have been compiled and described and possible solutions discussed.

The historical aspects of bear management have been examined and the present hunting regulations for the Northeast tabulated. The effect of various management programs on the bear's status is discussed and recommendations for Massachusetts are presented.

Target Date: 31 May 1976.

Status of Progress: On schedule.

Significant Deviations: None

Recommendations: Publish findings of this job, and accrued data from Job VI-1, as a Division research bulletin.

Cost: \$3,988.28 (project leader man days, 93.5)

Remarks: The first draft of a comprehensive report on the history and status of the bear in Massachusetts and adjacent states has nearly been completed. The outline for this report is as follows:

I. Introduction

II. Procedures

III. Regional background

A. Topography and vegetation

B. Indian population and hunting methods

C. Discovery and settlement

D. Civil history

E. Land use and general wildlife situation

IV. History and Status of the Black Bear

A. Pre-Colonial

B. Massachusetts

1. 1620-1750

2. 1750-1850

3. 1850-1950

4. Present (1950-date)

C. Connecticut

D. Rhode Island

E. New Hampshire

F. Vermont

V. Bears and Man

VI. Bear Management

VII. Summary and Conclusions

Prior to European settlement, bears were well distributed throughout New England with the possible exception of parts of southeastern Massachusetts and Rhode Island. Tribal hunting for food and clothing may have had some effect on those peripheral areas, where bears were scarce, but Indian numbers away from the coast and river valleys were low and the overall bear population was probably not threatened by native hunting.

After colonization, though, human populations climbed rapidly and bears were severely persecuted because of their depredations on crops and flocks and their reputed danger to man. The clearing of the land for farms and settlements restricted the bears' available habitat and brought them into closer contact with man who persisted in killing the shaggy varmints which dared to infringe on human territory. This widespread eradication of the forests, coupled with the unrestricted year-round shooting of bears resulted in a serious slump in populations. After 1800, bears were uncommon in southern New England, and by 1870-1880 were at a low point regionwide, being restricted to the more remote mountainous sections of Vermont, New Hampshire and extreme northwestern Massachusetts.

Persecution slackened as bear numbers dropped and agriculture declined, and after 1900 remnant populations began expanding as Bruin straggled back into the woodlands springing up on long abandoned farmsteads. After World War II, with bear numbers apparently increasing, its popularity grew and its status gradually shifted from despised vermin to favored game animal, with commensurate legislative protection and research and management programs being initiated. Bears are now found at least as stragglers, in all New Hampshire and mainland Vermont counties, and the western half of Massachusetts and adjacent areas of Connecticut.

Bear problems may arise, not so much from unencouraged attacks on man or depredations on crops or livestock, but from close contact situations where careless or heedless persons feed or harass wild bears, with subsequent potential for injury or property damage. Management programs should incorporate public education efforts aimed at decreasing this adverse conflict. Response to "nuisance" or depredation situations should be flexible, and solutions short of destroying the bear be employed wherever possible.

The first of these is the question of the origin of the human race. It is generally admitted that the human race is descended from a common ancestor, but the exact nature of this ancestor and the circumstances of his life are matters of dispute. Some authorities believe that the human race is descended from a single pair of individuals, while others believe that it is descended from a large number of individuals who lived at different times and in different places.

The second question is the question of the development of the human race. It is generally admitted that the human race has developed from a lower state to a higher state, but the exact nature of this development and the circumstances of it are matters of dispute. Some authorities believe that the human race has developed from a single pair of individuals, while others believe that it is descended from a large number of individuals who lived at different times and in different places.

The third question is the question of the distribution of the human race. It is generally admitted that the human race is distributed over the whole of the world, but the exact nature of this distribution and the circumstances of it are matters of dispute. Some authorities believe that the human race is descended from a single pair of individuals, while others believe that it is descended from a large number of individuals who lived at different times and in different places.

The fourth question is the question of the future of the human race. It is generally admitted that the human race will continue to exist, but the exact nature of its future and the circumstances of it are matters of dispute. Some authorities believe that the human race will continue to exist in its present state, while others believe that it will develop into a higher state.

Bears in the Northeast are maintaining populations at huntable levels, though seasonal changes may be necessary in some instances. In Massachusetts, bear populations are minimal and seasonal limits should be regulated so that the annual harvest does not greatly exceed present levels. Use of checking stations, hunter report forms, and distributional surveys should be employed as necessary to monitor and assess bear harvests and distribution.

MASSACHUSETTS DIVISION OF FISHERIES AND WILDLIFE
Bureau of Wildlife Research and Management

Approved: _____

Colton H. Bridges
Deputy Director

Prepared by _____

James E. Cardoza
Game Biologist

Date _____

 

ACME
BOOKBINDING CO., INC.

AUG 23 1984

100 CAMBRIDGE STREET
CHARLESTOWN, MASS.