Agriculture
Canada

Research
Branch

Direction générale
de la recherche

Technical Bulletin 1985-4E

Flora of the Manyberries Research Substation

630.72
C759
C 85-4
00Ag
c.3

Canada

The map on the cover has dots representing
Agriculture Canada research establishments.

Flora of the Manyberries Research Substation

S. SMOLIAK
Research Station
Lethbridge, Alberta

Lethbridge Research Station Contribution No. 6

Research Branch
Agriculture Canada
1985

Copies of this publication are available from:
Mr. S. Smoliak
Plant Science Section
Research Station
Research Branch, Agriculture Canada
Lethbridge, Alberta
T1J 4B1

Produced by Research Program Service

© Minister of Supply and Services Canada 1985
Cat. No. A54-8/1985-4E
ISBN 0-662-14227-6

CONTENTS

Summary/Résumé	i
Introduction	1
Plant Communities	4
Annotated List of the Vascular Plants	5
Bibliography	31

SUMMARY

Native and introduced plants have been collected at the Manyberries (Alberta) Research Substation since its inception in 1926. The plant collection, housed in the herbarium of the Lethbridge Research Station, represents the flora of the Mixed Prairie Association of the Northern Great Plains. Included in the 366 taxa are some rare vascular plants not found elsewhere on the prairies and some that are unique to the area. Several plant communities, characterized by one or more dominant species, have been identified. The plants collected are representative of 55 families but the bulk of the species are contained within a comparatively small number of families. The three largest families are Compositae, Gramineae and Leguminosae with 67, 62, and 29 species, respectively.

RESUME

La collection de plantes indigènes et introduites, à la Sous-station de recherches de Manyberries (Alberta), a remonté à l'ouverture de cette dernière en 1926. Elle constitue aujourd'hui l'herbier de la Station de recherches de Lethbridge et représente la flore de l'association mixte des Grandes plaines septentrionales des Prairies. Parmi les 366 taxons représentés, on trouve quelques plantes vasculaires qui ne poussent nulle part ailleurs dans les Prairies et d'autres qui sont endémiques dans la région. On a déterminé plusieurs groupements végétaux caractérisés par au moins une espèce dominante. Bien que 55 familles soient représentées, la plupart des espèces appartiennent à un nombre restreint de familles dont les trois plus importantes, les Composées, les Graminées et les Légumineuses, comprennent respectivement 67, 62 et 29 espèces.

Digitized by the Internet Archive
in 2013

<http://archive.org/details/floraofmanyberri19854smol>

INTRODUCTION

The Manyberries Research Substation is located at Onefour, Alberta. The southern portion adjoins the Canada - U.S.A. border between 49° and 49° 10' North and 110° 22' and 110° 32' West. The substation was established in 1927 on 6,300 ha (15,590 acres) of rangeland and was then called the Dominion Range Experiment Station, Manyberries, the first of its kind in Canada. Grazing and vegetation studies were initiated to improve the carrying capacity of the range. The station was increased to about 17,000 ha (42,000 acres) when the southern portion of the experimental range was acquired in 1951-1952. Some of this area was cultivated by settlers who arrived in 1910 and left in 1952.

The southwest corner of the Research Substation (Fig. 1) is traversed by the Lost River coulee which drains into the Milk River just south of the International Boundary. The northern portion of the substation is drained by several coulees. The coulees contain eroded slopes and badlands. All drainage is southward and joins the Missouri River System.

The geological formation that occurs at the surface or immediately below is Upper Cretaceous in age and is the Oldman formation. This formation consists chiefly of soft light-colored shales and sandstones, but clays or clayey shales predominate. The formation outcrops in the area usually as extensive badlands along both sides of the Lost River coulee (Fig. 2) and other eroded drainage systems. Near the top of the formation there are thin coal seams and coaly shales, as well as the occasional plant remains and invertebrate and vertebrate fossils.

The substation is located within the Brown Soil Zone characterized by shallow profiles, low amounts of organic matter, nitrogen and phosphorus, and a compact calcium carbonate layer averaging about 30 cm (12 inches) below the surface. The soils have developed from glacial till composed of locally derived materials deposited by continental glaciers or from underlying marine shales and sandstones where the drift deposits were thin.

The dominant soils are members of the Orthic Brown Subgroup of the Chernozemic Order, Brown Great Group and the Brown Solod Subgroup of the Solonetzic Order, Solod Great Group. The chernozemic soils are more common in the southern portion of the substation. They are loamy in texture and have formed largely from weathering of the underlying sandstone, although in some locations they have been deposited as alluvium or outwash from the higher lands to the north and reworked by wind. The solonetzic soils dominate the northern portion of the substation and vary in texture from loams to clay loams. These soils are primarily derived from the underlying shales in which clays and salts have not leached out owing to the low rainfall. They are characterized by a high sodium content, poor structure, and an impervious B horizon which forms a claypan. The layer of soil above the claypan may become powdery with drying and is very susceptible to erosion or soil "blowout" giving a pitted appearance to the landscape.

Fig. 1. Study area, about 35 km (22 miles) southeast of Manyberries, Alberta, showing the physical features and roads and fields. The Research Substation borders the U.S.A. boundary for 10 km (6 miles).

Fig. 2. Lost River coulee, with the Sweetgrass Hills in the background showing formations and terrain of the glacial drainage system.

The topography of the substation is undulating to gently rolling. The altitude varies from 915 m (3,000 ft) to 975 m (3,200 ft) and the area is surrounded by prominent landmarks. The Cypress Hills, to the north, reach a height of 1,465 m (4,800 ft) and can be seen from the northern part of the substation. The Bearpaw Mountains, to the southeast, and the Sweet Grass Hills, to the southwest, both rise to a height of 2,135 m (7,000 ft) above sea level and are in Montana.

The substation is located in one of the driest areas of the province. The climate is characterized by low precipitation, a high rate of evaporation, great extremes of temperature and abundant sunshine. Precipitation occurs chiefly in the form of rain, about 80 percent of which falls during the growing season, April to July, inclusive. Many of the rain showers are too light to be of any value to plant life and much of the heavier rains is lost by run-off. Hot, west or southwest, winds are frequent during midsummer, while in winter these warm winds or "chinooks" rarely reach across the Lost River coulee.

Long-term mean annual precipitation over a 30-year period (1951-1980) was 327 mm (12.9 inches) while mean annual temperature for the same period was 4.2°C (40.0°F). The lowest temperature of -49.0°C (-45°F) was recorded in January and the highest temperature of 40.6°C (105°F) was recorded in August. The mean annual evaporation was 1,060 mm (41.8 inches).

PLANT COMMUNITIES

The vegetation of the area is representative of the Mixed Prairie Association (Coupland 1950, 1961; Moss 1955; Looman 1980) which is characterized by an abundance of mid- and short-grasses, numerous forbs and few shrubs. Several plant communities have been identified on the substation. The communities are distinguished by one or more species commonly associated with certain soil characteristics.

The most extensive area is the upland prairie where the grassland vegetation types dominate. The *Stipa-Bouteloua-Agropyron* type is most common and occupies the light, sandy loam soils. The most abundant grasses are needle and thread (*Stipa comata*), blue grama (*Bouteloua gracilis*) and the two wheatgrasses, northern (*Agropyron dasystachyum*) and western (*A. smithii*), while June grass (*Koeleria cristata*) and Sandberg's bluegrass (*Poa sandbergii*) are also present. Common forbs are moss phlox (*Phlox hoodii*), broomweed (*Gutierrezia sarothrae*), golden aster (*Chrysopsis villosa*) and little club moss (*Selaginella densa*). Common shrubs are pasture sage (*Artemisia frigida*), winterfat (*Eurotia lanata*), salt sage (*Atriplex nuttallii*) and sagebrush (*A. cana*).

Other vegetation types on the upland prairie are the *Stipa-Bouteloua* type on the medium-textured soils, with needle and thread and blue grama the dominant grass species; the *Agropyron-Poa* type on clay soils; the *Stipa-Agropyron* type in the more sheltered areas; the *Calamovilfa-Artemisia* type on sandy soils; and the *Bouteloua-Carex* type along the dry coulee edges.

In the coulees, the common vegetation types include the *Agropyron-Muhlenbergia* types on the coulee slopes, the *Poa-Carex-Juncus* types in the more moist coulee bottoms, the *Distichlis-Sarcobatus* type along the dry coulee bottoms, and the *Calamovilfa-Stipa* type on the sandy coulee bottoms. The *Prunus-Amelanchier* type and the *Elaeagnus-Rosa* type are found on the more moist loam and sandy soils, respectively.

Dry, saline sloughs contain the *Agropyron-Hordeum* type while non-saline sloughs have the *Calamagrostis-Juncus* or the *Agrostis-Carex* types. Willows (*Salix* spp.) are common in wet places near dams and along roadsides while trees, mainly narrow-leaf cottonwood (*Populus angustifolia*) and western cottonwood (*P. deltoides*), are found occasionally in moist coulees.

Some rare and unique plants are found in the area. A natural population of soapweed (*Yucca glauca*) occurs along the exposed south-facing slopes of the Lost River coulee near the U.S.A. boundary. The flowering-quillwort (*Lilaea*

scilloides) discovered by N. A. Skoglund (Moss 1944) occurs in margins of some sloughs that retain water until mid- or late summer. Sand puffs (*Abronia micrantha*) grows only in the sand dune area (Johnson and Hallworth 1975) and has been mistaken for wild begonia (*Rumex venosus*). Although not on the substation, the Rocky Mountain juniper was found growing in a coulee (Smith and Wallis 1976) just 1 km west of the substation along the U.S.A. border.

A listing of vascular plants in the Milk River area west of the substation has been prepared (Wallis 1976). The list contains 362 species which were found in the upland, sagebrush flats, sandbars, and wooded areas.

ANNOTATED LIST OF THE VASCULAR PLANTS

In the following list, species of vascular plants are listed in the botanical order followed by Packer (1983). The genera and species, however, are listed in alphabetical order. The numbers following the listings refer to the collections made by the staff members. Except as otherwise stated, all specimens are contained in the herbarium of the Lethbridge Research Station. Critical specimens have been verified or identified by the staff of the Vascular Plant Section, Biosystematics Research Institute, Ottawa, Ontario.

The work of previous collectors has been very important in the compilation of the list. The plant collections were initiated by Dr. S. E. Clarke in 1926, and were continued by N. A. Skoglund, E. W. Tisdale, J. A. Campbell, J. B. Campbell, W. A. Hubbard and the author while at the Dominion Range Experiment Station. After transferring to the Agriculture Canada Research Station, Lethbridge, Alberta, in 1964, the author was assisted by T. D. Allen and R. G. Gschaid in continuing plant collections to make the list as complete as possible.

Nomenclature for the vascular plants follows that proposed by Packer (1983) in most instances. However, in retaining the nomenclature assigned by the Biosystematics Research Institute taxonomists, references such as Boivin (1967-1981), Hitchcock and Cronquist (1973), Looman and Best (1979) and Scoggan (1978-1979) were consulted. The above texts were also consulted for common names to represent the most commonly used or appropriate name for each species.

The list contains species from 55 families and includes some 366 specific or subspecific entities. The bulk of the species is contained within a comparatively small number of families. The three largest are Compositae (67), Gramineae (62) and Leguminosae (29).

SELAGINELLACEAE - Spike-moss family

Selaginella densa Rydb. Little club-moss. Very common, in dry grassland.
220, 446, 515, 755.

EQUISETACEAE - Horsetail family

Equisetum arvense L. Common horsetail. Occasional, in moist draws. 3384.

Equisetum laevigatum A. Br. Smooth scouring-rush. Common, in coulees and
moist alkaline areas. 116, 445, 2846, 3217, 3406.

MARSILEACEAE - Marsilea family

Marsilea vestita Hook. and Grev. Clover-fern. Rare, in dry slough bottoms.
2245, 3140.

CUPRESSACEAE - Cypress family

Juniperus communis L. var. *depressa* Pursh. Ground juniper. Rare, on
sandy hillsides. 3380.

Juniperus horizontalis Moench. Creeping juniper. Common, on coulee slopes,
sandy areas, and exposed rocky areas. 2302, 2970, 3360.

Juniperus scopulorum Sarg. Rocky Mountain juniper. Rare, in coulee one
kilometer west of Substation boundary. 5185.

TYPHACEAE - Cattail family

Typha latifolia L. Common cattail. Fairly common, in wet places and
shallow water. 3162.

JUNCAGINACEAE - Arrow-grass family

Triglochin maritima L. Seaside arrow-grass. Fairly common, in wet and
alkaline places in coulee bottoms. 233, 2847.

Triglochin palustris L. Slender arrow-grass. Occasional, in wet and
alkaline areas. 5307.

LILAEACEAE - Flowering-quillwort family

Lilaea scilloides (Poir) Haum. Flowering-quillwort. Rare, in saline
sloughs and wet places. 1326, 1565, 3091.

ALISMATACEAE - Water-plantain family

- Alisma gramineum* Gmel. Narrow-leaved water-plantain. Fairly common, in wet places. 3092, 3141.
- Alisma plantago-aquatica* L. Broad-leaved water-plantain. Common, in wet places. 104, 3163, 3427.
- Sagittaria cuneata* Sheld. Arrowhead. Fairly common, in muddy and wet areas. 2045, 2246.

GRAMINEAE - Grass family

- Agropyron albicans* Scribn. and Smith. Awned northern wheatgrass. Occasional, on upland and sandy prairie. 925, 1365, 1388, 2856, 3173.
- Agropyron cristatum* (L.) Gaertn. Crested wheatgrass. Introduced. Common, on disturbed areas and along roads. 2898, 3222.
- Agropyron dasystachyum* (Hook.) Scribn. Northern wheatgrass. Fairly common, on upland prairie. 245.
- Agropyron desertorum* (Fisch.) Schult. Desert wheatgrass. Introduced. Localized, on disturbed areas and along roads. 2854.
- Agropyron griffithsii* Scribn. and Smith. Griffith's wheatgrass. Occasional, on dry upland prairie. 2857, 3094.
- Agropyron intermedium* (Host) Beauv. Intermediate wheatgrass. Introduced. Localized on disturbed areas. 2859.
- Agropyron repens* (L.) Beauv. Couch grass. Introduced. Occasional, on disturbed areas and along roads. 3468, 5199.
- Agropyron riparium* Scribn. and Smith. Streambank wheatgrass. Occasional, on moist disturbed areas. 2742.
- Agropyron smithii* Rydb. Western wheatgrass. Common, on upland prairie, especially on low-lying areas and moist coulees. 132, 1213, 1366, 1367, 2858.
- Agropyron subsecundum* (Link) Hitchc. Awned wheatgrass. Rare, on coulee banks and moist areas. 1106, 5211.
- Agropyron trachycaulum* (Link) Malte. Slender wheatgrass. Occasional, in some sloughs and on disturbed areas. 131, 404, 1386, 1387, 2855, 3265.
- Agropyron trichophorum* (Link) Richt. Pubescent wheatgrass. Introduced. Localized, on disturbed areas. 2860.

GRAMINEAE - continued

- Agrostis scabra* Willd. Rough hairgrass. Fairly common, in moist areas and in sloughs. 237.
- Alopecurus aequalis* Sobol. Water foxtail. Occasional, in wet areas. 371, 823, 1987, 2186, 2250.
- Beckmannia syzigachne* (Steud.) Fern. Slough grass. Occasional, in wet areas. 124, 241.
- Bouteloua gracilis* (HBK.) Lag. Blue grama. Very common, on upland prairie. 212, 3370.
- Bromus inermis* Leyss. Awnless brome. Introduced. Localized, on disturbed areas. 2853, 3296.
- Bromus tectorum* L. Downy chess. Introduced. Rare, on disturbed areas. 2172, 3093.
- Calamagrostis montanensis* Scribn. Plains reed grass. Fairly common, on sandy upland prairie. 122, 1385, 3227.
- Calamovilfa longifolia* (Hook.) Scribn. Sand grass. Localized, on sandy areas on upland prairie. 123, 1397, 3364.
- Danthonia californica* Boland var. *americana* (Scribn.) Hitchc. California oat-grass. Rare, in moist coulee bottoms. 1099, 1275, 2251, 3228.
- Danthonia unispicata* (Thurb.) Munro. One-spike oat grass. Rare, in moist locations along coulee banks. 392, 2187.
- Deschampsia caespitosa* (L.) Beauv. Tufted hair grass. Common, in wet areas. 238, 1358.
- Distichlis stricta* (Torr.) Rydb. Salt grass. Common, in saline areas. 243, 1277, 1361.
- Echinochloa crusgalli* (L.) Beauv. Barnyard grass. Scarce, in moist disturbed areas. 2972.
- Elymus angustus* L. Altai wild rye. Introduced. Localized, on disturbed areas. 2861.
- Elymus canadensis* L. Canada wild rye. Occasional, on sandy areas along coulee edges. 483, 5338.
- Elymus junceus* Fisch. Russian wild rye. Introduced. Localized, on disturbed areas and along roadsides. 1369, 2699.

GRAMINEAE - continued

- Elymus macounii* Vasey. Macoun's wild rye. Occasional, in moist areas and slough edges. 5212.
- Festuca ovina* L. Sheep fescue. Introduced. Localized, on disturbed and sandy areas. 3513.
- Glyceria borealis* (Nash) Batchelder. Manna grass. Occasional, in wet places and shallow water. 528.
- Hordeum jubatum* L. Foxtail barley. Fairly common, in sloughs, roadsides, and disturbed areas. 1368, 3292.
- Hordeum jubatum* L. var. *caespitosum* (Scribn.) Hitchc. Short-awned foxtail barley. Fairly common, in low areas and saline flats. 531.
- Hordeum pusillum* Nutt. Little barley. Scarce, in open, alkaline areas. Listed by Scoggan.
- Koeleria cristata* (L.) Pers. June grass. Very common, on upland prairie. 126, 1360, 3211.
- Lolium perenne* L. Perennial ryegrass. Introduced. Scarce, on disturbed areas. 5207.
- Muhlenbergia asperifolia* (Nees and Mey.) Parodi. Scratchgrass. Occasional, in low sandy areas. 236.
- Muhlenbergia cuspidata* (Torr.) Rydb. Plains muhly. Fairly common, on dry prairie and coulees. 119.
- Muhlenbergia richardsonis* (Trin.) Rydb. Mat muhly. Fairly common, on dry prairie and hillsides. 120, 3425.
- Oryzopsis hymenoides* (R. and S.) Ricker. Indian rice grass. Common, on sandy soils and along roadsides. 105, 1384, 3210, 3445.
- Phalaris arundinacea* L. Reed canary grass. Occasional, in sloughs and wet areas. 3467.
- Poa bulbosa* L. Bulbous bluegrass. Introduced. Rare, on disturbed, wet areas. 825.
- Poa canbyi* (Scribn.) Piper. Canby bluegrass. Fairly common, on moist slopes and coulee banks. 524, 1363.
- Poa cusickii* Vasey. Early bluegrass. Occasional, on moist coulee banks. 790.
- Poa glaucifolia* Scribn. and Williams. Glaucous bluegrass. Occasional, in moist areas and coulee bottoms. 1055.

GRAMINEAE - continued

- Poa juncifolia* Scribn. Alkali bluegrass. Occasional, in saline sloughs and moist draws. 1243, 3229, 3402.
- Poa nervosa* (Hook.) Vasey. Wheeler's bluegrass. Occasional, on moist coulee banks. 1101.
- Poa nevadensis* Vasey. Nevada bluegrass. Occasional, in moist coulee draws. 2851.
- Poa palustris* L. Fowl bluegrass. Common, in coulees and wet places. 989, 1054.
- Poa pratensis* L. Kentucky bluegrass. Introduced. Occasional, in moist low areas. 727, 1362, 5310.
- Poa sandbergii* Vasey. Sandberg's bluegrass. Very common, on upland prairie. 128, 526, 1364, 3212.
- Puccinellia nuttalliana* (Schult.) Hitchc. Nuttall's alkali grass. Fairly common, on moist saline areas. 244, 2852.
- Schedonnardus paniculatus* (Nutt.) Trel. Tumble grass. Rare, on abandoned cropland and waste places. 242, 3517, 3810.
- Setaria viridis* (L.) Beauv. Green foxtail. Introduced. Occasional, on disturbed areas. 3298.
- Sitanion hystrix* (Nutt.) Smith. Squirrel tail. Scarce, on sandy soils. 1107, 1108, 3209.
- Spartina gracilis* Trin. Alkali cord grass. Fairly common, on saline, wet, and alkali areas. 239, 240, 1359.
- Sporobolus cryptandrus* (Torr.) Gray. Sand dropseed. Occasional, on sandy areas. 946.
- Stipa comata* Trin. and Rupr. Needle and thread. Very common, on dry upland prairie. 118, 1356, 2393.
- Stipa spartea* Trin. var. *curtiseta* Hitchc. Western porcupine grass. Occasional, on dry upland prairie. 724, 1396, 2247.
- Stipa viridula* Trin. Green needle grass. Occasional, in moist depressions and along dry roadsides. 234, 1357, 2248, 2849, 3214.
- Triticum aestivum* L. Common wheat. Introduced. Rare, on moist waste areas. 3297.

GRAMINEAE - continued

Vulpia octoflora (Walt.) Rydb. Six-weeks fescue. Occasional, on sandy areas and coulee bottoms. 130, 213, 615, 2188.

CYPERACEAE - Sedge family

Carex brevior (Dewey) Mack. Straw-colored sedge. Occasional, in sloughs and dams. 3415.

Carex eleocharis Bailey. Low sedge. Fairly common, on upland prairie. 112, 1348.

Carex filifolia Nutt. Thread-leaved sedge. Fairly common, on dry sandy areas. 113, 221, 397, 1555, 1578.

Carex heliophila Mack. Sun-loving sedge. Occasional, on upland prairie and coulee banks. 398, 827.

Carex lasiocarpa Ehrh. Hairy-fruited sedge. Occasional, in wet areas and dams. 2867, 3393, 3454.

Carex praegracilis Boott. Graceful sedge. Scarce, in moist slough areas. 2866, 3404.

Carex scoparia Schk. Broom sedge. Occasional, in moist slough areas. 3415.

Eleocharis acicularis (L.) R. and S. Needle spike-rush. Scarce, in alkaline meadows. 731.

Eleocharis palustris (L.) R. and S. Creeping spike-rush. Common, in sloughs and flooded areas. 246, 2862.

Scirpus americanus Pers. Three-square bulrush. Common, in sloughs, streams, and alkaline areas. 192, 2863, 5311.

Scirpus validus Vahl. Great bulrush. Fairly common, in sloughs and dams. 2864, 3403.

JUNCACEAE - Rush family

Juncus balticus Willd. var. *montanus* Engelm. Wire rush. Common, in sloughs and wet areas. 134, 2901, 3394.

Juncus bufonius L. Toad rush. Occasional, on low, moist areas and sloughs. 3034, 3219

Juncus castaneus Smith. Chestnut rush. Occasional, in dry coulee bottoms. 3443.

LILIACEAE - Lily family

- Allium textile* Nels. and Macbr. Prairie onion. Common, on dry upland prairie. 135, 222, 248, 406, 1349.
- Asparagus officinalis* L. Asparagus. Introduced. Rare, on disturbed areas. 5205.
- Fritillaria pudica* (Pursh) Spreng. Yellow-bell. Very rare, on moist slopes. 407.
- Smilacina stellata* (L.) Desf. Star-flowered Solomon's-seal. Fairly common, in moist areas and coulee bottoms. 485, 628, 1371, 2342, 3451.
- Yucca glauca* Nutt. Soapweed. Local, along dry exposed slopes of Lost River coulee. 409, 486, 2123, 2868.
- Zygadenus gramineus* Rydb. Death camas. Common on moist slopes. 133, 1370, 2190.

IRIDACEAE - Iris family

- Sisyrinchium montanum* Greene. Blue-eyed grass. Occasional, in moist draws and low-lying areas. 3142, 3208.

SALICACEAE - Willow family

- Populus angustifolia* James. Narrow-leaf cottonwood. Occasional, on moist areas along roadsides and in coulees. 3405.
- Populus deltoides* Marsh. Western cottonwood. Occasional, in moist coulee bottoms. 136, 3035.
- Salix amygdaloides* Anderss. Peach-leaved willow. Fairly common, in wet places near dams and along roadsides. 3036, 3456.
- Salix fluviatilis* Nutt. River willow. Fairly common, in wet places near dams and along roadsides. 2838, 3037.

ULMACEAE - Elm family

- Ulmus pumila* L. Dwarf elm. Introduced. Scarce, on farmyard. 5237.

SANTALACEAE - Sandalwood family

- Comandra umbellata* (L.) Nutt. var. *pallida* (DC.) Jones. Bastard toadflax. Fairly common, on upland prairie and long coulee edges. 310, 586, 1379, 1721, 2182, 2635, 3388.

POLYGONACEAE - Buckwheat family

Eriogonum cernuum Nutt. Nodding umbrella plant. Common, on badlands and dry coulee slopes. 2553, 2683.

Eriogonum flavum Nutt. Yellow umbrella plant. Common, along dry sandy coulee edges. 137, 1389, 2869, 3516.

Polygonum aviculare L. Prostrate knotweed. Common, on waste areas and roadways. 2346, 2474, 2745, 3039.

Polygonum coccineum Muhl. Water smartweed. Found in sloughs. 3165.

Polygonum convolvulus L. Wild buckwheat. Fairly common, on waste areas and cultivated ground. 2747, 3174.

Polygonum douglasii Greene. Douglas knotweed. Fairly common, along edges of saline sloughs. 996.

Polygonum lapathifolium L. Willow-weed. Fairly common, in sloughs and moist coulee bottoms. 253, 329, 410.

Polygonum scabrum Moench. Green smartweed. Fairly common, in waste areas. 2746.

Rumex acetosella L. Sheep sorrel. Introduced. Fairly common, on waste areas. 635.

Rumex maritimus L. var. *fueginus* (Phil.) Dusen. Golden dock. Common, along saline shores. 3038.

Rumex triangulivalvis (Dans.) Rech. Narrow-leaved dock. Common, in sloughs and moist areas. 252, 3368.

Rumex venosus Pursh. Wild begonia. Fairly common, along roadsides and sandy areas. 138, 250, 432.

CHENOPODIACEAE - Goosefoot family

Atriplex argentea Nutt. Silver saltbush. Fairly common, on eroded saline and waste areas. 2872, 3493.

Atriplex hortensis L. Garden orache. Introduced. Rare, on waste areas. 2982, 3492, 5181.

Atriplex nuttallii Wats. Salt sage. Common, in badlands and saline upland areas. 194, 332, 638, 1372, 3344, 3412.

Atriplex patula var. *hastata* (L.) Gray. Common orache. Occasional, on saline areas. 3462.

CHENOPODIACEAE - continued

- Atriplex suckleyi* (Torr.) Rydb. Endolepis. Occasional, on saline areas.
195, 333, 764, 2440.
- Chenopodium album* L. Lamb's-quarters. Introduced. Common, on waste areas
and roadsides. 3097.
- Chenopodium berlandieri* Moq. var. *farinosum* (Ludwig) Aellen. Stinking
goosefoot. Occasional, on low eroded areas. 2511, 3395.
- Chenopodium fremontii* Wats. Fremont's goosefoot. Occasional, on moist
areas and on sandy prairie areas. 2684, 3306, 3536.
- Chenopodium leptophyllum* (Moq.) Nutt. Narrow-leaved goosefoot. Fairly
common, on dry saline soil. 3428.
- Chenopodium pratericola* Rydb. Oblong-leaved goosefoot. Fairly common, on
dry soil. 2397, 2510, 2556.
- Chenopodium rubrum* L. Red goosefoot. Fairly common, in dry sloughs. 3040.
- Chenopodium salinum* Standl. Oak-leaved goosefoot. Occasional, on eroded
saline soil. 2675.
- Corispermum hyssopifolium* L. Bugseed. Occasional, on sandy areas. 2102.
- Corispermum orientale* Lam. var. *emarginatum* (Rydb.) Macbr. Villose
bugseed. Occasional, on sandy areas. 1058, 2103, 2685.
- Eurotia lanata* (Pursh) Moq. Winter fat. Common, on dry prairie and heavy
soils. 140, 1283, 1398, 4079.
- Kochia scoparia* (L.) Schrad. Summer cypress. Introduced. Common, along
roadsides and on disturbed areas. 2513, 3042.
- Monolepis nuttalliana* (Schultes) Greene. Spear-leaved monolepis. Common,
on saline and alkaline soils. 538, 2512, 3098, 3175.
- Salsola kali* L. Russian thistle. Introduced. Very common, on dry and
disturbed areas. 2983, 3043.
- Sarcobatus vermiculatus* (Hook.) Torr. Greasewood. Fairly common, along
coulee bottoms and on eroded saline areas. 254.
- Suaeda depressa* (Pursh) Wats. Western sea-blight. Fairly common, on saline
areas. 141, 3044.
- Suaeda depressa* (Pursh) Wats. var. *erecta* Wats. Common sea-blight.
Common, on saline areas. 5235.

AMARANTHACEAE - Amaranth family

Amaranthus albus L. Tumbleweed. Fairly common, along roadsides and on waste areas. 2986, 3045, 3144.

Amaranthus graecizans L. Prostrate amaranth. Common, on disturbed areas and along roadsides. 2985.

Amaranthus retroflexus L. Red-root pigweed. Introduced. Very common, on disturbed areas and cultivated fields. 2984.

NYCTAGINACEAE - Four-o'clock family

Abronia micrantha Torr. Sand-puffs. Rare, found only in sand dune area. 3464.

Mirabilis hirsuta (Pursh) MacM. Umbrellawort. Fairly common, on sandy soils. 1140, 2873.

CARYOPHYLLACEAE - Pink family

Cerastium arvense L. Field chickweed. Common, in draws and moist areas. 3100.

Paronychia sessiliflora Nutt. Low whitlow-wort. Common, on dry prairie and rocky areas. 255, 2607, 3424, 3542.

Silene cserei Baumg. Smooth catchfly. Fairly common, on disturbed areas and along roadsides. 3342.

Silene drummondii Hook. Drummond's cockle. Occasional, on sandy soils. 3501.

Silene pratensis (Rafn) Godron and Gren. White cockle. Introduced. Fairly common, on disturbed areas. 2800.

Spergularia marina (L.) Griseb. var. *leiosperma* (Kindb.) Gurke. Salt-marsh sand spurry. Occasional, along margins of saline sloughs. 3046.

RANUNCULACEAE - Crowfoot family

Myosurus aristatus Benth. Bristly mousetail. Occasional, in muddy slough margins. Also found in alkaline and saline sloughs and meadows. 974, 1929, 2126.

Ranunculus acris L. Tall buttercup. Introduced. Occasional, in wet areas near ponds. 258.

Ranunculus aquatilis L. var. *capillaceus* (Thuill.) DC. Large-leaved water crowfoot. Occasional, in wet areas near ponds. 3101.

RANUNCULACEAE - continued

Ranunculus cymbalaria Pursh. Creeping buttercup. Fairly common, in wet areas. 259, 449, 2201, 2875, 3047.

Ranunculus glaberrimus Hook. Early buttercup. Occasional, in moist depressions on prairie. 1558, 2163, 2198, 2333.

Ranunculus macounii Britt. Macoun's buttercup. Fairly common, in moist and wet places. 257.

CAPPARIDACEAE - Caper family

Cleome serrulata Pursh. Rocky mountain bee plant. Occasional, along roadsides. 264, 5309.

CRUCIFERAE - Mustard family

Arabis divaricarpa Nels. Purple rock cress. Occasional, on upland prairie. 5304.

Arabis holboellii Hornem. var. *retrofracta* (Graham) Rydb. Holboell's rock cress. Common, on upland prairie. 263, 375, 450, 546, 793, 833, 1350, 2164, 3396.

Brassica campestris L. Rape. Introduced. Occasional, on waste places. 3226, 3423.

Brassica juncea (L.) Cosson. Indian mustard. Introduced. Occasional, on waste places. 3225, 3494.

Camelina microcarpa Andrzej. Small-seeded false flax. Introduced. Scarce, in waste areas. 5312.

Capsella bursa-pastoris (L.) Medic. Shepherd's purse. Introduced. Scarce, in waste areas. 145.

Conringia orientalis (L.) Dum. Hare's-ear mustard. Introduced. Scarce, in waste areas. 215.

Descurainia pinnata (Walt.) Britt. Green tansy mustard. Occasional, on dry prairie. 867, 868, 2613.

Descurainia sophia (L.) Webb. Flixweed. Introduced. Common, on disturbed areas. 262, 2444, 2612, 3105.

Draba nemorosa L. Yellow whitlow-grass. Common, on moist slopes on upland prairie. 832, 1931, 2207.

CRUCIFERAE - continued

- Erysimum inconspicuum* (Wats.) MacM. Small-flowered prairie-rocket. Fairly common, on dry prairie and disturbed areas. 147, 2205, 2560, 3397.
- Lepidium densiflorum* Schrad. Common peppergrass. Common, along roadsides and fields. 261, 2203, 3522, 5308.
- Lepidium ramosissimum* Nels. Branched peppergrass. Fairly common, on dry prairie. 260, 3103.
- Lesquerella arenosa* (Richards) Rydb. Silvery bladderpod. Common, on eroded, exposed sites. 414, 643, 831, 1692, 2355, 2610, 2876, 5301.
- Rorippa islandica* (Oeder) Borbas. Marsh yellow cress. Fairly common, on moist areas. 3145.
- Rorippa sinuata* (Nutt.) Hitchc. Spreading yellow cress. Occasional, in dry stream beds. 2204.
- Sisymbrium altissimum* L. Tumbling mustard. Introduced. Common, on disturbed areas. 146, 2443, 2611.
- Thlaspi arvense* L. Stinkweed. Introduced. Common, on disturbed areas. 144, 2609.

GROSSULARIACEAE - Currant family

- Ribes aureum* Pursh. Golden currant. Scarce, in moist coulee bottoms. 2841.
- Ribes oxycanthoides* L. Wild gooseberry. Fairly common, in moist draws. 2840, 3385.

ROSACEAE - Rose family

- Amelanchier alnifolia* Nutt. Saskatoon. Scarce, on moist coulee slopes. 2367, 3392.
- Chamaerhodos erecta* (L.) Bunge var. *parviflora* (Nutt.) Hitchc. Ground rose. Fairly common, on gravelly soil. 149, 2366, 2486.
- Fragaria virginiana* Duchesne. Wild strawberry. Scarce, in moist draws. 3504.
- Geum allepicum* Jacq. Yellow avens. Rare, in moist coulee bottoms. 270.
- Geum triflorum* Pursh. Old man's whiskers. Fairly common, on flats and moist slopes on upland prairie. 224, 271, 399, 722, 1941, 2208.

ROSACEAE - continued

- Potentilla anserina* L. Silverweed. Fairly common, along coulee draws. 376, 452, 3409.
- Potentilla arguta* Pursh. White cinquefoil. Fairly common, on moist prairie. 269, 2760.
- Potentilla concinna* Richards. Early cinquefoil. Common, on dry prairie. 1351, 1562, 2175, 2842, 3389.
- Potentilla concinna* Richards var. *divisa* Rydb. Dissected cinquefoil. Fairly common, in moist draws and coulees. 834.
- Potentilla gracilis* Dougl. Graceful cinquefoil. Fairly common, on low moist areas. 2759, 2877, 3218, 3444.
- Potentilla norvegica* L. Rough cinquefoil. Scarce, in moist areas. 3048.
- Potentilla pensylvanica* L. Prairie cinquefoil. Fairly common, on moist slopes. 267, 721, 3515.
- Potentilla quinquefolia* Rydb. Five-fingered cinquefoil. Scarce, on dry prairie. 268, 835.
- Prunus virginiana* L. Choke cherry. Scarce, in moist draws. 491, 2878, 3293.
- Rosa acicularis* Lindl. Prickly rose. Scarce, along coulee draws. 3215.
- Rosa woodsii* Lindl. Common wild rose. Fairly common, on moist slopes. 2494, 3398.

LEGUMINOSAE - Pea family

- Astragalus agrestis* Dougl. Purple milkvetch. Fairly common, on moist grassland. 225, 279, 280, 434, 2882, 5222.
- Astragalus bisulcatus* (Hook.) Gray. Two-grooved milkvetch. Fairly common, on dry prairie. 276, 395, 1374, 1504, 1616.
- Astragalus cicer* L. Cicer milkvetch. Introduced. Scarce, on cultivated land. 1391, 3050.
- Astragalus crassicaupus* Nutt. Ground plum. Scarce, on moist locations on prairie. 156, 2015, 3382, 5306.
- Astragalus gilviflorus* Sheldon. Cushion milkvetch. Common, on eroded areas and dry slopes. 278, 841, 1418, 1945, 2167, 5131.

LEGUMINOSAE - continued

- Astragalus missouriensis* Nutt. Missouri milkvetch. Fairly common, on slopes and eroded hillsides. 455, 556, 787, 839, 939, 1505, 1693, 2013, 5130.
- Astragalus pectinatus* Dougl. Narrow-leaved milkvetch. Fairly common, on dry prairie. 155, 277, 1375, 1696.
- Astragalus purshii* Dougl. Pursh's milkvetch. Fairly common, on dry hillsides. 840, 1471, 1932, 2014, 2166.
- Astragalus spatulatus* Sheldon. Tufted milkvetch. Fairly common, along exposed coulee edges. 274, 1674, 2114, 2177, 3487, 5305.
- Astragalus striatus* Nutt. Ascending purple milkvetch. Fairly common, on dry prairie. 709, 1390, 1726, 2621, 2881.
- Astragalus tenellus* Pursh. Loose-flowered milkvetch. Fairly common, on eroded areas and dry slopes. 154, 275, 2274.
- Caragana arborescens* Lam. Common caragana. Introduced. Scarce, on old farmsteads. 3117.
- Glycyrrhiza lepidota* (Nutt.) Pursh. Wild licorice. Common, on sandy areas and valleys. 159.
- Hedysarum alpinum* L. var. *americanum* Michx. American hedysarum. Fairly common, in moist draws. 217.
- Hedysarum boreale* Nutt. Northern hedysarum. Scarce, on dry hillsides. 3486.
- Lupinus pusillus* Pursh. Annual lupine. Common, on sandy areas. 377, 1809.
- Medicago falcata* L. Yellow alfalfa. Introduced. Fairly common, along roadsides and in waste places. 3146.
- Medicago lupulina* L. Black medick. Introduced. Rare, on waste areas. 3049.
- Medicago sativa* L. Alfalfa. Introduced. Fairly common, along roadsides and waste places. 2449.
- Melilotus alba* Desr. White sweet clover. Introduced. Common, along roadsides and waste places. 2763, 3147.
- Melilotus officinalis* (L.) Lam. Yellow sweet clover. Introduced. Common, along roadsides and waste places. 2619, 2764, 2802.
- Oxytropis campestris* (L.) DC. var. *gracilis* (Nels.) Barneby. Late yellow locoweed. Common, on moist upland prairie. 157, 282, 283.

LEGUMINOSAE - continued

- Oxytropis sericea* Nutt. var. *spicata* (Hook.) Barneby. Early yellow locoweed. Common, on upland prairie and hillsides. 378, 1419.
- Oxytropis splendens* Dougl. Showy locoweed. Scarce, in moist areas. 416.
- Petalostemon candidum* (Willd.) Michx. White prairie clover. Fairly common, on dry coulee slopes. 273, 1399, 3303.
- Petalostemon purpureum* (Vent.) Rydb. Purple prairie clover. Scarce, in moist areas. 3532.
- Psoralea lanceolata* Pursh. Scurf pea. Common, on sandy areas. 2620, 2723, 2880.
- Thermopsis rhombifolia* (Nutt.) Richards. Golden bean. Common, along roadsides and sandy soils. 151, 1373, 2165, 2703.
- Vicia americana* Muhl. American vetch. Common, along roadsides and depressions in upland prairie. 160, 1376, 2215.

LINACEAE - Flax family

- Linum lewisii* Pursh. Wild blue flax. Common, on upland prairie and sandy areas. 286, 343, 344, 658, 2883, 3121, 3207.
- Linum rigidum* Pursh. Yellow flax. Common, on sandy soils. 287, 364, 381, 382, 741, 3177, 3302, 3362.

EUPHORBIACEAE - Spurge family

- Euphorbia serpyllifolia* Pers. Thyme-leaved spurge. Fairly common, on moist sandy soils. 1400, 2136, 2277, 2768.

ANACARDIACEAE - Sumach family

- Rhus trilobata* Nutt. Skunk bush. Common, on coulee slopes. 161, 288, 1509, 3446, 3530.

ACERACEAE - Maple family

- Acer negundo* L. Manitoba maple. Scarce, on farmyards and roadsides. 3122.

MALVACEAE - Mallow family

Malva rotundifolia L. Common mallow. Introduced. Rare, on waste areas.
2331, 3051.

Sphaeralcea coccinea (Pursh) Rydb. Scarlet globe-mallow. Common, on
disturbed areas along roadsides. 162, 218, 1401, 3447.

VIOLACEAE - Violet family

Viola adunca Smith. Early blue violet. Occasional, on moist prairie
slopes. 3383.

Viola nuttallii Pursh. Yellow prairie violet. Fairly common, in moist
locations. 444, 2168, 3386.

Viola nuttallii Pursh. var. *vallicola* (Nels.) St. John. Valley yellow
violet. Occasional, on moist prairie. 3123.

CACTACEAE - Cactus family

Coryphantha vivipara (Nutt.) Britt. and Brown. Cushion cactus. Fairly
common, on upland prairie. 164.

Opuntia polyacantha Haw. Prickly pear. Common, on upland prairie and
denuded areas. 163, 3124.

ELAEAGNACEAE - Oleaster family

Elaeagnus angustifolia L. Russian olive. Introduced. Scarce, on abandoned
farmyards. 2885.

Elaeagnus commutata Bernh. Wolf-willow. Common, on sandy soils and along
coulee slopes. 458, 2497, 2623.

Shepherdia argentea Nutt. Thorny buffalo-berry. Fairly common, in coulees
and along drainage channels. 289, 2886.

ONAGRACEAE - Evening primrose family

Boisduvalia glabella (Nutt.) Walp. Smooth spike-primrose. Rare, on moist,
alkaline mud flats. 2219, 2678.

Epilobium angustifolium L. Fireweed. Rare, in moist coulees. 383.

Epilobium ciliatum Raf. Northern willowherb. Occasional, in moist areas. 3052.

ONAGRACEAE - continued

Epilobium paniculatum Nutt. Annual willowherb. Occasional, on moist slopes. 1010, 2564, 3305.

Gaura coccinea Pursh. Scarlet butterfly-weed. Fairly common, on upland prairie. 167, 201, 2625.

Oenothera andina Nutt. Upland evening primrose. Scarce, on dry sandy soils. 2624, 2770, 5126.

Oenothera biennis L. Common evening primrose. Occasional, on moist slopes. 1118, 1408.

Oenothera caespitosa Nutt. Butte primrose. Common, on dry slopes of gumbo or clay soils and gumbo flats. 166, 219, 226, 459, 560, 2218, 3125.

Oenothera flava (Nels.) Garrett. Yellow evening primrose. Occasional, along drainage channels and clay flats. 1402, 2996, 3150.

Oenothera nuttallii Sweet. White evening primrose. Common, on sandy soil and along roadsides. 290, 291, 1895, 2450, 3361.

UMBELLIFERAE - Parsley family

Carum carvi L. Caraway. Introduced. Rare, on waste areas. 561.

Cymopterus acaulis (Pursh) Raf. Plains cymopterus. Common, on dry upland prairie. 796, 821, 842, 1730, 2337.

Lomatium foeniculaceum (Nutt.) Coult. and Rose. Hairy-fruited wild parsley. Common, on upland prairie. 293, 1219, 1352, 1731.

Lomatium macrocarpum (Hook. and Arn.) Coult. and Rose. Long-fruited wild parsley. Common, on upland prairie. 723, 843.

Lomatium triternatum (Pursh) Coult. and Rose. Western wild parsley. Rare, in moist coulees 3381, 5303.

Musineon divaricatum (Pursh) Nutt. Leafy musineon. Common, on dry upland prairie. 1729, 2169, 2336.

CORNACEAE - Dogwood family

Cornus stolonifera Michx. Red-osier dogwood. Occasional, in moist coulees. 5186.

PRIMULACEAE - Primrose family

Androsace septentrionalis L. Pygmy flower. Fairly common, on upland prairie. 1324, 3491.

Dodecatheon conjugens Greene. Shooting star. Occasional, in moist areas. 294, 3387.

Dodecatheon pulchellum (Raf.) Merr. Saline shooting star. Common, in wet areas and saline flats. 228, 401, 1353, 1564, 2170, 2596.

Glaux maritima L. Sea milkwort. Occasional, in moist, saline areas. 435, 494.

GENTIANACEAE - Gentian family

Gentiana affinis Griseb. Prairie gentian. Occasional, in moist grassland. 384, 2690, 3358.

Gentianella amarella (L.) Börner. Northern gentian. Scarce, in wet areas. 202.

APOCYNACEAE - Dogbane family

Apocynum androsaemifolium L. Spreading dogbane. Occasional, in moist creek beds. 3416.

Apocynum cannabinum L. Indian hemp. Occasional, on moist areas. 3410.

ASCLEPIADACEAE - Milkweed family

Asclepias speciosa Torr. Showy milkweed. Occasional, along roadsides and on moist areas. 744, 1392, 3407.

CONVOLVULACEAE - Morning-glory family

Convolvulus arvensis L. Field bindweed. Introduced. Occasional, on waste areas and roadsides. 3299.

Convolvulus sepium L. Hedge bindweed. Occasional, on waste places and roadsides. 2285, 3411.

POLEMONIACEAE - Phlox family

Collomia linearis Nutt. Narrow-leaved collomia. Fairly common, on dry soils and sandy places. 873, 2223, 2888, 3128.

POLEMONIACEAE - continued

Linanthus septentrionalis Mason. Northern linanthus. Rare, on sandy areas.
1191, 2628, 3519.

Navarretia minima Nutt. Pin-cushion plant. Fairly common, on coulee
slopes. 1124, 1393, 2141, 3155.

Phlox hoodii Richards. Moss phlox. Common, on upland prairie and eroded
soils. 114, 229, 402, 1935, 2105, 2339.

BORAGINACEAE - Borage family

Cryptantha fendleri (Gray) Greene. Fendler's crypthanthe. Common, on sandy
areas. 2500, 2630.

Cryptantha macounii (Eastw.) Payson. Macoun's crypthanthe. Occasional, on
upland prairie. 2629.

Cryptantha nubigena (Greene) Payson. Clustered oreocarya. Fairly common,
on upland prairie. 170, 298, 2230.

Heliotropium curassavicum L. Saline heliotrope. Occasional, on moist
saline flats. 169, 296, 1291, 4714.

Lappula echinata Gilib. Blue-bur. Introduced. Common, on waste areas.
297, 495, 2229, 2286, 2774, 3220.

Lappula redowskii (Hornem.) Greene. Western blue-bur. Fairly common, on
dry prairie and along roadsides. 462, 496, 570, 2142.

Lithospermum canescens (Michx.) Lehm. Hoary puccoon. Occasional, on dry or
sandy prairie. 418, 2181.

Lithospermum incisum Lehm. Narrow-leaved puccoon. Fairly common, on upland
prairie. 385, 419, 2381, 2631, 3221, 3503.

Plagiobothrys scouleri (H. and A.) Johnston. Scouler's popcorn-flower.
Occasional, on saline areas. 3440, 5153.

LABIATAE - Mint family

Hedeoma hispida Pursh. Rough penny royal. Fairly common, on upland
prairie. 577, 1727, 3514.

Mentha arvensis L. Wild mint. Common, in low-lying wet locations. 2288.

SOLANACEAE - Nightshade family

Lycium halimifolium Mill. Matrimony vine. Introduced. Rare, on abandoned farmsteads. 3291, 3437, 3465.

Solanum triflorum Nutt. Wild tomato. Occasional, on disturbed areas and roadsides. 299, 300, 3502.

SCROPHULARIACEAE - Figwort family

Castilleja sessiliflora Pursh. Downy paintbrush. Occasional, on sandy areas. 3213, 3399.

Limosella aquatica L. Mudwort. Occasional, in muddy or wet locations. 3131.

Linaria dalmatica (L.) Mill. Broad-leaved toad-flax. Introduced. Rare, on disturbed and waste areas. 3341.

Mimulus guttatus DC. Yellow monkey-flower. Very rare, in coulees. 306.

Orthocarpus luteus Nutt. Owl-clover. Very common, on upland prairie. 307, 499, 2776, 3266, 4866.

Penstemon albidus Nutt. White beard-tongue. Common, along hillsides and coulee edges. 749, 1377, 3400, 3448.

Penstemon confertus Dougl. Yellow beard-tongue. Fairly common, on upland prairie. 578, 2031, 2234, 2633, 3130, 3449.

Penstemon gracilis Nutt. Lilac-flowered beard-tongue. Rare, on moist grassland. 302.

Penstemon nitidus Dougl. Smooth blue beard-tongue. Occasional, on eroded areas and coulee banks. 301, 1473, 5300.

Penstemon procerus Dougl. Slender blue beard-tongue. Fairly common, around slough edges and low areas. 304, 3129.

Veronica peregrina L. var. *xalapensis* (HBK) St. John and Warren. Hairy speedwell. Common, on moist locations. 2146, 2235, 3132, 3524.

OROBANCHACEAE - Broom-rape family

Orobanche fasciculata Nutt. Clustered broom-rape. Fairly common, on upland prairie. 1084, 3531.

Orobanche ludoviciana Nutt. Broom-rape. Occasional, on sandy prairie. 3300, 3534.

LENTIBULARIACEAE - Bladderwort family

Utricularia vulgaris L. Common bladderwort. Fairly common, in sloughs and dams. 1327.

PLANTAGINACEAE - Plantain family

Plantago elongata Pursh. Linear-leaved plantain. Occasional, in moist locations. 2237.

Plantago eriopoda Torr. Saline plantain. Common, on saline or alkaline soils and slough margins. 309, 436, 2890.

Plantago major L. Common plantain. Introduced. Fairly common, in farmyards, roads, and waste areas. 308, 420, 3295.

Plantago patagonica Jacq. Pursh's plantain. Common, on upland prairie. 387, 438, 876, 1378.

Plantago patagonica Jacq. var. *spinulosa* (Dcne.) Gray. Spiny plantain. Fairly common, upland prairie and disturbed areas. 437, 2697, 2779, 3134.

RUBIACEAE - Madder family

Galium boreale L. Northern bedstraw. Fairly common, in moist low places. 500, 2780, 2891.

CAPRIFOLIACEAE - Honeysuckle family

Symphoricarpos occidentalis Hook. Western snowberry. Fairly common, in draws and coulees. 173, 2455.

CAMPANULACEAE - Bluebell family

Campanula rotundifolia L. Bluebell. Common, in low-lying areas and along roadsides. 388, 587, 3216.

COMPOSITAE - Composite family

Achillea millefolium L. Common yarrow. Introduced. Common, along roadsides and on waste places. 2785.

Achillea millefolium L. var. *lanulosa* (Nutt.) Piper. Woolly yarrow. Fairly common, in moist locations and along roadsides. 181, 206, 1382.

COMPOSITAE - continued

- Agoseris glauca* (Pursh) Raf. False dandelion. Fairly common, on moist prairie. 506, 1407, 3160.
- Agoseris glauca* (Pursh) Raf. var. *agrestis*. Long-leaved false dandelion. Occasional, on moist prairie. 5313.
- Agoseris glauca* (Pursh) Raf. var. *dasycephala* (T. and G.) Jeps. Pubescent false dandelion. Fairly common, on hillsides and moist prairie. 359, 505, 598, 1395.
- Antennaria dimorpha* (Nutt.) T. and G. Cushion pussy-toes. Scarce, on upland prairie. 1268, 1355.
- Antennaria nitida* Greene. Pussy-toes. Fairly common, on upland prairie. 78, 427, 5113.
- Antennaria parvifolia* Nutt. Small-leaf pussy-toes. Common, on upland prairie. 3390, 3505, 5127.
- Antennaria rosea* Greene. Rosy pussy-toes. Fairly common, on upland prairie. 426, 590, 3490, 5128.
- Arnica fulgens* Pursh. Shining arnica. Common, in moist locations. 324, 1521, 1914, 2240, 3224.
- Artemisia biennis* Willd. Biennial sage wort. Occasional, in moist areas and on disturbed areas. 595.
- Artemisia campestris* L. Plains wormwood. Fairly common, on sandy soils and dry slopes. 469, 909, 3363.
- Artemisia cana* Pursh. Silvery sagebrush. Common, on upland prairie and along stream banks. 183, 3023, 5047.
- Artemisia dracunculus* L. Linear-leaved wormwood. Fairly common, on upland prairie. 593.
- Artemisia frigida* Willd. Pasture sage. Common, on upland prairie and on disturbed areas. 182, 1621, 3304, 3526.
- Artemisia longifolia* Nutt. Long-leaved sage. Common, on eroded slopes and in badlands. 323, 880, 2161.
- Artemisia ludoviciana* Nutt. var. *gnaphalodes* (Nutt.) T. and G. Slender sage. Common, in moist areas. 322.
- Artemisia ludoviciana* Nutt. var. *pabularis* (Nels.) Fern. Prairie sage. Common, in saline areas. 3021.

COMPOSITAE - continued

- Aster brachyactis* Blake. Rayless aster. Fairly common, on moist saline areas. 3056.
- Aster ericoides* L. Many-flowered aster. Common, on upland prairie. 423, 2571, 3469.
- Aster falcatus* Lindl. Creeping white prairie aster. Fairly common, on upland prairie. 3356.
- Chrysopsis villosa* (Pursh) Nutt. Golden aster. Common, on upland prairie. 313, 422, 1405, 3357.
- Chrysothamnus nauseosus* (Pall.) Brit. Rabbitbrush. Occasional, on eroded slopes. 314, 352, 1294, 1409, 1733, 3426.
- Cirsium arvense* (L.) Scop. Canada thistle. Common, along roadsides and in moist disturbed areas. 3026, 3159, 3259.
- Cirsium flodmanii* (Rydb.) Arthur. Flodman's thistle. Occasional, on moist prairie. 3301.
- Cirsium undulatum* (Nutt.) Spreng. Wavy-leaved thistle. Fairly common, on upland prairie and roadsides. 3408.
- Coreopsis tinctoria* Nutt. Common tickseed. Common, along slough edges. 318, 2295, 3290.
- Crepis runcinata* (James) T. and G. Meadow hawk's-beard. Fairly common, in saline sloughs. 944.
- Erigeron caespitosus* Nutt. Tufted fleabane. Fairly common, on upland prairie. 425, 1949, 3450.
- Erigeron canadensis* L. Canada fleabane. Common, on disturbed areas. 316, 2293, 2582, 3061, 3181.
- Erigeron pumilus* Nutt. Hairy fleabane. Common, on upland prairie. 315, 1394, 3223.
- Franseria acanthicarpa* (Hook.) Coville. Bur ragweed. Occasional, in sandy areas. 1040, 2570, 2691.
- Gaillardia aristata* Pursh. Great-flowered gaillardia. Common, on upland prairie. 320, 1381, 2892, 3135.
- Gnaphalium palustre* Nutt. Western marsh cudweed. Common, on slough edges and moist areas. 3062.

COMPOSITAE - continued

- Grindelia squarrosa* (Pursh) Dunal. Gumweed. Common, on saline flats and slough margins. 312, 3359.
- Gutierrezia sarothrae* (Pursh) Britt. and Rusby. Broomweed. Common, on upland prairie. 203, 685, 1404, 2456.
- Haplopappus lanceolatus* (Hook.) T. and G. Lance-leaved ironplant. Fairly common, in moist saline areas. 176, 2807.
- Haplopappus nuttallii* T. and G. Toothed ironplant. Fairly common, on dry eroded areas. 956, 1295, 2636, 3498, 4948.
- Haplopappus spinulosus* (Pursh) DC. Spiny ironplant. Fairly common, on upland prairie and hillsides. 353.
- Helianthus couplandii* Boivin. Annual sunflower. Fairly common, in sandy areas and along roadsides. 2538, 2694.
- Helianthus nuttallii* T. and G. Common tall sunflower. Fairly common, in low moist areas. 3294.
- Hymenoxys acaulis* (Pursh) Parker. Butte marigold. Common, on eroded hillsides. 985, 1969, 2391, 2502.
- Hymenoxys richardsonii* (Hook.) Cockerell. Colorado rubber-plant. Common, on upland prairie and dry hillsides. 179, 319, 1380, 1518, 1903, 2392.
- Iva axillaris* Pursh. Poverty-weed. Fairly common, on saline heavy textured soils. 421, 2150.
- Iva xanthifolia* Nutt. False ragweed. Occasional, along roadsides and disturbed areas. 2568, 3267, 3346.
- Lactuca pulchella* (Pursh) DC. Common blue lettuce. Common, in moist areas and along roadsides. 2301.
- Lactuca serriola* L. Prickly lettuce. Introduced. Common, along roadsides and waste places. 3029.
- Liatris punctata* Hook. Dotted blazing star. Common, on upland prairie and dry hillsides. 311, 1403, 1541, 3355.
- Lygodesmia juncea* (Pursh) Don. Skeleton-weed. Common, on upland prairie and on sandy soils. 185, 1383, 2576.
- Lygodesmia rostrata* Gray. Annual skeleton-weed. Occasional, on sandy soils. 2104, 3544.

COMPOSITAE - continued

- Machaeranthera canescens* (Pursh) Gray. Hoary aster. Common, along roadsides and in moist areas. 424, 689, 1298, 2151, 3367.
- Madia glomerata* Hook. Tarweed. Occasional, on slough edges. 3015.
- Microseris cuspidata* (Pursh) Schultz-Bip. Prairie false dandelion. Rare, Lost River and along dry coulees. 2184, 3391, 5302.
- Ratibida columnifera* (Nutt.) Wooton and Standl. Prairie cone-flower. Fairly common, in moist areas and along roadsides. 205, 317, 2083, 3354.
- Senecio canus* Hook. Prairie groundsel. Common, on upland prairie. 325, 428, 1089, 2638, 3452.
- Senecio integerrimus* Nutt. Entire-leaved groundsel. Fairly common, on moist prairie. 2242.
- Senecio integerrimus* Nutt. var. *exaltatus* (Nutt.) Cronq. Western groundsel. Fairly common, on moist prairie. 1090.
- Solidago missouriensis* Nutt. Low goldenrod. Fairly common, on upland prairie. 204, 1406, 2809.
- Solidago rigida* L. var. *humilis* Porter. Stiff goldenrod. Fairly common, on upland prairie. 715, 3366, 3543.
- Solidago spathulata* DC. Mountain goldenrod. Fairly common, on upland prairie. 3365.
- Sonchus arvensis* L. Perennial sow thistle. Introduced. Fairly common, on moist disturbed areas. 3171, 3345.
- Sonchus asper* (L.) Hill. Annual sow thistle. Introduced. Fairly common, on moist disturbed areas. 3031.
- Sonchus uliginosus* Bieb. Smooth perennial sow thistle. Introduced. Fairly common, on moist disturbed areas. 3030, 3172.
- Taraxacum officinale* Weber. Dandelion. Introduced. Common, on moist disturbed areas. 470, 599, 2708.
- Townsendia exscapa* (Richards.) Porter. Low townsendia. Occasional, on eroded dry hillsides. 1965.
- Tragopogon dubius* Scop. Goat's-beard. Introduced. Common, on upland prairie, roadsides and waste places. 2640, 3343.
- Xanthium strumarium* L. Cocklebur. Occasional, on moist disturbed areas. 3369.

BIBLIOGRAPHY

- Boivin, B. 1967-1981. Flora of the Prairie Provinces, Vols. 1-5.
Provancheria, Laval University, Quebec, Quebec.
- Coupland, R. T. 1950. Ecology of Mixed Prairie in Canada. Ecol. Mono.
20: 271-315.
- Coupland, R. T. 1961. A reconsideration of grassland classification in the
Northern Great Plains of North America. J. Ecol. 49: 135-167.
- Hitchcock, C. L. and A. Cronquist. 1973. Flora of the Pacific Northwest.
Univ. of Washington Press, Seattle. 730 pp.
- Johnson, Hope and Beryl Hallworth. 1975. Further discoveries of sand verbena
in Alberta. Blue Jay. 33: 13-15.
- Looman, J. 1980. The vegetation of the Canadian Prairie Provinces II. The
grasslands, Part 1. Phytocoenologia 8: 153-190.
- Looman, J., and K. F. Best. 1979. Budd's flora of the Canadian Prairie
Provinces. Agr. Can. Publ. 1662. 863 pp.
- Moss, E. H. 1944. *Lilaea scillioides* in southeastern Alberta. Rhodora
46: 205-206.
- Moss, E. H. 1955. The vegetation of Alberta. Bot. Rev. 21: 493-567.
- Packer, J. G. 1983. Flora of Alberta. 2nd Ed. Univ. of Toronto Press.
Toronto, Ontario. 687 pp.
- Scoggan, H. J. 1978-1979. The Flora of Canada, vols. 1-4. National Museums
of Canada. Ottawa, Ontario.
- Smith, W. W., and C. A. Wallis. 1976. Extra-limital occurrence of Rocky
Mountain Juniper in southeastern Alberta. Blue Jay 34: 67-68.
- Wallis, C. A. 1976. Milk river canyon resource evaluation. Alberta
Recreation, Parks and Wildlife, Edmonton, Alberta 122 pp.

LIBRARY / BIBLIOTHEQUE

AGRICULTURE CANADA OTTAWA K1A 0C5

3 9073 00036147 9