A UNITED STATES
DEPARTMENT OF
COMMERCE
PUBLICATION

NOAA Technical Report NMFS CIRC-385

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

Fishery Publications, Calendar Year 1972: Lists and Indexes

LEE C. THORSON and MARY ELLEN ENGETT

SEATTLE, WA
NOVEMBER 1973

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Circulars

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyses, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS CIRC series continues a series that has been in existence since 1941. The Circulars are technical publications of general interest intended to aid conservation and management. Publications that review in considerable detail and at a high technical level certain broad areas of research appear in this series. Technical papers originating in economics studies and from management investigations appear in the Circular series.

NOAA Technical Reports NMFS CIRC are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from NOAA Publications Section, Rockville, Md. 20852. Recent Circulars are:

- 315. Synopsis of biological data on the chum salmon, *Oncorhynchus keta* (Walbaum) 1792. By Richard G. Bakkala. March 1970, iii + 89 pp., 15 figs., 51 tables.
- 319. Bureau of Commercial Fisheries Great Lakes Fishery Laboratory, Ann Arbor, Michigan. By Bureau of Commercial Fisheries. March 1970, 8 pp., 7 figs.
- 330. EASTROPAC Atlas: Vols. 4, 2. Catalog No. I 49.4:330/(vol.) 11 vols. (\$4.75 each). Available from the Superintendent of Documents, Washington, D.C. 20402.
- 331. Guidelines for the processing of hot-smoked chub. By H. L. Seagran, J. T. Graikoski, and J. A. Emerson. January 1970, iv + 23 pp., 8 figs., 2 tables.
- 332. Pacific hake. (12 articles by 20 authors.) March 1970, iii + 152 pp., 72 figs., 47 tables.
- 333. Recommended practices for vessel sanitation and fish handling. By Edgar W. Bowman and Alfred Larsen. March 1970, iv + 27 pp., 6 figs.
- 335. Progress report of the Bureau of Commercial Fisheries Center for Estuarine and Menhaden Research, Pesticide Field Station, Gulf Breeze, Fla., fiscal year 1969. By the Laboratory staff. August 1970, iii + 33 pp., 29 figs., 12 tables.
- 336. The northern fur seal. By Ralph C. Baker, Ford Wilke, and C. Howard Baltzo. April 1970, iii + 19 pp., 13 figs.
- 337. Program of Division of Economic Research, Bureau of Commercial Fisheries, fiscal year 1969. By Division of Economic Research. April 1970, iii + 29 pp., 12 figs., 7 tables.
- 338. Bureau of Commercial Fisheries Biological Laboratory, Auke Bay, Alaska. By Bureau of Commercial Fisheries. June 1970, 8 pp., 6 figs.
- 339. Salmon research at Ice Harbor Dam. By Wesley J. Ebel. April 1970, 6 pp., 4 figs.
- 340. Bureau of Commercial Fisheries Technological Laboratory, Gloucester, Massachusetts. By Bureau of Commercial Fisheries. June 1970, 8 pp., 8 figs.
- 341. Report of the Bureau of Commercial Fisheries Biological Laboratory, Beaufort, N.C., for the fiscal year ending June 30, 1968. By the Laboratory staff. August 1970, iii + 24 pp., 11 figs., 16 tables.
- 342. Report of the Bureau of Commercial Fisheries Biological Laboratory, St. Petersburg Beach, Florida, fiscal year 1969. By the Laboratory staff. August 1970, iii + 22 pp., 20 figs., 8 tables.
- 343. Report of the Bureau of Commercial Fisheries Biological Laboratory, Galveston, Texas, fiscal year 1969. By the Laboratory staff. August 1970, iii + 39 pp., 28 figs., 9 tables.
- 344. Bureau of Commercial Fisheries Tropical Atlantic Biological Laboratory progress in research 1965-69, Miami, Florida. By Ann Weeks. October 1970, iv + 65 pp., 53 figs.
- 346. Sportsman's guide to handling, smoking, and preserving Great Lakes coho salmon. By Shearon Dudley, J. T. Graikoski, H. L. Seagran, and Paul M. Earl. September 1970, iii + 28 pp., 15 figs.
- 347. Synopsis of biological data on Pacific ocean perch, *Sebastes alutus*. By Richard L. Major and Herbert H. Shippen. December 1970, iii + 38 pp., 31 figs., 11 tables.

Continued on inside back cover.

U.S. DEPARTMENT OF COMMERCE

Frederick B. Dent, Secretary

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

Robert M. White, Administrator

NATIONAL MARINE FISHERIES SERVICE

Robert W. Schoning, Director

NOAA Technical Report NMFS CIRC-385

**Fishery Publications,
Calendar Year 1972:
Lists and Indexes**

LEE C. THORSON and MARY ELLEN ENGETT

SEATTLE, WA
NOVEMBER 1973

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

	Page
ABSTRACT	1
INTRODUCTION	1
LISTS	2
Circular	2
NOAA Technical Report NMFS CIRC	3
Data Report	4
Fishery Facts	5
NOAA Technical Report NMFS SSRF	5
NOAA Technical Memorandum NMFS	9
AUTHOR INDEX	10
SUBJECT INDEX	11
INDEX BY MARSDEN SQUARES	22

FISHERY PUBLICATIONS, CALENDAR YEAR 1972: LISTS AND INDEXES

By

LEE C. THORSON and MARY ELLEN ENGETT

Scientific Publications Staff
National Marine Fisheries Service

ABSTRACT

The following series of fishery publications of the National Marine Fisheries Service, National Oceanic and Atmospheric Administration, in calendar year 1972 are listed numerically (with abstracts) and indexed by author, subject, and geographic area: NOAA Technical Report NMFS CIRC (formerly Circular); Data Report; Fishery Facts; NOAA Technical Report NMFS SSRF; and NOAA Technical Memorandum NMFS.

INTRODUCTION

This document provides for calendar year 1972 numerical lists (with abstracts) and indexes by author, subject, and geographical area, the following series of publications of the National Marine Fisheries Service, National Oceanic and Atmospheric Administration:

Circular
Data Report
Fishery Facts
Special Scientific Report—Fisheries
Technical Memorandum

The document is divided into four principal sections:

Numerical listing of series (with abstracts)
Author index
Subject index
Index by Marsden squares

The last section has been included to afford easy access to the publications for those persons interested in specific geographical areas. Figure 1 shows the Marsden squares treated in the several publications.

The series abbreviations used in the indexes are:

Circular	C
NOAA Technical Report NMFS CIRC	C
Data Report	D
Fishery Facts	FF
NOAA Technical Report NMFS SSRF	S
NOAA Technical Memorandum NMFS	TM

LISTS

Circular

- 330, Vol. 1. EASTROPAC Atlas: Physical Oceanographic and Meteorological Data from Principal Participating Ships, First Survey Cruise, February-March 1967. By Cuthbert M. Love, (editor). June 1972, xii + 157 pp., 255 figures. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402—Price \$4.75 per volume.

ABSTRACT

This atlas contains charts depicting the distribution of physical, chemical, and biological oceanographic properties and associated meteorological properties observed during EASTROPAC. EASTROPAC was an international cooperative investigation of the eastern tropical Pacific Ocean (20° N. to 20° S., and from the west coasts of the American continents to 119° W.) which was intended to provide data necessary for a more effective use of the marine resources of the area, especially tropical tunas, and also to increase knowledge of the ocean circulation, air-sea interaction, and ecology. The Bureau of Commercial Fisheries (now National Marine Fisheries Service) was the coordinating agency. The field work, from February 1967 through March 1968, was divided into seven 2-month cruise periods. During each cruise period one or more ships were operating in the study area.

On completion of the field work the data seemed too numerous for a classical data report. Instead, it was decided to produce an 11-volume atlas of the results, with 5 volumes containing physical oceanographic and meteorological data from the principal participating ships, 5 volumes containing biological and nutrient chemistry data from the same ships, and 1 volume containing all data from Latin American cooperating ships and ships of opportunity. Extensive use was made of a computer and automatic plotter in preparation of the atlas charts. Methods used to collect and process the data upon which the atlas is based are described in detail by the contributors of the following categories of charts: temperature, salinity, and derived quantities; thickness of the upper mixed layer; dissolved oxygen; meteorology; nutrient chemistry; phytoplankton standing stocks and production; zooplankton and fish larvae; micronekton; birds, fish schools, and marine mammals.

- 330, Vol. 5. EASTROPAC Atlas: Physical Oceanographic and Meteorological Data from Principal Participating Ships, Second Survey Cruise, August-September 1967. By Cuthbert M. Love, (editor). September 1972, viii + 100 pp., 143 figures. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402—Price \$4.75 per volume.

ABSTRACT

This atlas contains charts depicting the distribution of physical, chemical, and biological oceanographic properties and associated meteorological properties observed during EASTROPAC. EASTROPAC was an international cooperative investigation of the eastern tropical Pacific Ocean (20° N. to 20° S., and from the west coasts of the American continents to 119° W.) which was intended to provide data necessary for a more effective use of the marine resources of the area, especially tropical tunas, and also to increase knowledge of the ocean circulation, air-sea interaction, and ecology. The Bureau of Commercial Fisheries (now National Marine Fisheries Service) was the coordinating agency. The field work, from February 1967 through March 1968, was divided into seven 2-month cruise periods. During each cruise period one or more ships were operating in the study area.

On completion of the field work the data seemed too numerous for a classical data report. Instead, it was decided to produce an 11-volume atlas of the results, with 5 volumes containing physical oceanographic and meteorological data from the principal participating ships, 5 volumes containing biological and nutrient chemistry data from the same ships, and 1 volume containing all data from Latin American cooperating ships and ships of opportunity.

Extensive use was made of a computer and automatic plotter in preparation of the atlas charts. Methods used to collect and process the data upon which the atlas is based are described in detail by the contributors of the following categories of charts: temperature, salinity, and derived quantities; thickness of the upper mixed layer; dissolved oxygen; meteorology; nutrient chemistry; phytoplankton standing stocks and production; zooplankton and fish larvae; micronekton; birds, fish schools, and marine mammals.

- 330, Vol. 6. EASTROPAC Atlas: Biological and Nutrient Chemistry Data from Principal Participating Ships, Second Survey Cruise, August-September 1967. By Cuthbert M. Love, (editor). December 1972, vi + 80 pp., 149 figures. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402—Price \$4.75 per volume.

ABSTRACT

This atlas contains charts depicting the distribution of physical, chemical, and biological oceanographic properties and associated meteorological properties observed during EASTROPAC. EASTROPAC was an international cooperative investigation of the eastern tropical Pacific Ocean (20° N. to 20° S., and from the west coasts of the American continents to 119° W.) which was intended to provide data necessary for a more effective use of the marine resources of the area, especially tropical tunas, and also to increase knowledge of the ocean circulation, air-sea interaction, and ecology. The Bureau of Commercial Fisheries (now National Marine Fisheries Service) was the coordinating agency. The field work, from February 1967 through March 1968, was divided into

seven 2-month cruise periods. During each cruise period one or more ships were operating in the study area.

On completion of the field work the data seemed too numerous for a classical data report. Instead, it was decided to produce an 11-volume atlas of the results, with 5 volumes containing physical oceanographic and meteorological data from the principal participating ships, 5 volumes containing biological and nutrient chemistry data from the same ships, and 1 volume containing all data from Latin American cooperating ships and ships of opportunity.

Extensive use was made of a computer and automatic plotter in preparation of the atlas charts. Methods used to collect and process the data upon which the atlas is based are described in detail by the contributors of the following categories of charts: temperature, salinity, and derived quantities; thickness of the upper mixed layer; dissolved oxygen; meteorology; nutrient chemistry; phytoplankton standing stocks and production; zooplankton and fish larvae; micronekton; birds, fish schools, and marine mammals.

331-356. Issued before 1972.

357-360. Not issued.

361. Issued in 1971.

NOAA Technical Report NMFS CIRC

362. Issued in 1971.

363. Not issued.

364. Issued in 1971.

365. Processing EASTROPAC STD data and the construction of vertical temperature and salinity sections by computer, by Forrest R. Miller and Kenneth A. Bliss. February 1972, iv + 17 pp., 7 figs., 3 app. figs. Price 30 cents.

ABSTRACT

During the EASTROPAC expeditions the STD (salinity-temperature-depth recorder) was established as the principal instrument for obtaining detailed profiles of temperature and salinity to depths exceeding 1,000 m. The STD system recorded data in digital form on magnetic tape and required computer processing to provide accurate temperatures and salinities at 1-m intervals for each hydrographic station.

Procedures for processing STD data from pre-processed digital logger tapes are described. Also included is a discussion of processing temperatures and salinities from the STD analog charts or from Nansen cast data when the digital data logger was not operative. Essential calibration of STD data from Nansen cast data is outlined. Finally, a computer program which prepares a final, annotated STD tape, with corrected temperatures and salinities, is described in great detail.

Another computer program was developed to provide for the EASTROPAC Atlas representative vertical sections of temperatures and salinities extracted from the final STD tapes. Details of computer methods employed to analyze and contour STD data are presented, and they are followed by an outline of computer requirements and program limitations.

The STD provided many detailed profiles along cruise tracks in the eastern tropical Pacific with an accuracy approaching those taken by Nansen casts; but it created new problems in processing oceanographic data. The problems, their solutions and recommendations are presented for those who may benefit from many months of processing STD data from EASTROPAC cruises.

366. Key to field identification of anadromous juvenile salmonids in the Pacific Northwest, by Robert J. McConnell and George R. Snyder. January 1972, iv + 6 pp., 3 figs. Price 20 cents, stock number 0320-0021.

ABSTRACT

A key is presented with descriptive illustrations to help in field identification of live, juvenile salmonids in fresh waters of the Pacific Northwest. Other juvenile fish that may be mistakenly identified as salmonids are included.

367. Engineering economic model for fish protein concentration processes, by K.K. Almenas, L.C. Durilla, R.C. Ernst, J.W. Gentry, M.B. Hale, and J.M. Marchello. October 1972, iii + 175 pp., 6 figs., 6 tables, 1 app. Price \$1.50.

ABSTRACT

A process engineering economic model for fish protein concentration processes has been developed. The model predicts the construction and operating costs for fish meal plants and for plants producing fish protein concentrate (FPC) by: isopropyl alcohol extraction, biological, and press cake-isopropyl alcohol extraction.

Typical process flow sheets and a computer program were developed to be used in the design and cost computations. The model provides for each process to be studied both internally and externally in comparison with alternate processes. The program and model were prepared in such a way that changes and updating may be accomplished quite readily as new information becomes available. This report contains directions for users and descriptions of the processes. A listing of the computer program and example calculations for each process are presented in the Appendix to guide the user and to illustrate the nature of the model output.

While the model does develop construction, operating, and production aspects of the processes, it does not deal with the economics of selling the products and the resulting profit and return on investment. Problems of allocation of costs and marketing arrangements are not covered in this report, but must be considered in the final decisions relating to a complete evaluation of alternatives.

368. Cooperative Gulf of Mexico estuarine inventory and study, Florida: Phase I, area description, by J. Kneeland McNulty, William N. Lindall, Jr., and James E. Sykes. November 1972, vii + 126 pp., 46 figures, 62 tables. Price \$1.25.

ABSTRACT

Newly-developed tables and maps depict the dimensions, submerged vegetation, tidal marshes, mangrove swamps, commercial oyster beds, leased oyster-rearing areas, sources of pollution, drained tidal marshes, and filled areas of Florida's west coast estuaries. Published and unpublished information on temperature, salinity, geology, artificial fishing reefs, stream discharge, human population, commercial fishing, and economic development is presented in new form.

If the total area of estuaries (3,003,312 acres = 1,215,440 ha) is considered to be the area of open water (2,081,525 acres = 842,393 ha) plus the area of mangrove swamps (393,160 acres = 159,112 ha) and tidal marshes (528,528 acres = 213,895 ha), then roughly one-half of the total area of estuaries is unvegetated; the remaining half is about equally divided among mangroves, tidal marshes, and submerged vegetation.

Human population in coastal counties increased from 614,616 persons in 1930 to 3,320,226 persons in 1970, resulting in adverse effects from pollution to 43 percent of estuarine areas, filling of 23,521 acres (9,519 ha) mainly for residential and industrial development, and draining of 26,676 acres (10,796 ha) of tidal marshes for mosquito control. Increasing population correlates directly with the number of sources of pollution, filled area, and the area closed to shellfishing by public health authorities; thus, failure to control the adverse effects of population growth will clearly result in continued rapid degradation of estuarine habitat on Florida's west coast.

369. Field guide to the angelfishes (Pomacanthidae) in the western Atlantic, by Henry A. Feddern. November 1972, 10 pp., 17 figs. Price 25 cents.

ABSTRACT

A key illustrated by photographs and brief descriptions is presented to aid in identifying the six species of angelfishes, family Pomacanthidae, found in the western Atlantic.

370. Collecting and processing data on fish eggs and larvae in the California Current region, by David Kramer, Mary J. Kalin, Elizabeth G. Stevens, James R. Thraillkill, and James R. Zweifel. November 1972, iv + 38 pp., 38 figs., 2 tables. Price 50 cents.

ABSTRACT

Descriptions are given for the methods used by the California Cooperative Oceanic Fisheries Investigations to collect and process plankton. These include details of the design of the station pattern in the survey area, the gear and methods used for plankton hauls, measuring plankton, and sorting plankton for fish eggs and larvae; some procedures for identifying fish eggs and larvae;

details of "hand" processing data for standardization of numbers of organisms collected in all plankton hauls; calibration of flowmeters; and some new procedures for automatic data processing.

371. To be published in 1973.

372. Fishery publications, calendar year 1971: Lists and indexes, by Thomas A. Manar. October 1972, iv + 24 pp., 1 fig. Price 30 cents.

ABSTRACT

The following series of fishery publications of the National Marine Fisheries Service, National Oceanic and Atmospheric Administration, in calendar year 1971 are listed numerically (with abstracts) and indexed by author, subject, and geographic area: NOAA Technical Report NMFS CIRC (formerly Circular); Data Report; Fishery Leaflet; and NOAA Technical Report NMFS SSRF (formerly Special Scientific Report—Fisheries.)

- 373-376. To be published in 1973.

377. Fishery publications, calendar year 1970: Lists and indexes, by Mary Ellen Engett and Lee C. Thorson. December 1972, iv + 34 pp., 1 fig.

ABSTRACT

The following series of fishery publications of the National Marine Fisheries Service, National Oceanic and Atmospheric Administration (until October, 1970 the Bureau of Commercial Fisheries of the U.S. Fish and Wildlife Service) in calendar year 1970 are listed numerically (with abstracts) and indexed by author, subject, and geographic area: Circular, Data Report, Fishery Industrial Research, Fishery Leaflet, and Special Scientific Report—Fisheries.

Data Report

(Hard copies of Data Reports Nos. 71, 73 and 74 are for sale at \$3.00 each and microfiche copies for 95 cents each. No. 72 is for sale at \$9.00 and microfiche copies for 95 cents each by the U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22151.)

71. Predation, particularly by sculpins, on salmon fry in fresh waters of Washington, by Benjamin G. Patten. 1972, 21 pp. on 1 microfiche.

ABSTRACT

Stomach contents of 41 species of fish collected in one lake and 11 streams were examined to determine species that were predaceous on wild and cultured Pacific salmon, *Oncorhynchus* spp., fry. The only species with salmon in their stomachs were: sculpins, *Cottus* spp., coho salmon, *O. kisutch*, chinook salmon, *O. tshawytscha*, rainbow trout, *Salmo gairdneri*, and cutthroat trout, *S. clarki*. Presented are data on the number of salmon in the stomachs of the predators and on body lengths of the predator species.

Fishery Facts

72. Surface zooplankton from Auke Bay and vicinity, southeastern Alaska, August 1962 to January 1964, by Bruce L. Wing and Gerald M. Reid. 1972, 765 pp. on 12 microfiche.

ABSTRACT

A list of species, counts of each species per cubic meter, and volumes of zooplankton per 1,000 cubic meters from the marine surface waters of Auke Bay and vicinity are presented for monthly samples collected from August 1962 to January 1964. Meteorology, oceanography, and cycles of phytoplankton productivity for Auke Bay are described. A table for converting from counts to volume, wet weight, and dry weight for selected species is abstracted from the literature.

73. Hydrographic observations in Tampa Bay, Florida—1969, by Carl H. Saloman and John L. Taylor. 1972, 82 pp. on 2 microfiche.

ABSTRACT

Hydrographic data include water temperature, salinity, total phosphorus, total Kjeldahl nitrogen, pH, dissolved oxygen, turbidity, water transparency, chlorophyll *a*, *b*, and *c*, astacin and nonastacin carotenoids, and primary productivity based on the chlorophyll *a* extraction. Methods of collecting and analyzing samples are described. Tables summarize data collected from 30 permanent stations by mean, range, and number of observations according to month and area. Other tables summarize the mean, range, and number of observations of samples taken daily at the Laboratory dock.

74. Collections of larval Gulf menhaden, *Brevoortia patronus*, from Galveston Entrance (1959-1969) and Sabine Pass (1963-1967), Texas, by Paul L. Fore and Kenneth N. Baxter. 1972, 17 pp. on 1 microfiche.

ABSTRACT

The number of larvae, that were taken per tow with a Renfro beam trawl, and the dates of collection are given for two Texas inlets.

75. To be published in 1973.

76. Amphipoda (Hyperideae) distribution and abundance off the coast of central west Africa, by Joseph E. Tashiro and Jack W. Jossi. 1972, 38 pp. on 1 microfiche.

ABSTRACT

Distribution and abundance of pelagic Amphipoda (Hyperideae) are charted for 53 station positions off the coast of central west Africa. Systematics, zooplankton haul data, and species counts are shown.

1. Redfish, by George F. Kelly, Paul M. Earl, John D. Kaylor, Fred E. Lux, Henry R. McAvoy, and Ernest D. McRae. October 1972, 18 pp., 14 figs. Price 25 cents.

(No abstract.)

2. Alaska's fishery resources the Pacific herring, by Gerald M. Reid, June 1972, iii + 20 pp., 13 figs. Price 25 cents.

ABSTRACT

The Pacific herring, *Clupea harengus pallasii*, is a valuable natural resource in the coastal waters of Alaska, not only because of its direct commercial significance but also because of its importance as a component in the complex food cycle of other commercially valuable fishes.

Pacific herring are indigenous to the North Pacific rim and are closely related to the herring of the Atlantic Ocean.

Pacific herring generally spawn on intertidal vegetation in the spring. Although spawning is often successful in terms of density, subsequent mortality of eggs and larvae may exceed 99%.

Herring abundance fluctuates greatly. Biologists have attempted to understand and predict fluctuations by studying changes in the numerical strength of different year classes of herring taken by the commercial fisheries.

Commercial utilization of Alaska herring by American fishermen is at a low level because of various socioeconomic problems, although thousands of tons were taken in the past for food and industrial products. Foreign vessels are currently taking large quantities of herring in areas offshore from Alaska. A viable herring industry might develop if an economical method of producing fish protein concentrate from fatty fish is developed. Alaska herring reduced to the concentrate could provide an essential item in the diet of peoples of protein-deficient nations.

3. Dungeness crab pots, by Fred W. Hipkins. June 1972, 13 pp., 10 figs. Price 25 cents.

(No abstract.)

4. Inshore lobster fishing, by John T. Everett. October 1972, iv + 26 pp., 21 figs. Price 25 cents.

ABSTRACT

This document describes the equipment and methods now being used in the inshore lobster (*Homarus americanus*) fishery along the United States North Atlantic Coast.

NOAA Technical Report NMFS SSRF

644. Inhibition of flesh browning and skin color fading in frozen fillets of yelloweye snapper (*Lutjanus vivanus*), by Harold C. Thompson, Jr. and Mary H. Thompson. February 1972, iii + 6 pp., 3 tables. Price 25 cents.

ABSTRACT

Fresh yelloweye snapper (*Lutjanus vivanus*) which had been scaled and eviscerated were treated with the chemicals 3,3'-thiodipropionic acid, glutathione, disodium ethylenediaminetetraacetate dihydrate in combination with propyl gallate, and monotertiary butylhydroquinone. The inhibitory effects that these chemicals had on flesh browning (Maillard reaction) were studied over a 12-month frozen storage period. Also, the effects of vacuum packaging on snapper skin discoloration were studied.

645. Published in 1971.

646. Published in 1971.

647. Revised annotated list of parasites from sea mammals caught off the west coast of North America, by L. Margolis and M.D. Dailey. March 1972, iii + 23 pp. Price 35 cents.

ABSTRACT

Parasite-host and host-parasite lists, with supporting references, of the ecto- and endoparasites recorded from marine mammals of the North American west coast are provided. Excluding records in which the parasites have not been identified to the species level, there are known 15 trematodes, 3 cestodes, 10 nematodes, 7 acanthocephalans, 1 copepod, and 10 amphipods from 22 species of Cetacea; 8 trematodes, 10 cestodes, 12 nematodes, 8 acanthocephalans, 5 anoplurans, and 5 acarines from 8 species of Pinnipedia; and 4 trematodes, 2 cestodes, 1 nematode, 3 acanthocephalans, and 1 acarine from a single species of Carnivora.

Previously unpublished records which are included here are: *Anisakis simplex* from *Phocoenoides dalli* and *Orcinus orca*, British Columbia; *Cyamus balaenopterae* from *Balaenoptera musculus*, California; *Cyamus erraticus* from *Balaena glacialis*, Alaska; *Cyamus scammoni* from *Eschrichtius gibbosus*, Alaska; *Antarctophthirus trichechi* from *Odobenus rosmarus*, Alaska.

648. Published in 1971.

649. Distribution of forage of skipjack tuna (*Euthynnus pelamis*) in the eastern tropical Pacific, by Maurice Blackburn and R. Michael Laurs. January 1972, iii + 16 pp., 7 figs., 3 tables. Price 30 cents, stock number 0320-0036.

ABSTRACT

The fishery for skipjack tuna (*Euthynnus pelamis*) in the eastern Pacific Ocean might be extended if offshore areas of high skipjack abundance were known. One would expect the numbers of skipjack in these offshore areas to be related to the distribution of known skipjack forage organisms in the micronekton. The EASTROPAC oceanographic cruises yielded net-caught micronekton samples over large parts of the eastern tropical Pacific during seven successive 2-month periods. From these samples,

the occurrence of organisms known to be prey of skipjack was expressed as ml/100 m³. Charts of night and day concentrations in the upper 200 m were produced for each of the seven periods.

In the region from long 92° to 119° W the major areas of maximum concentration of potential skipjack forage remained essentially constant during most periods. Two of these areas lie parallel to the equatorial upwelling zone—one just to the north, the other just to the south. Another zonal area of abundant forage generally occurs between lat 6° and 14° N. Forage is also frequently abundant between lat 14° and 20° N and long 107° and 119° W.

The concentrations of skipjack forage in these areas are comparable with those in nearshore parts of the eastern tropical Pacific, where the present skipjack fishery occurs. The abundance of skipjack in the forage-rich offshore areas might therefore be sufficient to support commercial fishing operations. Sea-surface temperatures are generally suitable for skipjack in those areas. On three recent crossings of the equatorial region at about long 119° W, skipjack appeared to be abundant in the first three areas of high forage concentration mentioned above.

650. Effects of some antioxidants and EDTA on the development of rancidity in Spanish mackerel (*Scomberomorus maculatus*) during frozen storage, by Robert N. Farragut. February 1972, iv + 12 pp., 6 figs., 12 tables. Price 25 cents, stock number 0320-0032.

ABSTRACT

Spanish mackerel (*Scomberomorus maculatus*) were treated with antioxidant solutions containing BHA and BHT (Tenox 4); BHA, BHT, PG, citric acid, and propylene glycol (Tenox 6); Tenox 4 plus EDTA; Tenox 6 plus EDTA; Ca(Na)₂EDTA; (Na)₂EDTA; (Ca)₂EDTA; and (Na)₄EDTA both by dipping and injecting methods. Samples analyzed at 3-month intervals showed fillets packed in vacuum and treated with EDTA remained in good condition over the 12-month storage period. However, samples treated with (Na)₄EDTA remained superior to other samples throughout the storage period.

651. The effect of premortem stress, holding temperatures, and freezing on the biochemistry and quality of skipjack tuna, by Ladell Crawford. April 1972, iii + 23 pp., 3 figs., 4 tables. Price 35 cents.

ABSTRACT

This experiment was designed to determine if there were differences (biochemical and/or organoleptic) before and after canning rested and stressed skipjack tuna. The live fish were captured off Oahu and were placed in shoreside tanks in Honolulu, Hawaii. After having been under observation for 24 hr, the fish were sacrificed in a rested or stressed condition. Stress was induced by forcing fish to swim around a tank until they showed signs of exhaustion. The rested fish were kept in a separate tank and were agitated as little as possible before being sacrificed.

Some of the sacrificed tuna were canned immediately to serve as controls. Others were held in 32°, 60°, and 78° F seawater (SW) for 6 hr, and some were held in 78° F SW for 9 hr before canning. An equal number of fish from all treatments were brine frozen (for 20 hr), then thawed and canned. Sample wedges were taken before canning for measurements of glycolytic and purine degradation products. These measurements together with organoleptic evaluation were also determined on the canned product.

There were no commercially discernible differences between rested and stressed skipjack subjected to various time-temperature treatments. The relation of the measured biochemical parameters to the treatment of the fish and the subsequent relation to the quality of the canned product were studied. There were not sufficiently defined relations on which to base quality predictions.

652. Indexed bibliography of the eggs and young of tunas and other scombrids (*Pisces, Scombridae*), 1880-1970, by William J. Richards and Witold L. Klawe. September 1972, iv + 107 pp., 1 table. Price \$1.00.

ABSTRACT

This bibliography enumerates reports on the early life history of tunas and other scombrid fishes published before 1971. All the entries are indexed, usually by species, but on occasion by a larger taxonomic unit, and within each taxonomic unit the entries are indexed by one or more subjects.

653. The use of electricity in conjunction with a 12.5-meter (headrope) Gulf-of-Mexico shrimp trawl in Lake Michigan, by James E. Ellis. March 1972, iv + 10 pp., 11 figs. 4 tables. Price 25 cents.

ABSTRACT

The catching efficiency of a 12.5-meter standard shrimp trawl and the same trawl fitted with three different electrode array systems with power on and power off was investigated.

The standard trawl caught 1.54 times or 54.2% more kilograms of fish than the electrode-equipped trawl with power off. The electrode array hanging across the mouth area of the trawl acted as a visual stimulant and thus reduced the trawl's catch rate.

Overall the electrical trawl with power on caught 1.19 times or 19.0% more kilograms of fish than the electrical trawl with power off. Array 2 with power on had the best catch rate—1.86 times or 86.9% more kilograms of fish than the power off catch rate. The avoidance of fish to an electrode array was more than offset with the catch rate of array 2 with power on. The dominance patterns of the catches with each system tested did not change significantly with the exception of chub catches with array 2 with power on.

Length selectivity was highly significant for chubs caught with arrays 2 and 3 with power on. No significant length selectivity occurred with the other species landed.

654. An electronic detector system for recovering internally tagged menhaden, Genus *brevoortia*, by R.O. Parker, Jr. February 1972, iii + 7 pp., 3 figs., 1 app. Price 25 cents.

ABSTRACT

Operation and results are described of an electronic detector-recovery system for fish with internal ferromagnetic tags. The system does not interfere with the operation of fish reduction plants. Date and location of recapture can be obtained since tagged fish are detected and recovered as they are landed. Growth rates of 3.1 and 7.2 mm per month were obtained for two menhaden out 130 and 483 days. Scale analyses supported annulus formation assumption. The best location for tag injection in adult menhaden appears to be about 13 mm above and just forward of the origin of the pelvic fin. Tag incisions were healed in 85% of the fish recaptured after 10 days.

655. Immobilization of fingerling salmon and trout by decompression, by Doyle F. Sutherland. March 1972, iii + 7 pp., 3 figs., 2 tables. Price 25 cents.

ABSTRACT

Laboratory experiments revealed that some chinook salmon (*Oncorhynchus tshawytscha*) and coho salmon (*O. kisutch*) fingerlings are immobilized when decompressed from atmospheric pressure to high negative pressures. About 69% of the chinook salmon were partially or completely immobilized at 71 cm of mercury vacuum, 48% at 61 cm, 17% at 51 cm, and none at 41 cm. The effects developed rapidly, peaking 5 min after exposure. The coho salmon, in comparison, were less severely affected. Only 9% were immobilized at 71 and 61 cm of mercury vacuum (reached 10 min after exposure), 2% at 51 cm (5 min after exposure), and none at 41 cm.

Studies (with coho salmon and rainbow trout [*Salmo gairdneri*] to determine the effects of decompression within a turbine of a dam did not provide conclusive results. However, some fingerlings caught in the tailrace immediately after their release in the turbine intake were immobilized. Exposure to negative pressure below turbine runner blades is one possible cause of immobilization.

656. The calico scallop, *Argopecten gibbus*, by Donald M. Allen and T.J. Costello. May 1972, iii + 19 pp., 9 figs., 1 table. Price 35 cents.

ABSTRACT

The calico scallop, *Argopecten gibbus*, supports a developing fishery off the southeastern coast of the United States and in the Gulf of Mexico. Information is given on classification, description, distribution, environment, relative abundance, reproduction, age and growth, mortality, associated organisms, behavior, and the fishery.

657. Making fish protein concentrates by enzymatic hydrolysis. A status report on research and some

processes and products studied by NMFS, by Malcolm B. Hale. November 1972, v + 32 pp., 15 figs., 17 tables, 1 app. table. Price 50 cents.

ABSTRACT

Research into biological methods for fish protein concentrate (FPC) preparation which has been carried out within the National Marine Fisheries Service is summarized. The effects of various processing conditions and commercially available proteolytic enzymes on yields and characteristics of water-soluble fish protein hydrolysates are presented. Soluble FPC prepared from red hake (*Urophycis chuss*) tended to be deficient in either tryptophan or histidine, depending on the pH of hydrolysis. Hydrolysis of raw fish with an alkaline protease of *Bacillus subtilis* at pH 8.5 or above gave the best balance of essential amino acids and a high yield of soluble product. Pancreatin also gave very good results at pH 8.5. The protein efficiency ratio (PER) of a totally soluble FPC prepared from alewife (*Alosa pseudoharengus*) was equivalent to that of casein. Soluble products prepared from hake were equivalent to casein as a wheat supplement but not as a sole source of protein. Process outlines and preliminary cost estimates are presented for the production of two types of fish protein hydrolysates. Possible food uses and the flavor problem are discussed. This report includes a literature survey of fish protein modifications by fermentation and selected chemical hydrolysis methods as well as by enzymatic hydrolysis processes.

658. List of fishes of Alaska and adjacent waters with a guide to some of their literature, by Jay C. Quast and Elizabeth L. Hall. July 1972, iv + 47 pp. Price 50 cents.

ABSTRACT

The authors list 432 species known to occur in Alaska waters, supplemented by 137 species that have been recorded from neighboring waters and, in the authors' opinion, should be considered when new collections are identified. Species entries are annotated to include common names, recorded range, useful references, localities represented by specimens in the collection of the Auke Bay Fisheries Laboratory, and comments on Taxonomy. Recorded geographic ranges are extended for 26 species: Ophidiidae—*Spectrunculus radcliffei*; Scorpaenidae—*Sebastes emphaeus*, *S. nigrocinctus*, *S. wilsoni*; Cottidae—*Eurymen gyrinus*, *Gymnocanthus detrisus*, *G. pistilliger*, *Hemilepidotus zaplus*, *Icelus spatula*, *I. uncinialis*, *Myoxocephalus jaok*, *Nautichthys pribilovius*, *Triglops scepticus*; Agonidae—*Agonopsis emmelane*, *Aspidophoroides bartoni*, *Ocella verrucosa*; Cyclopteridae—*Careproctus melanurus*, *C. rastrinus*, *Cyclopteroopsis phrynoides*, *Liparis bristolense*, *L. ochotensis*, *Paraliparis caudatus*, *P. deani*, *Temnocora candida*; Scombridae—*Thunnus thynnus*; Pleuronectidae—*Limanda proboscidea*.

659. The Southeast Fisheries Center bionumeric code. Part 1: Fishes, by Harvey R. Bullis, Jr., Richard B. Roe, and Judith C. Gatlin. July 1972, xl + 95 pp., 2 figs. Price \$1.25.

ABSTRACT

The Southeast Fisheries Center, Pascagoula, Mississippi Laboratory uses a nine-digit numeric code to catalog marine organisms. The basic code was adapted from the FAO taxonomic code developed in 1960, to which generic and specific levels were added and considerable classification modifications adopted.

This publication provides the code listing for fishes. Succeeding publications will deal with such groups as crustacea, mollusca, and various invertebrates.

660. A freshwater fish electro-motivator (FFEM)-its characteristics and operation, by James E. Ellis and Charles C. Hoopes. November 1972, iii + 11 pp., 9 figs.

ABSTRACT

A prototype Freshwater Fish Electro-Motivator (FFEM) system was developed as a research tool to test the application of electricity for use with active and passive fishing gear for increasing the gear's catching efficiency. The system's basic characteristics and operating modes are explained. The prototype system is extremely sophisticated, and its versatility permits single or multiple time-sequenced electrode loading and various pulse patterns, and allows duty cycles over a wide dynamic electrode load range. A summary of the field testing is discussed.

- 661-662. To be published in 1973.

663. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967, by Kenneth D. Waldron. December 1972, iii + 16 pp., 2 figs., 1 table, 4 apps. tables. Price 30 cents.

ABSTRACT

Fish larvae belonging to 24 families and the suborder Blennioidea were collected from Puget Sound and the Pacific Ocean off British Columbia, Washington, and Oregon during April and May 1967. All families and the Blennioidea were present in oceanic waters, but only 13 families and the Blennioidea were present in Puget Sound. The most abundant families in the oceanic area were Scorpaenidae, Myctophidae, and Pleuronectidae, whereas in Puget Sound the most abundant families were Gadidae, Pleuronectidae, and Scorpaenidae. Variations in composition and numbers of larvae in the catch were associated with area, water depth, water temperature, and time of day at which the collections were made.

664. Tagging and tag-recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*, by Richard L. Kroger and Robert L. Dryfoos. December 1972, iv + 11 pp., 4 figs., 12 tables.

ABSTRACT

Laboratory tagging experiments with adult and juvenile Atlantic menhaden were conducted at Beaufort, N.C., in 1965 and 1969. Tag-recovery experiments were done at menhaden processing plants at Beaufort, N.C. Internal

ferromagnetic body tags of appropriate sizes are suitable for tagging adults and juveniles, and the tags can be recovered effectively on magnets in the processing plants.

665. Larval fish survey of Humboldt Bay, California, by Maxwell B. Eldridge and Charles F. Bryan. December 1972, iii + 8 pp., 8 figs., 1 table. Price 25 cents.

ABSTRACT

As part of a series of investigations of the marine resources of Humboldt Bay, Calif., a larval fish survey was conducted from January to December 1969. Bottom and oblique tows were made at five sampling stations with 1-m plankton nets on alternate biweekly intervals. Thirty-seven species of larval and juvenile fishes representing 17 families were collected. In terms of larval abundance, the dominant fish was the bay goby, *Lepidogobius lepidus*, followed by Pacific herring (*Clupea harengus pallasii*), Pacific staghorn sculpin (*Leptocottus armatus*), longfin smelt (*Spirinchus thaleichthys*), and the arrow goby (*Clevelandia ios*). These five species constituted 95% of all larvae captured.

The number of larvae captured increased with increasing distance from the mouth of the Bay. The lowest number of species captured was at a station which experienced the widest range of salinities and temperatures. Peaks of seasonal abundance occurred in January and February and in April and May. Relatively few fish were captured after June. Some notable appearances of offshore spawned fishes were found in Humboldt Bay.

NOAA Technical Memorandum NMFS (Atlantic Estuarine Fisheries Center)

- AEFC-1. Report of the National Marine Fisheries Service Atlantic Estuarine Fisheries Center, fiscal years 1970 and 1971, by T.R. Rice, Director and Staff. June 1972, iv + 16 pp., 14 figs.

ABSTRACT

Estuarine and radioecological research conducted during Fiscal Years 1970 and 1971 was concerned with energy relations in ecosystems, distribution and cycling of radionuclides and trace metals, and environmental stress on the physiology of marine organisms. Research on the status of Atlantic and Gulf menhaden resources included monitoring the purse seine fishery, predicting future abundance, and describing the role of menhaden in the coastal environment. Other activities reported are thread herring and blue crab programs and radiological consulting. A list of professional staff and their scientific publications is included.

NOAA Technical Memorandum NMFS (Southeast Fisheries Center)

- SEFC-1. Report of the National Marine Fisheries Service Southeast Fisheries Center, Miami Laboratory, fiscal years 1970 and 1971, by Ann

Weeks and Albert C. Jones. July 1972, ii + 21 pp., 16 figs.

ABSTRACT

The research program of the National Marine Fisheries Service, Southeast Fisheries Center, Miami Laboratory (U.S. Department of Commerce, National Oceanic and Atmospheric Administration), Miami, FL, is described. Progress in investigations of the tropical Atlantic Ocean during Fiscal Years 1970 and 1971 is reviewed.

- SEFC-2. Report of the National Marine Fisheries Service Southeast Fisheries Center, Pascagoula Laboratory, fiscal years 1970 and 1971, by Edward F. Klima and Richard B. Roe. June 1972, iv + 21 pp., 15 figs.

ABSTRACT

The National Marine Fisheries Service Southeast Fisheries Center, Pascagoula Laboratory (formerly the NMFS Exploratory Fishing and Gear Research Base) conducted research in a wide range of activities during Fiscal Years 1970 and 1971. Investigations into the application of remote sensors for resource detection were advanced using aerial photography, pulsed lasers, spectrophotometry, and low-light-level imagery. This program received national status in September 1970 with the establishment of a National Marine Fisheries Service Remote Sensing Program (now the Southeast Fisheries Center, Mississippi Test Facility Engineering Laboratory) at the Mississippi Test Facility, Bay St. Louis, Mississippi.

Assessment surveys were conducted along the outer Continental Shelf and upper Continental Slopes of the Gulf of Mexico and Caribbean Sea where deepsea prawns, crabs, and silver hake were often taken in quantity. Benthic shelf explorations were greatly facilitated by the development of a remote controlled underwater fisheries assessment vehicle (RUFAS) used successfully in assessing, monitoring, and predicting the calico scallop resource off the eastern seaboard.

A budding fishery for swordfish in the Gulf of Mexico suffered an untimely death with the discovery of high mercury concentration in swordfish.

Hydroacoustical assessment of pelagic marine resources was bolstered with the acquisition of a signal processing computer-echosounder unit (SAS) which prints out real-time information on the location and relative size of underwater targets. The system is currently undergoing extensive field testing.

New approaches were taken to sampling and harvesting coastal pelagic fishes. Underwater lights and light arrays were successfully used to attract and lead schooling fish. Artificial structures of various design and complexity were found highly successful in attracting large quantities of pelagic fishes. These applied behavior studies will provide a prime component in a new concept in harvesting coastal pelagic fishes, an automated harvesting platform.

Electrical harvesting gear is being developed at Pascagoula to increase the efficiency of available gear and to provide the technology for sampling resources presently impossible to harvest. An electrical shrimp trawl has proven highly successful and an electrical midwater trawl

is under construction. A 120 kva pulse generator is currently under construction for application in the automated fishing platform, electrical fish trawls, and electrical rough-bottom shrimp trawls.

NOAA Technical Memorandum NMFS (Southeast Region)

SER-1. Report of the National Marine Fisheries Service Gulf Coastal Fisheries Center, fiscal years 1970 and 1971. Anonymous. July 1972, iii + 26 pp., 14 figs., 4 tables.

ABSTRACT

Progress is reported at the National Marine Fisheries Service Gulf Coastal Fisheries Center (formerly the Biological Laboratory, Galveston, Texas). Emphasis is placed on shrimp, and the research involves the fields of mariculture, population dynamics, ecology, and oceanography.

SER-2. Report of the National Marine Fisheries Service Biological Laboratory, St. Petersburg Beach, fiscal years 1970 and 1971, by James E. Sykes. July 1972, 13 pp., 8 figs.

ABSTRACT

Most of the major coastal and offshore fisheries of the United States depend upon species related to rearing and nursery areas in estuaries and the nearshore zone. To maintain and increase coastal shelf fisheries, it is necessary to provide continuing biological production near shore. Such provision requires a thorough ecological knowledge of the nursery and rearing area. Currently over 6,000 engineering proposals for estuarine areas are reviewed by Federal agencies each year. In view of relentless pressures affecting estuaries, this Laboratory works with other Federal agencies and the Gulf States to provide data directly applicable to the preservation, maintenance, and enhancement of nursery areas that generate valuable commercial and recreational species.

SER-3. Report of the National Marine Fisheries Service Fishery Products Technology Laboratory, Pascagoula, fiscal years 1970 and 1971, by Travis D. Love, Mary H. Thompson, and Melvin E. Waters. June 1972, iii + 12 pp., 7 figs., 4 tables.
(No abstract.)

AUTHOR INDEX

Allen, Donald M., and T.J. Costello, S 656
Almenas, K.K., L.C. Durilla, R.C. Ernst, J.W. Gentry, M.B. Hale, and J.M. Marchello, C 367
Anonymous, TM SER-1

Baxter, Kenneth N.—see Fore and Baxter
Blackburn, Maurice, and R. Michael Laurs, S 649

Bliss, Kenneth A.—see Miller and Bliss
Bryan, Charles F.—see Eldridge and Bryan
Bullis, Harvey R., Jr., Richard B. Roe, and Judith C. Gatlin, S 659

Costello, T.J.—see Allen and Costello
Crawford, Ladell, S 651

Dailey, M.D.—see Margolis and Dailey
Dryfoos, Robert L.—see Kroger and Dryfoos
Durilla, L.C.—see Almenas et al.

Earl, Paul M.—see Kelly et al.
Eldridge, Maxwell B., and Charles F. Bryan, S 665
Ellis, James E., S 653
———, and Charles C. Hoopes, S 660
Engett, Mary Ellen, and Lee C. Thorson, C 377
Ernst, R.C.—see Almenas et al.
Everett, John T., FF 4

Farragut, Robert N., S 650
Feddern, Henry A., C 369
Fore, Paul L., and Kenneth N. Baxter, D 74

Gatlin, Judith C.—see Bullis et al.
Gentry, J.W.—see Almenas et al.

Hale, M.B.—see Almenas et al.
Hale, Malcolm B., S 657
Hall, Elizabeth L.—see Quast and Hall
Hipkins, Fred W., FF 3
Hoopes, Charles C.—see Ellis and Hoopes

Jones, Albert C.—see Weeks and Jones
Jossi, Jack W.—see Tashiro and Jossi

Kalin, Mary J.—see Kramer et al.
Kaylor, John D.—see Kelly et al.
Kelly, George F., Paul M. Earl, John D. Kaylor, Fred E. Lux, Henry R. McAvoy, and Ernest D. McRae, FF 1
Klawe, Witold L. see Richards and Klawe
Klima, Edward F., and Richard B. Roe, TM SEFC-2
Kramer, David, Mary J. Kalin, Elizabeth G. Stevens, James R. Thraikill, and James R. Zweifel, C 370
Kroger, Richard L., and Robert L. Dryfoos, S 664

Laurs, R. Michael—see Blackburn and Laurs
Lindall, William N., Jr.—see McNulty et al.
Love, Cuthbert M. (editor), C 330, v. 1; C 330, v. 5; C 330, v. 6
Love, Travis D., Mary H. Thompson, and Melvin E. Waters, TM SER-3
Lux, Fred E.—see Kelly et al.

Manar, Thomas A., C 372
Marchello, J.M.—see Almenas et al.
Margolis, L., and M.D. Dailey, S 647
McAvoy, Henry R.—see Kelly et al.

McConnell, Robert J., and George R. Snyder, C 366
 McNulty, J. Kneeland, William N. Lindall, Jr., and James E. Sykes, C 368
 McRae, Ernest D.—see Kelly et al.
 Miller, Forrest R., and Kenneth A. Bliss, C 365

 Parker, R.O., Jr., S 654
 Patten, Benjamin G., D 71

 Quast, Jay C., and Elizabeth L. Hall, S 658

 Reid, Gerald M., FF 2
 ———see Wing and Reid
 Rice, T.R., TM AEFC-1
 Richards, William J., and Witold L. Klawe, S 652
 Roe, Richard B.—see Bullis et al.
 ———see Klima and Roe

 Saloman, Carl H., and John L. Taylor, D 73
 Snyder, George R.—see McConnell and Snyder
 Stevens, Elizabeth G.—see Kramer et al.
 Sutherland, Doyle F., S 655
 Sykes, James E., TM SER-2
 ———see McNulty et al.

 Tashiro, Joseph E., and Jack W. Jossi, D 76
 Taylor, John L.—see Saloman and Taylor
 Thompson, Harold C., Jr., and Mary H. Thompson, S 644
 Thompson, Mary H.—see Love et al.
 ———see Thompson and Thompson
 Thorson, Lee C.—see Engett and Thorson
 Thrailkill, James R.—see Kramer et al.

 Waldron, Kenneth D., S 663
 Waters, Melvin E.—see Love et al.
 Weeks, Ann, and Albert C. Jones, TM SEFC-1
 Wing, Bruce L., and Gerald M. Reid, D 72

 Zweifel, James R.—see Kramer et al.

SUBJECT INDEX

Acanthocybium solandri—see Wahoo

 Africa
 coast of central west
 distribution and abundance of Amphipoda, D 76

Alaminos—see Vessels

 Alaska
 Auke Bay, D 72
 coastal waters
 Pacific herring, FF 2

Albacore
 eggs and young, S 652

Albatross—see Vessels

Albatross IV—see Vessels

 Alewife
 protein efficiency from soluble FPC concentrate, S 657.

Allothunnus fallai—see Slender tuna

Alosa pseudoharengus—see Alewife

Alvera C—see Vessels

American Beauty—see Vessels

Ammodytes hexapterus—see Pacific sand lance

 Ammodytidae
 fish larvae collected from Puget Sound during April and May 1967, S 663

 Amphipoda
 coast of central west Africa
 distribution and abundance off, D 76

 Anchovy—see Northern anchovy

 Angelfishes
 field guide to, in western Atlantic Ocean
 key, C 369
 in western Atlantic Ocean
 blue angelfish, C 369
 french angelfish, C 369
 gray angelfish, C 369
 pigmy angelfish, C 369
 queen angelfish, C 369
 rock beauty, C 369

Argo—see Vessels

Argopecten gibbus—see Calico scallop

 Arrow goby
 Humboldt Bay, California
 larval fish survey, S 665

Atherinopsis californiensis—see Jacksmelt

 Atlantic bonito
 eggs and young, S 652

- Atlantic Estuarine Fisheries Center
 blue crab program, TM AEFC-1
 cycling of elements in the estuarine environment, TM AEFC-1
 ecology of estuarine ecosystems, TM AEFC-1
 menhaden research, TM AEFC-1
 radiological consulting, TM AEFC-1
 thread herring program, TM AEFC-1
- Atlantic mackerel
 eggs and young, S 652
- Atlantic menhaden
 tagging and tag-recovery experiments with
 experimental tag recovery, S 664
 experimental tagging, S 664
 methods of handling experimental fish and
 recovering tags, S 664
 review of menhaden tagging, S 664
- Atlantic Ocean
 western
 field guide to angelfishes, C 369
- Atlas
 EASTROPAC, first survey cruise Feb.-Mar. 1967
 physical oceanographic and meteorological data,
 C 330, v. 1
 EASTROPAC, second survey cruise Aug.-Sept. 1967
 biological and nutrient chemistry data, C 330, v. 6
 physical oceanographic and meteorological data,
 C 330, v. 5
- Auke Bay, Alaska
 surface zooplankton, D 72
- Auxis rochei*—see Bullet mackerel
- Auxis thazard*—see Frigate mackerel
- Bacillus subtilis*
 effect on hydrolysis of raw fish, S 657
- Bay goby
 Humboldt Bay, California
 larval fish survey, S 665
- Beaufort, North Carolina, S 654, S 664
- Benttooth bristlemouth
 Humboldt Bay, California
 larval fish survey, S 665
- Brevoortia tyrannus*—see Atlantic menhaden
- Bigeye tuna
 eggs and young, S 652
- Biological data
 EASTROPAC atlas
 from principal participating ships, second survey
 cruise, Aug.-Sept. 1967, C 330, v. 6
- Black skipjack
 eggs and young, S 652
- Blackfin tuna
 eggs and young, S 652
- Blennioidea
 fish larvae collected from the Northeastern Pacific
 Ocean and Puget Sound during April and May 1967, S
 663
- Blue angelfish
 in western Atlantic Ocean, C 369
- Blue lampfish
 Humboldt Bay, California
 larval fish survey, S 665
- Bluefin tuna
 eggs and young, S 652
- Bonito—see Atlantic bonito; Pacific bonito; Striped
 bonito
- George M. Bowers*—see Vessels
- Brevoortia patronus*—see Gulf menhaden
- Bristlemouth—see Benttooth bristlemouth
- Bullet mackerel
 eggs and young, S 652
- Carl Burton*—see Vessels
- Calamar*—see Vessels
- Calico scallop
 biology
 age and growth, S 656
 associated organisms, S 656
 behavior, S 656
 classification, S 656
 description, S 656
 distribution, S 656
 environment, S 656
 mortality, S 656
 relative abundance, S 656
 reproduction, S 656
 fishery
 Florida east coast, S 656

- general, S 656
- North Carolina, S 656
- Northeastern Gulf of Mexico, S 656
- California
 - Humboldt Bay, S 665
- California halibut
 - Humboldt Bay, California
 - larval fish survey, S 665
- Cancer magister*—see Dungeness crab
- Caribbean Sea
 - assessment survey
 - crabs, TM SEFC-2
 - prawns, TM SEFC-2
 - silver hake, TM SEFC-2
- Centropyge argi*—see Pigmy angelfish
- Cero
 - eggs and young, S 652
- Chinook salmon
 - immobilization of fingerlings by decompression, S 655
- Chub mackerel
 - eggs and young, S 652
- Citharichthys stigmaeus*—see Speckled sanddab
- Clevelandia ios*—see Arrow goby
- Clupea harengus pallasii*—see Pacific herring
- John N. Cobb*—see Vessels
- Coho salmon
 - immobilization of fingerlings by decompression, S 655
- Combat*—see Vessels
- Crab—see Dungeness crab
- Crabs
 - Gulf of Mexico and Caribbean Sea
 - assessment survey, TM SEFC-2
- Cybiosarda elegans*
 - eggs and young, S 652
- Cyclothone acclinidens*—see Benttooth bristlemouth
- Cynoscion nobilis*—see White seabass
- Charles H. Davis*—see Vessels
- Decompression
 - immobilization of fingerlings by, S 655
- Defiance*—see Vessels
- Delaware II*—see Vessels
- Dungeness crab
 - commercial fishing regulations, FF 3
 - pots
 - brief history, FF 3
 - construction, FF 3
 - equipment, FF 3
 - longline crab-pot system, FF 3
 - vessels, FF 3
 - state of fishery along west coast of United States, FF 3
- EASTROPAC
 - atlas from principal participating ships, Feb.-Mar. 1967
 - physical oceanographic and meteorological data, C 330, v. 1
 - atlas from principal participating ships, Aug.-Sept. 1967
 - biological and nutrient chemistry data, C 330, v. 6
 - physical oceanographic and meteorological data, C 330, v. 5
 - construction of vertical temperature and salinity sections by computer, C 365
 - STD data processing, C 365
- English sole
 - Humboldt Bay, California
 - larval fish survey, S 665
- Engraulis mordax*—see Northern anchovy
- Enzymatic hydrolysis
 - effects on yields and characteristics of water-soluble fish protein hydrolysates, S 657
- Esmeralda*—see Vessels
- Eucyclogobius newberryi*—see Tidewater goby
- Euthynnus affinis*—see Kawakawa
- Euthynnus alletteratus*—see Little tunny
- Euthynnus lineatus*—see Black skipjack
- Euthynnus pelamis*—see Skipjack tuna
- Fingerlings
 - immobilization by decompression, S 655
- Fish
 - Alaska and adjacent waters
 - Acipenseridae, S 658

- Agonidae, S 658
 Alepisauridae, S 658
 Alepocephalidae, S 658
 Aloiidae, S 658
 Ammodytidae, S 658
 Anarhichadidae, S 658
 Anoplogasteridae, S 658
 Anoplopomatidae, S 658
 Anotopteridae, S 658
 Argentinidae, S 658
 Aulorhynchidae, S 658
 Bathylagidae, S 658
 Bathymasteridae, S 658
 Batrachoididae, S 658
 Bothidae, S 658
 Bramidae, S 658
 Carangidae, S 658
 Carcharhinidae, S 658
 Caristiidae, S 658
 Catostomidae, S 658
 Centrolophidae, S 658
 Chauliodontidae, S 658
 Chimaeridae, S 658
 Clupeidae, S 658
 Cottidae, S 658
 Cyclopteridae, S 658
 Cyemidae, S 658
 Cyprinidae, S 658
 Dasyatidae, S 658
 Embiotocidae, S 658
 Engraulidae, S 658
 Esocidae, S 658
 Gadidae, S 658
 Gasterosteidae, S 658
 Gobiesocidae, S 658
 Gonostomatidae, S 658
 Hexagrammidae, S 658
 Hexanchidae, S 658
 Icosteidae, S 658
 Lamnidae, S 658
 Lampridae, S 658
 Macrouridae, S 658
 Melamphaeidae, S 658
 Melanostomiidae, S 658
 Molidae, S 658
 Moridae, S 658
 Myctophidae, S 658
 Myxinidae, S 658
 Nemichthyidae, S 658
 Notacanthidae, S 658
 Oneirodidae, S 658
 Ophidiidae, S 658
 Opisthoproctidae, S 658
 Osmeridae, S 658
 Paralepididae, S 658
 Pentacerotidae, S 658
 Percopsidae, S 658
 Petromyzontidae, S 658
 Pholididae, S 658
 Pleuronectidae, S 658
 Psychrolutidae, S 658
 Ptilichthyidae, S 658
 Rajidae, S 658
 Salmonidae, S 658
 Sciaenidae, S 658
 Scomberesocidae, S 658
 Scombridae, S 658
 Scoperlarchidae, S 658
 Scopelosauridae, S 658
 Scorpaenidae, S 658
 Scyliorhinidae, S 658
 Scytalinidae, S 658
 Serrivomeridae, S 658
 Sphyracidae, S 658
 Squalidae, S 658
 Squatinidae, S 658
 Sternoptychidae, S 658
 Stichaeidae, S 658
 Synphobranchidae, S 658
 Syngnathidae, S 658
 Tetragonuridae, S 658
 Torpedinidae, S 658
 Trachipteridae, S 658
 Trichiuridae, S 658
 Trichodontidae, S 658
 Umbridae (Dallidae), S 658
 Zaprionidae, S 658
 Zeidae, S 658
 Zoarcidae, S 658
- bionumeric code
 application, S 659
 description, S 659
 index to families, S 659
 index to genera, S 659
 listing format, S 659
- freshwater, electro-motivator
 power supply, S 660
 pulser, S 660
 summary of field testing, S 660
- Fish eggs
 collecting and processing data on, in California Current region
 data collection, C 370
 identification of eggs and larvae, C 370
 plankton sorting, C 370
 plankton volume determination, C 370
 survey pattern and area, C 370
- Fish larvae
 collected from the Northeastern Pacific Ocean and Puget Sound during Apr.-May 1967, S 663
 collecting and processing data on, in California Current region

- data collection, C 370
 - identification of eggs and larvae, C 370
 - plankton sorting, C 370
 - plankton volume determination, C 370
 - survey pattern and area, C 370
- Fish protein concentration processes
- enzymatic hydrolysis, S 657
 - biological methods by NMFS, S 657
 - engineering economic model for
 - biological process, C 367
 - computer user information, C 367
 - cost estimation, C 367
 - fish meal process, C 367
 - IPA extraction process, C 367
 - press cake-biological process, C 367
 - press cake-IPA extraction process, C 367
 - process economics, C 367
- Fish protein hydrolysates
- possible food uses and flavor problems, S 657
 - process outlines and preliminary cost estimates for production, S 657
- Fishery publications
- lists and indexes, calendar year
 - 1970, C 377
 - 1971, C 372
- Florida
- Gulf of Mexico estuarine inventory and study,
 - area description
 - artificial fishing reefs, C 368
 - coast, C 368
 - dimensions, C 368
 - dredging, C 368
 - economic development, C 368
 - geology, C 368
 - mangroves, C 368
 - measurement of vegetated areas, C 368
 - oysters and clams, C 368
 - pollution, C 368
 - population, C 368
 - salinity, C 368
 - stream discharge, C 368
 - submerged vegetation, C 368
 - tidal marshes, C 368
 - water temperature, C 368
 - Miami, TM SEFC-1
 - St. Petersburg Beach, TM SER-2
 - Tampa Bay, D 73
- Flounder—see Starry flounder
- Ben Franklin*—see Vessels
- Fregata*—see Vessels
- French angelfish
- in western Atlantic, C 369
- Frigate mackerel
- eggs and young, S 652
- Gadidae
- fish larvae collected from Puget Sound during April and May 1967, S 663
- Galveston Entrance, Texas, D 74
- Gear
- electrical harvesting gear being developed in Pascagoula, Mississippi, TM SEFC-2
- Geronimo*—see Vessels
- Charles H. Gilbert*—see Vessels
- Goa*—see Vessels
- Goby—see Arrow goby; Bay goby; Tidewater goby
- Grammatorcynus bicarinatus*
- eggs and young, S 652
- Gray angelfish
- in western Atlantic, C 369
- Greenling—see Kelp greenling
- Gulf Coastal Fisheries Center
- effects of physical, chemical, and biological environments, TM SER-1
 - environmental research, TM SER-1
 - estuarine ecology, TM SER-1
 - population monitoring, TM SER-1
 - shrimp culture, TM SER-1
- Gulf menhaden
- collections of larval, from
 - Galveston Entrance, Texas (1959-1969), D 74
 - Sabine Pass, Texas (1963-1967), D 74
- Gulf of Mexico
- assessment surveys
 - crabs, TM SEFC-2
 - prawns, TM SEFC-2
 - death of swordfish industry, TM SEFC-2
 - estuarine inventory and study
 - area description, Florida, C 368
 - silver hake, TM SEFC-2
- Gulf Ranger*—see Vessels
- Gulf Stream*—see Vessels

- Gymnosarda unicolor*
eggs and young, S 652
- Hake—see Red hake; Silver hake
- Halibut—see California halibut
- Herring—see Pacific herring
- Hexagrammos decagrammus*—see Kelp greenling
- Hidalgo*—see Vessels
- Holacanthus bermudensis*—see Blue angelfish
- Holacanthus ciliaris*—see Queen angelfish
- Holacanthus tricolor*—see Rock beauty
- Homarus americanus*—see Lobster
- Huayaipé*—see Vessels
- Humboldt Bay, California
larval fish survey
areal distribution, S 665
environment, S 665
seasonal distribution, S 665
species composition, S 665
- Hydrographic observations
Tampa Bay, Florida—1969
astacin and nonastacin carotenoids, D 73
chlorophyll *a*, *b*, and *c*, D 73
dissolved oxygen, D 73
pH, D 73
primary productivity, D 73
salinity, D 73
total Kjeldahl nitrogen, D 73
total phosphorus, D 73
turbidity, D 73
water temperature, D 73
water transparency, D 73
- Hydrolysis—see Enzymatic hydrolysis
- Hypomesus pretiosus*—see Surf smelt
- Internal tags
electric detector system for recovering,
menhaden, S 654
- Jacksmelt
Humboldt Bay, California
larval fish survey, S 665
- David Starr Jordan*—see Vessels
- Kaho*—see Vessels
- Katsuwonus pelamis*—see Skipjack tuna
- Kawakawa
eggs and young, S 652
- George B. Kelez*—see Vessels
- Kelp greenling
Humboldt Bay, California
larval fish survey, S 665
- King mackerel
eggs and young, S 652
- Lampfish—see Blue lampfish; Northern lampfish
- Lance—see Pacific sand lance
- Larvae, fish—see Fish larvae
- Lepidogobius lepidus*—see Bay goby
- Leptocottus armatus*—see Pacific staghorn sculpin
- Lingcod
Humboldt Bay, California
larval fish survey, S 665
- Little tunny
eggs and young, S 652
- Lobster
inshore fishing
bait, FF 4
boat and other equipment, FF 4
costs, FF 4
fishing methods, FF 4
pot construction, FF 4
regulations, FF 4
rope and buoys, FF 4
- Longfin smelt
Humboldt Bay, California
larval fish survey, S 665
- Lutjanus vivanus*—see Yelloweye snapper
- Mackerel—see Atlantic mackerel; Bullet mackerel;
Chub mackerel; Frigate mackerel; King mackerel;
Monterey Spanish mackerel; Spanish mackerel
- Menhaden—see Atlantic menhaden; Gulf menhaden

- Menhaden
 electric detector system for recovering
 internally tagged
 description, S 654
 operational problems, S 654
 recoveries, S 654
- Meteorological data
 EASTROPAC atlas
 from principal participating ships, first survey cruise,
 Feb.-Mar. 1967, C 330, v. 1
 from principal participating ships, second survey
 cruise, Aug.-Sept. 1967, C 330, v. 5
- Miami, Florida, TM SEFC-1
- Mississippi
 Pascagoula, TM SEFC-2, TM SER-3
- Monterey Spanish mackerel
 eggs and young, S 652
- Dan Moore*—see Vessels
- Murre II*—see Vessels
- Myctophidae
 fish larvae collected from the Northeastern Pacific
 Ocean during April and May 1967, S 663
- National Marine Fisheries Service
 Atlantic Estuarine Fisheries Center report for fiscal
 years 1970 and 1971, TM AEFC-1
 Gulf Coastal Fisheries Center report for fiscal years
 1970 and 1971, TM SER-1
 Miami Laboratory report for fiscal years 1970 and 1971,
 TM SEFC-1
 Pascagoula Fishery Products Technology Laboratory
 report for fiscal years 1970 and 1971, TM SER-3
 research and processes studied
 fish protein concentrates made by enzymatic
 hydrolysis, S 657
 St. Petersburg Beach Biological Laboratory report for
 fiscal years 1970 and 1971, TM SER-2
 Southeast Fisheries Center
 Pascagoula Laboratory report for fiscal years
 1970-71, TM SEFC-2
- Night smelt
 Humboldt Bay, California
 larval fish survey, S 665
- North Carolina
 Beaufort, S 654, S 664
- Northeastern Pacific Ocean
 fish larvae collected during April and May 1967, S 663
- Northern anchovy
 Humboldt Bay, California
 larval fish survey, S 665
- Northern lampfish
 Humboldt Bay, California
 larval fish survey, S 665
- Nutrient chemistry data
 EASTROPAC atlas
 from principal participating ships, second survey
 cruise, Aug.-Sept. 1967, C 330, v. 6
- Oceanographer*—see Vessels
- Odontopyxis trispinosa*—see Pigmy poacher
- Ogon*—see Vessels
- Oncorhynchus kisutch*—see Coho salmon
- Oncorhynchus* spp.—see Salmon
- Oncorhynchus tshawytscha*—see Chinook salmon
- Ophidon elongatus*—see Lingcod
- Orcynopsis unicolor*
 eggs and young, S 652
- Oregon*—see Vessels
- Oregon II*—see Vessels
- Pacific bonito
 eggs and young, S 652
- Pacific herring
 Alaska coastal waters
 biological characteristics, FF 2
 commercial fisheries for, FF 2
 distribution, FF 2
 herring fishery in perspective, FF 2
 life history, FF 2
 population dynamics, FF 2
 Auke Bay, Alaska
 kinds and qualities of zooplankton, D 72
 Humboldt Bay, California
 larval fish survey, S 665
- Pacific sand lance
 Humboldt Bay, California
 larval fish survey, S 665
- Pacific staghorn sculpin
 Humboldt Bay, California
 larval fish survey, S 665

- Pancreatin
effect on hydrolysis of raw fish, S 657
- Paralichthys californicus*—see California halibut
- Parasites
from sea mammals caught off North American west coast
Acanthocephala, S 647
Acarina, S 647
Amphipoda, S 647
Anoplura, S 647
Cestoda, S 647
Copepoda, S 647
host-parasite list, S 647
Nematoda, S 647
Trematoda, S 647
- Parophrys vetulus*—see English sole
- Pascagoula, Mississippi, TM SEFC-2, TM SER-3
- Pelican—see Vessels
- Perry Cubmarine—see Vessels
- Physical oceanographic data
EASTROPAC atlas
from principal participating ships, first survey cruise,
Feb.-Mar. 1967, C 330, v. 1
from principal participating ships, second survey
cruise, Aug.-Sept. 1967, C 330, v. 5
- Pigmy angelfish
in western Atlantic Ocean, C 369
- Pigmy poacher
Humboldt Bay, California
larval fish survey, S 665
- PLADS
application of remote sensors to fishery assessment,
TM SEFC-2
- Platichthys stellatus*—see Starry flounder
- Pleuronectidae
fish larvae collected from the Northeastern Pacific
Ocean and Puget Sound during April and May 1967, S
663
- Poacher—see Pigmy poacher; Pricklebreast poacher
- Pomacanthidae—see Angelfishes
- Pomacanthus arcuatus*—see Gray angelfish
- Pomacanthus paru*—see French angelfish
- Prawns
assessment survey in Gulf of Mexico and Caribbean
Sea, TM SEFC-2
- Pricklebreast poacher
Humboldt Bay, California
larval fish survey, S 665
- Psetichthys melanostictus*—see Sand sole
- Puget Sound
fish larvae collected during April and May 1967, S 663
- Queen angelfish
in western Atlantic Ocean, C 369
- Queen Mary*—see Vessels
- Rainbow trout
immobilization of fingerlings by decompression, S 655
- Rastrelliger brachysoma*
eggs and young, S 652
- Rastrelliger fanghni*
eggs and young, S 652
- Rastrelliger kanagurta*
eggs and young, S 652
- Red hake
FPC effect by hydrolysis, S 657
- Redfish
distribution, FF 1
fishing gear, FF 1
growth, FF 1
history of catch
Canada, FF 1
Germany, FF 1
United States, FF 1
market demand, FF 1
reproduction, FF 1
tagging, FF 1
time of catch, FF 1
- Rock beauty
in western Atlantic, C 369
- Rockaway*—see Vessels
- Rockfish
Humboldt Bay, California
larval fish survey, S 665

RUFUS

application to fishery assessment, TM SEFC-2

Sabine Pass, Texas, D 74

St. Petersburg Beach, Florida, TM SER-2

Salinity

construction of vertical sections by computer, C 365

processing EASTROPAC STD data, C 365

sections by computer, C 365

Salmo gairdneri—see Rainbow trout

Salmon—see Chinook salmon; Coho salmon

Salmon

key to field identification of juveniles in fresh waters of Pacific Northwest, C 366

predation on fry, particularly by sculpins, in the fresh waters of Washington, D 71

Sand sole

Humboldt Bay, California

larval fish survey, S 665

Sanddab—see Speckled sanddab

Sarda chiliensis—see Pacific bonito

Sarda orientalis—see Striped bonito

Sarda sarda—see Atlantic bonito

Scallop—see Calico scallop

Scomber australasicus

eggs and young, S 652

Scomber japonicus—see Chub mackerel

Scomber scombrus—see Atlantic mackerel

Scomberomorus cavalla—see King mackerel

Scomberomorus chinense

eggs and young, S 652

Scomberomorus commerson

Eggs and young, S 652

Scomberomorus concolor—see Monterey Spanish mackerel

Scomberomorus guttatus

eggs and young, S 652

Scomberomorus lineolatus

eggs and young, S. 652

Scomberomorus maculatus—see Spanish mackerel

Scomberomorus niphonius

eggs and young, S 652

Scomberomorus queenslandicus

eggs and young, S 652

Scomberomorus regalis—see Cero

Scomberomorus semifasciatus

eggs and young, S 652

Scomberomorus sierra—see Sierra

Scomberomorus sinensis

eggs and young, S 652

Scomberomorus tritor

eggs and young, S 652

Scombridae

indexed bibliography of eggs and young, 1880-1970

indexing system, S 652

journal titles with abbreviations, S 652

list of genera and species with geographical distribution, S 652

references consulted, S 652

subject index by taxa, S 652

Scorpaenidae

fish larvae collected from the Northeastern Pacific Ocean and Puget Sound during April and May 1967, S 663

Sculpin—see Pacific staghorn sculpin

Sculpins

predation on salmon fry in fresh waters of Washington, D 71

Seabass—see White seabass

Lois Seaver—see Vessels

Sebastes marinus—see Redfish

Sebastes spp.—see Rockfish

Sensors

application of remote sensors to fishery assessment, TM SEFC-2

- Shrimp trawl
 - use of electricity in conjunction with a
 - Gulf-of-Mexico
 - equipment, S 653
 - method of testing, S 653
 - physical quality of water, S 653
 - results of testing, S 653
 - study area, S 653
- Sierra
 - eggs and young, S 652
- Silver Bay*—see Vessels
- Silver hake
 - assessment survey in Gulf of Mexico and Caribbean Sea, TM SEFC-2
- Skipjack tuna
 - distribution of forage in eastern tropical Pacific
 - area east of long 92°W, S 649
 - area west of long 92°W, S 649
 - effect of premortem stress, holding temperatures, and freezing on biochemistry and quality of
 - chemistry of raw and canned tuna, S 651
 - effect of premortem stress and postmortem biochemical changes on quality of canned, S 651
 - effect of refrigeration on yield, S 651
 - glycolytic degradation products, S 651
 - organoleptic analysis, S 651
 - purines and purine derivatives, S 651
 - quality in tuna, S 651
 - refrigeration of tuna: chilling and freezing, S 651
 - relation of refrigeration experiments to present practices, S 651
 - eggs and young, S 652
- Slender tuna
 - eggs and young, S 652
- Smelt—see Longfin smelt; Night smelt; Surf smelt
- Snapper—see Yelloweye snapper
- Sole—see English sole; Sand sole
- Spanish mackerel
 - development of rancidity in, during frozen storage
 - effect of various EDTA compounds upon texture and, S 650
 - effect of various treatments upon, S 650
 - eggs and young, S 652
- Speckled sanddab
 - Humboldt Bay, California
 - larval fish survey, S 665
- Spirinchus starksi*—see Night smelt
- Spirinchus thaleichthys*—see Longfin smelt
- Starry flounder
 - Humboldt Bay, California
 - larval fish survey, S 665
- Stellerina xyosterna*—see Pricklebreast poacher
- Stenobranchius lencopsarns*—see Northern lampfish
- Striped bonito
 - eggs and young, S 652
- Surf smelt
 - Humboldt Bay, California
 - larval fish survey, S 665
- Swordfish
 - death of industry in Gulf of Mexico, TM SEFC-2
- Tags—see Internal tags
- Tampa Bay, Florida, D 73
- Tarltonbeania crenularis*—see Blue lampfish
- Te Vega*—see Vessels
- Temperature
 - construction of vertical sections by computer, C 365
 - processing EASTROPAC STD data, C 365
- Texas
 - Galveston Entrance, D 74
 - Sabine Pass, D 74
- Thunnus alahunga*—see Albacore
- Thunnus albacares*—see Yellowfin tuna
- Thunnus atlanticus*—see Blackfin tuna
- Thunnus maccoyii*
 - eggs and young, S 652
- Thunnus obesus*—see Bigeye tuna
- Thunnus thynnus*—see Bluefin tuna
- Thunnus tonggol*
 - eggs and young, S 652

- Tidewater goby
Humboldt Bay, California
larval fish survey, S 665
- Trawl—see Shrimp trawl
- Trout—see Rainbow trout
- Tuna—see Bigeye tuna; Blackfin tuna; Bluefin tuna;
Skipjack tuna; Slender tuna; Yellowfin tuna
- Tuxpan*—see Vessels
- Unanue*—see Vessels
- Undaunted*—see Vessels
- Undersea Hunter*—see Vessels
- Urophycis chuss*—see Red hake
- Vessels
Alaminos, C 330, v. 1
Albatross, D 72
Albatross IV, TM AEFC-1, TM SEFC-2
Alvera C, TM SEFC-2
American Beauty, S 651
Argo, C 330, v. 1
George M. Bowers, S 659, TM SEFC-1, TM SEFC-2
Carl Burton, TM AEFC-1
Calamar, TM SEFC-1, TM SEFC-2
John N. Cobb, S 663
Combat, S 659
Charles H. Davis, C 330, v. 1; C 330, v. 5; C 330, v. 6
Defiance, C 330, v. 1; C 330, v. 5; C 330, v. 6
Delaware II, TM SEFC-2
Esmeralda, C 330, v. 1; C 330, v. 5; C 330, v. 6
Ben Franklin, TM SEFC-2
Fregata, TM SEFC-2
Geronimo, TM SEFC-1, TM SER-1
Charles H. Gilbert, S 651
Goa, TM SEFC-1
Gulf Ranger, TM SEFC-2
Gulf Stream, TM SEFC-2
Hidalgo, TM SER-1
Huayaipé, C 330, v. 1; C 330, v. 5; C 330, v. 6
David Starr Jordan, C 330, v. 1; C 365, C 370
Kaho, S 653, S 660
George B. Kelez, S 663
Dan Moore, TM SEFC-2
Murre II, D 72
Oceanographer, C 330, v. 1; C 330, v. 5; C 330, v. 6
Ogon, C 370
Oregon, S 659
Oregon II, S 659, TM SEFC-2, TM SER-1, TM SER-3
Pelican, S 659
Perry Cubmarine, TM SEFC-2
Pioneer, TM SEFC-2
Queen Mary, TM SEFC-2
Rockaway, C 330, v. 1; TM SEFC-2
Lois Seaver, S 651
Silver Bay, S 659
Te Vega, C 330, v. 1; C 330, v. 5; C 330, v. 6
Tuxpan, C 330, v. 1; C 330, v. 5; C 330, v. 6
Unanue, C 330, v. 1; C 330, v. 5; C 330, v. 6
Undaunted, C 330, v. 1; D 76, TM SEFC-1
Undersea Hunter, TM SEFC-2
Thomas Washington, C 330, v. 1
Westport, S 651
Yelcho, C 330, v. 1; C 330, v. 5; C 330, v. 6
Yolanda, C 330, v. 1; C 330, v. 5; C 330 v. 6
- Wahoo
eggs and young, S 652
- Thomas Washington*—see Vessels
- Washington
predation, particularly by sculpins, on salmon fry in
fresh water, D 71
- Westport*—see Vessels
- White seabass
Humboldt Bay, California
larval fish survey, S 665
- Yelcho*—see Vessels
- Yelloweye snapper
inhibition of flesh browning and skin color fading in
frozen fillets
chemical methods, S 644
organoleptic evaluation of fillets, S 644
preparation of samples, S 644
- Yellowfin tuna
eggs and young, S 652
- Yolanda*—see Vessels
- Zooplankton
from marine surface waters of Auke Bay, Alaska, D 72

INDEX BY MARSDEN SQUARES

(see Figure 1)

001	042	082	151
TM SEFC-1	TM SEFC-1	D 74	FF 1
002	043	S 656	153
TM SEFC-1	S 656	TM SEFC-2	S 653
003	TM SEFC-2	TM SER-1	156
TM SEFC-1	044	TM SER-2	C 366
004	S 656	083	S 655
TM SEFC-1	TM SEFC-2	C 330, v. 1	157
005	045	C 330, v. 5	C 366
TM SEFC-1	C 330, v. 1	C 330, v. 6	C 370
006	C 330, v. 5	C 370	D 71
TM SEFC-1	C 330, v. 6	S 649	FF 3
008	S 649	084	S 663
C 330, v. 1	TM SEFC-2	C 330, v. 1	S 665
C 330, v. 5	046	C 330, v. 5	181
C 330, v. 6	C 330, v. 1	C 330, v. 6	FF 1
S 649	C 330, v. 5	C 370	183
009	C 330, v. 6	S 649	FF 1
C 330, v. 1	S 649	085	184
C 330, v. 5	S 656	C 330, v. 1	FF 1
C 330, v. 6	047	C 370	185
S 649	C 330, v. 1	S 649	FF 1
010	C 330, v. 5	086	186
C 330, v. 1	C 330, v. 6	C 370	FF 1
C 330, v. 5	S 649	115	187
C 330, v. 6	048	FF 1	FF 1
S 649	C 330, v. 1	116	193
011	C 330, v. 5	S 654	D 72
C 330, v. 1	C 330, v. 6	S 656	FF 2
C 330, v. 5	C 365	S 664	FF 3
C 330, v. 6	S 649	TM AEFC-1	S 663
S 649	049	117	194
012	C 330, v. 1	C 368	FF 2
C 330, v. 1	C 330, v. 5	S 656	FF 3
C 330, v. 5	C 330, v. 6	TM SER-1	216
C 330, v. 6	C 365	TM SER-2	FF 1
C 365	S 649	TM SER-3	217
S 649	079	118	FF 1
013	S 656	TM SER-2	218
C 330, v. 1	080	120	FF 1
S 649	C 368	C 370	219
036	S 656	121	FF 1
TM SEFC-1	081	C 370	220
038	C 368	FF 3	FF 1
TM SEFC-1	D 73	122	221
039	S 656	C 370	FF 1
TM SEFC-1	TM SEFC-1	149	222
040	TM SEFC-2	FF 1	FF 1
TM SEFC-1	TM SER-1	150	223
041	TM SER-2	FF 1	FF 1
TM SEFC-1			

248	307	334	346
FF 1	C 330, v. 1	D 76	C 330, v. 1
249	C 330, v. 5	TM SEFC-1	S 649
FF 1	C 330, v. 6	335	347
251	S 649	D 76	C 330, v. 1
FF 1	308	TM SEFC-1	S 649
252	C 330, v. 1	336	348
FF 1	C 330, v. 5	TM SEFC-1	C 330, v. 1
284	C 330, v. 6	337	S 649
FF 1	S 649	TM SEFC-1	370
285	309	338	D 76
FF 1	C 330, v. 1	TM SEFC-1	371
286	C 330, v. 5	339	TM SEFC-1
FF 1	C 330, v. 6	TM SEFC-1	379
287	S 649	343	C 330, v. 1
FF 1	310	C 330, v. 1	C 330, v. 5
288	C 330, v. 1	C 330, v. 5	C 330, v. 6
FF 1	C 330, v. 5	C 330, v. 6	380
300	C 330, v. 6	S 649	C 330, v. 1
TM SEFC-1	S 649	344	C 330, v. 5
301	311	C 330, v. 1	C 330, v. 6
TM SEFC-1	C 330, v. 1	C 330, v. 5	415
302	C 330, v. 5	C 330, v. 6	C 330, v. 1
TM SEFC-1	C 330, v. 6	S 649	C 330, v. 5
303	C 365	345	C 330, v. 6
TM SEFC-1	S 649	C 330, v. 1	416
304	312	C 330, v. 5	C 330, v. 1
TM SEFC-1	C 330, v. 1	C 330, v. 6	C 330, v. 5
305	S 649	S 649	C 330, v. 6
TM SEFC-1			

349. Use of abstracts and summaries as communication devices in technical articles. By F. Bruce Sanford. February 1971, iii + 11 pp., 1 fig.
350. Research in fiscal year 1969 at the Bureau of Commercial Fisheries Biological Laboratory, Beaufort, N.C. By the Laboratory staff. November 1970, ii + 49 pp., 21 figs., 17 tables.
351. Bureau of Commercial Fisheries Exploratory Fishing and Gear Research Base, Pascagoula, Mississippi, July 1, 1967 to June 30, 1969. By Harvey R. Bullis, Jr., and John R. Thompson. November 1970, iv + 29 pp., 29 figs., 1 table.
352. Upstream passage of anadromous fish through navigation locks and use of the stream for spawning and nursery habitat, Cape Fear River, N.C., 1962-66. By Paul R. Nichols and Darrell E. Louder. October 1970, iv + 12 pp., 9 figs., 4 tables.
356. Floating laboratory for study of aquatic organisms and their environment. By George R. Snyder, Theodore H. Blahm, and Robert J. McConnell. May 1971, iii + 16 pp., 11 figs.
361. Regional and other related aspects of shellfish consumption — some preliminary findings from the 1969 Consumer Panel Survey. By Morton M. Miller and Darrel A. Nash. June 1971, iv + 18 pp., 19 figs., 3 tables, 10 apps.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
ROOM 450
1107 N.E. 45TH ST.
SEATTLE, WA 98105
OFFICIAL BUSINESS

FOURTH CLASS