**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1996

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☐ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☒ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☒ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x		14x		18x		22x		26x		30x	
							/				
	12x		16x		20x		24x		28x		32x

The copy filmed here has been reproduced thanks to the generosity of:

National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

1	2	3
---	---	---

1	2
4	5

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec la plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

MANITOBA AGRICULTURAL COLLEGE

Some Forage Crops for Manitoba

Part I.—Summer Forage—Especially for Dairy Cows

Part II.—Solling and Pasture Crops for Hogs.

T. J. HARRISON, B. S. A.
Professor of Field Husbandry,
Manitoba Agricultural College.
WINNIPEG, CANADA

10

PART I.

Summer Forage - Especially for Dairy Cows.

Crops Discussed:

1. Oats.	5. Rape.
2. Oats and Peas or Oats and Vetch.	6. Field Roots.
3. Corn.	7. Millets.
4. Spring Rye.	8. Winter Rye.

Throughout Western Canada the farmers are being asked to increase the number of live stock on the farms and to produce larger quantities of dairy products. If this is to be done, the summer feed that is used for live stock must be increased in quantity and improved in quality. The only means of accomplishing this at the present time is by planting annual crops that can be used either for pasture or soiling purposes.

WHY ANNUAL FORAGE CROPS SHOULD BE PLANTED.

1. To supply feed that will keep up the milk flow in August and September.
2. To provide feed when the regular pastures have winter killed.
3. To help feed the extra stock that was purchased in the winter or spring.
4. To assist in keeping down the weeds in fields that cannot be summer-fallowed.

REQUIREMENTS OF A GOOD ANNUAL FORAGE CROP.

Any crop that fulfils the following requirements can be used for annual pasture or soiling purposes:

1. It must thrive on poor soil.
2. The seed must be cheap and easily procured.
3. The plants must grow readily.
4. The crop must not be easily killed by grazing.
5. The crop must be readily eaten by the stock and not injure the flavor of the butter or cheese.
6. The soil should be left in good condition for the succeeding crop of which

CROPS THAT CAN BE USED.

The following crops fulfil most of the above requirements:

1. Oats.	5. Rape or Thousand-headed kale.
2. Oats and peas, or oats and vetch.	6. Field roots.
3. Corn.	7. Millets.
4. Spring rye.	8. Winter rye.

OATS.

The use of oats for this purpose is to be recommended because the seed is easily procured and it produces an abundance of palatable food.

Kind of Land.—Oats can be sown on summerfallow or corn land, but where this land can be sown to wheat it will be considered better farm practice to put the oats on stubble land. If there is a field that is badly infested with wild oats, it could be sown to advantage with oats for the crop will either be cut or pastured down before the wild oats have produced seed.

Date of Planting.—From May 10th to July 10th is the period during which oats can be sown for this purpose. Successive seedings about two weeks apart should be made. This will insure pasture or soiling crops from July 15th until the first killing frost.

Rate of Seeding.—If the Banner type of oats is used, sow two bushels per acre. If the Abundance type is selected, heavier seeding is recommended.

Method of Planting.—Use the ordinary grain drill and sow the seed three inches deep.

OATS AND PEAS OR OATS AND VETCH.

The introduction of peas or vetch into the mixture will greatly improve the feeding qualities, but will increase the cost of seeding. If it is desired, the following mixture gives good satisfaction:

(1) Banner oats, two bushels, and Canadian Field Pea, one bushel; or

(2) Banner oats, two bushels, and Common Vetch, forty pounds.

The date and method of seeding will be similar to that outlined for oats.

CORN.

Corn has not been used to a very large extent either for soiling or pasture purposes. It can be used for soiling to better advantage than for pasture. When utilized in this manner, Northwestern Dent, Minnesota No. 13 and Longfellow will prove to be the better varieties. If it is to be used as pasture, Improved Squaw, Gehu and some of the other early maturing varieties can be used.

Kind of Land.—Corn should be planted on land which was intended for summerfallow. It has been found that land that has been well cultivated with corn has given better yields of wheat the following year than the bare summerfallow. It can, therefore, be used as a substitute for the bare fallow.

Date of Planting.—If it is to be used as a soiling crop, the seed should be sown about May 24th. If it is to be used as pasture and the early maturing varieties planted, the planting should be done between May 24th and June 29th.

Method of Planting.—The seed can be planted with an ordinary grain drill by closing up the spouts in the bottom of the box, allowing one to run every 36 inches apart. This plants the corn in rows three feet apart. Where the soil is at all weedy, it will be found much cheaper in cultivating to plant the corn with the corn planter, checking the corn in hills three feet apart each way.

Rate of Planting.—When the corn is planted with the grain drill, it will require about 30 pounds of seed per acre. If it is planted in checked rows, it will require about 15 pounds of seed.

Cultivating.—After the corn is planted it should be harrowed three or four times, or until the plant is about 5 or 6 inches high, after which, if the weeds are bad, it will require intertilling.

SPRING RYE.

Spring rye is a crop that can be used to good advantage either for summer pasture or for soiling purposes because it will grow and thrive on light, sandy soil or soil in poor physical condition.

Kind of Land.—Spring rye will give best results on land in good condition, but since it can be grown on exhausted, sandy soils, this will be found the most profitable place to sow it. Any field that should be summerfallowed, but cannot be, could be utilized for this purpose.

Date of Planting.—The seed should be sown sometime between the 1st of May and 10th of July. If successive seedings similar to that recommended for oats are made, it will insure pasture or soiling crops from midsummer until the first killing frost.

Rate of Seeding.—The seed should be sown about $1\frac{1}{2}$ bushels per acre.

Method of Seeding.—Sow with an ordinary grain drill, running the disks or shoes about $2\frac{1}{2}$ inches deep.

RAPE OR THOUSAND-HEADED KALE.

Rape or Thousand-Headed Kale cannot be used to advantage as summer forage for dairy cows because it will have a tendency to taint the milk. It can be used to good advantage as feed for young dairy stock or beef cattle. It will probably be found more useful for late fall pasture than for early fall or late summer pasture because it can withstand considerable frost and still remain green.

Kind of Land.—These crops can be sown on land which has been treated as summerfallow until fairly late in the summer, but will, however, thrive on any rich, well cultivated piece of land.

Date of Planting.—If the crop is to be used for early fodder, early planting is necessary. If it is planted sometime between 10th and 15th of May it will produce pasture in July. As it is recommended for late fall pasture, it could be sown to advantage on land which had been treated as summerfallow until about July 10th, when the seed should be sown.

Method of Planting.—There are two methods of planting that can be used to advantage, depending largely upon the weeds. If the land is free from weeds, the seed can be sown broadcast, using about 6 pounds per acre. If the soil happens to be weedy, better results will be obtained by sowing in drills. The grain drill can be fitted up as for sowing corn and the seed sown in drills 36 inches apart.

Cultivation.—If the crop is sown in drills, it can be cultivated with the corn cultivator to keep down the weeds.

FIELD ROOTS.

Mangels and turnips are the two root crops that can be used for summer and fall forage. If it is dairy cows that the food is being raised for, mangels would prove more satisfactory than turnips, because they would not taint the milk.

Kind of Land.—For both these crops it will be found advantageous to plant the seed on land which has been summerfallowed. This insures a quick growth and a small amount of weeds.

Date of Planting.—The seed should be planted early if fall feed is desired. Some date between the 1st and 24th of May gives best results. If turnips are grown, later seeding can be practised. In favorable seasons seed sown as late as July 1st has produced good fall forage.

Rate of Seeding.—It will require about 4 pounds of turnip seed per acre and 6 to 8 pounds of mangel seed per acre.

Method of Planting.—The seed can be sown with an ordinary grain drill, closing up the spouts similar to that recommended for corn. This will put the roots in rows 36 inches apart. If turnips are sown late for late pasture, they can be sown broadcast and harrowed in.

Method of Feeding.—The early sown roots can be used for soiling and the late sown turnips for pasture.

MILLETS.

The millets have not proven very satisfactory as an annual pasture crop. They are sometimes, however, grown for soiling purposes to advantage.

Kind of Land.—The soil for millet should be rich, warm and moist. Summerfallow or corn land would give best results, but it is not practicable to use this kind of land for forage crops. It could be grown more profitably on first year spring plowed stubble. The land should be plowed, harrowed, packed and the seed sown the same day as plowed so that the soil will not dry out and the seed be slow in germinating.

Date of Planting.—The millets are warm climate plants; consequently, they should not be sown until the soil has become thoroughly warm. Best results are obtained from millets planted about June 10th.

Rate of Seeding.—The seed should be sown at the rate of 26 pounds per acre.

Method of Planting.—The best method of sowing is to use an ordinary grain drill, sowing the seed in every cup the same as wheat or oats and running the disks or shoes about 1 to 1½ inches deep.

WINTER RYE.

Winter rye makes the best early pasture or soiling crop. It can be pastured late in the fall and early in the spring before the other crops have started to grow.

Kind of Land.—Winter rye can be sown to advantage only on land which has been summerfallowed until about August 20th. It is not possible under ordinary conditions in Manitoba to take off a crop of grain and re-sow with winter rye.

Date of Planting.—Winter rye should be sown sometime between August 20th and September 1st.

Rate of Planting.—To get best results for pasture or soiling purposes, the seed should be sown 1½ bushels per acre.

Method of Planting.—The ordinary grain drill gives best satisfaction. If the seed is sown 2½ to 3 inches deep, the young rootlets will not be greatly injured by the tramping of the stock.

Soiling and Pasture Crops for Hogs.

It has been demonstrated by the Agricultural College, Experimental Farms and prominent hog raisers that hogs cannot be raised economically unless a large portion of their summer rations are composed of green crops. Whether these crops will be pastured or cut and fed green in the piggery will depend largely on the location of the hog pen and the fields where these crops can be grown. Where the piggery is convenient to the fields, or where hog cots are used, by far the cheapest method is to let the hogs pasture on the different crops. Where it is not convenient to arrange for this, it will be found profitable to soil the crops, that is, cut them and feed them to the hogs in the piggery.

ROTATION.

If the pasture is to be worked out on the most economical lines, a rotation should be arranged. The fields need not be very large; this will depend altogether on the number of hogs to be kept. For the average farmer the fields may be from one-half to one acre in size. The following crops have given good satisfaction and should be included in the hog pasture rotation: Alfalfa, oats and peas and rape. As alfalfa makes such a good pasture, it is advisable to have a large quantity of this, so that the main field should be divided into about five one-half or one acre plots with alfalfa growing on two of them, rape on one, oats and peas on another and the fifth one either summerfallowed or planted with Squaw corn. The arrangement of the rotation could be as follows:

- 1st year—Rape.
- 2nd year—Oats and peas.
- 3rd year—Squaw corn or sumb fallow.
- 4th year—Alfalfa.
- 5th year—Alfalfa.

RAPE.

The rape will follow the alfalfa. The land for this could be either plowed early in the fall if the alfalfa pasture was not required, or it could be left until spring and plowed in the spring. The plowing should be about 4 to 5 inches deep and the land cultivated with the disc and drag harrow to work up a good seed bed. The seed should be sown as soon as after the spring plowing as possible. To make a continuous pasture during the summer, different portions of the field could be sown at different dates. The rape is usually ready for pasture about five weeks after planting. It can be planted by two different methods—broadcast or in drills. If sown broadcast it will require from 4 to 6 pounds of seed per acre; if sown in drills from 3 to 4 pounds will be sufficient. Under most conditions sowing in drills is preferable because in this way weeds can be kept under con-

trol and the intertillage given will conserve the moisture and cause a better growth. After the rape is about 8 to 10 inches high, the hogs should be let into the field because if it becomes rank they do not care for it.

OATS AND PEAS.

The oats and peas will be sown on the field that was in rape the previous year. If the rape has been pastured off early, it might be fall plowed for the next crop but, if not, spring plowing will give good results. If fall plowed, the land should be cultivated from time to time in the spring to keep the weeds under control and conserve the moisture. The oats and peas should not be sown before the 15th of May, but can be sown at later dates if necessary to give continuous pasture throughout the season. They should be sown about two bushels of oats to one of peas. If there is sufficient green manure, the oats and peas may be allowed to ripen and hogged off, but if the green feed is short this crop may be used also as pasture.

CORN OR SUMMERFALLOW.

When Improved Squaw or Gehr corn can be obtained for seed, it would be advisable to use this instead of the bare fallow. The corn can be check rowed and given cultivation both ways, thus keeping down the weeds. After the seed has been matured, the hogs may be let in and the corn hogged off.

ALFALFA.

In this rotation alfalfa is sown on the summerfallow or corn land. If summerfallow is used, the crop should be sown similar to that outlined for the production of fodder in *Circular No. 18 of the Manitoba

*The leading points in connection with sowing alfalfa, as set forth in Circular No. 18 (copy of which can be had free by writing the Extension Department, Manitoba Agricultural College) are as follows:

1. Unless the alfalfa bacteria are known to be present in the soil, they should be introduced, either by scattering soil from an old alfalfa field (100 pounds per acre), or by using laboratory cultures containing alfalfa bacteria. Bottles of culture may be obtained from Manitoba Agricultural College (Bacteriological Department), enough culture to inoculate 60 pounds of seed for 25 cents. Full instructions accompany the bottle.
2. Grimm and Baltic are the hardiest alfalfa varieties with Turkestan also giving satisfactory results in most cases.
3. Good seed should be sown at the rate of 12 pounds per acre; 8 or 10 pounds in very dry districts.
4. Sow from May 15th to June 15.
5. The best plan is to sow with a grain drill, running one inch deep. Mix the seed with twice the quantity of coarsely cracked wheat or barley from which the fine flour has been blown with a fanning mill. If the seeder is set to sow two pecks of wheat, it will put in about 12 pounds of alfalfa seed.
6. Sow without a nurse crop. If soil drifting is feared, a bushel of oats per acre may be sown, but the oats should be clipped with a mower as soon as the alfalfa plants are established.
7. Clip the field with a mower when the alfalfa and weeds are about 6 inches high, and leave the clippings on the field. Have the cutting bar set high enough not to injure the crown of the plant. Clip two or three times during the summer, the last time not later than August 15th.

Agricultural College. If it is sown on the corn land, this should be well disced in the spring to work up a good, loose mulch on the surface and to kill the weeds and the seed sown according to the bulletin mentioned above. While it would not be advisable to pasture this closely in the fall, if the hogs' noses were well ringed to prevent rooting, pasturing lightly would not hurt it to any great extent. The second season, however, it could be used quite extensively.

CONCLUSION.

With the above crops and arranging the dates of seeding, a continuous pasture throughout the whole summer may be obtained which will greatly reduce the cost of producing pork. While the pasture will help materially in feeding the hogs, it will be necessary to give them some grain ration. If it is not possible to arrange the fields conveniently for pasture, the different crops may be cut green and fed in the piggery.