**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☒ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couvertur: 3 endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

**This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.**

10x			14x			18x			22x			26x			30x		
													✓				
12x			16x			20x			24x			28x			32x		

The copy filmed here has been reproduced thanks to the generosity of:

Stauffer Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\longrightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Stauffer Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\longrightarrow$ signifie "À SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME VIII: MOLLUSKS, ECHINODERMS,
COELENTERATES, ETC.

PART E: ROTATORIA

By H. K. HARRING

SOUTHERN PARTY--1913-1916

OTTAWA
F. A. ACLAND
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1921

Issued December 31, 1921.

G
670
1913
D2t
v.008
pt.E
c.3

Report of the Canadian Arctic Expedition, 1913-18.

VOLUME I: NARRATIVE OF THE EXPEDITION

- Part A. NORTHERN PARTY, 1913-16. By Vilhjalmur Stefansson. *In preparation.*
 Part B. SOUTHERN PARTY, 1913-16. By Rudolph Martin Anderson. *In preparation.*

VOLUME II: MAMMALS AND BIRDS

- Part A. MAMMALS OF WESTERN ARCTIC AMERICA. By Rudolph Martin Anderson. *(In preparation).*
 Part B. BIRDS OF WESTERN ARCTIC AMERICA. By R. M. Anderson and P. A. Taverner. *(In preparation).*

VOLUME III: INSECTS

- INTRODUCTION. By C. Gordon Hewitt. *(Issued December 10, 1920).*
 Part A: COLEMBOLA. By J. L. V. Upton. *(Issued July 10, 1919).*
 Part B: NEUROPTEROID INSECTS. By Nathan Banks. *(Issued July 11, 1919).*
 Part C: DIPTERA.
 Craneflies. By Charles P. Alexander.
 Mosquitoes. By Harrison G. Dyar.
 Diptera (excluding Tephritidae and Cypselids). By J. R. Malloch. *(Issued July 14, 1919).*
 Part D: MALLOPHAGA AND ANOPLURA.
 Mallophaga. By A. W. Baker.
 Anoplura. By G. F. Ferris and G. H. F. Nuttall. *(Issued September 12, 1919).*
 Part E: COLEOPTERA.
 Forest Insects, including Ipsidae, Cerambycidae, and Buprestidae. By J. M. Swaine.
 Curculionidae and Silphidae. By H. C. Fall.
 Cerambycidae, Elateridae, Chrysomelidae and Rhynchophora. (Excluding Ipsidae).
 By C. W. Leng. *(Issued December 12, 1919).*
 Dytiscidae. By J. D. Sherman, Jr. *(Issued July 14, 1919).*
 Part F: HEMIPTERA. By Edward P. VanDuzee. *(Issued July 14, 1919).*
 Part G: HYMENOPTERA AND PLANT GALLS.
 Sawflies. (Tenthredinoiden). By Alex. D. MacGillivray.
 Parasitic Hymenoptera. By Charles T. Brues.
 Wasps and Bees. By F. W. L. Scholten.
 Plant Galls. By E. Porrier Felt. *(Issued November 3, 1919).*
 Part H: SPIDERS, MITES AND MYRIAPODS.
 Spiders. By J. H. Emerton.
 Mites. By Nathan Banks.
 Myriapods. By Ralph V. Chamberlin. *(Issued July 14, 1919).*
 Part I: LEPIDOPTERA. By Arthur Gibson. *(Issued January 10, 1920).*
 Part J: ORTHOPTERA. By E. M. Walker. *(Issued September 4, 1920).*
 Part K: INSECT LIFE ON THE WESTERN ARCTIC COAST OF AMERICA. By Frits Johansen. *(In press).*

VOLUME IV: BOTANY

- Part A: FRESHWATER ALGAE AND FRESHWATER DIATOMS. By Charles W. Lowe. *(In preparation).*
 Part B: MARINE ALGAE. By F. S. Collins. *(In preparation).*
 Part C: FUNGI. By John Dearness. *(In preparation).*
 Part D: LICHENS. By G. K. Merrill. *(In preparation).*
 Part E: MOSSES. By R. S. Williams. *(Issued February 8, 1921).*

VOLUME V: BOTANY

- Part A: VASCULAR PLANTS. By James M. Macoun and Theo. Holm. *(Issued October 14, 1921).*
 Part B: CONTRIBUTIONS TO THE MORPHOLOGY, SYNONYMY AND GENERAL DISTRIBUTION OF ARCTIC PLANTS. By Theo. Holm. *(In press).*
 Part C: GENERAL NOTES ON ARCTIC VEGETATION. By Frits Johansen. *(In preparation).*

VOLUME VI: FISHES, TUNICATES, ETC.

- Part A: FISHES. By F. Johansen. *(In preparation).*
 Part B: ASCIDIANS, ETC. By A. G. Handmann. *(In preparation).*

VOLUME VII: CRUSTACEA

- Part A: DECAPOD CRUSTACEANS. By Mary J. Rathburn. *(Issued August 18, 1919).*
 Part B: SCHIZOPOD CRUSTACEANS. By Waldo L. Schmitt. *(Issued September 22, 1919).*
 Part C: CUMACEA. By W. T. Collins. *(Issued October 15, 1920).*
 Part D: ISOPODA. By P. L. Boone. *(Issued November 19, 1920).*
 Part E: AMPHIPODA. By Clarence R. Shoemaker. *(Issued September 7, 1920).*
 Part F: PYCNOGONIDA. By Leon L. Cole. *(Issued January 5, 1921).*
 Part G: EUPHYLLOPODA. By F. Johansen. *(In preparation).*
 Part H: CLADOCERA. By Chaney Juday. *(Issued June 23, 1920).*
 Part I: OSTRACODA. By R. W. Sharpe. *(In preparation).*
 Part J: FRESHWATER COPEPODA. By C. Dwight Marsh. *(Issued April 21, 1920).*
 Part K: MARINE COPEPODA. By A. Willey. *(Issued June 25, 1920).*
 Part L: PARASITIC COPEPODA. By Charles B. Wilson. *(Issued August 6, 1920).*
 Part M: CIRRIPEdia. By H. A. Purbury. *(In preparation).*

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME VIII: MOLLUSKS, ECHINODERMS,
COELENTERATES, ETC.

PART E: ROTATORIA

By H. K. HARRING

SOUTHERN PARTY—1913-1916

OTTAWA
F. A. ACLAND
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1921

Issued December 31, 1921.

The Rotatoria of the Canadian Arctic Expedition, 1913-1918.

By H. K. HAMMING,

Custodian of Rotatoria, U. S. National Museum.

When the Canadian Arctic Expedition was organized with Mr. James Murray as oceanographer and marine biologist, all students of the rotifers looked forward to a considerable increase of our knowledge of the group and hoped for a repetition of his success in the Antarctic as biologist of the Shackleton Expedition. These expectations were frustrated through his death following the loss of the *Karluq* and the career of an enthusiastic naturalist and tireless worker was brought to an untimely end. No other member of the expedition was able to make special collections of rotifers, but the general plankton collections made by the marine biologist of the southern party, Mr. Frits Johansen, contained a considerable number of rotifers, which were assigned to the writer for a report. Some collections made by Mr. J. M. Jessup while serving on the Alaskan Boundary Survey have been included, as they belong to the same faunal area and add somewhat to our scanty knowledge of the distribution of the Rotatoria in the Arctic. Virtually all that we know on this subject is to be found in Bergendal's *Zur Rotatorienfauna Grönlands* (1892), and the value of this is somewhat minimized by his unfamiliarity with the group prior to his visit to Greenland.

While the species reported on here are not very numerous, 64 in all, they furnish additional, even if superfluous, evidence that climate is not directly a factor in rotifer distribution. Four new species are described, among which a pelagic *Synchaeta* is of special interest, as it is an addition to the extremely small number of rotifers known to exist in the open ocean in water of normal salinity. The total absence of the genus *Brachionus*, so abundant elsewhere, is noteworthy; Bergendal mentions two species of this genus from Greenland, but his notes on these forms make it somewhat doubtful whether he really found any Brachionoids.

I am indebted to Mr. Frank J. Myers, of the American Museum of Natural History, New York City, for drawing the plates accompanying this report.

ORDER PLOIMA.

FAMILY NOTOMMATIDÆ.

Notommata copeus Ehrenberg.

A few specimens of this species were collected by Jessup in lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

Notommata cyrtopus Gosse.

Several specimens occurred in a collection from a pond near New Rampart House, at the International Boundary and Porcupine river, made by Jessup on June 12, 1911.

Proales sordida Gosse.

A few specimens in a collection made by Johansen among mosses and algae from a pond at Chantry island, Bernard harbor, Polkhu and Union strait, on June 17, 1916.

Diaschiza forficata (Ehrenberg).

Diaschiza cacca DIXON-NUTTALL and FREEMAN, Journ. Royal Micr. Soc., 1903, p. 134, pl. 4, fig. 11.

Abundant in a collection made by Johansen from a pond on the ridge at Bernard harbour, on July 3, 1916.

Diaschiza gracilis (Ehrenberg).

Common among algae growing on stones in river bed at Bernard harbour, August 16, 1915; abundant in ponds on the ridge at Bernard harbour, July 3, 1916. Both collections were made by Johansen.

Diaschiza gibba (Ehrenberg).

In a collection made by Johansen from ponds on the ridge at Bernard harbour, July 3, 1916, rare. In Jessup's collections from a muskeg lake, 28 miles north of New Rampart House, June 25, 1911, rare; lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911, few; lakes along the International Boundary, 48 miles north of New Rampart House, July 7, 1911, few; pools at Fort Yukon, May 24, 1912, rare.

Dicranophorus forcipatus (Müller).

Diglena forcipata HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 50, pl. 19, fig. 2.

Collected by Johansen among algae growing on stones in the river bed at Bernard harbour, August 16, 1915, few; by Jessup from lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911, few. The trophi of these Arctic specimens have only five large, relatively blunt, teeth in each ramus, while the typical form has eleven or twelve. It is possible that they may represent an undescribed species; the partly contracted material was not in such a condition as to make this clear.

Encentrum aligente, new species.

Plate I, figs. 1, 2.

The body is elongate and very slender, almost vermiform; the integument is soft and flexible and the animal highly contractile.

The head is small and cylindric, its diameter somewhat less than that of the abdomen, from which it is separated by a slight constriction at the level of the gastric glands. The abdomen is cylindric nearly its entire length; posteriorly it is slightly reduced at its junction with the foot, which has two joints of nearly equal length. The toes are short, about one twentieth of the entire length, blade-shaped and slightly decurved, with slender, acute points.

The corona is terminal; the lateral, marginal cilia are somewhat longer than the rest and form rudimentary auricles. The dorsal antenna is a small, ciliated pit in the normal position; the lateral antennae are on the posterior fourth of the abdomen.

The trophi are forcipate and of a rather unusual type. The rami are of the normal lyrate form, terminating in a strong, pointed tooth; on the inner edge, about mid-length, there is an additional fairly large, pointed tooth. The fulcrum is unusually well developed, its length being fully equal to the length of the rami. The unci are aberrant; a single, short and robust tooth, hinged to the ramus at mid-length on a knob-like epiphysis, appears to represent the uncus proper; it is connected to the manubrium through a rather slender bar, enlarged at the posterior end, and nearly as long as the tooth itself. The

manubria are long and strongly curved, so that their posterior ends meet in the median line. The bar intervening between the uncus and manubrium is probably only a local sclerification of the walls of the mastax, developed in response to a specialization of the typical forcipate trophi. A very similar structure is found in *Eucestrum ricciae* Harring; comparison may be made with *Eucestrum* (= *Diglena*) *hofsteti* de Beauchamp, which shows a simpler stage of the same development. The unci and the supplementary piece are no doubt closely joined to the rami, the several pieces moving together virtually as a unit.

The oesophagus is long and slender. The gastric glands are elongate oval and fairly large; they open into the stomach very close to the junction with the oesophagus. There is no constriction between the stomach and intestine. The ovary is fairly large and of somewhat irregular outline. A small bladder is present. The foot glands are pyriform and rather small; no mucus reservoir is present.

The ganglion is elongate saccate; no retrocerebral organ or eyespots are present.

Total length 360 μ ; toes, 22 μ ; trophi, 52 μ .

This species was found in abundance in a collection made by Johansen among algae in a brackish lagoon west of Martin point, on the arctic shore of Alaska, on July 28, 1914.

FAMILY BRACHIONIDÆ.

Platytias quadricornis (Ehrenberg).

Notus quadricornis HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 121, pl. 28, fig. 5.

A few specimens occurred in a collection from lakes on Old Crow river flats, 60 miles north of New Rampart House, visited by Jessup on July 11, 1911.

Keratella quadrata (Müller).

Anuraca aculeata HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 123, pl. 29 fig. 4.

This species is widely distributed in the Arctic; it was found in the following localities: among algae in a brackish lagoon west of Martin point, arctic Alaska, July 28, 1914; in freshwater plankton from the lake south of Bernard harbour, November 28, 1915; May 6 and 7, May 21, and June 12, 1916; all collections were made by Johansen. In Jessup's material it occurred in a small pool near the International Boundary line, lat. 69° 20' N., long. 141° W., July 23, 1912, and in a slough of Old Crow river, near New Rampart House.

Keratella cochlearis (Gosse).

Anuraca cochlearis HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 124, pl. 29, fig. 7.

Not common in the Arctic; a few specimens were collected by Johansen in the lake south of Bernard harbour, May 21 and June 12, 1916; by Jessup in a pool near the Boundary line, lat. 69° 20' N., July 23, 1912, and in a slough of Old Crow river, near New Rampart House, August 7, 1912.

Notholca striata (Müller).

This is the most abundant and widely distributed rotifer in the Arctic; it was collected by Johansen in a brackish lagoon at Martin point, Alaska, July 28, 1914; in the river-bed at Bernard harbour, August 16, 1915; ponds

on the ridge at Bernard Harbour, July 3, 1916; by Jessup in a pool at White Horse, Yukon Territory, June 11, 1911; pond near New Rampart House, International Boundary and Porcupine river, June 12, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911; lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911; lakes on Old Crow river flats, 60 miles north of New Rampart House, July 11, 1911; small muddy pool near the Boundary line, 25 miles north of New Rampart House, July 17, 1911; pools at Fort Yukon, May 24, 1912; slough of Old Crow river, near New Rampart House, August 7, 1912.

Notholca longispina (Kellicott).

Collected by Johansen in a brackish lagoon west of Martin point, arctic Alaska, July 28, 1914; abundant in the lake south of Bernard harbour, November 28, 1915; February 15, May 21 and June 12, 1916.

FAMILY EUCHLANIDÆ.

Mytilina ventralis (Ehrenberg).

Salpina macracantha HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 81, pl. 22, fig. 6.

In Jessup's collections from a pond near New Rampart House, at the International Boundary and Porcupine river, June 12, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911; lake 48 miles north of New Rampart House, July 7, 1911.

Mytilina mucronata (Müller).

Salpina mucronata HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 83, pl. 22, fig. 1.

A few specimens collected by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911.

Euchlanis dilatata Ehrenberg.

This species does not appear to be common in the Arctic: it was collected by Johansen among algae growing on stones in the river at Bernard harbour, on August 16, 1915, and in a pond at Chantry Island, Bernard harbour, on June 17, 1916; in Jessup's material it occurred in fair numbers in a collection from lakes on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911.

Euchlanis pellucida, new species.

Plate 2, figs. 1-5.

The corona agrees with that of other species of the genus. The body is triradiate in cross section and has a high dorsal keel and broad, lateral, wing-like expansions. The ventral plate is nearly circular, slightly constricted anteriorly at the opening for the head; it is joined directly to the lateral plates without the intervention of a longitudinal sulcus as in other species of the genus. The foot is obscurely two-jointed; two long setae project from the dorsal side of the first foot joint. The toes are long, slender and nearly straight, slightly enlarged posteriorly, and end in rather blunt points; their length is more than one fifth of the length of the body.

The dorsal antenna is very large in diameter and obliquely truncate; it bears a small tuft of sensory setae in a shallow, central depression. The lateral antennae are in the normal position.

The mastax, stomach, intestine, ovary and bladder are normal. The large retrocerebral sac is slightly opaque at the posterior end and without any median notch.

The width of the median and lateral ribs or keels is variable, as shown in figs. 3-5, plate 2. The lateral ribs usually project straight out from the body or slightly downwards; the form with strongly upcurved ribs is not common.

Total length 500-800 μ ; length of lorica, 300-180 μ ; width of lorica, 270-130 μ ; length of toes 90-140 μ .

This species has been known to the writer for some time from many localities in the United States; it was collected by President Birge, of the University of Wisconsin, and Mr. Juday, of the Wisconsin Geological and Natural History Survey, in ponds and bayous of the Mississippi river in southern Louisiana; by Mr. Frank J. Myers, around Atlantic City, New Jersey, and in Polk county, Florida; by Mr. Myers and the writer in sphagnum bogs and ponds throughout Vilas and Oneida counties, Wisconsin. It was common in a collection made by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911. The presence of this species in the Arctic is of no special significance, as the collections listed above show that it is widely distributed; it seems, however, to be limited to regions with very soft, non-calcareous waters, where it often occurs in great abundance.

Euchlanis pellucida looks superficially very much like *E. triquetra*, but it is considerably larger and readily distinguished from the latter by the absence of the lateral sulcus.

***Euchlanis deflexa* Gosse.**

This species, which is ordinarily considered quite rare, was common in Jessup's collections: from a pond near Yukon river, Yukon Territory, May 26, 1911; pool at White Horse, Yukon Territory, June 7, 1911; lakes on Old Crow river flats, 35 miles north of New Rampart House, June 23, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911; pools at Fort Yukon, May 24, 1912.

***Euchlanis triquetra* Ehrenberg.**

Collected in abundance by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911; pools at Fort Yukon, May 24, 1912, and in a slough of Old Crow river, near New Rampart House, August 7, 1912.

***Euchlanis eoropha* Gosse.**

A few specimens of this comparatively rare species were collected by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911, and in a small muddy pool 25 miles north of New Rampart House, July 17, 1911.

***Lecane ephestra*, new species.**

Plate 3, figs. 1, 2.

The outline of the lorica is slightly ovate; the anterior margins of the dorsal and ventral plates are coincident and straight. The anterior spines are very short and rather stout. The dorsal plate is broadly elliptic and rounded posteriorly; it is without markings. The ventral plate is considerably narrower

than the dorsal and nearly parallel-sided; the margins are ill-defined and no longitudinal sulci are present. The markings are confined to the central area of the ventral plate; the transverse fold is well marked. The posterior segment of the body is rounded and projects very slightly beyond the dorsal plate.

The first foot joint is narrowest at the posterior end; the second foot joint is large and sub-square; it projects beyond the lorica. The coxal plates are obtusely triangular, with their apices close to the second foot joint. The toes are long and slender; the outer edges are very slightly curved; a small claw is present, excavate on the inner margin.

Total length 125μ ; length of dorsal plate 81μ , width 88μ ; length of ventral plate 90μ , width 60μ ; width of anterior points 51μ ; length of toes including claws 36μ , length of claw 6μ .

Lecane ephestra was collected by Jessup in a muskeg lake, 28 miles north of New Rampart House, on June 25, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; ponds 48 miles north of New Rampart House, on July 7, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; it has been collected in the United States by Mr. Frank J. Myers around Atlantic City, New Jersey, and by Mr. Myers and the writer in Vilas and Oneida counties, Wisconsin.

***Lecane mira* Murray.**

Cathypna mira MURRAY, Journ. Royal Micr. Soc., 1913, p. 553, pl. 22, fig. 3.

Abundant in a collection made by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911; also in lakes 48 miles north of New Rampart House, July 7, 1911, and two lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911.

***Lecane ligona* (Dunlop).**

Cathypna ligona DUNLOP, Journ. Quekett Micr. Club, ser. 2, vol. 8, 1901, p. 29, pl. 2, figs. 4-6.

A few specimens of this rare species were collected by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, July 3, 1911, and in two lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911.

***Lecane hornemanni* (Ehrenberg).**

Cathypna hornemanni MURRAY, Journ. Royal Micr. Soc., 1913, p. 349, pl. 16, fig. 26.

Collected by Johansen among algae growing on stones in the river bed at Bernard harbour, August 16, 1915; by Jessup in lakes on Old Crow river flats, 10 miles north of New Rampart House, July 3, 1911.

***Lecane jessupi*, new species.**

Plate, 3 figs. 3, 4.

The outline of the lorica is slightly ovate, truncate posteriorly and the anterior margin cuspidate. The dorsal plate is ovate, rounded posteriorly and slightly narrower than the ventral plate, which is somewhat elliptic. The anterior margin of the dorsal plate is nearly straight; it is slightly convex for the greater portion of its width and excavate at the lateral cusps; the anterior margin of the ventral plate is lunate. There are no markings on either dorsal or ventral plate; the lateral sulci are deep. The lorica is strongly compressed dorso-ventrally.

The posterior segment of the body is roughly trapezoidal in outline and cuspidate at the posterior angles; the margin is convex in the median portion and has a slight concavity at the angles. There is a well marked constriction at the junction of the ventral plate and the posterior segment. The coxal plates are semi-ovate.

The first foot joint is well marked and widest posteriorly; the second foot joint is nearly square. The toes are short, cylindric for one half their length and end in acute, conical points.

Total length 126 μ ; length of lorica 108 μ ; length of dorsal plate 93 μ ; width 93 μ ; width of ventral plate 96 μ ; width of anterior points 58 μ ; length of toes 27 μ .

Lecane jessupi had some resemblance to *L. brachydactyla* (Stenroos), which is shown on plate 3, figs. 5, 6, for comparison. The differences are fairly evident: *L. brachydactyla* has anterior spines and a straight, somewhat flexible dorsal margin, a double-curved ventral margin and the posterior segment is very simple in outline. *L. jessupi* was collected by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; it was not abundant.

***Lecane luna* (Müller).**

Cathypna luna HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 91, pl. 24, fig. 4.

Apparently not common in the Arctic; it was collected by Jessup from two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911, and ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911.

***Lecane unguolata* (Gosse).**

Cathypna minnesotensis MURRAY, Journ. Royal Micr. Soc., 1913, p. 345, pl. 13, fig. 18.

A few specimens in a collection made by Jessup from two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

***Lecane clara* (Bryce).**

Distyla clara BRYCE, Science Gossip, vol. 28, 1892, p. 271, text fig.

Collected by Jessup from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911, and in a slough of Old Crow river, near New Rampart House, on August 7, 1912. Only a few specimens were found in each collection.

***Lecane depressa* (Bryce).**

Distyla depressa BRYCE, Science Gossip, vol. 28, 1892, p. 271, text fig.

Common in a collection made by Johansen among algae growing on stones in the river bed at Bernard harbour, on August 16, 1915. In Jessup's collections from lakes on Old Crow river flats, 35 miles north of New Rampart House, on June 23, 1911; muskeg lake, 28 miles north of New Rampart House, on June 25, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; lakes on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911; small muddy pool, 25 miles north of New Rampart House, on July 17, 1911.

***Lecane flexilis* (Gosse).**

Lecane flexilis HARRING, Proc. U. S. Nat. Mus., vol. 47, 1911, p. 538, pl. 19, figs. 1-3.

Collected by Johansen among algae growing on stones in the river bed at Bernard harbour, on August 16, 1915; in Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911, and lakes on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911.

***Monostyla hamata* Stokes.**

A few specimens in Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911.

***Monostyla bulla* Gosse.**

This species was found in small numbers in Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

***Monostyla closterocerca* Schmarda.**

Monostyla closterocerca MURRAY, Journ. Royal Micr. Soc., 1913, p. 357, pl. 15, fig. 39.

A few specimens in Jessup's collections from lakes on Old Crow river flats, 60 miles north of New Rampart House, July 11, 1911.

***Monostyla crenata* Harring.**

Collected by Jessup in two lakes on Old Crow river flats, 55 miles north of New Rampart House, July 10, 1911; not common.

***Monostyla lunaris* Ehrenberg.**

Abundant and widely distributed in the Arctic. It was collected by Johansen among algae growing on stones in the river bed at Bernard harbour, on August 16, 1915; by Jessup in a muskeg lake, 28 miles north of New Rampart House, on June 25, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; lakes 48 miles north of New Rampart House, on July 7, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911; small muddy pool, 25 miles north of New Rampart House, on July 17, 1911; slough of Old Crow river near New Rampart House, on August 7, 1912.

***Monostyla cornuta* (Müller).**

A few specimens collected by Johansen in the lake south of Bernard harbour, on May 21, 1916; by Jessup in a pond near New Rampart House, at the International Boundary and Porcupine River, on June 12, 1911.

***Lepadella ovalis* (Müller).**

Lepadella ovalis HARRING, Proc. U. S. Nat. Mus., vol. 51, 1916, p. 537, pl. 89, figs. 4-10.

Collected in small numbers by Jessup from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; pools at Fort Yukon, May 24, 1912.

Lepadella patella (Müller).

Lepadella patella HARRING, Proc. U. S. Nat. Mus., vol. 51, 1916, p. 539, pl. 90, figs. 1-12; pl. 91, figs. 1, 2.

This species is widely distributed in the Arctic, but apparently in small numbers; it was collected by Johansen among algae in the river bed at Bernard Harbour, on August 16, 1915; pond at Chantry Island, Bernard harbour, on June 17, 1916; by Jessup in a pond near New Rampart House, at the International Boundary and Porcupine river, on June 12, 1911; lakes on Old Crow river flats, 10 miles north of New Rampart House, on July 3, 1911; lakes 48 miles north of New Rampart House, on July 7, 1911; ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911; ponds at Fort Yukon, on May 21, 1912.

Lepadella acuminata (Ehrenberg).

Lepadella acuminata HARRING, Proc. U. S. Nat. Mus., vol. 51, 1916, p. 516, pl. 92, figs. 1-8.

A few specimens of this species were collected by Jessup in a pond near New Rampart House, at the International Boundary and Porcupine river, on June 12, 1911.

Lepadella ehrenbergii (Perty).

Lepadella ehrenbergii HARRING, Proc. U. S. Nat. Mus., vol. 51, 1916, p. 553, pl. 91, figs. 1-1.

In Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

Colurella colurus (Ehrenberg).

Colurus amblytelus HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 104, pl. 25, fig. 5.

Collected by Johansen in abundance from a brackish lagoon west of Martin point, Alaska, on July 28, 1911; among algae growing on stones in the river bed at Bernard harbour, on August 16, 1915; by Jessup in lakes in Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

Trichotria pocillum (Müller).

Trichotria pocillum HARRING, Bull. 81 U. S. Nat. Mus., 1913, p. 106.

A few specimens collected by Jessup in lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

Trichotria tetractis (Ehrenberg).

Trichotria tetractis HARRING, Bull. 81 U. S. Nat. Mus., 1913, p. 106.

Common in Jessup's collections; pond near New Rampart House, at the International Boundary and Porcupine river, on June 12, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; lakes 48 miles north of New Rampart House, on July 7, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; ponds on old Crow river flats 60 miles north of New Rampart House, on July 11, 1911; slough of Old Crow river, near New Rampart House, on August 7, 1912.

FAMILY TRICHOCERCIDÆ.

Trichocerca longiseta (Schränk).

Rattulus longiseta JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 341, pl. 8, figs. 67-72.

Collected by Jessup in a pond near New Rampart House, at the International Boundary and Porcupine river, on June 12, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

Trichocerca mucosa (Stokes).

Rattulus mucosus JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 331, pl. 10, figs. 86-91.

A few specimens in Jessup's collections from a slough of Old Crow river, near New Rampart House, on August 7, 1912.

Trichocerca rattus (Müller).

Rattulus rattus JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 333, pl. 11, figs. 100, 101.

Collected by Jessup from lakes 18 miles north of New Rampart House, on July 7, 1911; ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911; small muddy pool, 25 miles north of New Rampart House, on July 17, 1911; it was rare in all of these collections.

Trichocerca cristata Harring.

Rattulus carinatus JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 332, pl. 11, figs. 95-97.

A few specimens were collected by Jessup from pools at Fort Yukon, on May 24, 1912.

Trichocerca bicuspes (Pell.).

Rattulus bicuspes JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 336, pl. 8, figs. 73-76.

Rare in Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

Trichocerca scipio (Gosse).

Rattulus scipio JENNINGS, Bull. U. S. Fish Comm., vol. 22 (for 1902), 1903, p. 322, pl. 5, figs. 50-52, pl. 13, figs. 111, 112.

Collected by Jessup from a muskeg lake, 38 miles north of New Rampart House, on June 25, 1911.

Diurella porcellus (Gosse).

Collected by Johansen among algae growing on stones in the river bed at Bernard harbour, on August 16, 1915; by Jessup from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

Diurella tenuior (Gosse).

In Jessup's collections from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

Diurella collaris (Rousselet).

Collected by Jessup from lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911.

Diurella cavla (Gosse).

A few specimens in Jessup's collections from a pond near New Rampart House, at the International Boundary and Porcupine river, on June 12, 1911.

FAMILY SYNCHAETIDÆ.**Synchaeta oblonga** Ehrenberg.

Abundant in collections made from the lake at Bernard harbour by Johansen on May 6 and 7, 1916; same lake south of Bernard harbour, May 21, 1916.

Synchaeta johanseni, new species.

Plate 1, fig. 3.

The body is fairly slender, bell-shaped and very transparent. Its greatest width, about mid-length, is one third of the total length. The foot is well marked off from the body, large at the base and tapers gradually to the very small toes; its length is one fourth of the length of the body. The head is triangular and the auricles powerful; on the median line, between the anterior pair of tactile bristles, there is a tubular sensory organ as in *S. vorax* Rousselet. The dorsal antenna is in the normal position; the lateral antennae are near the posterior end of the body and well towards the ventral side; they are slender tubules, armed with a minute tuft of setae. The foot glands are very small. The form and position of the eyespot could not be made out from the preserved material.

Total length 350 μ ; width of body at mid-length 120 μ ; length of foot 70 μ ; length of toes 7 μ .

This species occurred in large numbers in a surface collection made by F. Johansen on August 23, 1914, at station 36, off Cape Lyon, in Amundsen gulf.

Synchaeta johanseni is closely related to *S. vorax* Rousselet, from which it differs in the more slender body, longer and stouter foot, very small foot glands and minute toes, as well as in the position of the lateral antennae. Its presence in Amundsen Gulf is of the greatest interest, as up to the present only two species of rotifers, *Synchaeta atlantica* and *Trichocerca* (= *Rattulus*) *henseni*, are known from oceanic waters; these were both found by Zelinka in the collections of the German Plankton Expedition from the Atlantic ocean, south of Iceland. While it would perhaps be incorrect to call Amundsen Gulf an ocean, the conditions where the collection was made are oceanic, at least as far as salinity and absence of admixture of fresh water are concerned; there are no rivers of any considerable volume discharging near Cape Lyon, and Mr. Johansen informs me that few of the rivers flowing into the Arctic ocean carry much water in the summer. How to account for the presence of this rotifer at a single station and its absence everywhere else is a problem for which no solution can be offered; it may be noted that the collection contained virtually no other zoöplankton, and it is possible that the absence of enemies may be an important factor in the maintenance of this rotifer in such a circumscribed area.

Filinia longiseta (Ehrenberg).

Triarthra longiseta HUDSON and GOSSE, Rotifera, 1886, vol. 2, p. 6, pl. 13, fig. 6.

Collected by Johansen in a brackish lagoon west of Martin point, Alaska, on July 28, 1914; lake south of Bernard harbour, November 28, 1915; on May 6, 7, and June 12, 1916.

FAMILY PLOESOMATIDÆ.

Ploesoma lenticulare Herriek.

A few specimens in Jessup's collections from lakes on Old Crow river flats, 10 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

FAMILY TESTUDINELLIDÆ.

Testudineella patina (Hermann).

In Jessup's collections from pools at White Horse, Yukon Territory, on June 7 and 11, 1911; lakes on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911; pools at Fort Yukon, on May 21, 1912.

Testudinella parva (Ternetzi).

Collected by Jessup from lakes on Old Crow river flats, 10 miles north of New Rampart House, on July 3, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911.

ORDER FLOSCULARIACEA

FAMILY CONOCHILIDÆ.

Conochilus hippocrepis (Schrank).

Conochilus color HUDSON and GOSSE, Rotifera, 1886, vol. 1, p. 89, pl. 8, fig. 3.

Abundant in Jessup's collections; lakes on Old Crow river flats, 35 miles north of New Rampart House, on June 23, 1911; muskeg lake, 28 miles north of New Rampart House, on June 25, 1911; lakes on Old Crow river flats, 40 miles north of New Rampart House, on July 3, 1911; lakes 18 miles north of New Rampart House, on July 7, 1911; two lakes on Old Crow river flats, 55 miles north of New Rampart House, on July 10, 1911; ponds on Old Crow river flats, 60 miles north of New Rampart House, on July 11, 1911.

Conochilus unicornis (Rousselet).

A few specimens in Jessup's collections from lakes on Old Crow river flats, 10 miles north of New Rampart House, on July 3, 1911.

ORDER BDELLIODA.

FAMILY PHILODINIDÆ.

Rotaria neptunia (Ehrenberg).

Actinurus neptunia HUDSON and GOSSE, Rotifera, 1886, vol. 1, p. 108, pl. 10, fig. 6.

Collected by Jessup in a small, muddy pool, 25 miles north of New Rampart House, on July 11, 1911.

Rotaria rotunda (Pallas).

Rotifer vulgaris HUDSON and GOSSE, Rotifera, 1886, vol. 1, p. 104, pl. 10, fig. 2.

Collected in great numbers by Johansen in a pond at Bernard harbour, on June 16, 1915.

Rotaria macrura (Ehrenberg).

Rotifer macrurus HUDSON and GOSSE, Rotifera, 1886, vol. 1, p. 107, pl. 10, fig. 4.

Very abundant in Johansen's collections from a tundra-swamp at Bernard harbour, on July 15, 1915, and May 9, 1916.

BIBLIOGRAPHY.

- Beauchamp, P. de.
1913. Rotifères, pp. 105-116, text figs., in: Deuxième expédition antarctique française. Sciences naturelles: Documents scientifiques. Polychètes et Tricladés - manéides. Pterobranches. Chironomides. Rotifères. Doc. Paris, 116 pp., 9 pls., 1 map, text figs.
- Bergendal, D.
1891. Kurzer Bericht über eine im Sommer d.J. 1890 unternommene Reise nach Grönland. *Bihang Svensk Vetensk. Akad. Handl.*, vol. 17, Abt. 4, No. 4, 29 pp.
1892. Beiträge zur Fauna Grönlands. I. Zur Rotatoria in Grönland. *Acta Univ. Lundensis*, vol. 28, sect. 2, No. 4, 189 pp., 6 pls.
- Brehm, V.
1910. Die Rotatorien des Sarggebietes. *Verh. Ges. Deutsch. Naturh. Ärzte*, vol. 81, pt. 2, 1. Hälfte, pp. 190-191.
- Bryce, D.
1892. On some moss-dwelling Cathypnidæ, with descriptions of five new species. *Science Gossip*, London, vol. 28, pp. 271-275, text figs.
1897. Contributions to the nonmarine fauna of Spitzbergen. Part II. Report on the Rotifera. *Proc. Zool. Soc. London*, pp. 793-799.
- Cleve, A.
1899. Notes on the plankton of some lakes in Lule Lappmark, Sweden. *Öfvers. Svensk. Vetensk. Akad. Förhandl.*, vol. 56, pp. 825-835, text figs.
- Dixon-Nuttall, F. R., and Freeman, R.
1903. The Rotatorian genus *Dioschiza*: a monographic study, with description of a new species. *Journ. Royal Micr. Soc.*, pp. 1-11, 129-131, pls. 1-4.
- Dunlop, M. F.
1901. On a new rotifer, *Cathypna ligona*. *Journ. Quekett Micr. Club*, ser. 2, vol. 8, pp. 29-32, pl. 2.
- Guerne, J. de, and Richard, J.
1889. Sur la faune des eaux douces du Grönland. *Compt. Rend. Acad. Sci., Paris*, vol. 108, pp. 630-632.
1892. Voyage de M. Charles Rabot en Islande. Sur la faune des eaux douces. *Bull. Soc. Zool. France*, vol. 17, pp. 75-80.
1892. Sur la faune pélagique des eaux douces de l'Islande. *Compt. Rend. Acad. Sci., Paris*, vol. 111, pp. 310-313.
- Harring, H. K.
1913. Synopsis of the Rotatoria. *Bull. S. I. S. Nat. Museum*, 226 pp.
1911. Report on Rotatoria from Panama with descriptions of new species. *Proc. U. S. Nat. Museum*, vol. 47, pp. 525-561, pls. 16-24.
1916. A revision of the Rotatorian genera *Lepadella* and *Lophocharis* with descriptions of five new species. *Proc. U. S. Nat. Museum*, vol. 51, pp. 527-568, pls. 89-97.
- Hudson, C. T., and Gosse, P. H.
1886. The Rotifera or wheel-animalcules, both British and foreign. Vol. 1, VI + 128 pp., 15 pls., vol. 2, 114 pp., pls. 16-30.
1889. The Rotifera or wheel-animalcules, both British and foreign. Supplement. 64 pp., pls. 31-34.
- Jennings, H. S.
1903. Rotatoria of the United States. II. A monograph of the Rotariidae. *Bull. U. S. Fish Comm.*, vol. 22, pp. 273-352, pls. 1-15.
- Levander, K. M.
1901. Beiträge zur Fauna und Algenflora der süßen Gewässer an der Murmanküste. *Acta Soc. Fauna et Flora Fennica*, vol. 20, No. 8, 35 pp., text figs.
1905. Zur Kenntnis des Planktons einiger Binnenseen in Russisch-Lappland. *Festschrift für Fäimén*, vol. 1, No. 11, 49 pp., 3 pls.
- Levinsen, G. M. R.
1882. Småa Bidrag til den grønlandske Fauna. *Vid. Medd. Naturh. For., Kjøbenhavn*, ser. 4, vol. 3, pp. 127-136, pl. 2, text figs.
1914. Rotatoria. *Medd. om Grønland*, Kjøbenhavn, vol. 23, No. 6, pp. 635-658.
- Murray, J.
1908. Arctic rotifers, collected by Dr. William S. Bruce. *Proc. Royal Phys. Soc. Edinburgh*, vol. 17, pp. 121-127.
1908. Note on microscopic life in Gough Island, South Atlantic Ocean. *Proc. Royal Phys. Soc. Edinburgh*, vol. 17, pp. 127-129.
1910. Antarctic Rotifera. British Antarctic Exped. 1907-1909. *Rep. Sci. Invest.*, vol. 1, pp. 41-65, pls. 9-13.
1913. South American Rotifera. Part II. *Journ. Royal Micr. Soc.*, 1913-1914, 13-15.
1913. Notes on the family Cathypnidæ. *Journ. Royal Micr. Soc.*, 1913-1914, A, pls. 22, 23.

- Pouchet, G.
1892. Sur la faune pélagique du Dyrefjord (Islande). Compt. Rend. Acad. Sci., Paris, vol. 114, pp. 191-192.
1893. Sur le plancton de la lagune nord de Jan-Mayen. Compt. Rend. Acad. Sci., Paris, vol. 116, pp. 1207-1208.
- Richard, J.
1898. Sur la faune des eaux douces explorées en 1898 pendant la campagne du yacht "Princesse Alice." (Lofoten, Spitzberg, Hes Beeren, Hope, de Barents et Færøer.) Mém. Soc. Zool. France, vol. 11, pp. 326-338, text figs.
- Richters, F.
1907. Die Fauna der Moosrasen des Gaussbergs und einiger südlicher Inseln. Deutsche Spdpolar-Expedition 1901-1903 (E. von Drygalski ed.), vol. 9, Zool., vol. 1 (pt. 1), pp. 259-302, pls. 16-20.
- Skorikov, A. S.
1901. Beitrag zur Planktonfauna arktischer Seen. Zool. Anz., vol. 27, pp. 209-212.
- Voronkov, N. V.
1911. Plankton vodnoy poluostrova Ia-mala. (Materialy, privezennye Ia-madskoi ekspeditsiei B. M. Zlatkova 19^o goda. Kolovratki i obshel'mia kharakteristika planktona.) (Sur le plancton des bassins de la presqu'île de Yamal. Rotifères et caractères généraux du plancton.) (Russian text) Ann. Mus. Zool. Acad. Sci., St.-Petersbourg, vol. 16, pp. 180-214, 3 maps, text figs.

PLATE I.

- Fig. 1. *Enccentrum algente*, lateral view. x170
2. *Enccentrum algente*, trophi, ventral view
3. *Synchaeta johanseni*, dorsal view. x170

PLATE I

2

1

3

PLATE II.

- Fig. 1. *Euchlanis pellucida*, dorsal view. x130.
2. *Euchlanis pellucida*, lateral view.
3. *Euchlanis pellucida*, variety A, cross section.
4. *Euchlanis pellucida*, variety B, cross section.
5. *Euchlanis pellucida*, variety C, cross section.

PLATE II.

PLATE III.

- Fig. 1. *Lecane ephestra*, dorsal view. x300.
2. *Lecane ephestra*, ventral view.
3. *Lecane jessupi*, dorsal view. x300.
4. *Lecane jessupi*, ventral view.
5. *Lecane brachydactyla*, dorsal view. x300.
6. *Lecane brachydactyla*, ventral view.

PLATE III.

PLATE IV.

- Fig. 1. Temporary pools in swamp, fed from snow drifts. Berland harbour, Northwest Territories, July 2, 1915. Photo by F. Johansen.
- Fig. 2. Valley of Firth river, Alaska-Yukon boundary, showing ponds and flood plain ice. Photograph by International Boundary survey, 141st Meridian, 1912.

PLATE IV

Fig. 1.

Fig. 2.

Report of the Canadian Arctic Expedition, 1913-18.

VOLUME VIII: MOLLUSKS, ECHINODERMS, COELENTERATES, ETC.

- Part A: MOLLUSKS, RECENT AND PLEISTOCENE. By William H. Dall. *Issued September 24, 1919.*
- Part B: CEPHALOPODA AND PTEROPODA.
Cephalopoda. By S. S. Berry.
Pteropoda. By W. F. Clapp. *In preparation.*
- Part C: ECHINODERMS. By Austin H. Clark. *Issued April 16, 1920.*
- Part D: BRYOZOA. By R. C. Osburn. *In preparation.*
- Part E: ROTATORIA. By H. K. Harring. *In press.*
- Part F: CHALITOGNATHA. By A. G. Huntsman. *In preparation.*
- Part G: ALCYONARIA AND ACTINARIA. By A. E. Verrill. *In press.*
- Part H: MILDUSAE AND CTENOPHORA. By H. B. Bigelow. *Issued June 24, 1921.*
- Part I: HYDROIDS. By C. McLean Fraser. *In preparation.*
- Part J: PORIFERA.

VOLUME IX: ANNELIDS, PARASITIC WORMS, PROTOZOANS, ETC.

- Part A: OLIGOCHAETA.
Lumbriculidae. By Frank Smith.
Enchytraeidae. By Paul S. Welch. *Issued September 29, 1919.*
- Part B: POLYCHAETA. By Ralph V. Chamberlain. *Issued November 29, 1920.*
- Part C: HIRUDINEA. By J. P. Moore. *Issued February 1, 1921.*
- Part D: GEPHYREA. By Ralph V. Chamberlain. *Issued June 24, 1920.*
- Part E: ACANTHOCEPHALA. By H. L. Van Cleave. *Issued April 2, 1920.*
- Part F: NEMATODA. By N. S. Cobb. *In preparation.*
- Part G-H: TREMATODA AND CESTODA. By A. R. Cooper. *Issued February 1, 1921.*
- Part I: TURBELLARIA. By A. Huxley. *In preparation.*
- Part J: GORDIACEA.
- Part K: NEMATINI. By Ralph V. Chamberlain. *In preparation.*
- Part L: SPOROZOA. By J. A. Mayr. *In preparation.*
- Part M: FORAMINIFERA. By J. A. Cushman. *Issued February 9, 1920.*

VOLUME X: PLANKTON, HYDROGRAPHY, TIDES, ETC.

- Part A: PLANKTON. By Albert Mann. *In preparation.*
- Part B: MARINE DIATOMS. By L. W. Bailey. *In preparation.*
- Part C: TIDAL OBSERVATIONS AND RESULTS. By W. Bell Dawson. *Issued October 1, 1920.*
- Part D: HYDROGRAPHY. *In preparation.*

VOLUME XI: GEOLOGY AND GEOGRAPHY

- Part A: THE GEOLOGY OF THE ARCTIC COAST OF CANADA, WEST OF THE KENT PENINSULA. By J. J. O'Neill. *In preparation.*
- Part B: MAPS AND GEOGRAPHICAL NOTES. By Kenneth G. Chiptman and John R. Cox. *In preparation.*

VOLUME XII: LIFE OF THE COPPER ESKIMOS

- THE LIFE OF THE COPPER ESKIMOS. By D. Jenness. *(Issued).*

VOLUME XIII: PHYSICAL CHARACTERISTICS AND TECHNOLOGY OF THE WESTERN AND CENTRAL ESKIMOS

- Part A: THE PHYSICAL CHARACTERISTICS OF THE WESTERN AND COPPER ESKIMOS. By D. Jenness. *In preparation.*
- Part B: THE OSTEOLOGY OF THE WESTERN AND CENTRAL ESKIMOS. By John Cameron. *In preparation.*
- Part C: TECHNOLOGY OF THE COPPER ESKIMOS. *To be prepared.*

VOLUME XIV: ESKIMO FOLK-LORE AND LANGUAGE

- Part A: FOLK-LORE, WITH TEXTS, FROM ALASKA, THE MACKENZIE DELTA, AND CORONATION GULF. By D. Jenness. *In preparation.*
- Part B: COMPARATIVE GRAMMAR AND VOCABULARY OF THE ESKIMO DIALECTS OF POINT BARROW, THE MACKENZIE DELTA, AND CORONATION GULF. By D. Jenness. *In preparation.*

VOLUME XV: ESKIMO STRING FIGURES AND SONGS

- Part A: STRING FIGURES OF THE ESKIMOS. By D. Jenness. *(Ready for Press).*
- Part B: SONGS OF THE COPPER ESKIMOS. By Helen H. Roberts and D. Jenness. *In preparation.*

VOLUME XVI: ARCHAEOLOGY

- CONTRIBUTIONS TO THE ARCHAEOLOGY OF WESTERN ARCTIC AMERICA. *(To be prepared).*