**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☒ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x
☐	☐	☐	☐	☒	☐
12x	16x	20x	24x	28x	32x

The copy filmed here has been reproduced thanks to the generosity of:

Stauffer Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Stauffer Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

1.45

1.50

1.56

1.60

1.65

1.70

1.75

1.80

1.85

1.90

1.95

2.00

2.05

2.10

2.15

2.20

2.25

2.30

2.35

2.40

2.45

2.50

2.55

2.60

2.65

2.70

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14603 CA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME III: INSECTS

PART J: ORTHOPTERA

By E. M. WALKER

SOUTHERN PARTY, 1913-16

G
670
1913
C2t
v.06
pt. J
c.3

OTTAWA
THOMAS MULVEY
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1920

Issued Sept. 4th, 1920

Report of the Canadian Arctic Expedition, 1913-18.

VOLUME III: INSECTS

- INTRODUCTION. By C. Gordon Hewitt. (*In press*).
- Part A: COLLEMBOLA. By Justus W. Tolson. (*Issued July 10, 1919*).
- Part B: NEUROPTEROID INSECTS. By Nathan Banks. (*Issued July 11, 1919*).
- Part C: DIPTERA.
Craneflies. By Charles P. Alexander.
Mosquitoes. By Harrison G. Dyer.
Diptera (excluding Tipulidae and Chironidae). By J. R. Malloch. (*Issued July 14, 1919*).
- Part D: MALLOPHAGA AND ANOPLURA.
Mallophaga. By A. W. Baker.
Anoplura. By G. F. Ferris and G. H. F. Nuttall. (*Issued September 12, 1919*).
- Part E: COLEOPTERA.
Forest Insects, including Ipsidae, Curculionidae, and Buprestidae. By J. M. Swaine.
Carabidae and Silphidae. By H. C. Fall.
Coccinellidae, Elateridae, Chrysomelidae, and Rhynchophora (excluding Ipsidae).
By C. W. Long.
Dytiscidae. By J. D. Sherman, Jr. (*Issued December 12, 1919*).
- Part F: HEMIPTERA. By Edward P. Van Duzee. (*Issued July 11, 1919*).
- Part G: HYMENOPTERA AND PLANT GALLS.
Sawflies. (Tenthredinoiden). By Alex. D. MacGillivray.
Parasitic Hymenoptera. By Charles T. Brues.
Wasps and Bees. By W. L. Shaden.
Plant Galls. By V. Porter Felt. (*Issued November 3, 1919*).
- Part H: SPIDERS, MITES, AND MYRIAPODS.
Spiders. By J. H. Emerton.
Mites. By Nathan Banks.
Myriapods. By Ralph V. Chamberlin. (*Issued July 11, 1919*).
- Part I: LEPIDOPTERA. By Arthur Gibson. (*Issued January 10, 1920*).
- Part J: ORTHOPTERA. By E. M. Walker. (*In press*).
- Part K: GENERAL OBSERVATIONS ON INSECT LIFE IN THE
ARCTIC. By Frits Johansen. (*In preparation*).

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME III: INSECTS

PART J: ORTHOPTERA

By E. M. WALKER

SOUTHERN PARTY, 1913-16

OTTAWA
THOMAS MULVEY
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1920

Orthoptera Collected in the Canadian Arctic.

By E. M. WALKER.

University of Toronto.

No orthoptera were received among the insects collected by the Canadian Arctic Expedition 1913-18, but we have received a single grasshopper collected by Mr. V. Stefánsson in 1911. The following is a note on this insect.

FAMILY ACRIDIDAE

Melanoplus frigidus (Bohemius).

A single female of this species in poor condition, bears label giving the following data:

"No. 1670 [E. J.], Langton bay (Franklin bay), Northwest Territories, summer of 1911. V. Stefánsson."

In a note to the writer Dr. R. M. Anderson states that "This specimen, brought to us with other insect specimens in 1914 from our old house at Langton bay by a former Eskimo employee, while taken in the Langton bay region, was most probably picked up from twenty to forty miles inland on the Horton river, south side of the Melville mountains, a range of hills about 1,000 feet high, skirting the south side of Franklin bay."

The capture of this Palearctic species in the above locality was not unexpected as several specimens were taken by Mr. J. M. Jessup on the International boundary, Alaska, lat. 69° 20' N., long. 141° W., on Aug. 8, 1912 (Caudell, Can. Ent., vol. XLVII, 1915, p. 160).

On account of the difficulty of determining species of this group from the female sex alone, the writer submitted this specimen to Mr. Morgan Hebard, who is engaged in a revision of the *Melanopli*. I had determined it as *Podisma frigidum* (Bohemius) with some doubt, as it differs slightly in the form of the valves of the ovipositor from the single female European specimen I have of this species, but Mr. Hebard has confirmed the determination. In a letter to the writer he says "You will note the transfer of this species to the genus *Melanoplus*. I am bringing out the data on this change in a paper which will be published shortly."¹ I have been likewise of the opinion, for some time, that this species is a true *Melanoplus*, and it is of special interest as being the only species of this genus known from the Old World, where it is widely distributed in northern regions, having been taken in Norway, Lapland and Siberia, and as a glacial relict in the Swiss Alps and the Tyrol.²

Three other species of Orthoptera are definitely recorded from the Arctic regions of North America. These are *Gomphoceris clavatus* Thomas, *Melanoplus borealis* (Fieber) and *M. fasciatus* (Barnston-Walker).

Gomphoceris clavatus was recorded by Caudell (*loc. cit.*) from the same locality in Alaska where *M. frigidus* was taken. It is a widely distributed species, ranging from eastern Manitoba to the Rocky Mountains and southward to Nebraska, Kansas, Colorado and New Mexico. It is found at high elevations in the mountains of Colorado, Wyoming, Idaho and New Mexico. The genus *Gomphoceris* is of Palearctic origin, *G. clavatus* being the only American species.

Melanoplus borealis has been until very recently considered as a typically arctic form. It has been recorded from Greenland (Fieber, *Notul.* 1853).

¹Since the above was written a preliminary discussion of this subject has appeared in the following paper by Mr. Hebard: "New Genera and Species of *Melanopli* found within the United States." (*Trans. Am. Ent. Soc.*, XIV, pp. 257-298, 1919).

²Hebard (*op. cit.*) states that the recently described *Podisma prosseni* Puschig from the Eisnhut in Carinthia is also a *Melanoplus*.

Labrador, Hudson Bay and Alaska. The specimens described by Scudder in his "Revision of the Melanopli" came from the Esquimaux village of Ramah, on the coast of Labrador, lat. 57° N., while the Alaska specimens recorded by Caudell (Pap. Harriman Alaska Exp., Proc. Wash. Acad. Sci., vol. II, pp. 511-512, 1900) were taken at Kukak Bay, Alaska Peninsula.

I have a large series of this species from Nain (lat. $56^{\circ} 30''$) and Hopedale (lat. $55^{\circ} 24''$), Coast of Labrador, an examination of which convinced me that they are not specifically distinct from *M. extremus* Walk., a species also recorded from Labrador and Arctic America, which was placed by Scudder in a different section of the genus. On submitting this question to Mr. Hebard, I learned that he had already placed *M. extremus* as a race of *borealis* in his unpublished notes on this group, and Messrs. Morse and Blatchley are also of the opinion that the two forms are conspecific. This being the case, *M. borealis*, as a species, is by no means exclusively arctic, but ranges over nearly the whole of Canada and extends also into various parts of the northern United States. It is, however, a typically boreal form and probably reaches its highest development in the Hudsonian zone.

Melanoplus fasciatus is a species of similar range and has also been recorded from Nain, Labrador, so that it may be fairly included in the arctic fauna.

Several other species of Acrididae have been reported from "Arctic America" chiefly by F. Walker (Cat. Derm. Salt. Brit. Mus., III-IV, 1869-71), but in the absence of definite localities it is uncertain whether they were taken in the Arctic zone proper or the Hudsonian. These are *Acrydium granulatum* Kirby, *Chorthippus curtipennis* (Harr.), *Hippiscus apiculatus* (Harr.) and *Melanoplus femur-rubrum* (DeGeer). The type locality of *A. granulatum* is in lat. 65° , and it was taken by Adam White (Richardson, Arctic Search. Exp., II, p. 360, 1851) at Fort Simpson (lat. $61^{\circ} 30'$) and at the junction of the Mackenzie and Slave rivers, so that it cannot be certainly regarded as arctic on the basis of these records. The other records are indefinite and that of *M. femur-rubrum* almost certainly erroneous, the species referred to being probably *M. borealis*.

A few other Orthoptera may be mentioned as very possibly ranging into the arctic regions. *Melanoplus kennicottii* Scudd. and *M. atlantis* (Riley) were taken by Kennicott on the Yukon river, Alaska. The former has also been recorded from Saskatchewan, Alberta and Montana, while the latter is widely distributed over the greater part of North America, extending into Mexico.

Melanoplus bruneri Scudd. (*M. alaskanus* Scudd.) is also recorded from Alaska and is widely distributed in Canada in the Boreal region.

Acrydium bruneri Boliver (which is perhaps the true *A. granulatum* (Kirby), and *Melanoplus birittatus* Say are recorded from the Hudson bay region, while *Circotettix verruculatus* (Kirby) ranges far north, the type specimen having come from lat. 57° N.

All the species mentioned above are locusts or "short-horned grasshoppers" (Acrididae), but there is one species of "long-horned grasshopper" (Tettigoniidae), which has some claim to membership in the Arctic Fauna. This is *Idionotus sphagnum* (Walk.) (syn. *I. brevipennis* Caud., *Platydes fletcheri* Caud.), which was originally described from St. Martin's Falls, Hudson Bay. It was redescribed by Caudell (Proc. U.S. Nat. Mus., XXXII, p. 396, 1907) as *Idionotus brevipennis* from a single male, taken by Kennicott in "Arctic America" (Caudell, l.c.). This same specimen had already been mentioned but not described by Scudder (Can. Ent., XXVI, p. 182, 1894) in his characterization of the genus *Idionotus*, and is stated to have been "collected by Kennicott somewhere on his explorations in or going to Alaska." This species is now known also from northwestern Ontario, Manitoba and Alberta.

¹I am indebted to Mr. W. S. Blatchley for the synonymy of *Idionotus brevipennis* Caud. with *Deeticus sphagnum* Walk. I had long suspected this to be the case, as we have only one northern Deeticine, as far as known.

Report of the Canadian Arctic Expedition, 1913-18.

Volume I: General Introduction, Narrative, Etc.

Part A: Northern Party, 1913-18.

Part B: Southern Party, 1913-16. By Rudolph Martin Anderson. (*In preparation*).

Volume II: Mammals and Birds.

Part A: Mammals. By Rudolph Martin Anderson. (*In preparation*).

Part B: Birds. By R. M. Anderson and P. A. Taverner. (*In preparation*).

Volume III: Insects.

Introduction. By C. Gordon Hewitt. (*In press*).

Part A: Collembola. By Justus W. Folsom. (*Issued*).

Part B: Neuropteroid Insects. By Nathan Banks. (*Issued*).

Part C: Diptera. By Chas. W. Alexander, Harrison G. Dyar, and J. R. Malloch. (*Issued*).

Part D: Mallophaga and Anoplura. By A. W. Baker, G. L. Ferris, and G. H. F. Nuttall. (*Issued*).

Part E: Coleoptera. By I. M. Swaine, H. C. Fall, C. W. Lere, and J. D. Sherman, Jr. (*Issued*).

Part F: Hemiptera. By E. P. Van Duzee. (*Issued*).

Part G: Hymenoptera and Plant Gallis. By Alex. D. MacGillivray, Charles T. Braes, F. W. L. Shaden,

and E. Porter Felt. (*Issued*).

Part H: Spiders, Mites, and Myriapoda. By J. H. Emerton, Nathan Banks, and Ralph V. Chamberlin.

(*Issued*).

Part I: Lepidoptera. By Arthur Gibson. (*Issued*).

Part J: Orthoptera. By L. M. Walker. (*In press*).

Part K: General Observations on Insect Life in the Arctic. By Frits Johansen. (*In preparation*).

Volume IV: Botany.

Part A: Freshwater Algae and Freshwater Diatoms. By Charles W. Lowe. (*In preparation*).

Part B: Marine Algae. By F. Collins. (*In preparation*).

Part C: Fungi. By John Dearness. (*In preparation*).

Part D: Lichens. By K. L. Merrill. (*In preparation*).

Part E: Mosses. By R. S. Williams. (*In press*).

Volume V: Botany.

Part A: Flowering Plants and Ferns. By James M. Macoun and Theo. Holm. (*In preparation*).

Part B: General Notes on Arctic Vegetation. By Frits Johansen. (*In preparation*).

Volume VI: Fishes, Tunicates, Etc.

Part A: Fishes. By F. Johansen. (*In preparation*).

Part B: Ascidians, etc. By A. G. Huntsman. (*In preparation*).

Volume VII: Crustacea.

Part A: Decapod Crustaceans. By Mary J. Rathbun. (*Issued*).

Part B: Schizopod Crustaceans. By Waldo L. Schmitt. (*Issued*).

Part C: Cumacea. By W. T. Calman. (*In press*).

Part D: Isopoda. By Miss P. L. Boone. (*In press*).

Part E: Amphipoda. By Clarence R. Shoemaker. (*In press*).

Part F: Pycnogonida. Leon J. Cole. (*In press*).

Part G: Euphyllipoda. By F. Johansen. (*In preparation*).

Part H: Cladocera. By Chancey Juday. (*Issued*).

Part I: Ostracoda. By R. W. Sharpe. (*In preparation*).

Part J: Freshwater Copepoda. By C. Dwight Marsh. (*Issued*).

Part K: Marine Copepoda. By A. Willey. (*Issued*).

Part L: Parasitic Copepoda. By Chas. B. Wilson. (*In press*).

Part M: Cirripedia. By H. A. Pilsbry. (*In preparation*).

Volume VIII: Mollusks, Echinoderms, Coelenterates, Etc.

Part A: Mollusks, Recent and Pleistocene. By Wm. H. Dall. (*Issued*).

Part B: Cephalopoda and Pteropoda. By S. S. Berry and W. F. Clapp. (*In preparation*).

Part C: Echinoderms. By Austin H. Clark. (*Issued*).

Part D: Bryozoa. By R. C. Osburn. (*In preparation*).

Part E: Rotatoria. By H. K. Harring. (*In preparation*).

Part F: Chetognatha. By A. G. Huntsman. (*In preparation*).

Part G: Actinozoa, and Aleyonaria. By A. E. Verrill. (*In press*).

Part H: Medusae and Ctenophora. By H. B. Bigelow. (*Issued*).

Part I: Hydroids. By McLean Fraser. (*In preparation*).

Part J: Porifera.

Volume IX: Annelids, Parasitic Worms, Protozoans, Etc.

Part A: Oligochaeta. By Frank Smith and Paul S. Welch. (*Issued*).

Part B: Polychaeta. By Ralph V. Chamberlin. (*In press*).

Part C: Hirudinea. By J. P. Moore. (*In press*).

Part D: Gephyrea. By Ralph V. Chamberlin. (*Issued*).

Part E: Acanthocephala. By H. J. Van Cleave. (*Issued*).

Part F: Nematoda. By N. A. Cobb. (*In preparation*).

Part G: Trematoda. By A. R. Cooper. (*In preparation*).

Part H: Cestoda. By A. R. Cooper. (*In preparation*).

Part I: Turbellaria. By A. Hassell. (*In preparation*).

Part J: Gordiacea.

Part K: Nemertina.

Part L: Sporozoa. By J. W. Mavor. (*In preparation*).

Part M: Foraminifera. By J. A. Cushman. (*Issued*).

Volume X: Plankton, Hydrography, Tides, Etc.

Part A: Plankton. By Albert Mann. (*In preparation*).

Part B: Marine Diatoms. By L. W. Bailey. (*In preparation*).

Part C: Tidal Observations and Results. By W. Bell Dawson. (*In press*).

Part D: Hydrography. (*In preparation*).