**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☒ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

**This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.**

10x 14x 18x 22x 26x 30x
12x 16x 20x 24x 28x 32x

The copy filmed here has been reproduced thanks to the generosity of:

MacOdrum Library
Carleton University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\longrightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

1	2	3
---	---	---

1	
4	

L'exemplaire filmé fut reproduit grâce à la générosité de:

MacOdrum Library
Carleton University

Les images suivantes ont été reproduites avec la plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

Ed Bruce
Carleton University
Library - Ottawa

Canada
Department of Mines

Hon. LOUIS CODERRE, Minister;
R. G. McCONNELL, Deputy Minister.

Geological Survey

Museum Bulletin No. 7

BIOLOGICAL SERIES, No. 4.

DECEMBER 20, 1914

**A NEW SPECIES OF DENDRAGAPUS (DENDRAGAPUS
OBSCURUS FLEMINGI) FROM SOUTHIERN
YUKON TERRITORY**

by

P. A. Taverner

33.5
104

OTTAWA
GOVERNMENT PRINTING BUREAU
1914

No. 1482

December 20, 1914

Canada
Geological Survey
Museum Bulletin No. 7.
BIOLOGICAL SERIES, No. 4.

*A New Species of Dendragapus (Dendragapus Obscurus Flemingi),
from Southern Yukon Territory.¹*

BY P. A. TAVERNER.

In the examination of a collection of *Dendragapus obscurus* recently acquired by the Canadian Geological Survey from the Teslin Lake region, on the Yukon-British Columbia boundary, it became evident that a new form of the species was probably represented. Two subspecies have heretofore been recognized as ranging into Canadian territory, *D. o. fuliginosus* and *richardsoni*. *Fuliginosus* is a dark bird with a light tail band and *richardsoni* a lighter one without a terminal band on the tail. The birds under consideration are without the tail band but are even darker than *fuliginosus*.

We had a fair series of both recognized forms from southern British Columbia, all in spring and early summer plumage and hence hard to distinguish from the new form which is represented entirely by late summer and autumn birds. The possibility existed that the darkness of the new birds but represented the new and unaged plumage of the post-nuptial moult; therefore application was made to the U. S. Biological Survey and its chief, Dr. Henshaw, was courteous enough to place at our disposal what summer and autumn specimens there were in the collection of that institution. A comparison of the specimens, including these, confirms our first surmise as to the racial differentiation of the

¹Republished from 'THE AUK,' Vol. XXXI, No. 3, July, 1914, pp. 385-388.

forms and I, therefore, present the following new subspecies for consideration.

Dendragapus obscurus flemingi new subspecies

FLEMING'S GROUSE.

Named after Mr. J. H. Fleming, C.M.Z.S., in recognition of the value of his ornithological work in Canada.

Subspecific Characters. Like *Dendragapus obscurus richardsoni*, without terminal tail band, but darker in general coloration even than *D. o. fuliginosus*.

D. o. richardsoni.¹

Back, interscapulars, and body color of rump. Olive-brown to fuscous.

Tail. Fuscous black. Tail feathers cut off sharply and squarely at end.

Throat patch. Body colour soiled creamy white.²

Sides of head and spotting of throat patch. Fuscous.

Around neck below throat patch. Between buffy brown and olive brown to clove brown, with sometimes a faint bluish tinge, deepening in front at lower neck to fuscous.

Breast and underparts. Light mouse gray.

Description of Type specimen. Type, adult male, No. 6651 Museum of the Geological Survey, Department of Mines, Canada. Near Teslin lake, Yukon Territory, October 10, 1912. Clement Lewis, Collector.

Back, rump, and tail, dusky neutral gray. Central feathers of tail and some lateral ones double rounded at ends.

Upper tail coverts, fuscous black, finely but sparsely vermiculated with buffish slate.³

Scapulars, wings, and upper wing coverts, fuscous, lightening towards wing tip. All, except flight feathers, finely marked with shades between rusty⁴ and grayish.⁵ Some scapulars with white shaft streaks, widening at

¹All colour terms except where otherwise stated, from Ridgway's "Colour Standards and Nomenclature," 1912 ed.

²These terms are not from "Ridgway's Colour Standards."

D. o. flemingi.

Back, interscapulars, and body colour of rump. Dusky neutral gray.

Tail. Dusky neutral gray. Tail feathers not having the same chopped off appearance. Middle and several lateral feathers slightly to markedly double-rounded at end.

Throat patch. Body colour pure white.

Sides of head and spotting of throat patch. Between neutral gray and deep neutral gray.

Around neck below throat patch. Between neutral gray and deep neutral gray, deepening in front at lower neck to dusky neutral gray.

Breast and underparts. Between neutral gray and deep neutral gray.

terminal to poorly defined marginal spots. The latter persisting on some of the lesser coverts as vague spots and narrow borders.

Below, between neutral gray and deep neutral gray, and obscured posteriorly by a white tipping of the feather ends that blends into the white of the basal under tail coverts.

Lower breast feathers with white shaft lines broadening towards the flanks and posteriorly where they develop into broad terminal spots, streaks, and bars. A fine vermiculation of buffy¹ and grayish¹ overlaying the body colour of the feather tips of the sides.

Under tail coverts dusky neutral gray or black with prominent white shaft lines and final borders. White increasing towards base of tail until body colour of feathers is hidden.

Head from deep neutral gray to neutral gray darkest on crown and ear coverts, overwashed with rusty¹ strongest on forehead at base of bill and more or less obscuring body colours of darker feathers and at last dying away at back and sides of upper neck as a faint olivaceous¹ tinge.

Throat patch, white with irregular deep neutral gray spotting, aggregated on malar region and each spotted feather terminated with small white border.

Lores, deep neutral gray, with white flecking.

Dark feathers of the breast band creamy white basally, the white gradually encroaching on the dark tips towards the sides, finally usurping most of the feather and making a semi-concealed white streak down the sides of the neck to the bend of the folded wing.

The characters of the female are less marked than of the male and without series for comparison may be difficult of recognition. However, they average in the same directions as the male being bluer underneath than *richardsoni* and darker dorsally, with the rufous¹ or rusty¹ markings bolder and more decided in character.

Measurements (in inches). Male. average of four specimens.

Wing, 8.25-8.75 (average 8.51); tail, 6.5-6.75 (average 6.33); tarsus, 1.70-1.80 (average 1.75); exposed culmen, 0.65-0.75 (average 0.71).

Female average of twelve specimens.

Wing, 7.45-8.10 (average 7.70); tail, 5.10-5.75 (average 5.25); tarsus 1.48-1.68 (average 1.55); exposed culmen, 0.65-0.71 (average 0.69).

Distribution. This description is based entirely on specimens taken within thirty miles of Teslin lake, on the boundary between British Columbia and Yukon Territory, longitude 123° 30', at the west base of the Cassiar mountains. The range of the form *cast.* therefore, be defined, but a study of the material examined is suggestive.

The male specimens in the series are as follows: British Columbia: Trail, Midway, and Revelstoke: 7 specimens.

Montana and Idaho: Preuss mts., Fiddle creek; and Nyak and Bear-tooth mts.: 4 specimens.

¹These terms are not from Ridgway's "Colour Standards."

Mackenzie; Mackenzie river: 1 specimen.

Yukon Territory, Teslin lake: 4 specimens.

	Wing	Tail	Tarsus	Culmen	General colour
Montana and Idaho	9.06	6.56	1.80	0.80	Light
British Columbia	8.90	6.96	1.77	0.77	Slightly darker
Mackenzie	8.75	6.55	1.70	0.80	Light
Teslin lake, Y. T.	8.51	6.33	1.75	0.71	Dark.

It will be seen that the most southern birds are the largest and lightest; the northern smallest and darkest, while those from lower British Columbia are intermediate though closer to the southern than the northern race. The Mackenzie form, being represented by only a single specimen in not very high plumage, cannot be definitely placed in the series though it agrees closely with the Montana and Idaho specimens. No specimens from the type locality of *richardsoni* (the northwest corner of Rocky Mountain park, lat. 53°, long. 115°) being obtainable I assume from the above that it ranges up the east side of the Rocky mountains and through them south of the International border in unmodified form, while the intermountain birds, *flemingi*, as they distribute northward from the boundary, become smaller and darker. It is possible that on examination, birds from farther north of Teslin lake will exhibit these characters in a still more marked degree

QH

301

C35

NO. 4

colour
darker

argest
from
to the
being
nage,
osely
n the
ocky
sume
ocky
order
, as
smaller
from
in a