Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

76
Circular No. 643

November 1942 • Washington, D. C.

UNITED STATES DEPARTMENT OF AGRICULTURE

LIBRARY
RECEIVED
DEC 23 1942

U. S. Department of Agriculture

Owlet Moths (Phalaenidae) Taken at Light Traps in Kansas and Nebraska¹

H. H. WALKDEN, *assistant entomologist, Division of Cereal and Forage Insect Investigations, Bureau of Entomology and Plant Quarantine*, and D. B. WHELAN, *assistant entomologist, Nebraska Agricultural Experiment Station*

United States Department of Agriculture, Bureau of Entomology and Plant Quarantine, in Cooperation with the Nebraska Agricultural Experiment Station.

CONTENTS

	Page		Page
Introduction.....	1	Seasonal flight records.....	14
Traps.....	2	Flight records of individual species.....	18
Collections.....	3	Summary.....	18
Proportion of the sexes.....	14		

INTRODUCTION

The information contained in this publication was obtained through the operation of light traps located at six widely separated points in the Missouri Basin, in areas typical of both the semiarid and humid phases of the agriculture of this region. It deals with the seasonal occurrence and abundance of the owlet moths (Phalaenidae (Noctuidae)), a group composed largely of such notoriously injurious pests as the corn earworm, the pale western cutworm, the western armyworm, the wheat-head armyworm, and many other seriously injurious cutworms inhabiting the region surveyed. This region is of exceptional interest to workers in applied entomology because it contains an owlet moth fauna representing not only the strictly subterranean species, controllable only by modification of cultural practices, but those possessing either intermediate or surface-inhabiting characteristics, and more or less amenable to insecticidal treatment.

The data herein contained, revealing the distribution, seasonal flight periods, and peaks of abundance of the various species, are basic to, or of value in, the consideration of cultural control and other methods for the suppression of the various pests involved.

The use of light traps offered an efficient means of obtaining this information. Accordingly, in the fall of 1934 traps were set up in

¹ Published as Paper No. 223, Journal Series, Nebraska Agricultural Experiment Station.

several locations and their operation was continued during the flight seasons of 1935-37. A record of the nightly catch was kept, the species, number of moths captured, and the proportion of the sexes being tabulated. The traps were put in operation early enough in the spring so that the first flights could be observed, and they were kept in operation until cold weather in the fall stopped all flight activity.

Observations were made in the following six localities: Cherryvale, Garden City, Hays, and Manhattan in Kansas and Lincoln and Scottsbluff in Nebraska (fig. 1). These localities ranged in elevation

FIGURE 1.—Location of the light traps.

as follows: Cherryvale 800 feet, Garden City 2,800 feet, Hays 2,000 feet, Mannattan 1,100 feet, Lincoln 1,200 feet, and Scottsbluff 3,900 feet.

TRAPS

With the exception of the one used at Manhattan, the traps were of similar type and were designed by the senior writer. They consisted of an inverted galvanized-iron cone, 2 feet in diameter, with a roof of the same material. The lamp (fig. 2, *B*) was suspended in the center just above the rim, and a fruit jar was attached to the bottom of the cone for receiving the catch. The killing agent, calcium cyanide, was put in a small sack and placed in the fruit jar. The cyanide was renewed daily.

The trap used at Manhattan (fig. 2, *A*) was designed by R. C. Smith, of the Kansas Agricultural Experiment Station. Except for a more shallow cone, and the use of glass buffers about the lamp, this trap was the same as those at the other localities. Five-hundred-watt clear lamps were used at Manhattan and Lincoln and 200-watt inside-frosted lamps elsewhere.

The locations of the traps were probably not ideal, but the availability of electric outlets was the deciding factor in the selection of sites. These were as follows:

(1) At Cherryvale the location was unsatisfactory. It was on a flour mill that was illuminated nearly every night and near a transformer station that was always brilliantly lighted. The competition of these other lights greatly affected the catches, and the results from this trap are not considered of much value.

(2) At Garden City the trap was located on the grounds of the Garden City Branch Station, Kansas Agricultural Experiment Station, about 5 miles northeast of the city. The light was visible in all directions, although a small grove of trees on the southwest somewhat obscured it in that direction. It was placed about 5 feet above the ground.

(3) At Hays the trap was placed in an open space between buildings on the grounds of the Fort Hays Branch Station, Kansas Agricultural Experiment Station. It was about 5 feet above the ground and was visible for a long distance to the east and south.

(4) At Manhattan the trap was placed on top of an outbuilding near the college power plant. The lamp was about 20 feet above the ground and could be seen for

FIGURE 2.—Light trap used at Manhattan, Kans. (A), and at the other locations (B).

some distance to the west, south, and north, but was obscured by buildings on the east.

(5) At Lincoln the trap was placed at the east edge of the Nebraska Agricultural Experiment Station orchard, well within the city limits. It commanded a view to the east for some distance, but in other directions the light was obscured by the orchard trees nearby.

(6) At Scottsbluff the trap was located on the grounds of the Scottsbluff Branch Station, Nebraska Agricultural Experiment Station, about 6 miles northwest of the city. The trap was elevated about 10 feet above the ground and was visible for a long distance in all directions.

COLLECTIONS

The several light traps were in operation for various periods during the 4 years. Except for certain unavoidable interruptions, they were operated each night throughout the flight season.² The catches from Cherryvale, Garden City, Hays, and Scottsbluff were forwarded to

² The traps at Manhattan and Lincoln were operated by the writers and those at Cherryvale, Garden City, Hays, and Scottsbluff by the N. Sauer Milling Co., F. A. Wagner, L. C. Aicher, and John Carter, respectively.

either Manhattan or Lincoln, where the material was sorted and the results were tabulated. The number of nights for which records were obtained are as follows: Cherryvale 202, Garden City 612, Hays 214, Manhattan 629, Lincoln 656, and Scottsbluff 406.

It was at first intended to record all species of Phalaenidae taken at the several traps. Limitations of time, however, made it necessary to omit some of the smaller forms and the species of the genus *Catocala*, and in 1937 only species of known economic importance were recorded from Manhattan and Garden City. Altogether 305 species were taken and 525,547 individuals. The total yearly catch of each species for the different localities, together with the flight period, is shown in table 1. The sequence of genera and species is in accordance with McDunnough's Check list of Lepidoptera.³ Much of the material was identified by the writers, largely by comparison with specimens determined by J. F. Gates Clarke, of the Division of Insect Identification, Bureau of Entomology and Plant Quarantine, Rowland R. McElvare, New York City, and the late F. H. Benjamin. Credit is due these men for their careful and painstaking work.

³ McDUNNOUGH, J. CHECK LIST OF THE LEPIDOPTERA OF CANADA AND THE UNITED STATES OF AMERICA. PART I. MACROLEPIDOPTERA. South. Calif. Acad. Sci. Mem. 1,275 pp. 1938.

	156	276	459	428	3,171	892	100	71	2	3	21	679	1,171	2,701	240	413	211
<i>A. venerabilis</i> Walk.																126	38
<i>A. solubilis</i> Harv.																	
<i>A. malefica</i> Guen.	1	2	2	3	2	2		26	34	73	13						
<i>A. ipsaton</i> (Rott.)	131	14	201	53	382	28	93	590	800	1,179	768	107	75	620	24	14	17
<i>F. fulta subnotata</i> (Haw.)	228	1,031	4,344	1,963	1,622	1,737	2,967	52	198	268	341	1,181	6,317	3,487	292	279	186
<i>F. annexa</i> (Tretl.)	90		92	74	53		3	10	3	49	10		9	24	3	61	1
<i>Copadopharum grandis</i> Stkr.									3	5							
<i>C. contraria</i> McD.																	
<i>C. longipenne</i> Grt.																	
<i>C. alba</i> (Harv.)																	
<i>Spaciotis claudestina</i> (Harr.)																	
<i>S. hawila</i> (Grt.)																	
<i>Choeophora fungorum</i> G. and R.	49	2	14						1								
<i>Anicla infecta</i> (Ochs.)																	
<i>Hemicruxa rudens</i> (Harv.)			1	2		1			6	2				1			
<i>and f. pelliculatis</i> (Grt.)			1	2			3			9				11			4
<i>Grassis tendrifera</i> (Walk.)																	
<i>Peridroma margaritosa</i> Haw. and f. <i>saucia</i> (Hbn.).	93	22	761	356	4,683	6	77	49	727	1,055	2,224	455	420	2,197	77	71	337
<i>Caradrina quadragula clemens</i> (Sm.).																	
<i>Amathes c-nigrum</i> (L.).	1	1	47	123	113			1	17	16	50	21	48	197	27	48	8
<i>A. badinodis</i> (Grt.)	39	24	41	26	41	2						12	30	25			
<i>Pseudoglossa olivata</i> (Harv.)																	
<i>Abagrotis placida</i> (Grt.)																	
<i>A. barnesi</i> (Beri.).	1			3	2				3						5	8	10
<i>Rhynchagrotis cupida</i> (Grt.)																	
<i>Uleus satyrus</i> Grt.								3									
<i>U. plicatus</i> Grt.														1			
Hadennae:																	
<i>Scotogramma trifolii</i> (Rott.)	153	861	3,677	1,100	984	89	3,615	498	86,661	75,021	42,325	970	2,492	8,295	1,396	742	403
<i>S. trifolii albifusa</i> (Walk.)																	
<i>Trichoclea postica</i> Sm.										1			2	1			
<i>T. artesia</i> Sm.																	
<i>Pola lepitima</i> (Grt.)	4	6	15	5								1	4			10	
<i>Laciniipolia meditata</i> (Grt.)		3	9	21			76					57	102	28	1	1	3
<i>L. anguina</i> (Grt.)			5														
<i>L. incurva</i> (Sm.)																	
<i>L. vicina</i> (Grt.)		32	43	7	1		101		2	1							
<i>L. renigera</i> (Steph.)		2	23	65	1							211	1,011	60	7	5	3
<i>L. erecia</i> (Walk.)	33	2															
<i>L. lorea</i> (Guen.)													1				
<i>L. olivacea</i> (Morr.)														2			
<i>L. maritima</i> (Harv.)	2	5	8	2													
<i>Anepea capsularis</i> (Guen.)			1	2													
<i>Tricholita signata</i> (Walk.)													1				
<i>Ulotanche culica</i> (Guen.)			2														
<i>U. modesta disticha</i> (Morr.)						2											
<i>Orthodes akalus</i> (Stkr.)						1											
<i>O. incincta</i> (Morr.)																	
<i>O. saturnus</i> (Stkr.)	16	51	104	83	141	10	16		1		36	91	103	117	117	98	9
<i>O. crenulata</i> (Buhl.)	3	2		6									3			8	

TABLE 1.—*Phalaenid moths caught in light traps at various localities in Kansas and Nebraska, 1934-37*—Continued

Species	Flight period	Number of moths caught															
		Cherry-vale, Kans.			Manhattan, Kans.			Hays, Kans.		Garden City, Kans.			Lincoln, Nebr.			Scottsbluff, Nebr.	
		Sept. 8 to Oct. 31, 1935	Mar. 13 to Dec. 5, 1934	Apr. 8 to Oct. 30, 1936	Apr. 12 to Oct. 15, 1937	Sept. 26 to Oct. 30, 1934	Sept. 24 to Nov. 23, 1934	Sept. 20, 1934	Mar. 24 to Nov. 31, 1935	Apr. 10 to Oct. 31, 1936	Apr. 12 to Nov. 12, 1937	Mar. 14 to Oct. 28, 1935	Apr. 9 to Oct. 31, 1936	Apr. 13 to Oct. 30, 1937	July 13 to Oct. 19, 1935	May 3 to Oct. 31, 1936	May 17 to Oct. 18, 1937
Hadoninae—Continued.																	
<i>O. furfurata</i> (Grt.)	May																
<i>Nephelodes emmedonia</i> (Cram.)	Sept.-Oct.	20	22	27	58	116	9	1			1	14	25	51	2	2	3
<i>Xylomyges</i> sp.	May																
<i>Orthostia garmani</i> (Grt.)	Apr				1					1			1	5		1	
<i>O. hibisci</i> f. <i>norm. inscens</i> Walk.	do																
<i>Ceramica picta</i> (Harr.)	Apr.-Oct.	1		9	39	12			6				9	33	102	119	27
<i>Protoparce rubripennis</i> (G. and R.)	Aug.-Sept.	1	15		1												32
<i>P. abilinea</i> (Hbn.)	Mar.-Oct.	139	63	1,440	477	327	33	32	1,023	403	649	339	1,120	413	71	55	15
<i>Leucania phragmatidicola</i> Guen.	Apr.-Oct.	20	8	211	159	266	5	18	98	104	15	36	107	108	17	16	3
<i>L. unipuncta</i> (Haw.)	Apr.-Nov.	1,029	381	4,107	273	3,357	145	613	577	450	2,250	1,035	822	3,140	113	48	252
Cucullinae:																	
<i>Cucullia speyeri</i> Lint	May-June													4		2	
<i>C. latifica</i> Lint.	Apr.-Oct.	4		172	41			71		24	173	3	9	10		2	
<i>C. astroides</i> Guen.	May-Sept.			34	15			1		1		1				2	
<i>Nyctophylla luna</i> (Morr.)	May-Aug.														3	13	59
<i>Oncocnemis ballenta</i> Sm.	Aug.																
<i>O. tricolor</i> Sm.	Sept.-Oct.																
<i>O. satina</i> Sm.	do			1												1	
<i>O. augustus</i> Harv.	do																
<i>O. saundersiana</i> Grt.	Oct.																11
<i>O. occulta</i> (Grt.)	Apr.-Oct.			36	3											481	143
<i>O. riparia</i> Morr.	May-June				10						7			11		146	3
<i>O. sp. nov.</i>	Sept.																2
<i>Homokadena badistriga</i> (Grt.)	June												1		1		
<i>H. stabilis</i> Sm.	May-June			9	43												
<i>H. nigra</i> dinadla Sm.	May-July			58	101										11	19	
<i>Adia chironanthi</i> (A. and S.)	Sept.-Oct.			11	6			18									
<i>Psaphida gressi</i> (Morr.)	Mar.-Aug.			29	16				7							30	1
<i>Eudrype depilis</i> (Grt.)	Apr	4			2				1							53	
<i>E. rolandi</i> (Grt.)	do																
<i>Grapholitha gorgii</i> (Grt.)	Apr.-Oct.												1			4	

[illegible]

TABLE 1.—*Phalaenid moths caught in light traps at various localities in Kansas and Nebraska, 1934-37*—Continued

Species	Flight period	Number of moths caught																			
		Manhattan, Kans.				Hays, Kans.				Garden City, Kans.				Lincoln, Nebr.				Scottsbluff, Nebr.			
		Cherry vale, Kans., Apr. 8 to Oct. 31, 1935	Sept. 8 to Dec. 5, 1934	Mar. 13 to Nov. 1, 1935	Apr. 8 to Oct. 30, 1936	Apr. 12 to Oct. 15, 1936	Sept. 26 to Oct. 30, 1934	Apr. 5 to Oct. 30, 1935	Sept. 23, 1934	Mar. 24 to Nov. 20, 1935	Apr. 10 to Oct. 31, 1936	Apr. 12 to Nov. 12, 1937	Mar. 14 to Oct. 28, 1935	Apr. 9 to Oct. 31, 1936	Apr. 13 to Oct. 30, 1937	July 13 to Oct. 19, 1935	May 3 to Oct. 31, 1936	May 17 to Oct. 18, 1937			
Amphipyridae—Continued.																					
<i>G. partita</i> f. <i>hepara</i> Guen.	Apr.-Oct.			2											11						
<i>Balsa malana</i> (Fitch)	May														55						
<i>Prodenia ornithogalli</i> Guen.	May-Oct.	273	177	1,616	503	1,669	14	197	62	631	165	240	1,683	262	2,352	15	5				
<i>Lophophanes frugiperda</i> (A. and S.).	Aug.-Oct.	23	12	133	31	28	3	5	29	43	17	49	21	3	82	1					
<i>L. ciziqua</i> (Hbn.).	May-Nov.																				
<i>Maqusa orbifera</i> (Walk.).	Aug.-Nov.	5	96	332	340	142	37	48	966	285	201	316	36	59	1,260	11	7	5			
<i>Enargia decolor</i> (Walk.).	July	2	1	3			1		7	6	1										
<i>Cosmia caesecens</i> f. <i>calami</i> (Harv.).	Aug.															2		1			
<i>Aethenia substa</i> Hbn.	Oct.			2																	
<i>Arzania obliqua</i> (Walk.).	July-Aug.																				
<i>Pyrithia umbra</i> (Hufn.).	May-Aug.			4																	
<i>Erythrocia stavis</i> (Hy. Edw.).	May-Sept.		2	15	30			33		60	21			4	1	14	12	3			
<i>Esocia clauda</i> Grt.	June-July																				
<i>Catocala lineolata</i> Walk.	Apr.-Sept.			29	242						1			3	358	8	36	9			
<i>Ogdoconta cinerella</i> (Guen.).	May-Oct.	3		24	163									6	2	1	6	6			
<i>Selicanis cinerella</i> Sm.	Oct.-Nov.	3	28	26	5		31	4	8		2		2	1	30	195	1	1			
<i>Silbidium spumatum</i> Grt.	June-Sept.	12	2	22	35		6	6			8			33	30	223	168	1			
<i>Plagiomimicus pitheochromus</i> Grt.	Aug.-Sept.	14		4	47									3	2						
<i>P. expallidus</i> Grt.	Sept.			1																	
<i>Stiria rugifrons</i> Grt.	May-Oct.	18	13	107	29			91		181	73		1	3	6	155	359	640			
<i>Basileodes pepita</i> Guen.	Aug.-Sept.			7						2	1										
<i>B. chiropsis</i> Grt.	Sept.																				
<i>B. catharops</i> Dyar	do																				
<i>Cirriophanus duplicatus</i> Sm.	June																8				
<i>Stirides perflava</i> (Harv.).	do																				
<i>Pseudocentra crustaria</i> (Morr.).	May-July													2			9	33			
Heliophaninae.																					
<i>Heliophana hirta</i> (Guen.).	Sept.				2																
<i>Heliophana hirta</i> (Guen.).	do		4	1	1																
<i>Heliophana chlorophia</i> (Hbn.).	May-Oct.	1	2	138	263			17	1	20	164		13	87	68	1	24	3			
<i>H. paratara</i> (Grt.).	Apr.-Sept.	2	1	37	59			4		1	8		2	49	33		4				

<i>H. armigera</i> (Hbn.)	Apr.-Nov	180	1,676	1,946	1,695	1,829	1,062	487	701	1,035	1,219	2,186	266	610	2,120	45	25	49
<i>H. virescens</i> (F.)	July-Sept			4	4			4		4	8			4	7	7	6	2
<i>Canthylidia scutosa</i> (Schiff.)	May-Oct		11	6	8		1	2			9		1	2	2	1	7	1
<i>Oryctes citrinellus</i> (G. and R.)	June-Sept			7	1			1		58	2			3	1	1	68	1
<i>Chlorocleptria simplex</i> Sm.	do		1	5	10			7		158	167		1		20	21	42	54
<i>Dasyus poudaeca lucens</i> (Morr.)	do			30	67									3			5	
<i>D. meadi</i> (Grt.)	May-Aug			3	11			4		140	195			4	3	11	233	209
<i>Rhodophora gaurae</i> (A. and S.)	May-Oct			3	11									4	4		5	3
<i>Thyreion rosea</i> Sm.	May-June			94	17			1		1			4	11	14		4	2
<i>Schinia cumatilis</i> (Grt.)	Aug.-Sept				131						6			1			1	2
<i>S. hulsia</i> Tapper	Sept.				7	13									2			4
<i>S. reniformis</i> Sm.	July-Aug																3	4
<i>S. chrysellæ</i> (Grt.)	Aug.-Oct			3	33	5		7			1			9	1		1	9
<i>S. trifascia</i> Hbn.	Aug.														7	4	3	1
<i>S. oleagina</i> Morr.	do					6									3			1
<i>S. gracilentæ</i> Hbn.	Aug.-Oct			42	120			179		13	27		1	3	2		7	4
<i>S. acutilinea</i> (Grt.)	Aug.																	4
<i>S. diffusa</i> Sm.	do					3												3
<i>S. walsinghami</i> (Hy. Edw.)	do																	3
<i>S. aundina</i> (Dru.)	Sept.			1														6
<i>S. tertia</i> (Grt.)	Aug.-Oct	36		2,043	6,624			1,024		17	106		33	334	79	1	4	1
<i>S. albaflascia</i> Sm.	Aug.																	4
<i>S. regia</i> (Stkr.)	July-Sept				1			3		2	50				1	4		4
<i>S. gloriosa</i> (Stkr.)	Sept.	3		5														
<i>S. sanguinea</i> (Geyer)	Aug.					1												
<i>S. saturata</i> (Grt.)	July-Oct			13	268			3			13				2			
<i>S. thoreani</i> (G. and R.)	July-Sept	21		10	53	10												
<i>S. marginata</i> (Haw.)	Aug.-Sept	30		8	4			1			2				2			
<i>S. lora</i> Stkr.	Sept.				6													
<i>S. bifascia</i> Hbn.	Aug.					5					1				4			
<i>S. ultima</i> (Stkr.)	Sept.				10								22					3
<i>S. iaguarina</i> (Guen.)	May-Sept	7		965	73			25		27	16			27	14	8	7	3
<i>S. lyra</i> (Guen.)	June-July													1	1			
<i>S. arcigera</i> (Guen.)	Aug.-Oct	2	1	6	11			1									3	
<i>S. bicuspida</i> Sm.	July-Sept			13	36	2		1			2			1			3	2
<i>S. mortua</i> (Grt.)	Aug.-Sept					3						2					3	3
<i>S. mortua packardii</i> (Grt.)	do	3		8	155			5		15	8				3	16	1	1
<i>S. mortua nobilis</i> (Grt.)	do		1															
<i>S. sordida</i> Sm.	do				27	22												
<i>S. siren</i> (Stkr.)	Aug.-Sept			4	48					3	15							
<i>S. roseincta</i> (Harv.)	June									6							4	1
<i>Trichosellus cupes</i> (Grt.)	June-July				1													
Acontinae:																		
<i>Erasmia synpochitis</i> G. and R.	June-July													2				
<i>E. carnea</i> Guen.	May-Sept			1	77						1			395	314			2
<i>Neorastria apicosa</i> (Haw.)	June-Sept				2									3				
<i>Chamyris cernitha</i> (Trett.)	June													1				
<i>Amynda octo</i> (Guen.)	Sept.																	
<i>Tarachidia erastrioides</i> (Guen.)	May-Oct			1														
<i>T. tenuescens</i> (Sm.)	Aug.														944		22	10
<i>T. semiflava</i> (Guen.)	June																1	1

TABLE 1.—*Phalaenid moths caught in light traps in various localities at Kansas and Nebraska, 1934-37*—Continued

Species	Flight period	Number of moths caught															
		Manhattan, Kans.				Hays, Kans.				Garden City, Kans.				Lincoln, Nebr.			
		Cherry-vale, Kans., Apr. 8 to Oct. 31, 1935	Mar. 13 to Oct. 31, 1935	Apr. 8 to Oct. 31, 1936	Apr. 12 to Oct. 15, 1937	Sept. 26 to Oct. 30, 1934	Apr. 5 to Oct. 30, 1935	Sept. 28 to Nov. 23, 1934	Mar. 24 to Oct. 31, 1936	Apr. 12 to Nov. 12, 1937	Mar. 14 to Oct. 31, 1937	Apr. 13 to Oct. 30, 1937	July 13 to Oct. 19, 1935	May 3 to Oct. 31, 1936	May 17 to Oct. 18, 1937	Scottsbluff, Nebr.	
Acontinae—Continued.	May-Aug											1		53	252		
<i>Theracsa angustipennis</i> (Grt.)	May-Oct		197	501			6					163		59	2		
Eutellinae:	May-June			2										4			
<i>Marathyssa inflexa</i> (Walk.)	Apr-Sept		3	109								95	7	46			
Plusiinae:	Apr											2					
<i>Autographa falcifera</i> (Kby.)	Mar-Oct	10	597	284			72		51	12	363	486	14	7	35		
<i>A. falcifera simplex</i> (Guen.)	Mar-Nov	95	3,512	611		5	124		219	45	1,240	262	195	20	36		
<i>A. brassicae</i> (Riley)	Sept			1			4		2	1	3	17	102		6		
<i>A. corymbifera</i> Geyer	Apr-Oct	6	105	13													
<i>A. biloba</i> (Steph.)	do		2	3													
<i>A. oo</i> (Cram.)	Apr-July		1	3											1		
<i>A. praeclonitis</i> (Guen.)	May-Oct			5													
<i>A. californica</i> (Speyer)	May-June			3													
<i>A. oo</i> (Guen.)	May			2													
<i>Plusia acer</i> (Hbn.)	Apr-Sept		28	154													
<i>Agrotis urens</i> Guen	July-Aug	4															
<i>Paratella bistriaria</i> Hbn.	Mar-Nov	249	63	2,225	636	9	1,447	28	1,069	872	1,997	4,827	67	58	117		
<i>Cucurpina crassiuscula</i> (Haw.) and <i>C. erechtha</i> (Cram.) mixed.	May-Aug			2													
<i>Agrostotris anilis</i> (Dru.)	May-Oct	5	18				1		1					1			
<i>Zale lunata</i> (Dru.)	Apr-July			15										2			
<i>Z. galbanata</i> (Morr.)	May-June													1			
<i>Z. lunifera</i> (Hbn.)	May-Sept	1	1	20	33		2							2			
<i>Matigramma pulverinea</i> Grt.	Sept																
<i>Hendix detrahens</i> (Walk.)	Apr		1														
<i>Phoberia atomaria</i> Hbn.	Mar			4													
<i>Medipolia idonipila</i> (Walk.)	Mar-Oct	6	47				6		117	371	1			1			
<i>M. jucunda</i> Hbn.	May-Aug		1	6		1	1		3	1				3			
<i>Infila deducta</i> (Morr.)			12	18					10	166				20			

	Apr.-Sept.									5	22							1	6	25	6
<i>Drasteria mirifica</i> (Hy. Edw.)																					
<i>Drasteria</i> sp.																				1	
<i>Anticarsia gemmatilis</i> (Hbn.)																					
<i>Srenoloma lunilinea</i> (Grt.)																				1	
<i>Hyporophia monilis</i> (F.)																					
<i>H. hormos</i> Hbn.																					
<i>H. sp. nov.</i>																					
<i>Plusiodonta compressipalpis</i> Guen.																					
<i>Alabama argillacea</i> (Hbn.)																					
Hypeninae:																					
<i>Tathorhynchus angustiorata</i> (Grt.)																					
<i>Plathypena scabra</i> (F.)																					
Hermiinae:																					
<i>Epizeuxis americanalis</i> (Guen.)																					
<i>E. lubricalis</i> Gey.																					
Total	3,674	7,265	37,090	21,094	28,874	5,723	16,703	10,963	113,256	111,686	70,847	11,213	21,957	43,711	5,455	9,599	6,437				
Number of species	76	70	139	151	41	42	73	37	76	88	29	69	113	126	86	157	112				

Although the response to light varies with different species, it is probable that the number of individuals recorded is a fairly reliable estimate of the relative abundance of the species. Approximately 90 percent of the total catch was referable to species of economic importance.

PROPORTION OF THE SEXES

During the course of this study an attempt was made to determine the proportion of the sexes in representative samples of the various species taken. In many cases, however, insufficient material was examined or captured for a reliable estimate to be obtained. The results are presented in table 2. Only those species represented by 25 or more specimens are tabulated, all others being included in one group at the end of the table. Altogether, 36,692 specimens were examined, and of this number 12,902, or 35 percent, were females. It will be seen that the proportion of females varied greatly with different species, ranging from 3 to 60 percent.

TABLE 2.—*Proportion of females among various species of phalaenid moths caught*

Species	Moths examined	Females	Species	Moths examined	Females
	Number	Percent		Number	Percent
<i>Acronicta lepusculina</i> Guen.....	28	36	<i>Nephelodoss emmedonia</i> (Cram.)	32	3
<i>A. parallela</i> (Grt.).....	34	35	<i>Proteoleucania albilinea</i> (Hbn.)	918	21
<i>Euxoa siccata</i> (Sm.).....	256	10	<i>Leucania phragmatidicola</i>		
<i>E. niveilinea</i> (Grt.).....	292	14	Guen.....	170	46
<i>E. immixta</i> (Grt.).....	50	22	<i>L. unipuncta</i> (Haw.).....	4,160	36
<i>E. scandens</i> (Riley).....	40	30	<i>Cucullia laetifica</i> Lint.....	28	25
<i>E. messoria</i> (Harr.).....	31	33	<i>Oncocnemis sanina</i> Sm.....	79	19
<i>E. recticincta</i> (Sm.).....	83	29	<i>O. augustus</i> Harv.....	31	35
<i>E. tessellata</i> (Harr.).....	32	44	<i>O. occata</i> (Grt.).....	25	60
<i>Chorizagrotis auxiliaris</i> (Grt.)	10,672	51	<i>Psaphidia grotei</i> (Morr.).....	26	23
<i>Loragrotis apicalis</i> (Grt.).....	32	31	<i>Crymodes burgesi</i> (Morr.).....	502	6
<i>Onychagrotis rileyaana</i> (Morr.)	262	13	<i>Platyperigea ertima</i> (Walk.)	30	7
<i>Agrotis tetusta</i> Walk.....	293	28	<i>Prodenia ornithogalli</i> Guen.....	1,501	25
<i>A. daedalus</i> (Sm.).....	62	15	<i>Laphygma frugiperda</i> (A. and		
<i>A. orthogonia</i> Morr.....	233	18	S.).....	44	48
<i>A. gladiaria</i> Morr.....	185	27	<i>L. exigua</i> (Hbn.).....	510	32
<i>A. venerabilis</i> Walk.....	984	3	<i>Selicanis cinereola</i> (Guen.).....	67	27
<i>A. malefida</i> Guen.....	35	43	<i>Heliothis paradoxa</i> (Grt.).....	32	56
<i>A. ypsilon</i> (Rott.).....	1,012	32	<i>H. armigera</i> (Hbn.).....	3,149	45
<i>Feltia subgithica</i> (Haw.).....	2,767	7	<i>Schinia mortua packardii</i> (Grt.)	29	21
<i>F. annexa</i> (Treit.).....	138	17	<i>Autographa falcifera simplex</i>		
<i>Peridroma margaritosa saucia</i>			(Guen.).....	59	8
(Hbn.).....	936	36	<i>A. brassicae</i> (Riley).....	528	39
<i>Amathes c-nigrum</i> (L.).....	44	52	<i>Cae n u r g i n a c r a s s i u s c u l a</i>		
<i>A. badinodis</i> (Grt.).....	27	11	(Haw.) and <i>C. erecta</i>		
<i>Scotogramma trifolii</i> (Rott.)	3,550	35	(Cram.) mixed.....	1,019	33
<i>Laciniipolia meditata</i> (Grt.)	74	12	All other species.....	1,133	33
<i>L. vicina</i> (Grt.).....	272	13			
<i>L. renigera</i> (Steph.).....	51	12			
<i>Orthodes incincta</i> (Morr.).....	145	17			
			Total.....	36,692	35

SEASONAL FLIGHT RECORDS

Taken as a group, the light-trap records show that Phalaenidae are in flight from early in March to late in November, with periods of heavy flight and scarcity interspersed irregularly throughout the season. The nightly flight of Phalaenidae in the different localities is shown graphically in figures 3, 4, and 5. The nightly flight at Garden City was much greater than at any of the other localities, owing chiefly to the enormous numbers of adults of *Scotogramma trifolii* captured. The numbers of moths taken varied greatly in the different localities, and from year to year, and also as to the time of the year when the

FIGURE 3.—Nightly flight of Phalaenidae at Garden City, Kans., 1935-37.

FIGURE 4.—Nightly flight of Phalaenidæ at Manhattan, Kans., 1934-37.

period of greatest flight occurs. For example, at Garden City in 1935 the peak flight occurred in July, in 1936 in May, and in 1937 in June.

FIGURE 5.—Nightly flight of Phalaenidae at Lincoln and Scottsbluff, Nebr., and at Hays and Cherryvale, Kans., 1934-37.

In other words, there is no definite pattern in the flight periods of the Phalaenidae as a group, largely because individual species vary greatly in numbers from year to year.

FLIGHT RECORDS OF INDIVIDUAL SPECIES

The species of Phalaenidae fall into two main groups based on their seasonal history, namely, those species producing a single generation annually and those producing more than one generation annually.

As noted previously, species of economic importance comprised 90 percent of the total number of individuals captured. The seasonal flight of several single-generation species of economic importance is shown in figures 6 and 7.

In general, the flight of individual species showed the same trend at all the localities, the chief difference being in numbers taken. Where no striking differences were noted only one locality was plotted on the charts. It will be seen that the species producing a single generation annually have a rather short flight period, usually late in August, extending through September and into the early part of October. The notable exception is *Chorizagrotis auxiliaris*. The enormous flight of this species in May and June suddenly subsides and the species reappears again in September, after an estivation period of about 2 months. The fall flight is greatly reduced in numbers. This species is normally confined to the semiarid regions, but occasionally occurs in abundance at Manhattan (fig. 6, *B*) and at Lincoln. The records for the individual species show the same trend, namely, years of abundance usually followed by years of scarcity.

The seasonal flight of several species producing multiple generations annually is shown in figures 8, 9, 10, and 11. The flight records alone do not delineate clearly the various generations owing to overlapping as the season progresses. The charts, however, bring out some points of interest with reference to the seasonal abundance of multiple-generation species. *Heliothis armigera* (fig. 10, *A*) is shown to be scarce early in the season, with a marked increase in abundance in the fall. With *Prodenia ornithogalli* (fig. 10, *B*) and *Peridroma margaritosa saucia* (fig. 11, *A*) the peak of abundance is reached in midseason, followed by comparative scarcity in the fall. *Proteleucania albilinea* (fig. 11, *B*) shows two distinct flight periods, one early in May and the other late in August or early in September.

SUMMARY

Light traps were operated for various periods in six localities in Kansas and Nebraska during the period 1934-37, in order to obtain information on the flight periods and abundance of owl moths (Phalaenidae). An attempt was made to operate the traps each night during the flight season, which extended from March to November. A total of 2,719 nightly records were made.

During the 4-year period 305 species of Phalaenidae were taken, totaling more than 525,000 individuals. Approximately 90 percent of the specimens taken were species of economic importance. The different species taken in each locality are shown, together with the total yearly flight, the flight period, and the proportion of the sexes. More than 36,000 individuals were examined for sex, and it was found that 35 percent were females. The sex ratio varied greatly with the species.

The seasonal flight records are presented graphically for the Phalaenidae as a group and for several species of economic importance. The numbers of moths taken differed greatly in the different localities,

FIGURE 6.—Seasonal flight of *Chorizagrotis auxiliaris* at Garden City, Kans. (A), and at Manhattan, Kans. (B), 1935-37.

FIGURE 7.—Seasonal flight of *Feltia subgothica* at Manhattan, Kans. (A), and at Lincoln, Nebr. (B), *Agrotis venerabilis* at Lincoln (C), *A. gladiaria* at Manhattan (D), *A. orthogonia* at Garden City, Kans. (E), and *Euxoa niveilinea* at Garden City (F), 1934-37.

FIGURE 8.—Seasonal flight of *Scotogramma trifolii* at Garden City, Kans., 1935-37.

FIGURE 9.—Seasonal flight of *Caenurgina erecta* and *C. crassiuscula* (A), and *Leucania unipuncta* (B), at Manhattan, Kans., 1935-37.

FIGURE 10.—Seasonal flight of *Heliothis armigera* at Manhattan, Kans. (A), and of *Prodenia ornithogalli* at Manhattan (B), and at Lincoln, Nebr. (C). 1935-37.

FIGURE 11.—Seasonal flight of *Peridroma margaritosa saucia* (A) and *Protoleucania albilinea* (B) at Manhattan, Kans., 1935-37.

and from year to year, and also as to the time of year when the greatest period of flight occurred. In general, however, flights of individual species showed the same trend in all the localities.

Species having a single generation annually had a rather short flight period, usually from late August to early October. *Chorizagrotis auxiliaris*, an exception to this rule, had an enormous flight in May and June and a much lighter flight in the fall.

Owing to overlapping of broods, the flights of the different generations of multiple-generation species were not clearly defined. *Heliothis armigera* was scarce early in the season, with a marked increase in abundance in the fall. Peak abundance of *Prodenia ornithogalli* and *Peridroma margaritosa saucia* occurred in midseason, followed by comparative scarcity in the fall. *Proteleucania albilinea* showed two distinct flight periods, one early in May and the other late in August or early in September.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE WHEN THIS PUBLICATION WAS EITHER FIRST PRINTED OR LAST REVISED

<i>Secretary of Agriculture</i>	CLAUDE R. WICKARD.
<i>Under Secretary</i>	PAUL H. APPLEBY.
<i>Assistant Secretary</i>	GROVER B. HILL.
<i>Chief, Bureau of Agricultural Economics</i>	HOWARD R. TOLLEY.
<i>Director of Agricultural War Relations</i>	SAM BLEDSOE.
<i>Director of Extension Work</i>	M. L. WILSON.
<i>Director of Finance</i>	W. A. JUMP.
<i>Director of Foreign Agricultural Relations</i>	L. A. WHEELER.
<i>Director of Information</i>	MORSE SALISBURY.
<i>Director of Personnel</i>	T. ROY REID.
<i>Land Use Coordinator</i>	ERNEST H. WICKING.
<i>Librarian</i>	RALPH R. SHAW.
<i>Solicitor</i>	ROBERT H. SHIELDS.
<i>Chief, Office of Civilian Conservation Corps Activities</i>	FRED MORRELL.
<i>Chief, Office of Plant and Operations</i>	ARTHUR B. THATCHER.
<i>Administrator of Agricultural Marketing</i>	ROY F. HENDRICKSON.
<i>Administrator of Agricultural Conservation and Adjustment.</i>	M. CLIFFORD TOWNSEND.
<i>Chief, Agricultural Adjustment Agency</i>	FRED S. WALLACE.
<i>Chief, Soil Conservation Service</i>	HUGH H. BENNETT.
<i>Manager, Federal Crop Insurance Corporation</i>	LEROY K. SMITH.
<i>Chief, Sugar Division</i>	JOSHUA BERNHARDT.
<i>Administrator of Agricultural Research</i>	E. C. AUCHTER.
<i>Chief, Bureau of Animal Industry</i>	JOHN R. MOHLER.
<i>Chief, Bureau of Agricultural Chemistry and Engineering.</i>	W. W. SKINNER, <i>Acting.</i>
<i>Chief, Bureau of Dairy Industry</i>	O. E. REED.
<i>Chief, Bureau of Entomology and Plant Quarantine.</i>	P. N. ANNAND.
<i>Chief, Office of Experiment Stations</i>	JAMES T. JARDINE.
<i>Chief, Bureau of Plant Industry</i>	R. M. SALTER.
<i>Chief, Bureau of Home Economics</i>	LOUISE STANLEY.
<i>Superintendent, Beltsville Research Center</i>	C. A. LOGAN.
<i>President, Commodity Credit Corporation</i>	J. B. HUTSON.
<i>Administrator of Farm Security Administration</i>	C. B. BALDWIN.
<i>Governor of Farm Credit Administration</i>	ALBERT G. BLACK.
<i>Chief, Forest Service</i>	EARLE H. CLAPP, <i>Acting.</i>
<i>Administrator, Rural Electrification Administration</i>	HARRY SLATTERY.

This bulletin is a contribution from

<i>Bureau of Entomology and Plant Quarantine</i> ..	P. N. ANNAND, <i>Chief.</i>
<i>Division of Cereal and Forage Insect Investigations.</i>	C. M. PACKARD, <i>Principal Entomologist, in Charge.</i>