**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☒ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☒ Additional comments / Commentaires supplémentaires:

Pagination is as follows: p.43-49.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x		14x		18x		22x		26x		30x	
	12x		16x		20x		24x		28x		32x

The copy filmed here has been reproduced thanks to the generosity of:

Legislative Library
Victoria

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

1	2	3
---	---	---

1	2
4	5

ks

L'exemplaire filmé fut reproduit grâce à la
générosité de:

Legislative Library
Victoria

Les images suivantes ont été reproduites avec le
plus grand soin, compte tenu de la condition et
de la netteté de l'exemplaire filmé, et en
conformité avec les conditions du contrat de
filmage.

d

-

9

Les exemplaires originaux dont la couverture en
papier est imprimée sont filmés en commençant
par le premier plat et en terminant soit par la
dernière page qui comporte une empreinte
d'impression ou d'illustration, soit par le second
plat, selon le cas. Tous les autres exemplaires
originaux sont filmés en commençant par la
première page qui comporte une empreinte
d'impression ou d'illustration et en terminant par
la dernière page qui comporte une telle
empreinte.

Un des symboles suivants apparaît sur la
dernière image de chaque microfiche, selon le
cas: le symbole → signifie "A SUIVRE", le
symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être
filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être
reproduit en un seul cliché, il est filmé à partir
de l'angle supérieur gauche, de gauche à droite,
et de haut en bas, en prenant le nombre
d'images nécessaire. Les diagrammes suivants
illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5000 - Fax

570
FROM THE TRANSACTIONS OF THE ROYAL SOCIETY OF CANADA
SERIES III 1915 VOLUME IX

The Swarming of *Odontosyllis*

by

C. McLEAN FRASER, Ph.D.

OTTAWA

PRINTED FOR THE ROYAL SOCIETY OF CANADA

5.14
41

595.14
F891

SECTION IV., 1915

[43]

TRANS. R.S.C.

The Swarming of Odontosyllis.

By C. McLEAN FRASER, PH.D.

(From the Pacific Coast Biological Station, Departure Bay, B.C.)

Presented by DR. A. B. MACCULLUM, F.R.S.C.

(Read May Meeting, 1915.)

On Feb. 10, 1913, Mr. F. A. Potts of Trinity Hall, Cambridge, read a paper on "The Swarming of *Odontosyllis*," at a meeting of the Cambridge Philosophical Society, which paper appeared in the Proceedings on April 23, following. Observations made on *Odontosyllis phosphorea* Moore, a species of Annelid found in the vicinity of the Biological Station, Departure Bay, B.C., provided material for the body of the paper, to which is added a comparison of the habits of this species with that of other Syllids and Nereids.

The observations made and recorded were of much interest and the conclusions appeared to be so but as is often the case when conclusions are based on very limited observations, further examination shows the necessity of considerable revision.

As far as the phenomenon of swarming is concerned, repeated observations confirm all of Mr. Potts' statements without producing any of value in addition. The time of day at which spawning takes place seems definite without doubt, but definite as regards position of the sun, not definite as to the time on the clock. Very few have been seen very much before sunset, but from sunset or possibly a little before it until almost dusk, that is for a period lasting from half an hour to an hour, they appear at the surface whether the sun sets early or late.

There is nothing further to indicate that the males are attracted to the females as they are in *O. enopla*. In every instance each individual whether male or female comes to the surface without any apparent regard to the position of any other individual. The movement at the surface seems just as liable to be away from as towards the nearest individual of the other sex. It would therefore seem that although the number of individuals is great, the chance for fertilizing all of the ova, or even a large portion of them, is very slight. The chances might be increased if the eggs were pelagic at or near the surface but they are not so. They may remain suspended for some time in water that is

107091

PACIFIC COAST BIOLOGICAL STATION
VICTORIA, B.C.

somewhat agitated but if placed in a receptacle where the water remains still they soon sink to the bottom. Since the water of the sea seems the most favorable for spawning when there is scarcely a ripple on the surface the eggs surely must be fertilized soon after ejection or not at all. No doubt this seeming waste is at least partially obviated by the movements of the Annelids themselves. As the female circles around with a motion that has been aptly described by an observer as a "wavy wiggle," the eggs are scattered right and left over an area relatively large. At the same time the male with a similar undulatory movement scatters the sperm, giving thereby a strong impetus to the movement of the very active spermatozoa. In the denser part of the swarm therefore, where these circles of distribution overlap, the chance for the fertilization of each ovum is not so slim as it would at first appear.

All of this then may be observed in the one evening if conditions are favorable and was observed by Mr. Potts. Numerous repetitions have merely substantiated the fact that the routine of the evening in which he was an interested spectator, is practically the same as that on any other evening during which the spawning occurs. With regard to the distribution of *Odontosyllis* in space and of its spawning in time, the limited extent of his observations did not serve as a satisfactory basis for his rather sweeping conclusions, but it is quite true that he makes a proviso for this in his paper.

Taking in the first place the geographic distribution of *Odontosyllis*, as Mr. Potts surmised, the limit was by no means reached in the much circumscribed area near Snake and Five Finger Islands. The Hexactinellid sponge referred to, as well as a number of other siliceous species are present over a wide area in the Strait of Georgia, an area of which we have not as yet found the limits. It has been traced in a south-easterly-northwesterly direction, i.e., running the same general direction as the strait and parallel to the eastern shore of Vancouver Island and the other outlying islands, to the northwest as far as Northwest Bay and Ballenas Islands, at least 15 miles from the original location and to the channels between Lasqueti and Texada Islands, 20 miles or more in a slightly different direction, while to the southeast they are just as plentiful around Gabriola Reefs and Breakwater Island, at least 15 miles away in that direction. Furthermore, *Odontosyllis* has been found in a somewhat different kind of bottom in the channels between the islands east of Vancouver Island to the south of the Station. In some of these instances the depth was much less than that indicated in the drawing of the originally described area. Dredging has not been done at a distance of much more than 25 miles from the Station in this direction but this Annelid has been found in the most distant of these dredgings. It would seem, therefore, that instead

of having a restricted distribution, it is one of the most widely dispersed species to be found in the vicinity. As to its presence or absence in Departure Bay I have nothing definite to record. If it is not to be found in the bay it is not because the bay is too shallow as the species has been found at a less depth than is found here in many places. The bottom in general and especially in the deeper parts is different from that where *Odontosyllis* has been found and it may be for that reason that it is not present if it is not. Since I have been especially interested in the distribution of this species we have dredged very little in the bay. Possibly if a diligent search were made it might be found. The reason for the swarming at the entrance to the bay it will be considered to better advantage after the time of swarming has received some attention. To this we shall now turn.

The swarms of *O. phosphorea* were first noticed in 1911 on August 15 and 16 and in 1912 on August 18. From this Mr. Potts concludes that there is a possibility of a periodicity in the case of *O. phosphorea* similar to that of the Palolo worms. If he had carefully considered the data before him he should not have come to such a hasty conclusion. According to his own remarks on *Nereis osawai* of Japan, after reading Izuka's paper on the subject "Their date of appearance is absolutely fixed for the days following the new moon." I believe the species found in the Gulf of Mexico has an even more restricted periodicity which is also absolutely fixed for a date definitely related to the time of the new moon. In 1911 full moon appeared on Aug. 9 and the last quarter on Aug. 17, Aug. 15 and 16, when the swarms of *Odontosyllis* were observed, were consequently nearly the end of the third phase. In 1912 there was new moon on Aug. 12, the first quarter falling on Aug. 19, consequently Aug. 18 is near the end of the first phase. Large swarms were also seen on Sept. 5. Since the last quarter of the moon came on Sept. 4, this date is near the beginning of the fourth phase.

In 1913 observations were made over somewhat the same period as in 1912, and dates nearly the same as those in 1912 were obtained. This approximation seemed to make it worth while to go into the matter much more fully in 1914. This was more readily accomplished since Dr. A. Willey, who was at the station for a portion of the summer, became interested in the outcome also. Either one or both of us made examination of the locality in which *Odontosyllis* had been found swarming, at more or less regular intervals from the middle of June until the middle of December. The accompanying table shows the results of these examinations as well as those for the two preceding years. As the phases of the moon are given it will be seen at a glance that there is no indication of periodicity in the swarming but appar-

ently it may take place at any time within a period of three or four months. There are gaps between the dates, it is true, but none are of very great length and the search in every case, particularly if none or few were found, was sufficiently prolonged to get accurate information as far as the area in question is concerned.

TABLE OF OBSERVATIONS.

Date	Result of Observation	Time	High tide	Low tide	Moon's Phases	
					New	Full
1912						
Aug. 18	Plentiful	19 30	16 37	23 24		
20	None		18 25	1 50		
21	None		19 30	3 02		
27						*
Sept. 5	Plentiful	19 00	18 13	0 39	*	
10						
1913						
Aug. 2					*	
11	Few	19 30	15 39	20 00		
14	Few		17 45	23 30		
15	None		18 08	23 51		
16						*
21	Numerous		20 06	13 48		
22	Numerous		20 27	14 22		
28	Few		16 26	21 24	*	
31						
Sept. 6	Numerous	19 00	20 47	15 34		
13	None		17 09	23 15		*
15						
1914						
June 18	None					
23	None					
July 1	None					
18	None				*	
22	None					
25	Few	20 00	20 00	12 48		
30	Numerous		22 29	16 42		
31	Numerous		23 11	17 56		
Aug. 1	None		15 42	19 36		
3	Few		17 15	22 23		*
5						
14	Numerous	19 30	21 47	9 16		
19	Numerous		17 27	22 44		
20	Few		17 49	23 26	*	
21						
29	Plentiful	19 00	22 29	18 30		*
Sept. 4						
5	Plentiful		18 10	0 34	*	
19						
21	Very plentiful	18 30	18 00	0 43		*
22	Three		18 30	1 28		*
Oct. 3						
6	Numerous	17 30	17 42	0 46		
8	Numerous		18 17	13 44	*	
18						
27	Few		13 19	20 21		*
Nov. 2						
4	Five (3m & 2f)	17 30	16 33	0 10		
11	One	17 00	12 22	19 10		
12	None		12 56	19 50		
17	One female		15 40	23 06	*	
24	None					
Dec. 7	None					
16	None					

With regard to the time of tide Mr. Potts says, "The tide was full or just falling." This is scarcely accurate. On Aug. 15, 1911, it was low tide at 14.53 and high at 21.10, on Aug. 16, low tide at 15.34 and high at 21.40. The 1912 tides are given in the table. Sunset in the two cases in 1911 occurs well on in the rising tide and on Aug. 18, 1912 at about half tide on the fall. It seems more nearly correct to say that the largest swarms are found at high tide or somewhat before that time, as far as the area at the entrance of Departure Bay is concerned (See table). A possible reason for this lies in the fact that the flood tide comes in from the south. The current is divided by Protection and Newcastle Islands, part going through Nanaimo Harbor and Newcastle Channel to reach Departure Bay. The outer current is less obstructed and as usual arrives at the north end of Newcastle Island before the other current, but the distance it works into the bay varies, apparently depending more upon the direction and force of the wind than on any other factor. Unless the surface of the water is very smooth the line where the two currents meet can readily be traced across the entrance of the bay. The outside current would tend to carry these annelids, coming from the deep in the Strait of Georgia, forward with it until the other current meets it and as there is practically still water here (the doldrums on a small scale) the individuals become collected into a swarm and it is here they are found the most plentifully. When the water is very smooth the area may be very much extended but on ordinary occasions searching for them outside of the entrance to the bay has always produced negative results. This would account for the greater numbers being found on the rising tide. If the water is very smooth during the height of the spawning period they are probably found at any time of the tide. A smooth surface seems to be one of the requisites for swarming. I have never been able to find any at the surface even when the surface was but moderately disturbed. After some days of storm, they have been found on the next smooth evening to be present in very large numbers. It may readily be that at times during the spawning period, when it has not been possible to find any or many at the entrance of the bay, on account of tide and current conditions, they may have drifted in some other direction and at some other smooth spot they may be plentiful.

I have no doubt since *Odontosyllis* is found over such a wide area in dredged material, that the phenomenon of swarming occurs at other places as well as at the entrance to Departure Bay, as especially among the islands to the south there is an endless variety of current conditions and in this variety there must be some that suit. The swarming takes place at such an awkward time of the day for making observations at a

distance from the station that none of these places have so far been discovered

The district along the east coast of Vancouver Island is very abundantly supplied with polychaetes, both as to variety of species and to number of individuals. It must be that many of these come to the surface to spawn and possibly some of them swarm as does *Odontosyllis*. There is not much chance of finding these otherwise than by accident unless a study is made of them throughout the year so that some idea may be obtained as to when the swarms should be looked for. As yet nothing has been done except for a short time in the summer. Only one case, apart from *O. phosphorea*, has come under my notice. On Sept. 30, 1913, late in the afternoon, probably about 5 o'clock, a male specimen of *Nephtysorca* (Fabricius) was found swimming at the surface near the station float. Apparently it came to the surface to spawn but no others were seen then or at any time since.

The polychaetes serve as a main article of diet for many of the flat-fish, hence the extensive and intensive study of their life-histories is of importance from an economic as well as from a purely biological point of view. It is quite true that little attention has yet been paid to any of the flatfish of the Pacific with the exception of the halibut but many other species are of just as good flavor (in my opinion, much better) as the halibut, although of course they are much smaller. Since they are probably abundant in various localities, they must receive attention some day and the sooner the better.