**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

A number line starting at $10x$ and ending at $32x$. The line is divided into 12 equal segments by 13 tick marks. Labels are placed above and below the line: $10x$, $12x$, $14x$, $16x$, $18x$, $20x$, $22x$, $24x$, $26x$, $28x$, $30x$, and $32x$. An arrow points to the tick mark between $22x$ and $24x$, which represents $23x$.

The copy filmed here has been reproduced thanks to the generosity of:

Stauffer Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Stauffer Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482-0300 - Phone
(716) 288-5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

Vol. VII: CRUSTACEA

PART A: DECAPOD CRUSTACEANS

By MARY J. RATHBURN.

SOUTHERN PARTY—1913-16

G
670
1913 C2
v. 6, Pt 1
c3

OTTAWA
J. de LABROQUERIE TACHÉ
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1919

Issued August 18, 1919

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

Vol. VII: CRUSTACEA

PART A: DECAPOD CRUSTACEANS

By MARY J. RATHBURN.

SOUTHERN PARTY—1913-16

OTTAWA
J. de LABROQUERIE TACHÉ
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1919

Vol. iii—46454—1

Issued August 18, 1910

The Decapod Crustaceans of the Canadian Arctic Expedition 1913-18.

By MARY J. RATHBUN,

Associate in Zoology, United States National Museum.

The known range of several species is extended by the specimens obtained by the Canadian Arctic Expedition. Five of the species of *Spirontocaris* are shown to have a more complete circumpolar distribution, while *Paralithodes camtschaticus* is recorded for the first time from the Arctic (occurrence possibly accidental), and *Pagurus brandti* seems to reach as far north in Bering sea as Norton sound. Attention is called to the distributional charts of von Hofsten (see bibliography) in some of which notable gaps are filled or reduced by the new data here published.

As nearly all of the species of shrimps collected by the Canadian Arctic Expedition were described and their synonymy worked out in the author's report on the decapod crustaceans of the Harriman Alaska expedition (see bibliography), a reference to that work is made under each of those species in the list below.

It may be noted that the care with which the specimens were obtained and preserved is due to Mr. Frits Johansen, naturalist of the expedition.

LIST OF THE SPECIES.

Order DECAPODA.

Suborder NATANTIA.

Superfamily PANDALOIDA.

Family PANDALIDÆ.

Genus *Pandalus* Leach.

Pandalus goniurus Stimpson.

Rathbun, 1904, p. 38, pl. 1, fig. 3.)

Off point Lav, Alaska lat. 69° 35' N., long. 163° 27' W.; 11-12 fathoms; rock and shells; August 17, 1913; station 22; twelve specimens.

DISTRIBUTION: East of Alaska southward through Bering sea to Okhotsk sea and to Japan. 3-100 fathoms.

Superfamily PALÆMONOIDA.

Family HIPPOLYTIDÆ.

Genus *Spirontocaris* Bate.

Spirontocaris groenlandica (J. C. Fabricius).

(90-101)

West of Cockburn point, 15-20 fathoms; sandy mud, with shells; three specimens.

Strait, Northwest Territories; September 14, 1915; station

Bernard harbour, Northwest Territories; from stomach of *Erigonathus barbatus*; October 22, 1915; station 42a; three specimens.

DISTRIBUTION.—Arctic coast of America; Bering sea to Puget sound; Kamchatka; Okhotsk sea; Atlantic coast of America from East and West Greenland to Narragansett bay, Rhode island; Norway. 1-117 fathoms.

***Spirontocaris solna* (Sowerby).**

(Rathbun, 1904, p. 64, pl. III, fig. 5.)

North of the Alaskan boundary; lat. $70^{\circ} 13' N.$, long. $140^{\circ} 50' W.$; about 30 fathoms mud; from stomach of *Phoca hispida*; April 4, 1914; station 29f. One specimen and fragments of three others.

Bernard harbour, Northwest Territories; from stomach of *Erigonathus barbatus*; October 22, 1915; station 42a; two specimens.

DISTRIBUTION.—From Arctic Alaska (off point Hope), through ring strait, Bering sea, Aleutian islands and Alaska peninsula to Lituya bay, Alaska. Arctic North America— from Ellesmere Land, Jones sound, and Melville peninsula, eastward to Greenland, Iceland, and Nova Zembla, and southward to Massachusetts bay, Faeroe islands, and Kattegat; 3-240 fathoms.

For table of distribution, see von Hofsten, 1916, fig. 1.

The two localities of the Canadian Arctic Expedition are at wide intervals between those previously recorded in Alaska and Arctic Canada.

***Spirontocaris arcuata* Rathbun.**

(Rathbun, 1904, p. 64, pl. III, fig. 4.)

West of Cockburn point, Dolphin and Union strait, Northwest Territories 15-20 fathoms; sandy mud, with stones and algae; September 14, 1915; station 43c; one specimen.

DISTRIBUTION.—Hitherto known only from Diabolof islands, Bering sea to strait of Juan de Fuca. Its occurrence in Arctic Canada considerably extends the range.

***Spirontocaris phiquetsii* (Kröyer).**

(Rathbun, 1904, p. 70.)

West of Cockburn point, Dolphin and Union strait, Northwest Territories 15-20 fathoms; sandy mud, with stones and algae; September 14, 1915; station 43c; three specimens.

Bernard harbour, Northwest Territories; from stomach of *Erigonathus barbatus*; October 22, 1915; station 42a; anterior half of one specimen.

DISTRIBUTION.—Arctic Alaska (west of point Franklin) southward to Plover bay, Siberia, and to the Shumagins, Alaska peninsula; Ellesmere Land and Jones sound to Eastern Greenland, southward via Labrador to cape Cod, Iceland, the Faeroes, and Spitzbergen, eastward to Kara strait and southward to Skagerak; Middle Arctic Siberia; shallow water to 200 fathoms.

For table of distribution, see von Hofsten, 1916, fig. 2.

The two localities of the Canadian Arctic Expedition are near together and remote from those on either side which have been previously recorded.

***Spirontocaris polaris* (Sabine).**

(Rathbun, 1904, p. 73.)

Off point Lay, Arctic Alaska; lat. $69^{\circ} 35' N.$, long. $163^{\circ} 27' W.$; 11-12 fathoms; rock and sand, with many algae; August 17, 1913; station 22; one specimen.

West of Cockburn point, Dolphin and Union strait, Northwest Territories; 15-20 fathoms; sandy mud, with stones and algae; September 11, 1915; station 43c; one specimen, with copepod parasite in left branchial chamber.

Outer harbour, Bernard harbour, Northwest Territories; about 5 fathoms; sandy mud, with algae; July 20, 1915; station 41; five specimens.

Outer harbour, Bernard harbour, Northwest Territories; about 5 fathoms; mud with algae (mostly loose); July 28, 1915; station 41c; three specimens (1 ♂ 2 ♀). The male and one female have purplish antennal scales and purple dot-markings on antennae, eye-stalks, and legs; telson more orange, carapace flammate.

Outer harbour, Bernard harbour, Northwest Territories; about 3 fathoms; mud with brown algae; August 1, 1915; station 41f, one ovigerous female, one young.

Bernard harbour, Northwest Territories; from stomach of big ♂ *Erignathus barbatus*; September 3, 1915; station 42c; one specimen.

Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; October 22, 1915; station 42a; one specimen.

DISTRIBUTION.—From Arctic ocean, north of Bering strait, southward to the Aleutian islands and Kodiak on the American side, and to Okhotsk sea on the Asiatic side; Arctic America from Melville island southward to Cape Cod, Massachusetts, and eastward via Greenland and Iceland to New Siberia islands; southward on the European coast to the Hebrides and Skagerrak; 1-553 fathoms.

For table of distribution, see von Hofsten, 1916, fig. 3.

The two general localities where this species was taken by the Canadian Expedition are intermediate between those previously recorded.

***Splrontocaris fabricii* (Kröyer).**

(Rathbun, 1904, p. 85.)

Off point Lay, Arctic Alaska; lat. 69° 35' N., long. 163° 27' W.; 11-12 fathoms; rock and sand, with many algae; August 17, 1913; station 22; two ♂ ovigerous.

West of Cockburn point, Dolphin and Union strait, Northwest Territories; 15-20 fathoms; sandy mud, with stones and algae; September 14, 1915; station 43c; twenty-three specimens.

DISTRIBUTION.—Arctic coast of Alaska southward through Bering sea to Okhotsk sea and Alaska (Cook inlet); Arctic Canada (see above); Atlantic coast of America from West Greenland southward to Massachusetts bay. Low water to 100 fathoms.

***Splrontocaris galmardii belcheri* (Bell).**

(Rathbun, 1904, p. 86, pl. III, figs. 3, 3a.)

Off Icy cape, Arctic Alaska; lat. 70° 24' N., long. 161° 25' W.; 9-10 fathoms; mud, with pebbles; August 19, 1913; station 23; one ovigerous ♀ and fragments of one ♂, one ♀.

Cape St. Stephen bay, Dolphin and Union strait, Northwest Territories; 25-30 fathoms; sandy mud, with pebbles; September 14, 1915; station 43b; one ♂.

DISTRIBUTION.—Nova Zembla, Kara sea, Arctic Siberia and Alaska, thence south to Sitka; Dolphin and Union strait, Arctic Canada (see above); Western Greenland and Labrador; 3½-37 fathoms.

The distribution of the species, *S. galmardii*, with all its variations is much more extensive (see von Hofsten, 1916, fig. 4), including Eastern Greenland, Iceland and Northern Europe, thence southward to Scotland and Kiel bay.

Superfamily CRAGONOIDÆ.

Family CRAGONIDÆ.

Genus *Crago* Lamarek.*Crago septemspinosus* (Say).(*Cragon septemspinosa* Rathbun, 1904, p. 116, text-fig. 55.)

Near Grantley harbour, Port Clarence, Alaska; 2-3 fathoms; mud, with many algæ; August 4, 1913; station 20*g*; one specimen.

DISTRIBUTION.—Arctic coast of Alaska at Eschscholtz bay southward along the eastern shore of Bering sea to the Shumagins; east coast of North America from eastern Florida (Say) northward; shallow water to 435 fathoms (off Delaware).

Genus *Sclerocrangon* Sars.*Sclerocrangon boreas* (Phipps).

(Rathbun, 1904, p. 133.)

Off point Lay, Arctic Alaska; lat. $69^{\circ} 35' N.$, long. $163^{\circ} 27' W.$; 11-12 fathoms; rock and sand, with many algæ; August 17, 1913; station 22; thirty-one specimens.

Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9-10 fathoms; mud, with pebbles; August 19, 1913; station 23; one small specimen.

West of Cockburn point, Dolphin and Union strait, Northwest Territories; 15-20 fathoms; sandy mud, with stones and algæ; September 14, 1915; station 43*c*; six specimens.

Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; July 6-8, 1915; station 40*a*; fragments of one specimen.

Bernard harbour, Northwest Territories; from stomach of big ♀ *Erignathus barbatus*; September 3, 1915; station 42*c*; fragments of three specimens.

Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; August 24, 1914; station 37*a*; third abdominal somite of a very large specimen.

DISTRIBUTION.—Arctic Siberia near Bering strait, and Arctic Alaska, southward via Bering sea to Killisnoo, Alaska, and strait of Georgia, British Columbia (Lenz); Sakhalin island and vicinity, Eastern Siberia; islands of Arctic America southward to cape Cod; Greenland, Iceland, Northern Europe, eastward to Kara sea and southward to the Faeroe islands and to lat. $67^{\circ} 15' N.$ on the coast of Norway; 0 to 200 fathoms.

For table of distribution, see von Hofsten, 1916, fig. 7.

Genus *Nectocrangon* Brandt.*Nectocrangon lar* (Owen).

(Rathbun, 1904, p. 137, text-figs. 74 and 75.)

Off point Lay, Arctic Alaska; lat. $69^{\circ} 35' N.$, long. $163^{\circ} 27' W.$; 11-12 fathoms; rock and sand, with many algæ; August 17, 1913; station 22; two specimens.

Near Grantley harbour, Port Clarence, Alaska; 2-3 fathoms; mud, with many algæ; August 4, 1913; station 20*g*; one specimen.

DISTRIBUTION.—Arctic coast of Alaska and Siberia southward to Sitka and Kurile islands; Western and Eastern Greenland southward to Nova Scotia; 0 to 220 fathoms.

Genus **Sabinea** Owen.

Sabinea septemcarinata Sabine.

(Hansen, 1908, p. 52.)

North of the Alaskan boundary; lat. $70^{\circ} 13' N.$, long. $140^{\circ} 50' W.$; from stomach of *Phoca foetida*; depth of water about 30 fathoms; mud; April 4, 1914; station 29f; fragments of two specimens.

Off Stapyllton bay, Dolphin and Union strait, Northwest Territories; 25-30 fathoms; sandy mud, with pebbles, no algæ; September 14, 1915; station 43b; four small specimens.

Off Cockburn point, Dolphin and Union strait, Northwest Territories; about 50 fathoms; mud, with pebbles, but no algæ; September 13, 1915; station 43a; five specimens.

Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; July 6-8, 1915; station 40a; one specimen.

Bernard harbour, Northwest Territories; from stomach of big ♀ *Erignathus barbatus*; September 3, 1915; station 42c; one specimen.

Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; October 22, 1915; station 42a; anterior half of one specimen.

DISTRIBUTION.—Arctic America from the Alaskan boundary eastward to the American islands, thence southward to cape Cod; western and eastern Greenland, Iceland, Faeroes, northern Europe and Siberia, eastward to long. $170^{\circ} 17' E.$, and southward to Saltenfjord, Norway; 0 to 246 fathoms.

For table of distribution, see von Hofsten, 1916, fig. 8.

Suborder REPTANTIA.

Tribe ANOMURA.

Superfamily PAGURIDEA.

Family PAGURIDÆ.

Genus **Pagurus** Fabricius.

Pagurus trigonocheirus (Stimpson).

(Stimpson, Proc. Acad. Nat. Sci. Philadelphia, vol. X, 1858, p. 249 [87].)

Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9-10 fathoms; mud, with pebbles, but no algæ; August 19, 1913; station 23; one ♂ ovigerous, one young.

DISTRIBUTION.—Arctic coast of Alaska and Siberia southward through Bering sea to Aleutian islands and Kamchatka; 3 to 100 fathoms.

Pagurus capillatus (Benedict).

(Benedict, Proc. U. S. Nat. Mus., vol. XV, 1892, p. 8. Rathbun, 1904, p. 157, pl. IV, fig. 3.)

Near Grantley harbour, port Clarence, Alaska; 2-3 fathoms; mud, with many algæ; August 4, 1913; station 20g; one specimen.

DISTRIBUTION.—Arctic ocean southward through Bering strait to Kamchatka and California (lat. $36^{\circ} 55' N.$); 3 to 240 fathoms.

Pagurus brandti (Benedict).

(Benedict, Proc. U. S. Nat. Mus., vol. XV, 1892, p. 9. Rathbun, 1904, p. 157, pl. IV, fig. 4.)

Norton sound, Alaska; lat. $63^{\circ} 43' N.$, long. $165^{\circ} 24' W.$; surface; July 8, 1913; station 19; one right chela.

DISTRIBUTION.—Bering sea southward to Oregon; 9 to 121 fathoms; hitherto the latitude of the Pribilof islands was the farthest north known for this species. The specimen from Norton sound, if the identification be correct, extends the range.

Pagurus splendescens Owen.

(Owen, in Zool. of Capt. Beechey's Voyage, 1825-28, London, 1839, p. 81, pl. XXV, figs. 1, 1a.)

Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9-10 fathoms; mud, with pebbles, but no algae; August 19, 1913; station 23; two small specimens.

DISTRIBUTION.—Arctic coast of Alaska (point Barrow) westward through Bering sea to Kamchatka and Washington; below low water to 225 fathoms.

Pagurus sp. indet.

Off cape Lisburne, Arctic Alaska; lat. $68^{\circ} 48' N.$, long. $165^{\circ} 10' W.$; surface; August 16, 1913; station 21*d, e, f*; two specimens of the megalops stage.

Family **LITHODIDÆ**.Genus **Paralithodes** Brandt.**Paralithodes camtschaticus** (Tilesius).

(Bouvier, Ann. Sci. Nat., Zool., ser. 8, vol. I, 1896, p. 23.)

Point Barrow, Alaska; on sand spit; beach; August 22, 1913; station 24; propodus of left cheliped of a specimen of medium size.

DISTRIBUTION.—Point Barrow, Arctic Alaska (see above); Bering sea (from Norton sound) to Aleutian islands to Orea, Prince William sound; Skidegate, Queen Charlotte islands, British Columbia (Geological Survey of Canada); Kamchatka; Okhotsk sea; Japan; low tide to 49 fathoms.

Tribe **BRACHYURA**.Subtribe **BRACHYGNATHA**.Superfamily **BRACHYRHYNCHA**.Family **ATELECYCLIDÆ**.Genus **Telmessus** White.**Telmessus cheiragonus** (Tilesius).

(Benedict, Proc. U. S. Nat. Mus., vol. XV, 1892, p. 224, pl. XXV, pl. XXVI, figs. 2-4.)

Beach at Teller, Port Clarence, Alaska; July 31, 1913; station 20*d*; one ♂, two ♀, and five carapaces.

Beach at Nome, Alaska; July 12, 1913; station 20*o*; one ♀ and five carapaces.

DISTRIBUTION.—Northeastern Siberia; Kamchatka; Kurile islands; Bering sea to California (Holmes); low water to 20 fathoms.

Family **CANCRIDÆ.**Genus **Cancer** Linnaeus.**Cancer magister** Dana.

(Dana, Proc. Acad. Nat. Sci. Philadelphia, 1852, p. 73; Crust. U. S. Exploring Exped., part 1, 1852 p. 151; atlas, 1855, pl. VII, figs. ta d.)

Beach at Orca (Cordova), Southeastern Alaska; September 5, 1916; station 60a; one carapace.

DISTRIBUTION.—Unalaska to Magdalena bay, Lower California; low water to 50 fathoms.

Superfamily *ONYRHYNCHIA.*

Family **INACHIDÆ.**Genus **Hyas** Leach.**Hyas coarctatus** Leach.

(Rathbun, Proc. U. S. Nat. Mus., vol. XVI, 1893, p. 69.)

Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9–10 fathoms; mud, with pebbles, but no algae; August 19, 1913; station 23; four ♂, four ♀.

Langton bay (East of Mackenzie river), Northwest Territories; October 26, 1910; Dr. R. M. Anderson; fifteen specimens (Amer. Mus. Nat. Hist.), two of which were taken in the net in 8 fathoms of water. A note accompanying the crabs states that they are known to the Kotzebue sound Eskimos as 'Pu-tu-ri-ak,' but are unknown to the people east of the Mackenzie.¹

DISTRIBUTION (including *Hyas coarctatus alutaceus* Brandt).—Arctic Siberia to Langton bay; Bering sea to Aleutian islands; Kamchatka; Okhotsk sea to Korea (lat. $37^{\circ} 02' N.$); Baffins bay and Eastern Greenland to Hudson strait and bay; and southward to off cape Hatteras, North Carolina; Iceland; Northern Europe southward to about lat. $49.5^{\circ} N.$ (Dons); shallow water to 906 fathoms.²

Various larvæ from the following stations have been referred to this species:

Off Cooper island, point Barrow, Alaska; surface; August 27, 1913; stations 25 b, c; three zoeæ.

Off cape Lisburne, Arctic Alaska; lat. $68^{\circ} 48' N.$, long. $165^{\circ} 10' W.$; surface; August 16, 1913; stations 21 c, d, e, f; three specimens of zoeæ, more than twenty-five specimens of megalopa, first form, with three-spined front, eight specimens of megalopa which may be a development from the preceding and from which the median or rostral spine has disappeared.

Off Nunivak island, Bering sea; lat. $60^{\circ} 09' N.$, long. $167^{\circ} 38' W.$; surface; July 6, 1913; about 50 zoeæ.

South of Shumagin islands; lat. $54^{\circ} 30' N.$, long. $159^{\circ} 42' W.$; surface; July 1, 1913; stations 13 a, b, c; three megalopa, second form.

The zoeæ and first megalops resemble those of *Hyas araneus* "araneus" described and figured by Williamson, Fisheries Board, Scotland, Sci. Invest., 1909, I (Dec., 1910), pp. 15–16, pl. 1, figs. 1 and 2. Williamson considered that his megalopa moulted into the first crab stage. The rostrum of a similar megalops, but with longer median spine, is figured as *H. coarctatus* by Bjorck, Acta Reg. Soc. Physiog. Lundensis, n.f., vol. XXIV, 1913, p. 22, text-figs. 1 and 2.

¹These specimens have been identified by Dr. W. G. Van Name and the information communicated by Dr. Roy W. Mearns, of the American Museum of Natural History.

²A station label indicating this great depth accompanies the specimen so recorded (see Hansen, Danish Ingolf-Expedition, III, 2, 1903, p. 16), but an error may, of course, have been made at the time of collection.

Genus *Chionæcetes* Krøyer.*Chionæcetes opilio* (O. Fabricius).

(Rathbun, Proc. U. S. Nat. Mus., vol. XVI, 1893, p. 74, pl. IV, figs. 5-7.)

Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9-10 fathoms; mud, with pebbles, but no algae; August 19, 1913; station 23; 2 young ♂, one young ♀.

According to Mr. Frits Johansen (in a letter) Dr. R. M. Anderson reports that several years ago one of the Eskimos caught a big "devil-crab" or "spider-crab" in a gill-net at Herschel island.

Dr. Roy W. Miner writes that there is in the American Museum of Natural History a specimen of *Chionæcetes opilio* collected at point Barrow, Alaska, by V. Stefansson in July or August, 1912.

DISTRIBUTION.—From British Columbia northward to Bering strait, thence eastward to Greenland, thence southward to Casco bay, Maine; Kamchatka; Okhotsk sea (off Robben island); Nagasaki, Japan; littoral to 350 fathoms.

LIST OF LOCALITIES WITH THE DECAPODS FOUND AT EACH.

Station 13a, b, c. South of Shumagin islands; lat. $54^{\circ} 30' N.$, long. $159^{\circ} 42' W.$; surface; July 1, 1913.

Hyas coarctatus Leach. Larvæ.

Station 19. Norton sound, Alaska; lat. $63^{\circ} 43' N.$, long. $165^{\circ} 24' W.$; surface; July 8, 1913.

Pagurus brandti (Benedict).

Station 20d. Beach at Teller, port Clarence, Alaska; July 31, 1913.

Telmessus cheiragonus (Tilesius).

Station 20g. Near Grantley harbour, port Clarence, Alaska; 2-3 fathoms; mud, with many algae; August 4, 1913.

Crago septemspinosus (Say).

Nectocrangon lar (Owen).

Pagurus capillatus (Benedict).

Station 20e. Beach at Nome, Alaska; July 12, 1913.

Telmessus cheiragonus (Tilesius).

Station 21c. Off cape Lisburne, Arctic Alaska; lat. $68^{\circ} 48' N.$, long. $165^{\circ} 10' W.$; surface; August 16, 1913.

Hyas coarctatus Leach. Larvæ.

Station 21d, e, f. Off cape Lisburne, Arctic Alaska; lat. $68^{\circ} 48' N.$; $165^{\circ} 10' W.$; surface; August 16, 1913.

Pagurus sp. indet. Larvæ.

Hyas coarctatus Leach. Larvæ.

Station 22. Off point Lay, Arctic Alaska; lat. $69^{\circ} 35' N.$, long. $163^{\circ} 27' W.$; 11-12 fathoms; rock and sand, with algae; August 17, 1913.

Pandalus gonurus Stimpson.

Spirontocaris polaris (Sabine).

Spirontocaris fabricii (Krøyer).

Sclerocrangon boreas (Phipps).

Nectocrangon lar (Owen).

Station 23. Off Icy cape, Arctic Alaska; lat. $70^{\circ} 24' N.$, long. $161^{\circ} 25' W.$; 9-10 fathoms; mud, with pebbles, but no algae; August 19, 1913.

Spirontocaris gaimardii belcheri (Bell).

Sclerocrangon boreas (Phipps).

Pagurus trigonocheirus (Stimpson).

Pagurus splendescens Owen.

Hyas coarctatus Leach.

Chionacetes opilio (O. Fabricius).

Station 24. Point Barrow, Alaska; on sand spit; beach; August 22, 1913.

Paralithodes camtschaticus (Tilesius).

Stations 25b, c. Off Cooper island, point Barrow, Alaska; surface; August 27, 1913.

Hyas coarctatus Leach. Larvæ.

Station 29f. North of Alaskan boundary; lat. $70^{\circ} 13' N.$, long. $110^{\circ} 50' W.$; from stomach of *Phoca hispida*; about 30 fathoms; mud; April 4, 1914.

Spirontocaris spina (Sowerby).

Sabinea septemcarinata Sabine.

Station 37a. Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; August 21, 1914.

Sclerocrangon boreas (Phipps).

Station 40a. Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; July 6-8, 1915.

Sclerocrangon boreas (Phipps).

Sabinea septemcarinata Sabine.

Station 41. Outer harbour, Bernard harbour, Northwest Territories; about 5 fathoms; sandy mud, with algæ; July 20, 1915.

Spirontocaris polaris (Sabine).

Station 41c. Outer harbour, Bernard harbour, Northwest Territories; about 5 fathoms; mud, with algæ (mostly loose); July 28, 1915.

Spirontocaris polaris (Sabine).

Station 41f. Outer harbour, Bernard harbour, Northwest Territories; about 3 fathoms; mud, with brown algæ; August 1, 1915.

Spirontocaris polaris (Sabine).

Station 42c. Bernard harbour, Northwest Territories; from stomach of big ♀ *Erignathus barbatus*; September 3, 1915.

Spirontocaris polaris (Sabine).

Sclerocrangon boreas (Phipps).

Sabinea septemcarinata Sabine.

Station 42a. Bernard harbour, Northwest Territories; from stomach of *Erignathus barbatus*; October 22, 1915.

Spirontocaris grælandica (J. C. Fabricius).

Spirontocaris spina (Sowerby).

Spirontocaris phippisii (Krøyer).

Spirontocaris polaris (Sabine).

Sabinea septemcarinata Sabine.

Station 43a. Off Cockburn point, Dolphin and Union strait, Northwest Territories; about 50 fathoms; mud, with pebbles, but no algæ; September 13, 1915.

Sabinea septemcarinata Sabine.

Station 43b. Off Stapylton bay, Dolphin and Union strait, North Territories; 25-30 fathoms; sandy mud, with pebbles, no algæ; September 14, 1915.

Spirontocaris gaimardii belcheri (Bell).

Sabinea septemcarinata Sabine.

Station 43c. West of Cockburn point, Dolphin and Union strait, Northwest Territories; 15-20 fathoms; sandy mud, with stones and algæ; September 14, 1915.

Spirontocaris grælandica (J. C. Fabricius).

Spirontocaris arcuata Rathbun.

Spirontocaris phippisii (Krøyer).

Spirontocaris polaris (Sabine).

Spirontocaris fabricii (Krøyer).

Sclerocrangon boreas (Phipps).

Station 60a. Beach at Orea (Cordova), Southeastern Alaska; September 5, 1916.

Cancer magister Dana.

Off Nuniyak island, Bering sea; lat. 60° 09' N., long. 167° 38' W., surface; July 6, 1913.

Hyas coarctatus Leach. Larva.

BIBLIOGRAPHY.

Appelhof, A.

Die Dekapoden Crustaceen.

Meeresfauna von Bergen, Heft 2 u. 3, 1906.

Birula, A.

Beiträge zur Kenntnis der Dekapoden-Krebse der eurasiatischen Arctis.

Mém. de l'Acad. Imp. Sci. St. Pétersbourg, ser. 8, vol. XXIX, 1910.

Dollein, F.

Die dekapoden Krebse der arktischen Meere.

Fauna Arctica, Band I, Lief. 2, Jena, 1900.

Dons, Carl.

Slegten Hyas.

Tromsø Museums Aarshefter 34, Tromsø, 1912-13.

Dons, Carl.

Nord-Norges Decapoder.

Tromsø Museums Aarshefter 37, Tromsø, 1914 (1915).

Hansen, H. J.

The Danish Ingolf-Expedition. Vol. III. 2. Crustacea Malacostraca. 1. Copenhagen, 1908.

Hofsten, Nils von.

Zoologische Ergebnisse der Schwedischen Expedition nach Spitzbergen 1908. Teil II. 7

Die Dekapoden Crustaceen des Eisfjords.

Kongl. Svenska Vetensk. Akad. Handl., Band 54, No. 7, Stockholm, 1916.

Murdoch, John.

Marine Invertebrates, in P. H. Ray, Report of the International Polar Expedition to Point Barrow, Alaska.

House of Representatives, 48th Congress, 2d Session, Ex. Doc. No. 44, Washington, 1885.

Ohlin, Axel.

Bidrag till Kännedom om Malakostrakfaunan i Baffin Bay och Smith Sound.

Acta Reg. Soc. Physiog. Lundensis, Handlingar, ny följd, Bd. 6, Lund, 1895.

Rathbun, Mary J.

Decapod Crustaceans of the Northwest Coast of North America.

Harriman Alaska Expedition, vol. X, New York, 1904.

Stappers, Louis.

Des d'Orléans. Campagne Arctique de 1907. Crustacés Malacostracés, par le Dr. Louis Stappers. Bruxelles, 1911.

Stephensen, K.

Report on the Malacostraca Pycnogonida and some Entomostraca collected by the "Danmark" Expedition to North-east Greenland.

Meddelelser om Grønland, vol. xlv, København, 1912.

Stephensen, K.

Grønlands Krebsdyr og Pycnogonider (Conspectus Crustaceorum et Pycnogonidorum Groenlandiae).

Meddelelser om Grønland, vol. XXII, København, 1913.

Hitherto unpublished Records of Decapod Crustaceans collected by other Canadian Expeditions and Explorers.¹

By MARY J. RATHBUN.

The material from which this list was made is in the Museum at Ottawa, and was sent to the United States National Museum for identification.

Spirontocaris grandaevia (J. C. Fabricius).

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4 12 specimens.
 Melville island (winter harbour), Northwest Territories, May, 1909. "Arctic"
 Expedition, 1908-9. 1 fragment.
 Fullerton, west side of Hudson bay, Northwest Territories, September 19, 1904.
 "Neptune" Expedition. 1 specimen.

Spirontocaris spina (Sowerby).

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4. 1 specimen.

Spirontocaris liljeborgii (Danielsson).

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4. 1 specimen.

Spirontocaris phippisii (Kröyer).

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4 23 specimens.
 Near mouth of Povungnituk river, east side of Hudson bay, August, 1898.
 A. P. Low. 1 specimen.

Spirontocaris polaris (Sabine).

- Fullerton, west side of Hudson bay, Northwest Territories, September 19, 1904.
 "Neptune" Expedition. 1 specimen.
 Port Burwell, Ungava, Northwest Territories, 1903-4. "Neptune" Expedition
 East of Blacklead island, Cumberland gulf, Northwest Territories; from stomach
 of *Cottus (Myoxocephalus) grandaevius* (Cuv. & Val.), September 4, 1904.
 "Neptune" Expedition. 6 specimens.
 Near mouth of Povungnituk river, east side of Hudson bay, August, 1898.
 A. P. Low. 1 specimen.
 2 specimens.

Spirontocaris fabricii (Kröyer).

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4. 17 specimens.
 Near mouth of Povungnituk river, east side Hudson bay, August, 1898. A. P. Low. 1 specimen.

Spirontocaris guinardii Milne Edwards, varying towards *S. g. bulcheri* Bell.

- Port Burwell, Ungava. "Neptune" Expedition, 1903-4. 20 specimens.
 Fullerton, west side of Hudson bay. "Neptune" Expedition, 1903-4 1 specimen.

Sclerocrangon boreas (Phipps).

- Ponds inlet, north side of Cockburn island, June, 1907. "Arctic" Expedition. 1 specimen.
 Winter harbour, Melville island, Northwest Territories, May 28, 1909. "Arctic"
 Expedition. 1 specimen.
 "Neptune" Expedition, (?), 1903-4 1 soft shell.
 Port Burwell, Ungava, Northwest Territories. "Neptune" Expedition, 1903-4. 1 specimen.

¹For additional records already published, see Whiteaves, J. F., Catalogue of the Marine Invertebrata of Eastern Canada. Published by the Geological Survey of Canada, Ottawa, 1901. Also various lists published in the "Diana" Expedition report and in the Geological Survey of Canada publications.

Canadian Arctic Expedition, 1913-18

Pagurus kroyeri Stimpson.

Ungava Bay, Northwest Territories, August or September, 1897. Hudson Bay Expedition. "Diana" Expedition 3 specimens.

Hyas coarctatus Leach.

Ungava Bay, Northwest Territories, August or September, 1897. Hudson Bay Expedition. "Diana" Expedition 1 male.

Some of the Decapod-Crustacea collected by the "Neptune" Expedition, 1903-4, were sent in 1905 to Prof. G. O. Sars, Christiania, Norway, for identification; and are published in the report of the Canadian Fisheries Museum, Ottawa, for 1905 (Department of Marine and Fisheries, Ottawa, Canada).

As these specimens are still in Christiania, they have not been examined in the U. S. National Museum, Washington, nor included in the list above.

Professor Sars's determinations follow:—

<i>Spirontocaris aculeata</i> (Fabricius).....	Fullerton, Northwest Territories.
<i>Spirontocaris guimardi</i> (Kröyer).....	"
<i>Nectoceraugon lar</i> (Kröyer).....	Port "Burwell, Ungava.

imens.

le.

lition,
iden-
seum,
ned in

itories.
"ngava.