**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☐ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☐ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☐ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	12x	14x	16x	18x	20x	22x	24x	26x	28x	30x	32x
								✓			

The copy filmed here has been reproduced thanks to the generosity of:

Stauffer Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

1	2	3
---	---	---

:

1	2
4	5

L'exemplaire filmé fut reproduit grâce à la générosité de:

Stauffer Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaît sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME III: INSECTS

PART B: NEUROPTEROID INSECTS

By NATHAN BANKS

SOUTHERN PARTY—1913-16

G
670
1913
C2t
v.003
pt.B
c.3

OTTAWA
J. de LABROQUERIE TACHÉ
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1918

Vol. iii—54539—1

Issued July 11th, 1919

The Neuropteroid Insects of the Canadian Arctic Expedition, 1913-18.

By NATHAN BANKS.

The Neuropteroid insects taken by the Canadian Arctic Expedition, 1913-16, belong to seven species, two of which are herewith described as new. Most of the specimens collected by Mr. Frits Johansen belong to one species which is common in the Arctic regions.

PERLIDAE.

***Alloperla pacifica* Banks.**

Two from Ketchikan, southern Alaska, September 10, 1916.

***Nemoura*, sp.**

Several specimens, quite possibly of a new species, from Bernard harbour, Northwest Territories, June 30, 1916, and July 30, 1916. They are in poor condition and cannot now be described.

***Capnia nearctica*, n. sp.**

Black. Body long and slender. Pronotum plainly broader than long, a little broader behind than in front, with a deep transverse groove in front parallel to the front margin. Wings in the male not reaching one-fourth way to the tip, in the female reaching to the tip of body. Setae about one-half the length of the body. In the male the third segment from the tip has a rather large, rounded, median elevation, the surface of which is deeply pitted and provided with short hairs, and behind over the last two segments is a median groove bordered by nearly parallel elevated ridges; in the posterior part the ridges swell out a little; the groove is fully two and a half times as long as broad.

Length, 3 mm.

Many specimens from Bernard harbour, Northwest Territories, Canadian Arctic Expedition, June 25, 1915 (F. Johansen, coll.). Type in Canadian National Collection of Insects, Ottawa, paratype in Museum of Comparative Zoology, Cambridge, Mass. Differs from other species in the shape and sculpture of the tubercle near tip of body.

TRICHOPTERA.

***Rhyacophila alberta* Banks.**

One from Ketchikan, southern Alaska, September 10, 1916.

***Chilostigma præterita* Walker.**

Many specimens. All taken at Bernard harbour, Northwest Territories, September 23, 1915. There is much variation in the extent of the markings on the front wings almost uniformly dark-coloured. Described from Canada. I have seen it from several northern localities and it also occurs in the more northern parts of Europe.

Anabolia emarginata, n. sp.

Black with black bristles, some yellowish bristles on vertex and thorax; antennae narrowly annulate with pale; legs yellowish brown, spines black. Fore-wings blackish, outer and posterior portions with scattered small pale spots, a distinct whitish hyaline spot over the thyridium and one below it near the end of the anal vein; the hind wings are greyish, darker on the anterior tip; genitalia dark. Venation much as in *A. nigricula* but the wing is rather shorter so that the apical cells are noticeably shorter than in that species. Male genitalia, seen from above show two upper appendages close together each with an excised tip; from the side the lower appendage is seen to be tipped with two little black teeth. In general similar to *A. nigricula* but the shape of the superior appendages is different, and they are shorter than in that species.

Expanse, 26 mm.

Two males from Teller, Alaska, July 29, 1913 (F. Johansen, coll.) Type in Canadian National Collection of Insects, Ottawa, paratype in Museum of Comparative Zoology, Cambridge, Mass.

Limnephilus, sp.

One female from Nome, Alaska, August 21, 1916 (F. Johansen, coll.) may be the female of *L. harrimani* Bks., known only from the male. *L. harrimani* is closely related to *L. sitchensis*, and so is this form. The body is greyish brown; vertex and thoracic vittae dull black, latter separated by a grey streak; vertex between lateral ocelli and eyes grey pollinose; palpi, antennae and legs yellowish, latter with black spines, basal joints of antennae dark; genitalia pale. Fore wings yellowish brown, marked with darker brown and whitish hyaline much as in *L. sitchensis*; an oblique mark across the thyridial cell, and a large area each side of the anastomosis hyaline, this spot extends over bases of apical and first subapical cells and the tips of discal and thyridial cells. The dark brown is mostly on the posterior part, and along the veins; veins pale, anastomosis and lower stigmal vein darker; hind wings grey hyaline, veins pale. Face with yellowish hair in the middle and black on the sides, vertex, pronotum and mesonotum mostly with yellowish hair; some macrochaetae on vertex dark. Fore wings have the shape of *L. sitchensis*; the discal cell is plainly longer than its pedicel; the first apical is wider at base than the second, the fourth is about one-third as wide at base as the third or fifth; in the hind wings the fourth apical cell is slightly pedicellate. The superior appendages of female are very long, slender, and divaricate, longer than in *L. sitchensis*, the inferior appendage is much shorter than in that species.

Expanse, 29 mm.

NOTE.—The collection also included a number of immature stages of Neuropteroids, the identification of which was not possible.

EXPLANATION OF PLATE.

- Fig. 1. Dorsal, posterior and lateral views of male genitalia of *Anabolia emeryi*.
 2. Dorsal view of tip of male abdomen of *Capnia nearctica*.
 3. Lateral view of tip of male abdomen of *Capnia nearctica*.
 4. Lateral and dorsal views of female genitalia of *Limnephilus* sp.