Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF ENTOMOLOGY—BULLETIN No. 67.

L. O. HOWARD, Entomologist and Chief of Bureau.

PROCEEDINGS

OF THE

NINETEENTH ANNUAL MEETING

OF THE

ASSOCIATION OF ECONOMIC ENTOMOLOGISTS.

ISSUED NOVEMBER 25, 1907.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1907.

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF ENTOMOLOGY—BULLETIN No. 67.

L. O. HOWARD, Entomologist and Chief of Bureau.

PROCEEDINGS
OF THE
NINETEENTH ANNUAL MEETING
OF THE
ASSOCIATION OF ECONOMIC ENTOMOLOGISTS.

ISSUED NOVEMBER 25, 1907.

WASHINGTON:
GOVERNMENT PRINTING OFFICE,
1907.

BUREAU OF ENTOMOLOGY.

L. O. HOWARD, *Entomologist and Chief of Bureau.*

C. L. MARLATT, *Entomologist and Acting Chief in absence of Chief.*

R. S. CLIFTON, *Chief Clerk.*

F. H. CHITTENDEN, *in charge of breeding experiments.*
A. D. HOPKINS, *in charge of forest insect investigations.*
W. D. HUNTER, *in charge of cotton boll weevil investigations.*
F. M. WEBSTER, *in charge of cereal and forage plant insect investigations.*
A. L. QUAINANCE, *in charge of deciduous fruit insect investigations.*
E. F. PHILLIPS, *in charge of apiculture.*
D. M. ROGERS, *in charge of gipsy moth and brown-tail moth work.*
A. W. MORRILL, *engaged in white fly investigations.*
W. F. FISKE, *in charge of gipsy moth laboratory.*
W. A. HOOKER, *engaged in cattle tick life history investigations.*
A. C. MORGAN, *engaged in tobacco insect investigations.*
R. S. WOGLUM, *engaged in hydrocyanic acid gas investigations.*
C. J. GILLISS, *engaged in silk investigations.*
R. P. CURRIE, *assistant in charge of editorial work.*
MABEL COLCORD, *librarian.*

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,

BUREAU OF ENTOMOLOGY,

Washington, D. C., July 20, 1907.

SIR: I have the honor to transmit herewith the manuscripts of the proceedings of the nineteenth annual meeting of the Association of Economic Entomologists, held at New York, N. Y., December 28 and 29, 1906. This association, made up largely of entomologists connected with the State experiment stations, the Bureau of Entomology of this Department, and others engaged officially in entomological work, is an important one, and the proceedings of its annual meetings contain papers and discussions on injurious and beneficial insects which should by all means be put in print. I therefore recommend that, following the custom of previous years, the proceedings of this meeting be published as Bulletin No. 67 of this Bureau. The two plates and six text figures are necessary for proper illustration of the text.

Respectfully,

L. O. HOWARD,

Entomologist and Chief of Bureau.

HON. JAMES WILSON,

Secretary of Agriculture.

CONTENTS.

NINETEENTH ANNUAL MEETING OF THE ASSOCIATION OF ECONOMIC ENTOMOLOGISTS.

	Page.
A Great Experiment in Applied Entomology.....A. H. Kirkland..	10
Report of Committee on Nomenclature.....	10
Report of Committee on Cooperative Testing of Insecticides.....	12
Insect Notes from Minnesota for 1906.....F. L. Washburn..	13
Report of Committee on National Control of Introduced Insect Pests....	19
A New Oriental Insect Pest (?) in Massachusetts ^aH. T. Fernald..	22
Occurrence of the Gipsy Moth in Connecticut.....W. E. Britton..	22
Notes on Fumigation and Dipping of Nursery Trees. ^a T. B. Symons and A. B. Gahan..	26
The Willow Borer as a Nursery Pest (<i>Cryptorhynchus lapathi</i> L.). W. J. Schoene..	27
Effects of Sprays on Aphis Eggs.....H. E. Hodgkins..	29
Manner of Birth of the Woolly Aphis of the Apple (<i>Schizoneura lanigera</i> Hausm.), and of other Aphididae.....W. E. Rumsey..	31
Unusual Insect Happenings in New Jersey in 1906.....John B. Smith..	34
Miscellaneous Insect Notes from Maryland for 1906. A. B. Gahan and G. P. Weldon..	37
Notes on Insects of the Year 1906 in New York State.....E. P. Felt..	39
The Pear Blister Mite (<i>Eriophyes pyri</i> (Pgst.) Nal.).....P. J. Parrott..	43
Some Observations on the Natural Checks of the Cottony Maple Scale (<i>Pulvinaria innumerabilis</i> Rathv.).....Edgar L. Dickerson..	48
Remarks on Methods Used in Codling Moth Experiments.....A. F. Burgess..	53
The Control of the Codling Moth in the Arid Regions.....E. D. Ball..	55
What Research in Economic Entomology is Legitimate under the Adams Act.....E. Dwight Sanderson..	77
Notes on a New Sawfly Attacking the Peach (<i>Pamphilius persicum</i> MacG.).....B. H. Walden..	85
Economic Work Against the Howard Scale in Colorado (<i>Aspidiotus</i> <i>howardii</i> Ckll.).....E. P. Taylor..	87
The Maple Leaf Stem-Borer (<i>Priophorus acericaulis</i> (MacG.)) ^a W. E. Britton..	94
The Value of Parasites in Cereal and Forage Crop Production. F. M. Webster..	94
Some Georgia Insects During 1906.....R. I. Smith..	101
Observations on Insect Enemies of Tobacco in Florida in 1905. W. A. Hooker..	106
Observations on Cecidomyiidae ^aE. P. Felt..	112
A Spray Nozzle for the Mechanical Mixture of Oil with Water or Other Liquids.....E. Dwight Sanderson..	112
The Beet Leaf-Hopper (<i>Eutettix stricta</i> Ball) ^aE. D. Ball..	117
Problems in Economic Entomology in the Philippines ^aC. S. Banks..	117

^a Withdrawn for publication elsewhere.

	Page.
Occurrence of the Throat Bot in Cuba (<i>Gastrophilus nasalis</i> L.).	
<i>C. F. Baker</i> __	117
The Remarkable Habits of an Important Predaceous Fly (<i>Ceratopogon</i> <i>eriophorus</i> Will.)-----	<i>C. F. Baker</i> __ 117
Some Notes on Leprosy in Havana-----	<i>C. F. Baker</i> __ 118
On the Eradication of the Black Currant Gall-Mite (<i>Eriophyes ribis</i> (Nalepa))-----	<i>Walter E. Collinge</i> __ 119
Destruction of Mosquitoes in Dwellings by the Powders of Chrysanthemum Spread Therein by Means of Hand Bellows or a Towel.	
<i>A. L. Herrera</i> __	123
Notes on Insects in Central Alberta-----	<i>P. B. Gregson</i> __ 125
Report of Committee on Nominations-----	127
Report of Committee on Membership-----	127
Report of Committee on Resolutions-----	127
List of Members of the Association of Economic Entomologists-----	130
Index-----	135

ILLUSTRATIONS.

PLATES.

	Page.
PLATE I. Fig. 1.—Peach branch, showing how leaves have been eaten by larvæ of the sawfly <i>Pamphilius persicum</i> . Fig. 2.—Peach leaves rolled by larvæ of the sawfly <i>Pamphilius persicum</i> . Eggs on midribs near base of leaves.....	86

TEXT FIGURES.

FIG. 1. Chart to illustrate relative value of early sprayings on the first brood of "worms" of the codling moth.....	58
2. Chart to illustrate relative value of early sprayings on the second brood of "worms" of the codling moth.....	60
3. Chart to illustrate relative value of early and late sprayings for the codling moth.....	64
4. The condition of the calyx cup of the apple in relation to spraying for the codling moth.....	71
5. Wheat straws injured by the joint-worm (<i>Isosoma tritici</i>) from which the joint-worms have been removed by some beneficial animal, perhaps the short-tail shrew (<i>Blarina brevicauda</i>).....	100
6. Diagram of spray nozzle for the mechanical mixture of oil with water or other liquids.....	114
7. Compressed-air pump, suitable for spraying a mechanical mixture of oil and water, when fitted with the nozzle herein described...	115

THE NINETEENTH ANNUAL MEETING OF THE ASSOCIATION OF ECONOMIC ENTOMOLOGISTS.

MORNING SESSION, FRIDAY, DECEMBER 28, 1906.

The association met in room 516, Schermerhorn Hall, Columbia University, New York, N. Y., at 10 a. m. An average attendance of over 75 members and visitors was present at each session on Friday and Saturday. The following members attended the meeting:

E. A. Back, Amherst, Mass.; E. D. Ball, Logan, Utah; Rev. C. J. S. Bethune, Guelph, Canada; William Beutenmuller, New York, N. Y.; F. C. Bishopp, Washington, D. C.; W. E. Britton, New Haven, Conn.; F. E. Brooks, Morgantown, W. Va.; A. F. Burgess, Columbus, Ohio; F. H. Chittenden, Washington, D. C.; R. S. Clifton, Washington, D. C.; J. H. Comstock, Ithaca, N. Y.; Mel. T. Cook, New York, N. Y.; D. W. Coquillett, Washington, D. C.; C. R. Crosby, Ithaca, N. Y.; E. L. Dickerson, New Brunswick, N. J.; H. G. Dyar, Washington, D. C.; E. P. Felt, Albany, N. Y.; H. T. Fernald, Amherst, Mass.; James Fletcher, Ottawa, Canada; H. L. Frost, Arlington, Mass.; A. B. Gahan, College Park, Md.; J. B. Garrett, Baton Rouge, La.; J. A. Grossbeck, New Brunswick, N. J.; T. J. Headlee, Durham, N. H.; W. E. Hinds, Washington, D. C.; H. E. Hodgkiss, Geneva, N. Y.; W. A. Hooker, Washington, D. C.; A. D. Hopkins, Washington, D. C.; C. O. Houghton, Newark, Del.; L. O. Howard, Washington, D. C.; W. D. Hunter, Washington, D. C.; G. B. King, Lawrence, Mass.; A. H. Kirkland, Boston, Mass.; D. K. McMillan, Harrisburg, Pa.; C. L. Marlatt, Washington, D. C.; A. C. Morgan, Washington, D. C.; Dudley Moulton, Washington, D. C.; Herbert Osborn, Columbus, Ohio; P. J. Parrott, Geneva, N. Y.; E. F. Phillips, Washington, D. C.; J. L. Phillips, Blacksburg, Va.; A. L. Quaintance, Washington, D. C.; F. W. Rane, Boston, Mass.; W. E. Rumsey, Morgantown, W. Va.; J. G. Sanders, Washington, D. C.; E. D. Sanderson, Durham, N. H.; William Saunders, Ottawa, Canada; Franklin Sherman, jr., Raleigh, N. C.; Henry Skinner, Philadelphia, Pa.; M. V. Slingerland, Ithaca, N. Y.; J. B. Smith, New Brunswick, N. J.; R. I. Smith, Atlanta, Ga.; E. B. Southwick, New York, N. Y.; H. E. Summers, Ames, Iowa; T. B. Symons, College Park, Md.; E. P. Taylor, Fort Collins, Colo.; E. S. G. Titus, Washington, D. C.; B. H. Walden, New Haven, Conn.; F. L. Washburn, St. Anthony Park, Minn.; F. M. Webster, Washington, D. C.; and G. P. Weldon, College Park, Md.

Among the visiting entomologists were the following:

H. G. Barber, Gustav Beyer, Henry Bird, E. A. Bischoff, H. H. Brehme, J. R. de la Torre Bueno, William T. Davis, Jacob Doll, G. P. Engelhardt, H. J. Erb, George Franck, C. F. Groth, C. W. Johnson, L. H. Joutel, Hugo

Kahl, W. D. Kearfott, G. J. Keller, H. H. Lyman, Ignaz Matausch, J. G. Needham, G. T. Rockwell, Charles Rummel, G. A. Runner, Charles Schaeffer, S. H. M. Seib, F. J. Smith, William Wasmuth, A. C. Weeks, W. M. Wheeler, and Rev. J. L. Zabriskie.

The meeting was called to order by President Kirkland, who called Mr. Hopkins to the chair while he delivered the annual address, entitled "A Great Experiment in Applied Entomology."^a

Immediately following the president's address the report of the treasurer was read, as follows:

Report of the treasurer.

Envelopes and paper-----	\$1. 90
Stamps -----	5. 00
Printing notices and programmes-----	9. 50
<hr/>	
Total -----	16. 40

Respectfully submitted.

A. F. BURGESS, *Treasurer.*

As no balance was left in the treasury after paying the expenses of the last annual meeting, action was called for on the amendment to the by-laws proposed by the committee on resolutions and printed in the proceedings of the eighteenth annual meeting. After a general discussion, the following amendment was adopted as a substitute for section 2, Article II, of the by-laws: "The annual dues of active members shall be \$1 and the dues of associate members 50 cents."

The report of the committee on nomenclature was presented by Mr. Herbert Osborn, chairman, as follows:

REPORT OF COMMITTEE ON NOMENCLATURE.

Your committee, in pursuance of the plan heretofore adopted for getting an expression of opinion as to the general use of common names, distributed, through the secretary, a list of 69 names, and in answer to these criticisms have been received from a number of members from widely different sections and out of the 69 in the list 41 have appeared acceptable to all who have replied, and these are submitted herewith for consideration or adoption at the present meeting. A number of others meet general acceptance, except in minor matters of form, and doubtless may be agreed to in the near future.

In the matter of use of scientific names the committee would recommend that the association adopt and recommend its members in practice to follow the rules of nomenclature adopted by the International Congress of Zoologists, and adhere to them strictly in all technical papers.

That in all economic papers these rules be followed in so far as practicable, and in accordance with the provision adopted in a previous meeting that in any doubtful cases as to generic reference the genus that has been in common use, or that may be adopted in the latest general catalogue of authority, shall be the preferred name.

^a Owing to the pressure of important work Mr. Kirkland has been unable to complete and supply the manuscript of his address.—Ed.

Apple curculio	<i>Anthonomus quadrigibbus</i> Say.
Apple twig-borer	<i>Schistoceros hamatus</i> Fab. ^a
Black cutworm	<i>Agrotis ypsilon</i> Rott.
Book-louse	<i>Troctes divinatoria</i> Müll.
Buck moth	<i>Hemileuca maia</i> Dru.
Cadelle	<i>Tenebroides mauritanicus</i> L.
Cabbage looper	<i>Autographa brassicae</i> Riley.
Chaff scale	<i>Parlatoria pergandei</i> Comst.
Cigarette beetle	<i>Lasioderma serricorne</i> Fab.
Catalpa sphinx	<i>Ceratomia catalpæ</i> Boisd.
Cherry scale	<i>Aspidiotus forbesi</i> Johns.
Clover cutworm	<i>Jaenestra trifolii</i> Rott.
Clover mite	<i>Bryobia pratensis</i> Garm.
Currant borer	<i>Ægeria tipuliformis</i> Clerck.
Fall armyworm	<i>Laphygma frugiperda</i> S. & A.
Forest tent-caterpillar	<i>Malacosoma disstria</i> Hbn.
Garden webworm	<i>Lorostege similalis</i> Guen.
Gray blister-beetle	<i>Epicauta cinerea</i> Forst.
House cricket	<i>Gryllus domesticus</i> L.
Imbricated snoutbeetle	<i>Epicarus imbricatus</i> Say.
Lappet moth	<i>Epicaptera americana</i> Harr.
Leaf crumpler	<i>Mineola indiginella</i> Zell.
Melon caterpillar	<i>Diaphania hyalinata</i> L.
Northern mole cricket	<i>Gryllotalpa borealis</i> Burm.
New York weevil	<i>Ithycerus noveboracensis</i> Forst.
Onion maggot	<i>Phorbia cepetorum</i> Meade. ^b
Orange scale	<i>Aonidiella aurantii</i> Mask.
Pickle worm	<i>Diaphania nitidalis</i> Cram.
Raspberry sawfly	<i>Monophadnoides rubi</i> Harr.
Rose sawfly	<i>Endelomyia c. rosa</i> Harr.
Salt-marsh caterpillar	<i>Estigmene acrea</i> Dru.
Sheep tick	<i>Melophagus ovinus</i> L.
Stable fly	<i>Stomoxys calcitrans</i> L.
Strawberry weevil	<i>Anthonomus signatus</i> Say.
Turkey gnat	<i>Simulium meridionale</i> Riley.
Variegated cutworm	<i>Peridroma saucia</i> Hbn.
Walking-stick	<i>Diapheromera femorata</i> Say.
White-lined sphinx	<i>Deilephila lineata</i> Fab.
Vagabond crambus	<i>Crambus vulgivagellus</i> Clem.
Yellow mealworm	<i>Tenebrio molitor</i> L.

HERBERT OSBORN, *Chairman*.

F. M. WEBSTER.

C. P. GILLETTE.

^a Synonym, *Amphicercus bicaudatus* Say. See Lesne, P.—Revision des Bostrychides. Ann. Soc. Ent. France, vol. lxxvii, pp. 513, 514, 1898,—Ed.

^b This species is placed by Coquillett in the genus *Pegomya* Desvoidy. See Chittenden, Cir. 63, 2d ed., Bur. Ent., U. S. Dept. Agric., p. 6, footnote 2, 1906.—Ed.

^c For reference of this species to the genus *Endelomyia* see Ashmead, Can. Ent., vol. xxx, No. 10, October, 1898, p. 256.—Ed.

Considerable discussion arose concerning the proper use of the hyphen in compound words, and Mr. Slingerland suggested that in the future some standard dictionary be followed as an authority. It was pointed out by Mr. Osborn that the main consideration was to secure the uniform use of common names when applied to insects of economic importance, and that the hyphenization should be considered a minor matter if the other object could be accomplished. Doctor Howard indorsed the part of the report which urged that the rules of nomenclature adopted by the International Congress of Zoologists be followed by the members of the Association, and after a brief discussion the report was adopted.

The report of the committee on cooperative testing of insecticides was read by Mr. H. T. Fernald, chairman.

REPORT OF COMMITTEE ON COOPERATIVE TESTING OF INSECTICIDES.

Your committee on the cooperative testing of insecticides begs leave to report as follows:

It seemed wise to the committee to learn how generally a desire existed for cooperation in this work. Accordingly a circular letter was sent to over 50 of the leading workers in economic entomology in all parts of the country, and answers were received from about half of this number. If these answers fairly represent the general feeling of the entomologists of the country, it seems that the idea of cooperation is almost universally indorsed, though with qualifications in many cases. The other questions asked in the circular were to bring out ideas as to the best methods to accomplish cooperation, and these brought widely divergent answers, hardly any two being in agreement on all points. From a study of all the correspondence, however, your committee recommends the following:

I. That all proprietary insecticides the use of which is restricted to particular localities should be tested, if at all, by the proper authorities in those localities. If they seem to promise wide application as a result, they should then be tested as below.

II. That all proprietary insecticides used or advertised in many parts of the country should be tested as follows:

(a) A standing Committee on Insecticides, to consist of five persons elected by this association, should obtain—from the manufacturer if possible—a statement of the composition of the insecticide in each case. If this proves that it could not possibly be of any value or do the work claimed, it should be dropped. If from a statement received from the manufacturers it seems possible that the material might be of some value, a sample should be sent to some chemist for qualitative and quantitative analysis.

(b) That, if possible, arrangements be made by which the chemist in charge of the insecticide work of the United States Department of Agriculture shall conduct this portion of the work, the material to be sent him by the standing committee.

(c) That if the report of the chemist shows that the material is worthless as an insecticide no further consideration be given to it.

(d) That if it appears probable that the material is of some value as an insecticide, samples be sent by the standing committee to several entomologists in different sections of the country, who may signify their willingness to do this work, for the purpose of making cooperative tests of it.

(e) That a plan of testing be worked out by the standing committee and sent by it to each entomologist, with the substance, together with a plan for reporting results, so that the factors entering into the test and the data in the reports may be uniform and full.

(f) That reports received be compiled and a general report made by the standing committee to this association at its annual meeting.

(g) That these reports, or such portions of them as seem advisable, be published by the association and then by others who may wish to publish them.

III. Your committee further recommends that if the above plan or any modification of it be adopted, the committee designated by this association to carry it out be requested to prepare the plans for testing and reporting on the tests and for assigning the materials to the entomologists selected by them to make the tests.

Respectfully submitted.

H. T. FERNALD. *Chairman.*

A. F. BURGESS.

H. A. SURFACE.

On motion this report was accepted and ordered published in the proceedings of the meeting. The appointment of the standing committee suggested was deferred for later consideration. A brief report of the programme committee was made by the secretary and related chiefly to the arrangement of the papers to be presented. A motion was made and carried that the time for the presentation of each paper be limited to ten minutes.

A paper was then read by Mr. Washburn, entitled:

INSECT NOTES FROM MINNESOTA FOR 1906.

By F. L. WASHBURN, *St. Anthony Park, Minn.*

THE CABBAGE MAGGOT.

Flies of the cabbage maggot (*Phorbia brassicae* Bouché) were first observed May 9. On May 16 egg laying was well under way. On May 31 first maggots were observed. On June 5 maggots were exceedingly abundant. By June 12 puparia were found in large numbers, and all the cauliflower of many market gardeners was reported destroyed. By July 7 maggots were transformed to pupæ. September 26 was the latest date at which flies emerged from puparia in the laboratory.

On July 2 several cabbage maggot flies which had been confined under a bell jar with a potted cauliflower plant laid fertilized eggs, which hatched on July 6, a few hours over three days from the time of laying. The eggs were laid on the soil near the plant. The same observation was repeated with other specimens, the flies in the latter

instances laying within four hours of the time they were confined and frequently placing their eggs in the axils of the leaves.

The hatching of two eggs was observed. In each case the larva appeared to emerge from the egg through the groove on the side. One egg was observed to hatch under the microscope, the active larva trying unsuccessfully for several hours to free itself from the shell, and finally dying. The smooth surface of the glass may have been, and probably was the cause of its failure to extricate itself from the egg. Two other eggs, which were placed under observation immediately after being laid, hatched in five days from the time of laying. Another egg hatched in three days and five hours. Evidently the egg stage lasts from three to five days. The two eggs referred to as hatching in five days had been considerably shaken in a vial in being transported from the field to the laboratory; hence disturbing the egg evidently does not necessarily prevent hatching. The larvæ, judging from observations made this season, evidently require twenty to twenty-one days, under ordinary conditions, to reach maturity.

A lot of maggots which changed to pupæ June 6 emerged as flies June 19 and 20. Evidently, then, the pupal stage in Minnesota may have from thirteen to fifteen days' duration.

We are losing faith in the use of carbolic emulsion as employed against this pest, and believe that cultural methods may possibly prove our best means of control. Young maggots lived in the laboratory for two hours and twenty minutes immersed in carbolic emulsion, 1 part to 30 parts of water, and adult maggots required three hours and forty minutes' immersion before dying. Some eggs hatched after thirty seconds' treatment in carbolic emulsion at the same strength. On the other hand, good results were obtained by immersing the roots in hellebore and water at time of setting, using 1 part of hellebore to 2 parts of hot water, allowing it to cool before being used. The plants were immersed deeply enough to coat the lower part of the stems and were immediately planted and made an excellent showing. Good results were also obtained, in a protective way, by the use of bran and glue and sawdust and glue, the sawdust being mixed with glue in the proportion of $\frac{1}{2}$ pound of the former to 1 quart of the latter. The glue was not at all thick, but represented 2 pounds of hard glue in 1 gallon of water. When bran was used the proportions were the same. The mixture was rather more sloppy than good chicken feed. It was applied by hand about the base of the plant, put well up on the stem, the diameter of the mass where it came in contact with the ground being about 4 inches. One quart was sufficient for 15 plants. The stuff quickly hardened and, though it softened somewhat during summer rains, did not disintegrate. Plants

so treated made an excellent showing. A man can treat 6 to 8 plants per minute. This treatment, of course, would not be practicable on a large acreage.

The small red mite *Trombidium scabrum* Say was observed to be extremely active in sucking the eggs of the Phorbia. On May 15 an assistant in the field reported this mite as very numerous, averaging about two to a plant, and occasionally four or five were observed about one plant. On this date a large number of eggs examined had evidently been sucked. Frequently there would not be a single good egg found around a plant, out of a lot of a dozen or more that had originally been laid there.

We have obtained from the burrow of a maggot a cynipid parasite, *Pseudeucala gillettei* Ashm.: we have also bred from a puparium *Plectiscus* sp.—identified by the American Entomological Society, of Philadelphia. We have also included among the predaceous enemies of Phorbia the carabid beetles *Pterostichus coracinus* Newm., *P. lucublandus* Say, *Agonoderus pallipes* Fab., and *Amara impuncticollis* Say, since immediately after being brought in from the field these beetles fed ravenously upon the maggot. These species were present in large numbers in almost all of our cabbage fields, as were also *Heterothops fumigatus* Lec., *Lathrobium anale* Lec., and *Bembidium quadrimaculatum* L., although these three latter beetles were not actually observed to eat the maggot. Plants in sandy soil appeared to suffer more, everything else being equal, than those planted in heavy soil.

RECENT OBSERVATIONS ON THE USE OF HYDROCYANIC ACID GAS.

One sometimes has occasion in fumigating mills of small size to drop packages of cyanide into jars containing acid, and at such times the question as to how much time elapses between the dropping of the cyanide, inclosed in a double sack, and the giving off of the deadly gas is an important one. In doing the work personally I have allowed fifteen seconds as a conservative estimate in this direction and acted accordingly. To place this matter beyond any doubt, however, we have this fall made several trials, timing the interval between the dropping of the double bag of cyanide into the jar and the first appearance of the fumes, with surprising results. A double manila sack was used in each case—that is, one sack inside another—and various makes of sacks. In each case the liquid was fairly warm, but no observation was made on its exact temperature. We found in a series of trials that this interval varies from twenty-nine seconds, the lowest period, to four minutes, the latter being the highest interval, the variation evidently being largely due to variations in the thickness and character of the paper of which the sacks

are made. Of course a variation in the warmth of the liquid would cause a difference. Further experiments, bringing in the temperature factor, are planned for next summer. The experiments cited herewith were performed late in the fall, and the steam caused by the union of the acid and water did not cease until the liquid was fairly cool.

In connection with these experiments the question arose, "Is it not possible that fumes which would be fatal, if breathed, rise from the jar before they are visible?" This was answered in the negative by placing a live guinea pig in a wire cage about 6 inches above the top of the jar immediately after the charge of cyanide had been lowered into the acid. Visible fumes began to rise two minutes after placing the charge, and a few seconds later the guinea pig succumbed, showing that the dangerous gas was not given off in a fatal amount until fumes were observed. It would seem, then, from these results, that, on a very conservative estimate, one can depend upon at least twenty seconds when double sacks are used, and much can be done in that time.

Another important question connected with the use of hydrocyanic acid gas is the distance it will penetrate into a semisolid mass—a sack of bran, for instance—or masses of cereals held together by the webbing made by the flour-moth larvæ. An apparatus used by this Department to ascertain this was found to be faulty too late in the year to remedy the defect, and the results of that particular experiment, therefore, can not be relied on. However, the matter was given a practical test by exposing a mass of webbing over 2 inches thick, containing live worms, pupæ (and probably eggs) of the flour moth to the fumes of the gas as commonly used at the above strength, for over ten hours, the time commonly occupied in a fumigation. As far as could be observed at the time, everything in this mass was killed, and after three months' observation of the same in our laboratory no sign of the flour moth in any stage was apparent, indicating, possibly, that this gas has a greater penetration than we had supposed. At the same time, this isolated experiment should not be relied on as certifying absolutely upon this point.

WORK AGAINST THE LITTLE RED ANT IN A RESIDENCE.

The little red ant (*Monomorium pharaonis* L.) has been a great pest in a large residence in Minneapolis for several years past. During the fall of 1905 these ants became so numerous that we were requested to try to eradicate them. At that time very few parts of the house seemed to be entirely free from the pests. Flowers, food, or candy left in almost any part of the house would be covered in a few hours. The ants were particularly annoying when they were

found in the pantries and kitchen, getting on food just before it was taken into the dining room.

Clusters of cocoons were located in the walls of the furnace room in the basement, this room being directly beneath the kitchen and pantries. At first carbon bisulphid was used with good results, the liquid being forced into crevices in the walls where the ants were observed, but the gas penetrated to all parts of the house and was particularly disagreeable to the members of the household. As far as could be observed these "nests" were never placed very deeply within the wall, and consequently when we had to resort to kerosene we found this liquid to be just as effective as was the carbon bisulphid. This was applied with a syringe. The ants that were out of the "nests" at the time of applying the liquid would, upon their return, seek another crevice, and would soon found a new colony; then these new colonies would be located and saturated with kerosene. Finally, in January or February, 1906, no ants were seen about the house; they had completely disappeared, and we were confident that they had all been destroyed. In the fall of 1906, however, they became very numerous again, and our attacks against them with kerosene were renewed. Several nests were found in practically the same places as in the previous year, but the most persistent colony was found, seemingly, in the chimney. The nest itself, however, was never seen. The extermination of the ants in this location was extremely difficult, but at the time of writing their numbers appear much lessened.

Many so-called "ant cures" were tried, such as "ant sugar," pennyroyal, and tartaric acid solution made according to the formula given on page 97 of Bulletin 30 of the Division of Entomology, U. S. Department of Agriculture, viz. tartaric acid, 10 grams; sugar, 100 grams; water, 1,000 grams. None of these remedies was effective.

We had planned during the winter, when the house was to be unoccupied, to fumigate with hydrocyanic acid gas, but on or about December 22 every ant had disappeared, and we are now awaiting their reappearance before taking active measures against them.

It was decided to make the discussion on this paper and the president's address the first order of business for the afternoon session.

On motion the following committees were appointed by the president:

Nominations: Messrs. Marlatt, Osborn, and Webster.

Resolutions: Messrs. Fletcher, Slingerland, and Sanderson.

Membership: Messrs. W. D. Hunter, Summers, and J. B. Smith.

AFTERNOON SESSION, FRIDAY, DECEMBER 28, 1906.

The meeting was opened with a discussion of the presidential address and of the paper presented by Mr. Washburn. In reply to an inquiry concerning the effect of fumigation on insects attacking grain, Mr. Kirkland stated that he had used hydrocyanic acid gas in fumigating a mill, and had secured excellent results. One hundred and twenty-five pounds of cyanide were used, and as the building was not very tight this was applied at the rate of 1 ounce of cyanide to 80 cubic feet of space, and the building was kept closed for six hours.

A discussion then followed on the use of arsenate of lead. Mr. Frost stated that shade trees of average size could be sprayed with this poison for from 40 to 60 cents each, and that in woodland areas the price could be reduced to from 25 to 40 cents per tree. Such treatment is applied in regions infested with the gipsy moth, and thousands of trees are being sprayed with this poison. In reply to a question by Mr. Slingerland he stated that arsenate of lead does not burn the foliage to any appreciable extent if it is properly prepared. Men in his employ used 14 tons of this poison during the past summer, and in such large lots it was secured at from 11 to 12 cents per pound. He did not consider it advisable to attempt its manufacture when good material could be bought at this price.

Mr. Britton remarked that his experience in Connecticut had led him to believe that peach and Japanese plum trees could not be sprayed while in foliage without causing injury.

Mr. Kirkland stated that in manufacturing arsenate of lead it was necessary to form the precipitate in large vats, and if made in this way there would be very slight chance of foliage injury. In some cases, however, delicate leaves might be burned.

Mr. Marlatt inquired if the yellow acetate of lead was as good as the white acetate for making arsenate of lead, and Mr. Kirkland replied that the arsenate of soda which is imported carries about 70 per cent As_2O_5 and could be bought at New York, duty paid, for about 8 cents per pound. This year white arsenic is selling for 8 to 9 cents a pound. The 50 per cent grade of arsenate of soda is made by fusing white arsenic and soda together. Acetate of lead is sold in different grades—known as crystallized, white granulated, brown, and dark brown. The best manufacturers of arsenate of lead use white granulated acetate of lead.

Mr. Burgess stated that he had observed considerable injury from using homemade arsenate of lead, and in several cases this proved to be due to the fact that the arsenate of soda had been adulterated with salt. In one case an analysis showed that the sample contained over 25 per cent of salt.

President Kirkland announced that Mr. Orlando Harrison, of Berlin, Md., president of the American Association of Nurserymen, Prof. John Craig, of Ithaca, N. Y., and Mr. Emory Albertson, of Bridgeport, Ind., representing the American Association of Nurserymen, were present on invitation of the Association of Economic Entomologists, as the former association was vitally interested in the report about to be read.

On motion of Mr. Sanderson these gentlemen were admitted to the privileges of the floor.

The report of the committee on national control of introduced insect pests was read by Mr. E. D. Sanderson, chairman.

REPORT OF COMMITTEE ON NATIONAL CONTROL OF INTRODUCED INSECT PESTS.

To the Association of Economic Entomologists:

Your committee, appointed at the last meeting of the association to consider the national control of introduced insect pests, beg to report as follows:

After preliminary correspondence the committee seemed to be generally agreed except upon the matter of uniformity of nursery inspection. A subcommittee, consisting of Messrs. Burgess, Forbes, and Gillette, was therefore appointed to consider this matter. This subcommittee met at Urbana, Ill., and, having had a large correspondence with inspection officials throughout the country, formulated a report which was presented to the full committee at a meeting held at Baton Rouge, La., November 14. This report is embodied in section C. below.

As instructed by the association at the meeting at Baton Rouge, your committee conferred with a similar committee of the Association of Horticultural Inspectors represented by Messrs. R. I. Smith, of Georgia, and S. A. Forbes, of Illinois, and with representatives of the National Nurserymen's Association, Messrs. Watrous, of Iowa, and Albertson, of Indiana. All present agreed upon the line of procedure outlined below. At the meeting of the Association of Horticultural Inspectors at Baton Rouge resolutions were adopted similar to those below, advocating that the Secretary of Agriculture be empowered to inspect all the imports for insects and plant diseases and that he be empowered to make regulations governing the certification and inspection of nursery stock for interstate commerce, and appointed their chairman, Dr. S. A. Forbes, to act as a member of a committee to be composed also of a representative of the Association of Economic Entomologists and a representative of the National Nurserymen's Association to push this legislation.

The committee would therefore suggest the following resolutions and procedure toward securing such legislation:

A. *Resolved*, That the Secretary of Agriculture should be empowered to inspect all imports and to make regulations governing importations liable to harbor insect pests and plant diseases, and that sufficient appropriation be made for this purpose.

B. That Congress should authorize the Secretary of Agriculture to proceed to exterminate or control imported insects or plant diseases or any insect previously native to a restricted locality, but which may become migratory and threaten the whole country, whenever, in his judgment, such action is practicable, and that an appropriation be made for this purpose as a reserve fund for emergency use against any such pest which may arise. Such legislation would

give the Secretary of Agriculture similar authority against plant enemies as now exists for procedure against animal diseases by the Bureau of Animal Industry.

C. (1) That the Congress of the United States be asked to enact a law empowering the Secretary of Agriculture to make such regulations as may be deemed necessary in order to secure uniform methods of nursery inspection and certification of all nursery stock which passes into interstate commerce.

(2) That all State or Territorial officials in charge of nursery inspection be urged to accept these certificates at their face value and that in States where laws are now in force which will not allow the acceptance of such certificates, the inspection departments be requested to endeavor to secure such State legislation as will make this possible.

(3) That each State should make and enforce such regulations as its local authorities may deem necessary, but that they be made as similar to those of the United States Department of Agriculture as practicable.

D. Your committee suggests that the Association elect a representative to form a committee with a representative of the Association of Horticultural Inspectors and a representative of the American Nurserymen's Association to push this legislation before Congress, as in the judgment of your committee this is the best method available for securing its passage. We would also suggest that topics A and C, above, be combined in one proposed law; and that topic B, involving the control of introduced species, be embodied in another law; and that, if the passage of both measures be impracticable, efforts be concentrated upon the law involving the inspection of importations and the control of nursery stock for interstate trade, and that the other measure be pushed later.

Respectfully submitted.

E. D. SANDERSON, *Chairman*.

C. P. GILLETTE.

H. A. MORGAN.

A. F. BURGESS.

S. A. FORBES.

Mr. Harrison stated that the nursery interests were being severely injured on account of the diverse and sometimes unreasonable requirements made for shipping stock into different States. He declared that the better class of nurserymen welcomed thorough inspection, and that under no circumstances would they be willing to have this work discontinued, as they considered it a benefit to themselves and the trade. Any movement which would bring about more uniform regulations and requirements, so that as little confusion as possible would result to the nurseryman, was very desirable, and he heartily favored the report.

Professor Craig stated that he believed that the principle outlined in the report was correct and that if workable legislation could be secured it would greatly benefit the nursery interests. He urged the necessity of such action as would prevent unjust discrimination, and which would help the nurseryman who was striving to do an honorable business to secure the delivery of his stock without expensive and objectionable delay. He therefore heartily indorsed the report.

Mr. Albertson remarked that he agreed with the statements made by the previous speakers and believed that the report submitted by

the committee was a step in the right direction. He also pointed out the fact that the entomologists and nurserymen were coming to a better understanding of the situation, and thanked the association for the courtesies extended to himself and the other representatives of the Nurserymen's Association.

Mr. Sanderson stated that the problem of bringing about uniform nursery inspection requirements was a large and difficult one and that it probably could not be solved in a single year. He felt, however, that the report of the committee indicated the most practical line of work to be taken up in this direction, and if the necessary legislation could be secured it would result in bringing about a condition that would be more satisfactory to nurserymen, horticulturists, and inspectors. The Association of Horticultural Inspectors had adopted a similar report at their annual meeting at Baton Rouge last month and had appointed a representative to act on the joint committee suggested in the report.

Mr. Marlatt said that Congressional action would be taken when the nurserymen of the country as a whole joined in a strong demand for it, and that a demand from this source would have great weight with Congress, especially with the indorsement of the official entomologists of the different States and the State horticultural inspectors. The remedy, therefore, lay largely in the hands of the nurserymen of the country, and without their united support relief could not be hoped for from Congress.

Mr. Burgess pointed out that for the first time in many years the entomologists interested in nursery inspection had, through a committee, proposed a definite scheme for handling the problem. Harmonious relations now exist with the nurserymen, and they and the inspectors appear to have come to an agreement as to the best measure to adopt. He expressed the hope that the entomologists would accept and adopt the report.

Mr. J. B. Smith stated that, although Congress might pass a National law, it would not be able to overrule the requirements or regulations of State officials, and that, this being the case, he could not see how the law would be enforced so that any great benefit would result.

Mr. Webster pointed out that if a National law had been passed years before when the matter was agitated, there would not now have been very many conflicting State laws to interfere with the work. In spite of this, he believed the adoption of the report would be a step in the right direction, and that it was not yet too late to take up and push forward the work that should have been done years ago.

Mr. Summers called attention to the fact that one of the reasons for stringent requirements in some States was the careless inspection of other State officials, and that if many of the State inspectors were

satisfied that a high standard of inspection requirements was maintained, such as would be the case if it were under Government control, many State requirements would, undoubtedly, be modified in such a way as to simplify interstate shipments.

After further general remarks a vote was called for, and the report was unanimously adopted. On motion, the election of the representative of the association to serve on the joint committee was referred to the committee on nominations.

The following paper was presented:

A NEW ORIENTAL INSECT PEST (?) IN MASSACHUSETTS.

By H. T. FERNALD, *Amherst, Mass.*

[Withdrawn for publication elsewhere.]

A paper was read, as follows:

OCCURRENCE OF THE GIPSY MOTH IN CONNECTICUT.

By W. E. BRITTON, *New Haven, Conn.*

During the season of 1906 my attention has been partly occupied in attempting to control, and to exterminate if possible, a small gipsy moth (*Porthetria dispar* L.) colony in Stonington, Conn. Stonington is the southeast corner town of the State, joining Rhode Island, and having an area of nearly 75 square miles. The infested portion covers, so far as we know, only about 1 square mile just north of the village, in the south part of the town.

For several years we have expected that the gipsy moth would appear in Connecticut, and have been on the watch for it. It has been reported several times from different places, but in each case upon investigation some other species was found to be the cause of alarm. The first real gipsy moth was taken at Stonington, July 30, 1905, by Mr. Ernst Frensch, a local collector, who recognized the insect because he had seen it in Germany. He noticed two males flying about in an apple tree, and, on looking closer, saw a female resting on the bark of the trunk, and put her in his cyanide jar. He put the specimen away in his collection, and forgot all about it until during the winter, when I wrote to the entomologists of the State asking for cooperation in furnishing records of their rare captures for use in our lists of Connecticut insects. Mr. Frensch sent me a number of records, including that of *Porthetria dispar*. I made an appointment with him, and visited the place March 6. He showed me where he had captured the female, and called my attention to an egg mass on a Norway maple tree near by. He also showed me an egg mass which he had found on a small bush. Suspecting that

these might be gipsy-moth eggs, he had, in order to make sure, cut off the abdomen of the female moth and, obtaining the eggs from her, compared them with those of the egg mass and concluded that they were identical. We inspected the region and found a number of egg masses in some low bushes near the velvet mill. This, as well as the place where Mr. Frensch found the adult moth, is near the railroad, and not far from the point where the spur track leading to the steamboat dock branches from the main track.

The next thing was to learn the extent of the infestation, or, in other words, the size of the infested area. We applied to Massachusetts for a trained scout, and through the kindness of Superintendent Kirkland we obtained the services of Mr. C. S. Mixter for two weeks. Mr. Mixter scouted nearly 5 square miles of territory, giving as his opinion that the infested territory had been well surrounded. The brush on about 5 acres of land around the pond by the velvet mill was cut and burned before May 1, the hatching time for the eggs. The egg masses found were destroyed by soaking them with creosote oil. Banding trees with burlap was commenced about the middle of May; only a few were banded at first, but the number was increased as fast as seemed desirable. We supposed, of course, that about all of the egg masses had been destroyed, but the number of caterpillars found showed that some had been overlooked, and during the summer we ran across a number of these old egg masses—more even than had previously been destroyed.

Most of the caterpillars were found on old apple trees, though cherry, quince, rose bushes, and red maple were infested. Many of the old apple trees had received no care for many years, if ever, and the tops were crowded with branches, some of which were dead, the bark was rough, and in many cases trees were hollow or had cavities caused by broken or rotting branches. All of these faults were serious hindrances to our work because they furnished hiding places out of our reach for the caterpillars, so that they would not go under the bands where we could find them. We therefore pruned and scraped many trees, and filled up the cavities with stone and cement. A few trees were sprayed with arsenate of lead, and sticky bands of "tree tanglefoot" were given a trial. In a few cases brush growing near stone walls was found infested. The brush was cut and the walls fired with fuel oil to kill any caterpillars that might be hiding in them.

We had men on the ground continuously from June 7 to September 1, and during the latter part of June and first half of July ten men were kept busy pruning, scraping, and banding trees, and destroying caterpillars. The bands were visited each day until after nearly all the caterpillars had pupated, when they were examined less frequently—perhaps every other day—and finally twice a week, until all

had transformed. All trees infested with caterpillars or egg masses were marked, as is done in Massachusetts. A breeding cage was made on the grounds, and in it were reared a number of adults for exhibition and illustrative purposes. During August a gang of laborers was employed to cut and burn brush, and the hedgerows through some of the fields were cleaned up. We interviewed the selectmen, interested them in our work, and they cooperated by cutting all the brush along the highways through and for some distance beyond the infested district. This brush was burned by our men. In four places caterpillars transformed in the stone walls, and egg masses were laid there. The walls were overhauled, eggs destroyed, and the walls relaid. All work was suspended September 1 to enable us to make the annual inspection of nurseries. Since November 16 five men have been employed cutting and burning brush and scouting for egg masses.

It was necessary to do considerable educational work, and immediately Bulletin 153 was issued from the Connecticut Agricultural Experiment Station giving brief accounts of the gipsy and brown-tail moths. Two months later the annual report, containing a further account of the gipsy moth, was distributed. Figures and brief descriptions of the insect and its injuries were printed on cards 11 by 14 inches in size, and nearly 2,000 of these have been distributed to schools. A number of life history sets in Riker mounts have been placed in drug-store windows in Stonington, Mystic, Noank, Groton, and New London. An illustrated lecture was given at New Haven May 9 and at Stonington November 26; specimens have been shown and talks given about the insect in about a dozen meetings in various other parts of the State.

Scouting for egg masses was done in April at Mystic, Midway, New London, Plainfield, Danielson, Putnam, and Willimantic. Nearly all portions of the State are visited during the year by some member of the office force on the lookout for such things, but no gipsy moths have been found anywhere outside of Stonington.

Up to the present time the results obtained may be expressed by the statistics in the following table:

Egg masses laid in 1905:

Number destroyed	29
Number hatched	36
Egg masses laid in 1906, number destroyed	47
Caterpillars destroyed	10,000
Pupæ destroyed	47
Number of trees banded, more than	1,300
Amount of money expended	\$1,700

A word as to funds: A few hundred dollars only could be spared at that time from our State appropriation for insect work, and the

Connecticut board of agriculture kindly appropriated \$2,000, to be used if needed, and Governor Roberts and his associates assured us that if after using this money at our disposal still more was needed to hold the pest in check it would be forthcoming. We called upon the board of agriculture for \$800, and the remaining \$900 has come from our own insect-pest appropriation. An attempt will be made to have the State legislature, which soon convenes, set aside a few thousand dollars to be used if needed in work against the gipsy and brown-tail moths. The brown-tail moth has not yet been found in Connecticut, though it must be very near its borders in Massachusetts. We shall endeavor to exterminate the gipsy-moth colony at Stonington, and this can be done if it has not spread beyond the area where we have found it. The village of Stonington is on a narrow point of land extending into the ocean. The infested territory extends from the village northward and slightly eastward; it is flanked on both sides by water—on the east by the Wequetequock River and on the west by an arm of the sea extending northward from Stonington Harbor. A line from the northernmost extremity of this salt water extending easterly to Wequetequock River cuts the mainland some distance north of where any caterpillars or egg masses have been found, although considerable scouting has been done in this section and many of the trees were banded in caterpillar time.

Two natural enemies of the gipsy moth have been observed in Connecticut. The "caterpillar hunter" or "searcher" (*Calosoma scrutator* Fab.) was quite common under the bands, and one of these in captivity devoured gipsy moth caterpillars with avidity. Out of the ten thousand or more caterpillars gathered and destroyed four diseased ones were observed. These shriveled and finally died, as if attacked by some bacterial disease. While in Massachusetts the last week in June I observed the same or a similar disease which killed many caterpillars, though of course only a small proportion. Dr. G. E. Stone, botanist of the Massachusetts experiment station at Amherst, was investigating the matter, and I sent him two of the diseased caterpillars from Stonington. At that time he was not ready to report on the nature of the disease, but stated that a number of different organisms had been isolated from the diseased caterpillars.

Just how the pest reached Stonington may perhaps never be known, but there is much speculation regarding it. Eggs or pupæ may have been brought on packing boxes to the velvet mill or upon freight cars left upon the spur track. Certainly the worst infestation was near the velvet mill and the railroad, and I feel that it must have reached Stonington on steam cars via the New York, New Haven and Hartford Railroad. Some think that it may have been a direct importation from Europe, as Germans live in the locality, work in the mill, and occasionally travel back and forth.

At the present time it is difficult to find egg masses or, except for the marked trees and the cutting of brush, any other indications that the area is infested by the gipsy moth. But the work must be kept up even after it is believed that the last one has been destroyed.

In reply to a question Mr. Britton stated that the chances for exterminating this insect in Connecticut appeared to be good, that the people were interested in the matter, and an urgent appeal would be made to the State legislature for funds to use in suppressing the moth.

A paper was read, entitled:

NOTES ON FUMIGATION AND DIPPING OF NURSERY TREES.

By T. B. SYMONS and A. B. GAHAN, *College Park, Md.*

[Withdrawn for publication elsewhere.]

Considerable interest was drawn to this paper, owing to the fact that the State of Oregon has recently required that nursery trees be dipped in the lime and sulphur wash rather than fumigated before they are planted.

Mr. Slingerland called attention to the fact that a temperature of 180° F. used in one series of the dipping experiments, was unnecessarily high, and this probably accounted for the serious injury to the trees.

Mr. J. L. Phillips stated that he had dipped one-year apple trees in cold lime-sulphur wash in the spring of 1906, as follows: 235 Black Twig, 205 York Imperial, and 20 Albemarle Pippin. The trees were dug, the tops cut off to within 3 feet of the crowns, and the tops dipped to the crowns in the wash, made by using 15 pounds of lime, 15 pounds of sulphur, and 5 pounds of salt to 50 gallons of water. The trees were immediately set in the orchard and all lived and grew well except one. The results with peach trees were not so good. They were cut back to within 2 feet of the crown and dipped as follows: 151 Smock, of which 124 lived; 142 Elberta, of which 76 lived; and 80 Salway, of which 66 lived. Many of the peach trees that lived died back a few inches from the top. As this was a commercial plantation, no checks were planted. There was considerable injury to both peach and apple trees in the experimental dipping tests, but the conditions under which the trees were handled may be partly at fault, as a number of untreated trees planted at the same time died also.

In reply to several questions, Mr. Symons stated that the lowest temperature used in the dipping experiments was such that a person could bear his hand in the solution. He could not say definitely whether all the trees were dug in the spring, but believed this was the case. No observations could be made as to whether this treatment killed the scale, as nearly all the treated trees died during the summer.

In the absence of Mr. Schoene, the following paper was read by Mr. Parrott:

THE WILLOW BORER AS A NURSERY PEST.

(*Cryptorhynchus lapathi* L.)

By W. J. SCHOENE, *Geneva, N. Y.*

The willows and poplars along the streams and canals, ornamental willows in the cities, and poplars and willows in the nursery are being seriously injured by this beetle. The first noticeable outbreak of the insect in this State occurred in a nursery at Rochester in 1902, and this species is now a serious pest in various parts of the State. In many poplar and willow plantations the beetle has been estimated to destroy 10 per cent of the stock and in some instances the entire plantation has been ruined. The species of willow and poplar that have been observed to sustain conspicuous injuries are: *Populus monilifera*, *Salix lucida*, *S. caprea*, *S. cordata*, *S. sericea*, *S. alba*, and *S. amygdaloides*.

This insect has been discussed in a comprehensive manner by Prof. F. M. Webster in a paper entitled "The Imported Willow and Poplar Curculio," which was presented before the Columbus Horticultural Society. This treatise also contains some observations made by Mr. A. H. Kirkland on the life history of the beetle and its destructiveness in Massachusetts.

The increasing importance of this species to the nursery interests of New York prompted an investigation to determine its life history in this State for the purpose of ascertaining a practical method for the control of the pest in poplar plantations. The aim of this paper is to call attention to the results that have been attained.

To understand clearly the trend of the work it is well at this time to review in brief the life history of this insect. Oviposition commences about August 1 and lasts through September. The eggs hatch in eighteen to twenty-one days, and the larva upon hatching begins to bore in the cambium layer, where it finds subsistence. As it approaches maturity it makes a channel in the heartwood. The larvæ commence to pupate about July 1 and the beetles begin to emerge two weeks later. For the next ten weeks the adults can be found in abundance. Before beginning to deposit eggs the beetles feed for a week or ten days on the bark of one-year shoots, after which they are more often found upon the older parts of the tree, especially in the injured portions of the bark and corky overgrowths caused by pruning. It is because of the large number of punctures on the young wood that the work of the adult is especially noticeable. This habit at once suggested the possibility of using arsenical sprays

as a means of control, and to test the value of these poisons some experiments were made, as follows:

A young poplar in a nursery row was headed back and sprayed with poison; a bag of mosquito netting was then put over the tree to inclose some beetles. As a check to each tree treated in this way, beetles were confined in a similar manner as mentioned above upon trees that had not been sprayed. With one exception this work was done upon nursery poplars that were 2 to 4 years old.

It is not necessary to give the data in detail, but it is sufficient to say that the results of a number of experiments made during the fall of 1905 indicated strongly the possibility of using arsenical sprays as a remedy. When beetles were confined upon trees sprayed with poison they died in a few days, while beetles confined in a similar manner upon unsprayed trees continued to feed and to oviposit.

In order to corroborate these results the work was continued during 1906. In addition, an effort was made to determine the length of time that the spray was effective and whether or not the poison acted as a repellent. The experiments were conducted in the same manner as in the previous year and the results were even more conclusive. The poison was found to be effective for thirty days, and by close observation it was found that the beetles, when feeding, failed to discriminate between sprayed and unsprayed bark.

While the experiments were conducted according to laboratory methods and the number of beetles involved was limited to about three hundred, the results are encouraging and indicate that thorough spraying of poplar plantations with an arsenical poison materially reduced the number of beetles and thereby lessened the number of eggs deposited in the trees.

Experiments are now under way in commercial poplar blocks to determine the value of this treatment. From the work that has been accomplished it is estimated that an application of an arsenical poison to nursery poplars will cost about one-fourth of a cent per tree for labor and poison.

Mr. J. B. Smith stated that this insect was not a nursery pest, although it was present in New Jersey.

In reply to a question, Mr. Parrott said that the insect was found in western New York.

Mr. Slingerland related a case which had come under his observation where infested trees had been treated with pure kerosene and the borers had been destroyed.

Mr. Washburn remarked that nursery trees from New York badly infested with this insect had been shipped into Minnesota, and he

thought that some requirements should be made to prevent such stock being transmitted.

Mr. Kirkland stated that the insect was common in Massachusetts and some of the nurseries were badly infested.

Professor Craig called attention to the fact that poplars were undesirable shade trees in the East, and raised the question as to whether this insect was not a blessing in disguise if it destroyed them. In the Northwest, where other trees would not grow, they were needed. In the East they should only be used for temporary planting: therefore it might be better to leave the growing of them to northwestern nurserymen.

Mr. Burgess remarked that poplars were undesirable shade trees, and that in East Cleveland, Ohio, where this borer had become established, and the San Jose and oyster-shell scales seriously attacked them, an ordinance had been passed prohibiting their planting.

The following paper was presented:

EFFECTS OF SPRAYS ON APHIS EGGS.

By H. E. HODGKISS, *Genora*, N. Y.

The apple aphides have been unusually abundant for several years in the orchards and nurseries of New York. The species represented are *Aphis mali* Fab., *Siphocoryne avenæ* Fab., and *Aphis malifolia* Fitch. The methods commonly used by our nurserymen in fighting the pests on apple blocks are, either dipping the stocks in oil emulsions or soapy solutions, or the direct application of these sprays to the foliage. In years when these pests are most abundant the treatment of the trees in this manner has not been entirely successful owing to belated applications and the protection derived by the insects from the curling of the leaves. As eggs on the nursery stock, especially the seedlings, have been numerous, and therefore quite conspicuous, nurserymen have often asked what would be the effects of contact sprays on eggs. As there was abundant opportunity for this work, experiments were conducted during the autumn and winter of 1904, 1905, and 1906 for the purpose of determining the comparative effects of different contact sprays upon aphis eggs.

For the purpose of the experiment, seedling apple stocks, upon which large numbers of eggs had been deposited, were selected. As it was necessary to have the conditions of the experiment under control, the trees, upon their removal from the nursery blocks, were grown in a greenhouse. The number of trees under observation was 322, and these divided into five lots. All the sprays were tested in each lot. The sprays employed were the sulphur washes, kerosene, kerosene emulsion, whale-oil soap, crude oil, Scalecide, Kil-o-Scale,

Rex solution, whitewash, K-L mixture, kerosene-whitewash, and caustic soda. The number of eggs per tree was variable, but there were not less than 400 eggs on each tree, and the maximum number on one tree was 4,800. The total number of eggs by actual count in the experiment was 223,920, of which 158,885 were firm, while the rest were more or less shrunk. The trees were planted in boxes of convenient size, and during the treatment were isolated to prevent the applications from reaching other stocks. Each tree was examined daily, and as each egg hatched the aphid was killed and a record was made of the daily hatching of the eggs.

The results of the last two tests, which are representative, are as follows:

Fourth experiment: Less than 1 per cent of the eggs hatched on trees sprayed with the sulphur wash, crude oil, kerosene-whitewash, and Rex solution. The percentages of eggs that hatched on the trees receiving other sprays are as follows:

	Per cent.		Per cent.
Scalecide	8.9	Caustic soda	7.3
Kerosene	6.7	Kil-o-Scale	26.
Kerosene emulsion	7.6	Checks	22.4
Whale-oil soap	7.8		

Fifth experiment: No eggs hatched on the trees that received applications of kerosene-whitewash and whitewash. On the trees treated with the other sprays the following percentages of eggs hatched:

	Per cent.		Per cent.
Sulphur washes	3.5	Caustic soda	5
Scalecide	40	Kil-o-Scale	4
Kerosene	1.5	Crude oil	10
Kerosene emulsion	6	Rex solution	1
Whale-oil soap	9	Checks	31.4

Mr. Titus mentioned the fact that aphid eggs are often deposited in such a manner on the twigs that they overlap, and this being the case it would be difficult, and in some cases impossible, to cover all the eggs with the spray material. In making an accurate statement as to the percentage of eggs destroyed by spraying, this point and parasitism should be considered.

Mr. Hodgkiss replied that the small size of the trees enabled a most thorough application to be made in each and every case. The purpose of the experiment being entirely to test the advisability of spraying nursery stock, the question raised by Mr. Titus would in no way affect the results obtained. Replying to a question on the value of

spraying these eggs he stated that the results show that spraying for the aphid in the egg stage is of doubtful utility, and the most effective work may be done soon after the aphides hatch.

A paper was read, entitled:

**MANNER OF BIRTH OF THE WOOLLY APHIS OF THE APPLE
(SCHIZONEURA LANIGERA HAUSM.) AND OF OTHER APHI-
DIDÆ.**

By W. E. RUMSEY, *Morgantown, W. Va.*

The exact manner of birth of the agamic forms of the woolly aphis of the apple seems to be a disputed point; at least, there is a variance in the published records that I have been able to find on the subject. In a study of this insect, now under way at the West Virginia Agricultural Experiment Station, some additional facts have been obtained along this line which may be of interest.

In the Eighth Report on the Noxious and Beneficial Insects of Illinois, by Dr. Cyrus Thomas, is a statement relating to the reproduction of this species, which says:

In so far as the method of propagation is concerned it has been shown by Dr. W. M. Smith, of New York, that it differs slightly from the true aphides, in that the young larvæ produced by the agamic females are inclosed in the thin egg-shaped covering heretofore mentioned, from which they have to free themselves in a manner analogous to hatching. The remains of this covering may often be seen attached to the tip of the abdomen, and is doubtless the supposed cottony secretion alluded to by Doctor Fitch in his description of the young larva.

A view diametrically opposed to that given by Doctor Thomas is found in the Report of the Department of Agriculture for 1879, where this insect is treated, and from which the following extract is taken:

Mr. Howard has repeatedly watched the birth of the young of the wingless agamic females and positively states that they are born without the enveloping pellicle or pseudovum. While the head and its appendages were still within the mother, he has seen the legs kicking vigorously outside.

These conflicting statements have led me to make careful observations along this line. While a student at Cornell, my study of this species seemed to corroborate the views of Dr. W. M. Smith. During my present study of the woolly aphis a large number of births have been carefully watched, which has added materially to the evidence sought.

To see the entire operation of birth satisfactorily it was necessary to devise some method by which the mother could be held in the position desired. To accomplish this a rectangular cell was made on a microscope slide with four small pieces of another slide. The cell was just wide enough to hold a mature insect when placed on its side, but of sufficient length to allow free extrusion of the young. The

pieces of glass were held in position with Canada balsam, and a cover glass was placed on top to prevent the escape of the insect. All observations of birth in this cell were made under a microscope, using a two-thirds or one-fourth inch objective.

The manner of birth, as repeatedly observed by me, is as follows: First there is a contorted movement of the abdomen of the mother, which is immediately followed by the appearance of the young aphid as an egg-like object at the anal opening. In about thirty seconds the body of the young is forced out until the eyes appear, when the movement is checked, then ceases entirely, with the mother retaining a hold on the crown of the head. At this time the dark spots that mark the eyes are the only means by which the object might at first be distinguished from an egg. The antennæ, together with the legs, are bent toward the tip of the abdomen and held closely against the ventral surface by a transparent membranous sac. This sac soon breaks at the crown of the head and is worked backward (caudad) by a continuous expansion and contraction of the body, accompanied by an interrupted pulling and pushing movement of the antennæ and legs. These motions are similar to those of an insect working its way out of a pupa case. The progress of the membrane as it moves backward (caudad) is at first indicated only by the hairs at the base of the antennæ and on the head springing into an erect position as they are freed. In from three to five minutes after the membrane begins moving off, the antennæ are liberated, and a small amount of whitish substance appears at the tip of the abdomen. Each pair of legs is then liberated in turn. As the sac works farther back its ragged edge becomes visible and the shriveled membrane at the tip of the abdomen increases in quantity. The last pair of legs is freed in from three to seven minutes after birth. As soon as the legs are freed they begin to kick vigorously, the kicking continuing from one to two minutes after the last pair is liberated; whereupon the mother lowers her abdomen and presses her offspring down, seemingly with the intention of assisting it in getting a firm footing, and at once loosens her hold. After the young is on its feet, the cast pellicle adheres to the tip of the abdomen for about two minutes, when it is worked off, leaving the insect smooth and glistening, but not yet entirely free, for a waxy thread still connects the setæ of the beak with the discarded pellicle. After several tugs and pulls the thread is broken and the insect crawls away.

Many times I have loosened the young insect from its mother, while she was holding it by the crown of its head, and carefully watched its subsequent movements. Only a slight touch with a camel's hair brush is necessary to break this hold. The motions of the body of the young and the freeing of antennæ and legs were iden-

tical to those already described, but in many such cases the young aphis died before freeing itself from the sac. This was probably due to the fact that the insect lay upon a microscope slide instead of being held aloof from the bark, as under normal conditions.

How are other species of Aphididæ born? In an attempt to answer this question I have made observations on several species, among which is the apple aphis (*Aphis mali* Fab.), which will be considered at some length. The birth of this species is similar in all the stages to that of the woolly aphis, except in one or two minor points. The legs, at least the last pair, are bent back upon themselves, so as not to extend beyond the abdomen, while being pressed to the ventral surface by the pellicle. When the young has all its legs free from the sac the mother does not in all cases, as observed with the woolly aphis, press her offspring down for a footing, but simply loosens her hold. Apparently the greater length of legs in this species makes it unnecessary for the mother to assist her young in this way. As soon as the young begin to free their antennæ and legs a delicate whitish substance appears on the tip of the abdomen, as with the other species described. The so-called honey tubes of the apple aphis, and kindred species, are bent toward the extremity of the abdomen and held closely against the body of the insect by the enveloping pellicle. As the edge of this membrane passes along over these tubes they bow up, and when the tips are released they straighten into nearly their normal position. Before the final release of the tubes the delicate transparent membrane, constituting the pellicle, can be readily seen stretched across the intervening space. Owing to the relatively much shorter beak of the apple aphis than that of the woolly aphis, I have not been able to see the waxy, thread-like connection between the setæ and the cast pellicle. If present, the separation probably takes place when the last pair of legs is being liberated, which would tend to prevent a view of the thread. When the legs of the apple aphis are free, the discarded pellicle adheres to the tip of the abdomen for a short time, whereupon it is worked off by the movements already described. In one instance, however, I distinctly saw a young aphis deliberately remove the shriveled pellicle from the tip of its abdomen with its hind legs.

In all the other species of Aphididæ which have been under consideration the manner of birth, in every case, was similar to that of those already described.

Mr. Quaintance remarked that his observations as to the mode of birth of aphides agreed with those of the speaker. He called attention to the fact of recent statements in literature to the effect that the so-called honeydew of aphides was excreted only from the anus.

the idea that some was secreted also from the cornicles being erroneous. He asked for observations of the members present on this point.

In reply to a question by Mr. Slingerland as to the number of aphides whose birth had been observed, Mr. Rumsey stated that he had watched the manner of birth of probably fifty individuals. He also stated that he had never seen honeydew secreted from the honey tubes, but always from the anus of the aphides.

Four papers were then presented, and discussed at the close of the last paper.

UNUSUAL INSECT HAPPENINGS IN NEW JERSEY IN 1906.

By JOHN B. SMITH, *New Brunswick, N. J.*

The summer of 1906 in New Jersey was remarkable for continuous rains during the middle of the season, favoring the development of some crops and insects and interfering seriously with others.

It seems contradictory to include the failure of the periodical cicada (*Tibicen septendecim* L.) to appear in New Jersey as scheduled as a "happening," yet it deserves to be recorded as such. In 1889 there had been no definite New Jersey localities for what was then known as Brood VIII; but in that year I secured four, at such points as to indicate a diagonal line of occurrence from the Palisades southwest to the Delaware a little south of Camden. The insect was nowhere very common and I doubted its reappearance in 1906. To cover the ground as thoroughly as possible I asked every correspondent of the office along this line to communicate with me, asked the members of the Newark, New York, and Philadelphia entomological societies to keep a lookout for specimens and records, and instructed Mr. E. L. Dickerson, one of my assistants, to cover the most likely area of appearance personally, besides keeping an open eye for "signs" in the course of his orchard and nursery inspection work. Mr. Dickerson was also asked to verify reports that were received, and this proved a wise precaution, since at least three records, apparently safe and in the proper localities, proved on investigation to be based on error. Not a single definite record did I get for New Jersey. Mr. Marlatt was good enough to send me a record from Bergen County, his correspondent claiming to have seen two specimens and to have heard of others from several localities. This record has not been verified, but may easily be correct, although my own correspondents in that same region failed to find anything. In any case nothing but a fragment of the brood remains in the State.

Incidentally, my attention was drawn to the other cicada species in the State, especially after the publications of Osborn, Davis, and

Joutel. It was found that all the forms occurred in New Jersey, and that we had also yet another which was readily identified as the real *pruinosa* of Say, leaving the species heretofore identified as *pruinosa* without a name. Mr. Grossbeck has proposed the name *sayi* for this form. To make certain just which form Linné intended as *tibicen*, the original descriptions and figures were studied, and it seems entirely certain that the original *tibicen* is a totally distinct species from the one we have heretofore determined as such. For the species heretofore known as *tibicen* the term *linnei* is proposed.

Perhaps the most interesting feature now in progress is the steady increase in numbers of the rose-chaffer (*Macrodactylus subspinosus* Fab.). When I first came into the State, seventeen years ago, a scourge was in progress that culminated in 1890 after a period of gradual rise, and I was told then that some sixteen to twenty years before there had been a similar trouble, followed by a series of years when little was seen of the insects. We have had our series of exempt years, and now for the four years last past the beetle has become increasingly abundant. In those same localities where it appeared as a pest sixteen to twenty years ago it is again a pest, but not yet quite as bad as it was in 1890. I anticipate another year of increase before the culmination is reached. As to the cause of the decrease I could find nothing. There were no apparent parasites; but the larvæ seemed simply to become less plentiful each year until little was seen of them. As to remedial measures practically no progress has been made. The insects are killed by arsenites, but, especially when they attack grape, the mischief is caused before the poison can do its work; to say nothing of the difficulty of getting a sufficient supply of it on the buds. I am advising our growers for the city market to bag the most valuable varieties and all others that it will pay to protect in that way. Bagging as a protection against rot has been entirely discarded in New Jersey in favor of spraying with the Bordeaux mixture.

The army worm (*Heliophila unipuncta* Haw.) has, for the first time in many years, appeared as a pest to field crops in a limited district in southern New Jersey. It was promptly dealt with and did very little harm, but the interesting feature was the practical absence of the tachinid parasites that usually attack so large a percentage of the caterpillars. Always on previous occasions I have found an injurious army with the seeds of its own destruction apparent, but in this case there was so little infestation that practically all the larvæ collected and placed in breeding cages in the laboratory came to maturity. It will be a matter of very great interest to watch developments in southern New Jersey in 1907. In this connection it might be said that some army-worm injury occurs almost every year

on our cranberry bogs and that, occasionally, there is so great an increase as to result in actually stripping a section of bog.

The cottony maple scale (*Pulvinaria innumerabilis* Rathv.), which has been the theme of papers through many localities in the north-western section of the State, reached its culmination in New Jersey in 1905, and in 1906 dropped out of sight completely in those places where it had been most abundant the year before. It was a curious repetition of an experience about eight years ago, although the agent of control this time was apparently different. Mr. E. L. Dickerson's paper covers that ground fully and it needs only a mere mention here.

The elm leaf-beetle (*Galerucella luteola* Müll.) also, after a period of two years during which no spraying was required at New Brunswick, has taken a new start, and in 1906 the unsprayed trees in parts of that city were almost completely defoliated. The trees on the campus of Rutgers College were protected with arsenate of lead, and I proved to my own satisfaction that the 12-cents-a-pound material made by one insecticide company was quite as effective and satisfactory in all respects as the 20-cent product of another company, while a 17-cent brand was inferior in arsenical content and was short weight besides. It should be added here that the controlling agent in this case is a disease that attacks the pupæ if the weather at that period of development is damp. In 1904 it was only moist, and the disease was not very prevalent; in 1905 the weather was hot and dry, the beetles all developed normally, and I prepared for the danger that I felt certain would come in 1906. I was not at home during the pupation of the brood last summer, so can say nothing as to probabilities for 1907.

The common oyster-shell scale (*Lepidosaphes ulmi* L.) has developed possibilities as a serious pest and has proved quite difficult of control in the more southern parts of the State. One of the Burlington County apple growers declares it more dangerous and difficult to deal with than the San Jose scale. There are two broods of it in that section of New Jersey, and in one of the towns it has developed as a serious pest to shade trees, especially maples.

Away off in one part of southern New Jersey is a little section of land especially adapted to peach growing and where fine trees bear excellent crops of good quality. In this corner and nowhere else in the State the peach soft scale (*Eulecanium nigrofasciatum* Pergande) has established itself, and our effort now is to prevent its getting away from there. Fortunately the area is completely isolated, and there is little or no chance of a natural spread, while no trees are grown for sale anywhere in the infested territory. Few trees are badly enough injured as yet to induce the growers to consider active measures, and matters will probably become a great deal worse before they become much better.

The catalpa sphinx (*Ceratomia catalpa* Bdv.) now covers about the entire State of New Jersey, positive records being absent from one county only. It always seems to be worse the second year of its appearance in a given locality, and it has been about as troublesome in nurseries as anywhere.

Another failure to establish the Chinese mantid (*Paratenodera sinensis* Sauss.) in New Jersey is to be recorded. A large number of egg masses were tied out in an ideal location on the southeastern slope of the Orange Mountains, and most of these fell a prey to field mice. It seems curious that the insects should do so well near Philadelphia and that they should fail so uniformly in all sections of New Jersey.

The Asiatic ladybird (*Chilocorus similis* Rossi) has not been found again, although the orchard in which the lots sent up from Georgia three years ago were freed still stands unsprayed—what is left of it.

Although not strictly entomological, mite infestation should be noted as among the most important happenings of the season. Trees and shrubs of the most diverse kinds were infested and a great deal of foliage was disfigured, if not seriously injured.

MISCELLANEOUS INSECT NOTES FROM MARYLAND FOR 1906.

By A. B. GAHAN and G. P. WELDON, *College Park, Md.*

Present indications are that the fruit growers of Maryland may have another serious scale pest to contend with in the near future. We refer to the terrapin scale (*Eulecanium nigrofasciatum* Pergande). From different localities in Washington County have come three complaints of very serious injury to peach trees by this scale. All told, several hundred trees have been killed or badly damaged by it. The growers report that the lime-sulphur-salt treatment is not effective against this pest, it being no uncommon thing to see full-grown scales in midsummer with a coating of the spray mixture still adhering to their backs, but apparently none the worse for it.

A more or less careful study of the life history of the insect was made at the Station the past season, and it brought out the following facts: The scales pass the winter as immature females, finishing their growth in the spring. The eggs are deposited beneath the female scale, and are very numerous. Hatching begins about June 1 in our latitude, and crawling young may be found from that time until the second week in August. The young apparently all go to the leaves immediately after hatching, and there settle along the midribs and veins, where they remain for a period of from six to eight weeks. The males then emerge, winged, while the females migrate back to the twigs, settling along the under side of the twigs and branches. This migration of the females began about July 20, and by September

1 practically all were gone from the leaves. Copulation was not observed, but probably took place during the migratory period.

Two important parasites were noted, one a braconid, the other a fungous disease (probably *Cordyceps clavulata*). The fungus first made its appearance in the latter part of July, when a few individual scales on a single tree were noticed with a whitish discoloration about the edges. In a few days the discoloration had spread over the whole bodies of these insects and had also infested many others, both young and old. Within three weeks the disease attacked a majority of the insects on the five infested trees that were under observation. By the 1st of November it had apparently cleared four of the trees of the scale, while the fifth tree showed only occasional healthy specimens, and these were usually isolated on the tip of a twig. The very damp season no doubt accounts to some extent for the activity of the fungus.

A small brood of periodical cicadas (*Tibicen septendecim* L.) appeared in Washington County in July. They belonged to Marlatt's Brood XIV, and are confined to a very limited portion of the country, principally in the neighborhood of Mapleville. While quite numerous in that locality, the brood did very little damage, and its occurrence is important only as a matter of record.

Some observations were made to determine the life history of the codling moth (*Carpocapsa pomonella* L.) for the latitude of Maryland. About 75 per cent of the first brood of larvæ were found to enter the fruit at the calyx end. It was found by banding trees that the greater number of larvæ of the first brood reach maturity and pupate between June 20 and July 10, but that quite a considerable number do not leave the fruit until later, some first-brood larvæ being taken as late as August 10. Eggs from the first-brood moths were collected quite plentifully during July on both leaves and fruit. The majority of the second-brood larvæ were found to enter the fruit at other than the calyx end. The larvæ of this brood begin coming down about August 1, and the number collected from beneath the bands showed a constant increase from that time until August 25, when it began to decrease. A few larvæ collected during August pupated, but the great majority spun cocoons and remained as larvæ. Undoubtedly the few pupating individuals were belated first-brood larvæ. An interesting case of destruction of codling-moth larvæ as they were pupating, by a small red ant, which was not identified, was observed. Scarcely a collection of larvæ was made from the banded trees without finding several worms that were being devoured by the ants.

In November a report was received of injury to chestnut telephone poles by a flat-headed borer of some kind. Specimens of the borers and their work were secured and taken to the Bureau of Entomology,

U. S. Department of Agriculture, Washington, D. C., for identification, but Doctor Hopkins reported that he was unable to identify it even generically. The larvæ collected were about an inch in length, white, with the first three or four segments much broader than the remaining. They work below the surface of the ground in the sapwood of the pole, some penetrating an inch or more into the wood. The pole from which our specimens were collected was very badly infested, there being probably four dozen borers in it. The borers weakened it so much at the surface of the ground that it had to be removed. The infested butt has been secured and placed in the insectary in the hope that adult specimens of the insect may be secured.

NOTES ON INSECTS OF THE YEAR 1906 IN NEW YORK STATE.

By E. P. FELT, Albany, N. Y.

The leaf feeders, such as the yellow-necked apple-tree caterpillar (*Datana ministra* Dru.), the red-humped apple-tree caterpillar (*Schizura concinna* S. and A.), the hickory tussock moth (*Halisidota caryæ* Harr.), and the black walnut worm (*Datana integerrima* Grt. and Rob.), have received more attention than usual owing, probably, to the wide-spread interest which led many to keep a close watch for the possible occurrence of either the gipsy or brown-tail moths (*Porthetria dispar* L. and *Euproctis chrysorrhæa* L.). A placard, describing these two insects briefly and figuring them in colors, was distributed in many sections of the State where there was likelihood of the pests becoming established. It is gratifying to state that no undoubted evidence of even their casual occurrence in New York State was secured, despite newspaper statements to the contrary. Every report regarding these species was followed up and in each instance found to be based upon insufficient information. An occurrence out of the ordinary was the capture in Albany of a large South American moth (*Thysania zenobia* Cramer).

The scurfy scale (*Chionaspis furfura* Fitch) has continued abundant in the Hudson Valley, being especially numerous on young fruit trees in the vicinity of Annandale.

Experiments with the San Jose scale (*Aspidiotus perniciosus* Comst.) have been continued and the weight of evidence is decidedly in favor of employing a lime-sulphur wash, which, if properly prepared and thoroughly applied, gives fully as satisfactory results as any other preparation. Several experiments were conducted largely for the purpose of determining whether this wash could be further modified to advantage. A lime-sulphur wash was made in the normal manner, except that the ordinary local burnt lime was replaced by a finely prepared hydrated magnesium lime known as

limoid, and the results were decidedly adverse to the employment of this material, despite its improved physical condition. The reaction between the limoid and the sulphur was not nearly so vigorous, even when hot water was employed. A sal-soda lime-sulphur wash was the subject of further tests and gave very satisfactory results. Oily preparations, known in a general way as "soluble" or miscible oils, have been used to some extent in the State. We have examined a number of trees treated with these materials and in each instance detected evidences of oil injury, though there was no doubt that, in some cases at least, a considerable proportion of the scale insects had been destroyed. It is impossible to say at present what would be the result of successive annual applications of such materials, though we would not be surprised were a considerable injury to develop with the advance of time.

The grape root-worm (*Fidia viticida* Walsh) continues abundant in the Chautauqua region, and during the past season has been exceptionally numerous in certain vineyards. It is an insect very local in its operations, and this fact renders it difficult to make any general statements. There is no question that it is becoming more abundant in certain vineyards here and there throughout the grape belt, and its numbers have increased materially the past year or two in the vineyards on the hills back from the lake. The grapevine or steely flea-beetle (*Haltica chalybea* Ill.) had an exceptional opportunity to injure grape buds, owing to the continued cold weather keeping the vines in check just as the foliage began to appear. The snowy tree-cricket or white flower cricket (*Ecanthus niveus* De G.) was so abundant in certain vineyards, where there was considerable weedy growth, as to injure many of the canes by depositing its eggs therein.

The sugar-maple borer (*Plagionotus speciosus* Say) continues to be a serious enemy of our sugar maples. An exceptionally interesting observation of the work of this insect was made in connection with a recent trip to Le Roy. Seven years before, namely, in September, 1899, we photographed a tree in that city which had been badly injured by this borer. The tree was about 18 inches in diameter and at the time of photographing was rather thrifty despite the fact that one side was completely girdled by the operations of the pest. The dead area at that time had commenced to enlarge and it was therefore not surprising, on examining this tree in November last, to find that the area of exposed wood had greatly increased. The original gallery was approximately 4 feet from the ground. At the present time the entire affected side, from the ground to 8 or 10 feet above, is dead, the bark has decayed or fallen away, and a large proportion of the magnificent limbs and branches on that side of the tree have disappeared. This illustrates in a striking manner the destructive nature of this insect's operations. It is very probable

that an injury of this kind could be helped by bridge grafting, and it is presumable that extremely beneficial results would have been obtained even if this means had not been employed until two or three years after the initial injury, provided the bridge grafts were inserted in rather vigorous tissues.

The white-marked tussock moth (*Hemerocampa leucostigma* S. & A.) has been the cause of extensive injury the past season to shade trees in Buffalo, Lockport, Geneva, Rochester, Syracuse, Utica, Albany, Troy, and Brooklyn. They were so abundant in many of these places as nearly to defoliate a large number of trees. An extremely interesting phenomenon was observed in Capitol Park, Albany, July 5. Many of the trees were then badly infested by the tussock moth, some being almost entirely defoliated. On that morning the ground was thickly strewn with leaves and leaf-bearing twigs. Most of the latter bore from 3 to 5 or 6 leaves, and the cut end had invariably been completely girdled for a distance ranging from one-fourth inch to nearly an inch in length. This peculiar form of injury was first observed by the late Doctor Lintner in 1883, he having actually seen the caterpillar engaged in the girdling operations. Subsequently questions arose as to whether the depredator was correctly identified. It is gratifying to state that the trees in Capitol Park, mentioned above, were infested by practically no other insect. The tussock moth caterpillars were in several instances observed upon the fallen twigs, and there is, therefore, every reason to believe that this larva was the author of this somewhat anomalous injury. The girdling was limited, as was also observed by Doctor Lintner, to the new growth, and as the past summer has been exceptionally moist it is barely possible that there is some connection between a rapid, succulent growth and this form of injury, since it is only occasionally that the larvæ girdle the twigs as reported above.

The elm leaf-beetle (*Galerucella luteola* Müll.) has been abundant and injurious in certain sections, such as Oyster Bay, Ossining, Albany, Troy, Fort Edward, and Ithaca.

The false maple scale (*Phenacoccus acericola* King) appears to be establishing its claim as one of the most serious pests of the hard maple in New York State, since it has been injurious to trees in the vicinity of New York City in particular during the past four or five years. It was very abundant last summer at Port Chester, Middletown, and probably in other villages in the southern part of the State. The elm bark-scale (*Gossyparia spuria* Mod.) continues quite destructive, displaying a marked preference for the Scotch elm.

The violet gall midge (*Contarinia violicola* Coq.) is a very serious pest of the extensive violet-growing industry in and about Rhinebeck, N. Y. The crop in infested houses, according to estimates of growers, is reduced from one-third to one-half by the operations of

this insect. Should this infestation become more general, the results might be exceedingly serious to the industry as a whole. An examination showed that the insect was distinctly local in its operations, since one-half of a house 150 feet long might be seriously injured, while the other half might be almost free from attack. Even in small houses there were distinct areas which suffered more severely than others, sometimes these being limited to only a square yard or two. The continued breeding of this species appears to be affected largely by temperature, since houses where the mercury was not kept below a certain point were decidedly more infested than others. The growers are almost unanimous in stating that when the temperature of a house can be kept down to 40° at night and does not rise to over 60° in the daytime there is little or no injury from the pest. The flies, according to growers, very rarely leave the plants, and can be observed only by flushing them with the hand. An examination was made in houses where there were flies and numerous larvæ on the plants, but none was to be found on the windows nor in the sheds at the ends of the houses, nor in cobwebs spun here and there about the structures. The insect displays a marked preference for recently opened leaves, apparently depositing its eggs in those which have just expanded fully; and, according to the growers, leaves perfectly straight one day may be badly curled the next. They note that leaves can be curled in a few hours, and are of the opinion that only a day or so elapses between the depositing of the egg and the curling of the leaves, an operation which protects the larvæ from most insecticides. Furthermore, several of them said that fumigating with hydrocyanic-acid gas apparently has no effect whatsoever in destroying the larvæ, though there is little doubt that the flies are killed. There is a marked periodicity in the appearance of the larvæ. Last summer they were first noticed in numbers early in July and then they became abundant again in August. Experience this year has shown that they may continue working in numbers even as late as the latter part of October. A number of infested leaves were placed on soil on the 12th, at which time no pupæ were manifest. The first adults appeared on the 22d and others emerged subsequently to the 26th, when about four were bred out. Another individual was obtained November 3 and lived to the 5th, at least. Owing to the fact that the plants could not be well cared for, it is probable that the period of the appearance of the flies was somewhat abridged by the unnatural conditions. The above data show that not over ten days is necessary from the time the larva forsakes the plant until the appearance of the perfect fly, and it may possibly be a little less. No pupæ were observed on the leaves, and there is no doubt that the insect normally undergoes its final transformations in the soil.

The periodical cicada (*Tibicen septendecim* L.) appeared in considerable numbers on Long Island during the past summer. The list of localities, compiled from various correspondents, is as follows: Wading River, Port Jefferson, St. James, Farmingville, Coram (on the road from Port Jefferson to Patchogue), Manorville, Eastport, East Moriches, Center Moriches, Commack, Brentwood, Cold Spring Harbor, Laurelton, Huntington, Oyster Bay, East Norwich, and Syosset. There is also a record of its appearance in very limited numbers on Staten Island.

THE PEAR BLISTER-MITE.

(*Eriophyes pyri* (Pgst.) Nal.)

By P. J. PARROTT, *General*, N. Y.

This mite was undoubtedly introduced at an early period into the United States in foreign importations of nursery stock and was probably well distributed in many fruit-growing areas long before its presence was recognized. The first writer to direct attention to its appearance as an orchard pest in this country was Mr. Townend Glover,^a Entomologist of U. S. Department of Agriculture, who in May, 1872, received specimens of its work from a correspondent. Under his direction Mr. Thomas Taylor, microscopist, examined some of the pear leaves covered with dark-brown blotches, which were said to be inhabited by myriads of small mites. The species was thought to be somewhat similar to the mite mentioned by Packard^b as "*Typhlodromus pyri* of Scheuten," known to infest pear trees in Europe.

Before the meeting of the American Association for the Advancement of Science, held at Saratoga, N. Y., August, 1879, Dr. W. S. Barnard^c presented a paper on "bud-blight insects," in which he attributed the brown and black blotches of pear leaves to the activities of mites. In 1880, Prof. T. J. Burrill^d called attention to a widespread disease of pear leaves in Illinois and in the country at large, which was ascribed to the work of the mite *Phytoptus pyri* Scheuten. He mentioned the fact of the hibernation of the mites under the bud scales and the probable dissemination of these creatures in cions and buds. In succeeding years the work of this species was recognized in many States and was given widespread mention. In 1883^e the mite was observed in large numbers upon imported Russian pear trees in experimental nurseries in Iowa, and by 1894

^a Report of U. S. Dept. of Agriculture, 1872, p. 113.

^b Guide to the Study of Insects. By A. S. Packard. 1869.

^c Scientific American, Dec., 1879, p. 3302.

^d Gardener's Monthly, v. 22, 1880, pp. 18-19.

^e Osborn, H., Iowa State Hort. Soc., 1883, pp. 127-135.

the mite was regarded as being generally distributed in the leading pear-growing States. The mite has attained its prominence in this country as an orchard pest because of its destructive work on pear foliage, and for this reason our literature upon the species is entirely concerned with the economy of the mite in its relationship to pear growing.

In recent years our attention has been called to the appearance of the mite in another rôle—as a pest of apple foliage. The mite has quite likely been active in this capacity for many years, but important injuries by it were not brought to our notice till 1902, when it was found to be very abundant in an apple orchard at Williamson, on Lake Ontario. In 1903 the infestation of these trees was much more conspicuous, and the strange appearance of the foliage attracted much attention from the fruit growers in attendance at the summer meeting of the State Fruit Growers' Association which was held in Geneva. Up to the present time its injurious numbers in this orchard have been maintained. During the same year Prof. M. V. Slingerland ^a observed, through the central portion of the State, numerous apple trees with many of their leaves showing the corky blisters characteristic of the work of this creature. The area in which the mite was present in conspicuous numbers has increased each year, and in 1905 marked infestation of many orchards in Wayne, Ontario, Monroe, and Niagara counties was noted. In his apple survey of Wayne County in 1903, Dr. G. F. Warren ^b recorded the presence of the mite in 53 orchards. It is stated that “the mites were not bad in more than one-half of a dozen orchards, but in a few orchards some trees had practically every leaf affected.” A like survey of Orleans County ^b in 1904 showed somewhat similar conditions of apple trees with respect to this pest. Of 19 orchards showing mite injury, 4 were recorded as seriously infested, 4 considerably infested, and 11 slightly infested. During 1906 the work of the mite again attracted much attention among fruit growers, and in addition to the above counties the species was also present in large numbers in apple orchards in Livingston, Wyoming, Seneca, and Yates counties. The mite may be said to be common in our leading apple-growing sections in western New York, and its work upon apple leaves has also been recognized in Pennsylvania and Illinois.

The host plants of this mite, in addition to the apple and pear, are, as recorded by Dr. Alfred Nalepa, the service-berry (*Amelanchier vulgaris*), the common cotoneaster (*Cotoneaster vulgaris*), the wild service tree (*Sorbus terminalis*), the white bean tree (*Sorbus aria*), and the European mountain ash (*Sorbus aucuparia*).

^a Bul. 46, Bureau of Entomology, U. S. Dept. of Agriculture, 1904, p. 72.

^b Cornell Bul. 226, 1905, p. 340; Bul. 229, 1905, p. 489.

The work of the mite on apple first shows on the upper surface of the leaf as distinct light-colored pimples, and on the underside as blisters or thickened areas of the same color as the leaf. The affected areas are of irregular size and are unevenly distributed, though the larger proportion of them are about the sides and the base of the leaf. Some of the blisters may have a reddish tinge, somewhat similar to the pear leaf-galls, but they are usually of a less brilliant color. As the galls become older they appear as corky spots of a reddish-brown color, and not black, as with the pear, which to the touch are very distinct from the healthy portions of the leaf. The individual galls average from one-twelfth to one-eighth of an inch in diameter and are usually oval or quadrangular in shape. Leading to the interior of each affected area there is usually one or more tiny openings. Often the spots coalesce, forming large irregular dead areas with smooth or slightly raised surfaces and of a dark-brown color, which rupture the leaves at the margins. About July 1 the most striking effects of the mites upon the leaves appear, especially if there is much yellowing of the foliage, as frequently occurs. Upon the upper surface of such leaves the mite-infested spots are of a light-brown or of a dark-green color and are uniformly brown beneath. These spots are usually thickly massed, forming a dark, broad band of irregular width along each side of the leaf, which contrasts conspicuously with the intervening light-yellow area about the main rib. The mites also cause pimples about the calyx cavity and on the stems of young apples. In several instances the work of the mites upon the leaves and fruit stems of the same cluster had so weakened the stems as to cause the falling of the fruit. Premature dropping of the apples by this means seems to be of rare occasion, and even on the worst-infested trees the loss of fruit is not appreciable. This seems incredible, as it does not seem possible that the foliage of the trees could be so completely overrun with mites without losses in crop production.

As has been stated by other observers, notably Prof. T. J. Burrill, in his study of the species on pear, the mites spend the winter in the buds, preferably under the second and third layers of the bud scales. As the buds burst there is a movement of the mites to the unfolding leaves, in which eggs are deposited. This migration takes place with the maturing of the bud scales, during the latter part of April and early May, depending on the season, soon after which the discoloration of the leaves by pale and red-colored spots occurs. On pear foliage the galls are largely grouped in a row on each side of the main rib, while on apple leaves the affected areas appear in the greatest numbers about the sides and the base of the leaf. This difference in the arrangement of the diseased spots on apple and pear foliage seems to be determined by the manner of the distribution of the pubes-

cence of the leaves and the condition of the leaves when unfolding, which differ slightly with these two kinds of fruits. Mature mites may be found in the leaf tissues during the first week in May, and from then on into September eggs and larvæ are present in the galls of the leaves. During the latter part of May and first part of June the mites in greatly increased numbers may be found in the pubescence of the new wood and in the fruit and leaf stems and upon the unfolding leaves of the new growth. During October they largely abandon the leaves and swell the numbers already in hiding in the buds and in the pubescence of the bark of the new wood. Hibernation occurs under the bud scales and apparently none of the mites passes the winter in the pubescence or in crevices of the bark.

Treatment for the mite is much more difficult on apple than on pear trees, mainly due to the larger size of the trees and the greater abundance of the pubescence on the buds and the new wood. In our experience the crude and refined oils, either clear or emulsified, have proved the most efficient sprays. Because of its comparatively safe qualities and cheapness, kerosene emulsion diluted with 5 parts of water appears to be the most practicable remedy for the spraying of apple orchards when treatment is advisable, the applications being made either in the late fall or early spring before the buds swell.

In the study of the mites on apple and pear foliage four other species of Eriophyidæ have been recognized. These are *Eriophyes pyri*, var. *variolata* Nal., *Eriophyes malifolia* Parr., *Phyllocoptes schlechtendali* Nal., and *Epitrimerus piri* Nal. With the exception of the first named, these are vagabond species and seem to thrive on the underside of the leaves. *Phyllocoptes schlechtendali* and *Epitrimerus piri* are foreign species and appear to be more common here than on the Continent. The behavior of these two species in the future is a matter of much interest, as both, because of their large numbers, seem to show possibilities of developing to greater economic importance.

In discussing these papers, Mr. J. B. Smith brought out the fact that experiments were being conducted for the purpose of preparing arsenate of lead by the action of electricity on lead. Some good results had been secured, and the process, if perfected, promised to cheapen the price of this insecticide. He had tried arsenate of lead made by several manufacturers.

Mr. Kirkland stated that about 200 tons of arsenate of lead had been used the past season for fighting the gipsy and brown-tail moths. It was applied at the rate of 1 pound to 10 gallons of water, and no burning of the foliage was observed.

Mr. Quaintance called attention to some tests of various arsenicals on peach foliage made by the Bureau of Entomology during 1906, and stated that all arsenicals used—as arsenate of lead (homemade) and the principal proprietary brands, Paris green and Scheele's green—were injurious either to the foliage or fruit. A new insecticide arsenical was tested, namely, arsenic sulphid, which was stated by chemists to be quite insoluble. Nevertheless this proves to be more injurious than any of the other arsenicals used—in fact, killing several of the trees outright.

Mr. Webster recalled the fact that in 1888, while in Tasmania, he found the pear mite and also a species of fungus in connection with it. Dr. J. C. Arthur had told him that this fungus was always associated with this particular mite.

Mr. Parrott stated that the attack of these mites on foliage was sometimes mistaken for the pear scab fungus (*Venturia pirina* Aderh.).

Mr. Taylor remarked that arsenate of lead was being used to a considerable extent in Colorado. During the past year 16 carloads had been applied with good results. He had used it on peach trees for the twig borer (*Anarsia lineatella* Zell.) and found that it worked satisfactorily. It can be used on these trees if it carries no free arsenic. In some cases the branches had been killed by burning. This trouble is usually indicated by the appearance of purple spots on the young wood. The rainfall in this section is 7 inches per year, and this may have some bearing on the effect of poison on the foliage.

Mr. Slingerland called attention to the use of Scalecide and other miscible oils. He stated that in an orchard badly infested with *Aspidiotus perniciosus* that had come under his observation excellent results had been secured when scalecide was applied.

Mr. Burgess remarked that he had used Scalecide and Kil-o-scale on a badly infested apple orchard last spring. The manufacturer's directions required the mixing of 1 gallon of these materials to 20 gallons of water. Tests were made using 1 gallon to 9, 14, 19, and 24 gallons of water. Satisfactory results were secured when Kil-o-scale was used at the rate of 1 gallon to 14 gallons of water, but when Scalecide was used at the rate of 1 gallon to 9 gallons of water the San Jose scale was simply held in check.

Mr. J. B. Smith stated that the difference in these two substances was not very great, although the Kil-o-scale carried more sulphur and 5 per cent less actual petroleum.

A general discussion of the subject followed, which was participated in by Messrs. Brooks, Taylor, Slingerland, J. B. Smith, Fernald, Quaintance, Britton, and others. This brought out the fact that variable results had been secured with these substances, as far as

killing the scale was concerned, but no injury to the trees was reported. It was considered advisable to use a stronger solution than the one recommended by the manufacturers.

Mr. Slingerland called attention to the fumigation of greenhouses for controlling the violet fly (*Contarinia violicola* Coq.). Some large violet growers had suffered severe loss from this insect. It had been found that less than one-half strength of cyanid, as often recommended, may destroy the foliage of violets. It therefore seemed quite necessary for some further tests to be made along this line.

Mr. Sherman gave some general notes from North Carolina, and stated that where Scalecide had been used at the rate of 1 gallon to 15 gallons of water in the fall, and lime and sulphur had been applied in the spring, good results had been secured. The fall armyworm (*Laphygma frugiperda* S. & A.) did considerable damage during the past season. Rose bugs (*Macrodactylus subspinosus* Fab.), although usually not an important pest, defoliated liquidambar trees over large areas.

Mr. Brooks stated that the larvæ of rose bugs are eaten by the short-tail shrew. He had observed this in West Virginia during the past season.

The meeting then adjourned until 10 a. m. Saturday.

MORNING SESSION, SATURDAY, DECEMBER 29, 1906.

The meeting was called to order by President Kirkland at 10 a. m., and the following paper was presented:

SOME OBSERVATIONS ON THE NATURAL CHECKS OF THE COTTONY MAPLE SCALE.

(*Pulvinaria innumerabilis* Rathv.)

By EDGAR L. DICKERSON, *New Brunswick, N. J.*

In the various accounts of the cottony maple scale (*Pulvinaria innumerabilis* Rathv.) several parasitic and predaceous enemies have been recorded as infesting this insect and aiding, to some extent at least, in keeping it in check. A notable incident is that mentioned by Dr. L. O. Howard in his account of this species, in which he states that, because of the parasite *Coccophagus lecanii* Fitch, "it was found almost impossible to carry the scale insect through the season at Washington in 1898." The forms acting as the most effective checks to the scale may differ in the various localities, and apparently even in the same locality at different periods. For example, it was stated that in Washington in 1879 the most effective enemy of this insect

was the larva of *Dakruma coccidivora* Comst. Some of the enemies, however, appear to be found wherever the scale occurs, and prominent among these are the parasite *Coccophagus lecanii* Fitch and the coccinellid beetle *Hyperaspis signata* Oliv. and its larva. It has been due largely to these species that the scale has been controlled in New Jersey during the past two years.

In 1904 *Pulvinaria innumerabilis* was noted rather plentifully in several places in Newark and near-by towns and at New Brunswick, and the indications were that it would be much more plentiful the following season. Our anticipations were realized, and in 1905 the scale occurred more abundantly than it had for several years. In order to watch its development the infested districts were visited from time to time and the condition of the insects and the trees noted. As the year progressed it was observed that the enemies previously mentioned were materially and effectively checking the scale, and it was predicted that the number of the insects would be considerably reduced and in some localities nearly exterminated in 1906. These predictions were likewise fulfilled; and in Montclair, where some of the worst infested districts occurred in 1905, there has been very little evidence of the *Pulvinaria* during the past year.

The first signs of parasitism were observed in the middle of April, after the fertilized female scales had started to develop. While most of the insects at that time were becoming enlarged, a few were observed to be quite convex, and an examination showed that they were infested with parasitic larvæ, a few of which were full grown. Only a single larva occurred in each of the scales, which were light in color at this time, but as the larvæ pupated the parasitized scales became dark and hard. From this time on the number of the latter increased, and by the middle of May two and one-half times as many parasitized as good scales could be observed on some of the leaves.

The adult parasites were first observed about the middle of May and continued in increasing numbers until early June, after which they began to decrease, and were last observed about the middle of that month. Examples were sent to Dr. William H. Ashmead, who determined them as *Coccophagus lecanii* Fitch. Whether this parasite hibernates as a larva or in the egg stage I am unaware. It is apparent, however, that the larva begins to develop some time in early April, and the species continues in evidence for a couple of months. In emerging it cuts an irregularly rounded hole in the dorsal surface of the scale, and, so far as I could observe, it was always nearest the posterior end.

Thus the parasites had destroyed many of the hibernating females, but the number of the latter which went into hibernation was so large

that, in spite of these parasites, there still remained a goodly number to oviposit and reproduce.

The egg masses began to develop early in June, and by the middle of the month many of them were apparently full sized. In Montclair, where the worst-infested district was observed, the insects were so abundant as to form continuous lines along the underside of many of the infested twigs and branches. An examination of the egg masses there and in other localities revealed the fact that some of them were infested by the larva of *Hyperaspis signata*.

This larva, because of the cottony-like waxy covering of the dorsal surface, is well protected among the egg masses, especially since it is often found feeding within the egg mass, when the latter may appear perfectly normal. Often only a single larva infests an egg mass; but sometimes more will be found, especially when they are young. On the other hand, a single larva may destroy a large proportion of the eggs in several masses.

On June 16, when the coccinellid larvæ were first observed at Montclair, a few of them were nearly, if not quite, full grown, while others were very small. From this time on their number increased until the early part of July, when they began to decrease. By June 28, in Montclair, scarcely an egg mass could be found which was not or had not been infested by them. The first pupæ were observed both in the laboratory and on the infested trees on June 24. A few of the larvæ at this time could be observed crawling on the trunks of the trees and getting beneath the loose bark, where the pupæ were found. The pupa is brown in color, but covered more or less with the white material from the cast larval skin. The earliest pupæ were found, as just stated, on the trunks of the trees under the loose bark and in the crevices, and it has been stated that the insect always goes to these places to pupate. But this is not the case, for most of the later pupæ were found within the eaten-out egg masses. The truth is that the insect desires to pupate in a protected situation, whether under the bark or elsewhere.

The first beetles to emerge in the laboratory were observed on July 7 and came from pupæ which had been in that stage for two weeks. It was about this time also that beetles were noted on the infested trees, and they continued in evidence until the early part of August, after which no more were observed. While the larva of the coccinellid fed on the eggs of the *Pulvinaria*, the beetle itself devoured the young scales which set on the leaves. The young scale adheres closely to the leaf, and it was interesting to watch the beetle in its efforts to secure it. If the latter was not too closely set the beetle would successfully pull it up and then rapidly devour it; but sometimes the scale was too securely fastened, and then the coccinellid,

after making several futile attempts to get its mandible under the side of the scale, would finally give it up and attack another specimen.

The great number of scale insects destroyed by the coccinellid and its larva can only be realized when we consider the large number of eggs deposited by the Pulvinaria. At Montclair, where these insects and beetles were most numerous, between 500 and 1,000 young scales were counted on each of several leaves and there were many leaves just as badly infested; but after the coccinellid had completed its work not a leaf could be found with more than a dozen scales set on it, while most of the leaves showed still fewer.

In looking over the long series of beetles bred from larvæ it was found that there were some variations in both the males and females in size and markings. The majority of the specimens were black in color, with the single reddish spot on the disk of each elytron, but a few examples were found which showed an extra smaller red spot near the tip of each elytron. This latter is in all probability the form which was originally described by Olivier as *signata*, while the form with the reddish spots, which occurs most commonly, is the one Say described later as *binotata*. Olivier's name, therefore, has precedence and should be the one by which the species is designated.

It is interesting to note also that while in 1905 this species was found almost exclusively feeding on the Pulvinaria and only occasionally attacking *Pseudococcus aceris* Sign., just the opposite has been true the past season (1906). The insect has been found wherever *Pseudococcus aceris* occurred, but only in small numbers attacking the Pulvinaria. This is due in part perhaps to the fact that the latter insect has not been so abundant. Apparently the Pulvinaria is the favorite food of the coccinellid, but being unable to secure a sufficient amount of this, it attacks the *Pseudococcus* and other species.

Although the beetle and its larva did such effective work in Montclair in checking the scale, there were other places where it had not been so abundant, and consequently a much larger number of scales set and developed in those localities.

On July 24, when, at New Brunswick, I examined some leaves fairly well set with young Pulvinaria, I found that a few of them were parasitized and, except for their smaller size, appeared precisely like the parasitized forms of the hibernating females. Upon inspecting the infested trees in other localities I found that conditions were similar—a few examples parasitized in every case. An examination of the latter showed larvæ and pupæ similar to those of the parasite of the hibernating form. Within a day or two adults of the latter made their appearance both in the laboratory and out of

doors; and except for their smaller size, being not more than one-half as large, they resembled very closely *Coccophagus lecanii*, the parasite of the hibernating scales.

From this time on the number of these insects increased and continued to appear until about the middle of September, after which very little was seen of them. Leaves collected August 11 showed 1 parasitized scale to 10 good ones, and leaves observed in September showed a still greater proportion. Two of the leaves carefully examined showed in one case 289 and in the other case 136 parasitized insects, and many of the leaves were in a similar condition. At Montclair, where the coccinellid had almost exterminated the *Pulvinaria*, the majority of the few remaining scale sets were parasitized. These insects bred and increased rapidly, and in all respects, except size, resembled *Coccophagus lecanii*.

Specimens were sent to Doctor Ashmead for examination, however, and he determined them as *Coccophagus flavoscutellum* Ashm. However this may be, I am strongly of the opinion that we have a single species which is dimorphic, the larger form determined as *lecanii* Fitch, bred from the larger hibernating female scales, and the smaller form, not more than half as large as the *lecanii* form, which may be the species described as *flavoscutellum* Ashm., bred from the smaller scales, the size of the parasite depending on the size of the host.

SUMMARY.

The trees were badly infested with *Pulvinaria innumerabilis* Rathv. in the winter of 1904-5. As the scales developed in the spring it became evident that they were infested with the parasite *Coccophagus lecanii* Fitch, which in some instances destroyed over two-thirds of the scales and continued in evidence until about the middle of June. By this time the egg masses of *Pulvinaria* were becoming large and conspicuous, and an examination revealed the fact that they were infested by the larva of *Hyperaspis signata* Oliv., which in some places, conspicuously at Montclair, destroyed nearly all the offspring of the scale, the coccinellid larvæ feeding on the eggs and the adult beetles destroying the scale sets. The coccinellid continued until the 1st of August, by which time the scale larvæ had all-set and were becoming well developed. These young scales were parasitized like the hibernating females, and the parasites, which continued until the middle of September, were apparently a smaller form of the spring parasite. The result of these combined attacks has been that in some places the scale has been nearly exterminated and in all the infested localities its numbers are considerably reduced.

Two papers were then presented on codling-moth investigations, after which they were discussed.

REMARKS ON METHODS USED IN CODLING MOTH EXPERIMENTS.

By A. F. BURGESS, *Columbus, Ohio.*

In the great mass of data already published concerning the codling moth (*Carpocapsa pomonella* L.) it will be found that numerous standards have been used for determining the results of spraying. Many entomologists have followed the system of basing results simply on the number of picked apples which are wormy or sound on treated and check trees, while in other cases samples of one or more bushels of picked fruit, supposed to represent average conditions, have been counted and have served for making the determinations of the benefit derived. Sometimes an estimate of the wormy and sound dropped fruit has been made at the time of harvesting, but in the majority of cases no accurate account has been kept of the wormy fruit that dropped early in the season.

Undoubtedly this lack of data has been largely due to the fact that most entomologists who have attempted such experiments have not had at their disposal the necessary time to make complete counts of fruit. In some sections, also, horticulturists pay little attention to the fruit which drops early in the season, considering it a necessary evil when the crop is light and a direct benefit when the crop is heavy, as it saves labor in thinning. It would appear, however, if an accurate statement of the benefit derived by treatment is to be made, that all fruit growing on the trees should be accounted for.

In order to illustrate the case more fully, two tables are submitted, showing the percentages of benefit from spraying that may be deduced from the two different methods. Table I gives the treatment, in 1903, of two rows of 10 trees and 10 check trees in the orchard of Mr. R. L. Hudson, at Delaware, Ohio, with Disparene and Bordeaux mixture, also the number of wormy dropped apples, the wormy and sound picked fruit, and the percentages of sound fruit, using all the apples as a basis or by using only the picked fruit.

Table II gives similar data taken in the orchard of Mr. Oscar Haise, at Birmingham, Ohio, in 1905. The crop in the Hudson orchard was short, while that in the other orchard was good, but nearly twice the number of trees were used in the former orchard.

In Table I no sound fruit was recorded when the count of the wind-falls was made in August. So few sound apples were found that no record was made, but in later experiments the numbers were carefully noted, as is shown in Table II.

TABLE I.—Comparison of sound and wormy fruit from sprayed and unsprayed trees in Mr. R. L. Hudson's orchard, 1903.

Number of trees.	First spraying.	Second spraying.	Third spraying.	Windfalls.		Picked fruit.		All apples.		Picked apples.		Difference in percentage of improvement between computations made from all apples and those made only from picked apples.
				Aug. 4; wormy.	Sept. 23; wormy.	Wormy.	Sound.	Per cent sound.	Per cent improvement from spraying.	Per cent sound.	Per cent improvement from spraying.	
10	Disparene			2,428	1,359	1,172	376	7		24		
10	Disparene		Disparene	156	187	229	1,934	77	70	90	66	4
10	Disparene and Bordeaux mixture.	Disparene and Bordeaux mixture.	Disparene	468	554	864	4,417	70	63	84	60	3

TABLE II.—Comparison of sound and wormy fruit from sprayed and unsprayed trees in Mr. Oscar Haise's orchard, 1905.

Number of trees.	First spraying.	Second spraying.	Third spraying.	Windfalls.				Picked fruit.		All apples.		Picked apples.		Difference in percentage of improvement between computations made from all apples and those made only from picked apples.
				Aug. 1.		Oct. 13-21.		Wormy.	Sound.	Per cent sound.	Per cent improvement from spraying.	Per cent sound.	Per cent improvement from spraying.	
				Wormy.	Sound.	Wormy.	Sound.							
5	Disparene			995	357	558	39	1,342	247	18		16		
6	Disparene	Disparene	Disparene	137	884	45	516	103	4,488	96	78	98	82	4
6	Disparene and Bordeaux mixture.	Disparene and Bordeaux mixture.	Disparene	82	916	68	676	169	4,086	94	76	96	80	4

It will be noted in Table I that on the check rows, where all the apples were computed, only 7 per cent were sound, and when only the picked fruit was used the sound apples amounted to 24 per cent. This was due to the large amount of wormy fruit found on August 4 and September 23. The percentage of improvement is secured by subtracting the percentage of sound fruit on the check row, made by each method of computation, from that secured on the treated rows. This gives a balance of 3 and 4 per cent as a result of the different methods of computation, the larger percentage in the results being in favor of considering the entire crop.

In Table II, if the same methods are followed, a 4 per cent difference is shown, but this increase is in favor of the method where only the picked fruit is used. If it is assumed that the correct method is to have all of the fruit grown on the tree considered in deciding the percentage value of treatment, then the other method shows 4 per cent decrease in sound fruit in Table I and 4 per cent increase in Table II. It is therefore evident, as shown by these tests, that the error made by using only the picked fruit may range from none to 8 per cent in any experiment.

THE CONTROL OF THE CODLING MOTH IN THE ARID REGIONS.

By E. D. BALL, *Logan, Utah.*

The codling moth (*Carpocapsa pomonella* L.) is by far the most serious pest with which the apple grower in the arid regions has to contend. The unsprayed orchards will average, taking one year with another, fully one worm for each apple. In a year of abundant crops there will be some apples remaining sound, but the inevitable light crop following this is likely to have three or four worms to an apple.

That thorough and persistent spraying will control the codling moth has been demonstrated so many times as to appear at first sight almost axiomatic, yet when the writer took up the work in Utah, in the fall of 1902, he found a very deplorable state of affairs existing throughout the State. The codling moth had evidently been increasing in number and destructiveness for a series of years, and many of the leading fruit growers, who had formerly handled it with ease, were now meeting with very indifferent success or failing entirely to control it. Numerous instances were cited where from 4 to 6 or even more sprayings had failed to save the crop, while those who succeeded in getting 75 to 85 per cent of the picked fruit sound were considered highly successful. To add to the confusion, the cry had gone forth that the poisons were adulterated, that early sprays were of little value, and that three or even four or more broods were to be

expected in a season, so even those who were disposed to profit by their failures knew not which way to turn. Further investigation revealed the fact that a somewhat similar condition prevailed throughout the entire intermountain region.

In view of this condition of excessive worminess in apples and the general uncertainty that prevailed with reference to so many of the factors involved, it was decided to take up but one factor at a time and try to work that out by exact methods, so that whatever was discovered would be the result of known factors and could be explained by the known variation in the factors according to the needs of the experiment. In this way it has been possible to get quite definite results on several factors in the course of the four years' work, and the most important ones of these are summarized below.

The poisons were first investigated and were found to be first class in every respect. A few samples of Paris green had been found in the West that contained some free arsenic, but that would rather increase than decrease their killing power, so the failure of the spraying could not be laid at that door.

NUMBER OF BROODS.

In the work on this insect in Colorado, during the time the writer was an assistant at the State agricultural experiment station, the fact that there were two definite generations in that region and that these generations occurred at fairly definite times was thoroughly established. These tests were carefully repeated for northern Utah conditions, and a life-history chart showing the times of appearance of all stages of the two generations was published in Bulletin 87 of the Utah Agricultural Experiment Station.

From these investigations it was found that there would be a period of a few days in which it would be possible to separate the few worms of the two broods then occurring in the apples. This period, as was shown by reference to the chart above mentioned, occurred during the first few days in August in a normal season. As on the accurate separation of the damage done by each brood depended much of the value of the other data obtained, this damage was carefully checked up each year, and in no case was there more than 1 per cent of the "total wormy" in doubt, and often almost no doubtful cases appeared.

In practical work methods must come before results; but in this summary it has been thought best to give results first, and then the full significance of the methods that gave them can be more fully appreciated.

RELATIVE VALUE OF THE EARLY SPRAYINGS.

(Fig. 1.)

Careful tests of the relative value of each of three early sprayings were first made in 1905. The first spraying was applied just after the blossoms fell, the second ten days later, and the third fifteen days after that. Each spraying was tested alone and again in combination with the others and the results compared. The tests were carried on in five complete series, three on different blocks of Ben Davis and one each on Esopus and Missouri Pippin. All five tests gave similar results, and they are summarized in the following table:

TABLE III.—*Worms killed in the first brood by early sprayings.*

First spraying.	Second spraying.	Third spraying.	Total number of wormy apples.	Number of worms killed.	Percent-age killed.	Number of apples with calyx wormy.	Percent-age killed.	Number of apples with side wormy.	Percent-age killed.
.....			72	0	0	31	0	41	0
.....	1		15	0	0		0		0
.....	1		8	57	79	5	84	10	76
1			4	64	89	1	96	7	83
1	1		8	68	94		98	3	93
1	1	1	3	69	96		99	2	95

In the plat where the third spraying alone was given a few more wormy apples are shown than in the unsprayed block. This of course meant that the third spraying alone was of little value and that the trees happened to average a few more worms than the unsprayed. They were therefore treated as an unsprayed lot and averaged in with this lot to get the 72 wormy. This spray, when applied with the others, killed one more worm: and, strange as it may seem, every set showed this same result, so it should be given full credit for that worm.

The second spray, when applied alone, killed almost four-fifths of the worms, and when applied after the first spraying killed half of what would have been left. The first spray proved to be the best, killing almost nine-tenths of the worms when applied alone and when applied with the second killed 94 per cent.

The difference between the two sprayings was shown in every set but one, and there they were equal, while the increased value by applying both together was shown in every case.

WHERE THE WORMS WERE KILLED.

Examining these results to see where the worms were killed, we found that of the 15 worms left by the second spraying only 5 went into the calyx; of the 8 left after the first spraying, only 1 went in at that point; and when these two sprayings were combined an aver-

age of less than 1 calyx worm to a tree was obtained. In general, this shows that such spraying practically exterminates the worms that go in at the calyx and also shows the superior value of the first spraying in bringing this about.

FIG. 1.—Chart to illustrate relative value of early sprayings on the first brood of "worms" of the codling moth.

Of those going in on the side a slightly larger percentage escaped, and as would be expected there was also less difference in killing power of the two sprayings, and more gained by combining them. Though lower than in the calyx, the killing power here indicated is very high.

RESULTS OF SPRAYINGS IN THE SECOND BROOD.

(Fig. 2.)

TABLE IV.—*Worms killed in the second brood by early sprayings.*

First spraying.	Second spraying.	Third spraying.	Total number of wormy apples.	Number of worms killed.	Percentage killed.	Number of apples with calyx wormy.	Percentage killed.	Number of apples with side wormy.	Percentage killed.
.....			216	0	0	101	0	115	0
.....	1	1		0	0		0		0
.....	1		120	96	44	41	59	79	31
1			78	138	64	8	92	70	39
1	1		48	168	78	3	97	45	61
1	1	1	48	168	78	4	96	44	62

The results of spraying in the second brood are an almost exact duplication of those in the first, with the number of worms to be killed trebled and the killing power somewhat reduced, the reduction, however, being almost entirely in the killing of the worms infesting the fruit at the side.

It is hard to believe that two sprayings applied before June 10 could kill over three-fourths of the worms appearing in August and September, but the results are so uniform throughout each set that there can be no question about their general accuracy.

These results were obtained on trees averaging about 3,500 apples apiece; so the 4 worms escaping in the first brood together with the 48 in the second would make a total of almost exactly $1\frac{1}{2}$ per cent of wormy fruit for the year remaining after two sprayings.

VALUE OF TWO EARLY SPRAYINGS.

The question of the value of two early sprayings was taken up first in 1904 and the results published in Bulletin 95 of the Utah Station. The comparative value of the two early sprayings in 1905 has been shown in the previous table. Some of the plats, however, did not contain trees enough for a complete test and so were not included in the first set.

Omitting in the second-brood tables the sets in which the results were obscured by lack of apples for the worms, in the first year we get the following summary for all plats:

FIG. 2.—Chart to illustrate relative value of early sprayings on the second brood of "worms" of the codling moth.

TABLE V.—*Worms killed in first brood by two early sprayings, Smart & Hatch orchard.*

First spraying.	Second spraying.	1904.	1905.	1905.	
		Total wormy apples.	Total wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.
.....		247	97	65	32
1	1	27	6	1	5
Worms killed		220	91	64	27
Per cent killed ..		89	92	98	84

TABLE VI.—*Worms killed in second brood by two early sprayings, Smart & Hatch orchard.*

First spraying.	Second spraying.	1904.	1905.	1905.	
		Total wormy apples.	Total wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.
.....		712	273	177	96
1	1	236	64	6	58
Worms killed		476	209	171	48
Per cent killed ..		67	76	97	40

Tables V and VI show clearly the reduction in the number of worms the second year of spraying and also show a corresponding increase in the killing power of the early sprays. This increase is probably not real; instead it can be taken to mean that in the first test some apples on the unsprayed trees harbored more than one worm. All of the records indicate that the more worms there are, up to nearly 1 to each apple, the larger the percentage that will be killed.

VALUE OF THREE LATE SPRAYINGS.

Three late sprayings were applied to the second brood of worms in 1904 and again in 1905. They were applied with the same nozzle used in the early sprayings, but the nozzle was held farther away and spraying stopped as soon as the trees began to drip. The first of the late sprayings was applied as soon as the second-brood worms began to enter and the other two at 15-day intervals.

Three separate tests were made in 1904. The first was on 12 trees in an orchard that had no early spraying. The trees were about one-third wormy in the first brood, an indication that the apples would be entirely destroyed unless sprayed. Results, counting only second-brood injury, were as follows:

TABLE VII.—*Results of three late sprayings alone, in Hoggan orchard, 1904.*

	Wormy apples.	Sound apples.
6 unsprayed trees averaged.....	267	0
6 late sprayed trees averaged	254	8

There were no sound apples on either set at picking time. The 8 sound apples on the sprayed trees were windfalls. In fact there were very few apples left at picking time. The only difference that could be noted was that the apples from the unsprayed trees had 1 calyx worm each and 3 or 4 on the sides, while the sprayed ones were also all calyx wormy, but had an average of only about 1 on the side.

The second test was on unsprayed checks in the twice-sprayed Smart & Hatch orchards. These trees were nearly one-third wormy in the first brood, but about two-thirds of these worms were caught under bands, and the resulting moths from the remaining worms would, of course, spread out over the sprayed trees, so that there would be many less worms in the second brood than if the orchard had been unsprayed. On two varieties many of the apples on the unsprayed checks had more than one worm, and the results in sound apples are again too low to show the real killing power of this late spray. They gave averages as follows:

TABLE VIII.—*Results of three late sprayings alone in badly infested blocks in Smart & Hatch orchards, 1904.*

	Wormy apples.	Sound apples.
Unsprayed checks	562	252
Late sprayed checks	375	519

Many of the sound apples on the unsprayed trees were windfalls and most of the rest were below and inside. The outer and upper apples were badly infested.

On the third variety there were apples enough for the worms and to spare, and these results should show the full killing power of the sprayings when applied alone.

TABLE IX.—*Results of three late sprayings alone in moderately infested blocks in Smart & Hatch orchard, 1904.*

	Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.	Percentage of sound apples.
Unsprayed checks	712	487	225	51
Late sprayed checks	193	120	73	76
Worms killed	519	367	152	
Percentage killed	73	75	68	

That the sound apples in Tables VIII and IX must be credited to the fact that the early two sprayings reduced the number of worms in the orchard, is evident on comparison with the first table, where there were no early sprayings. There was nearly the same number of

worms in the first brood in each orchard, and without spraying the results would, no doubt, have been nearly the same.

That the late sprayings alone are not to be considered as a means of controlling the codling moth under badly infested conditions is abundantly shown by the above results. Therefore their true value to us is their killing power when applied in connection with the early sprayings, as shown below.

In the third test the three late sprayings were applied to trees that had received the two early sprayings. Three varieties were used and gave similar results, which are averaged in Table X.

TABLE X.—*Results of three late sprayings on trees that had received two early ones, Smart & Hatch orchard, 1904.*

	Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.
Two early sprayings.....	417	106	311
Two early and three late sprayings.....	112	35	77
Worms killed.....	305	71	234
Percentage killed.....	71	67	75

The number of worms to be killed is much smaller than before, and the proportion of "calyx wormy" to "side wormy" is reversed as a result of the high killing power of the early spraying on the calyx worms. The killing power of the late sprayings, however, was only reduced 2 per cent below that shown when they were applied alone.

This experiment was repeated in 1905 with a still smaller number of worms left after the two early sprayings. A still further decrease in killing power of only 1 per cent is shown below:

TABLE XI.—*Result of three late sprayings following two early ones, Smart & Hatch orchard, 1905.*

	Total number of wormy apples, second brood.	Number of apples with calyx wormy, second brood.	Number of apples with side wormy, second brood.
Two early sprayings.....	64	6	58
Two early and three late sprayings.....	19	3	16
Worms killed.....	45	3	42
Percentage killed.....	70	50	72

In this case the number of "calyx wormy" was so small that the results would have had to be expressed in fractions to have been accurate. There were six sets in the experiment, and it is worthy of note that on the two wormiest ones the highest killing power was shown.

Examining to see where the worms were killed we found that there was little difference in the percentage killed in the calyx and side. Wherever the largest number of worms were, there the percentage killed was the highest.

THE RELATIVE VALUE OF EARLY AND LATE SPRAYINGS.

(Fig. 3.)

The entire value of the late sprayings can be measured by the number of worms killed in the second brood, and is easily obtained from the above tables. In the case of the early sprayings the problem is much more complex. A certain number are killed in the first brood,

FIG. 3.—Chart to illustrate relative value of early and late sprayings for the codling moth.

still more in the second, but by killing the worms in the first brood a larger number are prevented from appearing in the second. This latter factor is no doubt variable, and up to the present time no method has been devised by which the ratio of increase can be accurately determined. Estimating this increase at five times is doubtless

conservative, as all other authors have placed it higher; and by using that number in connection with the results found before we get the following summary:

TABLE XII.—*Relative value of early and late sprayings in 1904.*

Value of early sprayings:	
Worms killed in first brood.....	220
Worms killed in second brood.....	476
Worms killed in first brood would have increased to.....	1, 100
Total value of early sprayings.....	1, 796
Value of late sprayings:	
Worms killed when applied alone.....	579
Worms killed when following early ones.....	305
Total value of late sprayings either 579 or 305.	

This comparison shows that the early sprayings have a protecting power over three times as great as that of the late ones when applied alone and under the most favorable conditions, and almost six times as great as the late sprayings when following the early ones. Still it does not bring out the full value of the early sprayings in comparison with the late ones, as the results in the following year will show.

TABLE XIII.—*Relative value of early and late sprayings in 1905.*

Value of early sprayings:	
Worms killed in first brood.....	91
Worms killed in second brood.....	209
Worms prevented in second brood.....	455
Total value of early sprayings.....	755
Value of late sprayings:	
Worms killed when following early ones.....	45
Total value of late sprayings.....	45

Here we see that the effect of the early sprayings in 1904 was cumulative and resulted in a marked decrease in the number of worms appearing the following year. As a result, the early sprayings destroyed so large a percentage of the worms that there was a striking decrease in the number of worms killed by the late sprayings and a corresponding decrease in the comparative value of the late sprayings as compared with the early ones.

Comparing the results of two years we find that the late sprayings which followed the early ones killed one worm to the early ones' six the first year and only one to the early ones' 16 the second year. By the third year the worms were so reduced in numbers that no late sprayings were applied.

RESULTS IN OTHER YEARS.

No other tests of the late sprayings than these enumerated have been made. In the other tests the number of worms left after the early sprayings has not been sufficient to give accurate results, and none has been applied.

The first test of the relative value of the early sprayings was made on a single variety in 1904. The results in the first brood were good, but in the second brood most of the apples on the unsprayed trees and some of those on the once-sprayed trees fed more than one worm, so that the number of apples saved did not show the real killing power of the sprayings.

TABLE XIV.—*Worms killed in the first brood by early sprayings, Smart & Hatch orchard, 1904.*

First spraying.	Second spraying.	Total number of wormy apples.	Percentage killed.	Number of apples with calyx wormy.	Percentage killed.	Number of apples with side wormy.	Percentage killed.
0	0	439	0	289	0	150	0
1	0	192	66	110	62	82	45
1	1	37	92	10	97	27	82

This same orchard was used for the elaborate test of 1905, the result of which has been given. In 1906, although only about one tree in seven was bearing at all and those only about half a crop, three varieties were used in a third test. On account of the small number of trees bearing, no unsprayed checks were left and results can only be compared with those of former years.

Two other tests were made this year (1906) to see if the same results could be obtained in the earlier and warmer sections of the State. One test was made on three blocks of Ben Davis at Lehi, and the other one on two blocks of the same variety at Provo.

In the case of the Lehi orchard the trees were young and bearing only from 300 to 350 apples apiece. The unsprayed trees were nearly one-third wormy in the first brood, showing that the orchard would have been entirely wormy if unsprayed; consequently there were worms enough in both broods to give fairly accurate results. These results are averaged in Tables XV and XVI.

TABLE XV.—*Worms killed in the first brood by early sprayings, Lehi orchard, 1906.*

First spraying.	Second spraying.	Total number of wormy apples.	Percentage killed.	Number of apples with calyx wormy.	Percentage killed.	Number of apples with side wormy.	Percentage killed.
0	0	45	0	17	0	28	0
0	1	11	76	1	94	10	64
1	0	+ 4	90	+ 0	99	+ 4	84
1	1	8	82	+ 0	99	8	72

TABLE XVI.—Worms killed in the second brood by early sprayings, *Lehi orchard, 1906.*

First spraying.	Second spraying.	Total number of wormy apples.	Percentage killed.	Number of apples with calyx wormy.	Percentage killed.	Number of apples with side wormy.	Percentage killed.
0	0	75	0	28	0	47	0
0	1	31	59	2	93	29	38
1	0	21	72	1	97	20	57
1	1	20	73	+ 0	99	20	57

These results show an even higher killing power in the calyx than in the former tests, due, no doubt, to the fact that the small trees were easier to spray thoroughly and that more liquid was applied in proportion to the apples.

In the Smart & Hatch orchard in 1906 and in the Provo tests the number of worms present to begin with was much smaller, and those left after the early sprayings were so few in numbers that in some cases the individual variation in the trees was greater than the difference between the sprayings. The actual percentages are therefore of little value. This is readily understood if one considers that with so few moths in the orchards some trees might not be visited at all.

TABLE XVII.—Worms killed by early sprayings in *Smart & Hatch orchard, 1906.*

First spraying.	Second spraying.	First brood.			Second brood.		
		Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.	Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.
0	0	(?)	(?)	(?)	(?)	(?)	(?)
0	1	7	2	5	31	0	31
1	0	23	6	17	74	25	49
1	1	5	1	4	42	6	36

TABLE XVIII.—Worms killed by early sprayings in *Provo orchard, 1906.*

First spraying.	Second spraying.	First brood.			Second brood.		
		Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.	Total number of wormy apples.	Number of apples with calyx wormy.	Number of apples with side wormy.
0	0	17	9	8	21	5	16
0	1	8	2	6	8	1	7
1	0	5	1	4	8	1	7
1	1	4	0	4	11	1	10

The general average of Tables XVII and XVIII, with one exception, will be seen to be the same as that of the others. The actual percentage killed is less than where there were more worms; but as the Smart & Hatch orchards were only 3 per cent wormy for the season and the Provo orchard but a trifle over 1 per cent wormy, the results are entirely satisfactory.

The one exception to the general average is the relative value of the first and second sprayings as shown in the Smart orchard. In all other tests the first spraying has proved to be the best, and in all these cases the two sprayings have been applied alike. In this one case the writer, who has done most of the spraying himself, was away superintending the work in other parts of the State leaving the work on this orchard in charge of an assistant. On returning, the amount of poison found in the calyx cups was so small that an investigation was made, and it was found that the assistant had turned the work over to inexperienced students to carry out, with the inevitable results. In order to make the two sprayings "average up" with those of other years the second spray was applied with extra care and in larger amounts than usual, hence the reversal of the ordinary results.

AMOUNT OF SPRAY TO APPLY.

The above results, although not intended as an experiment, show that the relative value of a spray can be easily altered by a change in the method of applying it or in the amount applied.

In order to test the question as to whether it was possible to spray too much, two trees were sprayed in the ordinary way and two were soaked until the water stood on the ground in puddles, at least four times the ordinary amount of spray being applied, with the following results:

	Wormy apples, first brood.	Wormy apples, second brood.
Ordinary sprayed trees averaged.....	9	147
Soaked trees averaged.....	8	96

This shows that there is no danger of overspraying in the early sprays and indicates that even our liberal application might be increased with profit.

COMMERCIAL RESULTS.

All tests have been carried on in orchards of from 2 to 5 acres, and the results given for the twice-sprayed trees have represented the condition of the orchards as a whole. Besides this, more and more of the fruit raisers have adopted this method each year, until in 1906 the majority of the orchards of the State, as well as a large number in Idaho and Oregon, were sprayed by this method with the most satisfactory results ever obtained. In September the writer visited commercial orchards in which a calyx-wormy apple was such a rare thing that it was only after careful and prolonged search that one could be found.

METHODS USED.

In each experiment the trees selected for the different tests have been of the same variety, the same size, and bearing as nearly the same number of apples as possible. In every test the orchard has been sprayed twice and the spraying done before the trees were selected, except for those trees on which the given spraying was omitted, so that the spraying represents the average condition of a commercial orchard. Check trees have been scattered through the blocks so that one check would not influence another. All important tests have been made in parallel series on at least three winter varieties, and all have been or will be repeated through a series of years, and in the warm and cold sections of the State. Unsprayed trees have been left very sparingly, as one badly infested tree will noticeably influence half an acre. The check trees have been banded, the bands examined every three days, and the worms placed in cages and bred for the life history work. The remaining trees of the orchard have not been banded, and the worms caught on the check trees do not more than offset the larger number escaping from the unsprayed checks, thus leaving the orchard in about the same condition as if there were no check trees present.

All apples that set on the check trees were accounted for. Windfalls were picked up every three days and sorted into "sound," "calyx wormy" and "side wormy." All that were the least questionable were sliced open to determine whether the calyx had been entered. At the proper time to divide broods all the apples on the trees were examined and all wormy ones of the first brood recorded. These, with the windfalls to date, made up the damage from the first brood of worms.

Complete records were kept on 28 trees in 1904, 64 trees in 1905, and 50 trees in 1906. The 64 trees used in the comparison of early sprayings in 1905 carried 220,000 apples, nearly all of which were handled three times during the investigation.

COMPARISON OF RESULTS.

In all the experiments the results have been given in the actual number of wormy apples found in each brood under each different test, and the percentage given shows the actual killing power of the spray for that brood. In one case, for example, the unsprayed trees averaged in the first brood 72 wormy apples, while the twice-sprayed ones averaged only 4 wormy, showing that 68 worms, or 94 per cent of those that appeared, were killed. There can be no question of the accuracy of this method for the first brood, because in this brood the moths will be evenly distributed throughout the block without reference to sprayed or unsprayed trees. In the second brood the problem

of getting a fair standard of comparison was a little more difficult; but by banding the unsprayed trees and catching two-thirds or over of the first-brood worms it was thought that the few remaining worms in excess of the number on a sprayed tree would be so evenly scattered among the surrounding trees, especially where the unsprayed trees were well distributed through the block, that they would fairly represent the average distribution of the worms in the orchard. The results in the second brood have in every case agreed so remarkably with those in the first that there seems to be no question but that this method was fairly accurate.

All the tests have shown that the variation in the number of worms attacking trees of equal size is very slight, as compared with the variation in the number of apples that they may bear; therefore the method of comparing the actual number of wormy apples is much more accurate than the method of comparing percentage of sound and wormy fruit.

In these tests none of the wormy apples from the first brood has been found on the trees at picking time, and the percentage of those of the second brood remaining has varied so greatly, under different conditions and on different varieties, that results, based on percentage of wormy fruit at picking time, appear to be of doubtful value at best, and where they have included different varieties in the same test, of no value at all.

WHAT CONSTITUTES THOROUGH EARLY SPRAYING.

If the first spray is applied for the purpose of placing poison in the calyx cups—and all will agree to this, I think—then the test of the efficiency of such a spray will be the amount of poison found in the cup after it has closed. Now, paradoxical as it may sound, it is not as easy to spray into an open calyx as it is into one that has nearly closed. This will be made clear by referring to figure 4, subfigure 1, where a cross section of an open cup is shown with the pistil in the center and the circle of stamens forming a tight roof above, as seen at 1*a*. A week or ten days later the calyx lobes have nearly closed above, the opening being filled with the stamens and pistils, as seen in subfigure 2. By this time the stamen “bars,” as I have called them, have shrunk and become more or less twisted, so that there are many open spaces between them, as seen in 2*a*. Later the calyx lobes close tightly and the “bars” shrivel still more, as seen in subfigures 3 and 3*a*.

By reference to subfigures 2 and 3, it will be seen that there are two distinct cavities at first, one above and one below the stamens, either of which might be called the calyx cup, and which have been treated as a single cavity by most writers.

A large number of spraying tests have been made, and by using Paris green without lime the location of each grain of poison could be readily made out with a strong hand lens.

A fine mist spray applied only to the point of dripping was first tried. This left grains of poison well distributed over the inside of the calyx lobes and on the upper surface of the stamens when applied to the open calyx, as shown in subfigure 1. When this spray was continued until the drops ran together, the greater amount of the poison was deposited in a ring around the base of the stamens. When applied to the closing calyx of subfigure 2, it simply deposited a few grains in the outer part of the throat and on the tops of the stamens and pistils. In the hundreds of cases examined there were but few in which there was any poison to be seen in the lower cup and then only

FIG. 4.—The condition of the calyx cup of the apple in relation to spraying for the codling moth: *Fig. 1*—A calyx cup, five days after the petals fell, split open to show two cavities; *1a*, the roof of stamens as seen from above. *Fig. 2*—A calyx cup two weeks after blossoming, showing the calyx lobes above; *2a*, the stamens from above, to show spaces. *Fig. 3*—The relation of the two cavities in a nearly grown apple; *3a*, stamens from above.

in small amounts. The fine, mist-like drops did not seem to be able to force their way through the fleshy stamens in the open calyx nor through the hairy mass in the throat of the closing calyx. Simpson states that in his experiments he "was unable at any time to distinguish any particle of spray inside the tube," and he lays especial stress on the fine, mist-like spray. Slingerland states that the spray

lodges in the saucer-like calyx, but does not distinguish between the two cavities. He was able to find it two weeks later by chemical analysis.

A spray thrown up into the air and allowed to fall in large drops deposited poison much like the oversprayed mist, but with much less regularity.

A driving spray of fine drops thrown with sufficient force to go 6 or 8 feet before breaking into a mist was next tested. This was applied with the nozzle held so that it would throw the spray straight into each cup from a distance of from 1 to 3 feet, and was continued until each fruit cluster was freely dripping. When applied to the open calyx this spray left a rim of poison around the base of the stamens, scattered masses of it here and there in the central mass of stamens and pistils, and forced some down through the roof of stamens into the cup below; part of this, no doubt, passed through the minute openings and part was probably forced down through the center.

When applied to a partly closed calyx, the finer drops were driven straight down through the shrunken stamens' "bars" and gathered in the lower cups where the poison was deposited. Series of countings gave an average of about 10 grains of poison visible in the lower cups following this spray.

As will be explained later, this method of spraying was found to place the poison where the most worms entered, and was adopted for all early sprays, with 15 grains of poison in the lower cup as a standard for two thorough sprayings. Card notes that coarse spraying placed more poison in the calyx cups, but evidently referred to the outer one, as he continues that it was easier to fill them when wide open, though more remained when partly closed.

METHOD OF APPLYING A DRIVING SPRAY.

In order to apply such a spray in a uniform and thorough manner, it was found necessary for the operator to be on a level with the upper limbs of the trees. From such a position, by using a 10-foot extension pole with the nozzle set at an angle of 30° to the pole, one could spray from above down and from the sides in; and by spraying from all four quarters of the tree one could force the poison straight into each blossom at close range. Spraying was directed entirely at the blossoms, no attention being paid to anything else, and no spraying done where there were no blossoms. Each cluster of blossoms was gone over two or three times and left freely dripping. In practice it was found that this required about 1 gallon of liquid at each spraying for every 3 bushels of fruit expected, and cost three-fourths of a cent a bushel.

The spray was thrown through a Bordeaux nozzle set at almost its widest limit on the broadest side, and under a pressure of from 85 to 120 pounds. Paris green was used, without lime, at the rate of 1 pound to 120 gallons of water in all tests. (This could only be done in an arid climate.) Duplicate series were carried through with 6 pounds of lead arsenate to 120 gallons of water in three cases without showing any difference in results.

Heavy rains followed the sprayings in several cases, without lowering their killing value enough to be noticeable. This was no doubt due to the fact that the effective poison was placed in protected places.

HOW THE EARLY WORMS ARE KILLED.

In these experiments very few worms of the first brood have been observed to go in on the free surfaces of the apples, and those mostly the later ones. Over two-thirds of this brood, on an average, went in at the calyx and most of the rest were found under leaves where apples touched, or were going in at some scar. Of those going in at the latter situations, no definite observations have been made as to how they are killed. That three-fourths or more of them are killed by the two sprayings has been repeatedly demonstrated. Examinations have shown that where two apples touch, a driving spray will often leave a film of poison around the spot, due no doubt to capillary attraction holding the liquid there until it dries. The same thing was true, but to a less extent, with touching leaves and scars. In fact, more poison was found in such situations from the driving spray than from a mist spray applied so as to give the maximum amount of surface coating. This may well account for a considerable number of those killed, but the entire subject is one for further investigation.

Of the fate of those going in at the calyx we are more fortunate in having definite information. In cutting the thousands of apples examined for poison a number of small worms were found in the lower cups; most of these were found crawling around, but in several instances dead ones were found where there was no sign of their having started to penetrate. More cases were found where a hole had been eaten a short distance into the pulp before the worm had died. In every instance where a dead worm was found poison was found in the lower cup.

Later an orchard was found that had borne a large and very wormy crop the previous year, and was now bearing a very light one. It had received one driving spray, and was being examined to find out whether late ones would be necessary just as the greatest number of the first brood worms were appearing. Some of the apples already showed castings in the blossom end; of the rest, an average of about 1 in 3 or 4 contained a live worm crawling around in the calyx, and

almost all of these were in the lower cups. The spraying had not been well done, as less than half of the apples contained enough poison to be seen; but of those that did about one-third contained dead worms. Two "sick" worms and a number of fresh dead ones were found, but the great majority of the dead ones were only recognized by their shining black heads, and from that possibly a trace of a decayed body could be made out. In no case were two live worms found in the same cup, though in one case a live worm was found in both inner and outer cups. Live worms and dead ones were frequently found in the same cup; in many cases two, and in a few even more dead worms were found together. The climax was reached when three dead heads and a live worm were found in a single lower cup.

In only one case was a dead worm found above the stamens, and this one was on the shelf at their base. From the lack of dead worms above the stamens it was concluded that most of the live ones found on the outside were new arrivals, and had not yet made their way below. An examination of thousands of wormy apples with this point in mind showed that in 97 or 98 per cent of those classed as "calyx wormy" the entrance had been made from the lower part of the lower cup, from which it would seem that the method of spraying that would place the most poison in this region would be the best.

KILLING THE SECOND BROOD.

Many factors influence the number of worms entering the calyx in the second brood. Some always go in on the free surface of the apples, more go in where apples touch, but in our experiments, taking all the varieties through all the years, almost two-thirds of them have been found to go into the calyx. In slow-growing and stunted apples the calyx is often so tightly closed that it is hard for a worm to find an entrance, while in the fast-growing apples and the larger ones of all varieties the calyx often opens up again about the time the worms are appearing.

Only a little of the poison of the first sprayings can be found in even the most protected locations, and of course none would remain on the free surfaces of the apples even if it had been left there (and this sort of spraying leaves very little in the first place), so it was rather hard to explain how the worms were killed on the sides, yet every record has shown that some were killed. This is a subject that needs careful investigation.

In the case of the calyx worms, however, abundant evidence was at hand to show how they were killed. The poison placed in the calyx in June could be found in only slightly diminished quantities in August and September, and the record of 97 per cent of calyx worms

killed was no more than was to be expected. Apples from trees twice sprayed in June, 1905, were exhibited in February and March, 1906, at the Northwestern Fruit Growers' Convention and the Idaho Horticultural Society, and in the majority of them poison could still be detected.

CONCLUSIONS.

(1) The poisons used in the intermountain region have been satisfactory.

(2) The codling moth is definitely two-brooded in this region.

(3) It is so abundant that from 5 to 8 ordinary sprayings have been necessary to control it.

(4) The driving sprays force the poison into the calyx cups and into the cracks and spaces between the apples where the first-brood worms enter.

(5) The first early spraying is the best, the second is nearly as good, and the third is of little value.

(6) Two early driving sprayings will kill an average of 90 per cent of the first brood of worms.

(7) By killing these worms in the first brood the greater part of the second brood is prevented from appearing.

(8) Sufficient poison is retained from the early sprayings to kill an average of 74 per cent of the second brood of worms.

(9) Two early sprayings correctly applied are worth from 6 to 16 times as much as three late ones.

(10) Over two-thirds of the first brood and nearly two-thirds of the second brood of worms enter the calyx.

(11) Of these worms entering the calyx, an average of 98 per cent of the first brood and 97 per cent of the second were killed by the two early sprayings.

(12) Of the worms entering the sides, an average of 78 per cent of the first brood and 52 per cent of the second were killed by the same sprayings.

(13) Each succeeding year of such spraying will reduce the number of worms remaining in the orchard.

Mr. J. B. Smith inquired if there was any special explanation why an early and a late spraying of Disparene showed a larger percentage of perfect fruit than when three sprayings were applied and Bordeaux mixture was added to the first two; also, whether Bordeaux mixture seemed to affect the efficiency of the poison.

In reply, Mr. Burgess stated that he could give no reason for the better showing made where only two sprayings were applied. In

some cases Bordeaux mixture seemed to render the poison less effective, but in other experiments the opposite proved true. It did not seem advisable to leave out Bordeaux mixture, as in some regions its use was absolutely necessary for controlling apple scab and other fungous diseases.

Mr. Britton believed that in combining Bordeaux mixture with arsenate of lead the poison became less soluble, and this resulted in its killing fewer insects.

Mr. Sanderson, in speaking of the results secured by Mr. Ball, emphasized the desirability of having uniform methods of treatment, in order to secure experimental data which could be used for comparison.

Mr. Ball stated that in order to secure the results reported in his paper it was absolutely necessary to use a coarse driving spray, and that a mist spray, such as is usually recommended, would not drive the poison far enough into the calyx cups to be effective. He did not attempt to cover the fruit or foliage with the spray, but simply wished to fill the calyx with poison.

Mr. Fletcher pointed out the desirability of not casting any doubt on the efficiency of the methods now generally in vogue for controlling this insect. In Canada 70 per cent of the apple crop is saved by the present acknowledgedly imperfect spraying methods. He did not believe it necessary to lay so much stress on filling the calyx, and was decidedly in favor of delivering the spray in as mistlike condition as possible. With the pumps and nozzles in general use by farmers and fruit growers there was far too great a danger of going to the other extreme. Excellent paying results were now being secured by ordinary farmers with the mist spray which had been used for several years.

Mr. Quaintance pointed out that fruit-growing conditions in the Mississippi Valley and Eastern States were quite different from those in many sections of the West, as in Utah. The absence of rains there during the growing season largely obviated the necessity for the use of fungicides to control scab and other fungous diseases. While he did not doubt that it was entirely practicable to use a coarse spray for the codling moth in Utah and thoroughly drench the trees, this would be bad practice according to present ideas of spraying in the East, where a mist-like spray is desired to treat uniformly all parts of the foliage and fruit. Under present conditions of spraying, young apples are often russeted by the Bordeaux and arsenical treatment, especially by the one just after the petals have fallen, and a thorough drenching of the trees at this time, as advocated by Mr. Ball, would be likely to prove quite harmful in this way. He also called attention to excellent results which orchardists had been securing in the control of the codling moth in the East by the use of the

mist spray, and that success depended more on spraying with good pressure and making the applications thorough than on anything else.

Mr. Ball remarked that this system of spraying had been tried in not less than forty orchards by commercial growers, and they were able to save at least 96 per cent of their crop; other growers who used the mist spray and made five applications rarely secured more than 90 per cent of sound fruit. It was absolutely necessary in Utah to produce perfect fruit, as this was the only kind that could be profitably marketed on account of the long distance which it must be shipped.

Mr. Saunders asked concerning the date of applying the second spraying, and Mr. Ball replied that this must be applied within ten days or, at the longest, two weeks after the first spraying, but the exact time depended on the condition of the calyx lobes. It must be put on before the lobes have closed.

Mr. Hopkins stated that his phenological investigations in West Virginia and other sections of the country showed conclusively that the dates of periodical phenomena of plants and insects must be determined for different localities, latitudes, and altitudes. For instance, the dates at which the apple leaf and flower buds open not only differ greatly in different localities and seasons, but in different varieties of apples in the same locality. The rate of difference from a given locality may not be far from four days for each 400 feet of altitude and four days for each degree of latitude—later at localities north or higher, and earlier at those south or lower.

The following paper was presented:

WHAT RESEARCH IN ECONOMIC ENTOMOLOGY IS LEGITIMATE UNDER THE ADAMS ACT?

By E. DWIGHT SANDERSON, *Durham, N. H.*

The recent passage by Congress of "An act to provide for an increased annual appropriation for the agricultural experiment stations, and regulating the expenditure thereof," commonly termed the "Adams Act," in honor of its lamented author, the late Hon. H. C. Adams, of Wisconsin, should be made a notable milestone in the progress of economic entomology. This act provides that it shall "be applied only to paying the necessary expenses of conducting original researches or experiments bearing on the agricultural industry of the United States, having due regard to the varying conditions and needs of the respective States or Territories." The control of its proper expenditure has been placed in the hands of the Secretary of Agriculture, who has delegated his supervision to the Office of Experiment Stations of the Department of Agriculture. In endeavoring to secure the expenditure of the appropriation for original

research within the intent of the law, the Director of the Office of Experiment Stations has consulted with the various station directors and workers, and at the recent meeting of the Association of Agricultural Colleges and Experiment Stations the discussion of this question was one of the leading features. The main question is, "What is 'original research * * * bearing directly on the agricultural industry?'" In general, the principles which should govern the problems for such research seem to have been satisfactorily defined for all parties concerned, but in outlining his own work under this act the writer has felt that some discussion as to just what lines of work are most needed or most desirable in economic entomology might not be amiss at the present time.

The term "economic entomology" seems to be one of those which has come into common usage but which has not been carefully defined. In a somewhat careful search of the dictionaries and the proceedings of this association no definition of the term has been found. Might we not describe it as the science of entomology as applied or related to human welfare? It is therefore narrower in its scope than the general ethology of insect life, though it usually includes the ethology of all forms of economic importance. For a general discussion of the objects and field of the economic entomologist I have seen nothing better than the remarks of Dr. S. A. Forbes, quoted by Dr. J. W. Folsom in his recent *Entomology*.^a After reviewing with him the diverse sciences and arts with which the economic entomologist must deal, we might fairly question whether economic entomology is truly a science of itself or whether as economic entomologists, as we style ourselves, we are not really artisans applying the laws and knowledge gained from the true sciences. In any event, the science is a young one and a little definition of its objects and scope will not be out of place.

Economic entomology would therefore include the subjects of the relation of insects to human health, household insects, etc., while the Adams Act restricts us to work upon subjects related to agriculture. The economic study of mosquitoes is therefore not permissible under its terms. In general, our work will doubtless be restricted to work upon insects injurious to agriculture, for, although apiculture and sericulture are undoubtedly worthy objects for research by the economic entomologist, yet they have become so specialized that but few of us pretend having any practical knowledge of them such as would qualify for research.

Many of us have doubtless been surprised to find that, though constantly dealing with the term, our ideas of the true meaning of

^a *Entomology*, with special reference to its biological and economic aspects. By Justus Watson Folsom. Philadelphia, 1906.

research are possibly somewhat indefinite. The dictionaries define research as "continued and diligent investigation; studious and laborious inquiry; systematic scientific investigation." (Standard Dictionary.) Hitherto a large part of our work as station entomologists has been in the nature of miscellaneous observations upon various insects demanding our attention, often involving considerable study; but how often has our work upon them reached the position where it might properly be called research in the true sense? It is obvious that for the best development of our science research is greatly needed at the present time, and we therefore welcome this increased appropriation with its restrictions to such use.

In considering the branches of economic entomology which will furnish a field for such "diligent investigation" we may study individual insects, groups of insects, or general principles governing insect life and control.

In the past our work has necessarily been largely a study of individuals, more especially as regards their life histories. In this work marked advance has been made in the past few years. We have come to see that if an insect is of sufficient economic importance to warrant investigation, the study must cover all phases of the insect's ethology and the testing of all possible means of control. For example, consider the work of Hunter and Hinds on the boll weevil, Quaintance's investigation of the bollworm, Doctor Forbes's studies of the corn root-aphis, Professor Crandall's work on the life history of the plum curculio, and some of the investigations of the codling moth. These studies must always remain classic in the annals of economic entomology from the fact that they have raised a new standard for work upon such insects of primary economic importance. The study of an individual insect, therefore, furnishes a worthy object for research, but the work must be exhaustive. A mere breeding of a dozen specimens in an insectary, testing a remedy which seems to be satisfactory, and the publishing of a bulletin outlining the supposed facts, is not research. It may be that such work is all that the economic importance of an insect pest warrants in many instances, but it is hardly research in the true sense. Whether the insect warrants true research work upon it will depend largely upon its economic importance. Thus the rusty tussock moth (*Notolophus antiqua* L.) is hardly of sufficient importance to warrant any lengthy study, while its brother, the white-marked tussock moth (*Hemerocampa leucostigma* S. & A.), might well receive the most careful research. In some instances, however, an insect may be peculiarly adapted to some research work in an effort to establish certain principles of insect life; but in such a case it is merely a means to an end and is not in itself the object of study. The larger part of our life-history studies and observations can therefore hardly come

within the meaning of this act, though necessary and forming the foundation for the future development of the science.

In the study of groups of insects we deal with what is really a branch of systematic entomology, and which might properly be styled systematic economic entomology. Under this head comes the study of (1) taxonomic groups and (2) economic groups. If in his work the economic entomologist finds that he can not proceed further without a monographic revision of the family or genus with which he is most concerned, and if he is unable to secure the help of a satisfactory specialist for such work, why should he not himself undertake its revision, if he feels competent and able to do so? For instance, is not Doctor Hopkins's work on the Scolytidæ of fundamental economic importance? Would not careful systematic studies of the Aphididæ be of the greatest service to the economic entomologist, and can we ever be sure of just what we are dealing with in this most interesting and much neglected family until we do have such systematic work? Again, the economic entomologist constantly receives numerous caterpillars and larvæ of all orders, often of economic importance, which it is impossible for him to identify in the immature stages and which he must therefore rear before he can suggest remedial measures. These efforts at rearing are often unsuccessful, in which cases much time and labor is spent for naught. Systematic studies of insect larvæ, with monographic work enabling us to identify them, would be one of the greatest boons to the economic worker and save much really unnecessary work. Is not the systematic study of insect larvæ, therefore, a proper field for research by the economic entomologist, as related to agriculture?

Again, we may investigate economic groups as regards their economy, considering either taxonomic groups which are of general economic importance, such as the barkbeetles, or groups of insects affecting some one food plant, as corn insects, cotton insects, etc. When such studies cover the whole field in such a way that general methods of farm practice or procedure are worked out which will control all the common insect enemies of any important crop, are they not worthy objects of true research? Doctor Forbes's work on corn insects might furnish an example. It should be remembered, however, that mere general observations upon any class of insects and the compilation of a report upon them is not research; to come under that term the work must be clearly defined, systematic, and exhaustively studied. Probably much of such work upon groups of insects may well be deferred until more careful studies of many important individual pests have been made, though it will furnish an inviting field in the not far-distant future.

It is the study of the principles underlying insect life and control which, it seems to the writer, furnishes a field for research but little

tilled and with much virgin soil. In his presidential address at the fifth meeting of this association, Doctor Forbes noted a satisfaction with the present methods and a dearth of new methods of work. Has not this tendency continued to some extent, and do we not need more pioneer work along the lines of pure science which will furnish a basis for the future development of economic entomology and insect control? Much has been done, which will be noted below; but there is certainly vastly more to do if our science is to make any marked advance in the future, comparable with the advances in other branches of science, or if advance in the control of agricultural pests is to equal that now occurring in the control of insects affecting health.

Such new principles may be sought and furnish objects for research in various phases of economic entomology.

(1) Our studies of life histories are principally of value in revealing the vital time at which an insect may be attacked, and often the method of control, but the time factor is the dominant one. Would not exhaustive studies of the relation of temperature, moisture, latitude, altitude, etc., in short, climatic influences upon the various stages and processes of the transformations of insects, be of the greatest value in revealing the time at which certain pests might best be fought in certain localities, and the number of generations and consequent destructiveness of a pest in different latitudes? Something along this line was done by Dr. A. D. Hopkins and Prof. F. M. Webster in their studies of the Hessian fly in West Virginia and Ohio, and recently by the staff of the Bureau of Entomology in their work upon the boll weevil and bollworm. The importance of such work seems to be increasingly appreciated.

(2) The structure and physiology of an insect, including its habits, determine to a large extent the method of control. We know that a chewing insect can usually be poisoned with arsenicals and that a sucking bug must be handled otherwise, but have we carried our studies of anatomy and physiology far enough? Might we not learn more of vital importance? In his presidential address at the eighth meeting, Prof. C. H. Fernald called attention to the need of our securing the cooperation of the chemist and physiologist, and the work of the gipsy-moth committee furnished some examples of the wisdom of this advice. Of even simple features of anatomy we are often ignorant. For example, the writer recently had difficulty in ascertaining whether the common imbricated snout-beetle (*Epicurus imbricatus* Say) was winged or not, and the matter was only decided by the kindness of Mr. Schwarz, who examined specimens in the National Museum, proving the species to be apterous. A very similar beetle affecting peach foliage in Texas was controlled by us by banding the

trunks of trees with insect lime and thus preventing its ascent. It is obvious that the imbricated snout-beetle can probably be handled in the same way; but this fact of its anatomy seems to have entirely escaped observation, nor was the writer able to find reference to the wings in any monograph. Again, experiments recently conducted at Cornell University have shown that a liquid lice killer applied to the perches will kill the lice upon fowls and is exceedingly efficient. How is this done? Possibly Professor Slingerland can inform us, but it is exceedingly suggestive in showing that some things are possible at which most of us would have scoffed had we not the evidence of competent observers. Might we not find that through study of the nervous system of insects and experiments with substances or means of affecting the nervous apparatus, a means of control for some pests might be found? Would not a study of the alimentary tract and its physiology be of value? These are but suggestions. Who can say what might come from such research?

(3) As regards the artificial control of insects with insecticides and apparatus, we have to deal with a different problem. Is such work really that of a scientist or that of an artisan? Certainly the mere testing of one brand of an arsenical against another is not a matter of science. Doctor Forbes has aptly said that such work "really stands for the most part fairly across the boundary line of horticulture and agriculture." We need further work upon the origin and perfection of insecticides, and this will require much true research, but is not this within the chemist's field and should it not be turned over to him with the cooperation of the entomologist, who will aid in the testing of promising products of the chemical laboratory? We need better spray pumps and other insecticidal apparatus, but had we not better leave their construction to our mechanics or employ mechanics specially for such work under the employ or with the cooperation of the entomologist? Is it worth while for us to spend time perfecting apparatus if we can avoid it, and even if necessary is it true research in economic entomology?

(4) In the control of insects by means of general methods of farm practice, including such methods as rotation, planting, cultivation, trap crops, immune varieties, etc., we meet with a similar problem. Is it the function of the entomologist to ascertain the very best way to grow a crop to avoid insect depredations, how much fertilizer to use per acre, how to apply it, what sort of implements to use, etc., involving all the questions of agricultural practice, or should we seek to produce or find varieties immune to insect attack? Many of us have been forced into such work. Should it not be done by our horticultural and agricultural colleagues with our cooperation, and should not a rational policy of station management so divide such problems that each of us will be working on that phase of the

problem for which he is specially trained, and not encroaching upon the field of another where often the entomologist may flounder around to the amusement of those to whom it naturally belongs? Most of us are unable to fathom all the depths of all branches of agricultural science. Should we then attempt research in these phases of economic entomology except in a cooperative way?

(5) Coming to the study of the natural agencies in the control of insect pests, we again come to fertile soil for research. (A) The study of parasites is fast coming to the front. The mere recording of certain parasites bred from certain insects is often of no practical importance, but when we come to consider the general ethology of parasites in relation to injurious insects and the possibility of their importation either for the possible effect upon the normal host or upon other hosts, we may be led to the very best type of research concerning them. The work of attempting to propagate hymenopterous parasites in sufficient numbers to be used commercially, now being attempted at the Texas College, in cooperation, I believe, with the Bureau of Entomology, is a most valuable experiment. The work of Mr. W. Dwight Pierce, recently described by him at a meeting of the Association of the Cotton Belt Entomologists, in studying the interrelations of the parasites of the boll weevil with the native hosts of those parasites and the food plants of their natural hosts, is a field of research well worthy the careful work it is receiving. Or consider the influence of climatic conditions upon parasites. How often have we observed an outbreak of an insect pest with a partial parasitism and wondered whether to advise remedial measures or to advise leaving its control to the good offices of the parasites? Would not a more accurate knowledge of climatic influences on parasites greatly aid in our prognostications in such a case? (B) Similar remarks might be in order in regard to the study of predaceous insects. We have still much to learn concerning their economic importance, and more particularly as to the best means of making use of them. (C) So far we must confess to rather poor results with fungi and bacteria used as artificial means of insect control, but they are undoubtedly large factors in the control of Nature. Here we must have the cooperation of the mycologist, but is there not room for considerable more first-class research along this line? Has the last word been said upon the control of insect pests by diseases? (D) And, finally, what do we really know as regards the effect of climate upon the mortality of insects? Prof. Wilmon Newell, with his experiment with 28,000 boll weevils last winter, will undoubtedly be able to give us much valuable data on this subject, but I think he will admit that he has but made a start along the right track. In this subject of the relation of climate to insect mortality and control we must needs make use of meteorology and probably

of physiological chemistry; but such research seems to lie more clearly within the province of the entomologist than of any of his colleagues, unless perchance he is so fortunate as to have a professional meteorologist connected with the Station, which seldom occurs, in which case such work might be carried on cooperatively to better advantage. We are doing something along this line in New Hampshire and trust that in a few years we may know something definitely as to the influence of climate on various phases of insect life and mortality.

Thus there are abundant opportunities for research of the highest order in entomology, strictly economic and related to agriculture. Pardon me for their lengthy enumeration. They have doubtless occurred to most of those present. But sometimes it is well for us to consider these matters definitely and in systematic fashion, and this paper has been presented merely with the hope that it may form the point upon which discussion and thought concerning this whole question may be brought to focus. Will not more research along the lines above indicated tend to more clearly define the exact province and scope of economic entomology? Indeed, will not such work be the making of a distinct science of economic entomology and be the means of giving it a scientific standing in the front rank of modern sciences, thus commanding and attracting the best scientific ability?

Mr. Hopkins emphasized the importance of investigators of economic problems generally devoting a considerable part of their time to the determination and verification of fundamental technical data on which to base authoritative conclusions. He also referred to the confusion in our economic literature due to the continued quotation of unverified statements. As an example, he had found, by study of anatomical details, that the white-pine weevil (*Pissodes strobi* Peck) is specifically different from two or three undescribed species heretofore confused with it, not only in structural characters, but in life history and habits. Therefore much of the literature, especially that relating to remedies, is practically valueless, because it was founded on facts relating to several species with different habits. He stated further that in order to make the best progress in economic work governing boards and the public must realize more than they have heretofore that the economic entomologist must first have an authentic identification of the species, based on a knowledge of the systematic and technical facts, before he can determine the best methods of combating the injurious or of utilizing the beneficial ones.

Mr. J. B. Smith gave a short statement of the many difficulties which surrounded the earlier entomologists, particularly those who attempted to carry on systematic work. He related some incidents

showing the self-denial and perseverance of such men as Doctor Le Conte and Doctor Horn, and brought out the fact that if the work done by some of the older entomologists now proved defective, we should never lose sight of the poor equipment which they had and the many obstacles which they were obliged to overcome in order to do any work at all.

On request, Mr. Comstock gave a brief outline of the entomological work at Cornell. Both the Adams and the Hatch funds are being devoted to research. One assistant entomologist has been employed at a salary of \$1,000 a year and \$500 is held in reserve for his expenses. No teaching is required, and his entire time will be devoted to investigation. He stated that, in brief, future entomologists should be broadly educated.

Mr. Fernald gave as his opinion that it was now necessary for entomologists, particularly students who are expecting to make entomology their life work, to first become well acquainted with the fundamental principles of animal and plant life. The broader the man, as a general proposition, the better work he will be able to do.

Mr. Skinner agreed with the remarks of Mr. Smith, and cited a number of instances showing the extreme difficulties under which many of the older systematic entomologists worked.

Mr. Kirkland stated his belief that entomologists, particularly economic entomologists, should be familiar with business principles, and that many failed because they had little experience along this line.

The session then adjourned until 1.30 p. m.

AFTERNOON SESSION, SATURDAY, DECEMBER 29, 1906.

The meeting was called to order at 1.30 by President Kirkland, and the following paper was presented:

NOTES ON A NEW SAWFLY ATTACKING THE PEACH.

(*Pamphilus persicum* MacG.)^a

By B. H. WALDEN, *New Haven, Conn.*

(Plate I.)

On June 11 of the past season we received a letter from one of the leading Connecticut fruit growers, stating that an insect had appeared in a peach orchard at Yalesville and was eating the young fruit. He described the insect as being very active, and about one-half an inch long, head black and white, much like that of a hornet, body brown or

^a This insect was described under the above name by MacGillivray since the presentation of this paper (*Canadian Entomologist*, Vol. XXXIX, No. 9, Sept., 1907, p. 308).—Ed.

copper color, with rather large wings. He thought it the same insect that defoliated sections of the orchard the previous season.

Being unable to identify the insect from this description, a visit was made to the orchard on June 14. It was found, as expected, that the common rose-chaffer (*Macrodactylus subspinosus* Fab.) was eating the young peaches; but flying around among the trees were a number of sawflies which the owner pointed out as being the insects to which he had referred. Although not more than ten or twelve sawflies were seen around a tree, the owner stated that they were much more numerous a few days before. Some of these were caught to bring back for identification.

On examining the trees carefully, numerous eggs were found on the underside of the leaves along the midrib and usually on the basal portion of the leaf. These eggs were nearly white in color, about 1.6 mm. in length and 0.75 mm. in width. Some of these had hatched, and small greenish-white larvæ were feeding on the leaves. They eat a narrow section, beginning at the edge, and work toward the midrib of the leaf, then roll one of the corners over, thus forming a case within which they conceal themselves, as shown in the illustration (Pl. I, fig. 2). Some of this material was collected for rearing, but through neglect the larvæ all died.

The writer was unable to visit the orchard again until July 2. The larvæ had all disappeared, but the work of this insect was seen throughout the 80-acre orchard, though only in small sections were the trees badly defoliated. The owner thought that the injury was as severe as during the previous year, and that the insect had spread over a much larger area this season.

No remedy was tried against this pest. As peach foliage in Connecticut is usually injured by any spray, it seems to the writer that the most practicable treatment would be to keep the soil well stirred, especially near the base of the trees, for a few weeks after the larvæ begin to go into the ground to pupate. Spraying the ground around the trees with kerosene emulsion might also kill many of the larvæ.

It might be well to state that this is an old orchard, which was somewhat injured by the winter of 1904, and has not been cultivated as thoroughly as the owners usually cultivate their orchards.

Specimens of the sawfly were sent to Dr. A. D. MacGillivray, who pronounced it a new species. The female is about 9 mm. (three-eighths of an inch) long, with a wing expanse of 19 mm. (three-fourths of an inch). Head and thorax black, with pale-yellow markings. Antennæ about one-fourth of an inch long. Abdomen with two black basal segments, the remainder being brown-ocher in color. Legs with pale yellow femora, tibiæ and tarsi brown-ocher.

It is hoped that this insect can be studied another season and notes made regarding its life history.

FIG. 1.—PEACH BRANCH, SHOWING HOW LEAVES HAVE BEEN EATEN BY LARVÆ OF THE SAWFLY *PAMPHILIUS PERSICUM*. (ORIGINAL.)

FIG. 2.—PEACH LEAVES ROLLED BY LARVÆ OF THE SAWFLY *PAMPHILIUS PERSICUM*. EGGS ON MIDRIBS NEAR BASE OF LEAVES. NATURAL SIZE. (ORIGINAL.)

Mr. J. B. Smith stated that this sawfly occurs in New Jersey, as he has found it at South Orange, and Mr. Titus reported finding it in Pennsylvania.

A paper was read, entitled:

ECONOMIC WORK AGAINST THE HOWARD SCALE IN COLORADO.

(*Aspidiotus howardii* Ckll.)

By E. P. TAYLOR, *Fort Collins, Colo.*

It is my opportunity to report the first economic work against a new and important fruit-tree pest, *Aspidiotus howardii* Ckll., which is doing great damage to orchard trees in parts of Colorado and other points in the West.

The insect has been previously reported in various entomological publications, but up to a short time ago has received little attention from an economic standpoint. Upon discovery of its injurious work in the orchard sections of western Colorado I was enabled to undertake economic experiments for its control, and it is upon these preliminary tests that this paper is based.

As an introduction I shall briefly treat of the history of the insect from its first discovery at Canon City, Colo., by Prof. C. P. Gillette, on August 31, 1894. These first specimens were taken by Professor Gillette upon the fruit of plum, and specimens were sent to both Dr. L. O. Howard and to Prof. T. D. A. Cockerell, then of the New Mexico Agricultural Experiment Station, but now of the University of Colorado. The former pronounced the insect an entirely new species, and Professor Cockerell, after study and comparison, introduced the insect as a new species, *Aspidiotus howardii* Ckll., in an original description published in the Canadian Entomologist, Volume XXVII, page 16 (1895). Professor Cockerell himself encountered the scale at Albuquerque, N. Mex., in August, 1895, upon the fruit of silver prune, which determination was again verified by Mr. Pergande, of the U. S. Department of Agriculture, from material furnished him.

Prof. Wilmon Newell contributed, in 1899, from Iowa, an article upon The North American Species of the Subgenera Diaspidiotus and Hemiberlesia of the Genus Aspidiotus, including Professor Cockerell's original description of *Aspidiotus howardii* and giving as its habitat Colorado and New Mexico.

The next mention we have of it is from Professor Gillette, of Colorado, in his Entomological Report for 1901, when he reports its occurrence for the first time upon fruit trees in western Colorado on the western slope of the Rocky Mountain range. The following year he reported its discovery upon the under surface of the leaves and upon

the tender twigs of white ash shade trees growing upon the streets of Denver.

It may be stated that notwithstanding its occurrence upon shade as well as fruit trees it is primarily the pest of the latter. In my observations in Colorado this season I have taken the species upon pear, prune, plum, almond, and apple, and at elevations above sea level varying from a little over 4,000 feet to about 7,000 feet.

Injury from infestation follows from two sources, (1) the dwarfing and robbing of the trees of their sap substance, resulting in retarded growth and cracking of the bark, and (2) an unsightly pitting of the surface of the fruit itself, with more or less attendant discoloration about the points of scale attachment. In the case of yellow skinned plums these reddened blotches about the scale situs are most noticeable and objectionable. With dark-colored plums and prunes the scales appear as many small white specks scattered over the surface, and with the pear deep pits are found in the skin, some of these measuring nearly one-fourth inch deep and as wide across at the top. Bartlett pears are especially subject to attack, as are silver prunes, almonds, and wild goose plums. These varieties and some others when infested and allowed to go without spraying will in time become completely incrustated as in the case of the San Jose scale. Early descriptions of this insect gave it as a pest principally upon the fruit instead of the tree. The tendency to infest the fruit itself is perhaps greater than with other species of this genus, but the attack is also directed to bark, twigs, and leaves. A marked preference is shown for Grimes Golden apples, this variety becoming infested in orchards where all other varieties are exempt; in fact, Grimes Golden and Geneton are the only varieties of apples observed by me to be infested. By far the greatest injury, so far as my observations have extended, has been done to pear, and to the orchardist in certain sections it is popularly known as the "pear scale." From its variety of food plants the name Howard scale is more appropriate.

Though Colorado, so far as known, is entirely free from the San Jose scale, the life history and appearance of the scale has become more or less familiar to many of the more progressive orchardists of the State from literature published upon the pest. Upon the discovery of the Howard scale in injurious numbers, however, it was a natural mistake for them to assume the presence of the San Jose scale, and this was what occurred. The two are of rather close resemblance to the unpracticed eye, and both infest their host plants in about the same manner. Both are of the same genus and subgenus, *Diaspidiotus*. The Howard scale, however, carries a much closer resemblance to the Putnam scale (*Aspidiotus ancylus* Putn.) of maple and other trees than to the San Jose scale.

The female scale is circular and flat, about 1.5 millimeters in diameter or about the size of a pin head. It is pale grayish in color when matured, much lighter in hue than the partially matured or even the full-grown female of the San Jose scale. The scale exuvium in color is a dull orange when the secretion over it has been rubbed away, which is quite easily done. The female insect is broadly pyriform in shape. It belongs to that class of the genus *Aspidiotus* where the scales of the females are nearly circular, while those of the male are more elongate or nearly oval, though an examination of the female insect beneath the higher power lens is necessary to observe the typical second pair of large lobes at the caudal end of the female, which distinguishes it from that of the San Jose, Putnam, or other scales of its group. Its nature of attack and general appearance upon heavily infested trees are quite enough to enable ready recognition from others of these types. The pallidness of the female scale is quite characteristic, some individuals being nearly white.

At the beginning of my experiment with insecticides for the control of the pest, very little was known with certainty of the insect's exact life history. Observations, somewhat fragmentary, were, however, carried along through the season and some of the facts learned.

The first observations the past spring were begun upon March 19, at which time some females were found well grown and of pale gray color. Others among these were smaller in size, some circular and some oval in outline. All these smaller-sized scales at the date mentioned showed a whitish area in their center surrounded by an area much darker and in certain cases of a dusky gray. In the center of the white area, which occupied about one-third of the surface of the scale covering, a small whitish nipple was seen surrounded by a rather shallow or indistinct ring or furrow. On account of the weathering of these scales most of them showed reddish or orange-colored centers. Out of a large number of scales counted, about 31 per cent were found to contain no living insect, representing the average rate of mortality from natural causes. Some of the dead scales were the result of parasitism by insects which were observed later in the season, while others were dead from exposure to the winter climate.

The oval male scale is much darker in color than the female. These oval male scales were found to yield adults as early as April 30, at which time several winged specimens were seen in process of fertilizing the matured scales which had lived through the winter. The males are of very minute size, pale brown in color, with black eyes. Early in June many young scales were beginning to appear, crawling over the surface of the bark in much the same active manner as the young of the San Jose scale. By June 9 many had settled down, and

the bark was thickly covered with this early summer brood. Some were seen upon both upper and lower leaf surfaces and many at this date had well-developed scale coverings already secreted over them. Many oval eggs were found, showing the females to be oviparous rather than viviparous, as in the case of its close neighbor, the San Jose scale. The exact time required for hatching is evidently quite short, for beneath these female scale coverings and along with the small clusters of eggs are usually also to be seen the minute yellowish-orange-colored and newly hatched young. In western Colorado it is probable that at least three and perhaps four broods are developed throughout the season, including those living through the winter; and male insects are seemingly produced throughout the greater part of the summer season.

Early in June and again in the month of August adults of an interesting hymenopterous parasite were observed; specimens were determined as *Prospalta aurantii* How., which, according to Doctor Howard, is entirely new to this species. The adult parasites emerging left the small round circular apertures which I found in the centers of so large a percentage of the larger scales through the spring and summer. Adults and larvæ of the more common lady beetle, *Chilocorus bifulnerus* Muls., also played some part in the destruction of the scale.

A survey of the orchard section of western Colorado showed the scale's distribution to be quite general. The growers about the region of Grand Junction have thus far been most successful in withstanding the usual destruction occasioned by pear blight, accounting at this time for an extensive pear acreage. At this place the spraying experiments were carried out and upon pear trees previously unsprayed and consequently badly incrustated with the scale at the outset of the tests.

In all about 200 pear trees were included, and the insecticides used were principally those known to have been more or less successfully used against the San Jose scale. The sprays and their formulæ were as follows:

- (1) "Rex" lime and sulphur dip, diluted 1 to 11 with cold water.
- (2) "Rex" lime and sulphur dip, diluted 1 to 8 with cold water, with 15 pounds lime added per 50 gallons of spray.
- (3) Lime-sulphur-soda wash, prepared without use of external heat and boiled by the soda and heat of the slaking lime—30 pounds lump lime, 15 pounds sulphur, and 5 pounds caustic soda per 50 gallons of water.
- (4) Lime-sulphur wash, prepared in usual way by boiling 45 minutes with external heat, and composed of 15 pounds lime and 15 pounds sulphur per 50 gallons of water.

(5) Kerosene emulsion, 20 to 30 per cent kerosene, prepared with the use of slaked lime.

(6) The patent soluble petroleum spray "Scalecide," used with a 5 per cent dilution in cold water.

(7) "Chloroleum," a new commercially prepared creolin spray, used also with a 5 per cent dilution in cold water.

Both spring and fall applications were made with most of these preparations, the spring sprayings being given April 5 and 6 and the fall treatments on November 7.

Without going into the details of the experiment, it may be stated briefly that the lime-sulphur washes were most successful when cost and efficiency were considered. Comparing the results of my work against the Howard scale with my experience with the San Jose scale in Illinois, it was quite evident that the Howard scale was the more easily controlled. Examination of the structure of the scale covering protecting the Howard scale explains, perhaps, the cause of its greater susceptibility to contact sprays. With the Howard scale the coverings are much more loosely attached to the bark. This gives the spray less resistance in coming into direct contact with the body of the insect concealed beneath.

All of the lime and sulphur preparations were effective. "Chloroleum" did not promise well as an insecticide for this purpose and almost a complete failure was made in attempting to prepare the kerosene emulsion with the use of lime. A dry, finely slaked lime, discarded as a by-product from a local sugar factory, was used, which may have been the cause of its failure. The lime failed completely to properly absorb the oil, and when an attempt was made to agitate the mixture by forcing through a force pump the thick, doughy mess completely clogged strainer and nozzle.

"Scalecide," on the other hand, gave much promise. It was sprayed in November and consequently the final results are yet unknown. It must be said, however, that the soluble petroleum sprays are most deserving insecticides from the standpoint of both effectiveness and convenience of use, and could their cost and transportation expense to the West be lessened they would receive more universal use by orchard men.

As stated, the lime-sulphur washes were found best adapted for combating this new scale pest. The self-boiled mixtures and those boiled by caustic soda in the experiment were not so satisfactory as the ones prepared by boiling properly, but the inconvenience of cooking creates a certain demand for a material ready made and possible of immediate dilution with cold water. The "Rex" lime-sulphur used in the experiment was of the latter class. It is a concentrated lime-sulphur solution formerly used in the West as a stock

dip and first used in the Middle West against the San Jose scale by Professor Forbes in Illinois. The following comparison shows it slightly less effective than the orchard-boiled lime and sulphur wash, but successful enough to justify use under special conditions, and presents in brief a comparison of the "Rex" lime-sulphur wash and the homemade lime-sulphur spray as used in this test against the Howard scale of pear.

Spray.	Formulae.	Date sprayed.	Percentage of scales dead at examination, April 25.	Percentage of pears infested at picking time, August 17.
"Rex"	"Rex," 1 part; cold water, 8 parts. (Lime, 15 pounds per 50 gallons.)	April 6	84.4	2.8
Lime and sulphur.....	Lime, 15 pounds; sulphur, 15 pounds; water, 50 gallons; boil 45 minutes.	April 5	93.8	.6
Check	Not sprayed.....		55.0	96.1

Two means of comparison are to be had in the above table, the first as indicated in the count of the percentage of scales dead upon the bark of the tree upon April 25, at which time the Rex lime-sulphur gave 84.4 per cent dead, the lime-sulphur wash 93.8 per cent dead, and the plat given no spray 55 per cent scales dead of those counted. As will be noted, this count was made only twenty days following spraying, and a later examination would have yielded a higher percentage killed in both of the treated plats, due to longer action of the adhering spray.

The second means of comparison, and that of more practical bearing to the fruit grower, is shown in the table as a count of the percentage of pear fruit showing the pits from infestation, found on the surface in a count made on August 17, at time of harvest. This comparison shows but 2.8 per cent pears infested upon an average borne upon the tree in the plat sprayed with the Rex. The lime-sulphur plat gave 0.6 per cent infested pears or less than one per 100, while the check or unsprayed trees gave 96.1 per cent infested. In each case many hundreds of pears were counted in securing these ratios. They show good results with both sprays, but a slight advantage with the orchard-boiled mixture. It was also apparent that pears borne upon badly infested trees not treated would almost without exception show infestation sufficient to exclude them for market purposes.

With one proper dormant spray of lime-sulphur wash, it is shown that the Howard scale may be controlled, and, unlike the San Jose scale, it seems probable that one spraying in two years will accomplish this result.

The first spraying at Grand Junction against this pest was done with whale-oil soap. In one orchard observed at that place it was reported that three years ago over \$1,000 worth of fruit was rejected by the inspector of a local fruit growers' association on account of the pears being rendered unsightly by scale pits and blotches caused by the insect, aside from the dwarfed tree growth and cracked condition of the bark. The year following this a thorough spring spraying with lime-sulphur wash reduced the injury to tree and fruit to a point beneath practical notice. The treatment was repeated last March, with the remarkable result that of a crop of 2,200 boxes of choice and fancy pears packed this fall only 2 pears were encountered bearing scales. Last spring in the orchard section for a few miles surrounding Grand Junction, one of the fruit districts of western Colorado, 40,000 pounds of sulphur were used by the members of a single fruit growers' association, principally against the Howard scale of the pear, besides a carload of 30,000 pounds of the "Rex" lime and sulphur concentrate introduced as an experiment.

This season these same growers will consume a greatly increased amount of sulphur, and have already placed their order for 5 carloads of the "Rex" product, aggregating over 400 barrels of the concentrate, sufficient when diluted with 11 times its volume of water, to produce 4,400 barrels of spray. As a rule this will be applied with gasoline power-spray outfits, nearly 200 of which are owned and operated in the orchards about this one point mentioned.

Economic entomology is enthusiastically practiced by the progressive fruit growers of Colorado, and well it may be when her fancy grade lime-sulphur sprayed Búerré de Anjou pears net the growers \$5.10 per box, as they have done this season when placed upon the New York City market.

In reply to an inquiry, Mr. Taylor stated that the Howard scale was found on native trees, such as white ash, also on apple and plum. He considered it quite a serious pest and that it might possibly be shipped out on nursery stock. The danger, however, was somewhat remote, as no stock was shipped east from Colorado.

Mr. Titus remarked that this insect was probably a species native to Colorado, as he had found it on native wild plum.

Mr. Taylor stated that most of the spraying in Colorado was done with gasoline spraying outfits, and that each outfit would cover about 20 acres of bearing orchard. He had used the Rex lime and sulphur mixture with fair results. It was sold in Colorado for \$12 a barrel.

A paper was presented, as follows:

THE MAPLE LEAF STEM-BORER.

(*Priophorus acericaulis* MacG.)

By W. E. BRITTON, *New Haven, Conn.*

[Withdrawn for publication elsewhere.]

The following paper was read:

THE VALUE OF PARASITES IN CEREAL AND FORAGE CROP PRODUCTION.

By F. M. WEBSTER, *Washington, D. C.*

While there has been much said of late concerning the beneficial influences of parasites in affecting insect pests of the orchard, garden, and truck farm, we hear little of this in relation to such as attack grain and forage crops. It might almost be said that the average farmer, from his point of view, sees only the benefits derived by the fruit grower, while the aid that he himself receives from the good offices of parasites is quite outside his vision. The facts are, however, almost if not quite the reverse, and it is probable that the crops of the average farmer are more continually under the protection of, and greater losses are prevented by, beneficial insects than in any other field of husbandry. The real difference is that in the case of the farmer they are obscured or overlooked; while among the fruits and vegetables, where observation is less difficult, much more that transpires is seen and recorded in reports, bulletins, and the agricultural press. A case in point from my own observation many years ago will serve as an illustration.

There was a threatened outbreak of wheat midge (*Contarinia tritici* Kirby) in a field of wheat, the larvæ at the usual season being excessively abundant. During late afternoons and early forenoons the heads of wheat were visited by great numbers of predaceous insects that hunted about among the bracts, and greedily devoured the larvæ of the midge. Day after day this was observed, until it seemed to me that none of the victims could have escaped capture. Gradually and naturally the heads changed from green to golden, and the threatened disaster did not materialize, there being very few of the midge larvæ remaining in the heads, where a short time before there had been myriads. To me, who had watched the proceedings daily, it was clear that the crop had been saved from serious damage, if not, indeed, a loss so great as to render the field of too little value to harvest. One day at harvest the farmer himself was observed standing with a neighbor near the borders of his field, and I caught a few words of his conversation as I passed by, to the effect that his success in wheat growing that year had been due to good farming. I asked

him if he had noticed any insects on the heads of his wheat, and he gave me most conclusive proof to the contrary by stating that he had never looked to see if there were any there, but he thought not.

Another case in point will be appreciated by those who are familiar with the actions of the Hessian fly and its parasites. A field of wheat in northern Indiana back in the 80's was so seriously affected by the fly in the fall as to render the outlook for the owner getting even his seed back at harvest exceedingly dubious. With the coming of spring the wheat plants, or rather the belated tillers thereof, sprang up and the owner harvested a crop of 20 bushels per acre. There was almost no infestation of fly in spring. Now for the reason for this seemingly supernatural phenomenon. In fall I had taken the precaution to collect a quantity of dead wheat plants from this field and placed them in breeding in a warm room. Very few flies were reared, but swarms of *Polygnotus hiemalis* Forbes emerged from the "flaxseeds" ensconced among the dead wheat plants. I now know that of that yield of 20 bushels per acre secured from this field not less than 18 bushels per acre should go to the credit of *Polygnotus*. This was the year following Doctor Forbes's description of his *P. hiemalis*, and none of us at that time knew much about it. More than ten years after, in another State, almost a parallel case was presented. This time, however, a most perplexing matter developed. The infested wheat was of a variety that persons with seed to sell were endeavoring to boom, and one of the virtues claimed was immunity to attack of the Hessian fly. Although I reared fully 100 *Polygnotus* to one fly from the dead plants in the fall and knew absolutely that the parasite saved the crop by sweeping the fly almost out of existence, yet I found both myself and the *Polygnotus* alike discredited and the claim set up and sustained that the whole thing was due to certain virtues possessed by this particular variety of seed wheat, for which a fancy price was demanded and obtained.

Within the last two years, in a section where spring wheat only is grown, we have found a similar condition existing, and all facts so far obtainable go to show that but for the efficient aid offered by *Polygnotus* spring-wheat growing in the Northwest would cease to be profitable. These minute insect parasites save the country millions of dollars, yet they are unknown except among entomologists.

Again, Dr. Paul Marchal, who studied the development of *Polygnotus minutus* in France, but did not witness oviposition, states that there is every evidence of polyembryonic development, the egg being deposited in that of the Hessian fly and hatching in the body of the maggot of the latter after it has left the egg. That is to say, there may be several germ cells within a single ovum, and consequently several larvæ may develop from a single egg. We have not by any means cleared up the obscurity relative to an American species,

though all of my assistants have repeatedly observed the female *Polygnotus* ovipositing in the eggs of the Hessian fly, and Mr. G. I. Reeves has found larvæ to the number of 28 in the matured maggot. We have thus observed the oviposition in eggs of the fly and reared adults from the "flaxseeds." If Doctor Marchal's suppositions are correct, and, as is very probable, our American species have similar habits, their economic importance will be immense, for it will be possible for a single female *Polygnotus* to place her supply of eggs singly in those of the Hessian fly, and the number of her progeny would be limited only by the sustaining capacity of the host larvæ.

Another parasite with polyembryonic development is *Platygaster herrickii* Pack. Mr. Reeves has observed this also, ovipositing in the eggs of the Hessian fly, and at another time, in a locality several hundred miles distant, he counted over 40 larvæ in a single maggot.

Whether or not there are other species of *Platygaster* infesting the Hessian fly it is just now impossible to state, but there are certainly more than one species of *Polygnotus* engaged in this efficient work, and the saving to the wheat growers of North America through their influences is immense.

The whole problem of polyembryony among parasitic insects is exceedingly interesting and of the utmost economic importance. When Doctor Howard called attention to the enormous numbers of *Copidosoma* that he had reared from a larva of *Plusia brassicæ* in 1882, the number from a single individual, of 19 larvæ, being 2,528, he was unable to learn by dissection that the female *Copidosoma* was capable of laying more than 160 eggs. Later, in Ohio, I reared from a lot of 5 larvæ of this species the enormous number of over 4,800, and since that time Mr. C. W. Mally, in South Africa, reared 2,112 from a *Plusia* larva, while Monsieur E. Bugnion has reared over 3,000 individuals from a single host larva. Doctor Howard first thought that more than one female *Copidosoma* had oviposited in a single host larva, but this has since been found to be exceedingly unlikely. In the case of some of our American species of *Polygnotus* and *Platygaster herrickii* the female parasite positively refuses to oviposit in eggs that have already been visited by another female. Attention has also been called to the almost general phenomenon of all individuals from a single larva being of the same sex, all of which goes to imply that but a single ovum is placed in a single host, and the extent to which the parasitic larvæ will multiply from this is only limited by the size of the host larva. Thus we have an explanation of what has been a puzzle to entomologists, viz, the sudden and almost complete disappearance of the Hessian fly between broods. It also indicates the great value of these insects for use in introducing parasites in sections where an outbreak of the host insect is seemingly impending. How many of our parasites have this polyembry-

onic development we do not know, but where we rear such an abnormal number of parasites from a single host, and those from a single host are all of the same sex, the case will at least bear investigation.

These parasites have, however, one weak point. A sudden disaster to their host reduces them almost to the verge of extermination. For instance, over a considerable portion of the central Pacific States during the past autumn there was so much rain just prior to seeding time that the ground could not be prepared for wheat sowing until late. Hence there are but few of the fly even in the earliest sown fields, thus reducing the food supply of these parasites to the minimum and preventing their breeding in any considerable numbers.

There is another group of parasites of the Hessian fly whose habits are normal, but which have a greater number of host insects. Among these may be placed *Eupelmus allynii* French. This species ranges from New England to Montana, south to South Carolina, Georgia, and Texas. The insect has two advantages in that it is double brooded and is also a parasite of the several species of jointworms (*Isosoma*) that work in wheat, rye, and cheat, thus being present among these grains whether infested by the Hessian fly or not. We also rear it from *Isosoma*-infested timothy, orchard grass, quack grass, *Muhlenbergia*, *Elymus*, and in the West from *Bouteloua*, *Stipa*, *Sporobolus*, and *Hordeum jubatum*. So, while this parasite does not increase as rapidly as *Polygnotus* and *Platygaster*, it has the advantage of not being so dependent upon a single host for sustenance. This double-brooded feature is also found in *Semiotellus* (*Stictonotus*) *isosomatis* Riley, *Semiotellus chalcidiphagus* Walsh and Riley, and *Websterellus tritici* Ashmead, parasites of single-brooded jointworms, but not, so far as is known, of the Hessian fly.

Turning to the parasites of *Isosoma*, we find here also illustrations of the value of parasitism. While there are a few species that we expect to breed from the larvæ of any or all of them, there are others that seem to be restricted to a single species. As an illustration, we rear an undescribed species of *Cryptopristis* from *Isosoma tritici* Riley infesting wheat, but in over 400 rearings of *Isosoma* we have failed to get it from any other host. Its area of distribution is not that of *I. tritici*, however, as it seems to occur only from New York to Virginia, Kentucky, and eastern Illinois. This is also two brooded, while its host is single brooded, and from the enormous numbers that emerge from wheat stubble infested by the *Isosoma* larvæ, sometimes in the ratio of 35 or 50 *Cryptopristis* to one of the *Isosoma*, it will be seen that they are no small factor in holding the pest in check. But here, again, the entomologist does not know enough about it to be certain whether or not it is described, and the farmer is ignorant of its existence.

In the case of a species of *Isosoma* seriously affecting timothy, we have almost a parallel case, as we rear with it an undescribed species of *Websterellus* from New England to Tennessee and west to the Mississippi River, the absence of *Isosoma* apparently increasing and decreasing in proportion to the abundance of the parasite. As the pest decreases the yield of timothy seed from 5 to 18 per cent, and injures the hay crop as well, it will be seen that the parasite is, to a certain extent, saving the property of the farmer.

Take as another illustration the army worm (*Heliophila unipuncta* Haw.), which we know does not occur destructively in the same locality two years in succession. Anyone who has watched an outbreak has hardly failed to observe the havoc wrought by one or more species of tachinid flies. I remember distinctly an outbreak years ago in Indiana, where immediately after the outbreak of the army worm the fields were literally swarming with these flies, and the hum of their flight, as they flew about among the stubble and grasses, reminded me of bees. With a single exception, in the last twenty-five years, I have never observed an outbreak of this pest without these useful insects being present in abundance.

Perhaps the most serious insect pests of the arid regions, where irrigation is followed by alfalfa culture, are the several species of grasshoppers, and scarcely a year passes that numerous complaints do not come to us of their depredations. One of the most common species involved is *Melanoplus differentialis* Thos. Under date of August 7 a report was received from Fort Laramie, Wyo., stating that large numbers of these grasshoppers were dying and clinging to the alfalfa and weeds, over an area of about 6 acres. The material received was a mass of dead, disintegrating, and decaying bodies of this species of grasshopper, thickly populated with maggots from which *Sarcophaga georgina* Wied. was reared. Clearly it was this fly that had caused the mortality among the grasshoppers. June 6, previous, a similar complaint had been received from Lakin, Kans. In this last case the statement was made that in some instances over patches of nearly an acre, in an alfalfa field, the plants were literally covered with dead grasshoppers. While the grasshopper problem is not now, nor is it likely to be, settled by this species of *Sarcophaga*, it is clear that the restraining influences of these dipterous parasites are great; and we are not in a position to say what the grasshopper situation might not be if the parasites were not present.

The phenomenon that is most likely to attract the attention of the ordinary farmer is that of cutworms being devoured by *Calosoma* larvæ. I judge this to be the case from my own experience in connection with entomological correspondence, as there is hardly a year during which cutworms are especially numerous when our attention is not called to the matter by letters from farmers who have observed

cutworms to be attacked and eaten by their black enemies. Sometimes it is clear that a serious attack of cutworms is prevented or decidedly lessened in this manner. I distinctly remember, years ago, when there was a severe outbreak of the grain aphid *Macrosiphum granaria* Buckton in Indiana. After the wheat in badly infested fields was harvested and in the shock, standing with one of these shocks between me and the late afternoon sun I could see the emerging parasites swarming in the air just about and above the cap sheaves. A farmer might see them, but to him they would be nothing but "pissants." There is hardly an entomologist who has attempted to rear adult insects from the larvæ who has not failed again and again by reason of the presence of parasites.

Now, I wish to submit that the causes and effects in all of these phenomena are always present; whether they are observed or not does not in the least alter the situation. The more light we can get on the actions, habits, and life history of parasites the better we shall be able to utilize their force in applied agriculture. The forces of Niagara Falls ran to waste for ages because no man understood how to conserve and apply this power. The forces of insect parasitism are, clearly, not *all* going to waste in this manner; but we are too ignorant of this force or how best to apply it in agriculture to derive more than the crude benefits that naturally follow its primitive influences. How much might this power be improved and directed if we only understood how to do it! To the ordinary farmer, at the present time, it apparently makes little difference whether the development of a parasite is polyembryonic or otherwise; whether it is dimorphic or not. But we can never hope to put him in the way of deriving the greatest benefit from the interaction of parasites and hosts until we have sifted this matter over and to the bottom. Indeed, we shall not know ourselves how to search for and control this immense, natural, powerful element until we have done this. In the past we have perhaps been too much like the medical practitioner, who gave his patient not the treatment that he should have had for his well being, and it might even be detrimental, but that which tasted best and was the most pleasing. The patient, in return, eulogized the doctor, paid him a round fee, and told him what a great man he was, which was, of course, sweet and pleasing to that sort of a man of pills and capsules. There has been altogether too much of this interpolated into agricultural science already, and some one, perhaps at the risk of becoming temporarily unpopular among the unscientific, must accept the trust and work out such obscure problems from the beginning, securing for those who may or may not understand the best and most that is to be derived from a knowledge to be gained only by thorough painstaking investigations.

This paper was followed by a general discussion, which was participated in by Messrs. Titus, Fletcher, Hopkins, Bishopp, Kirkland, and others. It proved to be of exceeding interest to all present and served to show the value of thorough and careful investigation of problems which are not given the attention they deserve.

Mr. Webster added two interesting facts relative to *Isosoma tritici*. In the past it had always been thought necessary, as a precautionary measure, to burn the infested bits of hardened straw that break up in thrashing the wheat, many being carried out with the grain instead of going over in the straw. Several experiments in rearing adults from large numbers of these broken bits of straw, collected about elevators and threshing machines, has shown that almost all of the

FIG. 5.—Wheat straws injured by the jointworm (*Isosoma tritici*) from which the jointworms have been removed by some beneficial animal, perhaps the short-tail shrew (*Blarina brevicauda*). (Original.)

larvæ of both *Isosoma* and parasites are killed, probably by the concussion of the cylinder of the thresher. In some cases we have been able to verify these experiments by collections of stubble from fields in the vicinity of these elevators. So far as we have gone into the investigation everything indicates that the danger from these broken bits of hardened straw, or even the straw itself, is of too little importance to be worth consideration. Prof. R. H. Pettit, of Agricultural College, Mich., and also one of Mr. Webster's own assistants, Mr. W. J. Phillips, in

northern Indiana, have this year (1906) found great numbers of straws affected by the jointworm, where the enveloping sheath has been torn away and the galls formed by the larvæ deftly eaten away and the jointworms missing. In no case is the entire gall gnawed away, but just enough of the walls immediately over the larva to make possible the removal of the latter (fig. 5). While we have not been able to get definite information as to the identity of this decidedly beneficial animal, suspicion seems to point to the short-tail shrew (*Blarina brevicauda*) as the species to which credit should be given, and probably much of the work is done while the grain is in shock.

A paper was presented as follows:

SOME GEORGIA INSECTS DURING 1906.

By R. I. SMITH, *Atlanta, Ga.*

The year just closing has not been notable for any particular outbreak of insects, though certain crops have suffered more than commonly from the attack of common species. By that I mean that the weather conditions were favorable to the development of many field crop insects, such as the cotton bollworm, harlequin cabbage bug, army worm, corn weevil, and others. The record of insects of the year attacking staple crops is, however, much less interesting than that of last year.

COTTON INSECTS.

The bollworm (*Heliothis obsoleta* Fab.) was reported from Mitchell, Houston, Coffee, Lowndes, Baldwin, and Madison counties. Our correspondent from Valdosta, Lowndes County, reported his cotton damaged to an extent of nearly 90 per cent. It may be of interest in connection with bollworm injury to state that the feeding holes of the worms were in many instances the forerunners of anthracnose, a disease which caused the loss of thousands of bales of cotton in Georgia this year. Except for being an aid to the entrance of the anthracnose disease we do not believe that the bollworm caused its usual amount of damage.

The cotton leaf-worm (*Alabama argillacea* Hübn.), strange to say, was not once reported during the year. This meant that it did very little, if any, actual damage, but not that it did not appear. We are not so fortunate as to have become entirely rid of this insect.

The cotton aphid (*Aphis gossypii* Glov.) was present as usual on the young cotton.

Luperodes brunneus Cr., reported from Georgia last year at the meeting at New Orleans as the new cotton beetle, appeared again this season, but not in such great numbers. This insect was discovered at Lyons, Tatnall County, in a 10-acre field, and at Woodstock, Cherokee County, both reports coming in on July 9.

Chalcodermus aeneus Boh., the cowpea-pod weevil, was first reported from Statesboro, Bulloch County, on May 5, its earliest reported occurrence in Georgia. Our correspondent stated that young cotton was being destroyed by this insect as fast as the plants appeared above ground. Specimens from Metter, Bulloch County, were reported as ruining young cotton in a few days. Specimens were also received from Emanuel, Randolph, Berrien, and Clay counties. In nearly all instances the weevil was present in cotton fields following a crop of cowpeas. Rotation would undoubtedly prevent most of the damage caused by this insect, so far as cotton is concerned.

Euphoria melancholica Gory and *Euphoria sepulchralis* Fab. were found eating into cotton bolls, probably after the latter became injured by the anthracnose disease.

Pissodes strobi Peck and *Aramigus fulleri* Horn were collected in cotton fields, both being mistaken for the boll weevil by the farmers who collected them.

Carpophilus dimidiatus Fab. was frequently found in rotting cotton bolls, and *Conotelus obscurus* Er. was discovered feeding in a cotton bloom, evidently having emerged from a rotting cotton boll.

GRAIN AND GRASS INSECTS.

Mayetiola destructor Say, the Hessian fly, seems to have been less abundant than usual in the wheat fields. This condition is due largely to the fact that heavy and continuous rains prevented the sowing of wheat early in October, when the first fields are usually sown. Our observations show that wheat sown after October 15 is comparatively free from the fly in ordinary seasons. Volunteer wheat which came up during the past August and September was in nearly all cases found badly infested.

The jointworm (*Isosoma tritici* Riley) is present in wheat and certain grasses, but has not been studied in Georgia.

Sitotroga cerealella Oliv. was found in stored wheat at Austell, Ga. An interesting occurrence in connection with this record is that practically all the grain moths were killed by the parasite *Pediculoides ventricosus* Newp., which was found in countless numbers on the wheat and on the floor of the building where the grain was stored. A complaint came to us that some insect of almost microscopic size was very annoying to the persons who had the handling of the wheat. Upon investigation the trouble was found to be due to the parasite just mentioned. The grain moth was found also at Cornelia infesting stored corn.

Laphygma frugiperda S. and A., the fall army worm, was perhaps the insect that caused the greatest popular alarm in Georgia during 1906. On July 18 we investigated damage to corn at Cartersville, Bartow County, and found it to be the work of this army worm. The cornstalks were frequently eaten into in a manner resembling the work of the corn stalk-borer (*Diatraea saccharalis* Fab.). In fact, when we first received a report by letter we supposed the trouble must be caused by the latter insect. Young corn about knee-high was eaten down almost completely, and larger corn damaged severely. It was learned that the worms first appeared in a grass field, from which they migrated to the corn. One field of cotton near Cartersville was damaged slightly. This army worm was next reported on August 4, from Pinehurst, Dooly County, where it was devouring

grass and cotton—several fields of the former and one of the latter. Cotton was attacked in a peculiar manner. The leaves were not injured materially, but the young squares were devoured, and the worms had eaten into young bolls, destroying them completely.

This army worm appeared over considerable areas of grass fields at Albany, Vienna, and Americus. At Albany it was learned that the worms first became noticeable about August 1, and nearly all disappeared by August 10. My assistant, Mr. Lewis, who investigated the outbreak at all of the points mentioned, found great numbers of pupæ on August 14, from which adults emerged a few days later. The greatest injury caused by this insect in any one locality was at Albany, where 60 acres of grass were destroyed. The total loss in the State amounted to several thousand dollars. Cowpeas were injured slightly at Albany.

No other important insects of grain or grass have been reported or observed, except the usual cutworms and the white grubs attacking the roots.

FRUIT-TREE INSECTS.

Aspidiotus perniciosus Comst. was quite well controlled last winter and spring wherever lime-sulphur wash was properly applied. The soluble oil preparations have been used as a remedy with more or less success. Their value for Georgia conditions has not yet been fully demonstrated, though we feel certain that a strength greater than recommended by the manufacturers must be used to insure success. We have recently seen good results following the use of Scalecide at strengths of 1 to 10 and 1 to 12. The other soluble oil preparations have all been used in experiments made at Fort Valley during last October and November.

Scolytus rugulosus Ratz., *Sanninoidea exitiosa* Say, *Anarsia lineatella* Zell., and *Conotrachelus nenuphar* Hbst. all did about the usual amount of damage during the past season. The last named, the plum curculio, is undoubtedly an important factor in connection with brown rot of peaches. Nearly all the first rotten fruit is stung previously by the curculio, and it seems certain that the ruptures in the skin made by this insect offer a favorable entrance for the brown-rot fungus.

Aphis persicae-niger Sm. on peach foliage was reported as being quite injurious at Kensington, Silver Creek, and Augusta, Ga. Tobacco decoction was found to be the most effective remedy for this insect. We have not received any reports of occurrence of the root form of this aphid.

Diplotoxus frondicola Say was discovered at Cave Spring, Floyd County, eating foliage from June-budded peach stock.

Saperda candida Fab. and *Chrysobothris femorata* Fab. have caused some damage to apple orchards, but on account of the small number of apple orchards in the State these borers are not given sufficient attention.

The codling moth (*Carpocapsa pomonella*) has been the basis of some experimental work by the board of entomology in Georgia this past season. Our life history study, carried on by W. V. Reed, has shown quite conclusively that we have only two broods of this insect in Georgia. Our spraying experiment failed for lack of sufficient infestation in the orchard selected for the experiment.

Aphis mali Fab. causes great damage to young apple orchards and also in the nursery rows. Kerosene emulsion as it is commonly prepared has not been found to be a reliable remedy. Tobacco decoction has given better results.

Hoplia trivialis Har. destroyed opening buds of Japan plum at Adairsville. Specimens were received April 12 with a report that the beetles were preventing the better portion of a good-sized tree from putting out foliage. The buds were eaten out close down to the base.

The woolly aphis (*Schizoneura lanigera* Hausm.) still continues to be one of our worst apple pests. As I reported last year, we have successfully killed the root form by application of kerosene emulsion. It may be of interest to state here in advance of a bulletin which we hope soon to publish that 10 and 15 per cent kerosene emulsion have been used without injury on roots of apple trees from 2 to 10 years of age. We injured 2 trees by an application of 40 per cent emulsion, but have not observed any injury from lesser strengths, not even from 30 per cent, which was tested. We would, however, consider it unsafe to use over 20 per cent emulsion and 10 or 15 per cent seems to be amply sufficient. The aerial form of woolly aphis is best controlled by tobacco decoction.

GARDEN INSECTS.

Without giving a list of all the insects injurious to garden crops I wish to say that these pests have been rather more than commonly abundant, their increase being due, perhaps, to weather conditions. Harlequin cabbage bugs, cabbage worms, cutworms, aphides, squash bugs, etc., were very much in evidence during the past summer.

SHADE AND FOREST TREE INSECTS.

Hyphantria cunea Dru., the fall webworm, appeared in great numbers during August and September in a considerable portion of Georgia. In Columbia County I observed nests on September 4 on persimmon bushes that had been entirely stripped of foliage. In

that county it looked as though every persimmon bush and tree was inhabited by the fall webworm. Many farmers stated that this pest was more abundant this year than for many years past.

Datana integerrima G. and R. was found early in September in great numbers on a walnut tree in the city of Atlanta.

Saperda calcarata Say, the poplar borer, was found early in September severely injuring poplar trees at Lumpkin and Columbus. We were unable to breed specimens, but the samples of their work seemed conclusive evidence of the identity of this species.

Sesia tepperi Hy. Edw. and specimens of maple bark showing the work of the borer were received from Cordele on April 14. At that time we received 1 larva and 2 pupæ. Later we got more material, but only succeeded in rearing 1 adult, which emerged May 8. The injury to one maple tree was quite severe. Concerning the extent of injury on neighboring trees, we were not able to learn anything definite.

Pulvinaria innumerabilis Rathv. was discovered in small numbers May 1 on linden trees in Atlanta. At this date the egg sacs were fully developed, but no young were crawling. After being carried to the office the eggs commenced to hatch, about May 10.

Pulvinaria acericola W. and R. on sugar-maple leaves was received May 22. The egg sacs were numerous enough nearly to cover the leaves.

A plant-bug (*Neurocolpus nubilus* Say) was found May 1 in great numbers on two linden trees in Atlanta. At this date the trees were in full leaf, and the bugs, while abundant over the trees, did not appear to injure the full-grown leaves. They did, however, destroy new shoots on the trunk and about the base of the trees.

The pecan bud-worm and twig girdler have caused some injury to pecan nurseries and groves.

An aphid (*Callipterus carvella* Fitch) was found in abundance in a pecan grove near Valdosta, in south Georgia, but did not appear to cause much injury.

MISCELLANEOUS INSECTS.

The ragweed borer (*Eucosma strenuana* Walk.) was bred from ragweed collected at Augusta, August 22. Nearly every ragweed plant examined was found to have from one to several borers. The larvæ are easily located by the swollen spaces on the stems.

Sibine stimulea Clem., the saddle-back caterpillar, was found on rose at Middleton, and on pear nursery stock at Whitesburg, both during September. One larva collected on September 21 was barely half-grown.

Megalopyge opercularis S. and A. was frequently sent to the office last summer. Several correspondents reported great discomfort from being poisoned by touching the larvæ. This insect must have been much more common than usual to create so much attention.

The bee moth (*Galleria mellonella* L.) was bred from bee comb taken from hives at Cornelia and near Atlanta. While no study has been made to determine the distribution of this pest in Georgia, we feel sure that it is not uncommon.

No attempt has been made by the writer to mention all the insects that have come to our attention during the year. Many are too common to deserve mention. On the whole, the insects have appeared in greater variety, if not in greater numbers, than in recent years.

In reply to an inquiry, Mr. Smith (Georgia) stated that he had used 15 per cent kerosene emulsion for the woolly aphis and had secured good results. Thirty per cent had not injured the trees, while 40 per cent had caused some injury.

A paper was read, entitled:

OBSERVATIONS ON INSECT ENEMIES OF TOBACCO IN FLORIDA IN 1905.

By W. A. HOOKER, *Washington, D. C.*

While engaged, during April, May, and June, 1905, at Quincy, Fla., in studying the tobacco thrips (*Euthrips nicotianæ* Hinds)^a and methods for its control, the writer had opportunity to observe other insects affecting the tobacco crop and takes this occasion to present briefly the notes made.

THE BUDWORMS.

(*Heliothis obsoleta* Fab. and *Chloridea virescens* Fab.)

Aside from the tobacco thrips the budworms are the most destructive pests with which the Florida tobacco planter has to deal. The former species, the bollworm of cotton, is undoubtedly the more abundant, its moths having been observed in the tobacco field more commonly than those of the last named. On April 16 eggs were observed on the lower surface of young tobacco plants in the seed bed. As the plants get larger and after they are transplanted the eggs seem to be laid largely in the tender folded center leaves upon which the worms feed as they emerge. If upon hatching they are

^a The remedial treatment of the tobacco thrips has been considered in Circular No. 68, and a detailed account of the insect is published in Bulletin No. 65 of the Bureau of Entomology, U. S. Dept. of Agriculture.

allowed to feed for a single day, much injury is done, a small hole in the leaf bud at this stage developing later into numerous large holes in the mature leaf or leaves. As described and recommended by Prof. A. L. Quaintance,^a and by Dr. L. O. Howard,^b it is the practice to sift into the leaf bud Paris green, in corn meal as a diluent (1 tablespoonful of Paris green to 1 peck of sifted corn meal), by the use of perforated tomato cans or baking powder boxes, attached to handles. The leaf buds must be kept constantly covered with this mixture, two applications per week ordinarily being sufficient, but three being occasionally necessary. When the mixture is applied at a greater strength the buds are burnt and seriously injured. After the plants reach the height of 2 feet it is necessary to open with one hand and with the other to place a pinch of the mixture in the bud. While corn meal is largely used as a diluent it is also a common practice to add lime (air-slaked) and sand, 2 quarts of each to 4 of corn meal. The lime, however, seems to be objectionable in that it possesses the property of adhering to the tobacco after it is harvested. Again, road dust is occasionally used as a diluent. Corn meal, however, is now used by the majority of the tobacco planters. The worms seem to eat the meal and with it the Paris green.

The annual cost in treating the budworm, for labor and supplies, has been estimated by several planters as ranging from \$12 to \$15 per acre.

When the plants are permitted to bud, for seed, the worms are very destructive, as they bore into the green buds and seed pods and destroy a very large percentage. Applications of Paris green assist to some extent in preventing this injury.

Cultural methods in connection with this pest do not seem to have been recommended, nor are they practiced by the tobacco planters. As recommended for cotton, corn may be found of sufficient value as a trap crop, when planted around tobacco fields, to warrant its use in this manner. When clean culture between crops is followed, as recommended in connection with the tobacco thrips, the budworm will be in part destroyed in the pupal stage by the plowing of the fields.

As the pods of the cowpeas are the favorite food of budworms, when grown in the field between crops (as sometimes practiced), they should be plowed under or harvested and the field plowed late in the fall. In this way many of the pupæ will be destroyed, for it is in this stage that the winter is passed, from 2 to 7 inches below the surface of the ground.

^a Bulletin No. 48, Fla. Agrl. Exp. Sta., pp. 184-187 (1898).

^b Yearbook U. S. Dept. of Agric. for 1898, pp. 132-134, and Farmers' Bulletin 120, pp. 14-15 (1900).

As the early spring moths deposit from 400 to 1,200 eggs within a period of ten days, the importance of destroying the pupæ and as many of the early spring moths as possible can not be emphasized too strongly.

A four-winged fly belonging to the family Braconidæ was found quite numerous in some tobacco fields. The insect has been determined by Dr. Wm. H. Ashmead as *Toxoneuron seminigra* Cress. Doctor Ashmead states that it may be found to be parasitic on the budworm.

THE HORNWORMS OR TOBACCO WORMS.

(*Phlegethontius quinquemaculata* Haw., and *Phlegethontius sexta* Joh.)

A few eggs of the hornworms or tobacco worms were observed on the tobacco leaves, both in the seed bed and in the field, on April 20. They do but little injury before the middle of May, at which time it is the practice of many planters to commence using the powder gun with 1 pound of Paris green to 5 of lime, applications being made twice each week and continued until harvesting commences. Others practice picking in place of the powder guns, women and children going through the fields and examining the plants for worms. On one plantation visited when the moths begin to appear the superintendent offers a reward, usually 1 cent each, for the moths captured. In this way many are destroyed, being killed by the men, who find them on the plants and in the shade field on the posts when hoeing, the stimulus of the reward keeping them constantly on the lookout for the moths. This destruction of the moths before they lay is very important when we consider the large number of eggs that are deposited.

Paris green has been applied by spraying, but is now entirely replaced by dusting (1 pound of Paris green to 5 pounds of lime) with a blowgun. Aside from the effect of the Paris green (applied with a blowgun) on the hornworm, it aids in controlling the budworms, grasshoppers, flea-beetles, etc. The one disadvantage in its use lies in occasional burning when the application is followed by a light rain—one insufficient to wash off the Paris green before burning takes place. One field of sun tobacco was observed that had been burnt quite badly in this way. Care should be taken to avoid this, and also to see that the application is not made shortly before or closely following the spraying of emulsion for the thrips. It is a good plan to apply it as near as possible to the time of applications of Paris green and meal for the budworms. This will, of course, apply only to shade tobacco on which emulsion spray is used.

The hornworms of the second brood, which appear in July, are injurious in that they are likely to be taken into the barn on the tobacco when harvested and there consume the whole leaf and often

near-by leaves. Care when priming or the previous use of an arsenical application will, however, prevent such injury.

Much is gained in preventing the moths from entering by keeping the sides of the shade field patched and the gates closed.

THE TOBACCO FLEA-BEETLE.

(*Epitrix parvula* Melsh.)

The tobacco flea-beetle is a small but important enemy of Florida tobacco. It was observed during the summer of 1905 working especially on tobacco in the immediate vicinity and bordering the seed bed, from which it spreads through the field. Applications of Paris green with a blowgun are depended upon for control, and are as satisfactory as any remedy that can be recommended.

GRASSHOPPERS.

Grasshoppers were very numerous in the shade field, and with the flea-beetles cause much injury to the shade tobacco. They are a very hard and unsatisfactory pest to combat. Aside from the general application of Paris green with the blowgun, on one plantation many were killed by hand; this was done by going along the road through the field and crushing them while on the ground by use of a slat or narrow board used as a paddle. This, to be sure, is a slow and crude method of extermination, but much injury can be prevented in this way at small expense.

It is the practice of many planters to keep a flock of guinea hens in the shade field, as they catch and devour large numbers of the grasshoppers. Others, however, feel that the benefit is offset by injury done in tearing the leaves. Clean cultivation between crops will assist by causing the adults to look elsewhere for food and a place for egg deposition. Fall plowing will help in destroying the eggs.

THE SALT-MARSH CATERPILLAR.

(*Estigmene acerua* Dru.)

The salt-marsh caterpillar, commonly known at Quincy, Fla., as the "woolworm," was observed as the source of considerable injury on one plantation. In a field examined May 15 one section had been largely destroyed, only the midribs and stems remaining. From one plant the caterpillars pass to the next, injuring the young plants only. They can be controlled by hand picking or the use of arsenicals.

BETTER BEETLES WHICH INJURE NEWLY TRANSPLANTED PLANTS.

Several beetles were found to do more or less injury to the young newly transplanted tobacco plants, and the nature of their work was

much the same. When the young plants are first transplanted their leaves wilt and lie upon the ground. It is at this stage that the beetles do the injury; lying upon their backs upon the ground they perforate the leaves and feed thereon. A species belonging to the Tenebrionidae seems to be the chief offender. This has been determined as *Blapstinus metallicus* Fab. by Mr. E. A. Schwarz, who says that it is a species quite generally distributed through the South.

Two other species found similarly working, but in numbers insufficient to be of economic importance, have been determined as *Epicærus formidolosus* Boh., of the family Otiornychidae, and *Opatrinus notus* Say, another tenebrionid.

THE TOBACCO LEAF-MINER OR SPLITWORM.

(*Phthorimæa operculella* Zell.)

This insect seems to be of no economic importance whatever in Gadsden County, only four or five affected leaves being noticed while at Quincy. At Dade City, in Pasco County, it has, however, become the most important insect pest with which the tobacco planter has to deal. The writer has been informed by Mr. W. W. Cobey that during the season of 1906 an average of no less than two leaves per plant has been thus affected.

CUTWORMS AND WIREWORMS.

The cutworms (Noctuidæ) were observed as especially injurious on one plantation. They are not confined in their work to the newly transplanted plants, but even attack plants a foot high with stalks one-fourth to one-half of an inch in diameter. Neither do they confine their feeding to the portion of the plant at or below the surface, but often crawl up the stalk. A number of plants examined were found to have been cut off 2 inches above the ground. In freshly cut stalks the worm can always be found near the stalk and destroyed.

Wireworms belonging to a species of *Drasterius* were found very destructive in an 8-acre field of sun tobacco on which oats and cow-peas had been grown the previous season. On account of their feeding near the surface on the stems of the young plants, boring up and down in the pith, a nearly complete resetting of the field was necessary. A search in the soil about a wilted plant would reveal one and in some cases several white wireworms from one-half to 1 inch in length.

In Wisconsin, Paris green, and in Ohio, turpentine, has been used in the water when setting as a repellent for cutworms. As an experiment with the wireworms, 1 quart of kerosene emulsion to a cask of water was tried, each plant being watered. As no further injury

from the worms took place it was supposed that the emulsion had the desired effect.

It seems to the writer desirable that kerosene emulsion be used in the water when transplanting as a repellent to both cutworms and wireworms.

LEAFHOPPERS OR SHARPSHOOTERS.

There are two species known as sharpshooters of cotton that were commonly observed in the tobacco field. These have been determined by Mr. Otto Heidemann as *Aulacizes irrorata* Fab. and *Oncometopia lateralis* Fab. The latter species was supposed by one planter to injure the bud, specimens being collected and handed the writer. The work of the budworm, however, must have been mistaken for that of these insects, as the sharpshooter can not be the source of an appreciable injury to the plants.

THE CIGARETTE BEETLE.

(*Lasioderma testaceum* Dufts.)

This pest seems to be present in small numbers in nearly all packing houses, occasionally causing some considerable injury to the tobacco in bulk.

REMEDIAL PRACTICES.

Cheese cloth is used to furnish shade for tobacco on a considerable acreage in place of slats. In connection with the insect problem it has an advantage over slats in that it keeps the insects out to a large extent. The slats, however, do much to keep out the hornworm moths, the worms being much more numerous on sun tobacco. It is equally important, however, whether cloth or slat shade be used, that the sides be kept patched and the gates closed, which fact does not seem to be appreciated by many planters.

Commencing while the plants are in the seed bed, kerosene emulsion and Paris green should be applied. These applications must be continued, the emulsion twice a week until plants are half grown, the Paris green in the blowgun twice a week from the first appearance of hornworms. Paris green and corn meal must necessarily be applied at least twice a week, commencing as soon as the plants are transplanted.

Mr. Britton stated that the cucumber flea-beetle had caused considerable injury to tobacco plants in Connecticut; that he had dipped the plants, roots and all, in a wash made of 1 pound of arsenate of lead to 10 gallons of water. This treatment did not injure the plants.

Mr. Fletcher stated that Bordeaux mixture seems to repel the cucumber flea-beetle; but it is claimed by hop growers in British Columbia that neither Bordeaux nor Disparene is effective for controlling *Psylliodes punctulata* Melsh., which is a most serious pest in hopyards there. He asked for suggestions as to a remedy.

Mr. Titus stated that the species of this genus work on sugar beets, and that they can be driven away during irrigation by disturbing the beets, thus causing the beetles to jump into the water and be swept away.

Mr. Hooker stated that Lindeman had described *Thrips tabaci* from Russia several years ago, and reported that it attacked tobacco in that country. This species is distributed from New England to Texas, but feeds upon onions. He asked if anyone present had observed it on tobacco.

Mr. Quaintance thought it probable that Lindeman had confused two or more species in his account of the life history of *Euthrips tabaci* Lind. It is stated by him that this species infests tobacco and deposits its eggs along the veins. With the American insect designated under this name, the eggs are deposited beneath the epidermis of the leaf, and there are other equally important points of difference in the habits and life of the species. So far as he knew, they never attacked tobacco, but infested onions (producing the so-called "silver-top"), cabbage, cauliflower, etc. He had frequently seen these insects very abundant on the above-mentioned plants in proximity to tobacco fields in Florida, but the insects did not attack the tobacco.

Mr. Hinds stated that there was considerable confusion in regard to the species of thrips.

Mr. Washburn mentioned finding the cigarette beetle in boxes of fine-cut tobacco.

The following paper was presented:

OBSERVATIONS ON CECIDOMYIIDÆ.

By E. P. FELT, *Albany, N. Y.*

[Withdrawn for publication elsewhere.]

A paper was presented entitled:

A SPRAY NOZZLE FOR THE MECHANICAL MIXTURE OF OIL WITH WATER OR OTHER LIQUIDS.

By E. DWIGHT SANDERSON, *Durham, N. H.*

During the year 1900 the writer made a series of tests of the oil and water mixing pumps at the Delaware Experiment Station. Practically all of such pumps on the market were thoroughly tested, using various nozzles, single and double hose, simple and double

pipe extension rods, etc. Altogether many hundred individual tests were made, and the total result of the work tended to show that none of the pumps could be relied upon to give uniformly the percentage of oil desired. A detailed discussion of the mechanical faults of these pumps is not necessary at this time, but it may be stated that those pumps made with but a single pump drawing from the barrel and oil tank at once were found entirely unreliable, whereas those having two pumps, one in the oil and one in the water tank, and the percentage of oil regulated by the length of the stroke of the pumps, were found much more reliable, although in many cases they failed to deliver the proper percentages at the pump before the liquids mixed. In none of this type, even with the double-extension rod having an inner tube conveying the oil to the nozzle, was the unequal mixing of the oil and water in the hose and extension rod wholly prevented, so that variable percentages frequently occurred. The results of this work were never published, but the records exist in support of the above summary.

It soon became apparent that to eliminate the mixing of the oil and water in the hose and pipe before reaching the nozzle, some form of nozzle must be devised which would mix the two liquids as they left it. Such a nozzle was therefore devised and a working model was made by Queen & Co., a drawing of which was made by them as shown in fig. 6. In the Thirteenth Report of the Delaware Agricultural Experiment Station (1901, p. 196) the writer made mention of this nozzle as follows: "We have had a nozzle made according to plans originated by us which it is trusted will prove more satisfactory in making a uniform mixture of a given percentage of oil and water."

In his recent bulletin upon the San Jose scale,^a Mr. C. L. Marlatt remarks concerning the oil and water mixture, as follows:

The best outlook for good apparatus of this sort seems to be in carrying the oil and water in separate lines of hose to the nozzle, uniting them in the latter, and in maintaining an absolute equality of pressure on both oil and water tanks by employing compressed air as the motive force, kept up by an air pump, the air chamber communicating with both of the liquid receptacles. * * * One or more manufacturers are now working on an apparatus of this description.

This so closely describes the system originated by the writer in 1901 that it seemed wise to make a record of our work along this line in the proceedings of this association, as the writer was not aware that any manufacturers had taken up the idea or indeed that it had attracted any attention, though we had described the apparatus to several entomologists in a general way.

^a Bulletin 62, Bureau of Entomology, U. S. Dept. of Agriculture, p. 77.

FIG. 6.—Diagram of spray nozzle for the mechanical mixture of oil with water or other liquids.

The general structure of our apparatus has already been indicated by the quotation from Mr. Marlatt. A compressed-air pump was attached to two steel boilers, which were connected so that the same pressure was constant upon both of them. Oil was placed in one tank and water in the other, and a lead of hose carried each separately to the extension rod which kept them separate until after they had left the nozzle. The general appearance of the pump was much

Fig. 7.—Compressed-air pump, suitable for spraying a mechanical mixture of oil and water, when fitted with the nozzle herein described.

like that in figure 7, which is that of a compressed-air sprayer made by an Ohio manufacturer, but which was not designed for using oil and water. The extension rod and nozzle are shown in figure 6. The rod is constructed of brass, is of any desired base, and consists of an inner tube carrying the oil for the hose connecting it with the oil tank and an outer tube connected with hose to the water tank. At the end of the rod proper the base of the nozzle was screwed on, which

gave the inner oil stream a tangential motion, thus forming a cone-shaped spray, upon leaving the aperture, like that from a Vermorel nozzle. Testing the nozzle convinced us that this section was not necessary, and that it might be eliminated and the nozzle cap put directly upon the ends of the two pipes. The end of the inner oil pipe fits into grooves of the nozzle cap so that no connection between the oil and water is possible except after they have left the nozzle. The oil leaves the cap through a central, round, reamed hole, 0.02 inch in diameter. Around this central orifice are two or more holes, each pair being directly opposite each other and opening into the outer water pipe. The holes for the water are reamed so that the streams from them are directed to meet in front of the center of the nozzle the same as in a calla nozzle. Here they come into contact with the straight or cone-shaped oil jet and the whole breaks into a fine spray evenly composed of oil and water. It is evident that if the streams leave the nozzle under the same pressure and the apertures be of the same diameter, the spray must then consist of $33\frac{1}{3}$ per cent oil, if there be two water jets and one oil jet; 20 per cent oil, if there be four water jets, etc. The percentage of oil may, therefore, be readily regulated by having various nozzle caps with a different number of water holes or with water holes of various sizes which have been tested to give a greater percentage.

Several such caps were made and the whole apparatus was given a thorough test by us. Various minor troubles were found in its structure which it would require some little time and experiment to eliminate, but our tests showed that the apparatus did practically what it was designed to do, and that with proper mechanical execution, the principles upon which it was based were undoubtedly correct. While its perfection was under way, the writer was called to Texas, where circumstances did not permit the completion of the work, and the Delaware station has given it no further attention since that time.

It is the writer's belief that a nozzle may be perfected along this line, for there is nothing specially new in the structure of the pump; and that we might thus have a perfect mixture of oil and water or other mixture, which, if it could be secured, would be of the greatest service against many insects. This now seems to the writer more of a problem for the pump manufacturer than the entomologist, and our work toward the solution of the problem is therefore made public at this time.

A brief discussion of mechanical mixing pumps followed.

Mr. J. B. Smith stated that good work had been done by some of these pumps in certain sections in New Jersey. The general opinion

seemed to be that pumps of this character now on the market would not mix the oil and water accurately.

A paper was read, as follows:

THE BEET LEAF-HOPPER.

By E. D. BALL, *Logan, Utah.*

[Withdrawn for publication elsewhere.]

The following title was presented, and the paper will be published in the report of the Bureau of Science of the Philippines:

PROBLEMS IN ECONOMIC ENTOMOLOGY IN THE PHILIPPINES.

By C. S. BANKS, *Manila, Philippine Islands.*

The following papers were read by title and ordered printed in the "Proceedings:"

OCCURRENCE OF THE THROAT BOT IN CUBA.

(*Gastrophilus nasalis* L.)

By C. F. BAKER, *Santiago de las Vegas, Cuba.*

Although the horse bot (*Gastrophilus equi* Fab.) is commonly known in Cuba, I can not find that *nasalis* has yet been reported. I had had specimens of the latter from Texas, Kansas, and Wyoming. Now, it has been taken flying about the plow mules here on the grounds of the Estacion Agronomica.

This occurrence is especially interesting, because with it may be noted an apparent divergence in habit. Ordinarily it is said to annoy animals by striking for the vicinity of the neck for the deposition of its eggs. Here, as we have observed it, it flies almost entirely beneath the body. Whether this shows any definite indication of a difference in the egg-laying habit remains to be determined.

REMARKABLE HABITS OF AN IMPORTANT PREDACEOUS FLY.

(*Ceratopogon criophorus* Will.)

By C. F. BAKER, *Santiago de las Vegas, Cuba.*

A common Cuban fruit which occurs in gardens and patios is the tree gooseberry, or grosella, a species of *Phyllanthus*. This tree is commonly completely and rapidly defoliated by a most pernicious pest—the larva of *Melanchroia geometroides* Walk. I had observed this insect in passing as to its parasites, none of which appeared to exercise the least check upon it. One day, in visiting a tree swarming with the larvæ and almost defoliated, I discovered that great numbers of them were dead and dying, their blackened skins hanging

from the twigs or lying on the ground beneath. Drawing nearer, I found that the larvæ were being attacked by a very small sucking fly, from one to half a dozen being perched on each larva, on which they remained until it was sucked dry. Mr. Coquillett has kindly determined this fly as *Ceratopogon eriophorus* Will., previously reported only from the island of St. Vincent. This mosquito-like method of attack and its remarkable effectiveness were very novel and interesting to me.

SOME NOTES ON LEPROSY IN HAVANA.

By C. F. BAKER, *Santiago de las Vegas, Cuba.*

In doing what I could to investigate the possible relation of fleas to leprosy, I have examined the leper hospital of San Lazaro, in Havana; and in this connection have made some observations and had some experiences which possess some psychological as well as entomological interest.

In the first place, to the hospital physicians it seemed absurd that I should busy myself with a study of the fleas of rats; absolutely nothing was said in any of their books about any possible relation of rats to leprosy. Then my attention was called to the fact that they and the nurses had been there some time and had not yet acquired leprosy though they were commonly flea-bitten; so the possibilities of infection by this means could be set aside at the outset. My suggestion that the same result might be true also of tuberculosis did not seem to carry much point. However, my business was with the rats and the lepers, and I found them far more productive of facts. The old hospital swarms with rats, and the rats have fleas unlike rat fleas of the temperate regions, and which, as I have noted in Science, are very nearly related to *Pulex irritans*. Talking with the lepers brought to light immediately some remarkably significant points. They were greatly interested when a possible relation between the rats and their terrible disease was hinted at, and immediately united in saying that the rats frequently carried sores like theirs and were frequently minus ears or tail. On the cots there were no nets, nor were there screens on the windows. I had the general evidence of the lepers themselves that it was no uncommon thing for rats to climb on to the cots at night and gnaw at their insensible sores. Naturally my interest in all this rose by leaps and bounds, and I immediately set traps for the rats. The first one caught had a sore on its body, and I hurried with it to the president of the board of control of the hospital, who is also the director of the city bacteriological laboratory. Together we went at once to the laboratory and asked the bacteriologist to prepare and stain some cover glass smears from the sores, applying the ordinary tuberculosis test, which

is good also for leprosy. However, this man refused to have anything to do with the matter.

My contention was not that the rats had leprosy, nor that the fleas could carry it—though, as I have stated elsewhere, I should dislike to have a flea bite a leprous blotch and then turn its attention to me—but that the whole matter was one crying urgently for investigation and that the hospital in its present condition, within the city and open to visitors and rats and mosquitoes and flies, was a possible menace to the whole community. I told the authorities that they might easily undertake a piece of investigation there that would attract the attention of the whole scientific world and contribute much to scientific knowledge of these matters. But my efforts were sterile so far as exciting any further efforts toward investigation was concerned. Yet not wholly without results either, since somewhat later the junta of the patrones of this hospital, without further facts than those I had brought out, passed a resolution in which they state it as their belief that it is possible that leprosy may be communicated by flea bites. The matter was brought to the attention of the president, and he took steps toward the removal of the hospital. The matter was again laid before the Provisional Governor after his arrival, and it seems now as if the hospital might really be removed from the city at an early date.

If any insects are really concerned in the transmission of leprosy or bubonic plague, I consider that they are more likely to be fleas than either mosquitoes or flies, since the affected areas are usually kept covered. The great length of the usual period of incubation of leprosy and the lengthened course of the disease make direct observation and evidence very difficult in any case. In the meantime I believe that it would be by far the wisest course to rigidly exclude from leper hospitals like the San Lazaro in Havana all rats, cats, dogs, fleas, mosquitoes, and flies. This would not be a difficult matter to accomplish.

ON THE ERADICATION OF THE BLACK-CURRENT GALL-MITE.

(*Eriophyes ribis* Nalepa.)

By WALTER E. COLLINGE, M. Sc., F. E. S., Birmingham, England.

For upward of thirty-five years black-currant growers in the United Kingdom have waged war against a small mite of the genus *Eriophyes*, but with little or no avail.

Its rapid increase toward the close of the past century, which threatened the successful cultivation of black currants in this country, led me in 1899 to institute a careful investigation into the life history and preventive and remedial measures, which investigation has been continued up to the present time with excellent results.^a

^a Repts. on Inj. Insects for 1904 and for 1905.

Very briefly, the results which have accrued from this investigation are as follows:

(1) The life history has been repeatedly worked through and valuable data obtained as to the dates of egg laying, the first appearance of the mites, when the old infested buds dry up, when migration commences and ceases, when the mites were found in the new buds, and the eggs, the dates the mites were last seen, and the dates the new buds commenced to swell. These details have been recorded for eight consecutive years.

(2) Two points of considerable interest have, I think, received a satisfactory solution, viz, first, what becomes of the mites which fall off the trees? And, second, do the females continue to lay eggs throughout the year?

Hoping to obtain some solution of the first problem, I devised, early in May, 1903, a wooden frame in two halves, lined with black paper, which fit closely round the stem of a currant tree. From the end of May to the end of June thousands of mites were to be found on the tray. Many were dead, but the majority alive, and although a very careful watch was kept, none was ever found to return to the tree, but in some manner or other, possibly by attaching themselves to insects, they gradually disappeared from the tray. The percentage of dead mites was about 40 per cent.

A similar tray, filled with a fine layer of soil, gave similar results, only here the mites were much more difficult to trace. As a result of the above experiments, I conclude that of the mites which fall to the ground during the migration season many are distributed by insects, birds, leaves, etc., to other trees, the remainder dying.

About the middle of June eggs and mites in various stages of development are to be found in the center of the new buds. During the winter of 1902-3 one or more buds were examined once a week from September to March, and in nearly all cases eggs were found to be present. They were certainly very few in number from December, 1902, to March, 1903, and it seemed to me that many of them were dead. From October 14, 1903, to January 30, 1904, buds were taken from another tree and examined once a week, but in spite of the most careful searching no eggs were discovered after October.

The previous experiments ^a on the destruction of this pest, although very successful, were not of the nature that a fruit grower could apply at a cost which would repay him for the extra labor involved, although certain growers have given the spray fluid mentioned in my earlier report a trial with very satisfactory results.

The two main objections to the soap and sulphur spray fluid were, firstly, the large number of applications which were given in my

^a Report on Economic Zoology, No. 1, 1904.

experiments, and, secondly, the large quantity of soap used. But once having proved that the mite could be destroyed by the use of sulphur, it remained to be demonstrated in what form it was best applied and whether or not it could be shown that any benefit resulted from a smaller number of applications.

With these two objects in view a series of experiments was commenced early in 1905 on a piece of land set apart and prepared for the purpose by the council of the University of Birmingham.

On this plat seven rows of bushes were planted, consisting of Black Naples, Baldwins, and Boskoop Giant. All were as badly affected with "big bud" as any that could be obtained; indeed, I have never seen worse. The plat of land was far from an ideal one and the last that a fruit grower would have chosen, so that the bushes had no natural advantages in their favor.

The experiments carried out were as follows: Rows 1, 2, and 3 were dusted with equal parts of unslaked lime and flowers of sulphur. Rows 4, 5, and 6 were sprayed with a mixture consisting of 1 pound of lime, 1 pound of sulphur, and 20 gallons of water, while row 7 was sprayed with a mixture consisting of 1 pound of sulphur, 1 pound of soft soap, and 20 gallons of water.

Summary.—It is very evident that all the bushes benefited by the application of lime and sulphur. It would have been better, however, had a little less lime been used (1 part of lime to 2 parts of sulphur has acted as well). In the case of those bushes that received a single dusting, the big buds were considerably reduced in number, not more than one-fourth of the number being present in October as there were in February. Where two dustings were given, a distinctly marked diminution over those receiving one dusting was shown, while where three dustings were applied the mite was almost exterminated. It must be borne in mind that neither a spray fluid nor dry application will reach the eggs in the buds, and it seems clear that the number of adult mites which successfully migrated from the old buds into the new ones was very small indeed. In all cases, the mites found were immature specimens.

Spraying with lime and sulphur.—The results obtained by spraying were not so good as those by dusting. A large number of buds were affected, and in many of them there were adult and immature mites and eggs. The differences between the one, two, and three applications of the spray fluid were quite in keeping with those found to obtain where dusting had been done. The fewer the applications the more buds affected and the greater the number of mites.

Spraying with soft soap and sulphur.—Row 7 was sprayed twice with the spray fluid above mentioned. When the bushes were examined in October a fairly large number of big buds was noticed:

there were, however, not nearly so many as in the previous February; roughly estimated, I should think about one-third the number. In many of these buds 12 to 20 adult mites were present, many immature specimens, and a few eggs. The result of an examination of 86 suspected buds gave an average of 4 adults, 9 immature mites, and 3 eggs per bud.

SUMMARY AND CONCLUSION.

After the experiments which have been made I feel convinced that the application of lime and sulphur will keep this mite in check, and, if the dusting and spraying is continued, will eventually entirely eradicate the pest.

Various statements have appeared in a section of the horticultural and agricultural press, stating that there is no likelihood of a cure, or even of means whereby the mite can be kept in check; and, further, that its life history is very imperfectly understood. I would warn all fruit growers against such misleading statements. The life history is now practically fully known, and the experiments which I have conducted and which have now extended over eight years have yielded results, checked by many large fruit growers, which clearly point to the fact that the application of lime and sulphur offers an effective remedy.

It is interesting to note in this particular that the destructive rust mite of the orange and lemon has been combated in a similar manner. In 1889, according to Mr. Marlatt,^a large quantities of citrus trees were obtained from Florida, and a species of *Eriophyes* (*E. oleivorus* Ashmead) was undoubtedly introduced in the Rivera and San Diego Bay districts of California, where it did considerable harm in the orange and lemon groves. Mr. Marlatt states that "an estimate made from actual count, indicates that the mites and the eggs on a single leaf in midwinter may reach the enormous number of 75,000," indicating some billions of mites for each tree in the active breeding season. He further states that it "is readily destroyed by various insecticides. The eggs, however, are much more difficult to kill, and practically no wash can be relied upon to reach and destroy all the eggs of this mite. * * * The sovereign remedy for the rust mite is sulphur. * * * The advantage of the sulphur treatment arises from the fact that the sulphur adheres to the leaves and the young mites are killed as soon as they come in contact with it."

A further example of treating another species of the same genus of mites is offered in the case of the cotton-leaf blister mite (*E. gossypii* Banks.), which made its appearance in the West Indian cotton fields in 1903^b and spread quickly throughout the islands. Here the lime-

^a U. S. Dept. Agric., Farmers' Bul. No. 172, 1903, pp. 38-41.

^b West Indian Bul., 1903, Vol. IV, pp. 282 and 336.

and-sulphur treatment has proved most effective. Other species of Eriophyes have been treated with lime and sulphur, viz. *E. arellæna* Nal., *E. rudis* Canest., and *E. taxi* Murray, with equally satisfactory results.

DESTRUCTION OF MOSQUITOES IN DWELLINGS BY THE POWDERS OF CHRYSANTHEMUM, SPREAD THEREIN BY MEANS OF HAND BELLOWS OR A TOWEL.

By A. L. HERRERA, *Mexico City, Mexico.*

Ever since 1903 this commission has been commending the destruction of mosquitoes in dwellings by means of the chrysanthemum powders, spread or thrown around therein with hand bellows or a towel.^a

ESSENTIAL RECOMMENDATIONS.

(1) In each room or apartment there should be spread daily, half an hour or an hour before bedtime, the genuine chrysanthemum powders, by means of some hand bellows, cloth, or towel.

(2) The powders should be spread uniformly in the whole room, and to that effect they should be scattered with the bellows as high as possible, in various directions, care being taken besides that they penetrate under the beds and other pieces of furniture, behind the doors, etc. In case a towel is used, it should be shaken around in a very lively manner and for some time. Some people are wont to place the powders upon a table or a piece of pasteboard and blow said powders upwardly, and at once shake a bed sheet or a towel in the center of the room.

(3) It is convenient to employ a large spoonful of powder for every room of 20 to 30 cubic meters, and a larger quantity for larger apartments.

In order to find out whether the quantity of powder that has been used is sufficient, it will suffice to spread some bed sheets upon the floor, so that the mosquitoes may fall upon them. Some of the insects should then be kept under a glass to see, on the next day, if they are really dead. Should it not be so, or should no mosquito fall upon the sheets, after an hour, even though they should be in great numbers in the room, the quantity of powder scattered should be increased or a new lot should be bought in another drug store. The same should be done when the insects have only fallen into lethargy at the beginning of the night, and again become troublesome in the morning.

(4) To avoid inflammation of the throat the person who has to scatter the powder should cover the mouth and a part of the face with a handkerchief and leave the room as soon as possible.

^a Boletín de la Comisión de Parasitología Agrícola, Tomo II, pag. 139.

(5) The insecticide powder should be spread in the room before bedtime, it being possible for the people to enter the room after said powder has settled down.

(6) There are some kinds of the powders which are entirely inefficient; this is true of most of the powders sold by retailers in small quantities, in envelopes or in boxes, at 5 cents or less. To obtain good ones they should be bought at some reliable drug store and tried in the way above explained.

(7) In order to avoid a daily expense for powder, it is advisable to close the doors and windows early, so that the mosquitoes will not enter the apartments, and it will even be very useful to put a fine wire screen upon the windows. Should any mosquitoes, however, succeed in entering the rooms, they will be killed by the powder of chrysanthemum, an operation to be performed only when the insects happen accidentally to enter on account of the openings having been shut too late or because of any other circumstance.

(8) An excess of powder is quite unnecessary, and might even result in injury to the inhabitants.

NOTE.—It has been said that the fine powder, in large quantities, may become ignited when there is a candle in the room, but it has never been proved that such is the case when only genuine chrysanthemum powders are used. In a fumigating room made of canvas, with a capacity of 68 cubic decimeters, we blew a large quantity of genuine chrysanthemum powder, a part of which fell upon the flame of an alcohol lamp situated in the back of the room. The only powder that burned was that which touched the flame directly, whereby was produced a kind of rain of small sparks; but the fire was never communicated to the remainder of the powder, and no explosion occurred. This experiment may be made on a small scale with a candle, and thus the above statement will be confirmed, while at the same time it will be found out whether the powder contains any foreign matter which is inflammable and dangerous. Where electric light is used such a precaution is unnecessary.

This manner of destroying mosquitoes has already been put into practice. To that effect the commission of parasitology has distributed gratuitously several barrels of chrysanthemum powders under the form of small samples. The consumption of powders has been tripled, while the sale of the tablets, which only throw the insects into a state of lethargy but are very objectionable owing to their pungent fumes, was decreased.

The species of mosquito which is most frequently found in the City of Mexico and which invades the houses is *Culex pipiens* L.

NOTES ON INSECTS IN CENTRAL ALBERTA.

By P. B. GREGSON, *Blackfalds, Alberta, Canada.*

With so large and ever increasing immigration from the United States, a few notes from the grain section of central Alberta may not be without interest. The early season of 1906 in this district was notable for the phenomenal outbreak of "cutworms," chiefly *Noctua clandestina* Harr., *Chorizagrotis auxiliaris* Grt., and *Paragrotis ochrogaster* Guen.—extraordinary because climatic conditions had not seemed to warrant such an outbreak.

The preceding year (1905) was normal; the fall dry, the November mean minimum temperature being 20.15° F., snow 5.85 inches; and December mean minimum 8.30° F., snow 0.75 inch. The mean minimum temperature for January, 1906, was -0.70° F., and snow 4.40 inches. Hot suns on February 1, 2, and 3 dispelled nearly all of what little snow there was, and from the middle of February to the end of March was characterized by cold snowless weather, touching -19° F. on March 12, with all fields bare of snow, and being in fact the driest season for ten years. The total moisture from November, 1905, to May 16, 1906 (from snow and rain combined), did not exceed two-thirds of an inch. Ninety per cent of the local winter wheat was killed off. The total precipitation for the month of April and till May 16 was only 0.115 or 0.1 of an inch of moisture, a very warm and dry period, the mean maximum shade temperature for April being 60.68° F. and for May (up to the 18th) 66.10° F—a temperature above the average for ten years.

Studying these weather conditions—the reverse of a cold, wet spring—it would seem that parasites would at least have an equal chance with cutworms for surviving. But what was the result? The outbreak of cutworms was without precedent, and of parasites few could be discovered.

Every kind of vegetation seemed to be attacked by the cutworms. Among the instances of which the writer made special observation a few may be mentioned as showing the catholic nature of their food. A nursery gardener had planted in the fall of 1905 (in a brome-grass field which had been plowed up in the previous spring) an orchard of several hundred gooseberry, currant (red, white, and black), raspberry, and blackberry bushes and some hundreds of strawberry and rhubarb plants. In the spring of 1906 the young shoots of the raspberry and blackberry bushes were persistently cut through just below the soil and every bush died. The runners of the gooseberry and currant bushes shared the same fate, even the leaves being cleared off. Not one strawberry survived the attack, and the rhubarb also suffered severely. Circular pits had been dug round each bush in the

fall, but did not operate as any check to the pest. Several farmers lost their entire crops of potatoes, the caterpillars killing off each shoot from the tuber. Of grain farmers, many lost from 40 to 80 per cent of oats and spring wheat; one crop visited by the writer was sown with wheat a second time and again entirely destroyed. (It was sown a third time, but too late to harvest, though it escaped the cutworms.) And so with horticulturists: Peonies, columbines, pansies, foxgloves (*Digitalis*), wall flowers, candytuft, campanulæ—all were attacked indiscriminately by *Chorizagrotis auxiliaris* and *Noctua clandestina*, even two young Charles XII lilac bushes being killed by having their young buds devoured (this by *Peridroma occulta* L.) Carrots, parsnips, onions, cabbages, and other Cruciferæ grown in gardens were in many cases "cleaned out" by *Noctua clandestina*.

The magnitude of the attack seemed to discourage the farmers. Remedies were tried without much success, except in gardens where close attention could be given. Where tried in gardens poisoned bran succeeded very well, but farmers seem not to have the time or the pains for applying the poison-bran remedy to their large grain fields.

As a wholesale and very fairly efficient remedy for grain fields the writer recommended heavy rolling late in the evening. This was tried by one farmer whose field was infested, the rolling being done from 10.30 p. m. till about midnight, being repeated a time or two during the young growth of grain, and resulted not only in the virtual salvation of his crop, but in a strong stand of straw (with the summer the driest on record).

In short, the farmer should, in the opinion of the writer, consistently practice the following method with his grain:

(1) Late fall plowing (plowing can often be done in Alberta till the early days of November).

(2) Drilling the grain (in spring), not sowing broadcast. Drilling facilitates the operation of the harrow and seeds deeper than broadcast sowing.

(3) Roll in evening (10.30 p. m.) when grain is appearing and soil dry.

(4) Harrow when grain is a little more advanced.

(5) Roll once more in evening (on dry soil).

(6) Harrow finally (with grain about 4 inches high).

The writer has known crops to be harrowed three times after the first appearance of grain, with most beneficial results, and, except in years of unusual prevalence of cutworms, the second rolling might be dispensed with.

It was moved and carried that the Chair appoint a committee of five to take up the work outlined in the report of the committee on testing proprietary insecticides.

The following were appointed: Messrs. E. P. Felt, C. P. Gillette, E. D. Sanderson, R. I. Smith, and H. E. Summers.

A motion was made and carried that the next annual meeting of the association be held at the same time and place as the meeting of the American Association for the Advancement of Science, the exact dates to be fixed by the officers.

The report of the committee on nominations was presented, and is as follows:

REPORT OF COMMITTEE ON NOMINATIONS.

The nominating committee respectfully submit the following list for officers of the association for the ensuing year:

For president, H. A. Morgan, Knoxville, Tenn.

For first vice-president, H. E. Summers, Ames, Iowa.

For second vice-president, W. D. Hunter, Washington, D. C.

For secretary-treasurer, A. F. Burgess, Columbus, Ohio.

For member of joint committee on legislation, Wilmon Newell, Baton Rouge, La.

For member of standing committee on nomenclature, E. S. G. Titus, Washington, D. C.

For members of council, James Fletcher, Ottawa, Canada, and S. A. Forbes, Urbana, Ill.

Respectfully submitted,

C. L. MARLATT.

H. OSBORN.

F. M. WEBSTER.

On motion, the secretary was instructed to cast a ballot of the association for those mentioned, and they were declared elected.

The committee on membership submitted the following report:

REPORT OF COMMITTEE ON MEMBERSHIP.

The committee report that it has followed the plan of the last committee on membership, which was adopted by the association. We desire to reiterate the plan proposed and adopted at the last meeting, namely:

"That the association exercise greater care as to the qualifications of candidates for election to membership, and that a considerably more conservative policy than has prevailed should be adopted. In general, only those who are already associate members should be elected to active membership; and this should occur only after they have published a considerable amount of original matter on economic entomology, based on their own independent investigations. The privilege of associate members, however, may well be extended as a means of encouragement to young men who are expecting to pursue economic entomology as a profession, but who have not yet had time to show their capability by publication or otherwise."

In accordance with these principles, the committee make the following recommendations:

(1) For foreign membership: Dr. F. Silvestri, head of entomological department, Royal School of Agriculture, Portici, Italy.

(2) For active membership: Prof. J. Troop, State entomologist, La Fayette, Ind.

(3) For transfer from associate to active membership: Dr. E. F. Phillips, Bureau of Entomology, Washington, D. C.; J. G. Sanders, Bureau of Entomology, Washington, D. C.; H. E. Burke, Bureau of Entomology, Washington, D. C.; W. A. Hooker, Bureau of Entomology, Washington, D. C.; A. F. Conradi, State entomologist, College Station, Tex.

(4) For associate members: William Beutenmuller, American Museum of Natural History, New York, N. Y.; I. J. Condit, Bureau of Entomology, Washington, D. C.; R. A. Cushman, Bureau of Entomology, Washington, D. C.; H. M. Russell, Amherst, Mass.; W. V. Tower, entomologist Porto Rico Experiment Station, Mayaguez, P. R.; H. J. Franklin, Amherst, Mass.; E. A. Back, Amherst, Mass.; E. F. Hitchings, State entomologist, Augusta, Me.; Fred E. Brooks, Morgantown, W. Va.; D. K. McMillan, Harrisburg, Pa.; H. E. Hodgkiss, Geneva, N. Y.; W. J. Schoene, Geneva, N. Y.; G. P. Weldon, College Park, Md.; Dr. T. J. Headlee, Durham, N. H.; C. N. Ainslie, Department of Agriculture, Washington, D. C.; John A. Grossbeck, New Brunswick, N. J.; R. L. Webster, St. Anthony Park, Minn.; A. G. Ruggles, St. Anthony Park, Minn.; R. S. Woglum, Raleigh, N. C.; Dudley Moulton, Department of Agriculture, Washington, D. C.; Prof. T. D. Jarvis, Guelph, Ontario, Canada.

(5) That the following be dropped, since they have discontinued entomological work: J. H. Beattie; E. E. Bogue.

And that the resignation of Prof. J. H. Perkins be accepted.

(6) For transfer from active to associate membership: Prof. C. M. Weed, Lowell, Mass.

That a committee of three, including the president and secretary, be empowered to devise and have executed a suitable form of certificate of membership for the foreign members, and that the secretary be authorized to send the certificates to the entomologists concerned.

The committee also recommend that inasmuch as there is at present considerable confusion, owing to the various amendments to the constitution, the secretary be directed to codify the constitution and by-laws and present them to the association at the next meeting.

W. D. HUNTER.

H. E. SUMMERS.

JOHN B. SMITH.

On motion, it was accepted.

The report of the committee on resolutions was as follows:

REPORT OF COMMITTEE ON RESOLUTIONS.

Resolved, That the Association of Economic Entomologists expresses its appreciation for the courtesies extended by the president of Columbia University, the trustees of the American Museum of Natural History, and the council of the New York Academy of Science, and the successful efforts of the local committee of the A. A. A. S. for its entertainment; and

Resolved, That we express our gratitude to the entomological societies of New York, Brooklyn, and Newark for the most enjoyable reception tendered by them; and

Resolved, That the association, through its secretary, tender its thanks to the Honorable Secretary of Agriculture for the publication of its proceedings, and request that the proceedings of this meeting be published in like manner; and

Resolved, That this association places itself on record as indorsing the policy of the National Government in aiding the suppression of introduced pests of a serious nature, and asks for a continuation and increase of the Congressional appropriations to the Bureau of Entomology for such work and for the introduction of natural enemies of such pests; and

Resolved, That this association has learned with great interest, through the address of President Kirkland, of the attempt now being made in Massachusetts and other New England States to control the gipsy and brown-tail moths through systematic cooperation and liberal financial provision made for prosecuting field operations and for the introduction of parasites of these insects; and, being convinced that the successful prosecution of this work will protect the entire country from a serious menace, this effort has the unqualified indorsement of this association; and

Resolved, That this association extend its cordial greetings to the newly formed Entomological Society of America, and that the programme committee of this association be instructed to arrange its programme conjointly with the programme committee of that society.

JAMES FLETCHER.

M. V. SLINGERLAND.

E. D. SANDERSON.

At the request of the representatives of the American Nurserymen's Association, the following committee was appointed by the president to attend the annual meeting of that association at Detroit in June, 1907: Messrs. H. A. Morgan, Wilmon Newell, S. A. Forbes, A. F. Burgess, and E. D. Sanderson.

The committee on certificate of membership for foreign members was appointed, consisting of the president elect, the secretary, and Mr. W. E. Britton.

President Kirkland thanked the association for the honor of presiding at this meeting, which was the largest in its history, and, as there was no further business, the meeting was then adjourned.

A. F. BURGESS, *Secretary*.

LIST OF MEMBERS OF THE ASSOCIATION OF ECONOMIC
ENTOMOLOGISTS.

ACTIVE MEMBERS.

Aldrich, J. M., Agricultural Experiment Station, Moscow, Idaho.
Alwood, William B., Charlottesville, Va.
Ashmead, William H., U. S. National Museum, Washington, D. C.
Baker, C. F., Agricultural Experiment Station, Santiago de las Vegas, Cuba.
Ball, E. D., Agricultural Experiment Station, Logan, Utah.
Banks, C. S., Manila, P. I.
Banks, Nathan, U. S. Department of Agriculture, Washington, D. C.
Bethune, C. J. S., Guelph, Ontario, Canada.
Benton, Frank, 925 N street N. W., Washington, D. C.
Bishopp, F. C., U. S. Department of Agriculture, Washington, D. C.
Britton, W. E., New Haven, Conn.
Bruner, Lawrence, Agricultural Experiment Station, Lincoln, Nebr.
Burgess, Albert F., U. S. Department of Agriculture, Washington, D. C.
Burke, H. E., U. S. Department of Agriculture, Washington, D. C.
Busck, August, U. S. Department of Agriculture, Washington, D. C.
Caudell, A. N., U. S. Department of Agriculture, Washington, D. C.
Chambliss, C. E., Clemson College, S. C.
Chittenden, F. H., U. S. Department of Agriculture, Washington, D. C.
Cockerell, T. D. A., University of Colorado, Boulder, Colo.
Comstock, J. H., Cornell University, Ithaca, N. Y.
Conradi, A. F., College Station, Tex.
Cook, A. J., Pomona College, Claremont, Cal.
Cook, Mel T., Newark, Del.
Cooley, R. A., Agricultural Experiment Station, Bozeman, Mont.
Coquillett, D. W., U. S. Department of Agriculture, Washington, D. C.
Cordley, A. B., Agricultural Experiment Station, Corvallis, Oregon.
Crawford, J. C., U. S. Department of Agriculture, Washington, D. C.
Dickerson, Edgar L., Agricultural Experiment Station, New Brunswick, N. J.
Dyar, H. G., U. S. National Museum, Washington, D. C.
Ehrhorn, E. M., Room 611, Ferry Building, San Francisco, Cal.
Felt, E. P., Geological Hall, Albany, N. Y.
Fernald, C. H., Agricultural College, Amherst, Mass.
Fernald, H. T., Agricultural College, Amherst, Mass.
Fiske, W. F., U. S. Department of Agriculture, Washington, D. C.
Fletcher, James, Central Experimental Farm, Ottawa, Canada.
Forbes, S. A., University of Illinois, Urbana, Ill.
French, G. H., Normal avenue, Carbondale, Ill.
Garman, H., Agricultural Experiment Station, Lexington, Ky.
Gibson, Arthur, Central Experimental Farm, Ottawa, Canada.
Gillette, C. P., Agricultural Experiment Station, Fort Collins, Colo.
Girault, A. A., U. S. Department of Agriculture, Washington, D. C.
Gossard, H. A., Agricultural Experiment Station, Wooster, Ohio.
Gregson, P. B., Blackfalds, Alberta, Northwest Territory, Canada.
Hart, C. A., Illinois State Laboratory of Natural History, Urbana, Ill.

- Heidemann, Otto, U. S. Department of Agriculture, Washington, D. C.
Hinds, W. E., U. S. Department of Agriculture, Washington, D. C.
Hine, J. S., Ohio State University, Columbus, Ohio.
Holland, W. J., Carnegie Museum, Pittsburg, Pa.
Hooker, W. A., U. S. Department of Agriculture, Washington, D. C.
Hopkins, A. D., U. S. Department of Agriculture, Washington, D. C.
Houghton, C. O., Agricultural Experiment Station, Newark, Del.
Howard, L. O., U. S. Department of Agriculture, Washington, D. C.
Hunter, S. J., University of Kansas, Lawrence, Kans.
Hunter, W. D., U. S. Department of Agriculture, Washington, D. C.
Johnson, S. Arthur, State Agricultural College, Fort Collins, Colo.
Kellogg, Vernon L., Stanford University, Cal.
Kincaid, Trevor, University of Washington, Seattle, Wash.
Kirkland, A. H., 6 Beacon street, Boston, Mass.
Kotinsky, J., Honolulu, Hawaii.
Lochhead, William, Macdonald College of Agriculture, Montreal, Canada.
Marlatt, C. L., U. S. Department of Agriculture, Washington, D. C.
Morgan, H. A., University of Tennessee, Knoxville, Tenn.
Morrill, A. W., U. S. Department of Agriculture, Washington, D. C.
Murtfeldt, Miss M. E., Kirkwood, Mo.
Newell, Wilmon, La. Crop Pest Comm., Baton Rouge, La.
Osborn, Herbert, Ohio State University, Columbus, Ohio.
Parrott, P. J., Geneva, N. Y.
Pergande, Theodore, U. S. Department of Agriculture, Washington, D. C.
Pettit, R. H., Agricultural Experiment Station, Agricultural College, Mich.
Phillips, E. F., U. S. Department of Agriculture, Washington, D. C.
Phillips, J. L., Agricultural Experiment Station, Blacksburg, Va.
Phillips, W. J., U. S. Department of Agriculture, Washington, D. C.
Pierce, W. Dwight, U. S. Department of Agriculture, Washington, D. C.
Popenoe, E. A., R. F. D. No. 2, Topeka, Kans.
Pratt, F. C., U. S. Department of Agriculture, Washington, D. C.
Quaintance, A. L., U. S. Department of Agriculture, Washington, D. C.
Quayle, H. J., Agricultural Experiment Station, Berkeley, Cal.
Reeves, George I., U. S. Department of Agriculture, Washington, D. C.
Rumsey, W. E., Agricultural Experiment Station, Morgantown, W. Va.
Sanborn, C. E., College Station, Tex.
Sanders, J. G., U. S. Department of Agriculture, Washington, D. C.
Sanderson, E. Dwight, Agricultural Experiment Station, Durham, N. H.
Saunders, William, Central Experimental Farm, Ottawa, Canada.
Schwarz, E. A., U. S. Department of Agriculture, Washington, D. C.
Sherman, Franklin, jr., Div. of Entom., State Dept. of Agric., Raleigh, N. C.
Sirrinc, F. A., 124 Sound avenue, Riverhead, N. Y.
Skinner, Henry, Academy of Natural Sciences, Philadelphia, Pa.
Slingerland, M. V., Agricultural Experiment Station, Ithaca, N. Y.
Smith, J. B., Agricultural Experiment Station, New Brunswick, N. J.
Smith, R. I., Atlanta, Ga.
Snow, F. H., Lawrence, Kans.
Stedman, J. M., Agricultural Experiment Station, Columbia, Mo.
Summers, H. E., Agricultural Experiment Station, Ames, Iowa.
Surface, H. A., State Zoologist, Harrisburg, Pa.
Symons, T. B., Agricultural Experiment Station, College Park, Md.
Taylor, E. P., University of Illinois, Urbana, Ill.
Titus, E. S. G., Agricultural Experiment Station, Logan, Utah.
Troop, J., La Fayette, Ind.

- Van Dine, D. L., Government Entomologist, Hawaiian Exp. Sta., Honolulu, Hawaii.
- Viereck, H. L., Academy of Natural Sciences, Philadelphia, Pa.
- Walden, B. H., Agricultural Experiment Station, New Haven, Conn.
- Walker, C. M., Amherst, Mass.
- Washburn, F. L., Agricultural Experiment Station, St. Anthony Park, Minn.
- Webster, F. M., U. S. Department of Agriculture, Washington, D. C.
- Wilcox, E. V., U. S. Department of Agriculture, Washington, D. C.
- Woodworth, C. W., Agricultural Experiment Station, Berkeley, Cal.

ASSOCIATE MEMBERS.

- Adams, C. F., Fayetteville, Ark.
- Ainslie, C. N., U. S. Department of Agriculture, Washington, D. C.
- Back, E. A., Orlando, Fla.
- Beutenmüller, Wm., American Museum of Natural History, New York, N. Y.
- Barber, H. S., U. S. National Museum, Washington, D. C.
- Bartholomew, C. E., Iowa State College, Ames, Iowa.
- Beckwith, H. M., Elmira, N. Y.
- Bentley, Gordon M., University of Tennessee, Knoxville, Tenn.
- Braucher, R. W., Lincoln Park, Chicago, Ill.
- Brooks, Fred E., Morgantown, W. Va.
- Brues, C. T., Milwaukee Public Museum, Milwaukee, Wis.
- Bullard, W. S., 629 Water street, Bridgeport, Conn.
- Campbell, J. P., Athens, Ga.
- Clifton, R. S., U. S. Department of Agriculture, Washington, D. C.
- Condit, Ira J., California Polytechnic School, San Luis Obispo, Cal.
- Cotton, Edwin C., Ohio State Department of Agriculture, Columbus, Ohio.
- Couden, F. D., U. S. Department of Agriculture, Washington, D. C.
- Craw, Alexander, Board of Agriculture, Honolulu, Hawaii.
- Currie, Rolla P., U. S. Department of Agriculture, Washington, D. C.
- Cushman, R. A., U. S. Department of Agriculture, Washington, D. C.
- Dean, Harper, jr., State Board of Entomology, Atlanta, Ga.
- Doran, E. W., Belhaven College, Jackson, Miss.
- Engle, Enos B., Department of Agriculture, Harrisburg, Pa.
- Flynn, C. W., Assistant Entomologist, Louisiana Crop Pest Commission, Baton Rouge, La.
- Fowler, Carroll, Duarte, Cal.
- Franklin, H. J., Amherst, Mass.
- Frost, H. L., 6 Beacon street, Boston, Mass.
- Gahan, A. B., College Park, Md.
- Garrett, J. B., Assistant Entomologist, Louisiana Crop Pest Commission, Baton Rouge, La.
- Gifford, John, Princeton, N. J.
- Gould, H. P., U. S. Department of Agriculture, Washington, D. C.
- Green, E. C., College Station, Tex.
- Grossbeck, John A., New Brunswick, N. J.
- Hardy, E. S., Assistant Entomologist, Louisiana Crop Pest Commission, Baton Rouge, La.
- Hargitt, C. W., Syracuse University, Syracuse, N. Y.
- Harrington, W. H., Post-Office Department, Ottawa, Canada.
- Headlee, T. J., Durham, N. H.
- Hitchings, E. F., Waterville, Me.
- Hodgkiss, H. E., Geneva, N. Y.

- Houser, J. S., Assistant Entomologist, Ohio Agricultural Experiment Station, Wooster, Ohio.
- Hudson, G. H., State Normal and Training School, Plattsburg, N. Y.
- Isaac, John, Sacramento, Cal.
- Jarvis, T. D., Guelph, Ontario, Canada.
- Johnson, Fred, U. S. Department of Agriculture, Washington, D. C.
- Johnson, W. G., 52 Lafayette place, New York, N. Y.
- Jones, Charles R., U. S. Department of Agriculture, Washington, D. C.
- King, George B., Lawrence, Mass.
- Kirkaldy, G. W., Hawaiian Sugar Planters' Experiment Station, Honolulu, Hawaii.
- Koebele, Albert, Alameda, Cal.
- Mackintosh, R. S., State Board of Horticulture, Auburn, Ala.
- Mann, B. P., 1918 Sunderland place, Washington, D. C.
- Martin, Leslie, Lebanon, Tenn.
- Martin, George W., 1804 Grand avenue, Nashville, Tenn.
- Martin, W. O., Assistant Entomologist, Louisiana Crop Pest Commission, Baton Rouge, La.
- MacGillivray, A. D., Cornell University, Ithaca, N. Y.
- McCarthy, Gerald, Department of Agriculture, Raleigh, N. C.
- McMillan, D. K., Harrisburg, Pa.
- Morgan, A. C., U. S. Department of Agriculture, Washington, D. C.
- Mosher, F. H., Wakefield, R. F. D., Mass.
- Moulton, Dudley, U. S. Department of Agriculture, Washington, D. C.
- Nicholson, John F., Stillwater, Okla.
- Niswander, F. J., 519 East Seventeenth street, Cheyenne, Wyo.
- Palmer, R. M., Victoria, British Columbia.
- Perkins, R. C. L., Hawaiian Sugar Planters' Experiment Station, Honolulu, Hawaii.
- Piper, C. V., U. S. Department of Agriculture, Washington, D. C.
- Price, H. L., Agricultural Experiment Station, Blacksburg, Va.
- Randall, J. L., Durham, N. H.
- Rane, F. W., Agricultural Experiment Station, Durham, N. H.
- Rankin, John M., U. S. Department of Agriculture, Washington, D. C.
- Reed, E. B., Esquimault, British Columbia.
- Riley, W. A., Cornell University, Ithaca, N. Y.
- Rolfs, P. H., Miami, Fla.
- Ruggles, A. G., St. Anthony Park, Minn.
- Russell, H. M., U. S. Department of Agriculture, Washington, D. C.
- Sasscer, Ernest R., U. S. Department of Agriculture, Washington, D. C.
- Schoene, W. J., Geneva, N. Y.
- Scott, W. M., U. S. Department of Agriculture, Washington, D. C.
- Southwick, E. B., Arsenal Building, Central Park, New York, N. Y.
- Stimson, James, Santa Cruz, Cal.
- Swezey, O. H., Hawaiian Sugar Planters' Experiment Station, Honolulu, Hawaii.
- Thaxter, Roland, 7 Scott street, Cambridge, Mass.
- Toumey, J. W., Yale Forest School, New Haven, Conn.
- Tower, W. L., Porto Rico Experiment Station, Mayaguez, P. R.
- Townsend, C. H. T., U. S. Department of Agriculture, Washington, D. C.
- Webb, J. L., U. S. Department of Agriculture, Washington, D. C.
- Webster, R. L., St. Anthony Park, Minn.
- Weed, C. M., Lowell, Mass.
- Weldon, G. P., College Park, Md.

Woglum, R. S., U. S. Department of Agriculture, Washington, D. C.

Yothers, W. W., U. S. Department of Agriculture, Washington, D. C.

Young, D. B., Geological Hall, Albany, N. Y.

FOREIGN MEMBERS.

Ballou, H. A., Imperial Department of Agriculture, Barbados, West Indies.

Berlese, Dr. Antonio, Reale Stazione di Entomologia Agraria, Firenze, Italy.

Bordage, Edmond, Directeur de Musée, St. Denis, Reunion.

Carpenter, Dr. George H., Royal College of Science, Dublin, Ireland.

Cholodkosky, Prof. Dr. N., Militär-Medicinische Akademie, St. Petersburg, Russia.

Collinge, W. E., 55 Newhall street, Birmingham, England.

Danysz, J., Laboratoire de Parasitologie, Bourse de Commerce, Paris, France.

Enock, Fred, 42 Salisbury road, Bexley, London, SE., England.

French, Charles, Department of Agriculture, Melbourne, Australia.

Froggatt, W. W., Department of Agriculture, Sydney, New South Wales.

Fuller, Claude, Department of Agriculture, Pietermaritzburg, Natal, South Africa.

Giard, A., 14 Rue Stanislaus, Paris, France.

Goding, F. W., Newcastle, New South Wales.

Grasby, W. C., 6 West Australian Chambers, Perth, West Australia.

Green, E. E., Royal Botanic Gardens, Peradeniya, Ceylon.

Helms, Richard, 136 George street, North Sydney, New South Wales.

Herrera, A. L., Calle de Betlemitas No. 8, Mexico City, Mexico.

Horvath, Dr. G., Musée Nationale Hongroise, Budapest, Hungary.

Jablonowski, Josef, Entomological Station, Budapest, Hungary.

Lampa, Prof. Sven, Statens Entomologiska, Anstalt, Stockholm, Sweden.

Lea, A. M., Department of Agriculture, Hobart, Tasmania.

Leonardi, Gustavo, R. Scuola di Agricoltura, Portici, Italy.

Lounsbury, Charles P., Department of Agriculture, Cape Town, South Africa.

Mally, C. W., Department of Agriculture, Columbus, Ohio (formerly Grahams-town, Cape Colony).

Marchal, Dr. Paul, 16 Rue Claude Bernard, Paris, France.

Mussen, Charles T., Hawkesbury Agricultural College, Richmond, New South Wales.

Nawa, Yashushi, Entomological Laboratory, Kyomachi, Gifu, Japan.

Newstead, Robert, University School of Tropical Medicine, Liverpool, England.

Peal, H. W., Indian Museum, Calcutta, India.

Porchinski, Prof. A., Ministère de l'Agriculture, St. Petersburg, Russia.

Reed, E. C., Museo, Concepcion, Chile.

Reuter, Dr. Enzo, Agrikultur-Economiska Försöksanstalten, Helsingfors, Finland.

Ritzema Bos, Dr. J., Agricultural College, Wageningen, Netherlands.

Sajo, Prof. Karl, Gödöllő-Veresegyház, Hungary.

Schoyen, Prof. W. M., Zoological Museum, Christiania, Norway.

Shiple, Prof. Arthur E., Christ's College, Cambridge, England.

Silvestri, Dr. F., R. Scuola Superiore di Agricoltura, Portici, Italy.

Tepper, J. G. O., Norwood, South Australia.

Theobald, Frederick V., Wye Court, Wye, Kent, England.

Thompson, Rev. Edward H., Franklin, Tasmania.

Tryon, H., Queensland Museum, Brisbane, Queensland, Australia.

Urich, F. W., Victoria Institute, Port of Spain, Trinidad, West Indies.

Vermorel, V., Station Viticole, Villefranche, Rhone, France.

INDEX.

	Page.
Adams Act and economic entomology.....	77-84
<i>Agonoderus pallipes</i> , enemy of cabbage maggot.....	15
<i>Alabama argillacea</i> on cotton in Georgia.....	101
Alberta, central, notes on insects.....	125-126
Alfalfa, food plant of <i>Melanoplus differentialis</i>	98
Almond, food plant of <i>Aspidiotus howardii</i>	88
<i>Anara impuncticollis</i> , enemy of cabbage maggot.....	15
<i>Amelanchier vulgaris</i> , food plant of <i>Eriophyes pyri</i>	44
<i>Amphicerus bicaudatus</i> = <i>Schistoceros hamatus</i>	11
<i>Anarsia lineatella</i> , arsenate of lead a remedy.....	47
in Georgia.....	103
Anthraxnose following <i>Heliothis obsoleta</i> in cotton.....	101
Ant, little red, work against it in Minnesota.....	16-17
small red, enemy of <i>Carpocapsa pomonella</i>	38
sugar, ineffective against little red ant.....	17
Aphides, is honeydew excreted from anus or cornicles?.....	33-34
of apple in New York State.....	29
Aphididae, manner of birth.....	31-33
Aphis, apple. (See <i>Aphis mali</i> .)	
cotton. (See <i>Aphis gossypii</i> .)	
eggs, effects of sprays.....	29-31
<i>gossypii</i> on cotton in Georgia.....	101
<i>mali</i> , effects of sprays on eggs.....	29-30
in Georgia.....	104
New York State.....	29
manner of birth.....	33
<i>malifoliae</i> , effects of sprays on eggs.....	29-30
in New York State.....	29
<i>persicae-niger</i> in Georgia.....	103
woolly (see also <i>Schizoneura lanigera</i>).	
kerosene emulsion a remedy.....	106
Apple aphides in New York State.....	29
aphis. (See <i>Aphis mali</i> .)	
calyx cup in relation to spraying.....	70-72
caterpillar, red-humped. (See <i>Schizura concinna</i> .)	
yellow-necked. (See <i>Datana ministra</i> .)	
food plant of <i>Aphis mali</i>	104
<i>Aspidiotus howardii</i>	88, 93
<i>Chrysobothris femorata</i>	104
<i>Eriophyes pyri</i>	44, 45
gipsy moth.....	23
<i>Saperda candida</i>	104
<i>Schizoneura lanigera</i>	104
<i>Aramigus fulleri</i> in cotton fields in Georgia.....	102
Army worm, fall. (See <i>Laphygma frugiperda</i> .)	
(See <i>Heliothrips unipuncta</i> .)	
Arsenate of lead against <i>Anarsia lineatella</i>	47
brown-tail moth.....	46
cucumber flea-beetle.....	111
<i>Galerucella luteola</i>	36
gipsy moth.....	23, 46
experiments in cheaper preparation.....	46
Arsenicals against <i>Cryptorhynchus lapathi</i>	27-28
<i>Estigmene acrea</i> on tobacco.....	109
all injurious to foliage or fruit of peach.....	47

	Page.
Arsenic sulphid very injurious to peach trees.....	47
Arsenites against <i>Macroductylus subspinosus</i>	35
Ash, European mountain. (See <i>Sorbus aucuparia</i> .)	
white, food plant of <i>Aspidiotus howardii</i>	88, 93
<i>Aspidiotus ancylus</i> on maple.....	88
<i>howardii</i> in Colorado.....	87-93
<i>perniciosus</i> (see also Scale, San Jose).	
in Georgia.....	103
New York State, insecticide experiments.....	39-40
Association of Economic Entomologists. (See Entomologists, Association of Economic.)	
<i>Aulacizes irrorata</i> in tobacco fields in Florida.....	111
Bacteria in control of insects.....	83
Bagging grapes, protection against <i>Macroductylus subspinosus</i>	35
no longer employed against grape rot in New Jersey.....	35
Baker, C. F., paper, "Occurrence of the Throat Bot in Cuba (<i>Gastrophilus nasalis</i> L.)".....	117
"Remarkable Habits of an Important Predaceous Fly (<i>Ceratopogon eriophorus</i> Will.)".....	117-118
"Some Notes on Leprosy in Havana".....	118-119
Ball, E. D., paper, "The Beet Leaf-hopper".....	117
"The Control of the Codling Moth in the Arid Regions".....	55-75
Banding against gipsy moth.....	23
with insect lime against <i>Epicarus imbricatus</i>	81-82
Banks, C. S., paper, "Problems in Economic Entomology in the Philippines".....	117
Bark scale, elm. (See <i>Gossyparia spuria</i> .)	
Bean tree, white. (See <i>Sorbus aria</i> .)	
Bee comb, food of <i>Galleria mellonella</i>	106
moth. (See <i>Galleria mellonella</i> .)	
<i>Bembidium quadrimaculatum</i> , possible enemy of cabbage maggot.....	15
Blackberry, food plant of cutworms.....	125
<i>Blapstinus metallicus</i> on tobacco in Florida.....	110
<i>Blarina brevicauda</i> (see also Shrew, short-tailed).	
possible enemy of <i>Isosoma tritici</i>	100
Blister-mite, pear. (See <i>Eriophyes pyri</i> .)	
Bollworm. (See <i>Heliothis obsoleta</i> .)	
Bordeaux mixture against cucumber flea-beetle.....	112
rot of grape.....	35
and Disparene against codling moth.....	53-54, 75-76
not effective against <i>Psylliodes punctulata</i>	112
Borer, flat-headed, injury to chestnut telephone poles.....	38-39
sugar maple. (See <i>Plagionotus speciosus</i> .)	
twig. (See <i>Anarsia lineatella</i> .)	
Bot, horse. (See <i>Gastrophilus equi</i> .)	
throat. (See <i>Gastrophilus nasalis</i> .)	
Bouteloua, food plant of <i>Isosoma</i>	97
Bubonic plague, possible transmission by fleas.....	119
Budworms of tobacco. (See <i>Chloridea virescens</i> and <i>Heliothis obsoleta</i> .)	
Burgess, A. F., paper, "Remarks on Methods Used in Codling Moth Experiments".....	53-55
Braconid parasite of <i>Eulecanium nigrofasciatum</i>	38
Bran and glue, protection against cabbage maggot.....	14-15
Bridge grafting to offset injury by <i>Plagionotus speciosus</i>	41
Britton, W. E., paper, "Occurrence of the Gipsy Moth in Connecticut".....	22-26
"The Maple Leaf Stem-borer (<i>Priophorus acericaulis</i> MacG.)".....	94
Brown rot of peaches following work of <i>Conotrachelus nenuphar</i>	103
tail moth (see also <i>Euproctis chrysorrhæa</i>).	
not yet found in Connecticut.....	25
Cabbage bug, harlequin, in Georgia.....	104
food plant of <i>Noctua clandestina</i>	126
<i>Thrips tabaci</i>	112
maggot in Minnesota.....	13-15
worms in Georgia.....	104
<i>Callipterus carvella</i> on pecan in Georgia.....	105
Calosoma larvæ, enemies of cutworms.....	98
<i>scrutator</i> , enemy of gipsy moth.....	25

	Page.
Campanulæ, food plants of <i>Chorizagrotis auxiliaris</i> and <i>Noctua clandestina</i>	126
Candytuft, food plant of <i>Chorizagrotis auxiliaris</i> and <i>Noctua clandestina</i>	126
Carbolic emulsion against cabbage maggot	14
Carbon bisulphid against little red ant.	17
<i>Carpocapsa pomonella</i> (see also Codling moth).	
in Georgia	104
Maryland	38
<i>Carpophilus dimidiatus</i> in rotting cotton bolls in Georgia	102
Carrot, food plant of <i>Noctua clandestina</i>	126
"Caterpillar hunter." (See <i>Calosoma scrutator</i>).	
Cauliflower, food plant of cabbage maggot	13
<i>Thrips tabaci</i>	112
Caustic soda against aphid eggs	30
Cecidomyiidae, observations	112
<i>Ceratomia catalpæ</i> in New Jersey	37
<i>Ceratopogon eriophorus</i> , enemy of <i>Melanchroia geometroides</i>	117-118
Cereal and forage crop production, value of parasites	94-99
<i>Chalcodermus æneus</i> on cotton in Georgia	101
Cheat, food plant of <i>Isosoma</i>	97
Cherry, food plant of gipsy moth	23
Chestnut telephone poles, injury by flat-headed borer	38-39
<i>Chilocorus birulnerus</i> , enemy of <i>Aspidiotus howardii</i>	90
<i>similis</i> , introduction into New Jersey a failure	37
<i>Chionaspis furfura</i> in New York State	39
<i>Chloridea virescens</i> on tobacco in Florida, remedies	106-108
"Chloroleum" against <i>Aspidiotus howardii</i>	91
<i>Chorizagrotis auxiliaris</i> in central Alberta	125-126
Chrysanthemum powders against mosquitoes in dwellings	123-124
<i>Chrysobothris femorata</i> in Georgia	104
<i>Cicada linnei</i> , new name for species heretofore determined as <i>C. tibicen</i>	35
periodical. (See <i>Tibicen septendecim</i> .)	
<i>pruinosa</i> , identity	35
<i>sayi</i> , <i>C. pruinosa</i> of authors a synonym	35
<i>tibicen</i> , identity	35
Cigarette beetle. (See <i>Lasioderma testaceum</i> .)	
Climate as affecting insect control	83-84
Climatic influences as affecting life-history periods in insects	81
<i>Coccophagus flavoscutellum</i> , is it form of <i>C. lecanii</i> ?	52
parasite of <i>Pulvinaria innumerabilis</i>	51-52
<i>lecanii</i> , parasite of <i>Pulvinaria innumerabilis</i>	48-52
Codling moth. (See also <i>Carpocapsa pomonella</i> .)	
broods in Utah	56
control in arid regions (Utah)	55-75
experiments, methods	53-55
Collinge, Walter E., paper, "On the Eradication of the Black Currant Gall-mite (<i>Eriophyes ribis</i> Nalepa)"	119-123
Colorado, economic work against <i>Aspidiotus howardii</i>	87-93
Columbines, food plants of <i>Chorizagrotis auxiliaris</i> and <i>Noctua clandestina</i>	126
Connecticut, occurrence of gipsy moth	22-26
<i>Conotelus obscurus</i> in cotton bloom in Georgia	102
<i>Conotrachelus nenuphar</i> in Georgia	103
<i>Contarinia violicola</i> , control in greenhouses	48
in New York State	41-42
<i>tritici</i> , damage prevented by parasites	94
Copidosoma, parasite of <i>Plusia brassicae</i> , polyembryony	96
<i>Cordyceps clavulata</i> (?), fungous disease of <i>Eulecanium nigrofasciatum</i>	38
Cornicles of aphides, do they excrete honeydew?	33-34
Corn meal, diluent of Paris green against tobacco budworms	107
stored, food of <i>Sitotroga cerealella</i>	102
<i>Cotoneaster vulgaris</i> , food plant of <i>Eriophyes pyri</i>	44
Cotton aphid. (See <i>Aphis gossypii</i> .)	
beetle, new. (See <i>Luperodes brunneus</i> .)	
food plant of <i>Eriophyes gossypii</i>	122
insects in Georgia	101-102
leaf-worm. (See <i>Alabama argillacea</i> .)	
Cowpea, food plant of tobacco budworms	107
pod weevil. (See <i>Chalcodermus æneus</i> .)	

	Page.
Cranberry, food plant of <i>Heliophila unipuncta</i>	35-36
Creosote oil against eggs of gipsy moth.....	23
Cricket, white flower. (See <i>Ecanthus niveus</i> .).....	
Crude oil against aphid eggs.....	29-30
Cryptopristis, parasite of <i>Isosoma tritici</i>	97
<i>Cryptorhynchus lapathi</i> as a nursery pest.....	27-29
Cuba, occurrence of <i>Gastrophilus nasalis</i>	117
<i>Culex pipiens</i> , house mosquito in Mexico.....	124
Cultivation, remedy for <i>Pamphilus persicum</i>	86
Cultural remedies for cutworms.....	126
is their investigation a true part of economic entomology?.....	82-83
Curculio, plum. (See <i>Conotrachelus nenuphar</i> .).....	
Currant, black, food plant of <i>Eriophyes ribis</i>	119
food plant of cutworms.....	125
Cutworms (see also <i>Chorizagrotis auxiliaris</i> , <i>Noctua clandestina</i> , and <i>Paragrotis ochrogaster</i>). eaten by <i>Calosoma</i> larvæ.....	98
in Florida.....	110-111
Georgia.....	103, 104
remedies.....	126
<i>Dakrura coccivora</i> , enemy of <i>Pulvinaria innumerabilis</i>	49
<i>Datana integerrima</i> in Georgia.....	105
New York State.....	39
ministra in New York State.....	39
Dickerson, Edgar L., paper, "Some Observations on the Natural Checks of the Cottony Maple Scale (<i>Pulvinaria innumerabilis</i> Rathv.)".....	48-52
Digitalis, food plant of <i>Chorizagrotis auxiliaris</i> and <i>Noctua clandestina</i>	126
<i>Diplotaxis frondicola</i> on peach in Georgia.....	103
Dipping versus fumigation of nursery trees.....	26
Disease of gipsy-moth caterpillars.....	25
Disparene against codling moth.....	53-54, 75-76
and Bordeaux mixture against codling moth.....	53-54, 75-76
not effective against <i>Psyllhodes punctulata</i>	112
Drasterius. (See Wireworms.).....	
Elm bark-scale. (See <i>Gossyparia spuria</i> .).....	
Scotch, food plant of <i>Gossyparia spuria</i>	41
Elymus, food plant of <i>Isosoma</i>	97
Entomologists, Association of Economic, members, active.....	130-132
associate.....	132-134
foreign.....	134
officers for 1907.....	127
Entomology, applied, a great experiment therein.....	10
economic, importance of phenological observations.....	77
in the Philippines.....	117
under the Adams Act.....	77-84
systematic, often necessary part of economic entomology.....	80
<i>Epicærus formidosus</i> on tobacco in Florida.....	110
imbricatus an apterous species.....	81-82
<i>Epitrimerus piri</i> on apple and pear.....	46
<i>Epitrix parvula</i> on tobacco in Florida.....	109
<i>Eriophyes avellana</i> , lime and sulphur a remedy.....	123
gossypii in West Indies, lime and sulphur a remedy.....	122, 123
malifoliae on apple and pear.....	46
oleivorus in California, sulphur a remedy.....	122
pyri.....	43-46
var. variolata on apple and pear.....	46
ribis, eradication in United Kingdom.....	119-123
rudis and taxi, lime and sulphur a remedy.....	123
<i>Estigmene aceræa</i> on tobacco in Florida.....	109
<i>Eucosma strenuana</i> on ragweed in Georgia.....	105
<i>Eulecanium nigrofasciatum</i> on peach in Maryland.....	37-38
New Jersey.....	36
<i>Eupelmus allynii</i> , parasite of Hessian fly and <i>Isosoma</i>	97
<i>Euphoria melancholica</i> and <i>sepulchralis</i> on cotton in Georgia.....	102
<i>Euproctus chrysorrhæa</i> (see also Brown-tail moth). not yet in New York State.....	39

	Page.
<i>Euthrips nicotianæ</i> on tobacco in Florida.....	106
Felt, E. P., paper, "Notes on Insects of the Year 1906 in New York State".....	39-43
"Observations on Cecidomyiidae".....	112
Fernald, H. T., paper "A New Oriental Insect Pest (?) in Massachusetts".....	22
<i>Fidia viticida</i> in New York State.....	40
Flea-beetle, cucumber, Bordeaux mixture a remedy.....	112
on tobacco in Connecticut.....	111
grapevine. (See <i>Haltica chalybea</i> .)	
steely. (See <i>Haltica chalybea</i> .)	
tobacco. (See <i>Epitrix parvula</i> .)	
Flea-beetles on sugar beet, remedy.....	112
Fleas and leprosy in Havana.....	118-119
Florida, insect enemies of tobacco in 1905.....	106-111
Forage-crop production, value of parasites.....	94-99
Forest-tree insects in Georgia.....	104-105
Foxglove. (See <i>Digitalis</i> .)	
Fruit tree insects in Georgia.....	103-104
Fumigation versus dipping of nursery trees.....	26
Fungi in control of insects.....	83
Fungus associated with pear mite in Tasmania.....	47
enemy of <i>Eulecanium nigrofasciatum</i>	38
pear scab. (See <i>Venturia pirina</i> .)	
Gahan, A. B., and G. P. Weldon, paper, "Miscellaneous Insect Notes from Maryland for 1906".....	37-39
T. B. Symons, paper, "Notes on Fumigation and Dipping of Nursery Trees".....	26
<i>Galerucella luteola</i> in New Jersey.....	36
New York State.....	41
<i>Galleria mellonella</i> in bee comb in Georgia.....	106
Gall-midge, violet. (See <i>Contarinia violicola</i> .)	
mite, black currant. (See <i>Eriophyes ribis</i> .)	
Garden insects in Georgia.....	104
<i>Gastrophilus equi</i> and <i>nasalis</i> in Cuba.....	117
Georgia, some insects during 1906.....	101-106
Gipsy moth (see also <i>Porthetria dispar</i>). disease of caterpillars.....	25
in Connecticut.....	22-26
Glue and bran or sawdust, protection against cabbage maggot.....	14-15
Gooseberry, food plant of cutworms.....	125
tree. (See <i>Phyllanthus</i> .)	
<i>Gossyparia spuria</i> in New York State.....	41
Grain insects in Georgia.....	102-103
Grape root-worm. (See <i>Fidia viticida</i> .)	
Grapevine flea-beetle. (See <i>Haltica chalybea</i> .)	
Grasses, food plants of <i>Isosoma tritici</i> in Georgia.....	102
Grasshoppers on tobacco in Florida.....	109
Grass insects in Georgia.....	102-103
orchard, food plant of <i>Isosoma</i>	97
quack, food plant of <i>Isosoma</i>	97
Gregson, P. B., paper, "Notes on Insects in Central Alberta".....	125-126
Grosella. (See <i>Phyllanthus</i> .)	
Grubs, white, in Georgia.....	103
<i>Halisidota caryæ</i> in New York State.....	39
<i>Haltica chalybea</i> in New York State.....	40
Havana, notes on leprosy.....	118-119
<i>Heliophila unipuncta</i> , destruction by tachinid flies.....	98
in New Jersey.....	35-36
<i>Heliothis obsoleta</i> on cotton in Georgia.....	101
tobacco in Florida.....	106-108
Hellebore and water, remedy for cabbage maggot.....	14
<i>Hemerocampa leucostigma</i> , girdling habit.....	41
in New York State.....	41
Herrera, A. L., paper, "Destruction of Mosquitoes in Dwellings by the Powders of Chrysanthemum, Spread therein by Means of Hand-bellows or a Towel".....	123-124
Hessian fly (see also <i>Mayetiola destructor</i>). destruction by <i>Polygnotus himalis</i>	95
<i>Heterothops fumigatus</i> , possible enemy of cabbage maggot.....	15

	Page.
Hickory tussock-moth. (See <i>Halisidota caryæ</i> .)	
Hodgkiss, H. E., paper, "Effects of Sprays on Aphis Eggs".....	29-30
Honeydew, is it excreted from anus or cornicles?.....	33-34
Honey tubes. (See Cornicles.)	
Hooker, W. A., paper, "Observations on Insect Enemies of Tobacco in Florida in 1905".....	106-111
<i>Hoplia trivialis</i> on Japan plum in Georgia.....	104
Hops, food plant of <i>Psylliodes punctulata</i>	112
<i>Hordeum jubatum</i> , food plant of <i>Isosoma</i>	97
Hornworms of tobacco. (See <i>Phlegethontius quinque maculata</i> and <i>P. sexta</i> .)	
Hydrocyanic acid gas not effective against <i>Contarinia violicola</i> larvæ.....	42
recent observations on use.....	15-16
<i>Hyperaspis binotata</i> = <i>H. signata</i>	51
<i>signata</i> , enemy of <i>Pseudococcus aceris</i>	51
<i>Pulvinaria innumerabilis</i>	49, 52
<i>Hyphantria cunea</i> in Georgia.....	104-105
Insecticides and apparatus, is their investigation a true part of economic entomology?.....	82
report of committee on cooperative testing.....	12-13
Insect pest, oriental, in Massachusetts.....	22
pests, introduced, report of committee on national control, discussion..	19-22
<i>Isosoma</i> , <i>Eupelmus allynii</i> a parasite, food plants.....	97
of timothy, Websterellus a parasite.....	98
<i>tritici</i> , Cryptopristis a parasite.....	97
in straw, killed by thrashing machine.....	100
wheat and grasses in Georgia.....	102
probably eaten by <i>Blarina brevicauda</i>	100
Jointworm. (See <i>Isosoma tritici</i> .)	
Jointworms. (See <i>Isosoma</i> .)	
Kerosene against aphid eggs.....	29-30
little red ant.....	17
willow borer.....	28
emulsion against aphid eggs.....	29-30
<i>mali</i>	104
<i>Aspidiotus howardii</i>	91
cutworms and wireworms.....	110-111
<i>Eriophyes pyri</i>	46
pupæ of <i>Pamphilus persicum</i>	86
root form of <i>Schizoneura lanigera</i>	104
tobacco insects in Florida.....	111
whitewash against aphid eggs.....	30
Kil-o-Scale against aphid eggs.....	29-30
San Jose scale.....	47-48
Kirkland, A. H., presidential address, "A Great Experiment in Applied Entomology".....	10
K-L mixture against aphid eggs.....	30
Ladybird, Asiatic. (See <i>Chilocorus similis</i> .)	
<i>Laphygma frugiperda</i> in Georgia.....	102-103
North Carolina.....	48
<i>Lasioderma testaceum</i> in stored tobacco in Florida.....	111
<i>Lathrobium anale</i> , possible enemy of cabbage maggot.....	15
Leaf-beetle, elm. (See <i>Galerucella luteola</i> .)	
hopper, beet.....	117
Leaf-worm, cotton. (See <i>Alabama argillacea</i> .)	
Lemon, food plant of <i>Eriophyes oleivorus</i>	122
<i>Lepidosaphes ulmi</i> (see also Scale, oyster-shell).	
in New Jersey.....	36
Leprosy in Havana.....	118-119
Lilac, food plant of <i>Peridroma occulta</i>	126
Lime and sulphur against species of <i>Eriophyes</i>	121, 122-123
wash for dipping nursery stock.....	26
objectionable diluent of Paris green against tobacco budworms.....	107
sulphur-salt wash not effective against <i>Eulecanium nigrofasciatum</i>	37
soda wash against <i>Aspidiotus howardii</i>	90-91
wash against <i>Aspidiotus howardii</i>	90-93
<i>perniciösus</i> , experiments in New York State.....	39-40

	Page
Linden, food plant of <i>Neurocolpus nubilus</i>	105
<i>Pulvinaria innumerabilis</i>	105
Liquidambar, defoliation by <i>Macroductylus subspinosus</i>	48
<i>Luperodes brunneus</i> on cotton in Georgia.....	101
<i>Macroductylus subspinosus</i> (see also Rose bugs).	
in Connecticut.....	86
New Jersey.....	35
North Carolina.....	48
<i>Macrosiphum granaria</i> , destruction by parasites.....	99
Mantid, Chinese. (See <i>Paratenodera sinensis</i> .)	
Maple borer. (See <i>Plagionotus speciosus</i> .)	
food plant of <i>Aspidiotus ancyllus</i>	88
<i>Lepidosaphes ulmi</i>	36
<i>Sesia tepperi</i>	105
leaf stem-borer. (See <i>Priophorus accricaulis</i> .)	
red, food plant of gipsy moth.....	23
sugar, food plant of <i>Pulvinaria acercola</i>	105
Maryland, miscellaneous insect notes for 1906.....	37-39
Massachusetts, new oriental insect pest.....	22
<i>Mayetiola destructor</i> (see also Hessian fly).	
in Georgia.....	102
<i>Megalopyge opercularis</i> in Georgia.....	106
<i>Melanchroia geometroides</i> on grosella in Cuba.....	117
<i>Melanoplus differentialis</i> , destruction by <i>Sarcophaga georgina</i>	98
Membership, report of committee.....	127-128
Midge, wheat. (See <i>Contarinia tritici</i> .)	
Minnesota, insect notes for 1906.....	13-17
Mite, cotton-leaf blister. (See <i>Eriophyes gossypii</i> .)	
gall, of black currant. (See <i>Eriophyes ribis</i> .)	
infestation in New Jersey.....	37
pear blister. (See <i>Eriophyes pyri</i> .)	
rust. (See <i>Eriophyes oleivorus</i> .)	
<i>Monomorium pharaonis</i> . (See Ant, little red.)	
Mosquitoes, destruction in dwellings by chrysanthemum powders.....	123-124
Mountain ash, European. (See <i>Sorbus aucuparia</i> .)	
Mouse, field, devouring egg masses of <i>Paratenodera sinensis</i>	37
Muhlenbergia, food plant of <i>Isosoma</i>	97
National control of introduced insect pests, report of committee.....	19-20
Nectaries. (See Cornicles.)	
<i>Neurocolpus nubilus</i> on linden in Georgia.....	105
New Jersey, unusual insect happenings in 1906.....	34-37
New York State, insect notes of 1906.....	39-43
<i>Noctua clandestina</i> in central Alberta.....	125-126
Nomenclature, report of committee.....	10-11
Nominations, report of committee.....	127
Nursery inspection, desirability of uniform State laws.....	20-22
governmental, of imports and interstate shipments.....	19-22
pest, <i>Cryptorhynchus lapathi</i>	27-29
trees, notes on fumigation and dipping.....	26
Oats, food plant of cutworms.....	126
<i>Oecanthus niveus</i> in vineyards in New York State.....	40
<i>Oncometopia lateralis</i> in tobacco fields in Florida.....	111
Onion, food plant of <i>Noctua clandestina</i>	126
<i>Thrips tabaci</i> in United States.....	112
<i>Opatrinus notus</i> on tobacco in Florida.....	110
Orange, food plant of <i>Eriophyes oleivorus</i>	122
<i>Pamphilus persicum</i> in Connecticut.....	85-86
New Jersey and Pennsylvania.....	87
Pansies, food plants of <i>Chorizagrotis auxibaris</i> and <i>Noctua clandestina</i>	126
<i>Paragrotis ochrogaster</i> in central Alberta.....	125-126
Parasites in economic entomology, importance of study.....	83
value in cereal and forage crop production.....	94-99
<i>Paratenodera sinensis</i> , failure to establish it in New Jersey.....	37
Paris green against cutworms.....	110
<i>Epitrix parvula</i>	109
grasshoppers on tobacco.....	109

	Page
Pump, compressed-air, for spraying a mechanical mixture of oil with water or other liquids.....	115
Quince, food plant of gipsy moth.....	23
Ragweed borer. (See <i>Eucosma strenuana</i> .)	
Raspberry, food plant of cutworms.....	125
Rats and leprosy in Havana.....	118-119
Report of committee on cooperative testing of insecticides.....	12-13
membership.....	127-128
national control of introduced insect pests.....	19-20
nomenclature.....	10-11
nominations.....	127
resolutions.....	128-129
Research in economic entomology under the Adams Act.....	77-84
Resolutions, report of committee.....	128-129
"Rex" lime and sulphur wash against <i>Aspidiotus howardii</i>	90-93
solution against aphid eggs.....	30
Rhubarb food plant of cutworms.....	125
Root-worm, grape. (See <i>Fidia viticida</i> .)	
Rose bugs (see also <i>Macrodactylus subspinosus</i>).	
larvæ eaten by short-tailed shrew.....	48
chafer. (See <i>Macrodactylus subspinosus</i> and Rose bugs.)	
food plant of gipsy moth.....	23
<i>Sibine stimulea</i>	105
Rumsey, W. E., paper, "Manner of Birth of the Woolly Aphis of the Apple (<i>Schizoneura lanigera</i> Hausm.)".....	31-33
Rye, food plant of <i>Isosoma</i>	97
Saddle-back caterpillar. (See <i>Sibine stimulea</i> .)	
<i>Salix alba</i> , <i>amygdaloides</i> , <i>caprea</i> , <i>cordata</i> , <i>lucida</i> , <i>sericea</i> , food plants of <i>Cryptorhynchus lapathi</i>	27
Salt marsh caterpillar. (See <i>Estigmene acrea</i> .)	
Sanderson, E. Dwight, paper, "A Spray Nozzle for the Mechanical Mixture of Oil with Water or Other Liquids".....	112-116
"What Research in Economic Entomology is Legitimate under the Adams Act?".....	77-84
<i>Sanninoidea exitiosa</i> in Georgia.....	103
<i>Saperda calcarata</i> in poplar in Georgia.....	105
<i>candida</i> in Georgia.....	104
<i>Sarcophaga georgina</i> , parasite of <i>Melanoplus differentialis</i>	98
Sawdust and glue, protection against cabbage maggot.....	14-15
Sawfly, new, attacking the peach.....	85-86
Scalecide against aphid eggs.....	29-30
<i>Aspidiotus howardii</i>	91
<i>pernicius</i>	47-48, 103
Scale, cottony maple. (See <i>Pulvinaria innumerabilis</i> .)	
false maple. (See <i>Phenacoccus acericola</i> .)	
Howard. (See <i>Aspidiotus howardii</i> .)	
oyster-shell (see also <i>Lepidosaphes ulmi</i>).	
on poplars in Ohio.....	29
peach, soft. (See <i>Eulcanium nigrofasciatum</i> .)	
Putnam. (See <i>Aspidiotus ancyclus</i> .)	
San Jose (see also <i>Aspidiotus perniciosus</i>).	
on poplars in Ohio.....	29
scurfy. (See <i>Chionaspis furfura</i> .)	
terrapin. (See <i>Eulcanium nigrofasciatum</i> .)	
<i>Schizoneura lanigera</i> (see also Aphis, woolly).	
in Georgia.....	104
manner of birth.....	31-33
<i>Schizura concinna</i> in New York State.....	39
Schoene, W. J., paper, "The Willow Borer as a Nursery Pest".....	27-28
<i>Scolytus rugulosus</i> in Georgia.....	103
"Searcher." (See <i>Calosoma scrutator</i> .)	
<i>Semiotellus chalcidiphagus</i> , double brooded, parasite of jointworms.....	97
(<i>Stictonotus isosomatis</i> , parasite of jointworms.....	97
Service-berry. (See <i>Amelanchier vulgaris</i> .)	
tree, wild. (See <i>Sorbus terminalis</i> .)	
<i>Sesia tapperi</i> in maple in Georgia.....	105

	Page.
Shade tree insects in Georgia	104-105
trees, injury by <i>Lepidosaphes ulmi</i>	36
Sharpshooters. (See <i>Aulacizes irrorata</i> and <i>Oncometopia lateralis</i> .)	
Shrew, short-tailed (see also <i>Blarina brevicauda</i>). enemy of rose bugs.	48
<i>Sibine stimulea</i> on rose and pear in Georgia	105
<i>Siphocoryne avenæ</i> effect of sprays on eggs.	29-30
in New York State	29
<i>Sitotroga cerealella</i> in stored wheat in Georgia	102
Smith, J. B., paper, "Unusual Insect Happenings in New Jersey in 1906"	34-37
R. I., paper, "Some Georgia Insects during 1906"	101-106
Snout-beetle, imbricated. (See <i>Epicærus imbricatus</i> .)	
<i>Sorbus aria</i> , <i>aucuparia</i> , and <i>terminalis</i> food plants of <i>Eriophyes pyri</i>	44
South American moth. (See <i>Thysania zenobia</i> .)	
Sphinx, catalpa. (See <i>Ceratonia catalpæ</i> .)	
Splitworm of tobacco. (See <i>Phthorimæa operculella</i> .)	
Spraying for codling moth, discussion	75-77
in arid regions (Utah)	55-75
Ohio	53-55
Spray nozzle, for mechanical mixture of oil with water or other liquids	112-117
Sprays, effects on aphid eggs	29-31
Sporobolus, food plant of <i>Isosoma</i>	97
Squash bugs in Georgia	104
Stem-borer, maple leaf. (See <i>Priophorus acericaulis</i> .)	
Stipa, food plant of <i>Isosoma</i>	97
Strawberry, food plant of cutworms	125
Structural studies, importance in economic entomology	81-82
Sugar beet, food plant of flea-beetles	112
Sulphur against <i>Eriophyes oleivorus</i>	122
and lime against species of <i>Eriophyes</i>	121, 122-123
soft soap against <i>Eriophyes ribis</i>	121-122
washes against aphid eggs	29-30
Symons, T. B., and A. B. Gahan, paper, "Notes on Fumigation and Dipping of Nursery Trees"	26
Tachinid flies, parasites of <i>Helophila unipuncta</i>	98
parasites of <i>Heliophila unipuncta</i> , practical absence in New Jersey in 1906	35
Tartaric acid solution ineffective against little red ant	17
Taylor, E. P., paper, "Economic Work Against the Howard Scale in Colorado (<i>Aspidiotus howardii</i> Ckll.)"	87-93
<i>Thrips tabaci</i> not reported on tobacco in United States	112
tobacco. (See <i>Euthrips nicotianæ</i> .)	
<i>Thysania zenobia</i> in New York State	39
<i>Tibicen septendecim</i> in Maryland	38
New Jersey	34
New York State	43
Timothy, food plant of <i>Isosoma</i>	97, 98
Tobacco decoction against aerial form of <i>Schizoneura lanigera</i>	104
<i>Aphis mali</i>	104
<i>persicæ-niger</i>	103
flea-beetle. (See <i>Epitrix parvula</i> .)	
food plant of cucumber flea-beetle	111
<i>Thrips tabaci</i> in Russia	112
insects in Florida in 1905	106-111
remedial practices	111
worms. (See <i>Phlegethontius quinque-maculata</i> and <i>P. sexta</i> .)	
<i>Toxoneuron seminigra</i> , perhaps parasite of tobacco budworms	108
Tree-cricket, snowy. (See <i>Ecanthus niveus</i> .)	
tanglefoot against gipsy moth	23
<i>Trombidium scabrum</i> , enemy of cabbage maggot	15
Turpentine against cutworms	110
Tussock moth, hickory. (See <i>Holisidota caryæ</i> .)	
white-marked. (See <i>Hemerocampa leucostigma</i> .)	
<i>Typhlodromus pyri</i> in Europe	43
<i>Venturia pirina</i> , work of pear mites sometimes mistaken therefor	47
Violet fly. (See <i>Contarinia violicola</i> .)	
gall-midge. (See <i>Contarinia violicola</i> .)	

	Page.
Violets, injury by <i>Contarinia violicola</i>	41-42, 48
Walden, B. H., paper, "Notes on a New Sawfly Attacking the Peach (<i>Pamphilius persicum</i> MacG.)".....	85-86
Wall flowers, food plants of <i>Chorizagrotis auxiliaris</i> and <i>Noctua clandestina</i>	126
Walnut, food plant of <i>Datana integerrima</i>	105
worm, black. (See <i>Datania integerrima</i> .)	
Washburn, F. L., paper, "Insect Notes from Minnesota for 1906".....	13-17
Websterellus, parasite of <i>Isosoma</i> of timothy.....	98
<i>tritici</i> , double brooded, parasite of jointworms.....	97
Webster, F. M., paper, "The Value of Parasites in Cereal and Forage Crop Production".....	94-99
Webworm, fall. (See <i>Hyphantria caneu</i>)	
Weevil, white-pine. (See <i>Pissodes strobi</i> .)	
Weldon, G. P., paper, "Miscellaneous Insect Notes from Maryland for 1906" ..	37-39
Whale-oil soap against aphid eggs.....	29-30
<i>Aspidiotus howardii</i>	93
Wheat, food plant of cutworms.....	126
<i>Isosoma</i>	97
<i>tritici</i>	102
<i>Mayetiola destructor</i>	102
midge. (See <i>Contarinia tritici</i> .)	
stored, food of <i>Sitotroga cerealella</i>	102
Whitewash against aphid eggs.....	30
Willow borer. (See <i>Cryptorhynchus lapathi</i> .)	
Wireworms on tobacco in Florida.....	110-111
"Woolworm." (See <i>Estigmene acrea</i> .)

5/5

2