**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- ☐ Coloured covers / Couverture de couleur
- ☐ Covers damaged / Couverture endommagée
- ☐ Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- ☐ Cover title missing / Le titre de couverture manque
- ☐ Coloured maps / Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- ☒ Bound with other material / Relié avec d'autres documents
- ☐ Only edition available / Seule édition disponible
- ☒ Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- ☐ Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- ☐ Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages / Pages de couleur
- ☐ Pages damaged / Pages endommagées
- ☐ Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- ☐ Pages detached / Pages détachées
- ☒ Showthrough / Transparence
- ☐ Quality of print varies / Qualité inégale de l'impression
- ☐ Includes supplementary material / Comprend du matériel supplémentaire
- ☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- ☐ Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below /
Ce document est filmé au taux de réduction indiqué ci-dessous.

10x		14x		18x		22x		26x		30x	
	12x		16x		20x		24x		28x		32x

The copy filmed here has been reproduced thanks to the generosity of:

University of Calgary,
Arctic Institute of North America,
Library

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

University of Calgary,
Arctic Institute of North America,
Library

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)

APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOL. VII: CRUSTACEA
PART B: SCHIZOPOD CRUSTACEANS

By WALDO L. SCHMITT

SOUTHERN PARTY—1913-16

OTTAWA
J. de LABROQUERIE TACHE
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1919

The Schizopod Crustaceans of the Canadian Arctic Expedition, 1913-18.

By WALDO L. SCHMITT.

With three figures in the text.

In all, only three Mysids and three Euphausiids were taken. Of the former one is apparently new to science. It is described and named in honour of Dr. R. M. Anderson, leader of the southern party of the expedition.

Of these six species the most frequently encountered was the common *Mysis oculata* (Fabr.), found in all arctic seas and the northern part of sub-arctic regions.¹ This confirms at least one of the four schizopod records given by Richters for the Bering sea.² The other three not represented in the present collection were: *Stilomysis* > *Mysideis grandis* (Goës); *Pseudomma truncatum* Smith; and *Meganyctiphanes* > *Thysanopoda norvegica* (Sars)? Concerning the last of these Hansen says, "Richters' statement, with a query, about its occurrence in Bering sea is undoubtedly wrong"; of each of the other two he remarks that though the locality is not improbable, its correctness ought to be confirmed.⁴

Mysis relicta Lovén, the well-known relict of the fresh-water lakes of boreal Europe, heretofore reported from North America only from lakes Superior, Michigan (12-148 fathoms)⁵, and Erie⁶, Green lake, Wisconsin (17-49 meters)⁷, and Indian harbour, Labrador (fresh-water)⁸, is now to be listed from Bernard harbour, Northwest Territories, where it was taken by Mr. Frits Johansen "from the stomach of *Cristivomer namaycush* (Walbaum), caught in lake, inland."

Neomysis andersoni, the new species is known only from the type locality, Grantley harbour, port Clarence, Alaska.

Thysanoëssa longipes Brandt, originally described from the sea of Okhotsk and since "taken at no less than fourteen places in the colder temperate north Pacific and Bering Sea,"⁹ has its range extended into the Arctic ocean as far as cape *Barrow* (point Barrow), Alaska.

Thysanoëssa inermis (Krøyer), though not taken farther to the eastward along the coast of North America than 140° 51' W., by this expedition, is practically bipolar in its distribution, having been taken also at Spitzbergen, in the Kara Sea, and Franz Joseph Land. In addition, it is rather widely distributed in the northern north temperate regions of the Atlantic, from the British Isles and Vineyard Sound northward into the Arctic ocean and Davis strait, and in Bering sea and adjacent area of the North Pacific.¹⁰

Thysanoëssa raschii (Sars) in its distribution approximates that of *Thysanoëssa inermis*. It, however, does not range quite as far south in the North Atlantic, never having been taken south of Scotland, or cape Cod. On the other hand, it has been found both farther to the south in the North Pacific, as far as 39° 25' N. (150° 28' W.)¹¹ and farther to the eastward in the Pacific Arctic,

¹ Zimmer, Nordisches Plankton, VI, 162, 1909.

² Richters, Abh. senekensb. naturf. Ges., XIII, 406, 1884.

³ Hansen, Proc. U. S. Nat. Mus., 48, 70, 1915.

⁴ Hansen, Danish Ingolf, Exped., III, 111 and 114, 1908.

⁵ Smith, Rept. U. S. Fish Comm., 1872-73, 643, 1874.

⁶ Kellicott, cf. Journ. Royal Micros. Soc., II, 152, 1879. For a complete bibliography of this species to 1905, see Samter, Abh. K. preuss. Akad. Wissen., Berlin, 1905, Abh. V.

⁷ Marsh, Zool. Anz., XIV, 275, 1891, and Amer. Nat., XXVIII.

⁸ Rathbun, in "Labrador" by Grenfell et al., Appendix II, 452, 1909.

⁹ Hansen, Proc. U. S. Nat. Mus., 48, 90, 1915.

¹⁰ Hansen, Proc. U. S. Nat. Mus., 48, 94-96, 1915.

¹¹ Hansen, Proc. U. S. Nat. Mus., 48, 97, 1915.

where it was taken by this expedition as far east as Bernard harbour, Northwest Territories.

Strange to say, the large *Neomysis* > *Mysis rayi* (Murdoch)¹ was not obtained, though considerable collecting was undertaken near point Barrow, the type locality. Other than those mentioned above there are only two schizopods which have been reported from near the region covered by the expedition, but not taken by it: *Gnathophausia gigas* Willemoes-Suhm² and *Thysanoëssa spinifera* Holmes³, both from the southern Bering sea, off the coast of Alaska, latitudes 53° 56' 50" N. and 57° 40' N., respectively.

The former is a species ranging as far south as the Hawaiian islands in the Pacific and the Azores in the Atlantic, but in the Atlantic not known to occur north of the 50th parallel⁴; the latter is strictly a Pacific form, which ranges southward along the West American coast nearly to the southern end of California.⁵

Of course the commoner arctic forms, such as *Michtheimysis mixta* (Lillejborg)⁶ and *Thysanoëssa neglecta*⁴ and *longicaudata* (Krøyer)⁷ are to be expected in this region, as well as almost any of the thirty-odd species occurring in the Atlantic Arctic or subarctic ocean, a list too long to introduce here. Reference is suggested to: Zimmer, Fauna Arctien, III (Die arktischen Schizopoden), 1904, and Nordisches Plankton, VI (Schizopoden), 1909, in which all these species are illustrated and fully keyed out; and Hansen Danish Ingolf Expedition, III (Crustacea Malacostraca I), 1908, and Proc. U.S. Nat. Mus., 48 (The Crustacea Euphausiacea of the U.S. Nat. Mus.), 1915.

In the following list of the species, and the stations at which they were taken reference has been made, in the case of the Mysids, to a plate and description of the species, and in the case of the Euphausiids, to the latest review of the group. The drawings were made from balsam mounts by means of the Edinger drawing apparatus. All of the species, except one of *Mysis oculata*, so indicated, were collected by Mr. Frits Johansen, naturalist of the expedition.

MYSIDACEA

Mysis oculata (Fabr.)

G. O. Sars, Mon Norges Mysider, III, 69, pl. XXXI, 1879

Station 27 n: Collinson point, Alaska, September 20, 1913.

Pelagic, over about 1 foot of water; 8 inches ice.
4 juv. Taken in company with *Thysanoëssa raschii*.

Station 27 q: Collinson Point, Alaska, September 26, 1913.

Pelagic, over about 1 fathom of water.
1 juv.

Station 27 s: Collinson point, Alaska, October 3, 1913.

Depth about 3 fathoms; sandy mud with stones and algæ.
2 juv.

Station 27 y¹: Lagoon at Collinson point, Alaska, October 8, 1913.

Pelagic, over 2 feet of water.
2 (fragmentary, sex ?) and 6 juv. Taken in company with *Thysanoëssa raschii*.

¹ Murdoch, Proc. U. S. Nat. Mus., 7, 519, 1884.

² Ortmann, Proc. U. S. Nat. Mus., 31, 38, 1906.

³ Hansen, Proc. U. S. Nat. Mus., 48, 90, 1915.

⁴ Zimmer, Nordisches Plankton, VI, 33, 1909.

⁵ Zimmer, Nordisches Plankton, VI, 163, 1909.

⁶ Zimmer, Nordisches Plankton, VI, 20, 1909.

⁷ Hansen, Proc. U. S. Nat. Mus., 48, 97-100, 1915.

- Station 28 o: Collinson point, Alaska, July 8, 1914.
From stomach of *Salvelinus malma* W.
4 ♀, 6 juv.
- Station 28 p: Mouth of Marsh river, Collinson point, Alaska, July 12-13, 1914.
From stomach of *Coregonus* sp. ("White-fish").
1 ♂, 3 ♀, and 75+ juv.
- Station 37 g: Bernard harbour, Northwest Territories, September 3, 1914.
Pelagic, over 0-2 fathoms.
250+, ♀ and juv.
- Station 37 j: Bernard harbour, Northwest Territories, September 1, 1914.
Pelagic, over about 2 fathoms of water.
1 ♂.
- Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915.
Depth about 5 fathoms; bottom, sandy mud with algæ.
7 ♀.
- Station 41 c: Bernard harbour, Northwest Territories (outer harbour), July 28, 1915.
Depth, 5 fathoms; bottom, sandy mud with algæ.
1 ♀.
- Station 41 f: Bernard harbour, Northwest Territories (outer harbour), August 1, 1915.
Depth, about 3 fathoms; bottom, mud with brown algæ.
1 ♀.
- Station 41 s: Bernard harbour, Northwest Territories (inner harbour), August 24, 1915.
Surface.
3 juv.
- Station 41 u: Bernard harbour, Northwest Territories, end of August, 1915.
From stomach of *Salvelinus malma* W.
1 ♂, 1 ♀. Taken in company with *Thysanoëssa raschii*.
- Station 42 y: Dolphin and Union strait, Northwest Territories (Bernard harbour), December 6, 1915.
Pelagic, over 3 fathoms water.
1 ♀, 1 juv.
- Station 42 z: Dolphin and Union strait, Northwest Territories (Bernard harbour), December 12, 1915 (midnight).
Pelagic, over 3 fathoms water.
1 juv.
(Schizopoda, Schmitt).
- Station 43 b: Dolphin and Union strait, Northwest Territories (off Stapyllon bay), September 14, 1915.
Depth 25-30 fathoms; bottom, sandy mud with pebbles, but no algæ.
1 ♂, 2 ♀.
- Station 43 c: Dolphin and Union strait, Northwest Territories (west of Cockburn point), September 14, 1915.
Pelagic, over 15-20 fathoms; bottom, sandy mud with stones and algæ.
1 ♀.
- Station 59 a: 71° 55' North, about 125° West, off cape Kellett, Banks island, Northwest Territories, September 7, 1914.
Depth 5-6 fathoms; bottom, sand with Laminariæ and thread algæ; 34 F.; triangle-dredge, 2 miles an hour, ¼ mile distance. George H. Wilkins, collector.

Mysis relicta Lovén.

G. O. Sars, Mon. Norges Mysider, III, 73, pl. XXXXI, 1879.

Station 42 r¹: Bernard harbour, Northwest Territories, October 10, 1913.

From stomach of *Cristivomer namaycush* W., caught in lake inland.

2 ♂, 4 ♀.

Neomysis andersoni n. sp.

This apparently new species of *Neomysis* was collected by Mr. Frits Johansen of the Southern Party of the Canadian Arctic Expedition at Grantley harbour, Port Clarence, Alaska, August 3-4, 1913. (Stations 20 a, f, surface, 1 ♂, length 20 mm., 3 ♀, 20 + juv.). Catalogue, Nos. 1251 and 1252, Victoria Memorial Museum, Ottawa, Canada.

Though considerably smaller, it is similar to *Neomysis rayi*. It differs, however, from this and all other related species on the west coast in the armature of the telson. This consists of comparatively large spines, between which are intercalated groups of smaller ones rather regularly arranged, except for the interspace adjacent to the large terminal spines of the telson, which contains three of the smaller spines; the interspaces of the distal third of the telson have to be only two small spines to each interspace; proximally there seem to be only two small spines to each interspace; terminally there are two large spines with two small ones between. The telson is deeply sulcate above and about one-fourth as long as the rest of the abdomen; the inner branch of the uropods is about as long as the telson, while the outer exceeds the telson by a little more than one-third its length.

FIG. 1.—*Neomysis andersoni*, ♀, × about 15. a, dorsal view of antennal region; b, of tail fan.

From *Mysis? costata* Holmes (see Hansen, University California Pub. Zool., 11, 177, pl. 9, figs. 2 a-d), which has a telson somewhat similarly armed laterally it is at once distinguished by the fact that the two large terminal spines are not juxtaposed, but are separated by two smaller ones.

The frontal plate is proportioned like that of *N. franciscana* Holmes (Hansen, l. c., 178, pl. 9, figs. 3 a-b, 1913), and is subtruncate, and somewhat sinuous, forming on the frontal margin three very slight lobes, one at each

antero-lateral angle of the frontal plate and a median one which is slightly behind the line connecting the anterior margins of the lateral convexities; dorsally the frontal plate is slightly grooved or depressed between the median and lateral lobes.

The eyes are short and stout, as in *N. franciscana* and likewise reach the distal segment of the antennular peduncle, which, unlike that species, is nearly one-half and not only one-third as long as the antennal scale. The distal segment of the antennular peduncle is about, or a little more than, twice as long as the second and about two-thirds as long as the first. The antennal peduncle reaches about as far forward as the eyes. The antennal scale is about twelve times as long as wide at the widest part; the terminal portion beyond the suture is contained about six and one-half times in the total length of the scale.

FIG. 2.—*Neomysis andersoni*, fourth pleopod, $\times$ about 15

Fourth pleopod of male similar to that of *N. americana* (Smith) (Rept. U.S. Fish Comm., I, 552, 1973), ultimate segment of outer rami little more than half the length of the curved terminal stylets, penultimate segment between five and one-half and six times as long as the ultimate instead of four or five times, as in *N. americana*).

This species is most closely related to *N. americana*. But the telson is relatively not so broad, tapering less abruptly toward the tip, and the smaller spines are more numerous in the interspaces between the larger spines, at least

FIG. 3.—Dorsal view of terminal millimeter and a half of telson of: a, *Neomysis andersoni* juvenile; b, *N. andersoni* adult ♀; c, *N. americana* ♀; d, *N. rayi*, ♀; e, *N. kadiakensis*, ♂; all $\times$ about 30, (last three added for purposes of comparison.)

along the distal half of the margin of the telson. In *N. americana*, the number of small spines to the interspace does not exceed three, and they are very indifferently grouped, one, two, or three between any two large spines. Moreover the antennal peduncle of *N. americana* is relatively longer, extending from one-third to one-half the length of its terminal joint beyond the second segment of the antennular peduncle; and the antennal scale is shorter and stouter, being less than ten, "about nine times as long as broad" and exceeding the antennular peduncle by less than half its length. In *N. andersoni* the antennal scale exceeds the antennular peduncle by more than half the length of the scale.

From the same station there are numerous immature individuals which have the same general appearance as the adult form described above, and similarly proportioned. The distal half of the telson, however, is more attenuate; the spines fringing the tip are subequal, while those on the lateral margins are more or less irregularly large and small, usually two or three small ones between each of two larger ones. This difference, I think, is due rather to immaturity than to a possible difference in species (cf. *Mysis oculata*, adult juvenile, G. O. Sars, l.c.).

EUPHAUSIACEA

Thysanoëssa longipes Brandt.

Hansen, Proc. U. S. Nat. Mus., 48, 87, pl. I, figs. 3a-3d, pl. II, figs. 1a-1e, 1915.

Station 57 a: Cape Smyth (point Barrow), Alaska, August 8, 1916.

Pelagic, over about 1 fathom of water.

1 ♀, 1 juv., 1 fragmentary specimen. Taken in company with *Thysanoëssa inermis*.

Thysanoëssa inermis (Krøyer)

Hansen, Proc. U. S. Nat. Mus., 48, 93, pl. II, figs. 2a-2d, 1915.

Station 29 f: 70° 13' N., 140° 50' W., April 4, 1914.

From stomach of *Phoca hispida* Schreber. Depth, about 30 fathoms.

14 (partly digested, sex ?). Taken in company with *Thysanoëssa raschii*.

Station 57 a: Cape Smyth (point Barrow), Alaska, August 8, 1916.

Pelagic, over about 1 fathom of water.

1 ♀. Taken in company with *Thysanoëssa longipes*.

Thysanoëssa raschii (M. Sars)

Hansen, Proc. U. S. Nat. Mus., 48, 96, 1915.

Station 27 n: Collinson point, Alaska, September 20, 1913.

Pelagic, over about 1 foot of water; 8 inches ice.

1 ♂. Taken in company with *Mysis oculata*.

Station 27 y¹: Lagoon at Collinson point, Alaska, October 8, 1913.

Pelagic, over 2 feet of water.

1 ♂. Taken in company with *Mysis oculata*.

Station 29 f: 70° 13' N., 140° 50' W., April 4, 1914.

From stomach of *Phoca hispida* Schreber. Depth, about 30 fathoms.

8 (partly digested, sex ?). Taken in company with *Thysanoëssa inermis*.

Station 41 u: Bernard harbour, Northwest Territories, end of August, 1915.

From stomach of *Salvelinus malma* W.

1 ♀. Taken in company with *Mysis oculata*.