*NORTH AMERICAN
WILD FLOWERS*

NORTH AMERICAN WILD FLOWERS

BY

MARY VAUX WALCOTT

PUBLISHED BY

THE SMITHSONIAN INSTITUTION

WASHINGTON, D. C.

1925

THIS IS NUMBER

380

OF VOLUME ONE

2K
112
W3X
925
V. 1, C. 1
RE
Bot.
(SCH2B)

FOREWORD

WILD FLOWERS were a joy and inspiration in the happy days of childhood when I was taught to observe and sketch them under the direction of a skilled artist. Years passed before a botanical friend at Glacier, British Columbia, asked me to portray a rare and perishable alpine flower so as to preserve its beauty, color, and graceful outline as a living thing. During succeeding seasons I painted other rare specimens until many of the "living flowers that skirt the eternal frost" in the wildflower gardens of the Canadian Rockies were transferred in color and form to the East, where sketches of the native woodland and meadow blossoms soon began to join them.

During the past ten years I have spent from three to four months each season in the Canadian Rockies, where Dr. Walcott was carrying on geological explorations, covering in all more than five thousand miles on the mountain trails. This afforded me a wonderful opportunity for intimate study of the flora, my aim being to collect and paint the finest specimens obtainable, and to depict the natural grace and beauty of the plant without conventional design. Many of the western sketches were made under trying conditions. Often, on a mountain side or high pass, a fire was necessary to warm stiffened fingers and body. In camp, the diffused light of the white tent was a great handicap, and considerable ingenuity was required to obtain a proper combination of light and shade. The paint box and pads found safe conveyance on the back of the saddle, except in unusual storms of rain or snow, and many times while waiting for the pack train to be made ready, a sketch was begun or completed. The short lives of the blooming plants definitely limit the number of sketches that can be made during a single field season, for many hours of work are necessary to finish a single sketch, and wild flowers wither quickly. A sharp frost in July or early August will ruin them, or an unusually warm, dry season

or a cold, wet one will prevent flowering or kill the blossoms that have matured. For these reasons desirable specimens of many of the fragile alpine flowers are difficult to secure, and in some instances were seen in perfection but two or three times during the many seasons on the trail. The limited habitat of others made it necessary to take long rides and climb high above the timber line to procure them, and frequently no trails were available. Our sure-footed mountain ponies were a large factor in our success.

Both the bloom and the fruit of a few trees have been sketched with the hope that these exquisite forms may be more observed and appreciated by nature lovers. The illustrations of eastern plants have been made from specimens collected as opportunity offered and from those contributed by many friends. All the sketches are life size.

As time went on and the collection grew, botanists, artists, and others interested in flowers began to urge that the water-color sketches should be permanently preserved and made available for students and lovers of the beautiful in Nature, before the dust of time faded and browned them to the hues of the pressed flowers of the herbaria. A survey of wild flower publications led to the decision that there was need for a finely illustrated work that would be of service pictorially to all professional and amateur botanists and designers, and to the larger group of lovers of wild flowers and the great out-of-doors. To many of these the living flowers are inaccessible, and their real beauty is unknown. No attempt has been made to create a text book with technical descriptions, or to illustrate all native American wild flowers, and only native plants have been included.

The preparation of the work has been a labor of love and has been made possible by the sympathetic interest and inspiration of Dr. Walcott, who has been unfailing in his help and encouragement.

My sincere thanks are due to Dr. Frederick V. Coville, Dr. Edgar T. Wherry, Mr. Paul C. Standley, and Dr. Paul Bartsch, who have all given freely of their time and knowledge.

Washington, 1925

Mary Vaux Walcott

TO THE PATRONS
WHO HAVE MADE POSSIBLE ITS PUBLICATION
THIS WORK IS DEDICATED

MISS ANNA H. CHACE
Providence, Rhode Island

MR. ROBERT S. BROOKINGS
St. Louis, Missouri

MR. W. K. BIXBY
St. Louis, Missouri

MR. FREDERIC A. DELANO
Washington, District of Columbia

MRS. JOHN HAYS HAMMOND
Washington, District of Columbia

MRS. AVERY COONLEY
Washington, District of Columbia

MR. FREDERIC C. WALCOTT
New York, New York

MR. JAMES PARMELEE
Washington, District of Columbia

MR. WORCESTER R. WARNER
Tarrytown, New York

MR. WILLIAM H. CLARKE
New York, New York

MISS LYDIA T. MORRIS
Philadelphia, Pennsylvania

MR. EFFINGHAM B. MORRIS
Philadelphia, Pennsylvania

MR. CHARLES C. GLOVER
Washington, District of Columbia

MR. AND MRS. LARZ ANDERSON
Washington, District of Columbia

MRS. PIERRE S. DU PONT
Kennett Square, Pennsylvania

HON. AND MRS. HENRY WHITE
New York and Washington

MRS. ELON H. HOOKER
New York, New York

MR. JOHN D. ROCKEFELLER, JR.
New York, New York

DR. AND MRS. A. HAMILTON RICE
New York, New York

MISS ANNIE M. HEGEMAN
Washington, District of Columbia

MISS MARTHA C. CODMAN
Washington, District of Columbia

MRS. JULIUS ROSENWALD
Chicago, Illinois

MR. JOHN PIERPONT MORGAN
New York, New York

MR. ALBA B. JOHNSON
Philadelphia, Pennsylvania

MRS. E. HENRY HARRIMAN
New York, New York

MR. HENRY H. COLLINS
Bryn Mawr, Pennsylvania

MRS. CHARLES S. HINCHMAN
Philadelphia, Pennsylvania

HON. JOHN BARTON PAYNE
Washington, District of Columbia

MRS. ANDREW CARNEGIE
New York, New York

MRS. B. H. BUCKINGHAM
Washington, District of Columbia

PATRONS

MISS ISABEL C. FREEMAN
Washington, District of Columbia

MRS. CHARLES BOUGHTON WOOD
Simsbury, Connecticut

MRS. JAMES COUZENS
Detroit, Michigan

HON. ANDREW W. MELLON
Pittsburgh and Washington

MRS. CHARLES W. WETMORE
Washington, District of Columbia

MR. AND MRS. EUGENE MEYER, JR.
New York and Washington

MRS. ROBERT BACON
New York, New York

MR. CLEMENT S. HOUGHTON
Boston, Massachusetts

MISS E. A. ACHELIS
New York, New York

MRS. HENRY O. HAVEMEYER
New York, New York

MR. JAMES W. ELLSWORTH
New York, New York

MRS. HERBERT L. SATTERLEE
New York, New York

MRS. JOHN JACOB HOFF
Paris, France

MR. ERNEST B. DANE
Boston, Massachusetts

MR. EDWIN S. WEBSTER
Boston, Massachusetts

MR. HENRY HORNBLOWER
Boston, Massachusetts

NATIONAL GEOGRAPHIC SOCIETY
Washington, District of Columbia

MRS. WILLARD H. BROWNSON
Washington, District of Columbia

MR. CHARLES H. ATHERTON
Honolulu, Hawaii

COUNTESS L. SZÉCHÉNYI
Washington, District of Columbia

MRS. WILLARD SAULSBURY
Wilmington, Delaware

HON. JOHN W. WEEKS
West Newton, Massachusetts

CANADIAN PACIFIC RAILWAY
Montreal, Canada

MRS. BENJAMIN H. WARDER
Washington, District of Columbia

MRS. JOHN J. CARTY
New York, New York

COL. ROBERT M. THOMPSON
Washington, District of Columbia

MRS. J. SHERLOCK DAVIS
Brooklyn, New York

MR. JOHN GELLATLY
New York, New York

MRS. COLEMAN DU PONT
Wilmington, Delaware

COL. EDWARD A. DEEDS
Dayton, Ohio

MR. JOHN A. ROEBLING
Bernardsville, New Jersey

MISS ELLEN B. SCRIPPS
La Jolla, California

MR. CHAUNCEY J. HAMLIN
Buffalo, New York

MR. GEORGE G. BOOTH
Bloomfield Hills, Michigan

MRS. MARCIA FLAGG BUTLER
Washington, District of Columbia

MR. CHARLES F. JENKINS
Philadelphia, Pennsylvania

PATRONS

MR. CHARLES H. LUDINGTON
Philadelphia, Pennsylvania

MR. JOHN GRIBBEL
Philadelphia, Pennsylvania

GIRL SCOUTS, INCORPORATED
New York, New York

NEW YORK BOTANICAL GARDEN
New York, New York

MR. ROBERT W. DE FOREST
New York, New York

DR. AMBROSE SWASEY
Cleveland, Ohio

MR. HENRY W. DE FOREST
New York, New York

MR. FELIX M. WARBURG
New York, New York

MISS HELEN C. FRICK
New York, New York

MRS. NICHOLAS F. BRADY
New York, New York

MRS. ALFRED DUANE PELL
New York, New York

MR. PAUL D. CRAVATH
New York, New York

MR. CHARLES A. COFFIN
New York, New York

MR. ANTON G. HODENPYL
New York, New York

MR. CLEVELAND H. DODGE
New York, New York

MR. GANO DUNN
New York, New York

MR. HERBERT L. PRATT
New York, New York

DR. HENRY S. WELLCOME
London, England

MR. ROLAND L. TAYLOR
Philadelphia, Pennsylvania

MR. HENRY B. SPENCER
Washington, District of Columbia

DR. EDWARD D. ADAMS
New York, New York

DR. CHARLES D. WALCOTT
Washington, District of Columbia

MR. GEORGE B. MATHEWS
Buffalo, New York

MRS. IRÉNÉE DU PONT
Granogue, Delaware

MR. MARTIN A. RYERSON
Chicago, Illinois

AMERICAN MUSEUM OF NATURAL HISTORY
New York, New York

MR. GEORGE VAUX, JR.
Bryn Mawr, Pennsylvania

MR. CHARLES DEERING
Miami, Florida

MASSACHUSETTS HORTICULTURAL SOCIETY
Boston, Massachusetts

MISSOURI BOTANICAL GARDEN
St. Louis, Missouri

FIELD MUSEUM OF NATURAL HISTORY
Chicago, Illinois

MR. CYRUS H. MCCORMICK
Chicago, Illinois

MR. AND MRS. E. V. HUNTINGTON
Cambridge, Massachusetts

MRS. HERBERT WARD
London, England

MR. BENEDICT ERSTEIN
New York, New York

HON. THOMAS F. BAYARD
Wilmington, Delaware

PATRONS

MR. EUGENE H. ANGERT
St. Louis, Missouri

MR. HERMAN A. METZ
New York, New York

BROOKLYN BOTANIC GARDEN
Brooklyn, New York

MOUNT VERNON PUBLIC LIBRARY
Mount Vernon, New York

MRS. ROCKEFELLER McCORMICK
Chicago, Illinois

ESTATE OF SAMUEL BANCROFT, JR., INC.
Wilmington, Delaware

MR. JOHN RUSSELL GLADDING
Providence, Rhode Island

MR. EDWARD E. ARNOLD
Providence, Rhode Island

MR. ROBERT GOELET
New York, New York

MR. AUGUSTUS S. PEABODY
Chicago, Illinois

MR. L. W. DOMMERICH
New York, New York

MRS. WILLIAM H. FORBES
Milton, Massachusetts

JUDGE ROBERT W. BINGHAM
Louisville, Kentucky

MR. GEORGE M. SHRIVER
Baltimore, Maryland

DR. MICHAEL I. PUPIN
New York, New York

MR. H. B. CLOW
Chicago, Illinois

MRS. JESSE ISIDOR STRAUS
New York, New York

MRS. DUDLEY S. BLOSSOM
South Euclid, Ohio

DR. ANNA P. SHARPLESS
Philadelphia, Pennsylvania

MRS. CHESTER C. BOLTON
Cleveland, Ohio

MR. PARMELY W. HERRICK
Cleveland, Ohio

MR. J. J. BODELL
Providence, Rhode Island

ILLINOIS STATE LIBRARY
Springfield, Illinois

MR. HENRY F. DU PONT
Winterthur, Delaware

ELI LILLY AND COMPANY
Indianapolis, Indiana

MRS. CHARLES S. BIRD
East Walpole, Massachusetts

OHIO STATE LIBRARY
Columbus, Ohio

MASSACHUSETTS STATE LIBRARY
Boston, Massachusetts

MR. R. H. SCOTT
Lansing, Michigan

MR. AND MRS. HENRY SOLON GRAVES
New Haven, Connecticut

MR. ALBERT H. WIGGIN
New York, New York

MRS. EDWARD S. HARKNESS
New York, New York

MR. CHARLES R. MORLEY
Mentor, Ohio

MRS. STANLEY G. FLAGG, JR.
Bryn Mawr, Pennsylvania

MISS ELEANOR McC. CHALFANT
Pittsburgh, Pennsylvania

MRS. GEORGE W. PERKINS
Riverdale-on-Hudson, New York

PATRONS

MR. FREDERICK W. VANDERBILT
New York, New York

MRS. JOHN T. PRATT
New York, New York

MR. ROBERT McK. JONES
St. Louis, Missouri

MRS. W. A. HARRIMAN
New York, New York

MRS. MAUD WALKER AMES
Seattle, Washington

MR. ISRAEL PLATT PARDEE
Hazleton, Pennsylvania

MRS. ROBERT W. POMEROY
Buffalo, New York

MR. LOUIS LEE HAGGIN
Lexington, Kentucky

MRS. ERNEST A. BRYANT
Los Angeles, California

MR. LEWIS L. DELAFIELD
New York, New York

MRS. CHARLES COPELAND
Wilmington, Delaware

MR. AUGUST A. RUBEL
Santa Paula, California

MR. CARLL TUCKER
New York, New York

MR. C. C. AUCHINCLOSS
New York, New York

MRS. G. LISTER CARLISLE, JR.
New York, New York

MRS. MINTURN T. WRIGHT
Philadelphia, Pennsylvania

MRS. JAMES COY BRADY
New York, New York

MR. JAMES C. COLGATE
New York, New York

MRS. HAROLD IRVING PRATT
New York, New York

MRS. KING C. GILLETTE
Beverly Hills, California

DR. R. A. F. PENROSE, JR.
Philadelphia, Pennsylvania

MRS. MARGARET ROEBLING PERRINE
Trenton, New Jersey

MR. WALTER WOOD
Philadelphia, Pennsylvania

LIBRARY OF CONGRESS
Washington, District of Columbia

MR. AND MRS. J. OTIS WARDWELL
Haverhill, Massachusetts

MRS. LOUIS A. FROTHINGHAM
Washington, District of Columbia

MRS. JESSE H. METCALF
Providence, Rhode Island

MR. DWIGHT W. MORROW
New York, New York

MR. GEORGE W. MARSTON
San Diego, California

PENNSYLVANIA STATE LIBRARY
Harrisburg, Pennsylvania

MR. HUNT T. DICKINSON
New York, New York

E. P. DUTTON AND COMPANY
New York, New York

NEW YORK PUBLIC LIBRARY
New York, New York

WORTHY PAPER COMPANY ASSOCIATION
Mittineague, Massachusetts

PATRONS

HENRY E. HUNTINGTON LIBRARY
AND ART GALLERY
San Marino, California

SACRAMENTO CITY LIBRARY
Sacramento, California

DEPARTMENT OF AGRICULTURE
Washington, District of Columbia

LOS ANGELES PUBLIC LIBRARY
Los Angeles, California

PUBLIC LIBRARY OF TORONTO
Toronto, Canada

LOS ANGELES MUSEUM
Los Angeles, California

MRS. PAUL H. BASTEDO
Washington, District of Columbia

JAMES JEROME HILL REFERENCE
LIBRARY
St. Paul, Minnesota

MRS. PAUL D. I. MAIER
Bryn Mawr, Pennsylvania

MR. GEORGE D. PRATT
Glen Cove, New York

MR. JOSEPH LEITER
Chicago and Washington

MR. MOSES TAYLOR
Portsmouth, Rhode Island

HAVERHILL PUBLIC LIBRARY
Haverhill, Massachusetts

MR. DONALD WOODWARD
Le Roy, New York

MISS ELIZABETH CANBY MORRIS
Philadelphia, Pennsylvania

MR. JOSEPH R. NUTT
Cleveland, Ohio

HON. GEORGE WHARTON PEPPER
Philadelphia, Pennsylvania

MRS. IRWIN STRASBURGER
New York, New York

MR. ARTHUR N. PACK
Princeton, New Jersey

DR. WILLIAM H. WILMER
Baltimore, Maryland

MR. EMORY W. CLARK
Detroit, Michigan

J. W. ROBINSON COMPANY
Los Angeles, California

MISS HELEN R. BIDDLE
Bryn Mawr, Pennsylvania

MRS. GRACE PHILLIPS JOHNSON
New Castle, Pennsylvania

MR. DELOS A. BLODGETT, II
New York, New York

MRS. FRANK B. FOSTER
Haverford, Pennsylvania

MRS. CHARLES L. HUTCHINSON
Chicago, Illinois

MR. E. H. LUPTON, JR.
Topeka, Kansas

MRS. EDWIN C. JAMESON
New York, New York

EDWARD W. SMITH, M.D.
Meriden, Connecticut

MISS GERTRUDE B. WHITTEMORE
Naugatuck, Connecticut

MR. JOHN A. KEPPELMAN
Reading, Pennsylvania

MAJOR HENRY P. ERWIN
Washington, District of Columbia

TABLE OF CONTENTS

VOLUME I

NOTE: All sketches are life size. The system used in naming the plants is the American Code of Botanical Nomenclature. Descriptions of the plants illustrated may be found in Gray's New Manual, Britton and Brown's Illustrated Flora, Small's Flora of the Southeastern United States, or Rydberg's Flora of the Rocky Mountains.

PLATE

1. Mountain Ladyslipper. *Cypripedium montanum* DOUGLAS
2. Pussy-cars. *Calochortus elegans* LINDLEY
3. False Locoweed. *Oxytropis gracilis* (NELSON) JONES
4. Nodding Ladies-tresses. *Ibidium cernuum* (LINNAEUS) HOUSE
Slender Ladies-tresses. *Ibidium gracile* (BIGELOW) HOUSE
5. Pink Fumeroot. *Capnoides sempervirens* (LINNAEUS) BORKHAUSEN
6. Sun-dial Lupine. *Lupinus perennis* LINNAEUS
7. Big Whortleberry. *Vaccinium membranaceum* DOUGLAS
8. Pinebarren Gentian. *Gentiana porphyrio* GMELIN
9. Skeletonflower. *Lygodesmia juncea* (PURSH) DON
10. Lake Louise Arnica. *Arnica louiseana* FARR
11. Red Lily. *Lilium montanum* NELSON
12. Cottongrass. *Eriophorum chamissonis* MEYER
13. Vernal Iris. *Iris verna* LINNAEUS
14. Anemonella. *Synedesmon thalictroides* (LINNAEUS) HOFFMANNSEGG
15. White Troutlily. *Erythronium albidum* NUTTALL
16. Hazel Alder. *Alnus rugosa* (DU ROI) SPRENGEL
17. Tarflower. *Befaria racemosa* VENTENAT
18. Alpine Fir. *Abies lasiocarpa* (HOOKER) NUTTALL
19. Oconee-bells. *Shortia galacifolia* TORREY AND GRAY
20. Virginia Bluebells. *Mertensia virginica* (LINNAEUS) DE CANDOLLE
21. Burgess Milkvetch. *Astragalus bourgovii* GRAY
22. Greendragon. *Arisaema dracontium* (LINNAEUS) SCHOTT
23. Toad Trillium. *Trillium sessile* LINNAEUS
24. Southern Magnolia (flower). *Magnolia grandiflora* LINNAEUS

- 24a. Southern Magnolia (fruit). *Magnolia grandiflora* LINNAEUS
25. Trumpetleaf. *Sarracenia flava* LINNAEUS
26. Redbud. *Cercis canadensis* LINNAEUS
27. Tall Larkspur. *Delphinium elongatum* RYDBERG
28. Rhodora. *Rhodora canadensis* LINNAEUS
29. Pickerelweed. *Pontederia cordata* LINNAEUS
30. Creeping Hollygrape. *Berberis repens* LINDLEY
31. Red Chokeberry (flower). *Aronia arbutifolia* (LINNAEUS FILIUS) ELLIOTT
- 31a. Red Chokeberry (fruit). *Aronia arbutifolia* (LINNAEUS FILIUS) ELLIOTT
32. Devilsclub. *Echinopanax horridum* (SMITH) DECAISNE AND PLANCHON
33. Crested Iris. *Iris cristata* AITON
34. Lily Twayblade. *Liparis liliifolia* (LINNAEUS) RICHARD
35. Missouri Pricklypear. *Opuntia polyacantha* HAWORTH
36. Butterflyweed. *Asclepias tuberosa* LINNAEUS
37. Skunkcabbage. *Spathyema foetida* (LINNAEUS) RAFINESQUE
38. Red-osier Dogwood. *Cornus stolonifera* MICHAUX
39. Birdsfoot Violet. *Viola pedata* LINNAEUS
40. Virginia Spiderwort. *Tradescantia virginiana* LINNAEUS
41. Leatherflower. *Clematis viorna* LINNAEUS
42. Purple Saxifrage. *Saxifraga oppositifolia* LINNAEUS
43. Flame Azalea. *Azalea lutea* LINNAEUS
44. Rabbitbean. *Cracca virginiana* LINNAEUS
45. Tuliptree. *Liriodendron tulipifera* LINNAEUS
46. Trumpet Honeysuckle. *Lonicera sempervirens* LINNAEUS
47. Red Buckeye. *Aesculus pavia* LINNAEUS
48. Rose Paintbrush. *Castilleja pallida* (LINNAEUS) KUNTH
49. Shootingstar. *Dodecatheon meadia* LINNAEUS
50. Prairie Pentstemon. *Pentstemon erianthera* PURSH
51. Wild Sweet Crab. *Malus coronaria* (LINNAEUS) MILLER
52. Pitcherplant. *Sarracenia purpurea* LINNAEUS
53. Prairie-smoke. *Sieversia ciliata* (PURSH) DON
54. Winterberry. *Ilex verticillata* (LINNAEUS) GRAY

55. Sweet Azalea. *Azalea arborescens* PURSH
56. Kruhsea (flower). *Kruhsea streptopoides* (LEDEBOUR) KEARNEY
- 56a. Kruhsea (fruit). *Kruhsea streptopoides* (LEDEBOUR) KEARNEY
57. Arethusa. *Arethusa bulbosa* LINNAEUS
58. Pale Ladyslipper. *Cypripedium acaule* AITON
59. Quakerladies. *Houstonia caerulea* LINNAEUS
60. Bearberry Honeysuckle. *Lonicera involucrata* (RICHARDSON) BANKS
61. Pink Fleabane. *Erigeron caespitosus* NUTTALL
62. Labrador-tea. *Ledum groenlandicum* OEDER
63. Northern Bedstraw. *Galium boreale* LINNAEUS
64. Bronzebells. *Stenanthium occidentale* GRAY
65. Roundleaf Orchis. *Orchis rotundifolia* PURSH
66. Prickly Currant (flower). *Ribes lacustre* (PERSOON) POIRET
67. Prickly Currant (fruit). *Ribes lacustre* (PERSOON) POIRET
68. Glacierlily. *Erythronium grandiflorum* PURSH
69. Balsamroot. *Balsamorhiza sagittata* (PURSH) NUTTALL
70. Silverberry (flower). *Elaeagnus commutata* BERNHARDI
71. Silverberry (fruit). *Elaeagnus commutata* BERNHARDI
72. Twinleaf. *Jeffersonia diphylla* (LINNAEUS) PERSOON
73. Ivory Baneberry. *Actaea arguta* NUTTALL
74. Pink Mountainheather. *Phyllodoce empetriiformis* (SMITH) DON
75. Rocky Mountain Cassiope. *Cassiope mertensiana* (BONGARD) DON
76. One-leaf Bog-orchid. *Habenaria obtusata* (PURSH) RICHARDSON
77. Bush Cinquefoil. *Potentilla fruticosa* LINNAEUS
78. Refer to plate 24a
79. Refer to plate 31a
80. Refer to plate 56a

MOUNTAIN LADYSLIPPER

Cypripedium montanum Douglas

One lovely morning in early summer we left camp near Radium Hot Springs in the Columbia River Valley in British Columbia, our two saddle horses harnessed to the buckboard, the saddles securely tied behind the seat. After a drive of seven miles the horses were saddled, and we followed a disused trail four or five miles back into the range on the eastern side of the valley. There were many flowers, the dense growth of trees and shrubs keeping out the hot rays of the sun and conserving the moisture. Just as we turned across a partially shaded flat on the side of the little canyon we had been traversing, a wonderful sight burst upon us. Scattered among the low-growing bushes were great clumps of this splendid orchid in the perfection of bloom. A beautiful plant about eighteen inches tall, with a delicate perfume quite its own—no wonder that we quickly dismounted to pay homage to this queen of the forest.

Never before had we seen such a flower, and no book upon the plants of the Canadian Rockies mentioned its occurrence there. A bunch carried to camp survived many days, long after the sketch was completed.

The mountain ladyslipper is frequently found in the mountains of California, and northward along the coast as far as Vancouver, and northeastward along the Rocky Mountains to Saskatchewan.

PUSSY-EARS

Calochortus elegans Lindley

We were following the old stage road, now replaced by a new motor road, in the Columbia River Valley, twenty miles south of Canal Flats, at an altitude of 2,500 feet. The marvelous larches and pines of the primeval forest towered above us, and over the grassy forest floor were scattered many flowers with which we were familiar. Suddenly we saw the grass dotted with a white lily, which was unknown to us, and upon examination we quickly comprehended the reason for its common name of "pussy-ears," which our companion gave it. So graceful was it that we could only wonder that a flower so beautiful had not been enticed long ago to grow in our cultivated gardens.

Pussy-ears ranges from Utah and California to British Columbia and Montana.

FALSE LOCOWEED

Oxytropis gracilis (Nelson) Jones

This plant must not be confused with its relative, the true locoweed, that is so destructive to animals on the plains of the southwestern United States; it seemed entirely harmless to our horses, who ate it freely. In full sunshine its pale yellow or straw-colored flowers, borne on stems six or eight inches tall, are wonderfully attractive against their setting of sage-colored foliage. In limestone regions, where moisture is limited, the blossoms develop well only if an abundance of water is available at flowering time.

False locoweed is a member of the Pea Family. It ranges from Idaho to South Dakota and northward into Alberta, Canada.

The specimen sketched was obtained near Ghost River on the eastern slope of the Rocky Mountains, twenty-five miles from Banff, Alberta, at an altitude of 3,500 feet.

NODDING LADIES-TRESSES

Ibidium cernuum (Linnaeus) House

SLENDER LADIES-TRESSES

Ibidium gracile (Bigelow) House

These two fall-blooming members of the Orchid Family are frequently found growing in the grass of old pastures. Nodding ladies-tresses usually chooses a wet situation, but the slender one is more likely to be found in drier places. Both are inconspicuous in their chosen habitats, for their leaves and stalks are so grasslike that they seem a part of the general herbage. The genus name has arisen from the fancied resemblance of the anther to the head of an ibis.

Many botanists prefer to use the genus name *Spiranthes*, which means spiral-flower, but *Ibidium* has priority.

Slender ladies-tresses is notable for its green-centered lip, and for the fact that its leaves, which are evergreen, wither away before flowering time, the flower stalk bearing only inconspicuous bracts.

These two orchids have a similar range, occurring from Florida northward and westward, extending into southernmost Canada.

The specimens sketched were found near Mount Kisco, New York.

PINK FUMEROOT

Capnoides sempervirens (Linnaeus) Borkhausen

Pink fumeroot is a dainty wild beauty that loves rich rocky woods, where it grows in perfection. The leaves are a delicate shade of green, and the pale pink flowers, with a yellow lip, poised on slender stems, sway gently in the wind. The plant is a relative of the bleeding-heart of our gardens, and of the European fumitory. Like the latter, its roots exhale an intense nitrous odor, and the common name fumeroot refers to this characteristic feature.

Pink fumeroot has a wide range, from North Carolina to Nova Scotia and westward to British Columbia and Alaska.

We found it in the valley of the Kootenai River near the motor road between Banff and the Columbia River Valley, at an altitude of 4,000 feet.

SUN-DIAL LUPINE

Lupinus perennis Linnaeus

Sun-dial lupine in the East seldom equals in profusion of growth its sisters of the West, which often cover the hillsides with a carpet of bloom. Nevertheless, this common species of the Eastern States is as beautiful as any of the others, and deserves a place of honor in our wild gardens. In rich loam it will dwindle and die, but if planted in an acid, sandy soil, will thrive in cultivation. Although usually bright blue, the flowers are occasionally white or lavender, sometimes bordering on purple.

In Texas, one of the lupines commonly called blue-bonnets has been named the State flower.

The sun-dial lupine is found from Louisiana and Florida north to Maine and Minnesota.

The specimen sketched grew near Washington, District of Columbia.

BIG WHORTLEBERRY

Vaccinium membranaceum Douglas

Snow slides are not to be desired, but after they have mowed down the forest of the mountain slopes, exposing the ground to the full sunlight, a bushy growth springs up, three or four feet tall, with young trees interspersed. Among these shrubs we find whortleberry bushes, Rocky Mountain rhododendron, and other plants that love similar conditions of soil and exposure. Here in late summer a delightful feast is spread for the traveler, for the big whortleberry bushes are loaded with dark purple berries, some of them as large as small cherries. A large quantity may be gathered in a short time. The black bears also appreciate them, and often are met in these localities, sitting on their haunches and sweeping the ripe fruit into their mouths with their great paws. In autumn the leaves turn a deep red.

Big whortleberry is found occasionally in the Rockies, but reaches greatest perfection in the Selkirk Mountains, and is especially abundant at Rogers Pass and in the vicinity of Glacier House at Glacier. It ranges from California, Wyoming, and Michigan to British Columbia.

The specimen sketched was obtained near Hector Station on the Canadian Pacific Railway, British Columbia, at an altitude of 5,000 feet.

PINEBARREN GENTIAN

Gentiana porphyrio Gmelin

Pinebarren gentian is a graceful and attractive plant. Its deep blue corolla lobes, with peculiar fringes between them, and the curious yellow markings on the inside, make a pleasing color combination seldom seen in flowers. It frequents moist pine barrens, reaching a foot or more in height, and although the flowers grow singly on the branches, a number of stems usually arise from each root, so that the clumps are very showy.

This gentian blooms from late August into October, and can be cultivated if given the moist, acid, peaty soil in which it thrives in the wild state. It occurs in the Coastal Plain from Florida to southern New Jersey.

The specimen sketched was found near Wilmington, North Carolina.

SKELETONFLOWER

Lygodesmia juncea (Pursh) Don

This is a strange-looking plant, a foot tall, with pale pink, starlike flowers that close in the afternoon, or very soon after being gathered. Unconsciously, the observer is likely to notice only the flowers, for the rushlike stems and scalelike foliage are sage-green and inconspicuous. The plant belongs to the subdivision of the Aster Family that includes the dandelion, wild lettuce, and hawkweed, and is characterized by having all the flowers of the head furnished with petallike rays, and by a bitter milky juice.

The skeletonflower is found from Nevada and New Mexico north to Minnesota and Alberta.

We found it growing on a dry bank, high above the Kootenai River, near the junction of this stream with the Columbia River in British Columbia, at an altitude of 2,800 feet, and marveled that it could thrive in dusty soil, which remains damp only a short time after the mountain showers cease.

LAKE LOUISE ARNICA

Arnica louiseana Farr

The horses had traveled a hard and stony way, very steep in places, for we were riding above timber line to visit an alpine lake tucked away in a glaciated cirque near the headwaters of the Clearwater River in British Columbia. There had been no trail to follow, except for the first half mile from camp, and we had reveled in the constant succession of wild gardens, filled with mountain flowers in the perfection of bloom, through which we were passing. Many of the plants were crushed by the feet of the ponies, and we were continually on the watch for new varieties. A long grassy slope facing north was being conquered, when several yellow flowers appeared, quite different from any we had seen before in the mountains. Upon closer examination they were recognized as the rare Lake Louise arnica, a composite found only in the Canadian Rockies. The best specimens were carefully gathered, carried to camp, and sketched without delay, but the thrill of seeing them in their native habitat will always be recalled with delight.

RED LILY

Lilium montanum Nelson

Such hosts of red lilies are seen in the dry woods of the Canadian Rockies in some sections in July, that the prodigality of nature seems astounding. If perchance a richer soil and more sunlight are available, the perfectly developed flowers and the stouter leaves and stems indicate an appreciation of these favors. But the red lilies are tender things, withering speedily when gathered, especially when exposed to sun and wind, and they do not easily revive when placed in water.

The red lily is found in the Rocky Mountains from New Mexico to British Columbia, and eastward in the foothills and valleys.

The specimen painted was gathered in Sinclair Canyon, not far from Radium Hot Springs, British Columbia, at an altitude of 3,000 feet.

COTTONGRASS

Eriophorum chamissonis Meyer

In all parts of the Rocky Mountains of Canada one's attention is drawn in mid-July to the masses of sedge growing along the wet borders of alpine lakes, oftentimes out of the water itself. If there have been a few dry days, the cottony heads are waving in the breeze, as it fitfully blows over them. Farther away from the lake shore the plants become scattered and lose much of the beauty that comes from a massed effect. Buffeted by mountain showers and winds, they are in their prime only a few days.

This species of cottongrass is found usually about timber line in the Rockies. It is a circumpolar plant, occurring in northern Europe and Asia as well as North America. On this continent it perhaps survived the glacial period in Wyoming and adjoining States, where it still grows, but since the ice retreated it has spread far into Alaska, and eastward across Canada to Ontario and even to New Brunswick.

The sketch was made from specimens gathered at Cottongrass Lake, north of Ptarmigan Pass, ten miles north of Lake Louise Station, Alberta, at an altitude of 7,000 feet.

VERNAL IRIS

Iris verna Linnaeus

Although the vernal iris is not very well known, it is so lovely a wildling that it should find a home in our wildflower gardens. In his "Book of the Iris", Dykes reports his inability to cultivate it, but this was probably owing to his failure to recognize its peculiar soil requirements. It usually grows on wooded hillsides, preferring a mediacid soil. Though the whole plant is only about six inches tall, its violet-blue flowers with yellow markings at the base of the petals cannot be easily overlooked, for they are very large in proportion to the grasslike leaves. In early spring the blossoms dot the barren fields of the Southern States, where real blue violets do not occur, and are often called "violets." Curiously enough, their delicate fragrance adds to the deception, for it closely resembles the scent of the garden violet.

This iris ranges over most of Alabama and Georgia and the states adjoining them on the north, but farther northward it becomes rare, occurring in but a few favored localities in Kentucky, Maryland, and Pennsylvania.

The specimen painted was found near Beaufort, South Carolina.

ANEMONELLA

Syndesmon thalictroides (Linnaeus) Hoffmannsegg

Anemonella, often called rue-anemone, frequents open woods, where it is found in company with hepatica and bloodroot when the first warm days of spring call sleeping plants to awaken. The slender stems, bearing the fragile green leaves and delicate pink or white flowers, grow from a cluster of small tuberous roots. These hold so lightly to the soil that they are easily pulled up, unless care is taken in plucking the flowering stems. Anemonella, the only member of its genus, resembles somewhat its relatives, the true anemones or windflowers, but differs in having the flowers in clusters, and the leaves divided into rounded leaflets. It yields readily to cultivation and is a real desideratum for the wildflower garden.

Anemonella is a wide-ranging species, extending from Florida to New Hampshire and westward to Kansas and Minnesota.

The specimen sketched was found at Washington, District of Columbia.

WHITE TROUTLILY

Erythronium albidum Nuttall

White troutlily is one of the daintiest of our early spring flowers, and east of the Appalachian Mountains is much rarer than the yellow species, which it closely resembles in habit. The shining green leaves are marked with pale mottlings of brownish purple, or in exceptional plants are entirely green. The flowers open only in bright sunlight, and speedily close and wither when gathered. It is known in some sections as dogtooth violet, but as it has no relation to the Violet Family, the name troutlily, suggested by John Burroughs, seems most appropriate.

White troutlily is the common species in the Middle West, from Texas to Minnesota, and eastward to the Appalachian Mountain region. Here and there its seeds have chanced to cross the mountain barrier and, carried down the streams, have lodged along the banks and established colonies in the Piedmont regions from Georgia to northern New Jersey.

The specimen sketched was obtained in the Potomac Valley above Washington, District of Columbia.

HAZEL ALDER

Alnus rugosa (Du Roi) Sprengel

The alders are widespread shrubs known to all who welcome the coming of spring. With the exception of the skunkcabbage, their blooms are the earliest forerunners of spring, and almost every swamp and sheltered brook where the sun can penetrate is decorated with numerous bushes, hung with purple and yellow pendent catkins. The pollen, furnished by these in large quantities, is blown by the wind, lighting on the little green or purplish knobs of pistillate flowers, which grow into ripe cones in autumn. Several species of the genus *Alnus* contain in their bark a coloring principle of value, dyeing either yellow or orange. With copperas added, a good black may be made from it. Before the coming of the white man with his commercial colors, alder dye was frequently used by the Indians.

There are several species of alder in the United States, the one here represented ranging from Florida to Texas and from Maine to Minnesota.

The sketch was made from a specimen gathered near Washington, District of Columbia.

TARFLOWER

Befaria racemosa Ventenat

While traveling through the South in June, whether by rail or automobile, one finds the green fringe of low shrubbery by the roadsides dotted here and there with the pink starlike blossoms of the tarflower. These shrubs rarely occur in dense vegetation, but usually in such places as will permit them to raise their flowery heads above the grass, weeds, and low shrubbery, to beckon the insect sponsors necessary for their cross pollination.

The species ranges from southern Florida to southeastern Georgia.

The specimen here figured came from a woodland margin along the tracks of the Florida East Coast Railroad near Jacksonville, Florida.

ALPINE FIR

Abies lasiocarpa (Hooker) Nuttall

The alpine fir, commonly called balsam fir in the mountains, is the tree best known to those who follow the trail in the Canadian Rockies; for it is the substitute *par excellence* for a soft hair mattress, if the bed is properly made, and no sleep is more restful than that obtained on such an aromatic couch of "Rocky Mountain feathers". From the numerous blisters in the bark a thick juice may be collected which makes an effective dressing for cuts and bruises. The Canada balsam used by microscopists in mounting specimens is obtained from a related species, the balsam fir. By some of the Indian tribes the fir tree was called "cho-koh-tung", meaning blisters. The wood is soft and coarse-grained and, though it grows rather rapidly, is not of much value commercially. In the lower valleys of the Canadian Rockies the tree attains to some size, and at timber line holds the final outposts with the Lyall larches, often affording a striking exhibit of nature's adaptation to adverse conditions of soil, temperature, and wind.

This fir has a wide range. It survived the glacial period in the southern Rocky Mountains, in northern Arizona and New Mexico, and since the ice retreated has migrated northward along the coast ranges to the Olympic Mountains of Washington, thence into Alaska, and through the eastern Rockies into Alberta, Canada.

The specimen sketched was obtained north of Bow Pass, one day's ride from the Saskatchewan River, at the foot of Pyramid Peak, at 6,000 feet elevation.

OCONEE-BELLS

Shortia galacifolia Torrey and Gray

This evergreen, ground-covering plant was for many years one of America's "lost species". It had been discovered by Michaux, the French botanist, during his travels in the mountain wilderness of North Carolina in 1788, but the exact spot where he found it long remained unknown, and the dried specimen he had collected, preserved in Paris, was the only proof that such a plant existed. In 1877 the plant was rediscovered, and subsequently found to be fairly abundant in a few restricted areas. I have not seen it in the wild, but Dr. Edgar T. Wherry informs me that it thrives in acid soils on steep slopes along shaded mountain brooks, and that the primroselike white flowers, starring the mats of winter-bronzed foliage in March, produce a charmingly beautiful effect. If transplanted into ordinary garden loam it soon dwindles and dies, but if given acid humus soil it is not hard to grow in cultivation. Oconee-bells is known to the mountaineers as the one-flower coltsfoot, but the common name has been given from its abundance in the Oconee Valley.

Oconee-bells is known to grow only in the Blue Ridge and adjoining parts of the Piedmont region in North and South Carolina. A close relative occurs in Japan, however, indicating that before the glacial period the genus must have spread over both northeastern Asia and northern North America. The ice advances no doubt exterminated some of the species, but the two that chanced to migrate far enough to the southward of the line reached by the ice sheets were preserved.

The specimen painted was grown by Dr. Frederick V. Coville, in acid peat-sand soil, in a greenhouse of the United States Department of Agriculture at Washington.

VIRGINIA BLUEBELLS

Mertensia virginica (Linnaeus) De Candolle

The lovely Virginia bluebells, with their exquisite bright green leaves, burst forth in unsuspected places in spring. Each year, after the blooming and seeding season, the plants disappear before midsummer, and not until another winter has passed and the spring sun and rains have enticed them from their hiding places, do we again locate them by their tender bronze-green shoots pushing through the moist earth. They love the rich soil of river meadows or the banks of streams, and cover the ground with their masses of bloom and tender green leaves. Virginia bluebells yield easily to cultivation in a wildflower garden, but they are not so well known as other members of the Borage Family, the forget-me-nots and the heliotropes. In Europe they are much appreciated in cultivated gardens.

Virginia bluebells may be found from Georgia northward to New York and southern Ontario, and westward to Kansas and Minnesota.

The specimen sketched was obtained near Washington, District of Columbia.

BURGESS MILKVETCH

Astragalus bourgovii Gray

Among the numerous members of the Pea Family this dainty plant is one of the most appealing. A number of stems grow from a central point and, falling carelessly all around, give the impression of a loose bouquet, the flowers, on their delicate stems, projecting beyond the leaves. The ponies like to eat them, and perfect specimens are difficult to obtain where horses have had access to the colonies.

Burgess milkvetch has a rather restricted range in the northern Rocky Mountains, from South Dakota and Montana to British Columbia. Usually it is found among partially disintegrated rocks.

This specimen was gathered at Burgess Pass, seven miles by trail from Field, British Columbia, on a northwestern slope, at an elevation of 7,500 feet.

GREENDRAGON

Arisaema dracontium (Linnaeus) Schott

Greendragon is a vigorous plant and decidedly decorative, though its flower lacks the handsomer coloring of the jack-in-the-pulpit, its near relative. It is, however, of similar habit. Its deeply parted leaves growing on a long stem from the solid globular bulb or corm, reach a height of two feet or more. It loves the same rich, moist woods or low ground frequented by the jack-in-the-pulpit, but is rarely so abundant. The berrylike fruits of the greendragon are red-orange in color, and are clustered in heads that become conspicuous in early autumn. They are much more effective in calling attention to the plant than its green flowers or its foliage.

The range of the greendragon is rather wide, extending from Texas to Florida and northward to Minnesota, Ontario, and Maine.

This specimen grew along the Potomac River above Washington, District of Columbia.

TOAD TRILLIUM

Trillium sessile Linnaeus

Unlike most other members of its group, this trillium has a rather pleasant odor, but it does not approach other species in attractiveness, for its curious sessile flowers are dull in color. Normally the petals are of a deep maroon color, but in some plants, which show no differences, they are pale greenish yellow. It prefers deep, moist woods, and is easily overlooked among the fresh growths of April or May.

Toad trillium is a typical member of the flora of the Middle Western States, ranging from Mississippi to Florida, northward to Minnesota and western New York; occasionally it crosses the Appalachian Mountains and pushes down river valleys as far north as Maryland and southernmost Pennsylvania.

The specimen sketched grew on Plummers Island in the Potomac River near Washington, District of Columbia.

SOUTHERN MAGNOLIA

Magnolia grandiflora Linnaeus

This stately magnolia is one of the most conspicuous trees of the southern woods, occasionally attaining a height of seventy feet. It proves hardy in cultivation as far north as Washington. The wonderful creamy flowers, encircled by the large, glossy, evergreen leaves, spring from the ends of almost every branch. Their delightful fragrance immediately attracts attention when the tree is at the height of its glory in June. Several species of beetles patronize the blossoms, enticed by the abundant pollen and nectar.

Today the members of the Magnolia Family found in America are but a remnant of the numerous species that flourished in pre-glacial times from the mid-continental plains to the Arctic Circle. The fossilized trunks found at Amethyst Mountain in Yellowstone Park, with specimens of both fruit and leaves, give an indication of their grandeur and beauty in former ages.

Southern magnolia is native throughout the Southern States from Texas to Florida and northward to Arkansas and southeastern North Carolina. It has been officially adopted as the State flower of both Louisiana and Mississippi.

SOUTHERN MAGNOLIA

Magnolia grandiflora Linnaeus

FRUIT OF PLATE 24

As early autumn approaches, the fruit of the southern magnolia becomes conspicuous, and as it ripens the outer envelope cracks over each seed compartment and the crimson seeds protrude from their downy coverings. At this stage the tree affords a striking contrast of red fruit and glossy green leaves. As time passes, the seeds loosen from the shell, but are still attached to the conelike fruit by a slender white thread about an inch long, finally dropping to the ground when the wind blows them loose.

Southern magnolia is found native throughout the Southern States, from Texas to Florida, northward to Arkansas and southeastern North Carolina.

The specimen painted was taken from the large tree just south of the White House, and was given me by Mrs. Coolidge.

PLATE 24A

TRUMPETLEAF

Sarracenia flava Linnaeus

Anyone who has traveled through the South in spring and has been delighted by the fresh verdure of ferns and trees, cannot fail to have been attracted by the great numbers of these curious plants, which can be observed from the car windows. The trumpetleaf is the largest representative of the Pitcherplant Family, the yellow flowers measuring as much as five inches in diameter, and the yellowish-green tubular leaves, frequently veined with red on the pointed lid, growing two to three feet high. The inside of the leaves near the top has a very smooth and slippery waxed surface, so that flies and other insects that are attracted by an exudation just above the waxed surface, lose their hold and slide to their death in the digestive liquid which the pitchers contain. All the members of this family have arrangements for utilizing the dead insects to their own advantage.

The trumpetleaf grows in damp acid soils, from northern Florida to the southernmost counties of Virginia.

The specimen sketched was brought into flower by Dr. F. V. Coville in a greenhouse of the Department of Agriculture in Washington. These plants are easy to cultivate if placed in a pot filled with acid soil made from a mixture of peat and sand, and this put into another pot of two inches greater diameter, the space between being filled with sphagnum moss. They should be kept in a cool greenhouse.

REDBUD

Cercis canadensis Linnaeus

When the dogwood buds are bursting into bloom in early spring, a companion tree, the redbud, is seen throughout our southern woods, its knotted branches covered with clusters of purplish-pink, pea-shaped flowers. In favorable localities the tree attains a height of forty or fifty feet, though usually much smaller. The leaves, which develop after the flowers, are heart-shaped—quite unlike those of most legumes—and are a glossy green, turning to yellow in autumn. The flowers are well supplied with nectar, and are most attractive to bees, who visit them in large numbers. In some regions the redbud is known as Judas-tree, a name properly applied to the Old World species, *Cercis siliquastrum*.

The tree has a wide range, from Texas to Florida and northward. Though best developed in the South, it has been able to migrate north as far as Minnesota and northern New Jersey, and even a short distance into southern Ontario.

The specimen drawn was collected near Washington, District of Columbia.

TALL LARKSPUR

Delphinium elongatum Rydberg

Tall larkspur is found in rich mountain valleys, where it loves to grow among the willow clumps that partially shade the soil about its roots. It is usually from two to four feet in height, and if free from other plants has a striking clump of basal leaves, above which rise the long spikes of rich purplish-blue flowers, swaying in the breezes. The power to fertilize themselves having been lost, cross-pollination is effected by bees and butterflies, whose tongues reach into the deep recesses of the flowers where the nectar is hidden. The name *Delphinium* was given by Linnaeus, from a fancied resemblance of the parts of the flower to a dolphin.

Tall larkspur has a narrow range, being found only from Colorado to Alberta.

The specimen drawn was collected on the Clearwater River, fifty miles north of Lake Louise Station, Alberta, at an altitude of 4,500 feet.

RHODORA

Rhodora canadensis Linnaeus

Rhodora is surrounded with romantic interest, because of the attention drawn to it by Emerson's verses. It is the only plant of its genus. The rosy-purple flowers, usually appearing before the leaves, burst suddenly into bloom and form masses of color on wet hillsides or along the margins of acid swamps. Their shape suggests a relationship to the rhododendron, with which this plant is grouped by some botanists. Rhodora bushes grow from one to three feet high, and are inconspicuous except when in flower.

The range of rhodora is rather limited; presumably it survived the glacial period near the margins of the ice sheets, and it now occupies glaciated territory from northeastern Pennsylvania to Newfoundland.

The specimen sketched was collected at Pocono Manor, Pennsylvania.

THE RHODORA

In May, when sea-winds pierced our solitudes,
I found the fresh Rhodora in the woods,
Spreading its leafless blooms in a damp nook,
To please the desert and the sluggish brook.
The purple petals, fallen in the pool,
Made the black water with their beauty gay;
Here might the red-bird come his plumes to cool,
And court the flower that cheapens his array.
Rhodora! if the sages ask thee why
This charm is wasted on the earth and sky,
Tell them, dear, that if eyes were made for seeing,
Then Beauty is its own excuse for being:
Why thou wert there, O rival of the rose!
I never thought to ask, I never knew;
But in my simple ignorance suppose
The self-same Power that brought me there brought you.

RALPH WALDO EMERSON.

PICKERELWEED

Pontederia cordata Linnaeus

On the flats bordering the Anacostia River in Washington, the rank growths of the various marsh plants form a broad belt of vegetation of striking appearance. Perhaps because of their color, the flowers of pickerelweed are not so conspicuous as those of some other members of the colony, but in spite of this its glossy green leaves, borne well above the surface of the water, and its ragged spikes of small blue or lavender flowers, continuing in bloom through most of the summer, are a delight to behold. Without a boat it is almost impossible to gather the blooms, growing from the soft mud that supports the outer phalanxes of aquatic plants. The flowers of the pickerelweed, which last but a single day, are in three forms. They do not produce seeds without the aid of insects, and it is an interesting study to examine them in detail with a glass and note the complicated structure by which they are able to obtain the greatest amount of benefit from their insect visitors. Giulio Pontedera, professor of botany at Padua about 1730, is commemorated by the generic name.

Pickerelweed is found from Florida to Texas and northward to Minnesota and Nova Scotia, and grows also in tropical America.

The specimen sketched was obtained near Washington, District of Columbia.

CREEPING HOLLYGRAPE

Berberis repens Lindley

The shiny, prickly leaves of the hollygrape are so nearly like holly, that upon first glance we may think we have found that tree, reduced to a bush or low plant creeping over the ground. Closer inspection reveals many differences, for the berries, growing in a bunch, are of a lovely blue color—almost like small grapes—with plenty of bloom. They are rather pungent and sour to the taste, but are good for jelly or for making a refreshing drink, most welcome to quench the thirst when climbing a mountain side. As autumn approaches, the leaves change to red, either all over or around their borders, and are then most attractive in coloring. They remain upon the plant all winter.

Creeping hollygrape is found from California and New Mexico to Alberta and British Columbia. The Oregon hollygrape, *Berberis aquifolium*, often known as Oregon grape, has been designated the State flower of Oregon, by vote of the legislature. Three species of hollygrape occur abundantly in the Pacific Northwest.

The specimen sketched was gathered in Sinclair Canyon, Columbia River Valley in British Columbia, at an altitude of 3,500 feet.

RED CHOKEBERRY

Aronia arbutifolia (Linnaeus filius) Elliott

The red chokeberry is a small shrub found in low grounds, swamps, and dry open woods, and though not particular as to moisture in choosing a situation, usually prefers rather acid soils. The cymes and the under surface of the leaves are woolly, and in spring, when in full bloom, it is a graceful and handsome plant.

Examination of the individual flowers will show them to resemble those of the pear tree in structure, and the plant is referred by botanists to that group of the Rose Family which includes the pear, apple, and hawthorn.

Red chokeberry may be found from Louisiana and Florida to Minnesota and Nova Scotia.

The flowers sketched grew near Washington, District of Columbia.

RED CHOKEBERRY

Aronia arbutifolia (Linnaeus filius) Elliott

FRUIT OF PLATE 3 I

The fruit of the red chokeberry ripens in August or September, and later in the season the whole bush becomes a gorgeous mass of orange and scarlet color. The fruits are shaped like miniature pears, and this, together with their bright coloring, often tempts the stranger to taste them, but the result is sure to be a disappointment, for they are excessively sour as well as bitter. The birds do not seem to mind this feature, however, and consume them greedily during the winter months. A bush or two should be planted in the garden of every bird lover.

The red chokeberry ranges throughout the eastern United States and into southernmost Canada.

Like the flowers, the spray of berries sketched is from the vicinity of Washington, District of Columbia.

PLATE 3 I A

DEVILSCLUB

Echinopanax horridum (Smith) Decaisne and Planchon

The devilsclub is a sturdy shrub, rising above the ground four or five feet on stout prickly stems. The maplelike leaves spread from the stems about a foot below the red berries, and hide the ground, so that one is not aware of its terrible spines until he tries to walk through a thicket. No heavy woolen garment or leather shoe is stout enough to withstand the bristling, hard, gray spines. Without the help of an axe and a strong arm behind it, the thickets of devilsclub in the Selkirk Mountains are often almost impenetrable. The scarlet berries are very showy in late summer.

This plant belongs to the Aralia Family, which includes other spiny shrubs, such as the Herculesclub of the southeastern United States, as well as ginseng and wild-sarsaparilla.

The range of the devilsclub is from Oregon, Montana, and Michigan to Alaska.

The specimen sketched was obtained near Field, British Columbia, at an altitude of 4,000 feet.

CRESTED IRIS

Iris cristata Aiton

The crested iris is dwarf in habit and its lance-shaped leaves of bright green taper at each end. The flowers, which appear in May, seem out of proportion to the size of the plant. Their broad outer divisions or petals have yellow, raised flutings along the center, which give rise to the name. This lovely plant yields easily to cultivation in a wild garden, being relatively indifferent as to soil reaction, but requiring plenty of humus and enough rocks to insure good drainage.

The crested iris may be found from Georgia to Maryland and west to Missouri and southern Indiana.

This specimen was sketched at Plummers Island in the Potomac River near Washington, District of Columbia.

LILY TWAYBLADE

Liparis liliifolia (Linnaeus) Richard

This little orchid is inconspicuous where it grows, on account of its protective coloring, which blends so perfectly with its surroundings, though a fascinating plant when examined closely. It blooms in May and has two rich green leaves that clasp the flower stem and grow from a solid perennial bulb. It is the easiest to cultivate of all our native orchids, not being particular as to the reaction of the soil, although it prefers an abundance of humus.

The lily twayblade extends from Missouri to Georgia and northward, and since the ice of the glacial period retreated has succeeded in migrating as far north as Minnesota and Maine.

The specimen sketched grew on High Island in the Potomac River near Washington, District of Columbia.

MISSOURI PRICKLYPEAR

Opuntia polyacantha Haworth

The Missouri pricklypear is one of the commonest of the cactuses. The flowers are so beautiful that one forgets the wicked spines and the still more dangerous spicules, which are found on the thick stems or flat joints, until one endeavors to gather them. In the bright sunshine of early morning the dry prairies in some sections are dotted with masses of pale sulphur-yellow flowers, turning salmon in the late afternoon as they fade. The sensitive stamens are sometimes yellow and sometimes red, while the stigma-lobes, in the center, are always green. The irritable stamens are one of the provisions which nature makes to bring about cross-pollination. Bees visit the cactus flowers to obtain the nectar. When one of them settles upon the stamens, which spread widely apart in bright sunlight, these at once turn inward and downward, covering the insect and depositing the pollen on its back, legs, and head. This pollen is then carried by the insect to the next flower and dropped upon its stigma-lobes. Below the bright-colored petals is the spiny ovary, which ripens into a dry, many-seeded fruit, this pricklypear being one of the few which is not juicy.

The Missouri pricklypear may be found in dry places from New Mexico, Missouri, and Wisconsin to Alberta and British Columbia.

The specimen sketched was obtained near Medicine Hat, Alberta, Canada, at an altitude of 3,500 feet, a locality near the northern range of this species and also near the northern limit of the Cactus Family.

BUTTERFLYWEED

Asclepias tuberosa Linnaeus

The butterflyweed is probably our most beautiful orange-colored wild flower. Growing in poor sandy soil, it forms masses of brilliant color wherever it finds a congenial foothold. The plants are visited by hosts of butterflies, some of them especially adapted to the cross-pollination of the flowers, which have entirely lost the ability to fertilize themselves. Only insects with long tongues can reach the nectar hidden in the deep recesses of the complicated blossoms, whose structure definitely places the plant in the Milkweed Family. Curiously enough, however, its juice is not milky. The Indians are said to have used the long roots of this and other milkweeds as a remedy for various maladies, and herb doctors of a later day use them under the name of pleurisy-root. Linnaeus dedicated the family to Aesculapius on account of its alleged healing qualities, though the name, as he spelled it, is a Latinized corruption. The plant yields easily to cultivation, provided it is planted in dry, sterile soil, and should be grown in every wildflower garden with bright sunny exposure.

Butterflyweed has a wide range, extending from Florida to northern Mexico and north to Maine, Ontario, Minnesota, and Colorado.

The specimen sketched was found near Washington, District of Columbia.

SKUNKCABBAGE

Spathyema foetida (Linnaeus) Rafinesque

This first plant to flower in spring is easily found in swampy places, where the richly colored, hooded spathe is seen pushing its way through the moist earth, sometimes before the ground is free of snow. When one peeps inside the hood, the small flowers are found dotting the spadix more or less regularly. The rank insistent odor attracts many flesh flies. The leaves come through the earth before the flower has faded, and grow rapidly into showy, light green clumps, from one to three feet high. At this time the odor is very strong. The fruit ripens in September.

Like most of the members of the Arum Family, the skunkcabbage is rather southern in its distribution, ranging from Missouri to Florida and northward, locally reaching Minnesota and Maine, or even Nova Scotia.

The specimen sketched was found along Piney Branch, Washington, District of Columbia.

RED-OSIER DOGWOOD

Cornus stolonifera Michaux

The red-osier dogwood is an attractive shrub, not only when in bloom but also in the early fall, when its bunches of bluish-white berries are borne in abundance, and the leaves change in color to scarlet, purple, or gold. The Indians used the scraped inner bark for smoking purposes, and preferred it to any other plant, giving it the name "kinnikinnick". In winter the slender, graceful, purplish-red stems are conspicuous and distinctive.

Red-osier dogwood may be found from Virginia to Newfoundland, and from Mexico to Alaska, and at many places in the interior of the continent.

The specimen sketched grew at Radium Hot Springs, in the Columbia River Valley, British Columbia, at 3,000 feet elevation.

BIRDSFOOT VIOLET

Viola pedata Linnaeus

The birdsfoot violet is one of the most beautiful members of the Violet Family. The velvety purple color of the two upper petals, combined with the pale blue-violet shade of the three lower ones, contrasted with the bright orange anthers nestling in the center, immediately attracts the attention of all who love wild flowers. Bumblebees, also yellow butterflies, visit the flowers and partake of the sweets prepared for their enjoyment. When soil and exposure are favorable, the ground is purple with the lovely blossoms, the absence of scent being the only flaw in their perfection.

This plant prefers a dry situation where the soil is poor, sterile, and acid, and should not be transferred to a wildflower garden unless satisfied in this respect, for in ordinary rich loam it will promptly die. It will thrive best if planted in coarse gravel richly charged with decaying wood and so rendered thoroughly acid. The small bulblike rootstock is so poorly anchored in the soil and so easily lifted from its moorings, that frequently it comes up when one attempts to pick the flowers. Great care should therefore be taken in gathering them.

This violet is found from Louisiana to Florida, Minnesota, and Massachusetts, the dark purple variety being much more frequent in the southern portion of the range.

The specimen sketched was obtained near Washington, District of Columbia.

VIRGINIA SPIDERWORT

Tradescantia virginiana Linnaeus

The Virginia spiderwort loves the rich borders of woods and thickets, and is loveliest in the morning, when its blue petals open wide, showing the golden anthers in sharp contrast. The plant continues in bloom for some time, new flowers opening each morning, and enticing the bees to a fresh feast of nectar. These insect visitors are necessary to cross-fertilization, and carry from one blossom to the next a heavy load of pollen.

The genus is named for John Tradescant, gardener to King Charles I of England, and belongs in the same family as the dayflower. This and the Pickerelweed Family are closely related.

The Virginia spiderwort may be found from Virginia to Arkansas, northward to South Dakota and southern New York.

The specimen sketched grew near Washington, District of Columbia.

LEATHERFLOWER

Clematis viorna Linnaeus

The leatherflower is not so showy as the other members of the clematis group, nevertheless its graceful stems and leaves and reddish purple blossoms are not uninteresting. The sepals are remarkable in being a sixteenth of an inch thick, as if made of leather instead of the delicate tissue usual in flowers in which petals are lacking and the sepals are required to take their place. The feathery fruit, more attractive than the flowers, is erect and silky, each individual "seed" possessing a tail almost two inches long, by means of which it may be carried by the wind, when ripe, to a favorable situation.

This scrambling vine may be found in thickets where the soil is rocky but rather rich, from Alabama and Georgia northward to Ohio and Pennsylvania.

The sketch was made from a specimen gathered near Washington, District of Columbia.

PURPLE SAXIFRAGE

Saxifraga oppositifolia Linnaeus

The purple saxifrage is truly an alpine plant. In the Canadian Rockies it is found above tree line near the melting snows on southern slopes, where it comes quickly into bloom in the long days of early summer, and as quickly is past, leaving the matted rosettes of tiny, bright green leaves, dotted with dull red seed capsules, to reward our search, if, perchance, we are a few days too late to find it in bloom. It delights in disintegrated limestone, often growing in cracks between the rocks. Though plentiful in its chosen situation, it is not often seen by visitors, who must make a hard climb to reach its haunts. In Alaska and near the Arctic Circle it finds the most favorable conditions for perfect development, growing into large mats.

The purple saxifrage is typical of Arctic regions throughout the world, and no doubt survived the glacial period close to the margins of the great ice sheets. When these withdrew, it followed back on the bare rock surfaces. It is now found occasionally in the northern United States, from Wyoming to Vermont, and more abundantly northward.

The specimen sketched grew on the slopes of Fossil Mountain near Baker Lake, seven miles northeast of Lake Louise Station, Alberta, at an altitude of 8,000 feet.

M.Y.W.

FLAME AZALEA

Azalea lutea Linnaeus

The flame azalea is the most conspicuous of the native members of this genus. Its brilliant orange or yellow coloring, frequently suffused with red, attracts attention even from a distance, and when the mountain sides are flecked with it, the effect is particularly striking. The green leaves, which are well developed before the flowers open, add greatly to the beauty of the plant. It yields easily to cultivation and under favorable conditions, when grown with other plants requiring an acid soil, is one of the loveliest native shrubs for planting.

Flame azalea is a typical member of the flora of our Southern Appalachians, ranging from the uplands of Georgia northward, and being abundant as far as West Virginia. In the mountains of Pennsylvania it is rare, and though reported many years ago to grow in southern New York State, it has long since disappeared from there as a native plant.

The sketch was made from a specimen obtained near Linville, North Carolina.

RABBITBEAN

Cracca virginiana Linnaeus

The gray-green foliage of the rabbitbean makes a delightful background for its straw-colored, pea-shaped flowers, touched with red, and crowded at the end of a stiff stem. The plant blooms in early summer and is to be found in dry, sandy, acid soil. Should you wish to gather a bunch, you will find the stems and the unusually long roots surprisingly tough, and a serviceable substitute for twine, if necessity demands. It is, indeed, commonly known in the South as devil's shoestring.

The seeds are in bean-shaped pods, which are frequently rifled by weevils, so that few of them ever reach maturity.

The Indians used a tropical American species of this genus as fish poison in the same manner in which they employed many other plants, throwing the macerated stems into quiet streams or ponds, with the result that the fish became stupefied and floated on the surface of the water, so that they were easily taken.

Rabbitbean ranges from Texas to Florida and northward to Manitoba and Ontario.

The specimen sketched was collected near Washington, District of Columbia.

TULIPTREE

Liriodendron tulipifera Linnaeus

This beautiful tree is a joy to behold at any season of the year. Its bare gray branches outlined against the winter sky impress one with their vigor of growth and sturdy health, and when the warmer days and April showers swell the buds, and the tiny young leaves appear, the tree bursts into its supreme glory. If in a sunny situation, it is soon covered with green tulip-shaped flowers, each cup beautifully decorated with a brilliant orange bar, to entice the visiting bees. In autumn the leaves are a mass of orange and gold.

The tuliptree is the only member of the genus *Liriodendron* in America, though a closely related species flourishes in central China, these two being the sole survivals of the Cretaceous Period, when members of the genus were widely distributed in America and Europe. Though it reaches its greatest size in the lower Ohio Basin, sometimes growing almost 200 feet high, it is found in the eastern United States from Louisiana to Florida and northward to Michigan and southern Vermont.

The wood has a variety of commercial uses, while the bark, especially that of the roots, yields a tonic and heart stimulant. The blossom of the tuliptree is the State flower of Indiana.

The specimen sketched was obtained at Washington, District of Columbia.

TRUMPET HONEYSUCKLE

Lonicera sempervirens Linnaeus

While tramping through open woods or along streams in low ground, the attention may be attracted to a bit of red color quite in contrast to the brown leaves and gray tree trunks or low-growing bushes. Closer inspection reveals the gray stems that lead to the leafy shoots above, where the delicate trumpet-shaped flowers are clustered at the ends of the branches. Few vines are more attractive; the plant seems to poise itself in the most graceful way as it climbs from one supporting shrub to another.

The flower is a great favorite with humming birds, which frequently are seen probing the trumpets to obtain the delicious nectar to be found in them.

The fruit is a brilliant scarlet berry. The plant is a great addition to any wild garden.

Trumpet honeysuckle ranges from Texas to Florida, and northward to Nebraska, its northern limit being reached in New York and southern New England.

The specimen sketched was collected at Yemassee, South Carolina.

RED BUCKEYE

Aesculus pavia Linnaeus

The red buckeye is a straggling, inconspicuous shrub in the southern woods until it comes into bloom, when it immediately attracts attention by its bright red flowers which are borne in a loose spike. The stems of the new growths also are red at blooming time, and the tender green leaves with their red stems make an appropriate setting for the flower spikes. It is usually a shrub three or four feet high, but in favorable situations it sometimes attains tree size, although never approaching the stature of its relatives, the Ohio buckeye and the horsechestnut.

Red buckeye is a southern species, ranging from Texas to Florida, and extending northward as far as southern Missouri and southeastern Virginia.

The specimen sketched was collected near Beaufort, South Carolina.

ROSE PAINTBRUSH

Castilleja pallida (Linnaeus) Kunth

Crossing alpine meadows above tree line and on high passes in the Canadian Rockies, one often finds the drier ground covered with masses of rose paintbrush, growing in company with saxifrages and forget-me-nots, wherever soil resulting from the disintegrating rock has been deposited. The rose paintbrush varies from two to eight inches in height, according to altitude and local conditions. The leafy bracts, often mistaken for petals, range in color from greenish-white or pale yellow to various shades of mulberry or dull pink. The actual flowers have a dull-colored corolla, and are concealed between the bracts. The plant belongs to the Figwort Family.

Rose paintbrush grows from Alberta and British Columbia to Alaska, and also in Siberia.

The specimen sketched was obtained on the Clearwater River thirty miles north of Lake Louise Station in British Columbia, at an altitude of 8,000 feet.

SHOOTINGSTAR

Dodecatheon meadia Linnaeus

Shootingstar is not particular in choosing a habitat, for it is found in open woods, on moist hillsides, or where the meadows broaden into prairies. It belongs to the Primrose Family, and is distantly related to the Asiatic cyclamen familiar in cultivation, being in fact sometimes called wild cyclamen. It thrives in rich garden soil, where the flowers often become larger than in the wild state, and it should find a place in our gardens.

This species of shootingstar has a rather wide distribution, ranging through the prairies, where the flowers are often red, from Texas to Georgia, and northward into Manitoba; it also extends eastward through the Appalachians, reaching the Piedmont from North Carolina to Pennsylvania. The genus *Dodecatheon* is best represented in the West, where many species occur.

The specimen sketched was obtained near Washington, District of Columbia.

PRAIRIE PENTSTEMON

Pentstemon erianthera Pursh

Prairie pentstemon is one of a large genus of plants, embracing about one hundred and forty species, all of which are natives of North America. Wherever present they are likely to be found in great numbers, especially in the West, where they are very conspicuous. This species grows in small clumps on dry hillsides. Its large flowers are wonderfully varied in coloration, ranging from blue, through various shades of dull purple, to pale pink. They belong in the Figwort Family.

The prairie pentstemon may be found from Nevada, Nebraska, and North Dakota to British Columbia and Washington.

The specimen sketched was obtained near Sinclair Hot Springs, in the Columbia River Valley, British Columbia, at an elevation of 2,500 feet.

WILD SWEET CRAB

Malus coronaria (Linnaeus) Miller

What sweeter blossoms are to be found in spring than those of the wild sweet crab? It is usually a low bushy tree, growing perhaps twenty feet high, with tangled branches making a flat top, quite inconspicuous in its wild surroundings. But when early spring is past, and the bushes are masses of tender green, the pink buds appear, and soon the whole is covered with lovely flowers, whose sweet scent is wafted far on every gentle breeze. The fruit, also, is sweet-scented, and can be made into a delicious jelly.

The wild sweet crab is a native of the Central States, extending from Louisiana and Alabama northward to Michigan and Ontario, but in cultivation over a much wider range.

This specimen was collected near Washington, District of Columbia.

PITCHERPLANT

Sarracenia purpurea Linnaeus

The pitcherplant is found in peat bogs, where the acid soil provides ideal conditions for its successful growth. The pitcher-shaped leaves grow from a central crown, and are partially filled with water and a digestive substance. The inner surfaces of the leaves are lined with bristles which project downward, and these prevent many small insects that slide into the water from escaping. Thus they are trapped, drowned, and digested. These dead insects are also used as food by the larvae of several species of sarcophagous flies, which are instrumental in the cross-pollination of the flower, and are always found where the pitcher-plant grows.

The flowers are borne on a stem from six inches to two feet tall, a vigorous plant often yielding seven or eight blossoms.

A cool greenhouse is necessary for successful indoor cultivation. It is well to place the plant in a flower pot filled with acid soil made from a mixture of peat and sand, and to set this pot inside another of two inches greater diameter, the space between the two being filled with sphagnum moss, which should be kept moist.

The pitcherplant is one of the plants that was pushed southward by the ice cap during glacial times. Since the great ice sheet retreated it has been gradually moving northward, and now ranges from Florida to Kentucky and Iowa, and north to Labrador and Manitoba.

The specimen sketched was collected in eastern Maryland.

PRAIRIE-SMOKE

Sieversia ciliata (Pursh) Don

This pretty and graceful plant grows in grassy meadows, where its red flowers, which resemble buds, its crimson stems, and plumed fruit immediately attract attention. In late summer the silky fruit heads give to dense patches of the plants dispersed over the plains, a purplish hazy appearance, when viewed from a distance.

The plant belongs to that group of the Rose Family which includes the cinquefoils and barren-strawberries, and is most nearly related to the geums, being in fact listed in some books as *Geum triflorum*.

As its name suggests, this plant is typical of the prairie region of the United States, ranging from New Mexico to Missouri and northward. It extends well up into the eastern valleys of the Rocky Mountains, as far north as Alberta, and has also migrated across southern Canada and the northern United States as far as Maine.

The specimen sketched was collected near Banff, Alberta, Canada, at an altitude of 4,000 feet.

WINTERBERRY

Ilex verticillata (Linnaeus) Gray

In spring, when all nature is awakening, we easily overlook this inconspicuous shrub, from four to six feet tall, with tender leaves and tiny flowers, growing in damp acid ground among the other denizens of neglected places. But when autumn passes and the stems are stripped of their leaves, the fine red berries come into their own, and lend a delightful touch of color to the winter landscape. The whole top of the bush is covered with berries, and it then vies with its cousin, the holly, in beauty and interest, and in radiating Christmas cheer and good will.

The berries stay on the branches till late winter, and are rarely eaten by birds. A form with yellow fruit has been found in New England.

Another member of the genus *Ilex* is the well-known shrub, maté, whose leaves are largely used in South America for making a beverage similar to Chinese tea. From yaupon, an *Ilex* of our Southern States, the Indians make a stimulating drink, and it is now being placed on the market, under the name Cassina, as a substitute for tea.

Winterberry ranges from Missouri to Florida, and northward to Wisconsin and Nova Scotia.

The specimen sketched was collected near Washington, District of Columbia.

SWEET AZALEA

Azalea arborescens Pursh

The sweet azalea is one of the contributions to our flora from the Appalachian Mountains, where so many attractive plants were found by the early botanists. Its delightfully fragrant, spicy, white flowers appear in June when the leaves are well developed and afford a fine green background for the blossoms. The foliage often turns brilliant red in late autumn.

The plant loves a well drained, deep, moist, acid soil, and under favorable conditions becomes a spreading clump, which blooms freely. It is perhaps the largest of our native azaleas, occasionally attaining a height of twenty feet.

Sweet azalea occurs in the mountains from Georgia northward into Pennsylvania.

The sketch was made from a plant found in the vicinity of Linville, North Carolina, at the foot of Grandfather Mountain.

KRUHSEA

Kruhsea streptopoides (Ledebour) Kearney

Anyone who has visited Glacier House, and walked to the foot of the Illecillewaet Glacier, may have noticed beds of shiny green-leaved plants suggestive of a small Solomonseal. The tiny greenish, bell-shaped flowers hang from the under side of the stem, however, and are usually missed by the transient visitor, whose efforts are centered merely on climbing the trail to reach the edge of the ice. This rare member of the Lily-of-the-valley Family has a somewhat restricted range, and is especially interesting because it is one of those plants common to both the American and Asiatic shores of the Pacific Ocean.

It may be found from northern Washington to southern Alaska and in the Selkirks of British Columbia; also in Japan and Siberia.

The specimen sketched grew at Glacier, British Columbia, at an elevation of 3,500 feet.

KRUHSEA

Kruhsea streptopoides (Ledebour) Kearney

FRUIT OF PLATE 56

The change wrought in vegetation during a few weeks of growing weather is very great in mountain regions at the latitude of Glacier House, and when autumn approaches, the tiny green flowers of Kruhsea have developed rapidly into large, pulpy, red berries, which contrast strikingly with the yellow leaves and soon fall to the ground.

On this continent Kruhsea grows from northern Washington to southern Alaska, and in Asia it is recorded from both Japan and Siberia.

The specimen sketched was obtained at the same locality as the flowers shown in the preceding plate, at Glacier, British Columbia, at 3,500 feet elevation.

PLATE 56A

ARETHUSA

Arethusa bulbosa Linnaeus

The arethusa has a single large, one-sided flower with a delicate scent like that of fresh red raspberries. The recurved lip, with its fringes, forms a capital landing platform for visiting insects. As with many other orchids, the bees are frequent visitors to this plant. The flower is so constructed that the bee, in raising his head to depart after sipping the nectar, comes in contact with a few soft pellets of pollen, which are deposited upon his head from the helmet-shaped anther. Some of this pollen may be transferred to the stigmas of the next flower that he visits, although more often it is brushed off by other parts of the flower. Because of the infrequency of cross-pollination seeds rarely mature.

The name was given to the plant by Linnaeus, who recalled the myth of the nymph Arethusa, changed by Diana into a fountain, in order to protect her from the river god Alpheus, who fell deeply in love with her on seeing her at her bath.

Owing to the great demand for this orchid by European collectors, it has been nearly exterminated in many sphagnum bogs where it formerly grew in great abundance.

Arethusa may be found from North Carolina northward to Maine and Newfoundland, and westward to Indiana and Minnesota.

The specimen sketched was obtained from a swamp a few miles east of Washington, District of Columbia, where the plant is extremely rare.

PALE LADYSLIPPER

Cypripedium acaule Aiton

The pale ladyslipper is even lovelier than the common form of this beautiful orchid, which has pink flowers. The pale form is frequent in the North, but one who is so fortunate as to find it in the more southern part of its range, experiences a thrill that is not likely to be forgotten. If the soil is kept strongly acid, the plant will readily yield to culture in a wildflower garden, and will flourish in either dry or moist situations, and in sun or shade. A dressing of pine needles or oak leaves will help to conserve the moisture, as well as the acidity of the soil. No attempt should be made, however, to cultivate it in ordinary garden soil, for there it is sure to die.

The pale ladyslipper may be found from North Carolina and Tennessee northward to Manitoba and Newfoundland.

The specimen sketched grew near Washington, District of Columbia.

QUAKERLADIES

Houstonia caerulea Linnaeus

Quakerladies, sometimes called bluets and innocence, are among our earliest spring flowers, and delight flower lovers by their dainty growth and faint, sweet odor. Occurring plentifully in moist meadows and wayside places, they sometimes completely carpet the ground. They may be gathered freely, without fear of extermination. They continue blooming into early summer, and frequently put forth a second bloom in late fall. No wildflower garden would be complete without them, a rather sterile and acid soil and a fair amount of sunshine being all that they ask.

Linnaeus named this plant in honor of Dr. William Houston, a young English botanist, who died in South America after an exhausting collecting trip around the Gulf of Mexico in 1733.

Quakerladies are widely distributed, being found from Georgia and Alabama northward to eastern Canada and Michigan.

The plant sketched grew at Washington, District of Columbia.

M. Y. W.

BEARBERRY HONEYSUCKLE

Lonicera involucrata (Richardson) Banks

Bearberry honeysuckle is a plant that is conspicuous in July when the fruit ripens. The coarse, hairy leaves are of a dark rich green, and the twin berries, glossy black in color, are surrounded by a red frill, a combination which makes the bush very decorative. The unpretentious straw-colored flowers may sometimes be found on the younger branches along with the fruit produced by those that come into bloom earlier. If the bush is shaken, the ripe fruit falls easily. Its disagreeable flavor has gained for it the name of skunkberry in some regions.

Bearberry honeysuckle may be found from Quebec to Michigan, New Mexico, and California, and northward to Alaska.

The specimen sketched was collected near Hector, British Columbia, at an elevation of 4,000 feet.

PINK FLEABANE

Erigeron caespitosus Nuttall

The pink fleabane may be seen in great perfection in June in the upper Columbia River Valley. It seems to delight in dry, sandy soils, and especially after a shower, its graceful clumps of daisylike flowers, variously shaded from white to pink or pale purple, are a delight to the eye. It is a member of a group of plants widely distributed in North America, and represented in the Rocky Mountains by a hundred or more species.

Pink fleabane ranges from Colorado and Utah to the Yukon.

The specimen drawn was collected in the Saskatchewan River Valley, fifty miles north of Lake Louise, Alberta, at an altitude of 3,500 feet.

LABRADOR-TEA

Ledum groenlandicum Oeder

Who that has traveled by the Canadian Pacific Railway in June, along the north shore of Lake Superior and across the boggy country traversed by that railroad, has not noticed the masses of low bushes covered with feathery heads of white flowers? When the mountains are reached we still find the shrub growing luxuriantly in full sunshine or adapting itself to more shaded situations, provided the ground is sufficiently wet. The margins of the leaves are rolled, and their under surface is covered with brown wool.

The leaves have an aromatic fragrance, and were used by the early settlers as a substitute for tea, but the beverage is rather too much like turpentine to be palatable.

Labrador-tea is one of the members of the Heath Family and is at home in northern regions. During the glacial period it probably survived near the margins of the ice sheets, and when these melted back, followed them closely and became widespread in the glaciated territory, wherever acid soils developed. It now grows from the highlands of New Jersey northward and westward far into the Rocky Mountains and the Arctic regions. It was in fact given the specific name *groenlandicum* because it was first discovered in Greenland.

The specimen sketched came from the White Mountains in New Hampshire.

NORTHERN BEDSTRAW

Galium boreale Linnaeus

In many parts of North America northern bedstraw is a familiar plant. Although essentially northern in its distribution, as its name implies, it is not an Arctic plant, but seems to thrive best in regions of only moderately cold climate. In the central and southern Rockies it is abundant at middle and high altitudes, but farther north it frequents the foothills, extending out upon the plains. The plant sometimes forms dense clumps which afford a bouquet of feathery white flower sprays that are delicately scented. In the Rockies the name wild heliotrope is occasionally applied to the plant, although its fragrance is scarcely suggestive of our garden heliotrope.

Northern bedstraw grows in a great variety of situations, on open banks among rocks and grasses, in aspen thickets, or along streams, where the abundant moisture develops luxuriant plants that are sometimes two feet high. The species ranges from Pennsylvania, Missouri, and southern California, northward over the greater part of Canada and Alaska, and is widely distributed also in northern Europe and Asia.

This plant is one of a large group of the Madder Family, some members of which produce, in their roots, a red or purple dye. Most of our American bedstraws are unattractive plants, with weak rough stems and insignificant flowers.

The sketch was made from a specimen found near Banff, Alberta, Canada.

BRONZEBELLS

Stenanthium occidentale Gray

Bronzebells is so delicate and graceful a lily, and has such a modest coloring of green and dark maroon, that it is almost hidden among the vegetation of the moist rich woods, where it prefers to grow. Flowering in company with Rocky Mountain rhododendron and menziesia, it is always a delight to the lover of the beautiful who has eyes to see the variety in nature's handiwork, and will search for her hidden treasures.

Bronzebells ranges from Montana and Oregon to Alberta and British Columbia.

The specimen sketched was obtained in the Yoho Valley, ten miles from Field, British Columbia, at an altitude of 4,500 feet.

ROUNDLEAF ORCHIS

Orchis rotundifolia Pursh

The roundleaf orchis loves the wet shores of alpine lakes and frequently grows in sphagnum moss along the borders of mountain brooks in partially shaded, wet places. Where conditions are favorable it occurs in abundance, though often overlooked by the flower lover because of larger and ranker plants surrounding it. The single rounded leaf is a distinguishing characteristic, and the sweet scent of the flowers is so pervasive that it sometimes attracts attention before the bloom is discovered. Except for a single Alaskan species, it is the only American member of a group represented in Europe and Asia by eighty species or more.

The roundleaf orchis is a plant of northern range and is not known to grow south of the limits reached by the ice sheets of the glacial period. It must have survived close to the edge of the ice, and migrated back rapidly when this retreated. Although it died out from the places where it survived glaciation, it has subsequently spread across the continent, from northern Maine to British Columbia and northward to Greenland and Alaska.

The specimen sketched grew at Emerald Lake, seven miles from Field, British Columbia, at an altitude of 3,800 feet.

PRICKLY CURRANT

Ribes lacustre (Persoon) Poiret

The prickly currant is a beautiful bush when in bloom. Its graceful branches are then ornamented with dainty racemes of yellowish flowers, shaded with red, and the delicate green leaves are of a tint best suited to show them off to perfection. When growing on a steep slope, with full exposure to the sun, the plant is so pleasing that one wonders why it has not been transplanted to cultivated gardens.

Prickly currant is found from Pennsylvania to Newfoundland and from California to Alaska.

The specimen sketched was collected on the slopes of Mt. Wapta, ten miles by trail from Field, British Columbia, at an altitude of 7,000 feet.

PRICKLY CURRANT

Ribes lacustre (Persoon) Poiret

FRUIT OF PLATE 66

Though never very abundant, the prickly currant is found in some regions of the Rocky Mountains in sufficient quantity to furnish a delightful dish. When stewed and eaten with venison or wild mutton, it makes a deliciously spicy sauce. When raw, the berries are rather sour, and are eaten only for want of something better.

Prickly currant may be found from Pennsylvania to Newfoundland and from California to Alaska.

The specimen sketched was obtained near Glacier Lake, on the headwaters of the Saskatchewan River, fifty miles north of Lake Louise, British Columbia, at an altitude of 6,000 feet.

PLATE 67

GLACIERLILY

Erythronium grandiflorum Pursh

The glacierlily seems to radiate the spirit of the high places, and with bright sunshine and pure air helps to entice the lover of nature to the mountain tops. Along the edges of the melting snow the pointed, green, daggerlike leaves push upward, often through the snow itself, and soon the flower bursts into bloom, exhaling a delicious fragrance quite distinct from any other we experienced in the mountains. Carpeting the ground with gold, the plant may be found even in mid-summer, along with springbeauties, yellow violets, and buttercups, where the spring avalanches have lodged their load of snow and thus held the early flowers in cold storage. The name avalanche lily is often applied to it because of this fact. At lower altitudes it grows in greater numbers and with longer stems, but always with the same lovely coloring of brilliant green and gold. It is rarely found below an elevation of 4,000 feet.

The glacierlily is evidently adapted to grow in regions of heavy snowfall, and no doubt survived the glacial period close to the edge of the ice from Wyoming to Washington, having since pushed northward into the mountains of British Columbia and Alberta.

The specimen sketched was obtained on the slopes of Mt. Wapta, above Emerald Lake near Field, in the Canadian Rockies, at an altitude of 6,000 feet.

BALSAMROOT

Balsamorhiza sagittata (Pursh) Nuttall

Wherever it is found, balsamroot is a striking plant. The arrow-shaped leaves, blue-green on the upper surface and white on the under side, are borne on stalks five or six inches long, above which the handsome yellow flowers are poised on still longer stems. When in bloom these plants brighten whole mountain sides with gold. The plant prefers moist situations on partially shaded, steep slopes, but in the lower valleys it frequents the borders of swampy land, among coarse grass and alder and willow bushes. The horses love to feed upon it, and will never pass a fine clump in perfection of leaf and bloom unless urged on. The large fleshy roots are eaten by the Indians, and in Utah they are commonly called Mormon biscuit, because of their use by the early immigrants in times of scarcity.

Balsamroot is distributed from Colorado and California to British Columbia and South Dakota.

The specimen sketched was obtained near Radium Hot Springs, British Columbia, at an altitude of 3,500 feet.

SILVERBERRY

Elaeagnus commutata Bernhardt

After crossing the plains and coming into the foothill country, the traveler will observe many thickets of silvery-gray shrubs. In June, on approaching these clumps he is greeted by the peculiar sweet scent of the greenish funnel-shaped bells, with yellow petals, that hang from the under sides of the branches. The leaves, when examined under a lens, are seen to be covered with silvery scales. Similar scales covering the fruit have given the plant the name silverberry. It is nearly related to the buffaloberry, and often grows with it. Both are members of the group of plants known as the Oleaster Family, which is considered by botanists to belong in the same order as the loosestrifes and evening primroses.

Silverberry is found from Quebec to Minnesota and from Utah to the Yukon.

The specimen sketched was collected near Ghost River, twenty-five miles northeast of Banff, Alberta, at an altitude of 4,000 feet.

M. Y. W.

SILVERBERRY

Elaeagnus commutata Bernhardt

FRUIT OF PLATE 70

We had known the silverberry for years and had enjoyed its sweet fragrance, but had never seen fruit on the bushes until one occasion when we were traveling down the Kootenai River Valley, in the beginning of September. Here all the plants were growing in great perfection, and in riding across a flat, through which a mountain stream meandered, we came across some superb silverberry bushes, which were higher than our horses' heads and loaded with fruit. They were so beautiful that we carried a great bunch of them back to camp, tied to the pommel of the saddle. Since then we have learned that the berries are, in a sense, edible, though too dry and mealy to appeal to the taste of most people, and so the prairie chickens and other birds are allowed to enjoy them in peace, throughout their broad range from Quebec to Minnesota and Utah, and north to the Yukon.

The specimen sketched was obtained in the Kootenai River Valley, British Columbia, at an altitude of 3,000 feet.

PLATE 71

TWINLEAF

Jeffersonia diphylla (Linnaeus) Persoon

Jeffersonia, commonly called twinleaf, is one of the earliest of spring flowers. It blooms before the leaves are fully developed, and is somewhat like a bloodroot in appearance. The long-stemmed leaf blades are parted nearly to the base, and as they are blown about by the wind, remind one of a group of green butterflies, though they lose this resemblance when fully developed, for they are then rather stiff. This plant, which was named for Thomas Jefferson, can be easily cultivated in a wildflower garden under the same conditions that will render the bloodroot happy.

Jeffersonia is most abundant west of the Appalachian Mountains, from Tennessee northward to Minnesota, Ontario, and central New York; locally, however, it has crossed the mountain barrier and pushed down the river valleys, notably along the Potomac River in Virginia and Maryland.

The sketch was made from a beautiful clump growing on Plummers Island in the Potomac River near Washington, District of Columbia.

IVORY BANEERRY

Actaea arguta Nuttall

Ivory baneberry is a white-fruited form of the red-fruited western baneberry, *Actaea arguta*. It prefers shady situations, near the borders of mountain streams, where the soil is rich and the air damp. The flowers, which have a sweet, sickish odor, appear in spring in racemes two inches long at the ends of the slender stems. The white petals and sepals soon fall, and the raceme lengthens. When the berries develop and become heavy, the stems bearing them appear almost too weak to carry the load, and the fruits soon drop when ripe. The berries are said to be poisonous, like the seeds of many other members of the Buttercup Family, to which the plant belongs.

Ivory baneberry is found from Utah and Colorado to Alberta and British Columbia.

The specimen sketched was collected near Vermilion Pass on the motor road between Banff and the Columbia River Valley, at an altitude of 4,000 feet.

M.Y.W.

PINK MOUNTAINHEATHER

Phyllodoce empetriformis (Smith) Don

The mountainheathers of the Canadian Rockies are an inspiration and a delight to those who climb to the higher places. They are found near tree line and on the bare slopes just above it, and cover the mountain sides with their deep green foliage. In blossom time they make a wonderful carpet of color—pink, red, and white. Among them all, the pink mountainheather is in many respects the most beautiful.

On Burgess Pass, seven miles by trail from Field, British Columbia, where this specimen was gathered at an altitude of 7,000 feet, the coloring was almost like that of an India shawl. The seed-pods are deep red and covered with golden dots of resin.

Pink mountainheather ranges from California to Colorado and northward to Alaska.

M.Y.W.

ROCKY MOUNTAIN CASSIOPE

Cassiope mertensiana (Bongard) Don

With the exception of the rare orchids, the Rocky Mountain cassiope is perhaps the most romantic of American mountain flowers. It is found in perfection about tree line in the Canadian Rockies, which is usually at an altitude of 6,500 to 7,500 feet. Here, in favorable localities, it frequently forms dense, thick mats, and when in full bloom the plants are literally covered with delicate white bells. The tiny stems holding the blossoms may be either green or red, and when the flowers fall the seed capsules are usually red in color, becoming brown as the season advances. Growth in a single season is rather limited, and the stiff, woody stems underneath bear silent witness to the vicissitudes of plants at high altitudes. To the camper a bed of cassiope is most satisfying, for in addition to its springiness it possesses an elusive fragrance that persists even when the plants are quite dried out. The red, pink, and white mountainheathers are frequently found growing with the cassiope, and add color to the wild-flower carpet on the mountain sides.

Rocky Mountain cassiope is to be found from Montana to northern California and Alaska.

The specimen sketched was obtained at Burgess Pass near Field, British Columbia, Canada.

ONE-LEAF BOG-ORCHID

Habenaria obtusata (Pursh) Richardson

This orchid is exceptional in having but a single basal leaf. The plant grows on the mossy banks of streams or in boggy places, delighting in the peaty soil, where other members of the genus *Habenaria* flourish. Its green color renders it inconspicuous and easily overlooked, especially when the larger, white, sweet-scented varieties are its near neighbors.

This species ranges from Maine, New York, and Colorado and northward to Newfoundland and Alaska.

This specimen was found at Hillsdale, eleven miles northwest of Banff, Alberta, at an altitude of 4,000 feet.

M. Y. W.

BUSH CINQUEFOIL

Potentilla fruticosa Linnaeus

Bush cinquefoil is one of the showiest plants of the Canadian Rockies, where it seems to find congenial surroundings almost everywhere, flourishing in alpine meadows or on mountain slopes, and sometimes at high elevations, struggling against the mountain winds and other adverse conditions to obtain a foothold. In midsummer it is covered with clear yellow flowers, which appear to greater advantage by reason of the silver-gray foliage that forms their background. In moist limestone soils in New England it is classed as a weed, because of its invasion of idle fields and pastures.

This member of the Rose Family is found from California, New Mexico, and New Jersey northward to Alaska, central Canada, and Labrador, as well as in Europe and Asia.

The specimen sketched was collected on the trail below Burgess Pass, six miles from Field, British Columbia, at an altitude of 5,000 feet.

M. Y. W.

<u>Scientific Name</u>	<u>Volume</u>	<u>Plate No.</u>
Abies lasiocarpa	I	18
Acer carolinianum	II	138
Acer rubrum	II	137
Achillea lanulosa	II	151
Actaea arguta	I	73
Aesculus pavia	I	47
Agoseris Graminifolia	II	88
Agoseris gracilens	II	89
Agoseris villosa	III	195
Allium cernuum	IV	304
Allium sibericum	V	383
Alnus rugosa	I	16
Alnus sinuata	III	186
Amelanchier	II	117
Anaphalis margaritacea	IV	289
Androsace carinata	II	107
Androsace subumbellata	III	185
Anemone deltoidea	IV	306
Anemone globosa	IV	291
Anemone parviflora	V	371
Anisostichus capreolatus	IV	261
Antennaria howellii	II	104
Antennaria luzuloides	III	171
Antennaria rosea	IV	286
Aquilegia brevistyla	IV	292
Aquilegia canadensis	II	141
Aquilegia flavescens	III	201
Arctostaphylos uva-ursi	II	111
Arctostaphylos uva-ursi	II	112
Arctous alpina	V	355
Arethusa bulbosa	I	57
Arisaema dracontium	I	22
Arisaema triphyllum	V	331
Arnica louiseana	I	10
Arnica tomentosa	V	348
Aronia arbutifolia	I	31
Aronia arbutifolia	I	31a
Artemisia discolor	IV	288
Asarum canadense	II	127
Asclepias speciosa	II	90
Asclepias tuberosa	I	36
Asimina triloba	V	328
Aster squarrosus	II	160
Astragalus alpinus	IV	295
Astragalus bourgovii	I	21
Atamasco atamasco	IV	255
Aster campestris	II	118
Azalea arborescens	I	55
Azalea lutea	I	43
Azalea nudiflora	II	128
Azalea rosea	III	232
Azalea vaseyi	IV	253

Balsamorhiza sagittata	I	69
Befaria racemosa	I	17
Berberis repens	I	30
Bignonia radicans	III	227
Bikukulla canadensis	II	136
Bikukulla cucullaria	IV	247
Calla palustris	II	129
Callicarpa americana	III	210
Calochortus catalinae	III	205
Calochortus clavatus	IV	314
Calochortus elegans	I	2
Calochortus kennedyi	V	391
Calochortus macrocarpus	III	175
Calochortus splendens	IV	310
Calochortus weedii	III	199
Caltha leptosepala	IV	287
Caltha palustris	III	208
Campanula lasiocarpa	V	368
Campanula rotundifolia	V	369
Capnoides sempervirens	I	5
Carex aurea	IV	281
Cassiope mertensiana	I	75
Castalia odorata	III	223
Castilleja lancifolia	II	102
Castilleja miniata	V	372
Castilleja pallida	I	48
Centaurium venustum	IV	303
Centiana glauca	II	108
Cercis canadensis	I	26
Chelone glabra	IV	259
Chenopodium capitatum	V	349
Chimaphila umbellata	V	360
Chionanthus virginica	II	140
Chrosperma muscaetoxicum	II	147
Chrysamphora californica	V	390
Chrysogonum virginianum	II	145
Cirsium hookerianum	II	103
Cirsium undulatum	IV	309
Claytonia parvifolia	II	94
Claytonia virginica	III	234
Clematis crispa	II	150
Clematis columbiana (flower)	II	99
Clematis columbiana (fruit)	II	100
Clematis viorna	I	41
Clintonia borealis	V	338
Clintonia uniflora (flower)	III	203
Clintonia uniflora (fruit)	III	204
Collinsia verna	IV	246
Comandra livida	V	361
Conopholis americana	III	214
Cornus canadensis (flower)	IV	271
Cornus canadensis (fruit)	IV	272
Cornus florida (flower)	V	321
Cornus florida (fruit)	V	322
Cornus stolonifera	I	38
Cracca virginiana	I	44

Cypripedium acaule	I	58
Cypripedium acaule	V	327
Cypripedium arietinum	III	216
Cypripedium montanum	I	1
Cypripedium parviflorum	II	92
Cypripedium passerium	II	91
Cypripedium reginae	III	217
Cyrtopodium punctatum	III	212
Cytherea bulbosa	II	105
Delphinium depauperatum	V	384
Delphinium elogatum	I	27
Dendromecon rigidum	V	392
Dentaria laciniata	IV	249
Dionaea muscipula	III	219
Diplacus longiflora	III	200
Diplacus puniceus	IV	315
Dodecatheon meadia	I	49
Dodecatheon pauciflorum	IV	276
Dryas drummondii (flower)	V	364
Dryas drummondii (fruit)	V	365
Dryas octopetala (flower)	III	176
Dryas octopetala (fruit)	III	177
Echinocereus lloydii	II	155
Echinocereus viridiflorus	IV	308
Echinopanax horridum	I	32
Elaeagnus commutata (flower)	I	70
Elaeagnus commutata (fruit)	I	71
Empetrum nigrum	V	382
Epidendrum nocturnum	V	337
Epidendrum tampense	II	152
Epigaea repens	II	126
Epilobium angustifolium	IV	301
Epilobium latifolium	V	370
Epilobium luteum	IV	300
Erigeron aureus	IV	280
Erigeron caespitosus	I	61
Erigeron salsuginosus	IV	290
Erigeron speciosus	III	165
Erigeron unalaschensis	V	376
Eriophorum angustifolium	IV	312
Eriophorum chamissonis	I	12
Eritrichum elongatum	V	359
Erythronium albidum	I	15
Erythronium americanum	V	339
Erythronium grandiflorum	I	68
Erythronium montanum	III	202
Eschscholtzia mexicana	V	393
Fouquieria splendens	V	396
Fragaria glauca	V	362
Franklinia alatamaha	IV	244
Fremontodendron mexicanum	III	206

Gaillardia aristata	V	354
Galium boreale	I	63
Gaylussacia brachycera	III	229
Gelsemium sempervirens	III	220
Gentiana acuta	IV	294
Gentiana affinis	II	87
Gentiana calycosa	IV	318
Gentiana critita	V	336
Gentiana porphyrio	I	8
Gentiana prostrata	III	178
Gentiana saponaria	III	161
Geranium viscosissimum	IV	307
Habenaria ciliaris	V	340
Habenaria grandiflora	IV	243
Habenaria lacera	III	215
Habenaria obtusata	I	76
Hamamelis virginiana	V	323
Hedysarum mackenzii	II	97
Hepatica americana	II	125
Hookera pauciflora	V	389
Houstonia caerulea	I	59
Hymenocallis rotata	II	154
Hypopitys americana	II	157
Hypopitys lanuginosa	III	213
Ibidium cernuum	I	4
Ibidium gracile	I	4
Ibidium strictum	V	356
Ilex opaca	IV	266
Ilex verticillata	I	54
Ilex vomitoria	II	226
Iris cristata	I	33
Iris vericolor	V	332
Iris verna	I	13
Jeffersonia depphylla	I	72
Juniperus horizontalis	V	379
Juniperus sibirica	II	86
Kalmia angustifolia	V	326
Kalmia latifolia	III	221
Kalmia microphylla	IV	284
Kalmia polifolia	II	133
Kruhsea streptopoides (flower)	I	56
Kruhsea streptopoides (fruit)	I	56a
Kraunhia Frutescens	II	149
Lappula diffusa	V	351
Larix lyalli	V	381
Larix occidentalis	IV	269
Lathyrus ochroleucus	II	85
Ledum groenlandicum	I	62
Lepargyrea canadensis	II	115
Lilium canadense	II	148
Lilium columbianum	IV	316

<i>Lilium montanum</i>	I	11
<i>Lilium superbum</i>	IV	256
<i>Limodorum tuberosum</i>	II	131
<i>Linnaea borealis americana</i>	III	196
<i>Liparis lilifolia</i>	I	34
<i>Liriodendron tulipifera</i>	I	45
<i>Lobelia cardinalis</i>	IV	263
<i>Lobelia kalmii</i>	III	192
<i>Lonicera glaucescens</i> (flower)	II	81
<i>Lonicera glaucescens</i> (fruit)	II	82
<i>Lonicera involucrata</i>	I	60
<i>Lonicera serpervirens</i>	I	46
<i>Lupinus perennis</i>	I	6
<i>Lychnis apetala</i>	IV	279
<i>Lygodesmia juncea</i>	I	9
<i>Magnolia acuminata</i>	III	231
<i>Magnolia cordata</i>	V	330
<i>Magnolia grandiflora</i> (flower)	I	24
<i>Magnolia grandiflora</i> (fruit)	I	24a
<i>Magnolia virginiana</i>	V	325
<i>Malus coronaria</i>	I	51
<i>Menyanthes trifoliata</i>	III	225
<i>Menziesia glabella</i>	IV	298
<i>Mertensia paniculata</i>	III	173
<i>Mertensia virginica</i>	I	20
<i>Mimulus caepitosus</i>	V	375
<i>Mimulus cardinalis</i>	V	395
<i>Mimulus guttatus</i>	IV	313
<i>Mimulus lewisii</i>	V	374
<i>Mitchella repens</i>	III	207
<i>Monarda punctata</i>	III	233
<i>Moneses uniflora</i>	IV	273
<i>Monotropa uniflora</i>	IV	262
<i>Myosotis alpestris</i>	V	352
<i>Nymphaea advena</i>	II	159
<i>Ophrys nephrophylla</i>	II	109
<i>Opuntia polycantha</i>	I	35
<i>Orchis rotundifolia</i>	I	65
<i>Orchis spectabilis</i>	IV	241
<i>Orontium aquaticum</i>	III	222
<i>Orthocarpus tenuifolius</i>	II	119
<i>Oxycoccus palustris</i>	III	180
<i>Oxtropis gracilis</i>	I	3
<i>Oxytropis podocarpa</i> (flower)	IV	282
<i>Oxytropis podocarpa</i> (fruit)	IV	283
<i>Oxytropis spendens</i>	II	120
<i>Pachyloplus hirsutus</i>	V	388
<i>Pachyloplus marginatus</i>	V	387
<i>Parnassia fimbriata</i>	V	343
<i>Passiflora incarnata</i>	V	324
<i>Pedicularis bracteosa</i>	IV	258
<i>Pedicularis contorta</i>	II	121
<i>Pedicularis groenlandica</i>	V	373
<i>Pentstemon confertus</i>	V	357
<i>Pentstemon digitalis</i>	III	209

Pentstemon eatonii	V	394
Pentstemon erianthera	I	50
Pentstemon lyallii	V	346
Pentstemon menziesii	IV	319
Pentstemon wrightii	V	386
Peramium decipiens	V	350
Petalostemon purpureum	IV	320
Petasites hyperboreus	III	189
Phacelia linearis	III	197
Phacelia sericea	IV	278
Pjlox divaricata	IV	245
Phoradendron flavescens	IV	265
Phyllodoce empetrifolmis	I	74
Physaria didymocarpa (flower)	III	167
Physaria didymocarpa (fruit)	III	168
Picea engelmanni	V	378
Pinguicula elatior	III	235
Pinguicula vulgaris	III	198
Pinus albicaulis	V	377
Pinus contorta murrayana	II	101
Pinus flexilis	III	188
Pinus palustris	II	139
Pinus taeda	V	335
Podophyllum	II	143
Pogonia divaricata	IV	242
Pogonia ophioglossoides	III	218
Polycodium stamineum	II	132
Polygala lutea	IV	257
Polygala paucifolia	II	135
Pontederia cordata	I	29
Porteranthus trifolius	IV	252
Potentilla fruticosa	I	77
Potentilla glaucophylla	IV	296
Potentilla uniflora	III	182
Primula maccalliana	IV	274
Prunus angustifolia	II	130
Pseudotsuga mucronata	IV	270
Pulsatilla ludoviciana (flower)	II	95
Pulsatilla ludoviciana (fruit)	II	96
Pulsatilla occidentalis (flower)	III	163
Pulsatilla occidentalis (fruit)	III	164
Pyrola chlorantha	II	113
Pyrola minor	III	172
Pyrola secunda	IV	285
Pyxidanthera barbulata	II	124
Quamasia quamash	V	358
Ranunculus suksdorfii	II	114
Rhododendron albiflorum	IV	305
Rhododendron maximum	IV	254
Rhodora canadensis	I	28
Ribes lacustre (flower)	I	66
Ribes lacustre (fruit)	I	67
Romanzoffia sitchensis	II	98

Rosa bourgeauiana (flower)	V	344
Rosa bourgeauiana (fruit)	V	345
Rubus argutus	II	146
Rubus parviflorus	V	341
Rubus pedatus	IV	293
Sabbatia stellaris	V	342
Sagittaria cuneata	II	158
Salix discolor	II	122
Salix drummondiana	V	380
Salix nivalis	IV	277
Salix petrophila	II	106
Sambucus pubens	IV	260
Sanguinaria canadensis	II	123
Sarracenia catesbaei	V	400
Sarrezenia drummondii	V	329
Sarrezenia flava	I	25
Sarrezenia minor	IV	251
Sarracenia purpurea	I	52
Sarrezenia psittacina	III	236
Sarracenia rubra	IV	250
Saussurea densa	III	191
Saxifraga bronchialis	II	110
Saxifraga caespitosa	III	184
Saxifraga lyallii	V	363
Saxifraga oppositifolia	I	42
Senecia triangularis	V	385
Senecio lugens	IV	275
Senecio pauciflorus	II	93
Scutellaria serrata	II	237
Shortia galicifolia	I	19
Sieversia ciliata	I	53
Silene acaulis	V	367
Silene caroliniana	IV	248
Sisyrinchium angustifolium	III	238
Solidago ciliosa	III	183
Sorbus sambucifolia	III	162
Spathyema foetida	I	37
Sphaeralcea davidsonii	IV	311
Sphaeralcea grossulariaefolia	V	398
Stenanthium occidentale	I	64
Stewartia malachodendron	V	333
Streptopus amplexifolius	II	84
Streptopus curvipes	II	83
Syndesmon thalictroides	I	14
Thalesia uniflora	II	156
Thermopsis rhombifolia	IV	297
Thuja plicata	III	187
Tillandsia fasciculata	II	153
Tillandsia fasciculata	V	399
Tradescantia virginiana	I	40
Trillium chloropetalum	IV	299
Trillium album	V	334
Trillium erectum	III	239
Trillium grandiflorum	III	240
Trillium sessile	I	23
Trillium undulatum	II	134

Trollius albiflorus	V	353
Tsuga heterophylla	IV	268
Tsuga mertensiana	IV	267
Uvularia perfoliata	II	144
Vaccinium caepitosum	III	179
Vaccinium corymbosum	III	228
Vaccinium membranaceum	I	7
Vaccinium scoparium (flower)	III	169
Vaccinium scoparium (fruit)	III	170
Vaccinium tenellum	III	230
Vaccinium vitisidaea minus (flower)	III	193
Vaccinium vitisidaea minus (fruit)	III	194
Vagnera stellata	III	166
Valeriana sitchensis	V	366
Veratrum viride	III	174
Viburnum pauciflorum	IV	317
Vicia americana	III	190
Viola adunca	III	181
Viola canadensis	V	347
Viola eriocarpa	III	224
Viola pedata	I	39
Viola refinesquii	III	211
Viola septemloba	II	142
Xerophyllum tenax	IV	302
Yucca baileyi	V	397
Zenobia cassinifolia	IV	264
Zygadenus elegans	II	116

Common Name	Volume	Plate No.
Agoseris, Grassleaf	II	88
Agoseris, Slender	II	89
Agoseris, Woolly	III	195
Alaska Fleabane	IV	290
Alberta Paintbrush	V	372
Alberta Primrose	IV	274
Alder, Hazel	I	16
Alder, Western Green	III	186
Aleutian Fleabane	V	396
Alpine Fernleaf	II	121
Alpine Fir	I	18
Alpine Forget-me-not	V	352
Alpine Harebell	V	368
Alpine Milkvetch	IV	295
Alpine Monkeyflower	V	375
Alpine Pointvetch (flower)	IV	282
Alpine Pointvetch (fruit)	IV	283
American Columbine	II	141
American Holly	IV	266
American Mistletoe	IV	265
American Pasqueflower (flower)	II	95
American Pasqueflower (fruit)	II	96
American Twinflower	III	196
American Vetch	III	190
American Waterlily	III	223
American Wistaria	II	149
Androsace, Pygmy	III	185
Androsace, Sweet	II	107
Anemone, Forest	IV	306
Anemone, Globe	IV	291
Anemone, Northern	V	371
Anemone, Plume (flower)	III	163
Anemone, Plume (fruit)	III	164
Anemonella	I	14
Arborvitae, Giant	III	187
Arethusa	I	57
Arnica, Lake Louise	I	10
Arnica, Woolly	V	348
Arrowhead, Arum	II	158
Arrowleaf Groundsel	V	385
Arum Arrowhead	II	158
Aster, Pineland	II	160
Aster, Prairie	II	118
Atamasco-lily	IV	255
Avalanche Buttercup	II	14
Avalanche-lily	III	202
Azalea, Pinkshell	IV	253
Azalea, Sweet	I	55

Balsamroot	I	69
Baneberry, Ivory	I	73
Bearberry (flower)	II	111
Bearberry (fruit)	II	112
Bearberry Honeysuckle	I	60
Bearcabbage, Green	III	174
Beargrass	IV	302
Beautyberry	III	210
Bedstraw, Northern	I	63
Beebalm, Spotted	III	233
Birdsfoot Violet	I	39
Blackberry, Highbush	II	146
Bladderpod, Double (flower)	III	167
Bladderpod, Double (fruit)	III	168
Bloodroot	II	123
Bluebead	V	338
Bluebells, Virginia	I	20
Bluebells, Western	III	173
Blueberry, Highbush	III	228
Blueberry, Pineland	III	230
Blue-eyed-grass	III	238
Blue-eyed-mary	IV	246
Blueflag Iris	V	332
Bog Kalmia	II	133
Bogbean	III	225
Bog-orchid, One-leaf	I	76
Bottle Gentian	III	161
Bourgeau Rose (flower)	V	344
Bourgeau Rose (fruit)	V	345
Bowmansroot	IV	252
Box Huckleberry	III	229
Brook Lobelia	III	192
Bronzebells	I	64
Buckeye, Red	I	47
Buffaloberry, Canada	II	115
Bunchberry (flower)	IV	271
Bunchberry (fruit)	IV	272
Bur-forget-me-not	V	351
Burgess Milkvetch	I	21
Bush Cinquefoil	I	77
Bushpoppy	V	392
Butterbur, Northern	III	189
Buttercup, Avalanche	II	14
Butterflyweed	I	36
Butterwort, Northern	III	198
Butterwort, Purple	III	235
California Pitcherplant	V	390
Calla, Wild	II	129
Calypso	II	105
Campion, Nodding	IV	279
Camas	V	358
Campion, Moss	V	367

Canada Buffaloberry	II	115
Canada Lily	II	148
Canada Violet	V	347
Canada Wildginger	II	127
Cardinal Monkeyflower	V	395
Cardinalflower	IV	263
Carolina Jessamine	III	220
Carolina Maple	II	138
Cassiope, Rocky Mountain	I	75
Catalina Mariposa	III	205
Catesby Pitcherplant	V	400
Centaurium, Pink	IV	303
Chickasaw Plum	II	130
Chokeberry, Red (flower)	I	31
Chokeberry, Red (fruit)	I	31a
Cinquefoil, Bush	I	77
Cinquefoil, Rosette	III	182
Clematis, Columbia (flower)	II	99
Clematis, Columbia (fruit)	II	100
Clematis, Curly	II	150
Clusterlily	V	389
Columbia Clematis (flower)	II	99
Columbia Clematis (fruit)	II	100
Columbia Lily	IV	316
Columbine, American	II	141
Columbine, Lemon	III	201
Columbine, Shortspur	IV	292
Comandra, Red	V	361
Cottongrass	I	12
Cottongrass, Tassel	IV	312
Crab, Wild Sweet	I	51
Cranberry, Mountain (flower)	III	193
Cranberry, Mountain (fruit)	III	194
Cranberry, Small	III	180
Cranberrybush	IV	317
Cranesbill, Western	IV	307
Creeping Hollygrape	I	30
Creeping Juniper	V	379
Crested Iris	I	33
Crossvine	IV	261
Crowberry	V	382
Crowpoison	II	147
Cucumbertree	III	231
Cucumbertree, Yellow	V	330
Currant, Prickly (flower)	I	66
Currant, Prickly (fruit)	I	67
Cut Toothwort	IV	249
Cyrtopodium, Spotted	III	212
Dawnrose, White	V	387
Deathcamus	II	116
Deerberry	II	132
Devilsclub	I	32
Dewberry, Red	IV	293
Dogwood, Flowering (flower)	V	321
Dogwood, Flowering (fruit)	V	322
Dogwood, Red-osier	I	38

Double Bladderpod (flower)	III	167
Double Bladderpod (fruit)	III	168
Douglas-fir	IV	270
Douglas Honeysuckle (flower)	II	81
Douglas Honeysuckle (fruit)	II	82
Downy Pinxterbloom	III	232
Drummond Pitcherplant	V	329
Drummond Willow	V	380
Dutchmans-breeches	IV	247
Dryad, White (flower)	III	176
Dryad, White (fruit)	III	177
Dryad, Yellow (flower)	V	364
Dryad, Yellow (fruit)	V	365
Elder, Scarlet	IV	260
Elephanthead	V	373
Elkslip	IV	287
Engelmann Spruce	V	378
Epidendrum, Tampa	II	152
Epidendrum, White	V	337
Evening-primrose	V	388
Everlasting, Pearl	IV	289
False Locoweed	I	3
Fernleaf, Alpine	II	121
Field Violet	III	211
Fir, Alpine	I	18
Fire Pentstemon	V	394
Fireweed	IV	301
Fivefinger, Grayleaf	IV	296
Flame Azalea	I	43
Fleabane, Alaska	IV	290
Fleabane, Aleutian	V	376
Fleabane, Golden	IV	280
Fleabane, Meadow	III	165
Fleabane, Pink	I	61
Flowering Dogwood (flower)	V	321
Flowering Dogwood (fruit)	V	322
Flytrap, Venus	III	219
Forest Anemone	IV	306
Forget-me-not, Alpine	V	352
Forget-me-not, Moss	V	359
Foxglove Pentstemon	III	209
Franklinia	IV	244
Fremontia, Mexican	III	206
Fringed Gentian	V	336
Fringed Parnassia	V	343
Fringe-Orchid, Large Purple	IV	243
Fringe-orchid, Ragged	III	215
Fringeorchid, Yellow	V	340
Fringetree	II	140
Fumeroot, Pink	I	5

Gaillardia, Perennial	V	354
Gentian, Bluegreen	II	108
Gentian, Bottle	III	161
Gentian, Fringed	V	336
Gentian, Ladder	IV	294
Gentian, Moss	III	178
Gentian, Pinebarren	I	8
Gentian, Riverbank	II	87
Gentian, Ruff	IV	318
Ghostpipe	II	156
Giant Arborvitae	III	187
Giant Trillium	IV	299
Glacierlily	I	68
Globe Anemone	IV	291
Globeflower, White	V	353
Globemallow, Orange-eye	IV	311
Globe-mallow, Scarlet	V	398
Golden Fleabane	IV	280
Golden Sedge	IV	281
Goldenbowl Mariposa	IV	314
Goldclub	III	222
Goldenpea	IV	297
Goldenrod, Sidesaddle	III	183
Goldenstar	II	145
Grassleaf Agoseris	II	88
Grayleaf Fivefinger	IV	296
Greendragon	I	22
Groundsel, Arrowleaf	V	385
Groundsel, Mourning	IV	275
Groundsel, Rayless	II	93
Grouse Whortleberry (flower)	III	169
Grouse Whortleberry (fruit)	III	170
Harebell	V	369
Harebell, Alpine	V	368
Hazel Alder	I	16
Heliotrope Valerian	V	366
Hemlock, Mountain	IV	267
Hemlock, Western	IV	268
Hepatica	II	125
Highbush Blackberry	II	146
Highbush Blueberry	III	228
Holly, American	IV	266
Hollygrape, Creeping	I	30
Honeysuckle, Bearberry	I	60
Honeysuckle, Douglas (flower)	II	81
Honeysuckle, Douglas (fruit)	II	82
Honeysuckle, Trumpet	I	46
Hooded Ladies-tresses	V	356
Hooded Pitcherplant	IV	251
Huckleberry, Box	III	229
Indianpipe	IV	262
Iris, Blueflag	V	332
Iris, Crested	I	33
Iris, Vernal	I	13
Ivory Baneberry	I	73
Jack-in-the-pulpit	V	331

Jessamine, Caroline	III	220
Juniper, Creeping	V	379
Juniper, Mountain	II	86
Kalmia, Bog	II	133
Kalmia, Rocky Mountain	IV	284
Kruhsea (flower)	I	56
Krushea (fruit)	I	56a
Labrador-tea	I	62
Ladder Gentian	IV	294
Ladies-tresses, Hooded	V	356
Ladies-tresses, Nodding	I	4
Ladies-tresses, Slender	I	4
Ladyslipper, Mountain	I	1
Ladyslipper, Northern	II	91
Ladyslipper, Pale	I	58
Ladyslipper, Pink	V	327
Ladyslipper, Ramshead	III	216
Ladyslipper, Showy	III	217
Ladyslipper, Small Yellow	II	92
Lake Louise Arnica	I	10
Lambkill	V	326
Lanceleaf Paintbrush	II	102
Larch, Lyall	V	381
Larch, Western	IV	269
Large Purple Fringe-Orchid	IV	243
Larkspur, Slim	V	384
Larkspur, Tall	I	27
Leatherflower	I	41
Lemon Columbine	III	201
Lewis Monkeyflower	V	374
Lilac Mariposa	IV	210
Lily, Canada	II	148
Lily, Columbia	IV	316
Lily, Red	I	11
Lily, Turkscap	IV	256
Lily Twayblade	I	34
Limber Pine	III	188
Lloyds Strawberry-cactus	II	155
Lobelia, Brook	III	192
Loblolly Pine	V	335
Locoweed, False	I	3
Lodgepole Pine	II	101
Longleaf Pine	II	139
Low Whortleberry	III	179
Lupine, Sun-dial	I	6
Lyall Larch	V	381
Magnolia, Southern (flower)	I	24
Maple, Carolina	II	138
Maple, Red	II	137
Mariposa, Catalina	III	205
Mariposa, Goldenbowl	IV	314
Mariposa, Lilac	IV	310
Mariposa, Sagebrush	III	175
Mariposa, San Diego	III	199
Mariposa, Scarlet	V	391

Marshmarigold	III	208
Mayapple	II	143
Maypop	V	324
Meadow Fleabane	III	165
Menzies Pentstemon	IV	319
Menziesia, Western	IV	298
Marrybells, Wood	II	144
Mexican Fremontia	III	206
Mexican Poppy	V	393
Milkvetch, Alpine	IV	295
Milkvetch, Burgess	I	21
Milkweed, Showy	II	90
Missouri Pricklypear	I	35
Mistletoe, American	IV	265
Mistmaiden	II	98
Monkeyflower, Alpine	V	375
Monkeyflower, Buff	III	200
Monkeyflower, Cardinal	V	395
Monkeyflower, Lewis	V	374
Monkeyflower, Red	IV	315
Monkeyflower, Western	IV	313
Moss Champion	V	367
Moss Forget-me-not	V	359
Moss Gentian	III	178
Mountain-ash, Western	III	163
Mountain Cranberry (flower)	III	193
Mountain Cranberry (fruit)	III	194
Mountain Hemlock	IV	267
Mountain Juniper	II	86
Mountain Ladyslipper	I	1
Mountain-laurel	III	221
Mountainheather, Pink	I	74
Mourning Groundsel	IV	275
Naiad Springbeauty	II	94
Nodding Champion	IV	279
Nodding Ladies	I	4
Nodding Onion	IV	304
Northern Anemone	V	371
Northern Bedstraw	I	63
Northern Butterbur	III	189
Northern Butterwort	III	198
Northern Ladyslipper	II	91
Oconee-bells	I	19
Ocotillo	V	396
One-leaf Bog-orchid	I	76
Onion, Siberian	V	383
Orange-eye Globemallow	IV	311
Orchid, Grass-pink	II	131
Orchid, Rosebud	IV	242
Orchis, Roundleaf	I	65
Orchis, Showy	IV	241
Owl-clover	II	119
Oxytrope, Showy	II	120

Paintbrush, Alberta	V	372
Paintbrush, Lanceleaf	II	102
Paintbrush, Rose	I	48
Painted Trillium	II	134
Pale Ladyslipper	I	58
Pale Pinesap	II	157
Papaw	V	328
Parrot Pitcherplant	III	236
Partridgeberry	III	207
Pasqueflower, American (flower)	II	95
Pasqueflower, American (fruit)	II	96
Pea, White	II	85
Pearl Everlasting	IV	289
Peatpink	IV	248
Pentstemon, Fire	V	394
Pentstemon, Foxglove	III	209
Pentstemon, Menzies	IV	319
Pentstemon, Prairie	I	50
Pentstemon, Purple	V	346
Pentstemon, Wright	V	386
Pentstemon, Yellow	V	357
Perennial Gaillardia	V	354
Phacelia, Gray	IV	278
Phacelia, Sand	III	197
Phlox, Blue	IV	245
Pickernelweed	I	29
Pine, Limber	III	188
Pine, Loblolly	V	335
Pine, Lodgepole	II	101
Pine, Longleaf	II	139
Pine, Whitebark	V	377
Pinebarren Gentian	I	8
Pineland Aster	II	160
Pineland Blueberry	III	230
Pinesap, Pale	II	157
Pinesap, Red	III	213
Pinkshell Azalea	IV	253
Pinxterbloom	II	128
Pinxterbloom, Downy	III	232
Pipsissewa, Western	V	360
Pitcherplant	I	52
Pitcherplant, California	V	390
Pitcherplant, Catesby	V	400
Pitcherplant, Drummond	V	329
Pitcherplant, Hooded	IV	251
Pitcherplant, Parrot	III	236
Pitcherplant, Sweet	IV	250
Plum, Chickasaw	II	130
Plume Anemone (flower)	III	163
Plume Anemone (fruit)	III	164
Pogonia, Rose	III	218
Pointvetch, Alpine (flower)	IV	282
Pointvetch, Alpine (fruit)	IV	283
Polygala, Fringed	II	135
Polygala, Orange	IV	257
Poppy, Mexican	V	393

Prarie Aster	II	118
Prarie Pentstemon	I	50
Prarie-smoke	I	53
Prarie Thistle	IV	309
Prarieclover, Purple	IV	320
Prickly Currant (flower)	I	66
Prickly Currant (fruit)	I	67
Pricklypear, Missouri	I	35
Primrose, Alberta	IV	274
Ptarmiganberry	V	355
Purple Mountain Violet	III	181
Pussy-ears	I	2
Pussy Willow	II	122
Pussytoes, Buff	III	171
Pussytoes, Gray	II	104
Pussytoes, Pink	IV	286
Pygmy Androsace	III	185
Pyrola, Green	II	113
Pyrola, Sidebells	IV	285
Pyrola, Small	III	172
Pyxie	II	124
Quakerladies	I	59
Queencup (flower)	III	203
Queencup (fruit)	III	204
Quill-leaf Tillandsia	V	399
Rabbitbean	I	44
Ragged Fringe-orchid	III	215
Ramshead, Ladyslipper	III	216
Raspberry, Whiteflowering	V	341
Rattlesnakeplantain, Western	V	350
Rayless Groundsel	II	93
Red Buckeye	I	47
Red Chokeberry (flower)	I	31
Red Chokeberry (fruit)	I	31a
Red-helmet	IV	258
Red Lily	I	11
Red Maple	II	137
Red-osier Dogwood	I	38
Red Pinesap	III	213
Redbud	I	26
Redstem Saxifrage	V	363
Rhododendron, Rocky Mountain	IV	305
Rhododendron, Rosebay	IV	254
Rhodora	I	28
Riverbank Gentian	II	87
Rock Willow	II	106
Rock Wormwood	IV	288
Rocky Mountain Cassiope	I	75
Rocky Mountain Kalmia	IV	284
Rocky Mountain Rhododendron	IV	305
Rocky Mountain Twayblade	II	109
Rose, Bourgeau (flower)	V	344
Rose, Bourgeau (fruit)	V	345
Rose Paintbrush	I	48
Rose Pogonia	III	218

Rosebay Rhododendron	IV	254
Rosebud Orchis	IV	242
Rosegentian, Saltmarsh	V	342
Rosette Cinquefoil	III	182
Roundleaf Orchis	I	65
Ruff Gentian	IV	318
Sagebrush Mariposa	III	175
Saltmarsh Rosegentian	V	342
San Diego Mariposa	III	199
Sand Phacelia	III	197
Saskatoon	II	117
Saussurea	III	191
Saxifrage, Purple	I	42
Saxifrage, Redstem	V	363
Saxifrage, Spotted	II	110
Scarlet Elder	IV	260
Scarlet Mariposa	V	391
Sedge, Golden	IV	281
Shootingstar	I	49
Shootingstar, Slender	IV	276
Shortspur Columbine	IV	292
Showy Ladyslipper	III	217
Showy Milkweed	II	90
Showy Orchis	IV	241
Showy Oxytrope	II	210
Siberian Onion	V	383
Sidebells Pyrola	IV	285
Sidesaddle Goldenrod	III	183
Silverberry (flower)	I	70
Silverberry (fruit)	I	71
Skeletonflower	I	9
Skullcap, Wood	III	237
Skunkcabbage	I	37
Slender Agoseris	II	89
Slender Ladies-tresses	I	4
Slender Shootingstar	IV	276
Smooth Yellow Violet	III	224
Snow Trillium	III	240
Snow Willow	IV	277
Solomonplume, Star	III	166
Southern Coast Violet	II	142
Southern Magnolia (flower)	I	24
Southern Magnolia (fruit)	I	24a
Spatterdock	II	159
Spiderlily	II	154
Spiderwort, Virginia	I	40
Spotted Beebalm	III	233
Spotted Cyrtopodium	III	212
Springbeauty, Naiad	II	94
Springbeauty, Virginia	III	234
Spruce, Engelmann	V	378
Squawroot	III	214
Squirrelcorn	II	136
Star Solomonplume	III	166
Stewartia, Virginia	V	333

Strawberry-blite	V	349
Strawberry-cactus Green	IV	308
Strawberry-cactus, Lloyds	II	155
Strawberry, Pale	V	362
Sun-dial Lupine	I	6
Sweet Androsace	II	107
Sweet Azalea	I	55
Sweet Crab, Wild	I	51
Sweet Pitcherplant	IV	250
Sweetbay	V	325
Sweetvetch	II.	97
Tampa Epidendrum	II	152
Tarflower	I	17
Tassel Cottongrass	IV	312
Thistle, Prairie	IV	309
Thistle, White	II	103
Tillandsia	II	153
Tillandsia, Quill-leaf	V	399
Toad Trillium	I	23
Toothwort, Cut	IV	249
Trailing-arbutus	II	126
Trillium, Giant	IV	299
Trillium, Painted	II	134
Trillium, Red	III	239
Trillium, Snow	III	240
Trillium, Toad	I	23
Trillium, Wax	V	334
Troutlily, White	I	15
Troutlily, Yellow	V	399
Trumpet Honeysuckle	I	46
Trumpet creeper	III	227
Trumpetleaf	I	25
Tufted Saxifrage	III	184
Tuliptree	I	45
Turkscap Lily	IV	256
Turtlehead	IV	259
Twayblade, Lily	I	34
Twayblade, Rocky Mountain	II	109
Twinflower, American	III	196
Twinleaf	I	72
Twistedstalk, Clasping	II	84
Twistedstalk, Pink	II	83
Valerian, Heliotrope	V	366
Venus Flytrap	III	219
Vernal Iris	I	13
Vetch, American	III	190
Violet, Birdsfoot	I	39
Violet, Canada	V	347
Violet, Field	III	211
Violet, Purple Mountain	III	181
Violet, Smooth Yellow	III	224
Violet, Southern Coast	II	142
Virginia Bluebells	I	20
Virginia Spiderwort	I	40
Virginia Springbeauty	III	234
Virginia Stewartia	V	333

Waterlily, American	III	223
Wax Trillium	V	334
Western Bluebells	III	173
Western Cranesbill	IV	307
Western Green Alder	III	186
Western Hemlock	IV	268
Western Larch	IV	269
Western Menziesia	IV	298
Western Monkeyflower	IV	313
Western Mountain-ash	III	162
Western Pipsissewa	V	360
Western Rattlesnakeplantain	V	350
Western Yarrow	II	151
Whitebark Pine	V	377
Whiteflowering Raspberry	V	341
Whortleberry, Big	I	7
Whortleberry, Grouse (flower)	III	169
Whortleberry, Grouse (fruit)	III	170
Whortleberry, Low	III	179
Wild Sweet Crab	I	51
Wildginger, Canada	II	127
Willow, Drummond	V	380
Willow, Rock	II	106
Willow, Snow	IV	277
Willowweed, Red	V	370
Willow-weed, Yellow	IV	300
Winterberry	I	54
Wistaria, American	II	149
Witch-hazel	V	323
Wood Merrybells	II	144
Wood Skullcap	III	237
Woodnymph	IV	273
Woolly Agoseris	III	195
Woolly Arnica	V	348
Wormwood, Rock	IV	288
Wright Pentstemon	V	386
Yarrow, Western	II	151
Yaupon	III	226
Yucca	V	397
Zenobia	IV	264

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00640 3661