Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

11622
C3F76
Cap 5

FOREST RESOURCES OF ILLINOIS

CENTRAL STATES
FOREST EXPERIMENT STATION
Columbus 13, Ohio

Harold L. Mitchell, Director

FOREST RESOURCES OF ILLINOIS

BY

THE FOREST SURVEY ORGANIZATION
Central States Forest Experiment Station

in cooperation with

THE AGRICULTURAL EXPERIMENT STATION
University of Illinois

and

THE DIVISION OF FORESTRY
Illinois Department of Conservation

Personnel participating in the collection and compilation of data and preparation of this report.

Central States Forest Experiment Station

R. K. Winters, Chief, Division of Forest Economics

E. V. Roberts, In Charge, Forest Survey

Field Work - M. E. Becker, Field Supervisor

A. E. Block, J. L. Burkle, T. J. Schmitt,

J. E. Wiggins, D. A. Bernstein

Photo Interpretation - K. E. Moessner, In Charge

O. K. Hutchison, K. L. Quigley,

K. W. Chrisemer

Computations - Lake Compton, Margaret Peirsol, Mary L. Sterner

Volume Tables and Statistical Analysis - G. Luther Schnur

Stenographic Service - Virginia K. Tomlinson

Illinois Agricultural Experiment Station

J. Nelson Spaeth, E. P. Van Arsdel

Illinois Division of Forestry

E. E. Nuuttila

C O N T E N T S

	<u>Page</u>
Significant facts - - - - -	1
Forest survey procedure - - - - -	48
Accuracy of data - - - - -	49
Explanation of terms used - - - - -	50

<u>Table</u> <u>number</u>	<u>State</u>	<u>Southern</u> <u>Region</u>	<u>Claypan</u> <u>Region</u>	<u>Prairie</u> <u>Region</u>
	- - -	- - -	Page - - -	- - -
1 Forest and nonforest area by region or county	4	15	26	37
2 Commercial forest area by owner- ship class	5	16	27	39
3 Commercial forest area by forest type and stand-size class	6	17	28	40
4 Saw-timber volume by species and stand-size class	7	18	29	41
5 Saw-timber volume by species and tree diameter	8	19	30	42
6 Hardwood saw-timber volume by log grade	9	20	31	43
7 Total cubic volume by species and class of material	10	21	32	44
8 Cubic volume of growing stock by species and stand-size	11	22	33	45
9 Cubic volume of growing stock by stand-size class and tree diameter	12	23	34	46
10 Average volume per acre	13	24	35	47

FOREWORD

The Forest Survey is a nation-wide activity of the Forest Service. The fivefold purpose of the Forest Survey is: (1) to make a field inventory of the present supply of standing timber; (2) to ascertain the rate at which this supply is being increased through growth; (3) to determine the rate at which it is being diminished through industrial and domestic uses, windfall, fire, disease, and other causes; (4) to determine the present consumption and the probable future trend in requirements for forest products; and (5) to interpret and correlate these findings with existing and anticipated economic conditions, as an aid in the formulation of both private and public policies for use of land suitable for forest production.

The Forest Survey is conducted in the various regions by the forest experiment station of the Forest Service. In Illinois the project is directed by the Central States Forest Experiment Station with headquarters in Columbus, Ohio.

This Survey Release presents the more significant statistics on forest area and timber volume for the state of Illinois and for each of the three regions into which the state has been divided. Later an analytical report for the state will be published which will interpret forest area, timber volume, growth, and drain statistics in the light of existing and anticipated economic conditions.

LOCATION OF FOREST SURVEY REGIONS IN ILLINOIS AND THE PROPORTION OF THE AREA OF EACH COUNTY THAT IS FORESTED

SIGNIFICANT FACTS CONCERNING THE FOREST RESOURCES OF ILLINOIS

The forest area of Illinois, as determined by the Forest Survey, is 3,996,000 acres or 11 percent of the total land area. The proportion of forest land is highest in the Southern region where 26 percent of the land is in forest. Two counties in this region, Pope and Alexander, are more than 40 percent forested. The Claypan region averages 16 percent forest land with one county, Calhoun, reaching 43 percent. In the Prairie region much of the area was open grassland at the time of settlement. Most of the 1.6 million acres of forest land in this region is in the counties bordering the Mississippi and Illinois Rivers, and the remainder occurs principally in small scattered patches.

Ninety-five percent of the commercial forest land in the state is in private ownership. The Shawnee National Forest in the Southern region contains 3.7 percent of the forest land in the state and other federal and state agencies account for 1.1 percent. Most of the private forest land is in small farm woodlands, although some large areas of privately owned forest land occur along the Mississippi, Ohio, Wabash, and Illinois Rivers.

The oak-hickory type occurs on nearly half of the forest area in the state and predominates in the upland areas of all regions. The proportion of oak-hickory type is greatest in the Southern region and least in the Prairie region. The white oak type, which comprises 9 percent of the forest area of the state, is more prevalent in the north than in the south. Similarly the mixed hardwood type, in which elm, basswood, and maple are the principal species, occupies a larger proportion of the forest land in the northern portion of the state. The pin oak type is largely confined to poorly drained areas of the two southern regions. The bottomland hardwood type occurs along the major rivers and makes up nearly one-fifth of the forest area of the state. It is rather uniformly represented in all regions and is one of the most important timber-producing types in the state.

Timber stands with sufficient volume to be classed as saw timber occur on 46 percent of the commercial forest land in the state. More than two-thirds of the saw-timber area contains enough volume in trees 15 inches d.b.h. and larger to be classed as large saw-timber area. Pole-timber stands make up 25 percent of the forest area of the state, seedling and sapling 19 percent, and poorly stocked areas 10 percent. The distribution of forest area by stand-size class varies but little from region to region.

The net volume of saw timber in the state is 10.3 billion board feet. Nearly 70 percent of this volume is found in large saw-timber stands and 19 percent in small saw-timber stands. The remaining 11 percent is in scattered trees in the other stand-size classes. The distribution of volume by stand-size class in each of the regions is similar to the state average.

Oak species make up 56 percent of the board-foot volume in the state and white oak alone accounts for 20 percent of the total. Black oak and northern red oak each make up 10 percent of the total volume. Other species of importance are elm (10 percent), hickory (7 percent), and soft maple (6 percent). In general the proportion of white oak, northern red oak, and elm in the stands increases from south to north. Hickory, on the other hand, is less important in the northern part of the state. Softwood species are of minor importance throughout the state.

Less than one-third of the saw-timber volume in the state is in grade 1 and 2 logs. These logs produce the high-quality material required by the cooperage, veneer, and furniture industries. The large proportion of low-quality timber presents a serious problem in forest utilization and management.

The volume of the forest growing stock--the sawlog portion of merchantable trees of sawlog size and the stems of sound pole-timber trees--is 2.4 billion cubic feet. One-third of this volume is in trees of pole-timber size. In addition to the growing stock there is 1.7 billion cubic feet in tops and limbs of saw-timber trees and in the sound portion of cull trees. Although very little of this is now being utilized it is a large volume of wood awaiting an economic use.

The average volume per acre of all species on all commercial forest land in the state is 2,625 board feet. Areas classed as large saw timber averaged 5,726 board feet per acre and small saw-timber areas averaged 3,424 board feet. The average volume per acre in the three regions were similar to the state average. The average volume per acre of all growing stock was 608 cubic feet with a range from 75 cubic feet per acre in seedling and sapling areas to 1,052 cubic feet in large saw-timber areas.

STATE TABLES

STATE OF ILLINOIS

Table 1.--Forest and nonforest area by regions, 1948

Region	: Total land area ^{1/}	: Forest area	: Nonforest area
	<u>Thousand acres</u>	<u>Thousand acres</u> <u>Percent</u>	<u>Thousand acres</u> <u>Percent</u>
Southern	3,942	1,042 26	2,900 74
Claypan	8,487	1,356 16	7,131 84
Prairie	23,377	1,598 7	21,779 93
All regions	35,806	3,996 11	31,810 89

^{1/} Source: Area of United States 1940, U. S. Bureau of the Census.

STATE OF ILLINOIS

Table 2.--Commercial forest area by ownership class, 1948

Ownership class	Commercial forest area ^{1/}	
	<u>Thousand acres</u>	<u>Percent</u>
Federal:		
National forest	147	3.7
Other	32	.8
<hr/>		
Total	179	4.5
State	10	.3
Private	3,752	95.2
<hr/>		
All ownerships	3,941	100.0

^{1/} Does not include 55 thousand acres of forest land classified as noncommercial.

STATE OF ILLINOIS

Table 3.--Commercial forest area by forest type and stand-size class, 1948

	:		:	Large	:	Small	:	:	Seedling:	
	:		:	saw-	:	saw-	:	Pole-	:	and
Forest	:	Total	:	timber	:	timber	:	timber	:	sapling
type	:		:	area	:	area	:	area	:	area
		Thousand	Per-	-	-	-	-	Thousand acres	-	-
		acres	cent							
Oak-hickory		1,859	47.2	470		269		511		481
White oak		369	9.4	186		99		43		18
Mixed hdwd.		539	13.7	170		61		196		96
Pin oak flats		275	7.0	104		65		63		24
Bottomland hdwd.		767	19.4	321		79		163		90
Scrub hdwd.		132	3.3	0		0		7		20
All types		3,941		1,251		573		983		729
Percent			100.0	31.8		14.5		24.9		18.5
										10.3

STATE OF ILLINOIS

Table 4.--Saw-timber volume by species and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling and sapling area	Poorly stocked area
	<u>Million</u> <u>bd. ft.</u>	<u>Percent</u>	<u>Million board feet</u>			
Redcedar and cypress	40	0.4	17	22	0	0
White oak	2,117	20.5	1,392	513	115	48
Post-oak group	870	8.4	569	151	101	38
Black oak	1,095	10.6	759	227	82	13
Northern red oak	1,036	10.0	830	133	57	0
Other red oaks	714	6.9	428	184	60	16
Hickory	700	6.8	442	157	76	17
Elm	1,020	9.9	678	155	100	52
Soft maple	633	6.1	515	65	20	28
Sugar maple	231	2.2	161	35	6	3
Sycamore	314	3.0	232	25	12	41
Ash	273	2.6	201	61	10	0
Basswood and yellow-poplar	181	1.7	149	14	12	6
Cottonwood	299	2.9	196	85	14	2
Black walnut	167	1.6	130	24	4	5
Other hardwoods	656	6.4	464	111	69	10
All species	10,346		7,163	1,962	738	279
Percent		100.0	69.2	19.0	7.1	2.7

STATE OF ILLINOIS

Table 5.--Saw-timber volume by species and tree-diameter class, 1948

Species	Total	12-14 inches	16-18 inches	20-22 inches	24-26 inches	28 inches and larger
- - - - - Million board feet - - - - -						
Redcedar and cypress	40	21 ^{1/}	5	0	6	8
White oak	2,117	776	741	325	155	120
Post-oak group	870	230	280	217	73	70
Black oak	1,095	244	357	248	160	86
Northern red oak	1,036	208	301	190	168	169
Other red oaks	714	209	174	114	162	55
Hickory	700	325	229	104	24	18
Elm	1,020	335	320	182	91	92
Soft maple	633	157	153	130	105	88
Sugar maple	231	80	83	48	4	16
Sycamore	314	33	62	65	82	72
Ash	273	109	89	47	28	0
Basswood and yellow-poplar	181	37	46	45	19	34
Cottonwood	299	64	71	70	42	52
Black walnut	167	91	47	15	6	8
Other hardwoods	656	203	149	121	135	48
All species	10,346	3,122	3,107	1,921	1,260	936
Percent	100.0	30.2	30.0	18.6	12.2	9.0

^{1/} Includes 5 million board feet in 10-inch trees.

STATE OF ILLINOIS

Table 6.--Hardwood saw-timber volume by species group and percentage distribution in log grades, 1948

Species group	:	Log grade		
	:	1	2	3
	:	Percent		
	Volume	Percent		
	Million	Percent		
	bd. ft.	Percent		
White oaks ^{1/}	2,987	6.9	18.3	74.8
Red oaks ^{2/}	2,845	13.1	18.9	68.0
Other hardwoods	4,474	12.1	16.8	71.1
All hardwoods	10,306	11.0	17.9	71.1

^{1/} Includes white oak and post-oak group.

^{2/} Includes black oak, northern red oak, and other red oaks.

STATE OF ILLINOIS

Table 7.--Total cubic volume of sound wood by species
and class of material, 1948

Species	Growing stock					
	Total	Total	Saw-timber	Pole-timber	Tops &	Cull
			trees	trees	limbs	trees
	- - - - - Million cubic feet - - - - -					
Redcedar and cypress	11.8	11.7	10.2 ^{1/}	1.5	--	0.1
White oak	694.6	427.2	320.2	107.0	223.5	43.9
Post-oak group	320.4	185.6	131.9	53.7	93.4	41.4
Black oak	366.2	218.8	165.9	52.9	117.7	29.7
Northern red oak	304.3	177.8	152.6	25.2	106.3	20.2
Other red oaks	279.0	168.5	109.1	59.4	77.8	32.7
Hickory	309.3	214.7	111.6	103.1	79.3	15.3
Elm	483.7	286.6	160.1	126.5	113.2	83.9
Soft maple	315.2	147.9	99.3	48.6	73.7	93.6
Sugar maple	88.5	55.4	35.5	19.9	25.0	8.1
Sycamore	97.2	56.0	47.4	8.6	33.7	7.5
Ash	154.0	96.7	43.1	53.6	30.6	26.7
Basswood and yellow-poplar	76.0	34.8	27.0	7.8	19.0	22.2
Cottonwood	86.0	50.0	45.7	4.3	32.0	4.0
Black walnut	93.4	60.1	26.6	33.5	18.8	14.5
Other hardwoods	364.1	202.4	101.2	101.2	72.3	89.4
Noncommercial species	14.6	--	--	--	--	14.6
All species	4,058.3	2,394.2	1,587.4	806.8	1,116.3	547.8
Percent	100.0	59.0	39.1	19.9	27.5	13.5

^{1/} Includes tops of redcedar and cypress trees.

STATE OF ILLINOIS

Table 8.--Cubic volume of growing stock by species
and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling: and sapling area	Poorly stocked area
	Million cu. ft.	Percent	- - - -	Million cubic feet	- - - -	- - - -
Redcedar and cypress	11.7	0.5	3.3	7.5	0.2	0.3
White oak	427.2	17.8	237.7	113.7	57.3	9.9
Post-oak group	185.6	7.8	97.4	39.0	40.1	2.6
Black oak	218.8	9.1	118.8	50.2	45.0	2.9
Northern red oak	177.8	7.4	128.2	28.5	17.5	3.6
Other red oaks	168.5	7.0	75.0	47.9	38.0	4.9
Hickory	214.7	9.0	104.3	54.4	48.6	4.0
Elm	286.6	12.0	147.0	50.7	69.7	10.0
Soft maple	147.9	6.2	108.7	22.3	11.5	.9
Sugar maple	55.4	2.3	32.8	10.6	7.7	3.9
Sycamore	56.0	2.3	36.8	5.3	6.8	.6
Ash	96.7	4.0	48.5	21.4	26.1	.7
Basswood and yellow-poplar	34.8	1.5	25.8	3.7	3.7	.6
Cottonwood	50.0	2.1	29.9	16.6	2.7	.4
Black walnut	60.1	2.5	29.8	8.3	16.2	1.0
Other hardwoods	202.4	8.5	92.5	34.0	63.8	8.3
All species	2,394.2		1,316.5	514.1	454.9	54.6
Percent		100.0	55.0	21.5	19.0	2.3

STATE OF ILLINOIS

Table 9.--Cubic volume of growing stock by stand-size class
and tree-diameter class, 1948

Stand-size class	:	:	:	:	:	:	:	:	:28 inches
	: Total	: 6-8	: 10	:12-14	:16-18	:20-22	:24-26	: and	
	:	:inches	:inches	:inches	:inches	:inches	:inches	:larger	
	-	-	-	-	-	-	-	-	-
	<u>Million cubic feet</u>								
Large saw-timber area	1,316.6	125.8	108.8	217.8	345.4	243.3	148.2	127.3	
Small saw-timber area	514.0	97.2	104.2	203.4	80.6	18.5	7.2	2.9	
Pole-timber area	454.9	224.4	113.7	66.3	28.4	10.7	10.5	.9	
Seedling and sapling area	54.6	15.8	6.6	13.6	11.2	4.8	1.3	1.3	
Poorly stocked area	54.1	7.1	4.7	7.8	10.9	6.0	16.5	1.1	
All classes	2,394.2	470.3	338.0	508.9	476.5	283.3	183.7	133.5	
Percent	100.0	19.6	14.1	21.3	19.9	11.8	7.7	5.6	

STATE OF ILLINOIS

Table 10.--Average volume per acre by stand-size class and region, 1948

BOARD-FOOT VOLUME				
Stand-size class	: All regions	: Southern	: Claypan	: Prairie
Large saw-timber area	5,726	5,883	5,548	5,784
Small saw-timber area	3,424	3,385	3,244	3,598
Pole-timber area	751	668	846	721
Seedling and sapling area	280	291	248	301
Poorly stocked area	689	281	775	746
All classes	2,625	2,613	2,585	2,668
CUBIC-FOOT VOLUME <u>1/</u>				
Large saw-timber area	1,052.4	1,129.5	1,051.1	1,004.4
Small saw-timber area	897.2	930.8	898.3	871.0
Pole-timber area	462.8	513.3	401.7	484.4
Seedling and sapling area	74.9	87.7	55.6	83.0
Poorly stocked area	133.6	78.9	135.8	146.1
All classes	607.5	662.3	592.3	584.5

1/ Growing stock only.

SOUTHERN REGION

SOUTHERN REGION

Table 1.--Forest and nonforest area by county, 1948

County	: Total : land <u>1/</u> : area	: : :	Forest area	: : :	Nonforest area
	<u>Thousand acres</u>		<u>Thousand acres</u> <u>Percent</u>		<u>Thousand acres</u> <u>Percent</u>
Alexander	143		67 47		76 53
Franklin	278		62 22		216 78
Gallatin	210		55 26		155 74
Hamilton	278		53 19		225 81
Hardin	117		45 38		72 62
Jackson	386		123 32		263 68
Johnson	221		79 36		142 64
Massac	157		44 28		113 72
Perry	283		61 22		222 78
Pope	244		100 41		144 59
Pulaski	131		34 26		97 74
Randolph	380		85 22		295 78
Saline	246		43 18		203 82
Union	265		101 38		164 62
White	321		30 9		291 91
Williamson	282		60 21		222 79
Total	3,942		1,042 26		2,900 74

1/ Source. Area of United States 1940, U. S. Bureau of the Census.

SOUTHERN REGION

Table 2.--Commercial forest area by ownership class, 1948

Ownership class	Commercial forest area ^{1/}	
	<u>Thousand acres</u>	<u>Percent</u>
Federal:		
National forest	147	14.3
Other	10	1.0
	<hr/>	
Total	157	15.3
State	4	.4
Private	869	84.3
	<hr/>	
All ownerships	1,030	100.0

^{1/} Does not include 12 thousand acres of forest land classified as noncommercial.

SOUTHERN REGION

Table 3.--Commercial forest area by forest type and stand-size class, 1948

Forest type	:	:	Large	:	Small	:	Seedling:	:
	:	:	saw-	:	saw-	:	Pole- and	:Poorly
	:	Total	timber	:	timber:	timber	sapling	:stocked
	:	:	area	:	area	:	area	: area
		<u>Thousand</u>	<u>Per-</u>	- - - - -	<u>Thousand</u>	<u>acres</u>	- - - - -	
		<u>acres</u>	<u>cent</u>					
Oak-hickory		537	52.1	103	101	182	126	25
White oak		49	4.8	17	9	11	5	7
Mixed hdwd.		85	8.3	54	6	15	10	0
Pin oak flats		129	12.5	40	28	30	24	7
Bottomland hdwd.		214	20.8	101	25	48	33	7
Scrub hdwd.		16	1.5	0	0	0	5	11
All types		1,030		315	169	286	203	57
Percent			100.0	30.6	16.4	27.8	19.7	5.5

SOUTHERN REGION

Table 4.--Saw-timber volume by species and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling and sapling area	Poorly stocked area
	<u>Million</u> <u>bd. ft.</u>	<u>Percent</u>	<u>Million board feet</u>			
Redcedar and cypress	21	.8	17	3	0	0
White oak	410	15.2	263	97	40	5
Post-oak group	177	6.6	105	43	22	2
Black oak	293	10.9	180	76	24	2
Northern red oak	236	8.8	193	36	6	0
Other red oaks	366	13.6	251	77	14	4
Hickory	252	9.4	157	66	26	1/
Elm	140	5.2	90	22	22	1/
Soft maple	111	4.1	87	16	3	0
Sugar maple	33	1.2	26	4	3	0
Sycamore	105	3.9	77	12	12	0
Ash	63	2.3	52	6	4	0
Basswood and yellow-poplar ^{2/}	71	2.6	65	1	5	0
Cottonwood	120	4.5	67	53	0	0
Black walnut	22	.8	16	5	1	0
Other hardwoods	271	10.1	207	55	9	0
All species	2,691		1,853	572	191	16
Percent		100.0	68.9	21.2	7.1	.6

1/ Less than .5 million board feet.

2/ Chiefly yellow-poplar.

SOUTHERN REGION

Table 5.--Saw-timber volume by species and tree-diameter class 1948

Species	: Total	: 12-14	: 16-18	: 20-22	: 24-26	: 28 inches and larger
	: inches	: inches	: inches	: inches	: inches	: larger
	- - - - - Million board feet - - - - -					
Redcedar and cypress	21	7 ^{1/}	0	0	6	8
White oak	410	163	144	66	17	20
Post-oak group	177	63	54	33	19	8
Black oak	293	81	88	53	23	48
Northern red oak	236	55	65	36	43	37
Other red oaks	366	104	91	64	61	46
Hickory	252	123	60	51	11	7
Elm	140	52	44	19	14	11
Soft maple	111	31	29	29	22	0
Sugar maple	33	15	9	5	4	0
Sycamore	105	17	25	13	25	25
Ash	63	29	24	10	0	0
Basswood and yellow-poplar	71	11	21	18	11	10
Cottonwood	120	36	33	11	6	34
Black walnut	22	17	4	1	0	0
Other hardwoods	271	102	83	59	20	7
All species	2,691	906	774	468	282	261
Percent	100.0	33.7	28.7	17.4	10.5	9.7

^{1/} Includes 1 million board feet in 10-inch trees.

SOUTHERN REGION

Table 6.--Hardwood saw-timber volume by species group and percentage distribution in log grades, 1948

Species group	: Volume Million bd. ft.	: Log grade 1	: Log grade 2	: Log grade 3
		- - - - -	Percent - - - - -	
White oaks ^{1/}	587	8.4	15.2	76.4
Red oaks ^{2/}	895	18.0	16.4	65.6
Other hardwoods	1,188	12.9	11.6	75.5
All hardwoods	2,670	13.5	13.9	72.6

^{1/} Includes white oak and post-oak group.

^{2/} Includes black oak, northern red oak, and other red oaks.

SOUTHERN REGION

Table 7.--Total cubic volume of sound wood by species
and class of material, 1948

Species	:	:	Growing stock				:	:
	Total	:	Saw-timber	Pole-timber	Tops &	Cull		
	:	Total	trees	trees	limbs	trees		
<hr/>								
	-	-	-	-	-	-	-	-
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							
	<hr/>							

1/ Includes tops of redcedar and cypress trees.

SOUTHERN REGION

Table 8.--Cubic volume of growing stock by species
and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling: and sapling area	Poorly stocked area
	<u>Million</u> <u>cu. ft.</u>	<u>Percent</u>	- - - -	<u>Million cubic feet</u>	- - - -	- - - -
Redcedar and cypress	4.5	.7	3.3	.8	.1	.3
White oak	93.3	13.7	46.2	24.3	19.8	1.4
Post-oak group	46.7	6.8	20.3	10.3	14.3	1.4
Black oak	60.2	8.8	29.5	15.5	13.0	1.9
Northern red oak	41.2	6.0	29.4	7.9	3.5	.4
Other red oaks	91.3	13.4	45.2	20.6	20.9	4.0
Hickory	87.2	12.8	37.5	24.6	23.3	.7
Elm	50.5	7.4	24.6	10.6	12.4	2.8
Soft maple	25.6	3.8	17.0	5.5	2.0	.9
Sugar maple	10.9	1.6	7.6	1.8	1.4	.1
Sycamore	18.3	2.7	12.3	3.0	2.4	.6
Ash	25.7	3.8	14.0	4.4	7.0	.3
Basswood and yellow-poplar	12.8	1.9	10.8	.2	1.8	0
Cottonwood	21.9	3.2	10.1	11.3	.3	.1
Black walnut	9.2	1.3	4.3	1.9	3.0	0
Other hardwoods	82.9	12.1	43.7	14.6	21.6	2.9
All species	682.2		355.8	157.3	146.8	17.8
Percent		100.0	52.1	23.1	21.5	2.6

1/ Less than .05 million cubic feet.

SOUTHERN REGION

Table 9.--Cubic volume of growing stock by stand-size class
and tree-diameter class, 1948

Stand-size class	: Total	: 6-8	: 10	: 12-14	: 16-18	: 20-22	: 24-26	: 28 inches and larger
	: inches	: inches	: inches	: inches	: inches	: inches	: inches	: inches
	- - - - - Million cubic feet - - - - -							
Large saw-timber area	355.9	39.5	33.9	61.5	88.0	60.0	36.6	36.4
Small saw-timber area	157.2	33.1	32.4	63.2	21.2	6.2	1.1	0
Pole-timber area	146.8	82.4	34.0	19.6	6.0	1.5	2.4	.9
Seedling and sapling area	17.8	5.4	3.1	3.2	3.5	1.3	1.3	0
Poorly stocked area	4.5	1.5	.5	1.0	.6	.6	.3	0
All classes	682.2	161.9	103.9	148.5	119.3	69.6	41.7	37.3
Percent	100.0	23.7	15.2	21.8	17.5	10.2	6.1	5.5

SOUTHERN REGION

Table 10.--Average volume per acre by stand-size class, 1948

Stand-size class	Average volume per acre	
	<u>Board feet</u>	<u>Cubic feet</u> ^{1/}
Large saw-timber area	5,883	1,129.5
Small saw-timber area	3,385	930.8
Pole-timber area	668	513.3
Seedling and sapling area	291	87.7
Poorly stocked area	281	78.9
All classes	2,613	662.3

^{1/} Growing stock only.

CLAYPAN REGION

CLAYPAN REGION

Table 1.--Forest and nonforest area by county, 1948

County	: Total : land : area $\frac{1}{2}$	:	Forest area	:	Nonforest area
	<u>Thousand acres</u>		<u>Thousand acres</u> <u>Percent</u>		<u>Thousand acres</u> <u>Percent</u>
Bond	245		37 15		208 85
Calhoun	166		72 43		94 57
Clark	323		58 18		265 82
Clay	297		46 15		251 85
Clinton	319		62 19		257 81
Crawford	283		40 14		243 86
Cumberland	222		30 14		192 86
Edwards	144		20 14		124 86
Effingham	309		47 15		262 85
Fayette	460		84 18		376 82
Greene	348		54 16		294 84
Jasper	317		36 11		281 89
Jefferson	367		61 17		306 83
Jersey	239		66 28		173 72
Lawrence	239		34 14		205 86
Macoupin	558		95 17		463 83
Madison	468		57 12		411 88
Marion	371		62 17		309 83
Monroe	243		58 24		185 76
Montgomery	452		46 10		406 90
Richland	233		30 13		203 87
St. Clair	429		55 13		374 87
Shelby	494		55 11		439 89
Wabash	141		14 10		127 90
Washington	362		63 17		299 83
Wayne	458		74 16		384 84
Total	8,487		1,356 16		7,131 84

^{1/} Source: Area of United States 1940, U. S. Bureau of the Census.

CLAYPAN REGION

Table 2.--Commercial forest area by ownership class, 1948

Ownership class	Commercial forest area <u>1/</u>	
	<u>Thousand</u> <u>acres</u>	<u>Percent</u>
Federal:		
National forest	0	0
Other	6	.4
	<hr/>	
Total	6	.4
State	0	0
Private	1,341	99.6
	<hr/>	
All ownerships	1,347	100.0

1/ Does not include 9 thousand acres of forest land classified as noncommercial.

CLAYPAN REGION

Table 3.--Commercial forest area by forest type and stand-size class, 1948

Forest type	Total	: Large : saw- : timber : area	: Small : saw- : timber : area	: Pole- : timber : area	: Seedling: : and : sapling : area	: Poorly : stocked : area
	<u>Thousand</u> <u>acres</u>	<u>Per-</u> <u>cent</u>	- - - - -	<u>Thousand</u> <u>acres</u>	- - - - -	- - - - -
Oak-hickory	655	48.6	168	69	199	171 48
White oak	125	9.3	72	23	17	13 0
Mixed hdwd.	131	9.7	14	19	58	40 0
Pin oak flats	141	10.5	59	37	33	0 12
Bottomland hdwd.	271	20.1	127	32	50	26 36
Scrub hdwd.	24	1.8	0	0	0	0 24
All types	1,347		440	180	357	250 120
Percent		100.0	32.6	13.4	26.5	18.6 8.9

CLAYPAN REGION

Table 4.--Saw-timber volume by species and stand-size class, 1948

Species			: Large	: Small	:	: Seedling	:
			: saw-	: saw-	: Pole-	: and	: Poorly
			: timber	: timber	: timber	: sapling	: stocked
			: area	: area	: area	: area	: area
	<u>Million</u>	<u>Percent</u>	<u>Million board feet</u>				
	<u>bd. ft.</u>		-	-	-	-	-
White oak	690	19.8	485	129	40	30	6
Post-oak group	323	9.3	169	67	77	6	4
Black oak	418	12.0	329	59	30	0	0
Northern red oak	151	4.3	143	8	0	0	0
Other red oaks	334	9.6	169	101	46	6	12
Hickory	305	8.8	217	59	19	0	10
Elm	332	9.5	226	37	22	16	31
Soft maple	303	8.7	243	24	8	0	28
Sugar maple	31	.9	17	12	2	0	0
Sycamore	108	3.1	103	5	0	0	0
Ash	125	3.6	82	41	2	0	0
Basswood and yellow-poplar	5	.2	2	3	0	0	0
Cottonwood	46	1.3	30	0	12	2	2
Black walnut	78	2.2	64	9	3	2	0
Other hardwoods	233	6.7	162	30	41	0	0
All species	3,482		2,441	584	302	62	93
Percent		100.0	70.1	16.8	8.7	1.8	2.6

CLAYPAN REGION

Table 5.--Saw-timber volume by species and tree-diameter class, 1948

Species	: Total	: 12-14	: 16-18	: 20-22	: 24-26	: 28 inches
	: inches	: inches	: inches	: inches	: inches	: and
	:	:	:	:	:	: larger
	- - - - - Million board feet - - - - -					
White oak	690	251	270	95	44	30
Post-oak group	323	102	125	72	13	11
Black oak	418	78	145	96	65	34
Northern red oak	151	15	43	36	23	34
Other red oaks	334	100	79	45	101	9
Hickory	305	107	127	47	13	11
Elm	332	122	104	68	0	38
Soft maple	303	55	64	58	48	78
Sugar maple	31	8	23	0	0	0
Sycamore	108	13	28	34	12	21
Ash	125	53	34	14	24	0
Basswood and yellow-poplar	5	5	0	0	0	0
Cottonwood	46	9	0	9	10	18
Black walnut	78	51	16	5	6	0
Other hardwoods	233	38	49	37	75	34
All species	3,482	1,007	1,107	616	434	318
Percent	100.0	28.9	31.8	17.7	12.5	9.1

CLAYPAN REGION

Table 6.--Hardwood saw-timber volume by species group and percentage distribution in log grades, 1948

Species group	:	:	Log grade:	Log grade :	Log grade
	: Volume	:	1	:	2
	Million	- - - - -	Percent	- - - - -	3
	bd. ft.				
White oaks ^{1/}	1,013		3.5	28.0	68.5
Red oaks ^{2/}	903		5.8	21.4	72.8
Other hardwoods	1,566		13.4	27.2	59.4
<hr/>					
All hardwoods	3,482		8.0	25.8	66.2

^{1/} Includes white oak and post-oak group.

^{2/} Includes black oak, northern red oak, and other red oaks.

CLAYPAN REGION

Table 7.--Total cubic volume of sound wood by species
and class of material, 1948

Species	Growing stock					
	Total	Sav-timber:Pole-timber:Tops &				Cull
	Total:	trees	trees	limbs	trees	
<hr/>						
	<hr/>					
	<hr/>					
Redcedar	.1	.1	0	.1	0	0
White oak	225.9	140.7	104.9	35.8	73.6	11.6
Post-oak group	129.3	75.8	49.9	25.9	35.6	17.9
Black oak	129.7	81.3	63.4	17.9	45.3	3.1
Northern red oak	43.4	23.8	22.1	1.7	15.5	4.1
Other red oaks	126.3	72.6	51.2	21.4	36.5	17.2
Hickory	116.5	78.0	48.3	29.7	34.5	4.0
Elm	142.2	88.7	52.0	36.7	37.1	16.4
Soft maple	141.6	66.0	47.6	18.4	35.8	39.8
Sugar maple	12.2	8.2	5.0	3.2	3.5	.5
Sycamore	32.3	20.5	16.5	4.0	11.8	0
Ash	69.2	44.6	19.7	24.9	14.1	10.5
Basswood and yellow-poplar	5.9	2.8	.7	2.1	.5	2.6
Cottonwood	13.2	7.7	6.9	.8	4.8	.7
Black walnut	39.6	25.1	12.5	12.6	8.9	5.6
Other hardwoods	109.9	61.9	35.1	26.8	25.2	22.8
Noncommercial species	2.3	--	--	--	--	2.3
<hr/>						
All species	1,339.6	797.8	535.8	262.0	382.7	159.1
<hr/>						
Percent	100.0	59.5	40.0	19.5	28.6	11.9

CLAYPAN REGION

Table 8.--Cubic volume of growing stock by species
and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Seedling: Pole- timber area	and sapling area	Poorly stocked area
	<u>Million</u> <u>cu. ft.</u>	<u>Percent</u>	<u>Million cubic feet</u>			
Redcedar and cypress	.1	0	0	0	.1	0
White oak	140.7	17.6	86.9	30.6	17.4	4.9
Post-oak group	75.8	9.5	32.1	19.4	22.2	1.2
Black oak	81.3	10.2	51.1	15.5	14.6	.1
Northern red oak	23.8	3.0	21.2	1.9	.7	0
Other red oaks	72.6	9.1	27.1	26.0	16.5	.9
Hickory	78.0	9.8	49.0	16.0	11.3	0
Elm	88.7	11.1	52.6	14.6	13.5	3.0
Soft maple	66.0	8.3	52.7	5.8	3.2	0
Sugar maple	8.2	1.0	2.8	3.5	1.9	0
Sycamore	20.5	2.6	16.8	.9	2.8	0
Ash	44.6	5.6	17.8	14.1	12.5	.2
Basswood and yellow-poplar	2.8	.4	1.8	.7	0	.3
Cottonwood	7.7	1.0	5.0	0	2.1	.3
Black walnut	25.1	3.1	14.5	3.4	6.0	.7
Other hardwoods	61.9	7.7	31.1	9.3	18.6	2.3
All species	797.8		462.5	161.7	143.4	13.9
Percent		100.0	58.0	20.3	18.0	1.7

CLAYPAN REGION

Table 9.--Cubic volume of growing stock by stand-size class
and tree-diameter class, 1948

Stand-size class	:	:	:	:	:	:	:	:28 inches
	: Total	: 6-8	: 10	:12-14	: 16-18	:20-22	:24-26	: and
	:	:inches	:inches	:inches	:inches	:inches	:inches	: larger
	-	-	-	-	-	-	-	-
	<u>Million cubic feet</u>							
Large saw-timber area	462.5	51.7	39.2	75.9	118.4	85.3	47.3	44.7
Small saw-timber area	161.7	31.0	37.8	56.7	28.7	2.4	3.7	1.4
Pole-timber area	143.4	67.4	28.5	24.2	13.0	3.8	6.5	0
Seedling and sapling area	13.9	2.5	2.0	3.3	5.7	.4	0	0
Poorly stocked area	16.3	1.6	.3	3.8	4.6	0	6.0	0
All classes	797.8	154.2	107.8	163.9	170.4	91.9	63.5	46.1
Percent	100.0	19.3	13.5	20.5	21.4	11.5	8.0	5.8

CLAYPAN REGION

Table 10.--Average volume per acre by stand-size class, 1948

Stand-size class	Average volume per acre	
	Board feet	Cubic feet ^{1/}
Large saw-timber area	5,548	1,051.1
Small saw-timber area	3,244	898.3
Pole-timber area	846	401.7
Seedling and sapling area	248	55.6
Poorly stocked area	775	135.8
All classes	2,585	592.3

^{1/} Growing stock only.

PRAIRIE REGION

PRAIRIE REGION

Table 1.--Forest and nonforest area by county, 1948

County	: : Total : land : area $\frac{1}{2}$:	:	: : Forest area :	:	: : Nonforest area :
	<u>Thousand</u> <u>acres</u>	<u>Thousand</u> <u>acres</u>	<u>Percent</u>	<u>Thousand</u> <u>acres</u>	<u>Percent</u>
Adams	554	93	17	461	83
Boone	181	6	3	175	97
Brown	196	42	21	154	79
Bureau	555	35	6	520	94
Carroll	300	22	7	278	93
Cass	237	46	19	191	81
Champaign	640	7	1	633	99
Christian	454	13	3	441	97
Coles	324	23	7	301	93
Cook	611	30	5	581	95
DeKalb	407	5	1	402	99
DeWitt	255	11	4	244	96
Douglas	269	5	2	264	98
DuPage	212	11	5	201	95
Edgar	402	23	6	379	94
Ford	312	1	0	311	100
Fulton	559	96	17	463	83
Grundy	276	10	4	266	96
Hancock	510	46	9	464	91
Henderson	244	35	14	209	86
Henry	529	17	3	512	97
Iroquois	718	13	2	705	98
Jo Daviess	393	60	15	333	85
Kane	330	10	3	320	97
Kankakee	435	16	4	419	96
Kendall	205	8	4	197	96
Knox	466	44	9	422	91
Lake	292	20	7	272	93
LaSalle	738	26	4	712	96
Lee	467	9	2	458	98

PRAIRIE REGION

Table 1.--Forest and nonforest area by county, 1948, continued

County	: Total : land : area <u>1/</u>	:	Forest area	:	Nonforest area	
	<u>Thousand</u> <u>acres</u>		<u>Thousand</u> <u>acres</u>	<u>Percent</u>	<u>Thousand</u> <u>acres</u>	<u>Percent</u>
Livingston	668		9	1	659	99
Logan	398		11	3	387	97
McDonough	372		40	11	332	89
McHenry	391		16	4	375	96
McLean	751		18	2	733	98
Macon	369		12	3	357	97
Marshall	253		28	11	225	89
Mason	346		44	13	302	87
Menard	200		15	8	185	92
Mercer	356		28	8	328	92
Morgan	362		30	8	332	92
Moultrie	221		8	4	213	96
Ogle	484		29	6	455	94
Peoria	399		52	13	347	87
Piatt	280		7	2	273	98
Pike	530		97	18	433	82
Putnam	106		18	17	88	83
Rock Island	269		34	13	235	87
Sangamon	563		24	4	539	96
Schuyler	278		69	25	209	75
Scott	161		21	13	140	87
Stark	186		6	3	180	97
Stephenson	363		16	4	347	96
Tazewell	418		38	9	380	91
Vermilion	575		29	5	546	95
Warren	347		25	7	322	93
Whiteside	442		16	4	426	96
Will	541		23	4	518	96
Winnebago	333		22	7	311	93
Woodford	344		30	9	314	91
Total	23,377		1,598	7	21,779	93

1/ Source: Area of United States 1940, U. S. Bureau of the Census

PRAIRIE REGION

Table 2.--Commercial forest area by ownership class, 1948

Ownership class	Commercial forest area <u>1/</u>	
	<u>Thousand acres</u>	<u>Percent</u>
Federal:		
National forest	0	0
Other	16	1.0
<hr/>		
Total	16	1.0
State	6	.4
Private	1,542	98.6
<hr/>		
All ownerships	1,564	100.0

1/ Does not include 34 thousand acres of forest land classified as noncommercial.

PRAIRIE REGION

Table 3.--Commercial forest area by forest type and stand-size class, 1948

Forest type	:	:	:Large	: Small	:	:Seedling:	:
	:	:	: saw-	: saw-	: Pole-	: and	:Poorly
	:	Total	:timber:	timber	: timber	:sapling	:stocked
	:	:	: area	: area	: area	: area	: area
	<u>Thousand</u>	<u>Per-</u>	- - - -	- - -	<u>Thousand</u>	<u>acres</u>	- - - - -
	<u>acres</u>	<u>cent</u>					
Oak-hickory	667	42.6	199	99	130	184	55
White oak	195	12.5	97	67	15	0	16
Mixed hwd.	323	20.7	102	36	123	46	16
Pin oak flats	5	.3	5	0	0	0	0
Bottomland hwd.	282	18.0	93	22	65	31	71
Scrub hwd.	92	5.9	0	0	7	15	70
All types	1,564		496	224	340	276	228
Percent	100.0	31.7	14.3	21.8	17.6	14.6	

PRAIRIE REGION

Table 4.--Saw-timber volume by species and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling and sapling area	Poorly stocked area
	Million bd. ft.	Percent	- - - - - Million board feet - - - - -			
Redcedar	19	.5	0	19	0	0
White oak	1,017	24.4	644	287	35	14
Post-oak group <u>1/</u>	370	8.9	295	41	2	0
Black oak	384	9.2	250	92	28	3
Northern red oak	649	15.6	494	89	51	15
Other red oaks	14	.3	8	6	0	0
Hickory	143	3.4	68	32	31	8
Elm	548	13.1	362	96	56	13
Soft maple	219	5.3	185	25	9	0
Sugar maple	167	4.0	118	19	1	26
Sycamore	101	2.4	52	8	0	0
Ash	85	2.0	67	14	4	0
Basswood and yellow-poplar <u>2/</u>	105	2.5	82	10	7	0
Cottonwood	133	3.2	99	32	2	0
Black walnut	67	1.6	50	10	0	2
Other hardwoods	152	3.6	95	26	19	2
All species	4,173		2,869	806	245	83
Percent		100.0	68.7	19.3	5.9	2.0
						4.1

1/ Chiefly bur oak.
2/ Chiefly basswood.

PRAIRIE REGION

Table 5.--Saw-timber volume by species and tree-diameter class, 1948

Species	: :Total :	: : 12-14 : inches	: : 16-18 : inches	: : 20-22 : inches	: : 24-26 : inches	: : 28 inches : and : larger
	- - - - - Million board feet - - - - -					
Redcedar	19	14 ^{1/}	5	0	0	0
White oak	1,017	362	327	164	94	70
Post-oak group	370	65	101	112	41	51
Black oak	384	85	124	99	72	4
Northern red oak	649	138	193	118	102	98
Other red oaks	14	5	4	5	0	0
Hickory	143	95	42	6	0	0
Elm	548	161	172	95	77	43
Soft maple	219	71	60	43	35	10
Sugar maple	167	57	51	43	0	16
Sycamore	101	3	9	18	45	26
Ash	85	27	31	23	4	0
Basswood and yellow-poplar	105	21	25	27	8	24
Cottonwood	133	19	38	50	26	0
Black walnut	67	23	27	9	0	8
Other hardwoods	152	63	17	25	40	7
All species	4,173	1,209	1,226	837	544	357
Percent	100.0	29.0	29.4	20.1	13.0	8.5

^{1/} Includes 4 million board feet in 10-inch trees.

PRAIRIE REGION

Table 6.--Hardwood saw-timber volume by species group and percentage
distribution in log grades, 1948

Species group	: : Volume	:Log grade : 1	: Log grade : 2	: Log grade : 3
	<u>Million</u> <u>bd. ft.</u>	- - - - -	<u>Percent</u> - - - - -	- - - - -
White oaks ^{1/}	1,387	7.7	14.4	77.9
Red oaks ^{2/}	1,047	9.6	21.7	68.7
Other hardwoods	1,720	9.3	20.3	70.4
All hardwoods	4,154	8.9	18.8	72.3

^{1/} Includes white oak and post-oak group.

^{2/} Includes black oak, northern red oak, and other red oaks.

PRAIRIE REGION

Table 7.--Total cubic volume of sound wood by species
and class of material, 1948

Species	Growing stock					
	: Total :	: Saw-timber: Pole-timber: Tops & :			: Cull	
		: Total :	trees	trees	limbs	trees
	- - - - - Million cubic feet - - - - -					
Redcedar	7.2	7.1	5.8 ^{1/}	1.3	--	.1
White oak	320.8	193.2	152.6	40.6	105.9	21.7
Post-oak group	113.2	63.1	54.8	8.3	38.3	11.8
Black oak	138.1	77.3	57.8	19.5	40.5	20.3
Northern red oak	190.6	112.8	95.6	17.2	66.3	11.5
Other red oaks	7.1	4.6	2.1	2.5	1.5	1.0
Hickory	71.0	49.5	22.9	26.6	16.0	5.5
Elm	261.1	147.4	86.0	61.4	60.3	53.4
Soft maple	116.7	56.3	34.0	22.3	24.6	35.8
Sugar maple	58.5	36.3	25.5	10.8	17.9	4.3
Sycamore	30.5	17.2	15.0	2.2	10.5	2.8
Ash	43.7	26.4	13.2	13.2	9.3	8.0
Basswood and yellow-poplar	48.6	19.2	15.6	3.6	10.9	18.5
Cottonwood	35.8	20.4	20.2	.2	14.1	1.3
Black walnut	40.9	25.8	10.4	15.4	7.3	7.8
Other hardwoods	101.3	57.6	23.5	34.1	16.5	27.2
Noncommercial species	10.9	--	--	--	--	10.9
All species	1,596.0	914.2	635.0	279.2	439.9	241.9
Percent	100.0	57.3	39.8	17.5	27.6	15.1

^{1/} Includes tops of redcedar trees.

PRAIRIE REGION

Table 8.--Cubic volume of growing stock by species
and stand-size class, 1948

Species	Total	Large saw- timber area	Small saw- timber area	Pole- timber area	Seedling: and sapling area	Poorly stocked area
	Million cu. ft.	Percent	- - - - -	Million cubic feet	- - - - -	- - - - -
Redcedar	7.1	.8	0	6.7	0	.4
White oak	193.2	21.1	104.6	58.8	20.1	3.6
Post-oak group	63.1	6.9	45.0	9.3	3.6	0
Black oak	77.3	8.5	38.2	19.2	17.4	.9
Northern red oak	112.8	12.3	77.6	18.7	13.3	3.2
Other red oaks	4.6	.5	2.7	1.3	.6	0
Hickory	49.5	5.4	17.8	13.8	14.0	3.3
Elm	147.4	16.1	69.8	25.5	43.8	4.2
Soft maple	56.3	6.2	39.0	11.0	6.3	0
Sugar maple	36.3	4.0	22.4	5.3	4.4	3.8
Sycamore	17.2	1.9	7.7	1.4	1.6	0
Ash	26.4	2.9	16.7	2.9	6.6	.2
Basswood and yellow-poplar	19.2	2.1	13.2	2.8	1.9	.3
Cottonwood	20.4	2.2	14.8	5.3	.3	0
Black walnut	25.8	2.8	11.0	3.0	7.2	.3
Other hardwoods	57.6	6.3	17.7	10.1	23.6	3.1
All species	914.2		498.2	195.1	164.7	22.9
Percent		100.0	54.5	21.3	18.0	2.5

PRAIRIE REGION

Table 9.--Cubic volume of growing stock by stand-size class
and tree-diameter class, 1948

	:	:	:	:	:	:	:	:28 inches
Stand-size class:	Total	6-8	10	12-14	16-18	20-22	24-26	and
	:	inches	inches	inches	inches	inches	inches	larger
	-	-	-	-	-	-	-	-
	<u>Million cubic feet</u>							
Large saw-timber area	498.2	34.6	35.7	80.4	139.0	98.0	64.3	46.2
Small saw-timber area	195.1	33.1	34.0	83.5	30.7	9.9	2.4	1.5
Pole-timber area	164.7	74.6	51.2	22.5	9.4	5.4	1.6	0
Seedling and sapling area	22.9	7.9	1.5	7.1	2.0	3.1	0	1.3
Poorly stocked area	33.3	4.0	3.9	3.0	5.7	5.4	10.2	1.1
All classes	914.2	154.2	126.3	196.5	186.8	121.8	78.5	50.1
Percent	100.0	16.9	13.8	21.5	20.4	13.3	8.6	5.5

PRAIRIE REGION

Table 10.--Average volume per acre by stand-size class, 1948

Stand-size class	Average volume per acre	
	<u>Board feet</u>	<u>Cubic feet</u> <u>1/</u>
Large saw-timber area	5,784	1,004.4
Small saw-timber area	3,598	871.0
Pole-timber area	721	484.4
Seedling and sapling area	301	83.0
Poorly stocked area	746	146.1
<hr/>		
All classes	2,668	584.5

1/ Growing stock only.

FOREST SURVEY PROCEDURE

The inventory of the forest resources of Illinois was made during the period October 1947 to May 1948. A sampling procedure was used involving an office study of aerial photographs and a field examination of randomly selected forest and nonforest plots.

The proportion of forest land in each county was obtained by placing a transparent template marked with uniformly spaced dots over aerial photographs and counting the number of dots falling on forest and nonforest areas. The percentage of the forest dots in a county applied to the total land area gave a preliminary estimate of the forest area. This was later adjusted after field examination.

The location of a selected number of dots falling on forest land was marked on the photographs. The acre surrounding each marked dot was examined under stereoscope and was classified by stand-size class on the basis of the height, crown width, and density of trees on the plot. Plots to be examined in the field were randomly drawn from these classified under stereoscope. In making this selection the greatest weight was given to the stand-size classes containing the heaviest timber volume. In addition, a number of nonforest plots were selected for field examination to measure the movement of open land to forest since the date of photography.

Through the cooperation of the Illinois Agricultural Experiment Station and the Illinois Division of Forestry, the sampling intensity of the Southern region was increased to provide more accurate forest area and timber volume data for this important area.

The locations of the selected field plots were marked on the photographs, which were then sent to the field. Crews of two men each located these points on the ground. On forest land a 1/5-acre plot was established on which forest conditions were described and the species, size, quality, and growth of trees were recorded.

The following tabulation gives the number of dots and plots examined for each of the three regions and for the state.

	<u>Total</u>	<u>Southern</u> <u>region</u>	<u>Claypan</u> <u>region</u>	<u>Prairie</u> <u>region</u>
Number of dots counted to determine forest and nonforest area	191,350	35,702	41,006	114,642
Number of forest plots stereoscopically examined on photos	5,933	2,999	1,232	1,702
Number of forest plots field examined	1,033	542	212	279
Number of nonforest plots field examined	266	109	70	87

ACCURACY OF DATA

Statistical analysis of forest area and timber volume data shows the following sampling errors 1/ for each of three regions and for the state:

<u>Region</u>	<u>Total forest area</u>		<u>Total saw-timber volume</u>	
	<u>M acres</u>	<u>Percent</u>	<u>Million</u> <u>bd. ft.</u>	<u>Percent</u>
Southern	+ 24	+ 2.35	+ 129	+ 4.80
Claypan	+ 36	+ 2.66	+ 226	+ 6.50
Prairie	+ 45	+ 2.81	+ 277	+ 6.64
State	+ 63	+ 1.57	+ 380	+ 3.68

These estimates of sampling error do not include errors resulting from the development and application of volume tables and cull factors, or from mistakes in measurement or judgment. All phases of field and office work were closely supervised to keep these errors to a minimum. Since the percentage error increases with each subdivision of the total, small acreages or volumes may have large errors and may therefore indicate only relative magnitudes.

1/ At one standard deviation; i.e., the chances are two out of three that, if the survey were repeated, the total forest area or volume figures would not differ more than the errors shown in this table.

EXPLANATION OF TERMS USED

Forest land - Land bearing forest growth or land from which the forest has been removed but which shows evidence of past forest occupancy and which is not now in other use. To qualify as forest, an area must: (1) be at least 100 feet wide; (2) be at least one acre in area; (3) have a sufficient number of trees to provide 10 percent crown coverage; or (4) lacking 10 percent crown coverage, be likely to remain in forest use.

Commercial forest land - Forest land bearing or capable of bearing timber of commercial character and economically available now or prospectively for commercial use and not withdrawn from such use.

Noncommercial forest land - Forest land not qualifying as commercial forest land. Two classes of forest area are included: (1) commercially valuable forest land withdrawn from timber use for such purposes as parks, game refuges, or reservoir protection; and (2) forest land which because of poor growing conditions will not produce trees of commercial quality.

Forest types

Oak-hickory - Stands of hardwoods in which oaks and hickories comprise at least 60 percent of the dominant and codominant trees.

White oak - Stands in which white oak (*Quercus alba*) comprises at least 60 percent of the dominant and codominant trees.

Mixed hardwoods - Stands of mixed hardwood species not qualifying for other hardwood types. Principal species include elm, maple, basswood, and black walnut in mixture with oaks and hickories.

Pin oak flats - Stands of pin oak and other hardwoods occurring on poorly drained flats. Associate species include soft maple, elm, hickory, and sweetgum.

Bottomland hardwoods - Stands on the alluvial bottoms of rivers and streams. The principal species include sycamore, willow, elm, blackgum, sweetgum, soft maple, oaks, hickory, cottonwood, and cypress.

Scrub hardwoods - Stands in which scrub oak or other noncommercial tree species comprise at least 60 percent of the dominant and codominant trees.

Tree classes

Sound saw-timber tree - A coniferous tree at least 9.0 inches d.b.h. (diameter outside bark at 4.5 feet above ground), or a hardwood tree at least 11.0 inches d.b.h., with a sound butt log at least 8 feet long, or with at least half of the gross volume of the tree in sound material.

Sound pole-timber tree - A tree at least 5.0 inches d.b.h. but less than saw-timber size, which now is or gives promise of becoming a sound merchantable tree.

Cull tree - A tree that does not qualify as a sound pole-timber or saw-timber tree because of poor form, limbiness, rot, or other defect.

Volume estimates

Board-foot volume includes the volume of that portion of saw-timber trees merchantable for sawlogs. Volume deductions have been made for rot, crook, and other defects. Board-foot volumes are shown in the International 1/4-inch log rule, which approximates green lumber tally.

Cubic-foot volume

Total volume includes the sound wood inside bark in both sound and cull living trees 5.0 inches d.b.h. and larger from the stump to a minimum top diameter of 4.0 inches inside bark. It therefore includes the upper stems of coniferous trees and the upper stems and limbs of hardwoods.

Growing stock includes the volume of sound wood inside bark in: (1) the saw-timber portion of sound trees, (2) the upper stem of sound saw-timber-sized conifers to a minimum top diameter of 4.0 inches inside bark, and (3) the sound pole-timber trees to the same top diameter.

Stand-size class

Large saw timber - stands having a minimum net volume of 1500 board feet per acre in living merchantable trees, and having more than half of this volume in trees 15.0 inches d.b.h. and larger.

Small saw timber - stands having a net volume of 1500 board feet per acre in living merchantable trees, and having at least half of this volume in trees smaller than 15.0 inches in d.b.h.

Pole timber - stands having a net volume of less than 1500 board feet per acre in living merchantable trees, but which are at least 10 percent stocked with pole-sized and larger trees. At least half the stocking must be in pole-sized trees.

Seedlings and saplings - stands not qualifying either for saw timber or pole timber but having at least 300 seedlings and saplings of commercial species per acre.

Poorly stocked - commercial forest land not qualifying for any other class. Denuded areas that are not likely to be put to a nonforest use are classed as poorly stocked.

Hardwood log grades

Grade 1 - Logs at least 14.0 inches in diameter inside bark with five-sixths of the surface on the three best faces clear of defect in not more than two cuttings. Lumber from such logs will normally grade at least 60 percent No. 1 common and better.

Grade 2 - Logs at least 12 inches in diameter inside bark with two-thirds of the surface on the three best faces clear of defect in not more than three cuttings. Lumber from such logs will normally grade at least 35 percent No. 1 common and better.

Grade 3 - Merchantable logs at least 8.0 inches in diameter inside bark which do not meet the requirements of higher grades. Such logs will normally produce less than 35 percent No. 1 common and better lumber or will be suitable only for ties or timbers.

Species listed

Redcedar	-	<u>Juniperus virginiana</u>
Cypress	-	<u>Taxodium distichum</u>
White oak	-	<u>Quercus alba</u>
Post-oak group includes:		
Post oak	-	<u>Quercus stellata</u>
Swamp white oak	-	<u>Quercus bicolor</u>
Swamp chestnut oak	-	<u>Quercus prinus</u>
Overcup oak	-	<u>Quercus lyrata</u>
Bur oak	-	<u>Quercus macrocarpa</u>
Chinquapin oak	-	<u>Quercus muehlenbergii</u>
Chestnut oak	-	<u>Quercus montana</u>
Black oak	-	<u>Quercus velutina</u>
Northern red oak includes:		
Northern red oak	-	<u>Quercus borealis</u>
Swamp red oak	-	<u>Quercus falcata var. pagodaefolia</u>
Other red oaks include:		
Southern red oak	-	<u>Quercus falcata</u>
Pin oak	-	<u>Quercus palustris</u>
Scarlet oak	-	<u>Quercus coccinea</u>
Willow oak	-	<u>Quercus phellos</u>
Water oak	-	<u>Quercus nigra</u>
Shingle oak	-	<u>Quercus imbricaria</u>
Hickory	-	<u>Carya species</u>
Elm	-	<u>Ulmus species</u>
Soft maple includes:		
Red maple	-	<u>Acer rubrum</u>
Silver maple	-	<u>Acer saccharinum</u>
Boxelder	-	<u>Acer negundo</u>
Sugar maple	-	<u>Acer saccharophorum</u>
Sycamore	-	<u>Platanus occidentalis</u>
Ash	-	<u>Fraxinus species</u>
Basswood and yellow-poplar includes:		
Basswood	-	<u>Tilia species</u>
Yellow-poplar	-	<u>Liriodendron tulipifera</u>
Cucumbertree	-	<u>Magnolia acuminata</u>

Species listed (continued)

Cottonwood - Populus deltoides
Black walnut - Juglans nigra
Other hardwoods - include all other commercial hardwood species.

Noncommercial species - include species that do not normally have commercial value such as hawthorn, redbud, ironwood, alder, and service berry.

TERRITORY SERVED BY THE
CENTRAL STATES FOREST EXPERIMENT STATION
FOREST SERVICE
UNITED STATES DEPARTMENT OF AGRICULTURE

● RESEARCH CENTERS

▲ EXPERIMENTAL FORESTS

★ CENTRAL STATES HEADQUARTERS