1902

USNM

EX LIBRIS

William Healey Dall

Division of Mollusks
Sectional Library

I

Catlow

Division of Mollusks
Sectional Library

594.08

C 365

8mollusk

QL
403
C 36
1843
MOLL

POPULAR

CONCHOLOGY;

OR,

THE SHELL CABINET ARRANGED:

BEING

An Introduction

TO

THE MODERN SYSTEM OF CONCHOLOGY.

WITH A SKETCH OF

THE NATURAL HISTORY OF THE ANIMALS,

AN ACCOUNT OF THE FORMATION OF THE SHELLS,

AND

A COMPLETE DESCRIPTIVE LIST OF THE FAMILIES

AND GENERA.

BY

AGNES CATLOW.

LONDON:

PRINTED FOR

LONGMAN, BROWN, GREEN, AND LONGMANS,
PATERNOSTER-ROW.

1843.

LONDON:
Printed by A. SPOTTISWOODE,
New-Street-Square.

108

TO
MRS. CORRIE,
OF WOODVILLE, NEAR BIRMINGHAM;
WHOSE ACQUIREMENTS IN VARIOUS BRANCHES
OF SCIENCE ARE EQUALLED BY
THE GENEROSITY AND BENEVOLENCE WITH WHICH
SHE MAKES HER KNOWLEDGE AVAILABLE
TO OTHERS ;

This little Work,
INTENDED TO RENDER THE STUDY OF CONCHOLOGY
POPULAR,
AND WHICH OWES ITS PUBLICATION TO HER KIND
ENCOURAGEMENT,
IS
GRATEFULLY DEDICATED BY
THE AUTHOR.

P R E F A C E.

THE following pages are designed to facilitate the study of a branch of Natural History which is daily becoming more attractive and more important; not only from its intrinsic interest, but also from its multiplied relations with Geology. They will furnish the young Conchologist with an elementary manual, which, although scientific in its form, is, by the simplicity of its method and the familiarity of its style, calculated effectually to assist him in the early steps of his progress in this fascinating pursuit. The satisfaction he may have derived from possessing specimens of shells cannot fail to be greatly increased when, by the aid of this little volume, he has learned how to arrange them in proper scientific order, and discovered their names, localities, uses, and habits. When his cabinet has been thus

arranged, he will view it with a delight infinitely greater than that which could be afforded while it remained in confusion; and he will be further induced to enlarge his knowledge of the subject by consulting more extended and more elaborate treatises; nor will he rest satisfied until he becomes complete master of the science in all its bearings.

Among the many works which have been published on Conchology, there is none which appears exactly to answer the purpose intended by the present volume. Most of them are addressed exclusively to men of science, and are encumbered (to them enriched) with a vast mass of details, the attempt to embrace which would only confuse and embarrass a beginner; while others, which profess to be elementary, are inapplicable to the purposes of exact arrangement. The system which has been here followed is that adopted by Lamarck in his great work, entitled "*Les Animaux sans Vertèbres*;" but the genera established since that work was written have been inserted wherever it was possible. To each new genus is attached

the name of the naturalist who introduced it, in order that it may be distinguished from those mentioned by Lamarck himself. To avoid confusion, however, no great change has been made in the systematic arrangement.

The author wishes, in this place, to acknowledge the kind assistance, and valuable information, she has received from a gentleman well versed in the science, but to whom she is personally unknown; and to express her thanks to a kind friend for some of the drawings by which these pages are illustrated. She would be ungrateful, were she not also to acknowledge the instruction she has received from her friend Mr. Swainson, whose works on Natural History are so well known to the public, and who first turned her thoughts to the study of Conchology, which she trusts her humble efforts will have rendered interesting to those for whose information this volume was written.

CONTENTS.

			Page
INTRODUCTORY CHAPTER	-	-	1
PART I.—ANNULATA	-	-	28
FAMILY I.— <i>Dorsalidæ</i>	-	-	29
1. Arenicola	-	-	29
2. Siliquaria	-	-	30
FAMILY II.— <i>Maldanidæ</i>	-	-	31
3. Clymene	-	-	31
FAMILY III.— <i>Amphitritidæ</i>	-	-	31
4. Pectenaria	-	-	31
5. Sabellaria	-	-	32
6. Terebella	-	-	32
7. Amphitrite	-	-	33
FAMILY IV.— <i>Serpulidæ</i>	-	-	34
8. Spirorbis	-	-	34
9. Serpula	-	-	34
10. Vermilia	-	-	35
11. Galeolaria	-	-	35

	Page
PART II.—CIRRHIPEDA - -	36
FAMILY I.— <i>Cirrhipeda sedentaria</i> - -	37
12. Tubicinella - - -	37
13. Coronula - - -	37
14. Balanus - - -	38
15. Acasta - - -	39
16. Creusa - - -	39
17. Pyrgona - - -	39
FAMILY II.— <i>Cirrhipeda pedunculata</i> - -	40
18. Lithotrya - - -	40
19. Anatifa - - -	41
20. Scalpellum - - -	43
21. Pollicipes - - -	43
22. Cineras - - -	43
23. Otion - - -	44
PART III.—CONCHIFERA OR BIVALVES -	45
ORDER I.—BRACHIOPODA - - -	47
FAMILY I.— <i>Tendinosa</i> - - -	48
1. Lingula - - -	48
2. Terebratula - - -	49
FAMILY II.— <i>Adhærentia</i> - - -	50
3. Thecidium - - -	50
4. Crania - - -	51
5. Orbicula - - -	52
ORDER II.—DIMYARIA - - -	53
DIVISION I.—CONCHIFERA CRASSIPEDA - -	54
FAMILY I.— <i>Tubicolidæ</i> - - -	54
6. Aspergillum - - -	54

	Page
7. Clavagella - - -	56
8. Fistulana - - -	57
9. Gastrochæna - - -	58
10. Teredo - - -	58
FAMILY II.— <i>Pholadidæ</i> - - -	60
11. Xylophaga - - -	61
12. Pholas - - -	61
FAMILY III.— <i>Solenidæ</i> - - -	63
13. Solen - - -	63
14. Solecurtus - - -	65
15. Pholadomia - - -	66
16. Panopæa - - -	66
17. Glycimeris - - -	67
18. Solemya - - -	68
FAMILY IV.— <i>Myidæ</i> - - -	68
19. Mya - - -	68
20. Anatina - - -	69
21. Thracia - - -	70
22. Corbula - - -	71
23. Pandora - - -	71
24. Anatinella - - -	72
DIVISION II.—CONCHIFERA TENUIPEDA - - -	72
FAMILY V.— <i>Mactridæ</i> - - -	73
25. Lutraria - - -	73
26. Mactra - - -	73
27. Crassatella - - -	74
28. Mesodesma - - -	75
29. Erycina - - -	75
30. Ungulina - - -	76
31. Amphidesma - - -	76

				Page
FAMILY VI.— <i>Lithophagidæ</i>	-	-	-	77
32. Saxicava	-	-	-	77
33. Petricola	-	-	-	78
34. Venerupis	-	-	-	78
FAMILY VII.— <i>Nymphidæ</i>	-	-	-	79
35. Sanguinolaria	-	-	-	80
36. Psammobia	-	-	-	80
37. Galeomma	-	-	-	81
38. Tellina	-	-	-	81
39. Corbis	-	-	-	83
40. Lucina	-	-	-	83
41. Donax	-	-	-	84
42. Capsa	-	-	-	84
DIVISION III.—CONCHIFERA LAMELLIPEDA				85
FAMILY VIII.— <i>Conchidæ</i>	-	-	-	85
43. Cyclas	-	-	-	86
44. Cyrena	-	-	-	86
45. Galathea	-	-	-	87
46. Cyprina	-	-	-	87
47. Astarte	-	-	-	88
48. Cytheria	-	-	-	89
49. Venus	-	-	-	90
FAMILY IX.— <i>Cardiidæ</i>	-	-	-	91
50. Cardium	-	-	-	91
51. Cardita	-	-	-	92
52. Cypricardia	-	-	-	93
53. Isocardia	-	-	-	93
FAMILY X.— <i>Arcidæ</i>	-	-	-	94
54. Cucullæa	-	-	-	95
55. Arca	-	-	-	95
56. Pectunculus	-	-	-	96

				Page
57. Nucula	-	-	-	97
58. Trigonia	-	-	-	97
FAMILY XI.— <i>Naiidæ</i>	-	-	-	98
59. Unio	-	-	-	100
60. Hyria	-	-	-	101
61. Anodonta	-	-	-	101
62. Iridina	-	-	-	102
DIVISION IV.—CONCHIFERA AMBIGUA	-	-	-	103
FAMILY XII.— <i>Chamidæ</i>	-	-	-	103
63. Diceras	-	-	-	103
64. Chama	-	-	-	104
65. Etheria	-	-	-	105
ORDER III.—MONOMYARIA	-	-	-	106
DIVISION I.—LIGAMENT LONG AND MARGINAL	-	-	-	107
FAMILY I.— <i>Tridacnidæ</i>	-	-	-	107
66. Tridacna	-	-	-	107
67. Hippopus	-	-	-	109
FAMILY II.— <i>Mytilidæ</i>	-	-	-	109
68. Modiola	-	-	-	110
69. Lithodomus	-	-	-	110
70. Mytilus	-	-	-	111
71. Pinna	-	-	-	112
FAMILY III.— <i>Malleidæ</i>	-	-	-	114
72. Crenatula	-	-	-	114
73. Perna	-	-	-	115
74. Malleus	-	-	-	116
75. Vulsella	-	-	-	117
76. Avicula	-	-	-	118

	Page
DIVISION II.—LIGAMENT NOT MARGINAL -	121
FAMILY IV.— <i>Pectenidæ</i> - - -	121
77. <i>Pedum</i> - - -	121
78. <i>Lima</i> - - -	122
79. <i>Pecten</i> - - -	123
80. <i>Plicatula</i> - - -	125
81. <i>Spondylus</i> - - -	125
FAMILY V.— <i>Ostreidæ</i> - - -	126
82. <i>Ostrea</i> - - -	126
83. <i>Placuna</i> - - -	129
84. <i>Placunanomia</i> - - -	130
85. <i>Anomia</i> - - -	132
PART IV.—MOLLUSCA or UNIVALVES -	133
ORDER I.—PTEROPODA - - -	136
1. <i>Hyalæa</i> - - -	136
2. <i>Clio</i> - - -	136
3. <i>Cleodora</i> - - -	137
4. <i>Limacina</i> - - -	137
5. <i>Cymbulia</i> - - -	138
6. <i>Pneumodermon</i> - - -	138
ORDER II.—GASTEROPODA - - -	139
FAMILY I.— <i>Tritonidæ</i> - - -	139
7. <i>Glaucus</i> - - -	140
8. <i>Eolis</i> - - -	140
9. <i>Tritonia</i> - - -	140
10. <i>Scyllæa</i> - - -	140
11. <i>Tethys</i> - - -	140
12. <i>Doris</i> - - -	140
FAMILY II.— <i>Dentalidæ</i> - - -	140
13. <i>Dentalium</i> - - -	140

				Page
FAMILY III.— <i>Phyllididæ</i>	-	-	-	141
14. Phyllidea	-	-	-	141
15. Chitonellus	-	-	-	142
16. Chiton	-	-	-	142
17. Patella	-	-	-	143
18. Siphonaria	-	-	-	146
FAMILY IV.— <i>Semi-Phyllididæ</i>	-	-	-	147
19. Pleurobranchus	-	-	-	147
20. Umbrella	-	-	-	147
FAMILY V.— <i>Calyptræidæ</i>	-	-	-	148
21. Parmophorus	-	-	-	149
22. Emarginula	-	-	-	149
23. Fissurella	-	-	-	150
24. Crepidula	-	-	-	151
25. Calyptræa	-	-	-	152
26. Hipponyx	-	-	-	153
27. Pileopsis	-	-	-	154
FAMILY VI.— <i>Bullæidæ</i>	-	-	-	155
28. Acera	-	-	-	155
29. Bullæa	-	-	-	155
30. Bulla	-	-	-	156
FAMILY VII.— <i>Aplysidæ</i>	-	-	-	157
31. Aplysia	-	-	-	157
32. Dolabella	-	-	-	158
FAMILY VIII.— <i>Limacidæ</i>	-	-	-	159
33. Onchidium	-	-	-	159
34. Limax	-	-	-	160
35. Parmacella	-	-	-	160
36. Testacella	-	-	-	160
37. Vitrina	-	-	-	162

				Page
ORDER III.—TRACHELIPODA	-	-	-	163
DIVISION I.—PHYTOPHAGA	-	-	-	164
FAMILY I.— <i>Colimacidae</i>	-	-	-	164
38. <i>Helix</i>	-	-	-	165
39. <i>Carocolla</i>	-	-	-	170
40. <i>Anastoma</i>	-	-	-	171
41. <i>Helicina</i>	-	-	-	172
42. <i>Pupa</i>	-	-	-	172
43. <i>Clausilia</i>	-	-	-	173
44. <i>Bulimus</i>	-	-	-	174
45. <i>Achatina</i>	-	-	-	178
46. <i>Succinea</i>	-	-	-	179
47. <i>Auricula</i>	-	-	-	180
48. <i>Cyclostoma</i>	-	-	-	181
FAMILY II.— <i>Lymnæidae</i>	-	-	-	181
49. <i>Planorbis</i>	-	-	-	182
50. <i>Physa</i>	-	-	-	183
51. <i>Lymnea</i>	-	-	-	183
52. <i>Ancylus</i>	-	-	-	185
FAMILY III.— <i>Melanidae</i>	-	-	-	186
53. <i>Melania</i>	-	-	-	186
54. <i>Melanopsis</i>	-	-	-	187
FAMILY IV.— <i>Peristomidæ</i>	-	-	-	187
55. <i>Valvata</i>	-	-	-	188
56. <i>Paludina</i>	-	-	-	188
57. <i>Ampullaria</i>	-	-	-	189
58. <i>Ampullacera</i>	-	-	-	191
FAMILY V.— <i>Neritidæ</i>	-	-	-	192
59. <i>Navicella</i>	-	-	-	192

				Page
60.	Neritina	-	-	193
61.	Nerita	-	-	194
62.	Natica	-	-	195
FAMILY VI.—	<i>Janthinidæ</i>	-	-	196
63.	Janthina	-	-	196
FAMILY VII.—	<i>Macrostomidæ</i>	-	-	197
64.	Sigaretus	-	-	197
65.	Stomatella	-	-	198
66.	Stomatia	-	-	198
67.	Haliotis	-	-	199
FAMILY VIII.—	<i>Plicacidæ</i>	-	-	202
68.	Tornatella	-	-	202
69.	Melampus	-	-	203
70.	Pyramidella	-	-	203
FAMILY IX.—	<i>Scalaridæ</i>	-	-	204
71.	Vermetus	-	-	205
72.	Magilus	-	-	205
73.	Leptoconchus	-	-	207
74.	Scalaria	-	-	207
75.	Rissoa	-	-	209
76.	Delphinula	-	-	209
FAMILY X.—	<i>Turbinidæ</i>	-	-	210
77.	Solarium	-	-	210
78.	Rotella	-	-	211
79.	Trochus	-	-	211
80.	Monodonta	-	-	214
81.	Turbo	-	-	214
82.	Planaxis	-	-	216
83.	Phasianella	-	-	216
84.	Turritella	-	-	217

				Page
DIVISION II.—ZOOPHAGA	-	-	-	218
FAMILY XI.— <i>Canaliferidæ</i>	-	-	-	219
85. Cerithium	-	-	-	219
86. Pleurotoma	-	-	-	221
87. Turbinella	-	-	-	222
88. Cancellaria	-	-	-	223
89. Fasciolaria	-	-	-	223
90. Fusus	-	-	-	224
91. Pyrula	-	-	-	224
92. Struthiolaria	-	-	-	226
93. Ranella	-	-	-	227
94. Murex	-	-	-	227
95. Triton	-	-	-	228
FAMILY XII.— <i>Pteridæ</i>	-	-	-	229
96. Rostellaria	-	-	-	230
97. Pteroceras	-	-	-	231
98. Strombus	-	-	-	232
FAMILY XIII.— <i>Purpuridæ</i>	-	-	-	233
99. Cassidaria	-	-	-	234
100. Oniscia	-	-	-	235
101. Cassis	-	-	-	235
102. Ricinula	-	-	-	237
103. Purpura	-	-	-	237
104. Monoceras	-	-	-	238
105. Concholepas	-	-	-	239
106. Harpa	-	-	-	239
107. Dolium	-	-	-	241
108. Buccinum	-	-	-	242
109. Nassa	-	-	-	243
110. Eburna	-	-	-	243
111. Terebra	-	-	-	244

			Page
FAMILY XIV.— <i>Columellidæ</i>	-	-	245
112. Columbella	-	-	245
113. Mitra	-	-	246
114. Voluta	-	-	247
115. Melo	-	-	249
116. Cymba	-	-	249
117. Marginella	-	-	250
118. Volvaria	-	-	251
FAMILY XV.— <i>Convolutidæ</i>	-	-	251
119. Cypræa	-	-	252
120. Erato	-	-	255
121. Ovula	-	-	255
122. Terebellum	-	-	256
123. Ancillaria	-	-	256
124. Oliva	-	-	257
125. Conus	-	-	258
ORDER IV.—CEPHALOPODA	-	-	260
DIVISION I.—POLYTHALAMA	-	-	263
126. Belemnites	-	-	263
127. Orthoceras	-	-	264
128. Spirula	-	-	264
129. Nummulites	-	-	265
130. Nautilus	-	-	265
131. Ammonites	-	-	267
132. Turrilites	-	-	268
133. Baculites	-	-	269
DIVISION II.—MONOTHALAMA	-	-	269
134. Argonauta	-	-	269
DIVISION III.—NO EXTERNAL SHELL	-	-	272
135. Octopus	-	-	272

				Page
136. Loliopsis	-	-	-	273
137. Loligo	-	-	-	273
138. Sepia	-	-	-	273
ORDER V.—HETEROPODA	-	-	-	274
139. Carinaria	-	-	-	274
140. Atlanta	-	-	-	276
141. Pterotrachea	-	-	-	276
142. Phylliröe	-	-	-	276
GUIDE TO THE USE OF THE TABLES	-	-	-	278
AN ALPHABETICAL LIST OF ORDERS, DIVISIONS, FAMILIES, &c.	-	-	-	280
AN ALPHABETICAL LIST OF GENERA	-	-	-	283
ENGLISH NAMES USED IN THIS VOLUME	-	-	-	292
EXPLANATION OF TERMS	-	-	-	293

ERRATA.

- Page 29. line 4. for "*Dorsaliadæ*" read "*Dorsalidæ*."
31. line 1, for "*Maldaniadæ*" read "*Maldanidæ*."
36. line 1. of note, for "*Magillus*" read "*Magilus*."
49. line 10. for "*Cumming*" read "*Cuming*."
173. line 8. for "*CLAUSILEA*" read "*CLAUSILIA*."
line 15. for "*Clausileæ*" read "*Clausiliæ*."
188. under first wood-cut add "c, young state of
Valvata ; a, b, more advanced."

POPULAR CONCHOLOGY.

INTRODUCTORY CHAPTER.

THE study of shells is one of the most interesting that can be pursued by a lover of Natural History. If he confine himself to their external appearance, and compare them with the other orders of creation, with reference only to the beauties they present to the eye, he will find that they are inferior to none in form or colour, exhibiting the most exquisite contours, with the most varied and beautiful tints; if he pursue his examination further, and without limiting himself to the exterior, fix his attention upon the living animal, as well as upon the inanimate shell in which it resides, and which it constructs itself, he will be amply repaid by the search; he will find the same harmony existing in all its rela-

tions, the same adaptation of means to the end required, the same fitness of the creature for the element in which it is to reside, and the mode of life it is to pursue, as he will trace in the more complicated and apparently more perfect classes of animals; finally, should he enter into a more scrutinizing investigation, and make it his object to become acquainted with the physiology of the animal, apart from its shell, and to pursue its history from its production to its death, following its gradual increase and advancement to its perfect state, including in his researches the formation of its shell, and the mode by which it receives enlargement and colour, he will here again find the study of shells as interesting as that of any portion of the created kingdom.

The most familiar name by which the study of shells is known is *Conchology*.* In pursuing this pleasing branch of Natural History, whether an extended or limited view is taken, it is necessary, in the first place, to know something generally of the animal, its growth, the

* From *conchylion*, Gr., or *concha*, Lat., *a shell*, and *logos*, Gr., *a discourse*.

mode by which it forms its shell, its place of residence, and habits, before entering upon a minute and classified division. The present chapter will therefore have these objects in view.

The animals which inhabit and form shells are called *Mollusca* *, and, as their name signifies, have a soft body, most of them being entirely destitute of bones, though some are provided with internal shells, more or less perfectly formed; some species of slug, for instance, have internally a rudely shaped hard body, and others a mass of small grains. To compensate for the apparent deficiency in this respect, and to enable them to provide themselves with a defence against any injury to which they might be liable from so delicate a structure, they have been furnished with an external skin or envelope, covering them more or

* From *mollis*, Lat., *soft*, and *esca*, Lat., *flesh*. Lamarck gives the name of Mollusca only to those which inhabit univalve shells, but naturalists now generally comprise under that appellation the inhabitants of both univalves and bivalves.

less completely, and from which they are able to exude a liquid, which, on exposure to the air or water, hardens into shell, and thus most of them are enabled to cover themselves with a secure, and at the same time commodious and beautiful habitation, into which they can, by the aid of strong contractile muscles, retire far enough to escape from danger when it approaches. This outer skin is now usually called the *mantle*, and is common to most species of mollusca. As nearly all these animals live entirely in the water, from which they derive their nutriment, and which they breathe by the aid of *branchiæ**, resembling the gills of fish, it is necessary that a means should be provided for the admission of that fluid to the interior of the body, and in order to effect this, certain openings occur in the mantle, through which the water passes, and by which also the head and foot, when these limbs exist, are put forth and drawn back at the will of the animal.

Mollusca do not all possess heads, one entire class being quite destitute; when they do exist,

* The fringed appendages in the interior of the oyster are the *branchiæ* of that Mollusc.

they are usually provided with *tentacula*, by which the animals feel their way, and which they have the power of easily retracting when danger threatens; they have occasionally also

eyes, and other organs of sense, but they vary much in this respect, some possessing those of sight and hearing*, whilst others are devoid of both, and apparently enjoy only that of touch. That limb, which, from its being an organ of motion, has been called the *foot*, is much more universally found amongst mollusca than the head, and is a member of considerable importance. A few words only are necessary with regard to the internal construction of these animals; they possess the usual organs necessary for the support of life, a nervous system, a heart, with its attendant veins and arteries, furnished with a cold and white or blueish blood, and an apparatus for breathing, consisting, in those which live in water, of branchiæ placed externally or internally, and so arranged as to

* The latter sense, it is said, is confined to the *Cephalopoda*.

have a stream of water continually presented to them, and in those which live in air, of organs fit for its respiration.

Mollusca are to be found in all parts of the world, in the sea, in rivers, and on land, and differ in their construction and habits according to their locality. A great number have their residence in the sea, and are unable to exist out of salt water; of these some float on the surface, others have the power of moving through the deep waters, and others again are attached by various means to rocks and other substances below the surface. Many species are met with, living on the sea coast, in the sand, mud, or cliffs, and many in rivers: the land also has its peculiar varieties, either living under ground, in holes of trees, or in moist and marshy places. These different habitations give a general name to the *Mollusca* which are found in them, as *Pelagic**, to those which reside in the deep recesses of the ocean, *Litoral*†, to such as are found on the coast,

* From *pelagus*, Lat., *the depth of the sea*.

† From *litoris*, Lat., *of the shore*.

*Fluviatile**, to those which inhabit rivers, ponds, &c., and *Terrestrial*†, to the species which are met with on the land.

The methods of *locomotion* adopted by those Mollusca which move from one place to another, are very various. A great number, like the oyster, are perfectly stationary, and do not, when once fixed, move from the place they have selected; there are, however, several genera that are continually moving about, and various and beautiful are the contrivances by which they effect this object. Many move along the surface of the ground or bottom of the sea, by means of their foot, which they thrust out to a distance in advance, and fix to some solid object, after effecting which, a strong muscular contraction enables them to draw their body to it; it is then again protruded in a similar manner, and by this means, the animal moves slowly: others, using an action the reverse of this, push backwards with the foot, and thus impel themselves forward, and others, still however making use of the

* From *fluvius*, Lat., a river.

† From *terrestris*, Lat., earthly.

foot for the purpose, contrive to advance with considerable agility, by coiling it up tightly, and suddenly relaxing it, by which means, serving as a spring, it urges the body to some distance. There are many species of Mollusca which are unable to move along the rough surface of the ground, but manage to advance rapidly enough along the smooth marine plants, by a kind of sliding motion. Those tribes which do not creep along the ground, but move through the water, are obliged to have recourse to various methods to enable them to swim; some float along the surface, borne as the winds and waves direct; others, by squirting out suddenly a stream of water, which they have taken into their shells, send themselves forward in an opposite direction; and others swim, by using their foot as a fin, and moving it from one side to the other. In those Mollusca which affix themselves permanently to a submarine object, the method of attachment varies considerably; some, for instance, are fixed by the shell, some by a continuation of the foot either in a mass, or divided into separate fibres, some by a tube, and others

by a number of threads, which they spin and fix to different parts of the rock or other substance on which they dwell. In the locomotive Mollusca the foot is often serviceably employed for the purpose of burrowing in the sand and mud, to enable them to conceal themselves underneath, being used as a scoop with much effect, and thus enabling the animal to hide itself in a short time, and to emerge when all danger is over.

With regard to the *food* on which Mollusca live, there is as great a variety as exists among animals of a more complicated organization: some are carnivorous, or *Zoophaga**, others herbivorous, or *Phytophaga*†, and the substances on which they feed, whether animal or vegetable, are consumed by them in various states, some preferring the *living*, whilst others prefer the *dead* or *putrescent* form. To meet these different forms of nutrition, various and diversified organs are required; and it will be found that in these animals, as well as in the

* From *zo-on*, Gr., *an animal*, and *phago*, Gr., *to eat*.

† From *phyton*, Gr., *a vegetable*, and *phago*.

other ranks of creation, the appropriate apparatus is adapted in every case. Amongst those which are *carnivorous*, some that have the power of locomotion are furnished with appendages, by which they can capture their prey; others, that are fixed, by producing a slight current, obtain a constant stream of water supplied to their mouths, bringing with

it minute animals, which they retain. The mouth of the former is usually in the form of a proboscis, which is furnished with little hooks, with which they can penetrate the shells of those they feed on, and extract the contents. Those which are *herbivorous*, cut and masticate their food. In many cases they are provided with a single tooth, which they fix upon the substance they are about to devour. Many have teeth acting

in various ways, but of these the headless Molluscs are destitute.

Having thus given a short general description of the animals inhabiting shells, the next object of inquiry is the formation of *shell* itself. “The molluscous animals are *oviparous*, producing their young from eggs in the same way as birds, although many of the tribes do not, at the time of their exclusion, abandon them at once, but deposit them between the two membranes of the branchial laminae, where they undergo a species of incubation.”* “The animal has a small portion of shell attached to it, even before the egg is hatched: while enclosed, the shell is generally of a pale horn colour, and destitute of markings; when, therefore, they remain attached to the apex of the spire of adult shells, they may be easily distinguished by their appearance from the part formed after their exclusion.”† As the animal increases in bulk, it finds the shell is not sufficiently large to cover it, and it then adds to its

* From the Penny Cyclopædia, article *Conchifera*.

† From Mr. Gray, Philosophical Trans. 1833.

dimensions by the aid of its *mantle*. This portion of the body of the animal, as before mentioned, has the power of exuding a liquid matter, which, on exposure to the air or water, becomes shell, and the mollusc makes the addition to its original habitation by protruding the mantle beyond the edge, and suffering it in that manner to remain exposed: in a short time it is found to be covered with a fluid which has been secreted, and this gradually and slowly assumes consistence, fresh layers being added within, each projecting a little farther at the edge, and thus the new formation becomes similar to the shell already in existence. This operation is repeated at intervals, as the animal increases, so that by this beautiful contrivance the shell and its inhabitant are always adapted to each other, and preserve their relative proportions.

As the new portion of shell is always similar in shape, and, in fact, takes its form from that part of the mantle which is pushed out, it is evident that the conformation of every shell will depend upon the manner in which the Mollusc protrudes its mantle, and that it will

be spiral, cup-shaped, or otherwise, according to circumstances. This can be best explained by a few references : thus, the simplest form of shell is that of the *Limpet*, which has a very small portion (represented at A in the figure),

formed at the period of its birth ; as it increases in size, and finds this inadequate to protect its body, it expands the border of its mantle, and, by the process already described, enlarges its shell all round, until it presents the appearance of A B ; in like manner, at different periods, as its growth requires, it enlarges it to C, D, and E, at which time, having attained its full size, it ceases to require any farther enlargement, and the shell is then completed or mature.

The *Limpet* presents the most simple method of growth, as in it the mantle is extended equally at every part of the circumference of

the original shell, and the *size* merely is increased, still preserving the same shape; in the greater number of univalve shells, however, the method of enlargement is different, the mantle being more active on one side of the aperture than on the other: this gives rise to a *turbinated* or *spiral* shell, in which each successive portion of the spire, which forms an entire circle, is called a *whorl*. A shell of this nature is here shown. The new formations in this genus can easily be traced, each ridge represented consisting of a thick lip, which, at certain periods of its growth, the animal has formed at the edge of the shell then existing. These spiral shells are almost always formed with the whorls turning towards the *right*, which arises from the *left* side of the mantle being the most active, in consequence of the position of the heart and blood-vessels of the animal on that side*: in a few genera, however,

* Dr. Grant makes the following observations on this subject, in his *Lectures on Comparative Anatomy*, at the

the heart is situated on the *right* side, and then the whorls of the shell are turned to the *left*, and the shells in which this conformation takes place are called *reversed*, or *sinistral**, in opposition to the common, or *dextral*. †

The growth of bivalve shells is carried on in the same way, each valve being simultaneously enlarged, as will be seen in the annexed figure, where the additions may be easily traced.

London University :—"I am inclined to ascribe this spiral twist of shells to the action of the heart and the respiratory organs, which are situated on the left side, and to the descent which the foot of the animal makes over the columella, in order to reach a solid surface to creep upon. The dilatation of the heart bends the anterior free part of the body to the right side, and would cause the shell to revolve on the same plane, were that cause to operate alone ; but the constant descent of the foot over the surface of the pillar causes the plane of revolution to change at every instant, and by these two causes the spiral form of shells appears to be produced."

* From *sinister*, Lat., *left*.

† From *dexter*, Lat., *right*.

The power of forming shell does not appear to be confined to the *edge* of the mantle, but exists throughout its whole extent: this has been proved by removing part of the shell of a living snail, at a distance from the edge or lip: the animal immediately sets about filling up the gap, and effects it in the course of a few days by a succession of thin strata of liquid shelly matter, which exude from that portion of the mantle which has been laid bare.

The regularity of the growth of many shells appears to be interrupted at particular times by changes which arise in the shape of the mantle; this is observable in several species, which exhibit cones or spines projecting from their surface, and which have been produced by the mantle altering its form, and dividing into several long slender branches. On these processes shell is formed, constituting a row of hollow spines, which are gradually filled up by successive depositions of shelly matter, and eventually become solid; the mantle being then withdrawn from them, proceeds in the formation of new shell, branching out at regular intervals into another row of spines; and in this

manner are produced the prickles, ridges, and other excrescences visible in many genera, the *Murex*, for example. Analogous to these variations in the deposit of shell by the mantle, but differing, inasmuch as they only occur at one period, are the developements which take place when the mollusc has arrived at its full growth. In the cut are represented the immature and

mature states of the same shell. It will be seen that they differ widely; and the variation has been caused by the mantle forming the extension of shell with its accompanying projecting tubes, as a completion to the animal's habitation: it marks the termination of its labours, for the shell having now become perfect, and

the mollusc having arrived at its full growth, the secretion of shell is no longer required to be called into action for the purpose of increasing the size of its habitation ; and from this period the animal, if it continue to produce shell, only adds to the thickness and polish of its mansion, without extending its dimensions. “ It should be added, that all shells whose spires are exposed, and, being thin in their young state, would be liable to be broken off by the action of the sea, have that part strengthened by the internal deposition of shelly matter. Some few shells, however, as the *Bulimus decollatus*, for instance, lose the upper part of their spire ; for the animal, instead of lining the upper whorls, withdraws its body from them, and forms a division : the communication of the body and the *apex* of the shell being cut off, the latter part decays in the manner of a dead shell, and falls off in particles.”*

Many of the turbinated or spiral univalve Mollusca have the power of forming an additional protection to their bodies, by construct-

* From Mr. Gray, Philosophical Trans. 1833.

Opercula.

ing a kind of door, called the *operculum**, or *epiphragma*†, to the entrance of their shell. These are made more or less strongly, as circumstances require; some, as the snail and other land species, merely needing this extra covering during severe weather, at the end of that time throw it off, and on a recurrence of the necessity make another; these being, therefore, of a temporary nature only, are slight and easily formed, but there are other genera which have them permanently attached, and of a more solid description‡: these latter are fixed to the *foot*

* From *operculum*, a cover or lid.

† From *epiphrasso*, Gr., to stop up.

‡ The temporary clausilia, or doors, are called *epiphragmata*, the permanent, *opercula*.

of the animal, and are enlarged in size by the mollusc, as the shell is increased, in order that they may correspond with and close the aperture. They are found attached to the animal before the egg is hatched. In the foregoing figure are delineations of several varieties of *opercula*. Some are shelly, and others of a horny construction.

The beautiful and varied *colours* which shells present are produced by colouring matter contained in the pores, which exist in *the front part only* of the mantle, and which matter is made to penetrate the substance of the shell before it becomes hardened: the experiment before mentioned* will serve also to exemplify this. The portion of shell of which the animal has been deprived (and this may be done without injury to the snail,) will be reproduced by the action of the mantle, but the colours will be different according to the situation of the part of the shell which has been broken: if the piece be removed and reproduced *at the external lip* of the shell, where the front part of the mantle lies, the natural colours will appear in

* See page 16.

the new piece; but if the fracture has been made *at a distance from the aperture*, and in the body of the shell, a new portion will indeed be made by that part of the mantle which has been laid bare, but it will be white or nearly colourless; thus proving, that, although every part of the mantle can produce shell, *the external edge* alone can infuse colour into it. The brilliancy of the colours observed in shells is exceedingly influenced by the degree of health, and by the food, light, and heat which the mollusc has had the power of enjoying. Those which inhabit tropical countries have often a very great degree of brilliancy and beauty; whilst others which are less favoured by climate, and, still more remarkably, those which remain all their lives enclosed in wood, chalk, &c., have little or no colour. It is observable, also, in a similar manner, that a great difference frequently exists in the *size* of mature shells of the same species, particularly in such as have the strongly marked lip, or other final appendage: this may be accounted for by the strength or weakness of constitution existing in

different individuals, influenced undoubtedly by favourable or unfavourable situations, and abundance or scarcity of food. But this opinion can only be satisfactorily proved or disproved when shells are better known. Mr. Gray has the following remarks on the subject :— “ The *thickness, roughness, or smoothness* of shells appears to depend in a great degree on the stillness or agitation of the water which they inhabit. Two specimens of the same species of shell will often be found very different in appearance, according to the situation in which they have been placed. For instance, the common *Buccinum* of our coast is thick, solid, and heavy, when found in a rough sea, and thin, light, and smooth, when found in the still water of harbours. Boring shells are generally influenced in regard to their size, thickness, and form, by the hardness or softness of the rock in which they are found. Land shells are much influenced, as regards their size, by their temperature, altitude, and abundance of food. The shells of *Helix nemoralis*, and *Helix hortensis* found near London, are not above two thirds of the size of

those which occur in Portugal and the south of France.”*

Of the *uses* and *benefits* derived by man from shells and their inhabitants, a hasty mention may be made. First, as to their *inhabitants*; amongst civilised nations many species are highly prized as articles of food, such as the *oyster*, *mussel*, &c.; and among savage nations, particularly those inhabiting the coasts of Western Africa and New Holland, a great part of their food is derived from Mollusca of various kinds. Their principal use, however, in this respect appears to be, the serving as food for the classes of animals more immediately above them in the scale of organization, such as the amphibious mammalia, sea-birds, and fish, which in a great measure live upon them, destroying such vast quantities, that it is their extraordinary fecundity alone which prevents them from being totally annihilated. A silky thread is produced by the *Pinna*, which may be spun, and is occasionally made into gloves, &c., but more as a curiosity than for any useful

* Philosophical Trans., 1833.

purpose ; and from several species of Mollusca may be procured a strong colouring matter of a purple hue, which formerly was used as a dye, but has now been supplanted by other animal, mineral, and vegetable productions, which are easier of access. Secondly, as regards the *shell* ; in many countries, particularly in China and Japan, the valves of some of the genus *Placuna*, which are extremely thin and semi-transparent, are used instead of glass, to serve as windows. Lime may be burned from a great many varieties, and is so procured, in America and elsewhere, particularly in Holland, where many vessels are employed in fishing for shells, which are afterwards brought to Haarlem, where the lime-kilns are situated. In the Hebrides, Antilles, and many other islands, no other lime is known than that procured from shells. Some shells are used as money in different countries, particularly the *Cypræa* ; and in conclusion, pearls and mother of pearl, both the productions of this class of animals, are well known valuable and beautiful objects.

To counterbalance in some measure these benefits which are derived by mankind from

Mollusca, there are some which are *injurious*, but to no very serious extent: snails and slugs do much damage in our gardens by their voracity; and many of the marine species, which make their residence in wood and stone beneath the surface of the sea, occasionally do us much damage, by boring into dikes, piers, and even ships.

Having thus given an outline of the nature of *Mollusca*, or, in other words, of *shells and their inhabitants*, in order that the reader might be able to form a general idea of them, it is now time to enter into their classification and arrangement into different orders and genera, so that he may have the power of becoming more particularly acquainted with them, and of deciding upon the nature, habits, and peculiarities of any shell, and the mollusc which has resided in it, by a simple inspection of its form and appearance. It is to be observed, that it is rarely indeed that the *animal* is to be met with in collections, the shell only being retained as an object worthy of preservation, and, consequently, he will have to be guided in his researches by the latter alone.

Mollusca are divided into two great *classes*, *Univalves* and *Bivalves*: the first comprising all those whose shells are formed of one single valve or piece, as the *snail*, and the latter, those which have shells consisting of two valves, or distinct pieces, as the *oyster*. Each of these classes is again divided into *orders*, each order into *families*, each family into *genera*, and each genus into *species*. These will be treated on in succession, according to a systematic arrangement, extending as far as the different *genera*: the limits of this work, which is intended but as an introduction to the science, will not allow of a description of each *species*, and a mere dry list of their names would be uninteresting. Reference can easily be made by the reader, who has mastered the subject as far as it will be pursued here, to the many large and scientific works which have appeared, in which each *specific* variety is introduced: that they would extend these remarks much beyond a short treatise, is evident from the circumstance that one genus alone, the *Helix*, or snail, comprises nearly 250 different species.

The classes *Annulata* and *Cirrhipeda* form

the ninth and tenth of Lamarck's system of animal nature, the *Conchifera*, or *Bivalves*, and the *Mollusca*, or *Univalves*, being the eleventh and twelfth. The two former will be described (though the animals are not considered as possessing true shells) on account of specimens of the tubes or coverings being frequently found in collections of shells. Lamarck arranges his system in a regular succession, beginning with the inferior animals, and describing them in order as they gradually increase in powers and complicated formation. On this account the *Annulata* and *Cirrhipeda* come first, the animals forming those coverings being very inferior in organization to those of the other classes.*

* The information contained in this chapter is chiefly derived from Dr. Roget's Bridgewater Treatise, Mr. Gray's article in the Philosophical Transactions for 1833, and various articles in the Penny Cyclopædia, &c.

PART I.

ANNULATA.*

LAMARCK has divided his class *Annelides* into three parts: as the animals of two of these do not form any covering to their bodies, they will not be described here; but the third division, namely, the *Sedentaria*, will be admitted, inasmuch as specimens of the tubes they form are often met with in collections of shells.

The animals of the tribe *Annulata sedentaria* are soft, elongated, and worm-shaped, inhabiting a tube which they never entirely quit. The body has either transverse segments or wrinkles; the head, eyes, and antennæ are sometimes deficient, and they have no feet, but are furnished with retractile knobs, in lateral rows. They are usually found attached to marine substances, and inhabit tubes, either membranous or horny, more or less encrusted

* From *annulus*, Lat., a ring.

with grains of sand or fragments of shells, or solid and calcareous: there are 4 families, and the greater part of them are carnivorous.

FAMILY 1.—*Dorsaliadæ*. *

The *branchiæ* of the animals of this family are disposed lengthwise on the body; there are 2 genera: —

1. ARENICOLA. This genus is found on the sandy shores of Europe, and is much used as a bait for fish. “The animal bores for itself a passage through the sand, and secures the sides of the passage from falling in by applying to them a glutinous cement, which unites the particles of sand into a kind of wall or coating. This covering does not adhere to the body, but forms a detached tube, within which the animal moves with perfect freedom, and which it leaves behind it as it progressively advances; so that the passage is kept pervious throughout its whole length, by means of the lining, which may be compared to the brickwork of

* *Les Dorsalées*, Lamarck.

the shaft of a mine or tunnel.”*—There is but 1 species.

2. SILIQUARIA. *Tube*, irregularly twisted, tapering towards one end, which is sometimes spiral; the other end open, and a longitudinal fissure throughout its whole length. *Animal* very long and spiral, head distinct, having two small tentacula, with eyes at the base. — 7 species. †

Some naturalists consider this animal to come under the class Mollusca, and not Annulata. M. Audouin has examined the animal of one recently found in India, and appears to be of the same opinion. It is found also in the Mediterranean, near Sicily.

* Dr. Roget's Bridgewater Treatise.

† So many additions are made almost daily to the number of *species* of Mollusca, that those which are mentioned under each genus must only be taken as the number of well established species known at a certain period. Almost all those which are mentioned in this work are described by Lamarck and Deshayes in *Les Animaux sans Vertèbres*.

FAMILY 2.—*Maldaniadæ*.*

The *branchiæ* which this family possesses are in the form of long filaments, which, as well as the head, are enveloped in the mantle. — There is only 1 genus.

3. CLYMENE. *Tube*, slender, open at both ends, the exterior encrusted with sand and pieces of shell. — There is only 1 species, which inhabits the shores of the Red Sea.

FAMILY 3.—*Amphitritidæ*.†

The *branchiæ* are disposed near the anterior part of the body; the tube is membranous or corneous, and is more or less covered with sand. — This family consists of 4 genera.

4. PECTENARIA. *Tube*, membranous or papyraceous, in the form of a reversed cone, the

* *Les Maldanées*, Lamarck. Lamarck placed here the genus *Dentalium*, which is now proved to be a mollusc, and will be described in its proper place.

† *Les Amphitritées*, Lamarck.

exterior covered with sand. — There are 2 recent species.

This animal inhabits the seas of Europe and India, and is not confined to one spot.

5. SABELLARIA.

Tubes, numerous, united into a common mass, composed of agglutinated fragments

of shell and sand; the orifices are cup-shaped. — There are 2 recent species.

It is found on rocks on the coasts of Europe.

6. TEREbella. *Tube*, elongated, cylindrical, thin, and pointed at the base, membranous, and having grains of sand, and fragments of

shell adhering all round it; it is open only at the apex. — There are 3 recent species. This animal inhabits the Northern Seas. Dr.

Roget* gives an interesting account of the manner in which these little creatures form their tubes:—"If taken out and placed in water upon sand, its actions may be observed. It immediately drags towards its head fragments of shells, and larger particles of sand, where they are cemented together by the glutinous matter which exudes from that part of the surface. It has soon formed a kind of collar, which it lengthens at every part of the circumference. Having formed a tube of half an inch or an inch in length, it proceeds to burrow, and as it proceeds avails itself of the materials, till the whole tube is completed. The *Terebella* then turns itself in its tube, so that the head is next to the surface, ready to receive the water which brings its food. In summer the whole task is completed in four or five hours, but in cold weather the progress is much slower."

7. AMPHITRITE.

—*Tube* cylindrical, membranous or co-

* Bridgewater Treatise.

riaceous, thin towards one end, generally with adhesions.—There are 8 recent species.

The Amphitrite inhabits the coasts of Europe, Jamaica, &c.

FAMILY 4.—*Serpulidæ*. *

The *branchiæ* in this family are separated or covered by an operculum. The tube is solid and calcareous.—The genera are 4.

8. SPIROBIS. — *Tube* testaceous, turned spirally on a horizontal plane, the lower part of which is attached to marine substances; the opening is round or angular.—16 species, recent and fossil.

This little spiral tube is frequently seen on the fuci found on the shore, and also on shells or other marine bodies. It is often found in great numbers, but each individual is isolated.

9. SERPULA. — *Tube* solid, calcareous, irregularly twisted, fixed, the aperture round and smooth.—60 species, recent and fossil.

* *Les Serpulées*, Lamarck.

These tubes are fixed by the lower extremity, or more completely by other parts, to marine substances, and are either grouped or solitary. The greater number of species are found in the seas of Europe. The animal feeds on aquatic animalculæ.

10. VERMILIA. —

Tube testaceous, cylindrical, gradually lessening at one end, more or less twisted, and fixed at the base

to marine substances. Aperture round, and having several teeth. — There are 8 recent species.

The tube of the *Vermilia* is said to have a convex, orbicular, and sometimes conical operculum. It is found on stones, shells, fuci, &c.

11. GALEOLARIA. — *Tube*

testaceous, cylindrical, rather angular, fixed at the base, with the other end raised. Aperture orbicular, and formed of from five to nine little valves on the upper side, the whole

of them attached to the edge of the operculum on one side only. — 1 recent and 1 fossil species.

This tubular animal is found in New Holland.*

PART II.

CIRRHIPEDA.†

THESE animals are soft, destitute of head or eyes, testaceous, having the body fixed, and furnished with a mantle; they have also tentacula, with curled tufts. The number of arms varies, and they are unequal in size; the shell is either sessile or elevated on a flexible pedicle, and is composed of several valves, sometimes moveable, sometimes fixed.‡ They are divided into two families.

* *Magillus*, a genus placed here by Lamarck, is now found to be of the class *Mollusca*, and is described in the fourth part of the work.

† From *cirrus*, Lat., *a curled tuft of hair*, and *pedes*, *feet*.

‡ “All these animals are oviparous. As soon as the young one comes forth, it attaches itself to the nearest body by secreting shelly matter.” — Reeve’s *Conchologia Systematica*.

FAMILY 1.—*Cirrhipeda sedentaria*.*

The body of the animal possesses no pedicle, and is enclosed in a shell fixed to marine substances.—There are 6 genera.

12. TUBICINELLA.—*Shell* univalve, operculated, tubular, erect, slightly thinner towards the base, having annular transverse ribs, truncated at both ends, open at the summit, and closed at the base with a membrane. Operculum with four obtuse valves. *Animal* rather cylindrical and long.—1 species only.

This shell is found buried to the depth of several inches in the fat of the South Sea whale; its operculum and the upper part of the tube being visible.

13. CORONULA.—*Shell* rather orbicular, conoid, the extremities truncated, the sides very thick; inside hollowed into radiating cells.

* *Cirripèdes Sessiles*, Lamarck.

Operculum composed of four obtuse valves. *Animal* rather flat. — 4 recent species.

This genus, like the last, fixes itself on the backs of whales, turtle, and other marine animals, and likewise on stones and large shells.

14. BALANUS. — *Shell* conical, summit truncated, closed at the base by a solid testaceous substance, by which it adheres. Aperture rather triangular. Operculum internal, and having four valves. *Animal* either flattened or almost cylindrical. — 39 species, recent and fossil.

Some of the species of this genus are armed with spines on the outside. They are found in all seas, covering the rocks, and pressing close together, so as in many instances to alter their form. In Chili they are found five inches and a half long, and half an inch in diameter, and are highly esteemed by the natives as an article of food. When boiled, and eaten cold, the flesh much resembles the crab, and is equal to it in richness and delicacy.

15. *ACASTA*. — *Shell* oval, rather conical, composed of pieces which can be separated. Cone formed of six unequal valves united together, the base an orbicular valve, concave on the inside, resembling a small cup. Operculum formed of four pieces. — 4 recent species.

These animals are found lodging in sponge, and are never attached to hard bodies. They inhabit tropical seas.

16. *CREUSA*. — *Shell* orbicular, having four valves, which are unequal and united together. Operculum internal, and consisting of four valves. *Animal* rather globular, and having three or four pair of arms. — 4 recent species.

These small shells are found affixed to madrepores, and other substances in the Northern and Indian Seas.

17. *PYRGONA*. — *Shell* rather globular, ventricose, convex above, apex perforated, aperture small, and

oval. Operculum consisting of four valves.*
Animal unknown. — 3 recent species.

Found in madrepores.

Some modern writers have introduced into this family several new genera, but as the distinctions are founded upon characteristics of secondary importance, they will not be described in these pages.

FAMILY 2. — *Cirrhipeda pedunculata*.†

The body of the animals of this family is supported by a tubular peduncle, flexible and tough, the base of which is affixed to marine substances. The shell is variable in form, but always composed of several valves.—There are 6 genera.

18. LITHOTRYA (Sowerby).—
Shell irregular in form, compressed and pyramidal, consisting of eight unequal pieces; having at the base an irregular cap-shaped appendage, like the inverted shell of a Patella, and to which the lower part of the

* Some authors say only two.

† *Cirripèdes Pédonculés*, Lamarck.

peduncle is affixed. *Animal* compressed. —
1 (?) species.

These animals are always found fixed to the bottom of a deep irregular cavity in limestone, perforated by themselves.*

19. ANATIFA.†—*Shell* compressed at the sides, composed of five valves, which are con-

tiguous and unequal; the two side-pieces are rather striated, the others small, and all are united together by a thin membrane. *Animal*

* Reeve's *Conchologia Systematica*.

† Called by Leach *Pentelasmis*.

very flat, enclosed in a thin mantle; peduncle often very long. — 10 recent species.

These curious but common shells are found in all seas, particularly on the coasts of Africa. They are affixed to marine bodies, generally in numerous groups, and the peduncle is sometimes found a foot long. The inhabitants of Goree eat a large species of *Anatifa*, which is considered a great delicacy. Some species live at a distance from each other, others are often united in large masses; they prefer a situation where the sea is much agitated.

“ The *Anatifa*, as well as the other genera of this tribe, are found attached to ships, logs of wood, bottles, corks of nets, fuci, floating testaceous Mollusca, (the *Iantha*, for instance,) and even on whales, turtles, and serpents. A large log of timber covered with these animals, twisting and diverging in all directions, and so thick as entirely to hide the surface of the log, is a strange sight. They look like an enormous collection of serpents to the ignorant; and a living mass of this description, casually thrown into shallow water, and left by the tide, has been so termed. Their growth must

be exceeding rapid. A ship going out with a perfectly clean bottom, will often return, after a short voyage, covered with them.”*

20. SCALPELLUM (Leach). — *Shell* composed of thirteen unequal pieces, twelve of which are lateral and very much flattened or compressed, and one of which is dorsal, elongated, and acuminate at the base. They are all united together on the integument, and the peduncle is covered with small scales. Animal resembling the *Anatifa*. — 2 species.

21. POLLICIPES. — *Shell* compressed at the sides, valves rather contiguous, unequal, thirteen or more in number, the lower side valves the smallest; peduncle short and wrinkled. *Animal* like the *Anatifa*. — 10 species, recent and fossil.

Found in the Northern and European Seas, in groups upon rocks.

22. CINERAS. — *Shell* composed of five testaceous oblong valves, separate, and not cover-

* Penny Cyclopædia, art. *Cirrhipeda*.

ing the whole of the body, two at the sides of the aperture, the others on the back. They are enclosed in a membranous bag, a continuation of which forms the peduncle. *Animal* rather compressed, peduncle thick.— 1 recent species.

This genus is found near the coasts of Senegal, and also in the Northern Seas. It attaches itself to buildings covered by the sea.

23. OTION. *Shell* composed of five small testaceous valves, adhering near the sides of the aperture, enclosed in a membranous bag, but terminating in two tubes at the back; the centre aperture admits the passage of the tentacula.— 4 species.

Found in similar situations to the last genus.

Animal. Soft, without either head or eyes, and having the mouth concealed. A large mantle envelopes the whole of the body, which is entirely or partially covered by a bivalve shell.

CONCHIFERA. — BIVALVES.

DIVIDED INTO THREE ORDERS.

Shell always bivalve, the valves mostly joined together by a hinge or ligament. Sometimes free, sometimes fixed.

1. BRACHIOPODA.	2. DIMYARIA.	3. MONOMYARIA.
<p><i>Animal</i>, depressed, circular, or oval. <i>Shell</i> having no ligament to unite the valves, but a pedicle by which the shell is united to other substances.</p>	<p><i>Animal</i> attached to its shell by <i>two</i> muscles, which are at a distance from each other, and near the sides of the shell. <i>Shell</i> having <i>two</i> muscular impressions in each valve.</p>	<p><i>Animal</i> attached to its shell by <i>one</i> muscle. <i>Shell</i> having <i>one</i> muscular impression in each valve; near the centre.</p>
<p>FOUR DIVISIONS.</p>		
<p>1st Division, CRASSIPEDA. <i>Shell gaping.</i> 4 Families.</p>		
<p>1. TENOINOSA. (<i>Shell</i> attached by a pedicle.) <i>Lingula, Terebratula.</i></p> <p>2. ADHÆRENTIA. (<i>Shell</i> attached by one valve, as well as a pedicle.) <i>Thecidium, Crania, Orbicula.</i></p>	<p>1. TUBICOLIÆ. (<i>Shell</i>: valves either enclosed within a testaceous tube; encrusted entirely or partially in its substance; or projecting outwardly. Marine, and boring into wood, stone, and sand.) <i>Aspergillum, Clavagella, Fistulana, Gastrochæna, Teredo.</i></p> <p>2. PHOLADIDÆ. (<i>Shell</i> without a tubular sheath, but provided with accessory pieces, distinct from the valves. Marine, boring into stone, &c.) <i>Xylophaga, Pholas.</i></p> <p>3. SOLENIÆ. (<i>Shell</i> transversely long, gaping only at the ends, ligament external. Marine.) <i>Solen, Solecurtus, Pholadomya, Panopea, Glycimeris, Solemya.</i></p> <p>4. MYIÆ. (<i>Shell</i> having a large spoon-shaped tooth in one or both valves, to which the ligament is attached, gaping at one or both ends. Marine.) <i>Mya, Anatina, Thracia, Corbula, Pandora, Anatinella.</i></p>	<p>1st Division. <i>Ligament marginal and elongated.</i> 3 Families.</p> <p>1. TRIOACNIDÆ. (<i>Shell</i> transverse, equivalve, outer margin waved or lobed. Marine.) <i>Triacna, Hippopus.</i></p> <p>2. MYTILIDÆ. (<i>Shell</i> longitudinal, equivalve, rarely foliaceous. Marine.) <i>Modiola, Lithodomus, Mytilus, Pinna.</i></p> <p>3. MALLEIDÆ. (<i>Shell</i> foliaceous, interior iridescent. Marine.) <i>Crenatula, Perna, Malleus, Vulsella, Avicula.</i></p>
<p>2d Division, TENUPEOA. <i>Shell gaping slightly.</i> 3 Families.</p>		
<p>5. MACTRIÆ. (<i>Shell</i> most frequently gaped at the sides; ligament internal; valves equal. Marine.) <i>Lutraria, Mactra, Crassatella, Mesodesma, Erycina, Ungulina, Amphidesma.</i></p> <p>6. LITHOPHAGIÆ. (<i>Shell</i> more or less gaping on one side; ligament external. Marine.) <i>Saxicava, Petricola, Venerupis.</i></p> <p>7. NYMPHIDÆ. (<i>Shell</i> with at most two cardinal teeth in the same valve, gaping often slightly; ligament external; callosities at the hinge projecting. Litoral.) <i>Sanguinolitaria, Psammobia, Galeomma, Tellina, Corbis, Lucina, Donax, Capsa.</i></p>		
<p>3d Division, LAMELLIPEOA. <i>Shell closed.</i> 4 Families.</p>		
<p>8. CONCHIOÆ. (<i>Shell</i> having at least three cardinal teeth in one valve; as many or less in the other; sometimes lateral teeth. First three genera fluviatile; the rest marine.) <i>Cyclas, Cyrena, Galathea, Cyprina, Astarte, Cytherea, Venus.</i></p> <p>9. CARDIOÆ. (<i>Shell</i> having irregular cardinal teeth, and generally one or two lateral; usually ribbed; heart-shaped; some slightly gaping. Marine.) <i>Cardium, Cardita, Cypricardia, Isocardia.</i></p> <p>10. ARCIDÆ. (<i>Shell</i> with small cardinal teeth, numerous, and arranged either in a straight or arched line; ligament external. Marine.) <i>Cuculea, Arca, Pectunculus, Nucula, Trigonia.</i></p> <p>11. NARIDÆ. (<i>Shell</i> generally no teeth, or with an irregular, simple, or divided cardinal tooth, and a longitudinal one; or a row of irregular tubercles; ligament external. Fluviatile, with an epidermis.) <i>Unio, Hyria, Anodonta, Iridina.</i></p>		
<p>4th Division, ANBIGUA. <i>Shell closed.</i> 1 Family.</p>		
<p>12. CHAMIDÆ. (<i>Shell</i> inequivalve, affixed to other bodies; one large tooth or none. Marine or fluviatile.) <i>Diceras, Chama, Etheria.</i></p>		
<p>2d Division. <i>Ligament short, and placed under the Boss.</i> 2 Families.</p> <p>4. PECTENIÆ. (<i>Shell</i> of a solid substance, generally eared, and ribbed. Marine.) <i>Pecten, Lima, Pecten, Plicatula, Spondylus.</i></p> <p>5. OSTREIDÆ. (<i>Shell</i> mostly foliaceous, inequivalve, rarely eared. Marine.) <i>Ostrea, Placuna, Placunanomia, Anomia.</i></p>		

[To face page 45.]

Animal. Soft, without either head or eyes, and having the mouth concealed. A large mantle envelops the whole of the body, which is entirely or partially covered by a bivalve shell.

1. BRACHIOPODA.

Animal, depressed, circular, or oval.

Shell having no ligament to unite the valves, but a pedicle by which the shell is united to other substances.

Animal shell

sides lar

Shell have,

TWO FAMILIES.

1. TENDINOSA. (*Shell* attached by a pedicle.) *Lingula*, *Terebratula*.

2. ADHÆRENTIA. (*Shell* attached by one valve, as well as a pedicle.) *Thecidium*, *Crania*, *Orbicula*.

ent

1. Tur ed.

pa

sa bell

2. Phove,

th or

3. Sol ri-

So

4. Myon-

me,

Pla-

ho-

na.

5. Matia-

ries-

6. Litre-

Sdal-

7. Nyda.

lig

ment

der

8. Cor of

otge-

Cind

9. Cap-

rien,

Iss.

10. Arstly

ative,

11. Nae.)

atta-

au.

12. Ch
fl

PART III.

CONCHIFERA OR BIVALVES.

THE Mollusca inhabiting bivalve shells are chiefly distinguished from the other classes by the absence of a visible head or neck, and the consequent deprivation of the organs of sight and hearing; they possess a mouth, but it is a mere opening in the body, without jaws, teeth, or hard parts. The branchiæ are large, placed on each side, between the body and the mantle. The lobes of the mantle are fringed round the edge with numerous filaments, very sensitive, and irritable to the touch, and which are in constant activity, drawing a current of water within for the capture of their prey, which is supposed to consist of small particles of vegetable and animal substances. None of the genera are terrestrial, their construction not affording them sufficient powers of locomotion for finding their food on land, and confining them to the water, whether salt or fresh, or to the sands on the coasts.

Bivalves are separated into three orders: —

teeth, D D. E E are the *muscular impressions*, and F the *pallial impression*, which marks the size of the animal, being formed by the border of the pallium or mantle. G G are the *sides* of the shell, which is called an *equilateral* shell when they are equal in length, and *inequilateral* when they are unequal. H is the *ligament*, which is horny, and by its elasticity tends to keep open the valves. This is counteracted at the animal's pleasure, by the muscles attached within the shell. The ligament is sometimes internal, and sometimes external, or both at the same time.

In describing a bivalve shell, both Linnæus and Lamarck place the shell on the bosses, with the opening above and the ligament before; the valves are either called right or left, as they are nearest to the right hand or left hand of the observer.

ORDER I.

BRACHIOPODA.*

Authors have been much divided in opinion about the order *Brachiopoda*, not only as to its

* *Les Brachiopodes*, Lamarck.

divisions, but regarding its right place in the system. Some have placed these shells with the *Cirrhipeda*. Cuvier made them a separate class, but still arranged them next in order. Lamarck places them at the end of the *Monomyaria*, but merely as a division of that order. Deshayes thinks they should form a distinct order, and come before the *Dimyaria*: this plan will be followed here, but the number of genera will be the same as found in Reeve's *Conchologia Systematica*; the plan there pursued appearing more simple than that adopted or explained by other conchologists.

This order is divided into 2 families, distinguished by the mode of attachment.

FAMILY 1.—*Tendinosa*.*

The muscle or pedicle by which these animals are attached to rocks, &c. gives them but a limited sphere of action.—There are 2 genera.

1. LINGULA.—*Shell* thin, either horny or calcareous, equivalve, equilateral, somewhat

* From *tendo*, Lat., a tendon.

peaked at the apex, generally open at the base, rather acute, somewhat gaping. The valves are destitute of a hinge, being supported by their muscular adhesion. *Animal* having two long ciliated arms, curled up during repose. — 3 recent and 1 fossil species.

Mr. Cumming met with considerable numbers of these shells in the Philippine Islands: their inhabitants are often eaten by the natives, but he does not think them wholesome. They live buried in hard sand, and can only be found at low water.

2. TEREBRATULA. — *Shell* inequivalve, equilateral, oval or orbicular, adhering by a short gelatinous tendon. The upper valve has the apex often curved inwards, and perforated for the passage of the tendon. It has also a single tooth-like process on each side. The lower valve has two slender shelly processes, generally recurved and bent. *Animal* having a ciliated appendage

on each side the body.—16 recent and 98 fossil species.

These animals inhabit great depths, and fix themselves to other objects by filaments or a tendinous cord which proceeds from the hole in the boss. They have a curious kind of internal skeleton, as it may be termed, consisting of a slender, flattened, calcareous loop, with other pieces diverging from it: the arms of the animal seem to be attached to it. They are found in the European seas, those of N. Zealand, and the Southern Ocean.

FAMILY 2.—*Adhærentia*.*

This family attach themselves by the shell; there is still, however, a muscular tendon in most of them, passing through a perforation in the lower valve. The valves are not united by any ligament. The shell is thin, and rather horny than shelly.—There are 3 genera.

3. *THECIDIUM* (Defrance).—*Shell* ovate, small, inequivalve, nearly equilateral, imperforate, and attached by the concave valve, which

* From *adhæreo*, Lat., to adhere.

has two short cardinal processes, and a rather irregular space, extending from the hinge to the boss. The upper valve is rather flat, and has a small obtuse appendage near the base, and two processes closely locking into those of the lower valve.

Animal not known.—1 recent and 3 fossil species.

From the Mediterranean. The interior of the upper valve is very curious, being indented as if to fit certain parts of the body. These marks spread round in a semicircular direction from the hinge, and look exactly as if they were picked out in wax. The inside of both valves is of a very pale green colour, and finely granulated.

4. CRANIA.—*Shell* compressed, somewhat square, inequivalve, equilateral, the upper valve convex and patella-shaped, the lower flat and attached. The interior of the latter exhibits four very distinct muscular impressions. There are no car-

dinal processes nor any hinge. — 1 recent and 4 fossil species.

From the Indian seas. (Lamarck.)

5. ORBICULA. — *Shell* horny, orbicular, rather irregular, inequivalve, rather compressed, the upper valve like a patella, the lower flat. In the centre of the latter is a small oval depression, with an oblique fissure in it for the passage of a tendon; four muscular impressions in each valve. There is no hinge. *Animal* having two short ciliated arms. — 4 recent and 2 fossil species.

These shells are found in the Northern Seas, and those of Peru and Chili. The species from Peru is very abundant, the individuals are grouped one on the other, and form large masses.

ORDER II.

DIMYARIA.*

This order contains a great number of families, which may be grouped into four divisions, arising partly out of the more or less perfect manner in which the valves close upon each other, partly from the shape of the foot of the inhabitant. In the *first* division, the valves when shut do not come into contact in all parts of the circumference, but leave a space open at one or both ends; the foot of the animal, also, is large and thick. In the *second* the aperture left is much less, and the foot is small and compressed. In the *third* the contact of every part of the valves is more perfect, and the shells can most of them be completely closed, and the foot is flat or lamelliform. The *fourth* division contains shells of a doubtful nature: the muscular impressions are two, but the shells are fixed to rocks, so that they greatly resemble in that respect the order Monomyaria.

* Called sometimes *Bimusculosa*; the Latin form of expressing the sense of *Dimyaria*. *Les Conchifères dimyaires*, Lamarck.

DIVISION I.—CONCHIFERA CRASSIPEDA.*

Shell gaping.

This division contains four families, *Tubicolidæ*, *Pholadidæ*, *Solenidæ*, and *Myidæ*.

FAMILY 1.—*Tubicolidæ*.†

These singular shells have at first sight no appearance of being bivalves, the two valves being small, and enclosed in a testaceous tube, or encrusted entirely or partially in its substance; they even sometimes project outwardly. They bury themselves in stone, wood, sand, and in other shells. — There are 6 genera.

6. ASPERGILLUM. — *Shell*, tube resembling the spout of a watering-pot, being thicker at one end than the other, the smaller one open, and the larger one closed by a plate, which is perforated with small holes; valves situated near the plate, outside the tube. *Animal* unknown.‡ — 5 species, recent.

* *Conchifères crassipèdes*, Lamarck. *Crassipeda*, from *crassus*, Lat., *thick*, and *pedes*, Lat., *feet*.

† *Les Tubicolées*, Lamarck.

‡ M. Ruppel speaks of the animal of the species *vaginiferum*, from the Red Sea. It appears greatly to resemble that of the *Pholas*.

This very curious shell derives its name from its form, and varies in length from seven or eight inches to a foot; it is supposed by some authors that it attaches itself to the rocks by a byssus passing through the holes in the plate which terminates its tube, but it has generally been found on the sand: its colour is white, with a slight tinge of grey or pale red.

The *Aspergillum agglutinans*, from New Holland, has the disk or plate differently constituted from the other species: instead of the small holes are substituted slender tubes, separate and unequal, with a fissure in the centre.

Its specific name is derived from the circumstance of the tube being constantly covered with sand, fragments of shells, and madrepores. The *Aspergillum Javanum*, from New Zealand and Java, is the most common species.

“ The row of slender tubes round the disk

are supposed to be filled with fleshy filaments from the mantle. The upper end of the tube, which is rarely obtained perfect, is more or less attenuated, and terminating in smooth undulations, is reflected at the margins. This reflected extremity is left entire at certain periods of growth, and the tube is recommenced according to the necessities of the animal. Some specimens have been found with four or five of these entire reflections.”*

7. CLAVAGELLA.—*Shell*, valves very small, the one being placed on the exterior, the other in the interior of the tube, which is narrow and open at one end, club-shaped and spiny at the other. *Animal* unknown. — 2 species recent, and 5 fossil.

There are very few recent specimens of this shell to be met with in cabinets: some have been brought from the Australian and Polynesian islands, and one living individual from the bay of Guayaquil. It perforates stones, corals, and even other shells.

* Reeve's Con. Sys.

8. FISTULANA. — *Shell*, valves equal, free, and situated inside the tube, which is swollen and closed at one end, but thin and open at the other. *Animal* not well known. — 4 species recent, and 2 fossil.

This genus is found buried in wood or fruits which have been long in the sea, in the thick substance of other shells, and also in the sand. Its locality is the shores of India.

Lamarck here places two genera, *Septaria* and *Teredina*, which modern naturalists think unnecessary. Deshayes unites them to *Teredo*.

Reeve, in his *Conchologia Systematica*, arranges *Gastrochæna* in this family, as it lines with a testaceous tube the cavity in which it dwells. It also forms an unconnected tube for protection when not imbedded.

9. GASTROCHÆNA. — *Shell* equivalve, inequilateral, gaping very widely on one side,

united by a ligament, and having in the interior a small spoon-shaped curvature. The tube differs from that of *Fistulana* in being nearly divided into two. *Animal* having two tubes and a thick short foot.—3 recent species.

The *Gastrochæna* penetrates and makes its abode in hard substances: it seldom exceeds half an inch in length, and inhabits the coasts of Great Britain, the Isle of France, and America. *G. modiolina* is the English species.

10. TEREDO.—*Shell*, valves gaping, having a long recurved tooth under the boss of each, external and attached to the animal, not to the tube, which is long, irregular, and open at both ends. *Animal* long, worm-shaped, having two little tubes.—2 species, recent.

The *Teredo* is almost the only mollusc which is to any extent injurious to mankind, and this

is owing to the animal forming its abode in wood which it meets with in the sea, such as dead and floating trees, piles, and even ships: it is worm-shaped, and about six inches long. In making its excavations into the wood, which it does by boring into the substance in the direction of the grain, each individual is careful to avoid the tube formed by its neighbour, and often a very thin leaf alone of wood is left between; it also, when a knot occurs in its path, makes a turn to avoid it.

A great proof of the manner in which the operations of this little animal may become prejudicial to us was furnished some time ago (1730) in Holland. In that country a considerable portion of the land and coast is below the level of high water, and in order to prevent the irruption of the sea, immense dikes have been constructed along the shore, formed of large masses of sand, and strengthened by large piles driven into the ground: these piles were at-

tacked by the *Teredo*, and, on examination, were found to be pierced in all directions to such an extent, that, had it not been for a timely discovery of the mischief, the whole of that part of the country might have been overwhelmed by the sea breaking through the enfeebled and worm-eaten defences.*

Found in the European and Indian seas.

FAMILY 2. — *Pholadidæ*.†

In this family there is no shelly tube like that which characterises the *Tubicolidæ*, and the animal is contained entirely within the two valves. The distinguishing mark is that the latter gape widely, and have small shelly pieces to unite the hinge, instead of a ligament. From this cause the two valves are scarcely ever found together, except when the shell is taken with the animal alive, as, after the death of the mollusc, the body is soon washed out of the open shell, and the valves having nothing

* Sir Everard Home mentions a species called *T. gigantea*, which sometimes exceeds four feet in length and several inches in circumference.

† *Les Pholadaires*, Lamarck.

to keep them together, separate and are thrown in different directions. There are two genera, *Xylophaga* and *Pholas*, both of which bore into stones, wood, and madrepores.

11. XYLOPHAGA (Turton). *Shell*, valves equal, inequilateral, and very much gaping, very similar to those of *Teredo*.

They have no calcareous tube, but two small accessory testaceous pieces placed near the hinge, and one small tooth in each valve, with an internal rib running from the boss to the margin.

These shells are found in light wood or pieces of stick, which the animals penetrate to the depth of about an inch.*

12. PHOLAS. — *Shell* gaping at both ends, equivalve and transverse, having a flat, recurved, spoon-shaped process in each valve below the bosses, which are close to one end of the shell, covered by a callosity, and also having accessory pieces attached. *Animal* having two small united tubes at one end of the shell,

* Reeve's Con. Sys.

and a foot or short thick muscle, which is flat at the end, at the other.—10 species recent, and 1 fossil.

These animals inhabit indurated mud or clay, wood, and stone, into which they bore when young, and in which they continue to reside until their death; it is not decidedly known in what manner these soft-bodied animals, inhabiting so fragile a tenement, are able to penetrate hard substances, unless it be by aid of a corrosive agent: they are, from their habits, exceedingly destructive to piers or wooden buildings under water. The breakwater at Plymouth, soon after its erection, was attacked by *Pholades*; it is constructed of limestone.* In order to counteract their operations, nails are closely driven into piles that lie under water, as is the case with the pier at Southampton.

The generality of the species are milky white, and elegantly marked with rows of small prickles: they are not used in this country as food, but on the shores of the Mediterranean, where some large species are

* Penny Cyclopædia, art. *Pholas*.

found, they are not uncommonly eaten. When the animal is irritated, it is said to throw out a stream of water to a considerable distance.

This genus inhabits the European, American, and Indian seas.

The animal is phosphorescent. This fact is mentioned by Pliny, as well as by modern authors.

FAMILY 3. — *Solenidæ*.*

The shells of this family are usually long, gaping at both ends, and having the valves united by an external ligament. The animals do not bore into wood or stone, but bury themselves in the sand. — There are 6 genera.

13. SOLEN. — *Shell* generally very long, straight, or arched, both ends open, and equi-

valve; teeth small, and varying in number; ligament exterior; bosses small. *Animal* having a short tube, containing two others united,

* *Les Solenacées*, Lamarck.

at one end of the shell, and a cylindrical foot at the other. — 20 species, recent.*

These animals bury themselves in the sand, in which they form holes perpendicular to the surface, occasionally to the depth of a couple of feet, and which they ascend and descend as occasion requires; they are sometimes used as food, but in places where they abound, they are sought after as bait for fish, and are captured in a singular manner. The retreat of the creature having been observed, the fisherman throws into the entrance a small quantity of salt, which quickly causes it to make its appearance, that substance in its pure state appearing to have a very irritating effect upon the animal, although an inhabitant of salt water; as soon as it is visible at the surface, the person lying in wait for it snatches hastily at it, and if he succeeds in grasping it firmly, makes good his capture; should he, however, fail, and the animal escape by gaining its retreat again, the application of more salt produces no farther effect, either from the Solen

* One species is found in the fossil as well as the recent state, and one in the fossil only.

not being sensible to the additional irritation, or perhaps from the instinct of self-preservation causing it to suffer the inconvenience in preference to running the risk of capture: in this case, the only method of securing it is to dig it out of the sand with a spade or other instrument.

Some species are common on the English coast; others come from America, India, &c. One of the Indian varieties is of a beautiful colour; the shell under the epidermis being of a delicate violet, striped with white.

14. SOLECURTUS (De Blainville). — *Shell* equivalve, ovately oblong, transverse, gaping at both ends, and having the external surface

covered with fine undulating lines. Two cardinal teeth in one valve, and one, rarely two, in the other, but not locking into one another

as the *Solen*. Ligament thick and external. Pallial impression very deeply sinuous. *Animal* larger than its shell; foot large and thick.— 4 species recent, and 2 fossil.

Found in the Mediterranean and Atlantic Ocean.

15. PHOLADOMIA (Sowerby). — *Shell* thin, transparent, white or yellowish, the inside pearly, transverse and oval, inequilateral, and gaping at both ends; hinge having a long narrow hollow or pit; ligament external; bosses worn by being placed near each other. *Animal* unknown.— 1 species recent, and 3 fossil.

There is only one species that is known to be in existence at present, the *Pholadomia candida*, which is found after hurricanes at the island of Tortola; but there are fossil species. The recent species is supposed to be a native of deep seas.

16. PANOPÆA.—
— *Shell* equivalve, transverse, broader and shorter than the

Solen, unequally gaping at the sides ; one conical tooth in each valve, with a thick callosity on the side; ligament external; pallial impression deep. *Animal* unknown. — 4 species recent, 2 of which are also fossil.

This is a large and handsome shell, but it is not often met with in collections in a perfect state, the valves seldom being found together after the death of the animal : the locality of the genus is the Mediterranean and New Holland.

17. GLYCIMERIS.—*Shell* transverse, possessing no teeth, but thick projecting callosities, and gaping at both ends ; ligament external, placed on the short side of the valves ; bosses generally eroded. *Animal*, foot small and cylindrical, projecting from one side, and two united tubes, which cannot enter the shell, issuing from the other. — 2 species recent, and 1 fossil.

Found in the northern seas, on sands. It is covered with a thick, black, horny epidermis, passing over the edge of each valve.

18. SOLEMYA.—*Shell* transverse, equivalve, inequilateral, and blunt at the ends; bosses small; one cardinal tooth in each valve, much compressed and oblique; ligament, both exterior and interior; slightly gaping, and covered by a shining epidermis, which extends beyond the margin. *Animal* having a trunk-shaped foot.—2 species, recent.

From New Holland and the Mediterranean, found in the sand.

FAMILY 4.—*Myidæ*.*

This family is distinguished by having a large spoon-shaped tooth on one or both valves, which are gaping.—There are 6 genera.

19. MYA.—*Shell* strong, thick, and transverse, both ends gaping, a large spoon-shaped tooth on the left valve, and a hollow in the other; ligament internal. *Animal*, a large tube containing two smaller ones proceeds from one end of the shell, and a very

* *Les Myaires*, Lamarck.

small foot from the other.—4 species, recent.

One end of this shell gapes very much, and at the aperture the thick epidermis is lengthened out into a tube, which can be withdrawn at pleasure by the animal. This mollusc buries itself in the sand, allowing the tube just to reach the surface, and by the aid of the smaller tubes contained within it procures its food. It is found on the shores of the European, Asiatic, and African seas, and is used as an article of food in several places; it is also devoured by marine birds.

20. ANATINA.—*Shell* transverse, thin, and almost transparent, gaping at one or both ends, having a spoon-shaped tooth in each valve, and a small moveable appendage connected with the interior of the hinge. The tooth in each valve is supported by a thin small plate; ligament internal. *Animal*, foot tongue-shaped.—10 species, recent.

These shells are found in India, New Hol-

land, Africa, and the European seas; the Indian species is rare, and very beautiful. They are found generally in sands and shallow water.

21. THRACIA (Leach).—*Shell* oval oblong, sub-equilateral, inequivalve, a little gaping at the ends; hinge consists of a more or less prominent spoon-shaped tooth in each valve receiving the ligament. Pallial impression deeply marked. *Animal*, mantle thick, foot rather small and compressed.—3 species, recent and fossil.

These shells are thin, fragile, and generally covered with an epidermis. They are found in the Mediterranean Sea and on the coast of Senegal.

The two following genera Lamarck had formed into a family and placed after his *Mac-tracées*; but Deshayes and other authors place them amongst the *Myidæ*, as they agree in all necessary particulars.

22. CORBULA.—*Shell* regular, inequivalve, and inequilateral, scarcely gaping; one cardinal spoon-shaped tooth in each valve, but no lateral; ligament interior. *Animal* unknown.—9 species recent, 1 of which is also fossil, and 10 fossil only.

These small shells are met with in the seas of New Holland, China, &c., and the European seas; one fine species is brought from South America; some of them are found in the interior of stones, &c., and M. Dorbigny states that some species live in fresh waters.

23. PANDORA.—*Shell* regular, oblong, valves unequal and inequilateral, the upper one flat, and the lower convex; an obtuse, oblong tooth in one valve, and a cavity in the other; ligament internal. *Animal*, foot large and triangular.—10 species recent, and 1 fossil.

The *Pandora* inhabits the sand at a considerable depth, and is found principally in the European seas; it is also met with in the Per-

sian Gulf and Pacific Ocean. The shells are small, and pearly inside: they bury themselves very deep in the sand, and it requires some exertion to get them out.

24. ANATINELLA (Sowerby).—*Shell* ovate, nearly equal, one end rounded, the other truncated or beaked; ligament internal, inserted between two spoon-shaped processes, one in each valve, and on one side of each are two cardinal teeth. *Animal* unknown.—1 species (?).

This shell is thin, fragile, and rather pearly inside.

DIVISION II. — CONCHIFERA TENUIPEDA.*

Shell nearly closed.

This division contains three families, *Macridæ*, *Lithophagidæ*, and *Nymphidæ*; the first family has the ligament internal, and the others external. The foot of the animal is small and

* *Conchifères tenuipèdes*, Lamarck. *Tenuipeda*, from *tenuis*, Lat., *slender*, and *pedes*, *feet*.

compressed, and the gaping of the shells in most instances very inconsiderable.

FAMILY 5.—*Mactridæ*.*

In this family the valves are equal, generally gaping slightly at the sides, and having the ligament placed in the interior. There are 7 genera.

25. LUTRARIA.—*Shell* equivalve, inequilateral, transverse; two cardinal teeth in one valve, and a triangular pit; no lateral teeth; a spoon-shaped lobe, containing the hinge. *Animal* similar to the *Mactra*.—13 species recent, some of which are also fossil, and 3 fossil only.

Found in the sand at the mouth of rivers in temperate climates.

26. MACTRA.—*Shell* oval, transverse, with thin cardinal and lateral teeth, one of which is shaped like the letter

* *Les Mactracées*, Lamarck.

V, and has the point near the boss; valves slightly inequilateral, and gaping a little on each side; bosses protuberant. *Animal*, foot sharp, oval, and long.—33 species recent, and 3 fossil.

The *Mactræ* live in the sand, and are found in all quarters of the world. This genus includes many rare and beautiful species; the shells exhibit rather a diversity of form, but are generally more or less triangular.

27. CRASSATELLA.—*Shell* equivalve, inequilateral, solid, and heavy, not gaping; hinge very thick; cardinal teeth, two in one valve, one in the other; lateral teeth scarcely seen; a triangular cartilage under the bosses; ligament internal; muscular impressions very distinct. *Animal* unknown.—11 species recent, and 14 fossil.*

Mature specimens of this shell are very

* Deshayes divides this genus into two, describing seven or eight species under the genus *Crassatella*, and the rest with others under a new genus which he names *Mesodesma*.

thick; their locality is New Holland and the southern seas.

28. MESODESMA (Deshayes). — *Shell* ovate, transverse or triangular, equivalve, rather compressed, and generally closed; the hinge has two thick lateral teeth in each valve, between

which is situated a deep spoon-shaped cavity for the ligament. *Animal*, foot very flat, quadrangular. — 10 species.

29. ERYCINA. — *Shell* transverse, small, and resembling the Crassatella, but having oblong lateral teeth, and rarely gaping; ligament small, placed in a triangular pit. *Animal* unknown. — 2 species recent, and 10 fossil.

This genus is found in the sand on the shores of New Holland and the Mediterranean.

30. UNGULINA.—*Shell* transverse or almost round; valves nearly equilateral, and not gaping; one bifid cardinal tooth in each valve, and at the side a long pit; no lateral teeth.

Animal unknown.—2 species, recent.

The exterior of this shell is rayed, and the interior of a reddish colour; it is found in Senegal, New Holland, and the Mediterranean, and the animal perforates rocks, &c. The shells are small, thin, and transparent.

31. AMPHIDESMA.—*Shell* round, or rather oblong, slightly gaping and inequilateral; two cardinal teeth and a long narrow pit between, often lateral teeth; ligament both external and internal. *Animal* unknown.—16 species, recent.

These shells, which are small, are found in the sand on the sea-coasts of tropical countries, and also those of England, Holland, &c.

FAMILY 6.—*Lithophagidæ*.*

This family comprises a small number of genera, which, like the *Tubicolidæ* and *Pholididæ*, inhabit rocks, stones, &c., into which they bore: they possess an external ligament, and have no tube or shelly pieces uniting the valves, the characteristic mark of the genera mentioned above. The holes they make are tolerably deep, as they conceal nearly the whole of their shell; the entrance is usually small. There are three genera.

32. SAXICAVA. — *Shell* transverse, inequilateral, thick, and solid, gaping, rather variable in form; teeth and bosses obsolete or indistinct. *Animal*, foot small, thin, and pointed. — 6 species recent, and 6 fossil.

The *Saxicava* perforates and inhabits rocks, oyster-shells, roots of sea-weed, &c., and is known in the South Seas, Mediterranean, Northern Ocean, &c. It appears greatly to

* *Les Lithophages*, Lamarck.

resemble the *Pholas* in its habits, and is found in calcareous rocks, which it is supposed to pierce by means of an acid. Masses of rock on the coast of England are frequently found pierced with innumerable small holes, the entrances to the habitations of these animals. M. Fleuriau de Bellevue, who has examined this family particularly, finds they are phosphorescent.

33. PETRICOLA. — *Shell* inequilateral, variously oblong, transverse, and equivalve, no epidermis; hinge furnished with two teeth on one valve,

and one or two on the other; slightly gaping. *Animal*, foot small and tongue-shaped. — 11 species recent, and 4 fossil.

Found in many parts of the world in rocks, corals, &c., but most abundant in America. The shell is rather delicate, white, and radiated.

34. VENERUPIS. — *Shell* transverse, inequilateral, striated, and slightly gaping; three cardinal teeth placed close to each other, ge-

nerally in each valve ; ligament exterior. *Animal*, foot compressed and long.—8 species recent, and 2 fossil.

The *Venerupis* excavates stones and corals : it is found on the coasts of England, France, the countries washed by the Mediterranean, &c. The colour is a dirty white.

By some authors this genus is not admitted. Sowerby says they do not bore into the rocks, but are merely cast into the cavities in which they dwell, by accident. He unites them to the *Veneridæ*, under the name of *Pullastra*.

FAMILY 7.—*Nymphidæ*.*

Many of the genera of this family comprise very beautiful shells, being smooth, polished, and tinged with shades of pink, orange, &c. They are usually oblong, and rather more flattened than the preceding families. The ligament is always external, and supported by a thickened hinge-plate, (called a *nympha*, or

* *Les Nymphacées*, Lamarck.

callosity) which often projects, and the valves either do not gape, or so slightly that it is almost imperceptible. There are never more than two cardinal teeth on the same valve, and only some of the genera have lateral teeth. These shells are all *litoral*, and bury themselves in the sand.—There are 8 genera.

35. SANGUINOLARIA. — *Shell* transverse, equivalve, inequilateral, slightly gaping at both ends; hinge having in each valve two teeth approaching each other.

Animal unknown.—4 species recent. From Jamaica, New Holland, the Indian and American seas.

36. PSAMMOBIA.—*Shell* transverse, oval,

slightly gaping at each end; hinge having two teeth on one valve, and one on the other;

bosses prominent, thin epidermis. *Animal* unknown.—18 species recent.

This genus presents a good deal of beauty, the shells being often ornamented with lively colours. There are species from the Indian Ocean, North Sea, New Holland, &c.

Here follows in Lamarck's system a genus called *Psammotæa*, which is now discarded: it seems to have been formed on imperfect shells.

37. GALEOMMA (Turton).—*Shell* thin, equivalve, inequilateral, gaping, one cardinal tooth in each valve; the ligament is small and duplex, partly internal and partly external. *Animal* unknown.—2 species.

Found in the Atlantic Ocean. A small white and thin shell, covered with waved longitudinal stripes.

38. TELLINA.—*Shell* nearly equivalve, transverse or orbicular, and greatly compressed; the valves on the ligamental side irregularly undulated; cardinal and lateral teeth two, the latter sometimes wanting; ligament on the shortest side. *Animal*, foot compressed and pointed.—62 species recent, and 16 fossil.

The *Tellinas* are all attractive from their beautiful colour and elegant shape. The ani-

mal has the power of springing to a considerable distance, by folding a portion of its body, and then suddenly extending it; the shell being closed at the same time with a loud snap. They bore into the sand on the sea-shore, and often become the prey of carnivorous shell-fish, as the *Buccinum*, &c. The valves are not always similar, one being occasionally more convex than the other, the colours and bands also differing. They are found in all parts of the world. The valves in some species are much more twisted than in others.

The genus *Tellinides* of Lamarck is now united with *Tellina*.

39. CORBIS.—*Shell* transverse, equivalve, nearly equilateral, bosses curved inwardly, cardinal teeth two, lateral two, one of which is remote. *Animal* unknown. — 1 species recent, and 2 fossil.

The shell of this genus is thick and solid: there is but one species in a recent state, and that is rare, from India.

This shell may be easily distinguished by the longitudinal and transverse undulations or ridges crossing the external surface of the valves. It is always of a white colour, and tinged with pink occasionally.

40. LUCINA.—*Shell* nearly round, inequilateral, bosses small and pointed, the outer surface sculptured, cardinal and lateral teeth distinct, but variable in number. *Animal*, foot long and cylindrical. — 19 species recent, and 15 fossil.

From Europe, America, New Holland, &c.

The interior of this shell is often punctured with small holes or *striæ*.

41. *DONAX*. — *Shell* equivalve, inequilateral, transverse, wedge-shaped, cardinal teeth two, on one or both valves, lateral variable, ligament on the shortest side, which is very much flattened or truncated, valves more or less crenulated at the margin. *Animal*, foot compressed, sharp, and angular. — 30 species recent, and 9 fossil.

The *Donax* is found in all parts of the world, buried in the sand of the sea shore; many of the species are beautiful.

42. *CAPSA*. — *Shell* transverse, somewhat triangular, equivalve but rather inequilateral, ligament exterior and on the shortest side, two cardinal teeth on one valve, and one on the

other, none lateral, margins not crenulated. *Animal*, foot compressed and very large. — 2 species recent.

India and Brazil produce this shell, which lies buried at small depths in the sand; it has an epidermis.

The genus *Crassina*, which follows *Capsa* in Lamarck's system, has been removed to the next family, under the name of *Astarte*, which had been previously given to it by Sowerby.

DIVISION III. — CONCHIFERA LAMELLIPEDA.*

Shell closed.

This division contains four families: *Conchidæ*, *Cardiidæ*, *Arcidæ*, and *Naiidæ*.

FAMILY 8. — *Conchidæ*.†

This family is often considered one of the most beautiful amongst the Bivalves. It is composed of shells having three cardinal teeth at least in one valve; but not always lateral

* *Conchifères lamellipèdes*, Lamarck. *Lamellipeda*, from *lamella*, a thin plate, and *pedes*, feet.

† *Les Conques*, Lamarck.

teeth. They are generally orbicular, and closed all round. Three of the genera are fluviatile, have lateral teeth, and are covered with an epidermis. The rest are marine, and have for the most part no lateral teeth or epidermis. There are seven genera.

43. CYCLAS.—*Shell* thin, roundish, or transverse, equivalve, and inequilateral, two small cardinal teeth in each valve, one of which is folded in two, but rather uncertain, lateral teeth lengthened transversely, and compressed, ligament exterior; an epidermis. *Animal*, foot large.—11 species recent.

The shells of this genus are not larger than a hazel-nut, and some of the species are very thin and transparent, striped transversely with light colours. They inhabit rivers in various parts of Europe and America, and have projecting, not eroded bosses.

44. CYRENA.—*Shell* thick, inequilateral, and round in form, cardinal teeth three in each valve, lateral generally two, ligament external, and on the longest side,

bosses eroded; an epidermis. *Animal* similar to the *Cyclas*. — 16 species recent, and 11 fossil.

Natives of warm climates, particularly of Asia: they attain a much greater size than the last genus, and have the interior of a beautiful violet colour, the outside being occasionally tinged with the same. They are fluviatile.

45. GALATHEA. — *Shell* equivalve, thick, and solid, somewhat triangular, nearly equilateral, and covered by a greenish horny epidermis, cardinal teeth two in one valve, and three in the other, lateral teeth at some distance, ligament exterior, projecting, and swollen. *Animal*, foot large and oblong. — 1 species recent.

Only one species is known of the *Galathea*, which inhabits the rivers of Africa, Ceylon, and India, and is very rare. Under the epidermis the shell is milk-white, stained with violet near the base, and marked with rays also of a violet colour.

46. CYPRINA. — *Shell* equivalve, inequilateral, obliquely heart-shaped, covered with a

thick brown epidermis, ligament external, three cardinal teeth, and generally one lateral tooth

in each valve, bosses turned on one side. *Animal*, foot large. — 1 species recent, and 6 fossil.

Chiefly fossil, with one recent species, *Cyprina Islandica*, which is found in the Northern Seas. These shells are often very large.

47. ASTARTE (Sowerby). — *Shell* equivalve, inequilateral, nearly orbicular, but somewhat transverse; two diverging teeth in the right valve, with one distinct, and one obsolete tooth in the left, covered with a brown epidermis

One of the muscular impressions is divided into two; ligament external. *Animal* like the other genera of this family. — 4 recent, and 12 fossil.

Some of the species are English, and are generally found on the sandy mud at very variable depths.

48. CYTHERIA. — *Shell* equivalve, inequilateral, triangular or transverse, ligament on the longest side, four cardinal teeth in one valve, and three in the other, lateral teeth none. *Animal*, foot large.
— 78 species recent, and 9 fossil.

These shells resemble much the *Venus* in their beauty and colouring, but the number of teeth slightly distinguishes them. They inhabit the Indian and Atlantic Oceans, and one species, the *Cytheria lusoria*, is found in the Chinese seas; the latter is employed by the Japanese and Chinese in certain games, and is painted by them in the interior of various colours, and representing various figures.

This genus resembles so greatly the next that some authors class them together: the

fourth cardinal tooth, which is supposed to distinguish them, is sometimes scarcely visible.

49. VENUS. — *Shell* equivalve, inequilateral, nearly round or oval, sometimes rough on the outside, and every where closely shut. Three cardinal teeth in each valve, sometimes lateral teeth, bosses slightly turned on one side, ligament on the longest side. *Animal*, foot variable in form, but compressed. — 96 species recent, and 13 fossil.

This genus, which is one of the most beautiful amongst the Dimyaria, is found in all parts of the world. The species *Venus mercenaria* is cut by the North American Indians into beads, of which they construct their wampum or treaty belts, and the shells are also used among them as money, and are made into ornaments for their dresses. Most species serve as food for man. These Mollusca lie buried in the sand, at a short distance from the shore, and are most abundant in hot climates.

Lamarck's genus *Venericardia* is now sup-

pressed, and the species forming it united to the genus *Cardita* in the following family.

FAMILY 9. — *Cardiidæ*.*

The *Cardiidæ*, or Cockle family, are the next in order; there are four genera. The cardinal teeth are irregular both in form and situation, and there are generally one or two lateral teeth. The shells are usually globular, and are almost all marked with raised ribs; they often present also the form of a heart, when they are seen on the side upon which the ligament is placed.

50. CARDIUM. — *Shell* heart-shaped, equi-valve, nearly equilateral, having generally radiated ribs, and the borders of the valves toothed or plaited, two cardinal and two lateral teeth in each valve, ligament exterior, bosses prominent. *Animal*, foot large. — 48 species recent, and 30 fossil.

Some of the species are very beautiful, and

* *Les Cardiacées*, Lamarck.

others very singular in form; amongst the latter is the species *Cardium cardissa*, in which the valves are flattened, but in a contrary manner to the generality of flat Bivalves. This genus is found almost every where lying buried in the sand near the sea-shore; by means of their long elbow-shaped foot, the animals have the power of leaping to a considerable distance. *Cardium edule* is the common species, and is frequently used as an article of food.

51. CARDITA. — *Shell* equivalve, oblong, inequilateral, having in one valve two unequal

teeth, one short and straight under the bosses, the other oblique, and in the other one thick, oblique, elongated tooth. Valves more or less strongly ribbed or smooth, waved at the edge. *Animal*, foot large and thick. — 25 species recent, 7 fossil.

Some species are said to attach themselves

by a byssus to rocks. They are brought from Amboyna, New Holland, Africa, and the Northern Seas.

52. CYPRICARDIA. — *Shell* oblong, finely radiated from the summit to the base, equivalve, very inequilateral, bosses very near one end, having three cardinal and one lateral tooth in each valve. *Animal* unknown. — 4 species recent, 7 fossil.

Found principally in masses of coral in St. Domingo, and on the coasts of Guinea and New Holland. It also bores into soft stones.

The genus *Hiatella* of Lamarck is no longer used. The only species he mentions is found to be a *Saxicava*.

53. ISOCARDIA. — *Shell* heart-shaped, equivalve, inequilateral; bosses very prominent, rather spiral, recurved in the same direction, and not touching each other; cardinal teeth two, lateral, one in each valve;

ligament exterior. *Animal*, foot rather large. — 3 species recent, 1 of which is also fossil, and 5 fossil only.

The bosses of this shell are so peculiar that they easily indicate the genus. The *Isocardia* lives usually in the sand within reach of the waves, and is found in Europe, the East and West Indies, and New Holland; the *Isocardia cor* is frequently met with in Dublin Bay. At Torbay the animal is fried by the fishermen, and the dish called “red noses.”

FAMILY 10.—*Arcidæ*.*

The next family is the *Arcidæ*, or Ark shells, which are distinguished from all the others by the great number of teeth; they have the appearance of those of a fine saw, and form a straight or curved continuous line, without any distinction of cardinal or lateral. There are five genera, the greater number of which bury themselves in sand near the coasts, all being at the same time marine, and some attaching themselves to the rocks by a byssus.

* *Les Arcacées*, Lamarck.

54. CUCULLÆA. — *Shell* equivalve, inequilateral; bosses not touching each other; muscular impression forming a partly detached shelly plate within; hinge straight, with a row of small teeth; ligament external. *Animal*, foot large. — 1 species recent, 6 fossil.

This shell greatly resembles the Arca: there is but one species, which is found in the sand in the Indian Ocean: its colour is a deep cinnamon brown externally, and brown tinged with violet internally.

55. ARCA. — *Shell* transverse, nearly equivalve, inequilateral, heart-shaped, valves ribbed,

and in some species gaping at the lower part; hinge straight; bosses prominent and distant, and often a little recurved; ligament external.

Animal, foot short and thick. — 40 species recent, 18 fossil.

Some of the species bury themselves in the sand, and others moor themselves by a byssus to stones, coral, &c.; a few have one valve larger than the other, and many have a velvety or shelly epidermis, frequently ending in a deep fringe at the margin. They are found in the Atlantic, Pacific, and Indian Oceans, the Mediterranean, &c.

56. PECTUNCULUS. — *Shell* nearly round, equivalve, almost equilateral; hinge curved, with a line of teeth diverging on each side, those in the middle being sometimes incompletely formed; ligament external, hairy epidermis. *Animal*, foot large. — 20 species recent, 15 fossil.

The *Pectunculus* is found in the West Indies, Atlantic Ocean, and Mediterranean, where it lives on the sandy or muddy coasts, and moves with the aid of its foot. Many of the species have a soft downy epidermis. No byssus.

57. NUCULA. — *Shell* varying in shape, equivalve, inequilateral; teeth small, numerous, and prominent, with a large spoon-shaped one in the middle; hinge curved or straight; bosses contiguous and curved; ligament partly internal; epidermis green, or dark olive brown. *Animal*, foot large but thin. — 10 species recent, 7 fossil.

The localities of this genus are the Baltic and Mediterranean, the Indian Seas, the English Channel, &c., where they are met with on the sand and mud, either on the open coast or at the mouth of rivers. The shell is generally pearly inside.

58. TRIGONIA. — *Shell* pearly; cardinal teeth

oblong, transversely furrowed, two in the right valve and four in the other; equivalve, inequilateral, triangular, ribbed or furrowed, ligament external. *Animal*, foot sharp.—1 species recent, 17 fossil.

There is only one species, which is found in the deep seas of New Holland. The interior is lined with highly iridescent mother of pearl.

Lamarck had originally placed this genus amongst the *Arcacées* (where Deshayes thinks it should be placed); he afterwards made it, with another genus *Castalia*, into a family; but it has since been ascertained that the *Castalia* belongs to the *Unionidæ*, and it is therefore here left as was originally intended.

FAMILY 11.—*Naiidæ*.*

In this family there are four genera: they are all inhabitants of fresh water, and particularly of American rivers, of a dark brown colour outside, caused by an epidermis, and very pearly within, with teeth not very distinctly visible.

* *Les Nayades*, Lamarck.

The shells of this family (like all others which are lined with what is called mother-of-pearl) are sometimes found to contain pearls of a good quality, and in large numbers, as many as sixteen having been taken from one shell. The Esk and Conway are celebrated for them. Sir Richard Wynn, chamberlain to Catherine, queen of Charles II., is said to have presented her majesty with a Conway pearl, which is to this day honoured with a place in the regal crown. Some of the Scotch rivers have produced numerous specimens of large and beautiful pearls, which have borne a high price. A paper in the *Philosophical Transactions for 1693* gives a curious account of the fishery for these pearls in the river Omagh, county Tyrone, Ireland. "The poor people," the writer says, "in the summer months go into the water, and some with their toes, some with wooden tongs, and some by putting a sharpened stick into the opening of the shell, take them up; and although, by a common estimate, not above one shell in a hundred may have a pearl, and of these pearls not above one in a hundred be tolerably clear, yet a vast number of fair mer-

chantable pearls are offered for sale by these people every summer assizes. Some gentlemen of the country make good advantage thereof, and myself heard that a miller took out a pearl which he sold for 4*l.* 10*s.* to a man that sold it for 10*l.*; the last purchaser sold it to the late Lady Glenlealy for 30*l.*, with whom I saw it in a necklace: she refused 80*l.* for it from the late Duchess of Ormond."

Between the years 1761 and 1764 the pearls found in the mussels of the river Tay were worth 10,000*l.* It is not uncommon now to find pearls which are sold for one or even two pounds.

59. UNIO.—*Shell* thick and solid, trans-

verse, equivalve, inequilateral; bosses prominent and eroded; cardinal teeth solid, short, and oblique; ligament external. *Animal*, foot

very large, thick, and of a square form.—83 species recent, and two fossil.

The *Unio* is found in the rivers of Europe, both the Indies, and elsewhere; it is called the *fresh-water mussel*. It buries itself in the sand, with the bosses undermost, and sometimes produces tolerably large pearls: the valves in some species are not quite closed, and they are often highly iridescent within, sometimes white, sometimes pink; the colour, however, though constant in some species, frequently varies in different individuals of the same.

60. HYRIA.—*Shell* equivalve, obliquely triangular; hinge margin straight, both extremities elevated and winged; cardinal tooth long, and divided into many parts, lateral tooth long; ligament external. *Animal* resembling that of the *Unio*.—2 species recent.

Found in exotic lakes, covered with an olive-coloured epidermis, and pearly within.

61. ANODONTA.—*Shell* generally thin, equi-

valve, inequilateral; hinge straight, with either no teeth or mere rudiments; shell transverse;

ligament external. *Animal* resembling that of the *Unio*.—15 species recent.

They lie buried in the sand or mud of lakes and ponds, and are found distributed in every quarter of the world; the valves are thin and large, pearly, and covered with an epidermis. From their shape and lightness they are used in France for skimming milk.

62. IRIDINA.—*Shell* equivalve, inequilateral; hinge very long; teeth small, like little tubercles, and numerous; very iridescent; ligament external. *Animal*, foot compressed and sharp.—1 species recent.

This genus, which only possesses one species, the *Iridina exotica*, is found in the rivers of warm climates, the Nile amongst others; the shell is rather thick, very brilliant inside, and of a red cast. Caillaud says that the natives of Egypt use it in preparing flax, and as spoons in measuring oil, butter, and different kinds of provisions.

DIVISION IV.—CONCHIFERA AMBIGUA.*

This division contains but one family, *Chamidæ*.

FAMILY 12.—*Chamidæ*.†

The *Chamidæ* form the last family of the *Dimyaria*: the shells are large, rough, irregular, and attach themselves by one of their valves to rocks, corals, &c., or even to each other in a curious manner. There are three genera.

63. DICERAS.—*Shell* inequivalve; bosses very much lengthened, distorted, and turned different ways; one large tooth in the largest valve. *Animal* unknown.—2 species fossil.

* *Les Conchifères ambigus*, Lamarck. *Ambigua*, from *ambiguus*, Lat., doubtful.

† *Les Camacées*, Lamarck.

These shells are only found in the fossil state, and are very irregular in form, and by most naturalists are now included in the next genus.

64. CHAMA. — *Shell* irregular, spiny, or rough, inequivalve; bosses recurved and un-

equal, turning either to the right or left; one thick and long tooth in one valve, and a deep pit in the other; ligament external. *Animal*: foot small. — 17 species recent, and 13 fossil.

These shells appear to present almost every variety of shape and colour; the former fact is perhaps owing to the accidents of their position. They are confined to warm climates, the Mediterranean being the lowest temperature. They are found usually at small depths in the sea, affixed by the largest valve to rocks, &c., and occasionally united together in large

masses. They are so firmly attached that it is difficult to avoid breaking the shell when they are removed from rocks.

65. *ETHERIA*. — *Shell* inequivalve, very irregular; teeth none; bosses short and indistinct;

ligament external, penetrating partly into the shell. *Animal*, foot large. — 5 species recent.

This genus is found in the rivers of Africa and Madagascar, and is abundant in the Nile above the Cataracts, where the inhabitants collect them for the purposes of ornamenting their tombs with the shells, and feeding upon the animal. They are pearly within, foliaceous on the outside, large, and fix themselves by the under valve often to one another, presenting the most curious distortions of form. The in-

terior of the valves is often of a vivid green colour, and raised in small blisters, which are said to be formed by the accidental introduction of small particles of sand during the formation of the nacreous fluid.

ORDER III.

MONOMYARIA.*

Nearly the whole of the genera arranged under this order and its families, attach themselves to rocks and other substances, not possessing the power of locomotion enjoyed by the *Dimyaria*, still less that degree with which the *Univalves* are endowed. They are more limited in number than the *Bivalves* possessing *two* muscular impressions, but there are among them several well known, and to us useful genera, as the *Oyster*, the *Mussel*, and the *Pearl Oyster*.

* Called sometimes *Unimusculosa*, the Latin form of expressing the sense of *Monomyaria*. *Les Conchifères monomyaires*, Lamarck.

DIVISION I. — *Ligament long and marginal.*

3 families, *Tridacnidæ*, *Mytilidæ*, *Malleidæ*.

FAMILY 1. — *Tridacnidæ*.*

The *Tridacnidæ*, or *Clams*, are the giants of the acephalous Mollusca, nearly all the species attaining a large size; they are transverse and equivalve, and attach themselves by a fibrous muscle to rocks and corals. They are generally found in exposed situations.—Only two genera.

66. TRIDACNA. — *Shell* regular, equivalve,

inequilateral, transverse; two compressed blunt teeth in each valve; a wide opening on one side of the bosses for the passage of the byssus; ligament external. *Animal*, foot cylindrical and thick.—7 species recent.

These shells are brought from the Indian Ocean, where they attain so large a size that a single person can with difficulty carry one

* *Les Tridacnées*, Lamarck.

valve, and some, indeed, have been said to weigh 500 lbs. In Roman Catholic countries, the shells are sometimes used as receptacles for the holy water in the churches, and formerly they were considered sufficiently valuable to form a present to a monarch, those in the church of St. Sulpice at Paris having been presented to Francis the First by the republic of Venice. The animal is correspondingly large, and it has been stated that more than one hundred persons can make a meal upon that which inhabits the *Tridacna gigas*: this may be an exaggeration, but there is no doubt the architect of so extensive a dwelling must be exceedingly large.* The *Tridacna* adheres to the rocks by a stout and strong byssus with so much tenacity as to require the aid of a hatchet to separate them. A specimen of this shell, brought from Sumatra, is now in Arno's Vale, Ireland, and weighs 507 lbs., each valve being four feet six inches in length, and two feet five and a half inches in breadth. The interior of the shell is of an opaque white, and very beautiful.

* The valves are so large as probably to require centuries to complete their growth. Reeve's *Conchologia Systematica*.

67. HIPPOPUS. — *Shell* equivalve, regular, inequilateral; valves closed, transverse, two

compressed teeth in each valve; ligament external; shell imbricated with numerous tubercles. *Animal* similar to that of the *Tridacna*. — 1 species recent.

The *Hippopus* has no opening in the valves, and is therefore presumed to be destitute of a byssus. It is not nearly so large as the *Tridacna*, and is a common shell in collections, although there is only one species, the *Hippopus maculatus*, from the Indian Ocean.

FAMILY 2. — *Mytilidæ*.*

The *Mytilidæ*, or *Mussel* family, possesses

* *Les Mytilacées*, Lamarck.

four genera; they have all more or less the shape of the common genus, and some are pearly in the inside. Their habits vary, some attaching themselves by a byssus to the rocks, whilst others perforate them; in all the ligament is external, and they have rarely any teeth.

68. MODIOLA.—*Shell* rather transverse, equi-valve, inequilateral; bosses prominent and slightly removed from the end; valves gaping a little; no teeth; ligament almost internal. *Animal*, foot tongue-shaped.—27 species recent, and 20 fossil.

Attached by a byssus to the rocks, of which they sometimes almost cover the surface. Sometimes found in holes of stones, madrepores, &c. They are met with in the Indian Seas, New Holland, the Atlantic, &c.

69. LITHODOMUS (Cuvier). — *Shell* trans-

verse, cylindrically oblong, equivalve, and covered with a brown epidermis; extremities round; bosses scarcely prominent. The hinge is straight; ligament mostly internal. *Animal*, foot tongue-shaped.

The *Lithodomus*, when young, suspends itself to rocks by a byssus*, but, as it grows, it pierces a hole, and introduces itself, forming a cavity which thenceforward it never leaves: indeed, after a short time, as it merely enlarges the interior, without making the entrance any wider, its increasing bulk renders it unable to quit its cell, and in these circumstances its byssus dies away, being no longer necessary. It is common in the Mediterranean and the Antilles, where it often pierces other shells.

70. MYTILUS. *Shell* longitudinal, equivalve, ham-shaped, somewhat triangular, bosses placed at the extreme end, hinge mostly without teeth; fastened by a byssus;

* This assertion is from Cuvier; Sowerby thinks it has no byssus.

ligament partly internal. *Animal*, foot thin and cylindrical. — 38 species recent, and 9 fossil.

The *Mytilus* or *Mussel* is found in most parts of the world, and is abundant on rocky coasts, where it is met with, attached by its byssus so as to be covered at high tide: an instance is known of a lobster having been found covered with them. When the shell of the common English species, the *Mytilus edulis*, is freed from its epidermis and polished, it is of a beautiful deep purple colour; the animal is much sought after as an article of food, and, although not equal to the oyster, makes a palatable dish; but care is necessary, as it is often unwholesome. Frequently towards the end of autumn, small crabs are found in the shell of the *Mytilus*, which live there sheltered from danger without hurting the inhabitant. Small seed pearls are sometimes found in this species. One or two species are found in fresh water, the Danube, the Wolga, and Rhine, the London Docks, &c. &c. A new generic name, *Dris-sena*, has been given to them by M. Vanbeneden, but it has not been universally adopted.

71. PINNA. *Shell* equivalve, rather trian-

gular, ham-shaped; bosses terminal, the other extremity gaping; no teeth; a long and some-

what internal ligament; shell thin and fragile. *Animal*, foot thin, worm-shaped.—16 species recent, and 4 fossil.

The *Pinna* is usually found moored to the sand by its byssus; it is common to nearly all seas, and grows to a considerable size, some specimens being a foot and a half long; one species is found in our own seas on the coasts of Dorsetshire and Sussex. Pearls are occasionally found in the shell of the *Pinna*, but they are of an amber colour and of little value; sometimes a small species of crab takes up its abode inside, without prejudice to either party. A species found in the Mediterranean is much sought after by the inhabitants of Sicily and Calabria, not only as an article of food, but also for the sake of its byssus, which is

long, fine, shining, and abundant: of this they fabricate a species of cloth, very remarkable for its warmth and softness, but which will not take any dye. In the British Museum there is a pair of gloves made of this material, in a case with some large specimens of the shell; and it is recorded in history, that Pope Benedict XV. had a pair of stockings presented to him in 1754 made of this material; they were enclosed in a very small box, and were extremely fine in texture.

FAMILY 3. — *Malleidæ*.*

The shells of this family are foliaceous†, and have the interior highly iridescent and pearly; indeed they comprise amongst their number the pearl oyster: there are five genera, and nearly all adhere by a byssus.

72. CRENATULA. — *Shell* nearly equivalve, irregular, foliated, flattened; hinge straight, marked with a few semicircular excavations;

* *Les Malléacées*, Lamarck.

† Composed of thin plates of shell, laid one upon the other, each successive lamina projecting beyond its predecessor.

ligament internal;
no distinct opening
for the byssus. *Animal*
unknown.—7
species, recent.

This shell, which
is thin and fragile, lodges in sponges, &c. It
is brought from India and Australia. The bys-
sus has not been seen, and therefore it is not
positively ascertained whether it possesses one.

73. PERNA.—*Shell* nearly equivalve and
flat, irregular, gaping
for the passage of the
byssus; bosses small;
margins very brittle;
hinge straight and
broad, divided into
parallel grooves. *Animal*,
foot very small.
— 10 species recent, 6 fossil.

From India, the Cape de Verde Islands, &c.,
generally found moored to rocks deep in the
sea, and occasionally to mangrove trees. Con-
siderable clusters are frequently found attached
firmly to each other by the byssus.

74. *MALLEUS*.— *Shell* irregular, variously distorted, nearly equivalve; hinge margin generally very long, extended out into two ears; no

teeth, but a conical pit under the boss, which is small; an oblique notch for the passage of the byssus, near the boss; ligament external. *Animal*, foot very distinct.—6 species recent.

The *Malleus* is a native of the East and West Indies and Australia: it attaches itself by a byssus to submarine rocks. The shape is so very various, that scarcely two of a species

can be found alike, and the rarity of some causes them to be very valuable, and much sought after. Externally the appearance is very rude and irregular, but, on the other hand, the interior is exceedingly beautiful, being lined with the most brilliant mother-of-pearl.

75. VULSELLA. — *Shell* longitudinal, tongue-shaped, nearly equivalve, irregular; bosses equal; hinge, with a prominent callosity in each valve, showing an impression of a conical and arched pit for the ligament, which is partly external. *Animal* unknown, without byssus. — 6 recent and 1 fossil species.

These shells are not attached, but are found buried in sponge. They are iridescent within, and are brought from the Indian Ocean and the seas of New Holland.

This genus was placed by Lamarck in the *Ostreidæ* family, but more modern naturalists place it here.

76. AVICULA. — *Shell* inequivalve, fragile ; hinge margin straight and lengthened ; une-

qually eared ; minute teeth ; a large byssus. *Animal*, foot conical, worm-shaped, and rather long. — 21 species recent, and 8 fossil.

There are several beautiful species of the *Avicula* from the Indian Ocean, coast of Brazil, New Holland, the Red Sea, &c., and one from the Mediterranean. The valves are much in the shape of wings. The interior part is pearly in the centre : some species have a broad black border surrounding it, and the margin terminated in a fringe, occasioned either by the epidermis or the foliaceous texture of the exterior. The hinge of the *A. Macroptera* is used for making ornaments (bijoux) called peacock's eyes.

The *Avicula* and *Meleagrina* of Lamarck are now classed together, as the animals resemble each other.

The *Avicula Margaritifera*, or Pearl Oyster, is found chiefly in the Persian Gulf, and at Ceylon; it is lined with a pearly substance, which is formed by the animal in thin layers, and is imported into Europe in large quantities, for the purpose of making knife handles,

buttons, &c., under the name of *mother-of-pearl*. If by any accident the inner surface of the shell is injured, so as to cause a fracture in the mother-of-pearl, the layers in that place become for the future irregular, and a lump is gradually formed, which from accidental circumstances is sometimes oval, sometimes globular, and sometimes pear-shaped. Pearls, as these lumps are called, are occasionally found loose in the shell, and in this case the pearly substance has been deposited upon a loose grain of sand, or diseased ovum or egg of the animal.

“ Pearls are small nacrous balls, that become formed and hardened within the body of the

animal; they are found deposited in the most fleshy parts, particularly within and around the adductor muscle, and are said to be occasioned by the overcharge of those glands whose function it is to secrete the nacrous fluid destined for the internal lining of the shell. When the animal is thus diseased, this beautiful iridescent fluid is very irregularly discharged, being also deposited upon the inner surface of the shell in little excrescences; these are often detached, and form articles of commerce as pearls of inferior value, the former being considered more precious, both on account of their rotundity of form and the clearness and beauty of their complexion.”*

Excellent imitations of pearls have been made with hollow glass globules, the inside of which is covered with a liquid called pearl essence, and then filled with white wax: the essence is composed of the silver-coloured particles which adhere to the scales of the Bleak fish, and was first applied to this purpose, by a Frenchman named Jacquin, early in the

* Reeve's Con. Sys.

last century. The accounts of the pearl fishery are exceedingly interesting.

DIVISION II. — *Ligament not marginal, contracted into a short space under the bosses. Always visible. The shells are all inequivalve.*

Two families, *Pectenidæ*, *Ostreidæ*.

FAMILY 4. — *Pectenidæ*. *

The genera of this family are all more or less fan-shaped, and marked with rays from the bosses to the circumference; the substance of the shell is solid, not foliaceous. Some are attached, others free.—There are five genera.

77. PEDUM. — *Shell* irregular, longitudinal,

* *Les Pectinides*, Lamarck.

or hatchet-shaped, inequivalve, and slightly eared. A groove or pit in each valve, which increases in size with age, and bears the ligament; bosses unequal and distant, the lower valve rather convex, with the sides reflected over, upper valve flattish. A large yellow byssus. — *Animal*, foot small. — 1 recent species.

The only species known is found in the Indian Seas, at great depths, and is rare. It is white, slightly tinged with purple near the bosses; and buries itself partially in madrepores, in crevices of its own boring.

78. LIMA. — *Shell* longitudinal, nearly equivalve, obliquely fan-shaped, slightly eared;

valves gaping near the bosses, which are distant; hinge without teeth; no byssus. A tri-

angular pit for the ligament.—*Animal*, foot cylindrical.—6 species recent, and 18 fossil.

The *Lima* is almost always white; some of the species are rare. It is found on the coasts of America and the Isle of France. It has a brown horny epidermis.

The genus *Plagiostoma* of Lamarck, which he placed after *Lima*, and which contained six species, is by Deshayes divided between *Lima* and *Spondylus*. He thinks the genus unnecessary.

79. PECTEN.—*Shell* regular, nearly round, eared, ribbed or furrowed, inequivalve; ligament internal and triangular; hinge margin straight; no teeth; bosses contiguous.—*Animal*, foot small, dilated at the extremity.—61 species recent*, and 58 entirely fossil.

The animal of the *Pecten* has the power of making frequent and sudden contractions of its

* Several of the recent species are also found fossil.

muscles, by which means it moves rapidly through the water; and it requires some agility to catch the shell, as it flutters amongst the corals where it dwells. Aristotle mentions their power of leaping; and this observation has since been confirmed. A basket full of the common *Pecten* placed at the edge of the water has been speedily emptied, by the individuals springing from their confinement to their native element. Some of the species are stout and heavy, others thin and transparent, and many are ornamented with beautiful colours.

The *Pecten* is found almost every where; on our south coasts it is abundant, and much used as an article of food. In some countries the shells of the *Pecten maximus* are used as plates by the poor; in Paris the restaurateurs employ them for the same purpose, when serving up a certain preparation of mushrooms; and in England they are employed in cooking scalloped oysters, and the shell is consequently known as the *scallop shell*. The *Pecten Jacobæus* was formerly the badge of the pilgrims who had been to the Holy Land, and was worn on their caps and cloaks.

80. *PLICATULA*. *Shell* irregularly ovate, inequivalve, one valve being more convex than the other, sides strongly and transversely grooved; ligament internal, between the teeth; bosses unequal. *Animal* unknown. — 5 species recent, and 10 fossil.

Attached by the under valve to stones, &c.

81. *SPONDYLUS*. — *Shell* eared, inequivalve, rough, with spines and plaits; two strong teeth, uniting the valves closely; bosses unequal, that of the lower valve projecting; ligament internal. *Animal*, foot very indistinct. — 21 species recent, and 9 fossil.

The *Spondylus* lives at great depths in the sea, and attaches itself to corals, &c.: it has often been found fixed to anchors, cannons, and other iron articles that have been for some time at the bottom of the sea. Many of the species are very beautiful, and of very vivid colours, such as bright red, pink and yellow, or orange. A fine collection of these beautiful shells forms

at once a union of elegance and beauty hardly to be surpassed amongst the bivalves. They are found in the Indian Ocean, the Mediterranean, and on the American coasts.

The fossil genus *Podopsis* of Lamarck, is joined by Deshayes to *Spondylus*, and also some species of *Plagiostoma*.

FAMILY 5. — *Ostreidæ*.*

This family comprehends the oyster tribe, and the genera forming it are usually affixed by one valve to stones or rocks; some of the shells are foliaceous.—There are four genera.

82. — OSTREA. — *Shell* inequivalve, irregular, flattened; substance foliaceous; bosses distant, and very unequal, that of the lower valve becoming very large with age, upper valve smallest; no teeth. — *Animal*, no foot.—53 species recent, and 82 fossil.

Oysters have been found in most parts of the world in abundance, although none have as yet been discovered in the polar regions: they

* *Les Ostracées*, Lamarck.

are inhabitants of salt water only, and are always found on rocky ground, at no great

depth from the shore, attached by the under valve. The oyster, and indeed all bivalves which adhere by the shell, are covered at their birth with a mucilaginous liquid which attaches them to the surface of any object on which they rest. The animal strengthens this first adhesion in the same manner that it increases the size of its shell. At the mouths of several American, African, and Indian rivers, great quantities of oysters are found attached to the roots of trees, and even to their branches, where they are so situated as to be covered by the tide. Mrs. Lee, in her "Stories of Strange Lands" says, "The flavour of the tree-oyster is

delicious; they are small, and cover the lower branches of the mangroves. Two or three of these branches form an ample luncheon, and in the river Gaboon we had a daily supply." In the British Museum, there is a large specimen of a crab, to the back and claws of which a number of good-sized oysters have attached themselves.

Oysters generally spawn in May, and their growth is tolerably rapid; three days after the deposition of the spawn, the shell of the young oyster may be seen nearly a quarter of an inch broad; in three months it is larger than a shilling, and in six months it exceeds a half-crown piece in size. Oysters have been used as food almost from time immemorial; the Greeks, but more especially the Romans, held them in great estimation, those from the Dardanelles, Venice, and England being considered the best. The English fishery is principally carried on at Wivenhoe, near Colchester, Feversham and Milton in Kent, in the Medway, and at Tenby in Wales. When the oyster is first taken out of the sea, it has frequently a strong muddy taste, and it is therefore placed in what is called

a preserve, which is a kind of tank dug in the sand near the sea, and communicating with it by a narrow passage; this place is frequently cleared of mud and dirt by letting the water run off and refilling it; by this means the quality of the oyster is much improved. They are not considered wholesome during May, June, July, and August, and are not eaten until they are one year and a half old.

The animal of the oyster appears to be one of the most inanimate of the Mollusca; it remains fixed upon a rock, or other submarine object, enjoying only the nourishment brought it by the waves, and giving scarcely a sign of life, except the opening and shutting of its valves.

The genus *Gryphæa* of Lamarck is by Deshayes and other conchologists incorporated with the genus *Ostrea*.

83. PLACUNA.—*Shell* nearly equivalve, irregular, very much compressed, iridescent; two internal cardinal teeth in one valve, shaped like a V., and in the other

valve impressions to receive them; shell thin, and nearly transparent. *Animal* very little known.—3 species recent, 1 fossil.

Found on sand in India, China, the Red Sea, &c. The *Placuna Placentia*, commonly known as the Chinese window oyster, is sufficiently transparent to be used for windows, lanterns, &c., as horn is used in this country.* Though the animal has not been much examined by naturalists, it is obvious that it must be exceedingly flat; indeed, not much thicker than a shilling; for the valves, when closed, appear nearly to touch. The shells are sometimes waved, and bent in a curious manner.

84. PLACUNANOMIA (Broderip).—*Shell* irregular, nearly equivalve, smooth, plaited round the edge, and somewhat glossy inside. The hinge consists of two narrow ribs, unequal in length, and converging at the base, in one valve,

* Mr. Gray mentions that the Chinese use the powder of this shell as silver in their water-colour drawings. He has himself used it in colouring the figures of fish with good effect. It is not quite so brilliant as the powdered leaf metal, but it has the advantage of not changing colour by exposure.

and in the other two corresponding grooves. The lower valve has a longitudinal hole near

the hinge, and the opening is entirely filled up by the ossification of the tendon passing through it. Besides the muscular impression in the upper valve, there is a mark which shows the attachment of the other extremity of the tendon of adhesion.

Mr. Cuming found one species of this singular shell attached to dead bivalve shells and coral, at the depth of eleven fathoms in the Gulf of Dulce, in Costa Rica, Central America. It is intermediate, as its name denotes, between *Placuna* and *Anomia*: the hinge is like the former, and the opening in the lower valve for

the passage of the tendon causes its resemblance to the latter.

85. ANOMIA. — *Shell* inequivalve, irregular, attached to rocks, &c. by a calcareous mass affixed to a muscle, which passes through a hole in the lower valve, the latter being small and nearly flat. *Animal*, foot rudimental. — 9 species recent, and 1 fossil.

From the Mediterranean and other European seas. It is found adhering to oysters and other substances, rocks, &c., and takes the irregular form of the surface to which it is attached; so that if fixed to a *Pecten*, it will be striated, &c. It cannot be detached without destroying the animal. The calcareous mass by which this shell is attached is so constructed as to close the hole in the valve when the muscle is contracted. The animal is phosphorescent. It is eaten in France.

Animal. Soft, furnished with a head, most frequently with eyes and tentacula, or possessing arms disposed in the form of a coronet. Body sometimes naked, sometimes enclosing a shell, but generally having an external shell.

MOLLUSCA. — UNIVALVES.

Shell. Composed of one piece only.

DIVIDED INTO FIVE ORDERS.

1. PTEROPODA.	2. GASTEROPODA.	3. TRACHELIPODA.		4. CEPHALOPODA.	5. HETEROPODA.
<p><i>Animal.</i> Having no feet, but two opposite fins, adapted to swimming.</p> <p><i>Shell.</i> Thin; cartilaginous, or horny.</p>	<p><i>Animal.</i> Foot united to nearly the whole length of the body, and used for crawling. Body straight, never spiral.</p> <p><i>Shell.</i> Either cup-shaped, or slightly rolled up; in some genera not covering the body.</p>	<p><i>Animal.</i> Foot attached to the neck, and used for crawling. Part of the body spiral, and always enveloped in a shell.</p> <p><i>Shell.</i> Spiral, and the mouth either entire at the base, notched, or channelled.</p>		<p><i>Animal.</i> Feet or arms attached to the head.</p> <p><i>Shell.</i> Either polythalamous or monothalamous; many genera very flat; spire in or near the centre.</p>	<p><i>Animal.</i> Feet or fins attached in various parts. <i>Shell</i> slightly spiral, and nearly transparent.</p>
		PHYTOPHAGA.	ZOOPHAGA.		
		<p><i>Animal.</i> Feeding on vegetable substances, and being either marine, fluviatile, or terrestrial.</p> <p><i>Shell.</i> Aperture entire.</p>	<p><i>Animal.</i> Feeding on animal substances only, and being marine.</p> <p><i>Shell.</i> Aperture notched or channelled.</p>		
		TEN FAMILIES.	FIVE FAMILIES.	THREE DIVISIONS.	
<p><i>Hyalea, Clito</i> (no shell), <i>Cleodora</i>, <i>Limacina</i>, <i>Cymbulia</i>, <i>Pneumodermon</i> (no shell):</p>	<p>EIGHT FAMILIES.</p> <p>1. TRITONIDÆ. (No shell.) <i>Glaucaus, Eolis, Tritonia, Scyllæa, Tethys, Doris.</i></p> <p>2. DENTALIOÆ. (<i>Shell</i> tubular, open at both ends.) <i>Dentalium.</i></p> <p>3. PHYLIDIDÆ. (<i>Shell</i> either of one cup-shaped piece, or with moveable pieces at the back.) <i>Phyllidea</i> (no shell), <i>Chitonellus, Chiton, Patella, Siphonaria.</i></p> <p>4. SEMIPHYLIDIDÆ. (<i>Shell</i> cup-shaped, but very flat.) <i>Pleurobranchus, Umbrella.</i></p> <p>5. CALYPTRÆIDÆ. (<i>Shell</i> cup-shaped, though irregularly so; some genera with internal pieces.) <i>Parmophorus, Emarginula, Fissurella, Crepidula, Calyptræa, Hypponix, Pilopsis.</i></p> <p>6. BULLIDÆ. (<i>Shell</i> slightly rolled up.) <i>Acera</i> (no shell), <i>Bulla, Bulla.</i></p> <p>7. APLYSIDÆ. (<i>Shell</i> nearly flat and thin.) <i>Aplysia, Dolabella.</i></p> <p>8. LIMACINÆ. (<i>Shell</i> very slightly rolled or cup-shaped.) <i>Onchidium</i> (no shell), <i>Limax, Parmacella, Testacella, Vitrina.</i></p>	<p>1. COLIMACIDÆ. (<i>Shell</i> spiral, no projecting parts. The right lip often recurved outwards, globular or turreted, and generally thin. Terrestrial.) <i>Helix, Caracolla, Anastoma, Helicina, Pupa, Clausilia, Bulimus, Achatina, Succinea, Auricula, Cyclostoma.</i></p> <p>2. LYMNÆIDÆ. (<i>Shell</i> spiral, thin, smooth; right lip always sharp, never reflected. Fluviatile.) <i>Planorbis, Phgsa, Lymnaea, Ancylus.</i></p> <p>3. MELANIDÆ. (<i>Shell</i> turreted, lips disunited; right lip sharp. Fluviatile.) <i>Melania, Melanopsis.</i></p> <p>4. PERISTOMIDÆ. (<i>Shell</i> conoid, or rather discoid, lips united; shell rather thin. Fluviatile.) <i>Valvata, Patadina, Ampullaria, Ampullacera.</i></p> <p>5. NERITIDÆ. (<i>Shell</i> semi-globular, or rather flat; left lip resembling a partition, like the deck of a boat. Fluviatile and marine.) <i>Navicella, Neritina, Nerita, Natica.</i></p> <p>6. JANTHINIDÆ. (<i>Shell</i> thin, of a violet colour; mouth triangular. Marine.) <i>Janthina.</i></p> <p>7. MACROSTOMIDÆ. (<i>Shell</i> ear-shaped, flat; mouth very large; lips disunited. Marine.) <i>Sigarticus, Stomatella, Stomatia, Haliotis.</i></p> <p>8. PLOCACIDÆ. (<i>Shell</i> with the left lip plaited, right sharp. Marine.) <i>Tornatella, Melampus, Pyramidella.</i></p> <p>9. SCALARIDÆ. (<i>Shell</i> with whorls sometimes not touching, lips circularly united. Marine.) <i>Vermetus, Magilus, Leptocochnus, Scalaria, Rissos, Delphinula.</i></p> <p>10. TURBINIDÆ. (<i>Shell</i> turreted, conical, or pyramidal, substance thicker than the Colimacidae; mouth round or oblong; lips not united. Marine.) <i>Solarium, Rotella, Trochus, Monodonta, Turbo, Planaxis, Phasianella, Turritella.</i></p>	<p>11. CANALIFERIDÆ. (1st Division. <i>Shell</i> with a canal, more or less long, at the base of the mouth; right lip not thickened. Marine.) <i>Cerithium, Pleurotoma, Turbinella, Cancelaria, Fasciolaria, Fusus, Pyrala.</i> (2d Division. Right lip thickened.) <i>Struthiolaria, Ranella, Murex, Triton.</i></p> <p>12. PTERIDÆ. (<i>Shell</i> having the right lip dilated in the adult, either in one piece, or divided into fingers; a sinus at the lower part of the mouth. Marine.) <i>Rostellaria, Pteroceras, Strombus.</i></p> <p>13. PURPURIDÆ. (<i>Shell</i> with a short canal ascending and directed towards the back, or a notch; generally globular; left lip often spotted or rough. Marine.) <i>Cassidaria, Omiscia, Cassis, Ricinula, Purpura, Monoceras, Concholepas, Harpa, Dolium, Buccinum, Nassa, Eburna, Terebra.</i></p> <p>14. COLUMELLINÆ. (<i>Shell</i> notched only. Left lip plaited distinctly. Marine.) <i>Columbella, Mitra, Voluta, Melo, Cymba, Marginella, Volvaria.</i></p> <p>15. CONVOLUTIDÆ. (<i>Shell</i> notched, whorls large, compressed, the last nearly or quite covering the whole of the rest. Marine.) <i>Cypræa, Erato, Opula, Terebellum, Ancillaria, Oliva, Conus.</i></p>	<p>1st. POLYTHALAMA. (<i>Shell</i> divided into chambers.) <i>Belemnites, Orthoceras, Spirula, Nummulus, Nautilus, Ammonites, Turritiles, Bacculites.</i></p> <p>2d. MONOTHALAMA. (<i>Shell</i> not divided into chambers.) <i>Argonauta.</i></p> <p>3d. No shell. <i>Octopus, Loligopsis, Loligo, Sepia.</i></p>	<p><i>Carinaria, Almonia, Pterotrachea, Phyllirhoe.</i></p>

Animal. Soft, furnished with a head, most frequently with eyes and tentacula, or possessing arms disposed in the form of a coronet. Body sometimes naked, sometimes enclosing a shell, but generally having an external shell.

only.

1. PTEROPODA.	2. GASTEROPODA.		PODA.
<p><i>Animal.</i> Having no feet, but two opposite fins, adapted to swimming.</p> <p><i>Shell.</i> Thin; cartilaginous, or horny.</p>	<p><i>Animal.</i> Foot united to nearly the whole length of the body, and used for crawling. Body straight, never spiral.</p> <p><i>Shell.</i> Either cup-shaped, or slightly rolled up; in some genera not covering the body.</p>	<p><i>Animal.</i> Feet always enclosed in shell. Spiral.</p> <p><i>Animal.</i> Feet either marine or fresh-water.</p>	<p>Feet or tentacles. Spiral.</p> <p>trans-</p>
<p><i>Hyalæa, Clio</i> (no shell), <i>Cleodora, Limacina, Cymbulia, Pneumodermone</i> (no shell).</p>	<p>EIGHT FAMILIES.</p> <ol style="list-style-type: none"> 1. TRITONIDÆ. (No shell.) <i>Glaucus, Eolis, Tritonia, Scyllæa, Tethys, Doris.</i> 2. DENTALIDÆ. (Shell tubular, open at both ends.) <i>Dentalium.</i> 3. PHYLLIDIDÆ. (Shell either of one cup-shaped piece, or with moveable pieces at the back.) <i>Phyllidea</i> (no shell), <i>Chitonellus, Chiton, Patella, Siphonaria.</i> 4. SEMIPHYLLIDIDÆ. (Shell cup-shaped, but very flat.) <i>Pleurobranchus, Umbrella.</i> 5. CALYPTREIDÆ. (Shell cup-shaped, though irregularly so; some genera with internal pieces.) <i>Parmophorus, Emarginula, Fissurella, Crepidula, Calyptræa, Hyponix, Pileopsis.</i> 6. BULLÆIDÆ. (Shell slightly rolled up.) <i>Acera</i> (no shell), <i>Bullæa, Bulla.</i> 7. APLYSIDÆ. (Shell nearly flat and thin.) <i>Aplysia, Dolabella.</i> 8. LIMACIDÆ. (Shell very slightly rolled or cup-shaped.) <i>Onchidium</i> (no shell), <i>Limax, Parmacella, Testacella, Vitrina.</i> 	<ol style="list-style-type: none"> 1. COLIMACIDÆ. The right or turreted. <i>Helix, Clausilia, Cyclostoma.</i> 2. LYMNÆIDÆ. always shallow. <i>orbis, Physa.</i> 3. MELANIDÆ. lip sharp. 4. PERISTOMIDÆ. lips united. <i>vata, Paludina.</i> 5. NERITIDÆ. left lip retracted. <i>Fluviatilis, Neritina.</i> 6. JANTHINIDÆ. mouth triangular. 7. MACROSTOMIDÆ. very large. <i>Stomatella.</i> 8. PLOCIDÆ. sharp. <i>Meloidella.</i> 9. SCALARIDÆ. touching, <i>metus, Marginella, Delphinula.</i> 10. TURBINIDÆ. midal, subcylindrical. <i>Solarium, Planaxis,</i> 	<p>Atlan- tarchæa,</p>

PART IV.

MOLLUSCA OR UNIVALVES.*

THE molluscs of univalve shells are all furnished with a head, and a foot or fins, and are divided into five orders, which are distinguished from each other by the construction of the foot : they are,—1. *Pteropoda* †, having fin-like feet used for swimming. 2. *Gasteropoda* ‡, having the foot proceeding from or covering the stomach. 3. *Trachelipoda* §, having it attached to that part which lies between the head and body, or neck. 4. *Cephalopoda* ||, having them (for there are more than one) fixed to the head. And 5. *Heteropoda* ¶, with feet or fins varying in different genera.

In the annexed plan are shown, at one view, the divisions and subdivisions of the order Mollusca or Univalves ; but before entering upon

* *Les Mollusques*, Lamarck.

† From *pteron*, Gr., a wing, and *poda*, feet.

‡ From *gaster*, the stomach, and *poda*.

§ From *trachelo*, the neck, and *poda*.

|| From *cephale*, the head, and *poda*.

¶ From *heteros*, various, and *poda*.

them, it will be advisable to give representations of a few different kinds of univalve shells, with a description of the parts constituting them, and the names by which they are known, as in giving an account of the variations which exist in the different genera many of these will be referred to. In fig. 1. is delineated the simplest form, that of a *Patella*, in which the parts necessary for reference are but few. A is the *apex*, B the *mouth* or *aperture*, of which CC is the *margin*. The *exterior* and *interior* are sufficiently obvious. In figures 2. to 7., which are turbinated or spiral shells, and of a more complicated form, a greater number of terms have been adopted to mark the variations which occur. A is the *apex*, B the *mouth* or *aperture*, which in fig. 2. is *entire*; in fig. 3. *notched*; in fig. 4. *channelled*. D is the *spire*, formed of a greater or less number of *whorls*, the intersections of which, or where they wrap over, are called *sutures* E. F is the *right* or *outer lip*; G, the *left, inner, or columella lip*; H, in fig. 3., is the *notch*; I, in fig. 4., is the *channel* or *canal*; K, in figs. 5. and 6., the *umbilicus*; L, in fig. 7., is the *columella*, or *pillar*, which can only be

seen when the shell is cut open; M are the *plaits* on the left lip of fig. 3.

ORDER I.

PTEROPODA.*

The animals of this order have no feet to crawl with, but are free and floating, and have two opposite and similarly constructed fins adapted for swimming. Some genera only are furnished with a thin, cartilaginous, or horny shell. The greater part are animals of a small size. There are six genera, as follows:

1. HYALÆA. — *Shell* horny, transparent, ovate, globular; tridentated posteriorly, open at the summit and at the two posterior sides. *Animal*, body covered by a shell; head very indistinct; no eyes. — 17 species recent, 1 fossil.

These shells are brought from the Mediterranean and Atlantic. Their trivial name in England is Venus' Chariot.

2. CLIO.—No shell. *Animal*, body oblong; head provided with several tentacula; two eyes

* *Les Ptéropodes*, Lamarck.

on the upper part of the head.—7 species recent.

These mollusca inhabit the sea, where they are seen in calm weather and during the hottest part of the day. They appear and disappear continually on the surface. Found in the seas of the Cape of Good Hope, India, and the Northern Atlantic.

3. CLEODORA. — *Shell* gelatinous, cartilaginous, transparent, in the shape of a reversed pyramid, truncated, and open at the summit. *Animal*, body oblong; head distinct, having two eyes; no tentacula.—11 recent and 2 fossil species.

The Indian and Atlantic Oceans produce these curious animals.

4. LIMACINA. — *Shell* thin, fragile, papyraceous, and spiral; the whorls discoid or flattened. *Animal*, body long, and greatly resembling the Clio. — 1 species recent.

The animal can retire completely into its shell. It inhabits the Northern seas; and is said to be devoured by whales in vast quantities.

According to Mr. Scoresby, it is found in great profusion near Spitzbergen.

5. CYMBULIA. — *Shell* gelatinous, cartilaginous, very transparent, crystalline, oblong, in the shape of a shoe, truncated at the summit; aperture lateral. *Animal*, body oblong, transparent, enclosed in the shell; head having two eyes and two retractile tentacula. — 5 species recent.

This singular shell is as transparent as glass. It is found in the Mediterranean and Indian seas.

6. PNEUMODERMON. — No shell. *Animal*, body oblong, head distinct; tentacula placed on each side the mouth; no eyes. — 3 species.

Found in the Atlantic Ocean and the seas of Amboyna.

ORDER II.

GASTEROPODA.*

Some of the animals of this order have no shell; others have a dorsal shell not enveloping them; and others, again, have one more or less hidden in the mantle. The body is straight, not spiral, having a foot united to it nearly the whole length, and used for creeping. They are divided into eight families: — 1. *Tritonidæ*. 2. *Dentalidæ*. 3. *Phyllididæ*. 4. *Semi-phyl-lididæ*. 5. *Calyptræidæ*. 6. *Bullæidæ*. 7. *Aply-sidæ*. 8. *Limacidæ*. The seven first live in the sea, the last on land.

FAMILY 1. — *Tritonidæ*.†

The Mollusca which form this family have no shell, either external or internal. The body is long, and the respiratory organs distinct and placed above the mantle, either on the back or sides. They are all marine: six genera.

* *Les Gastéropodes*, Lamarck.

† *Les Tritoniens*, Lamarck.

7. Glaucus.	10. Scyllæa.
8. Eolis.	11. Tethys.
9. Tritonia.	12. Doris.

FAMILY 2. — *Dentalidæ*.

The branchiæ of this family are divided into two tufts of soft, flexible, club-shaped filaments, attached to a pedicle on the neck. — Only one genus is known. (See p. 30. note.)

13. DENTALIUM. — *Shell* tubular, regular, smooth or fluted, more or less curved, gradually

tapering to one end, open at both extremities. *Animal* long, slender, having a distinct head. — 21 recent, 1 fossil species.

Lamarck thought these animals belonged to the *Annulata*, but they are now placed amongst the *Mollusca*. The shell resembles in form an elephant's tusk. Some species are ribbed longitudinally, others transversely, but most of them are smooth. They inhabit the European and Indian seas.

FAMILY 3. — *Phyllididae*.*

Some of the shells comprised in this family differ from all other univalves, in being composed of moveable pieces at the back, formed to facilitate the motion of the animal. They are shaped like a boat, and adhere to rocks covered by the sea, the backs of turtle, and other marine objects; and so fast do they retain their situation, by exhausting the air beneath their bodies, that it requires a strong lever to detach them, frequently breaking the shell before the animal can be forced from its position. The others are particularly simple in form, presenting, in fact, scarcely any thing more than a conical cup or deep dish. They adhere to rocks and buildings washed by the sea; and although of a dull colour outside, are often very beautiful in the interior.—This family consists of five genera.

14. PHYLLIDEA. — *Shell* none. *Animal*, body long, four tentacula attached to the head. — 4 species.

* *Les Phyllidiens*, Lamarck.

They are found in the Indian seas.

15. CHITONELLUS. — *Shell* divided into several small, disunited pieces, placed in nearly a straight line down the back of the *animal*,

which, in form and manner of progression, resembles a caterpillar. — There are 4 recent species.

The animal of the *Chitonellus* can move with great rapidity, turning its body with ease both to the right and left, as it creeps on the stalks of marine plants. It is found only in the seas of Australia and the Philippine Islands.

16. CHITON. — *Shell* boat-shaped, composed

of about eight transverse pieces, folding over each other at their edges, and inserted into a

tough ligament. *Animal*, head not very distinct; no tentacula nor eyes, but having a long tongue rolled spirally, and armed with horny teeth; foot long.—70 species recent, 1 fossil.

The ligament of this shell is sometimes fleshy, smooth, or wrinkled, covered with scales or tufts of hair; and in one species, the *Chiton spinosus*, it is armed with numerous thin long black spines. The interior of the shell is generally white; but some are green, and one species is of a beautiful rose colour. They often attain a large size, measuring several inches in length; but do not usually exceed two inches. They are found in most seas, the larger species coming from warm latitudes. Several, of a small size, are found on the British coast. They have the power of rolling themselves up into a ball, like the woodlouse or the hedgehog.

17. PATELLA.—*Shell* simply conical, without any perforation or internal appendage, and with the margin undivided; apex inclined towards the front part of the shell, where the head of the animal is situated, the mark of which can generally be traced in the interior by its

differing in colour from the rest of the shell. *Animal*, head very distinct, supporting two ten-

tacula, having eyes at the lower extremity; foot large, and round or oval.—There are 57 species recent.

The genus *Patella* is to be found in all parts of the world, chiefly on rocky coasts, and often where the sea breaks with considerable violence; many species, (from the shape of the shell corresponding exactly with that of the rock on which they have lived,) appear never to move from the spot they have once selected, and even form in most cases cavities in the rock itself by a power they possess of absorbing its substance: the *Patella vulgata* of the British coast has this property.

The *Patella cochlea* is found at the Cape of Good Hope; it lives almost exclusively attached to a larger species of the same genus,

on the surface of which it forms a flat disk exactly the size of its mouth. To form these flat disks (of which there are so generally two, one on each side of the apex of the larger *Patella*, as almost to form a character of the species), and to assist in the increase of its size, the animal appears to absorb the coralline, or other similar substances, with which the larger shells are abundantly covered.*

There are other species, on the contrary, which do not remain entirely attached to the same place, but occasionally move; and others, again, which fix themselves to sea-weeds, which appear to be the food of the whole genus. They are particularly abundant at the Island of Cyprus and the Cape of Good Hope; where they are often found of a very large size, and are used as spoons by the Hottentots. The common English species is generally smooth on the outside; but many of the foreign ones are covered with ridges in rays from the apex to the circumference. Sometimes these ridges extend beyond the edge of the margin, and ap-

* Gray, Philosophical Transactions, 1833.

pear like the rays of a star. Some species were used for the table in former times, and the common species of the British coasts is still gathered by the poorer people for the same purpose.

18. SIPHONARIA (Sowerby).—*Shell* resembling the *Patella* in shape, but with an indented groove in the interior, extending from the apex

to the margin. *Animal*, head divided into two equal parts ; no tentacula, nor visible eyes ; the latter, at least, are very doubtful ; foot moderate.—There are 19 species recent, and 2 fossil.

It is difficult, at first sight, to distinguish this shell from a *Patella*, the groove not being very perceptible ; indeed, many authors class them together. They are found on the coasts of South America, New Holland, the Isle of France, and in the Mediterranean. The animals appear to be ornamented with various colours.

FAMILY 4.—*Semi-Phyllididæ*.*

The animals of this small family only differ from the last in the situation of the branchiæ, which are placed on the right side of the body. The shells are cup-shaped, but nearly flat. There are only two genera.

19. PLEUROBRANCHUS. — *Shell* thin, nearly flat and obliquely oval; it is internal, or covered by the animal. *Animal*, head distinct, with two cylindrical tentacula; foot large.—There are 7 species recent.

It is found in the Indian Seas and the Mediterranean.

20. UMBRELLA. — *Shell* rather irregular, slightly convex on the outside, and with a small, nearly central apex; margin sharp. *Animal*, head not very distinct, with four tentacula; foot very large. — There are 2 species recent.

The *Umbrella Indica* is met with in the Indian Ocean, and on the coasts of the Isle of France,

* *Les Semi-Phyllidiens*, Lamarck.

and is commonly called the Chinese Parasol : it is thin, rather transparent, and the interior

is marked with rays proceeding from the centre, and with a yellowish stain. The *Umbrella Mediterranea* is from the Gulf of Tarento, and is smaller than the preceding, also differing from it in not being marked with rays.

FAMILY 5. — *Calyptræidæ*.*

This family is composed of animals that differ principally from the two last in the situation of the branchiæ, or respiratory organs. They are placed in a cavity appropriated to

* *Les Calyptraciens*, Lamarck.

them on the back of the animal near the neck, projecting either within the cavity or above it. The shells, in many instances, resemble the *Patella* in form, though differing from it in other particulars. They are always external, and cover the animal.—7 genera.

21. PARMOPHORUS. — *Shell* oblong, depressed slightly, convex on the outside; apex small, and somewhat inclining to one end. *Animal*, head rather indistinct, with two tentacula, having eyes at the base; foot very large.—There are 3 species recent, and 2 fossil. This genus is found in New Holland and New Zealand.

The interior exhibits very strong muscular impressions, which are in some species marked with a blood red colour. It is often called the Duck's Bill Limpet.

22. EMARGINULA. — *Shell* rather oblong, with the apex curved, and having a narrow slit or hollow in the margin, generally rayed on the outside. *Animal*, head

having two short tentacula, with eyes at the base; foot large and very thick. — There are 11 species recent, and 5 fossil.

These are all small shells, and are found in European Seas, at Amboyna, the Isle of France, New Holland, &c.

23. FISSURELLA. — *Shell* resembling a *Patella*, but having a round or oval hole at the apex.

Animal, a large and distinct head, having two tentacula, with eyes at the base; foot large.

There are 30 species, of which several are fossil.

The hole at the apex of this shell, which serves as a passage for water to the branchiæ of the Mollusc, which are situated near it, differs in form, varying also with the age of the shell, being in some species round, whilst in others it is oblong, or resembling a key-hole. The shells are generally thick and strong, and often delicately marked with alternate dark and light colours. A few species are found on the coast of England, in the same situations as

the *Patella*; but none are so beautiful as those which are found in the Tropics.

24. CREPIDULA.—*Shell* irregular in shape, and often very much flattened; the inside partly covered with a plate, so as to resemble a half-decked boat; apex not in the centre. *Animal*, head forked, having two tentacula, with eyes at the base; foot small.—There are 25 recent species*, and 1 fossil.

This is a curious and often a very beautiful shell. The *Crepidula Onyx* is of the most brilliant black in the inside, with the little half-deck, as it may be called, of a beautiful white, and having the margin of the shell tinged with a rich brown. These shells are found on rocks, and are often of a very irregular outline at the circumference; like the *Patella*, they constantly remain upon the same portion of rock, and form the margin to agree with its shape.

* Several of the recent species are also found in a fossil state.

They are found in great abundance at the Cape of Good Hope. One species frequently fixes itself upon other living shells, particularly upon the *Purpura*, and of course follows it in all its movements.

25. CALYPTRÆA. — *Shell* variable in form, and often very brittle, with a cup or funnel-shaped appendage in the interior.

Animal, head large, having two long thin tentacula, pointed at the end, and furnished with eyes in the middle; foot not large. — 35 species recent, and 5 fossil.

The shell of the *Calyptræa* is often irregular in shape, similar to that of the last genus, and is often, like the *Crepidula*, found attached to large stones, shells, &c.: the outside is frequently covered with spines or ribs. It is generally thin, and not coloured. One species is called the Cup and Saucer Limpet, from its resemblance to those articles; it is brought from the South Seas. Another very curious species multiplies the little funnel-shaped appendage, so that it forms several little stages one above the other; it is called the *Calyptræa*

tectum sinense. Several species have the apex rather spiral. They are found in the seas of Europe, New Holland, &c.*

26. HIPPONYX (Defrance).—*Shell* obliquely cap-shaped, and sometimes, perhaps generally, supported on a solid shelly plate; the shell and

plate exhibit two strong muscular impressions, which are in the form of a horse-shoe. *Animal*, similar to *Pileopsis*.—9 species, recent and fossil.

These animals do not always form a plate; indeed, they are as often found without the sup-

* Mr. Cuming has lately found at the island of Zebu, one of the Philippines, several specimens of *Calyptraea equestris* on masses of dead coral about low water mark, having a testaceous plate attached to the mass to which the true shell adheres. The edges are finely crenulated, and turned up all round like a platter, as if to protect the margin of the cap-shaped shell which rests within it.—Reeve's *Con. Sys.* ,

port as with it; for when they adhere to other shells, the same object is gained by absorbing a suitable place of attachment, on which the same muscular impressions may be traced.

27. *PILEOPSIS*.—*Shell* irregular, conical; the apex lengthened, and re-curved backwards; inside dull. *Animal*, head distinct, having two tentacula, with eyes at the base; foot very large.—There are 10 species recent, and 12 fossil.

This shell is frequently called, from its shape, the Fool's Cap Limpet. It is found in the European Seas principally; but some very large species are inhabitants of tropical climates, and are met with in the Pacific Ocean, and in the East and West Indies. The *Pileopsis Ungarica* is the only one of which the animal is known. The shells of this genus, like many of the species of *Patella*, as before remarked, are supposed to remain constantly attached to the spot they have once fixed upon, from the circumstance that the form of the margin of the shell agrees exactly with the portion of rock

on which they live, and preserves the same form to adult age. The *P. Ungarica* of the British Seas has the inside bright, and often of a beautiful rose colour.

Lamarck's system includes the genus *Ancylus* amongst the Patelliform shells, but it is now placed with the *Lymnæidæ*.

FAMILY 6.—*Bullæidæ*.*

The shells forming this family are very simple in form, but differ from the preceding in some respects; they are all slightly rolled up, without being spiral. There are three genera forming this family, which are described as follows.

28. ACERA.—*Shell* none. *Animal*, oblong and flat; head, not distinct; foot very large.—1 species, found in the Mediterranean.

29. BULLÆA.—*Shell* less rolled up than the subsequent genus, very thin, and hidden in the folds of the animal's mantle, without adhering to the body. *Animal*, head not distinct; no

* *Les Bulléens*, Lamarck.

tentacula.—There are only 2 species recent, and 2 fossil.

The *Bullæa aperta* is found in the European Seas ; it climbs aquatic plants well, but swims badly. The *Bullæa hirundinina* is found at the Isle of France. The animal is of a beautiful blue colour.

30. BULLA.—*Shell* rolled up, aperture extending the whole length. *Animal*, head very indistinct ; no apparent tentacula ; hinder part of the body covered by the shell, which is external, and adheres by a muscle ; foot large. — 26 species recent*, and 11 fossil.

The *Bulla* can nearly, but not entirely, enter into its shell : it swims well, and is found in the seas of Europe, India, &c. The animal of the *B. lignaria* possesses a very singularly formed testaceous substance within its body, not no-

* Some of the recent species are also found in a fossil state.

ticed by Lamarck, but called by English collectors the gizzard. It is composed of three bony pieces, united by a muscular membrane; the two side pieces are triangular, the other is small.

The *B. Velum* is a very delicate shell, of a very light horn-colour, with a snow-white band about the middle, bordered on each side by one of a dark brown colour; the apex and base are white, both bordered by dark brown bands.

FAMILY 7.—*Aplysidae*.*

The Mollusca constituting this small family offer only two genera, and the shell with which they are furnished, if it can be called a shell at all, is a mere rudiment of one, forming only a covering for the gills; this at least is the case with the first genus; the other, in lieu of this plate of defence for its breathing apparatus, has a more defined shell, serving the same purpose.

31. APLYSIA.—*Shell*, or operculum for gills,

* *Les Laplysiens*, Lamarck.

an oval, thin, nearly transparent plate. *Animal*, body creeping, oblong; head with four tentacula, furnished with eyes; mantle large. — There are 22 recent species.

These animals resemble large slugs; they swim easily, but crawl slowly: they are found in the Mediterranean Sea. Darwin, in his *Journal**, says it feeds on delicate sea-weeds which grow among the stones in muddy and shallow water; and that he found in the stomach of several, small pebbles as in the gizzards of birds. He also mentions that, when irritated, it emits a fine purplish-red fluid, which stains the water for the space of a foot round, and that, besides this means of defence, an acrid secretion, which is spread over its body, causes a sharp stinging sensation. He found it at St. Jago.

32. DOLABELLA. — *Shell* oblong, thick, and slightly spiral on the posterior side, thin and flattened on the other. *Animal*, oblong, hav-

* *Journal of Researches into the Geology and Natural History of the Countries visited by the Beagle.*

ing four tentacula ; it is doubtful whether it has eyes ; foot large.—There are 11 recent species.

This animal resembles the *Aplysia* exceedingly, but has a more ample mantle ; it inhabits tranquil bays in India, and the Isle of France.

FAMILY 8.—*Limacidae*.*

This family, like the last, has a very slightly developed shell, forming merely a covering for the branchiæ, and into which the animal is unable to enter. All the genera are terrestrial, breathing air, but preferring moist places to those perfectly dry. There are five genera, two of which do not possess shells, although they have a substance protecting the organs of respiration.

33. ONCHIDIUM. — *Shell* none. *Animal*, long ; head with four tentacula ; foot long. Found

* *Les Limaciens*, Lamarck.

in the Islands of Bourbon and Martinique in woods and gardens.—8 recent species.

34. LIMAX.—*Shell* internal and rudimental. *Animal*, long; head tolerably distinct, having

two pair of tentacula; foot large and oblong.—19 recent species.

These animals are common everywhere, but particularly in the neighbourhood of ditches and gardens, where they often cause great damage. They are usually known by the name of *slug* in England.

35. PARMACELLA.—*Shell* merely rudimental, like the *Aplysia*, and defending the organs of respiration. *Animal*, oblong, having four tentacula, two of which have eyes, and all are retractile; foot large.—1 species.

This animal has been found in Mesopotamia.

36. TESTACELLA.—*Shell* very small, slightly ear-shaped, and spiral at the apex; mouth large, oval, having the left lip rolled inwards,

and the right sharp. *Animal*, body long like a slug, with the shell only covering the extre-

mity; head distinct, having four tentacula, the two largest of which have eyes at the summit; foot long.—2 species recent.

This animal is rarely observed, as it burrows under ground, where it lives upon earth-worms. One of the latter, an inch long, having been placed under a glass with three of these slugs, soon fell a prey to one of them; and another, three inches in length, although at first it succeeded in disengaging itself, was finally devoured by them, as well as two others of the same size which were subsequently introduced: two of the slugs were observed to make a simultaneous attack upon one of the worms, each commencing its repast at a different extremity.

The *Testacella haliotidea* is found in Europe; and M. Ferussac mentions a species, the *Testacella Maugei*, which was brought from Teneriffe

by Maugé to the Botanic Garden at Bristol, where it has become acclimated. As natives of England, they were first noticed by Mrs. Smith of Bristol, who found the shells casually scattered about her garden, and afterwards discovered the animal. They are common in Guernsey, and have been found at Lambeth. In winter they bury themselves from one to two feet deep in the earth, and are most above the surface from August to November.*

37. VITRINA.—*Shell* small, very thin, flattened, terminated by a short spire, the last whorl large; mouth large, and rather oval. *Animal*, body long, the shell only covering the extremity†; four tentacula, two of which have eyes at the summit.—7 species.

These mollusca live in shady and damp places among moss and short grass, and are found in Great Britain, Germany, Sweden, Celebes, &c. These little shells are distin-

* Gray.

† The French naturalists say that the animal cannot entirely enter its shell; but an English observer (Mr. Nilson) remarks that they not only enter it, but leave a space behind like the snails.

guished from young specimens of the *Helix* by their never being umbilicated or perforated; they greatly resemble them in other particulars.*

ORDER III.

TRACHELIPODA.†

The mollusca constituting this order differ from those which are comprehended in the last by having the foot attached to the neck, and the hinder part of the body coiled up and covered by a shell, which is always spiral, and to which the animal is fixed: this part of the body never leaves the shell, but the head and foot can also be drawn in at pleasure. The *Trachelipoda* are much more diversified in form than the last order, and are divided into a far greater number of genera, some of which inhabit the sea, others are found in fresh water, whilst others again are met with on land. Many of these are herbivorous, and many carnivorous; it will therefore, from the great

* Draparnaud.

† *Les Trachélipodes*, Lamarck.

extent of the order, be advisable to separate it into those two natural divisions, or into *Phytophaga* and *Zoophaga*.

DIVISION I. — PHYTOPHAGA.

The *Phytophaga*, or herbivorous *Trachelipods*, have the mouth of the shell entire; that is, without a notch or channel. Those which are marine procure their food from the seaweed which they find floating on the surface or at the bottom; and the land species obtain abundance of nourishment in our gardens and elsewhere.

They are divided into ten families: — *Colimacidae*, *Lymnæidae*, *Melanidae*, *Peristomidae*, *Neritidae*, *Janthinidae*, *Macrostomidae*, *Plicacidae*, *Scalaridae*, and *Turbinidae*.

FAMILY 1. — *Colimacidae*.*

The animals of this family are entirely terrestrial, though usually fond of damp situations. Only two of the genera have permanent opercula, the others forming merely a tem-

* *Les Colimacées*, Lamarck.

porary epiphragma to protect them during severe weather, which they throw off on the return of warmth. The shells are spiral, with no projecting parts on the exterior, excepting those caused by increase of growth; the right lip or margin of the aperture is often recurved, or reflected outwards. There are eleven genera. The animals of eight of these genera have four tentacula, their eyes being placed at the summit, the others only two.

38. *HELIX*. — *Shell* globular, convex, or flattened; spire not much elevated; mouth wide,

not long, placed very obliquely to the axis of the shell; lips disunited by the projection of the last whorl, never pearly, but polished; um-

bilicus either covered or visible. *Animal*, head furnished with four tentacula, the two longest having eyes at the summit. Foot large and oblong. — There are 225 species recent, and 16 fossil.

Throughout the whole of nature there are very few classes of animated beings more abundant, or more completely spread over the surface of the earth, than this genus, the *Helix*, or common snail. It is found even in the most barren and desert places, feeding upon the miserable specimens of vegetation which may be found, and existing in great numbers in places entirely deserted by the other ranks of creation. It is equally adapted to the hottest and the coldest climates, the most cultivated and most barren situations. The *Helix aspersa*, or common garden snail, may be found at the foot of Chimborazo, in the forests of Guiana and Brazil, and almost all over the continents of Europe, Asia, and Africa; even the great desert of Zahara is peopled by them. On the shores of the Mediterranean they are a valuable article of food, when boiled in the shell and eaten with rice; and it has been conjectured

that they formed a portion of the food of the Israelites both in the desert and on their way to the Red Sea, the country in that neighbourhood being described by travellers as having a herbage under the trees completely covered with snails of a large size. They form in some countries, as Switzerland, Britany, and some parts of Germany, a considerable article of commerce. They are fed by thousands in places made on purpose for them. It is supposed the Romans also fattened the *Helix pomatia* in the same manner. Even at the present day the *Helix aspersa* is sold in Covent Garden market. It is used, boiled in milk, for diseases of the lungs. They are also sent to America from this country as a delicacy.

These animals become torpid about October (at least in England), and remain so during the winter, having first closed the mouth of their shells with an epiphragma which they throw off, and emerge when warm weather returns. Some of them form two or three of these doors, one within the other, so as completely to exclude the cold; and others bury themselves in the earth. The *Helix aspersa*,

which is the pest of our gardens, is more influenced by the weather than many of the smaller sort; for upon the first appearance of cold these animals creep into crevices and under stones, clustering and clinging together as if they were capable of communicating warmth by association. In the centre of the epiphragma is an exceedingly minute orifice, communicating with the lungs; and this minute hole, though not large enough to admit a drop of water, is of sufficient capacity for the passage of air for the purpose of extremely slow, but not totally extinct respiration.*

Many instances have been known of the great tenacity of life which snails possess. A gentleman received some species closely packed with marine shells from the Mauritius; the day after they were unpacked they were discovered wandering about, apparently uninjured from their long and close confinement. Some lived ten days; but a want of proper food or the uncongenial climate then killed them. There are instances, also, of

* Gray's edition of Turton's Manual.

snails having been some years in a cabinet, and on the shells being put into warm water to be washed the animals made their appearance alive.

These mollusca have the power of repairing their bodies when injured as well as their shells: a horn, or even the whole head, if cut off, will be reproduced, the animal keeping within its shell until grown again. The head, however, requires some months to be replaced.

The *Helix pomatia* is said by some authors to have been introduced into this country from abroad, but others suppose it to be indigenous. It is almost peculiar to chalky and gravelly soils. The eggs of this species are globular, covered with a white opaque skin, and are about two and a half lines in diameter.*

Mr. Gray mentions having seen a specimen of a *Helix aspersa* curiously deformed by the obstruction of a young shell of the same species. The young specimen is attached to the spire, to which it had doubtless fixed itself during the dry season; and not waking from its torpor so early as its older companion, the

* Turton.

latter, when it commenced increasing the size of its shell, threw its new whorl partly over the smaller individual, which was thus enclosed in a prison formed by its own shell. In this instance the form of the larger specimen is not much altered; but about one half of the young shell projects above the spire.*

39. CAROCOLLA.—*Shell* flat; mouth wide, not long, with a thickened lip, and generally toothed within; circumference of the whorls

often angular; umbilicus either covered or visible. *Animal*, having a head furnished with four tentacula, the two largest having eyes at the summit; foot large and oblong.—18 species recent.

These shells, and the animals inhabiting them, much resemble the *Helix*; the latter, however, have not the edge of the whorls

* Philosophical Trans. 1833.

angular. They inhabit all parts of the world, living in dry as well as damp situations. Turton mentions one species as a native of England, *C. lapicida*, found in the fissures of limestone rocks, and in woods.

40. ANASTOMA.—*Shell* orbicular; spire obtuse; mouth toothed, and turned upwards in

the adult, nearly reaching the level of the apex. *Animal* like that of the *Helix*.—2 species, recent.

The structure of this extraordinary shell proves that in completing the last whorl, the animal carries it with the apex downwards, instead of its natural position upwards, which is the case with all other shells, and even with itself when young. It is very rare, and is brought from the East Indies.—Only 2 species are known, *Anastoma depressa* and *Anastoma globulosa*.

41. *HELICINA*. — *Shell* flattened; mouth semicircular, closed by a horny operculum; lip thickened, and columella often also thickened.

Animal, having a head furnished with a proboscis, and two tentacula, with eyes at the base on tubercles; foot rather short.—There are 19 species recent, and 1 fossil.

Some species feed on trees, and others on the ground; they are natives of the West Indies and other countries of the torrid zone. The operculum, which is permanently attached to the foot, is not spirally formed, but in concentric layers.—The *H. viridis*, from St. Domingo, is of a smooth shining green colour.

42. *PUPA*. — *Shell* cylindrical; spire long; aperture irregular, toothed, with the lips thickened, and unconnected at the upper part, the plaits on the left lip intervening. *Animal* like the *Helix*. —

There are 56 species recent, and 2 fossil.

They are found principally in Europe, but a few species are from America: they derive their name from the resemblance of the shell in shape to the *pupa* or chrysalis of an insect. One species is named the *Pupa chrysalis*.—Several species are found in England under stones, old walls, and in woods.

43. CLAUSILEA.—*Shell* fusiform or spindle-shaped, rather resembling the last genus, but more slender; aperture irregular, rounded or ovate; lips united; summit obtuse. *Animal* like the *Helix*.—There are nearly 150 species recent, and 1 fossil.*

The *Clausileæ* inhabit mosses at the foot of trees. Some species are English, and many are found in other parts of Europe, particularly on the borders of the Mediterranean; and Russia and the Crimea produce a great number of curious species, which have lately been discovered; so that it is thought there are more than 150 species: a few also are American.

* There are 44 species described by Lamarck and Deshayes, but many more species have been discovered lately.

They are all small shells, the largest scarcely exceeding an inch in length. Within the mouth, in the last whorl but one, there is a little elastic shelly plate attached to the shell, and called a *clausium*, from which the genus takes its name. It is used to close up the aperture when the animal has retreated within its shell, and in that respect resembles an operculum, or epiphragma; but differs, inasmuch as the latter are either attached to the animal, or are loose and thrown off, whereas the *clausium* is fixed permanently to the shell. Turton mentions five or six as natives of England. Many of the species are sinistral.

44. BULIMUS. — *Shell* oval, turreted, last whorl large; aperture having the margins dis-united at the upper part, long, not wide; lip thickened and reflected in the adult; spire obtuse, but much more prominent than in the *Helix*. *Animal* like the *Helix*. — 135 species recent, and 16 fossil.

These shells are found in almost every part of the world, from the torrid to the frigid zones; in the former they attain a large size, and are beautifully coloured, often with lemon, amber, and sea-green, which rarely exist amongst other shells. Many of the species are reversed. They are more commonly found on islands, or on the margin of the sea amongst rocks and trees, than in the interior of the country. The eggs of the *Bulimus hæmastoma* are large, even sometimes reaching the size of a pigeon's egg, and are represented in the annexed fig.

A specimen of the *Bulimus ovalis* was once brought from Rio Janeiro, and placed in a hot-house at Chiswick; it lived for more than a year, and fed principally on lettuces and cab-

bage, the former of which appeared to be its favourite food. It would sometimes, after devouring part of a large lettuce, remain for some days without touching food or moving from the spot. During the day it was usually in a dormant state, and remained in the shade; but towards evening, when the house became damp and warm, it would move about.

Some specimens that were brought from Chili continued in a state of torpidity for seventeen or eighteen months, and one even more than two years, and revived in this country. They were sent to Messrs. Loddiges' gardens, at Hackney, where they lived for eight months in a hot-house fitted for palm-trees: they had constructed a parchment like epiphragma, which they were observed partly to eat when they revived. They had been packed in cotton, in a box, from the period of their capture. Mr. Cuming brought many new species from South America. The *B. rosaceus* he found adhering to the under side of stones, generally among bushes, and close to the edge of the sea shore within reach of the spray. He also met with them at the top of hills about

one thousand feet above the sea; they were adhering to the lower leaves of an aloe-like plant. In the spring they burrow in the sand, and lay their eggs about two inches below the surface. When they hybernate, they adhere so fast to the stones and plants that they can hardly be taken without breaking the shell.* One species, the *B. decollatus*, as it advances in age always loses the upper volutions; but whether from accident or design appears uncertain. It, however, generally closes up the broken end; and the probability is that as the animal enlarges in growth it recedes from the smaller volutions, closes them up, and these neglected portions fall off. Turton mentions a colony of them being discovered in a greenhouse in Devonshire, lodged in the earth under the wood-work, whence they wandered abroad in the summer. No foreign earth was ever known to have been admitted into the house; and they were considered by the gardeners as natives.

Other species, particularly *B. articulatus*, are

* Penny Cyclopædia.

met with in such numbers in some parts of Cornwall, that it is impossible to take a step without crushing them. It is the prevailing opinion that they contribute much to fatten the sheep which browse upon the herbage where they are met with, which is principally on sandy maritime pastures.

45. *ACHATINA*. — *Shell* somewhat like the last genus, but longer; the lip thin, and ap-

parently notched at the base of the pillar, from its rolling over; aperture about half the length of the shell; no operculum. *Animal* like the *Helix*. — 35 species recent, 1 fossil.

Some of the *Achatinae* are the largest of all known land shells; they live always near water about trees, and are very plentiful in Africa, particularly at the Cape of Good Hope. Some are found in the West Indies; but very few in Europe. Two small species, *A. acicula* and *A.*

octona, are found in England, among the roots of trees at the base of limestone rocks. Many of them, as the *Achatina zebra*, are covered with a thick epidermis, but others are destitute: a few are constantly reversed or sinistral shells. The *Achatina columnaris* is one of the most remarkable of land shells; it is reversed, and the columella forms a winding pillar, visible within, quite to the summit of the spire.* A small species found in the South Sea Islands is strung by the natives, and used for an ornament.

46. SUCCINEA. — *Shell* oval, small, thin and transparent; spire very short, consisting of only two or three whorls; right lip thin, left smooth; aperture large; no operculum. *Animal* like the *Helix*. — There are 10 species, recent.

The animal cannot entirely enter its shell; it lives upon herbs and shrubs on the borders of rivulets, and appears to be almost amphibious. It is found in England and other parts

* Sowerby .

of Europe, Van Diemen's Land, Guadaloupe, &c.

47. AURICULA. — *Shell* oval; spire short and obtuse; right lip in some species thickened, in others sharp; mouth toothed and oblong, rather in the form of an ear, often very narrow. *Animal* having a head furnished with two tentacula, and eyes at their base; foot short and narrow.—There are 30 species recent, and 15 fossil.

Several species are European; others are found on the banks of rivers in Brazil and Madagascar, and the Indian and American islands. The name is derived from a fancied resemblance to the ears of some animals: they are for the most part covered with an epidermis, but some are frequently delicately sculptured.

Montford proposed a new name for those shells which have slightly raised varices, and the mouth toothed on both sides. He calls them *Scarabus*. They appear to be found in marshy places; though one species, *S. umbrum*, is said to have been found on the tops of mountains by Captain Freycinet.

48. CYCLOSTOMA.—*Shell* either flattened or turreted; mouth circular; the margin of both lips united, and slightly turned back; closed by a shelly or horny operculum. *Animal*, head having two tentacula with eyes at the base; foot oblong.—45 species recent, and 10 fossil.

These shells vary much in form, some being very flat, others having the apex lengthened out, and being more conical; they all have the mouth round.

The *Cyclostomata* are found principally in the torrid zone; but one species, the *Cyclostoma elegans*, is a native of Britain: they inhabit woods, and are found amongst moss, stones, &c.

FAMILY 2.—*Lymnæidæ*.*

This family consists of amphibious animals with two tentacula, having eyes at the base. They inhabit fresh water, and rise to breathe

* *Les Lymnéens*, Lamarck.

the air on its surface. The shell is spiral, with the exception of one genus; the external surface generally smooth, and the right margin of the aperture always sharp and not reflected. They do not possess an operculum. There are four genera.

49. PLANORBIS.—*Shell* flat and disk-shaped; mouth semicircular, and removed from the axis of the shell; whorls apparent on both sides; umbilicus large; no operculum; many species reversed. *Animal* having two very long tentacula, with eyes at the base; foot oval and short.—22 species recent, and 9 fossil.

This genus inhabits lakes and ditches in every part of the world; many species are common in England, and one or two are found in America with thickened rims. Some of the species are so much flattened that it is difficult to distinguish the upper side from the lower; but Deshayes says it may generally be done by observing that the upper part of the lip is more advanced than the lower. They are usually thin and fragile.

50. *PHYSA*.—*Shell* oval or oblong, smooth and thin; aperture contracted above; inner lip smooth, outer very thin; no operculum; shell generally reversed. *Animal* having two long tentacula, with eyes at the base; foot long; mantle large, so as to cover part of the shell, and very transparent.—10 species recent, and 1 fossil.

These animals are most frequently found on the under side of the leaves of aquatic plants: they have a very singular way of adhering to the surface of the water with the shell downwards, and crawl in that direction with as much apparent ease as on a solid surface, and they will occasionally let themselves down gradually by a thread. This power of crawling under water against its surface is not wholly confined to this and the following genus. Many species are found in Europe, others in Africa, America, &c.

51. *LYMNEA*.—*Shell* oval, inflated, thin; spire small and pointed; aperture oblong, sometimes very large; occasionally a fold on the columellar lip; no operculum. *Animal*

having two triangular tentacula, with eyes at the base; foot oval and thin.—20 species recent, 12 fossil.

This genus is often united with the last, as they are similar in most respects, differing chiefly in the direction of their spires; (the *Physæ* being generally sinistral, and the *Lymnæ* dextral shells;) and also in the oblique plait on the columella. They are abundantly found in our rivers and ponds

(particularly the latter, as they appear to prefer stagnant water), and in other parts of Europe, Africa, and America: one or two species have been mentioned as natives of India. They feed on aquatic plants, and come to the surface of the water for air, like the preceding genera; the number of their eggs is very great, and they are deposited on stones, stems of vegetables, &c. in long masses enveloped in a glairy substance. Darwin mentions finding *Lymnæ* in great numbers in a lake near Rio de Janeiro, into which the inhabitants assured him the sea annually, and sometimes oftener, entered, and made the water quite salt.

52. *ANCYLUS*. — *Shell* oblong, small and transparent; apex rather pointed, and bent to the right, somewhat resembling a *Patella* in form; but the apex is posterior, whereas that of the *Patella* is anterior. *Animal*, head very large, having two tentacula, with eyes at the base; foot large.—5 species, recent and fossil.

The *Ancylus* is found only in fresh water, in ponds and brooks, where it attaches itself to flags and other aquatic plants, and also to stones. It swims about on the surface of the water, with the back downwards. It is nearly amphibious; for in summer, when the brooks are dried up, it remains in the soft mud until the return of rain. It is also often found out of the water, and only within reach of the spray of a waterfall.*

* Turton's Manual. In Lamarck's system this shell is placed with the *Calyptraciens*; but it is now ascertained that the animal greatly resembles the *Physa* in form and habits, and it has therefore been removed to the family now described.

FAMILY 3.—*Melanidæ*.*

The *Melanidæ* are all fluviatile shells, mostly exotic, and covered by a black or greenish brown epidermis. The right lip is always sharp, and the lips disunited. The *animal* has two tentacula and an operculum, and breathes only water. There are two genera.

53. MELANIA.—*Shell* oblong, often turreted, exterior wrinkled, and the whorls sometimes surmounted with spires or knobs; aperture pear-shaped; operculum horny; almost all have a brown or black epidermis. *Animal* having two tentacula, with eyes at the base; foot generally short and thick.—36 species recent, 7 fossil.

Found in a great many parts of the world; abundant in Asia, but rare in Europe †, except in a fossil state. They inhabit rivers.

* *Les Mélaniens*, Lamarck.

† *M. helvetica* is found in fresh waters in Switzerland and Italy.

54. MELANOPSIS. — *Shell* turreted; mouth nearly entire, having a slight notch at the base, and occasionally one also at the upper part of the lip; spire having from five to fifteen whorls; inner lip thickened at the upper part; operculum horny. *Animal*, head furnished with two tentacula, having eyes at the base; foot very short.—There are 9 species recent, and 7 fossil.

These shells are found in great abundance in the fresh waters of the south of Europe and Africa; also in Syria, and in the rivers Tigris and Ohio.

The genus *Pirena* of Lamarck is now suppressed, and the species placed with the above.

FAMILY 4. — *Peristomidæ*.*

The animal of this family is similar to that of the last; but the shell is conoid or discoidal, and the margins of the aperture united. All the four genera have opercula, and a thin brownish green epidermis. The animal respires water.

* *Les Péristomiens*, Lamarck.

55. VALVATA. — *Shell* cone-shaped, or flattened; whorls rounded and distinct; mouth round, or nearly so, with the lips united and acute; an operculum.

Animal having a very distinct head, prolonged into a kind of proboscis, and furnished with two very long tentacula, with eyes at the base; foot short. — There are 6 species recent, and 2 fossil.

Found only in rivulets and ponds in Europe and North America.

56. PALUDINA. — *Shell* cone-shaped, having the whorls rounded; aperture roundish, angulated above; margins of the inner and outer lip united; operculum horny; shell covered by an epidermis.

Animal, head furnished with a proboscis, and two tentacula having eyes at the base; foot rather triangular. — 21 species recent, and 13 fossil.

This genus is distributed all over the world, and is usually found in fresh waters, rivers and ponds; there are, however, some species found in salt marshes, but none in the sea.

57. AMPULLARIA. — *Shell* generally large, thin and globular; spire very short; mouth semi-circular, with the lips united, the outer being acute; operculum thin and horny, and rarely calcareous; the shell has an epidermis, and an umbilicus varying in size. *Animal* having a large head furnished with four tentacula, with eyes at the base of the two longest; foot oval and large. — 27 species recent, and 16 fossil.

The Mollusca forming this genus are found in the rivers of Africa, India, and South America, but have not been discovered in Europe or North America. They are found in rivers and ditches; one species, the *Ampullaria ovata*, inhabits the Lake Mareotis, where the water is salt, and another the brackish water near Rio de Janeiro.* Some of the African species have reversed shells, and all that are natives of Africa and America have the operculum horny: those, on the contrary, which come from India, have it usually shelly,

* Darwin.

and are furnished with an internal groove for its reception parallel to the mouth, which is thickened.*

The animals possess the peculiarity of a large bag opening beneath, and situated on one side of the respiratory organs; this they fill with water, by which means they can exist for a considerable period out of their natural element. Specimens have been brought from Egypt to Paris alive, although packed up in sawdust; and, as a further instance, the following may be adduced. A box sent to M. Caillaud, containing a number of river shells from the Nile, was delayed four months on its road, and when it reached Paris was in such a state, from the putrefaction of the greater part of its contents, that they were all thrown into water. A few hours after a great number of living *Ampullariæ*, which had been shut up in this mass of putrefaction, were discovered creeping quietly about, and some of them lived four and six months after. The Indian species lay globular eggs, of a pale green colour, about the size of small peas, which are found in

* Swainson, Zoological Illustrations.

clusters attached to sticks or other things in the water; when dry, they have a beautiful appearance.

M. Quoy has proposed to form two species of Lamarck's genus *Ampullaria* into a distinct genus, which he calls *Ampullacera*. The animal is totally different, though the shells greatly resemble each other.

58. AMPULLACERA (Quoy).—*Shell* rather thick, globular; mouth round or oblique, having the lips united; spire short; umbilicus very deep; operculum horny, thin, and having sometimes a projection. *Animal*, head large, flat, divided into two lobes, and having two sessile eyes; no appearance of tentacula; foot short and square.—2 species recent.

This mollusc is found in great abundance in New Zealand, living in pools of brackish water, and buried in sandy mud. It is eaten by the natives. When the animal is touched it enters very deeply into its shell, at all times is much hidden by it, and sometimes even fills the entrance with earth.

FAMILY 5. — *Neritidæ*.*

The shells constituting this family are chiefly distinguished by the smallness and flatness of the spire, which appears placed on one side; the mouth is generally semicircular, being half closed by a flat partition, which forms the left lip. Some are marine, and others fluviatile; they have all an operculum of a peculiar form, having a little projection which slips under the partition of the left lip, and thus securely closes the mouth. There are four genera.

59. NAVICELLA. — *Shell* oblong, convex above, with the apex straight and bent down to the edge, not spiral; inner lip thin, and assuming the form of a plate across one end of the aperture; shell altogether much resembling a *Patella*, but having an operculum which is shelly.

Animal, head very large, having two tentacula, the eyes placed on the summit of two

* *Les Neritacées*, Lamarck.

small protuberances at their base; foot large.—There are 3 species, recent.

These Mollusca are fluviatile, inhabiting the clear rivers of India, the Isle of France, &c.: they creep well on the rocks, and do not continue fixed to one spot, as the *Patella*. In the Isle of France they are made into broth for invalids.

60. NERITINA.—*Shell* thin, having no umbilicus, and being globular or oval; mouth semicircular; left lip flattened and sharp, right lip destitute of teeth.

Animal, head large, having two tentacula, with eyes at the base; foot short.—54 species recent, and 9 fossil.

The *Neritinæ* are fluviatile, and are found in the East and West Indies, the Isle of France, &c.; they are exceedingly pretty shells, the lines and marks, when looked at through a microscope, being often very beautiful. They are worn as ornaments by the Indians. The *Neritina corona* is coated with a black epidermis, and has a row of long sharp spines: the *Neritina viridis*, a very small green species, found in

the West Indies, is doubtful as regards its habits, some authors supposing it to be marine.* Many of the species are covered with an epidermis. The *N. fluviatilis* is found in the rivers of England adhering to stones. This shell is often covered with incrustations.

61. *NERITA*. — *Shell* thick, having no umbilicus; spire nearly hid; mouth semicircular; inner lip flattened, and frequently toothed, as well as the outer; operculum shelly. *Animal*, head furnished with two pointed tentacula having eyes at the base; foot large. Marine. — There are 27 species recent, and 10 fossil.

The shells of the *Nerita* are often beautifully marked and sculptured; they inhabit the Eastern and American seas, a few species only being European. The *Nerita peloronta* is called the Bleeding Tooth, from the red appearance of the teeth on the inner lip. Many of the species

* Gray says it is only found in the sea.

are very common in the West Indies, Moluccas, &c. Some naturalists unite this and the preceding genus, as their form is very similar; but Mr. Gray describes their opercula as quite different. That of the *Neritina* is solid, shelly, and furnished with a thin, flexible, outer edge; the *Neritæ* have a horny operculum, covered on both sides with a hard shelly coat, which may be separated from the horny part.

62. NATICA.—*Shell* smooth, or without an epidermis, having an umbilicus, which, however, is in some species covered by a coating of shelly matter; outer lip sharp, inner lip and inside smooth; operculum shelly. *Animal*, head very large, having two tentacula with eyes at the base; foot large and thin.—There are 45 species recent, all marine, and 11 fossil.

Found on the coasts of England, and also those of India, China, New Holland, &c.

FAMILY 6.—*Janthinidæ*.*

This family consists only of one genus. The shell is marine, and the animal breathes water only.

63. JANTHINA.—*Shell* round, whorls slightly angulated; right lip forming an angle at the lower part of the mouth with the columella, which is produced so as to meet it; very fragile, thin,

and of a purple colour. *Animal*, head furnished with a round proboscis armed with little hooks; two tentacula with eyes at the base; the under surface of the foot furnished with a vesicular or bubble-like apparatus, which prevents the animal from crawling, but enables it to float on the surface of the water with the shell downwards. — 2 species, recent.

One species, *J. fragilis*, is found on our coasts

* *Les Janthines*, Lamarck.

and those of Ireland, (in consequence of the gulf stream and westward winds,) and in the European seas; the other, *J. globosa*, is from Madagascar, the tropical seas, and the Mediterranean. They are seen floating on the sea, suspended by the foot, from which a violet-coloured liquid is emitted when the animal is touched or injured. The shell is so extremely brittle that it is seldom met with unbroken. Sir Everard Home states that the animal carries its eggs on its shell for some time.

FAMILY 7. — *Macrostomidæ*.*

The *Macrostomidæ* are of a very simple construction as far as regards the shell: the spire is small; so much so, indeed, that the mouth of the shell is nearly as large as the shell itself. They are usually highly iridescent in the inside, and have no operculum. They are all marine, and are divided into four genera.

64. SIGARETUS. — *Shell* ear-shaped, not

* *Les Macrostomes*, Lamarck.

pearly inside, nor perforated with holes; right lip thin. *Animal*, head large, having two tentacula, with eyes at the base; foot very large.—4 species, recent.

The few species that are known are brought from the Indian seas, the Atlantic Ocean, and the Mediterranean. The shell is almost hidden within the mantle of the animal, and from this cause, probably, is devoid of colour: some species are solid and thick, others thin.

M. Adanson found abundance of these shells separated from the animal in the sand at the mouth of the Niger.

65. STOMATELLA.—*Shell* ear-shaped, oblong; aperture large, and also oblong, not perforated; interior very pearly. *Animal* not known.—There are 5 species, recent.

They are brought from Java, India, New Holland, &c.

66. STOMATIA.—*Shell* not perforated; a rib or raised line on the back; spire rather more

prominent than the last genus; ear-shaped; mouth large, oblong, interior pearly. *Animal* not known. — There are 2 species, recent.

These shells are brought from the East Indies and New Holland.

This genus and the preceding are now generally placed together under the name of *Stomatia*. The shells are distinguished from those of the *Haliotis* by being destitute of holes, and from the genus *Sigaretus* by being pearly.

67. HALIOTIS. — *Shell* flat, pearly inside, slightly spiral at one end, and having one side

perforated with small holes; aperture almost as large as the shell. *Animal*, head large, having two long round tentacula, with eyes at the base on footstalks; foot very large, having

the margin fringed all round, resembling a double row of festoons. — 15 species, recent.

The form of this shell somewhat resembles that of the human ear, and it is from that circumstance frequently called the Ear-shell; one species, from its length, having received the name of the Ass's Ear. The outside is generally rough, worn, or covered with marine substances; the inside presents the same enamelled appearance as mother-of-pearl, and exhibits the most beautiful colours. The holes with which the shell is perforated are for the passage of the lobes of the animal's mantle, and are made at regular intervals, as it increases in size. It is observable, however, that as each new one is formed the one nearest the spire is closed up; so that there are generally the same number open at a time, and seldom more than seven or eight. The annexed cut is a representation of one of the species of this curious genus, with its living inhabitant, which, notwithstanding its apparent size, can be covered by the shell. In its state of rest, it is able to adhere with such tenacity to the substance it is fixed upon as to be removed with

very great difficulty, although it can detach itself with ease. It is always found near the

surface of the water, and on serene nights feeds on the vegetation which springs up on the shore.

The *Halotis* is found in New Holland, China, the Cape of Good Hope, the Moluccas, &c.; but the most common species which is preserved in conchological collections is the *Halotis tuberculata*, represented in the last figure, which is found in great abundance on the coasts of Guernsey and Jersey, where the animal is used for food; and the shells are frequently arranged in patterns on the outside of the cottages, being cemented into the mortar, with the inside exposed: these have

a very beautiful appearance when the sun shines upon them. They are occasionally found upon our coasts; but as it is only after violent storms, they are supposed to have been brought from other shores.

In the islands of the South Seas these beautiful shells are used to ornament the boats of the natives, and to make a kind of musical instrument. They also form the thick left lip of the aperture into fish-hooks. The natives of New Zealand use the shell as spoons. Pearls are sometimes found in them.

FAMILY 8.—*Plicacidæ*. *

This family of the *Trachelipoda* is distinguished by having the pillar plaited, which gives the shells an appearance of having the mouth toothed. There are only three genera, all marine.

68. TORNATELLA.—*Shell* small, and oval; spire small, pointed, and usually striped transversely; pillar having one or two thick plaits; outer lip thin; no epidermis. *Animal* not known.—6 species, recent.

* *Les Plicacées*, Lamarck.

These shells are found on the shores of the Indian Ocean and Senegal; and one species, *Tornatella fasciata*, inhabits our own coasts. They form an operculum.

69. MELAMPUS (Montford).—*Shell* small, conical; spire depressed; pillar or inner lip having one or two plaits, outer lip thickened inside; no operculum. *Animal* having two tentacula, with eyes at the base.—*Species doubtful.*

This genus inhabits the shallow parts of the coast of the West Indian Isles, where it is found creeping on the mangrove roots. It is perfectly amphibious, and very tenacious of life.

70. PYRAMIDELLA.—*Shell* turreted; pillar having three plaits; outer lip thin; no epidermis. *Animal* not known.—There are 11 species recent, and 8 fossil.

This group of shells is found in the Indian and American seas.

FAMILY 9. — *Scalaridæ*.*

The shells forming this family are spiral, but the whorls of some of the species do not wrap over each other so decidedly as in many others; indeed, in many species, they do not touch. Modern authors introduce a few new genera. The third and fourth mentioned here have been placed in many different parts of the system. Lamarck mentions the first amongst his *Annelides*.† Rang says, in his *Manuel des Mollusques*, that when he was seeking the animal of the *Magilus* in India he was struck with its analogy to *Vermetus*. It will therefore be placed near that genus, as its long tube (caused by the circumstances hereafter mentioned) in some degree resembles the loosely twisted whorls of that shell, and the young student will on this account find the *Magilus* placed amongst those which resemble it in form. The family contains six genera.

* *Les Scalariens*, Lamarck.

† See page 36.

71. VERMETUS. — *Shell* thin, tubular, only slightly twisted; mouth round; lips united. *Animal*, head indistinct, having two tentacula with eyes at the base; foot cylindrical. — 1 species.

This is a very remarkable shell, both in appearance and habits. The spire is extremely irregular in formation, with the whorls perfectly separated from each other; and the animal, although of the tribe *Trachelipoda*, with a foot adapted for creeping, continues attached to one spot by the pointed extremity of the spire. They may be found in groups, twisted together in great numbers, in the seas near Senegal. There is only one species, the *Vermetus lumbricalis*.

72. MAGILUS. — *Shell* ventricose; spire short, consisting of three or four whorls; aperture longer than wide, without any notch, but an angle at the base. The last whorl, abandoning altogether the spiral form, produces a long tube, which is compressed laterally, especially on the side of the base of the shell; aperture elliptical. *Animal* but little known;

an operculum.* 1 species, recent.

When in the young state this curious shell presents all the characters of a regular spiral univalve. The animal establishes itself in the excavations of madrepores; and as the coral increases around it, the *Magilus* is obliged, in order to have its aperture on a level with the surrounding surface, or near it, to construct a tube, the growth of the coral determining its length. As this tube goes on increasing, the animal abandons the spiral for the tubular part of the shell; and in the operation it leaves behind no partitions, but secretes a compact calcareous matter, which reaches to the very summit of the spiral part; so that in an old specimen the posterior part of

* Dr. Rüppell.

the shell presents a solid mass. One species only, *Magilus antiquus*, is known. The colour is white, more or less pure.

73. LEPTOCONCHUS (*Rüppell*). — *Shell* subglobular, delicate, fragile, translucent; spire low, and nearly effaced by the encroachment of the last whorl; aperture large, and in form somewhat like the letter S reversed; the two margins not united; the right delicate at all ages, and a little expanded; no columella or umbilicus. *Animal*, head with a proboscis; two tentacula, with eyes in the middle; foot moderate in size; no operculum. — 1 species.

The shell of this genus is of a dirty milk-white, and is furrowed externally. It is found in the Red Sea, where it is embedded in calcareous masses of *Polyparia*, and having no communication with the water except by a moderate opening.*

74. SCALARIA. — *Shell* turreted, with or

* The descriptions of the two last genera are taken from the Penny Cyclopædia.

without a pillar ; mouth nearly round, lips united ; longitudinal bands the whole length of the shell ; operculum horny. *Animal* having two tentacula, with eyes at the base ; foot short and oval. — 7 species recent, and 5 fossil.

In these shells it is easy to distinguish the different periods of growth, as the mouth is always bordered by a thick lip, which is retained, and produces a band ; in the adult shell these are very numerous, and from their appearance have given the name to the genus, of *Scalaria*, or *Staircase*. The seas of China and Japan produce the rare species *Scalaria pretiosa*, for fine specimens of which, $3\frac{1}{2}$ inches in length, formerly the enormous price of one hundred guineas has been given ; they are now, however, much more frequently met with, and may be obtained for a few shillings. In this species the whorls only touch each other at the thickened lip, and occasionally a specimen is found in which they are perfectly separated. The *Scalaria communis* inhabits our own coasts.

The following genus, described both by *Deshayes* and *Rang*, will be introduced.

75. *RISOA* (*Fréminville*). — *Shell* oblong, turreted; no umbilicus; generally remarkable for its longitudinal bands; mouth oval, oblique; the two lips united, or nearly so; the right not reflected, but thickened; a calcareous or horny operculum. *Animal*, foot short and round.—44 species, recent and fossil.

The species of this small shell are brought from the Mediterranean, the seas of Senegal, and the European Ocean. They are generally found on sandy shores.

76. *DELPHINULA*. — *Shell* very thick; spire depressed; umbilicus large; exterior rough, with tubercles; mouth pearly; lips united; operculum shelly.

Animal, head distinct, having two tentacula, with eyes at the base; foot short and round.—3 species recent, and 10 fossil.

All the species forming this genus are found in the Indian Ocean. Some of them have the whorls entirely separated.

FAMILY 10. — *Turbinidæ*.*

This family constitutes the last of the *Phytophagous Trachelipoda*. The animals breathe only water, and all are confined to the sea. The shell is turreted, conoid, or pyramidal; the aperture round or oblong; and they appear all provided with an operculum. The three first genera are more pyramidal than the rest, and will stand upright on their base, with the axis of the shell scarcely, if at all, inclined. There are eight genera.

77. SOLARIUM. — *Shell* nearly discoid, not perlaceous; internal edge of the whorls visible

in the umbilicus, which is very wide; mouth almost square; margin thin. *Animal* resembles that of the *Trochus*. — There are 7 species recent, and 11 fossil.

The various species of this genus are brought

* *Les Turbinacées*, Lamarck.

from the coasts of New Holland, Tranquebar, and from the Indian and Chinese seas; they also frequent the shores of the Mediterranean, near Alexandria. The *Solarium perspectivum*, represented in the figure, is frequently called the Staircase-shell. Specimens have been found three inches in diameter; and as they are elegantly marked with black and straw colour, they are often very beautiful objects.

78. ROTELLA.—*Shell* small, outside smooth and polished; left lip very thick, and spreading over the under surface so as to form a callosity; right lip sharp at the edge; operculum horny. *Animal*, head having two very long and pointed tentacula, with eyes at the base; foot short.—5 species, recent.

From the Mediterranean, Indian, and Javanese seas.

79. TROCHUS.—*Shell* conical; spire elevated; circumference of the whorls more or less angulated, often quite sharp; mouth transversely depressed; edge of the outer lip disunited from the inner at the top; inner lip curved, rather oblique at the base; sometimes

an umbilicus; operculum generally horny, thin, and spiral, with many whorls. *Animal*, head

distinct, having two tentacula with eyes at the base; foot short and round.—There are 69 species recent, 9 fossil.

This genus is distributed all over the world, and presents a great many species. The *Trochus agglutinans**, from the Antilles, is remarkable for the singular habit of accumulating, during its formation, different substances which adhere to its shell. It is often found covered with stones, corals, small shells, and fragments of all kinds, and on account of this curious propensity it has been named the *Carrier*: it is supposed that it does this for the purpose of strengthening its shell, which is very thin and brittle, and that it causes the substances it collects to

* This species of *Trochus* has been formed into a genus by Montfort, and called *Phorus*.

adhere to it by bringing its shell into contact with them whilst the shelly matter is still in a partially liquid state. Mr. Swainson mentions* that amongst some fresh-water shells he received from Brazil he found two small ones resembling a *Helix*, and was led to examine them,

as their outside was covered with particles of dirt and sand. On placing them, however, under the magnifier, he was surprised to find

that the shells were actually composed of little stones and grains of sand only, agglutinated together with so much skill by the animal, that the regular turns of the spire and the form of the umbilicus were most accurately preserved. They were, in short, fresh-water Carriers. The *Trochus longispina*, from India, has the circumference ornamented by a row of long spines, of a silvery or gold colour, placed at regular distances, and somewhat resembling the rays of the sun as they are represented in carved work; the shell itself is also of a golden colour. The *Trochus jujubinus*, found in the seas of the Isle of France, is remarkable for its peculiar colour-

* Treatise on Shells and Shell-Fish.

ing, the upper whorls being blackish, whilst the apex and two last are red or flesh-coloured. Many species are very iridescent at the mouth and under the epidermis. There are some species common on our coasts, but none attain the beauty of the foreign varieties.

80. MONODONTA. — *Shell* oval or pyramidal; lips disunited at the upper part, the right being occasionally striped within, and the left having a tooth-like process, whence the name is derived; operculum horny and thin.

Animal, head distinct, having two tentacula, with eyes at the base; foot short. — 25 species, recent.

These shells are found in most seas; they seem to form a connecting link between the *Trochus* and the next genus, which they much resemble.

81. TURBO. — *Shell* rather turreted, base not flattened; mouth round; lips not united; outer lip thin; an operculum, shelly and solid. *Animal*, head having two pointed tentacula, with eyes at the base; foot short. — 34 species recent, and 4 fossil.

The shells of this genus, if placed upon their mouths, will stand steadily in that position, with the axis very much inclined. They are brought from China, India, Africa, &c. The *Turbo litoreus**, or common periwinkle, is used by mankind as an article of food, and is found on the shores of England in great numbers. In Sweden, where they also abound, they serve to prognosticate the approaching state of the weather; the peasants having observed that whenever the periwinkles ascend the rocks it is a sure sign of a storm being near, as their instinct teaches them to place themselves out of the reach of the dashing of the waves; on the contrary, when they make a descent upon the sand it is an indication of a calm. In hot countries some species are often seen on the trees near the coast, and on the rocks elevated above the surface of the water; they remain stationary on the latter during the hottest

* Férussac has introduced a new genus, of which the *T. litoreus* is a species. He names it *Littorina*. The shells are known from *Turbo* and *Phasianella* by having a horny operculum, and from *Trochus* by having few whorls. The shells are found on the shore, whence the name.

hours, even when it is painful to walk on them from their great heat: they leave the water early in the morning, but return at night. These circumstances prove that, although marine, many species are amphibious. These shells are often highly iridescent; and the mouth in some species, as the *Turbo chrysotomus*, is of a deep and beautiful golden colour.

82. PLANAXIS. — *Shell* small and oval; mouth of an oval shape, and having a half-formed notch; left lip flattened, right lip rayed or slightly toothed in the interior; an operculum. *Animal* supposed to be like that of the *Phasianella*. — There are only 2 species, recent.

These Mollusca are found in India, South America, and the Isle of France, where they abound in great numbers, sometimes almost covering the shores in particular spots; they are often found under stones. The shells are usually transversely grooved on the exterior.*

83. PHASIANELLA. — *Shell* oval, solid; mouth

* Some authors remove this genus to a Zoophagous family, but Rang says that the animals resemble the *Phasianella*.

oval, long, not wide; lips disunited at the upper part; right lip sharp, left smooth; the last whorl much larger than the others; a shelly operculum. *Animal*, head having two long and round tentacula, with eyes on two footstalks; foot oblong.—10 species recent, and 2 fossil.

Some species are beautifully coloured, smooth, and shining, and were formerly rare. The greater number have no epidermis. They are found in South America, India, New Holland, the Mediterranean, &c.

84. TURRITELLA.—*Shell* very long, needle-shaped, with numerous whorls, usually trans-

versely striped; no umbilicus; mouth round; lips disunited at the upper part; the right lip has a sinus or indentation about the centre. *Animal* with two tentacula, long and very fine

at the extremity, and having eyes at the base; an operculum.—13 species recent, and 12 fossil.

These shells are found in India, Africa, and Europe; and many of the foreign species attain a large size, some having been found four inches and a half long. They are all marine, more or less striated; but none of the species known possess vertical ribs, thickened bands, or tubercles. The right lip is seldom perfect, and the sinus therefore not always visible.

DIVISION II.—ZOOPHAGA.

The *Zoophaga*, or carnivorous *Trachelipods*, are divided into five families:—*Canaliferidæ*, *Pteridæ*, *Purpuridæ*, *Columellidæ*, and *Convolutidæ*. They all form the mouth of the shell with a notch or canal at the lower extremity, for the convenience of protruding a retractile trunk, with which they pierce the substance of other shells, and suck out the contents. It is this formation of the mouth which distinguishes the shells from those of the herbivorous members of the order, it being in the

latter entire. The *Zoophaga* are all marine, and all appear to have two tentacula; they are predaceous, as their name implies, and feed for the most part on other shell-fish, either living or in a decayed state.

FAMILY 11.—*Canaliferidæ*.*

The animals of this family breathe water only; they feed on animal substances by the help of their retractile trunk, and are confined entirely to the sea. The shells are spiral, enclosing the animal, the aperture having a canal either long or short. All have opercula. The family is divided into two parts: in the first the periodical growth of the shell is but slightly marked; in the second it is distinguished by thickened bands or varices, which remain on the external whorls. Seven genera belong to the first division, and four to the last.

DIVISION I.

85. CERITHIUM.—*Shell* turreted, and much elongated; mouth oblong and oblique, having

* *Les Canalifères*, Lamarck.

a short channel at the base, and a notch at the upper part of the right lip; operculum small and

norny. *Animal*, head furnished with a proboscis, and having two tentacula distant from each other, and eyes in the middle; foot short and oval.—87 species recent, and 60 fossil.*

The Mollusca forming these shells chiefly inhabit the Indian and Pacific Oceans, the Red Sea, &c.; they appear to have the power of remaining for a considerable period out of their natural element, and without food, as has been proved by the following fact:—Two specimens belonging to one of the species of this shell were brought from the Mauritius, and contained the animals, which had drawn themselves far within the shell: they remained in

* Or 220 (Deshayes).

that state during the voyage; but on arriving in England and being placed in sea water, immediately recovered from their inanimate state, apparently uninjured.

The *Cerithium giganteum* is more than a foot in length, and the only specimen which is in existence is in the cabinet of Lamarck*; it was fished up by an English sailor in the neighbourhood of New Holland, and is, of course, a recent shell. There are fossil specimens of this species, even larger than the recent *giganteum*, found at Grignon near Paris.

The surface of these shells is rarely smooth, being generally covered with spines, prickles, &c., or striped. The regularity of these projecting parts might furnish a sculptor with models for innumerable designs in ornamental architecture. They inhabit the sea; but such as have the canal straight live in salt marshes, or the mouths of rivers.

86. PLEUROTOMA. — *Shell* turreted or fusi-form; canal straight and long; right lip having a slit at the upper part, left lip smooth; oper-

* Now that of the Duc de Rivoli.

culum small and horny. *Animal* very little known. — 23 species recent, and 30 fossil.

This genus is found in the eastern and tropical seas, many species being abundant at Java. The *Pleurotoma buccinoides* has no canal; but it has a notch, and also possesses the slit in the right lip. *P. gracilis* is found on the British shores.

87. TURBINELLA. — Shell large, heavy, and rather fusiform; spire short; mouth oval, having a long and straight channel; left lip with from three to five distinct plaits, which are at right angles with the axis of the shell; operculum doubtful. *Animal* imperfectly known. — 23 species, recent.

Many of these shells are exceedingly heavy. They are found in the Indian and African Oceans. A reversed example of this genus was formerly

in the collection of Lord Tankerville, specimens of which are extremely rare and valuable.

88. CANCELLARIA. — *Shell* oval or turreted; spire slightly elevated, and pointed; mouth oval, having either a very short canal or a notch only; right lip sharp, and wrinkled within; left lip with several irregular plaits; operculum horny. *Animal* doubtful. — 12 species recent, and 7 fossil.

These shells are generally found on the sands of the shores washed by the Indian and African seas, and on the coast of America: they are rare, but not remarkable; and are usually rough to the touch, and striped.

89. FASCIOLARIA. — *Shell* fusiform, and not very thick; spire of a moderate length; mouth oval; canal rather long, sometimes slightly bent; right lip sharp, and often wrinkled inside; left lip having two or three oblique plaits; operculum doubtful. *Animal* unknown. — 8 species, recent.

From the Indian seas, the Antilles, and the Isle of France, and one species from the Mediterranean. They are not numerous, but the *Fasciolaria distans* is very beautiful.

90. FUSUS. — *Shell* fusiform or spindle-shaped, the channel and spire being both considerably elongated; left lip smooth; operculum horny. *Animal* unknown. — 37 species recent, and 36 fossil.

The localities of this genus are the Indian, Northern, and Atlantic Oceans, the Red Sea, &c. Many of the species are transversely striped, and are covered with an epidermis; on the removal of which they present very beautiful colours. The *Fusus colosseus* is often more than eleven inches long, and other species are five or six.

91. PYRULA. — *Shell* pear or fig-shaped; spire short, and sometimes flattened; mouth oval; left lip smooth; canal straight, sometimes narrow at the extremity, sometimes short and

wide; operculum horny. *Animal* unknown. — 28 species recent, and 6 fossil.

A common and numerous genus, chiefly from the Indian Ocean and Red Sea. The *Pyrula canaliculata* is found in the Northern Ocean; it is a large but light shell, six inches in length. The *Pyrula candelabrum* has the spire quite flat, and if placed upon that part will stand upright; it is very rare. The *Pyrula ficus* very nearly resembles a fig in shape, from whence its name is derived. Some of the species have an umbilicus. In the British Museum is a specimen of a *Pyrula bezoar* that appears to have grown with perfect regularity until the formation of its last half whorl, which is thrown considerably more than half an inch out of its proper position by a group of barnacles. These shells had probably attached themselves to the back of the *Pyrula* at an earlier stage, and as the latter increased in size at length filled the place that should have been occupied by the inner lip, which, on meeting with this interruption, di-

verged from its course, and was thrown over the barnacles. Had the shell not been taken until a later period, there can be little doubt that the animal would have at length destroyed the barnacles, and completely hidden them from view, although it would appear that it had not the power to remove them by absorption while they retained their vitality.*

DIVISION II.

92. STRUTHIOLARIA.—*Shell* oval; spire elevated; mouth oval, terminated by a very short straight canal; right lip thickened, left also thickened and spread out; no varices; operculum horny. *Animal* unknown. — 2 species, recent.

These shells are rare and singular; they are in shape like a *Buccinum*, but differ in having a thickened lip: they come from New Holland and New Zealand. Sowerby observes that the thickened aperture appears only to exist in adult specimens of this genus. The usual length of the shell is about two

* Gray.

inches, but some writers speak of specimens measuring four inches.

93. RANELLA.—*Shell* oval, rather flattened; canal short; two rows of varices situated at the distance of half a whorl from each other, and longitudinally united, forming a continued ridge on each side of the shell; a sinus or gutter at the upper part of the mouth. *Animal* and operculum unknown.—14 species recent, and 1 fossil.

From the Indian and American Oceans, and the Mediterranean Sea.

94. MUREX. — *Shell* oblong; spire short;

channel generally long, and sometimes closed:

varices three or more, spiny or rough, and connected; mouth round and small; operculum horny. *Animal*, head having two long tentacula furnished with eyes; foot round, and generally short.—66 species recent, and 15 fossil.

These shells are found in nearly every part of the world, and are remarkably curious: the finest species come from tropical countries. The *Murex tenuispina*, or Venus's Comb, as it is called, is entirely beset with long sharp spines, which the animal has the power of dissolving and replacing by a smooth and even surface, whenever it finds it necessary, in enlarging its shell, to remove them. This faculty is also possessed by other species, but it is very conspicuous in the *tenuispina*. In specimens in which the process is going on, there may be observed a notch formed by this means in the base of the spines, the completion of which eventually causes them to separate from the shell.*

95. TRITON.—*Shell* oblong; canal short; varices placed alternately on each whorl, often considerably distant from each other, and never

* Gray.

longitudinally united; right lip often wrinkled, and left occasionally thickened; operculum horny. *Animal* like the *Murex*.

—51 species recent, and 3 fossil.

This genus is found in the Mediterranean, Indian, and South Seas: in many of the countries near which it abounds it is used as an instrument of music. By breaking off the apex, or boring a hole in it, and blowing through the aperture, applied to the mouth of the performer, notes can be produced; and it is employed in this manner by some rude nations as a military horn. The *Triton variegatum* is often met with fifteen inches, or even two feet, in length: it is from the Asiatic seas, and is frequently beautifully coloured.

FAMILY 12.—*Pteridæ*.*

The shells of this family are frequently called Wing-shells, from the circumstance of the last whorl expanding more or less into the form of a wing when the animal has attained maturity;

* *Les Ailées*, Lamarck.

the adult shape of the shell is consequently different from its immature and younger form. There are three genera, all channelled.

96. ROSTELLARIA. — *Shell* rather long, the spire and channel being both elongated; right

lip dilating with age and slightly digitated, one digitation being carried near the apex of the spire and attached to the whorls; sinus near the base; operculum horny, and very small. *Animal* imperfectly known: — 3 species recent, 3 fossil.

These shells are more fusiform or turreted than the other genera of this family; they are found in the seas of China, the Moluccas, and Europe. The *R. rectirostis* is a very rare shell. *R. pes-pellicani* is one of the most common of the

genus, and is found plentifully in the Frith of Forth.

- Q 97. PTEROCERAS. — *Shell* oblong; spire small; mouth terminated by a rather long canal; right lip dilated in the adult into several claws, and having a sinus near the canal; operculum horny. *Animal* like the *Strombus*. — 7 species, recent.

The number of claws varies in different species; the *Pteroceras millipeda* has a great many, but they are small; in others they are straight, or nearly so, and smooth; whilst many of them have these appendages very much curved.

There is much singularity in the growth of these shells, and the appearance they present at various periods is strikingly different. In the young state of the animal the shell has no claws, and these gradually make their appearance, at first in the form of short and open canals, which by degrees assume the length and curve of the adult and completed shell, and eventually

are closed up with shelly matter and become solidified.

Nearly all the species are inhabitants of the Indian Ocean, and many of them exhibit the most beautiful colours on their internal surface.

98. STROMBUS. — *Shell* thick and oval; spire moderate; mouth long, and rather narrow, terminated by a canal more or less long and recurved; right lip dilated in the adult, and having a small notch or sinus near the canal; left lip sometimes thickened; operculum horny, long and narrow. *Animal*, head furnished with a proboscis and two tentacula, which are short; the eyes are situated on footstalks longer than the tentacula: foot small. — 32 species recent, and 1 fossil.

These shells are brought from India, Africa, and other hot countries. Many of them attain a large size; and the *Strombus gigas* is used for the purpose of paving the streets at Santa Cruz, and elsewhere. The spire is in many species quite hidden by the expansion of the

outer lip; the latter is sometimes reflected inwards towards the middle part.

The animals of the *Strombus* occasionally produce pearls. Mr. Wood, in his "Zoography," relates that he saw a pink pearl which was taken from the body of the animal of the *Strombus gigas*, which is caught for the table off the island of Barbadoes. The pearl was discovered by chance, whilst the men were employed in cleaning the fish. Its weight was 24 grains; but it would have been more valuable if it had been round. The same author states that only four of these pearls had been discovered in the vast numbers of fish that are annually brought to market in that part of the world, though he has reason to believe that this is in some measure owing to the carelessness of the negroes. This pearl was exactly of the same colour as the interior coat of the shell.*

FAMILY 13. — *Purpuridæ*.†

This family has the shell either notched at the base, or terminated by a short canal turned

* Penny Cyclopædia, article *Strombidæ*.

† *Les purpurifères*, Lamarck.

towards the back of the shell. Some of the genera have plaits or marks of various kinds on the inner lip; but the form of the shells is different from that of the *Columellidæ* family (which has the lip also plaited, but more distinctly), being much more globular.

The *Purpuridæ* derive their name from the circumstance of some of the genera, especially the *Purpura*, possessing a colouring matter, which was made use of by the Romans to produce their beautiful purple dye; their use has, however, now been discontinued, the cochineal insect having the same effect, and being more readily obtained. There are thirteen genera in this family, and all seem to possess an operculum.

99. CASSIDARIA.—*Shell* rather oblong; spire not much elevated; mouth long, and rather narrow, terminating in a canal, which is recurved, but does not touch the back; left lip often rough or wrinkled, and detached from the pillar; right lip thickened; exterior of shell transversely

grooved, and often spotted with round tubercles; operculum horny. *Animal* like that of the *Cassis*.—5 species recent, and 2 fossil.

These shells chiefly come from the seas of warm latitudes; one or two species are found in European countries.

100. ONISCIA (Sowerby).—*Shell* oblong, rather obtuse; spire short, base rather pointed; mouth longitudinal; canal very short; outer lip thickened, toothed within; inner lip expanded and covered with raised spots; outside ribbed. *Animal* not known.—3 species recent, and 1 fossil.

It is litoral in its habits, and is found in sand in the South Seas.

101. CASSIS.—*Shell* large and heavy; spire short; mouth either long and narrow, or rather oval, terminated by a short canal abruptly curved towards the back of the shell; both lips generally toothed and flattened; operculum horny. *Animal*, head large, with a very short proboscis, and two tentacula

with eyes in the middle; foot large. — 25 species recent, and 1 fossil.

These animals sometimes grow to a very large size, requiring of course a corresponding magnitude of shell. They are chiefly brought from the Indian seas, but a few species are found in the Mediterranean; they live at some distance from the shore on the sand, into which they occasionally burrow, and entirely hide themselves. In some species the columellar lip projects above the spire, giving a flat appearance to the lower part of the shell.

The shells of the *Cassis rufa* and other species are exquisitely sculptured by Italian artists in imitation of antique cameos, the different strata of colouring matter resembling those of the onyx and other precious stones. Of these a great variety of ornaments are made, and of late years a considerable trade has been carried on in them on the Continent. The substance of shells seems to be formed of three plates, the central generally being the thickest, and the outer one the thinnest. The Italian cameo cutters appear to be aware of this circumstance, and avail themselves of it in cutting the

cameos, the ground being always formed of the innermost layer of the three, which is also generally the most transparent.*

102. RICINULA.—*Shell* oval, generally covered externally with spines; mouth notched and narrow, strongly toothed on each lip, by which the aperture is contracted; spire very short; operculum horny. *Animal*, head large; proboscis short; two tentacula with eyes in the middle; foot rather large. — 9 species, recent.

These shells are usually small, and when young have no traces of teeth: they are all natives of foreign countries, and are found principally in the Indian seas on coral reefs and rocks. The *Ricinula horrida* is a remarkable species, being whitish, and covered with black pointed knobs; the mouth is full of teeth, and is tinged with a violet colour.

103. PURPURA.—*Shell* oval, thick, either smooth or tuberculated; spire short; mouth large and notched; left lip flattened; operculum

* Gray, Philosophical Trans.

horny. *Animal*, head large; proboscis either short or altogether wanting; two tentacula, having eyes in the middle; foot large. — 50 species, 1 of which is also fossil.

The *Purpura* is found most abundantly in warm seas, where the shells attain the largest size, a few only of the species being met with in Europe; they are chiefly litoral, or found on and near the sea coast. It was from the animal of this genus that the Roman purple dye was obtained. *P. patula* is supposed to be the species; it is found in the Atlantic Ocean and Mediterranean Sea. The species *P. lapillus*, found on the British coasts, affords a small quantity of this colouring matter.

104. MONOCERAS. — *Shell* oval; mouth rather long and notched; a sharp projecting tooth at the base of the right lip; left lip generally flattened. *Animal* like the *Purpura*. — 5 species, recent.

Found in the seas of southern

latitudes, particularly those of America; like the last genus they frequent the rocks.

105. CONCHOLEPAS. — *Shell* thick and rough; apex inclined towards the left lip; mouth very large, with a slight notch at the base; right lip having two teeth at the base; an operculum.

Animal like the *Purpura*. — Only 1 species, recent.

Of this singular genus the only species known is very abundant on the coasts of Peru and Chili, and is known by the name of *Concholepas Peruvianus*. It often grows to a large size, and although not beautiful is a curious shell. In its shape it has more analogy with the *Patella* family than any other; but from the different form of the animal, and the circumstance of its constructing a large operculum, it must be placed amongst the *Purpuridæ*. The shell is two or three inches in length.

106. HARPA. — *Shell* oval, smooth, and marked with parallel longitudinal ribs, the up-

per end of each of which, near the sutures, is projecting and pointed; spire short, last whorl large and deeply notched; outer lip thickened:

it is supposed to have no operculum. *Animal*, head large; mouth open below; destitute of a proboscis; but having two tentacula, with eyes in the middle; foot large.—8 species recent, and 1 fossil.

The principal localities of this genus are the Red Sea and the Indian and South American Oceans. The *Harpa imperialis*, from the Mauritius, is a remarkably beautiful shell, and exceedingly rare: at one time it could only be procured at an extravagant price, but now it is more common. It is distinguished by the very great number of ribs, which, indeed, are almost close together; and the colours and markings are very elegant and beautiful. There is a fishery for them at the Mauritius and the

neighbouring islands, and they are taken at night and at sunrise, when they are probably feeding. Some naturalists* affirm that the animal can, when attacked by an enemy, disembarass itself of part of its foot, and retire entirely within its shell. The ribs in this genus are evidently the remains of what has been the mouth of the shell at different periods. The *H. ventricosa* is the most common species of this genus, and perhaps the most beautiful.

107. DOLIUM. — *Shell* large, light, globular; mouth wide and notched, generally transversely banded, and the right lip in consequence waved; operculum horny. *Animal*, head large, having a short proboscis, and two tentacula with eyes in the middle; foot rather large. — 7 species, recent.

These shells are found, for the most part, in the seas of the Indies, Africa, and South America: one species, however, the *Dolium galea*, is met with in the Mediterranean. They are

* MM. Quoy and Gaimard, and M. Reynaud,

often found on reefs, and sometimes attain a very large size, specimens having been seen as large as a man's head ; even, then, however, they retain the fragility and thinness of texture which characterise most of the species.

108. **BUCCINUM.**—*Shell* rather oval ; spire and mouth equal in length ; inner lip not flat, but swollen at the upper part ; mouth notched ; shell slightly covered by an epidermis ; operculum horny. *Animal*, head furnished with a proboscis, and two tentacula having eyes in the middle ; foot short.—140 species recent, 5 fossil.

These shells are found in all parts of the world* ; and in many places, particularly in northern countries, the animals of the *Buccinum undatum* (under the name of *whelks*) form an article of food. The shells are not remarkable for brilliant colours ; but their great diversity of form, delicacy of sculpture, and pencilled

* Captain Parry found two species, *B. glaciale* and *B. Sabinii*, within the arctic circle.

markings, render them interesting. They never attain a large size, and the greater number are litoral shells. They are sometimes sinistral.

109. *NASSA*.—*Shell* globular or oval, according to the spire, which is sometimes short, sometimes rather long; mouth oblong, notched; inner lip thickened, and spread out, occasionally very large; right lip often wrinkled; operculum horny. *Animal*, head large; proboscis short, or altogether wanting; two tentacula, with eyes in the middle; foot very large.—9 species recent, and 2 fossil.

They abound in the South of Europe and the East Indies; and some are also found on our coasts. They are occasionally seen on the sand or reefs, feeding on the *Mactra*, which they effect by piercing the shell with their proboscis, making an extremely regular round hole, and extracting the contents through the aperture thus formed.

110. *EBURNA*.—*Shell* oval, smooth; mouth oblong and notched; left lip thickened, right thin; operculum horny; and an umbilicus. *Ani-*

mal, head furnished with a proboscis, and two tentacula having eyes in the middle; foot short. — 5 species, recent.

The shells of this genus are found in the Indian and Chinese seas. The *Eburna Zeylanica* is remarkable for the little spines with which the umbilicus is filled, and this peculiarity distinguishes it from the other species which are destitute of them. The *Eburna glabrata* very closely resembles the *Ancillaria**, in having the spire exceedingly smooth, and the sutures not in the least channelled; it differs, however, in having an umbilicus.

111. TEREBRA. — *Shell* long, turreted, and slender; spire pointed; whorls often strongly marked at the upper part, resembling a double suture; mouth oval and strongly notched; no epidermis nor operculum. *Animal*, head not furnished with a proboscis, but fringed, and having two tentacula with eyes at the base; foot oval. — 24 species recent, 2 fossil.

* It is now placed in that genus.

These shells are often called Needle-shells, on account of their sharp, lengthened, and spiral form; and although by no means rare, are often elegantly marked. Some of them exceed ten inches in length. The notched mouth distinguishes them from the *Turritella*, which they much resemble in shape. Very few of the species are European, the greater number inhabiting the warmer latitudes of the East and West Indies and Africa. They occasionally creep out of the water, still keeping, however, within reach of the spray.

FAMILY 14. — *Columellidæ*.*

The shells composing this family have no canal at the base of the aperture, but a notch, more or less distinct, and plaits on the columella or left lip. The beauty of many individuals of this family makes it interesting. There are five genera.

112. COLUMBELLA.—*Shell* oval; spire short; plaits small; right lip thickened, turned inwards, and swollen in the middle; shell often

* *Les Columellaires*, Lamarck

transversely striated, and much varied in colour; operculum small and horny.

Animal, head having two tentacula, with eyes near the base.

—18 species, recent.

Inhabits both the Atlantic and Pacific, in warm latitudes. The *C. rustica* is also found on the southern coast of Italy.

113. MITRA. — *Shell* turreted, rather fusiform; spire pointed at the end; the lowest plaits on the inner lip are the smallest; outer lip sometimes toothed; operculum doubtful.* *Animal* not much known; the trunk, however, is very long, and the foot small. — 80 species recent, and 14 fossil.

These shells are much smaller and more slender than the *Volutes* (the next genus), but are not inferior to them in beauty of sculpturing and colour. They exhibit an infinite variety of patterns; sometimes grooved, sometimes

* Existing in some species, not in others. — Gray.

smooth, and variegated with every kind of hue. They abound in the seas of hot climates, the greater number inhabiting the Pacific Ocean, where they are met with chiefly in shallow water, near coral reefs; others, however, are found at great depths. The animal has been supposed to be of a poisonous nature, and to wound, with its pointed trunk, those who touch it.

114. VOLUTA. — *Shell* oval, thick, generally tuberculated; spire short or conical; aperture generally angulated and large; outer lip thickened within; the lowest plaits on the inner lip, or those which are nearest the notch, are the largest; no operculum.*

Animal, head distinct, having two short triangular tentacula, with eyes at the base, and a long thick proboscis or trunk; foot very large. — There are 44 species recent, and 18 fossil.†

* Except *Voluta musica*, which has one of moderate size. — Gray.

† This number includes the species of the next two genera, as described by Lamarck.

This genus, as constituted by Lamarck, contains some of the most beautiful shells that are known: they are generally smooth, shining, and the colours bright and varied; and differ exceedingly in form and size, some being globular, others oval, some turreted, and others with only a very small spire. Some of them have spines at the upper part of each whorl, which form a kind of thorny crown; many are curiously marked with lines and spots, so as to form some resemblance to a line of printed music; and one very scarce species is marked with five or six transverse milk-white bands upon a dark ground, and spotted with reddish brown, forming a beautiful contrast of colours.

The *Volutes* are found principally in hot climates, none having been discovered in European seas, and but few in America. They often attain a very large size; but the fossil species are generally smaller than the recent. The most rare species is *V. Junonia*. Very few examples of it are known, not perhaps more than four. It is of a yellowish white, and covered with a multitude of red-brown spots; some round, others

almost square, and disposed in transverse rows near one another.

Mr. Broderip has separated some shells from the genus *Voluta* under the names of *Melo* and *Cymba*. Their generic distinctions will be understood by the following descriptions.

115. MELO (Broderip).—*Shell* light, ventricose, oval, with a light greenish-brown epidermis; spire short, regular, sometimes hidden by the last whorl; aperture large, nearly as long as the whole shell; outer lip thin; columella slightly curved, with four or five oblique prominent plaits.

The aperture of this shell is much larger than that of *Voluta*, and the shell and outer lip are thinner. They are very beautiful, and of a large size.

116. CYMBA (Broderip).—*Shell* smooth, ventricose, with a short rounded spire; a very large wide aperture, terminating at the upper part in a flat ledge, which separates the outer lip from the body whorl; columella with three or four oblique projecting folds, ter-

minating in a point; outer lip thin, with its spiral edge sharp; epidermis smooth, brown, covered partly or entirely by the glassy enamel, which, commencing with the outer lip, spreads over the body of the shell.

These shells are found in Africa, and are distinguished by the rude ill-shapen spire, and the curious ledge which separates the outer lip from the body whorl.

117. MARGINELLA. — *Shell* oval, smooth; spire short; right lip having a thick margin; plaits nearly equal in size; no operculum. *Animal*, head very distinct, with a small proboscis, and two tentacula having eyes at the base; foot very large. — 22 species recent, and 3 fossil.

These shells are often very beautifully coloured; many have the spire exceedingly short. They inhabit hot climates. The inhabitant is supposed to cover its shell with its mantle, depositing successive layers of shelly matter, as

the examples of this genus differ materially in thickness of substance.

118. VOLVARIA. — *Shell* oval ; spire scarcely visible ; mouth narrow, and as long as the shell ; plaits few, at the lower part of the inner lip ; outer lip thin, or only slightly thickened. *Animal* unknown. — 5 species recent, and 1 fossil.

These shells are found on the coasts of Africa and China, where the natives string them into necklaces.

FAMILY 15. — *Convolutidæ*.*

Shell without a canal, but having the base notched ; the whorls of the spire large, rolled round each other, so that the last nearly, and in some instances entirely, conceals all the others, rendering the aperture long and narrow ; the body of the animal is therefore very flat. The three first genera of this family have the right lip recurved inwardly in the adult. They do not possess an operculum. There are seven genera.

* *Les Enroulées*, Lamarck.

119. CYPRÆA.—*Shell* oval, convex; mouth linear, notched at both ends, thickened and toothed on each side in the adult shell; spire sometimes entirely hidden. *Animal*, head distinct, furnished with two tentacula, having eyes at the base on small projections; mantle and foot very large.—There are 68 species recent, and 18 fossil.

These shells are remarkable for the different appearance they present during the several stages of their growth. Whilst young they are very thin, almost colourless, and dull on the outside; the mouth is rather wide, the outer lip not rolled inwards, but having a sharp edge. At a more advanced period the lips are curved inwards; but the shell is still thin, and the colours not bright.* At a later stage, when

* The gradations of advancing growth may be easily remarked in the common *C. Mauritiana*.

it has attained its full growth, the outer lip is thickened and rolled inwards, the back marked with the most beautiful patterns, and bearing an exquisite polish, and the spire, at the same time, if not entirely hidden, scarcely projects. The animal itself undergoes a considerable change in appearance during its growth. Its mantle is at first small, but increases with the enlargement of the shell, and expands at its sides into two large wings; from these is deposited the final layer of shelly matter which forms the exterior coating, and which is caused by the animal enveloping the shell entirely in its mantle. At the point where the two expansions of the latter meet on the back of the shell a line is consequently formed, which is called the dorsal line; but occasionally these lobes are so much extended as not only to meet, but to wrap over each other, in which case there is no line.

This shell is generally found in the seas of hot climates, the Indian Ocean, and the seas of Africa and America. A few species are met with in temperate regions; but they possess no

great beauty, neither attaining the vivid tints nor the delicate markings of those more favoured in situation. The *Cypræa Europea*, or Nun Cowrie, is found in abundance on our own coasts.

When at rest the *Cypræa* remains buried under the sand, at the bottom of the sea, at a short distance from the shore; it occasionally traverses the rocks and coast, and may be found under stones and corals.

The *Cypræa moneta*, or Money Cowrie, is the current coin of Siam, Bengal, and Africa: it is collected in the latter by the negro women, and sent into different countries. In Bengal 3200 are reckoned to be equivalent to a rupee, or about two shillings of English money. In the Friendly Islands permission to wear the *Cypræa aurantia*, or Orange Cowrie, as an ornament, is only granted to persons of the highest rank.

The *C. aurora* is considered the most rare of this genus. The New Zealanders suspend it to their dress as an ornament, and it is therefore frequently found with an artificial perforation. *C. cervina* is supposed to be the

largest species, being more than four inches in length. It inhabits the American seas.

120. ERATO (Risso). — *Shell*, spire conical; apex rather round, blunt; shell, when young, smooth; the adult with both lips finely marked with lines; columella concave, plaited or smooth. A groove down the back of the shell. *Animal* like that of *Cypræa*. From the Pacific Ocean.

121. OVULA. — *Shell* oblong; inner lip without teeth; outer slightly toothed or wrinkled, the extremities generally projecting; spire quite hidden. *Animal* very similar to that of the *Cypræa*; the mantle, however, has only one lobe, which folds over the left lip. — 12 species recent, and 2 fossil.

In some species the ends of the mouth are so much lengthened as to make the shell fusiform or spindle-shaped. The *Ovula oviformis* is perfectly white outside, and of an orange brown within; the young shell is as thin as paper.

This genus chiefly inhabits the Indian and Chinese seas; it is also found in the Black Sea and the Mediterranean.

122. TEREHELLUM. — *Shell* oblong, smooth, and convolute; spire pointed; sutures not channelled; mouth narrow and long; right lip rather sharp, left lip quite smooth. *Animal* unknown. — 1 species recent, and 2 fossil.

These shells are brought from the Indian seas; they are thin, delicate, and prettily marked with bands and cloudy spots.

123. ANCILLARIA. — *Shell* oblong and very smooth; spire pointed; sutures not channelled, but nearly effaced; mouth wide and long; right lip sharp; left lip having the base thickened, with a shelly ridge. *Animal* almost unknown; the foot, however, is very large. — 4 species recent, and 5 fossil.

This genus is chiefly confined to tropical climates; the shells are smooth, and highly polished, presenting a surface infinitely more even than if produced by artificial means: some

of them have a small tooth at the lower part of the right lip. When the animal is alive, the shell is covered so much by the foot that only the middle of the back can be seen. The *Ancillaria cinnamonea* is exceedingly scarce.

124. OLIVA.—*Shell* oblong, smooth, and convolute; spire short; sutures channelled or

grooved; mouth narrow and long; right lip rather sharp; left lip thickened at the top, and striated; operculum horny and small in some species, in others not existing.* *Animal*, head small, terminated by a proboscis; two tentacula enlarged at the base, and having the eyes si-

* Gray.

tuated in the middle; foot very large; mantle large, covering great part of the shell. — There are 62 species recent, and 5 fossil.

These shells are very common, and are often very beautiful; many attain a large size. They are brought principally from Asia, but are also found in the African and American seas.

125. CONUS. — *Shell* shaped like a cone, the spire forming the base, and scarcely projecting; mouth narrow and long; left lip smooth; operculum very small and horny*; usually an epidermis. *Animal*, head very distinct, terminated by a trunk capable of being greatly extended; two tentacula, with eyes near the summit; foot oval, and rather long. — 181 species recent, and 9 fossil.

Many of the species of this genus are marked

* Adanson.

with the most beautiful and extraordinary figures, some of them resembling Hebrew, Greek, or Arabic characters, and bearing a most exquisite polish; in other varieties the colours are arranged in almost endless shapes, being clouded, veined, marbled, dotted, striped, and banded, in every kind of form. Many of them are very rare, and a specimen of the *Conus gloria maris* has been sold for as much as one hundred guineas.*

In some of the species the spire presents the appearance of having been pushed into the body of the shell, the apex being below the level of the outer whorl; a spire of this kind is called *retuse*.

When a section is made of a cone, it will appear that the sides of the internal whorls are much thinner than those of the external; it is supposed that this arises from the animal re-

* The famous *Conus cedo-nulli*, formerly in the cabinet of Lyonnet, at the Hague, is said to have been sold for three hundred guineas. There are very few specimens existing. Lamarck's specimen was considered by him the most precious in his collection.

ducing the thickness of these internal walls by absorption, either for the purpose of giving itself more room, or in order to diminish the weight of the shell.

They are found abundantly in all tropical countries, particularly in Asia, but become more rare as they approach the northern hemisphere; a few species only are found in the Mediterranean.

ORDER IV.

CEPHALOPODA.*

There is much difficulty attending the classification of this order, which derives its name from the position of the animal's apparatus for moving; it possesses for that purpose a circle of eight, ten, or more fleshy processes, which may be indifferently called arms, legs, or tentacula. These appendages are attached to the head of the animal, which is also furnished

* *Les Céphalopodes*, Lamarck.

with a mouth terminating in two strong mandibles, similar to the beak of a parrot; it likewise possesses a perceptible organ of hearing, and two large eyes, immoveable, and without eyelids.

The animals may be separated into three divisions,—*Polythalama*, in which the shell is chambered or divided into numerous cells; *Monothalama*, in which the shell has only one chamber, destitute of partitions; and a third, in which there is no shell whatever.

The mantle of the *Cephalopoda* is in the form of a bag, and covers the lower part of the body. Most of the recent species live at great depths in the sea, and are found nearly all over the world. Some swim, and attach themselves to marine bodies; others creep on the sand, by the aid of their tentacula, and often retire into holes in the rocks. They are all *zoophagous*, living on crabs and other animals, which they seize and devour.

The following is a list of the genera which Lamarck has introduced into this order:—

1. Belemnites.
2. Orthoceras.

- *3. Nodosaria.
4. Hippurites.

5. Conilites.	*20. Polystomella.
6. Spirula.	*21. Vorticialis.
*7. Spirolina.	22. Nummulites.
*8. Lituola.	23. Nautilus.
*9. Renulina.	24. Ammonites.
*10. Cristellaria.	25. Orbulites.
*11. Orbiculina.	26. Ammonoceras.
*12. Miliola.	27. Turrilites.
13. Gyrogonia.	28. Baculites.
*14. Melonia.	29. Argonauta.
*15. Rotalia.	30. Octopus.
*16. Lenticulina.	31. Lorigopsis.
*17. Placentula.	32. Loligo.
*18. Discorbis.	33. Sepia.
*19. Siderolites.	

Those genera which are marked thus * are now removed from the *Cephalopoda*, having been proved to differ from molluscos animals by Dujardin, and by him are placed in a distinct class. He says they are inferior in their organization to the *Radiata*, and endowed with locomotive power by the instrumentality of minute tentacular filaments. Ehrenburg makes them belong to the second great division (*Bryozoa*) of his coral animals.

No. 4. in the list proves to be a Bivalve, not unlike a *Chama*. No. 13. is the seed of the *Chara*. Nos. 5. 25. 26. are doubtful genera.

Of those left, *Spirula*, *Nautilus*, and *Argonauta* are perhaps the most interesting to the young conchologist, and will therefore be fully described. The remainder are nearly all fossils. Some of the generic names have been changed by modern naturalists; but as there is much uncertainty respecting them, they will only be described in a cursory manner.

DIVISION I.—POLYTHALAMA.*

126. BELEMNITES.—*Shell* straight, in the shape of a long cone; external part a sheath

having a conical cavity in the upper part, and containing an inner cone, which is chambered and perforated.—2 species described.

These shells are only found in a fossil state, and are very common, having been met with in all ages and countries, and having given rise to much speculation as to their real character. It

* *Les Céphalopodes-polythalamés*, Lamarck.

is but recently that their true origin has been discovered, and their place in creation ascertained; in former ages they have been described as petrified fingers, thunder-stones, stalactites, or as the teeth of an animal.— There are nearly 100 species at present known.

127. ORTHOCERAS. — *Shell* straight, with septa regularly concave towards the mouth, perforated by a simple and nearly cylindrical siphuncle, either in or not far removed from the centre of the disk.

All fossil? Found in strata mostly below the carboniferous or mountain limestone.

128. SPIRULA. — *Shell* cylindrical, thin, and almost transparent; partially spiral, the whorls not touching each other; perforation of the chambers at the side, which are equidistant; partitions externally concave. *Animal* with ten arms, two of which are longer than the others, and nearly conceal the shell. — 1 species, recent.

This shell, which is very curious, and not uncommon, is found in the Eastern Ocean. It is thin and fragile, and of a pearly white. The only species known is *S. Peronii*.

129. NUMMULITES.—*Shell* thin towards the margin; spire internal, discoid; whorls contiguous, and not apparent; cells numerous and small; partitions transverse, and not perforated.—4 species.

These shells are all fossil, and exist in immense numbers; they are very flat, and take their name from their resemblance to pieces of money. In many parts of the world they are accumulated in such vast masses as to form entire mountains, and many buildings have been constructed of limestone crowded with them. They vary in size from more than an inch in diameter to great minuteness.

130. NAUTILUS.—*Shell*, whorls contiguous, the last enveloping the others; chambers numerous, formed by transverse partitions, and perforated by a tube. *Animal* having numerous tentacula, and attached to the shell by two

muscles.—2 species recent, 1 of which is also found fossil.

This shell and its remarkable inhabitant are

found in the seas of warm latitudes, especially those of Africa, Asia with its islands, and Australia. They usually remain at the bottom of the water, and are able to creep along rather quickly, supporting themselves upon their tentacula, with the head downwards, and the shell raised above; they often get into the fishermen's nets when thus moving about at the bottom of the sea. After stormy weather, as it becomes more calm they may be observed,

in great numbers, floating upon the surface of the waves, with the head put out, and the tentacula resting upon the water, the shell at the same time being undermost; they do not, in a general way, remain long sailing in this manner, and have the power of easily returning to their situation at the bottom of the sea, by merely drawing in their tentacula and upsetting the shell, which immediately sinks with them. It was only in 1829 that this animal was known with any certainty, one having been caught alive, by Mr. G. Bennett, near the New Hebrides Islands; the animal, after drawings were made of it, was preserved in spirits, and is now in the museum of the College of Surgeons.

The ancients fabricated drinking-cups of the shell of the *Nautilus pompilius*; the other species, the *Umbilicatus*, is very rare, and is distinguished by a large umbilicus on each side: the former is often found in a fossil state.

131. AMMONITES.* — *Shell* discoid, spiral;

* This genus has been divided into three; *Ammonites*, *Goniatites*, and *Clymenia*.

whorls contiguous, and all apparent; inner surface (the exterior covering has generally disappeared) marked by the sinuous partitions, which are transverse, and lobed or cut, but not perforated; last chamber occupying the whole of the last whorl of the spire.—Between 200 and 300 species.

These shells are all fossil, and are occasionally two feet or more in diameter.* They occur in Europe, Asia, and America. Dr. Gerard found them in the Himalaya mountains, an elevation of 16,000 feet; M. Ménard met with one in the Maritime Alps. Their numbers must have been very great, as M. Dufresne informed Lamarck that the road from Auxerre to Avalon, in Burgundy, was absolutely paved with them; and it is no uncommon occurrence to find them used in some parts of Somersetshire, Gloucestershire, and Yorkshire, to pave the roads.

132. TURRILITES.—*Shell* spiral, turreted;

* Dr. Buckland speaks of some found in the chalk near Margate which exceeded four feet in diameter.

whorls contiguous and apparent; partitions sinuous; mouth round. — 1 species, fossil.

133. BACULITES. — *Shell* straight, cylindrical, sometimes slightly compressed, and

somewhat conical; partitions sinuous, transverse, a little distant. — 3 species.

All fossil; abundant in the limestone of Valognes in Normandy.

DIVISION II. — MONOTHALAMA.*

134. ARGONAUTA. — *Shell* monothalamous, symmetrical, external, and containing, (but not

* *Monothalames*, Lamarck.

attached to,) the body of the animal ; spire bica-
minated, tubercled, and curved back into the

mouth. *Animal* having eight tentacula; two of which are thin and dilated, and are often called the sails. — 3 species, recent.

These curious and beautiful shells are found in the seas of warm latitudes, and are both pelagic and litoral.* They are commonly called the Paper Nautilus.

An interesting series of experiments was made a few years ago by Madame Power, for the purpose of ascertaining the nature and habits of the *Argonauta*, about which much contro-

* Penny Cyclopædia, art. *Paper Nautilus*.

versy had existed. She constructed a kind of cage in the sea, near the coast of Messina, and enclosed in it several of the living animals, which she kept supplied with their natural food, consisting of small molluscs. The result of her observations went to prove that the animal is in the habit of floating, or rather sailing, on the surface of the water, using its dilated tentacula as sails, the remainder as oars, and aiding its movements by means of a kind of proboscis which it employs as a helm: the animal is not attached to the shell, but, when under water, it adheres firmly to it by its sail-arms, which are turned over and affixed to the outside, entirely covering it. The shell, which is remarkably brittle when exposed to the air, is perfectly pliable in water, and is thus enabled to escape the destruction which would otherwise appear likely to be met with by so thin and tender a fabric.

The animal of the *Argonauta*, during life, and when unconscious of being observed, is very beautiful, the sails and other visible parts presenting the most lively colours—purple, gold, and silver. At the approach of any object, it

takes in its tentacula, wraps its sails over the shell, and descends, blackening the water at the same time, if hard pressed, by a discharge of ink, to cover its escape. During the experiments referred to, the animals seemed very shy, and would descend to the bottom on the approach of any person; except when hungry, at which time they would take food from the hands of Madame Power.

DIVISION III.—*No external shell.*

The four following genera do not form an external shell, and will not therefore be particularly described.

135. OCTOPUS.—Has two little cartilaginous pieces in the interior of the body.

It is found in all seas.

136. *LOLIGOPSIS*.—Internal support unknown. The tentacula are enormously long.

137. *LOLIGO*.—*Shell* thin, transparent,

sometimes partially gelatinous, of variable form, but generally shaped like a pen.

Common on our coasts.

138. *SEPIA*.—*Shell* calcareous, oval, thick,

concave on both sides, terminated posteriorly by a point.

Brought principally from India, though found on our coast and the Mediterranean. Mr. F. D. Bennet states that the cuttle fish is considered a luxury by all classes of the Sandwich Islanders, and that when fresh and well cooked it is excellent, being in consistence and flavour not unlike the flesh of a lobster's claw.*

There are some very extraordinary accounts

* Narrative of a Whaling Voyage round the World.

given of the supposed size and power of these mollusca by De Montford, in Buffon's *Histoire Naturelle*, but, though exceedingly amusing, their accuracy is now doubtful. Some are described as large enough to draw down men-of-war at sea, and completely destroy them. But this is an exaggeration no doubt, though there are some species that grow to a very large size. Mr. Swainson remarks in his *Malacology* that he saw many caught on the shores of Sicily, and that two would be a heavy load, their arms being as thick as those of a man.

ORDER V.

HETEROPODA.

In this tribe, the foot is formed into a kind of fin, by the aid of which the animal moves; there are only four genera, *Carinaria*, *Atlanta*, *Pterotrachea*, and *Phyllirœe*, and of these the two former alone possess shells, and need a particular description.

139. CARINARIA (*Cuvier*).—*Shell*, conical, flat at the sides, transparent, and having the apex recurved. *Animal*, body long, gelatinous,

possessing one or more fins ; the head can be contracted within the body, and is provided with two tentacula, two eyes, and a retractile proboscis. — 3 species recent.

There are only three species known of this singular genus. The *Carinaria vitrea* is so called from its resemblance in substance to glass, and is found in the African seas : it is very rare. The *Carinaria fragilis* is extremely delicate in its texture, and inhabits the Mediterranean ; and the *Carinaria cymbium*, or *boat-shaped*, which is not larger than a grain of sand, and can only be seen with the assistance of a microscope, is brought from the Eastern Ocean and Mediterranean Sea ; it is found on the coast near Nice, and feeds on small *Medusæ* and other gelatinous substances, as also on minute fish, and occasionally the more feeble of its own species.

The mollusca which form these shells are very transparent, but are possessed of lively and brilliant colours ; they only appear on the surface of the sea during calms. The shells are comparatively small, as they only cover the circulating and respiratory organs ; they were

formerly very rare (specimens having been sold for one hundred guineas), but are now more common: they are seldom perfect.

140. ATLANTA.—*Shell*, somewhat resembling the *Carinaria*,

 but having the apex spirally whorled; transparent, and large enough to contain the animal entirely: a thin and glassy operculum.

Animal, head having two tentacula, and large eyes at the base; foot or fin large.—2 species recent.

This genus is found in all the seas of hot climates.

141. PTEROTRACHEA. }
 142. PHYLLIRÖE. } No shell.

In these two genera, the animal resembles the *Carinaria* in the position of the heart and gills; they are both found in the Mediterranean, and, owing to their extraordinary transparency, it is very difficult to perceive them when in the water. The *Pterotrachea* has four species, the *Phylliröe* only one.

Having brought this elementary work to a conclusion, it must be added, that the greater advance the reader makes in the study of Conchology, the more he will find to interest and delight him. The beauty of form in some shells, the grotesque shapes of others, the exquisite colouring and marking of many, and the habits and peculiarities of the inhabitants of all, will excite wonder and admiration at the care which has been bestowed on creatures, of which thousands are never seen by the eye of man. He will be led to reflect that they were formed by Him who has made nothing in vain, and that these, among the many exquisite productions of Nature, were created "for man to gaze on, wonder, and adore."

GUIDE

TO

THE USE OF THE TABLES.

THE young student will soon ascertain the generic name of any shell by adhering to the following method :— We will suppose an *oyster* to be a shell under examination. At one glance it will be found to be a Bivalve. The first table must, therefore, be consulted, which is found to be divided into three orders, each of which must be examined. The first has no ligament, but a pedicle to attach the shell to other substances. The oyster having a ligament (if the specimen be perfect), the description of the next order must be looked at ; but the two muscular impressions at once determine that the oyster does not belong to it. The specimen, therefore, must be placed amongst the Monomyaria, which is divided into two parts : the first has the ligament lengthened, which is not the case with the oyster, for it is thick and short. It may, therefore, be concluded that the shell belongs to the second division, and to one of its two families. The Pectenidæ are distinguished by the shells being solid, eared, and often ribbed. The oyster does not resemble this description, and therefore must be of the Ostreidæ family, for it is foliaceous. When the family is ascertained, the generic name is found by turning to the description of the former in the body of the work ; and the account of each genus, with the assistance of the drawing, will enable the student to determine that the

shell is an *Ostrea*. The next class (the Univalves) may be treated in the same way. A *whelk*, a shell common on our coasts, belongs to this class. The second table must be consulted. There are five orders, which must each be examined. The shell in question cannot be of the Order Pteropoda, because it is neither cartilaginous nor horny. It cannot belong to Gasteropoda, because it is not cup-shaped nor slightly rolled up. It may belong to Trachelipoda, because it is spiral. In the first division of that order the aperture is entire, which that of the *whelk* is not. The second division has the aperture with a canal or notch : this agrees with the specimen in hand. Each family of that division must now be examined. It cannot belong to Canaliferidæ, because it has not a canal, nor the right lip thickened. The Pteridæ have the right lip greatly dilated or lobed : that family may therefore be passed over. The Purpuridæ have a short canal turned back, or a notch, and the form of the shell is globular. The whelk is notched, and its form is rather round ; and as it could not belong to the other two remaining families, we may suppose it is one of the Purpuridæ. The genus is next found by consulting the account of that family in the work, and it will be with ease ascertained that the whelk is a *Buccinum*. By affixing a number to the specimen, corresponding to that in the book, it may be referred to at any time to refresh the memory.

The Annulata and Cirrhipeda have not been included in the tables, as their names are more easily ascertained than those of the other classes, from there being few genera, and their general form and appearance being very different.

AN
ALPHABETICAL LIST
OF
ORDERS, DIVISIONS, AND FAMILIES.

	Page
ADHÆRENTIA - - - -	50
AMPHITRITIDÆ - - - -	31
ANNULATA - - - -	28
APLYSIDÆ - - - -	157
ARCIDÆ - - - -	94
 BRACHIOPODA - - - -	 47
BULLÆIDÆ - - - -	155
 CALYPTRÆIDÆ - - - -	 148
CANALIFERIDÆ - - - -	219
CARDIIDÆ - - - -	91
CEPHALOPODA - - - -	260
CHAMIDÆ - - - -	103
CIRRHIPEDA - - - -	36
CIRRHIPEDA PEDUNCULATA - - - -	40
CIRRHIPEDA SEDENTARIA - - - -	37
COLIMACIDÆ - - - -	164
COLUMELLIDÆ - - - -	245
CONCHIDÆ - - - -	85
CONCHIFERA (<i>Bivalves</i>) - - - -	45
CONCHIFERA AMBIGUA - - - -	103

	Page
CONCHIFERA CRASSIPEDA - - -	54
CONCHIFERA LAMELLIPEDA - - -	85
CONCHIFERA TENUIPEDA - - -	72
CONVOLUTIDÆ - - -	251
DENTALIDÆ - - -	140
DIMYARIA - - -	53
DORSALIDÆ - - -	29
GASTEROPODA - - -	139
HETEROPODA - - -	274
JANTHINIDÆ - - -	196
LIMACIDÆ - - -	159
LITHOPHAGIDÆ - - -	77
LYMNÆIDÆ - - -	181
MACROSTOMIDÆ - - -	197
MACTRIDÆ - - -	73
MALDANIDÆ - - -	31
MALLEIDÆ - - -	114
MELANIDÆ - - -	186
MOLLUSCA (<i>Univalves</i>) - - -	133
MONOMYARIA - - -	106
MONOTHALAMA - - -	269
MYIDÆ - - -	68
MYTILIDÆ - - -	109
NAIIDÆ - - -	98
NERITIDÆ - - -	192
NYMPHIDÆ - - -	79

					Page
OSTREIDÆ	-	-	-	-	126
PECTINIDÆ	-	-	-	-	121
PERISTOMIDÆ	-	-	-	-	187
PLICACIDÆ	-	-	-	-	202
PHOLADIDÆ	-	-	-	-	60
PHYLLIDIDÆ	-	-	-	-	141
PHYTOPHAGA	-	-	-	-	164
POLYTHALAMA	-	-	-	-	263
PTERIDÆ	-	-	-	-	229
PTEROPODA	-	-	-	-	136
PURPURIDÆ	-	-	-	-	233
SCALARIDÆ	-	-	-	-	204
SEMIPHYLLIDIDÆ	-	-	-	-	147
SERPULIDÆ	-	-	-	-	34
SOLENIIDÆ	-	-	-	-	63
TENDINOSA	-	-	-	-	48
TRACHELIPODA	-	-	-	-	163
TRIDACNIDÆ	-	-	-	-	107
TRITONIDÆ	-	-	-	-	139
TUBICOLIDÆ	-	-	-	-	54
TURBINIDÆ	-	-	-	-	210
ZOOPHAGA	-	-	-	-	218

AN

ALPHABETICAL LIST OF GENERA,

WITH THEIR DERIVATIONS.

	Page
ACASTA - - - - -	39
ACERA - - - - -	155
ACHATINA. <i>Achates</i> , an agate - - -	178
AMMONITES. From Jupiter <i>Ammon</i> , who is represented with a ram's horn - - -	267
AMPHIDESMA. <i>Amphi</i> , Gr., in the middle; <i>desma</i> , a ligament - - -	76
AMPHITRITE - - - - -	33
AMPULLACERA. <i>Ampulla</i> , a rounded vessel -	191
AMPULLARIA. Do. - - - - -	189
ANASTOMA. <i>Ana</i> , Gr., backwards; <i>stoma</i> , mouth	171
ANATIFA. <i>Anas</i> , a duck; <i>fero</i> , to bear - -	41
ANATINA. <i>Anatinus</i> , that which belongs to a duck -	69
ANATINELLA. The diminutive of <i>Anatina</i> -	72
ANCILLARIA - - - - -	256
ANCYLUS. A little handle; probably from the apex somewhat resembling a handle - -	185
ANODONTA. <i>A</i> , without; <i>odos</i> , a tooth, Gr. -	101
ANOMIA - - - - -	132
APLYSIA - - - - -	157
ARCA. A boat - - - - -	95

					Page
ARENICOLA.	<i>Arena</i> , sand ; and <i>colo</i> , to inhabit	-	-	-	29
ARGONAUTA.	The ship <i>Argo</i> , and <i>nauta</i> , a sailor	-	-	-	269
ASPERGILLUM.	<i>Aspergo</i> , to sprinkle	-	-	-	54
ASTARTE	-	-	-	-	88
ATLANTA	-	-	-	-	276
AURICULA.	<i>Auris</i> , an ear	-	-	-	180
AVICULA.	<i>Avis</i> , a bird	-	-	-	118
BACULITES.	<i>Baculum</i> , a staff	-	-	-	269
BALANUS.	An acorn	-	-	-	38
BELEMNITES.	<i>Belemnion</i> , Gr., a dart or arrow	-	-	-	263
BUCCINUM.	A horn	-	-	-	242
BULIMUS	-	-	-	-	174
BULLA.	A bubble	-	-	-	156
BULLÆA.	A do.	-	-	-	155
CALYPTRÆA.	<i>Calyptra</i> , Gr., a covering	-	-	-	152
CANCELLARIA.	<i>Cancellatus</i> , cross-barred	-	-	-	223
CAPSA.	A box	-	-	-	84
CARDITA.	<i>Cardia</i> , Gr., a heart	-	-	-	92
CARDIUM.	Do.	-	-	-	91
CARINARIA.	<i>Carina</i> , a keel	-	-	-	274
CAROCOLLA	-	-	-	-	170
CASSIDARIA.	<i>Cassis</i> , a helmet	-	-	-	234
CASSIS.	Do.	-	-	-	235
CERITHIUM	-	-	-	-	219
CHAMA.	<i>Chamai</i> , Gr., on the ground	-	-	-	104
CHITON.	Gr., integument	-	-	-	142
CHITONELLUS.	From <i>Chiton</i>	-	-	-	142
CINERAS.	<i>Cinereus</i> , ash-coloured	-	-	-	43
CLAUSILIA.	<i>Clausium</i> , a valve or folding-door	-	-	-	173
CLAVAGELLA.	<i>Clava</i> , a club	-	-	-	56
CLIO.	One of the Muses	-	-	-	136

					Page
CLEODORA	-	-	-	-	137
CLYMENE	-	-	-	-	31
COLUMBELLA.	<i>Columba</i> , a dove	-	-	-	245
CONCHOLEPAS.	<i>Concha</i> , a shell ; <i>lepas</i> , a stone	-	-	-	239
CONUS.	<i>Conos</i> , Gr., a cone	-	-	-	258
CORBIS.	A basket	-	-	-	83
CORBULA.	A little basket	-	-	-	71
CORONULA.	<i>Corona</i> , a crown	-	-	-	37
CRANIA.	<i>Cranium</i> , a skull	-	-	-	51
CRASSATELLA.	<i>Crassus</i> , thick	-	-	-	74
CRENATULA	-	-	-	-	114
CREPIDULA.	A little slipper	-	-	-	151
CREUSA.	<i>Creux</i> , Fr., a cavity	-	-	-	39
CUCULLÆA.	<i>Cucullus</i> , a hood	-	-	-	95
CYCLAS.	<i>Cyclos</i> , round ?, Gr.	-	-	-	86
CYCLOSTOMA.	<i>Cyclos</i> , round ; <i>stoma</i> , mouth	-	-	-	181
CYMBÆ.	A boat or skiff	-	-	-	249
CYMBULIA.	Do.	-	-	-	138
CYPRÆA	-	-	-	-	252
CYPRICARDIA.	<i>Cyprina</i> , and <i>cardia</i> , a heart	-	-	-	93
CYPRINA	-	-	-	-	87
CYRENA	-	-	-	-	86
CYTHERIA	-	-	-	-	89
DELPHINULA.	<i>Delphinus</i> , a dolphin	-	-	-	209
DENTALIUM.	<i>Dens</i> , a tooth	-	-	-	140
DICERAS.	<i>Dis</i> , Gr., double ; <i>ceras</i> , a horn	-	-	-	103
DOLIUM.	A tun	-	-	-	241
DOLABELLA.	<i>Dolabra</i> , a hatchet	-	-	-	158
DONAX.	A wedge	-	-	-	84
DORIS	-	-	-	-	140
EBURNA.	<i>Ebur</i> , ivory	-	-	-	243

				Page
EMARGINULA.	<i>Margo</i> , border	-	-	149
EOLIS	-	-	-	140
ERATO.	<i>Eradere</i> , to scrape out	-	-	255
ERYCINA.	<i>Eruo</i> , Gr., to be enclosed	-	-	75
ETHERIA.	<i>Ether</i> , air	-	-	105
FASCIOLARIA.	<i>Fascia</i> , a band	-	-	223
FISSURELLA.	<i>Fissura</i> , a fissure	-	-	150
FISTULANA.	<i>Fistula</i> , a pipe	-	-	57
FUSUS.	A spindle	-	-	224
GALATHEA.	A sea nymph	-	-	87
GALEOLARIA.	<i>Galea</i> , a helmet	-	-	35
GALEOMMA	-	-	-	81
GASTROCHÆNA.	<i>Gaster</i> , Gr., belly ; <i>chaino</i> , to			
gape	-	-	-	58
GLAUCUS	-	-	-	140
GLYCIMERIS	-	-	-	67
HALIOTIS.	<i>Hals</i> , Gr., sea ; <i>os</i> , ear	-	-	199
HARPA.	A harp	-	-	239
HELICINA.	From <i>Helix</i>	-	-	172
HELIX.	A snail	-	-	165
HIPPONYX.	<i>Hippes</i> , Gr., a horse ; <i>onyx</i> , a nail or			
hoof	-	-	-	153
HIPPOPUS.	<i>Hippos</i> , Gr., a horse ; <i>pous</i> , a foot	-	-	109
HYALÆA.	<i>Hyalus</i> , glass	-	-	136
HYRIA	-	-	-	101
IRIDINA.	<i>Iris</i> , a rainbow	-	-	102
ISOCARDIA.	<i>Isos</i> , Gr., similar ; <i>cardia</i> , a heart	-	-	93

	Page
JANTHINA. <i>Janthium</i> , a violet - - -	196
LEPTOCONCHUS. <i>Leptos</i> , Gr., thin ; <i>conche</i> , shell	207
LIMA. A file - - - -	122
LIMACINA. From <i>limax</i> , a snail - - -	137
LIMAX. A snail - - - -	160
LINGULA. A tongue - - - -	48
LITHODOMUS. <i>Lithos</i> , Gr., a stone ; <i>domos</i> , house	110
LITHOTRYA. <i>Lithos</i> , <i>truo</i> , to bore through -	40
LOLIGO - - - - -	273
LOLIGOPSIS - - - - -	273
LUCINA. A name given to Juno and Diana -	83
LUTRARIA. <i>Lutrum</i> , mud - - -	73
LYMNEA. <i>Limus</i> , mud - - -	183
MACTRA. A kneading-trough - - -	73
MAGILUS - - - - -	205
MALLEUS. A hammer - - - -	116
MARGINELLA. A little rim or border - -	250
MELAMPUS. <i>Melas</i> , Gr., black ; <i>pous</i> , a foot -	203
MELANIA. <i>Melas</i> , Gr., black - - -	186
MELANOPSIS. Do. - - - -	187
MELO. A melon - - - - -	249
MESODESMA. <i>Mesos</i> , Gr., in the middle ; <i>desma</i> , a ligament - - - - -	75
MITRA. A head-dress - - - -	246
MODIOLA. <i>Modilus</i> , a little measure - -	110
MONOCERAS. <i>Monos</i> , Gr., single ; <i>ceras</i> , a horn -	238
MONODONTA. <i>Monos</i> , Gr., single ; <i>odos</i> , a tooth -	214
MYA. <i>Myon</i> , Gr., a muscle - - -	68
MYTILUS - - - - -	111
MUREX. A sharp rock - - - -	227

	Page
NASSA - - - - -	243
NATICA. <i>Nato</i> , to swim - - -	195
NAUTILUS. A little boat - - -	265
NAVICELLA. A little ship - - -	192
NERITA. The ancient name of a sea-shell -	194
NERITINA. From <i>Nerita</i> - - -	193
NUCULA. A small nut - - -	97
NUMMULITES. <i>Nummus</i> , money - - -	265
OCTOPUS. <i>Octo</i> , Gr., eight ; <i>pous</i> , foot -	272
OLIVA. An olive - - -	257
ONCHIDIUM - - - - -	159
ONISCIA - - - - -	235
ORBICULA. <i>Orbiculus</i> , a little globe -	52
ORTHOCERAS. <i>Orthos</i> , straight ; <i>ceras</i> , a horn -	264
OSTREA. - - - - -	126
OTION. A little ear, Gr. - - -	44
OVULA. <i>Ovum</i> , an egg - - -	255
PALUDINA. <i>Palus</i> , a marsh - - -	188
PANDORA - - - - -	71
PANOPÆA. <i>Pan</i> , Gr., all ; <i>opaion</i> , an aperture (open at both ends) - - -	66
PARMACELLA. A little cell - - -	160
PARMOPHORUS - - - - -	149
PATELLA. A dish or platter - - -	143
PECTEN. A comb - - -	123
PECTENARIA. Do. - - -	31
PECTUNCULUS. Do. - - -	96
PEDUM. A shepherd's crook - - -	121
PENTELASMIS. <i>Pente</i> , Gr., five ; <i>elasma</i> , a plate -	41
PERNA. A gammon of bacon - - -	115
PETRICOLA. <i>Petrus</i> , a stone ; <i>cola</i> , an inhabitant -	78

	Page
PHASIANELLA. <i>Phasianus</i> , a pheasant - -	216
PHOLADOMIA. <i>Pholas</i> and <i>mya</i> - -	66
PHOLAS. <i>Pholeo</i> , Gr., to lie hid in a cavity -	61
PHYLLIDEA - - - -	141
PHYLLIRÖE - - - -	276
PHYSA. A bubble - - - -	183
PILEOPSIS. <i>Pileos</i> , a bonnet - - - -	154
PINNA. The fin of a fish - - - -	112
PLACUNA. <i>Placous</i> , Gr., a cake - - - -	129
PLACUNANOMIA. <i>Placuna</i> and <i>anomia</i> - -	130
PLANAXIS. <i>Plana</i> , flat, and <i>axis</i> - - - -	216
PLANORBIS. <i>Plana</i> , flat ; <i>orbis</i> , an orb -	182
PLEUROBRANCHUS. <i>Pleura</i> , Gr., the side ; <i>branchæ</i> , the gills - - - -	147
PLEUROTOMA. Do., and <i>tome</i> , a slit - - - -	221
PLICATULA. <i>Plicatus</i> , of a fold - - - -	125
POLLICIPES. <i>Pollex</i> , a thumb's breadth ; <i>pous</i> , a foot - - - -	43
PNEUMODERMON - - - -	138
PSAMMOBIA. <i>Psammos</i> , sand - - - -	80
PTEROCERAS. <i>Pteron</i> , a wing ; <i>ceras</i> , a horn -	231
PTEROTRACHEA. Do., and <i>trachea</i> - - - -	276
PUPA. A chrysalis - - - -	172
PURPURA. The shell-fish from which the purple colour is taken - - - -	237
PYRAMIDELLA. A little pyramid - - - -	203
PYRGONA. <i>Pyrgos</i> , Gr., a tower - - - -	39
PYRULA. A little pear - - - -	224
RANELLA. <i>Rana</i> , a frog - - - -	227
RICINULA. Resembling the seed-vessel of the <i>Ricinus</i> - - - -	237
RISSEA. From M. Risso - - - -	209

				Page
ROSTELLARIA.	<i>Rostrum</i> , a beak	-	-	230
ROTELLA.	A little wheel	-	-	211
SABELLARIA.	<i>Sabula</i> , sand	-	-	32
SANGUINOLARIA.	<i>Sanguis</i> , blood	-	-	80
SAXICAVA.	<i>Saxum</i> , a stone; <i>cava</i> , hollow	-	-	77
SCALARIA.	<i>Scala</i> , a ladder	-	-	207
SCALPELLUM.	A little knife	-	-	43
SCYLLEA	-	-	-	140
SEPIA	-	-	-	273
SERPULA.	<i>Serpo</i> , to creep like a worm	-	-	34
SIGARETUS	-	-	-	197
SILICUARIA.	<i>Siliqua</i> , a pod	-	-	30
SIPHONARIA.	<i>Siphon</i> , Gr.	-	-	146
SOLARIUM.	A terrace or gallery	-	-	210
SOLECURTUS.	<i>Solen</i> , and <i>curtus</i> , short	-	-	65
SOLEMYA.	<i>Solen</i> and <i>mya</i>	-	-	68
SOLEN.	A kind of shell-fish (Pliny)	-	-	63
SPIRORBIS.	<i>Spira</i> , a fold; <i>orbis</i> , a globe	-	-	34
SPIRULA.	<i>Spira</i> , a winding compass	-	-	264
SPONDYLUS.	A shell-fish (ancients)	-	-	125
STOMATELLA.	<i>Stoma</i> , Gr., a mouth	-	-	198
STOMATIA.	Do.	-	-	198
STROMBUS	-	-	-	232
STRUTHIOLARIA.	<i>Struthio</i> , an ostrich	-	-	226
SUCCINEA.	<i>Succinum</i> , amber	-	-	179
TELLINA.	<i>Tellus</i> , the earth	-	-	81
TEREBELLA	-	-	-	32
TEREBELLUM.	<i>Terebra</i> , an augur	-	-	256
TEREBRA.	Do.	-	-	244
TEREBRATULA.	<i>Terebratus</i> , bored	-	-	49
TEREDO.	A piercer	-	-	58

				Page
TESTACELLA.	<i>Testa</i> , a shell	-	-	160
TETHYS	-	-	-	140
THECIDIUM.	<i>Theca</i> , a box	-	-	50
THRACIA.	From <i>Thrace</i>	-	-	70
TORNATELLA.	<i>Tornatus</i> , turned	-	-	202
TRIDACNA.	<i>Treis</i> , three, Gr.	-	-	107
TRIGONIA.	<i>Trigonos</i> , Gr., triangular	-	-	97
TRITON.	A Roman sea-god	-	-	228
TRITONIA	-	-	-	140
TROCHUS.	A top	-	-	211
TUBICINELLA.	<i>Tubicen</i> , a trumpeter	-	-	37
TURBINELLA.	A little top	-	-	222
TURBO.	A top	-	-	214
TURRILITES.	<i>Turris</i> , a tower	-	-	268
TURRITELLA.	A little tower	-	-	217
UMBRELLA.	A little shade	-	-	147
UNGULINA.	<i>Unguis</i> , a nail or hoof	-	-	76
UNIO.	A pearl	-	-	100
VALVATA.	<i>Valvata</i> , having valves	-	-	188
VENERUPIS.	<i>Venus</i> , and <i>rupis</i> , a rock	-	-	78
VENUS.	Goddess of beauty	-	-	90
VERMETUS.	<i>Vermis</i> , a worm	-	-	205
VERMILIA.	Do.	-	-	35
VITRINA.	<i>Vitreus</i> , glassy	-	-	162
VOLUTA.	<i>Volvo</i> , to revolve	-	-	247
VOLVARIA.	<i>Volva</i> , a shuttle	-	-	251
VULSELLA.	A little tongue	-	-	117
<i>Vernericardia</i> - see <i>Cardita</i>				
XYLOPHAGA.	<i>Xulon</i> , wood; <i>phago</i> , to eat	-	-	61

ENGLISH NAMES

IN THIS VOLUME, AND IN COMMON USE.

		Page			Page
<i>Barnacles</i>	-	- 38	<i>Oysters</i>	-	- 126
<i>Cockles</i>	-	- 91	<i>Pearls</i>	- -	99. 119
<i>Cowries</i>	-	- 252	<i>Periwinkles</i>	-	- 215
<i>Cuttle-fish</i>	-	- 273	<i>Scallop-shells</i>	-	- 124
<i>Ear-shells</i>	-	- 200	<i>Slugs</i>	-	- 160
<i>Fan-shells</i>	-	- 123	<i>Snails</i>	-	- 165
<i>Limpets</i>	-	- 143	<i>Staircase-shells</i>		208. 211
<i>Mussels</i>	-	100. 112	<i>Whelks</i>	-	- 242

EXPLANATION OF TERMS.

Acephalous. Mollusca without an apparent head ; comprehending the Bivalves of conchologists.

Acuminated. Ending in a graduated, and often acute, point.

Adductor Muscle. That which closes the two pieces of a Bivalve together, the base or insertion of which is indicated by an irregular depression in each valve, these being called the muscular impressions. (See *Muscular Impressions*.)

Annulated. Marked with rings.

Anterior. The side of the shell, in Bivalves, where the ligament is situated. (See *Posterior*.)

Aperture. The mouth or opening of the shell, in Univalves, from which the head of the animal protrudes.

Apex. The commencement or top of the spire in Univalves.

Articulated. Jointed.

Auriculated. (*Eared.*) A term given to a few Bivalves which have a flat angulated projection on one or both sides of the umbones or bosses : these processes are most developed in the *Pectens*.

Auriform. (*Ear-shaped.*) Shaped like an ear ; as the *Haliotis*.

Base. The extremity opposite to the *Apex*, in Univalves : in such Bivalves as adhere to other substances by one of their valves, as in *Spondylus*, *Anomia*, &c., that valve is termed the *basal*.

Bicarinated. Having two elevated or sharp ridges. (See *Carinated*.)

Bicornute. Like, or having two horns.

Bifid. Cleft in two.

Boss. The projecting point in each valve, near the hinge in Bivalves ; the commencement of the shell.

Byssus. A bunch of silk-like fibres, by which many species of Bivalves are attached to other substances.

Callosity. A thick mass of shelly matter.

Canal. A groove or gutter observable in different parts of certain spiral shells belonging to the *Zoophaga*, or carnivorous tribe. In these the canal is either lengthened, as in *Murex*, short, as in *Harpa*, or truncate, as in *Buccinum* : in the two latter it thus gives place to a notch, but in all these instances, it is that part fitted for the protrusion of the long cylindrical siphon possessed by all these animals.

Cancellated. When the surface is marked by striæ or lines, which cross each other at right angles.

Cardinal Teeth, in Bivalves, are those placed immediately beneath the bosses, and between the lateral teeth, where such exist, as in the common Cockle.

Carinated. Ridged like the keel of a boat.

Cartilaginous. Gristly, like a tendon.

Chambered. Divided internally into compartments, as in the testaceous *Cephalopoda*, or *Nautilus*. (*Polythalamous*.)

Columella (or *Pillar*). The internal support of most

spiral shells, round which the whorls convolute ; it is not present in *Solarium*, *Delphinula*, &c.

Concentric. Arranged in parallel curves or bands, having one common centre.

Convolute. The whorls rolled regularly one over the other.

Cordate. Heart-shaped.

Corneous. Resembling the colour or substance of horn ; the epidermis of some, and the operculum of other spiral shells often present this appearance.

Coronated. Crowned, towards the *apex*, by a row of spines.

Crenated. Notched.

Cuneiform. Wedge-shaped, as *Donax*.

Dentated. Toothed.

Dextral. Spiral shells are so called when the mouth faces the *right* hand of the observer, the shell being held with the *apex* upwards. The great majority of spiral shells are dextral, but frequently in the very same species (as in *Bulimus*) individuals occur whose aperture is reversed, or appears on the *left* side. (See *Sinistral*.)

Digitated. The expansion of any particular part, as in the outer lip of the *Pteroceras*, into finger-shaped processes.

Discoid. Disk-shaped, much flattened. The fresh-water genus *Planorbis*, and many of the *Helicidæ*, or land-shells, are of this form.

Dorsal. Belonging to the back.

Effuse. Having the lips separated by a gutter or channel.

Entire. Having the mouth undivided at the base by a channel.

Epidermis. The membranous covering of some shells.

Epiphragma. A covering to the aperture of some spiral shells, which (unlike the *operculum*) is not attached to the animal. (See *Operculum*.)

Equilateral. When the two sides of a Bivalve are equal, the bosses being nearly or quite in the middle; as in the *cockle*.

Equivalve. When the two valves of a Bivalve are equal in size; as in the *cockle*.

Eroded. Gnawed, or rubbed.

Fibrous. Resembling fibres: applied to the substance of a shell, it indicates those, like the *Pinna*, whose fracture presents perpendicular fibres.

Fluviatile. Living in fresh water.

Foliated. Composed of thin plates, lying on each other, as in the shell of the *oyster*.

Fusiform. Shaped like a spindle, swollen in the middle and tapering to each end.

Gaping. In Bivalves, when, the shell being closed, the valves do not touch each other in every part.

Gelatinous. Like jelly.

Hinge. The part where the valves of a Bivalve are united, consisting of ligament and teeth.

Hyaline. Of a glassy, thin, and semi-transparent substance.

Inequilateral. When the sides of a Bivalve are unequal in length. (See *Equilateral*.)

Inequivalve. When the valves of a Bivalve are unequal in size and shape, as the *oyster*. (See *Equivalve*.)

Inferior Valve. In adherent Bivalves, that by which the shell is attached to other substances.

Iridescent. Pearly, or having the appearance of pearl.

Irregular. In Bivalves, when the individuals of a species are not all of the same shape.

Lanceolate. Shaped like the head of a lance.

Lateral. Any thing on the sides of a shell. Thus the lateral teeth of Bivalves, where they exist, are on one or both sides of the cardinal teeth, which are always in the centre, under the bosses.

Length of Shells. Spiral shells are measured from the *apex*, or tip of the spire to the *base*, and therefore perpendicularly ; but the length of Bivalves is taken horizontally, or from the anterior to the posterior margin.

Ligament. A cartilage connecting the two valves of a Bivalve.

Lips. The two sides of the aperture of spiral shells are termed the lips. That which joins, and generally folds over the lower part of the columella is called the *inner* lip, while that part of the circumference opposite is called the *outer* lip.

Litoral. Belonging to, or found on the shore.

Lobed. Having broad finger-like divisions.

Longitudinal. In Univalves, stripes or marks from the *apex* to the *base* are called *longitudinal*, in opposition to *transverse*, which is applied to such as cross the shell, or are parallel to the whorls : in Bivalves, stripes from the anterior to the posterior margin are *transverse* ; from the bosses to the ventral margin *longitudinal*, or more properly *radiated*.

Marine. Belonging to, or found in the sea.

Monothalamous. Having only one chamber, in opposition to *Polythalamous* : the *Argonauta* is an instance.

Muscular Impressions. The indented marks in acephalous Bivalves, which indicate the insertion of the muscles, by which the animal is attached to its shell. Of these there are three principal sorts : lateral, central, and pallial. 1. The *lateral* impressions indicate the animal to have two adductor muscles, as in *Unio*, *Cardium*. 2. A *central* impression shows the presence of one adductor muscle, which is generally in the middle of the shell, as the *Oyster*. All these shells are, however, furnished with a third or *pallial* impression, which is indicated by a depressed line running parallel with the ventral margin. (See *Adductor Muscle*.)

Obsolete. This term implies a faint indication of particular characters. Thus, very slight or partially indistinct striæ upon a shell are called *obsolete* striæ, not because they are worn off by accident, but because they are only rudimentary, or very slightly developed.

Operculum. A hard shelly or horny lid, which covers the aperture of many spiral shells, and is attached to the foot of the animal. (See *Epiphragma*.)

Orbicular. Quite round.

Ovate. Egg-shaped.

Papyraceous. Excessively thin, like paper.

Peduncle. A stalk, or support.

Pelagic. Belonging to, or found in, deep seas.

Perforated. Having holes, as if bored by a sharp instrument, as the *Haliotis*.

Phosphorescent. Shining in the dark, like the glow-worm.

Plaits. Folds, or raised lines on the *Columella*.

Polythalamous. (See *Chambered* and *Monothalamous*.)

Posterior. The side of the shell, in Bivalves, opposite to

that on which the ligament is placed. (See *Anterior*.)

Recurved. Turned backwards. The canal of some of the genera *Strombus* and *Cerithium* is remarkably so.

Reflected. Turned or folded backwards, as in the thickened lip of the common *Helix*.

Retractile. Capable of being drawn backwards.

Retuse. When a spiral shell is retuse, the inner whorls appear to have been pressed into the body of the shell, and the *apex* is below the level of the last whorl.

Reversed. The same as *sinistral*. (See *Sinistral*.)

Sessile. Attached to any body, without a stalk or peduncle.

Sinistral. Having the aperture on the *left* side, as opposed to *dextral* (see *Dextral*), also called *reversed*.

Sinuuous. Waved.

Sinus. A gutter, or curvature, in the lip of Univalves.

Siphon. A fleshy sucker.

Spire. Those volutions of a spiral shell which are above the lowest or body whorl collectively form the *spire* of the shell, whatever shape it may assume. The variations in the form of this part are numerous. In some it is excessively lengthened ; in others it is not even raised above the body whorl. In *Planorbis* the spire is sunk, while in *Cypræa* it is so small as only to be seen when the shell is young, after which it is covered over by the enlargement of the body whorl or principal volution.

Striated. Marked with lines, either elevated or raised, at parallel and nearly equal distances.

Sub. Almost, or approaching to ; as sub-fusiform (nearly fusiform), &c.

Sutures. The corkscrew line formed in the spire, by the whorls wrapping over each other.

Tentacula. Arms or feelers.

Terebrating Testacea. Such as take up their habitations in other substances, as *Pholas*, &c.

Testaceous. Composed of the materials which constitute shell.

Transverse. See *Longitudinal*.

Truncated. Cut short off, or ending abruptly.

Tuberculated. Covered with small protuberances.

Turbinate. Having the body of the shell large or swollen, and the spire comparatively short.

Turreted. Having the whorls gradually decreasing, so as to form a fine point at the *apex*.

Umbilicus. A small hollow, or sometimes a deep hole, on the side of the inner lip in spiral shells, formed by the inner edges of the whorls not touching each other: in some genera, as *Scalaria* and *Solarium*, it is so deep as to extend to the *apex* of the shell; in others, as *Trochus*, it is small; in *Bulimus* it becomes nearly obsolete; and in *Natica* it is often entirely concealed by the spreading of the inner lip.

Varices. Longitudinal raised bands or ridges, which occur at regular distances in some Univalves, as *Murex*, *Triton*, &c. They mark the progressive enlargement of the shell, being the remnants of former apertures, beyond which the animal, when grown, forms another aperture.

The greater part of these explanations are taken from Mr. Swainson's *Malacology*.

THE END.

LONDON:

Printed by A. SPOTTISWOODE,
New-Street-Square.

May, 1846.

A CATALOGUE OF NEW WORKS AND NEW EDITIONS

PRINTED FOR

MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,
LONDON.

CLASSIFIED INDEX.

AGRICULTURE & RURAL AFFAIRS.

	Pages
Bayldon on Valuing Rents, etc.	6
Crocker's Land Surveying	9
Day's Agricultural Chemistry	9
Greenwood's (Col.) Tree-Lifter	12
Johnson's Farmer's Encyclopædia	15
Loudon's Encyclopædia of Agriculture	18
„ Self-Instruction for Farmers, etc.	17
(Mrs.) Lady's Country Companion	17
Low's Breeds of the Domesticated Animals	19
„ of Great Britain	19
„ Elements of Agriculture	19
„ On Landed Property	18
„ On the Domesticated Animals	18
Whitley's Agricultural Geology	32

ARTS, MANUFACTURES, AND ARCHITECTURE.

Brande's Dictionary of Science, Literature, and Art	6
Budge's Miner's Guide	7
De Burtin on the Knowledge of Pictures	9
Eastlake's History of Oil Painting	10
Gruner's Decorations of the Queen's Pavilion	12
Gwilt's Encyclopædia of Architecture	12
Haydon's Lectures on Painting & Design	13
Holland's Manufactures in Metal	13
Lerebours On Photography	17
Loudon's Encyclopædia of Cottage, Farm, and Villa Architecture and Furniture	18
Maitland's Church in the Catacombs	20
Porter's Manufacture of Silk	24
„ Porcelain and Glass	24
Reid (Dr.) on Warming and Ventilating	25
Steam Engine (The), by the Artisan Club	25
Ure's Dictionary of Arts, Manufactures, and Mines	31

BIOGRAPHY.

Aikin's Life of Addison	5
Bell's Lives of the British Poets	6
Dover's Life of the King of Prussia	10
Dunham's Early Writers of Britain	10
„ Lives of the British Dramatists	10
Forster's Statesmen of the Commonwealth of England	11
Gleig's Lives of the most Eminent British Military Commanders	11
Grant (Mrs.) Memoir and Correspondence	11
James's Life of the Black Prince	15
„ Eminent Foreign Statesmen	15
Lál's (M.) Life of Dost Mahomed	21
Leslie's Life of Constable	17
Life of a Travelling Physician	17
Mackintosh's Life of Sir T. More	19
Mauder's Biographical Treasury	21
Mignet's Antonio Perez and Philip II.	21
Roberts's Life of the Duke of Monmouth	25
Roscoe's Lives of Eminent British Lawyers	25
Russell's Bedford Correspondence	26
Shelley's Eminent Literary Men of Italy, etc.	27
„ Eminent French Writers	27

	Pages
Southey's Lives of the British Admirals	28
„ Life of Wesley	28
Townsend's Lives of Twelve eminent Judges	30
Waterton's Autobiography and Essays	31

BOOKS OF GENERAL UTILITY.

Acton's (Eliza) Cookery Book	5
Black's Treatise on Brewing	6
„ Supplement on Bavarian Beer	6
Collegian's Guide	8
Donovan's Domestic Economy	10
Hand-Book of Taste	12
Hints on Etiquette	13
Hudson's Parent's Hand-Book	14
„ Executor's Guide	14
„ On Making Wills	14
Loudon's Self Instruction	17
Mauder's Treasury of Knowledge	20
„ Scientific and Literary Treasury	21
„ Treasury of History	21
„ Biographical Treasury	21
„ Universal Class-Book	21
Parkes's Domestic Duties	23
Pycroft's Course of English Reading	24
Riddle's Eng.-Lat. and Lat.-Eng. Dict.	25
Robinson's Art of Curing, Pickling, etc.	25
Short Whist	27
Thomson's Management of Sick Room	30
„ Interest Tables	30
Tomlins' Law Dictionary	30
Webster's Encycl. of Domestic Economy	31

BOTANY AND GARDENING.

Abercrombie's Practical Gardener	5
„ and Main's Gardener's Companion	5
Calcott's Scripture Herbal	7
Conversations on Botany	8
Drummond's First Steps to Botany	10
Glendinning On the Pine Apple	11
Greenwood's (Col.) Tree-Lifter	12
Grimblot's William III. and Louis XIV.	12
Henslow's Botany	13
Hoare On the Grape Vine on Open Walls	13
„ On the Roots of Vines	13
Hooker's British Flora	13
„ and Taylor's Muscologia Britannica	13
Jackson's Pictorial Flora	15
Lindley's Theory of Horticulture	17
„ Orchard and Kitchen Garden	17
„ Introduction to Botany	17
„ Flora Medica	17
„ Synopsis of British Flora	17
Loudon's Hortus Britannicus	18
„ Hortus Lignosus Londinensis	18
„ Encyclopædia of Trees & Shrubs	18
„ Gardening	18
„ Plants	18
„ Suburban Gardener	18
„ Self-Instruction for Gardeners, etc.	17
Repton's Landscape Gardening and Landscape Architecture	25

	Pages
Rivers's Rose Amateur's Guide	25
Rogers's Vegetable Cultivator	25
Schleiden's Scientific Botany	26
Smith's Introduction to Botany	27
„ English Flora	27
„ Compendium of English Flora	27

CHRONOLOGY.

Blair's Chronological Tables	6
Calendar (Illuminated) and Diary	15
Nicolas's Chronology of History	23
Riddle's Ecclesiastical Chronology	25
Tate's Horatius Restitutus	29

COMMERCE AND MERCANTILE AFFAIRS.

Gilbart On Banking	11
Lorimer's Letters to a Master Mariner	17
McCulloch's Dictionary of Commerce	19
Steel's Shipmaster's Assistant	28
Thomson's Tables of Interest	30
Walford's Customs' Laws	31

GEOGRAPHY AND ATLASES.

Butler's Sketch of Ancient and Modern Geography	7
„ Atlas of Modern Geography	7
„ Ancient Geography	7
Cooley's World Surveyed	8
De Strzelecki's New South Wales	9
Forster's Historical Geography of Arabia	11
Hall's New General Atlas	12
McCulloch's Geographical Dictionary	19
Murray's Encyclopædia of Geography	22
Ordnance Maps, and Publications of the Geological Society	23
Parrot's Ascent of Mount Ararat	8

HISTORY AND CRITICISM.

Adair's (Sir R.) Mission to Vienna	5
„ Negotiations for the Peace of the Dardanelles	5
Addison's History of the Knights Templars	5
Bell's History of Russia	6
Blair's Chron. and Historical Tables	6
Bloomfield's Translation of Thucydides	6
„ Edition of Thucydides	6
Bunsen's Egypt	7
Cooley's Maritime and Inland Discovery	8
Crowe's History of France	9
Dahlmann's English Revolution	9
De Sismondi's Fall of the Roman Empire	9
„ Italian Republics	9
Dunham's History of Spain and Portugal	10
„ Europe in the Middle Ages	10
„ History of the German Empire	10
„ Denmark, Sweden, and Norway	10
„ History of Poland	10
Dunlop's History of Fiction	10
Eccleston's English Antiquities	10
Fergus's History of United States of America	10
Grant (Mrs.) Memoir and Correspondence	11
Grattan's History of Netherlands	11
Guicciardini's Hist. Maxims	12
Haisted's Life of Richard III.	12
Haydon's Lectures on Painting and Design	13
Historical Pictures of the Middle Ages	13
Horsley's (Bp.) Biblical Criticism	14
Jeffrey's (Lord) Contributions to the Edinburgh Review	15
Keightley's Outlines of History	15
Laing's Kings of Norway	16
Lemprière's Classical Dictionary	17
Macauley's Essays	19
Mackinnon's History of Civilisation	19
Mackintosh's History of England	19
„ Miscellaneous Works	19
McCulloch's Dictionary, Historical, Geographical, and Statistical	19
Maunder's Treasury of History	21

	Pages
Mignet's Antonio Perez and Philip II.	21
Milner's Church History	21
Moore's History of Ireland	21
Mosheim's Ecclesiastical History	22
Müller's Mythology	22
Nicolas's Chronology of History	23
Ranke's History of the Reformation	24
Roberts's Rebellion, etc. of the Duke of Monmouth	25
Rome, History of	25
Russell's Correspondence of the Duke of Bedford	6
Scott's History of Scotland	26
Stebbing's History of the Christian Church	28
„ History of the Reformation	28
„ Church History	23
Switzerland, History of	29
Sydney Smith's Works	27
Thirlwall's History of Greece	29
Tooke's History of Prices	30
Turner's History of England	30
Tytler's Elements of General History	31
Zumpt's Latin Grammar	32

JUVENILE BOOKS.

Including Mrs. Marcet's Works.

Boy's (the) Own Book	6
Conscience's Flemish Sketches	8
Hawes's Tales of the N. American Indians	13
Howitt's Boy's Country Book	14
Mackintosh's Life of Sir T. More	19
Marcet's Conversations—	
On the History of England	20
On Chemistry	20
On Natural Philosophy	20
On Political Economy	20
On Vegetable Physiology	20
On Land and Water	20
On Language	20
Marcet's Game of Grammar	20
„ Willy's Grammar	20
„ Lessons on Animals, etc.	20
Marryat's Masterman Ready	20
„ Settlers in Canada	20
„ Mission; or, Scenes in Africa	20
Maunder's Universal Class Book	21
Pycroft's (the Rev. J.), English Reading	24

MEDICINE.

Bull's Hints to Mothers	7
„ Management of Children	7
Copland's Dictionary of Medicine	8
Elliottson's Human Physiology	10
Holland's Medical Notes	13
Lefevre (Sir Geo.) on the Nerves	17
Pereira On Food and Diet	23
Reece's Medical Guide	24
Sandby On Mesmerism	26
Wigan (Dr.) On Insanity	32

MISCELLANEOUS

Adshead on Prisons and Prisoners	5
Bray's Philosophy of Necessity	7
Claviers's Forest Life	8
Cocks's Bordeaux, etc.	8
Collegian's Guide	8
Colton's Lacon	8
De Burtin On the Knowledge of Pictures	9
De Morgan On Probabilities	9
De Strzelecki's New South Wales	9
Dunlop's History of Fiction	10
Good's Book of Nature	11
Graham's English	11
Grant's Letters from the Mountains.	11
Guest's Mabinogion	12
Hand-Book of Taste	12
Hobbes's (Thos.) complete Works	13
Howitt's Rural Life of England	14
„ Visits to Remarkable Places	14
„ Student-Life of Germany	14

	Pages
Howitt's Rural and Social Life of Germany	14
„ Colonisation and Christianity	14
Humphreys' Illuminated Books	15
Illuminated Calendar and Diary for 1845	15
Jeffrey's (Lord) Contributions to the	
Edinburgh Review	15
Lane's Life at the Water Cure	16
Lefevre (Sir Geo.) On the Nerves	17
Life of a Travelling Physician	17
Loudon's (Mrs.) Lady's Country Companion	17
Macaulay's Critical and Historical Essays	19
Mackintosh's (Sir J.) Miscellaneous Works	19
Michelet's Priests, Women, and Families	
„ The People	21
Müller's Mythology	22
Necker De Sanssure's Progressive Education	22
Perry On German University Education	23
Peter Plymley's Letters	24
Pycroft's English Reading	24
Rowton's Debater	26
Sandby On Mesmerism	26
Sandford's Parochialia	26
Seaward's Narrative of his Shipwreck	26
Smith's (Rev. Sydney) Works	27
Southery's Common-Place Book	28
Taylor's Statesman	29
Walker's Chess Studies	31
Welsford On Language	32
Wigan (Dr.) On Insanity	32
Willoughby's (Lady) Diary	32
Zumpt's Latin Grammar	32

NATURAL HISTORY IN GENERAL.

Catlow's Popular Conchology	7
Doubleday's Butterflies and Moths	10
Drummond's Letters to a Naturalist	10
Gray's Figures of Molluscous Animals	12
„ Mammalia	12
„ and Mitchell's Ornithology	12
Kirby and Spence's Entomology	16
Lee's Taxidermy	17
„ Elements of Natural History	17
Marcet's Conversations on Animals, etc.	20
Newell's Zoology of the English Poets	22
Stephens' British Coleoptera	28
Swainson on the Study of Natural History	29
„ Animals	29
„ Quadrupeds	29
„ Birds	29
„ Animals in Menageries	29
„ Fish, Amphibians, & Reptiles	29
„ Insects	29
„ Malacology	29
„ the Habits and Instincts of	
„ Animals	29
„ Taxidermy	29
Turton's Shells of the British Islands	31
Waterton's Essays on Natural History	31
Westwood's Classification of Insects	32
Zoology of H.M. S.s' Erebus and Terror	32

NOVELS AND WORKS OF FICTION.

Bray's (Mrs.) Novels	7
Conscience's Flemish Sketches	8
Doctor (The)	9
Dunlop's History of Fiction	10
Margaret Russell	20
Marryat's Masterman Ready	20
„ Settlers in Canada	20
„ Mission; or, Scenes in Africa	20
Pericles, A Tale of Athens	23
Willis's (N.P.) Dashes at Life	32

ONE VOLUME ENCYCLOPÆDIAS AND DICTIONARIES.

Blaine's, of Rural Sports	6
Braude's, of Science, Literature, and Art	6
Copland's, of Medicine	8

	Pages
Cresy's, of Civil Engineering	9
Gwilt's, of Architecture	12
Johnson's Farmer	15
Loudon's, of Trees and Shrubs	18
„ of Gardening	18
„ of Agriculture	18
„ of Plants	18
„ of Rural Architecture	18
M'Culloch's Geographical, Statistical, and	
Historical Dictionary	19
„ Commerce	19
Murray's Encyclopedia of Geography	22
Ure's Arts, Manufactures, and Mines	31
Webster's Domestic Economy	31

POETRY AND THE DRAMA.

Aikin's (Dr.) British Poets	26
Bowdler's Family Shakspeare	26
Chaloner's Walter Gray	8
„ Poetical Remains	8
Collier's Roxburghie Ballads	8
Costello's Persian Rose Garden	8
Dante, translated by Wright	9
Goldsmith's Poems	11
Gray's Elegy, illuminated	11
Heron's Palestrina	13
Horace, by Tate	29
L. E. L.'s Poetical Works	16
Linwood's Anthologia Oxoniensis	17
Macaulay's Lays of Ancient Rome	19
Mackay's English Lakes	19
Montgomery's Poetical Works	21
Moore's Poetical Works	21
„ Lalla Rookh	22
„ Irish Melodies	22
Moral of Flowers	22
Poet's Pleasance	24
Pope's Works	24
Reynard the Fox	25
Sheldon's Minstrelsy	27
Sophocles, by Linwood	27
Southery's Poetical Works	28
„ Oliver Newman	28
„ British Poets	26
Spirit of the Woods	28
Thomson's Seasons	29
Turner's Richard III.	30
Watts's (A.A.) Lyrics of the Heart	31

POLITICAL ECONOMY AND STATISTICS.

Gilbart on Banking	11
M'Culloch's Geographical, Statistical, and	
Historical Dictionary	19
M'Culloch's Literature of Polit. Economy	19
„ On Taxation and Funding	19
„ Statistics of the British Empire	19
Strong's Greece as a Kingdom	28
Tooke's History of Prices	30
Twiss's Oregon Question Examined	31

RELIGIOUS AND MORAL WORKS, ETC.

Amy Herbert, edited by Rev. W. Sewell	5
Bailey's Essays on the Pursuit of Truth	5
Bloomfield's Greek Testament	6
„ College and School ditto	6
„ Lexicon to Greek Testament	6
Bunsen's Church of the Future	7
Burns's Christian Philosophy	7
„ Christian Fragments	7
Calcott's Scripture Herbal	7
Cooper's Sermons	8
Dale's Domestic Liturgy	9
Dibdin's Sunday Library	28
Doddridge's Family Expositor	10
Englishman's Hebrew and Chaldee Con-	
cordance of the Bible	16
„ Greek Concordance of the	
New Testament	10

	Pages		Pages
Fitzroy's (Lady) Scripture Conversations	11	Holland's Manufactures in Metal	13
Forster's Historical Geography of Arabia	11	Humboldt's Kosmos	14
" Life of Bishop Jebb	11	Hunt's Researches on Light	15
Gertrude, edited by the Rev. W. Sewell	11	Kane's Elements of Chemistry	15
Hook's (Dr.) Lectures on Passion Week	13	Kater and Lardner's Mechanics	15
Horne's Introduction to the Study of the		La Place's System of the World	16
Scriptures	14	Lardner's Cabinet Cyclopædia	16
Compendium of ditto	14	" Hydrostatics and Pneumatics	16
Horsley's (Bp.) Biblical Criticism	14	" and Walker's Electricity	16
Psalms	14	Lardner's Arithmetic	16
Jebb's Protestant Kempis	15	" Geometry	16
" Pastoral Instructions	15	" Treatise on Heat	16
" Correspondence with Knox	15	Lerebours On Photography	17
Knox's (Alexander) Remains	16	Lloyd On Light and Vision	17
Laing's Notes on the German Catholic		Mackenzie's Physiology of Vision	19
Schism	16	Marce's Conversations on Sciences, etc.	20
Laneton Parsonage	16	Moseley's Practical Mechanics	22
Maitland's Church in the Catacombs	20	" Engineering and Architecture	22
Marriage Gift	20	Narrien's Elements of Geometry	26
Michelet's Priests, Women, and Families	21	" Astronomy and Geodesy	26
" and Quinet's Jesuits	21	Nesbit's Mensuration	22
Milner's Church History	22	Owen's Lectures On Comparative Anatomy	23
Mosheim's Ecclesiastical History	21	Parnell On Roads	23
Parables (The)	23	Pearson's Practical Astronomy	23
Parkes's Domestic Duties	23	Peschel's Physics	24
Peter Plymley's Letters	24	Phillips's Palæozoic Fossils of Cornwall, etc.	24
Pitman's Sermons on the Psalms	24	" Guide to Geology	24
Riddle's Letters from a Godfather	25	" Treatise on Geology	24
Robinson's Greek and English Lexicon		" Introduction to Mineralogy	24
to the New Testament	25	Poisson's Mechanics	24
Sandford On Female Improvement	26	Portlock's Report on the Geology of	
" On Woman	26	Londonderry	24
" 's Parochialia	26	Powell's Natural Philosophy	24
Sermon on the Mount (The)	26	Quarterly Journal of the Geological Society	24
Shepherd's Horse Apostolicae	27	Ritchie (Robert) on Railways	25
Smith's Female Disciple	27	Roberts's Dictionary of Geology	25
" (G.) Perilous Times	27	Sandhurst Mathematical Course	26
" Religion of Ancient Britain	27	Scott's Arithmetic and Algebra	26
" (S.) Sermons	27	" Trigonometry	26
Southey's Life of Wesley	28	Thomson's Algebra	29
Stebbing's Church History	28	Wilkinson's Engines of War	32
Tate's History of St. Paul	29		
Taylor's (Rev. C. B.) Margaret; or, the Pearl	29		
" " Sermons	29		
" " Dora Melder	29		
" " Lady Mary	29		
Taylor's (Jeremy) Works	29		
Tomline's Christian Theology	30		
" Introduction to the Bible	30		
Trollope's Analecta Theologica	30		
Turner's Sacred History	30		
Wardlaw On Socinian Controversy	31		
Weil's Bible, Koran, and Talmud	31		
Whitley's Life Everlasting	32		
Wilberforce's View of Christianity	32		
Willoughby's (Lady) Diary	32		

RURAL SPORTS.

Blaine's Dictionary of Sports	6
Hansard's Fishing in Wales	12
Hawker's Instructions to Sportsmen	13
Loudon's (Mrs.) Lady's Country Companion	17
Stable Talk and Table Talk	28

THE SCIENCES IN GENERAL, AND MATHEMATICS.

Bakewell's Introduction to Geology	5
Balmain's Lessons on Chemistry	6
Brande's Dictionary of Science, Literature, and Art	6
Brewster's Optics	7
Conversations on Mineralogy	8
De la Beche on the Geology of Cornwall, etc.	9
Donovan's Chemistry	10
Elliot's Geometry	10
Farey on the Steam Engine	10
Fosbroke on the Arts of the Ancients	11
Greener on the Gun	12
Herschel's Natural Philosophy	13
" Astronomy	13

TRAVELS.

Allan's Mediterranean	5
Beale's Vale of the Towey	6
Cooley's World Surveyed	8
Costello's (Miss) North Wales	9
De Custine's Russia	9
De Strzelecki's New South Wales	9
Erman's Travels through Siberia	8
Harris's Highlands of Æthiopia	12
Howitt's (R.) Australia Felix	14
Laing's Notes of a Traveller	16
" Residence in Norway	16
" Tour in Sweden	16
Life of a Travelling Physician	17
Mackay's English Lakes	19
Montauban's Wanderings	21
Parrot's Ascent of Mount Ararat	8
Paton's (A. A.) Servia	23
" Modern Syrians	23
Pedestrian Reminiscences	23
Seaward's Narrative of his Shipwreck	26
Strong's Greece as a Kingdom	28
Von Örlich's Travels in India	31

VETERINARY MEDICINE

Centaur's Two Books on the Horse	8
Field's Veterinary Records	11
Morton's Veterinary Toxicological Chart	22
" Medicine	22
Miles On the Horse's Foot	21
Percivall's Hippopathology	23
" Anatomy of the Horse	23
Spooner on the Foot and Leg of the Horse	28
Stable Talk and Table Talk	28
Turner On the Foot of the Horse	30
Winter On the Horse	32

NEW WORKS AND NEW EDITIONS.

ABERCROMBIE.—ABERCROMBIE'S PRACTICAL GARDENER, AND IMPROVED SYSTEM OF MODERN HORTICULTURE, alphabetically arranged. 4th Edition, with an Introductory Treatise on Vegetable Physiology, and Plates by W. Salisbury. 12mo. 6s. boards.

ABERCROMBIE AND MAIN.—THE PRACTICAL GARDENER'S COMPANION; Or, Horticultural Calendar: to which is added, the Garden-Seed and Plant Estimate. Edited from a MS. of J. Abercrombie, by J. Main. 8th Edition. 32mo. 3s. 6d. sewed.

ACTON (MISS).—MODERN COOKERY, In all its Branches, reduced to a System of Easy Practice. For the use of Private Families. In a Series of Practical Receipts, all of which have been strictly tested, and are given with the most minute exactness. Dedicated to the Young Housekeepers of England. By Eliza Acton. New Edition, improved. Foolsap 8vo. with Woodcuts, 7s. 6d. cloth.

ADAIR (SIR ROBERT).—AN HISTORICAL MEMOIR OF A MISSION TO THE COURT OF VIENNA IN 1806. By the Right Honorable Sir Robert Adair, G.C.B. With a Selection from his Despatches, published by permission of the proper Authorities. 8vo. 18s. cloth.

ADAIR (SIR ROBERT).—THE NEGOTIATIONS FOR THE PEACE OF THE DARDANELLES, in 1808–9; with Despatches and Official Documents. By the Right Honorable Sir Robert Adair, G.C.B. Being a Sequel to the Memoir of his Mission to Vienna in 1806. 2 vols. 8vo. 28s. cloth.

ADDISON.—THE KNIGHTS TEMPLARS.

By C. G. Addison, of the Inner Temple. 2d Edition, enlarged. Square crown 8vo. with Illustrations, 18s. cloth.

ADDISON.—THE TEMPLE CHURCH IN LONDON:

Its History and Antiquities. By C. G. Addison, Esq., of the Inner Temple, author of "The History of the Knights Templars." Square crown 8vo. with 6 Plates, 5s. cloth.

Also,
A FULL AND COMPLETE GUIDE, HISTORICAL AND DESCRIPTIVE, TO THE TEMPLE CHURCH. (From Mr. Addison's "History of the Temple Church.") Square crown 8vo. 1s. sewed.

ADSHEAD (JOSEPH).—PRISONS AND PRISONERS.

By Joseph Adshead. 8vo. with Illustrations, 7s. 6d. cloth.

AIKIN.—THE LIFE OF JOSEPH ADDISON.

Illustrated by many of his Letters and Private Papers never before published. By Lucy Aikin. 2 vols. post 8vo. with Portrait from Sir Godfrey Kneller's Picture, 18s. cloth.

ALLAN (J. H.).—A PICTORIAL TOUR IN THE MEDITERRANEAN;

Comprising Malta, Dalmatia, Turkey, Asia Minor, Grecian Archipelago, Egypt, Nubia, Greece, Sicily, Italy, and Spain. By J. H. Allan, Member of the Athenian Archæological Society, and of the Egyptian Society of Cairo. 2d Edition. Imperial 4to. with upwards of 40 lithographed Drawings, and 70 Wood Engravings, 3l. 3s. cloth.

AMY HERBERT.

By a Lady. Edited by the Rev. William Sewell, B.D. of Exeter College, Oxford. 3d Edition. 2 vols. foolscap 8vo. 9s. cloth.

ARTISAN CLUB (THE).—A TREATISE ON THE STEAM ENGINE.

By the Artisan Club. 4to. with 30 Steel Plates, etc., and about 350 Wood Engravings, 27s. cloth.

[Early in June.]

, Nos. 1 to 23, 1s. each; No. 24 (on June 1st), 2s. sewed. Back Numbers to complete Sets may still be had.

BAILEY.—ESSAYS ON THE PURSUIT OF TRUTH,

And on the Progress of Knowledge. By Samuel Bailey, author of "Essays on the Formation and Publication of Opinions," "Berkeley's Theory of Vision," etc. 2d Edition, revised and enlarged. 8vo. 9s. 6d. cloth.

BAKEWELL.—AN INTRODUCTION TO GEOLOGY.

Intended to convey Practical Knowledge of the Science, and comprising the most important recent Discoveries; with Explanations of the Facts and Phenomena which serve to confirm or invalidate various Geological Theories. By Robert Bakewell. Fifth Edition, considerably enlarged. 8vo. with numerous Plates and Woodcuts, 21s. cloth.

BALMAIN.—LESSONS ON CHEMISTRY,

For the Use of Pupils in Schools, Junior Students in Universities, and Readers who wish to learn the fundamental Principles and leading Facts: with Questions for Examination, Glossaries of Chemical Terms and Chemical Symbols, and an Index. By William H. Balmain. With numerous Woodcuts, illustrative of the Decompositions. Foolscap 8vo. 6s. cloth.

BAYLDON.—THE ART OF VALUING RENTS AND TILLAGES,

And the Tenant's Right of Entering and Quitting Farms, explained by several Specimens of Valuations; and Remarks on the Cultivation pursued on Soils in different Situations. Adapted to the Use of Landlords, Land-Agents, Appraisers, Farmers, and Tenants. By J. S. Bayldon. 6th Edition, corrected and revised by John Donaldson, 8vo. 10s. 6d. cloth.

BEALE (ANNE).—THE VALE OF THE TOWEY;

Or, Sketches in South Wales. By Anne Beale. Post 8vo. 10s. 6d. cloth.

"The great merit of Miss Beale's book is its truth, which leaves a strong impression on the reader's mind."—Spectator.

BEDFORD CORRESPONDENCE.—CORRESPONDENCE OF JOHN,

FOURTH DUKE OF BEDFORD, selected from the Originals at Woburn Abbey: with Introductions by Lord John Russell. 8vo. vol. 1 (1742-48), 18s. cloth; vol. 2 (1749-60), 15s. cl.

**** Vol. III. to complete the work, is nearly ready.*

BELL.—LIVES OF THE MOST EMINENT ENGLISH POETS.

By Robert Bell, Esq. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

BELL.—THE HISTORY OF RUSSIA,

From the Earliest Period to the Treaty of Tilsit. By R. Bell, Esq. 3 vols. foolscap 8vo. 18s.

BLACK.—A PRACTICAL TREATISE ON BREWING.

Based on Chemical and Economical Principles: with Formulæ for Public Brewers, and Instructions for Private Families. By William Black. Third Edition, revised and corrected, with considerable Additions. The Additions revised by Professor Graham, of the London University. 8vo. 10s. 6d. cloth.

SUPPLEMENT, of REMARKS on BAVARIAN BEER, London Porter, the Influence of Electricity on Fermentation, and other Subjects. By William Black. 8vo. 2s. 6d. sewed.

BLAINE.—AN ENCYCLOPÆDIA OF RURAL SPORTS;

Or, a complete Account, Historical, Practical, and Descriptive, of Hunting, Shooting, Fishing, Racing, and other Field Sports and Athletic Amusements of the present day. By Delabere P. Blaine, Esq., author of "Outlines of the Veterinary Art," "Canine Pathology," etc. etc. From nearly 600 Engravings on Wood, by R. Branston, from Drawings by Alken, T. Landseer, Dickes, etc. 8vo. 2l. 10s. cloth.

BLAIR'S CHRONOLOGICAL AND HISTORICAL TABLES,

From the Creation to the present Time: with Additions and Corrections from the most authentic Writers; including the Computation of St. Paul, as connecting the Period from the Exode to the Temple. Under the revision of Sir Henry Ellis, K.H., Principal Librarian of the British Museum. Imperial 8vo. 3ls. 6d. half-bound morocco.

BLOOMFIELD.—THE HISTORY OF THE PELOPONNESIAN WAR.

By Thucydides. Newly Translated into English, and accompanied with very copious Notes, Philological and Explanatory, Historical and Geographical. By the Rev. S. T. Bloomfield, D.D. F.S.A. 3 vols. 8vo. with Maps and Plates, 2l. 5s. boards.

BLOOMFIELD.—THE HISTORY OF THE PELOPONNESIAN^o WAR.

By Thucydides. A New Recension of the Text, with a carefully amended Punctuation; and copious Notes, Critical, Philological, and Explanatory, almost entirely original, but partly selected and arranged from the best Expositors: accompanied with full Indexes, both of Greek Words and Phrases explained, and matters discussed in the Notes. Illustrated by Maps and Plans, mostly taken from actual Survey. By the Rev. S. T. Bloomfield, D.D. F.S.A. 2 vols. 8vo. 38s. cloth.

BLOOMFIELD.—THE GREEK TESTAMENT:

With copious English Notes, Critical, Philological, and Explanatory. By the Rev. S. T. Bloomfield, D.D. F.S.A. 6th Edit. improved. 2 vols. 8vo. with a Map of Palestine, 40s. cloth.

BLOOMFIELD.—THE GREEK TESTAMENT FOR COLLEGES AND

SCHOOLS; with English Notes. By the Rev. S. T. Bloomfield, D.D. Fourth Edition, enlarged and improved, accompanied with a New Map of Syria and Palestine, adapted to the New Testament and Josephus, and an Index of Greek Words and Phrases explained in the Notes. 12mo. 10s. 6d. cloth.

BLOOMFIELD.—GREEK AND ENGLISH LEXICON TO THE NEW

TESTAMENT: especially adapted to the use of Colleges, and the Higher Classes in Public Schools; but also intended as a convenient Manual for Biblical Students in general. By Dr. Bloomfield. 2d Edition, enlarged, and improved. 12mo. on wider paper, 10s. 6d. cloth.

BOY'S OWN BOOK (THE):

A Complete Encyclopædia of all the Diversions, Athletic, Scientific, and Recreative, of Boyhood and Youth. 20th Edition. Square 12mo. with many Engravings on Wood, 6s. boards.

BRANDE.—A DICTIONARY OF SCIENCE, LITERATURE, AND ART;

Comprising the History, Description, and Scientific Principles of every Branch of Human Knowledge; with the Derivation and Definition of all the Terms in general use. Edited by W. T. Brande, F.R.S.L. and E.; assisted by J. Cauvin. The various departments by Gentlemen of eminence in each. 8vo. with Wood-engravings, 3l. cloth.

BRAY (MRS.)—MRS. BRAY'S NOVELS AND ROMANCES,

Revised and corrected by Mrs. Bray. In 10 vols. fcap 8vo., uniformly with the "Standard Novels," with Frontispieces and Vignettes from Designs and Sketches by the late Thomas Stothard, R.A.; C. A. Stothard, F.S.A.; Henry Warren, Esq.; etc. 3l. cloth; or the Works separately 6s. each, as follows:—

Vol. I. "*The White Hoods*," with portrait of the Author, a view of her residence, and General Preface to the Series;— Vol. II. "*De Foix*;"—Vol. III. "*The Protestant*;"—Vol. IV. "*Fitz of Fitz-*

Ford;"—Vol. V. "*The Talba*;"—Vol. VI. "*Warleigh*;"—Vol. VII. "*Trelawny*;"—Vol. VIII. "*Trials of the Heart*;"—Vol. IX. "*Henry de Pomeroiy*;"—Vol. X. "*Courtenay of Walreddon*."

BRAY.—THE PHILOSOPHY OF NECESSITY;

Or, the Law of Consequences as applicable to Mental, Moral, and Social Science. By Charles Bray. 2 vols. 8vo. 15s. cloth.

BREWSTER.—A TREATISE ON OPTICS.

By Sir David Brewster, LL.D. F.R.S. etc. New Edition. Foolsap 8vo. with vignette title, and 176 Woodcuts, 6s. cloth.

BUDGE (J.)—THE PRACTICAL MINER'S GUIDE:

Comprising a Set of Trigonometrical Tables adapted to all the purposes of Oblique or Diagonal, Vertical, Horizontal, and Traverse Dialling; with their application to the Dial. Exercise of Drifts, Lodes, Slides, Levelling, Inaccessible Distances, Heights, etc. By J. Budge. New Edition, considerably enlarged, 8vo. with Portrait of the Author, 12s. cloth.

BULL.—THE MATERNAL MANAGEMENT OF CHILDREN,

In HEALTH and DISEASE. By Thomas Bull, M.D. Physician Accoucheur to the Finsbury Midwifery Institution, etc. 2d Edition, revised and enlarged. Foolsap 8vo. 7s. cloth.

BULL.—HINTS TO MOTHERS,

For the Management of Health during the Period of Pregnancy and in the Lying-in Room; with an Exposure of Popular Errors in connexion with those subjects. By Thomas Bull, M.D. Physician Accoucheur to the Finsbury Midwifery Institution, etc. 4th Edition, revised and considerably enlarged. Foolsap 8vo. 7s. cloth.

BUNSEN (C. C. J.)—THE CHURCH OF THE FUTURE:

Or, an Inquiry into its Prospects, Constitution, and Import, with especial reference to the Church of Prussia. To which is appended, a Correspondence with the Right Hon. William Gladstone. By the Chevalier C. C. J. Bunsen. Translated, under the superintendence of the Chevalier Bunsen, by the Rev. Leopold J. Bernays, Fellow of St. John's College, Oxford. Post 8vo. *[In the press.]*

BUNSEN.—AN INQUIRY INTO THE HISTORY, ARTS AND SCIENCES,

LANGUAGE, WRITING, MYTHOLOGY, and CHRONOLOGY of ANCIENT EGYPT: with the peculiar Position of that Nation in reference to the Universal History of Mankind. By the Chevalier C. C. J. Bunsen. Translated from the German, under the Author's superintendence, by C. H. Cottrell, Esq.; with additional matter, furnished by the Author. 2 vols. 8vo. with numerous Plates. *[In the press.]*

BURNS.—THE PRINCIPLES OF CHRISTIAN PHILOSOPHY;

Containing the Doctrines, Duties, Admonitions, and Consolations of the Christian Religion. By John Burns, M.D. F.R.S. 5th Edition. 12mo. 7s. boards.

BURNS.—CHRISTIAN FRAGMENTS;

Or, Remarks on the Nature, Precepts, and Comforts of Religion. By John Burns, M.D. F.R.S. Professor of Surgery in the University of Glasgow, author of "The Principles of Christian Philosophy." Foolsap 8vo. 5s. cloth.

BUTLER.—A SKETCH OF MODERN AND ANCIENT GEOGRAPHY.

By Samuel Butler, D.D., late Lord Bishop of Lichfield and Coventry; and formerly Head Master of Shrewsbury School. New Edition, revised by his Son. 8vo. 9s. boards.

BUTLER.—AN ATLAS OF MODERN GEOGRAPHY.

Consisting of Twenty-three coloured Maps, from a New Set of Plates; with an Index of all the Names of Places, referring to the Latitudes and Longitudes. By the late Dr. Butler, Bishop of Lichfield. New Edition, corrected. 8vo. 12s. half-bound.

BUTLER.—AN ATLAS OF ANCIENT GEOGRAPHY.

Consisting of Twenty-three coloured Maps: with an Index of all the Names of Places, referring to the Latitudes and Longitudes. By the late Dr. Butler, Bishop of Lichfield. New Edition, corrected. 8vo. 12s. half-bound.

BUTLER.—A GENERAL ATLAS OF MODERN AND ANCIENT GEOGRAPHY.

Consisting of Forty-five coloured Maps, and copious Indices referring to the Latitudes and Longitudes. By the late Dr. Butler, Bishop of Lichfield. New Edition, corrected. 4to. 24s. half-bound.

CALLCOTT.—A SCRIPTURE HERBAL:

With upwards of 120 Wood Engravings. By Lady Callcott. Square crown 8vo. 1l. 5s. cloth.

CATLOW.—POPULAR CONCHOLOGY;

Or, the Shell Cabinet Arranged: being an Introduction to the modern System of Conchology; with a sketch of the Natural History of the Animals, an account of the Formation of the Shells, and a complete Descriptive List of the Families and Genera. By Agnes Catlow. Foolsap. 8vo. with 312 Woodcuts, 10s. 6d. cloth.

CENTAUR.—THE HORSE: AND HOW TO CHOOSE HIM.

By Centaur. Fcp. 8vo.

*[In the press.]***CENTAUR.—THE HORSE: AND HOW TO USE HIM.**

By Centaur. Fcp. 8vo.

*[In the press.]***CHALENOR.—WALTER GRAY,**

A Ballad, and other Poems; including the Poetical Remains of Mary Chalenor. 2d Edition, with Additions. Fcp. 8vo. 6s. cloth.

CHALENOR.—POETICAL REMAINS OF MARY CHALENOR.

Fcp. 8vo. 4s. cloth.

CLAVERS.—FOREST LIFE.

By Mary Clavers, an Actual Settler; author of "A New Home, Who 'll Follow?" 2 vols. fcap. 8vo. 12s. cloth.

COCKS (C.)—BORDEAUX, ITS WINES, AND THE CLARET COUNTRY.

By C. Cocks, B.L., Professor of the Living Languages in the Royal Colleges of France; Translator of the Works of Michelet, Mignet, and Quinet.

*[In the press.]***COLLECIAN'S GUIDE (THE);**

Or, Recollections of College Days; setting forth the Advantages and Temptations of a University Education. By **** ***, M.A., ——— Coll. Oxon. Post 8vo. 10s. 6d. cloth.

COLLIER (J. PAYNE).—A BOOK OF ROXBURCHE BALLADS.

Edited by John Payne Collier, Esq. Post 4to.

*[In the press.]***COLTON.—LACON; OR, MANY THINGS IN FEW WORDS.**

By the Rev. C. C. Colton. New Edition, 8vo. 12s. cloth.

CONSCIENCE (HENDRIK).—SKETCHES FROM FLEMISH LIFE.

In Three Tales. Translated from the Flemish of Hendrik Conscience. Square 8vo. with 130 Engravings on Wood, from designs by Flemish artists, 6s. cloth.

CONVERSATIONS ON BOTANY.

9th Edition, improved. Foolscape 8vo. with 22 Plates, 7s. 6d. cloth; with coloured Plates, 12s.

CONVERSATIONS ON MINERALOGY.

With Plates, engraved by Mr. and Mrs. Lowry, from Original Drawings. 3d Edition, enlarged. 2 vols. 12mo. 14s. cloth.

COOLEY.—THE WORLD SURVEYED IN THE NINETEENTH CENTURY;

Or, Recent Narratives of Scientific and Exploring Expeditions (chiefly undertaken by command of Foreign Governments). Collected, translated, and, where necessary, abridged, by W. D. Cooley, Esq., author of "The History of Maritime and Inland Discovery" in the Cabinet Cyclopædia, etc.

The First Volume contains "The Ascent of Mount Ararat." By Dr. Friedrich Parrot, Professor of Natural Philosophy in the University of Dorpat, Russian Imperial Councillor of State, etc. 8vo. with a Map by Arrowsmith, and Woodcuts, 14s. cloth.

*** Each volume will form, for the most part, a Work complete in itself, and the whole Series will present an accurate and luminous picture of all the known portions of the earth. The Second Work of the Series, "Erman's Travels through Siberia," is in the press, in 2 vols. 8vo.

COOLEY.—THE HISTORY OF MARITIME AND INLAND DISCOVERY.

By W. D. Cooley, Esq. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

COOPER (REV. E.)—SERMONS,

Chiefly designed to elucidate some of the leading Doctrines of the Gospel. To which is added, an Appendix, containing Sermons preached on several Public Occasions, and printed by desire. By the Rev. Edward Cooper, Rector of Hamstall-Ridware, and of Yoxall. 7th Edition. 2 vols. 12mo. 10s. boards.

COOPER (REV. E.)—PRACTICAL AND FAMILIAR SERMONS,

Designed for Parochial and Domestic Instruction. By the Rev. Edward Cooper, Rector of Hamstall-Ridware, and of Yoxall, in the county of Stafford; and late Fellow of All-Souls College, Oxford. New Editions. 7 vols. 12mo. 1l. 18s. boards.

** Vols. 1 to 4, 5s. each; Vols. 5 to 7, 6s. each.

COPLAND.—A DICTIONARY OF PRACTICAL MEDICINE;

Comprising General Pathology, the Nature and Treatment of Diseases, Morbid Structures, and the Disorders especially incidental to Climates, to Sex, and to the different Epochs of Life, with numerous approved Formulæ of the Medicines recommended. By James Copland, M.D., etc. etc. In 3 vols. Vols. 1 and 2, 8vo. 3l. cloth; and Part 10, 4s. 6d. sewed.

COSTELLO (MISS).—THE ROSE GARDEN OF PERSIA.A Series of Translations from the Persian Poets. By Louisa Stuart Costello, author of "Specimens of the Early Poetry of France," etc. Long 8vo. with 12 Illuminated Titles, and Borders printed in Gold and Colours, 18s. boards; or 3ls. 6d. bound in morocco (*oriental style*) by Hayday.

COSTELLO (MISS)—FALLS, LAKES, AND MOUNTAINS OF NORTH WALES; being a Pictorial Tour through the most interesting parts of the Country. By Louisa Stuart Costello, author of "The Rose Garden of Persia," "Bearn and the Pyrenees," etc. Profusely illustrated with Views, from Original Sketches by D. H. M'Kewan, engraved on wood, and lithographed, by T. and E. Gilks. Square 8vo. with Map, 14s. cloth.

CRESY (E.)—AN ENCYCLOPÆDIA OF CIVIL ENGINEERING, HISTORICAL, THEORETICAL, and PRACTICAL. By Edward Cresy, F.S.A. C.E. Illustrated by many hundred Engravings on Wood, explanatory of the Principles, Machinery, and Constructions which come under the Direction of the Civil Engineer. In 1 large volume, 8vo. uniform with Messrs. Longman and Co.'s Series of One-Volume Encyclopædias and Dictionaries. *[In the press.]*

CROCKER'S ELEMENTS OF LAND SURVEYING.

Fifth Edition, corrected throughout, and considerably improved and modernised, by T. G. Bunt, Land Surveyor, Bristol. To which are added, TABLES OF SIX-FIGURE LOGARITHMS, etc., superintended by Richard Farley, of the Nautical Almanac Establishment. Post 8vo. 12s. cloth.

CROWE.—THE HISTORY OF FRANCE,

From the Earliest Period to the Abdication of Napoleon. By E. E. Crowe, Esq. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

DAHLMANN.—HISTORY OF THE ENGLISH REVOLUTION.

By F. C. Dahlmann, late Professor of History at the University of Göttingen. Translated from the German, by H. Evans Lloyd, Esq. 8vo. 10s. 6d. cloth.

DALE (THE REV. THOMAS).—THE DOMESTIC LITURGY AND

FAMILY CHAPLAIN, in Two Parts: the First Part being Church Services adapted for Domestic Use, with Prayers for every Day of the Week, selected exclusively from the Book of Common Prayer. Part II. comprising an appropriate Sermon for every Sunday in the Year. By the Rev. Thomas Dale, M.A. Canon Residentiary of St. Paul's, and Vicar of St. Bride's, London. Post 4to. handsomely printed, 21s. cloth: or, bound by Hayday, 31s. 6d. calf lettered; 50s. morocco, with goffered edges.

DANTE, TRANSLATED BY WRIGHT.—DANTE.

Translated by Ichabod Charles Wright, M.A. late Fellow of Magdalen College, Oxford. A New Edition, revised and corrected. 3 vols. foolscap 8vo. with Portrait, 7s. 6d. sewed.

* * * *Vol. I. contains the Inferno; Vol. II. the Purgatorio; Vol. III. the Paradiso.*

DAVY (SIR HUMPHRY).—ELEMENTS OF AGRICULTURAL CHEMISTRY

in a Course of Lectures. By Sir Humphry Davy. With Notes by Dr. John Davy. 6th Edition. 8vo. with 10 Plates, 15s. cloth.

DE BURTIN.—A TREATISE ON THE KNOWLEDGE NECESSARY TO

AMATEURS OF PICTURES. Translated and abridged from the French of M. Francis Xavier De Burtin, First Stipendiary Member of the Royal Academy of Brussels in the Class of Sciences, etc. By Robert White, Esq. 8vo. with Illustrations, 12s. cloth.

DE CUSTINE.—RUSSIA.

By the Marquis De Custine. Translated from the French. 2d Edition. 3 vols. post 8vo. 31s. 6d. cloth.

DE LA BECHE.—REPORT ON THE GEOLOGY OF CORNWALL, DEVON,

AND WESTSOMERSET. By Henry T. De la Beche, F.R.S. etc., Director of the Ordnance Geological Survey. Published by Order of the Lords Commissioners of H. M. Treasury. 8vo. with Maps, Woodcuts, and 12 large Plates, 14s. cloth.

DE MORGAN.—AN ESSAY ON PROBABILITIES,

And on their Application to Life Contingencies and Insurance Offices. By Aug. De Morgan, Esq., of Trinity College, Cambridge. Foolscap 8vo. with Vignette Title, 6s. cloth.

DE SISMONDI.—THE HISTORY OF THE ITALIAN REPUBLICS:

Or, of the Origin, Progress, and Fall of Freedom in Italy, from A.D. 476 to 1806. By J. C. L. Sismondi. Fcp. 8vo. with Vignette Title, 6s. cloth.

DE SISMONDI.—THE HISTORY OF THE FALL OF THE ROMAN EM-

PIRE. Comprising a View of the Invasion and Settlement of the Barbarians. By J. C. L. De Sismondi. 2 vols. Fcp. 8vo. with Vignette Titles, 12s. cloth.

DE STRZELECKI (P. E.)—PHYSICAL DESCRIPTION OF NEW SOUTH

WALES AND VAN DIEMAN'S LAND. Accompanied by a Geological Map, Sections, and Diagrams, and Figures of the Organic Remains. By P. E. De Strzelecki. 8vo. with coloured Map and numerous Plates, 24s. cloth.

DOCTOR (THE), ETC.

5 vols. post 8vo. 2l. 12s. 6d. cloth.

"Southey acknowledged the authorship of 'The Doctor' shortly before his last illness to his most confidential friend, an M.P. of high character. And in a private letter from Mrs. Southey, dated February 27, 1843, she not only states the fact, but adds that the greater part of a sixth volume had gone through the press, and that Southey looked forward to the pleasure of drawing her into it as a contributor; giving her full authority to affirm that her husband is the author."—Robert Bell, Esq., in The Story Teller.

DODDRIDGE.—THE FAMILY EXPOSITOR;

Or, a Paraphrase and Version of the New Testament: with Critical Notes, and a Practical Improvement of each Section. By P. Doddridge, D.D. To which is prefixed, a Life of the Author, by A. Kippis, D.D. F.R.S. and S.A. New Edition. 4 vols. 8vo. 1l. 16s. cloth.

DONOVAN.—TREATISE ON CHEMISTRY.

By Michael Donovan, Esq. M.R.I.A. 4th Edition. Fcap. 8vo. with Vignette Title, 6s. cloth.

DONOVAN.—A TREATISE ON DOMESTIC ECONOMY.

By M. Donovan, Esq. M.R.I.A., Professor of Chemistry to the Company of Apothecaries in Ireland. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

DOUBLEDAY'S BUTTERFLIES.—THE GENERA OF DIURNAL LEPI-

DOPTERA; comprising their Generic Characters—a Notice of the Habits and Transformations—and a Catalogue of the Species of each Genus. By Edward Doubleday, Esq. F.L.S. etc., Assistant in the Zoological Department of the British Museum. Imperial 4to. uniform with Gray and Mitchell's Ornithology; illustrated with 75 Coloured Plates.

* * * To be published in Monthly Parts, 5s. each; each Part to consist of two coloured Plates, with accompanying Letter-press. Publication will commence when 150 Subscribers' Names have been received.

DOVER.—LIFE OF FREDERICK II. KING OF PRUSSIA.

By Lord Dover. 2d Edition. 2 vols. 8vo. with Portrait, 28s. boards.

DRUMMOND (DR. J. L.)—LETTERS TO A YOUNG NATURALIST, ON

THE STUDY OF NATURE AND NATURAL THEOLOGY. By James L. Drummond, M.D. Author of "First Steps to Botany," etc. Second Edition. Post 8vo. with Wood Engravings, 7s. 6d. boards.

DRUMMOND.—FIRST STEPS TO BOTANY,

Intended as popular Illustrations of the Science, leading to its study as a branch of general education. By J. L. Drummond, M.D. 4th Edit. 12mo. with numerous Woodcuts, 9s. boards.

DUNHAM.—THE HISTORY OF THE GERMANIC EMPIRE.

By Dr. Dunham. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

By the same Author.

**THE HISTORY OF EUROPE DURING
THE MIDDLE AGES. 4 vols. 1l. 4s.****THE HISTORY OF SPAIN AND PORTU-
GAL. 5 vols. foolscap 8vo. 1l. 10s.****THE HISTORY OF SWEDEN, DENMARK,
AND NORWAY. 3 vols. foolscap 8vo. 18s.****THE HISTORY OF POLAND. 6s.****THE LIVES OF THE EARLY WRITERS
OF GREAT BRITAIN. Foolscap 8vo. 6s.****THE LIVES OF BRITISH DRAMATISTS.
2 vols. foolscap 8vo. 12s.****DUNLOP (JOHN).—THE HISTORY OF FICTION:**

Being a Critical Account of the most celebrated Prose Works of Fiction, from the earliest Greek Romances to the Novels of the Present Age. By John Dunlop. 3d Edition, complete in One Volume. Medium 8vo. 15s. cloth.

EASTLAKE (C. L.)—MATERIALS FOR A HISTORY OF OIL PAINTING.

By Charles Lock Eastlake, R.A. 8vo.

[In the press.]

ECCLESTON (JAMES).—A MANUAL OF ENGLISH ANTIQUITIES.

By James Eccleston, B.A. Head Master of Sutton Coldfield Grammar School. 8vo. with numerous Illustrations on Wood.

[In the press.]

ELLIOT (J.).—A COMPLETE TREATISE ON PRACTICAL GEOMETRY AND

MENSURATION. With numerous Exercises. By James Elliot, formerly Teacher of Mathematics in the Mechanics' Institution of Liverpool. 8vo 5s. cloth.

* * * Key, containing full Demonstrations and Solutions. 8vo. 6s. cloth.

ELLIOTSON.—HUMAN PHYSIOLOGY:

With which is incorporated much of the Elementary Part of the "Institutiones Physiologicae" of J. F. Blumenbach, Professor in the University of Gottingen. By John Elliotson, M.D. Cantab. F.R.S. Fifth Edition, 8vo. with numerous Woodcuts, 2l. 2s. cloth.

THE ENGLISHMAN'S GREEK CONCORDANCE OF THE NEW TESTA-

MENT; being an attempt at a Verbal Connexion between the Greek and the English Texts; including a Concordance to the Proper Names, with Indexes, Greek-English and English-Greek. 2d Edition, carefully revised, with a new Index, Greek and English. Royal 8vo. 42s.

THE ENGLISHMAN'S HEBREW AND CHALDEE CONCORDANCE OF

THE OLD TESTAMENT; being an attempt at a Verbal Connexion between the Original and the English Translations; with Indexes, a List of the Proper Names and their occurrences, etc. etc. 2 vols. royal 8vo., 3l. 13s. 6d. cloth; large paper, 4l. 14s. 6d.

FAREY.—A TREATISE ON THE STEAM-ENGINE,

Historical, Practical, and Descriptive. By John Farey, Engineer. 4to. illustrated by numerous Woodcuts, and 25 Copper-plates, 5l. 5s. in boards.

FERGUS.—THE HISTORY OF THE UNITED STATES OF AMERICA,

From the Discovery of America to the Election of General Jackson to the Presidency. By the Rev. H. Fergus. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

- FIELD.—POSTHUMOUS EXTRACTS FROM THE VETERINARY RECORDS OF THE LATE JOHN FIELD.** Edited by his Brother, William Field, Veterinary Surgeon, London. 8vo. 8s. boards.
- FITZROY (LADY).—SCRIPTURAL CONVERSATIONS BETWEEN CHARLES AND HIS MOTHER.** By Lady Charles Fitzroy. Foolsap 8vo. 4s. 6d. cloth.
- FORSTER.—STATESMEN OF THE COMMONWEALTH OF ENGLAND.** With an Introductory Treatise on the Popular Progress in English History. By John Forster, Esq. 5 vols. foolscap 8vo. with Original Portraits of Pym, Eliot, Hampden, Cromwell, and an Historical Scene after a Picture by Cattermole, 1l. 10s. cloth.
- The Introductory Treatise, intended as an Introduction to the Study of the Great Civil War in the Seventeenth Century, separately, price 2s. 6d. sewed.
- The above 5 vols. form Mr. Forster's Portion of the Lives of Eminent British Statesmen, by Sir James Mackintosh, the Right Hon. T. P. Courtenay, and John Forster, Esq. 7 vols. foolscap 8vo. with Vignette Titles, 2l. 2s. cloth.
- FORSTER (REV. C.).—THE HISTORICAL GEOGRAPHY OF ARABIA;** Or, the Patriarchal Evidences of Revealed Religion. A Memoir, with illustrative Maps and an Appendix, containing Translations, with an Alphabet and Glossary of the Hamyaritic Inscriptions recently discovered in Hadramaut. By the Rev. Charles Forster, B.D., Rector of Stisted, Essex; author of "Mahometanism Unveiled." 2 vols. 8vo. 30s. cloth.
- FORSTER (REV. C.).—THE LIFE OF JOHN JEBB. D.D. F.R.S.** Late Bishop of Limerick. With a Selection from his Letters. By the Rev. Charles Forster, B.D., Rector of Stisted, Essex, and one of the Six Preachers in the Cathedral of Christ, Canterbury, formerly Domestic Chaplain to the Bishop. 2d Edition. 8vo. with Portrait, etc. 16s. cloth.
- FOSBROKE.—A TREATISE ON THE ARTS, MANNERS, MANUFACTURES, and INSTITUTIONS OF THE GREEKS and ROMANS.** By the Rev. T. D. Fosbroke, etc. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.
- GERTRUDE.** A Tale. By the author of "Amy Herbert." Edited by the Rev. William Sewell, B.D., of Exeter College, Oxford. Second Edition. 2 vols. foolscap 8vo. 9s. cloth.
- GILBART (J. W.).—THE HISTORY AND PRINCIPLES OF BANKING.** By James William Gilbert, General Manager of the London and Westminster Bank. Third Edition. 8vo. 9s. boards.
- GLEIG.—LIVES OF THE MOST EMINENT BRITISH MILITARY COMMANDERS.** By the Rev. G. R. Gleig. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.
- GLENDINNING.—PRACTICAL HINTS ON THE CULTURE OF THE PINEAPPLE.** By R. Glendinning, Gardener to the Right Hon. Lord Rolle, Bicton. 12mo. with Plan of Pinery, 5s. cloth.
- GOLDSMITH.—THE POETICAL WORKS OF OLIVER GOLDSMITH.** Illustrated by Wood Engravings, from the Designs of G. W. Cope. A.R.A., Thomas Creswick, A.R.A., J. C. Horsley, R. Redgrave, A.R.A., and Frederick Tayler, Members of the Etching Club. With a Biographical Memoir, and Notes on the Poems. Edited by Belton Corney, Esq. Square crown 8vo., uniform with "Thomson's Seasons," 2ls. cloth; or 36s. bound in morocco, by Hayday.
- * One Hundred Copies, 2l. 2s. each, printed on prepared paper of great beauty.*
- GOOD.—THE BOOK OF NATURE.** A Popular Illustration of the Natural Laws and Phenomena of Creation. By John Mason Good, M.D. F.R.S. etc. 3d Edition, corrected. 3 vols. foolscap 8vo. 24s. cloth.
- GRAHAM.—ENGLISH; OR, THE ART OF COMPOSITION** explained in a Series of Instructions and Examples. By G. F. Graham. 2d Edition, revised and improved. Foolscap 8vo. 7s. cloth.
- GRANT (MRS.).—LETTERS FROM THE MOUNTAINS.** Being the Correspondence with her Friends, between the years 1773 and 1803. By Mrs. Grant, of Laggan. 6th Edition. Edited, with Notes and Additions, by her Son, J. P. Grant, Esq. 2 vols. post 8vo. 2ls. cloth.
- GRANT (MRS., OF LAGGAN).—MEMOIR AND CORRESPONDENCE** of the late Mrs. Grant, of Laggan, author of "Letters from the Mountains," etc. Edited by her Son, J. P. Grant, Esq. 2d Edition. 3 vols. post 8vo. Portrait, 1l. 11s. 6d. cloth.
- GRATTAN.—THE HISTORY OF THE NETHERLANDS,** From the Invasion by the Romans to the Belgian Revolution in 1830. By T. C. Grattan, Esq. Foolscap 8vo. with Vignette Titles, 6s. cloth.
- GRAY (JOHN).—GRAY'S ELEGY,** Written in a Country Churchyard. Illuminated in the Missal style. By Owen Jones, Architect. Imp. 8vo. 3ls. 6d. elegantly bound in patent relieve leather.
- GRAY.—FIGURES OF MOLLUSCOUS ANIMALS,** Selected from various Authors. Etched for the Use of Students. By Maria Emma Gray. Vol. I. 8vo. with 78 plates of Figures, 12s. cloth.

GRAY AND MITCHELL'S ORNITHOLOGY.—THE GENERA OF BIRDS;
Comprising their Generic Characters, a Notice of the Habits of each Genus, and an extensive List of Species, referred to their several Genera. By George Robert Gray, Acad. Imp. Georg. Florent. Soc. Corresp. Senior Assistant of the Zoological Department, British Museum; and author of the "List of the Genera of Birds," etc. etc. Imperial 4to. illustrated with 350 Plates, by David William Mitchell, B.A.

. *In course of publication in Monthly Parts, 10s. 6d. each; each Part consisting of Four coloured Plates and Three plain, with Letter-press, giving the Generic Characters, short Remarks on the Habits, and a List of Species of each Genus as complete as possible. The uncoloured Plates contain the Characters of all the Genera of the various Sub-families, consisting of numerous details of Heads, Wings, and Feet, as the case may require, for pointing out their distinguishing Characters. The Work will not exceed 50 Monthly Parts. No. 25 was published on 1st of May.*

GRAY (J. E.)—THE GENERA OF MAMMALIA;

Comprising their Generic Characters—a Notice of the Habits of each Genus—and a short Character of each of the well-established Species, referred to the several Genera. By John Edward Gray, Esq., Keeper of the Zoological Collection of the British Museum. Imperial 4to. uniform with Gray and Mitchell's Ornithology; illustrated with 175 Plates.

. *To be published in Monthly Parts, 12s. each; each Part to consist of Four coloured and Three plain Plates, with accompanying Letter-press. The Work will not exceed 25 Parts. Publication will commence when 150 Subscribers' Names have been received.*

GREENER.—THE CUN;

Or, a Treatise on the various Descriptions of Small Fire Arms. By W. Greener, Inventor of an Improved Method of Firing Cannon by Percussion, etc. 8vo. with Illustrations, 15s. boards.

GREENWOOD (COL.)—THE TREE-LIFTER;

Or a New Method of Transplanting Trees. By Col. Geo. Greenwood. 8vo. with an Illustrative Plate, 7s. cloth.

GRIMBLOT (P.)—LETTERS OF WILLIAM III. AND LOUIS XIV. AND OF THEIR MINISTERS. Illustrating the Domestic and Foreign Policy of England during the period which followed the Revolution of 1688. Extracted from the Archives of France and England, and from Foreign papers. Edited by P. Grimblot. 8vo. *[In the press.]*

GRUNER (L.)—THE DECORATIONS OF THE GARDEN PAVILION IN THE GROUNDS OF BUCKINGHAM PALACE. Fifteen Plates, by L. Gruner. With Descriptions by Mrs. Jameson. Published by Command of Her Majesty. Small folio, 31s. 6d. plain; coloured, 5l. 5s. cloth.

GUEST.—THE MABINOGION,

From the Llyfr Coch o Hergest, or Red Book of Hergest, and other ancient Welsh MSS. with an English Translation and Notes. By Lady Charlotte Guest. Parts 1 to 6. Royal 8vo. 8s. each sewed.

GUICCIARDINI (F.)—THE MAXIMS OF FRANCIS GUICCIARDINI, THE HISTORIAN. Translated by Emma Martin. With Notes, and Parallel Passages from the Works of Machiavelli, Lord Bacon, Pascal, Rochefoucault, Montesquieu, Burke, Prince Talleyrand, Guizot, and others; and a Sketch of the Life of Guicciardini. Square foolscap 8vo. with Portrait, 7s. ornamental boards; or 14s. bound in morocco (*old style*) by Hayday.

GWILT.—AN ENCYCLOPÆDIA OF ARCHITECTURE;

Historical, Theoretical, and Practical. By Joseph Gwilt, Esq., F.S.A. Illustrated with upwards of 1000 Engravings on Wood, from Designs by J. S. Gwilt. 8vo. 2l. 12s. 6d. cloth.

HALL.—NEW GENERAL LARGE LIBRARY ATLAS OF FIFTY-THREE

MAPS, on Colombier Paper; with the Divisions and Boundaries carefully coloured. Constructed entirely from New Drawings, and engraved by Sidney Hall. New Edition, thoroughly revised and corrected; including all the Alterations rendered necessary by the recent Official Surveys, the New Roads on the Continent, and a careful Comparison with the authenticated Discoveries published in the latest Voyages and Travels. Folded in half, Nine Guineas, half-bound in russia; full size of the Maps, Ten Pounds, half-bound in russia.

HALSTED.—LIFE AND TIMES OF RICHARD THE THIRD,

as Duke of Gloucester and King of England: in which all the Charges against him are carefully investigated and compared with the Statements of the Cotemporary Authorities. By Caroline A. Halsted, author of "The Life of Margaret Beaufort." 2 vols. 8vo. with Portrait from an Original Picture in the Possession of the Right Honorable Lord Stafford, never before engraved, and other Illustrations, 1l. 10s. cloth.

HAND-BOOK OF TASTE (THE);

Or, How to Observe Works of Art, especially Cartoons, Pictures, and Statues. By Fabius Pictor. 3d Edition. Foolscap 8vo. 3s. boards.

HANSARD.—TROUT AND SALMON FISHING IN WALES.

By G. A. Hansard, 12mo. 6s. 6d. cloth.

HARRIS.—THE HIGHLANDS OF ÆTHIOPIA;

Being the Account of Eighteen Months' Residence of a British Embassy to the Christian Court of Shoa. By Major Sir W. C. Harris, author of "Wild Sports in Southern Africa," etc. 2d Edition. 3 vols. 8vo. with Map and Illustrations, 2l. 2s. cloth.

HAWES (BARBARA).—TALES OF THE NORTH AMERICAN INDIANS,
and Adventures of the Early Settlers in America; from the Landing of the Pilgrim Fathers
in 1620, to the Time of the Declaration of Independence. By Barbara Hawes. Fcp. 8vo. 6s.

HAWKER.—INSTRUCTIONS TO YOUNG SPORTSMEN

In all that relates to Guns and Shooting. By Lieut. Col. P. Hawker. 9th edition, corrected,
enlarged, and improved, with Eighty-five Plates and Woodcuts, by Adlard and Branstons,
from Drawings by C. Varley, Dicks, etc. 8vo. 21s. cloth.

HAYDON (B. R.).—LECTURES ON PAINTING AND DESIGN,

Delivered at the London Institution, the Royal Institution, Albermarle Street, to the
University of Oxford, etc. By B. R. Haydon, Historical Painter. With Designs drawn on
Wood by the author, and Engraved by Edward Evans. 8vo. 12s. cloth.

**** The Second Volume is preparing for publication.*

**HENSLOW.—THE PRINCIPLES OF DESCRIPTIVE AND PHYSIOLO-
GICAL BOTANY.** By J. S. Henslow, M.A. F.L.S. etc. Foolsap 8vo. with Vignette Title,
and nearly 70 Woodcuts, 6s. cloth.

HERON (R. M.).—PALESTRINA :

A Metrical Romance. By Robert M. Heron, Esq. 8vo. 8s. 6d.

HERSCHEL.—A TREATISE ON ASTRONOMY.

By Sir John Herschel. New Edition. Fcap. 8vo. with Vignette Title, 6s. cloth.

**HERSCHEL.—A PRELIMINARY DISCOURSE ON THE STUDY OF
NATURAL PHILOSOPHY.** By Sir John Herschel. New Edition. Foolsap 8vo. 6s.

HINTS ON ETIQUETTE AND THE USAGES OF SOCIETY :

With a Glance at Bad Habits. By *Αγαγός*. "Manners make the Man." 24th Edition,
revised (with additions) by a Lady of Rank. Foolsap 8vo. 2s. 6d. cloth, gilt edges.

HISTORICAL PICTURES OF THE MIDDLE ACES,

In Black and White. Made on the spot, from Records in the Archives of Switzerland. By a
Wandering Artist. 2 vols post 8vo. 18s. cloth.

Contents.—The Nuns' War; the War of Two Abbots (Reichenau and St. Gall); the Passage
of the Great St. Bernard; and Bertha, Queen of Transjurane Burgundy.

**HOARE.—A DESCRIPTIVE ACCOUNT OF A NEW METHOD OF
PLANTING AND MANAGING THE ROOTS OF GRAPE VINES.** By Clement Hoare,
author of "A Treatise on the Cultivation of the Grape Vine on Open Walls." 12mo. 5s. cl.

**HOARE.—A PRACTICAL TREATISE ON THE CULTIVATION OF THE
GRAPE VINE ON OPEN WALLS.** By Clement Hoare. 3d Edition, 8vo. 7s. 6d. cloth.

HOBBS.—THE COMPLETE WORKS OF THOMAS HOBBS,

Of Malmesbury; now first collected and edited by Sir William Molesworth, Bart. 16 vols.
8vo. 8s. cloth.

**** Separately, the English Works, in 11 vols. 5l. 10s.; the Latin Works, in 5 vols. 2l. 10s.*

HOLLAND.—A TREATISE ON THE MANUFACTURES IN METAL.

By John Holland, Esq. 3 vols. foolsap 8vo. with about 300 Woodcuts, 18s. cloth.

HOLLAND.—MEDICAL NOTES AND REFLECTIONS.

By Henry Holland, M.D. F.R.S. etc. Fellow of the Royal College of Physicians, Physician
Extraordinary to the Queen, and Physician in Ordinary to His Royal Highness Prince Albert.
2d Edition. 8vo. 18s. cloth.

HOOK (DR. W. F.).—THE LAST DAYS OF OUR LORD'S MINISTRY;

A Course of Lectures on the principal Events of Passion Week. By Walter Farquhar Hood,
D.D., Vicar of Leeds, Prebendary of Lincoln, and Chaplain in Ordinary to the Queen. 4th
Edition. Foolsap 8vo. 6s. cloth.

HOOKE. —THE BRITISH FLORA.

In 2 vols. Vol. I.; comprising the Phænogamous or Flowering Plants, and the Ferns. By Sir
William Jackson Hooker, K.H. LL.D. F.R.A. and L.S. etc. etc. etc. 5th Edition, with
Additions and Corrections; and 173 Figures, illustrative of the Umbelliferous Plants, the
Composite Plants, the Grasses, and the Ferns. Vol. I. 8vo., with 12 Plates, 14s. plain; with
the plates coloured, 24s. cloth.

Vol. II, in Two Parts, comprising the Cryptogamia and the Fungi, completing the British
Flora, and forming Vol. V., Parts 1 and 2, of Smith's English Flora, 24s. boards.

HOOKE AND TAYLOR.—MUSCOLOGIA BRITANNICA.

Containing the Mosses of Great Britain and Ireland, systematically arranged and described;
with Plates, illustrative of the character of the Genera and Species. By Sir W. J. Hooker
and T. Taylor, M.D. F.L.S. etc. 2d Edition, enlarged, 8vo. 31s. 6d. plain; 3l. 3s. coloured.

HORNE (THE REV. T. H.).—AN INTRODUCTION TO THE CRITICAL STUDY AND KNOWLEDGE OF THE HOLY SCRIPTURES. By the Rev. Thomas Hartwell Horne, B.D. of St. John's College, Cambridge; Rector of the united Parishes of St. Edmund the King and Martyr, and St. Nicholas Acons, Lombard Street; Prebendary of St. Paul's. 9th Edition, revised and corrected. 5 vols. 8vo. with Maps and Fac-similes, 63s. cloth; or 5*l.* bound in calf half-extra, by Hayday.

SUPPLEMENTARY PAGES, for purchasers of the 7th and 8th Editions, 1*s.* 6*d.*

HORNE (THE REV. T. H.).—A COMPENDIOUS INTRODUCTION TO THE STUDY OF THE BIBLE. By the Rev. Thomas Hartwell Horne, B.D. of St. John's College, Cambridge. Being an Analysis of his "Introduction to the Critical Study and Knowledge of the Holy Scriptures." 7th Edition, corrected and enlarged. 12mo. with Maps and other Engravings, 9*s.* boards.

HORSLEY (BISHOP).—BIBLICAL CRITICISM ON THE FIRST FOURTEEN HISTORICAL BOOKS OF THE OLD TESTAMENT; AND ON THE FIRST NINE PROPHETICAL BOOKS. By Samuel Horsley, LL.D. F.R.S. F.A.S. Lord Bishop of St. Asaph. Second Edition, containing Translations by the Author, never before published, together with copious Indexes. 2 vols. 8vo. 30*s.* cloth.

By the same Author,

THE BOOK OF PSALMS; translated from the Hebrew: with Notes, explanatory and critical. 4th Edition. 8vo. 12*s.* cloth.

HOWITT.—THE RURAL LIFE OF ENGLAND.

By William Howitt. Third Edition, corrected and revised. Medium 8vo. with Engravings on Wood by Bewick and Williams, uniform with "Visits to Remarkable Places," 21*s.* cloth.

HOWITT.—VISITS TO REMARKABLE PLACES;

Old Halls, Battle-Fields, and Scenes illustrative of Striking Passages in English History and Poetry. By William Howitt. New Edition. Medium 8vo. with 40 Illustrations, 21*s.* cloth.

SECOND SERIES, chiefly in the Counties of DURHAM and NORTHUMBERLAND, with a Stroll along the BORDEEN. Medium 8vo. with upwards of 40 highly-finished Woodcuts, from Drawings made on the spot, 21*s.* cloth.

HOWITT.—THE RURAL AND DOMESTIC LIFE OF GERMANY:

With Characteristic Sketches of its chief Cities and Scenery. Collected in a General Tour, and during a Residence in that Country in the Years 1840-42. By William Howitt, author of "The Rural Life of England," etc. Medium 8vo., with above 50 Illustrations, 21*s.* cloth.

HOWITT.—THE STUDENT-LIFE OF GERMANY.

From the Unpublished MS. of Dr. Cornelius. By William Howitt. 8vo. with 24 Wood-Engravings, and 7 Steel Plates, 21*s.* cloth.

HOWITT.—COLONISATION AND CHRISTIANITY:

A Popular History of the Treatment of the Natives, in all their Colonies, by the Europeans. By William Howitt. Post 8vo. 10*s.* 6*d.* cloth.

HOWITT.—THE BOY'S COUNTRY BOOK:

Being the real Life of a Country Boy, written by Himself; exhibiting all the Amusements, Pleasures, and Pursuits of Children in the Country. Edited by William Howitt, author of "The Rural Life of England," etc. 2d Edition. Fcap. 8vo. with 40 Woodcuts, 8*s.* cloth.

HOWITT (RICHARD).—IMPRESSIONS OF AUSTRALIA FELIX,

During a Four Years' Residence in that Colony: with particular reference to the Prospects of Emigrants. With Notes of a Voyage round the World, Australian Poems, etc. By Richard Howitt. Foolscap 8vo. 7*s.* cloth.

HUDSON.—THE PARENT'S HAND-BOOK;

Or, Guide to the Choice of Professions, Employments, and Situations; containing useful and practical information on the subject of placing out Young Men, and of obtaining their Education with a view to particular occupations. By J. C. Hudson, Esq., author of "Plain Directions for Making Wills." Fcap. 8vo. 5*s.* cloth.

HUDSON.—PLAIN DIRECTIONS FOR MAKING WILLS

In conformity with the Law, and particularly with reference to the Act 7 Wm. IV. and 1 Vict. c. 26. To which is added, a clear Exposition of the Law relating to the Distribution of Personal Estate in the case of Intestacy, with two Forms of Wills, and much useful Information, etc. By J. C. Hudson, Esq. 13th Edition, corrected, with Notes of Cases judicially decided since the above Act came into operation. Fcap. 8vo. 2*s.* 6*d.*

HUDSON.—THE EXECUTOR'S GUIDE.

By J. C. Hudson, Esq., of the Legacy Duty Office, London; author of "Plain Directions for Making Wills," and "The Parent's Hand-Book." 4th Edition. Foolscap 8vo. 5*s.* cloth.

*** The above two works may be had in One volume, price 7*s.* cloth.*

HUMBOLDT (BARON).—KOSMOS:

A Physical Description of the Universe. Translated, with the Author's Sanction and Co-operation, under the superintendence of Lieutenant-Colonel Edward Sabine, F.R.S. of the Royal Artillery. Vols. 1 and 2, post 8vo. [In the press.]

*** This translation has been made at the request of Baron Von Humboldt; who will see the sheets before publication.*

HUMPHREYS.—THE ILLUMINATED BOOKS OF THE MIDDLE AGES.

A History of Illuminated Books, from the 14th to the 17th Century. By Henry Noel Humphreys. Illustrated by a Series of Fac-simile Specimens, each consisting of an entire Page, of the exact size of the Original, from the most celebrated and splendid MSS. in the Imperial and Royal Libraries of Vienna, Moscow, Paris, Naples, Copenhagen, and Madrid;—from the Vatican, Escorial, Ambrosian, and other great Libraries of the Continent;—and from the rich Public Collegiate, and Private Libraries of Great Britain.

In course of publication, in Parts. Parts 1 and 2, imperial 4to. each containing Three Plates, splendidly printed, in gold, silver, and colours, in imitation of the originals, as accurate as can be produced by mechanical means, with Descriptions, price 12s.

Large Paper, on Half Imperial (21½ in. by 15), to prevent folding the large Plates, 21s.

* * Six Parts to form a Volume, Four Volumes completing the work.

HUNT.—RESEARCHES ON LIGHT:

An Examination of all the Phenomena connected with the Chemical and Molecular Changes produced by the Influence of the Solar Rays; embracing all the known Photographic Processes, and new Discoveries in the Art. By Robert Hunt, Keeper of Mining Records, Museum of Economic Geology. 8vo. with Plate and Woodcuts, 10s. 6d. cloth.

ILLUMINATED CALENDAR (THE).—THE ILLUMINATED CALENDAR and HOME

DIARY for 1846; containing 12 pages of *fac-simile* from the Calendar of the rich MSS. "Hours" of the Duke of Anjou, styled King of Sicily and Jerusalem; also 24 pages of Diary, each illuminated with an elaborate Border taken from the same MS.; and an Illuminated Title. Imperial 8vo. 42s. in a binding composed from the miniature pictures of the same MS.

* * *The Illuminated Calendar and Home Diary, for 1845; copied from the Manuscript of the "Hours of Anne of Brittany."* Imp. 8vo. 42s. in emblazoned printing and binding.

JACKSON.—THE PICTORIAL FLORA;

Or, British Botany Delineated, in 1500 Lithographic Drawings of all the Species of Flowering Plants indigenous to Great Britain; illustrating the descriptive works on English Botany of Hooker, Lindley, Smith, etc. By Miss Jackson. 8vo. 15s. cloth.

JAMES.—A HISTORY OF THE LIFE OF EDWARD THE BLACK PRINCE,

and of various Events connected therewith, which occurred during the Reign of Edward III. King of England. By G. P. R. James, Esq. 2d Edition. 2 vols. foolscap 8vo. with Map, 15s.

JAMES.—LIVES OF THE MOST EMINENT FOREIGN STATESMEN.

By G. P. R. James, Esq., and E. E. Crowe, Esq. 5 vols. foolscap 8vo. 30s. cloth.

JEBB (BISHOP).—PASTORAL INSTRUCTIONS ON THE CHARACTER

AND PRINCIPLES OF THE CHURCH OF ENGLAND, selected from his former Publications. By John Jebb, D.D. F.R.S., late Bishop of Limerick, Ardfer, and Aghadoc. A New Edition. Foolscap 8vo. 6s. cloth.

By the same Author,

PIETY WITHOUT ASCETICISM; or, the Protestant Kempis: a Manual of Christian Faith and Practice, selected from the Writings of Scougal, Charles Howe, and Cudworth; with corrections and occasional Notes. Second Edition. Foolscap 8vo. 6s. cloth.

JEBB (BISHOP) AND KNOX (ALEXANDER).—THIRTY YEARS' COR-

RESPONDENCE between John Jebb, D.D. F.R.S., Bishop of Limerick, Ardfer, Aghadoc, and Alexander Knox, Esq. M.R.I.A. Edited by the Rev. Charles Forster, B.D. Rector of Stisted, Essex, and one of the Six Preachers in the Cathedral of Christ, Canterbury, formerly Domestic Chaplain to Bishop Jebb. 2d Edition. 2 vols. 8vo. 28s. cloth.

LORD JEFFREY.—CONTRIBUTIONS TO THE EDINBURGH REVIEW.

By Francis Jeffrey, now one of the Judges in the Court of Session in Scotland. 4 vols. 8vo. 48s. cloth.

JOHNSON.—THE FARMER'S ENCYCLOPÆDIA,

AND DICTIONARY OF RURAL AFFAIRS; embracing all the recent Discoveries in Agricultural Chemistry; adapted to the comprehension of unscientific Readers. By Cuthbert W. Johnson, Esq., F.R.S. Barrister-at-Law, Corresponding Member of the Agricultural Society of Königsberg, and of the Maryland Horticultural Society, author of several of the Prize Essays of the Royal Agricultural Society of England, and other Agricultural Works; Editor of the "Farmer's Almanack," etc. 8vo. with Wood Engravings of the best and most improved Agricultural Implements, 2l. 10s. cloth.

KANE.—ELEMENTS OF CHEMISTRY;

Including the most recent Discoveries and Applications of the Science to Medicine and Pharmacy, and to the Arts. By Robert Kane, M.D. M.R.I.A. Professor of Natural Philosophy to the Royal Dublin Society. 8vo. with 236 Woodcuts, 24s. cloth.

KATER AND LARDNER.—A TREATISE ON MECHANICS.

By Captain Kater and Dr. Lardner. New Edition. Foolscap 8vo. with Vignette Title, and 19 Plates, comprising 224 distinct figures, 6s. cloth.

KEIGHTLEY.—OUTLINES OF HISTORY,

From the Earliest Period. By Thomas Keightley, Esq. New Edition, corrected and considerably improved. Foolscap 8vo., 6s. cloth; or 6s. 6d. bound.

KIRBY AND SPENCE.—AN INTRODUCTION TO ENTOMOLOGY;
Or, Elements of the Natural History of Insects: comprising an account of noxious and useful Insects, of their Metamorphoses, Food, Stratagems, Habitations, Societies, Motions, Noises, Hybernation, Instinct, etc. By W. Kirby, M.A. F.R.S. & L.S. Rector of Barham; and W. Spence, Esq., F.R.S. & L.S. 6th Edition, corrected and considerably enlarged. 2 vols. 8vo. 1l. 11s. 6d. cloth.

* * * *The first two volumes of the "Introduction to Entomology" are published as a separate work, distinct from the third and fourth volumes, and, though much enlarged, at a considerable reduction of price, in order that the numerous class of readers who confine their study of insects to that of their manners and economy, need not be burthened with the cost of the technical portion of the work, relating to their anatomy, physiology, etc.*

KNOX (ALEXANDER).—REMAINS OF ALEXANDER KNOX, ESQ.
Of Dublin, M.R.I.A.; containing Essays, chiefly explanatory, of Christian Doctrine; and Confidential Letters, with Private Papers, illustrative of the Writer's Character, Sentiments, and Life. 3d Edition. 4 vols. 8vo. 2l. 8s. cloth.

LAING.—NOTES ON THE SCHISM FROM THE CHURCH OF ROME,
called the GERMAN CATHOLIC CHURCH, instituted by J. Ronge and I. Czerzki, in October 1844, on occasion of the Pilgrimage to the Holy Coat at Treves. By S. Laing, Esq., author of "Notes of a Traveller," etc. 2d Edition. Foolscep 8vo. 5s. cloth.

LAING.—THE CHRONICLE OF THE KINGS OF NORWAY,
From the Earliest Period of the History of the Northern Sea Kings to the Middle of the Twelfth Century, commonly called the Heimskringla. Translated from the Icelandic of Snorro Sturleson, with Notes, and a Preliminary Discourse, by Samuel Laing, author of "Notes of a Traveller," etc. 3 vols. 8vo. 36s. cloth.

LAING.—A TOUR IN SWEDEN

In 1838; comprising Observations on the Moral, Political, and Economical State of the Swedish Nation. By Samuel Laing, Esq. 8vo. 12s. cloth.

LAING.—NOTES OF A TRAVELLER

On the Social and Political State of France, Prussia, Switzerland, Italy, and other parts of Europe, during the present Century. By Samuel Laing, Esq. 2d Edition. 8vo. 16s. cloth.

LAING.—JOURNAL OF A RESIDENCE IN NORWAY

During the years 1834, 1835, and 1836; made with a view to inquire into the Rural and Political Economy of that Country, and the Condition of its Inhabitants. By Samuel Laing, Esq. 2d Edition. 8vo. 14s. cloth.

LANE (R. J.).—LIFE AT THE WATER CURE:

Or, a Month at Malvern. A Diary of Facts and Fancies. To which is added the Sequel. By Richard J. Lane, A.R.A., Lithographer in Ordinary to Her Majesty and His Royal Highness Prince Albert. 8vo. with many Illustrations. *[In the press.]*

LANETON PARSONAGE:

A Tale for Children, on the practical use of the Church Catechism. By the Author of "Amy Herbert," and "Gertrude," Fcp. 8vo. *[In the press.]*

LAPLACE (THE MARQUIS DE).—THE SYSTEM OF THE WORLD.

By M. Le Marquis De Laplace. Translated from the French, and elucidated with Explanatory Notes. By the Rev. Henry H. Harte, F.T.C.D. M.R.I.A. 2 vols. 8vo. 24s. boards.

LARDNER'S CABINET CYCLOPÆDIA;

Comprising a Series of Original Works on History, Biography, Literature, the Sciences, Arts, and Manufactures. Conducted and edited by Dr. Lardner.

The Series complete in One Hundred and Thirty-three Volumes, 39l. 18s. The Works separately, 6s. per volume.

LARDNER.—A TREATISE ON ARITHMETIC.

By Dr. Lardner, LL.D. F.R.S. Foolscep 8vo. with Vignette Title, 6s. cloth.

LARDNER.—A TREATISE ON GEOMETRY,

And its Application to the Arts. By Dr. Lardner. Foolscep 8vo., 6s. cloth.

LARDNER.—A TREATISE ON HEAT.

By Dr. Lardner, LL.D. etc. Fcp. 8vo. with Vignette Title and Woodcuts, 6s. cloth.

LARDNER.—A TREATISE ON HYDROSTATICS AND PNEUMATICS.

By Dr. Lardner. New Edition. Foolscep 8vo. with Vignette Title, 6s. cloth.

LARDNER AND WALKER.—A MANUAL ON ELECTRICITY, MAGNETISM, and METEOROLOGY.

By Dr. Lardner, LL.D. F.R.S., and C. V. Walker, Secretary of the Electrical Society. 2 vols. foolscep 8vo., with Vignette Titles, 12s. cloth.

L. E. L.—THE POETICAL WORKS OF LETITIA ELIZABETH LANDON.

New Edition, 4 vols. foolscep 8vo. with Illustrations by Howard, etc. 28s. cloth; or bound in morocco, with gilt edges, 2l. 4s.

The following Works separately:—

The IMPROVISATRICE	- - 10s. 6d.	The GOLDEN VIOLET	- - - 10s. 6d.
The VENETIAN BRACELET	10s. 6d.	The TROUBADOUR	- - - 10s. 6d.

LEE.—TAXIDERMY;

Or, the Art of Collecting, Preparing, and Mounting Objects of Natural History. For the use of Museums and Travellers. By Mrs. R. Lee (formerly Mrs. T. E. Bowdich), author of "Memoirs of Cuvier," etc. 6th Edition, improved, with an account of a Visit to Walton Hall, and Mr. Waterton's method of Preserving Animals. Fcap. 8vo. with Woodcuts, 7s.

LEE.—ELEMENTS OF NATURAL HISTORY,

For the Use of Schools and Young Persons: comprising the Principles of Classification, interspersed with amusing and instructive original Accounts of the most remarkable Animals. By Mrs. R. Lee (formerly Mrs. T. E. Bowdich), author of "Taxidermy," "Memoirs of Cuvier," etc. 12mo. with 55 Woodcuts, 7s. 6d. bound.

LEFEVRE (SIR GEORGE).—AN APOLOGY FOR THE NERVES;

Or, their Importance and Influence in Health and Disease. By Sir George Lefevre, M.D. author of "The Life of a Travelling Physician," "Thermal Comfort," etc. Post 8vo. 9s.

LEMPRIÈRE.—A CLASSICAL DICTIONARY;

Containing a copious Account of all the Proper Names mentioned in Ancient Authors; with the Value of Coins, Weights, and Measures, used amongst the Greeks and Romans; and a Chronological Table. By T. Lemprière, D.D. 20th Edition, corrected. 8vo. 9s. cloth.

LEREBOURS (N.P.).—A TREATISE ON PHOTOGRAPHY;

Containing the latest Discoveries appertaining to the Daguerreotype. Compiled from Communications by M.M. Daguerre and Arago, and other eminent Men of Science. By N. P. Lerebours, Optician to the Observatory, Paris, etc. Translated by J. Egerton. Post 8vo. with Plate, 7s. 6d. cloth.

LESLIE (C.R.).—MEMOIRS OF THE LIFE OF JOHN CONSTABLE, ESQ.

R. A. Composed chiefly of his Letters. By C. R. Leslie, R. A. Second Edition, with further Extracts from his Correspondence. Small 4to. with two Portraits (one from a new Sketch, by Mr. Leslie,) and a plate of "Spring," engraved by Lucas, 2ls. cloth.

LIFE OF A TRAVELLING PHYSICIAN,

From his first Introduction to Practice; including Twenty Years' Wanderings throughout the greater part of Europe. By the late Sir George Lefevre, M.D. 3 vols. post 8vo. with coloured Frontispieces, 3ls. 6d. cloth.

LINDLEY.—INTRODUCTION TO BOTANY.

By Prof. J. Lindley, Ph.D. F.R.S. L.S. etc. 3d Edition, with Corrections and considerable Additions. 8vo. with Six Plates and numerous Woodcuts, 18s. cloth.

LINDLEY.—FLORA MEDICA;

A Botanical Account of all the most important Plants used in Medicine in different Parts of the World. By John Lindley, Ph.D. F.R.S. etc. 8vo. 18s. cloth.

LINDLEY.—A SYNOPSIS OF THE BRITISH FLORA,

Arranged according to the Natural Orders. By Professor John Lindley, Ph. D., F.R.S., etc. Third Edition, with numerous Additions and Improvements. 12mo. 10s. 6d. cloth.

LINDLEY.—THE THEORY OF HORTICULTURE;

Or, an Attempt to Explain the Principal Operations of Gardening upon Physiological Principles. By John Lindley, Ph.D. F.R.S. 8vo. with Illustrations on Wood, 12s. cloth.

LINDLEY.—GUIDE TO THE ORCHARD AND KITCHEN GARDEN;

Or, an Account of the most valuable Fruits and Vegetables cultivated in Great Britain: with Kalendars of the Work required in the Orchard and Kitchen Garden during every month in the Year. By George Lindley, C.M.H.S. Edited by Professor Lindley. 8vo. 16s. boards.

LINWOOD (W.).—ANTHOLOGIA OXONIENSIS;

Sive, Florilegium e lusibus poeticis diversorum Oxoniensium Græcis et Latinis decerptum. Curante Gulielmo Linwood, M.A. Ædis Christi Alummo. 8vo. [In the press.]

LLOYD.—A TREATISE ON LIGHT AND VISION.

By the Rev. H. Lloyd, M.A., Fellow of Trin. Coll. Dublin. 8vo. 5s. boards.

LORIMER.—LETTERS TO A YOUNG MASTER MARINER,

On some Subjects connected with his Calling. By Charles Lorimer. 3d edition. 12mo. with an Appendix, 5s. 6d. cloth.

LOUDON (MRS.).—THE LADY'S COUNTRY COMPANION;

Or, How to Enjoy a Country Life Rationally. By Mrs. Loudon, author of "Gardening for Ladies," etc. New Edition. Foolscep 8vo., with an Engraving on Steel, and Illustrations on Wood, 7s. 6d. cloth.

LOUDON (J. C.).—SELF INSTRUCTION

For Young Gardeners, Foresters, Bailiffs, Land Stewards, and Farmers; in Arithmetic Book-keeping, Geometry, Mensuration, Practical Trigonometry, Mechanics, Land-Surveying, Leveling, Planning and Mapping, Architectural Drawing, and Isometrical Projection and Perspective; with Examples shewing their applications to Horticultural and Agricultural Purposes. By the late J. C. Loudon, F.L.S. H.S. etc. With a Portrait of Mr. Loudon, and a Memoir by Mrs. Loudon. 8vo. with Wood Engravings, 7s. 6d. cloth.

LOUDON.—AN ENCYCLOPÆDIA OF TREES AND SHRUBS ;

Being the "Arboretum et Fruticetum Britannicum" abridged: containing the Hardy Trees and Shrubs of Great Britain, Native and Foreign, scientifically and popularly described: with their Propagation, Culture, and Uses in the Arts; and with Engravings of nearly all the Species. For the use of Nurserymen, Gardeners, and Foresters. By J. C. Loudon, F.L.S. etc. 8vo. with upwards of 2000 Engravings on Wood, 2*l.* 10*s.* cloth.

The Original Work; a New Edition, in 8 vols. 8vo. with above 400 8vo. Plates of Trees, and upwards of 2500 Woodcuts, 10*l.* cloth.

LOUDON.—AN ENCYCLOPÆDIA OF GARDENING ;

Presenting in one systematic view, the History and Present State of Gardening in all Countries, and its Theory and Practice in Great Britain: with the Management of the Kitchen Garden, the Flower Garden, Laying-out Grounds, etc. By J. C. Loudon, F.L.S. etc. A new Edition, enlarged and much improved. 8vo. with nearly 1000 Engravings on Wood, 2*l.* 10*s.* cloth.

LOUDON.—AN ENCYCLOPÆDIA OF AGRICULTURE ;

Comprising the Theory and Practice of the Valuation, Transfer, Laying-out, Improvement, and Management of Landed Property, and of the cultivation and economy of the Animal and Vegetable productions of Agriculture, including all the latest improvements; a general History of Agriculture in all countries; a Statistical view of its present state, with suggestions for its future progress in the British Isles; and a Supplement, bringing down the work to the year 1844. By J. C. Loudon, F.L.G.Z. and H.S. etc. Fifth Edition. 8vo. with upwards of 1100 Engravings on Wood, by Branston, 2*l.* 10*s.* cloth.

The Supplement, with 65 Engravings, 5*s.* sewed.

LOUDON.—AN ENCYCLOPÆDIA OF PLANTS ;

Including all the Plants which are now found in, or have been introduced into, Great Britain; giving their Natural History, accompanied by such Descriptions, Engraved Figures, and Elementary Details, as may enable a beginner, who is a mere English reader, to discover the name of every Plant which he may find in flower, and acquire all the information respecting it which is useful and interesting. The Specific Characters by an Eminent Botanist; the Drawings by J. D. C. Sowerby, F.L.S. A new Edition (1841), with a new Supplement, comprising every desirable particular respecting all the Plants originated in, or introduced into, Britain between the first publication of the work, in 1829, and January 1840: with a new General Index to the whole work. Edited by J. C. Loudon, prepared by W. H. Baxter, Jun., and revised by George Don, F.L.S.; and 800 new Figures of Plants, on Wood, from Drawings by J. D. C. Sowerby, F.L.S. 8vo. with nearly 10,000 Wood Engravings, 7*3s.* 6*d.* cloth.

. The last Supplement, *separately*, 8vo. 15*s.* cloth.

LOUDON.—AN ENCYCLOPÆDIA OF COTTAGE, FARM, AND VILLA ARCHITECTURE AND FURNITURE.

Containing Designs for Cottages, Villas, Farm Houses, Farmeries, Country Inns, Public Houses, Parochial Schools, etc.; with the requisite Fittings-up, Fixtures, and Furniture, and appropriate Offices, Gardens, and Garden Scenery: each Design accompanied by Analytical and Critical Remarks illustrative of the Principles of Architectural Science and Taste on which it is composed, and General Estimates of the Expense. By J. C. Loudon, F.L.S. etc. New Edition, corrected, with a Supplement containing 160 additional pages of letter-press, and nearly 300 new engravings. 8vo. with more than 2000 Engravings on Wood, 6*3s.* cloth.

. The Supplement, *separately*, 8vo. 7*s.* 6*d.* sewed.

LOUDON.—HORTUS BRITANNICUS :

A Catalogue of all the Plants indigenous to or introduced into Britain. The 3d Edition, with a New Supplement, prepared, under the direction of J. C. Loudon, by W. H. Baxter, and revised by George Don, F.L.S. 8vo. 3*l.* 6*d.* cloth.

The Supplement *separately*, 8vo. 2*s.* 6*d.* sewed.

The later Supplement *separately*, 8vo. 8*s.* sewed.

LOUDON.—THE SUBURBAN GARDENER AND VILLA COMPANION :

Comprising the Choice of a Villa or Suburban Residence, or of a situation on which to form one; the Arrangement and Furnishing of the House; and the Laying-out, Planting, and general Management of the Garden and Grounds; the whole adapted for Grounds from one perch to fifty acres and upwards in extent; intended for the instruction of those who know little of Gardening or Rural Affairs, and more particularly for the use of Ladies. By J. C. Loudon, F.L.S., etc. 8vo. with above 300 Wood Engravings, 20*s.* cloth.

LOUDON.—HORTUS LIGNOSUS LONDINENSIS ;

Or, a Catalogue of all the Ligneous Plants cultivated in the neighbourhood of London. To which are added their usual Prices in Nurseries. By J. C. Loudon, F.L.S. etc. 8vo. 7*s.* 6*d.*

LOW.—ON LANDED PROPERTY, AND THE MANAGEMENT OF ESTATES ;

Comprehending the Relations between Landlord and Tenant, and the Principles and Forms of Leases; of Farm-buildings, Enclosures, Drains, Embankments, Roads, and other Rural Works, Minerals, and Woods. By David Low, Esq. F.R.S.E. etc., author of "Elements of Practical Agriculture," etc. 8vo. with numerous Wood Engravings, 2*l.* cloth.

LOW.—ON THE DOMESTICATED ANIMALS OF GREAT BRITAIN,

comprehending the Natural and Economical History of the Species and Breeds; Illustrations of the Properties of External Form; and Observations on the Principles and Practice of Breeding. By David Low, Esq., F.R.S.E. Professor of Agriculture in the University of Edinburgh, etc.; author of "Elements of Practical Agriculture," etc. 8vo. with Engravings on Wood, 25*s.* cloth.

LOW.—THE BREEDS OF THE DOMESTICATED ANIMALS OF GREAT BRITAIN described. By David Low, Esq. F.R.S.E., Professor of Agriculture in the University of Edinburgh, etc. etc. The Plates from drawings by W. Nicholson, R.S.A., reduced from a Series of Oil Paintings, executed for the Agricultural Museum of the University of Edinburgh, by W. Shiels, R.S.A. 2 vols. atlas quarto, with 56 plates of Animals, beautifully coloured after Nature, 16l. 16s. half-bound in morocco.

Or in four separate portions, as follow:—

The OX. 1 Vol. atlas quarto, with 22 Plates, price 6l. 16s. 6d. half-bound morocco.

The HORSE. 1 Vol. atlas quarto, with 8 Plates, price 3l. half-bound morocco.

The SHEEP. 1 Vol. atlas quarto, with 21 Plates, price 6l. 16s. 6d. half-bound morocco.

The HOG. 1 Vol. atlas quarto, with 5 Plates, price 2l. 2s. half-bound morocco.

LOW.—ELEMENTS OF PRACTICAL AGRICULTURE;

Comprehending the Cultivation of Plants, the Husbandry of the Domestic Animals, and the Economy of the Farm. By David Low, Esq. F.R.S.E., Professor of Agriculture in the University of Edinburgh. 4th Edition, with Alterations and Additions. 8vo. with above 200 Woodcuts, 2ls. cloth.

MACAULAY.—CRITICAL AND HISTORICAL ESSAYS CONTRIBUTED TO THE EDINBURGH REVIEW. By the Right Hon. Thomas Babington Macaulay, M.P. 3d Edition. 3 vols. 8vo. 36s. cloth.

MACAULAY.—LAYS OF ANCIENT ROME.

By the Right Honorable Thomas Babington Macaulay, M.P. 7th Edition. Crown 8vo. 10s. 6d. cloth.

MACKAY (CHARLES).—THE SCENERY AND POETRY OF THE ENGLISH LAKES; a Summer Ramble. By Charles Mackay, Esq. author of "Legends of the Isles," "The Salamandrine," "The Thames and its Tributaries," etc. 8vo. with beautiful Wood Engravings from Original Sketches. [Just ready.

MACKENZIE.—THE PHYSIOLOGY OF VISION.

By W. Mackenzie, M.D., Lecturer on the Eye in the University of Glasgow. 8vo. with Woodcuts, 10s. 6d. boards.

MACKINNON.—THE HISTORY OF CIVILISATION.

By Wm. Alexander Mackinnon, F.R.S. M.P. for Lymington. 2 vols. 8vo. 24s. cloth.

MACKINTOSH (SIR JAMES).—THE LIFE OF SIR THOMAS MORE.

By the Right Hon. Sir James Mackintosh. Reprinted from the Cabinet Cyclopædia; and intended for a Present-Book or School Prize. Foolsap 8vo. with Portrait, 5s. cloth; or bound in vellum gilt (*old style*), 8s.

MACKINTOSH'S (SIR JAMES) MISCELLANEOUS WORKS;

Including his Contributions to The EDINBURGH REVIEW. Edited by Robert James Mackintosh, Esq. 3 vols. 8vo. 42s. cloth.

MACKINTOSH, ETC.—THE HISTORY OF ENGLAND.

By Sir James Mackintosh; W. Wallace, Esq.; and Robert Bell, Esq. 10 vols. foolscap 8vo. with Vignette Titles, 3l. cloth.

M'CULLOCH.—THE LITERATURE OF POLITICAL ECONOMY;

Being a Classified Catalogue of the principal Works in the different departments of Political Economy, interspersed with Historical, Critical, and Biographical Notices. By J. R. M'Culloch, Esq. 8vo. 14s. cloth.

M'CULLOCH (J. R.).—AN ACCOUNT, DESCRIPTIVE, AND STATISTICAL,

of the BRITISH EMPIRE; exhibiting its Extent, Physical Capacities, Population, Industry, and Civil and Religious Institutions. By J. R. M'Culloch, Esq. 3d Edition, corrected, enlarged, and greatly improved. 2 thick vols. 8vo. [In the press.

M'CULLOCH.—A TREATISE ON THE PRINCIPLES AND PRACTICAL

INFLUENCE OF TAXATION AND THE FUNDING SYSTEM. By J. R. M'Culloch, Esq. 8vo. 15s. cloth.

M'CULLOCH.—A DICTIONARY, GEOGRAPHICAL, STATISTICAL, AND

HISTORICAL, of the various Countries, Places, and Principal Natural Objects in the World. By J. R. M'Culloch, Esq. A new Edition. 2 vols. 8vo. with Six large Maps, 4l. cloth.

* * The new Articles are printed separately as a Supplement to the former Edition. They comprise a full account of the present state of the United Kingdom, the Oregon Territory, etc. 8vo. 5s. sewed.

M'CULLOCH.—A DICTIONARY, PRACTICAL, THEORETICAL, AND

HISTORICAL, OF COMMERCE, AND COMMERCIAL NAVIGATION. By J. R. M'Culloch, Esq. A New Edition, corrected, enlarged, and improved. 8vo. with Maps and Plans, 50s. cloth; or 55s. strongly half-bound in russia, with flexible back.

* * This Edition, which has been carefully corrected, comprises, besides the New Tariff, the new Acts relating to Banking, the Sugar Trade, Navigation and Customs, the hiring of Seamen, etc.; and is further enriched with valuable information from all parts of the world.

A SUPPLEMENT, for the use of the purchasers of the last Edition, 8vo. price 3s. 6d. sewed.

MAITLAND (DR. CHARLES).—THE CHURCH IN THE CATACOMBS :

A Description of the primitive Church in Rome, Illustrated by its Sepulchral Remains. By Charles Maitland, M.D. 8vo. with very numerous Engravings on Wood.

[*Nearly ready.*]

MAR CET (MRS.)—CONVERSATIONS ON THE HISTORY OF ENGLAND.

For the Use of Children. By Mrs. Marcet, author of "Conversations on Chemistry," etc. 2d Edition, with Additions. 18mo. 5s. cloth.

MAR CET.—CONVERSATIONS ON CHEMISTRY ;

In which the Elements of that Science are familiarly Explained and Illustrated by Experiments. 15th Edition, enlarged and corrected. 2 vols. foolscap 8vo. 14s. cloth.

MAR CET.—CONVERSATIONS ON NATURAL PHILOSOPHY ;

In which the Elements of that Science are familiarly explained, and adapted to the comprehension of Young Persons. 10th Edition, enlarged and corrected. Fcap. 8vo. with 23 Plates, 10s. 6d. cloth.

MAR CET.—CONVERSATIONS ON POLITICAL ECONOMY ;

In which the Elements of that Science are familiarly explained. 7th Edition, revised and enlarged. Foolscap 8vo. 7s. 6d. cloth.

MAR CET.—CONVERSATIONS ON VEGETABLE PHYSIOLOGY ;

Comprehending the Elements of Botany, with their application to Agriculture. 3d Edition. Foolscap 8vo. with Four Plates, 9s. cloth.

MAR CET.—CONVERSATIONS FOR CHILDREN ;

On Land and Water. 3d Edition revised and corrected. Foolscap 8vo., with coloured Map, shewing the comparative Altitude of Mountains, 5s. 6d. cloth.

MAR CET.—CONVERSATIONS ON LANGUAGE,

For Children. By Mrs. Marcet, author of "Willy's Grammar," etc. 18mo. 4s. 6d. cloth.

MAR CET—THE GAME OF GRAMMAR,

With a Book of Conversations, shewing the Rules of the Game, and affording Examples of the manner of playing at it. In a varnished box, or done up as a post 8vo. volume, 8s.

MAR CET.—WILLY'S GRAMMAR ;

Interspersed with Stories, and intended for the Use of Boys. By Mrs. Marcet, author of "Mary's Grammar," etc. New edition. 18mo. 2s. 6d. cloth.

"*A sound and simple work for the earliest ages.*"—Quarterly Review.

MAR CET.—LESSONS ON ANIMALS, VEGETABLES, AND MINERALS.

By Mrs. Marcet, author of "Conversations on Chemistry," etc. New Edition. 18mo. 2s. cloth.

MARGARET RUSSELL :

An Autobiography. In one volume.

[*In the press.*]

MARRIAGE GIFT.

By a Mother. A Legacy to her Children. Post 8vo. 5s. cloth, with gilt edges.

MARRYAT (CAPT.)—THE MISSION ;

Or, Scenes in Africa. Written for Young People. By Captain Marryat, C.B., author of "Peter Simple," "Masterman Ready," "The Settlers in Canada," etc. 2 vols. fcap. 8vo. 12s. cloth.

MARRYAT (CAPT.)—THE SETTLERS IN CANADA.

Written for Young People. By Captain Marryat, C.B. author of "Peter Simple," "Masterman Ready," etc. 2 vols. fcap. 8vo. 12s. cloth.

MARRYAT (CAPT.)—MASTERMAN READY ;

Or, the Wreck of the Pacific. Written for Young People. By Captain Marryat, C.B. author of "Peter Simple," etc. 3 vols. fcap. 8vo. with numerous Engravings on Wood, 22s. 6d. cloth.

* * * The volumes, *separately*, 7s. 6d. each, cloth.

"*The best of Robinson Crusoe's numerous descendants, and one of the most captivating of modern children's books. The only danger is, lest parents should dispute with their children the possession of it.*"—Quarterly Review.

MAUNDER.—THE TREASURY OF KNOWLEDGE,

AND LIBRARY OF REFERENCE. By Samuel Maunder. 16th Edition, revised throughout and enlarged. Foolscap 8vo., with two Frontispieces, 10s. cloth; bound in roan, with gilt edges, 12s.

* * * The principal contents of the present new and thoroughly revised edition of "*The Treasury of Knowledge*," are—a new and enlarged *English Dictionary*, with a *Grammar*, *Verbal Distinctions*, and *Exercises*; a new *Universal Gazetteer*; a *compendious Classical Dictionary*; an *Analysis of History and Chronology*; a *Dictionary of Law Terms*; a new *Synopsis of the British Peerage*; and various useful *tabular Addenda*.

MAUNDER.—THE BIOGRAPHICAL TREASURY:

Consisting of Memoirs, Sketches, and brief Notices of above 12,000 Eminent Persons of all Ages and Nations, from the Earliest Period of History; forming a new and complete Dictionary of Universal Biography. By Samuel Maunder. 5th Edition, revised throughout, and containing a copious Supplement, brought down to 1845. Foolscep 8vo. with Frontispiece, 10s. cloth; bound in roan, with gilt edges, 12s.

MAUNDER.—THE SCIENTIFIC AND LITERARY TREASURY:

A New and Popular Encyclopædia of Science and the Belles Lettres; including all Branches of Science, and every Subject connected with Literature and Art. The whole written in a familiar style, adapted to the comprehension of all persons desirous of acquiring information on the subjects comprised in the work, and also adapted for a Manual of convenient Reference to the more instructed. By Samuel Maunder. 4th Edition. Fcap.8vo. with Frontispiece, 10s. cloth; bound in roan, with gilt edges, 12s.

MAUNDER.—THE TREASURY OF HISTORY;

Comprising a General Introductory Outline of Universal History, Ancient and Modern, and a Series of separate Histories of every principal Nation that exists; developing their Rise, Progress, and Present Condition, the Moral and Social Character of their respective Inhabitants, their Religion, Manners, and Customs, etc. etc. By Samuel Maunder. 2d Edit. Fcap. 8vo. with Frontispiece, 10s. cloth; bound in roan, with gilt edges, 12s.

MAUNDER.—THE UNIVERSAL CLASS-BOOK:

A new Series of Reading Lessons (original and selected) for Every Day in the Year. With Questions for Examination. By Samuel Maunder, author of "The Treasury of Knowledge," etc. 2d Edition, revised. 12mo. 5s. bound.

MICHELET (J.).—PRIESTS, WOMEN, AND FAMILIES.

By J. Michelet. Translated from the French (third edition), with the Author's permission, by C. Cocks, B.L. Professor of the Living Languages in the Royal Colleges of France. New Editions. Post 8vo. 9s. cloth. 16mo. 1s. 4d. sewed.

MICHELET (J.).—THE PEOPLE.

By M. Michelet, Member of the Institute of France, etc. Translated, with the approbation of the Author, by C. Cocks, B.L. Professor of the Living Languages in the Royal Colleges of France. New Edition. Post 8vo. 9s. cloth. 16mo. 1s. 6d. scwed.

* * Mr. Cocks's authorised translations of Michelet's "*Priests, Women, and Families*," and "*The People*," in one vol. 16mo. 3s. 6d. cloth.

MICHELET AND QUINET.—THE JESUITS

By J. Michelet, Member of the Institute of France; and E. Quinet, of the College of France. Translated, with the Authors' approbation, by C. Cocks, B.L. Professor of the Living Languages in the Royal Colleges of France. 16mo. uniform with the authorised editions of Michelet's "*Priests, Women, and Families*," and "*The People*." [Nearly ready.

MIGNET (M.).—ANTONIO PEREZ AND PHILIP II. OF SPAIN.

By M. Mignet, Member of the Institute of France, Perpetual Secretary of the Academy of Moral and Political Sciences, etc. Translated by C. Cocks, B. L. Professor of the Living Languages in the Royal Colleges of France. Post 8vo. 9s. cloth.

MILES (W.).—THE HORSE'S FOOT,

And How to Keep it Sound. By William Miles, Esq. Royal 8vo. with Engravings, 7s. cloth.

MILNER (REVS. J. AND I.).—THE HISTORY OF THE CHURCH OF

CHRIST. By the Rev. Joseph Milner, A.M. With Additions and Corrections by the late Rev. Isaac Milner, D.D. F.R.S. A New Edition. 4 vols. 8vo. 2l. 8s. boards.

MOHUN LÂL.—THE LIFE OF THE AMIR DOST MAHOMED KHAN OF

CABOUL. By Mohun Lâl, Knight of the Persian Order of the Sun; lately attached to the Mission at Caboul; author of "*Travels in the Punjab, Afghanistan*," etc. 8vo. with numerous Portraits. [2 vols. in the press.

MONTAUBAN (MRS. E).—A YEAR AND A DAY IN THE EAST;

Or, Wanderings over Land and Sea. By Mrs. Eliot Montauban. Post 8vo. 7s. cloth.

MONTGOMERY'S (JAMES) POETICAL WORKS.

New and only complete Edition. With some additional Poems and Autobiographical Prefaces. Collected and edited by Mr. Montgomery. 4 vols. foolscap 8vo. with Portrait, and seven other Plates, 20s. cloth; bound in morocco, 1l. 16s.

MOORE.—THE HISTORY OF IRELAND.

From the earliest Kings of that Realm, down to its latest Chief. By Thomas Moore, Esq. 4 vols. foolscap 8vo., with Vignette Titles, 24s. cloth.

MOORE'S POETICAL WORKS;

Containing the Author's recent Introduction and Notes. Complete in one volume, uniform with Lord Byron's Poems. With a New Portrait, by George Richmond, engraved in the line manner, and a View of Sloperton Cottage, the Residence of the Poet, by Thomas Creswick, A.R.A. Medium 8vo. 1l. 1s. cloth; or 42s. bound in morocco, by Hayday.

* * Also, an Edition in 10 vols. foolscap 8vo. with Portrait, and 19 Plates, 2l. 10s. cloth; morocco, 4l. 10s.

MOORE'S LALLA ROOKH. AN ORIENTAL ROMANCE.

New Edition. Medium 8vo. illustrated with 13 fine Engravings, 2ls. cloth; morocco, 35s.; with India Proof Plates, 35s. cloth.

MOORE'S LALLA ROOKH. AN ORIENTAL ROMANCE.

New Edition. Foolsap 8vo. with 4 Plates, by Westall, 10s. 6d. cloth; or 14s. bound in morocco.

MOORE'S IRISH MELODIES.

Illustrated by D. MacIise, R.A. Imp. 8vo. with 161 Designs, engraved on Steel, 3l. 3s. boards; or 4l. 14s. 6d. bound in morocco, by Hayday. Proof Impressions (only 200 copies printed, of which a few remain), 6l. 6s. boards.

*** India Proofs before letters of the 161 Designs, on Quarter Colombier, in Portfolio (only 25 copies printed, of which a few remain), 31l. 10s.*

India Proofs before letters of the 51 Large Designs, on Quarter Colombier, in Portfolio (only 25 copies printed, of which a few remain), 18l. 18s.

MOORE'S IRISH MELODIES.

New Edition. Fcap. 8vo. with Vignette Title, 10s. cloth; bound in morocco, 13s. 6d.

MORAL OF FLOWERS (THE).

3d Edition. Royal 8vo. with 24 beautifully coloured Engravings, 1l. 10s. half-bound.

MORTON.—A VETERINARY TOXICOLOGICAL CHART,

Containing those Agents known to cause Death in the Horse; with the Symptoms, Antidotes, Action on the Tissues, and Tests. By W. J. T. Morton. 12mo. 6s. in case; 8s. 6d. on rollers.

MORTON.—A MANUAL OF PHARMACY,

For the Student in Veterinary Medicine; containing the Substances employed at the Royal Veterinary College, with an Attempt at their Classification, and the Pharmacopœia of that Institution. By W. J. T. Morton. 3d Edition, 12mo. 10s. cloth.

MOSELEY.—ILLUSTRATIONS OF PRACTICAL MECHANICS.

By the Rev. H. Moseley, M.A., Professor of Natural Philosophy and Astronomy in King's College, London; being the First Volume of the Illustrations of Science by the Professors of King's College. Fcap. 8vo. with numerous Woodcuts, 8s. cloth.

MOSELEY.—THE MECHANICAL PRINCIPLES OF ENGINEERING AND ARCHITECTURE.

By the Rev. H. Moseley, M.A. F.R.S., Professor of Natural Philosophy and Astronomy in King's College, London; and author of "Illustrations of Practical Mechanics," etc. 8vo. with Woodcuts and Diagrams, 1l. 4s. cloth.

MOSHEIM'S ECCLESIASTICAL HISTORY,

Ancient and Modern. Translated, with copious Notes, by James Murdock, D.D. Edited, with Additions, by Henry Soames, M.A. Rector of Stapleford-Tawney, Essex. New Edition, revised, and continued to the present time. 4 vols. 8vo. 48s. cloth.

MÜLLER.—INTRODUCTION TO A SCIENTIFIC SYSTEM OF MYTHOLOGY.

By C. O. Müller, author of "The History and Antiquities of the Doric Race," etc. Translated from the German by John Leitch. 8vo. uniform with "Müller's Dorians," 12s. cloth.

MURRAY.—AN ENCYCLOPÆDIA OF GEOGRAPHY;

Comprising a complete Description of the Earth: exhibiting its Relation to the Heavenly Bodies, its Physical Structure, the Natural History of each Country, and the Industry, Commerce, Political Institutions, and Civil and Social State of all Nations. By Hugh Murray, F.R.S.E.: assisted in Astronomy, etc. by Professor Wallace; Geology, etc. by Professor Jameson; Botany, etc. by Sir W. J. Hooker; Zoology, etc. by W. Swainson, Esq. New Edition, with Supplement, bringing down the Statistical Information contained in the Work, to December 1843: with 82 Maps, drawn by Sidney Hall, and upwards of 1,000 other Engravings on Wood. 8vo. 3l. cloth.

NECKER DE SAUSSURE.—PROGRESSIVE EDUCATION;

Or, Considerations on the Course of Life. Translated and Abridged from the French of Madame Necker De Saussure, by Miss Holland. 3 vols. foolscap 8vo. 19s. 6d. cloth.

*** Separately, vols. I. and II. 12s.; vol. III. 7s. 6d.*

NESBIT (A.).—A TREATISE ON PRACTICAL MENSURATION;

Containing the most approved Methods of drawing Geometrical Figures; Mensuration of Superficies; Land-Surveying; Mensuration of Solids; the Use of the Carpenter's Rule; Timber Measure, etc. By A. Nesbit. 12th Edition. 12mo. with nearly 300 Woodcuts, 6s. bound.

KEY. 7th Edition. 12mo. 5s. bound.

NEWELL (REV. R.H.).—THE ZOOLOGY OF THE ENGLISH POETS,

Corrected by the Writings of Modern Naturalists. By the Rev. R.H. Newell, Rector of Little Horstead. Fcap. 8vo. with Engravings on Wood, 5s. 6d., cloth.

"This interesting little volume has for its object the pointing out some of the more prominent passages in which our English poets have erred with respect to the structure and habits of animals. Birds, reptiles, and mammalia, are successively spoken of by the author, and practical errors with regard to them are held up to the light of modern science. As a notebook for correcting false impressions, and the imparting sound ones, Mr. Newell's work will be found useful in the education of young people."—Athenæum.

NICOLAS—THE CHRONOLOGY OF HISTORY,

Containing Tables, Calculations, and Statements indispensable for ascertaining the Dates of Historical Events, and of Public and Private Documents, from the Earliest Period to the Present Time. By Sir Harris Nicolas, K.C. M.G. Second Edition, corrected throughout. Foolscep 8vo. with Vignette Title, 6s. cloth.

ORDNANCE MAPS (THE), AND PUBLICATIONS OF THE GEOLOGICAL SURVEY OF THE UNITED KINGDOM.—Messrs. Longman and Co. have been appointed by the Board of Ordnance Agents for the sale of the Maps of the Ordnance Survey of Great Britain. Also, sole Agents for the sale of the Maps, Sections, and Books of the Geological Survey of the United Kingdom, and of the Museum of Economic Geology, under the Chief Commissioner of Her Majesty's Woods, Works, and Land Revenues.

. *Complete detailed Catalogues of both Series may be had on application.*

OWEN.—LECTURES ON THE COMPARATIVE ANATOMY AND PHYSIOLOGY OF THE INVERTEBRATE ANIMALS, delivered at the Royal College of Surgeons in 1843. By Richard Owen, F.R.S. Hunterian Professor to the College. From Notes taken by William White Cooper, M.R.C.S. and revised by Professor Owen. With Glossary and Index. 8vo. with nearly 140 Illustrations on Wood, 14s. cloth.

. *A second and concluding Volume, being the Lectures on Vertebrata delivered by Professor Owen during the last session, is preparing for publication.*

PARABLES (THE).

The Parables of Our Lord, richly Illuminated with appropriate Borders, printed in Colours, and in Black and Gold; with a Design from one of the early German engravers. Square foolscep 8vo., uniform in size with the "Sermon on the Mount," 21s., in massive carved binding; or 30s. bound in morocco, by Hayday. [Just ready.

PARKES.—DOMESTIC DUTIES;

Or, Instructions to Young Married Ladies on the Management of their Households and the Regulation of their Conduct in the various Relations and Duties of Married Life. By Mrs. W. Parkes. 5th Edition. Foolscep 8vo. 9s. cloth.

PARNELL.—A TREATISE ON ROADS;

Wherein the Principles on which Roads should be made are explained and illustrated by the Plans, Specifications, and Contracts made use of by Thomas Telford, Esq. on the Holyhead Road. By the Right Hon. Sir Henry Parnell, Bart., Hon. Memb. Inst. Civ. Eng. London. Second Edition, greatly enlarged. 8vo. with 9 large Plates, 17. 1s. cloth.

PATON (A. A.)—SERVIA, THE YOUNGEST MEMBER OF THE EUROPEAN FAMILY; or, a Residence in Belgrade, and Travels through the Highlands and Woodlands of the Interior, during the years 1843 and 1844. By Andrew Archibald Paton, Esq. Post 8vo. with portrait and plate, 12s. cloth.

By the same Author.

THE MODERN SYRIANS; or, Native Society in Damascus, Aleppo, and the Mountains of the Druses. Post 8vo. 10s. 6d. cloth.

PEARSON.—AN INTRODUCTION TO PRACTICAL ASTRONOMY:

By the Rev. W. Pearson, LL.D. F.R.S. etc. Rector of South Kilworth, Leicestershire, and Treasurer to the Astronomical Society of London. 2 vols. 4to. with Plates, 7l. 7s. boards.

PEDESTRIAN AND OTHER REMINISCENCES AT HOME AND ABROAD, WITH SKETCHES OF COUNTRY LIFE. By Sylvanus. Post 8vo., with Frontispiece and Vignette Title, 10s. 6d. cloth.

PERCIVALL.—THE ANATOMY OF THE HORSE;

Embracing the Structure of the Foot. By W. Percivall, M.R.C.S. 8vo. 17. cloth.

PERCIVALL.—HIPPOPATHOLOGY:

A Systematic Treatise on the Disorders and Lameness of the Horse; with their Modern and most approved Methods of Cure; embracing the Doctrines of the English and French Veterinary Schools. By W. Percivall, M.R.C.S., Veterinary Surgeon in the First Life Guards. 3 vols. 8vo. Vol. 1, 10s. 6d.; vols. 2 and 3, 14s. each, boards.

PEREIRA.—A TREATISE ON FOOD AND DIET:

With Observations on the Dietetical Regimen suited for Disordered States of the Digestive Organs; and an Account of the Dietaries of some of the principal Metropolitan and other Establishments for Paupers, Lunatics, Criminals, Children, the Sick, etc. By Jon. Pereira, M.D. F.R.S., author of "Elements of Materia Medica." 8vo. 16s. cloth.

PERICLES:

A Tale of Athens in the 83d Olympiad. By the author of "A Brief Sketch of Greek Philosophy." 2 vols. post 8vo. 18s. cloth.

"All these classical scenes are vigorously drawn; and 'Pericles' must be pronounced one of the happiest examples ever yet produced of a graceful fiction, based on a profound living knowledge of ancient Greek history and manners."—Britannia.

PERRY (DR. W. C.)—GERMAN UNIVERSITY EDUCATION:

Or, the Professors and Students of Germany. By Walter C. Perry, Phil. D. of the University of Göttingen. Post 8vo. 4s. 6d. cloth.

PESCHEL (C. F.)—ELEMENTS OF PHYSICS,

Part I. Ponderable Bodies. By C. F. Peschel, Principal of the Royal Military College, Dresden, etc. etc. Translated from the German, with Notes, by E. West. Foolsap 8vo. with Diagrams and Woodcuts. 7s. 6d. cloth.

Part 2.—The Physics of Inponderable Bodies, is nearly ready.

PHILLIPS.—AN ELEMENTARY INTRODUCTION TO MINERALOGY:

Comprising a Notice of the Characters and Elements of Minerals; with Accounts of the Places and Circumstances in which they are found. By William Phillips, F.L.S. M.G.S. etc. 4th Edition, considerably augmented by R. Allan, F.R.S.E. 8vo. with woodcuts, 12s. cloth.

PHILLIPS.—FIGURES & DESCRIPTIONS OF THE PALÆOZOIC FOSSILS OF

CORNWALL, DEVON, and WEST SOMERSET; observed in the course of the Ordnance Geological Survey of that District. By John Phillips, F.R.S. F.G.S. etc. Published by Order of the Lords Commissioners of H. M. Treasury. 8vo. with 60 Plates, comprising very numerous Figures, 9s. cloth.

PHILLIPS.—A GUIDE TO GEOLOGY.

By John Phillips, F.R.S. G.S. etc. Foolsap 8vo. with Plates, 5s. cloth.

PHILLIPS.—A TREATISE ON GEOLOGY.

By John Phillips, F.R.S. G.S. etc. 2 vols. foolsap 8vo. with Woodcuts. 12s. cloth.

PITMAN (REV. J. R.)—SERMONS

On the principal Subjects comprised in the Book of Psalms, abridged from Eminent Divines. By the Rev. J. R. Pitman, A.M. Minister of St. Barnabas Church, Kensington. 8vo.

PLYMLEY (PETER).—LETTERS ON THE SUBJECT OF THE CATHOLICS TO MY BROTHER ABRAHAM, WHO LIVES IN THE COUNTRY.

By Peter Plymley. 21st Edition. Post 8vo. 7s. cloth.

POET'S PLEASANCE (THE);

Or, Garden Full of all Sorts of Pleasant Flowers, which our Pleasant Poets have in Past Time (for Pastime) Planted: with the right ordering of them. By Eden Warwick. In 1 vol. with Border Illustrations. *[In the press.]*

POISSON (S. D.)—A TREATISE ON MECHANICS.

By S. D. Poisson. Second Edition. Translated from the French, and illustrated with Explanatory Notes, by the Rev. Henry H. Harte, late Fellow of Trinity College, Dublin. 2 volumes, 8vo. 17. 8s. cloth.

POPE (ALEXANDER).—THE WORKS OF ALEXANDER POPE.

Edited by Thomas Roscoe, Esq. With the Author's Life. A New Edition. 8 vols. 8vo. *[In the press.]*

PORTER.—A TREATISE ON THE MANUFACTURE OF SILK.

By G. R. Porter, Esq. F.R.S., author of "The Progress of the Nation," etc. Fcap. 8vo. with Vignette Title, and 39 Engravings on Wood, 6s. cloth.

PORTER.—A TREATISE ON THE MANUFACTURES OF PORCELAIN AND GLASS.

By G. R. Porter, Esq. F.R.S. Foolsap 8vo. with 50 Woodcuts, 6s. cloth.

PORTLOCK. —REPORT ON THE GEOLOGY OF THE COUNTY OF

LONDONDERRY, and of Parts of Tyrone and Fermanagh, examined and described under the Authority of the Master-General and Board of Ordnance. By J. E. Portlock, F.R.S. etc. 8vo. with 48 Plates, 24s. cloth.

POWELL.—THE HISTORY OF NATURAL PHILOSOPHY.

From the Earliest Periods to the Present Time. By Baden Powell, M. A., Savilian Professor of Mathematics in the University of Oxford. Fcap. 8vo. with Vignette Title, 6s. cloth.

PYCROFT.—A COURSE OF ENGLISH READING;

Adapted to every Taste and Capacity. With Anecdotes of Men of Genius. By the Rev. James Pycroft, B.A., Trinity College, Oxford, author of "Greek Grammar Practice," "Latin Grammar Practice," etc. Foolsap 8vo. 6s. 6d. cloth.

QUARTERLY JOURNAL OF THE GEOLOGICAL SOCIETY OF LONDON.

Edited by David Thomas Ansted, M.A. F.R.S., Fellow of Jesus College, Cambridge; Professor of Geology in King's College, London; Vice-Secretary of the Geological Society. 8vo. 4s. each number, sewed. *[Published Quarterly.]*

RANKE (PROFESSOR).—RANKE'S HISTORY OF THE REFORMATION.

Translated by Sarah Austin, translator of Ranke's "History of the Popes." Vols. 1 and 2; 8vo. 30s. cloth. *[Vol. II. in the press.]*

REECE.—THE MEDICAL GUIDE:

For the use of the Clergy, Heads of Families, Seminaries, and Junior Practitioners in Medicine; comprising a complete Modern Dispensatory, and a Practical Treatise on the distinguishing Symptoms, Causes, Prevention, Cure, and Palliation of the Diseases incident to the Human Frame. By R. Reece, M.D., late Fellow of the Royal College of Surgeons of London, etc. 16th Edition. 8vo. 12s. boards.

REID (DR.)—ILLUSTRATIONS OF THE THEORY AND PRACTICE OF VENTILATION: with Remarks on Warming, Exclusive Lighting, and the Communication of Sound. By D. B. Reid, M.D. F.R.S.E. etc. 8vo. with Diagrams, and 320 Engravings on Wood, 16s. cloth.

REPTON.—THE LANDSCAPE GARDENING & LANDSCAPE ARCHITECTURE of the late Humphrey Repton, Esq.; being his entire Works on these subjects. A New Edition, with an Historical and Scientific Introduction, a systematic Analysis, a Biographical Notice, Notes, and a copious Alphabetical Index. By J. C. Loudon, F.L.S., etc. 8vo. with a Portrait and upwards of 250 Engravings, 30s. cloth; with coloured Plates, 3l. 6s. cloth.

REYNARD THE FOX;

A renowned Apologue of the Middle Age. Reproduced in Rhyme. Embellished throughout with Scroll Capitals, in Colours, from Wood-block Letters made expressly for this work, after Designs of the 12th and 13th Centuries. With an Introduction. By Samuel Naylor, late of Queen's College, Oxford. Large square 8vo. 18s. cloth.

RIDDLE—A COMPLETE ENGLISH-LATIN AND LATIN-ENGLISH DICTIONARY, from the best sources, chiefly German. By the Rev. J. E. Riddle, M.A. 4th Edition. 8vo. 3ls. 6d. cloth.

* * Separately—The English-Latin Dictionary, 10s. 6d.; the Latin-English Dictionary, 2ls.

RIDDLE.—A DIAMOND LATIN-ENGLISH DICTIONARY.

A Guide to the Meaning, Quality, and right Accentuation of Latin Classical Words. By the Rev. J. E. Riddle, M.A. Royal 32mo. 4s. bound.

RIDDLE.—LETTERS FROM AN ABSENT GODFATHER;

Or, a Compendium of Religious Instruction for Young Persons. By the Rev. J. E. Riddle, M.A. Foolsap 8vo. 6s. cloth.

RIDDLE.—ECCLESIASTICAL CHRONOLOGY;

Or, Annals of the Christian Church, from its Foundation to the present Time. Containing a View of General Church History, and the Course of Secular Events; the Limits of the Church and its Relations to the State; Controversies; Sects and Parties; Rites, Institutions, and Discipline; Ecclesiastical Writers. The whole arranged according to the order of Dates, and divided into Seven Periods. To which are added, Lists of Councils and Popes, Patriarchs, and Archbishops of Canterbury. By the Rev. J. E. Riddle, M.A. 8vo. 15s. cloth.

RITCHIE (ROBERT).—RAILWAYS: THEIR RISE AND PROGRESS, AND CONSTRUCTION, with Remarks on Railway Accidents, and Proposals for their prosecution. By Robert Ritchie, Esq., F.R.S., S.A., Civil Engineer, Associate of the Institution of Civil Engineers, etc. Fcap. 8vo. with Woodcuts and Diagrams, 9s. cloth.

RIVERS.—THE ROSE AMATEUR'S GUIDE:

Containing ample Descriptions of all the fine leading varieties of Roses, regularly classed in their respective Families; their History and mode of Culture. By T. Rivers, Jun. Third Edition, corrected and improved. Foolsap 8vo. 6s. cloth.

ROBERTS (GEORGE).—THE LIFE, PROGRESSES, AND REBELLION OF JAMES DUKE OF MONMOUTH, to his Capture and Execution: with a full Account of the Bloody Assizes, and copious Biographical Notices. By George Roberts. 2 vols. post 8vo. with Portrait, Maps, and other Illustrations, 24s. cloth.

ROBERTS.—AN ETYMOLOGICAL AND EXPLANATORY DICTIONARY OF THE TERMS AND LANGUAGE OF GEOLOGY; designed for the early Student, and those who have not made great progress in the Science. By G. Roberts. Foolsap 8vo. 6s. cloth.

ROBINSON (JAMES).—THE WHOLE ART OF CURING, PICKLING, and SMOKING EVERY DESCRIPTION OF MEAT AND FISH, according to both the British and Foreign Modes. To which is appended, a Description of the requisite Apparatus. By James Robinson, Eighteen Years a Practical Curer. Fcp. 8vo. [Just ready.]

ROBINSON—GREEK AND ENGLISH LEXICON TO THE NEW TESTAMENT. By E. Robinson, D.D., author of "Biblical Researches." Edited, with careful revision, corrections, etc., by the Rev. Dr. Bloomfield. 8vo. 18s. cloth.

ROGERS.—THE VEGETABLE CULTIVATOR;

Containing a plain and accurate Description of all the different Species of Culinary Vegetables, with the most approved Method of Cultivating them by Natural and Artificial Means, and the best Modes of Cooking them; alphabetically arranged. Together with a Description of the Physical Herbs in General Use. Also, some Recollections of the Life of Philip Miller, F.A.S., Gardener to the Worshipful Company of Apothecaries at Chelsea. By John Rogers, author of "The Fruit Cultivator." 2d Edition. Foolsap 8vo. 7s. cloth.

ROME.—THE HISTORY OF ROME (IN THE CABINET CYCLOPÆDIA). 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

ROSCOE.—LIVES OF EMINENT BRITISH LAWYERS.

By Henry Roscoe, Esq. Foolsap 8vo. with Vignette Title, 6s. cloth.

ROWTON (F.)—THE DEBATER;

A New Theory of the Art of Speaking. Being a Series of complete Debates, Questions for Discussion, and Rules of Debate. With ample references to the best sources of information upon each particular topic. For the use of Schools and Discussion Societies. By Frederick Rowton, Lecturer on General Literature. *[In the press.]*

SANDBY (REV. G.)—MESMERISM AND ITS OPPONENTS:

With a Narrative of Cases. By the Rev. George Sandby, Jun., Vicar of Flixton, and Rector of All Saints with St. Nicholas, South Elmham, Suffolk; Domestic Chaplain to the Right Hon. the Earl of Abergavenny. Foolscap 8vo. 6s. cloth.

SANDFORD (REV. JOHN).—PAROCHIALIA,

or Church, School, and Parish. By the Rev. John Sandford, M.A. Vicar of Dunchurch, Chaplain to the Lord Bishop of Worcester, Hon. Canon of Worcester, and Rural Dean. 8vo. with numerous Woodcuts, 16s. cloth.

SANDFORD.—WOMAN IN HER SOCIAL AND DOMESTIC CHARACTER.

By Mrs. John Sandford. 6th Edition. Foolscap 8vo. 6s. cloth.

SANDFORD.—FEMALE IMPROVEMENT.

By Mrs. John Sandford. 2d Edition. Foolscap 8vo. 7s. 6d. cloth.

SANDHURST COLLEGE MATHEMATICAL COURSE.

1. **ELEMENTS OF ARITHMETIC AND ALGEBRA.** By W. Scott, Esq., A.M. and F.R.A.S. Second Mathematical Professor at the Royal Military College, Sandhurst. 8vo. 16s. bound.
2. **ELEMENTS OF GEOMETRY**; consisting of the first Four and Sixth Books of Euclid, chiefly from the Text of Dr. Robert Simson; with the principal Theorems in Proportion, and a Course of Practical Geometry on the Ground; also, Four Tracts relating to Circles, Planes, and Solids, with one on Spherical Geometry. By John Narrien, Professor of Mathematics in the Royal Military College, Sandhurst. 8vo. with many Diagrams, 10s. 6d. bound.
3. **PLAIN TRIGONOMETRY AND MENSURATION**; for the use of the Royal Military College, Sandhurst. By W. Scott, Esq. A.M. and F.R.A.S., Second Mathematical Master in Institution. 8vo. 9s. 6d. bound.
5. **PRACTICAL ASTRONOMY and GEODESY**; Including the Projections of the Sphere, and Spherical Trigonometry. By John Narrien, F.R.S. and R.A.S. Professor of Mathematics in the Royal Military College, Sandhurst. 8vo. 14s. bound.

SCHLEIDEN (PROF.)—PRINCIPLES OF SCIENTIFIC BOTANY.

By M. J. Schleiden, Professor of Botany at Jena. Translated by E. Lankester, M.D. F.L.S. 8vo. with numerous wood engravings. *[In the press.]*

SCOTT.—THE HISTORY OF SCOTLAND.

By Sir Walter Scott, Bart. New Edition. 2 vols. fcap. 8vo. with Vignette Titles, 12s. cloth.

SEAWARD.—SIR EDWARD SEAWARD'S NARRATIVE OF HIS SHIPWRECK,

and consequent Discovery of certain Islands in the Caribbean Sea; with a Detail of many extraordinary and highly interesting Events in his Life, from 1733 to 1749, as written in his own Diary. Edited by Miss Jane Porter. 3d Edition, with a New Nautical and Geographical Introduction, containing Extracts from a Paper by Mr. C. F. Collett, of the Royal Navy, identifying the Islands described by Sir Edward Seaward. 2 vols. post 8vo. 21s. cloth.

SELECT WORKS OF THE BRITISH POETS:

From Chaucer to Withers. With Biographical Sketches, by R. Southey, LL.D. Medium 8vo. 30s. cloth; or with gilt edges, 31s. 6d.

SELECT WORKS OF THE BRITISH POETS:

With Biographical and Critical Prefaces by Dr. Aikin. A New Edition, with Supplement by Lucy Aikin, consisting of Selections from the Works of Crabbe, Scott, Coleridge, Pringle, Charlotte Smith, and Mrs. Barbauld. Medium 8vo. 18s. cloth.

* * *The peculiar feature of these two works is, that the Poems included are printed entire, without mutilation or abridgment.*

SERMON ON THE MOUNT (THE).

[St. Matthew v. vi. vii.] Intended for a Birthday-Present, or Gift-Book for all Seasons. Printed in Gold and Colours, in the Missal Style, with Ornamental Borders by Owen Jones, Architect, and an Illuminated Frontispiece by W. Boxall, Esq. A new edition. Foolscap 4to. in a rich brocaded silk cover, manufactured expressly, 21s.; or bound in morocco, in the Missal style, by Hayday, 25s.

SHAKSPEARE, BY BOWDLER.

THE FAMILY SHAKSPEARE, in which nothing is added to the Original Text; but those Words and Expressions are omitted which cannot with propriety be read aloud. By T. Bowdler, Esq. F.R.S. Seventh Edition, 8vo. with 36 Illustrations after Smirke, etc. 30s. cloth; with gilt edges, 31s. 6d.

* * A LIBRARY EDITION, without Illustrations, 8 vols. 8vo. 4l. 14s. 6d. boards.

SHELDON (F.)—THE MINSTRELSY OF THE ENGLISH BORDER :

Being a Collection of Ballads, Ancient, Re-modelled, and Original, founded on well-known Border Legends. With Illustrative Notes. By Frederick Sheldon. Square foolscap 8vo. [In the press.]

SHELLEY, ETC.—LIVES OF THE MOST EMINENT LITERARY MEN OF ITALY, SPAIN and PORTUGAL. By Mrs. Shelley, Sir D. Brewster, J. Montgomery, etc. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

SHELLEY.—LIVES OF THE MOST EMINENT FRENCH WRITERS.

By Mrs. Shelley and others. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

SHEPHERD (REV. W.)—HORÆ APOSTOLICÆ ;

Or, a Digested Narrative of the Acts, Lives, and Writings of the Apostles. By the Rev. William Shepherd, B.D. Rector of Margaret Roding, Essex, and Rural Dean. Fcp. 8vo. [In the press.]

SHORT WHIST :

Its Rise, Progress, and Laws ; with Observations to make any one a Whist Player ; containing also the Laws of Piquet, Cassino, Ecarte, Cribbage, Backgammon. By Major A * * * *. 9th Edition. To which are added, Precepts for Tyros. By Mrs. B * * * *. Foolscap 8vo. 3s. cloth, gilt edges.

SMITH (GEORGE).—THE RELIGION OF ANCIENT BRITAIN :

Or, a Succinct Account of the several Religious Systems which have obtained in this Island from the Earliest Times to the Norman Conquest : including an Investigation into the Early Progress of Error in the Christian Church, the Introduction of the Gospel into Britain, and the State of Religion in England till Popery had gained the Ascendency. By George Smith, F.A.S. Member of the Royal Asiatic Society, and of the Royal Society of Literature. Foolscap. 8vo. 7s. cloth.

SMITH (GEORGE).—PERILOUS TIMES :

Or, the Aggressions of Anti-Christian Error on Scriptural Christianity : considered in reference to the Dangers and Duties of Protestants. By George Smith, F.A.S. Member of the Royal Asiatic Society, and of the Royal Society, of Literature. Foolscap 8vo. 6s. cloth.

SMITH (MRS. H.)—THE FEMALE DISCIPLE OF THE FIRST THREE CENTURIES OF THE CHRISTIAN ERA : Her Trials and Her Mission. By Mrs. Henry Smith. Foolscap 8vo. 6s. cloth.

SMITH.—AN INTRODUCTION TO THE STUDY OF BOTANY.

By Sir J. E. Smith, late President of the Linnæan Society. 7th Edition, corrected ; in which the object of Smith's "Grammar of Botany" is combined with that of the "Introduction." By Sir William Jackson Hooker, K.H. LL.D. etc. 8vo. with 36 Steel Plates, 16s. cloth ; with coloured Plates, 2l. 12s. 6d. cloth.

SMITH—COMPENDIUM OF THE ENGLISH FLORA.

By Sir J. E. Smith. 2d Edition, with Additions and Corrections. By Sir W. J. Hooker. 12mo. 7s. 6d. cloth. THE SAME IN LATIN. 5th Edition, 12mo. 7s. 6d.

SMITH.—THE ENGLISH FLORA.

By Sir James Edward Smith, M.D. F.R.S., late President of the Linnæan Society, etc. 6 vols. 8vo. 3l. 12s. boards.

CONTENTS :

Vols. I. to IV. the Flowering Plants and the Ferns, 2l. 8s.

Vol. V. Part 1, 12s. —Cryptogamia ; comprising the Mosses, Hepaticæ, Lichens, Characeæ, and Algæ. By Sir W. J. Hooker.

Vol. V. Part 2, 12s. —The Fungi—completing the work, by Sir W. J. Hooker, and the Rev. M. J. Berkeley, F.L.S. etc.

SMITH (SYDNEY).—SERMONS PREACHED AT ST. PAUL'S CATHEDRAL. The Foundling Hospital, and several Churches in London ; together with others addressed to a Country Congregation. By the late Rev. Sydney Smith, Canon Residentiary of St. Paul's Cathedral. 8vo. 12s. cloth.

SMITH.—THE WORKS OF THE REV. SYDNEY SMITH.

3d Edition. 3 vols. 8vo. with Portrait, 36s. cloth.

* * * This collection consists of the Author's contributions to the "Edinburgh Review," Peter Plymley's "Letters on the Catholics," and other miscellaneous works ; to which are now first added—"Three Letters on Railways ;" "Letter to Mr. Horner ;" "Two Letters on American Debts ;" "A Prayer ;" "Changes" (never before published) ; "A Fragment on the Irish Roman-Catholic Church."

SOPHOCLES, BY LINWOOD.

SOPHOCLES TRAGÆDIÆ SUPERSTITES. Recensuit, et brevi adnotatione instruxit Gulielmus Linwood, A.M. Aedis Christi apud Oxonienses Alumnus. 8vo. 16s. cloth.

SOUTHEY (ROBERT).—THE LATE MR. SOUTHEY'S COMMON-PLACE BOOK; comprising his Readings and Collections in History, Biography, Manners and Literature, Voyages and Travels, etc. etc.; systematically arranged. [*In the press.*]

SOUTHEY (ROBERT).—SOUTHEY'S LIFE OF WESLEY.
A New Edition. Edited by his Son, the Rev. Cuthbert C. Southey. 8vo. [*In the press.*]

SOUTHEY, ETC.—LIVES OF THE BRITISH ADMIRALS;
With an Introductory View of the Naval History of England. By R. Southey, Esq. and R. Bell, Esq. 5 vols. foolscap 8vo., with Vignette Titles, 1*l.* 10*s.* cloth.

SOUTHEY (ROBERT).—OLIVER NEWMAN;
A New England Tale (unfinished): with other Poetical Remains. By the late Robert Southey. Foolscap 8vo. uniform with the Ten Volume Edition of Mr. Southey's Poetical Works, 5*s.* cloth.

SOUTHEY'S (ROBERT) COMPLETE POETICAL WORKS:
Containing all the Author's last Introductions and Notes. Complete in one volume, medium 8vo. with Portrait and View of the Poet's Residence at Keswick, uniform with Byron's Poems and Thomas Moore's Poetical Works, 2*l.*s.; or 4*2s.* bound in morocco, by Hayday.
Also, an Edition in 10 vols. foolscap 8vo. with Portrait and 19 Plates, 2*l.* 10*s.*; morocco, 4*l.* 10*s.*

The following Works separately:—

JOAN OF ARC - - -	Fcap. 8vo. 5 <i>s.</i> cloth.	THALABA - - -	Fcap. 8vo. 5 <i>s.</i> cloth.
MADOC - - - - -	" 5 <i>s.</i> "	BALLADS, etc. 2 vols.	" 10 <i>s.</i> "
CURSE OF KEHAMA	" 5 <i>s.</i> "	RODERICK - - - -	" 5 <i>s.</i> "

SPIRIT OF THE WOODS (THE).

By the author of "The Moral of Flowers." 2d Edition. Royal 8vo. with 23 beautifully coloured Engravings of the Forest Trees of Great Britain, 1*l.* 1*l.*s. 6*d.* cloth.

SPOONER.—A TREATISE ON THE STRUCTURE, FUNCTIONS, AND DISEASES OF THE FOOT AND LEG OF THE HORSE; comprehending the Comparative Anatomy of these Parts in other Animals; embracing the subject of Shoeing and the proper Treatment of the Foot; with the Rationale and Effects of various Important Operations, and the best Methods of performing them. By W. C. Spooner, M.R.V.C. 12mo. 7*s.* 6*d.* cloth.

STABLE TALK AND TABLE TALK; OR, SPECTACLES FOR YOUNG SPORTSMEN. By Harry Hieover. 8vo. 12*s.* cloth.

* * A Second Volume, with Portrait of the Author and complete Index to both volumes, is nearly ready.

STEBBING (REV. H.).—THE HISTORY OF THE CHURCH OF CHRIST,
From the Diet of Augsburg, 1530, to the Eighteenth Century; originally designed as a Continuation of Milner's "History of the Church of Christ." By the Rev. Henry Stebbing, D.D. 3 vols. 8vo. 36*s.* cloth.

STEBBING.—THE HISTORY OF THE CHRISTIAN CHURCH,
From its Foundation to A.D. 1492. By the Rev. H. Stebbing, M.A., etc. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

STEBBING.—THE HISTORY OF THE REFORMATION.
By the Rev. H. Stebbing. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

STEEL'S SHIPMASTER'S ASSISTANT.

Compiled for the use of Merchants, Owners and Masters of Ships, Officers of Customs, and all Persons connected with Shipping or Commerce; containing the Law and Local Regulations affecting the Ownership, Charge, and Management of Ships and their Cargoes; together with Notices of other Matters, and all necessary Information for Mariners. New Edition, rewritten throughout. Edited by Graham Willmore, Esq. M.A. Barrister-at-Law; the Customs and Shipping Department by George Clements, of the Customs, London, compiler of "The Customs Guide;" The Exchanges, etc. and Naval Book-keeping, by William Tate, author of "The Modern Cambist." 8vo. 28*s.* cloth; 29*s.* bound.

STEPHENS.—A MANUAL OF BRITISH COLEOPTERA;
Or, BEETLES: containing a Description of all the Species of Beetles hitherto ascertained to inhabit Great Britain and Ireland, etc. With a complete Index of the Genera. By J. F. Stephens, F.L.S., author of "Illustrations of Entomology." Post 8vo. 14*s.* cloth.

STRONG—GREECE AS A KINGDOM:
A Statistical Description of that Country—its Laws, Commerce, Resources, Public Institutions, Army, Navy, etc.—from the Arrival of King Otho, in 1833, down to the present time. From Official Documents and Authentic Sources. By Frederick Strong, Esq., Consul at Athens for the Kingdoms of Bavaria and Hanover. 8vo. 15*s.* cloth.

SUNDAY LIBRARY:

Containing nearly one hundred Sermons by eminent Divines. With Notes, etc. by the Rev. T. F. Dibdin, D.D. 6 vols. foolscap 8vo. with 6 Portraits, 30*s.* cloth; neatly half-bound in morocco, with gilt edges, 2*l.* 12*s.* 6*d.*

SWAINSON.—A PRELIMINARY DISCOURSE ON THE STUDY OF NATURAL HISTORY. By W. Swainson, Esq. Foolscape 8vo. 6s. cloth.

By the same Author.

A TREATISE ON THE NATURAL HISTORY & CLASSIFICATION OF ANIMALS. Fcap. 8vo. 6s.

NATURAL HISTORY AND CLASSIFICATION OF QUADRUPEDS. Fcap. 8vo. 6s.

NATURAL HISTORY AND CLASSIFICATION OF BIRDS. 2 vols. fcap. 8vo. 12s.

HABITS AND INSTINCT OF ANIMALS. Fcap. 8vo. 6s.

ANIMALS IN MENAGERIES. Fcap. 8vo. 6s.

NATURAL HISTORY ETC. OF FISH, AMPHIBIA, & REPTILES. 2 vols. fcap. 8vo. 12s.

A TREATISE ON MALACOLOGY; Or, the Natural Classification of Shells and Shell-Fish. Fcap. 8vo. 6s.

HISTORY AND NATURAL ARRANGEMENT OF INSECTS. Fcap. 8vo. 6s.

A TREATISE ON TAXIDERMY; with the Biography of Zoologists, and Notices of their Works. Fcap. 8vo. 6s.

SWITZERLAND.—THE HISTORY OF SWITZERLAND.

Foolscape 8vo. with Vignette Title, 6s. cloth.

TATE.—HORATIUS RESTITUTUS;

Or, the Books of Horace arranged in Chronological Order, according to the Scheme of Dr. Bentley, from the Text of Gesner, corrected and improved. With a Preliminary Dissertation, very much enlarged, on the Chronology of the Works, on the Localities, and on the Life and Character of that Poet. By James Tate, M.A. Second edition, to which is now added, an original Treatise on the Metres of Horace. 8vo. 12s. cloth.

TATE.—THE CONTINUOUS HISTORY OF THE LIFE AND WRITINGS OF ST. PAUL, on the basis of the Acts; with Intercalary Matter of Sacred Narrative, supplied from the Epistles, and elucidated in occasional Dissertations: with the *Horæ Paulinæ* of Dr. Paley, in a more correct edition, subjoined. By James Tate, M.A., Canon Residentiary of St. Paul's. 8vo. with Map, 13s. cloth.

TAYLER (REV. CHARLES B.)—MARGARET;

Or, the Pearl. By the Rev. Charles B. Tayler, M.A. Rector of St. Peter's, Chester, author of "Lady Mary; or, Not of the World;" etc. 2d Edition. Foolscape 8vo. 6s. cloth.

TAYLER (REV. CHARLES B.)—LADY MARY; OR, NOT OF THE WORLD.

By the Rev. Charles B. Tayler, Rector of St. Peter's, Chester; author of "Margaret, or the Pearl," etc. Foolscape 8vo. 6s. 6d. cloth.

TAYLER (REV. CHARLES B.)—TRACTARIANISM NOT OF COD.

Sermons. By the Rev. C. B. Tayler, Rector of St. Peter's, and Evening Lecturer at St. Mary's, Chester; author of "Lady Mary; or, Not of the World," etc. Fcap. 8vo. 6s. cloth.

TAYLER (REV. CHARLES B.)—DORA MELDER:

A Story of Alsace. By Meta Sander. A Translation. Edited by the Rev. C. B. Tayler, author of "Margaret; or, the Pearl," etc. Fcap. 8vo., with two Illustrations, 7s. cloth.

TAYLOR (JEREMY).—BISHOP JEREMY TAYLOR'S WORKS.

With the References verified. A New and thoroughly revised Edition.

** * This Work is in the hands of a competent Editor at Oxford, and will be published in volumes, price 10s. 6d. each; to be completed in 12 volumes, each of 600 closely printed pages, published at intervals of two months. The first volume will probably be published in January 1847. Subscribers' names received by the Proprietors, and all Booksellers.*

TAYLOR.—THE STATESMAN.

By Henry Taylor, Esq., author of "Philip Van Artevelde." 12mo. 6s. 6d. boards.

THIRLWALL.—THE HISTORY OF GREECE.

By the Right Rev. the Lord Bishop of St. David's. A new Edition, revised; with Notes. Vols. I. and II. demy 8vo. with Maps, 12s. each cloth. To be completed in 8 volumes.

[Vol. III. is in the press.

** * Also, an Edition in 8 vols. fcap. 8vo. with Vignette Titles, 2l. 8s. cloth.*

THOMSON'S SEASONS.

Edited by Bolton Corney, Esq. Illustrated with Seventy-seven Designs drawn on Wood by the Members of the Etching Club. Engraved by Thompson and other eminent Engravers. Square crown 8vo. uniform with "Goldsmith's Poems," 21s. cloth; in morocco, by Hayday, 36s.

THOMSON.—AN ELEMENTARY TREATISE ON ALGEBRA,

Theoretical and Practical. By James Thomson, LL.D. Professor of Mathematics in the University of Glasgow. 2d Edition. 12mo. 5s. cloth.

THOMSON (JOHN).—TABLES OF INTEREST,

At Three, Four, Four-and-a-half, and Five per Cent., from One Pound to Ten Thousand, and from One to Three Hundred and Sixty-five Days, in a regular progression of Single Days; with Interest at all the above Rates, from One to Twelve Months, and from One to Ten Years. Also, Tables shewing the Exchange on Bills, etc. etc. etc. By John Thomson, Accountant in Edinburgh. 12mo. 8s. bound.

THOMSON.—THE DOMESTIC MANAGEMENT OF THE SICK ROOM,

Necessary, in Aid of Medical Treatment, for the Cure of Diseases. By Anthony Todd Thomson, M.D. F.L.S. etc. 2d Edition. Post 8vo. 10s. 6d. cloth.

TOMLINE (BISHOP).—AN INTRODUCTION TO THE STUDY OF THE BIBLE:

Being the First Volume of the Elements of Christian Theology; containing Proofs of the Authenticity and Inspiration of the Holy Scriptures; a Summary of the History of the Jews; an Account of the Jewish Sects; and a Brief Statement of the Contents of the several Books of the Old Testament. By the late George Tomlin, D.D. F.R.S. Lord Bishop of Winchester. 20th Edition. Foolscap 8vo. 5s. 6d. cloth.

TOMLINE (BISHOP).—ELEMENTS OF CHRISTIAN THEOLOGY;

Containing Proofs of the Authenticity and Inspiration of the Holy Scriptures; a Summary of the History of the Jews; a Brief Statement of the Contents of the several Books of the Old and New Testaments; a Short Account of the English Translations of the Bible, and of the Liturgy of the Church of England; and a Scriptural Exposition of the Thirty-Nine Articles of Religion. By George Tomline, D.D. F.R.S., Lord Bishop of Winchester. 14th Edition. With Additional Notes, and a Summary of Ecclesiastical History. By Henry Stebbing, D.D., author of "A History of the Church of Christ, from the Confession of Augsburg," etc. etc. 2 vols. 8vo. 21s. cloth.

TOMLINS — A POPULAR LAW DICTIONARY;

Familiarly explaining the Terms and Nature of English Law; adapted to the comprehension of Persons not educated for the Legal Profession, and affording Information peculiarly useful to Magistrates, Merchants, Parochial Officers, and others. By Thomas Edlyne Tomlins, Attorney and Solicitor. Post 8vo. 18s. cloth.

The whole work has been revised by a Barrister.

TOOKE.—A HISTORY OF PRICES;

With reference to the Causes of their principal Variations, from 1792 to the Present Time. Preceded by a Sketch of the History of the Corn Trade in the last Two Centuries. By Thomas Tooke, Esq. F.R.S. 3 vols. 8vo. 21. 8s. cloth.

* * Separately, Vols. I. and II. 36s.; Vol. III. 12s.

TOWNSEND (CHARLES).—THE LIVES OF TWELVE EMINENT JUDGES.

OF THE PRESENT CENTURY. By W. Charles Townsend, Esq. A.M. Recorder of Macclesfield, author of "Memoirs of the House of Commons." 2 vols. 8vo. [*In the press.*]

TROLLOPE (REV. W.).—ANALECTA THEOLOGICA:

A Critical, Philological, and Exegetical Commentary on the New Testament, adapted to the Greek Text; compiled and digested from the most approved sources, British and Foreign, and so arranged as to exhibit the comparative weight of the different Opinions on Disputed Texts. By the Rev. William Trollope, M.A. of Pembroke College, Cambridge, and formerly one of the Masters of Christ's Hospital. New Edition. 2 vols. 8vo. 11. 12s. cloth.

TURNER.—THE SACRED HISTORY OF THE WORLD,

Philosophically considered. By S. Turner, F.S.A. R.A.S.L. New Edit. 3 vols. 8vo. 42s. bds.

TURNER (SHARON).—RICHARD III.:

A Poem. By Sharon Turner, Esq., F.S.A. and R.A.S.L., author of "The History of the Anglo-Saxons," "The Sacred History of the World," etc. Foolscap 8vo. 7s. 6d. cloth.

TURNER.—THE HISTORY OF ENGLAND,

From the Earliest Period to the Death of Elizabeth. By Sharon Turner, Esq. F.A.S. R.A.S.L. New Edition. 12 vols. 8vo. 81. 3s. cloth.

* * Or in four separate portions, as follows:—

THE HISTORY of the ANGLO-SAXONS; 6th Edition. 3 vols. 8vo. 21. 5s.

THE HISTORY of ENGLAND during the MIDDLE AGES; 3d Edition. 5 vols. 8vo. 31.

THE HISTORY of the REIGN of HENRY VIII.; 3d Edition. 2 vols. 8vo. 26s.

THE REIGNS of EDWARD VI., MARY, and ELIZABETH; 3d Edition 2 vols. 8vo. 32s.

TURNER.—A TREATISE ON THE FOOT OF THE HORSE,

And a New System of Shoeing, by One-sided Nailing; and on the Nature, Origin, and Symptoms of the Navicular Joint Lameness, with Preventive and Curative Treatment. By James Turner, M.R.V.C. Royal 8vo. 7s. 6d. boards.

TURTON'S (DR.) MANUAL OF THE LAND AND FRESHWATER SHELLS OF THE BRITISH ISLANDS. A new Edition, thoroughly revised, and with considerable Additions. By John Edward Gray, Keeper of the Zoological Collection in the British Museum. Post 8vo. with Woodcuts, and 12 coloured Plates, 15s. cloth.

TWISS (DR. T.)—THE OREGON QUESTION EXAMINED,

In respect to Facts and the Law of Nations. By Travers Twiss, D.C.L. F.R.S. Professor of Political Economy in the University of Oxford, and Advocate at Doctors' Commons. 8vo. with Maps of North America and the Oregon Territory (with the Columbian River on an enlarged scale, reduced from an American Government Survey), 12s. cloth.

TYTLER (PROFESSOR). — PROFESSOR TYTLER'S ELEMENTS OF GENERAL HISTORY, Ancient and Modern, with Dr. Nares' Continuation. A new Edition, revised and continued to the Death of William IV. 8vo. with 7 Maps. 14s. cloth.

URE.—DICTIONARY OF ARTS, MANUFACTURES, AND MINES;

Containing a clear Exposition of their Principles and Practice. By Andrew Ure, M.D. F.R.S. M.G.S. M.A.S. Lond.; M. Acad. N.S. Philad.; S. Ph. Soc. N. Germ. Hanov.; Mullii. etc. etc. Third Edition, corrected. 8vo. with 1240 Woodcuts, 50s. cloth.

By the same Author,

SUPPLEMENT OF RECENT IMPROVEMENTS. 2d Edition. 8vo. 14s. cloth.

VON ORLICH (CAPT.)—TRAVELS IN INDIA,

And the adjacent Countries, in 1842 and 1843. By Capt. Leopold Von Orlich. Translated from the German by H. Evans Lloyd, Esq. 2 vols. 8vo. with coloured Frontispieces, and numerous Illustrations on Wood, 25s. cloth.

WALFORD (J. E.)—THE LAWS OF THE CUSTOMS,

Compiled by Direction of the Lords Commissioners of Her Majesty's Treasury, and published under the Sanction of the Commissioners of Her Majesty's Customs; with Notes and a General Index. Edited by J. G. Walford, Esq. Solicitor for the Customs. Printed for Her Majesty's Stationery Office, and published by Authority. 8vo. 10s. 6d. cloth.

WALKER (GEO.)—CHESS STUDIES :

Comprising 1000 Games actually Played during the last Half Century; presenting a unique Collection of Classical and Brilliant Specimens of Chess Skill in every stage of the Game, and forming an Encyclopædia of Reference. By George Walker. Medium 8vo. 10s. 6d. sewed.

WATERTON.—ESSAYS ON NATURAL HISTORY,

Chiefly Ornithology. By Charles Waterton, Esq., author of "Wanderings in South America." With an Autobiography of the Author, and a View of Walton Hall. New Edition, foolscap 8vo. 8s. cloth.

SECOND SERIES. With Continuation of Mr. Waterton's Autobiography. New Edition, fcap. 8vo. with Vignette by T. Creswick, A.R.A. 6s. 6d. cloth.

WATTS (A. A.)—LYRICS OF THE HEART,

With other Poems. By Alaric A. Watts. Illustrated by 40 highly-finished Line Engravings, from the Designs of many of the eminent modern Painters, by the best Engravers of the age. Square crown 8vo. printed and embellished uniformly with Rogers's "Italy" and "Poems," 31s. 6d. boards; or proof impressions, 63s. boards; proofs before letters, on 4to. colombier, India paper (only 50 copies printed), price 51. 5s. [At Easter.

WEBSTER.—AN ENCYCLOPÆDIA OF DOMESTIC ECONOMY;

Comprising such subjects as are most immediately connected with Housekeeping; as, The Construction of Domestic Edifices, with the modes of Warming, Ventilating, and Lighting them—A description of the various articles of Furniture, with the nature of their Materials—Duties of Servants—A general account of the Animal and Vegetable Substances used as Food, and the methods of preserving and preparing them by Cooking—Making Bread—The Chemical Nature and the Preparation of all kinds of Fermented Liquors used as Beverage—Materials employed in Dress and the Toilet—Business of the Laundry—Description of the various Wheel Carriages—Preservation of Health—Domestic Medicine, etc. etc. etc. By Thomas Webster, F.G.S., etc.; assisted by the late Mrs. Parkes, author of "Domestic Duties." 8vo. with nearly 1000 Woodcuts, 50s. cloth.

WARDLAW.—DISCOURSES ON THE PRINCIPAL POINTS OF THE SOCINIAN

CONTROVERSY—the Unity of God, and the Trinity of Persons in the Godhead—the Supreme Divinity of Jesus Christ—the Doctrine of the Atonement—the Christian Character, etc. By Ralph Wardlaw, D.D. 5th Edition. 8vo. 15s. cloth.

WEIL (DR.)—THE BIBLE, THE KORAN, AND THE TALMUD;

Or, Biblical Legends of the Mussulmans, compiled from Arabic Sources, and compared with Jewish Traditions. By Dr. G. Weil, Librarian of the University of Heidelberg, etc. Translated from the German, with occasional Notes. Post 8vo. 7s. 6d. cloth.

"It has been the aim of the learned author of this curious book to collect the most esteemed of Mahomedan legends, as given in the Koran or Arabic MSS., that we may ascertain to what extent our Scripture narratives have been corrupted by human inventions, and know on what foundation the faith of Muhomedanism is built up. For giving an authentic notion of Muhomedan belief, Dr. Weil's volume, from its connected and simple form, may be made more useful than any other, not excepting even the Koran."—Britannia.

WELSFORD (HENRY).—ON THE ORIGIN AND RAMIFICATIONS OF THE ENGLISH LANGUAGE; preceded by an Inquiry into the Primitive Seats, Early Migrations, and Final Settlements, of the principal European Nations. By Henry Welsford. 8vo. 10s. 6d.

WESTWOOD (J. O.).—AN INTRODUCTION TO THE MODERN CLASSIFICATION OF INSECTS; founded on the Natural Habits and compounding Organisation of the different Families. By J. O. Westwood, F. L. S. etc. etc. etc. 2 vols. 8vo. with numerous Illustrations, 2l. 7s. cloth.

WHITLEY (DR. JOHN).—THE LIFE EVERLASTING:

In which are considered the Intermediate Life, the New Body and the New World, the Man in Heaven, Angels, the Final Consummate Life. By John Whitley, D.D. Rector of Ballymackey, and Chancellor of Killaloe. 8vo. 9s. cloth.

WHITLEY (NICHOLAS).—THE APPLICATION OF GEOLOGY TO AGRICULTURE, and to the Improvement and Valuation of Land; with the Nature and Properties of Soils, and the Principles of Cultivation. By Nicholas Whitley, Land-Surveyor. 8vo. 7s. 6d.

WIGAN (DR. A. L.).—THE DUALITY OF THE MIND,

Proved by the Structure, Functions, and Diseases of the Brain, and by the Phenomena of Mental Derangement; and shewn to be essential to Moral Responsibility. By A. L. Wigan, M.D. 8vo. 12s. cloth.

WILBERFORCE (W.).—A PRACTICAL VIEW OF THE PREVAILING RELIGIOUS SYSTEMS OF PROFESSED CHRISTIANS, in the Higher and Middle Classes in this Country, contrasted with Real Christianity. By William Wilberforce, Esq. M.P. for the County of York. New Editions. 8vo. 8s. boards. 12mo. 4s. 6d. cloth.

WILKINSON.—THE ENGINES OF WAR, ETC.

Being a History of Ancient and Modern Projectile Instruments and Engines of Warfare and Sporting; including the Manufacture of Fire Arms, the History and Manufacture of Gunpowder, of Swords, and of the cause of the Damascus Figure in Sword Blades, with some Observations on Bronze: to which are added, Remarks on some Peculiarities of Iron, and on the extraordinary Effect produced by the Action of Sea Water on Cast Iron; with Details of the various miscellaneous Experiments. By H. Wilkinson, M.R.A.S. 8vo. 9s. cloth.

WILLIS (N.P.).—DASHES AT LIFE WITH A FREE PENCIL.

By N. P. Willis, Esq., author of "Pencilings by the Way," "Inklings of Adventure," etc. 8 vols. post 8vo. 31s. 6d. boards.

WILLOUGHBY (LADY).—A DIARY.

Purporting to be by the LADY WILLOUGHBY of the Reign of Charles I., embracing some Passages of her Domestic History from 1635 to 1648. 3d edition. Square foolscap 8vo. 8s. boards, or 18s. bound in morocco (*old style*.)

. *This volume is printed and bound in the style of the period to which The Diary refers.*

WINTER (J. W.).—THE HORSE IN HEALTH AND DISEASE:

Or, Suggestions on his Natural and General History, Varieties, Conformation, Paces, Age, Soundness, Stabling, Condition, Training, and Shoeing. With a Digest of Veterinary Practice. By James W. Winter, M.R.C.V.S.L. 8vo. 10s. 6d. cloth.

"Mr. Winter is master of his subject, and hence writes with force and distinctness. He is thoroughly cognisant of all the varieties of the horse,—of the best modes of breeding, of all forms of disease, and of the most judicious modes of treatment. On these topics and various kindred ones he discourses in this elaborate treatise with great ability, giving the results of his enlarged knowledge and experience in a pleasing and intelligible manner. From the popular style of the book, and the great amount of information it contains, it may be read with profit by every one who is interested in the judicious treatment of the horse in 'health and disease.'"—*Britannia.*

ZOOLOGY OF THE VOYAGE OF H.M.SS. EREBUS AND TERROR.

Under the Command of Capt. Sir James Clark Ross, R.N. F.R.S. during the years 1839, 40, 41, 42, 43. Published by Authority of the Lords Commissioners of the Admiralty. Edited by John Richardson, M.D. F.R.S. etc.; and John Edward Gray, Esq. F.R.S. Parts I. to XI. Royal 4to. with numerous coloured and plain Plates, 10s. each, sewed.

. *To be completed in about 15 parts.*

ZUMPT (PROF.).—A GRAMMAR OF THE LATIN LANGUAGE.

By C. G. Zumpt, Ph. D. Professor in the University, and Member of the Royal Academy of Berlin. Translated from the 9th Edition of the original, and adapted to the use of English Students, by Leonhard Schmitz, Ph. D., Rector of the High School of Edinburgh; with numerous Additions and Corrections by the Author. 8vo. 14s. cloth.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00591 8156